Ch 4.3 - Logistic Regression

Lecture 10 - CMSE 381

Prof. Elizabeth Munch

Michigan State University
::
Dept of Computational Mathematics, Science & Engineering

Fri, Sep 20, 2024

Announcements

Lec #	Date			Reading	HW
1	Mon	8/26	Intro / First day stuff / Python Review Pt 1	1	
2	Wed	8/28	What is statistical learning?	2.1	
3	Wed	9/4	Assessing Model Accuracy	2.2.1, 2.2.2	
4	Fri	9/6	Linear Regression	3.1	HW #1 Due Sun 9/8
5	Mon	9/9	More Linear Regression	3.1	
6	Wed	9/11	Multi-linear regression	3.2	
7	Fri	9/13	Probably more linear regression	3.3	Hw #2 Due Dun 9/15
8	Mon	9/16	Last of the linear regression		
9	Wed	9/18	Intro to classification, Bayes classifier, KNN classifier	2.2.3	
10	Fri	9/20	Logistic Regression	4.1, 4.2, 4.3.1-3	Hw #3 Due Sun 9/22
11	Mon	9/23	Multiple Logistic Regression / Multinomial Logistic Regression	4.3.4-5	
	Wed	9/25	***Project Day & Review***		
	Fri	9/27	**Midterm #1**		

Announcements:

- Homework #3 Due Sunday on Crowdmark
- Wednesday - Review day
  - Nothing prepped
  - Bring your questions
- Friday - Exam #1
  - Bring 8.5x11 sheet of paper
  - Handwritten both sides
  - Anything you want on it, but must be your work
  - You will turn it in

Covered in this lecture

Last Time:

- Classification basics
- Bayes classifier
- KNN classifier

This time:

- Logistic Regression

Section 1

Review from last time

Error rate

- Training data:
  $\{(x_1, y_1), \cdots, (x_n, y_n)\}$ with y_i qualitative
- Estimate $\hat{y} = \hat{f}(x)$
- Indicator variable
  $\mathrm{I}(y_i \neq \hat{y}_i)$ *is 1 if not the same (misclassified) and 0 else*

Training error rate:

$$\frac{1}{n}\sum_{i=1}^{n} \mathrm{I}(y_i \neq \hat{y}_i)$$

Test error rate:

$$\mathrm{Ave}(\mathrm{I}(y_0 \neq \hat{y}_0))$$

Good classifier: one that minimizes test error rate, GET BETTER EXAMPLE FOR THIS

Best ever classifier

We can't have nice things

Bayes Classifier:
Give every observation the highest probability class given its predictor variables

$$\Pr(Y = j \mid X = x_0)$$

- It is possible to show (though the proof is outside of the scope of this book) that the test error rate is minimized, on average, by a very simple classifier that assigns each observation to the most likely class, given its

Bayes Decision Boundary



Bayes error rate

- Error at $X = x_0$

$$1 - \max_j \Pr(Y = j \mid X = x_0)$$

- Overall Bayes error:

$$1 - E\left(\max_j \Pr(Y = j \mid X = x_0)\right)$$

This is the irreducible error



- Game: do as good a job as possible estimating the Bayes decision boundary
- Classify given observation with highest estimated probability

K-Nearest Neighbors



$K = 3$



decision boundary

- Fix K positive integer
- $N(x)$ = the set of K closest neighbors to x
- Estimate conditional proability

$$\Pr(Y = j \mid X = x_0) = \frac{1}{K} \sum_{i \in N(x_0)} \mathrm{I}(y_i = j)$$

- Pick j with highest value





Section 2

Logistic Regression

Simulated `Default` data set





- 10,000 data points
- Individuals who defaulted shown in orange (reality is deafault rate is actualy 3% so only a fraction of those who didn't default are plotted)
- Goal: function $C(balance, income) = default/not$.

What is classification

- Classification: When the response variable is qualitative
- Goal: Model the probability that Y belongs to a particular category

$$p(\texttt{balance}) = \Pr(\texttt{default} = \texttt{yes} \mid \texttt{balance})$$



Goal for `Balance` data set



Goal: Model the probability that Y belongs to a particular category

Ex.

$\Pr(\texttt{default} = \texttt{yes} \mid \texttt{balance})$

- Use notation $p(\texttt{balance})$
- Choice of threshold depends on comfort with risk
- Note that highly contrived fake data means that largely decideable by balance, so we'll just look at that

Let's just use regression!

JK that's a bad idea

Bad idea:

- Set Y to be a dummy variable taking values in $\{0, 1, 2, \cdots\}$
- Run regression, and choose k based on what integer value $\hat{y}$ is closest to

Ex.

$$Y = \begin{cases} 1 & \text{if stroke} \\ 2 & \text{if drug overdose} \\ 3 & \text{if epileptic seizure} \end{cases}$$

vs.

$$Y = \begin{cases} 1 & \text{if mild} \\ 2 & \text{if moderate} \\ 3 & \text{if severe} \end{cases}$$

Ordering implies something about closeness. Draw a number line to emphasize this distinction.

Bad idea is still not a great idea for two levels

$$p(\texttt{balance}) = \Pr(\texttt{default} = \texttt{yes} \mid \texttt{balance})$$

$$Y = \begin{cases} 0 & \text{if not default} \\ 1 & \text{if default} \end{cases}$$

- Fit linear regression
- Predict default if $\hat{y} > 0.5$; not default otherwise
- Could spit out values outside of [0,1]
- Can't interpret this as probability
- Regression methods can't accomdate qualitative response with more than two classes
- Doesn't provide meaningful estimates of $\Pr(Y \mid X)$ even with two classes



$$p(\texttt{balance}) = \beta_0 + \beta_1 \texttt{balance}$$

Approximating the probability



- Trying to approximate the probability with a function
- Draw some example functions
- Issues with linear, density on the bottom over-compenasates
- issues going outside of the [0,1] bound
- Better job if we get this sharply increasing function to differentiate the boundary

Logistic function

$$y = \frac{e^x}{1 + e^x}$$



$$p(X) = \frac{e^{\beta_0 + \beta_1 X}}{1 + e^{\beta_0 + \beta_1 X}}$$

Try it out:
desmos.com/calculator/cw1pyzzqci

Logistic Regression

$$\Pr(\texttt{default} = \texttt{yes} \mid \texttt{balance}) = \frac{e^{\beta_0 + \beta_1 \texttt{balance}}}{1 + e^{\beta_0 + \beta_1 \texttt{balance}}}$$



What will the drawn logistic regression classifer predict for each of the following values of `Balance`



Balance	Prediction
0	*No*
500	*No*
1000	*No*
1500	*Yes*
2000	*Yes*
2500	*Yes*

Odds

$$\frac{p(x)}{1-p(x)} = \frac{\Pr(Y=1 \mid X=x)}{1-\Pr(Y=1 \mid X=x)} = \frac{\Pr(Y=1 \mid X=x)}{\Pr(Y=0 \mid X=x)}$$

Probability or risk $= \frac{p}{p+q}$

Odds $= p : q$

- Logistic function is chosen so that odds are linear
- Can take any value from 0 (low odds) to ∞ (high odds)

Examples:

- If the probability of default is 90% what are the odds?
  - $p(x) = 0.9$
  - $\frac{0.9}{1-0.9} = 9$
- If the odds are $1/3$, what is the probability of default?
  - $\frac{p}{1-p} = 1/3$
  - $3p = 1 - p$
  - $4p = 1$
  - $p = 1/4$

How to get logistic function

Assume the (natural) log odds (logits) follow a linear model

$$\log\left(\frac{p(x)}{1-p(x)}\right) = \beta_0 + \beta_1 x$$

Solve for $p(x)$:

$$p(x) = \frac{e^{\beta_0+\beta_1 x}}{1+e^{\beta_0+\beta_1 x}}$$

- Note that we can't use the linear interpretation from before since one unit increase in β_1 doesn't correspond to one unit of odds if that made sense
- But can say that positive β_1 corresponds to increasing $p(X)$ and vice versa

Playing with the logistic function: desmos.com/calculator/cw1pyzzqci

Using coefficients to make predictions

	Coefficient	Std. error	z-statistic	p-value
`Intercept`	-10.6513	0.3612	-29.5	<0.0001
`balance`	0.0055	0.0002	24.9	<0.0001

What is the estimated probability of default for someone with a balance of \$1,000?

$$\hat{p}(X) = \frac{e^{\hat{\beta}_0+\hat{\beta}_1 X}}{1+e^{\hat{\beta}_0+\hat{\beta}_1 X}} = \frac{e^{-10.6513+0.0055\times 1000}}{1+e^{-10.6513+0.0055\times 1000}} = 0.006$$

What is the estimated probability of default for someone with a balance of \$2,000:

$$\hat{p}(X) = \frac{e^{\hat{\beta}_0+\hat{\beta}_1 X}}{1+e^{\hat{\beta}_0+\hat{\beta}_1 X}} = \frac{e^{-10.6513+0.0055\times 2000}}{1+e^{-10.6513+0.0055\times 2000}} = 0.586$$

Interpreting the coefficients

$$p(x) = \frac{e^{\beta_0+\beta_1 x}}{1+e^{\beta_0+\beta_1 x}}$$

$$\log\left(\frac{p(x)}{1-p(x)}\right) = \beta_0 + \beta_1 x$$

	Coefficient	Std. error	z-statistic	p-value
`Intercept`	-10.6513	0.3612	-29.5	<0.0001
`balance`	0.0055	0.0002	24.9	<0.0001

- β_1 means increasing x by one unit increases the log odds by β_1 unit
- In this case, increasing the balance by 1 means increasing the log odds of default by 0.0055

Confusion Matrix: Predicting `default` from `balance`

		True default status		
		No	Yes	Total
Predicted default status	No	9644	252	9896
	Yes	23	81	104
	Total	9667	333	10000

- Overall error $(c+d)/N = 2.75\%$
- This table is from the LDA section, but we can use this for understanding confusion matrix

		True		
		Yes	No	Total
Predicted	Yes	a	b	$a+b$
	No	c	d	$c+d$
	Total	$a+c$	$b+d$	N

Do coding in jupyter notebook

Next time

Lec #	Date			Reading	HW
1	Mon	8/26	Intro / First day stuff / Python Review Pt 1	1	
2	Wed	8/28	What is statistical learning?	2.1	
3	Wed	9/4	Assessing Model Accuracy	2.2.1, 2.2.2	
4	Fri	9/6	Linear Regression	3.1	HW #1 Due Sun 9/8
5	Mon	9/9	More Linear Regression	3.1	
6	Wed	9/11	Multi-linear regression	3.2	
7	Fri	9/13	Probably more linear regression	3.3	Hw #2 Due Dun 9/15
8	Mon	9/16	Last of the linear regression		
9	Wed	9/18	Intro to classification, Bayes classifier, KNN classifier	2.2.3	
10	Fri	9/20	Logistic Regression	4.1, 4.2, 4.3.1-3	Hw #3 Due Sun 9/22
11	Mon	9/23	Multiple Logistic Regression / Multinomial Logistic Regression	4.3.4-5	
	Wed	9/25	***Project Day & Review***		
	Fri	9/27	**Midterm #1**		