Ch 6.3: PCR

Lecture 20 - CMSE 381

Prof. Elizabeth Munch

Michigan State University
::
Dept of Computational Mathematics, Science & Engineering

Fri, Oct 18, 2024

Announcements

Last time:

- PCA

This lecture:

- PCR

Announcements:

- Exam #2 on Friday!
  - Bring 8.5x11 sheet of paper
  - Handwritten both sides
  - Anything you want on it, but must be your work
  - You will turn it in
  - Non-internet calculator if you want it

Lec #	Date			Reading	HW
12	Mon	9/30	Leave one out CV	5.1.1, 5.1.2	
13	Wed	10/2	k-fold CV	5.1.3	
14	Fri	10/4	More k-fold CV,	5.1.4-5	
15	Mon	10/7	k-fold CV for classification	5.1.5	
16	Wed	10/9	Subset selection	6.1	HW #4 Due Weds 10/9
17	Fri	10/11	Shrinkage: Ridge	6.2.1	
18	Mon	10/14	Shrinkage: Lasso	6.2.2	
19	Wed	10/16	Dimension Reduction	6.3	
20	Fri	10/18	Overflow, Possibly more dimension reduction?		HW #5 Due Fri 10/18
	Mon	10/21	No class - Fall break		
	Wed	10/23	**Review**		
	Fri	10/25	**Midterm #2**		

Section 1

Last time

Shrinkage

Find β to minimize

$$RSS = \sum_{i=1}^{n} \left(y_i - \beta_0 - \sum_{j=1}^{p} \beta_j x_{ij} \right)^2$$

subject to:

Least Squares:
No constraints

Ridge:

$$\sum_{j=1}^{p} \beta_j^2 \leq s$$

The Lasso:

$$\sum_{j=1}^{p} |\beta_j| \leq s$$





Also, find best choice of λ or s using CV

Linear transformation of predictors

Original Predictors:

$$X_1, \cdots, X_p$$

New Predictors:

$$Z_1, \cdots, Z_M$$

$$Z_m = \sum_{j=1}^{p} \varphi_{jm} X_j$$

The goal:

- Find good φ's (PCA)
- Fit regression model on Z_i's using least squares (PLS)

$$y_i = \theta_0 + \sum_{m=1}^{M} \theta_m z_{im} + \varepsilon_i$$

- Hope that lower dimensions means less overfitting

PCA - First PC



higher variance of projection on the further right pic

Projection onto first PC



$$Z_1 = 0.839 \cdot (\mathtt{pop} - \overline{\mathtt{pop}}) + 0.544 \cdot (\mathtt{ad} - \overline{\mathtt{ad}})$$

Drawing points in PC space



Section 2

Principal Components Regression

So you've found your PCA coefficients

Now what?

- Do linear regression on $Z_1, \cdots, Z_M$
- Book calls this "principal components regression "
- Linear model is $y = \theta_0 + \theta_1 Z_1 + \cdots + \theta_M Z_M$

What are we assuming?

- The directions in which $X_1, \cdots, X_p$ have the most variation are the directions that are associated with Y
- If assumption holds, then fitting on Z_i better than fitting on X_i since fewer variables lessens chance of overfitting

Interpretation of PCR coefficients

Original Predictors:

$$X_1, \cdots, X_p$$

New Predictors:

$$Z_1, \cdots, Z_M$$

$$Z_m = \sum_{j=1}^{p} \varphi_{jm} X_j$$

Learned model:

$$y = \theta_0 + \theta_1 Z_1 + \cdots + \theta_M Z_M$$

- Do example showing that coefficients of Z_i can't be used to say anything about X_i's
- $p = 3, M = 2$
- $\varphi = \begin{pmatrix} 1 & -1 \\ 3 & 1/2 \\ 0 & 1/2 \end{pmatrix}$
- Learned model $y = 2Z_1 - Z_2$
- $y = 2(X_1 + 3X_2 + 0X_3) - (-X_1 + \frac{1}{2}X_2 + \frac{1}{2}X_3)$
- $= 3X_1 + 2.5X_2 - 0.5X_3$
- Conclusion: can't directly use coefficients learned to interpret original inputs!
- Each PC is a linear combination of the originals, so can go backwards to figure out whaat the coefficients would have been, see left figure

Picking M





- Credit data set
- Major drop in MSE at 10 compoentns, so barely a savings with the original 11 variables

Do PCR with hitters data

Doing better

Example with simulated data: $n = 50$ observations of $p = 45$ predictors

Y is a function of **2 predictors**

- Right side is usual least squares estimate
- Typical U shape for Test MSE
- PCR with good choice of number of components improvement over plain least squares



Doing better

Example with simulated data: $n = 50$ observations of $p = 45$ predictors
Y is a function of **all predictors**



- Right side is usual least squares estimate
- Even better improvement using PCR than previous
- note the chnage in y-axis values

Comparison to results on shrinkage

Y is a function of all predictors



Y is a function of 2 predictors



Shrinkage does better

Properties of PCR

- Note better in previous example because lots of PCs needed
- It does better when info contained in the first few PCs
- Number of PCs to use can be picked with CV
- Standardization needed to make sure big data values don't skew the results
- Not a feature selection method because we don't get a subset of features, we get a collection of linear combos of features

TL;DR

PCR

- Unsupervised dimensionality reduction + linear regression
- Choose component Z_1 in the direction of most variance using only X_i's information
- Choose Z_2 and beyond by the same method after "getting rid" of info in the directions already explained



Next time

Lec #	Date			Reading	HW
12	Mon	9/30	Leave one out CV	5.1.1, 5.1.2	
13	Wed	10/2	k-fold CV	5.1.3	
14	Fri	10/4	More k-fold CV,	5.1.4-5	
15	Mon	10/7	k-fold CV for classification	5.1.5	
16	Wed	10/9	Subset selection	6.1	HW #4 Due Weds 10/9
17	Fri	10/11	Shrinkage: Ridge	6.2.1	
18	Mon	10/14	Shrinkage: Lasso	6.2.2	
19	Wed	10/16	Dimension Reduction	6.3	
20	Fri	10/18	Overflow, Possibly more dimension reduction?		HW #5 Due Fri 10/18
	Mon	10/21	No class - Fall break		
	Wed	10/23	**Review**		
	Fri	10/25	**Midterm #2**		