Ch 7.2-7.4: Step Functions, Basis Functions, Start Splines

Lecture 22 - CMSE 381

Prof. Elizabeth Munch

Michigan State University
::
Dept of Computational Mathematics, Science & Engineering

Weds, Oct 30, 2024

Announcements

I split this class in two because it was too much last time, but now this is way too short. Or maybe I was just tired and skipping too much. Perhaps they should actually code the classification example?

Last time:

- 7.1 Polynomial regression
- 7.2 Step functions

This lecture:

- 7.2 Step functions
- 7.3 Basis functions
- 7.4 Regression Splines (Finish next lecture)

Announcements:

- HW #6 Due Sunday
- Projects

Section 1

Last time

Polynomial regression

Replace linear model

$$y_i = \beta_0 + \beta_1 x_1 + \varepsilon_i$$

with

$$y_i = \beta_0 + \beta_1 x_1 + \beta_2 x_i^2 + \cdots + \beta_d x_i^d + \varepsilon_i$$

- Can learn with linear models by passing in predictors x_i^ℓ
- Tend to not go higher than degree 3 or 4 because makes overly flexible

Example with wage data



- Plot of wage vs age for men in central Atlantic region of the US
- Dark line is degree 4 polynomial
- Have variance for each coefficient
- Assume you have a model $\hat{f}(x_0) = \hat{\beta}_0 + \hat{\beta}_1 x_0 + \cdots + \hat{\beta}_d x_0^d$
- Can use that to estimate the pointwise variance $\mathrm{Var}(\hat{f}(x_0))$
- Draw 2 std deviations away from the line, 95% confidence interval

$$-184.1542 + 21.24552 * age + -0.56386 * age^2 + 0.00681 * age^3 + -3e-05 * age^4$$

Step functions

- $\mathrm{I}(X < c)$
- $\mathrm{I}(c_1 \leq X < c_2)$
- $\mathrm{I}(c \leq X)$

$$
\begin{aligned}
C_0(X) &= I(X < c_1), \\
C_1(X) &= I(c_1 \leq X < c_2), \\
C_2(X) &= I(c_2 \leq X < c_3), \\
&\vdots \\
C_{K-1}(X) &= I(c_{K-1} \leq X < c_K), \\
C_K(X) &= I(c_K \leq X),
\end{aligned}
$$

Learned model:

$$y_i = \beta_0 + \beta_1 C_1(x_i) + \beta_2 C_2(x_i) + \cdots + \beta_K C_K(x_i) + \varepsilon_i$$

Example: Cut points at -4, -1, 3, 6

$C_1(X)$



$C_0(X)$



$C_2(X)$



$C_4(X)$



$C_3(X)$



Step function example



- Learned peak where the high earners are

Section 2

Classification versions

Remember logisitic regression?

$$y = \frac{e^x}{1 + e^x}$$



$$p(X) = \frac{e^{\beta_0 + \beta_1 X}}{1 + e^{\beta_0 + \beta_1 X}}$$

Multiple features:

$$p(X) = \frac{e^{\beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p}}{1 + e^{\beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p}}$$

Classification version: Polynomial regression

$$\Pr(y_i > 250 \mid x_i) = \frac{\exp(\beta_0 + \beta_1 x_i + \cdots + \beta_d x_i^d)}{1 + \exp(\beta_0 + \beta_1 x_i + \cdots + \beta_d x_i^d)}$$



- Note in previous fig that there is a distinct subcluster of high earners making more than \$250K
- Build a logistic regression model as above
- Note that the 95% confidence interval gets very wide on the right side
- Large sample size $n = 3,000$ but small number (79) of high earners makes for high variance in coeffs and wide confidence intervals

Classification version: Step functions

$$\Pr(y_i > 250 \mid x_i) = \frac{\exp(\beta_0 + \beta_1 C_1(x_i) + \beta_2 C_2(x_i) + \cdots + \beta_K C_K(x_i))}{1 + \exp(\beta_0 + \beta_1 C_1(x_i) + \beta_2 C_2(x_i) + \cdots + \beta_K C_K(x_i))}$$



- Again, learns that middle bit
- Still big confidence interval on the right, but likely because there's just less data over there

Coding bit: classification version



Just talk through on projector, there's nothing in there for them to code

A few more comments on step functions

- Gives the chance to break up the domain, avoid forcing global structure
- Need to make decisions about the c_i. A bit arbitrary unless your data has natural breakpoints.
- Popular in biostats and epidemiology

Section 3

Basis functions

Basis Functions Setup

Polynomial and piecewise-constant regression models are special cases of a *basis function* approach.

$$y_i = \beta_0 + \beta_1 b_1(x_i) + \beta_2 b_2(x_i) + \cdots + \beta_K b_K(x_i) + \varepsilon_i$$

- Pick a collection of basis functions $b_1(X), \cdots, b_K(X)$
- Use least squares to figure out the constants
- Explain the b_i's for polynomial and stepwise functions

- Lots of possible options for these
  - Some examples are wavelets or Fourier series
  - Next section is we'll talk about regression splines

Section 4

Regression Splines

Piecewise polynomials

- Fit a polynomial regression

$$y_i = \beta_0 + \beta_1 x_1 + \beta_2 x_i^2 + \cdots + \beta_d x_i^d + \varepsilon_i$$

- Let the β_i's be different at different locations of the range.

- Points where the coefficients change are called "knots"
- A piecewise cubic with no knots is just standard cubic polynomial
- More knots gives more flexible polynomial
- Piecewise constant of previous section is degree 0 version of this

Example of piecewise polynomial



Example:

$$y_i = \begin{cases} \beta_{01} + \beta_{11}x_i + \beta_{21}x_i^2 + \beta_{31}x_i^3 + \epsilon_i & \text{if } x_i < c \\ \beta_{02} + \beta_{12}x_i + \beta_{22}x_i^2 + \beta_{32}x_i^3 + \epsilon_i & \text{if } x_i \geq c. \end{cases}$$

- Uses 8 degrees of freedom to fit the model
- problem: we have a discontinuous function.

The fix

- Fit piecewise polynomial
- Require continuity at knots



Looks better, but pointy join is odd

The better fix: Cubic splines

- Fit piecewise polynomial
- Require continuity at knots
- Require the first and second derivatives to be continuous at knots
- Now we have smooth joins
- This is called a "cubic spline"
- 8 original degrees of freedom, plus three constraints (continuous, 1st deriv continuous, 2nd deriv continuous) leaves 5 degrees of freedom
- cubic spline with K knots uses $4 + K$ degrees of freedom



Example

We have the following piecewise cubic polynomial:

$$f(x) = \begin{cases} 2 + x + x^2 + 0.1x^3 & x \leq 1 \\ b_0 + b_1 x + b_2 x^2 - x^3 & x > 1 \end{cases}$$

What are b_1, b_1, and b_2 to make this a cubic spline?



Check your answers: desmos.com/calculator/ns4tr7mw0n

More space for work

$$f(x) = \begin{cases} 2 + x + x^2 + 0.1x^3 & x \leq 1 \\ b_0 + b_1 x + b_2 x^2 - x^3 & x > 1 \end{cases}$$

Instructor tool for generating solution while messing with b_3: `https://www.desmos.com/calculator/p9xzbkzqmf`

$$f(x) = \begin{cases} 2 + x + x^2 + 0.1x^3 & x \leq 1 \\ 3.1 + -2.3x + 4.3x^2 - x^3 & x > 1 \end{cases}$$



Next time

Lec #	Date			Reading	HW
21	Mon	10/28	Polynomial & Step Functions	7.1,7.2	
22	Wed	10/30	Step Functions; Basis functions; Start Splines	7.2 - 7.4	
23	Fri	11/1	Regression Splines	7.4	HW #7 Due Sun 11/3
24	Mon	11/4	Decision Trees	8.1	
25	Wed	11/6	Class Cancelled (Dr Munch out of town)		
26	Fri	11/8	Random Forests	8.2.1, 8.2.2	HW #8 Due Sun 11/10
27	Mon	11/11	Maximal Margin Classifier	9.1	
28	Wed	11/13	SVC	9.2	
29	Fri	11/15	SVM	9.3, 9.4	HW #9 Due Sun 11/17
30	Mon	11/18	Single layer NN	10.1	
31	Wed	11/20	Multi Layer NN	10.2	
32	Fri	11/22	CNN	10.3	HW #10 Due Sun 11/24
33	Mon	11/25	TBD: Unsupervised learning/clustering	12.1, 12.4?	
	Wed	11/27	Virtual: Project office hours		