Ch 9.1: Maximum Margin Classifier

Lecture 26 - CMSE 381

Prof. Elizabeth Munch. Substitute: Rachel Roca

Michigan State University
::
Dept of Computational Mathematics, Science & Engineering

Mon, Nov 11, 2024

Announcements

Last time:

- Ch 8: Random Forests

This lecture:

- Maximal Margin Classifier
-
- No jupyter notebook for this class

Announcements:

- HW #8 Due Sunday 11/17

Lec #	Date			Reading	HW
21	Mon	10/28	Polynomial & Step Functions	7.1,7.2	
22	Wed	10/30	Step Functions; Basis functions; Start Splines	7.2 - 7.4	
23	Fri	11/1	Regression Splines	7.4	HW #6 Due Sun 11/3
24	Mon	11/4	Decision Trees	8.1	
25	Wed	11/6	Class Cancelled (Dr Munch out of town)		
26	Fri	11/8	Random Forests	8.2.1, 8.2.2	HW #7 Due Sun 11/10
27	Mon	11/11	Maximal Margin Classifier	9.1	
28	Wed	11/13	SVC	9.2	
29	Fri	11/15	SVM	9.3, 9.4	HW #8 Due Sun 11/17
30	Mon	11/18	Single layer NN	10.1	
31	Wed	11/20	Multi Layer NN	10.2	
32	Fri	11/22	CNN	10.3	HW #11 Due Sun 11/24
33	Mon	11/25	TBD: Unsupervised learning/clustering	12.1, 12.4?	
	Wed	11/27	Virtual: Project office hours		
	Fri	11/29	No class - Thanksgiving		
	Mon	12/2	Review		
	Wed	12/4	**Midterm #3**		
	Fri	12/6	No class - EGR Design Day		Project due

Section 1

Maximal Margin Classifier

The goal



What is a hyperplane?

Mathematical definition of a hyperplane

$$\beta_0 + \beta_1 X_1 + \beta_2 X_2 + \cdots + \beta_p X_p = 0$$

Hyperplane for $p = 2$



$$1 + 2X_1 + 3X_2 = 0$$

There are two sides to every hyperplane

$$\beta_0 + \beta_1 X_1 + \beta_2 X_2 + \cdots + \beta_p X_p < 0 \qquad \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \cdots + \beta_p X_p > 0$$



Classification Setup

Data matrix:

$$X = \begin{pmatrix} - & x_1^T & - \\ - & x_2^T & - \\ & \vdots & \\ - & x_n^T & - \end{pmatrix}_{n \times p}$$

Observations in one of two classes, $y_i \in \{-1, 1\}$

$$Y = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix}$$

$$x_1 = \begin{pmatrix} x_{11} \\ \vdots \\ x_{1p} \end{pmatrix}, \cdots, x_n = \begin{pmatrix} x_{n1} \\ \vdots \\ x_{np} \end{pmatrix}$$

Separate out a test observation

$$x^* = (x_1^* \ \cdots \ x_p^*)^T$$

Separating Hyperplane

$$\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} > 0 \text{ if } y_i = 1$$
$$\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} < 0 \text{ if } y_i = -1$$



Another way to say it

$$\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} > 0 \text{ if } y_i = 1$$
$$\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} < 0 \text{ if } y_i = -1$$

For all i:

$$y_i(\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip}) > 0$$

Separating hyperplane becomes a classifier

If you have a separating hyperplane:

- Check

$$f(x^*) = \beta_0 + \beta_1 x_1^* + \beta_2 x_2^* + \cdots + \beta_p x_p^*$$

- If positive, assign $\hat{y} = 1$
- If negative, assign $\hat{y} = -1$



How do we pick?



Distance from an observation to a hyperplane





Maximal margin classifier



Example



- Sketch the maximal margin hyperplane.
- What is the equation of this line in the form $\beta_0 + \beta_1 X_1 + \beta_2 X_2 = 0$?
- Circle the support vectors. What is their distance from the line?

desmos.com/calculator/lqms253gfq

Extra work space



desmos.com/calculator/lqms253gfq

Mathematical Formulation

$$\underset{\beta_0,\beta_1,\ldots,\beta_p,M}{\text{maximize}} \; M$$

$$\text{subject to } \sum_{j=1}^{p} \beta_j^2 = 1,$$

$$y_i(\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip}) \geq M \;\; \forall\, i = 1, \ldots, n$$

First constraint



Second constraint



$$y_i(\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2}) \geq M$$

- Blue circles: $y_i = -1$
- Red Xs: $y_i = 1$
- $-2\sqrt{2} + \frac{\sqrt{2}}{2}X_1 + \frac{\sqrt{2}}{2}X_1 = 0$

An example with a bad choice of hyperplane



What is $y_i(\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2})$ for the point $x_i = (0, 3)$?

- Blue circles: $y_i = -1$
- Red Xs: $y_i = 1$
- $-\frac{4}{\sqrt{5}} + \frac{1}{\sqrt{5}} X_1 + \frac{2}{\sqrt{5}} X_1 = 0$

Second constraint extra space



$$y_i(\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2}) \geq M$$

- Blue circles: $y_i = -1$
- Red Xs: $y_i = 1$
- $-2\sqrt{2} + \frac{\sqrt{2}}{2}X_1 + \frac{\sqrt{2}}{2}X_1 = 0$

Mathematical Formulation

$$\underset{\beta_0,\beta_1,\ldots,\beta_p,M}{\text{maximize}} \; M$$

$$\text{subject to } \sum_{j=1}^{p} \beta_j^2 = 1,$$

$$y_i(\beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip}) \geq M \;\; \forall\, i = 1, \ldots, n$$

Section 2

Issues with Maximal Margin Classifier

But what if....



Sensitivity to new points



Next time

Lec #	Date			Reading	HW
21	Mon	10/28	Polynomial & Step Functions	7.1,7.2	
22	Wed	10/30	Step Functions; Basis functions; Start Splines	7.2 - 7.4	
23	Fri	11/1	Regression Splines	7.4	HW #6 Due Sun 11/3
24	Mon	11/4	Decision Trees	8.1	
25	Wed	11/6	Class Cancelled (Dr Munch out of town)		
26	Fri	11/8	Random Forests	8.2.1, 8.2.2	HW #7 Due Sun 11/10
27	Mon	11/11	Maximal Margin Classifier	9.1	
28	Wed	11/13	SVC	9.2	
29	Fri	11/15	SVM	9.3, 9.4	HW #8 Due Sun 11/17
30	Mon	11/18	Single layer NN	10.1	
31	Wed	11/20	Multi Layer NN	10.2	
32	Fri	11/22	CNN	10.3	HW #11 Due Sun 11/24
33	Mon	11/25	TBD: Unsupervised learning/clustering	12.1, 12.4?	
	Wed	11/27	Virtual: Project office hours		
	Fri	11/29	No class - Thanksgiving		
	Mon	12/2	Review		
	Wed	12/4	**Midterm #3**		