

SPARSE SAMPLING AND INVERSE CAPACITY IN SAFE FEEDBACK SYSTEMS

ZIQUING DING

ABSTRACT. We study the complexity of actual safe feedback names when a control period is fixed and observations are made at selected block times. The strategy class contains all legal Borel selectors and all finite Borel atom refinements. Two mechanisms give exact optimization results. For a finite closed cover of a mixing one-sided shift of finite type, every prescribed sampling sequence with diverging gaps detects the minimum cover cardinality in every Borel refinement. For branches with feedback-dependent successors, we introduce a criterion in terms of products of inverse-capacity operators over global label lists. Exponential decay of these products forces positive-density multivalued independence in the actual names of every feedback and gives the exact minimum maximal-pattern rate. Finite-cylinder systems provide explicit structural applications: contraction follows from source weights, cross-layer dispersion, or compensation of gain columns by unavoidable loss. The last class has no common positive nonstrict left budget in its original cylinder representation. We also retain the countable-language common-sampling theorem, symbolic safety realization, a direct Thue–Morse separation, and the collapse when control periods are optimized. Exact examples and conditional results delimit these conclusions. The inverse-capacity criterion does not itself provide a feedback-independent sampling sequence, and the general fixed-period Borel minimax problem remains open.

CONTENTS

1. Introduction	2
2. Entropy background and the control interface	4
3. Control systems and invariant name languages	5
4. Simultaneous attainment for pattern systems	7
5. A language-quotient minimax theorem	9
6. Realizing one-sided subshifts as safety systems	11
7. A strict Thue–Morse separation	13
8. Comparison, all periods, and limitations	14
8.1. Comparison with Kawan’s invariance entropy	14
8.2. Why the unrestricted all-period infimum collapses	15
8.3. Countability, trajectories, and physical time	15
9. Counting entropy and the closed-cover theorem	16
9.1. A category and concatenation lemma	17
9.2. Proof of the exact formula	18
10. A cover–partition gap at prescribed sparse times	18
11. Fixed-period safe feedbacks with a common successor	19
11.1. A positive overlapping example at period two	20
12. A feedback-dependent boundary with an exact full-class value	21
13. Inverse capacities and a full-feedback pattern theorem	22
13.1. Branch measures, lists, and actual fibers	22
13.2. The finite combinatorial input and its application	24
13.3. Single-step scalar and joint-density criteria	25

Date: October 8, 2026.

2020 Mathematics Subject Classification. 93C55, 37B10, 37B40, 94A17.

Key words and phrases. safe feedback, sequence entropy, maximal pattern entropy, Borel partition, inverse capacity, symbolic dynamics, matrix products.

13.4.	A smooth joint-density example	25
13.5.	An exact product separation from every single-step reference test	26
14.	Finite-cylinder safety and structural source budgets	27
14.1.	Exact finite representation without restricting feedbacks	27
14.2.	A strict source-weight criterion	28
14.3.	Cross-layer dispersion without a strict left budget	29
15.	Gain columns compensated by unavoidable cross-layer loss	30
15.1.	Original safety data and the count condition	30
15.2.	Complete estimate for every pair of global lists	31
15.3.	A family for arbitrary control counts	32
15.4.	A complete three-control gain witness	33
16.	Affine examples, common sampling, and sharp single-step boundaries	34
16.1.	An integer affine family with a genuine absorbing feedback	34
16.2.	A protected bridge giving a predetermined common sampling	35
16.3.	A critical boundary of the single-step density condition	36
16.4.	The noninteger four-branch system and what remains open	36
17.	Expansion, absence of continuous common factors, and phase obstructions	37
17.1.	Expansion extensions of the integer family	37
17.2.	An alternating open-loop language with no Borel realization	38
Appendix A.	Exact even-sampling entropy of the real greedy feedback	39
	Inverse paths furnishing the matching lower bound	40
Appendix B.	Arithmetic sampling and finite-block optimization	41
Appendix C.	Real itinerary closures and the unresolved low-entropy implementation	41
C.1.	A closed shift-invariant interval-covering language	42
C.2.	The exact forced-predecessor compatibility kernel	43
Appendix D.	A bounded-density obstruction to simultaneous scalar capacities	44
Appendix E.	Conditional real returns and closed viability boundaries	45
E.1.	A real two-branch certificate and one common sampling	45
E.2.	Nested viable sets and the limits of a growth route	46
Appendix F.	Scope, proof status, and open problems	47
	Use of AI tools	47
	References	47

1. INTRODUCTION

A safe feedback rule makes a legal control choice from the current state, repeats this rule at a fixed control period, and thereby produces an infinite instruction name. Counting these names at sparse observation times asks a different question from counting controls per unit of elapsed time. One observation records a whole control block; the lengths of the unobserved gaps do not enter the normalization. Two optimizations must therefore be distinguished. One first chooses a sampling sequence and then minimizes over feedbacks. The other first maximizes the pattern rate of each feedback over finite time windows and then minimizes over feedbacks. Throughout the paper the control period is fixed whenever these quantities are compared.

Write $L_\tau = \sup_A \inf_s h_A(s)$ and $M_\tau = \inf_s h^*(s)$, with precise definitions below. The inequality $L_\tau \leq M_\tau$ is immediate. Obtaining a lower bound for M_τ allows the useful coordinates to depend on the feedback. Obtaining the same lower bound for L_τ requires a sampling sequence chosen *before* every Borel feedback. This quantifier difference remains substantial even for finite controls and compact state spaces. Borel selectors may depend on the whole state, may split every overlap into arbitrary Borel pieces, and need not generate continuous closed-loop maps.

The principal feedback-dependent result is Theorem 13.2. For each legal control block u , the inverse-capacity operator L_u is the density of the safe pushforward of a reference measure. If every product obtained by retaining at most $k - 1$ global labels at each time has exponentially decreasing capacity, then the true length- n names of *every* legal Borel feedback have k -valued independent coordinates of density bounded below uniformly in that feedback. In particular $h^*(s) \geq \log k/\tau$. When $k = r$, the minimum number of closed safety domains needed to cover the state space, an ordered subcover selector gives the matching upper bound and $M_\tau = \log r/\tau$ with an actual minimizer.

The capacity estimate is connected to the feedback by its real, mutually disjoint initial-state fibers. Every one of these fibers is included, including nonempty fibers of zero reference measure. A label box may contain many unrealizable open-loop words; this causes no problem because the operators retain all safety restrictions. Conversely, an open-loop cover is never asserted to be a feedback realization. The finite combinatorial independence theorem used at this step is already contained in Huang and Ye [4, Section 4, Lemma 4.1 and Remark 4.2]. The new assertions concern its connection to safe inverse capacities and the resulting optimization over a full Borel feedback class, rather than a new Sauer lemma or a new theory of combinatorial independence.

For symbolic systems with branches $P_u\sigma$ and finite-cylinder safety domains, these capacities admit an exact finite-dimensional representation, although the actual feedbacks remain unrestricted Borel maps on an infinite state space. Sections 14 and 15 give three structural applications. The first uses source degrees and permutation landing sets to construct a common strict left budget. The second proves loss across layers even when no common strict left budget exists. The third permits gain columns and proves their compensation by the mass which a large allowed tail must send into the next list's loss class. That entire class has no positive common nonstrict left budget in the original first-coordinate representation. The distinction is exact; it is not inferred merely from seeing a column sum larger than one. A three-control example separates the product criterion from every direct single-step criterion under every full-support reference probability.

These estimates use established positive-operator and matrix-product methods. Shi et al. [11] study sub- and super-stochastic products, and Debauche et al. [1] develop comparisons of path-complete Lyapunov templates. The proofs here do not apply their theorems with missing hypotheses. They establish loss directly from the original allowed cylinders, for all histories of global lists. Neither a new general stability tool nor a historical priority claim for loss compensation is asserted.

A separate result, Theorem 9.1, deals with prescribed common sampling. For every finite closed cover of a mixing one-sided shift of finite type, every increasing sequence whose gaps tend to infinity gives the exact minimum $\log r$ over *all* Borel refinements. Its common-successor control application yields $L_\tau = M_\tau = \log r/\tau$. The proof combines nonmeager atoms, closed safety domains, openness, and finite-type concatenation. Romagnoli [10] already exhibits a cover-partition entropy gap for an excluded-letter cover. Our statement allows Borel refinements and a sparse sequence prescribed in advance. It does not resolve Romagnoli's consecutive-time Borel-refinement question.

Common sampling thresholds also have precedents. Huang, Maass, and Ye [3, Lemma 3.3 and Theorem 3.5] give one sequence with a common positive measure-entropy threshold for refinements of an appropriate measurable cover. Thus the mere existence of a common positive threshold is not claimed as new here. Their invariant-measure threshold and our exact combinatorial closed-cover value have different hypotheses and normalizations. Kerr and Li [8, 9] place these questions in the general independence framework. The finite combinatorial arguments do not require a continuous or measure-preserving Borel closed loop.

The classical sequence-entropy background is due to Goodman [2], with subsequence generators studied by Hulse [5]; maximal pattern complexity for discrete systems was developed by Kamae and Zamboni [6].

We include the foundational language results in full. A deletion argument constructs one sampling sequence attaining the maximal-pattern rates of any countable family of finite-alphabet pattern systems, with no shift or compactness assumption. It gives a minimax identity for countably many invariant languages up to relabeling. Every one-sided subshift can also be realized as a matching-control safety system, and a direct binary computation gives zero invariance entropy but period-one maximal-pattern infimum $\log 2$ for the Thue–Morse example. When all control periods are allowed, the pattern infimum instead collapses to the exact-containment invariance entropy of Kawan [7]. These statements explain why the period is fixed in the structural theorems.

The two principal mechanisms must not be conflated. The closed-cover theorem gives a predetermined common sampling sequence, but requires common successors. The product-capacity theorem allows feedback to change the future orbit, but supplies feedback-dependent independent positions. An exact feedback-dependent example shows that the cover number alone need not determine either optimized value. Further affine examples, a critical capacity boundary, and appendices retain the complete numerical and realization results needed to describe the limits of the theory. Unresolved implementation and common-sampling questions are stated as such, rather than promoted to entropy theorems.

All upper bounds used in feedback optimization come from a fully defined, safe, memoryless Borel rule. A finite refinement projects to its physical control labels and therefore inherits a lower bound. It may add patterns, so an upper bound for an unrefined rule is not assigned to its arbitrary refinements.

2. ENTROPY BACKGROUND AND THE CONTROL INTERFACE

Write $\mathbb{N} = \{1, 2, \dots\}$ and $\mathbb{N}_0 = \{0, 1, 2, \dots\}$. All logarithms in this paper are natural. For a finite open cover $\mathcal{V}$ of a compact space, let $N(\mathcal{V})$ be the minimum cardinality of a subcover. The join of finitely many covers consists of all intersections of one member from each cover, with empty intersections removed. For a continuous map $T : Z \rightarrow Z$ and a finite open cover $\mathcal{V}$, ordinary cover entropy is

$$h_{\text{top}}(T, \mathcal{V}) = \lim_{n \rightarrow \infty} \frac{1}{n} \log N \left(\bigvee_{i=0}^{n-1} T^{-i} \mathcal{V} \right).$$

Submultiplicativity of cover numbers gives this limit. Taking the supremum over finite open covers defines $h_{\text{top}}(T)$. For $A = (a_i)_{i \geq 1}$ with $0 \leq a_1 < a_2 < \dots$, set

$$h_{\text{top}}^A(T, \mathcal{V}) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log N \left(\bigvee_{i=1}^n T^{-a_i} \mathcal{V} \right),$$

$$p_{Z, \mathcal{V}}^*(n) = \max_{\substack{F \subset \mathbb{N}_0 \\ \#F=n}} N \left(\bigvee_{i \in F} T^{-i} \mathcal{V} \right),$$

$$h_{\text{top}}^*(T, \mathcal{V}) = \lim_{n \rightarrow \infty} \frac{1}{n} \log p_{Z, \mathcal{V}}^*(n).$$

The sequence in the last line is subadditive before division by n ; the arbitrary sequence in the first line need not have that property. This is why a limit and a limsup appear in different definitions.

Huang–Ye’s Theorem 2.1(1) states, under their convention that T is a homeomorphism of a compact metric space, that

$$h_{\text{top}}^*(T, \mathcal{V}) = \max_A h_{\text{top}}^A(T, \mathcal{V}). \quad (2.1)$$

Its proof selects long nearly maximizing time sets and separates them by translations. Surjectivity preserves the cover number under a common translation. The previous sets contribute a vanishing fraction of the selected observations. For a non-surjective map the translation step need not preserve cover numbers; our proof uses deletion instead.

If $\mathcal{V}$ is a clopen partition, the join count is exactly the number of names seen on the selected coordinates. An arbitrary open cover may have overlaps, so its minimum subcover count must not be replaced by the number of nonempty intersections. Similarly, the measurable version of (2.1) uses Shannon entropy $H_\mu(\alpha) = -\sum_{D \in \alpha} \mu(D) \log \mu(D)$, not the logarithm of a support cardinality. Our language counts use the latter quantity. Theorem 4.2 is a counting statement, not a measurable common-sequence theorem.

For a finite-alphabet subshift Y , the first-coordinate cylinders give one particular clopen partition. Its coordinate sequence rate is a cover-level quantity. Unlike ordinary topological entropy, sequence entropy need not be computed by a generating partition: replacing one-coordinate observations by fixed-length windows can change the normalized sparse rate. We therefore keep the subscript “coord” in the realization theorem and do not silently identify this rate with the supremum over all open covers.

Kawan’s Chapter 2 supplies the other interface. Definition 2.2 defines invariance entropy from finite-horizon spanning controls. Definitions 2.8 and 2.9 introduce invariant covers and their admissible word entropies. Proposition 2.21 proves word-count submultiplicativity; Proposition 2.22 bounds invariance entropy by invariant-cover entropy. Lemma 2.3 disjointizes a cover without increasing its name entropy. It does *not* turn arbitrary nonmeasurable sets into Borel sets. Borel measurability in Theorem 2.3 comes from the safe sets associated with finitely many spanning controls and their successive differences. The theorem takes an infimum over sampling periods as well as partitions. We provide the discrete-time safe-set argument in Section 8, where its application to pattern rates is also made explicit.

3. CONTROL SYSTEMS AND INVARIANT NAME LANGUAGES

Kawan’s invariance entropy and consecutive-name partition entropy are measured in nats per physical time unit; the sparse-time normalizations are specified separately below. Let X be a metrizable space, let U be a nonempty control set, and suppose that $f_u := f(\cdot, u) : X \rightarrow X$ is continuous for every $u \in U$. For a finite control word $\omega = (\omega_0, \dots, \omega_{N-1}) \in U^N$, set

$$\varphi(0, x, \omega) = x, \quad \varphi(n+1, x, \omega) = f(\varphi(n, x, \omega), \omega_n) \quad (0 \leq n < N).$$

A nonempty compact set $Q \subset X$ is *controlled invariant* if every $x \in Q$ admits an infinite control whose complete forward trajectory stays in Q .

For $N \in \mathbb{N}$, let $r_{\text{inv}}(N, Q)$ be the least cardinality of a set $S \subset U^N$ such that for every $x \in Q$ some $\omega \in S$ satisfies

$$\varphi(n, x, \omega) \in Q \quad (0 \leq n \leq N).$$

If no finite such set exists, put $r_{\text{inv}}(N, Q) = +\infty$. The discrete-time invariance entropy is

$$h_{\text{inv}}(Q) = \limsup_{N \rightarrow \infty} \frac{1}{N} \log r_{\text{inv}}(N, Q). \quad (3.1)$$

Throughout, the admissible control space is the full sequence space $\mathcal{U} = U^{\mathbb{N}_0}$; equivalently one may use Kawan’s two-sided full control space and retain its nonnegative coordinates. Every finite word therefore extends to an admissible infinite control, and finite-horizon dependence identifies the preceding spanning number with Kawan’s. Thus (3.1) is Kawan’s definition with the logarithm converted from base two to base e ; see [7, Definition 2.2, p. 44].

Fix a sampling period $\tau \in \mathbb{N}$. A *finite invariant cover* of Q with period τ is a triple $\mathcal{C} = (\mathcal{A}, \tau, \nu)$, where $\mathcal{A}$ is a finite cover of Q and $\nu(D) \in U^\tau$ is assigned to each $D \in \mathcal{A}$, such that

$$\varphi(s, x, \nu(D)) \in Q \quad (x \in D, 0 \leq s \leq \tau). \quad (3.2)$$

This is the discrete-time specialization of Kawan’s Definition 2.8 [7, p. 79]. An *invariant partition* is an invariant cover whose underlying family is a finite Borel partition of Q into

nonempty atoms. It is *clopen* if its atoms are clopen in Q . We write

$$\begin{aligned}\mathfrak{C}_{\text{Bor}}(Q, \tau) &= \{\text{finite Borel invariant partitions of } Q \text{ of period } \tau\}, \\ \mathfrak{C}_{\text{clop}}(Q, \tau) &= \{\mathcal{C} \in \mathfrak{C}_{\text{Bor}}(Q, \tau) : \mathcal{C} \text{ is clopen}\}, \\ \mathfrak{C}_{\text{Bor}}(Q) &= \bigcup_{\tau \in \mathbb{N}} \mathfrak{C}_{\text{Bor}}(Q, \tau).\end{aligned}$$

We use the extended-value convention $\inf \emptyset = +\infty$.

Let $\mathcal{C} = (\{D_i : i \in I_{\mathcal{C}}\}, \tau, \nu)$ be an invariant partition, with $I_{\mathcal{C}}$ a finite nonempty alphabet. For $a = (a_0, \dots, a_{N-1}) \in I_{\mathcal{C}}^N$, concatenate the assigned blocks in that order. The word a is *admissible* if some $x \in D_{a_0}$ follows this concatenated control and satisfies

$$\varphi(j\tau, x, \nu(D_{a_0}) \cdots \nu(D_{a_{N-1}})) \in D_{a_j} \quad (0 \leq j < N). \quad (3.3)$$

The set of such words is $W_N(\mathcal{C}; Q)$. These are the partition words of Kawan's Definition 2.9 [7, pp. 79–80].

Lemma 3.1 (Safety, suffixes, and right extension). *Let $\mathcal{C}$ be an invariant partition.*

- (i) *A witness for $a \in W_N(\mathcal{C}; Q)$ remains in Q at every physical time from 0 through $N\tau$ under the concatenated blocks.*
- (ii) *Every prefix and every suffix of an admissible word is admissible.*
- (iii) *Every $a \in W_N(\mathcal{C}; Q)$ has a right extension in $W_{N+1}(\mathcal{C}; Q)$. In particular, every finite admissible word has an infinite right extension.*
- (iv) *For $M, N \in \mathbb{N}$,*

$$\#W_{M+N}(\mathcal{C}; Q) \leq \#W_M(\mathcal{C}; Q) \#W_N(\mathcal{C}; Q). \quad (3.4)$$

Proof. On block j , condition (3.2), applied to the state in D_{a_j} , keeps the trajectory in Q throughout the next τ steps. Induction proves (i). Prefixes are immediate. A suffix beginning at block j is witnessed by the state at time $j\tau$, which proves (ii).

At time $N\tau$ the terminal state belongs to a unique atom D_b . Appending b and its assigned block gives an admissible word of length $N + 1$, proving (iii) without compactness. Finally, an admissible word of length $M + N$ is determined injectively by its length- M prefix and length- N suffix. Part (ii) places these in W_M and W_N , respectively, and gives (3.4). $\square$

Define the finite-prefix name system

$$\Sigma_{\mathcal{C}} = \{a \in I_{\mathcal{C}}^{\mathbb{N}_0} : a|_{[0, N-1]} \in W_N(\mathcal{C}; Q) \text{ for every } N \in \mathbb{N}\}. \quad (3.5)$$

We give $I_{\mathcal{C}}$ the discrete topology and $\Sigma_{\mathcal{C}}$ the subspace topology from $I_{\mathcal{C}}^{\mathbb{N}_0}$. Its complement is a union of cylinders determined by forbidden finite prefixes, so $\Sigma_{\mathcal{C}}$ is closed. Lemma 3.1(iii) shows that every word in W_N is the initial word of some element of $\Sigma_{\mathcal{C}}$. Lemma 3.1(ii) shows $\sigma(\Sigma_{\mathcal{C}}) \subset \Sigma_{\mathcal{C}}$; surjectivity need not hold. For Borel atoms, an element of $\Sigma_{\mathcal{C}}$ need not have one common state witness for all its prefixes. Our definitions use only finite projections and do not assume such a witness.

There is a useful concrete description of $\Sigma_{\mathcal{C}}$. The feedback block map and its consecutive atom names also appear in the numerical framework of [12, Section 2]. Define

$$T_{\mathcal{C}}(x) = \varphi(\tau, x, \nu(D_i)) \quad \text{if } x \in D_i,$$

and its atom itinerary $\iota_{\mathcal{C}}(x)$ by $(\iota_{\mathcal{C}}(x))_j = i$ if $T_{\mathcal{C}}^j(x) \in D_i$. Each branch of $T_{\mathcal{C}}$ is continuous, and $T_{\mathcal{C}}$ is Borel measurable; it need not be continuous when the atoms are merely Borel. Every finite admissible word is the prefix of the actual itinerary of its witness. Conversely every actual itinerary has admissible prefixes. It follows that

$$\Sigma_{\mathcal{C}} = \overline{\iota_{\mathcal{C}}(Q)}. \quad (3.6)$$

Indeed, each prefix cylinder meeting $\Sigma_{\mathcal{C}}$ meets $\iota_{\mathcal{C}}(Q)$, which proves density, and all actual itineraries belong to the closed set $\Sigma_{\mathcal{C}}$.

Remark 3.2 (Infinite witnesses). *If the atoms are clopen, T_C and ι_C are continuous. Compactness of Q makes $\iota_C(Q)$ closed, so every element of Σ_C has a common state witness. Equivalently, the sets of witnesses for its finite prefixes are nested nonempty compact sets.*

This conclusion can fail for a Borel partition even in a compact continuous system with one control. Take $Q = X = [0, 1]$, $f(x) = \min\{2x, 1\}$, $D_0 = (0, 1/2)$ and $D_1 = Q \setminus D_0$. Both atoms receive the unique control, so the partition is invariant. For every N , the point $x_N = 2^{-N-1}$ witnesses the word 0^N : for $0 \leq j < N$, $f^j(x_N) = 2^{j-N-1} \in (0, 1/2)$. Thus $0^\infty \in \Sigma_C$. No point witnesses this infinite name: zero is outside D_0 , and every positive point eventually leaves D_0 .

For a nonempty finite $F \subset \mathbb{N}_0$, put

$$P_C(F) = \#\{a|_F : a \in \Sigma_C\}. \quad (3.7)$$

Equivalently, if $m = \max F$, this is the number of projections of $W_{m+1}(C; Q)$ onto F . Thus a counted pattern always has a finite witness that remains in Q also between the selected block times.

Let

$$\mathcal{S} = \{A = (a_n)_{n \geq 1} : 0 \leq a_1 < a_2 < \dots, a_n \in \mathbb{N}_0\}.$$

Define

$$p_C^*(n) = \max_{\substack{F \subset \mathbb{N}_0 \\ \#F=n}} P_C(F), \quad (3.8)$$

$$h_{\text{pat}}^*(C; Q) = \lim_{n \rightarrow \infty} \frac{1}{n\tau} \log p_C^*(n), \quad (3.9)$$

$$h_{\text{seq}}^A(C; Q) = \limsup_{n \rightarrow \infty} \frac{1}{n\tau} \log P_C(\{a_1, \dots, a_n\}). \quad (3.10)$$

The existence of the limit in (3.9) follows from Lemma 4.1 below. Although there are infinitely many n -element subsets of $\mathbb{N}_0$, the maximum in (3.8) exists: its possible values form a nonempty subset of $\{1, \dots, (\#I_C)^n\}$.

The entries of A are *block indices*; the corresponding physical observation times are $\tau A = (\tau a_n)_{n \geq 1}$. The normalization in (3.10) is $n\tau$, not the elapsed horizon $a_n\tau$. For fixed τ , $A \mapsto \tau A$ is a bijection between block-index schedules and physical schedules contained in $\tau\mathbb{N}_0$. If sampling periods vary, the same block-index sequence represents different physical schedules.

The ordinary consecutive-name entropy is

$$h(C; Q) = \lim_{N \rightarrow \infty} \frac{1}{N\tau} \log \#W_N(C; Q), \quad (3.11)$$

whose limit exists by (3.4) and Fekete's lemma.

4. SIMULTANEOUS ATTAINMENT FOR PATTERN SYSTEMS

A *finite-alphabet pattern system* is a nonempty set $\Omega \subset E^{\mathbb{N}_0}$ with E finite and nonempty. No topology or shift invariance is imposed. For finite $F \subset \mathbb{N}_0$, write

$$P_\Omega(F) = \#\{x|_F : x \in \Omega\}, \quad p_\Omega^*(n) = \max_{\substack{F \subset \mathbb{N}_0 \\ \#F=n}} P_\Omega(F).$$

With a fixed scale $\tau > 0$, define

$$h_{\text{pat}}^*(\Omega; \tau) = \lim_{n \rightarrow \infty} \frac{1}{n\tau} \log p_\Omega^*(n),$$

$$h_{\text{seq}}^A(\Omega; \tau) = \limsup_{n \rightarrow \infty} \frac{1}{n\tau} \log P_\Omega(\{a_1, \dots, a_n\}).$$

The maximum exists: its possible values form a nonempty subset of the finite integer set $\{1, \dots, (\#E)^n\}$.

We use $P_\Omega(\emptyset) = 1$ when needed. Taking the product-topology closure of Ω changes none of its finite projections: a cylinder meeting $\overline{\Omega}$ already meets Ω . Thus dispensing with closedness is only a matter of formulation. The absence of shift invariance, in contrast, matters to the placement of the coordinate sets.

Lemma 4.1 (Pattern submultiplicativity and deletion). *Let $\Omega \subset E^{\mathbb{N}_0}$ be a pattern system and $q = \#E$. For $m, n \in \mathbb{N}$ and finite $B \subset F \subset \mathbb{N}_0$,*

$$p_\Omega^*(m+n) \leq p_\Omega^*(m)p_\Omega^*(n), \quad (4.1)$$

$$P_\Omega(F) \leq q^{\#(F \setminus B)} P_\Omega(B). \quad (4.2)$$

If $B \subset G$, then $P_\Omega(B) \leq P_\Omega(G)$. Consequently,

$$h_{\text{pat}}^*(\Omega; \tau) = \inf_{n \geq 1} \frac{1}{n\tau} \log p_\Omega^*(n), \quad \log p_\Omega^*(n) \geq n\tau h_{\text{pat}}^*(\Omega; \tau). \quad (4.3)$$

Proof. Split any $(m+n)$ -element set into disjoint subsets of sizes m and n . Restriction sends a pattern on the union injectively to the pair of its two restrictions, proving (4.1). Restriction from F to B has fibers of cardinality at most $q^{\#(F \setminus B)}$, which proves (4.2). If $B \subset G$, every pattern on B extends to a pattern on G using the same element of Ω , giving monotonicity. Fekete's lemma applied to $\log p_\Omega^*(n)$ proves (4.3). $\square$

For comparison, let $\Omega = \{0^\infty, 10^\infty\}$. This set is a closed one-sided subshift, but $P_\Omega(\{0\}) = 2$ and $P_\Omega(\{m\}) = 1$ for $m \geq 1$. One cannot simply translate a maximizing set. Deleting a finite initial interval has a different advantage: the number of lost coordinates is bounded independently of the size of the maximizing set. Inequality (4.2) turns this bound into a controlled information loss.

Theorem 4.2 (Simultaneous attainment). *Fix $\tau > 0$. For every countable family $(\Omega_j)_{j \geq 1}$ of finite-alphabet pattern systems, there is one $A \in \mathcal{S}$ such that*

$$h_{\text{seq}}^A(\Omega_j; \tau) = h_{\text{pat}}^*(\Omega_j; \tau) \quad (j \geq 1). \quad (4.4)$$

Thus the maximal-pattern entropy rate of each system is attained along the same sequence. No closedness, compactness, shift invariance, shift surjectivity, or backward extension is required.

Proof. Let E_j be the alphabet of Ω_j , put $q_j = \#E_j$ and $h_j^* = h_{\text{pat}}^*(\Omega_j; \tau)$. For every A and n ,

$$P_{\Omega_j}(\{a_1, \dots, a_n\}) \leq p_{\Omega_j}^*(n),$$

so (4.3) gives $h_{\text{seq}}^A(\Omega_j; \tau) \leq h_j^*$.

Choose an enumeration of stages

$$(j_k)_{k \geq 1} = (1; 1, 2; 1, 2, 3; 1, 2, 3, 4; \dots),$$

where semicolons only separate finite groups. Every positive integer occurs infinitely often. Choose $\eta_k > 0$ with $\eta_k \rightarrow 0$. We recursively construct nonempty finite sets B_k , each lying strictly to the right of all previous ones. Put

$$G_{k-1} = \bigcup_{i < k} B_i, \quad r_{k-1} = \#G_{k-1}, \quad M_{k-1} = \max G_{k-1},$$

with $r_0 = 0$ and $M_0 = -1$.

If $h_{j_k}^* > 0$, choose n_k so large that

$$n_k > M_{k-1} + 1, \quad \frac{(M_{k-1} + 1) \log q_{j_k}}{n_k \tau} < \eta_k, \quad \frac{r_{k-1}}{n_k} < \eta_k. \quad (4.5)$$

The quantities on the numerators are fixed before n_k is chosen, so these requirements can be satisfied even if the previously selected coordinates are extremely far apart. Notice that $M_{k-1} + 1$ bounds the number of *possible* old positions, whereas r_{k-1} counts the positions actually selected. The former controls deletion; the latter controls dilution by historical

observations. No uniform bound on q_j or on the coordinate gaps is required. Choose an n_k -element F_k attaining $p_{\Omega_{j_k}}^*(n_k)$ and set

$$B_k = F_k \cap \{M_{k-1} + 1, M_{k-1} + 2, \dots\}.$$

At most $M_{k-1} + 1$ coordinates are deleted, and the first condition in (4.5) makes B_k nonempty. If $h_{j_k}^* = 0$, set $B_k = \{M_{k-1} + 1\}$.

Let A be the increasing enumeration of $\bigcup_{k \geq 1} B_k$. Fix j with $h_j^* > 0$ and restrict to the infinitely many stages with $j_k = j$. Write $b_k = \#B_k$ and $R_k = r_{k-1} + b_k = \#G_k$; then G_k consists of the first R_k entries of A . By deletion, (4.3), and monotonicity,

$$\begin{aligned} \log P_{\Omega_j}(G_k) &\geq \log P_{\Omega_j}(B_k) \\ &\geq \log p_{\Omega_j}^*(n_k) - (M_{k-1} + 1) \log q_j \\ &\geq n_k \tau h_j^* - (M_{k-1} + 1) \log q_j. \end{aligned} \quad (4.6)$$

Also $R_k \leq r_{k-1} + n_k$. At all sufficiently late such stages the last line of (4.6) is positive. Dividing by $R_k \tau$ and using (4.5) yields the explicit endpoint estimate

$$\frac{1}{R_k \tau} \log P_{\Omega_j}(G_k) > \frac{h_j^* - \eta_k}{1 + \eta_k}. \quad (4.7)$$

For a fixed j , the indices k with $j_k = j$ tend to infinity, hence $\eta_k \rightarrow 0$ along those stages. Since every B_i is nonempty, $R_k \geq k \rightarrow \infty$. The limsup along these endpoints is at least h_j^* , proving the lower bound. If $h_j^* = 0$, the universal upper bound and nonnegativity give equality for every A . This proves (4.4) for all j . $\square$

Remark 4.3 (What is attained). *The theorem asserts equality of limsup rates. It does not assert a common subsequence of endpoints on which every system is simultaneously close to its maximal rate. The good endpoints used for different j can be different. Nor does it assert convergence of the normalized counts. For example, take $\tau = 1$, let $B \subset \mathbb{N}_0$ and its complement both be infinite, and let Ω_B consist of all binary sequences supported on B . For any schedule, the two normalized logarithmic counts for Ω_B and $\Omega_{\mathbb{N}_0 \setminus B}$ add to $\log 2$ at every n , although both maximal-pattern rates are $\log 2$. They cannot both converge to $\log 2$. The two limsups, however, can both equal $\log 2$.*

Exact finite-level maximization is also a stronger and generally false requirement. Put

$$\Omega = \{0x : x \in \{0, 1\}^{\mathbb{N}_0}\} \cup \{10^\infty, 20^\infty\} \subset \{0, 1, 2\}^{\mathbb{N}_0}.$$

The unique maximizing singleton is $\{0\}$, with three patterns. For a three-element set containing zero there are six patterns, whereas any three positive coordinates realize eight patterns. No increasing sequence maximizes the counts at both levels $n = 1$ and $n = 3$.

Corollary 4.4 (Fixed-language attainment). *For every finite-alphabet pattern system Ω and every $\tau > 0$,*

$$h_{\text{pat}}^*(\Omega; \tau) = \max_{A \in \mathcal{S}} h_{\text{seq}}^A(\Omega; \tau). \quad (4.8)$$

The same identity holds for every finite Borel invariant partition $\mathcal{C}$ after putting $\Omega = \Sigma_{\mathcal{C}}$.

5. A LANGUAGE-QUOTIENT MINIMAX THEOREM

Fix a compact controlled-invariant set Q and one period $\tau \in \mathbb{N}$. For two invariant partitions $\mathcal{C}$ and $\mathcal{D}$ of this period, write $\mathcal{C} \sim_{\text{lang}} \mathcal{D}$ if there is a bijection $\pi : I_{\mathcal{C}} \rightarrow I_{\mathcal{D}}$ such that, for every $N \in \mathbb{N}$,

$$\pi^N(W_N(\mathcal{C}; Q)) = W_N(\mathcal{D}; Q), \quad (5.1)$$

where π^N acts coordinatewise. This is *language equivalence*. Identity, inverse, and composition of the alphabet bijections show that $\sim_{\text{lang}}$ is an equivalence relation. If $F \subset \mathbb{N}_0$ is finite and $m = \max F$, then (5.1) at length $m+1$, followed by projection onto F , gives a bijection between the two F -pattern sets. Thus language-equivalent partitions have identical finite-coordinate pattern counts and identical sequence and maximal-pattern complexities.

For a nonempty class $\mathfrak{C} \subset \mathfrak{C}_{\text{Bor}}(Q, \tau)$, set

$$H_{\text{pat}}^*(Q; \mathfrak{C}) = \inf_{\mathcal{C} \in \mathfrak{C}} h_{\text{pat}}^*(\mathcal{C}; Q), \quad (5.2)$$

$$H_{\text{seq}}^A(Q; \mathfrak{C}) = \inf_{\mathcal{C} \in \mathfrak{C}} h_{\text{seq}}^A(\mathcal{C}; Q). \quad (5.3)$$

Every individual rate is nonnegative and finite, and because the class is nonempty both infima are finite; no uniform bound on the numbers of atoms is needed.

The elementary minimax inequality gives only

$$\sup_A \inf_{\mathcal{C} \in \mathfrak{C}} h_{\text{seq}}^A(\mathcal{C}; Q) \leq \inf_{\mathcal{C} \in \mathfrak{C}} \sup_A h_{\text{seq}}^A(\mathcal{C}; Q) = H_{\text{pat}}^*(Q; \mathfrak{C}).$$

Separate maximizing schedules for separate partitions do not give the opposite inequality. For instance, a two-by-two array with zero diagonal and unit off-diagonal has maximum row minimum zero and minimum column maximum one. The next theorem avoids this logical obstruction by attaining every individual rate at one schedule.

Theorem 5.1 (Minimax over countably many language types). *Let $\emptyset \neq \mathfrak{C} \subset \mathfrak{C}_{\text{Bor}}(Q, \tau)$ and suppose that the quotient $\mathfrak{C}/\sim_{\text{lang}}$ is finite or countable. There is one $A \in \mathcal{S}$ such that*

$$h_{\text{seq}}^A(\mathcal{C}; Q) = h_{\text{pat}}^*(\mathcal{C}; Q) \quad (\mathcal{C} \in \mathfrak{C}). \quad (5.4)$$

Consequently,

$$\max_{B \in \mathcal{S}} \inf_{\mathcal{C} \in \mathfrak{C}} h_{\text{seq}}^B(\mathcal{C}; Q) = \inf_{\mathcal{C} \in \mathfrak{C}} h_{\text{pat}}^*(\mathcal{C}; Q). \quad (5.5)$$

The outer maximum is attained. This proof does not require a partition minimizer.

Proof. Choose one representative $\mathcal{C}_j$ of each language-equivalence class. Enumerate the representatives, repeating them periodically when the quotient is finite, and apply Theorem 4.2 to the resulting countable family $\Sigma_{\mathcal{C}_j}$, using the common scale τ . The resulting A gives equality for each representative. Language equivalence preserves all finite pattern counts, so it gives (5.4) for every $\mathcal{C} \in \mathfrak{C}$. Taking the infimum over $\mathcal{C}$ gives equality at this A .

For any $B \in \mathcal{S}$, Corollary 4.4 gives $h_{\text{seq}}^B(\mathcal{C}; Q) \leq h_{\text{pat}}^*(\mathcal{C}; Q)$ for every $\mathcal{C}$. Taking first the infimum over $\mathcal{C}$ and then the supremum over B proves the reverse minimax inequality and hence (5.5). If the pattern infimum is zero, the pointwise upper bound and nonnegativity in fact give $\inf_{\mathcal{C} \in \mathfrak{C}} h_{\text{seq}}^B(\mathcal{C}; Q) = 0$ for every $B \in \mathcal{S}$, without any countability assumption. $\square$

The hypothesis concerns the symbolic behavior generated on Q , not the names of controls used to present a strategy. It identifies alphabet renumberings and strategies with the same admissible language after such a renumbering. It does not merge distinct atoms merely because their assigned controls agree. It is therefore closer to an intrinsic statement about the chosen partition class, although it remains relative to that class and to the fixed period.

Proposition 5.2 (A verifiable clopen sufficient condition). *Assume that Q is compact metrizable and that the control alphabet U is countable. Then $\mathfrak{C}_{\text{clon}}(Q, \tau)$ is countable. Hence, if this class is nonempty, Theorem 5.1 applies to it. In particular this holds when U is a finite digital alphabet.*

Proof. A compact metrizable space has a countable base. Every clopen subset is compact and hence is a finite union of basic open sets contained in it. Thus Q has only countably many clopen subsets and only countably many finite clopen partitions. Since τ is finite, U^τ is countable. A finite partition admits only countably many block assignments. The invariant assignments form a subcollection. $\square$

Theorem 5.1 fixes τ . It does not produce one common physical-time schedule for partitions of different periods and proves no minimax identity over $\mathfrak{C}_{\text{Bor}}(Q)$.

Example 5.3 (A purely combinatorial uncountable obstruction). *For every infinite $B \subset \mathbb{N}_0$, let*

$$\Omega_B = \{x \in \{0, 1\}^{\mathbb{N}_0} : \text{supp}(x) \subset B\}.$$

At scale $\tau = 1$,

$$P_{\Omega_B}(F) = 2^{\#(F \cap B)}, \quad h_{\text{pat}}^*(\Omega_B; 1) = \log 2.$$

For $A = (a_i) \in \mathcal{S}$,

$$h_{\text{seq}}^A(\Omega_B; 1) = \log 2 \cdot \limsup_{n \rightarrow \infty} \frac{\#(B \cap \{a_1, \dots, a_n\})}{n}.$$

If $\mathbb{N}_0 \setminus \{a_i : i \geq 1\}$ is infinite, choose an infinite B in that complement. Otherwise choose $B = \{a_{k^2} : k \geq 1\}$. In either case the displayed upper density is zero. Therefore

$$\sup_{A \in \mathcal{S}} \inf_{\substack{B \subset \mathbb{N}_0 \\ B \text{ infinite}}} h_{\text{seq}}^A(\Omega_B; 1) = 0 < \log 2 = \inf_{\substack{B \subset \mathbb{N}_0 \\ B \text{ infinite}}} h_{\text{pat}}^*(\Omega_B; 1). \quad (5.6)$$

This proves only that minimax exchange is not a formal consequence of pointwise attainment for arbitrary uncountable pattern families. Every invariant-partition name system satisfies $\sigma(\Sigma_C) \subset \Sigma_C$ by suffix closure. In contrast, $\sigma(\Omega_B) \subset \Omega_B$ forces $k-1 \in B$ whenever $k \in B \setminus \{0\}$; an infinite such B must equal $\mathbb{N}_0$. Hence the proper sparse sets producing (5.6) cannot occur as these autonomous control name systems. The equation is not a deterministic control counterexample and yields no failure statement for the unrestricted Borel invariant-partition class.

6. REALIZING ONE-SIDED SUBSHIFTS AS SAFETY SYSTEMS

Let E be a finite nonempty discrete alphabet and let $\emptyset \neq Y \subset E^{\mathbb{N}_0}$ be closed with $\sigma(Y) \subset Y$. Write

$$\mathcal{L}_N(Y) = \{y_0 \cdots y_{N-1} : y \in Y\}, \quad P_Y(F) = \#\{y|_F : y \in Y\}.$$

The coordinate complexities are

$$h_{\text{seq, coord}}^A(Y) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_Y(\{a_1, \dots, a_n\}),$$

$$h_{\text{pat, coord}}^*(Y) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \max_{\substack{F \subset \mathbb{N}_0 \\ \#F=n}} P_Y(F).$$

Adjoin an isolated cemetery point and give $X = Y \sqcup \{\dagger\}$ the topological disjoint-union topology. Put $U = E$ and define

$$f(y, u) = \begin{cases} \sigma y, & y_0 = u, \\ \dagger, & y_0 \neq u, \end{cases} \quad f(\dagger, u) = \dagger. \quad (6.1)$$

Theorem 6.1 (Symbolic-to-control realization). *For the matching-control system (6.1):*

- (i) *X is compact metrizable, $f : X \times U \rightarrow X$ is continuous, and Y is a compact controlled-invariant set.*
- (ii) *For every $N \in \mathbb{N}$,*

$$r_{\text{inv}}(N, Y) = \#\mathcal{L}_N(Y), \quad (6.2)$$

and therefore

$$h_{\text{inv}}(Y) = h_{\text{top}}(Y, \sigma) = \lim_{N \rightarrow \infty} \frac{1}{N} \log \#\mathcal{L}_N(Y). \quad (6.3)$$

- (iii) At period one identify U^1 with U . If $\mathcal{C} = (\{D_i : i \in I_{\mathcal{C}}\}, 1, \nu) \in \mathfrak{C}_{\text{Bor}}(Y, 1)$, set $\lambda_{\mathcal{C}}(i) = \nu(D_i) \in E$. Then every atom satisfies

$$D_i \subset [\lambda_{\mathcal{C}}(i)]_Y := \{y \in Y : y_0 = \lambda_{\mathcal{C}}(i)\}. \quad (6.4)$$

Moreover, the one-block label map

$$\Lambda_{\mathcal{C}} : \Sigma_{\mathcal{C}} \longrightarrow Y, \quad (a_j)_{j \geq 0} \longmapsto (\lambda_{\mathcal{C}}(a_j))_{j \geq 0}, \quad (6.5)$$

is a continuous, onto, shift-commuting map.

- (iv) For every $A \in \mathcal{S}$,

$$\inf_{\mathcal{C} \in \mathfrak{C}_{\text{Bor}}(Y, 1)} h_{\text{seq}}^A(\mathcal{C}; Y) = \inf_{\mathcal{C} \in \mathfrak{C}_{\text{clon}}(Y, 1)} h_{\text{seq}}^A(\mathcal{C}; Y) = h_{\text{seq, coord}}^A(Y), \quad (6.6)$$

$$\inf_{\mathcal{C} \in \mathfrak{C}_{\text{Bor}}(Y, 1)} h_{\text{pat}}^*(\mathcal{C}; Y) = \inf_{\mathcal{C} \in \mathfrak{C}_{\text{clon}}(Y, 1)} h_{\text{pat}}^*(\mathcal{C}; Y) = h_{\text{pat, coord}}^*(Y), \quad (6.7)$$

$$\inf_{\mathcal{C} \in \mathfrak{C}_{\text{Bor}}(Y, 1)} h(\mathcal{C}; Y) = \inf_{\mathcal{C} \in \mathfrak{C}_{\text{clon}}(Y, 1)} h(\mathcal{C}; Y) = h_{\text{top}}(Y, \sigma). \quad (6.8)$$

Every Borel and clopen infimum in (6.6)–(6.8) is attained by the same first-coordinate cylinder partition.

Proof. The space X is the union of the two clopen compact sets Y and $\{\dagger\}$, hence is compact metrizable. For fixed $u \in E$, the sets $[u]_Y$, $Y \setminus [u]_Y$, and $\{\dagger\}$ are clopen; on them f_u is, respectively, the shift or a constant. Thus f_u is continuous. Since U is finite discrete, f is jointly continuous. Choosing $u = y_0$ sends y to $\sigma y \in Y$, proving controlled invariance.

A direct induction gives

$$\varphi(n, y, \omega) \in Y \quad (0 \leq n \leq N) \iff \omega_j = y_j \quad (0 \leq j < N). \quad (6.9)$$

Indeed, after j matches the state is $\sigma^j y$, whereas a mismatch enters $\dagger$ permanently. Hence a length- N control word is safe from at least one state of Y exactly when it belongs to $\mathcal{L}_N(Y)$, and then it safely covers precisely its nonempty prefix cylinder. Distinct language words give disjoint cylinders. More explicitly:

Lower bound. For each $w \in \mathcal{L}_N(Y)$ choose $y^w \in Y$ with prefix w . If $S \subset U^N$ spans Y , some member of S is safe from y^w . By (6.9) that member must equal w . Thus $\mathcal{L}_N(Y) \subset S$, and $r_{\text{inv}}(N, Y) \geq \#\mathcal{L}_N(Y)$.

Upper bound. Take $S = \mathcal{L}_N(Y)$. For any $y \in Y$, its prefix $y_0 \cdots y_{N-1}$ belongs to S and is safe through time N by (6.9). Hence $r_{\text{inv}}(N, Y) \leq \#\mathcal{L}_N(Y)$. This proves (6.2). The language counts are submultiplicative, so their normalized logarithms converge.

For completeness, this limit is the usual topological entropy even when $\sigma|_Y$ is not onto. The join of the first-coordinate cylinder cover through times $0, \dots, N-1$ has exactly $\#\mathcal{L}_N(Y)$ nonempty atoms. Conversely, given a finite open cover of compact Y , the partition into cylinders of some fixed length m refines that cover; its N -fold dynamical join is then refined by the length- $(N+m-1)$ cylinders. Taking exponential rates proves (6.3).

Let $\mathcal{C} = (\{D_i : i \in I_{\mathcal{C}}\}, 1, \nu)$ be a Borel invariant partition. If D_i contained y with $y_0 \neq \lambda_{\mathcal{C}}(i)$, its assigned control would send y to $\dagger \notin Y$, contradicting safety. This proves (6.4).

Take $a = (a_j)_{j \geq 0} \in \Sigma_{\mathcal{C}}$ and put $z_j = \lambda_{\mathcal{C}}(a_j)$. For each N , choose a witness $y^{(N)}$ for the first N symbols of a . Induction using matching controls and (6.4) gives $y_j^{(N)} = \lambda_{\mathcal{C}}(a_j) = z_j$ for $0 \leq j < N$. Hence $y^{(N)} \rightarrow z$ in the product topology; since Y is closed, $z \in Y$. Thus $\Lambda_{\mathcal{C}}$ is well defined, and it is continuous: the preimage of a cylinder in $E^{\mathbb{N}_0}$ is a finite union of cylinders in $I_{\mathcal{C}}^{\mathbb{N}_0}$, restricted to $\Sigma_{\mathcal{C}}$. It is shift commuting because $(\Lambda_{\mathcal{C}}(\sigma a))_j = \lambda_{\mathcal{C}}(a_{j+1}) = (\sigma \Lambda_{\mathcal{C}}(a))_j$. Given $y \in Y$, let a_j be the unique index with $\sigma^j y \in D_{a_j}$. Its assigned label is y_j , and y witnesses every finite prefix of this atom itinerary. Thus (6.5) is onto. Here each prefix witness $y^{(N)}$ may be different. Its first N control labels nevertheless equal those of z , so convergence to $z \in Y$ follows. This does not prove that z follows the original atom name a ;

its orbit may meet different atoms with the same labels. Thus neither shift surjectivity nor a common atom-name witness is used.

For every finite F , the coordinate projection of $\Lambda_{\mathcal{C}}$ is onto, so

$$P_{\mathcal{C}}(F) \geq P_Y(F). \quad (6.10)$$

It follows that every Borel, and hence every clopen, invariant partition has sequence, maximal-pattern, and consecutive-name entropy at least the corresponding coordinate quantity.

Let $E_Y = \{e \in E : [e]_Y \neq \emptyset\}$ and define

$$\mathcal{C}_0 = (\{D_e = [e]_Y : e \in E_Y\}, 1, \nu_0), \quad \nu_0(D_e) = e.$$

This is a clopen invariant partition. Under the symbol identification, $W_N(\mathcal{C}_0; Y) = \mathcal{L}_N(Y)$ and $\Sigma_{\mathcal{C}_0} = Y$; the latter follows from closedness of Y . Equality holds in (6.10) for every F . Thus $\mathcal{C}_0$ attains all the lower bounds, proving (6.6)–(6.8). $\square$

7. A STRICT THUE–MORSE SEPARATION

Let $\vartheta(0) = 01$, $\vartheta(1) = 10$, and let

$$\mathbf{t} = \lim_{r \rightarrow \infty} \vartheta^r(0) = 0110100110010110 \dots$$

Equivalently, $t_n = s_2(n) \pmod{2}$, where $s_2(n)$ is the number of ones in the binary expansion of n . Let

$$Y = \overline{\{\sigma^m \mathbf{t} : m \geq 0\}} \subset \{0, 1\}^{\mathbb{N}_0}. \quad (7.1)$$

This is the one-sided Thue–Morse orbit closure. The proof below computes both rates used in the control corollary directly; no unverified infinite-word theorem is used as a dependency.

Proposition 7.1 (Exact symbolic rates). *For every $N \geq 1$,*

$$\#\mathcal{L}_N(Y) \leq 8N. \quad (7.2)$$

Hence $h_{\text{top}}(Y) = 0$. On the single schedule

$$A_{\text{TM}} = (a_i)_{i \geq 1}, \quad a_i = 4^{i-1} - 1, \quad (7.3)$$

one has, for every $k \geq 1$,

$$P_Y(\{a_1, \dots, a_k\}) = 2^k. \quad (7.4)$$

The chosen coordinate set contains zero. Since a cylinder is clopen, its patterns in the orbit closure are exactly the patterns obtained along shifts of $\mathbf{t}$; this calculation uses actual shifts of the fixed point. Consequently

$$h_{\text{seq, coord}}^{A_{\text{TM}}}(Y) = h_{\text{pat, coord}}^*(Y) = \log 2. \quad (7.5)$$

Proof. A finite word is in $\mathcal{L}_N(Y)$ exactly when it occurs as a factor of $\mathbf{t}$: one direction follows from the orbit, and the other from convergence in a prefix cylinder. Choose $\ell = \lceil \log_2 N \rceil$. The identity $\mathbf{t} = \vartheta^\ell(\mathbf{t})$ decomposes $\mathbf{t}$ into aligned supertiles of length 2^ℓ . Since $N \leq 2^\ell$, every length- N factor is contained in $\vartheta^\ell(ab)$ for an adjacent two-letter factor ab . There are at most four choices for ab and at most 2^ℓ starting offsets in the first supertile. Therefore $\#\mathcal{L}_N(Y) \leq 4 \cdot 2^\ell \leq 8N$, proving zero topological entropy.

Fix $k \geq 1$ and $\varepsilon = (\varepsilon_0, \dots, \varepsilon_{k-1}) \in \{0, 1\}^k$. Put

$$r = \#\{0 \leq i < k : \varepsilon_i = 0\}, \quad q = \sum_{\substack{0 \leq i < k \\ \varepsilon_i = 0}} 4^i + (r \bmod 2)4^k.$$

Only even binary positions occur in this sum. The binary digit sum of q is $r + (r \bmod 2)$, which is even, so $t_q = 0$. If $\varepsilon_i = 1$, bit $2i$ of q is zero, and adding $4^i = 2^{2i}$ changes the digit-sum parity. If $\varepsilon_i = 0$, bit $2i$ is one and bit $2i + 1$ is zero; adding 2^{2i} replaces these two bits by zero and one without a further carry, so the parity is unchanged. In both cases

$$t_{q+4^i} = \varepsilon_i \quad (0 \leq i < k). \quad (7.6)$$

The extra bit at position $2k$ only corrects the initial parity of q . For $i < k$, a carry stops at the empty odd position $2i + 1$, which is strictly below $2k$. The parity correction therefore interferes with none of these tests. Each addition $q + 4^i$ is performed separately; we are not adding all the 4^i 's successively to one integer.

Set $n = q + 1$. Since $a_{i+1} = 4^i - 1$,

$$(\sigma^n \mathbf{t})_{a_{i+1}} = t_{q+4^i} = \varepsilon_i.$$

Thus every binary k -pattern occurs on the first k members of A_{TM} , proving (7.4). The binary alphabet gives the reverse upper bound $p_Y^*(k) \leq 2^k$, and (7.5) follows. $\square$

Corollary 7.2 (Zero invariance entropy, full binary sparse-time complexity). *Apply the matching-control construction (6.1) to the Thue–Morse subshift (7.1). Then*

$$h_{\text{inv}}(Y) = 0 < \log 2 = \inf_{\mathcal{C} \in \mathfrak{C}_{\text{Bor}}(Y,1)} h_{\text{pat}}^*(\mathcal{C}; Y) = \inf_{\mathcal{C} \in \mathfrak{C}_{\text{clon}}(Y,1)} h_{\text{pat}}^*(\mathcal{C}; Y). \quad (7.7)$$

Moreover, the explicit schedule A_{TM} satisfies

$$\max_{A \in \mathcal{S}} \inf_{\mathcal{C} \in \mathfrak{C}_{\text{Bor}}(Y,1)} h_{\text{seq}}^A(\mathcal{C}; Y) = \inf_{\mathcal{C} \in \mathfrak{C}_{\text{Bor}}(Y,1)} h_{\text{seq}}^{A_{\text{TM}}}(\mathcal{C}; Y) = \log 2, \quad (7.8)$$

$$\max_{A \in \mathcal{S}} \inf_{\mathcal{C} \in \mathfrak{C}_{\text{clon}}(Y,1)} h_{\text{seq}}^A(\mathcal{C}; Y) = \inf_{\mathcal{C} \in \mathfrak{C}_{\text{clon}}(Y,1)} h_{\text{seq}}^{A_{\text{TM}}}(\mathcal{C}; Y) = \log 2. \quad (7.9)$$

Proof. Combine Theorem 6.1 with Proposition 7.1. For any schedule the binary coordinate complexity is at most $\log 2$, while A_{TM} attains $\log 2$. Equations (6.6)–(6.7) give the Borel and clopen infima. $\square$

Corollary 7.2 is a deterministic control-system statement, not merely a symbolic analogy. It also shows that the new sparse-time invariant-language infimum cannot in general be identified with Kawan's invariance entropy.

8. COMPARISON, ALL PERIODS, AND LIMITATIONS

8.1. Comparison with Kawan's invariance entropy. For every invariant partition, consecutive coordinates are one permissible choice in the maximal-pattern count. Therefore

$$h(\mathcal{C}; Q) \leq h_{\text{pat}}^*(\mathcal{C}; Q) \leq \frac{\log \#I_{\mathcal{C}}}{\tau}. \quad (8.1)$$

Kawan's Theorem 2.3, with natural logarithms, gives

$$h_{\text{inv}}(Q) = \inf_{\mathcal{C} \in \mathfrak{C}_{\text{Bor}}(Q)} h(\mathcal{C}; Q). \quad (8.2)$$

Its hypotheses are satisfied by the full-control discrete-time systems used here: X is metrizable, each controlled map is continuous, and Q is nonempty, compact and controlled invariant. The period in this formula is allowed to vary.

For clarity, the relevant argument can be given directly. Spanning families concatenate, so $r_{\text{inv}}(m + n, Q) \leq r_{\text{inv}}(m, Q)r_{\text{inv}}(n, Q)$ when the right-hand side is finite. Moreover $r_{\text{inv}}(n, Q)$ is nondecreasing. If any finite-horizon spanning number is finite, then $r_{\text{inv}}(1, Q)$ is finite, all of them are finite by concatenation, and Fekete's lemma gives a limit in (3.1). If none is finite, that limit equals $+\infty$.

Let $S_N = \{\omega^1, \dots, \omega^m\}$ be a minimal finite N -step spanning family. Its safe sets are

$$K_i = Q \cap \bigcap_{s=1}^N \{x \in X : \varphi(s, x, \omega^i) \in Q\}.$$

They are closed in Q , since finite control compositions are continuous and Q is closed. They cover Q . Set $D_1 = K_1$ and $D_i = K_i \setminus \bigcup_{j < i} K_j$, discard empty sets, and assign ω^i to D_i . This gives a Borel invariant partition $\mathcal{C}_N$ of period N with at most m atoms. This is the measurable construction underlying Kawan's Theorem 2.3; Lemma 2.3 alone only provides disjointization. The safe sets are already closed before that lemma is applied.

For the other inequality in (8.2), all admissible k -block words of any $\mathcal{C}$ form a spanning family after their assigned controls are concatenated. Indeed, from each initial state the feedback rule selects one such admissible word. Consequently $r_{\text{inv}}(k\tau, Q) \leq \#W_k(\mathcal{C}; Q)$, and taking limits gives $h_{\text{inv}}(Q) \leq h(\mathcal{C}; Q)$. The reverse inequality follows from $h(\mathcal{C}_N; Q) \leq N^{-1} \log r_{\text{inv}}(N, Q)$ and $N \rightarrow \infty$. If no finite spanning family exists, no finite invariant partition exists, and both sides are $+\infty$.

Thus, for every nonempty fixed-period subclass $\mathfrak{C} \subset \mathfrak{C}_{\text{Bor}}(Q, \tau)$, the general comparison is

$$h_{\text{inv}}(Q) \leq \inf_{\mathcal{C} \in \mathfrak{C}} h(\mathcal{C}; Q) \leq \inf_{\mathcal{C} \in \mathfrak{C}} h_{\text{pat}}^*(\mathcal{C}; Q). \quad (8.3)$$

The minimax theorem concerns the last quantity. The Thue–Morse realization proves that it can strictly exceed $h_{\text{inv}}(Q)$.

8.2. Why the unrestricted all-period infimum collapses. There is a natural candidate for an all-period Borel quantity,

$$H_{\text{pat,all}}^{*,\text{Bor}}(Q) := \inf_{\tau \in \mathbb{N}} \inf_{\mathcal{C} \in \mathfrak{C}_{\text{Bor}}(Q, \tau)} h_{\text{pat}}^*(\mathcal{C}; Q).$$

The following proposition records the consequence of Kawan’s spanning construction for this all-period quantity.

Proposition 8.1 (All-period collapse). *In the full-control discrete-time setting of Section 3, every nonempty compact controlled-invariant set Q satisfies*

$$H_{\text{pat,all}}^{*,\text{Bor}}(Q) = h_{\text{inv}}(Q). \quad (8.4)$$

Proof. The lower bound follows from (8.3). For the upper bound use the same $\mathcal{C}_N$ just constructed. Since its alphabet has at most $r_{\text{inv}}(N, Q)$ symbols, every pattern on k coordinates has at most $r_{\text{inv}}(N, Q)^k$ possibilities. Hence

$$H_{\text{pat,all}}^{*,\text{Bor}}(Q) \leq h_{\text{pat}}^*(\mathcal{C}_N; Q) \leq \frac{\log r_{\text{inv}}(N, Q)}{N}.$$

Let $N \rightarrow \infty$. The infinite-entropy case was already covered by nonexistence of finite invariant partitions. This proves (8.4). $\square$

For the matching-control Thue–Morse example, the collapse is particularly visible. At period N , the length- N cylinder partition with its matching control blocks is clopen, has at most $8N$ atoms and has pattern rate at most $N^{-1} \log(8N)$. Thus even the all-period clopen infimum is zero in this example, while the period-one Borel infimum is $\log 2$. Changing the sampling period changes which information is packed into one observed symbol and the cost assigned to that symbol.

8.3. Countability, trajectories, and physical time. The countability hypothesis in Theorem 5.1 is used exactly when one representative language from each class is listed and assigned infinitely many construction stages. Compact metrizability gives countably many clopen sets; it does not give countably many Borel sets or Borel name languages. No approximation theorem allowing that substitution is proved here. Example 5.3 rules out a formal uncountable extension for arbitrary pattern sets, but its non-shift-invariant examples do not settle the unrestricted autonomous control problem. Theorem 6.1 bypasses countability in its specific realization by exhibiting a least language under the label projections. The countable-language theorem does not settle the general Borel minimax problem. The later closed-cover theorem resolves a particular uncountable class; the inverse-capacity theorem resolves maximal-pattern optimization, but does not by itself resolve the corresponding common-sampling problem.

The patterns count admissible language projections. Every finite pattern has a finite trajectory witness which stays safe between observations. An infinite point in the completed Borel name system may have only separate finite-prefix witnesses, as Remark 3.2 shows. This distinction cannot be removed by compactness alone when the atoms are not closed. The

factor in Theorem 6.1 is a factor of compact *name systems*; it is not a continuous factor defined by an arbitrary Borel feedback map on the original state space.

For one fixed τ , A denotes block indices and τA denotes physical observation times. The normalization $n\tau$ charges one block length per observation and disregards the lengths of the gaps. A common block-index sequence for different periods would represent different physical schedules. Theorem 4.2 could be applied to countably many languages with different positive normalizing constants by rescaling each equality, but that would not solve the distinct physical-time minimax problem. Formula (8.4) is an infimum identity and makes no common-schedule assertion.

Finally, if one safe control block works for every point of Q , the one-atom invariant partition makes all three fixed-period infima zero for any class containing it. On a connected Q , every finite clopen partition has one atom. The full clopen class is therefore either empty or has zero pattern infimum, while Borel strategies can remain nontrivial. These observations explain both the usefulness and the limits of the clopen sufficient condition. The nontrivial separation studied here records the richness of safe symbolic instructions at a specified sampling period, despite a zero asymptotic count per elapsed time.

9. COUNTING ENTROPY AND THE CLOSED-COVER THEOREM

Let Q be a nonempty compact metric space and let $T : Q \rightarrow Q$ be a Borel map. For a finite Borel partition α of Q and a finite $F \subset \mathbb{N}_0$, put

$$P_\alpha(F) = \left| \left\{ (\alpha(T^k x))_{k \in F} : x \in Q \right\} \right|,$$

where $\alpha(y)$ denotes the atom containing y and empty atoms are discarded. Thus $P_\alpha(F)$ is the number of nonempty atoms in $\bigvee_{k \in F} T^{-k} \alpha$. For a strictly increasing sequence $A = (a_i)_{i \geq 1}$ in $\mathbb{N}_0$, define

$$h_A(T, \alpha) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_\alpha(\{a_1, \dots, a_n\}), \quad (9.1)$$

$$p_\alpha^*(n) = \max_{\substack{F \subset \mathbb{N}_0 \\ |F|=n}} P_\alpha(F), \quad h_{\text{pat}}^*(T, \alpha) = \lim_{n \rightarrow \infty} \frac{1}{n} \log p_\alpha^*(n). \quad (9.2)$$

All logarithms are natural. These are combinatorial counting quantities; for Borel atoms they are not Shannon entropies of an invariant measure. The maximum in (9.2) exists because the counts are integers between 1 and $|\alpha|^n$. Splitting a window into two parts gives $p_\alpha^*(n+m) \leq p_\alpha^*(n)p_\alpha^*(m)$. Subadditivity proves the existence of the limit, and $h_A(T, \alpha) \leq h_{\text{pat}}^*(T, \alpha)$. No invertibility or continuity of the observation is used here.

Let $\mathcal{K} = (K_w)_{w \in W}$ be a finite closed cover of Q , and write

$$r = N(\mathcal{K}) = \min\{|I| : I \subset W, \bigcup_{w \in I} K_w = Q\}.$$

The set $\mathcal{B}(\mathcal{K})$ consists of all finite Borel partitions refining $\mathcal{K}$: each atom must be contained in at least one K_w . Equivalently, one may first consider all Borel maps $s : Q \rightarrow W$ with $x \in K_{s(x)}$, and then allow arbitrary finite Borel refinements of their fibers.

Let H be a primitive finite 0–1 matrix, with vertex set V , and let

$$Y_H = \{y \in V^{\mathbb{N}_0} : H_{y_j y_{j+1}} = 1 \text{ for all } j \geq 0\}.$$

Primitivity means that all sufficiently large powers of H have strictly positive entries. Write σ for the one-sided shift.

Theorem 9.1. *Suppose $T : Q \rightarrow Q$ is topologically conjugate to $\sigma^p : Y_H \rightarrow Y_H$ for some primitive H and integer $p \geq 1$. Let $\mathcal{K}$ be any finite closed cover of Q . For every strictly increasing $A = (a_i)$ satisfying*

$$a_{i+1} - a_i \longrightarrow \infty, \quad (9.3)$$

one has

$$\min_{\alpha \in \mathcal{B}(\mathcal{K})} h_A(T, \alpha) = \min_{\alpha \in \mathcal{B}(\mathcal{K})} h_{\text{pat}}^*(T, \alpha) = \log r. \quad (9.4)$$

An ordered difference partition of any minimum subcover attains both minima, simultaneously for every A satisfying (9.3). Consequently,

$$\max_A \inf_{\alpha \in \mathcal{B}(\mathcal{K})} h_A(T, \alpha) = \log r, \quad (9.5)$$

where the maximum ranges over all strictly increasing infinite sequences and is attained, for example, by $a_i = (i - 1)^2$.

The order of quantifiers is part of the assertion. Every A satisfying (9.3) works for every allowed partition, although the finite initial segment discarded in the proof can depend on that partition. The proof does not interchange an infimum over partitions with a finite-window limit.

9.1. A category and concatenation lemma. We use meagerness relative to Q . A Borel set has the Baire property, so a nonmeager Borel set is comeager in some nonempty open set. Indeed, the sets differing from open sets by meager sets form a sigma-algebra: for complements, the additional difference is contained in the nowhere dense boundary of an open set. A compact metric space is a Baire space. In particular, a nonempty open set cannot be covered by countably many nowhere dense sets, as is also seen by choosing nested nonempty open sets with closures avoiding those sets and diameters tending to zero.

If $F : Q \rightarrow Q$ is continuous and open, preimages of meager sets are meager. For a closed nowhere dense N , the closed set $F^{-1}N$ has empty interior: otherwise the image of a nonempty open subset would be a nonempty open subset of N . Passing to countable unions proves the assertion.

Lemma 9.2. *Under the dynamical assumptions of Theorem 9.1, T is continuous, open, and onto. Given finitely many nonempty cylinder sets $C_1, \dots, C_q$ in symbolic coordinates, there is an integer $b \geq 1$ such that, for every $k \geq 1$, every $t_1, \dots, t_k \in \{1, \dots, q\}$, and all $0 \leq n_1 < \dots < n_k$ with $n_{j+1} - n_j \geq b$,*

$$\bigcap_{j=1}^k T^{-n_j} C_{t_j}$$

is nonempty and open.

Proof. Work on Y_H through the conjugacy. The image under σ of a prefix cylinder of length at least two is the cylinder obtained by removing its first symbol. The image of a one-symbol cylinder is the union of the permitted successor cylinders. Hence σ is open. Every vertex has a predecessor, so every infinite path has a predecessor; thus σ is onto. These properties pass to powers and conjugates.

Choose within each C_j a nonempty prefix cylinder, and extend their defining words, if necessary, to a common length ℓ . Choose N with $H^n > 0$ for every $n \geq N$, and then choose b with $pb - \ell + 1 \geq N$. Place the word for C_{t_j} at coordinate pn_j . Between the end of one placed word and the start of the next there are $p(n_{j+1} - n_j) - \ell + 1$ edges to fill. Primitivity supplies a path of that length between any two endpoint vertices. Predecessors fill the finite segment before the first placed word, and successors provide an infinite continuation after the last. This constructs an actual point in the intersection. The intersection is open because it is a finite intersection of preimages of open sets under continuous maps. $\square$

Lemma 9.3. *Under the same assumptions, let α be a finite Borel partition with q nonmeager atoms. For every A satisfying (9.3) there exists $m \geq 1$ such that*

$$P_\alpha(\{a_1, \dots, a_n\}) \geq q^{n-m+1} \quad (n \geq m). \quad (9.6)$$

In particular $h_A(T, \alpha) \geq \log q$.

Proof. List the nonmeager atoms as $D_1, \dots, D_q$. The Baire property provides a nonempty cylinder C_j for which $C_j \setminus D_j$ is meager. Let b be supplied by Lemma 9.2, and choose m so that $a_{i+1} - a_i \geq b$ for all $i \geq m$.

Fix $n \geq m$ and any target labels $t_m, \dots, t_n$. The set

$$G = \bigcap_{i=m}^n T^{-a_i} C_{t_i}$$

is nonempty and open. Each power T^{a_i} is continuous and open, so

$$\bigcup_{i=m}^n T^{-a_i} (C_{t_i} \setminus D_{t_i})$$

is meager and cannot cover G . A point outside this union realizes all the specified atoms at the sampled times. Distinct target label strings give distinct patterns; restriction from the full window to this tail is onto. This proves (9.6). Dividing its logarithm by n and taking a limit inferior proves the asserted bound. $\square$

9.2. Proof of the exact formula.

Proof of Theorem 9.1. Let $\alpha \in \mathcal{B}(\mathcal{K})$. Assign to each atom D one index $w(D)$ with $D \subset K_{w(D)}$. Let J be the set of indices assigned to the nonmeager atoms. If the closed sets $(K_w)_{w \in J}$ did not cover Q , their complement would be a nonempty open set covered by the finitely many meager atoms of α , which is impossible. Thus $|J| \geq r$, and the number q of nonmeager atoms is at least r . Lemma 9.3 gives

$$h_A(T, \alpha) \geq \log q \geq \log r$$

for every A satisfying (9.3). Closedness of the cover members is used precisely in this passage from category to a full cover.

Choose a minimum subcover $K_{w_1}, \dots, K_{w_r}$, and set

$$E_j = K_{w_j} \setminus \bigcup_{k < j} K_{w_k}, \quad \alpha_0 = \{E_1, \dots, E_r\}. \quad (9.7)$$

Every E_j is Borel and is nonempty, since a minimum subcover has no redundant member. The partition refines $\mathcal{K}$ and has r atoms. Hence $P_{\alpha_0}(F) \leq r^{|F|}$ for all finite F , giving $h_A(T, \alpha_0) \leq h_{\text{pat}}^*(T, \alpha_0) \leq \log r$ for every A . The established lower bound and $h_{\text{pat}}^* \geq h_A$ yield (9.4) and simultaneous attainment. For arbitrary A , the partition α_0 bounds the inner infimum in (9.5) by $\log r$. Any A satisfying (9.3) gives the reverse bound and attains the maximum. $\square$

10. A COVER-PARTITION GAP AT PRESCRIBED SPARSE TIMES

For any finite cover $\mathcal{K}$ define its combinatorial sequence entropy by

$$h_A(T, \mathcal{K}) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log N \left(\bigvee_{i=1}^n T^{-a_i} \mathcal{K} \right).$$

For a partition this is (9.1). For overlapping sets, covering each finite orbit window need not correspond to one fixed refining partition used at every time.

Corollary 10.1. *Let $q \geq 3$, $Q = \{0, \dots, q-1\}^{\mathbb{N}_0}$, $T = \sigma$, and*

$$K_j = \{x : x_0 \neq j\} \quad (0 \leq j < q), \quad \mathcal{K} = (K_j)_{j=0}^{q-1}.$$

For every increasing infinite A ,

$$h_A(\sigma, \mathcal{K}) = \log \frac{q}{q-1}. \quad (10.1)$$

For every such A with diverging successive gaps,

$$\min_{\alpha \in \mathcal{B}(\mathcal{K})} h_A(\sigma, \alpha) = \log 2 > h_A(\sigma, \mathcal{K}). \quad (10.2)$$

Proof. The cover has no single member equal to Q , and any two distinct members cover Q . Thus $r = 2$, and (10.2) follows from Theorem 9.1 once (10.1) is proved.

For a window F of size n , the full shift has all q^n possible input vectors. Each member of the joined cover specifies one excluded letter at each coordinate, and covers $(q-1)^n$ of these vectors. Consequently the minimum number $c(n)$ of such boxes is at least $(q/(q-1))^n$. For completeness, the following elementary probabilistic covering bound gives the matching exponent. Choose R excluded-letter vectors independently and uniformly from the q^n possibilities. A fixed input vector is left uncovered with probability at most

$$\left(1 - \left(\frac{q-1}{q}\right)^n\right)^R \leq \exp\left(-R \left(\frac{q-1}{q}\right)^n\right).$$

Taking $R = \lceil (q/(q-1))^n (n \log q + 1) \rceil$ makes the sum of these probabilities over all input vectors smaller than one. Some choice therefore covers every vector. We have proved

$$\left(\frac{q}{q-1}\right)^n \leq c(n) \leq \left\lceil \left(\frac{q}{q-1}\right)^n (n \log q + 1) \right\rceil.$$

The count depends only on n , not on the positions of the sampled coordinates, so its exponential rate proves (10.1). $\square$

The excluded-letter cover and its covering exponent are already present in Romagnoli [10]; the calculation above is included to identify the sparse-time quantity without an external dependency. The consequence of Theorem 9.1 is the exact minimum over *all* Borel refinements for a sequence fixed in advance. It does not assert that every Borel refinement has consecutive-time entropy at least $\log 2$, or that every sampling sequence gives that lower bound.

11. FIXED-PERIOD SAFE FEEDBACKS WITH A COMMON SUCCESSOR

Let X be a compact metric space, U a nonempty finite control set, and $f_u : X \rightarrow X$ continuous for every $u \in U$. Let $Q \subset X$ be a nonempty compact controlled-invariant set: at each $x \in Q$ at least one control keeps the next state in Q . All infinite control strings are allowed. Fix an integer period $\tau \geq 1$ once and for all, and put $W = U^\tau$. For $w = (u_0, \dots, u_{\tau-1})$ write

$$\varphi(0, x, w) = x, \quad \varphi(j, x, w) = f_{u_{j-1}} \circ \dots \circ f_{u_0}(x),$$

and define

$$K_w = \{x \in Q : \varphi(j, x, w) \in Q \text{ } (0 \leq j \leq \tau)\}, \quad g_w(x) = \varphi(\tau, x, w). \quad (11.1)$$

The K_w are closed and cover Q . The coverage follows by choosing legal single-step controls successively for the fixed finite horizon.

The class $\mathcal{S}_\tau$ consists of *all* Borel selectors $s : Q \rightarrow W$ such that $x \in K_{s(x)}$. Its closed-loop block map is $T_s(x) = g_{s(x)}(x)$. All Borel mixtures of legal selectors belong to $\mathcal{S}_\tau$. No continuity or dependence on finitely many coordinates is imposed. A selector is used without an additional clock, register, or feedback memory.

For a finite window F , let

$$P_s(F) = \left| \{(s(T_s^k x))_{k \in F} : x \in Q\} \right|.$$

Define $h_A(s)$ and $h_{\text{pat}}^*(s)$ from these counts as in (9.1)–(9.2), but dividing by $n\tau$ rather than n . Thus normalization is per sampled control block, divided by its fixed length; it is *not* division by elapsed time τa_n . Put

$$L_\tau = \sup_A \inf_{s \in \mathcal{S}_\tau} h_A(s), \quad M_\tau = \inf_{s \in \mathcal{S}_\tau} h_{\text{pat}}^*(s). \quad (11.2)$$

Always $L_\tau \leq M_\tau$.

The counts use actual closed-loop trajectories. If instead one takes the closure of their one-sided name set, the finite patterns are unchanged: a cylinder specifying finitely many coordinates is open, so if it meets that closure, it meets the original name set. Taking a name closure therefore introduces no extra finite-window witnesses.

Lemma 11.1. *The infima in (11.2), and the inner infimum at any fixed A , are unchanged if one allows every finite Borel safe partition with a control block assigned to each atom, and counts atom names rather than control-block names.*

Proof. Merge all atoms assigned the same block. Their finite union is Borel and remains inside that block's safety domain. The resulting selector has exactly the same closed-loop state map. The map from original atom names to block names is coordinatewise and onto the actual block-name patterns, so every block-pattern count is no larger than the original atom-pattern count. Conversely, the nonempty fibers of any legal selector form a legal finite Borel safe partition whose atom and block counts agree. Both directions of the infimum equality follow. $\square$

Theorem 11.2. *Assume in addition that there is a map $T : Q \rightarrow Q$ such that*

$$g_w(x) = T(x) \quad \text{for every } w \in W \text{ and every } x \in K_w, \quad (11.3)$$

and that T is conjugate to σ^p on a mixing one-sided shift of finite type. With $r = N((K_w)_{w \in W})$, for every A satisfying (9.3) one has

$$\min_{s \in \mathcal{S}_\tau} h_A(s) = \min_{s \in \mathcal{S}_\tau} h_{\text{pat}}^*(s) = L_\tau = M_\tau = \frac{\log r}{\tau}. \quad (11.4)$$

The maximum defining L_τ is attained by every such A , and an ordered minimum-subcover selector attains all the indicated minima. The same minimum and minimax values hold when all finite Borel safe partitions, including every finite Borel refinement, are allowed. Every refinement satisfies the stated lower bound; no upper bound for each individual refinement is asserted.

Proof. For every legal selector, (11.3) gives $T_s = T$ pointwise. Its fibers form a finite Borel refinement of $(K_w)_{w \in W}$. Theorem 9.1, with normalization by τ , gives the lower bound at every A with diverging gaps. If $w_1, \dots, w_r$ form a minimum subcover, define $s_0(x) = w_j$ on the sets E_j of (9.7). This is Borel and legal at every point. Each selected block stays in Q at its intermediate times by (11.1), and the endpoint again lies in Q , so recursion gives an infinite safe closed-loop orbit for every initial state. The selector uses only r blocks and thus gives the upper bound for every window and every sampling sequence. This proves attainment, (11.4), and the outer maximum. Lemma 11.1 proves the last assertion. $\square$

The standard safe-partition framework is described by Kawan [7, Chapter 2]. His minimization over control periods in invariance entropy is a different optimization. Here τ is fixed, intermediate safety is retained, and the exact value comes from the additional common-successor and finite-type hypotheses.

11.1. A positive overlapping example at period two. Take

$$Q = \{0, 1, 2\}^{\mathbb{N}_0}, \quad X = Q \sqcup \{\dagger\}, \quad U = \{0, 1, 2\}, \quad \tau = 2,$$

where the isolated point $\dagger$ lies outside Q . Define

$$f_u(x) = \begin{cases} \sigma x, & x \in Q, x_0 \neq u, \\ \dagger, & x \in Q, x_0 = u, \end{cases} \quad f_u(\dagger) = \dagger.$$

Each f_u is continuous since the cases are clopen. For $w = (a, b)$,

$$K_{(a,b)} = \{x : x_0 \neq a, x_1 \neq b\}, \quad g_{(a,b)}|_{K_{(a,b)}} = \sigma^2.$$

All nine control blocks are available to Borel feedbacks, and each state permits four of them. A safety domain covers four of the nine two-symbol input blocks, so two domains cannot cover Q . The three diagonal blocks $(0,0), (1,1), (2,2)$ do cover Q , since some symbol differs from both x_0 and x_1 . Hence $r = 3$, and

$$L_2 = M_2 = \frac{\log 3}{2}.$$

An attaining selector is

$$c(x) = \min(\{0, 1, 2\} \setminus \{x_0, x_1\}), \quad s_0(x) = (c(x), c(x)).$$

It is clopen and legal everywhere. Its actual block-boundary states are $\sigma^{2k}x$. To prescribe any sequence of diagonal labels, use successive input blocks 11, 00, and 01 for the desired labels 0, 1, and 2, respectively. Therefore its actual name set is the full shift on these three blocks. The lower bound, however, holds for every legal Borel rule using any of the nine blocks.

This example does not preassign a nonconstant common control-letter observation to every feedback. More precisely, suppose a map $\rho : U^2 \rightarrow V'$ and an observation $F : Q \rightarrow V'$ satisfy $\rho(s(x)) = F(x)$ for every legal Borel selector and every x . Any two blocks (a, b) and (a', b') are colegal at a state beginning with ij , where $i \notin \{a, a'\}$ and $j \notin \{b, b'\}$. A legal baseline selector may be changed at this single state to use either block, while remaining Borel and legal. Hence their ρ values agree. All pairs of blocks are colegal somewhere, so ρ is constant. This rules out this particular forced-letter explanation; it makes no claim about every possible dynamical factor.

12. A FEEDBACK-DEPENDENT BOUNDARY WITH AN EXACT FULL-CLASS VALUE

The common-successor assumption cannot be removed solely because each branch is continuous and open. We give a system with individual prefix cylinders as safety domains and compute its full-class value.

Let $Q = \{0, 1\}^{\mathbb{N}_0}$, let $\mathcal{C} = \{0, 10, 11\}$ be a set of finite words, and take $U = \mathcal{C} \times \{0, 1\}$ and $\tau = 1$. For $e \in \{0, 1\}$ let P_e add e modulo two to every coordinate. On $X = Q \sqcup \{\dagger\}$ define

$$f_{(c,e)}(x) = \begin{cases} P_e \sigma x, & x \in [c], \\ \dagger, & x \in Q \setminus [c], \end{cases} \quad f_{(c,e)}(\dagger) = \dagger. \quad (12.1)$$

The cylinders $[0], [10], [11]$ are disjoint and cover Q . Each state has exactly two legal controls with different successors. The global maps $f_{(c,e)}$ are continuous, and their legal restrictions are open. The safety domains are $K_{(c,e)} = [c]$, and $r = 3$.

Proposition 12.1. *For the system (12.1), over all legal Borel feedbacks and arbitrary Borel mixtures,*

$$\min_s h_{A_0}(s) = \min_s h_{\text{pat}}^*(s) = \max_A \inf_s h_A(s) = \log 2 < \log r, \quad A_0 = (0, 1, 2, \dots). \quad (12.2)$$

There is one clopen legal feedback s_f satisfying $h_A(s_f) = \log 2$ for every increasing infinite A . The same full-class values hold if arbitrary finite Borel atom refinements are allowed.

Proof. Let $c(x)$ be the unique member of $\mathcal{C}$ with $x \in [c(x)]$, and set

$$s_f(x) = (c(x), x_1), \quad T_f x = P_{x_1} \sigma x.$$

This is a clopen selector, legal at every state. Its four nonempty initial control fibers, for labels $(0,0), (0,1), (10,0), (11,1)$, are $[00], [01], [10], [11]$, respectively. For all $n \geq 1$, direct induction gives

$$T_f^n x = P_{x_n} \sigma^n x, \quad s_f(T_f^n x) = (0, x_{n+1} \oplus x_n), \quad (12.3)$$

where $\oplus$ is addition modulo two. In the induction step the chosen flip is $x_{n+1} \oplus x_n$; it cancels the previous flip and leaves $P_{x_{n+1}} \sigma^{n+1} x$. The first coordinate is always zero after time zero, which also proves all-time safety.

For every finite nonempty window F , with $n = |F|$, one has

$$P_{s_f}(F) = \begin{cases} 2^n, & 0 \notin F, \\ 2^{n+1}, & 0 \in F. \end{cases} \quad (12.4)$$

The four initial labels and two later labels give the upper bounds. To prove the lower bounds, prescribe any of the four initial labels if $0 \in F$, thereby fixing (x_0, x_1) ; if $0 \notin F$, set $x_0 = x_1 = 0$. At positive sampled times prescribe the desired second components d_k , and put $d_k = 0$ at the other positive times. Recursively set $x_{k+1} = x_k \oplus d_k$ for every $k \geq 1$. This constructs a full state in Q whose actual trajectory under the same selector has the required sampled controls by (12.3). Thus all counted patterns are realized by true infinite closed-loop trajectories. Equation (12.4) implies $h_A(s_f) = h_{\text{pat}}^*(s_f) = \log 2$ for every A .

For the opposite bound fix any legal Borel selector s . If its first n actual controls from x are (c_k, e_k) , let $E_0 = 0$ and $E_k = e_0 \oplus \cdots \oplus e_{k-1}$. Since shifts commute with flips,

$$T_s^k x = P_{E_k} \sigma^k x.$$

Let $b(c)$ be the first bit of the word c . Safety at time k gives

$$x_k = b(c_k) \oplus E_k \quad (0 \leq k < n). \quad (12.5)$$

Thus one actual length- n control word determines at most one input prefix of length n . Every binary input prefix has an extension in Q , and the feedback is defined and safe on every such extension. Distinct prefixes must yield distinct actual control words. Hence

$$P_s(\{0, \dots, n-1\}) \geq 2^n$$

for every n and every legal s . This proves $h_{A_0}(s) \geq \log 2$ and $h_{\text{pat}}^*(s) \geq \log 2$. Combining these inequalities with s_f proves (12.2), including attainment. Lemma 11.1 handles atom refinements. $\square$

The obstruction has a precise orbit interpretation. In this example $T_f(Q) = [0]$: for the reverse inclusion, given $y \in [0]$, take $x = 0y$. Therefore neither $[10]$ nor $[11]$ is visited at a positive time. The initial fibers with those labels are nonmeager, but their labels cannot be freely concatenated in the future. The map T_f is itself continuous and open, as follows by restricting to the four clopen two-symbol cylinders. It is not onto Q .

Even preservation of meager preimages does not repair this problem. For any selector s and any n , the value of $T_s^n x$ is either $\sigma^n x$ or $P_1 \sigma^n x$. Hence, for a meager set N ,

$$(T_s^n)^{-1} N \subset (\sigma^n)^{-1} N \cup (P_1 \sigma^n)^{-1} N$$

is meager. What fails is the nonempty open intersection used in Lemma 9.2 for the *selected* dynamics. Also, the same s_f gives

$$\inf_s h_A(s) \leq \log 2 = \log r - \log(3/2) \quad \text{for every } A.$$

The failure of the cover formula is therefore a uniform numerical gap over all samplings, rather than a failure at one finite window. It does not refute $L_\tau = M_\tau$: equality holds in this very example. Nor does the proof determine the inner infimum for every other fixed sampling sequence.

13. INVERSE CAPACITIES AND A FULL-FEEDBACK PATTERN THEOREM

13.1. Branch measures, lists, and actual fibers. In this and the following sections the shorter notation $h^*(s) = h_{\text{pat}}^*(s)$ denotes the already defined maximal-pattern rate, and $p_s^*(n) = \max_{|F|=n} P_s(F)$. The notation $h_A(s)$ retains the same sampled rate and the same $n\tau$ normalization. Only notation is shortened.

Fix a period $\tau \geq 1$ and a finite collection W of admissible control blocks, with $q = |W|$. For each $u \in W$ let $K_u \subset Q$ be its closed safety domain, including all intermediate-time safety conditions, and $g_u : K_u \rightarrow Q$ its continuous endpoint map. Assume these domains cover the nonempty compact metric space Q . The legal class $\mathcal{S}_\tau$ consists of all Borel maps $s : Q \rightarrow W$

with $x \in K_{s(x)}$. Define $T_s(x) = g_{s(x)}(x)$. Reusing this same map at each block boundary gives an infinite safe trajectory for every state. No continuity of s or of T_s is imposed.

Let μ be a full-support Borel probability on Q and suppose

$$\nu_u = (g_u)_*(\mu|_{K_u}) \ll \mu, \quad \rho_u = \frac{d\nu_u}{d\mu} \in L^\infty(\mu). \quad (13.1)$$

The positive bounded operator $L_u : L^\infty(\mu) \rightarrow L^\infty(\mu)$ is specified by

$$\int h L_u f d\mu = \int_{K_u} h(g_u x) f(x) d\mu(x) \quad (h, f \text{ bounded Borel}). \quad (13.2)$$

Indeed, the signed pushforward of $f\mu|_{K_u}$ is dominated by $\|f\|_\infty \nu_u$; its density defines $L_u f$ and $|L_u f| \leq \|f\|_\infty \rho_u$. Changing f on a μ -null set changes no pushforward measure, so this is well defined on L^∞ classes. In particular $L_u \mathbf{1} = \rho_u$.

Fix $2 \leq k \leq q$ and put $l = k - 1$. A list $S \subset W$ is a *global* set of l labels, not a state-dependent choice. Write $A_S = \sum_{u \in S} L_u$, and define

$$b_0 = 1, \quad b_n = \max_{|S_0|=\dots=|S_{n-1}|=l} \|A_{S_{n-1}} \cdots A_{S_0} \mathbf{1}\|_\infty. \quad (13.3)$$

Products are ordered in the forward pushforward order. Positivity gives $\|P\|_{\infty \rightarrow \infty} = \|P\mathbf{1}\|_\infty$ for every positive product P . Consequently $b_{n+m} \leq b_n b_m$. If $b_h < 1$ for some $h \geq 1$, choose $b_h < \theta^h < 1$ and set

$$H = \max_{0 \leq j < h} b_j \theta^{-j}. \quad b_n \leq H \theta^n \quad (n \geq 0). \quad (13.4)$$

This follows by writing $n = ah + j$ and using submultiplicativity. The proof length h is not a change of control period.

For a fixed feedback let

$$E_w(s) = \{x : s(T_s^j x) = w_j \ (0 \leq j < n)\}, \quad F_n(s) = \{w \in W^n : E_w(s) \neq \emptyset\}.$$

The $E_w(s)$ are Borel, pairwise disjoint, and cover Q . For $w = w_0 \cdots w_{n-1}$ put $L_w = L_{w_{n-1}} \cdots L_{w_0}$ and $P_F = \sum_{w \in F} L_w$.

Lemma 13.1 (Actual-fiber capacity). *For every legal Borel feedback and every $n \geq 1$,*

$$1 \leq \int P_{F_n(s)} \mathbf{1} d\mu. \quad (13.5)$$

If $F_n(s)$ is covered by t boxes $S_0^a \times \cdots \times S_{n-1}^a$ whose lists have size at most l , then $1 \leq t b_n$.

Proof. For a nonempty fiber E_w , push $\mu|_{E_w}$ along its actual trajectory. Initially the density is $\mathbf{1}_{E_w}$. At its j th block the entire pushed measure is supported on K_{w_j} , because all these states really use that safe block. Thus (13.2), successively tested against bounded Borel functions, gives the terminal density $L_w \mathbf{1}_{E_w}$, of mass $\mu(E_w)$. Positivity gives $L_w \mathbf{1}_{E_w} \leq L_w \mathbf{1}$. Summing over all nonempty actual fibers proves (13.5). Null but nonempty fibers remain in $F_n(s)$; they have not been removed from the combinatorial names.

Complete every nonempty list to size l . Distributivity and positivity give

$$P_{F_n(s)} \mathbf{1} \leq \sum_{a=1}^t A_{S_{n-1}^a} \cdots A_{S_0^a} \mathbf{1}.$$

Repeated boxes and overlaps only enlarge this upper bound. Unsafe words in a box are subject to the original safety gates in L_u . Integrating and using (13.3) yields $1 \leq t b_n$. $\square$

13.2. The finite combinatorial input and its application. For $F \subset W^n$, let $C_l(F)$ be its minimum covering number by l -list boxes. Given a k -element set $V \subset W$, let $C_V(F)$ be the minimum number of boxes of the form

$$\prod_{j=0}^{n-1} (W \setminus \{v_j\}), \quad v_j \in V,$$

needed to cover F . This is a cover, since $k \geq 2$.

We use the following already established finite-set lemma. For every $k \geq 2$ and $b > 0$ there are $c > 0$ and N such that if $G \subset \{0, 1, \dots, k\}^n$, $n \geq N$, needs at least e^{bn} boxes omitting one nonzero letter per coordinate, then some $J \subset \{0, \dots, n-1\}$ with $|J| \geq cn$ has

$$G|_J \supset \{1, \dots, k\}^J. \quad (13.6)$$

This is the positive-density conclusion of Huang–Ye [4, Lemma 4.1 and Remark 4.2, pp. 1701–1705]. It is a statement about finite sets; no continuity, invertibility, or invariant measure of a closed loop is required. Its constants depend only on b, k .

For clarity, the entire reduction from list boxes to that lemma is included. Choose a minimum omitted-letter cover for every V and intersect all these covers. The intersections cover F , with at most $\prod_V C_V(F)$ members. In a nonempty intersection each coordinate contains at most $k-1$ letters: if it contained a k -set V_0 , its factor from the V_0 cover would have omitted a letter of V_0 . Hence

$$C_l(F) \leq \prod_{V \in \binom{W}{k}} C_V(F). \quad (13.7)$$

For actual names Lemma 13.1 gives $C_l(F_n(s)) \geq b_n^{-1}$. Under (13.4), for all sufficiently large n this lower bound is at least $e^{\eta m/2}$, where $\eta = -\log \theta > 0$. With $J_0 = \binom{q}{k}$, (13.7) implies that some V satisfies

$$C_V(F_n(s)) \geq e^{\eta m/(2J_0)}.$$

Code the members of V by $1, \dots, k$ and all other letters by 0. Omitted-letter boxes pull back exactly to the boxes defining C_V , so the coded set has precisely that covering number. Applying (13.6) and undoing this coding yields $F_n(s)|_J \supset V^J$. No distinct members of V were merged.

Theorem 13.2 (Product inverse-capacity criterion). *Assume (13.1). If $b_h < 1$ for some $h \geq 1$, then there are $\alpha > 0$ and N , depending only on the branches, μ, k , such that, for every legal Borel feedback s and every $n \geq N$, its actual length- n names have a k -valued cube projection on at least αn coordinates. Consequently*

$$p_s^*(m) \geq k^m \quad (m \geq 1), \quad h^*(s) \geq \frac{\log k}{\tau}. \quad (13.8)$$

The same lower bound holds for all finite Borel atom refinements. If $k = r \geq 2$ is the minimum number of safety domains covering Q , then

$$M_\tau = \min_s h^*(s) = \frac{\log r}{\tau}, \quad (13.9)$$

both over unrefined selectors and over all finite Borel safe partitions.

Proof. The preceding argument supplies α, N independent of s . Given m , choose n so large that $\alpha n \geq m$ and retain m of the independent positions. All k^m patterns are actual feedback names, proving (13.8) with the prescribed normalization by $m\tau$.

For $k = r$, choose a minimum subcover $K_{u_1}, \dots, K_{u_r}$ and select u_j on $K_{u_j} \setminus \bigcup_{i < j} K_{u_i}$. This is a Borel state function, legal at every point. Each chosen block stays safe at its intermediate times and ends in Q ; applying the same rule at the next state gives infinite safety. It uses r labels and has at most r^m patterns on any m positions. This proves (13.9) and attainment. For a refined partition, projection to its physical control labels is onto the actual coarse patterns and preserves the dynamics. The lower bound therefore passes to

every refinement, while the unrefined selector remains an available upper-bound witness in the enlarged class. $\square$

The theorem asserts a uniform *density*, not a common set of independent positions. The sets J and V may depend on both s and n . In particular it proves no equality $L_\tau = M_\tau$ without an additional common-sampling argument.

13.3. Single-step scalar and joint-density criteria. Define

$$\Theta_k = \left\| \max_{|S|=k-1} \sum_{u \in S} \rho_u \right\|_\infty. \quad (13.10)$$

If $\Theta_k < 1$, then $b_1 = \Theta_k < 1$ and Theorem 13.2 applies. If the branch capacities obey $\mu(K_u \cap g_u^{-1}E) \leq c_u \mu(E)$ for every Borel E , then $\rho_u \leq c_u$ almost everywhere, and the sum of the largest $k-1$ of the c_u bounds Θ_k . Thus a scalar-capacity sum smaller than one is a sufficient special case. The order of taking the branch supremum and summing matters: joint densities may pass the test while the individual optimal scalar bounds do not.

The joint-density case also has a direct quantitative proof worth recording. Let $d_k(F)$ be the largest size of a coordinate set on which $F \subset W^n$ contains a product of k letters per coordinate. Put $\theta = \Theta_k < 1$, $C_0 = \|\sum_u \rho_u\|_\infty$ and $B = C_0 \binom{n}{k}$. For nonnegative f let $\Phi f = \max_{|S|=k-1} \sum_{u \in S} L_u f$ and $\mathcal{T}f = \sum_u L_u f$. Their norms are bounded by θ and C_0 , respectively.

Proposition 13.3 (Weighted shattering estimate). *For every $F \subset W^n$,*

$$\|P_F\|_{\infty \rightarrow \infty} \leq H(n, d_k(F)), \quad H(n, d) = \sum_{j=0}^{\min(d, n)} \binom{n}{j} B^j \theta^{n-j}. \quad (13.11)$$

For actual names this gives $d_k(F_n(s)) > \alpha n$, uniformly in every legal feedback, whenever $0 < \alpha < 1/2$ satisfies

$$\log \theta + \mathfrak{h}(\alpha) + \alpha \log(B/\theta) < 0, \quad \mathfrak{h}(t) = -t \log t - (1-t) \log(1-t). \quad (13.12)$$

Proof. Take $H(n, -1) = 0$ and treat the empty word at $n = 0$ separately. Split F at its *first* letter: $F_a = \{v : av \in F\}$, $U_0 = \bigcup_a F_a$, and $I_S = \bigcap_{a \in S} F_a$ for $|S| = k$. If $d = d_k(F)$, then $d_k(U_0) \leq d$ and $d_k(I_S) \leq d-1$: independence of d tail positions in the latter would also make the first position independent and give $d+1$. For each $v \in U_0$ the set of its permitted first letters has either at most $k-1$ members, or contains a k -set S . Applying the positive tail product L_v and summing gives

$$P_F f \leq P_{U_0} \Phi f + \sum_{|S|=k} P_{I_S} \mathcal{T} f.$$

This respects the order $L_{av} = L_v L_a$; no operators are commuted. Induction and Pascal's identity give $\theta H(n-1, d) + B H(n-1, d-1) = H(n, d)$. Positive domination handles signed f and proves (13.11).

By actual-fiber mass, $1 \leq H(n, d_k(F_n(s)))$. For $d \leq \alpha n$, $B/\theta \geq 1$ and $\sum_{j \leq \alpha n} \binom{n}{j} \leq e^{n\mathfrak{h}(\alpha)}$ give $H(n, d) \leq \exp(n[\log \theta + \mathfrak{h}(\alpha) + \alpha \log(B/\theta)]) < 1$, a contradiction. The binomial bound follows by comparing each term to its probability under independent Bernoulli(α) trials; those probabilities sum to at most one. Condition (13.12) is possible because its last two terms tend to zero as α tends to zero. $\square$

13.4. A smooth joint-density example. On $Q = [0, 1]$ take three full inverse branches with

$$\begin{aligned} l_u &= \frac{19u}{60}, & \rho_u(y) &= \frac{11}{30} + \frac{7}{30} \cos(2\pi(y - u/3)), \\ \psi_u(y) &= l_u + \frac{11}{30}y + \frac{7}{60\pi} [\sin(2\pi(y - u/3)) - \sin(-2\pi u/3)]. \end{aligned} \quad (13.13)$$

Here $g_u = \psi_u^{-1}$, for $u = 0, 1, 2$. Their derivatives lie between $2/15$ and $3/5$, so the branches are strictly increasing and expanding in the forward direction. The closed safety intervals are $[0, 22/60]$, $[19/60, 41/60]$, and $[38/60, 1]$. They cover Q and each is needed: the middle interval alone covers the open gap between $22/60$ and $38/60$, while the two endpoints require the first and third intervals. Hence $r = 3$.

With Lebesgue probability, the densities are exactly $\psi'_u = \rho_u$. The three cosine terms sum to zero. The sum of two is the negative of the remaining cosine, whose maximum is one; therefore

$$\Theta_3 = \frac{22}{30} + \frac{7}{30} = \frac{29}{30} < 1.$$

Each branch has optimal scalar bound $3/5$, since its continuous density reaches that value. The largest two scalar bounds sum to $6/5$. The joint criterion consequently gives $M_1 = \log 3$ although that Lebesgue scalar criterion fails.

To place these interval branches in a continuous compact ambient system, take $X = [-1, 2]$. Extend each g_u to the left of its safety interval $[a_u, b_u]$ by $-\min\{a_u - x, 1\}$ and to its right by $1 + \min\{x - b_u, 1\}$. At the endpoints these formulas match 0 and 1. Outside the closed safety interval the image lies outside Q , so this extension changes neither safety domains nor legal dynamics. The same construction applies to full monotone interval branches elsewhere in the paper.

The fixed-reference separation proved here is not, by itself, a separation from every full-support reference probability. The bounded positive-density obstruction is recorded in Appendix D; its extension to arbitrary full-support probabilities remains unresolved for this smooth example.

13.5. An exact product separation from every single-step reference test. Let $Q = (\mathbb{Z}/3\mathbb{Z})^{\mathbb{N}_0}$, let P_u add u to each coordinate, and use $g_u = P_u\sigma$ at period one. Put

$$K_0 = [02] \cup [12] \cup [22], \quad K_1 = [00] \cup [20], \quad K_2 = [01] \cup [10] \cup [11] \cup [12] \cup [21].$$

The only overlap is $[12]$. The cylinders $[02], [00], [01]$ require controls 0, 1, 2, respectively, so $r = 3$. On $[12]$, choosing 0 or 2 changes the successor's first letter from 2 to 1. Every Borel mixture on this entire overlap is allowed. Illegal controls go to an isolated fixed cemetery point; the resulting maps on $Q \sqcup \{\dagger\}$ are continuous.

For fair Bernoulli probability, the operators leave first-coordinate functions invariant and have matrices $M_u/3$, where

$$M_0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix}, \quad M_1 = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad M_2 = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

This follows from splitting the source measure by its first letter, and retaining exactly the allowed two-letter cylinders. For a positive product the full L^∞ norm equals its norm on $\mathbf{1}$, whose entire orbit stays in this invariant space. Thus this finite representation computes the full product norms, not an approximation to arbitrary Borel functions.

Let $A = M_0 + M_1$, $B = M_0 + M_2$, $C = M_1 + M_2$. The vectors after two of these unnormalized factors are

$$\begin{array}{ccc} (0, 3, 5) & (0, 7, 8) & (0, 4, 7) \\ (5, 2, 7) & (8, 1, 9) & (7, 3, 10) \\ (5, 5, 2) & (8, 8, 1) & (7, 7, 3). \end{array}$$

They are coordinatewise dominated by one of the vectors in

$$V = \{(0, 7, 8), (7, 3, 10), (7, 7, 3), (8, 1, 9), (8, 8, 1)\}.$$

For $v = (p, q', r')$, $Av = (0, p + r', p + q' + r')$, $Bv = (p + q' + r', q', p + 2q' + r')$, and $Cv = (p + q' + r', p + q' + r', q')$. For the five vectors of V , their largest resulting coordinate is, respectively, 22, 23, 24, 19, 25. Also $BCB\mathbf{1} = (17, 8, 25)$. Consequently $b_3 = 25/27 < 1$. Submultiplicativity, rather than an extrapolation from finite calculations, controls all lengths.

Theorem 13.2 gives $M_1 = \log 3$, and every unrefined feedback has that maximal-pattern rate. An explicit safe selector chooses 0 if $x_1 = 2$, otherwise 1 if $x_1 = 0$ and $x_0 \neq 1$, and otherwise 2.

On the three target first-letter cylinders the densities are $(0, 0, 1)$, $(0, 2/3, 1/3)$, and $(1, 0, 1/3)$. Hence $\Theta_3 = 4/3$. This failure persists under *every* full-support probability ν . If one of the safe branch measures has no bounded ν -density, the single-step criterion already fails its prerequisites. Otherwise $g_0(K_0) \subset [2]$ and $g_1(K_1) \subset [1]$ give disjoint density supports. At most two densities are nonzero at a point, so their largest two sum to their total. Its integral is

$$\sum_{u=0}^2 \nu(K_u) = 1 + \nu([12]) > 1.$$

Thus $\Theta_3(\nu) > 1$ for every such ν , including singular or atomic probabilities. This excludes both original single-step tests, but not all possible changes of presentation or iteration.

There is also a distinction from iterating the pointwise envelope Φ . On nonnegative functions the disjoint supports give $\Phi = \sum_u L_u$. Under fair measure its first-letter matrix is $\frac{1}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 2 & 1 \end{pmatrix}$, with positive eigenvector $(1, 1, (\sqrt{13} - 1)/2)$ and eigenvalue $(3 + \sqrt{13})/6 > 1$. Hence $\|\Phi^n \mathbf{1}\|_\infty$ grows, while every history of *global* two-label lists decays. Maximization by state and composition cannot be interchanged.

14. FINITE-CYLINDER SAFETY AND STRUCTURAL SOURCE BUDGETS

14.1. Exact finite representation without restricting feedbacks. Let D be a finite alphabet of size $d \geq 2$, $Q = D^{\mathbb{N}_0}$, and μ its fair Bernoulli probability. The finite controls have legal maps $g_u = P_u \sigma$, where p_u permutes D and P_u acts in every coordinate. Suppose

$$K_u = \bigcup_{\xi \in R_u} [\xi], \quad R_u \subset D^h, \quad h \geq 2,$$

and the K_u cover Q . Shorter cylinders can be completed to the same length h . The continuous cemetery extension used above applies to these clopen domains. The control period is always one.

For every bounded Borel function the full operator is

$$L_u f(y) = \frac{1}{d} \sum_{a \in D} \mathbf{1}_{R_u}(a, p_u^{-1} y_0, \dots, p_u^{-1} y_{h-2}) f(a P_u^{-1} y). \quad (14.1)$$

Indeed, conditioned on $x_0 = a$ the tail of x still has law μ , and P_u preserves that law. Substitution in (13.2) gives this formula with the exact safety gate.

Index source and target vectors by D^{h-1} . For target η and source ξ , set

$$T_u(\eta, \xi) = \frac{1}{d} \# \left\{ a \in D : \begin{array}{l} (a, p_u^{-1} \eta_0, \dots, p_u^{-1} \eta_{h-2}) \in R_u, \\ (a, p_u^{-1} \eta_0, \dots, p_u^{-1} \eta_{h-3}) = \xi \end{array} \right\}. \quad (14.2)$$

When $h = 2$ the second tuple is just a . Functions of the first $h - 1$ coordinates form an invariant subspace of every L_u , represented by T_u . Starting at $\mathbf{1}$, every list product stays in this space. Every such cylinder has positive measure and a positive product has norm equal to its value on $\mathbf{1}$. Therefore, for every length and every history,

$$\|A_{S_{n-1}} \cdots A_{S_0}\|_{\infty \rightarrow \infty} = \|T_{S_{n-1}} \cdots T_{S_0} \mathbf{1}\|_\infty, \quad T_S = \sum_{u \in S} T_u. \quad (14.3)$$

This equality is on the full L^∞ space. It does not claim that an arbitrary Borel function is approximated by a fixed finite space, or that a Borel feedback uses a finite-state rule. The finite indices are proof tools, not new state coordinates or a clock.

For two-letter domains, rows are target letters and columns are source letters, with

$$T_u(j, i) = d^{-1} \mathbf{1}_{R_u}(i, p_u^{-1} j), \quad d(w^\top T_u)_i = \sum_{z: (i, z) \in R_u} w(p_u z). \quad (14.4)$$

The structural conditions below use this source-column identity. A left budget $w^\top T_S \leq \theta w^\top$ is different from changing a reference density, which leads instead to a right inequality $T_S a \leq \theta a$. They cannot be interchanged.

14.2. A strict source-weight criterion. Partition $D = A \sqcup B \sqcup C$ into nonempty sets of sizes a, b, c , so $d = a + b + c$. Use three controls with

$$p_0(C) \subset D \setminus A, \quad p_1(B) \subset A, \quad p_2(A) \subset D \setminus A. \quad (14.5)$$

Choose arbitrary relations $E_0 \subset A \times C$ and $E_1 \subset A \times B$, with $E_0 \cup E_1 \neq \emptyset$, and define

$$\begin{aligned} R_0 &= ((D \setminus A) \times C) \cup E_0, \\ R_1 &= ((D \setminus A) \times B) \cup E_1, \\ R_2 &= (A \times D) \cup (D \times A), \quad K_u = [R_u]. \end{aligned} \quad (14.6)$$

Put $e = \max_{i \in A} |\{z : (i, z) \in E_1\}|$. Require only the following counts and landing conditions:

$$s_- := \max \left\{ \frac{c}{b+c}, \frac{e}{b+c-e} \right\} < s_+ := \frac{\min\{a, c\}}{b}. \quad (14.7)$$

These are direct conditions on the original allowed cylinders.

Theorem 14.1 (Strict source budget). *For every system satisfying (14.5)–(14.7), $r = 3$, all histories of global two-label lists have exponentially decaying inverse capacity, and every legal unrefined Borel feedback has $h^*(s) = M_1 = \log 3$. Every finite Borel atom refinement has $h^* \geq \log 3$. Moreover, no full-support reference probability can directly satisfy the joint-density or scalar single-step criterion for $k = 3$ on these original branches.*

Proof. Take $s = (s_- + s_+)/2$, $\gamma = 1 + s$, and let $w_i = \gamma$ on A , $w_i = 1$ elsewhere. Write $H = \sum_i w_i = d + sa$. Before division by d , the weighted contributions of controls 0, 1, 2 in a source column are

	0	1	2
$i \in A$	$e_0(i) \leq c$	$\gamma e_1(i) \leq \gamma e$	H
$i \notin A$	c	γb	a .

Each entry follows from the original domains and (14.5). For example, on A control 2 permits every tail, hence the sum of all weights irrespective of its permutation. Off A , its permitted tails lie in A and land outside A , each with weight one.

Since $sb < a, c$, both c and $\gamma e \leq \gamma b$ are smaller than H . Thus every two-label sum on A is bounded by $H + \max\{c, \gamma e\}$. Define

$$\theta = \max \left\{ \frac{H + \max\{c, \gamma e\}}{d\gamma}, \frac{c + \gamma b}{d}, \frac{c + a}{d}, \frac{\gamma b + a}{d} \right\}.$$

We have $d\gamma - H = s(b + c) > c, \gamma e$ because $s > s_-$. The other three terms are smaller than one by $sb < a, c$ and $b > 0$. Hence $0 < \theta < 1$ and $w^\top T_S \leq \theta w^\top$ for every global two-label list. Repeatedly applying this single inequality gives

$$\|T_{S_{n-1}} \cdots T_{S_0} \mathbf{1}\|_\infty \leq w^\top T_{S_{n-1}} \cdots T_{S_0} \mathbf{1} \leq H\theta^n,$$

since $\min_i w_i = 1$. Equation (14.3) controls the complete capacity for all lengths.

If $x_0 \in A$, control 2 is legal; otherwise its next letter in A, B, C requires, respectively, 2, 1, 0. Thus the domains cover Q . Taking $x_0 \notin A$ and x_1 successively in C, B, A gives cylinders on which each of 0, 1, 2 alone is legal, so $r = 3$. A clopen attaining selector chooses 2 when $x_0 \in A$ or $x_1 \in A$, otherwise 1 when $x_1 \in B$, otherwise 0. It is safe, single-valued, and stays in Q at each step; the same rule at every successor gives all-time safety. Applying Theorem 13.2 proves all pattern conclusions, including the lower bound for arbitrary refinements.

Let $O = [E_0 \cup E_1]$. It is nonempty and clopen, and safety multiplicity is exactly $1 + \mathbf{1}_O$. For a full-support probability ν , either some branch fails the bounded-density prerequisite, or all densities exist. In the latter case the images of controls 0 and 1 lie, respectively, in $[D \setminus A]$ and $[A]$; their supports are disjoint. The largest two densities consequently sum to all

three almost everywhere. Their integral is $\sum_u \nu(K_u) = 1 + \nu(O) > 1$. Thus $\Theta_3(\nu) > 1$. The scalar condition implies $\Theta_3 < 1$ and also fails. $\square$

This is a system family, not one matrix calculation. For instance $1 \leq b \leq a \leq b + c$ and $c > b$ imply (14.7) for every possible $e \leq b$: $e/(b + c - e) \leq b/c < 1$, $c/(b + c) < 1$, while $s_+ \geq 1$. Every allowed overlapping relation and permutation landing pattern is covered. Constants can be uniform when these counts have fixed bounds.

For a concrete example take $D = \{0, 1, 2, 3, 4\}$, $A = \{0, 1\}$, $B = \{2\}$, $C = \{3, 4\}$, $E_0 = \{(0, 3), (1, 4)\}$, $E_1 = \{(0, 2)\}$. Let p_0 be the identity, $p_1 = (0\ 2)$, and specify p_2 by $0 \mapsto 2$, $1 \mapsto 3$, $2 \mapsto 0$, $3 \mapsto 4$, $4 \mapsto 1$. Here $s = 4/3$, $\gamma = 7/3$, $H = 23/3$, $\theta = 13/15$, and $b_n \leq (23/3)(13/15)^n$. The target densities at first letters 0, 1, 2, 3, 4 are, respectively,

$$(0, 4/5, 2/5), \quad (0, 0, 2/5), \quad (0, 0, 1), \quad (4/5, 0, 1), \quad (4/5, 0, 2/5).$$

Thus $\Theta_3 = 9/5$. In [14] the two legal successors have first letters 4 and 1. No common successor is present. The natural $A/B/C$ merging also fails to commute with the listed permutations; this observation does not exclude every possible factor.

14.3. Cross-layer dispersion without a strict left budget. Let U have size $q \geq 2$. Partition $D = \bigsqcup_{t \in U} I_t$ into nonempty source classes, and let $m(i) = t$ on I_t . For each i partition the tail alphabet by

$$D = \bigsqcup_{u \neq m(i)} V_{i,u}, \quad F_i \subset D.$$

Control $u \neq m(i)$ is the primary legal control on $V_{i,u}$, while control $m(i)$ is also legal on F_i . Precisely,

$$R_u = \{(i, z) : u \neq m(i), z \in V_{i,u}\} \cup \{(i, z) : u = m(i), z \in F_i\}. \quad (14.8)$$

The permutations p_u are arbitrary subject to the count below. Assume

$$a_0 = \min_{i, u \neq m(i)} (|V_{i,u}| - |F_i|) > 0, \quad (14.9)$$

$$a_1 = \min_{i,t} \left[d - \sum_{u \neq m(i)} |p_u(V_{i,u}) \cap I_t| \right] > 0. \quad (14.10)$$

The second count includes multiplicities of coincident landings. It says that the primary successor mass cannot be confined to any one source class. Set $\varepsilon = a_0/d$, $\eta = a_1/d$, and $\zeta = 1 - \varepsilon\eta \in (0, 1)$.

Theorem 14.2 (Dispersion criterion). *Under (14.8)–(14.10), $r = q$ and*

$$b_n \leq d\zeta^{\lfloor n/2 \rfloor} \leq \frac{d}{\sqrt{\zeta}} (\sqrt{\zeta})^n \quad (n \geq 0). \quad (14.11)$$

Every legal unrefined Borel feedback has $h^(s) = M_1 = \log q$; all finite Borel atom refinements have $h^* \geq \log q$. There is no $w \in (0, \infty)^d$ and $\vartheta < 1$ such that $w^\top T_S \leq \vartheta w^\top$ for every global $q - 1$ label list. A common nonstrict left budget does exist.*

Proof. Write $S_t = U \setminus \{t\}$ and $T^{(t)} = T_{S_t}$. Its column mass is exactly

$$c_i(t) = \sum_j T^{(t)}(j, i) = \begin{cases} 1, & m(i) = t, \\ 1 - |V_{i,t}|/d + |F_i|/d \leq 1 - \varepsilon, & m(i) \neq t. \end{cases}$$

Thus every $T^{(t)}$ has induced l^1 norm at most one. Let $Q_i(j) = d^{-1} \sum_{u \neq m(i)} \mathbf{1}_{p_u(V_{i,u})}(j)$. It has total mass one, and by (14.10) $\sum_{j \notin I_t} Q_i(j) \geq \eta$ for every i, t .

Fix two global lists S_s, S_t . The source- i column mass of $T^{(t)}T^{(s)}$ is $\sum_j c_j(t)T^{(s)}(j, i)$. If $i \notin I_s$, it is at most $1 - \varepsilon \leq 1 - \varepsilon\eta$. If $i \in I_s$, its first column is exactly Q_i , and at least η of that mass lands outside I_t , where the next list loses at least ε . Its mass is again at most ζ . Hence every pair has l^1 norm at most ζ . Grouping arbitrary histories into pairs proves their

l^1 norm is at most $\zeta^{\lfloor n/2 \rfloor}$. Applying the product to the all-one vector and using (14.3) gives (14.11). No restrictions on switching order or on the feedback are used.

For every control u choose i with $m(i) \neq u$. Condition (14.9) gives a $z \in V_{i,u} \setminus F_i$, on whose cylinder only u is legal. This proves $r = q$. The unique primary control on each two-letter state is a clopen selector, legal at every state and at each of its future successors. The capacity theorem now proves the entropy statements.

If a strict common left budget existed, take a coordinate i with minimum w_i and select the already assumed inequality with $t = m(i)$. That column is Q_i , of mass one, and

$$(w^\top T^{(m(i))})_i \geq w_i > \vartheta w_i,$$

a contradiction. This checks an inequality that was assumed for every global list; it does not alter lists by source in an actual product. Finally $\mathbf{1}^\top T^{(t)} \leq \mathbf{1}^\top$ gives the nonstrict budget, with strict contraction after two steps. $\square$

An explicit family is obtained with $q \geq 2$, $m \geq 2$, and $d = (q - 1)m$. Partition tails into $q - 1$ blocks of size m . For each source, biject its $q - 1$ primary labels with these blocks, and let every p_u preserve each tail block. Choose arbitrary F_i of size at most $m - 1$, and any partition into q nonempty source classes whenever such a partition exists. The primary landing mass is uniform, so $a_0 \geq 1$ and $a_1 = d - \max_t |I_t| > 0$. This supplies arbitrary overlapping choices and nontrivial permutations, rather than one computed product.

For a small witness use $D = \{0, 1, 2, 3\}$, $U = \{0, 1, 2\}$, $I_0 = \{0, 1\}$, $I_1 = \{2\}$, $I_2 = \{3\}$, and $F_i = \{0\}$ for every source. At sources 0, 1 the primary labels 1, 2 use tails $\{0, 1\}$, $\{2, 3\}$; at source 2 the labels 0, 2 use those tails; at source 3 the labels 0, 1 do so. Let p_0 be the identity, $p_1 = (0\ 1)$, $p_2 = (2\ 3)$. The densities at target first letters are

$$(1, 1/2, 1/4), \quad (1/2, 3/4, 0), \quad (0, 1/4, 3/4), \quad (0, 1/4, 3/4).$$

Thus $\Theta_3 = 3/2$, but $\varepsilon = 1/4$, $\eta = 1/2$ give $b_n \leq 4(7/8)^{\lfloor n/2 \rfloor}$ and $M_1 = \log 3$. On $[00]$ controls 0, 1 have different successors. The preceding minimum-coordinate proof excludes all strict left budgets, not just a numerically tested weight. This example does not inherit the all-reference-probability exclusion of Theorem 14.1; its proved exclusion concerns the specified matrix representation.

15. GAIN COLUMNS COMPENSATED BY UNAVOIDABLE CROSS-LAYER LOSS

The preceding dispersion theorem retains a common nonstrict column budget. We now describe a class in which every source has a gain list, no positive common nonstrict left budget exists, and all global list histories nevertheless contract. The condition is on original safety sets and their cardinalities, not on a computed product norm.

15.1. Original safety data and the count condition. Let $q = |U| \geq 3$ and partition $D = \bigsqcup_{t \in U} L_t$ into nonempty source classes. Write $d = |D|$, $\kappa(i) = t$ on L_t , and $M = \min_t |L_t|$. For every source i choose a spare label $m(i) \neq \kappa(i)$. For each $u \in A_i := U \setminus \{\kappa(i), m(i)\}$ choose mutually disjoint tail sets $H_{i,u}$, all of size $a \geq 1$. Put

$$H_i = \bigsqcup_{u \in A_i} H_{i,u}, \quad b = (q - 2)a, \quad B_i = D \setminus H_i.$$

Also choose an arbitrary spare tail set F_i of size f . It may overlap any of the primary tail sets. Define the original allowed relation by

$$\begin{aligned} R_u = & \{(i, z) : u = \kappa(i), z \in B_i\} \\ & \cup \{(i, z) : u \in A_i, z \in H_{i,u}\} \\ & \cup \{(i, z) : u = m(i), z \in F_i\}, \quad K_u = [R_u]. \end{aligned} \tag{15.1}$$

Every two-letter state has one primary control and possibly a spare. The permutations p_u may be arbitrary. Use $g_u = P_u \sigma$ with the same continuous cemetery extension and period one. Require

$$M > b, \quad a < f, \quad b + f < d, \quad (15.2)$$

and

$$2d(f - a) + (f - a)^2 < [d - (q - 1)a][M - (q - 2)a]. \quad (15.3)$$

These conditions contain only cardinalities of the original source classes, small primary tails, and spare tails. The theorem applies to every placement of these tails and every permutation, not only to matrices previously known to be stable.

Set

$$\begin{aligned} \gamma &= (f - a)/d, & C &= 1 + \gamma, & \varepsilon &= 1 - (b + f)/d, & \eta &= (M - b)/d, \\ \zeta &= \max\{C(1 - \varepsilon), C^2 - (\varepsilon + \gamma)\eta\}, & \theta &= \sqrt{\zeta}, & H &= dC/\sqrt{\zeta}. \end{aligned} \quad (15.4)$$

Theorem 15.1 (Gain-loss safety criterion). *For every system satisfying (15.1)–(15.3), $r = q$, $0 < \zeta < 1$, and the complete inverse capacities satisfy*

$$b_n \leq dC^{n \bmod 2} \zeta^{\lfloor n/2 \rfloor} \leq H\theta^n \quad (n \geq 0). \quad (15.5)$$

Every source has an original $q - 1$ label list whose column mass is $C > 1$. There is no $w \in (0, \infty)^d$ satisfying $w^\top T_S \leq w^\top$ for all original $q - 1$ label lists. Nevertheless every unrefined legal Borel feedback has

$$h^*(s) = M_1 = \log q = \log r, \quad (15.6)$$

and every finite Borel atom refinement has $h^ \geq \log q$.*

15.2. Complete estimate for every pair of global lists.

Proof. Write $S_t = U \setminus \{t\}$ and $T^{(t)} = T_{S_t}$. The exact source-column masses, counted with edge multiplicities, are

$$c_i(t) = \sum_j T^{(t)}(j, i) = \begin{cases} (b + f)/d = 1 - \varepsilon, & t = \kappa(i), \\ 1 + (f - a)/d = C, & t \in A_i, \\ 1, & t = m(i). \end{cases} \quad (15.7)$$

Indeed, omitting the large primary control leaves all small tails and the spare; omitting a small primary tail removes a edges and adds f spare edges to the primary total d ; omitting the spare leaves exactly the primary partition. Distinct controls landing on the same target are still distinct counted edges.

Fix two whole-space lists S_s, S_t . The source- i column mass of their product is

$$\sum_j (T^{(t)} T^{(s)})(j, i) = \sum_j c_j(t) T^{(s)}(j, i) = c_i(s) + \sum_j [c_j(t) - 1] T^{(s)}(j, i). \quad (15.8)$$

If $s = \kappa(i)$, then $c_i(s) = 1 - \varepsilon$ and every next-column mass is at most C . The product column is at most $C(1 - \varepsilon)$.

If $s \neq \kappa(i)$, the whole large primary tail is retained. Its permutation image misses just b letters. Hence, for every next loss class L_t ,

$$\sum_{j \in L_t} T^{(s)}(j, i) \geq \frac{|p_{\kappa(i)}(B_i) \cap L_t|}{d} \geq \frac{|L_t| - b}{d} \geq \eta. \quad (15.9)$$

Small and spare tails can only increase this mass. On L_t the next list has mass $1 - \varepsilon$, whereas elsewhere its mass is at most C . Thus

$$\begin{aligned} \sum_j c_j(t) T^{(s)}(j, i) &\leq C c_i(s) - [C - (1 - \varepsilon)] \sum_{j \in L_t} T^{(s)}(j, i) \\ &\leq C^2 - (\varepsilon + \gamma)\eta. \end{aligned} \quad (15.10)$$

This includes both the current gain and the worst next gain, rather than subtracting a loss while omitting the cost of spare labels. The lists s, t remain fixed over the state space throughout.

Dividing (15.3) by d^2 yields

$$2\gamma + \gamma^2 < (\varepsilon + \gamma)\eta, \quad \varepsilon + \gamma = [d - (q - 1)a]/d > 0.$$

Since $M \leq d/q$ and $M > b$, $0 < \eta < 1$. The last inequality therefore implies $\varepsilon > \gamma + \gamma^2 > \gamma$. It follows that $C(1 - \varepsilon) < 1$; it also gives $C^2 - (\varepsilon + \gamma)\eta < 1$ directly. Both quantities are positive: the first by $b + f > 0$, the second because $\varepsilon + \gamma < 1$, $\eta < 1$, and $C^2 > 1$. Thus $0 < \zeta < 1$ and every list pair has induced l^1 norm at most ζ .

For every history, group consecutive factors into pairs. The possible odd remaining factor has norm at most C , not one. Consequently

$$\|T^{(t_{n-1})} \dots T^{(t_0)}\|_1 \leq C^{n \bmod 2} \zeta^{\lfloor n/2 \rfloor}.$$

Equation (14.3) and $\|\mathbf{1}\|_1 = d$ prove (15.5), including $b_0 = 1 \leq d$.

To exclude *all* common positive nonstrict left budgets, suppose one existed and take a coordinate i of smallest w_i . Since $q \geq 3$, choose $t \in A_i$. The already assumed inequality for this global list would give

$$w_i \geq (w^\top T^{(t)})_i = \sum_j w_j T^{(t)}(j, i) \geq w_i c_i(t) = C w_i > w_i,$$

a contradiction. This is a complete positivity certificate and uses no floating-point feasibility test.

All states have a primary legal control. For each u choose $i \in L_u$. As $|B_i| = d - b > f$, a $z \in B_i \setminus F_i$ exists. Only u is legal on $[iz]$. Thus every control is needed in a subcover, giving $r = q$. The actual primary selector is

$$s_0(x) = \begin{cases} \kappa(x_0), & x_1 \in B_{x_0}, \\ u, & x_1 \in H_{x_0, u}, \quad u \in A_{x_0}. \end{cases} \quad (15.11)$$

Its primary domains partition all states, so it is single-valued and clopen. At every successor it again uses the same legal state function; all infinite trajectories, including periodic and constant states, stay safe. The capacity theorem gives the lower bound for every Borel rule, not only this selector. With only q physical labels every unrefined rule has at most q^m patterns on m positions. This proves (15.6) and the refinement statement. $\square$

The unavailable common budget is the *original first-letter, one-step, positive left* budget. Two-step products do have a strict common budget. No claim excludes higher-dimensional lifts, right budgets, arbitrary changes of reference measure, or path-dependent Lyapunov functions. In particular no conclusion about L_1 follows from (15.6).

15.3. A family for arbitrary control counts. For any $q \geq 3$ and integer $m \geq q + 1$, take $d = qm$, all $|L_t| = m$, $a = 1$, and $f = 2$. Each source may assign its spare label arbitrarily, choose its $q - 2$ distinct small-tail letters arbitrarily, choose any two spare-tail letters, and use arbitrary permutations. Conditions (15.2) hold. The right side minus the left side of (15.3) is

$$P(m) = qm^2 - (q^2 + q - 1)m + q^2 - 3q + 1.$$

Here $P(q + 1) = q^2 - 2q + 2 > 0$ and $P(m + 1) - P(m) = 2qm - q^2 + 1 > 0$ for $m \geq q + 1$. Thus the entire family satisfies the theorem, with constants depending only on these counts. The condition allows arbitrary safe-edge arrangements and not just repetitions of a small matrix. It does not assert that every member lacks all protected symbolic subsystems; no such subsystem is a hypothesis of the proof.

15.4. A complete three-control gain witness. Let $D = \{0, \dots, 23\}$ and $U = \{0, 1, 2\}$. Take $L_0 = \{0, \dots, 7\}$, $L_1 = \{8, \dots, 15\}$, $L_2 = \{16, \dots, 23\}$. For $i \in L_t$ the large label is t , the small label is $t + 1$, and the spare is $t + 2$ modulo three. The complete tails are specified in the following table; every source in a row has that row's tails.

Source class	Large label	Small tail H_t	Spare tail F_t
L_0	0	$\{0, 1, 2, 3\}$	$\{4, 5, 6, 7, 8\}$
L_1	1	$\{8, 9, 10, 11\}$	$\{0, 1, 2, 3, 4\}$
L_2	2	$\{8, 9, 10, 11\}$	$\{0, 1, 2, 3, 4\}$

The large tail is $D \setminus H_t$. Let p_0 be the identity, $p_1 = (0\ 1)$, $p_2 = (16\ 17)$, fixing all other letters. These permutations preserve each of their actually used tail sets. Thus the exact matrix description, with subscripts modulo three, is

$$T_u(j, i) = \frac{1}{24} [\mathbf{1}_{L_u}(i) \mathbf{1}_{D \setminus H_u}(j) + \mathbf{1}_{L_{u-1}}(i) \mathbf{1}_{H_{u-1}}(j) + \mathbf{1}_{L_{u-2}}(i) \mathbf{1}_{F_{u-2}}(j)].$$

It computes capacity without replacing the actual permutations by common successor maps. Its target densities are

Target letter	ρ_0	ρ_1	ρ_2	Largest two sum
0, 1, 2, 3	1/3	1	1/3	4/3
4	2/3	2/3	2/3	4/3
5, 6, 7	1/3	1/3	2/3	1
8	2/3	0	2/3	4/3
9, 10, 11	2/3	0	1/3	1
12, ..., 23	1/3	1/3	1/3	2/3

In particular $\Theta_3 = 4/3$. Here $d = 24$, $M = 8$, $a = b = 4$, $f = 5$, and the strict count margin is $64 - 49 = 15$. The constants are

$$\varepsilon = 5/8, \quad \gamma = 1/24, \quad C = 25/24, \quad \eta = 1/6,$$

$$C(1 - \varepsilon) = 25/64, \quad C^2 - (\varepsilon + \gamma)\eta = 187/192.$$

The full-history estimate is therefore

$$b_n \leq 24(25/24)^{n \bmod 2} (187/192)^{\lfloor n/2 \rfloor}. \quad (15.12)$$

Source columns have masses $3/8$, $25/24$, or 1 , depending on which of their large, small, or spare labels is omitted. The cylinders $[0, 12]$, $[8, 12]$, $[16, 12]$ require controls $0, 1, 2$. On $[16, 0]$ controls $1, 2$ are legal but have successor first letters $1, 0$. Consequently the full Borel class includes choices genuinely changing the future trajectory.

There is no consecutive 000 trajectory with uniquely legal controls at every step. A unique 0 at a source in L_0 has its next first letter in $\{9, \dots, 23\}$; a unique 0 in L_2 has its next first letter in $\{8, \dots, 11\} \subset L_1$. In L_1 , control 0 is only spare and is never unique. Thus from L_0 there can be at most one further unique 0 , and from L_2 none. This excludes a direct consecutively forced three-symbol full language. It does not exclude sparse bridges, other factors, or every protected coding construction.

The gain–loss argument is elementary positive-matrix reasoning, applied here through exact allowed-tail quotas and actual feedback fibers. Shi et al. [11, Definition 2.5 and Theorems 2.6, 2.8, pp. 8–11 of arXiv:2004.01867v2] already allow losses to compensate growth in matrix products. Their positive-diagonal and ordered-window assumptions do not automatically hold here: for example $T^{(1)}(0, 0) = 0$ in this witness. Our proof uses the mass quota (15.9) for every next global list and does not depend on that theorem. The separation concerns a specified common-left template; it is not a new general Lyapunov theory.

16. AFFINE EXAMPLES, COMMON SAMPLING, AND SHARP SINGLE-STEP BOUNDARIES

These examples separate fixed-period maximal-pattern optimization, common sampling, and a fixed arithmetic sampling. Their proofs are included because a matching upper bound must be supplied by a real feedback. Volume estimates, integer digit coding, and protected inverse branches are classical mechanisms, not new general tools.

16.1. An integer affine family with a genuine absorbing feedback. Let $m \geq 2$, $D \geq m$ be integers, $R = D/(m-1)$, $Q = [0, R]$, and controls $u = 0, \dots, D$ act legally by $mx - u$. On $X = [-1, R+1]$ define the continuous ambient maps

$$f_u(x) = \max\{-1, \min\{R+1, mx - u\}\}.$$

For $x \in Q$, this is in Q exactly when $mx - u$ is in Q , and then truncation has no effect. The safety intervals $K_u = [u/m, (u+R)/m]$ cover Q . Fix any period $\tau \geq 1$.

Proposition 16.1. *For this system, over all legal Borel block feedbacks and finite Borel atom refinements,*

$$\min_s h_{(0,1,\dots)}(s) = L_\tau = M_\tau = \log m.$$

There is an actual Borel selector s_τ with $h_A(s_\tau) = h^(s_\tau) = \log m$ for every infinite increasing A , although its absorption time has no uniform bound.*

Proof. For any actual length- n block word, concatenate its controls into $w_0 \cdots w_{n\tau-1}$. Safety gives

$$x_{n\tau} = m^{n\tau}x - \sum_{j=0}^{n\tau-1} w_j m^{n\tau-1-j} \in [0, R].$$

Its initial-state fiber is contained in an interval of length $R/m^{n\tau}$, whether or not that fiber itself is an interval. These containing intervals cover Q , so there are at least $m^{n\tau}$ actual consecutive block words. This gives the full-class lower bound at the common sequence $(0, 1, \dots)$.

Define the one-step rule $v(x) = \min\{D, \lfloor mx \rfloor\}$ and $T(x) = mx - v(x)$. If $x < D/m$, the next state is in $[0, 1)$; if $x \geq D/m$, it is $mx - D \in [0, R]$. The core $[0, 1)$ is invariant and uses only $\Gamma = \{0, \dots, m-1\}$. Both 0 and R are fixed; D/m uses D and goes to zero. For $x < R$, while D is used the state is $R - m^j(R - x)$. Since $R - x > 0$, eventually it is below D/m , then one transitional control $v < D$ sends it into the core. Only R uses D forever. The absorption times are unbounded as x tends to R .

From the current state define $s_\tau(x) = (v(x), v(Tx), \dots, v(T^{\tau-1}x))$. It is Borel, each intermediate step is safe, and its actual endpoint map is T^τ . Put $\mathcal{B} = \Gamma^\tau$, $b = m^\tau$, and $d_0 = (D, \dots, D)$. All its real block names are contained in

$$\bigcup_{j \geq 0} d_0^j \mathcal{C}_\tau \mathcal{B}^{\mathbb{N}_0} \cup \{d_0^\infty\}, \quad |\mathcal{C}_\tau| \leq D \frac{m^\tau - 1}{m - 1} =: C_\tau.$$

The transitional block has some initial D 's, one control below D , and then core controls; this proves the stated count. For any window of n block positions, an unobserved transition leaves just one of $n+1$ cuts between constant initial labels and core labels. A sampled transition has n possible locations and at most C_τ labels. Thus

$$\mathbf{P}_{s_\tau}(F) \leq [n+1 + nC_\tau]m^{\tau n}, \quad |F| = n.$$

There is no factor depending on the last observation time or on the unbounded absorption time.

For the matching lower count prescribe any n core blocks at the positions of F , fill all other one-step digits with zero, and let $x = \sum_{j \geq 0} e_j m^{-j-1} < 1$. There are only finitely many nonzero digits. Every tail is smaller than one, so $\lfloor mT^j x \rfloor = e_j$ and these are real controls of the same rule. Hence $\mathbf{P}_{s_\tau}(F) \geq m^{\tau n}$ for every window. The all-window bounds and the

full-class consecutive lower bound prove the proposition, with minima actually attained. Control-label projection treats arbitrary atom refinements. $\square$

When $m \geq 3$ and $D = m$, the minimum one-step safe subcover has $r = m + 1$, although $L_1 = M_1 = \log m$. Indeed, m intervals of length R/m could cover an interval of length R only by abutting without positive-length overlap. Their left endpoints are multiples of $1/m$, so this would force their consecutive control differences to equal the noninteger $R = m/(m - 1)$. All $m + 1$ intervals are therefore needed. There is a common state factor $x \mapsto e^{2\pi i x}$, with induced map $z \mapsto z^m$; it is preserved by every legal branch. The result cannot be presented as eliminating common-factor explanations.

16.2. A protected bridge giving a predetermined common sampling. On $Q = [0, 1]$, at period one, use

$$g_a(x) = \frac{5}{4}x, \quad g_b(x) = \frac{20}{3}x - \frac{16}{3}, \quad g_c(x) = 20x - 19, \quad g_d(x) = 100x.$$

Their safety domains are $[0, 4/5]$, $[4/5, 19/20]$, $[19/20, 1]$, and $[0, 1/100]$, respectively. Continuous ambient extensions are obtained by truncating these affine maps to $[-1, 2]$, with no change on legal paths. The minimum subcover is three and the largest two inverse slopes sum to $4/5 + 3/20 = 19/20 < 1$; hence $M_1 = \log 3$ by the scalar special case of the capacity theorem.

Proposition 16.2. *For $A_2 = (0, 2, 4, \dots)$, every legal Borel feedback and every finite Borel atom refinement has at least 3^n actual patterns on its first n observations. Consequently*

$$\min_s h_{A_2}(s) = L_1 = M_1 = \log 3.$$

Proof. The full inverse branches are $\psi_a(y) = 4y/5$, $\psi_b(y) = 4/5 + 3y/20$, $\psi_c(y) = 19/20 + y/20$. Let $J = [1/2, 999/1000]$ and $\theta_u = \psi_u \psi_b$. They satisfy

$$\theta_a(y) = 16/25 + 3y/25, \quad \theta_b(y) = 23/25 + 9y/400, \quad \theta_c(y) = 99/100 + 3y/400.$$

Their complete images and the intermediate image are

image	exact endpoints	only legal control
$\psi_b(J)$	$[7/8, 18997/20000]$	b
$\theta_a(J)$	$[7/10, 18997/25000]$	a
$\theta_b(J)$	$[149/160, 376991/400000]$	b
$\theta_c(J)$	$[159/160, 398997/400000]$	c

Each $\theta_u(J)$ lies in J . Every displayed closed interval is strictly inside the appropriate uniquely legal region; in particular $\theta_a(J)$ avoids the extra control d .

Given any $u_0 \cdots u_{n-1} \in \{a, b, c\}^n$, set $x_{2n} = 1/2$ and recursively $x_{2j+1} = \psi_b(x_{2j+2})$, $x_{2j} = \theta_{u_j}(x_{2j+2})$. The table shows that at every even state only u_j is legal and at every odd state only b is legal. Thus *every* feedback must execute the actual period-one prefix $u_0, b, \dots, u_{n-1}, b$ from the same constructed initial state. The endpoint remains in Q and the globally legal rule continues safely forever. There is no added phase or period-two control. All three-symbol sampled words are therefore actual patterns. Refinement only increases their count.

The Borel selector using a below $4/5$, b from $4/5$ to $19/20$, and c at and above $19/20$, is legal at all endpoints and uses only three labels. Its upper count is 3^n on every window. This matches the lower bound, including attainment, and $L_1 \leq M_1$ completes the equalities. The normalization is $n\tau = n$, not $2n$. $\square$

The closed contracting branches $\theta_a, \theta_b, \theta_c$ have disjoint first-level images in J . Their nested images give a compact set homeomorphic to the three-symbol full shift, and every T_s^2 on that set is the same shift under this coding. To verify this, a word's image has diameter at most $(3/25)^n \text{diam } J$; nested images intersect in one point, different first differing letters give disjoint images, and the inverse identities used above give the shift relation. The union of

this set with its ψ_b image is protected under every legal feedback. These sets can overlap: the all- b state $16/17$ belongs to both and uniquely uses b . Hence the proof does not introduce an implicit disjoint phase register. This is precisely a protected-subsystem mechanism, even if the rest of the state space has feedback-dependent successors.

16.3. A critical boundary of the single-step density condition. For $3/2 < \lambda \leq 2$, on $Q = [0, 1]$ use

$$g_a(x) = \lambda x, \quad g_b(x) = \lambda x - (\lambda - 1)/2, \quad g_c(x) = 3x - 2.$$

The safety intervals are

$$K_a = [0, 1/\lambda], \quad K_b = [(\lambda - 1)/(2\lambda), (\lambda + 1)/(2\lambda)], \quad K_c = [2/3, 1].$$

They cover Q and $r = 3$. Both a and c are forced into every subcover by endpoints $0, 1$. Since $1/\lambda < 2/3$, they leave an open gap, so b is also needed. The inverse densities are constants $1/\lambda, 1/\lambda, 1/3$; therefore $\Theta_2 = 1/\lambda < 1$ and $\Theta_3 = 2/\lambda$.

Proposition 16.3. *For every $3/2 < \lambda \leq 2$, this full Borel class has $M_1 = \log 2 < \log 3$. Thus the condition $\Theta_3 < 1$ cannot be uniformly replaced by $\Theta_3 \leq 1$ or by $\Theta_3 < 1 + \eta$ for any $\eta > 0$, while retaining the conclusion $M_1 = \log 3$.*

Proof. The $k = 2$ single-step criterion gives every feedback $h^* \geq \log 2$. Put $t = (\lambda - 1)/(2\lambda)$ and define

$$s_\lambda(x) = \begin{cases} a, & 0 \leq x < t, \\ b, & t \leq x < 2/3, \\ c, & 2/3 \leq x \leq 1. \end{cases}$$

It is Borel and legal: $t < 1/\lambda$ and $(\lambda + 1)/(2\lambda) \geq 3/4 > 2/3$. For $x < t$, $g_a(x) < (\lambda - 1)/2 \leq 1/2$. For $t \leq x \leq 1/2$, $g_b(x) = \lambda x - (\lambda - 1)/2 \in [0, 1/2]$. Hence $J = [0, 1/2]$ is invariant. Before entering J only b, c occur; once in J only a, b occur. This remains true for states which never enter and at all thresholds, including the fixed points $1/2$ and 1 .

At any n selected times a real name has some cut $0 \leq j \leq n$ between the two groups, each of size two. Thus its pattern count is at most $(n + 1)2^n$ for *every* window. This actual feedback gives $h^* \leq \log 2$ and completes the equality. At $\lambda = 2$, $\Theta_3 = 1$; as λ increases to 2 from below, Θ_3 decreases to 1 from above. These values give both claimed failures of uniform relaxation. The example does not make $\Theta_3 < 1$ a necessary condition for every system, and does not determine L_1 for this family. $\square$

16.4. The noninteger four-branch system and what remains open. Let $Q = [0, 6]$ and use, at period one,

$$g_{A0}(x) = \frac{3}{2}x, \quad g_{A1}(x) = \frac{3}{2}x - 3, \quad g_{B0}(x) = \frac{5}{3}x, \quad g_{B1}(x) = \frac{5}{3}x - 4.$$

The original safety intervals are $[0, 4]$, $[2, 6]$, $[0, 18/5]$, $[12/5, 6]$. Truncation to the ambient interval $[-1, 7]$ retains exactly these safe transitions. All legal Borel feedbacks, arbitrary mixtures, and finite atom refinements are allowed. Two A domains cover Q and no single one does, so $r = 2$. The largest scalar inverse capacity is $2/3 < 1$. Thus $M_1 = \log 2$. Every actual length- N control prefix has its fiber inside an interval of length at most $6(3/2)^{-N}$; these intervals cover Q . Its number $D_N(s)$ is therefore at least $(3/2)^N$. This gives

$$\log(3/2) \leq L_1 \leq \log 2. \tag{16.1}$$

For $A_2 = (0, 2, \dots)$ let $\ell_2 = \inf_s h_{A_2}(s)$, and let ℓ_2^A restrict to all two- A Borel feedbacks, without restricting their partitions. The unrefined two- A rules have the exact description

$$s_E(x) = A0 \text{ on } [0, 2) \cup E, \quad s_E(x) = A1 \text{ elsewhere,} \quad E \subset [2, 4] \text{ Borel.}$$

Indeed the two overlap domains are exactly $[2, 4]$. For any feedback, an actual $2n$ prefix is determined by its even and odd sampled patterns. Odd patterns occur among the even patterns from states $T_s x \in Q$. Hence $D_{2n}(s) \leq N_n(s)^2$ and the volume lower bound gives $N_n(s) \geq (3/2)^n$. Consequently $\ell_2 \geq \log(3/2)$.

Proposition 16.4 (Density-one samplings). *If $A = (a_i)$ satisfies $a_n/n \rightarrow 1$, then $\inf_s h_A(s) = \log(3/2)$ over the full class. The greedy feedback $s_g = s_\emptyset$ is an actual minimizer for every such A .*

Proof. Set $\beta = 3/2$. On the invariant core $[0, 3)$, scale $x = 3t$. The greedy dynamics becomes $B(t) = \beta t - \lfloor \beta t \rfloor$ on $[0, 1)$. Every nonempty length- n digit cylinder is a left-closed right-open interval whose B^n image is $[0, l)$ for some $0 < l \leq 1$. This is proved by splitting the current image at $1/\beta$: it has a single child if $\beta l \leq 1$, and otherwise a full-image child and a nonfull-image child. Equality creates no extra singleton at the excluded right endpoint.

Let F_n be the number of full-image cylinders, with $F_0 = 1$. The sum of image lengths is β^n , since each source length is l/β^n and the sources partition $[0, 1)$. Thus $F_n \leq \beta^n$. Each nonfull cylinder has a last full-image ancestor, and each ancestor has at most one descendant at a given later level without another full-image step. The total core count C_n consequently satisfies

$$C_n \leq \sum_{j=0}^n F_j \leq 3\beta^n.$$

For $x \in [3, 6)$ the real greedy control remains $A1$ until entering the core; while waiting its state is $6 - \beta^j(6 - x)$, so entry is finite. State 6 uses $A1$ forever. Therefore the full-space count satisfies

$$D_n(s_g) \leq 1 + \sum_{j=1}^n C_j \leq 10\beta^n.$$

This includes all thresholds and the exceptional endpoint.

Let $N = a_n + 1$, so $N - n = o(n)$. Each sampled word has at most 4^{N-n} completions to a consecutive word, giving

$$P_s(\{a_1, \dots, a_n\}) \geq \frac{D_N(s)}{4^{N-n}} \geq \frac{\beta^N}{4^{N-n}}.$$

For the greedy rule the same count is at most $D_N(s_g) \leq 10\beta^N$. After division by n the two rates tend to $\log \beta$. Refinements preserve the lower bound and the unrefined rule gives the matching upper bound. $\square$

Let $\rho > 1$ be the unique solution

$$\rho^6 = \rho^5 + \rho^4 + \rho + 1, \quad \log \rho = 0.55600938749\dots \quad (16.2)$$

Appendix A proves the exact real-feedback value $h_{A_2}(s_g) = \log \rho$, with full upper and lower counts. Thus the currently established intervals are

$$\log(3/2) \leq \ell_2 \leq \ell_2^A \leq \log \rho. \quad (16.3)$$

Neither greedy optimality in the full class nor greedy optimality in the two- A class is established. These fixed- A_2 questions are distinct from (16.1). A lower entropy abstract open-loop language is constructed in Appendix C; its full Borel feedback implementation remains unresolved.

17. EXPANSION, ABSENCE OF CONTINUOUS COMMON FACTORS, AND PHASE OBSTRUCTIONS

17.1. Expansion extensions of the integer family. The next statement retains a matching real feedback while allowing other feedbacks to change the dynamics inside its absorbing core. It is a geometric extension of Proposition 16.1, rather than a new digit-coding theorem.

Proposition 17.1. *Start with the integer family of Proposition 16.1. Add finitely many continuous branches $g_v : Q \rightarrow \mathbb{R}$ satisfying*

$$|g_v(x) - g_v(y)| \geq m|x - y| \quad (x, y \in Q). \quad (17.1)$$

A branch is legal precisely when its value is in Q . For every fixed period τ , over the entire enlarged Borel class,

$$\min_s h_{(0,1,\dots)}(s) = L_\tau = M_\tau = \log m.$$

If among the added branches is $g_e(x) = kmx$, with an integer $k \geq 2$ and $R \geq k$, there is no nonconstant continuous map $\pi : Q \rightarrow Z$, to any compact metric space, and no map $H : Z \rightarrow Z$ such that $\pi(T_s x) = H(\pi(x))$ for all legal period- τ feedbacks and all states. Continuity of H is not required.

Proof. Use the continuous ambient extension $f_v = c_X \circ g_v \circ c_Q$ on $X = [-1, R+1]$, where c denotes interval truncation. A value outside Q remains outside Q after truncation to X . The original branches still cover Q . Two states in the same real length- n block fiber follow the same $n\tau$ legal controls, so (17.1) gives $m^{n\tau}|x-y| \leq R$. Their fiber, however fragmented, has diameter at most $Rm^{-n\tau}$. Its containing interval, together with the other containing intervals, covers Q . There must be at least $m^{n\tau}$ real consecutive words. The original actual feedback s_τ is still available and retains every all-window count proved in Proposition 16.1. These matching bounds prove the equalities and treat refinements by projection.

Suppose the asserted common continuous factor exists. Any legal block at a given state can be chosen there by changing a globally safe Borel selector at that singleton. Thus all legal blocks at that state have the same projected endpoint. At $x = y/m^\tau$, for $1 \leq y \leq R$, the legal blocks 0^τ and $0^{\tau-1}1$ end at $y, y-1$. All intermediate states are in Q . Hence $\pi(y) = \pi(y-1)$ throughout $[1, R]$. The restriction to $[0, 1]$ induces a continuous $F : \mathbb{R}/\mathbb{Z} \rightarrow Z$, and repeated subtraction of integers gives $\pi(y) = F(y \bmod 1)$ on Q . At the same initial state the blocks 0^τ and $0^{\tau-1}e$ end at y, ky , provided $0 \leq y \leq R/k$. Since $R \geq k$, this implies $F(t) = F(kt \bmod 1)$ for all t on the circle. Iterating gives $F(j/k^n) = F(0)$ for every $0 \leq j < k^n$. Their union is dense, so continuity makes F , and then π , constant, a contradiction. $\square$

For example, $m = 2, D = 3, Q = [0, 3]$ admits both $g_e = 4x$ and $g_v = 2x + x^2$ as additional branches. At $x = 1/8$ the legal successors under $0, e, v$ are $1/4, 1/2, 17/64$, all in the original core. A rule choosing e on $(0, 1/4)$ and otherwise the original greedy control is globally Borel and safe. These are genuine feedback-dependent core dynamics, and the proposition excludes every nonconstant continuous common factor. It does not exclude Borel factors: for integer affine branches and e , rationality of the state is a nonconstant Borel common factor.

Condition (17.1) cannot be replaced by expansion on each monotonicity interval. In the $m = 2, D = 3$ system add $g_b(x) = 2x$ on $[0, 3/2]$ and $g_b(x) = 6 - 2x$ on $[3/2, 3]$. Both slopes have absolute value two, but this single branch maps all of Q into Q . The constant real block b^τ has one instruction name; both optimized rates are zero. Folding hides the spatial branching from the control-label alphabet.

17.2. An alternating open-loop language with no Borel realization. Let $R > 0$, $1 < a, b < 2$, and $\log a / \log b \notin \mathbb{Q}$. On $[0, R]$ take four legal affine branches

$$g_{Ai}(x) = ax - i(a-1)R, \quad g_{Bi}(x) = bx - i(b-1)R \quad (i = 0, 1).$$

Each type's two full inverse intervals cover Q . Let Λ_{alt} be the two-phase language whose A, B types alternate strictly and whose digits are otherwise arbitrary. Let

$$\hat{\Lambda} = \Lambda_{\text{alt}} \cup \{wc^\infty : w \text{ is a finite alternating-type word, } c \in U\}.$$

The empty prefix is allowed. This compact shift-invariant language includes all constant endpoint tails and their finite alternating prehistories. The inverse contractions $\psi_{Ci}(y) = (y + i(\lambda_C - 1)R)/\lambda_C$, with $\lambda_A = a, \lambda_B = b$, define a continuous coding $\pi(d) = \lim_n \psi_{d_0} \cdots \psi_{d_{n-1}}(0)$. Every state has a coding in Λ_{alt} : in each prescribed type choose a safe digit and repeat. The contraction error is at most $R \min(a, b)^{-n}$. This proves full open-loop coverage, not feedback consistency.

Proposition 17.2. *No globally safe Borel period-one feedback has all its real names in $\widehat{\Lambda}$. This remains true with arbitrary Borel mixtures and finite atom refinements. Nevertheless for every n -position window F ,*

$$2^{n+1} \leq P_{\widehat{\Lambda}}(F) \leq 8(n+1)2^n.$$

Proof. The alternating part has two distinct type patterns and 2^n independent digits. For a constant-tail member there are at most $n+1$ cuts on F , two prefix phases, 2^j prefix digits, and four constant tails. Summing by cuts bounds the count by $8(n+1)2^n$ (and also covers the infinite alternating case).

For nonrealizability, let C be the union of all finite affine preimages of $\{0, R\}$. This set is countable and forward invariant in the sense that a legal successor of a state outside C remains outside C . A constant safe tail under any one branch forces its tail state to be the corresponding endpoint, by expansion about 0 or R . Thus any hypothetical feedback has strictly alternating types forever on $Q \setminus C$. Choose

$$0 < \eta < \min\{R/a, R/b, (a-1)R/a, (b-1)R/b\}.$$

On $(0, \eta)$ only zero-digit controls are legal. For $0 < x < \eta/(ab)$ outside C , the true first two steps are x, ax, abx if the first type is A , and x, bx, abx otherwise. The third type is the first type. Put $p = \log a, q = \log b, h = p + q$ and encode the feedback type at $\eta e^t, t < 0$, by a Borel $c(t) \in \{0, 1\}$, with A represented by one. Outside a countable exceptional set,

$$c(t+h) = c(t), \quad c(t) = 1 \Rightarrow c(t+p) = 0, \quad c(t) = 0 \Rightarrow c(t+q) = 1 \quad (t < -h).$$

Translating each negative interval of length h to $[-h, 0)$ gives a measurable circle function, with all countably many translated exceptions still null. Apply the last two implications at $t = r - h, -h \leq r < 0$. For its set E of type one, circle rotations satisfy $R_p E \subset E^c$ and $R_q E^c \subset E$ modulo null sets. Since $q = -p$ modulo h , these imply $R_p E = E^c$, measure $1/2$, and $R_{2p} E = E$ modulo null sets. The number $2p/h$ is irrational. For $G = 2\mathbf{1}_E - 1$, invariance under that rotation and Fourier coefficients give $\widehat{G}(j) = e^{2\pi i j(2p/h)} \widehat{G}(j)$. Every nonzero coefficient vanishes; the mean vanishes as well. Completeness of the trigonometric system gives $G = 0$ almost everywhere, contradicting $|G| = 1$. This proves impossibility. Refinements project to the same physical control rule and cannot avoid the obstruction. $\square$

The noninteger system in Section 16.4 has $a = 3/2, b = 5/3, R = 6$. Its log ratio is irrational by unique prime factorization: $(3/2)^j = (5/3)^k$ would imply $3^{j+k} = 2^j 5^k$. None of its nonempty finite affine composites has an integer slope. The proposition is therefore a substantive implementation boundary for this system; it is not a proof of a strict optimized entropy gap.

APPENDIX A. EXACT EVEN-SAMPLING ENTROPY OF THE REAL GREEDY FEEDBACK

The system and s_g are those of Section 16.4. We prove the exact value, including all initial states, rather than identifying an auxiliary language with a feedback without proof. On $C_0 = [0, 3)$ the actual map $R = T_{s_g}^2$ is

$$R(x) = \begin{cases} 9x/4, & 0 \leq x < 4/3, \\ 9x/4 - 3, & 4/3 \leq x < 2, \\ 9x/4 - 9/2, & 2 \leq x < 3. \end{cases} \quad (\text{A.1})$$

The middle state really uses $A0$ or $A1$ according to the original threshold, so these are actual two-step trajectories of a period-one rule. Let $0, 1$ denote its even labels $A0, A1$.

If a core even name begins 11 , its second state is in $[2, 9/4)$, its third in $[0, 9/16)$, and the next two images are in $[0, 81/64)$ and $[0, 729/256)$. The first of these is below $4/3$. If the last image is at least 2 , its next image is smaller than $1953/1024 < 2$. Consequently every core even name avoids

$$111, \quad 1101, \quad 110011. \quad (\text{A.2})$$

All interval upper endpoints in these calculations are excluded by the core or its actual branch convention; $4/3$ and 2 both map to zero under R .

Every word avoiding (A.2) has a unique left-to-right parsing into the prefix-code blocks $a = 0$, $b = 10$, $c = 1100$, with no adjacent cc , followed by one of $\emptyset, 1, 11, 110$. The first two exclusions bound one-runs and force two zeros after a full double-one run; the third prevents two complete c blocks. Let H_m count complete block strings of bit length m with no cc , with $H_m = 0$ for $m < 0$. If X and Z count strings ending in a/b and in c , respectively, their formal generating series satisfy

$$X = (z + z^2)H, \quad Z = z^4(1 + X), \quad H = 1 + X + Z.$$

Thus

$$H(z) = \frac{1 + z^4}{1 - z - z^2 - z^5 - z^6}, \quad H_m = H_{m-1} + H_{m-2} + H_{m-5} + H_{m-6} \quad (m \geq 5), \quad (\text{A.3})$$

with $H_0, \dots, H_4 = 1, 1, 2, 3, 6$ and $H_5 = 10$. The equation for ρ in (16.2) is equivalent to $\rho^{-1} + \rho^{-2} + \rho^{-5} + \rho^{-6} = 1$, whose left side is strictly decreasing. Its unique root obeys $5/3 < \rho < 7/4$. The initial counts are at most ρ^m , and the recurrence proves $H_m \leq \rho^m$ at all lengths. The incomplete-tail count is at most $H_m + H_{m-1} + H_{m-2} + H_{m-3} \leq D\rho^m$, where $D = 1 + \rho^{-1} + \rho^{-2} + \rho^{-3}$.

For a real full-space even word, either all labels are 1, or its first 0 occurs at some position j . At that moment the state is below 2 and lies in the core; its remaining word is governed by (A.2). This includes arbitrarily long waiting outside the core and the fixed endpoint 6. Therefore

$$N_n(s_g) \leq \sum_{m=0}^n D\rho^m \leq \frac{D\rho}{\rho-1}\rho^n. \quad (\text{A.4})$$

This is a real feedback upper bound for every length, not an equality of its language with the larger forbidden-block language.

Inverse paths furnishing the matching lower bound. Put $\eta = 4/9$, $L = 162/65$, $d_0 = 729/256$, and define

$$P_0 = [0, 16/27] \cup [72/65, 4/3), \quad P_1 = [L, 3), \\ C = P_0 \cup P_1, \quad D_0 = P_0 \cup [2, 550/243] \cup [L, d_0).$$

These are subsets of the original interval used only to prove realizability, not feedback states or phase variables. The following inverse paths realize the indicated even blocks:

Even block	Target y	Source state	Real one-step controls
$a = 0$	$y \in C$	ηy	00
$b = 10$	$y \in P_1$	$2 + \eta^2 y$	1000
$b = 10$	$y \in C$	$2 + \eta^2 y$	1000
$b = 10$	$y \in P_0$	$70/27 + \eta^2 y$	1001
$c = 1100$	$y \in D_0$	$26/9 + \eta^4 y$	10100000

The duplicate left b row has source in $C \cap D_0$ when $y \in P_1$, and source only required in D_0 for general $y \in C$.

Here are all intermediate checks. For a , the source ηy is in P_0 , the intermediate state is $2y/3 < 2$, and the endpoint is y ; both controls are zero. For left b , the states after its source are $8y/27, \eta y, 2y/3, y$. If $y \in P_0$, the source is in $[2, 550/243]$; if $y \in P_1$, it is in $[L, 70/27]$. Each source is above 2 and outside $I = [12/5, L)$, while the three subsequent control states are below 2. This gives 1000 with the stated source memberships.

For right b the states are $8/9 + 8y/27, 4/3 + \eta y, 2 + 2y/3, y$. The source belongs to $[70/27, 694/243] \subset P_1$. The first two subsequent control states are below 2. If $y \in [0, 16/27]$, the last control state is at most $194/81 < 12/5$; if $y \in [72/65, 4/3)$, it is at least $178/65 > L$ and below $26/9$. Thus it is above 2 and outside I , proving 1001 at all included endpoints.

For c , its source is in $[26/9, 3) \subset P_1$, since $26/9 + \eta^4 d_0 = 3$ and d_0 is excluded from D_0 . Its successive states are

$$\begin{aligned} 4/3 + (3/2)\eta^4 y &< 3/2, & 2 + \eta^3 y &< 9/4, & (3/2)\eta^3 y &< 3/8, & \eta^2 y &< 9/16, \\ (3/2)\eta^2 y &< 27/32, & \eta y &< 81/64, & (3/2)\eta y &< 243/128 < 2, & y. \end{aligned}$$

Only its source and its second subsequent state are above 2; both avoid I . This proves 10100000 with every intermediate state safe. We have obtained true inverse relations

$$C \xrightarrow{a,b} C, \quad C \xrightarrow{c} D_0, \quad D_0 \xrightarrow{a,b} C,$$

where each indicated target set is covered in its entirety by the specified inverse rule.

For any complete block string without cc , start with proof label C and update to D_0 after c , and to C after a, b . End at $0 \in C \cap D_0$ and build the path backwards using the table: for b with source label D_0 use its left rule; with source label C use the left rule for a target in P_1 and the right rule for a target in P_0 . The checked memberships justify every preceding target requirement. Forward execution by the same greedy state rule now realizes the prescribed even word, ends at zero, and continues safely forever. Different block strings give different words because the blocks are a prefix code. Thus $N_m(s_g) \geq H_m$.

Let $c_* = \min_{0 \leq j \leq 5} H_j / \rho^j > 0$. The full recurrence for $m \geq 6$ proves $H_m \geq c_* \rho^m$. This and (A.4) yield

$$h_{A_2}(s_g) = \log \rho. \quad (\text{A.5})$$

All inverse paths above avoid I . Therefore the same lower construction applies when an arbitrary Borel subset $E \subset [12/5, 162/65)$ is changed to control $B1$, with all other states greedy. Such a feedback is globally legal and Borel, but has $h_{A_2} \geq \log \rho$; this entire short-reset template cannot lower the greedy value. This is not a lower bound for the full four-control class or for arbitrary two- A Borel rules.

APPENDIX B. ARITHMETIC SAMPLING AND FINITE-BLOCK OPTIMIZATION

For a fixed globally safe Borel rule and arithmetic sampling $(0, p, 2p, \dots)$, a length- $n + m$ sampled pattern is determined by its first n entries and the length- m pattern starting from $T_s^{pm} x \in Q$. Consequently $N_{n+m}(s) \leq N_n(s) N_m(s)$. The logarithmic rate is a limit and equals $\inf_n n^{-1} \log N_n(s)$ (with division by τ when the control period is τ). This uses forward safety, not surjectivity or unrestricted concatenation.

For the full four-control class at A_2 , let K_n be the least actual count $N_n(s)$ among unrefined legal Borel feedbacks. It exists as a least member of a nonempty subset of $\{1, \dots, 4^n\}$. Merging refinements cannot increase counts, and unrefined rules are themselves allowed, so

$$\ell_2 = \inf_s \inf_n \frac{\log N_n(s)}{n} = \inf_n \frac{\log K_n}{n}. \quad (\text{B.1})$$

Only two infima are exchanged. No common feedback is asserted to attain every K_n , no minimizer of the asymptotic infimum is deduced, and no supremum over sampling sequences is interchanged. This formula turns a *proved real-feedback* finite-block upper bound into an asymptotic upper bound. An open-loop word table or a finite enumeration without global implementation is insufficient.

APPENDIX C. REAL ITINERARY CLOSURES AND THE UNRESOLVED LOW-ENTROPY IMPLEMENTATION

This appendix supplies a complete counterexample to a *relaxed* symbolic lower-bound claim. It is not a low-entropy feedback. For an arbitrary two- A rule s_E in Section 16.4, write its real binary name as $B_E(x)_j = \mathbf{1}_{\{s_E(T_E^j x) = A1\}}$ and put

$$Y_E = \text{cl } B_E(Q), \quad \pi(d) = 2 \sum_{j \geq 0} d_j (2/3)^j, \quad P_2(d) = (d_0, d_2, d_4, \dots).$$

Every coordinate of B_E is Borel, and the original state rule gives $\sigma B_E(x) = B_E(T_E x)$. Thus Y_E is nonempty, compact, and $\sigma Y_E \subset Y_E$, without continuity of the closed loop. The true affine recurrence gives

$$x = 2 \sum_{j=0}^{n-1} B_E(x)_j (2/3)^j + (2/3)^n T_E^n x.$$

The remainder is at most $6(2/3)^n$, so $\pi(B_E(x)) = x$. Continuity of π and the range $[0, 6]$ imply $\pi(Y_E) = Q$.

A cylinder fixing the first n even digits is open. If it meets the closure Y_E , it meets $B_E(Q)$; therefore closure adds no finite even patterns. Since $\sigma P_2 = P_2 \sigma^2$, $P_2(Y_E)$ is a one-sided subshift with exactly these real prefixes. Its prefix entropy is its topological entropy, by the same finite-cylinder refinement argument used in Theorem 6.1. Consequently

$$h_{\text{top}}(P_2(Y_E)) = h_{A_2}(s_E). \quad (\text{C.1})$$

No factor $1/2$ is inserted, because the rate is per sampled symbol at control period one.

C.1. A closed shift-invariant interval-covering language. Let

$$W_{10} = \{(a, b, a, c, d, e, f, g, h, g) : a, \dots, h \in \{0, 1\}\},$$

let Y_0 be arbitrary infinite concatenations of these ten-digit blocks, and set $Y = \bigcup_{i=0}^9 \sigma^i Y_0$. The equality conditions $d_{10j} = d_{10j+2}$ and $d_{10j+7} = d_{10j+9}$ make Y_0 closed. Its compact shift images have a finite closed union; adding or removing one full block gives $\sigma^{10} Y_0 = Y_0$ and $\sigma Y = Y$.

Proposition C.1. *The language Y satisfies*

$$\pi(Y) = Q, \quad h_{\text{top}}(P_2(Y)) = \frac{4}{5} \log 2 < \log \rho.$$

It therefore disproves the assertion that closedness, $\sigma Y \subset Y$, and full interval projection alone force the greedy even entropy. It does not provide an implemented Borel feedback or improve ℓ_2^A .

Proof. Put $r_0 = 2/3$, $\vartheta = r_0^{10} = 1024/59049$, $\Delta = 6\vartheta = 2048/19683$. The block projection $t_w = 2 \sum_{j=0}^9 w_j r_0^j$ is a subset sum of the eight weights

$$26/9, \quad 4/3, \quad 16/27, \quad 32/81, \quad 64/243, \quad 128/729, \quad 3328/19683, \quad 512/6561.$$

In increasing order their numerators over denominator 19683 are

$$1536, 3328, 3456, 5184, 7776, 11664, 26244, 56862.$$

The corresponding values of $19683(\Delta + \sum_{j < i} \kappa_j - \kappa_i)$ are

$$512, 256, 3456, 5184, 7776, 11664, 8748, 4374,$$

all positive. Thus each new weight is at most the earlier total plus Δ . Inductively, the intervals $[t, t + \Delta]$ over all subset sums fill the single interval $[0, \Delta + \sum_i \kappa_i] = [0, 6]$: adding a weight unites the preceding interval with its overlapping translate. Hence

$$Q = \bigcup_{w \in W_{10}} (t_w + \vartheta Q).$$

Given $x_0 \in Q$, choose a containing interval and write $x_0 = t_{w_0} + \vartheta x_1$, $x_1 \in Q$; repeat forever. Then $x_0 = \sum_{j < n} \vartheta^j t_{w_j} + \vartheta^n x_n$, with remainder at most $6\vartheta^n$. The infinite block concatenation belongs to Y_0 and projects to x_0 . This proves genuine infinite coverage, including all endpoints, not just finite-depth coverage. The reverse inclusion is automatic for binary sequences.

The even and odd block projections of Y_0 are (a, a, d, f, h) and (b, c, e, g, g) : each has exactly four free bits among five output positions. For a shift of a block concatenation, an even prefix consists of complete five-position outputs and at most two partial boundary blocks. There is a constant C , independent of the length, with prefix count at most $10 \cdot 2^{4n/5+C}$. On

lengths $n = 5m$, $P_2(Y_0)$ has 2^{4m} distinct prefixes. Since $P_2(Y)$ is a subshift, its entropy limit is exactly $(4/5) \log 2$. The strict comparison with ρ can be made without decimals: integer arithmetic gives $\rho > 1743/1000 > 2^{4/5}$ by substituting $1743/1000$ in the decreasing reciprocal equation for ρ , and checking $1743^5 > 16 \cdot 1000^5$. $\square$

The coding constructed in this proof is open loop. Its successive tails are safe states, since $\pi(\sigma d) = (3/2)\pi(d) - 3d_0$, but different codings of the same state may require incompatible first controls. The exact unresolved implementation question is whether there is a Borel $B : Q \rightarrow Y$ with, for *every* x ,

$$\pi(B(x)) = x, \quad B((3/2)x - 3B(x)_0) = \sigma B(x). \quad (\text{C.2})$$

A Borel right inverse of π alone does not meet this requirement. If both equations were proved, the real rule would have $h_{A_2} \leq (4/5) \log 2$ and would disprove greedy optimality. Neither that implementation nor its global impossibility is asserted.

C.2. The exact forced-predecessor compatibility kernel. For $a \in \{0, 1\}$ write $\kappa_a(d) = ad$ and define

$$Z_0 = Y, \quad Z_{n+1} = \{d \in Z_n : \sigma d \in Z_n, \pi(d) < 3 \Rightarrow 0d \in Z_n, \pi(d) > 3 \Rightarrow 1d \in Z_n\}.$$

These are closed and nested. For example the excluded condition $\pi(d) < 3$ together with $0d \notin Z_n$ is open; the other strict-threshold exclusion is likewise open. Every implementation in (C.2) has $B(Q) \subset Z_n$ for every n . Indeed, when $x < 3$, its original predecessor $\psi_0(x) = 2x/3 < 2$ has only control 0 legal, so $B(\psi_0 x) = 0B(x)$. When $x > 3$, $\psi_1(x) = 2 + 2x/3 > 4$ has only 1 legal and $B(\psi_1 x) = 1B(x)$. This, forward consistency, and induction give the claim. At $x = 3$, its predecessors 2, 4 are in overlap safety domains and supply neither compulsory condition.

The limit kernel has the following exact form:

$$H = \left\{ d \in Y : \forall j \geq 0, \begin{array}{l} \pi(\sigma^j d) < 3 \Rightarrow 00\sigma^j d \in Y, \\ \pi(\sigma^j d) > 3 \Rightarrow 11\sigma^j d \in Y \end{array} \right\} = \bigcap_n Z_n. \quad (\text{C.3})$$

Here each tested member may use its own phase in the union Y . To verify the equality, first note the elementary prefix fact: if $aad \in Y$, then $a^m d \in Y$ for every $m \geq 2$. In a fixed member phase, all visible copying pairs have distance two. On prepending one equal digit, the old pairs shift by one; the only newly exposed pair compares two of the equal initial digits. The phase is adjusted cyclically among the ten possible phases. Conversely, the visible pair conditions characterize a shift image because missing prefix partners can be filled freely. This proves the prefix fact with all phases allowed.

The set H is closed, is forward shift-invariant, and is closed under $d \mapsto 0d$ when $\pi(d) < 3$ and under $d \mapsto 1d$ when $\pi(d) > 3$. For example $00d \in Y$ gives $0d \in Y$; the new zero-tail test requires $000d \in Y$, supplied by the prefix fact, while the later tests are those of d . Thus $H \subset Z_n$ inductively. Conversely, the nested intersection is closed under shift and these forced prefixes: membership of d in every Z_{n+1} places the required image in every Z_n . For $\pi(d) < 3$, $\pi(0d) = 2\pi(d)/3 < 3$, so two such prefixes belong to the intersection, hence to Y ; the same argument holds above three. Applying it to every shifted tail proves the reverse inclusion in (C.3).

For each x , the fibers $Z_n \cap \pi^{-1}\{x\}$ are nested compact sets. Therefore

$$x \in \pi(H) \iff Z_n \cap \pi^{-1}\{x\} \neq \emptyset \text{ for every } n. \quad (\text{C.4})$$

This is an exact compactness interface, not a proof that the fibers are nonempty.

There is also the exact central reduction

$$[2, 3] \subset \pi(H) \iff \pi(H) = [0, 6]. \quad (\text{C.5})$$

Complementation $d \mapsto 1 - d$ preserves Y and H and sends the projection to $6 - \pi(d)$. If $0 < x < 2$, take the least k with $y = (3/2)^k x \geq 2$; then $2 \leq y < 3$. Starting with an H coding of y , repeated forced zero prefixes give a coding of x . The endpoints zero and six are the

constant names, and 2, 3, 4 are supplied by the central interval and complementation. This proves the nontrivial direction; the other is immediate. No condition is added at projection three.

Neither (C.4) nor (C.5) decides the remaining coverage of $[2, 3]$. Even full coverage of H would not by itself give a single-step-consistent Borel section. A missing fiber would exclude this language's global implementation but would not prove greedy optimality or exclude other lower-entropy feedbacks. Accordingly the bounds (16.1) and (16.3) are unchanged.

APPENDIX D. A BOUNDED-DENSITY OBSTRUCTION TO SIMULTANEOUS SCALAR CAPACITIES

The smooth branches in Section 13.4 cannot all be scalarized by an equivalent probability with a density bounded both above and away from zero. The proof treats every such Borel density, without inserting it at a null periodic point.

Proposition D.1. *There are no Borel a and positive c_0, c_1, c_2 with $0 < \text{ess inf } a \leq \text{ess sup } a < \infty$,*

$$\rho_u(y)a(\psi_u(y)) \leq c_u a(y) \quad \text{almost everywhere,} \quad \max_{u \neq v} (c_u + c_v) < 1.$$

Here all three branches use the same density.

Proof. Choose $0 < m \leq M < \infty$ bounding a almost everywhere, and remove the null exceptions for these bounds and the three inequalities. Also remove their preimages under every finite composition of the ψ_u . These are null because every composition has a Lipschitz inverse on its image: its derivative is bounded below by $(2/15)^n$. The remaining Borel set G has full measure and is preserved by every inverse branch. For $y_0 \in G$ and $y_{j+1} = \psi_{u_j}(y_j)$, multiplication gives

$$\left(\prod_{j < n} \rho_{u_j}(y_j) \right) a(y_n) \leq \left(\prod_{j < n} c_{u_j} \right) a(y_0). \quad (\text{D.1})$$

Since $\psi_0(0) = 0$, $\psi_0(y) \leq 3y/5$, and $\rho_0(0) = 3/5$, every $\eta > 0$ has a small invariant interval on which $\rho_0 \geq 3/5 - \eta$. Choose a point of G in its interior, iterate, and take n th roots in $m(3/5 - \eta)^n \leq M c_0^n$. Letting n grow and then η decrease proves $c_0 \geq 3/5$.

For a second constraint use the positive-length closed intervals

$$\begin{aligned} I_0 &= [9/10, 23/25], & I_1 &= [3/10, 33/100], \\ I_2 &= [9/20, 47/100], & I_3 &= [53/100, 14/25], \\ I_4 &= [19/25, 79/100], & I_5 &= I_0. \end{aligned}$$

The inverse word is $(0, 1, 1, 2, 2)$. The following bounds are rational containing intervals for the *complete* images:

Input and branch	Containing interval	Next interval
$\psi_0(I_0)$	$[3081/10000, 3195/10000]$	int I_1
$\psi_1(I_1)$	$[4510/10000, 4692/10000]$	int I_2
$\psi_1(I_2)$	$[5385/10000, 5494/10000]$	int I_3
$\psi_2(I_3)$	$[7672/10000, 7836/10000]$	int I_4
$\psi_2(I_4)$	$[9002/10000, 9170/10000]$	int I_0

For an explicit verification of every endpoint bound, use $314159/100000 < \pi < 314160/100000$ and $173205/100000 < \sqrt{3} < 173206/100000$. The latter bounds follow by squaring; the former follow from Machin's identity with the alternating arctangent sums through degrees 11 and 3 at $1/5$ and $1/239$. At the two control-zero endpoints use $z = 2\pi(y - 1)$, and at the others use $z = 2\pi(y - u/3)$. In every case $|z| < 1$. Substitute the rational interval for z in

$$S_7(z) = z - z^3/6 + z^5/120 - z^7/5040, \quad |\sin z - S_7(z)| \leq 1/362880.$$

The constant sine terms are $0, -\sqrt{3}/2, \sqrt{3}/2$, and $7/(60\pi)$ is bounded by the reciprocal rational interval. Interval addition and multiplication at each endpoint lie strictly inside the displayed containing bounds. Monotonicity of ψ_u then proves the full-image inclusions, including endpoints.

On each input interval the distance to its corresponding cosine center $(1, 1/3, 2/3)$ is at most $41/300$. Thus

$$\rho_{u_j}(y) \geq \frac{3}{5} - \frac{7}{60}(41\pi/150)^2 > \frac{1214099}{2362500} > \frac{1}{2}.$$

The composite $\psi_2\psi_2\psi_1\psi_1\psi_0$ preserves I_0 . Starting at any $y \in G \cap I_0$, repeat its five-step word n times in (D.1). This yields $m(1/32)^n \leq M(c_0c_1^2c_2^2)^n$, and hence $c_0c_1^2c_2^2 \geq 1/32$. But the pairwise sums and $3/5 \leq c_0 < 1$ give $c_1, c_2 < 1 - c_0$. The function $t(1-t)^4$ decreases on $[3/5, 1)$, so

$$c_0c_1^2c_2^2 < \frac{3}{5}(2/5)^4 = \frac{48}{3125} < \frac{1}{32},$$

a contradiction. All iterations occur in the same full-measure set; no regularity of a was assumed. $\square$

For $d\mu_a = a dy / \int a$, the density of the branch measure relative to μ_a is exactly

$$\rho_u^a(y) = \rho_u(y)a(\psi_u(y))/a(y),$$

by the change of variables $x = \psi_u(y)$. The proposition excludes this bounded positive density class. It does not exclude arbitrary full-support singular probabilities or densities without uniform bounds; that question remains open for this smooth example. It does not alter its already proved value $M_1 = \log 3$.

There is an exact reduction of that remaining question, without a density assumption. If a full-support probability μ satisfied $\mu(\psi_u(E)) \leq c_u\mu(E)$ for all Borel sets and the same target pair sums, then $q_0 = \max c_u < 1$ would force it to be nonatomic. Indeed, an atom of mass $m > 0$ can be followed along any successive legal controls, since the safety intervals cover Q . Applying the set inequality to the successor singleton raises its mass to at least m/q_0 at each step, eventually exceeding one. Repetition of states does not avoid that contradiction.

Its distribution function $h(x) = \mu([0, x])$ would therefore be a continuous strictly increasing homeomorphism of $[0, 1]$, sending μ to Lebesgue probability. Applying the inequalities to $(x, y]$ shows that every $h\psi_u h^{-1}$ is c_u -Lipschitz. Conversely, any one increasing homeomorphism with these simultaneous Lipschitz bounds defines a full-support nonatomic measure $\mu = (h^{-1})_* \text{Leb}$ with the required inequalities on every Borel set. To see the last assertion, cover a Borel set in the h coordinate by an open set of arbitrarily close measure. Decompose that open set into countably many intervals; each image has length at most c_u times its length. Subadditivity and outer regularity give the full Borel estimate, not merely an interval estimate. This equivalence leaves existence of h undecided and is not an extension of the preceding exclusion.

APPENDIX E. CONDITIONAL REAL RETURNS AND CLOSED VIABILITY BOUNDARIES

E.1. A real two-branch certificate and one common sampling. The affine system below is the four-branch system of Section 16.4. A sufficient certificate is stated with its true fibers; extracting it for every Borel feedback is an unresolved step.

Proposition E.1. *Suppose one legal feedback has distinct real control words w^0, w^1 of the same length k , and a nonempty open interval $I \subset (0, 6)$, such that its real prefix fiber E_{w^i} is comeager in the complete inverse interval $J_i = G_{w^i}^{-1}(I)$, and $J_i \subset I$. For the single predetermined sequence $A = (j!)_{j \geq 1}$ this feedback has $h_A(s) \geq \log 2$.*

Proof. Each inverse $\psi_i : I \rightarrow J_i \subset I$ is an affine homeomorphism, and its bad subset $J_i \setminus E_{w^i}$ is meager. For a prescribed finite sequence of the two words, compose their complete inverse branches backwards from I . The resulting initial interval is nonempty and open; the inverse images of every bad set along the finite affine maps are meager. A point outside their finite

union executes precisely the prescribed sequence by the same feedback. It remains safe thereafter by global legality.

Choose an offset $0 \leq t < k$ where the two words differ. For all $j \geq k$, $j!$ is divisible by k . Prescribe independently the word in block $j!/k$ of a sufficiently long concatenation, fill all other blocks arbitrarily, and start observation at the actual state $T_s^t x$. Its control at time $j!$ is the offset- t letter of that chosen block. Thus the first n factorial samples have at least $2^{n-O(k)}$ patterns. The ignored head is fixed and the normalized limsup gives the claim. The word length k is a proof device; the feedback period is still one. $\square$

For any expanding affine word $G_w(x) = \lambda_w x - t_w$ and $I = (\alpha, \beta)$, elementary endpoint inequalities give

$$G_w^{-1}(I) \subset I \iff t_w/(\lambda_w - 1) \in [\alpha, \beta].$$

The interval must be the complete inverse of I ; a locally feasible subinterval is insufficient. The proposition would give $L_1 = M_1 = \log 2$ if its certificate were extracted for every legal Borel rule, even with k depending on the rule. Such an extraction is not proved here. Compactness of the state interval does not make the true return graph finite, and openness of each individual affine branch does not imply Borel closed-loop stitching.

E.2. Nested viable sets and the limits of a growth route. For a closed $K \subset Q = [0, 6]$, let

$$\Gamma(K) = \{x : \exists \text{ a legal } u, g_u(x) \in K\}, \quad c(K) = \sup\{\text{diam } C : C \text{ a component of } K\}.$$

For closed C , define $C^{(0)} = C$, $C^{(n+1)} = C^{(n)} \cap \Gamma(C^{(n)})$, and $V(C) = \bigcap_n C^{(n)}$. All these sets are compact. Finite controls and nested compactness show $V(C) \subset \Gamma(V(C))$: the nested nonempty sets of legal controls carrying a point into $C^{(n)}$ have a common control. Conversely every closed viable $K \subset C$ remains in every iterate and hence in $V(C)$.

Proposition E.2. *Every decreasing sequence of closed viable sets K_j has one globally safe Borel selector preserving all K_j . If also $\text{int}_Q K_j \neq \emptyset$ and $c(K_j) \rightarrow 0$, this selector fails complete-branch growth at every positive fixed scale. No sequence satisfying these additional interior and component-scale conditions can have a finite intersection.*

Proof. On $K_j \setminus K_{j+1}$ choose the first original control carrying the state into K_j . On $K_\infty = \bigcap_j K_j$ choose the first control carrying it into all K_j ; a control exists because the finite nonempty admissible-control sets are decreasing. Outside K_1 use any ordered safe selector. The selection domains are Borel, the choices are pointwise consistent, and the resulting rule preserves each K_j for every state, hence along infinite real trajectories.

Fix $\delta > 0$, choose j with $c(K_j) < \min\{\delta, 6\}$, and an open interval $V \subset \text{int}_Q K_j$. A complete actual inverse branch starting in V and ending onto a relative open ball $B_Q(y, \delta)$ would send a comeager subset into K_j . Its affine homeomorphism makes the image dense in that ball. Closedness forces the entire ball into K_j , but this connected ball has diameter at least $\min\{\delta, 6\}$, contradicting the component bound. Length zero gives the same contradiction. This is precisely the complete-branch growth obstruction, not an entropy counterexample.

For the last assertion, impose the additional conditions just stated. Nested nonempty compactness makes $F = \bigcap K_j$ nonempty. Suppose that F is finite. Choose a closed η -neighborhood of F small enough that its nearest center is unique and every actual safe step staying in it has $g_u(p_j) = p_{j+1}$ for the corresponding centers. Such an η exists: for the finitely many nonzero distances $|g_u(p) - q|$, choose η smaller than their minimum divided by $1 + \max_u \lambda_u$, and also smaller than one third of the minimum center separation. If all relevant distances vanish, only the separation restriction is needed. Along a trajectory remaining in this neighborhood, $|x_n - p_n| \geq (3/2)^n |x_0 - p_0|$, whereas the left side is at most η . Therefore $x_0 \in F$, and the full viability kernel of the neighborhood is contained in F . Nested compactness places some K_j inside this neighborhood. Its viability then places it in F , forcing eventual equality with F . It cannot have nonempty interior, a contradiction. $\square$

There is a further exact restriction: a nondegenerate closed interval $[a, b] \subset [0, 6]$ is viable if and only if $b - a \geq 3$. The allowable $A0$ source interval inside it is $[a, 2b/3]$, and its $A1$ source interval is $[2 + 2a/3, b]$, with empty intervals discarded. The corresponding $B0, B1$ source pieces are contained in these, because $3b/5 \leq 2b/3$ and $(3a + 12)/5 \geq 2 + 2a/3$ for $a \leq 6$. Thus they cover the interval exactly when $2b/3 \geq 2 + 2a/3$, equivalently $b - a \geq 3$. Nonconvex viable sets need not preserve their separate components, so this statement is not extended to their convex hulls.

An infinite nested viable chain with nonempty interior and vanishing component scale has not been constructed or excluded. The conditional growth obstruction above neither excludes all real return certificates nor changes L_1 . These unresolved steps are kept separate from the complete maximum-pattern theorems.

APPENDIX F. SCOPE, PROOF STATUS, AND OPEN PROBLEMS

The primary new structural conclusion is the inverse-list product criterion and its source-geometry applications, particularly the gain-loss class without a common nonstrict positive left budget. The closed-cover theorem answers a different common-sampling question. The countable-language theorem, symbolic realization, and period collapse provide the framework in which both are stated.

The main remaining question is whether hypotheses that force $M_\tau = \log r/\tau$ through product capacities also force $L_\tau = M_\tau$, or whether there is a genuine strict numeric gap. The positive-density independent positions furnished here may vary with the feedback, so they do not answer this question. For the four-branch affine system, both fixed-even-sampling infima and greedy optimality remain open within the stated intervals. The ten-block language's full Borel implementation and the full-support scalarization of the smooth example are also unresolved.

No theorem is inferred from a finite search, an open-loop word table, or a merely almost-everywhere implementation. Every optimization upper bound is furnished by a globally defined real feedback.

Use of AI tools. OpenAI ChatGPT/Codex tools assisted the argument exploration, candidate proofs, comparison of primary literature, manuscript integration, and automated source, compilation, and PDF checks. These activities included mathematical work and were not limited to language editing. Model versions for the complete earlier research history cannot be reliably reconstructed from the record.

REFERENCES

- [1] Debauche, V., Della Rossa, M., & Jungers, R. M. (2022). Comparison of path-complete Lyapunov functions via template-dependent lifts. *Nonlinear Analysis: Hybrid Systems*, 46, 101237. doi:10.1016/j.nahs.2022.101237. The comparison used here is with the author version arXiv:2110.13474v1 (2021).
- [2] Goodman, T. N. T. (1974). Topological sequence entropy. *Proceedings of the London Mathematical Society*, 29(2), 331–350. doi:10.1112/plms/s3-29.2.331.
- [3] Huang, W., Maass, A., & Ye, X. (2004). Sequence entropy pairs and complexity pairs for a measure. *Annales de l'Institut Fourier*, 54(4), 1005–1028. doi:10.5802/aif.2041.
- [4] Huang, W., & Ye, X. (2009). Combinatorial lemmas and applications to dynamics. *Advances in Mathematics*, 220(6), 1689–1716. doi:10.1016/j.aim.2008.11.009.
- [5] Hulse, P. (1982). Sequence entropy and subsequence generators. *Journal of the London Mathematical Society*, 26(3), 441–450. doi:10.1112/jlms/s2-26.3.441.
- [6] Kamae, T., & Zamboni, L. (2002). Maximal pattern complexity for discrete systems. *Ergodic Theory and Dynamical Systems*, 22(4), 1201–1214. doi:10.1017/S0143385702000585.
- [7] Kawan, C. (2013). *Invariance entropy for deterministic control systems: An introduction*. Lecture Notes in Mathematics, Vol. 2089. Springer. doi:10.1007/978-3-319-01288-9.
- [8] Kerr, D., & Li, H. (2007). Independence in topological and C^* -dynamics. *Mathematische Annalen*, 338, 869–926. doi:10.1007/s00208-007-0097-z.
- [9] Kerr, D., & Li, H. (2009). Combinatorial independence in measurable dynamics. *Journal of Functional Analysis*, 256(5), 1341–1386. doi:10.1016/j.jfa.2008.12.014.

- [10] Romagnoli, P.-P. (2007). A remark on the topological entropies of covers and partitions. *Studia Mathematica*, 182(3), 273–281. doi:10.4064/sm182-3-6.
- [11] Shi, L., Zheng, W. X., Shao, J., & Cheng, Y. (2021). Sub-Super-stochastic matrix with applications to bipartite tracking control over signed networks. *SIAM Journal on Control and Optimization*, 59(6), 4563–4589. doi:10.1137/19M1239982. The product assumptions discussed here were read in the author version arXiv:2004.01867v2 (2020), Definitions 2.5 and Theorems 2.6 and 2.8, pp. 8–11; they are not asserted to apply directly to our list products.
- [12] Tomar, M. S., Kawan, C., & Zamani, M. (2022). Numerical over-approximation of invariance entropy via finite abstractions. *Systems & Control Letters*, 170, 105395. doi:10.1016/j.sysconle.2022.105395.

SCHOOL OF MATHEMATICAL SCIENCES, NANJING NORMAL UNIVERSITY, NANJING 210023, CHINA