

Search for New Physics in Events with Jets and Missing Transverse Momentum at the CMS Experiment

Dissertation

zur Erlangung des Doktorgrades

des Fachbereichs Physik

der Universität Hamburg

vorgelegt von

Dipl.- Phys. Kristin Goebel geb. Heine

aus Gifhorn

Hamburg

2014

Gutachter der Dissertation:	Prof. Dr. bla Prof. Dr. bla
Gutachter der Disputation:	Prof. Dr. bla Prof. Dr. bla
Datum der Disputation:	?? . ?? 2014
Vorsitzender des Prüfungsausschusses:	Dr. bla
Vorsitzender des Promotionsausschusses:	Prof. Dr. bla
Leiterin des Fachbereichs Physik:	Prof. Dr. bla
Dekan der Fakultät für Mathematik, Informatik und Naturwissenschaften:	Prof. Dr. bla

Abstract

bla

Kurzfassung

bla

Contents

1	Introduction	1
2	Theoretical Background	3
2.1	The Standard Model of Particle Physics	3
2.1.1	Shortcomings of the Standard Model	3
2.2	Supersymmetry	3
2.2.1	Searches for Supersymmetry at Collider Experiments	3
3	Experimental Setup	5
3.1	The Large Hadron Collider	5
3.2	The CMS Experiment	6
3.2.1	Coordinate System and Kinematic Variables	7
3.2.2	Inner Tracking System	7
3.2.3	Electromagnetic Calorimeter	7
3.2.4	Hadronic Calorimeter	7
3.2.5	Muon System	7
3.2.6	Trigger System	7
3.3	Data Taking and Event Simulation	7
4	Reconstruction Algorithms and Object Definition	9
4.1	Global Event Description with the Particle-Flow Algorithm at CMS	9
4.2	Reconstruction of Jets	9
4.2.1	Jet Algorithms	9
4.2.2	Jet Types at CMS	9
4.2.3	Jet Energy Calibration	9
4.3	Identification of Boosted Top Quark Decays	9
4.3.1	The CMS Top Tagger	9
4.3.2	The HEP Top Tagger	9
5	Measurement of the Jet Transverse-Momentum Resolution	11
5.1	Basic Concept of the Dijet Asymmetry Method	11
5.2	Application to Realistic Collision Events	11
5.3	Samples and Event Selection	11
5.3.1	Datasets and Triggers	11
5.3.2	Selection Criteria	11
5.4	Corrections to the Dijet Asymmetry	11
5.4.1	Correction for Additional Jet Activity	11
5.4.2	Correction for Particle-Level Imbalance	11
5.4.3	Results of the Corrections to the Asymmetry	11
5.5	Determination of the Data-to-Simulation Ratio	11

5.6	Validation of the Method	11
5.6.1	Validation in Simulated Events	11
5.6.2	Validation of the Measured Data-to-Simulation Ratio	11
5.7	Systematic Uncertainties	11
5.8	Extension of the Method to the Forward Detector Region	11
5.9	Results	11
5.9.1	Comparison to Other Measurements	11
6	Search for New Physics with Jets and Missing Transverse Momentum	13
6.1	Event Selection	13
6.1.1	Data samples and trigger	13
6.1.2	Event Cleaning	13
6.1.3	Baseline Selection	13
6.1.4	Exclusive Search Regions	13
6.2	QCD Background Estimation with the Rebalance-And-Smear Method . . .	13
6.2.1	Rebalance Procedure using Kinematic Fits	13
6.2.2	Response Smearing	13
6.2.3	Validation Tests	13
6.2.4	Systematic Uncertainties	13
6.2.5	QCD Background Prediction	13
6.3	Estimation of Non-QCD Backgrounds	13
6.3.1	Invisible Z Background	13
6.3.2	Hadronic τ Background	13
6.3.3	Lost-Lepton Background	13
6.4	Results and Interpretation	13
7	Prospect Studies for a Search for Top Squarks	15
7.1	Data samples	15
7.2	Top Tagging Efficiency Studies	15
7.2.1	Top Tag Efficiency	15
7.2.2	Misidentification Rate	15
7.3	Identification of Selection Criteria	15
7.4	Comparison of the Performance of Various Selections	15
7.5	Impact of Specific Assumptions in the Analysis	15
8	Conclusions	17

1 Introduction

blabla bla [1]

2 Theoretical Background

2.1 The Standard Model of Particle Physics

2.1.1 Shortcomings of the Standard Model

2.2 Supersymmetry

2.2.1 Searches for Supersymmetry at Collider Experiments

3 Experimental Setup

In order to probe the various aspects of the well-established standard model or search for hints of new physics beyond the SM, particle physics experiments preferentially make use of powerful particle accelerators where particles of a certain type are collided in order to probe the constituents of matter and interactions between them. The analyses presented in this thesis are all performed in the context of the CMS experiment located at the Large Hadron Collider (LHC) at CERN near Geneva.

The first part of this chapter provides an introduction to the LHC. This is followed by an overview of the detector system of the CMS experiment. Afterwards the hitherto periods of collision data taking at the LHC are discussed together with an introduction to the generation of simulated events which are used in the analysis of real data events.

3.1 The Large Hadron Collider

The Large Hadron Collider [2, 3] is a ring-accelerator designed to provide particle collisions of hadrons. It is built in the tunnel of the former LEP [4] collider 45 – 170 m below the ground and has a circumference of 26.7 km. The LHC is a particle-particle collider and thus composed of two rings with counter-rotating beams. The operation can be performed in different modes with either proton beams or heavy ions like e.g. lead ¹.

In each beam, protons are grouped together in bunches and accelerated in two evacuated beam pipes using superconducting radio-frequency cavities. With a nominal bunch spacing of 25 ns the bunch revolution frequency is 40 MHz. Each of the 2808 individual bunches per beam contains at design conditions 1.15×10^{11} protons. In order to bend the beams around the LHC ring superconducting dipole magnets are used with an operation temperature of 1.9 K. They provide a magnetic field of up to 8.33 T while additional quadrupole and sextupole magnets are utilized to squeeze and focus the beams.

Before the protons are injected into the LHC they are already pre-accelerated in various smaller accelerators up to a beam energy of 450 GeV while passing through the injector chain Linac2 – Proton Synchrotron Booster (PSB) – Proton Synchrotron (PS) – Super Proton Synchrotron (SPS). An overview of the accelerator complex at CERN is given in Fig. 3.1.

The main goal of the LHC is to provide proton-proton collisions to the experiments with center of mass energies up to 14 TeV in order to explore physics processes at novel energy regimes. The expected number of events N for a certain type of process is given by the product of the specific cross section σ of that process and the integral $L = \int \mathcal{L} dt$ of the instantaneous luminosity \mathcal{L} over time such that

$$N = \sigma \cdot L. \tag{3.1}$$

¹All studies presented in this thesis are based on proton-proton collisions. Thus the operation with heavy ions is not discussed.




Figure 3.1: Illustration of the CERN accelerator complex. Numbers below the names of individual machines indicate the year of their first operation. For ring accelerators also the circumference is given. Taken from [5].

The luminosity is a machine parameter and can be expressed for beams with Gaussian-shaped profiles as

$$\mathcal{L} = f \frac{n_1 n_2}{4\pi \sigma_x \sigma_y} \quad (3.2)$$

with the revolution frequency f , the number of particles n_1 and n_2 contained in the two colliding bunches and the transverse beam sizes σ_x (σ_y) in the horizontal (vertical) directions. The nominal peak luminosity of the LHC is $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$.

The LHC beams cross at four locations along the ring. At these interaction points the four main experiments of the LHC are located in order to measure the delivered particle collisions. The two high luminosity experiments ATLAS [6] and CMS [7,8] are designed for multiple purposes like precision measurements of SM quantities, search for the standard model Higgs Boson or searches for signals indicating new physics processes. The LHCb detector [9] however is a specialised experiment focusing on the measurement of CP violation in the interactions of hadrons containing b-quarks. The only experiment designed especially for the analysis of heavy ion collisions is the ALICE [10] detector with the main emphasis on the physics of strongly interacting matter at extreme energy densities like for instance quark-gluon plasma.

3.2 The CMS Experiment

The CMS detector is one of the two experiments at the LHC designed to address a multitude of physics questions. In addition to tests of the SM at the TeV scale, studies of the nature of elektroweak symmetry breaking which might show up in the presence of a Higgs boson and searches for so far unknown particles pointing to e.g. new symmetries in nature are the primary targets of these experiments. These ambitious physics goals can only be achieved by fully exploiting the by now unprecedented collision energy and luminosity.

Since the total inelastic proton-proton cross-section at a center of mass energy of 14 TeV is expected to be around 100 mb, the experiments have to deal with an event rate of approximately 10^9 events per second. This is resulting in high experimental challenges. The CMS detector with its typical cylindrical design of different sub-detector components around the beam line is designed to perfectly meet these particular conditions. As a typical high-energy particle experiment the CMS detector makes mainly use of tracking detectors and calorimeters to measure particles' momenta, energy depositions and flight directions in order to identify the objects emerging from the particle collisions. Table ... gives an overview of the performance goals of the various sub-detectors.

Bild + Ref

Performance
Table

The following sections comprise a description of the CMS detector and individual sub-detector components focusing on the detector parts most relevant for the analyses presented in this thesis. A detailed discussion of the detector design can be found in [7, 8].

3.2.1 Coordinate System and Kinematic Variables

3.2.2 Inner Tracking System

3.2.3 Electromagnetic Calorimeter

3.2.4 Hadronic Calorimeter

3.2.5 Muon System

3.2.6 Trigger System

3.3 Data Taking and Event Simulation

4 Reconstruction Algorithms and Object Definition

4.1 Global Event Description with the Particle-Flow Algorithm at CMS

4.2 Reconstruction of Jets

4.2.1 Jet Algorithms

4.2.2 Jet Types at CMS

4.2.3 Jet Energy Calibration

4.3 Identification of Boosted Top Quark Decays

4.3.1 The CMS Top Tagger

4.3.2 The HEP Top Tagger

5 Measurement of the Jet Transverse-Momentum Resolution

5.1 Basic Concept of the Dijet Asymmetry Method

5.2 Application to Realistic Collision Events

5.3 Samples and Event Selection

5.3.1 Datasets and Triggers

5.3.2 Selection Criteria

5.4 Corrections to the Dijet Asymmetry

5.4.1 Correction for Additional Jet Activity

5.4.2 Correction for Particle-Level Imbalance

5.4.3 Results of the Corrections to the Asymmetry

5.5 Determination of the Data-to-Simulation Ratio of the Jet Transverse Momentum Resolution

5.6 Validation of the Method

5.6.1 Validation in Simulated Events

5.6.2 Validation of the Measured Data-to-Simulation Ratio

5.7 Systematic Uncertainties

5.8 Extension of the Method to the Forward Detector Region

5.9 Results

5.9.1 Comparison to Other Measurements

6 Search for New Physics in the Multijet and Missing Transverse Momentum Final State at $\sqrt{s} = 8$ TeV

6.1 Event Selection

6.1.1 Data samples and trigger

6.1.2 Event Cleaning

6.1.3 Baseline Selection

6.1.4 Exclusive Search Regions

6.2 QCD Background Estimation with the Rebalance-And-Smear Method

6.2.1 Rebalance Procedure using Kinematic Fits

6.2.2 Response Smearing

6.2.3 Validation Tests

6.2.4 Systematic Uncertainties

6.2.5 QCD Background Prediction

6.3 Estimation of Non-QCD Backgrounds

6.3.1 Invisible Z Background

6.3.2 Hadronic τ Background

6.3.3 Lost-Lepton Background

6.4 Results and Interpretation

7 Prospect Studies for a Search for Top Squarks in Events with Jets and Missing Transverse Momentum at $\sqrt{s} = 13$ TeV

7.1 Data samples

7.2 Top Tagging Efficiency Studies

7.2.1 Top Tag Efficiency

7.2.2 Misidentification Rate

7.3 Identification of Selection Criteria

7.4 Comparison of the Performance of Various Selections

7.5 Impact of Specific Assumptions in the Analysis

8 Conclusions

bla

Bibliography

- [1] K. Nakamura and et al. (Particle Data Group), “Review of particle physics”, *J. Phys. G* **37** (2010, and 2011 partial update for the 2012 edition) 075021. doi:10.1088/0954-3899/37/7A/075021.
- [2] O. S. Brüning, P. Collier, P. Lebrun et al., “LHC Design Report”, volume 1: The LHC Main Ring. CERN, Geneva, 2004. CERN-2004-003-V-1.
- [3] L. Evans and P. Bryant, “LHC Machine”, *Journal of Instrumentation* **3** (2008), no. 08, S08001. doi:10.1088/1748-0221/3/08/S08001.
- [4] CERN, “LEP Design Report”, volume 2: The LEP Main Ring. CERN, Geneva, 1984. CERN-LEP-84-01.
- [5] C. Lefèvre, “The CERN accelerator complex”. CERN-DI-0812015, 2008.
- [6] ATLAS Collaboration, “The ATLAS Experiment at the CERN Large Hadron Collider”, *JINST* **3** (2008) S08003. doi:10.1088/1748-0221/3/08/S08003.
- [7] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004. doi:10.1088/1748-0221/3/08/S08004.
- [8] CMS Collaboration, “CMS Detector Performance and Software”, volume 1 of *CMS Physics Technical Design Report*. CERN, Geneva, 2006. CMS-TDR-008-1. CERN-LHCC-2006-001.
- [9] LHCb Collaboration, “The LHCb Detector at the LHC”, *JINST* **3** (2008) S08005. doi:10.1088/1748-0221/3/08/S08005.
- [10] ALICE Collaboration, “The ALICE experiment at the CERN LHC”, *JINST* **3** (2008) S08002. doi:10.1088/1748-0221/3/08/S08002.

