1. The probability of a leap year selected at random contain 53					
Sunday is:					
(a) 53/366 (b) 1/7 (c) 2/7 (d) 53/365					
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at					
random. The probability of drawing a black ball is:					
(a) 3/5 (b) 2/5 (c) 0/5 (d) 1/5					
3. The probability that it will rain tomorrow is 0.85. What is the					
probability that it will not rain tomorrow					
(a) 0.25 (b) 0.145 (c) 3/20 (d) none of these					
4. What is the probability that a number selected from the number	rs				
(1, 2, 3,,15) is a multiple of 4?					
(a) 1/5 (b) 4/5 (c) 2/15 (d) 1/3					
5. What are the total outcomes when we throw three coins?					
(a) 4 (b) 5 (c) 8 (d) 7					
6. The probability that a prime number selected at random from t	he				
numbers (1,2,3,35) is :					
(a) 12/35 (b) 11/35 (c) 13/35 (d) none of these	:				
7. The sum of the probability of an event and non event is :					
(a) 2 (b) 1 (c) 0 (d) none of these.					
8. The following probabilities are given; choose the correct answ	er				
for that which is not possible.					
(a) 0.15 (b) 2/7 (c) 7/5 (d) none of these.					
9. If three coins are tossed simultaneously, than the probability of	f				
getting at least two heads, is:					
(a) 1/4 (b) 3/8 (c) ½ (d) 1/8					
10. A letter is chosen at random from the letters of the word					
ASSASSINATION. The probability that the letter chosen has:					
(a) 6/13 (b) 7/13 (c) 1 (d) none of thes	e.				
11. A dice is thrown. Find the probability of getting an even number	r.				
(A) 2/3 (B) 1 (C) 5/6 (D) 1/2					
12. Two coins are thrown at the same time. Find the probability of					
getting both heads.					
(A) 3/4 (B) 1/4 (C) 1/2 (D) 0					
13. Two dice are thrown simultaneously. The probability of getting a					
and the same and an arminimized and the probability or gotting	, –				

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/9				
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
	3) 27/50	(C) 1/4	(D) 2	9/100			
of drawing a b	lue ball is do bag is:	uble that of a r		the probability the number of			
	andom from t bulb is:	tains 12 defecthis box. Then to the contract of the contract o	he probabili				
17. Cards mar mixed thoroug	ked with num ghly. One card that the num	nbers 2 to 101 d is drawn from nber on card is	are placed ir n this box rar	n a box and ndomly, then			
		of getting 53 N					
probability of	getting a king	well shuffled of of red suit. 7/52 (D)		ards. Find the			
equally likely to 1,2,312 ,th	to come to res en the probab	sists of spinning to continuous that it will (C) 7/12	ne of the nu I point to an				
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) $3/4$ (B) $1/2$ (C) 1 (D) $1/4$							

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
(A) 364/365		The second secon	(D) 1/133225				
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$							
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is $2/3$, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7							
Then the probab	25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) $7/50$ (B) $4/25$ (C) $1/25$ (D) $2/25$						
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$				
			nes are 50, 70, 82,				
93, and 20. The sand 25.79) 25.49	c) 25.29	d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
a) 13, 15	b) 13, 18	c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) 1/2 30. X is a variate	b) $\frac{1}{3}$	c) $^1\!/_4$ d 3. The value of	$^{\mathrm{d}})^{1}/_{6}$ E(X ²) is				
a) 8 b) 7	22) 27 d)	9				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

a) 3	b) 4	c) 5	d) 7					
32.Out of the following values, which one is not possible in probability?								
a) $P(x) = 1$ b) $\sum x P(x) = 3$ c) $P(x) = 0.5$ d) $P(x) = -0.5$								
33.If E(x) =	2 and E(z) = 4 , b) 6	then E(z - x c) 0		fficient data				
		,						
34.The cov	ariance of two	independent	random variable	is				
a) 1	b) 0	c) - 1	d) Und	efined				
35.If $\Sigma P(x) = k^2 - 8$ then, the value of k is? a) 0 b) 1 c) 3 d) Insufficient data								
36.If $P(x) = 0.5$ and $x = 4$, then $E(x) = ?$ a) 1 b) 0.5 c) 4 d) 2								
37.In a discrete probability distribution, the sum of all probabilities is always?								
a) 0	b) Infinite	c) 1	d) Unde	efined				
38.If the probability of hitting the target is 0.4, find mean and								
variance. a) 0.4, 0.24	b) 0.6, 0	0.24	c) 0.4, 0.16	d) 0.6, 0.16				
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6 , 0.24 b) 6 , 2.4 c) 0.4 , 0.16 d) 4 , 1.6								
 40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution? 								

		and varia		•				
a) E 43 .	(X) Mean of	a random	2) variable	c) E(X2	en by)2	d) (E(X))2	
44.N	a) E(X) b) E(X2) c) E(X2) - (E(X))2 d) (E(X))2 44.Mean of a constant 'a' is a) 0 b) a c) a/2 d) 1							
a) 0	45. Variance of a constant 'a' is a) 0 b) a c) a/2 d) 1 46. Find the mean and variance of X?							
	x f(x)	0	2/9	2 3/9	3 2/9	1/9		
a) 2	4/3		. 4/3		c) 2. 2/3		d) 3. 2/3	

47. Find the expectation of a random variable X?

	Х	0	1	2	3		
	f(x)	1/6	2/6	2/6	1/6		
a) ().5	13	b) 1.5			c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

5

- a) np
- b) npq
- c) np2q
- d) npq2

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**

- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}