

a
fraction outside the reference range (5% expected)
bacterial families
other reference adults (28 studies, n = 1,809)
US participants of the Hadza study (n = 12)
Hadza, Tanzania (n = 350)
gene families (KOs)
other reference adults (28 studies, n = 1,809)
US participants of the Hadza study (n = 12)
Hadza, Tanzania (n = 350)
0%
20%
40%
60%
fraction outside, per sample
b
each country in the reference
grey bar spans that country's own studies
median
4.4%
11.5%
20.6%
3.7%
2.0%
9.8%
MDG
MNG
IND
KAZ
JPN
IRL
FJI
SWE
ESP
ISR
FRA
CHN
NLD
DNK
DEU
GBR
USA
0%
5%
10%
fraction outside, per sample
family
genus
species
c
the genera and gene families behind the shift
one dot per Hadza sample; open ring at left, some samples lack it
Genera
Treponema
Sarcina
Brachyspira
Anaerovibrio
Rikenella
Sanguibacteroides
Parasutterella
71% above
47% above
39% above
42% above
26% below
33% below
31% absent
Gene families
ATP-dependent DNA helicase RecG [EC:3.6.4.12]
lysozyme [EC:3.2.1.17]
signal peptidase I [EC:3.4.21.89]
aspartate kinase [EC:2.7.2.4]
tRNA 2-thiouridine synthesizing protein A [EC:2.8.1.-]
32% above
33% above
29% above
62% below
60% below
2.5
25
50
75
97.5
percentile of the reference population
below the range
above the range