

a
one cohort: 56 C. difficile patients, one dot per patient
bacterial families
Morganellaceae
Lactobacillaceae
Streptococcaceae
Enterococcaceae
Atopobiaceae
Barnesiellaceae
Prevotellaceae
Akkermansiaceae
Rikenellaceae
Eubacteriaceae
Butyricicoccaceae
Oscillospiraceae
5% above
12% above
11% above
11% above
59% absent
55% absent
27% absent
34% absent
34% absent
25% below
32% below
54% below
gene families
glycyl-tRNA synthetase beta chain
glycyl-tRNA synthetase
phosphoserine / homoserine phosphotransferase
oligosaccharide reducing-end xylanase
polyhydroxybutyrate depolymerase
41% above
36% above
52% absent
62% absent
48% absent
2.5
25
50
75
97.5
percentile of the reference population
below the range
within
above the range
absent from some patients
controls' median
b
nine colorectal cancer cohorts: one line per cohort, cases against its own controls
label: median shift in percentile points, and the number of cohorts holding it
taxa
Eggerthellaceae
Fusicatenibacter
Oscillospiraceae
Bifidobacteriaceae
Lachnospiraceae
Lachnospira
–23 in 3
–25 in 3
–33 in 3
–29 in 3
–35 in 4
–25 in 3
pathways
pyruvate fermentation to acetate and lactate II
pentose phosphate pathway (non-oxidative branch)
thiamin formation from pyrithiamine and oxythiamine (yeast)
methylerythritol phosphate pathway II
S-adenosyl-L-methionine cycle I
myo-, chiro- and scillo-inositol degradation
–30 in 3
–28 in 3
–46 in 3
–29 in 3
–39 in 3
–32 in 3
2.5
25
50
75
97.5
median percentile of the reference population
controls' median
cases' median, below controls
cases' median, above controls