

(a) One cohort: 56 C. difficile patients, one dot per patient
diamond, the 14 controls at their median; open ring at left, the feature is absent from some pa
Bacterial families
Morganellaceae
Lactobacillaceae
Streptococcaceae
Enterococcaceae
Atopobiaceae
Barnesiellaceae
Prevotellaceae
Akkermansiaceae
Rikenellaceae
Eubacteriaceae
Butyricicoccaceae
Oscillospiraceae
5% above
12% above
11% above
11% above
59% absent
55% absent
27% absent
34% absent
34% absent
25% below
32% below
54% below
Gene families
glycyl-tRNA synthetase beta chain [EC:6.1.1.14]
glycyl-tRNA synthetase [EC:6.1.1.14]
phosphoserine / homoserine phosphotransferase [EC:3.1.3.3 2.7.1.39]
oligosaccharide reducing-end xylanase [EC:3.2.1.156]
polyhydroxybutyrate depolymerase
41% above
36% above
52% absent
62% absent
48% absent
2.5
25
50
75
97.5
percentile of the reference population
below the range
above the range
(b) Nine colorectal cancer cohorts: each cohort's median, cases against its own contr
diamond, a cohort's controls; dot, its cases; label, the median shift and the cohorts it holds in
Taxa
Eggerthellaceae
Fusicatenibacter
Oscillospiraceae
Bifidobacteriaceae
Lachnospiraceae
Lachnospira
–23 in 3
–25 in 3
–33 in 3
–29 in 3
–35 in 4
–25 in 3
Function
pyruvate fermentation to acetate and lactate II
pentose phosphate pathway (non-oxidative branch)
thiamin formation from pyrithiamine and oxythiamine (yeast)
methylerythritol phosphate pathway II
S-adenosyl-L-methionine cycle I
myo-, chiro- and scillo-inositol degradation
–30 in 3
–28 in 3
–46 in 3
–29 in 3
–39 in 3
–32 in 3
2.5
25
50
75
97.5
median percentile of the reference population