

a
share of each sample outside the range (5 % expected)
bacterial families
other healthy adults (28 studies, n = 1,809)
US participants of the Hadza study (n = 12)
Hadza, Tanzania (n = 350)
median
4.4%
11.5%
20.6%
gene families (KOs)
other healthy adults (28 studies, n = 1,809)
US participants of the Hadza study (n = 12)
Hadza, Tanzania (n = 350)
3.7%
2.0%
9.8%
0%
20%
40%
60%
features outside the reference range, per sample
b
every population the baseline is built from
grey bar spans that country's own studies
MDG
MNG
IND
KAZ
JPN
IRL
FJI
SWE
ESP
ISR
FRA
CHN
NLD
DNK
DEU
GBR
USA
0%
5%
10%
outside the range, per sample
family
genus
species
c
the genera and gene families behind the shift
one dot per Hadza sample; open ring at left, some samples lack it
Treponema
Sarcina
Brachyspira
Anaerovibrio
Rikenella
Sanguibacteroides
Parasutterella
71% above
47% above
39% above
42% above
26% below
33% below
31% absent
Common gut genera
ATP-dependent DNA helicase RecG [EC:3.6.4.12]
lysozyme [EC:3.2.1.17]
signal peptidase I [EC:3.4.21.89]
aspartate kinase [EC:2.7.2.4]
tRNA 2-thiouridine synthesizing protein A [EC:2.8.1.-]
32% above
33% above
29% above
62% below
60% below
Gene families
2.5
25
50
75
97.5
percentile of the reference population
below the range
above the range