



Problem R-07J (C₁₀H₂₃NO) Interpret the ¹H NMR spectrum and determine the structure of **R-07J**.

- (a) DBE:____
- (b) The singlet at δ 4.05 disappears on addition of D₂O. What does this tell you about the structure?
- (c) Suggest part structures for the signals at δ 0.9 and δ 1.0.
- (d) Do a complete interpretation of the 6-proton multiplet between δ 2.2 and δ 2.7. Measure all couplings and chemical shifts, report them in the standard format, and assign them to the structure reported in part (e). Draw a "coupling tree" on the spectrum. You may use first order analysis if applicable.



(e) Draw the structure of **R-07J** below. If more than one structure fits the data, draw them, but circle your best choice.

(f) Are there features of the NMR spectrum which provide some hints as to the conformation of R-07J?




10

Problem R-07J (C₁₀H₂₃NO) Interpret the ¹H NMR spectrum and determine the structure of **R-07J**.

- (a) DBE: 0 1
 - (b) The singlet at δ 4.05 disappears on addition of D₂O. What does this tell you about the structure?
- This is probably an OH (or possibly NH) signal 2
 - (c) Suggest part structures for the signals at δ 0.9 and δ 1.0.

The 9-proton singlet at δ 0.90 is very likely a t-butyl group -C(CH₃)₃

- The 6-proton triplet is probably the CH₃ of a double ethyl group (also possible: diastereotopic CH(CH₃)₂ 4 group with accidental coincidence of middle peaks.
 - (d) Do a complete interpretation of the 6-proton multiplet between δ 2.2 and δ 2.7. Measure all couplings and chemical shifts, report them in the standard format, and assign them to the structure reported in part (e). Draw a "coupling tree" on the spectrum. You may use first order analysis if applicable.



(e) Draw the structure of R-07J below. If more than one structure fits the data, draw them, but circle your best choice.


OH δ calc CH 4.0 (obs 3.2) δ calc CH₂ 2.8 (obs 2.4) Σ Δδ: 1.2

HO δ calc CH 2.9 (obs 3.2) δ calc CH₂ 3.65 (obs 2.4) $\Sigma \Delta \delta$: 1.55

"Nutty compound" (aminal) ΗÒ δ calc CH 5.2 (obs 3.2) δ calc CH₂ 1.55 (obs 2.4) $\Sigma \Delta \delta$: 2.85

Hz

- (f) Are there features of the NMR spectrum which provide some hints as to the conformation of R-07J?
- The large difference in the two ³J couplings suggests 3 a strongly favored conformation - not surprising with a t-butyl group as anchor



7