

# Getting Started with Bering10K Level 2 & 3 indices

K. Holsman and K. Aydin (Tutorial), A. Hermann, K. Kearney, W. Cheng, I. Ortiz (Bering10K)

## Contents

<b>1. Overview</b>	<b>1</b>
1.1. Resources . . . . .	2
1.2 Guidelines for use and citation of the data . . . . .	2
<b>2. Installation</b>	<b>3</b>
2.1 Minimal Install . . . . .	3
2.2 Full install . . . . .	3
2.3 Set up environment and get shapefiles (full install) . . . . .	4
<b>3. Get ROMSNPZ data</b>	<b>5</b>
3.1 Available data sources . . . . .	5
3.2 Access and save the data . . . . .	6
<b>4. Explore indices &amp; plot the data</b>	<b>12</b>
4.1. A quick intro to Level 2 and 3 data . . . . .	12
4.2. Level 3 indices: . . . . .	12
<b>5. Hindcasts:</b>	<b>15</b>
5.1. Level 3 hindcasts . . . . .	15
<b>6. Projections:</b>	<b>29</b>
Table 1: Summary of ROMSNPZ downscaled model runs . . . . .	30
6.1. Level 3 projections . . . . .	30
6.2. Level 2 projections . . . . .	38
<b>7. Funding and acknowledgments (needs updating):</b>	<b>41</b>
7.1 Acknowledgements suggestion: projections . . . . .	41
<b>8. Helpful links and further reading:</b>	<b>42</b>
8.1 Citations for GCMs and carbon scenarios: . . . . .	42
8.2 Weblinks for further reading: . . . . .	42
8.3 Additional information on Hindcast and Projection Models (needs updating) . . . . .	42

## 1. Overview

This repository contains R code and Rdata files for working with netcdf-format data generated from the **downscaled ROMSNPZ modeling** of the ROMSNPZ Bering Sea Ocean Modeling team; Drs. Hermann, Cheng, Kearney, Pilcher, Ortiz, and Aydin. The code and R resources described in this tutorial are publicly available through the **ACLIM2 github repository** maintained by Kirstin Holsman as part of NOAA's **ACLIM project** for the Bering Sea. See *Hollowed et al. 2020* for more information about the **ACLIM project**



Figure 1: The ACLIM Repository [github.com/kholsman/ACLIM2](https://github.com/kholsman/ACLIM2) is maintained by **Kirstin Holsman**, Alaska Fisheries Science Center, NOAA Fisheries, Seattle WA. Multiple programs and projects have supported the production and sharing of the suite of Bering10K hindcasts and projections. *Last updated: Mar 09, 2021*

## 1.1. Resources

We **strongly recommend** reviewing the following documentation before using the data in order to understand the origin of the indices and their present level of skill and validation, which varies considerably across indices and in space and time:

- **The Bering10K Dataset documentation:** A pdf describing the dataset, including full model descriptions, inputs for specific results, and a tutorial for working directly with the ROMS native grid (Level 1 outputs).
- **Bering10K Simulation Variables:** A spreadsheet listing all simulations and the archived output variables associated with each, updated periodically as new simulations are run or new variables are made available.
- A **collection** of Bering10K ROMSNPZ model documentation (including the above files) is maintained by Kelly Kearney and will be regularly updated with new documentation and publications.

## 1.2 Guidelines for use and citation of the data

The data described here are published and publicly available for use, except as explicitly noted. However, for novel uses of the data, it is **strongly recommended** that you consult with and consider including at least one author from the ROMSNPZ team (Drs. Hermann, Cheng, Kearney, Pilcher, Aydin, Ortiz). There are multiple spatial and temporal caveats that are best described in discussions with the authors of these data and inclusion as co-authors will facilitate appropriate application and interpretation.

### 1.2.1. The Bering 10K Model (v. H16) with 10 depth layers:

The H16 model is the original BSIERP era 10 depth layer model with a 10 Km grid. This version was used in ACLIM1.0 to dynamically downscaled 3 global scale general circulation models (GCMs) under two CMIP (Coupled Model Intercomparison Project) phase 5 representative carbon pathways (RCP): RCP 4.5 or “moderate global carbon mitigation” and RCP 8.5 “high baseline global carbon emissions”. Details of the model and projections can be found in:

- **Hindcast (1979-2012; updated to 2018 during ACLIM 1.0):**

Hermann, A. J., G. A. Gibson, N. A. Bond, E. N. Curchitser, K. Hedstrom, W. Cheng, M. Wang, E. D. Cokelet, P. J. Stabeno, and K. Aydin. 2016. Projected future biophysical states of the Bering Sea. Deep Sea Research Part II: Topical Studies in Oceanography 134:30–47.doi:10.1016/j.dsr2.2015.11.001

- **Projections of the H16 10 layer model using CMIP5 scenarios:**

Hermann, A. J., G. A. Gibson, W. Cheng, I. Ortiz, K. Aydin, M. Wang, A. B. Hollowed, K. K. Holsman, and S. Sathyendranath. 2019. Projected biophysical conditions of the Bering Sea to 2100 under multiple emission scenarios. ICES Journal of Marine Science 76:1280–1304.doi:10.1093/icesjms/fsz043)

### 1.2.2. The Bering 10K Model (v. K20) with 30 depth layers and other advancements:

The Bering10K model was subsequently updated by Kearney et al. 2020 (30 layer and other NPZ updates) and Pilcher et al .2019 (OA and O<sub>2</sub> dynamics) and this version is used for the projections in ACLIM2.0 under the most recent CMIP phase 6.

- **Hindcast (1979-2020 hindcast with OA dynamics used in ACLIM 2.0):**

Kearney, K., A. Hermann, W. Cheng, I. Ortiz, and K. Aydin. 2020. A coupled pelagic-benthic-sympagic biogeochemical model for the Bering Sea: documentation and validation of the BESTNPZ model (v2019.08.23) within a high-resolution regional ocean model. Geoscientific Model Development 13:597–650.

Pilcher, D. J., D. M. Naiman, J. N. Cross, A. J. Hermann, S. A. Siedlecki, G. A. Gibson, and J. T. Mathis. 2019. Modeled Effect of Coastal Biogeochemical Processes, Climate Variability, and Ocean Acidification on Aragonite Saturation State in the Bering Sea. Frontiers in Marine Science 5:1–18.

- **Projections of the K20 30 layer model using CMIP6 scenarios:**

Hermann et al. in prep

Cheng et al. in prep

Kearney et al. in prep

Pilcher et al. in prep (CMIP5 K20 projections) (ACLIM indices avail by permission only)

## 2. Installation

### 2.1 Minimal Install

A minimal R install (for example XXX only) requires installing the `ncdf4`, `devtools` libraries (available on CRAN), and `thredds` R library through its github site:

```
install.packages(devtools)
install.packages(ncdf4)
devtools::install_github("bocinsky/thredds")
```

Note that each of these has multiple sub-dependent libraries and may take several minutes to install. *The full install below includes installation of these packages, so you don't need to perform this step if you perform the full install.*

### 2.2 Full install

The full install consists of the full directory structure in the ACLIM2 Repo; this includes a substantial set of resource files including shape files and data for performing Bering Sea spatial analysis in R. This will eventually become a library package, but currently requires manual downloading of the full directory structure from github. The full install may take up to **1GB of disk space** (initial download ~12MB).

### Option 1: Clone the repository

If you have git installed and can work with it, this is the preferred method as it preserves all directory structure and can aid in future updating. Use this from a **terminal command line, not in R**, to clone the full ACLIM2 directory and subdirectories:

```
git clone https://github.com/kholsman/ACLIM2.git
```

### Option 2: Download the repository

Download the full zip archive directly from the **ACLIM2 Repo** using this link: <https://github.com/kholsman/ACLIM2> and unzip its contents while preserving directory structure. **Important:** if downloading from zip, please **rename the root folder** from ACLIM2-main (in the zipfile) to ACLIM2 (name used in cloned copies) after unzipping, for consistency in the following examples.

### Option 3: Use R to download the repository

This set of commands, run within R, downloads the ACLIM2 repository and unpacks it, with the ACLIM2 directory structure being located in the specified `download_path`. This also performs the folder renaming mentioned in Option 2.

```
# Specify the download directory
main_nm      <- "ACLIM2"

# Note: Edit download_path for preference
download_path <- path.expand("~/desktop")
dest_fldr     <- file.path(download_path,main_nm)

url          <- "https://github.com/kholsman/ACLIM2/archive/main.zip"
dest_file    <- file.path(download_path,paste0(main_nm,".zip"))
download.file(url=url, destfile=dest_file)

# unzip the .zip file
setwd(download_path)
unzip(dest_file, exdir = "./",overwrite = T)

#rename the unzipped folder from ACLIM2-main to ACLIM2
file.rename(paste0(main_nm,"-main"), main_nm)
setwd(main_nm)
```

## 2.3 Set up environment and get shapefiles (full install)

The remainder of this tutorial was tested in RStudio. This may work in “plain” R, but is untested. If you are using RStudio, open `ACLIM2.Rproj` in Rstudio. If using R, use `setwd()` to get to the main ACLIM2 directory. Then run:

```
# -----
# SETUP WORKSPACE
# rm(list=ls())
tmstp  <- format(Sys.time(), "%Y_%m_%d")
main   <- getwd()  #~/GitHub_new/ACLIM2
source("R/make.R")
```

```

source("R/sub_scripts/load_maps.R") # skip this for faster load
# -----

```

The R/make.R command will install missing libraries (including those listed under the minimal install) and download and process multiple shapefiles for geographic analysis, it takes several minutes depending on bandwidth.

## 3. Get ROMSNPZ data

### 3.1 Available data sources

There are presently two sources of ROMSNPZ level 2 and level 3 post-processed datasets:

1) Public web-based ACLIM CMIP5 datasets

- **Level1** : (Empty; data not copied from Mox)
- **Level2** : (full grid, rotated to lat lon from the native ROMSNPZ grid, weekly averages)
  - **Bottom 5m** : subset of variables from the bottom 5 m of the water column
  - **Bottom 5m** : subset of variables for the surface 5 m of the water column
  - **Integrated**: watercolumn integrated averages or totals for various variables
  - **Level3**: two post-processed datasets
    - \* **ACLIMsurveyrep-x.nc**.: Survey replicated (variables “sampled” at the average location and date that each groundfish survey is sampled)(*Note that the resampling stations need to be removed before creating bottom temperature maps*)
    - **ACLIMregion-xnc**.: weekly variables averaged for each survey strata (*Note that area (km2) weighting should be used to combine values across multiple strata*)

For all files the general naming convention of the folders is: B10K-[ROMSNPZ version]\_[CMIP]\_[GCM]\_[carbon scenario]. For example, the CMIP5 set of indices was downscaled using the H16 (Hermann et al. 2016) version of the ROMSNPZ. Three models were used to force boundary conditions( MIROC, CESM, and GFDL) under 2 carbon scenarios RCP 8.5 and RCP 4.5. So to see an individual trajectory we might look in the level3 (timeseries indices) folder under B10K-H16\_CMIP5\_CESM\_rcp45, which would be the B10K version H16 of the CMIP5 CESM model under RCP4.5.

#### 3.1.1 Option 1: Public web-based data (hindcasts & CMIP5 projections)

The public web-based data (hindcasts & CMIP5 projections) option is available for Level3 and Level2 CMIP5 public data, it is not yet available for the embargoed CMIP6 data but through ACLIM2.0 will eventually be used to host that as well.

The ROMSNPZ team has been working with Roland Schweitzer and Peggy Sullivan to develop the ACLIM Live Access Server (LAS) to publicly host the published CMIP5 hindcasts and downscaled projections. This server is in beta testing phase and can be accessed at the following links:

- LAS custom ROMSNPZ data exploration, query, mapping, and plotting tool
- ERDAPP ACLIM data access tool
- THREDDS ACLIM direct access to Level 2 and 3 netcdf files

## 3.2 Access and save the data

The below code will extract variables from the Level 2 and Level 3 netcdf files (.nc) and save them as compressed .Rdata files on your local Data/in/Newest/Rdata folder.

### 3.2.1 Setup up the R worksace

First let's get the workspace set up, will we step through an example downloading the hindcast and a single projection (CMIP5 MIROC rcp8.5) but you can loop the code below to download the full set of CMIP5 projections.

```
# -----
# SETUP WORKSPACE
# rm(list=ls())
tmstp <- format(Sys.time(), "%Y_%m_%d")
main <- getwd() #~/GitHub_new/ACLIM
source("R/make.R")
# -----
```

Let's take a look at the available online datasets:

```
# preview the datasets on the server:
url_list <- tds_list_datasets(thredds_url = ACLIM_data_url)

#display the full set of datasets:
cat(paste(url_list$dataset, "\n"))

## Constants/
## B1OK-H16_CMIP5_CESM_B10_rcp85/
## B1OK-H16_CMIP5_CESM_rcp45/
## B1OK-H16_CMIP5_CESM_rcp85/
## B1OK-H16_CMIP5_GFDL_B10_rcp85/
## B1OK-H16_CMIP5_GFDL_rcp45/
## B1OK-H16_CMIP5_GFDL_rcp85/
## B1OK-H16_CMIP5_MIROC_rcp45/
## B1OK-H16_CMIP5_MIROC_rcp85/
## B1OK-H16_CORECFS/
## B1OK-K20_CORECFS/
## files/
```

### 3.2.2 Download Level 2 data

First we will explore the Level 2 bottom temperature data on the ACLIM Thredds server using the H16 hindcast and the H16 (CMIP5) projection for MIROC under rcp8.5. The first step is to get the data urls:

```
# define the simulation to download:
cmip <- "CMIP5"      # Coupled Model Intercomparison Phase
GCM  <- "MIROC"       # Global Circulation Model
rcp  <- "rcp85"        # future carbon scenario
mod  <- "B1OK-H16"     # ROMSNPZ model
hind <- "CORECFS"      # Hindcast

# define the projection simulation:
proj  <- paste0(mod, "_", cmip, "_", GCM, "_", rcp)
hind  <- paste0(mod, "_", hind)
```

```

# get the url for the projection and hindcast datasets:
proj_url      <- url_list[url_list$dataset == paste0(proj, "/"),]$path
hind_url      <- url_list[url_list$dataset == paste0(hind, "/"),]$path

# preview the projection and hindcast data and data catalogs (Level 1, 2, and 3):
proj_datasets <- tds_list_datasets(thredds_url = proj_url)
hind_datasets  <- tds_list_datasets(thredds_url = hind_url)

# get url for the projection and hindcast Level 2 and Level 3 catalogs
proj_12_cat    <- proj_datasets[proj_datasets$dataset == "Level 2/",]$path
proj_13_cat    <- proj_datasets[proj_datasets$dataset == "Level 3/",]$path
hind_12_cat    <- hind_datasets[hind_datasets$dataset == "Level 2/",]$path
hind_13_cat    <- hind_datasets[hind_datasets$dataset == "Level 3/",]$path
hind_12_cat

```

```
## [1] "https://data.pmel.noaa.gov/aclim/thredds/B10K-H16_CORECFS/Level2.html"
```

Now that we have the URLs let's take a look at the available Level2 datasets (currently temperature only, other variables available by request to Kelly Kearney:

- Bottom 5m : bottom water temperature at 5 meters
- Surface 5m : surface water temperature in the first 5 meters
- Integrated : Integrated water column averages for various NPZ variables

```

# preview the projection and hindcast Level 2 datasets:
proj_12_datasets <- tds_list_datasets(proj_12_cat)
hind_12_datasets <- tds_list_datasets(hind_12_cat)
proj_12_datasets$dataset

## [1] "Bottom 5m" "Surface 5m" "Integrated"

# get url for bottom temperature:
proj_12_BT_url   <- proj_12_datasets[proj_12_datasets$dataset == "Bottom 5m",]$path
hind_12_BT_url   <- hind_12_datasets[hind_12_datasets$dataset == "Bottom 5m",]$path
proj_12_BT_url

```

```
## [1] "https://data.pmel.noaa.gov/aclim/thredds/B10K-H16_CMIP5_MIROC_rcp85/Level2.html?dataset=B10K-H16_CMIP5_MIROC_rcp85_Bottom_5m"
```

We can't preview the Level 3 datasets in the same way but they are identical to those in the google drive and include two datasets

- ACLIMsurveyrep\_B10K-H16\_CMIP5\_CESM\_BIO\_rcp85.nc : NMFS Groundfish summer NBS and EBS survey replicated values for 60+ variables
- ACLIMregion\_B10K-H16\_CMIP5\_CESM\_BIO\_rcp85.nc : weekly strata averages for 60+ variables

```
weekly_vars # list of possible variables in the ACLIMregion_ files
```

```

## [1] "region_area"           "Ben"                  "DetBen"
## [4] "Hsbl"                 "IceNH4"               "IceN03"
## [7] "IcePhL"                "aice"                 "hice"
## [10] "shflux"                "ssflux"                "Cop_integrated"
## [13] "Cop_surface5m"         "Eup0_integrated"     "Eup0_surface5m"
## [16] "EupS_integrated"        "EupS_surface5m"      "Iron_bottom5m"
## [19] "Iron_integrated"        "Iron_surface5m"       "Jel_integrated"
## [22] "Jel_surface5m"         "MZL_integrated"       "MZL_surface5m"
## [25] "NCa0_integrated"        "NCa0_surface5m"       "NCaS_integrated"
## [28] "NCaS_surface5m"         "NH4_bottom5m"         "NH4_integrated"
## [31] "NH4_surface5m"          "NO3_bottom5m"         "NO3_integrated"

```

```

## [34] "N03_surface5m"      "PhL_integrated"      "PhL_surface5m"
## [37] "PhS_integrated"      "PhS_surface5m"      "prod_Cop_integrated"
## [40] "prod_Eup0_integrated" "prod_EupS_integrated" "prod_Eup_integrated"
## [43] "prod_Jel_integrated"  "prod_MZL_integrated" "prod_NCa0_integrated"
## [46] "prod_NCaS_integrated" "prod_NCa_integrated"  "prod_PhL_integrated"
## [49] "prod_PhS_integrated"  "salt_surface5m"     "temp_bottom5m"
## [52] "temp_integrated"      "temp_surface5m"     "uEast_bottom5m"
## [55] "uEast_surface5m"      "vNorth_bottom5m"    "vNorth_surface5m"
## [58] "fracbelow0"           "fracbelow1"          "fracbelow2"

```

Now we can download a subset of the Level2 data (full 10KM Lat Lon re-gridded data), here with an example of sampling on Aug 1 of each year:

```

# Currently available Level 2 variables
dl      <- proj_12_datasets$dataset # datasets

# variable list
svl <- list(
  'Bottom 5m' = "temp",
  'Surface 5m' = "temp",
  'Integrated' = c("EupS", "Cop", "NCaS") )

# preview the variables, timesteps, and lat lon in each dataset:
l2_info <- scan_l2(ds_list = dl, sim_list = "B10K-H16_CORECFS" )

names(l2_info)
l2_info[["Bottom 5m"]]$vars
l2_info[["Surface 5m"]]$vars
l2_info[["Integrated"]]$vars
max(l2_info[["Integrated"]]$time_steps)
l2_info[["Integrated"]]$years

# Simulation list:
# --> --> Tinker: add additional projection scenarios here
sl <- c(hind, proj)

# Currently available Level 2 variables
dl      <- proj_12_datasets$dataset # datasets

# variables to pull from each data set
# --> --> Tinker: try subbing in other Integrated variables
# (l2_info[["Integrated"]]$vars) into the third list vector
svl <- list(
  'Bottom 5m' = "temp",
  'Surface 5m' = "temp",
  'Integrated' = c("EupS", "Cop", "NCaS") )

# Let's sample the model years as close to Aug 1 as the model timesteps run:
# --> --> Tinker - try a different date
tr      <- c("-08-1 12:00:00 GMT")

# grab nc files from the aclim server and convert to rdatafiles with the ID Aug1

```

```

get_l2(
  ID          = "_Aug1",
  ds_list     = dl,
  trIN        = tr,
  sub_varlist = svl,
  sim_list    = sl )

```

### 3.2.3 Download Level 3 data

Now let's grab some of the Level 3 data and store it in the Data/in/Newest/Rdata folder. This is comparatively faster because Level 3 files are already post-processed to be in the ACLIM indices format and are relatively small:

```

# Simulation list:
# --> --> Tinker:add additional projection scenarios here
sl <- c(hind, proj)

# variable list
# --> --> Tinker:add additional variables to varlist
vl <- c(
  "temp_bottom5m",      # bottom temperature,
  "NCaS_integrated",   # Large Cop
  "Cop_integrated",    # Small Cop
  "EupS_integrated")   # Shelf euphausiids

# convert nc files into a long data.frame for each variable
# three options are:
# -----
# opt 1: access nc files remotely (fast, less local storage needed)
get_l3(web_nc = TRUE, download_nc = F,
       varlist = vl, sim_list = sl)

# opt 2: download nc files then access locally:
get_l3(web_nc = TRUE, download_nc = T,
       local_path = file.path(local_f1, "aclim_thredds"),
       varlist = vl, sim_list = sl)

# opt 3: access existing nc files locally:
get_l3(web_nc = F, download_nc = F,
       local_path = file.path(local_f1, "aclim_thredds"),
       varlist = vl, sim_list = sl)

```

### 3.2.4 ACLIM only: Convert CMIP6 (embargoed) Level 3 .nc -> .Rdata

Public CMIP5 and embargoed CMIP6 Level 3 netcdf (.nc) files are saved in the shared ACLIM data folder (note: Level 2 files are too large for the google drive but are available by request from Kelly Kearney).

**IMPORTANT** Please note that while the CMIP5 set is now public (Hermann et al. 2019; section 2.2) **the CMIP6 suite is under embargo for QAQC and should not be shared outside of the ACLIM group.** The ROMSNPZ team (Drs. Hermann, Cheng, Kearney, Pilcher, Adyin) are in the process of synthesizing and publishing the CMIP6 data (goal is spring 2021 for submission), following those publications the data will be made accessible to the public via the PMEL data portal, as is the case for the CMIP5 data

and public hindcasts. The ROMSNPZ team has made these runs available to ACLIM2 members in order to accelerate coupling to biological and social and economic models, thus out of professional courtesy please do not publish the data without permission from all ROMSNPZ team members, it is strongly advised that some or multiple ROMSNPZ team members be included as co-authors to ensure proper application and use of the ROMSNPZ data.

For most applications you can use the ACLIM level3 post-processed indices available on the shared ACLIM drive in the root google drive data folder: **00\_ACLIM\_shared>02\_DATA**.

The **Newest** folder is organized by Bering10K version, General Circulation Model (GCM) and carbon scenario, e.g. B10K-H16\_CMIP5\_CESM\_rcp45. Within each folder the following subfolders are:

ACLIM google drive CMIP6 datasets (embargoed; *ACLIM2 Only*) \* **Level1**: (Empty; data not copied from Mox) \* **Level2**: (Empty; data not copied from Mox) \* **Level3**: two post-processed datasets \* **ACLIMsurveyrep-x.nc**: Survey replicated (variables “sampled” at the average location and date that each groundfish survey is sampled) (*Note that the resampling stations need to be removed before creating bottom temperature maps*)

\* **ACLIMregion-xnc**: weekly variables averaged for each survey strata (*Note that area (km<sup>2</sup>) weighting should be used to combine values across multiple strata*)

For all files the general naming convention of the folders is: B10K-[ROMSNPZ version]\_[CMIP]\_[GCM]\_[carbon scenario]. For example, the CMIP5 set of indices was downscaled using the H16 (Hermann et al. 2016) version of the ROMSNPZ. Three models were used to force boundary conditions( MIROC, CESM, and GFDL) under 2 carbon scenarios RCP 8.5 and RCP 4.5. So to see an individual trajectory we might look in the level3 (timeseries indices) folder under B10K-H16\_CMIP5\_CESM\_rcp45, which would be the B10K version H16 of the CMIP5 CESM model under RCP4.5.

- 1) **ACLIMsurveyrep\_B10K-x.nc** contains summer groundfish trawl “survey replicated” indices (using mean date and lat lon) (*Note that the resampling stations need to be removed before creating bottom temperature maps*)
- 2) **ACLIMregion\_B10K-x.nc**: contains weekly “strata” values (*Note that area (km<sup>2</sup>) weighting should be used to combine values across multiple strata*)

There are two folders that need to be copied into the ACLIM2 folder on your computer under ‘~[YOURPATH]/ACLIM2/Data/in/’:

- 1) **00\_ACLIM\_shared>02\_DATA>Newest**. This folder contains a folder called **roms\_for\_aclim** with all the ACLIM Level3 indices for model simulations available to ACLIM members.
- 2) **00\_ACLIM\_shared>02\_DATA>Map\_layers.zip**. This file needs to be unzipped after you download it to your local folder. It contains (large) base maps for the code below including **shp\_files** and **geo\_tif** folders.

Now let’s convert the CMIP6 Level3 .nc files to .Rdata files (as in section 3.2.3)

```
# preview the available CMIP6 data
dir(file.path(local_f1,"roms_for_aclim"))

# variable list
vl      <- c(
  "temp_bottom5m",
  "NCaS_integrated", # Large Cop
  "Cop_integrated", # Small Cop
  "EupS_integrated") # Euphausiids

# define the simulation to download:
cmip <- "CMIP6"      # Coupled Model Intercomparison Phase
GCM   <- "MIROC"       # Global Circulation Model
```




Figure 2: Your local ACLIM2/Data directory should look something like this when you are done downloading the data and unzipping it.




Figure 3: The final folder structure on your local drive in Data/in/Newest should look something like this.

```

rcp  <- "ssp585"      # future carbon scenario
mod  <- "B10K-K20P19"  # ROMSNPZ model
hind <- "CORECFS"       # Hindcast

# define the projection simulation:
proj  <- paste0(mod,"_",cmip,"_",GCM,"_",rcp)
hind  <- paste0("B10K-K20","_",hind)
sl    <- c(hind, proj)

# opt 3: convert subset of nc files to rdata files for analysis:
get_l3(web_nc = F, download_nc = F,
       local_path = file.path(local_f1,"roms_for_aclim"),
       varlist = vl,sim_list = sl)

```

## 4. Explore indices & plot the data

### 4.1. A quick intro to Level 2 and 3 data

KERIM's text and example here

### 4.2. Level 3 indices:

Level 3 indices can be used to generate seasonal, monthly, and annual indices (like those reported in Reum et al. 2020), Holsman et al. 2020). In the section below we explore these indices in more detail using R, including using (2) above to generate weekly, monthly, and seasonal indices (e.g. Fall Zooplankton) for use in biological models. In section 3 below we explore these indices in more detail using R, including using (2) above to generate weekly, monthly, and seasonal indices (e.g. Fall Zooplankton) for use in biological models. The following examples show how to analyze and plot the ACLIM indices from the .Rdata files created in the previous step 3.

Please be sure to coordinate with ROMSNPZ modeling team members to ensure data is applied appropriately. If you need access to the raw ROMSNPZ files (netcdf, non-regridded large files located on MOX). Please contact **Al Hermann** or **Kelly Kearney**. Please note that while the CMIP5 set is now public (Hermann et al. 2019) **the CMIP6 suite is under embargo for QAQC and should not be shared outside of the ACLIM group**. See Section 1 above for more detail.

#### 4.1.1 Explore Level 3 data catalog

Once the base files and setup are loaded you can explore the index types. Recall that in each scenario folder there are two indices saved within the Level3 subfolders:

- 1) **ACLIMsurveyrep\_B10K-x.nc** contains summer groundfish trawl “survey replicated” indices (using mean date and lat lon) (*Note that the resampling stations need to be removed before creating bottom temperature maps*)
- 2) **ACLIMregion\_B10K-x.nc**: contains weekly “strata” values (*Note that area weighting should be used to combine values across multiple strata*)

First run the below set of code to set up the workspace:

```

# -----
# SETUP WORKSPACE
tmstp  <- format(Sys.time(), "%Y_%m_%d")
main   <- getwd()  #~/GitHub_new/ACLIM2
source("R/make.R")
# -----


# list of the scenario x GCM downscaled ACLIM indices
for(k in aclim)
  cat(paste(k, "\n"))

embargoed # not yet public or published
public    # published runs (CMIP5)

# get some info about a scenario:
all_info1
all_info2

# variables in each of the two files:
srvy_vars
weekly_vars


#summary tables for variables
srvy_var_def
weekly_var_def

# explore stations in the survey replicated data:
head(station_info)

```

#### 4.1.2 Level 3: Spatial indices (survey replicated)

Let's start by exploring the survey replicated values for each variable. Steps 2 and 3 generated the Rdata files that are stored in the `ACLIMsurveyrep_B10K-[version_CMIPx_GCM_RCP].Rdata` in each corresponding simulation folder.



The code segment below will recreate the above figures. Note that if this is the first time through it may take 3-5 mins to load the spatial packages and download the files from the web (first time through only).

```
# if load_gis is set to FALSE in R/setup.R (default)
# we will need to load the gis layers and packages
# if this is the first time through this would be a good time
# to grab a coffee...

source("R/sub_scripts/load_maps.R")

# first convert the station_info object into a shapefile for mapping:
station_sf      <- convert2shp(station_info)
station_sf$stratum <- factor(station_sf$stratum)

# plot the stations:
p <- plot_stations_basemap(sfIN = station_sf,
                           fillIN = "subregion",
                           colorIN = "subregion") +
  scale_color_viridis_d(begin = .2,end=.6) +
  scale_fill_viridis_d(begin = .2,end=.6)

if(update.figs){
  p
  ggsave(file=file.path(main,"Figs/stations_NS.jpg"),width=5,height=5)
}

p2 <- plot_stations_basemap(sfIN = station_sf,fillIN = "stratum",colorIN = "stratum") +
  scale_color_viridis_d() +
  scale_fill_viridis_d()

if(update.figs){
  p2
  ggsave(file=file.path(main,"Figs/stations.jpg"),width=5,height=5)}
```

## 5. Hindcasts:

There are two model versions of hindcasts available for comparison. The Hermann et al. 2016 H16 10 depth layer model and the Kearney et al. 2020 30 depth layer model. Both are resolved spatially at a ~10km grid cell.

### 5.1. Level 3 hindcasts

Level 3 hindcast products include survey replicated station data and strata averaged weekly values. The code below will explore these in more detail.

#### 5.1.1. Level 3 hindcasts: spatial patterns

Now let's explore the survey replicated data in more detail and use to plot bottom temperature.

```
# run this line if load_gis is set to F in R/setup.R:
source("R/sub_scripts/load_maps.R")

# preview the l3 data for the hindcast:
tt <- all_info1%>%filter(name == "B10K-K20_CORECFS")
tt <- seq(as.numeric(substring(tt$Start,1,4)),as.numeric(substring(tt$End,1,4)),10)

# now create plots of average BT during four time periods
time_seg <- list(
  '1970-1980' = c(1970:1980),
  '1980-1990' = c(1980:1990),
  '1990-2000' = c(1990:2000),
  '2000-2010' = c(2000:2010),
  '2010-2020' = c(2010:2020))

# lists the possible variables
srvy_vars # lists the possible variables

# specify the variables to plot
vl <- c(
  "temp_bottom5m",
  "NCaS_integrated", # Large Cop
  "Cop_integrated", # Small Cop
  "EupS_integrated") # Euphausiids

# assign the simulation to download
# --> Tinker: try selecting a different set of models to compare
sim <- "B10K-K20_CORECFS"

# open a "region" or strata specific nc file
fl <- file.path(sim,paste0(srvy_txt,sim,".Rdata"))

# create local rdata files (opt 1)
if(!file.exists(file.path(Rdata_path,fl)))
  get_l3(web_nc = TRUE, download_nc = F,
         varlist = vl,sim_list = sim )

# load object 'ACLIMsurveyrep'
```

```

load(file.path(main,Rdata_path,f1))

# Collate mean values across timeperiods and simulations
# -----
ms <- c("B1OK-H16_CORECFS","B1OK-K20_CORECFS" )

# Loop over model set
for(sim in ms){
  f1           <- file.path(sim,paste0(srvy_txt,sim,".Rdata"))

  if(!file.exists( file.path(Rdata_path,f1)) )
    get_l3(web_nc = TRUE, download_nc = F,
            varlist = vl,sim_list =sim )
}

# get the mean values for the time blocks from the rdata versions
# will throw "implicit NA" errors that can be ignored
mn_var_all <- get_mn_rd(modset = ms,
                         names  = c("H16","K20") ,
                         varUSE = "temp_bottom5m")
# --> Tinker:           varUSE = "EupS_integrated")

# convert results to a shapefile
mn_var_sf  <- convert2shp(mn_var_all%>%filter(!is.na(mnval)))
lab_t       <- "Bering1OK CORECFS hindcast"


p_hind_3      <- plot_stations_basemap(sfIN = mn_var_sf,
                                         fillIN = "mnval",
                                         colorIN = "mnval",
                                         sizeIN=.3) +
  facet_grid(simulation~time_period)+
  scale_color_viridis_c()+
  scale_fill_viridis_c()+
  guides(
    color =  guide_legend(title="Bottom T (degC)" ),
    fill  =  guide_legend(title="Bottom T (degC)") ) +
  ggtitle(lab_t)

# This is slow but it works (repeat dev.new() twice if in Rstudio)...
dev.new()
p_hind_3

if(update.figs)
  ggsave(file=file.path(main,"Figs/mn_hindcast_BT.jpg"),width=8,height=6)

```

## Bering10K CORECFS hindcast



Now let's look at the Marine Heatwave conditions in 2018 and compare that to the average conditions prior to 2010:

```
# now create plots of average BT during four time periods
time_seg    <- list( '1970-2010' = c(1970:2010),
                     '2018-2018' = c(2018:2018))

# assign the simulation to download
sim         <- "B10K-K20_CORECFS"

# open a "region" or strata specific nc file
fl          <- file.path(sim,paste0(srwy_txt,sim,".Rdata"))

# load object 'ACLIMsurveyrep'
load(file.path(main,Rdata_path,fl))

# get the mean values for the time blocks from the rdata versions
mn_var_all <- get_mn_rd(modset = "B10K-K20_CORECFS",
                         varUSE = "temp_bottom5m")

# convert results to a shapefile
mn_var_sf   <- convert2shp(mn_var_all%>%filter(!is.na(mnval)))
lab_t       <- "Bering10K CORECFS hindcast"

p_mhw      <- plot_stations_basemap(sfIN = mn_var_sf,
```

```

      fillIN = "mnval",
      colorIN = "mnval",
      sizeIN=.3) +
facet_grid(simulation~time_period)+
scale_color_viridis_c()+
scale_fill_viridis_c()+
guides(
  color = guide_legend(title="Bottom T (degC)"),
  fill = guide_legend(title="Bottom T (degC))") +
ggtitle(lab_t)

# This is slow but it works (repeat dev.new() twice if in Rstudio)...
dev.new(width=4,height=3)
p_mhw

if(update.figs)
  ggsave(file=file.path(main,"Figs/mn_hindcast_mhw.jpg"),width=4,height=3)

```




Figure 4: Decadal averages of bottom temperature from the two hindcast models.

### 5.1.2. Level 3 hindcasts: Weekly strata averages

The next set of indices to will explore are the weekly strata-specific values for each variable. These are stored in the `ACLIMregion_B10K-[version_CMIPx_GCM_RCP].nc` in each scenario folder.

```

# View an individual variable (e.g., Bottom Temp)
# -----
weekly_vars

# assign the simulation to download
sim      <- "B10K-K20_CORECFS"

# define a "region" or strata specific nc file
fl       <- file.path(sim,paste0(reg_txt,sim,".Rdata"))

vl      <- c(
  "temp_bottom5m",
  "NCaS_integrated", # Large Cop
  "Cop_integrated", # Small Cop
  "EupS_integrated") # Euphausiids

# create local rdata files (opt 1)
if(!file.exists(file.path(Rdata_path,fl)))
  get_l3(web_nc = TRUE, download_nc = F,
         varlist = vl,sim_list = sim)

# load object 'ACLIMregion' for bottom temperature
load(file.path(main,Rdata_path,fl))
tmp_var   <- ACLIMregion%>%filter(var == "temp_bottom5m")

# now plot the data:
p4 Hind <- ggplot(data = tmp_var) +
  geom_line(aes(x=time,y=val,color= strata),alpha=.8)+ 
  facet_grid(basin~.)+
  ylab(tmp_var$units[1])+ 
  ggtitle( paste(sim,tmp_var$var[1]))+
  theme_minimal()
p4 Hind

if(update.figs)
  ggsave(file=file.path(main,"Figs/hind_weekly_bystrata.jpg"),width=8,height=5)

# To get the average value for a set of strata, weight the val by the area:
mn_NEBS <- getAVGnSUM(strataIN = NEBS_strata, dataIN = tmp_var)
mn_NEBS$basin = "NEBS"
mn_SEBS <-getAVGnSUM(strataIN = SEBS_strata, dataIN = tmp_var)
mn_SEBS$basin = "SEBS"

p5 Hind <- ggplot(data = rbind(mn_NEBS,mn_SEBS)) +
  geom_line(aes(x=time,y=mn_val,color=basin),alpha=.8)+ 
  geom_smooth(aes(x=time,y=mn_val,color=basin),

```

```

        formula = y ~ x, se = T) +
facet_grid(basin~.) +
scale_color_viridis_d(begin=.4,end=.8) +
ylab(tmp_var$units[1]) +
ggtitle( paste(sim,mn_NEBS$var[1])) +
theme_minimal()
p5_hind
if(update.figs)
ggsave(file=file.path(main,"Figs/hind_weekly_byreg.jpg"),width=8,height=5)

```




Figure 5: Weekly indices by sub-region

### 5.1.3. Level 3 hindcasts: Seasonal averages

Now using a similar approach get the monthly mean values for a variable:

```

# assign the simulation to download
sim      <- "B10K-K20_CORECFS"

# Set up seasons (this follows Holsman et al. 2020)
seasons <- data.frame(mo = 1:12,
                      season = factor("", 
                                      levels=c("Winter","Spring","Summer","Fall")))
seasons$season[1:3]    <- "Winter"
seasons$season[4:6]    <- "Spring"
seasons$season[7:9]    <- "Summer"
seasons$season[10:12]  <- "Fall"

```




Figure 6: Weekly indices by sub-region

```

vl <- c(
    "temp_bottom5m",
    "NCaS_integrated", # Large Cop
    "Cop_integrated", # Small Cop
    "EupS_integrated") # Euphausiids

# create local rdata files (opt 1)
if(!file.exists(file.path(Rdata_path,f1)))
  get_l3(web_nc = TRUE, download_nc = F,
         varlist = vl,sim_list = sim)

# open a "region" or strata specific file
f1      <- file.path(sim,paste0(reg_txt,sim,".Rdata"))
load(file.path(main,Rdata_path,f1))

# get large zooplankton as the sum of euph and NCaS
tmp_var   <- ACLIMregion%>%
  filter(var%in%vl[c(2,3)])%>%
  group_by(time,strata,strata_area_km2,basin)%>%
  group_by(time,
           strata,
           strata_area_km2,
           basin,
           units)%>%
  summarise(val =sum(val))%>%
  mutate(var       = "Zoop_integrated",
        long_name ="Total On-shelf

```

```

    large zooplankton concentration,
    integrated over depth (NCa, Eup)")

rm(ACLIMregion)
head(tmp_var)

# define some columns for year mo and julian day
tmp_var$yr      <- strftime(as.Date(tmp_var$time),
                           format="%Y-%m-%d")$year + 1900
tmp_var$mo       <- strftime(as.Date(tmp_var$time),
                           format="%Y-%m-%d")$mon  + 1
tmp_var$jday    <- strftime(as.Date(tmp_var$time),
                           format="%Y-%m-%d")$yday + 1
tmp_var$season  <- seasons[tmp_var$mo,2]

# To get the average value for a set of strata, weight the val by the area: (slow...)
mn_NEBS_season <- getAVGnSUM(
  strataIN = NEBS_strata,
  dataIN = tmp_var,
  tblock=c("yr","season"))
mn_NEBS_season$basin = "NEBS"


mn_SEBS_season <- getAVGnSUM(
  strataIN = SEBS_strata,
  dataIN = tmp_var,
  tblock=c("yr","season"))
mn_SEBS_season$basin = "SEBS"

plot_data       <- rbind(mn_NEBS_season,mn_SEBS_season)

# plot Fall values:
p6 Hind <- ggplot(data = plot_data%>%filter(season=="Fall")) +
  geom_line(aes(x = yr,y = mn_val,color=basin),alpha=.8) +
  geom_smooth(aes(x = yr,y = mn_val,color=basin),
              formula = y ~ x, se = T) +
  facet_grid(basin~.) +
  scale_color_viridis_d(begin=.4,end=.8) +
  ylab(tmp_var$units[1]) +
  ggtitle(paste(sim,"Fall",mn_NEBS_season$var[1])) +
  theme_minimal()
p6 Hind

if(update.figs)
  ggsave(file=file.path(main,"Figs/Hind_Fall_large_Zoop.jpg"),width=8,height=5)

```



### 5.1.4. Level 3 hind-

casts: Monthly averages

Using the same approach we can get monthly averages for a given variable:


```
# To get the average value for a set of strata, weight the val by the area: (slow...)
mn_NEBS_season <- getAVGnSUM(
  strataIN = NEBS_strata,
  dataIN   = tmp_var,
  tblock   = c("yr", "mo"))
mn_NEBS_season$basin = "NEBS"

mn_SEBS_season <- getAVGnSUM(
  strataIN = SEBS_strata,
  dataIN   = tmp_var,
  tblock=c("yr", "mo"))
mn_SEBS_season$basin = "SEBS"

plot_data      <- rbind(mn_NEBS_season,mn_SEBS_season)

# plot Fall values:
p7_hind <- ggplot(data = plot_data%>%filter(mo==9) ) +
  geom_line( aes(x = yr,y = mn_val,color=basin),alpha=.8)+ 
  geom_smooth( aes(x = yr,y = mn_val,color=basin),
               formula = y ~ x, se = T)+ 
  facet_grid(basin~.)+
  scale_color_viridis_d(begin=.4,end=.8)+ 
  ylab(tmp_var$units[1])+ 
  ggtitle( paste(aclim[2],"Sept.",mn_NEBS_season$var[1]))+ 
  theme_minimal()

if(update.figs)
  ggsave(file=file.path(main,"Figs/Hind_Sept_large_Zoop.jpg"),width=8,height=5)
```



## ## 5.2. Level 2 hindcasts

Level 2 data can be explored in the same way as the above indices but we will focus in the section below on a simple spatial plot and temporal index. The advantage of Level2 indices is in the spatial resolution and values outside of the survey area.

### 5.2.1. Level 2 hindcasts: Custom spatial indices

As we did in section 5.1.1. let's create spatial plots of hindcast time periods for Aug 1 of each year:

```
# run this line if load_gis is set to F in R/setup.R:
source("R/sub_scripts/load_maps.R")

# now create plots of average BT during four time periods
time_seg <- list( '1970-1980' = c(1970:1980),
                   '1980-1990' = c(1980:1990),
                   '1990-2000' = c(1990:2000),
                   '2000-2010' = c(2000:2010),
                   '2010-2020' = c(2010:2020))

# preview the datasets on the server:
tds_list_datasets(thredds_url = ACLIM_data_url)

# assign the simulation to download
# --> Tinker: try selecting a different set of models to compare
sim <- "B10K-K20_CORECFS"
#ms <- c("B10K-H16_CORECFS", "B10K-K20_CORECFS" )

# Currently available Level 2 variables
dl <- proj_l2_datasets$dataset # datasets

svl <- list(
  'Bottom 5m' = "temp",
  'Surface 5m' = "temp",
  'Integrated' = c("EupS", "Cop", "NCaS") )

# Let's sample the model years as close to Aug 1 as the model timesteps run:
```

```

tr           <- c("-08-1 12:00:00 GMT")

# the full grid is large and takes a longtime to plot, so let's subsample the grid every 4 cells

IDin        <- "_Aug1_subgrid"
var_use     <- "_bottom5m_temp"

# open a "region" or strata specific nc file
fl          <- file.path(main,Rdata_path,sim,"Level2",
                         paste0(sim,var_use,IDin,".Rdata"))

# load data from level 2 nc files (approx <10sec)
startTime = Sys.time()
if(!file.exists(file.path(Rdata_path,fl))){
  get_l2(
    ID          = "_1990_subgrid",
    overwrite   = T,
    xi_rangeIN = seq(1,182,10),
    eta_rangeIN = seq(1,258,10),
    ds_list     = dl[1], # must be same length as sub_varlist
    trIN        = tr,
    yearsIN    = 1990,
    sub_varlist = list('Bottom 5m' = "temp" ),
    sim_list    = sim  )
}
endTime   = Sys.time()
endTime - startTime

# load data from level 2 nc files for all years and vars (yearsIN = NULL by default)
#       NOTE: THIS IS SLOWWWWW..~ 2 min
startTime2 = Sys.time()
if(!file.exists(file.path(Rdata_path,fl))){
  get_l2(
    ID          = IDin,
    overwrite   = T,
    xi_rangeIN = seq(1,182,10),
    eta_rangeIN = seq(1,258,10),
    ds_list     = dl,
    trIN        = tr,
    sub_varlist = svl,
    sim_list    = sim  )
}
endTime2  = Sys.time()
endTime2 - startTime2

# load R data file
load(f1)  # temp

# there are smarter ways to do this; looping because
# we don't want to mess it up but this is slow...
i <-1
data_long <- data.frame(latitude = as.vector(temp$lat),
                         longitude = as.vector(temp$lon),

```

```

    val = as.vector(temp$val[,,i]),
    time = temp$time[i],
    year = substr( temp$time[i],1,4),stringsAsFactors = F
  )

for(i in 2:dim(temp$val)[3])
  data_long <- rbind(data_long,
    data.frame(latitude = as.vector(temp$lat),
    longitude = as.vector(temp$lon),
    val = as.vector(temp$val[,,i]),
    time = temp$time[i],
    year = substr( temp$time[i],1,4),stringsAsFactors = F)
  )

# get the mean values for the time blocks from the rdata versions
# may throw "implicit NA" errors that can be ignored
tmp_var <-data_long # get mean var val for each time segment
j<-0
for(i in 1:length(time_seg)){
  if(length( which(as.numeric(tmp_var$year)%in%time_seg[[i]]) ))>0){
    j <- j +1
    mn_tmp_var <- tmp_var%>%
      filter(year%in%time_seg[[i]],!is.na(val))%>%
      group_by(latitude, longitude)%>%
      summarise(mnval = mean(val,rm.na=T))

    mn_tmp_var$time_period <- factor(names(time_seg)[i],levels=names(time_seg))

    if(j == 1) mn_var <- mn_tmp_var
    if(j > 1) mn_var <- rbind(mn_var,mn_tmp_var)
    rm(mn_tmp_var)
  }
}

# convert results to a shapefile
L2_sf <- convert2shp(mn_var%>%filter(!is.na(mnval)))

p9 Hind     <- plot_stations_basemap(sfIN = L2_sf,
                                         fillIN = "mnval",
                                         colorIN = "mnval",
                                         sizeIN=.6) +
  #facet_wrap(~time_period,nrow=2,ncol=3) +
  facet_grid(~time_period) +
  scale_color_viridis_c() +
  scale_fill_viridis_c() +
  guides(
    color = guide_legend(title="Bottom T (degC)"),
    fill = guide_legend(title="Bottom T (degC)")) +
  ggtitle(paste(sim,var_use,IDin))

# This is slow but it works (repeat dev.new() twice if in Rstudio)...
dev.new()

```

```

p9 Hind

if(update.figs)  ggsave(file=file.path(main,"Figs/Hind_sub_grid_mn_BT_Aug1.jpg"),width=8,height=4)

# graphics.off()

```




Figure 7: Aug 1 Bottom temperature from Level 2 dataset

### 5.2.2. Level 2 hindcasts: M2 mooring comparison

As final hindcast comparison, let's look a surface temperature from observations vs the H16 and K20 model versions of the hindcast:

```

# M2_lat <- (56.87°N, -164.06°W)
# 56.8777    -164.06 xi = 99      eta= 62
IDin       <- "_2013_M2"
var_use     <- "_surface5m_temp"

# get data from M2 data page:
pmelM2_url <-"https://www.ncei.noaa.gov/data/oceans/ncei/ocads/data/0157599/"
yr_dat      <- "M2_164W_57N_Apr2019_May2019.csv"
yr_dat      <- "M2_164W_57N_May2013_Sep2013.csv"

# preview the datasets on the server:
temp <- tempfile()
download.file(paste0(pmelm2_url, yr_dat), temp)
#M2data <- read.csv(temp, skip=4, stringsAsFactors = F)
M2data <- read.csv(temp, skip=0, stringsAsFactors = F)

unlink(temp)

```

```

# convert date and time to t
M2data$t <- as.POSIXct(paste0(M2data$Date, " ", M2data$Time, ":00"), "%m/%d/%Y %H:%M:%S",
                        origin = "1900-01-01 00:00:00",
                        tz = "GMT")

# open a "region" or strata specific nc file
f1           <- file.path(main,Rdata_path,sim,"Level2",
                           paste0(sim,var_use,IDin,".Rdata"))

# assign the simulation to download
sim         <- "B10K-K20_CORECFS"

# Let's sample the model years as close to Aug 1 as the model timesteps run:
#tr          <- c("-08-1 12:00:00 GMT")
tr          <- substring(M2data$t,5,20)
# the full grid is large and takes a longtime to plot, so let's subsample the grid every 4 cells

# load data from level 2 nc files (approx <10sec)
if(!file.exists(file.path(Rdata_path,f1))){
  get_l2(
    ID          = IDin,
    overwrite   = T,
    xi_rangeIN = 99,
    eta_rangeIN = 62,
    ds_list     = dl[2],  # must be same length as sub_varlist
    trIN        = tr,
    yearsIN     = 2013,
    sub_varlist = list('Surface 5m' = "temp" ),
    sim_list    = c("B10K-H16_CORECFS","B10K-K20_CORECFS" ) )
}

# load R data file
# open a "region" or strata specific nc file
sim <- "B10K-H16_CORECFS"
f1           <- file.path(main,Rdata_path,sim,"Level2",
                           paste0(sim,var_use,IDin,".Rdata"))
load(f1)    # temp

# there are smarter ways to do this; looping because
# we don't want to mess it up but this is slow...
i <-1
data_long <- data.frame(latitude = as.vector(temp$lat),
                         longitude = as.vector(temp$lon),
                         val = as.vector(temp$val[, , i]),
                         sim = sim,
                         time = temp$time[i],
                         year = substr( temp$time[i] ,1,4),stringsAsFactors = F
                         )

for(i in 2:dim(temp$val)[3])
  data_long <- rbind(data_long,
                      data.frame(latitude = as.vector(temp$lat),

```

```

longitude = as.vector(temp$lon),
val = as.vector(temp$val[,,i]),
sim = sim,
time = temp$time[i],
year = substr( temp$time[i],1,4),stringsAsFactors = F)
)

# open a "region" or strata specific nc file
sim <- "B10K-K20_CORECFS"
f12 <- file.path(main,Rdata_path,sim,"Level2",
paste0(sim,var_use,IDin, ".Rdata"))

load(f12) # temp
for(i in 1:dim(temp$val)[3])
  data_long <- rbind(data_long,
    data.frame(latitude = as.vector(temp$lat),
    longitude = as.vector(temp$lon),
    val = as.vector(temp$val[,,i]),
    sim = sim,
    time = temp$time[i],
    year = substr( temp$time[i],1,4),stringsAsFactors = F)
  )

plotM2_dat <- M2data%>%dplyr::select(SST = SST..C.,Date = t)
plotM2_dat$sim <- factor("Obs",levels=c("Obs","B10K-H16_CORECFS","B10K-K20_CORECFS"))
plotM2_dat <- plotM2_dat%>%filter(SST>-99)
plotroms_dat <- data_long%>%dplyr::select(SST = val,Date = time,sim)
plotroms_dat$sim <- factor(plotroms_dat$sim,levels=c("Obs","B10K-H16_CORECFS","B10K-K20_CORECFS"))
plotdat <- rbind(plotM2_dat,plotroms_dat)

p10 Hind <- ggplot(plotdat) +
  geom_line( aes(x=Date,y=SST,color=sim),alpha=.8)+ 
  # geom_smooth( aes(x = Date,y = SST,color=sim),
  # formula = y ~ x, se = T)+ 
  scale_color_viridis_d(begin=.9,end=.2)+ 
  ylab(tmp_var$units[1])+ 
  ggtitle( "Bering M2 Mooring: 2013 SST")+
  theme_minimal()

# This is slow but it works (repeat dev.new() twice if in Rstudio)...
dev.new()
p10 Hind

if(update.figs)
  ggsave(file=file.path(main,"Figs/Hind_M2_SST.jpg"),width=8,height=4)

# graphics.off()

```

## 6. Projections:

The ACLIM project utilizes the full “suite” of Bering10K model hindcasts and projections, summarized in the following table. These represent downscaled models hindcast and projections whereby boundary conditions of the high resolution Bering10K model are forced by the coarser resolution General Circulation Models (GCM) run under Coupled Model Intercomparison Project (CMIP) phase 5 (5th IPCC Assessment Report)

Bering M2 Mooring: 2013 SST




Figure 8: M2 mooring SST in 2013.

or phase 6 (6th IPCC Assessment Report; “AR”) global carbon mitigation scenarios. Hindcasts are similarly forced at the boundaries from global scale climate reanalysis CORE and CFS products (see sections 1-5). For full details see the Kearney 2021 Tech. Memo.

**Table 1: Summary of ROMSNPZ downscaled model runs**

CMIP	GCM	Scenario	Def	Years	Model	Source	Status
5	CORECFS	Reanalysis	Hindcast	1970 - 2018	H16	IEA/ACLIM	Public
	CORECFS	Reanalysis	Hindcast	1970 - 2020	K20	MAPP/IEA/ACLIM	Public
	GFDL	RCP 4.5	Med. mitigation	2006 - 2099	H16	ACLIM/FATE	Public
	GFDL	RCP 8.5	High baseline	2006 - 2099	H16	ACLIM/FATE	Public
	GFDL	RCP 8.5bio*	High baseline	2006 - 2099	H16	ACLIM/FATE	Public
	MIROC	RCP 4.5	Med. mitigation	2006 - 2099	H16	ACLIM/FATE	Public
	MIROC	RCP 8.5	High baseline	2006 - 2099	H16	ACLIM/FATE	Public
	CESM	RCP 4.5	Med. mitigation	2006 - 2099	H16	ACLIM/FATE	Public
	CESM	RCP 8.5	High baseline	2006 - 2080	H16	ACLIM/FATE	Public
	CESM	RCP 8.5bio*	High baseline	2006 - 2099	H16	ACLIM/FATE	Public
6	CESM	SSP585	High baseline	2014 - 2099	K20P19	ACLIM2/RTAP	Embargo
6	CESM	SSP126	High Mitigation	2014 - 2099	K20P19	ACLIM2/RTAP	Embargo
6	CESM	Historical	Historical	1980 - 2014	K20P19	ACLIM2/RTAP	Embargo
6	GFDL	SSP585	High baseline	2014 - 2099	K20P19	ACLIM2/RTAP	Embargo
6	GFDL	SSP126	High Mitigation	2014 - 2099	K20P19	ACLIM2/RTAP	Embargo
6	GFDL	Historical	Historical	1980 - 2014	K20P19	ACLIM2/RTAP	Embargo
6	MIROC	SSP585	High baseline	2014 - 2099	K20P19	ACLIM2/RTAP	Embargo
6	MIROC	SSP126	High Mitigation	2014 - 2099	K20P19	ACLIM2/RTAP	Embargo
6	MIROC	Historical	Historical	1980 - 2014	K20P19	ACLIM2/RTAP	Embargo

\*“bio” = nutrient forcing on boundary conditions

## 6.1. Level 3 projections

some text here

### 6.1.1. Level 3 projections: spatial patterns

```

    "NCaS_integrated", # Large Cop
    "Cop_integrated", # Small Cop
    "EupS_integrated") # Euphausiids

# View possible simulations:
head(aclim)

# assign the simulation to download
# --> Tinker: try selecting a different set of models to compare
sim      <- "B10K-H16_CMIP5_MIROC_rcp85"

# open a "region" or strata specific nc file
fl       <- file.path(sim,paste0(srwy_txt,sim,".Rdata"))

# load object 'ACLIMsurveyrep'
load(file.path(main,Rdata_path,fl))

# create local rdata files
if(!file.exists(file.path(Rdata_path,fl)))
  get_l3(web_nc = TRUE, download_nc = F,
         varlist = vl,sim_list =sim )

# Collate mean values across timeperiods and simulations
# -----
m_set     <- c(9,7,8)
ms        <- aclim[m_set]

# Loop over model set
for(sim in ms){
  fl       <- file.path(sim,paste0(srwy_txt,sim,".Rdata"))

  # download & convert .nc files that are not already in Rdata folder
  if(!file.exists( file.path(Rdata_path,fl)) )
    get_l3(web_nc = TRUE, download_nc = F,
           varlist = vl,sim_list =sim )
}

# get the mean values for the time blocks from the rdata versions
# will throw "implicit NA" errors that can be ignored
mn_var_all <- get_mn_rd(modset = ms ,varUSE="temp_bottom5m")

# convert results to a shapefile
mn_var_sf  <- convert2shp(mn_var_all%>%filter(!is.na(mnval)))
lab_t      <- ms[2]%>%stringr::str_remove("(^-]")

p3         <- plot_stations_basemap(sfIN = mn_var_sf,
                                         fillIN = "mnval",
                                         colorIN = "mnval",
                                         sizeIN=.3) +
  facet_grid(simulation~time_period) +
  scale_color_viridis_c() +
  scale_fill_viridis_c()

```

```

guides(
  color = guide_legend(title="Bottom T (degC")),
  fill = guide_legend(title="Bottom T (degC))") +
ggttitle(lab_t)

# This is slow but it works (repeat dev.new() twice if in Rstudio)...
dev.new()
p3

if(update.figs) ggsave(file=file.path(main,"Figs/mn_BT.jpg"),width=8,height=6)

# graphics.off()

```




Figure 9: Bottom temperature projections under differing SSP126 (top row) and SSP585 (bottom row)

### 6.1.2. Level 3 projections: Weekly strata averages

The next set of indices to will explore are the weekly strata-specific values for each variable. These are stored in the `ACLIMregion_B10K-[version_CMIPx_GCM_RCP].nc` in each scenario folder.

```

# View an individual variable (e.g., Bottom Temp)
# -----
weekly_vars
aclim
sim      <-"B10K-H16_CMIP5_MIROC_rcp85"

```

```

# open a "region" or strata specific nc file
f1      <- file.path(sim,paste0(reg_txt,sim,".Rdata"))

var_use   <- "temp_bottom5m"
vl       <- c(
    "temp_bottom5m",
    "NCaS_integrated", # Large Cop
    "Cop_integrated", # Small Cop
    "EupS_integrated") # Euphausiids

# create local rdata files (opt 1)
if(!file.exists(file.path(Rdata_path,f1)))
  get_l3(web_nc = TRUE, download_nc = F,
         varlist = vl,sim_list = sim)

# load object 'ACLIMregion'
load(file.path(main,Rdata_path,f1))
tmp_var   <- ACLIMregion%>%filter(var == var_use)

# now plot the data:

p4 <- ggplot(data = tmp_var) +
  geom_line(aes(x=time,y=val,color= strata),alpha=.8)+ 
  facet_grid(basin~.)+
  ylab(tmp_var$units[1])+ 
  ggtitle( paste(sim,tmp_var$var[1]))+
  theme_minimal()
p4
if(update.fig)  ggsave(file=file.path(main,"Figs/weekly_bystrata.jpg"),width=8,height=5)

# To get the average value for a set of strata, weight the val by the area:
mn_NEBS <- getAVGnSUM(strataIN = NEBS_strata, dataIN = tmp_var)
mn_NEBS$basin = "NEBS"
mn_SEBS <-getAVGnSUM(strataIN = SEBS_strata, dataIN = tmp_var)
mn_SEBS$basin = "SEBS"

p5 <- ggplot(data = rbind(mn_NEBS,mn_SEBS)) +
  geom_line(aes(x=time,y=mn_val,color=basin),alpha=.8)+ 
  geom_smooth(aes(x=time,y=mn_val,color=basin),
              formula = y ~ x, se = T)+ 
  facet_grid(basin~.)+
  scale_color_viridis_d(begin=.4,end=.8)+ 
  ylab(tmp_var$units[1])+ 
  ggtitle( paste(sim,mn_NEBS$var[1]))+
  theme_minimal()
p5
if(update.fig)  ggsave(file=file.path(main,"Figs/weekly_byreg.jpg"),width=8,height=5)

```





Figure 10: Weekly indices by sub-region



### 6.1.3. Level 3 projections: Seasonal averages

Now using a similar approach get the monthly mean values for a variable:

```
sim <- "B10K-H16_CMIP5_MIROC_rcp85"
# Set up seasons (this follows Holsman et al. 2020)
```

```

seasons <- data.frame(mo = 1:12,
                      season = factor("", 
                                      levels=c("Winter", "Spring", "Summer", "Fall")))
seasons$season[1:3]   <- "Winter"
seasons$season[4:6]   <- "Spring"
seasons$season[7:9]   <- "Summer"
seasons$season[10:12] <- "Fall"

vl <- c(
    "temp_bottom5m",
    "NCaS_integrated", # Large Cop
    "Cop_integrated", # Small Cop
    "EupS_integrated") # Euphausiids

# create local rdata files (opt 1)
if(!file.exists(file.path(Rdata_path,f1)))
  get_l3(web_nc = TRUE, download_nc = F,
         varlist = vl,sim_list = sim)

# open a "region" or strata specific file
f1      <- file.path(sim,paste0(reg_txt,sim,".Rdata"))
load(file.path(main,Rdata_path,f1))

# get large zooplankton as the sum of euph and NCaS
tmp_var   <- ACLIMregion%>%
  filter(var%in%vl[c(2,3)])%>%
  group_by(time,strata,strata_area_km2,basin)%>%
  group_by(time,
            strata,
            strata_area_km2,
            basin,
            units)%>%
  summarise(val =sum(val))%>%
  mutate(var       = "Zoop_integrated",
         long_name ="Total On-shelf
large zooplankton concentration,
integrated over depth (NCa, Eup)")

rm(ACLIMregion)
head(tmp_var)

tmp_var$yr     <- strftime(as.Date(tmp_var$time),
                           format="%Y-%m-%d")$year + 1900
tmp_var$mo     <- strftime(as.Date(tmp_var$time),
                           format="%Y-%m-%d")$mon + 1
tmp_var$jday   <- strftime(as.Date(tmp_var$time),
                           format="%Y-%m-%d")$yday + 1
tmp_var$season <- seasons[tmp_var$mo,2]

# To get the average value for a set of strata, weight the val by the area: (slow...)
mn_NEBS_season <- getAVGnSUM(
  strataIN = NEBS_strata,

```

```

    dataIN = tmp_var,
    tblock=c("yr","season"))
mn_NEBS_season$basin = "NEBS"
mn_SEBS_season <- getAVGnSUM(
  strataIN = SEBS_strata,
  dataIN = tmp_var,
  tblock=c("yr","season"))
mn_SEBS_season$basin = "SEBS"

plot_data      <- rbind(mn_NEBS_season,mn_SEBS_season)

# plot Fall values:
p6 <- ggplot(data = plot_data%>%filter(season=="Fall") ) +
  geom_line(   aes(x = yr,y = mn_val,color=basin),alpha=.8)+ 
  geom_smooth( aes(x = yr,y = mn_val,color=basin),
                formula = y ~ x, se = T)+ 
  facet_grid(basin~.)+
  scale_color_viridis_d(begin=.4,end=.8)+ 
  ylab(tmp_var$units[1])+ 
  ggttitle( paste(sim,"Fall",mn_NEBS_season$var[1]))+ 
  theme_minimal()
p6

if(update.figs)
  ggsave(file=file.path(main,"Figs/Fall_large_Zoop.jpg"),width=8,height=5)

```



6.1.4. Level 3 Projections: Monthly averages Using the same approach we can get monthly averages for a given variable:

```

# To get the average value for a set of strata, weight the val by the area: (slow...)
mn_NEBS_season <- getAVGnSUM(

```

```

strataIN = NEBS_strata,
dataIN   = tmp_var,
tblock   = c("yr","mo"))
mn_NEBS_season$basin = "NEBS"
mn_SEBS_season <- getAVGnSUM(
  strataIN = SEBS_strata,
  dataIN = tmp_var,
  tblock=c("yr","mo"))
mn_SEBS_season$basin = "SEBS"

plot_data      <- rbind(mn_NEBS_season,mn_SEBS_season)

# plot Fall values:
p7 <- ggplot(data = plot_data%>%filter(mo==9) ) +
  geom_line( aes(x = yr,y = mn_val,color=basin),alpha=.8)+ 
  geom_smooth( aes(x = yr,y = mn_val,color=basin),
                formula = y ~ x, se = T)+ 
  facet_grid(basin~.)+
  scale_color_viridis_d(begin=.4,end=.8)+ 
  ylab(tmp_var$units[1])+ 
  ggtitle( paste(aclim[2],"Sept.",mn_NEBS_season$var[1]))+ 
  theme_minimal()

p7

if(update.figs)
  ggsave(file=file.path(main,"Figs/Sept_large_Zoop.jpg"),width=8,height=5)

```




Figure 11: September large zooplankton integrated concentration

Finally we can use this approach to plot the monthly averages and look for phenological shifts:

```

# or average in 4 time slices by mo:
# now create plots of average BT during four time periods
time_seg <- list( '2010-2020' = c(2010:2020),
                   '2021-2040' = c(2021:2040),
                   '2041-2060' = c(2041:2060),
                   '2061-2080' = c(2061:2080),
                   '2081-2099' = c(2081:2099))

plot_data$ts <- names(time_seg)[1]
for(tt in 1:length((time_seg)))
  plot_data$ts[plot_data$yr%in%(time_seg[[tt]][1]:time_seg[[tt]][2])]<-names(time_seg)[tt]

plot_data2 <- plot_data%>%
  group_by(var,mo,units,long_name,basin, ts)%>%
  summarize(mn_val2 = mean(mn_val))

# now plot phenological shift:
p8 <- ggplot(data = plot_data2 ) +
  geom_line( aes(x = mo,y = mn_val2,color=ts),alpha=.8,size=0) +
  geom_smooth( aes(x = mo,y = mn_val2,color=ts),
               formula = y ~ x, se = F) +
  facet_grid(basin~.) +
  scale_color_viridis_d(begin=.9,end=.2) +
  ylab(tmp_var$units[1]) +
  ggtitle( paste(aclim[2],mn_NEBS_season$var[1])) +
  theme_minimal()

p8
if(update.figs)
  ggsave(file=file.path(main,"Figs/PhenShift_large_Zoop.jpg"),width=8,height=5)

```

## 6.2. Level 2 projections

Level 2 data can be explored in the same way as the above indices but we will focus in the section below on a simple spatial plot and temporal index. The advantage of Level2 inidces is in the spatial resolution and values outside of the survey area.

### 6.2.1 Level 2 projections: Custom spatial indices

```

# define four time periods
time_seg <- list( '2010-2020' = c(2000:2020),
                   '2021-2040' = c(2021:2040),
                   '2041-2060' = c(2041:2060),
                   '2061-2080' = c(2061:2080),
                   '2081-2099' = c(2081:2099))

# View an individual variable (e.g., Bottom Temp)
# -----
head(srvy_vars)
head(aclim)

# assign the simulation to download

```




Figure 12: September large zooplankton integrated concentration

```

# --> --> Tinker: try selecting a different set of models to compare
sim      <- "B10K-H16_CMIP5_MIROC_rcp85"

svl <- list(
  'Bottom 5m' = "temp",
  'Surface 5m' = "temp",
  'Integrated' = c("EupS", "Cop", "NCaS") )

# Currently available Level 2 variables
dl       <- proj_l2_datasets$dataset # datasets

# Let's sample the model years as close to Aug 1 as the model timesteps run:
tr      <- c("-08-1 12:00:00 GMT")

# the full grid is large and takes a longtime to plot, so let's subsample the grid every 4 cells

IDin      <- "_Aug1_subgrid"
var_use   <- "_bottom5m_temp"

# open a "region" or strata specific nc file
fl        <- file.path(main,Rdata_path,sim,"Level2",
                      paste0(sim,var_use,IDin,".Rdata"))

# load object 'ACLIMsurveyrep'
if(!file.exists(file.path(Rdata_path,fl)))
  get_l2(
    ID          = IDin,

```

```

xi_rangeIN  = seq(1,182,10),
eta_rangeIN = seq(1,258,10),
ds_list      = dl,
trIN         = tr,
sub_varlist  = svl,
sim_list     = sim )

# load R data file
load(f1)    # temp

# there are smarter ways to do this; looping because
# we don't want to mess it up but this is slow...
i <-1
data_long <- data.frame(latitude = as.vector(temp$lat),
                         longitude = as.vector(temp$lon),
                         val = as.vector(temp$val[,,i]),
                         time = temp$time[i],
                         year = substr( temp$time[i],1,4),stringsAsFactors = F
                         )
for(i in 2:dim(temp$val)[3])
  data_long <- rbind(data_long,
                      data.frame(latitude = as.vector(temp$lat),
                                 longitude = as.vector(temp$lon),
                                 val = as.vector(temp$val[,,i]),
                                 time = temp$time[i],
                                 year = substr( temp$time[i],1,4),stringsAsFactors = F
                                 )
  )

# get the mean values for the time blocks from the rdata versions
# will throw "implicit NA" errors that can be ignored
tmp_var <-data_long # get mean var val for each time segment
j<-0
for(i in 1:length(time_seg)){
  if(length( which(as.numeric(tmp_var$year)%in%time_seg[[i]]) )>0){
    j <- j +1
    mn_tmp_var <- tmp_var%>%
      filter(year%in%time_seg[[i]],!is.na(val))%>%
      group_by(latitude, longitude)%>%
      summarise(mnval = mean(val,rm.na=T))

    mn_tmp_var$time_period = factor(names(time_seg)[i],levels=names(time_seg))
    if(j == 1) mn_var <- mn_tmp_var
    if(j > 1) mn_var <- rbind(mn_var,mn_tmp_var)
    rm(mn_tmp_var)
  }
}

# convert results to a shapefile
L2_sf  <- convert2shp(mn_var%>%filter(!is.na(mnval)))

p9      <- plot_stations_basemap(sfIN = L2_sf,
                                    fillIN = "mnval",

```

```


        colorIN = "mnval",
        sizeIN=.6) +
facet_grid(.~time_period)+
scale_color_viridis_c()+
scale_fill_viridis_c()+
guides(
  color = guide_legend(title="Bottom T (degC)", 
  fill = guide_legend(title="Bottom T (degC)")) +
ggtitle(paste(sim,var_use,IDin))

# This is slow but it works (repeat dev.new() twice if in Rstudio)...
dev.new()
p9

if(update.figs) ggsave(file=file.path(main,"Figs/sub_grid_mn_BT_Aug1.jpg"),width=8,height=6)

# graphics.off()

```



#### 6.2.2 Level 2 projections: Projections at M2 mooring

## 7. Funding and acknowledgments (needs updating):

### 7.1 Acknowledgements suggestion: projections

**PLEASE** Include a statement like the following one in your acknowledgements section:

*This study is part of NOAA's Alaska Climate Integrated Modeling project (ACLIM) and FATE project XXXX. We would like to thank the entire ACLIM team including [add specific names] for feedback and discussions on the broader application of this work. Multiple NOAA National Marine Fisheries programs provided support for ACLIM including Fisheries and the Environment (FATE), Stock Assessment Analytical Methods (SAAM) Science and Technology North Pacific Climate Regimes and Ecosystem Productivity, the Integrated Ecosystem Assessment Program (IEA), the NOAA Economics and Social Analysis Division, NOAA Research Transition Acceleration Program (RTAP), the Alaska Fisheries Science Center (ASFC), the Office of Oceanic and Atmospheric Research (OAR) and the National Marine Fisheries Service (NMFS). The scientific views, opinions, and conclusions expressed herein are solely those of the authors and do not represent the views, opinions, or conclusions of NOAA or the Department of Commerce.*

**For some of the integrated papers the following maybe should also be added:**

*Additionally, the International Council for the Exploration of the Sea (ICES) and the North Pacific Marine Science Organization (PICES) provided support for Strategic Initiative for the Study of Climate Impacts on Marine Ecosystems (SI-CCME) workshops, which facilitated development of the ideas presented in this paper. The scientific views, opinions, and conclusions expressed herein are solely those of the authors and do not represent the views, opinions, or conclusions of NOAA, the Department of Commerce, ICES, or PICES.*

## **8. Helpful links and further reading:**

### **8.1 Citations for GCMs and carbon scenarios:**

#### **CMIP3 (BSIERP global climate model runs):**

Meehl, G. A., C. Covey, T. Delworth, M. Latif, B. McAvaney, J. F. B. Mitchell, R. J. Stouffer, and K. E. Taylor, 2007: The WCRP CMIP3 multimodel dataset: A new era in climate change research. Bull. Amer. Meteor. Soc., 88, 1383–1394.

#### **CMIP5 (ACLIM global climate model runs):**

Taylor, K. E., R. J. Stouffer, and G. A. Meehl, 2012: An overview of CMIP5 and the experiment design. Bull. Amer. Meteor. Soc., 93, 485–498.

#### **CMIP6 and SSPs (ACLIM2 global climate model runs):**

ONeill, B. C., C. Tebaldi, D. P. van Vuuren, V. Eyring, P. Friedlingstein, G. Hurtt, R. Knutti, E. Kriegler, J.-F. Lamarque, J. Lowe, G. A. Meehl, R. Moss, K. Riahi, and B. M. Sanderson. 2016. The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. Geoscientific Model Development 9:3461–3482.

### **8.2 Weblinks for further reading:**

- Explore annual indices of downscaled projections for the EBS: **ACLIM indices**
- To view climate change projections from CMIP5 (eventually CMIP6):**ESRL climate change portal**

### **8.3 Additional information on Hindcast and Projection Models (needs updating)**

#### **CORE-CFSR (1976-2012)**

This is the hindcast for the Bering Sea and is a combination of the reconstructed climatology from the **CLIVAR** Co-ordinated Ocean-Ice Reference Experiments (CORE) Climate Model (1969-2006) the **NCEP** Climate Forecast System Reanalysis is a set of re-forecasts carried out by NOAA's National Center for Environmental Prediction (NCEP). See **CFS-R** for more info.

### **CCCMA(2006-2039; AR4 SRES A1B)**

Developed by the Canadian Centre for Climate Modelling and Analysis, this is also known as the CGCM3/T47 model. This model showed the greatest warming over time compared to other models tested by PMEL. See more data the [AOOS:CCCMA portal](#).

### **ECHOG(2006-2039; AR4 SRES A1B)**

The ECHO-G model from the Max Planck Institute in Germany This model showed the least warming over time compared to other models tested by PMEL. See more data the [AOOS:ECHO-G portal](#).

### **GFDL (2006-2100; AR5 RCP 4.5, 8.5, SSP126,SSP585)**

The NOAA Geophysical Fluid Dynamics Laboratory **GFDL** has lead development of the first Earth System Models (ESMs), which like physical climate models, are based on an atmospheric circulation model coupled with an oceanic circulation model, with representations of land, sea ice and iceberg dynamics; ESMs additionally incorporate interactive biogeochemistry, including the carbon cycle. The ESM2M model used in this project is an evolution of the prototype EMS2.1 model, where pressure-based vertical coordinates are used along the developmental path of GFDL's Modular Ocean Model version 4.1 and where the land model is more advanced (LM3) than in the previous ESM2.1

### **MIROC(2006-2039; AR4 SRES A1B; 2006-2100 RCP4.5, RCP8.5, SSP585, SSP126)**

The Model for Interdisciplinary Research on Climate (MIROC)-M model developed by a consortium of agencies in Japan []. Compared to other models tested by PMEL, MIROC-M was intermediate in degree of warming over the Bering Sea shelf for the first half of the 21st century. See more data the [AOOS:MIROC portal](#).