

Assignment: Asymptotic Notations and Correctness of Algorithms

1. **Identify and compare the order of growth:** Use the definitions of O , Ω and Θ to identify if the following statements are true or false. Prove your assertion. Draw a rough graph marking c and n_0 values if the statement is True. [You don't have to find the values of c and n_0].

- a. $n(n+1)/2 \in O(n^3)$ **True**

$$\lim_{n \rightarrow \infty} \frac{\frac{n(n+1)}{2}}{n^3}$$

$$= \lim_{n \rightarrow \infty} \frac{n^2+n}{2n^3}$$

$$= \lim_{n \rightarrow \infty} \frac{2n+1}{6n^2} \quad \text{Applying L'Hospital rule}$$

$$= \lim_{n \rightarrow \infty} \frac{2}{12n} \quad \text{Applying L'Hospital rule}$$

$$= 0$$

$$\Rightarrow f(n) \in O(g(n))$$

- b. $n(n+1)/2 \in \Theta(n^2)$ **True**

$$\lim_{n \rightarrow \infty} \frac{\frac{n(n+1)}{2}}{n^2}$$

$$= \lim_{n \rightarrow \infty} \frac{n^2+n}{2n^2}$$

$$= \lim_{n \rightarrow \infty} \frac{2n+1}{4n} \quad \text{Applying L'Hospital rule}$$

$$= \lim_{n \rightarrow \infty} \frac{2}{4} \quad \text{Applying L'Hospital rule}$$

$$= \text{Constant}$$

$$\Rightarrow f(n) \in \Theta(g(n))$$

- c. $10n-6 \in \Omega(78n+2020)$ **True**

$$\lim_{n \rightarrow \infty} \frac{10n-6}{78n+2020}$$

$$= \lim_{n \rightarrow \infty} \frac{10}{78} \quad \text{Applying L'Hospital rule}$$

= Constant

$\Rightarrow f(n) \in \Theta(g(n))$

Since Θ is stronger notation than Ω -notation. Θ -notation implies Ω -notation. Hence,

True

d. $n! \in \Omega(0.00001n)$ True

$$\lim_{n \rightarrow \infty} \frac{n!}{0.00001n}$$

$$= \lim_{n \rightarrow \infty} \frac{\sqrt{2\pi n} \left(\frac{n}{e}\right)^n}{0.00001n} \quad \text{using Stirling's Formula}$$

$$= \frac{\sqrt{2\pi}}{0.00001} \lim_{n \rightarrow \infty} \frac{\sqrt{n}}{n} \left(\frac{n}{e}\right)^n$$

$$= \frac{\sqrt{2\pi}}{0.00001} \lim_{n \rightarrow \infty} \frac{\sqrt{n}}{(e)^n} x \frac{n^n}{n}$$

$$= \frac{\sqrt{2\pi}}{0.00001} \lim_{n \rightarrow \infty} \frac{\sqrt{n}}{e^n} x^{n^{n-1}}$$

$$= (\text{something}) * (\infty)^\infty = \infty$$

$\Rightarrow f(n) \in \Omega(g(n))$

Or

$$\lim_{n \rightarrow \infty} \frac{n!}{0.00001n}$$

$$= \lim_{n \rightarrow \infty} \frac{n(n-1)(n-2) \dots}{0.00001n}$$

$$= \lim_{n \rightarrow \infty} \frac{(n-1)(n-2) \dots}{0.00001} \quad [\text{simplifying}]$$

$$= \infty$$

2. Read and Analyze Pseudocode: Consider the following algorithm

```
Classified(A...n-1):  
    minval = A[0]  
    maxval = A[0]  
    for i = 1 to n-1:  
        if A[i] < minval:  
            minval = A[i]
```

```

        if A[i] > maxval
            maxval = A[i]
    return maxval - minval

```

- What does this algorithm compute? **The difference between the largest and the smallest element in an array.**
- What is its basic operation (i.e. the line of code or operation that is executed maximum number of times)?

The if conditions

```

if A[i] < minval; if A[i] > maxval

```

- How many times is the basic operation executed?

$$\sum_{i=1}^{n-1} 2 = 2 \cdot (n - 1)$$

- What is the time complexity of this algorithm?

$\theta(n)$

3. Using mathematical induction prove below non-recursive algorithm:

```

def reverse_array(Arr):
    n = len(Arr)
    i = (n-1)//2
    j = n//2
    while(i >= 0 and j <= (n-1)):
        temp = Arr[i]
        Arr[i] = Arr[j]
        Arr[j] = temp
        i = i-1
        j = j+1

```

- Write the loop invariant of the reverse_array function.
Before the iteration of the while loop, for values i and j, Arr[i+1 : j-1] is reversed. That is values in the Arr from 'i+1' to 'j-1' are reversed.
- Prove correctness of reverse_array function using induction.

Loop invariant: For i and j, Arr[i+1 : j-1] is reversed

Initialization: Before the entry into the while loop $j-1 < i+1$, hence the part of the Arr[i+1 : j-1] does not contain any elements. The invariant is true by default.

Maintenance: At the start of k^{th} iteration, assume that Arr[i+1 : j-1] is reversed.

In the previous iteration i is decremented and j is incremented. In the k^{th} iteration, in the body of the loop Arr[i] and Arr[j] are swapped.

Thus at the start of $(k+1)^{\text{th}}$ iteration Arr[i : j] is reversed.

Termination: After the loop terminates, $i = -1$ and $j = n$.

Based on loop invariant Arr[i+1 : j-1] is reversed. Therefore, Arr[0:n-1], which is our entire array, is reversed.

