



The Dentistry Insight

CLINICAL EDUCATION SERIES

Australian Dental Therapeutic Guideline (TG4) **Topics: Detailed Explanations with Mock Questions**

Author: **Dr Hussain Ahmad**
Founder, The Dentistry Insight
www.thedentistryinsight.com

A note on this book: This resource has been prepared for educational purposes only, to help students and clinicians revise key topics from the Australian dental therapeutic guidelines in plain, easy-to-understand language, supported by original mock questions written by the author. Every effort has been made to ensure accuracy, but errors and omissions are possible; readers should always verify clinical information against the original source material published by the Australian Dental Association (ADA) before applying it in practice.

This publication is an independent educational work and is not affiliated with, endorsed by, or produced in association with the Australian Dental Association, the Australian Dental Council, or any dental academy or institute in Australia.

Chapter 1 — Infection Control: Standard Precautions and the Patient with HIV/AIDS

Infection Control: Standard Precautions (ADA/ADC Style)

The ADA Guidelines for Infection Prevention and Control (5th Edition) emphasize that Standard Precautions are the primary strategy for successful infection control in the dental office.

Cause & Pathogenesis in the Clinic

Cause of Cross-Infection: Transmission of blood-borne viruses (HIV, HBV, HCV) via percutaneous injury (needlesticks) or mucous membrane exposure.

Pathogenesis of Exposure: HIV is a fragile virus outside the body, but a deep sharps injury with contaminated blood poses a documented risk of seroconversion (approx. 0.3%).

ADA Standard Precautions Protocol

Hand Hygiene: The single most important measure. Perform before and after every patient contact and immediately after removing gloves.

Personal Protective Equipment (PPE):

Gloves: Single-use, non-sterile (for routine) or sterile (for surgery).

Masks & Eye Protection: Level 2 or 3 surgical masks and wrap-around goggles/shields to protect against splashes/splatter.

Gowns: Fluid-resistant, long-sleeved.

Sharps Management: Never recap needles using two hands (use the "single-handed scoop" or a mechanical device). Dispose of sharps at the point of use.

Environmental Cleaning: Use hospital-grade disinfectants for non-critical surfaces.

Step 1: Pre-Treatment Risk Assessment & Patient Management

No Segregation: The patient is treated in the same clinical environment as any other patient. No special room or "last appointment of the day" is required.

Medical History: A thorough, confidential medical history is taken. You must ask about HIV status, current CD4 count, viral load, and any antiretroviral therapy (ART). This is not for discrimination, but to assess for potential complications (e.g., bleeding tendencies from thrombocytopenia, risk of opportunistic infections, drug interactions).

Consent & Confidentiality: The patient's HIV status is protected by strict privacy laws. You do not disclose it to other staff unless they have a direct clinical need to know.

Step 2: Personal Protective Equipment (PPE) - The Non-Negotiable Barrier

This is where the "standard" part is most critical. For any procedure on any patient, the ADA mandates:

Gloves: Single-use, sterile or non-sterile examination gloves. Double-gloving is not a standard ADA recommendation for HIV alone, but is recommended for high-risk, prolonged, or surgical procedures (e.g., oral surgery, implant placement) where glove failure is more likely.

Protective Eyewear: Must be worn by the clinician, assistant, and patient. This protects against splash or aerosolized blood.

Mask: A surgical mask (Type IIR) is standard. For aerosol-generating procedures (AGPs) like using a high-speed handpiece or ultrasonic scaler, an N95/P2 respirator is recommended, especially if the patient has a high viral load or is not on effective ART.

Gown/Protective Clothing: A long-sleeved, fluid-resistant gown or clinic coat. It must be changed if it becomes visibly soiled or penetrated.

Step 3: Environmental & Surface Management

Surface Barriers: Use impervious plastic barriers on all high-touch surfaces (light handles, bracket table, x-ray unit, computer mouse). This is standard for all patients.

Cleaning & Disinfection: After the procedure, remove barriers. Then, clean and disinfect all surfaces with a hospital-grade disinfectant with a label claim against HIV (e.g., sodium hypochlorite 0.1-1%, or a TGA-approved disinfectant). HIV is an enveloped virus and is very susceptible to most disinfectants.

Spills: Blood or body fluid spills are managed with a 1:10 dilution of sodium hypochlorite (0.5% available chlorine) for a 10-minute contact time.

Step 4: Instrument Reprocessing - The Critical Loop

This is where many errors occur. The ADA follows the Spaulding Classification:

Critical Instruments (penetrate soft tissue/bone, e.g., scalpel, forceps, burs): Must be sterilized after each use. This means cleaning, then steam autoclaving at 121°C for 15 minutes or 134°C for 3-4 minutes. HIV is completely inactivated by standard autoclaving.

Semi-Critical Instruments (contact mucous membranes but not penetrate, e.g., mouth mirror, impression trays): Must be sterilized or, at minimum, high-level disinfected.

Non-Critical Instruments (contact intact skin, e.g., x-ray head, pulse oximeter): Cleaned and low-level disinfected.

Key Point: There is no special "HIV protocol" for instrument reprocessing. Standard sterilization kills HIV, Hepatitis B, and Hepatitis C effectively.

Step 5: Aerosol Management & Procedure-Specific Considerations

Pre-procedural Mouth Rinse: The ADA strongly recommends a 1% povidone-iodine or 0.2% chlorhexidine mouth rinse for all patients before any procedure to reduce the microbial load in aerosols. This is especially prudent for immunocompromised patients.

High-Volume Evacuation (HVE): Use HVE for every procedure generating aerosol. This is standard, not special.

Rubber Dam: Use a rubber dam whenever possible to minimize aerosol generation and blood exposure. This is a best practice for all restorative and endodontic procedures.

Step 6: Sharps Safety - The Highest Priority

One-Handed Recapping: Never use two hands to recap a needle. Use a one-handed scoop technique or a recapping device.

Disposal: All sharps (needles, scalpel blades, burs) are immediately placed into a puncture-resistant, yellow, labeled sharps container at the point of use.

Needlestick Protocol: If a needlestick or splash occurs, the ADA mandates an immediate, documented protocol: wash the site with soap and water (or irrigate mucous membranes with saline), report to the designated infection control officer, and seek immediate post-exposure prophylaxis (PEP) within 2 hours, ideally.

Step 7: Post-Treatment & Waste Management

Clinical Waste: All waste contaminated with blood or body fluids (e.g., gloves, gauze, barriers) is disposed of as clinical waste in a designated, lined bin.

Hand Hygiene: Perform hand hygiene immediately after glove removal. Use soap and water if hands are visibly soiled, or an alcohol-based hand rub if not.

The Professor's Summary for Your Exam:

The ADA's approach is universal. The patient with HIV/AIDS is treated with the same rigorous standard precautions as every other patient. The key differences are not in the type of precautions, but in the meticulousness of their application. You must be hyper-vigilant about:

Sharps safety (highest risk of transmission).

PPE integrity (no shortcuts).

Sterilization validation (use biological indicators regularly).

Confidentiality (ethical and legal obligation).

Now, let's go deeper. A question for you:

You are treating a 35-year-old patient with well-controlled HIV (undetectable viral load, CD4 >500). During a routine scaling, you accidentally sustain a needlestick injury from a used anesthetic needle. The patient's last viral load test was 3 months ago and was undetectable. What is your immediate step-by-step action plan, and what is the current evidence-based risk of seroconversion in this specific scenario?

Mock Questions (Author-written, for self-assessment)

1. **What is the approximate risk of HIV seroconversion after a single percutaneous sharps injury with contaminated blood?**

- A. 0.03%
- B. 0.3%
- C. 3%
- D. 30%

Answer: B

2. **Which Spaulding category of instrument (e.g., scalpels, forceps) must be sterilised after every single use?**

- A. Non-critical
- B. Semi-critical
- C. Critical
- D. None of the above

Answer: C

3. **What is the recommended contact time for a 1:10 dilution of sodium hypochlorite when managing a blood or body-fluid spill?**

- A. 1 minute
- B. 5 minutes
- C. 10 minutes
- D. 30 minutes

Answer: C

4. **Post-exposure prophylaxis (PEP) after a needlestick injury should ideally be commenced within:**

- A. 24 hours
- B. 2 hours
- C. 72 hours
- D. 1 week

Answer: B

5. **Which of the following best reflects the ADA position on disclosure of a patient's HIV status?**

- A. It must be shared with all clinic staff
- B. It is protected by privacy law and disclosed only on a genuine clinical need-to-know basis
- C. It should be written on the outside of the patient file
- D. The patient must disclose it to the receptionist

Answer: B

6. **Which pre-procedural mouth rinse is recommended to reduce microbial load in aerosols, especially before aerosol-generating procedures?**

- A. Sodium bicarbonate rinse
- B. 0.2% chlorhexidine or 1% povidone-iodine rinse
- C. 3% hydrogen peroxide rinse
- D. Plain tap water only

Answer: B

7. **What is the correct technique for recapping a used local anaesthetic needle?**

- A. Two-handed recapping for better control
- B. A one-handed scoop technique or a mechanical recapping device
- C. Bending the needle by hand first
- D. Asking the dental assistant to hold the cap

Answer: B



The Dentistry
— I N S I G H T —

Chapter 2 — Infection Control for Cancer and Chemotherapy Patients

Step-by-Step: Infection Control for Cancer & Chemotherapy Patients

Step 1: Pre-Treatment Risk Stratification (The Critical First Step)

This is the most important difference from the HIV patient. You must determine the patient's neutrophil count (ANC - Absolute Neutrophil Count) and platelet count.

Low Risk (ANC > 1500 cells/ μ L, Platelets > 50,000): Standard precautions apply. Routine care is generally safe.

Moderate Risk (ANC 500-1500, Platelets 20,000-50,000): Increased vigilance. Avoid aerosol-generating procedures if possible. Use atraumatic techniques.

High Risk (ANC < 500, Platelets < 20,000): Defer all non-emergency dental care. The patient is at extreme risk of sepsis from oral bacteria and uncontrolled bleeding. Only emergency care (e.g., drainage of an abscess, extraction of a source of sepsis) is performed, and only in a hospital setting with hematology support.

Key Question: You must ask: "When was your last chemotherapy cycle?" The nadir (lowest point) of neutrophils is typically 7-14 days after chemo. This is the highest-risk period.

Step 2: Personal Protective Equipment (PPE) - Enhanced for Patient Protection

Gloves: Standard, single-use. Double-gloving is recommended for surgical procedures (extractions, biopsies) due to the high risk of bleeding and the need for prolonged barrier integrity.

Mask: N95/P2 respirator is strongly recommended for all aerosol-generating procedures (AGPs). This is not to protect you from the patient's cancer, but to protect the patient from your respiratory flora (e.g., Aspergillus, Pseudomonas). A standard surgical mask is insufficient for this purpose.

Eye Protection: Standard.

Gown: Standard fluid-resistant gown.

Mock Questions (Author-written, for self-assessment)

1. **An Absolute Neutrophil Count (ANC) below which value places a chemotherapy patient in the 'high risk' category, requiring deferral of non-emergency dental care?**

- A. <1500 cells/ μ L
- B. <1000 cells/ μ L
- C. <500 cells/ μ L
- D. <2000 cells/ μ L

Answer: C

2. **The neutrophil nadir (lowest point) after a chemotherapy cycle typically occurs:**

- A. 1-2 days after chemotherapy
- B. 7-14 days after chemotherapy
- C. 3-4 weeks after chemotherapy
- D. Immediately during the infusion

Answer: B

3. **A platelet count of 20,000-50,000/ μ L in a chemotherapy patient is classified as:**

- A. >50,000 (low risk)
- B. Moderate risk
- C. <20,000 (high risk)
- D. Not clinically relevant

Answer: B

4. **Why is an N95/P2 respirator recommended for aerosol-generating procedures in chemotherapy patients?**


- A. To protect the clinician from the patient's malignancy
- B. To protect the immunocompromised patient from the clinician's own respiratory flora
- C. To reduce clinic noise
- D. It is a billing requirement only

Answer: B

5. **Double gloving is specifically recommended in chemotherapy patients for:**

- A. Routine oral examinations
- B. Surgical procedures such as extractions or biopsies, due to bleeding risk
- C. Denture try-ins
- D. Fluoride varnish application

Answer: B



The Dentistry
— INSIGHT —

6. **An ANC under 500 cells/ μ L with active infection risk means emergency dental treatment should occur:**

- A. In the regular dental clinic as usual
- B. In a hospital setting with haematology support
- C. At the patient's home
- D. Only after chemotherapy is fully completed

Answer: B

7. **The single most important question to ask a chemotherapy patient before any dental procedure is:**

- A. "Do you brush twice daily?"
- B. "When was your last chemotherapy cycle?"
- C. "Do you floss regularly?"
- D. "What toothpaste do you use?"

Answer: B



The Dentistry
— I N S I G H T —

Chapter 3 — Stroke (Cerebrovascular Accident): Recognition and Dental Management

1. Pathogenesis and Etiology

A stroke, or Cerebrovascular Accident (CVA), occurs when the blood supply to part of the brain is interrupted or reduced, preventing brain tissue from getting oxygen and nutrients.

Ischaemic Stroke (approx. 85%): Caused by a blockage (thrombus or embolus) in an artery supplying the brain. Often related to atherosclerosis or atrial fibrillation.

Haemorrhagic Stroke (approx. 15%): Results from the rupture of a blood vessel (e.g., aneurysm or poorly controlled hypertension) causing bleeding into or around the brain.

Transient Ischaemic Attack (TIA): A "mini-stroke" where neurological deficit resolves within 24 hours. It is a significant warning sign for a future major stroke.

2. Clinical Features (Signs & Symptoms)

Recognition in the dental chair often follows the F.A.S.T. protocol:

Facial Droop: Asymmetry of the mouth or eyes.

Arm Weakness: Inability to raise both arms equally.

Speech Difficulty: Slurring or inability to find words (aphasia).

Time: Immediate emergency intervention is required.

Other features: Sudden severe headache ("thunderclap" headache), loss of vision, dizziness, or loss of consciousness.

3. Diagnosis

Clinical Presentation: Sudden onset of focal neurological deficits.

Medical History Review: Identifying risk factors (Hypertension, Diabetes, Smoking, Atrial Fibrillation, Anticoagulant use).

Hospital Diagnosis: Non-contrast CT scan (to differentiate between ischaemic and haemorrhagic) and MRI.

4. Dental Management (ADC TG 4 Guidelines)

A. Pre-operative Management

Timing: Elective dental treatment should be deferred for at least 6 months following a stroke to allow for neurological stabilization and reduction of recurrence risk.

Stress Reduction: Short morning appointments, effective local anesthesia, and potentially conscious sedation.

Medication Review: Most stroke patients are on antiplatelets (Aspirin, Clopidogrel) or anticoagulants (Warfarin, NOACs like Rivaroxaban). Do not stop these medications for routine dental procedures; manage bleeding locally using sutures and haemostatic agents (e.g., oxidized cellulose).

B. In-Chair Emergency Protocol

Stop Procedure: Remove all instruments from the mouth.

Positioning: Keep the patient in a comfortable position (usually semi-reclined). If unconscious, use the recovery position.

Emergency Services: Call 000 (Australia) or local emergency services immediately.

Oxygen: Administer oxygen only if the patient is hypoxic ($SpO_2 < 94\%$).

Monitor: Record vital signs (Blood Pressure, Heart Rate) until help arrives.

C. Oral Health Considerations

Physical Impairment: Patients may have hemiplegia (paralysis on one side), making oral hygiene difficult. Recommend electric toothbrushes or modified handles.

Xerostomia: Often a side effect of antihypertensive medications.

Dysphagia: Be cautious with water spray and high-volume suction to prevent aspiration.

Step 1: Definition & Pathophysiology (The "What")

A stroke is a neurological deficit caused by an acute focal injury to the central nervous system (CNS) due to a vascular cause. It is a medical emergency.

Two Main Types:

Ischemic Stroke (87%): Blockage of a blood vessel.

Thrombotic: Clot forms locally (e.g., atherosclerosis).

Embolic: Clot travels from elsewhere (e.g., heart in Atrial Fibrillation).

Hemorrhagic Stroke (13%): Rupture of a blood vessel.

Intracerebral: Bleeding within brain tissue (often due to hypertension).

Subarachnoid: Bleeding into space between brain and meninges (often due to aneurysm).

Transient Ischemic Attack (TIA): A "mini-stroke" where symptoms resolve within 24 hours. This is a major warning sign for a future full stroke.

Step 2: Clinical Presentation (The "Signs & Symptoms")

The FAST Acronym (Crucial for Emergency Recognition):

F (Face Drooping): One side of the face is numb or drooping. Ask the patient to smile.

A (Arm Weakness): One arm is weak or numb. Ask the patient to raise both arms.

S (Speech Difficulty): Slurred speech or inability to speak/understand.

T (Time to call emergency services): Note the time symptoms began.

Other Key Signs:

Sudden severe headache (especially in hemorrhagic).

Sudden trouble walking, dizziness, loss of balance.

Sudden vision loss in one or both eyes.

Step 3: Dental Implications (The "Why We Care")

This is the core of your dental education. A stroke history changes everything.

A. Pre-Treatment Assessment (Medical History & Risk Stratification)

Identify the Type & Timeframe:

Acute Stroke (< 3 months): ABSOLUTE CONTRAINDICATION for elective dental care. The patient is at high risk for another stroke, bleeding, or aspiration. Only emergency, life-saving care (e.g., airway management) is permitted.

Chronic Stroke (> 6 months): Generally safe for elective care, but with modifications.

TIA History: Treat with extreme caution; patient is high-risk.

Medication Review (The "Big Three"):

Antiplatelets (e.g., Aspirin, Clopidogrel, Ticagrelor): Increased bleeding risk. Do NOT stop for routine dental procedures (extractions, scaling, fillings). Bleeding is usually manageable with local measures (gauze, sutures, hemostatic agents).

Anticoagulants (e.g., Warfarin, Apixaban, Rivaroxaban, Dabigatran): High bleeding risk.

Warfarin: Check INR. Safe for most procedures if INR < 3.5 (or < 4.0 per some guidelines). Do NOT stop without consulting the physician.

DOACs (Direct Oral Anticoagulants): Do not require routine INR monitoring. Do NOT stop for most dental procedures. Consult physician for major surgery.

Antihypertensives: Ensure blood pressure is controlled (< 180/110 mmHg for elective care). Uncontrolled hypertension increases risk of hemorrhagic stroke.

Assess Residual Deficits:

Motor: Hemiparesis (weakness on one side). Position the patient for comfort and safety. Use a wheelchair or support.

Sensory: Numbness or paresthesia. Be careful with injections and instruments.

Cognitive: Aphasia (difficulty speaking/understanding), dysarthria (slurred speech), memory loss, depression. Use simple language, write instructions, involve a caregiver.

Swallowing (Dysphagia): CRITICAL. Risk of aspiration pneumonia. Use a rubber dam, high-volume suction, and keep the patient upright. Avoid supine position.

Visual: Hemianopsia (blindness in half of visual field). Approach from the unaffected side.

B. Intra-Operative Management (During the Procedure)

Positioning: Semi-supine or upright. Support the affected side. Avoid prolonged supine position (reduces cerebral perfusion).

Anesthesia:

Use local anesthetic WITH vasoconstrictor (e.g., Lidocaine 1:100,000 epinephrine). The amount of epinephrine is safe and helps control bleeding.

Avoid epinephrine in patients with uncontrolled hypertension or recent stroke (< 6 months).

Use aspiration technique to prevent intravascular injection.

Stress Reduction: Anxiety increases blood pressure and heart rate. Use pre-medication (e.g., benzodiazepine) if needed, but be cautious with sedation (respiratory depression).

Monitoring: Monitor vital signs (BP, HR, O2 saturation) throughout the procedure.

Bleeding Control: Be prepared for prolonged bleeding. Use local hemostatic agents (Gelfoam, Surgicel, Tranexamic acid mouthwash).

C. Post-Operative Care

Observe for Stroke Symptoms: Monitor for any new neurological deficits (FAST signs) for at least 30 minutes post-op.

Pain Management: Avoid NSAIDs (e.g., Ibuprofen, Naproxen) in patients on anticoagulants/antiplatelets (increases bleeding risk). Use Acetaminophen (Paracetamol) as first-line.

Follow-up: Ensure the patient has a caregiver to take them home. Provide written post-op instructions in large font.

Step 4: Emergency Protocol (If a Stroke Occurs in the Dental Chair)

Stop the procedure immediately.

Call emergency services (911/112). Note the time of symptom onset.

Position the patient: Supine with head elevated 30 degrees (if no airway compromise) or recovery position (if unconscious).

Maintain airway: Suction, remove dental materials, insert an oral airway if needed.

Administer oxygen (2-4 L/min).

Monitor vital signs.

Do NOT give aspirin (if hemorrhagic, it will worsen bleeding).

Mock Questions (Author-written, for self-assessment)

1. Approximately what proportion of strokes are ischaemic in nature?

- A. 50%
- B. 65%
- C. 85%
- D. 95%

Answer: C

2. A Transient Ischaemic Attack (TIA) is defined as a neurological deficit that resolves within:

- A. 1 hour
- B. 6 hours
- C. 24 hours
- D. 72 hours

Answer: C

3. In the FAST acronym for stroke recognition, the letters stand for Face, Arm, Speech and:

- A. Temperature
- B. Time
- C. Tone
- D. Tremor

Answer: B

4. Elective dental treatment following a stroke should generally be deferred for at least:

- A. 1 month
- B. 6 months
- C. 2 years
- D. Indefinitely

Answer: B

5. A stroke patient taking warfarin needs a single, straightforward tooth extraction. The recommended approach is to:

- A. Stop warfarin 5 days before the procedure
- B. Continue warfarin (if INR is therapeutic) and manage bleeding with local measures
- C. Switch the patient to aspirin instead
- D. Refer only once the INR reaches zero

Answer: B

6. Which finding helps differentiate a stroke from hypoglycaemia in the dental chair?


- A. Both conditions are always accompanied by profuse sweating
- B. A stroke typically produces a focal neurological deficit, whereas hypoglycaemia produces more global/whole-body symptoms
- C. Hypoglycaemia always causes facial droop
- D. There is no clinically useful difference

Answer: B

7. The first action if a stroke is suspected mid-treatment is to:

- A. Administer aspirin immediately
- B. Stop the procedure and call emergency services
- C. Continue treatment while monitoring vital signs
- D. Give high-flow oxygen regardless of saturation

Answer: B





The Dentistry

— I N S I G H T —

Chapter 4 — Seizures and Antiepileptic Drugs: Dental Management

Definition: A seizure is a transient occurrence of signs and/or symptoms due to abnormal, excessive, or synchronous neuronal activity in the brain. It's a symptom, not a disease.

Classification (Simplified for Dental Context):

- **Generalized Seizures:** Involve both hemispheres. Tonic-Clonic (Grand Mal) is the most critical for us. Loss of consciousness, stiffening (tonic), then rhythmic jerking (clonic).
- **Focal (Partial) Seizures:** Start in one area. Can be simple (no loss of consciousness) or complex (impaired consciousness). A focal seizure can secondarily generalize into a tonic-clonic seizure.
- **Absence Seizures (Petit Mal):** Brief staring spells, common in children. Usually not a major dental emergency, but the patient may appear unresponsive.

Dental Significance:

- **Emergency:** A tonic-clonic seizure in the dental chair is a true medical emergency.
- **Triggers:** Stress, pain, infection, sleep deprivation, flashing lights (rare), and local anesthetic overdose (especially with prilocaine or articaine in high doses, or accidental intravascular injection).
- **Medication Interactions:** Antiepileptic drugs (AEDs) have significant interactions with dental drugs.

Part 2: Antiepileptic Drugs (AEDs) - The Core Pharmacology

This is the "meat" of your question. We'll cover the most common ones you will encounter.

1. Phenytoin (Dilantin)

- **Mechanism:** Stabilizes neuronal membranes by blocking voltage-gated sodium channels.
- **Indication:** Generalized tonic-clonic and focal seizures. Not for absence seizures (can worsen them).
- **Dental Contraindication:** Gingival Hyperplasia (overgrowth). This is the most famous dental side effect. It's not a contraindication to using the drug, but a contraindication to poor oral hygiene. It requires meticulous plaque control, frequent recalls, and sometimes gingivectomy.
- **Systemic Side Effects:** Nystagmus, ataxia, drowsiness, hirsutism, megaloblastic anemia (due to folate deficiency), osteomalacia (due to Vitamin D metabolism interference), Stevens-Johnson syndrome (rare, severe).
- **Drug Interactions (Critical for Dentists):**
- **Metronidazole & Fluconazole:** MAJOR. These drugs inhibit phenytoin metabolism, leading to toxic levels (ataxia, nystagmus, sedation). Avoid or use with extreme caution.
- **NSAIDs:** Can increase phenytoin levels.
- **Doxycycline:** Phenytoin reduces doxycycline's half-life.
- **Oral Contraceptives:** Phenytoin reduces their efficacy.

2. Carbamazepine (Tegretol)

- **Mechanism:** Sodium channel blocker.
- **Indication:** Focal seizures, generalized tonic-clonic, and trigeminal neuralgia (a key dental indication!).
- **Contraindications:** History of bone marrow depression (agranulocytosis, aplastic anemia), AV heart block, MAOI use.
- **Side Effects:** Drowsiness, dizziness, ataxia, diplopia, hyponatremia (low sodium), leukopenia (low WBC). Agranulocytosis is a rare but fatal risk.
- **Drug Interactions (Critical):**
- **Macrolide Antibiotics (Erythromycin, Clarithromycin):** MAJOR. Inhibit carbamazepine metabolism → toxicity. Avoid.
- **Propoxyphene (Darvon):** Contraindicated. Increases carbamazepine levels.
- **Acetaminophen:** Carbamazepine reduces its efficacy.
- **Oral Contraceptives:** Reduces efficacy.

3. Valproic Acid (Depakote, Depakene)

- **Mechanism:** Multiple: increases GABA, blocks sodium channels.
- **Indication:** Broad spectrum: generalized tonic-clonic, absence, myoclonic, focal, and bipolar disorder.
- **Contraindications:** Liver disease, pancreatitis, pregnancy (high teratogenicity - neural tube defects).
- **Side Effects:** Hepatotoxicity (especially in children <2 years on polytherapy), pancreatitis, thrombocytopenia (low platelets → bleeding risk), weight gain, tremor, alopecia.
- **Dental Implications:** Bleeding risk due to thrombocytopenia. Check platelet count before surgery. Also, it can cause gingival hyperplasia (less common than phenytoin).
- **Drug Interactions:**
- **Aspirin:** Displaces valproate from protein binding → increased free valproate → toxicity. Avoid aspirin.
- **NSAIDs:** Increased risk of bleeding (additive effect with thrombocytopenia).
- **Benzodiazepines:** Increased sedation.

4. Phenobarbital

- Mechanism: Enhances GABA activity.
- Indication: Generalized tonic-clonic, focal seizures. (Less common now due to side effects).
- Contraindications: Porphyria, severe liver disease, respiratory depression.
- Side Effects: Sedation (significant), cognitive impairment, depression, physical dependence, tolerance.
- Dental Implications: Profound sedation. Enzyme inducer – it accelerates the metabolism of many drugs.
- Drug Interactions:
 - Metronidazole, Fluconazole, Doxycycline, Oral Contraceptives, Corticosteroids: All are metabolized faster, reducing their efficacy.
 - Warfarin: Reduced efficacy.

5. Newer AEDs (Gabapentin, Lamotrigine, Levetiracetam, Topiramate)

- General Advantage: Fewer drug interactions and less enzyme induction.
- Gabapentin (Neurontin): Used for focal seizures and neuropathic pain (e.g., trigeminal neuralgia). Minimal drug interactions. Side effects: sedation, dizziness.
- Lamotrigine (Lamictal): Broad spectrum. Risk of Stevens-Johnson syndrome (severe skin reaction) if dose is escalated too quickly.
- Levetiracetam (Keppra): Very safe profile. Minimal drug interactions. Side effects: irritability, dizziness.
- Topiramate (Topamax): Weight loss, kidney stones, glaucoma (acute angle closure – dental emergency? No, but be aware), cognitive slowing.

Part 3: Dental Management Protocol (Step-by-Step)

Pre-Treatment (The Consultation):

History: Type of seizure, frequency, last seizure, triggers, current medications (name, dose, compliance).

Medical Clearance: If uncontrolled or on complex polytherapy, contact the physician.

Blood Work: If on valproate (platelets, LFTs), carbamazepine (CBC, LFTs, sodium), or phenytoin (levels, folate, Vitamin D).

Appointment Timing: Schedule in the morning (patient is less fatigued). Ensure they have taken their medication as usual. Never tell them to stop their AEDs.

Stress Reduction: Pre-op sedation (e.g., nitrous oxide, oral benzodiazepine like triazolam – but be aware of additive sedation with AEDs). Good local anesthesia to prevent pain.

Intra-Treatment (The Procedure):

Local Anesthesia: Use lidocaine or mepivacaine with epinephrine. Avoid prilocaine and articaine in high doses (lower seizure threshold). Use aspirating syringe to prevent intravascular injection.

Drug Interactions:

- Avoid: Erythromycin, clarithromycin, metronidazole, fluconazole, propoxyphene.
- Use with Caution: NSAIDs (bleeding risk with valproate), aspirin (avoid with valproate), tetracyclines (reduced efficacy with phenytoin/phenobarbital).
- Safe: Penicillin VK, amoxicillin, clindamycin, cephalexin.

Analgesics:

- Acetaminophen: Safe, but may be less effective in patients on enzyme-inducing AEDs.
- NSAIDs (Ibuprofen, Naproxen): Use with caution in valproate patients (bleeding). Avoid aspirin.
- Opioids: Codeine and tramadol can lower the seizure threshold. Avoid tramadol. Use morphine or hydromorphone if needed.

Antibiotics: Amoxicillin is first choice. Avoid metronidazole and macrolides.

Post-Treatment (The Emergency Protocol):

If a seizure occurs in the chair:

STOP the procedure. Remove all instruments from the mouth.

Protect the patient: Lower the chair, turn them on their side (recovery position) to prevent aspiration.

Do NOT force anything into the mouth (no bite blocks, no fingers).

Time the seizure. Most are self-limiting (<2 minutes).

If >5 minutes or repeated seizures (Status Epilepticus):

- Call 911/EMS.
- Administer emergency medication:
 - Midazolam (Versed): 5-10 mg intranasal (preferred in dental setting) or intramuscular.
 - Diazepam (Valium): 5-10 mg rectal gel (Diastat) or IV (if trained).
 - Lorazepam (Ativan): 2-4 mg IV/IM.

Post-ictal state: Patient will be confused, drowsy, and may have a headache. Do not discharge them alone. Monitor until fully oriented.

Summary Table for Quick Reference

Drug

Key Dental Side Effect
Key Drug Interaction (Dental)
Contraindication
Phenytoin
Gingival Hyperplasia
Metronidazole, Fluconazole (toxicity)
Absence seizures
Carbamazepine
Hyponatremia, Leukopenia
Erythromycin, Clarithromycin (toxicity)
Bone marrow depression
Valproic Acid
Bleeding (Thrombocytopenia)
Aspirin (toxicity), NSAIDs (bleeding)
Liver disease, Pregnancy
Phenobarbital
Sedation
Metronidazole, Fluconazole (reduced efficacy)
Porphyria, Respiratory depression
Levetiracetam
Irritability
Minimal
None significant

Final Question to Go Deeper

You now have the framework. Let's test your clinical reasoning.

Scenario: A 35-year-old female patient with well-controlled epilepsy on Carbamazepine presents for extraction of a mandibular third molar. She has a history of trigeminal neuralgia (also on

Step 2: Classification (Simplified for Dental Context) We don't need the full ILAE classification, but we must distinguish between the two main types you will encounter:

A. Generalized Seizures (Involve both hemispheres)

Tonic-Clonic (Grand Mal): The classic, most dramatic. Tonic phase: Rigid muscle contraction, loss of consciousness, cyanosis, possible cry. Clonic phase: Rhythmic jerking of limbs. Post-ictal phase: Confusion, fatigue, headache, amnesia.

Absence (Petit Mal): Brief (seconds) loss of awareness. Staring, blinking. No post-ictal phase. Common in children. Often missed in the dental chair.

B. Focal (Partial) Seizures (Start in one hemisphere)

Simple Focal: No loss of consciousness. Symptoms depend on the brain area (e.g., motor: jerking of one arm; sensory: strange smell or taste).

Complex Focal: Impaired consciousness. Patient may appear confused, perform repetitive movements (automatisms like lip-smacking, picking at clothes).

Step 2: Pre-Operative Preparation

Appointment Timing: Schedule in the morning when the patient is well-rested and has eaten.

Stress Reduction: Profound local anesthesia, short appointments, consider nitrous oxide sedation (if safe and indicated).

Emergency Kit: ALWAYS have a fully stocked emergency kit. For seizures, the key drugs are:

Midazolam (IV/IM/Intranasal/Buccal) – First-line for prolonged seizures.

Diazepam (IV/rectal gel) – Alternative.

Oxygen tank & mask.

Suction.

Oral airway (nasopharyngeal is safer than oropharyngeal in a seizing patient).

Inform Staff: Ensure your team knows the protocol.

Phase 2: Intra-Operative (During the Seizure)

Step 1: Recognize & Stop Treatment

Stop all dental procedures immediately.

Remove all instruments from the mouth. This is the #1 priority to prevent aspiration or laceration.

Remove the rubber dam, cotton rolls, and any loose objects.

Step 2: Protect the Patient (The "ABCs" of Seizure Management)

A - Airway: Do NOT force anything into the mouth. Do NOT put your fingers in. Do NOT use a bite block. The patient's jaw is clenched. Trying to force it open can break teeth or your fingers. Wait for the clonic phase to pass.

B - Breathing: Once the clonic jerking stops, assess breathing. If not breathing, start rescue breathing.

C - Circulation: Check pulse. If no pulse, start CPR.

Position: Gently lower the chair to a flat position. Turn the patient on their left side (recovery position). This allows saliva/blood to drain out, preventing aspiration.

Protect from Injury: Clear the area of sharp objects, dental unit arms, etc. Do NOT restrain.

Step 4: Administer Medication (If Indicated)

When to give medication?

Seizure lasting > 5 minutes (Status Epilepticus).

Seizure lasting > 2 minutes in a patient with a known history of status epilepticus.

Multiple seizures without regaining consciousness.

Which drug?

First-line: Midazolam (5-10 mg intranasal or buccal, or 5 mg IM). This is the safest and most effective in the dental office.

Alternative: Diazepam (5-10 mg IV or 0.2-0.5 mg/kg rectal gel).

How to give?

Intranasal: Use a mucosal atomizer device (MAD). Half the dose in each nostril.

Buccal: Inject the solution into the buccal mucosa.

IM: Inject into the deltoid or vastus lateralis.

After medication: Administer 100% oxygen via a non-rebreather mask.

Phase 3: Post-Operative (After the Seizure)

Step 1: Post-Ictal Care

The patient will be confused, drowsy, and possibly combative. Do not leave them alone.

Reassure them calmly. "You had a seizure. You are safe now."

Monitor vital signs (BP, HR, O2 saturation).

Do not attempt to restart dental treatment.

Step 2: Disposition

When to call 911 (Emergency Medical Services)?

First-time seizure.

Seizure lasting > 5 minutes (Status Epilepticus).

Seizure in a pregnant patient.

Seizure in a diabetic patient (possible hypoglycemia).

Patient does not regain consciousness.

Patient has multiple seizures.

Patient is injured (e.g., head trauma from fall).

When to discharge home?

Patient has a known seizure disorder.

Seizure was brief (< 2 minutes).

Patient is fully conscious and oriented.

Patient has a responsible adult to drive them home.

Advise them to see their neurologist.

Mock Questions (Author-written, for self-assessment)

1. **The classic dental side effect most strongly associated with phenytoin is:**

- A. Xerostomia
- B. Gingival hyperplasia
- C. Black hairy tongue
- D. Angular cheilitis

Answer: B

2. **Which antibiotic class should be avoided in a patient taking carbamazepine because of a major toxicity**

interaction?

- A. Penicillin V
- B. Macrolides (e.g., erythromycin, clarithromycin)
- C. Amoxicillin
- D. Cephalexin

Answer: B

3. **Which analgesic should be avoided in a patient on valproic acid, as it displaces the drug from protein binding and raises toxicity risk?**

- A. Paracetamol
- B. Aspirin
- C. Codeine
- D. Amoxicillin

Answer: B

4. **The first-line drug for managing a seizure lasting longer than five minutes in the dental chair is:**

- A. Oral diazepam
- B. Intranasal, buccal or intramuscular midazolam
- C. Intramuscular adrenaline
- D. Salbutamol

Answer: B

5. **During the tonic-clonic phase of a seizure, the correct airway management is to:**

- A. Insert a bite block between the teeth
- B. Never force anything into the mouth
- C. Hold the tongue with gauze
- D. Give oral glucose immediately

Answer: B

6. **For a patient with a known seizure disorder, which local anaesthetic combination is preferred, and which should be limited in high doses?**


- A. Prefer high-dose prilocaine; avoid lidocaine
- B. Prefer lidocaine or mepivacaine; use high-dose prilocaine/articaine cautiously
- C. Both are equally recommended without restriction
- D. All local anaesthetics must be avoided

Answer: B

7. **A patient with known epilepsy has a brief, typical tonic-clonic seizure lasting 90 seconds in the waiting room and then recovers fully and is oriented. The appropriate disposition is to:**

- A. Call an ambulance regardless of recovery
- B. Allow discharge with a responsible adult present and advise neurology follow-up
- C. Admit to hospital in every case
- D. Proceed immediately with the planned dental treatment

Answer: B



Chapter 5 — Dementia: Comprehensive Dental Management

The Caregiver is Your Co-Pilot: You must establish a direct line of communication with the primary caregiver. They are your historian, your behavior interpreter, and your home-care executor. Get written consent from the patient (if they have capacity) or the legal guardian.

Assess the Stage of Dementia: This dictates everything.

Early Stage (Mild): Patient can communicate needs, follow instructions, and cooperate. Focus on education, intensive prevention, and complex treatment if tolerated.

Middle Stage (Moderate): Significant cognitive decline, memory loss, possible agitation, and difficulty with new routines. Focus on simplification, desensitization, and prevention. Complex procedures are risky.

Late Stage (Severe): Non-verbal, dependent for all care, often bedridden. Focus on palliative care, pain relief, and oral hygiene maintenance to prevent aspiration pneumonia.

Medical History Deep Dive: Dementia patients are often elderly with multiple comorbidities (CVD, diabetes, hypertension) and polypharmacy. Look for:

Anticholinergics: Worsen cognitive decline and cause xerostomia (dry mouth).

Antipsychotics/Sedatives: Risk of extrapyramidal side effects (tardive dyskinesia affecting jaw movement) and orthostatic hypotension.

Anticoagulants/Antiplatelets: Increased bleeding risk.

Bisphosphonates/Denosumab: Risk of Medication-Related Osteonecrosis of the Jaw (MRONJ).

Step 2: The Appointment Environment & Communication (The Art)

This is where we apply the "Tell-Show-Do" technique on steroids.

The Environment:

Sensory Overload is the Enemy: Dim the lights. Turn off background music. Minimize staff traffic. Use a quiet, familiar operatory.

Time of Day: Schedule appointments during the patient's "best" time of day (often mid-morning, after breakfast). Avoid late afternoon "sundowning" periods.

Short Appointments: 15-20 minutes maximum. The goal is a positive interaction, not a full quadrant of dentistry.

Communication (The "Dementia-Speak"):

Approach from the Front: Never startle them from behind.

Eye Level: Sit or kneel to be at their eye level.

Simple, One-Step Commands: "Please open your mouth." Not "Open your mouth so I can look at your teeth and clean them."

Positive Language: "Let's put this little mirror in to see your pretty smile." Avoid "This won't hurt."

Validate, Don't Correct: If they say "I want to go home," don't say "You are home." Say "I know you want to go home. We'll be done soon, and then you can go."

Use Non-Verbal Cues: A gentle touch on the hand, a calm smile, and a soft tone are more powerful than words.

Step 3: Clinical Adjustments (The Science & Technique)

Positioning:

Semi-Upright is Safer: Reduces risk of aspiration, which is a major concern due to dysphagia (swallowing difficulty) common in late stages.

Support the Head: Use a headrest and pillows to prevent involuntary head movements.

The "Hand-Over-Hand" Technique: For oral examination or hygiene, place your hand over the patient's hand and guide it to their mouth. This can reduce the feeling of invasion.

Examination:

Be Fast and Gentle: Use a mouth prop (e.g., a soft bite block) to maintain access without fatigue. A finger sweep is often better than a mirror for initial assessment.

Look for Pain Signs: In non-verbal patients, look for grimacing, moaning, pulling at the face, refusal to eat, or changes in behavior (agitation). These are often the only signs of a dental abscess or toothache.

Treatment Modifications (The "KISS" Principle - Keep It Simple, Safe):

Prevention is the ONLY Cure:

Fluoride Varnish: Apply at every visit. It's fast, painless, and highly effective.

High-Fluoride Toothpaste (5000 ppm): Prescribe for caregiver to apply.

Chlorhexidine Swabs/Gel: For chemical plaque control when brushing is impossible.

Restorative Dentistry:

Avoid Complex Procedures: No crowns, bridges, or implants if possible. The patient cannot maintain them, and the procedures are too long.

Glass Ionomer Cement (GIC) is Your Best Friend: It releases fluoride, bonds to tooth structure, and is easy to place. Use it for

all restorations. It's a "therapeutic" restoration, not a "cosmetic" one.

Atraumatic Restorative Treatment (ART): Hand instruments only. No drill. Perfect for this population.

Extractions:

Simple, Atraumatic Extractions: Avoid surgical extractions if possible. The goal is to remove the source of infection with minimal trauma.

Suture with Care: Use resorbable sutures to avoid a return visit for removal.

Prosthodontics:

Dentures: Often poorly tolerated in moderate-severe dementia. The patient may forget they have them, lose them, or refuse to wear them. Label dentures with the patient's name.

If they have dentures: Check for fit. Ill-fitting dentures cause sores, which lead to pain and refusal to eat. Reline them frequently.

Anesthesia & Sedation:

Local Anesthesia: Use a topical gel first. Inject slowly. Be prepared for a sudden movement.

Sedation: Use with extreme caution. Benzodiazepines can cause paradoxical agitation in dementia patients. Nitrous oxide can be helpful for mild anxiety but can cause confusion. General anesthesia in a hospital setting is often the safest option for extensive treatment in a patient who cannot cooperate.

Step 4: Post-Operative & Home Care (The Long Game)

Pain Management: Dementia patients cannot reliably report pain. Assume they are in pain after any procedure. Prescribe acetaminophen (paracetamol) as a first line. Avoid NSAIDs if they are on anticoagulants or have renal issues. Avoid opioids if possible (risk of constipation, delirium, and respiratory depression).

Caregiver Instructions:

Written, Simple Instructions: "Brush teeth twice a day. Use this toothpaste. Call if they refuse to eat."

Demonstrate: Show the caregiver how to brush the patient's teeth (e.g., knee-to-knee position, using a three-headed toothbrush, or a suction toothbrush).

Oral Hygiene Aids:

Electric Toothbrush: Often better tolerated than manual.

Suction Toothbrush: For patients with dysphagia, to prevent aspiration of saliva and toothpaste.

Foam Swabs: For applying

Step 1: The Legal & Ethical Framework (The Australian Mandate)

Before any clinical intervention, you must establish the legal basis for treatment. This is the most critical step in Australia.

Capacity Assessment (The "Gillick" Competence Test): You, as the dentist, must make a prima facie assessment of the patient's capacity to consent.

Can they understand the nature and purpose of the proposed treatment?

Can they retain that information?

Can they weigh it up to arrive at a decision?

Can they communicate that decision?

If NO: The patient lacks capacity for that specific decision.

The Hierarchy of Decision-Making (Australian Law):

Advance Care Directive (ACD): If the patient has a legally binding ACD (e.g., "No dental extractions"), you must follow it.

Enduring Guardian / Medical Power of Attorney: The person legally appointed to make health decisions. You must verify their authority (e.g., NSW Guardianship Act 1987).

Person Responsible (in order): Spouse/partner, carer, close friend/relative. This is a default position, not a substitute for a formal guardian.

Guardianship Tribunal: For major, irreversible decisions (e.g., multiple extractions under GA).

ADA Policy Stance: The ADA states that treatment must be in the patient's best interests. This is a beneficence model, not a paternalistic one. You cannot treat simply because it is "good for them" if it causes distress without clear benefit.

Step 2: The Clinical Risk Assessment (ADA-Aligned)

The ADA's Guidelines for Infection Control and Policy on Radiographs are directly relevant.

Radiographs (The "Justification" Principle): The ADA is strict on radiation justification.

Early Dementia: Standard radiographs (BW, OPG) are acceptable if the patient can cooperate.

Moderate/Severe Dementia: Avoid intraoral radiographs if the patient cannot tolerate the sensor/film. The risk of a bitewing is outweighed by the risk of agitation, aspiration, or injury. An OPG (orthopantomogram) is the preferred modality if a radiograph is essential. If the patient cannot tolerate an OPG, clinical examination alone is acceptable.

Medical History & Polypharmacy: The ADA's Prescribing Guidelines are crucial.

Anticoagulants (Warfarin, DOACs): The ADA states do not routinely cease anticoagulants for dental procedures. For simple

extractions (1-3 teeth), continue warfarin if INR is within therapeutic range (2.0-3.0). For complex surgery, liaise with the physician.

Anticholinergics: Be aware of xerostomia risk. The ADA recommends saliva substitutes (e.g., GC Dry Mouth Gel) and high-fluoride toothpaste (5000ppm) as first-line management.

Infection Control: The ADA's Infection Control Guidelines are non-negotiable.

Aspiration Risk: Dementia patients have a high risk of aspiration. Use a rubber dam for any restorative procedure. If a rubber dam is not tolerated, use a high-volume suction and a throat pack (with a visible gauze tail).

Aerosol Generation: Minimise aerosols. Use hand instruments (ART) where possible. If a high-speed handpiece is needed, use extra-oral suction and pre-procedural mouth rinse (0.2% chlorhexidine).

Step 3: The Clinical Management Plan (ADA-Recommended)

The ADA's Policy on Oral Health and Older Persons explicitly recommends a preventive-first, minimally invasive approach.

Prevention (The Core):

Fluoride Varnish (5% NaF): Apply at every visit. The ADA supports this as a standard of care.

High-Fluoride Toothpaste (5000ppm): Prescribe via PBS (Authority Required) for patients with high caries risk.

Chlorhexidine Gel (0.2%): For chemical plaque control when brushing is impossible.

Restorative (The "ART" Approach):

Atraumatic Restorative Treatment (ART): The ADA endorses ART for patients with special needs. Use Glass Ionomer Cement (GIC) (e.g., Fuji IX, Equia Forte). No drill, no local anaesthetic needed for small lesions.

Avoid Complex Restorations: No crowns, bridges, or implants. The ADA states that "complex treatment should only be considered if the patient can tolerate the procedure and maintain the restoration."

Extractions (The "Last Resort"):

Simple, Atraumatic Extractions: Use local anaesthetic (with vasoconstrictor if no contraindication). Avoid surgical extractions if possible.

Sutures: Use resorbable sutures (e.g., Vicryl Rapide) to avoid a return visit for removal.

Post-operative Care: Prescribe paracetamol (1g QID) as first-line analgesia. Avoid NSAIDs if on anticoagulants. Avoid opioids (risk of delirium, constipation, respiratory depression).

Prosthodontics (The "Pragmatic" Approach):

Dentures: The ADA recommends simple, well-fitting dentures. Label dentures with the patient's name (e.g., using a denture marking kit).

Implant-Retained Overdentures: Not recommended for patients with moderate-severe dementia due to inability to maintain oral hygiene and risk of peri-implantitis.

Sedation & General Anaesthesia (The "Hospital" Setting):

Oral Sedation (Benzodiazepines): Use with extreme caution. The ADA warns that benzodiazepines can cause paradoxical agitation in dementia patients.

Nitrous Oxide: Can be helpful for mild anxiety but can cause confusion in moderate-severe dementia.

General Anaesthesia (GA): The ADA recommends GA in a hospital setting for extensive treatment in a patient who cannot cooperate. This requires a pre-anaesthetic assessment and liaison with the geriatrician.

Step 4: The Appointment Structure (ADA-Recommended)

The ADA's Guidelines for the Management of Patients with Special Needs recommend a structured, predictable approach.

Pre-Appointment Communication:

Contact the caregiver to confirm the appointment time, medication list, and any recent changes in behaviour.

Send a reminder letter (simple, large font) to the patient/caregiver.

The Appointment Itself:

Time of Day: Schedule for the patient's "best" time (usually mid-morning).

Environment: Quiet, dimly lit, minimal staff traffic.

Communication: Use the "Tell-Show-Do" technique. Use simple, one-step commands. Validate, don't correct.

Duration: 15-20 minutes maximum.

Post-Appointment:

Written Instructions: Provide a simple, written care plan for the caregiver.

Follow-Up: Schedule a recall in 3-6 months (depending on risk).

Step 5: The Medico-Legal Documentation (The Australian Standard)

This is where the ADA is most strict. Your records must be defensible.

Consent: Document the capacity assessment. If the patient lacks capacity, document who provided consent (e.g., "Consent obtained from Mrs. Smith, Enduring Guardian, as per NSW Guardianship Act").

Treatment Plan: Document the rationale for

Mock Questions (Author-written, for self-assessment)

1. **For most patients with dementia, dental appointments are best scheduled:**

- A. Late in the afternoon
- B. Mid-morning, in the patient's usual best period of alertness
- C. In the evening
- D. Timing makes no difference

Answer: B

2. **Which restorative material is generally favoured in dementia patients because it releases fluoride, bonds to tooth structure and is simple to place?**

- A. Composite resin
- B. Glass ionomer cement
- C. Amalgam only
- D. Zirconia

Answer: B

3. **The technique using hand instruments only, avoiding the drill where possible, is known as:**

- A. General anaesthesia
- B. Atraumatic Restorative Treatment (ART)
- C. Rubber dam isolation
- D. Vital bleaching

Answer: B

4. **In a non-verbal patient with late-stage dementia, dental pain is most often first recognised through:**

- A. A clear verbal complaint
- B. A behavioural change, such as agitation or refusal to eat
- C. A completely normal radiograph
- D. Increased appetite

Answer: B

5. **Which class of sedative drug can paradoxically increase agitation in patients with dementia?**

- A. NSAIDs
- B. Benzodiazepines
- C. Antibiotics
- D. Fluoride varnish

Answer: B

6. **If a patient with dementia lacks capacity for a specific decision and there is no formally appointed guardian, who is generally next in line to provide substitute consent?**

- A. Any available staff member
- B. The legally recognised "Person Responsible" (e.g., spouse, carer, close relative), per the state guardianship hierarchy
- C. The treating dentist alone
- D. No one can provide consent

Answer: B

7. **The recommended communication style for a patient with dementia is best summarised as:**

- A. "Correct their confusion firmly and repeatedly"
- B. "Validate, don't correct"
- C. "Ignore all requests to leave"
- D. "Use long, detailed explanations"

Answer: B



The Dentistry
— INSIGHT —

Chapter 6 — Periodontal Charting: Probing Depths, CAL, BOP, Furcation and Mobility

1. Probing Depths (PD) - The Core Measurement

What it is: The distance from the gingival margin (GM) to the base of the sulcus/pocket.

How to measure:

Use a periodontal probe (e.g., UNC-15, Williams, or Michigan "O" probe with Williams markings).

The probe has markings at 1, 2, 3, 5, 7, 8, 9, and 10 mm (UNC-15) or 1, 2, 3, 5, 7, 8, 9, 10 mm (Williams).

Walk the probe circumferentially around each tooth.

Record six points per tooth:

Disto-buccal (DB)

Mid-buccal (B)

Mesi-buccal (MB)

Disto-lingual (DL)

Mid-lingual (L)

Mesi-lingual (ML)

Normal values: 1-3 mm.

Clinical significance:

4-5 mm: Shallow pocket. May be due to gingival inflammation (pseudopocket) or early attachment loss.

6 mm or more: Deep pocket. Indicates significant attachment loss and periodontitis.

2. Gingival Margin (GM) or Recession (REC)

What it is: The position of the gingival margin relative to the cemento-enamel junction (CEJ).

How to measure:

If the GM is apical to the CEJ (recession), measure the distance from the CEJ to the GM. Record as a positive number (e.g., +2 mm).

If the GM is coronal to the CEJ (overgrowth/pseudopocket), measure the distance from the CEJ to the GM. Record as a negative number (e.g., -1 mm).

If the GM is at the CEJ, record as 0.

Clinical significance: Recession exposes root surfaces, leading to sensitivity, caries, and esthetic concerns.

3. Clinical Attachment Level (CAL) - The True Measure of Disease

What it is: The distance from the CEJ to the base of the sulcus/pocket. This is the true measure of periodontal destruction.

How to calculate: $CAL = \text{Probing Depth (PD)} + \text{Recession (REC)}$

Example: PD = 6 mm, REC = +2 mm. $CAL = 6 + 2 = 8$ mm.

Example: PD = 4 mm, REC = -1 mm (overgrowth). $CAL = 4 + (-1) = 3$ mm.

Why it matters: Probing depth alone can be misleading. A 6 mm pocket with 3 mm of recession (CAL = 9 mm) is much worse than a 6 mm pocket with no recession (CAL = 6 mm).

4. Bleeding on Probing (BOP)

What it is: The presence or absence of bleeding within 15-30 seconds after probing.

How to record: Simply mark a dot or a "B" at the site where bleeding occurs.

Clinical significance:

Positive BOP: Indicates active inflammation in the pocket. It is a sign of disease activity.

Negative BOP: Indicates a healthy or inactive site.

BOP < 10%: Healthy.

BOP > 30%: Significant inflammation.

5. Furcation Involvement

What it is: Bone loss in the area between the roots of multi-rooted teeth (molars).

How to assess: Use a Nabers probe (curved, with markings).

How to grade (Glickman's classification):

Grade I: Incipient. The probe can feel a concavity but cannot enter the furcation.

Grade II: The probe can enter the furcation but cannot pass through to the other side (through-and-through).

Grade III: Through-and-through. The probe passes completely from one side to the other.

Grade IV: Same as Grade III, but the furcation is visible clinically due to recession.

Clinical significance: Furcation involvement significantly worsens the prognosis of a tooth. It is difficult to clean and often

requires surgical intervention.

6. Mobility

What it is: The degree of horizontal or vertical movement of a tooth.

How to assess: Use the handles of two instruments (e.g., mirror and probe) to apply gentle pressure.

How to grade (Miller's classification):

Grade 0: No detectable movement.

Grade I: 0.2-1 mm of horizontal movement.

Grade II: 1-2 mm of horizontal movement.

Grade III: > 2 mm of horizontal movement OR any vertical movement (depressibility).

Clinical significance: Mobility indicates loss of supporting bone and periodontal ligament. Grade III mobility often indicates a hopeless tooth.

Part 3: The "How" - Step-by-Step Charting Protocol

Start with the Maxillary Right Quadrant (Tooth #1-16).

Probe the Disto-buccal (DB) of Tooth #1. Record the number.

Walk the probe to the Mid-buccal (B). Record.

Walk to the Mesio-buccal (MB). Record.

Repeat for the Lingual side: DL, L, ML.

Move to Tooth #2. Repeat.

Continue through all teeth.

After probing, assess and record:

BOP at each site.

Recession at each site.

Furcation on all molars.

Mobility on all teeth.

Calculate CAL for any site with PD > 3 mm or recession.

Part 4: Common Errors and How to Avoid Them

Error

Consequence

How to Avoid

Probing too lightly

Underestimates pocket depth

Use firm, consistent pressure (20-25 grams).

Probing at an angle

Inaccurate depth

Keep the probe parallel to the long axis of the tooth.

Not walking the probe

Misses the deepest point

Walk the probe in a "stepping" motion around the tooth.

Forgetting to record BOP

Misses inflammation

Record BOP immediately after probing each site.

Not assessing furcations

Misses a key prognostic factor

Always use a Nabers probe on molars.

Confusing PD with CAL

Misdiagnosis

Always calculate CAL for deep pockets.


Part 5: The "So What?" - Interpreting the Chart

A basic chart is not just a collection of numbers. It tells a story.

Generalized PD 4-5 mm with BOP: Generalized gingivitis or early periodontitis.

Localized PD 6-8 mm on molars with furcation Grade II: Localized severe periodontitis. These teeth have a guarded prognosis.

Generalized PD 7-10 mm with mobility Grade II and furcation Grade III: Generalized severe periodontitis. Many teeth may be hopeless.



PD 3 mm with recession +5 mm (CAL = 8 mm): This is a case of recession with attachment loss, not a deep pocket. The disease is "burned out" but the damage is done.

Mock Questions (Author-written, for self-assessment)

1. **Clinical Attachment Level (CAL) is calculated as:**

- A. $CAL = \text{Probing Depth} - \text{Recession}$
- B. $CAL = \text{Probing Depth} + \text{Recession}$
- C. $CAL = \text{Gingival Margin} - \text{CEJ only}$
- D. $CAL = \text{Bleeding on Probing} + \text{Probing Depth}$

Answer: B

2. **A full periodontal chart records how many sites per tooth?**

- A. Two (mesial and distal only)
- B. Six (three buccal, three lingual/palatal)
- C. One per tooth
- D. Four, all on the buccal surface

Answer: B

3. **A furcation grade in which the probe passes completely from one side of the tooth to the other is:**

- A. Grade I
- B. Grade II
- C. Grade III
- D. Grade 0

Answer: C

4. **Grade III mobility is defined as horizontal movement of:**

- A. 0–0.2 mm
- B. 0.2–1 mm
- C. 1–2 mm
- D. More than 2 mm, or any vertical (depressible) movement

Answer: D

5. **A tooth with a probing depth of 4 mm and recession of 3 mm has a Clinical Attachment Level of:**

- A. 1 mm
- B. 4 mm
- C. 7 mm
- D. 3 mm

Answer: C

6. **Bleeding on probing (BOP) recorded at more than 30% of sites generally indicates:**

- A. A healthy periodontium
- B. Significant, generalised gingival inflammation
- C. A measurement error requiring no action
- D. No further treatment is required

Answer: B

7. **Which charting error most commonly leads to underestimation of true pocket depth?**

- A. Probing with too much pressure
- B. Probing too lightly
- C. Using a Nabers probe on a molar
- D. Walking the probe circumferentially

Answer: B



Chapter 7 — Consent and Capacity: The Gillick Test and the "Person Responsible"

Step 1: The Foundational Australian Principle - The "Therapeutic Relationship" and the Gillick Competence Test

The ADA emphasizes that consent is not a one-off event but a process within a therapeutic relationship. The key legal test for a minor or a person with a cognitive disability in Australia is often the Gillick competence test (from the UK House of Lords, adopted in Australian common law).

The Gillick Test: A person is competent to consent if they have sufficient understanding and intelligence to enable them to understand fully what is proposed. This is a functional test, not an age or diagnosis-based test.

The ADA's Stance: The ADA explicitly states that a person's disability does not, of itself, mean they lack capacity to consent. You must assess their functional ability to understand the specific dental decision at hand.

Step 2: The Critical First Step - Capacity Assessment (The Australian "Functional" Approach)

The ADA directs you to assess capacity for the specific decision at the specific time. This is identical to the global standard but is codified in Australian law. The test is:

Can the person understand the information relevant to the decision? (e.g., "This tooth has a hole. I need to clean it and put a filling in.")

Can the person retain that information? (Even for a short time.)

Can the person use or weigh that information as part of the process of making the decision? (e.g., "If I get the filling, the pain stops. If I don't, it will get worse.")

Can the person communicate their decision (by any means)? (Verbal, non-verbal, assistive technology.)

Crucial ADA Point: If the patient can do all four, they have capacity. You must obtain their direct, informed consent. You cannot override their decision simply because you disagree with it or because a family member disagrees.

Step 3: When Capacity is Lacking - The Australian "Person Responsible" Hierarchy

If, after your best efforts (using simple language, visual aids, etc.), the patient lacks capacity for that specific decision, you must turn to a "Person Responsible". This is a legal term defined in each state's Guardianship and Administration Act. The hierarchy is generally:

A legally appointed guardian (by the Victorian Civil and Administrative Tribunal (VCAT), NCAT, QCAT, etc.).

An enduring guardian (appointed by the person before they lost capacity).

A spouse or de facto partner (including same-sex partner).

A carer (who provides unpaid care and is not a paid health professional).

A close friend or relative (who has a close personal relationship).

Critical ADA Rule: The "Person Responsible" must make decisions based on the person's known wishes (substituted judgment). If those wishes are unknown, they must act in the person's best interests. They cannot make a decision based on their own convenience or preferences.

Step 4: The ADA's Specific Dental Practice Protocol

Here is the exact workflow the ADA recommends for your clinic:

1. Pre-Appointment:

Review the patient's history. Note any diagnosis, but do not pre-judge capacity.

Prepare your environment: quiet, no time pressure, visual aids (models, intraoral photos, simple diagrams).

If possible, speak with the carer or family before the appointment to understand the patient's communication style, fears, and triggers.

2. The Initial Interaction (Patient First):

Always address the patient directly first. Introduce yourself, explain your role.

Use simple, concrete language. "I'm going to look at your teeth. Open wide."

Assess their ability to engage. Do they make eye contact? Do they respond to simple commands?

3. The Capacity Assessment (For the Specific Procedure):

Explain the proposed treatment using the "Chunk and Check" method.

Chunk: "I see a cavity in this tooth. It's a hole. It can get bigger and hurt."

Check: "Can you tell me what I just said?"

Explain the risks and benefits simply. "The filling will fix the hole. The needle for freezing will pinch for a few seconds."

Ask for their choice. "Do you want me to fix this tooth today?"

4. Decision Point:

If YES and they demonstrate capacity: Proceed with obtaining their informed consent. Document the conversation, their

responses, and your assessment of capacity in the chart.

If NO or UNCLEAR: Do not proceed. If they refuse, that is their right, even if you disagree. Document the refusal.

If they cannot demonstrate capacity: Stop. Do not proceed. Explain to the patient that you need to speak with their helper/family.

5. Engaging the "Person Responsible":

Confirm their legal authority. Ask for documentation (guardianship papers, enduring power of attorney).

Explain the patient's lack of capacity for this specific decision.

Provide the "Person Responsible" with the same full disclosure you would give a capacitated patient.

Ask the "Person Responsible": "Based on what you know of [Patient's Name]'s values and wishes, what do you believe they would choose?"

Document everything: Your capacity assessment, the "Person Responsible's" identity and authority, the discussion, and their decision.

Step 5: Special Considerations & "Deeper" Knowledge (Australian Context)

Assent vs. Consent: Even when a "Person Responsible" provides legal consent, you should always seek the assent (agreement) of the patient. If the patient is actively resisting, you must stop and re-evaluate. Forcing treatment on a resisting patient, even with legal consent, is ethically problematic and can be traumatic.

Fluctuating Capacity: For patients with dementia, schedule appointments for their "best time of day." Re-assess capacity at each visit.

Minimizing Restraint: The use of physical or chemical restraint (sedation) to perform treatment on a non-consenting patient is a last resort and requires a separate, rigorous consent process with the "Person Responsible," often involving a second opinion and a clear medical necessity.

Emergency Exception: In a true, life-threatening dental emergency (e.g., severe facial space infection causing airway compromise), you can provide necessary treatment without consent to save life or prevent serious harm. This is a very narrow exception.

Documentation is Your Shield: Meticulous documentation is your only defense against a future complaint. Record:

Your capacity assessment (the four abilities).

The specific information you disclosed.

The patient's response.

The name and relationship of the "Person Responsible."

Their decision and the basis for it (substituted judgment or best interests).

Summary for the Busy Clinician (Australian Version)

Scenario

Action

Patient has capacity

Obtain direct informed consent from the patient.

Patient lacks capacity

Identify the "Person Responsible" (legal hierarchy).

"Person Responsible" is identified

Obtain informed consent from them, based on substituted judgment.

No "Person Responsible" is available

Do not proceed with non-emergency treatment. Seek legal advice or apply to the state guardianship tribunal (e.g., VCAT, NCAT).

Patient resists (even with "Person Responsible" consent)

Stop. Re-assess. Do not force treatment.

Mock Questions (Author-written, for self-assessment)

1. **The Gillick competence test primarily assesses capacity to consent based on:**

- A. The patient's chronological age
- B. The patient's medical diagnosis
- C. Functional understanding of the specific decision at hand
- D. The opinion of the patient's family

Answer: C

2. **Which of the following is NOT one of the four functional elements of a capacity assessment?**

- A. Understanding the relevant information
- B. Retaining the information

- C. Weighing the information to reach a decision
- D. Having a formal diagnosis of intellectual disability

Answer: D

3. **A patient is assessed as having capacity, but a family member disagrees with the patient's treatment choice. The dentist should:**
- A. Follow the family member's wishes instead
 - B. Respect the patient's own autonomous decision
 - C. Decline to treat either party
 - D. Refer the matter to police

Answer: B

4. **When a "Person Responsible" provides substitute consent, their decision should be based on:**
- A. Whatever is most convenient for them
 - B. The patient's known wishes (substituted judgment) or the patient's best interests
 - C. Cost considerations alone
 - D. Whichever option is quickest to complete

Answer: B

5. **The Australian "Person Responsible" hierarchy typically places which of the following first?**
- A. A distant relative met that day
 - B. A legally appointed or enduring guardian
 - C. The dental receptionist
 - D. The treating dentist

Answer: B

6. **Using physical or chemical restraint to treat a resisting patient who lacks capacity is considered:**
- A. A routine step requiring no special process
 - B. A last resort requiring a separate, rigorous consent process, often with a second opinion
 - C. Never permissible under any circumstances
 - D. Automatically approved by any relative present

Answer: B

7. **In a genuine, life-threatening dental emergency where consent cannot be obtained in time, a dentist may:**
- A. Never provide any treatment
 - B. Provide the treatment necessary to save life or prevent serious harm
 - C. Wait for the guardianship tribunal to convene
 - D. Always contact police before treating

Answer: B



Chapter 8 — Acute Severe Asthma: Emergency Management in the Dental Chair

Your Exact Next Steps (ADA Protocol)

Step 1: Activate Emergency Medical Services (EMS) Immediately

Action: Call Triple Zero (000).

What to say: "Medical emergency. Severe asthma attack. Patient is unresponsive to initial Salbutamol. Cannot speak in full sentences. We need an ambulance immediately."

Provide: Your practice address, floor level, and any specific entry instructions.

Step 2: Continue Salbutamol (Reliever)

Action: Administer another 4 puffs of Salbutamol via spacer.

Frequency: Repeat 4 puffs every 4 minutes while waiting for the ambulance.

Why: You must continue bronchodilation even while escalating care.

Step 3: Administer High-Flow Oxygen

Action: Give oxygen at 8–10 L/min via a non-rebreather mask.

Why: The patient is severely hypoxic. High-flow oxygen is critical to maintain oxygen saturation until paramedics arrive.

Step 4: Administer Adrenaline (Epinephrine) - The Critical Escalation Drug

Drug: Adrenaline (Epinephrine) 1:1000

Dose: 0.3 mg intramuscularly (IM) (which is 0.3 mL of the 1:1000 solution)

Site: Anterolateral thigh (mid-thigh, outer aspect)

Device: Use an EpiPen (auto-injector) or draw from an ampoule.

Repeat: Can be repeated every 5–15 minutes if no improvement.

Why: This is the life-saving drug for severe, unresponsive asthma. It is a potent bronchodilator and also reduces airway edema.

Step 5: Monitor and Prepare for Handover

Monitor: Consciousness, breathing rate, pulse, oxygen saturation (if pulse oximeter available).

Position: Keep upright or in tripod position.

Prepare: Have your documentation ready (time of onset, drugs given, doses, times, patient response).

Summary of Your Actions (Quick Reference)

Action

Drug/Dose

Route

Frequency

1

Stop procedure, position upright.

•

2

Administer Salbutamol via spacer.

4 puffs. Wait 4 minutes.

3

Assess. If improved, continue monitoring.

Repeat 4 puffs every 4 min if needed.

4

If NOT improved, call 000.

•

5

While waiting: Continue Salbutamol + give Oxygen.

4 puffs every 4 min + 8-10 L/min O2.

6


If life-threatening (silent chest, cyanosis, exhaustion): Give Adrenaline IM.

0.3 mg (0.3 mL of 1:1000) IM. Repeat every 5-15 min.

7

Monitor and hand

Document



Why This is the Correct Answer (According to ADA)

The "4 + 4 + 4" Rule: The ADA uses 4 puffs as the standard dose, not 2. You already gave the first 4. Now you repeat 4 puffs every 4 minutes.

Adrenaline is for Life-Threatening Asthma: In Australia, IM adrenaline is explicitly recommended for severe, life-threatening asthma that is not responding to Salbutamol. This patient qualifies.

Oxygen is Mandatory: The ADA mandates oxygen for severe respiratory distress. You must have it in your emergency kit.

Call 000 Early: The moment you recognize the attack is not responding to initial therapy, you call for an ambulance. Do not wait.

Question to Go Deeper

You have administered 4 puffs of Salbutamol, high-flow oxygen, and 0.3 mg of IM adrenaline. The ambulance is still 10 minutes away. The patient is now becoming lethargic, cyanotic, and has a "silent chest" (no wheezing, no breath sounds).

According to the Australian Resuscitation Council (ARC) and ADA guidelines, what is your immediate next action, and what specific

Mock Questions (Author-written, for self-assessment)

1. **In the ADA emergency asthma protocol, how many puffs of salbutamol via spacer are given per dose?**

- A. 2 puffs
- B. 4 puffs
- C. 6 puffs
- D. 10 puffs

Answer: B

2. **The dose and route of adrenaline recommended for life-threatening asthma unresponsive to salbutamol is:**

- A. 0.3 mg intramuscularly (1:1000)
- B. 1 mg intravenous bolus
- C. 0.1 mg subcutaneously only
- D. An oral tablet

Answer: A

3. **The preferred injection site for intramuscular adrenaline in an asthma emergency is the:**

- A. Deltoid muscle
- B. Anterolateral thigh
- C. Gluteal region
- D. Forearm

Answer: B

4. **The target oxygen flow rate via a non-rebreather mask in severe asthma is approximately:**

- A. 1-2 L/min
- B. 8-10 L/min
- C. 15-20 L/min
- D. No oxygen is required

Answer: B

5. **A "silent chest" on auscultation during an asthma attack indicates:**

- A. Mild asthma that is resolving
- B. Life-threatening asthma with minimal air movement
- C. A normal recovering airway
- D. Anxiety alone, with no respiratory compromise

Answer: B


6. **If there is no improvement, intramuscular adrenaline may be repeated:**

- A. Every 1 minute
- B. Every 5-15 minutes
- C. Once only, never repeated
- D. Only once every hour

Answer: B

7. **The correct first action when a patient fails to improve after the initial dose of salbutamol is to:**

- A. Wait and observe for another hour
- B. Call 000 and continue salbutamol plus oxygen while awaiting the ambulance
- C. Offer the patient some food
- D. Discharge the patient to go home and rest



The Dentistry
— I N S I G H T —



The Dentistry

— I N S I G H T —

Chapter 9 — COPD: Management in the Dental Chair

■ Comprehensive Step-by-Step Overview: Management of COPD in the Dental Chair (ADA Guidelines)

1. Pre-Treatment Assessment (The "ADA Triage")

This is the most critical step. The ADA mandates a structured assessment before any procedure.

Confirm Diagnosis & Severity:

Ask: "Do you have COPD, emphysema, or chronic bronchitis?"

Ask about FEV1 (Forced Expiratory Volume in 1 second). If the patient doesn't know, ask about their MRC Dyspnoea Scale (e.g., "Do you get short of breath walking on flat ground?").

ADA Severity Classification (Simplified):

Mild: FEV1 $\geq$ 80% predicted. No oxygen.

Moderate: FEV1 50-79%. May use inhalers.

Severe: FEV1 30-49%. Often on home oxygen.

Very Severe: FEV1 < 30% or on long-term oxygen.

Identify "Red Flags" (ADA Contraindications to Routine Care):

Acute exacerbation: Increased cough, sputum purulence, fever, or dyspnoea. Defer all non-emergency treatment.

Unstable COPD: Recent hospitalization (within 3 months), frequent exacerbations (>2/year), or change in medication.

Cor Pulmonale: Signs of right heart failure (ankle oedema, raised JVP). High risk for general anaesthesia/sedation.

Medication Reconciliation (The "Inhaler Check"):

Identify all inhalers: SABA (e.g., salbutamol), LABA (e.g., salmeterol), LAMA (e.g., tiotropium), ICS (e.g., fluticasone).

Check for oral steroids: Prednisolone (risk of adrenal suppression).

Check for antibiotics: Azithromycin (risk of QT prolongation).

Ask about oxygen use: Flow rate (L/min) and hours per day.

2. Treatment Planning & Modifications (The "ADA Chairside Protocol")

Appointment Timing:

Morning appointments are preferred. COPD patients are often more fatigued and breathless later in the day.

Short appointments (30-45 minutes max). Avoid prolonged procedures.

Positioning:

Semi-supine (45 degrees) or upright. Never fully supine in severe COPD. This can worsen ventilation-perfusion mismatch and cause orthopnoea.

Allow the patient to sit upright for 2-3 minutes between position changes.

Anaesthesia & Pain Control:

Local Anaesthesia is SAFE. Use lidocaine 2% with 1:100,000 epinephrine or articaine 4% with 1:100,000 epinephrine.

Epinephrine is NOT contraindicated. It provides profound anaesthesia, reduces bleeding, and prolongs effect. The ADA explicitly states: "The use of epinephrine in local anaesthetics is safe in stable COPD."

Avoid prilocaine with felypressin (can cause methemoglobinemia in patients with low oxygen saturation).

Sedation (The "ADA Caution"):

Minimal sedation (anxiolysis) is preferred. Use oral midazolam (2.5-5 mg) or nitrous oxide/oxygen (max 50% N2O, 50% O2).

Avoid benzodiazepines in severe COPD (risk of respiratory depression). If needed, use flumazenil as reversal agent.

Avoid opioids (morphine, fentanyl) unless absolutely necessary and with monitoring.

General anaesthesia is high-risk and should only be performed in a hospital setting with an anaesthetist.

Oxygen Therapy (The "ADA Rule"):

Routine oxygen is NOT recommended for stable COPD patients.

Indications for oxygen:

SpO2 < 90% on room air.

Patient is on home oxygen (maintain their usual flow rate).

Signs of respiratory distress (tachypnoea, accessory muscle use).

Target SpO2: 88-92%. Do NOT over-oxygenate (risk of CO2 retention and hypercapnic respiratory failure).

Use nasal prongs (not a mask) to allow the patient to talk and breathe easily.

3. Chairside Monitoring (The "ADA Safety Net")

Continuous Pulse Oximetry: Mandatory for all moderate-to-severe COPD patients.

Visual Monitoring: Watch for:

Accessory muscle use (sternocleidomastoid, intercostals).

Pursed-lip breathing (a sign of distress).

Cyanosis (late sign).

Blood Pressure: Monitor every 5-10 minutes.

Respiratory Rate: Count for 30 seconds. Normal is 12-20. Tachypnoea (>25) is a warning sign.

4. Emergency Management (The "ADA Crisis Protocol")

If the patient develops acute dyspnoea, wheezing, or desaturation:

Stop the procedure.

Position upright.

Administer their own SABA inhaler (salbutamol) via spacer. Give 2-4 puffs every 20 seconds.

Administer oxygen via nasal prongs at 2-4 L/min. Target SpO₂ 88-92%.

Call for medical assistance if no improvement in 5 minutes.

Consider nebulised salbutamol (5 mg) if available.

Do NOT give high-flow oxygen (risk of CO₂ narcosis).

5. Post-Treatment Care (The "ADA Follow-up")

Observe for 15-30 minutes after the procedure.

Ensure the patient is stable (SpO₂ > 90%, no dyspnoea).

Provide written post-op instructions including:

Continue all regular medications.

Avoid strenuous activity for 24 hours.

Contact their GP if they develop increased cough, fever, or shortness of breath.

Document everything: SpO₂ readings, oxygen used, any adverse events.

🔑 Key Knowledge Points for Australian Dental Practice (ADA Specific)

Epinephrine is SAFE. The ADA explicitly states that epinephrine in local anaesthesia is not contraindicated in stable COPD. It is actually beneficial for haemostasis and prolonged anaesthesia.

Avoid high-flow oxygen. The ADA warns against over-oxygenation in COPD patients due to the risk of Haldane effect and CO₂ retention.

Nitrous oxide is acceptable but only at ≤50% concentration. Higher concentrations can cause diffusion hypoxia.

Oral midazolam is the preferred sedative for mild-to-moderate COPD. For severe COPD, avoid sedation altogether.

Azithromycin interaction: If the patient is on prophylactic azithromycin, be aware of QT prolongation risk. Avoid using other QT-prolonging drugs (e.g., erythromycin, some antifungals).

Corticosteroid cover: If the patient has been on oral prednisolone (>7.5 mg/day for >3 weeks), they may need stress-dose steroids (e.g., 100 mg hydrocortisone IV) before major surgery. For routine extractions, this is usually not required.

? Question to Go Deeper

You have a 72-year-old patient with severe COPD (FEV₁ 32%, on home oxygen 2 L/min, and using a LAMA/LABA/ICS inhaler). They present for a routine scale and clean. During the procedure, they become acutely dyspnoeic, and their SpO₂ drops to 85%...

📖 Mock Questions (Author-written, for self-assessment)

1. In a patient with stable COPD, using a local anaesthetic containing epinephrine is:

- A. Absolutely contraindicated
- B. Safe, and beneficial for haemostasis and duration of anaesthesia
- C. Reserved for emergencies only
- D. Always replaced by plain articaine

Answer: B

2. The target SpO₂ range for a COPD patient receiving supplemental oxygen is:

- A. 95-100%
- B. 88-92%
- C. 80-85%
- D. No target is needed

Answer: B

3. High-flow oxygen is avoided in COPD patients mainly because it risks:

- A. Being too costly
- B. CO₂ retention and hypercapnic respiratory failure
- C. An unpleasant taste

D. No risk; it is actually preferred

Answer: B

4. **The maximum recommended nitrous oxide concentration when sedating a COPD patient is approximately:**

- A. 70%
- B. 50%
- C. 100%
- D. 30%

Answer: B

5. **The preferred oxygen delivery device for a stable COPD patient requiring supplemental oxygen in the chair is:**

- A. A non-rebreather mask
- B. Nasal prongs
- C. A venturi mask at maximum flow
- D. No device; mouth breathing only

Answer: B

6. **A red flag that contraindicates routine dental care in a COPD patient is:**

- A. Stable, regular inhaler use
- B. An acute exacerbation with increased sputum purulence and fever
- C. Use of a spacer device
- D. A history of having previously quit smoking

Answer: B

7. **The preferred chair position for a patient with severe COPD is:**

- A. Fully supine
- B. Semi-supine (~45°) or upright
- C. Trendelenburg (head down)
- D. Left lateral position only

Answer: B



The Dentistry
— I N S I G H T —

Chapter 10 — Tobacco and Alcohol: Effects in Dentistry

Here's a comprehensive overview:

Effects of Tobacco & Alcohol in Dentistry

🚬 TOBACCO

Oral Hard Tissue Effects

Tooth staining — extrinsic brown/black discolouration, difficult to remove

Increased caries risk — reduced salivary flow, altered oral flora

Alveolar bone loss — accelerated periodontal destruction

Impaired healing — vasoconstriction reduces blood supply to healing sockets

Periodontal Effects

Masking of inflammation — nicotine causes vasoconstriction, reducing classic signs (bleeding on probing). Disease is worse than it appears clinically.

Increased pocket depth and attachment loss

Poorer response to periodontal therapy — both surgical and non-surgical

Higher implant failure rates

Soft Tissue Effects

Smoker's melanosis — diffuse brown pigmentation of gingiva

Nicotinic stomatitis — white, keratotic palatal mucosa with red dots (minor salivary gland openings)

Hairy tongue — elongated filiform papillae, brown/black discolouration

Leukoplakia — white patch; potentially malignant disorder

Erythroplakia — red patch; high malignant transformation risk

Oral submucous fibrosis — associated with smokeless tobacco/betel nut

Oral cancer — strong independent risk factor. Synergistic with alcohol.

Salivary Effects

Reduced salivary flow → xerostomia

Altered salivary composition → increased caries risk

Impaired antimicrobial function

Implant & Surgical Implications

Relative contraindication for implants in heavy smokers

Higher rates of peri-implantitis

Impaired wound healing post-extraction → dry socket risk significantly increased

Delayed healing after periodontal surgery

Anaesthetic Considerations

Increased airway reactivity → bronchospasm risk

Higher LA dose may be needed due to altered tissue pH in inflamed/infected sites

🍷 ALCOHOL

Oral Hard Tissue Effects

Dental erosion — acidic pH of alcoholic beverages (wine, beer, ciders) directly erodes enamel

Increased caries — sugary mixers, reduced salivary protection, poor diet

Bruxism association — alcohol disrupts sleep architecture, increases parafunctional activity

Soft Tissue Effects

Oral cancer — alcohol is an independent risk factor; combined with tobacco the risk is multiplicative, not additive

Candidiasis — immunosuppression and xerostomia increase susceptibility

Glossitis — nutritional deficiencies (B12, folate) from chronic alcohol use

Angular cheilitis — nutritional deficiency related

Salivary Effects

Chronic use → reduced salivary flow (xerostomia)

Altered salivary proteins

Parotid gland enlargement (sialosis) in chronic alcoholism

Haemostatic & Surgical Considerations

Antiplatelet effect — inhibits thromboxane A₂ → prolonged bleeding time

Hepatic dysfunction in chronic users → reduced clotting factor synthesis (II, VII, IX, X)

Warfarin interaction — alcohol potentiates anticoagulation



Always take bleeding history before extractions or surgery

Drug Interactions Relevant to Dentistry

Metronidazole + alcohol → disulfiram-like reaction (nausea, vomiting, flushing, tachycardia). Counsel patients strictly — avoid alcohol during course and 48 hours after.

Tinidazole — same interaction, same counselling

NSAIDs + alcohol → increased GI bleeding risk

Paracetamol + chronic alcohol use → hepatotoxicity risk (use with caution, reduce dose)

Benzodiazepines + alcohol → additive CNS/respiratory depression — dangerous in sedation patients

Behavioural & Medical Considerations

Poor compliance with post-op instructions

Nutritional deficiencies → impaired healing

Liver disease → altered drug metabolism (LA, analgesics, antibiotics)

Wernicke's encephalopathy / Korsakoff syndrome in severe chronic users

⚠ **COMBINED TOBACCO + ALCOHOL**

Risk Factor

Tobacco Alone

Alcohol Alone

Combined

Oral Cancer Risk

↑ ↑

↑ ↑

↑ ↑ ↑ ↑ (multiplicative)

Periodontal Disease

↑ ↑ ↑

↑

↑ ↑ ↑

Wound Healing

Impaired

Impaired

Severely impaired

Candidiasis

↑

↑

↑ ↑

🦷 **Dental Management Principles**

Before treatment:

Thorough medical and social history — quantify tobacco (pack years) and alcohol (standard drinks/week)

Assess bleeding risk (LFTs if chronic alcoholic)

Screen all mucosal surfaces for potentially malignant disorders

During treatment:

Use vasoconstrictor-containing LA cautiously if cardiovascular compromise

Avoid metronidazole if patient drinks alcohol

Extra care with haemostasis in alcoholic patients

After treatment:

Smoking cessation counselling — brief intervention (Ask, Advise, Assist)

Alcohol reduction counselling — refer to GP if dependent

Emphasise dry socket risk with smoking post-extraction

Regular recall and mucosal surveillance

🔴 **High-Yield ADC Exam Points**

Metronidazole + alcohol = disulfiram-like reaction — always counsel

Tobacco masks periodontal bleeding — disease worse than signs suggest

Tobacco + alcohol = multiplicative oral cancer risk

Nicotinic stomatitis = palate, pipe smokers, red dots = inflamed salivary gland openings



Smoker's melanosis = gingiva, brown, not premalignant

Dry socket risk significantly higher in smokers

Target SpO₂ 88-92% if COPD coexists (smoker)

Paracetamol in chronic alcoholics — hepatotoxicity risk, use minimum effective dose

Mock Questions (Author-written, for self-assessment)

1. **Which drug produces a disulfiram-like reaction when combined with alcohol?**

- A. Amoxicillin
- B. Metronidazole
- C. Paracetamol
- D. Lidocaine

Answer: B

2. **Why can periodontal disease appear less severe clinically than it truly is in smokers?**

- A. Smoking improves gingival health
- B. Nicotine-induced vasoconstriction masks bleeding on probing
- C. Smoking has no periodontal effect
- D. Smokers do not develop periodontitis

Answer: B

3. **Combined tobacco and alcohol use raises oral cancer risk in a manner best described as:**

- A. Unchanged compared to either alone
- B. Simply additive
- C. Multiplicative
- D. Protective

Answer: C

4. **A white, keratotic palatal lesion with red dots at minor salivary gland openings, seen in heavy smokers, is called:**

- A. Leukoplakia
- B. Nicotinic stomatitis
- C. Erythroplakia
- D. Lichen planus

Answer: B

5. **Alcohol increases bleeding risk mainly by:**

- A. Stimulating platelet aggregation
- B. Inhibiting thromboxane A₂, producing an antiplatelet effect
- C. Increasing clotting factor synthesis
- D. Having no effect on bleeding

Answer: B

6. **Which post-extraction complication occurs significantly more often in smokers?**

- A. Rapid, uneventful healing
- B. Dry socket (alveolar osteitis)
- C. Excess salivation
- D. Tooth whitening

Answer: B

7. **Which analgesic requires cautious, minimum-effective dosing in chronic heavy alcohol users because of hepatotoxicity risk?**

- A. Ibuprofen
- B. Paracetamol
- C. Amoxicillin
- D. Chlorhexidine

Answer: B



The Dentistry
— INSIGHT —

Chapter 11 — Hepatitis: A Complete Overview

🦠 Hepatitis - Complete Overview

Hepatitis = inflammation of the liver. Most commonly caused by viruses, but also drugs, alcohol, and autoimmune disease.

🦠 Types of Hepatitis

1. Viral Hepatitis (Most important)

Ⓐ Hepatitis A

Cause: Fecal-oral (contaminated food/water)

Type: Acute only (no chronic)

Clinical features:

Fever, malaise, nausea

Jaundice

Diagnosis: Anti-HAV IgM

Treatment: Supportive

Prevention: Vaccine

Ⓑ Hepatitis B

Cause: Blood, sexual, perinatal

Can be acute or chronic

Clinical features:

Jaundice, fatigue

Can lead to cirrhosis & liver cancer

Diagnosis markers:

HBsAg → infection

Anti-HBs → immunity

Treatment:

Acute: Supportive

Chronic: Antivirals (e.g. Tenofovir, Entecavir)

Prevention: Vaccine

Ⓒ Hepatitis C

Cause: Blood (IV drugs, transfusion)

Usually chronic

Clinical features:

Often asymptomatic early

Leads to cirrhosis

Diagnosis: Anti-HCV + PCR

Treatment:

Direct-acting antivirals (DAAs) → CURABLE

No vaccine

Ⓓ Hepatitis D

Requires Hepatitis B to survive

More severe disease

Ⓔ Hepatitis E

Fecal-oral

Dangerous in pregnancy (high mortality)

🦠 Non-Viral Hepatitis

1. Alcoholic Hepatitis

Cause: Chronic alcohol use

Features:

Jaundice, hepatomegaly

AST > ALT (classic)

2. Drug-induced Hepatitis

Common drug: Paracetamol overdose



The Dentistry
— I N S I G H T —

Others: antibiotics, anti-TB drugs

3. Autoimmune Hepatitis

Immune system attacks liver

Seen in young females

Treatment: steroids

△ Clinical Features (General)

Jaundice (yellow eyes/skin)

Dark urine

Pale stools

Fatigue

Right upper quadrant pain

🔍 Diagnosis

Liver function tests (ALT, AST ↑)

Bilirubin ↑

Viral serology

Ultrasound

! Complications

Cirrhosis

Liver failure

Hepatocellular Carcinoma

🦷 Dental Management (Important for You)

Before Treatment:

Check liver function

Check bleeding risk (INR)

Avoid elective treatment in acute hepatitis

Drug considerations:

Avoid high doses of Paracetamol

Avoid hepatotoxic drugs

Infection control:

Strict universal precautions

💊 Drug Summary (Mechanism + Key Points)

Tenofovir / Entecavir (HBV)

MOA: Inhibit viral DNA polymerase

Indication: Chronic HBV

Side effects: Renal toxicity (Tenofovir)

Paracetamol toxicity

MOA of toxicity: NAPQI accumulation → liver damage

Antidote: N-acetylcysteine

📊 Summary Table

Type

Transmission

Chronic?

Vaccine

A

Fecal-oral

✖

✔

B

Blood/sex

✔

✔



The Dentistry
— I N S I G H T —

C

Blood



✗

D

With B



Indirect

E

Fecal-oral

✗

Limited

🧠 Quick Memory Trick

A & E = "Eat" → food/water

B, C, D = Blood & Body fluids

If you want, I can give you: ✓💡Hepatitis serology interpretation (very important for exams)

✓💡OSCE-style dental management

✓💡MCQs & SEQs for ADC exam preparation

📖 Mock Questions (Author-written, for self-assessment)

1. **Which hepatitis virus is transmitted via the faecal-oral route and has no chronic form?**

- A. Hepatitis B
- B. Hepatitis A
- C. Hepatitis C
- D. Hepatitis D

Answer: B

2. **Hepatitis D can only establish infection in the presence of:**

- A. Hepatitis A
- B. Hepatitis B
- C. Hepatitis C
- D. No other virus is required

Answer: B

3. **Which hepatitis virus is potentially curable with direct-acting antivirals but currently has no vaccine?**

- A. Hepatitis A
- B. Hepatitis B
- C. Hepatitis C
- D. Hepatitis E

Answer: C

4. **Which serological marker indicates active Hepatitis B infection?**

- A. Anti-HBs
- B. HBsAg
- C. Anti-HAV IgM
- D. Anti-HCV


Answer: B

5. **Which hepatitis virus carries a particularly high mortality risk if contracted during pregnancy?**

- A. Hepatitis A
- B. Hepatitis C
- C. Hepatitis E
- D. Hepatitis D

Answer: C

6. **The classic laboratory pattern seen in alcoholic hepatitis is:**



- A. ALT greater than AST
- B. AST greater than ALT
- C. Both transaminases normal
- D. Decreased bilirubin

Answer: B

7. **The antidote for paracetamol-induced hepatotoxicity is:**

- A. Naloxone
- B. N-acetylcysteine
- C. Flumazenil
- D. Vitamin K

Answer: B



The Dentistry
— I N S I G H T —

Chapter 12 — Standard Precautions in Dental Infection Control (ADA/NHMRC Summary)

🦷 Infection Control in Dentistry (According to the Australian Dental Association & Australian Guidelines)

Based on ADA + NHMRC “Standard Precautions” (very high-yield for ADC exam)

📋 1. Standard Precautions (Apply to ALL patients)

- Assume every patient may carry infections like Hepatitis B, Hepatitis C, or Human Immunodeficiency Virus

Includes:

Hand hygiene

PPE

Sharps safety

Cleaning & sterilization

📋 2. Hand Hygiene (MOST IMPORTANT)

When?

Before & after every patient

After removing gloves

After contact with blood/saliva

How?

Soap + water (if visibly dirty)

Alcohol-based hand rub (routine)

📋 3. Personal Protective Equipment (PPE)

Gloves (single use)

Surgical mask

Protective eyewear / face shield

Gown (if splashing expected)

📌 4. Sharps Safety

Never recap needles (or use one-handed scoop)

Dispose immediately in sharps container

Prevent needle-stick injuries

🦷 5. Instrument Sterilization

Critical instruments (penetrate tissue)

MUST be sterilized (autoclave)

Semi-critical

Sterilize or high-level disinfection

Non-critical

Clean + disinfect

- Preferred method: Steam sterilization (autoclave)

🧼 6. Environmental Cleaning

Clean dental chair after each patient

Use disinfectants on surfaces

Barrier protection (covers for handles, light, etc.)

💧 7. Waterline Management

Flush dental unit waterlines

Use treated water

Prevent biofilm formation

🗑️ 8. Waste Management

Clinical waste → special disposal

Sharps → puncture-proof containers

Follow local regulations

📌 9. Vaccination of Dental Staff

Mandatory: Hepatitis B



The Dentistry
— I N S I G H T —

Recommended: Influenza, COVID-19

🦷 10. Post-Exposure Protocol

If needle-stick injury:

Wash immediately

Report incident

Assess risk

HBV → vaccine + HBIG if needed

△ 11. Transmission-Based Precautions (if needed)

For known infections:

Airborne (e.g. TB) → N95 mask

Droplet/contact precautions

🦷 ADC Exam Memory Trick

“HIPS SWaP”

Hand hygiene

Instruments sterilization

PPE

Sharps safety

Waste management

air/water control

Post-exposure care

🦷 Key Exam Points

Standard precautions for ALL patients

Autoclave = gold standard

Hand hygiene = most important step

HBV vaccination mandatory

Never recap needles

If you want next:

✓ 💎 OSCE checklist (step-by-step examiner marking scheme)

✓ 💎 MCQs on infection control

✓ 💎 Real clinical mistakes students make (very high yield)



📖 Mock Questions (Author-written, for self-assessment)

1. **The single most important infection control measure in the dental setting is:**

- A. Wearing gloves
- B. Hand hygiene
- C. Wearing a gown
- D. Using mouthwash

Answer: B

2. **Standard precautions should be applied to:**

- A. Only patients with a known infectious disease
- B. All patients, regardless of known infectious status
- C. Only surgical patients
- D. Only elderly patients

Answer: B

3. **The preferred sterilisation method for critical dental instruments is:**

- A. A chemical wipe-down
- B. Steam autoclaving
- C. UV light exposure alone
- D. An alcohol soak

Answer: B

4. **Which vaccination is considered mandatory for dental staff because of occupational blood exposure risk?**

- A. Hepatitis A

- B. Hepatitis B
- C. Influenza only
- D. No vaccination is required

Answer: B

5. **For a patient with a known airborne infection such as tuberculosis, which additional precaution is required?**
- A. A standard surgical mask
 - B. An N95/P2 respirator
 - C. Gloves alone
 - D. No additional precaution

Answer: B

6. **The correct method to avoid a needle-stick injury when recapping is to use:**
- A. A two-handed recapping technique
 - B. A one-handed scoop technique or a safety recapping device
 - C. No recapping tool is needed
 - D. Bending the needle first

Answer: B

7. **The first step after sustaining a needle-stick injury is to:**
- A. Ignore it and continue treatment
 - B. Wash the site immediately and report the incident
 - C. Wait until the end of the day to report it
 - D. Simply discard the needle

Answer: B



The Dentistry
— I N S I G H T —

Chapter 13 — Antiviral Medications in Dentistry

Below is a high-yield, exam-focused overview of major antiviral medications, with indications, contraindications, and side effects (ADC + clinical dentistry relevance).

1. Anti-Herpes Viruses

Acyclovir

★ Indications

Herpes simplex (HSV-1, HSV-2)

Herpes labialis (cold sores)

Herpes zoster (shingles)

Immunocompromised viral infections

⊗ Contraindications

Hypersensitivity

Severe renal impairment (dose adjust)

△ Side effects

Nausea, vomiting

Nephrotoxicity (crystal nephropathy)

Neurotoxicity (confusion, tremors in high dose)

Valacyclovir

(prodrug of acyclovir)

★ Indications

HSV infections

Zoster (better absorption than acyclovir)

⊗ Contraindications

Severe renal disease (dose adjustment needed)

△ Side effects

Headache

Nausea

Rare renal toxicity

Famciclovir

★ Indications

HSV and VZV infections

⊗ Contraindications

Hypersensitivity

△ Side effects

Headache

GI upset

2. Anti-Hepatitis B Drugs

Tenofovir

★ Indications

Chronic Hepatitis B

HIV (combination therapy)

⊗ Contraindications

Severe renal failure (relative)

Hypersensitivity

△ Side effects



Nephrotoxicity
Reduced bone mineral density
GI upset

📖 **Entecavir**

✦ **Indications**

Chronic HBV infection

🚫 **Contraindications**

HIV monotherapy (resistance risk)

⚠️ **Side effects**

Headache
Fatigue
Lactic acidosis (rare)

📖 **3. Anti-Hepatitis C Drugs (DAAs)**

📖 **Sofosbuvir**

✦ **Indications**

Chronic Hepatitis C (curative therapy)

🚫 **Contraindications**

Severe liver failure (depends on combo regimen)

⚠️ **Side effects**

Fatigue
Headache
Nausea
📖 Other DAAs (e.g. Ledipasvir, Velpatasvir)

✦ **Indications**

HCV combination therapy

⚠️ **Side effects**

Mild GI upset
Headache

📖 **4. Anti-HIV Drugs**

📖 **Zidovudine (AZT)**

✦ **Indications**

HIV infection
Post-exposure prophylaxis

🚫 **Contraindications**

Severe bone marrow suppression

⚠️ **Side effects**

Anemia
Neutropenia
Myopathy
📖 Tenofovir (also used in HIV)

✦ **Indications**

HIV, HBV

⚠️ **Side effects**

Renal toxicity
Bone loss

📖 **5. Anti-Influenza Drugs**



The Dentistry
— I N S I G H T —

📖 Oseltamivir

✦ Indications

Influenza A & B

🚫 Contraindications

Hypersensitivity

⚠️ Side effects

Nausea, vomiting

Rare neuropsychiatric effects

💎 QUICK EXAM SUMMARY TABLE

Drug

Use

Key Side Effect

Acyclovir

HSV

Nephrotoxicity

Valacyclovir

HSV

Headache

Tenofovir

HBV/HIV

Kidney damage

Entecavir

HBV

Lactic acidosis

Sofosbuvir

HCV

Fatigue

Zidovudine

HIV

Anemia

Oseltamivir

Flu

GI upset

🔑 DENTAL CLINIC PEARLS (ADC IMPORTANT)

Delay elective treatment in active HSV lesions

HBV patients may be on Tenofovir/Entecavir

Adjust doses in liver/renal disease

Avoid nephrotoxic combinations (important with acyclovir + NSAIDs)

Always apply standard precautions

💎 MEMORY TRICK

“A V T E S Z O”

Acyclovir

Valacyclovir

Tenofovir

Entecavir

Sofosbuvir

Zidovudine

Oseltamivir

If you want next, I can give: ✔️ 💎 MCQs + SEQs (ADC style antiviral exam set)

✔️ 💎 Drug interactions in dentistry (very high yield)

✔️ 💎 Viral diseases + dental management combined OSCE cases



The Dentistry
— I N S I G H T —

 Mock Questions (Author-written, for self-assessment)

1. **Acyclovir's main dose-related toxicity concern is:**

- A. Hepatotoxicity
- B. Nephrotoxicity (crystal nephropathy)
- C. Cardiotoxicity
- D. Ototoxicity

Answer: B

2. **Valacyclovir is best described as:**

- A. An entirely new, independent antiviral
- B. A prodrug of acyclovir with improved oral absorption
- C. An antibacterial agent
- D. An antifungal agent

Answer: B

3. **First-line drugs for chronic Hepatitis B include:**

- A. Sofosbuvir and Ledipasvir
- B. Tenofovir and Entecavir
- C. Zidovudine and Oseltamivir
- D. Famciclovir and Valacyclovir

Answer: B

4. **Direct-acting antivirals such as sofosbuvir are used to treat, and can potentially cure:**

- A. Hepatitis B
- B. Hepatitis C
- C. HIV
- D. Influenza

Answer: B

5. **Zidovudine (AZT), used in HIV management, has a notable side effect of:**

- A. Hepatotoxicity
- B. Anaemia and neutropenia (bone marrow suppression)
- C. Nephrotoxicity
- D. Dental staining

Answer: B

6. **Oseltamivir is indicated for the treatment of:**

- A. Herpes zoster
- B. Influenza A and B
- C. Hepatitis C
- D. HIV

Answer: B

7. **A key dental precaution with acyclovir is avoiding concurrent use with which nephrotoxic drug class?**

- A. Paracetamol
- B. NSAIDs
- C. Amoxicillin
- D. Chlorhexidine

Answer: B



Chapter 14 — Liver Cirrhosis: A High-Yield Overview

📖 Liver Cirrhosis (Cirrhosis) — Complete High-Yield Overview (ADC + Dentistry Focus)

Cirrhosis is a chronic, irreversible scarring of the liver where normal liver tissue is replaced by fibrous tissue → leading to liver failure and portal hypertension.

🔑 1. Causes (Etiology)

🍷 Alcohol-related

Chronic alcohol abuse (most common in many countries)

🦠 Viral hepatitis

Chronic Hepatitis B

Chronic Hepatitis C

🍖 Other causes

Fatty liver disease (NAFLD/NASH)

Autoimmune hepatitis

Biliary diseases (e.g. primary biliary cholangitis)

Drugs/toxins

⚙️ 2. Pathophysiology (Simple)

Chronic liver injury

Inflammation

Fibrosis (scar formation)

Nodular regeneration

Cirrhosis → liver failure + portal hypertension

⚠️ 3. Clinical Features

🕒 Early (compensated)

Fatigue

Mild hepatomegaly

Mild abnormal LFTs

🕒 Advanced (decompensated)

Jaundice

Ascites (fluid in abdomen)

Easy bruising / bleeding

Peripheral edema

Hepatic encephalopathy (confusion, drowsiness)

Spider angiomas

Palmar erythema

🔑 4. Major Complications

🚰 Portal hypertension

Esophageal varices → life-threatening bleeding

Splenomegaly

🧠 Hepatic encephalopathy

↑ ammonia → brain dysfunction

🦠 Liver cancer

Hepatocellular Carcinoma

🩸 Coagulopathy

↓ clotting factors → bleeding tendency

🔑 5. Diagnosis

LFTs (AST, ALT ↑ early; may normalize later)

Low albumin

↑ INR (very important)

Ultrasound / CT



Liver biopsy (confirmatory)

6. Treatment

General

Treat cause (stop alcohol, antivirals for HBV/HCV)

Nutritional support

Complication management

Ascites → diuretics (spironolactone)

Varices → beta blockers / band ligation

Encephalopathy → lactulose

7. Dental Management (VERY IMPORTANT FOR ADC)

BEFORE TREATMENT

Check:

INR (bleeding risk)

Platelets

Liver function

- Avoid elective procedures in:

Decompensated cirrhosis

DURING TREATMENT

Use atraumatic technique

Good local hemostasis:

Pressure packs

Sutures

Tranexamic acid mouthwash if needed

DRUG CONSIDERATIONS

⊘ Avoid / reduce dose:

Paracetamol (limit dose)

NSAIDs (↑ bleeding + kidney risk)

Sedatives (prolonged effect due to poor metabolism)

△ Key issue:

Liver cannot metabolize drugs properly → toxicity risk

POST-OPERATIVE CARE

Monitor bleeding

Avoid hepatotoxic drugs

Ensure follow-up

8. Key Exam Points (ADC High Yield)

Cirrhosis = irreversible liver scarring

Main risks:

Bleeding (↑ INR)

Infection

Drug toxicity

HBV/HCV are major causes

Always check coagulation profile before dental surgery

MEMORY TRICK

“CIRRHOSIS = C.I.R.R.H.O.S.I.S”

Chronic liver damage

Inflammation

Regeneration nodules

Risk bleeding

Hepatic failure

Oedema/ascites



The Dentistry
— I N S I G H T —

Serious infection risk

Increased INR

Survival reduced

If you want next, I can give: ✓ ♦ MCQs + SEQs (ADC exam cirrhosis set)

✓ ♦ OSCE dental management script (step-by-step examiner answer)

✓ ♦ Hepatitis → cirrhosis → cancer progression diagram (very high yield)

📖 Mock Questions (Author-written, for self-assessment)

1. **Which laboratory value is most important to check before dental surgery in a patient with liver cirrhosis?**

- A. Blood glucose
- B. INR (International Normalised Ratio)
- C. Amylase
- D. TSH

Answer: B

2. **Ascites, jaundice and hepatic encephalopathy are features of which stage of cirrhosis?**

- A. Compensated
- B. Decompensated
- C. Early/subclinical
- D. None of the above

Answer: B

3. **Which analgesic requires strict dose limitation in patients with cirrhosis?**

- A. Amoxicillin
- B. Paracetamol
- C. Chlorhexidine
- D. Fluoride varnish

Answer: B

4. **Elevated blood ammonia levels in cirrhosis are linked to which complication?**

- A. Gingival hyperplasia
- B. Hepatic encephalopathy
- C. Xerostomia
- D. Dental erosion

Answer: B

5. **NSAIDs are generally cautioned against in cirrhotic patients mainly because of:**

- A. Taste alteration
- B. Increased bleeding risk and renal impairment
- C. Tooth staining
- D. Allergic reactions only

Answer: B

6. **Elective dental treatment should generally be avoided in which state of cirrhosis?**


- A. Compensated, stable cirrhosis
- B. Decompensated cirrhosis
- C. Any cirrhosis, regardless of severity
- D. It is always safe, regardless of stage

Answer: B

7. **A useful local haemostatic measure after an extraction in a cirrhotic patient is:**

- A. Tranexamic acid mouthwash
- B. An aspirin rinse
- C. Loading with warfarin
- D. No local measures are needed

Answer: A



Chapter 15 — Chronic Kidney Disease (ESKD) and the Dialysis Patient

Step 1: The Pathophysiological Foundation (The "Why")

What is ESKD? It's the final stage of chronic kidney disease (CKD Stage 5), where the kidneys have lost ~85-90% of their function. The glomerular filtration rate (GFR) is $<15 \text{ mL/min/1.73m}^2$. The patient is likely on dialysis (hemodialysis or peritoneal dialysis) or awaiting a transplant.

The Systemic Derangements that Affect the Mouth:

Uremia: Accumulation of nitrogenous waste products (urea, creatinine) in the blood.

Metabolic Acidosis: Kidneys can't excrete acid or reabsorb bicarbonate.

Electrolyte Imbalances: Hyperkalemia (high K^+), hyperphosphatemia (high PO_4), hypocalcemia (low Ca^{++}).

Anemia: Due to decreased erythropoietin (EPO) production.

Platelet Dysfunction: Uremic toxins impair platelet adhesion and aggregation (NOT a low platelet count, but poor function).

Immune Dysfunction: Impaired chemotaxis and phagocytosis of neutrophils and macrophages.

Altered Drug Metabolism & Excretion: Many drugs are renally cleared.

Step 2: The Oral & Dental Complications (The "What")

These are the direct consequences of the systemic derangements above. Memorize this list.

A. Soft Tissue & Mucosal Complications (Uremic Stomatitis)

Uremic Stomatitis: A painful, erythematous, and sometimes ulcerative or pseudomembranous lesion on the buccal mucosa, floor of mouth, and tongue. It's a direct toxic effect of high salivary urea, which is converted to ammonia by bacterial urease. The ammonia burns the tissue.

Clinical Pearl: It often resolves dramatically after dialysis or a kidney transplant.

Xerostomia (Dry Mouth): Extremely common. Caused by:

Fluid restriction for dialysis patients.

Medications (antihypertensives, diuretics).

Direct effect of uremia on salivary glands.

Consequence: Rampant caries, candidiasis, dysgeusia (bad taste).

Dysgeusia / Metallic Taste: "Uremic fetor" – a urine-like or ammonia-like breath and taste.

Pale Mucosa: Due to renal anemia.

Petechiae, Ecchymoses, Gingival Bleeding: Due to platelet dysfunction. This is a major surgical concern.

Oral Candidiasis: Common due to immune dysfunction, xerostomia, and frequent antibiotic use.

Lichenoid Reactions: Can be drug-induced (e.g., from antihypertensives) or related to the underlying disease.

B. Hard Tissue & Periodontal Complications

Periodontal Disease: This is the most significant dental complication. ESKD patients have a higher prevalence and severity of periodontitis. Why?

Immune Dysfunction: Poor host response to bacterial plaque.

Altered Bone Metabolism: Renal osteodystrophy (see below) weakens the alveolar bone.

Xerostomia: Loss of salivary buffering and antimicrobial protection.

Systemic Inflammation: Periodontitis is a source of systemic inflammation that worsens cardiovascular risk in these patients.

Renal Osteodystrophy (ROD): A complex bone disease. Two main types:

High-Turnover (Osteitis Fibrosa Cystica): Due to secondary hyperparathyroidism (high PTH from low Ca^{++} and high PO_4). Leads to bone resorption, "brown tumors" (giant cell lesions in jaws), and loss of lamina dura (classic radiographic sign).

Low-Turnover (Adynamic Bone Disease): Due to oversuppression of PTH (from medications like calcitriol or cinacalcet). Bone is brittle and heals poorly.

Clinical Implication: Pathologic jaw fracture is a real risk during extractions or implant placement.

Enamel Hypoplasia: If the patient had CKD during childhood (when teeth were developing), you may see horizontal lines, pits, or discoloration on the permanent teeth.

Delayed Eruption: Common in pediatric CKD patients.

Pulp Calcifications: Increased incidence of pulp stones and obliteration of the pulp chamber.

C. Complications Related to Dialysis

Heparinization: During hemodialysis, patients receive heparin to prevent clotting in the circuit. Do not perform any invasive dental procedure within 4-6 hours after dialysis due to the risk of prolonged bleeding.

Arteriovenous (AV) Fistula/Graft: This is the patient's lifeline. Never take blood pressure, draw blood, or place an IV on the arm with the fistula. Protect it from pressure during dental procedures.

Hepatitis B & C: Higher prevalence in dialysis patients. Use universal precautions and consider vaccination status.

Step 3: The Clinical Management Protocol (The "How")

This is the step-by-step approach you must follow for every ESKD patient.

Phase 1: Pre-Treatment (The Consultation)

Medical History & Communication:

Contact the Nephrologist. This is non-negotiable for any invasive procedure.

Confirm: Dialysis schedule, dry weight, medications (especially anticoagulants, antihypertensives, immunosuppressants), and recent labs.

Essential Lab Work (Order these):

CBC: Check for anemia (Hgb < 10 g/dL may require pre-op transfusion or EPO).

BUN & Creatinine: To assess uremia level.

Electrolytes: K⁺ (risk of arrhythmia), PO₄, Ca⁺⁺.

Coagulation Profile: PT/INR and PTT are often normal. The bleeding risk is from platelet dysfunction, not clotting factors. Bleeding Time is the best test, but it's rarely done. A surrogate is the Platelet Function Assay (PFA-100).

Hepatitis B/C & HIV Serology.

Timing of Treatment:

Ideal Day: The day after dialysis. The patient is well-dialyzed (lower uremia, better platelet function) and the heparin has worn off.

Never treat on the day of dialysis (heparin risk) or the day before (high uremia, fluid overload).

Phase 2: The Dental Procedure

Antibiotic Prophylaxis?

Not routinely indicated for dental procedures in ESKD patients unless they have a specific cardiac condition (e.g., prosthetic valve, history of infective endocarditis, cardiac transplant with valvulopathy).

However, consider prophylaxis for patients with peritoneal dialysis catheters or central venous catheters (e.g., tunneled dialysis catheter) due to risk of seeding the catheter. Consult the nephrologist.

Choice: Amoxicillin 2g PO 1 hour before. If allergic, Clindamycin 600mg.

Bleeding Management (The Critical Step):

For minor procedures (scaling, simple filling): Usually no special measures.

For surgical procedures (extraction, biopsy, implant):

Local Hemostasis: Use local anesthetic with vasoconstrictor (epinephrine) – it's safe and helps control bleeding.

Surgical Technique: Atraumatic surgery. Use a periosteal elevator carefully. Suture the socket (resorbable sutures like Vicryl).

Topical Agents: Place Gelfoam, Surgicel, or Collaplug in the socket.

Tranexamic Acid (TXA): This is your best friend. 5% TXA solution. Rinse mouth for 2 minutes, 4 times a day for 5-7 days post-op. It inhibits fibrinolysis and stabilizes the clot.

Desmopressin (DDAVP): For severe bleeding risk. 0.3 mcg/kg IV or intranasal. It releases von Willebrand factor and factor VIII, temporarily correcting platelet dysfunction. Must be ordered by the nephrologist.

Anesthetic Considerations:

Lidocaine: Safe. Max dose is standard (7 mg/kg with epi).

Articaine: Safe.

Avoid Prilocaine: Risk of methemoglobinemia in renal patients.

Avoid Mepivacaine 3% (plain): Metabolized in the liver, but its metabolites are renally cleared. Use with caution.

Avoid NSAIDs (Ibuprofen, Naproxen, Ketorolac): ABSOLUTELY CONTRAINDICATED. They reduce renal blood flow and worsen kidney function. They also increase bleeding risk. Use Acetaminophen (Paracetamol) or Opioids (e.g., Tramadol, Oxycodone) for pain. Note: Tramadol dose must be adjusted for

Mock Questions (Author-written, for self-assessment)

1. **Dental treatment for a haemodialysis patient is best scheduled:**

- A. On the day of dialysis
- B. The day before dialysis
- C. The day after dialysis
- D. Any day; timing makes no difference

Answer: C

2. **The primary mechanism behind the bleeding risk in end-stage kidney disease (ESKD) is:**

- A. A low platelet count
- B. Qualitative platelet dysfunction caused by uraemic toxins
- C. Excess clotting factors

D. Vitamin K deficiency

Answer: B

3. **Which structure must never be used for blood pressure measurement or venepuncture in a dialysis patient?**

- A. The dominant hand
- B. The arm containing the arteriovenous (AV) fistula
- C. The leg
- D. The earlobe

Answer: B

4. **Uraemic stomatitis is thought to result from:**

- A. A vitamin deficiency
- B. Bacterial conversion of salivary urea to ammonia
- C. Excess fluoride exposure
- D. An allergy to dental materials

Answer: B

5. **High-turnover renal osteodystrophy (from secondary hyperparathyroidism) can cause which classic radiographic finding in the jaws?**

- A. Increased lamina dura
- B. Loss of lamina dura and "brown tumours"
- C. Enamel hypoplasia only
- D. No radiographic changes

Answer: B

6. **Which drug class is contraindicated in ESKD patients due to worsening renal function and increased bleeding risk?**

- A. Paracetamol
- B. NSAIDs
- C. Amoxicillin
- D. Lidocaine

Answer: B

7. **Which local anaesthetic should be avoided in renal patients because of the risk of methemoglobinemia?**

- A. Lidocaine
- B. Prilocaine
- C. Articaine
- D. Mepivacaine with epinephrine

Answer: B



The Dentistry
—INSIGHT—

Chapter 16 — Allergy Management in Dentistry

The Universal Management Algorithm (The "4 R's")

Before we dive into specifics, understand this core framework. Every allergy management case follows these steps:

Recognize: Identify the allergen (via history, clinical exam, and patch testing).

Remove: Eliminate the offending material from contact with the patient.

Replace: Choose a biocompatible alternative material.

Re-evaluate: Confirm the lesion resolves and the new material is tolerated.

Now, let's apply this to each major allergen.

1. Management of Nickel Allergy

This is the most common metal allergy you will manage.

Step 1: Confirm the Diagnosis. The lichenoid reaction directly adjacent to a metal restoration is highly suggestive. Refer for patch testing to confirm nickel (and rule out cobalt/chromium co-sensitivity).

Step 2: The "Remove" Decision. This is critical.

Asymptomatic Lichenoid Reaction (White, lacy, no pain): You do NOT need to remove the restoration. The risk of a severe reaction is low. Monitor annually. Explain to the patient that it's a benign, localized immune response.

Symptomatic Lichenoid Reaction (Burning, pain, erosion, ulceration): You MUST remove the restoration. The patient's quality of life is affected.

Step 3: The "Replace" Strategy.

For a Crown/Bridge: Replace with a nickel-free alternative.

Gold Alloy (Type III or IV): Excellent biocompatibility, but expensive and aesthetic.

Zirconia: My preferred choice. Full-contour zirconia or zirconia-based crown. No metal, no allergy risk.

Lithium Disilicate (e.g., e.max): Excellent aesthetics, but requires a resin cement. Ensure the patient is not also allergic to acrylates (see below).

Titanium: For implant abutments or crowns, but less common for routine crowns.

For a Partial Denture Framework: Replace with a flexible, metal-free alternative.

Valplast (Nylon-based): Excellent for small partials.

Acetal Resin: More rigid than nylon.

Zirconia Framework: The ultimate, but expensive.

For Orthodontic Brackets: Switch to ceramic brackets (polycrystalline or monocrystalline alumina) or gold-plated brackets (if the patient is not allergic to gold).

For Stainless Steel Crowns (Pediatric): Use zirconia pediatric crowns or pre-veneered stainless steel crowns (but ensure the veneer material is also biocompatible).

Step 4: Re-evaluate. The lesion should resolve within 2-4 weeks. If it persists, consider a co-allergy (e.g., to the cement or the new material).

2. Management of Acrylate (Methacrylate) Allergy

This is the most challenging to manage, both for the patient and the clinician.

Step 1: Confirm the Diagnosis. Patch testing with a specific acrylate series is mandatory. Do not use the patient's own composite for

Glass Ionomer Cement (GIC): Can be used to cement zirconia or lithium disilicate crowns. It bonds chemically to the tooth and mechanically to the ceramic.

Zinc Phosphate Cement: The oldest, most biocompatible cement. No monomers. Excellent for metal or zirconia crowns.

Resin Cement: AVOID. All resin cements contain acrylates.

The "Temporary" Solution:

Pre-formed Aluminum or Stainless Steel Crowns: For posterior teeth.

GIC-based Temporary Crowns: Some newer materials are GIC-based and acrylate-free.

Step 4: The "Occupational" Management (For the Dental Professional).

Prevention is everything. Once sensitized, you cannot be cured.

No-touch technique: Use brush tips, capsules, and syringes. Never touch uncured monomer with your gloved hand.

Gloves: Nitrile gloves only. Latex and vinyl offer zero protection against acrylates. Change gloves frequently.

Ventilation: Use a high-volume evacuator (HVE) and a room air purifier.

Barrier Creams: Some are available, but not a substitute for physical barriers.

If you are already sensitized: You may need to change your career path (e.g., move to oral surgery, prosthodontics with a focus on metal-free, or academia).

3. Management of Fragrance/Balsam of Peru Allergy

This is often overlooked but very common.

Step 1: Confirm the Diagnosis. Patch testing.

Step 2: The "Remove" Decision. You don't remove a restoration. You remove the product.

Step 3: The "Replace" Strategy.

Toothpaste: Switch to an unflavored, fragrance-free toothpaste (e.g., Biotene, Sensodyne ProNamel, or a prescription fluoride gel).

Mouthwash: Use unflavored, alcohol-free mouthwash.

Prophylaxis Paste: Use unflavored pumice or a non-flavored prophylaxis paste.

Topical Anesthetic: Use benzocaine-free or eugenol-free alternatives. Lidocaine gel is usually safe.

Temporary Cement: Avoid ZOE (Zinc Oxide Eugenol). Use a non-eugenol temporary cement (e.g., TempBond NE, RelyX Temp NE).

Impression Materials: Avoid those with flavoring. Use unflavored silicone or polyether.

Step 4: Re-evaluate. The cheilitis and stomatitis should resolve within 1-2 weeks.

4. Management of Latex Allergy

This is a medical emergency risk.

Step 1: Identify the Type.

Type I (Immediate, IgE-mediated): LIFE-THREATENING. Strict avoidance is mandatory.

Type IV (Delayed, Contact Dermatitis): Less dangerous, but still requires avoidance.

Step 2: The "Remove" Decision. Remove all latex products from the operatory.

Step 3: The "Replace" Strategy.

Gloves: Use powder-free, low-protein nitrile gloves or neoprene gloves. Never use latex gloves on a latex-allergic patient.

Rubber Dam: Use silicone rubber dam (e.g., OptiDam, Dental Dam from Coltène/Whaledent).

Prophylaxis Cups: Use silicone or non-latex prophylaxis cups.

Impression Materials: Use silicone or polyether. Avoid latex-containing impression materials.

Other Items: Check all materials (e.g., some matrix bands, some orthodontic elastics). Use a latex-free emergency kit.

Step 4: The "Emergency" Protocol.

Have a latex-free emergency kit with epinephrine, antihistamines, and a bronchodilator.

Train your staff on how to recognize and manage anaphylaxis.

Document the allergy prominently in the patient's chart and on a medical alert bracelet.

5. Management of Gold Allergy

Step 1: Confirm the Diagnosis. Patch testing.

Step 2: The "Remove" Decision. If symptomatic (lichenoid reaction, pain), remove the gold restoration.

Step 3: The "Replace" Strategy.

Zirconia: My top choice. Full-contour zirconia or zirconia-based crown.

Lithium Disilicate: Excellent aesthetics.

Titanium: For implant abutments.

Precious Metal Alloy (e.g., Platinum-Palladium): If the patient is not allergic to these metals.

Step 4: Re-evaluate. The lesion should resolve.

6. Management of Cobalt/Chromium Allergy

Step 1: Confirm the Diagnosis. Patch testing (often co-positive with nickel).

Step 2: The "Remove" Decision. If symptomatic, remove the Co-Cr restoration.

Step 3: The "Replace" Strategy.

Titanium: The best alternative for frameworks and implant abutments.

Zirconia: For crowns and bridges.

Gold Alloy: If the patient is not allergic to gold.

Mock Questions (Author-written, for self-assessment)

1. **An asymptomatic lichenoid reaction seen adjacent to a nickel-containing restoration should generally be managed by:**

- A. Immediate removal of the restoration
- B. Monitoring only, since removal is not required if the lesion is symptom-free
- C. Extraction of the tooth
- D. A course of antibiotics

Answer: B

2. **A preferred nickel-free crown material offering excellent aesthetics and biocompatibility is:**

- A. Nickel-chromium alloy
- B. Zirconia
- C. Amalgam
- D. Stainless steel

Answer: B

3. **For a patient allergic to acrylates (methacrylates), which cement should be avoided?**

- A. Glass ionomer cement
- B. Resin cement
- C. Zinc phosphate cement
- D. None; all are equally safe

Answer: B

4. **The correct glove choice for a patient with a Type I (IgE-mediated) latex allergy is:**

- A. Latex gloves
- B. Nitrile or neoprene gloves
- C. Vinyl gloves only
- D. No gloves at all

Answer: B

5. **The best replacement for a latex rubber dam in a latex-allergic patient is:**

- A. Vinyl sheeting
- B. A silicone rubber dam
- C. A paper dam
- D. No isolation is possible

Answer: B

6. **A fragrance/Balsam of Peru allergy is best managed by switching the patient to:**


- A. A more strongly flavoured mouthwash
- B. Unflavoured, fragrance-free dental products
- C. A stronger antiseptic mouthwash
- D. No change is needed

Answer: B

7. **Cobalt/chromium allergy most commonly co-occurs with sensitivity to:**

- A. Gold
- B. Nickel
- C. Titanium
- D. Zirconia

Answer: B



The Dentistry
— INSIGHT —

Chapter 17 — Severe Odontogenic Infection and Airway Emergencies (Ludwig's Angina)

Image 1:

Step 1: Recognition - Signs and Symptoms of Severe Infection

You must differentiate a severe infection from a simple, localized abscess. The key is systemic involvement and anatomical spread.

A. Systemic Signs (SIRS Criteria - Systemic Inflammatory Response Syndrome)

- Fever: Core temperature $>38^{\circ}\text{C}$ or $<36^{\circ}\text{C}$.
- Tachycardia: Heart rate >90 bpm.
- Tachypnoea: Respiratory rate >20 breaths/min or $\text{PaCO}_2 <32$ mmHg.
- Leukocytosis or Leukopenia: WBC $>12,000/\mu\text{L}$ or $<4,000/\mu\text{L}$, or $>10\%$ immature bands.
- General malaise, rigors, and confusion (especially in elderly).

B. Local Signs of Severity (Anatomical Danger Zones)

- Trismus: Inability to open mouth $>20\text{mm}$ (indicates deep masticator space involvement).
- Dysphagia: Difficulty swallowing (indicates lateral pharyngeal or retropharyngeal space).
- Dysphonia: "Hot potato" voice or muffled voice (indicates peritonsillar or parapharyngeal space).

Image 2:

Step 3: Definitive Medical Management (According to Australian Therapeutic Guidelines)

A. Antibiotic Therapy (Empiric, IV, High Dose)

- First Line (Non-Penicillin Allergic):
 - Benzylpenicillin 1.2g (2 million units) IV every 4 hours PLUS
 - Metronidazole 500mg IV every 8 hours.
- Penicillin Allergy (Non-Anaphylactic):
 - Cephalothin 2g IV every 6 hours PLUS Metronidazole 500mg IV every 8 hours.
- Penicillin Allergy (Anaphylactic):
 - Clindamycin 600mg IV every 8 hours PLUS
 - Gentamicin (single daily dose, e.g., 5-7mg/kg IV) - requires renal monitoring.
- If severe sepsis or immunocompromised:
 - Piperacillin + Tazobactam (Tazocin) 4.5g IV every 6 hours OR
 - Meropenem 1g IV every 8 hours.

B. Surgical Intervention (The Definitive Treatment)

- Timing: Urgent (within 2-4 hours of diagnosis) in an operating theatre under general anaesthesia.
- Procedure:

Image 3:

B. Local Signs of Severity (Anatomical Danger Zones)

- Trismus: Inability to open mouth $>20\text{mm}$ (indicates deep masticator space involvement).
- Dysphagia: Difficulty swallowing (indicates lateral pharyngeal or retropharyngeal space).
- Dysphonia: "Hot potato" voice or muffled voice (indicates peritonsillar or parapharyngeal space).
- Drooling: Inability to manage secretions (impending airway obstruction).
- Floor of mouth elevation: "Woody" or brawny induration of the submental/submandibular region (Ludwig's Angina).
- Neck swelling: Extending below the mandible or into the anterior neck.
- Cervical lymphadenopathy: Tender, firm, matted nodes.

C. Red Flags for Airway Compromise (The "Can't Intubate, Can't Ventilate" scenario)

- Stridor (high-pitched breathing sound).
- Tripod positioning (patient sitting upright, leaning forward, hands on knees).
- Tachypnoea with accessory muscle use (intercostal, supraclavicular retractions).

Mock Questions (Author-written, for self-assessment)

1. **Trismus limiting mouth opening to less than 20 mm suggests involvement of which anatomical space?**
 - A. Buccal space only
 - B. The deep masticator space

- C. The vestibular space
- D. No specific space

Answer: B

2. **The first-line empiric IV antibiotic regimen for a severe odontogenic infection in a non-penicillin-allergic patient is:**
- A. Oral amoxicillin alone
 - B. Benzylpenicillin IV plus metronidazole IV
 - C. Oral doxycycline
 - D. A chlorhexidine rinse alone

Answer: B

3. **Bilateral submandibular and sublingual space infection with "woody" induration of the floor of the mouth describes:**
- A. A simple periapical abscess
 - B. Ludwig's angina
 - C. Dry socket
 - D. Pericoronitis

Answer: B

4. **A "hot potato", muffled voice in a patient with a spreading dental infection suggests involvement of the:**
- A. Buccal space
 - B. Peritonsillar/parapharyngeal space
 - C. Vestibular space
 - D. No specific space

Answer: B

5. **The definitive treatment for a severe, spreading odontogenic infection is:**
- A. Antibiotics alone
 - B. Urgent surgical drainage under general anaesthesia, combined with antibiotics
 - C. Review again in one week
 - D. Extraction is contraindicated in all cases


Answer: B

6. **Which clinical sign indicates impending airway obstruction and requires immediate escalation of care?**
- A. Mild facial swelling
 - B. Stridor and tripod positioning
 - C. A slight fever
 - D. A normal voice

Answer: B

7. **For a patient with a severe infection and a history of anaphylactic penicillin allergy, an appropriate antibiotic regimen includes:**
- A. Amoxicillin
 - B. Clindamycin plus gentamicin
 - C. Metronidazole alone
 - D. No antibiotics are required

Answer: B



The Dentistry

— INSIGHT —

Chapter 18 — Hyperventilation Syndrome

Hyperventilation syndrome (HVS) is a condition where a person breathes faster or deeper than normal without a clear physical need (like exercise). This leads to excessive loss of carbon dioxide (CO₂) from the blood, which can cause a range of symptoms.

🦋 What's happening in the body?

When you breathe too rapidly:

CO₂ levels drop (called hypocapnia)

Blood pH becomes more alkaline (respiratory alkalosis)

This affects nerve and muscle function

⚠️ Common symptoms

Shortness of breath (even though oxygen is usually normal)

Dizziness or lightheadedness

Tingling in fingers, toes, or around the mouth

Chest tightness or pain

Rapid heartbeat

Feeling anxious or panicky

HVS is often linked with stress or anxiety, but it can also occur without obvious emotional triggers.

🦋 Causes & triggers

Anxiety or panic attacks

Stress or emotional distress

Pain or illness

Habitual over-breathing

Sometimes no clear cause

🦋 Diagnosis

Doctors usually diagnose HVS by:

Reviewing symptoms and medical history

Ruling out other serious conditions (like asthma, heart problems, or pulmonary embolism)

Sometimes measuring blood gases

🦋 Treatment & management

Breathing control techniques (slow, diaphragmatic breathing)

Relaxation strategies (meditation, mindfulness)

Cognitive behavioral therapy (CBT) if anxiety is a factor

In some cases, treatment of underlying conditions

🦋 When to seek help

Get medical attention if symptoms:

Occur suddenly and severely

Include chest pain, fainting, or confusion

Happen for the first time (to rule out serious causes)

If you want, I can show you simple breathing exercises to manage hyperventilation in real time.



The Dentistry
INSIGHT —

📖 Mock Questions (Author-written, for self-assessment)

1. **Hyperventilation leads to excessive loss of which gas from the blood?**

- A. Oxygen
- B. Carbon dioxide
- C. Nitrogen
- D. Nitrous oxide

Answer: B

2. **The acid-base disturbance that results from hyperventilation is:**

- A. Metabolic acidosis
- B. Respiratory alkalosis
- C. Metabolic alkalosis
- D. Respiratory acidosis

Answer: B

3. **A classic symptom combination in hyperventilation syndrome includes dizziness, chest tightness and:**

- A. Bradycardia
- B. Perioral or peripheral tingling
- C. Bleeding gums
- D. Sudden deafness

Answer: B

4. **The first-line management of an acute hyperventilation episode in the dental chair is:**

- A. High-flow oxygen
- B. Calm reassurance and slow, controlled diaphragmatic breathing
- C. Intramuscular adrenaline
- D. Immediate hospital transfer

Answer: B

5. **Hyperventilation syndrome is most commonly associated with:**

- A. An underlying infection
- B. Anxiety or panic
- C. Diabetes
- D. An allergy to local anaesthetic

Answer: B

6. **A first-time episode of hyperventilation should prompt medical review, rather than reassurance alone, if it is accompanied by:**

- A. Nothing in particular; reassurance is always sufficient
- B. Chest pain, fainting, or confusion
- C. Simply asking for a glass of water
- D. Any anxiety at all

Answer: B

7. **A key differential diagnosis to exclude in a patient who is hyperventilating in the dental chair is:**

- A. Dental caries
- B. Asthma, a cardiac event, or pulmonary embolism
- C. Gingivitis
- D. Bruxism

Answer: B



The Dentistry
— I N S I G H T —

Chapter 19 — Pharmacology of Type 2 Diabetes Medications

Major Drug Classes for Type 2 Diabetes

1. Biguanides

Main Drug

Metformin

Mechanism



Actions

Reduces liver glucose production

Improves insulin sensitivity

Decreases intestinal glucose absorption

Indications

First-line treatment for type 2 diabetes

Prediabetes

Polycystic ovary syndrome

Contraindications

Severe renal failure

Severe liver disease

Alcohol abuse

Risk of lactic acidosis

Side Effects

Diarrhea

Nausea

Metallic taste

Vitamin B12 deficiency

Rare lactic acidosis

2. Sulfonylureas

Drugs

Gliclazide

Glimepiride

Glibenclamide

Mechanism



Side Effects

Hypoglycemia

Weight gain

Contraindications

Severe liver disease

Severe renal disease

Type 1 diabetes

3. Meglitinides

Drugs

Repaglinide

Nateglinide

Mechanism

Stimulate rapid insulin release from pancreas.

Important Point

Short acting

Taken before meals

Side Effects

Hypoglycemia

Weight gain



4. Thiazolidinediones (TZDs)

Drugs

Pioglitazone

Mechanism



Side Effects

Weight gain

Fluid retention

Heart failure worsening

Fracture risk

Contraindications

Heart Failure

Liver disease

5. DPP-4 Inhibitors (Gliptins)

Drugs

Sitagliptin

Linagliptin

Vildagliptin

Mechanism



Advantages

Weight neutral

Low hypoglycemia risk

Side Effects

Nasopharyngitis

Pancreatitis (rare)

6. GLP-1 Receptor Agonists

Drugs

Semaglutide

Liraglutide

Dulaglutide

Mechanism



Benefits

Weight loss

Cardiovascular protection

Side Effects

Nausea

Vomiting

Pancreatitis

7. Alpha-Glucosidase Inhibitors

Drug

Acarbose

Mechanism



Side Effects

Flatulence

Diarrhea

Abdominal discomfort

8. Insulin

Types

Rapid Acting

Insulin Lispro



The Dentistry
— I N S I G H T —

Insulin Aspart
Short Acting
Regular Insulin
Intermediate Acting
NPH Insulin
Long Acting
Insulin Glargine
Insulin Detemir
Mechanism



Side Effects
Hypoglycemia
Weight gain
Lipodystrophy

9. Amylin Analog

Drug
Pramlintide
Mechanism
Slows gastric emptying
Suppresses glucagon
Increases satiety
Simplified Comparison Table
Drug Class
Weight Effect
Hypoglycemia Risk
Main Benefit
Metformin
Loss/neutral
Low
First-line
Sulfonylureas
Gain
High
Cheap
SGLT2 inhibitors
Loss
Low
Heart/kidney benefit
GLP-1 agonists
Major loss
Low
Weight reduction
DPP4 inhibitors
Neutral
Low
Safe in elderly
Insulin
Gain
High
Strong glucose lowering
3 MCQs

MCQ 1

Which antidiabetic drug commonly causes vitamin B12 deficiency?



The Dentistry
— I N S I G H T —

- A. Insulin**
- B. Metformin**
- C. Acarbose**
- D. Sitagliptin**

✓ **Answer: B. Metformin**

MCQ 2

Which diabetic drug class commonly causes weight loss?

- A. Sulfonylureas**
- B. Insulin**
- C. GLP-1 agonists**
- D. Pioglitazone**

✓ **Answer: C. GLP-1 agonists**

MCQ 3

Which drug has highest risk of hypoglycemia?

- A. Metformin**
- B. DPP4 inhibitors**
- C. Sulfonylureas**
- D. Acarbose**

✓ **Answer: C. Sulfonylureas**

6 SEQs (Scenario-Based)

SEQ 1

A 55-year-old obese diabetic patient needs weight reduction and cardiovascular benefit.

Best drug?

✓ **Semaglutide**

Why?

Causes significant weight loss

Cardiovascular protection

Extracted MCQs

GLP-1 agonists mainly cause:

- A. Weight gain**
- B. Weight loss** ✓
- C. Hypercalcemia**
- D. Bradycardia**

Which is injectable?


- A. Metformin**
- B. Semaglutide** ✓
- C. Acarbose**
- D. Gliclazide**

SEQ 2

An elderly diabetic develops repeated hypoglycemia on glimepiride.

Cause?

✓ **Sulfonylurea-induced insulin excess**



The Dentistry
— I N S I G H T —

Management?

Reduce dose

Switch to safer drug like DPP4 inhibitor

Extracted MCQs

Sulfonylureas increase:

A. Glucagon

B. Insulin secretion ✓

C. Ketones

D. Cortisol

Major side effect:

A. Weight loss

B. Hypoglycemia ✓

C. Dry mouth

D. Constipation

SEQ 3

A diabetic with heart failure requires glucose-lowering therapy.

Avoid?

✓ **Pioglitazone**

Why?

Fluid retention worsens heart failure.

Extracted MCQs

Pioglitazone acts on:

A. GLP receptors

B. PPAR-gamma ✓

C. SGLT2

D. DPP4

TZDs may cause:

A. Fluid retention ✓

B. Hearing loss

C. Hyperthyroidism

D. Bradycardia

SEQ 4

A patient on metformin develops severe renal failure.

Concern?

✓ **Lactic acidosis risk**

Action?

Stop metformin.

Extracted MCQs

Metformin is contraindicated in:

A. Renal failure ✓

B. Asthma

C. Migraine

D. Psoriasis



The Dentistry
— I N S I G H T —

Rare severe complication:

A. Ketoacidosis

B. Lactic acidosis ✓

C. Hypernatremia

D. Hypercalcemia

SEQ 5

A diabetic patient on insulin becomes sweaty and confused during dental extraction.

Diagnosis?

✓ **Hypoglycemia**

Immediate treatment?

Oral glucose if conscious

Glucagon/IV dextrose if unconscious

Extracted MCQs

Most dangerous insulin side effect:

A. Hypertension

B. Hypoglycemia ✓

C. Hyperpigmentation

D. Hyperthermia

Sweating and confusion suggest:

A. Stroke

B. Hypoglycemia ✓

C. Asthma

D. Allergy

SEQ 6

A patient taking acarbose complains of bloating and flatulence.

Cause?

Undigested carbohydrates fermented in colon.

Extracted MCQs

Acarbose acts in:

A. Liver

B. Intestine ✓

C. Kidney

D. Pancreas

Common side effect:

A. Hair loss

B. Flatulence ✓

C. Bradycardia

D. Hematuria



The Dentistry
— I N S I G H T —

Chapter 20 — Hypoglycaemia: Recognition, Emergency Management and Drug Considerations in Diabetic Patients

Experienced excessive physical exertion or stress (dental anxiety can burn glucose).

Less Common: Non-diabetic patients with reactive hypoglycemia, liver disease, or starvation.

Step 2: The Clinical Presentation (The "How to Spot It")

This is the most critical step for you. You must differentiate hypoglycemia from other emergencies (e.g., syncope, stroke, panic attack).

The "Classic" Triad (Whipple's Triad):

Symptoms of hypoglycemia.

Low plasma glucose.

Resolution of symptoms after glucose administration.

Signs & Symptoms (in order of progression):

Stage | Autonomic (Early) | Neuroglycopenic (Late - Danger!)

Subjective | Hunger, weakness, sweating, palpitations, anxiety, tremor | Confusion, difficulty speaking, blurred vision, dizziness
| Pallor, | Slurred speech, dis

Do not assume it's a panic attack. Panic attacks are usually dry, not diaphoretic.

Step 3: The Immediate Management Protocol (The "What to Do")

This is a step-by-step, time-sensitive protocol. Do not deviate.

Phase 1: Recognition & Assessment (0-30 seconds)

Stop the procedure.

Remove all instruments from the mouth.

Assess the patient's level of consciousness (AVPU):

Alert?

Verbal response?

Pain response?

Unresponsive?

Check vital signs (pulse, respiratory rate). In hypoglycemia, pulse is usually rapid and weak (tachycardia).

Phase 2: Intervention (30 seconds - 2 minutes)

Scenario A: Patient is Conscious, Alert, and Can Swallow (Mild-Moderate)

Goal: Administer fast-acting oral glucose.

The "15-15 Rule":

Give 15 grams of fast-acting carbohydrate.

Wait 15 minutes.

Re-assess symptoms.

Sources of 15g Glucose (choose one):

4 oz (120 ml) of fruit juice or regular soda (not diet).

1 tablespoon of sugar or honey (placed under the tongue or in the buccal pouch).

3-4 glucose tablets (chewed).

1 tube of glucose gel (e.g., Insta-Glucose).

Do NOT give: Chocolate, cake, ice cream (fat slows absorption).

Scenario B: Patient is Confused, Drowsy, or Unconscious (Severe)

Goal: Do NOT give anything by mouth. Risk of aspiration is high.

Action:

Activate Emergency Medical Services (EMS) / Call 911.

Position: Place patient in the recovery position (left lateral recumbent) to protect airway.

Administer Glucagon (if available in your emergency kit):


Glucagon (1 mg) : Given intramuscularly (IM) or subcutaneously (SQ). It stimulates the liver to release stored glucose.

Intranasal Glucagon (Baqsimi) : A newer, easier option. Administer 3 mg into one nostril.

Monitor: Check pulse and breathing continuously. Be prepared to start CPR if they become pulseless.

Phase 3: Post-Intervention (After 15 minutes)

If symptoms resolve: The patient is stable. Offer a longer-acting snack (e.g., crackers with peanut butter, a sandwich) to prevent



recurrence. Do not dismiss them. Monitor for 30 minutes.

If symptoms do NOT resolve: This is a medical emergency. Repeat the 15g glucose (if conscious) or activate EMS again. The patient may need IV dextrose.

Step 4: The Post-Grad / Clinical Pearls (The "Expert Level")

Prevention is Paramount:

Medical History: Always ask diabetic patients: "When did you last eat? When did you take your medication?"

Morning Appointments: Schedule diabetic patients in the morning, shortly after breakfast and their usual medication.

The "Sick Day" Rule: If a patient is NPO for a procedure, they should not take their usual dose of insulin or sulfonylurea. Consult with their physician.

The "Dental Chair" Differential:

Hypoglycemia vs. Syncope (Fainting): Syncope is usually preceded by a prodrome of warmth, nausea, and "tunnel vision." The patient is pale but not sweaty (they are vasovagal, not adrenergic). Pulse is slow and weak (bradycardia), not fast.

Hypoglycemia vs. Stroke: A stroke is usually a sudden, focal neurological deficit (e.g., one-sided weakness, facial droop). Hypoglycemia is global (confusion, whole-body tremor). If in doubt, give glucose. It won't hurt a stroke patient, but it will save a hypoglycemic one.

The "Hidden" Hypoglycemic:

Patients on beta-blockers (e.g., propranolol, metoprolol) may not exhibit the classic autonomic symptoms

3. Drug Considerations in Diabetic Patients

Analgesics

Paracetamol — safe, first line

NSAIDs — use with caution — renal impairment common in long-standing diabetes

Avoid NSAIDs in diabetics with nephropathy

Antibiotics

Diabetics have impaired immune response — infections spread more rapidly

Lower threshold for antibiotic prescribing in poorly controlled diabetics

Amoxicillin first line — standard doses

Local Anaesthesia

Lignocaine with adrenaline — safe in well-controlled diabetics

Adrenaline in LA causes transient increase in blood glucose — clinically insignificant at dental doses

Do not withhold adrenaline — adequate anaesthesia reduces stress response which is more harmful than the adrenaline itself

Corticosteroids

Systemic corticosteroids cause significant hyperglycaemia in diabetics

If prescribing dexamethasone for swelling — warn patient and physician

Monitor BGL if corticosteroids used

4. Specific Drug Classes — Dental Implications

Metformin

Most common Type 2 diabetes drug

Lactic acidosis risk with contrast media — relevant for radiology referrals

No direct dental procedural implications

Safe to continue for routine dental treatment

Sulfonylureas (e.g., glipizide, gliclazide)

Stimulate insulin release — hypoglycaemia risk especially if patient skips meals

If NPO — do not take sulfonylurea on day of procedure

Higher hypoglycaemia risk than metformin

Insulin

Multiple types — rapid, intermediate, long-acting

Morning appointment after breakfast and usual insulin dose — safest

If fasting for procedure — physician must adjust insulin dose

Always have glucose available chairside

SGLT2 Inhibitors (e.g., empagliflozin, dapagliflozin, canagliflozin)

Withhold on day of surgery — risk of euglycaemic DKA

Euglycaemic DKA — DKA occurs with normal blood glucose — easily missed

Increased UTI and genital fungal infection risk — relevant for immunocompromised state

Consult physician before any surgical procedure



GLP-1 Receptor Agonists (e.g., semaglutide, liraglutide, dulaglutide)

Slow gastric emptying — aspiration risk under sedation or GA

Relevant when planning IV sedation or general anaesthesia

Inform anaesthetist — extended fasting may be required

No direct hypoglycaemia risk as monotherapy

Mock Questions (Author-written, for self-assessment)

1. **Whipple's triad, used to confirm hypoglycaemia, includes symptoms of hypoglycaemia, low plasma glucose and:**

- A. Resolution of symptoms after glucose administration
- B. A normal ECG reading
- C. A raised body temperature
- D. A normal blood pressure

Answer: A

2. **The "15-15 rule" for managing mild-to-moderate hypoglycaemia means:**

- A. Wait 15 minutes, then give 15 puffs of a bronchodilator
- B. Give 15 g of fast-acting carbohydrate, wait 15 minutes, then reassess
- C. Give 15 units of insulin
- D. Check blood glucose every 15 hours

Answer: B

3. **In an unconscious or severely confused hypoglycaemic patient, the correct action is to:**

- A. Give oral glucose gel
- B. Give nothing by mouth; consider intramuscular/intranasal glucagon and call emergency services
- C. Offer a chocolate bar
- D. Administer an antihistamine

Answer: B

4. **The typical dose of intramuscular glucagon carried in an emergency kit is:**

- A. 1 mg
- B. 10 mg
- C. 100 mg
- D. 0.1 mg

Answer: A

5. **A key feature distinguishing hypoglycaemia from vasovagal syncope in the dental chair is that hypoglycaemia typically causes:**

- A. Bradycardia and pallor, just like syncope
- B. Tachycardia and profuse sweating (diaphoresis), unlike the bradycardia and pallor of syncope
- C. Exactly the same presentation as syncope
- D. A slower onset than syncope

Answer: B

6. **SGLT2 inhibitors are typically withheld before surgery mainly because of the risk of:**


- A. Hypoglycaemia
- B. Euglycaemic diabetic ketoacidosis
- C. Hypertension
- D. Bradycardia

Answer: B

7. **The best time to schedule a diabetic patient's dental appointment, to reduce hypoglycaemia risk, is:**

- A. Late afternoon, after skipping lunch
- B. Morning, shortly after breakfast and their usual medication
- C. Evening, on an empty stomach
- D. Timing makes no difference

Answer: B



The Dentistry
— INSIGHT —

Chapter 21 — The ALARA Principle and Radiation Safety in Dental Radiology

ALARA Principle in Dental Radiology

ALARA = As Low As Reasonably Achievable

It means radiation exposure to the patient, dentist, and staff should be kept as low as possible while still obtaining a diagnostic-quality radiograph.

It is a core principle in dental radiology and radiation protection.

Main ALARA Principles in Dentistry

1. Justification

Only take radiographs when clinically necessary.

Examples:

Caries detection

Impacted teeth

Periapical pathology

Trauma assessment

Periodontal bone loss

Do NOT take unnecessary routine X-rays without indication.

2. Optimization

Use techniques that reduce radiation dose while maintaining image quality.

Methods include:

Patient Protection

Thyroid collar

Lead apron (especially in children/pregnancy where indicated)

Correct positioning

Avoid repeat exposures

Equipment Measures

Rectangular collimation

High-speed receptors/digital sensors

Proper filtration

Long cone paralleling technique

Lowest effective exposure settings

Operator Protection

Stand at least 2 meters away

Position at 90–135° from primary beam

Never hold the receptor during exposure



3. Dose Limitation

Radiation dose should remain below recommended safety limits for staff and public.

Dental patients have no strict dose limit because exposure is based on clinical need, but exposure must still follow ALARA.

Important Radiation Reduction Methods

Method

Radiation Reduction

Digital radiography

Lower dose than film

Rectangular collimation

Major dose reduction

Fast film (F-speed)

Lower exposure

Thyroid collar

Protects thyroid gland

Proper technique

Prevents retakes

CBCT only when necessary

Avoid unnecessary high dose

According to Australian Dental Guidelines

Radiographs should be based on individual patient assessment.

Routine “screening” X-rays without indication are discouraged.

Use smallest field of view in CBCT.

Children require extra radiation caution due to higher sensitivity.

Biological Effects of Radiation

Deterministic Effects

Threshold-dependent:

Skin burns

Cataracts

Tissue damage

Rare in dental radiology.

Stochastic Effects

No threshold:

Cancer risk

Genetic mutation

Even small doses may theoretically increase risk, therefore ALARA is essential.

Dental Considerations

Before taking radiographs:

Check pregnancy status when relevant

Confirm clinical indication

Use previous radiographs if adequate

Minimize repeat images

Explain benefits vs risks to patient

MCQs

MCQ 1

What does ALARA stand for?

A. As Long As Radiation Allows

B. As Low As Reasonably Achievable

C. As Little As Radiation Applies

D. As Long As Required Always

Answer:

B. As Low As Reasonably Achievable

MCQ 2

Which technique significantly reduces patient radiation dose?

A. Round collimation

B. Increased exposure time

C. Rectangular collimation

D. Holding sensor by hand

Answer:

C. Rectangular collimation

MCQ 3

Which imaging modality usually gives lower radiation dose?

A. Conventional CT

B. CBCT

C. Digital intraoral radiograph

D. MRI



The Dentistry
— I N S I G H T —

Answer:

C. Digital intraoral radiograph

SBQs (Scenario-Based Questions)

SBQ 1

A dentist takes bitewing radiographs at every visit without examination.

Question:

Which ALARA principle is violated?

Answer:

Justification — radiographs should only be taken when clinically indicated.

SBQ 2

A radiograph must be repeated because of poor positioning.

Question:

Which ALARA concept was not followed?

Answer:

Optimization — proper technique reduces repeat exposure.

SBQ 3

A dentist orders a large-field CBCT for a simple periapical lesion.

Question:

What should have been done?

Answer:

Use the smallest appropriate field of view and lower-dose imaging if adequate.

Additional MCQs from SBQs

MCQ 4

Unnecessary routine radiographs violate:

A. Optimization

B. Justification

C. Dose limitation

D. Filtration

Answer:

B. Justification

MCQ 5

Repeat exposures mainly increase:

A. Contrast

B. Radiation dose

C. Sensor speed

D. Resolution only

Answer:

B. Radiation dose

MCQ 6

For CBCT imaging, ALARA recommends:

A. Largest field possible

B. Maximum exposure

C. Smallest suitable field of view

D. Multiple repeat scans

Answer:





The Dentistry
— I N S I G H T —