

Some amasing formulas

These days I study Hilbnert Theorem of non-immersion of Lobachevsky plane in \mathbf{E}^3 . To my surprise it has so many mathphysical applications. Here I just give some very preliminary results. (see the file in Etudes/Geometry/Lobachevsky)

Consider on \mathbf{R}^2 Riemannian metric

$$G = dx^2 + 2\cos\Theta(x,y)dxdy + dy^2$$

This metric naturally appears in the proof of Hilbert Theorem.

Performing routine calculations we come to

$$\Gamma_{xx}^x = \Theta_x \cotan \Theta, \quad \Gamma_{xx}^y = -\Theta_x \frac{1}{\sin \Theta},$$

$$\Gamma_{yy}^y = \Theta_y \cotan \Theta, \quad \Gamma_{yy}^x = -\Theta_y \frac{1}{\sin \Theta},$$

and

$$\Gamma_{xy}^x = \Gamma_{yx}^x = \Gamma_{xy}^y = \Gamma_{yx}^y = 0$$

and

$$K = \frac{R}{2} = R_{1212} \det g = \frac{-\Theta_{xy} \sin \Theta}{1 - \cos^2 \Theta} = \frac{-\Theta_{xy}}{\sin \Theta},$$

and in particular the condition

$$\Theta_{xy} = \sin \Theta$$

means that this manifold is (whole?) Lobachesky plane (this function has to exist)

E.g. function

$$\theta = \arctan Ce^{(x+y)}$$