

Homework 2

1

- a) Show that $(\mathbf{x}, \mathbf{y}) = x^1 y^1 + x^2 y^2 + x^3 y^3$ defines a scalar product in \mathbf{R}^3 .
- b) Show that $(\mathbf{x}, \mathbf{y}) = x^1 y^1 + x^2 y^2$ does not define a scalar product in \mathbf{R}^3 .
- c) Show that $(\mathbf{x}, \mathbf{y}) = x^1 y^1 + x^2 y^2 - x^3 y^3$ does not define a scalar product in \mathbf{R}^3 .
- d) Show that $(\mathbf{x}, \mathbf{y}) = x^1 y^1 + 3x^2 y^2 + 5x^3 y^3$ defines a scalar product in \mathbf{R}^3 .

e[†]) Find necessary and sufficient conditions for entries a, b, c of symmetrical matrix $\begin{pmatrix} a & b \\ b & c \end{pmatrix}$ such that the formula

$$(\mathbf{x}, \mathbf{y}) = (x^1, x^2) \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} y^1 \\ y^2 \end{pmatrix} = ax^1 y^1 + b(x^1 y^2 + x^2 y^1) + cx^2 y^2$$

defines scalar product in \mathbf{R}^2 .

2 The matrix $A = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ obeys the conditions $A^T A = I$. Show that

- a) $\det A = \pm 1$
- b) if $\det A = 1$ then there exists an angle $\varphi : 0 \leq \varphi < 2\pi$ such that $A = A_\varphi$ where

$$A_\varphi = \begin{pmatrix} \cos \varphi & -\sin \varphi \\ \sin \varphi & \cos \varphi \end{pmatrix} \quad (\text{rotation matrix})$$

c) if $\det A = -1$ then there exists an angle $\varphi : 0 \leq \varphi < 2\pi$ such that $A = A_\varphi R$, where $R = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ (a reflection matrix).

3 Show that for matrix A_φ defined in the previous exercise the following relations are satisfied:

$$A_\varphi^{-1} = A_\varphi^T = A_{-\varphi}, \quad A_{\varphi+\theta} = A_\varphi \cdot A_\theta.$$

4 Show that under the transformation $(\mathbf{e}'_1, \mathbf{e}'_2) = (\mathbf{e}_1, \mathbf{e}_2) A_\varphi$ an orthonormal basis transforms to an orthonormal one.

How coordinates of vectors change if we rotate the orthonormal basis $(\mathbf{e}_1, \mathbf{e}_2)$ on the angle $\varphi = \frac{\pi}{3}$ anticlockwise?

5 Let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ be an orthonormal basis of Euclidean space \mathbf{E}^3 . Consider the ordered set of vectors $\{\mathbf{e}'_1, \mathbf{e}'_2, \mathbf{e}'_3\}$ which is expressed via basis $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ as in the exercise 6 of the Homework 1.

Write down explicitly transition matrix from the basis $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ to the ordered set of the vectors $\{\mathbf{e}'_1, \mathbf{e}'_2, \mathbf{e}'_3\}$. What is the rank of this matrix? Is this matrix orthogonal?

Find out is the ordered set of vectors $\{\mathbf{e}'_1, \mathbf{e}'_2, \mathbf{e}'_3\}$ a basis in \mathbf{E}^3 . Is this basis an orthonormal basis of \mathbf{E}^3 ? (you have to consider all cases a), b) c) and d)).

6[†]. Show that an arbitrary orthogonal transformation of two-dimensional Euclidean space can be considered as a composition of reflections.

7[†] Prove the Cauchy–Bunyakovsky–Schwarz inequality

$$(\mathbf{x}, \mathbf{y})^2 \leq (\mathbf{x}, \mathbf{x})(\mathbf{y}, \mathbf{y}),$$

where \mathbf{x}, \mathbf{y} are arbitrary two vectors and $(\ , \)$ is a scalar product in Euclidean space.

Hint: For any two given vectors \mathbf{x}, \mathbf{y} consider the quadratic polynomial $At^2 + 2Bt + C$ where $A = (\mathbf{x}, \mathbf{x})$, $B = (\mathbf{x}, \mathbf{y})$, $C = (\mathbf{y}, \mathbf{y})$. Show that this polynomial has at most one real root and consider its discriminant.