

Homework 3

In all exercises we assume by default that Riemannian metric on embedded surfaces is induced by the Euclidean metric.

1 a) Show that surface of the cone $\begin{cases} x^2 + y^2 - k^2 z^2 = 0 \\ z > 0 \end{cases}$ in \mathbf{E}^3 is locally isometric to Euclidean plane.

2 a) Consider the domain D on the conic surface $x^2 + y^2 - z^2$ defined by the conditions

$$\begin{cases} 0 < z < H \\ y \neq 0 \quad \text{if } x > 0 \end{cases}.$$

(The second condition means that the line $x = z, y = 0$ is removed from the surface of the cone)

Find a domain D' in Euclidean plane such that it is isometric to the surface D .

b) Find a shortest distance between points $A = (0, 1, 1)$ and $B = (0, -1, 1)$ for an ant living on the surface D (we assume that $H > 1$).

3 Consider plane with Riemannian metric given in cartesian coordinates (x, y) by the formula

$$G = \frac{4(dx)^2 + 4(dy)^2}{(1 + x^2 + y^2)^2}.$$

Show that this Riemannian manifold is locally isometric to the sphere.

4 Consider catenoid: $x^2 + y^2 = \cosh^2 z$ and helicoid: $y - x \tan z = 0$.

Find induced Riemannian metrics on these surfaces.

Show that these surfaces are locally isomorphic.

5 a) Consider the domain D on the cone $x^2 + y^2 - k^2 z^2$ defined by the condition $0 < z < H$. Find an area of this domain using induced Riemannian metric. Compare with the answer when using standard formulae.

6 a) Find an area of 2-dimensional sphere of radius R using explicit formulae for induced Riemannian metric in stereographic coordinates.

b)[†] Find a volume of n -dimensional sphere of radius R . (You may use Riemannian metric in stereographic coordinates, or you may do it in other way... You just have to calculate the answer.)