

Let $S(x, q) = S^a(x)q_a + \dots$, then for arbitrary function $g(y)$, the inverse image under the thick morphism Φ_S , $f(x) = \Phi_S^*(g)$ is equal to

$$f(x) = \lim_{\hbar \rightarrow 0} \frac{\hbar}{i} \log \left[\int e^{\frac{i}{\hbar}(g(y) + S(x, q) - y^a q_a)} \mathcal{D}(y, q) \right].$$

Thus

$$\begin{aligned} \Phi_S^*(g + \varepsilon h) &= \lim_{\hbar \rightarrow 0} \frac{\hbar}{i} \log \left[\int e^{\frac{i}{\hbar}(g(y) + S(x, q) - y^a q_a + \varepsilon \mathbf{h}(\mathbf{y}))} \mathcal{D}(y, q) \right] = \\ &= \lim_{\hbar \rightarrow 0} \frac{\hbar}{i} \log \left[\int e^{\frac{i}{\hbar}(g(y) + S(x, q) - y^a q_a)} \left(1 + \frac{i\varepsilon}{\hbar} h(y) \right) \mathcal{D}(y, q) \right] = \\ &\quad \Phi_S^*(g) + \frac{\int e^{\frac{i}{\hbar}(g(y) + S(x, q) - y^a q_a)} \varepsilon h(y) \mathcal{D}(y, q)}{\int e^{\frac{i}{\hbar}(g(y) + S(x, q) - y^a q_a)} \mathcal{D}(y, q)}. \end{aligned}$$

We come to

$$\Phi_S^*(g + \varepsilon h) - \Phi_S^*(g) = \varepsilon \frac{\int e^{\frac{i}{\hbar}(g(y) + S(x, q) - y^a q_a)} h(y) \mathcal{D}(y, q)}{\int e^{\frac{i}{\hbar}(g(y) + S(x, q) - y^a q_a)} \mathcal{D}(y, q)}.$$

Here we have to be carefull with "concurrence" between small ε and big $\frac{1}{\hbar}$. We hae to care about the cobntribution of amplitudes...

Hovik 28 November 2018.