

# The Future of Air Quality in the Developing World

Jessica Neu<sup>1</sup>, Daniel Jacob<sup>2</sup>, and Vivienne Payne<sup>1</sup> on behalf of the GEO-CAPE Atmospheric Science Working Group<sup>3</sup>

<sup>1</sup> Jet Propulsion Laboratory, California Institute of Technology, <sup>2</sup> Harvard University, <sup>3</sup><http://geo-cape.larc.nasa.gov/docs/team/AtmosphericScienceWorkingGroup.pdf>

## ***What are the key challenges or questions for Earth System Science across the spectrum of basic research, applied research, applications, and/or operations in the coming decade?***

Attendees of the 2014 NASA workshop on Outstanding Questions in Atmospheric Composition identified the future of air quality in the developing world as the number one priority for tropospheric composition research over the next few decades. As noted in the workshop report [[https://espo.nasa.gov/home/content/NASA\\_SMD\\_Workshop](https://espo.nasa.gov/home/content/NASA_SMD_Workshop)], the developing world is experiencing rapid industrialization in combination with high population density and limited emission controls, leading to serious air quality problems [Cohen *et al.*, 2005]. A recent study showed that more than 4000 people die every day in China from particulate matter (PM) exposure [Rohde and Muller, 2015], and given that ozone has also been linked to premature mortality [Jerrett *et al.*, 2009, 2015; Lelieveld *et al.*, 2015], the total death toll associated with air pollution is likely even higher. While China receives much of the international attention on this issue, other locations in South Asia and Africa also have dangerously poor air quality [e.g. Tawari and Abowei, 2012; Lioussé *et al.*, 2014]. In the tropics, smoke from biomass burning is becoming a major public health issue as it affects population centers [Pavagadhi *et al.*, 2013]. Worldwide, air pollution is believed to kill more people than AIDS, malaria, breast cancer, or tuberculosis [WHO, 2014; WHO, 2012; O’Keefe, 2013; Yang *et al.*, 2013]. Underscoring the importance of this topic, the GEO-CAPE Atmospheric Science Working Group (ASWG) endorsed it as a critical research and applications need at a recent community workshop. This ASWG represents a broad cross-section of ~35 scientists from universities and government laboratories who have contributed to pre-formulation studies for the air quality component of the GEO-CAPE mission recommended by the 2007 Decadal Survey [<http://geo-cape.larc.nasa.gov/>].

The 2014 NASA workshop report identified four science questions associated with air quality in the developing world, paraphrased here and listed in order of importance:

1. What are the emissions associated with major sources of pollution in the developing world and how are these emissions related to near-surface ozone and PM?
2. How will changing patterns of energy-related and agricultural emissions driven by urbanization affect local and regional air quality and what are the consequences for global tropospheric composition?
3. Can we develop a space-based observing system for global air quality, with spatial and temporal resolution relevant for determining public exposure to ozone and PM, that can be used to issue air quality warnings?
4. How do extreme air quality events form and evolve in these understudied regions and how do they affect global composition?

### ***Why are these challenges/questions timely to address now especially with respect to readiness?***

The issue of air quality in the developing world is extremely timely because the next few decades are crucial in terms of providing scientists and policy makers with the information they need to monitor, understand, and reduce air pollution in order to protect populations and to achieve concomitant short-term climate benefits. There has been much progress made in recent years toward the application of satellite measurements to air quality issues, as exemplified by the NASA AqAST team [<http://acmg.seas.harvard.edu/aqast>], which has pioneered new techniques and approaches for utilizing measurements from the current generation of low-Earth orbiting (LEO) satellites to aid air quality managers at the local, state, and national level. Further, advances in multispectral retrievals have demonstrated capability for near-surface measurements of carbon monoxide and ozone [Worden *et al.*, 2010, 2013; Fu *et al.*, 2013; Cuesta *et al.*, 2013], with the potential to provide better understanding of the effects of air pollution on humans and ecosystems. Most critically, NASA (through the GEO-CAPE ASWG) and ESA have started to work with the air quality community to prepare for and exploit the expected high volume of high spatiotemporal resolution measurements of ozone, aerosols, and precursor species from the upcoming geostationary Earth orbit (GEO) missions TEMPO (NASA), GEMS (KARI), and Sentinel 4 (ESA). This “virtual constellation” has led to a set of well-defined measurement requirements for space-based observations of air quality [e.g. Fishman *et al.*, 2012; GEO-CAPE, 2015] and is expected to revolutionize the use of satellite measurements for air quality applications and operational services. However, it will be focused on the developed regions of the Northern midlatitudes. There is a critical need for similar high spatiotemporal resolution measurements focused on developing nations in the tropics and subtropics.

Precedent for partnership between NASA and governments in the developing world has already been established through the NASA – U.S. Agency for International Development (USAID) SERVIR program [[http://www.nasa.gov/mission\\_pages/servir/overview.html](http://www.nasa.gov/mission_pages/servir/overview.html)]. SERVIR provides satellite data, predictive models and science applications to help improve environmental decision-making among developing nations in parts of Africa, the Hindu-Kush region, and Southeast Asia. SERVIR strengthens the ability of governments and other stakeholders to incorporate Earth observations and geospatial technologies to respond to natural disasters, improve food security, safeguard human health, and manage natural resources. While the quantitative application of satellite data to air quality decision-making is currently out of the scope of SERVIR, the US-based input to air quality management for the Beijing Olympic Games [Streets *et al.*, 2007] provides an effective model for implementation.

### ***Why are space-based observations fundamental to addressing these challenges / questions?***

To answer the science questions described above, we must be able to quantify emissions from various sources and connect them to chemical processes and the subsequent production of ozone and PM near the surface, where they impact human health. Sparse surface measurements are insufficient for this purpose, particularly in the developing world where surface networks are rudimentary at best. Aircraft campaigns, which can be difficult to organize in developing regions, provide anecdotal data that cannot easily be generalized to larger spatiotemporal scales. Space-based measurements are thus critical because they provide the spatial coverage and long, continuous record required to address these science questions under a variety of environmental conditions.

The current generation of low-earth orbiting (LEO) satellites have provided a wealth of valuable measurements for quantifying emissions [e.g. Jiang *et al.*, 2015; Marais *et al.*, 2014; Gu *et al.*, 2014;

*Huang et al., 2014; Ghude et al., 2013; Barkley et al., 2015; Richards et al., 2013; Kim et al., 2013]* and understanding the processes controlling mid-tropospheric ozone [e.g. *Verstraeten et al., 2015; Neu et al., 2014; Huang et al., 2015*]. The GEO-CAPE pre-formulation studies, however, have shown that once-per-day LEO measurements are insufficient for monitoring and understanding air quality. These studies indicate that observations must be made multiple times a day, at horizontal resolutions of ~10 km or finer, and with as much near-surface sensitivity as possible in order to separate emissions sources at the urban scale, quantify the contribution of transported pollution, understand diurnal and day-to-day variability in emissions and chemical processing, and provide sufficient measurement statistics to enable quantitative studies in cloudy regions [e.g. *Fishman et al., 2012*].

The TEMPO, Sentinel-4, and GEMS geostationary satellites will achieve many of the measurements needed to address these goals over the regions surrounding the United States, Europe, and Korea, respectively, but large swaths of the developing world will remain uncovered. Furthermore, none of these missions will incorporate TIR measurements that help to identify biomass burning plumes (e.g. peroxyacetyl nitrate and formic acid [*Alvarado et al., 2011*]) and constrain agricultural emissions (e.g. ammonia [*Zhu et al., 2013*] and, when combined with UV/VIS measurements, provide the most robust sensitivity to near-surface ozone [*Natraj et al., 2011; Zoogman et al., 2011*].

The Sentinel 5P/5, METOP, and JPSS satellites will provide global LEO coverage of many of the relevant species once per day, but will lack the spatiotemporal resolution needed to understand and quantify air pollution processes. Therefore a space-based mission is needed that measures critical air quality and biomass burning species multiple times per day, with as much coverage as possible over South and Central America, Africa, and South Asia (where many current and emerging megacities are located). There are several potential implementation approaches for such a mission, including additional GEO instruments to complement the spatial coverage of Sentinel 4, TEMPO, and GEMS; companion LEO instruments to complement the temporal coverage of Sentinel 5, METOP, and JPSS; and a set of 2 LEO orbiters in opposing near-equatorial orbits covering 30°S-30°N. The geostationary option offers the advantage of a hosted payload approach using planned geostationary communications satellites. Since these satellites provide commercial services, operation of air quality instruments for the benefit of the developing world could conceivably be supported by public/private partnerships including philanthropic organizations. NASA and other space agencies could partner with governments in the developing world to build native capacity for air quality monitoring and prediction featuring the operational use of these measurements.

A space-based mission to study air quality in the developing world would serve the air quality, public health, and chemistry-climate interactions communities and provide a wealth of scientific and public health-relevant information to allow us to answer the critical science questions described here. Space-based observations will foster interest in aircraft campaigns and ground-based measurements to obtain additional correlative data. While such a mission would not be focused on the US, it serves national interests by providing information that can help to improve societal and political stability around the world and can be used to quantify and help reduce the long-range transport of air pollution. With the addition of CH<sub>4</sub> and CO<sub>2</sub> measurements, it could help to identify optimal solutions for mitigating climate warming through reduction of climate forcers (see companion white paper on “Understanding the Co-Evolution of Air Quality and Climate Forcing in a Rapidly Changing World”) or, when combined with ecosystem measurements, can contribute to ensuring global food security (see companion paper on “Air Quality Effects on Crop and Ecosystem Health”).

## References

- Alvarado, M. J., K. E. Cady-Pereira, Y. Xiao, D. B. Millet, and V. H. Payne (2011), Emission Ratios for Ammonia and Formic Acid and Observations of Peroxy Acetyl Nitrate (PAN) and Ethylene in Biomass Burning Smoke as Seen by the Tropospheric Emission Spectrometer (TES), *Atmosphere*, 2(4), 633-654, doi:10.3390/atmos2040633.
- Barkley, M. P., I. De Smedt, M. Van Roozendaal, T. Kurosu, K. Chance, A. Arneth, D. Hagberg, A. Guenther, F. Paulot, E. Marais, and J. Mao (2013), Top-down isoprene emissions over tropical South America inferred from SCIAMACHY and OMI formaldehyde columns, *J. Geophys. Res.*, 118, 6849-6868, doi:10.1002/jgrd.50552.
- Cohen, A. J., H. Ross Anderson, B. Ostro, K. D. Pandey, M. Krzyzanowski, N. Künzli, K. Gutschmidt, A. Pope, I. Romieu, J. M. Samet, and K. Smith (2005), The Global Burden of Disease Due to Outdoor Air Pollution, *Journal of Toxicology and Environmental Health, Part A*, 68(13-14), 1301-1307, doi:10.1080/15287390590936166.
- Cuesta, J., M. Emerenko, X. Liu, G. Dufour, Z. Cai, M. Hoepfner, T. von Clarmann, P. Sellitto, G. Foret, B. Gaubert, M. Beekman, J. Orphal, K. Chance, R. Spurr and J.-M. Flaud (2013), Satellite observation of lowermost tropospheric ozone by multispectral synergism of IASI thermal infrared and GOME-2 ultraviolet measurements over Europe, *Atmos. Chem. Phys.*, 13, 9675-9693.
- Fishman, J., L. T. Iraci, J. Al-Saadi, K. Chance, F. Chavez, M. Chin, P. Coble, C. Davis, P. M. DiGiacomo, D. Edwards, A. Eldering, J. Goes, J. Herman, C. Hu, D. J. Jacob, C. Jordan, S. R. KAWA, R. Key, X. Liu, S. Lohrenz, A. Mannino, V. Natraj, D. Neil, J. Neu, M. Newchurch, K. Pickering, J. Salisbury, H. Sosik, A. Subramaniam, M. Tzortziou, J. Wang and M. Wang (2012), The United States next generation of atmospheric composition and coastal ecosystem measurements: NASA's Geostationary Coastal and Air Pollution Events (GEO-CAPE) mission, *B. Am. Meteorol. Soc.*, 93 (10), 1547-1566.
- Fu D., J. R. Worden, X. Liu, S. S. Kulawik, K. W. Bowman and V. Natraj (2013), Characterization of ozone profiles derived from Aura TES and OMI radiances, *Atmos. Chem. Phys.*, 13, 3445-3462.
- GEO-CAPE (2015), Advancing the science of both coastal ocean biophysics and atmospheric pollution chemistry: A White Paper report to the NASA Earth Science Division by the GEO-CAPE Team. Available: <http://geo-cape.larc.nasa.gov/>
- Ghude, S. D., G. G. Pfister, C. Jena, R. J. van der A, L. K. Emmons and R. Kumar (2013), Satellite constraints of nitrogen dioxide emissions from India based on OMI observations and WRF-Chem simulations, *Geophys. Res. Lett.*, 40 (2), 423-428.
- Gu, d., Y. Wang, C. Smeltzer and K. F. Boersma (2014), Anthropogenic emissions of NO<sub>x</sub> over China: Reconciling the difference of inverse modeling results using GOME-2 and OMI measurements, *J. Geophys. Res.*, 119(12), 7732-7740.
- Huang, M., K. W. Bowman, G. R. Carmichael, T. Chai, R. B. Pierce, J. R. Worden, M. Luo, I. B. Pollack, T. B. Ryerson, J. B. Nowak, J. A. Neuman, J. M. Roberts, E. L. Atlas and D. R. Blake (2014), Changes in nitrogen oxides emissions in California during 2005-2010 indicated from top-down and bottom-up emission estimates, *J. Geophys. Res.*, 119(22), 12928-12952.

- Huang, M., K. W. Bowman, G. R. Carmichael, M. Lee, T. Chai, S. N. Spak, D. K. Henze, A. S. Darmenov, and A. M. Silva (2015), Improved western US background ozone estimates via constraining nonlocal and local source contributions using Aura TES and OMI observations, *J. Geophys. Res.*, **120**, 3572-3592.
- Jerrett, M., R. T. Burnett, C. A. Pope, K. Ito, G. Thurston, D. Krewski, Y. Shi, E. Calle, and M. Thun (2009), *N. Engl. J. Med.*, **360**, 1085-109, DOI: 10.1056/NEJMoa0803894.
- Jerrett, M. (2015), The death toll from air-pollution sources, *Nature*, **525**, 330-331.
- Jiang, Z., J. R. Worden, D. B. A. Jones, J.-T. Lin, W. W. Verstraeten and D. K. Henze (2015), Constraints on Asian ozone using Aura TES, OMI and Terra MOPITT, *Atmos. Chem. Phys.*, **15**, 99-112.
- Kim, P. S., D. J. Jacob, X. Liu, J. X. Warner, K. Yang, K. Chance, V. Thouret and P. Nedelec (2013), Global ozone-CO correlations from OMI and AIRS: constraints on tropospheric ozone sources, *Atmos. Chem. Phys.*, **13**, 9321-9335.
- Lelieveld, J., J. S. Evans, M. Fnais, D. Giannadaki, and A. Pozzer (2015), The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, **525**, 367-371.
- Liousse, C., E. Assamoi, P. Criqui, C. Granier and R. Rosset, Explosive growth in African combustion emissions from 2005 to 2030, *Environ. Res. Lett.*, **9**, doi:10.1088/1748-9326/9/3/035003, 2014.
- Marais, E. A., D. J. Jacob, K. Wecht, C. Lerot, L. Zhang, K. Yu, T. P. Kurosu, K. Chance and B. Sauvage (2014), Anthropogenic emissions in Nigeria and implications for atmospheric ozone pollution: A view from space, *Atmos. Environ.*, **99**, 32-40.
- Natraj, V., X. Liu et al. (2011), Multi-spectral sensitivity studies for the retrieval of tropospheric and lowermost tropospheric ozone from simulated clear-sky GEO-CAPE measurements, *Atmos. Environ.* **45**(39), 7151-7165.
- Neu, J.L., T. Flury, G. L. Manney, M. L. Santee, N. J. Livesey and J. Worden (2014), Tropospheric ozone variations governed by changes in stratospheric circulation, *Nature Geoscience*, **7**, 340-344.
- Pavagadhi, S., R. Betha, S. Venkatesan, R. Balasubramanian, and M. Hande (2013), Physicochemical and toxicological characteristics of urban aerosols during a recent Indonesian biomass burning episode, *Environmental Science and Pollution Research*, **20**(4), 2569-2578, 10.1007/s11356-012-1157-9.
- Rohde, R. A. and R. A. Muller (2015) Air Pollution in China: Mapping of Concentrations and Sources. *PLoS ONE* **10**(8): e0135749. doi:10.1371/journal.pone.0135749.
- Richards, N. A. D., S. R. Arnold, M. P. Chipperfield, G. Miles, R. Siddans, S. A. Monks and M. J. Holloway (2013), The Mediterranean summertime ozone maximum: global emission sensitivities and radiative impacts, *Atmos. Chem. Phys.* **13**, 2331-2345.
- Streets, D. G., J. S. Fu, C. J. Jang, et al. (2007), Air quality during the 2008 Beijing Olympic Games, *Atmos. Environ.*, **41**, 480-492.
- Tawari, C. C., and J. F. N. Abowei (2012), Air Pollution in the Niger Delta Area of Nigeria, *International Journal of Fisheries and Aquatic Sciences*, **1**(2), 94-117.

Verstraeten, W. W., J. L. Neu, J. E. Williams, K. W. Bowman, J. R. Worden and K. F. Boersma (2015), Rapid increases in tropospheric ozone production and export from China, *Nature Geoscience*, 8, 690-695

WHO (2012), World Health Organization, Burden of disease from the joint effects of Household and Ambient Air Pollution for 2012. WHO Technical Report. 2012. Available: [http://www.who.int/phe/health\\_topics/outdoorair/databases/AP\\_jointeffect\\_BoD\\_results\\_March2014.pdf](http://www.who.int/phe/health_topics/outdoorair/databases/AP_jointeffect_BoD_results_March2014.pdf).

WHO (2014), World Health Organization methods and data sources for global causes of death 2000–2012. Global Health Estimates Technical Paper WHO/HIS/HSI/GHE/2014.7. 2014. Available: [http://www.who.int/entity/healthinfo/global\\_burden\\_disease/GlobalCOD\\_method\\_2000\\_2012.pdf](http://www.who.int/entity/healthinfo/global_burden_disease/GlobalCOD_method_2000_2012.pdf).

Worden, H. M., M. N. Deeter, D. P. Edwards, J. C. Gille, J. R. Drummond, and P. Nédélec (2010), Observations of near surface carbon monoxide from space using MOPITT multispectral retrievals, *J. Geophys. Res.*, 115, D18314.

Worden H.M., D. P. Edwards, M. N. Deeter, D. Fu, Kulawik S.S., Worden J.R., Arellano A (2013), “Averaging kernel prediction from atmospheric and surface state parameters based on multiple regression with MOPITT CO and TES-OMI O<sub>3</sub> multispectral observations”, *Atmos. Meas. Tech.*, 6, 1633-1646.

Yang, I. A. and S. T. Holgate (2013), Air pollution and lung health: An epilogue, *Respirology*, 18 (1), 3-4.

Zhu, L., D. K. Henze, K. E. Cady-Pereira, M. W. Shephard, M. Luo, R. W. Pinder, J. O. Bash, and G.-R. Jeong (2013), Constraining U.S. ammonia emissions using TES remote sensing observations and the GEOS-Chem adjoint model, *J. Geophys. Res. Atmos.*, 118, 3355–3368, doi:10.1002/jgrd.50166.

Zoogman, P., D. J. Jacob, K. Chance, L. Zhang, P. Le Sager, A. M. Fiore, A. Eldering, X. Liu, V. Natraj, and S. S. Kulawik (2011), Ozone air quality measurement requirements for a geostationary satellite mission, *Atmos. Environ.*, 45, 7143-7150.