

Development of Comprehensive Models for Black Carbon Scavenging in Arctic Clouds: Integrating Size and Mixing State for Enhanced Climate Predictions

Author Name

January 26, 2026

Abstract

This paper aims to develop comprehensive models that integrate the size and mixing state of Black Carbon (BC) to enhance predictions of its atmospheric impact and removal processes in Arctic environments. By analyzing observational data and utilizing advanced modeling techniques, we seek to improve the accuracy of climate models regarding BC interactions with clouds and its overall influence on climate change.

1 Introduction

Black Carbon (BC) is a significant contributor to climate change, particularly in the Arctic, where it absorbs sunlight and reduces albedo when deposited on snow and ice. Despite its short atmospheric lifetime, BC's radiative properties can influence the climate system profoundly, leading to accelerated warming in sensitive regions. Current climate models often fail to accurately represent BC scavenging processes, primarily due to a lack of consideration for the size distribution and mixing states of BC particles. This study aims to develop more sophisticated models that incorporate these factors to better predict BC's roles in atmospheric processes.

2 Methodology

2.1 Data Collection and Analysis

The primary data source for this study will be the in-cloud single-particle BC dataset collected over 18 months at the Zeppelin Observatory in Svalbard. Statistical analyses will be performed on size distributions and mixing states of BC to establish relationships with scavenging efficiencies.

2.2 Model Development

A new modeling framework will be developed based on:

- **Size Distribution Functions:** Utilizing a power-law function to represent the aerosol size distribution:

$$N(D) = C \cdot D^{-\gamma} \quad (1)$$

where $N(D)$ is the number concentration of BC particles of diameter D , C is a normalization constant, and γ is the size distribution exponent.

- **Mixing State Representation:** Core-shell models to account for the internal structure of BC particles, where the core represents the BC and the shell denotes the surrounding soluble materials.
- **Scavenging Efficiency:**

$$E = 1 - e^{-\lambda \cdot L} \quad (2)$$

where E is the scavenging efficiency, λ is the scavenging coefficient, and L is the cloud liquid water content determining the contact time.

2.3 Validation

The model will be validated against observed data, ensuring its accuracy in predicting scavenging processes. Calibration of the model parameters will be conducted by comparing simulation results with observational data to refine predictions.

3 Results

Outcomes will include model simulations predicting BC scavenging efficiencies based on integrated size and mixing state factors. The new model's predictions will be compared to existing traditional models, demonstrating improvement in capturing the scavenging dynamics under varying environmental conditions.

4 Discussion

The implications of improved modeling for climate predictions will be assessed, particularly regarding long-term climate policy and strategies for reducing BC emissions. Highlight the necessity for continued refinement and validation of the model against real-world observations to reduce uncertainties in forecasts.

5 Conclusion

This study emphasizes the importance of integrating size and mixing state into models of BC scavenging. As BC's role in Arctic warming expands, accurately modeling its interactions with clouds will be crucial for understanding its broader climatic effects. Future research should focus on enhancing model accuracy and integrating these findings into larger climate models.

6 References