

Medical Al Ensemble Clinical Decision Report

Case ID: Generated: 2025-09-05

Primary Diagnostic Consensus

| Diagnosis | ICD-10 | Agreement | Confidence | Status |
|--|--------|-----------|------------|---------|
| ST-Elevation Myocardial Infarction (STEMI) Evidence: ST elevation in leads II, III, aVF indicating inferior wall MI, Elevated troponin I (0.8), Elevated CK-MB, Classic crushing substernal chest pain | I21.0 | 0.0% | Very Low | PRIMARY |

Alternative & Minority Diagnoses

| Diagnosis | ICD-10 | Support | Туре |
|--|--------|---------|----------------------|
| Unstable Angina Evidence: Chest pain pattern, Cardiovascular risk factors | 120.0 | 25.9% | Alternative (10-29%) |
| Aortic Dissection Evidence: Severe chest pain, Hypertension, Sudden onset | 171.0 | 22.2% | Alternative (10-29%) |
| Pulmonary Embolism Evidence: Chest pain, Shortness of breath, Cardiovascular risk factors | 126.99 | 18.5% | Alternative (10-29%) |
| Acute Pericarditis Evidence: Chest pain, ECG changes | 130.9 | 14.8% | Alternative (10-29%) |
| Non-ST-Elevation Myocardial Infarction (NSTEMI) Evidence: Elevated cardiac enzymes, Chest pain | l21.4 | 7.4% | Minority (<10%) |
| Acute Coronary Syndrome Evidence: Chest pain, Elevated cardiac markers | 120.1 | 7.4% | Minority (<10%) |
| Esophageal Spasm Evidence: Chest pain mimicking cardiac pain | K22.8 | 3.7% | Minority (<10%) |
| Angina Pectoris Evidence: Chest pain, Cardiovascular risk factors | 120.0 | 3.7% | Minority (<10%) |

| Diagnosis | ICD-10 | Support | Туре |
|--|--------|---------|-----------------|
| Acute Inferior Wall Myocardial Infarction Evidence: ST elevation in inferior leads, Elevated cardiac enzymes | l21.1 | 7.4% | Minority (<10%) |
| Acute Myocardial Infarction Evidence: Elevated cardiac enzymes, Classic presentation | I21.9 | 7.4% | Minority (<10%) |

| Analysis Overview | |
|-------------------------------|--|
| Models Queried: 23 | |
| Successful Responses: 23 | |
| Consensus Level: High | |
| Total Estimated Cost: \$0.286 | |

Critical Decision Points & Evidence Synthesis

Critical Decision Points

Key areas where models showed significant divergence in diagnostic or management approach:

Evidence Synthesis & Clinical Correlation

Symptom-Diagnosis Correlation Matrix

| Symptom | STEMI | Unstable | Aortic D |
|-----------------|--------|----------|----------|
| crushing chest | Strong | - | - |
| ST elevation | Strong | - | - |
| elevated tropon | Strong | - | - |

Legend: +++ Strong association, ++ Moderate, + Weak, - Not typical

Diagnostic Decision Tree

| Step | Action | If Positive | If Negative |
|------|--------------------------|--------------------------------|--------------------------------|
| 1 | Initial Laboratory Tests | → Confirm suspicion | ightarrow Broaden differential |
| 2 | Imaging Studies | → Identify pathology | → Consider specialized tests |
| 3 | Specialized Testing | ightarrow Definitive diagnosis | → Empiric treatment |
| 4 | Treatment Trial | → Continue if effective | → Reconsider diagnosis |

Executive Summary

Case Description

A 45-year-old male presents to the emergency department with sudden onset chest pain that started 2 hours ago. The pain is described as crushing, substernal, and radiates to the left arm. He has been sweating profusely and feels nauseous.

Past Medical History:

- Hypertension (diagnosed 5 years ago, on amlodipine)
- Type 2 Diabetes Mellitus (diagnosed 3 years ago, on metformin)
- Hyperlipidemia (on atorvastatin)
- 20 pack-year smoking history, quit 2 years ago
- Family history of coronary artery disease (father had MI at age 55)

Physical Examination:

- BP: 150/90 mmHg
- HR: 95 bpm, regular
- RR: 22/min
- O2 Sat: 94% on room air
- Temperature: 37.2°C
- Diaphoretic, appears anxious
- Cardiovascular: Normal S1/S2, no murmurs or gallops
- Lungs: Clear bilaterally
- Abdomen: Soft, non-tender

Initial Tests:

- ECG: ST elevation in leads II, III, and aVF
- Troponin I: 0.8 ng/mL (elevated)
- CK-MB: 25 ng/mL (elevated)

Key Clinical Findings

· Positive family history of similar episodes

Primary Recommendations

- Consider ST-Elevation Myocardial Infarction (STEMI) among differential diagnoses
- Activate cardiac catheterization lab for primary PCI
- Administer dual antiplatelet therapy (aspirin + P2Y12 inhibitor)
- Obtain IV access and continuous cardiac monitoring
- Obtain 12-lead ECG for diagnostic confirmation

Primary Diagnosis Clinical Summaries

■ Key Clinical Findings

| Finding | Supporting Evidence | Clinical Reasoning |
|------------------------------------|-----------------------|--------------------------|
| ST elevation in leads II, III, aVF | Clinical presentation | Key diagnostic indicator |
| Elevated troponin I (0.8 ng/mL) | Clinical presentation | Key diagnostic indicator |
| Elevated CK-MB | Clinical presentation | Key diagnostic indicator |
| Crushing substernal chest pain | Clinical presentation | Key diagnostic indicator |
| Pain radiating to left arm | Clinical presentation | Key diagnostic indicator |

■ Recommended Tests

| Test Name | Туре | Priority | Rationale |
|-------------------------------------|------------|----------|-------------------------|
| 12-lead ECG | Laboratory | Urgent | Diagnostic confirmation |
| Cardiac troponin levels | Laboratory | Urgent | Diagnostic confirmation |
| Complete blood count with platelets | Laboratory | Urgent | Diagnostic confirmation |
| Basic metabolic panel | Laboratory | Urgent | Diagnostic confirmation |
| PT/PTT/INR | Laboratory | Urgent | Diagnostic confirmation |

■ Immediate Management

| Intervention | Category | Urgency | Clinical Reasoning |
|--|----------|-----------|-----------------------|
| Activate cardiac catheterization lab for primary PCI | Medical | Immediate | Critical intervention |
| Administer dual antiplatelet therapy (aspirin + P2Y12 inhibitor) | Medical | Immediate | Critical intervention |
| Obtain IV access and continuous cardiac monitoring | Medical | Immediate | Critical intervention |
| Administer oxygen if SpO2 < 90% | Medical | Immediate | Critical intervention |

| Intervention | Category | Urgency | Clinical Reasoning |
|---|----------|-----------|-----------------------|
| Pain management with morphine if needed | Medical | Immediate | Critical intervention |

■ Medications

| Medication | Dosage | Route/Frequency | Indication |
|--------------|--------|------------------|-----------------------------------|
| Aspirin | 325mg | PO / once | antiplatelet therapy for STEMI |
| Clopidogrel | 600mg | PO / once | P2Y12 inhibition for STEMI |
| Atorvastatin | 80mg | PO / daily | high-intensity statin therapy |
| Metoprolol | 25mg | PO / twice daily | beta-blocker for cardioprotection |

Diagnostic Landscape Analysis

Detailed Diagnostic Analysis

The ensemble analysis identified **ST-Elevation Myocardial Infarction (STEMI)** as the primary diagnosis with 0.0% consensus among 11 models.

Detailed Alternative Analysis

| Diagnosis | Support | Key Evidence | Clinical Significance |
|---|---------|--------------|-----------------------|
| Unstable Angina Evidence: Chest pain pattern, Cardiovascular risk factors | 25.9% | 7 models | Less likely |
| Aortic Dissection Evidence: Severe chest pain, Hypertension, Sudden onset | 22.2% | 6 models | Less likely |
| Pulmonary Embolism Evidence: Chest pain, Shortness of breath, Cardiovascular risk factors | 18.5% | 5 models | Less likely |
| Acute Pericarditis Evidence: Chest pain, ECG changes | 14.8% | 4 models | Less likely |
| Non-ST-Elevation Myocardial Infarction (NSTEMI) Evidence: Elevated cardiac enzymes, Chest pain | 7.4% | 2 models | Unlikely |
| Acute Coronary Syndrome Evidence: Chest pain, Elevated cardiac markers | 7.4% | 2 models | Unlikely |
| Esophageal Spasm Evidence: Chest pain mimicking cardiac pain | 3.7% | 1 models | Unlikely |
| Angina Pectoris Evidence: Chest pain, Cardiovascular risk factors | 3.7% | 1 models | Unlikely |

Minority Opinions

All alternative diagnoses suggested by any models with their clinical rationale:

• Non-ST-Elevation Myocardial Infarction (NSTEMI) (ICD-10: Unknown) - 7.4% agreement (2 models)

Supporting Models: Model1, Model15

• Acute Coronary Syndrome (ICD-10: Unknown) - 7.4% agreement (2 models)

Supporting Models: Model10, Model19

• Esophageal Spasm (ICD-10: Unknown) - 3.7% agreement (1 models)

Supporting Models: Model9

• Angina Pectoris (ICD-10: Unknown) - 3.7% agreement (1 models)

Supporting Models: Model10

• Acute Inferior Wall Myocardial Infarction (ICD-10: Unknown) - 7.4% agreement (2 models)

Supporting Models: Model13, Model15

• Acute Myocardial Infarction (ICD-10: Unknown) - 7.4% agreement (2 models)

Supporting Models: Model19, Model22

Additional Diagnoses Considered:

• Unstable Angina (ICD-10: I20.0) - 34.8% (8 models)

Evidence: Chest pain pattern, Cardiovascular risk factors

• Aortic Dissection (ICD-10: I71.0) - 21.7% (5 models)

Evidence: Severe chest pain, Hypertension, Sudden onset

• **Pulmonary Embolism** (ICD-10: I26.99) - 21.7% (5 models)

Evidence: Chest pain, Shortness of breath, Cardiovascular risk factors

• Acute Pericarditis (ICD-10: I30.9) - 17.4% (4 models)

Evidence: Chest pain, ECG changes

Management Strategies & Clinical Pathways

Immediate Actions Required

| Priority | Action | Rationale | Consensus |
|----------|--|---------------------|-----------|
| 1 | Activate cardiac catheterization lab for primary PCI | Clinical indication | 50% |
| 2 | Administer dual antiplatelet therapy (aspirin + P2Y12 inhibitor) | Clinical indication | 50% |
| 3 | Obtain IV access and continuous cardiac monitoring | Clinical indication | 50% |
| 4 | Administer oxygen if SpO2 < 90% | Clinical indication | 50% |
| 5 | Pain management with morphine if needed | Clinical indication | 50% |

Recommended Diagnostic Tests

| Test | Purpose | Priority | Timing |
|-------------------------------------|-------------------------|----------|--------------|
| 12-lead ECG | Diagnostic confirmation | Routine | As indicated |
| Cardiac troponin levels | Diagnostic confirmation | Routine | As indicated |
| Complete blood count with platelets | Diagnostic confirmation | Routine | As indicated |
| Basic metabolic panel | Diagnostic confirmation | Routine | As indicated |
| PT/PTT/INR | Diagnostic confirmation | Routine | As indicated |
| Chest X-ray | Diagnostic confirmation | Routine | As indicated |

Treatment Recommendations

Treatment recommendations pending diagnostic confirmation.

Model Diversity & Bias Analysis

Model Response Overview & Cost Analysis

| Model | Origin | Tier | Cost | Diagnosis | Training Profile |
|-----------------|--------|-----------|---------|---|------------------|
| gpt-4o | USA | Premium | \$0.018 | Inferior ST-Elevation Myocardial Infarction (STEMI) | Comprehensive |
| gpt-4o-mini | USA | Budget | <\$0.01 | Acute Myocardial Infarction (ST-Elevation) | General |
| gpt-oss-120b | USA | Mid-Range | <\$0.01 | Acute inferior ST-elevation myocardial infarction (STEMI) | Standard |
| claude-3-opus-2 | USA | Premium | \$0.109 | Acute ST-elevation myocardial infarction (STEMI) | Comprehensive |
| gemini-2.5-pro | USA | Premium | \$0.029 | Not specified | General |
| gemini-2.5-flas | USA | Budget | <\$0.01 | ST-Elevation Myocardial Infarction (STEMI) | General |
| gemini-2.5-flas | USA | Budget | <\$0.01 | ST-Elevation Myocardial Infarction (STEMI) | General |
| gemma-2-9b-it | USA | Budget | <\$0.01 | Acute Myocardial Infarction (STEMI) | Standard |
| gemma-3-12b-it | USA | Unknown | <\$0.01 | Acute ST-Elevation Myocardial Infarction (STEMI) | Standard |
| llama-3.2-3b-in | USA | Budget | <\$0.01 | Acute Coronary Syndrome | General |
| mistral-large-2 | France | Premium | \$0.015 | ST-Elevation Myocardial Infarction (STEMI) | Standard |
| mistral-7b-inst | France | Budget | <\$0.01 | Acute Myocardial Infarction | General |
| deepseek-chat | China | Budget | <\$0.01 | Acute Inferior Wall Myocardial Infarction | Regional |
| deepseek-r1 | China | Budget | <\$0.01 | ST-Elevation Myocardial Infarction (Inferior Wall) | Regional |
| deepseek-chat-v | China | Unknown | <\$0.01 | Acute Inferior Wall Myocardial Infarction | Regional |
| qwen-2.5-coder- | China | Mid-Range | <\$0.01 | Acute Myocardial Infarction (STEMI) | Regional |
| command-r-plus | Canada | Premium | \$0.020 | Acute Myocardial Infarction (Heart Attack) | Standard |
| command-r | Canada | Mid-Range | <\$0.01 | Acute Myocardial Infarction (STEMI) | Standard |
| jamba-large-1.7 | Israel | Premium | \$0.012 | Acute Myocardial Infarction (AMI) | Standard |
| sonar-deep-rese | USA | Premium | \$0.012 | ST elevation myocardial infarction of inferior wall | Standard |
| wizardlm-2-8x22 | USA | Mid-Range | <\$0.01 | ST-Elevation Myocardial Infarction (STEMI) | Standard |
| grok-2-1212 | USA | Premium | \$0.012 | Acute Myocardial Infarction | Standard |

| Model | Origin | Tier | Cost | Diagnosis | Training Profile |
|--------|--------|---------|---------|--|------------------|
| grok-4 | USA | Premium | \$0.040 | ST-Elevation Myocardial Infarction (STEMI) | Alternative |

^{**}Total Estimated Cost: \$0.286**

Understanding Training Profiles

Training profiles indicate the type and depth of medical knowledge in each model:

Comprehensive: Extensive medical literature training with broad clinical knowledge

Standard: Standard medical knowledge base with general clinical training

Regional: Region-specific medical training reflecting local practices and conditions

General: Broad general knowledge, not specifically trained on medical literature

Alternative: Alternative medical perspectives and non-conventional approaches

Al Model Bias Analysis

Al model bias analysis is generated during orchestration (Step 2). This comprehensive analysis examines cultural, geographic, and training data biases across the Al models used.

Detailed Model Responses

Complete diagnostic assessments from each model:

1. gpt-4o (USA, Released: 2024-05-13)

Primary Diagnosis: Inferior ST-Elevation Myocardial Infarction (STEMI) (ICD-10: I21.19) - Confidence: 0.95

Differential Diagnoses:

- Unstable Angina (ICD: I20.0) 0.6
- Non-ST-Elevation Myocardial Infarction (NSTEMI) (ICD: I21.4) 0.5
- Aortic Dissection (ICD: I71.01) 0.4

Key Clinical Findings:

- Sudden onset crushing chest pain
- · Radiating pain to left arm
- Profuse sweating
- ST elevation in leads II, III, and aVF

2. gpt-4o-mini (USA, Released: 2024-07-18)

Primary Diagnosis: Acute Myocardial Infarction (ST-Elevation) (ICD-10: I21.01) - Confidence: 0.95 **Differential Diagnoses:**

- Unstable Angina (ICD: I20.0) 0.75
- Aortic Dissection (ICD: I71.0) 0.5
- Pulmonary Embolism (ICD: I26.99) 0.4

Key Clinical Findings:

- Crushing substernal chest pain
- · Radiation to left arm
- ST elevation in leads II, III, and aVF
- Elevated Troponin I and CK-MB levels

3. gpt-oss-120b (USA, Released: 2025-08-05)

4. claude-3-opus-2 (USA, Released: 2024-02-29)

Primary Diagnosis: Acute ST-elevation myocardial infarction (STEMI) (ICD-10: I21.3) - Confidence: 0.95

Differential Diagnoses:

Unstable angina (ICD: I20.0) - 0.4
Acute pericarditis (ICD: I30.0) - 0.2
Aortic dissection (ICD: I71.0) - 0.1

Key Clinical Findings:

- Sudden onset chest pain (crushing, substernal, radiating to left arm)
- · Diaphoresis and nausea
- ST elevation on ECG in leads II, III, and aVF
- Elevated troponin I and CK-MB

5. gemini-2.5-pro (USA, Released: 2024-12-19)

6. gemini-2.5-flas (USA, Released: 2024-12-19)

Primary Diagnosis: ST-Elevation Myocardial Infarction (STEMI) (ICD-10: I21.0) - Confidence: 0.98 **Differential Diagnoses:**

Unstable Angina (ICD: I20.0) - 0.7
Acute Pericarditis (ICD: I30.9) - 0.4
Aortic Dissection (ICD: I71.0) - 0.3

Key Clinical Findings:

- Sudden onset crushing substernal chest pain radiating to left arm
- Profuse sweating and nausea
- Hypertension, Type 2 Diabetes Mellitus, Hyperlipidemia
- 20 pack-year smoking history, family history of CAD

7. gemini-2.5-flas (USA, Released: 2024-12-19)

Primary Diagnosis: ST-Elevation Myocardial Infarction (STEMI) (ICD-10: I21.19) - Confidence: 0.95 **Differential Diagnoses:**

- Unstable Angina (ICD: I20.0) 0.05
- Pulmonary Embolism (ICD: I26.99) 0.02
- Aortic Dissection (ICD: I71.00) 0.01

- Sudden onset crushing substernal chest pain radiating to left arm
- Diaphoresis and nausea
- ST elevation in leads II, III, and aVF on ECG
- Elevated Troponin I

8. gemma-2-9b-it (USA, Released: 2024-06-27)

Primary Diagnosis: Acute Myocardial Infarction (STEMI) (ICD-10: I21.0) - Confidence: 0.95

Differential Diagnoses:

• Aortic Dissection (ICD: I71.0) - 0.05

• Pericarditis (ICD: I30.0) - 0.05

• Severe Angina (ICD: I20.9) - 0.03

Key Clinical Findings:

- · Crushing substernal chest pain radiating to left arm
- · Diaphoresis and nausea
- ST elevation in leads II, III, and aVF on ECG
- Elevated Troponin I and CK-MB levels

9. gemma-3-12b-it (USA, Released: 2024-12-11)

Primary Diagnosis: Acute ST-Elevation Myocardial Infarction (STEMI) (ICD-10: I21.0) - Confidence: 0.95

Differential Diagnoses:

• Unstable Angina (ICD: I20.0) - 0.2

• Esophageal Spasm (ICD: K22.8) - 0.1

• Pericarditis (ICD: I30.0) - 0.05

Key Clinical Findings:

- Crushing substernal chest pain radiating to the left arm
- Profuse sweating (diaphoresis)
- Nausea
- ST elevation in leads II, III, and aVF on ECG

10. Ilama-3.2-3b-in (USA, Released: 2024-09-25)

Primary Diagnosis: Acute Coronary Syndrome (ICD-10: I21.0) - Confidence: 0.9

Differential Diagnoses:

• Myocardial Infarction (ICD: I21.0) - 0.8

• Angina Pectoris (ICD: I20.0) - 0.7

• Pericarditis (ICD: K89.0) - 0.6

- Sudden onset chest pain radiating to the left arm
- Elevated troponin and CK-MB
- ECG changes (ST elevation in leads II, III, and aVF)

11. mistral-large-2 (France, Released: 2024-11-01)

Primary Diagnosis: ST-Elevation Myocardial Infarction (STEMI) (ICD-10: I21.02) - Confidence: 0.9 **Differential Diagnoses:**

- Unstable Angina (ICD: I20.0) 0.7
- Aortic Dissection (ICD: I71.0) 0.3
- Pulmonary Embolism (ICD: I26.9) 0.2

Key Clinical Findings:

- Crushing substernal chest pain radiating to the left arm
- Diaphoresis and nausea
- ST elevation in leads II, III, and aVF
- Elevated Troponin I and CK-MB

12. mistral-7b-inst (France, Released: 2023-09-27)

Primary Diagnosis: Acute Myocardial Infarction (ICD-10: I21.9 (Acute coronary heart disease, unspecified, without mention of acute MI, not in acute stage)) - Confidence: 0.95

Differential Diagnoses:

- Atypical Angina Pectoris (ICD: I20.0 (Angina pectoris, stable angina pectoris)) 0.05
- Pericarditis (ICD: I30.0 (Acute pericarditis)) 0.03
- Esophageal Spasm (ICD: K21.3 (Esophageal spasm without obstruction)) 0.02

Key Clinical Findings:

· Sudden onset chest pain, crushing, substernal, radiating to the left arm, diaphoresis, nausea

13. deepseek-chat (China, Released: 2024-12-20)

Primary Diagnosis: Acute Inferior Wall Myocardial Infarction (ICD-10: I21.1) - Confidence: 0.95 **Differential Diagnoses:**

- Aortic Dissection (ICD: I71.00) 0.2
- Pulmonary Embolism (ICD: I26.99) 0.15
- Pericarditis (ICD: I30.9) 0.1

- Sudden onset crushing substernal chest pain radiating to the left arm
- ST elevation in leads II, III, and aVF on ECG
- Elevated Troponin I and CK-MB
- Hypertension, diabetes, hyperlipidemia, and smoking history

14. deepseek-r1 (China, Released: 2025-01-20)

15. deepseek-chat-v (China, Released: 2024-12-26)

Primary Diagnosis: Acute Inferior Wall Myocardial Infarction (ICD-10: I21.4) - Confidence: 0.98 **Differential Diagnoses:**

- Aortic Dissection (ICD: I71.01) 0.15
- Pericarditis (ICD: I30.9) 0.1
- Pulmonary Embolism (ICD: I26.99) 0.08

Key Clinical Findings:

- ST elevation in leads II, III, aVF (inferior wall pattern)
- Elevated troponin I (0.8 ng/mL)
- Elevated CK-MB (25 ng/mL)
- · Crushing substernal chest pain radiating to left arm

16. qwen-2.5-coder- (China, Released: 2024-11-12)

Primary Diagnosis: Acute Myocardial Infarction (STEMI) (ICD-10: I21.0) - Confidence: 0.95 **Differential Diagnoses:**

- Unstable Angina (ICD: I20.1) 0.03
- Pulmonary Embolism (ICD: I26.9) 0.01
- Pneumothorax (ICD: J93.0) 0.005

Key Clinical Findings:

- Sudden onset chest pain, crushing, substernal, radiating to left arm
- · Profuse sweating, nausea
- ST elevation in leads II, III, and aVF on ECG
- Elevated Troponin I (0.8 ng/mL)

17. command-r-plus (Canada, Released: 2024-04-04)

Primary Diagnosis: Acute Myocardial Infarction (Heart Attack) (ICD-10: I21.0) - Confidence: 0.9 **Differential Diagnoses:**

- Unstable Angina (ICD: I20.0) 0.7
- Aortic Dissection (ICD: I71.0) 0.4
- Pulmonary Embolism (ICD: I26.9) 0.3

- · Chest pain with radiation to left arm
- Diaphoresis and nausea
- · Elevated troponin and CK-MB levels
- ST elevation on ECG

18. command-r (Canada, Released: 2024-03-11)

Primary Diagnosis: Acute Myocardial Infarction (STEMI) (ICD-10: I21.1) - Confidence: 0.95 **Differential Diagnoses:**

- Anginal Syndrome (ICD: I20.9) 0.08
- Non-ST Elevation Myocardial Infarction (NSTEMI) (ICD: I21.2) 0.05
- Acute Coronary Syndrome (ACS) (ICD: I25.1) 0.05

Key Clinical Findings:

- Chest pain: Crushing, substernal, radiates to left arm
- Elevated cardiac enzymes: Troponin I 0.8 ng/mL, CK-MB 25 ng/mL
- ST elevation on ECG in leads II, III, and aVF
- History of cardiovascular risk factors: Hypertension, Diabetes, Hyperlipidemia, Smoking, Family Hx of CAD

19. jamba-large-1.7 (Israel, Released: 2025-07-01)

Primary Diagnosis: Acute Myocardial Infarction (AMI) (ICD-10: I21.9) - Confidence: 0.95 **Differential Diagnoses:**

- Acute Coronary Syndrome (ICD: I20.1) 0.85
- Pulmonary Embolism (ICD: I26.90) 0.7
- Aortic Dissection (ICD: I71.00) 0.65

Key Clinical Findings:

- Substernal chest pain radiating to left arm
- ST elevation in leads II, III, and aVF on ECG
- Elevated troponin I and CK-MB
- Hypertension and diabetes mellitus as risk factors

20. sonar-deep-rese (USA, Released: 2025-03-07)

21. wizardlm-2-8x22 (USA, Released: 2024-04-15)

Primary Diagnosis: ST-Elevation Myocardial Infarction (STEMI) (ICD-10: I21.3) - Confidence: 0.9

Differential Diagnoses:

- Unstable Angina (ICD: I20.0) 0.6
- Acute Pericarditis (ICD: I30.9) 0.2
- Pulmonary Embolism (ICD: I26.99) 0.1

Key Clinical Findings:

- Sudden onset crushing substernal chest pain
- Chest pain radiating to the left arm
- · Diaphoresis and nausea
- ST elevation in leads II, III, and aVF on ECG

22. grok-2-1212 (USA, Released: 2024-12-12)

Primary Diagnosis: Acute Myocardial Infarction (ICD-10: I21.1) - Confidence: 0.95

Differential Diagnoses:

- Unstable Angina (ICD: I20.0) 0.7
- Pulmonary Embolism (ICD: I26.99) 0.5
- Aortic Dissection (ICD: I71.0) 0.4

Key Clinical Findings:

- Sudden onset crushing chest pain
- · Pain radiates to left arm
- Profuse sweating and nausea
- ST elevation in leads II, III, and aVF

23. grok-4 (USA, Released: 2024-12-26)

Primary Diagnosis: ST-Elevation Myocardial Infarction (STEMI) (ICD-10: I21.19) - Confidence: 0.95 **Differential Diagnoses:**

- Aortic Dissection (ICD: I71.00) 0.2
- Pulmonary Embolism (ICD: I26.99) 0.15
- Pericarditis (ICD: I30.9) 0.1

- Sudden onset crushing substernal chest pain radiating to left arm
- Profuse sweating and nausea
- ST elevation in leads II, III, aVF on ECG
- Elevated troponin I (0.8 ng/mL) and CK-MB (25 ng/mL)