MEDLEY Clinical Decision Report

Case tmpgpg3f1s4: Medical Case from tmpgpg3f1s4.txt

Case Overview

Presentation:	A 28-year-old male of Mediterranean descent presents with: - Recurrent episodes of fever lasting 1-3 days - Severe abdominal pain during episodes - Chest pain
	with breathing difficulties - Joint pain affecting knees and ankles - Family history: Father and paternal uncle have similar symptoms - Ep

Diagnostic Landscape

PRIMARY DIAGNOSIS	Familial Mediterranean Fever	17% Agreement
Minority Opinion	Familial Mediterranean Fever	By Mistral 7B, Mistral 7B
Minority Opinion	Inflammatory Bowel Disease	By Mistral 7B

Management Strategies

Critical Decision Points

- Low consensus indicates complex presentation consider specialist consultation
- ■■ Multiple minority diagnoses suggest atypical presentation
- ■■ Geographic variation in diagnoses consider regional disease patterns

Models: 8	Responded: 6	Unique Diagnoses: 7	Consensus: None
-----------	--------------	---------------------	-----------------

Model Diversity & Bias Analysis

Model Response Patterns

Model	Origin	Release	Primary Diagnosis	Confidence
Mistral 7B	France	2023-09	Familial Mediterranean Feve	Varied
Gemma 2 9B	USA	2024-06	Appendicitis	Varied
Llama 3.2 3B	USA	2024-09	Unknown	Varied
Qwen 2.5 32B	China	2024-11	Unknown	Varied
Gemini 2.0 Flash	USA	2024-12	Unknown	Varied
DeepSeek V3	China	2024-03	Unknown	Varied

Identified Bias Sources

Geographic	France vs USA	Divergent diagnoses based on regional training	
Geographic	France vs China	Divergent diagnoses based on regional training	
Temporal	Release dates vary	Model release dates span from 2023-09 to 2024-12, potential	ly affecting knowle
Training Data	Multilingual web data	Broad knowledge but variable quality	
Training Data	Curated datasets	Standard medical training data	

Unique Model Perspectives

Model	Unique Diagnosis	Reasoning
Mistral 7B	Inflammatory Bowel Disease	ikely." }, { "name": , "confidence": 0.2, "icd_code": "K50.XX", "reasoning":
Mistral 7B	Systemic Lupus Erythematosus	tial_diagnoses": [{ "name": , "confidence": 0.2, "icd_code": "M32.0", "leas
Mistral 7B	Rheumatoid Arthritis	bable." }, { "name": " (RA)", "confidence": 0.1, "icd_code": "M05.XX", "rea
Gemma 2 9B	Appendicitis	No reasoning
Gemma 2 9B	Gastroenteritis	No reasoning

Diagnostic Categories

- Autoimmune/Inflammatory: Inflammatory Bowel Disease, Systemic Lupus Erythematosus, Rheumatoid Arthritis
- Infectious: Familial Mediterranean Fever

• Other: Appendicitis, Gastroenteritis, Constipation

Detailed Model Responses

Mistral 7B (France)

 Familial Mediterranean Fever (FMF) primary_diagnosis": { "name": , "confidence": 0.8, "icd_code": "M39.0", "reasoning": "Recurrent episodes of fever, abdominal pain, chest pain,

joint pain, and elevate

2. FMF name": "Familial Mediterranean Fever ()", "confidence": 0.8,

"icd_code": "M39.0", "reasoning": "Recurrent episodes of fever,

abdominal pain, chest pain, joint pain, and elevate

3. Inflammatory Bowel

Disease (IBD)

ikely." }, { "name": , "confidence": 0.2, "icd_code": "K50.XX", "reasoning": "Recurrent abdominal pain, fever, and elevated

inflammato

Gemma 2 9B (USA)

Appendicitis
Gastroenteritis
Constipation
No reasoning provided
No reasoning provided

Llama 3.2 3B (USA)

Unknown The patient's presentation could also be consistent with

undifferentiated arthritis, as the diagnosis is not yet specific enough to

assign an ICD-10 code.

Unknown The patient's joint pain and family history could suggest ankylosing

spondylitis, but the lack of specific symptoms and radiographic

evidence makes this a less likely diagnosis.

3. Unknown The patient's episodes of fever and abdominal pain could also be

consistent with gastroenteritis, but the duration and frequency of

episodes suggest a more chronic condition.

Qwen 2.5 32B (China)

Unknown Recurrent episodes of fever with elevated WBC and neutropenia can

mimic PFAPA, though the abdominal and chest pain are less typical.

2. Unknown Recurrent fever with abdominal pain and arthritis in a person of

Mediterranean descent, but FMF typically presents with peritonitis,

chest pain is less common, and no pharyngitis or adenitis.

3. Unknown Recurrent fever and chest pain can occur in Takayasu Arteritis, but it is

less common in children and usually presents with other symptoms like

fatigue and weight loss.

Gemini 2.0 Flash (USA)

Unknown Shares similar symptoms of recurrent fever, abdominal pain, and joint

pain. However, FMF is more common in Mediterranean populations.

Genetic testing is crucial to differentiate.

2. Unknown TRAPS presents with recurrent fever, abdominal pain, and arthritis, but

episodes tend to last longer (several days to weeks) than in FMF.

Family history also supports considering this diagnosis.

3. Unknown Can present with fever, joint pain, and elevated inflammatory markers.

However, the specific pattern of symptoms and family history make

FMF more likely. Usually presents earlier in childhood.

DeepSeek V3 (China)

1. Unknown Recurrent fever episodes and family history of similar symptoms could

fit this diagnosis, but absence of aphthous ulcers and pharyngitis

makes it less likely.

2. Unknown Episodic fever and joint pain are common in SJIA, but the absence of

rash and the family history make FMF more likely.

3. Unknown Other hereditary periodic fever syndromes (e.g., TRAPS, MKD) could

present similarly, but FMF is most common in Mediterranean

populations.

Report generated: 2025-08-11 12:40:41 | Medley Medical Al Ensemble System | Developed by Farhad Abtahi, SMAILE at Karolinska Institutet