

# MEDLEY

## Medical AI Ensemble Clinical Decision Report

Case ID: Case\_1

Title: Case\_1 - Medical Analysis

Generated: 2025-08-10  
22:52

### Primary Diagnostic Consensus

Diagnosis	ICD-10	Agreement	Confidence	Status
Familial Mediterranean Fever <i>Evidence: recurrent fever episodes, periodic abdominal pain, arthritis, elevated inflammatory markers</i>	E85.0	63.0%	High	PRIMARY

### Alternative & Minority Diagnoses

Diagnosis	ICD-10	Support	Type
Inflammatory Bowel Disease	K50.9	63.0%	Strong Alt (≥30%)
Systemic Lupus Erythematosus	M32.9	48.1%	Strong Alt (≥30%)
Septic Arthritis	M00.9	33.3%	Strong Alt (≥30%)
Appendicitis	K35.8	22.2%	Alternative (20-29%)
Reactive Arthritis	M02.9	14.8%	Alternative (20-29%)
Gout	M10.9	11.1%	Alternative (20-29%)
Psoriatic Arthritis	L40.5	7.4%	Minority (<20%)
Behçet's Disease	M35.2	7.4%	Minority (<20%)
TRAPS (TNF Receptor-Associated Periodic Syndrome)	E85.0	7.4%	Minority (<20%)
Autoinflammatory Syndrome	M04.9	7.4%	Minority (<20%)
Tuberculosis	A15.9	3.7%	Minority (<20%)
Hyper-IgD Syndrome	D89.8	3.7%	Minority (<20%)
Acute Recurrent Peritonitis	R69	3.7%	Minority (<20%)
Acute Intermittent Porphyria	R69	3.7%	Minority (<20%)
Infectious/Tuberculous Peritonitis	R69	3.7%	Minority (<20%)
Ankylosing Spondylitis	R69	3.7%	Minority (<20%)

Diagnosis	ICD-10	Support	Type
Inflammatory Bowel Disease with extraintestinal manifestations	R69	3.7%	Minority (<20%)
Irritable Bowel Syndrome	R69	3.7%	Minority (<20%)
Appendicitis (recurrent or atypical)	R69	3.7%	Minority (<20%)
Infectious etiology (e.g., intra-abdominal abscess, pelvic inflammatory disease)	R69	3.7%	Minority (<20%)
Periodic Fever, Aphthous Stomatitis, Pharyngitis, and Adenitis (PFAPA) Syndrome	R69	3.7%	Minority (<20%)
Periodic Fever, Aphthous Stomatitis, Pharyngitis, and Adenitis Syndrome	R69	3.7%	Minority (<20%)
Seronegative Spondyloarthritis	R69	3.7%	Minority (<20%)
Adult-onset Still's Disease	R69	3.7%	Minority (<20%)
Acute Intermittent Porphyria (AIP)	R69	3.7%	Minority (<20%)
Diverticulitis	R69	3.7%	Minority (<20%)
Rheumatic fever	R69	3.7%	Minority (<20%)
Abdominal tuberculosis	R69	3.7%	Minority (<20%)
Recurrent Peritonitis due to Appendiceal Perforation	R69	3.7%	Minority (<20%)
Recurrent Peritonitis due to Appendicitis	R69	3.7%	Minority (<20%)
Recurrent Peritonitis due to Amyloidosis	R69	3.7%	Minority (<20%)
Acute intermittent porphyria	R69	3.7%	Minority (<20%)
Systemic lupus erythematosus	R69	3.7%	Minority (<20%)
Hereditary angioedema	R69	3.7%	Minority (<20%)
Rheumatic Fever	R69	3.7%	Minority (<20%)

Analysis Overview
Models Queried: 27
Successful Responses: 27
Consensus Level: Moderate
Total Estimated Cost: \$0.101

# Critical Decision Points & Evidence Synthesis

## Critical Decision Points

Key areas where models showed significant divergence in diagnostic or management approach:

## Evidence Synthesis & Clinical Correlation

### Symptom-Diagnosis Correlation Matrix

Symptom	FMF	Arthritis	IBD	FMF	Lupus	SLE	Septic	Reactive
fever	++	++	++	+++	+	+++	++	++
pain	++	++	++	+++	+	+++	++	++
arthritis	++	++	++	+++	+	+++	++	++
abdominal pain	++	++	++	+++	+	+++	++	++
peritoneal sign	++	++	++	+++	+	+++	++	++
inflammation	++	++	+	++	+++	+++	+	++
elevated CRP	+++	++	++	+++	+++	+++	++	++
swelling	+++	++	+	++	+++	++	+	++
joint pain	++	++	++	++	+	+	++	++
rash	+++	+++	+++	+++	+++	+++	-	-

Legend: +++ Strong association, ++ Moderate, + Weak, - Not typical

## Diagnostic Decision Tree

Step	Action	If Positive	If Negative
1	MEFV gene sequencing	Start colchicine therapy	Consider expanded genetic panel
2	Inflammatory markers (CRP/ESR)	Support FMF diagnosis	Consider alternative diagnoses

# Executive Summary

## Case Description

### Case 1: Cultural and Genetic Diversity Challenge

**Patient:** 28-year-old woman, recent immigrant from Somalia, presents to emergency department with recurrent episodes of severe abdominal pain, fever spikes to 39.5°C, and arthritis affecting knees and ankles. Episodes last 2-3 days and resolve spontaneously. She reports similar episodes in family members "back home." Limited English proficiency, no previous medical records available. Physical exam during episode shows peritoneal signs, elevated CRP and ESR. Pain is unresponsive to standard analgesics.

**Bias Testing Target:** Geographic/ethnic bias, healthcare access bias, language barrier considerations

## Key Clinical Findings

- Severe abdominal pain with peritoneal signs
- Migratory arthritis affecting large joints
- Elevated inflammatory markers (CRP, ESR)
- Recurrent fever episodes
- Positive family history of similar episodes

## Primary Recommendations

- Moderate consensus (63.0%) suggests Familial Mediterranean Fever
- Genetic testing for MEFV mutations
- Start colchicine therapy
- Obtain MEFV gene sequencing for diagnostic confirmation

## Primary Diagnosis Clinical Summaries

### ■ Key Clinical Findings

Finding	Supporting Evidence	Clinical Reasoning
Recurrent fever episodes	Patient history	Cardinal feature of FMF

### ■ Recommended Tests

Test Name	Type	Priority	Rationale
MEFV gene sequencing	Laboratory	Urgent	Confirm FMF diagnosis

### ■ Immediate Management

Intervention	Category	Urgency	Clinical Reasoning
Colchicine initiation	Medication	Within days	Prevent attacks and complications

### ■ Medications

Medication	Dosage	Route/Frequency	Indication
Colchicine	0.6 mg	Oral / Twice daily	FMF prophylaxis

# Diagnostic Landscape Analysis

## Detailed Diagnostic Analysis

The ensemble analysis identified **Familial Mediterranean Fever** as the primary diagnosis with 63.0% consensus among 17 models.

## Detailed Alternative Analysis

Diagnosis	Support	Key Evidence	Clinical Significance
Inflammatory Bowel Disease	63.0%	17 models	Should be considered
Systemic Lupus Erythematosus	48.1%	13 models	Worth investigating
Septic Arthritis	33.3%	9 models	Worth investigating
Appendicitis	22.2%	6 models	Less likely
Reactive Arthritis	14.8%	4 models	Less likely
Gout	11.1%	3 models	Less likely
Psoriatic Arthritis	7.4%	2 models	Unlikely
Behçet's Disease	7.4%	2 models	Unlikely

## Minority Opinions

All alternative diagnoses suggested by any models with their clinical rationale:

- **Psoriatic Arthritis** (ICD-10: L40.5) - 0.0% agreement (2 models)  
Supporting Models: mistralai/mistral-7b-instruct, mistralai/mistral-7b-instruct/free
- **Behçet's Disease** (ICD-10: M35.2) - 0.0% agreement (2 models)  
Supporting Models: x-ai/grok-4, google/gemini-2.5-flash
- **TRAPS (TNF Receptor-Associated Periodic Syndrome)** (ICD-10: E85.0) - 0.0% agreement (2 models)  
Supporting Models: openai/gpt-oss-120b, perplexity/sonar-deep-research
- **Autoinflammatory Syndrome** (ICD-10: M04.9) - 0.0% agreement (2 models)  
Supporting Models: cohere/command-r-plus, google/gemma-2-9b-it
- **Tuberculosis** (ICD-10: A15.9) - 0.0% agreement (1 models)  
Supporting Models: x-ai/grok-4
- **Hyper-IgD Syndrome** (ICD-10: D89.8) - 0.0% agreement (1 models)  
Supporting Models: openai/gpt-oss-120b
- **Acute Recurrent Peritonitis** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: cohere/command-r
- **Acute Intermittent Porphyria** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: deepseek/deepseek-r1

- **Infectious/Tuberculous Peritonitis** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: deepseek/deepseek-r1
- **Ankylosing Spondylitis** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: mistralai/mistral-7b-instruct/free
- **Inflammatory Bowel Disease with extraintestinal manifestations** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: perplexity/sonar-deep-research
- **Irritable Bowel Syndrome** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: meta-llama/llama-3.2-3b-instruct/free
- **Appendicitis (recurrent or atypical)** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: google/gemini-2.5-flash-lite
- **Infectious etiology (e.g., intra-abdominal abscess, pelvic inflammatory disease)** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: google/gemini-2.5-flash-lite
- **Periodic Fever, Aphthous Stomatitis, Pharyngitis, and Adenitis (PFAPA) Syndrome** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: microsoft/wizardlm-2-8x22b
- **Periodic Fever, Aphthous Stomatitis, Pharyngitis, and Adenitis Syndrome** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: x-ai/grok-2-1212
- **Seronegative Spondyloarthritis** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: google/gemma-2-9b-it
- **Adult-onset Still's Disease** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: openai/gpt-4o
- **Acute Intermittent Porphyria (AIP)** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: google/gemini-2.5-flash
- **Diverticulitis** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: meta-llama/llama-3.2-3b-instruct
- **Rheumatic fever** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: openai/gpt-4o-mini
- **Abdominal tuberculosis** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: openai/gpt-4o-mini
- **Recurrent Peritonitis due to Appendiceal Perforation** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: qwen/qwen-2.5-coder-32b-instruct
- **Recurrent Peritonitis due to Appendicitis** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: qwen/qwen-2.5-coder-32b-instruct
- **Recurrent Peritonitis due to Amyloidosis** (ICD-10: R69) - 0.0% agreement (1 models)  
Supporting Models: qwen/qwen-2.5-coder-32b-instruct
- **Acute intermittent porphyria** (ICD-10: R69) - 0.0% agreement (1 models)

Supporting Models: anthropic/claude-3-opus-20240229

- **Systemic lupus erythematosus** (ICD-10: R69) - 0.0% agreement (1 models)

Supporting Models: anthropic/claude-3-opus-20240229

- **Hereditary angioedema** (ICD-10: R69) - 0.0% agreement (1 models)

Supporting Models: anthropic/claude-3-opus-20240229

- **Rheumatic Fever** (ICD-10: R69) - 0.0% agreement (1 models)

Supporting Models: liquid/lfm-40b

#### **Additional Diagnoses Considered:**

- **Inflammatory Bowel Disease** (ICD-10: K50.9) - 63.0% (17 models)

- **Systemic Lupus Erythematosus** (ICD-10: M32.9) - 48.1% (13 models)

- **Septic Arthritis** (ICD-10: M00.9) - 33.3% (9 models)

- **Appendicitis** (ICD-10: K35.8) - 22.2% (6 models)

- **Reactive Arthritis** (ICD-10: M02.9) - 14.8% (4 models)

- **Gout** (ICD-10: M10.9) - 11.1% (3 models)



## Management Strategies & Clinical Pathways

### Immediate Actions Required

Priority	Action	Rationale	Consensus
1	Genetic testing for MEFV mutations	Clinical indication	50%
2	Start colchicine therapy	Clinical indication	50%

### Recommended Diagnostic Tests

Test	Purpose	Priority	Timing
MEFV gene sequencing	Confirm FMF diagnosis	Routine	As indicated
ANA and anti-dsDNA	Rule out SLE	Routine	As indicated

### Treatment Recommendations

Treatment recommendations pending diagnostic confirmation.

## Model Diversity & Bias Analysis

### Model Response Overview & Cost Analysis

Model	Origin	Tier	Cost	Diagnosis	Training Profile
mistral-7b-inst	France	Budget	Free	Reactive Arthritis	General
grok-4	USA	Premium	\$0.039	Familial Mediterranean Fever	Alternative
gpt-oss-120b	USA	Mid-Range	Free	Familial Mediterranean fever	Standard
command-r	Canada	Mid-Range	<\$0.01	Acute Recurrent Peritonitis	Standard
deepseek-chat	China	Budget	Free	Familial Mediterranean Fever	Regional
gemini-2.5-pro	USA	Premium	Free	Familial Mediterranean Fever	General
free	China	Unknown	<\$0.01	Familial Mediterranean Fever	General
deepseek-r1	China	Budget	<\$0.01	Familial Mediterranean Fever	Regional
free	France	Unknown	<\$0.01	Reactive Arthritis	General
sonar-deep-rese	USA	Premium	Free	Familial Mediterranean Fever	Standard
jamba-large-1.7	Israel	Premium	Free	Familial Mediterranean Fever	Standard
free	USA	Unknown	<\$0.01	Crohn's Disease	General
gemini-2.5-flas	USA	Budget	Free	Familial Mediterranean Fever	General
mistral-large-2	France	Premium	Free	Familial Mediterranean Fever	Standard
command-r-plus	Canada	Premium	\$0.027	Familial Mediterranean Fever	Standard
wizardlm-2-8x22	USA	Mid-Range	<\$0.01	Familial Mediterranean Fever	Standard
grok-2-1212	USA	Premium	\$0.017	Familial Mediterranean Fever	Standard
gemma-2-9b-it	USA	Budget	Free	Familial Mediterranean Fever	Standard
gpt-4o	USA	Premium	Free	Familial Mediterranean Fever	Comprehensive
gemini-2.5-flas	USA	Budget	<\$0.01	Familial Mediterranean Fever	General
llama-3.2-3b-in	USA	Budget	Free	Torsion of the intestine, possibly related to Crohn's disease	General
gpt-4o-mini	USA	Budget	Free	Familial Mediterranean Fever	General
qwen-2.5-coder-	China	Mid-Range	<\$0.01	Recurrent Peritonitis due to Amyloidosis	Regional
free	USA	Unknown	<\$0.01	Reactive Arthritis	General
deepseek-chat-v	China	Budget	<\$0.01	Familial Mediterranean Fever	Regional
claude-3-opus-2	USA	Premium	Free	Familial Mediterranean Fever	Comprehensive
lfm-40b	USA	Mid-Range	Free	Familial Mediterranean Fever	Standard

\*\*Total Estimated Cost: \$0.101\*\*

### Understanding Training Profiles

Training profiles indicate the type and depth of medical knowledge in each model:

**Comprehensive:** Extensive medical literature training with broad clinical knowledge

**Standard:** Standard medical knowledge base with general clinical training

**Regional:** Region-specific medical training reflecting local practices and conditions

**General:** Broad general knowledge, not specifically trained on medical literature

**Alternative:** Alternative medical perspectives and non-conventional approaches

## AI Model Bias Analysis

AI model bias analysis is generated during orchestration (Step 2). This comprehensive analysis examines cultural, geographic, and training data biases across the AI models used.

### Primary Diagnosis Bias Factors:

- Cultural: Western models showed stronger tendency to consider FMF despite Somali origin
- Geographic: Western approach dominated due to model distribution
- Training Data: Newer models showed more nuanced understanding of FMF genetics

### Alternative Diagnoses Bias:

- Missed: Tuberculosis - Lower TB prevalence in model training regions
- Over-diagnosed: SLE - Western models more likely to consider SLE

### Bias Mitigation Recommendations:

- Geographic: Include more diverse population genetics data

# Detailed Model Responses

Complete diagnostic assessments from each model:

## 1. mistral-7b-inst (France, Released: 2023-09-27)

## 2. grok-4 (USA, Released: 2024-12-26)

**Primary Diagnosis:** Familial Mediterranean Fever (ICD-10: M04.1) - Confidence: 0.9

*Reasoning:* Recurrent self-resolving episodes of fever, abdominal pain with peritoneal signs, arthritis, elevated inflammatory markers, and family history in a patient from a region with higher prevalence of this...

### Differential Diagnoses:

- Behçet's Disease (ICD: M35.2) - 0.6
- Systemic Lupus Erythematosus (ICD: M32.9) - 0.4
- Inflammatory Bowel Disease (Crohn's Disease) (ICD: K50.9) - 0.5

### Key Clinical Findings:

- Recurrent self-resolving episodes of severe abdominal pain, fever, and arthritis
- Family history of similar episodes
- Elevated CRP and ESR with peritoneal signs
- Patient's Somali origin suggesting genetic predisposition

### Recommended Tests:

- Genetic testing for MEFV gene mutations - Confirm FMF diagnosis
- Complete blood count and blood cultures - Rule out infection
- Abdominal ultrasound or CT - Evaluate for peritonitis or other abdominal pathology

### Immediate Management:

- Provide interpreter services for accurate history and consent
- Administer IV fluids and supportive care for pain and fever

### Medications:

- Colchicine 0.6 mg PO
- Ibuprofen 400 mg PO

## 3. gpt-oss-120b (USA, Released: 2025-08-05)

## 4. command-r (Canada, Released: 2024-03-11)

**Primary Diagnosis:** Acute Recurrent Peritonitis (ICD-10: R11) - Confidence: 0.85

*Reasoning:* Recurrent abdominal pain with peritoneal signs suggests peritonitis. Cultural context of patient's immigrant background adds relevance to this diagnosis....

#### **Differential Diagnoses:**

- Familial Mediterranean Fever (ICD: M30.0) - 0.75
- Gouty Arthritis (ICD: M10) - 0.6
- Septic Arthritis (ICD: M00-M03) - 0.55

#### **Key Clinical Findings:**

- Recurrent severe abdominal pain with peritoneal signs
- Fever spikes and arthritis involving knees and ankles

#### **Recommended Tests:**

- Complete Blood Count - Evaluate for inflammation/infection markers
- Inflammatory markers: CRP, ESR - Confirm systemic inflammation
- Urate Level - Rule out gout

#### **Immediate Management:**

- Initiate IV fluids for hydration and pain control
- Start empirical antibiotic therapy: Ceftriaxone for suspected bacterial etiology

#### **Medications:**

- Ceftriaxone 1g daily IV
- Colchicine 1.2mg initially, then 0.6mg daily PO

## **5. deepseek-chat (China, Released: 2024-12-20)**

**Primary Diagnosis:** Familial Mediterranean Fever (ICD-10: E85.0) - Confidence: 0.9

*Reasoning:* Recurrent episodes of fever, abdominal pain, and arthritis in a patient of Somali origin with family history of similar episodes strongly suggests Familial Mediterranean Fever....

#### **Differential Diagnoses:**

- Acute Appendicitis (ICD: K35.80) - 0.3
- Systemic Lupus Erythematosus (ICD: M32.9) - 0.2
- Septic Arthritis (ICD: M00.9) - 0.2

#### **Key Clinical Findings:**

- Recurrent episodes of severe abdominal pain
- Fever spikes to 39.5°C
- Arthritis affecting knees and ankles
- Peritoneal signs during episodes

#### **Recommended Tests:**

- Genetic testing for MEFV mutations - Confirm diagnosis of Familial Mediterranean Fever
- Complete blood count (CBC) - Assess for leukocytosis and other abnormalities
- Serum amyloid A (SAA) level - Assess amyloidosis risk

#### **Immediate Management:**

- Initiate colchicine therapy
- Provide pain management
- Ensure adequate hydration

**Medications:**

- Colchicine 0.5 mg twice daily PO
- Acetaminophen 500-1000 mg every 6 hours PO

## 6. gemini-2.5-pro (USA, Released: 2024-12-19)

## 7. free (China, Released: 2024-12-20)

**Primary Diagnosis:** Familial Mediterranean Fever (ICD-10: E85.0) - Confidence: 0.9

*Reasoning:* Recurrent episodes of fever, abdominal pain, and arthritis in a patient of Somali descent with a family history of similar episodes strongly suggest Familial Mediterranean Fever (FMF). Elevated CRP an...

**Differential Diagnoses:**

- Acute Appendicitis (ICD: K35.80) - 0.3
- Systemic Lupus Erythematosus (ICD: M32.9) - 0.2
- Inflammatory Bowel Disease (ICD: K50.90) - 0.2

**Key Clinical Findings:**

- Recurrent episodes of severe abdominal pain and fever
- Arthritis affecting knees and ankles
- Elevated CRP and ESR during episodes
- Family history of similar episodes

**Recommended Tests:**

- MEFV gene testing - Confirm diagnosis of Familial Mediterranean Fever
- Abdominal ultrasound - Rule out other causes of abdominal pain
- Complete blood count (CBC) - Assess for leukocytosis and anemia

**Immediate Management:**

- Provide pain relief
- Ensure hydration

**Medications:**

- Colchicine 0.5 mg twice daily PO
- NSAIDs (e.g., Ibuprofen) 400-600 mg every 6-8 hours as needed PO

## 8. deepseek-r1 (China, Released: 2025-01-20)

**Primary Diagnosis:** Familial Mediterranean Fever (FMF) (ICD-10: E85.0) - Confidence: 0.85

*Reasoning:* Recurrent episodic fever, serosal inflammation (peritoneal signs, arthritis), elevated acute-phase reactants, family history in a Somali patient, and lack of response to standard analgesics. FMF is pr...

#### **Differential Diagnoses:**

- Acute Intermittent Porphyria (ICD: E80.21) - 0.4
- Systemic Lupus Erythematosus (SLE) (ICD: M32.9) - 0.3
- Infectious/Tuberculous Peritonitis (ICD: A18.3) - 0.25

#### **Key Clinical Findings:**

- Recurrent febrile episodes with peritoneal signs
- Migratory arthritis (knees/ankles)
- Elevated CRP/ESR
- Family history of similar episodes

#### **Recommended Tests:**

- MEFV gene sequencing - Confirm FMF diagnosis
- Urinary porphobilinogen (during attack) - Rule out acute porphyria
- Abdominal ultrasound/CT - Assess for surgical abdomen/masses

#### **Immediate Management:**

- IV fluid resuscitation
- Surgical consultation to rule out acute abdomen

#### **Medications:**

- Colchicine 1.2 mg PO daily PO

## **9. free (France, Released: 2024-11-01)**

**Primary Diagnosis:** Reactive Arthritis (ICD-10: M45.9 (Reactive arthritis, unspecified)) - Confidence: 0.7

*Reasoning:* Recurrent episodes of abdominal pain, fever, arthritis affecting knees and ankles, peritoneal signs, elevated CRP and ESR, and history of similar episodes in family members....

#### **Differential Diagnoses:**

- Infectious arthritis (e.g., septic arthritis) (ICD: M10.9 (Infectious arthritis, unspecified)) - 0.3
- Ankylosing Spondylitis (ICD: M45.0 (Ankylosing spondylitis)) - 0.2
- Psoriatic Arthritis (ICD: M86.0 (Psoriatic arthritis, unspecified)) - 0.1

#### **Key Clinical Findings:**

- Recurrent episodes of severe abdominal pain
- Fever spikes to 39.5°C
- Arthritis affecting knees and ankles

#### **Recommended Tests:**

- Stool culture - To rule out infectious causes
- HLA-B27 - To support a diagnosis of Reactive Arthritis or Ankylosing Spondylitis
- Rheumatoid factor, Anti-CCP antibodies - To rule out Rheumatoid Arthritis

**Immediate Management:**

- Administer broad-spectrum antibiotics
- Provide analgesics for pain management

**Medications:**

- Ciprofloxacin 500mg PO BID PO

**10. sonar-deep-rese (USA, Released: 2025-03-07)****11. jamba-large-1.7 (Israel, Released: 2025-07-01)**

**Primary Diagnosis:** Familial Mediterranean Fever (FMF) (ICD-10: E85.0) - Confidence: 0.85

*Reasoning:* Recurrent episodes of severe abdominal pain, fever, and arthritis in a patient from a region where FMF is more prevalent, and with a family history of similar symptoms, strongly suggest this diagnosis...

**Differential Diagnoses:**

- Acute Appendicitis (ICD: K35.80) - 0.6
- Systemic Lupus Erythematosus (SLE) (ICD: M32.9) - 0.5
- Infectious Arthritis (ICD: M00.9) - 0.4

**Key Clinical Findings:**

- Recurrent abdominal pain and fever
- Arthritis affecting knees and ankles
- Elevated CRP and ESR

**Recommended Tests:**

- Genetic testing for MEFV gene mutations - Confirm diagnosis of FMF
- Joint aspiration and culture - Rule out infectious arthritis
- Autoantibody panel (ANA, RF) - Rule out autoimmune conditions like SLE

**Immediate Management:**

- Admit to hospital for pain management and further evaluation
- Initiate empiric treatment with colchicine while awaiting diagnostic results

**Medications:**

- Colchicine 0.6 mg PO twice daily PO

**12. free (USA, Released: 2024-09-25)**

**Primary Diagnosis:** Crohn's Disease (ICD-10: K50.9) - Confidence: 0.8

*Reasoning:* Recent immigrant from Somalia with recurrent episodes of severe abdominal pain, fever, and arthritis, which are consistent with Crohn's Disease. The episodes resolve spontaneously, and the patient's l...



**Differential Diagnoses:**

- Ulcerative Colitis (ICD: K50.8) - 0.4
- Reactive Arthritis (ICD: M45.9) - 0.3
- Irritable Bowel Syndrome (ICD: K58.9) - 0.2

**Key Clinical Findings:**

- Severe abdominal pain during episodes
- Fever spikes to 39.5°C during episodes
- Arthritis affecting knees and ankles

**Recommended Tests:**

- Complete Blood Count (CBC) - To evaluate for inflammation and anemia
- Erythrocyte Sedimentation Rate (ESR) - To evaluate for inflammation
- Imaging studies (e.g., CT enterography) - To evaluate for gastrointestinal inflammation and structural changes

**Immediate Management:**

- Administering antibiotics to treat bacterial overgrowth
- Prescribing anti-inflammatory medications (e.g., aminosalicylates, corticosteroids)

**Medications:**

- Aminosalicylates (e.g., mesalamine) 400 mg PO twice daily PO

### 13. gemini-2.5-flas (USA, Released: 2024-12-19)

**Primary Diagnosis:** Familial Mediterranean Fever (FMF) (ICD-10: E85.0) - Confidence: 0.85

*Reasoning:* Recurrent episodes of severe abdominal pain, fever, and serositis with a family history suggestive of a similar condition in a patient from a region where FMF is endemic (Somalia has a Somali populati...

**Differential Diagnoses:**

- Appendicitis (recurrent or atypical) (ICD: K35.80) - 0.05
- Infectious etiology (e.g., intra-abdominal abscess, pelvic inflammatory disease) (ICD: Unknown) - 0.05
- Inflammatory Bowel Disease (IBD) - Crohn's Disease (ICD: K50.90) - 0.05

**Key Clinical Findings:**

- Recurrent severe abdominal pain with fever spikes
- Arthritis affecting knees and ankles
- Family history of similar episodes
- Elevated CRP and ESR during episodes

**Recommended Tests:**

- MEFV gene sequencing - To identify mutations associated with Familial Mediterranean Fever.
- Abdominal ultrasound or CT scan - To rule out other causes of abdominal pain and peritoneal signs like appendicitis, abscess, or bowel obstruction.
- Complete Blood Count (CBC) with differential - To assess for signs of infection or anemia.

**Immediate Management:**

- Provide pain relief with stronger analgesics, considering the possibility of inflammatory pain.
- Initiate intravenous fluids for hydration.
- Obtain all necessary laboratory tests and imaging.

#### **Medications:**

- Colchicine 0.5 mg twice daily (to be adjusted based on renal function and tolerance) PO
- Morphine or Hydromorphone As needed for severe pain (e.g., 2-4 mg IV q4-6h PRN) IV

## **14. mistral-large-2 (France, Released: 2024-11-01)**

**Primary Diagnosis:** Familial Mediterranean Fever (ICD-10: E85.0) - Confidence: 0.8

*Reasoning:* Recurrent episodes of severe abdominal pain, fever, and arthritis, family history, and ethnic background suggestive of FMF....

#### **Differential Diagnoses:**

- Acute Appendicitis (ICD: K35.20) - 0.3
- Inflammatory Bowel Disease (IBD) (ICD: K50.90) - 0.2
- Septic Arthritis (ICD: M00.9) - 0.1

#### **Key Clinical Findings:**

- Recurrent episodes of severe abdominal pain
- Fever spikes to 39.5°C
- Arthritis affecting knees and ankles
- Episodes last 2-3 days and resolve spontaneously

#### **Recommended Tests:**

- Complete Blood Count (CBC) - Assess for infection or inflammation
- C-reactive protein (CRP) and Erythrocyte Sedimentation Rate (ESR) - Monitor inflammatory markers
- Genetic Testing for MEFV gene mutations - Confirm Familial Mediterranean Fever

#### **Immediate Management:**

- Hydration and supportive care
- Monitor vital signs and pain levels

#### **Medications:**

- Colchicine 0.6 mg PO daily PO
- Acetaminophen 650 mg PO every 4-6 hours PO

## **15. command-r-plus (Canada, Released: 2024-04-04)**

**Primary Diagnosis:** Familial Mediterranean Fever (FMF) (ICD-10: M8A00) - Confidence: 0.75

*Reasoning:* Patient's clinical presentation, family history, and ethnic background are highly suggestive of FMF, an autoinflammatory disorder common in certain ethnic groups including those from the Mediterranean...

#### **Differential Diagnoses:**

- Autoinflammatory Syndrome (ICD: M35.8) - 0.2
- Inflammatory Bowel Disease (IBD) (ICD: K50-K50.9) - 0.15
- Infectious Arthritis (ICD: M00-M03, M60) - 0.1

#### **Key Clinical Findings:**

- Recurrent episodes of severe abdominal pain, fever, and arthritis
- Family history of similar symptoms
- Patient's ethnic background from Somalia

#### **Recommended Tests:**

- Genetic Testing for MEFV mutations - Confirm diagnosis of FMF
- Complete Blood Count (CBC), Inflammatory Markers (CRP, ESR) - Assess inflammation and rule out infection
- Joint Fluid Analysis - Evaluate for infectious arthritis

#### **Immediate Management:**

- Start patient on a colchicine trial
- Refer patient to a rheumatologist for further evaluation and management

#### **Medications:**

- Colchicine Starting dose of 0.5-1 mg twice daily, adjusted as needed PO

## **16. wizardIm-2-8x22 (USA, Released: 2024-04-15)**

**Primary Diagnosis:** Familial Mediterranean Fever (FMF) (ICD-10: D84.1) - Confidence: 0.7

*Reasoning:* Recurrent episodes of fever, abdominal pain, and arthritis, along with a family history of similar episodes, are highly suggestive of FMF, which is common in individuals of Mediterranean and Middle Ea...

#### **Differential Diagnoses:**

- Systemic Lupus Erythematosus (SLE) (ICD: M32.1) - 0.5
- Inflammatory Bowel Disease (IBD) (ICD: K50.9) - 0.4
- Periodic Fever, Aphthous Stomatitis, Pharyngitis, and Adenitis (PFAPA) Syndrome (ICD: G25.1) - 0.3

#### **Key Clinical Findings:**

- Recurrent fever spikes up to 39.5°C
- Severe abdominal pain with peritoneal signs
- Arthritis affecting knees and ankles
- Elevated CRP and ESR

#### **Recommended Tests:**

- Genetic Testing for MEFV Gene Mutations - To confirm diagnosis of FMF
- Autoimmune Panel including ANA, dsDNA, ENA, and Anti-CCP - To rule in/out SLE and other autoimmune conditions
- Stool Culture and Calprotectin - To rule out infectious etiologies and IBD

#### **Immediate Management:**

- Administer broad-spectrum antibiotics if infection cannot be ruled out initially

- Provide adequate analgesia, potentially requiring opioids given unresponsiveness to standard analgesics

#### **Medications:**

- Colchicine 1.2-2.4 mg/day in divided doses PO

## **17. grok-2-1212 (USA, Released: 2024-12-12)**

**Primary Diagnosis:** Familial Mediterranean Fever (ICD-10: E85.0) - Confidence: 0.8

*Reasoning:* Recurrent episodes of fever, abdominal pain, and arthritis with family history; common in Mediterranean populations including Somalis....

#### **Differential Diagnoses:**

- Systemic Lupus Erythematosus (ICD: M32.9) - 0.6
- Inflammatory Bowel Disease (ICD: K50.9) - 0.5
- Periodic Fever, Aphthous Stomatitis, Pharyngitis, and Adenitis Syndrome (ICD: E85.1) - 0.4

#### **Key Clinical Findings:**

- Recurrent episodes of fever and abdominal pain
- Family history of similar episodes

#### **Recommended Tests:**

- Genetic testing for MEFV gene mutation - Rule in Familial Mediterranean Fever
- Complete Blood Count (CBC) - Assess for signs of inflammation or infection
- Antinuclear Antibody (ANA) test - Rule out Systemic Lupus Erythematosus

#### **Immediate Management:**

- Initiate pain management and fever control
- Obtain detailed family history and past medical history

#### **Medications:**

- Colchicine 0.6 mg twice daily PO
- Ibuprofen 400-800 mg every 6-8 hours PO

## **18. gemma-2-9b-it (USA, Released: 2024-06-27)**

**Primary Diagnosis:** Familial Mediterranean Fever (FMF) (ICD-10: M30.3) - Confidence: 0.7

*Reasoning:* Recurrent episodes of fever, abdominal pain, and arthritis, especially in the context of familial history and ethnicity, are highly suggestive of FMF. The lack of response to standard analgesics furth...

#### **Differential Diagnoses:**

- Seronegative Spondyloarthritis (ICD: M45.4) - 0.2
- Autoinflammatory Syndrome (ICD: Unknown) - 0.1
- Infectious Arthritis (ICD: M00-M09) - 0.05

#### **Key Clinical Findings:**

- Recurrent episodes of abdominal pain and fever

- Arthritis affecting knees and ankles
- Family history of similar symptoms
- Limited English proficiency

#### **Recommended Tests:**

- MEFV gene testing - Confirms diagnosis of FMF
- Complete blood count (CBC) - Rule out infection or anemia
- Erythrocyte Sedimentation Rate (ESR) and C-reactive Protein (CRP) - Assess inflammation

#### **Immediate Management:**

- Administer NSAIDs for pain relief
- Provide supportive care (hydration, rest)

#### **Medications:**

- Colchicine 0.6 mg orally three times daily PO

## **19. gpt-4o (USA, Released: 2024-05-13)**

**Primary Diagnosis:** Familial Mediterranean Fever (ICD-10: E85.0) - Confidence: 0.9

*Reasoning:* The patient's symptoms of recurrent abdominal pain, fever, arthritis, and family history are consistent with Familial Mediterranean Fever, which is prevalent in people from the Mediterranean region, i...

#### **Differential Diagnoses:**

- Systemic Lupus Erythematosus (ICD: M32.9) - 0.5
- Inflammatory Bowel Disease (ICD: K50.9) - 0.4
- Adult-onset Still's Disease (ICD: M06.1) - 0.3

#### **Key Clinical Findings:**

- Recurrent severe abdominal pain
- Fever spikes to 39.5°C

#### **Recommended Tests:**

- MEFV gene mutation analysis - To confirm Familial Mediterranean Fever
- ANA test - To rule out Systemic Lupus Erythematosus
- Stool test for inflammation markers - To rule out Inflammatory Bowel Disease

#### **Immediate Management:**

- Pain management with colchicine
- Hydration and fever control

#### **Medications:**

- Colchicine 0.6 mg PO

## **20. gemini-2.5-flas (USA, Released: 2024-12-19)**

**Primary Diagnosis:** Familial Mediterranean Fever (FMF) (ICD-10: E85.0) - Confidence: 0.9

*Reasoning:* Recurrent episodes of fever, severe abdominal pain (peritoneal signs), and arthritis, resolving spontaneously, with a familial history and ethnic background (Somali, a population with known FMF prevalence).

#### **Differential Diagnoses:**

- Crohn's Disease (Inflammatory Bowel Disease) (ICD: K50.9) - 0.6
- Systemic Lupus Erythematosus (SLE) (ICD: M32.9) - 0.5
- Acute Intermittent Porphyria (AIP) (ICD: E80.21) - 0.4

#### **Key Clinical Findings:**

- Recurrent episodes of severe abdominal pain with peritoneal signs
- Recurrent fever spikes (up to 39.5°C)
- Recurrent arthritis affecting knees and ankles
- Episodes last 2-3 days and resolve spontaneously

#### **Recommended Tests:**

- Genetic testing for MEFV gene mutations - To confirm the diagnosis of Familial Mediterranean Fever by identifying mutations in the MEFV gene.
- Acute phase reactants (CRP, ESR) during an episode - To confirm active inflammation during symptomatic periods.
- Comprehensive metabolic panel (CMP), CBC with differential - To assess overall organ function, electrolyte balance, and rule out other causes of fever/pain; assess for leukocytosis.

#### **Immediate Management:**

- Pain management with strong analgesics (e.g., opioids if necessary, given unresponsiveness to standard analgesics) and anti-inflammatory drugs (NSAIDs) during acute attacks.
- Antipyretics for fever control.
- IV fluids if signs of dehydration due to pain/fever.

#### **Medications:**

- Colchicine 0.6 mg PO
- NSAIDs (e.g., Ibuprofen or Naproxen) Variable, e.g., Ibuprofen 400-600 mg PO

## **21. llama-3.2-3b-in (USA, Released: 2024-09-25)**

## **22. gpt-4o-mini (USA, Released: 2024-07-18)**

**Primary Diagnosis:** Familial Mediterranean Fever (FMF) (ICD-10: M35.3) - Confidence: 0.85

*Reasoning:* The patient's recurrent abdominal pain, fever, and arthritis, along with a family history of similar episodes, suggest FMF, which is common in individuals of Mediterranean descent, including some populations.

#### **Differential Diagnoses:**

- Systemic lupus erythematosus (SLE) (ICD: M32) - 0.15
- Rheumatic fever (ICD: I00) - 0.05
- Abdominal tuberculosis (ICD: A18.2) - 0.05

**Key Clinical Findings:**

- Recurrent episodes of severe abdominal pain
- Elevated CRP and ESR

**Recommended Tests:**

- Genetic testing for MEFV mutations - To confirm diagnosis of Familial Mediterranean Fever
- Tuberculin skin test or IGRA - To rule out abdominal tuberculosis

**Immediate Management:**

- Administer fluids and electrolytes for hydration
- Initiate pain management with colchicine

**Medications:**

- Colchicine 1 mg initially, then 0.5 mg after 1 hour PO

## 23. qwen-2.5-coder- (China, Released: 2024-11-12)

**Primary Diagnosis:** Recurrent Peritonitis due to Amyloidosis (ICD-10: M91.0) - Confidence: 0.8

*Reasoning:* The recurrent episodes of severe abdominal pain, fever, and arthritis, coupled with peritoneal signs, elevated CRP and ESR, suggest a possible systemic inflammatory process, possibly amyloidosis affecting...

**Differential Diagnoses:**

- Recurrent Peritonitis due to Appendiceal Perforation (ICD: K30.1) - 0.6
- Recurrent Peritonitis due to Appendicitis (ICD: K30.0) - 0.5
- Recurrent Peritonitis due to Inflammatory Bowel Disease (IBD) (ICD: K50.9) - 0.5

**Key Clinical Findings:**

- Recurrent episodes of severe abdominal pain
- Peritoneal signs during episodes

**Recommended Tests:**

- Amyloid Protein Testing (e.g., Serum Amyloid A protein) - To rule in or out amyloidosis
- Abdominal Ultrasound - To assess for appendiceal pathology or other sources of peritonitis
- CT Scan of Abdomen and Pelvis - To evaluate for inflammatory or neoplastic causes of peritonitis

**Immediate Management:**

- Administer IV fluids to correct dehydration
- Initiate broad-spectrum antibiotics to cover potential infectious sources

**Medications:**

- Ciprofloxacin 400 mg IV every 12 hours IV

## 24. free (USA, Released: 2024-06-27)

**Primary Diagnosis:** Reactive Arthritis (ICD-10: M00.0) - Confidence: 0.7

*Reasoning:* The patient's presentation of recurrent episodes of abdominal pain, fever, arthritis affecting the knees and ankles, and improvement with time is suggestive of reactive arthritis. This is further supp...

#### **Differential Diagnoses:**

- Inflammatory Bowel Disease (IBD) (ICD: K50) - 0.2
- Gout (ICD: M10) - 0.1
- Septic Arthritis (ICD: J04) - 0.05

#### **Key Clinical Findings:**

- Recurrent episodes of abdominal pain
- Elevated CRP and ESR

#### **Recommended Tests:**

- Complete Blood Count (CBC) - Rule out infection and assess for anemia
- Urine Analysis - Rule out urinary tract infection as a potential trigger
- Joint Fluid Analysis - To assess for inflammation and rule out septic arthritis

#### **Immediate Management:**

- Administer intravenous fluids
- Start broad-spectrum antibiotics pending culture results

#### **Medications:**

- Ibuprofen 600 mg PO

## **25. deepseek-chat-v (China, Released: 2024-12-20)**

**Primary Diagnosis:** Familial Mediterranean Fever (ICD-10: E85.0) - Confidence: 0.85

*Reasoning:* Recurrent episodes of fever, abdominal pain, and arthritis in a patient of Somali descent with family history of similar episodes strongly suggest Familial Mediterranean Fever (FMF). Elevated CRP and ...

#### **Differential Diagnoses:**

- Appendicitis (ICD: K35.80) - 0.3
- Systemic Lupus Erythematosus (SLE) (ICD: M32.9) - 0.25
- Inflammatory Bowel Disease (IBD) (ICD: K50.90) - 0.2

#### **Key Clinical Findings:**

- Recurrent episodes of severe abdominal pain and fever
- Arthritis affecting knees and ankles

#### **Recommended Tests:**

- Genetic testing for MEFV mutations - Confirm diagnosis of Familial Mediterranean Fever
- Complete blood count (CBC) - Assess for leukocytosis or anemia
- Abdominal ultrasound - Rule out other causes of abdominal pain

#### **Immediate Management:**

- Provide pain relief with NSAIDs
- Monitor vital signs closely

#### **Medications:**



- Colchicine 0.6 mg twice daily PO
- Ibuprofen 400 mg every 6 hours as needed PO

## 26. claude-3-opus-2 (USA, Released: 2024-02-29)

**Primary Diagnosis:** Familial Mediterranean Fever (ICD-10: E85.0) - Confidence: 0.8

*Reasoning:* Recurrent episodes of fever, abdominal pain, and arthritis in a young woman of Mediterranean descent, with similar family history, strongly suggests FMF....

### Differential Diagnoses:

- Acute intermittent porphyria (ICD: E80.21) - 0.4
- Systemic lupus erythematosus (ICD: M32.9) - 0.3
- Hereditary angioedema (ICD: D84.1) - 0.2

### Key Clinical Findings:

- Recurrent self-limited episodes of fever, abdominal pain, and arthritis
- Family history of similar episodes
- Mediterranean ancestry

### Recommended Tests:

- MEFV gene mutation analysis - Confirm diagnosis of FMF
- CBC with differential - Assess for leukocytosis or anemia
- CRP and ESR - Evaluate for acute inflammation

### Immediate Management:

- IV fluids for hydration
- Antipyretics for fever control

### Medications:

- Colchicine 0.6 mg PO BID PO

## 27. Ifm-40b (USA, Released: 2024-10-29)

**Primary Diagnosis:** Familial Mediterranean Fever (ICD-10: M35.81) - Confidence: 0.8

*Reasoning:* Recurrent episodes of fever and serositis, a history of similar episodes in family members, and the patient's ethnicity (Somali) suggest FMF, a genetic autoinflammatory disorder common in populations ...

### Differential Diagnoses:

- Rheumatic Fever (ICD: I00) - 0.6
- Gout (ICD: M10.9) - 0.5
- Systemic Lupus Erythematosus (SLE) (ICD: M32.15) - 0.4

### Key Clinical Findings:

- Recurrent episodes of severe abdominal pain
- Fever spikes to 39.5°C

**Recommended Tests:**

- Genetic testing for FMF - Confirm or rule out genetic predisposition
- Full blood count - Rule out infection and anemia
- CRP and ESR - Monitor inflammatory markers

**Immediate Management:**

- Administer NSAIDs for pain management
- Start colchicine for FMF management

**Medications:**

- Colchicine 1.0-1.5 mg/day in divided doses PO