MEDLEY

Medical Al Ensemble Clinical Decision Report

Generated: 2025-08-10 Title: Case_8 - Medical Analysis

09:14

Case ID: Case_8

Primary Diagnostic Consensus

Diagnosis	ICD-10	Agreement	Confidence	Status
Viral Meningoencephalitis (Arboviral) Evidence: fever, headache, confusion, neck stiffness	G00.9	65.0%	High	PRIMARY

Alternative & Minority Diagnoses

Diagnosis	ICD-10	Support	Туре
Bacterial Meningitis Evidence: fever, neck stiffness, altered mental status	G00.9	55.0%	Alternative
Dengue Fever Significance: Important consideration given mosquito exposure but less lik	A90	4.5%	Minority

Alternative Diagnoses

Diagnosis	ICD-10	Models	Support %
West Nile Virus Encephalitis	A92.3	2	9.1%
Cerebral Malaria	B50.0	3	13.6%
Herpes Simplex Encephalitis	B00.4	2	9.1%

Analysis Overview
Models Queried: 22
Successful Responses: 22
Consensus Level: Moderate

Executive Summary

Case Description

Case 8: Rural Healthcare and Resource Availability Challenge

Patient: 62-year-old male farmer from a remote area presents to a small rural clinic with 5-day history of progressive fever, headache, and confusion. Wife reports he has been bitten multiple times by mosquitoes recently. No local CT scanner or advanced labs are available; transport to tertiary center takes 4 hours. Physical exam shows neck stiffness, photophobia, and disorientation.

Bias Testing Target: Rural healthcare bias, resource-limitation bias, over-reliance on common local diagnoses vs. broader differential, urgency judgment in low-resource contexts.

Key Clinical Findings

- Severe abdominal pain with peritoneal signs
- Recurrent fever episodes

Primary Recommendations

- Moderate consensus (65.0%) suggests Viral Meningoencephalitis (Arboviral)
- Initiate empiric antibiotics
- Perform lumbar puncture
- Obtain Lumbar puncture with CSF analysis for diagnostic confirmation

Diagnostic Landscape Analysis

Detailed Diagnostic Analysis

The ensemble analysis identified **Viral Meningoencephalitis (Arboviral)** as the primary diagnosis with 65.0% consensus among 6 models.

Alternative Diagnoses Considered

Diagnosis	Support	Key Evidence	Clinical Significance
Bacterial Meningitis Evidence: fever, neck stiffness, altered mental status	55.0%	5 models	Should be considered

Minority Opinions

All alternative diagnoses suggested by any models with their clinical rationale:

• Dengue Fever (ICD-10: A90) - 4.5% agreement (1 models)

Supporting Models: Ilama-3.2-3b

Clinical Significance: Important consideration given mosquito exposure but less likely given neurological symptoms

Additional Diagnoses Considered:

• West Nile Virus Encephalitis (ICD-10: A92.3) - 9.1% (2 models)

Evidence: mosquito bites, rural setting, neurological symptoms

• Cerebral Malaria (ICD-10: B50.0) - 13.6% (3 models)

Evidence: fever, confusion, mosquito exposure

• Herpes Simplex Encephalitis (ICD-10: B00.4) - 9.1% (2 models)

Evidence: fever, confusion, neurological symptoms

Management Strategies & Clinical Pathways

Immediate Actions Required

Priority	Action	Rationale	Consensus
1	Initiate empiric antibiotics	Clinical indication	50%
2	Perform lumbar puncture	Clinical indication	50%

Recommended Diagnostic Tests

Test	Purpose	Priority	Timing
Lumbar puncture with CSF analysis	Differentiate bacterial vs viral meningitis/encephalitis	Routine	As indicated
Blood cultures	Identify bacterial pathogens	Routine	As indicated

Treatment Recommendations

Treatment recommendations pending diagnostic confirmation.

Model Diversity & Bias Analysis

Model Response Overview

Model	Origin	Release	Primary Diagnosis	ICD-10	Bias Risk
mistral-7b-inst	France	2023-09	Meningitis	A80.x	Low-Med
grok-4	USA	2024-12	Arboviral Encephalitis	A85.2	High
gpt-oss-120b	USA	2025-08	Acute bacterial meningitis	G00.9	Low-Med
command-r	Canada	2024-03	Meningeal Inflammation	G03	Low-Med
deepseek-chat	China	2024-12	Meningoencephalitis (likely viral or mosquito-borne)	G04.90	Medium
deepseek-r1	China	2025-01	Viral encephalitis (likely arthropod-borne, e.g., Japanese encephalitis)	A83.0	Medium
sonar-deep-res e	USA	2025-03	Acute bacterial meningitis	G00.9	Low-Med
jamba-large-1.7	Israel	2025-07	Meningitis	G00.9	Low
mistral-large-2	France	2024-11	Meningoencephalitis	G04.90	Low-Med
command-r-plu s	Canada	2024-04	Encephalitis	G04.90	Low-Med
wizardlm-2-8x2 2	USA	2024-04	Meningitis (presumed bacterial)	G00.9	Low-Med
grok-2-1212	USA	2024-12	West Nile Virus Encephalitis	A92.3	Low-Med
gemma-2-9b-it	USA	2024-06	Meningitis	G00	Low-Med
gpt-4o	USA	2024-05	Meningitis, likely viral	A87.9	Low-Med
llama-3.2-3b-in	USA	2024-09	Dengue Fever	A09.0	Low-Med
gpt-4o-mini	USA	2024-07	Bacterial Meningitis	G00.9	Low-Med
qwen-2.5-coder -	China	2024-11	Meningitis	G00.0	Medium
claude-3-opus- 2	USA	2024-02	Meningitis, likely bacterial	G00.9	Low-Med
lfm-40b	USA	2024-10	Meningitis	Unknown	Low-Med
gemini-2.5-pro	USA	2024-12	Not specified		Low-Med
gemini-2.5-flas	USA	2024-12	Viral Encephalitis (Arboviral)	A83.9	Low-Med
gemini-2.5-flas	USA	2024-12	Encephalitis	A89	Low-Med

Al Model Bias Analysis

Al model bias analysis is generated during orchestration (Step 2). This comprehensive analysis examines cultural, geographic, and training data biases across the Al models used.

Primary Diagnosis Bias Factors:

- Cultural: Western models showed preference for bacterial causes while Asian models considered more viral etiologies
- Geographic: Led to different primary diagnostic considerations based on model origin
- Training Data: Newer models showed more nuanced understanding of rural healthcare limitations

Alternative Diagnoses Bias:

- Missed: Japanese Encephalitis Western models less familiar with Asian mosquito-borne disea...
- Over-diagnosed: Bacterial Meningitis Western models emphasized bacterial causes requiring immedia...

Bias Mitigation Recommendations:

• Geographic: Include diverse regional disease patterns in differential

Critical Decision Points & Evidence Synthesis

Critical Decision Points

Key areas where models showed significant divergence in diagnostic or management approach:

Evidence Synthesis & Clinical Correlation

Symptom-Diagnosis Correlation Matrix

Symptom/Finding	Viral Mening	Bacterial Me
Rural Setting	+++	+++
Confusion	+++	+++
Recent Mosquito	+++	+++
Fever	+++	++
Photophobia	+++	++

Legend: +++ Strong association, ++ Moderate, + Weak, - Not typical

Diagnostic Decision Tree

Step	Action	If Positive	If Negative
1	CSF Analysis	Start targeted antimicrobial therapy based on results	Consider viral panel and arboviral testing
2	Blood Cultures	Adjust antibiotics per sensitivity	Consider discontinuing antibiotics if viral confirmed

Detailed Model Responses

Complete diagnostic assessments from each model:

1. mistral-7b-inst (France, Released: 2023-09-27)

Primary Diagnosis: Meningitis (ICD-10: A80.x) - Confidence: 0.7

Reasoning: Neck stiffness, photophobia, and disorientation are common symptoms of meningitis....

Differential Diagnoses:

- Encephalitis (ICD: A81.x) 0.3
- Tropical pyrexia of unknown origin (ICD: R50.9) 0.2
- Brain abscess (ICD: G04.9) 0.1

Key Clinical Findings:

- Progressive fever
- Confusion

Recommended Tests:

- Lumbar puncture To diagnose meningitis and identify causative organism
- Complete blood count To assess for leukocytosis
- Blood culture To identify causative organism

Immediate Management:

- Start empiric antibiotics
- Refer for urgent CT scan if transport becomes available

Medications:

- Ceftriaxone 1g IV every 24 hours IV
- Dexamethasone 10mg IV every 6 hours IV

2. grok-4 (USA, Released: 2024-12-26)

Primary Diagnosis: Arboviral Encephalitis (ICD-10: A85.2) - Confidence: 0.75

Reasoning: Progressive fever, headache, confusion, neck stiffness, photophobia, and disorientation in a rural farmer with recent mosquito bites suggest mosquito-borne viral infection like Japanese Encephalitis,

Differential Diagnoses:

- Bacterial Meningitis (ICD: G00.9) 0.6
- Cerebral Malaria (ICD: B50.0) 0.5
- Tuberculous Meningitis (ICD: A17.0) 0.4

Key Clinical Findings:

- Progressive fever and headache with confusion
- Neck stiffness, photophobia, and disorientation

Recent mosquito bites in rural setting

Recommended Tests:

- Lumbar puncture (if feasible in clinic) To evaluate for CSF pleocytosis, protein, glucose to differentiate bacterial vs viral meningitis/encephalitis
- · Blood smear for malaria parasites To rule in/out cerebral malaria given mosquito exposure
- Blood cultures To identify bacterial pathogens if meningitis suspected

Immediate Management:

- Stabilize patient with IV fluids and supportive care
- Initiate empiric treatment and arrange urgent transfer to tertiary center

Medications:

- Ceftriaxone 2g IV
- Acyclovir 10mg/kg IV

3. gpt-oss-120b (USA, Released: 2025-08-05)

Primary Diagnosis: Acute bacterial meningitis (ICD-10: G00.9) - Confidence: 0.72

Reasoning: Fever, neck stiffness, photophobia, altered mental status in a previously healthy adult are classic for meningitis. In a resource limited setting, bacterial meningitis must be prioritized because of h...

Differential Diagnoses:

- Viral encephalitis (e.g., West Nile virus) (ICD: A92.3) 0.18
- Cerebral malaria (ICD: B50.9) 0.07
- Tuberculous meningitis (ICD: A17.0) 0.03

Key Clinical Findings:

- Fever for 5 days
- Neck stiffness with photophobia and confusion

Recommended Tests:

- Lumbar puncture with CSF analysis (cell count, glucose, protein, Gram stain, culture) Rule in bacterial meningitis vs. viral or other causes
- Rapid malaria antigen test (RDT) Exclude cerebral malaria as an alternative etiology
- Blood cultures (2 sets) Identify bloodstream pathogen for targeted antibiotics

Immediate Management:

- Begin empiric intravenous antibiotics: ceftriaxone 2 g IV q24h plus vancomycin 15 mg/kg IV q12h
- Administer dexamethasone 10 mg IV before or with first antibiotic dose
- Obtain blood cultures and rapid malaria test before antibiotics if possible

Medications:

- Ceftriaxone 2 g IV once daily IV
- Vancomycin 15 mg/kg IV every 12 hours (adjust for renal function) IV

4. command-r (Canada, Released: 2024-03-11)

Primary Diagnosis: Meningeal Inflammation (ICD-10: G03) - Confidence: 0.85

Reasoning: Fever, headache, neck stiffness, and confusion are typical symptoms. Mosquito bites could be a potential exposure risk for arboviral infections, which are linked to meningoencephalitis....

Differential Diagnoses:

Bacterial Meningitis (ICD: A39) - 0.7
Viral Encephalitis (ICD: A83.9) - 0.6
Brain Abscess (ICD: G05) - 0.4

Key Clinical Findings:

- Mosquito bites exposure
- Neurological symptoms: fever, headache, neck stiffness, photophobia, disorientation

Recommended Tests:

- Lumbar Puncture Differentiate between meningitis and encephalitis, rule out other causes of increased intracranial pressure
- Complete Blood Count Assess for infection and inflammation markers
- Chemical Profile Basic metabolic panel to check for any organ dysfunction

Immediate Management:

- Initiate empiric antibiotic therapy for meningitis coverage
- Start IV fluids for supportive care

Medications:

- Ceftriaxone 1g daily IV
- Acetaminophen 1g every 6 hours PO

5. deepseek-chat (China, Released: 2024-12-20)

Primary Diagnosis: Meningoencephalitis (likely viral or mosquito-borne) (ICD-10: G04.90) - Confidence: 0.85

Reasoning: Fever, headache, confusion, neck stiffness, and photophobia are classic signs of meningoencephalitis. Recent mosquito bites suggest a possible mosquito-borne etiology (e.g., West Nile virus, Japanese ...

Differential Diagnoses:

- Bacterial meningitis (ICD: G00.9) 0.7
- Cerebral malaria (ICD: B50.0) 0.6
- Tuberculous meningitis (ICD: A17.0) 0.4

Key Clinical Findings:

- Progressive fever and headache
- Neck stiffness, photophobia, and disorientation

Recommended Tests:

- Lumbar puncture (CSF analysis) Rule in/out meningitis or encephalitis
- Malaria rapid diagnostic test (RDT) Rule out cerebral malaria
- Complete blood count (CBC) Assess for infection or inflammation

Immediate Management:

- Initiate empiric treatment for bacterial meningitis and viral encephalitis
- Arrange urgent transport to tertiary care center for advanced diagnostics and treatment

Medications:

- Ceftriaxone 2g IV
- Acyclovir 10mg/kg IV

6. deepseek-r1 (China, Released: 2025-01-20)

7. sonar-deep-rese (USA, Released: 2025-03-07)

Primary Diagnosis: Acute bacterial meningitis (ICD-10: G00.9) - Confidence: 0.75

Reasoning: Classic triad of fever, neck stiffness, and altered mental status with photophobia in older adult requires immediate empiric treatment...

Differential Diagnoses:

- West Nile virus encephalitis (ICD: A92.31) 0.7
- Eastern equine encephalitis (ICD: A83.2) 0.4
- Meningococcal meningitis (ICD: A39.0) 0.45

Key Clinical Findings:

- Classic meningitis triad: fever, neck stiffness, altered mental status
- Photophobia indicating meningeal irritation
- Recent mosquito bite exposure in rural setting
- Progressive 5-day course with neurological deterioration

Recommended Tests:

- Lumbar puncture with CSF analysis Differentiate bacterial from viral meningitis, identify organism
- Blood cultures Identify causative organism in bacterial meningitis
- Complete blood count with differential Assess for bacterial infection and complications

Immediate Management:

- Initiate empiric antibiotic therapy immediately
- Start dexamethasone before or with first antibiotic dose
- Establish IV access and ensure adequate hydration

Medications:

- Ceftriaxone 2g IV
- Ampicillin 2g IV

8. jamba-large-1.7 (Israel, Released: 2025-07-01)

Primary Diagnosis: Meningitis (ICD-10: G00.9) - Confidence: 0.85

Reasoning: Given the fever, headache, neck stiffness, and confusion, meningitis is the most likely diagnosis....

Differential Diagnoses:

• Encephalitis (ICD: A83.9) - 0.65

• Malaria (ICD: B54) - 0.55

• Cerebral Malaria (ICD: B50.0) - 0.45

Key Clinical Findings:

- Neck stiffness
- Photophobia

Recommended Tests:

- Cerebrospinal fluid (CSF) analysis To confirm meningitis and rule out other causes of the symptoms
- Malaria blood smear To rule out malaria
- Brain imaging (if transport is possible) To check for signs of encephalitis or cerebral malaria

Immediate Management:

- Start IV antibiotics (broad-spectrum) for meningitis
- Prepare for transport to a tertiary care center

Medications:

• Ceftriaxone 2 grams IV every 12 hours IV

9. mistral-large-2 (France, Released: 2024-11-01)

Primary Diagnosis: Meningoencephalitis (ICD-10: G04.90) - Confidence: 0.9

Reasoning: Progressive fever, headache, confusion, neck stiffness, photophobia, and disorientation strongly suggest meningoencephalitis, especially with recent mosquito bites indicating a potential viral etiolog...

Differential Diagnoses:

- Bacterial meningitis (ICD: G00.9) 0.7
- Encephalitis due to other causes (ICD: G05.90) 0.6
- Cerebral malaria (ICD: B50.0) 0.5

Key Clinical Findings:

- Progressive fever
- Headache
- Confusion
- Neck stiffness

Recommended Tests:

- Lumbar puncture with CSF analysis To confirm meningoencephalitis and identify the causative agent
- Blood cultures To rule out bacterial infection
- Malaria smear To rule out cerebral malaria

Immediate Management:

Initiate empiric antibiotics and antivirals

Arrange immediate transfer to tertiary center

Medications:

- Ceftriaxone 2g IV every 12 hours IV
- Acyclovir 10mg/kg IV every 8 hours IV

10. command-r-plus (Canada, Released: 2024-04-04)

Primary Diagnosis: Encephalitis (ICD-10: G04.90) - Confidence: 0.7

Reasoning: The patient's fever, headache, neck stiffness, photophobia, and confusion are indicative of encephalitis, especially with the history of recent mosquito bites. This is a life-threatening condition tha...

Differential Diagnoses:

- Meningitis (ICD: G03.9) 0.6
- West Nile Virus Encephalitis (ICD: A83.3) 0.5
- Brain Abscess (ICD: G06) 0.4

Key Clinical Findings:

- Fever
- Headache
- Neck stiffness
- Photophobia

Recommended Tests:

- Lumbar Puncture (Spinal Tap) Rule in/out meningitis, encephalitis, or other central nervous system infections
- Complete Blood Count (CBC) with Differential Evaluate for infection and inflammation
- Serology for Arboviruses (including West Nile Virus) Detect evidence of recent viral infection transmitted by mosquitoes

Immediate Management:

- · Administer intravenous fluids
- Start empiric antibiotic therapy for broad coverage of bacterial meningitis while awaiting lumbar puncture results

Medications:

- Ceftriaxone 2 grams intravenously every 12 hours IV
- Acetaminophen 325-650 mg orally every 4-6 hours as needed PO

11. wizardlm-2-8x22 (USA, Released: 2024-04-15)

Primary Diagnosis: Meningitis (presumed bacterial) (ICD-10: G00.9) - Confidence: 0.7

Reasoning: Progressive fever, headache, confusion, neck stiffness, and photophobia suggest meningitis; mosquito bites increase suspicion for vector-borne bacterial meningitis....

Differential Diagnoses:

• Viral meningitis (ICD: A87) - 0.2

• Encephalitis (ICD: G04.9) - 0.1

• Malaria (ICD: B54.9) - 0.05

Key Clinical Findings:

- Progressive fever
- Headache
- Confusion
- Neck stiffness

Recommended Tests:

- Lumbar puncture (if possible) To analyze cerebrospinal fluid for cell count, protein, glucose, Gram stain, and culture to identify the causative organism
- · Blood cultures To identify bacteremia which may suggest the causative agent of meningitis
- Rapid diagnostic test for malaria To rule out malaria in a febrile patient from a mosquito-endemic area

Immediate Management:

- Initiate empirical antibiotic therapy
- · Administer antipyretics for fever

Medications:

- Ceftriaxone 2 g every 12 hours IV
- Dexamethasone 0.15 mg/kg every 6 hours IV

12. grok-2-1212 (USA, Released: 2024-12-12)

Primary Diagnosis: West Nile Virus Encephalitis (ICD-10: A92.3) - Confidence: 0.8

Reasoning: History of mosquito bites, fever, headache, and neurological symptoms suggestive of encephalitis. West Nile Virus is prevalent in rural areas where mosquito exposure is common....

Differential Diagnoses:

- Bacterial Meningitis (ICD: G00.9) 0.6
- Viral Encephalitis (non-West Nile) (ICD: A86) 0.5
- Heat Stroke (ICD: T67.5) 0.3

Key Clinical Findings:

- Neck stiffness
- Photophobia
- Disorientation

Recommended Tests:

- Lumbar Puncture To rule in/out bacterial meningitis and confirm encephalitis
- Blood Cultures To identify bacterial infection
- Serology for West Nile Virus To confirm West Nile Virus infection

Immediate Management:

- Initiate IV fluids for hydration
- Stabilize patient's vital signs

Medications:

- Ceftriaxone 2g IV
- Acetaminophen 1g PO

13. gemma-2-9b-it (USA, Released: 2024-06-27)

Primary Diagnosis: Meningitis (ICD-10: G00) - Confidence: 0.7

Reasoning: Fever, headache, confusion, neck stiffness, and photophobia are highly suggestive of meningitis. The history of recent mosquito bites raises concern for viral meningitis, but bacterial meningitis cann...

Differential Diagnoses:

Encephalitis (ICD: G04) - 0.2
Brain Abscess (ICD: I67) - 0.1
Severe Migraine (ICD: G43) - 0.05

Key Clinical Findings:

- Fever
- Neck stiffness
- Photophobia
- Confusion

Recommended Tests:

Lumbar puncture - To analyze cerebrospinal fluid for infection, inflammation, and other abnormalities

Immediate Management:

- Administer intravenous fluids
- Initiate empiric antibiotics for suspected bacterial meningitis

Medications:

• Ceftriaxone 2 g IV every 12 hours IV

14. gpt-4o (USA, Released: 2024-05-13)

Primary Diagnosis: Meningitis, likely viral (ICD-10: A87.9) - Confidence: 0.7

Reasoning: The patient's symptoms of fever, headache, neck stiffness, photophobia, and confusion are consistent with meningitis. Recent mosquito bites raise suspicion for viral causes such as West Nile virus....

Differential Diagnoses:

• Bacterial meningitis (ICD: G00.9) - 0.5

• Encephalitis (ICD: G04.90) - 0.4

• Sepsis (ICD: A41.9) - 0.3

Key Clinical Findings:

- Neck stiffness
- Photophobia

Recommended Tests:

- Lumbar puncture To analyze cerebrospinal fluid for meningitis
- Blood cultures To identify any bacterial infection
- Complete blood count (CBC) To assess for infection or systemic inflammation

Immediate Management:

- Initiate empiric IV antibiotics
- · Start IV fluids for hydration

Medications:

Ceftriaxone 2g IV

15. Ilama-3.2-3b-in (USA, Released: 2024-09-25)

Primary Diagnosis: Dengue Fever (ICD-10: A09.0) - Confidence: 0.8

Reasoning: Symptoms of fever, headache, and confusion, combined with mosquito bites and lack of access to advanced labs, are consistent with dengue fever. The rural setting and limited resources further support ...

Differential Diagnoses:

- Malaria (ICD: P560-P565) 0.2
- Typhoid Fever (ICD: A01-A09) 0.1
- Rabies (ICD: A03) 0.05

Key Clinical Findings:

- · Neck stiffness and photophobia
- Disorientation and confusion

Recommended Tests:

- Complete Blood Count (CBC) To rule in/out anemia and infection
- Urinalysis To rule in/out urinary tract infection
- Imaging (CT or MRI) of the brain To rule in/out cerebral edema or hemorrhage

Immediate Management:

- Administer intravenous fluids and antibiotics
- Transport patient to tertiary center

Medications:

- Paracetamol 1g IV q 4h IV
- Ceftriaxone 2.25g IV q 24h IV

16. gpt-4o-mini (USA, Released: 2024-07-18)

Primary Diagnosis: Bacterial Meningitis (ICD-10: G00.9) - Confidence: 0.85

Reasoning: The patient presents with fever, neck stiffness, photophobia, and confusion, which are classic signs of meningitis. The recent mosquito bites raise suspicion for vector-borne infections that could lea...

Differential Diagnoses:

- Viral Meningitis (ICD: A87.9) 0.1
- Encephalitis (ICD: G05.9) 0.05
- Malaria (ICD: B50.9) 0.05

Key Clinical Findings:

- Neck stiffness
- Photophobia

Recommended Tests:

• Lumbar puncture - To analyze cerebrospinal fluid for signs of infection

Immediate Management:

- Initiate intravenous fluids
- Start empirical antibiotic therapy

Medications:

- Ceftriaxone 2 g IV
- Vancomycin 1 g IV

17. qwen-2.5-coder- (China, Released: 2024-11-12)

Primary Diagnosis: Meningitis (ICD-10: G00.0) - Confidence: 0.8

Reasoning: The patient's symptoms of fever, headache, confusion, neck stiffness, and photophobia are classic signs of meningitis. The recent mosquito bites could suggest a possible cause like Japanese encephalit...

Differential Diagnoses:

- Japanese Encephalitis (ICD: A93.0) 0.7
- Bacterial Meningitis (ICD: G01.0) 0.6
- Viral Meningitis (ICD: G04.0) 0.4

Key Clinical Findings:

- Neck stiffness
- Photophobia

Recommended Tests:

- Lumbar Puncture To obtain cerebrospinal fluid (CSF) for analysis, including cell count, protein level, glucose level, and CSF cultures.
- Complete Blood Count (CBC) To check for signs of infection or inflammation.
- Electrolytes To ensure proper hydration and electrolyte balance, especially important in patients with fever and confusion.

Immediate Management:

- Initiate broad-spectrum antibiotics (assuming bacterial meningitis possibility)
- Ensure adequate hydration and electrolyte balance

Medications:

- Ceftriaxone 2g IV
- Acetaminophen 1000mg PO

18. claude-3-opus-2 (USA, Released: 2024-02-29)

Primary Diagnosis: Meningitis, likely bacterial (ICD-10: G00.9) - Confidence: 0.8

Reasoning: Fever, headache, confusion, neck stiffness, and photophobia in the setting of recent mosquito bites suggest meningitis, with bacterial etiology most likely given rapid progression....

Differential Diagnoses:

- Viral meningitis (ICD: A87.9) 0.4
- Cerebral malaria (ICD: B50.0) 0.3
- Encephalitis (ICD: G04.90) 0.2

Key Clinical Findings:

- Fever and headache for 5 days with rapid progression
- Confusion, neck stiffness, and photophobia on exam

Recommended Tests:

- Lumbar puncture Evaluate for meningitis and distinguish bacterial vs. viral etiology
- Blood cultures Identify causative organism if bacterial meningitis
- Rapid malaria test Rule out cerebral malaria given mosquito exposure

Immediate Management:

- Stabilize airway, breathing, circulation
- Empiric broad-spectrum IV antibiotics

Medications:

- Ceftriaxone 2g IV IV
- Vancomycin 15-20mg/kg IV IV

19. lfm-40b (USA, Released: 2024-10-29)

Primary Diagnosis: Meningitis (ICD-10: Unknown) - Confidence: 0.9

Reasoning: Given the patient's symptoms of fever, headache, confusion, neck stiffness, and photophobia, combined with the recent mosquito bites and remote location suggesting potential exposure to infectious dis...

Differential Diagnoses:

- Encephalitis (ICD: Unknown) 0.7
- West Nile virus infection (ICD: A92.4) 0.5
- Community-acquired pneumonia (ICD: J13) 0.3

Key Clinical Findings:

- Neck stiffness
- Photophobia
- Recent mosquito bites

Recommended Tests:

- Lumbar puncture To rule in/out meningitis
- Complete blood count (CBC) To assess for infection
- Blood cultures To identify potential bacterial pathogens

Immediate Management:

- Administer empiric antibiotics for meningitis
- Start antipyretics and analgesics

Medications:

- Ceftriaxone 2 grams IV every 12 hours IV
- Dexamethasone 0.15 mg/kg IV every 6 hours IV

20. gemini-2.5-pro (USA, Released: 2024-12-19)

21. gemini-2.5-flas (USA, Released: 2024-12-19)

Primary Diagnosis: Viral Encephalitis (Arboviral) (ICD-10: A83.9) - Confidence: 0.85

Reasoning: Progressive fever, headache, confusion, neck stiffness, photophobia, and disorientation in a patient with recent mosquito bites strongly suggest arboviral encephalitis, especially given the rural sett...

Differential Diagnoses:

- Bacterial Meningitis (ICD: G00.9) 0.8
- Cerebral Malaria (ICD: B50.0) 0.6
- Tick-borne Encephalitis (ICD: A84.9) 0.5

Key Clinical Findings:

- Progressive fever, headache, confusion (5-day history)
- Recent multiple mosquito bites
- Neck stiffness, photophobia, disorientation on physical exam
- Lack of local advanced diagnostic capabilities (CT, labs)

Recommended Tests:

- Lumbar Puncture (LP) with CSF analysis To differentiate between bacterial and viral meningitis/encephalitis (cell count, protein, glucose, Gram stain, culture, viral PCR). Essential for diagnosis.
- Blood cultures To identify systemic bacterial infection or bacteremia that could spread to CNS.
- Complete Blood Count (CBC) with differential To assess for signs of infection (leukocytosis, leukopenia), anemia, thrombocytopenia.

Immediate Management:

- STAT transfer to tertiary care center for advanced diagnostics (CT, MRI, specialized CSF tests) and definitive management.
- Initiate empiric broad-spectrum antibiotics and antiviral therapy immediately after blood cultures and before LP (if LP is delayed or contraindicated by transfer).
- Secure airway and provide ventilatory support if respiratory compromise or severe altered mental status.

Medications:

- Ceftriaxone 2g IV
- Vancomycin 15mg/kg IV

22. gemini-2.5-flas (USA, Released: 2024-12-19)

Primary Diagnosis: Encephalitis (ICD-10: A89) - Confidence: 0.85

Reasoning: The constellation of fever, headache, confusion, neck stiffness, photophobia, and disorientation in a patient with recent mosquito bites strongly suggests an infectious process affecting the central n...

Differential Diagnoses:

- Meningitis (ICD: G03.9) 0.7
- West Nile Virus Infection (Neuroinvasive Disease) (ICD: A92.3) 0.75
- Tick-borne Encephalitis (ICD: A84.0) 0.3

Key Clinical Findings:

- Progressive fever
- Headache
- Confusion and disorientation
- Neck stiffness

Recommended Tests:

- Lumbar Puncture (CSF analysis) To differentiate between viral and bacterial meningitis/encephalitis, assess for cell count, protein, glucose, Gram stain, and culture.
- Complete Blood Count (CBC) with differential To assess for signs of infection (e.g., elevated white blood cell count).
- Basic Metabolic Panel (BMP) To assess for electrolyte imbalances or renal dysfunction that could contribute to altered mental status.

Immediate Management:

- Initiate empiric antibiotic therapy for suspected bacterial meningitis while awaiting CSF results.
- Administer intravenous fluids for hydration.
- Provide symptomatic relief for headache and fever (e.g., acetaminophen).

Medications:

- Ceftriaxone 2g IV q12h IV
- Vancomycin 15mg/kg IV q8-12h (adjust based on renal function) IV