

CAS 760
Simple Type Theory
Winter 2026

7 Inductive Sets and Types

William M. Farmer

Department of Computing and Software
McMaster University

December 31, 2025



Outline

1. Inductive sets.
2. Inductive types.
3. Inductive type theory extensions.

1. Inductive Sets

Inductive Sets [1/5]

- Let A be a nonempty set of values.
- A **generator for A** is a value g that satisfies one of the following four conditions:
 1. $g \in A$.
 2. $g : B \rightarrow A$ is a total, injective function such that B is a nonempty set of values.
 3. $g : (A \times \cdots \times A) \rightarrow A$ is a total, injective function such that the length of the Cartesian product is $m \geq 1$.
 4. $g : (B \times A \times \cdots \times A) \rightarrow A$ is a total, injective function such that B is a nonempty set of values and the length of the Cartesian product is $m + 1 \geq 2$.
- g is **nullary** if g is the first kind of generator.
- g is **pure** if g is the first or third kind of generator.
- g is **recursive** if g is the third or fourth kind of generator.

Inductive Sets [2/5]

- Let X_g be the formula, in which $p : A \rightarrow \mathbb{B}$ is a predicate, defined as follows:
 1. If g is the first kind of generator, then X_g is $p(g)$.
 2. If g is the second of generator, then X_g is $\forall y \in B . p(g(y))$.
 3. If g is the third kind of generator, then X_g is $\forall x_1 \in A, \dots, x_m \in A . (p(x_1) \wedge \dots \wedge p(x_m)) \Rightarrow p(g(x_1, \dots, x_m))$.
 4. If g is the fourth kind of generator, then X_g is $\forall x_1 \in A, \dots, x_m \in A . (p(x_1) \wedge \dots \wedge p(x_m)) \Rightarrow \forall y \in B . p(g(y, x_1, \dots, x_m))$.

Inductive Sets [3/5]

- Let G be a finite set of generators for A .
- The **set of values generated by G** , written $\text{gen}(G)$, is the set $A' \subseteq A$ defined inductively by the following membership rules:
 1. For each $g \in G$ of the first kind, $g \in A'$.
 2. For each $g \in G$ of the second kind with $g : B \rightarrow A$, if $y \in B$, then $g(y) \in A'$.
 3. For each $g \in G$ of the third kind with $g : (A \times \cdots \times A) \rightarrow A$ such that the length of the Cartesian product is $m \geq 1$, if $x_1, \dots, x_m \in A'$, then $g(x_1, \dots, x_m) \in A'$.
 4. For each $g \in G$ of the fourth kind with $g : (B \times A \times \cdots \times A) \rightarrow A$ such that the length of the Cartesian product is $m + 1 \geq 2$, if $x_1, \dots, x_m \in A'$ and $y \in B$, then $g(y, x_1, \dots, x_m) \in A'$.

Inductive Sets [4/5]

- A is **freely generated by G** if:

1. The sentences

$$g \neq g',$$

$$g \notin \text{ran}(h), \text{ and}$$

$$\text{ran}(h) \cap \text{ran}(h') = \emptyset$$

are true for all distinct nullary $g, g' \in G$ and distinct nonnullary $h, h' \in G$. These sentences, together with the injectivity of the nonnullary generators, ensure that the generators in G produce distinct members of A .

2. The sentence

$$\forall p : A \rightarrow \mathbb{B} . (X_{g_1} \wedge \cdots \wedge X_{g_n}) \Rightarrow \forall x \in A . p(x)$$

is true where $G = \{g_1, \dots, g_n\}$. This sentence is called the **structural induction principle** for A with respect to g_1, \dots, g_n . It ensures $A = \text{gen}(G)$.

Inductive Sets [5/5]

- An **inductive set** is a pair (A, G) where A is a nonempty set and G is a finite set of generators for A such that A is freely generated by G .
- **Theorem (No Junk)**. Let (A, G) be an inductive set. Then $A = \text{gen}(G)$.
- **Theorem (No Confusion)**. Let (A, G) be an inductive set. Then each member of A is generated by G in exactly one way.

Example: Inductive Set of Boolean Values

- Let $\mathbb{B} = \{F, T\}$, the set of Boolean values.
- Then $(\mathbb{B}, \{F, T\})$ is an inductive set.
- The structural induction principle for \mathbb{B} with respect to F, T is
$$\forall p : \mathbb{B} \rightarrow \mathbb{B} . (p(F)) \wedge p(T) \Rightarrow \forall x \in \mathbb{B} . p(x).$$
- $(\mathbb{B}, \{F, T\})$ is an example of an inductive set with an enumerated structure.

Example: Inductive Set of Natural Numbers

- Let $\mathbb{N} = \{0, 1, 2, \dots\}$, the set of natural numbers, and let $S : \mathbb{N} \rightarrow \mathbb{N}$ be the successor function.
- Then $(\mathbb{N}, \{0, S\})$ is an inductive set.
- The structural induction principle for \mathbb{N} with respect to $0, S$ is

$$\forall p : \mathbb{N} \rightarrow \mathbb{B} .$$

$$(p(0) \wedge \forall x_1 \in \mathbb{N} . p(x_1) \Rightarrow p(S(x_1))) \Rightarrow \forall x \in \mathbb{N} . p(x).$$

- $(\mathbb{N}, \{0, S\})$ is the standard example of an inductive set that has a hierarchical structure.

2. Inductive Types

Inductive Types [1/5]

- Let \mathbf{a} be a base type.
- A **constructor for \mathbf{a}** is one of the following:
 1. A constant $\mathbf{c}_{\mathbf{a}}$.
 2. A pair $(\mathbf{c}_{\beta \rightarrow \mathbf{a}}, \mathbf{B}_{\{\beta\}})$ where β does not contain \mathbf{a} and $\mathbf{B}_{\{\beta\}}$ is closed.
 3. A constant \mathbf{c}_{α} where $\alpha = (\mathbf{a} \times \cdots \times \mathbf{a}) \rightarrow \mathbf{a}$ and the length in the extended product type is $m \geq 1$.
 4. A pair $(\mathbf{c}_{\alpha}, \mathbf{B}_{\{\beta\}})$ where $\alpha = (\beta \times \mathbf{a} \times \cdots \times \mathbf{a}) \rightarrow \mathbf{a}$ and the length of the extended product type is $m + 1 \geq 2$, β does not contain \mathbf{a} , and $\mathbf{B}_{\{\beta\}}$ is closed.
- Let c be a constructor for \mathbf{a} .
- Define

$$\bar{c} = \begin{cases} \mathbf{c}_{\alpha} & \text{if } c = \mathbf{c}_{\alpha}; \\ \mathbf{c}_{\alpha} & \text{if } c = (\mathbf{c}_{\alpha}, \mathbf{B}_{\{\beta\}}). \end{cases}$$

- c is **nullary** if c is the first kind of constructor.

Inductive Types [2/5]

- c is **pure** if c is the first or third kind of constructor.
- c is **recursive** if c is the third or fourth kind of constructor.
- Let \mathbf{X}_o^c be the formula of Alonzo defined as follows:

1. If $c = \mathbf{c}_a$ is the first kind of constructor, \mathbf{X}_o^c is
$$(p : \mathbf{a} \rightarrow o) \mathbf{c}_a,$$
2. If $c = (\mathbf{c}_{\beta \rightarrow \mathbf{a}}, \mathbf{B}_{\{\beta\}})$ is the second kind of constructor, then \mathbf{X}_o^c is
$$\forall y : \mathbf{B}_{\{\beta\}} . (p : \mathbf{a} \rightarrow o) (\mathbf{c}_{\beta \rightarrow \mathbf{a}} y).$$
3. If $c = \mathbf{c}_\alpha$ is the third kind of constructor, \mathbf{X}_o^c is
$$\begin{aligned} &\forall x_1, \dots, x_m : \mathbf{a} . \\ &((p : \mathbf{a} \rightarrow o) x_1 \wedge \dots \wedge (p : \mathbf{a} \rightarrow o) x_m) \Rightarrow \\ &(p : \mathbf{a} \rightarrow o) (\mathbf{c}_\alpha (x_1, \dots, x_m)). \end{aligned}$$
4. If $c = (\mathbf{c}_\alpha, \mathbf{B}_{\{\beta\}})$ is the fourth kind of constructor, then \mathbf{X}_o^c is
$$\begin{aligned} &\forall x_1, \dots, x_m : \mathbf{a} . \\ &((p : \mathbf{a} \rightarrow o) x_1 \wedge \dots \wedge (p : \mathbf{a} \rightarrow o) x_m) \Rightarrow \\ &\forall y : \mathbf{B}_{\{\beta\}} . (p : \mathbf{a} \rightarrow o) (\mathbf{c}_\alpha (y, x_1, \dots, x_m)). \end{aligned}$$

Inductive Types [3/5]

- Let C be a finite set of constructors for \mathbf{a} .
- The **set of obligations for C** , written $\text{obl}(C)$, is the set of sentences defined by:
 1. For each $(\mathbf{c}_{\beta \rightarrow \mathbf{a}}, \mathbf{B}_{\{\beta\}}) \in C$, the obligations of C are:
 - a. $\mathbf{B}_{\{\beta\}} \neq \emptyset_{\{\beta\}}$.
 - b. $\mathbf{c}_{\beta \rightarrow \mathbf{a}} \downarrow (\mathbf{B}_{\{\beta\}} \rightarrow U_{\{\mathbf{a}\}})$.
 - c. $\text{TOTAL-ON}(\mathbf{c}_{\beta \rightarrow \mathbf{a}}, \mathbf{B}_{\{\beta\}}, U_{\{\mathbf{a}\}})$.
 - d. $\text{INJ}(\mathbf{c}_{\beta \rightarrow \mathbf{a}})$.
 2. For each $\mathbf{c}_{\alpha} \in C$ where $\alpha = (\mathbf{a} \times \cdots \times \mathbf{a}) \rightarrow \mathbf{a}$, the obligations of C are:
 - a. $\text{TOTAL}(\mathbf{c}_{\alpha})$.
 - b. $\text{INJ}(\mathbf{c}_{\alpha})$.

Inductive Types [4/5]

3. For each $(\mathbf{c}_\alpha, \mathbf{B}_{\{\beta\}}) \in C$ where $\alpha = (\beta \times \mathbf{a} \times \cdots \times \mathbf{a}) \rightarrow \mathbf{a}$, the obligations of C are:
- a. $\mathbf{B}_{\{\beta\}} \neq \emptyset_{\{\beta\}}$.
 - b. $\mathbf{c}_\alpha \downarrow ((\mathbf{B}_{\{\beta\}} \times U_{\{\mathbf{a}\}} \times \cdots \times U_{\{\mathbf{a}\}}) \rightarrow U_{\{\mathbf{a}\}})$.
 - c. $\text{TOTAL-ON}(\mathbf{c}_\alpha, \mathbf{B}_{\{\beta\}} \times U_{\{\mathbf{a}\}} \times \cdots \times U_{\{\mathbf{a}\}}, U_{\{\mathbf{a}\}})$.
 - d. $\text{INJ}(\mathbf{c}_\alpha)$.
4. For all distinct nullary $c, c' \in C$ and distinct nonnullary $d, d' \in C$ where \bar{d} has type $\alpha \rightarrow \mathbf{a}$ and \bar{d}' has type $\alpha' \rightarrow \mathbf{a}$, the obligations of C are:
- a. $c \neq c'$,
 - b. $c \notin (\text{ran}_{(\alpha \rightarrow \mathbf{a}) \rightarrow \{\mathbf{a}\}} \bar{d})$.
 - c. $(\text{ran}_{(\alpha \rightarrow \mathbf{a}) \rightarrow \{\mathbf{a}\}} \bar{d}) \cap (\text{ran}_{(\alpha' \rightarrow \mathbf{a}) \rightarrow \{\mathbf{a}\}} \bar{d}') = \emptyset_{\{\mathbf{a}\}}$.
5. The sentence
- $$\forall p : \mathbf{a} \rightarrow o . (\mathbf{X}_o^{c_1} \wedge \cdots \wedge \mathbf{X}_o^{c_n}) \Rightarrow \forall x : \mathbf{a} . p\,x,$$
- where $C = \{c_1, \dots, c_n\}$, is an obligation of C . This sentence is called the **structural induction principle** for \mathbf{a} with respect to c_1, \dots, c_n .

Inductive Types [5/5]

- Let $T = (L, \Gamma)$ be a theory where $L = (\mathcal{B}, \mathcal{C})$, \mathbf{a} is a base type, and C is a finite set of constructors for \mathbf{a} .
- (\mathbf{a}, C) belongs to T if $\mathbf{a} \in \mathcal{B}$, $\{\bar{c} \mid c \in C\} \subseteq \mathcal{C}$, and $\mathbf{B}_\beta \in \mathcal{E}(L)$ for all $(\mathbf{c}_\alpha, \mathbf{B}_\beta) \in C$.
- \mathbf{a} is freely constructed by C in T if:
 1. (\mathbf{a}, C) belongs to T .
 2. $T \models \mathbf{A}_o$ for all $\mathbf{A}_o \in \text{obl}(C)$.
- An inductive type in T is a pair (\mathbf{a}, C) where \mathbf{a} is a base type and C is a finite set of constructors for \mathbf{a} such that \mathbf{a} is freely constructed by C in T .
- Theorem (Inductive Types Theorem). Let T be a theory and M be a standard model of T . If (\mathbf{a}, C) is an inductive type in T , then $(D_{\mathbf{a}}^M, G)$ is an inductive set where
$$G = \{I^M(c) \mid c \in C \text{ is pure}\} \cup \{\text{gts}(I^M(\bar{c})) \mid c \in C \text{ is nonpure}\}.$$

Example: Theory of Boolean Values

Theory Definition (Boolean Values)

Name: BOOLE.

Base types: B .

Constants: f_B, t_B .

Axioms:

1. $f_B \neq t_B$ (Boolean values are distinct).
2. $\forall b : B . b = f_B \vee b = t_B$. (all truth values are Boolean).

Example: Theory of Natural Numbers

Theory Definition (Peano Arithmetic)

Name: PA.

Base types: N .

Constants: 0_N , $S_{N \rightarrow N}$.

Axioms:

1. $\text{TOTAL}(S)$ (S is total).
2. $\forall x : N . 0 \neq S x$ (0 has no predecessor).
3. $\forall x, y : N . S x = S y \Rightarrow x = y$. (S is injective).
4. $\forall p : N \rightarrow o .$
 $(p 0 \wedge \forall x : N . (p x \Rightarrow p (S x))) \Rightarrow \forall x : N . p x$
(induction principle).

3. Inductive Type Theory Extensions

Inductive Type Theory Extensions [1/2]

- Let $L_i = (\mathcal{B}_i, \mathcal{C}_i)$ be a language and $T_i = (L_i, \Gamma_i)$ be a theory for $i \in \{1, 2\}$.
- Let \mathbf{a} be a base type and C be a finite set of constructors for \mathbf{a} .
- T_2 is an **inductive type theory extension** of T_1 by (\mathbf{a}, C) if:
 1. $\mathcal{B}_2 = \mathcal{B}_1 \cup \{\mathbf{a}\}$ with $\mathbf{a} \notin \mathcal{B}_1$.
 2. $\mathcal{C}_2 = \mathcal{C}_1 \cup \{\bar{c} \mid c \in C\}$ with $\{\bar{c} \mid c \in C\} \cap \mathcal{C}_1 = \emptyset$.
 3. $\mathbf{B}_\beta \in \mathcal{E}(L_1)$ for all $(\mathbf{c}_\alpha, \mathbf{B}_\beta) \in C$.
 4. $\Gamma_2 = \Gamma_1 \cup \text{obl}(C)$.
- **Proposition.** Let T_2 be an inductive type theory extension of T_1 by (\mathbf{a}, C) . Then (\mathbf{a}, C) is an inductive type in T_2 .

Inductive Type Theory Extensions [2/2]

- **Theorem.** Let T_2 be an inductive type theory extension of T_1 by (\mathbf{a}, C) . Then the following statements hold:
 1. If each $c \in C$ is pure, then $T_1 \trianglelefteq_m T_2$.
 2. If $T_1 \models \mathbf{B}_{\{\beta\}} \neq \emptyset_{\{\beta\}}$ for all $(\mathbf{c}_\alpha, \mathbf{B}_{\{\beta\}}) \in C$, then $T_1 \trianglelefteq_m T_2$.
- **Remark.** The proof of this theorem shows that, if (\mathbf{a}, C) is an inductive type, then \mathbf{a} can be interpreted, roughly speaking, as the set of expressions constructed from the members of C .

Inductive Type Theory Extension Module

Inductive Type Theory Extension X.Y

Name: Name.

Extends Name-of-subtheory.

New base type: **a**.

Constructors:

1. c_1 with the form \mathbf{c}_α or $(\mathbf{c}_\alpha, \mathbf{B}_{\{\beta\}})$.

\vdots

p. c_p with the form \mathbf{c}_α or $(\mathbf{c}_\alpha, \mathbf{B}_{\{\beta\}})$.

Example: Ind. Type Thy. Ext. of Boolean Values

Inductive Type Theory Extension (Boolean Values)

Name: BOOLE-ALT.

Extends: MIN.

New Base Type: B .

Constructors:

1. f_B .
2. t_B .

Example: Ind. Type Thy. Ext. of Nat. Numbers

Inductive Type Theory Extension (Natural Numbers)

Name: PA-ALT.

Extends: MIN.

New Base Type: N .

Constructors:

1. 0_N .
2. $S_{N \rightarrow N}$.

The End.