

# Stacks in Mathematical Physics

Kadri İlker Berktav

Department of Mathematics  
Bilkent University

# Outline

1. Field theories and their phase spaces
2. Why higher categorical objects emerge
3. A glimpse to higher structures in physics
  - Stacks in gauge theory
  - Stacks in gravity

# Field Theories and Their Phase Spaces

Some terminology:

# Field Theories and Their Phase Spaces

Some terminology:

- By a **classical field theory**, we mean the data of

$$(M, F_M, \mathcal{S} : F_M \rightarrow \mathbb{C}, G). \quad (1)$$

# Field Theories and Their Phase Spaces

Some terminology:

- By a **classical field theory**, we mean the data of

$$(M, F_M, \mathcal{S} : F_M \rightarrow \mathbb{C}, G). \quad (1)$$

- The **critical locus**  $\text{crit}(\mathcal{S})$  of  $\mathcal{S}$ , modulo symmetries:

$$\text{crit}(\mathcal{S}) = \{p \in F_M : d\mathcal{S}|_p = 0\} / \sim, \quad (2)$$

which is the moduli space of solutions to the **Euler-Lagrange equations (or EOM)**.

# Field Theories and Their Phase Spaces

Some terminology:

- By a **classical field theory**, we mean the data of

$$(M, F_M, \mathcal{S} : F_M \rightarrow \mathbb{C}, G). \quad (1)$$

- The **critical locus**  $\text{crit}(\mathcal{S})$  of  $\mathcal{S}$ , modulo symmetries:

$$\text{crit}(\mathcal{S}) = \{p \in F_M : d\mathcal{S}|_p = 0\} / \sim, \quad (2)$$

which is the moduli space of solutions to the  
**Euler-Lagrange equations (or EOM)**.

- Want to understand EOMs for each open  $U \subset M$  and organize this data coherently...

Classical field theories  $\equiv$  (Sheaves of) Moduli spaces

# Field Theories and Their Phase Spaces

**Examples...**

# Field Theories and Their Phase Spaces

## Example (1)

A classical free particle of mass  $m$  moving in  $\mathbb{R}^n$  together with a potential energy  $V : \mathbb{R}^n \rightarrow \mathbb{R}$ : Let



# Field Theories and Their Phase Spaces

## Example (1)

A classical free particle of mass  $m$  moving in  $\mathbb{R}^n$  together with a potential energy  $V : \mathbb{R}^n \rightarrow \mathbb{R}$ : Let

- $F_M := \text{Maps}(M, \mathbb{R}^n)$  for  $M := [0, 1]$ , and  $G = \text{ISO}(\mathbb{R}^n)$ ;

# Field Theories and Their Phase Spaces

## Example (1)

A classical free particle of mass  $m$  moving in  $\mathbb{R}^n$  together with a potential energy  $V : \mathbb{R}^n \rightarrow \mathbb{R}$ : Let

- $F_M := Maps(M, \mathbb{R}^n)$  for  $M := [0, 1]$ , and  $G = ISO(\mathbb{R}^n)$ ;
- $\mathcal{S}(q) := \int_{[0,1]} \left( \frac{m \|\dot{q}\|^2}{2} - V(q) \right), \quad q : [0, 1] \longrightarrow \mathbb{R}^n.$

# Field Theories and Their Phase Spaces

## Example (1)

A classical free particle of mass  $m$  moving in  $\mathbb{R}^n$  together with a potential energy  $V : \mathbb{R}^n \rightarrow \mathbb{R}$ : Let

- $F_M := Maps(M, \mathbb{R}^n)$  for  $M := [0, 1]$ , and  $G = ISO(\mathbb{R}^n)$ ;
- $\mathcal{S}(q) := \int_{[0,1]} \left( \frac{m \|\dot{q}\|^2}{2} - V(q) \right), \quad q : [0, 1] \longrightarrow \mathbb{R}^n.$
- EOM:  $m\ddot{q} = -grad V(q).$
- $crit(\mathcal{S}) = \{q \in F_M : m\ddot{q} = -grad V(q)\} / \sim.$

# Field Theories and Their Phase Spaces

## Example (2)

Consider the  $SU(2)$ -Chern-Simons gauge theory on a closed, orientable 3-manifold  $M$ :

# Field Theories and Their Phase Spaces

## Example (2)

Consider the  $SU(2)$ -Chern-Simons gauge theory on a closed, orientable 3-manifold  $M$ :

- $F_M := \Omega^1(M) \otimes \mathfrak{su}(2)$  and  $G = \text{Maps}(M, SU(2))$  -gauge transformations-

# Field Theories and Their Phase Spaces

## Example (2)

Consider the  $SU(2)$ -Chern-Simons gauge theory on a closed, orientable 3-manifold  $M$ :

- $F_M := \Omega^1(M) \otimes \mathfrak{su}(2)$  and  $G = \text{Maps}(M, SU(2))$  -*gauge transformations*-
- $\mathcal{S}(A) := \frac{k}{4\pi} \int_M \text{Tr}(A \wedge dA + \frac{2}{3} A \wedge A \wedge A), \quad k \in \mathbb{Z}.$

# Field Theories and Their Phase Spaces

## Example (2)

Consider the  $SU(2)$ -Chern-Simons gauge theory on a closed, orientable 3-manifold  $M$ :

- $F_M := \Omega^1(M) \otimes \mathfrak{su}(2)$  and  $G = \text{Maps}(M, SU(2))$  -gauge transformations-
- $\mathcal{S}(A) := \frac{k}{4\pi} \int_M \text{Tr}(A \wedge dA + \frac{2}{3} A \wedge A \wedge A), \quad k \in \mathbb{Z}.$
- EOM:  $dA + A \wedge A = 0$  ( $F_A = 0$ ).
- $\text{crit}(\mathcal{S}) := \{A \in \Omega^1(M) \otimes \mathfrak{su}(2) : F_A = 0\} / \sim.$

# Field Theories and Their Phase Spaces

## Example (3)

Consider (vacuum,  $\Lambda = 0$ ) Einstein gravity on Lorentzian 3-manifolds, say  $M = \Sigma \times \mathbb{R}$ :



# Field Theories and Their Phase Spaces

## Example (3)

Consider (vacuum,  $\Lambda = 0$ ) Einstein gravity on Lorentzian 3-manifolds, say  $M = \Sigma \times \mathbb{R}$ :

- $F_M := \{\text{Lorentzian metrics on } M\}$  and  $G = \text{Diff}(M)$ .

# Field Theories and Their Phase Spaces

## Example (3)

Consider (vacuum,  $\Lambda = 0$ ) Einstein gravity on Lorentzian 3-manifolds, say  $M = \Sigma \times \mathbb{R}$ :

- $F_M := \{\text{Lorentzian metrics on } M\}$  and  $G = \text{Diff}(M)$ .
- $\mathcal{S}(g) := \int_M dx^3 \sqrt{-\det(g)} R$ .

# Field Theories and Their Phase Spaces

## Example (3)

Consider (vacuum,  $\Lambda = 0$ ) Einstein gravity on Lorentzian 3-manifolds, say  $M = \Sigma \times \mathbb{R}$ :

- $F_M := \{\text{Lorentzian metrics on } M\}$  and  $G = \text{Diff}(M)$ .
- $\mathcal{S}(g) := \int_M dx^3 \sqrt{-\det(g)} R$ .
- EOM:  $R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 0$ .
- $\text{crit}(\mathcal{S}) := \{g \in F_M : R_{\mu\nu} = 0\} / \sim$  (*Ricci-flat*).

## Some issues with $\text{crit}(\mathcal{S})$

We may have some structural problems about  $\text{crit}(\mathcal{S})$  :

## Some issues with $\text{crit}(\mathcal{S})$

We may have some structural problems about  $\text{crit}(\mathcal{S})$  :

- Worst case: It may only have the structure of **a set**.

## Some issues with $\text{crit}(\mathcal{S})$

We may have some structural problems about  $\text{crit}(\mathcal{S})$  :

- Worst case: It may only have the structure of **a set**.
- Best case: A manifold + some extra geometric structures.  
(e.g. symplectic, Poisson)

## Some issues with $\text{crit}(\mathcal{S})$

We may have some structural problems about  $\text{crit}(\mathcal{S})$  :

- Worst case: It may only have the structure of **a set**.
- Best case: A manifold + some extra geometric structures.  
(e.g. symplectic, Poisson)

**Reason:** Given  $(M, F_M, \mathcal{S}, G)$ , we perform two problematic operations to define  $\text{crit}(\mathcal{S})$ :

## Some issues with $\text{crit}(\mathcal{S})$

We may have some structural problems about  $\text{crit}(\mathcal{S})$  :

- Worst case: It may only have the structure of **a set**.
- Best case: A manifold + some extra geometric structures.  
(e.g. symplectic, Poisson)

**Reason:** Given  $(M, F_M, \mathcal{S}, G)$ , we perform two problematic operations to define  $\text{crit}(\mathcal{S})$ :

1.  $\text{crit}(\mathcal{S})$  : the intersection of the graph of  $d\mathcal{S} : F_M \rightarrow T^*F_M$  with the zero section  $\Theta : F_M \hookrightarrow T^*F_M$  inside  $T^*F_M$ .



## Some issues with $\text{crit}(\mathcal{S})$

We may have some structural problems about  $\text{crit}(\mathcal{S})$  :

- Worst case: It may only have the structure of **a set**.
- Best case: A manifold + some extra geometric structures.  
(e.g. symplectic, Poisson)

**Reason:** Given  $(M, F_M, \mathcal{S}, G)$ , we perform two problematic operations to define  $\text{crit}(\mathcal{S})$ :

1.  $\text{crit}(\mathcal{S})$  : the intersection of the graph of  $d\mathcal{S} : F_M \rightarrow T^*F_M$  with the zero section  $\Theta : F_M \hookrightarrow T^*F_M$  inside  $T^*F_M$ .  
i.e. a *fibered product*  $F_M \times_{T^*F_M} F_M$ ,

## Some issues with $\text{crit}(\mathcal{S})$

We may have some structural problems about  $\text{crit}(\mathcal{S})$  :

- Worst case: It may only have the structure of **a set**.
- Best case: A manifold + some extra geometric structures.  
(e.g. symplectic, Poisson)

**Reason:** Given  $(M, F_M, \mathcal{S}, G)$ , we perform two problematic operations to define  $\text{crit}(\mathcal{S})$ :

1.  $\text{crit}(\mathcal{S})$  : the intersection of the graph of  $d\mathcal{S} : F_M \rightarrow T^*F_M$  with the zero section  $\Theta : F_M \hookrightarrow T^*F_M$  inside  $T^*F_M$ .

i.e. a *fibred product*  $F_M \times_{T^*F_M} F_M$ ,

which may not exist in the same category!!

**[Bad intersections]**

## Some issues with $\text{crit}(\mathcal{S})$




Figure:  $\text{crit}(f)$  as the intersection of two Lagrangians in  $T^*X$

1.  $\text{crit}(\mathcal{S})$ , as the *fibered product/pullback* may not exist in the same category!! [**Bad intersections**]

## Some issues with $\text{crit}(\mathcal{S})$




Figure:  $\text{crit}(f)$  as the intersection of two Lagrangians in  $T^*X$

1.  $\text{crit}(\mathcal{S})$ , as the *fibred product/pullback* may not exist in the same category!! [**Bad intersections**]
2.  $\text{crit}(\mathcal{S}) = \text{crit}(\mathcal{S})/G$  may not exist in the same category!! [**Bad quotients**]

## Some issues with $\text{crit}(\mathcal{S})$

**Hidden problem:** Ordinary categories of spaces (manifolds, varieties, schemes,...) are **not** structured enough!!

e.g. existence of pullbacks/pushouts (limits/colimits):

$$\begin{array}{ccc} \text{"}L_1 \times_Y L_2\text{"} & \longrightarrow & L_2 \\ \downarrow & & \downarrow \\ L_1 & \longrightarrow & Y \end{array} \quad \text{and} \quad \begin{array}{ccc} X \times G & \xrightarrow{\text{act}} & X \\ \downarrow \text{proj} & & \downarrow \\ X & \longrightarrow & \text{"}X/G\text{"} \end{array}$$

## Some issues with $\text{crit}(\mathcal{S})$

**Hidden problem:** Ordinary categories of spaces (manifolds, varieties, schemes,...) are **not** structured enough!!

e.g. existence of pullbacks/pushouts (limits/colimits):

$$\begin{array}{ccc} "L_1 \times_Y L_2" & \longrightarrow & L_2 \\ \downarrow & & \downarrow \\ L_1 & \longrightarrow & Y \end{array} \quad \text{and} \quad \begin{array}{ccc} X \times G & \xrightarrow{\text{act}} & X \\ \downarrow \text{proj} & & \downarrow \\ X & \longrightarrow & "X/G" \end{array}$$

Possible solution: Need something **bigger/more than ordinary categories or spaces !!**

## Some issues with $\text{crit}(\mathcal{S})$

**Hidden problem:** Ordinary categories of spaces (manifolds, varieties, schemes,...) are **not** structured enough!!

e.g. existence of pullbacks/pushouts (limits/colimits):

$$\begin{array}{ccc} "L_1 \times_Y L_2" & \longrightarrow & L_2 \\ \downarrow & & \downarrow \\ L_1 & \longrightarrow & Y \end{array} \quad \text{and} \quad \begin{array}{ccc} X \times G & \xrightarrow{\text{act}} & X \\ \downarrow \text{proj} & & \downarrow \\ X & \longrightarrow & "X/G" \end{array}$$

Possible solution: Need something **bigger/more than ordinary categories or spaces !!**

$\rightsquigarrow$  **Higher spaces + Derived geometry**

( $\infty$ -categories of ) Higher Spaces: stacks, higher stacks, derived schemes, derived stacks

# Higher spaces in the context of derived algebraic geometry (DAG)

- $DAG := AG \oplus_{CT} HT \rightsquigarrow$  a homotopical refinement of AG



# Higher spaces in the context of derived algebraic geometry (DAG)

- $DAG := AG \oplus_{CT} HT \rightsquigarrow$  a homotopical refinement of AG
- Higher spaces as *homotopy sheaves*:

$$\begin{array}{ccc} CAlg_{\mathbb{K}} & \xrightarrow{\text{schemas}} & Sets \\ \downarrow & \searrow \text{stacks} & \downarrow \\ & Grpd & \\ \downarrow h. \text{ stacks} & & \downarrow \\ cdga_{\mathbb{K}}^{\leq 0} & \xrightarrow{d. \text{ stacks}} & Grpd_{\infty} \end{array}$$

# Higher spaces in the context of derived algebraic geometry (DAG)

- $DAG := AG \oplus_{CT} HT \rightsquigarrow$  a homotopical refinement of AG
- Higher spaces as *homotopy sheaves*:

$$\begin{array}{ccc} CAlg_{\mathbb{K}} & \xrightarrow{\text{schemas}} & Sets \\ \downarrow & \searrow \text{stacks} & \downarrow \\ & Grpd & \\ \downarrow h. \text{ stacks} & & \downarrow \\ cdga_{\mathbb{K}}^{\leq 0} & \xrightarrow{d. \text{ stacks}} & Grpd_{\infty} \end{array}$$

- RHS encodes higher symmetries/equivalences/relations (more non-trivial higher morphisms are emerged).
- Get an infinite tower of equivalences

## Punchlines:

- DAG essentially provides a new setup to deal with non-generic situations in geometry. E.g. non-transversal intersections and “bad” quotients.

## Punchlines:

- DAG essentially provides a new setup to deal with non-generic situations in geometry. E.g. non-transversal intersections and “bad” quotients.
  1. The intersection of two Lagrangians  $L_1 \times_Y L_2$  is a stack.
  2. The *quotient stack* defined by

$$[X/G] = \operatorname{colim} \left( X \rightrightarrows X \times G \rightrightarrows X \times G \times G \rightrightarrows \cdots \right),$$

where the maps are the action *act* and projection *pr*.

## Punchlines:

- DAG essentially provides a new setup to deal with non-generic situations in geometry. E.g. non-transversal intersections and “bad” quotients.
  1. The intersection of two Lagrangians  $L_1 \times_Y L_2$  is a stack.
  2. The *quotient stack* defined by

$$[X/G] = \operatorname{colim} \left( X \rightrightarrows X \times G \rightrightarrows X \times G \times G \rightrightarrows \cdots \right),$$

where the maps are the action *act* and projection *pr*.

- DAG also offers generalized versions of certain familiar geometric structures and studies their properties.

## Punchlines:

- DAG essentially provides a new setup to deal with non-generic situations in geometry. E.g. non-transversal intersections and “bad” quotients.
  1. The intersection of two Lagrangians  $L_1 \times_Y L_2$  is a stack.
  2. The *quotient stack* defined by

$$[X/G] = \operatorname{colim} \left( X \rightrightarrows X \times G \rightrightarrows X \times G \times G \rightrightarrows \cdots \right),$$

where the maps are the action *act* and projection *pr*.

- DAG also offers generalized versions of certain familiar geometric structures and studies their properties.
  1. Symplectic, Poisson (well-known)
  2. Contact (uncharted area)

# Stacks in physics

1. Stacks and groupoids naturally appear **in gauge theories**:

# Stacks in physics

## 1. Stacks and groupoids naturally appear **in gauge theories**:

- The phase space of gauge fields is modeled as a groupoid (not as a set!).
- The stack of gauge fields is “richer” than the usual orbit space.
- DAG is more suitable for higher gauge theories.



# Stacks in physics

1. Stacks and groupoids naturally appear **in gauge theories**:
  - The phase space of gauge fields is modeled as a groupoid (not as a set!).
  - The stack of gauge fields is “richer” than the usual orbit space.
  - DAG is more suitable for higher gauge theories.
2. *Our work*: Stacks **in Einstein gravity**. We construct

# Stacks in physics

## 1. Stacks and groupoids naturally appear **in gauge theories**:

- The phase space of gauge fields is modeled as a groupoid (not as a set!).
- The stack of gauge fields is “richer” than the usual orbit space.
- DAG is more suitable for higher gauge theories.

## 2. *Our work*: Stacks **in Einstein gravity**. We construct

- the stacks of Ricci-flat metrics on a fixed manifold and on families of manifolds;
- a gauge theoretical stack of 3D gravity; and
- a natural transformation between the two stacks arising from the different models of 3D gravity.

# Stacks in physics

## 1. Stacks and groupoids naturally appear **in gauge theories**:

- The phase space of gauge fields is modeled as a groupoid (not as a set!).
- The stack of gauge fields is “richer” than the usual orbit space.
- DAG is more suitable for higher gauge theories.

## 2. *Our work*: Stacks **in Einstein gravity**. We construct

- the stacks of Ricci-flat metrics on a fixed manifold and on families of manifolds;
- a gauge theoretical stack of 3D gravity; and
- a natural transformation between the two stacks arising from the different models of 3D gravity.

## 3. Why important? New formulations, tools and methods...

Cost: new dictionary, different levels of abstraction, new math...

**Thank you!**