



Genesis Biotech Inc.

website: <http://www.genesisbio.com.tw> e-mail : info@genesisbio.com.tw

Anti-M1 protein of Influenza Virus A, Rabbit-Monoclonal Antibody

Catalog No. GB-60087 **Quantity:** 1.2 ml **Antigen species:** Influenza virus A
Reactivity: human, swine, and chicken Influenza virus A **Applications tested:** ELISA
Host species: rabbit **Type:** monoclonal antibody **Form:** hybridoma culture supernatant

Target description

The M1 protein of influenza A virus has multiple regulatory functions during the infectious cycle, which include mediation of nuclear export of viral ribonucleoproteins, inhibition of viral transcription and a crucial role in virus assembly and budding. The filamentous phenotype was lost when the amino acid at position 41 was switched from A to V. This observation suggests that an interaction among these regions of M1 may occur during assembly.

Antigen

This monoclonal antibody was raised by immunizing rabbit with synthetic peptide mixture containing amino acids on the N-terminal domain (4-54) of matrix protein (M1) of Influenza virus A. The antigen contained the epitope defined by Epitoscree™ (Genesis Biotech Inc.).

Application

The antibody specificity was assayed by ELISA with the synthetic peptide of matrix protein (M1) of Influenza virus A. It has not been tested in the other applications. However, for the first testing, we recommend 1/50 dilution for ELISA, 1/50 dilution for Western blot analysis (WB) of recombinant protein, 1/10 dilution for immunohistochemistry (IHC) staining on frozen cryosections or paraffin embedded sections.

Related Products

1. Anti-Matrix protein M1 of Influenza virus A rabbit pAb (GB- 10083M)
2. Anti-Matrix protein M1 of Influenza virus A rabbit mAb (GB- 60083)

Ab dilution	prebleed	mAb culture medium
1:1	0.045	1.421
1:10	0.042	0.582
1:100	0.041	0.066
Titer		~50

ELISA Procedure

Antigen is coated on EIA strips at 1µg per well. Add 200µl of blocking buffer and then wash wells with PBST buffer. Monoclonal antibody in culture medium GB-60087 is diluted in series as 10^0 ~ 10^2 folds and added in separate wells. Incubate antibody for 1hr. Wash unbound antibodies and add anti-rabbit IgG-HRP conjugate. Wash the plates and add substrate to develop color for 5 min. Read absorbance (ABS) at 650 nm. Amount of color is directly proportional to the amount of antibodies. Antibody is positive at >2 folds of ABS of control/Pre-Immune serum.

Storage

It is supplied as culture medium of monoclonal antibody. Store at 4°C for short term application. For long-term storage, aliquot and store at -20°C.

References

1. Elleman CJ, Barclay WS. The M1 matrix protein controls the filamentous phenotype of influenza A virus. *Virology*. 2004 Mar 30;321(1):144-53.
2. Reinhardt J, Wolff T. The influenza A virus M1 protein interacts with the cellular receptor of activated C kinase (RACK) 1 and can be phosphorylated by protein kinase C. *Vet Microbiol*. 2000 May 22;74(1-2):87-100.
3. Roberts PC, Lamb RA, Compans RW. The M1 and M2 proteins of influenza A virus are important determinants in filamentous particle formation. *Virology*. 1998 Jan 5;240(1):127-37.

FOR RESEARCH USE ONLY AND NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE