



Anti-Growth Hormone, Chicken-Polyclonal Antibody

Catalog No. PY-10103

Antigen species: Human

Host species: Chicken

Quantity: 100 μ g

Reactivity: Human

Form: Antigen affinity purified Antibody

Applications tested: ELISA, Western Blot

Target description

Single chain polypeptide hormone essential for growth of all tissues. Increases fat mobilization. M.W: 21,700 Human serum protein.

Antigen


This polyclonal antibody was raised by immunizing chicken with Full Length recombinant protein.

Application

ELISA, Western blotting, tissue or cell immunostaining. Recommend concentration for ELISA is 0.1 to 1 μ g/ml. Optimal working dilutions must be determined by the end user.

Related Products

1. Anti-Insulin, pAb (PY-10104)
2. Anti-Albumin, pAb(PY-10105)
3. Anti-CRP, pAb (PY-10106)
4. Anti-IgE, pAb (PY-10111)
5. Anti-Fibronectin, pAb (PY-10112)



ELISA Standard Curve:

Recombinant protein as test antigen. Affinity-Purified anti-hGH IgY as 1st antibody (0.1 μ g/mL), and Rabbit anti-IgY-HRP as 2nd antibody.



Western blot Protocol

1. Block membrane with 5% non-fat milk in PBS-T for 1 hour at room temperature or longer at 4°C.
2. Incubate membrane with IgY antibodies at dilution of 1: 2,000 with 1% milk in PBS-T at R.T. for 1 h.
3. Rinse 3 times with PBS-T, then wash membrane with PBS-T, 5 min each, total of 3 times.
4. Incubate with 2nd antibody (goat-anti-IgY/Fc-HRP) at dilution 1:1,000 for ECL (with 1% milk PBS-T) at R.T. for 1 h.
5. Rinse 3 times with PBS-T, then wash with PBS-T, 5 min each with shaking, total of 3 times.
6. Perform ECL detection of signal using Pierce ECL kit.

Storage

It is supplied as antigen affinity purified antibody in lyophilized powder. Redissolve the powder with 100 microliter sterile water will restore to the original concentration 1mg/ml (1 $\times$ PBS). Store at 4°C for short-term application. For long-term storage, aliquot and store at -20°C.

References

1. Sundstrom, M. *et al.* Crystal structure of an antagonist mutant of human growth hormone, G120R, in complex with its receptor at 2.9 Å resolution. *J. Biol. Chem.* 271 (50), 32197-32203 (1996)

FOR RESEARCH USE ONLY AND NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE