



Genesis Biotech Inc.

website: <http://www.genesisbio.com.tw> e-mail : info@genesisbio.com.tw

Anti-Denature β -Lactoglobulin, Mouse-monoclonal Antibody

Catalog No. PG-20015

Applications tested: WB, ELISA

Type: IgG1

Quantity: 100 μ g

Host species: Mouse

Clone #: 1D8F8

Reactivity: bovine

Form: DEAE purified Ab

Target description

Using dry milk as an immunogen, we have produced monoclonal antibodies (mAb) that can distinguish dry milk from raw milk (Chen et al., 2004). Antigenic mapping shows that the epitope is located at the residues 66-76 in strand D of thermally denatured β -LG (Song et al., 2005). When heated above 80 °C, the epitope region of β -LG is exposed and bound by the mAb (3E7G7). The antibody is therefore defined as "denatured" mAb. On the other hand, the native mAb specifically reacted with the native β -LG, but not heat denatured β -LG.

Antigen


β -lactoglobulin

Application

ELISA, Western blotting

Related Products

Anti-native β -LG mAb, denature mAb	PG-20012
Anti-denature β -LG mAb, denature mAb	PG-20014



Western Blot Procedure

1. Block with 1% Gelatin buffer for 1 hour at RT.
2. Wash with 0.05% PBST 3 X 10 minutes.
3. Add 3000X dilution of monoclonal antibody and Incubate for 1 hour at RT.
4. Wash with 0.05% PBST 3 X 10 minutes.
5. Add appropriate amount of correct secondary antibody, goat anti-mouse antibody conjugated with HRP). Incubate for 1 hour at RT.
6. Wash 3 X 10 minutes with 0.05% PBST at RT.
7. Add HRP substrate and develop

ELISA Procedure

About 0.5 μ g of raw or dry milk protein in 50 μ L of PBS was coated onto a respective microtiter plate. Unbound β -LG was washed and subsequently blocked by an addition of 350 μ L of 1% gelatin (wt/vol) for 30 min. Following washes with PBS, 50 μ L of 3000X dilution of monoclonal antibody were added and incubated at room temperature for 60 min. Each well was then washed 3 x with gelatin-PBS containing 0.1% gelatin and 0.05% Tween-20. Bound β -LG antibodies were detected using a goat anti-mouse IgG conjugated with HRP in gelatin-PBS for 30 min. Finally, each well was washed and developed with 0.04% (wt/vol) 2,2-Azino-bis(3-ethylbenz-thiazoline-6-sulfonic acid) (ABTS) in PBS containing 0.01% H_2O_2 (vol/vol).

Storage

It is supplied as DEAE purified mouse monoclonal antibody (mAb) in lyophilized form. Redissolve the powder with 100 μ L sterile water will restore to the original condition. Store at 4°C for short-term application. For long-term storage, aliquot and store at -20°C.

FOR RESEARCH USE ONLY AND NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE