



Genesis Biotech Inc.

website: <http://www.genesisbio.com.tw> e-mail : info@genesisbio.com.tw

Anti-IgG , Rabbit, Chicken-Polyclonal Antibody

Catalog No. PY-10129

Quantity: 100 µg

Reactivity: Rabbit, chicken

Applications tested: W B

Conjugate: unconjugated

Type: polyclonal antibody

Antigen species: Rabbit

Host species: chicken

Form: Affinity purified Ab

Target description

IgGr : IgG, Rabbit .

IgG is the most abundant class of antibodies found in blood serum and lymph. It is active against bacteria, fungi, viruses, and foreign particles. Immunoglobulin G antibodies trigger action of the complement system.

Antigen


This polyclonal antibody was raised by immunizing chicken with rabbit IgG protein that be derived by *E.coli*

Application

The antibody specificity was assayed by Western blotting with IgG protein. The antibody titer is more than 5000 for Western blot analysis. It has not been tested in the other applications. However, for the first testing, we recommend 1/10000 dilution for ELISA, 1/5000 dilution for WB of recombinant protein, 1/400 dilution for tissue extracts or cell lysates, 1/100 dilution for immunohistochemistry (IHC) staining on frozen cryosections, 1/50 dilution for IHC staining on paraffin embedded sections.

Related Products

1. Anti-IgA , Chicken-Poly (PY-10126)
2. Anti-IgG Fc, Chicken-Poly (PY-10127)
3. Anti-IgG, Mouse, Chicken-Poly (PY-10128)
4. Anti-IgM, Human, Chicken-Poly (PY-10130)
5. Anti-IgE, Chicken-Poly (PY-10111)
6. Anti-Immunoglobulin, light chain κ , Human, Chicken-Poly (PY-10131)



Western blot Procedure

1. Block membrane with 5% non-fat milk in PBS-T for 1 hour at room temperature or longer at 4°C.
2. Incubate membrane with IgY antibodies at dilution of 1: 5,000 with 1% milk in PBS-T @ R.T. for 1 h.
3. Rinse 3 times with PBS-T, then wash membrane with PBS-T, 5 min each, total of 3 times.
4. Incubate with 2nd antibody (goat-anti-IgY/Fc-HRP) at dilution 1:1,000 for ECL (with 1% milk PBS-T) @ R.T. for 1 h.
5. Rinse 3 times with PBS-T, then wash with PBS-T, 5 min each with shaking, total of 3 times.
6. Perform ECL detection of signal using Pierce ECL kit.

Storage

It is supplied as affinity purified antibody/ PBS buffer in lyophilized powder. Redissolve the powder with 100µL sterile water will restore to the original concentration 1mg/mL. Store at 4°C for short-term application. For long-term storage, aliquot and store at -20°C.

References

1. McKelvey EM, Fahey JL. Immunoglobulin changes in disease: quantitation on the basis of heavy polypeptide chains, IgG (gammaG), IgA (gammaA), and IgM (gammaM), and of light polypeptide chains, type K (I) and type L (II). J Clin Invest. 44(11):1778-87(1965).
2. Prahl JW. Enzymic degradation of the Fc fragment of rabbit immunoglobulin IgG. Biochem J. 104(2):647-55(1967).
3. Alepa FP. Antigenic factors characteristic of human immunoglobulin G, detected in the sera of non-human primates. Primates Med. 1(0): 1-9(1968).

FOR RESEARCH USE ONLY AND NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE