



Genesis Biotech Inc.

website: <http://www.genesisbio.com.tw> e-mail : [info@genesisbio.com.tw](mailto:info@genesisbio.com.tw)

## Anti-Adiponectin 1 (a.a.72-245), Rabbit-Polyclonal Antibody

**Catalog No.** PY-10208  
**Antigen species:** Human  
**Conjugate:** unconjugated

**Quantity:** 100 µg  
**Applications tested:** Western, ELISA  
**Type:** polyclonal antibody

**Reactivity:** Human  
**Host species:** Rabbit  
**Form:** Affinity-purified antibody

### Target description

Adiponectin is a adipose tissue-specific protein that has structural homology to collagen VIII and X and complement factor C1q, and that circulates in human plasma at high levels. A reduction in adiponectin expression is associated with insulin resistance in animal models. Administration of adiponectin has been accompanied by a reduction in plasma glucose and an increase in insulin sensitivity.

### Antigen


This polyclonal antibody was raised by immunizing chicken with a human Adiponectin fusion protein (72-245 amino acids).

### Application

The antibody specificity was assayed by Western blotting with the human Adiponectin protein (72-245 amino acids). The antibody titer is more than 3000 for Western blot analysis. It has not been tested in the other applications. However, for the first testing, we recommend 1/4,000 dilution for ELISA, 1/3000 dilution for WB of recombinant protein, 1/200 dilution for tissue extracts or cell lysates, 1/100 dilution for immunohistochemistry (IHC) staining on frozen cryosections, 1/50 dilution for IHC staining on paraffin embedded sections.


### Related Products

1. Anti-C-Reactive Protein, Chicken-Poly (PY-10106)
2. Anti-Plasmin, Chicken-Poly (PY-10107)
3. Anti-Macroglobulin, Chicken-Poly (PY-10108)
4. Anti-Transferrin, Chicken-Poly (PY-10109)
5. Anti-Long EGF, Chicken-Poly (PY-10110)
6. Anti-IgE, Chicken-Poly (PY-10111)
7. Anti-Fibronectin, Chicken-Poly (PY-10112)
8. Anti-Microglobulin, Chicken-Poly (PY-10113)
9. Anti-Antithrombin III, Chicken-Poly (PY-10115)



### Western blot Procedure

1. Block membrane with 5% non-fat milk in PBS-T for 1 hour at room temperature or longer at 4°C.
2. Incubate membrane with IgY antibodies at dilution of 1:3,000 with 1% milk in PBS-T @ R.T. for 1 h.
3. Rinse 3 times with PBS-T, then wash membrane with PBS-T, 5 min each, total of 3 times.
4. Incubate with 2nd antibody (goat-anti-Rb-HRP) at dilution 1:10,000 for ECL (with 1% milk PBS-T) @ R.T. for 1 h.
5. Rinse 3 times with PBS-T, then wash with PBS-T, 5 min each with shaking, total of 3 times.
6. Perform ECL detection of signal using Pierce ECL kit.



### Sandwich ELISA:

ELISA plate is coated with affinity-purified Standard serum as test antigen.  
Affinity-purified IgG as detecting antibody  
Goat anti-Rb-HRP dilution: 1:4000.

TMB method for signal development.

### Storage

It is supplied as affinity purified antibody/ PBS buffer in lyophilized powder. Redissolve the powder with 100µL sterile water will restore to the original concentration 1mg/mL. Store at 4°C for short-term application. For long-term storage, aliquot and store at -20°C.

### References

1. Sasaki, H., Yoshioka, N., Takagi, Y. and Sakaki, Y. Structure of the chromosomal gene for human serum prealbumin. Gene 37 (1-3), 191-197 (1985)

FOR RESEARCH USE ONLY AND NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE