



Genesis Biotech Inc.

website: <http://www.genesisbio.com.tw> e-mail : info@genesisbio.com.tw

Anti-RA-regulated nuclear matrix-associated protein (L2DTL), Chicken-Polyclonal Antibody

Catalog No. PY-10234 **Quantity:** 1mg **Reactivity:** Human
Antigen species: Human **Applications tested:** Western blot, IHC **Host species:** Chicken
Conjugate: unconjugated **Type:** polyclonal antibody **Form:** total IgY

Target description

RA-regulated nuclear matrix-associated protein, L2DTL, CDW1, RAMP, DTL (Human).

Human homologue of *Drosophila* lethal(2) denticleless protein (L2DTL) was found to associate with CUL4 and DDB1 and cell cycle-regulated nuclear and centrosome protein, in aggressive hepatocellular carcinoma. L2DTL also interacts with replication licensing protein CDT1 in vivo. Loss of L2DTL in *Drosophila* S2 and human cells suppressed proteolysis of CDT1 in response to DNA damage.

Antigen

This polyclonal antibody was raised by immunizing chicken with fusion protein (1-231 amino acids).


Application

Western blotting, tissue or cell immunostaining. Recommended starting dilution for Western blot analysis is 1:3000, for tissue or cell staining is 1:150. Optimal working dilutions must be determined by the end user.

References

Pan HW, Chou HY, Liu SH, Peng SY, Liu CL, Hsu HC. *Cell Cycle*. 2006 Nov;5(22):2676-87. Epub 2006 Nov 15.

Western blot



E coli-derived fusion protein as test antigen. Total IgY dilution: 1:5000, Goat anti-IgY-HRP dilution: 1:10000. ECL method for signal development.

Storage

It is supplied as total IgY/ PBS buffer in lyophilized powder. Redissolve the powder with 100μL sterile water will restore to the original concentration 10mg/mL. Store at 4°C for short-term application. For long-term storage, aliquot and store at -20°C.

FOR RESEARCH USE ONLY AND NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE