微積分

習題詳解

Diagnostic tests	0002-0009
Ch 01	0010-0075
Ch 02	0076-0179
Ch 03	0180-0325
Ch 04	0326-0515
Ch 05	0516-0593
Ch 06	0594-0673
Ch 07	0674-0809
Ch 08	0810-0871
Ch 09	0872-0935
Ch 10	0936-1039
Ch 11	1040-1191
Ch 12	1192-1271
Ch 13	1272-1341
Ch 14	1342-1491
Ch 15	1492-1611
Ch 16	1612-1707
Appendixes	1708-1737
Concept check answers	1738-1754

☐ DIAGNOSTIC TESTS

Test A Algebra

1. (a) $(-3)^4 = (-3)(-3)(-3)(-3) = 81$

(b) $-3^4 = -(3)(3)(3)(3) = -81$

(c) $3^{-4} = \dfrac{1}{3^4} = \dfrac{1}{81}$

(d) $\dfrac{5^{23}}{5^{21}} = 5^{23-21} = 5^2 = 25$

(e) $\left(\frac{2}{3}\right)^{-2} = \left(\frac{3}{2}\right)^2 = \frac{9}{4}$

(f) $16^{-3/4} = \dfrac{1}{16^{3/4}} = \dfrac{1}{\left(\sqrt[4]{16}\right)^3} = \dfrac{1}{2^3} = \dfrac{1}{8}$

2. (a) Note that $\sqrt{200} = \sqrt{100 \cdot 2} = 10\sqrt{2}$ and $\sqrt{32} = \sqrt{16 \cdot 2} = 4\sqrt{2}$. Thus $\sqrt{200} - \sqrt{32} = 10\sqrt{2} - 4\sqrt{2} = 6\sqrt{2}$.

(b) $(3a^3b^3)(4ab^2)^2 = 3a^3b^3 16a^2b^4 = 48a^5b^7$

(c) $\left(\dfrac{3x^{3/2}y^3}{x^2y^{-1/2}}\right)^{-2} = \left(\dfrac{x^2y^{-1/2}}{3x^{3/2}y^3}\right)^2 = \dfrac{(x^2y^{-1/2})^2}{(3x^{3/2}y^3)^2} = \dfrac{x^4y^{-1}}{9x^3y^6} = \dfrac{x^4}{9x^3y^6y} = \dfrac{x}{9y^7}$

3. (a) $3(x+6) + 4(2x-5) = 3x + 18 + 8x - 20 = 11x - 2$

(b) $(x+3)(4x-5) = 4x^2 - 5x + 12x - 15 = 4x^2 + 7x - 15$

(c) $\left(\sqrt{a} + \sqrt{b}\right)\left(\sqrt{a} - \sqrt{b}\right) = \left(\sqrt{a}\right)^2 - \sqrt{a}\sqrt{b} + \sqrt{a}\sqrt{b} - \left(\sqrt{b}\right)^2 = a - b$

Or: Use the formula for the difference of two squares to see that $\left(\sqrt{a} + \sqrt{b}\right)\left(\sqrt{a} - \sqrt{b}\right) = \left(\sqrt{a}\right)^2 - \left(\sqrt{b}\right)^2 = a - b$.

(d) $(2x+3)^2 = (2x+3)(2x+3) = 4x^2 + 6x + 6x + 9 = 4x^2 + 12x + 9$.

Note: A quicker way to expand this binomial is to use the formula $(a+b)^2 = a^2 + 2ab + b^2$ with $a = 2x$ and $b = 3$:
$(2x+3)^2 = (2x)^2 + 2(2x)(3) + 3^2 = 4x^2 + 12x + 9$

(e) See Reference Page 1 for the binomial formula $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$. Using it, we get
$(x+2)^3 = x^3 + 3x^2(2) + 3x(2^2) + 2^3 = x^3 + 6x^2 + 12x + 8$.

4. (a) Using the difference of two squares formula, $a^2 - b^2 = (a+b)(a-b)$, we have
$4x^2 - 25 = (2x)^2 - 5^2 = (2x+5)(2x-5)$.

(b) Factoring by trial and error, we get $2x^2 + 5x - 12 = (2x-3)(x+4)$.

(c) Using factoring by grouping and the difference of two squares formula, we have
$x^3 - 3x^2 - 4x + 12 = x^2(x-3) - 4(x-3) = (x^2-4)(x-3) = (x-2)(x+2)(x-3)$.

(d) $x^4 + 27x = x(x^3 + 27) = x(x+3)(x^2 - 3x + 9)$

This last expression was obtained using the sum of two cubes formula, $a^3 + b^3 = (a+b)(a^2 - ab + b^2)$ with $a = x$ and $b = 3$. [See Reference Page 1 in the textbook.]

(e) The smallest exponent on x is $-\frac{1}{2}$, so we will factor out $x^{-1/2}$.
$3x^{3/2} - 9x^{1/2} + 6x^{-1/2} = 3x^{-1/2}(x^2 - 3x + 2) = 3x^{-1/2}(x-1)(x-2)$

(f) $x^3y - 4xy = xy(x^2 - 4) = xy(x-2)(x+2)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. (a) $\dfrac{x^2+3x+2}{x^2-x-2} = \dfrac{(x+1)(x+2)}{(x+1)(x-2)} = \dfrac{x+2}{x-2}$

(b) $\dfrac{2x^2-x-1}{x^2-9} \cdot \dfrac{x+3}{2x+1} = \dfrac{(2x+1)(x-1)}{(x-3)(x+3)} \cdot \dfrac{x+3}{2x+1} = \dfrac{x-1}{x-3}$

(c) $\dfrac{x^2}{x^2-4} - \dfrac{x+1}{x+2} = \dfrac{x^2}{(x-2)(x+2)} - \dfrac{x+1}{x+2} = \dfrac{x^2}{(x-2)(x+2)} - \dfrac{x+1}{x+2} \cdot \dfrac{x-2}{x-2} = \dfrac{x^2-(x+1)(x-2)}{(x-2)(x+2)}$

$$= \frac{x^2-(x^2-x-2)}{(x+2)(x-2)} = \frac{x+2}{(x+2)(x-2)} = \frac{1}{x-2}$$

(d) $\dfrac{\dfrac{y}{x}-\dfrac{x}{y}}{\dfrac{1}{y}-\dfrac{1}{x}} = \dfrac{\dfrac{y}{x}-\dfrac{x}{y}}{\dfrac{1}{y}-\dfrac{1}{x}} \cdot \dfrac{xy}{xy} = \dfrac{y^2-x^2}{x-y} = \dfrac{(y-x)(y+x)}{-(y-x)} = \dfrac{y+x}{-1} = -(x+y)$

6. (a) $\dfrac{\sqrt{10}}{\sqrt{5}-2} = \dfrac{\sqrt{10}}{\sqrt{5}-2} \cdot \dfrac{\sqrt{5}+2}{\sqrt{5}+2} = \dfrac{\sqrt{50}+2\sqrt{10}}{\left(\sqrt{5}\right)^2-2^2} = \dfrac{5\sqrt{2}+2\sqrt{10}}{5-4} = 5\sqrt{2}+2\sqrt{10}$

(b) $\dfrac{\sqrt{4+h}-2}{h} = \dfrac{\sqrt{4+h}-2}{h} \cdot \dfrac{\sqrt{4+h}+2}{\sqrt{4+h}+2} = \dfrac{4+h-4}{h\left(\sqrt{4+h}+2\right)} = \dfrac{h}{h\left(\sqrt{4+h}+2\right)} = \dfrac{1}{\sqrt{4+h}+2}$

7. (a) $x^2+x+1 = \left(x^2+x+\frac{1}{4}\right)+1-\frac{1}{4} = \left(x+\frac{1}{2}\right)^2+\frac{3}{4}$

(b) $2x^2-12x+11 = 2(x^2-6x)+11 = 2(x^2-6x+9-9)+11 = 2(x^2-6x+9)-18+11 = 2(x-3)^2-7$

8. (a) $x+5 = 14-\frac{1}{2}x \quad\Leftrightarrow\quad x+\frac{1}{2}x = 14-5 \quad\Leftrightarrow\quad \frac{3}{2}x = 9 \quad\Leftrightarrow\quad x = \frac{2}{3}\cdot 9 \quad\Leftrightarrow\quad x = 6$

(b) $\dfrac{2x}{x+1} = \dfrac{2x-1}{x} \quad\Rightarrow\quad 2x^2 = (2x-1)(x+1) \quad\Leftrightarrow\quad 2x^2 = 2x^2+x-1 \quad\Leftrightarrow\quad x = 1$

(c) $x^2-x-12 = 0 \quad\Leftrightarrow\quad (x+3)(x-4) = 0 \quad\Leftrightarrow\quad x+3 = 0$ or $x-4 = 0 \quad\Leftrightarrow\quad x = -3$ or $x = 4$

(d) By the quadratic formula, $2x^2+4x+1 = 0 \quad\Leftrightarrow$

$$x = \frac{-4\pm\sqrt{4^2-4(2)(1)}}{2(2)} = \frac{-4\pm\sqrt{8}}{4} = \frac{-4\pm 2\sqrt{2}}{4} = \frac{2\left(-2\pm\sqrt{2}\right)}{4} = \frac{-2\pm\sqrt{2}}{2} = -1\pm\tfrac{1}{2}\sqrt{2}.$$

(e) $x^4-3x^2+2 = 0 \quad\Leftrightarrow\quad (x^2-1)(x^2-2) = 0 \quad\Leftrightarrow\quad x^2-1 = 0$ or $x^2-2 = 0 \quad\Leftrightarrow\quad x^2 = 1$ or $x^2 = 2 \quad\Leftrightarrow$
$x = \pm 1$ or $x = \pm\sqrt{2}$

(f) $3\,|x-4| = 10 \quad\Leftrightarrow\quad |x-4| = \frac{10}{3} \quad\Leftrightarrow\quad x-4 = -\frac{10}{3}$ or $x-4 = \frac{10}{3} \quad\Leftrightarrow\quad x = \frac{2}{3}$ or $x = \frac{22}{3}$

(g) Multiplying through $2x(4-x)^{-1/2}-3\sqrt{4-x} = 0$ by $(4-x)^{1/2}$ gives $2x-3(4-x) = 0 \quad\Leftrightarrow$
$2x-12+3x = 0 \quad\Leftrightarrow\quad 5x-12 = 0 \quad\Leftrightarrow\quad 5x = 12 \quad\Leftrightarrow\quad x = \frac{12}{5}$.

9. (a) $-4 < 5-3x \le 17 \quad\Leftrightarrow\quad -9 < -3x \le 12 \quad\Leftrightarrow\quad 3 > x \ge -4$ or $-4 \le x < 3$.
In interval notation, the answer is $[-4, 3)$.

(b) $x^2 < 2x+8 \quad\Leftrightarrow\quad x^2-2x-8 < 0 \quad\Leftrightarrow\quad (x+2)(x-4) < 0$. Now, $(x+2)(x-4)$ will change sign at the critical values $x = -2$ and $x = 4$. Thus the possible intervals of solution are $(-\infty, -2)$, $(-2, 4)$, and $(4, \infty)$. By choosing a single test value from each interval, we see that $(-2, 4)$ is the only interval that satisfies the inequality.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) The inequality $x(x-1)(x+2) > 0$ has critical values of $-2, 0$, and 1. The corresponding possible intervals of solution are $(-\infty, -2)$, $(-2, 0)$, $(0, 1)$ and $(1, \infty)$. By choosing a single test value from each interval, we see that both intervals $(-2, 0)$ and $(1, \infty)$ satisfy the inequality. Thus, the solution is the union of these two intervals: $(-2, 0) \cup (1, \infty)$.

(d) $|x-4| < 3 \quad \Leftrightarrow \quad -3 < x-4 < 3 \quad \Leftrightarrow \quad 1 < x < 7$. In interval notation, the answer is $(1, 7)$.

(e) $\dfrac{2x-3}{x+1} \le 1 \quad \Leftrightarrow \quad \dfrac{2x-3}{x+1} - 1 \le 0 \quad \Leftrightarrow \quad \dfrac{2x-3}{x+1} - \dfrac{x+1}{x+1} \le 0 \quad \Leftrightarrow \quad \dfrac{2x-3-x-1}{x+1} \le 0 \quad \Leftrightarrow \quad \dfrac{x-4}{x+1} \le 0.$

Now, the expression $\dfrac{x-4}{x+1}$ may change signs at the critical values $x = -1$ and $x = 4$, so the possible intervals of solution are $(-\infty, -1)$, $(-1, 4]$, and $[4, \infty)$. By choosing a single test value from each interval, we see that $(-1, 4]$ is the only interval that satisfies the inequality.

10. (a) False. In order for the statement to be true, it must hold for all real numbers, so, to show that the statement is false, pick $p = 1$ and $q = 2$ and observe that $(1+2)^2 \ne 1^2 + 2^2$. In general, $(p+q)^2 = p^2 + 2pq + q^2$.

(b) True as long as a and b are nonnegative real numbers. To see this, think in terms of the laws of exponents: $\sqrt{ab} = (ab)^{1/2} = a^{1/2}b^{1/2} = \sqrt{a}\,\sqrt{b}$.

(c) False. To see this, let $p = 1$ and $q = 2$, then $\sqrt{1^2 + 2^2} \ne 1 + 2$.

(d) False. To see this, let $T = 1$ and $C = 2$, then $\dfrac{1 + 1(2)}{2} \ne 1 + 1$.

(e) False. To see this, let $x = 2$ and $y = 3$, then $\dfrac{1}{2-3} \ne \dfrac{1}{2} - \dfrac{1}{3}$.

(f) True since $\dfrac{1/x}{a/x - b/x} \cdot \dfrac{x}{x} = \dfrac{1}{a-b}$, as long as $x \ne 0$ and $a - b \ne 0$.

Test B Analytic Geometry

1. (a) Using the point $(2, -5)$ and $m = -3$ in the point-slope equation of a line, $y - y_1 = m(x - x_1)$, we get $y - (-5) = -3(x-2) \quad \Rightarrow \quad y + 5 = -3x + 6 \quad \Rightarrow \quad y = -3x + 1$.

(b) A line parallel to the x-axis must be horizontal and thus have a slope of 0. Since the line passes through the point $(2, -5)$, the y-coordinate of every point on the line is -5, so the equation is $y = -5$.

(c) A line parallel to the y-axis is vertical with undefined slope. So the x-coordinate of every point on the line is 2 and so the equation is $x = 2$.

(d) Note that $2x - 4y = 3 \quad \Rightarrow \quad -4y = -2x + 3 \quad \Rightarrow \quad y = \frac{1}{2}x - \frac{3}{4}$. Thus the slope of the given line is $m = \frac{1}{2}$. Hence, the slope of the line we're looking for is also $\frac{1}{2}$ (since the line we're looking for is required to be parallel to the given line). So the equation of the line is $y - (-5) = \frac{1}{2}(x-2) \quad \Rightarrow \quad y + 5 = \frac{1}{2}x - 1 \quad \Rightarrow \quad y = \frac{1}{2}x - 6$.

2. First we'll find the distance between the two given points in order to obtain the radius, r, of the circle: $r = \sqrt{[3-(-1)]^2 + (-2-4)^2} = \sqrt{4^2 + (-6)^2} = \sqrt{52}$. Next use the standard equation of a circle, $(x-h)^2 + (y-k)^2 = r^2$, where (h, k) is the center, to get $(x+1)^2 + (y-4)^2 = 52$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. We must rewrite the equation in standard form in order to identify the center and radius. Note that $x^2 + y^2 - 6x + 10y + 9 = 0 \quad \Rightarrow \quad x^2 - 6x + 9 + y^2 + 10y = 0$. For the left-hand side of the latter equation, we factor the first three terms and complete the square on the last two terms as follows: $x^2 - 6x + 9 + y^2 + 10y = 0 \quad \Rightarrow$ $(x-3)^2 + y^2 + 10y + 25 = 25 \quad \Rightarrow \quad (x-3)^2 + (y+5)^2 = 25$. Thus, the center of the circle is $(3, -5)$ and the radius is 5.

4. (a) $A(-7, 4)$ and $B(5, -12) \quad \Rightarrow \quad m_{AB} = \dfrac{-12-4}{5-(-7)} = \dfrac{-16}{12} = -\dfrac{4}{3}$

(b) $y - 4 = -\frac{4}{3}[x - (-7)] \quad \Rightarrow \quad y - 4 = -\frac{4}{3}x - \frac{28}{3} \quad \Rightarrow \quad 3y - 12 = -4x - 28 \quad \Rightarrow \quad 4x + 3y + 16 = 0$. Putting $y = 0$, we get $4x + 16 = 0$, so the x-intercept is -4, and substituting 0 for x results in a y-intercept of $-\frac{16}{3}$.

(c) The midpoint is obtained by averaging the corresponding coordinates of both points: $\left(\frac{-7+5}{2}, \frac{4+(-12)}{2}\right) = (-1, -4)$.

(d) $d = \sqrt{[5-(-7)]^2 + (-12-4)^2} = \sqrt{12^2 + (-16)^2} = \sqrt{144 + 256} = \sqrt{400} = 20$

(e) The perpendicular bisector is the line that intersects the line segment $\overline{AB}$ at a right angle through its midpoint. Thus the perpendicular bisector passes through $(-1, -4)$ and has slope $\frac{3}{4}$ [the slope is obtained by taking the negative reciprocal of the answer from part (a)]. So the perpendicular bisector is given by $y + 4 = \frac{3}{4}[x - (-1)]$ or $3x - 4y = 13$.

(f) The center of the required circle is the midpoint of $\overline{AB}$, and the radius is half the length of $\overline{AB}$, which is 10. Thus, the equation is $(x+1)^2 + (y+4)^2 = 100$.

5. (a) Graph the corresponding horizontal lines (given by the equations $y = -1$ and $y = 3$) as solid lines. The inequality $y \geq -1$ describes the points (x, y) that lie on or *above* the line $y = -1$. The inequality $y \leq 3$ describes the points (x, y) that lie on or *below* the line $y = 3$. So the pair of inequalities $-1 \leq y \leq 3$ describes the points that lie on or *between* the lines $y = -1$ and $y = 3$.

(b) Note that the given inequalities can be written as $-4 < x < 4$ and $-2 < y < 2$, respectively. So the region lies between the vertical lines $x = -4$ and $x = 4$ and between the horizontal lines $y = -2$ and $y = 2$. As shown in the graph, the region common to both graphs is a rectangle (minus its edges) centered at the origin.

(c) We first graph $y = 1 - \frac{1}{2}x$ as a dotted line. Since $y < 1 - \frac{1}{2}x$, the points in the region lie *below* this line.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) We first graph the parabola $y = x^2 - 1$ using a solid curve. Since $y \geq x^2 - 1$, the points in the region lie on or *above* the parabola.

(e) We graph the circle $x^2 + y^2 = 4$ using a dotted curve. Since $\sqrt{x^2 + y^2} < 2$, the region consists of points whose distance from the origin is less than 2, that is, the points that lie *inside* the circle.

(f) The equation $9x^2 + 16y^2 = 144$ is an ellipse centered at $(0, 0)$. We put it in standard form by dividing by 144 and get $\dfrac{x^2}{16} + \dfrac{y^2}{9} = 1$. The x-intercepts are located at a distance of $\sqrt{16} = 4$ from the center while the y-intercepts are a distance of $\sqrt{9} = 3$ from the center (see the graph).

Test C Functions

1. (a) Locate -1 on the x-axis and then go down to the point on the graph with an x-coordinate of -1. The corresponding y-coordinate is the value of the function at $x = -1$, which is -2. So, $f(-1) = -2$.

(b) Using the same technique as in part (a), we get $f(2) \approx 2.8$.

(c) Locate 2 on the y-axis and then go left and right to find all points on the graph with a y-coordinate of 2. The corresponding x-coordinates are the x-values we are searching for. So $x = -3$ and $x = 1$.

(d) Using the same technique as in part (c), we get $x \approx -2.5$ and $x \approx 0.3$.

(e) The domain is all the x-values for which the graph exists, and the range is all the y-values for which the graph exists. Thus, the domain is $[-3, 3]$, and the range is $[-2, 3]$.

2. Note that $f(2 + h) = (2 + h)^3$ and $f(2) = 2^3 = 8$. So the difference quotient becomes

$$\frac{f(2+h) - f(2)}{h} = \frac{(2+h)^3 - 8}{h} = \frac{8 + 12h + 6h^2 + h^3 - 8}{h} = \frac{12h + 6h^2 + h^3}{h} = \frac{h(12 + 6h + h^2)}{h} = 12 + 6h + h^2.$$

3. (a) Set the denominator equal to 0 and solve to find restrictions on the domain: $x^2 + x - 2 = 0 \quad \Rightarrow$ $(x - 1)(x + 2) = 0 \quad \Rightarrow \quad x = 1$ or $x = -2$. Thus, the domain is all real numbers except 1 or -2 or, in interval notation, $(-\infty, -2) \cup (-2, 1) \cup (1, \infty)$.

(b) Note that the denominator is always greater than or equal to 1, and the numerator is defined for all real numbers. Thus, the domain is $(-\infty, \infty)$.

(c) Note that the function h is the sum of two root functions. So h is defined on the intersection of the domains of these two root functions. The domain of a square root function is found by setting its radicand greater than or equal to 0. Now,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$4 - x \geq 0 \quad \Rightarrow \quad x \leq 4$ and $x^2 - 1 \geq 0 \quad \Rightarrow \quad (x-1)(x+1) \geq 0 \quad \Rightarrow \quad x \leq -1$ or $x \geq 1$. Thus, the domain of h is $(-\infty, -1] \cup [1, 4]$.

4. (a) Reflect the graph of f about the x-axis.

(b) Stretch the graph of f vertically by a factor of 2, then shift 1 unit downward.

(c) Shift the graph of f right 3 units, then up 2 units.

5. (a) Make a table and then connect the points with a smooth curve:

x	-2	-1	0	1	2
y	-8	-1	0	1	8

(b) Shift the graph from part (a) left 1 unit.

(c) Shift the graph from part (a) right 2 units and up 3 units.

(d) First plot $y = x^2$. Next, to get the graph of $f(x) = 4 - x^2$, reflect f about the x-axis and then shift it upward 4 units.

(e) Make a table and then connect the points with a smooth curve:

x	0	1	4	9
y	0	1	2	3

(f) Stretch the graph from part (e) vertically by a factor of two.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(g) First plot $y = 2^x$. Next, get the graph of $y = -2^x$ by reflecting the graph of $y = 2^x$ about the x-axis.

(h) Note that $y = 1 + x^{-1} = 1 + 1/x$. So first plot $y = 1/x$ and then shift it upward 1 unit.

6. (a) $f(-2) = 1 - (-2)^2 = -3$ and $f(1) = 2(1) + 1 = 3$

(b) For $x \le 0$ plot $f(x) = 1 - x^2$ and, on the same plane, for $x > 0$ plot the graph of $f(x) = 2x + 1$.

7. (a) $(f \circ g)(x) = f(g(x)) = f(2x-3) = (2x-3)^2 + 2(2x-3) - 1 = 4x^2 - 12x + 9 + 4x - 6 - 1 = 4x^2 - 8x + 2$

(b) $(g \circ f)(x) = g(f(x)) = g(x^2 + 2x - 1) = 2(x^2 + 2x - 1) - 3 = 2x^2 + 4x - 2 - 3 = 2x^2 + 4x - 5$

(c) $(g \circ g \circ g)(x) = g(g(g(x))) = g(g(2x-3)) = g(2(2x-3) - 3) = g(4x-9) = 2(4x-9) - 3$

$= 8x - 18 - 3 = 8x - 21$

Test D Trigonometry

1. (a) $300^\circ = 300^\circ\left(\dfrac{\pi}{180^\circ}\right) = \dfrac{300\pi}{180} = \dfrac{5\pi}{3}$ (b) $-18^\circ = -18^\circ\left(\dfrac{\pi}{180^\circ}\right) = -\dfrac{18\pi}{180} = -\dfrac{\pi}{10}$

2. (a) $\dfrac{5\pi}{6} = \dfrac{5\pi}{6}\left(\dfrac{180}{\pi}\right)^\circ = 150^\circ$ (b) $2 = 2\left(\dfrac{180}{\pi}\right)^\circ = \left(\dfrac{360}{\pi}\right)^\circ \approx 114.6^\circ$

3. We will use the arc length formula, $s = r\theta$, where s is arc length, r is the radius of the circle, and θ is the measure of the central angle in radians. First, note that $30^\circ = 30^\circ\left(\dfrac{\pi}{180^\circ}\right) = \dfrac{\pi}{6}$. So $s = (12)\left(\dfrac{\pi}{6}\right) = 2\pi$ cm.

4. (a) $\tan(\pi/3) = \sqrt{3}$ [You can read the value from a right triangle with sides 1, 2, and $\sqrt{3}$.]

(b) Note that $7\pi/6$ can be thought of as an angle in the third quadrant with reference angle $\pi/6$. Thus, $\sin(7\pi/6) = -\frac{1}{2}$, since the sine function is negative in the third quadrant.

(c) Note that $5\pi/3$ can be thought of as an angle in the fourth quadrant with reference angle $\pi/3$. Thus, $\sec(5\pi/3) = \dfrac{1}{\cos(5\pi/3)} = \dfrac{1}{1/2} = 2$, since the cosine function is positive in the fourth quadrant.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $\sin\theta = a/24 \quad\Rightarrow\quad a = 24\sin\theta$ and $\cos\theta = b/24 \quad\Rightarrow\quad b = 24\cos\theta$

6. $\sin x = \frac{1}{3}$ and $\sin^2 x + \cos^2 x = 1 \quad\Rightarrow\quad \cos x = \sqrt{1-\frac{1}{9}} = \dfrac{2\sqrt{2}}{3}$. Also, $\cos y = \frac{4}{5} \quad\Rightarrow\quad \sin y = \sqrt{1-\frac{16}{25}} = \frac{3}{5}$.

So, using the sum identity for the sine, we have

$$\sin(x+y) = \sin x\,\cos y + \cos x\,\sin y = \frac{1}{3}\cdot\frac{4}{5} + \frac{2\sqrt{2}}{3}\cdot\frac{3}{5} = \frac{4+6\sqrt{2}}{15} = \frac{1}{15}\left(4+6\sqrt{2}\right)$$

7. (a) $\tan\theta\,\sin\theta + \cos\theta = \dfrac{\sin\theta}{\cos\theta}\sin\theta + \cos\theta = \dfrac{\sin^2\theta}{\cos\theta} + \dfrac{\cos^2\theta}{\cos\theta} = \dfrac{1}{\cos\theta} = \sec\theta$

(b) $\dfrac{2\tan x}{1+\tan^2 x} = \dfrac{2\sin x/(\cos x)}{\sec^2 x} = 2\,\dfrac{\sin x}{\cos x}\,\cos^2 x = 2\sin x\,\cos x = \sin 2x$

8. $\sin 2x = \sin x \quad\Leftrightarrow\quad 2\sin x\,\cos x = \sin x \quad\Leftrightarrow\quad 2\sin x\,\cos x - \sin x = 0 \quad\Leftrightarrow\quad \sin x\,(2\cos x - 1) = 0 \quad\Leftrightarrow$

$\sin x = 0$ or $\cos x = \frac{1}{2} \quad\Rightarrow\quad x = 0, \frac{\pi}{3}, \pi, \frac{5\pi}{3}, 2\pi$.

9. We first graph $y = \sin 2x$ (by compressing the graph of $\sin x$ by a factor of 2) and then shift it upward 1 unit.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

1 □ FUNCTIONS AND MODELS

1.1 Four Ways to Represent a Function

1. The functions $f(x) = x + \sqrt{2-x}$ and $g(u) = u + \sqrt{2-u}$ give exactly the same output values for every input value, so f and g are equal.

2. $f(x) = \dfrac{x^2 - x}{x-1} = \dfrac{x(x-1)}{x-1} = x$ for $x - 1 \neq 0$, so f and g [where $g(x) = x$] are not equal because $f(1)$ is undefined and $g(1) = 1$.

3. (a) The point $(-2, 2)$ lies on the graph of g, so $g(-2) = 2$. Similarly, $g(0) = -2$, $g(2) = 1$, and $g(3) \approx 2.5$.

(b) Only the point $(-4, 3)$ on the graph has a y-value of 3, so the only value of x for which $g(x) = 3$ is -4.

(c) The function outputs $g(x)$ are never greater than 3, so $g(x) \leq 3$ for the entire domain of the function. Thus, $g(x) \leq 3$ for $-4 \leq x \leq 4$ (or, equivalently, on the interval $[-4, 4]$).

(d) The domain consists of all x-values on the graph of g: $\{x \mid -4 \leq x \leq 4\} = [-4, 4]$. The range of g consists of all the y-values on the graph of g: $\{y \mid -2 \leq y \leq 3\} = [-2, 3]$.

(e) For any $x_1 < x_2$ in the interval $[0, 2]$, we have $g(x_1) < g(x_2)$. [The graph rises from $(0, -2)$ to $(2, 1)$.] Thus, $g(x)$ is increasing on $[0, 2]$.

4. (a) From the graph, we have $f(-4) = -2$ and $g(3) = 4$.

(b) Since $f(-3) = -1$ and $g(-3) = 2$, or by observing that the graph of g is above the graph of f at $x = -3$, $g(-3)$ is larger than $f(-3)$.

(c) The graphs of f and g intersect at $x = -2$ and $x = 2$, so $f(x) = g(x)$ at these two values of x.

(d) The graph of f lies below or on the graph of g for $-4 \leq x \leq -2$ and for $2 \leq x \leq 3$. Thus, the intervals on which $f(x) \leq g(x)$ are $[-4, -2]$ and $[2, 3]$.

(e) $f(x) = -1$ is equivalent to $y = -1$, and the points on the graph of f with y-values of -1 are $(-3, -1)$ and $(4, -1)$, so the solution of the equation $f(x) = -1$ is $x = -3$ or $x = 4$.

(f) For any $x_1 < x_2$ in the interval $[-4, 0]$, we have $g(x_1) > g(x_2)$. Thus, $g(x)$ is decreasing on $[-4, 0]$.

(g) The domain of f is $\{x \mid -4 \leq x \leq 4\} = [-4, 4]$. The range of f is $\{y \mid -2 \leq y \leq 3\} = [-2, 3]$.

(h) The domain of g is $\{x \mid -4 \leq x \leq 3\} = [-4, 3]$. Estimating the lowest point of the graph of g as having coordinates $(0, 0.5)$, the range of g is approximately $\{y \mid 0.5 \leq y \leq 4\} = [0.5, 4]$.

5. From Figure 1 in the text, the lowest point occurs at about $(t, a) = (12, -85)$. The highest point occurs at about $(17, 115)$. Thus, the range of the vertical ground acceleration is $-85 \leq a \leq 115$. Written in interval notation, the range is $[-85, 115]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. *Example 1:* A car is driven at 60 mi/h for 2 hours. The distance d traveled by the car is a function of the time t. The domain of the function is $\{t \mid 0 \le t \le 2\}$, where t is measured in hours. The range of the function is $\{d \mid 0 \le d \le 120\}$, where d is measured in miles.

Example 2: At a certain university, the number of students N on campus at any time on a particular day is a function of the time t after midnight. The domain of the function is $\{t \mid 0 \le t \le 24\}$, where t is measured in hours. The range of the function is $\{N \mid 0 \le N \le k\}$, where N is an integer and k is the largest number of students on campus at once.

Example 3: A certain employee is paid \$8.00 per hour and works a maximum of 30 hours per week. The number of hours worked is rounded down to the nearest quarter of an hour. This employee's gross weekly pay P is a function of the number of hours worked h. The domain of the function is $[0, 30]$ and the range of the function is $\{0, 2.00, 4.00, \ldots, 238.00, 240.00\}$.

7. We solve $3x - 5y = 7$ for y: $3x - 5y = 7 \quad \Leftrightarrow \quad -5y = -3x + 7 \quad \Leftrightarrow \quad y = \frac{3}{5}x - \frac{7}{5}$. Since the equation determines exactly one value of y for each value of x, the equation defines y as a function of x.

8. We solve $3x^2 - 2y = 5$ for y: $3x^2 - 2y = 5 \quad \Leftrightarrow \quad -2y = -3x^2 + 5 \quad \Leftrightarrow \quad y = \frac{3}{2}x^2 - \frac{5}{2}$. Since the equation determines exactly one value of y for each value of x, the equation defines y as a function of x.

9. We solve $x^2 + (y-3)^2 = 5$ for y: $x^2 + (y-3)^2 = 5 \quad \Leftrightarrow \quad (y-3)^2 = 5 - x^2 \quad \Leftrightarrow \quad y - 3 = \pm\sqrt{5 - x^2} \quad \Leftrightarrow$ $y = 3 \pm \sqrt{5 - x^2}$. Some input values x correspond to more than one output y. (For instance, $x = 1$ corresponds to $y = 1$ and to $y = 5$.) Thus, the equation does *not* define y as a function of x.

10. We solve $2xy + 5y^2 = 4$ for y: $2xy + 5y^2 = 4 \quad \Leftrightarrow \quad 5y^2 + (2x)\,y - 4 = 0 \quad \Leftrightarrow$

$y = \dfrac{-2x \pm \sqrt{(2x)^2 - 4(5)(-4)}}{2(5)} = \dfrac{-2x \pm \sqrt{4x^2 + 80}}{10} = \dfrac{-x \pm \sqrt{x^2 + 20}}{5}$ (using the quadratic formula). Some input values x correspond to more than one output y. (For instance, $x = 4$ corresponds to $y = -2$ and to $y = 2/5$.) Thus, the equation does *not* define y as a function of x.

11. We solve $(y+3)^3 + 1 = 2x$ for y: $(y+3)^3 + 1 = 2x \quad \Leftrightarrow \quad (y+3)^3 = 2x - 1 \quad \Leftrightarrow \quad y + 3 = \sqrt[3]{2x - 1} \quad \Leftrightarrow$ $y = -3 + \sqrt[3]{2x - 1}$. Since the equation determines exactly one value of y for each value of x, the equation defines y as a function of x.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. We solve $2x - |y| = 0$ for y: $2x - |y| = 0 \quad \Leftrightarrow \quad |y| = 2x \quad \Leftrightarrow \quad y = \pm 2x$. Some input values x correspond to more than one output y. (For instance, $x = 1$ corresponds to $y = -2$ and to $y = 2$.) Thus, the equation does *not* define y as a function of x.

13. The height 60 in $(x = 60)$ corresponds to shoe sizes 7 and 8 $(y = 7$ and $y = 8)$. Since an input value x corresponds to more than output value y, the table does *not* define y as a function of x.

14. Each year x corresponds to exactly one tuition cost y. Thus, the table defines y as a function of x.

15. No, the curve is not the graph of a function because a vertical line intersects the curve more than once. Hence, the curve fails the Vertical Line Test.

16. Yes, the curve is the graph of a function because it passes the Vertical Line Test. The domain is $[-2, 2]$ and the range is $[-1, 2]$.

17. Yes, the curve is the graph of a function because it passes the Vertical Line Test. The domain is $[-3, 2]$ and the range is $[-3, -2) \cup [-1, 3]$.

18. No, the curve is not the graph of a function since for $x = 0$, ± 1, and ± 2, there are infinitely many points on the curve.

19. (a) When $t = 1950$, $T \approx 13.8°\text{C}$, so the global average temperature in 1950 was about $13.8°\text{C}$.

(b) When $T = 14.2°\text{C}$, $t \approx 1990$.

(c) The global average temperature was smallest in 1910 (the year corresponding to the lowest point on the graph) and largest in 2000 (the year corresponding to the highest point on the graph).

(d) When $t = 1910$, $T \approx 13.5°\text{C}$, and when $t = 2000$, $T \approx 14.4°\text{C}$. Thus, the range of T is about $[13.5, 14.4]$.

20. (a) The ring width varies from near 0 mm to about 1.6 mm, so the range of the ring width function is approximately $[0, 1.6]$.

(b) According to the graph, the earth gradually cooled from 1550 to 1700, warmed into the late 1700s, cooled again into the late 1800s, and has been steadily warming since then. In the mid-19th century, there was variation that could have been associated with volcanic eruptions.

21. The water will cool down almost to freezing as the ice melts. Then, when the ice has melted, the water will slowly warm up to room temperature.

22. The temperature of the pie would increase rapidly, level off to oven temperature, decrease rapidly, and then level off to room temperature.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. (a) The power consumption at 6 AM is 500 MW, which is obtained by reading the value of power P when $t = 6$ from the graph. At 6 PM we read the value of P when $t = 18$, obtaining approximately 730 MW.

(b) The minimum power consumption is determined by finding the time for the lowest point on the graph, $t = 4$, or 4 AM. The maximum power consumption corresponds to the highest point on the graph, which occurs just before $t = 12$, or right before noon. These times are reasonable, considering the power consumption schedules of most individuals and businesses.

24. Runner A won the race, reaching the finish line at 100 meters in about 15 seconds, followed by runner B with a time of about 19 seconds, and then by runner C who finished in around 23 seconds. B initially led the race, followed by C, and then A. C then passed B to lead for a while. Then A passed first B, and then passed C to take the lead and finish first. Finally, B passed C to finish in second place. All three runners completed the race.

25. Of course, this graph depends strongly on the geographical location!

26. The summer solstice (the longest day of the year) is around June 21, and the winter solstice (the shortest day) is around December 22. (Exchange the dates for the southern hemisphere.)

27. As the price increases, the amount sold decreases.

28. The value of the car decreases fairly rapidly initially, then somewhat less rapidly.

29.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. (a)

(b)

(c)

(d)

31. (a)

(b) 9:00 AM corresponds to $t = 9$. When $t = 9$, the temperature T is about $74°\text{F}$.

32. (a)

(b) The blood alcohol concentration rises rapidly, then slowly decreases to near zero.

33. $f(x) = 3x^2 - x + 2$.

$f(2) = 3(2)^2 - 2 + 2 = 12 - 2 + 2 = 12$.

$f(-2) = 3(-2)^2 - (-2) + 2 = 12 + 2 + 2 = 16$.

$f(a) = 3a^2 - a + 2$.

$f(-a) = 3(-a)^2 - (-a) + 2 = 3a^2 + a + 2$.

$f(a+1) = 3(a+1)^2 - (a+1) + 2 = 3(a^2 + 2a + 1) - a - 1 + 2 = 3a^2 + 6a + 3 - a + 1 = 3a^2 + 5a + 4$.

$2f(a) = 2 \cdot f(a) = 2(3a^2 - a + 2) = 6a^2 - 2a + 4$.

$f(2a) = 3(2a)^2 - (2a) + 2 = 3(4a^2) - 2a + 2 = 12a^2 - 2a + 2$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$f(a^2) = 3(a^2)^2 - (a^2) + 2 = 3(a^4) - a^2 + 2 = 3a^4 - a^2 + 2.$

$$[f(a)]^2 = \left[3a^2 - a + 2\right]^2 = \left(3a^2 - a + 2\right)\left(3a^2 - a + 2\right)$$
$$= 9a^4 - 3a^3 + 6a^2 - 3a^3 + a^2 - 2a + 6a^2 - 2a + 4 = 9a^4 - 6a^3 + 13a^2 - 4a + 4.$$

$f(a+h) = 3(a+h)^2 - (a+h) + 2 = 3(a^2 + 2ah + h^2) - a - h + 2 = 3a^2 + 6ah + 3h^2 - a - h + 2.$

34. $g(x) = \dfrac{x}{\sqrt{x+1}}.$

$g(0) = \dfrac{0}{\sqrt{0+1}} = 0.$

$g(3) = \dfrac{3}{\sqrt{3+1}} = \dfrac{3}{2}.$

$5g(a) = 5 \cdot \dfrac{a}{\sqrt{a+1}} = \dfrac{5a}{\sqrt{a+1}}.$

$\dfrac{1}{2}g(4a) = \dfrac{1}{2} \cdot g(4a) = \dfrac{1}{2} \cdot \dfrac{4a}{\sqrt{4a+1}} = \dfrac{2a}{\sqrt{4a+1}}.$

$g(a^2) = \dfrac{a^2}{\sqrt{a^2+1}}$; $[g(a)]^2 = \left(\dfrac{a}{\sqrt{a+1}}\right)^2 = \dfrac{a^2}{a+1}.$

$g(a+h) = \dfrac{(a+h)}{\sqrt{(a+h)+1}} = \dfrac{a+h}{\sqrt{a+h+1}}.$

$g(x-a) = \dfrac{(x-a)}{\sqrt{(x-a)+1}} = \dfrac{x-a}{\sqrt{x-a+1}}.$

35. $f(x) = 4 + 3x - x^2$, so $f(3+h) = 4 + 3(3+h) - (3+h)^2 = 4 + 9 + 3h - (9 + 6h + h^2) = 4 - 3h - h^2$,

and $\dfrac{f(3+h) - f(3)}{h} = \dfrac{(4 - 3h - h^2) - 4}{h} = \dfrac{h(-3-h)}{h} = -3 - h.$

36. $f(x) = x^3$, so $f(a+h) = (a+h)^3 = a^3 + 3a^2h + 3ah^2 + h^3$,

and $\dfrac{f(a+h) - f(a)}{h} = \dfrac{(a^3 + 3a^2h + 3ah^2 + h^3) - a^3}{h} = \dfrac{h(3a^2 + 3ah + h^2)}{h} = 3a^2 + 3ah + h^2.$

37. $f(x) = \dfrac{1}{x}$, so $\dfrac{f(x) - f(a)}{x - a} = \dfrac{\dfrac{1}{x} - \dfrac{1}{a}}{x - a} = \dfrac{\dfrac{a-x}{xa}}{x-a} = \dfrac{a-x}{xa(x-a)} = \dfrac{-1(x-a)}{xa(x-a)} = -\dfrac{1}{ax}.$

38. $f(x) = \sqrt{x+2}$, so $\dfrac{f(x) - f(1)}{x-1} = \dfrac{\sqrt{x+2} - \sqrt{3}}{x-1}$. Depending upon the context, this may be considered simplified.

Note: We may also rationalize the numerator:

$$\frac{\sqrt{x+2} - \sqrt{3}}{x-1} = \frac{\sqrt{x+2} - \sqrt{3}}{x-1} \cdot \frac{\sqrt{x+2} + \sqrt{3}}{\sqrt{x+2} + \sqrt{3}} = \frac{(x+2) - 3}{(x-1)\left(\sqrt{x-2} + \sqrt{3}\right)}$$
$$= \frac{x-1}{(x-1)\left(\sqrt{x-2} + \sqrt{3}\right)} = \frac{1}{\sqrt{x+2} + \sqrt{3}}$$

39. $f(x) = (x+4)/(x^2 - 9)$ is defined for all x except when $0 = x^2 - 9 \quad \Leftrightarrow \quad 0 = (x+3)(x-3) \quad \Leftrightarrow \quad x = -3$ or 3, so the domain is $\{x \in \mathbb{R} \mid x \neq -3, 3\} = (-\infty, -3) \cup (-3, 3) \cup (3, \infty)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. The function $f(x) = \dfrac{x^2+1}{x^2+4x-21}$ is defined for all values of x except those for which $x^2+4x-21=0 \quad \Leftrightarrow$ $(x+7)(x-3)=0 \quad \Leftrightarrow \quad x=-7$ or $x=3$. Thus, the domain is $\{x \in \mathbb{R} \mid x \neq -7, 3\} = (-\infty,-7) \cup (-7,3) \cup (3,\infty)$.

41. $f(t) = \sqrt[3]{2t-1}$ is defined for all real numbers. In fact $\sqrt[3]{p(t)}$, where $p(t)$ is a polynomial, is defined for all real numbers. Thus, the domain is $\mathbb{R}$, or $(-\infty,\infty)$.

42. $g(t) = \sqrt{3-t} - \sqrt{2+t}$ is defined when $3-t \geq 0 \quad \Leftrightarrow \quad t \leq 3$ and $2+t \geq 0 \quad \Leftrightarrow \quad t \geq -2$. Thus, the domain is $-2 \leq t \leq 3$, or $[-2,3]$.

43. $h(x) = 1 \,/\, \sqrt[4]{x^2-5x}$ is defined when $x^2-5x>0 \quad \Leftrightarrow \quad x(x-5)>0$. Note that $x^2-5x \neq 0$ since that would result in division by zero. The expression $x(x-5)$ is positive if $x<0$ or $x>5$. (See Appendix A for methods for solving inequalities.) Thus, the domain is $(-\infty,0) \cup (5,\infty)$.

44. $f(u) = \dfrac{u+1}{1+\dfrac{1}{u+1}}$ is defined when $u+1 \neq 0$ $[u \neq -1]$ and $1+\dfrac{1}{u+1} \neq 0$. Since $1+\dfrac{1}{u+1}=0 \quad \Leftrightarrow$ $\dfrac{1}{u+1}=-1 \quad \Leftrightarrow \quad 1=-u-1 \quad \Leftrightarrow \quad u=-2$, the domain is $\{u \mid u \neq -2, u \neq -1\} = (-\infty,-2) \cup (-2,-1) \cup (-1,\infty)$.

45. $F(p) = \sqrt{2-\sqrt{p}}$ is defined when $p \geq 0$ and $2-\sqrt{p} \geq 0$. Since $2-\sqrt{p} \geq 0 \quad \Leftrightarrow \quad 2 \geq \sqrt{p} \quad \Leftrightarrow \quad \sqrt{p} \leq 2 \quad \Leftrightarrow$ $0 \leq p \leq 4$, the domain is $[0,4]$.

46. The function $h(x) = \sqrt{x^2-4x-5}$ is defined when $x^2-4x-5 \geq 0 \quad \Leftrightarrow \quad (x+1)(x-5) \geq 0$. The polynomial $p(x) = x^2-4x-5$ may change signs only at its zeros, so we test values of x on the intervals separated by $x=-1$ and $x=5$: $p(-2)=7>0$, $p(0)=-5<0$, and $p(6)=7>0$. Thus, the domain of h, equivalent to the solution intervals of $p(x) \geq 0$, is $\{x \mid x \leq -1 \text{ or } x \geq 5\} = (-\infty,-1] \cup [5,\infty)$.

47. $h(x) = \sqrt{4-x^2}$. Now $y = \sqrt{4-x^2} \quad \Rightarrow \quad y^2 = 4-x^2 \quad \Leftrightarrow \quad x^2+y^2=4$, so the graph is the top half of a circle of radius 2 with center at the origin. The domain is $\{x \mid 4-x^2 \geq 0\} = \{x \mid 4 \geq x^2\} = \{x \mid 2 \geq |x|\} = [-2,2]$. From the graph, the range is $0 \leq y \leq 2$, or $[0,2]$.

48. The function $f(x) = \dfrac{x^2-4}{x-2}$ is defined when $x-2 \neq 0 \Leftrightarrow x \neq 2$, so the domain is $\{x \mid x \neq 2\} = (-\infty,2) \cup (2,\infty)$. On its domain, $f(x) = \dfrac{x^2-4}{x-2} = \dfrac{(x-2)(x+2)}{x-2} = x+2$. Thus, the graph of f is the line $y=x+2$ with a hole at $(2,4)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. $f(x) = \begin{cases} x^2 + 2 & \text{if } x < 0 \\ x & \text{if } x \geq 0 \end{cases}$

$f(-3) = (-3)^2 + 2 = 11$, $f(0) = 0$, and $f(2) = 2$.

50. $f(x) = \begin{cases} 5 & \text{if } x < 2 \\ \frac{1}{2}x - 3 & \text{if } x \geq 2 \end{cases}$

$f(-3) = 5$, $f(0) = 5$, and $f(2) = \frac{1}{2}(2) - 3 = -2$.

51. $f(x) = \begin{cases} x + 1 & \text{if } x \leq -1 \\ x^2 & \text{if } x > -1 \end{cases}$

$f(-3) = -3 + 1 = -2$, $f(0) = 0^2 = 0$, and $f(2) = 2^2 = 4$.

52. $f(x) = \begin{cases} -1 & \text{if } x \leq 1 \\ 7 - 2x & \text{if } x > 1 \end{cases}$

$f(-3) = -1$, $f(0) = -1$, and $f(2) = 7 - 2(2) = 3$.

53. $|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$

so $f(x) = x + |x| = \begin{cases} 2x & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases}$

Graph the line $y = 2x$ for $x \geq 0$ and graph $y = 0$ (the x-axis) for $x < 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. $f(x) = |x+2| = \begin{cases} x+2 & \text{if } x+2 \geq 0 \\ -(x+2) & \text{if } x+2 < 0 \end{cases}$

$= \begin{cases} x+2 & \text{if } x \geq -2 \\ -x-2 & \text{if } x < -2 \end{cases}$

55. $g(t) = |1-3t| = \begin{cases} 1-3t & \text{if } 1-3t \geq 0 \\ -(1-3t) & \text{if } 1-3t < 0 \end{cases}$

$= \begin{cases} 1-3t & \text{if } t \leq \frac{1}{3} \\ 3t-1 & \text{if } t > \frac{1}{3} \end{cases}$

56. $f(x) = \dfrac{|x|}{x}$

The domain of f is $\{x \mid x \neq 0\}$ and $|x| = x$ if $x > 0$, $|x| = -x$ if $x < 0$.

So we can write

$$f(x) = \begin{cases} \dfrac{-x}{x} = -1 & \text{if } x < 0 \\ \dfrac{x}{x} = 1 & \text{if } x > 0 \end{cases}$$

57. To graph $f(x) = \begin{cases} |x| & \text{if } |x| \leq 1 \\ 1 & \text{if } |x| > 1 \end{cases}$, graph $y = |x|$ [Figure 16]

for $-1 \leq x \leq 1$ and graph $y = 1$ for $x > 1$ and for $x < -1$.

We could rewrite f as $f(x) = \begin{cases} 1 & \text{if } x < -1 \\ -x & \text{if } -1 \leq x < 0 \\ x & \text{if } 0 \leq x \leq 1 \\ 1 & \text{if } x > 1 \end{cases}$.

58. $g(x) = \big||x|-1\big| = \begin{cases} |x|-1 & \text{if } |x|-1 \geq 0 \\ -(|x|-1) & \text{if } |x|-1 < 0 \end{cases}$

$= \begin{cases} |x|-1 & \text{if } |x| \geq 1 \\ -|x|+1 & \text{if } |x| < 1 \end{cases}$

$= \begin{cases} x-1 & \text{if } |x| \geq 1 \text{ and } x \geq 0 \\ -x-1 & \text{if } |x| \geq 1 \text{ and } x < 0 \\ -x+1 & \text{if } |x| < 1 \text{ and } x \geq 0 \\ -(-x)+1 & \text{if } |x| < 1 \text{ and } x < 0 \end{cases} = \begin{cases} x-1 & \text{if } x \geq 1 \\ -x-1 & \text{if } x \leq -1 \\ -x+1 & \text{if } 0 \leq x < 1 \\ x+1 & \text{if } -1 < x < 0 \end{cases}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. Recall that the slope m of a line between the two points (x_1, y_1) and (x_2, y_2) is $m = \dfrac{y_2 - y_1}{x_2 - x_1}$ and an equation of the line connecting those two points is $y - y_1 = m(x - x_1)$. The slope of the line segment joining the points $(1, -3)$ and $(5, 7)$ is $\dfrac{7-(-3)}{5-1} = \dfrac{5}{2}$, so an equation is $y - (-3) = \frac{5}{2}(x - 1)$. The function is $f(x) = \frac{5}{2}x - \frac{11}{2}$, $1 \le x \le 5$.

60. The slope of the line segment joining the points $(-5, 10)$ and $(7, -10)$ is $\dfrac{-10-10}{7-(-5)} = -\dfrac{5}{3}$, so an equation is $y - 10 = -\frac{5}{3}[x - (-5)]$. The function is $f(x) = -\frac{5}{3}x + \frac{5}{3}$, $-5 \le x \le 7$.

61. We need to solve the given equation for y. $x + (y-1)^2 = 0 \;\Leftrightarrow\; (y-1)^2 = -x \;\Leftrightarrow\; y - 1 = \pm\sqrt{-x} \;\Leftrightarrow\; y = 1 \pm \sqrt{-x}$. The expression with the positive radical represents the top half of the parabola, and the one with the negative radical represents the bottom half. Hence, we want $f(x) = 1 - \sqrt{-x}$. Note that the domain is $x \le 0$.

62. $x^2 + (y-2)^2 = 4 \;\Leftrightarrow\; (y-2)^2 = 4 - x^2 \;\Leftrightarrow\; y - 2 = \pm\sqrt{4-x^2} \;\Leftrightarrow\; y = 2 \pm \sqrt{4-x^2}$. The top half is given by the function $f(x) = 2 + \sqrt{4-x^2}$, $-2 \le x \le 2$.

63. For $0 \le x \le 3$, the graph is the line with slope -1 and y-intercept 3, that is, $y = -x + 3$. For $3 < x \le 5$, the graph is the line with slope 2 passing through $(3, 0)$; that is, $y - 0 = 2(x - 3)$, or $y = 2x - 6$. So the function is

$$f(x) = \begin{cases} -x + 3 & \text{if } 0 \le x \le 3 \\ 2x - 6 & \text{if } 3 < x \le 5 \end{cases}$$

64. For $-4 \le x \le -2$, the graph is the line with slope $-\frac{3}{2}$ passing through $(-2, 0)$; that is, $y - 0 = -\frac{3}{2}[x - (-2)]$, or $y = -\frac{3}{2}x - 3$. For $-2 < x < 2$, the graph is the top half of the circle with center $(0, 0)$ and radius 2. An equation of the circle is $x^2 + y^2 = 4$, so an equation of the top half is $y = \sqrt{4 - x^2}$. For $2 \le x \le 4$, the graph is the line with slope $\frac{3}{2}$ passing through $(2, 0)$; that is, $y - 0 = \frac{3}{2}(x - 2)$, or $y = \frac{3}{2}x - 3$. So the function is

$$f(x) = \begin{cases} -\frac{3}{2}x - 3 & \text{if } -4 \le x \le -2 \\ \sqrt{4 - x^2} & \text{if } -2 < x < 2 \\ \frac{3}{2}x - 3 & \text{if } \;\; 2 \le x \le 4 \end{cases}$$

65. Let the length and width of the rectangle be L and W. Then the perimeter is $2L + 2W = 20$ and the area is $A = LW$. Solving the first equation for W in terms of L gives $W = \dfrac{20 - 2L}{2} = 10 - L$. Thus, $A(L) = L(10 - L) = 10L - L^2$. Since lengths are positive, the domain of A is $0 < L < 10$. If we further restrict L to be larger than W, then $5 < L < 10$ would be the domain.

66. Let the length and width of the rectangle be L and W. Then the area is $LW = 16$, so that $W = 16/L$. The perimeter is $P = 2L + 2W$, so $P(L) = 2L + 2(16/L) = 2L + 32/L$, and the domain of P is $L > 0$, since lengths must be positive quantities. If we further restrict L to be larger than W, then $L > 4$ would be the domain.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. Let the length of a side of the equilateral triangle be x. Then by the Pythagorean Theorem, the height y of the triangle satisfies $y^2 + \left(\frac{1}{2}x\right)^2 = x^2$, so that $y^2 = x^2 - \frac{1}{4}x^2 = \frac{3}{4}x^2$ and $y = \frac{\sqrt{3}}{2}x$. Using the formula for the area A of a triangle, $A = \frac{1}{2}(\text{base})(\text{height})$, we obtain $A(x) = \frac{1}{2}(x)\left(\frac{\sqrt{3}}{2}x\right) = \frac{\sqrt{3}}{4}x^2$, with domain $x > 0$.

68. Let the length, width, and height of the closed rectangular box be denoted by L, W, and H, respectively. The length is twice the width, so $L = 2W$. The volume V of the box is given by $V = LWH$. Since $V = 8$, we have $8 = (2W)WH \quad \Rightarrow$ $8 = 2W^2H \quad \Rightarrow \quad H = \dfrac{8}{2W^2} = \dfrac{4}{W^2}$, and so $H = f(W) = \dfrac{4}{W^2}$.

69. Let each side of the base of the box have length x, and let the height of the box be h. Since the volume is 2, we know that $2 = hx^2$, so that $h = 2/x^2$, and the surface area is $S = x^2 + 4xh$. Thus, $S(x) = x^2 + 4x(2/x^2) = x^2 + (8/x)$, with domain $x > 0$.

70. Let r and h denote the radius and the height of the right circular cylinder, respectively. Then the volume V is given by $V = \pi r^2 h$, and for this particular cylinder we have $\pi r^2 h = 25 \quad \Leftrightarrow \quad r^2 = \dfrac{25}{\pi h}$. Solving for r and rejecting the negative solution gives $r = \dfrac{5}{\sqrt{\pi h}}$, so $r = f(h) = \dfrac{5}{\sqrt{\pi h}}$ in.

71. The height of the box is x and the length and width are $L = 20 - 2x$, $W = 12 - 2x$. Then $V = LWx$ and so $V(x) = (20 - 2x)(12 - 2x)(x) = 4(10 - x)(6 - x)(x) = 4x(60 - 16x + x^2) = 4x^3 - 64x^2 + 240x$.
The sides L, W, and x must be positive. Thus, $L > 0 \quad \Leftrightarrow \quad 20 - 2x > 0 \quad \Leftrightarrow \quad x < 10$;
$W > 0 \quad \Leftrightarrow \quad 12 - 2x > 0 \quad \Leftrightarrow \quad x < 6$; and $x > 0$. Combining these restrictions gives us the domain $0 < x < 6$.

72. The area of the window is $A = xh + \frac{1}{2}\pi\left(\frac{1}{2}x\right)^2 = xh + \dfrac{\pi x^2}{8}$, where h is the height of the rectangular portion of the window. The perimeter is $P = 2h + x + \frac{1}{2}\pi x = 30 \quad \Leftrightarrow \quad 2h = 30 - x - \frac{1}{2}\pi x \quad \Leftrightarrow \quad h = \frac{1}{4}(60 - 2x - \pi x)$. Thus,

$$A(x) = x\,\frac{60 - 2x - \pi x}{4} + \frac{\pi x^2}{8} = 15x - \frac{1}{2}x^2 - \frac{\pi}{4}x^2 + \frac{\pi}{8}x^2 = 15x - \frac{4}{8}x^2 - \frac{\pi}{8}x^2 = 15x - x^2\left(\frac{\pi + 4}{8}\right).$$

Since the lengths x and h must be positive quantities, we have $x > 0$ and $h > 0$. For $h > 0$, we have $2h > 0 \quad \Leftrightarrow$ $30 - x - \frac{1}{2}\pi x > 0 \quad \Leftrightarrow \quad 60 > 2x + \pi x \quad \Leftrightarrow \quad x < \dfrac{60}{2 + \pi}$. Hence, the domain of A is $0 < x < \dfrac{60}{2 + \pi}$.

73. We can summarize the amount of the fine with a piecewise defined function.

$$F(x) = \begin{cases} 15(40 - x) & \text{if } 0 \le x < 40 \\ 0 & \text{if } 40 \le x \le 65 \\ 15(x - 65) & \text{if } x > 65 \end{cases}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

74. For the first 1200 kWh, $E(x) = 10 + 0.06x$.

For usage over 1200 kWh, the cost is

$E(x) = 10 + 0.06(1200) + 0.07(x - 1200) = 82 + 0.07(x - 1200)$.

Thus,

$$E(x) = \begin{cases} 10 + 0.06x & \text{if } 0 \le x \le 1200 \\ 82 + 0.07(x - 1200) & \text{if } x > 1200 \end{cases}$$

75. (a)

(b) On \$14,000, tax is assessed on \$4000, and $10\%(\$4000) = \$400$.

On \$26,000, tax is assessed on \$16,000, and

$10\%(\$10{,}000) + 15\%(\$6000) = \$1000 + \$900 = \$1900$.

(c) As in part (b), there is \$1000 tax assessed on \$20,000 of income, so the graph of T is a line segment from $(10{,}000, 0)$ to $(20{,}000, 1000)$. The tax on \$30,000 is \$2500, so the graph of T for $x > 20{,}000$ is the ray with initial point $(20{,}000, 1000)$ that passes through $(30{,}000, 2500)$.

76. (a) Because an even function is symmetric with respect to the y-axis, and the point $(5, 3)$ is on the graph of this even function, the point $(-5, 3)$ must also be on its graph.

(b) Because an odd function is symmetric with respect to the origin, and the point $(5, 3)$ is on the graph of this odd function, the point $(-5, -3)$ must also be on its graph.

77. f is an odd function because its graph is symmetric about the origin. g is an even function because its graph is symmetric with respect to the y-axis.

78. f is not an even function since it is not symmetric with respect to the y-axis. f is not an odd function since it is not symmetric about the origin. Hence, f is *neither* even nor odd. g is an even function because its graph is symmetric with respect to the y-axis.

79. (a) The graph of an even function is symmetric about the y-axis. We reflect the given portion of the graph of f about the y-axis in order to complete it.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) For an odd function, $f(-x) = -f(x)$. The graph of an odd function is symmetric about the origin. We rotate the given portion of the graph of f through $180°$ about the origin in order to complete it.

80. (a) The graph of an even function is symmetric about the y-axis. We reflect the given portion of the graph of f about the y-axis in order to complete it.

(b) The graph of an odd function is symmetric about the origin. We rotate the given portion of the graph of f through $180°$ about the origin in order to complete it.

81. $f(x) = \dfrac{x}{x^2+1}$.

$f(-x) = \dfrac{-x}{(-x)^2+1} = \dfrac{-x}{x^2+1} = -\dfrac{x}{x^2+1} = -f(x)$.

Since $f(-x) = -f(x)$, f is an odd function.

82. $f(x) = \dfrac{x^2}{x^4+1}$.

$f(-x) = \dfrac{(-x)^2}{(-x)^4+1} = \dfrac{x^2}{x^4+1} = f(x)$.

Since $f(-x) = f(x)$, f is an even function.

83. $f(x) = \dfrac{x}{x+1}$, so $f(-x) = \dfrac{-x}{-x+1} = \dfrac{x}{x-1}$.

Since this is neither $f(x)$ nor $-f(x)$, the function f is neither even nor odd.

84. $f(x) = x\,|x|$.

$f(-x) = (-x)\,|-x| = (-x)\,|x| = -(x\,|x|)$
$= -f(x)$

Since $f(-x) = -f(x)$, f is an odd function.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

85. $f(x) = 1 + 3x^2 - x^4$.

$f(-x) = 1+3(-x)^2-(-x)^4 = 1+3x^2-x^4 = f(x)$.

Since $f(-x) = f(x)$, f is an even function.

86. $f(x) = 1 + 3x^3 - x^5$, so

$$f(-x) = 1 + 3(-x)^3 - (-x)^5 = 1 + 3(-x^3) - (-x^5)$$
$$= 1 - 3x^3 + x^5$$

Since this is neither $f(x)$ nor $-f(x)$, the function f is neither even nor odd.

87. (i) If f and g are both even functions, then $f(-x) = f(x)$ and $g(-x) = g(x)$. Now $(f+g)(-x) = f(-x) + g(-x) = f(x) + g(x) = (f+g)(x)$, so $f+g$ is an *even* function.

(ii) If f and g are both odd functions, then $f(-x) = -f(x)$ and $g(-x) = -g(x)$. Now $(f+g)(-x) = f(-x) + g(-x) = -f(x) + [-g(x)] = -[f(x) + g(x)] = -(f+g)(x)$, so $f+g$ is an *odd* function.

(iii) If f is an even function and g is an odd function, then $(f+g)(-x) = f(-x) + g(-x) = f(x) + [-g(x)] = f(x) - g(x)$, which is not $(f+g)(x)$ nor $-(f+g)(x)$, so $f+g$ is *neither* even nor odd. (Exception: if f is the zero function, then $f+g$ will be *odd*. If g is the zero function, then $f+g$ will be *even*.)

88. (i) If f and g are both even functions, then $f(-x) = f(x)$ and $g(-x) = g(x)$. Now $(fg)(-x) = f(-x)g(-x) = f(x)g(x) = (fg)(x)$, so fg is an *even* function.

(ii) If f and g are both odd functions, then $f(-x) = -f(x)$ and $g(-x) = -g(x)$. Now $(fg)(-x) = f(-x)g(-x) = [-f(x)][-g(x)] = f(x)g(x) = (fg)(x)$, so fg is an *even* function.

(iii) If f is an even function and g is an odd function, then $(fg)(-x) = f(-x)g(-x) = f(x)[-g(x)] = -[f(x)g(x)] = -(fg)(x)$, so fg is an *odd* function.

1.2 Mathematical Models: A Catalog of Essential Functions

1. (a) $f(x) = x^3 + 3x^2$ is a polynomial function of degree 3. (This function is also an algebraic function.)

(b) $g(t) = \cos^2 t - \sin t$ is a trigonometric function.

(c) $r(t) = t^{\sqrt{3}}$ is a power function.

(d) $v(t) = 8^t$ is an exponential function.

(e) $y = \dfrac{\sqrt{x}}{x^2+1}$ is an algebraic function. It is the quotient of a root of a polynomial and a polynomial of degree 2.

(f) $g(u) = \log_{10} u$ is a logarithmic function.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. (a) $f(t) = \dfrac{3t^2+2}{t}$ is a rational function. (This function is also an algebraic function.)

(b) $h(r) = 2.3^r$ is an exponential function.

(c) $s(t) = \sqrt{t+4}$ is an algebraic function. It is a root of a polynomial.

(d) $y = x^4 + 5$ is a polynomial function of degree 4.

(e) $g(x) = \sqrt[3]{x}$ is a root function. Rewriting $g(x)$ as $x^{1/3}$, we recognize the function also as a power function. (This function is, further, an algebraic function because it is a root of a polynomial.)

(f) $y = \dfrac{1}{x^2}$ is a rational function. Rewriting y as x^{-2}, we recognize the function also as a power function. (This function is, further, an algebraic function because it is the quotient of two polynomials.)

3. We notice from the figure that g and h are even functions (symmetric with respect to the y-axis) and that f is an odd function (symmetric with respect to the origin). So (b) $\left[y = x^5\right]$ must be f. Since g is flatter than h near the origin, we must have (c) $\left[y = x^8\right]$ matched with g and (a) $\left[y = x^2\right]$ matched with h.

4. (a) The graph of $y = 3x$ is a line (choice G).

(b) $y = 3^x$ is an exponential function (choice f).

(c) $y = x^3$ is an odd polynomial function or power function (choice F).

(d) $y = \sqrt[3]{x} = x^{1/3}$ is a root function (choice g).

5. The denominator cannot equal 0, so $1 - \sin x \neq 0 \quad \Leftrightarrow \quad \sin x \neq 1 \quad \Leftrightarrow \quad x \neq \frac{\pi}{2} + 2n\pi$. Thus, the domain of $f(x) = \dfrac{\cos x}{1 - \sin x}$ is $\left\{x \mid x \neq \frac{\pi}{2} + 2n\pi,\ n \text{ an integer}\right\}$.

6. The denominator cannot equal 0, so $1 - \tan x \neq 0 \quad \Leftrightarrow \quad \tan x \neq 1 \quad \Leftrightarrow \quad x \neq \frac{\pi}{4} + n\pi$. The tangent function is not defined if $x \neq \frac{\pi}{2} + n\pi$. Thus, the domain of $g(x) = \dfrac{1}{1 - \tan x}$ is $\left\{x \mid x \neq \frac{\pi}{4} + n\pi,\ x \neq \frac{\pi}{2} + n\pi,\ n \text{ an integer}\right\}$.

7. (a) An equation for the family of linear functions with slope 2 is $y = f(x) = 2x + b$, where b is the y-intercept.

(b) $f(2) = 1$ means that the point $(2, 1)$ is on the graph of f. We can use the point-slope form of a line to obtain an equation for the family of linear functions through the point $(2, 1)$. $\quad y - 1 = m(x - 2)$, which is equivalent to $y = mx + (1 - 2m)$ in slope-intercept form.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) To belong to both families, an equation must have slope $m = 2$, so the equation in part (b), $y = mx + (1 - 2m)$, becomes $y = 2x - 3$. It is the *only* function that belongs to both families.

8. All members of the family of linear functions $f(x) = 1 + m(x + 3)$ have graphs that are lines passing through the point $(-3, 1)$.

9. All members of the family of linear functions $f(x) = c - x$ have graphs that are lines with slope -1. The y-intercept is c.

10. We graph $P(x) = x^3 - cx^2$ for $c = -2, 0, 1$, and 3. For $c \neq 0$, $P(x) = x^3 - cx^2 = x^2(x - c)$ has two x-intercepts, 0 and c. The curve has one decreasing portion that begins or ends at the origin and increases in length as $|c|$ increases; the decreasing portion is in quadrant II for $c < 0$ and in quadrant IV for $c > 0$.

11. Because f is a quadratic function, we know it is of the form $f(x) = ax^2 + bx + c$. The y-intercept is 18, so $f(0) = 18 \Rightarrow c = 18$ and $f(x) = ax^2 + bx + 18$. Since the points $(3, 0)$ and $(4, 2)$ lie on the graph of f, we have

$$f(3) = 0 \Rightarrow 9a + 3b + 18 = 0 \Rightarrow 3a + b = -6 \quad \textbf{(1)}$$

$$f(4) = 2 \Rightarrow 16a + 4b + 18 = 2 \Rightarrow 4a + b = -4 \quad \textbf{(2)}$$

This is a system of two equations in the unknowns a and b, and subtracting **(1)** from **(2)** gives $a = 2$. From **(1)**, $3(2) + b = -6 \Rightarrow b = -12$, so a formula for f is $f(x) = 2x^2 - 12x + 18$.

12. g is a quadratic function so $g(x) = ax^2 + bx + c$. The y-intercept is 1, so $g(0) = 1 \Rightarrow c = 1$ and $g(x) = ax^2 + bx + 1$. Since the points $(-2, 2)$ and $(1, -2.5)$ lie on the graph of g, we have

$$g(-2) = 2 \Rightarrow 4a - 2b + 1 = 2 \Rightarrow 4a - 2b = 1 \quad \textbf{(1)}$$

$$g(1) = -2.5 \Rightarrow a + b + 1 = -2.5 \Rightarrow a + b = -3.5 \quad \textbf{(2)}$$

Then **(1)** $+ 2 \cdot$ **(2)** gives us $6a = -6 \Rightarrow a = -1$ and from **(2)**, we have $-1 + b = -3.5 \Rightarrow b = -2.5$, so a formula for g is $g(x) = -x^2 - 2.5x + 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. Since $f(-1) = f(0) = f(2) = 0$, f has zeros of -1, 0, and 2, so an equation for f is $f(x) = a[x - (-1)](x - 0)(x - 2)$, or $f(x) = ax(x + 1)(x - 2)$. Because $f(1) = 6$, we'll substitute 1 for x and 6 for $f(x)$.

$6 = a(1)(2)(-1) \quad \Rightarrow \quad -2a = 6 \quad \Rightarrow \quad a = -3$, so an equation for f is $f(x) = -3x(x + 1)(x - 2)$.

14. (a) For $T = 0.02t + 8.50$, the slope is 0.02, which means that the average surface temperature of the world is increasing at a rate of $0.02\ ^\circ\mathrm{C}$ per year. The T-intercept is 8.50, which represents the average surface temperature in $^\circ\mathrm{C}$ in the year 1900.

(b) $t = 2100 - 1900 = 200 \quad \Rightarrow \quad T = 0.02(200) + 8.50 = 12.50\ ^\circ\mathrm{C}$

15. (a) $D = 200$, so $c = 0.0417D(a + 1) = 0.0417(200)(a + 1) = 8.34a + 8.34$. The slope is 8.34, which represents the change in mg of the dosage for a child for each change of 1 year in age.

(b) For a newborn, $a = 0$, so $c = 8.34$ mg.

16. (a)

(b) The slope of -4 means that for each increase of 1 dollar for a rental space, the number of spaces rented *decreases* by 4. The y-intercept of 200 is the number of spaces that would be occupied if there were no charge for each space. The x-intercept of 50 is the smallest rental fee that results in no spaces rented.

17. (a)

(b) The slope of $\frac{9}{5}$ means that F increases $\frac{9}{5}$ degrees for each increase of $1^\circ\mathrm{C}$. (Equivalently, F increases by 9 when C increases by 5 and F decreases by 9 when C decreases by 5.) The F-intercept of 32 is the Fahrenheit temperature corresponding to a Celsius temperature of 0.

18. (a) Jari is traveling faster since the line representing her distance versus time is steeper than the corresponding line for Jade.

(b) At $t = 0$, Jade has traveled 10 miles. At $t = 6$, Jade has traveled 16 miles. Thus, Jade's speed is

$$\frac{16 \text{ miles} - 10 \text{ miles}}{6 \text{ minutes} - 0 \text{ minutes}} = 1 \text{ mi/min. This is } \frac{1 \text{ mile}}{1 \text{ minute}} \times \frac{60 \text{ minutes}}{1 \text{ hour}} = 60 \text{ mi/h}$$

At $t = 0$, Jari has traveled 0 miles. At $t = 6$, Jari has traveled 7 miles. Thus, Jari's speed is

$$\frac{7 \text{ miles} - 0 \text{ miles}}{6 \text{ minutes} - 0 \text{ minutes}} = \frac{7}{6} \text{ mi/min or } \frac{7 \text{ miles}}{6 \text{ minutes}} \times \frac{60 \text{ minutes}}{1 \text{ hour}} = 70 \text{ mi/h}$$

(c) From part (b), we have a slope of 1 (mile/minute) for the linear function f modeling the distance traveled by Jade and from the graph the y-intercept is 10. Thus, $f(t) = 1t + 10 = t + 10$. Similarly, we have a slope of $\frac{7}{6}$ miles/minute for

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Jari and a y-intercept of 0. Thus, the distance traveled by Jari as a function of time t (in minutes) is modeled by $g(t) = \frac{7}{6}t + 0 = \frac{7}{6}t$.

19. (a) Let x denote the number of chairs produced in one day and y the associated cost. Using the points $(100, 2200)$ and $(300, 4800)$, we get the slope $\frac{4800-2200}{300-100} = \frac{2600}{200} = 13$. So $y - 2200 = 13(x - 100) \quad \Leftrightarrow$ $y = 13x + 900$.

(b) The slope of the line in part (a) is 13 and it represents the cost (in dollars) of producing each additional chair.

(c) The y-intercept is 900 and it represents the fixed daily costs of operating the factory.

20. (a) Using d in place of x and C in place of y, we find the slope to be $\dfrac{C_2 - C_1}{d_2 - d_1} = \dfrac{460 - 380}{800 - 480} = \dfrac{80}{320} = \dfrac{1}{4}$.

So a linear equation is $C - 460 = \frac{1}{4}(d - 800) \quad \Leftrightarrow \quad C - 460 = \frac{1}{4}d - 200 \quad \Leftrightarrow \quad C = \frac{1}{4}d + 260$.

(b) Letting $d = 1500$ we get $C = \frac{1}{4}(1500) + 260 = 635$.

The cost of driving 1500 miles is \$635.

(c)

The slope of the line represents the cost per mile, \$0.25.

(d) The y-intercept represents the fixed cost, \$260.

(e) A linear function gives a suitable model in this situation because you have fixed monthly costs such as insurance and car payments, as well as costs that increase as you drive, such as gasoline, oil, and tires, and the cost of these for each additional mile driven is a constant.

21. (a) We are given $\dfrac{\text{change in pressure}}{\text{10 feet change in depth}} = \dfrac{4.34}{10} = 0.434$. Using P for pressure and d for depth with the point $(d, P) = (0, 15)$, we have the slope-intercept form of the line, $P = 0.434d + 15$.

(b) When $P = 100$, then $100 = 0.434d + 15 \quad \Leftrightarrow \quad 0.434d = 85 \quad \Leftrightarrow \quad d = \frac{85}{0.434} \approx 195.85$ feet. Thus, the pressure is $100\ \text{lb/in}^2$ at a depth of approximately 196 feet.

22. (a) $R(x) = kx^{-2}$ and $R(0.005) = 140$, so $140 = k(0.005)^{-2} \quad \Leftrightarrow \quad k = 140(0.005)^2 = 0.0035$.

(b) $R(x) = 0.0035x^{-2}$, so for a diameter of 0.008 m the resistance is $R(0.008) = 0.0035(0.008)^{-2} \approx 54.7$ ohms.

23. If x is the original distance from the source, then the illumination is $f(x) = kx^{-2} = k/x^2$. Moving halfway to the lamp gives an illumination of $f\left(\frac{1}{2}x\right) = k\left(\frac{1}{2}x\right)^{-2} = k(2/x)^2 = 4(k/x^2)$, so the light is four times as bright.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. (a) $P = k/V$ and $P = 39$ kPa when $V = 0.671\ \text{m}^3$, so $39 = k/0.671 \quad \Leftrightarrow \quad k = 39(0.671) = 26.169$.

(b) When $V = 0.916$, $P = 26.169/V = 26.169/0.916 \approx 28.6$, so the pressure is reduced to approximately 28.6 kPa.

25. (a) $P = kAv^3$ so doubling the windspeed v gives $P = kA(2v)^3 = 8(kAv^3)$. Thus, the power output is increased by a factor of eight.

(b) The area swept out by the blades is given by $A = \pi l^2$, where l is the blade length, so the power output is $P = kAv^3 = k\pi l^2 v^3$. Doubling the blade length gives $P = k\pi(2l)^2 v^3 = 4(k\pi l^2 v^3)$. Thus, the power output is increased by a factor of four.

(c) From part (b) we have $P = k\pi l^2 v^3$, and $k = 0.214\ \text{kg/m}^3$, $l = 30$ m gives

$$P = 0.214\ \frac{\text{kg}}{\text{m}^3} \cdot 900\pi\ \text{m}^2 \cdot v^3 = 192.6\pi v^3\ \frac{\text{kg}}{\text{m}}$$

For $v = 10\ \text{m/s}$, we have

$$P = 192.6\pi\left(10\ \frac{\text{m}}{\text{s}}\right)^3\ \frac{\text{kg}}{\text{m}} = 192{,}600\pi\ \frac{\text{m}^2 \cdot \text{kg}}{\text{s}^3} \approx 605{,}000\ \text{W}$$

Similarly, $v = 15\ \text{m/s}$ gives $P = 650{,}025\pi \approx 2{,}042{,}000$ W and $v = 25\ \text{m/s}$ gives $P = 3{,}009{,}375\pi \approx 9{,}454{,}000$ W.

26. (a) We graph $E(T) = (5.67 \times 10^{-8})T^4$ for $100 \le T \le 300$:

(b) From the graph, we see that as temperature increases, energy increases—slowly at first, but then at an increasing rate.

27. (a) The data appear to be periodic and a sine or cosine function would make the best model. A model of the form $f(x) = a\cos(bx) + c$ seems appropriate.

(b) The data appear to be decreasing in a linear fashion. A model of the form $f(x) = mx + b$ seems appropriate.

28. (a) The data appear to be increasing exponentially. A model of the form $f(x) = a \cdot b^x$ or $f(x) = a \cdot b^x + c$ seems appropriate.

(b) The data appear to be decreasing similarly to the values of the reciprocal function. A model of the form $f(x) = a/x$ seems appropriate.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Exercises 29–33: Some values are given to many decimal places. The results may depend on the technology used—rounding is left to the reader.

29. (a)

A linear model does seem appropriate.

(b) Using the points $(4000, 14.1)$ and $(60{,}000, 8.2)$, we obtain

$$y - 14.1 = \frac{8.2 - 14.1}{60{,}000 - 4000}(x - 4000) \text{ or, equivalently,}$$

$$y \approx -0.000105357x + 14.521429.$$

(c) Using a computing device, we obtain the regression line $y = -0.0000997855x + 13.950764$.

The following commands and screens illustrate how to find the regression line on a TI-84 Plus calculator.

Enter the data into list one (L1) and list two (L2). Press STAT 1 to enter the editor.

Find the regession line and store it in Y_1. Press 2nd QUIT STAT ▸ 4 VARS ▸ 1 1 ENTER.

Note from the last figure that the regression line has been stored in Y_1 and that Plot1 has been turned on (Plot1 is highlighted). You can turn on Plot1 from the Y= menu by placing the cursor on Plot1 and pressing ENTER or by pressing 2nd STAT PLOT 1 ENTER.

Now press ZOOM 9 to produce a graph of the data and the regression line. Note that choice 9 of the ZOOM menu automatically selects a window that displays all of the data.

(d) When $x = 25{,}000$, $y \approx 11.456$; or about 11.5 per 100 population.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(e) When $x = 80{,}000$, $y \approx 5.968$; or about a 6% chance.

(f) When $x = 200{,}000$, y is negative, so the model does not apply.

30. (a) Using a computing device, we obtain the regression line $y = 0.01879x + 0.30480$.

(b) The regression line appears to be a suitable model for the data.

(c) The y-intercept represents the percentage of laboratory rats that develop lung tumors when *not* exposed to asbestos fibers.

31. (a)

(b) Using a computing device, we obtain the regression line $y = 1.88074x + 82.64974$.

(c) When $x = 53$ cm, $y \approx 182.3$ cm.

32. (a) See the scatter plot in part (b). A linear model seems appropriate.

(b) Using a computing device, we obtain the regression line $y = 0.31567x + 8.15578$.

(c) For 2005, $x = 5$ and $y \approx 9.73$ cents/kWh. For 2017, $x = 17$ and $y \approx 13.52$ cents/kWh.

33. (a) See the scatter plot in part (b). A linear model seems appropriate.

(b) Using a computing device, we obtain the regression line $y = 1124.86x + 60{,}119.86$.

(c) For 2002, $x = 17$ and $y \approx 79{,}242$ thousands of barrels per day. For 2017, $x = 32$ and $y \approx 96{,}115$ thousands of barrels per day.

34. (a) $T = 1.000\,431\,227 d^{\,1.499\,528\,750}$

(b) The power model in part (a) is approximately $T = d^{1.5}$. Squaring both sides gives us $T^2 = d^3$, so the model matches Kepler's Third Law, $T^2 = kd^3$.

35. (a) If $A = 60$, then $S = 0.7A^{0.3} \approx 2.39$, so you would expect to find 2 species of bats in that cave.

(b) $S = 4 \;\Rightarrow\; 4 = 0.7A^{0.3} \;\Rightarrow\; \frac{40}{7} = A^{3/10} \;\Rightarrow\; A = \left(\frac{40}{7}\right)^{10/3} \approx 333.6$, so we estimate the surface area of the cave to be $334\ \mathrm{m}^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. (a) Using a computing device, we obtain a power function $N = cA^b$, where $c \approx 3.1046$ and $b \approx 0.308$.

(b) If $A = 291$, then $N = cA^b \approx 17.8$, so you would expect to find 18 species of reptiles and amphibians on Dominica.

37. We have $I = \dfrac{S}{4\pi r^2} = \left(\dfrac{S}{4\pi}\right)\left(\dfrac{1}{r^2}\right) = \dfrac{S/(4\pi)}{r^2}$. Thus, $I = \dfrac{k}{r^2}$ with $k = \dfrac{S}{4\pi}$.

1.3 New Functions from Old Functions

1. (a) If the graph of f is shifted 3 units upward, its equation becomes $y = f(x) + 3$.

(b) If the graph of f is shifted 3 units downward, its equation becomes $y = f(x) - 3$.

(c) If the graph of f is shifted 3 units to the right, its equation becomes $y = f(x - 3)$.

(d) If the graph of f is shifted 3 units to the left, its equation becomes $y = f(x + 3)$.

(e) If the graph of f is reflected about the x-axis, its equation becomes $y = -f(x)$.

(f) If the graph of f is reflected about the y-axis, its equation becomes $y = f(-x)$.

(g) If the graph of f is stretched vertically by a factor of 3, its equation becomes $y = 3f(x)$.

(h) If the graph of f is shrunk vertically by a factor of 3, its equation becomes $y = \frac{1}{3}f(x)$.

2. (a) To obtain the graph of $y = f(x) + 8$ from the graph of $y = f(x)$, shift the graph 8 units upward.

(b) To obtain the graph of $y = f(x + 8)$ from the graph of $y = f(x)$, shift the graph 8 units to the left.

(c) To obtain the graph of $y = 8f(x)$ from the graph of $y = f(x)$, stretch the graph vertically by a factor of 8.

(d) To obtain the graph of $y = f(8x)$ from the graph of $y = f(x)$, shrink the graph horizontally by a factor of 8.

(e) To obtain the graph of $y = -f(x) - 1$ from the graph of $y = f(x)$, first reflect the graph about the x-axis, and then shift it 1 unit downward.

(f) To obtain the graph of $y = 8f(\frac{1}{8}x)$ from the graph of $y = f(x)$, stretch the graph horizontally and vertically by a factor of 8.

3. (a) *Graph 3:* The graph of f is shifted 4 units to the right and has equation $y = f(x - 4)$.

(b) *Graph 1:* The graph of f is shifted 3 units upward and has equation $y = f(x) + 3$.

(c) *Graph 4:* The graph of f is shrunk vertically by a factor of 3 and has equation $y = \frac{1}{3}f(x)$.

(d) *Graph 5:* The graph of f is shifted 4 units to the left and reflected about the x-axis. Its equation is $y = -f(x + 4)$.

(e) *Graph 2:* The graph of f is shifted 6 units to the left and stretched vertically by a factor of 2. Its equation is $y = 2f(x + 6)$.

4. (a) $y = f(x) - 3$: Shift the graph of f 3 units down.

(b) $y = f(x + 1)$: Shift the graph of f 1 unit to the left.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $y = \frac{1}{2}f(x)$: Shrink the graph of f vertically by a factor of 2.

(d) $y = -f(x)$: Reflect the graph of f about the x-axis.

5. (a) To graph $y = f(2x)$ we shrink the graph of f horizontally by a factor of 2.

The point $(4, -1)$ on the graph of f corresponds to the point $\left(\frac{1}{2} \cdot 4, -1\right) = (2, -1)$.

(b) To graph $y = f\left(\frac{1}{2}x\right)$ we stretch the graph of f horizontally by a factor of 2.

The point $(4, -1)$ on the graph of f corresponds to the point $(2 \cdot 4, -1) = (8, -1)$.

(c) To graph $y = f(-x)$ we reflect the graph of f about the y-axis.

The point $(4, -1)$ on the graph of f corresponds to the point $(-1 \cdot 4, -1) = (-4, -1)$.

(d) To graph $y = -f(-x)$ we reflect the graph of f about the y-axis, then about the x-axis.

The point $(4, -1)$ on the graph of f corresponds to the point $(-1 \cdot 4, -1 \cdot -1) = (-4, 1)$.

6. The graph of $y = f(x) = \sqrt{3x - x^2}$ has been shifted 2 units to the right and stretched vertically by a factor of 2. Thus, a function describing the graph is

$$y = 2f(x-2) = 2\sqrt{3(x-2) - (x-2)^2} = 2\sqrt{3x - 6 - (x^2 - 4x + 4)} = 2\sqrt{-x^2 + 7x - 10}$$

7. The graph of $y = f(x) = \sqrt{3x - x^2}$ has been shifted 4 units to the left, reflected about the x-axis, and shifted downward 1 unit. Thus, a function describing the graph is

$$y = \underbrace{-1 \cdot}_{\substack{\text{reflect} \\ \text{about } x\text{-axis}}} f\underbrace{(x+4)}_{\substack{\text{shift} \\ \text{4 units left}}} \underbrace{-\ 1}_{\substack{\text{shift} \\ \text{1 unit left}}}$$

This function can be written as

$$\begin{aligned} y &= -f(x+4) - 1 = -\sqrt{3(x+4) - (x+4)^2} - 1 \\ &= -\sqrt{3x + 12 - (x^2 + 8x + 16)} - 1 = -\sqrt{-x^2 - 5x - 4} - 1 \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. (a) The graph of $y = 1 + \sqrt{x}$ can be obtained from the graph of $y = \sqrt{x}$ by shifting it upward 1 unit.

(b) The graph of $y = \sin \pi x$ can be obtained from the graph of $y = \sin x$ by compressing horizontally by a factor of π, giving a period of $2\pi/\pi = 2$. The graph of $y = 5 \sin \pi x$ is then obtained by stretching vertically by a factor of 5.

9. $y = 1 + x^2$. Start with the graph of $y = x^2$ and shift 1 unit upward

10. $y = (x + 1)^2$. Start with the graph of $y = x^2$ and shift 1 unit to the left.

11. $y = |x + 2|$. Start with the graph of $y = |x|$ and shift 2 units to the left.

12. $y = 1 - x^3$. Start with the graph of $y = x^3$, reflect about the x-axis, and then shift 1 unit upward.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $y = \dfrac{1}{x} + 2$. Start with the graph of $y = \dfrac{1}{x}$ and shift 2 units upward.

14. $y = -\sqrt{x} - 1$. Start with the graph of $y = \sqrt{x}$, reflect about the x-axis, and then shift 1 unit downward.

15. $y = \sin 4x$. Start with the graph of $y = \sin x$ and compress horizontally by a factor of 4. The period becomes $2\pi/4 = \pi/2$.

16. $y = 1 + \dfrac{1}{x^2}$. Start with the graph of $y = \dfrac{1}{x^2}$ and shift 1 unit upward.

17. $y = 2 + \sqrt{x+1}$. Start with the graph of $y = \sqrt{x}$, shift 1 unit to the left, and then shift 2 units upward.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $y = -(x-1)^2 + 3$. Start with the graph of $y = x^2$, shift 1 unit to the right, reflect about the x-axis, and then shift 3 units upward.

$y = x^2$ $\quad$ $y = (x-1)^2$ $\quad$ $y = -(x-1)^2$ $\quad$ $y = -(x-1)^2 + 3$

19. $y = x^2 - 2x + 5 = (x^2 - 2x + 1) + 4 = (x-1)^2 + 4$. Start with the graph of $y = x^2$, shift 1 unit to the right, and then shift 4 units upward.

20. $y = (x+1)^3 + 2$. Start with the graph of $y = x^3$, shift 1 unit to the left, and then shift 2 units upward.

21. $y = 2 - |x|$. Start with the graph of $y = |x|$, reflect about the x-axis, and then shift 2 units upward.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $y = 2 - 2\cos x$. Start with the graph of $y = \cos x$, reflect about the x-axis, stretch vertically by a factor of 2, and then shift 2 units upward.

23. $y = 3\sin\frac{1}{2}x + 1$. Start with the graph of $y = \sin x$, stretch horizontally by a factor of 2, stretch vertically by a factor of 3, and then shift 1 unit upward.

24. $y = \frac{1}{4}\tan\left(x - \frac{\pi}{4}\right)$. Start with the graph of $y = \tan x$, shift $\frac{\pi}{4}$ units to the right, and then compress vertically by a factor of 4.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $y = |\cos \pi x|$. Start with the graph of $y = \cos x$, shrink horizontally by a factor of π, and reflect all the parts of the graph below the x-axis about the x-axis.

26. $y = \left|\sqrt{x} - 1\right|$. Start with the graph of $y = \sqrt{x}$, shift 1 unit downward, and then reflect the portion of the graph below the x-axis about the x-axis.

27. This is just like the solution to Example 4 except the amplitude of the curve (the 30°N curve in Figure 9 on June 21) is $14 - 12 = 2$. So the function is $L(t) = 12 + 2\sin\left[\frac{2\pi}{365}(t - 80)\right]$. March 31 is the 90th day of the year, so the model gives $L(90) \approx 12.34$ h. The daylight time (5:51 AM to 6:18 PM) is 12 hours and 27 minutes, or 12.45 h. The model value differs from the actual value by $\frac{12.45 - 12.34}{12.45} \approx 0.009$, less than 1%.

28. Using a sine function to model the brightness of Delta Cephei as a function of time, we take its period to be 5.4 days, its amplitude to be 0.35 (on the scale of magnitude), and its average magnitude to be 4.0. If we take $t = 0$ at a time of average brightness, then the magnitude (brightness) as a function of time t in days can be modeled by the formula $M(t) = 4.0 + 0.35\sin\left(\frac{2\pi}{5.4}t\right)$.

29. The water depth $D(t)$ can be modeled by a cosine function with amplitude $\dfrac{12 - 2}{2} = 5$ m, average magnitude $\dfrac{12 + 2}{2} = 7$ m, and period 12 hours. High tide occurred at time 6:45 AM ($t = 6.75$ h), so the curve begins a cycle at time $t = 6.75$ h (shift 6.75 units to the right). Thus, $D(t) = 5\cos\left[\frac{2\pi}{12}(t - 6.75)\right] + 7 = 5\cos\left[\frac{\pi}{6}(t - 6.75)\right] + 7$, where D is in meters and t is the number of hours after midnight.

30. The total volume of air $V(t)$ in the lungs can be modeled by a sine function with amplitude $\dfrac{2500 - 2000}{2} = 250$ mL, average volume $\dfrac{2500 + 2000}{2} = 2250$ mL, and period 4 seconds. Thus, $V(t) = 250\sin\frac{2\pi}{4}t + 2250 = 250\sin\frac{\pi}{2}t + 2250$, where V is in mL and t is in seconds.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. (a) To obtain $y = f(|x|)$, the portion of the graph of $y = f(x)$ to the right of the y-axis is reflected about the y-axis.

(b) $y = \sin|x|$

(c) $y = \sqrt{|x|}$

32. The most important features of the given graph are the x-intercepts and the maximum and minimum points. The graph of $y = 1/f(x)$ has vertical asymptotes at the x-values where there are x-intercepts on the graph of $y = f(x)$. The maximum of 1 on the graph of $y = f(x)$ corresponds to a minimum of $1/1 = 1$ on $y = 1/f(x)$. Similarly, the minimum on the graph of $y = f(x)$ corresponds to a maximum on the graph of $y = 1/f(x)$. As the values of y get large (positively or negatively) on the graph of $y = f(x)$, the values of y get close to zero on the graph of $y = 1/f(x)$.

33. $f(x) = \sqrt{25 - x^2}$ is defined only when $25 - x^2 \geq 0 \quad \Leftrightarrow \quad x^2 \leq 25 \quad \Leftrightarrow \quad -5 \leq x \leq 5$, so the domain of f is $[-5, 5]$.

For $g(x) = \sqrt{x+1}$, we must have $x + 1 \geq 0 \quad \Leftrightarrow \quad x \geq -1$, so the domain of g is $[-1, \infty)$.

(a) $(f+g)(x) = \sqrt{25 - x^2} + \sqrt{x+1}$. The domain of $f + g$ is found by intersecting the domains of f and g: $[-1, 5]$.

(b) $(f-g)(x) = \sqrt{25 - x^2} - \sqrt{x+1}$. The domain of $f - g$ is found by intersecting the domains of f and g: $[-1, 5]$.

(c) $(fg)(x) = \sqrt{25 - x^2} \cdot \sqrt{x+1} = \sqrt{-x^3 - x^2 + 25x + 25}$. The domain of fg is found by intersecting the domains of f and g: $[-1, 5]$.

(d) $\left(\dfrac{f}{g}\right)(x) = \dfrac{\sqrt{25 - x^2}}{\sqrt{x+1}} = \sqrt{\dfrac{25 - x^2}{x+1}}$. Notice that we must have $x + 1 \neq 0$ in addition to any previous restrictions. Thus, the domain of f/g is $(-1, 5]$.

34. For $f(x) = \dfrac{1}{x-1}$, we must have $x - 1 \neq 0 \quad \Leftrightarrow \quad x \neq 1$. For $g(x) = \dfrac{1}{x} - 2$, we must have $x \neq 0$.

(a) $(f+g)(x) = \dfrac{1}{x-1} + \dfrac{1}{x} - 2 = \dfrac{x + x - 1 - 2x(x-1)}{x(x-1)} = \dfrac{2x - 1 - 2x^2 + 2x}{x^2 - x} = -\dfrac{2x^2 - 4x + 1}{x^2 - x}, \quad \{x \mid x \neq 0, 1\}$

(b) $(f-g)(x) = \dfrac{1}{x-1} - \left(\dfrac{1}{x} - 2\right) = \dfrac{x - (x-1) + 2x(x-1)}{x(x-1)} = \dfrac{1 + 2x^2 - 2x}{x^2 - x} = \dfrac{2x^2 - 2x + 1}{x^2 - x}, \quad \{x \mid x \neq 0, 1\}$

(c) $(fg)(x) = \dfrac{1}{x-1}\left(\dfrac{1}{x} - 2\right) = \dfrac{1}{x^2 - x} - \dfrac{2}{x-1} = \dfrac{1 - 2x}{x^2 - x}, \quad \{x \mid x \neq 0, 1\}$

(d) $\left(\dfrac{f}{g}\right)(x) = \dfrac{\dfrac{1}{x-1}}{\dfrac{1}{x} - 2} = \dfrac{\dfrac{1}{x-1}}{\dfrac{1-2x}{x}} = \dfrac{1}{x-1} \cdot \dfrac{x}{1-2x} = \dfrac{x}{(x-1)(1-2x)} = -\dfrac{x}{(x-1)(2x-1)}$

$= -\dfrac{x}{2x^2 - 3x + 1}, \quad \left\{x \mid x \neq 0, \tfrac{1}{2}, 1\right\}$

[Note the additional domain restriction $g(x) \neq 0 \quad \Rightarrow \quad x \neq \frac{1}{2}$.]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. $f(x) = x^3 + 5$ and $g(x) = \sqrt[3]{x}$. The domain of each function is $(-\infty, \infty)$.

(a) $(f \circ g)(x) = f(g(x)) = f\left(\sqrt[3]{x}\right) = \left(\sqrt[3]{x}\right)^3 + 5 = x + 5$. The domain is $(-\infty, \infty)$.

(b) $(g \circ f)(x) = g(f(x)) = g(x^3 + 5) = \sqrt[3]{x^3 + 5}$. The domain is $(-\infty, \infty)$.

(c) $(f \circ f)(x) = f(f(x)) = f(x^3 + 5) = (x^3 + 5)^3 + 5$. The domain is $(-\infty, \infty)$.

(d) $(g \circ g)(x) = g(g(x)) = g\left(\sqrt[3]{x}\right) = \sqrt[3]{\sqrt[3]{x}} = \sqrt[9]{x}$. The domain is $(-\infty, \infty)$.

36. $f(x) = 1/x$ and $g(x) = 2x + 1$. The domain of f is $(-\infty, 0) \cup (0, \infty)$. The domain of g is $(-\infty, \infty)$.

(a) $(f \circ g)(x) = f(g(x)) = f(2x + 1) = \dfrac{1}{2x + 1}$. The domain is
$\{x \mid 2x + 1 \neq 0\} = \left\{x \mid x \neq -\frac{1}{2}\right\} = \left(-\infty, -\frac{1}{2}\right) \cup \left(-\frac{1}{2}, \infty\right)$.

(b) $(g \circ f)(x) = g(f(x)) = g\left(\dfrac{1}{x}\right) = 2\left(\dfrac{1}{x}\right) + 1 = \dfrac{2}{x} + 1$. We must have $x \neq 0$, so the domain is $(-\infty, 0) \cup (0, \infty)$.

(c) $(f \circ f)(x) = f(f(x)) = f\left(\dfrac{1}{x}\right) = \dfrac{1}{1/x} = x$. Since f requires $x \neq 0$, the domain is $(-\infty, 0) \cup (0, \infty)$.

(d) $(g \circ g)(x) = g(g(x)) = g(2x + 1) = 2(2x + 1) + 1 = 4x + 3$. The domain is $(-\infty, \infty)$.

37. $f(x) = \dfrac{1}{\sqrt{x}}$ and $g(x) = x + 1$. The domain of f is $(0, \infty)$. The domain of g is $(-\infty, \infty)$.

(a) $(f \circ g)(x) = f(g(x)) = f(x + 1) = \dfrac{1}{\sqrt{x + 1}}$. We must have $x + 1 > 0$, or $x > -1$, so the domain is $(-1, \infty)$.

(b) $(g \circ f)(x) = g(f(x)) = g\left(\dfrac{1}{\sqrt{x}}\right) = \dfrac{1}{\sqrt{x}} + 1$. We must have $x > 0$, so the domain is $(0, \infty)$.

(c) $(f \circ f)(x) = f(f(x)) = f\left(\dfrac{1}{\sqrt{x}}\right) = \dfrac{1}{\sqrt{1/\sqrt{x}}} = \dfrac{1}{1/\sqrt{\sqrt{x}}} = \sqrt{\sqrt{x}} = \sqrt[4]{x}$. We must have $x > 0$, so the domain is $(0, \infty)$.

(d) $(g \circ g)(x) = g(g(x)) = g(x + 1) = (x + 1) + 1 = x + 2$. The domain is $(-\infty, \infty)$.

38. $f(x) = \dfrac{x}{x + 1}$ and $g(x) = 2x - 1$. The domain of f is $(-\infty, -1) \cup (-1, \infty)$. The domain of g is $(-\infty, \infty)$.

(a) $(f \circ g)(x) = f(g(x)) = f(2x - 1) = \dfrac{2x - 1}{(2x - 1) + 1} = \dfrac{2x - 1}{2x}$. We must have $2x \neq 0 \quad \Leftrightarrow \quad x \neq 0$. Thus, the domain is $(-\infty, 0) \cup (0, \infty)$.

(b) $(g \circ f)(x) = g(f(x)) = 2\left(\dfrac{x}{x + 1}\right) - 1 = \dfrac{2x}{x + 1} - 1 = \dfrac{2x - 1(x + 1)}{x + 1} = \dfrac{x - 1}{x + 1}$. We must have $x + 1 \neq 0 \quad \Leftrightarrow$ $x \neq -1$. Thus, the domain is $(-\infty, -1) \cup (-1, \infty)$.

(c) $(f \circ f)(x) = f(f(x)) = \dfrac{\dfrac{x}{x + 1}}{\dfrac{x}{x + 1} + 1} = \dfrac{\dfrac{x}{x + 1}}{\dfrac{x}{x + 1} + 1} \cdot \dfrac{x + 1}{x + 1} = \dfrac{x}{x + (x + 1)} = \dfrac{x}{2x + 1}$. We must have both $x + 1 \neq 0$ and $2x + 1 \neq 0$, so the domain excludes both -1 and $-\frac{1}{2}$. Thus, the domain is $(-\infty, -1) \cup \left(-1, -\frac{1}{2}\right) \cup \left(-\frac{1}{2}, \infty\right)$.

(d) $(g \circ g)(x) = g(g(x)) = g(2x - 1) = 2(2x - 1) - 1 = 4x - 3$. The domain is $(-\infty, \infty)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. $f(x) = \dfrac{2}{x}$ and $g(x) = \sin x$. The domain of f is $(-\infty, 0) \cup (0, \infty)$. The domain of g is $(-\infty, \infty)$.

(a) $(f \circ g)(x) = f(g(x)) = f(\sin x) = \dfrac{2}{\sin x} = 2\csc x$. We must have $\sin x \neq 0$, so the domain is $\{x \mid x \neq k\pi,\ k \text{ an integer}\}$.

(b) $(g \circ f)(x) = g(f(x)) = g\left(\dfrac{2}{x}\right) = \sin\left(\dfrac{2}{x}\right)$. We must have $x \neq 0$, so the domain is $(-\infty, 0) \cup (0, \infty)$.

(c) $(f \circ f)(x) = f(f(x)) = f\left(\dfrac{2}{x}\right) = \dfrac{2}{\dfrac{2}{x}} = x$. Since f requires $x \neq 0$, the domain is $(-\infty, 0) \cup (0, \infty)$.

(d) $(g \circ g)(x) = g(g(x)) = g(\sin x) = \sin(\sin x)$. The domain is $(-\infty, \infty)$.

40. $f(x) = \sqrt{5 - x}$ and $g(x) = \sqrt{x - 1}$. The domain of f is $(-\infty, 5]$ and the domain of g is $[1, \infty)$.

(a) $(f \circ g)(x) = f(g(x)) = f\left(\sqrt{x-1}\right) = \sqrt{5 - \sqrt{x-1}}$. We must have $x - 1 \geq 0 \quad \Leftrightarrow \quad x \geq 1$ and $5 - \sqrt{x-1} \geq 0 \quad \Leftrightarrow \quad \sqrt{x-1} \leq 5 \quad \Leftrightarrow \quad 0 \leq x - 1 \leq 25 \quad \Leftrightarrow \quad 1 \leq x \leq 26$. Thus, the domain is $[1, 26]$.

(b) $(g \circ f)(x) = g(f(x)) = g\left(\sqrt{5-x}\right) = \sqrt{\sqrt{5-x} - 1}$. We must have $5 - x \geq 0 \quad \Leftrightarrow \quad x \leq 5$ and $\sqrt{5-x} - 1 \geq 0 \quad \Leftrightarrow \quad \sqrt{5-x} \geq 1 \quad \Leftrightarrow \quad 5 - x \geq 1 \quad \Leftrightarrow \quad x \leq 4$. Intersecting the restrictions on x gives a domain of $(-\infty, 4]$.

(c) $(f \circ f)(x) = f(f(x)) = f\left(\sqrt{5-x}\right) = \sqrt{5 - \sqrt{5-x}}$. We must have $5 - x \geq 0 \quad \Leftrightarrow \quad x \leq 5$ and $5 - \sqrt{5-x} \geq 0 \quad \Leftrightarrow \quad \sqrt{5-x} \leq 5 \quad \Leftrightarrow \quad 0 \leq 5 - x \leq 25 \quad \Leftrightarrow \quad -5 \leq -x \leq 20 \quad \Leftrightarrow \quad -20 \leq x \leq 5$. Intersecting the restrictions on x gives a domain of $[-20, 5]$.

(d) $(g \circ g)(x) = g(g(x)) = g\left(\sqrt{x-1}\right) = \sqrt{\sqrt{x-1} - 1}$. We must have $x - 1 \geq 0 \quad \Leftrightarrow \quad x \geq 1$ and $\sqrt{x-1} - 1 \geq 0 \quad \Leftrightarrow \quad \sqrt{x-1} \geq 1 \quad \Leftrightarrow \quad x - 1 \geq 1 \quad \Leftrightarrow \quad x \geq 2$. Intersecting the restrictons on x gives a domain of $[2, \infty)$.

41. $(f \circ g \circ h)(x) = f(g(h(x))) = f(g(x^2)) = f(\sin(x^2)) = 3\sin(x^2) - 2$

42. $(f \circ g \circ h)(x) = f(g(h(x))) = f\left(g\left(\sqrt{x}\right)\right) = f\left(2^{\sqrt{x}}\right) = \left|2^{\sqrt{x}} - 4\right|$

43. $(f \circ g \circ h)(x) = f(g(h(x))) = f(g(x^3 + 2)) = f[(x^3 + 2)^2] = f(x^6 + 4x^3 + 4)$
$= \sqrt{(x^6 + 4x^3 + 4) - 3} = \sqrt{x^6 + 4x^3 + 1}$

44. $(f \circ g \circ h)(x) = f(g(h(x))) = f\left(g\left(\sqrt[3]{x}\right)\right) = f\left(\dfrac{\sqrt[3]{x}}{\sqrt[3]{x} - 1}\right) = \tan\left(\dfrac{\sqrt[3]{x}}{\sqrt[3]{x} - 1}\right)$

45. Let $g(x) = 2x + x^2$ and $f(x) = x^4$. Then $(f \circ g)(x) = f(g(x)) = f(2x + x^2) = (2x + x^2)^4 = F(x)$.

46. Let $g(x) = \cos x$ and $f(x) = x^2$. Then $(f \circ g)(x) = f(g(x)) = f(\cos x) = (\cos x)^2 = \cos^2 x = F(x)$.

47. Let $g(x) = \sqrt[3]{x}$ and $f(x) = \dfrac{x}{1 + x}$. Then $(f \circ g)(x) = f(g(x)) = f\left(\sqrt[3]{x}\right) = \dfrac{\sqrt[3]{x}}{1 + \sqrt[3]{x}} = F(x)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. Let $g(x) = \dfrac{x}{1+x}$ and $f(x) = \sqrt[3]{x}$. Then $(f \circ g)(x) = f(g(x)) = f\left(\dfrac{x}{1+x}\right) = \sqrt[3]{\dfrac{x}{1+x}} = G(x)$.

49. Let $g(t) = t^2$ and $f(t) = \sec t \, \tan t$. Then $(f \circ g)(t) = f(g(t)) = f(t^2) = \sec(t^2) \, \tan(t^2) = v(t)$.

50. Let $g(x) = \sqrt{x}$ and $f(x) = \sqrt{1+x}$. Then $(f \circ g)(x) = f(g(x)) = f\left(\sqrt{x}\right) = \sqrt{1+\sqrt{x}} = H(x)$.

51. Let $h(x) = \sqrt{x}$, $g(x) = x - 1$, and $f(x) = \sqrt{x}$. Then

$(f \circ g \circ h)(x) = f(g(h(x))) = f\left(g\left(\sqrt{x}\right)\right) = f\left(\sqrt{x} - 1\right) = \sqrt{\sqrt{x} - 1} = R(x)$.

52. Let $h(x) = |x|$, $g(x) = 2 + x$, and $f(x) = \sqrt[8]{x}$. Then

$(f \circ g \circ h)(x) = f(g(h(x))) = f(g(|x|)) = f(2 + |x|) = \sqrt[8]{2 + |x|} = H(x)$.

53. Let $h(t) = \cos t$, $g(t) = \sin t$, and $f(t) = t^2$. Then

$(f \circ g \circ h)(t) = f(g(h(t))) = f(g(\cos t)) = f(\sin(\cos t)) = [\sin(\cos t)]^2 = \sin^2(\cos t) = S(t)$.

54. Let $h(t) = \tan t$, $g(t) = \sqrt{t} + 1$, and $f(t) = \cos t$. Then

$(f \circ g \circ h)(t) = f(g(h(t))) = f(g(\tan t)) = f\left(\sqrt{\tan t} + 1\right) = \cos\left(\sqrt{\tan t} + 1\right) = H(t)$.

55. (a) $f(g(3)) = f(4) = 6$.

(b) $g(f(2)) = g(1) = 5$.

(c) $(f \circ g)(5) = f(g(5)) = f(3) = 5$.

(d) $(g \circ f)(5) = g(f(5)) = g(2) = 3$.

56. (a) $g(g(g(2))) = g(g(3)) = g(4) = 1$.

(b) $(f \circ f \circ f)(1) = f(f(f(1))) = f(f(3)) = f(5) = 2$.

(c) $(f \circ f \circ g)(1) = f(f(g(1))) = f(f(5)) = f(2) = 1$.

(d) $(g \circ f \circ g)(3) = g(f(g(3))) = g(f(4)) = g(6) = 2$.

57. (a) $g(2) = 5$, because the point $(2, 5)$ is on the graph of g. Thus, $f(g(2)) = f(5) = 4$, because the point $(5, 4)$ is on the graph of f.

(b) $g(f(0)) = g(0) = 3$

(c) $(f \circ g)(0) = f(g(0)) = f(3) = 0$

(d) $(g \circ f)(6) = g(f(6)) = g(6)$. This value is not defined, because there is no point on the graph of g that has x-coordinate 6.

(e) $(g \circ g)(-2) = g(g(-2)) = g(1) = 4$

(f) $(f \circ f)(4) = f(f(4)) = f(2) = -2$

58. To find a particular value of $f(g(x))$, say for $x = 0$, we note from the graph that $g(0) \approx 2.8$ and $f(2.8) \approx -0.5$. Thus, $f(g(0)) \approx f(2.8) \approx -0.5$. The other values listed in the table were obtained in a similar fashion.

x	$g(x)$	$f(g(x))$
-5	-0.2	-4
-4	1.2	-3.3
-3	2.2	-1.7
-2	2.8	-0.5
-1	3	-0.2

x	$g(x)$	$f(g(x))$
0	2.8	-0.5
1	2.2	-1.7
2	1.2	-3.3
3	-0.2	-4
4	-1.9	-2.2
5	-4.1	1.9

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. (a) Using the relationship *distance* = *rate* · *time* with the radius r as the distance, we have $r(t) = 60t$.

(b) $A = \pi r^2 \quad \Rightarrow \quad (A \circ r)(t) = A(r(t)) = \pi(60t)^2 = 3600\pi t^2$. This formula gives us the extent of the rippled area (in cm^2) at any time t.

60. (a) The radius r of the balloon is increasing at a rate of $2\ \text{cm/s}$, so $r(t) = (2\ \text{cm/s})(t\ \text{s}) = 2t$ (in cm).

(b) Using $V = \frac{4}{3}\pi r^3$, we get $(V \circ r)(t) = V(r(t)) = V(2t) = \frac{4}{3}\pi(2t)^3 = \frac{32}{3}\pi t^3$.

The result, $V = \frac{32}{3}\pi t^3$, gives the volume of the balloon (in cm^3) as a function of time (in s).

61. (a) From the figure, we have a right triangle with legs 6 and d, and hypotenuse s.

By the Pythagorean Theorem, $d^2 + 6^2 = s^2 \quad \Rightarrow \quad s = f(d) = \sqrt{d^2 + 36}$.

(b) Using $d = rt$, we get $d = (30\ \text{km/h})(t\ \text{hours}) = 30t$ (in km). Thus, $d = g(t) = 30t$.

(c) $(f \circ g)(t) = f(g(t)) = f(30t) = \sqrt{(30t)^2 + 36} = \sqrt{900t^2 + 36}$. This function represents the distance between the lighthouse and the ship as a function of the time elapsed since noon.

62. (a) $d = rt \quad \Rightarrow \quad d(t) = 350t$

(b) There is a Pythagorean relationship involving the legs with lengths d and 1 and the hypotenuse with length s: $d^2 + 1^2 = s^2$. Thus, $s(d) = \sqrt{d^2 + 1}$.

(c) $(s \circ d)(t) = s(d(t)) = s(350t) = \sqrt{(350t)^2 + 1}$

63. (a)

$$H(t) = \begin{cases} 0 & \text{if } t < 0 \\ 1 & \text{if } t \geq 0 \end{cases}$$

(b)

$$V(t) = \begin{cases} 0 & \text{if } t < 0 \\ 120 & \text{if } t \geq 0 \end{cases} \quad \text{so } V(t) = 120H(t).$$

(c)

Starting with the formula in part (b), we replace 120 with 240 to reflect the different voltage. Also, because we are starting 5 units to the right of $t = 0$, we replace t with $t - 5$. Thus, the formula is $V(t) = 240H(t - 5)$.

64. (a) $R(t) = tH(t)$

$$= \begin{cases} 0 & \text{if } t < 0 \\ t & \text{if } t \geq 0 \end{cases}$$

(b) $V(t) = \begin{cases} 0 & \text{if } t < 0 \\ 2t & \text{if } 0 \leq t \leq 60 \end{cases}$

so $V(t) = 2tH(t)$, $t \leq 60$.

(c) $V(t) = \begin{cases} 0 & \text{if } t < 7 \\ 4(t-7) & \text{if } 7 \leq t \leq 32 \end{cases}$

so $V(t) = 4(t - 7)H(t - 7)$, $t \leq 32$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

65. If $f(x) = m_1x + b_1$ and $g(x) = m_2x + b_2$, then

$(f \circ g)(x) = f(g(x)) = f(m_2x + b_2) = m_1(m_2x + b_2) + b_1 = m_1m_2x + m_1b_2 + b_1$.

So $f \circ g$ is a linear function with slope m_1m_2.

66. If $A(x) = 1.04x$, then

$(A \circ A)(x) = A(A(x)) = A(1.04x) = 1.04(1.04x) = (1.04)^2x$,

$(A \circ A \circ A)(x) = A((A \circ A)(x)) = A((1.04)^2x) = 1.04(1.04)^2x = (1.04)^3x$, and

$(A \circ A \circ A \circ A)(x) = A((A \circ A \circ A)(x)) = A((1.04)^3x) = 1.04(1.04)^3x, = (1.04)^4x$.

These compositions represent the amount of the investment after 2, 3, and 4 years.

Based on this pattern, when we compose n copies of A, we get the formula $\underbrace{(A \circ A \circ \cdots \circ A)}_{n\ A\text{'s}}(x) = (1.04)^nx$.

67. (a) By examining the variable terms in g and h, we deduce that we must square g to get the terms $4x^2$ and $4x$ in h. If we let $f(x) = x^2 + c$, then $(f \circ g)(x) = f(g(x)) = f(2x + 1) = (2x + 1)^2 + c = 4x^2 + 4x + (1 + c)$. Since $h(x) = 4x^2 + 4x + 7$, we must have $1 + c = 7$. So $c = 6$ and $f(x) = x^2 + 6$.

(b) We need a function g so that $f(g(x)) = 3(g(x)) + 5 = h(x)$. But $h(x) = 3x^2 + 3x + 2 = 3(x^2 + x) + 2 = 3(x^2 + x - 1) + 5$, so we see that $g(x) = x^2 + x - 1$.

68. We need a function g so that $g(f(x)) = g(x + 4) = h(x) = 4x - 1 = 4(x + 4) - 17$. So we see that the function g must be $g(x) = 4x - 17$.

69. We need to examine $h(-x)$.

$$h(-x) = (f \circ g)(-x) = f(g(-x)) = f(g(x)) \quad \text{[because } g \text{ is even]} \quad = h(x)$$

Because $h(-x) = h(x)$, h is an even function.

70. $h(-x) = f(g(-x)) = f(-g(x))$. At this point, we can't simplify the expression, so we might try to find a counterexample to show that h is not an odd function. Let $g(x) = x$, an odd function, and $f(x) = x^2 + x$. Then $h(x) = x^2 + x$, which is neither even nor odd.

Now suppose f is an odd function. Then $f(-g(x)) = -f(g(x)) = -h(x)$. Hence, $h(-x) = -h(x)$, and so h is odd if both f and g are odd.

Now suppose f is an even function. Then $f(-g(x)) = f(g(x)) = h(x)$. Hence, $h(-x) = h(x)$, and so h is even if g is odd and f is even.

71. (a) $E(x) = f(x) + f(-x) \quad \Rightarrow \quad E(-x) = f(-x) + f(-(-x)) = f(-x) + f(x) = E(x)$. Since $E(-x) = E(x)$, E is an even function.

(b) $O(x) = f(x) - f(-x) \quad \Rightarrow \quad O(-x) = f(-x) - f(-(-x)) = f(-x) - f(x) = -[f(x) - f(-x)] = -O(x)$.

Since $O(-x) = -O(x)$, O is an odd function.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) For any function f with domain $\mathbb{R}$, define functions E and O as in parts (a) and (b). Then $\frac{1}{2}E$ is even, $\frac{1}{2}O$ is odd, and we show that $f(x) = \frac{1}{2}E(x) + \frac{1}{2}O(x)$:

$$\begin{aligned}\tfrac{1}{2}E(x) + \tfrac{1}{2}O(x) &= \tfrac{1}{2}\left[f(x) + f(-x)\right] + \tfrac{1}{2}\left[f(x) - f(-x)\right] \\ &= \tfrac{1}{2}[f(x) + f(-x) + f(x) - f(-x)] \\ &= \tfrac{1}{2}[2f(x)] = f(x)\end{aligned}$$

as desired.

(d) $f(x) = 2^x + (x-3)^2$ has domain $\mathbb{R}$, so we know from part (c) that $f(x) = \frac{1}{2}E(x) + \frac{1}{2}O(x)$, where

$$\begin{aligned}E(x) = f(x) + f(-x) &= 2^x + (x-3)^2 + 2^{-x} + (-x-3)^2 \\ &= 2^x + 2^{-x} + (x-3)^2 + (x+3)^2\end{aligned}$$

and

$$\begin{aligned}O(x) = f(x) - f(-x) &= 2^x + (x-3)^2 - [2^{-x} + (-x-3)^2] \\ &= 2^x - 2^{-x} + (x-3)^2 - (x+3)^2\end{aligned}$$

1.4 Exponential Functions

1. (a) $\dfrac{-2^6}{4^3} = \dfrac{-2^6}{(2^2)^3} = -\dfrac{2^6}{2^6} = -1$

(b) $\dfrac{(-3)^6}{9^6} = \left(\dfrac{-3}{9}\right)^6 = \left(-\dfrac{1}{3}\right)^6 = \dfrac{1}{3^6}$

(c) $\dfrac{1}{\sqrt[4]{x^5}} = \dfrac{1}{\sqrt[4]{x^4 \cdot x}} = \dfrac{1}{x\sqrt[4]{x}}$

(d) $\dfrac{x^3 \cdot x^n}{x^{n+1}} = \dfrac{x^{3+n}}{x^{n+1}} = x^{(3+n)-(n+1)} = x^2$

(e) $b^3(3b^{-1})^{-2} = b^3 3^{-2}(b^{-1})^{-2} = \dfrac{b^3 \cdot b^2}{3^2} = \dfrac{b^5}{9}$

(f) $\dfrac{2x^2y}{(3x^{-2}y)^2} = \dfrac{2x^2y}{3^2(x^{-2})^2y^2} = \dfrac{2x^2y}{9x^{-4}y^2} = \dfrac{2}{9}x^{2-(-4)}y^{1-2} = \dfrac{2}{9}x^6y^{-1} = \dfrac{2x^6}{9y}$

2. (a) $\dfrac{\sqrt[3]{4}}{\sqrt[3]{108}} = \dfrac{\sqrt[3]{4}}{\sqrt[3]{4 \cdot 27}} = \dfrac{\sqrt[3]{4}}{\sqrt[3]{4} \cdot \sqrt[3]{27}} = \dfrac{1}{\sqrt[3]{27}} = \dfrac{1}{3}$

(b) $27^{2/3} = (27^{1/3})^2 = \left(\sqrt[3]{27}\right)^2 = 3^2 = 9$

(c) $2x^2(3x^5)^2 = 2x^2 \cdot 3^2(x^5)^2 = 2x^2 \cdot 9x^{10} = 2 \cdot 9x^{2+10} = 18x^{12}$

(d) $(2x^{-2})^{-3}x^{-3} = 2^{-3}(x^{-2})^{-3}x^{-3} = \dfrac{x^6 \cdot x^{-3}}{2^3} = \dfrac{x^{6+(-3)}}{8} = \dfrac{x^3}{8}$

(e) $\dfrac{3a^{3/2} \cdot a^{1/2}}{a^{-1}} = 3a^{3/2+1/2} \cdot a^1 = 3a^2 \cdot a = 3a^3$

(f) $\dfrac{\sqrt{a\sqrt{b}}}{\sqrt[3]{ab}} = \dfrac{(ab^{1/2})^{1/2}}{(ab)^{1/3}} = \dfrac{a^{1/2}\,(b^{1/2})^{1/2}}{a^{1/3}\,b^{1/3}} = \dfrac{a^{1/2}\,b^{1/4}}{a^{1/3}\,b^{1/3}} = a^{1/2-1/3}\,b^{1/4-1/3} = a^{1/6}\,b^{-1/12} = \dfrac{a^{1/6}}{b^{1/12}} = \dfrac{\sqrt[6]{a}}{\sqrt[12]{b}}$

3. (a) $f(x) = b^x, \;\; b > 0$ (b) $\mathbb{R}$ (c) $(0, \infty)$ (d) See Figures 4(c), 4(b), and 4(a), respectively.

4. (a) The number e is the value of a such that the slope of the tangent line at $x = 0$ on the graph of $y = a^x$ is exactly 1.

(b) $e \approx 2.71828$ (c) $f(x) = e^x$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. All of these graphs approach 0 as $x \to -\infty$, all of them pass through the point $(0, 1)$, and all of them are increasing and approach ∞ as $x \to \infty$. The larger the base, the faster the function increases for $x > 0$, and the faster it approaches 0 as $x \to -\infty$.

Note: The notation "$x \to \infty$" can be thought of as "x becomes large" at this point. More details on this notation are given in Chapter 2.

6. The graph of e^{-x} is the reflection of the graph of e^x about the y-axis, and the graph of 8^{-x} is the reflection of that of 8^x about the y-axis. The graph of 8^x increases more quickly than that of e^x for $x > 0$, and approaches 0 faster as $x \to -\infty$.

7. The functions with base greater than 1 (3^x and 10^x) are increasing, while those with base less than 1 $\left[\left(\frac{1}{3}\right)^x \text{ and } \left(\frac{1}{10}\right)^x\right]$ are decreasing. The graph of $\left(\frac{1}{3}\right)^x$ is the reflection of that of 3^x about the y-axis, and the graph of $\left(\frac{1}{10}\right)^x$ is the reflection of that of 10^x about the y-axis. The graph of 10^x increases more quickly than that of 3^x for $x > 0$, and approaches 0 faster as $x \to -\infty$.

8. Each of the graphs approaches ∞ as $x \to -\infty$, and each approaches 0 as $x \to \infty$. The smaller the base, the faster the function grows as $x \to -\infty$, and the faster it approaches 0 as $x \to \infty$.

9. We start with the graph of $y = 3^x$ (Figure 15) and shift 1 unit upward to get the graph of $g(x) = 3^x + 1$.

10. We start with the graph of $y = \left(\frac{1}{2}\right)^x$ (Figure 3) and stretch vertically by a factor of 2 to obtain the graph of $y = 2\left(\frac{1}{2}\right)^x$. Then we shift the graph 3 units downward to get the graph of $h(x) = 2\left(\frac{1}{2}\right)^x - 3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. We start with the graph of $y = e^x$ (Figure 15) and reflect about the y-axis to get the graph of $y = e^{-x}$. Then we reflect the graph about the x-axis to get the graph of $y = -e^{-x}$.

12. We start with the graph of $y = 4^x$ (Figure 3) and shift 2 units to the left to get the graph of $y = 4^{x+2}$.

13. We start with the graph of $y = e^x$ (Figure 15) and reflect about the y-axis to get the graph of $y = e^{-x}$. Then we compress the graph vertically by a factor of 2 to obtain the graph of $y = \frac{1}{2}e^{-x}$ and then reflect about the x-axis to get the graph of $y = -\frac{1}{2}e^{-x}$. Finally, we shift the graph one unit upward to get the graph of $y = 1 - \frac{1}{2}e^{-x}$.

14. We start with the graph of $y = e^x$ (Figure 15) and reflect the portion of the graph in the first quadrant about the y-axis to obtain the graph of $y = e^{|x|}$.

15. (a) To find the equation of the graph that results from shifting the graph of $y = e^x$ two units downward, we subtract 2 from the original function to get $y = e^x - 2$.

(b) To find the equation of the graph that results from shifting the graph of $y = e^x$ two units to the right, we replace x with $x - 2$ in the original function to get $y = e^{x-2}$.

(c) To find the equation of the graph that results from reflecting the graph of $y = e^x$ about the x-axis, we multiply the original function by -1 to get $y = -e^x$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) To find the equation of the graph that results from reflecting the graph of $y = e^x$ about the y-axis, we replace x with $-x$ in the original function to get $y = e^{-x}$.

(e) To find the equation of the graph that results from reflecting the graph of $y = e^x$ about the x-axis and then about the y-axis, we first multiply the original function by -1 (to get $y = -e^x$) and then replace x with $-x$ in this equation to get $y = -e^{-x}$.

16. (a) This reflection consists of first reflecting the graph about the x-axis (giving the graph with equation $y = -e^x$) and then shifting this graph $2 \cdot 4 = 8$ units upward. So the equation is $y = -e^x + 8$.

(b) This reflection consists of first reflecting the graph about the y-axis (giving the graph with equation $y = e^{-x}$) and then shifting this graph $2 \cdot 2 = 4$ units to the right. So the equation is $y = e^{-(x-4)}$.

17. (a) The denominator is zero when $1 - e^{1-x^2} = 0 \quad \Leftrightarrow \quad e^{1-x^2} = 1 \quad \Leftrightarrow \quad 1 - x^2 = 0 \quad \Leftrightarrow \quad x = \pm 1$. Thus, the function $f(x) = \dfrac{1 - e^{x^2}}{1 - e^{1-x^2}}$ has domain $\{x \mid x \neq \pm 1\} = (-\infty, -1) \cup (-1, 1) \cup (1, \infty)$.

(b) The denominator is never equal to zero, so the function $f(x) = \dfrac{1 + x}{e^{\cos x}}$ has domain $\mathbb{R}$, or $(-\infty, \infty)$.

18. (a) The function $g(t) = \sqrt{10^t - 100}$ has domain $\left\{t \mid 10^t - 100 \geq 0\right\} = \left\{t \mid 10^t \geq 10^2\right\} = \{t \mid t \geq 2\} = [2, \infty)$.

(b) The sine and exponential functions have domain $\mathbb{R}$, so $g(t) = \sin(e^t - 1)$ also has domain $\mathbb{R}$.

19. Use $y = Cb^x$ with the points $(1, 6)$ and $(3, 24)$. $\quad 6 = Cb^1 \quad \left[C = \frac{6}{b}\right] \quad$ and $24 = Cb^3 \quad \Rightarrow \quad 24 = \left(\dfrac{6}{b}\right) b^3 \quad \Rightarrow$ $4 = b^2 \quad \Rightarrow \quad b = 2 \quad$ [since $b > 0$] $\quad$ and $C = \frac{6}{2} = 3$. The function is $f(x) = 3 \cdot 2^x$.

20. Use $y = Cb^x$ with the points $(-1, 3)$ and $\left(1, \frac{4}{3}\right)$. From the point $(-1, 3)$, we have $3 = Cb^{-1}$, hence $C = 3b$. Using this and the point $\left(1, \frac{4}{3}\right)$, we get $\frac{4}{3} = Cb^1 \quad \Rightarrow \quad \frac{4}{3} = (3b)b \quad \Rightarrow \quad \frac{4}{9} = b^2 \quad \Rightarrow \quad b = \frac{2}{3} \quad$ [since $b > 0$] $\quad$ and $C = 3(\frac{2}{3}) = 2$. The function is $f(x) = 2(\frac{2}{3})^x$.

21. If $f(x) = 5^x$, then $\dfrac{f(x+h) - f(x)}{h} = \dfrac{5^{x+h} - 5^x}{h} = \dfrac{5^x 5^h - 5^x}{h} = \dfrac{5^x\left(5^h - 1\right)}{h} = 5^x\left(\dfrac{5^h - 1}{h}\right)$.

22. Suppose the month is February. Your payment on the 28th day would be $2^{28-1} = 2^{27} = 134{,}217{,}728$ cents, or \$1,342,177.28. Clearly, the second method of payment results in a larger amount for any month.

23. 2 ft $= 24$ in, $f(24) = 24^2$ in $= 576$ in $= 48$ ft. $\quad g(24) = 2^{24}$ in $= 2^{24}/(12 \cdot 5280)$ mi ≈ 265 mi

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. We see from the graphs that for x less than about 1.8, $g(x) = 5^x > f(x) = x^5$, and then near the point $(1.8, 17.1)$ the curves intersect. Then $f(x) > g(x)$ from $x \approx 1.8$ until $x = 5$. At $(5, 3125)$ there is another point of intersection, and for $x > 5$ we see that $g(x) > f(x)$. In fact, g increases much more rapidly than f beyond that point.

25. The graph of g finally surpasses that of f at $x \approx 35.8$.

26. We graph $y = e^x$ and $y = 1{,}000{,}000{,}000$ and determine where $e^x = 1 \times 10^9$. This seems to be true at $x \approx 20.723$, so $e^x > 1 \times 10^9$ for $x > 20.723$.

27. (a)

(b) Using a graphing calculator, we obtain the exponential curve $f(t) = 36.89301(1.06614)^t$.

(c) Using the TRACE and zooming in, we find that the bacteria count doubles from 37 to 74 in about 10.87 hours.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. Let $t = 0$ correspond to 1900 to get the model $P = ab^t$, where $a \approx 80.8498$ and $b \approx 1.01269$. To estimate the population in 1925, let $t = 25$ to obtain $P \approx 111$ million. To predict the population in 2020, let $t = 120$ to obtain $P \approx 367$ million.

29. (a) Three hours represents 6 doubling periods (one doubling period is 30 minutes). Thus, $500 \cdot 2^6 = 32{,}000$.

(b) In t hours, there will be $2t$ doubling periods. The initial population is 500, so the population y at time t is $y = 500 \cdot 2^{2t}$.

(c) $t = \frac{40}{60} = \frac{2}{3} \quad \Rightarrow \quad y = 500 \cdot 2^{2(2/3)} \approx 1260$

(d) We graph $y_1 = 500 \cdot 2^{2t}$ and $y_2 = 100{,}000$. The two curves intersect at $t \approx 3.82$, so the population reaches 100,000 in about 3.82 hours.

30. (a) Let a be the initial population. Since 18 years is 3 doubling periods, $a \cdot 2^3 = 600 \quad \Rightarrow \quad a = \dfrac{600}{8} = 75$. The initial squirrel popluation was 75.

(b) A period of t years corresponds to $t/6$ doubling periods, so the expected squirrel population t years after introduction is $P = 75 \cdot 2^{t/6}$.

(c) Ten years from now will be $18 + 10 = 28$ years from introduction. The population is estimated to be $P = 75 \cdot 2^{28/6} \approx 1905$ squirrels.

31. Half of 76.0 RNA copies per mL, corresponding to $t = 1$, is 38.0 RNA copies per mL. Using the graph of V in Figure 11, we estimate that it takes about 3.5 additional days for the patient's viral load to decrease to 38 RNA copies per mL.

32. (a) The exponential decay model has the form $C(t) = a\left(\frac{1}{2}\right)^{t/1.5}$, where t is the number of hours after midnight and $C(t)$ is the BAC. We are given that $C(0) = 0.14$, so $a = 0.14$, and the model is $C(t) = 0.14\left(\frac{1}{2}\right)^{t/1.5}$.

(b) From the graph, we estimate that the BAC is 0.08 g/dL when $t \approx 1.2$ hours.

33.

From the graph, it appears that f is an odd function (f is undefined for $x = 0$). To prove this, we must show that $f(-x) = -f(x)$.

$$f(-x) = \frac{1 - e^{1/(-x)}}{1 + e^{1/(-x)}} = \frac{1 - e^{(-1/x)}}{1 + e^{(-1/x)}} = \frac{1 - \dfrac{1}{e^{1/x}}}{1 + \dfrac{1}{e^{1/x}}} \cdot \frac{e^{1/x}}{e^{1/x}} = \frac{e^{1/x} - 1}{e^{1/x} + 1}$$

$$= -\frac{1 - e^{1/x}}{1 + e^{1/x}} = -f(x)$$

so f is an odd function.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. We'll start with $b = -1$ and graph $f(x) = \dfrac{1}{1 + ae^{bx}}$ for $a = 0.1$, 1, and 5.

From the graph, we see that there is a horizontal asymptote $y = 0$ as $x \to -\infty$ and a horizontal asymptote $y = 1$ as $x \to \infty$. If $a = 1$, the y-intercept is $\left(0, \frac{1}{2}\right)$. As a gets smaller (close to 0), the graph of f moves left. As a gets larger, the graph of f moves right.

As b changes from -1 to 0, the graph of f is stretched horizontally. As b changes through large negative values, the graph of f is compressed horizontally. (This takes care of negatives values of b.)

If b is positive, the graph of f is reflected through the y-axis.

Last, if $b = 0$, the graph of f is the horizontal line $y = 1/(1 + a)$.

35. We graph the function $f(x) = \dfrac{a}{2}(e^{x/a} + e^{-x/a})$ for $a = 1$, 2, and 5. Because $f(0) = a$, the y-intercept is a, so the y-intercept moves upward as a increases. Notice that the graph also widens, becoming flatter near the y-axis as a increases.

1.5 Inverse Functions and Logarithms

1. (a) See Definition 1.

(b) It must pass the Horizontal Line Test.

2. (a) $f^{-1}(y) = x \quad \Leftrightarrow \quad f(x) = y$ for any y in B. The domain of f^{-1} is B and the range of f^{-1} is A.

(b) See the steps in Box 5.

(c) Reflect the graph of f about the line $y = x$.

3. f is not one-to-one because $2 \neq 6$, but $f(2) = 2.0 = f(6)$.

4. f is one-to-one because it never takes on the same value twice.

5. We could draw a horizontal line that intersects the graph in more than one point. Thus, by the Horizontal Line Test, the function is not one-to-one.

6. No horizontal line intersects the graph more than once. Thus, by the Horizontal Line Test, the function is one-to-one.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. No horizontal line intersects the graph more than once. Thus, by the Horizontal Line Test, the function is one-to-one.

8. We could draw a horizontal line that intersects the graph in more than one point. Thus, by the Horizontal Line Test, the function is not one-to-one.

9. The graph of $f(x) = 2x - 3$ is a line with slope 2. It passes the Horizontal Line Test, so f is one-to-one.
Algebraic solution: If $x_1 \neq x_2$, then $2x_1 \neq 2x_2 \quad \Rightarrow \quad 2x_1 - 3 \neq 2x_2 - 3 \quad \Rightarrow \quad f(x_1) \neq f(x_2)$, so f is one-to-one.

10. The graph of $f(x) = x^4 - 16$ is symmetric with respect to the y-axis. Pick any x-values equidistant from 0 to find two equal function values. For example, $f(-1) = -15$ and $f(1) = -15$, so f is not one-to-one.

11. No horizontal line intersects the graph of $r(t) = t^3 + 4$ more than once. Thus, by the Horizontal Line Test, the function is one-to-one.
Algebraic solution: If $t_1 \neq t_2$, then $t_1^3 \neq t_2^3 \quad \Rightarrow \quad t_1^3 + 4 \neq t_2^3 + 4 \quad \Rightarrow \quad r(t_1) \neq r(t_2)$, so r is one-to-one.

12. The graph of $g(x) = \sqrt[3]{x}$ passes the Horizontal Line Test, so g is one-to-one.

13. $g(x) = 1 - \sin x. \quad g(0) = 1$ and $g(\pi) = 1$, so g is not one-to-one.

14. The graph of $f(x) = x^4 - 1$ passes the Horizontal Line Test when x is restricted to the interval [0,10], so f is one-to-one.

15. A football will attain every height h up to its maximum height twice: once on the way up, and again on the way down. Thus, even if t_1 does not equal t_2, $f(t_1)$ may equal $f(t_2)$, so f is not 1-1.

16. f is not 1-1 because eventually we all stop growing and therefore, there are two times at which we have the same height.

17. (a) Since f is 1-1, $f(6) = 17 \quad \Leftrightarrow \quad f^{-1}(17) = 6$.
(b) Since f is 1-1, $f^{-1}(3) = 2 \quad \Leftrightarrow \quad f(2) = 3$.

18. First, we must determine x such that $f(x) = 3$. By inspection, we see that if $x = 1$, then $f(1) = 3$. Since f is 1-1 (f is an increasing function), it has an inverse, and $f^{-1}(3) = 1$. If f is a 1-1 function, then $f(f^{-1}(a)) = a$, so $f(f^{-1}(2)) = 2$.

19. First, we must determine x such that $g(x) = 4$. By inspection, we see that if $x = 0$, then $g(x) = 4$. Since g is 1-1 (g is an increasing function), it has an inverse, and $g^{-1}(4) = 0$.

20. (a) f is 1-1 because it passes the Horizontal Line Test.
(b) Domain of $f = [-3, 3] =$ Range of f^{-1}. Range of $f = [-1, 3] =$ Domain of f^{-1}.
(c) Since $f(0) = 2$, $f^{-1}(2) = 0$.
(d) Since $f(-1.7) \approx 0$, $f^{-1}(0) \approx -1.7$.

21. We solve $C = \frac{5}{9}(F - 32)$ for F: $\frac{9}{5}C = F - 32 \quad \Rightarrow \quad F = \frac{9}{5}C + 32$. This gives us a formula for the inverse function, that is, the Fahrenheit temperature F as a function of the Celsius temperature C. $F \geq -459.67 \quad \Rightarrow \quad \frac{9}{5}C + 32 \geq -459.67 \quad \Rightarrow \quad \frac{9}{5}C \geq -491.67 \quad \Rightarrow \quad C \geq -273.15$, the domain of the inverse function.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $m = \dfrac{m_0}{\sqrt{1 - v^2/c^2}} \quad \Rightarrow \quad 1 - \dfrac{v^2}{c^2} = \dfrac{m_0^2}{m^2} \quad \Rightarrow \quad \dfrac{v^2}{c^2} = 1 - \dfrac{m_0^2}{m^2} \quad \Rightarrow \quad v^2 = c^2\left(1 - \dfrac{m_0^2}{m^2}\right) \quad \Rightarrow \quad v = c\sqrt{1 - \dfrac{m_0^2}{m^2}}$.

This formula gives us the speed v of the particle in terms of its mass m, that is, $v = f^{-1}(m)$.

23. First note that $f(x) = 1 - x^2$, $x \geq 0$, is one-to-one. We first write $y = 1 - x^2$, $x \geq 0$, and solve for x: $x^2 = 1 - y \quad \Rightarrow \quad x = \sqrt{1 - y}$ (since $x \geq 0$). Interchanging x and y gives $y = \sqrt{1 - x}$, so the inverse function is $f^{-1}(x) = \sqrt{1 - x}$.

24. Completing the square, we have $g(x) = x^2 - 2x = (x^2 - 2x + 1) - 1 = (x - 1)^2 - 1$ and, with the restriction $x \geq 1$, g is one-to-one. We write $y = (x - 1)^2 - 1$, $x \geq 1$, and solve for x: $x - 1 = \sqrt{y + 1}$ (since $x \geq 1 \quad \Leftrightarrow \quad x - 1 \geq 0$), so $x = 1 + \sqrt{y + 1}$. Interchanging x and y gives $y = 1 + \sqrt{x + 1}$, so $g^{-1}(x) = 1 + \sqrt{x + 1}$.

25. First write $y = g(x) = 2 + \sqrt{x + 1}$ and note that $y \geq 2$. Solve for x: $y - 2 = \sqrt{x + 1} \quad \Rightarrow \quad (y - 2)^2 = x + 1 \quad \Rightarrow$ $x = (y - 2)^2 - 1$ $(y \geq 2)$. Interchanging x and y gives $y = (x - 2)^2 - 1$, so $g^{-1}(x) = (x - 2)^2 - 1$ with domain $x \geq 2$.

26. We write $y = h(x) = \dfrac{6 - 3x}{5x + 7}$ and solve for x: $y(5x + 7) = 6 - 3x \quad \Rightarrow \quad 5xy + 7y = 6 - 3x \quad \Rightarrow$ $5xy + 3x = 6 - 7y \quad \Rightarrow \quad x(5y + 3) = 6 - 7y \quad \Rightarrow \quad x = \dfrac{6 - 7y}{5y + 3}$. Interchanging x and y gives $y = \dfrac{6 - 7x}{5x + 3}$, so $h^{-1}(x) = \dfrac{6 - 7x}{5x + 3}$.

27. We solve $y = e^{1-x}$ for x: $\ln y = \ln e^{1-x} \quad \Rightarrow \quad \ln y = 1 - x \quad \Rightarrow \quad x = 1 - \ln y$. Interchanging x and y gives the inverse function $y = 1 - \ln x$.

28. We solve $y = 3\ln(x - 2)$ for x: $y/3 = \ln(x - 2) \quad \Rightarrow \quad e^{y/3} = x - 2 \quad \Rightarrow \quad x = 2 + e^{y/3}$. Interchanging x and y gives the inverse function $y = 2 + e^{x/3}$.

29. We solve $y = \left(2 + \sqrt[3]{x}\right)^5$ for x: $\sqrt[5]{y} = 2 + \sqrt[3]{x} \quad \Rightarrow \quad \sqrt[3]{x} = \sqrt[5]{y} - 2 \quad \Rightarrow \quad x = \left(\sqrt[5]{y} - 2\right)^3$. Interchanging x and y gives the inverse function $y = \left(\sqrt[5]{x} - 2\right)^3$.

30. We solve $y = \dfrac{1 - e^{-x}}{1 + e^{-x}}$ for x: $y(1 + e^{-x}) = 1 - e^{-x} \quad \Rightarrow \quad y + ye^{-x} = 1 - e^{-x} \quad \Rightarrow \quad e^{-x} + ye^{-x} = 1 - y \quad \Rightarrow$ $e^{-x}(1 + y) = 1 - y \quad \Rightarrow \quad e^{-x} = \dfrac{1 - y}{1 + y} \quad \Rightarrow \quad -x = \ln\dfrac{1 - y}{1 + y} \quad \Rightarrow \quad x = -\ln\dfrac{1 - y}{1 + y}$ or, equivalently, $x = \ln\left(\dfrac{1 - y}{1 + y}\right)^{-1} = \ln\dfrac{1 + y}{1 - y}$. Interchanging x and y gives the inverse function $y = \ln\dfrac{1 + x}{1 - x}$.

31. $y = f(x) = \sqrt{4x + 3}$ $(y \geq 0) \quad \Rightarrow \quad y^2 = 4x + 3 \quad \Rightarrow \quad x = \dfrac{y^2 - 3}{4}$.

Interchange x and y: $y = \dfrac{x^2 - 3}{4}$. So $f^{-1}(x) = \dfrac{x^2 - 3}{4}$ $(x \geq 0)$. From the graph, we see that f and f^{-1} are reflections about the line $y = x$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. $y = f(x) = 1 + e^{-x} \quad \Rightarrow \quad e^{-x} = y - 1 \quad \Rightarrow \quad -x = \ln(y-1) \quad \Rightarrow$ $x = -\ln(y-1)$. Interchange x and y: $y = -\ln(x-1)$.

So $f^{-1}(x) = -\ln(x-1)$. From the graph, we see that f and f^{-1} are reflections about the line $y = x$.

33. Reflect the graph of f about the line $y = x$. The points $(-1,-2)$, $(1,-1)$, $(2,2)$, and $(3,3)$ on f are reflected to $(-2,-1)$, $(-1,1)$, $(2,2)$, and $(3,3)$ on f^{-1}.

34. Reflect the graph of f about the line $y = x$.

35. (a) $y = f(x) = \sqrt{1-x^2} \quad (0 \le x \le 1$ and note that $y \ge 0) \quad \Rightarrow$ $y^2 = 1 - x^2 \quad \Rightarrow \quad x^2 = 1 - y^2 \quad \Rightarrow \quad x = \sqrt{1-y^2}$. So $f^{-1}(x) = \sqrt{1-x^2},\ 0 \le x \le 1$. We see that $f^{-1}$and f are the same function.

(b) The graph of f is the portion of the circle $x^2 + y^2 = 1$ with $0 \le x \le 1$ and $0 \le y \le 1$ (quarter-circle in the first quadrant). The graph of f is symmetric with respect to the line $y = x$, so its reflection about $y = x$ is itself, that is, $f^{-1} = f$.

36. (a) $y = g(x) = \sqrt[3]{1-x^3} \quad \Rightarrow \quad y^3 = 1 - x^3 \quad \Rightarrow \quad x^3 = 1 - y^3 \quad \Rightarrow$ $x = \sqrt[3]{1-y^3}$. So $g^{-1}(x) = \sqrt[3]{1-x^3}$. We see that g and g^{-1} are the same function.

(b) The graph of g is symmetric with respect to the line $y = x$, so its reflection about $y = x$ is itself, that is, $g^{-1} = g$.

37. (a) It is defined as the inverse of the exponential function with base b, that is, $\log_b x = y \quad \Leftrightarrow \quad b^y = x$.
(b) $(0, \infty)$ (c) $\mathbb{R}$ (d) See Figure 11.

38. (a) The natural logarithm is the logarithm with base e, denoted $\ln x$.

(b) The common logarithm is the logarithm with base 10, denoted $\log x$.

(c) See Figure 13.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. (a) $\log_3 81 = \log_3 3^4 = 4$ (b) $\log_3\left(\frac{1}{81}\right) = \log_3 3^{-4} = -4$ (c) $\log_9 3 = \log_9 9^{1/2} = \frac{1}{2}$

40. (a) $\ln \dfrac{1}{e^2} = \ln e^{-2} = -2$ (b) $\ln \sqrt{e} = \ln e^{1/2} = \frac{1}{2}$ (c) $\ln\left(\ln e^{e^{50}}\right) = \ln(e^{50}) = 50$

41. (a) $\log_2 30 - \log_2 15 = \log_2\left(\dfrac{30}{15}\right) = \log_2 2 = 1$

(b) $\log_3 10 - \log_3 5 - \log_3 18 = \log_3\left(\dfrac{10}{5}\right) - \log_3 18 = \log_3 2 - \log_3 18 = \log_3\left(\dfrac{2}{18}\right) = \log_3\left(\dfrac{1}{9}\right)$

$= \log_3 3^{-2} = -2$

(c) $2\log_5 100 - 4\log_5 50 = \log_5 100^2 - \log_5 50^4 = \log_5\left(\dfrac{100^2}{50^4}\right) = \log_5\left(\dfrac{10^4}{5^4 \cdot 10^4}\right) = \log_5 5^{-4} = -4$

42. (a) $e^{3\ln 2} = e^{\ln 2^3} = 2^3 = 8$ (b) $e^{-2\ln 5} = e^{\ln 5^{-2}} = 5^{-2} = \frac{1}{25}$ (c) $e^{\ln(\ln e^3)} = e^{\ln(3)} = 3$

43. (a) $\log_{10}\left(x^2y^3z\right) = \log_{10} x^2 + \log_{10} y^3 + \log_{10} z$ [Law 1]

$= 2\log_{10} x + 3\log_{10} y + \log_{10} z$ [Law 3]

(b) $\ln\left(\dfrac{x^4}{\sqrt{x^2-4}}\right) = \ln x^4 - \ln(x^2-4)^{1/2}$ [Law 2]

$= 4\ln x - \frac{1}{2}\ln[(x+2)(x-2)]$ [Law 3]

$= 4\ln x - \frac{1}{2}[\ln(x+2) + \ln(x-2)]$ [Law 1]

$= 4\ln x - \frac{1}{2}\ln(x+2) - \frac{1}{2}\ln(x-2)$

44. (a) $\ln\sqrt{\dfrac{3x}{x-3}} = \ln\left(\dfrac{3x}{x-3}\right)^{1/2} = \dfrac{1}{2}\ln\left(\dfrac{3x}{x-3}\right)$ [Law 3]

$= \frac{1}{2}[\ln 3 + \ln x - \ln(x-3)]$ [Laws 1 and 2]

$= \frac{1}{2}\ln 3 + \frac{1}{2}\ln x - \frac{1}{2}\ln(x-3)$

(b) $\log_2\left[(x^3+1)\sqrt[3]{(x-3)^2}\right] = \log_2(x^3+1) + \log_2\sqrt[3]{(x-3)^2}$ [Law 1]

$= \log_2(x^3+1) + \log_2(x-3)^{2/3}$

$= \log_2(x^3+1) + \frac{2}{3}\log_2(x-3)$ [Law 3]

45. (a) $\log_{10} 20 - \frac{1}{3}\log_{10} 1000 = \log_{10} 20 - \log_{10} 1000^{1/3} = \log_{10} 20 - \log_{10}\sqrt[3]{1000}$

$= \log_{10} 20 - \log_{10} 10 = \log_{10}\left(\frac{20}{10}\right) = \log_{10} 2$

(b) $\ln a - 2\ln b + 3\ln c = \ln a - \ln b^2 + \ln c^3 = \ln\dfrac{a}{b^2} + \ln c^3 = \ln\dfrac{ac^3}{b^2}$

46. (a) $3\ln(x-2) - \ln(x^2-5x+6) + 2\ln(x-3) = \ln(x-2)^3 - \ln[(x-2)(x-3)] + \ln(x-3)^2$

$= \ln\left[\dfrac{(x-2)^3(x-3)^2}{(x-2)(x-3)}\right] = \ln[(x-2)^2(x-3)]$

(b) $c\log_a x - d\log_a y + \log_a z = \log_a x^c - \log_a y^d + \log_a z = \log_a\left(\dfrac{x^c z}{y^d}\right)$

47. (a) $\log_5 10 = \dfrac{\ln 10}{\ln 5} \approx 1.430677$ (b) $\log_{15} 12 = \dfrac{\ln 12}{\ln 15} \approx 0.917600$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. (a) $\log_3 12 = \dfrac{\ln 12}{\ln 3} \approx 2.261860$ (b) $\log_{12} 6 = \dfrac{\ln 6}{\ln 12} \approx 0.721057$

49. To graph these functions, we use $\log_{1.5} x = \dfrac{\ln x}{\ln 1.5}$ and $\log_{50} x = \dfrac{\ln x}{\ln 50}$.

These graphs all approach $-\infty$ as $x \to 0^+$, and they all pass through the point $(1, 0)$. Also, they are all increasing, and all approach ∞ as $x \to \infty$. The functions with larger bases increase extremely slowly, and the ones with smaller bases do so somewhat more quickly. The functions with large bases approach the y-axis more closely as $x \to 0^+$.

50. We see that the graph of $\ln x$ is the reflection of the graph of e^x about the line $y = x$, and that the graph of $\log_8 x$ is the reflection of the graph of 8^x about the same line. The graph of 8^x increases more quickly than that of e^x. Also note that $\log_8 x \to \infty$ as $x \to \infty$ more slowly than $\ln x$.

51. 3 ft $= 36$ in, so we need x such that $\log_2 x = 36 \quad \Leftrightarrow \quad x = 2^{36} = 68{,}719{,}476{,}736$. In miles, this is

$68{,}719{,}476{,}736 \text{ in} \cdot \dfrac{1 \text{ ft}}{12 \text{ in}} \cdot \dfrac{1 \text{ mi}}{5280 \text{ ft}} \approx 1{,}084{,}587.7 \text{ mi}.$

52.

From the graphs, we see that $f(x) = x^{0.1} > g(x) = \ln x$ for approximately $0 < x < 3.06$, and then $g(x) > f(x)$ for $3.06 < x < 3.43 \times 10^{15}$ (approximately). At that point, the graph of f finally surpasses the graph of g for good.

53. (a) Shift the graph of $y = \log_{10} x$ five units to the left to obtain the graph of $y = \log_{10}(x + 5)$. Note the vertical asymptote of $x = -5$.

(b) Reflect the graph of $y = \ln x$ about the x-axis to obtain the graph of $y = -\ln x$.

$y = \log_{10} x$

$y = \log_{10}(x + 5)$

$y = \ln x$

$y = -\ln x$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. (a) Reflect the graph of $y = \ln x$ about the y-axis to obtain the graph of $y = \ln(-x)$.

$y = \ln x$

$y = \ln(-x)$

(b) Reflect the portion of the graph of $y = \ln x$ to the right of the y-axis about the y-axis. The graph of $y = \ln|x|$ is that reflection in addition to the original portion.

$y = \ln x$

$y = \ln|x|$

55. (a) The domain of $f(x) = \ln x + 2$ is $x > 0$ and the range is $\mathbb{R}$.

(b) $y = 0 \quad \Rightarrow \quad \ln x + 2 = 0 \quad \Rightarrow \quad \ln x = -2 \quad \Rightarrow \quad x = e^{-2}$

(c) We shift the graph of $y = \ln x$ two units upward.

56. (a) The domain of $f(x) = \ln(x-1) - 1$ is $x > 1$ and the range is $\mathbb{R}$.

(b) $y = 0 \quad \Rightarrow \quad \ln(x-1) - 1 = 0 \quad \Rightarrow \quad \ln(x-1) = 1 \quad \Rightarrow$
$x - 1 = e^1 \quad \Rightarrow \quad x = e + 1$

(c) We shift the graph of $y = \ln x$ one unit to the right and one unit downward.

57. (a) $\ln(4x+2) = 3 \quad \Rightarrow \quad e^{\ln(4x+2)} = e^3 \quad \Rightarrow \quad 4x + 2 = e^3 \quad \Rightarrow \quad 4x = e^3 - 2 \quad \Rightarrow \quad x = \frac{1}{4}(e^3 - 2) \approx 4.521$

(b) $e^{2x-3} = 12 \quad \Rightarrow \quad \ln e^{2x-3} = \ln 12 \quad \Rightarrow \quad 2x - 3 = \ln 12 \quad \Rightarrow \quad 2x = 3 + \ln 12 \quad \Rightarrow \quad x = \frac{1}{2}(3 + \ln 12) \approx 2.742$

58. (a) $\log_2(x^2 - x - 1) = 2 \quad \Rightarrow \quad x^2 - x - 1 = 2^2 = 4 \quad \Rightarrow \quad x^2 - x - 5 = 0 \quad \Rightarrow$

$$x = \frac{1 \pm \sqrt{(-1)^2 - 4(1)(-5)}}{2(1)} = \frac{1 \pm \sqrt{21}}{2}.$$

Solutions are $x_1 = \dfrac{1 - \sqrt{21}}{2} \approx -1.791$ and $x_2 = \dfrac{1 + \sqrt{21}}{2} \approx 2.791$.

(b) $1 + e^{4x+1} = 20 \quad \Rightarrow \quad e^{4x+1} = 19 \quad \Rightarrow \quad \ln e^{4x+1} = \ln 19 \quad \Rightarrow \quad 4x + 1 = \ln 19 \quad \Rightarrow \quad 4x = -1 + \ln 19 \quad \Rightarrow$
$x = \frac{1}{4}(-1 + \ln 19) \approx 0.486$

59. (a) $\ln x + \ln(x-1) = 0 \quad \Rightarrow \quad \ln[x(x-1)] = 0 \quad \Rightarrow \quad e^{\ln[x^2 - x]} = e^0 \quad \Rightarrow \quad x^2 - x = 1 \quad \Rightarrow \quad x^2 - x - 1 = 0$. The quadratic formula gives $x = \dfrac{1 \pm \sqrt{(-1)^2 - 4(1)(-1)}}{2(1)} = \dfrac{1 \pm \sqrt{5}}{2}$, but we note that $\ln \dfrac{1 - \sqrt{5}}{2}$ is undefined because $\dfrac{1 - \sqrt{5}}{2} < 0$. Thus, $x = \dfrac{1 + \sqrt{5}}{2} \approx 1.618$.

(b) $5^{1-2x} = 9 \quad \Rightarrow \quad \ln 5^{1-2x} = \ln 9 \quad \Rightarrow \quad (1 - 2x)\ln 5 = \ln 9 \quad \Rightarrow \quad 1 - 2x = \dfrac{\ln 9}{\ln 5} \quad \Rightarrow \quad x = \dfrac{1}{2} - \dfrac{\ln 9}{2\ln 5} \approx -0.183$

60. (a) $\ln(\ln x) = 0 \quad \Rightarrow \quad e^{\ln(\ln x)} = e^0 \quad \Rightarrow \quad \ln x = 1 \quad \Rightarrow \quad x = e \approx 2.718$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\dfrac{60}{1+e^{-x}} = 4 \quad\Rightarrow\quad 60 = 4(1+e^{-x}) \quad\Rightarrow\quad 15 = 1+e^{-x} \quad\Rightarrow\quad 14 = e^{-x} \quad\Rightarrow\quad \ln 14 = \ln e^{-x} \quad\Rightarrow$

$\ln 14 = -x \quad\Rightarrow\quad x = -\ln 14 \approx -2.639$

61. (a) $\ln x < 0 \quad\Rightarrow\quad x < e^0 \quad\Rightarrow\quad x < 1$. Since the domain of $f(x) = \ln x$ is $x > 0$, the solution of the original inequality is $0 < x < 1$.

(b) $e^x > 5 \quad\Rightarrow\quad \ln e^x > \ln 5 \quad\Rightarrow\quad x > \ln 5$

62. (a) $1 < e^{3x-1} < 2 \quad\Rightarrow\quad \ln 1 < 3x-1 < \ln 2 \quad\Rightarrow\quad 0 < 3x-1 < \ln 2 \quad\Rightarrow\quad 1 < 3x < 1+\ln 2 \quad\Rightarrow$

$\frac{1}{3} < x < \frac{1}{3}(1+\ln 2)$

(b) $1 - 2\ln x < 3 \quad\Rightarrow\quad -2\ln x < 2 \quad\Rightarrow\quad \ln x > -1 \quad\Rightarrow\quad x > e^{-1}$

63. (a) We must have $e^x - 3 > 0 \quad\Leftrightarrow\quad e^x > 3 \quad\Leftrightarrow\quad x > \ln 3$. Thus, the domain of $f(x) = \ln(e^x - 3)$ is $(\ln 3, \infty)$.

(b) $y = \ln(e^x - 3) \quad\Rightarrow\quad e^y = e^x - 3 \quad\Rightarrow\quad e^x = e^y + 3 \quad\Rightarrow\quad x = \ln(e^y + 3)$, so $f^{-1}(x) = \ln(e^x + 3)$.

Now $e^x + 3 > 0 \quad\Rightarrow\quad e^x > -3$, which is true for any real x, so the domain of f^{-1} is $\mathbb{R}$.

64. (a) By (9), $e^{\ln 300} = 300$ and $\ln(e^{300}) = 300$.

(b) A calculator gives $e^{\ln 300} = 300$ and an error message for $\ln(e^{300})$ because e^{300} is larger than most calculators can evaluate.

65. We see that the graph of $y = f(x) = \sqrt{x^3 + x^2 + x + 1}$ is increasing, so f is 1-1.

Enter $x = \sqrt{y^3 + y^2 + y + 1}$ and use your CAS to solve the equation for y. You will likely get two (irrelevant) solutions involving imaginary expressions, as well as one which can be simplified to

$y = f^{-1}(x) = -\frac{\sqrt[3]{4}}{6}\left(\sqrt[3]{D - 27x^2 + 20} - \sqrt[3]{D + 27x^2 - 20} + \sqrt[3]{2}\right)$

where $D = 3\sqrt{3}\sqrt{27x^4 - 40x^2 + 16}$ or, equivalently, $\dfrac{1}{6}\dfrac{M^{2/3} - 8 - 2M^{1/3}}{2M^{1/3}}$,

where $M = 108x^2 + 12\sqrt{48 - 120x^2 + 81x^4} - 80$.

66. (a) Depending on the software used, solving $x = y^6 + y^4$ for y may give six solutions of the form $y = \pm\frac{\sqrt{3}}{3}\sqrt{B-1}$, where $B \in \left\{-2\sin\dfrac{A}{3}, 2\sin\left(\dfrac{A}{3} + \dfrac{\pi}{3}\right), -2\cos\left(\dfrac{A}{3} + \dfrac{\pi}{6}\right)\right\}$ and $A = \sin^{-1}\left(\dfrac{27x-2}{2}\right)$. The inverse for $y = x^6 + x^4$ $(x \geq 0)$ is $y = \frac{\sqrt{3}}{3}\sqrt{B-1}$ with $B = 2\sin\left(\dfrac{A}{3} + \dfrac{\pi}{3}\right)$, but because the domain of A is $\left[0, \frac{4}{27}\right]$, this expression is only valid for $x \in \left[0, \frac{4}{27}\right]$.

If we solve $x = y^6 + y^4$ for y using Maple, we get the two real solutions $\pm\dfrac{\sqrt{6}}{6}\dfrac{\sqrt{C^{1/3}\left(C^{2/3} - 2C^{1/3} + 4\right)}}{C^{1/3}}$,

where $C = 108x + 12\sqrt{3}\sqrt{x(27x-4)}$, and the inverse for $y = x^6 + x^4$ $(x \geq 0)$ is the positive solution, whose domain is $\left[\frac{4}{27}, \infty\right)$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Mathematica also gives two real solutions, equivalent to those of Maple. The positive one is $\frac{\sqrt{6}}{6}\left(\sqrt[3]{4}D^{1/3} + 2\sqrt[3]{2}D^{-1/3} - 2\right)$, where $D = -2 + 27x + 3\sqrt{3}\sqrt{x}\sqrt{27x-4}$. Although this expression also has domain $\left[\frac{4}{27}, \infty\right)$, Mathematica is mysteriously able to plot the solution for all $x \geq 0$.

(b)

67. (a) $n = f(t) = 100 \cdot 2^{t/3} \quad \Rightarrow \quad \frac{n}{100} = 2^{t/3} \quad \Rightarrow \quad \log_2\left(\frac{n}{100}\right) = \frac{t}{3} \quad \Rightarrow \quad t = 3\log_2\left(\frac{n}{100}\right)$. Using the Change of Base Formula, we can write this as $t = f^{-1}(n) = 3 \cdot \frac{\ln(n/100)}{\ln 2}$. This function tells us how long it will take to obtain n bacteria (given the number n).

(b) $n = 50{,}000 \quad \Rightarrow \quad t = f^{-1}(50{,}000) = 3 \cdot \frac{\ln\left(\frac{50{,}000}{100}\right)}{\ln 2} = 3\left(\frac{\ln 500}{\ln 2}\right) \approx 26.9$ hours

68. (a) We write $Q = Q_0(1 - e^{-t/a})$ and solve for t: $\frac{Q}{Q_0} = 1 - e^{-t/a} \quad \Rightarrow \quad e^{-t/a} = 1 - \frac{Q}{Q_0} \quad \Rightarrow$ $-\frac{t}{a} = \ln\left(1 - \frac{Q}{Q_0}\right) \quad \Rightarrow \quad t = -a\ln\left(1 - \frac{Q}{Q_0}\right)$. This formula gives the time (in seconds) needed after a discharge to obtain a given charge Q.

(b) We set $Q = 0.9Q_0$ and $a = 50$ to get $t = -50\ln\left(1 - \frac{0.9Q_0}{Q_0}\right) = -50\ln(0.1) \approx 115.1$ seconds. It will take approximately 115 seconds—just shy of two minutes—to recharge the capacitors to 90% of capacity.

69. (a) $\cos^{-1}(-1) = \pi$ because $\cos\pi = -1$ and π is in the interval $[0, \pi]$ (the range of $\cos^{-1}$).

(b) $\sin^{-1}(0.5) = \frac{\pi}{6}$ because $\sin\frac{\pi}{6} = 0.5$ and $\frac{\pi}{6}$ is in the interval $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ (the range of $\sin^{-1}$).

70. (a) $\tan^{-1}\sqrt{3} = \frac{\pi}{3}$ because $\tan\frac{\pi}{3} = \sqrt{3}$ and $\frac{\pi}{3}$ is in the interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (the range of $\tan^{-1}$).

(b) $\arctan(-1) = -\frac{\pi}{4}$ because $\tan\left(-\frac{\pi}{4}\right) = -1$ and $-\frac{\pi}{4}$ is in the interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (the range of $\arctan$).

71. (a) $\csc^{-1}\sqrt{2} = \frac{\pi}{4}$ because $\csc\frac{\pi}{4} = \sqrt{2}$ and $\frac{\pi}{4}$ is in $\left(0, \frac{\pi}{2}\right] \cup \left(\pi, \frac{3\pi}{2}\right]$ (the range of $\csc^{-1}$).

(b) $\arcsin 1 = \frac{\pi}{2}$ because $\sin\frac{\pi}{2} = 1$ and $\frac{\pi}{2}$ is in $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ (the range of arcsin).

72. (a) $\sin^{-1}(-1/\sqrt{2}) = -\frac{\pi}{4}$ because $\sin\left(-\frac{\pi}{4}\right) = -1/\sqrt{2}$ and $-\frac{\pi}{4}$ is in $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.

(b) $\cos^{-1}\left(\sqrt{3}/2\right) = \frac{\pi}{6}$ because $\cos\frac{\pi}{6} = \sqrt{3}/2$ and $\frac{\pi}{6}$ is in $[0, \pi]$.

73. (a) $\cot^{-1}\left(-\sqrt{3}\right) = \frac{5\pi}{6}$ because $\cot\frac{5\pi}{6} = -\sqrt{3}$ and $\frac{5\pi}{6}$ is in $(0, \pi)$ (the range of $\cot^{-1}$).

(b) $\sec^{-1} 2 = \frac{\pi}{3}$ because $\sec\frac{\pi}{3} = 2$ and $\frac{\pi}{3}$ is in $\left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ (the range of $\sec^{-1}$).

74. (a) $\arcsin(\sin(5\pi/4)) = \arcsin\left(-1/\sqrt{2}\right) = -\frac{\pi}{4}$ because $\sin\left(-\frac{\pi}{4}\right) = -1/\sqrt{2}$ and $-\frac{\pi}{4}$ is in $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Let $\theta = \sin^{-1}\left(\frac{5}{13}\right)$ [see the figure].

$$\cos\left(2\sin^{-1}\left(\tfrac{5}{13}\right)\right) = \cos 2\theta = \cos^2\theta - \sin^2\theta$$
$$= \left(\tfrac{12}{13}\right)^2 - \left(\tfrac{5}{13}\right)^2 = \tfrac{144}{169} - \tfrac{25}{169} = \tfrac{119}{169}$$

75. Let $y = \sin^{-1} x$. Then $-\frac{\pi}{2} \le y \le \frac{\pi}{2} \quad \Rightarrow \quad \cos y \ge 0$, so $\cos(\sin^{-1} x) = \cos y = \sqrt{1 - \sin^2 y} = \sqrt{1 - x^2}$.

76. Let $y = \sin^{-1} x$. Then $\sin y = x$, so from the triangle (which illustrates the case $y > 0$), we see that

$$\tan(\sin^{-1} x) = \tan y = \frac{x}{\sqrt{1 - x^2}}.$$

77. Let $y = \tan^{-1} x$. Then $\tan y = x$, so from the triangle (which illustrates the case $y > 0$), we see that

$$\sin(\tan^{-1} x) = \sin y = \frac{x}{\sqrt{1 + x^2}}.$$

78. Let $y = \arccos x$. Then $\cos y = x$, so from the triangle (which illustrates the case $y > 0$), we see that

$$\sin(2\arccos x) = \sin 2y = 2\sin y\,\cos y$$
$$= 2(\sqrt{1 - x^2})(x) = 2x\sqrt{1 - x^2}$$

79.

The graph of $\sin^{-1} x$ is the reflection of the graph of $\sin x$ about the line $y = x$.

80.

The graph of $\tan^{-1} x$ is the reflection of the graph of $\tan x$ about the line $y = x$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

81. $g(x) = \sin^{-1}(3x+1)$.

Domain $(g) = \{x \mid -1 \le 3x+1 \le 1\} = \{x \mid -2 \le 3x \le 0\} = \left\{x \mid -\frac{2}{3} \le x \le 0\right\} = \left[-\frac{2}{3}, 0\right]$.

Range $(g) = \left\{y \mid -\frac{\pi}{2} \le y \le \frac{\pi}{2}\right\} = \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.

82. (a) $f(x) = \sin\left(\sin^{-1}x\right)$

Since one function undoes what the other one does, we get the identity function, $y = x$, on the restricted domain $-1 \le x \le 1$.

(b) $g(x) = \sin^{-1}(\sin x)$

This is similar to part (a), but with domain $\mathbb{R}$.

Equations for g on intervals of the form $\left(-\frac{\pi}{2} + \pi n, \frac{\pi}{2} + \pi n\right)$, for any integer n, can be found using $g(x) = (-1)^n x + (-1)^{n+1} n\pi$.

The sine function is monotonic on each of these intervals, and hence, so is g (but in a linear fashion).

83. (a) If the point (x, y) is on the graph of $y = f(x)$, then the point $(x - c, y)$ is that point shifted c units to the left. Since f is 1-1, the point (y, x) is on the graph of $y = f^{-1}(x)$ and the point corresponding to $(x - c, y)$ on the graph of f is $(y, x - c)$ on the graph of f^{-1}. Thus, the curve's reflection is shifted *down* the same number of units as the curve itself is shifted to the left. So an expression for the inverse function is $g^{-1}(x) = f^{-1}(x) - c$.

(b) If we compress (or stretch) a curve horizontally, the curve's reflection in the line $y = x$ is compressed (or stretched) *vertically* by the same factor. Using this geometric principle, we see that the inverse of $h(x) = f(cx)$ can be expressed as $h^{-1}(x) = (1/c)\, f^{-1}(x)$.

1 Review

TRUE-FALSE QUIZ

1. False. Let $f(x) = x^2$, $s = -1$, and $t = 1$. Then $f(s+t) = (-1+1)^2 = 0^2 = 0$, but $f(s) + f(t) = (-1)^2 + 1^2 = 2 \ne 0 = f(s+t)$.

2. False. Let $f(x) = x^2$. Then $f(-2) = 4 = f(2)$, but $-2 \ne 2$.

3. False. Let $f(x) = x^2$. Then $f(3x) = (3x)^2 = 9x^2$ and $3f(x) = 3x^2$. So $f(3x) \ne 3f(x)$.

4. True. The inverse function f^{-1} of a one-to-one function f is *defined* by $f^{-1}(y) = x \Leftrightarrow f(x) = y$.

5. True. See the Vertical Line Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. False. Let $f(x) = x^2$ and $g(x) = 2x$. Then $(f \circ g)(x) = f(g(x)) = f(2x) = (2x)^2 = 4x^2$ and $(g \circ f)(x) = g(f(x)) = g(x^2) = 2x^2$. So $f \circ g \neq g \circ f$.

7. False. Let $f(x) = x^3$. Then f is one-to-one and $f^{-1}(x) = \sqrt[3]{x}$. But $1/f(x) = 1/x^3$, which is not equal to $f^{-1}(x)$.

8. True. We can divide by e^x since $e^x \neq 0$ for every x.

9. True. The function $\ln x$ is an increasing function on $(0, \infty)$.

10. False. Let $x = e$. Then $(\ln x)^6 = (\ln e)^6 = 1^6 = 1$, but $6 \ln x = 6 \ln e = 6 \cdot 1 = 6 \neq 1 = (\ln x)^6$. What *is* true, however, is that $\ln(x^6) = 6 \ln x$ for $x > 0$.

11. False. Let $x = e^2$ and $a = e$. Then $\dfrac{\ln x}{\ln a} = \dfrac{\ln e^2}{\ln e} = \dfrac{2 \ln e}{\ln e} = 2$ and $\ln \dfrac{x}{a} = \ln \dfrac{e^2}{e} = \ln e = 1$, so in general the statement is false. What *is* true, however, is that $\ln \dfrac{x}{a} = \ln x - \ln a$.

12. False. It is true that $\tan \frac{3\pi}{4} = -1$, but since the range of $\tan^{-1}$ is $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, we must have $\tan^{-1}(-1) = -\frac{\pi}{4}$.

13. False. For example, $\tan^{-1} 20$ is defined; $\sin^{-1} 20$ and $\cos^{-1} 20$ are not.

14. False. For example, if $x = -3$, then $\sqrt{(-3)^2} = \sqrt{9} = 3$, not -3.

EXERCISES

1. (a) When $x = 2$, $y \approx 2.7$. Thus, $f(2) \approx 2.7$.

(b) $f(x) = 3 \quad \Rightarrow \quad x \approx 2.3, 5.6$

(c) The domain of f is $-6 \leq x \leq 6$, or $[-6, 6]$.

(d) The range of f is $-4 \leq y \leq 4$, or $[-4, 4]$.

(e) f is increasing on $[-4, 4]$, that is, on $-4 \leq x \leq 4$.

(f) f is not one-to-one because it fails the Horizontal Line Test.

(g) f is odd because its graph is symmetric about the origin.

2. (a) When $x = 2$, $y = 3$. Thus, $g(2) = 3$.

(b) g is one-to-one because it passes the Horizontal Line Test.

(c) When $y = 2$, $x \approx 0.2$. So $g^{-1}(2) \approx 0.2$.

(d) The range of g is $[-1, 3.5]$, which is the same as the domain of g^{-1}.

(e) We reflect the graph of g through the line $y = x$ to obtain the graph of g^{-1}.

3. $f(x) = x^2 - 2x + 3$, so $f(a + h) = (a + h)^2 - 2(a + h) + 3 = a^2 + 2ah + h^2 - 2a - 2h + 3$, and

$$\frac{f(a+h) - f(a)}{h} = \frac{(a^2 + 2ah + h^2 - 2a - 2h + 3) - (a^2 - 2a + 3)}{h} = \frac{h(2a + h - 2)}{h} = 2a + h - 2.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. There will be some yield with no fertilizer, increasing yields with increasing fertilizer use, a leveling-off of yields at some point, and disaster with too much fertilizer use.

5. $f(x) = 2/(3x-1)$.

Domain: $3x - 1 \neq 0 \quad \Rightarrow \quad 3x \neq 1 \quad \Rightarrow \quad x \neq \frac{1}{3}$. $\quad D = \left(-\infty, \frac{1}{3}\right) \cup \left(\frac{1}{3}, \infty\right)$

Range: all reals except 0 ($y = 0$ is the horizontal asymptote for f.)

$R = (-\infty, 0) \cup (0, \infty)$

6. $g(x) = \sqrt{16 - x^4}$.

Domain: $16 - x^4 \geq 0 \quad \Rightarrow \quad x^4 \leq 16 \quad \Rightarrow \quad |x| \leq \sqrt[4]{16} \quad \Rightarrow \quad |x| \leq 2$. $\quad D = [-2, 2]$

Range: $y \geq 0$ and $y \leq \sqrt{16} \quad \Rightarrow \quad 0 \leq y \leq 4$.

$R = [0, 4]$

7. $h(x) = \ln(x + 6)$.

Domain: $x + 6 > 0 \quad \Rightarrow \quad x > -6$. $\quad D = (-6, \infty)$

Range: $x + 6 > 0$, so $\ln(x + 6)$ takes on all real numbers and, hence, the range is $\mathbb{R}$.

$R = (-\infty, \infty)$

8. $y = F(t) = 3 + \cos 2t$.

Domain: $\mathbb{R}$. $\quad D = (-\infty, \infty)$

Range: $-1 \leq \cos 2t \leq 1 \quad \Rightarrow \quad 2 \leq 3 + \cos 2t \leq 4 \quad \Rightarrow \quad 2 \leq y \leq 4$.

$R = [2, 4]$

9. (a) To obtain the graph of $y = f(x) + 5$, we shift the graph of $y = f(x)$ 5 units upward.

(b) To obtain the graph of $y = f(x + 5)$, we shift the graph of $y = f(x)$ 5 units to the left.

(c) To obtain the graph of $y = 1 + 2f(x)$, we stretch the graph of $y = f(x)$ vertically by a factor of 2, and then shift the resulting graph 1 unit upward.

(d) To obtain the graph of $y = f(x - 2) - 2$, we shift the graph of $y = f(x)$ 2 units to the right (for the "-2" inside the parentheses), and then shift the resulting graph 2 units downward.

(e) To obtain the graph of $y = -f(x)$, we reflect the graph of $y = f(x)$ about the x-axis.

(f) To obtain the graph of $y = f^{-1}(x)$, we reflect the graph of $y = f(x)$ about the line $y = x$ (assuming f is one–to-one).

10. (a) To obtain the graph of $y = f(x - 8)$, we shift the graph of $y = f(x)$ right 8 units.

(b) To obtain the graph of $y = -f(x)$, we reflect the graph of $y = f(x)$ about the x-axis.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) To obtain the graph of $y = 2 - f(x)$, we reflect the graph of $y = f(x)$ about the x-axis, and then shift the resulting graph 2 units upward.

(d) To obtain the graph of $y = \frac{1}{2}f(x) - 1$, we shrink the graph of $y = f(x)$ by a factor of 2, and then shift the resulting graph 1 unit downward.

(e) To obtain the graph of $y = f^{-1}(x)$, we reflect the graph of $y = f(x)$ about the line $y = x$.

(f) To obtain the graph of $y = f^{-1}(x + 3)$, we reflect the graph of $y = f(x)$ about the line $y = x$ [see part (e)], and then shift the resulting graph left 3 units.

11. $f(x) = x^3 + 2$. Start with the graph of $y = x^3$ and shift 2 units upward.

12. $f(x) = (x - 3)^2$. Start with the graph of $y = x^2$ and shift 3 units to the right.

13. $y = \sqrt{x + 2}$. Start with the graph of $y = \sqrt{x}$ and shift 2 units to the left.

14. $y = \ln(x + 5)$. Start with the graph of $y = \ln x$ and shift 5 units to the left.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $g(x) = 1 + \cos 2x$. Start with the graph of $y = \cos x$, compress horizontally by a factor of 2, and then shift 1 unit upward.

16. $h(x) = -e^x + 2$. Start with the graph of $y = e^x$, reflect about the x-axis, and then shift 2 units upward.

17. $s(x) = 1 + 0.5^x$. Start with the graph of $y = 0.5^x = \left(\frac{1}{2}\right)^x$ and shift 1 unit upward.

18. $f(x) = \begin{cases} -x & \text{if } x < 0 \\ e^x - 1 & \text{if } x \geq 0 \end{cases}$

On $(-\infty, 0)$, graph $y = -x$ (the line with slope -1 and y-intercept 0) with open endpoint $(0, 0)$.

On $[0, \infty)$, graph $y = e^x - 1$ (the graph of $y = e^x$ shifted 1 unit downward) with closed endpoint $(0, 0)$.

19. (a) $f(x) = 2x^5 - 3x^2 + 2 \quad \Rightarrow \quad f(-x) = 2(-x)^5 - 3(-x)^2 + 2 = -2x^5 - 3x^2 + 2$. Since $f(-x) \neq f(x)$ and $f(-x) \neq -f(x)$, f is neither even nor odd.

(b) $f(x) = x^3 - x^7 \quad \Rightarrow \quad f(-x) = (-x)^3 - (-x)^7 = -x^3 + x^7 = -(x^3 - x^7) = -f(x)$, so f is odd.

(c) $f(x) = e^{-x^2} \quad \Rightarrow \quad f(-x) = e^{-(-x)^2} = e^{-x^2} = f(x)$, so f is even.

(d) $f(x) = 1 + \sin x \quad \Rightarrow \quad f(-x) = 1 + \sin(-x) = 1 - \sin x$. Now $f(-x) \neq f(x)$ and $f(-x) \neq -f(x)$, so f is neither even nor odd.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(e) $f(x) = 1 - \cos 2x \quad \Rightarrow \quad f(-x) = 1 - \cos[2(-x)] = 1 - \cos(-2x) = 1 - \cos 2x = f(x)$, so f is even.

(f) $f(x) = (x+1)^2 = x^2 + 2x + 1$. Now $f(-x) = (-x)^2 + 2(-x) + 1 = x^2 - 2x + 1$. Since $f(-x) \neq f(x)$ and $f(-x) \neq -f(x)$, f is neither even nor odd.

20. For the line segment from $(-2, 2)$ to $(-1, 0)$, the slope is $\dfrac{0-2}{-1+2} = -2$, and an equation is $y - 0 = -2(x+1)$ or, equivalently, $y = -2x - 2$. The circle has equation $x^2 + y^2 = 1$; the top half has equation $y = \sqrt{1 - x^2}$ (we have solved for positive y). Thus, $f(x) = \begin{cases} -2x - 2 & \text{if } -2 \le x \le -1 \\ \sqrt{1 - x^2} & \text{if } -1 < x \le 1 \end{cases}$.

21. $f(x) = \ln x, \quad D = (0, \infty); \quad g(x) = x^2 - 9, \quad D = \mathbb{R}$.

(a) $(f \circ g)(x) = f(g(x)) = f(x^2 - 9) = \ln(x^2 - 9)$.
Domain: $x^2 - 9 > 0 \quad \Rightarrow \quad x^2 > 9 \quad \Rightarrow \quad |x| > 3 \quad \Rightarrow \quad x \in (-\infty, -3) \cup (3, \infty)$

(b) $(g \circ f)(x) = g(f(x)) = g(\ln x) = (\ln x)^2 - 9$. Domain: $x > 0$, or $(0, \infty)$

(c) $(f \circ f)(x) = f(f(x)) = f(\ln x) = \ln \ln x$. Domain: $\ln x > 0 \quad \Rightarrow \quad x > e^0 = 1$, or $(1, \infty)$

(d) $(g \circ g)(x) = g(g(x)) = g(x^2 - 9) = (x^2 - 9)^2 - 9$. Domain: $x \in \mathbb{R}$, or $(-\infty, \infty)$

22. Let $h(x) = x + \sqrt{x}$, $g(x) = \sqrt{x}$, and $f(x) = 1/x$. Then $(f \circ g \circ h)(x) = \dfrac{1}{\sqrt{x + \sqrt{x}}} = F(x)$.

23.

More than one model appears to be plausible. Your choice of model depends on whether you think medical advances will keep increasing life expectancy, or if there is bound to be a natural leveling-off of life expectancy. A linear model, $y = 0.2441x - 413.3960$, gives us an estimate of 82.1 years for the year 2030.

24. (a) Let x denote the number of toaster ovens produced in one week and y the associated cost. Using the points $(1000, 9000)$ and $(1500, 12{,}000)$, we get an equation of a line:

$$y - 9000 = \frac{12{,}000 - 9000}{1500 - 1000}(x - 1000) \quad \Rightarrow$$

$$y = 6(x - 1000) + 9000 \quad \Rightarrow \quad y = 6x + 3000.$$

(b) The slope of 6 means that each additional toaster oven produced adds \$6 to the weekly production cost.

(c) The y-intercept of 3000 represents the overhead cost—the cost incurred without producing anything.

25. The value of x for which $f(x) = 2x + 4^x$ equals 6 will be $f^{-1}(6)$. To solve $2x + 4^x = 6$, we either observe that letting $x = 1$ gives us equality, or we graph $y_1 = 2x + 4^x$ and $y_2 = 6$ to find the intersection at $x = 1$. Since $f(1) = 6$, $f^{-1}(6) = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. We write $y = \dfrac{2x+3}{1-5x}$ and solve for x: $y(1-5x) = 2x+3 \quad\Rightarrow\quad y - 5xy = 2x+3 \quad\Rightarrow\quad y-3 = 2x+5xy \quad\Rightarrow$

$y-3 = x(2+5y) \quad\Rightarrow\quad x = \dfrac{y-3}{2+5y}$. Interchanging x and y gives $y = \dfrac{x-3}{2+5x}$, so $f^{-1}(x) = \dfrac{x-3}{2+5x}$.

27. (a) $\ln x\sqrt{x+1} = \ln x + \ln\sqrt{x+1}$ [Law 1]

$= \ln x + \ln(x+1)^{1/2} = \ln x + \frac{1}{2}\ln(x+1)$ [Law 3]

(b) $\log_2\sqrt{\dfrac{x^2+1}{x-1}} = \log_2\left(\dfrac{x^2+1}{x-1}\right)^{1/2}$

$= \dfrac{1}{2}\log_2\left(\dfrac{x^2+1}{x-1}\right)$ [Law 3]

$= \frac{1}{2}\left[\log_2(x^2+1) - \log_2(x-1)\right]$ [Law 2]

$= \frac{1}{2}\log_2(x^2+1) - \frac{1}{2}\log_2(x-1)$

28. (a) $\frac{1}{2}\ln x - 2\ln(x^2+1) = \ln x^{1/2} - \ln(x^2+1)^2 = \ln\dfrac{\sqrt{x}}{(x^2+1)^2}$

(b) $\ln(x-3) + \ln(x+3) - 2\ln(x^2-9) = \ln[(x-3)(x+3)] - \ln(x^2-9)^2$

$= \ln\dfrac{(x-3)(x+3)}{(x^2-9)^2} = \ln\dfrac{x^2-9}{(x^2-9)^2} = \ln\dfrac{1}{x^2-9}$

29. (a) $e^{2\ln 5} = e^{\ln 5^2} = 5^2 = 25$

(b) $\log_6 4 + \log_6 54 = \log_6(4\cdot 54) = \log_6 216 = \log_6 6^3 = 3$

(c) Let $\theta = \arcsin\frac{4}{5}$, so $\sin\theta = \frac{4}{5}$. Draw a right triangle with angle θ as shown in the figure. By the Pythagorean Theorem, the adjacent side has length 3, and $\tan\left(\arcsin\dfrac{4}{5}\right) = \tan\theta = \dfrac{\text{opp}}{\text{adj}} = \dfrac{4}{3}$.

30. (a) $\ln\dfrac{1}{e^3} = \ln e^{-3} = -3$

(b) $\sin(\tan^{-1}1) = \sin\dfrac{\pi}{4} = \dfrac{\sqrt{2}}{2}$

(c) $10^{-3\log 4} = 10^{\log 4^{-3}} = 4^{-3} = \dfrac{1}{4^3} = \dfrac{1}{64}$

31. $e^{2x} = 3 \quad\Rightarrow\quad \ln(e^{2x}) = \ln 3 \quad\Rightarrow\quad 2x = \ln 3 \quad\Rightarrow\quad x = \frac{1}{2}\ln 3 \approx 0.549$

32. $\ln x^2 = 5 \quad\Rightarrow\quad e^{\ln x^2} = e^5 \quad\Rightarrow\quad x^2 = e^5 \quad\Rightarrow\quad x = \pm\sqrt{e^5} \approx \pm 12.182$

33. $e^{e^x} = 10 \quad\Rightarrow\quad \ln\left(e^{e^x}\right) = \ln 10 \quad\Rightarrow\quad e^x = \ln 10 \quad\Rightarrow\quad \ln e^x = \ln(\ln 10) \quad\Rightarrow\quad x = \ln(\ln 10) \approx 0.834$

34. $\cos^{-1}x = 2 \quad\Rightarrow\quad \cos(\cos^{-1}x) = \cos 2 \quad\Rightarrow\quad x = \cos 2 \approx -0.416$

35. $\tan^{-1}(3x^2) = \dfrac{\pi}{4} \quad\Rightarrow\quad \tan(\tan^{-1}(3x^2)) = \tan\dfrac{\pi}{4} \quad\Rightarrow\quad 3x^2 = 1 \quad\Rightarrow\quad x^2 = \dfrac{1}{3} \quad\Rightarrow\quad x = \pm\dfrac{1}{\sqrt{3}} \approx \pm 0.577$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. $\ln x - 1 = \ln(5+x) - 4 \quad \Rightarrow \quad \ln x - \ln(5+x) = -4+1 \quad \Rightarrow \quad \ln \dfrac{x}{5+x} = -3 \quad \Rightarrow \quad e^{\ln(x/(5+x))} = e^{-3} \quad \Rightarrow$

$\dfrac{x}{5+x} = e^{-3} \quad \Rightarrow \quad x = 5e^{-3} + xe^{-3} \quad \Rightarrow \quad x - xe^{-3} = 5e^{-3} \quad \Rightarrow \quad x(1-e^{-3}) = 5e^{-3} \quad \Rightarrow \quad x = \dfrac{5e^{-3}}{1-e^{-3}}$

or, multiplying by $\dfrac{e^3}{e^3}$, we have $x = \dfrac{5}{e^3 - 1} \approx 0.262$.

37. (a) The half-life of the virus with this treatment is eight days and 24 days is 3 half-lives, so the viral load after 24 days is

$52.0\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)\left(\frac{1}{2}\right) = 52.0\left(\frac{1}{2}\right)^3 = 6.5$ RNA copies/mL.

(b) The viral load is halved every $t/8$ days, so $V(t) = 52.0\left(\frac{1}{2}\right)^{t/8}$.

(c) $V = V(t) = 52.0\left(\frac{1}{2}\right)^{t/8} \quad \Rightarrow \quad \dfrac{V}{52.0} = \left(\frac{1}{2}\right)^{t/8} = 2^{-t/8} \quad \Rightarrow \quad \log_2\left(\dfrac{V}{52.0}\right) = \log_2\left(2^{-t/8}\right) = -\dfrac{t}{8} \quad \Rightarrow$

$t = t(V) - 8\log_2\left(\dfrac{V}{52.0}\right)$. This gives the number of days t needed after treatment begins for the viral load to be reduced to V RNA copies/mL.

(d) Using the function from part (c), we have $t(2.0) = -8\log_2\left(\dfrac{2.0}{52.0}\right) = -8 \cdot \dfrac{\ln \frac{1}{26}}{\ln 2} \approx 37.6$ days.

38. (a)

The population would reach 900 in about 4.4 years.

(b) $P = \dfrac{100{,}000}{100 + 900e^{-t}} \Rightarrow \quad 100P + 900Pe^{-t} = 100{,}000 \Rightarrow \quad 900Pe^{-t} = 100{,}000 - 100P \quad \Rightarrow$

$e^{-t} = \dfrac{100{,}000 - 100P}{900P} \quad \Rightarrow \quad -t = \ln\left(\dfrac{1000-P}{9P}\right) \quad \Rightarrow \quad t = -\ln\left(\dfrac{1000-P}{9P}\right)$, or $\ln\left(\dfrac{9P}{1000-P}\right)$;

this is the time required for the population to reach a given number P.

(c) $P = 900 \quad \Rightarrow \quad t = \ln\left(\dfrac{9 \cdot 900}{1000 - 900}\right) = \ln 81 \approx 4.4$ years, as in part (a).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PRINCIPLES OF PROBLEM SOLVING

1.

By using the area formula for a triangle, $\frac{1}{2}$ (base) (height), in two ways, we see that $\frac{1}{2}(4)(y) = \frac{1}{2}(h)(a)$, so $a = \dfrac{4y}{h}$. Since $4^2 + y^2 = h^2$, $y = \sqrt{h^2 - 16}$, and $a = \dfrac{4\sqrt{h^2 - 16}}{h}$.

2.

Refer to Example 1, where we obtained $h = \dfrac{P^2 - 100}{2P}$. The 100 came from 4 times the area of the triangle. In this case, the area of the triangle is $\frac{1}{2}(h)(12) = 6h$. Thus, $h = \dfrac{P^2 - 4\,(6h)}{2P} \quad \Rightarrow \quad 2Ph = P^2 - 24h \quad \Rightarrow$

$2Ph + 24h = P^2 \quad \Rightarrow \quad h\,(2P + 24) = P^2 \quad \Rightarrow \quad h = \dfrac{P^2}{2P + 24}$.

3. $\big|4x - |x + 1|\,\big| = 3 \quad \Rightarrow \quad 4x - |x + 1| = -3$ (Equation 1) or $4x - |x + 1| = 3$ (Equation 2).

If $x + 1 < 0$, or $x < -1$, then $|x + 1| = -(x + 1) = -x - 1$. If $x + 1 \geq 0$, or $x \geq -1$, then $|x + 1| = x + 1$. We thus consider two cases, $x < -1$ (Case 1) and $x \geq -1$ (Case 2), for each of Equations 1 and 2.

Equation 1, Case 1: $4x - |x + 1| = -3 \quad \Rightarrow \quad 4x - (-x - 1) = -3 \quad \Rightarrow \quad 5x + 1 = -3 \quad \Rightarrow$
$5x = -4 \quad \Rightarrow \quad x = -\frac{4}{5}$ which is invalid since $x < -1$.

Equation 1, Case 2: $4x - |x + 1| = -3 \quad \Rightarrow \quad 4x - (x - 1) = -3 \quad \Rightarrow \quad 3x - 1 = -3 \quad \Rightarrow$
$3x = -2 \quad \Rightarrow \quad x = -\frac{2}{3}$, which is valid since $x \geq -1$.

Equation 2, Case 1: $4x - |x + 1| = 3 \quad \Rightarrow \quad 4x - (-x - 1) = 3 \quad \Rightarrow \quad 5x + 1 = 3 \quad \Rightarrow$
$5x = 2 \quad \Rightarrow \quad x = \frac{2}{5}$, which is invalid since $x < -1$.

Equation 2, Case 2: $4x - |x + 1| = 3 \quad \Rightarrow \quad 4x - (x + 1) = 3 \quad \Rightarrow \quad 3x - 1 = 3 \quad \Rightarrow$
$3x = 4 \quad \Rightarrow \quad x = \frac{4}{3}$, which is valid since $x \geq -1$.

Thus, the solution set is $\left\{-\frac{2}{3}, \frac{4}{3}\right\}$.

4. $|x - 1| = \begin{cases} x - 1 & \text{if } x \geq 1 \\ 1 - x & \text{if } x < 1 \end{cases}$ and $|x - 3| = \begin{cases} x - 3 & \text{if } x \geq 3 \\ 3 - x & \text{if } x < 3 \end{cases}$

Therefore, we consider the three cases $x < 1$, $1 \leq x < 3$, and $x \geq 3$.

If $x < 1$, we must have $1 - x - (3 - x) \geq 5 \quad \Leftrightarrow \quad 0 \geq 7$, which is false.

If $1 \leq x < 3$, we must have $x - 1 - (3 - x) \geq 5 \quad \Leftrightarrow \quad x \geq \frac{9}{2}$, which is false because $x < 3$.

If $x \geq 3$, we must have $x - 1 - (x - 3) \geq 5 \quad \Leftrightarrow \quad 2 \geq 5$, which is false.

All three cases lead to falsehoods, so the inequality has no solution.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $f(x) = \left|x^2 - 4\,|x| + 3\right|$. If $x \geq 0$, then $f(x) = \left|x^2 - 4x + 3\right| = |(x-1)(x-3)|$.

Case (i): If $0 < x \leq 1$, then $f(x) = x^2 - 4x + 3$.

Case (ii): If $1 < x \leq 3$, then $f(x) = -(x^2 - 4x + 3) = -x^2 + 4x - 3$.

Case (iii): If $x > 3$, then $f(x) = x^2 - 4x + 3$.

This enables us to sketch the graph for $x \geq 0$. Then we use the fact that f is an even function to reflect this part of the graph about the y-axis to obtain the entire graph. Or, we could consider also the cases $x < -3$, $-3 \leq x < -1$, and $-1 \leq x < 0$.

6. $g(x) = \left|x^2 - 1\right| - \left|x^2 - 4\right|$.

$$\left|x^2 - 1\right| = \begin{cases} x^2 - 1 & \text{if } |x| \geq 1 \\ 1 - x^2 & \text{if } |x| < 1 \end{cases} \quad \text{and} \quad \left|x^2 - 4\right| = \begin{cases} x^2 - 4 & \text{if } |x| \geq 2 \\ 4 - x^2 & \text{if } |x| < 2 \end{cases}$$

So for $0 \leq |x| < 1, g(x) = 1 - x^2 - (4 - x^2) = -3$, for

$1 \leq |x| < 2, g(x) = x^2 - 1 - (4 - x^2) = 2\,x^2 - 5$, and for

$|x| \geq 2, g(x) = x^2 - 1 - (x^2 - 4) = 3$.

7. Remember that $|a| = a$ if $a \geq 0$ and that $|a| = -a$ if $a < 0$. Thus,

$$x + |x| = \begin{cases} 2x & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases} \qquad \text{and} \qquad y + |y| = \begin{cases} 2y & \text{if } y \geq 0 \\ 0 & \text{if } y < 0 \end{cases}$$

We will consider the equation $x + |x| = y + |y|$ in four cases.

(1) $x \geq 0, y \geq 0$	(2) $x \geq 0, y < 0$	(3) $x < 0, y \geq 0$	(4) $x < 0, y < 0$
$2x = 2y$	$2x = 0$	$0 = 2y$	$0 = 0$
$x = y$	$x = 0$	$0 = y$	

Case 1 gives us the line $y = x$ with nonnegative x and y.

Case 2 gives us the portion of the y-axis with y negative.

Case 3 gives us the portion of the x-axis with x negative.

Case 4 gives us the entire third quadrant.

8. $|x - y| + |x| - |y| \leq 2$ [call this inequality $(\star)$]

Case (i):	$x \geq y \geq 0$.	Then $(\star)$	$\Leftrightarrow$	$x - y + x - y \leq 2$	$\Leftrightarrow$	$x - y \leq 1$	$\Leftrightarrow$	$y \geq x - 1$.
Case (ii):	$y \geq x \geq 0$.	Then $(\star)$	$\Leftrightarrow$	$y - x + x - y \leq 2$	$\Leftrightarrow$	$0 \leq 2$ (true).		
Case (iii):	$x \geq 0$ and $y \leq 0$.	Then $(\star)$	$\Leftrightarrow$	$x - y + x + y \leq 2$	$\Leftrightarrow$	$2x \leq 2$	$\Leftrightarrow$	$x \leq 1$.
Case (iv):	$x \leq 0$ and $y \geq 0$.	Then $(\star)$	$\Leftrightarrow$	$y - x - x - y \leq 2$	$\Leftrightarrow$	$-2x \leq 2$	$\Leftrightarrow$	$x \geq -1$.
Case (v):	$y \leq x \leq 0$.	Then $(\star)$	$\Leftrightarrow$	$x - y - x + y \leq 2$	$\Leftrightarrow$	$0 \leq 2$ (true).		
Case (vi):	$x \leq y \leq 0$.	Then $(\star)$	$\Leftrightarrow$	$y - x - x + y \leq 2$	$\Leftrightarrow$	$y - x \leq 1$	$\Leftrightarrow$	$y \leq x + 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Note: Instead of considering cases (iv), (v), and (vi), we could have noted that the region is unchanged if x and y are replaced by $-x$ and $-y$, so the region is symmetric about the origin. Therefore, we need only draw cases (i), (ii), and (iii), and rotate through $180°$ about the origin.

9. (a) To sketch the graph of $f(x) = \max\{x, 1/x\}$, we first graph $g(x) = x$ and $h(x) = 1/x$ on the same coordinate axes. Then create the graph of f by plotting the largest y-value of g and h for every value of x.

(b)

(c)

On the TI-84 Plus, max is found under `LIST`, then under `MATH`. To graph $f(x) = \max\left\{x^2, 2+x, 2-x\right\}$, use $\texttt{Y} = \max(x^2, \max(2+x, 2-x))$.

10. (a) If $\max\{x, 2y\} = 1$, then either $x = 1$ *and* $2y \le 1$ ***or*** $x \le 1$ *and* $2y = 1$. Thus, we obtain the set of points such that $x = 1$ *and* $y \le \frac{1}{2}$ $\left[\text{a vertical line with highest point } (1, \frac{1}{2})\right]$ ***or*** $x \le 1$ *and* $y = \frac{1}{2}$ $\left[\text{a horizontal line with rightmost point } (1, \frac{1}{2})\right]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The graph of $\max\{x, 2y\} = 1$ is shown in part (a), and the graph of $\max\{x, 2y\} = -1$ can be found in a similar manner. The inequalities in $-1 \le \max\{x, 2y\} \le 1$ give us all the points on or inside the boundaries.

(c) $\max\{x, y^2\} = 1 \quad \Leftrightarrow \quad x = 1$ *and* $y^2 \le 1$ $[-1 \le y \le 1]$

or $x \le 1$ *and* $y^2 = 1$ $[y = \pm 1]$.

11.
$$\begin{aligned}
\frac{1}{\log_2 x} + \frac{1}{\log_3 x} + \frac{1}{\log_5 x} &= \frac{1}{\dfrac{\log x}{\log 2}} + \frac{1}{\dfrac{\log x}{\log 3}} + \frac{1}{\dfrac{\log x}{\log 5}} && \text{[Change of Base formula]} \\
&= \frac{\log 2}{\log x} + \frac{\log 3}{\log x} + \frac{\log 5}{\log x} \\
&= \frac{\log 2 + \log 3 + \log 5}{\log x} = \frac{\log(2 \cdot 3 \cdot 5)}{\log x} && \text{[Law 1 of Lograithms]} \\
&= \frac{\log 30}{\log x} = \frac{1}{\dfrac{\log x}{\log 30}} = \frac{1}{\log_{30} x} && \text{[Change of Base formula]}
\end{aligned}$$

12. We note that $-1 \le \sin x \le 1$ for all x. Thus, any solution of $\sin x = x/100$ will have $-1 \le x/100 \le 1$, or $-100 \le x \le 100$. We next observe that the period of $\sin x$ is 2π, and $\sin x$ takes on each value in its range, except for -1 and 1, twice each cycle. We observe that $x = 0$ is a solution. Finally, we note that because $\sin x$ and $x/100$ are both odd functions, every solution on $0 \le x \le 100$ gives us a corresponding solution on $-100 \le x \le 0$.

$100/2\pi \approx 15.9$, so there 15 full cycles of $\sin x$ on $[0, 100]$ .Each of the 15 intervals $[0, 2\pi]$, $[2\pi, 4\pi]$, . . ., $[28\pi, 30\pi]$ must contain two solutions of $\sin x = x/100$, as the graph of $\sin x$ will intersect the graph of $x/100$ twice each cycle. We must be careful with the next (16th) interval $[30\pi, 32\pi]$, because 100 is contained in the interval. A graph of $y_1 = \sin x$ and $y_2 = x/100$ over this interval reveals that two intersections occur within the interval with $x \le 100$.

Thus, there are $16 \cdot 2 = 32$ solutions of $\sin x = x/100$ on $[0, 100]$. There are also 32 solutions of the equation on $[-100, 0]$. Being careful to not count the solution $x = 0$ twice, we find that there are $32 + 32 - 1 = 63$ solutions of the equation $\sin x = x/100$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. By rearranging terms, we write the given expression as

$$\left(\sin\frac{\pi}{100}+\sin\frac{199\pi}{100}\right)+\left(\sin\frac{2\pi}{100}+\sin\frac{198\pi}{100}\right)+\cdots+\left(\sin\frac{99\pi}{100}+\sin\frac{101\pi}{100}\right)+\sin\frac{100\pi}{100}+\sin\frac{200\pi}{100}$$

Each grouped sum is of the form $\sin x+\sin y$ with $x+y=2\pi$ so that $\dfrac{x+y}{2}=\dfrac{2\pi}{2}=\pi$. We now derive a useful identity from the product-to-sum identity $\sin x\,\cos y=\frac{1}{2}\left[\sin(x+y)+\sin(x-y)\right]$. If in this identity we replace x with $\dfrac{x+y}{2}$ and y with $\dfrac{x-y}{2}$, we have

$$\sin\left(\frac{x+y}{2}\right)\cos\left(\frac{x-y}{2}\right)=\frac{1}{2}\left[\sin\left(\frac{x+y}{2}+\frac{x-y}{2}\right)+\sin\left(\frac{x+y}{2}-\frac{x-y}{2}\right)\right]=\frac{1}{2}(\sin x+\sin y)$$

Multiplication of the left and right members of this equality by 2 gives the sum-to-product identity $\sin x+\sin y=2\sin\left(\dfrac{x+y}{2}\right)\cos\left(\dfrac{x+y}{2}\right)$. Using this sum-to-product identity, we have each grouped sum equal to 0, since $\sin\left(\dfrac{x+y}{2}\right)=\sin\pi=0$ is always a factor of the right side. Since $\sin\dfrac{100\pi}{100}=\sin\pi=0$ and $\sin\dfrac{200\pi}{100}=\sin 2\pi=0$, the sum of the given expression is 0.

Another approach: Since the sine function is odd, $\sin(-x)=-\sin x$. Because the period of the sine function is 2π, we have $\sin(-x+2\pi)=-\sin x$. Multiplying each side by -1 and rearranging, we have $\sin x=-\sin(2\pi-x)$. This means that $\sin\dfrac{\pi}{100}=-\sin\left(2\pi-\dfrac{\pi}{100}\right)=-\sin\dfrac{199\pi}{100}$, $\sin\dfrac{2\pi}{100}=-\sin\left(2\pi-\dfrac{2\pi}{100}\right)=\sin\dfrac{198\pi}{100}$, and so on, until we have $\sin\dfrac{99\pi}{100}=-\sin\left(2\pi-\dfrac{99\pi}{100}\right)=-\sin\dfrac{101\pi}{100}$. As before we rearrange terms to write the given expression as

$$\left(\sin\frac{\pi}{100}+\sin\frac{199\pi}{100}\right)+\left(\sin\frac{2\pi}{100}+\sin\frac{198\pi}{100}\right)+\cdots+\left(\sin\frac{99\pi}{100}+\sin\frac{101\pi}{100}\right)+\sin\frac{100\pi}{100}+\sin\frac{200\pi}{100}$$

Each sum in parentheses is 0 since the two terms are opposites, and the last two terms again reduce to $\sin\pi$ and $\sin 2\pi$, respectively, each also 0. Thus, the value of the original expression is 0.

14. (a)
$$\begin{aligned}f(-x)&=\ln\left(-x+\sqrt{(-x)^2+1}\right)=\ln\left(-x+\sqrt{x^2+1}\cdot\frac{-x-\sqrt{x^2+1}}{-x-\sqrt{x^2+1}}\right)\\&=\ln\left(\frac{x^2-(x^2+1)}{-x-\sqrt{x^2+1}}\right)=\ln\left(\frac{-1}{-x-\sqrt{x^2+1}}\right)=\ln\left(\frac{1}{x+\sqrt{x^2+1}}\right)\\&=\ln 1-\ln\left(x+\sqrt{x^2+1}\right)=-\ln\left(x+\sqrt{x^2-1}\right)=-f(x)\end{aligned}$$

(b) $y=\ln\left(x+\sqrt{x^2+1}\right)$. Interchanging x and y, we get $x=\ln\left(y+\sqrt{y^2+1}\right)\;\Rightarrow\;e^x=y+\sqrt{y^2+1}\;\Rightarrow$

$$e^x-y=\sqrt{y^2+1}\;\Rightarrow\;e^{2x}-2ye^x+y^2=y^2+1\;\Rightarrow\;e^{2x}-1=2ye^x\;\Rightarrow\;y=\frac{e^{2x}-1}{2e^x}=f^{-1}(x).$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $\ln\left(x^2-2x-2\right) \le 0 \quad \Rightarrow \quad x^2-2x-2 \le e^0 = 1 \quad \Rightarrow \quad x^2-2x-3 \le 0 \quad \Rightarrow \quad (x-3)(x+1) \le 0 \quad \Rightarrow \quad x \in [-1,3]$.

Since the argument must be positive, $x^2-2x-2>0 \quad \Rightarrow \quad \left[x-\left(1-\sqrt{3}\right)\right]\left[x-\left(1+\sqrt{3}\right)\right]>0 \quad \Rightarrow$

$x \in \left(-\infty, 1-\sqrt{3}\right) \cup \left(1+\sqrt{3}, \infty\right)$. The intersection of these intervals is $\left[-1, 1-\sqrt{3}\right) \cup \left(1+\sqrt{3}, 3\right]$.

16. Assume that $\log_2 5$ is rational. Then $\log_2 5 = m/n$ for natural numbers m and n. Changing to exponential form gives us $2^{m/n} = 5$ and then raising both sides to the nth power gives $2^m = 5^n$. But 2^m is even and 5^n is odd. We have arrived at a contradiction, so we conclude that our hypothesis, that $\log_2 5$ is rational, is false. Thus, $\log_2 5$ is irrational.

17. Let d be the distance traveled on each half of the trip. Let t_1 and t_2 be the times taken for the first and second halves of the trip. For the first half of the trip we have $t_1 = d/30$ and for the second half we have $t_2 = d/60$. Thus, the average speed for the entire trip is $\dfrac{\text{total distance}}{\text{total time}} = \dfrac{2d}{t_1+t_2} = \dfrac{2d}{\dfrac{d}{30}+\dfrac{d}{60}} \cdot \dfrac{60}{60} = \dfrac{120d}{2d+d} = \dfrac{120d}{3d} = 40$. The average speed for the entire trip is 40 mi/h.

18. Let $f(x) = \sin x$, $g(x) = x$, and $h(x) = x$. Then the left-hand side of the equation is $[f \circ (g+h)](x) = \sin(x+x) = \sin 2x = 2\sin x \cos x$; and the right-hand side is $(f \circ g)(x) + (f \circ h)(x) = \sin x + \sin x = 2\sin x$. The two sides are not equal, so the given statement is false.

19. Let S_n be the statement that $7^n - 1$ is divisible by 6.

- S_1 is true because $7^1 - 1 = 6$ is divisible by 6.
- Assume S_k is true, that is, $7^k - 1$ is divisible by 6. In other words, $7^k - 1 = 6m$ for some positive integer m. Then $7^{k+1} - 1 = 7^k \cdot 7 - 1 = (6m+1)\cdot 7 - 1 = 42m + 6 = 6(7m+1)$, which is divisible by 6, so S_{k+1} is true.
- Therefore, by mathematical induction, $7^n - 1$ is divisible by 6 for every positive integer n.

20. Let S_n be the statement that $1+3+5+\cdots+(2n-1) = n^2$.

- S_1 is true because $[2(1)-1] = 1 = 1^2$.
- Assume S_k is true, that is, $1+3+5+\cdots+(2k-1) = k^2$. Then

  $1+3+5+\cdots+(2k-1)+[2(k+1)-1] = 1+3+5+\cdots+(2k-1)+(2k+1) = k^2+(2k+1) = (k+1)^2$

  which shows that S_{k+1} is true.
- Therefore, by mathematical induction, $1+3+5+\cdots+(2n-1) = n^2$ for every positive integer n.

21. $f_0(x) = x^2$ and $f_{n+1}(x) = f_0(f_n(x))$ for $n = 0, 1, 2, \ldots$.

$f_1(x) = f_0(f_0(x)) = f_0\left(x^2\right) = \left(x^2\right)^2 = x^4$, $f_2(x) = f_0(f_1(x)) = f_0(x^4) = (x^4)^2 = x^8$,

$f_3(x) = f_0(f_2(x)) = f_0(x^8) = (x^8)^2 = x^{16}, \ldots$. Thus, a general formula is $f_n(x) = x^{2^{n+1}}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. (a) $f_0(x) = 1/(2-x)$ and $f_{n+1} = f_0 \circ f_n$ for $n = 0, 1, 2, \ldots$.

$$f_1(x) = f_0\left(\frac{1}{2-x}\right) = \frac{1}{2 - \dfrac{1}{2-x}} = \frac{2-x}{2(2-x)-1} = \frac{2-x}{3-2x},$$

$$f_2(x) = f_0\left(\frac{2-x}{3-2x}\right) = \frac{1}{2 - \dfrac{2-x}{3-2x}} = \frac{3-2x}{2(3-2x)-(2-x)} = \frac{3-2x}{4-3x},$$

$$f_3(x) = f_0\left(\frac{3-2x}{4-3x}\right) = \frac{1}{2 - \dfrac{3-2x}{4-3x}} = \frac{4-3x}{2(4-3x)-(3-2x)} = \frac{4-3x}{5-4x}, \ldots$$

Thus, we conjecture that the general formula is $f_n(x) = \dfrac{n+1-nx}{n+2-(n+1)x}$.

To prove this, we use the Principle of Mathematical Induction. We have already verified that f_n is true for $n = 1$.

Assume that the formula is true for $n = k$; that is, $f_k(x) = \dfrac{k+1-kx}{k+2-(k+1)x}$. Then

$$f_{k+1}(x) = (f_0 \circ f_k)(x) = f_0(f_k(x)) = f_0\left(\frac{k+1-kx}{k+2-(k+1)x}\right) = \frac{1}{2 - \dfrac{k+1-kx}{k+2-(k+1)x}}$$

$$= \frac{k+2-(k+1)x}{2\,[k+2-(k+1)x] - (k+1-kx)} = \frac{k+2-(k+1)x}{k+3-(k+2)x}$$

This shows that the formula for f_n is true for $n = k+1$. Therefore, by mathematical induction, the formula is true for all positive integers n.

(b) From the graph, we can make several observations:

- The values at each fixed $x = a$ keep increasing as n increases.
- The vertical asymptote gets closer to $x = 1$ as n increases.
- The horizontal asymptote gets closer to $y = 1$ as n increases.
- The x-intercept for f_{n+1} is the value of the vertical asymptote for f_n.
- The y-intercept for f_n is the value of the horizontal asymptote for f_{n+1}.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2 ☐ LIMITS AND DERIVATIVES

2.1 The Tangent and Velocity Problems

1. (a) Using $P(15, 250)$, we construct the following table:

t	Q	slope $= m_{PQ}$
5	$(5, 694)$	$\frac{694-250}{5-15} = -\frac{444}{10} = -44.4$
10	$(10, 444)$	$\frac{444-250}{10-15} = -\frac{194}{5} = -38.8$
20	$(20, 111)$	$\frac{111-250}{20-15} = -\frac{139}{5} = -27.8$
25	$(25, 28)$	$\frac{28-250}{25-15} = -\frac{222}{10} = -22.2$
30	$(30, 0)$	$\frac{0-250}{30-15} = -\frac{250}{15} = -16.\overline{6}$

(b) Using the values of t that correspond to the points closest to P ($t = 10$ and $t = 20$), we have

$$\frac{-38.8 + (-27.8)}{2} = -33.3$$

(c) From the graph, we can estimate the slope of the tangent line at P to be $\frac{-300}{9} = -33.\overline{3}$.

2. (a) (i) On the interval $[0, 40]$, slope $= \dfrac{7398 - 3438}{40 - 0} = 99$.

(ii) On the interval $[10, 20]$, slope $= \dfrac{5622 - 4559}{20 - 10} = 106.3$.

(iii) On the interval $[20, 30]$, slope $= \dfrac{6536 - 5622}{30 - 20} = 91.4$.

The slopes represent the average number of steps per minute the student walked during the respective time intervals.

(b) Averaging the slopes of the secant lines corresponding to the intervals immediately before and after $t = 20$, we have

$$\frac{106.3 + 91.4}{2} = 98.85$$

The student's walking pace is approximately 99 steps per minute at 3:20 PM.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a) $y = \dfrac{1}{1-x}$, $P(2, -1)$

	x	$Q(x, 1/(1-x))$	m_{PQ}
(i)	1.5	$(1.5, -2)$	2
(ii)	1.9	$(1.9, -1.111\,111)$	1.111 111
(iii)	1.99	$(1.99, -1.010\,101)$	1.010 101
(iv)	1.999	$(1.999, -1.001\,001)$	1.001 001
(v)	2.5	$(2.5, -0.666\,667)$	0.666 667
(vi)	2.1	$(2.1, -0.909\,091)$	0.909 091
(vii)	2.01	$(2.01, -0.990\,099)$	0.990 099
(viii)	2.001	$(2.001, -0.999\,001)$	0.999 001

(b) The slope appears to be 1.

(c) Using $m = 1$, an equation of the tangent line to the curve at $P(2, -1)$ is $y - (-1) = 1(x - 2)$, or $y = x - 3$.

4. (a) $y = \cos \pi x$, $P(0.5, 0)$

	x	Q	m_{PQ}
(i)	0	$(0, 1)$	-2
(ii)	0.4	$(0.4, 0.309017)$	-3.090170
(iii)	0.49	$(0.49, 0.031411)$	-3.141076
(iv)	0.499	$(0.499, 0.003142)$	-3.141587
(v)	1	$(1, -1)$	-2
(vi)	0.6	$(0.6, -0.309017)$	-3.090170
(vii)	0.51	$(0.51, -0.031411)$	-3.141076
(viii)	0.501	$(0.501, -0.003142)$	-3.141587

(b) The slope appears to be $-\pi$.

(c) $y - 0 = -\pi(x - 0.5)$ or $y = -\pi x + \frac{1}{2}\pi$.

(d)

5. (a) $y = y(t) = 275 - 16t^2$. At $t = 4$, $y = 275 - 16(4)^2 = 19$. The average velocity between times 4 and $4 + h$ is

$$v_{\text{avg}} = \frac{y(4+h) - y(4)}{(4+h) - 4} = \frac{\left[275 - 16(4+h)^2\right] - 19}{h} = \frac{-128h - 16h^2}{h} = -128 - 16h \quad \text{if } h \neq 0$$

(i) 0.1 seconds: $h = 0.1$, $v_{\text{avg}} = -129.6$ ft/s

(ii) 0.05 seconds: $h = 0.05$, $v_{\text{avg}} = -128.8$ ft/s

(iii) 0.01 seconds: $h = 0.01$, $v_{\text{avg}} = -128.16$ ft/s

(b) The instantaneous velocity when $t = 4$ (h approaches 0) is -128 ft/s.

6. (a) $y = y(t) = 10t - 1.86t^2$. At $t = 1$, $y = 10(1) - 1.86(1)^2 = 8.14$. The average velocity between times 1 and $1 + h$ is

$$v_{\text{avg}} = \frac{y(1+h) - y(1)}{(1+h) - 1} = \frac{\left[10(1+h) - 1.86(1+h)^2\right] - 8.14}{h} = \frac{6.28h - 1.86h^2}{h} = 6.28 - 1.86h, \text{ if } h \neq 0.$$

(i) $[1, 2]$: $h = 1$, $v_{\text{avg}} = 4.42$ m/s

(ii) $[1, 1.5]$: $h = 0.5$, $v_{\text{avg}} = 5.35$ m/s

(iii) $[1, 1.1]$: $h = 0.1$, $v_{\text{avg}} = 6.094$ m/s

(iv) $[1, 1.01]$: $h = 0.01$, $v_{\text{avg}} = 6.2614$ m/s

(v) $[1, 1.001]$: $h = 0.001$, $v_{\text{avg}} = 6.27814$ m/s

(b) The instantaneous velocity when $t = 1$ (h approaches 0) is 6.28 m/s.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. (a) (i) On the interval $[2,4]$, $v_{\text{avg}} = \dfrac{s(4)-s(2)}{4-2} = \dfrac{79.2-20.6}{2} = 29.3$ ft/s.

(ii) On the interval $[3,4]$, $v_{\text{avg}} = \dfrac{s(4)-s(3)}{4-3} = \dfrac{79.2-46.5}{1} = 32.7$ ft/s.

(iii) On the interval $[4,5]$, $v_{\text{avg}} = \dfrac{s(5)-s(4)}{5-4} = \dfrac{124.8-79.2}{1} = 45.6$ ft/s.

(iv) On the interval $[4,6]$, $v_{\text{avg}} = \dfrac{s(6)-s(4)}{6-4} = \dfrac{176.7-79.2}{2} = 48.75$ ft/s.

(b) Using the points $(2,16)$ and $(5,105)$ from the approximate tangent line, the instantaneous velocity at $t=3$ is about $\dfrac{105-16}{5-2} = \dfrac{89}{3} \approx 29.7$ ft/s.

8. (a) (i) $s = s(t) = 2\sin \pi t + 3\cos \pi t$. On the interval $[1,2]$, $v_{\text{avg}} = \dfrac{s(2)-s(1)}{2-1} = \dfrac{3-(-3)}{1} = 6$ cm/s.

(ii) On the interval $[1,1.1]$, $v_{\text{avg}} = \dfrac{s(1.1)-s(1)}{1.1-1} \approx \dfrac{-3.471-(-3)}{0.1} = -4.71$ cm/s.

(iii) On the interval $[1,1.01]$, $v_{\text{avg}} = \dfrac{s(1.01)-s(1)}{1.01-1} \approx \dfrac{-3.0613-(-3)}{0.01} = -6.13$ cm/s.

(iv) On the interval $[1,1.001]$, $v_{\text{avg}} = \dfrac{s(1.001)-s(1)}{1.001-1} \approx \dfrac{-3.00627-(-3)}{0.001} = -6.27$ cm/s.

(b) The instantaneous velocity of the particle when $t=1$ appears to be about -6.3 cm/s.

9. (a) For the curve $y = \sin(10\pi/x)$ and the point $P(1,0)$:

x	Q	m_{PQ}
2	$(2,0)$	0
1.5	$(1.5, 0.8660)$	1.7321
1.4	$(1.4, -0.4339)$	-1.0847
1.3	$(1.3, -0.8230)$	-2.7433
1.2	$(1.2, 0.8660)$	4.3301
1.1	$(1.1, -0.2817)$	-2.8173

x	Q	m_{PQ}
0.5	$(0.5, 0)$	0
0.6	$(0.6, 0.8660)$	-2.1651
0.7	$(0.7, 0.7818)$	-2.6061
0.8	$(0.8, 1)$	-5
0.9	$(0.9, -0.3420)$	3.4202

As x approaches 1, the slopes do not appear to be approaching any particular value.

(b)

We see that problems with estimation are caused by the frequent oscillations of the graph. The tangent is so steep at P that we need to take x-values much closer to 1 in order to get accurate estimates of its slope.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) If we choose $x = 1.001$, then the point Q is $(1.001, -0.0314)$ and $m_{PQ} \approx -31.3794$. If $x = 0.999$, then Q is $(0.999, 0.0314)$ and $m_{PQ} = -31.4422$. The average of these slopes is -31.4108. So we estimate that the slope of the tangent line at P is about -31.4.

2.2 The Limit of a Function

1. As x approaches 2, $f(x)$ approaches 5. [Or, the values of $f(x)$ can be made as close to 5 as we like by taking x sufficiently close to 2 (but $x \neq 2$).] Yes, the graph could have a hole at $(2, 5)$ and be defined such that $f(2) = 3$.

2. As x approaches 1 from the left, $f(x)$ approaches 3; and as x approaches 1 from the right, $f(x)$ approaches 7. No, the limit does not exist because the left- and right-hand limits are different.

3. (a) $\lim\limits_{x \to -3} f(x) = \infty$ means that the values of $f(x)$ can be made arbitrarily large (as large as we please) by taking x sufficiently close to -3 (but not equal to -3).

(b) $\lim\limits_{x \to 4^+} f(x) = -\infty$ means that the values of $f(x)$ can be made arbitrarily large negative by taking x sufficiently close to 4 through values larger than 4.

4. (a) As x approaches 2 from the left, the values of $f(x)$ approach 3, so $\lim\limits_{x \to 2^-} f(x) = 3$.

(b) As x approaches 2 from the right, the values of $f(x)$ approach 1, so $\lim\limits_{x \to 2^+} f(x) = 1$.

(c) $\lim\limits_{x \to 2} f(x)$ does not exist since the left-hand limit does not equal the right-hand limit.

(d) When $x = 2$, $y = 3$, so $f(2) = 3$.

(e) As x approaches 4, the values of $f(x)$ approach 4, so $\lim\limits_{x \to 4} f(x) = 4$.

(f) There is no value of $f(x)$ when $x = 4$, so $f(4)$ does not exist.

5. (a) As x approaches 1, the values of $f(x)$ approach 2, so $\lim\limits_{x \to 1} f(x) = 2$.

(b) As x approaches 3 from the left, the values of $f(x)$ approach 1, so $\lim\limits_{x \to 3^-} f(x) = 1$.

(c) As x approaches 3 from the right, the values of $f(x)$ approach 4, so $\lim\limits_{x \to 3^+} f(x) = 4$.

(d) $\lim\limits_{x \to 3} f(x)$ does not exist since the left-hand limit does not equal the right-hand limit.

(e) When $x = 3$, $y = 3$, so $f(3) = 3$.

6. (a) $h(x)$ approaches 4 as x approaches -3 from the left, so $\lim\limits_{x \to -3^-} h(x) = 4$.

(b) $h(x)$ approaches 4 as x approaches -3 from the right, so $\lim\limits_{x \to -3^+} h(x) = 4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\lim_{x \to -3} h(x) = 4$ because the limits in part (a) and part (b) are equal.

(d) $h(-3)$ is not defined, so it doesn't exist.

(e) $h(x)$ approaches 1 as x approaches 0 from the left, so $\lim_{x \to 0^-} h(x) = 1$.

(f) $h(x)$ approaches -1 as x approaches 0 from the right, so $\lim_{x \to 0^+} h(x) = -1$.

(g) $\lim_{x \to 0} h(x)$ does not exist because the limits in part (e) and part (f) are not equal.

(h) $h(0) = 1$ since the point $(0, 1)$ is on the graph of h.

(i) Since $\lim_{x \to 2^-} h(x) = 2$ and $\lim_{x \to 2^+} h(x) = 2$, we have $\lim_{x \to 2} h(x) = 2$.

(j) $h(2)$ is not defined, so it doesn't exist.

(k) $h(x)$ approaches 3 as x approaches 5 from the right, so $\lim_{x \to 5^+} h(x) = 3$.

(l) $h(x)$ does not approach any one number as x approaches 5 from the left, so $\lim_{x \to 5^-} h(x)$ does not exist.

7. (a) $\lim_{x \to 4^-} g(x) \neq \lim_{x \to 4^+} g(x)$, so $\lim_{x \to 4} g(x)$ does not exist. However, there is a point on the graph representing $g(4)$. Thus, $a = 4$ satisfies the given description.

(b) $\lim_{x \to 5^-} g(x) = \lim_{x \to 5^+} g(x)$, so $\lim_{x \to 5} g(x)$ exists. However, $g(5)$ is not defined. Thus, $a = 5$ satisfies the given description.

(c) From part (a), $a = 4$ satisfies the given description. Also, $\lim_{x \to 2^-} g(x)$ and $\lim_{x \to 2^+} g(x)$ exist, but $\lim_{x \to 2^-} g(x) \neq \lim_{x \to 2^+} g(x)$. Thus, $\lim_{x \to 2} g(x)$ does not exist, and $a = 2$ also satisfies the given description.

(d) $\lim_{x \to 4^+} g(x) = g(4)$, but $\lim_{x \to 4^-} g(x) \neq g(4)$. Thus, $a = 4$ satisfies the given description.

8. (a) $\lim_{x \to -3} A(x) = \infty$ (b) $\lim_{x \to 2^-} A(x) = -\infty$

(c) $\lim_{x \to 2^+} A(x) = \infty$ (d) $\lim_{x \to -1} A(x) = -\infty$

(e) The equations of the vertical asymptotes are $x = -3$, $x = -1$ and $x = 2$.

9. (a) $\lim_{x \to -7} f(x) = -\infty$ (b) $\lim_{x \to -3} f(x) = \infty$ (c) $\lim_{x \to 0} f(x) = \infty$

(d) $\lim_{x \to 6^-} f(x) = -\infty$ (e) $\lim_{x \to 6^+} f(x) = \infty$

(f) The equations of the vertical asymptotes are $x = -7$, $x = -3$, $x = 0$, and $x = 6$.

10. $\lim_{t \to 12^-} f(t) = 150$ mg and $\lim_{t \to 12^+} f(t) = 300$ mg. These limits show that there is an abrupt change in the amount of drug in the patient's bloodstream at $t = 12$ h. The left-hand limit represents the amount of the drug just before the fourth injection. The right-hand limit represents the amount of the drug just after the fourth injection.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11.

From the graph of f we see that $\lim\limits_{x\to 0^-} f(x) = 1$, but $\lim\limits_{x\to 0^+} f(x) = -1$, so $\lim\limits_{x\to a} f(x)$ does not exist for $a = 0$. However, $\lim\limits_{x\to a} f(x)$ exists for all other values of a. Thus, $\lim\limits_{x\to a} f(x)$ exists for all a in $(-\infty, 0) \cup (0, \infty)$.

12.

From the graph of f we see that $\lim\limits_{x\to 2^-} f(x) = 2$, but $\lim\limits_{x\to 2^+} f(x) = 1$, so $\lim\limits_{x\to a} f(x)$ does not exist for $a = 2$. However, $\lim\limits_{x\to a} f(x)$ exists for all other values of a. Thus, $\lim\limits_{x\to a} f(x)$ exists for all a in $(-\infty, 2) \cup (2, \infty)$.

13. (a) From the graph, $\lim\limits_{x\to 0^-} f(x) = -1$.

(b) From the graph, $\lim\limits_{x\to 0^+} f(x) = 1$.

(c) Since $\lim\limits_{x\to 0^-} f(x) \neq \lim\limits_{x\to 0^+} f(x)$, $\lim\limits_{x\to 0} f(x)$ does not exist.

$f(x) = x\sqrt{1 + x^{-2}}$

14. (a) From the graph, $\lim\limits_{x\to 0^-} f(x) = -2$.

(b) From the graph, $\lim\limits_{x\to 0^+} f(x) = 1$.

(c) Since $\lim\limits_{x\to 0^-} f(x) \neq \lim\limits_{x\to 0^+} f(x)$, $\lim\limits_{x\to 0} f(x)$ does not exist.

$f(x) = \dfrac{e^{1/x} - 2}{e^{1/x} + 1}$

15. $\lim\limits_{x\to 1^-} f(x) = 3$, $\lim\limits_{x\to 1^+} f(x) = 0$, $f(1) = 2$

16. $\lim\limits_{x\to 0} f(x) = 4$, $\lim\limits_{x\to 8^-} f(x) = 1$, $\lim\limits_{x\to 8^+} f(x) = -3$,

$f(0) = 6$, $f(8) = -1$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\lim_{x\to -1^-} f(x) = 0$, $\lim_{x\to -1^+} f(x) = 1$, $\lim_{x\to 2} f(x) = 3$,

$f(-1) = 2$, $f(2) = 1$

18. $\lim_{x\to -3^-} f(x) = 3$, $\lim_{x\to -3^+} f(x) = 2$, $\lim_{x\to 3^-} f(x) = -1$,

$\lim_{x\to 3^+} f(x) = 2$, $f(-3) = 2$, $f(3) = 0$

19. For $f(x) = \dfrac{x^2 - 3x}{x^2 - 9}$:

x	$f(x)$
3.1	0.508 197
3.05	0.504 132
3.01	0.500 832
3.001	0.500 083
3.0001	0.500 008

x	$f(x)$
2.9	0.491 525
2.95	0.495 798
2.99	0.499 165
2.999	0.499 917
2.9999	0.499 992

It appears that $\lim_{x\to 3} \dfrac{x^2 - 3x}{x^2 - 9} = \dfrac{1}{2}$.

20. For $f(x) = \dfrac{x^2 - 3x}{x^2 - 9}$:

x	$f(x)$
−2.5	−5
−2.9	−29
−2.95	−59
−2.99	−299
−2.999	−2999
−2.9999	−29,999

x	$f(x)$
−3.5	7
−3.1	31
−3.05	61
−3.01	301
−3.001	3001
−3.0001	30,001

It appears that $\lim_{x\to -3^+} f(x) = -\infty$ and that

$\lim_{x\to -3^-} f(x) = \infty$, so $\lim_{x\to -3} \dfrac{x^2 - 3x}{x^2 - 9}$ does not exist.

21. For $f(t) = \dfrac{e^{5t} - 1}{t}$:

t	$f(t)$
0.5	22.364 988
0.1	6.487 213
0.01	5.127 110
0.001	5.012 521
0.0001	5.001 250

t	$f(t)$
−0.5	1.835 830
−0.1	3.934 693
−0.01	4.877 058
−0.001	4.987 521
−0.0001	4.998 750

It appears that $\lim_{t\to 0} \dfrac{e^{5t} - 1}{t} = 5$.

22. For $f(h) = \dfrac{(2+h)^5 - 32}{h}$:

h	$f(h)$
0.5	131.312 500
0.1	88.410 100
0.01	80.804 010
0.001	80.080 040
0.0001	80.008 000

h	$f(h)$
−0.5	48.812 500
−0.1	72.390 100
−0.01	79.203 990
−0.001	79.920 040
−0.0001	79.992 000

It appears that $\lim_{h\to 0} \dfrac{(2+h)^5 - 32}{h} = 80$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. For $f(x) = \dfrac{\ln x - \ln 4}{x - 4}$:

x	$f(x)$
3.9	0.253 178
3.99	0.250 313
3.999	0.250 031
3.9999	0.250 003

x	$f(x)$
4.1	0.246 926
4.01	0.249 688
4.001	0.249 969
4.0001	0.249 997

It appears that $\lim\limits_{x \to 4} f(x) = 0.25$. The graph confirms that result.

24. For $f(p) = \dfrac{1 + p^9}{1 + p^{15}}$:

p	$f(p)$
-1.1	0.427 397
-1.01	0.582 008
-1.001	0.598 200
-1.0001	0.599 820

p	$f(p)$
-0.9	0.771 405
-0.99	0.617 992
-0.999	0.601 800
-0.9999	0.600 180

It appears that $\lim\limits_{p \to -1} f(p) = 0.6$. The graph confirms that result.

25. For $f(\theta) = \dfrac{\sin 3\theta}{\tan 2\theta}$:

θ	$f(\theta)$
± 0.1	1.457 847
± 0.01	1.499 575
± 0.001	1.499 996
± 0.0001	1.500 000

It appears that $\lim\limits_{\theta \to 0} \dfrac{\sin 3\theta}{\tan 2\theta} = 1.5$.

The graph confirms that result.

26. For $f(t) = \dfrac{5^t - 1}{t}$:

t	$f(t)$
0.1	1.746 189
0.01	1.622 459
0.001	1.610 734
0.0001	1.609 567

t	$f(t)$
-0.1	1.486 601
-0.01	1.596 556
-0.001	1.608 143
-0.0001	1.609 308

It appears that $\lim\limits_{t \to 0} f(t) \approx 1.6094$. The graph confirms that result.

27. For $f(x) = x^x$:

x	$f(x)$
0.1	0.794 328
0.01	0.954 993
0.001	0.993 116
0.0001	0.999 079

It appears that $\lim\limits_{x \to 0^+} f(x) = 1$.

The graph confirms that result.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. For $f(x) = x^2 \ln x$:

x	$f(x)$
0.1	$-0.023\,026$
0.01	$-0.000\,461$
0.001	$-0.000\,007$
0.0001	$-0.000\,000$

It appears that $\lim\limits_{x \to 0^+} f(x) = 0$.

The graph confirms that result.

29. $\lim\limits_{x \to 5^+} \dfrac{x+1}{x-5} = \infty$ since the numerator is positive and the denominator approaches 0 from the positive side as $x \to 5^+$.

30. $\lim\limits_{x \to 5^-} \dfrac{x+1}{x-5} = -\infty$ since the numerator is positive and the denominator approaches 0 from the negative side as $x \to 5^-$.

31. $\lim\limits_{x \to 2} \dfrac{x^2}{(x-2)^2} = \infty$ since the numerator is positive and the denominator approaches 0 through positive values as $x \to 2$.

32. $\lim\limits_{x \to 3^-} \dfrac{\sqrt{x}}{(x-3)^5} = -\infty$ since the numerator is positive and the denominator approaches 0 from the negative side as $x \to 3^-$.

33. $\lim\limits_{x \to 1^+} \ln(\sqrt{x} - 1) = -\infty$ since $\sqrt{x} - 1 \to 0^+$ as $x \to 1^+$.

34. $\lim\limits_{x \to 0^+} \ln(\sin x) = -\infty$ since $\sin x \to 0^+$ as $x \to 0^+$.

35. $\lim\limits_{x \to (\pi/2)^+} \dfrac{1}{x} \sec x = -\infty$ since $\dfrac{1}{x}$ is positive and $\sec x \to -\infty$ as $x \to (\pi/2)^+$.

36. $\lim\limits_{x \to \pi^-} x \cot x = -\infty$ since x is positive and $\cot x \to -\infty$ as $x \to \pi^-$.

37. $\lim\limits_{x \to 1} \dfrac{x^2 + 2x}{x^2 - 2x + 1} = \lim\limits_{x \to 1} \dfrac{x^2 + 2x}{(x-1)^2} = \infty$ since the numerator is positive and the denominator approaches 0 through positive values as $x \to 1$.

38. $\lim\limits_{x \to 3^-} \dfrac{x^2 + 4x}{x^2 - 2x - 3} = \lim\limits_{x \to 3^-} \dfrac{x^2 + 4x}{(x-3)(x+1)} = -\infty$ since the numerator is positive and the denominator approaches 0 through negative values as $x \to 3^-$.

39. $\lim\limits_{x \to 0} (\ln x^2 - x^{-2}) = -\infty$ since $\ln x^2 \to -\infty$ and $x^{-2} \to \infty$ as $x \to 0$.

40. $\lim\limits_{x \to 0^+} \left(\dfrac{1}{x} - \ln x \right) = \infty$ since $\dfrac{1}{x} \to \infty$ and $\ln x \to -\infty$ as $x \to 0^+$.

41. The denominator of $f(x) = \dfrac{x-1}{2x+4}$ is equal to 0 when $x = -2$ (and the numerator is not), so $x = -2$ is the vertical asymptote of the function.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. (a) The denominator of $y = \dfrac{x^2+1}{3x-2x^2} = \dfrac{x^2+1}{x(3-2x)}$ is equal to zero when $x = 0$ and $x = \frac{3}{2}$ (and the numerator is not), so $x = 0$ and $x = 1.5$ are vertical asymptotes of the function.

(b)

43. (a) $f(x) = \dfrac{1}{x^3-1}$.

From these calculations, it seems that

$\lim\limits_{x\to 1^-} f(x) = -\infty$ and $\lim\limits_{x\to 1^+} f(x) = \infty$.

x	$f(x)$
0.5	−1.14
0.9	−3.69
0.99	−33.7
0.999	−333.7
0.9999	−3333.7
0.99999	−33,333.7

x	$f(x)$
1.5	0.42
1.1	3.02
1.01	33.0
1.001	333.0
1.0001	3333.0
1.00001	33,333.3

(b) If x is slightly smaller than 1, then $x^3 - 1$ will be a negative number close to 0, and the reciprocal of $x^3 - 1$, that is, $f(x)$, will be a negative number with large absolute value. So $\lim\limits_{x\to 1^-} f(x) = -\infty$.

If x is slightly larger than 1, then $x^3 - 1$ will be a small positive number, and its reciprocal, $f(x)$, will be a large positive number. So $\lim\limits_{x\to 1^+} f(x) = \infty$.

(c) It appears from the graph of f that

$\lim\limits_{x\to 1^-} f(x) = -\infty$ and $\lim\limits_{x\to 1^+} f(x) = \infty$.

44. (a) From the graphs, it seems that $\lim\limits_{x\to 0} \dfrac{\cos 2x - \cos x}{x^2} = -1.5$.

(b)

x	$f(x)$
±0.1	−1.493 759
±0.01	−1.499 938
±0.001	−1.499 999
±0.0001	−1.500 000

45. (a) Let $h(x) = (1+x)^{1/x}$.

x	$h(x)$
−0.001	2.71964
−0.0001	2.71842
−0.00001	2.71830
−0.000001	2.71828
0.000001	2.71828
0.00001	2.71827
0.0001	2.71815
0.001	2.71692

(b)

It appears that $\lim\limits_{x\to 0} (1+x)^{1/x} \approx 2.71828$, which is approximately e.

In Section 3.6 we will see that the value of the limit is exactly e.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. (a)

No, because the calculator-produced graph of $f(x) = e^x + \ln|x-4|$ looks like an exponential function, but the graph of f has an infinite discontinuity at $x = 4$. A second graph, obtained by increasing the `numpoints` option in Maple, begins to reveal the discontinuity at $x = 4$.

(b) There isn't a single graph that shows all the features of f. Several graphs are needed since f looks like $\ln|x-4|$ for large negative values of x and like e^x for $x > 5$, but yet has the infinite discontiuity at $x = 4$.

A hand-drawn graph, though distorted, might be better at revealing the main features of this function.

47. For $f(x) = x^2 - (2^x/1000)$:

(a)

x	$f(x)$
1	0.998 000
0.8	0.638 259
0.6	0.358 484
0.4	0.158 680
0.2	0.038 851
0.1	0.008 928
0.05	0.001 465

It appears that $\lim_{x\to 0} f(x) = 0$.

(b)

x	$f(x)$
0.04	0.000 572
0.02	−0.000 614
0.01	−0.000 907
0.005	−0.000 978
0.003	−0.000 993
0.001	−0.001 000

It appears that $\lim_{x\to 0} f(x) = -0.001$.

48. For $h(x) = \dfrac{\tan x - x}{x^3}$:

(a)

x	$h(x)$
1.0	0.557 407 73
0.5	0.370 419 92
0.1	0.334 672 09
0.05	0.333 667 00
0.01	0.333 346 67
0.005	0.333 336 67

(b) It seems that $\lim_{x\to 0} h(x) = \frac{1}{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c)

x	$h(x)$
0.001	0.333 333 50
0.0005	0.333 333 44
0.0001	0.333 330 00
0.00005	0.333 336 00
0.00001	0.333 000 00
0.000001	0.000 000 00

Here the values will vary from one calculator to another. Every calculator will eventually give *false values*.

(d) As in part (c), when we take a small enough viewing rectangle we get incorrect output.

49.

There appear to be vertical asymptotes of the curve $y = \tan(2\sin x)$ at $x \approx \pm 0.90$ and $x \approx \pm 2.24$. To find the exact equations of these asymptotes, we note that the graph of the tangent function has vertical asymptotes at $x = \frac{\pi}{2} + \pi n$. Thus, we must have $2\sin x = \frac{\pi}{2} + \pi n$, or equivalently, $\sin x = \frac{\pi}{4} + \frac{\pi}{2}n$. Since $-1 \le \sin x \le 1$, we must have $\sin x = \pm\frac{\pi}{4}$ and so $x = \pm\sin^{-1}\frac{\pi}{4}$ (corresponding to $x \approx \pm 0.90$). Just as $150°$ is the reference angle for $30°$, $\pi - \sin^{-1}\frac{\pi}{4}$ is the reference angle for $\sin^{-1}\frac{\pi}{4}$. So $x = \pm\left(\pi - \sin^{-1}\frac{\pi}{4}\right)$ are also equations of vertical asymptotes (corresponding to $x \approx \pm 2.24$).

50. (a) For any positive integer n, if $x = \dfrac{1}{n\pi}$, then $f(x) = \tan\dfrac{1}{x} = \tan(n\pi) = 0$. (Remember that the tangent function has period π.)

(b) For any nonnegative number n, if $x = \dfrac{4}{(4n+1)\pi}$, then

$$f(x) = \tan\frac{1}{x} = \tan\frac{(4n+1)\pi}{4} = \tan\left(\frac{4n\pi}{4} + \frac{\pi}{4}\right) = \tan\left(n\pi + \frac{\pi}{4}\right) = \tan\frac{\pi}{4} = 1$$

(c) From part (a), $f(x) = 0$ infinitely often as $x \to 0$. From part (b), $f(x) = 1$ infinitely often as $x \to 0$. Thus, $\lim\limits_{x\to 0}\tan\dfrac{1}{x}$ does not exist since $f(x)$ does not get close to a fixed number as $x \to 0$.

51. $\lim\limits_{v\to c^-} m = \lim\limits_{v\to c^-}\dfrac{m_0}{\sqrt{1 - v^2/c^2}}$. As $v \to c^-$, $\sqrt{1 - v^2/c^2} \to 0^+$, and $m \to \infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2.3 Calculating Limits Using the Limit Laws

1. (a) $\lim\limits_{x\to 2}[f(x)+5g(x)] = \lim\limits_{x\to 2} f(x) + \lim\limits_{x\to 2}[5g(x)]$ [Limit Law 1]

$= \lim\limits_{x\to 2} f(x) + 5\lim\limits_{x\to 2} g(x)$ [Limit Law 3]

$= 4 + 5(-2) = -6$

(b) $\lim\limits_{x\to 2}[g(x)]^3 = \left[\lim\limits_{x\to 2} g(x)\right]^3$ [Limit Law 6]

$= (-2)^3 = -8$

(c) $\lim\limits_{x\to 2}\sqrt{f(x)} = \sqrt{\lim\limits_{x\to 2} f(x)}$ [Limit Law 7]

$= \sqrt{4} = 2$

(d) $\lim\limits_{x\to 2}\dfrac{3f(x)}{g(x)} = \dfrac{\lim\limits_{x\to 2}[3f(x)]}{\lim\limits_{x\to 2} g(x)}$ [Limit Law 5]

$= \dfrac{3\lim\limits_{x\to 2} f(x)}{\lim\limits_{x\to 2} g(x)}$ [Limit Law 3]

$= \dfrac{3(4)}{-2} = -6$

(e) Because the limit of the denominator is 0, we can't use Limit Law 5. The given limit, $\lim\limits_{x\to 2}\dfrac{g(x)}{h(x)}$, does not exist because the denominator approaches 0 while the numerator approaches a nonzero number.

(f) $\lim\limits_{x\to 2}\dfrac{g(x)\,h(x)}{f(x)} = \dfrac{\lim\limits_{x\to 2}[g(x)\,h(x)]}{\lim\limits_{x\to 2} f(x)}$ [Limit Law 5]

$= \dfrac{\lim\limits_{x\to 2} g(x)\cdot\lim\limits_{x\to 2} h(x)}{\lim\limits_{x\to 2} f(x)}$ [Limit Law 4]

$= \dfrac{-2\cdot 0}{4} = 0$

2. (a) $\lim\limits_{x\to 2}[f(x)+g(x)] = \lim\limits_{x\to 2} f(x) + \lim\limits_{x\to 2} g(x)$ [Limit Law 1]

$= -1 + 2$

$= 1$

(b) $\lim\limits_{x\to 0} f(x)$ exists, but $\lim\limits_{x\to 0} g(x)$ does not exist, so we cannot apply Limit Law 2 to $\lim\limits_{x\to 0}[f(x)-g(x)]$.

The limit does not exist.

(c) $\lim\limits_{x\to -1}[f(x)\,g(x)] = \lim\limits_{x\to -1} f(x)\cdot\lim\limits_{x\to -1} g(x)$ [Limit Law 4]

$= 1\cdot 2$

$= 2$

(d) $\lim\limits_{x\to 3} f(x) = 1$, but $\lim\limits_{x\to 3} g(x) = 0$, so we cannot apply Limit Law 5 to $\lim\limits_{x\to 3}\dfrac{f(x)}{g(x)}$. The limit does not exist.

Note: $\lim\limits_{x\to 3^-}\dfrac{f(x)}{g(x)} = \infty$ since $g(x)\to 0^+$ as $x\to 3^-$ and $\lim\limits_{x\to 3^+}\dfrac{f(x)}{g(x)} = -\infty$ since $g(x)\to 0^-$as $x\to 3^+$.

Therefore, the limit does not exist, even as an infinite limit.

(e) $\lim\limits_{x\to 2}\left[x^2 f(x)\right] = \lim\limits_{x\to 2} x^2\cdot\lim\limits_{x\to 2} f(x)$ [Limit Law 4]

$= 2^2\cdot(-1)$

$= -4$

(f) $f(-1) + \lim\limits_{x\to -1} g(x)$ is undefined since $f(-1)$ is not defined.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. $\lim_{x\to5}(4x^2-5x)=\lim_{x\to5}(4x^2)-\lim_{x\to5}(5x)$ [Limit Law 2]

$=4\lim_{x\to5}x^2-5\lim_{x\to5}x$ [3]

$=4(5^2)-5(5)$ [10, 9]

$=75$

4. $\lim_{x\to-3}(2x^3+6x^2-9)=\lim_{x\to-3}(2x^3)+\lim_{x\to-3}(6x^2)-\lim_{x\to-3}9$ [Limits Laws 1 and 2]

$=2\lim_{x\to-3}x^3+6\lim_{x\to-3}x^2-\lim_{x\to-3}9$ [3]

$=2(-3)^3+6(-3)^2-9$ [10, 8]

$=-9$

5. $\lim_{v\to2}(v^2+2v)(2v^3-5)=\lim_{v\to2}(v^2+2v)\cdot\lim_{v\to2}(2v^3-5)$ [Limit Law 4]

$=\left(\lim_{v\to2}v^2+\lim_{v\to2}2v\right)\left(\lim_{v\to2}2v^3-\lim_{v\to2}5\right)$ [1 and 2]

$=\left(\lim_{v\to2}v^2+2\lim_{v\to2}v\right)\left(2\lim_{v\to2}v^3-\lim_{v\to2}5\right)$ [3]

$=\left[2^2+2(2)\right]\left[2(2)^3-5\right]$ [10, 9, and 8]

$=(8)(11)=88$

6. $\lim_{t\to7}\dfrac{3t^2+1}{t^2-5t+2}=\dfrac{\lim_{t\to7}(3t^2+1)}{\lim_{t\to7}(t^2-5t+2)}$ [Limit Law 5]

$=\dfrac{\lim_{t\to7}3t^2+\lim_{t\to7}1}{\lim_{t\to7}t^2-\lim_{t\to7}5t+\lim_{t\to7}2}$ [1 and 2]

$=\dfrac{3\lim_{t\to7}t^2+\lim_{t\to7}1}{\lim_{t\to7}t^2-5\lim_{t\to7}t+\lim_{t\to7}2}$ [3]

$=\dfrac{3(7^2)+1}{7^2-5(7)+2}$ [10, 9, and 8]

$=\dfrac{148}{16}=\dfrac{37}{4}$

7. $\lim_{u\to-2}\sqrt{9-u^3+2u^2}=\sqrt{\lim_{u\to-2}(9-u^3+2u^2)}$ [Limit Law 7]

$=\sqrt{\lim_{u\to-2}9-\lim_{u\to-2}u^3+\lim_{u\to-2}2u^2}$ [2 and 1]

$=\sqrt{\lim_{u\to-2}9-\lim_{u\to-2}u^3+2\lim_{u\to-2}u^2}$ [3]

$=\sqrt{9-(-2)^3+2(-2)^2}$ [8 and 10]

$=\sqrt{25}=5$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $\lim\limits_{x\to 3} \sqrt[3]{x+5}\,(2x^2-3x) = \lim\limits_{x\to 3} \sqrt[3]{x+5} \cdot \lim\limits_{x\to 3} \left(2x^2-3x\right)$ [Limit Law 4]

$= \sqrt[3]{\lim\limits_{x\to 3}(x+5)} \cdot \lim\limits_{x\to 3}(2x^2-3x)$ [7]

$= \sqrt[3]{\lim\limits_{x\to 3} x + \lim\limits_{x\to 3} 5} \cdot \left(\lim\limits_{x\to 3} 2x^2 - \lim\limits_{x\to 3} 3x\right)$ [1 and 2]

$= \sqrt[3]{\lim\limits_{x\to 3} x + \lim\limits_{x\to 3} 5} \cdot \left(2\lim\limits_{x\to 3} x^2 - 3\lim\limits_{x\to 3} x\right)$ [3]

$= \sqrt[3]{3+5} \cdot \left[2(3^2) - 3(3)\right]$ [9, 8, and 10]

$= 2 \cdot (18-9) = 18$

9. $\lim\limits_{t\to -1} \left(\dfrac{2t^5-t^4}{5t^2+4}\right)^3 = \left(\lim\limits_{t\to -1} \dfrac{2t^5-t^4}{5t^2+4}\right)^3$ [Limit Law 6]

$= \left(\dfrac{\lim\limits_{t\to -1}\left(2t^5-t^4\right)}{\lim\limits_{t\to -1}\left(5t^2+4\right)}\right)^3$ [5]

$= \left(\dfrac{2\lim\limits_{t\to -1} t^5 - \lim\limits_{t\to -1} t^4}{5\lim\limits_{t\to -1} t^2 + \lim\limits_{t\to -1} 4}\right)^3$ [3, 2, and 1]

$= \left(\dfrac{2(-1)^5-(-1)^4}{5(-1)^2+4}\right)^3$ [10 and 8]

$= \left(-\dfrac{3}{9}\right)^3 = -\dfrac{1}{27}$

10. (a) The left-hand side of the equation is not defined for $x = 2$, but the right-hand side is.

(b) Since the equation holds for all $x \neq 2$, it follows that both sides of the equation approach the same limit as $x \to 2$, just as in Example 3. Remember that in finding $\lim\limits_{x\to a} f(x)$, we never consider $x = a$.

11. $\lim\limits_{x\to -2}(3x-7) = 3(-2)-7 = -13$

12. $\lim\limits_{x\to 6}\left(8-\frac{1}{2}x\right) = 8-\frac{1}{2}(6) = 5$

13. $\lim\limits_{t\to 4} \dfrac{t^2-2t-8}{t-4} = \lim\limits_{t\to 4} \dfrac{(t-4)(t+2)}{t-4} = \lim\limits_{t\to 4}(t+2) = 4+2 = 6$

14. $\lim\limits_{x\to -3} \dfrac{x^2+3x}{x^2-x-12} = \lim\limits_{x\to -3} \dfrac{x(x+3)}{(x-4)(x+3)} = \lim\limits_{x\to -3} \dfrac{x}{x-4} = \dfrac{-3}{-3-4} = \dfrac{3}{7}$

15. $\lim\limits_{x\to 2} \dfrac{x^2+5x+4}{x-2}$ does not exist since $x-2 \to 0$, but $x^2+5x+4 \to 18$ as $x \to 2$.

16. $\lim\limits_{x\to 4} \dfrac{x^2+3x}{x^2-x-12} = \lim\limits_{x\to 4} \dfrac{x(x+3)}{(x-4)(x+3)} = \lim\limits_{x\to 4} \dfrac{x}{x-4}$. The last limit does not exist since $\lim\limits_{x\to 4^-} \dfrac{x}{x-4} = -\infty$ and $\lim\limits_{x\to 4^+} \dfrac{x}{x-4} = \infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\lim\limits_{x\to-2}\dfrac{x^2-x-6}{3x^2+5x-2}=\lim\limits_{x\to-2}\dfrac{(x-3)(x+2)}{(3x-1)(x+2)}=\lim\limits_{x\to-2}\dfrac{x-3}{3x-1}=\dfrac{-2-3}{3(-2)-1}=\dfrac{-5}{-7}=\dfrac{5}{7}$

18. $\lim\limits_{x\to-5}\dfrac{2x^2+9x-5}{x^2-25}=\lim\limits_{x\to-5}\dfrac{(2x-1)(x+5)}{(x-5)(x+5)}=\lim\limits_{x\to-5}\dfrac{2x-1}{x-5}=\dfrac{2(-5)-1}{-5-5}=\dfrac{-11}{-10}=\dfrac{11}{10}$

19. Factoring t^3-27 as the difference of two cubes, we have

$$\lim_{t\to3}\frac{t^3-27}{t^2-9}=\lim_{t\to3}\frac{(t-3)(t^2+3t+9)}{(t-3)(t+3)}=\lim_{t\to3}\frac{t^2+3t+9}{t+3}=\frac{3^2+3(3)+9}{3+3}=\frac{27}{6}=\frac{9}{2}.$$

20. Factoring u^3+1 as the sum of two cubes, we have

$$\lim_{u\to-1}\frac{u+1}{u^3+1}=\lim_{u\to-1}\frac{u+1}{(u+1)(u^2-u+1)}=\lim_{u\to-1}\frac{1}{u^2-u+1}=\frac{1}{(-1)^2-(-1)+1}=\frac{1}{3}.$$

21. $\lim\limits_{h\to0}\dfrac{(h-3)^2-9}{h}=\lim\limits_{h\to0}\dfrac{h^2-6h+9-9}{h}=\lim\limits_{h\to0}\dfrac{h^2-6h}{h}=\lim\limits_{h\to0}\dfrac{h(h-6)}{h}=\lim\limits_{h\to0}(h-6)=0-6=-6$

22. $\lim\limits_{x\to9}\dfrac{9-x}{3-\sqrt{x}}=\lim\limits_{x\to9}\dfrac{9-x}{3-\sqrt{x}}\cdot\dfrac{3+\sqrt{x}}{3+\sqrt{x}}=\lim\limits_{x\to9}\dfrac{(9-x)(3+\sqrt{x}\,)}{9-x}=\lim\limits_{x\to9}(3+\sqrt{x}\,)=3+\sqrt{9}=6$

23.
$$\begin{aligned}\lim_{h\to0}\frac{\sqrt{9+h}-3}{h}&=\lim_{h\to0}\frac{\sqrt{9+h}-3}{h}\cdot\frac{\sqrt{9+h}+3}{\sqrt{9+h}+3}=\lim_{h\to0}\frac{\left(\sqrt{9+h}\,\right)^2-3^2}{h\left(\sqrt{9+h}+3\right)}=\lim_{h\to0}\frac{(9+h)-9}{h\left(\sqrt{9+h}+3\right)}\\&=\lim_{h\to0}\frac{h}{h\left(\sqrt{9+h}+3\right)}=\lim_{h\to0}\frac{1}{\sqrt{9+h}+3}=\frac{1}{3+3}=\frac{1}{6}\end{aligned}$$

24.
$$\begin{aligned}\lim_{x\to2}\frac{2-x}{\sqrt{x+2}-2}&=\lim_{x\to2}\frac{2-x}{\sqrt{x+2}-2}\cdot\frac{\sqrt{x+2}+2}{\sqrt{x+2}+2}=\lim_{x\to2}\frac{(2-x)\left(\sqrt{x+2}+2\right)}{\left(\sqrt{x+2}\right)^2-4}=\lim_{x\to2}\frac{-(x-2)\left(\sqrt{x+2}+2\right)}{x-2}\\&=\lim_{x\to2}\left[-\left(\sqrt{x+2}+2\right)\right]=-\left(\sqrt{4}+2\right)=-4\end{aligned}$$

25. $\lim\limits_{x\to3}\dfrac{\dfrac{1}{x}-\dfrac{1}{3}}{x-3}=\lim\limits_{x\to3}\dfrac{\dfrac{1}{x}-\dfrac{1}{3}}{x-3}\cdot\dfrac{3x}{3x}=\lim\limits_{x\to3}\dfrac{3-x}{3x(x-3)}=\lim\limits_{x\to3}\dfrac{-1}{3x}=-\dfrac{1}{9}$

26.
$$\begin{aligned}\lim_{h\to0}\frac{(-2+h)^{-1}+2^{-1}}{h}&=\lim_{h\to0}\frac{\dfrac{1}{h-2}+\dfrac{1}{2}}{h}=\lim_{h\to0}\frac{\dfrac{2+(h-2)}{2(h-2)}}{h}=\lim_{h\to0}\frac{2+(h-2)}{2h(h-2)}\\&=\lim_{h\to0}\frac{h}{2h(h-2)}=\lim_{h\to0}\frac{1}{2(h-2)}=\frac{1}{2(0-2)}=-\frac{1}{4}\end{aligned}$$

27.
$$\begin{aligned}\lim_{t\to0}\frac{\sqrt{1+t}-\sqrt{1-t}}{t}&=\lim_{t\to0}\frac{\sqrt{1+t}-\sqrt{1-t}}{t}\cdot\frac{\sqrt{1+t}+\sqrt{1-t}}{\sqrt{1+t}+\sqrt{1-t}}=\lim_{t\to0}\frac{\left(\sqrt{1+t}\,\right)^2-\left(\sqrt{1-t}\,\right)^2}{t\left(\sqrt{1+t}+\sqrt{1-t}\,\right)}\\&=\lim_{t\to0}\frac{(1+t)-(1-t)}{t\left(\sqrt{1+t}+\sqrt{1-t}\,\right)}=\lim_{t\to0}\frac{2t}{t\left(\sqrt{1+t}+\sqrt{1-t}\,\right)}=\lim_{t\to0}\frac{2}{\sqrt{1+t}+\sqrt{1-t}}\\&=\frac{2}{\sqrt{1}+\sqrt{1}}=\frac{2}{2}=1\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $\lim\limits_{t\to 0}\left(\dfrac{1}{t}-\dfrac{1}{t^2+t}\right)=\lim\limits_{t\to 0}\left(\dfrac{1}{t}-\dfrac{1}{t(t+1)}\right)=\lim\limits_{t\to 0}\dfrac{t+1-1}{t(t+1)}=\lim\limits_{t\to 0}\dfrac{1}{t+1}=\dfrac{1}{0+1}=1$

29. $$\lim_{x\to 16}\frac{4-\sqrt{x}}{16x-x^2}=\lim_{x\to 16}\frac{(4-\sqrt{x})(4+\sqrt{x})}{(16x-x^2)(4+\sqrt{x})}=\lim_{x\to 16}\frac{16-x}{x(16-x)(4+\sqrt{x})}$$
$$=\lim_{x\to 16}\frac{1}{x(4+\sqrt{x})}=\frac{1}{16\left(4+\sqrt{16}\right)}=\frac{1}{16(8)}=\frac{1}{128}$$

30. $$\lim_{x\to 2}\frac{x^2-4x+4}{x^4-3x^2-4}=\lim_{x\to 2}\frac{(x-2)^2}{(x^2-4)(x^2+1)}=\lim_{x\to 2}\frac{(x-2)^2}{(x+2)(x-2)(x^2+1)}$$
$$=\lim_{x\to 2}\frac{x-2}{(x+2)(x^2+1)}=\frac{0}{4\cdot 5}=0$$

31. $$\lim_{t\to 0}\left(\frac{1}{t\sqrt{1+t}}-\frac{1}{t}\right)=\lim_{t\to 0}\frac{1-\sqrt{1+t}}{t\sqrt{1+t}}=\lim_{t\to 0}\frac{\left(1-\sqrt{1+t}\right)\left(1+\sqrt{1+t}\right)}{t\sqrt{t+1}\left(1+\sqrt{1+t}\right)}=\lim_{t\to 0}\frac{-t}{t\sqrt{1+t}\left(1+\sqrt{1+t}\right)}$$
$$=\lim_{t\to 0}\frac{-1}{\sqrt{1+t}\left(1+\sqrt{1+t}\right)}=\frac{-1}{\sqrt{1+0}\left(1+\sqrt{1+0}\right)}=-\frac{1}{2}$$

32. $$\lim_{x\to -4}\frac{\sqrt{x^2+9}-5}{x+4}=\lim_{x\to -4}\frac{\left(\sqrt{x^2+9}-5\right)\left(\sqrt{x^2+9}+5\right)}{(x+4)\left(\sqrt{x^2+9}+5\right)}=\lim_{x\to -4}\frac{(x^2+9)-25}{(x+4)\left(\sqrt{x^2+9}+5\right)}$$
$$=\lim_{x\to -4}\frac{x^2-16}{(x+4)\left(\sqrt{x^2+9}+5\right)}=\lim_{x\to -4}\frac{(x+4)(x-4)}{(x+4)\left(\sqrt{x^2+9}+5\right)}$$
$$=\lim_{x\to -4}\frac{x-4}{\sqrt{x^2+9}+5}=\frac{-4-4}{\sqrt{16+9}+5}=\frac{-8}{5+5}=-\frac{4}{5}$$

33. $$\lim_{h\to 0}\frac{(x+h)^3-x^3}{h}=\lim_{h\to 0}\frac{(x^3+3x^2h+3xh^2+h^3)-x^3}{h}=\lim_{h\to 0}\frac{3x^2h+3xh^2+h^3}{h}$$
$$=\lim_{h\to 0}\frac{h(3x^2+3xh+h^2)}{h}=\lim_{h\to 0}(3x^2+3xh+h^2)=3x^2$$

34. $$\lim_{h\to 0}\frac{\dfrac{1}{(x+h)^2}-\dfrac{1}{x^2}}{h}=\lim_{h\to 0}\frac{\dfrac{x^2-(x+h)^2}{(x+h)^2x^2}}{h}=\lim_{h\to 0}\frac{x^2-(x^2+2xh+h^2)}{hx^2(x+h)^2}=\lim_{h\to 0}\frac{-h(2x+h)}{hx^2(x+h)^2}$$
$$=\lim_{h\to 0}\frac{-(2x+h)}{x^2(x+h)^2}=\frac{-2x}{x^2\cdot x^2}=-\frac{2}{x^3}$$

35. (a)

$$\lim_{x\to 0}\frac{x}{\sqrt{1+3x}-1}\approx\frac{2}{3}$$

(b)

x	$f(x)$
−0.001	0.666 166 3
−0.000 1	0.666 616 7
−0.000 01	0.666 661 7
−0.000 001	0.666 666 2
0.000 001	0.666 667 2
0.000 01	0.666 671 7
0.000 1	0.666 716 7
0.001	0.667 166 3

The limit appears to be $\dfrac{2}{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\lim\limits_{x\to 0}\left(\dfrac{x}{\sqrt{1+3x}-1}\cdot\dfrac{\sqrt{1+3x}+1}{\sqrt{1+3x}+1}\right)=\lim\limits_{x\to 0}\dfrac{x\left(\sqrt{1+3x}+1\right)}{(1+3x)-1}=\lim\limits_{x\to 0}\dfrac{x\left(\sqrt{1+3x}+1\right)}{3x}$

$$=\frac{1}{3}\lim_{x\to 0}\left(\sqrt{1+3x}+1\right) \qquad \text{[Limit Law 3]}$$

$$=\frac{1}{3}\left[\sqrt{\lim_{x\to 0}(1+3x)}+\lim_{x\to 0}1\right] \qquad \text{[1 and 7]}$$

$$=\frac{1}{3}\left(\sqrt{\lim_{x\to 0}1+3\lim_{x\to 0}x}+1\right) \qquad \text{[1, 3, and 8]}$$

$$=\frac{1}{3}\left(\sqrt{1+3\cdot 0}+1\right) \qquad \text{[8 and 9]}$$

$$=\frac{1}{3}(1+1)=\frac{2}{3}$$

36. (a)

$$\lim_{x\to 0}\frac{\sqrt{3+x}-\sqrt{3}}{x}\approx 0.29$$

(b)

x	$f(x)$
-0.001	0.288 699 2
$-0.000\,1$	0.288 677 5
$-0.000\,01$	0.288 675 4
$-0.000\,001$	0.288 675 2
$0.000\,001$	0.288 675 1
$0.000\,01$	0.288 674 9
$0.000\,1$	0.288 672 7
0.001	0.288 651 1

The limit appears to be approximately 0.2887.

(c) $\lim\limits_{x\to 0}\left(\dfrac{\sqrt{3+x}-\sqrt{3}}{x}\cdot\dfrac{\sqrt{3+x}+\sqrt{3}}{\sqrt{3+x}+\sqrt{3}}\right)=\lim\limits_{x\to 0}\dfrac{(3+x)-3}{x\left(\sqrt{3+x}+\sqrt{3}\right)}=\lim\limits_{x\to 0}\dfrac{1}{\sqrt{3+x}+\sqrt{3}}$

$$=\frac{\lim\limits_{x\to 0}1}{\lim\limits_{x\to 0}\sqrt{3+x}+\lim\limits_{x\to 0}\sqrt{3}} \qquad \text{[Limit Laws 5 and 1]}$$

$$=\frac{1}{\sqrt{\lim\limits_{x\to 0}(3+x)}+\sqrt{3}} \qquad \text{[7 and 8]}$$

$$=\frac{1}{\sqrt{3+0}+\sqrt{3}} \qquad \text{[1, 8, and 9]}$$

$$=\frac{1}{2\sqrt{3}}$$

37. Let $f(x)=-x^2$, $g(x)=x^2\cos 20\pi x$ and $h(x)=x^2$. Then

$-1\le\cos 20\pi x\le 1 \quad\Rightarrow\quad -x^2\le x^2\cos 20\pi x\le x^2 \quad\Rightarrow\quad f(x)\le g(x)\le h(x)$.

So since $\lim\limits_{x\to 0}f(x)=\lim\limits_{x\to 0}h(x)=0$, by the Squeeze Theorem we have

$\lim\limits_{x\to 0}g(x)=0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. Let $f(x) = -\sqrt{x^3 + x^2}$, $g(x) = \sqrt{x^3 + x^2}\,\sin(\pi/x)$, and $h(x) = \sqrt{x^3 + x^2}$. Then $-1 \le \sin(\pi/x) \le 1 \quad \Rightarrow \quad -\sqrt{x^3 + x^2} \le \sqrt{x^3 + x^2}\sin(\pi/x) \le \sqrt{x^3 + x^2} \quad \Rightarrow$ $f(x) \le g(x) \le h(x)$. So since $\lim\limits_{x\to 0} f(x) = \lim\limits_{x\to 0} h(x) = 0$, by the Squeeze Theorem we have $\lim\limits_{x\to 0} g(x) = 0$.

39. We have $\lim\limits_{x\to 4}(4x - 9) = 4(4) - 9 = 7$ and $\lim\limits_{x\to 4}\left(x^2 - 4x + 7\right) = 4^2 - 4(4) + 7 = 7$. Since $4x - 9 \le f(x) \le x^2 - 4x + 7$ for $x \ge 0$, $\lim\limits_{x\to 4} f(x) = 7$ by the Squeeze Theorem.

40. We have $\lim\limits_{x\to 1}(2x) = 2(1) = 2$ and $\lim\limits_{x\to 1}(x^4 - x^2 + 2) = 1^4 - 1^2 + 2 = 2$. Since $2x \le g(x) \le x^4 - x^2 + 2$ for all x, $\lim\limits_{x\to 1} g(x) = 2$ by the Squeeze Theorem.

41. $-1 \le \cos(2/x) \le 1 \quad \Rightarrow \quad -x^4 \le x^4\cos(2/x) \le x^4$. Since $\lim\limits_{x\to 0}\left(-x^4\right) = 0$ and $\lim\limits_{x\to 0} x^4 = 0$, we have $\lim\limits_{x\to 0}\left[x^4\cos(2/x)\right] = 0$ by the Squeeze Theorem.

42. $-1 \le \sin(\pi/x) \le 1 \quad \Rightarrow \quad e^{-1} \le e^{\sin(\pi/x)} \le e^1 \quad \Rightarrow \quad \sqrt{x}/e \le \sqrt{x}\,e^{\sin(\pi/x)} \le \sqrt{x}\,e$. Since $\lim\limits_{x\to 0^+}(\sqrt{x}/e) = 0$ and $\lim\limits_{x\to 0^+}(\sqrt{x}\,e) = 0$, we have $\lim\limits_{x\to 0^+}\left[\sqrt{x}\,e^{\sin(\pi/x)}\right] = 0$ by the Squeeze Theorem.

43. $|x + 4| = \begin{cases} x + 4 & \text{if } x + 4 \ge 0 \\ -(x + 4) & \text{if } x + 4 < 0 \end{cases} = \begin{cases} x + 4 & \text{if } x \ge -4 \\ -(x + 4) & \text{if } x < -4 \end{cases}$

Thus, $\lim\limits_{x\to -4^+}(|x + 4| - 2x) = \lim\limits_{x\to -4^+}(x + 4 - 2x) = \lim\limits_{x\to -4^+}(-x + 4) = 4 + 4 = 8$ and

$\lim\limits_{x\to -4^-}(|x + 4| - 2x) = \lim\limits_{x\to -4^-}(-(x + 4) - 2x) = \lim\limits_{x\to -4^-}(-3x - 4) = 12 - 4 = 8.$

The left and right limits are equal, so $\lim\limits_{x\to -4}(|x + 4| - 2x) = 8$.

44. $|x + 4| = \begin{cases} x + 4 & \text{if } x + 4 \ge 0 \\ -(x + 4) & \text{if } x + 4 < 0 \end{cases} = \begin{cases} x + 4 & \text{if } x \ge -4 \\ -(x + 4) & \text{if } x < -4 \end{cases}$

Thus, $\lim\limits_{x\to -4^+}\dfrac{|x + 4|}{2x + 8} = \lim\limits_{x\to -4^+}\dfrac{x + 4}{2x + 8} = \lim\limits_{x\to -4^+}\dfrac{x + 4}{2(x + 4)} = \lim\limits_{x\to -4^+}\dfrac{1}{2} = \dfrac{1}{2}$ and

$\lim\limits_{x\to -4^-}\dfrac{|x + 4|}{2x + 8} = \lim\limits_{x\to -4^-}\dfrac{-(x + 4)}{2x + 8} = \lim\limits_{x\to -4^-}\dfrac{-(x + 4)}{2(x + 4)} = \lim\limits_{x\to -4^-}\dfrac{-1}{2} = -\dfrac{1}{2}.$

The left and right limits are different, so $\lim\limits_{x\to -4}\dfrac{|x + 4|}{2x + 8}$ does not exist.

45. $\left|2x^3 - x^2\right| = \left|x^2(2x - 1)\right| = \left|x^2\right| \cdot |2x - 1| = x^2\,|2x - 1|$

$|2x - 1| = \begin{cases} 2x - 1 & \text{if } 2x - 1 \ge 0 \\ -(2x - 1) & \text{if } 2x - 1 < 0 \end{cases} = \begin{cases} 2x - 1 & \text{if } x \ge 0.5 \\ -(2x - 1) & \text{if } x < 0.5 \end{cases}$

So $\left|2x^3 - x^2\right| = x^2[-(2x - 1)]$ for $x < 0.5$.

Thus, $\lim\limits_{x\to 0.5^-}\dfrac{2x - 1}{|2x^3 - x^2|} = \lim\limits_{x\to 0.5^-}\dfrac{2x - 1}{x^2[-(2x - 1)]} = \lim\limits_{x\to 0.5^-}\dfrac{-1}{x^2} = \dfrac{-1}{(0.5)^2} = \dfrac{-1}{0.25} = -4.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. Since $|x| = -x$ for $x < 0$, we have $\lim\limits_{x\to -2} \dfrac{2-|x|}{2+x} = \lim\limits_{x\to -2} \dfrac{2-(-x)}{2+x} = \lim\limits_{x\to -2} \dfrac{2+x}{2+x} = \lim\limits_{x\to -2} 1 = 1.$

47. Since $|x| = -x$ for $x < 0$, we have $\lim\limits_{x\to 0^-} \left(\dfrac{1}{x} - \dfrac{1}{|x|}\right) = \lim\limits_{x\to 0^-} \left(\dfrac{1}{x} - \dfrac{1}{-x}\right) = \lim\limits_{x\to 0^-} \dfrac{2}{x}$, which does not exist since the denominator approaches 0 and the numerator does not.

48. Since $|x| = x$ for $x > 0$, we have $\lim\limits_{x\to 0^+} \left(\dfrac{1}{x} - \dfrac{1}{|x|}\right) = \lim\limits_{x\to 0^+} \left(\dfrac{1}{x} - \dfrac{1}{x}\right) = \lim\limits_{x\to 0^+} 0 = 0.$

49. (a)

(b) (i) Since $\operatorname{sgn} x = 1$ for $x > 0$, $\lim\limits_{x\to 0^+} \operatorname{sgn} x = \lim\limits_{x\to 0^+} 1 = 1$.

(ii) Since $\operatorname{sgn} x = -1$ for $x < 0$, $\lim\limits_{x\to 0^-} \operatorname{sgn} x = \lim\limits_{x\to 0^-} -1 = -1$.

(iii) Since $\lim\limits_{x\to 0^-} \operatorname{sgn} x \neq \lim\limits_{x\to 0^+} \operatorname{sgn} x$, $\lim\limits_{x\to 0} \operatorname{sgn} x$ does not exist.

(iv) Since $|\operatorname{sgn} x| = 1$ for $x \neq 0$, $\lim\limits_{x\to 0} |\operatorname{sgn} x| = \lim\limits_{x\to 0} 1 = 1$.

50. (a) $g(x) = \operatorname{sgn}(\sin x) = \begin{cases} -1 & \text{if } \sin x < 0 \\ 0 & \text{if } \sin x = 0 \\ 1 & \text{if } \sin x > 0 \end{cases}$

(i) $\lim\limits_{x\to 0^+} g(x) = \lim\limits_{x\to 0^+} \operatorname{sgn}(\sin x) = 1$ since $\sin x$ is positive for small positive values of x.

(ii) $\lim\limits_{x\to 0^-} g(x) = \lim\limits_{x\to 0^-} \operatorname{sgn}(\sin x) = -1$ since $\sin x$ is negative for small negative values of x.

(iii) $\lim\limits_{x\to 0} g(x)$ does not exist since $\lim\limits_{x\to 0^+} g(x) \neq \lim\limits_{x\to 0^-} g(x)$.

(iv) $\lim\limits_{x\to \pi^+} g(x) = \lim\limits_{x\to \pi^+} \operatorname{sgn}(\sin x) = -1$ since $\sin x$ is negative for values of x slightly greater than π.

(v) $\lim\limits_{x\to \pi^-} g(x) = \lim\limits_{x\to \pi^-} \operatorname{sgn}(\sin x) = 1$ since $\sin x$ is positive for values of x slightly less than π.

(vi) $\lim\limits_{x\to \pi} g(x)$ does not exist since $\lim\limits_{x\to \pi^+} g(x) \neq \lim\limits_{x\to \pi^-} g(x)$.

(b) The sine function changes sign at every integer multiple of π, so the signum function equals 1 on one side and -1 on the other side of $n\pi$, n an integer. Thus, $\lim\limits_{x\to a} g(x)$ does not exist for $a = n\pi$, n an integer.

(c)

51. (a) (i)
$$\begin{aligned} \lim_{x\to 2^+} g(x) &= \lim_{x\to 2^+} \frac{x^2+x-6}{|x-2|} = \lim_{x\to 2^+} \frac{(x+3)(x-2)}{|x-2|} \\ &= \lim_{x\to 2^+} \frac{(x+3)(x-2)}{x-2} \quad [\text{since } x-2>0 \text{ if } x \to 2^+] \\ &= \lim_{x\to 2^+} (x+3) = 5 \end{aligned}$$

(ii) The solution is similar to the solution in part (i), but now $|x-2| = 2-x$ since $x-2<0$ if $x \to 2^-$.

Thus, $\lim\limits_{x\to 2^-} g(x) = \lim\limits_{x\to 2^-} -(x+3) = -5$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Since the right-hand and left-hand limits of g at $x = 2$ are not equal, $\lim_{x \to 2} g(x)$ does not exist.

(c)

52. (a) $f(x) = \begin{cases} x^2 + 1 & \text{if } x < 1 \\ (x-2)^2 & \text{if } x \geq 1 \end{cases}$

$$\lim_{x \to 1^-} f(x) = \lim_{x \to 1^-} (x^2 + 1) = 1^2 + 1 = 2, \quad \lim_{x \to 1^+} f(x) = \lim_{x \to 1+} (x-2)^2 = (-1)^2 = 1$$

(b) Since the right-hand and left-hand limits of f at $x = 1$ are not equal, $\lim_{x \to 1} f(x)$ does not exist.

(c)

53. For the $\lim_{t \to 2} B(t)$ to exist, the one-sided limits at $t = 2$ must be equal. $\lim_{t \to 2^-} B(t) = \lim_{t \to 2^-} \left(4 - \frac{1}{2}t\right) = 4 - 1 = 3$ and $\lim_{t \to 2^+} B(t) = \lim_{t \to 2^+} \sqrt{t + c} = \sqrt{2 + c}$. Now $3 = \sqrt{2 + c} \;\Rightarrow\; 9 = 2 + c \;\Leftrightarrow\; c = 7$.

54. (a) (i) $\lim_{x \to 1^-} g(x) = \lim_{x \to 1^-} x = 1$

(ii) $\lim_{x \to 1^+} g(x) = \lim_{x \to 1^+} (2 - x^2) = 2 - 1^2 = 1$. Since $\lim_{x \to 1^-} g(x) = 1$ and $\lim_{x \to 1^+} g(x) = 1$, we have $\lim_{x \to 1} g(x) = 1$.

Note that the fact $g(1) = 3$ does not affect the value of the limit.

(iii) When $x = 1$, $g(x) = 3$, so $g(1) = 3$.

(iv) $\lim_{x \to 2^-} g(x) = \lim_{x \to 2^-} (2 - x^2) = 2 - 2^2 = 2 - 4 = -2$

(v) $\lim_{x \to 2^+} g(x) = \lim_{x \to 2^+} (x - 3) = 2 - 3 = -1$

(vi) $\lim_{x \to 2} g(x)$ does not exist since $\lim_{x \to 2^-} g(x) \neq \lim_{x \to 2^+} g(x)$.

(b)

$$g(x) = \begin{cases} x & \text{if } x < 1 \\ 3 & \text{if } x = 1 \\ 2 - x^2 & \text{if } 1 < x \leq 2 \\ x - 3 & \text{if } x > 2 \end{cases}$$

55. (a) (i) $[\![x]\!] = -2$ for $-2 \leq x < -1$, so $\lim_{x \to -2^+} [\![x]\!] = \lim_{x \to -2^+} (-2) = -2$

(ii) $[\![x]\!] = -3$ for $-3 \leq x < -2$, so $\lim_{x \to -2^-} [\![x]\!] = \lim_{x \to -2^-} (-3) = -3$.

The right and left limits are different, so $\lim_{x \to -2} [\![x]\!]$ does not exist.

(iii) $[\![x]\!] = -3$ for $-3 \leq x < -2$, so $\lim_{x \to -2.4} [\![x]\!] = \lim_{x \to -2.4} (-3) = -3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) (i) $[\![x]\!] = n - 1$ for $n - 1 \le x < n$, so $\lim\limits_{x \to n^-} [\![x]\!] = \lim\limits_{x \to n^-} (n - 1) = n - 1$.

(ii) $[\![x]\!] = n$ for $n \le x < n + 1$, so $\lim\limits_{x \to n^+} [\![x]\!] = \lim\limits_{x \to n^+} n = n$.

(c) $\lim\limits_{x \to a} [\![x]\!]$ exists $\Leftrightarrow$ a is not an integer.

56. (a) See the graph of $y = \cos x$.

Since $-1 \le \cos x < 0$ on $[-\pi, -\pi/2)$, we have $y = f(x) = [\![\cos x]\!] = -1$ on $[-\pi, -\pi/2)$.

Since $0 \le \cos x < 1$ on $[-\pi/2, 0) \cup (0, \pi/2]$, we have $f(x) = 0$ on $[-\pi/2, 0) \cup (0, \pi/2]$.

Since $-1 \le \cos x < 0$ on $(\pi/2, \pi]$, we have $f(x) = -1$ on $(\pi/2, \pi]$.

Note that $f(0) = 1$.

(b) (i) $\lim\limits_{x \to 0^-} f(x) = 0$ and $\lim\limits_{x \to 0^+} f(x) = 0$, so $\lim\limits_{x \to 0} f(x) = 0$.

(ii) As $x \to (\pi/2)^-$, $f(x) \to 0$, so $\lim\limits_{x \to (\pi/2)^-} f(x) = 0$.

(iii) As $x \to (\pi/2)^+$, $f(x) \to -1$, so $\lim\limits_{x \to (\pi/2)^+} f(x) = -1$.

(iv) Since the answers in parts (ii) and (iii) are not equal, $\lim\limits_{x \to \pi/2} f(x)$ does not exist.

(c) $\lim\limits_{x \to a} f(x)$ exists for all a in the open interval $(-\pi, \pi)$ except $a = -\pi/2$ and $a = \pi/2$.

57. The graph of $f(x) = [\![x]\!] + [\![-x]\!]$ is the same as the graph of $g(x) = -1$ with holes at each integer, since $f(a) = 0$ for any integer a. Thus, $\lim\limits_{x \to 2^-} f(x) = -1$ and $\lim\limits_{x \to 2^+} f(x) = -1$, so $\lim\limits_{x \to 2} f(x) = -1$. However, $f(2) = [\![2]\!] + [\![-2]\!] = 2 + (-2) = 0$, so $\lim\limits_{x \to 2} f(x) \ne f(2)$.

58. $\lim\limits_{v \to c^-} \left(L_0 \sqrt{1 - \dfrac{v^2}{c^2}} \right) = L_0 \sqrt{1 - 1} = 0$. As the velocity approaches the speed of light, the length approaches 0.

A left-hand limit is necessary since L is not defined for $v > c$.

59. Since $p(x)$ is a polynomial, $p(x) = a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n$. Thus, by the Limit Laws,

$$\begin{aligned} \lim_{x \to a} p(x) &= \lim_{x \to a} \left(a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n \right) = a_0 + a_1 \lim_{x \to a} x + a_2 \lim_{x \to a} x^2 + \cdots + a_n \lim_{x \to a} x^n \\ &= a_0 + a_1 a + a_2 a^2 + \cdots + a_n a^n = p(a) \end{aligned}$$

Thus, for any polynomial p, $\lim\limits_{x \to a} p(x) = p(a)$.

60. Let $r(x) = \dfrac{p(x)}{q(x)}$ where $p(x)$ and $q(x)$ are any polynomials, and suppose that $q(a) \ne 0$. Then

$$\lim_{x \to a} r(x) = \lim_{x \to a} \frac{p(x)}{q(x)} = \frac{\lim\limits_{x \to a} p(x)}{\lim\limits_{x \to a} q(x)} \quad \text{[Limit Law 5]} \quad = \frac{p(a)}{q(a)} \quad \text{[Exercise 59]} \quad = r(a).$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. $\lim\limits_{x\to1}[f(x)-8]=\lim\limits_{x\to1}\left[\dfrac{f(x)-8}{x-1}\cdot(x-1)\right]=\lim\limits_{x\to1}\dfrac{f(x)-8}{x-1}\cdot\lim\limits_{x\to1}(x-1)=10\cdot0=0.$

Thus, $\lim\limits_{x\to1}f(x)=\lim\limits_{x\to1}\{[f(x)-8]+8\}=\lim\limits_{x\to1}[f(x)-8]+\lim\limits_{x\to1}8=0+8=8.$

Note: The value of $\lim\limits_{x\to1}\dfrac{f(x)-8}{x-1}$ does not affect the answer since it's multiplied by 0. What's important is that $\lim\limits_{x\to1}\dfrac{f(x)-8}{x-1}$ exists.

62. (a) $\lim\limits_{x\to0}f(x)=\lim\limits_{x\to0}\left[\dfrac{f(x)}{x^2}\cdot x^2\right]=\lim\limits_{x\to0}\dfrac{f(x)}{x^2}\cdot\lim\limits_{x\to0}x^2=5\cdot0=0$

(b) $\lim\limits_{x\to0}\dfrac{f(x)}{x}=\lim\limits_{x\to0}\left[\dfrac{f(x)}{x^2}\cdot x\right]=\lim\limits_{x\to0}\dfrac{f(x)}{x^2}\cdot\lim\limits_{x\to0}x=5\cdot0=0$

63. Observe that $0\le f(x)\le x^2$ for all x, and $\lim\limits_{x\to0}0=0=\lim\limits_{x\to0}x^2$. So, by the Squeeze Theorem, $\lim\limits_{x\to0}f(x)=0$.

64. Let $f(x)=[\![x]\!]$ and $g(x)=-[\![x]\!]$. Then $\lim\limits_{x\to3}f(x)$ and $\lim\limits_{x\to3}g(x)$ do not exist [Example 10]

but $\lim\limits_{x\to3}[f(x)+g(x)]=\lim\limits_{x\to3}([\![x]\!]-[\![x]\!])=\lim\limits_{x\to3}0=0.$

65. Let $f(x)=H(x)$ and $g(x)=1-H(x)$, where H is the Heaviside function defined in Exercise 1.3.63.

Thus, either f or g is 0 for any value of x. Then $\lim\limits_{x\to0}f(x)$ and $\lim\limits_{x\to0}g(x)$ do not exist, but $\lim\limits_{x\to0}[f(x)g(x)]=\lim\limits_{x\to0}0=0.$

66.
$$\lim_{x\to2}\frac{\sqrt{6-x}-2}{\sqrt{3-x}-1}=\lim_{x\to2}\left(\frac{\sqrt{6-x}-2}{\sqrt{3-x}-1}\cdot\frac{\sqrt{6-x}+2}{\sqrt{6-x}+2}\cdot\frac{\sqrt{3-x}+1}{\sqrt{3-x}+1}\right)$$
$$=\lim_{x\to2}\left[\frac{\left(\sqrt{6-x}\right)^2-2^2}{\left(\sqrt{3-x}\right)^2-1^2}\cdot\frac{\sqrt{3-x}+1}{\sqrt{6-x}+2}\right]=\lim_{x\to2}\left(\frac{6-x-4}{3-x-1}\cdot\frac{\sqrt{3-x}+1}{\sqrt{6-x}+2}\right)$$
$$=\lim_{x\to2}\frac{(2-x)\left(\sqrt{3-x}+1\right)}{(2-x)\left(\sqrt{6-x}+2\right)}=\lim_{x\to2}\frac{\sqrt{3-x}+1}{\sqrt{6-x}+2}=\frac{1}{2}$$

67. Since the denominator approaches 0 as $x\to-2$, the limit will exist only if the numerator also approaches 0 as $x\to-2$. In order for this to happen, we need $\lim\limits_{x\to-2}\left(3x^2+ax+a+3\right)=0\quad\Leftrightarrow$

$3(-2)^2+a(-2)+a+3=0\quad\Leftrightarrow\quad12-2a+a+3=0\quad\Leftrightarrow\quad a=15$. With $a=15$, the limit becomes

$$\lim_{x\to-2}\frac{3x^2+15x+18}{x^2+x-2}=\lim_{x\to-2}\frac{3(x+2)(x+3)}{(x-1)(x+2)}=\lim_{x\to-2}\frac{3(x+3)}{x-1}=\frac{3(-2+3)}{-2-1}=\frac{3}{-3}=-1.$$

68. *Solution 1:* First, we find the coordinates of P and Q as functions of r. Then we can find the equation of the line determined by these two points, and thus find the x-intercept (the point R), and take the limit as $r\to0$. The coordinates of P are $(0,r)$. The point Q is the point of intersection of the two circles $x^2+y^2=r^2$ and $(x-1)^2+y^2=1$. Eliminating y from these equations, we get $r^2-x^2=1-(x-1)^2\quad\Leftrightarrow\quad r^2=1+2x-1\quad\Leftrightarrow\quad x=\frac{1}{2}r^2$. Substituting back into the equation of the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

shrinking circle to find the y-coordinate, we get $\left(\frac{1}{2}r^2\right)^2 + y^2 = r^2 \quad \Leftrightarrow \quad y^2 = r^2\left(1 - \frac{1}{4}r^2\right) \quad \Leftrightarrow \quad y = r\sqrt{1 - \frac{1}{4}r^2}$ (the positive y-value). So the coordinates of Q are $\left(\frac{1}{2}r^2, r\sqrt{1 - \frac{1}{4}r^2}\right)$. The equation of the line joining P and Q is thus $y - r = \dfrac{r\sqrt{1 - \frac{1}{4}r^2} - r}{\frac{1}{2}r^2 - 0}(x - 0)$. We set $y = 0$ in order to find the x-intercept, and get

$$x = -r\frac{\frac{1}{2}r^2}{r\left(\sqrt{1 - \frac{1}{4}r^2} - 1\right)} = \frac{-\frac{1}{2}r^2\left(\sqrt{1 - \frac{1}{4}r^2} + 1\right)}{1 - \frac{1}{4}r^2 - 1} = 2\left(\sqrt{1 - \frac{1}{4}r^2} + 1\right)$$

Now we take the limit as $r \to 0^+$: $\lim\limits_{r\to 0^+} x = \lim\limits_{r\to 0^+} 2\left(\sqrt{1 - \frac{1}{4}r^2} + 1\right) = \lim\limits_{r\to 0^+} 2\left(\sqrt{1} + 1\right) = 4$.

So the limiting position of R is the point $(4, 0)$.

Solution 2: We add a few lines to the diagram, as shown. Note that $\angle PQS = 90°$ (subtended by diameter PS). So $\angle SQR = 90° = \angle OQT$ (subtended by diameter OT). It follows that $\angle OQS = \angle TQR$. Also $\angle PSQ = 90° - \angle SPQ = \angle ORP$. Since $\triangle QOS$ is isosceles, so is $\triangle QTR$, implying that $QT = TR$. As the circle C_2 shrinks, the point Q plainly approaches the origin, so the point R must approach a point twice as far from the origin as T, that is, the point $(4, 0)$, as above.

2.4 The Precise Definition of a Limit

1. If $|f(x) - 1| < 0.2$, then $-0.2 < f(x) - 1 < 0.2 \quad \Rightarrow \quad 0.8 < f(x) < 1.2$. From the graph, we see that the last inequality is true if $0.7 < x < 1.1$, so we can choose $\delta = \min\{1 - 0.7, 1.1 - 1\} = \min\{0.3, 0.1\} = 0.1$ (or any smaller positive number).

2. If $|f(x) - 2| < 0.5$, then $-0.5 < f(x) - 2 < 0.5 \quad \Rightarrow \quad 1.5 < f(x) < 2.5$. From the graph, we see that the last inequality is true if $2.6 < x < 3.8$, so we can take $\delta = \min\{3 - 2.6, 3.8 - 3\} = \min\{0.4, 0.8\} = 0.4$ (or any smaller positive number). Note that $x \neq 3$.

3. The leftmost question mark is the solution of $\sqrt{x} = 1.6$ and the rightmost, $\sqrt{x} = 2.4$. So the values are $1.6^2 = 2.56$ and $2.4^2 = 5.76$. On the left side, we need $|x - 4| < |2.56 - 4| = 1.44$. On the right side, we need $|x - 4| < |5.76 - 4| = 1.76$. To satisfy both conditions, we need the more restrictive condition to hold—namely, $|x - 4| < 1.44$. Thus, we can choose $\delta = 1.44$, or any smaller positive number.

4. The leftmost question mark is the positive solution of $x^2 = \frac{1}{2}$, that is, $x = \frac{1}{\sqrt{2}}$, and the rightmost question mark is the positive solution of $x^2 = \frac{3}{2}$, that is, $x = \sqrt{\frac{3}{2}}$. On the left side, we need $|x - 1| < \left|\frac{1}{\sqrt{2}} - 1\right| \approx 0.292$ (rounding down to be safe). On the right side, we need $|x - 1| < \left|\sqrt{\frac{3}{2}} - 1\right| \approx 0.224$. The more restrictive of these two conditions must apply, so we choose $\delta = 0.224$ (or any smaller positive number).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5.

From the graph, we find that $y = \sqrt{x^2+5} = 2.7\ [3-0.3]$ when $x \approx 1.513$, so $2-\delta_1 \approx 1.513 \quad \Rightarrow \quad \delta_1 \approx 2-1.513 = 0.487$. Also, $y = \sqrt{x^2+5} = 3.3\ [3+0.3]$ when $x \approx 2.426$, so $2+\delta_2 \approx 2.426 \quad \Rightarrow$ $\delta_2 \approx 2.426 - 2 = 0.426$. Thus, we choose $\delta = 0.426$ (or any smaller positive number) since this is the smaller of δ_1 and δ_2.

6.

From the graph, we find that $y = \cos^2 x = 0.85\ [0.75+0.10]$ when $x \approx 0.398$, so $\frac{\pi}{6} - \delta_1 \approx 0.398 \quad \Rightarrow \quad \delta_1 \approx \frac{\pi}{6} - 0.398 \approx 0.126$. Also, $y = \cos^2 x = 0.65\ [0.75-0.10]$ when $x \approx 0.633$, so $\frac{\pi}{6} + \delta_2 \approx 0.633 \quad \Rightarrow$ $\delta_2 \approx 0.633 - \frac{\pi}{6} \approx 0.109$. Thus, we choose $\delta = 0.109$ (or any smaller positive number) since this is the smaller of δ_1 and δ_2.

7.

From the graph with $\varepsilon = 0.2$, we find that $y = x^3 - 3x + 4 = 5.8\ [6-\varepsilon]$ when $x \approx 1.9774$, so $2-\delta_1 \approx 1.9774 \quad \Rightarrow \quad \delta_1 \approx 0.0226$. Also, $y = x^3 - 3x + 4 = 6.2\ [6+\varepsilon]$ when $x \approx 2.022$, so $2+\delta_2 \approx 2.0219 \quad \Rightarrow$ $\delta_2 \approx 0.0219$. Thus, we choose $\delta = 0.0219$ (or any smaller positive number) since this is the smaller of δ_1 and δ_2.

For $\varepsilon = 0.1$, we get $\delta_1 \approx 0.0112$ and $\delta_2 \approx 0.0110$, so we choose $\delta = 0.011$ (or any smaller positive number).

8.

From the graph with $\varepsilon = 0.5$, we find that $y = (e^{2x}-1)/x = 1.5\ [2-\varepsilon]$ when $x \approx -0.303$, so $\delta_1 \approx 0.303$. Also, $y = (e^{2x}-1)/x = 2.5\ [2+\varepsilon]$ when $x \approx 0.215$, so $\delta_2 \approx 0.215$. Thus, we choose $\delta = 0.215$ (or any smaller positive number) since this is the smaller of δ_1 and δ_2.

For $\varepsilon = 0.1$, we get $\delta_1 \approx 0.052$ and $\delta_2 \approx 0.048$, so we choose $\delta = 0.048$ (or any smaller positive number).

9. (a)

The first graph of $y = \dfrac{1}{\ln(x-1)}$ shows a vertical asymptote at $x = 2$. The second graph shows that $y = 100$ when $x \approx 2.01$ (more accurately, 2.01005). Thus, we choose $\delta = 0.01$ (or any smaller positive number).

(b) From part (a), we see that as x gets closer to 2 from the right, y increases without bound. In symbols,

$$\lim_{x \to 2^+} \frac{1}{\ln(x-1)} = \infty.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. We graph $y = \csc^2 x$ and $y = 500$. The graphs intersect at $x \approx 3.186$, so we choose $\delta = 3.186 - \pi \approx 0.044$. Thus, if $0 < |x - \pi| < 0.044$, then $\csc^2 x > 500$. Similarly, for $M = 1000$, we get $\delta = 3.173 - \pi \approx 0.031$.

11. (a) $A = \pi r^2$ and $A = 1000 \text{ cm}^2 \quad \Rightarrow \quad \pi r^2 = 1000 \quad \Rightarrow \quad r^2 = \frac{1000}{\pi} \quad \Rightarrow \quad r = \sqrt{\frac{1000}{\pi}} \quad (r > 0) \quad \approx 17.8412$ cm.

(b) $|A - 1000| \le 5 \quad \Rightarrow \quad -5 \le \pi r^2 - 1000 \le 5 \quad \Rightarrow \quad 1000 - 5 \le \pi r^2 \le 1000 + 5 \quad \Rightarrow$

$\sqrt{\frac{995}{\pi}} \le r \le \sqrt{\frac{1005}{\pi}} \quad \Rightarrow \quad 17.7966 \le r \le 17.8858$. $\sqrt{\frac{1000}{\pi}} - \sqrt{\frac{995}{\pi}} \approx 0.04466$ and $\sqrt{\frac{1005}{\pi}} - \sqrt{\frac{1000}{\pi}} \approx 0.04455$. So if the machinist gets the radius within 0.0445 cm of 17.8412, the area will be within 5 cm^2 of 1000.

(c) x is the radius, $f(x)$ is the area, a is the target radius given in part (a), L is the target area (1000 cm^2), ε is the magnitude of the error tolerance in the area (5 cm^2), and δ is the tolerance in the radius given in part (b).

12. (a) $T = 0.1w^2 + 2.155w + 20$ and $T = 200 \quad \Rightarrow$

$0.1w^2 + 2.155w + 20 = 200 \quad \Rightarrow$ [by the quadratic formula or from the graph] $\quad w \approx 33.0$ watts $(w > 0)$

(b) From the graph, $199 \le T \le 201 \quad \Rightarrow \quad 32.89 < w < 33.11$.

(c) x is the input power, $f(x)$ is the temperature, a is the target input power given in part (a), L is the target temperature (200), ε is the tolerance in the temperature (1), and δ is the tolerance in the power input in watts indicated in part (b) (0.11 watts).

13. (a) $|4x - 8| = 4\,|x - 2| < 0.1 \quad \Leftrightarrow \quad |x - 2| < \dfrac{0.1}{4}$, so $\delta = \dfrac{0.1}{4} = 0.025$.

(b) $|4x - 8| = 4\,|x - 2| < 0.01 \quad \Leftrightarrow \quad |x - 2| < \dfrac{0.01}{4}$, so $\delta = \dfrac{0.01}{4} = 0.0025$.

14. $|(5x - 7) - 3| = |5x - 10| = |5(x - 2)| = 5\,|x - 2|$. We must have $|f(x) - L| < \varepsilon$, so $5\,|x - 2| < \varepsilon \quad \Leftrightarrow$ $|x - 2| < \varepsilon/5$. Thus, choose $\delta = \varepsilon/5$. For $\varepsilon = 0.1$, $\delta = 0.02$; for $\varepsilon = 0.05$, $\delta = 0.01$; for $\varepsilon = 0.01$, $\delta = 0.002$.

15. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x - 4| < \delta$, then $\left|\left(\frac{1}{2}x - 1\right) - 1\right| < \varepsilon$. But $\left|\left(\frac{1}{2}x - 1\right) - 1\right| < \varepsilon \quad \Leftrightarrow \quad \left|\frac{1}{2}x - 2\right| < \varepsilon \quad \Leftrightarrow$ $\left|\frac{1}{2}\right| |x - 4| < \varepsilon \quad \Leftrightarrow \quad |x - 4| < 2\varepsilon$. So if we choose $\delta = 2\varepsilon$, then $0 < |x - 4| < \delta \quad \Rightarrow \quad \left|\left(\frac{1}{2}x - 1\right) - 1\right| < \varepsilon$. Thus, $\lim\limits_{x \to 4} \left(\frac{1}{2}x - 1\right) = 1$ by the definition of a limit.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-2| < \delta$, then $|(2-3x)-(-4)| < \varepsilon$. But $|(2-3x)-(-4)| < \varepsilon \quad \Leftrightarrow$ $|6-3x| < \varepsilon \quad \Leftrightarrow \quad |-3|\,|x-2| < \varepsilon \quad \Leftrightarrow \quad |x-2| < \frac{1}{3}\varepsilon$. So if we choose $\delta = \frac{1}{3}\varepsilon$, then $0 < |x-2| < \delta \quad \Rightarrow \quad |(2-3x)-(-4)| < \varepsilon$. Thus, $\lim\limits_{x\to 2}(2-3x) = -4$ by the definition of a limit.

17. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-(-2)| < \delta$, then $|(-2x+1)-5| < \varepsilon$. But $|(-2x+1)-5| < \varepsilon \quad \Leftrightarrow$ $|-2x-4| < \varepsilon \quad \Leftrightarrow \quad |-2|\,|x-(-2)| < \varepsilon \quad \Leftrightarrow \quad |x-(-2)| < \frac{1}{2}\varepsilon$. So if we choose $\delta = \frac{1}{2}\varepsilon$, then $0 < |x-(-2)| < \delta \quad \Rightarrow$ $|(-2x+1)-5| < \varepsilon$. Thus, $\lim\limits_{x\to -2}(-2x+1) = 5$ by the definition of a limit.

18. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-1| < \delta$, then $|(2x-5)-(-3)| < \varepsilon$. But $|(2x-5)-(-3)| < \varepsilon \quad \Leftrightarrow$ $|2x-2| < \varepsilon \quad \Leftrightarrow \quad |2|\,|x-1| < \varepsilon \quad \Leftrightarrow \quad |x-1| < \frac{1}{2}\varepsilon$. So if we choose $\delta = \frac{1}{2}\varepsilon$, then $0 < |x-1| < \delta \quad \Rightarrow \quad |(2x-5)-(-3)| < \varepsilon$. Thus, $\lim\limits_{x\to 1}(2x-5) = -3$ by the definition of a limit.

19. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-9| < \delta$, then $\left|\left(1-\frac{1}{3}x\right)-(-2)\right| < \varepsilon$. But $\left|\left(1-\frac{1}{3}x\right)-(-2)\right| < \varepsilon \quad \Leftrightarrow$ $\left|3-\frac{1}{3}x\right| < \varepsilon \quad \Leftrightarrow \quad \left|-\frac{1}{3}\right|\,|x-9| < \varepsilon \quad \Leftrightarrow \quad |x-9| < 3\varepsilon$. So if we choose $\delta = 3\varepsilon$, then $0 < |x-9| < \delta \quad \Rightarrow$ $\left|\left(1-\frac{1}{3}x\right)-(-2)\right| < \varepsilon$. Thus, $\lim\limits_{x\to 9}\left(1-\frac{1}{3}x\right) = -2$ by the definition of a limit.

20. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-5| < \delta$, then $\left|\left(\frac{3}{2}x-\frac{1}{2}\right)-7\right| < \varepsilon$. But $\left|\left(\frac{3}{2}x-\frac{1}{2}\right)-7\right| < \varepsilon \quad \Leftrightarrow$ $\left|\frac{3}{2}x-\frac{15}{2}\right| < \varepsilon \quad \Leftrightarrow \quad \left|\frac{3}{2}\right|\,|x-5| < \varepsilon \quad \Leftrightarrow \quad |x-5| < \frac{2}{3}\varepsilon$. So if we choose $\delta = \frac{2}{3}\varepsilon$, then $0 < |x-5| < \delta \quad \Rightarrow$ $\left|\left(\frac{3}{2}x-\frac{1}{2}\right)-7\right| < \varepsilon$. Thus, $\lim\limits_{x\to 5}\left(\frac{3}{2}x-\frac{1}{2}\right) = 7$ by the definition of a limit.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-4| < \delta$, then $\left|\dfrac{x^2-2x-8}{x-4} - 6\right| < \varepsilon \quad \Leftrightarrow$

$\left|\dfrac{(x-4)(x+2)}{x-4} - 6\right| < \varepsilon \quad \Leftrightarrow \quad |x+2-6| < \varepsilon \quad [x \neq 4] \quad \Leftrightarrow \quad |x-4| < \varepsilon$. So choose $\delta = \varepsilon$. Then

$0 < |x-4| < \delta \quad \Rightarrow \quad |x-4| < \varepsilon \quad \Rightarrow \quad |x+2-6| < \varepsilon \quad \Rightarrow \quad \left|\dfrac{(x-4)(x+2)}{x-4} - 6\right| < \varepsilon \quad [x \neq 4] \quad \Rightarrow$

$\left|\dfrac{x^2-2x-8}{x-4} - 6\right| < \varepsilon$. By the definition of a limit, $\displaystyle\lim_{x\to 4} \frac{x^2-2x-8}{x-4} = 6$.

22. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x+1.5| < \delta$, then $\left|\dfrac{9-4x^2}{3+2x} - 6\right| < \varepsilon \quad \Leftrightarrow$

$\left|\dfrac{(3+2x)(3-2x)}{3+2x} - 6\right| < \varepsilon \quad \Leftrightarrow \quad |3-2x-6| < \varepsilon \quad [x \neq -1.5] \quad \Leftrightarrow \quad |-2x-3| < \varepsilon \quad \Leftrightarrow \quad |-2|\,|x+1.5| < \varepsilon \quad \Leftrightarrow$

$|x+1.5| < \varepsilon/2$. So choose $\delta = \varepsilon/2$. Then $0 < |x+1.5| < \delta \quad \Rightarrow \quad |x+1.5| < \varepsilon/2 \quad \Rightarrow \quad |-2|\,|x+1.5| < \varepsilon \quad \Rightarrow$

$|-2x-3| < \varepsilon \quad \Rightarrow \quad |3-2x-6| < \varepsilon \quad \Rightarrow \quad \left|\dfrac{(3+2x)(3-2x)}{3+2x} - 6\right| < \varepsilon \quad [x \neq -1.5] \quad \Rightarrow \quad \left|\dfrac{9-4x^2}{3+2x} - 6\right| < \varepsilon$.

By the definition of a limit, $\displaystyle\lim_{x\to -1.5} \frac{9-4x^2}{3+2x} = 6$.

23. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-a| < \delta$, then $|x-a| < \varepsilon$. So $\delta = \varepsilon$ will work.

24. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-a| < \delta$, then $|c-c| < \varepsilon$. But $|c-c| = 0$, so this will be true no matter what δ we pick.

25. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-0| < \delta$, then $\left|x^2-0\right| < \varepsilon \quad \Leftrightarrow \quad x^2 < \varepsilon \quad \Leftrightarrow \quad |x| < \sqrt{\varepsilon}$. Take $\delta = \sqrt{\varepsilon}$. Then $0 < |x-0| < \delta \quad \Rightarrow \quad \left|x^2-0\right| < \varepsilon$. Thus, $\displaystyle\lim_{x\to 0} x^2 = 0$ by the definition of a limit.

26. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-0| < \delta$, then $\left|x^3-0\right| < \varepsilon \quad \Leftrightarrow \quad |x|^3 < \varepsilon \quad \Leftrightarrow \quad |x| < \sqrt[3]{\varepsilon}$. Take $\delta = \sqrt[3]{\varepsilon}$. Then $0 < |x-0| < \delta \quad \Rightarrow \quad \left|x^3-0\right| < \delta^3 = \varepsilon$. Thus, $\displaystyle\lim_{x\to 0} x^3 = 0$ by the definition of a limit.

27. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-0| < \delta$, then $\big||x|-0\big| < \varepsilon$. But $\big||x|\big| = |x|$. So this is true if we pick $\delta = \varepsilon$. Thus, $\displaystyle\lim_{x\to 0} |x| = 0$ by the definition of a limit.

28. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < x-(-6) < \delta$, then $\left|\sqrt[8]{6+x} - 0\right| < \varepsilon$. But $\left|\sqrt[8]{6+x} - 0\right| < \varepsilon \quad \Leftrightarrow$

$\sqrt[8]{6+x} < \varepsilon \quad \Leftrightarrow \quad 6+x < \varepsilon^8 \quad \Leftrightarrow \quad x-(-6) < \varepsilon^8$. So if we choose $\delta = \varepsilon^8$, then $0 < x-(-6) < \delta \quad \Rightarrow$

$\left|\sqrt[8]{6+x} - 0\right| < \varepsilon$. Thus, $\displaystyle\lim_{x\to -6^+} \sqrt[8]{6+x} = 0$ by the definition of a right-hand limit.

29. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-2| < \delta$, then $\left|\left(x^2-4x+5\right)-1\right| < \varepsilon \quad \Leftrightarrow \quad \left|x^2-4x+4\right| < \varepsilon \quad \Leftrightarrow$

$\left|(x-2)^2\right| < \varepsilon$. So take $\delta = \sqrt{\varepsilon}$. Then $0 < |x-2| < \delta \quad \Leftrightarrow \quad |x-2| < \sqrt{\varepsilon} \quad \Leftrightarrow \quad \left|(x-2)^2\right| < \varepsilon$. Thus,

$\displaystyle\lim_{x\to 2} \left(x^2-4x+5\right) = 1$ by the definition of a limit.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-2| < \delta$, then $\left|(x^2+2x-7)-1\right| < \varepsilon$. But $\left|(x^2+2x-7)-1\right| < \varepsilon \quad \Leftrightarrow$ $\left|x^2+2x-8\right| < \varepsilon \quad \Leftrightarrow \quad |x+4|\,|x-2| < \varepsilon$. Thus our goal is to make $|x-2|$ small enough so that its product with $|x+4|$ is less than ε. Suppose we first require that $|x-2| < 1$. Then $-1 < x-2 < 1 \quad \Rightarrow \quad 1 < x < 3 \quad \Rightarrow \quad 5 < x+4 < 7 \quad \Rightarrow$ $|x+4| < 7$, and this gives us $7\,|x-2| < \varepsilon \quad \Rightarrow \quad |x-2| < \varepsilon/7$. Choose $\delta = \min\{1, \varepsilon/7\}$. Then if $0 < |x-2| < \delta$, we have $|x-2| < \varepsilon/7$ and $|x+4| < 7$, so $\left|(x^2+2x-7)-1\right| = |(x+4)(x-2)| = |x+4|\,|x-2| < 7(\varepsilon/7) = \varepsilon$, as desired. Thus, $\lim\limits_{x\to 2}(x^2+2x-7) = 1$ by the definition of a limit.

31. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-(-2)| < \delta$, then $\left|\left(x^2-1\right)-3\right| < \varepsilon$ or upon simplifying we need $\left|x^2-4\right| < \varepsilon$ whenever $0 < |x+2| < \delta$. Notice that if $|x+2| < 1$, then $-1 < x+2 < 1 \quad \Rightarrow \quad -5 < x-2 < -3 \quad \Rightarrow$ $|x-2| < 5$. So take $\delta = \min\{\varepsilon/5, 1\}$. Then $0 < |x+2| < \delta \quad \Rightarrow \quad |x-2| < 5$ and $|x+2| < \varepsilon/5$, so $\left|\left(x^2-1\right)-3\right| = |(x+2)(x-2)| = |x+2|\,|x-2| < (\varepsilon/5)(5) = \varepsilon$. Thus, by the definition of a limit, $\lim\limits_{x\to -2}(x^2-1) = 3$.

32. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-2| < \delta$, then $\left|x^3-8\right| < \varepsilon$. Now $\left|x^3-8\right| = \left|(x-2)\left(x^2+2x+4\right)\right|$. If $|x-2| < 1$, that is, $1 < x < 3$, then $x^2+2x+4 \;<\; 3^2+2(3)+4 = 19$ and so $\left|x^3-8\right| = |x-2|\left(x^2+2x+4\right) < 19\,|x-2|$. So if we take $\delta = \min\left\{1, \frac{\varepsilon}{19}\right\}$, then $0 < |x-2| < \delta \quad \Rightarrow$ $\left|x^3-8\right| = |x-2|\left(x^2+2x+4\right) < \frac{\varepsilon}{19}\cdot 19 = \varepsilon$. Thus, by the definition of a limit, $\lim\limits_{x\to 2} x^3 = 8$.

33. Given $\varepsilon > 0$, we let $\delta = \min\left\{2, \frac{\varepsilon}{8}\right\}$. If $0 < |x-3| < \delta$, then $|x-3| < 2 \quad \Rightarrow \quad -2 < x-3 < 2 \quad \Rightarrow$ $4 < x+3 < 8 \quad \Rightarrow \quad |x+3| < 8$. Also $|x-3| < \frac{\varepsilon}{8}$, so $\left|x^2-9\right| = |x+3|\,|x-3| < 8\cdot\frac{\varepsilon}{8} = \varepsilon$. Thus, $\lim\limits_{x\to 3} x^2 = 9$.

34. From the figure, our choices for δ are $\delta_1 = 3-\sqrt{9-\varepsilon}$ and $\delta_2 = \sqrt{9+\varepsilon}-3$. The *largest* possible choice for δ is the minimum value of $\{\delta_1, \delta_2\}$; that is, $\delta = \min\{\delta_1, \delta_2\} = \delta_2 = \sqrt{9+\varepsilon}-3$.

35. (a) The points of intersection in the graph are $(x_1, 2.6)$ and $(x_2, 3.4)$ with $x_1 \approx 0.891$ and $x_2 \approx 1.093$. Thus, we can take δ to be the smaller of $1-x_1$ and x_2-1. So $\delta = x_2 - 1 \approx 0.093$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Solving $x^3 + x + 1 = 3 + \varepsilon$ with a CAS gives us two nonreal complex solutions and one real solution, which is

$$x(\varepsilon) = \frac{\left(216 + 108\varepsilon + 12\sqrt{336 + 324\varepsilon + 81\varepsilon^2}\,\right)^{2/3} - 12}{6\left(216 + 108\varepsilon + 12\sqrt{336 + 324\varepsilon + 81\varepsilon^2}\,\right)^{1/3}}. \text{ Thus, } \delta = x(\varepsilon) - 1.$$

(c) If $\varepsilon = 0.4$, then $x(\varepsilon) \approx 1.093\,272\,342$ and $\delta = x(\varepsilon) - 1 \approx 0.093$, which agrees with our answer in part (a).

36. *1. Guessing a value for δ* Let $\varepsilon > 0$ be given. We have to find a number $\delta > 0$ such that $\left|\dfrac{1}{x} - \dfrac{1}{2}\right| < \varepsilon$ whenever $0 < |x - 2| < \delta$. But $\left|\dfrac{1}{x} - \dfrac{1}{2}\right| = \left|\dfrac{2 - x}{2x}\right| = \dfrac{|x - 2|}{|2x|} < \varepsilon$. We find a positive constant C such that $\dfrac{1}{|2x|} < C \quad \Rightarrow$ $\dfrac{|x - 2|}{|2x|} < C\,|x - 2|$ and we can make $C\,|x - 2| < \varepsilon$ by taking $|x - 2| < \dfrac{\varepsilon}{C} = \delta$. We restrict x to lie in the interval $|x - 2| < 1 \quad \Rightarrow \quad 1 < x < 3$ so $1 > \dfrac{1}{x} > \dfrac{1}{3} \quad \Rightarrow \quad \dfrac{1}{6} < \dfrac{1}{2x} < \dfrac{1}{2} \quad \Rightarrow \quad \dfrac{1}{|2x|} < \dfrac{1}{2}$. So $C = \dfrac{1}{2}$ is suitable. Thus, we should choose $\delta = \min\{1, 2\varepsilon\}$.

2. Showing that δ works Given $\varepsilon > 0$ we let $\delta = \min\{1, 2\varepsilon\}$. If $0 < |x - 2| < \delta$, then $|x - 2| < 1 \quad \Rightarrow \quad 1 < x < 3 \quad \Rightarrow$ $\dfrac{1}{|2x|} < \dfrac{1}{2}$ (as in part 1). Also $|x - 2| < 2\varepsilon$, so $\left|\dfrac{1}{x} - \dfrac{1}{2}\right| = \dfrac{|x - 2|}{|2x|} < \dfrac{1}{2} \cdot 2\varepsilon = \varepsilon$. This shows that $\lim\limits_{x \to 2} (1/x) = \frac{1}{2}$.

37. *1. Guessing a value for δ* Given $\varepsilon > 0$, we must find $\delta > 0$ such that $|\sqrt{x} - \sqrt{a}| < \varepsilon$ whenever $0 < |x - a| < \delta$. But $|\sqrt{x} - \sqrt{a}| = \dfrac{|x - a|}{\sqrt{x} + \sqrt{a}} < \varepsilon$ (from the hint). Now if we can find a positive constant C such that $\sqrt{x} + \sqrt{a} > C$ then $\dfrac{|x - a|}{\sqrt{x} + \sqrt{a}} < \dfrac{|x - a|}{C} < \varepsilon$, and we take $|x - a| < C\varepsilon$. We can find this number by restricting x to lie in some interval centered at a. If $|x - a| < \frac{1}{2}a$, then $-\frac{1}{2}a < x - a < \frac{1}{2}a \quad \Rightarrow \quad \frac{1}{2}a < x < \frac{3}{2}a \quad \Rightarrow \quad \sqrt{x} + \sqrt{a} > \sqrt{\frac{1}{2}a} + \sqrt{a}$, and so $C = \sqrt{\frac{1}{2}a} + \sqrt{a}$ is a suitable choice for the constant. So $|x - a| < \left(\sqrt{\frac{1}{2}a} + \sqrt{a}\right)\varepsilon$. This suggests that we let $\delta = \min\left\{\frac{1}{2}a, \left(\sqrt{\frac{1}{2}a} + \sqrt{a}\right)\varepsilon\right\}$.

2. Showing that δ works Given $\varepsilon > 0$, we let $\delta = \min\left\{\frac{1}{2}a, \left(\sqrt{\frac{1}{2}a} + \sqrt{a}\right)\varepsilon\right\}$. If $0 < |x - a| < \delta$, then $|x - a| < \frac{1}{2}a \quad \Rightarrow \quad \sqrt{x} + \sqrt{a} > \sqrt{\frac{1}{2}a} + \sqrt{a}$ (as in part 1). Also $|x - a| < \left(\sqrt{\frac{1}{2}a} + \sqrt{a}\right)\varepsilon$, so

$$|\sqrt{x} - \sqrt{a}\,| = \frac{|x - a|}{\sqrt{x} + \sqrt{a}} < \frac{\left(\sqrt{a/2} + \sqrt{a}\right)\varepsilon}{\left(\sqrt{a/2} + \sqrt{a}\right)} = \varepsilon. \text{ Therefore, } \lim_{x \to a} \sqrt{x} = \sqrt{a} \text{ by the definition of a limit.}$$

38. Suppose that $\lim\limits_{t \to 0} H(t) = L$. Given $\varepsilon = \frac{1}{2}$, there exists $\delta > 0$ such that $0 < |t| < \delta \quad \Rightarrow \quad |H(t) - L| < \frac{1}{2} \quad \Leftrightarrow$ $L - \frac{1}{2} < H(t) < L + \frac{1}{2}$. For $0 < t < \delta$, $H(t) = 1$, so $1 < L + \frac{1}{2} \quad \Rightarrow \quad L > \frac{1}{2}$. For $-\delta < t < 0$, $H(t) = 0$, so $L - \frac{1}{2} < 0 \quad \Rightarrow \quad L < \frac{1}{2}$. This contradicts $L > \frac{1}{2}$. Therefore, $\lim\limits_{t \to 0} H(t)$ does not exist.

39. Suppose that $\lim\limits_{x \to 0} f(x) = L$. Given $\varepsilon = \frac{1}{2}$, there exists $\delta > 0$ such that $0 < |x| < \delta \quad \Rightarrow \quad |f(x) - L| < \frac{1}{2}$. Take any rational number r with $0 < |r| < \delta$. Then $f(r) = 0$, so $|0 - L| < \frac{1}{2}$, so $L \le |L| < \frac{1}{2}$. Now take any irrational number s with

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$0 < |s| < \delta$. Then $f(s) = 1$, so $|1 - L| < \frac{1}{2}$. Hence, $1 - L < \frac{1}{2}$, so $L > \frac{1}{2}$. This contradicts $L < \frac{1}{2}$, so $\lim_{x \to 0} f(x)$ does not exist.

40. First suppose that $\lim_{x \to a} f(x) = L$. Then, given $\varepsilon > 0$ there exists $\delta > 0$ so that $0 < |x - a| < \delta \quad \Rightarrow \quad |f(x) - L| < \varepsilon$. Then $a - \delta < x < a \quad \Rightarrow \quad 0 < |x - a| < \delta$ so $|f(x) - L| < \varepsilon$. Thus, $\lim_{x \to a^-} f(x) = L$. Also $a < x < a + \delta \quad \Rightarrow$ $0 < |x - a| < \delta$ so $|f(x) - L| < \varepsilon$. Hence, $\lim_{x \to a^+} f(x) = L$.

Now suppose $\lim_{x \to a^-} f(x) = L = \lim_{x \to a^+} f(x)$. Let $\varepsilon > 0$ be given. Since $\lim_{x \to a^-} f(x) = L$, there exists $\delta_1 > 0$ so that $a - \delta_1 < x < a \quad \Rightarrow \quad |f(x) - L| < \varepsilon$. Since $\lim_{x \to a^+} f(x) = L$, there exists $\delta_2 > 0$ so that $a < x < a + \delta_2 \quad \Rightarrow$ $|f(x) - L| < \varepsilon$. Let δ be the smaller of δ_1 and δ_2. Then $0 < |x - a| < \delta \quad \Rightarrow \quad a - \delta_1 < x < a$ or $a < x < a + \delta_2$ so $|f(x) - L| < \varepsilon$. Hence, $\lim_{x \to a} f(x) = L$. So we have proved that $\lim_{x \to a} f(x) = L \quad \Leftrightarrow \quad \lim_{x \to a^-} f(x) = L = \lim_{x \to a^+} f(x)$.

41. $\dfrac{1}{(x+3)^4} > 10{,}000 \quad \Leftrightarrow \quad (x+3)^4 < \dfrac{1}{10{,}000} \quad \Leftrightarrow \quad |x+3| < \dfrac{1}{\sqrt[4]{10{,}000}} \quad \Leftrightarrow \quad |x - (-3)| < \dfrac{1}{10}$

42. Given $M > 0$, we need $\delta > 0$ such that $0 < |x+3| < \delta \quad \Rightarrow \quad 1/(x+3)^4 > M$. Now $\dfrac{1}{(x+3)^4} > M \quad \Leftrightarrow$ $(x+3)^4 < \dfrac{1}{M} \quad \Leftrightarrow \quad |x+3| < \dfrac{1}{\sqrt[4]{M}}$. So take $\delta = \dfrac{1}{\sqrt[4]{M}}$. Then $0 < |x+3| < \delta = \dfrac{1}{\sqrt[4]{M}} \quad \Rightarrow \quad \dfrac{1}{(x+3)^4} > M$, so $\lim_{x \to -3} \dfrac{1}{(x+3)^4} = \infty$.

43. Given $M < 0$ we need $\delta > 0$ so that $\ln x < M$ whenever $0 < x < \delta$; that is, $x = e^{\ln x} < e^M$ whenever $0 < x < \delta$. This suggests that we take $\delta = e^M$. If $0 < x < e^M$, then $\ln x < \ln e^M = M$. By the definition of a limit, $\lim_{x \to 0^+} \ln x = -\infty$.

44. (a) Let M be given. Since $\lim_{x \to a} f(x) = \infty$, there exists $\delta_1 > 0$ such that $0 < |x - a| < \delta_1 \quad \Rightarrow \quad f(x) > M + 1 - c$. Since $\lim_{x \to a} g(x) = c$, there exists $\delta_2 > 0$ such that $0 < |x - a| < \delta_2 \quad \Rightarrow \quad |g(x) - c| < 1 \quad \Rightarrow \quad g(x) > c - 1$. Let δ be the smaller of δ_1 and δ_2. Then $0 < |x - a| < \delta \quad \Rightarrow \quad f(x) + g(x) > (M + 1 - c) + (c - 1) = M$. Thus, $\lim_{x \to a} [f(x) + g(x)] = \infty$.

(b) Let $M > 0$ be given. Since $\lim_{x \to a} g(x) = c > 0$, there exists $\delta_1 > 0$ such that $0 < |x - a| < \delta_1 \quad \Rightarrow$ $|g(x) - c| < c/2 \quad \Rightarrow \quad g(x) > c/2$. Since $\lim_{x \to a} f(x) = \infty$, there exists $\delta_2 > 0$ such that $0 < |x - a| < \delta_2 \quad \Rightarrow$ $f(x) > 2M/c$. Let $\delta = \min\{\delta_1, \delta_2\}$. Then $0 < |x - a| < \delta \quad \Rightarrow \quad f(x)\,g(x) > \dfrac{2M}{c}\,\dfrac{c}{2} = M$, so $\lim_{x \to a} f(x)\,g(x) = \infty$.

(c) Let $N < 0$ be given. Since $\lim_{x \to a} g(x) = c < 0$, there exists $\delta_1 > 0$ such that $0 < |x - a| < \delta_1 \quad \Rightarrow$ $|g(x) - c| < -c/2 \quad \Rightarrow \quad g(x) < c/2$. Since $\lim_{x \to a} f(x) = \infty$, there exists $\delta_2 > 0$ such that $0 < |x - a| < \delta_2 \quad \Rightarrow$ $f(x) > 2N/c$. (Note that $c < 0$ and $N < 0 \quad \Rightarrow \quad 2N/c > 0$.) Let $\delta = \min\{\delta_1, \delta_2\}$. Then $0 < |x - a| < \delta \quad \Rightarrow$ $f(x) > 2N/c \quad \Rightarrow \quad f(x)\,g(x) < \dfrac{2N}{c} \cdot \dfrac{c}{2} = N$, so $\lim_{x \to a} f(x)\,g(x) = -\infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2.5 Continuity

1. From Definition 1, $\lim_{x \to 4} f(x) = f(4)$.

2. The graph of f has no hole, jump, or vertical asymptote.

3. (a) f is discontinuous at -4 since $f(-4)$ is not defined and at -2, 2, and 4 since the limit does not exist (the left and right limits are not the same).

(b) f is continuous from the left at -2 since $\lim_{x \to -2^-} f(x) = f(-2)$. f is continuous from the right at 2 and 4 since $\lim_{x \to 2^+} f(x) = f(2)$ and $\lim_{x \to 4^+} f(x) = f(4)$. The function is not continuous from either side at -4 since $f(-4)$ is undefined.

4. g is not continuous at -2 since $g(-2)$ is not defined. g is not continuous at $a = -1$ since the limit does not exist (the left and right limits are $-\infty$). g is not continuous at $a = 0$ and $a = 1$ since the limit does not exist (the left and right limits are not equal).

5. (a) From the graph we see that $\lim_{x \to a} f(x)$ does not exist at $a = 1$ since the left and right limits are not the same.

(b) f is not continuous at $a = 1$ since $\lim_{x \to 1} f(x)$ does not exist by part (a). Also, f is not continuous at $a = 3$ since $\lim_{x \to 3} f(x) \neq f(3)$.

(c) From the graph we see that $\lim_{x \to 3} f(x) = 3$, but $f(3) = 2$. Since the limit is not equal to $f(3)$, f is not continuous at $a = 3$.

6. (a) From the graph we see that $\lim_{x \to a} f(x)$ does not exist at $a = 1$ since the function increases without bound from the left and from the right. Also, $\lim_{x \to a} f(x)$ does not exist at $a = 5$ since the left and right limits are not the same.

(b) f is not continuous at $a = 1$ and at $a = 5$ since the limits do not exist by part (a). Also, f is not continuous at $a = 3$ since $\lim_{x \to 3} f(x) \neq f(3)$.

(c) From the graph we see that $\lim_{x \to 3} f(x)$ exists, but the limit is not equal to $f(3)$, so f is not continuous at $a = 3$.

7.

The graph of $y = f(x)$ must have a removable discontinuity (a hole) at $x = -2$ and an infinite discontinuity (a vertical asymptote) at $x = 2$.

8.

The graph of $y = f(x)$ must have a jump discontinuity at $x = -3$ and a removable discontinuity (a hole) at $x = 4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9.

The graph of $y = f(x)$ must have discontinuities at $x = 0$ and $x = 3$. It must show that $\lim_{x \to 0^+} f(x) = f(0)$ and $\lim_{x \to 3^-} f(x) = f(3)$.

10.

The graph of $y = f(x)$ must have a discontinuity at $x = -1$ with $\lim_{x \to -1^-} f(x) = f(-1)$ and $\lim_{x \to -1^+} f(x) \neq f(-1)$. The graph must also show that $\lim_{x \to 3^-} f(x) \neq f(3)$ and $\lim_{x \to 3^+} f(x) \neq f(3)$.

11. (a) The toll is \$5 except between 7:00 AM and 10:00 AM and between 4:00 PM and 7:00 PM, when the toll is \$7.

(b) The function T has jump discontinuities at $t = 7$, 10, 16, and 19. Their significance to someone who uses the road is that, because of the sudden jumps in the toll, they may want to avoid the higher rates between $t = 7$ and $t = 10$ and between $t = 16$ and $t = 19$ if feasible.

12. (a) Continuous; at the location in question, the temperature changes smoothly as time passes, without any instantaneous jumps from one temperature to another.

(b) Continuous; the temperature at a specific time changes smoothly as the distance due west from New York City increases, without any instantaneous jumps.

(c) Discontinuous; as the distance due west from New York City increases, the altitude above sea level may jump from one height to another without going through all of the intermediate values — at a cliff, for example.

(d) Discontinuous; as the distance traveled increases, the cost of the ride jumps in small increments.

(e) Discontinuous; when the lights are switched on (or off), the current suddenly changes between 0 and some nonzero value, without passing through all of the intermediate values. This is debatable, though, depending on your definition of current.

13. $$\lim_{x \to -1} f(x) = \lim_{x \to -1} \left[3x^2 + (x+2)^5\right] = \lim_{x \to -1} 3x^2 + \lim_{x \to -1} (x+2)^5 = 3 \lim_{x \to -1} x^2 + \lim_{x \to -1} (x+2)^5$$
$$= 3(-1)^2 + (-1+2)^5 = 4 = f(-1)$$

By the definition of continuity, f is continuous at $a = -1$.

14. $$\lim_{t \to 2} g(t) = \lim_{t \to 2} \frac{t^2 + 5t}{2t + 1} = \frac{\lim_{t \to 2}(t^2 + 5t)}{\lim_{t \to 2}(2t + 1)} = \frac{\lim_{t \to 2} t^2 + 5 \lim_{t \to 2} t}{2 \lim_{t \to 2} t + \lim_{t \to 2} 1} = \frac{2^2 + 5(2)}{2(2) + 1} = \frac{14}{5} = g(2).$$

By the definition of continuity, g is continuous at $a = 2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $\lim\limits_{v\to 1} p(v) = \lim\limits_{v\to 1} 2\sqrt{3v^2+1} = 2\lim\limits_{v\to 1}\sqrt{3v^2+1} = 2\sqrt{\lim\limits_{v\to 1}(3v^2+1)} = 2\sqrt{3\lim\limits_{v\to 1} v^2 + \lim\limits_{v\to 1} 1}$

$= 2\sqrt{3(1)^2+1} = 2\sqrt{4} = 4 = p(1)$

By the definition of continuity, p is continuous at $a = 1$.

16. $\lim\limits_{r\to -2} f(r) = \lim\limits_{r\to -2} \sqrt[3]{4r^2-2r+7} = \sqrt[3]{\lim\limits_{r\to -2}(4r^2-2r+7)} = \sqrt[3]{4(-2)^2-2(-2)+7} = \sqrt[3]{27} = 3 = f(-2)$

By the definition of continuity, f is continuous at $a = -2$.

17. For $a > 4$, we have

$$\begin{aligned}
\lim_{x\to a} f(x) &= \lim_{x\to a}(x+\sqrt{x-4}) = \lim_{x\to a} x + \lim_{x\to a}\sqrt{x-4} && \text{[Limit Law 1]}\\
&= a + \sqrt{\lim_{x\to a} x - \lim_{x\to a} 4} && \text{[8, 11, and 2]}\\
&= a + \sqrt{a-4} && \text{[8 and 7]}\\
&= f(a)
\end{aligned}$$

So f is continuous at $x = a$ for every a in $(4, \infty)$. Also, $\lim\limits_{x\to 4^+} f(x) = 4 = f(4)$, so f is continuous from the right at 4.

Thus, f is continuous on $[4, \infty)$.

18. For $a < -2$, we have

$$\begin{aligned}
\lim_{x\to a} g(x) &= \lim_{x\to a}\frac{x-1}{3x+6} = \frac{\lim\limits_{x\to a}(x-1)}{\lim\limits_{x\to a}(3x+6)} && \text{[Limit Law 5]}\\
&= \frac{\lim\limits_{x\to a} x - \lim\limits_{x\to a} 1}{3\lim\limits_{x\to a} x + \lim\limits_{x\to a} 6} && \text{[2, 1, and 3]}\\
&= \frac{a-1}{3a+6} && \text{[8 and 7]}
\end{aligned}$$

Thus, g is continuous at $x = a$ for every a in $(-\infty, -2)$; that is, g is continuous on $(-\infty, -2)$.

19. $f(x) = \dfrac{1}{x+2}$ is discontinuous at $a = -2$ because $f(-2)$ is undefined.

20. $f(x) = \begin{cases} \dfrac{1}{x+2} & \text{if } x \neq -2 \\ 1 & \text{if } x = -2 \end{cases}$

Here $f(-2) = 1$, but $\lim\limits_{x\to -2^-} f(x) = -\infty$ and $\lim\limits_{x\to -2^+} f(x) = \infty$,

so $\lim\limits_{x\to -2} f(x)$ does not exist and f is discontinuous at -2.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. $f(x) = \begin{cases} x+3 & \text{if } x \le -1 \\ 2^x & \text{if } x > -1 \end{cases}$

$\lim\limits_{x \to -1^-} f(x) = \lim\limits_{x \to -1^-} (x+3) = -1 + 3 = 2$ and

$\lim\limits_{x \to -1^+} f(x) = \lim\limits_{x \to -1^+} 2^x = 2^{-1} = \frac{1}{2}$. Since the left-hand and the right-hand limits of f at -1 are not equal, $\lim\limits_{x \to -1} f(x)$ does not exist, and f is discontinuous at -1.

22. $f(x) = \begin{cases} \dfrac{x^2 - x}{x^2 - 1} & \text{if } x \ne 1 \\ 1 & \text{if } x = 1 \end{cases}$

$\lim\limits_{x \to 1} f(x) = \lim\limits_{x \to 1} \dfrac{x^2 - x}{x^2 - 1} = \lim\limits_{x \to 1} \dfrac{x(x-1)}{(x+1)(x-1)} = \lim\limits_{x \to 1} \dfrac{x}{x+1} = \dfrac{1}{2}$,
but $f(1) = 1$, so f is discontinous at 1.

23. $f(x) = \begin{cases} \cos x & \text{if } x < 0 \\ 0 & \text{if } x = 0 \\ 1 - x^2 & \text{if } x > 0 \end{cases}$

$\lim\limits_{x \to 0} f(x) = 1$, but $f(0) = 0 \ne 1$, so f is discontinuous at 0.

24. $f(x) = \begin{cases} \dfrac{2x^2 - 5x - 3}{x - 3} & \text{if } x \ne 3 \\ 6 & \text{if } x = 3 \end{cases}$

$\lim\limits_{x \to 3} f(x) = \lim\limits_{x \to 3} \dfrac{2x^2 - 5x - 3}{x - 3} = \lim\limits_{x \to 3} \dfrac{(2x+1)(x-3)}{x-3} = \lim\limits_{x \to 3} (2x+1) = 7$,
but $f(3) = 6$, so f is discontinuous at 3.

25. (a) $f(x) = \dfrac{x-3}{x^2 - 9} = \dfrac{x-3}{(x-3)(x+3)} = \dfrac{1}{x+3}$ for $x \ne 3$. $f(3)$ is undefined, so f is discontinuous at $x = 3$. Further, $\lim\limits_{x \to 3} f(x) = \dfrac{1}{3+3} = \dfrac{1}{6}$. Since f is discontinuous at $x = 3$, but $\lim\limits_{x \to 3} f(x)$ exists, f has a removable discontinuity at $x = 3$.

(b) If f is redefined to be $\frac{1}{6}$ at $x = 3$, then f will be equivalent to the function $g(x) = \dfrac{1}{x+3}$, which is continuous at $x = 3$.

26. (a) $f(x) = \dfrac{x^2 - 7x + 12}{x - 3} = \dfrac{(x-3)(x-4)}{x-3} = x - 4$ for $x \ne 3$. $f(3)$ is undefined, so f is discontinuous at $x = 3$. Further, $\lim\limits_{x \to 3} f(x) = 3 - 4 = -1$. Since f is discontinuous at $x = 3$, but $\lim\limits_{x \to 3} f(x)$ exists, f has a removable discontinuity at $x = 3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) If f is redefined to be -1 at $x = 3$, then f will be equivalent to the function $g(x) = x - 4$, which is continuous everywhere (and is thus continuous at $x = 3$).

27. The domain of $f(x) = \dfrac{x^2}{\sqrt{x^4+2}}$ is $(-\infty, \infty)$ since the denominator is never 0. By Theorem 5(a), the polynomial x^2 is continuous everywhere. By Theorems 5(a), 7, and 9, $\sqrt{x^4+2}$ is continuous everywhere. Finally, by part 5 of Theorem 4, $f(x)$ is continuous everywhere.

28. $g(v) = \dfrac{3v-1}{v^2+2v-15} = \dfrac{3v-1}{(v+5)(v-3)}$ is a rational function, so it is continuous on its domain, $(-\infty, -5) \cup (-5, 3) \cup (3, \infty)$, by Theorem 5(b).

29. The domain of $h(t) = \dfrac{\cos(t^2)}{1-e^t}$ must exclude any value of t for which $1 - e^t = 0$. $1 - e^t = 0 \;\Rightarrow\; e^t = 1 \;\Rightarrow\; \ln(e^t) = \ln 1 \;\Rightarrow\; t = 0$, so the domain of $h(t)$ is $(-\infty, 0) \cup (0, \infty)$. By Theorems 7 and 9, $\cos(t^2)$ is continuous on $\mathbb{R}$. By Theorems 5 and 7 and part 2 of Theorem 4, $1 - e^t$ is continuous everywhere. Finally, by part 5 of Theorem 4, $h(t)$ is continuous on its domain.

30. $B(u) = \sqrt{3u-2} + \sqrt[3]{2u-3}$ is defined when $3u - 2 \geq 0 \;\Rightarrow\; 3u \geq 2 \;\Rightarrow\; u \geq \frac{2}{3}$. (Note that $\sqrt[3]{2u-3}$ is defined everywhere.) So B has domain $\left[\frac{2}{3}, \infty\right)$. By Theorems 7 and 9, $\sqrt{3u-2}$ and $\sqrt[3]{2u-3}$ are each continuous on their domain because each is the composite of a root function and a polynomial function. B is the sum of these two functions, so it is continuous at every number in its domain by part 1 of Theorem 4.

31. $L(v) = v\ln(1-v^2)$ is defined when $1 - v^2 > 0 \;\Leftrightarrow\; v^2 < 1 \;\Leftrightarrow\; |v| < 1 \;\Leftrightarrow\; -1 < v < 1$. Thus, L has domain $(-1, 1)$. Now v and the composite function $\ln(1-v^2)$ are continuous on their domains by Theorems 7 and 9. Thus, by part 4 of Theorem 4, $L(v)$ is continuous on its domain.

32. $f(t) = e^{-t^2}\ln(1+t^2)$ has domain $(-\infty, \infty)$ since $1 + t^2 > 0$. By Theorems 7 and 9, e^{-t^2} and $\ln(1+t^2)$ are continuous everywhere. Finally, by part 4 of Theorem 4, $f(t)$ is continuous everywhere.

33. $M(x) = \sqrt{1 + \dfrac{1}{x}} = \sqrt{\dfrac{x+1}{x}}$ is defined when $\dfrac{x+1}{x} \geq 0 \;\Rightarrow\; x + 1 \geq 0$ and $x > 0$ or $x + 1 \leq 0$ and $x < 0 \;\Rightarrow\; x > 0$ or $x \leq -1$, so M has domain $(-\infty, -1] \cup (0, \infty)$. M is the composite of a root function and a rational function, so it is continuous at every number in its domain by Theorems 7 and 9.

34. The function $g(t) = \cos^{-1}(e^t - 1)$ is defined for $-1 \leq e^t - 1 \leq 1 \;\Rightarrow\; 0 \leq e^t \leq 2 \;\Rightarrow\; \ln(e^t) \leq \ln 2$ [since e^t is always positive] $\;\Rightarrow\; t \leq \ln 2$, so the domain of g is $(-\infty, \ln 2]$. The function $e^t - 1$ is the difference of an exponential and a constant (polynomial) function, so it is continuous on its domain by Theorem 7 and part 2 of Theorem 4. The inverse trigonometric function $\cos^{-1} t$ is continuous on its domain by Theorem 7. The function g is then the composite of continuous functions, so by Theorem 9 it is continuous on its domain.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. Because x is continuous on $\mathbb{R}$ and $\sqrt{20-x^2}$ is continuous on its domain, $-\sqrt{20} \le x \le \sqrt{20}$, the product $f(x) = x\sqrt{20-x^2}$ is continuous on $-\sqrt{20} \le x \le \sqrt{20}$. The number 2 is in that domain, so f is continuous at 2, and $\lim\limits_{x\to 2} f(x) = f(2) = 2\sqrt{16} = 8$.

36. The function $f(\theta) = \sin(\tan(\cos\theta))$ is the composite of trigonometric functions, so it is continuous throughout its domain. Now the domain of $\cos\theta$ is $\mathbb{R}$, $-1 \le \cos\theta \le 1$, the domain of $\tan\theta$ includes $[-1, 1]$, and the domain of $\sin\theta$ is $\mathbb{R}$, so the domain of f is $\mathbb{R}$. Thus f is continuous at $\frac{\pi}{2}$, and $\lim\limits_{0\to\pi/2} \sin(\tan(\cos\theta)) = \sin(\tan(\cos\frac{\pi}{2})) = \sin(\tan(0)) = \sin(0) = 0$.

37. The function $f(x) = \ln\left(\dfrac{5-x^2}{1+x}\right)$ is continuous throughout its domain because it is the composite of a logarithm function and a rational function. For the domain of f, we must have $\dfrac{5-x^2}{1+x} > 0$, so the numerator and denominator must have the same sign, that is, the domain is $(-\infty, -\sqrt{5}\,] \cup (-1, \sqrt{5}\,]$. The number 1 is in that domain, so f is continuous at 1, and $\lim\limits_{x\to 1} f(x) = f(1) = \ln\dfrac{5-1}{1+1} = \ln 2$.

38. The function $f(x) = 3^{\sqrt{x^2-2x-4}}$ is continuous throughout its domain because it is the composite of an exponential function, a root function, and a polynomial. Its domain is

$$\begin{aligned}\{x \mid x^2-2x-4 \ge 0\} &= \{x \mid x^2-2x+1 \ge 5\} = \{x \mid (x-1)^2 \ge 5\} \\ &= \left\{x \,\middle|\, |x-1| \ge \sqrt{5}\,\right\} = (-\infty, 1-\sqrt{5}\,] \cup [1+\sqrt{5}, \infty)\end{aligned}$$

The number 4 is in that domain, so f is continuous at 4, and $\lim\limits_{x\to 4} f(x) = f(4) = 3^{\sqrt{16-8-4}} = 3^2 = 9$.

39. The function $f(x) = \dfrac{1}{\sqrt{1-\sin x}}$ is discontinuous wherever $1 - \sin x = 0 \;\Rightarrow\; \sin x = 1 \;\Rightarrow\; x = \frac{\pi}{2} + 2n\pi$, where n is any integer. The graph shows the discontinuities for $n = -1$, 0, and 1.

40. The function $y = \arctan(1/x)$ is discontinuous only where $1/x$ is undefined. Thus $y = \arctan(1/x)$ is discontinuous at $x = 0$. (From the graph, note also that the left- and right-hand limits at $x = 0$ are different.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. $f(x) = \begin{cases} 1 - x^2 & \text{if } x \le 1 \\ \ln x & \text{if } x > 1 \end{cases}$

By Theorem 5, since $f(x)$ equals the polynomial $1 - x^2$ on $(-\infty, 1]$, f is continuous on $(-\infty, 1]$.

By Theorem 7, since $f(x)$ equals the logarithm function $\ln x$ on $(1, \infty)$, f is continuous on $(1, \infty)$.

At $x = 1$, $\lim\limits_{x \to 1^-} f(x) = \lim\limits_{x \to 1^-} (1 - x^2) = 1 - 1^2 = 0$ and $\lim\limits_{x \to 1^+} f(x) = \lim\limits_{x \to 1^+} \ln x = \ln 1 = 0$. Thus, $\lim\limits_{x \to 1} f(x)$ exists and equals 0. Also, $f(1) = 1 - 1^2 = 0$. Thus, f is continuous at $x = 1$. We conclude that f is continuous on $(-\infty, \infty)$.

42. $f(x) = \begin{cases} \sin x & \text{if } x < \pi/4 \\ \cos x & \text{if } x \ge \pi/4 \end{cases}$

By Theorem 7, the trigonometric functions are continuous. Since $f(x) = \sin x$ on $(-\infty, \pi/4)$ and $f(x) = \cos x$ on $(\pi/4, \infty)$, f is continuous on $(-\infty, \pi/4) \cup (\pi/4, \infty)$. $\lim\limits_{x \to (\pi/4)^-} f(x) = \lim\limits_{x \to (\pi/4)^-} \sin x = \sin \frac{\pi}{4} = 1/\sqrt{2}$ since the sine function is continuous at $\pi/4$. Similarly, $\lim\limits_{x \to (\pi/4)^+} f(x) = \lim\limits_{x \to (\pi/4)^+} \cos x = 1/\sqrt{2}$ by continuity of the cosine function at $\pi/4$. Thus, $\lim\limits_{x \to (\pi/4)} f(x)$ exists and equals $1/\sqrt{2}$, which agrees with the value $f(\pi/4)$. Therefore, f is continuous at $\pi/4$, so f is continuous on $(-\infty, \infty)$.

43. $f(x) = \begin{cases} x^2 & \text{if } x < -1 \\ x & \text{if } -1 \le x < 1 \\ 1/x & \text{if } x \ge 1 \end{cases}$

f is continuous on $(-\infty, -1)$, $(-1, 1)$, and $(1, \infty)$, where it is a polynomial, a polynomial, and a rational function, respectively.

Now $\lim\limits_{x \to -1^-} f(x) = \lim\limits_{x \to -1^-} x^2 = 1$ and $\lim\limits_{x \to -1^+} f(x) = \lim\limits_{x \to -1^+} x = -1$, so f is discontinuous at -1. Since $f(-1) = -1$, f is continuous from the right at -1. Also, $\lim\limits_{x \to 1^-} f(x) = \lim\limits_{x \to 1^-} x = 1$ and $\lim\limits_{x \to 1^+} f(x) = \lim\limits_{x \to 1^+} \dfrac{1}{x} = 1 = f(1)$, so f is continuous at 1.

44. $f(x) = \begin{cases} 2^x & \text{if } x \le 1 \\ 3 - x & \text{if } 1 < x \le 4 \\ \sqrt{x} & \text{if } x > 4 \end{cases}$

f is continuous on $(-\infty, 1)$, $(1, 4)$, and $(4, \infty)$, where it is an exponential, a polynomial, and a root function, respectively.

Now $\lim\limits_{x \to 1^-} f(x) = \lim\limits_{x \to 1^-} 2^x = 2$ and $\lim\limits_{x \to 1^+} f(x) = \lim\limits_{x \to 1^+} (3 - x) = 2$. Since $f(1) = 2$ we have continuity at 1. Also, $\lim\limits_{x \to 4^-} f(x) = \lim\limits_{x \to 4^-} (3 - x) = -1 = f(4)$ and $\lim\limits_{x \to 4^+} f(x) = \lim\limits_{x \to 4^+} \sqrt{x} = 2$, so f is discontinuous at 4, but it is continuous from the left at 4.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. $f(x) = \begin{cases} x+2 & \text{if } x<0 \\ e^x & \text{if } 0 \le x \le 1 \\ 2-x & \text{if } x>1 \end{cases}$

f is continuous on $(-\infty, 0)$ and $(1, \infty)$ since on each of these intervals it is a polynomial; it is continuous on $(0, 1)$ since it is an exponential.

Now $\lim\limits_{x\to 0^-} f(x) = \lim\limits_{x\to 0^-} (x+2) = 2$ and $\lim\limits_{x\to 0^+} f(x) = \lim\limits_{x\to 0^+} e^x = 1$, so f is discontinuous at 0. Since $f(0) = 1$, f is continuous from the right at 0. Also $\lim\limits_{x\to 1^-} f(x) = \lim\limits_{x\to 1^-} e^x = e$ and $\lim\limits_{x\to 1^+} f(x) = \lim\limits_{x\to 1^+} (2-x) = 1$, so f is discontinuous at 1. Since $f(1) = e$, f is continuous from the left at 1.

46. By Theorem 5, each piece of F is continuous on its domain. We need to check for continuity at $r = R$. $\lim\limits_{r\to R^-} F(r) = \lim\limits_{r\to R^-} \dfrac{GMr}{R^3} = \dfrac{GM}{R^2}$ and $\lim\limits_{r\to R^+} F(r) = \lim\limits_{r\to R^+} \dfrac{GM}{r^2} = \dfrac{GM}{R^2}$, so $\lim\limits_{r\to R} F(r) = \dfrac{GM}{R^2}$. Since $F(R) = \dfrac{GM}{R^2}$, F is continuous at R. Therefore, F is a continuous function of r.

47. $f(x) = \begin{cases} cx^2+2x & \text{if } x<2 \\ x^3-cx & \text{if } x \ge 2 \end{cases}$

f is continuous on $(-\infty, 2)$ and $(2, \infty)$. Now $\lim\limits_{x\to 2^-} f(x) = \lim\limits_{x\to 2^-} \left(cx^2+2x\right) = 4c+4$ and $\lim\limits_{x\to 2^+} f(x) = \lim\limits_{x\to 2^+} \left(x^3-cx\right) = 8-2c$. So f is continuous $\Leftrightarrow$ $4c+4 = 8-2c$ $\Leftrightarrow$ $6c = 4$ $\Leftrightarrow$ $c = \frac{2}{3}$. Thus, for f to be continuous on $(-\infty, \infty)$, $c = \frac{2}{3}$.

48. $f(x) = \begin{cases} \dfrac{x^2-4}{x-2} & \text{if } x<2 \\ ax^2-bx+3 & \text{if } 2 \le x < 3 \\ 2x-a+b & \text{if } x \ge 3 \end{cases}$

At $x = 2$: $\lim\limits_{x\to 2^-} f(x) = \lim\limits_{x\to 2^-} \dfrac{x^2-4}{x-2} = \lim\limits_{x\to 2^-} \dfrac{(x+2)(x-2)}{x-2} = \lim\limits_{x\to 2^-} (x+2) = 2+2 = 4$

$\lim\limits_{x\to 2^+} f(x) = \lim\limits_{x\to 2^+} (ax^2-bx+3) = 4a-2b+3$

We must have $4a-2b+3 = 4$, or $\boldsymbol{4a-2b=1}$ **(1)**.

At $x = 3$: $\lim\limits_{x\to 3^-} f(x) = \lim\limits_{x\to 3^-} (ax^2-bx+3) = 9a-3b+3$

$\lim\limits_{x\to 3^+} f(x) = \lim\limits_{x\to 3^+} (2x-a+b) = 6-a+b$

We must have $9a-3b+3 = 6-a+b$, or $\boldsymbol{10a-4b=3}$ **(2)**.

Now solve the system of equations by adding -2 times equation **(1)** to equation **(2)**.

$$\begin{aligned} -8a+4b &= -2 \\ 10a-4b &= 3 \\ \hline 2a &= 1 \end{aligned}$$

So $a = \frac{1}{2}$. Substituting $\frac{1}{2}$ for a in **(1)** gives us $-2b = -1$, so $b = \frac{1}{2}$ as well. Thus, for f to be continuous on $(-\infty, \infty)$, $a = b = \frac{1}{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. If f and g are continuous and $g(2) = 6$, then $\lim\limits_{x\to 2}[3f(x) + f(x)\,g(x)] = 36 \quad \Rightarrow$

$3\lim\limits_{x\to 2} f(x) + \lim\limits_{x\to 2} f(x) \cdot \lim\limits_{x\to 2} g(x) = 36 \quad \Rightarrow \quad 3f(2) + f(2)\cdot 6 = 36 \quad \Rightarrow \quad 9f(2) = 36 \quad \Rightarrow \quad f(2) = 4.$

50. (a) $f(x) = \dfrac{1}{x}$ and $g(x) = \dfrac{1}{x^2}$, so $(f \circ g)(x) = f(g(x)) = f(1/x^2) = 1\,/(1/x^2) = x^2$.

(b) The domain of $f \circ g$ is the set of numbers x in the domain of g (all nonzero reals) such that $g(x)$ is in the domain of f (also all nonzero reals). Thus, the domain is $\left\{x \;\middle|\; x \neq 0 \text{ and } \dfrac{1}{x^2} \neq 0\right\} = \{x \mid x \neq 0\}$ or $(-\infty, 0) \cup (0, \infty)$. Since $f \circ g$ is the composite of two rational functions, it is continuous throughout its domain; that is, everywhere except $x = 0$.

51. (a) $f(x) = \dfrac{x^4 - 1}{x - 1} = \dfrac{(x^2+1)(x^2-1)}{x-1} = \dfrac{(x^2+1)(x+1)(x-1)}{x-1} = (x^2+1)(x+1) \quad [\text{or } x^3 + x^2 + x + 1]$

for $x \neq 1$. The discontinuity is removable and $g(x) = x^3 + x^2 + x + 1$ agrees with f for $x \neq 1$ and is continuous on $\mathbb{R}$.

(b) $f(x) = \dfrac{x^3 - x^2 - 2x}{x - 2} = \dfrac{x(x^2 - x - 2)}{x - 2} = \dfrac{x(x-2)(x+1)}{x-2} = x(x+1) \quad [\text{or } x^2 + x] \quad$ for $x \neq 2$. The discontinuity is removable and $g(x) = x^2 + x$ agrees with f for $x \neq 2$ and is continuous on $\mathbb{R}$.

(c) $\lim\limits_{x\to\pi^-} f(x) = \lim\limits_{x\to\pi^-} [\![\sin x]\!] = \lim\limits_{x\to\pi^-} 0 = 0$ and $\lim\limits_{x\to\pi^+} f(x) = \lim\limits_{x\to\pi^+} [\![\sin x]\!] = \lim\limits_{x\to\pi^+} (-1) = -1$, so $\lim\limits_{x\to\pi} f(x)$ does not exist. The discontinuity at $x = \pi$ is a jump discontinuity.

52.

f does not satisfy the conclusion of the Intermediate Value Theorem.

f does satisfy the conclusion of the Intermediate Value Theorem.

53. $f(x) = x^2 + 10\sin x$ is continuous on the interval $[31, 32]$, $f(31) \approx 957$, and $f(32) \approx 1030$. Since $957 < 1000 < 1030$, there is a number c in $(31, 32)$ such that $f(c) = 1000$ by the Intermediate Value Theorem. *Note:* There is also a number c in $(-32, -31)$ such that $f(c) = 1000$.

54. Suppose that $f(3) < 6$. By the Intermediate Value Theorem applied to the continuous function f on the closed interval $[2, 3]$, the fact that $f(2) = 8 > 6$ and $f(3) < 6$ implies that there is a number c in $(2, 3)$ such that $f(c) = 6$. This contradicts the fact that the only solutions of the equation $f(x) = 6$ are $x = 1$ and $x = 4$. Hence, our supposition that $f(3) < 6$ was incorrect. It follows that $f(3) \geq 6$. But $f(3) \neq 6$ because the only solutions of $f(x) = 6$ are $x = 1$ and $x = 4$. Therefore, $f(3) > 6$.

55. $f(x) = -x^3 + 4x + 1$ is continuous on the interval $[-1, 0]$, $f(-1) = -2$, and $f(0) = 1$. Since $-2 < 0 < 1$, there is a number c in $(-1, 0)$ such that $f(c) = 0$ by the Intermediate Value Theorem. Thus, there is a solution of the equation $-x^3 + 4x + 1 = 0$ in the interval $(-1, 0)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

56. The equation $\ln x = x - \sqrt{x}$ is equivalent to the equation $\ln x - x + \sqrt{x} = 0$. $f(x) = \ln x - x + \sqrt{x}$ is continuous on the interval $[2, 3]$, $f(2) = \ln 2 - 2 + \sqrt{2} \approx 0.107$, and $f(3) = \ln 3 - 3 + \sqrt{3} \approx -0.169$. Since $f(2) > 0 > f(3)$, there is a number c in $(2, 3)$ such that $f(c) = 0$ by the Intermediate Value Theorem. Thus, there is a solution of the equation $\ln x - x + \sqrt{x} = 0$, or $\ln x = x - \sqrt{x}$, in the interval $(2, 3)$.

57. The equation $e^x = 3 - 2x$ is equivalent to the equation $e^x + 2x - 3 = 0$. $f(x) = e^x + 2x - 3$ is continuous on the interval $[0, 1]$, $f(0) = -2$, and $f(1) = e - 1 \approx 1.72$. Since $-2 < 0 < e - 1$, there is a number c in $(0, 1)$ such that $f(c) = 0$ by the Intermediate Value Theorem. Thus, there is a solution of the equation $e^x + 2x - 3 = 0$, or $e^x = 3 - 2x$, in the interval $(0, 1)$.

58. The equation $\sin x = x^2 - x$ is equivalent to the equation $\sin x - x^2 + x = 0$. $f(x) = \sin x - x^2 + x$ is continuous on the interval $[1, 2]$, $f(1) = \sin 1 \approx 0.84$, and $f(2) = \sin 2 - 2 \approx -1.09$. Since $\sin 1 > 0 > \sin 2 - 2$, there is a number c in $(1, 2)$ such that $f(c) = 0$ by the Intermediate Value Theorem. Thus, there is a solution of the equation $\sin x - x^2 + x = 0$, or $\sin x = x^2 - x$, in the interval $(1, 2)$.

59. (a) $f(x) = \cos x - x^3$ is continuous on the interval $[0, 1]$, $f(0) = 1 > 0$, and $f(1) = \cos 1 - 1 \approx -0.46 < 0$. Since $1 > 0 > -0.46$, there is a number c in $(0, 1)$ such that $f(c) = 0$ by the Intermediate Value Theorem. Thus, there is a solution of the equation $\cos x - x^3 = 0$, or $\cos x = x^3$, in the interval $(0, 1)$.

(b) $f(0.86) \approx 0.016 > 0$ and $f(0.87) \approx -0.014 < 0$, so there is a solution between 0.86 and 0.87, that is, in the interval $(0.86, 0.87)$.

60. (a) $f(x) = \ln x - 3 + 2x$ is continuous on the interval $[1, 2]$, $f(1) = -1 < 0$, and $f(2) = \ln 2 + 1 \approx 1.7 > 0$. Since $-1 < 0 < 1.7$, there is a number c in $(1, 2)$ such that $f(c) = 0$ by the Intermediate Value Theorem. Thus, there is a solution of the equation $\ln x - 3 + 2x = 0$, or $\ln x = 3 - 2x$, in the interval $(1, 2)$.

(b) $f(1.34) \approx -0.03 < 0$ and $f(1.35) \approx 0.0001 > 0$, so there is a solution between 1.34 and 1.35, that is, in the interval $(1.34, 1.35)$.

61. (a) Let $f(x) = 100e^{-x/100} - 0.01x^2$. Then $f(0) = 100 > 0$ and $f(100) = 100e^{-1} - 100 \approx -63.2 < 0$. So by the Intermediate Value Theorem, there is a number c in $(0, 100)$ such that $f(c) = 0$. This implies that $100e^{-c/100} = 0.01c^2$.

(b) Using the intersect feature of the graphing device, we find that the solution of the equation is $x = 70.347$, correct to three decimal places.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

62. (a) Let $f(x) = \arctan x + x - 1$. Then $f(0) = -1 < 0$ and $f(1) = \frac{\pi}{4} > 0$. So by the Intermediate Value Theorem, there is a number c in $(0, 1)$ such that $f(c) = 0$. This implies that $\arctan c = 1 - c$.

(b) Using the intersect feature of the graphing device, we find that the solution of the equation is $x = 0.520$, correct to three decimal places.

63. Let $f(x) = \sin x^3$. Then f is continuous on $[1, 2]$ since f is the composite of the sine function and the cubing function, both of which are continuous on $\mathbb{R}$. The zeros of the sine are at $n\pi$, so we note that $0 < 1 < \pi < \frac{3}{2}\pi < 2\pi < 8 < 3\pi$, and that the pertinent cube roots are related by $1 < \sqrt[3]{\frac{3}{2}\pi}$ [call this value A] < 2. [By observation, we might notice that $x = \sqrt[3]{\pi}$ and $x = \sqrt[3]{2\pi}$ are zeros of f.]

Now $f(1) = \sin 1 > 0$, $f(A) = \sin \frac{3}{2}\pi = -1 < 0$, and $f(2) = \sin 8 > 0$. Applying the Intermediate Value Theorem on $[1, A]$ and then on $[A, 2]$, we see there are numbers c and d in $(1, A)$ and $(A, 2)$ such that $f(c) = f(d) = 0$. Thus, f has at least two x-intercepts in $(1, 2)$.

64. Let $f(x) = x^2 - 3 + 1/x$. Then f is continuous on $(0, 2]$ since f is a rational function whose domain is $(0, \infty)$. By inspection, we see that $f\left(\frac{1}{4}\right) = \frac{17}{16} > 0$, $f(1) = -1 < 0$, and $f(2) = \frac{3}{2} > 0$. Appling the Intermediate Value Theorem on $\left[\frac{1}{4}, 1\right]$ and then on $[1, 2]$, we see there are numbers c and d in $\left(\frac{1}{4}, 1\right)$ and $(1, 2)$ such that $f(c) = f(d) = 0$. Thus, f has at least two x-intercepts in $(0, 2)$.

65. ($\Rightarrow$) If f is continuous at a, then by Theorem 8 with $g(h) = a + h$, we have

$$\lim_{h\to 0} f(a+h) = f\Big(\lim_{h\to 0}(a+h)\Big) = f(a).$$

($\Leftarrow$) Let $\varepsilon > 0$. Since $\lim_{h\to 0} f(a+h) = f(a)$, there exists $\delta > 0$ such that $0 < |h| < \delta \quad \Rightarrow$ $|f(a+h) - f(a)| < \varepsilon$. So if $0 < |x - a| < \delta$, then $|f(x) - f(a)| = |f(a + (x - a)) - f(a)| < \varepsilon$. Thus, $\lim_{x\to a} f(x) = f(a)$ and so f is continuous at a.

66.
$$\begin{aligned}\lim_{h\to 0}\sin(a+h) &= \lim_{h\to 0}(\sin a\cos h + \cos a\sin h) = \lim_{h\to 0}(\sin a\cos h) + \lim_{h\to 0}(\cos a\sin h)\\ &= \Big(\lim_{h\to 0}\sin a\Big)\Big(\lim_{h\to 0}\cos h\Big) + \Big(\lim_{h\to 0}\cos a\Big)\Big(\lim_{h\to 0}\sin h\Big) = (\sin a)(1) + (\cos a)(0) = \sin a\end{aligned}$$

67. As in the previous exercise, we must show that $\lim_{h\to 0}\cos(a+h) = \cos a$ to prove that the cosine function is continuous.

$$\begin{aligned}\lim_{h\to 0}\cos(a+h) &= \lim_{h\to 0}(\cos a\cos h - \sin a\sin h) = \lim_{h\to 0}(\cos a\cos h) - \lim_{h\to 0}(\sin a\sin h)\\ &= \Big(\lim_{h\to 0}\cos a\Big)\Big(\lim_{h\to 0}\cos h\Big) - \Big(\lim_{h\to 0}\sin a\Big)\Big(\lim_{h\to 0}\sin h\Big) = (\cos a)(1) - (\sin a)(0) = \cos a\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

68. (a) Since f is continuous at a, $\lim_{x\to a} f(x) = f(a)$. Thus, using the Constant Multiple Law of Limits, we have

$\lim_{x\to a}(cf)(x) = \lim_{x\to a} cf(x) = c\lim_{x\to a} f(x) = cf(a) = (cf)(a)$. Therefore, cf is continuous at a.

(b) Since f and g are continuous at a, $\lim_{x\to a} f(x) = f(a)$ and $\lim_{x\to a} g(x) = g(a)$. Since $g(a) \neq 0$, we can use the Quotient Law of Limits: $\lim_{x\to a}\left(\dfrac{f}{g}\right)(x) = \lim_{x\to a}\dfrac{f(x)}{g(x)} = \dfrac{\lim_{x\to a} f(x)}{\lim_{x\to a} g(x)} = \dfrac{f(a)}{g(a)} = \left(\dfrac{f}{g}\right)(a)$. Thus, $\dfrac{f}{g}$ is continuous at a.

69. *Proof of Law 6*: Let n be a positive integer. By Theorem 8 with $f(x) = x^n$, we have

$$\lim_{x\to a}[g(x)]^n = \lim_{x\to a} f(g(x)) = f\left(\lim_{x\to a} g(x)\right) = \left[\lim_{x\to a} g(x)\right]^n$$

Proof of Law 7: Let n be a positive integer. By Theorem 8 with $f(x) = \sqrt[n]{x}$, we have

$$\lim_{x\to a}\sqrt[n]{g(x)} = \lim_{x\to a} f(g(x)) = f\left(\lim_{x\to a} g(x)\right) = \sqrt[n]{\lim_{x\to a} g(x)}$$

70. If there is such a number, it satisfies the equation $x^3 + 1 = x \quad \Leftrightarrow \quad x^3 - x + 1 = 0$. Let the left-hand side of this equation be called $f(x)$. Now $f(-2) = -5 < 0$, and $f(-1) = 1 > 0$. Note also that $f(x)$ is a polynomial, and thus continuous. So by the Intermediate Value Theorem, there is a number c between -2 and -1 such that $f(c) = 0$, so that $c = c^3 + 1$.

71. $f(x) = \begin{cases} 0 & \text{if } x \text{ is rational} \\ 1 & \text{if } x \text{ is irrational} \end{cases}$ is continuous nowhere. For, given any number a and any $\delta > 0$, the interval $(a-\delta, a+\delta)$ contains both infinitely many rational and infinitely many irrational numbers. Since $f(a) = 0$ or 1, there are infinitely many numbers x with $0 < |x-a| < \delta$ and $|f(x) - f(a)| = 1$. Thus, $\lim_{x\to a} f(x) \neq f(a)$. [In fact, $\lim_{x\to a} f(x)$ does not even exist.]

72. $g(x) = \begin{cases} 0 & \text{if } x \text{ is rational} \\ x & \text{if } x \text{ is irrational} \end{cases}$ is continuous at 0. To see why, note that $-|x| \leq g(x) \leq |x|$, so by the Squeeze Theorem $\lim_{x\to 0} g(x) = 0 = g(0)$. But g is continuous nowhere else. For if $a \neq 0$ and $\delta > 0$, the interval $(a-\delta, a+\delta)$ contains both infinitely many rational and infinitely many irrational numbers. Since $g(a) = 0$ or a, there are infinitely many numbers x with $0 < |x-a| < \delta$ and $|g(x) - g(a)| > |a|/2$. Thus, $\lim_{x\to a} g(x) \neq g(a)$.

73. $f(x) = x^4\sin(1/x)$ is continuous on $(-\infty, 0) \cup (0, \infty)$ since it is the product of a polynomial and a composite of a trigonometric function and a rational function. Now since $-1 \leq \sin(1/x) \leq 1$, we have $-x^4 \leq x^4\sin(1/x) \leq x^4$. Because $\lim_{x\to 0}(-x^4) = 0$ and $\lim_{x\to 0} x^4 = 0$, the Squeeze Theorem gives us $\lim_{x\to 0}(x^4\sin(1/x)) = 0$, which equals $f(0)$. Thus, f is continuous at 0 and, hence, on $(-\infty, \infty)$.

74. $\dfrac{a}{x^3+2x^2-1} + \dfrac{b}{x^3+x-2} = 0 \quad \Rightarrow \quad a(x^3+x-2) + b(x^3+2x^2-1) = 0$. Let $p(x)$ denote the left side of the last equation. Since p is continuous on $[-1, 1]$, $p(-1) = -4a < 0$, and $p(1) = 2b > 0$, there exists a c in $(-1, 1)$ such that

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$p(c) = 0$ by the Intermediate Value Theorem. Note that the only solution of either denominator that is in $(-1, 1)$ is $(-1 + \sqrt{5})/2 = r$, but $p(r) = (3\sqrt{5} - 9)a/2 \neq 0$. Thus, c is not a solution of either denominator, so $p(c) = 0 \quad \Rightarrow$ $x = c$ is a solution of the given equation.

75. Define $u(t)$ to be the monk's distance from the monastery, as a function of time t (in hours), on the first day, and define $d(t)$ to be his distance from the monastery, as a function of time, on the second day. Let D be the distance from the monastery to the top of the mountain. From the given information we know that $u(0) = 0$, $u(12) = D$, $d(0) = D$ and $d(12) = 0$. Now consider the function $u - d$, which is clearly continuous. We calculate that $(u - d)(0) = -D$ and $(u - d)(12) = D$. So by the Intermediate Value Theorem, there must be some time t_0 between 0 and 12 such that $(u - d)(t_0) = 0 \quad \Leftrightarrow$ $u(t_0) = d(t_0)$. So at time t_0 after 7:00 AM, the monk will be at the same place on both days.

76. (a) $\lim\limits_{x \to 0^+} F(x) = 0$ and $\lim\limits_{x \to 0^-} F(x) = 0$, so $\lim\limits_{x \to 0} F(x) = 0$, which is $F(0)$, and hence F is continuous at $x = a$ if $a = 0$. For $a > 0$, $\lim\limits_{x \to a} F(x) = \lim\limits_{x \to a} x = a = F(a)$. For $a < 0$, $\lim\limits_{x \to a} F(x) = \lim\limits_{x \to a} (-x) = -a = F(a)$. Thus, F is continuous at $x = a$; that is, continuous everywhere.

(b) Assume that f is continuous on the interval I. Then for $a \in I$, $\lim\limits_{x \to a} |f(x)| = \left| \lim\limits_{x \to a} f(x) \right| = |f(a)|$ by Theorem 8. (If a is an endpoint of I, use the appropriate one-sided limit.) So $|f|$ is continuous on I.

(c) No, the converse is false. For example, the function $f(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ -1 & \text{if } x < 0 \end{cases}$ is not continuous at $x = 0$, but $|f(x)| = 1$ is continuous on $\mathbb{R}$.

2.6 Limits at Infinity; Horizontal Asymptotes

1. (a) As x becomes large, the values of $f(x)$ approach 5.

(b) As x becomes large negative, the values of $f(x)$ approach 3.

2. (a) The graph of a function can intersect a vertical asymptote in the sense that it can meet but not cross it.

The graph of a function can intersect a horizontal asymptote. It can even intersect its horizontal asymptote an infinite number of times.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The graph of a function can have 0, 1, or 2 horizontal asymptotes. Representative examples are shown.

No horizontal asymptote

One horizontal asymptote

Two horizontal asymptotes

3. (a) $\lim_{x\to\infty} f(x) = -2$ (b) $\lim_{x\to-\infty} f(x) = 2$ (c) $\lim_{x\to 1} f(x) = \infty$

(d) $\lim_{x\to 3} f(x) = -\infty$ (e) Vertical: $x = 1$, $x = 3$; horizontal: $y = -2$, $y = 2$

4. (a) $\lim_{x\to\infty} g(x) = 2$ (b) $\lim_{x\to-\infty} g(x) = -1$ (c) $\lim_{x\to 0} g(x) = -\infty$

(d) $\lim_{x\to 2^-} g(x) = -\infty$ (e) $\lim_{x\to 2^+} g(x) = \infty$ (f) Vertical: $x = 0$, $x = 2$; horizontal: $y = -1$, $y = 2$

5. $f(2) = 4$, $f(-2) = -4$, $\lim_{x\to-\infty} f(x) = 0$, $\lim_{x\to\infty} f(x) = 2$

6. $f(0) = 0$, $\lim_{x\to 1^-} f(x) = \infty$, $\lim_{x\to 1^+} f(x) = -\infty$, $\lim_{x\to-\infty} f(x) = -2$, $\lim_{x\to\infty} f(x) = -2$

7. $\lim_{x\to 0} f(x) = \infty$, $\lim_{x\to 3^-} f(x) = -\infty$, $\lim_{x\to 3^+} f(x) = \infty$, $\lim_{x\to-\infty} f(x) = 1$, $\lim_{x\to\infty} f(x) = -1$

8. $\lim_{x\to-\infty} f(x) = -\infty$, $\lim_{x\to -2^-} f(x) = \infty$, $\lim_{x\to -2^+} f(x) = -\infty$, $\lim_{x\to 2} f(x) = \infty$, $\lim_{x\to\infty} f(x) = \infty$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $f(0) = 0$, $\lim\limits_{x \to 1} f(x) = -\infty$, $\lim\limits_{x \to \infty} f(x) = -\infty$,

f is odd

10. $\lim\limits_{x \to -\infty} f(x) = -1$, $\lim\limits_{x \to 0^-} f(x) = \infty$,

$\lim\limits_{x \to 0^+} f(x) = -\infty$, $\lim\limits_{x \to 3^-} f(x) = 1$, $f(3) = 4$,

$\lim\limits_{x \to 3^+} f(x) = 4$, $\lim\limits_{x \to \infty} f(x) = 1$

11. If $f(x) = x^2/2^x$, then a calculator gives $f(0) = 0$, $f(1) = 0.5$, $f(2) = 1$, $f(3) = 1.125$, $f(4) = 1$, $f(5) = 0.78125$, $f(6) = 0.5625$, $f(7) = 0.3828125$, $f(8) = 0.25$, $f(9) = 0.158203125$, $f(10) = 0.09765625$, $f(20) \approx 0.00038147$, $f(50) \approx 2.2204 \times 10^{-12}$, $f(100) \approx 7.8886 \times 10^{-27}$. It appears that $\lim\limits_{x \to \infty} \left(x^2/2^x\right) = 0$.

12. (a) From a graph of $f(x) = (1 - 2/x)^x$ in a window of $[0, 10{,}000]$ by $[0, 0.2]$, we estimate that $\lim\limits_{x \to \infty} f(x) = 0.14$ (to two decimal places).

(b)

x	$f(x)$
10,000	0.135 308
100,000	0.135 333
1,000,000	0.135 335

From the table, we estimate that $\lim\limits_{x \to \infty} f(x) = 0.1353$ (to four decimal places).

13.
$$\begin{aligned}
\lim_{x \to \infty} \frac{2x^2 - 7}{5x^2 + x - 3} &= \lim_{x \to \infty} \frac{(2x^2 - 7)/x^2}{(5x^2 + x - 3)/x^2} && \text{[Divide both the numerator and denominator by } x^2 \text{ (the highest power of } x \text{ that appears in the denominator)]} \\
&= \frac{\lim\limits_{x \to \infty} (2 - 7/x^2)}{\lim\limits_{x \to \infty} (5 + 1/x - 3/x^2)} && \text{[Limit Law 5]} \\
&= \frac{\lim\limits_{x \to \infty} 2 - \lim\limits_{x \to \infty} (7/x^2)}{\lim\limits_{x \to \infty} 5 + \lim\limits_{x \to \infty} (1/x) - \lim\limits_{x \to \infty} (3/x^2)} && \text{[Limit Laws 1 and 2]} \\
&= \frac{2 - 7 \lim\limits_{x \to \infty} (1/x^2)}{5 + \lim\limits_{x \to \infty} (1/x) - 3 \lim\limits_{x \to \infty} (1/x^2)} && \text{[Limit Laws 8 and 3]} \\
&= \frac{2 - 7(0)}{5 + 0 + 3(0)} && \text{[Theorem 5]} \\
&= \frac{2}{5}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $$\begin{aligned}
\lim_{x\to\infty}\sqrt{\frac{9x^3+8x-4}{3-5x+x^3}} &= \sqrt{\lim_{x\to\infty}\frac{9x^3+8x-4}{3-5x+x^3}} && \text{[Limit Law 7]}\\
&= \sqrt{\lim_{x\to\infty}\frac{9+8/x^2-4/x^3}{3/x^3-5/x^2+1}} && \text{[Divide by } x^3\text{]}\\
&= \sqrt{\frac{\lim_{x\to\infty}(9+8/x^2-4/x^3)}{\lim_{x\to\infty}(3/x^3-5/x^2+1)}} && \text{[Limit Law 5]}\\
&= \sqrt{\frac{\lim_{x\to\infty}9+\lim_{x\to\infty}(8/x^2)-\lim_{x\to\infty}(4/x^3)}{\lim_{x\to\infty}(3/x^3)-\lim_{x\to\infty}(5/x^2)+\lim_{x\to\infty}1}} && \text{[Limit Laws 1 and 2]}\\
&= \sqrt{\frac{9+8\lim_{x\to\infty}(1/x^2)-4\lim_{x\to\infty}(1/x^3)}{3\lim_{x\to\infty}(1/x^3)-5\lim_{x\to\infty}(1/x^2)+1}} && \text{[Limit Laws 8 and 3]}\\
&= \sqrt{\frac{9+8(0)-4(0)}{3(0)-5(0)+1}} && \text{[Theorem 5]}\\
&= \sqrt{\frac{9}{1}} = \sqrt{9} = 3
\end{aligned}$$

15. $$\lim_{x\to\infty}\frac{4x+3}{5x-1} = \lim_{x\to\infty}\frac{(4x+3)/x}{(5x-1)/x} = \lim_{x\to\infty}\frac{4+3/x}{5-1/x} = \frac{\lim_{x\to\infty}4+3\lim_{x\to\infty}(1/x)}{\lim_{x\to\infty}5-\lim_{x\to\infty}(1/x)} = \frac{4+3(0)}{5-0} = \frac{4}{5}$$

16. $$\lim_{x\to\infty}\frac{-2}{3x+7} = \lim_{x\to\infty}\frac{-2/x}{(3x+7)/x} = \lim_{x\to\infty}\frac{-2/x}{3+7/x} = \frac{-2\lim_{x\to\infty}(1/x)}{\lim_{x\to\infty}3+7\lim_{x\to\infty}(1/x)} = \frac{0}{3+0} = 0$$

17. $$\begin{aligned}
\lim_{t\to-\infty}\frac{3t^2+t}{t^3-4t+1} &= \lim_{t\to-\infty}\frac{(3t^2+t)/t^3}{(t^3-4t+1)/t^3} = \lim_{t\to-\infty}\frac{3/t+1/t^2}{1-4/t^2+1/t^3}\\
&= \frac{3\lim_{t\to-\infty}(1/t)+\lim_{t\to-\infty}(1/t^2)}{\lim_{t\to-\infty}1-4\lim_{t\to-\infty}(1/t^2)+\lim_{t\to-\infty}(1/t^3)} = \frac{3(0)+0}{1-4(0)+0} = 0
\end{aligned}$$

18. $$\lim_{t\to-\infty}\frac{6t^2+t-5}{9-2t^2} = \lim_{t\to-\infty}\frac{(6t^2+t-5)/t^2}{(9-2t^2)/t^2} = \lim_{t\to-\infty}\frac{6+1/t-5/t^2}{9/t^2-2} = \frac{6+0-0}{0-2} = -3$$

19. $$\lim_{r\to\infty}\frac{r-r^3}{2-r^2+3r^3} = \lim_{r\to\infty}\frac{(r-r^3)/r^3}{(2-r^2+3r^3)/r^3} = \lim_{r\to\infty}\frac{1/r^2-1}{2/r^3-1/r+3} = \frac{0-1}{0-0+3} = -\frac{1}{3}$$

20. $$\lim_{x\to\infty}\frac{3x^3-8x+2}{4x^3-5x^2-2} = \lim_{x\to\infty}\frac{(3x^3-8x+2)/x^3}{(4x^3-5x^2-2)/x^3} = \lim_{x\to\infty}\frac{3-8/x^2+2/x^3}{4-5/x-2/x^3} = \frac{3-0+0}{4-0-0} = \frac{3}{4}$$

21. $$\lim_{x\to\infty}\frac{4-\sqrt{x}}{2+\sqrt{x}} = \lim_{x\to\infty}\frac{(4-\sqrt{x})/\sqrt{x}}{(2+\sqrt{x})/\sqrt{x}} = \lim_{x\to\infty}\frac{4/\sqrt{x}-1}{2/\sqrt{x}+1} = \frac{0-1}{0+1} = -1$$

22. $$\begin{aligned}
\lim_{u\to-\infty}\frac{(u^2+1)(2u^2-1)}{(u^2+2)^2} &= \lim_{u\to-\infty}\frac{\left[(u^2+1)(2u^2-1)\right]/u^4}{(u^2+2)^2/u^4} = \lim_{u\to-\infty}\frac{\left[(u^2+1)/u^2\right]\left[(2u^2-1)/u^2\right]}{(u^4+4u^2+4)/u^4}\\
&= \lim_{u\to-\infty}\frac{(1+1/u^2)(2-1/u^2)}{(1+4/u^2+4/u^4)} = \frac{(1+0)(2-0)}{1+0+0} = 2
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. $$\lim_{x\to\infty}\frac{\sqrt{x+3x^2}}{4x-1}=\lim_{x\to\infty}\frac{\sqrt{x+3x^2}/x}{(4x-1)/x}=\frac{\lim\limits_{x\to\infty}\sqrt{(x+3x^2)/x^2}}{\lim\limits_{x\to\infty}(4-1/x)}\qquad[\text{since } x=\sqrt{x^2}\text{ for } x>0]$$

$$=\frac{\lim\limits_{x\to\infty}\sqrt{1/x+3}}{\lim\limits_{x\to\infty}4-\lim\limits_{x\to\infty}(1/x)}=\frac{\sqrt{\lim\limits_{x\to\infty}(1/x)+\lim\limits_{x\to\infty}3}}{4-0}=\frac{\sqrt{0+3}}{4}=\frac{\sqrt{3}}{4}$$

24. $$\lim_{t\to\infty}\frac{t+3}{\sqrt{2t^2-1}}=\lim_{t\to\infty}\frac{(t+3)/t}{\sqrt{2t^2-1}/t}=\lim_{t\to\infty}\frac{1+3/t}{\sqrt{2-1/t^2}}\qquad\left[\text{since } t=\sqrt{t^2}\text{ for } t>0\right]$$

$$=\frac{\lim\limits_{t\to\infty}(1+3/t)}{\lim\limits_{t\to\infty}\sqrt{2-1/t^2}}=\frac{\lim\limits_{t\to\infty}1+\lim\limits_{t\to\infty}(3/t)}{\sqrt{\lim\limits_{t\to\infty}2-\lim\limits_{t\to\infty}(1/t^2)}}=\frac{1+0}{\sqrt{2-0}}=\frac{1}{\sqrt{2}},\text{ or }\frac{\sqrt{2}}{2}$$

25. $$\lim_{x\to\infty}\frac{\sqrt{1+4x^6}}{2-x^3}=\lim_{x\to\infty}\frac{\sqrt{1+4x^6}/x^3}{(2-x^3)/x^3}=\frac{\lim\limits_{x\to\infty}\sqrt{(1+4x^6)/x^6}}{\lim\limits_{x\to\infty}(2/x^3-1)}\qquad\left[\text{since } x^3=\sqrt{x^6}\text{ for } x>0\right]$$

$$=\frac{\lim\limits_{x\to\infty}\sqrt{1/x^6+4}}{\lim\limits_{x\to\infty}(2/x^3)-\lim\limits_{x\to\infty}1}=\frac{\sqrt{\lim\limits_{x\to\infty}(1/x^6)+\lim\limits_{x\to\infty}4}}{0-1}$$

$$=\frac{\sqrt{0+4}}{-1}=\frac{2}{-1}=-2$$

26. $$\lim_{x\to-\infty}\frac{\sqrt{1+4x^6}}{2-x^3}=\lim_{x\to-\infty}\frac{\sqrt{1+4x^6}/x^3}{(2-x^3)/x^3}=\frac{\lim\limits_{x\to-\infty}-\sqrt{(1+4x^6)/x^6}}{\lim\limits_{x\to-\infty}(2/x^3-1)}\qquad\left[\text{since } x^3=-\sqrt{x^6}\text{ for } x<0\right]$$

$$=\frac{\lim\limits_{x\to-\infty}-\sqrt{1/x^6+4}}{2\lim\limits_{x\to-\infty}(1/x^3)-\lim\limits_{x\to-\infty}1}=\frac{-\sqrt{\lim\limits_{x\to-\infty}(1/x^6)+\lim\limits_{x\to-\infty}4}}{2(0)-1}$$

$$=\frac{-\sqrt{0+4}}{-1}=\frac{-2}{-1}=2$$

27. $$\lim_{x\to-\infty}\frac{2x^5-x}{x^4+3}=\lim_{x\to-\infty}\frac{\left(2x^5-x\right)/x^4}{\left(x^4+3\right)/x^4}=\lim_{x\to-\infty}\frac{2x-1/x^3}{1+3/x^4}$$

$$=-\infty\text{ since } 2x-1/x^3\to-\infty\text{ and } 1+3/x^4\to1\text{ as } x\to-\infty$$

28. $$\lim_{q\to\infty}\frac{q^3+6q-4}{4q^2-3q+3}=\lim_{q\to\infty}\frac{\left(q^3+6q-4\right)/q^2}{\left(4q^2-3q+3\right)/q^2}=\lim_{q\to\infty}\frac{q+6/q-4/q^2}{4-3/q+3/q^2}$$

$$=\infty\text{ since } q+6/q-4/q^2\to\infty\text{ and } 4-3/q+3/q+3/q^2\to4\text{ as } q\to\infty$$

29. $$\lim_{t\to\infty}\left(\sqrt{25t^2+2}-5t\right)=\lim_{t\to\infty}\left(\sqrt{25t^2+2}-5t\right)\left(\frac{\sqrt{25t^2+2}+5t}{\sqrt{25t^2+2}+5t}\right)=\lim_{t\to\infty}\frac{\left(25t^2+2\right)-(5t)^2}{\sqrt{25t^2+2}+5t}$$

$$=\lim_{t\to\infty}\frac{2}{\sqrt{25t^2+2}+5t}=\lim_{t\to\infty}\frac{2/t}{\left(\sqrt{25t^2+2}+5t\right)/t}$$

$$=\lim_{t\to\infty}\frac{2/t}{\sqrt{25+2/t^2}+5}\qquad[\text{since } t=\sqrt{t^2}\text{ for } t>0]$$

$$=\frac{0}{\sqrt{25+0}+5}=0$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $$\lim_{x\to-\infty}\left(\sqrt{4x^2+3x}+2x\right)=\lim_{x\to-\infty}\left(\sqrt{4x^2+3x}+2x\right)\left[\frac{\sqrt{4x^2+3x}-2x}{\sqrt{4x^2+3x}-2x}\right]=\lim_{x\to-\infty}\frac{\left(4x^2+3x\right)-(2x)^2}{\sqrt{4x^2+3x}-2x}$$
$$=\lim_{x\to-\infty}\frac{3x}{\sqrt{4x^2+3x}-2x}=\lim_{x\to-\infty}\frac{3x/x}{\left(\sqrt{4x^2+3x}-2x\right)/x}$$
$$=\lim_{x\to-\infty}\frac{3}{-\sqrt{4+3/x}-2}\qquad\left[\text{since } x=-\sqrt{x^2} \text{ for } x<0\right]$$
$$=\frac{3}{-\sqrt{4+0}-2}=-\frac{3}{4}$$

31. $$\lim_{x\to\infty}\left(\sqrt{x^2+ax}-\sqrt{x^2+bx}\right)=\lim_{x\to\infty}\frac{\left(\sqrt{x^2+ax}-\sqrt{x^2+bx}\right)\left(\sqrt{x^2+ax}+\sqrt{x^2+bx}\right)}{\sqrt{x^2+ax}+\sqrt{x^2+bx}}$$
$$=\lim_{x\to\infty}\frac{(x^2+ax)-(x^2+bx)}{\sqrt{x^2+ax}+\sqrt{x^2+bx}}=\lim_{x\to\infty}\frac{[(a-b)x]/x}{\left(\sqrt{x^2+ax}+\sqrt{x^2+bx}\right)/\sqrt{x^2}}$$
$$=\lim_{x\to\infty}\frac{a-b}{\sqrt{1+a/x}+\sqrt{1+b/x}}=\frac{a-b}{\sqrt{1+0}+\sqrt{1+0}}=\frac{a-b}{2}$$

32. $$\lim_{x\to\infty}\left(x-\sqrt{x}\right)=\lim_{x\to\infty}\left(x-\sqrt{x}\right)\left[\frac{x+\sqrt{x}}{x+\sqrt{x}}\right]=\lim_{x\to\infty}\frac{x^2-\left(\sqrt{x}\right)^2}{x+\sqrt{x}}=\lim_{x\to\infty}\frac{x^2-x}{x+\sqrt{x}}=\lim_{x\to\infty}\frac{(x^2-x)/x}{\left(x+\sqrt{x}\right)/x}$$
$$=\lim_{x\to\infty}\frac{x-1}{1+1/\sqrt{x}}=\infty\quad\text{since } x-1\to\infty \text{ and } 1+1/\sqrt{x}\to 1 \text{ as } x\to\infty$$

33. $\lim_{x\to-\infty}(x^2+2x^7)=\lim_{x\to-\infty}x^7\left(\frac{1}{x^5}+2\right)$ [factor out the largest power of x] $=-\infty$ because $x^7\to-\infty$ and $1/x^5+2\to 2$ as $x\to-\infty$.

Or: $\lim_{x\to-\infty}\left(x^2+2x^7\right)=\lim_{x\to-\infty}x^2\left(1+2x^5\right)=-\infty$.

34. $\lim_{x\to\infty}(e^{-x}+2\cos 3x)$ does not exist. $\lim_{x\to\infty}e^{-x}=0$, but $\lim_{x\to\infty}(2\cos 3x)$ does not exist because the values of $2\cos 3x$ oscillate between the values of -2 and 2 infinitely often, so the given limit does not exist.

35. Since $-1\le\cos x\le 1$ and $e^{-2x}>0$, we have $-e^{-2x}\le e^{-2x}\cos x\le e^{-2x}$. We know that $\lim_{x\to\infty}(-e^{-2x})=0$ and $\lim_{x\to\infty}\left(e^{-2x}\right)=0$, so by the Squeeze Theorem, $\lim_{x\to\infty}(e^{-2x}\cos x)=0$.

36. Since $0\le\sin^2 x\le 1$, we have $0\le\frac{\sin^2 x}{x^2+1}\le\frac{1}{x^2+1}$. We know that $\lim_{x\to\infty}0=0$ and $\lim_{x\to\infty}\frac{1}{x^2+1}=0$, so by the Squeeze Theorem, $\lim_{x\to\infty}\frac{\sin^2 x}{x^2+1}=0$.

37. $$\lim_{x\to\infty}\frac{1-e^x}{1+2e^x}=\lim_{x\to\infty}\frac{(1-e^x)/e^x}{(1+2e^x)/e^x}=\lim_{x\to\infty}\frac{1/e^x-1}{1/e^x+2}=\frac{0-1}{0+2}=-\frac{1}{2}$$

38. Divide numerator and denominator by e^{3x}: $\lim_{x\to\infty}\frac{e^{3x}-e^{-3x}}{e^{3x}+e^{-3x}}=\lim_{x\to\infty}\frac{1-e^{-6x}}{1+e^{-6x}}=\frac{1-0}{1+0}=1$

39. $\lim_{x\to(\pi/2)^+}e^{\sec x}=0$ since $\sec x\to-\infty$ as $x\to(\pi/2)^+$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. Let $t = \ln x$. As $x \to 0^+$, $t \to -\infty$. $\lim\limits_{x\to 0^+} \tan^{-1}(\ln x) = \lim\limits_{t\to -\infty} \tan^{-1} t = -\frac{\pi}{2}$ by (4).

41. $\lim\limits_{x\to\infty} [\ln(1+x^2) - \ln(1+x)] = \lim\limits_{x\to\infty} \ln \dfrac{1+x^2}{1+x} = \ln\left(\lim\limits_{x\to\infty} \dfrac{1+x^2}{1+x}\right) = \ln\left(\lim\limits_{x\to\infty} \dfrac{\frac{1}{x}+x}{\frac{1}{x}+1}\right) = \infty$, since the limit in parentheses is ∞.

42. $\lim\limits_{x\to\infty} [\ln(2+x) - \ln(1+x)] = \lim\limits_{x\to\infty} \ln\left(\dfrac{2+x}{1+x}\right) = \lim\limits_{x\to\infty} \ln\left(\dfrac{2/x+1}{1/x+1}\right) = \ln\dfrac{1}{1} = \ln 1 = 0$

43. (a) (i) $\lim\limits_{x\to 0^+} f(x) = \lim\limits_{x\to 0^+} \dfrac{x}{\ln x} = 0$ since $x \to 0^+$ and $\ln x \to -\infty$ as $x \to 0^+$.

(ii) $\lim\limits_{x\to 1^-} f(x) = \lim\limits_{x\to 1^-} \dfrac{x}{\ln x} = -\infty$ since $x \to 1$ and $\ln x \to 0^-$ as $x \to 1^-$.

(iii) $\lim\limits_{x\to 1^+} f(x) = \lim\limits_{x\to 1^+} \dfrac{x}{\ln x} = \infty$ since $x \to 1$ and $\ln x \to 0^+$ as $x \to 1^+$.

(b)

x	$f(x)$
10,000	1085.7
100,000	8685.9
1,000,000	72,382.4

It appears that $\lim\limits_{x\to\infty} f(x) = \infty$.

(c)

44. (a) (i) $\lim\limits_{x\to\infty} f(x) = \lim\limits_{x\to\infty} \left(\dfrac{2}{x} - \dfrac{1}{\ln x}\right) = 0$ since $\dfrac{2}{x} \to 0$ and $\dfrac{1}{\ln x} \to 0$ as $x \to \infty$.

(ii) $\lim\limits_{x\to 0^+} f(x) = \lim\limits_{x\to 0^+} \left(\dfrac{2}{x} - \dfrac{1}{\ln x}\right) = \infty$ since $\dfrac{2}{x} \to \infty$ and $\dfrac{1}{\ln x} \to 0$ as $x \to 0^+$.

(iii) $\lim\limits_{x\to 1^-} f(x) = \lim\limits_{x\to 1^-} \left(\dfrac{2}{x} - \dfrac{1}{\ln x}\right) = \infty$ since $\dfrac{2}{x} \to 2$ and $\dfrac{1}{\ln x} \to -\infty$ as $x \to 1^-$.

(iv) $\lim\limits_{x\to 1^+} f(x) = \lim\limits_{x\to 1^+} \left(\dfrac{2}{x} - \dfrac{1}{\ln x}\right) = -\infty$ since $\dfrac{2}{x} \to 2$ and $\dfrac{1}{\ln x} \to \infty$ as $x \to 1^+$.

(b)

45. (a)

From the graph of $f(x) = \sqrt{x^2+x+1} + x$, we estimate the value of $\lim\limits_{x\to -\infty} f(x)$ to be -0.5.

(b)

x	$f(x)$
−10,000	−0.499 962 5
−100,000	−0.499 996 2
−1,000,000	−0.499 999 6

From the table, we estimate the limit to be -0.5.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\lim\limits_{x\to-\infty}\left(\sqrt{x^2+x+1}+x\right) = \lim\limits_{x\to-\infty}\left(\sqrt{x^2+x+1}+x\right)\left[\dfrac{\sqrt{x^2+x+1}-x}{\sqrt{x^2+x+1}-x}\right] = \lim\limits_{x\to-\infty}\dfrac{\left(x^2+x+1\right)-x^2}{\sqrt{x^2+x+1}-x}$

$$= \lim_{x\to-\infty}\frac{(x+1)(1/x)}{\left(\sqrt{x^2+x+1}-x\right)(1/x)} = \lim_{x\to-\infty}\frac{1+(1/x)}{-\sqrt{1+(1/x)+(1/x^2)}-1}$$

$$= \frac{1+0}{-\sqrt{1+0+0}-1} = -\frac{1}{2}$$

Note that for $x<0$, we have $\sqrt{x^2}=|x|=-x$, so when we divide the radical by x, with $x<0$, we get $\dfrac{1}{x}\sqrt{x^2+x+1} = -\dfrac{1}{\sqrt{x^2}}\sqrt{x^2+x+1} = -\sqrt{1+(1/x)+(1/x^2)}$.

46. (a)

From the graph of $f(x)=\sqrt{3x^2+8x+6}-\sqrt{3x^2+3x+1}$, we estimate (to one decimal place) the value of $\lim\limits_{x\to\infty} f(x)$ to be 1.4.

(b)

x	$f(x)$
10,000	1.443 39
100,000	1.443 38
1,000,000	1.443 38

From the table, we estimate (to four decimal places) the limit to be 1.4434.

(c) $\lim\limits_{x\to\infty} f(x) = \lim\limits_{x\to\infty}\dfrac{\left(\sqrt{3x^2+8x+6}-\sqrt{3x^2+3x+1}\right)\left(\sqrt{3x^2+8x+6}+\sqrt{3x^2+3x+1}\right)}{\sqrt{3x^2+8x+6}+\sqrt{3x^2+3x+1}}$

$$= \lim_{x\to\infty}\frac{\left(3x^2+8x+6\right)-\left(3x^2+3x+1\right)}{\sqrt{3x^2+8x+6}+\sqrt{3x^2+3x+1}} = \lim_{x\to\infty}\frac{(5x+5)(1/x)}{\left(\sqrt{3x^2+8x+6}+\sqrt{3x^2+3x+1}\right)(1/x)}$$

$$= \lim_{x\to\infty}\frac{5+5/x}{\sqrt{3+8/x+6/x^2}+\sqrt{3+3/x+1/x^2}} = \frac{5}{\sqrt{3}+\sqrt{3}} = \frac{5}{2\sqrt{3}} = \frac{5\sqrt{3}}{6} \approx 1.443376$$

47. $\lim\limits_{x\to\pm\infty}\dfrac{5+4x}{x+3} = \lim\limits_{x\to\pm\infty}\dfrac{(5+4x)/x}{(x+3)/x} = \lim\limits_{x\to\pm\infty}\dfrac{5/x+4}{1+3/x} = \dfrac{0+4}{1+0} = 4$, so $y=4$ is a horizontal asymptote. $y=f(x)=\dfrac{5+4x}{x+3}$, so $\lim\limits_{x\to-3^+} f(x)=-\infty$ since $5+4x\to-7$ and $x+3\to0^+$ as $x\to-3^+$. Thus, $x=-3$ is a vertical asymptote. The graph confirms our work.

48. $\lim\limits_{x\to\pm\infty}\dfrac{2x^2+1}{3x^2+2x-1} = \lim\limits_{x\to\pm\infty}\dfrac{(2x^2+1)/x^2}{(3x^2+2x-1)/x^2}$

$$= \lim_{x\to\pm\infty}\frac{2+1/x^2}{3+2/x-1/x^2} = \frac{2}{3}$$

so $y=\dfrac{2}{3}$ is a horizontal asymptote. $y=f(x)=\dfrac{2x^2+1}{3x^2+2x-1}=\dfrac{2x^2+1}{(3x-1)(x+1)}$.

The denominator is zero when $x=\frac{1}{3}$ and -1, but the numerator is nonzero, so $x=\frac{1}{3}$ and $x=-1$ are vertical asymptotes. The graph confirms our work.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. $\displaystyle\lim_{x\to\pm\infty}\frac{2x^2+x-1}{x^2+x-2}=\lim_{x\to\pm\infty}\frac{\dfrac{2x^2+x-1}{x^2}}{\dfrac{x^2+x-2}{x^2}}=\lim_{x\to\pm\infty}\frac{2+\dfrac{1}{x}-\dfrac{1}{x^2}}{1+\dfrac{1}{x}-\dfrac{2}{x^2}}=\frac{\displaystyle\lim_{x\to\pm\infty}\left(2+\frac{1}{x}-\frac{1}{x^2}\right)}{\displaystyle\lim_{x\to\pm\infty}\left(1+\frac{1}{x}-\frac{2}{x^2}\right)}$

$$=\frac{\displaystyle\lim_{x\to\pm\infty}2+\lim_{x\to\pm\infty}\frac{1}{x}-\lim_{x\to\pm\infty}\frac{1}{x^2}}{\displaystyle\lim_{x\to\pm\infty}1+\lim_{x\to\pm\infty}\frac{1}{x}-2\lim_{x\to\pm\infty}\frac{1}{x^2}}=\frac{2+0-0}{1+0-2(0)}=2,\quad \text{so } y=2 \text{ is a horizontal asymptote.}$$

$y=f(x)=\dfrac{2x^2+x-1}{x^2+x-2}=\dfrac{(2x-1)(x+1)}{(x+2)(x-1)}$, so $\displaystyle\lim_{x\to-2^-}f(x)=\infty$, $\displaystyle\lim_{x\to-2^+}f(x)=-\infty$, $\displaystyle\lim_{x\to1^-}f(x)=-\infty$, and $\displaystyle\lim_{x\to1^+}f(x)=\infty$. Thus, $x=-2$ and $x=1$ are vertical asymptotes. The graph confirms our work.

50. $\displaystyle\lim_{x\to\pm\infty}\frac{1+x^4}{x^2-x^4}=\lim_{x\to\pm\infty}\frac{\dfrac{1+x^4}{x^4}}{\dfrac{x^2-x^4}{x^4}}=\lim_{x\to\pm\infty}\frac{\dfrac{1}{x^4}+1}{\dfrac{1}{x^2}-1}=\frac{\displaystyle\lim_{x\to\pm\infty}\left(\frac{1}{x^4}+1\right)}{\displaystyle\lim_{x\to\pm\infty}\left(\frac{1}{x^2}-1\right)}=\frac{\displaystyle\lim_{x\to\pm\infty}\frac{1}{x^4}+\lim_{x\to\pm\infty}1}{\displaystyle\lim_{x\to\pm\infty}\frac{1}{x^2}-\lim_{x\to\pm\infty}1}$

$$=\frac{0+1}{0-1}=-1,\quad \text{so } y=-1 \text{ is a horizontal asymptote.}$$

$y=f(x)=\dfrac{1+x^4}{x^2-x^4}=\dfrac{1+x^4}{x^2(1-x^2)}=\dfrac{1+x^4}{x^2(1+x)(1-x)}$. The denominator is zero when $x=0$, -1, and 1, but the numerator is nonzero, so $x=0$, $x=-1$, and $x=1$ are vertical asymptotes. Notice that as $x\to0$, the numerator and denominator are both positive, so $\displaystyle\lim_{x\to0}f(x)=\infty$. The graph confirms our work.

51. $y=f(x)=\dfrac{x^3-x}{x^2-6x+5}=\dfrac{x(x^2-1)}{(x-1)(x-5)}=\dfrac{x(x+1)(x-1)}{(x-1)(x-5)}=\dfrac{x(x+1)}{x-5}=g(x)$ for $x\neq1$.

The graph of g is the same as the graph of f with the exception of a hole in the graph of f at $x=1$. By long division, $g(x)=\dfrac{x^2+x}{x-5}=x+6+\dfrac{30}{x-5}$. As $x\to\pm\infty$, $g(x)\to\pm\infty$, so there is no horizontal asymptote. The denominator of g is zero when $x=5$. $\displaystyle\lim_{x\to5^-}g(x)=-\infty$ and $\displaystyle\lim_{x\to5^+}g(x)=\infty$, so $x=5$ is a vertical asymptote. The graph confirms our work.

52. $\displaystyle\lim_{x\to\infty}\frac{2e^x}{e^x-5}=\lim_{x\to\infty}\frac{2e^x}{e^x-5}\cdot\frac{1/e^x}{1/e^x}=\lim_{x\to\infty}\frac{2}{1-(5/e^x)}=\frac{2}{1-0}=2$, so $y=2$ is a horizontal asymptote.

$\displaystyle\lim_{x\to-\infty}\frac{2e^x}{e^x-5}=\frac{2(0)}{0-5}=0$, so $y=0$ is a horizontal asymptote. The denominator is zero (and the numerator isn't) when $e^x-5=0\quad\Rightarrow\quad e^x=5\quad\Rightarrow\quad x=\ln5$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\lim\limits_{x\to(\ln 5)^+} \dfrac{2e^x}{e^x-5} = \infty$ since the numerator approaches 10 and the denominator approaches 0 through positive values as $x \to (\ln 5)^+$. Similarly, $\lim\limits_{x\to(\ln 5)^-} \dfrac{2e^x}{e^x-5} = -\infty$. Thus, $x = \ln 5$ is a vertical asymptote. The graph confirms our work.

53. From the graph, it appears $y = 1$ is a horizontal asymptote.

$$\lim_{x\to\pm\infty} \frac{3x^3+500x^2}{x^3+500x^2+100x+2000} = \lim_{x\to\pm\infty} \frac{\dfrac{3x^3+500x^2}{x^3}}{\dfrac{x^3+500x^2+100x+2000}{x^3}}$$

$$= \lim_{x\to\pm\infty} \frac{3+(500/x)}{1+(500/x)+(100/x^2)+(2000/x^3)}$$

$$= \frac{3+0}{1+0+0+0} = 3, \quad \text{so } y = 3 \text{ is a horizontal asymptote.}$$

The discrepancy can be explained by the choice of the viewing window. Try $[-100{,}000,\ 100{,}000]$ by $[-1, 4]$ to get a graph that lends credibility to our calculation that $y = 3$ is a horizontal asymptote.

54. (a)

From the graph, it appears at first that there is only one horizontal asymptote, at $y \approx 0$, and a vertical asymptote at $x \approx 1.7$. However, if we graph the function with a wider and shorter viewing rectangle, we see that in fact there seem to be two horizontal asymptotes: one at $y \approx 0.5$ and one at $y \approx -0.5$. So we estimate that

$$\lim_{x\to\infty} \frac{\sqrt{2x^2+1}}{3x-5} \approx 0.5 \qquad \text{and} \qquad \lim_{x\to-\infty} \frac{\sqrt{2x^2+1}}{3x-5} \approx -0.5$$

(b) $f(1000) \approx 0.4722$ and $f(10{,}000) \approx 0.4715$, so we estimate that $\lim\limits_{x\to\infty} \dfrac{\sqrt{2x^2+1}}{3x-5} \approx 0.47$.

$f(-1000) \approx -0.4706$ and $f(-10{,}000) \approx -0.4713$, so we estimate that $\lim\limits_{x\to-\infty} \dfrac{\sqrt{2x^2+1}}{3x-5} \approx -0.47$.

(c) $\lim\limits_{x\to\infty} \dfrac{\sqrt{2x^2+1}}{3x-5} = \lim\limits_{x\to\infty} \dfrac{\sqrt{2+1/x^2}}{3-5/x}$ [since $\sqrt{x^2} = x$ for $x > 0$] $= \dfrac{\sqrt{2}}{3} \approx 0.471404$.

For $x < 0$, we have $\sqrt{x^2} = |x| = -x$, so when we divide the numerator by x, with $x < 0$, we

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

get $\dfrac{1}{x}\sqrt{2x^2+1} = -\dfrac{1}{\sqrt{x^2}}\sqrt{2x^2+1} = -\sqrt{2+1/x^2}$. Therefore,

$$\lim_{x\to-\infty}\frac{\sqrt{2x^2+1}}{3x-5} = \lim_{x\to-\infty}\frac{-\sqrt{2+1/x^2}}{3-5/x} = -\frac{\sqrt{2}}{3} \approx -0.471404.$$

55. Divide the numerator and the denominator by the highest power of x in $Q(x)$.

(a) If $\deg P < \deg Q$, then the numerator $\to 0$ but the denominator doesn't. So $\lim\limits_{x\to\infty}[P(x)/Q(x)] = 0$.

(b) If $\deg P > \deg Q$, then the numerator $\to \pm\infty$ but the denominator doesn't, so $\lim\limits_{x\to\infty}[P(x)/Q(x)] = \pm\infty$ (depending on the ratio of the leading coefficients of P and Q).

56.

(i) $n = 0$ (ii) $n > 0$ (n odd) (iii) $n > 0$ (n even) (iv) $n < 0$ (n odd) (v) $n < 0$ (n even)

From these sketches we see that

(a) $\lim\limits_{x\to0^+} x^n = \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{if } n > 0 \\ \infty & \text{if } n < 0 \end{cases}$

(b) $\lim\limits_{x\to0^-} x^n = \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{if } n > 0 \\ -\infty & \text{if } n < 0,\ n \text{ odd} \\ \infty & \text{if } n < 0,\ n \text{ even} \end{cases}$

(c) $\lim\limits_{x\to\infty} x^n = \begin{cases} 1 & \text{if } n = 0 \\ \infty & \text{if } n > 0 \\ 0 & \text{if } n < 0 \end{cases}$

(d) $\lim\limits_{x\to-\infty} x^n = \begin{cases} 1 & \text{if } n = 0 \\ -\infty & \text{if } n > 0,\ n \text{ odd} \\ \infty & \text{if } n > 0,\ n \text{ even} \\ 0 & \text{if } n < 0 \end{cases}$

57. Let's look for a rational function.

(1) $\lim\limits_{x\to\pm\infty} f(x) = 0 \quad\Rightarrow\quad$ degree of numerator $<$ degree of denominator

(2) $\lim\limits_{x\to0} f(x) = -\infty \quad\Rightarrow\quad$ there is a factor of x^2 in the denominator (not just x, since that would produce a sign change at $x = 0$), and the function is negative near $x = 0$.

(3) $\lim\limits_{x\to3^-} f(x) = \infty$ and $\lim\limits_{x\to3^+} f(x) = -\infty \quad\Rightarrow\quad$ vertical asymptote at $x = 3$; there is a factor of $(x-3)$ in the denominator.

(4) $f(2) = 0 \quad\Rightarrow\quad$ 2 is an x-intercept; there is at least one factor of $(x-2)$ in the numerator.

Combining all of this information and putting in a negative sign to give us the desired left- and right-hand limits gives us $f(x) = \dfrac{2-x}{x^2(x-3)}$ as one possibility.

58. Since the function has vertical asymptotes $x = 1$ and $x = 3$, the denominator of the rational function we are looking for must have factors $(x-1)$ and $(x-3)$. Because the horizontal asymptote is $y = 1$, the degree of the numerator must equal the degree of the denominator, and the ratio of the leading coefficients must be 1. One possibility is $f(x) = \dfrac{x^2}{(x-1)(x-3)}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. (a) We must first find the function f. Since f has a vertical asymptote $x = 4$ and x-intercept $x = 1$, $x - 4$ is a factor of the denominator and $x - 1$ is a factor of the numerator. There is a removable discontinuity at $x = -1$, so $x - (-1) = x + 1$ is a factor of both the numerator and denominator. Thus, f now looks like this: $f(x) = \dfrac{a(x-1)(x+1)}{(x-4)(x+1)}$, where a is still to be determined. Then $\lim\limits_{x\to-1} f(x) = \lim\limits_{x\to-1} \dfrac{a(x-1)(x+1)}{(x-4)(x+1)} = \lim\limits_{x\to-1} \dfrac{a(x-1)}{x-4} = \dfrac{a(-1-1)}{(-1-4)} = \dfrac{2}{5}a$, so $\dfrac{2}{5}a = 2$, and $a = 5$. Thus $f(x) = \dfrac{5(x-1)(x+1)}{(x-4)(x+1)}$ is a ratio of quadratic functions satisfying all the given conditions and $f(0) = \dfrac{5(-1)(1)}{(-4)(1)} = \dfrac{5}{4}$.

(b) $\lim\limits_{x\to\infty} f(x) = 5 \lim\limits_{x\to\infty} \dfrac{x^2-1}{x^2-3x-4} = 5 \lim\limits_{x\to\infty} \dfrac{(x^2/x^2)-(1/x^2)}{(x^2/x^2)-(3x/x^2)-(4/x^2)} = 5\dfrac{1-0}{1-0-0} = 5(1) = 5$

60. $y = f(x) = 2x^3 - x^4 = x^3(2-x)$. The y-intercept is $f(0) = 0$. The x-intercepts are 0 and 2. There are sign changes at 0 and 2 (odd exponents on x and $2 - x$). As $x \to \infty$, $f(x) \to -\infty$ because $x^3 \to \infty$ and $2 - x \to -\infty$. As $x \to -\infty$, $f(x) \to -\infty$ because $x^3 \to -\infty$ and $2 - x \to \infty$. Note that the graph of f near $x = 0$ flattens out (looks like $y = x^3$).

61. $y = f(x) = x^4 - x^6 = x^4(1-x^2) = x^4(1+x)(1-x)$. The y-intercept is $f(0) = 0$. The x-intercepts are 0, -1, and 1 [found by solving $f(x) = 0$ for x]. Since $x^4 > 0$ for $x \neq 0$, f doesn't change sign at $x = 0$. The function does change sign at $x = -1$ and $x = 1$. As $x \to \pm\infty$, $f(x) = x^4(1-x^2)$ approaches $-\infty$ because $x^4 \to \infty$ and $(1-x^2) \to -\infty$.

62. $y = f(x) = x^3(x+2)^2(x-1)$. The y-intercept is $f(0) = 0$. The x-intercepts are 0, -2, and 1. There are sign changes at 0 and 1 (odd exponents on x and $x - 1$). There is no sign change at -2. Also, $f(x) \to \infty$ as $x \to \infty$ because all three factors are large. And $f(x) \to \infty$ as $x \to -\infty$ because $x^3 \to -\infty$, $(x+2)^2 \to \infty$, and $(x-1) \to -\infty$. Note that the graph of f at $x = 0$ flattens out (looks like $y = -x^3$).

63. $y = f(x) = (3-x)(1+x)^2(1-x)^4$. The y-intercept is $f(0) = 3(1)^2(1)^4 = 3$. The x-intercepts are 3, -1, and 1. There is a sign change at 3, but not at -1 and 1. When x is large positive, $3 - x$ is negative and the other factors are positive, so $\lim\limits_{x\to\infty} f(x) = -\infty$. When x is large negative, $3 - x$ is positive, so $\lim\limits_{x\to-\infty} f(x) = \infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. $y = f(x) = x^2(x^2-1)^2(x+2) = x^2(x+1)^2(x-1)^2(x+2)$. The y-intercept is $f(0) = 0$. The x-intercepts are 0, -1, 1, and -2. There is a sign change at -2, but not at 0, -1, and 1. When x is large positive, all the factors are positive, so $\lim\limits_{x\to\infty} f(x) = \infty$. When x is large negative, only $x+2$ is negative, so $\lim\limits_{x\to-\infty} f(x) = -\infty$.

65. (a) Since $-1 \le \sin x \le 1$ for all x, $-\dfrac{1}{x} \le \dfrac{\sin x}{x} \le \dfrac{1}{x}$ for $x > 0$. As $x \to \infty$, $-1/x \to 0$ and $1/x \to 0$, so by the Squeeze Theorem, $(\sin x)/x \to 0$. Thus, $\lim\limits_{x\to\infty} \dfrac{\sin x}{x} = 0$.

(b) From part (a), the horizontal asymptote is $y = 0$. The function $y = (\sin x)/x$ crosses the horizontal asymptote whenever $\sin x = 0$; that is, at $x = \pi n$ for every integer n. Thus, the graph crosses the asymptote *an infinite number of times.*

66. (a) In both viewing rectangles, $\lim\limits_{x\to\infty} P(x) = \lim\limits_{x\to\infty} Q(x) = \infty$ and $\lim\limits_{x\to-\infty} P(x) = \lim\limits_{x\to-\infty} Q(x) = -\infty$. In the larger viewing rectangle, P and Q become less distinguishable.

(b) $\lim\limits_{x\to\infty} \dfrac{P(x)}{Q(x)} = \lim\limits_{x\to\infty} \dfrac{3x^5 - 5x^3 + 2x}{3x^5} = \lim\limits_{x\to\infty} \left(1 - \dfrac{5}{3}\cdot\dfrac{1}{x^2} + \dfrac{2}{3}\cdot\dfrac{1}{x^4}\right) = 1 - \frac{5}{3}(0) + \frac{2}{3}(0) = 1 \quad \Rightarrow$ P and Q have the same end behavior.

67. $\lim\limits_{x\to\infty} \dfrac{5\sqrt{x}}{\sqrt{x-1}} \cdot \dfrac{1/\sqrt{x}}{1/\sqrt{x}} = \lim\limits_{x\to\infty} \dfrac{5}{\sqrt{1-(1/x)}} = \dfrac{5}{\sqrt{1-0}} = 5$ and

$\lim\limits_{x\to\infty} \dfrac{10e^x - 21}{2e^x} \cdot \dfrac{1/e^x}{1/e^x} = \lim\limits_{x\to\infty} \dfrac{10 - (21/e^x)}{2} = \dfrac{10-0}{2} = 5$. Since $\dfrac{10e^x - 21}{2e^x} < f(x) < \dfrac{5\sqrt{x}}{\sqrt{x-1}}$,

we have $\lim\limits_{x\to\infty} f(x) = 5$ by the Squeeze Theorem.

68. (a) After t minutes, $25t$ liters of brine with 30 g of salt per liter has been pumped into the tank, so it contains $(5000 + 25t)$ liters of water and $25t \cdot 30 = 750t$ grams of salt. Therefore, the salt concentration at time t will be

$C(t) = \dfrac{750t}{5000 + 25t} = \dfrac{30t}{200 + t}\ \dfrac{\text{g}}{\text{L}}$.

(b) $\lim\limits_{t\to\infty} C(t) = \lim\limits_{t\to\infty} \dfrac{30t}{200+t} = \lim\limits_{t\to\infty} \dfrac{30t/t}{200/t + t/t} = \dfrac{30}{0+1} = 30$. So the salt concentration approaches that of the brine being pumped into the tank.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

69. (a) $\lim\limits_{t\to\infty} v(t) = \lim\limits_{t\to\infty} v^*\left(1 - e^{-gt/v^*}\right) = v^*(1-0) = v^*$

(b) We graph $v(t) = 1 - e^{-9.8t}$ and $v(t) = 0.99v^*$, or in this case, $v(t) = 0.99$. Using an intersect feature or zooming in on the point of intersection, we find that $t \approx 0.47$ s.

70. (a) $y = e^{-x/10}$ and $y = 0.1$ intersect at $x_1 \approx 23.03$.

If $x > x_1$, then $e^{-x/10} < 0.1$.

(b) $e^{-x/10} < 0.1 \quad \Rightarrow \quad -x/10 < \ln 0.1 \quad \Rightarrow$

$x > -10 \ln \frac{1}{10} = -10 \ln 10^{-1} = 10 \ln 10 \approx 23.03$

71. Let $g(x) = \dfrac{3x^2+1}{2x^2+x+1}$ and $f(x) = |g(x) - 1.5|$. Note that $\lim\limits_{x\to\infty} g(x) = \frac{3}{2}$ and $\lim\limits_{x\to\infty} f(x) = 0$. We are interested in finding the x-value at which $f(x) < 0.05$. From the graph, we find that $x \approx 14.804$, so we choose $N = 15$ (or any larger number).

72. We want to find a value of N such that $x > N \quad \Rightarrow \quad \left|\dfrac{1-3x}{\sqrt{x^2+1}} - (-3)\right| < \varepsilon$, or equivalently, $-3 - \varepsilon < \dfrac{1-3x}{\sqrt{x^2+1}} < -3 + \varepsilon$. When $\varepsilon = 0.1$, we graph $y = f(x) = \dfrac{1-3x}{\sqrt{x^2+1}}$, $y = -3.1$, and $y = -2.9$. From the graph, we find that $f(x) = -2.9$ at about $x = 11.283$, so we choose $N = 12$ (or any larger number). Similarly for $\varepsilon = 0.05$, we find that $f(x) = -2.95$ at about $x = 21.379$, so we choose $N = 22$ (or any larger number).

73. We want a value of N such that $x < N \Rightarrow \left|\dfrac{1-3x}{\sqrt{x^2+1}} - 3\right| < \varepsilon$, or equivalently, $3 - \varepsilon < \dfrac{1-3x}{\sqrt{x^2+1}} < 3 + \varepsilon$. When $\varepsilon = 0.1$, we graph $y = f(x) = \dfrac{1-3x}{\sqrt{x^2+1}}$, $y = 3.1$, and $y = 2.9$. From the graph, we find that $f(x) = 3.1$ at about $x = -8.092$, so we choose $N = -9$ (or any lesser number). Similarly for $\varepsilon = 0.05$, we find that $f(x) = 3.05$ at about $x = -18.338$, so we choose $N = -19$ (or any lesser number).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

74. We want to find a value of N such that $x > N \quad \Rightarrow \quad \sqrt{x \ln x} > 100$.

We graph $y = f(x) = \sqrt{x \ln x}$ and $y = 100$. From the graph, we find that $f(x) = 100$ at about $x = 1382.773$, so we choose $N = 1383$ (or any larger number).

75. (a) $1/x^2 < 0.0001 \quad \Leftrightarrow \quad x^2 > 1/0.0001 = 10\,000 \quad \Leftrightarrow \quad x > 100 \quad (x > 0)$

(b) If $\varepsilon > 0$ is given, then $1/x^2 < \varepsilon \quad \Leftrightarrow \quad x^2 > 1/\varepsilon \quad \Leftrightarrow \quad x > 1/\sqrt{\varepsilon}$. Let $N = 1/\sqrt{\varepsilon}$.

Then $x > N \quad \Rightarrow \quad x > \dfrac{1}{\sqrt{\varepsilon}} \quad \Rightarrow \quad \left|\dfrac{1}{x^2} - 0\right| = \dfrac{1}{x^2} < \varepsilon$, so $\displaystyle\lim_{x\to\infty} \frac{1}{x^2} = 0$.

76. (a) $1/\sqrt{x} < 0.0001 \quad \Leftrightarrow \quad \sqrt{x} > 1/0.0001 = 10^4 \quad \Leftrightarrow \quad x > 10^8$

(b) If $\varepsilon > 0$ is given, then $1/\sqrt{x} < \varepsilon \quad \Leftrightarrow \quad \sqrt{x} > 1/\varepsilon \quad \Leftrightarrow \quad x > 1/\varepsilon^2$. Let $N = 1/\varepsilon^2$.

Then $x > N \quad \Rightarrow \quad x > \dfrac{1}{\varepsilon^2} \quad \Rightarrow \quad \left|\dfrac{1}{\sqrt{x}} - 0\right| = \dfrac{1}{\sqrt{x}} < \varepsilon$, so $\displaystyle\lim_{x\to\infty} \frac{1}{\sqrt{x}} = 0$.

77. For $x < 0$, $|1/x - 0| = -1/x$. If $\varepsilon > 0$ is given, then $-1/x < \varepsilon \quad \Leftrightarrow \quad x < -1/\varepsilon$.

Take $N = -1/\varepsilon$. Then $x < N \quad \Rightarrow \quad x < -1/\varepsilon \quad \Rightarrow \quad |(1/x) - 0| = -1/x < \varepsilon$, so $\displaystyle\lim_{x\to-\infty} (1/x) = 0$.

78. Given $M > 0$, we need $N > 0$ such that $x > N \quad \Rightarrow \quad x^3 > M$. Now $x^3 > M \quad \Leftrightarrow \quad x > \sqrt[3]{M}$, so take $N = \sqrt[3]{M}$. Then $x > N = \sqrt[3]{M} \quad \Rightarrow \quad x^3 > M$, so $\displaystyle\lim_{x\to\infty} x^3 = \infty$.

79. Given $M > 0$, we need $N > 0$ such that $x > N \quad \Rightarrow \quad e^x > M$. Now $e^x > M \quad \Leftrightarrow \quad x > \ln M$, so take $N = \max(1, \ln M)$. (This ensures that $N > 0$.) Then $x > N = \max(1, \ln M) \quad \Rightarrow \quad e^x > \max(e, M) \geq M$, so $\displaystyle\lim_{x\to\infty} e^x = \infty$.

80. **Definition** Let f be a function defined on some interval $(-\infty, a)$. Then $\displaystyle\lim_{x\to-\infty} f(x) = -\infty$ means that for every negative number M there is a corresponding negative number N such that $f(x) < M$ whenever $x < N$.

Now we use the definition to prove that $\displaystyle\lim_{x\to-\infty} \left(1 + x^3\right) = -\infty$. Given a negative number M, we need a negative number N such that $x < N \quad \Rightarrow \quad 1 + x^3 < M$. Now $1 + x^3 < M \quad \Leftrightarrow \quad x^3 < M - 1 \quad \Leftrightarrow \quad x < \sqrt[3]{M-1}$. Thus, we take $N = \sqrt[3]{M-1}$ and find that $x < N \quad \Rightarrow \quad 1 + x^3 < M$. This proves that $\displaystyle\lim_{x\to-\infty} \left(1 + x^3\right) = -\infty$.

81. (a) Suppose that $\displaystyle\lim_{x\to\infty} f(x) = L$. Then for every $\varepsilon > 0$ there is a corresponding positive number N such that $|f(x) - L| < \varepsilon$ whenever $x > N$. If $t = 1/x$, then $x > N \quad \Leftrightarrow \quad 0 < 1/x < 1/N \quad \Leftrightarrow \quad 0 < t < 1/N$. Thus, for every $\varepsilon > 0$ there is a corresponding $\delta > 0$ (namely $1/N$) such that $|f(1/t) - L| < \varepsilon$ whenever $0 < t < \delta$. This proves that $\displaystyle\lim_{t\to0^+} f(1/t) = L = \lim_{x\to\infty} f(x)$.

Now suppose that $\displaystyle\lim_{x\to-\infty} f(x) = L$. Then for every $\varepsilon > 0$ there is a corresponding negative number N such that $|f(x) - L| < \varepsilon$ whenever $x < N$. If $t = 1/x$, then $x < N \quad \Leftrightarrow \quad 1/N < 1/x < 0 \quad \Leftrightarrow \quad 1/N < t < 0$. Thus, for every

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\varepsilon > 0$ there is a corresponding $\delta > 0$ (namely $-1/N$) such that $|f(1/t) - L| < \varepsilon$ whenever $-\delta < t < 0$. This proves that $\lim\limits_{t \to 0^-} f(1/t) = L = \lim\limits_{x \to -\infty} f(x)$.

(b)
$$\begin{aligned}
\lim_{x \to 0^+} x \sin \frac{1}{x} &= \lim_{t \to 0^+} t \sin \frac{1}{t} && [\text{let } x = t] \\
&= \lim_{y \to \infty} \frac{1}{y} \sin y && [\text{part (a) with } y = 1/t] \\
&= \lim_{x \to \infty} \frac{\sin x}{x} && [\text{let } y = x] \\
&= 0 && [\text{by Exercise 65}]
\end{aligned}$$

2.7 Derivatives and Rates of Change

1. (a) This is just the slope of the line through two points: $m_{PQ} = \dfrac{\Delta y}{\Delta x} = \dfrac{f(x) - f(3)}{x - 3}$.

(b) This is the limit of the slope of the secant line PQ as Q approaches P: $m = \lim\limits_{x \to 3} \dfrac{f(x) - f(3)}{x - 3}$.

2. The curve looks more like a line as the viewing rectangle gets smaller.

3. (a) (i) Using Definition 1 with $f(x) = x^2 + 3x$ and $P(-1, -2)$, the slope of the tangent line is

$$\begin{aligned}
m &= \lim_{x \to a} \frac{f(x) - f(a)}{x - a} = \lim_{x \to -1} \frac{(x^2 + 3x) - (-2)}{x - (-1)} = \lim_{x \to -1} \frac{x^2 + 3x + 2}{x + 1} = \lim_{x \to -1} \frac{(x + 2)(x + 1)}{x + 1} \\
&= \lim_{x \to -1} (x + 2) = -1 + 2 = 1
\end{aligned}$$

(ii) Using Equation 2 with $f(x) = x^2 + 3x$ and $P(-1, -2)$, the slope of the tangent line is

$$\begin{aligned}
m &= \lim_{h \to 0} \frac{f(a + h) - f(a)}{h} = \lim_{h \to 0} \frac{f(-1 + h) - f(-1)}{h} = \lim_{h \to 0} \frac{\left[(-1 + h)^2 + 3(-1 + h)\right] - (-2)}{h} \\
&= \lim_{h \to 0} \frac{1 - 2h + h^2 - 3 + 3h + 2}{h} = \lim_{h \to 0} \frac{h^2 + h}{h} = \lim_{h \to 0} \frac{h(h + 1)}{h} = \lim_{h \to 0} (h + 1) = 1
\end{aligned}$$

(b) An equation of the tangent line is $y - f(a) = f'(a)(x - a) \;\Rightarrow\; y - f(-1) = f'(-1)(x - (-1)) \;\Rightarrow$ $y - (-2) = 1(x + 1) \;\Rightarrow\; y + 2 = x + 1$, or $y = x - 1$.

(c)

The graph of $y = x - 1$ is tangent to the graph of $y = x^2 + 3x$ at the point $(-1, -2)$. Now zoom in toward the point $(-1, -2)$ until the parabola and the tangent line are indistinguishable.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a) (i) Using Definition 1 with $f(x) = x^3 + 1$ and $P(1, 2)$, the slope of the tangent line is

$$m = \lim_{x \to a} \frac{f(x) - f(a)}{x - a} = \lim_{x \to 1} \frac{(x^3 + 1) - 2}{x - 1} = \lim_{x \to 1} \frac{x^3 - 1}{x - 1} = \lim_{x \to 1} \frac{(x - 1)\left(x^2 + x + 1\right)}{x - 1}$$
$$= \lim_{x \to 1} \left(x^2 + x + 1\right) = 1^2 + 1 + 1 = 3$$

(ii) Using Equation 2 with $f(x) = x^3 + 1$ and $P(1, 2)$, the slope of the tangent line is

$$m = \lim_{h \to 0} \frac{f(a + h) - f(a)}{h} = \lim_{h \to 0} \frac{f(1 + h) - f(1)}{h} = \lim_{h \to 0} \frac{\left[(1 + h)^3 + 1\right] - 2}{h}$$
$$= \lim_{h \to 0} \frac{1 + 3h + 3h^2 + h^3 + 1 - 2}{h} = \lim_{h \to 0} \frac{h^3 + 3h^2 + 3h}{h} = \lim_{h \to 0} \frac{h(h^2 + 3h + 3)}{h}$$
$$= \lim_{h \to 0} (h^2 + 3h + 3) = 3$$

(b) An equation of the tangent line is $y - f(a) = f'(a)(x - a) \quad \Rightarrow \quad y - f(1) = f'(1)(x - 1) \quad \Rightarrow \quad y - 2 = 3\,(x - 1)$, or $y = 3x - 1$.

(c)

The graph of $y = 3x - 1$ is tangent to the graph of $y = x^3 + 1$ at the point $(1, 2)$. Now zoom in toward the point $(1, 2)$ until the cubic and the tangent line appear to coincide.

5. Using (1) with $f(x) = 2x^2 - 5x + 1$ and $P(3, 4)$ [we could also use Equation (2)], the slope of the tangent line is

$$m = \lim_{x \to a} \frac{f(x) - f(a)}{x - a} = \lim_{x \to 3} \frac{(2x^2 - 5x + 1) - 4}{x - 3} = \lim_{x \to 3} \frac{2x^2 - 5x - 3}{x - 3} = \lim_{x \to 3} \frac{(2x + 1)(x - 3)}{x - 3}$$
$$= \lim_{x \to 3} (2x + 1) = 2(3) + 1 = 7$$

Tangent line: $y - 4 = 7(x - 3) \quad \Leftrightarrow \quad y - 4 = 7x - 21 \quad \Leftrightarrow \quad y = 7x - 17$

6. Using (2) with $f(x) = x^2 - 2x^3$ and $P(1, -1)$, the slope of the tangent line is

$$m = \lim_{h \to 0} \frac{f(a + h) - f(a)}{h} = \lim_{h \to 0} \frac{f(1 + h) - f(1)}{h} = \lim_{h \to 0} \frac{\left[(1 + h)^2 - 2(1 + h)^3\right] - (-1)}{h}$$
$$= \lim_{h \to 0} \frac{1 + 2h + h^2 - 2 - 6h - 6h^2 - 2h^3 + 1}{h} = \lim_{h \to 0} \frac{-2h^3 - 5h^2 - 4h}{h} = \lim_{h \to 0} \frac{-h(2h^2 + 5h + 4)}{h}$$
$$= \lim_{h \to 0} \left[-(2h^2 + 5h + 4)\right] = -4$$

Tangent line: $y - (-1) = -4(x - 1) \quad \Leftrightarrow \quad y + 1 = -4x + 4 \quad \Leftrightarrow \quad y = -4x + 3$

7. Using (1) with $f(x) = \dfrac{x + 2}{x - 3}$ and $P(2, -4)$, the slope of the tangent line is

$$m = \lim_{x \to a} \frac{f(x) - f(a)}{x - a} = \lim_{x \to 2} \frac{\dfrac{x + 2}{x - 3} - (-4)}{x - 2} = \lim_{x \to 2} \frac{\dfrac{x + 2 + 4(x - 3)}{x - 3}}{x - 2} = \lim_{x \to 2} \frac{5x - 10}{(x - 2)(x - 3)}$$
$$= \lim_{x \to 2} \frac{5(x - 2)}{(x - 2)(x - 3)} = \lim_{x \to 2} \frac{5}{x - 3} = \frac{5}{2 - 3} = -5$$

Tangent line: $y - (-4) = -5(x - 2) \quad \Leftrightarrow \quad y + 4 = -5x + 10 \quad \Leftrightarrow \quad y = -5x + 6$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. Using (1) with $f(x) = \sqrt{1-3x}$ and $P(-1, 2)$, the slope of the tangent line is

$$m = \lim_{x\to a} \frac{f(x)-f(a)}{x-a} = \lim_{x\to -1} \frac{\sqrt{1-3x}-2}{x-(-1)} = \lim_{x\to -1} \frac{\left(\sqrt{1-3x}-2\right)\left(\sqrt{1-3x}+2\right)}{(x+1)\left(\sqrt{1-3x}+2\right)} = \lim_{x\to -1} \frac{(1-3x)-4}{(x+1)\left(\sqrt{1-3x}+2\right)}$$

$$= \lim_{x\to -1} \frac{-3x-3}{(x+1)\left(\sqrt{1-3x}+2\right)} = \lim_{x\to -1} \frac{-3(x+1)}{(x+1)\left(\sqrt{1-3x}+2\right)} = \lim_{x\to -1} \frac{-3}{\sqrt{1-3x}+2} = \frac{-3}{2+2} = -\frac{3}{4}$$

Tangent line: $y - 2 = -\frac{3}{4}[x-(-1)] \quad \Leftrightarrow \quad y - 2 = -\frac{3}{4}x - \frac{3}{4} \quad \Leftrightarrow \quad y = -\frac{3}{4}x + \frac{5}{4}$

9. (a) Using (2) with $y = f(x) = 3 + 4x^2 - 2x^3$, the slope of the tangent line is

$$\begin{aligned} m &= \lim_{h\to 0} \frac{f(a+h)-f(a)}{h} = \lim_{h\to 0} \frac{3+4(a+h)^2-2(a+h)^3-(3+4a^2-2a^3)}{h} \\ &= \lim_{h\to 0} \frac{3+4(a^2+2ah+h^2)-2(a^3+3a^2h+3ah^2+h^3)-3-4a^2+2a^3}{h} \\ &= \lim_{h\to 0} \frac{3+4a^2+8ah+4h^2-2a^3-6a^2h-6ah^2-2h^3-3-4a^2+2a^3}{h} \\ &= \lim_{h\to 0} \frac{8ah+4h^2-6a^2h-6ah^2-2h^3}{h} = \lim_{h\to 0} \frac{h(8a+4h-6a^2-6ah-2h^2)}{h} \\ &= \lim_{h\to 0} (8a+4h-6a^2-6ah-2h^2) = 8a-6a^2 \end{aligned}$$

(b) At $(1, 5)$: $m = 8(1) - 6(1)^2 = 2$, so an equation of the tangent line is $y - 5 = 2(x-1) \quad \Leftrightarrow \quad y = 2x + 3$.

At $(2, 3)$: $m = 8(2) - 6(2)^2 = -8$, so an equation of the tangent line is $y - 3 = -8(x-2) \quad \Leftrightarrow \quad y = -8x + 19$.

(c)

10. (a) Using (1) with $f(x) = 2\sqrt{x}$ and $P\left(a, 2\sqrt{a}\right)$, the slope of the tangent line is

$$m = \lim_{x\to a} \frac{2\sqrt{x}-2\sqrt{a}}{x-a} = \lim_{x\to a} \frac{\left(2\sqrt{x}-2\sqrt{a}\right)\left(2\sqrt{x}+2\sqrt{a}\right)}{(x-a)\left(2\sqrt{x}+2\sqrt{a}\right)} = \lim_{x\to a} \frac{4x-4a}{(x-a)\left(2\sqrt{x}+2\sqrt{a}\right)}$$

$$= \lim_{x\to a} \frac{4(x-a)}{(x-a)\left(2\sqrt{x}+2\sqrt{a}\right)} = \lim_{x\to a} \frac{4}{2\sqrt{x}+2\sqrt{a}} = \frac{4}{2\sqrt{a}+2\sqrt{a}} = \frac{4}{4\sqrt{a}} = \frac{1}{\sqrt{a}}, \text{ or } \frac{\sqrt{a}}{a} \quad [a > 0]$$

(b) At $(1, 2)$: $m = \dfrac{1}{\sqrt{1}} = 1$, so an equation of the tangent line is $y - 2 = 1(x-1) \quad \Leftrightarrow \quad y = x + 1$.

At $(9, 6)$: $m = \dfrac{1}{\sqrt{9}} = \dfrac{1}{3}$, so an equation of the tangent line is $y - 6 = \frac{1}{3}(x-9) \quad \Leftrightarrow \quad y = \frac{1}{3}x + 3$.

(c)

11. (a) We have $d(t) = 16t^2$. The diver will hit the water when $d(t) = 100 \quad \Leftrightarrow \quad 16t^2 = 100 \quad \Leftrightarrow \quad t^2 = \frac{25}{4} \quad \Leftrightarrow$ $t = \frac{5}{2}$ $(t > 0)$. The diver will hit the water after 2.5 seconds.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) By Definition 3, the instantaneous velocity of an object with position function $d(t)$ at time $t = 2.5$ is

$$v(2.5) = \lim_{h\to 0} \frac{d(2.5+h) - d(2.5)}{h} = \lim_{h\to 0} \frac{16(2.5+h)^2 - 100}{h} = \lim_{h\to 0} \frac{100 + 80h + 16h^2 - 100}{h}$$

$$= \lim_{h\to 0} \frac{80h + 16h^2}{h} = \lim_{h\to 0} \frac{h(80+16h)}{h} = \lim_{h\to 0}(80+16h) = 80$$

The diver will hit the water with a velocity of 80 ft/s.

12. (a) Let $H(t) = 10t - 1.86t^2$.

$$v(1) = \lim_{h\to 0} \frac{H(1+h) - H(1)}{h} = \lim_{h\to 0} \frac{\left[10(1+h) - 1.86(1+h)^2\right] - (10 - 1.86)}{h}$$

$$= \lim_{h\to 0} \frac{10 + 10h - 1.86(1 + 2h + h^2) - 10 + 1.86}{h} = \lim_{h\to 0} \frac{10 + 10h - 1.86 - 3.72h - 1.86h^2 - 10 + 1.86}{h}$$

$$= \lim_{h\to 0} \frac{6.28h - 1.86h^2}{h} = \lim_{h\to 0}(6.28 - 1.86h) = 6.28$$

The velocity of the rock after one second is 6.28 m/s.

(b) $$v(a) = \lim_{h\to 0} \frac{H(a+h) - H(a)}{h} = \lim_{h\to 0} \frac{\left[10(a+h) - 1.86(a+h)^2\right] - (10a - 1.86a^2)}{h}$$

$$= \lim_{h\to 0} \frac{10a + 10h - 1.86(a^2 + 2ah + h^2) - 10a + 1.86a^2}{h}$$

$$= \lim_{h\to 0} \frac{10a + 10h - 1.86a^2 - 3.72ah - 1.86h^2 - 10a + 1.86a^2}{h} = \lim_{h\to 0} \frac{10h - 3.72ah - 1.86h^2}{h}$$

$$= \lim_{h\to 0} \frac{h(10 - 3.72a - 1.86h)}{h} = \lim_{h\to 0}(10 - 3.72a - 1.86h) = 10 - 3.72a$$

The velocity of the rock when $t = a$ is $(10 - 3.72a)$ m/s.

(c) The rock will hit the surface when $H = 0 \quad \Leftrightarrow \quad 10t - 1.86t^2 = 0 \quad \Leftrightarrow \quad t(10 - 1.86t) = 0 \quad \Leftrightarrow \quad t = 0$ or $1.86t = 10$.

The rock hits the surface when $t = 10/1.86 \approx 5.4$ s.

(d) The velocity of the rock when it hits the surface is $v\left(\frac{10}{1.86}\right) = 10 - 3.72\left(\frac{10}{1.86}\right) = 10 - 20 = -10$ m/s.

13. $$v(a) = \lim_{h\to 0} \frac{s(a+h) - s(a)}{h} = \lim_{h\to 0} \frac{\dfrac{1}{(a+h)^2} - \dfrac{1}{a^2}}{h} = \lim_{h\to 0} \frac{\dfrac{a^2 - (a+h)^2}{a^2(a+h)^2}}{h}$$

$$= \lim_{h\to 0} \frac{a^2 - (a^2 + 2ah + h^2)}{ha^2(a+h)^2} = \lim_{h\to 0} \frac{-(2ah + h^2)}{ha^2(a+h)^2}$$

$$= \lim_{h\to 0} \frac{-h(2a+h)}{ha^2(a+h)^2} = \lim_{h\to 0} \frac{-(2a+h)}{a^2(a+h)^2} = \frac{-2a}{a^2 \cdot a^2} = \frac{-2}{a^3} \text{ m/s}$$

So $v(1) = \dfrac{-2}{1^3} = -2$ m/s, $v(2) = \dfrac{-2}{2^3} = -\dfrac{1}{4}$m/s, and $v(3) = \dfrac{-2}{3^3} = -\dfrac{2}{27}$ m/s.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. (a) The average velocity between times t and $t+h$ is

$$\begin{aligned}\frac{s(t+h)-s(t)}{(t+h)-t} &= \frac{\frac{1}{2}(t+h)^2-6(t+h)+23-\left(\frac{1}{2}t^2-6t+23\right)}{h}\\ &= \frac{\frac{1}{2}t^2+th+\frac{1}{2}h^2-6t-6h+23-\frac{1}{2}t^2+6t-23}{h}\\ &= \frac{th+\frac{1}{2}h^2-6h}{h} = \frac{h\left(t+\frac{1}{2}h-6\right)}{h} = \left(t+\tfrac{1}{2}h-6\right)\ \text{ft/s}\end{aligned}$$

(i) $[4,8]$: $t=4$, $h=8-4=4$, so the average velocity is $4+\frac{1}{2}(4)-6=0$ ft/s.

(ii) $[6,8]$: $t=6$, $h=8-6=2$, so the average velocity is $6+\frac{1}{2}(2)-6=1$ ft/s.

(iii) $[8,10]$: $t=8$, $h=10-8=2$, so the average velocity is $8+\frac{1}{2}(2)-6=3$ ft/s.

(iv) $[8,12]$: $t=8$, $h=12-8=4$, so the average velocity is $8+\frac{1}{2}(4)-6=4$ ft/s.

(b) $v(t)=\lim\limits_{h\to 0}\dfrac{s(t+h)-s(t)}{h}=\lim\limits_{h\to 0}\left(t+\frac{1}{2}h-6\right)$
$=t-6$, so $v(8)=2$ ft/s.

(c)

15. (a) The particle is moving to the right when s is increasing; that is, on the intervals $(0,1)$ and $(4,6)$. The particle is moving to the left when s is decreasing; that is, on the interval $(2,3)$. The particle is standing still when s is constant; that is, on the intervals $(1,2)$ and $(3,4)$.

(b) The velocity of the particle is equal to the slope of the tangent line of the graph. Note that there is no slope at the corner points on the graph. On the interval $(0,1)$, the slope is $\dfrac{3-0}{1-0}=3$. On the interval $(2,3)$, the slope is $\dfrac{1-3}{3-2}=-2$. On the interval $(4,6)$, the slope is $\dfrac{3-1}{6-4}=1$.

16. (a) **Runner A** runs the entire 100-meter race at the same velocity since the slope of the position function is constant. **Runner B** starts the race at a slower velocity than runner A, but finishes the race at a faster velocity.

(b) The distance between the runners is the greatest at the time when the largest vertical line segment fits between the two graphs—this appears to be somewhere between 9 and 10 seconds.

(c) The runners had the same velocity when the slopes of their respective position functions are equal—this also appears to be at about 9.5 s. Note that the answers for parts (b) and (c) must be the same for these graphs because as soon as the velocity for runner B overtakes the velocity for runner A, the distance between the runners starts to decrease.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $g'(0)$ is the only negative value. The slope at $x = 4$ is smaller than the slope at $x = 2$ and both are smaller than the slope at $x = -2$. Thus, $g'(0) < 0 < g'(4) < g'(2) < g'(-2)$.

18. (a) On $[20, 60]$: $\dfrac{f(60) - f(20)}{60 - 20} = \dfrac{700 - 300}{40} = \dfrac{400}{40} = 10$

(b) Pick any interval that has the same y-value at both endpoints. $[10, 50]$ is such an interval since $f(10) = 400$ and $f(50) = 400$.

(c) $\dfrac{f(40) - f(10)}{40 - 10} = \dfrac{200 - 400}{30} = \dfrac{-200}{30} = -\dfrac{20}{3}$

This value represents the slope of the line segment from $(10, f(10))$ to $(40, f(40))$.

(d) The tangent line at $x = 50$ appears to pass through the points $(40, 200)$ and $(60, 700)$, so

$f'(50) \approx \dfrac{700 - 200}{60 - 40} = \dfrac{500}{20} = 25.$

(e) The tangent line at $x = 10$ is steeper than the tangent line at $x = 30$, so it is larger in magnitude, but less in numerical value, that is, $f'(10) < f'(30)$.

(f) The slope of the tangent line at $x = 60$, $f'(60)$, is greater than the slope of the line through $(40, f(40))$ and $(80, f(80))$.

So yes, $f'(60) > \dfrac{f(80) - f(40)}{80 - 40}$.

19. Using Definition 4 with $f(x) = \sqrt{4x + 1}$ and $a = 6$,

$$\begin{aligned} f'(6) &= \lim_{h \to 0} \frac{f(6 + h) - f(6)}{h} = \lim_{h \to 0} \frac{\sqrt{4(6 + h) + 1} - 5}{h} = \lim_{h \to 0} \frac{\sqrt{25 + 4h} - 5}{h} \\ &= \lim_{h \to 0} \frac{\left(\sqrt{25 + 4h} - 5\right)\left(\sqrt{25 + 4h} + 5\right)}{h\left(\sqrt{25 + 4h} + 5\right)} = \lim_{h \to 0} \frac{(25 + 4h) - 25}{h\left(\sqrt{25 + 4h} + 5\right)} = \lim_{h \to 0} \frac{4h}{h\left(\sqrt{25 + 4h} + 5\right)} \\ &= \lim_{h \to 0} \frac{4}{\sqrt{25 + 4h} + 5} = \frac{4}{5 + 5} = \frac{2}{5} \end{aligned}$$

20. Using Definition 4 with $f(x) = 5x^4$ and $a = -1$,

$$\begin{aligned} f'(-1) &= \lim_{h \to 0} \frac{f(-1 + h) - f(-1)}{h} = \lim_{h \to 0} \frac{5(-1 + h)^4 - 5}{h} = \lim_{h \to 0} \frac{5(1 - 4h + 6h^2 - 4h^3 + h^4) - 5}{h} \\ &= \lim_{h \to 0} \frac{-20h + 30h^2 - 20h^3 + 5h^4}{h} = \lim_{h \to 0} \frac{-h(20 - 30h + 20h^2 - 5h^3)}{h} \\ &= \lim_{h \to 0} \left[-(20 - 30h + 20h^2 - 5h^3)\right] = -20 \end{aligned}$$

21. Using Equation 5 with $f(x) = \dfrac{x^2}{x + 6}$ and $a = 3$,

$$\begin{aligned} f'(3) &= \lim_{x \to 3} \frac{f(x) - f(3)}{x - 3} = \lim_{x \to 3} \frac{\dfrac{x^2}{x + 6} - 1}{x - 3} = \lim_{x \to 3} \frac{\dfrac{x^2 - (x + 6)}{x + 6}}{x - 3} = \lim_{x \to 3} \frac{x^2 - x - 6}{(x + 6)(x - 3)} \\ &= \lim_{x \to 3} \frac{(x + 2)(x - 3)}{(x + 6)(x - 3)} = \lim_{x \to 3} \frac{x + 2}{x + 6} = \frac{3 + 2}{3 + 6} = \frac{5}{9} \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. Using Equation 5 with $f(x) = \dfrac{1}{\sqrt{2x+2}}$ and $a = 1$,

$$f'(1) = \lim_{x\to 1} \frac{f(x)-f(1)}{x-1} = \lim_{x\to 1} \frac{\dfrac{1}{\sqrt{2x+2}} - \dfrac{1}{2}}{x-1} = \lim_{x\to 1} \frac{\dfrac{2-\sqrt{2x+2}}{2\sqrt{2x+2}}}{x-1} = \lim_{x\to 1} \frac{2-\sqrt{2x+2}}{2\sqrt{2x+2}\,(x-1)}$$

$$= \lim_{x\to 1} \frac{\left(2-\sqrt{2x+2}\right)\left(2+\sqrt{2x+2}\right)}{2\sqrt{2x+2}\,(x-1)\left(2+\sqrt{2x+2}\right)} = \lim_{x\to 1} \frac{4-(2x+2)}{2\sqrt{2x+2}\,(x-1)\left(2+\sqrt{2x+2}\right)}$$

$$= \lim_{x\to 1} \frac{-2x+2}{2\sqrt{2x+2}\,(x-1)\left(2+\sqrt{2x+2}\right)} = \lim_{x\to 1} \frac{-2(x-1)}{2\sqrt{2x+2}\,(x-1)\left(2+\sqrt{2x+2}\right)}$$

$$= \lim_{x\to 1} \frac{-1}{\sqrt{2x+2}\left(2+\sqrt{2x+2}\right)} = \frac{-1}{\sqrt{4}\left(2+\sqrt{4}\right)} = -\frac{1}{8}$$

23. Using Definition 4 with $f(x) = 2x^2 - 5x + 3$,

$$f'(a) = \lim_{h\to 0} \frac{f(a+h)-f(a)}{h} = \lim_{h\to 0} \frac{\left[2(a+h)^2 - 5(a+h) + 3\right] - (2a^2 - 5a + 3)}{h}$$

$$= \lim_{h\to 0} \frac{2a^2 + 4ah + 2h^2 - 5a - 5h + 3 - 2a^2 + 5a - 3}{h} = \lim_{h\to 0} \frac{4ah + 2h^2 - 5h}{h}$$

$$= \lim_{h\to 0} \frac{h(4a+2h-5)}{h} = \lim_{h\to 0}(4a+2h-5) = 4a-5$$

24. Using Definition 4 with $f(t) = t^3 - 3t$,

$$f'(a) = \lim_{h\to 0} \frac{f(a+h)-f(a)}{h} = \lim_{h\to 0} \frac{\left[(a+h)^3 - 3(a+h)\right] - (a^3 - 3a)}{h}$$

$$= \lim_{h\to 0} \frac{a^3 + 3a^2h + 3ah^2 + h^3 - 3a - 3h - a^3 + 3a}{h} = \lim_{h\to 0} \frac{3a^2h + 3ah^2 + h^3 - 3h}{h}$$

$$= \lim_{h\to 0} \frac{h(3a^2 + 3ah + h^2 - 3)}{h} = \lim_{h\to 0}(3a^2 + 3ah + h^2 - 3) = 3a^2 - 3$$

25. Using Definition 4 with $f(t) = \dfrac{1}{t^2+1}$.

$$f'(a) = \lim_{h\to 0} \frac{f(a+h)-f(a)}{h} = \lim_{h\to 0} \frac{\dfrac{1}{(a+h)^2+1} - \dfrac{1}{a^2+1}}{h} = \lim_{h\to 0} \frac{\dfrac{(a^2+1) - \left[(a+h)^2+1\right]}{\left[(a+h)^2+1\right](a^2+1)}}{h}$$

$$= \lim_{h\to 0} \frac{\left(a^2+1\right) - (a^2 + 2ah + h^2 + 1)}{h[(a+h)^2+1](a^2+1)} = \lim_{h\to 0} \frac{-(2ah+h^2)}{h[(a+h)^2+1](a^2+1)} = \lim_{h\to 0} \frac{-h(2a+h)}{h[(a+h)^2+1](a^2+1)}$$

$$= \lim_{h\to 0} \frac{-(2a+h)}{[(a+h)^2+1](a^2+1)} = \frac{-2a}{(a^2+1)(a^2+1)} = -\frac{2a}{(a^2+1)^2}$$

26. Use Definition 4 with $f(x) = \dfrac{x}{1-4x}$.

$$f'(a) = \lim_{h\to 0} \frac{f(a+h)-f(a)}{h} = \lim_{h\to 0} \frac{\dfrac{a+h}{1-4(a+h)} - \dfrac{a}{1-4a}}{h} = \lim_{h\to 0} \frac{\dfrac{(a+h)(1-4a) - a\left[1-4(a+h)\right]}{[1-4(a+h)](1-4a)}}{h}$$

$$= \lim_{h\to 0} \frac{a - 4a^2 + h - 4ah - a + 4a^2 + 4ah}{h[1-4(a+h)](1-4a)} = \lim_{h\to 0} \frac{h}{h[1-4(a+h)](1-4a)}$$

$$= \lim_{h\to 0} \frac{1}{[1-4(a+h)](1-4a)} = \frac{1}{(1-4a)(1-4a)} = \frac{1}{(1-4a)^2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. Since $B(6) = 0$, the point $(6, 0)$ is on the graph of B. Since $B'(6) = -\frac{1}{2}$, the slope of the tangent line at $x = 6$ is $-\frac{1}{2}$.

Using the point-slope form of a line gives us $y - 0 = -\frac{1}{2}(x - 6)$, or $y = -\frac{1}{2}x + 3$.

28. Since $g(5) = -3$, the point $(5, -3)$ is on the graph of g. Since $g'(5) = 4$, the slope of the tangent line at $x = 5$ is 4.

Using the point-slope form of a line gives us $y - (-3) = 4(x - 5)$, or $y = 4x - 23$.

29. Using Definition 4 with $f(x) = 3x^2 - x^3$ and $a = 1$,

$$\begin{aligned} f'(1) &= \lim_{h\to 0} \frac{f(1+h) - f(1)}{h} = \lim_{h\to 0} \frac{[3(1+h)^2 - (1+h)^3] - 2}{h} \\ &= \lim_{h\to 0} \frac{(3 + 6h + 3h^2) - (1 + 3h + 3h^2 + h^3) - 2}{h} = \lim_{h\to 0} \frac{3h - h^3}{h} = \lim_{h\to 0} \frac{h(3 - h^2)}{h} \\ &= \lim_{h\to 0} (3 - h^2) = 3 - 0 = 3 \end{aligned}$$

Tangent line: $y - 2 = 3(x - 1) \quad \Leftrightarrow \quad y - 2 = 3x - 3 \quad \Leftrightarrow \quad y = 3x - 1$

30. Using Equation 5 with $g(x) = x^4 - 2$ and $a = 1$,

$$\begin{aligned} g'(1) &= \lim_{x\to 1} \frac{g(x) - g(1)}{x - 1} = \lim_{x\to 1} \frac{(x^4 - 2) - (-1)}{x - 1} = \lim_{x\to 1} \frac{x^4 - 1}{x - 1} = \lim_{x\to 1} \frac{(x^2 + 1)(x^2 - 1)}{x - 1} \\ &= \lim_{x\to 1} \frac{(x^2 + 1)(x + 1)(x - 1)}{x - 1} = \lim_{x\to 1} [(x^2 + 1)(x + 1)] = 2(2) = 4 \end{aligned}$$

Tangent line: $y - (-1) = 4(x - 1) \quad \Leftrightarrow \quad y + 1 = 4x - 4 \quad \Leftrightarrow \quad y = 4x - 5$

31. (a) Using Definition 4 with $F(x) = 5x/(1 + x^2)$ and the point $(2, 2)$, we have

$$\begin{aligned} F'(2) &= \lim_{h\to 0} \frac{F(2+h) - F(2)}{h} = \lim_{h\to 0} \frac{\dfrac{5(2+h)}{1 + (2+h)^2} - 2}{h} \\ &= \lim_{h\to 0} \frac{\dfrac{5h + 10}{h^2 + 4h + 5} - 2}{h} = \lim_{h\to 0} \frac{\dfrac{5h + 10 - 2(h^2 + 4h + 5)}{h^2 + 4h + 5}}{h} \\ &= \lim_{h\to 0} \frac{-2h^2 - 3h}{h(h^2 + 4h + 5)} = \lim_{h\to 0} \frac{h(-2h - 3)}{h(h^2 + 4h + 5)} = \lim_{h\to 0} \frac{-2h - 3}{h^2 + 4h + 5} = \frac{-3}{5} \end{aligned}$$

So an equation of the tangent line at $(2, 2)$ is $y - 2 = -\frac{3}{5}(x - 2)$ or $y = -\frac{3}{5}x + \frac{16}{5}$.

(b)

32. (a) Using Definition 4 with $G(x) = 4x^2 - x^3$, we have

$$\begin{aligned} G'(a) &= \lim_{h\to 0} \frac{G(a+h) - G(a)}{h} = \lim_{h\to 0} \frac{[4(a+h)^2 - (a+h)^3] - (4a^2 - a^3)}{h} \\ &= \lim_{h\to 0} \frac{4a^2 + 8ah + 4h^2 - (a^3 + 3a^2h + 3ah^2 + h^3) - 4a^2 + a^3}{h} \\ &= \lim_{h\to 0} \frac{8ah + 4h^2 - 3a^2h - 3ah^2 - h^3}{h} = \lim_{h\to 0} \frac{h(8a + 4h - 3a^2 - 3ah - h^2)}{h} \\ &= \lim_{h\to 0} (8a + 4h - 3a^2 - 3ah - h^2) = 8a - 3a^2 \end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

At the point $(2, 8)$, $G'(2) = 16 - 12 = 4$, and an equation of the tangent line is $y - 8 = 4(x - 2)$, or $y = 4x$. At the point $(3, 9)$, $G'(3) = 24 - 27 = -3$, and an equation of the tangent line is $y - 9 = -3(x - 3)$, or $y = -3x + 18$.

(b)

33. For the tangent line $y = 4x - 5$: when $x = 2$, $y = 4(2) - 5 = 3$ and its slope is 4 (the coefficient of x). At the point of tangency, these values are shared with the curve $y = f(x)$; that is, $f(2) = 3$ and $f'(2) = 4$.

34. Since $(4, 3)$ is on $y = f(x)$, $f(4) = 3$. The slope of the tangent line between $(0, 2)$ and $(4, 3)$ is $\frac{1}{4}$, so $f'(4) = \frac{1}{4}$.

35.
$$\begin{aligned} v(4) = f'(4) &= \lim_{h\to 0} \frac{f(4+h) - f(4)}{h} = \lim_{h\to 0} \frac{\left[80(4+h) - 6(4+h)^2\right] - \left[80(4) - 6(4)^2\right]}{h} \\ &= \lim_{h\to 0} \frac{(320 + 80h - 96 - 48h - 6h^2) - (320 - 96)}{h} = \lim_{h\to 0} \frac{32h - 6h^2}{h} \\ &= \lim_{h\to 0} \frac{h(32 - 6h)}{h} = \lim_{h\to 0}(32 - 6h) = 32 \text{ m/s} \end{aligned}$$

The speed when $t = 4$ is $|32| = 32$ m/s.

36.
$$\begin{aligned} v(4) = f'(4) &= \lim_{h\to 0} \frac{f(4+h) - f(4)}{h} = \lim_{h\to 0} \frac{\left(10 + \dfrac{45}{4+h+1}\right) - \left(10 + \dfrac{45}{4+1}\right)}{h} = \lim_{h\to 0} \frac{\dfrac{45}{5+h} - 9}{h} \\ &= \lim_{h\to 0} \frac{45 - 9(5+h)}{h(5+h)} = \lim_{h\to 0} \frac{-9h}{h(5+h)} = \lim_{h\to 0} \frac{-9}{5+h} = -\frac{9}{5} \text{ m/s.} \end{aligned}$$

The speed when $t = 4$ is $\left|-\frac{9}{5}\right| = \frac{9}{5}$ m/s.

37. The sketch shows the graph for a room temperature of $72°$ and a refrigerator temperature of $38°$. The initial rate of change is greater in magnitude than the rate of change after an hour.

38. The slope of the tangent (that is, the rate of change of temperature with respect to time) at $t = 1$ h seems to be about $\dfrac{75 - 168}{132 - 0} \approx -0.7\,°\text{F/min}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. We begin by drawing a curve through the origin with a slope of 3 to satisfy $f(0) = 0$ and $f'(0) = 3$. Since $f'(1) = 0$, we will round off our figure so that there is a horizontal tangent directly over $x = 1$. Last, we make sure that the curve has a slope of -1 as we pass over $x = 2$. Two of the many possibilities are shown.

40. We begin by drawing a curve through the origin with a slope of 1 to satisfy $g(0) = 0$ and $g'(0) = 1$. We round off our figure at $x = 1$ to satisfy $g'(1) = 0$, and then pass through $(2, 0)$ with slope -1 to satisfy $g(2) = 0$ and $g'(2) = -1$. We round the figure at $x = 3$ to satisfy $g'(3) = 0$, and then pass through $(4, 0)$ with slope 1 to satisfy $g(4) = 0$ and $g'(4) = 1$. Finally we extend the curve on both ends to satisfy $\lim_{x\to\infty} g(x) = \infty$ and $\lim_{x\to-\infty} g(x) = -\infty$.

41. We begin by drawing a curve through $(0, 1)$ with a slope of 1 to satisfy $g(0) = 1$ and $g'(0) = 1$. We round off our figure at $x = -2$ to satisfy $g'(-2) = 0$. As $x \to -5^+$, $y \to \infty$, so we draw a vertical asymptote at $x = -5$. As $x \to 5^-$, $y \to 3$, so we draw a dot at $(5, 3)$ [the dot could be open or closed].

42. We begin by drawing an odd function (symmetric with respect to the origin) through the origin with slope -2 to satisfy $f'(0) = -2$. Now draw a curve starting at $x = 1$ and increasing without bound as $x \to 2^-$ since $\lim_{x\to 2^-} f(x) = \infty$. Lastly, reflect the last curve through the origin (rotate 180°) since f is an odd function.

43. By Definition 4, $\lim_{h\to 0} \dfrac{\sqrt{9+h} - 3}{h} = f'(9)$, where $f(x) = \sqrt{x}$ and $a = 9$.

44. By Definition 4, $\lim_{h\to 0} \dfrac{e^{-2+h} - e^{-2}}{h} = f'(-2)$, where $f(x) = e^x$ and $a = -2$.

45. By Equation 5, $\lim_{x\to 2} \dfrac{x^6 - 64}{x - 2} = f'(2)$, where $f(x) = x^6$ and $a = 2$.

46. By Equation 5, $\lim_{x\to 1/4} \dfrac{\frac{1}{x} - 4}{x - \frac{1}{4}} = f'(4)$, where $f(x) = \dfrac{1}{x}$ and $a = \dfrac{1}{4}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. By Definition 4, $\lim\limits_{h\to 0} \dfrac{\tan\left(\dfrac{\pi}{4}+h\right)-1}{h} = f'\left(\dfrac{\pi}{4}\right)$, where $f(x)=\tan x$ and $a=\dfrac{\pi}{4}$.

48. By Equation 5, $\lim\limits_{\theta\to\pi/6} \dfrac{\sin\theta-\frac{1}{2}}{\theta-\dfrac{\pi}{6}} = f'\left(\dfrac{\pi}{6}\right)$, where $f(\theta)=\sin\theta$ and $a=\dfrac{\pi}{6}$.

49. (a) (i) $\dfrac{\Delta C}{\Delta x} = \dfrac{C(105)-C(100)}{105-100} = \dfrac{6601.25-6500}{5} = \$20.25/\text{unit}.$

(ii) $\dfrac{\Delta C}{\Delta x} = \dfrac{C(101)-C(100)}{101-100} = \dfrac{6520.05-6500}{1} = \$20.05/\text{unit}.$

(b) $$\frac{C(100+h)-C(100)}{h} = \frac{\left[5000+10(100+h)+0.05(100+h)^2\right]-6500}{h} = \frac{20h+0.05h^2}{h}$$
$$= 20+0.05h,\ h\neq 0$$

So the instantaneous rate of change is $\lim\limits_{h\to 0}\dfrac{C(100+h)-C(100)}{h} = \lim\limits_{h\to 0}(20+0.05h) = \$20/\text{unit}.$

50. (a) $H'(58)$ is the rate at which the daily heating cost changes with respect to temperature when the outside temperature is $58\,^\circ\text{F}$. The units are dollars/$^\circ$F.

(b) If the outside temperature increases, the building should require less heating, so we would expect $H'(58)$ to be negative.

51. (a) $f'(x)$ is the rate of change of the production cost with respect to the number of ounces of gold produced. Its units are dollars per ounce.

(b) After 800 ounces of gold have been produced, the rate at which the production cost is increasing is \$17/ounce. So the cost of producing the 800th (or 801st) ounce is about \$17.

(c) In the short term, the values of $f'(x)$ will decrease because more efficient use is made of start-up costs as x increases. But eventually $f'(x)$ might increase due to large-scale operations.

52. (a) $f'(8)$ is the rate of change of the quantity of coffee sold with respect to the price per pound when the price is \$8 per pound. The units for $f'(8)$ are pounds/(dollars/pound).

(b) $f'(8)$ is negative since the quantity of coffee sold will decrease as the price charged for it increases. People are generally less willing to buy a product when its price increases.

53. (a) $S'(T)$ is the rate at which the oxygen solubility changes with respect to the water temperature. Its units are $(\text{mg/L})/^\circ\text{C}$.

(b) For $T=16^\circ\text{C}$, it appears that the tangent line to the curve goes through the points $(0,14)$ and $(32,6)$. So $S'(16)\approx\dfrac{6-14}{32-0}=-\dfrac{8}{32}=-0.25\ (\text{mg/L})/^\circ\text{C}$. This means that as the temperature increases past 16°C, the oxygen solubility is decreasing at a rate of $0.25\ (\text{mg/L})/^\circ\text{C}$.

54. (a) $S'(T)$ is the rate of change of the maximum sustainable speed of Coho salmon with respect to the temperature. Its units are $(\text{cm/s})/^\circ\text{C}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) For $T = 15°\text{C}$, it appears the tangent line to the curve goes through the points $(10, 25)$ and $(20, 32)$. So $S'(15) \approx \dfrac{32-25}{20-10} = 0.7\ (\text{cm/s})/°\text{C}$. This tells us that at $T = 15°\text{C}$, the maximum sustainable speed of Coho salmon is changing at a rate of $0.7\ (\text{cm/s})/°\text{C}$. In a similar fashion for $T = 25°\text{C}$, we can use the points $(20, 35)$ and $(25, 25)$ to obtain $S'(25) \approx \dfrac{25-35}{25-20} = -2\ (\text{cm/s})/°\text{C}$. As it gets warmer than $20°\text{C}$, the maximum sustainable speed decreases rapidly.

55. (a) (i) $[1.0, 2.0]$: $\dfrac{C(2)-C(1)}{2-1} = \dfrac{0.018-0.033}{1} = -0.015\ \dfrac{\text{g/dL}}{\text{h}}$

(ii) $[1.5, 2.0]$: $\dfrac{C(2)-C(1.5)}{2-1.5} = \dfrac{0.018-0.024}{0.5} = \dfrac{-0.006}{0.5} = -0.012\ \dfrac{\text{g/dL}}{\text{h}}$

(iii) $[2.0, 2.5]$: $\dfrac{C(2.5)-C(2)}{2.5-2} = \dfrac{0.012-0.018}{0.5} = \dfrac{-0.006}{0.5} = -0.012\ \dfrac{\text{g/dL}}{\text{h}}$

(iv) $[2.0, 3.0]$: $\dfrac{C(3)-C(2)}{3-2} = \dfrac{0.007-0.018}{1} = -0.011\ \dfrac{\text{g/dL}}{\text{h}}$

(b) We estimate the instantaneous rate of change at $t = 2$ by averaging the average rates of change for $[1.5, 2.0]$ and $[2.0, 2.5]$: $\dfrac{-0.012+(-0.012)}{2} = -0.012\ \dfrac{\text{g/dL}}{\text{h}}$. After two hours, the BAC is decreasing at a rate of $0.012\ \dfrac{\text{g/dL}}{\text{h}}$.

56. (a) (i) $[2008, 2010]$: $\dfrac{N(2010)-N(2008)}{2010-2008} = \dfrac{16{,}858-16{,}680}{2} = \dfrac{178}{2} = 89$ locations/year.

(ii) $[2010, 2012]$: $\dfrac{N(2012)-N(2010)}{2012-2010} = \dfrac{18{,}066-16{,}858}{2} = \dfrac{1208}{2} = 604$ locations/year.

The rate of growth increased over the period 2008 to 2012.

(b) Averaging the values from parts (i) and (ii) of (a), we have $\dfrac{89+604}{2} = \dfrac{693}{2} = 346.5$ locations/year.

(c) We plot the function N and estimate the slope of the tangent line at $x = 2010$. The tangent segment has endpoints $(2008, 16{,}250)$ and $(2012, 17{,}500)$. An estimate of the instantaneous rate of growth in 2010 is $\dfrac{17{,}500-16{,}250}{2012-2008} = \dfrac{1250}{4} = 312.5$ locations/year.

57. Since $f(x) = x\sin(1/x)$ when $x \neq 0$ and $f(0) = 0$, we have $f'(0) = \lim\limits_{h\to 0} \dfrac{f(0+h)-f(0)}{h} = \lim\limits_{h\to 0} \dfrac{h\sin(1/h)-0}{h} = \lim\limits_{h\to 0} \sin(1/h)$. This limit does not exist since $\sin(1/h)$ takes the values -1 and 1 on any interval containing 0. (Compare with Example 2.2.5.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

58. Since $f(x) = x^2 \sin(1/x)$ when $x \neq 0$ and $f(0) = 0$, we have

$f'(0) = \lim\limits_{h \to 0} \dfrac{f(0+h) - f(0)}{h} = \lim\limits_{h \to 0} \dfrac{h^2 \sin(1/h) - 0}{h} = \lim\limits_{h \to 0} h \sin(1/h)$. Since $-1 \leq \sin \dfrac{1}{h} \leq 1$, we have

$-|h| \leq |h| \sin \dfrac{1}{h} \leq |h| \quad \Rightarrow \quad -|h| \leq h \sin \dfrac{1}{h} \leq |h|$. Because $\lim\limits_{h \to 0} (-|h|) = 0$ and $\lim\limits_{h \to 0} |h| = 0$, we know that

$\lim\limits_{h \to 0} \left(h \sin \dfrac{1}{h} \right) = 0$ by the Squeeze Theorem. Thus, $f'(0) = 0$.

59. (a) The slope at the origin appears to be 1.

(b) The slope at the origin still appears to be 1.

(c) Yes, the slope at the origin now appears to be 0.

60. (a) The symmetric difference quotient on $[2004, 2012]$ is (with $a = 2008$ and $d = 4$)

$$\begin{aligned} \frac{f(2008+4) - f(2008-4)}{2(4)} &= \frac{f(2012) - f(2004)}{8} \\ &= \frac{16{,}432.7 - 7596.1}{8} \\ &= 1104.575 \approx 1105 \text{ billion dollars per year} \end{aligned}$$

This result agrees with the estimate for $D'(2008)$ computed in the example.

(b) Averaging the average rates of change of f over the intervals $[a-d, a]$ and $[a, a+d]$ gives

$$\begin{aligned} \frac{\dfrac{f(a) - f(a-d)}{a-(a-d)} + \dfrac{f(a+d) - f(a)}{(a+d)-a}}{2} &= \frac{\dfrac{f(a) - f(a-d)}{d} + \dfrac{f(a+d) - f(a)}{d}}{2} \\ &= \frac{f(a+d) - f(a-d)}{2d} \end{aligned}$$

which is the symmeric difference quotient.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) For $f(x) = x^3 - 2x^2 + 2$, $a = 1$, and $d = 0.4$, we have

$$f'(1) \approx \frac{f(1+0.4) - f(1-0.4)}{2(0.4)} = \frac{f(1.4) - f(0.6)}{0.8}$$
$$= \frac{0.824 - 1.496}{0.8} = -0.84$$

On the graph, (i)–(v) correspond to:

(i) $f(x) = x^3 - 2x^2 + 2$

(ii) secant line corresponding to average rate of change over $[1 - 0.4, 1] = [0.6, 1]$

(iii) secant line corresponding to average rate of change over $[1, 1 + 0.4] = [1, 1.4]$

(iv) secant line corresponding to average rate of change over $[1 - 0.4, 1 + 0.4] = [0.6, 1.4]$

(v) tangent line at $x = 1$

The secant line corresponding to the average rate of change over $[0.6, 1.4]$ — that is, graph (iv) — appears to have slope closest to that of the tangent line at $x = 1$.

2.8 The Derivative as a Function

1. We estimate the slopes of tangent lines on the graph of f to determine the derivative approximations that follow. Your answers may vary depending on your estimates.

(a) $f'(0) = \frac{1}{2}$ (b) $f'(1) \approx 0$

(c) $f'(2) \approx -1$ (d) $f'(3) \approx -\frac{3}{2}$

(e) $f'(4) \approx -1$ (f) $f'(5) \approx 0$

(g) $f'(6) \approx 1$ (h) $f'(7) \approx 1$

2. We estimate the slopes of tangent lines on the graph of f to determine the derivative approximations that follow. Your answers may vary depending on your estimates.

(a) $f'(-3) \approx -1$ (b) $f'(-2) \approx 0$

(c) $f'(-1) \approx \frac{1}{2}$ (d) $f'(0) \approx \frac{3}{2}$

(e) $f'(1) \approx 3$ (f) $f'(2) \approx 0$

(g) $f'(3) \approx -\frac{3}{2}$

3. (a)′ = II, since from left to right, the slopes of the tangents to graph (a) start out negative, become 0, then positive, then 0, then negative again. The actual function values in graph II follow the same pattern.

(b)′ = IV, since from left to right, the slopes of the tangents to graph (b) start out at a fixed positive quantity, then suddenly become negative, then positive again. The discontinuities in graph IV indicate sudden changes in the slopes of the tangents.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c)$'$ = I, since the slopes of the tangents to graph (c) are negative for $x < 0$ and positive for $x > 0$, as are the function values of graph I.

(d)$'$ = III, since from left to right, the slopes of the tangents to graph (d) are positive, then 0, then negative, then 0, then positive, then 0, then negative again, and the function values in graph III follow the same pattern.

Hints for Exercises 4–11: First plot x-intercepts on the graph of f' for any horizontal tangents on the graph of f. Look for any corners on the graph of f—there will be a discontinuity on the graph of f'. On any interval where f has a tangent with positive (or negative) slope, the graph of f' will be positive (or negative). If the graph of the function is linear, the graph of f' will be a horizontal line.

4.

5.

6.

7.

8.

9.

10.

11.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. The slopes of the tangent lines on the graph of $y = P(t)$ are always positive, so the y-values of $y = P'(t)$ are always positive. These values start out relatively small and keep increasing, reaching a maximum at about $t = 6$. Then the y-values of $y = P'(t)$ decrease and get close to zero. The graph of P' tells us that the yeast culture grows most rapidly after 6 hours and then the growth rate declines.

13. (a) $C'(t)$ is the instantaneous rate of change of percentage of full capacity with respect to elapsed time in hours.

(b) The graph of $C'(t)$ tells us that the rate of change of percentage of full capacity is decreasing and approaching 0.

14. (a) $F'(v)$ is the instantaneous rate of change of fuel economy with respect to speed.

(b) Graphs will vary depending on estimates of F', but will change from positive to negative at about $v = 50$.

(c) To save on gas, drive at the speed where F is a maximum and F' is 0, which is about 50 mi/h.

15. It appears that there are horizontal tangents on the graph of f for $t = 2$ and for $t \approx 7.5$. Thus, there are zeros for those values of t on the graph of f'. The derivative is negative for values of t between 0 and 2 and for values of t between approximately 7.5 and 12. The value of $f'(t)$ appears to be largest at $t \approx 5.25$.

16.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17.

The slope at 0 appears to be 1 and the slope at 1 appears to be 2.7. As x decreases, the slope gets closer to 0. Since the graphs are so similar, we might guess that $f'(x) = e^x$.

18.

As x increases toward 1, $f'(x)$ decreases from very large numbers to 1. As x becomes large, $f'(x)$ gets closer to 0. As a guess, $f'(x) = 1/x^2$ or $f'(x) = 1/x$ makes sense.

19. (a) By zooming in, we estimate that $f'(0) = 0$, $f'\left(\frac{1}{2}\right) = 1$, $f'(1) = 2$, and $f'(2) = 4$.

(b) By symmetry, $f'(-x) = -f'(x)$. So $f'\left(-\frac{1}{2}\right) = -1$, $f'(-1) = -2$, and $f'(-2) = -4$.

(c) It appears that $f'(x)$ is twice the value of x, so we guess that $f'(x) = 2x$.

(d) $$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \to 0} \frac{(x+h)^2 - x^2}{h}$$

$$= \lim_{h \to 0} \frac{\left(x^2 + 2hx + h^2\right) - x^2}{h} = \lim_{h \to 0} \frac{2hx + h^2}{h} = \lim_{h \to 0} \frac{h(2x+h)}{h} = \lim_{h \to 0} (2x + h) = 2x$$

20. (a) By zooming in, we estimate that $f'(0) = 0$, $f'\left(\frac{1}{2}\right) \approx 0.75$, $f'(1) \approx 3$, $f'(2) \approx 12$, and $f'(3) \approx 27$.

(b) By symmetry, $f'(-x) = f'(x)$. So $f'\left(-\frac{1}{2}\right) \approx 0.75$, $f'(-1) \approx 3$, $f'(-2) \approx 12$, and $f'(-3) \approx 27$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c)

(d) Since $f'(0) = 0$, it appears that f' may have the form $f'(x) = ax^2$.

Using $f'(1) = 3$, we have $a = 3$, so $f'(x) = 3x^2$.

(e) $f'(x) = \lim\limits_{h\to 0} \dfrac{f(x+h) - f(x)}{h} = \lim\limits_{h\to 0} \dfrac{(x+h)^3 - x^3}{h} = \lim\limits_{h\to 0} \dfrac{(x^3 + 3x^2h + 3xh^2 + h^3) - x^3}{h}$

$= \lim\limits_{h\to 0} \dfrac{3x^2h + 3xh^2 + h^3}{h} = \lim\limits_{h\to 0} \dfrac{h(3x^2 + 3xh + h^2)}{h} = \lim\limits_{h\to 0}(3x^2 + 3xh + h^2) = 3x^2$

21. $f'(x) = \lim\limits_{h\to 0} \dfrac{f(x+h) - f(x)}{h} = \lim\limits_{h\to 0} \dfrac{[3(x+h) - 8] - (3x - 8)}{h} = \lim\limits_{h\to 0} \dfrac{3x + 3h - 8 - 3x + 8}{h}$

$= \lim\limits_{h\to 0} \dfrac{3h}{h} = \lim\limits_{h\to 0} 3 = 3$

Domain of f = domain of $f' = \mathbb{R}$.

22. $f'(x) = \lim\limits_{h\to 0} \dfrac{f(x+h) - f(x)}{h} = \lim\limits_{h\to 0} \dfrac{[m(x+h) + b] - (mx + b)}{h} = \lim\limits_{h\to 0} \dfrac{mx + mh + b - mx - b}{h}$

$= \lim\limits_{h\to 0} \dfrac{mh}{h} = \lim\limits_{h\to 0} m = m$

Domain of f = domain of $f' = \mathbb{R}$.

23. $f'(t) = \lim\limits_{h\to 0} \dfrac{f(t+h) - f(t)}{h} = \lim\limits_{h\to 0} \dfrac{\left[2.5(t+h)^2 + 6(t+h)\right] - \left(2.5t^2 + 6t\right)}{h}$

$= \lim\limits_{h\to 0} \dfrac{2.5(t^2 + 2th + h^2) + 6t + 6h - 2.5t^2 - 6t}{h} = \lim\limits_{h\to 0} \dfrac{2.5t^2 + 5th + 2.5h^2 + 6h - 2.5t^2}{h}$

$= \lim\limits_{h\to 0} \dfrac{5th + 2.5h^2 + 6h}{h} = \lim\limits_{h\to 0} \dfrac{h\,(5t + 2.5h + 6)}{h} = \lim\limits_{h\to 0} (5t + 2.5h + 6)$

$= 5t + 6$

Domain of f = domain of $f' = \mathbb{R}$.

24. $f'(x) = \lim\limits_{h\to 0} \dfrac{f(x+h) - f(x)}{h} = \lim\limits_{h\to 0} \dfrac{\left[4 + 8(x+h) - 5(x+h)^2\right] - (4 + 8x - 5x^2)}{h}$

$= \lim\limits_{h\to 0} \dfrac{4 + 8x + 8h - 5(x^2 + 2xh + h^2) - 4 - 8x + 5x^2}{h} = \lim\limits_{h\to 0} \dfrac{8h - 5x^2 - 10xh - 5h^2 + 5x^2}{h}$

$= \lim\limits_{h\to 0} \dfrac{8h - 10xh - 5h^2}{h} = \lim\limits_{h\to 0} \dfrac{h(8 - 10x - 5h)}{h} = \lim\limits_{h\to 0} (8 - 10x - 5h)$

$= 8 - 10x$

Domain of f = domain of $f' = \mathbb{R}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $$A'(p) = \lim_{h\to 0} \frac{A(p+h) - A(p)}{h} = \lim_{h\to 0} \frac{\left[4(p+h)^3 + 3(p+h)\right] - (4p^3 + 3p)}{h}$$
$$= \lim_{h\to 0} \frac{4p^3 + 12p^2h + 12ph^2 + 4h^3 + 3p + 3h - 4p^3 - 3p}{h} = \lim_{h\to 0} \frac{12p^2h + 12ph^2 + 4h^3 + 3h}{h}$$
$$= \lim_{h\to 0} \frac{h(12p^2 + 12ph + 4h^2 + 3)}{h} = \lim_{h\to 0} \left(12p^2 + 12ph + 4h^2 + 3\right) = 12p^2 + 3$$

Domain of A = Domain of $A' = \mathbb{R}$.

26. $$F'(t) = \lim_{h\to 0} \frac{F(t+h) - F(t)}{h} = \lim_{h\to 0} \frac{\left[(t+h)^3 - 5(t+h) + 1\right] - (t^3 - 5t + 1)}{h}$$
$$= \lim_{h\to 0} \frac{t^3 + 3t^2h + 3th^2 + h^3 - 5t - 5h + 1 - t^3 + 5t - 1}{h} = \lim_{h\to 0} \frac{3t^2h + 3th^2 + h^3 - 5h}{h}$$
$$= \lim_{h\to 0} \frac{h(3t^2 + 3th + h^2 - 5)}{h} = \lim_{h\to 0} \left(3t^2 + 3th + h^2 - 5\right) = 3t^2 - 5$$

Domain of F = Domain of $F' = \mathbb{R}$.

27. $$f'(x) = \lim_{h\to 0} \frac{f(x+h) - f(x)}{h} = \lim_{h\to 0} \frac{\dfrac{1}{(x+h)^2 - 4} - \dfrac{1}{x^2 - 4}}{h} = \lim_{h\to 0} \frac{\dfrac{(x^2 - 4) - \left[(x+h)^2 - 4\right]}{\left[(x+h)^2 - 4\right](x^2 - 4)}}{h}$$
$$= \lim_{h\to 0} \frac{(x^2 - 4) - \left(x^2 + 2xh + h^2 - 4\right)}{h\left[(x+h)^2 - 4\right](x^2 - 4)} = \lim_{h\to 0} \frac{x^2 - 4 - x^2 - 2xh - h^2 + 4}{h[(x+h)^2 - 4](x^2 - 4)} = \lim_{h\to 0} \frac{-2xh - h^2}{h[(x+h)^2 - 4](x^2 - 4)}$$
$$= \lim_{h\to 0} \frac{h(-2x - h)}{h[(x+h)^2 - 4](x^2 - 4)} = \lim_{h\to 0} \frac{-2x - h}{[(x+h)^2 - 4](x^2 - 4)} = \frac{-2x}{(x^2 - 4)(x^2 - 4)} = -\frac{2x}{(x^2 - 4)^2}$$

Domain of f = Domain of $f' = (-\infty, -2) \cup (-2, 2) \cup (2, \infty)$.

28. $$F'(v) = \lim_{h\to 0} \frac{F(v+h) - F(v)}{h} = \lim_{h\to 0} \frac{\dfrac{v+h}{(v+h)+2} - \dfrac{v}{v+2}}{h} = \lim_{h\to 0} \frac{\dfrac{(v+h)(v+2) - v\left[(v+h)+2\right]}{\left[(v+h)+2\right](v+2)}}{h}$$
$$= \lim_{h\to 0} \frac{v^2 + 2v + vh + 2h - v^2 - vh - 2v}{h[(v+h)+2](v+2)} = \lim_{h\to 0} \frac{2h}{h[(v+h)+2](v+2)} = \lim_{h\to 0} \frac{2}{[(v+h)+2](v+2)}$$
$$= \frac{2}{(v+2)(v+2)} = \frac{2}{(v+2)^2}$$

Domain of F = Domain of $f' = (-\infty, -2) \cup (-2, \infty)$.

29. $$g'(u) = \lim_{h\to 0} \frac{g(u+h) - g(u)}{h} = \lim_{h\to 0} \frac{\dfrac{(u+h)+1}{4(u+h)-1} - \dfrac{u+1}{4u-1}}{h} = \lim_{h\to 0} \frac{\dfrac{\left[(u+h)+1\right](4u-1) - (u+1)\left[4(u+h)-1\right]}{\left[4(u+h)-1\right](4u-1)}}{h}$$
$$= \lim_{h\to 0} \frac{\dfrac{(u+h+1)(4u-1) - (u+1)(4u+4h-1)}{\left[4(u+h)-1\right](4u-1)}}{h}$$
$$= \lim_{h\to 0} \frac{4u^2 + 4uh + 4u - u - h - 1 - 4u^2 - 4uh + u - 4u - 4h + 1}{h\left[4(u+h)-1\right](4u-1)}$$
$$= \lim_{h\to 0} \frac{-5h}{h\left[4(u+h)-1\right](4u-1)} = \lim_{h\to 0} \frac{-5}{\left[4(u+h)-1\right](4u-1)} = \frac{-5}{(4u-1)(4u-1)} = -\frac{5}{(4u-1)^2}$$

Domain of g = Domain of $g' = \left(-\infty, \frac{1}{4}\right) \cup \left(\frac{1}{4}, \infty\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $f'(x) = \lim\limits_{h\to 0} \dfrac{f(x+h)-f(x)}{h} = \lim\limits_{h\to 0} \dfrac{(x+h)^4 - x^4}{h} = \lim\limits_{h\to 0} \dfrac{(x^4+4x^3h+6x^2h^2+4xh^3+h^4)-x^4}{h}$

$$= \lim_{h\to 0} \frac{4x^3h+6x^2h^2+4xh^3+h^4}{h} = \lim_{h\to 0} \frac{h(4x^3+6x^2h+4xh^2+h^3)}{h}$$

$$= \lim_{h\to 0} (4x^3+6x^2h+4xh^2+h^3) = 4x^3$$

Domain of f = domain of $f' = \mathbb{R}$.

31. $f'(x) = \lim\limits_{h\to 0} \dfrac{f(x+h)-f(x)}{h} = \lim\limits_{h\to 0} \dfrac{\dfrac{1}{\sqrt{1+(x+h)}} - \dfrac{1}{\sqrt{1+x}}}{h}$

$$= \lim_{h\to 0} \frac{\dfrac{1}{\sqrt{1+(x+h)}} - \dfrac{1}{\sqrt{1+x}}}{h} \cdot \frac{\sqrt{1+(x+h)}\sqrt{1+x}}{\sqrt{1+(x+h)}\sqrt{1+x}}$$

$$= \lim_{h\to 0} \frac{\sqrt{1+x}-\sqrt{1+(x+h)}}{h\sqrt{1+(x+h)}\sqrt{1+x}} \cdot \frac{\sqrt{1+x}+\sqrt{1+(x+h)}}{\sqrt{1+x}+\sqrt{1+(x+h)}}$$

$$= \lim_{h\to 0} \frac{(1+x)-[1+(x+h)]}{h\sqrt{1+(x+h)}\sqrt{1+x}\left(\sqrt{1+x}+\sqrt{1+(x+h)}\right)}$$

$$= \lim_{h\to 0} \frac{-h}{h\sqrt{1+x+h}\sqrt{1+x}\left(\sqrt{1+x}+\sqrt{1+x+h}\right)} = \lim_{h\to 0} \frac{-1}{\sqrt{1+x+h}\sqrt{1+x}\left(\sqrt{1+x}+\sqrt{1+x+h}\right)}$$

$$= \frac{-1}{\sqrt{1+x}\sqrt{1+x}\left(\sqrt{1+x}+\sqrt{1+x}\right)} = \frac{-1}{(1+x)\left(2\sqrt{1+x}\right)} = -\frac{1}{2(1+x)^{3/2}}$$

Domain of f = Domain of $f' = (-1, \infty)$.

32. $g'(x) = \lim\limits_{h\to 0} \dfrac{g(x+h)-g(x)}{h} = \lim\limits_{h\to 0} \dfrac{\dfrac{1}{1+\sqrt{x+h}} - \dfrac{1}{1+\sqrt{x}}}{h} = \lim\limits_{h\to 0} \dfrac{\dfrac{\left(1+\sqrt{x}\right)-\left(1+\sqrt{x+h}\right)}{\left(1+\sqrt{x+h}\right)\left(1+\sqrt{x}\right)}}{h}$

$$= \lim_{h\to 0} \frac{1+\sqrt{x}-1-\sqrt{x+h}}{h\left(1+\sqrt{x+h}\right)\left(1+\sqrt{x}\right)} = \lim_{h\to 0} \frac{\sqrt{x}-\sqrt{x+h}}{h\left(1+\sqrt{x+h}\right)\left(1+\sqrt{x}\right)} \cdot \frac{\sqrt{x}+\sqrt{x+h}}{\sqrt{x}+\sqrt{x+h}}$$

$$= \lim_{h\to 0} \frac{x-(x+h)}{h\left(1+\sqrt{x+h}\right)\left(1+\sqrt{x}\right)\left(\sqrt{x}+\sqrt{x+h}\right)}$$

$$= \lim_{h\to 0} \frac{-h}{h\left(1+\sqrt{x+h}\right)\left(1+\sqrt{x}\right)\left(\sqrt{x}+\sqrt{x+h}\right)}$$

$$= \lim_{h\to 0} \frac{-1}{\left(1+\sqrt{x+h}\right)\left(1+\sqrt{x}\right)\left(\sqrt{x}+\sqrt{x+h}\right)}$$

$$= \frac{-1}{\left(1+\sqrt{x}\right)\left(1+\sqrt{x}\right)\left(\sqrt{x}+\sqrt{x}\right)} = \frac{-1}{\left(1+\sqrt{x}\right)^2 \cdot 2\sqrt{x}} = -\frac{1}{2\sqrt{x}\left(1+\sqrt{x}\right)^2}$$

Domain of $g = [0, \infty)$, domain of $g' = (0, \infty)$.

33. (a)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Note that the third graph in part (a) generally has small positive values for its slope, f'; but as $x \to -3^+$, $f' \to \infty$. See the graph in part (d).

(c) $$f'(x) = \lim_{h\to 0} \frac{f(x+h)-f(x)}{h} = \lim_{h\to 0} \frac{1+\sqrt{(x+h)+3}-\left(1+\sqrt{x+3}\right)}{h}$$

$$= \lim_{h\to 0} \frac{\sqrt{(x+h)+3}-\sqrt{x+3}}{h} = \lim_{h\to 0} \frac{\sqrt{(x+h)+3}-\sqrt{x+3}}{h}\left[\frac{\sqrt{(x+h)+3}+\sqrt{x+3}}{\sqrt{(x+h)+3}+\sqrt{x+3}}\right]$$

$$= \lim_{h\to 0} \frac{[(x+h)+3]-(x+3)}{h\sqrt{(x+h)+3}+\sqrt{x+3}} = \lim_{h\to 0} \frac{x+h+3-x-3}{h\left(\sqrt{(x+h)+3}+\sqrt{x+3}\right)}$$

$$= \lim_{h\to 0} \frac{h}{h\left(\sqrt{(x+h)+3}+\sqrt{x+3}\right)} = \lim_{h\to 0} \frac{1}{\sqrt{(x+h)+3}+\sqrt{x+3}} = \frac{1}{2\sqrt{x+3}}$$

Domain of $f = [-3, \infty)$, Domain of $f' = (-3, \infty)$.

(d)

34. (a) $$f'(x) = \lim_{h\to 0} \frac{f(x+h)-f(x)}{h} = \lim_{h\to 0} \frac{[(x+h)+1/(x+h)]-(x+1/x)}{h} = \lim_{h\to 0} \frac{\dfrac{(x+h)^2+1}{x+h} - \dfrac{x^2+1}{x}}{h}$$

$$= \lim_{h\to 0} \frac{x[(x+h)^2+1]-(x+h)(x^2+1)}{h(x+h)x} = \lim_{h\to 0} \frac{(x^3+2hx^2+xh^2+x)-(x^3+x+hx^2+h)}{h(x+h)x}$$

$$= \lim_{h\to 0} \frac{hx^2+xh^2-h}{h(x+h)x} = \lim_{h\to 0} \frac{h(x^2+xh-1)}{h(x+h)x} = \lim_{h\to 0} \frac{x^2+xh-1}{(x+h)x} = \frac{x^2-1}{x^2}, \text{ or } 1-\frac{1}{x^2}$$

(b) Notice that $f'(x) = 0$ when f has a horizontal tangent, $f'(x)$ is positive when the tangents have positive slope, and $f'(x)$ is negative when the tangents have negative slope. Both functions are discontinuous at $x = 0$.

35. (a) $$f'(x) = \lim_{h\to 0} \frac{f(x+h)-f(x)}{h} = \lim_{h\to 0} \frac{[(x+h)^4+2(x+h)]-(x^4+2x)}{h}$$

$$= \lim_{h\to 0} \frac{x^4+4x^3h+6x^2h^2+4xh^3+h^4+2x+2h-x^4-2x}{h}$$

$$= \lim_{h\to 0} \frac{4x^3h+6x^2h^2+4xh^3+h^4+2h}{h} = \lim_{h\to 0} \frac{h(4x^3+6x^2h+4xh^2+h^3+2)}{h}$$

$$= \lim_{h\to 0}(4x^3+6x^2h+4xh^2+h^3+2) = 4x^3+2$$

(b) Notice that $f'(x) = 0$ when f has a horizontal tangent, $f'(x)$ is positive when the tangents have positive slope, and $f'(x)$ is negative when the tangents have negative slope.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. (a) $N'(t)$ is the rate at which the number of minimally invasive cosmetic surgery procedures performed in the United States is changing with respect to time. Its units are thousands of surgeries per year.

(b) To find $N'(t)$, we use $\lim\limits_{h\to 0} \dfrac{N(t+h)-N(t)}{(t+h)-t} \approx \dfrac{N(t+h)-N(t)}{h}$ for small values of h.

For 2000: $N'(2000) \approx \dfrac{N(2002)-N(2000)}{2002-2000} = \dfrac{4897-5500}{2} = -301.5$

For 2002: We estimate $N'(2002)$ by using $h=-2$ and $h=2$, and then average the two results to obtain a final estimate.

$h=-2 \quad\Rightarrow\quad N'(2002) \approx \dfrac{N(2000)-N(2002)}{2000-2002} = \dfrac{5500-4897}{-2} = -301.5$

$h=2 \quad\Rightarrow\quad N'(2002) \approx \dfrac{N(2004)-N(2002)}{2004-2002} = \dfrac{7470-4897}{2} = 1286.5$

So we estimate that $N'(2002) \approx \frac{1}{2}[-301.5+1286.5] = 492.5$.

t	2000	2002	2004	2006	2008	2010	2012	2014
$N'(t)$	-301.5	492.5	1060.25	856.75	605.75	534.5	596	455

(c)

(d) We could get more accurate values for $N'(t)$ by obtaining data for more values of t.

37. As in Exercise 36, we use one-sided difference quotients for the first and last values, and average two difference quotients for all other values.

t	14	21	28	35	42	49
$H(t)$	41	54	64	72	78	83
$H'(t)$	$\frac{13}{7}$	$\frac{23}{14}$	$\frac{18}{14}$	$\frac{14}{14}$	$\frac{11}{14}$	$\frac{5}{7}$

38. As in Exercise 36, we use one-sided difference quotients for the first and last values, and average two difference quotients for all other values. The units for $W'(x)$ are grams per degree (g/°C).

x	15.5	17.7	20.0	22.4	24.4
$W(x)$	37.2	31.0	19.8	9.7	-9.8
$W'(x)$	-2.82	-3.87	-4.53	-6.73	-9.75

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. (a) dP/dt is the rate at which the percentage of the city's electrical power produced by solar panels changes with respect to time t, measured in percentage points per year.

(b) 2 years after January 1, 2020 (January 1, 2022), the percentage of electrical power produced by solar panels was increasing at a rate of 3.5 percentage points per year.

40. dN/dp is the rate at which the number of people who travel by car to another state for a vacation changes with respect to the price of gasoline. If the price of gasoline goes up, we would expect fewer people to travel, so we would expect dN/dp to be negative.

41. f is not differentiable at $x = -4$, because the graph has a corner there, and at $x = 0$, because there is a discontinuity there.

42. f is not differentiable at $x = -1$, because there is a discontinuity there, and at $x = 2$, because the graph has a corner there.

43. f is not differentiable at $x = 1$, because f is not defined there, and at $x = 5$, because the graph has a vertical tangent there.

44. f is not differentiable at $x = -2$ and $x = 3$, because the graph has corners there, and at $x = 1$, because there is a discontinuity there.

45. As we zoom in toward $(-1, 0)$, the curve appears more and more like a straight line, so $f(x) = x + \sqrt{|x|}$ is differentiable at $x = -1$. But no matter how much we zoom in toward the origin, the curve doesn't straighten out—we can't eliminate the sharp point (a cusp). So f is not differentiable at $x = 0$.

46. As we zoom in toward $(0, 1)$, the curve appears more and more like a straight line, so $g(x) = (x^2 - 1)^{2/3}$ is differentiable at $x = 0$. But no matter how much we zoom in toward $(1, 0)$ or $(-1, 0)$, the curve doesn't straighten out—we can't eliminate the sharp point (a cusp). So g is not differentiable at $x = \pm 1$.

47. Call the curve with the positive y-intercept g and the other curve h. Notice that g has a maximum (horizontal tangent) at $x = 0$, but $h \neq 0$, so h cannot be the derivative of g. Also notice that where g is positive, h is increasing. Thus, $h = f$ and $g = f'$. Now $f'(-1)$ is negative since f' is below the x-axis there and $f''(1)$ is positive since f is concave upward at $x = 1$. Therefore, $f''(1)$ is greater than $f'(-1)$.

48. Call the curve with the smallest positive x-intercept g and the other curve h. Notice that where g is positive in the first quadrant, h is increasing. Thus, $h = f$ and $g = f'$. Now $f'(-1)$ is positive since f' is above the x-axis there and $f''(1)$ appears to be zero since f has an inflection point at $x = 1$. Therefore, $f'(1)$ is greater than $f''(-1)$.

49. $a = f$, $b = f'$, $c = f''$. We can see this because where a has a horizontal tangent, $b = 0$, and where b has a horizontal tangent, $c = 0$. We can immediately see that c can be neither f nor f', since at the points where c has a horizontal tangent, neither a nor b is equal to 0.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50. Where d has horizontal tangents, only c is 0, so $d' = c$. c has negative tangents for $x < 0$ and b is the only graph that is negative for $x < 0$, so $c' = b$. b has positive tangents on $\mathbb{R}$ (except at $x = 0$), and the only graph that is positive on the same domain is a, so $b' = a$. We conclude that $d = f$, $c = f'$, $b = f''$, and $a = f'''$.

51. We can immediately see that a is the graph of the acceleration function, since at the points where a has a horizontal tangent, neither c nor b is equal to 0. Next, we note that $a = 0$ at the point where b has a horizontal tangent, so b must be the graph of the velocity function, and hence, $b' = a$. We conclude that c is the graph of the position function.

52. a must be the jerk since none of the graphs are 0 at its high and low points. a is 0 where b has a maximum, so $b' = a$. b is 0 where c has a maximum, so $c' = b$. We conclude that d is the position function, c is the velocity, b is the acceleration, and a is the jerk.

53. $$f'(x) = \lim_{h\to 0} \frac{f(x+h) - f(x)}{h} = \lim_{h\to 0} \frac{[3(x+h)^2 + 2(x+h) + 1] - (3x^2 + 2x + 1)}{h}$$
$$= \lim_{h\to 0} \frac{(3x^2 + 6xh + 3h^2 + 2x + 2h + 1) - (3x^2 + 2x + 1)}{h} = \lim_{h\to 0} \frac{6xh + 3h^2 + 2h}{h}$$
$$= \lim_{h\to 0} \frac{h(6x + 3h + 2)}{h} = \lim_{h\to 0} (6x + 3h + 2) = 6x + 2$$

$$f''(x) = \lim_{h\to 0} \frac{f'(x+h) - f'(x)}{h} = \lim_{h\to 0} \frac{[6(x+h) + 2] - (6x + 2)}{h} = \lim_{h\to 0} \frac{(6x + 6h + 2) - (6x + 2)}{h}$$
$$= \lim_{h\to 0} \frac{6h}{h} = \lim_{h\to 0} 6 = 6$$

We see from the graph that our answers are reasonable because the graph of f' is that of a linear function and the graph of f'' is that of a constant function.

54. $$f'(x) = \lim_{h\to 0} \frac{f(x+h) - f(x)}{h} = \lim_{h\to 0} \frac{[(x+h)^3 - 3(x+h)] - (x^3 - 3x)}{h}$$
$$= \lim_{h\to 0} \frac{(x^3 + 3x^2h + 3xh^2 + h^3 - 3x - 3h) - (x^3 - 3x)}{h} = \lim_{h\to 0} \frac{3x^2h + 3xh^2 + h^3 - 3h}{h}$$
$$= \lim_{h\to 0} \frac{h(3x^2 + 3xh + h^2 - 3)}{h} = \lim_{h\to 0} (3x^2 + 3xh + h^2 - 3) = 3x^2 - 3$$

$$f''(x) = \lim_{h\to 0} \frac{f'(x+h) - f'(x)}{h} = \lim_{h\to 0} \frac{[3(x+h)^2 - 3] - (3x^2 - 3)}{h} = \lim_{h\to 0} \frac{(3x^2 + 6xh + 3h^2 - 3) - (3x^2 - 3)}{h}$$
$$= \lim_{h\to 0} \frac{6xh + 3h^2}{h} = \lim_{h\to 0} \frac{h(6x + 3h)}{h} = \lim_{h\to 0} (6x + 3h) = 6x$$

We see from the graph that our answers are reasonable because the graph of f' is that of an even function (f is an odd function) and the graph of f'' is that of an odd function. Furthermore, $f' = 0$ when f has a horizontal tangent and $f'' = 0$ when f' has a horizontal tangent.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

55. $f'(x) = \lim\limits_{h\to 0} \dfrac{f(x+h)-f(x)}{h} = \lim\limits_{h\to 0} \dfrac{\left[2(x+h)^2-(x+h)^3\right]-(2x^2-x^3)}{h}$

$= \lim\limits_{h\to 0} \dfrac{h(4x+2h-3x^2-3xh-h^2)}{h} = \lim\limits_{h\to 0} (4x+2h-3x^2-3xh-h^2) = 4x-3x^2$

$f''(x) = \lim\limits_{h\to 0} \dfrac{f'(x+h)-f'(x)}{h} = \lim\limits_{h\to 0} \dfrac{\left[4(x+h)-3(x+h)^2\right]-(4x-3x^2)}{h} = \lim\limits_{h\to 0} \dfrac{h(4-6x-3h)}{h}$

$= \lim\limits_{h\to 0}(4-6x-3h) = 4-6x$

$f'''(x) = \lim\limits_{h\to 0} \dfrac{f''(x+h)-f''(x)}{h} = \lim\limits_{h\to 0} \dfrac{[4-6(x+h)]-(4-6x)}{h} = \lim\limits_{h\to 0} \dfrac{-6h}{h} = \lim\limits_{h\to 0} (-6) = -6$

$f^{(4)}(x) = \lim\limits_{h\to 0} \dfrac{f'''(x+h)-f'''(x)}{h} = \lim\limits_{h\to 0} \dfrac{-6-(-6)}{h} = \lim\limits_{h\to 0} \dfrac{0}{h} = \lim\limits_{h\to 0} (0) = 0$

The graphs are consistent with the geometric interpretations of the derivatives because f' has zeros where f has a local minimum and a local maximum, f'' has a zero where f' has a local maximum, and f''' is a constant function equal to the slope of f''.

56. (a) Since we estimate the velocity to be a maximum at $t = 10$, the acceleration is 0 at $t = 10$.

(b) Drawing a tangent line at $t = 10$ on the graph of a, a appears to decrease by $10\ \text{ft/s}^2$ over a period of 20 s.

So at $t = 10$ s, the jerk is approximately $-10/20 = -0.5\ (\text{ft/s}^2)/\text{s}$ or ft/s^3.

57. (a) Note that we have factored $x - a$ as the difference of two cubes in the third step.

$f'(a) = \lim\limits_{x\to a} \dfrac{f(x)-f(a)}{x-a} = \lim\limits_{x\to a} \dfrac{x^{1/3}-a^{1/3}}{x-a} = \lim\limits_{x\to a} \dfrac{x^{1/3}-a^{1/3}}{(x^{1/3}-a^{1/3})(x^{2/3}+x^{1/3}a^{1/3}+a^{2/3})}$

$= \lim\limits_{x\to a} \dfrac{1}{x^{2/3}+x^{1/3}a^{1/3}+a^{2/3}} = \dfrac{1}{3a^{2/3}}$ or $\frac{1}{3}a^{-2/3}$

(b) $f'(0) = \lim\limits_{h\to 0} \dfrac{f(0+h)-f(0)}{h} = \lim\limits_{h\to 0} \dfrac{\sqrt[3]{h}-0}{h} = \lim\limits_{h\to 0} \dfrac{1}{h^{2/3}}$. This function increases without bound, so the limit does not exist, and therefore $f'(0)$ does not exist.

(c) $\lim\limits_{x\to 0} |f'(x)| = \lim\limits_{x\to 0} \dfrac{1}{3x^{2/3}} = \infty$ and f is continuous at $x = 0$ (root function), so f has a vertical tangent at $x = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

58. (a) $g'(0) = \lim\limits_{x\to 0} \dfrac{g(x)-g(0)}{x-0} = \lim\limits_{x\to 0} \dfrac{x^{2/3}-0}{x} = \lim\limits_{x\to 0} \dfrac{1}{x^{1/3}}$, which does not exist.

(b) $g'(a) = \lim\limits_{x\to a} \dfrac{g(x)-g(a)}{x-a} = \lim\limits_{x\to a} \dfrac{x^{2/3}-a^{2/3}}{x-a} = \lim\limits_{x\to a} \dfrac{(x^{1/3}-a^{1/3})(x^{1/3}+a^{1/3})}{(x^{1/3}-a^{1/3})(x^{2/3}+x^{1/3}a^{1/3}+a^{2/3})}$

$= \lim\limits_{x\to a} \dfrac{x^{1/3}+a^{1/3}}{x^{2/3}+x^{1/3}a^{1/3}+a^{2/3}} = \dfrac{2a^{1/3}}{3a^{2/3}} = \dfrac{2}{3a^{1/3}}$ or $\frac{2}{3}a^{-1/3}$

(c) $g(x) = x^{2/3}$ is continuous at $x = 0$ and

$\lim\limits_{x\to 0} |g'(x)| = \lim\limits_{x\to 0} \dfrac{2}{3\,|x|^{1/3}} = \infty$. This shows that

g has a vertical tangent line at $x = 0$.

(d)

59. $f(x) = |x-6| = \begin{cases} x-6 & \text{if } x-6 \ge 6 \\ -(x-6) & \text{if } x-6 < 0 \end{cases} = \begin{cases} x-6 & \text{if } x \ge 6 \\ 6-x & \text{if } x < 6 \end{cases}$

So the right-hand limit is $\lim\limits_{x\to 6^+} \dfrac{f(x)-f(6)}{x-6} = \lim\limits_{x\to 6^+} \dfrac{|x-6|-0}{x-6} = \lim\limits_{x\to 6^+} \dfrac{x-6}{x-6} = \lim\limits_{x\to 6^+} 1 = 1$, and the left-hand limit is $\lim\limits_{x\to 6^-} \dfrac{f(x)-f(6)}{x-6} = \lim\limits_{x\to 6^-} \dfrac{|x-6|-0}{x-6} = \lim\limits_{x\to 6^-} \dfrac{6-x}{x-6} = \lim\limits_{x\to 6^-} (-1) = -1$. Since these limits are not equal,

$f'(6) = \lim\limits_{x\to 6} \dfrac{f(x)-f(6)}{x-6}$ does not exist and f is not differentiable at 6.

However, a formula for f' is $f'(x) = \begin{cases} -1 & \text{if } x < 6 \\ 1 & \text{if } x > 6 \end{cases}$

Another way of writing the formula is $f'(x) = \dfrac{x-6}{|x-6|}$.

60. $f(x) = [\![x]\!]$ is not continuous at any integer n, so f is not differentiable at n by the contrapositive of Theorem 4. If a is not an integer, then f is constant on an open interval containing a, so $f'(a) = 0$. Thus, $f'(x) = 0$, x not an integer.

61. (a) $f(x) = x\,|x| = \begin{cases} x^2 & \text{if } x \ge 0 \\ -x^2 & \text{if } x < 0 \end{cases}$

(b) Since $f(x) = x^2$ for $x \ge 0$, we have $f'(x) = 2x$ for $x > 0$. [See Exercise 19(d).] Similarly, since $f(x) = -x^2$ for $x < 0$, we have $f'(x) = -2x$ for $x < 0$. At $x = 0$, we have

$f'(0) = \lim\limits_{x\to 0} \dfrac{f(x)-f(0)}{x-0} = \lim\limits_{x\to 0} \dfrac{x\,|x|}{x} = \lim\limits_{x\to 0} |x| = 0.$

So f is differentiable at 0. Thus, f is differentiable for all x.

(c) From part (b), we have $f'(x) = \left\{ \begin{matrix} 2x & \text{if } x \ge 0 \\ -2x & \text{if } x < 0 \end{matrix} \right\} = 2\,|x|$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

62. (a) $|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$

so $g(x) = x + |x| = \begin{cases} 2x & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases}$.

Graph the line $y = 2x$ for $x \geq 0$ and graph $y = 0$ (the x-axis) for $x < 0$.

(b) g is not differentiable at $x = 0$ because the graph has a corner there, but is differentiable at all other values; that is, g is differentiable on $(-\infty, 0) \cup (0, \infty)$.

(c) $g(x) = \begin{cases} 2x & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases} \quad \Rightarrow \quad g'(x) = \begin{cases} 2 & \text{if } x > 0 \\ 0 & \text{if } x < 0 \end{cases}$

Another way of writing the formula is $g'(x) = 1 + \operatorname{sgn} x$ for $x \neq 0$.

63. (a) If f is even, then

$$\begin{aligned} f'(-x) &= \lim_{h \to 0} \frac{f(-x+h) - f(-x)}{h} = \lim_{h \to 0} \frac{f[-(x-h)] - f(-x)}{h} \\ &= \lim_{h \to 0} \frac{f(x-h) - f(x)}{h} = -\lim_{h \to 0} \frac{f(x-h) - f(x)}{-h} \qquad [\text{let } \Delta x = -h] \\ &= -\lim_{\Delta x \to 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = -f'(x) \end{aligned}$$

Therefore, f' is odd.

(b) If f is odd, then

$$\begin{aligned} f'(-x) &= \lim_{h \to 0} \frac{f(-x+h) - f(-x)}{h} = \lim_{h \to 0} \frac{f[-(x-h)] - f(-x)}{h} \\ &= \lim_{h \to 0} \frac{-f(x-h) + f(x)}{h} = \lim_{h \to 0} \frac{f(x-h) - f(x)}{-h} \qquad [\text{let } \Delta x = -h] \\ &= \lim_{\Delta x \to 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = f'(x) \end{aligned}$$

Therefore, f' is even.

64. (a) $f(x) = \begin{cases} 0 & \text{if } x \leq 0 \\ x & \text{if } x > 0 \end{cases}$

$$f'_-(0) = \lim_{h \to 0^-} \frac{f(0+h) - f(0)}{h} = \lim_{h \to 0^-} \frac{0-0}{h} = \lim_{h \to 0^-} \frac{0}{h} = \lim_{h \to 0^-} 0 = 0$$

$$f'_+(0) = \lim_{h \to 0^+} \frac{f(0+h) - f(0)}{h} = \lim_{h \to 0^+} \frac{(0+h) - 0}{h} = \lim_{h \to 0^+} \frac{h}{h} = \lim_{h \to 0^+} 1 = 1$$

Since these one-sided derivatives are not equal, $f'(0)$ does not exist, so f is not differentiable at 0.

(b) $f(x) = \begin{cases} 0 & \text{if } x \leq 0 \\ x^2 & \text{if } x > 0 \end{cases}$

$$f'_-(0) = \lim_{h \to 0^-} \frac{f(0+h) - f(0)}{h} = \lim_{h \to 0^-} \frac{0-0}{h} = \lim_{h \to 0^-} \frac{0}{h} = \lim_{h \to 0^-} 0 = 0$$

$$f'_+(0) = \lim_{h \to 0^+} \frac{f(0+h) - f(0)}{h} = \lim_{h \to 0^+} \frac{(0+h)^2 - 0^2}{h} = \lim_{h \to 0^+} \frac{h^2}{h} = \lim_{h \to 0^+} h = 0$$

Since these one-sided derivatives are equal, $f'(0)$ exists (and equals 0), so f is differentiable at 0.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

65. (a) $f(x) = \begin{cases} 0 & \text{if } x \le 0 \\ 5 - x & \text{if } 0 < x < 4 \\ \dfrac{1}{5-x} & \text{if } x \ge 4 \end{cases}$

Note that as $h \to 0^-$, $4 + h < 4$, so $f(4+h) = 5 - (4+h)$. As $h \to 0^+$, $4 + h > 4$, so $f(4+h) = \dfrac{1}{5-(4+h)}$.

$$f'_-(4) = \lim_{h\to 0^-} \frac{f(4+h) - f(4)}{h} = \lim_{h\to 0^-} \frac{[5-(4+h)] - (5-4)}{h} = \lim_{h\to 0^-} \frac{(5-4-h)-1}{h}$$

$$= \lim_{h\to 0^-} \frac{-h}{h} = \lim_{h\to 0^-} (-1) = -1$$

$$f'_+(4) = \lim_{h\to 0^+} \frac{f(4+h) - f(4)}{h} = \lim_{h\to 0^+} \frac{\dfrac{1}{5-(4+h)} - \dfrac{1}{5-4}}{h} = \lim_{h\to 0^+} \frac{\dfrac{1}{1-h} - 1}{h}$$

$$= \lim_{h\to 0^+} \frac{\dfrac{1}{1-h} - \dfrac{1-h}{1-h}}{h} = \lim_{h\to 0^+} \frac{1-(1-h)}{h(1-h)} = \lim_{h\to 0^+} \frac{h}{h(1-h)} = \lim_{h\to 0^+} \frac{1}{1-h} = 1$$

(b)

(c) f is discontinuous at $x = 0$ (jump discontinuity) and at $x = 5$ (infinite discontinuity).

(d) f is not differentiable at $x = 0$ [discontinuous, from part (c)], $x = 4$ [one-sided derivatives are not equal, from part (a)], and at $x = 5$ [discontinuous, from part (c)].

66. (a) The initial temperature of the water is close to room temperature because of the water that was in the pipes. When the water from the hot water tank starts coming out, dT/dt is large and positive as T increases to the temperature of the water in the tank. In the next phase, $dT/dt = 0$ as the water comes out at a constant, high temperature. After some time, dT/dt becomes small and negative as the contents of the hot water tank are exhausted. Finally, when the hot water has run out, dT/dt is once again 0 as the water maintains its (cold) temperature.

(b)

67. These graphs are idealizations conveying the spirit of the problem. In reality, changes in speed are not instantaneous, so the graph in (a) would not have corners and the graph in (b) would be continuous.

(a)

(b)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

68.

In the right triangle in the diagram, let Δy be the side opposite angle ϕ and Δx the side adjacent to angle ϕ. Then the slope of the tangent line ℓ is $m = \Delta y/\Delta x = \tan\phi$. Note that $0 < \phi < \frac{\pi}{2}$. We know (see Exercise 19) that the derivative of $f(x) = x^2$ is $f'(x) = 2x$. So the slope of the tangent to the curve at the point $(1, 1)$ is 2. Thus, ϕ is the angle between 0 and $\frac{\pi}{2}$ whose tangent is 2; that is, $\phi = \tan^{-1} 2 \approx 63°$.

2 Review

TRUE-FALSE QUIZ

1. False. Limit Law 2 applies only if the individual limits exist (these don't).

2. False. Limit Law 5 cannot be applied if the limit of the denominator is 0 (it is).

3. True. Limit Law 5 applies.

4. False. $\dfrac{x^2-9}{x-3}$ is not defined when $x = 3$, but $x + 3$ is.

5. True. $\displaystyle\lim_{x\to 3}\frac{x^2-9}{x-3} = \lim_{x\to 3}\frac{(x+3)(x-3)}{(x-3)} = \lim_{x\to 3}(x+3)$

6. True. The limit doesn't exist since $f(x)/g(x)$ doesn't approach any real number as x approaches 5. (The denominator approaches 0 and the numerator doesn't.)

7. False. Consider $\displaystyle\lim_{x\to 5}\frac{x(x-5)}{x-5}$ or $\displaystyle\lim_{x\to 5}\frac{\sin(x-5)}{x-5}$. The first limit exists and is equal to 5. By Example 2.2.2, we know that the latter limit exists (and it is equal to 1).

8. False. If $f(x) = 1/x$, $g(x) = -1/x$, and $a = 0$, then $\displaystyle\lim_{x\to 0} f(x)$ does not exist, $\displaystyle\lim_{x\to 0} g(x)$ does not exist, but $\displaystyle\lim_{x\to 0}[f(x) + g(x)] = \lim_{x\to 0} 0 = 0$ exists.

9. True. Suppose that $\displaystyle\lim_{x\to a}[f(x) + g(x)]$ exists. Now $\displaystyle\lim_{x\to a} f(x)$ exists and $\displaystyle\lim_{x\to a} g(x)$ does not exist, but $\displaystyle\lim_{x\to a} g(x) = \lim_{x\to a}\{[f(x) + g(x)] - f(x)\} = \lim_{x\to a}[f(x) + g(x)] - \lim_{x\to a} f(x)$ [by Limit Law 2], which exists, and we have a contradiction. Thus, $\displaystyle\lim_{x\to a}[f(x) + g(x)]$ does not exist.

10. True. A polynomial is continuous everywhere, so $\displaystyle\lim_{x\to b} p(x)$ exists and is equal to $p(b)$.

11. False. Consider $\displaystyle\lim_{x\to 0}[f(x) - g(x)] = \lim_{x\to 0}\left(\frac{1}{x^2} - \frac{1}{x^4}\right)$. This limit is $-\infty$ (not 0), but each of the individual functions approaches ∞.

12. True. See Figure 2.6.8.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. False. Consider $f(x) = \sin x$ for $x \geq 0$. $\lim\limits_{x\to\infty} f(x) \neq \pm\infty$ and f has no horizontal asymptote.

14. False. Consider $f(x) = \begin{cases} 1/(x-1) & \text{if } x \neq 1 \\ 2 & \text{if } x = 1 \end{cases}$

15. False. The function f must be *continuous* in order to use the Intermediate Value Theorem. For example, let $f(x) = \begin{cases} 1 & \text{if } 0 \leq x < 3 \\ -1 & \text{if } x = 3 \end{cases}$ There is no number $c \in [0, 3]$ with $f(c) = 0$.

16. True. Use Theorem 2.5.8 with $a = 2$, $b = 5$, and $g(x) = 4x^2 - 11$. Note that $f(4) = 3$ is not needed.

17. True. Use the Intermediate Value Theorem with $a = -1$, $b = 1$, and $N = \pi$, since $3 < \pi < 4$.

18. True, by the definition of a limit with $\varepsilon = 1$.

19. False. For example, let $f(x) = \begin{cases} x^2 + 1 & \text{if } x \neq 0 \\ 2 & \text{if } x = 0 \end{cases}$

Then $f(x) > 1$ for all x, but $\lim\limits_{x\to 0} f(x) = \lim\limits_{x\to 0} \left(x^2 + 1\right) = 1$.

20. False. See the note after Theorem 2.8.4.

21. True. $f'(r)$ exists $\Rightarrow$ f is differentiable at r $\Rightarrow$ f is continuous at r $\Rightarrow$ $\lim\limits_{x\to r} f(x) = f(r)$.

22. False. $\dfrac{d^2y}{dx^2}$ is the second derivative while $\left(\dfrac{dy}{dx}\right)^2$ is the first derivative squared. For example, if $y = x$, then $\dfrac{d^2y}{dx^2} = 0$, but $\left(\dfrac{dy}{dx}\right)^2 = 1$.

23. True. $f(x) = x^{10} - 10x^2 + 5$ is continuous on the interval $[0, 2]$, $f(0) = 5$, $f(1) = -4$, and $f(2) = 989$. Since $-4 < 0 < 5$, there is a number c in $(0, 1)$ such that $f(c) = 0$ by the Intermediate Value Theorem. Thus, there is a solution of the equation $x^{10} - 10x^2 + 5 = 0$ in the interval $(0, 1)$. Similarly, there is a solution in $(1, 2)$.

24. True. See Exercise 2.5.76(b).

25. False. See Exercise 2.5.76(c).

26. False. For example, let $f(x) = x$ and $a = 0$. Then f is differentiable at a, but $|f| = |x|$ is not.

EXERCISES

1. (a) (i) $\lim\limits_{x\to 2^+} f(x) = 3$ (ii) $\lim\limits_{x\to -3^+} f(x) = 0$

(iii) $\lim\limits_{x\to -3} f(x)$ does not exist since the left and right limits are not equal. (The left limit is -2.)

(iv) $\lim\limits_{x\to 4} f(x) = 2$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(v) $\lim_{x\to 0} f(x) = \infty$ (vi) $\lim_{x\to 2^-} f(x) = -\infty$

(vii) $\lim_{x\to\infty} f(x) = 4$ (viii) $\lim_{x\to-\infty} f(x) = -1$

(b) The equations of the horizontal asymptotes are $y = -1$ and $y = 4$.

(c) The equations of the vertical asymptotes are $x = 0$ and $x = 2$.

(d) f is discontinuous at $x = -3$, 0, 2, and 4. The discontinuities are jump, infinite, infinite, and removable, respectively.

2. $\lim_{x\to-\infty} f(x) = -2$, $\lim_{x\to\infty} f(x) = 0$, $\lim_{x\to-3} f(x) = \infty$,

$\lim_{x\to 3^-} f(x) = -\infty$, $\lim_{x\to 3^+} f(x) = 2$,

f is continuous from the right at 3

3. Since the cosine function is continuous on $(-\infty, \infty)$, $\lim_{x\to 0} \cos(x^3 + 3x) = \cos(0^3 + 3\cdot 0) = \cos 0 = 1$.

4. Since a rational function is continuous on its domain, $\lim_{x\to 3} \dfrac{x^2-9}{x^2+2x-3} = \dfrac{3^2-9}{3^2+2(3)-3} = \dfrac{0}{12} = 0$.

5. $\lim_{x\to-3} \dfrac{x^2-9}{x^2+2x-3} = \lim_{x\to-3} \dfrac{(x+3)(x-3)}{(x+3)(x-1)} = \lim_{x\to-3} \dfrac{x-3}{x-1} = \dfrac{-3-3}{-3-1} = \dfrac{-6}{-4} = \dfrac{3}{2}$

6. $\lim_{x\to 1^+} \dfrac{x^2-9}{x^2+2x-3} = -\infty$ since $x^2+2x-3 \to 0^+$ as $x \to 1^+$ and $\dfrac{x^2-9}{x^2+2x-3} < 0$ for $1 < x < 3$.

7. $\lim_{h\to 0} \dfrac{(h-1)^3+1}{h} = \lim_{h\to 0} \dfrac{\left(h^3-3h^2+3h-1\right)+1}{h} = \lim_{h\to 0} \dfrac{h^3-3h^2+3h}{h} = \lim_{h\to 0} \left(h^2-3h+3\right) = 3$

Another solution: Factor the numerator as a sum of two cubes and then simplify.

$$\begin{aligned}\lim_{h\to 0} \frac{(h-1)^3+1}{h} &= \lim_{h\to 0} \frac{(h-1)^3+1^3}{h} = \lim_{h\to 0} \frac{[(h-1)+1]\left[(h-1)^2-1(h-1)+1^2\right]}{h} \\ &= \lim_{h\to 0} \left[(h-1)^2-h+2\right] = 1-0+2 = 3\end{aligned}$$

8. $\lim_{t\to 2} \dfrac{t^2-4}{t^3-8} = \lim_{t\to 2} \dfrac{(t+2)(t-2)}{(t-2)(t^2+2t+4)} = \lim_{t\to 2} \dfrac{t+2}{t^2+2t+4} = \dfrac{2+2}{4+4+4} = \dfrac{4}{12} = \dfrac{1}{3}$

9. $\lim_{r\to 9} \dfrac{\sqrt{r}}{(r-9)^4} = \infty$ since $(r-9)^4 \to 0^+$ as $r \to 9$ and $\dfrac{\sqrt{r}}{(r-9)^4} > 0$ for $r \neq 9$.

10. $\lim_{v\to 4^+} \dfrac{4-v}{|4-v|} = \lim_{v\to 4^+} \dfrac{4-v}{-(4-v)} = \lim_{v\to 4^+} \dfrac{1}{-1} = -1$

11. $\lim_{r\to -1} \dfrac{r^2-3r-4}{4r^2+r-3} = \lim_{r\to -1} \dfrac{(r-4)(r+1)}{(4r-3)(r+1)} = \lim_{r\to -1} \dfrac{r-4}{4r-3} = \dfrac{-1-4}{4(-1)-3} = \dfrac{-5}{-7} = \dfrac{5}{7}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $\lim\limits_{t\to5}\dfrac{3-\sqrt{t+4}}{t-5}=\lim\limits_{t\to5}\dfrac{3-\sqrt{t+4}}{t-5}\left(\dfrac{3+\sqrt{t+4}}{3+\sqrt{t+4}}\right)=\lim\limits_{t\to5}\dfrac{9-(t+4)}{(t-5)\big(3+\sqrt{t+4}\big)}$

$$=\lim_{t\to5}\frac{5-t}{(t-5)\big(3+\sqrt{t+4}\big)}=\lim_{t\to5}\frac{-1}{3+\sqrt{t+4}}=\frac{-1}{3+\sqrt{5+4}}=-\frac{1}{6}$$

13. Since x is positive, $\sqrt{x^2}=|x|=x$. Thus,

$$\lim_{x\to\infty}\frac{\sqrt{x^2-9}}{2x-6}=\lim_{x\to\infty}\frac{\sqrt{x^2-9}/\sqrt{x^2}}{(2x-6)/x}=\lim_{x\to\infty}\frac{\sqrt{1-9/x^2}}{2-6/x}=\frac{\sqrt{1-0}}{2-0}=\frac{1}{2}$$

14. Since x is negative, $\sqrt{x^2}=|x|=-x$. Thus,

$$\lim_{x\to-\infty}\frac{\sqrt{x^2-9}}{2x-6}=\lim_{x\to-\infty}\frac{\sqrt{x^2-9}/\sqrt{x^2}}{(2x-6)/(-x)}=\lim_{x\to-\infty}\frac{\sqrt{1-9/x^2}}{-2+6/x}=\frac{\sqrt{1-0}}{-2+0}=-\frac{1}{2}$$

15. Let $t=\sin x$. Then as $x\to\pi^-$, $\sin x\to0^+$, so $t\to0^+$. Thus, $\lim\limits_{x\to\pi^-}\ln(\sin x)=\lim\limits_{t\to0^+}\ln t=-\infty$.

16. $\lim\limits_{x\to-\infty}\dfrac{1-2x^2-x^4}{5+x-3x^4}=\lim\limits_{x\to-\infty}\dfrac{(1-2x^2-x^4)/x^4}{(5+x-3x^4)/x^4}=\lim\limits_{x\to-\infty}\dfrac{1/x^4-2/x^2-1}{5/x^4+1/x^3-3}=\dfrac{0-0-1}{0+0-3}=\dfrac{-1}{-3}=\dfrac{1}{3}$

17. $\lim\limits_{x\to\infty}\left(\sqrt{x^2+4x+1}-x\right)=\lim\limits_{x\to\infty}\left(\dfrac{\sqrt{x^2+4x+1}-x}{1}\cdot\dfrac{\sqrt{x^2+4x+1}+x}{\sqrt{x^2+4x+1}+x}\right)=\lim\limits_{x\to\infty}\dfrac{(x^2+4x+1)-x^2}{\sqrt{x^2+4x+1}+x}$

$$=\lim_{x\to\infty}\frac{(4x+1)/x}{(\sqrt{x^2+4x+1}+x)/x}\qquad\left[\text{divide by } x=\sqrt{x^2} \text{ for } x>0\right]$$

$$=\lim_{x\to\infty}\frac{4+1/x}{\sqrt{1+4/x+1/x^2}+1}=\frac{4+0}{\sqrt{1+0+0}+1}=\frac{4}{2}=2$$

18. Let $t=x-x^2=x(1-x)$. Then as $x\to\infty$, $t\to-\infty$, and $\lim\limits_{x\to\infty}e^{x-x^2}=\lim\limits_{t\to-\infty}e^t=0$.

19. Let $t=1/x$. Then as $x\to0^+$, $t\to\infty$, and $\lim\limits_{x\to0^+}\tan^{-1}(1/x)=\lim\limits_{t\to\infty}\tan^{-1}t=\dfrac{\pi}{2}$.

20. $\lim\limits_{x\to1}\left(\dfrac{1}{x-1}+\dfrac{1}{x^2-3x+2}\right)=\lim\limits_{x\to1}\left[\dfrac{1}{x-1}+\dfrac{1}{(x-1)(x-2)}\right]=\lim\limits_{x\to1}\left[\dfrac{x-2}{(x-1)(x-2)}+\dfrac{1}{(x-1)(x-2)}\right]$

$$=\lim_{x\to1}\left[\frac{x-1}{(x-1)(x-2)}\right]=\lim_{x\to1}\frac{1}{x-2}=\frac{1}{1-2}=-1$$

21. From the graph of $y=\big(\cos^2x\big)/x^2$, it appears that $y=0$ is the horizontal asymptote and $x=0$ is the vertical asymptote. Now $0\le(\cos x)^2\le1\ \Rightarrow$ $\dfrac{0}{x^2}\le\dfrac{\cos^2x}{x^2}\le\dfrac{1}{x^2}\ \Rightarrow\ 0\le\dfrac{\cos^2x}{x^2}\le\dfrac{1}{x^2}$. But $\lim\limits_{x\to\pm\infty}0=0$ and $\lim\limits_{x\to\pm\infty}\dfrac{1}{x^2}=0$, so by the Squeeze Theorem, $\lim\limits_{x\to\pm\infty}\dfrac{\cos^2x}{x^2}=0$.

Thus, $y=0$ is the horizontal asymptote. $\lim\limits_{x\to0}\dfrac{\cos^2x}{x^2}=\infty$ because $\cos^2x\to1$ and $x^2\to0^+$ as $x\to0$, so $x=0$ is the vertical asymptote.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. From the graph of $y = f(x) = \sqrt{x^2+x+1} - \sqrt{x^2-x}$, it appears that there are 2 horizontal asymptotes and possibly 2 vertical asymptotes. To obtain a different form for f, let's multiply and divide it by its conjugate.

$$f_1(x) = \left(\sqrt{x^2+x+1} - \sqrt{x^2-x}\right) \frac{\sqrt{x^2+x+1}+\sqrt{x^2-x}}{\sqrt{x^2+x+1}+\sqrt{x^2-x}} = \frac{(x^2+x+1)-(x^2-x)}{\sqrt{x^2+x+1}+\sqrt{x^2-x}}$$
$$= \frac{2x+1}{\sqrt{x^2+x+1}+\sqrt{x^2-x}}$$

Now

$$\lim_{x\to\infty} f_1(x) = \lim_{x\to\infty} \frac{2x+1}{\sqrt{x^2+x+1}+\sqrt{x^2-x}}$$
$$= \lim_{x\to\infty} \frac{2+(1/x)}{\sqrt{1+(1/x)+(1/x^2)}+\sqrt{1-(1/x)}} \qquad [\text{since } \sqrt{x^2} = x \text{ for } x > 0]$$
$$= \frac{2}{1+1} = 1,$$

so $y = 1$ is a horizontal asymptote. For $x < 0$, we have $\sqrt{x^2} = |x| = -x$, so when we divide the denominator by x, with $x < 0$, we get

$$\frac{\sqrt{x^2+x+1}+\sqrt{x^2-x}}{x} = -\frac{\sqrt{x^2+x+1}+\sqrt{x^2-x}}{\sqrt{x^2}} = -\left[\sqrt{1+\frac{1}{x}+\frac{1}{x^2}}+\sqrt{1-\frac{1}{x}}\right]$$

Therefore,

$$\lim_{x\to-\infty} f_1(x) = \lim_{x\to-\infty} \frac{2x+1}{\sqrt{x^2+x+1}+\sqrt{x^2-x}} = \lim_{x\to\infty} \frac{2+(1/x)}{-\left[\sqrt{1+(1/x)+(1/x^2)}+\sqrt{1-(1/x)}\right]}$$
$$= \frac{2}{-(1+1)} = -1,$$

so $y = -1$ is a horizontal asymptote.

The domain of f is $(-\infty, 0] \cup [1, \infty)$. As $x \to 0^-$, $f(x) \to 1$, so $x = 0$ is *not* a vertical asymptote. As $x \to 1^+$, $f(x) \to \sqrt{3}$, so $x = 1$ is *not* a vertical asymptote and hence there are no vertical asymptotes.

23. Since $2x - 1 \le f(x) \le x^2$ for $0 < x < 3$ and $\lim_{x\to1}(2x-1) = 1 = \lim_{x\to1} x^2$, we have $\lim_{x\to1} f(x) = 1$ by the Squeeze Theorem.

24. Let $f(x) = -x^2$, $g(x) = x^2\cos(1/x^2)$ and $h(x) = x^2$. Then since $\left|\cos(1/x^2)\right| \le 1$ for $x \ne 0$, we have $f(x) \le g(x) \le h(x)$ for $x \ne 0$, and so $\lim_{x\to0} f(x) = \lim_{x\to0} h(x) = 0 \;\Rightarrow\; \lim_{x\to0} g(x) = 0$ by the Squeeze Theorem.

25. Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < |x-2| < \delta$, then $|(14-5x)-4| < \varepsilon$. But $|(14-5x)-4| < \varepsilon \;\Leftrightarrow\; |-5x+10| < \varepsilon \;\Leftrightarrow\; |-5|\,|x-2| < \varepsilon \;\Leftrightarrow\; |x-2| < \varepsilon/5$. So if we choose $\delta = \varepsilon/5$, then $0 < |x-2| < \delta \;\Rightarrow\; |(14-5x)-4| < \varepsilon$. Thus, $\lim_{x\to2}(14-5x) = 4$ by the definition of a limit.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. Given $\varepsilon > 0$ we must find $\delta > 0$ so that if $0 < |x - 0| < \delta$, then $|\sqrt[3]{x} - 0| < \varepsilon$. Now $|\sqrt[3]{x} - 0| = |\sqrt[3]{x}| < \varepsilon \quad \Rightarrow$ $|x| = |\sqrt[3]{x}|^3 < \varepsilon^3$. So take $\delta = \varepsilon^3$. Then $0 < |x - 0| = |x| < \varepsilon^3 \quad \Rightarrow \quad |\sqrt[3]{x} - 0| = |\sqrt[3]{x}| = \sqrt[3]{|x|} < \sqrt[3]{\varepsilon^3} = \varepsilon$.

Therefore, by the definition of a limit, $\lim\limits_{x \to 0} \sqrt[3]{x} = 0$.

27. Given $\varepsilon > 0$, we need $\delta > 0$ so that if $0 < |x - 2| < \delta$, then $\left|x^2 - 3x - (-2)\right| < \varepsilon$. First, note that if $|x - 2| < 1$, then $-1 < x - 2 < 1$, so $0 < x - 1 < 2 \quad \Rightarrow \quad |x - 1| < 2$. Now let $\delta = \min\{\varepsilon/2, 1\}$. Then $0 < |x - 2| < \delta \quad \Rightarrow$ $\left|x^2 - 3x - (-2)\right| = |(x - 2)(x - 1)| = |x - 2|\,|x - 1| < (\varepsilon/2)(2) = \varepsilon$.

Thus, $\lim\limits_{x \to 2}(x^2 - 3x) = -2$ by the definition of a limit.

28. Given $M > 0$, we need $\delta > 0$ such that if $0 < x - 4 < \delta$, then $2/\sqrt{x - 4} > M$. This is true $\quad \Leftrightarrow \quad \sqrt{x - 4} < 2/M \quad \Leftrightarrow$ $x - 4 < 4/M^2$. So if we choose $\delta = 4/M^2$, then $0 < x - 4 < \delta \quad \Rightarrow \quad 2/\sqrt{x - 4} > M$. So by the definition of a limit, $\lim\limits_{x \to 4^+} \left(2/\sqrt{x - 4}\right) = \infty$.

29. (a) $f(x) = \sqrt{-x}$ if $x < 0$, $f(x) = 3 - x$ if $0 \le x < 3$, $f(x) = (x - 3)^2$ if $x > 3$.

(i) $\lim\limits_{x \to 0^+} f(x) = \lim\limits_{x \to 0^+} (3 - x) = 3$

(ii) $\lim\limits_{x \to 0^-} f(x) = \lim\limits_{x \to 0^-} \sqrt{-x} = 0$

(iii) Because of (i) and (ii), $\lim\limits_{x \to 0} f(x)$ does not exist.

(iv) $\lim\limits_{x \to 3^-} f(x) = \lim\limits_{x \to 3^-} (3 - x) = 0$

(v) $\lim\limits_{x \to 3^+} f(x) = \lim\limits_{x \to 3^+} (x - 3)^2 = 0$

(vi) Because of (iv) and (v), $\lim\limits_{x \to 3} f(x) = 0$.

(b) f is discontinuous at 0 since $\lim\limits_{x \to 0} f(x)$ does not exist.

f is discontinuous at 3 since $f(3)$ does not exist.

(c)

30. (a) $g(x) = 2x - x^2$ if $0 \le x \le 2$, $g(x) = 2 - x$ if $2 < x \le 3$, $g(x) = x - 4$ if $3 < x < 4$, $g(x) = \pi$ if $x \ge 4$.

Therefore, $\lim\limits_{x \to 2^-} g(x) = \lim\limits_{x \to 2^-} \left(2x - x^2\right) = 0$ and $\lim\limits_{x \to 2^+} g(x) = \lim\limits_{x \to 2^+} (2 - x) = 0$. Thus, $\lim\limits_{x \to 2} g(x) = 0 = g(2)$, so g is continuous at 2. $\lim\limits_{x \to 3^-} g(x) = \lim\limits_{x \to 3^-} (2 - x) = -1$ and $\lim\limits_{x \to 3^+} g(x) = \lim\limits_{x \to 3^+} (x - 4) = -1$. Thus, $\lim\limits_{x \to 3} g(x) = -1 = g(3)$, so g is continuous at 3.

$\lim\limits_{x \to 4^-} g(x) = \lim\limits_{x \to 4^-} (x - 4) = 0$ and $\lim\limits_{x \to 4^+} g(x) = \lim\limits_{x \to 4^+} \pi = \pi$.

Thus, $\lim\limits_{x \to 4} g(x)$ does not exist, so g is discontinuous at 4. But $\lim\limits_{x \to 4^+} g(x) = \pi = g(4)$, so g is continuous from the right at 4.

(b)

31. $\sin x$ and e^x are continuous on $\mathbb{R}$ by Theorem 2.5.7. Since e^x is continuous on $\mathbb{R}$, $e^{\sin x}$ is continuous on $\mathbb{R}$ by Theorem 2.5.9. Lastly, x is continuous on $\mathbb{R}$ since it's a polynomial and the product $xe^{\sin x}$ is continuous on its domain $\mathbb{R}$ by Theorem 2.5.4.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. $x^2 - 9$ is continuous on $\mathbb{R}$ since it is a polynomial and $\sqrt{x}$ is continuous on $[0, \infty)$ by Theorem 2.5.7, so the composition $\sqrt{x^2 - 9}$ is continuous on $\{x \mid x^2 - 9 \geq 0\} = (-\infty, -3] \cup [3, \infty)$ by Theorem 2.5.9. Note that $x^2 - 2 \neq 0$ on this set and so the quotient function $g(x) = \dfrac{\sqrt{x^2 - 9}}{x^2 - 2}$ is continuous on its domain, $(-\infty, -3] \cup [3, \infty)$ by Theorem 2.5.4.

33. $f(x) = x^5 - x^3 + 3x - 5$ is continuous on the interval $[1, 2]$, $f(1) = -2$, and $f(2) = 25$. Since $-2 < 0 < 25$, there is a number c in $(1, 2)$ such that $f(c) = 0$ by the Intermediate Value Theorem. Thus, there is a solution of the equation $x^5 - x^3 + 3x - 5 = 0$ in the interval $(1, 2)$.

34. $f(x) = \cos\sqrt{x} - e^x + 2$ is continuous on the interval $[0, 1]$, $f(0) = 2$, and $f(1) \approx -0.2$. Since $-0.2 < 0 < 2$, there is a number c in $(0, 1)$ such that $f(c) = 0$ by the Intermediate Value Theorem. Thus, there is a solution of the equation $\cos\sqrt{x} - e^x + 2 = 0$, or $\cos\sqrt{x} = e^x - 2$, in the interval $(0, 1)$.

35. (a) The slope of the tangent line at $(2, 1)$ is

$$\lim_{x\to 2} \frac{f(x) - f(2)}{x - 2} = \lim_{x\to 2} \frac{9 - 2x^2 - 1}{x - 2} = \lim_{x\to 2} \frac{8 - 2x^2}{x - 2} = \lim_{x\to 2} \frac{-2(x^2 - 4)}{x - 2} = \lim_{x\to 2} \frac{-2(x - 2)(x + 2)}{x - 2}$$
$$= \lim_{x\to 2} [-2(x + 2)] = -2 \cdot 4 = -8$$

(b) An equation of this tangent line is $y - 1 = -8(x - 2)$ or $y = -8x + 17$.

36. For a general point with x-coordinate a, we have

$$m = \lim_{x\to a} \frac{f(x) - f(a)}{x - a} = \lim_{x\to a} \frac{2/(1 - 3x) - 2/(1 - 3a)}{x - a} = \lim_{x\to a} \frac{2(1 - 3a) - 2(1 - 3x)}{(1 - 3a)(1 - 3x)(x - a)}$$
$$= \lim_{x\to a} \frac{6(x - a)}{(1 - 3a)(1 - 3x)(x - a)} = \lim_{x\to a} \frac{6}{(1 - 3a)(1 - 3x)} = \frac{6}{(1 - 3a)^2}$$

For $a = 0$, $m = 6$ and $f(0) = 2$, so an equation of the tangent line is $y - 2 = 6(x - 0)$ or $y = 6x + 2$. For $a = -1$, $m = \frac{3}{8}$ and $f(-1) = \frac{1}{2}$, so an equation of the tangent line is $y - \frac{1}{2} = \frac{3}{8}(x + 1)$ or $y = \frac{3}{8}x + \frac{7}{8}$.

37. (a) $s = s(t) = 1 + 2t + t^2/4$. The average velocity over the time interval $[1, 1 + h]$ is

$$v_{\text{ave}} = \frac{s(1 + h) - s(1)}{(1 + h) - 1} = \frac{1 + 2(1 + h) + (1 + h)^2/4 - 13/4}{h} = \frac{10h + h^2}{4h} = \frac{10 + h}{4}$$

So for the following intervals the average velocities are:

(i) $[1, 3]$: $h = 2$, $v_{\text{ave}} = (10 + 2)/4 = 3$ m/s (ii) $[1, 2]$: $h = 1$, $v_{\text{ave}} = (10 + 1)/4 = 2.75$ m/s

(iii) $[1, 1.5]$: $h = 0.5$, $v_{\text{ave}} = (10 + 0.5)/4 = 2.625$ m/s (iv) $[1, 1.1]$: $h = 0.1$, $v_{\text{ave}} = (10 + 0.1)/4 = 2.525$ m/s

(b) When $t = 1$, the instantaneous velocity is $\displaystyle\lim_{h\to 0} \frac{s(1 + h) - s(1)}{h} = \lim_{h\to 0} \frac{10 + h}{4} = \frac{10}{4} = 2.5$ m/s.

38. (a) When V increases from 200 in^3 to 250 in^3, we have $\Delta V = 250 - 200 = 50$ in^3, and since $P = 800/V$, $\Delta P = P(250) - P(200) = \dfrac{800}{250} - \dfrac{800}{200} = 3.2 - 4 = -0.8$ lb/in^2. So the average rate of change is $\dfrac{\Delta P}{\Delta V} = \dfrac{-0.8}{50} = -0.016\ \dfrac{\text{lb/in}^2}{\text{in}^3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Since $V = 800/P$, the instantaneous rate of change of V with respect to P is

$$\lim_{h\to 0}\frac{\Delta V}{\Delta P} = \lim_{h\to 0}\frac{V(P+h)-V(P)}{h} = \lim_{h\to 0}\frac{800/(P+h)-800/P}{h} = \lim_{h\to 0}\frac{800\,[P-(P+h)]}{h(P+h)P}$$
$$= \lim_{h\to 0}\frac{-800}{(P+h)P} = -\frac{800}{P^2}$$

which is inversely proportional to the square of P.

39. (a) $f'(2) = \lim_{x\to 2}\dfrac{f(x)-f(2)}{x-2} = \lim_{x\to 2}\dfrac{x^3-2x-4}{x-2}$

$$= \lim_{x\to 2}\frac{(x-2)(x^2+2x+2)}{x-2} = \lim_{x\to 2}(x^2+2x+2) = 10$$

(b) $y - 4 = 10(x-2)$ or $y = 10x - 16$

(c)

40. $2^6 = 64$, so $f(x) = x^6$ and $a = 2$.

41. (a) $f'(r)$ is the rate at which the total cost changes with respect to the interest rate. Its units are dollars/(percent per year).

(b) The total cost of paying off the loan is increasing by \$1200/(percent per year) as the interest rate reaches 10%. So if the interest rate goes up from 10% to 11%, the cost goes up approximately \$1200.

(c) As r increases, C increases. So $f'(r)$ will always be positive.

42.

43.

44.

45. $f'(x) = \lim_{h\to 0}\dfrac{f(x+h)-f(x)}{h} = \lim_{h\to 0}\dfrac{\dfrac{2}{(x+h)^2}-\dfrac{2}{x^2}}{h} = \lim_{h\to 0}\dfrac{\dfrac{2x^2-2(x+h)^2}{x^2(x+h)^2}}{h}$

$$= \lim_{h\to 0}\frac{2x^2-2x^2-4xh-2h^2}{hx^2(x+h)^2} = \lim_{h\to 0}\frac{-4xh-2h^2}{hx^2(x+h)^2} = \lim_{h\to 0}\frac{h\,(-4x-2h)}{hx^2(x+h)^2}$$
$$= \lim_{h\to 0}\frac{-4x-2h}{x^2(x+h)^2} = \frac{-4x}{x^2\cdot x^2} = -\frac{4}{x^3}$$

Domain of $f' = (-\infty, 0) \cup (0, \infty)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. $$f'(t) = \lim_{h\to 0} \frac{f(t+h)-f(t)}{h} = \lim_{h\to 0} \frac{\dfrac{1}{\sqrt{(t+h)+1}} - \dfrac{1}{\sqrt{t+1}}}{h} = \lim_{h\to 0} \frac{\dfrac{\sqrt{t+1}-\sqrt{(t+h)+1}}{\sqrt{(t+h)+1}\sqrt{t+1}}}{h}$$

$$= \lim_{h\to 0} \frac{\sqrt{t+1}-\sqrt{(t+h)+1}}{h\sqrt{(t+h)+1}\sqrt{t+1}} \left[\frac{\sqrt{t+1}+\sqrt{(t+h)+1}}{\sqrt{t+1}+\sqrt{(t+h)+1}}\right]$$

$$= \lim_{h\to 0} \frac{(t+1)-[(t+h)+1]}{h\sqrt{(t+h)+1}\sqrt{t+1}\left(\sqrt{t+1}+\sqrt{(t+h)+1}\right)}$$

$$= \lim_{h\to 0} \frac{-h}{h\sqrt{(t+h)+1}\sqrt{t+1}\left(\sqrt{t+1}+\sqrt{(t+h)+1}\right)}$$

$$= \lim_{h\to 0} \frac{-1}{\sqrt{(t+h)+1}\sqrt{t+1}\left(\sqrt{t+1}+\sqrt{(t+h)+1}\right)}$$

$$= \frac{-1}{\sqrt{t+1}\sqrt{t+1}\left(\sqrt{t+1}+\sqrt{t+1}\right)} = -\frac{1}{(t+1)\,2\sqrt{t+1}} = -\frac{1}{2\,(t+1)^{3/2}}$$

Domain of $f' = (-1, \infty)$.

47. (a) $$f'(x) = \lim_{h\to 0} \frac{f(x+h)-f(x)}{h} = \lim_{h\to 0} \frac{\sqrt{3-5(x+h)}-\sqrt{3-5x}}{h} \frac{\sqrt{3-5(x+h)}+\sqrt{3-5x}}{\sqrt{3-5(x+h)}+\sqrt{3-5x}}$$

$$= \lim_{h\to 0} \frac{[3-5(x+h)]-(3-5x)}{h\left(\sqrt{3-5(x+h)}+\sqrt{3-5x}\right)} = \lim_{h\to 0} \frac{-5}{\sqrt{3-5(x+h)}+\sqrt{3-5x}} = \frac{-5}{2\sqrt{3-5x}}$$

(b) Domain of f: (the radicand must be nonnegative) $3-5x \geq 0 \quad \Rightarrow$

$5x \leq 3 \quad \Rightarrow \quad x \in \left(-\infty, \frac{3}{5}\right]$

Domain of f': exclude $\frac{3}{5}$ because it makes the denominator zero;

$x \in \left(-\infty, \frac{3}{5}\right)$

(c) Our answer to part (a) is reasonable because $f'(x)$ is always negative and f is always decreasing.

48. (a) As $x \to \pm\infty$, $f(x) = (4-x)/(3+x) \to -1$, so there is a horizontal asymptote at $y = -1$. As $x \to -3^+$, $f(x) \to \infty$, and as $x \to -3^-$, $f(x) \to -\infty$. Thus, there is a vertical asymptote at $x = -3$.

(b) Note that f is decreasing on $(-\infty, -3)$ and $(-3, \infty)$, so f' is negative on those intervals. As $x \to \pm\infty$, $f' \to 0$. As $x \to -3^-$ and as $x \to -3^+$, $f' \to -\infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $f'(x) = \lim\limits_{h\to 0} \dfrac{f(x+h)-f(x)}{h} = \lim\limits_{h\to 0} \dfrac{\dfrac{4-(x+h)}{3+(x+h)} - \dfrac{4-x}{3+x}}{h} = \lim\limits_{h\to 0} \dfrac{(3+x)\,[4-(x+h)] - (4-x)\,[3+(x+h)]}{h\,[3+(x+h)]\,(3+x)}$

$$= \lim_{h\to 0} \frac{(12-3x-3h+4x-x^2-hx)-(12+4x+4h-3x-x^2-hx)}{h[3+(x+h)](3+x)}$$

$$= \lim_{h\to 0} \frac{-7h}{h\,[3+(x+h)]\,(3+x)} = \lim_{h\to 0} \frac{-7}{[3+(x+h)]\,(3+x)} = -\frac{7}{(3+x)^2}$$

(d) The graphing device confirms our graph in part (b).

49. f is not differentiable: at $x=-4$ because f is not continuous, at $x=-1$ because f has a corner, at $x=2$ because f is not continuous, and at $x=5$ because f has a vertical tangent.

50. The graph of a has tangent lines with positive slope for $x<0$ and negative slope for $x>0$, and the values of c fit this pattern, so c must be the graph of the derivative of the function for a. The graph of c has horizontal tangent lines to the left and right of the x-axis and b has zeros at these points. Hence, b is the graph of the derivative of the function for c. Therefore, a is the graph of f, c is the graph of f', and b is the graph of f''.

51. Domain: $(-\infty,0)\cup(0,\infty)$; $\lim\limits_{x\to 0^-} f(x)=1$; $\lim\limits_{x\to 0^+} f(x)=0$;

$f'(x)>0$ for all x in the domain; $\lim\limits_{x\to -\infty} f'(x)=0$; $\lim\limits_{x\to\infty} f'(x)=1$

52. (a) $P'(t)$ is the rate at which the percentage of Americans under the age of 18 is changing with respect to time. Its units are percent per year (%/yr).

(b) To find $P'(t)$, we use $\lim\limits_{h\to 0} \dfrac{P(t+h)-P(t)}{h} \approx \dfrac{P(t+h)-P(t)}{h}$ for small values of h.

For 1950: $P'(1950) \approx \dfrac{P(1960)-P(1950)}{1960-1950} = \dfrac{35.7-31.1}{10} = 0.46$

For 1960: We estimate $P'(1960)$ by using $h=-10$ and $h=10$, and then average the two results to obtain a final estimate.

$h=-10 \quad\Rightarrow\quad P'(1960) \approx \dfrac{P(1950)-P(1960)}{1950-1960} = \dfrac{31.1-35.7}{-10} = 0.46$

$h=10 \quad\Rightarrow\quad P'(1960) \approx \dfrac{P(1970)-P(1960)}{1970-1960} = \dfrac{34.0-35.7}{10} = -0.17$

So we estimate that $P'(1960) \approx \frac{1}{2}[0.46+(-0.17)] = 0.145$.

t	1950	1960	1970	1980	1990	2000	2010
$P'(t)$	0.460	0.145	−0.385	−0.415	−0.115	−0.085	−0.170

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c)

(d) We could get more accurate values for $P'(t)$ by obtaining data for the mid-decade years 1955, 1965, 1975, 1985, 1995, and 2005.

53. $B'(t)$ is the rate at which the number of US \$20 bills in circulation is changing with respect to time. Its units are billions of bills per year. We use a symmetric difference quotient (see Exercise 2.7.60) to estimate $B'(2010)$.

$$B'(2010) \approx \frac{B(2010+5)-B(2010-5)}{(2010+5)-(2010-5)} = \frac{B(2015)-B(2005)}{2(5)} = \frac{8.57-5.77}{10} = 0.280 \text{ billion of bills per year}$$

(or 280 million bills per year).

54. (a) Drawing slope triangles, we obtain the following estimates: $F'(1950) \approx \frac{1.1}{10} = 0.11$, $F'(1965) \approx \frac{-1.6}{10} = -0.16$, and $F'(1987) \approx \frac{0.2}{10} = 0.02$.

(b) The rate of change of the average number of children born to each woman was increasing by 0.11 in 1950, decreasing by 0.16 in 1965, and increasing by 0.02 in 1987.

(c) There are many possible reasons:

- In the baby-boom era (post-WWII), there was optimism about the economy and family size was rising.
- In the baby-bust era, there was less economic optimism, and it was considered less socially responsible to have a large family.
- In the baby-boomlet era, there was increased economic optimism and a return to more conservative attitudes.

55. $|f(x)| \le g(x) \quad \Leftrightarrow \quad -g(x) \le f(x) \le g(x)$ and $\lim\limits_{x\to a} g(x) = 0 = \lim\limits_{x\to a} -g(x)$.

Thus, by the Squeeze Theorem, $\lim\limits_{x\to a} f(x) = 0$.

56. (a) Note that f is an even function since $f(x) = f(-x)$. Now for any integer n, $[\![n]\!] + [\![-n]\!] = n - n = 0$, and for any real number k which is not an integer, $[\![k]\!] + [\![-k]\!] = [\![k]\!] + (-[\![k]\!] - 1) = -1$. So $\lim\limits_{x\to a} f(x)$ exists (and is equal to -1) for all values of a.

(b) f is discontinuous at all integers.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1. Let $t = \sqrt[6]{x}$, so $x = t^6$. Then $t \to 1$ as $x \to 1$, so

$$\lim_{x\to 1} \frac{\sqrt[3]{x}-1}{\sqrt{x}-1} = \lim_{t\to 1} \frac{t^2-1}{t^3-1} = \lim_{t\to 1} \frac{(t-1)(t+1)}{(t-1)\,(t^2+t+1)} = \lim_{t\to 1} \frac{t+1}{t^2+t+1} = \frac{1+1}{1^2+1+1} = \frac{2}{3}.$$

Another method: Multiply both the numerator and the denominator by $(\sqrt{x}+1)\left(\sqrt[3]{x^2}+\sqrt[3]{x}+1\right)$.

2. First rationalize the numerator: $\displaystyle\lim_{x\to 0} \frac{\sqrt{ax+b}-2}{x} \cdot \frac{\sqrt{ax+b}+2}{\sqrt{ax+b}+2} = \lim_{x\to 0} \frac{ax+b-4}{x\left(\sqrt{ax+b}+2\right)}$. Now since the denominator approaches 0 as $x \to 0$, the limit will exist only if the numerator also approaches 0 as $x \to 0$. So we require that $a(0)+b-4=0 \;\Rightarrow\; b=4$. So the equation becomes $\displaystyle\lim_{x\to 0} \frac{a}{\sqrt{ax+4}+2} = 1 \;\Rightarrow\; \frac{a}{\sqrt{4}+2} = 1 \;\Rightarrow\; a=4$.

Therefore, $a=b=4$.

3. For $-\frac{1}{2} < x < \frac{1}{2}$, we have $2x-1<0$ and $2x+1>0$, so $|2x-1| = -(2x-1)$ and $|2x+1| = 2x+1$.

Therefore, $\displaystyle\lim_{x\to 0} \frac{|2x-1|-|2x+1|}{x} = \lim_{x\to 0} \frac{-(2x-1)-(2x+1)}{x} = \lim_{x\to 0} \frac{-4x}{x} = \lim_{x\to 0}(-4) = -4$.

4. Let R be the midpoint of OP, so the coordinates of R are $\left(\frac{1}{2}x, \frac{1}{2}x^2\right)$ since the coordinates of P are $\left(x, x^2\right)$. Let $Q=(0,a)$.

Since the slope $m_{OP} = \dfrac{x^2}{x} = x$, $m_{QR} = -\dfrac{1}{x}$ (negative reciprocal). But $m_{QR} = \dfrac{\frac{1}{2}x^2-a}{\frac{1}{2}x-0} = \dfrac{x^2-2a}{x}$, so we conclude that

$-1 = x^2-2a \;\Rightarrow\; 2a = x^2+1 \;\Rightarrow\; a = \frac{1}{2}x^2+\frac{1}{2}$. As $x\to 0$, $a \to \frac{1}{2}$, and the limiting position of Q is $\left(0, \frac{1}{2}\right)$.

5. (a) For $0<x<1$, $[\![x]\!]=0$, so $\dfrac{[\![x]\!]}{x}=0$, and $\displaystyle\lim_{x\to 0^+} \frac{[\![x]\!]}{x} = 0$. For $-1<x<0$, $[\![x]\!]=-1$, so $\dfrac{[\![x]\!]}{x} = \dfrac{-1}{x}$, and

$\displaystyle\lim_{x\to 0^-} \frac{[\![x]\!]}{x} = \lim_{x\to 0^-}\left(\frac{-1}{x}\right) = \infty$. Since the one-sided limits are not equal, $\displaystyle\lim_{x\to 0} \frac{[\![x]\!]}{x}$ does not exist.

(b) For $x>0$, $1/x-1 \le [\![1/x]\!] \le 1/x \;\Rightarrow\; x(1/x-1) \le x[\![1/x]\!] \le x(1/x) \;\Rightarrow\; 1-x \le x[\![1/x]\!] \le 1$.

As $x\to 0^+$, $1-x\to 1$, so by the Squeeze Theorem, $\displaystyle\lim_{x\to 0^+} x[\![1/x]\!] = 1$.

For $x<0$, $1/x-1 \le [\![1/x]\!] \le 1/x \;\Rightarrow\; x(1/x-1) \ge x[\![1/x]\!] \ge x(1/x) \;\Rightarrow\; 1-x \ge x[\![1/x]\!] \ge 1$.

As $x\to 0^-$, $1-x\to 1$, so by the Squeeze Theorem, $\displaystyle\lim_{x\to 0^-} x[\![1/x]\!] = 1$.

Since the one-sided limits are equal, $\displaystyle\lim_{x\to 0} x[\![1/x]\!] = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. (a) $[\![x]\!]^2 + [\![y]\!]^2 = 1$. Since $[\![x]\!]^2$ and $[\![y]\!]^2$ are positive integers or 0, there are only 4 cases:

Case (i): $[\![x]\!] = 1, [\![y]\!] = 0 \quad \Rightarrow 1 \le x < 2$ and $0 \le y < 1$

Case (ii): $[\![x]\!] = -1, [\![y]\!] = 0 \Rightarrow -1 \le x < 0$ and $0 \le y < 1$

Case (iii): $[\![x]\!] = 0, [\![y]\!] = 1 \quad \Rightarrow 0 \le x < 1$ and $1 \le y < 2$

Case (iv): $[\![x]\!] = 0, [\![y]\!] = -1 \Rightarrow 0 \le x < 1$ and $-1 \le y < 0$

(b) $[\![x]\!]^2 - [\![y]\!]^2 = 3$. The only integral solution of $n^2 - m^2 = 3$ is $n = \pm 2$ and $m = \pm 1$. So the graph is

$$\{(x, y) \mid [\![x]\!] = \pm 2,\ [\![y]\!] = \pm 1\} = \left\{ (x, y) \;\middle|\; \begin{array}{l} 2 \le x \le 3 \text{ or } -2 \le x < 1, \\ 1 \le y < 2 \text{ or } -1 \le y < 0 \end{array} \right\}.$$

(c) $[\![x + y]\!]^2 = 1 \quad \Rightarrow \quad [\![x + y]\!] = \pm 1 \quad \Rightarrow \quad 1 \le x + y < 2$ or $-1 \le x + y < 0$

(d) For $n \le x < n + 1$, $[\![x]\!] = n$. Then $[\![x]\!] + [\![y]\!] = 1 \quad \Rightarrow \quad [\![y]\!] = 1 - n \quad \Rightarrow$ $1 - n \le y < 2 - n$. Choosing integer values for n produces the graph.

7. (a) The function $f(x) = x / [\![x]\!]$ is defined whenever $[\![x]\!] \ne 0$. Since $[\![x]\!] = 0$ for $x \in [0, 1)$, it follows that the domain of f is $(-\infty, 0) \cup [1, \infty)$.

To determine the range we examine the values of f on the intervals $(-\infty, 0)$ and $[1, \infty)$ separately. A graph of f is helpful here.

On $(-\infty, 0)$, consider the intervals $[-a, -a + 1)$ for each positive integer a. On each such interval, f is decreasing, $f(a) = 1$, and

$$\lim_{x \to (-a+1)^-} f(x) = \frac{\lim\limits_{x \to (-a+1)^-} x}{\lim\limits_{x \to (-a+1)^-} [\![x]\!]} = \frac{-a + 1}{-a} = 1 - \frac{1}{a}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

So the range of f on the interval $[-a, -a+1)$ is $(1-1/a, 1]$. The intervals $(1-1/a, 1]$ are nested and their union is just the largest one, which occurs when $a = 1$. So the range of f on $(-\infty, 0)$ is $(0, 1]$.

On $[1, \infty)$, consider the intervals $[a, a+1)$, for each positive integer a. On each such interval, f is increasing, $f(a) = 1$, and

$$\lim_{x \to (a+1)^-} f(x) = \frac{\lim\limits_{x \to (a+1)^-} x}{\lim\limits_{x \to (a+1)^-} [\![x]\!]} = \frac{a+1}{a} = 1 + \frac{1}{a}$$

So the range of f on the interval $[a, a+1)$ is $[1, 1+1/a)$. The intervals $[1, 1+1/a)$ are nested and their union is the largest one, which occurs when $a = 1$. So the range of f on $[1, \infty)$ is $[1, 2)$.

Finally, combining the preceding results, we see that the range of f is $(0, 1] \cup [1, 2)$, or $(0, 2)$.

(b) First note that $x - 1 \le [\![x]\!] \le x$. For $x > 0$, $\dfrac{x-1}{x} \le \dfrac{[\![x]\!]}{x} \le \dfrac{x}{x}$. For $x > 2$, taking reciprocals, we have $\dfrac{x}{x-1} \ge \dfrac{x}{[\![x]\!]} \ge 1$. Now $\lim\limits_{x \to \infty} \dfrac{x}{x-1} = 1$ and $\lim\limits_{x \to \infty} 1 = 1$. It follows by the Squeeze Theorem that $\lim\limits_{x \to \infty} \dfrac{x}{[\![x]\!]} = 1$.

8. (a) Here are a few possibilities:

(b) The "obstacle" is the line $x = y$ (see diagram). Any intersection of the graph of f with the line $y = x$ constitutes a fixed point, and if the graph of the function does not cross the line somewhere in $(0, 1)$, then it must either start at $(0, 0)$ (in which case 0 is a fixed point) or finish at $(1, 1)$ (in which case 1 is a fixed point).

(c) Consider the function $F(x) = f(x) - x$, where f is any continuous function with domain $[0, 1]$ and range in $[0, 1]$. We shall prove that f has a fixed point. Now if $f(0) = 0$ then we are done: f has a fixed point (the number 0), which is what we are trying to prove. So assume $f(0) \ne 0$. For the same reason we can assume that $f(1) \ne 1$. Then $F(0) = f(0) > 0$ and $F(1) = f(1) - 1 < 0$. So by the Intermediate Value Theorem, there exists some number c in the interval $(0, 1)$ such that $F(c) = f(c) - c = 0$. So $f(c) = c$, and therefore f has a fixed point.

9. $$\begin{cases} \lim\limits_{x \to a} [f(x) + g(x)] = 2 \\ \lim\limits_{x \to a} [f(x) - g(x)] = 1 \end{cases} \Rightarrow \begin{cases} \lim\limits_{x \to a} f(x) + \lim\limits_{x \to a} g(x) = 2 & \textbf{(1)} \\ \lim\limits_{x \to a} f(x) - \lim\limits_{x \to a} g(x) = 1 & \textbf{(2)} \end{cases}$$

Adding equations **(1)** and **(2)** gives us $2 \lim\limits_{x \to a} f(x) = 3 \quad \Rightarrow \quad \lim\limits_{x \to a} f(x) = \frac{3}{2}$. From equation **(1)**, $\lim\limits_{x \to a} g(x) = \frac{1}{2}$. Thus, $\lim\limits_{x \to a} [f(x)\, g(x)] = \lim\limits_{x \to a} f(x) \cdot \lim\limits_{x \to a} g(x) = \frac{3}{2} \cdot \frac{1}{2} = \frac{3}{4}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. (a) *Solution 1:* We introduce a coordinate system and drop a perpendicular from P, as shown. We see from $\angle NCP$ that $\tan\, 2\theta = \dfrac{y}{1-x}$, and from $\angle NBP$ that $\tan\, \theta = y/x$. Using the double-angle formula for tangents, we get $\dfrac{y}{1-x} = \tan 2\theta = \dfrac{2\,\tan\,\theta}{1-\tan^2\theta} = \dfrac{2(\,y/x)}{1-(\,y/x)^2}$. After a bit of simplification, this becomes $\dfrac{1}{1-x} = \dfrac{2x}{x^2-y^2} \quad \Leftrightarrow \quad y^2 = x\,(3x-2)$.

As the altitude AM decreases in length, the point P will approach the x-axis, that is, $y \to 0$, so the limiting location of P must be one of the roots of the equation $x(3x-2) = 0$. Obviously it is not $x = 0$ (the point P can never be to the left of the altitude AM, which it would have to be in order to approach 0) so it must be $3x - 2 = 0$, that is, $x = \frac{2}{3}$.

Solution 2: We add a few lines to the original diagram, as shown. Now note that $\angle BPQ = \angle PBC$ (alternate angles; $QP \parallel BC$ by symmetry) and similarly $\angle CQP = \angle QCB$. So ΔBPQ and ΔCQP are isosceles, and the line segments BQ, QP and PC are all of equal length. As $|AM| \to 0$, P and Q approach points on the base, and the point P is seen to approach a position two-thirds of the way between B and C, as above.

(b) The equation $y^2 = x(3x-2)$ calculated in part (a) is the equation of the curve traced out by P. Now as $|AM| \to \infty$, $2\,\theta \to \frac{\pi}{2}$, $\theta \to \frac{\pi}{4}$, $x \to 1$, and since $\tan\theta = y/x$, $y \to 1$. Thus, P only traces out the part of the curve with $0 \le y < 1$.

11. (a) Consider $G(x) = T(x + 180°) - T(x)$. Fix any number a. If $G(a) = 0$, we are done: Temperature at $a =$ Temperature at $a + 180°$. If $G(a) > 0$, then $G(a + 180°) = T(a + 360°) - T(a + 180°) = T(a) - T(a + 180°) = -G(a) < 0$. Also, G is continuous since temperature varies continuously. So, by the Intermediate Value Theorem, G has a zero on the interval $[a, a + 180°]$. If $G(a) < 0$, then a similar argument applies.

(b) Yes. The same argument applies.

(c) The same argument applies for quantities that vary continuously, such as barometric pressure. But one could argue that altitude above sea level is sometimes discontinuous, so the result might not always hold for that quantity.

12. $$g'(x) = \lim_{h\to 0} \frac{g(x+h)-g(x)}{h} = \lim_{h\to 0} \frac{(x+h)f(x+h)-xf(x)}{h} = \lim_{h\to 0}\left[\frac{xf(x+h)-xf\,(x)}{h} + \frac{hf(x+h)}{h}\right]$$
$$= x\lim_{h\to 0}\frac{f(x+h)-f(x)}{h} + \lim_{h\to 0} f(x+h) = xf'(x) + f(x)$$

because f is differentiable and therefore continuous.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. (a) Put $x = 0$ and $y = 0$ in the equation: $f(0+0) = f(0) + f(0) + 0^2 \cdot 0 + 0 \cdot 0^2 \quad \Rightarrow \quad f(0) = 2f(0)$.

Subtracting $f(0)$ from each side of this equation gives $f(0) = 0$.

(b) $f'(0) = \lim\limits_{h \to 0} \dfrac{f(0+h) - f(0)}{h} = \lim\limits_{h \to 0} \dfrac{\left[f(0) + f(h) + 0^2 h + 0h^2\right] - f(0)}{h} = \lim\limits_{h \to 0} \dfrac{f(h)}{h} = \lim\limits_{x \to 0} \dfrac{f(x)}{x} = 1$

(c) $f'(x) = \lim\limits_{h \to 0} \dfrac{f(x+h) - f(x)}{h} = \lim\limits_{h \to 0} \dfrac{\left[f(x) + f(h) + x^2 h + xh^2\right] - f(x)}{h} = \lim\limits_{h \to 0} \dfrac{f(h) + x^2 h + xh^2}{h}$

$= \lim\limits_{h \to 0} \left[\dfrac{f(h)}{h} + x^2 + xh\right] = 1 + x^2$

14. We are given that $|f(x)| \le x^2$ for all x. In particular, $|f(0)| \le 0$, but $|a| \ge 0$ for all a. The only conclusion is that $f(0) = 0$. Now $\left|\dfrac{f(x) - f(0)}{x - 0}\right| = \left|\dfrac{f(x)}{x}\right| = \dfrac{|f(x)|}{|x|} \le \dfrac{x^2}{|x|} = \dfrac{|x^2|}{|x|} = |x| \quad \Rightarrow \quad -|x| \le \dfrac{f(x) - f(0)}{x - 0} \le |x|$.

But $\lim\limits_{x \to 0} (-|x|) = 0 = \lim\limits_{x \to 0} |x|$, so by the Squeeze Theorem, $\lim\limits_{x \to 0} \dfrac{f(x) - f(0)}{x - 0} = 0$. So by the definition of a derivative, f is differentiable at 0 and, furthermore, $f'(0) = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3 □ DIFFERENTIATION RULES

3.1 Derivatives of Polynomials and Exponential Functions

1. (a) e is the number such that $\lim\limits_{h\to 0} \dfrac{e^h - 1}{h} = 1$.

(b)

x	$\dfrac{2.7^x - 1}{x}$
-0.001	0.9928
-0.0001	0.9932
0.001	0.9937
0.0001	0.9933

x	$\dfrac{2.8^x - 1}{x}$
-0.001	1.0291
-0.0001	1.0296
0.001	1.0301
0.0001	1.0297

From the tables (to two decimal places),

$\lim\limits_{h\to 0} \dfrac{2.7^h - 1}{h} = 0.99$ and $\lim\limits_{h\to 0} \dfrac{2.8^h - 1}{h} = 1.03$.

Since $0.99 < 1 < 1.03$, $2.7 < e < 2.8$.

2. (a)

The function value at $x = 0$ is 1 and the slope at $x = 0$ is 1.

(b) $f(x) = e^x$ is an exponential function and $g(x) = x^e$ is a power function. $\dfrac{d}{dx}(e^x) = e^x$ and $\dfrac{d}{dx}(x^e) = ex^{e-1}$.

(c) $f(x) = e^x$ grows more rapidly than $g(x) = x^e$ when x is large.

3. $g(x) = 4x + 7 \quad\Rightarrow\quad g'(x) = 4(1) + 0 = 4$

4. $g(t) = 5t + 4t^2 \quad\Rightarrow\quad g'(t) = 5(1) + 4(2t^{2-1}) = 5(1) + 4(2t) = 5 + 8t$

5. $f(x) = x^{75} - x + 3 \quad\Rightarrow\quad f'(x) = 75x^{75-1} - 1(1) + 0 = 75x^{74} - 1$

6. $g(x) = \frac{7}{4}x^2 - 3x + 12 \quad\Rightarrow\quad g'(x) = \frac{7}{4}(2x^{2-1}) - 3(1) + 0 = \frac{7}{4}(2x) - 3 = \frac{7}{2}x - 3$

7. $f(t) = -2e^t \quad\Rightarrow\quad f'(t) = -2(e^t) = -2e^t$

8. $F(t) = t^3 + e^3 \quad\Rightarrow\quad F'(t) = 3t^2 + 0 = 3t^2$ [Note that e^3 is constant, so its derivative is zero.]

9. $W(v) = 1.8v^{-3} \quad\Rightarrow\quad W'(v) = 1.8(-3v^{-3-1}) = 1.8(-3v^{-4}) = -5.4v^{-4}$

10. $r(z) = z^{-5} - z^{1/2} \quad\Rightarrow\quad r'(z) = -5z^{-6} - \frac{1}{2}z^{-1/2}$

11. $f(x) = x^{3/2} + x^{-3} \quad\Rightarrow\quad f'(x) = \frac{3}{2}x^{1/2} + (-3x^{-4}) = \frac{3}{2}x^{1/2} - 3x^{-4}$

12. $V(t) = t^{-3/5} + t^4 \quad\Rightarrow\quad V'(t) = -\frac{3}{5}t^{-8/5} + 4t^3$

13. $s(t) = \dfrac{1}{t} + \dfrac{1}{t^2} = t^{-1} + t^{-2} \quad\Rightarrow\quad s'(t) = -t^{-2} + \left(-2t^{-3}\right) = -t^{-2} - 2t^{-3} = -\dfrac{1}{t^2} - \dfrac{2}{t^3}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $r(t) = \dfrac{a}{t^2} + \dfrac{b}{t^4} = at^{-2} + bt^{-4} \quad \Rightarrow \quad r'(t) = a(-2t^{-3}) + b(-4t^{-5}) = -2at^{-3} - 4bt^{-5} = -\dfrac{2a}{t^3} - \dfrac{4b}{t^5}$

15. $y = 2x + \sqrt{x} = 2x + x^{1/2} \quad \Rightarrow \quad y' = 2(1) + \frac{1}{2}x^{-1/2} = 2 + \frac{1}{2}x^{-1/2}$ or $2 + \dfrac{1}{2\sqrt{x}}$

16. $h(w) = \sqrt{2}\,w - \sqrt{2} \quad \Rightarrow \quad h'(w) = \sqrt{2}\,(1) - 0 = \sqrt{2}$

17. $g(x) = \dfrac{1}{\sqrt{x}} + \sqrt[4]{x} = x^{-1/2} + x^{1/4} \quad \Rightarrow \quad g'(x) = -\frac{1}{2}x^{-3/2} + \frac{1}{4}x^{-3/4}$ or $-\dfrac{1}{2x\sqrt{x}} + \dfrac{1}{4\sqrt[4]{x^3}}$

18. $W(t) = \sqrt{t} - 2e^t = t^{1/2} - 2e^t \quad \Rightarrow \quad W'(t) = \frac{1}{2}t^{-1/2} - 2(e^t) = \frac{1}{2}t^{-1/2} - 2e^t$ or $\dfrac{1}{2\sqrt{t}} - 2e^t$

19. $f(x) = x^3(x+3) = x^4 + 3x^3 \quad \Rightarrow \quad f'(x) = 4x^3 + 3(3x^2) = 4x^3 + 9x^2$

20. $F(t) = (2t-3)^2 = 4t^2 - 12t + 9 \quad \Rightarrow \quad F'(t) = 4(2t) - 12(1) + 0 = 8t - 12$

21. $y = 3e^x + \dfrac{4}{\sqrt[3]{x}} = 3e^x + 4x^{-1/3} \quad \Rightarrow \quad y' = 3(e^x) + 4(-\frac{1}{3})x^{-4/3} = 3e^x - \frac{4}{3}x^{-4/3}$ or $3e^x - \dfrac{4}{3x\sqrt[3]{x}}$

22. $S(R) = 4\pi R^2 \quad \Rightarrow \quad S'(R) = 4\pi(2R) = 8\pi R$

23. $f(x) = \dfrac{3x^2 + x^3}{x} = \dfrac{3x^2}{x} + \dfrac{x^3}{x} = 3x + x^2 \quad \Rightarrow \quad f'(x) = 3(1) + 2x = 3 + 2x$

24. $y = \dfrac{\sqrt{x} + x}{x^2} = \dfrac{\sqrt{x}}{x^2} + \dfrac{x}{x^2} = x^{1/2-2} + x^{1-2} = x^{-3/2} + x^{-1} \quad \Rightarrow \quad y' = -\frac{3}{2}x^{-5/2} + (-1x^{-2}) = -\frac{3}{2}x^{-5/2} - x^{-2}$

25. $G(r) = \dfrac{3r^{3/2} + r^{5/2}}{r} = \dfrac{3r^{3/2}}{r} + \dfrac{r^{5/2}}{r} = 3r^{3/2-2/2} + r^{5/2-2/2} = 3r^{1/2} + r^{3/2} \quad \Rightarrow$

$G'(r) = 3\left(\frac{1}{2}r^{-1/2}\right) + \frac{3}{2}r^{1/2} = \frac{3}{2}r^{-1/2} + \frac{3}{2}r^{1/2}$ or $\dfrac{3}{2\sqrt{r}} + \dfrac{3}{2}\sqrt{r}$

26. $G(t) = \sqrt{5t} + \dfrac{\sqrt{7}}{t} = \sqrt{5}\,t^{1/2} + \sqrt{7}\,t^{-1} \quad \Rightarrow \quad G'(t) = \sqrt{5}\left(\frac{1}{2}t^{-1/2}\right) + \sqrt{7}\left(-1t^{-2}\right) = \dfrac{\sqrt{5}}{2\sqrt{t}} - \dfrac{\sqrt{7}}{t^2}$

27. $j(x) = x^{2.4} + e^{2.4} \quad \Rightarrow \quad j'(x) = 2.4x^{1.4} + 0 = 2.4x^{1.4}$

28. $k(r) = e^r + r^e \quad \Rightarrow \quad k'(r) = e^r + er^{e-1}$

29. $F(z) = \dfrac{A + Bz + Cz^2}{z^2} = \dfrac{A}{z^2} + \dfrac{Bz}{z^2} + \dfrac{Cz^2}{z^2} = Az^{-2} + Bz^{-1} + C \quad \Rightarrow$

$F'(z) = A(-2z^{-3}) + B(-1z^{-2}) + 0 = -2Az^{-3} - Bz^{-2} = -\dfrac{2A}{z^3} - \dfrac{B}{z^2}$ or $-\dfrac{2A + Bz}{z^3}$

30. $G(q) = \left(1 + q^{-1}\right)^2 = 1 + 2q^{-1} + q^{-2} \quad \Rightarrow \quad G'(q) = 0 + 2(-1q^{-2}) + (-2q^{-3}) = -2q^{-2} - 2q^{-3}$

31. $D(t) = \dfrac{1 + 16t^2}{(4t)^3} = \dfrac{1 + 16t^2}{64t^3} = \dfrac{1}{64}t^{-3} + \dfrac{1}{4}t^{-1} \quad \Rightarrow$

$D'(t) = \frac{1}{64}(-3t^{-4}) + \frac{1}{4}(-1t^{-2}) = -\frac{3}{64}t^{-4} - \frac{1}{4}t^{-2}$ or $-\dfrac{3}{64t^4} - \dfrac{1}{4t^2}$ or $-\dfrac{3 + 16t^2}{64t^4}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. $f(v) = \dfrac{\sqrt[3]{v} - 2ve^v}{v} = \dfrac{\sqrt[3]{v}}{v} - \dfrac{2ve^v}{v} = v^{-2/3} - 2e^v \quad \Rightarrow \quad f'(v) = -\frac{2}{3}v^{-5/3} - 2e^v$

33. $P(w) = \dfrac{2w^2 - w + 4}{\sqrt{w}} = \dfrac{2w^2}{\sqrt{w}} - \dfrac{w}{\sqrt{w}} + \dfrac{4}{\sqrt{w}} = 2w^{2-1/2} - w^{1-1/2} + 4w^{-1/2} = 2w^{3/2} - w^{1/2} + 4w^{-1/2} \quad \Rightarrow$

$P'(w) = 2\left(\frac{3}{2}w^{1/2}\right) - \frac{1}{2}w^{-1/2} + 4\left(-\frac{1}{2}w^{-3/2}\right) = 3w^{1/2} - \frac{1}{2}w^{-1/2} - 2w^{-3/2}$ or $3\sqrt{w} - \dfrac{1}{2\sqrt{w}} - \dfrac{2}{w\sqrt{w}}$

34. $y = e^{x+1} + 1 = e^x e^1 + 1 = e \cdot e^x + 1 \quad \Rightarrow \quad y' = e \cdot e^x = e^{x+1}$

35. $y = tx^2 + t^3x$.

To find dy/dx, we treat t as a constant and x as a variable to get $dy/dx = t(2x) + t^3(1) = 2tx + t^3$.

To find dy/dt, we treat x as a constant and t as a variable to get $dy/dt = (1)x^2 + (3t^2)x = x^2 + 3t^2x$.

36. $y = \dfrac{t}{x^2} + \dfrac{x}{t} = tx^{-2} + xt^{-1}$.

To find dy/dx, we treat t as a constant and x as a variable to get $dy/dx = t(-2x^{-3}) + (1)t^{-1} = -2tx^{-3} + t^{-1}$ or $-\dfrac{2t}{x^3} + \dfrac{1}{t}$.

To find dy/dt, we treat x as a constant and t as a variable to get $dy/dt = (1)x^{-2} + x(-1t^{-2}) = x^{-2} - xt^{-2}$ or $\dfrac{1}{x^2} - \dfrac{x}{t^2}$.

37. $y = 2x^3 - x^2 + 2 \quad \Rightarrow \quad y' = 6x^2 - 2x$. At $(1, 3)$, $y' = 6(1)^2 - 2(1) = 4$ and an equation of the tangent line is $y - 3 = 4(x - 1)$ or $y = 4x - 1$.

38. $y = 2e^x + x \quad \Rightarrow \quad y' = 2e^x + 1$. At $(0, 2)$, $y' = 2e^0 + 1 = 3$ and an equation of the tangent line is $y - 2 = 3(x - 0)$ or $y = 3x + 2$.

39. $y = x + \dfrac{2}{x} = x + 2x^{-1} \quad \Rightarrow \quad y' = 1 - 2x^{-2}$. At $(2, 3)$, $y' = 1 - 2(2)^{-2} = \frac{1}{2}$ and an equation of the tangent line is $y - 3 = \frac{1}{2}(x - 2)$ or $y = \frac{1}{2}x + 2$.

40. $y = \sqrt[4]{x} - x = x^{1/4} - x \quad \Rightarrow \quad y' = \frac{1}{4}x^{-3/4} - 1 = \dfrac{1}{4\sqrt[4]{x^3}} - 1$. At $(1, 0)$, $y' = \frac{1}{4} - 1 = -\frac{3}{4}$ and an equation of the tangent line is $y - 0 = -\frac{3}{4}(x - 1)$ or $y = -\frac{3}{4}x + \frac{3}{4}$.

41. $y = x^4 + 2e^x \quad \Rightarrow \quad y' = 4x^3 + 2e^x$. At $(0, 2)$, $y' = 2$ and an equation of the tangent line is $y - 2 = 2(x - 0)$ or $y = 2x + 2$. The slope of the normal line is $-\frac{1}{2}$ (the negative reciprocal of 2) and an equation of the normal line is $y - 2 = -\frac{1}{2}(x - 0)$ or $y = -\frac{1}{2}x + 2$.

42. $y = x^{3/2} \quad \Rightarrow \quad y' = \frac{3}{2}x^{1/2}$. At $(1, 1)$, $y' = \frac{3}{2}$ and an equation of the tangent line is $y - 1 = \frac{3}{2}(x - 1)$ or $y = \frac{3}{2}x - \frac{1}{2}$. The slope of the normal line is $-\frac{2}{3}$ $\left(\text{the negative reciprocal of } \frac{3}{2}\right)$ and an equation of the normal line is $y - 1 = -\frac{2}{3}(x - 1)$ or $y = -\frac{2}{3}x + \frac{5}{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. $y = 3x^2 - x^3 \quad \Rightarrow \quad y' = 6x - 3x^2$.

At $(1, 2)$, $y' = 6 - 3 = 3$, so an equation of the tangent line is $y - 2 = 3(x - 1)$ or $y = 3x - 1$.

44. $y = x - \sqrt{x} \quad \Rightarrow \quad y' = 1 - \frac{1}{2}x^{-1/2} = 1 - \dfrac{1}{2\sqrt{x}}$.

At $(1, 0)$, $y' = \frac{1}{2}$, so an equation of the tangent line is $y - 0 = \frac{1}{2}(x - 1)$ or $y = \frac{1}{2}x - \frac{1}{2}$.

45. $f(x) = x^4 - 2x^3 + x^2 \quad \Rightarrow \quad f'(x) = 4x^3 - 6x^2 + 2x$

Note that $f'(x) = 0$ when f has a horizontal tangent, f' is positive when f is increasing, and f' is negative when f is decreasing.

46. $f(x) = x^5 - 2x^3 + x - 1 \quad \Rightarrow \quad f'(x) = 5x^4 - 6x^2 + 1$

Note that $f'(x) = 0$ when f has a horizontal tangent, f' is positive when f is increasing, and f' is negative when f is decreasing.

47. (a)

(b) From the graph in part (a), it appears that f' is zero at $x_1 \approx -1.25$, $x_2 \approx 0.5$, and $x_3 \approx 3$. The slopes are negative (so f' is negative) on $(-\infty, x_1)$ and (x_2, x_3). The slopes are positive (so f' is positive) on (x_1, x_2) and (x_3, ∞).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $f(x) = x^4 - 3x^3 - 6x^2 + 7x + 30 \quad \Rightarrow$

$f'(x) = 4x^3 - 9x^2 - 12x + 7$

48. (a)

(b) From the graph in part (a), it appears that f' is zero at $x_1 \approx 0.2$ and $x_2 \approx 2.8$. The slopes are positive (so f' is positive) on $(-\infty, x_1)$ and (x_2, ∞). The slopes are negative (so f' is negative) on (x_1, x_2).

(c) $g(x) = e^x - 3x^2 \quad \Rightarrow \quad g'(x) = e^x - 6x$

49. $f(x) = 0.001x^5 - 0.02x^3 \quad \Rightarrow \quad f'(x) = 0.005x^4 - 0.06x^2 \quad \Rightarrow \quad f''(x) = 0.02x^3 - 0.12x$

50. $G(r) = \sqrt{r} + \sqrt[3]{r} \quad \Rightarrow \quad G'(r) = \frac{1}{2}r^{-1/2} + \frac{1}{3}r^{-2/3} \quad \Rightarrow \quad G''(r) = -\frac{1}{4}r^{-3/2} - \frac{2}{9}r^{-5/3}$

51. $f(x) = 2x - 5x^{3/4} \quad \Rightarrow \quad f'(x) = 2 - \frac{15}{4}x^{-1/4} \quad \Rightarrow \quad f''(x) = \frac{15}{16}x^{-5/4}$

Note that f' is negative when f is decreasing and positive when f is increasing. f'' is always positive since f' is always increasing.

52. $f(x) = e^x - x^3 \quad \Rightarrow \quad f'(x) = e^x - 3x^2 \quad \Rightarrow \quad f''(x) = e^x - 6x$

Note that $f'(x) = 0$ when f has a horizontal tangent and that $f''(x) = 0$ when f' has a horizontal tangent.

53. (a) $s = t^3 - 3t \quad \Rightarrow \quad v(t) = s'(t) = 3t^2 - 3 \quad \Rightarrow \quad a(t) = v'(t) = 6t$

(b) $a(2) = 6(2) = 12 \text{ m/s}^2$

(c) $v(t) = 3t^2 - 3 = 0$ when $t^2 = 1$, that is, $t = 1 \ \ [t \geq 0]$ and $a(1) = 6 \text{ m/s}^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. (a) $s = t^4 - 2t^3 + t^2 - t \quad \Rightarrow$

$v(t) = s'(t) = 4t^3 - 6t^2 + 2t - 1 \quad \Rightarrow$

$a(t) = v'(t) = 12t^2 - 12t + 2$

(b) $a(1) = 12(1)^2 - 12(1) + 2 = 2\,\mathrm{m/s^2}$

(c)

55. $L = 0.0155A^3 - 0.372A^2 + 3.95A + 1.21 \quad \Rightarrow \quad \dfrac{dL}{dA} = 0.0465A^2 - 0.744A + 3.95$, so

$\left.\dfrac{dL}{dA}\right|_{A=12} = 0.0465(12)^2 - 0.744(12) + 3.95 = 1.718$. The derivative is the instantaneous rate of change of the length (in inches) of an Alaskan rockfish with respect to its age when its age is 12 years. Its units are $\mathrm{inches/year}$.

56. $S(A) = 0.882A^{0.842} \quad \Rightarrow \quad S'(A) = 0.882(0.842A^{-0.158}) = 0.742644A^{-0.158}$, so

$S'(100) = 0.742644(100)^{-0.158} \approx 0.36$. The derivative is the instantaneous rate of change of the number of tree species with respect to area. Its units are number of species per square meter.

57. (a) $P = \dfrac{k}{V}$ and $P = 50$ when $V = 0.106$, so $k = PV = 50(0.106) = 5.3$. Thus, $P = \dfrac{5.3}{V}$ and $V = \dfrac{5.3}{P}$.

(b) $V = 5.3P^{-1} \quad \Rightarrow \quad \dfrac{dV}{dP} = 5.3(-1P^{-2}) = -\dfrac{5.3}{P^2}$. When $P = 50$, $\dfrac{dV}{dP} = -\dfrac{5.3}{50^2} = -0.00212$. The derivative is the instantaneous rate of change of the volume with respect to the pressure at $25^\circ\mathrm{C}$. Its units are $\mathrm{m^3/kPa}$.

58. (a) $L = aP^2 + bP + c$, where $a \approx -0.275428$, $b \approx 19.74853$, and $c \approx -273.55234$.

(b) $\dfrac{dL}{dP} = 2aP + b$. When $P = 30$, $\dfrac{dL}{dP} \approx 3.2$, and when $P = 40$, $\dfrac{dL}{dP} \approx -2.3$. The derivative is the instantaneous rate of change of tire life with respect to pressure. Its units are (thousands of miles)/$(\mathrm{lb/in^2})$. When $\dfrac{dL}{dP}$ is positive, tire life is increasing, and when $\dfrac{dL}{dP} < 0$, tire life is decreasing.

59. $y = x^3 + 3x^2 - 9x + 10 \quad \Rightarrow \quad y' = 3x^2 + 3(2x) - 9(1) + 0 = 3x^2 + 6x - 9$. Horizontal tangents occur where $y' = 0$. Thus, $3x^2 + 6x - 9 = 0 \quad \Rightarrow \quad 3(x^2 + 2x - 3) = 0 \quad \Rightarrow \quad 3(x+3)(x-1) = 0 \quad \Rightarrow \quad x = -3$ or $x = 1$. The corresponding points are $(-3, 37)$ and $(1, 5)$.

60. $f(x) = e^x - 2x \quad \Rightarrow \quad f'(x) = e^x - 2$. $f'(x) = 0 \quad \Rightarrow \quad e^x = 2 \quad \Rightarrow \quad x = \ln 2$, so f has a horizontal tangent when $x = \ln 2$.

61. $y = 2e^x + 3x + 5x^3 \quad \Rightarrow \quad y' = 2e^x + 3 + 15x^2$. Since $2e^x > 0$ and $15x^2 \geq 0$, we must have $y' > 0 + 3 + 0 = 3$, so no tangent line can have slope 2.

62. $y = x^4 + 1 \quad \Rightarrow \quad y' = 4x^3$. The slope of the line $32x - y = 15$ (or $y = 32x - 15$) is 32, so the slope of any line parallel to it is also 32. Thus, $y' = 32 \quad \Leftrightarrow \quad 4x^3 = 32 \quad \Leftrightarrow \quad x^3 = 8 \quad \Leftrightarrow \quad x = 2$, which is the x-coordinate of the point on the curve at which the slope is 32. The y-coordinate is $2^4 + 1 = 17$, so an equation of the tangent line is $y - 17 = 32(x - 2)$ or $y = 32x - 47$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

63. The slope of the line $3x - y = 15$ (or $y = 3x - 15$) is 3, so the slope of both tangent lines to the curve is 3. $y = x^3 - 3x^2 + 3x - 3 \;\Rightarrow\; y' = 3x^2 - 6x + 3 = 3(x^2 - 2x + 1) = 3(x-1)^2$. Thus, $3(x-1)^2 = 3 \;\Rightarrow\; (x-1)^2 = 1 \;\Rightarrow\; x - 1 = \pm 1 \;\Rightarrow\; x = 0$ or 2, which are the x-coordinates at which the tangent lines have slope 3. The points on the curve are $(0, -3)$ and $(2, -1)$, so the tangent line equations are $y - (-3) = 3(x - 0)$ or $y = 3x - 3$ and $y - (-1) = 3(x - 2)$ or $y = 3x - 7$.

64. The slope of $y = 1 + 2e^x - 3x$ is given by $m = y' = 2e^x - 3$.

The slope of $3x - y = 5 \;\Leftrightarrow\; y = 3x - 5$ is 3.

$m = 3 \;\Rightarrow\; 2e^x - 3 = 3 \;\Rightarrow\; e^x = 3 \;\Rightarrow\; x = \ln 3$.

This occurs at the point $(\ln 3, 7 - 3\ln 3) \approx (1.1, 3.7)$.

65. The slope of $y = \sqrt{x}$ is given by $y = \frac{1}{2}x^{-1/2} = \dfrac{1}{2\sqrt{x}}$. The slope of $2x + y = 1$ (or $y = -2x + 1$) is -2, so the desired normal line must have slope -2, and hence, the tangent line to the curve must have slope $\frac{1}{2}$. This occurs if $\dfrac{1}{2\sqrt{x}} = \dfrac{1}{2} \;\Rightarrow\; \sqrt{x} = 1 \;\Rightarrow\; x = 1$. When $x = 1$, $y = \sqrt{1} = 1$, and an equation of the normal line is $y - 1 = -2(x - 1)$ or $y = -2x + 3$.

66. $y = f(x) = x^2 - 1 \;\Rightarrow\; f'(x) = 2x$. So $f'(-1) = -2$, and the slope of the normal line is $\frac{1}{2}$. The equation of the normal line at $(-1, 0)$ is $y - 0 = \frac{1}{2}[x - (-1)]$ or $y = \frac{1}{2}x + \frac{1}{2}$. Substituting this into the equation of the parabola, we obtain $\frac{1}{2}x + \frac{1}{2} = x^2 - 1 \;\Leftrightarrow\; x + 1 = 2x^2 - 2 \;\Leftrightarrow\; 2x^2 - x - 3 = 0 \;\Leftrightarrow\; (2x - 3)(x + 1) = 0 \;\Leftrightarrow\; x = \frac{3}{2}$ or -1. Substituting $\frac{3}{2}$ into the equation of the normal line gives us $y = \frac{5}{4}$. Thus, the second point of intersection is $\left(\frac{3}{2}, \frac{5}{4}\right)$, as shown in the sketch.

67.

Let (a, a^2) be a point on the parabola at which the tangent line passes through the point $(0, -4)$. The tangent line has slope $2a$ and equation $y - (-4) = 2a(x - 0) \;\Leftrightarrow\; y = 2ax - 4$. Since (a, a^2) also lies on the line, $a^2 = 2a(a) - 4$, or $a^2 = 4$. So $a = \pm 2$ and the points are $(2, 4)$ and $(-2, 4)$.

68. (a) If $y = x^2 + x$, then $y' = 2x + 1$. If the point at which a tangent meets the parabola is $(a, a^2 + a)$, then the slope of the tangent is $2a + 1$. But since it passes through $(2, -3)$, the slope must also be $\dfrac{\Delta y}{\Delta x} = \dfrac{a^2 + a + 3}{a - 2}$.

Therefore, $2a + 1 = \dfrac{a^2 + a + 3}{a - 2}$. Solving this equation for a we get $a^2 + a + 3 = 2a^2 - 3a - 2 \;\Leftrightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$a^2 - 4a - 5 = (a-5)(a+1) = 0 \quad \Leftrightarrow \quad a = 5$ or -1. If $a = -1$, the point is $(-1, 0)$ and the slope is -1, so the equation is $y - 0 = (-1)(x+1)$ or $y = -x - 1$. If $a = 5$, the point is $(5, 30)$ and the slope is 11, so the equation is $y - 30 = 11(x-5)$ or $y = 11x - 25$.

(b) As in part (a), but using the point $(2, 7)$, we get the equation

$$2a + 1 = \frac{a^2 + a - 7}{a - 2} \quad \Rightarrow \quad 2a^2 - 3a - 2 = a^2 + a - 7 \quad \Leftrightarrow \quad a^2 - 4a + 5 = 0.$$

The last equation has no real solution (discriminant $= -16 < 0$), so there is no line through the point $(2, 7)$ that is tangent to the parabola. The diagram shows that the point $(2, 7)$ is "inside" the parabola, but tangent lines to the parabola do not pass through points inside the parabola.

69. $f'(x) = \lim\limits_{h \to 0} \dfrac{f(x+h) - f(x)}{h} = \lim\limits_{h \to 0} \dfrac{\dfrac{1}{x+h} - \dfrac{1}{x}}{h} = \lim\limits_{h \to 0} \dfrac{x - (x+h)}{hx(x+h)} = \lim\limits_{h \to 0} \dfrac{-h}{hx(x+h)} = \lim\limits_{h \to 0} \dfrac{-1}{x(x+h)} = -\dfrac{1}{x^2}$

70. (a) $f(x) = x^n \quad \Rightarrow \quad f'(x) = nx^{n-1} \quad \Rightarrow \quad f''(x) = n(n-1)x^{n-2} \quad \Rightarrow \quad \cdots \quad \Rightarrow$

$f^{(n)}(x) = n(n-1)(n-2)\cdots 2 \cdot 1x^{n-n} = n!$

(b) $f(x) = x^{-1} \quad \Rightarrow \quad f'(x) = (-1)x^{-2} \quad \Rightarrow \quad f''(x) = (-1)(-2)x^{-3} \quad \Rightarrow \quad \cdots \quad \Rightarrow$

$f^{(n)}(x) = (-1)(-2)(-3)\cdots(-n)x^{-(n+1)} = (-1)^n n! x^{-(n+1)}$ or $\dfrac{(-1)^n\, n!}{x^{n+1}}$

71. Let $P(x) = ax^2 + bx + c$. Then $P'(x) = 2ax + b$ and $P''(x) = 2a$. $P''(2) = 2 \quad \Rightarrow \quad 2a = 2 \quad \Rightarrow \quad a = 1$.
$P'(2) = 3 \quad \Rightarrow \quad 2(1)(2) + b = 3 \quad \Rightarrow \quad 4 + b = 3 \quad \Rightarrow \quad b = -1$.
$P(2) = 5 \quad \Rightarrow \quad 1(2)^2 + (-1)(2) + c = 5 \quad \Rightarrow \quad 2 + c = 5 \quad \Rightarrow \quad c = 3$. So $P(x) = x^2 - x + 3$.

72. $y = Ax^2 + Bx + C \quad \Rightarrow \quad y' = 2Ax + B \quad \Rightarrow \quad y'' = 2A$. We substitute these expressions into the equation $y'' + y' - 2y = x^2$ to get

$$\begin{aligned} (2A) + (2Ax + B) - 2(Ax^2 + Bx + C) &= x^2 \\ 2A + 2Ax + B - 2Ax^2 - 2Bx - 2C &= x^2 \\ (-2A)x^2 + (2A - 2B)x + (2A + B - 2C) &= (1)x^2 + (0)x + (0) \end{aligned}$$

The coefficients of x^2 on each side must be equal, so $-2A = 1 \quad \Rightarrow \quad A = -\frac{1}{2}$. Similarly, $2A - 2B = 0 \quad \Rightarrow$ $A = B = -\frac{1}{2}$ and $2A + B - 2C = 0 \quad \Rightarrow \quad -1 - \frac{1}{2} - 2C = 0 \quad \Rightarrow \quad C = -\frac{3}{4}$.

73. $y = f(x) = ax^3 + bx^2 + cx + d \quad \Rightarrow \quad f'(x) = 3ax^2 + 2bx + c$. The point $(-2, 6)$ is on f, so $f(-2) = 6 \quad \Rightarrow$ $-8a + 4b - 2c + d = 6$ **(1)**. The point $(2, 0)$ is on f, so $f(2) = 0 \quad \Rightarrow \quad 8a + 4b + 2c + d = 0$ **(2)**. Since there are horizontal tangents at $(-2, 6)$ and $(2, 0)$, $f'(\pm 2) = 0$. $f'(-2) = 0 \quad \Rightarrow \quad 12a - 4b + c = 0$ **(3)** and $f'(2) = 0 \quad \Rightarrow$ $12a + 4b + c = 0$ **(4)**. Subtracting equation **(3)** from **(4)** gives $8b = 0 \quad \Rightarrow \quad b = 0$. Adding **(1)** and **(2)** gives $8b + 2d = 6$, so $d = 3$ since $b = 0$. From **(3)** we have $c = -12a$, so **(2)** becomes $8a + 4(0) + 2(-12a) + 3 = 0 \quad \Rightarrow \quad 3 = 16a \quad \Rightarrow$ $a = \frac{3}{16}$. Now $c = -12a = -12\left(\frac{3}{16}\right) = -\frac{9}{4}$ and the desired cubic function is $y = \frac{3}{16}x^3 - \frac{9}{4}x + 3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

74. $y = ax^2 + bx + c \quad \Rightarrow \quad y'(x) = 2ax + b$. The parabola has slope 4 at $x = 1$ and slope -8 at $x = -1$, so $y'(1) = 4 \quad \Rightarrow$ $2a + b = 4$ **(1)** and $y'(-1) = -8 \quad \Rightarrow \quad -2a + b = -8$ **(2)**. Adding **(1)** and **(2)** gives us $2b = -4 \quad \Leftrightarrow \quad b = -2$. From **(1)**, $2a - 2 = 4 \quad \Leftrightarrow \quad a = 3$. Thus, the equation of the parabola is $y = 3x^2 - 2x + c$. Since it passes through the point $(2, 15)$, we have $15 = 3(2)^2 - 2(2) + c \quad \Rightarrow \quad c = 7$, so the equation is $y = 3x^2 - 2x + 7$.

75. $f(x) = \begin{cases} x^2 + 1 & \text{if } x < 1 \\ x + 1 & \text{if } x \geq 1 \end{cases}$

Calculate the left- and right-hand derivatives as defined in Exercise 2.8.64:

$$f'_-(1) = \lim_{h \to 0^-} \frac{f(1+h) - f(1)}{h} = \lim_{h \to 0^-} \frac{[(1+h)^2 + 1] - (1+1)}{h} = \lim_{h \to 0^-} \frac{h^2 + 2h}{h} = \lim_{h \to 0^-} (h + 2) = 2 \text{ and}$$

$$f'_+(1) = \lim_{h \to 0^+} \frac{f(1+h) - f(1)}{h} = \lim_{h \to 0^+} \frac{[(1+h) + 1] - (1+1)}{h} = \lim_{h \to 0^+} \frac{h}{h} = \lim_{h \to 0^+} 1 = 1.$$

Since the left and right limits are different, $\lim\limits_{h \to 0} \dfrac{f(1+h) - f(1)}{h}$ does not exist, that is, $f'(1)$ does not exist. Therefore, f is not differentiable at 1.

76. $g(x) = \begin{cases} 2x & \text{if } x \leq 0 \\ 2x - x^2 & \text{if } 0 < x < 2 \\ 2 - x & \text{if } x \geq 2 \end{cases}$

Investigate the left- and right-hand derivatives at $x = 0$ and $x = 2$:

$$g'_-(0) = \lim_{h \to 0^-} \frac{g(0+h) - g(0)}{h} = \lim_{h \to 0^-} \frac{2h - 2(0)}{h} = 2 \text{ and}$$

$$g'_+(0) = \lim_{h \to 0^+} \frac{g(0+h) - g(0)}{h} = \lim_{h \to 0^+} \frac{(2h - h^2) - 2(0)}{h} = \lim_{h \to 0^+} (2 - h) = 2, \text{ so } g \text{ is differentiable at } x = 0.$$

$$g'_-(2) = \lim_{h \to 0^-} \frac{g(2+h) - g(2)}{h} = \lim_{h \to 0^-} \frac{2(2+h) - (2+h)^2 - (2-2)}{h} = \lim_{h \to 0^-} \frac{-2h - h^2}{h} = \lim_{h \to 0^-} (-2 - h) = -2$$

and

$$g'_+(2) = \lim_{h \to 0^+} \frac{g(2+h) - g(2)}{h} = \lim_{h \to 0^+} \frac{[2 - (2+h)] - (2-2)}{h} = \lim_{h \to 0^+} \frac{-h}{h} = \lim_{h \to 0^+} (-1) = -1,$$

so g is not differentiable at $x = 2$. Thus, a formula for g' is

$$g'(x) = \begin{cases} 2 & \text{if } x \leq 0 \\ 2 - 2x & \text{if } 0 < x < 2 \\ -1 & \text{if } x > 2 \end{cases}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

77. (a) Note that $x^2 - 9 < 0$ for $x^2 < 9 \quad \Leftrightarrow \quad |x| < 3 \quad \Leftrightarrow \quad -3 < x < 3$. So

$$f(x) = \begin{cases} x^2 - 9 & \text{if } x \le -3 \\ -x^2 + 9 & \text{if } -3 < x < 3 \\ x^2 - 9 & \text{if } x \ge 3 \end{cases} \quad \Rightarrow \quad f'(x) = \begin{cases} 2x & \text{if } x < -3 \\ -2x & \text{if } -3 < x < 3 \\ 2x & \text{if } x > 3 \end{cases} = \begin{cases} 2x & \text{if } |x| > 3 \\ -2x & \text{if } |x| < 3 \end{cases}$$

To show that $f'(3)$ does not exist we investigate $\lim\limits_{h \to 0} \dfrac{f(3+h) - f(3)}{h}$ by computing the left- and right-hand derivatives defined in Exercise 2.8.64.

$$f'_-(3) = \lim_{h \to 0^-} \frac{f(3+h) - f(3)}{h} = \lim_{h \to 0^-} \frac{[-(3+h)^2 + 9] - 0}{h} = \lim_{h \to 0^-} (-6 - h) = -6 \quad \text{and}$$

$$f'_+(3) = \lim_{h \to 0^+} \frac{f(3+h) - f(3)}{h} = \lim_{h \to 0^+} \frac{\left[(3+h)^2 - 9\right] - 0}{h} = \lim_{h \to 0^+} \frac{6h + h^2}{h} = \lim_{h \to 0^+} (6 + h) = 6.$$

Since the left and right limits are different, $\lim\limits_{h \to 0} \dfrac{f(3+h) - f(3)}{h}$ does not exist, that is, $f'(3)$ does not exist. Similarly, $f'(-3)$ does not exist. Therefore, f is not differentiable at 3 or at -3.

(b)

78. If $x \ge 1$, then $h(x) = |x - 1| + |x + 2| = x - 1 + x + 2 = 2x + 1$.

If $-2 < x < 1$, then $h(x) = -(x - 1) + x + 2 = 3$.

If $x \le -2$, then $h(x) = -(x - 1) - (x + 2) = -2x - 1$. Therefore,

$$h(x) = \begin{cases} -2x - 1 & \text{if } x \le -2 \\ 3 & \text{if } -2 < x < 1 \\ 2x + 1 & \text{if } x \ge 1 \end{cases} \quad \Rightarrow \quad h'(x) = \begin{cases} -2 & \text{if } x < -2 \\ 0 & \text{if } -2 < x < 1 \\ 2 & \text{if } x > 1 \end{cases}$$

To see that $h'(1) = \lim\limits_{x \to 1} \dfrac{h(x) - h(1)}{x - 1}$ does not exist, observe that $\lim\limits_{x \to 1^-} \dfrac{h(x) - h(1)}{x - 1} = \lim\limits_{x \to 1^-} \dfrac{3 - 3}{3 - 1} = 0$ but $\lim\limits_{x \to 1^+} \dfrac{h(x) - h(1)}{x - 1} = \lim\limits_{x \to 1^+} \dfrac{2x - 2}{x - 1} = 2$. Similarly, $h'(-2)$ does not exist.

79. Substituting $x = 1$ and $y = 1$ into $y = ax^2 + bx$ gives us $a + b = 1$ **(1)**. The slope of the tangent line $y = 3x - 2$ is 3 and the slope of the tangent to the parabola at (x, y) is $y' = 2ax + b$. At $x = 1$, $y' = 3 \quad \Rightarrow \quad 3 = 2a + b$ **(2)**. Subtracting **(1)** from **(2)** gives us $2 = a$ and it follows that $b = -1$. The parabola has equation $y = 2x^2 - x$.

80. $y = x^4 + ax^3 + bx^2 + cx + d \quad \Rightarrow \quad y(0) = d$. Since the tangent line $y = 2x + 1$ is equal to 1 at $x = 0$, we must have $d = 1$. $\quad y' = 4x^3 + 3ax^2 + 2bx + c \quad \Rightarrow \quad y'(0) = c$. Since the slope of the tangent line $y = 2x + 1$ at $x = 0$ is 2, we must have $c = 2$. Now $y(1) = 1 + a + b + c + d = a + b + 4$ and the tangent line $y = 2 - 3x$ at $x = 1$ has y-coordinate -1,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

so $a+b+4=-1$ or $a+b=-5$ **(1)**. Also, $y'(1)=4+3a+2b+c=3a+2b+6$ and the slope of the tangent line $y=2-3x$ at $x=1$ is -3, so $3a+2b+6=-3$ or $3a+2b=-9$ **(2)**. Adding -2 times **(1)** to **(2)** gives us $a=1$ and hence, $b=-6$. The curve has equation $y=x^4+x^3-6x^2+2x+1$.

81. $y=f(x)=ax^2 \quad\Rightarrow\quad f'(x)=2ax.$ So the slope of the tangent to the parabola at $x=2$ is $m=2a(2)=4a$. The slope of the given line, $2x+y=b \quad\Leftrightarrow\quad y=-2x+b$, is seen to be -2, so we must have $4a=-2 \quad\Leftrightarrow\quad a=-\frac{1}{2}$. So when $x=2$, the point in question has y-coordinate $-\frac{1}{2}\cdot 2^2=-2$. Now we simply require that the given line, whose equation is $2x+y=b$, pass through the point $(2,-2)$: $2(2)+(-2)=b \quad\Leftrightarrow\quad b=2$. So we must have $a=-\frac{1}{2}$ and $b=2$.

82. The slope of the curve $y=c\sqrt{x}$ is $y'=\dfrac{c}{2\sqrt{x}}$ and the slope of the tangent line $y=\frac{3}{2}x+6$ is $\frac{3}{2}$. These must be equal at the point of tangency $\left(a, c\sqrt{a}\right)$, so $\dfrac{c}{2\sqrt{a}}=\dfrac{3}{2} \quad\Rightarrow\quad c=3\sqrt{a}$. The y-coordinates must be equal at $x=a$, so $c\sqrt{a}=\frac{3}{2}a+6 \quad\Rightarrow\quad \left(3\sqrt{a}\right)\sqrt{a}=\frac{3}{2}a+6 \quad\Rightarrow\quad 3a=\frac{3}{2}a+6 \quad\Rightarrow\quad \frac{3}{2}a=6 \quad\Rightarrow\quad a=4$. Since $c=3\sqrt{a}$, we have $c=3\sqrt{4}=6$.

83. The line $y=2x+3$ has slope 2. The parabola $y=cx^2 \quad\Rightarrow\quad y'=2cx$ has slope $2ca$ at $x=a$. Equating slopes gives us $2ca=2$, or $ca=1$. Equating y-coordinates at $x=a$ gives us $ca^2=2a+3 \quad\Leftrightarrow\quad (ca)a=2a+3 \quad\Leftrightarrow\quad 1a=2a+3 \quad\Leftrightarrow\quad a=-3$. Thus, $c=\dfrac{1}{a}=-\dfrac{1}{3}$.

84. $f(x)=ax^2+bx+c \quad\Rightarrow\quad f'(x)=2ax+b$. The slope of the tangent line at $x=p$ is $2ap+b$, the slope of the tangent line at $x=q$ is $2aq+b$, and the average of those slopes is $\dfrac{(2ap+b)+(2aq+b)}{2}=ap+aq+b$. The midpoint of the interval $[p,q]$ is $\dfrac{p+q}{2}$ and the slope of the tangent line at the midpoint is $2a\left(\dfrac{p+q}{2}\right)+b=a(p+q)+b$. This is equal to $ap+aq+b$, as required.

85. f is clearly differentiable for $x<2$ and for $x>2$. For $x<2$, $f'(x)=2x$, so $f'_-(2)=4$. For $x>2$, $f'(x)=m$, so $f'_+(2)=m$. For f to be differentiable at $x=2$, we need $4=f'_-(2)=f'_+(2)=m$. So $f(x)=4x+b$. We must also have continuity at $x=2$, so $4=f(2)=\lim\limits_{x\to 2^+} f(x)=\lim\limits_{x\to 2^+}(4x+b)=8+b$. Hence, $b=-4$.

86. We have $g(x)=\begin{cases} ax^3-3x & \text{if} \quad x\le 1 \\ bx^2+2 & \text{if} \quad x>1 \end{cases}$

For $x<1$, $g'(x)=a(3x^2)-3(1)=3ax^2-3$, so $g'_-(1)=3a(1)^2-3=3a-3$. For $x>1$, $g'(x)=b(2x)+0=2bx$, so $g'_+(1)=2b(1)=2b$. For g to be differentiable at $x=1$, we need $g'_-(1)=g'_+(1)$, so $3a-3=2b$, or $b=\dfrac{3a-3}{2}$. For g to be continuous at $x=1$, we need $g_-(1)=a-3$ equal to $g_+(1)=b+2$. So we have the system of two equations: $a-3=b+2$, $b=\dfrac{3a-3}{2}$. Substituting the second equation into the first equation we have $a-3=\dfrac{3a-3}{2}+2 \quad\Rightarrow$ $2a-6=3a-3+4 \quad\Rightarrow\quad a=-7$ and $b=\dfrac{3(-7)-3}{2}=-12$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

87. *Solution 1:* Let $f(x) = x^{1000}$. Then, by the definition of a derivative, $f'(1) = \lim\limits_{x\to 1} \dfrac{f(x) - f(1)}{x - 1} = \lim\limits_{x\to 1} \dfrac{x^{1000} - 1}{x - 1}$.

But this is just the limit we want to find, and we know (from the Power Rule) that $f'(x) = 1000x^{999}$, so

$f'(1) = 1000(1)^{999} = 1000$. So $\lim\limits_{x\to 1} \dfrac{x^{1000} - 1}{x - 1} = 1000$.

Solution 2: Note that $(x^{1000} - 1) = (x - 1)(x^{999} + x^{998} + x^{997} + \cdots + x^2 + x + 1)$. So

$$\lim_{x\to 1} \frac{x^{1000} - 1}{x - 1} = \lim_{x\to 1} \frac{(x-1)(x^{999} + x^{998} + x^{997} + \cdots + x^2 + x + 1)}{x - 1} = \lim_{x\to 1}(x^{999} + x^{998} + x^{997} + \cdots + x^2 + x + 1)$$
$$= \underbrace{1 + 1 + 1 + \cdots + 1 + 1 + 1}_{1000 \text{ ones}} = 1000, \text{ as above.}$$

88. (a) $xy = c \quad \Rightarrow \quad y = \dfrac{c}{x}$. Let $P = \left(a, \dfrac{c}{a}\right)$. The slope of the tangent line at $x = a$ is $y'(a) = -\dfrac{c}{a^2}$. Its equation is

$y - \dfrac{c}{a} = -\dfrac{c}{a^2}(x - a)$ or $y = -\dfrac{c}{a^2}x + \dfrac{2c}{a}$, so its y-intercept is $\dfrac{2c}{a}$. Setting $y = 0$ gives $x = 2a$, so the x-intercept is $2a$.

The midpoint of the line segment joining $\left(0, \dfrac{2c}{a}\right)$ and $(2a, 0)$ is $\left(a, \dfrac{c}{a}\right) = P$.

(b) We know the x- and y-intercepts of the tangent line from part (a), so the area of the triangle bounded by the axes and the tangent is $\frac{1}{2}(\text{base})(\text{height}) = \frac{1}{2}xy = \frac{1}{2}(2a)(2c/a) = 2c$, a constant.

89. In order for the two tangents to intersect on the y-axis, the points of tangency must be at equal distances from the y-axis, since the parabola $y = x^2$ is symmetric about the y-axis. Say the points of tangency are (a, a^2) and $(-a, a^2)$, for some $a > 0$. Then since the derivative of $y = x^2$ is $dy/dx = 2x$, the left-hand tangent has slope $-2a$ and equation $y - a^2 = -2a(x + a)$, or $y = -2ax - a^2$, and similarly the right-hand tangent line has equation $y - a^2 = 2a(x - a)$, or $y = 2ax - a^2$. So the two lines intersect at $(0, -a^2)$. Now if the lines are perpendicular, then the product of their slopes is -1, so $(-2a)(2a) = -1 \quad \Leftrightarrow \quad a^2 = \frac{1}{4} \quad \Leftrightarrow \quad a = \frac{1}{2}$. So the lines intersect at $(0, -\frac{1}{4})$.

90.

From the sketch, it appears that there may be a line that is tangent to both curves. The slope of the line through the points $P(a, a^2)$ and $Q(b, b^2 - 2b + 2)$ is $\dfrac{b^2 - 2b + 2 - a^2}{b - a}$. The slope of the tangent line at P is $2a \quad [y' = 2x]$ and at Q is $2b - 2 \quad [y' = 2x - 2]$. All three slopes are equal, so $2a = 2b - 2 \quad \Leftrightarrow \quad a = b - 1$.

Also, $2b - 2 = \dfrac{b^2 - 2b + 2 - a^2}{b - a} \quad \Rightarrow \quad 2b - 2 = \dfrac{b^2 - 2b + 2 - (b-1)^2}{b - (b-1)} \quad \Rightarrow \quad 2b - 2 = b^2 - 2b + 2 - b^2 + 2b - 1 \quad \Rightarrow$

$2b = 3 \quad \Rightarrow \quad b = \frac{3}{2}$ and $a = \frac{3}{2} - 1 = \frac{1}{2}$. Thus, an equation of the tangent line at P is $y - \left(\frac{1}{2}\right)^2 = 2\left(\frac{1}{2}\right)\left(x - \frac{1}{2}\right)$ or $y = x - \frac{1}{4}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

91. $y = x^2 \quad \Rightarrow \quad y' = 2x$, so the slope of a tangent line at the point (a, a^2) is $y' = 2a$ and the slope of a normal line is $-1/(2a)$, for $a \neq 0$. The slope of the normal line through the points (a, a^2) and $(0, c)$ is $\dfrac{a^2 - c}{a - 0}$, so $\dfrac{a^2 - c}{a} = -\dfrac{1}{2a} \quad \Rightarrow$ $a^2 - c = -\frac{1}{2} \quad \Rightarrow \quad a^2 = c - \frac{1}{2}$. The last equation has two solutions if $c > \frac{1}{2}$, one solution if $c = \frac{1}{2}$, and no solution if $c < \frac{1}{2}$. Since the y-axis is normal to $y = x^2$ regardless of the value of c (this is the case for $a = 0$), we have *three* normal lines if $c > \frac{1}{2}$ and *one* normal line if $c \leq \frac{1}{2}$.

APPLIED PROJECT Building a Better Roller Coaster

1. (a) $f(x) = ax^2 + bx + c \quad \Rightarrow \quad f'(x) = 2ax + b$.

The origin is at P:	$f(0) = 0$	$\Rightarrow$	$c = 0$
The slope of the ascent is 0.8:	$f'(0) = 0.8$	$\Rightarrow$	$b = 0.8$
The slope of the drop is -1.6:	$f'(100) = -1.6$	$\Rightarrow$	$200a + b = -1.6$

(b) $b = 0.8$, so $200a + b = -1.6 \quad \Rightarrow \quad 200a + 0.8 = -1.6 \quad \Rightarrow \quad 200a = -2.4 \quad \Rightarrow \quad a = -\dfrac{2.4}{200} = -0.012$.

Thus, $f(x) = -0.012x^2 + 0.8x$.

(c) Since L_1 passes through the origin with slope 0.8, it has equation $y = 0.8x$. The horizontal distance between P and Q is 100, so the y-coordinate at Q is $f(100) = -0.012(100)^2 + 0.8(100) = -40$. Since L_2 passes through the point $(100, -40)$ and has slope -1.6, it has equation $y + 40 = -1.6(x - 100)$ or $y = -1.6x + 120$.

(d) The difference in elevation between $P(0, 0)$ and $Q(100, -40)$ is $0 - (-40) = 40$ feet.

2. (a)

Interval	Function	First Derivative	Second Derivative
$(-\infty, 0)$	$L_1(x) = 0.8x$	$L_1'(x) = 0.8$	$L_1''(x) = 0$
$[0, 10)$	$g(x) = kx^3 + lx^2 + mx + n$	$g'(x) = 3kx^2 + 2lx + m$	$g''(x) = 6kx + 2l$
$[10, 90]$	$q(x) = ax^2 + bx + c$	$q'(x) = 2ax + b$	$q''(x) = 2a$
$(90, 100]$	$h(x) = px^3 + qx^2 + rx + s$	$h'(x) = 3px^2 + 2qx + r$	$h''(x) = 6px + 2q$
$(100, \infty)$	$L_2(x) = -1.6x + 120$	$L_2'(x) = -1.6$	$L_2''(x) = 0$

There are 4 values of x (0, 10, 90, and 100) for which we must make sure the function values are equal, the first derivative values are equal, and the second derivative values are equal. The third column in the following table contains the value of each side of the condition — these are found after solving the system in part (b).

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

At $x =$	Condition	Value	Resulting Equation
0	$g(0) = L_1(0)$	0	$n = 0$
	$g'(0) = L_1'(0)$	$\frac{4}{5}$	$m = 0.8$
	$g''(0) = L_1''(0)$	0	$2l = 0$
10	$g(10) = q(10)$	$\frac{68}{9}$	$1000k + 100l + 10m + n = 100a + 10b + c$
	$g'(10) = q'(10)$	$\frac{2}{3}$	$300k + 20l + m = 20a + b$
	$g''(10) = q''(10)$	$-\frac{2}{75}$	$60k + 2l = 2a$
90	$h(90) = q(90)$	$-\frac{220}{9}$	$729{,}000p + 8100q + 90r + s = 8100a + 90b + c$
	$h'(90) = q'(90)$	$-\frac{22}{15}$	$24{,}300p + 180q + r = 180a + b$
	$h''(90) = q''(90)$	$-\frac{2}{75}$	$540p + 2q = 2a$
100	$h(100) = L_2(100)$	-40	$1{,}000{,}000p + 10{,}000q + 100r + s = -40$
	$h'(100) = L_2'(100)$	$-\frac{8}{5}$	$30{,}000p + 200q + r = -1.6$
	$h''(100) = L_2''(100)$	0	$600p + 2q = 0$

(b) We can arrange our work in a 12×12 matrix as follows.

a	b	c	k	l	m	n	p	q	r	s	constant
0	0	0	0	0	0	1	0	0	0	0	0
0	0	0	0	0	1	0	0	0	0	0	0.8
0	0	0	0	2	0	0	0	0	0	0	0
−100	−10	−1	1000	100	10	1	0	0	0	0	0
−20	−1	0	300	20	1	0	0	0	0	0	0
−2	0	0	60	2	0	0	0	0	0	0	0
−8100	−90	−1	0	0	0	0	729,000	8100	90	1	0
−180	−1	0	0	0	0	0	24,300	180	1	0	0
−2	0	0	0	0	0	0	540	2	0	0	0
0	0	0	0	0	0	0	1,000,000	10,000	100	1	−40
0	0	0	0	0	0	0	30,000	200	1	0	−1.6
0	0	0	0	0	0	0	600	2	0	0	0

Solving the system gives us the formulas for q, g, and h.

$$\left.\begin{aligned} a &= -0.01\overline{3} = -\tfrac{1}{75} \\ b &= 0.9\overline{3} = \tfrac{14}{15} \\ c &= -0.\overline{4} = -\tfrac{4}{9} \end{aligned}\right\} q(x) = -\tfrac{1}{75}x^2 + \tfrac{14}{15}x - \tfrac{4}{9} \qquad \left.\begin{aligned} k &= -0.000\overline{4} = -\tfrac{1}{2250} \\ l &= 0 \\ m &= 0.8 = \tfrac{4}{5} \\ n &= 0 \end{aligned}\right\} g(x) = -\tfrac{1}{2250}x^3 + \tfrac{4}{5}x$$

$$\left.\begin{aligned} p &= 0.000\overline{4} = \tfrac{1}{2250} \\ q &= -0.1\overline{3} = -\tfrac{2}{15} \\ r &= 11.7\overline{3} = \tfrac{176}{15} \\ s &= -324.\overline{4} = -\tfrac{2920}{9} \end{aligned}\right\} h(x) = \tfrac{1}{2250}x^3 - \tfrac{2}{15}x^2 + \tfrac{176}{15}x - \tfrac{2920}{9}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) Graph of L_1, q, g, h, and L_2:

This is the piecewise-defined function assignment on a TI-83/4 Plus calculator, where $Y_2 = L_1$, $Y_6 = g$, $Y_5 = q$, $Y_7 = h$, and $Y_3 = L_2$.

The graph of the five functions as a piecewise-defined function:

A comparison of the graphs in part 1(c) and part 2(c):

3.2 The Product and Quotient Rules

1. Product Rule: $f(x) = (1 + 2x^2)(x - x^2) \quad \Rightarrow$

$$f'(x) = (1 + 2x^2)(1 - 2x) + (x - x^2)(4x) = 1 - 2x + 2x^2 - 4x^3 + 4x^2 - 4x^3 = 1 - 2x + 6x^2 - 8x^3.$$

Multiplying first: $f(x) = (1 + 2x^2)(x - x^2) = x - x^2 + 2x^3 - 2x^4 \quad \Rightarrow \quad f'(x) = 1 - 2x + 6x^2 - 8x^3$ (equivalent).

2. Quotient Rule: $F(x) = \dfrac{x^4 - 5x^3 + \sqrt{x}}{x^2} = \dfrac{x^4 - 5x^3 + x^{1/2}}{x^2} \quad \Rightarrow$

$$F'(x) = \frac{x^2(4x^3 - 15x^2 + \frac{1}{2}x^{-1/2}) - (x^4 - 5x^3 + x^{1/2})(2x)}{(x^2)^2} = \frac{4x^5 - 15x^4 + \frac{1}{2}x^{3/2} - 2x^5 + 10x^4 - 2x^{3/2}}{x^4}$$

$$= \frac{2x^5 - 5x^4 - \frac{3}{2}x^{3/2}}{x^4} = 2x - 5 - \tfrac{3}{2}x^{-5/2}$$

Simplifying first: $F(x) = \dfrac{x^4 - 5x^3 + \sqrt{x}}{x^2} = x^2 - 5x + x^{-3/2} \quad \Rightarrow \quad F'(x) = 2x - 5 - \frac{3}{2}x^{-5/2}$ (equivalent).

For this problem, simplifying first seems to be the better method.

3. By the Product Rule, $y = (4x^2 + 3)(2x + 5) \quad \Rightarrow$

$$y' = (4x^2 + 3)(2x + 5)' + (2x + 5)(4x^2 + 3)' = (4x^2 + 3)(2) + (2x + 5)(8x)$$

$$= 8x^2 + 6 + 16x^2 + 40x = 24x^2 + 40x + 6$$

4. By the Product Rule, $y = (10x^2 + 7x - 2)(2 - x^2) \quad \Rightarrow$

$$y' = (10x^2 + 7x - 2)(2 - x^2)' + (2 - x^2)(10x^2 + 7x - 2)' = (10x^2 + 7x - 2)(-2x) + (2 - x^2)(20x + 7)$$

$$= -20x^3 - 14x^2 + 4x + 40x + 14 - 20x^3 - 7x^2 = -40x^3 - 21x^2 + 44x + 14$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The notations $\overset{\text{PR}}{\Rightarrow}$ and $\overset{\text{QR}}{\Rightarrow}$ indicate the use of the Product and Quotient Rules, respectively.

5. $y = x^3 e^x \overset{\text{PR}}{\Rightarrow} y' = x^3(e^x)' + e^x(x^3)' = x^3 e^x + e^x \cdot 3x^2 = e^x(x^3 + 3x^2)$

6. $y = (e^x + 2)(2e^x - 1) \overset{\text{PR}}{\Rightarrow}$

$$y' = (e^x + 2)(2e^x - 1)' + (2e^x - 1)(e^x + 2)' = (e^x + 2)(2e^x) + (2e^x - 1)(e^x)$$
$$= 2e^{2x} + 4e^x + 2e^{2x} - e^x = 4e^{2x} + 3e^x \text{ or } e^x(4e^x + 3)$$

7. $f(x) = (3x^2 - 5x)e^x \overset{\text{PR}}{\Rightarrow}$

$$f'(x) = (3x^2 - 5x)(e^x)' + e^x(3x^2 - 5x)' = (3x^2 - 5x)e^x + e^x(6x - 5)$$
$$= e^x[(3x^2 - 5x) + (6x - 5)] = e^x(3x^2 + x - 5)$$

8. $g(x) = (x + 2\sqrt{x}\,)\,e^x \overset{\text{PR}}{\Rightarrow}$

$$g'(x) = (x + 2\sqrt{x}\,)(e^x)' + e^x(x + 2\sqrt{x}\,)' = (x + 2\sqrt{x}\,)e^x + e^x\left(1 + 2 \cdot \tfrac{1}{2}x^{-1/2}\right)$$
$$= e^x\left[(x + 2\sqrt{x}\,) + \left(1 + 1/\sqrt{x}\right)\right] = e^x\left(x + 2\sqrt{x} + 1 + 1/\sqrt{x}\right)$$

9. By the Quotient Rule, $y = \dfrac{x}{e^x} \Rightarrow y' = \dfrac{e^x(1) - x(e^x)}{(e^x)^2} = \dfrac{e^x(1-x)}{(e^x)^2} = \dfrac{1-x}{e^x}$.

10. By the Quotient Rule, $y = \dfrac{e^x}{1 - e^x} \Rightarrow y' = \dfrac{(1 - e^x)e^x - e^x(-e^x)}{(1 - e^x)^2} = \dfrac{e^x - e^{2x} + e^{2x}}{(1 - e^x)^2} = \dfrac{e^x}{(1 - e^x)^2}$.

11. $g(t) = \dfrac{3 - 2t}{5t + 1} \overset{\text{QR}}{\Rightarrow} g'(t) = \dfrac{(5t + 1)(-2) - (3 - 2t)(5)}{(5t + 1)^2} = \dfrac{-10t - 2 - 15 + 10t}{(5t + 1)^2} = -\dfrac{17}{(5t + 1)^2}$

12. $G(u) = \dfrac{6u^4 - 5u}{u + 1} \overset{\text{QR}}{\Rightarrow}$

$$G'(u) = \frac{(u + 1)(24u^3 - 5) - (6u^4 - 5u)(1)}{(u + 1)^2} = \frac{24u^4 - 5u + 24u^3 - 5 - 6u^4 + 5u}{(u + 1)^2} = \frac{18u^4 + 24u^3 - 5}{(u + 1)^2}$$

13. $f(t) = \dfrac{5t}{t^3 - t - 1} \overset{\text{QR}}{\Rightarrow} f'(t) = \dfrac{(t^3 - t - 1)(5) - (5t)(3t^2 - 1)}{(t^3 - t - 1)^2} = \dfrac{5t^3 - 5t - 5 - 15t^3 + 5t}{(t^3 - t - 1)^2} = -\dfrac{10t^3 + 5}{(t^3 - t - 1)^2}$

14. $F(x) = \dfrac{1}{2x^3 - 6x^2 + 5} \overset{\text{QR}}{\Rightarrow} F'(x) = \dfrac{(2x^3 - 6x^2 + 5)(0) - 1(6x^2 - 12x)}{(2x^3 - 6x^2 + 5)^2} = -\dfrac{6x^2 - 12x}{(2x^3 - 6x^2 + 5)^2}$

15. $y = \dfrac{s - \sqrt{s}}{s^2} = \dfrac{s}{s^2} - \dfrac{\sqrt{s}}{s^2} = s^{-1} - s^{-3/2} \Rightarrow y' = -s^{-2} + \tfrac{3}{2}s^{-5/2} = \dfrac{-1}{s^2} + \dfrac{3}{2s^{5/2}} = \dfrac{3 - 2\sqrt{s}}{2s^{5/2}}$

16. $y = \dfrac{\sqrt{x}}{\sqrt{x} + 1} \overset{\text{QR}}{\Rightarrow} y' = \dfrac{\left(\sqrt{x} + 1\right)\left(\dfrac{1}{2\sqrt{x}}\right) - \sqrt{x}\left(\dfrac{1}{2\sqrt{x}}\right)}{\left(\sqrt{x} + 1\right)^2} = \dfrac{\dfrac{1}{2} + \dfrac{1}{2\sqrt{x}} - \dfrac{1}{2}}{\left(\sqrt{x} + 1\right)^2} = \dfrac{1}{2\sqrt{x}\left(\sqrt{x} + 1\right)^2}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $J(u) = \left(\dfrac{1}{u} + \dfrac{1}{u^2}\right)\left(u + \dfrac{1}{u}\right) = (u^{-1} + u^{-2})(u + u^{-1}) \overset{\text{PR}}{\Rightarrow}$

$$J'(u) = (u^{-1} + u^{-2})(u + u^{-1})' + (u + u^{-1})(u^{-1} + u^{-2})' = (u^{-1} + u^{-2})(1 - u^{-2}) + (u + u^{-1})(-u^{-2} - 2u^{-3})$$

$$= u^{-1} - u^{-3} + u^{-2} - u^{-4} - u^{-1} - 2u^{-2} - u^{-3} - 2u^{-4} = -u^{-2} - 2u^{-3} - 3u^{-4} = -\left(\frac{1}{u^2} + \frac{2}{u^3} + \frac{3}{u^4}\right)$$

18. $h(w) = (w^2 + 3w)(w^{-1} - w^{-4}) \overset{\text{PR}}{\Rightarrow}$

$$h'(w) = (w^2 + 3w)(-w^{-2} + 4w^{-5}) + (w^{-1} - w^{-4})(2w + 3)$$

$$= -1 + 4w^{-3} - 3w^{-1} + 12w^{-4} + 2 + 3w^{-1} - 2w^{-3} - 3w^{-4} = 1 + 2w^{-3} + 9w^{-4}$$

Alternate solution: An easier method is to simplify first and then differentiate as follows:

$$h(w) = (w^2 + 3w)(w^{-1} - w^{-4}) = w - w^{-2} + 3 - 3w^{-3} \quad \Rightarrow \quad h'(w) = 1 + 2w^{-3} + 9w^{-4}$$

19. $H(u) = (u - \sqrt{u})(u + \sqrt{u}) \overset{\text{PR}}{\Rightarrow}$

$$H'(u) = \left(u - \sqrt{u}\right)\left(1 + \frac{1}{2\sqrt{u}}\right) + \left(u + \sqrt{u}\right)\left(1 - \frac{1}{2\sqrt{u}}\right) = u + \tfrac{1}{2}\sqrt{u} - \sqrt{u} - \tfrac{1}{2} + u - \tfrac{1}{2}\sqrt{u} + \sqrt{u} - \tfrac{1}{2} = 2u - 1.$$

Alternate solution: An easier method is to simplify first and then differentiate as follows:

$$H(u) = (u - \sqrt{u})(u + \sqrt{u}) = u^2 - (\sqrt{u})^2 = u^2 - u \quad \Rightarrow \quad H'(u) = 2u - 1$$

20. $f(z) = (1 - e^z)(z + e^z) \overset{\text{PR}}{\Rightarrow}$

$$f'(z) = (1 - e^z)(1 + e^z) + (z + e^z)(-e^z) = 1^2 - (e^z)^2 - ze^z - (e^z)^2 = 1 - ze^z - 2e^{2z}$$

21. $V(t) = (t + 2e^t)\sqrt{t} \overset{\text{PR}}{\Rightarrow}$

$$V'(t) = (t + 2e^t)\frac{1}{2\sqrt{t}} + \sqrt{t}\,(1 + 2e^t) = \frac{t}{2\sqrt{t}} + \frac{2e^t}{2\sqrt{t}} + \sqrt{t} + 2\sqrt{t}\,e^t = \frac{t + 2e^t + 2t + 4te^t}{2\sqrt{t}} = \frac{3t + 2e^t + 4te^t}{2\sqrt{t}}$$

22. $W(t) = e^t(1 + te^t) \overset{\text{PR}}{\Rightarrow}$

$$W'(t) = e^t[0 + (te^t + e^t(1))] + (1 + te^t)(e^t) \qquad [\text{factor out } e^t]$$

$$= e^t(te^t + e^t + 1 + te^t) = e^t(1 + e^t + 2te^t)$$

23. $y = e^p(p + p\sqrt{p}) = e^p(p + p^{3/2}) \overset{\text{PR}}{\Rightarrow} y' = e^p\left(1 + \tfrac{3}{2}p^{1/2}\right) + (p + p^{3/2})e^p = e^p\left(1 + \tfrac{3}{2}\sqrt{p} + p + p\sqrt{p}\right)$

24. $h(r) = \dfrac{ae^r}{b + e^r} \overset{\text{QR}}{\Rightarrow} h'(r) = \dfrac{(b + e^r)(ae^r) - (ae^r)(e^r)}{(b + e^r)^2} = \dfrac{abe^r + ae^{2r} - ae^{2r}}{(b + e^r)^2} = \dfrac{abe^r}{(b + e^r)^2}$

25. $f(t) = \dfrac{\sqrt[3]{t}}{t - 3} \overset{\text{QR}}{\Rightarrow}$

$$f'(t) = \frac{(t - 3)\left(\frac{1}{3}t^{-2/3}\right) - t^{1/3}(1)}{(t - 3)^2} = \frac{\frac{1}{3}t^{1/3} - t^{-2/3} - t^{1/3}}{(t - 3)^2} = \frac{-\frac{2}{3}t^{1/3} - t^{-2/3}}{(t - 3)^2}$$

$$= \frac{\dfrac{-2t}{3t^{2/3}} - \dfrac{3}{3t^{2/3}}}{(t - 3)^2} = \frac{-2t - 3}{3t^{2/3}(t - 3)^2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $y = (z^2 + e^z)\sqrt{z} \overset{\text{PR}}{\Rightarrow}$

$$y' = (z^2 + e^z)\left(\frac{1}{2\sqrt{z}}\right) + \sqrt{z}\,(2z + e^z) = \frac{z^2}{2\sqrt{z}} + \frac{e^z}{2\sqrt{z}} + 2z\sqrt{z} + \sqrt{z}\,e^z$$

$$= \frac{z^2 + e^z + 4z^2 + 2ze^z}{2\sqrt{z}} = \frac{5z^2 + e^z + 2ze^z}{2\sqrt{z}}$$

27. $f(x) = \dfrac{x^2e^x}{x^2 + e^x} \overset{\text{QR}}{\Rightarrow}$

$$f'(x) = \frac{(x^2 + e^x)\left[x^2e^x + e^x(2x)\right] - x^2e^x(2x + e^x)}{(x^2 + e^x)^2} = \frac{x^4e^x + 2x^3e^x + x^2e^{2x} + 2xe^{2x} - 2x^3e^x - x^2e^{2x}}{(x^2 + e^x)^2}$$

$$= \frac{x^4e^x + 2xe^{2x}}{(x^2 + e^x)^2} = \frac{xe^x(x^3 + 2e^x)}{(x^2 + e^x)^2}$$

28. $F(t) = \dfrac{At}{Bt^2 + Ct^3} = \dfrac{A}{Bt + Ct^2} \overset{\text{QR}}{\Rightarrow}$

$$F'(t) = \frac{(Bt + Ct^2)(0) - A(B + 2Ct)}{(Bt + Ct^2)^2} = \frac{-A(B + 2Ct)}{(t)^2(B + Ct)^2} = -\frac{A(B + 2Ct)}{t^2(B + Ct)^2}$$

29. $f(x) = \dfrac{x}{x + \dfrac{c}{x}} \overset{\text{QR}}{\Rightarrow} f'(x) = \dfrac{(x + c/x)(1) - x(1 - c/x^2)}{\left(x + \dfrac{c}{x}\right)^2} = \dfrac{x + c/x - x + c/x}{\left(\dfrac{x^2 + c}{x}\right)^2} = \dfrac{2c/x}{\dfrac{(x^2 + c)^2}{x^2}} \cdot \dfrac{x^2}{x^2} = \dfrac{2cx}{(x^2 + c)^2}$

30. $f(x) = \dfrac{ax + b}{cx + d} \overset{\text{QR}}{\Rightarrow} f'(x) = \dfrac{(cx + d)(a) - (ax + b)(c)}{(cx + d)^2} = \dfrac{acx + ad - acx - bc}{(cx + d)^2} = \dfrac{ad - bc}{(cx + d)^2}$

31. $f(x) = x^2e^x \overset{\text{PR}}{\Rightarrow} f'(x) = x^2e^x + e^x(2x) = e^x(x^2 + 2x)$

Using the Product Rule and $f'(x) = e^x(x^2 + 2x)$, we get

$$f''(x) = e^x(2x + 2) + (x^2 + 2x)e^x = e^x(2x + 2 + x^2 + 2x) = e^x(x^2 + 4x + 2)$$

32. $f(x) = \sqrt{x}\,e^x \overset{\text{PR}}{\Rightarrow} f'(x) = \sqrt{x}\,e^x + e^x\left(\dfrac{1}{2\sqrt{x}}\right) = \left(\sqrt{x} + \dfrac{1}{2\sqrt{x}}\right)e^x = \dfrac{2x + 1}{2\sqrt{x}}\,e^x.$

Using the Product Rule and $f'(x) = \left(x^{1/2} + \frac{1}{2}x^{-1/2}\right)e^x$, we get

$$f''(x) = \left(x^{1/2} + \tfrac{1}{2}x^{-1/2}\right)e^x + e^x\left(\tfrac{1}{2}x^{-1/2} - \tfrac{1}{4}x^{-3/2}\right) = \left(x^{1/2} + x^{-1/2} - \tfrac{1}{4}x^{-3/2}\right)e^x = \frac{4x^2 + 4x - 1}{4x^{3/2}}\,e^x$$

33. $f(x) = \dfrac{x}{x^2 - 1} \overset{\text{QR}}{\Rightarrow} f'(x) = \dfrac{(x^2 - 1)(1) - x(2x)}{(x^2 - 1)^2} = \dfrac{x^2 - 1 - 2x^2}{(x^2 - 1)^2} = \dfrac{-x^2 - 1}{(x^2 - 1)^2} \Rightarrow$

$$f''(x) = \frac{(x^2 - 1)^2(-2x) - (-x^2 - 1)(x^4 - 2x^2 + 1)'}{[(x^2 - 1)^2]^2} = \frac{(x^2 - 1)^2(-2x) + (x^2 + 1)(4x^3 - 4x)}{(x^2 - 1)^4}$$

$$= \frac{(x^2 - 1)^2(-2x) + (x^2 + 1)(4x)(x^2 - 1)}{(x^2 - 1)^4} = \frac{(x^2 - 1)[(x^2 - 1)(-2x) + (x^2 + 1)(4x)]}{(x^2 - 1)^4}$$

$$= \frac{-2x^3 + 2x + 4x^3 + 4x}{(x^2 - 1)^3} = \frac{2x^3 + 6x}{(x^2 - 1)^3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. $f(x) = \dfrac{x}{1+\sqrt{x}} \overset{\text{QR}}{\Rightarrow}$

$$f'(x) = \frac{\left(1+\sqrt{x}\right)(1) - x\left(\dfrac{1}{2\sqrt{x}}\right)}{\left(1+\sqrt{x}\right)^2} = \frac{1+\sqrt{x}-\frac{1}{2}\sqrt{x}}{1+2\sqrt{x}+x} = \frac{1+\frac{1}{2}\sqrt{x}}{1+2\sqrt{x}+x} = \frac{\dfrac{2+\sqrt{x}}{2}}{1+2\sqrt{x}+x}$$

$$= \frac{2+\sqrt{x}}{2\left(1+2\sqrt{x}+x\right)} = \frac{2+\sqrt{x}}{2+4\sqrt{x}+2x}$$

Using the Quotient Rule and $f'(x) = \dfrac{2+\sqrt{x}}{2+4\sqrt{x}+2x}$, we get

$$f''(x) = \frac{\left(2+4\sqrt{x}+2x\right)\left(\dfrac{1}{2\sqrt{x}}\right) - \left(2+\sqrt{x}\right)\left(\dfrac{4}{2\sqrt{x}}+2\right)}{\left(2+4\sqrt{x}+2x\right)^2}$$

$$= \frac{\dfrac{1}{\sqrt{x}} + 2 + \sqrt{x} - \dfrac{4}{\sqrt{x}} - 4 - 2 - 2\sqrt{x}}{\left(2+4\sqrt{x}+2x\right)^2}$$

$$= \frac{\dfrac{-3-4\sqrt{x}-x}{\sqrt{x}}}{\left(2+4\sqrt{x}+2x\right)^2} = -\frac{3+4\sqrt{x}+x}{\sqrt{x}\left(2+4\sqrt{x}2x\right)^2}$$

35. $y = \dfrac{x^2}{1+x} \quad \Rightarrow \quad y' = \dfrac{(1+x)(2x) - x^2(1)}{(1+x)^2} = \dfrac{2x+2x^2-x^2}{(1+x)^2} = \dfrac{x^2+2x}{(1+x)^2}$

At $\left(1, \frac{1}{2}\right)$, $y' = \dfrac{1^2+2\,(1)}{(1+1)^2} = \dfrac{3}{4}$, and an equation of the tangent line is $y - \frac{1}{2} = \frac{3}{4}(x-1)$, or $y = \frac{3}{4}x - \frac{1}{4}$.

36. $y = \dfrac{1+x}{1+e^x} \quad \Rightarrow \quad y' = \dfrac{(1+e^x)(1) - (1+x)e^x}{(1+e^x)^2} = \dfrac{1+e^x-e^x-xe^x}{(1+e^x)^2} = \dfrac{1-xe^x}{(1+e^x)^2}$

At $\left(0, \frac{1}{2}\right)$, $y' = \dfrac{1}{(1+1)^2} = \dfrac{1}{4}$, and an equation of the tangent line is $y - \frac{1}{2} = \frac{1}{4}(x-0)$ or $y = \frac{1}{4}x + \frac{1}{2}$.

37. $y = \dfrac{3x}{1+5x^2} \quad \Rightarrow \quad y' = \dfrac{(1+5x^2)(3) - 3x(10x)}{(1+5x^2)^2} = \dfrac{3+15x^2-30x^2}{(1+5x^2)^2} = \dfrac{3-15x^2}{(1+5x^2)^2}$

At $\left(1, \frac{1}{2}\right)$, $y' = \dfrac{3-15(1^2)}{(1+5\cdot 1^2)^2} = \dfrac{-12}{6^2} = -\dfrac{1}{3}$, and an equation of the tangent line is $y - \frac{1}{2} = -\frac{1}{3}(x-1)$, or $y = -\frac{1}{3}x + \frac{5}{6}$.

The slope of the normal line is 3, so an equation of the normal line is $y - \frac{1}{2} = 3(x-1)$, or $y = 3x - \frac{5}{2}$.

38. $y = x + xe^x \quad \Rightarrow \quad y' = 1 + (xe^x + e^x \cdot 1) = 1 + e^x(x+1)$

At $(0,0)$, $y' = 1 + e^0(0+1) = 1 + 1\cdot 1 = 2$, and an equation of the tangent line is $y - 0 = 2(x-0)$, or $y = 2x$.

The slope of the normal line is $-\frac{1}{2}$, so an equation of the normal line is $y - 0 = -\frac{1}{2}(x-0)$, or $y = -\frac{1}{2}x$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. (a) $y = f(x) = \dfrac{1}{1+x^2} \quad \Rightarrow$

$f'(x) = \dfrac{(1+x^2)(0) - 1(2x)}{(1+x^2)^2} = \dfrac{-2x}{(1+x^2)^2}$. So the slope of the tangent line at the point $\left(-1, \frac{1}{2}\right)$ is $f'(-1) = \dfrac{2}{2^2} = \frac{1}{2}$ and its equation is $y - \frac{1}{2} = \frac{1}{2}(x+1)$ or $y = \frac{1}{2}x + 1$.

(b)

40. (a) $y = f(x) = \dfrac{x}{1+x^2} \quad \Rightarrow$

$f'(x) = \dfrac{(1+x^2)1 - x(2x)}{(1+x^2)^2} = \dfrac{1-x^2}{(1+x^2)^2}$. So the slope of the tangent line at the point $(3, 0.3)$ is $f'(3) = \frac{-8}{100}$ and its equation is $y - 0.3 = -0.08(x-3)$ or $y = -0.08x + 0.54$.

(b)

41. (a) $f(x) = (x^3 - x)e^x \quad \Rightarrow \quad f'(x) = (x^3 - x)e^x + e^x(3x^2 - 1) = e^x(x^3 + 3x^2 - x - 1)$

(b)

$f' = 0$ when f has a horizontal tangent line, f' is negative when f is decreasing, and f' is positive when f is increasing.

42. (a) $f(x) = \dfrac{x^2-1}{x^2+1} \quad \Rightarrow$

$$f'(x) = \frac{(x^2+1)(2x) - (x^2-1)(2x)}{(x^2+1)^2} = \frac{(2x)[(x^2+1)-(x^2-1)]}{(x^2+1)^2} = \frac{(2x)(2)}{(x^2+1)^2} = \frac{4x}{(x^2+1)^2} \quad \Rightarrow$$

$$f''(x) = \frac{(x^2+1)^2(4) - 4x(x^4+2x^2+1)'}{[(x^2+1)^2]^2}$$

$$= \frac{4(x^2+1)^2 - 4x(4x^3+4x)}{(x^2+1)^4} = \frac{4(x^2+1)^2 - 16x^2(x^2+1)}{(x^2+1)^4}$$

$$= \frac{4(x^2+1)[(x^2+1)-4x^2]}{(x^2+1)^4} = \frac{4(1-3x^2)}{(x^2+1)^3}$$

(b)

$f' = 0$ when f has a horizontal tangent and $f'' = 0$ when f' has a horizontal tangent. f' is negative when f is decreasing and positive when f is increasing. f'' is negative when f' is decreasing and positive when f' is increasing. f'' is negative when f is concave down and positive when f is concave up.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. $f(x) = \dfrac{x^2}{1+x} \quad \Rightarrow \quad f'(x) = \dfrac{(1+x)(2x) - x^2(1)}{(1+x)^2} = \dfrac{2x + 2x^2 - x^2}{(1+x)^2} = \dfrac{x^2+2x}{x^2+2x+1} \quad \Rightarrow$

$$f''(x) = \frac{(x^2+2x+1)(2x+2) - (x^2+2x)(2x+2)}{(x^2+2x+1)^2} = \frac{(2x+2)(x^2+2x+1-x^2-2x)}{[(x+1)^2]^2}$$

$$= \frac{2(x+1)(1)}{(x+1)^4} = \frac{2}{(x+1)^3},$$

so $f''(1) = \dfrac{2}{(1+1)^3} = \dfrac{2}{8} = \dfrac{1}{4}$.

44. $g(x) = \dfrac{x}{e^x} \quad \Rightarrow \quad g'(x) = \dfrac{e^x \cdot 1 - x \cdot e^x}{(e^x)^2} = \dfrac{e^x(1-x)}{(e^x)^2} = \dfrac{1-x}{e^x} \quad \Rightarrow$

$$g''(x) = \frac{e^x \cdot (-1) - (1-x)e^x}{(e^x)^2} = \frac{e^x[-1-(1-x)]}{(e^x)^2} = \frac{x-2}{e^x} \quad \Rightarrow$$

$$g'''(x) = \frac{e^x \cdot 1 - (x-2)e^x}{(e^x)^2} = \frac{e^x[1-(x-2)]}{(e^x)^2} = \frac{3-x}{e^x} \quad \Rightarrow$$

$$g^{(4)}(x) = \frac{e^x \cdot (-1) - (3-x)e^x}{(e^x)^2} = \frac{e^x[-1-(3-x)]}{(e^x)^2} = \frac{x-4}{e^x}.$$

The pattern suggests that $g^{(n)}(x) = \dfrac{(x-n)(-1)^n}{e^x}$. (We could use mathematical induction to prove this formula.)

45. We are given that $f(5) = 1$, $f'(5) = 6$, $g(5) = -3$, and $g'(5) = 2$.

(a) $(fg)'(5) = f(5)g'(5) + g(5)f'(5) = (1)(2) + (-3)(6) = 2 - 18 = -16$

(b) $\left(\dfrac{f}{g}\right)'(5) = \dfrac{g(5)f'(5) - f(5)g'(5)}{[g(5)]^2} = \dfrac{(-3)(6)-(1)(2)}{(-3)^2} = -\dfrac{20}{9}$

(c) $\left(\dfrac{g}{f}\right)'(5) = \dfrac{f(5)g'(5) - g(5)f'(5)}{[f(5)]^2} = \dfrac{(1)(2)-(-3)(6)}{(1)^2} = 20$

46. We are given that $f(4) = 2$, $g(4) = 5$, $f'(4) = 6$, and $g'(4) = -3$.

(a) $h(x) = 3f(x) + 8g(x) \quad \Rightarrow \quad h'(x) = 3f'(x) + 8g'(x)$, so

$h'(4) = 3f'(4) + 8g'(4) = 3(6) + 8(-3) = 18 - 24 = -6$.

(b) $h(x) = f(x)\,g(x) \quad \Rightarrow \quad h'(x) = f(x)\,g'(x) + g(x)\,f'(x)$, so

$h'(4) = f(4)\,g'(4) + g(4)\,f'(4) = 2(-3) + 5(6) = -6 + 30 = 24$.

(c) $h(x) = \dfrac{f(x)}{g(x)} \quad \Rightarrow \quad h'(x) = \dfrac{g(x)\,f'(x) - f(x)\,g'(x)}{[g(x)]^2}$, so

$h'(4) = \dfrac{g(4)\,f'(4) - f(4)\,g'(4)}{[g(4)]^2} = \dfrac{5(6) - 2(-3)}{5^2} = \dfrac{30+6}{25} = \dfrac{36}{25}$.

(d) $h(x) = \dfrac{g(x)}{f(x)+g(x)} \quad \Rightarrow$

$$h'(4) = \frac{[f(4)+g(4)]\,g'(4) - g(4)\,[f'(4)+g'(4)]}{[f(4)+g(4)]^2} = \frac{(2+5)\,(-3) - 5\,[6+(-3)]}{(2+5)^2} = \frac{-21-15}{7^2} = -\frac{36}{49}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $f(x) = e^x g(x) \quad \Rightarrow \quad f'(x) = e^x g'(x) + g(x)e^x = e^x[g'(x) + g(x)]$. $f'(0) = e^0[g'(0) + g(0)] = 1(5 + 2) = 7$

48. $\dfrac{d}{dx}\left[\dfrac{h(x)}{x}\right] = \dfrac{xh'(x) - h(x) \cdot 1}{x^2} \quad \Rightarrow \quad \dfrac{d}{dx}\left[\dfrac{h(x)}{x}\right]_{x=2} = \dfrac{2h'(2) - h(2)}{2^2} = \dfrac{2(-3) - (4)}{4} = \dfrac{-10}{4} = -2.5$

49. $g(x) = xf(x) \quad \Rightarrow \quad g'(x) = xf'(x) + f(x) \cdot 1$. Now $g(3) = 3f(3) = 3 \cdot 4 = 12$ and $g'(3) = 3f'(3) + f(3) = 3(-2) + 4 = -2$. Thus, an equation of the tangent line to the graph of g at the point where $x = 3$ is $y - 12 = -2(x - 3)$, or $y = -2x + 18$.

50. $f'(x) = x^2 f(x) \quad \Rightarrow \quad f''(x) = x^2 f'(x) + f(x) \cdot 2x$. Now $f'(2) = 2^2 f(2) = 4(10) = 40$, so $f''(2) = 2^2(40) + 10(4) = 200$.

51. (a) From the graphs of f and g, we obtain the following values: $f(1) = 2$ since the point $(1, 2)$ is on the graph of f; $g(1) = 3$ since the point $(1, 3)$ is on the graph of g; $f'(1) = \frac{1}{3}$ since the slope of the line segment between $(1, 2)$ and $(4, 3)$ is $\dfrac{3-2}{4-1} = \dfrac{1}{3}$; $g'(1) = 1$ since the slope of the line segment between $(-2, 0)$ and $(2, 4)$ is $\dfrac{4-0}{2-(-2)} = \dfrac{4}{4} = 1$.

Now $u(x) = f(x)\,g(x)$, so $u'(1) = f(1)\,g'(1) + g(1)\,f'(1) = 2 \cdot 1 + 3 \cdot \frac{1}{3} = 3$.

(b) From the graphs of f and g, we obtain the following values: $f(4) = 3$ since the point $(4, 3)$ is on the graph of f; $g(4) = 2$ since the point $(4, 2)$ is on the graph of g; $f'(4) = f'(1) = \frac{1}{3}$ from the part (a); $g'(4) = 1$ since the slope of the line segment between $(3, 1)$ and $(5, 3)$ is $\dfrac{3-1}{5-3} = \dfrac{2}{2} = 1$.

$v(x) = \dfrac{f(x)}{g(x)}$, so $v'(4) = \dfrac{g(4)f'(4) - f(4)g'(4)}{[g(4)]^2} = \dfrac{2 \cdot \frac{1}{3} - 3 \cdot 1}{2^2} = \dfrac{-\frac{7}{3}}{4} = -\dfrac{7}{12}$

52. (a) $P(x) = F(x)\,G(x)$, so $P'(2) = F(2)\,G'(2) + G(2)\,F'(2) = 3 \cdot \frac{2}{4} + 2 \cdot 0 = \frac{3}{2}$.

(b) $Q(x) = \dfrac{F(x)}{G(x)}$, so $Q'(7) = \dfrac{G(7)\,F'(7) - F(7)\,G'(7)}{[G(7)]^2} = \dfrac{1 \cdot \frac{1}{4} - 5 \cdot \left(-\frac{2}{3}\right)}{1^2} = \dfrac{1}{4} + \dfrac{10}{3} = \dfrac{43}{12}$

53. (a) $y = xg(x) \quad \Rightarrow \quad y' = xg'(x) + g(x) \cdot 1 = xg'(x) + g(x)$

(b) $y = \dfrac{x}{g(x)} \quad \Rightarrow \quad y' = \dfrac{g(x) \cdot 1 - xg'(x)}{[g(x)]^2} = \dfrac{g(x) - xg'(x)}{[g(x)]^2}$

(c) $y = \dfrac{g(x)}{x} \quad \Rightarrow \quad y' = \dfrac{xg'(x) - g(x) \cdot 1}{(x)^2} = \dfrac{xg'(x) - g(x)}{x^2}$

54. (a) $y = x^2 f(x) \quad \Rightarrow \quad y' = x^2 f'(x) + f(x)(2x)$

(b) $y = \dfrac{f(x)}{x^2} \quad \Rightarrow \quad y' = \dfrac{x^2 f'(x) - f(x)(2x)}{(x^2)^2} = \dfrac{xf'(x) - 2f(x)}{x^3}$

(c) $y = \dfrac{x^2}{f(x)} \quad \Rightarrow \quad y' = \dfrac{f(x)(2x) - x^2 f'(x)}{[f(x)]^2}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $y = \dfrac{1 + xf(x)}{\sqrt{x}} \quad \Rightarrow$

$$y' = \frac{\sqrt{x}\,[xf'(x) + f(x)] - [1 + xf(x)]\,\dfrac{1}{2\sqrt{x}}}{\left(\sqrt{x}\right)^2}$$

$$= \frac{x^{3/2}f'(x) + x^{1/2}f(x) - \frac{1}{2}x^{-1/2} - \frac{1}{2}x^{1/2}f(x)}{x} \cdot \frac{2x^{1/2}}{2x^{1/2}} = \frac{xf(x) + 2x^2f'(x) - 1}{2x^{3/2}}$$

55. If $y = f(x) = \dfrac{x}{x+1}$, then $f'(x) = \dfrac{(x+1)(1) - x(1)}{(x+1)^2} = \dfrac{1}{(x+1)^2}$. When $x = a$, the equation of the tangent line is $y - \dfrac{a}{a+1} = \dfrac{1}{(a+1)^2}(x - a)$. This line passes through $(1, 2)$ when $2 - \dfrac{a}{a+1} = \dfrac{1}{(a+1)^2}(1 - a) \quad \Leftrightarrow$ $2(a+1)^2 - a(a+1) = 1 - a \quad \Leftrightarrow \quad 2a^2 + 4a + 2 - a^2 - a - 1 + a = 0 \quad \Leftrightarrow \quad a^2 + 4a + 1 = 0$.

The quadratic formula gives the solutions of this equation as $a = \dfrac{-4 \pm \sqrt{4^2 - 4(1)(1)}}{2(1)} = \dfrac{-4 \pm \sqrt{12}}{2} = -2 \pm \sqrt{3}$, so there are two such tangent lines. Since

$$f\left(-2 \pm \sqrt{3}\right) = \frac{-2 \pm \sqrt{3}}{-2 \pm \sqrt{3} + 1} = \frac{-2 \pm \sqrt{3}}{-1 \pm \sqrt{3}} \cdot \frac{-1 \mp \sqrt{3}}{-1 \mp \sqrt{3}}$$

$$= \frac{2 \pm 2\sqrt{3} \mp \sqrt{3} - 3}{1 - 3} = \frac{-1 \pm \sqrt{3}}{-2} = \frac{1 \mp \sqrt{3}}{2},$$

the lines touch the curve at $A\left(-2 + \sqrt{3}, \frac{1 - \sqrt{3}}{2}\right) \approx (-0.27, -0.37)$ and $B\left(-2 - \sqrt{3}, \frac{1 + \sqrt{3}}{2}\right) \approx (-3.73, 1.37)$.

56. $y = \dfrac{x-1}{x+1} \quad \Rightarrow \quad y' = \dfrac{(x+1)(1) - (x-1)(1)}{(x+1)^2} = \dfrac{2}{(x+1)^2}$. If the tangent intersects the curve when $x = a$, then its slope is $2/(a+1)^2$. But if the tangent is parallel to $x - 2y = 2$, that is, $y = \frac{1}{2}x - 1$, then its slope is $\frac{1}{2}$. Thus, $\dfrac{2}{(a+1)^2} = \dfrac{1}{2} \quad \Rightarrow$ $(a+1)^2 = 4 \quad \Rightarrow \quad a + 1 = \pm 2 \quad \Rightarrow \quad a = 1$ or -3. When $a = 1$, $y = 0$ and the equation of the tangent is $y - 0 = \frac{1}{2}(x - 1)$ or $y = \frac{1}{2}x - \frac{1}{2}$. When $a = -3$, $y = 2$ and the equation of the tangent is $y - 2 = \frac{1}{2}(x + 3)$ or $y = \frac{1}{2}x + \frac{7}{2}$.

57. $R = \dfrac{f}{g} \quad \Rightarrow \quad R' = \dfrac{gf' - fg'}{g^2}$. For $f(x) = x - 3x^3 + 5x^5$, $f'(x) = 1 - 9x^2 + 25x^4$, and for $g(x) = 1 + 3x^3 + 6x^6 + 9x^9$, $g'(x) = 9x^2 + 36x^5 + 81x^8$.

Thus, $R'(0) = \dfrac{g(0)f'(0) - f(0)g'(0)}{[g(0)]^2} = \dfrac{1 \cdot 1 - 0 \cdot 0}{1^2} = \dfrac{1}{1} = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

58. $Q = \dfrac{f}{g} \quad \Rightarrow \quad Q' = \dfrac{gf' - fg'}{g^2}$. For $f(x) = 1 + x + x^2 + xe^x$, $f'(x) = 1 + 2x + xe^x + e^x$,

and for $g(x) = 1 - x + x^2 - xe^x$, $g'(x) = -1 + 2x - xe^x - e^x$.

Thus, $Q'(0) = \dfrac{g(0)f'(0) - f(0)g'(0)}{[g(0)]^2} = \dfrac{1 \cdot 2 - 1 \cdot (-2)}{1^2} = \dfrac{4}{1} = 4$.

59. If $P(t)$ denotes the population at time t and $A(t)$ denotes the average annual income, then $T(t) = P(t)\,A(t)$ is the total personal income. The rate at which $T(t)$ is rising is given by $T'(t) = P(t)\,A'(t) + A(t)\,P'(t) \quad \Rightarrow$

$$\begin{aligned} T'(2015) &= P(2015)\,A'(2015) + A(2015)\,P'(2015) = (107{,}350)\,(\$2250/\text{year}) + (\$60{,}220)\,(1960/\text{year}) \\ &= \$241{,}537{,}500/\text{year} + \$118{,}031{,}200/\text{year} = \$359{,}568{,}700/\text{year} \end{aligned}$$

So the total personal income in Boulder was rising by about \$360 million per year in 2015.

The term $P(t)\,A'(t) \approx \$242$ million represents the portion of the rate of change of total income due to the existing population's increasing income. The term $A(t)\,P'(t) \approx \$118$ million represents the portion of the rate of change of total income due to increasing population.

60. (a) $f(20) = 10{,}000$ means that when the price of the fabric is $\$20/\text{yard}$, 10,000 yards will be sold.

$f'(20) = -350$ means that as the price of the fabric increases past $\$20/\text{yard}$, the amount of fabric which will be sold is decreasing at a rate of 350 yards per (dollar per yard).

(b) $R(p) = pf(p) \quad \Rightarrow \quad R'(p) = pf'(p) + f(p) \cdot 1 \quad \Rightarrow \quad R'(20) = 20f'(20) + f(20) \cdot 1 = 20(-350) + 10{,}000 = 3000$.

This means that as the price of the fabric increases past $\$20/\text{yard}$, the total revenue is increasing at $\$3000/(\$/\text{yard})$. Note that the Product Rule indicates that we will lose $\$7000/(\$/\text{yard})$ due to selling less fabric, but this loss is more than made up for by the additional revenue due to the increase in price.

61. $v = \dfrac{0.14[\text{S}]}{0.015 + [\text{S}]} \quad \Rightarrow \quad \dfrac{dv}{d[\text{S}]} = \dfrac{(0.015 + [\text{S}])(0.14) - (0.14[\text{S}])(1)}{(0.015 + [\text{S}])^2} = \dfrac{0.0021}{(0.015 + [\text{S}])^2}$.

$dv/d[\text{S}]$ represents the rate of change of the rate of an enzymatic reaction with respect to the concentration of a substrate S.

62. $B(t) = N(t)\,M(t) \quad \Rightarrow \quad B'(t) = N(t)\,M'(t) + M(t)\,N'(t)$, so

$B'(4) = N(4)\,M'(4) + M(4)\,N'(4) = 820(0.14) + 1.2(50) = 174.8$ g/week.

63. (a) $(fgh)' = [(fg)h]' = (fg)'h + (fg)h' = (f'g + fg')h + (fg)h' = f'gh + fg'h + fgh'$

(b) Putting $f = g = h$ in part (a), we have $\dfrac{d}{dx}[f(x)]^3 = (fff)' = f'ff + ff'f + fff' = 3fff' = 3[f(x)]^2 f'(x)$.

(c) $\dfrac{d}{dx}\,(e^{3x}) = \dfrac{d}{dx}\,(e^x)^3 = 3(e^x)^2 e^x = 3e^{2x}e^x = 3e^{3x}$

64. (a) We use the Product Rule repeatedly: $F = fg \quad \Rightarrow \quad F' = f'g + fg' \quad \Rightarrow$

$F'' = (f''g + f'g') + (f'g' + fg'') = f''g + 2f'g' + fg''$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $F''' = f'''g + f''g' + 2\left(f''g' + f'g''\right) + f'g'' + fg''' = f'''g + 3f''g' + 3f'g'' + fg''' \quad \Rightarrow$

$$F^{(4)} = f^{(4)}g + f'''g' + 3\left(f'''g' + f''g''\right) + 3\left(f''g'' + f'g'''\right) + f'g''' + fg^{(4)}$$
$$= f^{(4)}g + 4f'''g' + 6f''g'' + 4f'g''' + fg^{(4)}$$

(c) By analogy with the Binomial Theorem, we make the guess:

$$F^{(n)} = f^{(n)}g + nf^{(n-1)}g' + \binom{n}{2}f^{(n-2)}g'' + \cdots + \binom{n}{k}f^{(n-k)}g^{(k)} + \cdots + nf'g^{(n-1)} + fg^{(n)},$$

where $\dfrac{\binom{n}{k}}{} = \dfrac{n!}{k!\,(n-k)!} = \dfrac{n(n-1)(n-2)\cdots(n-k+1)}{k!}$.

65. For $f(x) = x^2e^x$, $f'(x) = x^2e^x + e^x(2x) = e^x(x^2+2x)$. Similarly, we have

$$f''(x) = e^x(x^2+4x+2)$$
$$f'''(x) = e^x(x^2+6x+6)$$
$$f^{(4)}(x) = e^x(x^2+8x+12)$$
$$f^{(5)}(x) = e^x(x^2+10x+20)$$

It appears that the coefficient of x in the quadratic term increases by 2 with each differentiation. The pattern for the constant terms seems to be $0 = 1\cdot 0$, $2 = 2\cdot 1$, $6 = 3\cdot 2$, $12 = 4\cdot 3$, $20 = 5\cdot 4$. So a reasonable guess is that $f^{(n)}(x) = e^x[x^2+2nx+n(n-1)]$.

Proof: Let S_n be the statement that $f^{(n)}(x) = e^x[x^2+2nx+n(n-1)]$.

1. S_1 is true because $f'(x) = e^x(x^2+2x)$.
2. Assume that S_k is true; that is, $f^{(k)}(x) = e^x[x^2+2kx+k(k-1)]$. Then
$$f^{(k+1)}(x) = \frac{d}{dx}\left[f^{(k)}(x)\right] = e^x(2x+2k) + [x^2+2kx+k(k-1)]e^x$$
$$= e^x[x^2+(2k+2)x+(k^2+k)] = e^x[x^2+2(k+1)x+(k+1)k]$$
This shows that S_{k+1} is true.
3. Therefore, by mathematical induction, S_n is true for all n; that is, $f^{(n)}(x) = e^x[x^2+2nx+n(n-1)]$ for every positive integer n.

66. (a) $\dfrac{d}{dx}\left(\dfrac{1}{g(x)}\right) = \dfrac{g(x)\cdot\frac{d}{dx}(1) - 1\cdot\frac{d}{dx}[g(x)]}{[g(x)]^2}$ [Quotient Rule] $= \dfrac{g(x)\cdot 0 - 1\cdot g'(x)}{[g(x)]^2} = \dfrac{0-g'(x)}{[g(x)]^2} = -\dfrac{g'(x)}{[g(x)]^2}$

(b) $\dfrac{d}{dx}\left(\dfrac{1}{2x^3-6x^2+5}\right) = -\dfrac{(2x^3-6x^2+5)'}{(2x^3-6x^2+5)^2} = -\dfrac{6x^2-12x}{(2x^3-6x^2+5)^2}$

(c) $\dfrac{d}{dx}\left(x^{-n}\right) = \dfrac{d}{dx}\left(\dfrac{1}{x^n}\right) = -\dfrac{(x^n)'}{(x^n)^2}$ [by the Reciprocal Rule] $= -\dfrac{nx^{n-1}}{x^{2n}} = -nx^{n-1-2n} = -nx^{-n-1}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3.3 Derivatives of Trigonometric Functions

1. $f(x) = 3\sin x - 2\cos x \quad \Rightarrow \quad f'(x) = 3(\cos x) - 2(-\sin x) = 3\cos x + 2\sin x$

2. $f(x) = \tan x - 4\sin x \quad \Rightarrow \quad f'(x) = \sec^2 x - 4(\cos x) = \sec^2 x - 4\cos x$

3. $y = x^2 + \cot x \quad \Rightarrow \quad y' = 2x + (-\csc^2 x) = 2x - \csc^2 x$

4. $y = 2\sec x - \csc x \quad \Rightarrow \quad y' = 2(\sec x\ \tan x) - (-\csc x\ \cot x) = 2\sec x\ \tan x + \csc x\ \cot x$

5. $h(\theta) = \theta^2 \sin\theta \overset{\text{PR}}{\Rightarrow} h'(\theta) = \theta^2(\cos\theta) + (\sin\theta)(2\theta) = \theta(\theta\cos\theta + 2\sin\theta)$

6. $g(x) = 3x + x^2\cos x \quad \Rightarrow \quad g'(x) = 3 + x^2(-\sin x) + (\cos x)(2x) = 3 - x^2\sin x + 2x\cos x$

7. $y = \sec\theta\ \tan\theta \quad \Rightarrow \quad y' = \sec\theta\,(\sec^2\theta) + \tan\theta\,(\sec\theta\tan\theta) = \sec\theta\,(\sec^2\theta + \tan^2\theta)$. Using the identity $1 + \tan^2\theta = \sec^2\theta$, we can write alternative forms of the answer as $\sec\theta\,(1 + 2\tan^2\theta)$ or $\sec\theta\,(2\sec^2\theta - 1)$.

8. $y = \sin\theta\ \cos\theta \quad \Rightarrow \quad y' = \sin\theta(-\sin\theta) + \cos\theta(\cos\theta) = \cos^2\theta - \sin^2\theta \quad [\text{or } \cos 2\theta]$

9. $f(\theta) = (\theta - \cos\theta)\ \sin\theta \overset{\text{PR}}{\Rightarrow} f'(\theta) = (\theta - \cos\theta)(\cos\theta) + (\sin\theta)(1 + \sin\theta) = \theta\cos\theta - \cos^2\theta + \sin\theta + \sin^2\theta$

10. $g(\theta) = e^\theta(\tan\theta - \theta) \quad \Rightarrow \quad g'(\theta) = e^\theta(\sec^2\theta - 1) + (\tan\theta - \theta)e^\theta = e^\theta(\sec^2\theta - 1 + \tan\theta - \theta)$

11. $H(t) = \cos^2 t = \cos t \cdot \cos t \overset{\text{PR}}{\Rightarrow} H'(t) = \cos t\,(-\sin t) + \cos t\,(-\sin t) = -2\sin t\ \cos t$. Using the identity $\sin 2t = 2\sin t\ \cos t$, we can write an alternative form of the answer as $-\sin 2t$.

12. $f(x) = e^x\sin x + \cos x \Rightarrow f'(x) = e^x(\cos x) + \sin x \cdot e^x + (-\sin x) = e^x(\cos x + \sin x) - \sin x$

13. $f(\theta) = \dfrac{\sin\theta}{1 + \cos\theta} \quad \Rightarrow$

$$f'(\theta) = \frac{(1 + \cos\theta)\cos\theta - (\sin\theta)(-\sin\theta)}{(1 + \cos\theta)^2} = \frac{\cos\theta + \cos^2\theta + \sin^2\theta}{(1 + \cos\theta)^2} = \frac{\cos\theta + 1}{(1 + \cos\theta)^2} = \frac{1}{1 + \cos\theta}$$

14. $y = \dfrac{\cos x}{1 - \sin x} \quad \Rightarrow$

$$y' = \frac{(1 - \sin x)(-\sin x) - \cos x(-\cos x)}{(1 - \sin x)^2} = \frac{-\sin x + \sin^2 x + \cos^2 x}{(1 - \sin x)^2} = \frac{-\sin x + 1}{(1 - \sin x)^2} = \frac{1}{1 - \sin x}$$

15. $y = \dfrac{x}{2 - \tan x} \quad \Rightarrow \quad y' = \dfrac{(2 - \tan x)(1) - x(-\sec^2 x)}{(2 - \tan x)^2} = \dfrac{2 - \tan x + x\sec^2 x}{(2 - \tan x)^2}$

16. $f(t) = \dfrac{\cot t}{e^t} \quad \Rightarrow \quad f'(t) = \dfrac{e^t(-\csc^2 t) - (\cot t)e^t}{(e^t)^2} = \dfrac{e^t(-\csc^2 t - \cot t)}{(e^t)^2} = -\dfrac{\csc^2 t + \cot t}{e^t}$

17. $f(w) = \dfrac{1 + \sec w}{1 - \sec w} \overset{\text{QR}}{\Rightarrow}$

$$f'(w) = \frac{(1 - \sec w)(\sec w\ \tan w) - (1 + \sec w)(-\sec w\ \tan w)}{(1 - \sec w)^2}$$

$$= \frac{\sec w\ \tan w - \sec^2 w\ \tan w + \sec w\ \tan w + \sec^2 w\ \tan w}{(1 - \sec w)^2} = \frac{2\sec w\ \tan w}{(1 - \sec w)^2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $y = \dfrac{\sin t}{1+\tan t} \quad \Rightarrow$

$$y' = \frac{(1+\tan t)\cos t - (\sin t)\sec^2 t}{(1+\tan t)^2} = \frac{\cos t + \sin t - \dfrac{\sin t}{\cos^2 t}}{(1+\tan t)^2} = \frac{\cos t + \sin t - \tan t \sec t}{(1+\tan t)^2}$$

19. $y = \dfrac{t\sin t}{1+t} \quad \Rightarrow$

$$y' = \frac{(1+t)(t\cos t + \sin t) - t\sin t(1)}{(1+t)^2} = \frac{t\cos t + \sin t + t^2\cos t + t\sin t - t\sin t}{(1+t)^2} = \frac{(t^2+t)\cos t + \sin t}{(1+t)^2}$$

20. $g(z) = \dfrac{z}{\sec z + \tan z} \quad \overset{\text{QR}}{\Rightarrow}$

$$g'(z) = \frac{(\sec z + \tan z)(1) - z(\sec z \tan z + \sec^2 z)}{(\sec z + \tan z)^2} = \frac{(\sec z + \tan z)(1) - z\sec z(\tan z + \sec z)}{(\sec z + \tan z)^2}$$

$$= \frac{(1 - z\sec z)(\sec z + \tan z)}{(\sec z + \tan z)^2} = \frac{1 - z\sec z}{\sec z + \tan z}$$

21. Using Exercise 3.2.63(a), $f(\theta) = \theta\cos\theta\,\sin\theta \quad \Rightarrow$

$$f'(\theta) = 1\cos\theta\,\sin\theta + \theta(-\sin\theta)\sin\theta + \theta\cos\theta(\cos\theta) = \cos\theta\,\sin\theta - \theta\sin^2\theta + \theta\cos^2\theta$$

$$= \sin\theta\,\cos\theta + \theta(\cos^2\theta - \sin^2\theta) = \tfrac{1}{2}\sin 2\theta + \theta\cos 2\theta \quad \text{[using double-angle formulas]}$$

22. Using Exercise 3.2.63(a), $f(t) = te^t\cot t \quad \Rightarrow$

$$f'(t) = 1e^t\cot t + te^t\cot t + te^t(-\csc^2 t) = e^t(\cot t + t\cot t - t\csc^2 t)$$

23. $\dfrac{d}{dx}(\csc x) = \dfrac{d}{dx}\left(\dfrac{1}{\sin x}\right) = \dfrac{(\sin x)(0) - 1(\cos x)}{\sin^2 x} = \dfrac{-\cos x}{\sin^2 x} = -\dfrac{1}{\sin x}\cdot\dfrac{\cos x}{\sin x} = -\csc x\,\cot x$

24. $\dfrac{d}{dx}(\sec x) = \dfrac{d}{dx}\left(\dfrac{1}{\cos x}\right) = \dfrac{(\cos x)(0) - 1(-\sin x)}{\cos^2 x} = \dfrac{\sin x}{\cos^2 x} = \dfrac{1}{\cos x}\cdot\dfrac{\sin x}{\cos x} = \sec x\tan x$

25. $\dfrac{d}{dx}(\cot x) = \dfrac{d}{dx}\left(\dfrac{\cos x}{\sin x}\right) = \dfrac{(\sin x)(-\sin x) - (\cos x)(\cos x)}{\sin^2 x} = -\dfrac{\sin^2 x + \cos^2 x}{\sin^2 x} = -\dfrac{1}{\sin^2 x} = -\csc^2 x$

26. $f(x) = \cos x \quad \Rightarrow$

$$f'(x) = \lim_{h\to 0}\frac{f(x+h) - f(x)}{h} = \lim_{h\to 0}\frac{\cos(x+h) - \cos x}{h} = \lim_{h\to 0}\frac{\cos x\,\cos h - \sin x\,\sin h - \cos x}{h}$$

$$= \lim_{h\to 0}\left(\cos x\,\frac{\cos h - 1}{h} - \sin x\,\frac{\sin h}{h}\right) = \cos x\,\lim_{h\to 0}\frac{\cos h - 1}{h} - \sin x\,\lim_{h\to 0}\frac{\sin h}{h}$$

$$= (\cos x)(0) - (\sin x)(1) = -\sin x$$

27. $y = \sin x + \cos x \quad \Rightarrow \quad y' = \cos x - \sin x$, so $y'(0) = \cos 0 - \sin 0 = 1 - 0 = 1$. An equation of the tangent line to the curve $y = \sin x + \cos x$ at the point $(0, 1)$ is $y - 1 = 1(x - 0)$ or $y = x + 1$.

28. $y = x + \sin x \quad \Rightarrow \quad y' = 1 + \cos x$, so $y'(\pi) = 1 + \cos\pi = 1 + (-1) = 0$. An equation of the tangent line to the curve $y = x + \sin x$ at the point (π, π) is $y - \pi = 0(x - \pi)$ or $y = \pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. $y = e^x \cos x + \sin x \quad \Rightarrow \quad y' = e^x(-\sin x) + (\cos x)(e^x) + \cos x = e^x(\cos x - \sin x) + \cos x$, so $y'(0) = e^0(\cos 0 - \sin 0) + \cos 0 = 1(1-0) + 1 = 2$. An equation of the tangent line to the curve $y = e^x \cos x + \sin x$ at the point $(0, 1)$ is $y - 1 = 2(x - 0)$ or $y = 2x + 1$.

30. $y = \dfrac{1 + \sin x}{\cos x} \overset{\text{QR}}{\Rightarrow} y' = \dfrac{\cos x(\cos x) - (1 + \sin x)(-\sin x)}{\cos^2 x} = \dfrac{\cos^2 x + \sin x + \sin^2 x}{\cos^2 x}$

Using the identity $\cos^2 x + \sin^2 x = 1$, we can write $y' = \dfrac{1 + \sin x}{\cos^2 x}$.

$y'(\pi) = \dfrac{1 + \sin \pi}{\cos^2 \pi} = \dfrac{1 + 0}{(-1)^2} = 1$. An equation of the tangent line to the curve $y = \dfrac{1 + \sin x}{\cos x}$ at the point $(\pi, -1)$ is $y - (-1) = 1(x - \pi)$ or $y = x - \pi - 1$.

31. (a) $y = 2x \sin x \quad \Rightarrow \quad y' = 2(x \cos x + \sin x \cdot 1)$. At $\left(\frac{\pi}{2}, \pi\right)$, $y' = 2\left(\frac{\pi}{2} \cos \frac{\pi}{2} + \sin \frac{\pi}{2}\right) = 2(0 + 1) = 2$, and an equation of the tangent line is $y - \pi = 2\left(x - \frac{\pi}{2}\right)$, or $y = 2x$.

(b)

32. (a) $y = 3x + 6 \cos x \quad \Rightarrow \quad y' = 3 - 6 \sin x$. At $\left(\frac{\pi}{3}, \pi + 3\right)$, $y' = 3 - 6 \sin \frac{\pi}{3} = 3 - 6\,\frac{\sqrt{3}}{2} = 3 - 3\sqrt{3}$, and an equation of the tangent line is $y - (\pi + 3) = \left(3 - 3\sqrt{3}\right)\left(x - \frac{\pi}{3}\right)$, or $y = \left(3 - 3\sqrt{3}\right)x + 3 + \pi\sqrt{3}$.

(b)

33. (a) $f(x) = \sec x - x \quad \Rightarrow \quad f'(x) = \sec x \tan x - 1$

(b)

Note that $f' = 0$ where f has a minimum. Also note that f' is negative when f is decreasing and f' is positive when f is increasing.

34. (a) $f(x) = e^x \cos x \quad \Rightarrow \quad f'(x) = e^x(-\sin x) + (\cos x)\, e^x = e^x(\cos x - \sin x) \quad \Rightarrow$

$f''(x) = e^x(-\sin x - \cos x) + (\cos x - \sin x)\, e^x = e^x(-\sin x - \cos x + \cos x - \sin x) = -2e^x \sin x$

(b)

Note that $f' = 0$ where f has a minimum and $f'' = 0$ where f' has a minimum. Also note that f' is negative when f is decreasing and f'' is negative when f' is decreasing.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. $g(\theta) = \dfrac{\sin\theta}{\theta} \overset{\text{QR}}{\Rightarrow} g'(\theta) = \dfrac{\theta(\cos\theta) - (\sin\theta)(1)}{\theta^2} = \dfrac{\theta\cos\theta - \sin\theta}{\theta^2}$

Using the Quotient Rule and $g'(\theta) = \dfrac{\theta\cos\theta - \sin\theta}{\theta^2}$, we get

$$g''(\theta) = \frac{\theta^2\{[\theta(-\sin\theta) + (\cos\theta)(1)] - \cos\theta\} - (\theta\cos\theta - \sin\theta)(2\theta)}{(\theta^2)^2}$$

$$= \frac{-\theta^3\sin\theta + \theta^2\cos\theta - \theta^2\cos\theta - 2\theta^2\cos\theta + 2\theta\sin\theta}{\theta^4} = \frac{\theta(-\theta^2\sin\theta - 2\theta\cos\theta + 2\sin\theta)}{\theta\cdot\theta^3}$$

$$= \frac{-\theta^2\sin\theta - 2\theta\cos\theta + 2\sin\theta}{\theta^3}$$

36. $f(t) = \sec t \;\Rightarrow\; f'(t) = \sec t\,\tan t \;\Rightarrow\; f''(t) = (\sec t)\sec^2 t + (\tan t)\sec t\,\tan t = \sec^3 t + \sec t\,\tan^2 t$, so $f''\left(\frac{\pi}{4}\right) = \left(\sqrt{2}\right)^3 + \sqrt{2}(1)^2 = 2\sqrt{2} + \sqrt{2} = 3\sqrt{2}$.

37. (a) $f(x) = \dfrac{\tan x - 1}{\sec x} \;\Rightarrow$

$$f'(x) = \frac{\sec x(\sec^2 x) - (\tan x - 1)(\sec x\tan x)}{(\sec x)^2} = \frac{\sec x(\sec^2 x - \tan^2 x + \tan x)}{\sec^2 x} = \frac{1 + \tan x}{\sec x}$$

(b) $f(x) = \dfrac{\tan x - 1}{\sec x} = \dfrac{\dfrac{\sin x}{\cos x} - 1}{\dfrac{1}{\cos x}} = \dfrac{\dfrac{\sin x - \cos x}{\cos x}}{\dfrac{1}{\cos x}} = \sin x - \cos x \;\Rightarrow\; f'(x) = \cos x - (-\sin x) = \cos x + \sin x$

(c) From part (a), $f'(x) = \dfrac{1 + \tan x}{\sec x} = \dfrac{1}{\sec x} + \dfrac{\tan x}{\sec x} = \cos x + \sin x$, which is the expression for $f'(x)$ in part (b).

38. (a) $g(x) = f(x)\sin x \;\Rightarrow\; g'(x) = f(x)\cos x + \sin x\cdot f'(x)$, so

$g'\left(\frac{\pi}{3}\right) = f\left(\frac{\pi}{3}\right)\cos\frac{\pi}{3} + \sin\frac{\pi}{3}\cdot f'\left(\frac{\pi}{3}\right) = 4\cdot\frac{1}{2} + \frac{\sqrt{3}}{2}\cdot(-2) = 2 - \sqrt{3}$

(b) $h(x) = \dfrac{\cos x}{f(x)} \;\Rightarrow\; h'(x) = \dfrac{f(x)\cdot(-\sin x) - \cos x\cdot f'(x)}{[f(x)]^2}$, so

$$h'\left(\tfrac{\pi}{3}\right) = \frac{f\left(\frac{\pi}{3}\right)\cdot\left(-\sin\frac{\pi}{3}\right) - \cos\frac{\pi}{3}\cdot f'\left(\frac{\pi}{3}\right)}{\left[f\left(\frac{\pi}{3}\right)\right]^2} = \frac{4\left(-\frac{\sqrt{3}}{2}\right) - \left(\frac{1}{2}\right)(-2)}{4^2} = \frac{-2\sqrt{3} + 1}{16} = \frac{1 - 2\sqrt{3}}{16}$$

39. $f(x) = x + 2\sin x$ has a horizontal tangent when $f'(x) = 0 \;\Leftrightarrow\; 1 + 2\cos x = 0 \;\Leftrightarrow\; \cos x = -\frac{1}{2} \;\Leftrightarrow$ $x = \frac{2\pi}{3} + 2\pi n$ or $\frac{4\pi}{3} + 2\pi n$, where n is an integer. Note that $\frac{4\pi}{3}$ and $\frac{2\pi}{3}$ are $\pm\frac{\pi}{3}$ units from π. This allows us to write the solutions in the more compact equivalent form $(2n+1)\pi \pm \frac{\pi}{3}$, n an integer.

40. $f(x) = e^x\cos x$ has a horizontal tangent when $f'(x) = 0$. $f'(x) = e^x(-\sin x) + (\cos x)e^x = e^x(\cos x - \sin x)$.
$f'(x) = 0 \;\Leftrightarrow\; \cos x - \sin x = 0 \;\Leftrightarrow\; \cos x = \sin x \;\Leftrightarrow\; \tan x = 1 \;\Leftrightarrow\; x = \frac{\pi}{4} + n\pi$, n an integer.

41. (a) $x(t) = 8\sin t \;\Rightarrow\; v(t) = x'(t) = 8\cos t \;\Rightarrow\; a(t) = x''(t) = -8\sin t$

(b) The mass at time $t = \frac{2\pi}{3}$ has position $x\left(\frac{2\pi}{3}\right) = 8\sin\frac{2\pi}{3} = 8\left(\frac{\sqrt{3}}{2}\right) = 4\sqrt{3}$, velocity $v\left(\frac{2\pi}{3}\right) = 8\cos\frac{2\pi}{3} = 8\left(-\frac{1}{2}\right) = -4$, and acceleration $a\left(\frac{2\pi}{3}\right) = -8\sin\frac{2\pi}{3} = -8\left(\frac{\sqrt{3}}{2}\right) = -4\sqrt{3}$. Since $v\left(\frac{2\pi}{3}\right) < 0$, the particle is moving to the left.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. (a) $s(t) = 2\cos t + 3\sin t \quad \Rightarrow \quad v(t) = -2\sin t + 3\cos t \quad \Rightarrow$

$a(t) = -2\cos t - 3\sin t$

(b)

(c) $s = 0 \quad \Rightarrow \quad t_2 \approx 2.55$. So the mass passes through the equilibrium position for the first time when $t \approx 2.55$ s.

(d) $v = 0 \quad \Rightarrow \quad t_1 \approx 0.98$, $s(t_1) \approx 3.61$ cm. So the mass travels a maximum of about 3.6 cm (upward and downward) from its equilibrium position.

(e) The speed $|v|$ is greatest when $s = 0$, that is, when $t = t_2 + n\pi$, n a positive integer.

43.

From the diagram we can see that $\sin\theta = x/10 \quad \Leftrightarrow \quad x = 10\sin\theta$. We want to find the rate of change of x with respect to θ, that is, $dx/d\theta$. Taking the derivative of $x = 10\sin\theta$, we get $dx/d\theta = 10(\cos\theta)$. So when $\theta = \frac{\pi}{3}$, $\frac{dx}{d\theta} = 10\cos\frac{\pi}{3} = 10\left(\frac{1}{2}\right) = 5$ ft/rad.

44. (a) $F = \dfrac{\mu W}{\mu\sin\theta + \cos\theta} \quad \Rightarrow \quad \dfrac{dF}{d\theta} = \dfrac{(\mu\sin\theta + \cos\theta)(0) - \mu W(\mu\cos\theta - \sin\theta)}{(\mu\sin\theta + \cos\theta)^2} = \dfrac{\mu W(\sin\theta - \mu\cos\theta)}{(\mu\sin\theta + \cos\theta)^2}$

(b) $\dfrac{dF}{d\theta} = 0 \quad \Leftrightarrow \quad \mu W(\sin\theta - \mu\cos\theta) = 0 \quad \Leftrightarrow \quad \sin\theta = \mu\cos\theta \quad \Leftrightarrow \quad \tan\theta = \mu \quad \Leftrightarrow \quad \theta = \tan^{-1}\mu$

(c)

From the graph of $F = \dfrac{0.6(50)}{0.6\sin\theta + \cos\theta}$ for $0 \le \theta \le 1$, we see that $\dfrac{dF}{d\theta} = 0 \quad \Rightarrow \quad \theta \approx 0.54$. Checking this with part (b) and $\mu = 0.6$, we calculate $\theta = \tan^{-1} 0.6 \approx 0.54$. So the value from the graph is consistent with the value in part (b).

45. $\displaystyle\lim_{x\to 0}\frac{\sin 5x}{3x} = \lim_{x\to 0}\frac{5}{3}\left(\frac{\sin 5x}{5x}\right) = \frac{5}{3}\lim_{x\to 0}\frac{\sin 5x}{5x} = \frac{5}{3}\lim_{\theta\to 0}\frac{\sin\theta}{\theta} \quad \left[\begin{array}{l}\text{where } \theta = 5x,\\ \text{using Equation 5}\end{array}\right] \quad = \frac{5}{3}\cdot 1 = \frac{5}{3}$

46. $\displaystyle\lim_{x\to 0}\frac{\sin x}{\sin\pi x} = \lim_{x\to 0}\frac{\sin x}{x}\cdot\frac{\pi x}{\sin\pi x}\cdot\frac{1}{\pi} = \lim_{x\to 0}\frac{\sin x}{x}\cdot\lim_{\theta\to 0}\frac{\theta}{\sin\theta}\cdot\frac{1}{\pi} \quad \left[\begin{array}{l}\text{where } \theta = \pi x,\\ \text{using Equation 5}\end{array}\right]$

$\displaystyle = 1\cdot\lim_{\theta\to 0}\frac{1}{\dfrac{\sin\theta}{\theta}}\cdot\frac{1}{\pi} = 1\cdot 1\cdot\frac{1}{\pi} = \frac{1}{\pi}$

47. $\displaystyle\lim_{t\to 0}\frac{\sin 3t}{\sin t} = \lim_{t\to 0}\frac{\sin 3t}{3t}\cdot\frac{t}{\sin t}\cdot 3 = \lim_{t\to 0}\frac{\sin 3t}{3t}\cdot\lim_{t\to 0}\frac{t}{\sin t}\cdot\lim_{t\to 0}3 = \lim_{\theta\to 0}\frac{\sin\theta}{\theta}\cdot\lim_{t\to 0}\frac{1}{\dfrac{\sin t}{t}}\cdot\lim_{t\to 0}3 \qquad [\theta = 3t]$

$= 1\cdot 1\cdot 3 = 3$

48. $\displaystyle\lim_{x\to 0}\frac{\sin^2 3x}{x} = \lim_{x\to 0}\frac{\sin 3x}{3x}\cdot\frac{\sin 3x}{3x}\cdot 3\cdot 3\cdot x = \lim_{x\to 0}\frac{\sin 3x}{3x}\cdot\lim_{x\to 0}\frac{\sin 3x}{3x}\cdot\lim_{x\to 0}3\cdot 3\cdot x$

$\displaystyle = \lim_{\theta\to 0}\frac{\sin\theta}{\theta}\cdot\lim_{\theta\to 0}\frac{\sin\theta}{\theta}\cdot\lim_{x\to 0}9x \qquad [\theta = 3x]$

$= 1\cdot 1\cdot 0 = 0$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. $$\lim_{x\to 0}\frac{\sin x-\sin x\cos x}{x^2}=\lim_{x\to 0}\frac{\sin x\,(1-\cos x)}{x^2}=\lim_{x\to 0}\frac{\sin x}{x}\cdot\lim_{x\to 0}\frac{1-\cos x}{x}$$
$$=1\cdot 0 \quad \text{[by Equations 5 and 6]} \quad =0$$

50. $$\lim_{x\to 0}\frac{1-\sec x}{2x}=\lim_{x\to 0}\frac{1-(1/\cos x)}{2x}\cdot\frac{\cos x}{\cos x}=\lim_{x\to 0}\frac{\cos x-1}{2x\cos x}=\lim_{x\to 0}\frac{\cos x-1}{x}\cdot\lim_{x\to 0}\frac{1}{2\cos x}$$
$$=0\cdot\frac{1}{2} \quad \text{[using Equation 6]} \quad =0$$

51. $$\lim_{x\to 0}\frac{\tan 2x}{x}=\lim_{x\to 0}\frac{\dfrac{\sin 2x}{\cos 2x}}{x}=\lim_{x\to 0}\frac{\sin 2x}{x\cos 2x}=\lim_{x\to 0}\frac{\sin 2x}{2x}\cdot\frac{2}{\cos 2x}$$
$$=\lim_{x\to 0}\frac{\sin 2x}{2x}\cdot\lim_{x\to 0}\frac{2}{\cos 2x}=\lim_{x\to 0}\frac{\sin\theta}{\theta}\cdot\lim_{x\to 0}\frac{2}{\cos 2x} \qquad [\theta=2x]$$
$$=1\cdot\frac{2}{1}=2$$

52. $$\lim_{\theta\to 0}\frac{\sin\theta}{\tan 7\theta}=\lim_{\theta\to 0}\frac{\sin\theta}{\dfrac{\sin 7\theta}{\cos 7\theta}}=\lim_{\theta\to 0}\sin\theta\cdot\frac{\cos 7\theta}{\sin 7\theta}=\lim_{\theta\to 0}\frac{\sin\theta}{\theta}\cdot\cos 7\theta\cdot\frac{7\theta}{\sin 7\theta}\cdot\frac{1}{7}$$
$$=\lim_{\theta\to 0}\frac{\sin\theta}{\theta}\cdot\lim_{\theta\to 0}\cos 7\theta\cdot\lim_{\theta\to 0}\frac{1}{\dfrac{\sin 7\theta}{7\theta}}\cdot\frac{1}{7}=1\cdot 1\cdot\lim_{x\to 0}\frac{1}{\dfrac{\sin x}{x}}\cdot\frac{1}{7} \qquad [x=7\theta]$$
$$=1\cdot 1\cdot 1\cdot\frac{1}{7}=\frac{1}{7}$$

53. $$\lim_{x\to 0}\frac{\sin 3x}{5x^3-4x}=\lim_{x\to 0}\left(\frac{\sin 3x}{3x}\cdot\frac{3}{5x^2-4}\right)=\lim_{x\to 0}\frac{\sin 3x}{3x}\cdot\lim_{x\to 0}\frac{3}{5x^2-4}=1\cdot\left(\frac{3}{-4}\right)=-\frac{3}{4}$$

54. $$\lim_{x\to 0}\frac{\sin 3x\sin 5x}{x^2}=\lim_{x\to 0}\left(\frac{3\sin 3x}{3x}\cdot\frac{5\sin 5x}{5x}\right)=\lim_{x\to 0}\frac{3\sin 3x}{3x}\cdot\lim_{x\to 0}\frac{5\sin 5x}{5x}$$
$$=3\lim_{x\to 0}\frac{\sin 3x}{3x}\cdot 5\lim_{x\to 0}\frac{\sin 5x}{5x}=3(1)\cdot 5(1)=15$$

55. Divide numerator and denominator by θ. [$\sin\theta$ also works.]

$$\lim_{\theta\to 0}\frac{\sin\theta}{\theta+\tan\theta}=\lim_{\theta\to 0}\frac{\dfrac{\sin\theta}{\theta}}{1+\dfrac{\sin\theta}{\theta}\cdot\dfrac{1}{\cos\theta}}=\frac{\lim\limits_{\theta\to 0}\dfrac{\sin\theta}{\theta}}{1+\lim\limits_{\theta\to 0}\dfrac{\sin\theta}{\theta}\lim\limits_{\theta\to 0}\dfrac{1}{\cos\theta}}=\frac{1}{1+1\cdot 1}=\frac{1}{2}$$

56. $$\lim_{x\to 0}\csc x\,\sin(\sin x)=\lim_{x\to 0}\frac{\sin(\sin x)}{\sin x}=\lim_{\theta\to 0}\frac{\sin\theta}{\theta} \quad [\text{As } x\to 0,\ \theta=\sin x\to 0.] \quad =1$$

57. $$\lim_{\theta\to 0}\frac{\cos\theta-1}{2\theta^2}=\lim_{\theta\to 0}\frac{\cos\theta-1}{2\theta^2}\cdot\frac{\cos\theta+1}{\cos\theta+1}=\lim_{\theta\to 0}\frac{\cos^2\theta-1}{2\theta^2(\cos\theta+1)}=\lim_{\theta\to 0}\frac{-\sin^2\theta}{2\theta^2(\cos\theta+1)}$$
$$=-\frac{1}{2}\lim_{\theta\to 0}\frac{\sin\theta}{\theta}\cdot\frac{\sin\theta}{\theta}\cdot\frac{1}{\cos\theta+1}=-\frac{1}{2}\lim_{\theta\to 0}\frac{\sin\theta}{\theta}\cdot\lim_{\theta\to 0}\frac{\sin\theta}{\theta}\cdot\lim_{\theta\to 0}\frac{1}{\cos\theta+1}$$
$$=-\frac{1}{2}\cdot 1\cdot 1\cdot\frac{1}{1+1}=-\frac{1}{4}$$

58. $$\lim_{x\to 0}\frac{\sin(x^2)}{x}=\lim_{x\to 0}\left[x\cdot\frac{\sin(x^2)}{x\cdot x}\right]=\lim_{x\to 0}x\cdot\lim_{x\to 0}\frac{\sin(x^2)}{x^2}=0\cdot\lim_{y\to 0^+}\frac{\sin y}{y} \quad \left[\text{where } y=x^2\right] \quad =0\cdot 1=0$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. $\lim\limits_{x\to\pi/4} \dfrac{1-\tan x}{\sin x-\cos x} = \lim\limits_{x\to\pi/4} \dfrac{\left(1-\dfrac{\sin x}{\cos x}\right)\cdot\cos x}{(\sin x-\cos x)\cdot\cos x} = \lim\limits_{x\to\pi/4} \dfrac{\cos x-\sin x}{(\sin x-\cos x)\cos x} = \lim\limits_{x\to\pi/4} \dfrac{-1}{\cos x} = \dfrac{-1}{1/\sqrt{2}} = -\sqrt{2}$

60. $\lim\limits_{x\to1} \dfrac{\sin(x-1)}{x^2+x-2} = \lim\limits_{x\to1} \dfrac{\sin(x-1)}{(x+2)(x-1)} = \lim\limits_{x\to1} \dfrac{1}{x+2} \lim\limits_{x\to1} \dfrac{\sin(x-1)}{x-1} = \dfrac{1}{3}\cdot 1 = \dfrac{1}{3}$

61. $\dfrac{d}{dx}(\sin x) = \cos x \quad\Rightarrow\quad \dfrac{d^2}{dx^2}(\sin x) = -\sin x \quad\Rightarrow\quad \dfrac{d^3}{dx^3}(\sin x) = -\cos x \quad\Rightarrow\quad \dfrac{d^4}{dx^4}(\sin x) = \sin x.$

The derivatives of $\sin x$ occur in a cycle of four. Since $99 = 4(24)+3$, we have $\dfrac{d^{99}}{dx^{99}}(\sin x) = \dfrac{d^3}{dx^3}(\sin x) = -\cos x$.

62. Let $f(x) = x\sin x$ and $h(x) = \sin x$, so $f(x) = xh(x)$. Then $f'(x) = h(x) + xh'(x)$,

$f''(x) = h'(x) + h'(x) + xh''(x) = 2h'(x) + xh''(x)$,

$f'''(x) = 2h''(x) + h''(x) + xh'''(x) = 3h''(x) + xh'''(x), \cdots, f^{(n)}(x) = nh^{(n-1)}(x) + xh^{(n)}(x)$.

Since $34 = 4(8)+2$, we have $h^{(34)}(x) = h^{(2)}(x) = \dfrac{d^2}{dx^2}(\sin x) = -\sin x$ and $h^{(35)}(x) = -\cos x$.

Thus, $\dfrac{d^{35}}{dx^{35}}(x\sin x) = 35h^{(34)}(x) + xh^{(35)}(x) = -35\sin x - x\cos x$.

63. $y = A\sin x + B\cos x \quad\Rightarrow\quad y' = A\cos x - B\sin x \quad\Rightarrow\quad y'' = -A\sin x - B\cos x$. Substituting these expressions for y, y', and y'' into the given differential equation $y'' + y' - 2y = \sin x$ gives us

$(-A\sin x - B\cos x) + (A\cos x - B\sin x) - 2(A\sin x + B\cos x) = \sin x \quad\Leftrightarrow$

$-3A\sin x - B\sin x + A\cos x - 3B\cos x = \sin x \quad\Leftrightarrow\quad (-3A-B)\sin x + (A-3B)\cos x = 1\sin x$, so we must have $-3A - B = 1$ and $A - 3B = 0$ (since 0 is the coefficient of $\cos x$ on the right side). Solving for A and B, we add the first equation to three times the second to get $B = -\frac{1}{10}$ and $A = -\frac{3}{10}$.

64. (a) Let $\theta = \dfrac{1}{x}$. Then as $x\to\infty$, $\theta\to0^+$, and $\lim\limits_{x\to\infty} x\sin\dfrac{1}{x} = \lim\limits_{\theta\to0^+} \dfrac{1}{\theta}\sin\theta = \lim\limits_{\theta\to0} \dfrac{\sin\theta}{\theta} = 1$.

(b) Since $-1 \le \sin(1/x) \le 1$, we have (as illustrated in the figure) $-|x| \le x\sin(1/x) \le |x|$. We know that $\lim\limits_{x\to0}(|x|) = 0$ and $\lim\limits_{x\to0}(-|x|) = 0$; so by the Squeeze Theorem, $\lim\limits_{x\to0} x\sin(1/x) = 0$.

(c)

65. (a) $\dfrac{d}{dx}\tan x = \dfrac{d}{dx}\dfrac{\sin x}{\cos x} \quad\Rightarrow\quad \sec^2 x = \dfrac{\cos x\,\cos x - \sin x\,(-\sin x)}{\cos^2 x} = \dfrac{\cos^2 x + \sin^2 x}{\cos^2 x}$. So $\sec^2 x = \dfrac{1}{\cos^2 x}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\dfrac{d}{dx}\sec x = \dfrac{d}{dx}\dfrac{1}{\cos x} \quad\Rightarrow\quad \sec x\,\tan x = \dfrac{(\cos x)(0) - 1(-\sin x)}{\cos^2 x}$. So $\sec x\,\tan x = \dfrac{\sin x}{\cos^2 x}$.

(c) $\dfrac{d}{dx}(\sin x + \cos x) = \dfrac{d}{dx}\dfrac{1+\cot x}{\csc x} \quad\Rightarrow$

$$\cos x - \sin x = \frac{\csc x\,(-\csc^2 x) - (1+\cot x)(-\csc x\,\cot x)}{\csc^2 x} = \frac{\csc x\,[-\csc^2 x + (1+\cot x)\,\cot x]}{\csc^2 x}$$

$$= \frac{-\csc^2 x + \cot^2 x + \cot x}{\csc x} = \frac{-1+\cot x}{\csc x}$$

So $\cos x - \sin x = \dfrac{\cot x - 1}{\csc x}$.

66. We get the following formulas for r and h in terms of θ:

$$\sin\frac{\theta}{2} = \frac{r}{10} \quad\Rightarrow\quad r = 10\sin\frac{\theta}{2} \quad\text{and}\quad \cos\frac{\theta}{2} = \frac{h}{10} \quad\Rightarrow\quad h = 10\cos\frac{\theta}{2}$$

Now $A(\theta) = \frac{1}{2}\pi r^2$ and $B(\theta) = \frac{1}{2}(2r)h = rh$. So

$$\lim_{\theta\to 0^+}\frac{A(\theta)}{B(\theta)} = \lim_{\theta\to 0^+}\frac{\frac{1}{2}\pi r^2}{rh} = \tfrac{1}{2}\pi\lim_{\theta\to 0^+}\frac{r}{h} = \tfrac{1}{2}\pi\lim_{\theta\to 0^+}\frac{10\sin(\theta/2)}{10\cos(\theta/2)}$$

$$= \tfrac{1}{2}\pi\lim_{\theta\to 0^+}\tan(\theta/2) = 0$$

67. By the definition of radian measure, $s = r\theta$, where r is the radius of the circle. By drawing the bisector of the angle θ, we can see that $\sin\dfrac{\theta}{2} = \dfrac{d/2}{r} \quad\Rightarrow\quad d = 2r\sin\dfrac{\theta}{2}$. So $\displaystyle\lim_{\theta\to 0^+}\frac{s}{d} = \lim_{\theta\to 0^+}\frac{r\theta}{2r\sin(\theta/2)} = \lim_{\theta\to 0^+}\frac{2\cdot(\theta/2)}{2\sin(\theta/2)} = \lim_{\theta\to 0}\frac{\theta/2}{\sin(\theta/2)} = 1$.

[This is just the reciprocal of the limit $\displaystyle\lim_{x\to 0}\frac{\sin x}{x} = 1$ combined with the fact that as $\theta\to 0$, $\frac{\theta}{2}\to 0$ also.]

68. (a)

It appears that $f(x) = \dfrac{x}{\sqrt{1-\cos 2x}}$ has a jump discontinuity at $x = 0$.

(b) Using the identity $\cos 2x = 1 - \sin^2 x$, we have $\dfrac{x}{\sqrt{1-\cos 2x}} = \dfrac{x}{\sqrt{1-(1-2\sin^2 x)}} = \dfrac{x}{\sqrt{2\sin^2 x}} = \dfrac{x}{\sqrt{2}\,|\sin x|}$.

Thus,
$$\lim_{x\to 0^-}\frac{x}{\sqrt{1-\cos 2x}} = \lim_{x\to 0^-}\frac{x}{\sqrt{2}\,|\sin x|} = \frac{1}{\sqrt{2}}\lim_{x\to 0^-}\frac{x}{-(\sin x)}$$

$$= -\frac{1}{\sqrt{2}}\lim_{x\to 0^-}\frac{1}{\sin x/x} = -\frac{1}{\sqrt{2}}\cdot\frac{1}{1} = -\frac{\sqrt{2}}{2}$$

Evaluating $\displaystyle\lim_{x\to 0^+} f(x)$ is similar, but $|\sin x| = +\sin x$, so we get $\frac{1}{2}\sqrt{2}$. These values appear to be reasonable values for the graph, so they confirm our answer to part (a).

Another method: Multiply numerator and denominator by $\sqrt{1+\cos 2x}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3.4 The Chain Rule

1. Let $u = g(x) = 5 - x^4$ and $y = f(u) = u^3$. Then $\dfrac{dy}{dx} = \dfrac{dy}{du}\dfrac{du}{dx} = (3u^2)(-4x^3) = 3(5-x^4)^2(-4x^3) = -12x^3(5-x^4)^2$.

2. Let $u = g(x) = x^3 + 2$ and $y = f(u) = \sqrt{u}$. Then $\dfrac{dy}{dx} = \dfrac{dy}{du}\dfrac{du}{dx} = \dfrac{1}{2\sqrt{u}} \cdot 3x^2 = \dfrac{1}{2\sqrt{x^3+2}} \cdot 3x^2 = \dfrac{3x^2}{2\sqrt{x^3+2}}$.

3. Let $u = g(x) = \cos x$ and $y = f(u) = \sin u$. Then

$$\frac{dy}{dx} = \frac{dy}{du}\frac{du}{dx} = (\cos u)(-\sin x) = (\cos(\cos x))(-\sin x) = -\sin x\,\cos(\cos x).$$

4. Let $u = g(x) = x^2$ and $y = f(u) = \tan u$. Then $\dfrac{dy}{dx} = \dfrac{dy}{du}\dfrac{du}{dx} = (\sec^2 u)(2x) = (\sec^2(x^2))(2x) = 2x\sec^2(x^2)$.

5. Let $u = g(x) = \sqrt{x}$ and $y = f(u) = e^u$. Then $\dfrac{dy}{dx} = \dfrac{dy}{du}\dfrac{du}{dx} = (e^u)\left(\frac{1}{2}x^{-1/2}\right) = e^{\sqrt{x}} \cdot \dfrac{1}{2\sqrt{x}} = \dfrac{e^{\sqrt{x}}}{2\sqrt{x}}$.

6. Let $u = g(x) = e^x + 1$ and $y = f(u) = \sqrt[3]{u} = u^{1/3}$. Then

$$\frac{dy}{dx} = \frac{dy}{du}\frac{du}{dx} = \left(\frac{1}{3}u^{-2/3}\right)(e^x) = \left(\frac{1}{3\sqrt[3]{(e^x+1)^2}}\right)(e^x) = \frac{e^x}{3\sqrt[3]{(e^x+1)^2}}.$$

Note: The notation $\overset{\text{CR}}{\Rightarrow}$ indicates the use of the Chain Rule.

7. $f(x) = (2x^3 - 5x^2 + 4)^5 \quad \overset{\text{CR}}{\Rightarrow}$

$$\begin{aligned} f'(x) &= 5(2x^3 - 5x^2 + 4)^4 \cdot \frac{d}{dx}(2x^3 - 5x^2 + 4) = 5(2x^3 - 5x^2 + 4)^4(6x^2 - 10x) \\ &= 5(2x^3 - 5x^2 + 4)^4 \cdot 2x(3x - 5) = 10x(2x^3 - 5x^2 + 4)^4(3x - 5) \end{aligned}$$

8. $f(x) = (x^5 + 3x^2 - x)^{50} \quad \overset{\text{CR}}{\Rightarrow}$

$$f'(x) = 50(x^5 + 3x^2 - x)^{49} \cdot \frac{d}{dx}(x^5 + 3x^2 - x) = 50(x^5 + 3x^2 - x)^{49}(5x^4 + 6x - 1)$$

9. $f(x) = \sqrt{5x+1} = (5x+1)^{1/2} \quad \overset{\text{CR}}{\Rightarrow} \quad f'(x) = \frac{1}{2}(5x+1)^{-1/2} \cdot \dfrac{d}{dx}(5x+1) = \frac{1}{2}(5x+1)^{-1/2}(5) = \dfrac{5}{2\sqrt{5x+1}}$

10. $f(x) = \dfrac{1}{\sqrt[3]{x^2-1}} = (x^2-1)^{-1/3} \quad \overset{\text{CR}}{\Rightarrow} \quad f'(x) = -\dfrac{1}{3}(x^2-1)^{-4/3}(2x) = \dfrac{-2x}{3(x^2-1)^{4/3}}$

11. $g(t) = \dfrac{1}{(2t+1)^2} = (2t+1)^{-2} \quad \overset{\text{CR}}{\Rightarrow} \quad g'(t) = -2(2t+1)^{-3} \cdot \dfrac{d}{dt}(2t+1) = -2(2t+1)^{-3}(2) = -\dfrac{4}{(2t+1)^3}$

12. $F(t) = \left(\dfrac{1}{2t+1}\right)^4 = [(2t+1)^{-1}]^4 = (2t+1)^{-4} \quad \overset{\text{CR}}{\Rightarrow}$

$$F'(t) = -4(2t+1)^{-5} \cdot \frac{d}{dt}(2t+1) = -4(2t+1)^{-5}(2) = -\frac{8}{(2t+1)^5}$$

13. $f(\theta) = \cos(\theta^2) \quad \overset{\text{CR}}{\Rightarrow} \quad f'(\theta) = -\sin(\theta^2) \cdot \dfrac{d}{d\theta}(\theta^2) = -\sin(\theta^2) \cdot (2\theta) = -2\theta\sin(\theta^2)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $g(\theta) = \cos^2\theta = (\cos\theta)^2 \overset{\text{CR}}{\Rightarrow} g'(\theta) = 2(\cos\theta)^1(-\sin\theta) = -2\sin\theta\cos\theta = -\sin 2\theta$

15. $g(x) = e^{x^2-x} \overset{\text{CR}}{\Rightarrow} g'(x) = e^{x^2-x}\cdot\dfrac{d}{dx}(x^2-x) = e^{x^2-x}(2x-1)$

16. Using Formula 5 and the Chain Rule, $y = 5^{\sqrt{x}} \Rightarrow y' = 5^{\sqrt{x}}\ln 5\cdot\dfrac{d}{dx}\left(\sqrt{x}\right) = 5^{\sqrt{x}}\ln 5\cdot\dfrac{1}{2\sqrt{x}} = \dfrac{5^{\sqrt{x}}\ln 5}{2\sqrt{x}}$.

17. Using the Product Rule and the Chain Rule, $y = x^2e^{-3x} \Rightarrow$

$y' = x^2e^{-3x}(-3) + e^{-3x}(2x) = e^{-3x}(-3x^2+2x) = xe^{-3x}(2-3x)$.

18. Using the Product Rule and the Chain Rule, $f(t) = t\sin\pi t \Rightarrow f'(t) = t(\cos\pi t)\cdot\pi + (\sin\pi t)\cdot 1 = \pi t\cos\pi t + \sin\pi t$.

19. $f(t) = e^{at}\sin bt \Rightarrow f'(t) = e^{at}(\cos bt)\cdot b + (\sin bt)e^{at}\cdot a = e^{at}(b\cos bt + a\sin bt)$

20. $A(r) = \sqrt{r}\cdot e^{r^2+1} \Rightarrow$

$$A'(r) = \sqrt{r}\cdot e^{r^2+1}\cdot\frac{d}{dr}\left(r^2+1\right) + e^{r^2+1}\cdot\frac{d}{dr}\left(\sqrt{r}\right) = \sqrt{r}\cdot e^{r^2+1}\cdot 2r + e^{r^2+1}\cdot\frac{1}{2\sqrt{r}}$$
$$= e^{r^2+1}\left(2r\sqrt{r} + \frac{1}{2\sqrt{r}}\right) \text{ or } e^{r^2+1}\left(\frac{4r^2+1}{2\sqrt{r}}\right)$$

21. $F(x) = (4x+5)^3(x^2-2x+5)^4 \Rightarrow$

$$\begin{aligned}F'(x) &= (4x+5)^3\cdot 4(x^2-2x+5)^3(2x-2) + (x^2-2x+5)^4\cdot 3(4x+5)^2\cdot 4\\ &= 4(4x+5)^2(x^2-2x+5)^3\left[(4x+5)(2x-2) + (x^2-2x+5)\cdot 3\right]\\ &= 4(4x+5)^2(x^2-2x+5)^3(8x^2+2x-10+3x^2-6x+15)\\ &= 4(4x+5)^2(x^2-2x+5)^3(11x^2-4x+5)\end{aligned}$$

22. $G(z) = (1-4z)^2\sqrt{z^2+1} \Rightarrow$

$$G'(z) = (1-4z)^2\cdot\frac{1}{2\sqrt{z^2+1}}\cdot 2z + \sqrt{z^2+1}\cdot 2(1-4z)^1(-4) = 2(1-4z)\left[\frac{(1-4z)z}{2\sqrt{z^2+1}} - 4\sqrt{z^2+1}\right]$$
$$= 2(1-4z)\left[\frac{(1-4z)z}{2\sqrt{z^2+1}} - \frac{8(z^2+1)}{2\sqrt{z^2+1}}\right] = 2(1-4z)\left(\frac{z-4z^2-8z^2-8}{2\sqrt{z^2+1}}\right)$$
$$= (1-4z)\left(\frac{-12z^2+z-8}{\sqrt{z^2+1}}\right) \text{ or } (4z-1)\left(\frac{12z^2-z+8}{\sqrt{z^2+1}}\right)$$

23. $y = \sqrt{\dfrac{x}{x+1}} = \left(\dfrac{x}{x+1}\right)^{1/2} \Rightarrow$

$$y' = \frac{1}{2}\left(\frac{x}{x+1}\right)^{-1/2}\frac{d}{dx}\left(\frac{x}{x+1}\right) = \frac{1}{2}\frac{x^{-1/2}}{(x+1)^{-1/2}}\frac{(x+1)(1)-x(1)}{(x+1)^2}$$
$$= \frac{1}{2}\frac{(x+1)^{1/2}}{x^{1/2}}\frac{1}{(x+1)^2} = \frac{1}{2\sqrt{x}(x+1)^{3/2}}$$

24. $y = \left(x+\dfrac{1}{x}\right)^5 \Rightarrow y' = 5\left(x+\dfrac{1}{x}\right)^4\dfrac{d}{dx}\left(x+\dfrac{1}{x}\right) = 5\left(x+\dfrac{1}{x}\right)^4\left(1-\dfrac{1}{x^2}\right)$.

Another form of the answer is $\dfrac{5(x^2+1)^4(x^2-1)}{x^6}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $y = e^{\tan\theta} \quad \Rightarrow \quad y' = e^{\tan\theta}\dfrac{d}{d\theta}(\tan\theta) = (\sec^2\theta)e^{\tan\theta}$

26. Using Formula 5 and the Chain Rule, $f(t) = 2^{t^3} \quad \Rightarrow \quad f'(t) = 2^{t^3}\ln 2\,\dfrac{d}{dt}\left(t^3\right) = 3(\ln 2)t^2 2^{t^3}$.

27. $g(u) = \left(\dfrac{u^3-1}{u^3+1}\right)^8 \quad \Rightarrow$

$$g'(u) = 8\left(\frac{u^3-1}{u^3+1}\right)^7 \frac{d}{du}\frac{u^3-1}{u^3+1} = 8\,\frac{\left(u^3-1\right)^7}{(u^3+1)^7}\,\frac{(u^3+1)(3u^2)-(u^3-1)(3u^2)}{(u^3+1)^2}$$

$$= 8\,\frac{\left(u^3-1\right)^7}{(u^3+1)^7}\,\frac{3u^2[(u^3+1)-(u^3-1)]}{(u^3+1)^2} = 8\frac{\left(u^3-1\right)^7}{(u^3+1)^7}\,\frac{3u^2(2)}{(u^3+1)^2} = \frac{48u^2\left(u^3-1\right)^7}{(u^3+1)^9}$$

28. $s(t) = \sqrt{\dfrac{1+\sin t}{1+\cos t}} = \left(\dfrac{1+\sin t}{1+\cos t}\right)^{1/2} \quad \Rightarrow$

$$s'(t) = \frac{1}{2}\left(\frac{1+\sin t}{1+\cos t}\right)^{-1/2}\frac{(1+\cos t)\cos t-(1+\sin t)(-\sin t)}{(1+\cos t)^2}$$

$$= \frac{1}{2}\,\frac{(1+\sin t)^{-1/2}}{(1+\cos t)^{-1/2}}\,\frac{\cos t+\cos^2 t+\sin t+\sin^2 t}{(1+\cos t)^2} = \frac{\cos t+\sin t+1}{2\sqrt{1+\sin t}\,(1+\cos t)^{3/2}}$$

29. Using Formula 5 and the Chain Rule, $r(t) = 10^{2\sqrt{t}} \quad \Rightarrow$

$$r'(t) = 10^{2\sqrt{t}}\ln 10\,\frac{d}{dt}\left(2\sqrt{t}\right) = 10^{2\sqrt{t}}\ln 10\left(2\cdot\tfrac{1}{2}t^{-1/2}\right) = \frac{(\ln 10)\,10^{2\sqrt{t}}}{\sqrt{t}}.$$

30. $f(z) = e^{z/(z-1)} \quad \Rightarrow \quad f'(z) = e^{z/(z-1)}\dfrac{d}{dz}\dfrac{z}{z-1} = e^{z/(z-1)}\dfrac{(z-1)(1)-z(1)}{(z-1)^2} = -\dfrac{e^{z/(z-1)}}{(z-1)^2}$

31. $H(r) = \dfrac{(r^2-1)^3}{(2r+1)^5} \quad \Rightarrow$

$$H'(r) = \frac{(2r+1)^5\cdot 3(r^2-1)^2(2r)-(r^2-1)^3\cdot 5(2r+1)^4(2)}{[(2r+1)^5]^2} = \frac{2(2r+1)^4(r^2-1)^2[3r(2r+1)-5(r^2-1)]}{(2r+1)^{10}}$$

$$= \frac{2(r^2-1)^2(6r^2+3r-5r^2+5)}{(2r+1)^6} = \frac{2(r^2-1)^2(r^2+3r+5)}{(2r+1)^6}$$

32. $J(\theta) = \tan^2(n\theta) = [\tan(n\theta)]^2 \quad \Rightarrow$

$$J'(\theta) = 2\,[\tan(n\theta)]^1\,\frac{d}{d\theta}\tan(n\theta) = 2\tan(n\theta)\,\sec^2(n\theta)\cdot n = 2n\tan(n\theta)\,\sec^2(n\theta)$$

33. $F(t) = e^{t\sin 2t} \quad \Rightarrow$

$$F'(t) = e^{t\sin 2t}(t\sin 2t)' = e^{t\sin 2t}(t\cdot 2\cos 2t+\sin 2t\cdot 1) = e^{t\sin 2t}(2t\cos 2t+\sin 2t)$$

34. $F(t) = \dfrac{t^2}{\sqrt{t^3+1}} \quad \Rightarrow$

$$F'(t) = \frac{(t^3+1)^{1/2}(2t)-t^2\cdot\frac{1}{2}(t^3+1)^{-1/2}(3t^2)}{\left(\sqrt{t^3+1}\right)^2} = \frac{t(t^3+1)^{-1/2}\left[2(t^3+1)-\frac{3}{2}t^3\right]}{(t^3+1)^1} = \frac{t\left(\frac{1}{2}t^3+2\right)}{(t^3+1)^{3/2}} = \frac{t(t^3+4)}{2(t^3+1)^{3/2}}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. Using Formula 5 and the Chain Rule, $G(x) = 4^{C/x} \quad \Rightarrow$

$$G'(x) = 4^{C/x}\,(\ln 4)\,\frac{d}{dx}\frac{C}{x} \quad \left[\frac{C}{x} = Cx^{-1}\right] \quad = 4^{C/x}\,(\ln 4)\left(-Cx^{-2}\right) = -C\,(\ln 4)\,\frac{4^{C/x}}{x^2}.$$

36. $U(y) = \left(\dfrac{y^4+1}{y^2+1}\right)^5 \quad \Rightarrow$

$$U'(y) = 5\left(\frac{y^4+1}{y^2+1}\right)^4 \frac{(y^2+1)(4y^3)-(y^4+1)(2y)}{(y^2+1)^2} = \frac{5(y^4+1)^4 2y[2y^2(y^2+1)-(y^4+1)]}{(y^2+1)^4(y^2+1)^2}$$

$$= \frac{10y(y^4+1)^4(y^4+2y^2-1)}{(y^2+1)^6}$$

37. $f\,(x) = \sin x\,\cos(1-x^2) \quad \Rightarrow$

$$f'(x) = \sin x\left[-\sin(1-x^2)(-2x)\right] + \cos(1-x^2)\cdot\cos x = 2x\sin x\,\sin(1-x^2) + \cos x\,\cos(1-x^2)$$

38. $g(x) = e^{-x}\cos(x^2) \quad \Rightarrow \quad g'(x) = e^{-x}[-\sin(x^2)]\cdot 2x + \cos(x^2)\cdot e^{-x}(-1) = -e^{-x}[2x\sin(x^2)+\cos(x^2)]$

39. $F(t) = \tan\sqrt{1+t^2} \Rightarrow F'(t) = \sec^2\sqrt{1+t^2}\cdot\dfrac{1}{2\sqrt{1+t^2}}\cdot 2t = \dfrac{t\sec^2\sqrt{1+t^2}}{\sqrt{1+t^2}}$

40. $G(z) = (1+\cos^2 z)^3 \Rightarrow G'(z) = 3(1+\cos^2 z)^2\,[2(\cos z)(-\sin z)] = -6\cos z\,\sin z(1+\cos^2 z)^2$

41. $y = \sin^2(x^2+1) \quad \Rightarrow \quad y' = 2\sin(x^2+1)\cdot\cos(x^2+1)\cdot 2x = 4x\sin(x^2+1)\,\cos(x^2+1)$

42. $y = e^{\sin 2x} + \sin(e^{2x}) \quad \Rightarrow$

$$y' = e^{\sin 2x}\frac{d}{dx}\sin 2x + \cos(e^{2x})\frac{d}{dx}e^{2x} = e^{\sin 2x}(\cos 2x)\cdot 2 + \cos(e^{2x})\,e^{2x}\cdot 2 = 2\cos 2x\,e^{\sin 2x} + 2e^{2x}\cos(e^{2x})$$

43. $g(x) = \sin\left(\dfrac{e^x}{1+e^x}\right) \quad \Rightarrow$

$$g'(x) = \cos\left(\frac{e^x}{1+e^x}\right)\cdot\frac{(1+e^x)e^x - e^x(e^x)}{(1+e^x)^2} = \cos\left(\frac{e^x}{1+e^x}\right)\cdot\frac{e^x(1+e^x-e^x)}{(1+e^x)^2} = \frac{e^x}{(1+e^x)^2}\cos\left(\frac{e^x}{1+e^x}\right)$$

44. $f(t) = e^{1/t}\sqrt{t^2-1} \quad \Rightarrow$

$$f'(t) = e^{1/t}\cdot\frac{1}{2\sqrt{t^2-1}}\cdot 2t + \sqrt{t^2-1}\cdot e^{1/t}\cdot\left(-\frac{1}{t^2}\right) \qquad \left[\frac{1}{t} = t^{-1};\ \frac{d}{dt}\left(t^{-1}\right) = -t^{-2} = -\frac{1}{t^2}\right]$$

$$= e^{1/t}\left(\frac{t}{\sqrt{t^2-1}} - \frac{\sqrt{t^2-1}}{t^2}\right) \text{ or } e^{1/t}\left(\frac{t^3-t^2+1}{t^2\sqrt{t^2-1}}\right)$$

45. $f(t) = \tan(\sec(\cos t)) \quad \Rightarrow$

$$f'(t) = \sec^2(\sec(\cos t))\,\frac{d}{dt}\,\sec(\cos t) = \sec^2(\sec(\cos t))[\sec(\cos t)\,\tan(\cos t)]\,\frac{d}{dt}\,\cos t$$

$$= -\sec^2(\sec(\cos t))\,\sec(\cos t)\,\tan(\cos t)\,\sin t$$

46. $y = \sqrt{x+\sqrt{x+\sqrt{x}}} \quad \Rightarrow \quad y' = \frac{1}{2}\left(x+\sqrt{x+\sqrt{x}}\right)^{-1/2}\left[1+\frac{1}{2}\left(x+\sqrt{x}\right)^{-1/2}\left(1+\frac{1}{2}x^{-1/2}\right)\right]$

47. $f(x) = e^{\sin^2(x^2)} \Rightarrow f'(x) = e^{\sin^2(x^2)}\cdot 2\sin(x^2)\cdot\cos(x^2)\cdot 2x = 4x\sin(x^2)\,\cos(x^2)e^{\sin^2(x^2)}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. $y = 2^{3^{4^x}} \quad \Rightarrow$

$$y' = 2^{3^{4^x}}(\ln 2)\frac{d}{dx}3^{4^x} = 2^{3^{4^x}}(\ln 2)3^{4^x}(\ln 3)\frac{d}{dx}4^x = 2^{3^{4^x}}(\ln 2)3^{4^x}(\ln 3)4^x(\ln 4) = (\ln 2)(\ln 3)(\ln 4)4^x 3^{4^x} 2^{3^{4^x}}$$

49. $y = \left(3^{\cos(x^2)} - 1\right)^4 \quad \Rightarrow$

$$y' = 4\left(3^{\cos(x^2)} - 1\right)^3 \cdot 3^{\cos(x^2)} \ln 3 \cdot (-\sin(x^2)) \cdot 2x = -8x(\ln 3)\sin(x^2)\, 3^{\cos(x^2)}\left(3^{\cos(x^2)} - 1\right)^3$$

50. $y = \sin(\theta + \tan(\theta + \cos\theta)) \quad \Rightarrow \quad y' = \cos(\theta + \tan(\theta + \cos\theta)) \cdot [1 + \sec^2(\theta + \cos\theta) \cdot (1 - \sin\theta)]$

51. $y = \cos\sqrt{\sin(\tan \pi x)} = \cos(\sin(\tan \pi x))^{1/2} \quad \Rightarrow$

$$y' = -\sin(\sin(\tan \pi x))^{1/2} \cdot \frac{d}{dx}(\sin(\tan \pi x))^{1/2} = -\sin(\sin(\tan \pi x))^{1/2} \cdot \tfrac{1}{2}(\sin(\tan \pi x))^{-1/2} \cdot \frac{d}{dx}(\sin(\tan \pi x))$$

$$= \frac{-\sin\sqrt{\sin(\tan \pi x)}}{2\sqrt{\sin(\tan \pi x)}} \cdot \cos(\tan \pi x) \cdot \frac{d}{dx}\tan \pi x = \frac{-\sin\sqrt{\sin(\tan \pi x)}}{2\sqrt{\sin(\tan \pi x)}} \cdot \cos(\tan \pi x) \cdot \sec^2(\pi x) \cdot \pi$$

$$= \frac{-\pi\cos(\tan \pi x)\sec^2(\pi x)\sin\sqrt{\sin(\tan \pi x)}}{2\sqrt{\sin(\tan \pi x)}}$$

52. $y = \sin^3(\cos(x^2)) \quad \Rightarrow$

$$y' = 3\sin^2(\cos(x^2)) \cdot \cos(\cos(x^2)) \cdot [-\sin(x^2) \cdot 2x] = -6x\sin(x^2)\ \sin^2(\cos(x^2))\ \cos(\cos(x^2))$$

53. $y = \cos(\sin 3\theta) \quad \Rightarrow \quad y' = -\sin(\sin 3\theta) \cdot (\cos 3\theta) \cdot 3 = -3\cos 3\theta\ \sin(\sin 3\theta) \quad \Rightarrow$

$$y'' = -3\,[(\cos 3\theta)\cos(\sin 3\theta)(\cos 3\theta) \cdot 3 + \sin(\sin 3\theta)(-\sin 3\theta) \cdot 3] = -9\cos^2(3\theta)\cos(\sin 3\theta) + 9(\sin 3\theta)\sin(\sin 3\theta)$$

54. $y = \left(1 + \sqrt{x}\right)^3 \quad \Rightarrow \quad y' = 3\left(1 + \sqrt{x}\right)^2\left(\frac{1}{2\sqrt{x}}\right) = \frac{3\left(1 + \sqrt{x}\right)^2}{2\sqrt{x}} \quad \Rightarrow$

$$y'' = \frac{2\sqrt{x} \cdot 3 \cdot 2\left(1 + \sqrt{x}\right)^1 \cdot \frac{1}{2\sqrt{x}} - 3\left(1 + \sqrt{x}\right)^2 \cdot 2 \cdot \frac{1}{2\sqrt{x}}}{\left(2\sqrt{x}\right)^2}$$

$$= \frac{6\sqrt{x}\left(1 + \sqrt{x}\right) \cdot \frac{1}{\sqrt{x}} - 3\left(1 + 2\sqrt{x} + x\right) \cdot \frac{1}{\sqrt{x}}}{4x} \cdot \frac{\sqrt{x}}{\sqrt{x}} = \frac{6\sqrt{x} + 6x - 3 - 6\sqrt{x} - 3x}{4x\sqrt{x}}$$

$$= \frac{3x - 3}{4x\sqrt{x}} \text{ or } \frac{3(x-1)}{4x^{3/2}}$$

55. $y = \sqrt{\cos x} \quad \Rightarrow \quad y' = \frac{1}{2\sqrt{\cos x}}(-\sin x) = -\frac{\sin x}{2\sqrt{\cos x}}$. With $y' = \frac{-\sin x}{2\sqrt{\cos x}}$, we get

$$y'' = \frac{2\sqrt{\cos x} \cdot (-\cos x) - (-\sin x)\left(2 \cdot \frac{1}{2\sqrt{\cos x}}(-\sin x)\right)}{\left(2\sqrt{\cos x}\right)^2} = \frac{-2\cos x\sqrt{\cos x} - \frac{\sin^2 x}{\sqrt{\cos x}}}{4\cos x} \cdot \frac{\sqrt{\cos x}}{\sqrt{\cos x}}$$

$$= \frac{-2\cos x \cdot \cos x - \sin^2 x}{4\cos x\sqrt{\cos x}} = -\frac{2\cos^2 x + \sin^2 x}{4(\cos x)^{3/2}}$$

Using the identity $\sin^2 x + \cos^2 x = 1$, the answer may be written as $-\frac{1 + \cos^2 x}{4(\cos x)^{3/2}}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

56. $y = e^{e^x} \quad \Rightarrow \quad y' = e^{e^x} \cdot (e^x)' = e^{e^x} \cdot e^x \quad \Rightarrow$

$y'' = e^{e^x} \cdot (e^x)' + e^x \cdot \left(e^{e^x}\right)' = e^{e^x} \cdot e^x + e^x \cdot e^{e^x} \cdot e^x = e^{e^x} \cdot e^x(1 + e^x) \quad \text{or} \quad e^{e^x + x}(1 + e^x)$

57. $y = 2^x \quad \Rightarrow \quad y' = 2^x \ln 2$. At $(0, 1)$, $y' = 2^0 \ln 2 = \ln 2$, and an equation of the tangent line is $y - 1 = (\ln 2)(x - 0)$ or $y = (\ln 2)x + 1$.

58. $y = \sqrt{1 + x^3} = (1 + x^3)^{1/2} \quad \Rightarrow \quad y' = \frac{1}{2}(1 + x^3)^{-1/2} \cdot 3x^2 = \dfrac{3x^2}{2\sqrt{1 + x^3}}$. At $(2, 3)$, $y' = \dfrac{3 \cdot 4}{2\sqrt{9}} = 2$, and an equation of the tangent line is $y - 3 = 2(x - 2)$, or $y = 2x - 1$.

59. $y = \sin(\sin x) \quad \Rightarrow \quad y' = \cos(\sin x) \cdot \cos x$. At $(\pi, 0)$, $y' = \cos(\sin \pi) \cdot \cos \pi = \cos(0) \cdot (-1) = 1(-1) = -1$, and an equation of the tangent line is $y - 0 = -1(x - \pi)$, or $y = -x + \pi$.

60. $y = xe^{-x^2} \quad \Rightarrow \quad y' = xe^{-x^2}(-2x) + e^{-x^2}(1) = e^{-x^2}(-2x^2 + 1)$. At $(0, 0)$, $y' = e^0(1) = 1$, and an equation of the tangent line is $y - 0 = 1(x - 0)$ or $y = x$.

61. (a) $y = \dfrac{2}{1 + e^{-x}} \quad \Rightarrow \quad y' = \dfrac{(1 + e^{-x})(0) - 2(-e^{-x})}{(1 + e^{-x})^2} = \dfrac{2e^{-x}}{(1 + e^{-x})^2}$.

At $(0, 1)$, $y' = \dfrac{2e^0}{(1 + e^0)^2} = \dfrac{2(1)}{(1 + 1)^2} = \dfrac{2}{2^2} = \dfrac{1}{2}$. So an equation of the tangent line is $y - 1 = \frac{1}{2}(x - 0)$ or $y = \frac{1}{2}x + 1$.

(b)

62. (a) For $x > 0$, $|x| = x$, and $y = f(x) = \dfrac{x}{\sqrt{2 - x^2}} \quad \Rightarrow$

$$f'(x) = \frac{\sqrt{2 - x^2}\,(1) - x\left(\frac{1}{2}\right)(2 - x^2)^{-1/2}(-2x)}{\left(\sqrt{2 - x^2}\right)^2} \cdot \frac{(2 - x^2)^{1/2}}{(2 - x^2)^{1/2}}$$

$$= \frac{(2 - x^2) + x^2}{(2 - x^2)^{3/2}} = \frac{2}{(2 - x^2)^{3/2}}$$

So at $(1, 1)$, the slope of the tangent line is $f'(1) = 2$ and its equation is $y - 1 = 2(x - 1)$ or $y = 2x - 1$.

(b)

63. (a) $f(x) = x\sqrt{2 - x^2} = x(2 - x^2)^{1/2} \quad \Rightarrow$

$$f'(x) = x \cdot \tfrac{1}{2}(2 - x^2)^{-1/2}(-2x) + (2 - x^2)^{1/2} \cdot 1 = (2 - x^2)^{-1/2}\left[-x^2 + (2 - x^2)\right] = \frac{2 - 2x^2}{\sqrt{2 - x^2}}$$

(b)

$f' = 0$ when f has a horizontal tangent line, f' is negative when f is decreasing, and f' is positive when f is increasing.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. (a)

From the graph of f, we see that there are 5 horizontal tangents, so there must be 5 zeros on the graph of f'. From the symmetry of the graph of f, we must have the graph of f' as high at $x = 0$ as it is low at $x = \pi$. The intervals of increase and decrease as well as the signs of f' are indicated in the figure.

(b) $f(x) = \sin(x + \sin 2x) \quad \Rightarrow$

$$f'(x) = \cos(x+\sin 2x) \cdot \frac{d}{dx}(x+\sin 2x) = \cos(x+\sin 2x)(1+2\cos 2x)$$

65. For the tangent line to be horizontal, $f'(x) = 0$. $f(x) = 2\sin x + \sin^2 x \quad \Rightarrow \quad f'(x) = 2\cos x + 2\sin x\,\cos x = 0 \quad \Leftrightarrow$ $2\cos x(1+\sin x) = 0 \quad \Leftrightarrow \quad \cos x = 0$ or $\sin x = -1$, so $x = \frac{\pi}{2} + 2n\pi$ or $\frac{3\pi}{2} + 2n\pi$, where n is any integer. Now $f\left(\frac{\pi}{2}\right) = 3$ and $f\left(\frac{3\pi}{2}\right) = -1$, so the points on the curve with a horizontal tangent are $\left(\frac{\pi}{2} + 2n\pi, 3\right)$ and $\left(\frac{3\pi}{2} + 2n\pi, -1\right)$, where n is any integer.

66. $y = \sqrt{1+2x} \quad \Rightarrow \quad y' = \frac{1}{2}(1+2x)^{-1/2} \cdot 2 = \dfrac{1}{\sqrt{1+2x}}$. The line $6x + 2y = 1$ $\left(\text{or } y = -3x + \frac{1}{2}\right)$ has slope -3, so the tangent line perpendicular to it must have slope $\frac{1}{3}$. Thus, $\dfrac{1}{3} = \dfrac{1}{\sqrt{1+2x}} \quad \Leftrightarrow \quad \sqrt{1+2x} = 3 \quad \Rightarrow \quad 1 + 2x = 9 \quad \Leftrightarrow$ $2x = 8 \quad \Leftrightarrow \quad x = 4$. When $x = 4$, $y = \sqrt{1+2(4)} = 3$, so the point is $(4, 3)$.

67. $F(x) = f(g(x)) \quad \Rightarrow \quad F'(x) = f'(g(x)) \cdot g'(x)$, so $F'(5) = f'(g(5)) \cdot g'(5) = f'(-2) \cdot 6 = 4 \cdot 6 = 24$.

68. $h(x) = \sqrt{4 + 3f(x)} \quad \Rightarrow \quad h'(x) = \frac{1}{2}(4 + 3f(x))^{-1/2} \cdot 3f'(x)$, so
$h'(1) = \frac{1}{2}(4 + 3f(1))^{-1/2} \cdot 3f'(1) = \frac{1}{2}(4 + 3 \cdot 7)^{-1/2} \cdot 3 \cdot 4 = \frac{6}{\sqrt{25}} = \frac{6}{5}$.

69. (a) $h(x) = f(g(x)) \quad \Rightarrow \quad h'(x) = f'(g(x)) \cdot g'(x)$, so $h'(1) = f'(g(1)) \cdot g'(1) = f'(2) \cdot 6 = 5 \cdot 6 = 30$.

(b) $H(x) = g(f(x)) \quad \Rightarrow \quad H'(x) = g'(f(x)) \cdot f'(x)$, so $H'(1) = g'(f(1)) \cdot f'(1) = g'(3) \cdot 4 = 9 \cdot 4 = 36$.

70. (a) $F(x) = f(f(x)) \quad \Rightarrow \quad F'(x) = f'(f(x)) \cdot f'(x)$, so $F'(2) = f'(f(2)) \cdot f'(2) = f'(1) \cdot 5 = 4 \cdot 5 = 20$.

(b) $G(x) = g(g(x)) \quad \Rightarrow \quad G'(x) = g'(g(x)) \cdot g'(x)$, so $G'(3) = g'(g(3)) \cdot g'(3) = g'(2) \cdot 9 = 7 \cdot 9 = 63$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

71. (a) From the graphs of f and g, we obtain the following values: $g(1) = 4$ since the point $(1, 4)$ is on the graph of g; $f'(4) = -\frac{1}{4}$ since the slope of the line segment between $(2, 4)$ and $(6, 3)$ is $\dfrac{3-4}{6-2} = -\dfrac{1}{4}$; and $g'(1) = -1$ since the slope of the line segment between $(0, 5)$ and $(3, 2)$ is $\dfrac{2-5}{3-0} = -1$. Now $u(x) = f(g(x))$, so $u'(1) = f'(g(1))\,g'(1) = f'(4)\,g'(1) = -\frac{1}{4}(-1) = \frac{1}{4}$.

(b) From the graphs of f and g, we obtain the following values: $f(1) = 2$ since the point $(1, 2)$ is on the graph of f; $g'(2) = g'(1) = -1$ [see part (a)]; and $f'(1) = 2$ since the slope of the line segment between $(0, 0)$ and $(2, 4)$ is $\dfrac{4-0}{2-0} = 2$. Now $v(x) = g(f(x))$, so $v'(1) = g'(f(1))f'(1) = g'(2)f'(1) = -1(2) = -2$.

(c) From part (a), we have $g(1) = 4$ and $g'(1) = -1$. From the graph of g we obtain $g'(4) = \frac{1}{2}$ since the slope of the line segment between $(3, 2)$ and $(7, 4)$ is $\dfrac{4-2}{7-3} = \dfrac{1}{2}$. Now $w(x) = g(g(x))$, so $w'(1) = g'(g(1))\,g'(1) = g'(4)\,g'(1) = \frac{1}{2}(-1) = -\frac{1}{2}$.

72. (a) $h(x) = f(f(x)) \;\Rightarrow\; h'(x) = f'(f(x))f'(x)$. So $h'(2) = f'(f(2))f'(2) = f'(1)f'(2) \approx (-1)(-1) = 1$.

(b) $g(x) = f(x^2) \;\Rightarrow\; g'(x) = f'(x^2) \cdot \dfrac{d}{dx}\left(x^2\right) = f'(x^2)(2x)$. So $g'(2) = f'(2^2)(2 \cdot 2) = 4f'(4) \approx 4(2) = 8$.

73. The point $(3, 2)$ is on the graph of f, so $f(3) = 2$. The tangent line at $(3, 2)$ has slope $\dfrac{\Delta y}{\Delta x} = \dfrac{-4}{6} = -\dfrac{2}{3}$.

$g(x) = \sqrt{f(x)} \;\Rightarrow\; g'(x) = \frac{1}{2}[f(x)]^{-1/2} \cdot f'(x) \;\Rightarrow$

$g'(3) = \frac{1}{2}[f(3)]^{-1/2} \cdot f'(3) = \frac{1}{2}(2)^{-1/2}(-\frac{2}{3}) = -\dfrac{1}{3\sqrt{2}}$ or $-\dfrac{1}{6}\sqrt{2}$.

74. (a) $F(x) = f(x^\alpha) \;\Rightarrow\; F'(x) = f'(x^\alpha)\dfrac{d}{dx}\left(x^\alpha\right) = f'(x^\alpha)\alpha x^{\alpha-1}$

(b) $G(x) = [f(x)]^\alpha \;\Rightarrow\; G'(x) = \alpha\,[f(x)]^{\alpha-1}\,f'(x)$

75. (a) $F(x) = f(e^x) \;\Rightarrow\; F'(x) = f'(e^x)\dfrac{d}{dx}\left(e^x\right) = f'(e^x)e^x$

(b) $G(x) = e^{f(x)} \;\Rightarrow\; G'(x) = e^{f(x)}\dfrac{d}{dx}\,f(x) = e^{f(x)}f'(x)$

76. (a) $g(x) = e^{cx} + f(x) \;\Rightarrow\; g'(x) = e^{cx} \cdot c + f'(x) \;\Rightarrow\; g'(0) = e^0 \cdot c + f'(0) = c + 5$.

$g'(x) = ce^{cx} + f'(x) \;\Rightarrow\; g''(x) = ce^{cx} \cdot c + f''(x) \;\Rightarrow\; g''(0) = c^2e^0 + f''(0) = c^2 - 2$.

(b) $h(x) = e^{kx}f(x) \;\Rightarrow\; h'(x) = e^{kx}f'(x) + f(x) \cdot ke^{kx} \;\Rightarrow\; h'(0) = e^0f'(0) + f(0) \cdot ke^0 = 5 + 3k$.

An equation of the tangent line to the graph of h at the point $(0, h(0)) = (0, f(0)) = (0, 3)$ is $y - 3 = (5 + 3k)(x - 0)$ or $y = (5 + 3k)x + 3$.

77. $r(x) = f(g(h(x))) \;\Rightarrow\; r'(x) = f'(g(h(x))) \cdot g'(h(x)) \cdot h'(x)$, so

$r'(1) = f'(g(h(1))) \cdot g'(h(1)) \cdot h'(1) = f'(g(2)) \cdot g'(2) \cdot 4 = f'(3) \cdot 5 \cdot 4 = 6 \cdot 5 \cdot 4 = 120$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

78. $f(x) = xg(x^2) \quad \Rightarrow \quad f'(x) = xg'(x^2)\,2x + g(x^2)\cdot 1 = 2x^2g'(x^2) + g(x^2) \quad \Rightarrow$

$f''(x) = 2x^2g''(x^2)\,2x + g'(x^2)\,4x + g'(x^2)\,2x = 4x^3g''(x^2) + 4xg'(x^2) + 2xg'(x^2) = 6xg'(x^2) + 4x^3g''(x^2)$

79. $F(x) = f(3f(4f(x))) \quad \Rightarrow$

$$F'(x) = f'(3f(4f(x)))\cdot\frac{d}{dx}(3f(4f(x))) = f'(3f(4f(x)))\cdot 3f'(4f(x))\cdot\frac{d}{dx}(4f(x))$$
$$= f'(3f(4f(x)))\cdot 3f'(4f(x))\cdot 4f'(x), \quad \text{so}$$

$F'(0) = f'(3f(4f(0)))\cdot 3f'(4f(0))\cdot 4f'(0) = f'(3f(4\cdot 0))\cdot 3f'(4\cdot 0)\cdot 4\cdot 2 = f'(3\cdot 0)\cdot 3\cdot 2\cdot 4\cdot 2 = 2\cdot 3\cdot 2\cdot 4\cdot 2 = 96.$

80. $F(x) = f(xf(xf(x))) \quad \Rightarrow$

$$F'(x) = f'(xf(xf(x)))\cdot\frac{d}{dx}(xf(xf(x))) = f'(xf(xf(x)))\cdot\left[x\cdot f'(xf(x))\cdot\frac{d}{dx}(xf(x)) + f(xf(x))\cdot 1\right]$$
$$= f'(xf(xf(x)))\cdot[xf'(xf(x))\cdot(xf'(x) + f(x)\cdot 1) + f(xf(x))], \quad \text{so}$$

$$F'(1) = f'(f(f(1)))\cdot[f'(f(1))\cdot(f'(1) + f(1)) + f(f(1))] = f'(f(2))\cdot[f'(2)\cdot(4+2) + f(2)]$$
$$= f'(3)\cdot[5\cdot 6 + 3] = 6\cdot 33 = 198.$$

81. $y = e^{2x}(A\cos 3x + B\sin 3x) \quad \Rightarrow$

$$y' = e^{2x}(-3A\sin 3x + 3B\cos 3x) + (A\cos 3x + B\sin 3x)\cdot 2e^{2x}$$
$$= e^{2x}(-3A\sin 3x + 3B\cos 3x + 2A\cos 3x + 2B\sin 3x)$$
$$= e^{2x}[(2A+3B)\cos 3x + (2B-3A)\sin 3x] \quad \Rightarrow$$

$$y'' = e^{2x}[-3(2A+3B)\sin 3x + 3(2B-3A)\cos 3x] + [(2A+3B)\cos 3x + (2B-3A)\sin 3x]\cdot 2e^{2x}$$
$$= e^{2x}\{[-3(2A+3B) + 2(2B-3A)]\sin 3x + [3(2B-3A) + 2(2A+3B)]\cos 3x\}$$
$$= e^{2x}[(-12A-5B)\sin 3x + (-5A+12B)\cos 3x]$$

Substitute the expressions for y, y', and y'' in $y'' - 4y' + 13y$ to get

$$y'' - 4y' + 13y = e^{2x}[(-12A-5B)\sin 3x + (-5A+12B)\cos 3x]$$
$$- 4e^{2x}[(2A+3B)\cos 3x + (2B-3A)\sin 3x] + 13e^{2x}(A\cos 3x + B\sin 3x)$$
$$= e^{2x}[(-12A-5B-8B+12A+13B)\sin 3x + (-5A+12B-8A-12B+13A)\cos 3x]$$
$$= e^{2x}[(0)\sin 3x + (0)\cos 3x] = 0$$

Thus, the function y satisfies the differential equation $y'' - 4y' + 13y = 0$.

82. $y = e^{rx} \quad \Rightarrow \quad y' = re^{rx} \quad \Rightarrow \quad y'' = r^2e^{rx}$. Substituting y, y', and y'' into $y'' - 4y' + y = 0$ gives us

$r^2e^{rx} - 4re^{rx} + e^{rx} = 0 \quad \Rightarrow \quad e^{rx}(r^2 - 4r + 1) = 0$. Since $e^{rx} \neq 0$, we must have

$$r^2 - 4r + 1 = 0 \quad \Rightarrow \quad r = \frac{4 \pm \sqrt{16-4}}{2} = 2 \pm \sqrt{3}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

83. The use of D, D^2, …, D^n is just a derivative notation (see text page 159). In general, $Df(2x) = 2f'(2x)$, $D^2 f(2x) = 4f''(2x)$, …, $D^n f(2x) = 2^n f^{(n)}(2x)$. Since $f(x) = \cos x$ and $50 = 4(12) + 2$, we have $f^{(50)}(x) = f^{(2)}(x) = -\cos x$, so $D^{50} \cos 2x = -2^{50} \cos 2x$.

84. $f(x) = xe^{-x}$, $f'(x) = e^{-x} - xe^{-x} = (1-x)e^{-x}$, $f''(x) = -e^{-x} + (1-x)(-e^{-x}) = (x-2)e^{-x}$. Similarly, $f'''(x) = (3-x)e^{-x}$, $f^{(4)}(x) = (x-4)e^{-x}$, …, $f^{(1000)}(x) = (x-1000)e^{-x}$.

85. $s(t) = 10 + \frac{1}{4}\sin(10\pi t) \quad \Rightarrow$ the velocity after t seconds is $v(t) = s'(t) = \frac{1}{4}\cos(10\pi t)(10\pi) = \frac{5\pi}{2}\cos(10\pi t)$ cm/s.

86. (a) $s = A\cos(\omega t + \delta) \quad \Rightarrow \quad$ velocity $= s' = -\omega A \sin(\omega t + \delta)$.

(b) If $A \neq 0$ and $\omega \neq 0$, then $s' = 0 \quad \Leftrightarrow \quad \sin(\omega t + \delta) = 0 \quad \Leftrightarrow \quad \omega t + \delta = n\pi \quad \Leftrightarrow \quad t = \dfrac{n\pi - \delta}{\omega}$, n an integer.

87. (a) $B(t) = 4.0 + 0.35\sin\left(\dfrac{2\pi t}{5.4}\right) \quad \Rightarrow \quad \dfrac{dB}{dt} = \left(0.35\cos\dfrac{2\pi t}{5.4}\right)\left(\dfrac{2\pi}{5.4}\right) = \dfrac{0.7\pi}{5.4}\cos\dfrac{2\pi t}{5.4} = \dfrac{7\pi}{54}\cos\dfrac{2\pi t}{5.4}$

(b) At $t = 1$, $\dfrac{dB}{dt} = \dfrac{7\pi}{54}\cos\dfrac{2\pi}{5.4} \approx 0.16$.

88. $L(t) = 12 + 2.8\sin\left[\frac{2\pi}{365}(t-80)\right] \quad \Rightarrow \quad L'(t) = 2.8\cos\left[\frac{2\pi}{365}(t-80)\right]\left(\frac{2\pi}{365}\right)$.

On March 21, $t = 80$, and $L'(80) \approx 0.0482$ hours per day. On May 21, $t = 141$, and $L'(141) \approx 0.02398$, which is approximately one-half of $L'(80)$.

89. $s(t) = 2e^{-1.5t}\sin 2\pi t \quad \Rightarrow$

$v(t) = s'(t) = 2[e^{-1.5t}(\cos 2\pi t)(2\pi) + (\sin 2\pi t)e^{-1.5t}(-1.5)] = 2e^{-1.5t}(2\pi\cos 2\pi t - 1.5\sin 2\pi t)$

Graph of position

Graph of velocity

90. (a) $\lim\limits_{t\to\infty} p(t) = \lim\limits_{t\to\infty} \dfrac{1}{1 + ae^{-kt}} = \dfrac{1}{1 + a \cdot 0} = 1$, since $k > 0 \quad \Rightarrow \quad -kt \to -\infty \quad \Rightarrow \quad e^{-kt} \to 0$. As time increases, the proportion of the population that has heard the rumor approaches 1; that is, everyone in the population has heard the rumor.

(b) $p(t) = (1 + ae^{-kt})^{-1} \quad \Rightarrow \quad \dfrac{dp}{dt} = -(1 + ae^{-kt})^{-2}(-kae^{-kt}) = \dfrac{kae^{-kt}}{(1 + ae^{-kt})^2}$

(c)

From the graph of $p(t) = (1 + 10e^{-0.5t})^{-1}$, it seems that $p(t) = 0.8$ (indicating that 80% of the population has heard the rumor) when $t \approx 7.4$ hours.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

91. (a) Use $C(t) = ate^{bt}$ with $a = 0.00225$ and $b = -0.0467$ to get $C'(t) = a(t \cdot e^{bt} \cdot b + e^{bt} \cdot 1) = a(bt+1)e^{bt}$.

$C'(10) = 0.00225(0.533)e^{-0.467} \approx 0.00075$, so the BAC was increasing at approximately 0.00075 (g/dL)/min after 10 minutes.

(b) A half an hour later gives us $t = 10 + 30 = 40$. $C'(40) = 0.00225(-0.868)e^{-1.868} \approx -0.00030$, so the BAC was decreasing at approximately 0.00030 (g/dL)/min after 40 minutes.

92. (a) The derivative dV/dr represents the rate of change of the volume with respect to the radius and the derivative dV/dt represents the rate of change of the volume with respect to time.

(b) Since $V = \dfrac{4}{3}\pi r^3$, $\dfrac{dV}{dt} = \dfrac{dV}{dr}\dfrac{dr}{dt} = 4\pi r^2 \dfrac{dr}{dt}$.

93. By the Chain Rule, $a(t) = \dfrac{dv}{dt} = \dfrac{dv}{ds}\dfrac{ds}{dt} = \dfrac{dv}{ds}\,v(t) = v(t)\,\dfrac{dv}{ds}$. The derivative dv/dt is the rate of change of the velocity with respect to time (in other words, the acceleration) whereas the derivative dv/ds is the rate of change of the velocity with respect to the displacement.

94. (a) $P = ab^t$ with $a = 4.502714 \times 10^{-20}$ and $b = 1.029953851$, where P is measured in thousands of people. The fit appears to be very good.

(b) **For 1800:** $m_1 = \dfrac{5308 - 3929}{1800 - 1790} = 137.9$, $m_2 = \dfrac{7240 - 5308}{1810 - 1800} = 193.2$.

So $P'(1800) \approx (m_1 + m_2)/2 = 165.55$ thousand people/year.

For 1850: $m_1 = \dfrac{23{,}192 - 17{,}063}{1850 - 1840} = 612.9$, $m_2 = \dfrac{31{,}443 - 23{,}192}{1860 - 1850} = 825.1$.

So $P'(1850) \approx (m_1 + m_2)/2 = 719$ thousand people/year.

(c) Using $P'(t) = ab^t \ln b$ (from Formula 5) with the values of a and b from part (a), we get $P'(1800) \approx 156.85$ and $P'(1850) \approx 686.07$. These estimates are somewhat less than the ones in part (b).

(d) $P(1870) \approx 41{,}946.56$. The difference of 3.4 million people is most likely due to the Civil War (1861–1865).

95. (a) If f is even, then $f(x) = f(-x)$. Using the Chain Rule to differentiate this equation, we get

$f'(x) = f'(-x)\dfrac{d}{dx}(-x) = -f'(-x)$. Thus, $f'(-x) = -f'(x)$, so f' is odd.

(b) If f is odd, then $f(x) = -f(-x)$. Differentiating this equation, we get $f'(x) = -f'(-x)(-1) = f'(-x)$, so f' is even.

96.
$$\left[\frac{f(x)}{g(x)}\right]' = \left\{f(x)\,[g(x)]^{-1}\right\}' = f'(x)\,[g(x)]^{-1} + (-1)\,[g(x)]^{-2}\,g'(x)f(x)$$
$$= \frac{f'(x)}{g(x)} - \frac{f(x)g'(x)}{[g(x)]^2} = \frac{g(x)f'(x) - f(x)g'(x)}{[g(x)]^2}$$

This is an alternative derivation of the *formula* in the Quotient Rule. But part of the purpose of the Quotient Rule is to show that if f and g are differentiable, so is f/g. The proof in Section 3.2 does that; this one doesn't.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

97. Since $\theta^\circ = \left(\frac{\pi}{180}\right)\theta$ rad, we have $\dfrac{d}{d\theta}\left(\sin\theta^\circ\right) = \dfrac{d}{d\theta}\left(\sin\frac{\pi}{180}\theta\right) = \frac{\pi}{180}\cos\frac{\pi}{180}\theta = \frac{\pi}{180}\cos\theta^\circ$.

98. (a) $f(x) = |x| = \sqrt{x^2} = (x^2)^{1/2} \quad\Rightarrow\quad f'(x) = \frac{1}{2}(x^2)^{-1/2}(2x) = x/\sqrt{x^2} = x/|x|$ for $x \neq 0$.

f is not differentiable at $x = 0$.

(b) $f(x) = |\sin x| = \sqrt{\sin^2 x} \quad\Rightarrow$

$$f'(x) = \tfrac{1}{2}(\sin^2 x)^{-1/2}\,2\sin x\,\cos x = \frac{\sin x}{|\sin x|}\cos x$$
$$= \begin{cases} \cos x & \text{if } \sin x > 0 \\ -\cos x & \text{if } \sin x < 0 \end{cases}$$

f is not differentiable when $x = n\pi$, n an integer.

(c) $g(x) = \sin|x| = \sin\sqrt{x^2} \quad\Rightarrow$

$$g'(x) = \cos|x| \cdot \frac{x}{|x|} = \frac{x}{|x|}\cos x = \begin{cases} \cos x & \text{if } x > 0 \\ -\cos x & \text{if } x < 0 \end{cases}$$

g is not differentiable at 0.

99. $y = b^x \quad\Rightarrow\quad y' = b^x \ln b$, so the slope of the tangent line to the curve $y = b^x$ at the point (a, b^a) is $b^a \ln b$. An equation of this tangent line is then $y - b^a = b^a \ln b\,(x - a)$. If c is the x-intercept of this tangent line, then $0 - b^a = b^a \ln b\,(c - a) \quad\Rightarrow$ $-1 = \ln b\,(c - a) \quad\Rightarrow\quad \dfrac{-1}{\ln b} = c - a \quad\Rightarrow\quad |c - a| = \left|\dfrac{-1}{\ln b}\right| = \dfrac{1}{|\ln b|}$. The distance between $(a, 0)$ and $(c, 0)$ is $|c - a|$, and this distance is the constant $\dfrac{1}{|\ln b|}$ for any a. [*Note*: The absolute value is needed for the case $0 < b < 1$ because $\ln b$ is negative there. If $b > 1$, we can write $a - c = 1/(\ln b)$ as the constant distance between $(a, 0)$ and $(c, 0)$.]

100. $y = b^x \quad\Rightarrow\quad y' = b^x \ln b$, so the slope of the tangent line to the curve $y = b^x$ at the point (x_0, y_0) is $b^{x_0} \ln b$. An equation of this tangent line is then $y - y_0 = b^{x_0} \ln b\,(x - x_0)$. Since this tangent line must pass through $(0, 0)$, we have $0 - y_0 = b^{x_0} \ln b\,(0 - x_0)$, or $y_0 = b^{x_0}\,(\ln b)\,x_0$. Since (x_0, y_0) is a point on the exponential curve $y = b^x$, we also have $y_0 = b^{x_0}$. Equating the expressions for y_0 gives $b^{x_0} = b^{x_0}\,(\ln b)\,x_0 \quad\Rightarrow\quad 1 = (\ln b)\,x_0 \quad\Rightarrow\quad x_0 = 1/(\ln b)$.

So $y_0 = b^{x_0} = e^{x_0 \ln b}$ [by Formula 1.5.10] $= e^{(1/(\ln b))\ln b} = e^1 = e$.

101. Let $j(x) = g(h(x))$ so that $F(x) = f(g(h(x))) = f(j(x))$. By the Chain Rule, we have $j'(x) = g'(h(x)) \cdot h'(x)$ and, by the Chain Rule and substitution, we have $F'(x) = f'(j(x)) \cdot j'(x) = f'(g(h(x))) \cdot g'(h(x)) \cdot h'(x)$.

102. $F(x) = f(g(x)) \quad\Rightarrow\quad F'(x) = f'(g(x)) \cdot g'(x)$ by the Chain Rule. By the Product Rule and Chain Rule we have

$$F''(x) = f'(g(x)) \cdot g''(x) + g'(x) \cdot \frac{d}{dx} f'(g(x)) = f'(g(x)) \cdot g''(x) + g'(x) \cdot f''(g(x))\,g'(x)$$
$$= f''(g(x))[g'(x)]^2 + f'(g(x)) \cdot g''(x)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

APPLIED PROJECT Where Should a Pilot Start Descent?

1. Condition (i) will hold if and only if all of the following four conditions hold:

(α) $P(0) = 0$

(β) $P'(0) = 0$ (for a smooth landing)

(γ) $P'(\ell) = 0$ (since the plane is cruising horizontally when it begins its descent)

(δ) $P(\ell) = h$.

First of all, condition α implies that $P(0) = d = 0$, so $P(x) = ax^3 + bx^2 + cx \quad \Rightarrow \quad P'(x) = 3ax^2 + 2bx + c$. But $P'(0) = c = 0$ by condition β. So $P'(\ell) = 3a\ell^2 + 2b\ell = \ell\,(3a\ell + 2b)$. Now by condition γ, $3a\ell + 2b = 0 \quad \Rightarrow \quad a = -\dfrac{2b}{3\ell}$. Therefore, $P(x) = -\dfrac{2b}{3\ell}x^3 + bx^2$. Setting $P(\ell) = h$ for condition δ, we get $P(\ell) = -\dfrac{2b}{3\ell}\ell^3 + b\ell^2 = h \quad \Rightarrow$ $-\dfrac{2}{3}b\ell^2 + b\ell^2 = h \quad \Rightarrow \quad \dfrac{1}{3}b\ell^2 = h \quad \Rightarrow \quad b = \dfrac{3h}{\ell^2} \quad \Rightarrow \quad a = -\dfrac{2h}{\ell^3}$. So $y = P(x) = -\dfrac{2h}{\ell^3}x^3 + \dfrac{3h}{\ell^2}x^2$.

2. By condition (ii), $\dfrac{dx}{dt} = -v$ for all t, so $x\,(t) = \ell - vt$. Condition (iii) states that $\left|\dfrac{d^2y}{dt^2}\right| \le k$. By the Chain Rule, we have $\dfrac{dy}{dt} = \dfrac{dy}{dx}\dfrac{dx}{dt} = -\dfrac{2h}{\ell^3}\left(3x^2\right)\dfrac{dx}{dt} + \dfrac{3h}{\ell^2}\,(2x)\,\dfrac{dx}{dt} = \dfrac{6hx^2v}{\ell^3} - \dfrac{6hxv}{\ell^2}$ (for $x \le \ell$) $\Rightarrow$ $\dfrac{d^2y}{dt^2} = \dfrac{6hv}{\ell^3}\,(2x)\,\dfrac{dx}{dt} - \dfrac{6hv}{\ell^2}\dfrac{dx}{dt} = -\dfrac{12hv^2}{\ell^3}x + \dfrac{6hv^2}{\ell^2}$. In particular, when $t = 0$, $x = \ell$ and so $\left.\dfrac{d^2y}{dt^2}\right|_{t=0} = -\dfrac{12hv^2}{\ell^3}\ell + \dfrac{6hv^2}{\ell^2} = -\dfrac{6hv^2}{\ell^2}$. Thus, $\left|\left.\dfrac{d^2y}{dt^2}\right|_{t=0}\right| = \dfrac{6hv^2}{\ell^2} \le k$. (This condition also follows from taking $x = 0$.)

3. We substitute $k = 860$ mi/h^2, $h = 35{,}000\text{ ft} \times \dfrac{1\text{ mi}}{5280\text{ ft}}$, and $v = 300$ mi/h into the result of part (b):

$$\frac{6\left(35{,}000 \cdot \frac{1}{5280}\right)(300)^2}{\ell^2} \le 860 \quad \Rightarrow \quad \ell \ge 300\sqrt{6 \cdot \frac{35{,}000}{5280 \cdot 860}} \approx 64.5 \text{ miles.}$$

4. Substituting the values of h and ℓ in Problem 3 into $P(x) = -\dfrac{2h}{\ell^3}x^3 + \dfrac{3h}{\ell^2}x^2$ gives us $P(x) = ax^3 + bx^2$, where $a \approx -4.937 \times 10^{-5}$ and $b \approx 4.78 \times 10^{-3}$.

3.5 Implicit Differentiation

1. (a) $\dfrac{d}{dx}\left(5x^2 - y^3\right) = \dfrac{d}{dx}(7) \quad \Rightarrow \quad 10x - 3y^2\,y' = 0 \quad \Rightarrow \quad 3y^2\,y' = 10x \quad \Rightarrow \quad y' = \dfrac{10x}{3y^2}$

(b) $5x^2 - y^3 = 7 \quad \Rightarrow \quad y^3 = 5x^2 - 7 \quad \Rightarrow \quad y = \sqrt[3]{5x^2 - 7}$, so $y' = \frac{1}{3}(5x^2 - 7)^{-2/3}(10x) = \dfrac{10x}{3(5x^2 - 7)^{2/3}}$

(c) From part (a), $y' = \dfrac{10x}{3y^2} = \dfrac{10x}{3(y^3)^{2/3}} = \dfrac{10x}{3(5x^2 - 7)^{2/3}}$, which agrees with part (b).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. (a) $\dfrac{d}{dx}\left(6x^4+y^5\right)=\dfrac{d}{dx}(2) \quad\Rightarrow\quad 24x^3+5y^4y'=2 \quad\Rightarrow\quad 5y^4\,y'=2-24x^3 \quad\Rightarrow\quad y'=\dfrac{2-24x^3}{5y^4}$

(b) $6x^4+y^5=2x \quad\Rightarrow\quad y^5=2x-6x^4 \quad\Rightarrow\quad y=\sqrt[5]{2x-6x^4}$, so $y'=\frac{1}{5}(2x-6x^4)^{-4/5}(2-24x^3)=\dfrac{2-24x^3}{5(2x-6x^4)^{4/5}}$

(c) From part (a), $y'=\dfrac{2-24x^3}{5y^4}=\dfrac{2-24x^2}{5(y^5)^{4/5}}=\dfrac{2-24x^3}{5(2x-6x^4)^{4/5}}$, which agrees with part (b).

3. (a) $\dfrac{d}{dx}\left(\sqrt{x}+\sqrt{y}\right)=\dfrac{d}{dx}(1) \quad\Rightarrow\quad \dfrac{1}{2}x^{-1/2}+\dfrac{1}{2}y^{-1/2}y'=0 \quad\Rightarrow\quad \dfrac{1}{2\sqrt{y}}\,y'=-\dfrac{1}{2\sqrt{x}} \quad\Rightarrow\quad y'=-\dfrac{\sqrt{y}}{\sqrt{x}}$

(b) $\sqrt{x}+\sqrt{y}=1 \quad\Rightarrow\quad \sqrt{y}=1-\sqrt{x} \quad\Rightarrow\quad y=\left(1-\sqrt{x}\,\right)^2 \quad\Rightarrow\quad y=1-2\sqrt{x}+x$, so

$y'=-2\cdot\dfrac{1}{2}x^{-1/2}+1=1-\dfrac{1}{\sqrt{x}}$.

(c) From part (a), $y'=-\dfrac{\sqrt{y}}{\sqrt{x}}=-\dfrac{1-\sqrt{x}}{\sqrt{x}}$ [from part (b)] $=-\dfrac{1}{\sqrt{x}}+1$, which agrees with part (b).

4. (a) $\dfrac{d}{dx}\left(\dfrac{2}{x}-\dfrac{1}{y}\right)=\dfrac{d}{dx}(4) \quad\Rightarrow\quad -2x^{-2}+y^{-2}y'=0 \quad\Rightarrow\quad \dfrac{1}{y^2}\,y'=\dfrac{2}{x^2} \quad\Rightarrow\quad y'=\dfrac{2y^2}{x^2}$

(b) $\dfrac{2}{x}-\dfrac{1}{y}=4 \quad\Rightarrow\quad \dfrac{1}{y}=\dfrac{2}{x}-4 \quad\Rightarrow\quad \dfrac{1}{y}=\dfrac{2-4x}{x} \quad\Rightarrow\quad y=\dfrac{x}{2-4x}$, so

$y'=\dfrac{(2-4x)(1)-x(-4)}{(2-4x)^2}=\dfrac{2}{(2-4x)^2}\ \left[\text{or}\ \dfrac{1}{2(1-2x)^2}\right]$.

(c) From part (a), $y'=\dfrac{2y^2}{x^2}=\dfrac{2\left(\dfrac{x}{2-4x}\right)^2}{x^2}$ [from part (b)] $=\dfrac{2x^2}{x^2(2-4x)^2}=\dfrac{2}{(2-4x)^2}$, which agrees with part (b).

5. $\dfrac{d}{dx}\left(x^2-4xy+y^2\right)=\dfrac{d}{dx}(4) \quad\Rightarrow\quad 2x-4[xy'+y(1)]+2y\,y'=0 \quad\Rightarrow\quad 2y\,y'-4xy'=4y-2x \quad\Rightarrow$

$y'(y-2x)=2y-x \quad\Rightarrow\quad y'=\dfrac{2y-x}{y-2x}$

6. $\dfrac{d}{dx}\left(2x^2+xy-y^2\right)=\dfrac{d}{dx}(2) \quad\Rightarrow\quad 4x+xy'+y(1)-2y\,y'=0 \quad\Rightarrow\quad xy'-2y\,y'=-4x-y \quad\Rightarrow$

$(x-2y)y'=-4x-y \quad\Rightarrow\quad y'=\dfrac{-4x-y}{x-2y}$

7. $\dfrac{d}{dx}\left(x^4+x^2y^2+y^3\right)=\dfrac{d}{dx}(5) \quad\Rightarrow\quad 4x^3+x^2\cdot 2y\,y'+y^2\cdot 2x+3y^2y'=0 \quad\Rightarrow\quad 2x^2y\,y'+3y^2y'=-4x^3-2xy^2 \quad\Rightarrow$

$(2x^2y+3y^2)y'=-4x^3-2xy^2 \quad\Rightarrow\quad y'=\dfrac{-4x^3-2xy^2}{2x^2y+3y^2}=-\dfrac{2x(2x^2+y^2)}{y(2x^2+3y)}$

8. $\dfrac{d}{dx}\left(x^3-xy^2+y^3\right)=\dfrac{d}{dx}(1) \quad\Rightarrow\quad 3x^2-x\cdot 2y\,y'-y^2\cdot 1+3y^2y'=0 \quad\Rightarrow\quad 3y^2y'-2x\,y\,y'=y^2-3x^2 \quad\Rightarrow$

$(3y^2-2xy)\,y'=y^2-3x^2 \quad\Rightarrow\quad y'=\dfrac{y^2-3x^2}{3y^2-2xy}=\dfrac{y^2-3x^2}{y(3y-2x)}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $\dfrac{d}{dx}\left(\dfrac{x^2}{x+y}\right) = \dfrac{d}{dx}(y^2+1) \quad \Rightarrow \quad \dfrac{(x+y)(2x) - x^2(1+y')}{(x+y)^2} = 2y\,y' \quad \Rightarrow$

$2x^2 + 2xy - x^2 - x^2\,y' = 2y(x+y)^2\,y' \quad \Rightarrow \quad x^2 + 2xy = 2y(x+y)^2\,y' + x^2\,y' \quad \Rightarrow$

$x(x+2y) = [2y(x^2+2xy+y^2) + x^2]\,y' \quad \Rightarrow \quad y' = \dfrac{x(x+2y)}{2x^2y + 4xy^2 + 2y^3 + x^2}$

Or: Start by clearing fractions and then differentiate implicitly.

10. $\dfrac{d}{dx}(xe^y) = \dfrac{d}{dx}(x-y) \quad \Rightarrow \quad xe^y\,y' + e^y \cdot 1 = 1 - y' \quad \Rightarrow \quad xe^y\,y' + y' = 1 - e^y \quad \Rightarrow \quad y'(xe^y+1) = 1 - e^y \quad \Rightarrow$

$y' = \dfrac{1-e^y}{xe^y+1}$

11. $\dfrac{d}{dx}(\sin x + \cos y) = \dfrac{d}{dx}(2x-3y) \quad \Rightarrow \quad \cos x - \sin y \cdot y' = 2 - 3y' \quad \Rightarrow \quad 3y' - \sin y \cdot y' = 2 - \cos x \quad \Rightarrow$

$y'(3 - \sin y) = 2 - \cos x \quad \Rightarrow \quad y' = \dfrac{2-\cos x}{3-\sin y}$

12. $\dfrac{d}{dx}(e^x \sin y) = \dfrac{d}{dx}(x+y) \quad \Rightarrow \quad e^x \cos y \cdot y' + \sin y \cdot e^x = 1 + y' \quad \Rightarrow e^x \cos y \cdot y' - y' = 1 - e^x \sin y \quad \Rightarrow$

$y'(e^x \cos y - 1) = 1 - e^x \sin y \quad \Rightarrow \quad y' = \dfrac{1 - e^x \sin y}{e^x \cos y - 1}$

13. $\dfrac{d}{dx}\sin(x+y) = \dfrac{d}{dx}(\cos x + \cos y) \quad \Rightarrow \quad \cos(x+y)\cdot(1+y') = -\sin x - \sin y \cdot y' \quad \Rightarrow$

$\cos(x+y) + y'\cos(x+y) = -\sin x - \sin y \cdot y' \quad \Rightarrow \quad y'\cos(x+y) + \sin y \cdot y' = -\sin x - \cos(x+y) \quad \Rightarrow$

$y'[\cos(x+y) + \sin y] = -[\sin x + \cos(x+y)] \quad \Rightarrow \quad y' = -\dfrac{\cos(x+y) + \sin x}{\cos(x+y) + \sin y}$

14. $\dfrac{d}{dx}\tan(x-y) = \dfrac{d}{dx}(2xy^3+1) \quad \Rightarrow \quad \sec^2(x-y)\cdot(1-y') = 2x(3y^2\,y') + y^3 \cdot 2 \quad \Rightarrow$

$\sec^2(x-y) - y'\sec^2(x-y) = 6xy^2y' + 2y^3 \quad \Rightarrow \quad 6xy^2y' + y'\sec^2(x-y) = \sec^2(x-y) - 2y^3 \quad \Rightarrow$

$y'[6xy^2 + \sec^2(x-y)] = \sec^2(x-y) - 2y^3 \quad \Rightarrow \quad y' = \dfrac{\sec^2(x-y) - 2y^3}{\sec^2(x-y) + 6xy^2}$

15. $\dfrac{d}{dx}(y\cos x) = \dfrac{d}{dx}(x^2+y^2) \quad \Rightarrow \quad y(-\sin x) + \cos x \cdot y' = 2x + 2y\,y' \quad \Rightarrow \quad \cos x \cdot y' - 2y\,y' = 2x + y\sin x \quad \Rightarrow$

$y'(\cos x - 2y) = 2x + y\sin x \quad \Rightarrow \quad y' = \dfrac{2x + y\sin x}{\cos x - 2y}$

16. $\dfrac{d}{dx}\sin(xy) = \dfrac{d}{dx}\cos(x+y) \quad \Rightarrow \quad \cos(xy)\cdot(xy' + y\cdot 1) = -\sin(x+y)\cdot(1+y') \quad \Rightarrow$

$x\cos(xy)\,y' + y\cos(xy) = -\sin(x+y) - y'\sin(x+y) \quad \Rightarrow$

$x\cos(xy)\,y' + y'\sin(x+y) = -y\cos(xy) - \sin(x+y) \quad \Rightarrow$

$[x\cos(xy) + \sin(x+y)]\,y' = -1\,[y\cos(xy) + \sin(x+y)] \quad \Rightarrow \quad y' = -\dfrac{y\cos(xy) + \sin(x+y)}{x\cos(xy) + \sin(x+y)}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\dfrac{d}{dx}\left(2xe^y + ye^x\right) = \dfrac{d}{dx}(3) \quad \Rightarrow \quad (2x \cdot e^y y' + e^y \cdot 2) + (ye^x + e^x\, y') = 0 \quad \Rightarrow \quad 2xe^y\, y' + e^x\, y' = -2e^y - ye^x \quad \Rightarrow$

$y'\left(2xe^y + e^x\right) = -(2e^y + ye^x) \quad \Rightarrow \quad y' = -\dfrac{2e^y + ye^x}{2xe^y + e^x}$

18. $\dfrac{d}{dx}(\sin x\, \cos y) = \dfrac{d}{dx}\left(x^2 - 5y\right) \quad \Rightarrow \quad \sin x\,(-\sin y) \cdot y' + \cos y\,(\cos x) = 2x - 5y' \quad \Rightarrow$

$5y' - \sin x\, \sin y \cdot y' = 2x - \cos x\, \cos y \quad \Rightarrow \quad y'\,(5 - \sin x\, \sin y) = 2x - \cos x\, \cos y \quad \Rightarrow \quad y' = \dfrac{2x - \cos x\, \cos y}{5 - \sin x\, \sin y}$

19. $\dfrac{d}{dx}\sqrt{x+y} = \dfrac{d}{dx}\left(x^4 + y^4\right) \quad \Rightarrow \quad \frac{1}{2}\,(x+y)^{-1/2}\,(1 + y') = 4x^3 + 4y^3 y' \quad \Rightarrow$

$\dfrac{1}{2\sqrt{x+y}} + \dfrac{1}{2\sqrt{x+y}}\, y' = 4x^3 + 4y^3 y' \quad \Rightarrow \quad \dfrac{1}{2\sqrt{x+y}} - 4x^3 = 4y^3 y' - \dfrac{1}{2\sqrt{x+y}} y' \quad \Rightarrow$

$\dfrac{1 - 8x^3\sqrt{x+y}}{2\sqrt{x+y}} = \dfrac{8y^3\sqrt{x+y} - 1}{2\sqrt{x+y}}\, y' \quad \Rightarrow \quad y' = \dfrac{1 - 8x^3\sqrt{x+y}}{8y^3\sqrt{x+y} - 1}$

20. $\dfrac{d}{dx}(xy) = \dfrac{d}{dx}\sqrt{x^2 + y^2} \quad \Rightarrow \quad xy' + y(1) = \frac{1}{2}\left(x^2 + y^2\right)^{-1/2}(2x + 2y\, y') \quad \Rightarrow$

$xy' + y = \dfrac{x}{\sqrt{x^2+y^2}} + \dfrac{y}{\sqrt{x^2+y^2}}\, y' \quad \Rightarrow \quad xy' - \dfrac{y}{\sqrt{x^2+y^2}}\, y' = \dfrac{x}{\sqrt{x^2+y^2}} - y \quad \Rightarrow$

$\dfrac{x\sqrt{x^2+y^2} - y}{\sqrt{x^2+y^2}}\, y' = \dfrac{x - y\sqrt{x^2+y^2}}{\sqrt{x^2+y^2}} \quad \Rightarrow \quad y' = \dfrac{x - y\sqrt{x^2+y^2}}{x\sqrt{x^2+y^2} - y}$

21. $\dfrac{d}{dx}(e^{x/y}) = \dfrac{d}{dx}(x - y) \quad \Rightarrow \quad e^{x/y} \cdot \dfrac{d}{dx}\left(\dfrac{x}{y}\right) = 1 - y' \quad \Rightarrow \quad e^{x/y} \cdot \dfrac{y \cdot 1 - x \cdot y'}{y^2} = 1 - y' \quad \Rightarrow$

$e^{x/y} \cdot \dfrac{1}{y} - \dfrac{xe^{x/y}}{y^2} \cdot y' = 1 - y' \quad \Rightarrow \quad y' - \dfrac{xe^{x/y}}{y^2} \cdot y' = 1 - \dfrac{e^{x/y}}{y} \quad \Rightarrow \quad y'\left(1 - \dfrac{xe^{x/y}}{y^2}\right) = \dfrac{y - e^{x/y}}{y} \quad \Rightarrow$

$y' = \dfrac{\dfrac{y - e^{x/y}}{y}}{\dfrac{y^2 - xe^{x/y}}{y^2}} = \dfrac{y(y - e^{x/y})}{y^2 - xe^{x/y}}$

22. $\dfrac{d}{dx}\cos(x^2 + y^2) = \dfrac{d}{dx}\left(xe^y\right) \quad \Rightarrow \quad -\sin(x^2 + y^2) \cdot (2x + 2y\, y') = xe^y\, y' + e^y \cdot 1 \quad \Rightarrow$

$-2x\sin(x^2 + y^2) - 2y\, y'\sin(x^2 + y^2) = xe^y\, y' + e^y \quad \Rightarrow \quad -e^y - 2x\sin(x^2 + y^2) = xe^y\, y' + 2y\, y'\sin(x^2 + y^2) \quad \Rightarrow$

$-[e^y + 2x\sin(x^2 + y^2)] = y'[xe^y + 2y\sin(x^2 + y^2)] \quad \Rightarrow \quad y' = -\dfrac{e^y + 2x\sin(x^2 + y^2)}{xe^y + 2y\sin(x^2 + y^2)}$

23. $\dfrac{d}{dx}\left\{f(x) + x^2[f(x)]^3\right\} = \dfrac{d}{dx}(10) \quad \Rightarrow \quad f'(x) + x^2 \cdot 3[f(x)]^2 \cdot f'(x) + [f(x)]^3 \cdot 2x = 0.$ If $x = 1$, we have

$f'(1) + 1^2 \cdot 3[f(1)]^2 \cdot f'(1) + [f(1)]^3 \cdot 2(1) = 0 \quad \Rightarrow \quad f'(1) + 1 \cdot 3 \cdot 2^2 \cdot f'(1) + 2^3 \cdot 2 = 0 \quad \Rightarrow$

$f'(1) + 12f'(1) = -16 \quad \Rightarrow \quad 13f'(1) = -16 \quad \Rightarrow \quad f'(1) = -\frac{16}{13}.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $\dfrac{d}{dx}\left[g(x)+x\sin g(x)\right]=\dfrac{d}{dx}\left(x^2\right) \quad\Rightarrow\quad g'(x)+x\cos g(x)\cdot g'(x)+\sin g(x)\cdot 1=2x.$ If $x=0$, we have

$g'(0)+0+\sin g(0)=2(0) \quad\Rightarrow\quad g'(0)+\sin 0=0 \quad\Rightarrow\quad g'(0)+0=0 \quad\Rightarrow\quad g'(0)=0.$

25. $\dfrac{d}{dy}(x^4y^2-x^3y+2xy^3)=\dfrac{d}{dy}(0) \quad\Rightarrow\quad x^4\cdot 2y+y^2\cdot 4x^3\,x'-(x^3\cdot 1+y\cdot 3x^2\,x')+2(x\cdot 3y^2+y^3\cdot x')=0 \quad\Rightarrow$

$4x^3y^2\,x'-3x^2y\,x'+2y^3\,x'=-2x^4y+x^3-6xy^2 \quad\Rightarrow\quad (4x^3y^2-3x^2y+2y^3)\,x'=-2x^4y+x^3-6xy^2 \quad\Rightarrow$

$x'=\dfrac{dx}{dy}=\dfrac{-2x^4y+x^3-6xy^2}{4x^3y^2-3x^2y+2y^3}$

26. $\dfrac{d}{dy}(y\sec x)=\dfrac{d}{dy}(x\tan y) \quad\Rightarrow\quad y\cdot\sec x\,\tan x\cdot x'+\sec x\cdot 1=x\cdot\sec^2 y+\tan y\cdot x' \quad\Rightarrow$

$y\sec x\,\tan x\cdot x'-\tan y\cdot x'=x\sec^2 y-\sec x \quad\Rightarrow\quad (y\sec x\,\tan x-\tan y)\,x'=x\sec^2 y-\sec x \quad\Rightarrow$

$x'=\dfrac{dx}{dy}=\dfrac{x\sec^2 y-\sec x}{y\sec x\tan x-\tan y}$

27. $ye^{\sin x}=x\cos y \quad\Rightarrow\quad ye^{\sin x}\cdot\cos x+e^{\sin x}\cdot y'=x(-\sin y\cdot y')+\cos y\cdot 1 \quad\Rightarrow$

$y'\,e^{\sin x}+y'\,x\sin y=\cos y-y\cos x\,e^{\sin x} \quad\Rightarrow\quad y'\left(e^{\sin x}+x\sin y\right)=\cos y-y\cos x\,e^{\sin x} \quad\Rightarrow$

$y'=\dfrac{\cos y-y\cos x\,e^{\sin x}}{x\sin y+e^{\sin x}}.$

When $x=0$ and $y=0$, we have $y'=\dfrac{\cos 0-0\cdot\cos 0\cdot e^{\sin 0}}{0\cdot\sin 0+e^{\sin 0}}=\dfrac{1-0}{0+1}=1$, so an equation of the tangent line is

$y-0=1\,(x-0)$, or $y=x$.

28. $\tan(x+y)+\sec(x-y)=2 \quad\Rightarrow\quad \sec^2(x+y)\cdot(1+y')+\sec(x-y)\tan(x-y)\cdot(1-y')=0 \quad\Rightarrow$

$\sec^2(x+y)+y'\cdot\sec^2(x+y)+\sec(x-y)\tan(x-y)-y'\cdot\sec(x-y)\tan(x-y)=0 \quad\Rightarrow$

$\sec^2(x+y)+\sec(x-y)\tan(x-y)=y'\cdot\sec(x-y)\tan(x-y)-y'\cdot\sec^2(x+y) \quad\Rightarrow$

$\sec^2(x+y)+\sec(x-y)\tan(x-y)=y'\left[\sec(x-y)\tan(x-y)-\sec^2(x+y)\right] \quad\Rightarrow$

$y'=\dfrac{\sec(x-y)\,\tan(x-y)+\sec^2(x+y)}{\sec(x-y)\,\tan(x-y)-\sec^2(x+y)}.$

When $x=\frac{\pi}{8}$ and $y=\frac{\pi}{8}$, we have

$y'=\dfrac{\sec\left(\frac{\pi}{8}-\frac{\pi}{8}\right)\tan\left(\frac{\pi}{8}-\frac{\pi}{8}\right)+\sec^2\left(\frac{\pi}{8}+\frac{\pi}{8}\right)}{\sec\left(\frac{\pi}{8}-\frac{\pi}{8}\right)\tan\left(\frac{\pi}{8}-\frac{\pi}{8}\right)-\sec^2\left(\frac{\pi}{8}+\frac{\pi}{8}\right)}=\dfrac{\sec 0\,\tan 0+\sec^2\left(\frac{\pi}{4}\right)}{\sec 0\,\tan 0-\sec^2\left(\frac{\pi}{4}\right)}=\dfrac{0+\left(\sqrt{2}\right)^2}{0-\left(\sqrt{2}\right)^2}=-1$, so an equation of the

tangent line is $y-\frac{\pi}{8}=-1\left(x-\frac{\pi}{8}\right)$, or $y=-x+\frac{\pi}{4}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. $x^{2/3} + y^{2/3} = 4 \quad \Rightarrow \quad \frac{2}{3}x^{-1/3} + \frac{2}{3}y^{-1/3}y' = 0 \quad \Rightarrow \quad \dfrac{1}{\sqrt[3]{x}} + \dfrac{y'}{\sqrt[3]{y}} = 0 \quad \Rightarrow \quad y' = -\dfrac{\sqrt[3]{y}}{\sqrt[3]{x}}.$

When $x = -3\sqrt{3}$ and $y = 1$, we have $y' = -\dfrac{1}{\left(-3\sqrt{3}\right)^{1/3}} = -\dfrac{1}{(-3^{3/2})^{1/3}} = \dfrac{1}{3^{1/2}} = \dfrac{1}{\sqrt{3}}$, so an equation of the tangent line is $y - 1 = \frac{1}{\sqrt{3}}\left(x + 3\sqrt{3}\right)$ or $y = \frac{1}{\sqrt{3}}x + 4$.

30. $y^2(6 - x) = x^3 \quad \Rightarrow \quad y^2(-1) + (6 - x) \cdot 2yy' = 3x^2 \quad \Rightarrow \quad y' \cdot 2y(6 - x) = 3x^2 + y^2 \quad \Rightarrow \quad y' = \dfrac{3x^2 + y^2}{2y(6 - x)}.$

When $x = 2$ and $y = \sqrt{2}$, we have $y' = \dfrac{3(2)^2 + \left(\sqrt{2}\right)^2}{2\sqrt{2}(6 - 2)} = \dfrac{12 + 2}{2\sqrt{2} \cdot 4} = \dfrac{14}{8\sqrt{2}} = \dfrac{7}{4\sqrt{2}}$, so an equation of the tangent line is $y - \sqrt{2} = \dfrac{7}{4\sqrt{2}}(x - 2)$, or $y = \dfrac{7}{4\sqrt{2}}x - \dfrac{7}{2\sqrt{2}} + \dfrac{4}{2\sqrt{2}} = \dfrac{7}{4\sqrt{2}}x - \dfrac{3}{2\sqrt{2}}$.

31. $x^2 - xy - y^2 = 1 \quad \Rightarrow \quad 2x - (xy' + y \cdot 1) - 2y\,y' = 0 \quad \Rightarrow \quad 2x - xy' - y - 2y\,y' = 0 \quad \Rightarrow \quad 2x - y = xy' + 2y\,y' \quad \Rightarrow$

$2x - y = (x + 2y)\,y' \quad \Rightarrow \quad y' = \dfrac{2x - y}{x + 2y}.$

When $x = 2$ and $y = 1$, we have $y' = \dfrac{4 - 1}{2 + 2} = \dfrac{3}{4}$, so an equation of the tangent line is $y - 1 = \frac{3}{4}(x - 2)$, or $y = \frac{3}{4}x - \frac{1}{2}$.

32. $x^2 + 2xy + 4y^2 = 12 \quad \Rightarrow \quad 2x + 2x\,y' + 2y + 8y\,y' = 0 \quad \Rightarrow \quad 2x\,y' + 8y\,y' = -2x - 2y \quad \Rightarrow$

$(x + 4y)\,y' = -x - y \quad \Rightarrow \quad y' = -\dfrac{x + y}{x + 4y}.$

When $x = 2$ and $y = 1$, we have $y' = -\dfrac{2 + 1}{2 + 4} = -\dfrac{1}{2}$, so an equation of the tangent line is $y - 1 = -\frac{1}{2}(x - 2)$ or $y = -\frac{1}{2}x + 2$.

33. $x^2 + y^2 = (2x^2 + 2y^2 - x)^2 \quad \Rightarrow \quad 2x + 2y\,y' = 2(2x^2 + 2y^2 - x)(4x + 4y\,y' - 1).$

When $x = 0$ and $y = \frac{1}{2}$, we have $0 + y' = 2(\frac{1}{2})(2y' - 1) \quad \Rightarrow \quad y' = 2y' - 1 \quad \Rightarrow \quad y' = 1$, so an equation of the tangent line is $y - \frac{1}{2} = 1(x - 0)$ or $y = x + \frac{1}{2}$.

34. $x^2y^2 = (y + 1)^2(4 - y^2) \quad \Rightarrow \quad x^2 \cdot 2y\,y' + y^2 \cdot 2x = (y + 1)^2(-2y\,y') + (4 - y^2) \cdot 2(y + 1)\,y' \quad \Rightarrow$

$2x^2y\,y' + 2xy^2 = -2y\,y'(y + 1)^2 + 2y'(4 - y^2)(y + 1) \quad \Rightarrow$

$2xy^2 = 2y'(4 - y^2)(y + 1) - 2y\,y'\,(y + 1)^2 - 2x^2y\,y' \quad \Rightarrow \quad 2xy^2 = y'[2(4 - y^2)(y + 1) - 2y(y + 1)^2 - 2x^2y] \quad \Rightarrow$

$y' = \dfrac{2xy^2}{2(4 - y^2)(y + 1) - 2y(y + 1)^2 - 2x^2y} = \dfrac{xy^2}{(4 - y^2)(y + 1) - y(y + 1)^2 - x^2y}$

When $x = 2\sqrt{3}$ and $y = 1$, we have $y' = \dfrac{2\sqrt{3} \cdot 1^2}{(4 - 1^2)(1 + 1) - 1(1 + 1)^2 - \left(2\sqrt{3}\right)^2 \cdot 1} = \dfrac{2\sqrt{3}}{6 - 4 - 12} = -\dfrac{2\sqrt{3}}{10} = -\dfrac{\sqrt{3}}{5}$,

so an equation of the tangent line is $y - 1 = -\dfrac{\sqrt{3}}{5}\left(x - 2\sqrt{3}\right)$, or $y = -\dfrac{\sqrt{3}}{5}x + \dfrac{11}{5}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. $2(x^2+y^2)^2 = 25(x^2-y^2) \quad \Rightarrow \quad 4(x^2+y^2)(2x+2y\,y') = 25(2x-2y\,y') \quad \Rightarrow$

$4(x+y\,y')(x^2+y^2) = 25(x-y\,y') \quad \Rightarrow \quad 4y\,y'(x^2+y^2)+25y\,y' = 25x-4x(x^2+y^2) \quad \Rightarrow$

$y' = \dfrac{25x-4x(x^2+y^2)}{25y+4y(x^2+y^2)}$.

When $x=3$ and $y=1$, we have $y' = \frac{75-120}{25+40} = -\frac{45}{65} = -\frac{9}{13}$, so an equation of the tangent line is $y-1 = -\frac{9}{13}(x-3)$ or $y = -\frac{9}{13}x + \frac{40}{13}$.

36. $y^2(y^2-4) = x^2(x^2-5) \quad \Rightarrow \quad y^4-4y^2 = x^4-5x^2 \quad \Rightarrow \quad 4y^3\,y'-8y\,y' = 4x^3-10x$.

When $x=0$ and $y=-2$, we have $-32y'+16y'=0 \quad \Rightarrow \quad -16y'=0 \quad \Rightarrow \quad y'=0$, so an equation of the tangent line is $y+2=0(x-0)$ or $y=-2$.

37. (a) $y^2 = 5x^4-x^2 \quad \Rightarrow \quad 2y\,y' = 5(4x^3)-2x \quad \Rightarrow \quad y' = \dfrac{10x^3-x}{y}$.

So at the point $(1,2)$ we have $y' = \dfrac{10(1)^3-1}{2} = \dfrac{9}{2}$, and an equation of the tangent line is $y-2 = \frac{9}{2}(x-1)$ or $y = \frac{9}{2}x - \frac{5}{2}$.

(b)

38. (a) $y^2 = x^3+3x^2 \quad \Rightarrow \quad 2y\,y' = 3x^2+3(2x) \quad \Rightarrow \quad y' = \dfrac{3x^2+6x}{2y}$. So at the point $(1,-2)$ we have

$y' = \dfrac{3(1)^2+6(1)}{2(-2)} = -\dfrac{9}{4}$, and an equation of the tangent line is $y+2 = -\frac{9}{4}(x-1)$ or $y = -\frac{9}{4}x + \frac{1}{4}$.

(b) The curve has a horizontal tangent where $y'=0 \quad \Leftrightarrow$

$3x^2+6x=0 \quad \Leftrightarrow \quad 3x(x+2)=0 \quad \Leftrightarrow \quad x=0$ or $x=-2$.

But note that at $x=0$, $y=0$ also, so the derivative does not exist.

At $x=-2$, $y^2 = (-2)^3+3(-2)^2 = -8+12 = 4$, so $y=\pm 2$.

So the two points at which the curve has a horizontal tangent are $(-2,-2)$ and $(-2,2)$.

(c)

39. $x^2+4y^2 = 4 \quad \Rightarrow \quad 2x+8y\,y' = 0 \quad \Rightarrow \quad y' = -x/(4y) \quad \Rightarrow$

$y'' = -\dfrac{1}{4}\dfrac{y\cdot 1 - x\cdot y'}{y^2} = -\dfrac{1}{4}\dfrac{y-x[-x/(4y)]}{y^2} = -\dfrac{1}{4}\dfrac{4y^2+x^2}{4y^3} = -\dfrac{1}{4}\dfrac{4}{4y^3}$ $\quad$ [since x and y must satisfy the original equation $x^2+4y^2=4$]

Thus, $y'' = -\dfrac{1}{4y^3}$.

40. $x^2+xy+y^2 = 3 \quad \Rightarrow \quad 2x+xy'+y+2y\,y' = 0 \quad \Rightarrow \quad (x+2y)y' = -2x-y \quad \Rightarrow \quad y' = \dfrac{-2x-y}{x+2y}$.

Differentiating $2x+xy'+y+2y\,y' = 0$ to find y'' gives $2+xy''+y'+y'+2y\,y''+2y'y' = 0 \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$(x+2y)\,y'' = -2 - 2y' - 2(y')^2 = -2\left[1 - \frac{2x+y}{x+2y} + \left(\frac{2x+y}{x+2y}\right)^2\right] \quad \Rightarrow$$

$$\begin{aligned} y'' &= -\frac{2}{x+2y}\left[\frac{(x+2y)^2 - (2x+y)(x+2y) + (2x+y)^2}{(x+2y)^2}\right] \\ &= -\frac{2}{(x+2y)^3}\,(x^2 + 4xy + 4y^2 - 2x^2 - 4xy - xy - 2y^2 + 4x^2 + 4xy + y^2) \\ &= -\frac{2}{(x+2y)^3}\,(3x^2 + 3xy + 3y^2) = -\frac{2}{(x+2y)^3}\,(9) \qquad \left[\begin{array}{c}\text{since } x \text{ and } y \text{ must satisfy the} \\ \text{original equation } x^2 + xy + y^2 = 3\end{array}\right] \end{aligned}$$

Thus, $y'' = -\dfrac{18}{(x+2y)^3}$.

41. $\sin y + \cos x = 1 \quad \Rightarrow \quad \cos y \cdot y' - \sin x = 0 \quad \Rightarrow \quad y' = \dfrac{\sin x}{\cos y} \quad \Rightarrow$

$$\begin{aligned} y'' &= \frac{\cos y \cos x - \sin x(-\sin y)\,y'}{(\cos y)^2} = \frac{\cos y \cos x + \sin x\,\sin y(\sin x/\cos y)}{\cos^2 y} \\ &= \frac{\cos^2 y \cos x + \sin^2 x \sin y}{\cos^2 y \cos y} = \frac{\cos^2 y \cos x + \sin^2 x \sin y}{\cos^3 y} \end{aligned}$$

42. $x^3 - y^3 = 7 \quad \Rightarrow \quad 3x^2 - 3y^2 y' = 0 \quad \Rightarrow \quad y' = \dfrac{x^2}{y^2} \quad \Rightarrow$

$$y'' = \frac{y^2(2x) - x^2(2y\,y')}{(y^2)^2} = \frac{2xy[y - x(x^2/y^2)]}{y^4} = \frac{2x(y - x^3/y^2)}{y^3} = \frac{2x(y^3 - x^3)}{y^3 y^2} = \frac{2x(-7)}{y^5} = \frac{-14x}{y^5}$$

43. If $x = 0$ in $xy + e^y = e$, then we get $0 + e^y = e$, so $y = 1$ and the point where $x = 0$ is $(0, 1)$. Differentiating implicitly with respect to x gives us $x\,y' + y \cdot 1 + e^y\,y' = 0$. Substituting 0 for x and 1 for y gives us $0 + 1 + e\,y' = 0 \quad \Rightarrow$ $e\,y' = -1 \quad \Rightarrow \quad y' = -1/e$. Differentiating $x\,y' + y + e^y\,y' = 0$ implicitly with respect to x gives us $x\,y'' + y' \cdot 1 + y' + e^y\,y'' + y' \cdot e^y\,y' = 0$. Now substitute 0 for x, 1 for y, and $-1/e$ for y'.

$$0 + \left(-\frac{1}{e}\right) + \left(-\frac{1}{e}\right) + e\,y'' + \left(-\frac{1}{e}\right)(e)\left(-\frac{1}{e}\right) = 0 \quad \Rightarrow \quad -\frac{2}{e} + e\,y'' + \frac{1}{e} = 0 \quad \Rightarrow \quad e\,y'' = \frac{1}{e} \quad \Rightarrow \quad y'' = \frac{1}{e^2}.$$

44. If $x = 1$ in $x^2 + xy + y^3 = 1$, then we get $1 + y + y^3 = 1 \quad \Rightarrow \quad y^3 + y = 0 \quad \Rightarrow \quad y(y^2 + 1) \quad \Rightarrow \quad y = 0$, so the point where $x = 1$ is $(1, 0)$. Differentiating implicitly with respect to x gives us $2x + x\,y' + y \cdot 1 + 3y^2 \cdot y' = 0$. Substituting 1 for x and 0 for y gives us $2 + y' + 0 + 0 = 0 \quad \Rightarrow \quad y' = -2$. Differentiating $2x + x\,y' + y + 3y^2\,y' = 0$ implicitly with respect to x gives us $2 + x\,y'' + y' \cdot 1 + y' + 3(y^2\,y'' + y' \cdot 2yy') = 0$. Now substitute 1 for x, 0 for y, and -2 for y'.

$2 + y'' + (-2) + (-2) + 3(0 + 0) = 0 \quad \Rightarrow \quad y'' = 2$. Differentiating $2 + x\,y'' + 2\,y' + 3y^2\,y'' + 6y(y')^2 = 0$ implicitly with respect to x gives us $x\,y''' + y'' \cdot 1 + 2\,y'' + 3(y^2\,y''' + y'' \cdot 2y\,y') + 6[y \cdot 2y'\,y'' + (y')^2\,y'] = 0$. Now substitute 1 for x, 0 for y, -2 for y', and 2 for y''. $y''' + 2 + 4 + 3(0 + 0) + 6[0 + (-8)] = 0 \quad \Rightarrow \quad y''' = -2 - 4 + 48 = 42$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. (a) There are eight points with horizontal tangents: four at $x \approx 1.57735$ and four at $x \approx 0.42265$.

(b) $y' = \dfrac{3x^2 - 6x + 2}{2(2y^3 - 3y^2 - y + 1)} \quad \Rightarrow \quad y' = -1$ at $(0, 1)$ and $y' = \frac{1}{3}$ at $(0, 2)$.

Equations of the tangent lines are $y = -x + 1$ and $y = \frac{1}{3}x + 2$.

(c) $y' = 0 \quad \Rightarrow \quad 3x^2 - 6x + 2 = 0 \quad \Rightarrow \quad x = 1 \pm \frac{1}{3}\sqrt{3}$

(d) By multiplying the right side of the equation by $x - 3$, we obtain the first graph. By modifying the equation in other ways, we can generate the other graphs.

$y(y^2 - 1)(y - 2)$
$= x(x - 1)(x - 2)(x - 3)$

$y(y^2 - 4)(y - 2)$
$= x(x - 1)(x - 2)$

$y(y + 1)(y^2 - 1)(y - 2)$
$= x(x - 1)(x - 2)$

$(y + 1)(y^2 - 1)(y - 2)$
$= (x - 1)(x - 2)$

$x(y + 1)(y^2 - 1)(y - 2)$
$= y(x - 1)(x - 2)$

$y(y^2 + 1)(y - 2)$
$= x(x^2 - 1)(x - 2)$

$y(y + 1)(y^2 - 2)$
$= x(x - 1)(x^2 - 2)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. (a)

(b) $\dfrac{d}{dx}(2y^3 + y^2 - y^5) = \dfrac{d}{dx}(x^4 - 2x^3 + x^2) \quad \Rightarrow$

$6y^2y' + 2y\,y' - 5y^4y' = 4x^3 - 6x^2 + 2x \quad \Rightarrow$

$y' = \dfrac{2x(2x^2 - 3x + 1)}{6y^2 + 2y - 5y^4} = \dfrac{2x(2x-1)(x-1)}{y(6y + 2 - 5y^3)}$. From the graph and the values for which $y' = 0$, we speculate that there are 9 points with horizontal tangents: 3 at $x = 0$, 3 at $x = \frac{1}{2}$, and 3 at $x = 1$. The three horizontal tangents along the top of the wagon are hard to find, but by limiting the y-range of the graph (to $[1.6, 1.7]$, for example) they are distinguishable.

47. From Exercise 35, a tangent to the lemniscate will be horizontal if $y' = 0 \quad \Rightarrow \quad 25x - 4x(x^2 + y^2) = 0 \quad \Rightarrow$ $x[25 - 4(x^2 + y^2)] = 0 \quad \Rightarrow \quad x^2 + y^2 = \frac{25}{4}$ (**1**). (Note that when x is 0, y is also 0, and there is no horizontal tangent at the origin.) Substituting $\frac{25}{4}$ for $x^2 + y^2$ in the equation of the lemniscate, $2(x^2 + y^2)^2 = 25(x^2 - y^2)$, we get $x^2 - y^2 = \frac{25}{8}$ (**2**). Solving (**1**) and (**2**), we have $x^2 = \frac{75}{16}$ and $y^2 = \frac{25}{16}$, so the four points are $\left(\pm\frac{5\sqrt{3}}{4}, \pm\frac{5}{4}\right)$.

48. $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 \quad \Rightarrow \quad \dfrac{2x}{a^2} + \dfrac{2yy'}{b^2} = 0 \quad \Rightarrow \quad y' = -\dfrac{b^2x}{a^2y} \quad \Rightarrow$ an equation of the tangent line at (x_0, y_0) is $y - y_0 = \dfrac{-b^2x_0}{a^2y_0}(x - x_0)$. Multiplying both sides by $\dfrac{y_0}{b^2}$ gives $\dfrac{y_0y}{b^2} - \dfrac{y_0^2}{b^2} = -\dfrac{x_0x}{a^2} + \dfrac{x_0^2}{a^2}$. Since (x_0, y_0) lies on the ellipse, we have $\dfrac{x_0x}{a^2} + \dfrac{y_0y}{b^2} = \dfrac{x_0^2}{a^2} + \dfrac{y_0^2}{b^2} = 1$.

49. $\dfrac{x^2}{a^2} - \dfrac{y^2}{b^2} = 1 \quad \Rightarrow \quad \dfrac{2x}{a^2} - \dfrac{2yy'}{b^2} = 0 \quad \Rightarrow \quad y' = \dfrac{b^2x}{a^2y} \quad \Rightarrow$ an equation of the tangent line at (x_0, y_0) is $y - y_0 = \dfrac{b^2x_0}{a^2y_0}(x - x_0)$. Multiplying both sides by $\dfrac{y_0}{b^2}$ gives $\dfrac{y_0y}{b^2} - \dfrac{y_0^2}{b^2} = \dfrac{x_0x}{a^2} - \dfrac{x_0^2}{a^2}$. Since (x_0, y_0) lies on the hyperbola, we have $\dfrac{x_0x}{a^2} - \dfrac{y_0y}{b^2} = \dfrac{x_0^2}{a^2} - \dfrac{y_0^2}{b^2} = 1$.

50. $\sqrt{x} + \sqrt{y} = \sqrt{c} \quad \Rightarrow \quad \dfrac{1}{2\sqrt{x}} + \dfrac{y'}{2\sqrt{y}} = 0 \quad \Rightarrow \quad y' = -\dfrac{\sqrt{y}}{\sqrt{x}} \quad \Rightarrow$ an equation of the tangent line at (x_0, y_0) is $y - y_0 = -\dfrac{\sqrt{y_0}}{\sqrt{x_0}}(x - x_0)$. Now $x = 0 \quad \Rightarrow \quad y = y_0 - \dfrac{\sqrt{y_0}}{\sqrt{x_0}}(-x_0) = y_0 + \sqrt{x_0}\sqrt{y_0}$, so the y-intercept is $y_0 + \sqrt{x_0}\sqrt{y_0}$. And $y = 0 \quad \Rightarrow \quad -y_0 = -\dfrac{\sqrt{y_0}}{\sqrt{x_0}}(x - x_0) \quad \Rightarrow \quad x - x_0 = \dfrac{y_0\sqrt{x_0}}{\sqrt{y_0}} \quad \Rightarrow \quad x = x_0 + \sqrt{x_0}\sqrt{y_0}$, so the x-intercept is $x_0 + \sqrt{x_0}\sqrt{y_0}$. The sum of the intercepts is

$$\left(y_0 + \sqrt{x_0}\sqrt{y_0}\right) + \left(x_0 + \sqrt{x_0}\sqrt{y_0}\right) = x_0 + 2\sqrt{x_0}\sqrt{y_0} + y_0 = \left(\sqrt{x_0} + \sqrt{y_0}\right)^2 = \left(\sqrt{c}\right)^2 = c.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. If the circle has radius r, its equation is $x^2 + y^2 = r^2 \quad \Rightarrow \quad 2x + 2yy' = 0 \quad \Rightarrow \quad y' = -\dfrac{x}{y}$, so the slope of the tangent line at $P(x_0, y_0)$ is $-\dfrac{x_0}{y_0}$. The negative reciprocal of that slope is $\dfrac{-1}{-x_0/y_0} = \dfrac{y_0}{x_0}$, which is the slope of OP, so the tangent line at P is perpendicular to the radius OP.

52. $y^q = x^p \quad \Rightarrow \quad qy^{q-1}y' = px^{p-1} \quad \Rightarrow \quad y' = \dfrac{px^{p-1}}{qy^{q-1}} = \dfrac{px^{p-1}y}{qy^q} = \dfrac{px^{p-1}x^{p/q}}{qx^p} = \dfrac{p}{q}x^{(p/q)-1}$

53. $x^2 + y^2 = r^2$ is a circle with center O and $ax + by = 0$ is a line through O [assume a and b are not both zero]. $x^2 + y^2 = r^2 \quad \Rightarrow \quad 2x + 2yy' = 0 \quad \Rightarrow \quad y' = -x/y$, so the slope of the tangent line at $P_0\,(x_0, y_0)$ is $-x_0/y_0$. The slope of the line OP_0 is y_0/x_0, which is the negative reciprocal of $-x_0/y_0$. Hence, the curves are orthogonal, and the families of curves are orthogonal trajectories of each other.

54. The circles $x^2 + y^2 = ax$ and $x^2 + y^2 = by$ intersect at the origin where the tangents are vertical and horizontal [assume a and b are both nonzero]. If (x_0, y_0) is the other point of intersection, then $x_0^2 + y_0^2 = ax_0$ (**1**) and $x_0^2 + y_0^2 = by_0$ (**2**).

Now $x^2 + y^2 = ax \quad \Rightarrow \quad 2x + 2yy' = a \quad \Rightarrow \quad y' = \dfrac{a - 2x}{2y}$ and $x^2 + y^2 = by \quad \Rightarrow$ $2x + 2yy' = by' \quad \Rightarrow \quad y' = \dfrac{2x}{b - 2y}$. Thus, the curves are orthogonal at $(x_0, y_0) \quad \Leftrightarrow$ $\dfrac{a - 2x_0}{2y_0} = -\dfrac{b - 2y_0}{2x_0} \quad \Leftrightarrow \quad 2ax_0 - 4x_0^2 = 4y_0^2 - 2by_0 \quad \Leftrightarrow \quad ax_0 + by_0 = 2(x_0^2 + y_0^2)$, which is true by (**1**) and (**2**).

55. $y = cx^2 \quad \Rightarrow \quad y' = 2cx$ and $x^2 + 2y^2 = k$ [assume $k > 0$] $\quad \Rightarrow \quad 2x + 4yy' = 0 \quad \Rightarrow$ $2yy' = -x \quad \Rightarrow \quad y' = -\dfrac{x}{2(y)} = -\dfrac{x}{2(cx^2)} = -\dfrac{1}{2cx}$, so the curves are orthogonal if $c \neq 0$. If $c = 0$, then the horizontal line $y = cx^2 = 0$ intersects $x^2 + 2y^2 = k$ orthogonally at $\left(\pm\sqrt{k}, 0\right)$, since the ellipse $x^2 + 2y^2 = k$ has vertical tangents at those two points.

56. $y = ax^3 \quad \Rightarrow \quad y' = 3ax^2$ and $x^2 + 3y^2 = b$ [assume $b > 0$] $\quad \Rightarrow \quad 2x + 6yy' = 0 \quad \Rightarrow$ $3yy' = -x \quad \Rightarrow \quad y' = -\dfrac{x}{3(y)} = -\dfrac{x}{3(ax^3)} = -\dfrac{1}{3ax^2}$, so the curves are orthogonal if $a \neq 0$. If $a = 0$, then the horizontal line $y = ax^3 = 0$ intesects $x^2 + 3y^2 = b$ orthogonally at $\left(\pm\sqrt{b}, 0\right)$, since the ellipse $x^2 + 3y^2 = b$ has vertical tangents at those two points.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. Since $A^2 < a^2$, we are assured that there are four points of intersection.

(1) $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 \quad \Rightarrow \quad \dfrac{2x}{a^2} + \dfrac{2yy'}{b^2} = 0 \quad \Rightarrow \quad \dfrac{yy'}{b^2} = -\dfrac{x}{a^2} \quad \Rightarrow$

$y' = m_1 = -\dfrac{xb^2}{ya^2}.$

(2) $\dfrac{x^2}{A^2} - \dfrac{y^2}{B^2} = 1 \quad \Rightarrow \quad \dfrac{2x}{A^2} - \dfrac{2yy'}{B^2} = 0 \quad \Rightarrow \quad \dfrac{yy'}{B^2} = \dfrac{x}{A^2} \quad \Rightarrow$

$y' = m_2 = \dfrac{xB^2}{yA^2}.$

Now $m_1 m_2 = -\dfrac{xb^2}{ya^2} \cdot \dfrac{xB^2}{yA^2} = -\dfrac{b^2B^2}{a^2A^2} \cdot \dfrac{x^2}{y^2}$ **(3)**. Subtracting equations, **(1)** − **(2)**, gives us $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} - \dfrac{x^2}{A^2} + \dfrac{y^2}{B^2} = 0 \quad \Rightarrow$

$\dfrac{y^2}{b^2} + \dfrac{y^2}{B^2} = \dfrac{x^2}{A^2} - \dfrac{x^2}{a^2} \quad \Rightarrow \quad \dfrac{y^2B^2 + y^2b^2}{b^2B^2} = \dfrac{x^2a^2 - x^2A^2}{A^2a^2} \quad \Rightarrow \quad \dfrac{y^2(b^2+B^2)}{b^2B^2} = \dfrac{x^2(a^2-A^2)}{a^2A^2}$ **(4)**. Since

$a^2 - b^2 = A^2 + B^2$, we have $a^2 - A^2 = b^2 + B^2$. Thus, equation **(4)** becomes $\dfrac{y^2}{b^2B^2} = \dfrac{x^2}{A^2a^2} \quad \Rightarrow \quad \dfrac{x^2}{y^2} = \dfrac{A^2a^2}{b^2B^2}$, and

substituting for $\dfrac{x^2}{y^2}$ in equation **(3)** gives us $m_1 m_2 = -\dfrac{b^2B^2}{a^2A^2} \cdot \dfrac{a^2A^2}{b^2B^2} = -1$. Hence, the ellipse and hyperbola are orthogonal trajectories.

58. $y = (x+c)^{-1} \quad \Rightarrow \quad y' = -(x+c)^{-2}$ and $y = a(x+k)^{1/3} \quad \Rightarrow \quad y' = \frac{1}{3}a(x+k)^{-2/3}$, so the curves are othogonal if the product of the slopes is -1, that is, $\dfrac{-1}{(x+c)^2} \cdot \dfrac{a}{3(x+k)^{2/3}} = -1 \quad \Rightarrow \quad a = 3(x+c)^2(x+k)^{2/3} \quad \Rightarrow$

$a = 3\left(\dfrac{1}{y}\right)^2 \left(\dfrac{y}{a}\right)^2$ [since $y^2 = (x+c)^{-2}$ and $y^2 = a^2(x+k)^{2/3}$] $\quad \Rightarrow \quad a = 3\left(\dfrac{1}{a^2}\right) \quad \Rightarrow \quad a^3 = 3 \quad \Rightarrow \quad a = \sqrt[3]{3}.$

59. (a) $\left(P + \dfrac{n^2a}{V^2}\right)(V - nb) = nRT \quad \Rightarrow \quad PV - Pnb + \dfrac{n^2a}{V} - \dfrac{n^3ab}{V^2} = nRT \quad \Rightarrow$

$\dfrac{d}{dP}(PV - Pnb + n^2aV^{-1} - n^3abV^{-2}) = \dfrac{d}{dP}(nRT) \quad \Rightarrow$

$PV' + V \cdot 1 - nb - n^2aV^{-2} \cdot V' + 2n^3abV^{-3} \cdot V' = 0 \quad \Rightarrow \quad V'(P - n^2aV^{-2} + 2n^3abV^{-3}) = nb - V \quad \Rightarrow$

$V' = \dfrac{nb - V}{P - n^2aV^{-2} + 2n^3abV^{-3}}$ or $\dfrac{dV}{dP} = \dfrac{V^3(nb - V)}{PV^3 - n^2aV + 2n^3ab}$

(b) Using the last expression for dV/dP from part (a), we get

$$\frac{dV}{dP} = \frac{(10\text{ L})^3[(1\text{ mole})(0.04267\text{ L/mole}) - 10\text{ L}]}{\left[\begin{array}{l}(2.5\text{ atm})(10\text{ L})^3 - (1\text{ mole})^2(3.592\text{ L}^2\text{-atm/ mole}^2)(10\text{ L}) \\ \qquad\qquad + 2(1\text{ mole})^3(3.592\text{ L}^2\text{-atm/ mole}^2)(0.04267\text{ L/ mole})\end{array}\right]}$$

$$= \frac{-9957.33\text{ L}^4}{2464.386541\text{ L}^3\text{-atm}} \approx -4.04\text{ L/ atm}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. (a) $x^2 + xy + y^2 + 1 = 0 \quad \Rightarrow \quad 2x + xy' + y \cdot 1 + 2yy' + 0 = 0 \quad \Rightarrow \quad y'(x + 2y) = -2x - y \quad \Rightarrow \quad y' = \dfrac{-2x - y}{x + 2y}$

(b) Plotting the curve in part (a) gives us an empty graph, that is, there are no points that satisfy the equation. If there were any points that satisfied the equation, then x and y would have opposite signs; otherwise, all the terms are positive and their sum can not equal 0. $x^2 + xy + y^2 + 1 = 0 \quad \Rightarrow \quad x^2 + 2xy + y^2 - xy + 1 = 0 \quad \Rightarrow \quad (x + y)^2 = xy - 1$. The left side of the last equation is nonnegative, but the right side is at most -1, so that proves there are no points that satisfy the equation.

Another solution: $x^2 + xy + y^2 + 1 = \frac{1}{2}x^2 + xy + \frac{1}{2}y^2 + \frac{1}{2}x^2 + \frac{1}{2}y^2 + 1 = \frac{1}{2}(x^2 + 2xy + y^2) + \frac{1}{2}(x^2 + y^2) + 1$

$$= \tfrac{1}{2}(x + y)^2 + \tfrac{1}{2}(x^2 + y^2) + 1 \geq 1$$

Another solution: Regarding $x^2 + xy + y^2 + 1 = 0$ as a quadratic in x, the discriminant is $y^2 - 4(y^2 + 1) = -3y^2 - 4$. This is negative, so there are no real solutions.

(c) The expression for y' in part (a) is meaningless; that is, since the equation in part (a) has no solution, it does not implicitly define a function y of x, and therefore it is meaningless to consider y'.

61. To find the points at which the ellipse $x^2 - xy + y^2 = 3$ crosses the x-axis, let $y = 0$ and solve for x.

$y = 0 \quad \Rightarrow \quad x^2 - x(0) + 0^2 = 3 \quad \Leftrightarrow \quad x = \pm\sqrt{3}$. So the graph of the ellipse crosses the x-axis at the points $\left(\pm\sqrt{3}, 0\right)$.

Using implicit differentiation to find y', we get $2x - xy' - y + 2yy' = 0 \quad \Rightarrow \quad y'(2y - x) = y - 2x \quad \Leftrightarrow \quad y' = \dfrac{y - 2x}{2y - x}$.

So y' at $\left(\sqrt{3}, 0\right)$ is $\dfrac{0 - 2\sqrt{3}}{2(0) - \sqrt{3}} = 2$ and y' at $\left(-\sqrt{3}, 0\right)$ is $\dfrac{0 + 2\sqrt{3}}{2(0) + \sqrt{3}} = 2$. Thus, the tangent lines at these points are parallel.

62. (a) We use implicit differentiation to find $y' = \dfrac{y - 2x}{2y - x}$ as in Exercise 61. The slope of the tangent line at $(-1, 1)$ is $m = \dfrac{1 - 2(-1)}{2(1) - (-1)} = \dfrac{3}{3} = 1$, so the slope of the normal line is $-\dfrac{1}{m} = -1$, and its equation is $y - 1 = -1(x + 1) \quad \Leftrightarrow$ $y = -x$. Substituting $-x$ for y in the equation of the ellipse, we get $x^2 - x(-x) + (-x)^2 = 3 \quad \Rightarrow \quad 3x^2 = 3 \quad \Leftrightarrow \quad x = \pm 1$. So the normal line must intersect the ellipse again at $x = 1$, and since the equation of the line is $y = -x$, the other point of intersection must be $(1, -1)$.

(b)

63. $x^2y^2 + xy = 2 \quad \Rightarrow \quad x^2 \cdot 2yy' + y^2 \cdot 2x + x \cdot y' + y \cdot 1 = 0 \quad \Leftrightarrow \quad y'(2x^2y + x) = -2xy^2 - y \quad \Leftrightarrow$

$y' = -\dfrac{2xy^2 + y}{2x^2y + x}$. So $-\dfrac{2xy^2 + y}{2x^2y + x} = -1 \quad \Leftrightarrow \quad 2xy^2 + y = 2x^2y + x \quad \Leftrightarrow \quad y(2xy + 1) = x(2xy + 1) \quad \Leftrightarrow$

$y(2xy + 1) - x(2xy + 1) = 0 \quad \Leftrightarrow \quad (2xy + 1)(y - x) = 0 \quad \Leftrightarrow \quad xy = -\frac{1}{2}$ or $y = x$. But $xy = -\frac{1}{2} \quad \Rightarrow$

$x^2y^2 + xy = \frac{1}{4} - \frac{1}{2} \neq 2$, so we must have $x = y$. Then $x^2y^2 + xy = 2 \quad \Rightarrow \quad x^4 + x^2 = 2 \quad \Leftrightarrow \quad x^4 + x^2 - 2 = 0 \quad \Leftrightarrow$

$(x^2 + 2)(x^2 - 1) = 0$. So $x^2 = -2$, which is impossible, or $x^2 = 1 \quad \Leftrightarrow \quad x = \pm 1$. Since $x = y$, the points on the curve where the tangent line has a slope of -1 are $(-1, -1)$ and $(1, 1)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. $x^2 + 4y^2 = 36 \quad \Rightarrow \quad 2x + 8yy' = 0 \quad \Rightarrow \quad y' = -\dfrac{x}{4y}$. Let (a, b) be a point on $x^2 + 4y^2 = 36$ whose tangent line passes through $(12, 3)$. The tangent line is then $y - 3 = -\dfrac{a}{4b}(x - 12)$, so $b - 3 = -\dfrac{a}{4b}(a - 12)$. Multiplying both sides by $4b$ gives $4b^2 - 12b = -a^2 + 12a$, so $4b^2 + a^2 = 12(a + b)$. But $4b^2 + a^2 = 36$, so $36 = 12(a + b) \quad \Rightarrow \quad a + b = 3 \quad \Rightarrow$ $b = 3 - a$. Substituting $3 - a$ for b into $a^2 + 4b^2 = 36$ gives $a^2 + 4(3 - a)^2 = 36 \quad \Leftrightarrow \quad a^2 + 36 - 24a + 4a^2 = 36 \quad \Leftrightarrow$ $5a^2 - 24a = 0 \quad \Leftrightarrow \quad a(5a - 24) = 0$, so $a = 0$ or $a = \frac{24}{5}$. If $a = 0$, $b = 3 - 0 = 3$, and if $a = \frac{24}{5}$, $b = 3 - \frac{24}{5} = -\frac{9}{5}$. So the two points on the ellipse are $(0, 3)$ and $\left(\frac{24}{5}, -\frac{9}{5}\right)$. Using $y - 3 = -\dfrac{a}{4b}(x - 12)$ with $(a, b) = (0, 3)$ gives us the tangent line $y - 3 = 0$ or $y = 3$. With $(a, b) = \left(\frac{24}{5}, -\frac{9}{5}\right)$, we have $y - 3 = -\frac{24/5}{4(-9/5)}(x - 12) \quad \Leftrightarrow \quad y - 3 = \frac{2}{3}(x - 12) \quad \Leftrightarrow \quad y = \frac{2}{3}x - 5$. A graph of the ellipse and the tangent lines confirms our results.

65. For $\dfrac{x}{y} = y^2 + 1$, $y \neq 0$, we have $\dfrac{d}{dx}\left(\dfrac{x}{y}\right) = \dfrac{d}{dx}(y^2 + 1) \quad \Rightarrow \quad \dfrac{y \cdot 1 - x \cdot y'}{y^2} = 2y\,y' \quad \Rightarrow \quad y - x\,y' = 2y^3\,y' \quad \Rightarrow$

$2y^3\,y' + x\,y' = y \quad \Rightarrow \quad y'\,(2y^3 + x) = y \quad \Rightarrow \quad y' = \dfrac{y}{2y^3 + x}$.

For $x = y^3 + y$, $y \neq 0$, we have $\dfrac{d}{dx}(x) = \dfrac{d}{dx}(y^3 + y) \quad \Rightarrow \quad 1 = 3y^2 y' + y' \quad \Rightarrow \quad 1 = y'\,(3y^2 + 1) \quad \Rightarrow$

$y' = \dfrac{1}{3y^2 + 1}$.

From part (a), $y' = \dfrac{y}{2y^3 + x}$. Since $y \neq 0$, we substitute $y^3 + y$ for x to get

$\dfrac{y}{2y^3 + x} = \dfrac{y}{2y^3 + (y^3 + y)} = \dfrac{y}{3y^3 + y} = \dfrac{y}{y(3y^2 + 1)} = \dfrac{1}{3y^2 + 1}$, which agrees with part (b).

66. (a) $y = J(x)$ and $xy'' + y' + xy = 0 \quad \Rightarrow \quad xJ''(x) + J'(x) + xJ(x) = 0$. If $x = 0$, we have $0 + J'(0) + 0 = 0$, so $J'(0) = 0$.

(b) Differentiating $xy'' + y' + xy = 0$ implicitly, we get $xy''' + y'' \cdot 1 + y'' + xy' + y \cdot 1 = 0 \quad \Rightarrow$ $xy''' + 2y'' + xy' + y = 0$, so $xJ'''(x) + 2J''(x) + xJ'(x) + J(x) = 0$. If $x = 0$, we have $0 + 2J''(0) + 0 + 1 \quad [J(0) = 1 \text{ is given}] \quad = 0 \quad \Rightarrow \quad 2J''(0) = -1 \quad \Rightarrow \quad J''(0) = -\frac{1}{2}$.

67. $x^2 + 4y^2 = 5 \quad \Rightarrow \quad 2x + 4(2yy') = 0 \quad \Rightarrow \quad y' = -\dfrac{x}{4y}$. Now let h be the height of the lamp, and let (a, b) be the point of tangency of the line passing through the points $(3, h)$ and $(-5, 0)$. This line has slope $(h - 0)/[3 - (-5)] = \frac{1}{8}h$. But the slope of the tangent line through the point (a, b) can be expressed as $y' = -\dfrac{a}{4b}$, or as $\dfrac{b - 0}{a - (-5)} = \dfrac{b}{a + 5}$ [since the line passes through $(-5, 0)$ and (a, b)], so $-\dfrac{a}{4b} = \dfrac{b}{a + 5} \quad \Leftrightarrow \quad 4b^2 = -a^2 - 5a \quad \Leftrightarrow \quad a^2 + 4b^2 = -5a$. But $a^2 + 4b^2 = 5$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

[since (a, b) is on the ellipse], so $5 = -5a \quad \Leftrightarrow \quad a = -1$. Then $4b^2 = -a^2 - 5a = -1 - 5(-1) = 4 \quad \Rightarrow \quad b = 1$, since the point is on the top half of the ellipse. So $\dfrac{h}{8} = \dfrac{b}{a+5} = \dfrac{1}{-1+5} = \dfrac{1}{4} \quad \Rightarrow \quad h = 2$. So the lamp is located 2 units above the x-axis.

DISCOVERY PROJECT Families of Implicit Curves

1. (a) There appear to be nine points of intersection. The "inner four" near the origin are about $(\pm 0.2, -0.9)$ and $(\pm 0.3, -1.1)$. The "outer five" are about $(2.0, -8.9)$, $(-2.8, -8.8)$, $(-7.5, -7.7)$, $(-7.8, -4.7)$, and $(-8.0, 1.5)$.

(b) We see from the graphs with $c = 5$ and $c = 10$, and for other values of c, that the curves change shape but the nine points of intersection are the same.

2. (a) If $c = 0$, the graph is the unit circle. As c increases, the graph looks more diamondlike and then more crosslike (see the graph for $c \geq 0$).

For $-1 < c < 0$ (see the graph), there are four hyperboliclike branches as well as an ellipticlike curve bounded by $|x| \leq 1$ and $|y| \leq 1$ for values of c close to 0. As c gets closer to -1, the branches and the curve become more rectangular, approaching the lines $|x| = 1$ and $|y| = 1$.

For $c = -1$, we get the lines $x = \pm 1$ and $y = \pm 1$. As c decreases, we get four test-tubelike curves (see the graph) that are bounded by $|x| = 1$ and $|y| = 1$, and get thinner as $|c|$ gets larger.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The curve for $c = -1$ is described in part (a). When $c = -1$, we get

$x^2 + y^2 - x^2y^2 = 1 \quad \Leftrightarrow \quad 0 = x^2y^2 - x^2 - y^2 + 1 \quad \Leftrightarrow \quad 0 = (x^2 - 1)(y^2 - 1) \quad \Leftrightarrow \quad x = \pm 1$ or $y = \pm 1$, which algebraically proves that the graph consists of the stated lines.

(c) $\dfrac{d}{dx}(x^2 + y^2 + cx^2y^2) = \dfrac{d}{dx}(1) \quad \Rightarrow \quad 2x + 2y\,y' + c(x^2 \cdot 2y\,y' + y^2 \cdot 2x) = 0 \quad \Rightarrow$

$2y\,y' + 2cx^2y\,y' = -2x - 2cxy^2 \quad \Rightarrow \quad 2y(1 + cx^2)y' = -2x(1 + cy^2) \quad \Rightarrow \quad y' = -\dfrac{x(1 + cy^2)}{y(1 + cx^2)}$.

For $c = -1$, $y' = -\dfrac{x(1 - y^2)}{y(1 - x^2)} = -\dfrac{x(1 + y)(1 - y)}{y(1 + x)(1 - x)}$, so $y' = 0$ when $y = \pm 1$ or $x = 0$ (which leads to $y = \pm 1$) and y' is undefined when $x = \pm 1$ or $y = 0$ (which leads to $x = \pm 1$). Since the graph consists of the lines $x = \pm 1$ and $y = \pm 1$, the slope at any point on the graph is undefined or 0, which is consistent with the expression found for y'.

3.6 Derivatives of Logarithmic and Inverse Trigonometric Functions

1. The differentiation formula for logarithmic functions, $\dfrac{d}{dx}(\log_a x) = \dfrac{1}{x \ln a}$, is simplest when $a = e$ because $\ln e = 1$.

2. $g(t) = \ln(3 + t^2) \quad \Rightarrow \quad g'(t) = \dfrac{1}{3 + t^2} \cdot \dfrac{d}{dt}(3 + t^2) = \dfrac{1}{3 + t^2} \cdot 2t = \dfrac{2t}{3 + t^2}$

3. $f(x) = \ln(x^2 + 3x + 5) \quad \Rightarrow \quad f'(x) = \dfrac{1}{x^2 + 3x + 5} \cdot \dfrac{d}{dx}(x^2 + 3x + 5) = \dfrac{1}{x^2 + 3x + 5} \cdot (2x + 3) = \dfrac{2x + 3}{x^2 + 3x + 5}$

4. $f(x) = x \ln x - x \quad \Rightarrow \quad f'(x) = x \cdot \dfrac{1}{x} + (\ln x) \cdot 1 - 1 = 1 + \ln x - 1 = \ln x$

5. $f(x) = \sin(\ln x) \quad \Rightarrow \quad f'(x) = \cos(\ln x) \cdot \dfrac{d}{dx} \ln x = \cos(\ln x) \cdot \dfrac{1}{x} = \dfrac{\cos(\ln x)}{x}$

6. $f(x) = \ln(\sin^2 x) = \ln(\sin x)^2 = 2 \ln|\sin x| \quad \Rightarrow \quad f'(x) = 2 \cdot \dfrac{1}{\sin x} \cdot \cos x = 2 \cot x$

7. $f(x) = \ln \dfrac{1}{x} \quad \Rightarrow \quad f'(x) = \dfrac{1}{1/x} \dfrac{d}{dx}\left(\dfrac{1}{x}\right) = x\left(-\dfrac{1}{x^2}\right) = -\dfrac{1}{x}$.

Another solution: $f(x) = \ln \dfrac{1}{x} = \ln 1 - \ln x = -\ln x \quad \Rightarrow \quad f'(x) = -\dfrac{1}{x}$.

8. $y = \dfrac{1}{\ln x} = (\ln x)^{-1} \quad \Rightarrow \quad y' = -1(\ln x)^{-2} \cdot \dfrac{1}{x} = \dfrac{-1}{x(\ln x)^2}$

9. $g(x) = \ln(xe^{-2x}) = \ln x + \ln e^{-2x} = \ln x - 2x \quad \Rightarrow \quad g'(x) = \dfrac{1}{x} - 2$

10. $g(t) = \sqrt{1 + \ln t} \quad \Rightarrow \quad g'(t) = \frac{1}{2}(1 + \ln t)^{-1/2} \dfrac{d}{dt}(1 + \ln t) = \dfrac{1}{2\sqrt{1 + \ln t}} \cdot \dfrac{1}{t} = \dfrac{1}{2t\sqrt{1 + \ln t}}$

11. $F(t) = (\ln t)^2 \sin t \quad \Rightarrow \quad F'(t) = (\ln t)^2 \cos t + \sin t \cdot 2 \ln t \cdot \dfrac{1}{t} = \ln t \left(\ln t \cos t + \dfrac{2 \sin t}{t}\right)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $p(t) = \ln\sqrt{t^2+1} \quad \Rightarrow \quad p'(t) = \dfrac{1}{\sqrt{t^2+1}} \cdot \dfrac{d}{dt}\left(\sqrt{t^2+1}\right) = \dfrac{1}{\sqrt{t^2+1}} \cdot \dfrac{2t}{2\sqrt{t^2+1}} = \dfrac{t}{t^2+1}$

Or: $p(t) = \ln\sqrt{t^2+1} = \ln(t^2+1)^{1/2} = \frac{1}{2}\ln(t^2+1) \quad \Rightarrow \quad p'(t) = \dfrac{1}{2} \cdot \dfrac{1}{t^2+1} \cdot 2t = \dfrac{t}{t^2+1}$

13. $y = \log_8(x^2+3x) \quad \Rightarrow \quad y' = \dfrac{1}{(x^2+3x)\ln 8} \cdot \dfrac{d}{dx}(x^2+3x) = \dfrac{1}{(x^2+3x)\ln 8} \cdot (2x+3) = \dfrac{2x+3}{(x^2+3x)\ln 8}$

14. $y = \log_{10}\sec x \quad \Rightarrow \quad y' = \dfrac{1}{\sec x\,(\ln 10)} \cdot \dfrac{d}{dx}(\sec x) = \dfrac{1}{\sec x\,(\ln 10)} \cdot \sec x\,\tan x = \dfrac{\tan x}{\ln 10}$

15. $F(s) = \ln\ln s \quad \Rightarrow \quad F'(s) = \dfrac{1}{\ln s}\dfrac{d}{ds}\ln s = \dfrac{1}{\ln s} \cdot \dfrac{1}{s} = \dfrac{1}{s\ln s}$

16. $P(v) = \dfrac{\ln v}{1-v} \quad \Rightarrow \quad P'(v) = \dfrac{(1-v)(1/v) - (\ln v)(-1)}{(1-v)^2} \cdot \dfrac{v}{v} = \dfrac{1 - v + v\ln v}{v(1-v)^2}$

17. $T(z) = 2^z\log_2 z \quad \Rightarrow \quad T'(z) = 2^z\dfrac{1}{z\ln 2} + \log_2 z \cdot 2^z\ln 2 = 2^z\left(\dfrac{1}{z\ln 2} + \log_2 z\,(\ln 2)\right)$.

Note that $\log_2 z\,(\ln 2) = \dfrac{\ln z}{\ln 2}(\ln 2) = \ln z$ by the change of base formula. Thus, $T'(z) = 2^z\left(\dfrac{1}{z\ln 2} + \ln z\right)$.

18. $g(t) = \ln\dfrac{t(t^2+1)^4}{\sqrt[3]{2t-1}} = \ln t + \ln(t^2+1)^4 - \ln\sqrt[3]{2t-1} = \ln t + 4\ln(t^2+1) - \frac{1}{3}\ln(2t-1) \quad \Rightarrow$

$g'(t) = \dfrac{1}{t} + 4 \cdot \dfrac{1}{t^2+1} \cdot 2t - \dfrac{1}{3} \cdot \dfrac{1}{2t-1} \cdot 2 = \dfrac{1}{t} + \dfrac{8t}{t^2+1} - \dfrac{2}{3(2t-1)}$

19. $y = \ln\left|3 - 2x^5\right| \quad \Rightarrow \quad y' = \dfrac{1}{3-2x^5} \cdot (-10x^4) = \dfrac{-10x^4}{3-2x^5}$

20. $y = \ln(\csc x - \cot x) \quad \Rightarrow$

$y' = \dfrac{1}{\csc x - \cot x}\dfrac{d}{dx}(\csc x - \cot x) = \dfrac{1}{\csc x - \cot x}(-\csc x\,\cot x + \csc^2 x) = \dfrac{\csc x(\csc x - \cot x)}{\csc x - \cot x} = \csc x$

21. $y = \ln(e^{-x} + xe^{-x}) = \ln(e^{-x}(1+x)) = \ln(e^{-x}) + \ln(1+x) = -x + \ln(1+x) \quad \Rightarrow$

$y' = -1 + \dfrac{1}{1+x} = \dfrac{-1-x+1}{1+x} = -\dfrac{x}{1+x}$

22. $g(x) = e^{x^2\ln x} \quad \Rightarrow$

$g'(x) = e^{x^2\ln x} \cdot \dfrac{d}{dx}(x^2\ln x) = e^{x^2\ln x}\left[x^2 \cdot \dfrac{1}{x} + (\ln x) \cdot 2x\right] = e^{x^2\ln x}(x + 2x\ln x) = x\,e^{x^2\ln x}(1 + 2\ln x)$

23. $h(x) = e^{x^2+\ln x} = e^{x^2} \cdot e^{\ln x} = e^{x^2} \cdot x = xe^{x^2} \Rightarrow$

$$h'(x) = x \cdot \frac{d}{dx}\left(e^{x^2}\right) + e^{x^2} \cdot \frac{d}{dx}(x) = x \cdot e^{x^2} \cdot \frac{d}{dx}(x^2) + e^{x^2} \cdot 1 = x \cdot e^{x^2} \cdot 2x + e^{x^2}$$
$$= 2x^2e^{x^2} + e^{x^2} = e^{x^2}(2x^2+1)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $y = \ln\sqrt{\dfrac{1+2x}{1-2x}} = \ln\sqrt{1+2x} - \ln\sqrt{1-2x} = \frac{1}{2}\ln(1+2x) - \frac{1}{2}\ln(1-2x) \quad \Rightarrow$

$$y' = \frac{1}{2}\cdot\frac{1}{1+2x}\cdot 2 - \frac{1}{2}\cdot\frac{1}{1-2x}\cdot(-2) = \frac{1}{1+2x} + \frac{1}{1-2x}$$

25. $y = \ln\dfrac{x^a}{b^x} = \ln x^a - \ln b^x = a\ln x - x\ln b \quad \Rightarrow \quad y' = a\cdot\dfrac{1}{x} - \ln b = \dfrac{a}{x} - \ln b$

26. $y = \log_2(x\log_5 x) \quad \Rightarrow$

$$y' = \frac{1}{(x\log_5 x)(\ln 2)}\frac{d}{dx}(x\log_5 x) = \frac{1}{(x\log_5 x)(\ln 2)}\left(x\cdot\frac{1}{x\ln 5} + \log_5 x\right) = \frac{1}{(x\log_5 x)(\ln 5)(\ln 2)} + \frac{1}{x(\ln 2)}.$$

Note that $\log_5 x(\ln 5) = \dfrac{\ln x}{\ln 5}(\ln 5) = \ln x$ by the change of base formula. Thus, $y' = \dfrac{1}{x\ln x\ln 2} + \dfrac{1}{x\ln 2} = \dfrac{1+\ln x}{x\ln x\ln 2}$.

27. $\dfrac{d}{dx}\ln\left(x+\sqrt{x^2+1}\right) = \dfrac{1}{x+\sqrt{x^2+1}}\cdot\dfrac{d}{dx}\left(x+\sqrt{x^2+1}\right) = \dfrac{1}{x+\sqrt{x^2+1}}\cdot\left(1+\dfrac{2x}{2\sqrt{x^2+1}}\right)$

$$= \frac{1}{x+\sqrt{x^2+1}}\cdot\left(\frac{\sqrt{x^2+1}}{\sqrt{x^2+1}} + \frac{x}{\sqrt{x^2+1}}\right) = \frac{1}{x+\sqrt{x^2+1}}\cdot\left(\frac{x+\sqrt{x^2+1}}{\sqrt{x^2+1}}\right) = \frac{1}{\sqrt{x^2+1}}$$

28. $\dfrac{d}{dx}\ln\sqrt{\dfrac{1-\cos x}{1+\cos x}} = \dfrac{d}{dx}\left(\ln\sqrt{1-\cos x} - \ln\sqrt{1+\cos x}\right) = \dfrac{d}{dx}\left[\dfrac{1}{2}\ln(1-\cos x) - \dfrac{1}{2}\ln(1+\cos x)\right]$

$$= \frac{1}{2}\cdot\frac{1}{1-\cos x}\cdot\sin x - \frac{1}{2}\cdot\frac{1}{1+\cos x}\cdot(-\sin x)$$

$$= \frac{1}{2}\left(\frac{\sin x}{1-\cos x} + \frac{\sin x}{1+\cos x}\right) = \frac{1}{2}\left[\frac{\sin x\,(1+\cos x) + \sin x\,(1-\cos x)}{(1-\cos x)(1+\cos x)}\right]$$

$$= \frac{1}{2}\left(\frac{\sin x + \sin x\,\cos x + \sin x - \sin x\,\cos x}{1-\cos^2 x}\right) = \frac{1}{2}\left(\frac{2\sin x}{\sin^2 x}\right) = \frac{1}{\sin x} = \csc x$$

29. $y = \sqrt{x}\,\ln x \quad \Rightarrow \quad y' = \sqrt{x}\cdot\dfrac{1}{x} + (\ln x)\dfrac{1}{2\sqrt{x}} = \dfrac{2+\ln x}{2\sqrt{x}} \quad \Rightarrow$

$$y'' = \frac{2\sqrt{x}\,(1/x) - (2+\ln x)(1/\sqrt{x}\,)}{(2\sqrt{x}\,)^2} = \frac{2/\sqrt{x} - (2+\ln x)(1/\sqrt{x}\,)}{4x} = \frac{2-(2+\ln x)}{\sqrt{x}(4x)} = -\frac{\ln x}{4x\sqrt{x}}$$

30. $y = \dfrac{\ln x}{1+\ln x} \quad \Rightarrow \quad y' = \dfrac{(1+\ln x)(1/x) - (\ln x)(1/x)}{(1+\ln x)^2} = \dfrac{1}{x(1+\ln x)^2} \quad \Rightarrow$

$$y'' = -\frac{\dfrac{d}{dx}[x(1+\ln x)^2]}{[x(1+\ln x)^2]^2} \quad \text{[Reciprocal Rule]} \quad = -\frac{x\cdot 2(1+\ln x)\cdot(1/x) + (1+\ln x)^2}{x^2(1+\ln x)^4}$$

$$= -\frac{(1+\ln x)[2+(1+\ln x)]}{x^2(1+\ln x)^4} = -\frac{3+\ln x}{x^2(1+\ln x)^3}$$

31. $y = \ln|\sec x| \quad \Rightarrow \quad y' = \dfrac{1}{\sec x}\dfrac{d}{dx}\sec x = \dfrac{1}{\sec x}\sec x\,\tan x = \tan x \quad \Rightarrow \quad y'' = \sec^2 x$

32. $y = \ln(1+\ln x) \quad \Rightarrow \quad y' = \dfrac{1}{1+\ln x}\cdot\dfrac{1}{x} = \dfrac{1}{x(1+\ln x)} \quad \Rightarrow$

$$y'' = -\frac{\dfrac{d}{dx}[x(1+\ln x)]}{[x(1+\ln x)]^2} \quad \text{[Reciprocal Rule]} \quad = -\frac{x(1/x) + (1+\ln x)(1)}{x^2(1+\ln x)^2} = -\frac{1+1+\ln x}{x^2(1+\ln x)^2} = -\frac{2+\ln x}{x^2(1+\ln x)^2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. $f(x) = \dfrac{x}{1 - \ln(x-1)} \quad \Rightarrow$

$$f'(x) = \frac{[1-\ln(x-1)]\cdot 1 - x\cdot\dfrac{-1}{x-1}}{[1-\ln(x-1)]^2} = \frac{\dfrac{(x-1)[1-\ln(x-1)]+x}{x-1}}{[1-\ln(x-1)]^2} = \frac{x-1-(x-1)\ln(x-1)+x}{(x-1)[1-\ln(x-1)]^2}$$
$$= \frac{2x-1-(x-1)\ln(x-1)}{(x-1)[1-\ln(x-1)]^2}$$

$\mathrm{Dom}(f) = \{x \mid x-1>0 \quad \text{and} \quad 1-\ln(x-1)\neq 0\} = \{x \mid x>1 \quad \text{and} \quad \ln(x-1)\neq 1\}$
$= \left\{x \mid x>1 \quad \text{and} \quad x-1\neq e^1\right\} = \{x \mid x>1 \quad \text{and} \quad x\neq 1+e\} = (1,1+e)\cup(1+e,\infty)$

34. $f(x) = \sqrt{2+\ln x} = (2+\ln x)^{1/2} \quad \Rightarrow \quad f'(x) = \dfrac{1}{2}(2+\ln x)^{-1/2}\cdot\dfrac{1}{x} = \dfrac{1}{2x\sqrt{2+\ln x}}$

$\mathrm{Dom}(f) = \{x \mid 2+\ln x\geq 0\} = \{x \mid \ln x\geq -2\} = \{x \mid x\geq e^{-2}\} = [e^{-2},\infty)$.

35. $f(x) = \ln(x^2-2x) \quad \Rightarrow \quad f'(x) = \dfrac{1}{x^2-2x}(2x-2) = \dfrac{2(x-1)}{x(x-2)}$.

$\mathrm{Dom}(f) = \{x \mid x(x-2)>0\} = (-\infty,0)\cup(2,\infty)$.

36. $f(x) = \ln\ln\ln x \quad \Rightarrow \quad f'(x) = \dfrac{1}{\ln\ln x}\cdot\dfrac{1}{\ln x}\cdot\dfrac{1}{x}$.

$\mathrm{Dom}(f) = \{x \mid \ln\ln x>0\} = \{x \mid \ln x>1\} = \{x \mid x>e\} = (e,\infty)$.

37. $f(x) = \ln(x+\ln x) \quad \Rightarrow \quad f'(x) = \dfrac{1}{x+\ln x}\dfrac{d}{dx}(x+\ln x) = \dfrac{1}{x+\ln x}\left(1+\dfrac{1}{x}\right)$.

Substitute 1 for x to get $f'(1) = \dfrac{1}{1+\ln 1}\left(1+\dfrac{1}{1}\right) = \dfrac{1}{1+0}(1+1) = 1\cdot 2 = 2$.

38. $f(x) = \cos(\ln x^2) \quad \Rightarrow \quad f'(x) = -\sin(\ln x^2)\dfrac{d}{dx}\ln x^2 = -\sin(\ln x^2)\dfrac{1}{x^2}(2x) = -\dfrac{2\sin(\ln x^2)}{x}$.

Substitute 1 for x to get $f'(1) = -\dfrac{2\sin(\ln 1^2)}{1} = -2\sin 0 = 0$.

39. $y = \ln(x^2-3x+1) \quad \Rightarrow \quad y' = \dfrac{1}{x^2-3x+1}\cdot(2x-3) \quad \Rightarrow \quad y'(3) = \frac{1}{1}\cdot 3 = 3$, so an equation of a tangent line at $(3,0)$ is $y-0 = 3(x-3)$, or $y = 3x-9$.

40. $y = x^2\ln x \quad \Rightarrow \quad y' = x^2\cdot\dfrac{1}{x} + (\ln x)(2x) \quad \Rightarrow \quad y'(1) = 1+0 = 1$, so an equation of a tangent line at $(1,0)$ is $y-0 = 1(x-1)$, or $y = x-1$.

41. $f(x) = \sin x + \ln x \quad \Rightarrow \quad f'(x) = \cos x + 1/x$.

This is reasonable, because the graph shows that f increases when f' is positive, and $f'(x) = 0$ when f has a horizontal tangent.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. $y = \dfrac{\ln x}{x} \quad \Rightarrow \quad y' = \dfrac{x(1/x) - \ln x}{x^2} = \dfrac{1 - \ln x}{x^2}$.

$y'(1) = \dfrac{1-0}{1^2} = 1$ and $y'(e) = \dfrac{1-1}{e^2} = 0 \quad \Rightarrow$ equations of tangent lines are $y - 0 = 1(x-1)$ or $y = x - 1$ and $y - 1/e = 0(x - e)$ or $y = 1/e$.

43. $f(x) = cx + \ln(\cos x) \quad \Rightarrow \quad f'(x) = c + \dfrac{1}{\cos x} \cdot (-\sin x) = c - \tan x$.

$f'(\frac{\pi}{4}) = 6 \quad \Rightarrow \quad c - \tan\frac{\pi}{4} = 6 \quad \Rightarrow \quad c - 1 = 6 \quad \Rightarrow \quad c = 7$.

44. $f(x) = \log_b(3x^2 - 2) \quad \Rightarrow \quad f'(x) = \dfrac{1}{(3x^2-2)\ln b} \cdot 6x$. $f'(1) = 3 \quad \Rightarrow \quad \dfrac{1}{\ln b} \cdot 6 = 3 \quad \Rightarrow \quad 2 = \ln b \quad \Rightarrow \quad b = e^2$.

45. $y = (x^2+2)^2(x^4+4)^4 \quad \Rightarrow \quad \ln y = \ln[(x^2+2)^2(x^4+4)^4] \quad \Rightarrow \quad \ln y = 2\ln(x^2+2) + 4\ln(x^4+4) \quad \Rightarrow$

$\dfrac{1}{y}y' = 2 \cdot \dfrac{1}{x^2+2} \cdot 2x + 4 \cdot \dfrac{1}{x^4+4} \cdot 4x^3 \quad \Rightarrow \quad y' = y\left(\dfrac{4x}{x^2+2} + \dfrac{16x^3}{x^4+4}\right) \quad \Rightarrow$

$y' = (x^2+2)^2(x^4+4)^4\left(\dfrac{4x}{x^2+2} + \dfrac{16x^3}{x^4+4}\right)$

46. $y = \dfrac{e^{-x}\cos^2 x}{x^2+x+1} \quad \Rightarrow \quad \ln y = \ln \dfrac{e^{-x}\cos^2 x}{x^2+x+1} \quad \Rightarrow$

$\ln y = \ln e^{-x} + \ln|\cos x|^2 - \ln(x^2+x+1) = -x + 2\ln|\cos x| - \ln(x^2+x+1) \quad \Rightarrow$

$\dfrac{1}{y}y' = -1 + 2 \cdot \dfrac{1}{\cos x}(-\sin x) - \dfrac{1}{x^2+x+1}(2x+1) \quad \Rightarrow \quad y' = y\left(-1 - 2\tan x - \dfrac{2x+1}{x^2+x+1}\right) \quad \Rightarrow$

$y' = -\dfrac{e^{-x}\cos^2 x}{x^2+x+1}\left(1 + 2\tan x + \dfrac{2x+1}{x^2+x+1}\right)$

47. $y = \sqrt{\dfrac{x-1}{x^4+1}} \quad \Rightarrow \quad \ln y = \ln\left(\dfrac{x-1}{x^4+1}\right)^{1/2} \quad \Rightarrow \quad \ln y = \dfrac{1}{2}\ln(x-1) - \dfrac{1}{2}\ln(x^4+1) \quad \Rightarrow$

$\dfrac{1}{y}y' = \dfrac{1}{2}\dfrac{1}{x-1} - \dfrac{1}{2}\dfrac{1}{x^4+1} \cdot 4x^3 \quad \Rightarrow \quad y' = y\left(\dfrac{1}{2(x-1)} - \dfrac{2x^3}{x^4+1}\right) \quad \Rightarrow \quad y' = \sqrt{\dfrac{x-1}{x^4+1}}\left(\dfrac{1}{2x-2} - \dfrac{2x^3}{x^4+1}\right)$

48. $y = \sqrt{x}\,e^{x^2-x}(x+1)^{2/3} \quad \Rightarrow \quad \ln y = \ln\left[x^{1/2}e^{x^2-x}(x+1)^{2/3}\right] \quad \Rightarrow$

$\ln y = \frac{1}{2}\ln x + (x^2 - x) + \frac{2}{3}\ln(x+1) \quad \Rightarrow \quad \dfrac{1}{y}y' = \dfrac{1}{2} \cdot \dfrac{1}{x} + 2x - 1 + \dfrac{2}{3} \cdot \dfrac{1}{x+1} \quad \Rightarrow$

$y' = y\left(\dfrac{1}{2x} + 2x - 1 + \dfrac{2}{3x+3}\right) \quad \Rightarrow \quad y' = \sqrt{x}\,e^{x^2-x}(x+1)^{2/3}\left(\dfrac{1}{2x} + 2x - 1 + \dfrac{2}{3x+3}\right)$

49. $y = x^x \quad \Rightarrow \quad \ln y = \ln x^x \quad \Rightarrow \quad \ln y = x\ln x \quad \Rightarrow \quad y'/y = x(1/x) + (\ln x) \cdot 1 \quad \Rightarrow \quad y' = y(1 + \ln x) \quad \Rightarrow$

$y' = x^x(1 + \ln x)$

50. $y = x^{1/x} \quad \Rightarrow \quad \ln y = \dfrac{1}{x}\ln x \quad \Rightarrow \quad \dfrac{y'}{y} = -\dfrac{1}{x^2}\ln x + \dfrac{1}{x}\left(\dfrac{1}{x}\right) \quad \Rightarrow \quad y' = x^{1/x}\dfrac{1 - \ln x}{x^2}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. $y = x^{\sin x} \quad \Rightarrow \quad \ln y = \ln x^{\sin x} \quad \Rightarrow \quad \ln y = \sin x \ln x \quad \Rightarrow \quad \dfrac{y'}{y} = (\sin x) \cdot \dfrac{1}{x} + (\ln x)(\cos x) \quad \Rightarrow$

$y' = y\left(\dfrac{\sin x}{x} + \ln x \cos x\right) \quad \Rightarrow \quad y' = x^{\sin x}\left(\dfrac{\sin x}{x} + \ln x \, \cos x\right)$

52. $y = \left(\sqrt{x}\right)^x \quad \Rightarrow \quad \ln y = \ln\left(\sqrt{x}\right)^x \quad \Rightarrow \quad \ln y = x \ln x^{1/2} \quad \Rightarrow \quad \ln y = \frac{1}{2} x \ln x \quad \Rightarrow \quad \dfrac{1}{y}\, y' = \dfrac{1}{2} x \cdot \dfrac{1}{x} + \ln x \cdot \dfrac{1}{2} \quad \Rightarrow$

$y' = y\left(\frac{1}{2} + \frac{1}{2} \ln x\right) \quad \Rightarrow \quad y' = \frac{1}{2}\left(\sqrt{x}\right)^x (1 + \ln x)$

53. $y = (\cos x)^x \quad \Rightarrow \quad \ln y = \ln(\cos x)^x \quad \Rightarrow \quad \ln y = x \ln \cos x \quad \Rightarrow \quad \dfrac{1}{y}\, y' = x \cdot \dfrac{1}{\cos x} \cdot (-\sin x) + \ln \cos x \cdot 1 \quad \Rightarrow$

$y' = y\left(\ln \cos x - \dfrac{x \sin x}{\cos x}\right) \quad \Rightarrow \quad y' = (\cos x)^x (\ln \cos x - x \tan x)$

54. $y = (\sin x)^{\ln x} \quad \Rightarrow \quad \ln y = \ln(\sin x)^{\ln x} \quad \Rightarrow \quad \ln y = \ln x \cdot \ln \sin x \quad \Rightarrow \quad \dfrac{1}{y}\, y' = \ln x \cdot \dfrac{1}{\sin x} \cdot \cos x + \ln \sin x \cdot \dfrac{1}{x} \quad \Rightarrow$

$y' = y\left(\ln x \cdot \dfrac{\cos x}{\sin x} + \dfrac{\ln \sin x}{x}\right) \quad \Rightarrow \quad y' = (\sin x)^{\ln x}\left(\ln x \cot x + \dfrac{\ln \sin x}{x}\right)$

55. $y = x^{\ln x} \quad \Rightarrow \quad \ln y = \ln x \ln x = (\ln x)^2 \quad \Rightarrow \quad \dfrac{y'}{y} = 2 \ln x \left(\dfrac{1}{x}\right) \quad \Rightarrow \quad y' = x^{\ln x}\left(\dfrac{2 \ln x}{x}\right)$

56. $y = (\ln x)^{\cos x} \quad \Rightarrow \quad \ln y = \cos x \ln(\ln x) \quad \Rightarrow \quad \dfrac{y'}{y} = \cos x \cdot \dfrac{1}{\ln x} \cdot \dfrac{1}{x} + (\ln \ln x)(-\sin x) \quad \Rightarrow$

$y' = (\ln x)^{\cos x}\left(\dfrac{\cos x}{x \ln x} - \sin x \ln \ln x\right)$

57. $y = \ln(x^2 + y^2) \quad \Rightarrow \quad y' = \dfrac{1}{x^2 + y^2} \dfrac{d}{dx}(x^2 + y^2) \quad \Rightarrow \quad y' = \dfrac{2x + 2yy'}{x^2 + y^2} \quad \Rightarrow \quad x^2 y' + y^2 y' = 2x + 2yy' \quad \Rightarrow$

$x^2 y' + y^2 y' - 2yy' = 2x \quad \Rightarrow \quad (x^2 + y^2 - 2y) y' = 2x \quad \Rightarrow \quad y' = \dfrac{2x}{x^2 + y^2 - 2y}$

58. $x^y = y^x \quad \Rightarrow \quad y \ln x = x \ln y \quad \Rightarrow \quad y \cdot \dfrac{1}{x} + (\ln x) \cdot y' = x \cdot \dfrac{1}{y} \cdot y' + \ln y \quad \Rightarrow \quad y' \ln x - \dfrac{x}{y}\, y' = \ln y - \dfrac{y}{x} \quad \Rightarrow$

$y' = \dfrac{\ln y - y/x}{\ln x - x/y}$

59. $f(x) = \ln(x - 1) \quad \Rightarrow \quad f'(x) = \dfrac{1}{(x - 1)} = (x - 1)^{-1} \quad \Rightarrow \quad f''(x) = -(x - 1)^{-2} \quad \Rightarrow \quad f'''(x) = 2(x - 1)^{-3} \quad \Rightarrow$

$f^{(4)}(x) = -2 \cdot 3(x - 1)^{-4} \quad \Rightarrow \quad \cdots \quad \Rightarrow \quad f^{(n)}(x) = (-1)^{n-1} \cdot 2 \cdot 3 \cdot 4 \cdot \cdots \cdot (n - 1)(x - 1)^{-n} = (-1)^{n-1} \dfrac{(n - 1)!}{(x - 1)^n}$

60. $y = x^8 \ln x$, so $D^9 y = D^8 y' = D^8(8x^7 \ln x + x^7)$. But the eighth derivative of x^7 is 0, so we now have

$D^8(8x^7 \ln x) = D^7(8 \cdot 7x^6 \ln x + 8x^6) = D^7(8 \cdot 7x^6 \ln x) = D^6(8 \cdot 7 \cdot 6x^5 \ln x) = \cdots = D(8!\, x^0 \ln x) = 8!/x.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. If $f(x) = \ln(1+x)$, then $f'(x) = \dfrac{1}{1+x}$, so $f'(0) = 1$.

Thus, $\displaystyle\lim_{x\to 0} \frac{\ln(1+x)}{x} = \lim_{x\to 0} \frac{f(x)}{x} = \lim_{x\to 0} \frac{f(x)-f(0)}{x-0} = f'(0) = 1.$

62. Let $m = n/x$. Then $n = xm$, and as $n \to \infty$, $m \to \infty$.

Therefore, $\displaystyle\lim_{n\to\infty}\left(1+\frac{x}{n}\right)^n = \lim_{m\to\infty}\left(1+\frac{1}{m}\right)^{mx} = \left[\lim_{m\to\infty}\left(1+\frac{1}{m}\right)^m\right]^x = e^x$ by Equation 6.

63. $f(x) = \sin^{-1}(5x) \Rightarrow f'(x) = \dfrac{1}{\sqrt{1-(5x)^2}} \cdot \dfrac{d}{dx}(5x) = \dfrac{5}{\sqrt{1-25x^2}}$

64. $g(x) = \sec^{-1}(e^x) \Rightarrow g'(x) = \dfrac{1}{e^x\sqrt{(e^x)^2-1}} \cdot \dfrac{d}{dx}(e^x) = \dfrac{1}{e^x\sqrt{e^{2x}-1}} \cdot e^x = \dfrac{1}{\sqrt{e^{2x}-1}}$

65. $y = \tan^{-1}\sqrt{x-1} \quad \Rightarrow$

$$y' = \frac{1}{1+\left(\sqrt{x-1}\right)^2} \cdot \frac{d}{dt}\left(\sqrt{x-1}\right) = \frac{1}{1+(x-1)} \cdot \frac{1}{2\sqrt{x-1}} = \frac{1}{x} \cdot \frac{1}{2\sqrt{x-1}} = \frac{1}{2x\sqrt{x-1}}$$

66. $y = \tan^{-1}(x^2) \quad \Rightarrow \quad y' = \dfrac{1}{1+(x^2)^2} \cdot \dfrac{d}{dx}(x^2) = \dfrac{1}{1+x^4} \cdot 2x = \dfrac{2x}{1+x^4}$

67. $y = (\tan^{-1}x)^2 \quad \Rightarrow \quad y' = 2(\tan^{-1}x)^1 \cdot \dfrac{d}{dx}(\tan^{-1}x) = 2\tan^{-1}x \cdot \dfrac{1}{1+x^2} = \dfrac{2\tan^{-1}x}{1+x^2}$

68. $g(x) = \arccos\sqrt{x} \quad \Rightarrow \quad g'(x) = -\dfrac{1}{\sqrt{1-\left(\sqrt{x}\right)^2}}\dfrac{d}{dx}\sqrt{x} = -\dfrac{1}{\sqrt{1-x}}\left(\dfrac{1}{2}x^{-1/2}\right) = -\dfrac{1}{2\sqrt{x}\sqrt{1-x}}$

69. $h(x) = (\arcsin x)\ln x \quad \Rightarrow \quad h'(x) = (\arcsin x) \cdot \dfrac{1}{x} + (\ln x) \cdot \dfrac{1}{\sqrt{1-x^2}} = \dfrac{\arcsin x}{x} + \dfrac{\ln x}{\sqrt{1-x^2}}$

70. $g(t) = \ln(\arctan(t^4)) \quad \Rightarrow$

$$\begin{aligned} g'(t) &= \frac{1}{\arctan(t^4)} \cdot \frac{d}{dt}(\arctan(t^4)) = \frac{1}{\arctan(t^4)} \cdot \frac{1}{1+(t^4)^2} \cdot \frac{d}{dt}(t^4) \\ &= \frac{1}{\arctan(t^4)} \cdot \frac{1}{1+t^8} \cdot 4t^3 = \frac{4t^3}{(1+t^8)\arctan(t^4)} \end{aligned}$$

71. $f(z) = e^{\arcsin(z^2)} \quad \Rightarrow \quad f'(z) = e^{\arcsin(z^2)} \cdot \dfrac{d}{dz}[\arcsin(z^2)] = e^{\arcsin(z^2)} \cdot \dfrac{1}{\sqrt{1-(z^2)^2}} \cdot 2z = \dfrac{2ze^{\arcsin(z^2)}}{\sqrt{1-z^4}}$

72. $y = \tan^{-1}\left(x - \sqrt{1+x^2}\right) \quad \Rightarrow$

$$\begin{aligned} y' &= \frac{1}{1+\left(x-\sqrt{x^2+1}\right)^2}\left(1-\frac{x}{\sqrt{x^2+1}}\right) = \frac{1}{1+x^2-2x\sqrt{x^2+1}+x^2+1}\left(\frac{\sqrt{x^2+1}-x}{\sqrt{x^2+1}}\right) \\ &= \frac{\sqrt{x^2+1}-x}{2\left(1+x^2-x\sqrt{x^2+1}\right)\sqrt{x^2+1}} = \frac{\sqrt{x^2+1}-x}{2\left[\sqrt{x^2+1}\,(1+x^2)-x(x^2+1)\right]} = \frac{\sqrt{x^2+1}-x}{2\left[(1+x^2)\left(\sqrt{x^2+1}-x\right)\right]} \\ &= \frac{1}{2(1+x^2)} \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

73. $h(t) = \cot^{-1}(t) + \cot^{-1}(1/t) \quad \Rightarrow$

$$h'(t) = -\frac{1}{1+t^2} - \frac{1}{1+(1/t)^2} \cdot \frac{d}{dt}\frac{1}{t} = -\frac{1}{1+t^2} - \frac{t^2}{t^2+1} \cdot \left(-\frac{1}{t^2}\right) = -\frac{1}{1+t^2} + \frac{1}{t^2+1} = 0.$$

Note that this makes sense because $h(t) = \dfrac{\pi}{2}$ for $t > 0$ and $h(t) = \dfrac{3\pi}{2}$ for $t < 0$.

74. $R(t) = \arcsin(1/t) \quad \Rightarrow$

$$R'(t) = \frac{1}{\sqrt{1-(1/t)^2}}\,\frac{d}{dt}\frac{1}{t} = \frac{1}{\sqrt{1-1/t^2}}\left(-\frac{1}{t^2}\right) = -\frac{1}{\sqrt{1-1/t^2}}\,\frac{1}{\sqrt{t^4}} = -\frac{1}{\sqrt{t^4-t^2}}$$

$$= -\frac{1}{\sqrt{t^2(t^2-1)}} = -\frac{1}{|\,t\,|\sqrt{t^2-1}}$$

75. $y = x\sin^{-1}x + \sqrt{1-x^2} \quad \Rightarrow$

$$y' = x \cdot \frac{1}{\sqrt{1-x^2}} + (\sin^{-1}x)(1) + \frac{1}{2}(1-x^2)^{-1/2}(-2x) = \frac{x}{\sqrt{1-x^2}} + \sin^{-1}x - \frac{x}{\sqrt{1-x^2}} = \sin^{-1}x$$

76. $y = \cos^{-1}(\sin^{-1}t) \quad \Rightarrow \quad y' = -\dfrac{1}{\sqrt{1-(\sin^{-1}t)^2}} \cdot \dfrac{d}{dt}\sin^{-1}t = -\dfrac{1}{\sqrt{1-(\sin^{-1}t)^2}} \cdot \dfrac{1}{\sqrt{1-t^2}}$

77. $y = \tan^{-1}\left(\dfrac{x}{a}\right) + \ln\sqrt{\dfrac{x-a}{x+a}} = \tan^{-1}\left(\dfrac{x}{a}\right) + \dfrac{1}{2}\ln\left(\dfrac{x-a}{x+a}\right) \quad \Rightarrow$

$$y' = \frac{1}{1+\left(\frac{x}{a}\right)^2} \cdot \frac{1}{a} + \frac{1}{2} \cdot \frac{1}{\frac{x-a}{x+a}} \cdot \frac{(x+a)\cdot 1 - (x-a)\cdot 1}{(x+a)^2} = \frac{1}{a+\frac{x^2}{a}} + \frac{1}{2} \cdot \frac{x+a}{x-a} \cdot \frac{2a}{(x+a)^2}$$

$$= \frac{1}{a+\frac{x^2}{a}} \cdot \frac{a}{a} + \frac{a}{(x-a)(x+a)} = \frac{a}{x^2+a^2} + \frac{a}{x^2-a^2}$$

78. $y = \arctan\sqrt{\dfrac{1-x}{1+x}} = \arctan\left(\dfrac{1-x}{1+x}\right)^{1/2} \quad \Rightarrow$

$$y' = \frac{1}{1+\left(\sqrt{\frac{1-x}{1+x}}\right)^2} \cdot \frac{d}{dx}\left(\frac{1-x}{1+x}\right)^{1/2} = \frac{1}{1+\frac{1-x}{1+x}} \cdot \frac{1}{2}\left(\frac{1-x}{1+x}\right)^{-1/2} \cdot \frac{(1+x)(-1)-(1-x)(1)}{(1+x)^2}$$

$$= \frac{1}{\frac{1+x}{1+x}+\frac{1-x}{1+x}} \cdot \frac{1}{2}\left(\frac{1+x}{1-x}\right)^{1/2} \cdot \frac{-2}{(1+x)^2} = \frac{1+x}{2} \cdot \frac{1}{2} \cdot \frac{(1+x)^{1/2}}{(1-x)^{1/2}} \cdot \frac{-2}{(1+x)^2}$$

$$= \frac{-1}{2(1-x)^{1/2}(1+x)^{1/2}} = \frac{-1}{2\sqrt{1-x^2}}$$

79. $f(x) = \sqrt{1-x^2}\arcsin x \quad \Rightarrow \quad f'(x) = \sqrt{1-x^2} \cdot \dfrac{1}{\sqrt{1-x^2}} + \arcsin x \cdot \dfrac{1}{2}\left(1-x^2\right)^{-1/2}(-2x) = 1 - \dfrac{x\arcsin x}{\sqrt{1-x^2}}$

Note that $f' = 0$ where the graph of f has a horizontal tangent. Also note that f' is negative when f is decreasing and f' is positive when f is increasing.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

80. $f(x) = \arctan(x^2 - x) \quad \Rightarrow \quad f'(x) = \dfrac{1}{1 + (x^2 - x)^2} \cdot \dfrac{d}{dx}(x^2 - x) = \dfrac{2x - 1}{1 + (x^2 - x)^2}$

Note that $f' = 0$ where the graph of f has a horizontal tangent. Also note that f' is negative when f is decreasing and f' is positive when f is increasing.

81. Let $y = \cos^{-1} x$. Then $\cos y = x$ and $0 \le y \le \pi \quad \Rightarrow \quad -\sin y \dfrac{dy}{dx} = 1 \quad \Rightarrow$

$\dfrac{dy}{dx} = -\dfrac{1}{\sin y} = -\dfrac{1}{\sqrt{1 - \cos^2 y}} = -\dfrac{1}{\sqrt{1 - x^2}}$. [Note that $\sin y \ge 0$ for $0 \le y \le \pi$.]

82. (a) Let $y = \sec^{-1} x$. Then $\sec y = x$ and $y \in \left(0, \frac{\pi}{2}\right] \cup \left(\pi, \frac{3\pi}{2}\right]$. Differentiate with respect to x: $\sec y \tan y \left(\dfrac{dy}{dx}\right) = 1 \quad \Rightarrow$

$\dfrac{dy}{dx} = \dfrac{1}{\sec y \tan y} = \dfrac{1}{\sec y \sqrt{\sec^2 y - 1}} = \dfrac{1}{x\sqrt{x^2 - 1}}$. Note that $\tan^2 y = \sec^2 y - 1 \quad \Rightarrow \quad \tan y = \sqrt{\sec^2 y - 1}$

since $\tan y > 0$ when $0 < y < \frac{\pi}{2}$ or $\pi < y < \frac{3\pi}{2}$.

(b) $y = \sec^{-1} x \quad \Rightarrow \quad \sec y = x \quad \Rightarrow \quad \sec y \tan y \dfrac{dy}{dx} = 1 \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{1}{\sec y \tan y}$. Now $\tan^2 y = \sec^2 y - 1 = x^2 - 1$,

so $\tan y = \pm\sqrt{x^2 - 1}$. For $y \in \left[0, \frac{\pi}{2}\right)$, $x \ge 1$, so $\sec y = x = |x|$ and $\tan y \ge 0 \quad \Rightarrow$

$\dfrac{dy}{dx} = \dfrac{1}{x\sqrt{x^2 - 1}} = \dfrac{1}{|x|\sqrt{x^2 - 1}}$. For $y \in \left(\frac{\pi}{2}, \pi\right]$, $x \le -1$, so $|x| = -x$ and $\tan y = -\sqrt{x^2 - 1} \quad \Rightarrow$

$\dfrac{dy}{dx} = \dfrac{1}{\sec y \tan y} = \dfrac{1}{x\left(-\sqrt{x^2 - 1}\right)} = \dfrac{1}{(-x)\sqrt{x^2 - 1}} = \dfrac{1}{|x|\sqrt{x^2 - 1}}$.

83. If $y = f^{-1}(x)$, then $f(y) = x$. Differentiating implicitly with respect to x and remembering that y is a function of x,

we get $f'(y)\dfrac{dy}{dx} = 1$, so $\dfrac{dy}{dx} = \dfrac{1}{f'(y)} \quad \Rightarrow \quad \left(f^{-1}\right)'(x) = \dfrac{1}{f'(f^{-1}(x))}$.

84. $f(4) = 5 \quad \Rightarrow \quad f^{-1}(5) = 4$. By Exercise 83, $\left(f^{-1}\right)'(5) = \dfrac{1}{f'(f^{-1}(5))} = \dfrac{1}{f'(4)} = \dfrac{1}{2/3} = \dfrac{3}{2}$.

85. $f(x) = x + e^x \quad \Rightarrow \quad f'(x) = 1 + e^x$. Observe that $f(0) = 1$, so that $f^{-1}(1) = 0$. By Exercise 83, we have

$(f^{-1})'(1) = \dfrac{1}{f'(f^{-1}(1))} = \dfrac{1}{f'(0)} = \dfrac{1}{1 + e^0} = \dfrac{1}{1 + 1} = \dfrac{1}{2}$.

86. $f(x) = x^3 + 3\sin x + 2\cos x \quad \Rightarrow \quad f'(x) = 3x^2 + 3\cos x - 2\sin x$. Observe that $f(0) = 2$, so that $f^{-1}(2) = 0$.

By Exercise 83, we have $(f^{-1})'(2) = \dfrac{1}{f'(f^{-1}(2))} = \dfrac{1}{f'(0)} = \dfrac{1}{3(0)^2 + 3\cos 0 - 2\sin 0} = \dfrac{1}{3(1)} = \dfrac{1}{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

87. $h = f^g \quad \Rightarrow \quad \ln h = \ln f^g \quad \Rightarrow \quad \ln h = g \ln f \quad \Rightarrow \quad \dfrac{1}{h}\, h' = g \cdot \dfrac{1}{f}\, f' + (\ln f)\, g' \quad \Rightarrow$

$h' = h\left(g \cdot \dfrac{1}{f}\, f' + (\ln f)\, g'\right) = f^g\left(g \cdot \dfrac{1}{f}\, f' + (\ln f)\, g'\right) = g \cdot \dfrac{f^g}{f} \cdot f' + (\ln f) \cdot f^g \cdot g' \quad \Rightarrow$

$h' = g \cdot f^{g-1} \cdot f' + (\ln f) \cdot f^g \cdot g'$

88. (a) With $h(x) = x^3$, we have $f(x) = x$ and $g(x) = 3$ in the formula in Exercise 87. That formula gives

$h'(x) = 3 \cdot x^{3-1} \cdot 1 + (\ln x) \cdot x^3 \cdot 0 = 3x^2$.

(b) With $h(x) = 3^x$, we have $f(x) = 3$ and $g(x) = x$ in the formula in Exercise 87. That formula gives

$h'(x) = x \cdot 3^{x-1} \cdot 0 + (\ln 3) \cdot 3^x \cdot 1 = 3^x(\ln 3)$.

(c) With $h(x) = (\sin x)^x$, we have $f(x) = \sin x$ and $g(x) = x$ in the formula in Exercise 87. That formula gives

$h'(x) = x \cdot (\sin x)^{x-1} \cdot \cos x + (\ln \sin x) \cdot (\sin x)^x \cdot 1 = x \cos x\, (\sin x)^{x-1} + (\sin x)^x \ln \sin x$. Further simplification gives $x \cos x\, (\sin x)^{x-1} + (\sin x)^x \ln \sin x = \dfrac{x \cos x\, (\sin x)^x}{\sin x} + (\sin x)^x \ln \sin x = (\sin x)^x\, (x \cot x + \ln \sin x)$.

3.7 Rates of Change in the Natural and Social Sciences

1. (a) $s = f(t) = t^3 - 9t^2 + 24t$ (in feet) $\quad \Rightarrow \quad v(t) = f'(t) = 3t^2 - 18t + 24$ (in ft/s)

(b) $v(1) = 3(1)^2 - 18(1) + 24 = 9$ ft/s

(c) The particle is at rest when $v(t) = 0$: $3t^2 - 18t + 24 = 0 \quad \Leftrightarrow \quad 3(t-2)(t-4) = 0 \quad \Leftrightarrow \quad t = 2$ s or $t = 4$ s.

(d) The particle is moving in the positive direction when $v(t) > 0$. $3(t-2)(t-4) > 0 \quad \Rightarrow \quad 0 \le t < 2$ or $t > 4$.

(e) Because the particle changes direction when $t = 2$ and $t = 4$, we need to calculate the distances traveled in the intervals $[0, 2]$, $[2, 4]$, and $[4, 6]$ separately.

$$|f(2) - f(0)| = |20 - 0| = 20$$
$$|f(4) - f(2)| = |16 - 20| = 4$$
$$|f(6) - f(4)| = |36 - 16| = 20$$

The total distance is $20 + 4 + 20 = 44$ ft.

(f)

(g) $v(t) = 3t^2 - 18t + 24 \quad \Rightarrow$

$a(t) = v'(t) = 6t - 18$ (in (ft/s)/s or ft/s^2).

$a(1) = 6(1) - 18 = -12$ ft/s^2

(h)

(i) The particle is speeding up when v and a have the same sign. This occurs when $2 < t < 3$ (v and a are both negative) and when $t > 4$ (v and a are both positive). It is slowing down when v and a have opposite signs; that is, when $0 \le t < 2$ and when $3 < t < 4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. (a) $s = f(t) = \dfrac{9t}{t^2+9}$ (in feet) $\Rightarrow$ $v(t) = f'(t) = \dfrac{(t^2+9)(9) - 9t(2t)}{(t^2+9)^2} = \dfrac{-9t^2+81}{(t^2+9)^2} = \dfrac{-9(t^2-9)}{(t^2+9)^2}$ (in ft/s)

(b) $v(1) = \dfrac{-9(1-9)}{(1+9)^2} = \dfrac{72}{100} = 0.72$ ft/s

(c) The particle is at rest when $v(t) = 0$. $\dfrac{-9(t^2-9)}{(t^2+9)^2} = 0 \quad \Leftrightarrow \quad t^2 - 9 = 0 \quad \Rightarrow \quad t = 3$ s [since $t \geq 0$].

(d) The particle is moving in the positive direction when $v(t) > 0$.

$$\frac{-9(t^2-9)}{(t^2+9)^2} > 0 \quad \Rightarrow \quad -9(t^2-9) > 0 \quad \Rightarrow \quad t^2 - 9 < 0 \quad \Rightarrow \quad t^2 < 9 \quad \Rightarrow \quad 0 \leq t < 3.$$

(e) Since the particle is moving in the positive direction and in the negative direction, we need to calculate the distance traveled in the intervals $[0, 3]$ and$[3, 6]$, respectively.

$|f(3) - f(0)| = \left|\frac{27}{18} - 0\right| = \frac{3}{2}$

$|f(6) - f(3)| = \left|\frac{54}{45} - \frac{27}{18}\right| = \frac{3}{10}$

The total distance is $\frac{3}{2} + \frac{3}{10} = \frac{9}{5}$ or 1.8 ft.

(f)

(g) $v(t) = -9\dfrac{t^2-9}{(t^2+9)^2} \quad \Rightarrow$

$$a(t) = v'(t) = -9\,\frac{(t^2+9)^2(2t) - (t^2-9)2(t^2+9)(2t)}{[(t^2+9)^2]^2} = -9\,\frac{2t(t^2+9)[(t^2+9) - 2(t^2-9)]}{(t^2+9)^4} = \frac{18t(t^2-27)}{(t^2+9)^3}.$$

$a(1) = \dfrac{18(-26)}{10^3} = -0.468$ ft/s^2

(h)

(i) The particle is speeding up when v and a have the same sign. a is negative for $0 < t < \sqrt{27}$ [≈ 5.2], so from the figure in part (h), we see that v and a are both negative for $3 < t < 3\sqrt{3}$. The particle is slowing down when v and a have opposite signs. This occurs when $0 < t < 3$ and when $t > 3\sqrt{3}$.

3. (a) $s = f(t) = \sin(\pi t/2)$ (in feet) $\Rightarrow$ $v(t) = f'(t) = \cos(\pi t/2) \cdot (\pi/2) = \frac{\pi}{2}\cos(\pi t/2)$ (in ft/s)

(b) $v(1) = \frac{\pi}{2}\cos\frac{\pi}{2} = \frac{\pi}{2}(0) = 0$ ft/s

(c) The particle is at rest when $v(t) = 0$. $\frac{\pi}{2}\cos\frac{\pi}{2}t = 0 \quad \Leftrightarrow \quad \cos\frac{\pi}{2}t = 0 \quad \Leftrightarrow \quad \frac{\pi}{2}t = \frac{\pi}{2} + n\pi \quad \Leftrightarrow \quad t = 1 + 2n$, where n is a nonnegative integer since $t \geq 0$.

(d) The particle is moving in the positive direction when $v(t) > 0$. From part (c), we see that v changes sign at every positive odd integer. v is positive when $0 < t < 1$, $3 < t < 5$, $7 < t < 9$, and so on.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(e) v changes sign at $t = 1$, 3, and 5 in the interval $[0, 6]$. The total distance traveled during the first 6 seconds is

$$\begin{aligned}|f(1) - f(0)| + |f(3) - f(1)| + |f(5) - f(3)| + |f(6) - f(5)| &= |1 - 0| + |-1 - 1| + |1 - (-1)| + |0 - 1| \\ &= 1 + 2 + 2 + 1 = 6 \text{ ft}\end{aligned}$$

(f)

(g) $v(t) = \frac{\pi}{2}\cos(\pi t/2) \quad \Rightarrow$

$$\begin{aligned}a(t) = v'(t) &= \tfrac{\pi}{2}\left[-\sin(\pi t/2) \cdot (\pi/2)\right] \\ &= (-\pi^2/4)\sin(\pi t/2) \text{ ft/s}^2\end{aligned}$$

$$a(1) = (-\pi^2/4)\sin(\pi/2) = -\pi^2/4 \text{ ft/s}^2$$

(h)

(i) The particle is speeding up when v and a have the same sign. From the figure in part (h), we see that v and a are both positive when $3 < t < 4$ and both negative when $1 < t < 2$ and $5 < t < 6$. Thus, the particle is speeding up when $1 < t < 2$, $3 < t < 4$, and $5 < t < 6$. The particle is slowing down when v and a have opposite signs; that is, when $0 < t < 1$, $2 < t < 3$, and $4 < t < 5$.

4. (a) $s = f(t) = t^2 e^{-t}$ (in feet) $\quad\Rightarrow\quad v(t) = f'(t) = t^2(-e^{-t}) + e^{-t}(2t) = te^{-t}(-t + 2)$ (in ft/s)

(b) $v(1) = (1)e^{-1}(-1 + 2) = 1/e$ ft/s

(c) The particle is at rest when $v(t) = 0$. $v(t) = 0 \quad\Leftrightarrow\quad t = 0$ or 2 s.

(d) The particle is moving in the positive direction when $v(t) > 0 \quad\Leftrightarrow\quad te^{-t}(-t + 2) > 0 \quad\Leftrightarrow\quad t(-t + 2) > 0 \quad\Leftrightarrow$ $0 < t < 2$.

(e) v changes sign at $t = 2$ in the interval $[0, 6]$. The total distance traveled during the first 6 seconds is

$$\begin{aligned}|f(2) - f(0)| + |f(6) - f(2)| &= \left|4e^{-2} - 0\right| + \left|36e^{-6} - 4e^{-2}\right| = 4e^{-2} + 4e^{-2} - 36e^{-6} \\ &= 8e^{-2} - 36e^{-6} \approx 0.99 \text{ ft}\end{aligned}$$

(f)

(g) $v(t) = (2t - t^2)e^{-t} \quad\Rightarrow$

$$\begin{aligned}a(t) = v'(t) &= (2t - t^2)(-e^{-t}) + e^{-t}(2 - 2t) \\ &= e^{-t}\left[-(2t - t^2) + (2 - 2t)\right] \\ &= e^{-t}(t^2 - 4t + 2) \text{ ft/s}^2\end{aligned}$$

$$a(1) = e^{-1}(1 - 4 + 2) = -1/e \text{ ft/s}^2$$

(h)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(i) $a(t) = 0 \quad \Leftrightarrow \quad t^2 - 4t + 2 = 0 \quad [e^{-t} \neq 0] \quad \Leftrightarrow \quad t = \dfrac{4 \pm \sqrt{8}}{2} = 2 \pm \sqrt{2}$ $[\approx 0.6$ and $3.4]$. The particle is speeding up when v and a have the same sign. Using the previous information and the figure in part (h), we see that v and a are both positive when $0 < t < 2 - \sqrt{2}$ and both negative when $2 < t < 2 + \sqrt{2}$. The particle is slowing down when v and a have opposite signs. This occurs when $2 - \sqrt{2} < t < 2$ and $t > 2 + \sqrt{2}$.

5. (a) From the figure, the velocity v is positive on the interval $(0, 2)$ and negative on the interval $(2, 3)$. The acceleration a is positive (negative) when the slope of the tangent line is positive (negative), so the acceleration is positive on the interval $(0, 1)$, and negative on the interval $(1, 3)$. The particle is speeding up when v and a have the same sign, that is, on the interval $(0, 1)$ when $v > 0$ and $a > 0$, and on the interval $(2, 3)$ when $v < 0$ and $a < 0$. The particle is slowing down when v and a have opposite signs, that is, on the interval $(1, 2)$ when $v > 0$ and $a < 0$.

(b) $v > 0$ on $(0, 3)$ and $v < 0$ on $(3, 4)$. $a > 0$ on $(1, 2)$ and $a < 0$ on $(0, 1)$ and $(2, 4)$. The particle is speeding up on $(1, 2)$ $[v > 0, a > 0]$ and on $(3, 4)$ $[v < 0, a < 0]$. The particle is slowing down on $(0, 1)$ and $(2, 3)$ $[v > 0, a < 0]$.

6. (a) The velocity v is positive when s is increasing, that is, on the intervals $(0, 1)$ and $(3, 4)$; and it is negative when s is decreasing, that is, on the interval $(1, 3)$. The acceleration a is positive when the graph of s is concave upward (v is increasing), that is, on the interval $(2, 4)$; and it is negative when the graph of s is concave downward (v is decreasing), that is, on the interval $(0, 2)$. The particle is speeding up on the interval $(1, 2)$ $[v < 0, a < 0]$ and on $(3, 4)$ $[v > 0, a > 0]$. The particle is slowing down on the interval $(0, 1)$ $[v > 0, a < 0]$ and on $(2, 3)$ $[v < 0, a > 0]$.

(b) The velocity v is positive on $(3, 4)$ and negative on $(0, 3)$. The acceleration a is positive on $(0, 1)$ and $(2, 4)$ and negative on $(1, 2)$. The particle is speeding up on the interval $(1, 2)$ $[v < 0, a < 0]$ and on $(3, 4)$ $[v > 0, a > 0]$. The particle is slowing down on the interval $(0, 1)$ $[v < 0, a > 0]$ and on $(2, 3)$ $[v < 0, a > 0]$.

7. The particle is traveling forward when its velocity is positive. From the graph, this occurs when $0 < t < 5$. The particle is traveling backward when its velocity is negative. From the graph, this occurs when $7 < t < 8$. When $5 < t < 7$, its velocity is zero and the particle is not moving.

8. The graph of the acceleration function is the graph of the derivative of the velocity function. Since the velocity function is piecewise linear, its derivative, where it exists, equals the slope of the corresponding piece of the velocity graph. Thus, we obtain the following table and graph of the acceleration function.

Interval	Acceleration (slope of velocity graph)
$0 < t < 1$	3
$1 < t < 3$	0
$3 < t < 5$	$-\frac{3}{2}$
$5 < t < 7$	0
$7 < t < 8$	-3

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The particle speeds up when its velocity and acceleration have the same sign. This occurs when $0 < t < 1$ ($v > 0$ and $a > 0$) and when $7 < t < 8$ ($v < 0$ and $a < 0$). The particle slows down when its velocity and acceleration have opposite signs. This occurs when $3 < t < 5$ ($v > 0$ and $a < 0$). The particle travels at a constant speed when its acceleration is zero. This occurs when $1 < t < 3$ and when $5 < t < 7$.

9. (a) $h(t) = 2 + 24.5t - 4.9t^2 \quad \Rightarrow \quad v(t) = h'(t) = 24.5 - 9.8t$. The velocity after 2 s is $v(2) = 24.5 - 9.8(2) = 4.9$ m/s and after 4 s is $v(4) = 24.5 - 9.8(4) = -14.7$ m/s.

(b) The projectile reaches its maximum height when the velocity is zero. $v(t) = 0 \quad \Leftrightarrow \quad 24.5 - 9.8t = 0 \quad \Leftrightarrow$

$t = \dfrac{24.5}{9.8} = 2.5$ s.

(c) The maximum height occurs when $t = 2.5$. $h(2.5) = 2 + 24.5(2.5) - 4.9(2.5)^2 = 32.625$ m $\left[\text{or } 32\frac{5}{8} \text{ m}\right]$.

(d) The projectile hits the ground when $h = 0 \quad \Leftrightarrow \quad 2 + 24.5t - 4.9t^2 = 0 \quad \Leftrightarrow$

$t = \dfrac{-24.5 \pm \sqrt{24.5^2 - 4(-4.9)(2)}}{2(-4.9)} \quad \Rightarrow \quad t = t_f \approx 5.08$ s [since $t \geq 0$].

(e) The projectile hits the ground when $t = t_f$. Its velocity is $v(t_f) = 24.5 - 9.8t_f \approx -25.3$ m/s [downward].

10. (a) At maximum height the velocity of the ball is 0 ft/s. $v(t) = s'(t) = 80 - 32t = 0 \quad \Leftrightarrow \quad 32t = 80 \quad \Leftrightarrow \quad t = \frac{5}{2}$.

So the maximum height is $s\left(\frac{5}{2}\right) = 80\left(\frac{5}{2}\right) - 16\left(\frac{5}{2}\right)^2 = 200 - 100 = 100$ ft.

(b) $s(t) = 80t - 16t^2 = 96 \quad \Leftrightarrow \quad 16t^2 - 80t + 96 = 0 \quad \Leftrightarrow \quad 16(t^2 - 5t + 6) = 0 \quad \Leftrightarrow \quad 16(t-3)(t-2) = 0$.

So the ball has a height of 96 ft on the way up at $t = 2$ and on the way down at $t = 3$. At these times the velocities are $v(2) = 80 - 32(2) = 16$ ft/s and $v(3) = 80 - 32(3) = -16$ ft/s, respectively.

11. (a) $h(t) = 15t - 1.86t^2 \quad \Rightarrow \quad v(t) = h'(t) = 15 - 3.72t$. The velocity after 2 s is $v(2) = 15 - 3.72(2) = 7.56$ m/s.

(b) $25 = h \quad \Leftrightarrow \quad 1.86t^2 - 15t + 25 = 0 \quad \Leftrightarrow \quad t = \dfrac{15 \pm \sqrt{15^2 - 4(1.86)(25)}}{2(1.86)} \quad \Leftrightarrow \quad t = t_1 \approx 2.35$ or $t = t_2 \approx 5.71$.

The velocities are $v(t_1) = 15 - 3.72t_1 \approx 6.24$ m/s [upward] and $v(t_2) = 15 - 3.72t_2 \approx -6.24$ m/s [downward].

12. (a) $s(t) = t^4 - 4t^3 - 20t^2 + 20t \quad \Rightarrow \quad v(t) = s'(t) = 4t^3 - 12t^2 - 40t + 20$. $v = 20 \quad \Leftrightarrow$

$4t^3 - 12t^2 - 40t + 20 = 20 \quad \Leftrightarrow \quad 4t^3 - 12t^2 - 40t = 0 \quad \Leftrightarrow \quad 4t(t^2 - 3t - 10) = 0 \quad \Leftrightarrow$

$4t(t-5)(t+2) = 0 \quad \Leftrightarrow \quad t = 0$ s or 5 s [for $t \geq 0$].

(b) $a(t) = v'(t) = 12t^2 - 24t - 40$. $a = 0 \quad \Leftrightarrow \quad 12t^2 - 24t - 40 = 0 \quad \Leftrightarrow \quad 4(3t^2 - 6t - 10) = 0 \quad \Leftrightarrow$

$t = \dfrac{6 \pm \sqrt{6^2 - 4(3)(-10)}}{2(3)} = 1 \pm \dfrac{1}{3}\sqrt{39} \approx 3.08$ s [for $t \geq 0$]. At this time, the acceleration changes from negative to positive and the velocity attains its minimum value.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. (a) $A(x) = x^2 \quad \Rightarrow \quad A'(x) = 2x$. $A'(15) = 30$ mm^2/mm is the rate at which the area is increasing with respect to the side length as x reaches 15 mm.

(b) The perimeter is $P(x) = 4x$, so $A'(x) = 2x = \frac{1}{2}(4x) = \frac{1}{2}P(x)$. The figure suggests that if Δx is small, then the change in the area of the square is approximately half of its perimeter (2 of the 4 sides) times Δx. From the figure, $\Delta A = 2x\,(\Delta x) + (\Delta x)^2$. If Δx is small, then $\Delta A \approx 2x\,(\Delta x)$ and so $\Delta A/\Delta x \approx 2x$.

14. (a) $V(x) = x^3 \quad \Rightarrow \quad \dfrac{dV}{dx} = 3x^2$. $\left.\dfrac{dV}{dx}\right|_{x=3} = 3(3)^2 = 27$ mm^3/mm is the rate at which the volume is increasing as x increases past 3 mm.

(b) The surface area is $S(x) = 6x^2$, so $V'(x) = 3x^2 = \frac{1}{2}(6x^2) = \frac{1}{2}S(x)$. The figure suggests that if Δx is small, then the change in the volume of the cube is approximately half of its surface area (the area of 3 of the 6 faces) times Δx. From the figure, $\Delta V = 3x^2(\Delta x) + 3x(\Delta x)^2 + (\Delta x)^3$. If Δx is small, then $\Delta V \approx 3x^2(\Delta x)$ and so $\Delta V/\Delta x \approx 3x^2$.

15. (a) Using $A(r) = \pi r^2$, we find that the average rate of change is:

(i) $\dfrac{A(3) - A(2)}{3 - 2} = \dfrac{9\pi - 4\pi}{1} = 5\pi$ (ii) $\dfrac{A(2.5) - A(2)}{2.5 - 2} = \dfrac{6.25\pi - 4\pi}{0.5} = 4.5\pi$

(iii) $\dfrac{A(2.1) - A(2)}{2.1 - 2} = \dfrac{4.41\pi - 4\pi}{0.1} = 4.1\pi$

(b) $A(r) = \pi r^2 \quad \Rightarrow \quad A'(r) = 2\pi r$, so $A'(2) = 4\pi$.

(c) The circumference is $C(r) = 2\pi r = A'(r)$. The figure suggests that if Δr is small, then the change in the area of the circle (a ring around the outside) is approximately equal to its circumference times Δr. Straightening out this ring gives us a shape that is approximately rectangular with length $2\pi r$ and width Δr, so $\Delta A \approx 2\pi r(\Delta r)$. Algebraically, $\Delta A = A(r + \Delta r) - A(r) = \pi(r + \Delta r)^2 - \pi r^2 = 2\pi r(\Delta r) + \pi(\Delta r)^2$. So we see that if Δr is small, then $\Delta A \approx 2\pi r(\Delta r)$ and therefore, $\Delta A/\Delta r \approx 2\pi r$.

16. After t seconds the radius is $r = 60t$, so the area is $A(t) = \pi(60t)^2 = 3600\pi t^2 \quad \Rightarrow \quad A'(t) = 7200\pi t \quad \Rightarrow$

(a) $A'(1) = 7200\pi$ cm^2/s (b) $A'(3) = 21{,}600\pi$ cm^2/s (c) $A'(5) = 36{,}000\pi$ cm^2/s

As time goes by, the area grows at an increasing rate. In fact, the rate of change is linear with respect to time.

17. $S(r) = 4\pi r^2 \quad \Rightarrow \quad S'(r) = 8\pi r \quad \Rightarrow$

(a) $S'(1) = 8\pi$ ft^2/ft (b) $S'(2) = 16\pi$ ft^2/ft (c) $S'(3) = 24\pi$ ft^2/ft

As the radius increases, the surface area grows at an increasing rate. In fact, the rate of change is linear with respect to the radius.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. (a) Using $V(r) = \frac{4}{3}\pi r^3$, we find that the average rate of change is:

(i) $\dfrac{V(8) - V(5)}{8 - 5} = \dfrac{\frac{4}{3}\pi(512) - \frac{4}{3}\pi(125)}{3} = 172\pi\ \mu\text{m}^3/\mu\text{m}$

(ii) $\dfrac{V(6) - V(5)}{6 - 5} = \dfrac{\frac{4}{3}\pi(216) - \frac{4}{3}\pi(125)}{1} = 121.\overline{3}\pi\ \mu\text{m}^3/\mu\text{m}$

(iii) $\dfrac{V(5.1) - V(5)}{5.1 - 5} = \dfrac{\frac{4}{3}\pi(5.1)^3 - \frac{4}{3}\pi(5)^3}{0.1} = 102.01\overline{3}\pi\ \mu\text{m}^3/\mu\text{m}$

(b) $V'(r) = 4\pi r^2$, so $V'(5) = 100\pi\ \mu\text{m}^3/\mu\text{m}$.

(c) $V(r) = \frac{4}{3}\pi r^3 \quad \Rightarrow \quad V'(r) = 4\pi r^2 = S(r)$. By analogy with Exercise 15(c), we can say that the change in the volume of the spherical shell, ΔV, is approximately equal to its thickness, Δr, times the surface area of the inner sphere. Thus, $\Delta V \approx 4\pi r^2(\Delta r)$ and so $\Delta V/\Delta r \approx 4\pi r^2$.

19. The mass is $f(x) = 3x^2$, so the linear density at x is $\rho(x) = f'(x) = 6x$.

(a) $\rho(1) = 6$ kg/m (b) $\rho(2) = 12$ kg/m (c) $\rho(3) = 18$ kg/m

Since ρ is an increasing function, the density will be the highest at the right end of the rod and lowest at the left end.

20. $V(t) = 5000\left(1 - \frac{1}{40}t\right)^2 \quad \Rightarrow \quad V'(t) = 5000 \cdot 2\left(1 - \frac{1}{40}t\right)\left(-\frac{1}{40}\right) = -250\left(1 - \frac{1}{40}t\right)$

(a) $V'(5) = -250\left(1 - \frac{5}{40}\right) = -218.75$ gal/min (b) $V'(10) = -250\left(1 - \frac{10}{40}\right) = -187.5$ gal/min

(c) $V'(20) = -250\left(1 - \frac{20}{40}\right) = -125$ gal/min (d) $V'(40) = -250\left(1 - \frac{40}{40}\right) = 0$ gal/min

The water is flowing out the fastest at the beginning — when $t = 0$, $V'(t) = -250$ gal/min. The water is flowing out the slowest at the end — when $t = 40$, $V'(t) = 0$. As the tank empties, the water flows out more slowly.

21. The quantity of charge is $Q(t) = t^3 - 2t^2 + 6t + 2$, so the current is $Q'(t) = 3t^2 - 4t + 6$.

(a) $Q'(0.5) = 3(0.5)^2 - 4(0.5) + 6 = 4.75$ A (b) $Q'(1) = 3(1)^2 - 4(1) + 6 = 5$ A

The current is lowest when Q' has a minimum. $Q''(t) = 6t - 4 < 0$ when $t < \frac{2}{3}$. So the current decreases when $t < \frac{2}{3}$ and increases when $t > \frac{2}{3}$. Thus, the current is lowest at $t = \frac{2}{3}$ s.

22. (a) $F = \dfrac{GmM}{r^2} = (GmM)r^{-2} \quad \Rightarrow \quad \dfrac{dF}{dr} = -2(GmM)r^{-3} = -\dfrac{2GmM}{r^3}$, which is the rate of change of the force with respect to the distance between the bodies. The minus sign indicates that as the distance r between the bodies increases, the magnitude of the force F exerted by the body of mass m on the body of mass M is decreasing.

(b) Given $F'(20{,}000) = -2$, find $F'(10{,}000)$. $-2 = -\dfrac{2GmM}{20{,}000^3} \quad \Rightarrow \quad GmM = 20{,}000^3$.

$$F'(10{,}000) = -\frac{2(20{,}000^3)}{10{,}000^3} = -2 \cdot 2^3 = -16 \text{ N/km}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. With $m = m_0\left(1 - \dfrac{v^2}{c^2}\right)^{-1/2}$,

$$F = \frac{d}{dt}(mv) = m\frac{d}{dt}(v) + v\frac{d}{dt}(m) = m_0\left(1 - \frac{v^2}{c^2}\right)^{-1/2} \cdot a + v \cdot m_0\left[-\frac{1}{2}\left(1 - \frac{v^2}{c^2}\right)^{-3/2}\right]\left(-\frac{2v}{c^2}\right)\frac{d}{dt}(v)$$

$$= m_0\left(1 - \frac{v^2}{c^2}\right)^{-3/2} \cdot a\left[\left(1 - \frac{v^2}{c^2}\right) + \frac{v^2}{c^2}\right] = \frac{m_0 a}{(1 - v^2/c^2)^{3/2}}$$

Note that we factored out $(1 - v^2/c^2)^{-3/2}$ since $-3/2$ was the lesser exponent. Also note that $\dfrac{d}{dt}(v) = a$.

24. (a) $D(t) = 7 + 5\cos[0.503(t - 6.75)] \quad \Rightarrow \quad D'(t) = -5\sin[0.503(t - 6.75)](0.503) = -2.515\sin[0.503(t - 6.75)]$.

At 3:00 AM, $t = 3$, and $D'(3) = -2.515\sin[0.503(-3.75)] \approx 2.39$ m/h (rising).

(b) At 6:00 AM, $t = 6$, and $D'(6) = -2.515\sin[0.503(-0.75)] \approx 0.93$ m/h (rising).

(c) At 9:00 AM, $t = 9$, and $D'(9) = -2.515\sin[0.503(2.25)] \approx -2.28$ m/h (falling).

(d) At noon, $t = 12$, and $D'(12) = -2.515\sin[0.503(5.25)] \approx -1.21$ m/h (falling).

25. (a) To find the rate of change of volume with respect to pressure, we first solve for V in terms of P.

$$PV = C \quad \Rightarrow \quad V = \frac{C}{P} \quad \Rightarrow \quad \frac{dV}{dP} = -\frac{C}{P^2}.$$

(b) From the formula for dV/dP in part (a), we see that as P increases, the absolute value of dV/dP decreases.

Thus, the volume is decreasing more rapidly at the beginning of the 10 minutes.

(c) $\beta = -\dfrac{1}{V}\dfrac{dV}{dP} = -\dfrac{1}{V}\left(-\dfrac{C}{P^2}\right)$ [from part (a)] $= \dfrac{C}{(PV)P} = \dfrac{C}{CP} = \dfrac{1}{P}$

26. (a) $[\mathrm{C}] = \dfrac{a^2kt}{akt + 1} \quad \Rightarrow \quad$ rate of reaction $= \dfrac{d[\mathrm{C}]}{dt} = \dfrac{(akt + 1)(a^2k) - (a^2kt)(ak)}{(akt + 1)^2} = \dfrac{a^2k(akt + 1 - akt)}{(akt + 1)^2} = \dfrac{a^2k}{(akt + 1)^2}$

(b) If $x = [\mathrm{C}]$, then $a - x = a - \dfrac{a^2kt}{akt + 1} = \dfrac{a^2kt + a - a^2kt}{akt + 1} = \dfrac{a}{akt + 1}$.

So $k(a - x)^2 = k\left(\dfrac{a}{akt + 1}\right)^2 = \dfrac{a^2k}{(akt + 1)^2} = \dfrac{d[\mathrm{C}]}{dt}$ [from part (a)] $= \dfrac{dx}{dt}$.

(c) As $t \to \infty$, $[\mathrm{C}] = \dfrac{a^2kt}{akt + 1} = \dfrac{(a^2kt)/t}{(akt + 1)/t} = \dfrac{a^2k}{ak + (1/t)} \to \dfrac{a^2k}{ak} = a$ moles/L.

(d) As $t \to \infty$, $\dfrac{d[\mathrm{C}]}{dt} = \dfrac{a^2k}{(akt + 1)^2} \to 0$.

(e) As t increases, nearly all of the reactants A and B are converted into product C. In practical terms, the reaction virtually stops.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. In Example 6, the population function was $n = 2^t\, n_0$. Since we are tripling instead of doubling and the initial population is 400, the population function is $n(t) = 400 \cdot 3^t$. The rate of growth is $n'(t) = 400 \cdot 3^t \cdot \ln 3$, so the rate of growth after 2.5 hours is $n'(2.5) = 400 \cdot 3^{2.5} \cdot \ln 3 \approx 6850$ bacteria/hour.

28. $n = f(t) = \dfrac{a}{1 + be^{-0.7t}} \quad \Rightarrow \quad n' = -\dfrac{a \cdot be^{-0.7t}(-0.7)}{(1 + be^{-0.7t})^2}$ [Reciprocal Rule]. When $t = 0$, $n = 20$ and $n' = 12$.

$f(0) = 20 \quad \Rightarrow \quad 20 = \dfrac{a}{1+b} \quad \Rightarrow \quad a = 20(1+b)$. $\quad f'(0) = 12 \quad \Rightarrow \quad 12 = \dfrac{0.7ab}{(1+b)^2} \quad \Rightarrow \quad 12 = \dfrac{0.7(20)(1+b)b}{(1+b)^2} \quad \Rightarrow$

$\dfrac{12}{14} = \dfrac{b}{1+b} \quad \Rightarrow \quad 6(1+b) = 7b \quad \Rightarrow \quad 6 + 6b = 7b \quad \Rightarrow \quad b = 6$ and $a = 20(1+6) = 140$. For the long run, we let t increase without bound. $\displaystyle\lim_{t\to\infty} f(t) = \lim_{t\to\infty} \frac{140}{1 + 6e^{-0.7t}} = \frac{140}{1 + 6 \cdot 0} = 140$, indicating that in the long run, the yeast population stabilizes at 140 cells.

29. (a) **1920:** $m_1 = \dfrac{1860 - 1750}{1920 - 1910} = \dfrac{110}{10} = 11$, $m_2 = \dfrac{2070 - 1860}{1930 - 1920} = \dfrac{210}{10} = 21$,

$(m_1 + m_2)/2 = (11 + 21)/2 = 16$ million/year

1980: $m_1 = \dfrac{4450 - 3710}{1980 - 1970} = \dfrac{740}{10} = 74$, $m_2 = \dfrac{5280 - 4450}{1990 - 1980} = \dfrac{830}{10} = 83$,

$(m_1 + m_2)/2 = (74 + 83)/2 = 78.5$ million/year

(b) $P(t) = at^3 + bt^2 + ct + d$ (in millions of people), where $a \approx -0.000\,284\,900\,3$, $b \approx 0.522\,433\,122\,43$, $c \approx -6.395\,641\,396$, and $d \approx 1720.586\,081$.

(c) $P(t) = at^3 + bt^2 + ct + d \quad \Rightarrow \quad P'(t) = 3at^2 + 2bt + c$ (in millions of people per year)

(d) 1920 corresponds to $t = 20$ and $P'(20) \approx 14.16$ million/year. 1980 corresponds to $t = 80$ and $P'(80) \approx 71.72$ million/year. These estimates are smaller than the estimates in part (a).

(e) $f(t) = pq^t$ (where $p = 1.43653 \times 10^9$ and $q = 1.01395$) $\quad \Rightarrow \quad f'(t) = pq^t \ln q$ (in millions of people per year)

(f) $f'(20) \approx 26.25$ million/year [much larger than the estimates in part (a) and (d)].

$f'(80) \approx 60.28$ million/year [much smaller than the estimates in parts (a) and (d)].

(g) $P'(85) \approx 76.24$ million/year and $f'(85) \approx 64.61$ million/year. The first estimate is probably more accurate.

30. (a) $A(t) = at^4 + bt^3 + ct^2 + dt + e$ years of age, where $a \approx -1.404\,771\,699 \times 10^{-6}$, $b \approx 0.011\,167\,331\,7$, $c \approx -33.288\,096\,21$, $d \approx 44{,}097.25101$, $e \approx -21{,}904{,}396.36$.

(b) $A(t) = at^4 + bt^3 + ct^2 + dt + e \quad \Rightarrow \quad A'(t) = 4at^3 + 3bt^2 + 2ct + d$ (in years of age per year).

(c) $A'(1990) \approx 0.11$, so the rate of change of first marriage age for Japanese women in 1990 was approximately 0.11 years of age per year.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) The model for A and the data points are shown on the left, and a graph for $A'(t)$ is shown on the right:

31. (a) Using $v = \dfrac{P}{4\eta l}(R^2 - r^2)$ with $R = 0.01$, $l = 3$, $P = 3000$, and $\eta = 0.027$, we have v as a function of r:

$$v(r) = \frac{3000}{4(0.027)3}(0.01^2 - r^2).\ \ v(0) = 0.\overline{925}\text{ cm/s},\ v(0.005) = 0.69\overline{4}\text{ cm/s},\ v(0.01) = 0.$$

(b) $v(r) = \dfrac{P}{4\eta l}(R^2 - r^2) \quad \Rightarrow \quad v'(r) = \dfrac{P}{4\eta l}(-2r) = -\dfrac{Pr}{2\eta l}.$ When $l = 3$, $P = 3000$, and $\eta = 0.027$, we have

$$v'(r) = -\frac{3000r}{2(0.027)3}.\ \ v'(0) = 0,\ v'(0.005) = -92.\overline{592}\ (\text{cm/s})/\text{cm, and } v'(0.01) = -185.\overline{185}\ (\text{cm/s})/\text{cm}.$$

(c) The velocity is greatest where $r = 0$ (at the center) and the velocity is changing most where $r = R = 0.01$ cm (at the edge).

32. (a) (i) $f = \dfrac{1}{2L}\sqrt{\dfrac{T}{\rho}} = \left(\dfrac{1}{2}\sqrt{\dfrac{T}{\rho}}\right)L^{-1} \quad \Rightarrow \quad \dfrac{df}{dL} = -\left(\dfrac{1}{2}\sqrt{\dfrac{T}{\rho}}\right)L^{-2} = -\dfrac{1}{2L^2}\sqrt{\dfrac{T}{\rho}}$

(ii) $f = \dfrac{1}{2L}\sqrt{\dfrac{T}{\rho}} = \left(\dfrac{1}{2L\sqrt{\rho}}\right)T^{1/2} \quad \Rightarrow \quad \dfrac{df}{dT} = \dfrac{1}{2}\left(\dfrac{1}{2L\sqrt{\rho}}\right)T^{-1/2} = \dfrac{1}{4L\sqrt{T\rho}}$

(iii) $f = \dfrac{1}{2L}\sqrt{\dfrac{T}{\rho}} = \left(\dfrac{\sqrt{T}}{2L}\right)\rho^{-1/2} \quad \Rightarrow \quad \dfrac{df}{d\rho} = -\dfrac{1}{2}\left(\dfrac{\sqrt{T}}{2L}\right)\rho^{-3/2} = -\dfrac{\sqrt{T}}{4L\rho^{3/2}}$

(b) *Note:* Illustrating tangent lines on the generic figures may help to explain the results.

(i) $\dfrac{df}{dL} < 0$ and L is decreasing $\Rightarrow$ f is increasing $\Rightarrow$ higher note

(ii) $\dfrac{df}{dT} > 0$ and T is increasing $\Rightarrow$ f is increasing $\Rightarrow$ higher note

(iii) $\dfrac{df}{d\rho} < 0$ and ρ is increasing $\Rightarrow$ f is decreasing $\Rightarrow$ lower note

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. (a) $C(x) = 2000 + 3x + 0.01x^2 + 0.0002x^3 \quad \Rightarrow \quad C'(x) = 0 + 3(1) + 0.01(2x) + 0.0002(3x^2) = 3 + 0.02x + 0.0006x^2$

(b) $C'(100) = 3 + 0.02(100) + 0.0006(100)^2 = 3 + 2 + 6 = \$11/\text{pair}$. $C'(100)$ is the rate at which the cost is increasing as the 100th pair of jeans is produced. It predicts the (approximate) cost of the 101st pair.

(c) The cost of manufacturing the 101st pair of jeans is

$C(101) - C(100) = 2611.0702 - 2600 = 11.0702 \approx \$11.07$. This is close to the marginal cost from part (b).

34. (a) $C(q) = 84 + 0.16q - 0.0006q^2 + 0.000003q^3 \quad \Rightarrow \quad C'(q) = 0.16 - 0.0012q + 0.000009q^2$, and

$C'(100) = 0.16 - 0.0012(100) + 0.000009(100)^2 = 0.13$. This is the rate at which the cost is increasing as the 100th item is produced.

(b) The actual cost of producing the 101st item is $C(101) - C(100) = 97.13030299 - 97 \approx \$0.13$

35. (a) $A(x) = \dfrac{p(x)}{x} \quad \Rightarrow \quad A'(x) = \dfrac{xp'(x) - p(x) \cdot 1}{x^2} = \dfrac{xp'(x) - p(x)}{x^2}$.

$A'(x) > 0 \quad \Rightarrow \quad A(x)$ is increasing; that is, the average productivity increases as the size of the workforce increases.

(b) $p'(x)$ is greater than the average productivity $\quad \Rightarrow \quad p'(x) > A(x) \quad \Rightarrow \quad p'(x) > \dfrac{p(x)}{x} \quad \Rightarrow \quad xp'(x) > p(x) \quad \Rightarrow$

$xp'(x) - p(x) > 0 \quad \Rightarrow \quad \dfrac{xp'(x) - p(x)}{x^2} > 0 \quad \Rightarrow \quad A'(x) > 0.$

36. (a) $R = \dfrac{40 + 24x^{0.4}}{1 + 4x^{0.4}} \quad \Rightarrow \quad S = \dfrac{dR}{dx} = \dfrac{(1 + 4x^{0.4})(9.6x^{-0.6}) - (40 + 24x^{0.4})(1.6x^{-0.6})}{(1 + 4x^{0.4})^2}$

$$= \frac{9.6x^{-0.6} + 38.4x^{-0.2} - 64x^{-0.6} - 38.4x^{-0.2}}{(1 + 4x^{0.4})^2} = -\frac{54.4x^{-0.6}}{(1 + 4x^{0.4})^2}$$

(b)

At low levels of brightness, R is quite large $[R(0) = 40]$ and is quickly decreasing, that is, S is negative with large absolute value. This is to be expected: at low levels of brightness, the eye is more sensitive to slight changes than it is at higher levels of brightness.

37. $t = \ln\left(\dfrac{3c + \sqrt{9c^2 - 8c}}{2}\right) = \ln\left(3c + \sqrt{9c^2 - 8c}\right) - \ln 2 \quad \Rightarrow$

$$\frac{dt}{dc} = \frac{1}{3c + \sqrt{9c^2 - 8c}} \frac{d}{dc}\left(3c + \sqrt{9c^2 - 8c}\right) - 0 = \frac{3 + \frac{1}{2}(9c^2 - 8c)^{-1/2}(18c - 8)}{3c + \sqrt{9c^2 - 8c}}$$

$$= \frac{3 + \dfrac{9c - 4}{\sqrt{9c^2 - 8c}}}{3c + \sqrt{9c^2 - 8c}} = \frac{3\sqrt{9c^2 - 8c} + 9c - 4}{\sqrt{9c^2 - 8c}\left(3c + \sqrt{9c^2 - 8c}\right)}.$$

This derivative represents the rate of change of duration of dialysis required with respect to the initial urea concentration.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. $f(r) = 2\sqrt{Dr} \quad \Rightarrow \quad f'(r) = 2 \cdot \frac{1}{2}(Dr)^{-1/2} \cdot D = \dfrac{D}{\sqrt{Dr}} = \sqrt{\dfrac{D}{r}}$. $f'(r)$ is the rate of change of the wave speed with respect to the reproductive rate.

39. $PV = nRT \quad \Rightarrow \quad T = \dfrac{PV}{nR} = \dfrac{PV}{(10)(0.0821)} = \dfrac{1}{0.821}(PV)$. Using the Product Rule, we have

$$\frac{dT}{dt} = \frac{1}{0.821}\left[P(t)V'(t) + V(t)P'(t)\right] = \frac{1}{0.821}\left[(8)(-0.15) + (10)(0.10)\right] \approx -0.2436 \text{ K/min.}$$

40. (a) If $dP/dt = 0$, the population is stable (it is constant).

(b) $\dfrac{dP}{dt} = 0 \quad \Rightarrow \quad \beta P = r_0\left(1 - \dfrac{P}{P_c}\right)P \quad \Rightarrow \quad \dfrac{\beta}{r_0} = 1 - \dfrac{P}{P_c} \quad \Rightarrow \quad \dfrac{P}{P_c} = 1 - \dfrac{\beta}{r_0} \quad \Rightarrow \quad P = P_c\left(1 - \dfrac{\beta}{r_0}\right)$.

If $P_c = 10{,}000$, $r_0 = 5\% = 0.05$, and $\beta = 4\% = 0.04$, then $P = 10{,}000\left(1 - \frac{4}{5}\right) = 2000$.

(c) If $\beta = 0.05$, then $P = 10{,}000\left(1 - \frac{5}{5}\right) = 0$. There is no stable population.

41. (a) If the populations are stable, then the growth rates are neither positive nor negative; that is, $\dfrac{dC}{dt} = 0$ and $\dfrac{dW}{dt} = 0$.

(b) "The caribou go extinct" means that the population is zero, or mathematically, $C = 0$.

(c) We have the equations $\dfrac{dC}{dt} = aC - bCW$ and $\dfrac{dW}{dt} = -cW + dCW$. Let $dC/dt = dW/dt = 0$, $a = 0.05$, $b = 0.001$, $c = 0.05$, and $d = 0.0001$ to obtain $0.05C - 0.001CW = 0$ **(1)** and $-0.05W + 0.0001CW = 0$ **(2)**. Adding 10 times **(2)** to **(1)** eliminates the CW-terms and gives us $0.05C - 0.5W = 0 \quad \Rightarrow \quad C = 10W$. Substituting $C = 10W$ into **(1)** results in $0.05(10W) - 0.001(10W)W = 0 \quad \Leftrightarrow \quad 0.5W - 0.01W^2 = 0 \quad \Leftrightarrow \quad 50W - W^2 = 0 \quad \Leftrightarrow$ $W(50 - W) = 0 \quad \Leftrightarrow \quad W = 0$ or 50. Since $C = 10W$, $C = 0$ or 500. Thus, the population pairs (C, W) that lead to stable populations are $(0, 0)$ and $(500, 50)$. So it is possible for the two species to live in harmony.

42. (a) The rate of change of retention t days after a task is learned is given by $R'(t)$. $R(t) = a + b(1 + ct)^{-\beta} \quad \Rightarrow$ $R'(t) = b \cdot (-\beta)(1 + ct)^{-\beta-1} \cdot c = -\beta bc(1 + ct)^{-\beta-1}$ (expressed as a fraction of memory per day).

(b) We may write the rate of change as $R'(t) = -\dfrac{\beta bc}{(1 + ct)^{\beta+1}}$. The magnitude of this quantity decreases as t increases. Thus, you forget how to perform a task faster soon after learning it than a long time after you have learned it.

(c) $\lim\limits_{t\to\infty} R(t) = \lim\limits_{t\to\infty}\left[a + \dfrac{b}{(1 + ct)^{\beta}}\right] = a + 0 = a$, so the fraction of memory that is permanent is a.

43. (a) $I = \log_2\left(\dfrac{2D}{W}\right) \quad \Rightarrow \quad \dfrac{dI}{dD} \quad [W \text{ constant}] \quad = \dfrac{1}{\left(\dfrac{2D}{W}\right)\ln 2} \cdot \dfrac{2}{W} = \dfrac{1}{D \ln 2}$

As D increases, the rate of change of difficulty decreases.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $I = \log_2\left(\dfrac{2D}{W}\right) \Rightarrow \dfrac{dI}{dW}$ [D constant] $= \dfrac{1}{\left(\dfrac{2D}{W}\right)\ln 2} \cdot \left(-2DW^{-2}\right) = \dfrac{W}{2D\ln 2} \cdot \dfrac{-2D}{W^2} = -\dfrac{1}{W\ln 2}$

The negative sign indicates that difficulty decreases with increasing width. While the magnitude of the rate of change decreases with increasing width $\left(\text{that is, } \left|-\dfrac{1}{W\ln 2}\right| = \dfrac{1}{W\ln 2} \text{ decreases as } W \text{ increases}\right)$, the rate of change itself increases (gets closer to zero from the negative side) with increasing values of W.

(c) The answers to (a) and (b) agree with intuition. For fixed width, the difficulty of acquiring a target increases, but less and less so, as the distance to the target increases. Similarly, for a fixed distance to a target, the difficulty of acquiring the target decreases, but less and less so, as the width of the target increases.

3.8 Exponential Growth and Decay

1. The relative growth rate is $\dfrac{1}{P}\dfrac{dP}{dt} = 0.4159$, so $\dfrac{dP}{dt} = 0.4159P$ and by Theorem 2, $P(t) = P(0)e^{0.4159t} = 3.8e^{0.4159t}$ million cells. Thus, $P(2) = 3.8e^{0.4159(2)} \approx 8.7$ million cells.

2. (a) By Theorem 2, $P(t) = P(0)e^{kt} = 50e^{kt}$. In 20 minutes $\left(\frac{1}{3}\text{ hour}\right)$, there are 100 cells, so $P\left(\frac{1}{3}\right) = 50e^{k/3} = 100 \quad \Rightarrow$ $e^{k/3} = 2 \quad \Rightarrow \quad k/3 = \ln 2 \quad \Rightarrow \quad k = 3\ln 2 = \ln(2^3) = \ln 8$.

(b) $P(t) = 50e^{(\ln 8)t} = 50 \cdot 8^t$

(c) $P(6) = 50 \cdot 8^6 = 50 \cdot 2^{18} = 13{,}107{,}200$ cells

(d) $\dfrac{dP}{dt} = kP \quad \Rightarrow \quad P'(6) = kP(6) = (\ln 8)P(6) \approx 27{,}255{,}656$ cells/h

(e) $P(t) = 10^6 \quad \Leftrightarrow \quad 50 \cdot 8^t = 1{,}000{,}000 \quad \Leftrightarrow \quad 8^t = 20{,}000 \quad \Leftrightarrow \quad t\ln 8 = \ln 20{,}000 \quad \Leftrightarrow \quad t = \dfrac{\ln 20{,}000}{\ln 8} \approx 4.76$ h

3. (a) By Theorem 2, $P(t) = P(0)e^{kt} = 50e^{kt}$. Now $P(1.5) = 50e^{k(1.5)} = 975 \quad \Rightarrow \quad e^{1.5k} = \frac{975}{50} \quad \Rightarrow$ $1.5k = \ln 19.5 \quad \Rightarrow \quad k = \frac{1}{1.5}\ln 19.5 \approx 1.9803$. So $P(t) \approx 50e^{1.9803t}$ cells.

(b) Using 1.9803 for k, we get $P(3) = 50e^{1.9803(3)} = 19{,}013.85 \approx 19{,}014$ cells.

(c) $\dfrac{dP}{dt} = kP \quad \Rightarrow \quad P'(3) = k \cdot P(3) = 1.9803 \cdot 19{,}014$ [from parts (a) and (b)] $= 37{,}653.4 \approx 37{,}653$ cells/h

(d) $P(t) = 50e^{1.9803t} = 250{,}000 \quad \Rightarrow \quad e^{1.9803t} = \dfrac{250{,}000}{50} \quad \Rightarrow \quad e^{1.9803t} = 5000 \quad \Rightarrow \quad 1.9803t = \ln 5000 \quad \Rightarrow$ $t = \dfrac{\ln 5000}{1.9803} \approx 4.30$ h

4. (a) $y(t) = y(0)e^{kt} \quad \Rightarrow \quad y(2) = y(0)e^{2k} = 400$ and $y(6) = y(0)e^{6k} = 25{,}600$. Dividing these equations, we get $e^{6k}/e^{2k} = 25{,}600/400 \quad \Rightarrow \quad e^{4k} = 64 \quad \Rightarrow \quad 4k = \ln 2^6 = 6\ln 2 \quad \Rightarrow \quad k = \frac{3}{2}\ln 2 \approx 1.0397$, about 104% per hour.

(b) $400 = y(0)e^{2k} \quad \Rightarrow \quad y(0) = 400/e^{2k} \quad \Rightarrow \quad y(0) = 400/e^{3\ln 2} = 400/\left(e^{\ln 2}\right)^3 = 400/2^3 = 50$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $y(t) = y(0)e^{kt} = 50e^{(3/2)(\ln 2)t} = 50(e^{\ln 2})^{(3/2)t} \quad \Rightarrow \quad y(t) = 50(2)^{1.5t}$

(d) $y(4.5) = 50(2)^{1.5(4.5)} = 50(2)^{6.75} \approx 5382$ bacteria

(e) $\dfrac{dy}{dt} = ky = \left(\dfrac{3}{2}\ln 2\right)(50(2)^{6.75})$ [from parts (a) and (b)] ≈ 5596 bacteria/h

(f) $y(t) = 50{,}000 \quad \Rightarrow \quad 50{,}000 = 50(2)^{1.5t} \quad \Rightarrow \quad 1000 = (2)^{1.5t} \quad \Rightarrow \quad \ln 1000 = 1.5t \ln 2 \quad \Rightarrow$

$t = \dfrac{\ln 1000}{1.5 \ln 2} \approx 6.64$ h

5. (a) Let the population (in millions) in the year t be $P(t)$. Since the initial time is the year 1750, we substitute $t - 1750$ for t in Theorem 2, so the exponential model gives $P(t) = P(1750)e^{k(t-1750)}$. Then $P(1800) = 980 = 790e^{k(1800-1750)} \quad \Rightarrow$ $\frac{980}{790} = e^{k(50)} \quad \Rightarrow \quad \ln \frac{980}{790} = 50k \quad \Rightarrow \quad k = \frac{1}{50} \ln \frac{980}{790} \approx 0.0043104$. So with this model, we have $P(1900) = 790e^{k(1900-1750)} \approx 1508$ million, and $P(1950) = 790e^{k(1950-1750)} \approx 1871$ million. Both of these estimates are much too low.

(b) In this case, the exponential model gives $P(t) = P(1850)e^{k(t-1850)} \quad \Rightarrow \quad P(1900) = 1650 = 1260e^{k(1900-1850)} \quad \Rightarrow$ $\ln \frac{1650}{1260} = k(50) \quad \Rightarrow \quad k = \frac{1}{50} \ln \frac{1650}{1260} \approx 0.005393$. So with this model, we estimate $P(1950) = 1260e^{k(1950-1850)} \approx 2161$ million. This is still too low, but closer than the estimate of $P(1950)$ in part (a).

(c) The exponential model gives $P(t) = P(1900)e^{k(t-1900)} \quad \Rightarrow \quad P(1950) = 2560 = 1650e^{k(1950-1900)} \quad \Rightarrow$ $\ln \frac{2560}{1650} = k(50) \quad \Rightarrow \quad k = \frac{1}{50} \ln \frac{2560}{1650} \approx 0.008785$. With this model, we estimate $P(2000) = 1650e^{k(2000-1900)} \approx 3972$ million. This is much too low. The discrepancy is explained by the fact that the world birth rate (average yearly number of births per person) is about the same as always, whereas the mortality rate (especially the infant mortality rate) is much lower, owing mostly to advances in medical science and to the wars in the first part of the twentieth century. The exponential model assumes, among other things, that the birth and mortality rates will remain constant.

6. (a) Let $P(t)$ be the population (in millions) in the year t. Since the initial time is the year 1950, we substitute $t - 1950$ for t in Theorem 2, and find that the exponential model gives $P(t) = P(1950)e^{k(t-1950)} \quad \Rightarrow$ $P(1960) = 100 = 83e^{k(1960-1950)} \quad \Rightarrow \quad \frac{100}{83} = e^{10k} \quad \Rightarrow \quad k = \frac{1}{10} \ln \frac{100}{83} \approx 0.0186$. With this model, we estimate $P(1980) = 83e^{k(1980-1950)} = 83e^{30k} \approx 145$ million, which is an underestimate of the actual population of 150 million.

(b) As in part (a), $P(t) = P(1960)e^{k(t-1960)} \quad \Rightarrow \quad P(1980) = 150 = 100e^{20k} \quad \Rightarrow \quad 20k = \ln \frac{150}{100} \quad \Rightarrow$ $k = \frac{1}{20} \ln \frac{3}{2} \approx 0.0203$. Thus, $P(2000) = 100e^{40k} = 225$ million, which is an overestimate of the actual population of 214 million.

(c) As in part (a), $P(t) = P(1980)e^{k(t-1980)} \quad \Rightarrow \quad P(2000) = 214 = 150e^{20k} \quad \Rightarrow \quad 20k = \ln \frac{214}{150} \quad \Rightarrow$ $k = \frac{1}{20} \ln \frac{214}{150} \approx 0.0178$. Thus, $P(2010) = 150e^{30k} \approx 256$, which is an overestimate of the actual population of 243 million.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) Using the model in part (c), $P(2025) = 150e^{(2025-1980)k} = 150e^{45k} \approx 334$ million. This prediction is likely too high. The model gave an overestimate for 2010, and the amount of overestimation is likely to compound as time increases.

7. (a) If $y = [N_2O_5]$ then by Theorem 2, $\dfrac{dy}{dt} = -0.0005y \quad \Rightarrow \quad y(t) = y(0)e^{-0.0005t} = Ce^{-0.0005t}$.

(b) $y(t) = Ce^{-0.0005t} = 0.9C \quad \Rightarrow \quad e^{-0.0005t} = 0.9 \quad \Rightarrow \quad -0.0005t = \ln 0.9 \quad \Rightarrow \quad t = -2000 \ln 0.9 \approx 211$ s

8. (a) The mass remaining after t days is $y(t) = y(0)\,e^{kt} = 50e^{kt}$. Since the half-life is 28 days, $y(28) = 50e^{28k} = 25 \quad \Rightarrow$ $e^{28k} = \frac{1}{2} \quad \Rightarrow \quad 28k = \ln \frac{1}{2} \quad \Rightarrow \quad k = -(\ln 2)/28$, so $y(t) = 50e^{-(\ln 2)t/28} = 50 \cdot 2^{-t/28}$.

(b) $y(40) = 50 \cdot 2^{-40/28} \approx 18.6$ mg

(c) $y(t) = 2 \quad \Rightarrow \quad 2 = 50 \cdot 2^{-t/28} \quad \Rightarrow \quad \frac{2}{50} = 2^{-t/28} \quad \Rightarrow$ $(-t/28)\ln 2 = \ln \frac{1}{25} \quad \Rightarrow \quad t = \left(-28 \ln \frac{1}{25}\right) / \ln 2 \approx 130$ days

(d)

9. (a) If $y(t)$ is the mass (in mg) remaining after t years, then $y(t) = y(0)e^{kt} = 100e^{kt}$.

$y(30) = 100e^{30k} = \frac{1}{2}(100) \quad \Rightarrow \quad e^{30k} = \frac{1}{2} \quad \Rightarrow \quad k = -(\ln 2)/30 \quad \Rightarrow \quad y(t) = 100e^{-(\ln 2)t/30} = 100 \cdot 2^{-t/30}$

(b) $y(100) = 100 \cdot 2^{-100/30} \approx 9.92$ mg

(c) $100e^{-(\ln 2)t/30} = 1 \quad \Rightarrow \quad -(\ln 2)t/30 = \ln \frac{1}{100} \quad \Rightarrow \quad t = -30\,\frac{\ln 0.01}{\ln 2} \approx 199.3$ years

10. (a) If $y(t)$ is the mass after t days and $y(0) = A$, then $y(t) = Ae^{kt}$.
$y(300) = Ae^{300k} = 0.643A \quad \Rightarrow \quad e^{300k} = 0.643 \quad \Rightarrow \quad k = \frac{1}{300}\ln 0.643$. To find the half-life, we set the mass after t days equal to one-half of the original mass. Hence, $Ae^{(1/300)(\ln 0.643)t} = \frac{1}{2}A \quad \Leftrightarrow \quad \frac{1}{300}(\ln 0.643)\,t = \ln \frac{1}{2} \quad \Leftrightarrow$ $t = \dfrac{300 \ln \frac{1}{2}}{\ln 0.643} \approx 471$ days.

(b) $Ae^{(1/300)(\ln 0.643)t} = \frac{1}{3}A \quad \Leftrightarrow \quad \frac{1}{300}(\ln 0.643)\,t = \ln \frac{1}{3} \quad \Leftrightarrow \quad t = \dfrac{300 \ln \frac{1}{3}}{\ln 0.643} \approx 746$ days

11. Let $y(t)$ be the level of radioactivity. Thus, $y(t) = y(0)e^{-kt}$ and k is determined by using the half-life:

$$y(5730) = \tfrac{1}{2}y(0) \quad \Rightarrow \quad y(0)e^{-k(5730)} = \tfrac{1}{2}y(0) \quad \Rightarrow \quad e^{-5730k} = \tfrac{1}{2} \quad \Rightarrow \quad -5730k = \ln \tfrac{1}{2} \quad \Rightarrow \quad k = -\frac{\ln \frac{1}{2}}{5730} = \frac{\ln 2}{5730}.$$

If 74% of the ^{14}C remains, then we know that $y(t) = 0.74y(0) \quad \Rightarrow \quad 0.74 = e^{-t(\ln 2)/5730} \quad \Rightarrow \quad \ln 0.74 = -\dfrac{t \ln 2}{5730} \quad \Rightarrow$

$t = -\dfrac{5730(\ln 0.74)}{\ln 2} \approx 2489 \approx 2500$ years.

12. From Exercise 11, we have the model $y(t) = y(0)e^{-kt}$ with $k = (\ln 2)/5730$. Thus,

$y(68{,}000{,}000) = y(0)e^{-68{,}000{,}000k} \approx y(0) \cdot 0 = 0$. There would be an undetectable amount of ^{14}C remaining for a 68-million-year-old dinosaur.

Now let $y(t) = 0.1\%\,y(0)$, so $0.001y(0) = y(0)e^{-kt} \quad \Rightarrow \quad 0.001 = e^{-kt} \quad \Rightarrow \quad \ln 0.001 = -kt \quad \Rightarrow$

$t = \dfrac{\ln 0.001}{-k} = \dfrac{\ln 0.001}{-(\ln 2)/5730} \approx 57{,}104$, which is the maximum age of a fossil that we could date using ^{14}C.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. Let t measure time since a dinosaur died in millions of years, and let $y(t)$ be the amount of ^{40}K in the dinosaur's bones at time t. Then $y(t) = y(0)e^{-kt}$ and k is determined by the half-life: $y(1250) = \frac{1}{2}y(0) \quad \Rightarrow \quad y(0)e^{-k(1250)} = \frac{1}{2}y(0) \quad \Rightarrow$ $e^{-1250k} = \frac{1}{2} \quad \Rightarrow \quad -1250k = \ln\frac{1}{2} \quad \Rightarrow \quad k = -\dfrac{\ln\frac{1}{2}}{1250} = \dfrac{\ln 2}{1250}$. To determine if a dinosaur dating of 68 million years is possible, we find that $y(68) = y(0)e^{-k(68)} \approx 0.963y(0)$, indicating that about 96% of the ^{40}K is remaining, which is clearly detectable. To determine the maximum age of a fossil by using ^{40}K, we solve $y(t) = 0.1\%y(0)$ for t.

$y(0)e^{-kt} = 0.001y(0) \quad \Leftrightarrow \quad e^{-kt} = 0.001 \quad \Leftrightarrow \quad -kt = \ln 0.001 \quad \Leftrightarrow \quad t = \dfrac{\ln 0.001}{-(\ln 2)/1250} \approx 12{,}457$ million, or 12.457 billion years.

14. From the information given, we know that $\dfrac{dy}{dx} = 2y \quad \Rightarrow \quad y = Ce^{2x}$ by Theorem 2. To calculate C we use the point $(0, 5)$: $5 = Ce^{2(0)} \quad \Rightarrow \quad C = 5$. Thus, the equation of the curve is $y = 5e^{2x}$.

15. (a) Using Newton's Law of Cooling, $\dfrac{dT}{dt} = k(T - T_s)$, we have $\dfrac{dT}{dt} = k(T - 75)$. Now let $y = T - 75$, so $y(0) = T(0) - 75 = 185 - 75 = 110$, so y is a solution of the initial-value problem $dy/dt = ky$ with $y(0) = 110$ and by Theorem 2 we have $y(t) = y(0)e^{kt} = 110e^{kt}$.

$y(30) = 110e^{30k} = 150 - 75 \quad \Rightarrow \quad e^{30k} = \frac{75}{110} = \frac{15}{22} \quad \Rightarrow \quad k = \frac{1}{30}\ln\frac{15}{22}$, so $y(t) = 110e^{\frac{1}{30}t\ln\left(\frac{15}{22}\right)}$ and $y(45) = 110e^{\frac{45}{30}\ln\left(\frac{15}{22}\right)} \approx 62°\text{F}$. Thus, $T(45) \approx 62 + 75 = 137°\text{F}$.

(b) $T(t) = 100 \quad \Rightarrow \quad y(t) = 25$. $\quad y(t) = 110e^{\frac{1}{30}t\ln\left(\frac{15}{22}\right)} = 25 \quad \Rightarrow \quad e^{\frac{1}{30}t\ln\left(\frac{15}{22}\right)} = \frac{25}{110} \quad \Rightarrow \quad \frac{1}{30}t\ln\frac{15}{22} = \ln\frac{25}{110} \quad \Rightarrow$ $t = \dfrac{30\ln\frac{25}{110}}{\ln\frac{15}{22}} \approx 116$ min.

16. Let $T(t)$ be the temperature of the body t hours after 1:30 PM. Then $T(0) = 32.5$ and $T(1) = 30.3$. Using Newton's Law of Cooling, $\dfrac{dT}{dt} = k(T - T_s)$, we have $\dfrac{dT}{dt} = k(T - 20)$. Now let $y = T - 20$, so $y(0) = T(0) - 20 = 32.5 - 20 = 12.5$, so y is a solution to the initial value problem $dy/dt = ky$ with $y(0) = 12.5$ and by Theorem 2 we have $y(t) = y(0)e^{kt} = 12.5e^{kt}$.

$y(1) = 30.3 - 20 \quad \Rightarrow \quad 10.3 = 12.5e^{k(1)} \quad \Rightarrow \quad e^k = \frac{10.3}{12.5} \quad \Rightarrow \quad k = \ln\frac{10.3}{12.5}$. The murder occurred when $y(t) = 37 - 20 \quad \Rightarrow \quad 12.5e^{kt} = 17 \quad \Rightarrow \quad e^{kt} = \frac{17}{12.5} \quad \Rightarrow \quad kt = \ln\frac{17}{12.5} \quad \Rightarrow \quad t = \left(\ln\frac{17}{12.5}\right)/\ln\frac{10.3}{12.5} \approx -1.588$ h ≈ -95 minutes. Thus, the murder took place about 95 minutes before 1:30 PM, or 11:55 AM.

17. $\dfrac{dT}{dt} = k(T - 20)$. Letting $y = T - 20$, we get $\dfrac{dy}{dt} = ky$, so $y(t) = y(0)e^{kt}$. $\quad y(0) = T(0) - 20 = 5 - 20 = -15$, so $y(25) = y(0)e^{25k} = -15e^{25k}$, and $y(25) = T(25) - 20 = 10 - 20 = -10$, so $-15e^{25k} = -10 \quad \Rightarrow \quad e^{25k} = \frac{2}{3}$. Thus,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$25k = \ln\left(\frac{2}{3}\right)$ and $k = \frac{1}{25}\ln\left(\frac{2}{3}\right)$, so $y(t) = y(0)e^{kt} = -15e^{(1/25)\ln(2/3)t}$. More simply, $e^{25k} = \frac{2}{3} \quad\Rightarrow\quad e^k = \left(\frac{2}{3}\right)^{1/25} \quad\Rightarrow$ $e^{kt} = \left(\frac{2}{3}\right)^{t/25} \quad\Rightarrow\quad y(t) = -15\cdot\left(\frac{2}{3}\right)^{t/25}$.

(a) $T(50) = 20 + y(50) = 20 - 15\cdot\left(\frac{2}{3}\right)^{50/25} = 20 - 15\cdot\left(\frac{2}{3}\right)^2 = 20 - \frac{20}{3} = 13.\bar{3}\,^\circ\text{C}$

(b) $15 = T(t) = 20 + y(t) = 20 - 15\cdot\left(\frac{2}{3}\right)^{t/25} \quad\Rightarrow\quad 15\cdot\left(\frac{2}{3}\right)^{t/25} = 5 \quad\Rightarrow\quad \left(\frac{2}{3}\right)^{t/25} = \frac{1}{3} \quad\Rightarrow$

$(t/25)\,\ln\left(\frac{2}{3}\right) = \ln\left(\frac{1}{3}\right) \quad\Rightarrow\quad t = 25\ln\left(\frac{1}{3}\right)/\ln\left(\frac{2}{3}\right) \approx 67.74$ min.

18. $\dfrac{dT}{dt} = k(T-20)$. Let $y = T - 20$. Then $\dfrac{dy}{dt} = ky$, so $y(t) = y(0)e^{kt}$. $\quad y(0) = T(0) - 20 = 95 - 20 = 75$, so $y(t) = 75e^{kt}$. When $T(t) = 70$, $\dfrac{dT}{dt} = -1^\circ\text{C/min}$. Equivalently, $\dfrac{dy}{dt} = -1$ when $y(t) = 50$. Thus, $-1 = \dfrac{dy}{dt} = ky(t) = 50k$ and $50 = y(t) = 75e^{kt}$. The first relation implies $k = -1/50$, so the second relation says $50 = 75e^{-t/50}$. Thus, $e^{-t/50} = \frac{2}{3} \quad\Rightarrow\quad -t/50 = \ln\left(\frac{2}{3}\right) \quad\Rightarrow\quad t = -50\ln\left(\frac{2}{3}\right) \approx 20.27$ min.

19. (a) Let $P(h)$ be the pressure at altitude h. Then $dP/dh = kP \quad\Rightarrow\quad P(h) = P(0)e^{kh} = 101.3e^{kh}$.

$P(1000) = 101.3e^{1000k} = 87.14 \quad\Rightarrow\quad 1000k = \ln\left(\frac{87.14}{101.3}\right) \quad\Rightarrow\quad k = \frac{1}{1000}\ln\left(\frac{87.14}{101.3}\right) \quad\Rightarrow$

$P(h) = 101.3\,e^{\frac{1}{1000}h\ln\left(\frac{87.14}{101.3}\right)}$, so $P(3000) = 101.3e^{3\ln\left(\frac{87.14}{101.3}\right)} \approx 64.5$ kPa.

(b) $P(6187) = 101.3\,e^{\frac{6187}{1000}\ln\left(\frac{87.14}{101.3}\right)} \approx 39.9$ kPa

20. (a) Using $A = A_0\left(1 + \dfrac{r}{n}\right)^{nt}$ with $A_0 = 2500$, $r = 0.045$, and $t = 3$, we have:

(i) Annually: $n = 1$ $\qquad A = 2500\left(1 + \dfrac{0.045}{1}\right)^{1\cdot 3} = \$2852.92$

(ii) Quarterly: $n = 4$ $\qquad A = 2500\left(1 + \dfrac{0.045}{4}\right)^{4\cdot 3} = \$2859.19$

(iii) Monthly: $n = 12$ $\qquad A = 2500\left(1 + \dfrac{0.045}{12}\right)^{12\cdot 3} = \$2860.62$

(iv) Weekly: $n = 52$ $\qquad A = 2500\left(1 + \dfrac{0.045}{52}\right)^{52\cdot 3} = \$2861.17$

(v) Daily: $n = 365$ $\qquad A = 2500\left(1 + \dfrac{0.045}{365}\right)^{365\cdot 3} = \$2861.32$

(vi) Hourly: $n = 365\cdot 24$ $\qquad A = 2500\left(1 + \dfrac{0.045}{365\cdot 24}\right)^{365\cdot 24\cdot 3} = \$2861.34$

(vii) Continuously: $\qquad A = 2500e^{(0.045)^3} = \$2861.34$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b)

21. (a) Using $A = A_0\left(1 + \dfrac{r}{n}\right)^{nt}$ with $A_0 = 4000$, $r = 0.0175$, and $t = 5$, we have:

(i) Annually: $n = 1$ $\quad A = 4000\left(1 + \dfrac{0.0175}{1}\right)^{1\cdot 5} = \$4362.47$

(ii) Semiannually: $n = 2$ $\quad A = 4000\left(1 + \dfrac{0.0175}{2}\right)^{2\cdot 5} = \$4364.11$

(iii) Monthly: $n = 12$ $\quad A = 4000\left(1 + \dfrac{0.0175}{12}\right)^{12\cdot 5} = \$4365.49$

(iv) Weekly: $n = 52$ $\quad A = 4000\left(1 + \dfrac{0.0175}{52}\right)^{52\cdot 5} = \$4365.70$

(v) Daily: $n = 365$ $\quad A = 4000\left(1 + \dfrac{0.0175}{365}\right)^{365\cdot 5} = \$4365.76$

(vi) Continuously: $\quad A = 4000e^{(0.0175)5} = \$4365.77$

(b) $dA/dt = 0.0175A$ and $A(0) = 4000$.

22. (a) $A_0e^{0.03t} = 2A_0 \quad \Leftrightarrow \quad e^{0.03t} = 2 \quad \Leftrightarrow \quad 0.03t = \ln 2 \quad \Leftrightarrow \quad t = \frac{100}{3}\ln 2 \approx 23.10$, so the investment will double in about 23.10 years.

(b) The annual interest rate in $A = A_0(1 + r)^t$ is r. From part (a), we have $A = A_0e^{0.03t}$. These amounts must be equal, so $(1 + r)^t = e^{0.03t} \quad \Rightarrow \quad 1 + r = e^{0.03} \quad \Rightarrow \quad r = e^{0.03} - 1 \approx 0.0305 = 3.05\%$, which is the equivalent annual interest rate.

APPLIED PROJECT Controlling Red Blood Cell Loss During Surgery

1. Let $R(t)$ be the volume of RBCs (in liters) at time t (in hours). Since the total volume of blood is 5 L, the concentration of RBCs is $R/5$. The patient bleeds 2 L of blood in 4 hours, so

$$\frac{dR}{dt} = -\frac{2L}{4h} \cdot \frac{R}{5} = -\frac{1}{10}R$$

From Section 3.8, we know that $dR/dt = kR$ has solution $R(t) = R(0)e^{kt}$. In this case, $R(0) = 45\%$ of $5 = \frac{9}{4}$ and $k = -\frac{1}{10}$, so $R(t) = \frac{9}{4}e^{-t/10}$. At the end of the operation, the volume of RBCs is $R(4) = \frac{9}{4}e^{-0.4} \approx 1.51$ L.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. Let V be the volume of blood that is extracted and replaced with saline solution. Let $R_A(t)$ be the volume of RBCs with the ANH procedure. Then $R_A(0)$ is 45% of $(5-V)$, or $\frac{9}{20}(5-V)$, and hence $R_A(t)=\frac{9}{20}(5-V)e^{-t/10}$. We want $R_A(4)\geq 25\%$ of $5 \quad\Leftrightarrow\quad \frac{9}{20}(5-V)e^{-0.4}\geq\frac{5}{4} \quad\Leftrightarrow\quad 5-V\geq\frac{25}{9}e^{0.4} \quad\Leftrightarrow\quad V\leq 5-\frac{25}{9}e^{0.4}\approx 0.86$ L. To maximize the effect of the ANH procedure, the surgeon should remove 0.86 L of blood and replace it with saline solution.

3. The RBC loss *without* the ANH procedure is $R(0)-R(4)=\frac{9}{4}-\frac{9}{4}e^{-0.4}\approx 0.74$ L. The RBC loss *with* the ANH procedure is $R_A(0)-R_A(4)=\frac{9}{20}(5-V)-\frac{9}{20}(5-V)e^{-0.4}=\frac{9}{20}(5-V)(1-e^{-0.4})$. Now let $V=5-\frac{25}{9}e^{0.4}$ [from Problem 2] to get $R_A(0)-R_A(4)=\frac{9}{20}\left[5-\left(5-\frac{25}{9}e^{0.4}\right)\right](1-e^{0.4})=\frac{9}{20}\cdot\frac{25}{9}e^{0.4}(1-e^{0.4})=\frac{5}{4}(e^{0.4}-1)\approx 0.61$ L. Thus, the ANH procedure reduces the RBC loss by about $0.74-0.61=0.13$ L (about 4.4 fluid ounces).

3.9 Related Rates

1. (a) $V=x^3 \quad\Rightarrow\quad \dfrac{dV}{dt}=\dfrac{dV}{dx}\dfrac{dx}{dt}=3x^2\dfrac{dx}{dt}$

(b) With $\dfrac{dx}{dt}=4$ cm/s and $x=15$ cm, we have $\dfrac{dV}{dt}=3(15)^2\cdot 4=2700$ cm^3/s.

2. (a) $A=\pi r^2 \quad\Rightarrow\quad \dfrac{dA}{dt}=\dfrac{dA}{dr}\dfrac{dr}{dt}=2\pi r\dfrac{dr}{dt}$

(b) With $\dfrac{dr}{dt}=2$ m/s and $r=30$ m, we have $\dfrac{dA}{dt}=2\pi\cdot 30\cdot 2=120\pi$ m^2/s.

3. Let s denote the side of a square. The square's area A is given by $A=s^2$. Differentiating with respect to t gives us $\dfrac{dA}{dt}=2s\dfrac{ds}{dt}$. When $A=16$, $s=4$. Substituting 4 for s and 6 for $\dfrac{ds}{dt}$ gives us $\dfrac{dA}{dt}=2(4)(6)=48$ cm^2/s.

4. $V=\frac{4}{3}\pi r^3 \quad\Rightarrow\quad \dfrac{dV}{dt}=\frac{4}{3}\pi\cdot 3r^2\dfrac{dr}{dt} \quad\Rightarrow\quad \dfrac{dV}{dt}=4\pi\left(\frac{1}{2}\cdot 80\right)^2(4)=25{,}600\pi$ mm^3/s.

5. $S=4\pi r^2 \quad\Rightarrow\quad \dfrac{dS}{dt}=4\pi\cdot 2r\dfrac{dr}{dt} \quad\Rightarrow\quad \dfrac{dS}{dt}=4\pi\cdot 2\cdot 8\cdot 2=128\pi$ cm^2/min.

6. $A=\ell w \quad\Rightarrow\quad \dfrac{dA}{dt}=\ell\cdot\dfrac{dw}{dt}+w\cdot\dfrac{d\ell}{dt}=20(3)+10(8)=140$ cm^2/s.

7. $V=\pi r^2h=\pi(5)^2h=25\pi h \quad\Rightarrow\quad \dfrac{dV}{dt}=25\pi\dfrac{dh}{dt} \quad\Rightarrow\quad 3=25\pi\dfrac{dh}{dt} \quad\Rightarrow\quad \dfrac{dh}{dt}=\dfrac{3}{25\pi}$ m/min.

8. (a) $A=\frac{1}{2}ab\sin\theta \quad\Rightarrow\quad \dfrac{dA}{dt}=\frac{1}{2}ab\cos\theta\dfrac{d\theta}{dt}=\frac{1}{2}(2)(3)\left(\cos\frac{\pi}{3}\right)(0.2)=3\left(\frac{1}{2}\right)(0.2)=0.3$ cm^2/min.

(b) $A=\frac{1}{2}ab\sin\theta \quad\Rightarrow$

$$\frac{dA}{dt}=\frac{1}{2}a\left(b\cos\theta\frac{d\theta}{dt}+\sin\theta\frac{db}{dt}\right)=\tfrac{1}{2}(2)\left[3\left(\cos\tfrac{\pi}{3}\right)(0.2)+\left(\sin\tfrac{\pi}{3}\right)(1.5)\right]$$
$$=3\left(\tfrac{1}{2}\right)(0.2)+\tfrac{1}{2}\sqrt{3}\left(\tfrac{3}{2}\right)=0.3+\tfrac{3}{4}\sqrt{3}\text{ cm}^2/\text{min}\quad[\approx 1.6]$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $A = \frac{1}{2}ab\sin\theta \quad \Rightarrow$

$$\frac{dA}{dt} = \frac{1}{2}\left(\frac{da}{dt}\,b\sin\theta + a\,\frac{db}{dt}\sin\theta + ab\cos\theta\,\frac{d\theta}{dt}\right) \qquad \text{[by Exercise 3.2.63(a)]}$$

$$= \tfrac{1}{2}\left[(2.5)(3)\left(\tfrac{1}{2}\sqrt{3}\right) + (2)(1.5)\left(\tfrac{1}{2}\sqrt{3}\right) + (2)(3)\left(\tfrac{1}{2}\right)(0.2)\right]$$

$$= \left(\tfrac{15}{8}\sqrt{3} + \tfrac{3}{4}\sqrt{3} + 0.3\right) = \left(\tfrac{21}{8}\sqrt{3} + 0.3\right)\ \text{cm}^2/\text{min} \quad [\approx 4.85]$$

Note how this answer relates to the answer in part (a) [θ changing] and part (b) [b and θ changing].

9. (a) $\dfrac{d}{dt}\left(4x^2 + 9y^2\right) = \dfrac{d}{dt}(25) \quad \Rightarrow \quad 8x\,\dfrac{dx}{dt} + 18y\,\dfrac{dy}{dt} = 0 \quad \Rightarrow \quad 4x\,\dfrac{dx}{dt} + 9y\,\dfrac{dy}{dt} = 0 \quad \Rightarrow$

$4(2)\,\dfrac{dx}{dt} + 9(1)\cdot\frac{1}{3} = 0 \quad \Rightarrow \quad 8\,\dfrac{dx}{dt} + 3 = 0 \quad \Rightarrow \quad \dfrac{dx}{dt} = -\dfrac{3}{8}$

(b) $4x\,\dfrac{dx}{dt} + 9y\,\dfrac{dy}{dt} = 0 \quad \Rightarrow \quad 4(-2)(3) + 9(1)\cdot\dfrac{dy}{dt} = 0 \quad \Rightarrow \quad -24 + 9\,\dfrac{dy}{dt} = 0 \quad \Rightarrow \quad \dfrac{dy}{dt} = \dfrac{24}{9} = \dfrac{8}{3}$

10. $\dfrac{d}{dt}\left(x^2 + y^2 + z^2\right) = \dfrac{d}{dt}(9) \quad \Rightarrow \quad 2x\,\dfrac{dx}{dt} + 2y\,\dfrac{dy}{dt} + 2z\,\dfrac{dz}{dt} = 0 \quad \Rightarrow \quad x\,\dfrac{dx}{dt} + y\,\dfrac{dy}{dt} + z\,\dfrac{dz}{dt} = 0.$

If $\dfrac{dx}{dt} = 5$, $\dfrac{dy}{dt} = 4$ and $(x, y, z) = (2, 2, 1)$, then $2(5) + 2(4) + 1\,\dfrac{dz}{dt} = 0 \quad \Rightarrow \quad \dfrac{dz}{dt} = -18.$

11. $w = w_0\left(\dfrac{3960}{3960 + h}\right)^2 = w_0 \cdot 3960^2(3960 + h)^{-2} \quad \Rightarrow \quad \dfrac{dw}{dt} = w_0 \cdot 3960^2(-2)(3960 + h)^{-3} \cdot \dfrac{dh}{dt}$. Then $w_0 = 130$ lb,

$h = 40$ mi, and $dh/dt = 12$ mi/s $\quad \Rightarrow \quad \dfrac{dw}{dt} = 130 \cdot 3960^2(-2)(3960 + 40)^{-3}(12) = -0.764478 \approx -0.7645$ lb/s.

12. $\dfrac{d}{dt}(xy) = \dfrac{d}{dt}(8) \quad \Rightarrow \quad x\,\dfrac{dy}{dt} + y\,\dfrac{dx}{dt} = 0.$ If $\dfrac{dy}{dt} = -3$ cm/s and $(x, y) = (4, 2)$, then $4(-3) + 2\,\dfrac{dx}{dt} = 0 \quad \Rightarrow$

$\dfrac{dx}{dt} = 6$. Thus, the x-coordinate is increasing at a rate of 6 cm/s.

13. (a) Given: a plane flying horizontally at an altitude of 1 mi and a speed of 500 mi/h passes directly over a radar station.

If we let t be time (in hours) and x be the horizontal distance traveled by the plane (in mi), then we are given that $dx/dt = 500$ mi/h.

(b) Unknown: the rate at which the distance from the plane to the station is increasing when it is 2 mi from the station. If we let y be the distance from the plane to the station, then we want to find dy/dt when $y = 2$ mi.

(c)

(d) By the Pythagorean Theorem, $y^2 = x^2 + 1 \quad \Rightarrow \quad 2y\,(dy/dt) = 2x\,(dx/dt)$.

(e) $\dfrac{dy}{dt} = \dfrac{x}{y}\dfrac{dx}{dt} = \dfrac{x}{y}(500)$. Since $y^2 = x^2 + 1$, when $y = 2$, $x = \sqrt{3}$, so $\dfrac{dy}{dt} = \dfrac{\sqrt{3}}{2}(500) = 250\sqrt{3} \approx 433$ mi/h.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. (a) Given: the rate of decrease of the surface area is 1 cm^2/min. If we let t be time (in minutes) and S be the surface area (in cm^2), then we are given that $dS/dt = -1\ \text{cm}^2/\text{s}$.

(b) Unknown: the rate of decrease of the diameter when the diameter is 10 cm. If we let x be the diameter, then we want to find dx/dt when $x = 10$ cm.

(c)

(d) If the radius is r and the diameter $x = 2r$, then $r = \frac{1}{2}x$ and

$$S = 4\pi r^2 = 4\pi\left(\tfrac{1}{2}x\right)^2 = \pi x^2 \quad \Rightarrow \quad \frac{dS}{dt} = \frac{dS}{dx}\frac{dx}{dt} = 2\pi x\,\frac{dx}{dt}.$$

(e) $-1 = \dfrac{dS}{dt} = 2\pi x\,\dfrac{dx}{dt} \quad \Rightarrow \quad \dfrac{dx}{dt} = -\dfrac{1}{2\pi x}$. When $x = 10$, $\dfrac{dx}{dt} = -\dfrac{1}{20\pi}$. So the rate of decrease is $\dfrac{1}{20\pi}$ cm/min.

15. (a) Given: a man 6 ft tall walks away from a street light mounted on a 15-ft-tall pole at a rate of 5 ft/s. If we let t be time (in s) and x be the distance from the pole to the man (in ft), then we are given that $dx/dt = 5$ ft/s.

(b) Unknown: the rate at which the tip of his shadow is moving when he is 40 ft from the pole. If we let y be the distance from the man to the tip of his shadow (in ft), then we want to find $\dfrac{d}{dt}(x+y)$ when $x = 40$ ft.

(c)

(d) By similar triangles, $\dfrac{15}{6} = \dfrac{x+y}{y} \quad \Rightarrow \quad 15y = 6x + 6y \quad \Rightarrow \quad 9y = 6x \quad \Rightarrow \quad y = \frac{2}{3}x$.

(e) The tip of the shadow moves at a rate of $\dfrac{d}{dt}(x+y) = \dfrac{d}{dt}\left(x + \dfrac{2}{3}x\right) = \dfrac{5}{3}\dfrac{dx}{dt} = \frac{5}{3}(5) = \frac{25}{3}$ ft/s.

16. (a) Given: at noon, ship A is 150 km west of ship B; ship A is sailing east at 35 km/h, and ship B is sailing north at 25 km/h. If we let t be time (in hours), x be the distance traveled by ship A (in km), and y be the distance traveled by ship B (in km), then we are given that $dx/dt = 35$ km/h and $dy/dt = 25$ km/h.

(b) Unknown: the rate at which the distance between the ships is changing at 4:00 PM. If we let z be the distance between the ships, then we want to find dz/dt when $t = 4$ h.

(c)

(d) $z^2 = (150 - x)^2 + y^2 \quad \Rightarrow \quad 2z\,\dfrac{dz}{dt} = 2(150 - x)\left(-\dfrac{dx}{dt}\right) + 2y\,\dfrac{dy}{dt}$

(e) At 4:00 PM, $x = 4(35) = 140$ and $y = 4(25) = 100 \quad \Rightarrow \quad z = \sqrt{(150-140)^2 + 100^2} = \sqrt{10{,}100}$.

So $\dfrac{dz}{dt} = \dfrac{1}{z}\left[(x - 150)\,\dfrac{dx}{dt} + y\,\dfrac{dy}{dt}\right] = \dfrac{-10(35) + 100(25)}{\sqrt{10{,}100}} = \dfrac{215}{\sqrt{101}} \approx 21.4$ km/h.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17.

We are given that $\dfrac{dx}{dt} = 60$ mi/h and $\dfrac{dy}{dt} = 25$ mi/h. $z^2 = x^2 + y^2 \quad \Rightarrow$

$2z\,\dfrac{dz}{dt} = 2x\,\dfrac{dx}{dt} + 2y\,\dfrac{dy}{dt} \quad \Rightarrow \quad z\,\dfrac{dz}{dt} = x\,\dfrac{dx}{dt} + y\,\dfrac{dy}{dt} \quad \Rightarrow \quad \dfrac{dz}{dt} = \dfrac{1}{z}\left(x\,\dfrac{dx}{dt} + y\,\dfrac{dy}{dt}\right)$.

After 2 hours, $x = 2\,(60) = 120$ and $y = 2\,(25) = 50 \quad \Rightarrow \quad z = \sqrt{120^2 + 50^2} = 130$,

so $\dfrac{dz}{dt} = \dfrac{1}{z}\left(x\,\dfrac{dx}{dt} + y\,\dfrac{dy}{dt}\right) = \dfrac{120(60) + 50(25)}{130} = 65$ mi/h.

18.

We are given that $\dfrac{dx}{dt} = 1.6$ m/s. By similar triangles, $\dfrac{y}{12} = \dfrac{2}{x} \quad \Rightarrow \quad y = \dfrac{24}{x} \quad \Rightarrow$

$\dfrac{dy}{dt} = -\dfrac{24}{x^2}\dfrac{dx}{dt} = -\dfrac{24}{x^2}(1.6)$. When $x = 8$, $\dfrac{dy}{dt} = -\dfrac{24(1.6)}{64} = -0.6$ m/s, so the shadow is decreasing at a rate of 0.6 m/s.

19.

We are given that $\dfrac{dx}{dt} = 4$ ft/s and $\dfrac{dy}{dt} = 5$ ft/s. $z^2 = (x + y)^2 + 500^2 \quad \Rightarrow$

$2z\,\dfrac{dz}{dt} = 2(x + y)\left(\dfrac{dx}{dt} + \dfrac{dy}{dt}\right)$. 15 minutes after the woman starts, we have

$x = (4 \text{ ft/s})(20 \text{ min})(60 \text{ s/min}) = 4800$ ft and $y = 5 \cdot 15 \cdot 60 = 4500 \quad \Rightarrow$

$z = \sqrt{(4800 + 4500)^2 + 500^2} = \sqrt{86{,}740{,}000}$, so

$\dfrac{dz}{dt} = \dfrac{x + y}{z}\left(\dfrac{dx}{dt} + \dfrac{dy}{dt}\right) = \dfrac{4800 + 4500}{\sqrt{86{,}740{,}000}}(4 + 5) = \dfrac{837}{\sqrt{8674}} \approx 8.99$ ft/s.

20. We are given that $\dfrac{dx}{dt} = 24$ ft/s.

(a)

$y^2 = (90 - x)^2 + 90^2 \quad \Rightarrow \quad 2y\,\dfrac{dy}{dt} = 2(90 - x)\left(-\dfrac{dx}{dt}\right)$. When $x = 45$,

$y = \sqrt{45^2 + 90^2} = 45\sqrt{5}$, so $\dfrac{dy}{dt} = \dfrac{90 - x}{y}\left(-\dfrac{dx}{dt}\right) = \dfrac{45}{45\sqrt{5}}(-24) = -\dfrac{24}{\sqrt{5}}$,

so the distance from second base is decreasing at a rate of $\frac{24}{\sqrt{5}} \approx 10.7$ ft/s.

(b) Due to the symmetric nature of the problem in part (a), we expect to get the same answer—and we do.

$z^2 = x^2 + 90^2 \quad \Rightarrow \quad 2z\,\dfrac{dz}{dt} = 2x\,\dfrac{dx}{dt}$. When $x = 45$, $z = 45\sqrt{5}$, so $\dfrac{dz}{dt} = \dfrac{45}{45\sqrt{5}}(24) = \dfrac{24}{\sqrt{5}} \approx 10.7$ ft/s.

21. $A = \frac{1}{2}bh$, where b is the base and h is the altitude. We are given that $\dfrac{dh}{dt} = 1$ cm/min and $\dfrac{dA}{dt} = 2$ cm^2/min. Using the Product Rule, we have $\dfrac{dA}{dt} = \dfrac{1}{2}\left(b\,\dfrac{dh}{dt} + h\,\dfrac{db}{dt}\right)$. When $h = 10$ and $A = 100$, we have $100 = \frac{1}{2}b(10) \quad \Rightarrow \quad \frac{1}{2}b = 10 \quad \Rightarrow$

$b = 20$, so $2 = \dfrac{1}{2}\left(20 \cdot 1 + 10\,\dfrac{db}{dt}\right) \quad \Rightarrow \quad 4 = 20 + 10\,\dfrac{db}{dt} \quad \Rightarrow \quad \dfrac{db}{dt} = \dfrac{4 - 20}{10} = -1.6$ cm/min.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22.

Given $\dfrac{dy}{dt} = -1$ m/s, find $\dfrac{dx}{dt}$ when $x = 8$ m. $y^2 = x^2 + 1 \;\Rightarrow\; 2y\dfrac{dy}{dt} = 2x\dfrac{dx}{dt} \;\Rightarrow$ $\dfrac{dx}{dt} = \dfrac{y}{x}\dfrac{dy}{dt} = -\dfrac{y}{x}$. When $x = 8$, $y = \sqrt{65}$, so $\dfrac{dx}{dt} = -\dfrac{\sqrt{65}}{8}$. Thus, the boat approaches the dock at $\dfrac{\sqrt{65}}{8} \approx 1.01$ m/s.

23. Let x be the distance (in meters) the first dropped stone has traveled, and let y be the distance (in meters) the stone dropped one second later has traveled. Let t be the time (in seconds) since the woman drops the second stone. Using $d = 4.9t^2$, we have $x = 4.9(t+1)^2$ and $y = 4.9t^2$. Let z be the distance between the stones. Then $z = x - y$ and we have $\dfrac{dz}{dt} = \dfrac{dx}{dt} - \dfrac{dy}{dt} \;\Rightarrow\; \dfrac{dz}{dt} = 9.8(t+1) - 9.8t = 9.8$ m/s.

24. *Given:* Two men 10 m apart each drop a stone, the second one, one minute after the first. Let x be the distance (in meters) the first dropped stone has traveled, and let y be the distance (in meters) the second stone has traveled. Let t be the time (in seconds) since the man drops the second stone. Using $d = 4.9t^2$, we have

$x = 4.9(t+1)^2$ and $y = 4.9t^2$. Let z be the vertical distance between the stones. Then $z = x - y \;\Rightarrow$ $\dfrac{dz}{dt} = \dfrac{dx}{dt} - \dfrac{dy}{dt} \;\Rightarrow\; \dfrac{dz}{dt} = 9.8(t+1) - 9.8t = 9.8$ m/s.

By the Pythagorean Theorem, $w^2 = 10^2 + z^2$. Differentiating with respect to t, we obtain $2w\dfrac{dw}{dt} = 2z\dfrac{dz}{dt} \;\Rightarrow\; \dfrac{dw}{dt} = \dfrac{z\,(dz/dt)}{w}$. One second after the second stone is dropped, $t = 1$, so $z = x - y = 4.9(1+1)^2 - 4.9(1)^2 = 14.7$ m, and $w = \sqrt{10^2 + (14.7)^2} = \sqrt{316.09}$, so $\dfrac{dw}{dt} = \dfrac{14.7\,(9.8)}{\sqrt{316.09}} \approx 8.10$ m/s.

25. If $C =$ the rate at which water is pumped in, then $\dfrac{dV}{dt} = C - 10{,}000$, where $V = \frac{1}{3}\pi r^2 h$ is the volume at time t. By similar triangles, $\dfrac{r}{2} = \dfrac{h}{6} \;\Rightarrow\; r = \dfrac{1}{3}h \;\Rightarrow$ $V = \frac{1}{3}\pi\left(\frac{1}{3}h\right)^2 h = \frac{\pi}{27}h^3 \;\Rightarrow\; \dfrac{dV}{dt} = \dfrac{\pi}{9}h^2\dfrac{dh}{dt}$. When $h = 200$ cm, $\dfrac{dh}{dt} = 20$ cm/min, so $C - 10{,}000 = \dfrac{\pi}{9}(200)^2(20) \;\Rightarrow\; C = 10{,}000 + \dfrac{800{,}000}{9}\pi \approx 289{,}253$ cm^3/min.

26. The distance z of the particle to the origin is given by $z = \sqrt{x^2 + y^2}$, so $z^2 = x^2 + [2\sin(\pi x/2)]^2 \;\Rightarrow$ $2z\dfrac{dz}{dt} = 2x\dfrac{dx}{dt} + 4\cdot 2\sin\left(\dfrac{\pi}{2}x\right)\cos\left(\dfrac{\pi}{2}x\right)\cdot\dfrac{\pi}{2}\dfrac{dx}{dt} \;\Rightarrow\; z\dfrac{dz}{dt} = x\dfrac{dx}{dt} + 2\pi\sin\left(\dfrac{\pi}{2}x\right)\cos\left(\dfrac{\pi}{2}x\right)\dfrac{dx}{dt}$. When $(x, y) = \left(\dfrac{1}{3}, 1\right)$, $z = \sqrt{\left(\dfrac{1}{3}\right)^2 + 1^2} = \sqrt{\dfrac{10}{9}} = \dfrac{1}{3}\sqrt{10}$, so $\dfrac{1}{3}\sqrt{10}\dfrac{dz}{dt} = \dfrac{1}{3}\sqrt{10} + 2\pi\,\sin\dfrac{\pi}{6}\,\cos\dfrac{\pi}{6}\cdot\sqrt{10} \;\Rightarrow$ $\dfrac{1}{3}\dfrac{dz}{dt} = \dfrac{1}{3} + 2\pi\left(\dfrac{1}{2}\right)\left(\dfrac{1}{2}\sqrt{3}\right) \;\Rightarrow\; \dfrac{dz}{dt} = 1 + \dfrac{3\sqrt{3}\,\pi}{2}$ cm/s.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27.

The figure is labeled in meters. The area A of a trapezoid is $\frac{1}{2}(\text{base}_1 + \text{base}_2)(\text{height})$, and the volume V of the 10-meter-long trough is $10A$. Thus, the volume of the trapezoid with height h is $V = (10)\frac{1}{2}[0.3 + (0.3 + 2a)]h$.

By similar triangles, $\dfrac{a}{h} = \dfrac{0.25}{0.5} = \dfrac{1}{2}$, so $2a = h \;\Rightarrow\; V = 5(0.6 + h)h = 3h + 5h^2$.

Now $\dfrac{dV}{dt} = \dfrac{dV}{dh}\dfrac{dh}{dt} \;\Rightarrow\; 0.2 = (3 + 10h)\dfrac{dh}{dt} \;\Rightarrow\; \dfrac{dh}{dt} = \dfrac{0.2}{3 + 10h}$. When $h = 0.3$,

$\dfrac{dh}{dt} = \dfrac{0.2}{3 + 10(0.3)} = \dfrac{0.2}{6}$ m/min $= \dfrac{1}{30}$ m/min or $\dfrac{10}{3}$ cm/min.

28. By similar triangles, $\dfrac{3}{1} = \dfrac{b}{h}$, so $b = 3h$. The trough has volume

$V = \frac{1}{2}bh(10) = 5(3h)h = 15h^2 \;\Rightarrow\; 12 = \dfrac{dV}{dt} = 30h\,\dfrac{dh}{dt} \;\Rightarrow\; \dfrac{dh}{dt} = \dfrac{2}{5h}$.

When $h = \frac{1}{2}$, $\dfrac{dh}{dt} = \dfrac{2}{5 \cdot \frac{1}{2}} = \dfrac{4}{5}$ ft/min.

29. We are given that $\dfrac{dV}{dt} = 30$ ft^3/min. $V = \dfrac{1}{3}\pi r^2 h = \dfrac{1}{3}\pi\left(\dfrac{h}{2}\right)^2 h = \dfrac{\pi h^3}{12} \;\Rightarrow$

$\dfrac{dV}{dt} = \dfrac{dV}{dh}\dfrac{dh}{dt} \;\Rightarrow\; 30 = \dfrac{\pi h^2}{4}\dfrac{dh}{dt} \;\Rightarrow\; \dfrac{dh}{dt} = \dfrac{120}{\pi h^2}$.

When $h = 10$ ft, $\dfrac{dh}{dt} = \dfrac{120}{10^2\pi} = \dfrac{6}{5\pi} \approx 0.38$ ft/min.

30. The figure is drawn without the top 3 feet.

$V = \frac{1}{2}(b + 12)h(20) = 10(b + 12)h$ and, from similar triangles,

$\dfrac{x}{h} = \dfrac{6}{6}$ and $\dfrac{y}{h} = \dfrac{16}{6} = \dfrac{8}{3}$, so $b = x + 12 + y = h + 12 + \dfrac{8h}{3} = 12 + \dfrac{11h}{3}$.

Thus, $V = 10\left(24 + \dfrac{11h}{3}\right)h = 240h + \dfrac{110h^2}{3}$ and so $0.8 = \dfrac{dV}{dt} = \left(240 + \dfrac{220}{3}h\right)\dfrac{dh}{dt}$.

When $h = 5$, $\dfrac{dh}{dt} = \dfrac{0.8}{240 + 5(220/3)} = \dfrac{3}{2275} \approx 0.00132$ ft/min.

31. The area A of an equilateral triangle with side s is given by $A = \frac{1}{4}\sqrt{3}\,s^2$.

$\dfrac{dA}{dt} = \frac{1}{4}\sqrt{3} \cdot 2s\,\dfrac{ds}{dt} = \frac{1}{4}\sqrt{3} \cdot 2(30)(10) = 150\sqrt{3}$ cm^2/min.

32. We are given $dx/dt = 8$ ft/s. $\cot\theta = \dfrac{x}{100} \;\Rightarrow\; x = 100\cot\theta \;\Rightarrow$

$\dfrac{dx}{dt} = -100\csc^2\theta\,\dfrac{d\theta}{dt} \;\Rightarrow\; \dfrac{d\theta}{dt} = -\dfrac{\sin^2\theta}{100} \cdot 8$. When $y = 200$, $\sin\theta = \dfrac{100}{200} = \dfrac{1}{2} \;\Rightarrow$

$\dfrac{d\theta}{dt} = -\dfrac{(1/2)^2}{100} \cdot 8 = -\dfrac{1}{50}$ rad/s. The angle is decreasing at a rate of $\frac{1}{50}$ rad/s.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33.

Let t be the time, in seconds, after the drone passes directly over the car. Given $\dfrac{dx}{dt} = 20$ m/s, $x = 20t$ m, $\dfrac{dy}{dt} = 6$ m/s, and $y = 6t$ m, find $\dfrac{dz}{dt}$ when $t = 5$.

By the Pythagorean Theorem, $w^2 = x^2 + y^2$ and $z^2 = 25^2 + w^2$. This gives

$$z^2 = 25^2 + x^2 + y^2 \quad \Rightarrow$$

$$2z\,\frac{dz}{dt} = 2x\,\frac{dx}{dt} + 2y\,\frac{dy}{dt} \quad \Rightarrow \quad \frac{dz}{dt} = \frac{x\,(dx/dt) + y\,(dy/dt)}{z}$$

When $t = 5$, $x = 20(5) = 100$ and $y = 6(5) = 30$, so $z^2 = 25^2 + 100^2 + 30^2 \quad \Rightarrow \quad z = \sqrt{11{,}525}$ m.

$$\frac{dz}{dt} = \frac{100(20) + 30(6)}{\sqrt{11{,}525}} \approx 20.3 \text{ m/s}.$$

34. The area A of a sector of a circle with radius r and angle θ is given by $A = \frac{1}{2}r^2\theta$. Here r is constant and θ varies, so $\dfrac{dA}{dt} = \dfrac{1}{2}r^2\,\dfrac{d\theta}{dt}$. The minute hand rotates through $360° = 2\pi$ radians each hour, so $\dfrac{d\theta}{dt} = 2\pi$ and $\dfrac{dA}{dt} = \dfrac{1}{2}r^2(2\pi) = \pi r^2$ cm^2/h. This answer makes sense because the minute hand sweeps through the full area of a circle, πr^2, each hour.

35. $\cos\theta = \dfrac{x}{10} \quad \Rightarrow \quad -\sin\theta\,\dfrac{d\theta}{dt} = \dfrac{1}{10}\,\dfrac{dx}{dt}$. From Example 2, $\dfrac{dx}{dt} = 4$ and when $x = 6$, $y = 8$, so $\sin\theta = \dfrac{8}{10}$.

Thus, $-\dfrac{8}{10}\,\dfrac{d\theta}{dt} = \dfrac{1}{10}(4) \quad \Rightarrow \quad \dfrac{d\theta}{dt} = -\dfrac{1}{2}$ rad/s.

36. According to the model in Example 2, $\dfrac{dy}{dt} = -\dfrac{x}{y}\,\dfrac{dx}{dt} \to -\infty$ as $y \to 0$, which doesn't make physical sense. For example, the model predicts that for sufficiently small y, the tip of the ladder moves at a speed greater than the speed of light. Therefore the model is not appropriate for small values of y. What actually happens is that the tip of the ladder leaves the wall at some point in its descent. For a discussion of the true situation see the article "The Falling Ladder Paradox" by Paul Scholten and Andrew Simoson in *The College Mathematics Journal,* 27, (1), January 1996, pages 49–54. Also see "On Mathematical and Physical Ladders" by M. Freeman and P. Palffy-Muhoray in the *American Journal of Physics,* 53 (3), March 1985, pages 276–277.

37. Differentiating both sides of $PV = C$ with respect to t and using the Product Rule gives us $P\,\dfrac{dV}{dt} + V\,\dfrac{dP}{dt} = 0 \quad \Rightarrow$ $\dfrac{dV}{dt} = -\dfrac{V}{P}\,\dfrac{dP}{dt}$. When $V = 600$, $P = 150$ and $\dfrac{dP}{dt} = 20$, so we have $\dfrac{dV}{dt} = -\dfrac{600}{150}(20) = -80$. Thus, the volume is decreasing at a rate of 80 cm^3/min.

38. The volume of a hemisphere is $\frac{2}{3}\pi r^3$, so the volume of a hemispherical basin of radius 30 cm is $\frac{2}{3}\pi(30)^3 = 18{,}000\pi$ cm^3. If the basin is half full, then $V = \pi\left(rh^2 - \frac{1}{3}h^3\right) \quad \Rightarrow \quad 9000\pi = \pi\left(30h^2 - \frac{1}{3}h^3\right) \quad \Rightarrow \quad \frac{1}{3}h^3 - 30h^2 + 9000 = 0 \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$h = H \approx 19.58$ [from a graph or numerical rootfinder; the other two solutions are less than 0 and greater than 30].

$V = \pi\left(30h^2 - \frac{1}{3}h^3\right) \;\Rightarrow\; \dfrac{dV}{dt} = \pi\left(60h\,\dfrac{dh}{dt} - h^2\,\dfrac{dh}{dt}\right) \;\Rightarrow\; \left(2\,\dfrac{\text{L}}{\text{min}}\right)\left(1000\,\dfrac{\text{cm}^3}{\text{L}}\right) = \pi(60h - h^2)\,\dfrac{dh}{dt} \;\Rightarrow$

$\dfrac{dh}{dt} = \dfrac{2000}{\pi(60H - H^2)} \approx 0.804 \text{ cm/min}.$

39. With $R_1 = 80$ and $R_2 = 100$, $\dfrac{1}{R} = \dfrac{1}{R_1} + \dfrac{1}{R_2} = \dfrac{1}{80} + \dfrac{1}{100} = \dfrac{180}{8000} = \dfrac{9}{400}$, so $R = \dfrac{400}{9}$. Differentiating $\dfrac{1}{R} = \dfrac{1}{R_1} + \dfrac{1}{R_2}$ with respect to t, we have $-\dfrac{1}{R^2}\dfrac{dR}{dt} = -\dfrac{1}{R_1^2}\dfrac{dR_1}{dt} - \dfrac{1}{R_2^2}\dfrac{dR_2}{dt} \;\Rightarrow\; \dfrac{dR}{dt} = R^2\left(\dfrac{1}{R_1^2}\dfrac{dR_1}{dt} + \dfrac{1}{R_2^2}\dfrac{dR_2}{dt}\right)$. When $R_1 = 80$ and $R_2 = 100$, $\dfrac{dR}{dt} = \dfrac{400^2}{9^2}\left[\dfrac{1}{80^2}(0.3) + \dfrac{1}{100^2}(0.2)\right] = \dfrac{107}{810} \approx 0.132\ \Omega/\text{s}$.

40. $PV^{1.4} = C \;\Rightarrow\; P \cdot 1.4V^{0.4}\,\dfrac{dV}{dt} + V^{1.4}\,\dfrac{dP}{dt} = 0 \;\Rightarrow\; \dfrac{dV}{dt} = -\dfrac{V^{1.4}}{P \cdot 1.4V^{0.4}}\,\dfrac{dP}{dt} = -\dfrac{V}{1.4P}\,\dfrac{dP}{dt}$.

When $V = 400$, $P = 80$ and $\dfrac{dP}{dt} = -10$, so we have $\dfrac{dV}{dt} = -\dfrac{400}{1.4(80)}(-10) = \dfrac{250}{7}$. Thus, the volume is increasing at a rate of $\frac{250}{7} \approx 36 \text{ cm}^3/\text{min}$.

41.

We are given that $\dfrac{dx}{dt} = 40$ mi/h and $\dfrac{dy}{dt} = 60$ mi/h. By the Law of Cosines,

$z^2 = x^2 + y^2 - 2xy\cos 60^\circ = x^2 + y^2 - xy \;\Rightarrow\; 2z\,\dfrac{dz}{dt} = 2x\,\dfrac{dx}{dt} + 2y\,\dfrac{dy}{dt} - x\,\dfrac{dy}{dt} - y\,\dfrac{dx}{dt}$.

At $t = \frac{1}{2}$ h, we have $x = 40\left(\frac{1}{2}\right) = 20$ and $y = 60\left(\frac{1}{2}\right) = 30 \;\Rightarrow$

$z^2 = 20^2 + 30^2 - 20(30) = 700 \;\Rightarrow\; z = \sqrt{700}$ and

$\dfrac{dz}{dt} = \dfrac{2(20)(40) + 2(30)(60) - 20(60) - 30(40)}{2\sqrt{700}} = \dfrac{2800}{2\sqrt{700}} \approx 52.9$ mi/h.

42. We want to find $\dfrac{dB}{dt}$ when $L = 18$ using $B = 0.007W^{2/3}$ and $W = 0.12L^{2.53}$.

$$\frac{dB}{dt} = \frac{dB}{dW}\frac{dW}{dL}\frac{dL}{dt} = \left(0.007 \cdot \tfrac{2}{3}W^{-1/3}\right)(0.12 \cdot 2.53 \cdot L^{1.53})\left(\frac{20 - 15}{10{,}000{,}000}\right)$$

$$= \left[0.007 \cdot \tfrac{2}{3}(0.12 \cdot 18^{2.53})^{-1/3}\right]\left(0.12 \cdot 2.53 \cdot 18^{1.53}\right)\left(\frac{5}{10^7}\right) \approx 1.045 \times 10^{-8} \text{ g/yr}$$

43. We are given $d\theta/dt = 2^\circ/\text{min} = \frac{\pi}{90}$ rad/min. By the Law of Cosines,

$x^2 = 12^2 + 15^2 - 2(12)(15)\cos\theta = 369 - 360\cos\theta \;\Rightarrow$

$2x\,\dfrac{dx}{dt} = 360\sin\theta\,\dfrac{d\theta}{dt} \;\Rightarrow\; \dfrac{dx}{dt} = \dfrac{180\sin\theta}{x}\,\dfrac{d\theta}{dt}$. When $\theta = 60^\circ$,

$x = \sqrt{369 - 360\cos 60^\circ} = \sqrt{189} = 3\sqrt{21}$, so $\dfrac{dx}{dt} = \dfrac{180\sin 60^\circ}{3\sqrt{21}}\,\dfrac{\pi}{90} = \dfrac{\pi\sqrt{3}}{3\sqrt{21}} = \dfrac{\sqrt{7}\,\pi}{21} \approx 0.396$ m/min.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. Using Q for the origin, we are given $\dfrac{dx}{dt} = -2$ ft/s and need to find $\dfrac{dy}{dt}$ when $x = -5$.

Using the Pythagorean Theorem twice, we have $\sqrt{x^2+12^2} + \sqrt{y^2+12^2} = 39$, the total length of the rope. Differentiating with respect to t, we get

$$\frac{x}{\sqrt{x^2+12^2}}\frac{dx}{dt} + \frac{y}{\sqrt{y^2+12^2}}\frac{dy}{dt} = 0, \text{ so } \frac{dy}{dt} = -\frac{x\sqrt{y^2+12^2}}{y\sqrt{x^2+12^2}}\frac{dx}{dt}.$$

Now when $x = -5$, $39 = \sqrt{(-5)^2+12^2} + \sqrt{y^2+12^2} = 13 + \sqrt{y^2+12^2} \quad \Leftrightarrow \quad \sqrt{y^2+12^2} = 26$, and

$y = \sqrt{26^2-12^2} = \sqrt{532}$. So when $x = -5$, $\dfrac{dy}{dt} = -\dfrac{(-5)(26)}{\sqrt{532}\,(13)}(-2) = -\dfrac{10}{\sqrt{133}} \approx -0.87$ ft/s.

So cart B is moving towards Q at about 0.87 ft/s.

45. (a) By the Pythagorean Theorem, $4000^2 + y^2 = \ell^2$. Differentiating with respect to t,

we obtain $2y\dfrac{dy}{dt} = 2\ell\dfrac{d\ell}{dt}$. We know that $\dfrac{dy}{dt} = 600$ ft/s, so when $y = 3000$ ft,

$\ell = \sqrt{4000^2+3000^2} = \sqrt{25{,}000{,}000} = 5000$ ft

and $\dfrac{d\ell}{dt} = \dfrac{y}{\ell}\dfrac{dy}{dt} = \dfrac{3000}{5000}(600) = \dfrac{1800}{5} = 360$ ft/s.

(b) Here $\tan\theta = \dfrac{y}{4000} \quad \Rightarrow \quad \dfrac{d}{dt}(\tan\theta) = \dfrac{d}{dt}\left(\dfrac{y}{4000}\right) \quad \Rightarrow \quad \sec^2\theta\,\dfrac{d\theta}{dt} = \dfrac{1}{4000}\dfrac{dy}{dt} \quad \Rightarrow \quad \dfrac{d\theta}{dt} = \dfrac{\cos^2\theta}{4000}\dfrac{dy}{dt}$. When

$y = 3000$ ft, $\dfrac{dy}{dt} = 600$ ft/s, $\ell = 5000$ and $\cos\theta = \dfrac{4000}{\ell} = \dfrac{4000}{5000} = \dfrac{4}{5}$, so $\dfrac{d\theta}{dt} = \dfrac{(4/5)^2}{4000}(600) = 0.096$ rad/s.

46. We are given that $\dfrac{d\theta}{dt} = 4(2\pi) = 8\pi$ rad/min. $x = 3\tan\theta \quad \Rightarrow$

$\dfrac{dx}{dt} = 3\sec^2\theta\,\dfrac{d\theta}{dt}$. When $x = 1$, $\tan\theta = \frac{1}{3}$, so $\sec^2\theta = 1 + \left(\frac{1}{3}\right)^2 = \frac{10}{9}$

and $\dfrac{dx}{dt} = 3\left(\frac{10}{9}\right)(8\pi) = \frac{80}{3}\pi \approx 83.8$ km/min.

47. $\cot\theta = \dfrac{x}{5} \quad \Rightarrow \quad -\csc^2\theta\,\dfrac{d\theta}{dt} = \dfrac{1}{5}\dfrac{dx}{dt} \quad \Rightarrow \quad -\left(\csc\dfrac{\pi}{3}\right)^2\left(-\dfrac{\pi}{6}\right) = \dfrac{1}{5}\dfrac{dx}{dt} \quad \Rightarrow$

$\dfrac{dx}{dt} = \dfrac{5\pi}{6}\left(\dfrac{2}{\sqrt{3}}\right)^2 = \dfrac{10}{9}\pi$ km/min $[\approx 130$ mi/h$]$

48. We are given that $\dfrac{d\theta}{dt} = \dfrac{2\pi \text{ rad}}{2 \text{ min}} = \pi$ rad/min. By the Pythagorean Theorem, when

$h = 6$, $x = 8$, so $\sin\theta = \frac{6}{10}$ and $\cos\theta = \frac{8}{10}$. From the figure, $\sin\theta = \dfrac{h}{10} \quad \Rightarrow$

$h = 10\sin\theta$, so $\dfrac{dh}{dt} = 10\cos\theta\,\dfrac{d\theta}{dt} = 10\left(\dfrac{8}{10}\right)\pi = 8\pi$ m/min.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. We are given that $\dfrac{dx}{dt} = 300$ km/h. By the Law of Cosines,

$y^2 = x^2 + 1^2 - 2(1)(x)\cos 120^\circ = x^2 + 1 - 2x\left(-\frac{1}{2}\right) = x^2 + x + 1$, so

$2y\,\dfrac{dy}{dt} = 2x\,\dfrac{dx}{dt} + \dfrac{dx}{dt} \quad\Rightarrow\quad \dfrac{dy}{dt} = \dfrac{2x+1}{2y}\,\dfrac{dx}{dt}$. After 1 minute, $x = \frac{300}{60} = 5$ km $\quad\Rightarrow$

$y = \sqrt{5^2 + 5 + 1} = \sqrt{31}$ km $\quad\Rightarrow\quad \dfrac{dy}{dt} = \dfrac{2(5)+1}{2\sqrt{31}}(300) = \dfrac{1650}{\sqrt{31}} \approx 296$ km/h.

50. We are given that $\dfrac{dx}{dt} = 3$ mi/h and $\dfrac{dy}{dt} = 2$ mi/h. By the Law of Cosines,

$z^2 = x^2 + y^2 - 2xy\cos 45^\circ = x^2 + y^2 - \sqrt{2}\,xy \quad\Rightarrow$

$2z\dfrac{dz}{dt} = 2x\,\dfrac{dx}{dt} + 2y\,\dfrac{dy}{dt} - \sqrt{2}\,x\,\dfrac{dy}{dt} - \sqrt{2}\,y\,\dfrac{dx}{dt}$. After 15 minutes $\left[= \frac{1}{4}\text{ h}\right]$,

we have $x = \frac{3}{4}$ and $y = \frac{2}{4} = \frac{1}{2} \quad\Rightarrow\quad z^2 = \left(\frac{3}{4}\right)^2 + \left(\frac{2}{4}\right)^2 - \sqrt{2}\left(\frac{3}{4}\right)\left(\frac{2}{4}\right) \quad\Rightarrow\quad z = \dfrac{\sqrt{13 - 6\sqrt{2}}}{4}$ and

$$\frac{dz}{dt} = \frac{2}{\sqrt{13-6\sqrt{2}}}\left[2\left(\tfrac{3}{4}\right)3 + 2\left(\tfrac{1}{2}\right)2 - \sqrt{2}\left(\tfrac{3}{4}\right)2 - \sqrt{2}\left(\tfrac{1}{2}\right)3\right] = \frac{2}{\sqrt{13-6\sqrt{2}}}\,\frac{13-6\sqrt{2}}{2} = \sqrt{13-6\sqrt{2}} \approx 2.125 \text{ mi/h}.$$

51. Let the distance between the runner and the friend be ℓ. Then by the Law of Cosines,

$\ell^2 = 200^2 + 100^2 - 2 \cdot 200 \cdot 100 \cdot \cos\theta = 50{,}000 - 40{,}000\cos\theta$ $(\star)$. Differentiating implicitly with respect to t, we obtain $2\ell\,\dfrac{d\ell}{dt} = -40{,}000(-\sin\theta)\,\dfrac{d\theta}{dt}$. Now if D is the distance run when the angle is θ radians, then by the formula for the length of an arc on a circle, $s = r\theta$, we have $D = 100\theta$, so $\theta = \dfrac{1}{100}D \quad\Rightarrow\quad \dfrac{d\theta}{dt} = \dfrac{1}{100}\,\dfrac{dD}{dt} = \dfrac{7}{100}$. To substitute into the expression for $\dfrac{d\ell}{dt}$, we must know $\sin\theta$ at the time when $\ell = 200$, which we find from $(\star)$: $200^2 = 50{,}000 - 40{,}000\cos\theta \quad\Leftrightarrow$

$\cos\theta = \frac{1}{4} \quad\Rightarrow\quad \sin\theta = \sqrt{1 - \left(\frac{1}{4}\right)^2} = \frac{\sqrt{15}}{4}$. Substituting, we get $2(200)\,\dfrac{d\ell}{dt} = 40{,}000\frac{\sqrt{15}}{4}\left(\frac{7}{100}\right) \quad\Rightarrow$

$d\ell/dt = \frac{7\sqrt{15}}{4} \approx 6.78$ m/s. Whether the distance between them is increasing or decreasing depends on the direction in which the runner is running.

52. The hour hand of a clock goes around once every 12 hours or, in radians per hour,

$\frac{2\pi}{12} = \frac{\pi}{6}$ rad/h. The minute hand goes around once an hour, or at the rate of 2π rad/h. So the angle θ between them (measuring clockwise from the minute hand to the hour hand) is changing at the rate of $d\theta/dt = \frac{\pi}{6} - 2\pi = -\frac{11\pi}{6}$ rad/h. Now, to relate θ to ℓ, we use the Law of Cosines: $\ell^2 = 4^2 + 8^2 - 2 \cdot 4 \cdot 8 \cdot \cos\theta = 80 - 64\cos\theta$ $(\star)$.

Differentiating implicitly with respect to t, we get $2\ell\,\dfrac{d\ell}{dt} = -64(-\sin\theta)\dfrac{d\theta}{dt}$. At 1:00, the angle between the two hands is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

one-twelfth of the circle, that is, $\frac{2\pi}{12} = \frac{\pi}{6}$ radians. We use ($\star$) to find ℓ at 1:00: $\ell = \sqrt{80 - 64\cos\frac{\pi}{6}} = \sqrt{80 - 32\sqrt{3}}$.

Substituting, we get $2\ell\,\dfrac{d\ell}{dt} = 64\sin\dfrac{\pi}{6}\left(-\dfrac{11\pi}{6}\right) \quad\Rightarrow\quad \dfrac{d\ell}{dt} = \dfrac{64\left(\frac{1}{2}\right)\left(-\frac{11\pi}{6}\right)}{2\sqrt{80-32\sqrt{3}}} = -\dfrac{88\pi}{3\sqrt{80-32\sqrt{3}}} \approx -18.6.$

So at 1:00, the distance between the tips of the hands is decreasing at a rate of 18.6 mm/h ≈ 0.005 mm/s.

53. The volume of the snowball is given by $V = \dfrac{4}{3}\pi r^3$, so $\dfrac{dV}{dt} = \dfrac{4}{3}\pi\cdot 3r^2\,\dfrac{dr}{dt} = 4\pi r^2\,\dfrac{dr}{dt}$. Since the volume is proportional to the surface area S, with $S = 4\pi r^2$, we also have $\dfrac{dV}{dt} = k\cdot 4\pi r^2$ for some constant k. Equating the two expressions for $\dfrac{dV}{dt}$ gives $4\pi r^2\,\dfrac{dr}{dt} = k\cdot 4\pi r^2 \quad\Rightarrow\quad \dfrac{dr}{dt} = k$, that is, dr/dt is constant.

3.10 Linear Approximations and Differentials

1. $f(x) = x^3 - x^2 + 3 \quad\Rightarrow\quad f'(x) = 3x^2 - 2x$, so $f(-2) = -9$ and $f'(-2) = 16$. Thus,

$L(x) = f(-2) + f'(-2)(x - (-2)) = -9 + 16(x+2) = 16x + 23.$

2. $f(x) = e^{3x} \quad\Rightarrow\quad f'(x) = 3e^{3x}$, so $f(0) = 1$ and $f'(0) = 3$. Thus, $L(x) = f(0) + f'(0)(x-0) = 1 + 3(x-0) = 3x + 1$.

3. $f(x) = \sqrt[3]{x} \quad\Rightarrow\quad f'(x) = \dfrac{1}{3\sqrt[3]{x^2}}$, so $f(8) = 2$ and $f'(8) = \dfrac{1}{12}$. Thus,

$L(x) = f(8) + f'(8)(x-8) = 2 + \frac{1}{12}(x-8) = \frac{1}{12}x + \frac{4}{3}.$

4. $f(x) = \cos 2x \quad\Rightarrow\quad f'(x) = -2\sin 2x$, so $f\left(\frac{\pi}{6}\right) = \frac{1}{2}$ and $f'\left(\frac{\pi}{6}\right) = -\sqrt{3}$. Thus,

$L(x) = f\left(\frac{\pi}{6}\right) + f'\left(\frac{\pi}{6}\right)\left(x - \frac{\pi}{6}\right) = \frac{1}{2} - \sqrt{3}\left(x - \frac{\pi}{6}\right) = -\sqrt{3}\,x + (\sqrt{3}\,\pi)/6 + \frac{1}{2}.$

5. $f(x) = \sqrt{1-x} \quad\Rightarrow\quad f'(x) = \dfrac{-1}{2\sqrt{1-x}}$, so $f(0) = 1$ and $f'(0) = -\frac{1}{2}$.

Therefore,

$\sqrt{1-x} = f(x) \approx f(0) + f'(0)(x-0) = 1 + \left(-\frac{1}{2}\right)(x-0) = 1 - \frac{1}{2}x.$

So $\sqrt{0.9} = \sqrt{1-0.1} \approx 1 - \frac{1}{2}(0.1) = 0.95$

and $\sqrt{0.99} = \sqrt{1-0.01} \approx 1 - \frac{1}{2}(0.01) = 0.995.$

6. $g(x) = \sqrt[3]{1+x} = (1+x)^{1/3} \quad\Rightarrow\quad g'(x) = \frac{1}{3}(1+x)^{-2/3}$, so $g(0) = 1$ and $g'(0) = \frac{1}{3}$. Therefore, $\sqrt[3]{1+x} = g(x) \approx g(0) + g'(0)(x-0) = 1 + \frac{1}{3}x$.

So $\sqrt[3]{0.95} = \sqrt[3]{1+(-0.05)} \approx 1 + \frac{1}{3}(-0.05) = 0.98\overline{3}$,

and $\sqrt[3]{1.1} = \sqrt[3]{1+0.1} \approx 1 + \frac{1}{3}(0.1) = 1.0\overline{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. $f(x) = \tan^{-1} x \quad \Rightarrow \quad f'(x) = \dfrac{1}{1+x^2}$, so $f(0) = 0$ and $f'(0) = 1$.

Thus, $f(x) \approx f(0) + f'(0)(x - 0) = x$. We need $\tan^{-1} x - 0.1 < x < \tan^{-1} x + 0.1$, which is true when $-0.732 < x < 0.732$. Note that to ensure the accuracy, we have rounded the smaller value up and the larger value down.

8. $f(x) = (1+x)^{-3} \quad \Rightarrow \quad f'(x) = -3(1+x)^{-4}$, so $f(0) = 1$ and $f'(0) = -3$. Thus, $f(x) \approx f(0) + f'(0)(x - 0) = 1 - 3x$. We need $(1+x)^{-3} - 0.1 < 1 - 3x < (1+x)^{-3} + 0.1$, which is true when $-0.116 < x < 0.144$. Note that to ensure the accuracy, we have rounded the smaller value up and the larger value down.

9. $f(x) = \sqrt[4]{1+2x} \quad \Rightarrow \quad f'(x) = \frac{1}{4}(1+2x)^{-3/4}(2) = \frac{1}{2}(1+2x)^{-3/4}$, so $f(0) = 1$ and $f'(0) = \frac{1}{2}$. Thus, $f(x) \approx f(0) + f'(0)(x - 0) = 1 + \frac{1}{2}x$. We need $\sqrt[4]{1+2x} - 0.1 < 1 + \frac{1}{2}x < \sqrt[4]{1+2x} + 0.1$, which is true when $-0.368 < x < 0.677$. Note that to ensure the accuracy, we have rounded the smaller value up and the larger value down.

10. $f(x) = \dfrac{2}{1+e^x} \quad \Rightarrow \quad f'(x) = -\dfrac{2e^x}{(1+e^x)^2}$, so $f(0) = 1$ and $f'(0) = -\dfrac{1}{2}$.

Thus, $f(x) \approx f(0) + f'(0)(x - 0) = 1 - \frac{1}{2}x$. We need $\dfrac{2}{1+e^x} - 0.1 < 1 - \dfrac{1}{2}x < \dfrac{2}{1+e^x} + 0.1$, which is true when $-1.423 < x < 1.423$. Note that to ensure the accuracy, we have rounded the smaller value up and the larger value down.

11. The differential dy is defined in terms of dx by the equation $dy = f'(x)\,dx$. For $y = f(x) = e^{5x}$, $f'(x) = 5e^{5x}$, so $dy = 5e^{5x}\,dx$.

12. For $y = f(t) = \sqrt{1-t^4}$, $f'(t) = \frac{1}{2}(1-t^4)^{-1/2}(-4t^3) = -\dfrac{2t^3}{\sqrt{1-t^4}}$, so $dy = -\dfrac{2t^3}{\sqrt{1-t^4}}\,dt$.

13. For $y = f(u) = \dfrac{1+2u}{1+3u}$, $f'(u) = \dfrac{(1+3u)(2) - (1+2u)(3)}{(1+3u)^2} = \dfrac{-1}{(1+3u)^2}$, so $dy = \dfrac{-1}{(1+3u)^2}\,du$.

14. For $y = f(\theta) = \theta^2 \sin 2\theta$, $f'(\theta) = \theta^2(\cos 2\theta)(2) + (\sin 2\theta)(2\theta)$, so $dy = 2\theta(\theta\cos 2\theta + \sin 2\theta)\,d\theta$.

15. For $y = f(x) = \dfrac{1}{x^2 - 3x} = (x^2 - 3x)^{-1}$, $f'(x) = -(x^2 - 3x)^{-2} \cdot (2x - 3) = -\dfrac{2x-3}{(x^2-3x)^2}$, so $dy = -\dfrac{2x-3}{(x^2-3x)^2}\,dx$.

16. For $y = f(\theta) = \sqrt{1+\cos\theta}$, $f'(\theta) = \frac{1}{2}(1+\cos\theta)^{-1/2}(-\sin\theta) = -\dfrac{\sin\theta}{2\sqrt{1+\cos\theta}}$, so $dy = -\dfrac{\sin\theta}{2\sqrt{1+\cos\theta}}\,d\theta$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. For $y = f(\theta) = \ln(\sin\theta)$, $f'(\theta) = \dfrac{1}{\sin\theta}\cos\theta = \cot\theta$, so $dy = \cot\theta\, d\theta$.

18. For $y = f(x) = \dfrac{e^x}{1-e^x}$, $f'(x) = \dfrac{(1-e^x)e^x - e^x(-e^x)}{(1-e^x)^2} = \dfrac{e^x[(1-e^x)-(-e^x)]}{(1-e^x)^2} = \dfrac{e^x}{(1-e^x)^2}$, so $dy = \dfrac{e^x}{(1-e^x)^2}\,dx$.

19. (a) $y = e^{x/10} \quad\Rightarrow\quad dy = e^{x/10}\cdot\frac{1}{10}\,dx = \frac{1}{10}e^{x/10}dx$

(b) $x = 0$ and $dx = 0.1 \quad\Rightarrow\quad dy = \frac{1}{10}e^{0/10}(0.1) = 0.01$.

20. (a) $y = \cos\,\pi x \quad\Rightarrow\quad dy = -\sin\,\pi x\cdot\pi\,dx = -\pi\,\sin\pi x\,dx$

(b) $x = \frac{1}{3}$ and $dx = -0.02 \quad\Rightarrow\quad dy = -\pi\,\sin\frac{\pi}{3}(-0.02) = \pi\left(\sqrt{3}/2\right)(0.02) = 0.01\pi\sqrt{3} \approx 0.054$.

21. (a) $y = \sqrt{3+x^2} \quad\Rightarrow\quad dy = \dfrac{1}{2}(3+x^2)^{-1/2}(2x)\,dx = \dfrac{x}{\sqrt{3+x^2}}\,dx$

(b) $x = 1$ and $dx = -0.1 \quad\Rightarrow\quad dy = \dfrac{1}{\sqrt{3+1^2}}(-0.1) = \dfrac{1}{2}(-0.1) = -0.05$.

22. (a) $y = \dfrac{x+1}{x-1} \quad\Rightarrow\quad dy = \dfrac{(x-1)(1)-(x+1)(1)}{(x-1)^2}\,dx = \dfrac{-2}{(x-1)^2}\,dx$

(b) $x = 2$ and $dx = 0.05 \quad\Rightarrow\quad dy = \dfrac{-2}{(2-1)^2}(0.05) = -2(0.05) = -0.1$.

23. $y = f(x) = x^2 - 4x,\ x = 3,\ \Delta x = 0.5 \quad\Rightarrow$

$\Delta y = f(3.5) - f(3) = -1.75 - (-3) = 1.25$

$dy = f'(x)\,dx = (2x-4)\,dx = (6-4)(0.5) = 1$

24. $y = f(x) = x - x^3,\ x = 0,\ \Delta x = -0.3 \quad\Rightarrow$

$\Delta y = f(-0.3) - f(0) = -0.273 - 0 = -0.273$

$dy = f'(x)\,dx = (1-3x^2)\,dx = (1-0)(-0.3) = -0.3$

25. $y = f(x) = \sqrt{x-2},\ x = 3,\ \Delta x = 0.8 \quad\Rightarrow$

$\Delta y = f(3.8) - f(3) = \sqrt{1.8} - 1 \approx 0.34$

$dy = f'(x)\,dx = \dfrac{1}{2\sqrt{x-2}}\,dx = \dfrac{1}{2(1)}(0.8) = 0.4$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $y = f(x) = e^x,\ x = 0,\ \Delta x = 0.5 \quad \Rightarrow$

$\Delta y = f(0.5) - f(0) = \sqrt{e} - 1\ [\approx 0.65]$

$dy = e^x\,dx = e^0(0.5) = 0.5$

27. $y = f(x) = x^4 - x + 1$. If x changes from 1 to 1.05, $dx = \Delta x = 1.05 - 1 = 0.05 \quad \Rightarrow$

$\Delta y = f(1.05) - f(1) = 1.16550625 - 1 \approx 0.1655$ and $dy = (4x^3 - 1)\,dx = (4 \cdot 1^3 - 1)(0.05) = 3(0.05) = 0.15$.

If x changes from 1 to 1.01, $dx = \Delta x = 1.01 - 1 = 0.01 \quad \Rightarrow \quad \Delta y = f(1.01) - f(1) = 1.03060401 - 1 \approx 0.0306$

and $dy = (4x^3 - 1)\,dx = 3(0.01) = 0.03$.

With $\Delta x = 0.05$, $|dy - \Delta y| \approx |0.15 - 0.1655| = 0.0155$. With $\Delta x = 0.01$, $|dy - \Delta y| \approx |0.03 - 0.0306| = 0.0006$.

Since $|dy - \Delta y|$ is smaller for $\Delta x = 0.01$ than for $\Delta x = 0.05$, yes, the approximation $\Delta y \approx dy$ becomes better as Δx gets smaller.

28. $y = f(x) = e^{2x-2}$. If x changes from 1 to 1.05, $dx = \Delta x = 1.05 - 1 = 0.05 \quad \Rightarrow$

$\Delta y = f(1.05) - f(1) = e^{0.10} - 1 \approx 0.105$ and $dy = 2e^{2x-2}dx = 2e^{2(1)-2}(0.05) = 2(0.05) = 0.1$.

If x changes from 1 to 1.01, $dx = \Delta x = 1.01 - 1 = 0.01 \quad \Rightarrow \quad \Delta y = f(1.01) - f(1) = e^{0.02} - 1 \approx 0.0202$

and $dy = 2e^{2x-2}dx = 2(0.01) = 0.02$.

With $\Delta x = 0.05$, $|dy - \Delta y| \approx |0.1 - 0.105| = 0.005$. With $\Delta x = 0.01$, $|dy - \Delta y| \approx |0.02 - 0.0202| = 0.0002$.

Since $|dy - \Delta y|$ is smaller for $\Delta x = 0.01$ than for $\Delta x = 0.05$, yes, the approximation $\Delta y \approx dy$ becomes better as Δx gets smaller.

29. $y = f(x) = \sqrt{5 - x}$. If x changes from 1 to 1.05, $dx = \Delta x = 1.05 - 1 = 0.05 \quad \Rightarrow$

$\Delta y = f(1.05) - f(1) = \sqrt{3.95} - 2 \approx -0.012539$ and $dy = -\dfrac{1}{2\sqrt{5-x}}dx = -\dfrac{1}{2\sqrt{5-1}}dx = -\dfrac{1}{4}(0.05) = -0.0125$.

If x changes from 1 to 1.01, $dx = \Delta x = 1.01 - 1 = 0.01 \quad \Rightarrow \quad \Delta y = f(1.01) - f(1) = \sqrt{3.99} - 2 \approx -0.002502$

and $dy = -\frac{1}{4}(0.01) = -0.0025$.

With $\Delta x = 0.05$, $|dy - \Delta y| \approx |-0.0125 - (-0.012539)| = 0.000039$. With $\Delta x = 0.01$,

$|dy - \Delta y| \approx |-0.0025 - (-0.002502)| = 0.000002$. Since $|dy - \Delta y|$ is smaller for $\Delta x = 0.01$ than for $\Delta x = 0.05$,

yes, the approximation $\Delta y \approx dy$ becomes better as Δx gets smaller.

30. $y = f(x) = \dfrac{1}{x^2 + 1}$. If x changes from 1 to 1.05, $dx = \Delta x = 1.05 - 1 = 0.05 \quad \Rightarrow$

$\Delta y = f(1.05) - f(1) = \dfrac{1}{(1.05)^2 - 1} - \dfrac{1}{2} \approx -0.02438$ and

$dy = -\dfrac{2x}{(x^2+1)^2}dx = -\dfrac{2(1)}{(1^2+1)^2}\,(0.05) = -\dfrac{1}{2}\,(0.05) = -0.025$.

If x changes from 1 to 1.01, $dx = \Delta x = 1.01 - 1 = 0.01 \quad \Rightarrow \quad \Delta y = f(1.01) - f(1) = \dfrac{1}{1.01^2 + 1} - \dfrac{1}{2} \approx -0.00498$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and $dy = -\frac{1}{2}(0.01) = -0.005$.

With $\Delta x = 0.05$, $|dy - \Delta y| \approx |-0.025 - (-0.02438)| = 0.00038$. With $\Delta x = 0.01$, $|dy - \Delta y| \approx |-0.005 - (-0.00498)| = 0.00002$. Since $|dy - \Delta y|$ is smaller for $\Delta x = 0.01$ than for $\Delta x = 0.05$, yes, the approximation $\Delta y \approx dy$ becomes better as Δx gets smaller.

31. To estimate $(1.999)^4$, we'll find the linearization of $f(x) = x^4$ at $a = 2$. Since $f'(x) = 4x^3$, $f(2) = 16$, and $f'(2) = 32$, we have $L(x) = 16 + 32(x - 2)$. Thus, $x^4 \approx 16 + 32(x - 2)$ when x is near 2, so $(1.999)^4 \approx 16 + 32(1.999 - 2) = 16 - 0.032 = 15.968$.

32. $y = f(x) = 1/x \quad \Rightarrow \quad dy = -1/x^2\,dx$. When $x = 4$ and $dx = 0.002$, $dy = -\frac{1}{16}(0.002) = -\frac{1}{8000}$, so $\frac{1}{4.002} \approx f(4) + dy = \frac{1}{4} - \frac{1}{8000} = \frac{1999}{8000} = 0.249875$.

33. $y = f(x) = \sqrt[3]{x} \quad \Rightarrow \quad dy = \frac{1}{3}x^{-2/3}\,dx$. When $x = 1000$ and $dx = 1$, $dy = \frac{1}{3}(1000)^{-2/3}(1) = \frac{1}{300}$, so $\sqrt[3]{1001} = f(1001) \approx f(1000) + dy = 10 + \frac{1}{300} = 10.00\overline{3} \approx 10.003$.

34. $y = f(x) = \sqrt{x} \quad \Rightarrow \quad dy = \frac{1}{2}x^{-1/2}\,dx$. When $x = 100$ and $dx = 0.5$, $dy = \frac{1}{2}(100)^{-1/2}\left(\frac{1}{2}\right) = \frac{1}{40}$, so $\sqrt{100.5} = f(100.5) \approx f(100) + dy = 10 + \frac{1}{40} = 10.025$.

35. $y = f(x) = e^x \quad \Rightarrow \quad dy = e^x\,dx$. When $x = 0$ and $dx = 0.1$, $dy = e^0(0.1) = 0.1$, so $e^{0.1} = f(0.1) \approx f(0) + dy = 1 + 0.1 = 1.1$.

36. $y = f(x) = \cos x \quad \Rightarrow \quad dy = -\sin x\,dx$. When $x = 30^\circ$ $[\pi/6]$ and $dx = -1^\circ$ $[-\pi/180]$, $dy = \left(-\sin\frac{\pi}{6}\right)\left(-\frac{\pi}{180}\right) = -\frac{1}{2}\left(-\frac{\pi}{180}\right) = \frac{\pi}{360}$, so $\cos 29^\circ = f(29^\circ) \approx f(30^\circ) + dy = \frac{1}{2}\sqrt{3} + \frac{\pi}{360} \approx 0.875$.

37. $y = f(x) = \ln x \quad \Rightarrow \quad f'(x) = 1/x$, so $f(1) = 0$ and $f'(1) = 1$. The linear approximation of f at 1 is $f(1) + f'(1)(x - 1) = x - 1$. Now $f(1.04) = \ln 1.04 \approx 1.04 - 1 = 0.04$, so the approximation is reasonable.

38. $y = f(x) = \sqrt{x} \quad \Rightarrow \quad f'(x) = 1/(2\sqrt{x}\,)$, so $f(4) = 2$ and $f'(4) = \frac{1}{4}$. The linear approximation of f at 4 is $f(4) + f'(4)(x - 4) = 2 + \frac{1}{4}(x - 4)$. Now $f(4.02) = \sqrt{4.02} \approx 2 + \frac{1}{4}(0.02) = 2 + 0.005 = 2.005$, so the approximation is reasonable.

39. $y = f(x) = 1/x \quad \Rightarrow \quad f'(x) = -1/x^2$, so $f(10) = 0.1$ and $f'(10) = -0.01$. The linear approximation of f at 10 is $f(10) + f'(10)(x - 10) = 0.1 - 0.01(x - 10)$. Now $f(9.98) = 1/9.98 \approx 0.1 - 0.01(-0.02) = 0.1 + 0.0002 = 0.1002$, so the approximation is reasonable.

40. (a) $f(x) = (x - 1)^2 \quad \Rightarrow \quad f'(x) = 2(x - 1)$, so $f(0) = 1$ and $f'(0) = -2$.

Thus, $f(x) \approx L_f(x) = f(0) + f'(0)(x - 0) = 1 - 2x$.

$g(x) = e^{-2x} \quad \Rightarrow \quad g'(x) = -2e^{-2x}$, so $g(0) = 1$ and $g'(0) = -2$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Thus, $g(x) \approx L_g(x) = g(0) + g'(0)(x-0) = 1 - 2x$.

$h(x) = 1 + \ln(1-2x) \quad \Rightarrow \quad h'(x) = \dfrac{-2}{1-2x}$, so $h(0) = 1$ and $h'(0) = -2$.

Thus, $h(x) \approx L_h(x) = h(0) + h'(0)(x-0) = 1 - 2x$.

Notice that $L_f = L_g = L_h$. This happens because f, g, and h have the same function values and the same derivative values at $a = 0$.

(b) The linear approximation appears to be the best for the function f since it is closer to f for a larger domain than it is to g and h. The approximation looks worst for h since h moves away from L faster than f and g do.

41. (a) If x is the edge length, then $V = x^3 \quad \Rightarrow \quad dV = 3x^2\,dx$. When $x = 30$ and $dx = 0.1$, $dV = 3(30)^2(0.1) = 270$, so the maximum possible error in computing the volume of the cube is about 270 cm^3. The relative error is calculated by dividing the change in V, ΔV, by V. We approximate ΔV with dV.

Relative error $= \dfrac{\Delta V}{V} \approx \dfrac{dV}{V} = \dfrac{3x^2\,dx}{x^3} = 3\,\dfrac{dx}{x} = 3\left(\dfrac{0.1}{30}\right) = 0.01$.

Percentage error = relative error $\times\, 100\% = 0.01 \times 100\% = 1\%$.

(b) $S = 6x^2 \quad \Rightarrow \quad dS = 12x\,dx$. When $x = 30$ and $dx = 0.1$, $dS = 12(30)(0.1) = 36$, so the maximum possible error in computing the surface area of the cube is about 36 cm^2.

Relative error $= \dfrac{\Delta S}{S} \approx \dfrac{dS}{S} = \dfrac{12x\,dx}{6x^2} = 2\,\dfrac{dx}{x} = 2\left(\dfrac{0.1}{30}\right) = 0.00\overline{6}$.

Percentage error = relative error $\times\, 100\% = 0.00\overline{6} \times 100\% = 0.\overline{6}\%$.

42. (a) $A = \pi r^2 \quad \Rightarrow \quad dA = 2\pi r\,dr$. When $r = 24$ and $dr = 0.2$, $dA = 2\pi(24)(0.2) = 9.6\pi$, so the maximum possible error in the calculated area of the disk is about $9.6\pi \approx 30$ cm^2.

(b) Relative error $= \dfrac{\Delta A}{A} \approx \dfrac{dA}{A} = \dfrac{2\pi r\,dr}{\pi r^2} = \dfrac{2\,dr}{r} = \dfrac{2(0.2)}{24} = \dfrac{0.2}{12} = \dfrac{1}{60} = 0.01\overline{6}$.

Percentage error = relative error $\times 100\% = 0.01\overline{6} \times 100\% = 1.\overline{6}\%$.

43. (a) For a sphere of radius r, the circumference is $C = 2\pi r$ and the surface area is $S = 4\pi r^2$, so

$r = \dfrac{C}{2\pi} \quad \Rightarrow \quad S = 4\pi\left(\dfrac{C}{2\pi}\right)^2 = \dfrac{C^2}{\pi} \quad \Rightarrow \quad dS = \dfrac{2}{\pi}C\,dC$. When $C = 84$ and $dC = 0.5$, $dS = \dfrac{2}{\pi}(84)(0.5) = \dfrac{84}{\pi}$,

so the maximum error is about $\dfrac{84}{\pi} \approx 27$ cm^2. Relative error $\approx \dfrac{dS}{S} = \dfrac{84/\pi}{84^2/\pi} = \dfrac{1}{84} \approx 0.012 = 1.2\%$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $V = \dfrac{4}{3}\pi r^3 = \dfrac{4}{3}\pi\left(\dfrac{C}{2\pi}\right)^3 = \dfrac{C^3}{6\pi^2} \quad \Rightarrow \quad dV = \dfrac{1}{2\pi^2}C^2\,dC$. When $C = 84$ and $dC = 0.5$,

$dV = \dfrac{1}{2\pi^2}(84)^2(0.5) = \dfrac{1764}{\pi^2}$, so the maximum error is about $\dfrac{1764}{\pi^2} \approx 179\text{ cm}^3$.

The relative error is approximately $\dfrac{dV}{V} = \dfrac{1764/\pi^2}{(84)^3/(6\pi^2)} = \dfrac{1}{56} \approx 0.018 = 1.8\%$.

44. For a hemispherical dome, $V = \frac{2}{3}\pi r^3 \quad \Rightarrow \quad dV = 2\pi r^2\,dr$. When $r = \frac{1}{2}(50) = 25$ m and $dr = 0.05$ cm $= 0.0005$ m,

$dV = 2\pi(25)^2(0.0005) = \frac{5\pi}{8}$, so the amount of paint needed is about $\frac{5\pi}{8} \approx 2\text{ m}^3$.

45. (a) $V = \pi r^2 h \quad \Rightarrow \quad \Delta V \approx dV = 2\pi rh\,dr = 2\pi rh\,\Delta r$

(b) The error is

$\Delta V - dV = [\pi(r+\Delta r)^2 h - \pi r^2 h] - 2\pi rh\,\Delta r = \pi r^2 h + 2\pi rh\,\Delta r + \pi(\Delta r)^2 h - \pi r^2 h - 2\pi rh\,\Delta r = \pi(\Delta r)^2 h.$

46. (a) $\sin\theta = \dfrac{20}{x} \quad \Rightarrow \quad x = 20\csc\theta \quad \Rightarrow$

$dx = 20(-\csc\theta\,\cot\theta)\,d\theta = -20\csc 30^\circ\,\cot 30^\circ\,(\pm 1^\circ)$

$= -20(2)(\sqrt{3})\left(\pm\dfrac{\pi}{180}\right) = \pm\dfrac{2\sqrt{3}}{9}\pi$

So the maximum error is about $\pm\frac{2}{9}\sqrt{3}\,\pi \approx \pm 1.21$ cm.

(b) The relative error is $\dfrac{\Delta x}{x} \approx \dfrac{dx}{x} = \dfrac{\pm\frac{2}{9}\sqrt{3}\,\pi}{20(2)} = \pm\dfrac{\sqrt{3}}{180}\pi \approx \pm 0.03$, so the percentage error is approximately $\pm 3\%$.

47. $V = RI \quad \Rightarrow \quad I = \dfrac{V}{R} \quad \Rightarrow \quad dI = -\dfrac{V}{R^2}\,dR$. The relative error in calculating I is $\dfrac{\Delta I}{I} \approx \dfrac{dI}{I} = \dfrac{-(V/R^2)\,dR}{V/R} = -\dfrac{dR}{R}$.

Hence, the relative error in calculating I is approximately the same (in magnitude) as the relative error in R.

48. $F = kR^4 \quad \Rightarrow \quad dF = 4kR^3\,dR \quad \Rightarrow \quad \dfrac{dF}{F} = \dfrac{4kR^3\,dR}{kR^4} = 4\left(\dfrac{dR}{R}\right)$. Thus, the relative change in F is about 4 times the relative change in R. So a 5% increase in the radius corresponds to a 20% increase in blood flow.

49. (a) $dc = \dfrac{dc}{dx}\,dx = 0\,dx = 0$

(b) $d(cu) = \dfrac{d}{dx}(cu)\,dx = c\,\dfrac{du}{dx}\,dx = c\,du$

(c) $d(u+v) = \dfrac{d}{dx}(u+v)\,dx = \left(\dfrac{du}{dx} + \dfrac{dv}{dx}\right)dx = \dfrac{du}{dx}\,dx + \dfrac{dv}{dx}\,dx = du + dv$

(d) $d(uv) = \dfrac{d}{dx}(uv)\,dx = \left(u\,\dfrac{dv}{dx} + v\,\dfrac{du}{dx}\right)dx = u\,\dfrac{dv}{dx}\,dx + v\,\dfrac{du}{dx}\,dx = u\,dv + v\,du$

(e) $d\left(\dfrac{u}{v}\right) = \dfrac{d}{dx}\left(\dfrac{u}{v}\right)dx = \dfrac{v\,\dfrac{du}{dx} - u\,\dfrac{dv}{dx}}{v^2}\,dx = \dfrac{v\,\dfrac{du}{dx}\,dx - u\,\dfrac{dv}{dx}\,dx}{v^2} = \dfrac{v\,du - u\,dv}{v^2}$

(f) $d\left(x^n\right) = \dfrac{d}{dx}\left(x^n\right)dx = nx^{n-1}\,dx$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50. (a) $f(\theta) = \sin\theta \quad \Rightarrow \quad f'(\theta) = \cos\theta$, so $f(0) = 0$ and $f'(0) = 1$.

Thus, $f(\theta) \approx f(0) + f'(0)(\theta - 0) = 0 + 1(\theta - 0) = \theta$.

(b) The relative error in approximating $\sin\theta$ by θ for $\theta = \pi/18$ is $\dfrac{\pi/18 - \sin(\pi/18)}{\sin(\pi/18)} \approx 0.0051$, so the percentage error is 0.51%.

(c)

We want to know the values of θ for which $y = \theta$ approximates $y = \sin\theta$ with less than a 2% difference; that is, the values of θ for which

$$\left|\frac{\theta - \sin\theta}{\sin\theta}\right| < 0.02 \quad \Leftrightarrow \quad -0.02 < \frac{\theta - \sin\theta}{\sin\theta} < 0.02 \quad \Leftrightarrow$$

$$\begin{cases} -0.02\sin\theta < \theta - \sin\theta < 0.02\sin\theta & \text{if } \sin\theta > 0 \\ -0.02\sin\theta > \theta - \sin\theta > 0.02\sin\theta & \text{if } \sin\theta < 0 \end{cases} \quad \Leftrightarrow \quad \begin{cases} 0.98\sin\theta < \theta < 1.02\sin\theta & \text{if } \sin\theta > 0 \\ 1.02\sin\theta < \theta < 0.98\sin\theta & \text{if } \sin\theta < 0 \end{cases}$$

In the first figure, we see that the graphs are very close to each other near $\theta = 0$. Changing the viewing rectangle and using an intersect feature (see the second figure) we find that $y = \theta$ intersects $y = 1.02\sin\theta$ at $\theta \approx 0.344$. By symmetry, they also intersect at $\theta \approx -0.344$ (see the third figure).

Thus, with θ measured in radians, $\sin\theta$ and θ differ by less than 2% for $-0.344 < \theta < 0.344$. Converting 0.344 radians to degrees, we get $0.344(180°/\pi) \approx 19.7°$, so the corresponding interval in degrees is approximately $-19.7° < \theta < 19.7°$.

51. (a) The graph shows that $f'(1) = 2$, so $L(x) = f(1) + f'(1)(x - 1) = 5 + 2(x - 1) = 2x + 3$.

$f(0.9) \approx L(0.9) = 4.8$ and $f(1.1) \approx L(1.1) = 5.2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) From the graph, we see that $f'(x)$ is positive and decreasing. This means that the slopes of the tangent lines are positive, but the tangents are becoming less steep. So the tangent lines lie *above* the curve. Thus, the estimates in part (a) are too large.

52. (a) $g'(x) = \sqrt{x^2+5} \quad \Rightarrow \quad g'(2) = \sqrt{9} = 3$. $g(1.95) \approx g(2) + g'(2)(1.95-2) = -4 + 3(-0.05) = -4.15$.
$g(2.05) \approx g(2) + g'(2)(2.05-2) = -4 + 3(0.05) = -3.85$.

(b) The formula $g'(x) = \sqrt{x^2+5}$ shows that $g'(x)$ is positive and increasing. This means that the slopes of the tangent lines are positive and the tangents are getting steeper. So the tangent lines lie *below* the graph of g. Hence, the estimates in part (a) are too small.

DISCOVERY PROJECT Polynomial Approximations

1. We first write the functions described in conditions (i), (ii), and (iii):

$$\begin{aligned} P(x) &= A + Bx + Cx^2 & f(x) &= \cos x \\ P'(x) &= B + 2Cx & f'(x) &= -\sin x \\ P''(x) &= 2C & f''(x) &= -\cos x \end{aligned}$$

So, taking $a = 0$, our three conditions become

$$\begin{aligned} P(0) = f(0)&: \quad A = \cos 0 = 1 \\ P'(0) = f'(0)&: \quad B = -\sin 0 = 0 \\ P''(0) = f''(0)&: \quad 2C = -\cos 0 = -1 \quad \Rightarrow \quad C = -\tfrac{1}{2} \end{aligned}$$

The desired quadratic function is $P(x) = 1 - \frac{1}{2}x^2$, so the quadratic approximation is $\cos x \approx 1 - \frac{1}{2}x^2$.

The figure shows a graph of the cosine function together with its linear approximation $L(x) = 1$ and quadratic approximation $P(x) = 1 - \frac{1}{2}x^2$ near 0. You can see that the quadratic approximation is much better than the linear one.

2. Accuracy to within 0.1 means that $\left|\cos x - \left(1 - \frac{1}{2}x^2\right)\right| < 0.1 \quad \Leftrightarrow \quad -0.1 < \cos x - \left(1 - \frac{1}{2}x^2\right) < 0.1 \quad \Leftrightarrow$
$0.1 > \left(1 - \frac{1}{2}x^2\right) - \cos x > -0.1 \quad \Leftrightarrow \quad \cos x + 0.1 > 1 - \frac{1}{2}x^2 > \cos x - 0.1 \quad \Leftrightarrow \quad \cos x - 0.1 < 1 - \frac{1}{2}x^2 < \cos x + 0.1$.

From the figure we see that this is true between A and B. Zooming in or using an intersect feature, we find that the x-coordinates of B and A are about ± 1.26. Thus, the approximation $\cos x \approx 1 - \frac{1}{2}x^2$ is accurate to within 0.1 when $-1.26 < x < 1.26$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. If $P(x) = A + B(x-a) + C(x-a)^2$, then $P'(x) = B + 2C(x-a)$ and $P''(x) = 2C$. Applying the conditions (i), (ii), and (iii), we get

$$\begin{aligned} P(a) = f(a)&: \quad A = f(a) \\ P'(a) = f'(a)&: \quad B = f'(a) \\ P''(a) = f''(a)&: \quad 2C = f''(a) \quad \Rightarrow \quad C = \tfrac{1}{2}f''(a) \end{aligned}$$

Thus, $P(x) = A + B(x-a) + C(x-a)^2$ can be written in the form $P(x) = f(a) + f'(a)(x-a) + \frac{1}{2}f''(a)(x-a)^2$.

4. From Example 3.10.1, we have $f(1) = 2$, $f'(1) = \frac{1}{4}$, and $f'(x) = \frac{1}{2}(x+3)^{-1/2}$.

So $f''(x) = -\frac{1}{4}(x+3)^{-3/2} \quad \Rightarrow \quad f''(1) = -\frac{1}{32}$.

From Problem 3, the quadratic approximation $P(x)$ is

$\sqrt{x+3} \approx f(1) + f'(1)(x-1) + \frac{1}{2}f''(1)(x-1)^2 = 2 + \frac{1}{4}(x-1) - \frac{1}{64}(x-1)^2$.

The figure shows the function $f(x) = \sqrt{x+3}$ together with its linear approximation $L(x) = \frac{1}{4}x + \frac{7}{4}$ and its quadratic approximation $P(x)$. You can see that $P(x)$ is a better approximation than $L(x)$ and this is borne out by the numerical values in the following chart.

	from $L(x)$	actual value	from $P(x)$
$\sqrt{3.98}$	1.9950	1.99499373...	1.99499375
$\sqrt{4.05}$	2.0125	2.01246118...	2.01246094
$\sqrt{4.2}$	2.0500	2.04939015...	2.04937500

5. $T_n(x) = c_0 + c_1(x-a) + c_2(x-a)^2 + c_3(x-a)^3 + \cdots + c_n(x-a)^n$. If we put $x = a$ in this equation, then all terms after the first are 0 and we get $T_n(a) = c_0$. Now we differentiate $T_n(x)$ and obtain $T_n'(x) = c_1 + 2c_2(x-a) + 3c_3(x-a)^2 + 4c_4(x-a)^3 + \cdots + nc_n(x-a)^{n-1}$. Substituting $x = a$ gives $T_n'(a) = c_1$. Differentiating again, we have $T_n''(x) = 2c_2 + 2 \cdot 3c_3(x-a) + 3 \cdot 4c_4(x-a^2) + \cdots + (n-1)nc_n(x-a)^{n-2}$ and so $T_n''(a) = 2c_2$. Continuing in this manner, we get $T_n'''(x) = 2 \cdot 3c_3 + 2 \cdot 3 \cdot 4c_4(x-a) + \cdots + (n-2)(n-1)nc_n(x-a)^{n-3}$ and $T_n'''(a) = 2 \cdot 3c_3$. By now we see the pattern. If we continue to differentiate and substitute $x = a$, we obtain $T_n^{(4)}(a) = 2 \cdot 3 \cdot 4c_4$ and in general, for any integer k between 1 and n, $T_n^{(k)}(a) = 2 \cdot 3 \cdot 4 \cdot 5 \cdot \cdots \cdot kc_k = k!\,c_k \quad \Rightarrow$ $c_k = \dfrac{T_n^{(k)}(a)}{k!}$. Because we want T_n and f to have the same derivatives at a, we require that $c_k = \dfrac{f^{(k)}(a)}{k!}$ for $k = 1, 2, \ldots, n$.

6. $T_n(x) = f(a) + f'(a)(x-a) + \dfrac{f''(a)}{2!}(x-a)^2 + \cdots + \dfrac{f^{(n)}(a)}{n!}(x-a)^n$. To compute the coefficients in this equation we need to calculate the derivatives of f at 0:

$$\begin{aligned} f(x) &= \cos x & f(0) &= \cos 0 = 1 \\ f'(x) &= -\sin x & f'(0) &= -\sin 0 = 0 \\ f''(x) &= -\cos x & f''(0) &= -1 \\ f'''(x) &= \sin x & f'''(0) &= 0 \\ f^{(4)}(x) &= \cos x & f^{(4)}(0) &= 1 \end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

We see that the derivatives repeat in a cycle of length 4, so $f^{(5)}(0) = 0$, $f^{(6)}(0) = -1$, $f^{(7)}(0) = 0$, and $f^{(8)}(0) = 1$.

From the original expression for $T_n(x)$, with $n = 8$ and $a = 0$, we have

$$T_8(x) = f(0) + f'(0)(x-0) + \frac{f''(0)}{2!}(x-0)^2 + \frac{f'''(0)}{3!}(x-0)^3 + \cdots + \frac{f^{(8)}(0)}{8!}(x-0)^8$$

$$= 1 + 0 \cdot x + \frac{-1}{2!}x^2 + 0 \cdot x^3 + \frac{1}{4!}x^4 + 0 \cdot x^5 + \frac{-1}{6!}x^6 + 0 \cdot x^7 + \frac{1}{8!}x^8 = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!}$$

and the desired approximation is $\cos x \approx 1 - \dfrac{x^2}{2!} + \dfrac{x^4}{4!} - \dfrac{x^6}{6!} + \dfrac{x^8}{8!}$. The Taylor polynomials T_2, T_4, and T_6 consist of the initial terms of T_8 up through degree 2, 4, and 6, respectively. Therefore, $T_2(x) = 1 - \dfrac{x^2}{2!}$, $T_4(x) = 1 - \dfrac{x^2}{2!} + \dfrac{x^4}{4!}$, and $T_6(x) = 1 - \dfrac{x^2}{2!} + \dfrac{x^4}{4!} - \dfrac{x^6}{6!}$. We graph T_2, T_4, T_6, T_8, and f:

Notice that $T_2(x)$ is a good approximation to $\cos x$ near 0, $T_4(x)$ is a good approximation on a larger interval, $T_6(x)$ is a better approximation, and $T_8(x)$ is better still. Each successive Taylor polynomial is a good approximation on a larger interval than the previous one.

3.11 Hyperbolic Functions

1. (a) $\sinh 0 = \frac{1}{2}(e^0 - e^{-0}) = 0$ (b) $\cosh 0 = \frac{1}{2}(e^0 + e^{-0}) = \frac{1}{2}(1+1) = 1$

2. (a) $\tanh 0 = \dfrac{(e^0 - e^{-0})/2}{(e^0 + e^{-0})/2} = 0$ (b) $\tanh 1 = \dfrac{e^1 - e^{-1}}{e^1 + e^{-1}} = \dfrac{e^2 - 1}{e^2 + 1} \approx 0.76159$

3. (a) $\cosh(\ln 5) = \frac{1}{2}(e^{\ln 5} + e^{-\ln 5}) = \frac{1}{2}\left(5 + (e^{\ln 5})^{-1}\right) = \frac{1}{2}(5 + 5^{-1}) = \frac{1}{2}\left(5 + \frac{1}{5}\right) = \frac{13}{5}$

(b) $\cosh 5 = \frac{1}{2}(e^5 + e^{-5}) \approx 74.20995$

4. (a) $\sinh 4 = \frac{1}{2}(e^4 - e^{-4}) \approx 27.28992$

(b) $\sinh(\ln 4) = \frac{1}{2}(e^{\ln 4} - e^{-\ln 4}) = \frac{1}{2}\left(4 - (e^{\ln 4})^{-1}\right) = \frac{1}{2}(4 - 4^{-1}) = \frac{1}{2}\left(4 - \frac{1}{4}\right) = \frac{15}{8}$

5. (a) $\operatorname{sech} 0 = \dfrac{1}{\cosh 0} = \dfrac{1}{1} = 1$ (b) $\cosh^{-1} 1 = 0$ because $\cosh 0 = 1$.

6. (a) $\sinh 1 = \frac{1}{2}(e^1 - e^{-1}) \approx 1.17520$

(b) Using Equation 3, we have $\sinh^{-1} 1 = \ln\left(1 + \sqrt{1^2 + 1}\right) = \ln\left(1 + \sqrt{2}\right) \approx 0.88137$.

7. $8 \sinh x + 5 \cosh x = 8\left(\dfrac{e^x - e^{-x}}{2}\right) + 5\left(\dfrac{e^x + e^{-x}}{2}\right) = \dfrac{8}{2}e^x - \dfrac{8}{2}e^{-x} + \dfrac{5}{2}e^x + \dfrac{5}{2}e^{-x} = \dfrac{13}{2}e^x - \dfrac{3}{2}e^{-x}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $2e^{2x}+3e^{-2x}=a\sinh 2x+b\cosh 2x \quad\Rightarrow\quad 2e^{2x}+3e^{-2x}=a\left(\dfrac{e^{2x}-e^{-2x}}{2}\right)+b\left(\dfrac{e^{2x}+e^{-2x}}{2}\right) \quad\Rightarrow$

$2e^{2x}+3e^{-2x}=\dfrac{a}{2}e^{2x}-\dfrac{a}{2}e^{-2x}+\dfrac{b}{2}e^{2x}+\dfrac{b}{2}e^{-2x} \quad\Rightarrow\quad 2e^{2x}+3e^{-2x}=\dfrac{a+b}{2}e^{2x}+\dfrac{-a+b}{2}e^{-2x} \quad\Rightarrow$

$\dfrac{a+b}{2}=2$ and $\dfrac{-a+b}{2}=3 \quad\Rightarrow\quad a+b=4$ and $-a+b=6 \quad\Rightarrow\quad 2b=10 \quad\Rightarrow\quad b=5$ and $a=-1$.

Thus, $2e^{2x}+3e^{-2x}=-\sinh 2x+5\cosh 2x$.

9. $\sinh(\ln x)=\dfrac{1}{2}(e^{\ln x}-e^{-\ln x})=\dfrac{1}{2}\left(x-e^{\ln x^{-1}}\right)=\dfrac{1}{2}(x-x^{-1})=\dfrac{1}{2}\left(x-\dfrac{1}{x}\right)=\dfrac{1}{2}\left(\dfrac{x^2-1}{x}\right)=\dfrac{x^2-1}{2x}$

10. $\cosh(4\ln x)=\cosh(\ln x^4)=\dfrac{1}{2}\left(e^{\ln x^4}+e^{-\ln x^4}\right)=\dfrac{1}{2}\left(x^4+e^{\ln x^{-4}}\right)=\dfrac{1}{2}(x^4+x^{-4})$

$$=\frac{1}{2}\left(x^4+\frac{1}{x^4}\right)=\frac{1}{2}\left(\frac{x^8+1}{x^4}\right)=\frac{x^8+1}{2x^4}$$

11. $\sinh(-x)=\frac{1}{2}[e^{-x}-e^{-(-x)}]=\frac{1}{2}(e^{-x}-e^{x})=-\frac{1}{2}(e^{-x}-e^{x})=-\sinh x$

12. $\cosh(-x)=\frac{1}{2}[e^{-x}+e^{-(-x)}]=\frac{1}{2}(e^{-x}+e^{x})=\frac{1}{2}(e^{x}+e^{-x})=\cosh x$

13. $\cosh x+\sinh x=\frac{1}{2}(e^{x}+e^{-x})+\frac{1}{2}(e^{x}-e^{-x})=\frac{1}{2}(2e^{x})=e^{x}$

14. $\cosh x-\sinh x=\frac{1}{2}(e^{x}+e^{-x})-\frac{1}{2}(e^{x}-e^{-x})=\frac{1}{2}(2e^{-x})=e^{-x}$

15. $\sinh x\,\cosh y+\cosh x\,\sinh y=\left[\frac{1}{2}(e^{x}-e^{-x})\right]\left[\frac{1}{2}(e^{y}+e^{-y})\right]+\left[\frac{1}{2}(e^{x}+e^{-x})\right]\left[\frac{1}{2}(e^{y}-e^{-y})\right]$

$$=\tfrac{1}{4}[(e^{x+y}+e^{x-y}-e^{-x+y}-e^{-x-y})+(e^{x+y}-e^{x-y}+e^{-x+y}-e^{-x-y})]$$

$$=\tfrac{1}{4}(2e^{x+y}-2e^{-x-y})=\tfrac{1}{2}[e^{x+y}-e^{-(x+y)}]=\sinh(x+y)$$

16. $\cosh x\,\cosh y+\sinh x\,\sinh y=\left[\frac{1}{2}(e^{x}+e^{-x})\right]\left[\frac{1}{2}(e^{y}+e^{-y})\right]+\left[\frac{1}{2}(e^{x}-e^{-x})\right]\left[\frac{1}{2}(e^{y}-e^{-y})\right]$

$$=\tfrac{1}{4}\left[(e^{x+y}+e^{x-y}+e^{-x+y}+e^{-x-y})+(e^{x+y}-e^{x-y}-e^{-x+y}+e^{-x-y})\right]$$

$$=\tfrac{1}{4}(2e^{x+y}+2e^{-x-y})=\tfrac{1}{2}\left[e^{x+y}+e^{-(x+y)}\right]=\cosh(x+y)$$

17. Divide both sides of the identity $\cosh^2 x-\sinh^2 x=1$ by $\sinh^2 x$:

$\dfrac{\cosh^2 x}{\sinh^2 x}-\dfrac{\sinh^2 x}{\sinh^2 x}=\dfrac{1}{\sinh^2 x} \quad\Leftrightarrow\quad \coth^2 x-1=\operatorname{csch}^2 x.$

18. $\tanh(x+y)=\dfrac{\sinh(x+y)}{\cosh(x+y)}=\dfrac{\sinh x\,\cosh y+\cosh x\,\sinh y}{\cosh x\,\cosh y+\sinh x\,\sinh y}=\dfrac{\dfrac{\sinh x\,\cosh y}{\cosh x\,\cosh y}+\dfrac{\cosh x\,\sinh y}{\cosh x\,\cosh y}}{\dfrac{\cosh x\,\cosh y}{\cosh x\,\cosh y}+\dfrac{\sinh x\,\sinh y}{\cosh x\,\cosh y}}$

$$=\frac{\tanh x+\tanh y}{1+\tanh x\,\tanh y}$$

19. Putting $y=x$ in the result from Exercise 15, we have

$\sinh 2x=\sinh(x+x)=\sinh x\,\cosh x+\cosh x\,\sinh x=2\sinh x\,\cosh x.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. Putting $y = x$ in the result from Exercise 16, we have

$\cosh 2x = \cosh(x + x) = \cosh x \cosh x + \sinh x \sinh x = \cosh^2 x + \sinh^2 x.$

21. $\tanh(\ln x) = \dfrac{\sinh(\ln x)}{\cosh(\ln x)} = \dfrac{(e^{\ln x} - e^{-\ln x})/2}{(e^{\ln x} + e^{-\ln x})/2} = \dfrac{x - (e^{\ln x})^{-1}}{x + (e^{\ln x})^{-1}} = \dfrac{x - x^{-1}}{x + x^{-1}} = \dfrac{x - 1/x}{x + 1/x} = \dfrac{(x^2 - 1)/x}{(x^2 + 1)/x} = \dfrac{x^2 - 1}{x^2 + 1}$

22. $\dfrac{1 + \tanh x}{1 - \tanh x} = \dfrac{1 + (\sinh x)/\cosh x}{1 - (\sinh x)/\cosh x} = \dfrac{\cosh x + \sinh x}{\cosh x - \sinh x} = \dfrac{\frac{1}{2}(e^x + e^{-x}) + \frac{1}{2}(e^x - e^{-x})}{\frac{1}{2}(e^x + e^{-x}) - \frac{1}{2}(e^x - e^{-x})} = \dfrac{e^x}{e^{-x}} = e^{2x}$

Or: Using the results of Exercises 13 and 14, $\dfrac{\cosh x + \sinh x}{\cosh x - \sinh x} = \dfrac{e^x}{e^{-x}} = e^{2x}$

23. By Exercise 13, $(\cosh x + \sinh x)^n = (e^x)^n = e^{nx} = \cosh nx + \sinh nx.$

24. $\coth x = \dfrac{1}{\tanh x} \quad \Rightarrow \quad \coth x = \dfrac{1}{\tanh x} = \dfrac{1}{12/13} = \dfrac{13}{12}.$

$\operatorname{sech}^2 x = 1 - \tanh^2 x = 1 - \left(\frac{12}{13}\right)^2 = \frac{25}{169} \quad \Rightarrow \quad \operatorname{sech} x = \frac{5}{13}$ [sech, like cosh, is positive].

$\cosh x = \dfrac{1}{\operatorname{sech} x} \quad \Rightarrow \quad \cosh x = \dfrac{1}{5/13} = \dfrac{13}{5}.$

$\tanh x = \dfrac{\sinh x}{\cosh x} \quad \Rightarrow \quad \sinh x = \tanh x \cosh x \quad \Rightarrow \quad \sinh x = \dfrac{12}{13} \cdot \dfrac{13}{5} = \dfrac{12}{5}.$

$\operatorname{csch} x = \dfrac{1}{\sinh x} \quad \Rightarrow \quad \operatorname{csch} x = \dfrac{1}{12/5} = \dfrac{5}{12}.$

25. $\operatorname{sech} x = \dfrac{1}{\cosh x} \quad \Rightarrow \quad \operatorname{sech} x = \dfrac{1}{5/3} = \dfrac{3}{5}.$

$\cosh^2 x - \sinh^2 x = 1 \quad \Rightarrow \quad \sinh^2 x = \cosh^2 x - 1 = \left(\frac{5}{3}\right)^2 - 1 = \frac{16}{9} \quad \Rightarrow \quad \sinh x = \frac{4}{3}$ [because $x > 0$].

$\operatorname{csch} x = \dfrac{1}{\sinh x} \quad \Rightarrow \quad \operatorname{csch} x = \dfrac{1}{4/3} = \dfrac{3}{4}.$

$\tanh x = \dfrac{\sinh x}{\cosh x} \quad \Rightarrow \quad \tanh x = \dfrac{4/3}{5/3} = \dfrac{4}{5}.$

$\coth x = \dfrac{1}{\tanh x} \quad \Rightarrow \quad \coth x = \dfrac{1}{4/5} = \dfrac{5}{4}.$

26. (a)

27. (a) $\displaystyle\lim_{x\to\infty} \tanh x = \lim_{x\to\infty} \frac{e^x - e^{-x}}{e^x + e^{-x}} \cdot \frac{e^{-x}}{e^{-x}} = \lim_{x\to\infty} \frac{1 - e^{-2x}}{1 + e^{-2x}} = \frac{1 - 0}{1 + 0} = 1$

(b) $\displaystyle\lim_{x\to-\infty} \tanh x = \lim_{x\to-\infty} \frac{e^x - e^{-x}}{e^x + e^{-x}} \cdot \frac{e^x}{e^x} = \lim_{x\to-\infty} \frac{e^{2x} - 1}{e^{2x} + 1} = \frac{0 - 1}{0 + 1} = -1$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\lim\limits_{x\to\infty} \sinh x = \lim\limits_{x\to\infty} \dfrac{e^x - e^{-x}}{2} = \infty$

(d) $\lim\limits_{x\to-\infty} \sinh x = \lim\limits_{x\to-\infty} \dfrac{e^x - e^{-x}}{2} = -\infty$

(e) $\lim\limits_{x\to\infty} \operatorname{sech} x = \lim\limits_{x\to\infty} \dfrac{2}{e^x + e^{-x}} = 0$

(f) $\lim\limits_{x\to\infty} \coth x = \lim\limits_{x\to\infty} \dfrac{e^x + e^{-x}}{e^x - e^{-x}} \cdot \dfrac{e^{-x}}{e^{-x}} = \lim\limits_{x\to\infty} \dfrac{1 + e^{-2x}}{1 - e^{-2x}} = \dfrac{1+0}{1-0} = 1$ [*Or:* Use part (a).]

(g) $\lim\limits_{x\to 0^+} \coth x = \lim\limits_{x\to 0^+} \dfrac{\cosh x}{\sinh x} = \infty$, since $\sinh x \to 0$ through positive values and $\cosh x \to 1$.

(h) $\lim\limits_{x\to 0^-} \coth x = \lim\limits_{x\to 0^-} \dfrac{\cosh x}{\sinh x} = -\infty$, since $\sinh x \to 0$ through negative values and $\cosh x \to 1$.

(i) $\lim\limits_{x\to-\infty} \operatorname{csch} x = \lim\limits_{x\to-\infty} \dfrac{2}{e^x - e^{-x}} = 0$

(j) $\lim\limits_{x\to\infty} \dfrac{\sinh x}{e^x} = \lim\limits_{x\to\infty} \dfrac{e^x - e^{-x}}{2e^x} = \lim\limits_{x\to\infty} \dfrac{1 - e^{-2x}}{2} = \dfrac{1-0}{2} = \dfrac{1}{2}$

28. (a) $\dfrac{d}{dx}(\cosh x) = \dfrac{d}{dx}\left[\frac{1}{2}(e^x + e^{-x})\right] = \frac{1}{2}(e^x - e^{-x}) = \sinh x$

(b) $\dfrac{d}{dx}(\tanh x) = \dfrac{d}{dx}\left(\dfrac{\sinh x}{\cosh x}\right) = \dfrac{\cosh x \cosh x - \sinh x \sinh x}{\cosh^2 x} = \dfrac{\cosh^2 x - \sinh^2 x}{\cosh^2 x} = \dfrac{1}{\cosh^2 x} = \operatorname{sech}^2 x$

(c) $\dfrac{d}{dx}(\operatorname{csch} x) = \dfrac{d}{dx}\left(\dfrac{1}{\sinh x}\right) = -\dfrac{\cosh x}{\sinh^2 x} = -\dfrac{1}{\sinh x} \cdot \dfrac{\cosh x}{\sinh x} = -\operatorname{csch} x \coth x$

(d) $\dfrac{d}{dx}(\operatorname{sech} x) = \dfrac{d}{dx}\left(\dfrac{1}{\cosh x}\right) = -\dfrac{\sinh x}{\cosh^2 x} = -\dfrac{1}{\cosh x} \cdot \dfrac{\sinh x}{\cosh x} = -\operatorname{sech} x \tanh x$

(e) $\dfrac{d}{dx}(\coth x) = \dfrac{d}{dx}\left(\dfrac{\cosh x}{\sinh x}\right) = \dfrac{\sinh x \sinh x - \cosh x \cosh x}{\sinh^2 x} = \dfrac{\sinh^2 x - \cosh^2 x}{\sinh^2 x} = -\dfrac{1}{\sinh^2 x} = -\operatorname{csch}^2 x$

29. Let $y = \sinh^{-1} x$. Then $\sinh y = x$ and, by Example 1(a), $\cosh^2 y - \sinh^2 y = 1 \;\Rightarrow\;$ [with $\cosh y > 0$] $\cosh y = \sqrt{1 + \sinh^2 y} = \sqrt{1 + x^2}$. So by Exercise 13, $e^y = \sinh y + \cosh y = x + \sqrt{1 + x^2} \;\Rightarrow$ $y = \ln\left(x + \sqrt{1 + x^2}\right)$.

30. Let $y = \cosh^{-1} x$. Then $\cosh y = x$ and $y \geq 0$, so $\sinh y = \sqrt{\cosh^2 y - 1} = \sqrt{x^2 - 1}$. So, by Exercise 13, $e^y = \cosh y + \sinh y = x + \sqrt{x^2 - 1} \;\Rightarrow\; y = \ln\left(x + \sqrt{x^2 - 1}\right)$.

Another method: Write $x = \cosh y = \frac{1}{2}\left(e^y + e^{-y}\right)$ and solve a quadratic, as in Example 3.

31. (a) Let $y = \tanh^{-1} x$. Then $x = \tanh y = \dfrac{\sinh y}{\cosh y} = \dfrac{(e^y - e^{-y})/2}{(e^y + e^{-y})/2} \cdot \dfrac{e^y}{e^y} = \dfrac{e^{2y} - 1}{e^{2y} + 1} \;\Rightarrow\; xe^{2y} + x = e^{2y} - 1 \;\Rightarrow$

$1 + x = e^{2y} - xe^{2y} \;\Rightarrow\; 1 + x = e^{2y}(1 - x) \;\Rightarrow\; e^{2y} = \dfrac{1+x}{1-x} \;\Rightarrow\; 2y = \ln\left(\dfrac{1+x}{1-x}\right) \;\Rightarrow$

$y = \dfrac{1}{2}\ln\left(\dfrac{1+x}{1-x}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Let $y = \tanh^{-1} x$. Then $x = \tanh y$, so from Exercise 22 we have

$$e^{2y} = \frac{1+\tanh y}{1-\tanh y} = \frac{1+x}{1-x} \quad \Rightarrow \quad 2y = \ln\left(\frac{1+x}{1-x}\right) \quad \Rightarrow \quad y = \frac{1}{2}\ln\left(\frac{1+x}{1-x}\right).$$

32. (a) (i) $y = \operatorname{csch}^{-1} x \quad \Leftrightarrow \quad \operatorname{csch} y = x \quad (x \neq 0)$

(ii) We sketch the graph of csch^{-1} by reflecting the graph of csch (see Exercise 26) about the line $y = x$.

(iii) Let $y = \operatorname{csch}^{-1} x$. Then $x = \operatorname{csch} y = \dfrac{2}{e^y - e^{-y}} \quad \Rightarrow \quad xe^y - xe^{-y} = 2 \quad \Rightarrow$

$x(e^y)^2 - 2e^y - x = 0 \quad \Rightarrow \quad e^y = \dfrac{1 \pm \sqrt{x^2+1}}{x}$. But $e^y > 0$, so for $x > 0$,

$e^y = \dfrac{1+\sqrt{x^2+1}}{x}$ and for $x < 0$, $e^y = \dfrac{1-\sqrt{x^2+1}}{x}$. Thus, $\operatorname{csch}^{-1} x = \ln\left(\dfrac{1}{x} + \dfrac{\sqrt{x^2+1}}{|x|}\right)$.

(b) (i) $y = \operatorname{sech}^{-1} x \quad \Leftrightarrow \quad \operatorname{sech} y = x$ and $y > 0$.

(ii) We sketch the graph of sech^{-1} by reflecting the graph of sech (see Exercise 26) about the line $y = x$.

(iii) Let $y = \operatorname{sech}^{-1} x$, so $x = \operatorname{sech} y = \dfrac{2}{e^y + e^{-y}} \quad \Rightarrow \quad xe^y + xe^{-y} = 2 \quad \Rightarrow$

$x\,(e^y)^2 - 2e^y + x = 0 \quad \Leftrightarrow \quad e^y = \dfrac{1 \pm \sqrt{1-x^2}}{x}$. But $y > 0 \quad \Rightarrow \quad e^y > 1$.

This rules out the minus sign because $\dfrac{1-\sqrt{1-x^2}}{x} > 1 \quad \Leftrightarrow \quad 1 - \sqrt{1-x^2} > x \quad \Leftrightarrow \quad 1 - x > \sqrt{1-x^2} \quad \Leftrightarrow$

$1 - 2x + x^2 > 1 - x^2 \quad \Leftrightarrow \quad x^2 > x \quad \Leftrightarrow \quad x > 1$, but $x = \operatorname{sech} y \leq 1$.

Thus, $e^y = \dfrac{1+\sqrt{1-x^2}}{x} \quad \Rightarrow \quad \operatorname{sech}^{-1} x = \ln\left(\dfrac{1+\sqrt{1-x^2}}{x}\right)$.

(c) (i) $y = \coth^{-1} x \quad \Leftrightarrow \quad \coth y = x$

(ii) We sketch the graph of $\coth^{-1}$ by reflecting the graph of coth (see Exercise 26) about the line $y = x$.

(iii) Let $y = \coth^{-1} x$. Then $x = \coth y = \dfrac{e^y + e^{-y}}{e^y - e^{-y}} \quad \Rightarrow$

$xe^y - xe^{-y} = e^y + e^{-y} \quad \Rightarrow \quad (x-1)e^y = (x+1)e^{-y} \quad \Rightarrow \quad e^{2y} = \dfrac{x+1}{x-1} \quad \Rightarrow$

$2y = \ln\dfrac{x+1}{x-1} \quad \Rightarrow \quad \coth^{-1} x = \dfrac{1}{2}\ln\dfrac{x+1}{x-1}$

33. (a) Let $y = \cosh^{-1} x$. Then $\cosh y = x$ and $y \geq 0 \quad \Rightarrow \quad \sinh y\,\dfrac{dy}{dx} = 1 \quad \Rightarrow$

$\dfrac{dy}{dx} = \dfrac{1}{\sinh y} = \dfrac{1}{\sqrt{\cosh^2 y - 1}} = \dfrac{1}{\sqrt{x^2-1}}$ [since $\sinh y \geq 0$ for $y \geq 0$]. *Or:* Use Formula 4.

(b) Let $y = \tanh^{-1} x$. Then $\tanh y = x \quad \Rightarrow \quad \operatorname{sech}^2 y\,\dfrac{dy}{dx} = 1 \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{1}{\operatorname{sech}^2 y} = \dfrac{1}{1-\tanh^2 y} = \dfrac{1}{1-x^2}$.

Or: Use Formula 5.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) Let $y = \coth^{-1} x$. Then $\coth y = x \quad \Rightarrow \quad -\operatorname{csch}^2 y \dfrac{dy}{dx} = 1 \quad \Rightarrow \quad \dfrac{dy}{dx} = -\dfrac{1}{\operatorname{csch}^2 y} = \dfrac{1}{1-\coth^2 y} = \dfrac{1}{1-x^2}$

by Exercise 17.

34. (a) Let $y = \operatorname{sech}^{-1} x$. Then $\operatorname{sech} y = x \quad \Rightarrow \quad -\operatorname{sech} y \tanh y \dfrac{dy}{dx} = 1 \quad \Rightarrow$

$\dfrac{dy}{dx} = -\dfrac{1}{\operatorname{sech} y \tanh y} = -\dfrac{1}{\operatorname{sech} y \sqrt{1-\operatorname{sech}^2 y}} = -\dfrac{1}{x\sqrt{1-x^2}}$. [Note that $y > 0$ and so $\tanh y > 0$.]

(b) Let $y = \operatorname{csch}^{-1} x$. Then $\operatorname{csch} y = x \quad \Rightarrow \quad -\operatorname{csch} y \coth y \dfrac{dy}{dx} = 1 \quad \Rightarrow \quad \dfrac{dy}{dx} = -\dfrac{1}{\operatorname{csch} y \coth y}$. By Exercise 17,

$\coth y = \pm\sqrt{\operatorname{csch}^2 y + 1} = \pm\sqrt{x^2+1}$. If $x > 0$, then $\coth y > 0$, so $\coth y = \sqrt{x^2+1}$. If $x < 0$, then $\coth y < 0$,

so $\coth y = -\sqrt{x^2+1}$. In either case we have $\dfrac{dy}{dx} = -\dfrac{1}{\operatorname{csch} y \coth y} = -\dfrac{1}{|x|\sqrt{x^2+1}}$.

35. $f(x) = \cosh 3x \quad \Rightarrow \quad f'(x) = \sinh(3x) \cdot \dfrac{d}{dx}(3x) = \sinh(3x) \cdot 3 = 3\sinh 3x$

36. $f(x) = e^x \cosh x \overset{\text{PR}}{\Rightarrow} f'(x) = e^x \sinh x + (\cosh x)e^x = e^x(\sinh x + \cosh x)$, or, using Exercise 13, $e^x(e^x) = e^{2x}$.

37. $h(x) = \sinh(x^2) \quad \Rightarrow \quad h'(x) = \cosh(x^2)\dfrac{d}{dx}(x^2) = 2x\cosh(x^2)$

38. $g(x) = \sinh^2 x = (\sinh x)^2 \quad \Rightarrow \quad g'(x) = 2(\sinh x)^1 \dfrac{d}{dx}(\sinh x) = 2\sinh x \cosh x$, or, using Exercise 19, $\sinh 2x$.

39. $G(t) = \sinh(\ln t) \quad \Rightarrow \quad G'(t) = \cosh(\ln t)\dfrac{d}{dt}\ln t = \dfrac{1}{2}\left(e^{\ln t} + e^{-\ln t}\right)\left(\dfrac{1}{t}\right) = \dfrac{1}{2t}\left(t + \dfrac{1}{t}\right) = \dfrac{1}{2t}\left(\dfrac{t^2+1}{t}\right) = \dfrac{t^2+1}{2t^2}$

Or: $G(t) = \sinh(\ln t) = \dfrac{1}{2}(e^{\ln t} - e^{-\ln t}) = \dfrac{1}{2}\left(t - \dfrac{1}{t}\right) \quad \Rightarrow \quad G'(t) = \dfrac{1}{2}\left(1 + \dfrac{1}{t^2}\right) = \dfrac{t^2+1}{2t^2}$

40. $F(t) = \ln(\sinh t) \quad \Rightarrow \quad F'(t) = \dfrac{1}{\sinh t}\dfrac{d}{dt}\sinh t = \dfrac{1}{\sinh t}\cosh t = \coth t$

41. $f(x) = \tanh\sqrt{x} \quad \Rightarrow \quad f'(x) = \operatorname{sech}^2\sqrt{x}\dfrac{d}{dx}\sqrt{x} = \operatorname{sech}^2\sqrt{x}\left(\dfrac{1}{2\sqrt{x}}\right) = \dfrac{\operatorname{sech}^2\sqrt{x}}{2\sqrt{x}}$

42. $H(v) = e^{\tanh 2v} \quad \Rightarrow \quad H'(v) = e^{\tanh 2v} \cdot \dfrac{d}{dv}\tanh 2v = e^{\tanh 2v} \cdot \operatorname{sech}^2(2v) \cdot 2 = 2e^{\tanh 2v}\operatorname{sech}^2(2v)$

43. $y = \operatorname{sech} x \tanh x \overset{\text{PR}}{\Rightarrow} y' = \operatorname{sech} x \cdot \operatorname{sech}^2 x + \tanh x \cdot (-\operatorname{sech} x \tanh x) = \operatorname{sech}^3 x - \operatorname{sech} x \tanh^2 x$

44. $y = \operatorname{sech}(\tanh x) \quad \Rightarrow \quad y' = -\operatorname{sech}(\tanh x)\tanh(\tanh x) \cdot \dfrac{d}{dx}(\tanh x) = -\operatorname{sech}(\tanh x)\tanh(\tanh x) \cdot \operatorname{sech}^2 x$

45. $g(t) = t\coth\sqrt{t^2+1} \overset{\text{PR}}{\Rightarrow}$

$g'(t) = t\left[-\operatorname{csch}^2\sqrt{t^2+1}\left(\tfrac{1}{2}(t^2+1)^{-1/2} \cdot 2t\right)\right] + \left(\coth\sqrt{t^2+1}\right)(1) = \coth\sqrt{t^2+1} - \dfrac{t^2}{\sqrt{t^2+1}}\operatorname{csch}^2\sqrt{t^2+1}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. $f(t) = \dfrac{1+\sinh t}{1-\sinh t} \overset{\text{QR}}{\Rightarrow}$

$$f'(t) = \frac{(1-\sinh t)\,\cosh t - (1+\sinh t)(-\cosh t)}{(1-\sinh t)^2} = \frac{\cosh t - \sinh t\,\cosh t + \cosh t + \sinh t\,\cosh t}{(1-\sinh t)^2}$$

$$= \frac{2\cosh t}{(1-\sinh t)^2}$$

47. $f(x) = \sinh^{-1}(-2x) \quad \Rightarrow \quad f'(x) = \dfrac{1}{\sqrt{1+(-2x)^2}} \cdot \dfrac{d}{dx}(-2x) = -\dfrac{2}{\sqrt{1+4x^2}}$

48. $g(x) = \tanh^{-1}(x^3) \quad \Rightarrow \quad g'(x) = \dfrac{1}{1-(x^3)^2} \cdot \dfrac{d}{dx}(x^3) = \dfrac{3x^2}{1-x^6}$

49. $y = \cosh^{-1}(\sec\theta) \quad \Rightarrow$

$$y' = \frac{1}{\sqrt{\sec^2\theta - 1}} \cdot \frac{d}{d\theta}(\sec\theta) = \frac{1}{\sqrt{\tan^2\theta}} \cdot \sec\theta\,\tan\theta = \frac{1}{\tan\theta} \cdot \sec\theta\,\tan\theta \quad [\text{since } 0 \le \theta < \pi/2] \quad = \sec\theta$$

50. $y = \operatorname{sech}^{-1}(\sin\theta) \quad \Rightarrow$

$$y' = -\frac{1}{\sin\theta\,\sqrt{1-\sin^2\theta}} \cdot \frac{d}{d\theta}(\sin\theta) = -\frac{1}{\sin\theta\,\sqrt{\cos^2\theta}} \cdot \cos\theta$$

$$= -\frac{1}{\sin\theta \cdot \cos\theta} \cdot \cos\theta \quad [\text{since } 0 < \theta < \pi/2] \quad = -\frac{1}{\sin\theta} = -\csc\theta$$

51. $G(u) = \cosh^{-1}\sqrt{1+u^2} \quad \Rightarrow$

$$G'(u) = \frac{1}{\sqrt{\left(\sqrt{1+u^2}\right)^2 - 1}} \cdot \frac{d}{du}\left(\sqrt{1+u^2}\right) = \frac{1}{\sqrt{(1+u^2)-1}} \cdot \frac{2u}{2\sqrt{1+u^2}} = \frac{u}{\sqrt{u^2}\cdot\sqrt{1+u^2}}$$

$$= \frac{u}{u\sqrt{1+u^2}} \quad [\text{since } u > 0] \quad = \frac{1}{\sqrt{1+u^2}}$$

52. $y = x\tanh^{-1}x + \ln\sqrt{1-x^2} = x\tanh^{-1}x + \frac{1}{2}\ln(1-x^2) \quad \Rightarrow$

$$y' = \tanh^{-1}x + \frac{x}{1-x^2} + \frac{1}{2}\left(\frac{1}{1-x^2}\right)(-2x) = \tanh^{-1}x$$

53. $y = x\sinh^{-1}(x/3) - \sqrt{9+x^2} \quad \Rightarrow$

$$y' = \sinh^{-1}\left(\frac{x}{3}\right) + x\frac{1/3}{\sqrt{1+(x/3)^2}} - \frac{2x}{2\sqrt{9+x^2}} = \sinh^{-1}\left(\frac{x}{3}\right) + \frac{x}{\sqrt{9+x^2}} - \frac{x}{\sqrt{9+x^2}} = \sinh^{-1}\left(\frac{x}{3}\right)$$

54. $\dfrac{1+\tanh x}{1-\tanh x} = \dfrac{1+(\sinh x)/\cosh x}{1-(\sinh x)/\cosh x} = \dfrac{\cosh x + \sinh x}{\cosh x - \sinh x} = \dfrac{e^x}{e^{-x}}$ [by Exercises 13 and 14] $= e^{2x}$, so

$\sqrt[4]{\dfrac{1+\tanh x}{1-\tanh x}} = \sqrt[4]{e^{2x}} = e^{x/2}$. Thus, $\dfrac{d}{dx}\sqrt[4]{\dfrac{1+\tanh x}{1-\tanh x}} = \dfrac{d}{dx}(e^{x/2}) = \dfrac{1}{2}e^{x/2}$.

55. $\dfrac{d}{dx}\arctan(\tanh x) = \dfrac{1}{1+(\tanh x)^2}\dfrac{d}{dx}(\tanh x) = \dfrac{\operatorname{sech}^2 x}{1+\tanh^2 x} = \dfrac{1/\cosh^2 x}{1+(\sinh^2 x)/\cosh^2 x}$

$$= \frac{1}{\cosh^2 x + \sinh^2 x} = \frac{1}{\cosh 2x} \quad [\text{by Exercise 20}] \quad = \operatorname{sech} 2x$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

56. (a) Let $a = 0.03291765$. A graph of the central curve,

$y = f(x) = 211.49 - 20.96\cosh ax$, is shown.

(b) $f(0) = 211.49 - 20.96\cosh 0 = 211.49 - 20.96(1) = 190.53$ m.

(c) $y = 100 \quad\Rightarrow\quad 100 = 211.49 - 20.96\cosh ax \quad\Rightarrow$

$20.96\cosh ax = 111.49 \quad\Rightarrow\quad \cosh ax = \dfrac{111.49}{20.96} \quad\Rightarrow$

$ax = \pm\cosh^{-1}\dfrac{111.49}{20.96} \quad\Rightarrow\quad x = \pm\dfrac{1}{a}\cosh^{-1}\dfrac{111.49}{20.96} \approx \pm 71.56$ m. The points are approximately $(\pm 71.56, 100)$.

(d) $f(x) = 211.49 - 20.96\cosh ax \quad\Rightarrow\quad f'(x) = -20.96\sinh ax \cdot a$.

$$f'\left(\pm\frac{1}{a}\cosh^{-1}\frac{111.49}{20.96}\right) = -20.96a\sinh\left[a\left(\pm\frac{1}{a}\cosh^{-1}\frac{111.49}{20.96}\right)\right] = -20.96a\sinh\left(\pm\cosh^{-1}\frac{111.49}{20.96}\right) \approx \mp 3.6.$$

So the slope at $(71.56, 100)$ is about -3.6 and the slope at $(-71.56, 100)$ is about 3.6.

57. As the depth d of the water gets large, the fraction $\dfrac{2\pi d}{L}$ gets large, and from Figure 5 or Exercise 27(a), $\tanh\left(\dfrac{2\pi d}{L}\right)$ approaches 1. Thus, $v = \sqrt{\dfrac{gL}{2\pi}\tanh\left(\dfrac{2\pi d}{L}\right)} \approx \sqrt{\dfrac{gL}{2\pi}(1)} = \sqrt{\dfrac{gL}{2\pi}}$.

58.

For $y = a\cosh(x/a)$ with $a > 0$, we have the y-intercept equal to a. As a increases, the graph flattens.

59. (a) $y = 20\cosh(x/20) - 15 \quad\Rightarrow\quad y' = 20\sinh(x/20)\cdot\frac{1}{20} = \sinh(x/20)$. Since the right pole is positioned at $x = 7$, we have $y'(7) = \sinh\frac{7}{20} \approx 0.3572$.

(b) If α is the angle between the tangent line and the x-axis, then $\tan\alpha =$ slope of the line $= \sinh\frac{7}{20}$, so $\alpha = \tan^{-1}\left(\sinh\frac{7}{20}\right) \approx 0.343$ rad $\approx 19.66°$. Thus, the angle between the line and the pole is $\theta = 90° - \alpha \approx 70.34°$.

60. We differentiate the function twice, then substitute into the differential equation: $y = \dfrac{T}{\rho g}\cosh\dfrac{\rho g x}{T} \quad\Rightarrow$

$\dfrac{dy}{dx} = \dfrac{T}{\rho g}\sinh\left(\dfrac{\rho g x}{T}\right)\dfrac{\rho g}{T} = \sinh\dfrac{\rho g x}{T} \quad\Rightarrow\quad \dfrac{d^2y}{dx^2} = \cosh\left(\dfrac{\rho g x}{T}\right)\dfrac{\rho g}{T} = \dfrac{\rho g}{T}\cosh\dfrac{\rho g x}{T}$. We evaluate the two sides separately: LHS $= \dfrac{d^2y}{dx^2} = \dfrac{\rho g}{T}\cosh\dfrac{\rho g x}{T}$ and RHS $= \dfrac{\rho g}{T}\sqrt{1 + \left(\dfrac{dy}{dx}\right)^2} = \dfrac{\rho g}{T}\sqrt{1 + \sinh^2\dfrac{\rho g x}{T}} = \dfrac{\rho g}{T}\cosh\dfrac{\rho g x}{T}$, by the identity proved in Example 1(a).

61. (a) From Exercise 60, the shape of the cable is given by $y = f(x) = \dfrac{T}{\rho g}\cosh\left(\dfrac{\rho g x}{T}\right)$. The shape is symmetric about the y-axis, so the lowest point is $(0, f(0)) = \left(0, \dfrac{T}{\rho g}\right)$ and the poles are at $x = \pm 100$. We want to find T when the lowest

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

point is 60 m, so $\dfrac{T}{\rho g} = 60 \quad \Rightarrow \quad T = 60\rho g = (60\text{ m})(2\text{ kg/m})(9.8\text{ m/s}^2) = 1176\ \dfrac{\text{kg-m}}{\text{s}^2}$, or 1176 N (newtons).

The height of each pole is $f(100) = \dfrac{T}{\rho g}\cosh\left(\dfrac{\rho g \cdot 100}{T}\right) = 60\cosh\left(\dfrac{100}{60}\right) \approx 164.50$ m.

(b) If the tension is doubled from T to $2T$, then the low point is doubled since $\dfrac{T}{\rho g} = 60 \quad \Rightarrow \quad \dfrac{2T}{\rho g} = 120$. The height of the poles is now $f(100) = \dfrac{2T}{\rho g}\cosh\left(\dfrac{\rho g \cdot 100}{2T}\right) = 120\cosh\left(\dfrac{100}{120}\right) \approx 164.13$ m, just a slight decrease.

62. (a) $\lim\limits_{t\to\infty} v(t) = \lim\limits_{t\to\infty}\sqrt{\dfrac{mg}{k}}\tanh\left(t\sqrt{\dfrac{gk}{m}}\right) = \sqrt{\dfrac{mg}{k}}\lim\limits_{t\to\infty}\tanh\left(t\sqrt{\dfrac{gk}{m}}\right) = \sqrt{\dfrac{mg}{k}}\cdot 1 \quad \left[\begin{array}{c}\text{as } t\to\infty,\\ t\sqrt{gk/m}\to\infty\end{array}\right] \quad = \sqrt{\dfrac{mg}{k}}$

(b) Belly-to-earth: $g = 9.8$, $k = 0.515$, $m = 60$, so the terminal velocity is $\sqrt{\dfrac{60(9.8)}{0.515}} \approx 33.79$ m/s.

Feet-first: $g = 9.8$, $k = 0.067$, $m = 60$, so the terminal velocity is $\sqrt{\dfrac{60(9.8)}{0.067}} \approx 93.68$ m/s.

63. (a) $y = A\sinh mx + B\cosh mx \quad \Rightarrow \quad y' = mA\cosh mx + mB\sinh mx \quad \Rightarrow$

$y'' = m^2 A\sinh mx + m^2 B\cosh mx = m^2(A\sinh mx + B\cosh mx) = m^2 y$

(b) From part (a), a solution of $y'' = 9y$ is $y(x) = A\sinh 3x + B\cosh 3x$. Now $-4 = y(0) = A\sinh 0 + B\cosh 0 = B$, so $B = -4$. Also, $y'(x) = 3A\cosh 3x - 12\sinh 3x$, so $6 = y'(0) = 3A \quad \Rightarrow \quad A = 2$. Thus, $y = 2\sinh 3x - 4\cosh 3x$.

64. $\cosh x = \cosh[\ln(\sec\theta + \tan\theta)] = \dfrac{1}{2}\left[e^{\ln(\sec\theta+\tan\theta)} + e^{-\ln(\sec\theta+\tan\theta)}\right] = \dfrac{1}{2}\left[\sec\theta + \tan\theta + \dfrac{1}{\sec\theta+\tan\theta}\right]$

$= \dfrac{1}{2}\left[\sec\theta + \tan\theta + \dfrac{\sec\theta - \tan\theta}{(\sec\theta+\tan\theta)(\sec\theta-\tan\theta)}\right] = \dfrac{1}{2}\left[\sec\theta + \tan\theta + \dfrac{\sec\theta - \tan\theta}{\sec^2\theta - \tan^2\theta}\right]$

$= \frac{1}{2}(\sec\theta + \tan\theta + \sec\theta - \tan\theta) = \sec\theta$

65. The tangent to $y = \cosh x$ has slope 1 when $y' = \sinh x = 1 \quad \Rightarrow \quad x = \sinh^{-1} 1 = \ln\left(1+\sqrt{2}\right)$, by Equation 3.

Since $\sinh x = 1$ and $y = \cosh x = \sqrt{1+\sinh^2 x}$, we have $\cosh x = \sqrt{2}$. The point is $\left(\ln\left(1+\sqrt{2}\right), \sqrt{2}\right)$.

66. $f_n(x) = \tanh(n\sin x)$, where n is a positive integer. Note that $f_n(x+2\pi) = f_n(x)$; that is, f_n is periodic with period 2π. Also, from Figure 3, $-1 < \tanh x < 1$, so we can choose a viewing rectangle of $[0, 2\pi] \times [-1, 1]$. From the graph, we see that $f_n(x)$ becomes more rectangular looking as n increases. As n becomes large, the graph of f_n approaches the graph of $y = 1$ on the intervals $(2k\pi, (2k+1)\pi)$ and $y = -1$ on the intervals $((2k-1)\pi, 2k\pi)$.

67. If $ae^x + be^{-x} = \alpha\cosh(x+\beta)$ [or $\alpha\sinh(x+\beta)$], then

$ae^x + be^{-x} = \frac{\alpha}{2}\left(e^{x+\beta} \pm e^{-x-\beta}\right) = \frac{\alpha}{2}\left(e^x e^\beta \pm e^{-x}e^{-\beta}\right) = \left(\frac{\alpha}{2}e^\beta\right)e^x \pm \left(\frac{\alpha}{2}e^{-\beta}\right)e^{-x}$. Comparing coefficients of e^x and e^{-x}, we have $a = \frac{\alpha}{2}e^\beta$ **(1)** and $b = \pm\frac{\alpha}{2}e^{-\beta}$ **(2)**. We need to find α and β. Dividing equation **(1)** by equation **(2)**

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

gives us $\frac{a}{b} = \pm e^{2\beta} \quad \Rightarrow \quad (\star) \quad 2\beta = \ln\left(\pm\frac{a}{b}\right) \quad \Rightarrow \quad \beta = \frac{1}{2}\ln\left(\pm\frac{a}{b}\right)$. Solving equations **(1)** and **(2)** for e^{β} gives us $e^{\beta} = \dfrac{2a}{\alpha}$ and $e^{\beta} = \pm\dfrac{\alpha}{2b}$, so $\dfrac{2a}{\alpha} = \pm\dfrac{\alpha}{2b} \quad \Rightarrow \quad \alpha^2 = \pm 4ab \quad \Rightarrow \quad \alpha = 2\sqrt{\pm ab}$.

$(\star)$ If $\frac{a}{b} > 0$, we use the $+$ sign and obtain a $\cosh$ function, whereas if $\frac{a}{b} < 0$, we use the $-$ sign and obtain a $\sinh$ function.

In summary, if a and b have the same sign, we have $ae^x + be^{-x} = 2\sqrt{ab}\cosh\left(x + \frac{1}{2}\ln\frac{a}{b}\right)$, whereas, if a and b have the opposite sign, then $ae^x + be^{-x} = 2\sqrt{-ab}\sinh\left(x + \frac{1}{2}\ln\left(-\frac{a}{b}\right)\right)$.

3 Review

TRUE-FALSE QUIZ

1. True. This is the Sum Rule.

2. False. See the warning before the Product Rule.

3. True. This is the Chain Rule.

4. True. $\dfrac{d}{dx}\sqrt{f(x)} = \dfrac{d}{dx}[f(x)]^{1/2} = \dfrac{1}{2}[f(x)]^{-1/2}\,f'(x) = \dfrac{f'(x)}{2\sqrt{f(x)}}$

5. False. $\dfrac{d}{dx}\,f(\sqrt{x}\,) = f'(\sqrt{x}\,)\cdot\frac{1}{2}x^{-1/2} = \dfrac{f'(\sqrt{x}\,)}{2\sqrt{x}}$, which is not $\dfrac{f'(x)}{2\sqrt{x}}$.

6. False. $y = e^2$ is a constant, so $y' = 0$, not $2e$.

7. False. $\dfrac{d}{dx}\,(10^x) = 10^x \ln 10$, which is not equal to $x10^{x-1}$.

8. False. $\ln 10$ is a constant, so its derivative, $\dfrac{d}{dx}\,(\ln 10)$, is 0, not $\frac{1}{10}$.

9. True. $\dfrac{d}{dx}\,(\tan^2 x) = 2\,\tan x\sec^2 x$, and $\dfrac{d}{dx}\,(\sec^2 x) = 2\,\sec x\,(\sec x\tan x) = 2\,\tan x\sec^2 x$.

Or: $\dfrac{d}{dx}\,(\sec^2 x) = \dfrac{d}{dx}\,(1 + \tan^2 x) = \dfrac{d}{dx}\,(\tan^2 x)$.

10. False. $f(x) = \left|x^2 + x\right| = x^2 + x$ for $x \geq 0$ or $x \leq -1$ and $\left|x^2 + x\right| = -(x^2 + x)$ for $-1 < x < 0$. So $f'(x) = 2x + 1$ for $x > 0$ or $x < -1$ and $f'(x) = -(2x + 1)$ for $-1 < x < 0$. But $|2x + 1| = 2x + 1$ for $x \geq -\frac{1}{2}$ and $|2x + 1| = -2x - 1$ for $x < -\frac{1}{2}$.

11. True. If $p(x) = a_n x^n + a_{n-1}x^{n-1} + \cdots + a_1 x + a_0$, then $p'(x) = na_n x^{n-1} + (n-1)a_{n-1}x^{n-2} + \cdots + a_1$, which is a polynomial.

12. True. $f(x) = (x^6 - x^4)^5$ is a polynomial of degree 30, so its 31st derivative, $f^{(31)}(x)$, is 0.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. True. If $r(x) = \dfrac{p(x)}{q(x)}$, then $r'(x) = \dfrac{q(x)p'(x) - p(x)q'(x)}{[q(x)]^2}$, which is a quotient of polynomials, that is, a rational function.

14. False. A tangent line to the parabola $y = x^2$ has slope $dy/dx = 2x$, so at $(-2, 4)$ the slope of the tangent is $2(-2) = -4$ and an equation of the tangent line is $y - 4 = -4(x + 2)$. [The given equation, $y - 4 = 2x(x + 2)$, is not even linear!]

15. True. $g(x) = x^5 \;\Rightarrow\; g'(x) = 5x^4 \;\Rightarrow\; g'(2) = 5(2)^4 = 80$, and by the definition of the derivative,

$$\lim_{x\to 2} \frac{g(x) - g(2)}{x - 2} = g'(2) = 5(2)^4 = 80.$$

EXERCISES

1. $y = (x^2 + x^3)^4 \;\Rightarrow\; y' = 4(x^2 + x^3)^3(2x + 3x^2) = 4(x^2)^3(1 + x)^3 x(2 + 3x) = 4x^7(x + 1)^3(3x + 2)$

2. $y = \dfrac{1}{\sqrt{x}} - \dfrac{1}{\sqrt[5]{x^3}} = x^{-1/2} - x^{-3/5} \;\Rightarrow\; y' = -\dfrac{1}{2}x^{-3/2} + \dfrac{3}{5}x^{-8/5}$ or $\dfrac{3}{5x\sqrt[5]{x^3}} - \dfrac{1}{2x\sqrt{x}}$ or $\dfrac{1}{10}x^{-8/5}(-5x^{1/10} + 6)$

3. $y = \dfrac{x^2 - x + 2}{\sqrt{x}} = x^{3/2} - x^{1/2} + 2x^{-1/2} \;\Rightarrow\; y' = \dfrac{3}{2}x^{1/2} - \dfrac{1}{2}x^{-1/2} - x^{-3/2} = \dfrac{3}{2}\sqrt{x} - \dfrac{1}{2\sqrt{x}} - \dfrac{1}{\sqrt{x^3}}$

4. $y = \dfrac{\tan x}{1 + \cos x} \;\Rightarrow\; y' = \dfrac{(1 + \cos x)\sec^2 x - \tan x(-\sin x)}{(1 + \cos x)^2} = \dfrac{(1 + \cos x)\sec^2 x + \tan x \sin x}{(1 + \cos x)^2}$

5. $y = x^2 \sin \pi x \;\Rightarrow\; y' = x^2(\cos \pi x)\pi + (\sin \pi x)(2x) = x(\pi x \cos \pi x + 2 \sin \pi x)$

6. $y = x \cos^{-1} x \;\Rightarrow\; y' = x\left(-\dfrac{1}{\sqrt{1 - x^2}}\right) + (\cos^{-1} x)(1) = \cos^{-1} x - \dfrac{x}{\sqrt{1 - x^2}}$

7. $y = \dfrac{t^4 - 1}{t^4 + 1} \;\Rightarrow\; y' = \dfrac{(t^4 + 1)4t^3 - (t^4 - 1)4t^3}{(t^4 + 1)^2} = \dfrac{4t^3[(t^4 + 1) - (t^4 - 1)]}{(t^4 + 1)^2} = \dfrac{8t^3}{(t^4 + 1)^2}$

8. $\dfrac{d}{dx}(xe^y) = \dfrac{d}{dx}(y \sin x) \;\Rightarrow\; xe^y y' + e^y \cdot 1 = y \cos x + \sin x \cdot y' \;\Rightarrow\; xe^y y' - \sin x \cdot y' = y \cos x - e^y \;\Rightarrow$

$(xe^y - \sin x)y' = y \cos x - e^y \;\Rightarrow\; y' = \dfrac{y \cos x - e^y}{xe^y - \sin x}$

9. $y = \ln(x \ln x) \;\Rightarrow\; y' = \dfrac{1}{x \ln x}(x \ln x)' = \dfrac{1}{x \ln x}\left(x \cdot \dfrac{1}{x} + \ln x \cdot 1\right) = \dfrac{1 + \ln x}{x \ln x}$

Another method: $y = \ln(x \ln x) = \ln x + \ln \ln x \;\Rightarrow\; y' = \dfrac{1}{x} + \dfrac{1}{\ln x} \cdot \dfrac{1}{x} = \dfrac{\ln x + 1}{x \ln x}$

10. $y = e^{mx} \cos nx \;\Rightarrow$

$y' = e^{mx}(\cos nx)' + \cos nx\,(e^{mx})' = e^{mx}(-\sin nx \cdot n) + \cos nx\,(e^{mx} \cdot m) = e^{mx}(m \cos nx - n \sin nx)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. $y = \sqrt{x}\cos\sqrt{x} \quad \Rightarrow$

$$y' = \sqrt{x}\left(\cos\sqrt{x}\right)' + \cos\sqrt{x}\left(\sqrt{x}\right)' = \sqrt{x}\left[-\sin\sqrt{x}\left(\tfrac{1}{2}x^{-1/2}\right)\right] + \cos\sqrt{x}\left(\tfrac{1}{2}x^{-1/2}\right)$$

$$= \tfrac{1}{2}x^{-1/2}\left(-\sqrt{x}\,\sin\sqrt{x} + \cos\sqrt{x}\right) = \frac{\cos\sqrt{x} - \sqrt{x}\,\sin\sqrt{x}}{2\sqrt{x}}$$

12. $y = (\arcsin 2x)^2 \quad \Rightarrow \quad y' = 2(\arcsin 2x)\cdot(\arcsin 2x)' = 2\arcsin 2x \cdot \dfrac{1}{\sqrt{1-(2x)^2}}\cdot 2 = \dfrac{4\arcsin 2x}{\sqrt{1-4x^2}}$

13. $y = \dfrac{e^{1/x}}{x^2} \quad \Rightarrow \quad y' = \dfrac{x^2(e^{1/x})' - e^{1/x}\left(x^2\right)'}{(x^2)^2} = \dfrac{x^2(e^{1/x})(-1/x^2) - e^{1/x}(2x)}{x^4} = \dfrac{-e^{1/x}(1+2x)}{x^4}$

14. $y = \ln\sec x \quad \Rightarrow \quad y' = \dfrac{1}{\sec x}\dfrac{d}{dx}(\sec x) = \dfrac{1}{\sec x}(\sec x\,\tan x) = \tan x$

15. $\dfrac{d}{dx}(y + x\cos y) = \dfrac{d}{dx}(x^2 y) \quad \Rightarrow \quad y' + x(-\sin y\cdot y') + \cos y\cdot 1 = x^2 y' + y\cdot 2x \quad \Rightarrow$

$$y' - x\sin y\cdot y' - x^2 y' = 2xy - \cos y \quad \Rightarrow \quad (1 - x\sin y - x^2)y' = 2xy - \cos y \quad \Rightarrow \quad y' = \frac{2xy - \cos y}{1 - x\sin y - x^2}$$

16. $y = \left(\dfrac{u-1}{u^2+u+1}\right)^4 \quad \Rightarrow$

$$y' = 4\left(\frac{u-1}{u^2+u+1}\right)^3 \frac{d}{du}\left(\frac{u-1}{u^2+u+1}\right) = 4\left(\frac{u-1}{u^2+u+1}\right)^3 \frac{(u^2+u+1)(1) - (u-1)(2u+1)}{(u^2+u+1)^2}$$

$$= \frac{4(u-1)^3}{(u^2+u+1)^3}\,\frac{u^2+u+1-2u^2+u+1}{(u^2+u+1)^2} = \frac{4(u-1)^3(-u^2+2u+2)}{(u^2+u+1)^5}$$

17. $y = \sqrt{\arctan x} \quad \Rightarrow \quad y' = \dfrac{1}{2}(\arctan x)^{-1/2}\dfrac{d}{dx}(\arctan x) = \dfrac{1}{2\sqrt{\arctan x}\,(1+x^2)}$

18. $y = \cot(\csc x) \quad \Rightarrow \quad y' = -\csc^2(\csc x)\dfrac{d}{dx}(\csc x) = -\csc^2(\csc x)\cdot(-\csc x\,\cot x) = \csc^2(\csc x)\,\csc x\,\cot x$

19. $y = \tan\left(\dfrac{t}{1+t^2}\right) \quad \Rightarrow$

$$y' = \sec^2\left(\frac{t}{1+t^2}\right)\frac{d}{dt}\left(\frac{t}{1+t^2}\right) = \sec^2\left(\frac{t}{1+t^2}\right)\cdot\frac{(1+t^2)(1) - t(2t)}{(1+t^2)^2} = \frac{1-t^2}{(1+t^2)^2}\sec^2\left(\frac{t}{1+t^2}\right)$$

20. $y = e^{x\sec x} \quad \Rightarrow \quad y' = e^{x\sec x}\dfrac{d}{dx}(x\sec x) = e^{x\sec x}(x\sec x\,\tan x + \sec x\cdot 1) = \sec x\, e^{x\sec x}(x\tan x + 1)$

21. $y = 3^{x\ln x} \quad \Rightarrow \quad y' = 3^{x\ln x}(\ln 3)\dfrac{d}{dx}(x\ln x) = 3^{x\ln x}(\ln 3)\left(x\cdot\dfrac{1}{x} + \ln x\cdot 1\right) = 3^{x\ln x}(\ln 3)(1+\ln x)$

22. $y = \sec(1+x^2) \quad \Rightarrow \quad y' = 2x\sec(1+x^2)\,\tan(1+x^2)$

23. $y = (1-x^{-1})^{-1} \quad \Rightarrow$

$$y' = -1(1-x^{-1})^{-2}[-(-1x^{-2})] = -(1-1/x)^{-2}x^{-2} = -((x-1)/x)^{-2}x^{-2} = -(x-1)^{-2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $y = \dfrac{1}{\sqrt[3]{x+\sqrt{x}}} = \left(x+\sqrt{x}\right)^{-1/3} \quad \Rightarrow \quad y' = -\frac{1}{3}\left(x+\sqrt{x}\right)^{-4/3}\left(1+\dfrac{1}{2\sqrt{x}}\right)$

25. $\sin(xy) = x^2 - y \quad \Rightarrow \quad \cos(xy)(xy' + y\cdot 1) = 2x - y' \quad \Rightarrow \quad x\cos(xy)y' + y' = 2x - y\cos(xy) \quad \Rightarrow$

$y'[x\cos(xy)+1] = 2x - y\cos(xy) \quad \Rightarrow \quad y' = \dfrac{2x - y\cos(xy)}{x\cos(xy)+1}$

26. $y = \sqrt{\sin\sqrt{x}} \quad \Rightarrow \quad y' = \frac{1}{2}\left(\sin\sqrt{x}\right)^{-1/2}\left(\cos\sqrt{x}\right)\left(\dfrac{1}{2\sqrt{x}}\right) = \dfrac{\cos\sqrt{x}}{4\sqrt{x\sin\sqrt{x}}}$

27. $y = \log_5(1+2x) \quad \Rightarrow \quad y' = \dfrac{1}{(1+2x)\,\ln 5}\,\dfrac{d}{dx}\,(1+2x) = \dfrac{2}{(1+2x)\,\ln 5}$

28. $y = (\cos x)^x \quad \Rightarrow \quad \ln y = \ln(\cos x)^x = x\ln\cos x \quad \Rightarrow \quad \dfrac{y'}{y} = x\cdot\dfrac{1}{\cos x}\cdot(-\sin x) + \ln\cos x\cdot 1 \quad \Rightarrow$

$y' = (\cos x)^x(\ln\cos x - x\tan x)$

29. $y = \ln\sin x - \frac{1}{2}\sin^2 x \quad \Rightarrow \quad y' = \dfrac{1}{\sin x}\cdot\cos x - \frac{1}{2}\cdot 2\sin x\cdot\cos x = \cot x - \sin x\,\cos x$

30. $y = \dfrac{(x^2+1)^4}{(2x+1)^3(3x-1)^5} \quad \Rightarrow$

$\ln y = \ln\dfrac{(x^2+1)^4}{(2x+1)^3(3x-1)^5} = \ln(x^2+1)^4 - \ln[(2x+1)^3(3x-1)^5] = 4\ln(x^2+1) - [\ln(2x+1)^3 + \ln(3x-1)^5]$

$= 4\ln(x^2+1) - 3\ln(2x+1) - 5\ln(3x-1) \quad \Rightarrow$

$\dfrac{y'}{y} = 4\cdot\dfrac{1}{x^2+1}\cdot 2x - 3\cdot\dfrac{1}{2x+1}\cdot 2 - 5\cdot\dfrac{1}{3x-1}\cdot 3 \quad \Rightarrow \quad y' = \dfrac{(x^2+1)^4}{(2x+1)^3(3x-1)^5}\left(\dfrac{8x}{x^2+1} - \dfrac{6}{2x+1} - \dfrac{15}{3x-1}\right).$

[The answer could be simplified to $y' = -\dfrac{(x^2+56x+9)(x^2+1)^3}{(2x+1)^4(3x-1)^6}$, but this is unnecessary.]

31. $y = x\tan^{-1}(4x) \quad \Rightarrow \quad y' = x\cdot\dfrac{1}{1+(4x)^2}\cdot 4 + \tan^{-1}(4x)\cdot 1 = \dfrac{4x}{1+16x^2} + \tan^{-1}(4x)$

32. $y = e^{\cos x} + \cos(e^x) \quad \Rightarrow \quad y' = e^{\cos x}(-\sin x) + [-\sin(e^x)\cdot e^x] = -\sin x\, e^{\cos x} - e^x\sin(e^x)$

33. $y = \ln|\sec 5x + \tan 5x| \quad \Rightarrow$

$y' = \dfrac{1}{\sec 5x + \tan 5x}(\sec 5x\,\tan 5x\cdot 5 + \sec^2 5x\cdot 5) = \dfrac{5\sec 5x\,(\tan 5x + \sec 5x)}{\sec 5x + \tan 5x} = 5\sec 5x$

34. $y = 10^{\tan\pi\theta} \quad \Rightarrow \quad y' = 10^{\tan\pi\theta}\cdot\ln 10\cdot\sec^2\pi\theta\cdot\pi = \pi(\ln 10)10^{\tan\pi\theta}\sec^2\pi\theta$

35. $y = \cot(3x^2+5) \quad \Rightarrow \quad y' = -\csc^2(3x^2+5)(6x) = -6x\csc^2(3x^2+5)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. $y = \sqrt{t\ln(t^4)} \quad \Rightarrow$

$$y' = \frac{1}{2}[t\ln(t^4)]^{-1/2}\,\frac{d}{dt}\,[t\ln(t^4)] = \frac{1}{2\sqrt{t\ln(t^4)}}\cdot\left[1\cdot\ln(t^4) + t\cdot\frac{1}{t^4}\cdot 4t^3\right] = \frac{1}{2\sqrt{t\ln(t^4)}}\cdot[\ln(t^4)+4] = \frac{\ln(t^4)+4}{2\sqrt{t\ln(t^4)}}$$

Or: Since y is only defined for $t > 0$, we can write $y = \sqrt{t\cdot 4\ln t} = 2\sqrt{t\ln t}$. Then

$y' = 2\cdot\dfrac{1}{2\sqrt{t\ln t}}\cdot\left(1\cdot\ln t + t\cdot\dfrac{1}{t}\right) = \dfrac{\ln t+1}{\sqrt{t\ln t}}$. This agrees with our first answer since

$$\frac{\ln(t^4)+4}{2\sqrt{t\ln(t^4)}} = \frac{4\ln t+4}{2\sqrt{t\cdot 4\ln t}} = \frac{4(\ln t+1)}{2\cdot 2\sqrt{t\ln t}} = \frac{\ln t+1}{\sqrt{t\ln t}}.$$

37. $y = \sin\left(\tan\sqrt{1+x^3}\right) \quad\Rightarrow\quad y' = \cos\left(\tan\sqrt{1+x^3}\right)\left(\sec^2\sqrt{1+x^3}\right)\left[3x^2/\left(2\sqrt{1+x^3}\right)\right]$

38. $y = x\sec^{-1}x \quad\Rightarrow\quad y' = x\cdot\dfrac{1}{x\sqrt{x^2-1}} + \left(\sec^{-1}x\right)\cdot 1 = \dfrac{1}{\sqrt{x^2-1}} + \sec^{-1}x$

39. $y = 5\arctan\dfrac{1}{x} \quad\Rightarrow\quad y' = 5\cdot\dfrac{1}{1+\left(\dfrac{1}{x}\right)^2}\cdot\dfrac{d}{dx}\left(\dfrac{1}{x}\right) = \dfrac{5}{1+\dfrac{1}{x^2}}\left(-\dfrac{1}{x^2}\right) = -\dfrac{5}{x^2+1}$

40. $y = \sin^{-1}(\cos\theta) \quad\Rightarrow$

$$y' = \frac{1}{\sqrt{1-\cos^2\theta}}\cdot\frac{d}{d\theta}(\cos\theta) = \frac{1}{\sqrt{1-\cos^2\theta}}\cdot(-\sin\theta) = -\frac{\sin\theta}{\sqrt{1-\cos^2\theta}}$$

$$= -\frac{\sin\theta}{\sin\theta} \qquad [\sin^2\theta+\cos^2\theta = 1 \quad\Rightarrow\quad \sin\theta = \sqrt{1-\cos^2\theta} \text{ for } 0<\theta<\pi]$$

$$= -1$$

41. $y = x\tan^{-1}x - \frac{1}{2}\ln(1+x^2) \quad\Rightarrow$

$$y' = x\cdot\frac{1}{1+x^2} + \left(\tan^{-1}x\right)\cdot 1 - \frac{1}{2}\left(\frac{2x}{1+x^2}\right) = \frac{x}{1+x^2} + \tan^{-1}x - \frac{x}{1+x^2} = \tan^{-1}x$$

42. $y = \ln(\arcsin x^2) \quad\Rightarrow\quad y' = \dfrac{1}{\arcsin x^2}\cdot\dfrac{d}{dx}\left(\arcsin x^2\right) = \dfrac{1}{\arcsin x^2}\cdot\dfrac{1}{\sqrt{1-(x^2)^2}}\cdot 2x = \dfrac{2x}{(\arcsin x^2)\sqrt{1-x^4}}$

43. $y = \tan^2(\sin\theta) = [\tan(\sin\theta)]^2 \quad\Rightarrow\quad y' = 2[\tan(\sin\theta)]\cdot\sec^2(\sin\theta)\cdot\cos\theta$

44. $y + \ln y = xy^2 \quad\Rightarrow\quad y' + (1/y)\,y' = x\cdot 2y\,y' + y^2\cdot 1 \quad\Rightarrow\quad y' + (1/y)\,y' - 2xy\,y' = y^2 \quad\Rightarrow$

$[1+(1/y)-2xy]\,y' = y^2 \quad\Rightarrow\quad y' = \dfrac{y^2}{1+(1/y)-2xy}\cdot\dfrac{y}{y} = \dfrac{y^3}{y+1-2xy^2}$

45. $y = \dfrac{\sqrt{x+1}\,(2-x)^5}{(x+3)^7} \quad\Rightarrow\quad \ln y = \frac{1}{2}\ln(x+1) + 5\ln(2-x) - 7\ln(x+3) \quad\Rightarrow\quad \dfrac{y'}{y} = \dfrac{1}{2(x+1)} + \dfrac{-5}{2-x} - \dfrac{7}{x+3} \quad\Rightarrow$

$y' = \dfrac{\sqrt{x+1}\,(2-x)^5}{(x+3)^7}\left[\dfrac{1}{2(x+1)} - \dfrac{5}{2-x} - \dfrac{7}{x+3}\right]$ or $y' = \dfrac{(2-x)^4(3x^2-55x-52)}{2\sqrt{x+1}\,(x+3)^8}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. $y = \dfrac{(x+\lambda)^4}{x^4+\lambda^4} \quad \Rightarrow \quad y' = \dfrac{(x^4+\lambda^4)(4)(x+\lambda)^3 - (x+\lambda)^4(4x^3)}{(x^4+\lambda^4)^2} = \dfrac{4(x+\lambda)^3(\lambda^4 - \lambda x^3)}{(x^4+\lambda^4)^2}$

47. $y = x\sinh(x^2) \quad \Rightarrow \quad y' = x\cosh(x^2)\cdot 2x + \sinh(x^2)\cdot 1 = 2x^2\cosh(x^2) + \sinh(x^2)$

48. $y = \dfrac{\sin mx}{x} \quad \Rightarrow \quad y' = \dfrac{x\cdot\cos mx\cdot m - \sin mx\cdot 1}{(x)^2} = \dfrac{mx\cos mx - \sin mx}{x^2}$

49. $y = \ln(\cosh 3x) \quad \Rightarrow \quad y' = (1/\cosh 3x)(\sinh 3x)(3) = 3\tanh 3x$

50. $y = \ln\left|\dfrac{x^2-4}{2x+5}\right| = \ln|x^2-4| - \ln|2x+5| \quad \Rightarrow \quad y' = \dfrac{2x}{x^2-4} - \dfrac{2}{2x+5}$ or $\dfrac{2(x+1)(x+4)}{(x+2)(x-2)(2x+5)}$

51. $y = \cosh^{-1}(\sinh x) \quad \Rightarrow \quad y' = \dfrac{1}{\sqrt{(\sinh x)^2 - 1}}\cdot\cosh x = \dfrac{\cosh x}{\sqrt{\sinh^2 x - 1}}$

52. $y = x\tanh^{-1}\sqrt{x} \quad \Rightarrow \quad y' = \tanh^{-1}\sqrt{x} + x\dfrac{1}{1-\left(\sqrt{x}\right)^2}\dfrac{1}{2\sqrt{x}} = \tanh^{-1}\sqrt{x} + \dfrac{\sqrt{x}}{2(1-x)}$

53. $y = \cos\left(e^{\sqrt{\tan 3x}}\right) \quad \Rightarrow$

$$y' = -\sin\left(e^{\sqrt{\tan 3x}}\right)\cdot\left(e^{\sqrt{\tan 3x}}\right)' = -\sin\left(e^{\sqrt{\tan 3x}}\right)\, e^{\sqrt{\tan 3x}}\cdot\tfrac{1}{2}(\tan 3x)^{-1/2}\cdot\sec^2(3x)\cdot 3$$

$$= \frac{-3\sin\left(e^{\sqrt{\tan 3x}}\right)e^{\sqrt{\tan 3x}}\,\sec^2(3x)}{2\sqrt{\tan 3x}}$$

54. $y = \sin^2\left(\cos\sqrt{\sin\pi x}\right) = \left[\sin\left(\cos\sqrt{\sin\pi x}\right)\right]^2 \quad \Rightarrow$

$$y' = 2\left[\sin\left(\cos\sqrt{\sin\pi x}\right)\right]\left[\sin\left(\cos\sqrt{\sin\pi x}\right)\right]' = 2\sin\left(\cos\sqrt{\sin\pi x}\right)\cos\left(\cos\sqrt{\sin\pi x}\right)\left(\cos\sqrt{\sin\pi x}\right)'$$

$$= 2\sin\left(\cos\sqrt{\sin\pi x}\right)\cos\left(\cos\sqrt{\sin\pi x}\right)\left(-\sin\sqrt{\sin\pi x}\right)\left(\sqrt{\sin\pi x}\right)'$$

$$= -2\sin\left(\cos\sqrt{\sin\pi x}\right)\cos\left(\cos\sqrt{\sin\pi x}\right)\sin\sqrt{\sin\pi x}\cdot\tfrac{1}{2}(\sin\pi x)^{-1/2}(\sin\pi x)'$$

$$= \frac{-\sin\left(\cos\sqrt{\sin\pi x}\right)\cos\left(\cos\sqrt{\sin\pi x}\right)\sin\sqrt{\sin\pi x}}{\sqrt{\sin\pi x}}\cdot\cos\pi x\cdot\pi$$

$$= \frac{-\pi\sin\left(\cos\sqrt{\sin\pi x}\right)\cos\left(\cos\sqrt{\sin\pi x}\right)\sin\sqrt{\sin\pi x}\,\cos\pi x}{\sqrt{\sin\pi x}}$$

55. $f(t) = \sqrt{4t+1} \quad \Rightarrow \quad f'(t) = \frac{1}{2}(4t+1)^{-1/2}\cdot 4 = 2(4t+1)^{-1/2} \quad \Rightarrow$

$f''(t) = 2(-\frac{1}{2})(4t+1)^{-3/2}\cdot 4 = -4/(4t+1)^{3/2}$, so $f''(2) = -4/9^{3/2} = -\frac{4}{27}$.

56. $g(\theta) = \theta\sin\theta \quad \Rightarrow \quad g'(\theta) = \theta\cos\theta + \sin\theta\cdot 1 \quad \Rightarrow \quad g''(\theta) = \theta(-\sin\theta) + \cos\theta\cdot 1 + \cos\theta = 2\cos\theta - \theta\sin\theta$,

so $g''(\pi/6) = 2\cos(\pi/6) - (\pi/6)\sin(\pi/6) = 2\left(\sqrt{3}/2\right) - (\pi/6)(1/2) = \sqrt{3} - \pi/12$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. $x^6 + y^6 = 1 \quad \Rightarrow \quad 6x^5 + 6y^5 y' = 0 \quad \Rightarrow \quad y' = -x^5/y^5 \quad \Rightarrow$

$$y'' = -\frac{y^5(5x^4) - x^5(5y^4 y')}{(y^5)^2} = -\frac{5x^4 y^4\left[y - x(-x^5/y^5)\right]}{y^{10}} = -\frac{5x^4\left[(y^6 + x^6)/y^5\right]}{y^6} = -\frac{5x^4}{y^{11}}$$

58. $f(x) = (2-x)^{-1} \quad \Rightarrow \quad f'(x) = (2-x)^{-2} \quad \Rightarrow \quad f''(x) = 2(2-x)^{-3} \quad \Rightarrow \quad f'''(x) = 2 \cdot 3(2-x)^{-4} \quad \Rightarrow$

$f^{(4)}(x) = 2 \cdot 3 \cdot 4(2-x)^{-5}$. In general, $f^{(n)}(x) = 2 \cdot 3 \cdot 4 \cdot \cdots \cdot n(2-x)^{-(n+1)} = \dfrac{n!}{(2-x)^{(n+1)}}$.

59. We first show it is true for $n = 1$: $f(x) = xe^x \quad \Rightarrow \quad f'(x) = xe^x + e^x = (x+1)e^x$. We now assume it is true for $n = k$: $f^{(k)}(x) = (x+k)e^x$. With this assumption, we must show it is true for $n = k+1$:

$f^{(k+1)}(x) = \dfrac{d}{dx}\left[f^{(k)}(x)\right] = \dfrac{d}{dx}\left[(x+k)e^x\right] = (x+k)e^x + e^x = [(x+k)+1]e^x = [x+(k+1)]e^x$.

Therefore, $f^{(n)}(x) = (x+n)e^x$ by mathematical induction.

60. $$\lim_{t\to 0} \frac{t^3}{\tan^3(2t)} = \lim_{t\to 0} \frac{t^3\cos^3(2t)}{\sin^3(2t)} = \lim_{t\to 0} \cos^3(2t) \cdot \frac{1}{8\dfrac{\sin^3(2t)}{(2t)^3}} = \lim_{t\to 0} \frac{\cos^3(2t)}{8\left(\lim\limits_{t\to 0} \dfrac{\sin(2t)}{2t}\right)^3} = \frac{1}{8 \cdot 1^3} = \frac{1}{8}$$

61. $y = 4\sin^2 x \quad \Rightarrow \quad y' = 4 \cdot 2\sin x \cos x$. At $\left(\frac{\pi}{6}, 1\right)$, $y' = 8 \cdot \frac{1}{2} \cdot \frac{\sqrt{3}}{2} = 2\sqrt{3}$, so an equation of the tangent line is $y - 1 = 2\sqrt{3}\left(x - \frac{\pi}{6}\right)$, or $y = 2\sqrt{3}\,x + 1 - \pi\sqrt{3}/3$.

62. $y = \dfrac{x^2-1}{x^2+1} \quad \Rightarrow \quad y' = \dfrac{(x^2+1)(2x) - (x^2-1)(2x)}{(x^2+1)^2} = \dfrac{4x}{(x^2+1)^2}$.

At $(0, -1)$, $y' = 0$, so an equation of the tangent line is $y + 1 = 0(x - 0)$, or $y = -1$.

63. $y = \sqrt{1 + 4\sin x} \quad \Rightarrow \quad y' = \frac{1}{2}(1 + 4\sin x)^{-1/2} \cdot 4\cos x = \dfrac{2\cos x}{\sqrt{1 + 4\sin x}}$.

At $(0, 1)$, $y' = \dfrac{2}{\sqrt{1}} = 2$, so an equation of the tangent line is $y - 1 = 2(x - 0)$, or $y = 2x + 1$.

64. $x^2 + 4xy + y^2 = 13 \quad \Rightarrow \quad 2x + 4(xy' + y \cdot 1) + 2yy' = 0 \quad \Rightarrow \quad x + 2xy' + 2y + yy' = 0 \quad \Rightarrow$

$2xy' + yy' = -x - 2y \quad \Rightarrow \quad y'(2x + y) = -x - 2y \quad \Rightarrow \quad y' = \dfrac{-x-2y}{2x+y}$.

At $(2, 1)$, $y' = \dfrac{-2-2}{4+1} = -\dfrac{4}{5}$, so an equation of the tangent line is $y - 1 = -\frac{4}{5}(x - 2)$, or $y = -\frac{4}{5}x + \frac{13}{5}$.

The slope of the normal line is $\frac{5}{4}$, so an equation of the normal line is $y - 1 = \frac{5}{4}(x - 2)$, or $y = \frac{5}{4}x - \frac{3}{2}$.

65. $y = (2+x)e^{-x} \quad \Rightarrow \quad y' = (2+x)(-e^{-x}) + e^{-x} \cdot 1 = e^{-x}[-(2+x) + 1] = e^{-x}(-x - 1)$.

At $(0, 2)$, $y' = 1(-1) = -1$, so an equation of the tangent line is $y - 2 = -1(x - 0)$, or $y = -x + 2$.

The slope of the normal line is 1, so an equation of the normal line is $y - 2 = 1(x - 0)$, or $y = x + 2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

66. $f(x) = xe^{\sin x} \quad \Rightarrow \quad f'(x) = x[e^{\sin x}(\cos x)] + e^{\sin x}(1) = e^{\sin x}(x\cos x + 1)$. As a check on our work, we notice from the graphs that $f'(x) > 0$ when f is increasing. Also, we see in the larger viewing rectangle a certain similarity in the graphs of f and f': the sizes of the oscillations of f and f' are linked.

67. (a) $f(x) = x\sqrt{5-x} \quad \Rightarrow$

$$f'(x) = x\left[\frac{1}{2}(5-x)^{-1/2}(-1)\right] + \sqrt{5-x} = \frac{-x}{2\sqrt{5-x}} + \sqrt{5-x}\cdot\frac{2\sqrt{5-x}}{2\sqrt{5-x}} = \frac{-x}{2\sqrt{5-x}} + \frac{2(5-x)}{2\sqrt{5-x}}$$

$$= \frac{-x+10-2x}{2\sqrt{5-x}} = \frac{10-3x}{2\sqrt{5-x}}$$

(b) At $(1, 2)$: $f'(1) = \frac{7}{4}$.

So an equation of the tangent line is $y - 2 = \frac{7}{4}(x-1)$ or $y = \frac{7}{4}x + \frac{1}{4}$.

At $(4, 4)$: $f'(4) = -\frac{2}{2} = -1$.

So an equation of the tangent line is $y - 4 = -1(x-4)$ or $y = -x + 8$.

(c)

(d)

The graphs look reasonable, since f' is positive where f has tangents with positive slope, and f' is negative where f has tangents with negative slope.

68. (a) $f(x) = 4x - \tan x \quad \Rightarrow \quad f'(x) = 4 - \sec^2 x \quad \Rightarrow \quad f''(x) = -2\sec x\,(\sec x \tan x) = -2\sec^2 x \tan x$.

(b)

We can see that our answers are reasonable, since the graph of f' is 0 where f has a horizontal tangent, and the graph of f' is positive where f has tangents with positive slope and negative where f has tangents with negative slope. The same correspondence holds between the graphs of f' and f''.

69. $y = \sin x + \cos x \quad \Rightarrow \quad y' = \cos x - \sin x = 0 \quad \Leftrightarrow \quad \cos x = \sin x$ and $0 \le x \le 2\pi \quad \Leftrightarrow \quad x = \frac{\pi}{4}$ or $\frac{5\pi}{4}$, so the points are $\left(\frac{\pi}{4}, \sqrt{2}\right)$ and $\left(\frac{5\pi}{4}, -\sqrt{2}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

70. $x^2 + 2y^2 = 1 \quad \Rightarrow \quad 2x + 4yy' = 0 \quad \Rightarrow \quad y' = -x/(2y) = 1 \quad \Leftrightarrow \quad x = -2y$. Since the points lie on the ellipse, we have $(-2y)^2 + 2y^2 = 1 \quad \Rightarrow \quad 6y^2 = 1 \quad \Rightarrow \quad y = \pm\frac{1}{\sqrt{6}}$. The points are $\left(-\frac{2}{\sqrt{6}}, \frac{1}{\sqrt{6}}\right)$ and $\left(\frac{2}{\sqrt{6}}, -\frac{1}{\sqrt{6}}\right)$.

71. $f(x) = (x-a)(x-b)(x-c) \quad \Rightarrow \quad f'(x) = (x-b)(x-c) + (x-a)(x-c) + (x-a)(x-b)$.

So $\dfrac{f'(x)}{f(x)} = \dfrac{(x-b)(x-c) + (x-a)(x-c) + (x-a)(x-b)}{(x-a)(x-b)(x-c)} = \dfrac{1}{x-a} + \dfrac{1}{x-b} + \dfrac{1}{x-c}$.

Or: $f(x) = (x-a)(x-b)(x-c) \quad \Rightarrow \quad \ln|f(x)| = \ln|x-a| + \ln|x-b| + \ln|x-c| \quad \Rightarrow$

$\dfrac{f'(x)}{f(x)} = \dfrac{1}{x-a} + \dfrac{1}{x-b} + \dfrac{1}{x-c}$

72. (a) $\cos 2x = \cos^2 x - \sin^2 x \quad \Rightarrow \quad -2\sin 2x = -2\cos x \sin x - 2\sin x \cos x \quad \Leftrightarrow \quad \sin 2x = 2\sin x \cos x$

(b) $\sin(x+a) = \sin x \cos a + \cos x \sin a \quad \Rightarrow \quad \cos(x+a) = \cos x \cos a - \sin x \sin a$.

73. (a) $S(x) = f(x) + g(x) \quad \Rightarrow \quad S'(x) = f'(x) + g'(x) \quad \Rightarrow \quad S'(1) = f'(1) + g'(1) = 3 + 1 = 4$

(b) $P(x) = f(x)\,g(x) \quad \Rightarrow \quad P'(x) = f(x)\,g'(x) + g(x)\,f'(x) \quad \Rightarrow$

$P'(2) = f(2)\,g'(2) + g(2)f'(2) = 1(4) + 1(2) = 4 + 2 = 6$

(c) $Q(x) = \dfrac{f(x)}{g(x)} \quad \Rightarrow \quad Q'(x) = \dfrac{g(x)\,f'(x) - f(x)\,g'(x)}{[g(x)]^2} \quad \Rightarrow$

$Q'(1) = \dfrac{g(1)\,f'(1) - f(1)\,g'(1)}{[g(1)]^2} = \dfrac{3(3) - 2(1)}{3^2} = \dfrac{9-2}{9} = \dfrac{7}{9}$

(d) $C(x) = f(g(x)) \quad \Rightarrow \quad C'(x) = f'(g(x))\,g'(x) \quad \Rightarrow \quad C'(2) = f'(g(2))\,g'(2) = f'(1) \cdot 4 = 3 \cdot 4 = 12$

74. (a) $P(x) = f(x)\,g(x) \quad \Rightarrow \quad P'(x) = f(x)\,g'(x) + g(x)\,f'(x) \quad \Rightarrow$

$P'(2) = f(2)\,g'(2) + g(2)\,f'(2) = (1)\left(\frac{6-0}{3-0}\right) + (4)\left(\frac{0-3}{3-0}\right) = (1)(2) + (4)(-1) = 2 - 4 = -2$

(b) $Q(x) = \dfrac{f(x)}{g(x)} \quad \Rightarrow \quad Q'(x) = \dfrac{g(x)\,f'(x) - f(x)\,g'(x)}{[g(x)]^2} \quad \Rightarrow$

$Q'(2) = \dfrac{g(2)\,f'(2) - f(2)\,g'(2)}{[g(2)]^2} = \dfrac{(4)(-1) - (1)(2)}{4^2} = \dfrac{-6}{16} = -\dfrac{3}{8}$

(c) $C(x) = f(g(x)) \quad \Rightarrow \quad C'(x) = f'(g(x))g'(x) \quad \Rightarrow$

$C'(2) = f'(g(2))g'(2) = f'(4)g'(2) = \left(\frac{6-0}{5-3}\right)(2) = (3)(2) = 6$

75. $f(x) = x^2 g(x) \quad \Rightarrow \quad f'(x) = x^2 g'(x) + g(x)(2x)$ or $x[xg'(x) + 2g(x)]$

76. $f(x) = g(x^2) \quad \Rightarrow \quad f'(x) = g'(x^2)(2x) = 2xg'(x^2)$

77. $f(x) = [\,g(x)]^2 \quad \Rightarrow \quad f'(x) = 2[\,g(x)] \cdot g'(x) = 2g(x)\,g'(x)$

78. $f(x) = g(g(x)) \quad \Rightarrow \quad f'(x) = g'(g(x))\,g'(x)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

79. $f(x) = g(e^x) \quad \Rightarrow \quad f'(x) = g'(e^x)\, e^x$

80. $f(x) = e^{g(x)} \quad \Rightarrow \quad f'(x) = e^{g(x)} g'(x)$

81. $f(x) = \ln|g(x)| \quad \Rightarrow \quad f'(x) = \dfrac{1}{g(x)} g'(x) = \dfrac{g'(x)}{g(x)}$

82. $f(x) = g(\ln x) \quad \Rightarrow \quad f'(x) = g'(\ln x) \cdot \dfrac{1}{x} = \dfrac{g'(\ln x)}{x}$

83. $h(x) = \dfrac{f(x)\, g(x)}{f(x) + g(x)} \quad \Rightarrow$

$$\begin{aligned} h'(x) &= \frac{[f(x)+g(x)]\,[f(x)\,g'(x) + g(x)\,f'(x)] - f(x)\,g(x)\,[f'(x)+g'(x)]}{[f(x)+g(x)]^2} \\ &= \frac{[f(x)]^2\,g'(x) + f(x)\,g(x)\,f'(x) + f(x)\,g(x)\,g'(x) + [g(x)]^2\,f'(x) - f(x)\,g(x)\,f'(x) - f(x)\,g(x)\,g'(x)}{[f(x)+g(x)]^2} \\ &= \frac{f'(x)\,[g(x)]^2 + g'(x)\,[f(x)]^2}{[f(x)+g(x)]^2} \end{aligned}$$

84. $h(x) = \sqrt{\dfrac{f(x)}{g(x)}} \quad \Rightarrow \quad h'(x) = \dfrac{f'(x)\,g(x) - f(x)\,g'(x)}{2\sqrt{f(x)/g(x)}\,[g(x)]^2} = \dfrac{f'(x)\,g(x) - f(x)\,g'(x)}{2[g(x)]^{3/2}\sqrt{f(x)}}$

85. Using the Chain Rule repeatedly, $h(x) = f(g(\sin 4x)) \quad \Rightarrow$

$$h'(x) = f'(g(\sin 4x)) \cdot \frac{d}{dx}(g(\sin 4x)) = f'(g(\sin 4x)) \cdot g'(\sin 4x) \cdot \frac{d}{dx}(\sin 4x) = f'(g(\sin 4x))g'(\sin 4x)(\cos 4x)(4).$$

86. (a)

(b) The average rate of change is larger on $[2, 3]$.

(c) The instantaneous rate of change (the slope of the tangent) is larger at $x = 2$.

(d) $f(x) = x - 2\sin x \quad \Rightarrow \quad f'(x) = 1 - 2\cos x$,

so $f'(2) = 1 - 2\cos 2 \approx 1.8323$ and $f'(5) = 1 - 2\cos 5 \approx 0.4327$.

So $f'(2) > f'(5)$, as predicted in part (c).

87. $y = [\ln(x+4)]^2 \quad \Rightarrow \quad y' = 2[\ln(x+4)]^1 \cdot \dfrac{1}{x+4} \cdot 1 = 2\,\dfrac{\ln(x+4)}{x+4}$ and $y' = 0 \quad \Leftrightarrow \quad \ln(x+4) = 0 \quad \Leftrightarrow$

$x + 4 = e^0 \quad \Rightarrow \quad x + 4 = 1 \quad \Leftrightarrow \quad x = -3$, so the tangent is horizontal at the point $(-3, 0)$.

88. (a) The line $x - 4y = 1$ has slope $\frac{1}{4}$. A tangent to $y = e^x$ has slope $\frac{1}{4}$ when $y' = e^x = \frac{1}{4} \quad \Rightarrow \quad x = \ln\frac{1}{4} = -\ln 4$.

Since $y = e^x$, the y-coordinate is $\frac{1}{4}$ and the point of tangency is $\left(-\ln 4, \frac{1}{4}\right)$. Thus, an equation of the tangent line is $y - \frac{1}{4} = \frac{1}{4}(x + \ln 4)$ or $y = \frac{1}{4}x + \frac{1}{4}(\ln 4 + 1)$.

(b) The slope of the tangent at the point (a, e^a) is $\dfrac{d}{dx}\, e^x \Big|_{x\,=\,a} = e^a$. Thus, an equation of the tangent line is

$y - e^a = e^a(x - a)$. We substitute $x = 0$, $y = 0$ into this equation, since we want the line to pass through the origin:

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$0 - e^a = e^a(0 - a) \quad \Leftrightarrow \quad -e^a = e^a(-a) \quad \Leftrightarrow \quad a = 1$. So an equation of the tangent line at the point $(a, e^a) = (1, e)$ is $y - e = e(x - 1)$ or $y = ex$.

89. $y = f(x) = ax^2 + bx + c \quad \Rightarrow \quad f'(x) = 2ax + b$. We know that $f'(-1) = 6$ and $f'(5) = -2$, so $-2a + b = 6$ and $10a + b = -2$. Subtracting the first equation from the second gives $12a = -8 \quad \Rightarrow \quad a = -\frac{2}{3}$. Substituting $-\frac{2}{3}$ for a in the first equation gives $b = \frac{14}{3}$. Now $f(1) = 4 \quad \Rightarrow \quad 4 = a + b + c$, so $c = 4 + \frac{2}{3} - \frac{14}{3} = 0$ and hence, $f(x) = -\frac{2}{3}x^2 + \frac{14}{3}x$.

90. (a) $\lim\limits_{t\to\infty} C(t) = \lim\limits_{t\to\infty} [K(e^{-at} - e^{-bt})] = K \lim\limits_{t\to\infty} (e^{-at} - e^{-bt}) = K(0 - 0) = 0$ because $-at \to -\infty$ and $-bt \to -\infty$ as $t \to \infty$.

(b) $C(t) = K(e^{-at} - e^{-bt}) \quad \Rightarrow \quad C'(t) = K(e^{-at}(-a) - e^{-bt}(-b)) = K(-ae^{-at} + be^{-bt})$

(c) $C'(t) = 0 \quad \Leftrightarrow \quad be^{-bt} = ae^{-at} \quad \Leftrightarrow \quad \dfrac{b}{a} = e^{(-a+b)t} \quad \Leftrightarrow \quad \ln\dfrac{b}{a} = (b - a)t \quad \Leftrightarrow \quad t = \dfrac{\ln(b/a)}{b - a}$

91. $s(t) = Ae^{-ct}\cos(\omega t + \delta) \quad \Rightarrow$

$$v(t) = s'(t) = A\{e^{-ct}[-\omega\sin(\omega t + \delta)] + \cos(\omega t + \delta)(-ce^{-ct})\} = -Ae^{-ct}[\omega\sin(\omega t + \delta) + c\cos(\omega t + \delta)] \quad \Rightarrow$$

$$\begin{aligned} a(t) = v'(t) &= -A\{e^{-ct}[\omega^2\cos(\omega t + \delta) - c\omega\sin(\omega t + \delta)] + [\omega\sin(\omega t + \delta) + c\cos(\omega t + \delta)](-ce^{-ct})\} \\ &= -Ae^{-ct}[\omega^2\cos(\omega t + \delta) - c\omega\sin(\omega t + \delta) - c\omega\sin(\omega t + \delta) - c^2\cos(\omega t + \delta)] \\ &= -Ae^{-ct}[(\omega^2 - c^2)\cos(\omega t + \delta) - 2c\omega\sin(\omega t + \delta)] = Ae^{-ct}[(c^2 - \omega^2)\cos(\omega t + \delta) + 2c\omega\sin(\omega t + \delta)] \end{aligned}$$

92. (a) $x = \sqrt{b^2 + c^2t^2} \quad \Rightarrow \quad v(t) = x' = \left[1/\left(2\sqrt{b^2 + c^2t^2}\right)\right] 2c^2t = c^2t/\sqrt{b^2 + c^2t^2} \quad \Rightarrow$

$$a(t) = v'(t) = \frac{c^2\sqrt{b^2 + c^2t^2} - c^2t\left(c^2t/\sqrt{b^2 + c^2t^2}\right)}{b^2 + c^2t^2} = \frac{b^2c^2}{(b^2 + c^2t^2)^{3/2}}$$

(b) $v(t) > 0$ for $t > 0$, so the particle always moves in the positive direction.

93. (a) $y = t^3 - 12t + 3 \quad \Rightarrow \quad v(t) = y' = 3t^2 - 12 \quad \Rightarrow \quad a(t) = v'(t) = 6t$

(b) $v(t) = 3(t^2 - 4) > 0$ when $t > 2$, so it moves upward when $t > 2$ and downward when $0 \le t < 2$.

(c) Distance upward $= y(3) - y(2) = -6 - (-13) = 7$,

Distance downward $= y(0) - y(2) = 3 - (-13) = 16$. Total distance $= 7 + 16 = 23$.

(d)

(e) The particle is speeding up when v and a have the same sign, that is, when $t > 2$. The particle is slowing down when v and a have opposite signs; that is, when $0 < t < 2$.

94. (a) $V = \frac{1}{3}\pi r^2 h \quad \Rightarrow \quad dV/dh = \frac{1}{3}\pi r^2 \quad [r \text{ constant}]$

(b) $V = \frac{1}{3}\pi r^2 h \quad \Rightarrow \quad dV/dr = \frac{2}{3}\pi rh \quad [h \text{ constant}]$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

95. The linear density ρ is the rate of change of mass m with respect to length x.

$m = x\left(1+\sqrt{x}\right) = x + x^{3/2} \quad \Rightarrow \quad \rho = dm/dx = 1 + \frac{3}{2}\sqrt{x}$, so the linear density when $x = 4$ is $1 + \frac{3}{2}\sqrt{4} = 4$ kg/m.

96. (a) $C(x) = 920 + 2x - 0.02x^2 + 0.00007x^3 \quad \Rightarrow \quad C'(x) = 2 - 0.04x + 0.00021x^2$

(b) $C'(100) = 2 - 4 + 2.1 = \$0.10$/unit. This value represents the rate at which costs are increasing as the hundredth unit is produced, and is the approximate cost of producing the 101st unit.

(c) The cost of producing the 101st item is $C(101) - C(100) = 990.10107 - 990 = \$0.10107$, slightly larger than $C'(100)$.

97. (a) $y(t) = y(0)e^{kt} = 200e^{kt} \quad \Rightarrow \quad y(0.5) = 200e^{0.5k} = 360 \quad \Rightarrow \quad e^{0.5k} = 1.8 \quad \Rightarrow \quad 0.5k = \ln 1.8 \quad \Rightarrow$

$k = 2\ln 1.8 = \ln(1.8)^2 = \ln 3.24 \quad \Rightarrow \quad y(t) = 200e^{(\ln 3.24)t} = 200(3.24)^t$

(b) $y(4) = 200(3.24)^4 \approx 22{,}040$ cells

(c) $y'(t) = 200(3.24)^t \cdot \ln 3.24$, so $y'(4) = 200(3.24)^4 \cdot \ln 3.24 \approx 25{,}910$ cells per hour

(d) $200(3.24)^t = 10{,}000 \quad \Rightarrow \quad (3.24)^t = 50 \quad \Rightarrow \quad t\ln 3.24 = \ln 50 \quad \Rightarrow \quad t = \ln 50/\ln 3.24 \approx 3.33$ hours

98. (a) If $y(t)$ is the mass remaining after t years, then $y(t) = y(0)e^{kt} = 100e^{kt}$. $y(5.24) = 100e^{5.24k} = \frac{1}{2}\cdot 100 \quad \Rightarrow$

$e^{5.24k} = \frac{1}{2} \quad \Rightarrow \quad 5.24k = -\ln 2 \quad \Rightarrow \quad k = -\frac{1}{5.24}\ln 2 \quad \Rightarrow \quad y(t) = 100e^{-(\ln 2)t/5.24} = 100\cdot 2^{-t/5.24}$. Thus,

$y(20) = 100\cdot 2^{-20/5.24} \approx 7.1$ mg.

(b) $100\cdot 2^{-t/5.24} = 1 \quad \Rightarrow \quad 2^{-t/5.24} = \dfrac{1}{100} \quad \Rightarrow \quad -\dfrac{t}{5.24}\ln 2 = \ln\dfrac{1}{100} \quad \Rightarrow \quad t = 5.24\,\dfrac{\ln 100}{\ln 2} \approx 34.8$ years

99. (a) $C'(t) = -kC(t) \quad \Rightarrow \quad C(t) = C(0)e^{-kt}$ by Theorem 3.8.2. But $C(0) = C_0$, so $C(t) = C_0e^{-kt}$.

(b) $C(30) = \frac{1}{2}C_0$ since the concentration is reduced by half. Thus, $\frac{1}{2}C_0 = C_0e^{-30k} \quad \Rightarrow \ln\frac{1}{2} = -30k \quad \Rightarrow$

$k = -\frac{1}{30}\ln\frac{1}{2} = \frac{1}{30}\ln 2$. Since 10% of the original concentration remains if 90% is eliminated, we want the value of t such that $C(t) = \frac{1}{10}C_0$. Therefore, $\frac{1}{10}C_0 = C_0e^{-t(\ln 2)/30} \quad \Rightarrow \quad \ln 0.1 = -t(\ln 2)/30 \quad \Rightarrow \quad t = -\frac{30}{\ln 2}\ln 0.1 \approx 100$ h.

100. (a) If $y = u - 20$, $u(0) = 80 \quad \Rightarrow \quad y(0) = 80 - 20 = 60$, and the initial-value problem is $dy/dt = ky$ with $y(0) = 60$.

So the solution is $y(t) = 60e^{kt}$. Now $y(0.5) = 60e^{k(0.5)} = 60 - 20 \quad \Rightarrow \quad e^{0.5k} = \frac{40}{60} = \frac{2}{3} \quad \Rightarrow \quad k = 2\ln\frac{2}{3} = \ln\frac{4}{9}$,

so $y(t) = 60e^{(\ln 4/9)t} = 60(\frac{4}{9})^t$. Thus, $y(1) = 60(\frac{4}{9})^1 = \frac{80}{3} = 26\frac{2}{3}$ °C and $u(1) = 46\frac{2}{3}$ °C.

(b) $u(t) = 40 \quad \Rightarrow \quad y(t) = 20$. $y(t) = 60\left(\dfrac{4}{9}\right)^t = 20 \quad \Rightarrow \quad \left(\dfrac{4}{9}\right)^t = \dfrac{1}{3} \quad \Rightarrow \quad t\ln\dfrac{4}{9} = \ln\dfrac{1}{3} \quad \Rightarrow \quad t = \dfrac{\ln\frac{1}{3}}{\ln\frac{4}{9}} \approx 1.35$ h

or 81.3 min.

101. If $x =$ edge length, then $V = x^3 \quad \Rightarrow \quad dV/dt = 3x^2\,dx/dt = 10 \quad \Rightarrow \quad dx/dt = 10/(3x^2)$ and $S = 6x^2 \quad \Rightarrow$

$dS/dt = (12x)\,dx/dt = 12x[10/(3x^2)] = 40/x$. When $x = 30$, $dS/dt = \frac{40}{30} = \frac{4}{3}$ cm^2/min.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

102. Given $dV/dt = 2$, find dh/dt when $h = 5$. $V = \frac{1}{3}\pi r^2 h$ and, from similar triangles, $\dfrac{r}{h} = \dfrac{3}{10} \quad \Rightarrow \quad V = \dfrac{\pi}{3}\left(\dfrac{3h}{10}\right)^2 h = \dfrac{3\pi}{100}h^3$, so

$$2 = \frac{dV}{dt} = \frac{9\pi}{100}h^2\,\frac{dh}{dt} \quad \Rightarrow \quad \frac{dh}{dt} = \frac{200}{9\pi h^2} = \frac{200}{9\pi\,(5)^2} = \frac{8}{9\pi}\ \text{cm/s}$$

when $h = 5$.

103. Given $dh/dt = 5$ and $dx/dt = 15$, find dz/dt. $z^2 = x^2 + h^2 \quad \Rightarrow$

$2z\,\dfrac{dz}{dt} = 2x\,\dfrac{dx}{dt} + 2h\,\dfrac{dh}{dt} \quad \Rightarrow \quad \dfrac{dz}{dt} = \dfrac{1}{z}(15x + 5h)$. When $t = 3$,

$h = 45 + 3(5) = 60$ and $x = 15(3) = 45 \quad \Rightarrow \quad z = \sqrt{45^2 + 60^2} = 75$,

so $\dfrac{dz}{dt} = \dfrac{1}{75}[15(45) + 5(60)] = 13$ ft/s.

104. We are given $dz/dt = 30$ ft/s. By similar triangles, $\dfrac{y}{z} = \dfrac{4}{\sqrt{241}} \quad \Rightarrow$

$y = \dfrac{4}{\sqrt{241}}z$, so $\dfrac{dy}{dt} = \dfrac{4}{\sqrt{241}}\,\dfrac{dz}{dt} = \dfrac{120}{\sqrt{241}} \approx 7.7$ ft/s.

105. We are given $d\theta/dt = -0.25$ rad/h. $\tan\theta = 400/x \quad \Rightarrow$

$x = 400\cot\theta \quad \Rightarrow \quad \dfrac{dx}{dt} = -400\csc^2\theta\,\dfrac{d\theta}{dt}$. When $\theta = \frac{\pi}{6}$,

$\dfrac{dx}{dt} = -400(2)^2(-0.25) = 400$ ft/h.

106. (a) $f(x) = \sqrt{25 - x^2} \quad \Rightarrow \quad f'(x) = \dfrac{-2x}{2\sqrt{25 - x^2}} = -x(25 - x^2)^{-1/2}$.

So the linear approximation to $f(x)$ near 3 is $f(x) \approx f(3) + f'(3)(x - 3) = 4 - \frac{3}{4}(x - 3)$.

(b)

(c) For the required accuracy, we want $\sqrt{25 - x^2} - 0.1 < 4 - \frac{3}{4}(x - 3)$ and $4 - \frac{3}{4}(x - 3) < \sqrt{25 - x^2} + 0.1$. From the graph, it appears that these both hold for $2.24 < x < 3.66$.

107. (a) $f(x) = \sqrt[3]{1 + 3x} = (1 + 3x)^{1/3} \quad \Rightarrow \quad f'(x) = (1 + 3x)^{-2/3}$, so the linearization of f at $a = 0$ is

$L(x) = f(0) + f'(0)(x - 0) = 1^{1/3} + 1^{-2/3}x = 1 + x$. Thus, $\sqrt[3]{1 + 3x} \approx 1 + x \quad \Rightarrow$

$\sqrt[3]{1.03} = \sqrt[3]{1 + 3(0.01)} \approx 1 + (0.01) = 1.01$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The linear approximation is $\sqrt[3]{1+3x} \approx 1+x$, so for the required accuracy we want $\sqrt[3]{1+3x} - 0.1 < 1+x < \sqrt[3]{1+3x} + 0.1$. From the graph, it appears that this is true when $-0.235 < x < 0.401$.

108. $y = x^3 - 2x^2 + 1 \quad\Rightarrow\quad dy = (3x^2 - 4x)\,dx$. When $x = 2$ and $dx = 0.2$, $dy = \left[3(2)^2 - 4(2)\right](0.2) = 0.8$.

109. $A = x^2 + \frac{1}{2}\pi\left(\frac{1}{2}x\right)^2 = \left(1 + \frac{\pi}{8}\right)x^2 \quad\Rightarrow\quad dA = \left(2 + \frac{\pi}{4}\right)x\,dx$. When $x = 60$ and $dx = 0.1$, $dA = \left(2 + \frac{\pi}{4}\right)60(0.1) = 12 + \frac{3\pi}{2}$, so the maximum error is approximately $12 + \frac{3\pi}{2} \approx 16.7\ \text{cm}^2$.

110. $\displaystyle\lim_{x\to 1} \frac{x^{17} - 1}{x - 1} = \left[\frac{d}{dx}\,x^{17}\right]_{x=1} = 17(1)^{16} = 17$

111. $\displaystyle\lim_{h\to 0} \frac{\sqrt[4]{16+h} - 2}{h} = \left[\frac{d}{dx}\,\sqrt[4]{x}\right]_{x=16} = \frac{1}{4}x^{-3/4}\bigg|_{x=16} = \frac{1}{4\left(\sqrt[4]{16}\right)^3} = \frac{1}{32}$

112. $\displaystyle\lim_{\theta\to\pi/3} \frac{\cos\theta - 0.5}{\theta - \pi/3} = \left[\frac{d}{d\theta}\cos\theta\right]_{\theta=\pi/3} = -\sin\frac{\pi}{3} = -\frac{\sqrt{3}}{2}$

113.
$$\begin{aligned}
\lim_{x\to 0} \frac{\sqrt{1+\tan x} - \sqrt{1+\sin x}}{x^3} &= \lim_{x\to 0} \frac{\left(\sqrt{1+\tan x} - \sqrt{1+\sin x}\right)\left(\sqrt{1+\tan x} + \sqrt{1+\sin x}\right)}{x^3\left(\sqrt{1+\tan x} + \sqrt{1+\sin x}\right)} \\
&= \lim_{x\to 0} \frac{(1+\tan x) - (1+\sin x)}{x^3\left(\sqrt{1+\tan x} + \sqrt{1+\sin x}\right)} = \lim_{x\to 0} \frac{\sin x\,(1/\cos x - 1)}{x^3\left(\sqrt{1+\tan x} + \sqrt{1+\sin x}\right)} \cdot \frac{\cos x}{\cos x} \\
&= \lim_{x\to 0} \frac{\sin x\,(1-\cos x)}{x^3\left(\sqrt{1+\tan x} + \sqrt{1+\sin x}\right)\cos x} \cdot \frac{1+\cos x}{1+\cos x} \\
&= \lim_{x\to 0} \frac{\sin x \cdot \sin^2 x}{x^3\left(\sqrt{1+\tan x} + \sqrt{1+\sin x}\right)\cos x\,(1+\cos x)} \\
&= \left(\lim_{x\to 0} \frac{\sin x}{x}\right)^3 \lim_{x\to 0} \frac{1}{\left(\sqrt{1+\tan x} + \sqrt{1+\sin x}\right)\cos x\,(1+\cos x)} \\
&= 1^3 \cdot \frac{1}{\left(\sqrt{1} + \sqrt{1}\right)\cdot 1 \cdot (1+1)} = \frac{1}{4}
\end{aligned}$$

114. Differentiating the first given equation implicitly with respect to x and using the Chain Rule, we obtain $f(g(x)) = x \quad\Rightarrow$ $f'(g(x))\,g'(x) = 1 \quad\Rightarrow\quad g'(x) = \dfrac{1}{f'(g(x))}$. Using the second given equation to expand the denominator of this expression gives $g'(x) = \dfrac{1}{1 + [f(g(x))]^2}$. But the first given equation states that $f(g(x)) = x$, so $g'(x) = \dfrac{1}{1+x^2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

115. $\dfrac{d}{dx}\left[f(2x)\right] = x^2 \quad \Rightarrow \quad f'(2x)\cdot 2 = x^2 \quad \Rightarrow \quad f'(2x) = \frac{1}{2}x^2$. Let $t = 2x$. Then $f'(t) = \frac{1}{2}\left(\frac{1}{2}t\right)^2 = \frac{1}{8}t^2$, so $f'(x) = \frac{1}{8}x^2$.

116. Let (b, c) be on the curve, that is, $b^{2/3} + c^{2/3} = a^{2/3}$. Now $x^{2/3} + y^{2/3} = a^{2/3} \quad \Rightarrow \quad \frac{2}{3}x^{-1/3} + \frac{2}{3}y^{-1/3}\dfrac{dy}{dx} = 0$, so

$\dfrac{dy}{dx} = -\dfrac{y^{1/3}}{x^{1/3}} = -\left(\dfrac{y}{x}\right)^{1/3}$, so at (b, c) the slope of the tangent line is $-(c/b)^{1/3}$ and an equation of the tangent line is

$y - c = -(c/b)^{1/3}(x - b)$ or $y = -(c/b)^{1/3}x + (c + b^{2/3}c^{1/3})$. Setting $y = 0$, we find that the x-intercept is

$b^{1/3}c^{2/3} + b = b^{1/3}(c^{2/3} + b^{2/3}) = b^{1/3}a^{2/3}$ and setting $x = 0$ we find that the y-intercept is

$c + b^{2/3}c^{1/3} = c^{1/3}(c^{2/3} + b^{2/3}) = c^{1/3}a^{2/3}$. So the length of the tangent line between these two points is

$$\begin{aligned}\sqrt{(b^{1/3}a^{2/3})^2 + (c^{1/3}a^{2/3})^2} &= \sqrt{b^{2/3}a^{4/3} + c^{2/3}a^{4/3}} = \sqrt{(b^{2/3} + c^{2/3})a^{4/3}} \\ &= \sqrt{a^{2/3}a^{4/3}} = \sqrt{a^2} = a = \text{constant}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1. Let a be the x-coordinate of Q. Since the derivative of $y = 1 - x^2$ is $y' = -2x$, the slope at Q is $-2a$. But since the triangle is equilateral, $\overline{AO}/\overline{OC} = \sqrt{3}/1$, so the slope at Q is $-\sqrt{3}$. Therefore, we must have that $-2a = -\sqrt{3} \quad \Rightarrow \quad a = \frac{\sqrt{3}}{2}$.

Thus, the point Q has coordinates $\left(\frac{\sqrt{3}}{2}, 1 - \left(\frac{\sqrt{3}}{2}\right)^2\right) = \left(\frac{\sqrt{3}}{2}, \frac{1}{4}\right)$ and by symmetry, P has coordinates $\left(-\frac{\sqrt{3}}{2}, \frac{1}{4}\right)$.

2. $y = x^3 - 3x + 4 \quad \Rightarrow \quad y' = 3x^2 - 3$, and $y = 3(x^2 - x) \quad \Rightarrow \quad y' = 6x - 3$.

The slopes of the tangents of the two curves are equal when $3x^2 - 3 = 6x - 3$; that is, when $x = 0$ or 2. At $x = 0$, both tangents have slope -3, but the curves do not intersect. At $x = 2$, both tangents have slope 9 and the curves intersect at $(2, 6)$. So there is a common tangent line at $(2, 6)$, $y = 9x - 12$.

3.

We must show that r (in the figure) is halfway between p and q, that is, $r = (p + q)/2$. For the parabola $y = ax^2 + bx + c$, the slope of the tangent line is given by $y' = 2ax + b$. An equation of the tangent line at $x = p$ is $y - (ap^2 + bp + c) = (2ap + b)(x - p)$. Solving for y gives us

$$y = (2ap + b)x - 2ap^2 - bp + (ap^2 + bp + c)$$

or $$y = (2ap + b)x + c - ap^2 \qquad \textbf{(1)}$$

Similarly, an equation of the tangent line at $x = q$ is

$$y = (2aq + b)x + c - aq^2 \qquad \textbf{(2)}$$

We can eliminate y and solve for x by subtracting equation **(1)** from equation **(2)**.

$$\begin{aligned}
[(2aq + b) - (2ap + b)]x - aq^2 + ap^2 &= 0 \\
(2aq - 2ap)x &= aq^2 - ap^2 \\
2a(q - p)x &= a(q^2 - p^2) \\
x &= \frac{a(q + p)(q - p)}{2a(q - p)} = \frac{p + q}{2}
\end{aligned}$$

Thus, the x-coordinate of the point of intersection of the two tangent lines, namely r, is $(p + q)/2$.

4. We could differentiate and then simplify or we can simplify and then differentiate. The latter seems to be the simpler method.

$$\begin{aligned}
\frac{\sin^2 x}{1 + \cot x} + \frac{\cos^2 x}{1 + \tan x} &= \frac{\sin^2 x}{1 + \dfrac{\cos x}{\sin x}} \cdot \frac{\sin x}{\sin x} + \frac{\cos^2 x}{1 + \dfrac{\sin x}{\cos x}} \cdot \frac{\cos x}{\cos x} = \frac{\sin^3 x}{\sin x + \cos x} + \frac{\cos^3 x}{\cos x + \sin x} \\
&= \frac{\sin^3 x + \cos^3 x}{\sin x + \cos x} \quad [\text{factor sum of cubes}] \quad = \frac{(\sin x + \cos x)(\sin^2 x - \sin x \cos x + \cos^2 x)}{\sin x + \cos x} \\
&= \sin^2 x - \sin x \cos x + \cos^2 x = 1 - \sin x \cos x = 1 - \tfrac{1}{2}(2 \sin x \cos x) = 1 - \tfrac{1}{2} \sin 2x
\end{aligned}$$

Thus, $\dfrac{d}{dx}\left(\dfrac{\sin^2 x}{1 + \cot x} + \dfrac{\cos^2 x}{1 + \tan x}\right) = \dfrac{d}{dx}\left(1 - \tfrac{1}{2} \sin 2x\right) = -\tfrac{1}{2} \cos 2x \cdot 2 = -\cos 2x.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. Using $f'(a) = \lim\limits_{x \to a} \dfrac{f(x) - f(a)}{x - a}$, we recognize the given expression, $f(x) = \lim\limits_{t \to x} \dfrac{\sec t - \sec x}{t - x}$, as $g'(x)$ with $g(x) = \sec x$. Now $f'(\frac{\pi}{4}) = g''(\frac{\pi}{4})$, so we will find $g''(x)$. $g'(x) = \sec x \tan x \quad \Rightarrow$ $g''(x) = \sec x \sec^2 x + \tan x \sec x \tan x = \sec x(\sec^2 x + \tan^2 x)$, so $g''(\frac{\pi}{4}) = \sqrt{2}(\sqrt{2}^2 + 1^2) = \sqrt{2}(2+1) = 3\sqrt{2}$.

6. Using $f'(0) = \lim\limits_{x \to 0} \dfrac{f(x) - f(0)}{x - 0}$, we see that for the given equation, $\lim\limits_{x \to 0} \dfrac{\sqrt[3]{ax+b} - 2}{x} = \dfrac{5}{12}$, we have $f(x) = \sqrt[3]{ax+b}$, $f(0) = 2$, and $f'(0) = \frac{5}{12}$. Now $f(0) = 2 \quad \Leftrightarrow \quad \sqrt[3]{b} = 2 \quad \Leftrightarrow \quad b = 8$. Also $f'(x) = \frac{1}{3}(ax+b)^{-2/3} \cdot a$, so $f'(0) = \frac{5}{12} \quad \Leftrightarrow \quad \frac{1}{3}(8)^{-2/3} \cdot a = \frac{5}{12} \quad \Leftrightarrow \quad \frac{1}{3}(\frac{1}{4})a = \frac{5}{12} \quad \Leftrightarrow \quad a = 5$.

7. Let $y = \tan^{-1} x$. Then $\tan y = x$, so from the triangle we see that $\sin(\tan^{-1} x) = \sin y = \dfrac{x}{\sqrt{1+x^2}}$. Using this fact we have that

$$\sin(\tan^{-1}(\sinh x)) = \frac{\sinh x}{\sqrt{1 + \sinh^2 x}} = \frac{\sinh x}{\cosh x} = \tanh x.$$

Hence, $\sin^{-1}(\tanh x) = \sin^{-1}(\sin(\tan^{-1}(\sinh x))) = \tan^{-1}(\sinh x)$.

8. We find the equation of the parabola by substituting the point $(-100, 100)$, at which the car is situated, into the general equation $y = ax^2$: $100 = a(-100)^2 \quad \Rightarrow \quad a = \frac{1}{100}$. Now we find the equation of a tangent to the parabola at the point (x_0, y_0). We can show that $y' = a(2x) = \frac{1}{100}(2x) = \frac{1}{50}x$, so an equation of the tangent is $y - y_0 = \frac{1}{50}x_0(x - x_0)$. Since the point (x_0, y_0) is on the parabola, we must have $y_0 = \frac{1}{100}x_0^2$, so our equation of the tangent can be simplified to $y = \frac{1}{100}x_0^2 + \frac{1}{50}x_0(x - x_0)$. We want the statue to be located on the tangent line, so we substitute its coordinates $(100, 50)$ into this equation: $50 = \frac{1}{100}x_0^2 + \frac{1}{50}x_0(100 - x_0) \quad \Rightarrow \quad x_0^2 - 200x_0 + 5000 = 0 \quad \Rightarrow$ $x_0 = \frac{1}{2}\left[200 \pm \sqrt{200^2 - 4(5000)}\right] \quad \Rightarrow \quad x_0 = 100 \pm 50\sqrt{2}$. But $x_0 < 100$, so the car's headlights illuminate the statue when it is located at the point $\left(100 - 50\sqrt{2}, 150 - 100\sqrt{2}\right) \approx (29.3, 8.6)$, that is, about 29.3 m east and 8.6 m north of the origin.

9. We use mathematical induction. Let S_n be the statement that $\dfrac{d^n}{dx^n}(\sin^4 x + \cos^4 x) = 4^{n-1}\cos(4x + n\pi/2)$.

S_1 is true because

$$\begin{aligned}\frac{d}{dx}(\sin^4 x + \cos^4 x) &= 4\sin^3 x\,\cos x - 4\cos^3 x\,\sin x = 4\sin x\,\cos x\left(\sin^2 x - \cos^2 x\right)x \\ &= -4\sin x\,\cos x\,\cos 2x = -2\sin 2x\,\cos 2 = -\sin 4x = \sin(-4x) \\ &= \cos\left(\tfrac{\pi}{2} - (-4x)\right) = \cos\left(\tfrac{\pi}{2} + 4x\right) = 4^{n-1}\cos\left(4x + n\tfrac{\pi}{2}\right) \text{ when } n = 1\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Now assume S_k is true, that is, $\dfrac{d^k}{dx^k}\left(\sin^4 x+\cos^4 x\right)=4^{k-1}\cos\left(4x+k\frac{\pi}{2}\right)$. Then

$$\begin{aligned}\frac{d^{k+1}}{dx^{k+1}}\left(\sin^4 x+\cos^4 x\right)&=\frac{d}{dx}\left[\frac{d^k}{dx^k}(\sin^4 x+\cos^4 x)\right]=\frac{d}{dx}\left[4^{k-1}\cos\left(4x+k\tfrac{\pi}{2}\right)\right]\\&=-4^{k-1}\sin\left(4x+k\tfrac{\pi}{2}\right)\cdot\frac{d}{dx}\left(4x+k\tfrac{\pi}{2}\right)=-4^k\sin\left(4x+k\tfrac{\pi}{2}\right)\\&=4^k\sin\left(-4x-k\tfrac{\pi}{2}\right)=4^k\cos\left(\tfrac{\pi}{2}-\left(-4x-k\tfrac{\pi}{2}\right)\right)=4^k\cos\left(4x+(k+1)\tfrac{\pi}{2}\right)\end{aligned}$$

which shows that S_{k+1} is true.

Therefore, $\dfrac{d^n}{dx^n}\left(\sin^4 x+\cos^4 x\right)=4^{n-1}\cos\left(4x+n\frac{\pi}{2}\right)$ for every positive integer n, by mathematical induction.

Another proof: First write

$$\sin^4 x+\cos^4 x=(\sin^2 x+\cos^2 x)^2-2\sin^2 x\,\cos^2 x=1-\tfrac{1}{2}\sin^2 2x=1-\tfrac{1}{4}(1-\cos 4x)=\tfrac{3}{4}+\tfrac{1}{4}\cos 4x$$

Then we have $\dfrac{d^n}{dx^n}\left(\sin^4 x+\cos^4 x\right)=\dfrac{d^n}{dx^n}\left(\dfrac{3}{4}+\dfrac{1}{4}\cos 4x\right)=\dfrac{1}{4}\cdot 4^n\cos\left(4x+n\dfrac{\pi}{2}\right)=4^{n-1}\cos\left(4x+n\dfrac{\pi}{2}\right)$.

10. $$\begin{aligned}\lim_{x\to a}\frac{f(x)-f(a)}{\sqrt{x}-\sqrt{a}}&=\lim_{x\to a}\left[\frac{f(x)-f(a)}{\sqrt{x}-\sqrt{a}}\cdot\frac{\sqrt{x}+\sqrt{a}}{\sqrt{x}+\sqrt{a}}\right]=\lim_{x\to a}\left[\frac{f(x)-f(a)}{x-a}\cdot\left(\sqrt{x}+\sqrt{a}\right)\right]\\&=\lim_{x\to a}\frac{f(x)-f(a)}{x-a}\cdot\lim_{x\to a}\left(\sqrt{x}+\sqrt{a}\right)=f'(a)\cdot\left(\sqrt{a}+\sqrt{a}\right)=2\sqrt{a}\,f'(a)\end{aligned}$$

11. We must find a value x_0 such that the normal lines to the parabola $y=x^2$ at $x=\pm x_0$ intersect at a point one unit from the points $\left(\pm x_0, x_0^2\right)$. The normals to $y=x^2$ at $x=\pm x_0$ have slopes $-\dfrac{1}{\pm 2x_0}$ and pass through $\left(\pm x_0, x_0^2\right)$ respectively, so the normals have the equations $y-x_0^2=-\dfrac{1}{2x_0}(x-x_0)$ and $y-x_0^2=\dfrac{1}{2x_0}(x+x_0)$. The common y-intercept is $x_0^2+\dfrac{1}{2}$. We want to find the value of x_0 for which the distance from $\left(0, x_0^2+\frac{1}{2}\right)$ to $\left(x_0, x_0^2\right)$ equals 1. The square of the distance is $(x_0-0)^2+\left[x_0^2-\left(x_0^2+\frac{1}{2}\right)\right]^2=x_0^2+\frac{1}{4}=1 \quad\Leftrightarrow\quad x_0=\pm\frac{\sqrt{3}}{2}$. For these values of x_0, the y-intercept is $x_0^2+\frac{1}{2}=\frac{5}{4}$, so the center of the circle is at $\left(0,\frac{5}{4}\right)$.

Another solution: Let the center of the circle be $(0,a)$. Then the equation of the circle is $x^2+(y-a)^2=1$. Solving with the equation of the parabola, $y=x^2$, we get $x^2+(x^2-a)^2=1 \quad\Leftrightarrow\quad x^2+x^4-2ax^2+a^2=1 \quad\Leftrightarrow$ $x^4+(1-2a)x^2+a^2-1=0$. The parabola and the circle will be tangent to each other when this quadratic equation in x^2 has equal roots; that is, when the discriminant is 0. Thus, $(1-2a)^2-4(a^2-1)=0 \quad\Leftrightarrow$ $1-4a+4a^2-4a^2+4=0 \quad\Leftrightarrow\quad 4a=5$, so $a=\frac{5}{4}$. The center of the circle is $\left(0,\frac{5}{4}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. See the figure. The parabolas $y = 4x^2$ and $x = c + 2y^2$ intersect each other at right angles at the point (a, b) if and only if (a, b) satisfies both equations and the tangent lines at (a, b) are perpendicular. $y = 4x^2 \quad \Rightarrow \quad y' = 8x$ and $x = c + 2y^2 \quad \Rightarrow \quad 1 = 4y\,y' \quad \Rightarrow \quad y' = \dfrac{1}{4y}$, so at (a, b) we must

have $8a = -\dfrac{1}{1/(4b)} \quad \Rightarrow \quad 8a = -4b \quad \Rightarrow \quad b = -2a$. Since (a, b) is on both parabolas, we have **(1)** $b = 4a^2$ and **(2)** $a = c + 2b^2$. Substituting $-2a$ for b in **(1)** gives us $-2a = 4a^2 \quad \Rightarrow \quad 4a^2 + 2a = 0 \quad \Rightarrow \quad 2a(2a + 1) = 0 \quad \Rightarrow \quad a = 0$ or $a = -\frac{1}{2}$.

If $a = 0$, then $b = 0$ and $c = 0$, and the tangent lines at $(0, 0)$ are $y = 0$ and $x = 0$.

If $a = -\frac{1}{2}$, then $b = -2\left(-\frac{1}{2}\right) = 1$ and $-\frac{1}{2} = c + 2(1)^2 \quad \Rightarrow \quad c = -\frac{5}{2}$, and the tangent lines at $\left(-\frac{1}{2}, 1\right)$ are $y - 1 = -4\left(x + \frac{1}{2}\right)$ [or $y = -4x - 1$] and $y - 1 = \frac{1}{4}\left(x + \frac{1}{2}\right)$ $\left[\text{or } y = \frac{1}{4}x + \frac{9}{8}\right]$.

13. See the figure. Clearly, the line $y = 2$ is tangent to both circles at the point $(0, 2)$. We'll look for a tangent line L through the points (a, b) and (c, d), and if such a line exists, then its reflection through the y-axis is another such line. The slope of L is the same at (a, b) and (c, d). Find those slopes: $x^2 + y^2 = 4 \quad \Rightarrow$ $2x + 2y\,y' = 0 \quad \Rightarrow \quad y' = -\dfrac{x}{y} \quad \left[= -\dfrac{a}{b}\right]$ and $x^2 + (y - 3)^2 = 1 \quad \Rightarrow$ $2x + 2(y - 3)y' = 0 \quad \Rightarrow \quad y' = -\dfrac{x}{y - 3} \quad \left[= -\dfrac{c}{d - 3}\right]$.

Now an equation for L can be written using either point-slope pair, so we get $y - b = -\dfrac{a}{b}(x - a) \quad \left[\text{or } y = -\dfrac{a}{b}x + \dfrac{a^2}{b} + b\right]$ and $y - d = -\dfrac{c}{d - 3}(x - c) \quad \left[\text{or } y = -\dfrac{c}{d - 3}\,x + \dfrac{c^2}{d - 3} + d\right]$. The slopes are equal, so $-\dfrac{a}{b} = -\dfrac{c}{d - 3} \quad \Leftrightarrow$ $d - 3 = \dfrac{bc}{a}$. Since (c, d) is a solution of $x^2 + (y - 3)^2 = 1$, we have $c^2 + (d - 3)^2 = 1$, so $c^2 + \left(\dfrac{bc}{a}\right)^2 = 1 \quad \Rightarrow$ $a^2c^2 + b^2c^2 = a^2 \quad \Rightarrow \quad c^2(a^2 + b^2) = a^2 \quad \Rightarrow \quad 4c^2 = a^2$ [since (a, b) is a solution of $x^2 + y^2 = 4$] $\Rightarrow \quad a = 2c$.

Now $d - 3 = \dfrac{bc}{a} \quad \Rightarrow \quad d = 3 + \dfrac{bc}{2c}$, so $d = 3 + \dfrac{b}{2}$. The y-intercepts are equal, so $\dfrac{a^2}{b} + b = \dfrac{c^2}{d - 3} + d \quad \Leftrightarrow$ $\dfrac{a^2}{b} + b = \dfrac{(a/2)^2}{b/2} + \left(3 + \dfrac{b}{2}\right) \quad \Leftrightarrow \quad \left[\dfrac{a^2}{b} + b = \dfrac{a^2}{2b} + 3 + \dfrac{b}{2}\right](2b) \quad \Leftrightarrow \quad 2a^2 + 2b^2 = a^2 + 6b + b^2 \quad \Leftrightarrow$ $a^2 + b^2 = 6b \quad \Leftrightarrow \quad 4 = 6b \quad \Leftrightarrow \quad b = \frac{2}{3}$. It follows that $d = 3 + \dfrac{b}{2} = \dfrac{10}{3}$, $a^2 = 4 - b^2 = 4 - \frac{4}{9} = \frac{32}{9} \quad \Rightarrow \quad a = \frac{4}{3}\sqrt{2}$, and $c^2 = 1 - (d - 3)^2 = 1 - \left(\frac{1}{3}\right)^2 = \frac{8}{9} \quad \Rightarrow \quad c = \frac{2}{3}\sqrt{2}$. Thus, L has equation $y - \dfrac{2}{3} = -\dfrac{(4/3)\sqrt{2}}{2/3}\left(x - \dfrac{4}{3}\sqrt{2}\right) \quad \Leftrightarrow$ $y - \frac{2}{3} = -2\sqrt{2}\left(x - \frac{4}{3}\sqrt{2}\right) \quad \Leftrightarrow \quad y = -2\sqrt{2}\,x + 6$. Its reflection has equation $y = 2\sqrt{2}\,x + 6$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

In summary, there are three lines tangent to both circles: $y = 2$ touches at $(0, 2)$, L touches at $\left(\frac{4}{3}\sqrt{2}, \frac{2}{3}\right)$ and $\left(\frac{2}{3}\sqrt{2}, \frac{10}{3}\right)$, and its reflection through the y-axis touches at $\left(-\frac{4}{3}\sqrt{2}, \frac{2}{3}\right)$ and $\left(-\frac{2}{3}\sqrt{2}, \frac{10}{3}\right)$.

14. $f(x) = \dfrac{x^{46} + x^{45} + 2}{1 + x} = \dfrac{x^{45}(x+1) + 2}{x+1} = \dfrac{x^{45}(x+1)}{x+1} + \dfrac{2}{x+1} = x^{45} + 2(x+1)^{-1}$, so

$f^{(46)}(x) = (x^{45})^{(46)} + 2\left[(x+1)^{-1}\right]^{(46)}$. The forty-sixth derivative of any forty-fifth degree polynomial is 0, so $(x^{45})^{46} = 0$. Thus, $f^{(46)}(x) = 2\left[(-1)(-2)(-3)\cdots(-46)(x+1)^{-47}\right] = 2(46!)(x+1)^{-47}$ and $f^{(46)}(3) = 2(46!)(4)^{-47}$ or $(46!)2^{-93}$.

15. We can assume without loss of generality that $\theta = 0$ at time $t = 0$, so that $\theta = 12\pi t$ rad. [The angular velocity of the wheel is 360 rpm $= 360 \cdot (2\pi \text{ rad})/(60 \text{ s}) = 12\pi$ rad/s.] Then the position of A as a function of time is

$A = (40\cos\theta, 40\sin\theta) = (40\cos 12\pi t, 40\sin 12\pi t)$, so $\sin\alpha = \dfrac{y}{1.2 \text{ m}} = \dfrac{40\sin\theta}{120} = \dfrac{\sin\theta}{3} = \dfrac{1}{3}\sin 12\pi t$.

(a) Differentiating the expression for $\sin\alpha$, we get $\cos\alpha \cdot \dfrac{d\alpha}{dt} = \dfrac{1}{3} \cdot 12\pi \cdot \cos 12\pi t = 4\pi\cos\theta$. When $\theta = \dfrac{\pi}{3}$, we have

$\sin\alpha = \dfrac{1}{3}\sin\theta = \dfrac{\sqrt{3}}{6}$, so $\cos\alpha = \sqrt{1 - \left(\dfrac{\sqrt{3}}{6}\right)^2} = \sqrt{\dfrac{11}{12}}$ and $\dfrac{d\alpha}{dt} = \dfrac{4\pi\cos\frac{\pi}{3}}{\cos\alpha} = \dfrac{2\pi}{\sqrt{11/12}} = \dfrac{4\pi\sqrt{3}}{\sqrt{11}} \approx 6.56$ rad/s.

(b) By the Law of Cosines, $|AP|^2 = |OA|^2 + |OP|^2 - 2\,|OA|\,|OP|\cos\theta \quad \Rightarrow$

$120^2 = 40^2 + |OP|^2 - 2 \cdot 40\,|OP|\cos\theta \quad \Rightarrow \quad |OP|^2 - (80\cos\theta)\,|OP| - 12{,}800 = 0 \quad \Rightarrow$

$|OP| = \frac{1}{2}\left(80\cos\theta \pm \sqrt{6400\cos^2\theta + 51{,}200}\right) = 40\cos\theta \pm 40\sqrt{\cos^2\theta + 8} = 40\left(\cos\theta + \sqrt{8 + \cos^2\theta}\right)$ cm

[since $|OP| > 0$]. As a check, note that $|OP| = 160$ cm when $\theta = 0$ and $|OP| = 80\sqrt{2}$ cm when $\theta = \frac{\pi}{2}$.

(c) By part (b), the x-coordinate of P is given by $x = 40\left(\cos\theta + \sqrt{8 + \cos^2\theta}\right)$, so

$$\frac{dx}{dt} = \frac{dx}{d\theta}\frac{d\theta}{dt} = 40\left(-\sin\theta - \frac{2\cos\theta\sin\theta}{2\sqrt{8 + \cos^2\theta}}\right) \cdot 12\pi = -480\pi\sin\theta\left(1 + \frac{\cos\theta}{\sqrt{8 + \cos^2\theta}}\right) \text{ cm/s.}$$

In particular, $dx/dt = 0$ cm/s when $\theta = 0$ and $dx/dt = -480\pi$ cm/s when $\theta = \frac{\pi}{2}$.

16. The equation of T_1 is $y - x_1^2 = 2x_1(x - x_1) = 2x_1x - 2x_1^2$ or $y = 2x_1x - x_1^2$. The equation of T_2 is $y = 2x_2x - x_2^2$. Solving for the point of intersection, we get $2x(x_1 - x_2) = x_1^2 - x_2^2 \quad \Rightarrow \quad x = \frac{1}{2}(x_1 + x_2)$. Therefore, the coordinates of P are $\left(\frac{1}{2}(x_1 + x_2), x_1x_2\right)$. So if the point of contact of T is $\left(a, a^2\right)$, then Q_1 is $\left(\frac{1}{2}(a + x_1), ax_1\right)$ and Q_2 is $\left(\frac{1}{2}(a + x_2), ax_2\right)$. Therefore,

$|PQ_1|^2 = \frac{1}{4}(a - x_2)^2 + x_1^2(a - x_2)^2 = (a - x_2)^2\left(\frac{1}{4} + x_1^2\right)$ and

$|PP_1|^2 = \frac{1}{4}(x_1 - x_2)^2 + x_1^2(x_1 - x_2)^2 = (x_1 - x_2)^2\left(\frac{1}{4} + x_1^2\right)$.

So $\dfrac{|PQ_1|^2}{|PP_1|^2} = \dfrac{(a - x_2)^2}{(x_1 - x_2)^2}$, and similarly $\dfrac{|PQ_2|^2}{|PP_2|^2} = \dfrac{(x_1 - a)^2}{(x_1 - x_2)^2}$. Finally, $\dfrac{|PQ_1|}{|PP_1|} + \dfrac{|PQ_2|}{|PP_2|} = \dfrac{a - x_2}{x_1 - x_2} + \dfrac{x_1 - a}{x_1 - x_2} = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. Consider the statement that $\dfrac{d^n}{dx^n}(e^{ax}\sin bx) = r^n e^{ax}\sin(bx+n\theta)$. For $n=1$,

$\dfrac{d}{dx}(e^{ax}\sin bx) = ae^{ax}\sin bx + be^{ax}\cos bx$, and

$re^{ax}\sin(bx+\theta) = re^{ax}[\sin bx\cos\theta + \cos bx\sin\theta] = re^{ax}\left(\dfrac{a}{r}\sin bx + \dfrac{b}{r}\cos bx\right) = ae^{ax}\sin bx + be^{ax}\cos bx$

since $\tan\theta = \dfrac{b}{a} \quad\Rightarrow\quad \sin\theta = \dfrac{b}{r}$ and $\cos\theta = \dfrac{a}{r}$. So the statement is true for $n=1$.

Assume it is true for $n=k$. Then

$$\begin{aligned}\frac{d^{k+1}}{dx^{k+1}}(e^{ax}\sin bx) &= \frac{d}{dx}\left[r^k e^{ax}\sin(bx+k\theta)\right] = r^k ae^{ax}\sin(bx+k\theta) + r^k e^{ax} b\cos(bx+k\theta)\\ &= r^k e^{ax}[a\sin(bx+k\theta) + b\cos(bx+k\theta)]\end{aligned}$$

But

$\sin[bx+(k+1)\theta] = \sin[(bx+k\theta)+\theta] = \sin(bx+k\theta)\cos\theta + \sin\theta\cos(bx+k\theta) = \frac{a}{r}\sin(bx+k\theta) + \frac{b}{r}\cos(bx+k\theta)$.

Hence, $a\sin(bx+k\theta) + b\cos(bx+k\theta) = r\sin[bx+(k+1)\theta]$. So

$\dfrac{d^{k+1}}{dx^{k+1}}(e^{ax}\sin bx) = r^k e^{ax}[a\sin(bx+k\theta)+b\cos(bx+k\theta)] = r^k e^{ax}[r\sin(bx+(k+1)\theta)] = r^{k+1}e^{ax}[\sin(bx+(k+1)\theta)]$.

Therefore, the statement is true for all n by mathematical induction.

18. We recognize this limit as the definition of the derivative of the function $f(x) = e^{\sin x}$ at $x=\pi$, since it is of the form

$\displaystyle\lim_{x\to\pi}\frac{f(x)-f(\pi)}{x-\pi}$. Therefore, the limit is equal to $f'(\pi) = (\cos\pi)e^{\sin\pi} = -1\cdot e^0 = -1$.

19. It seems from the figure that as P approaches the point $(0,2)$ from the right, $x_T\to\infty$ and $y_T\to 2^+$. As P approaches the point $(3,0)$ from the left, it appears that $x_T\to 3^+$ and $y_T\to\infty$. So we guess that $x_T\in(3,\infty)$ and $y_T\in(2,\infty)$. It is more difficult to estimate the range of values for x_N and y_N. We might perhaps guess that $x_N\in(0,3)$, and $y_N\in(-\infty,0)$ or $(-2,0)$.

In order to actually solve the problem, we implicitly differentiate the equation of the ellipse to find the equation of the tangent line: $\dfrac{x^2}{9}+\dfrac{y^2}{4}=1 \quad\Rightarrow\quad \dfrac{2x}{9}+\dfrac{2y}{4}y'=0$, so $y' = -\dfrac{4}{9}\dfrac{x}{y}$. So at the point (x_0,y_0) on the ellipse, an equation of the tangent line is $y-y_0 = -\dfrac{4}{9}\dfrac{x_0}{y_0}(x-x_0)$ or $4x_0x+9y_0y = 4x_0^2+9y_0^2$. This can be written as $\dfrac{x_0x}{9}+\dfrac{y_0y}{4} = \dfrac{x_0^2}{9}+\dfrac{y_0^2}{4} = 1$, because (x_0,y_0) lies on the ellipse. So an equation of the tangent line is $\dfrac{x_0x}{9}+\dfrac{y_0y}{4}=1$.

Therefore, the x-intercept x_T for the tangent line is given by $\dfrac{x_0x_T}{9}=1 \quad\Leftrightarrow\quad x_T = \dfrac{9}{x_0}$, and the y-intercept y_T is given by $\dfrac{y_0y_T}{4}=1 \quad\Leftrightarrow\quad y_T = \dfrac{4}{y_0}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

So as x_0 takes on all values in $(0,3)$, x_T takes on all values in $(3,\infty)$, and as y_0 takes on all values in $(0,2)$, y_T takes on all values in $(2,\infty)$. At the point (x_0,y_0) on the ellipse, the slope of the normal line is $-\dfrac{1}{y'(x_0,y_0)} = \dfrac{9}{4}\dfrac{y_0}{x_0}$, and its equation is $y - y_0 = \dfrac{9}{4}\dfrac{y_0}{x_0}(x - x_0)$. So the x-intercept x_N for the normal line is given by $0 - y_0 = \dfrac{9}{4}\dfrac{y_0}{x_0}(x_N - x_0) \quad\Rightarrow$ $x_N = -\dfrac{4x_0}{9} + x_0 = \dfrac{5x_0}{9}$, and the y-intercept y_N is given by $y_N - y_0 = \dfrac{9}{4}\dfrac{y_0}{x_0}(0 - x_0) \quad\Rightarrow\quad y_N = -\dfrac{9y_0}{4} + y_0 = -\dfrac{5y_0}{4}$.

So as x_0 takes on all values in $(0,3)$, x_N takes on all values in $\left(0,\frac{5}{3}\right)$, and as y_0 takes on all values in $(0,2)$, y_N takes on all values in $\left(-\frac{5}{2},0\right)$.

20. $\displaystyle\lim_{x\to 0}\frac{\sin(3+x)^2 - \sin 9}{x} = f'(3)$ where $f(x) = \sin x^2$. Now $f'(x) = (\cos x^2)(2x)$, so $f'(3) = 6\cos 9$.

21. (a) If the two lines L_1 and L_2 have slopes m_1 and m_2 and angles of inclination ϕ_1 and ϕ_2, then $m_1 = \tan\,\phi_1$ and $m_2 = \tan\,\phi_2$. The triangle in the figure shows that $\phi_1 + \alpha + (180^\circ - \phi_2) = 180^\circ$ and so $\alpha = \phi_2 - \phi_1$. Therefore, using the identity for $\tan(x-y)$, we have $\tan\alpha = \tan(\phi_2 - \phi_1) = \dfrac{\tan\phi_2 - \tan\phi_1}{1 + \tan\phi_2\tan\phi_1}$ and so $\tan\,\alpha = \dfrac{m_2 - m_1}{1 + m_1 m_2}$.

(b) (i) The parabolas intersect when $x^2 = (x-2)^2 \quad\Rightarrow\quad x = 1$. If $y = x^2$, then $y' = 2x$, so the slope of the tangent to $y = x^2$ at $(1,1)$ is $m_1 = 2(1) = 2$. If $y = (x-2)^2$, then $y' = 2(x-2)$, so the slope of the tangent to $y = (x-2)^2$ at $(1,1)$ is $m_2 = 2(1-2) = -2$. Therefore, $\tan\alpha = \dfrac{m_2 - m_1}{1 + m_1 m_2} = \dfrac{-2-2}{1+2(-2)} = \dfrac{4}{3}$ and so $\alpha = \tan^{-1}\left(\frac{4}{3}\right) \approx 53^\circ$ [or 127°].

(ii) $x^2 - y^2 = 3$ and $x^2 - 4x + y^2 + 3 = 0$ intersect when $x^2 - 4x + (x^2 - 3) + 3 = 0 \quad\Leftrightarrow\quad 2x(x-2) = 0 \quad\Rightarrow$ $x = 0$ or 2, but 0 is extraneous. If $x = 2$, then $y = \pm 1$. If $x^2 - y^2 = 3$ then $2x - 2yy' = 0 \quad\Rightarrow\quad y' = x/y$ and $x^2 - 4x + y^2 + 3 = 0 \quad\Rightarrow\quad 2x - 4 + 2yy' = 0 \quad\Rightarrow\quad y' = \dfrac{2-x}{y}$. At $(2,1)$ the slopes are $m_1 = 2$ and $m_2 = 0$, so $\tan\alpha = \frac{0-2}{1+2\cdot 0} = -2 \quad\Rightarrow\quad \alpha \approx 117^\circ$. At $(2,-1)$ the slopes are $m_1 = -2$ and $m_2 = 0$, so $\tan\alpha = \dfrac{0-(-2)}{1+(-2)(0)} = 2 \quad\Rightarrow\quad \alpha \approx 63^\circ$ [or 117°].

22. $y^2 = 4px \quad\Rightarrow\quad 2yy' = 4p \quad\Rightarrow\quad y' = 2p/y \quad\Rightarrow$ slope of tangent at $P(x_1,y_1)$ is $m_1 = 2p/y_1$. The slope of FP is $m_2 = \dfrac{y_1}{x_1 - p}$, so by the formula from Problem 21(a),

$$\tan\alpha = \frac{\dfrac{y_1}{x_1-p} - \dfrac{2p}{y_1}}{1 + \left(\dfrac{2p}{y_1}\right)\left(\dfrac{y_1}{x_1-p}\right)} \cdot \frac{y_1\,(x_1-p)}{y_1\,(x_1-p)} = \frac{y_1^2 - 2p(x_1-p)}{y_1(x_1-p) + 2py_1} = \frac{4px_1 - 2px_1 + 2p^2}{x_1y_1 - py_1 + 2py_1}$$

$$= \frac{2p(p+x_1)}{y_1(p+x_1)} = \frac{2p}{y_1} = \text{slope of tangent at } P = \tan\beta$$

Since $0 \le \alpha, \beta \le \frac{\pi}{2}$, this proves that $\alpha = \beta$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. Since $\angle ROQ = \angle OQP = \theta$, the triangle QOR is isosceles, so $|QR| = |RO| = x$. By the Law of Cosines, $x^2 = x^2 + r^2 - 2rx\cos\theta$. Hence, $2rx\cos\theta = r^2$, so $x = \dfrac{r^2}{2r\cos\theta} = \dfrac{r}{2\cos\theta}$. Note that as $y \to 0^+$, $\theta \to 0^+$ (since $\sin\theta = y/r$), and hence $x \to \dfrac{r}{2\cos 0} = \dfrac{r}{2}$. Thus, as P is taken closer and closer to the x-axis, the point R approaches the midpoint of the radius AO.

24. $$\lim_{x\to 0}\frac{f(x)}{g(x)} = \lim_{x\to 0}\frac{f(x)-0}{g(x)-0} = \lim_{x\to 0}\frac{f(x)-f(0)}{g(x)-g(0)} = \lim_{x\to 0}\frac{\dfrac{f(x)-f(0)}{x-0}}{\dfrac{g(x)-g(0)}{x-0}} = \frac{\lim\limits_{x\to 0}\dfrac{f(x)-f(0)}{x-0}}{\lim\limits_{x\to 0}\dfrac{g(x)-g(0)}{x-0}} = \frac{f'(0)}{g'(0)}$$

25. $$\lim_{x\to 0}\frac{\sin(a+2x) - 2\sin(a+x) + \sin a}{x^2}$$

$$= \lim_{x\to 0}\frac{\sin a\,\cos 2x + \cos a\,\sin 2x - 2\sin a\cos x - 2\cos a\sin x + \sin a}{x^2}$$

$$= \lim_{x\to 0}\frac{\sin a\,(\cos 2x - 2\cos x + 1) + \cos a\,(\sin 2x - 2\sin x)}{x^2}$$

$$= \lim_{x\to 0}\frac{\sin a\,(2\cos^2 x - 1 - 2\cos x + 1) + \cos a\,(2\sin x\cos x - 2\sin x)}{x^2}$$

$$= \lim_{x\to 0}\frac{\sin a\,(2\cos x)(\cos x - 1) + \cos a\,(2\sin x)(\cos x - 1)}{x^2}$$

$$= \lim_{x\to 0}\frac{2(\cos x - 1)[\sin a\,\cos x + \cos a\,\sin x](\cos x + 1)}{x^2(\cos x + 1)}$$

$$= \lim_{x\to 0}\frac{-2\sin^2 x\,[\sin(a+x)]}{x^2(\cos x + 1)} = -2\lim_{x\to 0}\left(\frac{\sin x}{x}\right)^2\cdot\frac{\sin(a+x)}{\cos x + 1} = -2(1)^2\,\frac{\sin(a+0)}{\cos 0 + 1} = -\sin a$$

26. (a) $f(x) = x(x-2)(x-6) = x^3 - 8x^2 + 12x \quad\Rightarrow\quad f'(x) = 3x^2 - 16x + 12$. The average of the first pair of zeros is $(0+2)/2 = 1$. At $x = 1$, the slope of the tangent line is $f'(1) = -1$, so an equation of the tangent line has the form $y = -1x + b$. Since $f(1) = 5$, we have $5 = -1 + b \quad\Rightarrow\quad b = 6$ and the tangent has equation $y = -x + 6$.

Similarly, at $x = \dfrac{0+6}{2} = 3$, $y = -9x + 18$; at $x = \dfrac{2+6}{2} = 4$, $y = -4x$. From the graph, we see that each tangent line drawn at the average of two zeros intersects the graph of f at the third zero.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Using Exercise 3.2.63(a), $f(x) = (x-a)(x-b)(x-c) \quad \Rightarrow$

$f'(x) = (x-b)(x-c) + (x-a)(x-c) + (x-a)(x-b)$.

$$\begin{aligned} f'\left(\frac{a+b}{2}\right) &= \left(\frac{a+b}{2} - b\right)\left(\frac{a+b}{2} - c\right) + \left(\frac{a+b}{2} - a\right)\left(\frac{a+b}{2} - c\right) + \left(\frac{a+b}{2} - a\right)\left(\frac{a+b}{2} - b\right) \\ &= \left(\frac{a-b}{2}\right)\left(\frac{a+b-2c}{2}\right) + \left(\frac{-a+b}{2}\right)\left(\frac{a+b-2c}{2}\right) + \left(\frac{-a+b}{2}\right)\left(\frac{a-b}{2}\right) \\ &= \left(\frac{a-b}{2}\right)\left(\frac{a+b-2c}{2}\right) - \left(\frac{a-b}{2}\right)\left(\frac{a+b-2c}{2}\right) - \left(\frac{a-b}{2}\right)\left(\frac{a-b}{2}\right) \\ &= -\frac{(a-b)^2}{4} \end{aligned}$$

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &= \left(\frac{a+b}{2} - a\right)\left(\frac{a+b}{2} - b\right)\left(\frac{a+b}{2} - c\right) \\ &= \left(\frac{b-a}{2}\right)\left(\frac{a-b}{2}\right)\left(\frac{a+b-2c}{2}\right) \\ &= -\frac{(a-b)^2}{8}(a+b-2c) \end{aligned}$$

An equation of the tangent line at $x = \dfrac{a+b}{2}$ is $y = -\dfrac{(a-b)^2}{4}\left(x - \dfrac{a+b}{2}\right) - \dfrac{(a-b)^2}{8}(a+b-2c)$.

To find the x-intercept, let $y = 0$ and solve for x.

$$\begin{aligned} \frac{(a-b)^2}{4}\left(x - \frac{a+b}{2}\right) &= -\frac{(a-b)^2}{8}(a+b-2c) \\ x - \frac{a+b}{2} &= -\frac{1}{2}(a+b-2c) \\ x &= \tfrac{1}{2}a + \tfrac{1}{2}b - \tfrac{1}{2}a - \tfrac{1}{2}b + c = c \end{aligned}$$

The result is $x = c$, which is the third zero.

27.

Let $f(x) = e^{2x}$ and $g(x) = k\sqrt{x} \quad [k > 0]$. From the graphs of f and g, we see that f will intersect g exactly once when f and g share a tangent line. Thus, we must have $f = g$ and $f' = g'$ at $x = a$.

$$f(a) = g(a) \quad \Rightarrow \quad e^{2a} = k\sqrt{a} \quad (\star)$$

and $$f'(a) = g'(a) \quad \Rightarrow \quad 2e^{2a} = \frac{k}{2\sqrt{a}} \quad \Rightarrow \quad e^{2a} = \frac{k}{4\sqrt{a}}.$$

So we must have $k\sqrt{a} = \dfrac{k}{4\sqrt{a}} \quad \Rightarrow \quad \left(\sqrt{a}\right)^2 = \dfrac{k}{4k} \quad \Rightarrow \quad a = \frac{1}{4}$. From $(\star)$, $e^{2(1/4)} = k\sqrt{1/4} \quad \Rightarrow$

$k = 2e^{1/2} = 2\sqrt{e} \approx 3.297$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. We see that at $x = 0$, $f(x) = a^x = 1 + x = 1$, so if $y = a^x$ is to lie above $y = 1 + x$, the two curves must just touch at $(0, 1)$, that is, we must have $f'(0) = 1$. [To see this analytically, note that $a^x \geq 1 + x \quad \Rightarrow \quad a^x - 1 \geq x \quad \Rightarrow \quad \dfrac{a^x - 1}{x} \geq 1$ for $x > 0$, so $f'(0) = \lim\limits_{x \to 0^+} \dfrac{a^x - 1}{x} \geq 1$. Similarly, for $x < 0$, $a^x - 1 \geq x \quad \Rightarrow \quad \dfrac{a^x - 1}{x} \leq 1$, so $f'(0) = \lim\limits_{x \to 0^-} \dfrac{a^x - 1}{x} \leq 1$.

Since $1 \leq f'(0) \leq 1$, we must have $f'(0) = 1$.] But $f'(x) = a^x \ln a \quad \Rightarrow \quad f'(0) = \ln a$, so we have $\ln a = 1 \quad \Leftrightarrow \quad a = e$.

Another method: The inequality certainly holds for $x \leq -1$, so consider $x > -1$, $x \neq 0$. Then $a^x \geq 1 + x \quad \Rightarrow \quad a \geq (1 + x)^{1/x}$ for $x > 0 \quad \Rightarrow \quad a \geq \lim\limits_{x \to 0^+} (1 + x)^{1/x} = e$, by Equation 3.6.5. Also, $a^x \geq 1 + x \quad \Rightarrow \quad a \leq (1 + x)^{1/x}$ for $x < 0 \quad \Rightarrow \quad a \leq \lim\limits_{x \to 0^-} (1 + x)^{1/x} = e$. So since $e \leq a \leq e$, we must have $a = e$.

29. $y = \dfrac{x}{\sqrt{a^2 - 1}} - \dfrac{2}{\sqrt{a^2 - 1}} \arctan \dfrac{\sin x}{a + \sqrt{a^2 - 1} + \cos x}$. Let $k = a + \sqrt{a^2 - 1}$. Then

$$\begin{aligned} y' &= \frac{1}{\sqrt{a^2 - 1}} - \frac{2}{\sqrt{a^2 - 1}} \cdot \frac{1}{1 + \sin^2 x/(k + \cos x)^2} \cdot \frac{\cos x(k + \cos x) + \sin^2 x}{(k + \cos x)^2} \\ &= \frac{1}{\sqrt{a^2 - 1}} - \frac{2}{\sqrt{a^2 - 1}} \cdot \frac{k \cos x + \cos^2 x + \sin^2 x}{(k + \cos x)^2 + \sin^2 x} = \frac{1}{\sqrt{a^2 - 1}} - \frac{2}{\sqrt{a^2 - 1}} \cdot \frac{k \cos x + 1}{k^2 + 2k \cos x + 1} \\ &= \frac{k^2 + 2k \cos x + 1 - 2k \cos x - 2}{\sqrt{a^2 - 1}\,(k^2 + 2k \cos x + 1)} = \frac{k^2 - 1}{\sqrt{a^2 - 1}\,(k^2 + 2k \cos x + 1)} \end{aligned}$$

But $k^2 = 2a^2 + 2a\sqrt{a^2 - 1} - 1 = 2a\big(a + \sqrt{a^2 - 1}\big) - 1 = 2ak - 1$, so $k^2 + 1 = 2ak$, and $k^2 - 1 = 2(ak - 1)$.

So $y' = \dfrac{2(ak - 1)}{\sqrt{a^2 - 1}\,(2ak + 2k \cos x)} = \dfrac{ak - 1}{\sqrt{a^2 - 1}k\,(a + \cos x)}$. But $ak - 1 = a^2 + a\sqrt{a^2 - 1} - 1 = k\sqrt{a^2 - 1}$, so $y' = 1/(a + \cos x)$.

30. Suppose that $y = mx + c$ is a tangent line to the ellipse. Then it intersects the ellipse at only one point, so the discriminant of the equation $\dfrac{x^2}{a^2} + \dfrac{(mx + c)^2}{b^2} = 1 \quad \Leftrightarrow \quad (b^2 + a^2m^2)x^2 + 2mca^2x + a^2c^2 - a^2b^2 = 0$ must be 0; that is,

$$\begin{aligned} 0 &= (2mca^2)^2 - 4(b^2 + a^2m^2)(a^2c^2 - a^2b^2) = 4a^4c^2m^2 - 4a^2b^2c^2 + 4a^2b^4 - 4a^4c^2m^2 + 4a^4b^2m^2 \\ &= 4a^2b^2(a^2m^2 + b^2 - c^2) \end{aligned}$$

Therefore, $a^2m^2 + b^2 - c^2 = 0$.

Now if a point (α, β) lies on the line $y = mx + c$, then $c = \beta - m\alpha$, so from above,

$$0 = a^2m^2 + b^2 - (\beta - m\alpha)^2 = (a^2 - \alpha^2)m^2 + 2\alpha\beta m + b^2 - \beta^2 \quad \Leftrightarrow \quad m^2 + \frac{2\alpha\beta}{a^2 - \alpha^2}m + \frac{b^2 - \beta^2}{a^2 - \alpha^2} = 0.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(a) Suppose that the two tangent lines from the point (α, β) to the ellipse have slopes m and $\dfrac{1}{m}$. Then m and $\dfrac{1}{m}$ are roots of the equation $z^2 + \dfrac{2\alpha\beta}{a^2 - \alpha^2} z + \dfrac{b^2 - \beta^2}{a^2 - \alpha^2} = 0$. This implies that $(z - m)\left(z - \dfrac{1}{m}\right) = 0 \quad \Leftrightarrow$ $z^2 - \left(m + \dfrac{1}{m}\right) z + m\left(\dfrac{1}{m}\right) = 0$, so equating the constant terms in the two quadratic equations, we get $\dfrac{b^2 - \beta^2}{a^2 - \alpha^2} = m\left(\dfrac{1}{m}\right) = 1$, and hence $b^2 - \beta^2 = a^2 - \alpha^2$. So (α, β) lies on the hyperbola $x^2 - y^2 = a^2 - b^2$.

(b) If the two tangent lines from the point (α, β) to the ellipse have slopes m and $-\dfrac{1}{m}$, then m and $-\dfrac{1}{m}$ are roots of the quadratic equation, and so $(z - m)\left(z + \dfrac{1}{m}\right) = 0$, and equating the constant terms as in part (a), we get $\dfrac{b^2 - \beta^2}{a^2 - \alpha^2} = -1$, and hence $b^2 - \beta^2 = \alpha^2 - a^2$. So the point (α, β) lies on the circle $x^2 + y^2 = a^2 + b^2$.

31. $y = x^4 - 2x^2 - x \quad \Rightarrow \quad y' = 4x^3 - 4x - 1$. The equation of the tangent line at $x = a$ is $y - (a^4 - 2a^2 - a) = (4a^3 - 4a - 1)(x - a)$ or $y = (4a^3 - 4a - 1)x + (-3a^4 + 2a^2)$ and similarly for $x = b$. So if at $x = a$ and $x = b$ we have the same tangent line, then $4a^3 - 4a - 1 = 4b^3 - 4b - 1$ and $-3a^4 + 2a^2 = -3b^4 + 2b^2$. The first equation gives $a^3 - b^3 = a - b \quad \Rightarrow \quad (a - b)(a^2 + ab + b^2) = (a - b)$. Assuming $a \neq b$, we have $1 = a^2 + ab + b^2$. The second equation gives $3(a^4 - b^4) = 2(a^2 - b^2) \quad \Rightarrow \quad 3(a^2 - b^2)(a^2 + b^2) = 2(a^2 - b^2)$ which is true if $a = -b$. Substituting into $1 = a^2 + ab + b^2$ gives $1 = a^2 - a^2 + a^2 \quad \Rightarrow \quad a = \pm 1$ so that $a = 1$ and $b = -1$ or vice versa. Thus, the points $(1, -2)$ and $(-1, 0)$ have a common tangent line.

As long as there are only two such points, we are done. So we show that these are in fact the only two such points. Suppose that $a^2 - b^2 \neq 0$. Then $3(a^2 - b^2)(a^2 + b^2) = 2(a^2 - b^2)$ gives $3(a^2 + b^2) = 2$ or $a^2 + b^2 = \frac{2}{3}$. Thus, $ab = (a^2 + ab + b^2) - (a^2 + b^2) = 1 - \dfrac{2}{3} = \dfrac{1}{3}$, so $b = \dfrac{1}{3a}$. Hence, $a^2 + \dfrac{1}{9a^2} = \dfrac{2}{3}$, so $9a^4 + 1 = 6a^2 \quad \Rightarrow$ $0 = 9a^4 - 6a^2 + 1 = (3a^2 - 1)^2$. So $3a^2 - 1 = 0 \quad \Rightarrow \quad a^2 = \dfrac{1}{3} \quad \Rightarrow \quad b^2 = \dfrac{1}{9a^2} = \dfrac{1}{3} = a^2$, contradicting our assumption that $a^2 \neq b^2$.

32. Suppose that the normal lines at the three points (a_1, a_1^2), (a_2, a_2^2), and (a_3, a_3^2) intersect at a common point. Now if one of the a_i is 0 (suppose $a_1 = 0$) then by symmetry $a_2 = -a_3$, so $a_1 + a_2 + a_3 = 0$. So we can assume that none of the a_i is 0.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The slope of the tangent line at $\left(a_i, a_i^2\right)$ is $2a_i$, so the slope of the normal line is $-\dfrac{1}{2a_i}$ and its equation is $y - a_i^2 = -\dfrac{1}{2a_i}(x - a_i)$. We solve for the x-coordinate of the intersection of the normal lines from $\left(a_1, a_1^2\right)$ and $\left(a_2, a_2^2\right)$:

$$y = a_1^2 - \frac{1}{2a_1}(x - a_1) = a_2^2 - \frac{1}{2a_2}(x - a_2) \quad \Rightarrow \quad x\left(\frac{1}{2a_2} - \frac{1}{2a_1}\right) = a_2^2 - a_1^2 \quad \Rightarrow$$

$x\left(\dfrac{a_1 - a_2}{2a_1a_2}\right) = (-a_1 - a_2)(a_1 + a_2) \quad \Leftrightarrow \quad x = -2a_1a_2(a_1 + a_2)$ **(1)**. Similarly, solving for the x-coordinate of the intersections of the normal lines from $\left(a_1, a_1^2\right)$ and $\left(a_3, a_3^2\right)$ gives $x = -2a_1a_3(a_1 + a_3)$ **(2)**.

Equating **(1)** and **(2)** gives $a_2(a_1 + a_2) = a_3(a_1 + a_3) \quad \Leftrightarrow \quad a_1(a_2 - a_3) = a_3^2 - a_2^2 = -(a_2 + a_3)(a_2 - a_3) \quad \Leftrightarrow$ $a_1 = -(a_2 + a_3) \quad \Leftrightarrow \quad a_1 + a_2 + a_3 = 0$.

33. Because of the periodic nature of the lattice points, it suffices to consider the points in the 5×2 grid shown. We can see that the minimum value of r occurs when there is a line with slope $\frac{2}{5}$ which touches the circle centered at $(3, 1)$ and the circles centered at $(0, 0)$ and $(5, 2)$.

To find P, the point at which the line is tangent to the circle at $(0, 0)$, we simultaneously solve $x^2 + y^2 = r^2$ and $y = -\frac{5}{2}x \quad \Rightarrow \quad x^2 + \frac{25}{4}x^2 = r^2 \quad \Rightarrow \quad x^2 = \frac{4}{29}r^2 \quad \Rightarrow \quad x = \frac{2}{\sqrt{29}}r, y = -\frac{5}{\sqrt{29}}r$. To find Q, we either use symmetry or solve $(x - 3)^2 + (y - 1)^2 = r^2$ and $y - 1 = -\frac{5}{2}(x - 3)$. As above, we get $x = 3 - \frac{2}{\sqrt{29}}r, y = 1 + \frac{5}{\sqrt{29}}r$. Now the slope of the line PQ is $\frac{2}{5}$, so $m_{PQ} = \dfrac{1 + \frac{5}{\sqrt{29}}r - \left(-\frac{5}{\sqrt{29}}r\right)}{3 - \frac{2}{\sqrt{29}}r - \frac{2}{\sqrt{29}}r} = \dfrac{1 + \frac{10}{\sqrt{29}}r}{3 - \frac{4}{\sqrt{29}}r} = \dfrac{\sqrt{29} + 10r}{3\sqrt{29} - 4r} = \dfrac{2}{5} \quad \Rightarrow$

$5\sqrt{29} + 50r = 6\sqrt{29} - 8r \quad \Leftrightarrow \quad 58r = \sqrt{29} \quad \Leftrightarrow \quad r = \frac{\sqrt{29}}{58}$. So the minimum value of r for which any line with slope $\frac{2}{5}$ intersects circles with radius r centered at the lattice points on the plane is $r = \frac{\sqrt{29}}{58} \approx 0.093$.

34.

Assume the axes of the cone and the cylinder are parallel. Let H denote the initial height of the water. When the cone has been dropping for t seconds, the water level has risen x centimeters, so the tip of the cone is $x + 1t$ centimeters below the water line. We want to find dx/dt when $x + t = h$ (when the cone is completely submerged). Using similar triangles, $\dfrac{r_1}{x + t} = \dfrac{r}{h} \quad \Rightarrow \quad r_1 = \dfrac{r}{h}(x + t)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\begin{array}{ccccc}
\text{volume of water and cone at time } t & = & \text{original volume of water} & + & \text{volume of submerged part of cone} \\
\pi R^2(H+x) & = & \pi R^2 H & + & \frac{1}{3}\pi r_1^2(x+t) \\
\pi R^2 H + \pi R^2 x & = & \pi R^2 H & + & \frac{1}{3}\pi \dfrac{r^2}{h^2}(x+t)^3 \\
3h^2R^2x & = & r^2(x+t)^3 & &
\end{array}$$

Differentiating implicitly with respect to t gives us $3h^2R^2\,\dfrac{dx}{dt} = r^2\left[3(x+t)^2\,\dfrac{dx}{dt} + 3(x+t)^2\,\dfrac{dt}{dt}\right] \quad \Rightarrow$

$\dfrac{dx}{dt} = \dfrac{r^2(x+t)^2}{h^2R^2 - r^2(x+t)^2} \quad \Rightarrow \quad \left.\dfrac{dx}{dt}\right|_{x+t=h} = \dfrac{r^2h^2}{h^2R^2 - r^2h^2} = \dfrac{r^2}{R^2 - r^2}$. Thus, the water level is rising at a rate of $\dfrac{r^2}{R^2 - r^2}$ cm/s at the instant the cone is completely submerged.

35.

By similar triangles, $\dfrac{r}{5} = \dfrac{h}{16} \quad \Rightarrow \quad r = \dfrac{5h}{16}$. The volume of the cone is $V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi\left(\dfrac{5h}{16}\right)^2 h = \dfrac{25\pi}{768}h^3$, so $\dfrac{dV}{dt} = \dfrac{25\pi}{256}h^2\,\dfrac{dh}{dt}$. Now the rate of change of the volume is also equal to the difference of what is being added (2 cm^3/min) and what is oozing out ($k\pi rl$, where πrl is the area of the cone and k is a proportionality constant). Thus, $\dfrac{dV}{dt} = 2 - k\pi rl$.

Equating the two expressions for $\dfrac{dV}{dt}$ and substituting $h = 10$, $\dfrac{dh}{dt} = -0.3$, $r = \dfrac{5(10)}{16} = \dfrac{25}{8}$, and $\dfrac{l}{\sqrt{281}} = \dfrac{10}{16} \quad \Leftrightarrow$

$l = \dfrac{5}{8}\sqrt{281}$, we get $\dfrac{25\pi}{256}(10)^2(-0.3) = 2 - k\pi\dfrac{25}{8}\cdot\dfrac{5}{8}\sqrt{281} \quad \Leftrightarrow \quad \dfrac{125k\pi\sqrt{281}}{64} = 2 + \dfrac{750\pi}{256}$. Solving for k gives us $k = \dfrac{256 + 375\pi}{250\pi\sqrt{281}}$. To maintain a certain height, the rate of oozing, $k\pi rl$, must equal the rate of the liquid being poured in; that is, $\dfrac{dV}{dt} = 0$. Thus, the rate at which we should pour the liquid into the container is

$$k\pi rl = \frac{256 + 375\pi}{250\pi\sqrt{281}}\cdot\pi\cdot\frac{25}{8}\cdot\frac{5\sqrt{281}}{8} = \frac{256 + 375\pi}{128} \approx 11.204 \text{ cm}^3/\text{min}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4 □ APPLICATIONS OF DIFFERENTIATION

4.1 Maximum and Minimum Values

1. A function f has an **absolute minimum** at $x = c$ if $f(c)$ is the smallest function value on the entire domain of f, whereas f has a **local minimum** at c if $f(c)$ is the smallest function value when x is near c.

2. (a) The Extreme Value Theorem

(b) See the Closed Interval Method.

3. Absolute maximum at s, absolute minimum at r, local maximum at c, local minima at b and r, neither a maximum nor a minimum at a and d.

4. Absolute maximum at r; absolute minimum at a; local maxima at b and r; local minimum at d; neither a maximum nor a minimum at c and s.

5. Absolute maximum value is $f(4) = 5$; there is no absolute minimum value; local maximum values are $f(4) = 5$ and $f(6) = 4$; local minimum values are $f(2) = 2$ and $f(1) = f(5) = 3$.

6. There is no absolute maximum value; absolute minimum value is $g(4) = 1$; local maximum values are $g(3) = 4$ and $g(6) = 3$; local minimum values are $g(2) = 2$ and $g(4) = 1$.

7. Absolute maximum at 5, absolute minimum at 2, local maximum at 3, local minima at 2 and 4

8. Absolute maximum at 4, absolute minimum at 5, local maximum at 2, local minimum at 3

9. Absolute minimum at 3, absolute maximum at 4, local maximum at 2

10. Absolute maximum at 2, absolute minimum at 5, 4 is a critial number but there is no local maximum or minimum there.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. (a)

(b)

(c)

12. (a) Note that a local maximum cannot occur at an endpoint.

(b)

Note: By the Extreme Value Theorem, f must *not* be continuous.

13. (a) *Note:* By the Extreme Value Theorem, f must *not* be continuous; because if it were, it would attain an absolute minimum.

(b)

14. (a)

(b)

15. $f(x) = 3 - 2x$, $x \geq -1$. Absolute maximum $f(-1) = 5$; no local maximum. No absolute or local minimum.

16. $f(x) = x^2$, $-1 \leq x < 2$. No absolute or local maximum. Absolute and local minimum $f(0) = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $f(x) = 1/x$, $x \geq 1$. Absolute maximum $f(1) = 1$; no local maximum. No absolute or local minimum.

18. $f(x) = 1/x$, $1 < x < 3$. No absolute or local maximum. No absolute or local minimum.

19. $f(x) = \sin x$, $0 \leq x < \pi/2$. No absolute or local maximum. Absolute minimum $f(0) = 0$; no local minimum.

20. $f(x) = \sin x$, $0 < x \leq \pi/2$. Absolute maximum $f\left(\frac{\pi}{2}\right) = 1$; no local maximum. No absolute or local minimum.

21. $f(x) = \sin x$, $-\pi/2 \leq x \leq \pi/2$. Absolute maximum $f\left(\frac{\pi}{2}\right) = 1$; no local maximum. Absolute minimum $f\left(-\frac{\pi}{2}\right) = -1$; no local minimum.

22. $f(t) = \cos t$, $-\frac{3\pi}{2} \leq t \leq \frac{3\pi}{2}$. Absolute and local maximum $f(0) = 1$; absolute and local minima $f(\pm\pi, -1)$.

23. $f(x) = \ln x$, $0 < x \leq 2$. Absolute maximum $f(2) = \ln 2 \approx 0.69$; no local maximum. No absolute or local minimum.

24. $f(x) = |\ x\ |$. No absolute or local maximum. Absolute and local minimum $f(0) = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $f(x) = 1 - \sqrt{x}$. Absolute maximum $f(0) = 1$; no local maximum. No absolute or local minimum.

26. $f(x) = e^x$. No absolute or local maximum or minimum value.

27. $f(x) = \begin{cases} x^2 & \text{if } -1 \le x \le 0 \\ 2 - 3x & \text{if } 0 < x \le 1 \end{cases}$

No absolute or local maximum.
Absolute minimum $f(1) = -1$.
Local minimum $f(0) = 0$.

28. $f(x) = \begin{cases} 2x + 1 & \text{if } 0 \le x < 1 \\ 4 - 2x & \text{if } 1 \le x \le 3 \end{cases}$

No absolute or local maximum.
Absolute minimum $f(3) = -2$; no local minimum.

29. $f(x) = 3x^2 + x - 2 \quad \Rightarrow \quad f'(x) = 6x + 1$. $f'(x) = 0 \quad \Rightarrow \quad x = -\frac{1}{6}$. This is the only critical number.

30. $g(v) = v^3 - 12v + 4 \quad \Rightarrow \quad g'(v) = 3v^2 - 12 = 3(v^2 - 4) = 3(v + 2)(v - 2)$. $g'(v) = 0 \quad \Rightarrow \quad v = -2, 2$. These are the only critical numbers.

31. $f(x) = 3x^4 + 8x^3 - 48x^2 \quad \Rightarrow \quad f'(x) = 12x^3 + 24x^2 - 96x = 12x(x^2 + 2x - 8) = 12x(x + 4)(x - 2)$.
$f'(x) = 0 \quad \Rightarrow \quad x = -4, 0, 2$. These are the only critical numbers.

32. $f(x) = 2x^3 + x^2 + 8x \quad \Rightarrow \quad f'(x) = 6x^2 + 2x + 8 = 2(3x^2 + x + 4)$. Using the quadratic formula, $f'(x) = 0 \quad \Leftrightarrow$ $x = \dfrac{-1 \pm \sqrt{-47}}{6}$. Since the discriminant, -47, is negative, there are no real solutions, and hence, there are no critical numbers.

33. $g(t) = t^5 + 5t^3 + 50t \quad \Rightarrow \quad g'(t) = 5t^4 + 15t^2 + 50 = 5(t^4 + 3t^2 + 10)$. Using the quadratic formula to solve for t^2, $g'(t) = 0 \quad \Leftrightarrow \quad t^2 = \dfrac{-3 \pm \sqrt{3^2 - 4(1)(10)}}{2(1)} = \dfrac{-3 \pm \sqrt{-31}}{2}$. Since the discriminant, -31, is negative, there are no real solutions, and hence, there are no critical numbers.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. $A(x) = |3-2x| = \begin{cases} 3-2x & \text{if } 3-2x \geq 0 \\ -(3-2x) & \text{if } 3-2x < 0 \end{cases} = \begin{cases} 3-2x & \text{if } x \leq \frac{3}{2} \\ 2x-3 & \text{if } x > \frac{3}{2} \end{cases}$

$A'(x) = \begin{cases} -2 & \text{if } x < \frac{3}{2} \\ 2 & \text{if } x > \frac{3}{2} \end{cases}$ and $A'(x)$ does not exist at $x = \dfrac{3}{2}$, so $x = \dfrac{3}{2}$ is a critical number.

35. $g(y) = \dfrac{y-1}{y^2-y+1} \quad \Rightarrow$

$$g'(y) = \frac{(y^2-y+1)(1)-(y-1)(2y-1)}{(y^2-y+1)^2} = \frac{y^2-y+1-(2y^2-3y+1)}{(y^2-y+1)^2} = \frac{-y^2+2y}{(y^2-y+1)^2} = \frac{y(2-y)}{(y^2-y+1)^2}.$$

$g'(y) = 0 \quad \Rightarrow \quad y = 0, 2$. The expression y^2-y+1 is never equal to 0, so $g'(y)$ exists for all real numbers.

The critical numbers are 0 and 2.

36. $h(p) = \dfrac{p-1}{p^2+4} \quad \Rightarrow \quad h'(p) = \dfrac{(p^2+4)(1)-(p-1)(2p)}{(p^2+4)^2} = \dfrac{p^2+4-2p^2+2p}{(p^2+4)^2} = \dfrac{-p^2+2p+4}{(p^2+4)^2}.$

$h'(p) = 0 \quad \Rightarrow \quad p = \dfrac{-2 \pm \sqrt{4+16}}{-2} = 1 \pm \sqrt{5}$. The critical numbers are $1 \pm \sqrt{5}$. [$h'(p)$ exists for all real numbers.]

37. $p(x) = \dfrac{x^2+2}{2x-1} \quad \Rightarrow$

$$p'(x) = \frac{(2x-1)(2x)-(x^2+2)(2)}{(2x-1)^2} = \frac{4x^2-2x-2x^2-4}{(2x-1)^2} = \frac{2x^2-2x-4}{(2x-1)^2} = \frac{2(x^2-x-2)}{(2x-1)^2} = \frac{2(x-2)(x+1)}{(2x-1)^2}.$$

$p'(x) = 0 \quad \Rightarrow \quad x = -1$ or 2. $p'(x)$ does not exist at $x = \frac{1}{2}$, but $\frac{1}{2}$ is not in the domain of p, so the critical numbers are -1 and 2.

38. $q(t) = \dfrac{t^2+9}{t^2-9} \quad \Rightarrow \quad q'(t) = \dfrac{(t^2-9)(2t)-(t^2+9)(2t)}{(t^2-9)^2} = \dfrac{2t^3-18t-2t^3-18t}{(t^2-9)^2} = -\dfrac{36t}{(t^2-9)^2}$. $q'(t) = 0 \quad \Rightarrow$

$36t = 0 \quad \Rightarrow \quad t = 0$. $q'(t)$ does not exist at $t = \pm 3$, but 3 and -3 are not in the domain of q, so 0 is the only critical number.

39. $h(t) = t^{3/4} - 2t^{1/4} \quad \Rightarrow \quad h'(t) = \frac{3}{4}t^{-1/4} - \frac{2}{4}t^{-3/4} = \frac{1}{4}t^{-3/4}(3t^{1/2}-2) = \dfrac{3\sqrt{t}-2}{4\sqrt[4]{t^3}}.$

$h'(t) = 0 \quad \Rightarrow \quad 3\sqrt{t} = 2 \quad \Rightarrow \quad \sqrt{t} = \frac{2}{3} \quad \Rightarrow \quad t = \frac{4}{9}$. $h'(t)$ does not exist at $t = 0$, so the critical numbers are 0 and $\frac{4}{9}$.

40. $g(x) = \sqrt[3]{4-x^2} = (4-x^2)^{1/3} \quad \Rightarrow \quad g'(x) = \frac{1}{3}(4-x^2)^{-2/3}(-2x) = \dfrac{-2x}{3(4-x^2)^{2/3}}$. $g'(x) = 0 \quad \Rightarrow \quad x = 0$.

$g'(\pm 2)$ do not exist. Thus, the three critical numbers are -2, 0, and 2.

41. $F(x) = x^{4/5}(x-4)^2 \quad \Rightarrow$

$$F'(x) = x^{4/5} \cdot 2(x-4) + (x-4)^2 \cdot \tfrac{4}{5}x^{-1/5} = \tfrac{1}{5}x^{-1/5}(x-4)[5 \cdot x \cdot 2 + (x-4) \cdot 4]$$
$$= \frac{(x-4)(14x-16)}{5x^{1/5}} = \frac{2(x-4)(7x-8)}{5x^{1/5}}$$

$F'(x) = 0 \quad \Rightarrow \quad x = 4, \frac{8}{7}$. $F'(0)$ does not exist. Thus, the three critical numbers are 0, $\frac{8}{7}$, and 4.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. $h(x) = x^{-1/3}(x-2) \quad \Rightarrow$

$h'(x) = x^{-1/3} \cdot 1 + (x-2) \cdot \left(-\frac{1}{3}x^{-4/3}\right) = \frac{1}{3}x^{-4/3}\,[3x + (x-2)(-1)] = \dfrac{2x+2}{3x^{4/3}} = \dfrac{2(x+1)}{3x^{4/3}}$. $h'(x) = 0 \quad \Rightarrow$

$2(x+1) = 0 \quad \Rightarrow \quad x = -1$. $h'(x)$ does not exist at $x = 0$, but 0 is not in the domain of h, so -1 is the only critical number.

43. $f(x) = x^{1/3}(4-x)^{2/3} \quad \Rightarrow$

$f'(x) = x^{1/3} \cdot \frac{2}{3}\,(4-x)^{-1/3} \cdot (-1) + (4-x)^{2/3} \cdot \frac{1}{3}x^{-2/3} = \frac{1}{3}x^{-2/3}(4-x)^{-1/3}[-2x + (4-x)] = \dfrac{4-3x}{3x^{2/3}(4-x)^{1/3}}$.

$f'(x) = 0 \quad \Rightarrow \quad 4 - 3x = 0 \quad \Rightarrow \quad x = \frac{4}{3}$. $f'(0)$ and $f'(4)$ are undefined. Thus, the three critical numbers are 0, $\frac{4}{3}$, and 4.

44. $f(\theta) = \theta + \sqrt{2}\cos\theta \quad \Rightarrow \quad f'(\theta) = 1 - \sqrt{2}\sin\theta$. $f'(0) = 0 \quad \Rightarrow \quad 1 - \sqrt{2}\sin\theta = 0 \quad \Rightarrow \quad \sin\theta = \dfrac{1}{\sqrt{2}} \quad \Rightarrow$

$\theta = \frac{\pi}{4} + 2n\pi$ [n an integer], $\theta = \frac{3\pi}{4} + 2n\pi$ are critical numbers. [$f'(\theta)$ exists for all real numbers.]

45. $f(\theta) = 2\cos\theta + \sin^2\theta \quad \Rightarrow \quad f'(\theta) = -2\sin\theta + 2\sin\theta\,\cos\theta$. $f'(\theta) = 0 \quad \Rightarrow \quad 2\sin\theta\,(\cos\theta - 1) = 0 \quad \Rightarrow \quad \sin\theta = 0$ or $\cos\theta = 1 \quad \Rightarrow \quad \theta = n\pi$ [n an integer] or $\theta = 2n\pi$. The solutions $\theta = n\pi$ include the solutions $\theta = 2n\pi$, so the critical numbers are $\theta = n\pi$.

46. $p(t) = te^{4t} \quad \Rightarrow \quad p'(t) = t(4e^{4t}) + e^{4t}(1) = e^{4t}(4t+1)$. $p'(t) = 0 \quad \Rightarrow \quad t = -\frac{1}{4}$ [e^{4t} is never equal to 0]. $p'(t)$ always exists, so the only critical number is $-\frac{1}{4}$.

47. $g(x) = x^2 \ln x \quad \Rightarrow \quad g'(x) = x^2(1/x) + (\ln x)(2x) = x + 2x\ln x = x(1 + 2\ln x)$. $g'(x) = 0 \quad \Rightarrow \quad 1 + 2\ln x = 0$ [or $x = 0$, but 0 is not in the domain of g] $\quad \Rightarrow \quad \ln x = -\frac{1}{2} \quad \Rightarrow \quad x = e^{-1/2}$. $g'(0)$ does not exist, but 0 is not in the domain of g, so the only critical number is $e^{-1/2} = 1/\sqrt{e}$.

48. $B(u) = 4\tan^{-1} u - u \quad \Rightarrow \quad B'(u) = \dfrac{4}{1+u^2} - 1 = \dfrac{4}{1+u^2} - \dfrac{1+u^2}{1+u^2} = \dfrac{3-u^2}{1+u^2}$. $B'(u) = 0 \quad \Rightarrow \quad 3 - u^2 = 0 \quad \Rightarrow$

$u = -\sqrt{3}, \sqrt{3}$. The expression $1 + u^2$ is never equal to 0, so $B'(u)$ exists for all real numbers. Thus, the critical numbers are $-\sqrt{3}$ and $\sqrt{3}$.

49. The graph of $f'(x) = 5e^{-0.1|x|}\sin x - 1$ has 10 zeros and exists everywhere, so f has 10 critical numbers.

50. A graph of $f'(x) = \dfrac{100\cos^2 x}{10 + x^2} - 1$ is shown. There are 7 zeros between 0 and 10, and 7 more zeros since f' is an even function. f' exists everywhere, so f has 14 critical numbers.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. $f(x) = 12 + 4x - x^2,\ [0, 5]$. $f'(x) = 4 - 2x = 0 \quad \Leftrightarrow \quad x = 2$. $f(0) = 12$, $f(2) = 16$, and $f(5) = 7$. So $f(2) = 16$ is the absolute maximum value and $f(5) = 7$ is the absolute minimum value.

52. $f(x) = 5 + 54x - 2x^3,\ [0, 4]$. $f'(x) = 54 - 6x^2 = 6(9 - x^2) = 6(3 + x)(3 - x) = 0 \quad \Leftrightarrow \quad x = -3, 3$. $f(0) = 5$, $f(3) = 113$, and $f(4) = 93$. So $f(3) = 113$ is the absolute maximum value and $f(0) = 5$ is the absolute minimum value.

53. $f(x) = 2x^3 - 3x^2 - 12x + 1,\ [-2, 3]$. $f'(x) = 6x^2 - 6x - 12 = 6(x^2 - x - 2) = 6(x - 2)(x + 1) = 0 \quad \Leftrightarrow$ $x = 2, -1$. $f(-2) = -3$, $f(-1) = 8$, $f(2) = -19$, and $f(3) = -8$. So $f(-1) = 8$ is the absolute maximum value and $f(2) = -19$ is the absolute minimum value.

54. $f(x) = x^3 - 6x^2 + 5,\ [-3, 5]$. $f'(x) = 3x^2 - 12x = 3x(x - 4) = 0 \quad \Leftrightarrow \quad x = 0, 4$. $f(-3) = -76$, $f(0) = 5$, $f(4) = -27$, and $f(5) = -20$. So $f(0) = 5$ is the absolute maximum value and $f(-3) = -76$ is the absolute minimum value.

55. $f(x) = 3x^4 - 4x^3 - 12x^2 + 1,\ [-2, 3]$. $f'(x) = 12x^3 - 12x^2 - 24x = 12x(x^2 - x - 2) = 12x(x + 1)(x - 2) = 0 \quad \Leftrightarrow$ $x = -1, 0, 2$. $f(-2) = 33$, $f(-1) = -4$, $f(0) = 1$, $f(2) = -31$, and $f(3) = 28$. So $f(-2) = 33$ is the absolute maximum value and $f(2) = -31$ is the absolute minimum value.

56. $f(t) = (t^2 - 4)^3,\ [-2, 3]$. $f'(t) = 3(t^2 - 4)^2(2t) = 6t(t + 2)^2(t - 2)^2 = 0 \quad \Leftrightarrow \quad t = -2, 0, 2$. $f(\pm 2) = 0$, $f(0) = -64$, and $f(3) = 5^3 = 125$. So $f(3) = 125$ is the absolute maximum value and $f(0) = -64$ is the absolute minimum value.

57. $f(x) = x + \dfrac{1}{x},\ [0.2, 4]$. $f'(x) = 1 - \dfrac{1}{x^2} = \dfrac{x^2 - 1}{x^2} = \dfrac{(x + 1)(x - 1)}{x^2} = 0 \quad \Leftrightarrow \quad x = \pm 1$, but $x = -1$ is not in the given interval, $[0.2, 4]$. $f'(x)$ does not exist when $x = 0$, but 0 is not in the given interval, so 1 is the only critical nuumber. $f(0.2) = 5.2$, $f(1) = 2$, and $f(4) = 4.25$. So $f(0.2) = 5.2$ is the absolute maximum value and $f(1) = 2$ is the absolute minimum value.

58. $f(x) = \dfrac{x}{x^2 - x + 1},\ [0, 3]$.

$$f'(x) = \frac{(x^2 - x + 1) - x(2x - 1)}{(x^2 - x + 1)^2} = \frac{x^2 - x + 1 - 2x^2 + x}{(x^2 - x + 1)^2} = \frac{1 - x^2}{(x^2 - x + 1)^2} = \frac{(1 + x)(1 - x)}{(x^2 - x + 1)^2} = 0 \quad \Leftrightarrow$$

$x = \pm 1$, but $x = -1$ is not in the given interval, $[0, 3]$. $f(0) = 0$, $f(1) = 1$, and $f(3) = \frac{3}{7}$. So $f(1) = 1$ is the absolute maximum value and $f(0) = 0$ is the absolute minimum value.

59. $f(t) = t - \sqrt[3]{t},\ [-1, 4]$. $f'(t) = 1 - \frac{1}{3}t^{-2/3} = 1 - \dfrac{1}{3t^{2/3}}$. $f'(t) = 0 \quad \Leftrightarrow \quad 1 = \dfrac{1}{3t^{2/3}} \quad \Leftrightarrow \quad t^{2/3} = \dfrac{1}{3} \quad \Leftrightarrow$

$t = \pm\left(\dfrac{1}{3}\right)^{3/2} = \pm\sqrt{\dfrac{1}{27}} = \pm\dfrac{1}{3\sqrt{3}} = \pm\dfrac{\sqrt{3}}{9}$. $f'(t)$ does not exist when $t = 0$. $f(-1) = 0$, $f(0) = 0$,

$f\left(\dfrac{-1}{3\sqrt{3}}\right) = \dfrac{-1}{3\sqrt{3}} - \dfrac{-1}{\sqrt{3}} = \dfrac{-1 + 3}{3\sqrt{3}} = \dfrac{2\sqrt{3}}{9} \approx 0.3849$, $f\left(\dfrac{1}{3\sqrt{3}}\right) = \dfrac{1}{3\sqrt{3}} - \dfrac{1}{\sqrt{3}} = -\dfrac{2\sqrt{3}}{9}$, and

$f(4) = 4 - \sqrt[3]{4} \approx 2.413$. So $f(4) = 4 - \sqrt[3]{4}$ is the absolute maximum value and $f\left(\dfrac{\sqrt{3}}{9}\right) = -\dfrac{2\sqrt{3}}{9}$ is the absolute minimum value.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. $f(x) = \dfrac{e^x}{1+x^2}$, $[0,3]$. $f'(x) = \dfrac{(1+x^2)e^x - e^x(2x)}{(1+x^2)^2} = \dfrac{e^x(x^2-2x+1)}{(1+x^2)^2} = \dfrac{e^x(x-1)^2}{(1+x^2)^2}$. $f'(x) = 0 \;\Rightarrow$ $(x-1)^2 = 0 \;\Leftrightarrow\; x = 1$. $f'(x)$ exists for all real numbers since $1+x^2$ is never equal to 0. $f(0) = 1$, $f(1) = e/2 \approx 1.359$, and $f(3) = e^3/10 \approx 2.009$. So $f(3) = e^3/10$ is the absolute maximum value and $f(0) = 1$ is the absolute minimum value.

61. $f(t) = 2\cos t + \sin 2t$, $[0, \pi/2]$.

$f'(t) = -2\sin t + \cos 2t \cdot 2 = -2\sin t + 2(1 - 2\sin^2 t) = -2(2\sin^2 t + \sin t - 1) = -2(2\sin t - 1)(\sin t + 1)$.

$f'(t) = 0 \;\Rightarrow\; \sin t = \frac{1}{2}$ or $\sin t = -1 \;\Rightarrow\; t = \frac{\pi}{6}$. $f(0) = 2$, $f(\frac{\pi}{6}) = \sqrt{3} + \frac{1}{2}\sqrt{3} = \frac{3}{2}\sqrt{3} \approx 2.60$, and $f(\frac{\pi}{2}) = 0$. So $f(\frac{\pi}{6}) = \frac{3}{2}\sqrt{3}$ is the absolute maximum value and $f(\frac{\pi}{2}) = 0$ is the absolute minimum value.

62. $f(\theta) = 1 + \cos^2\theta$, $[\pi/4, \pi]$. $f'(\theta) = 2\cos\theta(-\sin\theta) = -2\sin\theta\,\cos\theta = -\sin 2\theta$. $f'(\theta) = 0 \;\Rightarrow\; -\sin 2\theta = 0 \;\Rightarrow$ $2\theta = n\pi \;\Rightarrow\; \theta = \frac{n\pi}{2}$. Only $\theta = \frac{\pi}{2}$ $[n = 1]$ is in the interval $(\pi/4, \pi)$. $f(\pi/4) = 1 + \left(\dfrac{\sqrt{2}}{2}\right)^2 = \dfrac{3}{2}$, $f(\pi/2) = 1 + 0^2 = 1$, and $f(\pi) = 1 + (-1)^2 = 2$. So $f(\pi) = 2$ is the absolute maximum value and $f(\pi/2) = 1$ is the absolute minimum value.

63. $f(x) = x^{-2}\ln x$, $\left[\frac{1}{2}, 4\right]$. $f'(x) = x^{-2} \cdot \dfrac{1}{x} + (\ln x)(-2x^{-3}) = x^{-3} - 2x^{-3}\ln x = x^{-3}(1 - 2\ln x) = \dfrac{1 - 2\ln x}{x^3}$.

$f'(x) = 0 \;\Leftrightarrow\; 1 - 2\ln x = 0 \;\Leftrightarrow\; 2\ln x = 1 \;\Leftrightarrow\; \ln x = \frac{1}{2} \;\Leftrightarrow\; x = e^{1/2} \approx 1.65$. $f'(x)$ does not exist when $x = 0$, which is not in the given interval, $\left[\frac{1}{2}, 4\right]$. $f\left(\frac{1}{2}\right) = \dfrac{\ln 1/2}{(1/2)^2} = \dfrac{\ln 1 - \ln 2}{1/4} = -4\ln 2 \approx -2.773$,

$f(e^{1/2}) = \dfrac{\ln e^{1/2}}{(e^{1/2})^2} = \dfrac{1/2}{e} = \dfrac{1}{2e} \approx 0.184$, and $f(4) = \dfrac{\ln 4}{4^2} = \dfrac{\ln 4}{16} \approx 0.087$. So $f(e^{1/2}) = \dfrac{1}{2e}$ is the absolute maximum value and $f\left(\frac{1}{2}\right) = -4\ln 2$ is the absolute minimum value.

64. $f(x) = xe^{x/2}$, $[-3, 1]$. $f'(x) = xe^{x/2}\left(\frac{1}{2}\right) + e^{x/2}(1) = e^{x/2}\left(\frac{1}{2}x + 1\right)$. $f'(x) = 0 \;\Leftrightarrow\; \frac{1}{2}x + 1 = 0 \;\Leftrightarrow\; x = -2$. $f(-3) = -3e^{-3/2} \approx -0.669$, $f(-2) = -2e^{-1} \approx -0.736$, and $f(1) = e^{1/2} \approx 1.649$. So $f(1) = e^{1/2}$ is the absolute maximum value and $f(-2) = -2/e$ is the absolute minimum value.

65. $f(x) = \ln(x^2 + x + 1)$, $[-1, 1]$. $f'(x) = \dfrac{1}{x^2 + x + 1} \cdot (2x + 1) = 0 \;\Leftrightarrow\; x = -\frac{1}{2}$. Since $x^2 + x + 1 > 0$ for all x, the domain of f and f' is $\mathbb{R}$. $f(-1) = \ln 1 = 0$, $f\left(-\frac{1}{2}\right) = \ln\frac{3}{4} \approx -0.29$, and $f(1) = \ln 3 \approx 1.10$. So $f(1) = \ln 3 \approx 1.10$ is the absolute maximum value and $f\left(-\frac{1}{2}\right) = \ln\frac{3}{4} \approx -0.29$ is the absolute minimum value.

66. $f(x) = x - 2\tan^{-1} x$, $[0, 4]$. $f'(x) = 1 - 2 \cdot \dfrac{1}{1+x^2} = 0 \;\Leftrightarrow\; 1 = \dfrac{2}{1+x^2} \;\Leftrightarrow\; 1 + x^2 = 2 \;\Leftrightarrow\; x^2 = 1 \;\Leftrightarrow$ $x = \pm 1$. $f(0) = 0$, $f(1) = 1 - \frac{\pi}{2} \approx -0.57$, and $f(4) = 4 - 2\tan^{-1} 4 \approx 1.35$. So $f(4) = 4 - 2\tan^{-1} 4$ is the absolute maximum value and $f(1) = 1 - \frac{\pi}{2}$ is the absolute minimum value.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. $f(x) = x^a(1-x)^b, \quad 0 \le x \le 1, a > 0, b > 0.$

$$f'(x) = x^a \cdot b(1-x)^{b-1}(-1) + (1-x)^b \cdot ax^{a-1} = x^{a-1}(1-x)^{b-1}[x \cdot b(-1) + (1-x) \cdot a]$$
$$= x^{a-1}(1-x)^{b-1}(a - ax - bx)$$

At the endpoints, we have $f(0) = f(1) = 0$ [the minimum value of f]. In the interval $(0, 1)$, $f'(x) = 0 \quad \Leftrightarrow \quad x = \dfrac{a}{a+b}$.

$$f\left(\frac{a}{a+b}\right) = \left(\frac{a}{a+b}\right)^a \left(1 - \frac{a}{a+b}\right)^b = \frac{a^a}{(a+b)^a}\left(\frac{a+b-a}{a+b}\right)^b = \frac{a^a}{(a+b)^a} \cdot \frac{b^b}{(a+b)^b} = \frac{a^a b^b}{(a+b)^{a+b}}.$$

So $f\left(\dfrac{a}{a+b}\right) = \dfrac{a^a b^b}{(a+b)^{a+b}}$ is the absolute maximum value.

68.

The graph of $f(x) = \left|1 + 5x - x^3\right|$ indicates that $f'(x) = 0$ at $x \approx \pm 1.3$ and that $f'(x)$ does not exist at $x \approx -2.1$, -0.2, and 2.3. Those five values of x are the critical numbers of f.

69. (a)

From the graph, it appears that the absolute maximum value is about $f(-0.77) = 2.19$, and the absolute minimum value is about $f(0.77) = 1.81$.

(b) $f(x) = x^5 - x^3 + 2 \quad \Rightarrow \quad f'(x) = 5x^4 - 3x^2 = x^2(5x^2 - 3)$. So $f'(x) = 0 \quad \Rightarrow \quad x = 0, \pm\sqrt{\frac{3}{5}}$.

$$f\left(-\sqrt{\tfrac{3}{5}}\right) = \left(-\sqrt{\tfrac{3}{5}}\right)^5 - \left(-\sqrt{\tfrac{3}{5}}\right)^3 + 2 = -\left(\tfrac{3}{5}\right)^2\sqrt{\tfrac{3}{5}} + \tfrac{3}{5}\sqrt{\tfrac{3}{5}} + 2$$
$$= \left(\tfrac{3}{5} - \tfrac{9}{25}\right)\sqrt{\tfrac{3}{5}} + 2 = \tfrac{6}{25}\sqrt{\tfrac{3}{5}} + 2 \quad \text{(maximum)}$$

and similarly, $f\left(\sqrt{\frac{3}{5}}\right) = -\frac{6}{25}\sqrt{\frac{3}{5}} + 2$ (minimum).

70. (a)

From the graph, it appears that the absolute maximum value is about $f(1) = 2.85$, and the absolute minimum value is about $f(0.23) = 1.89$.

(b) $f(x) = e^x + e^{-2x} \quad \Rightarrow \quad f'(x) = e^x - 2e^{-2x} = e^{-2x}(e^{3x} - 2)$. So $f'(x) = 0 \quad \Leftrightarrow \quad e^{3x} = 2 \quad \Leftrightarrow \quad 3x = \ln 2 \quad \Leftrightarrow$ $x = \frac{1}{3}\ln 2$ [≈ 0.23]. $f\left(\frac{1}{3}\ln 2\right) = (e^{\ln 2})^{1/3} + (e^{\ln 2})^{-2/3} = 2^{1/3} + 2^{-2/3}$ [≈ 1.89], the minimum value. $f(1) = e^1 + e^{-2}$ [≈ 2.85], the maximum.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

71. (a)

From the graph, it appears that the absolute maximum value is about $f(0.75) = 0.32$, and the absolute minimum value is $f(0) = f(1) = 0$; that is, at both endpoints.

(b) $f(x) = x\sqrt{x - x^2} \quad \Rightarrow \quad f'(x) = x \cdot \dfrac{1 - 2x}{2\sqrt{x - x^2}} + \sqrt{x - x^2} = \dfrac{(x - 2x^2) + (2x - 2x^2)}{2\sqrt{x - x^2}} = \dfrac{3x - 4x^2}{2\sqrt{x - x^2}}$.

So $f'(x) = 0 \quad \Rightarrow \quad 3x - 4x^2 = 0 \quad \Rightarrow \quad x(3 - 4x) = 0 \quad \Rightarrow \quad x = 0$ or $\frac{3}{4}$.

$f(0) = f(1) = 0$ (minimum), and $f\left(\frac{3}{4}\right) = \frac{3}{4}\sqrt{\frac{3}{4} - \left(\frac{3}{4}\right)^2} = \frac{3}{4}\sqrt{\frac{3}{16}} = \frac{3\sqrt{3}}{16}$ (maximum).

72. (a)

From the graph, it appears that the absolute maximum value is about $f(-2) = -1.17$, and the absolute minimum value is about $f(-0.52) = -2.26$.

(b) $f(x) = x - 2\cos x \quad \Rightarrow \quad f'(x) = 1 + 2\sin x$. So $f'(x) = 0 \quad \Rightarrow \quad \sin x = -\frac{1}{2} \quad \Rightarrow \quad x = -\frac{\pi}{6}$ on $[-2, 0]$.

$f(-2) = -2 - 2\cos(-2)$ (maximum) and $f\left(-\frac{\pi}{6}\right) = -\frac{\pi}{6} - 2\cos\left(-\frac{\pi}{6}\right) = -\frac{\pi}{6} - 2\left(\frac{\sqrt{3}}{2}\right) = -\frac{\pi}{6} - \sqrt{3}$ (minimum).

73. Let $a = 0.135$ and $b = -2.802$. Then $C(t) = ate^{bt} \quad \Rightarrow \quad C'(t) = a(t \cdot e^{bt} \cdot b + e^{bt} \cdot 1) = ae^{bt}(bt + 1)$. $\quad C'(t) = 0 \quad \Leftrightarrow$

$bt + 1 = 0 \quad \Leftrightarrow \quad t = -\dfrac{1}{b} \approx 0.36$ h. $\quad C(0) = 0$, $C(-1/b) = -\dfrac{a}{b}e^{-1} = -\dfrac{a}{be} \approx 0.0177$, and $C(3) = 3ae^{3b} \approx 0.00009$.

The maximum average BAC during the first three hours is about 0.0177 g/dL and it occurs at approximately 0.36 h (21.4 min).

74. $C(t) = 8(e^{-0.4t} - e^{-0.6t}) \quad \Rightarrow \quad C'(t) = 8(-0.4e^{-0.4t} + 0.6e^{-0.6t})$. $\quad C'(t) = 0 \quad \Leftrightarrow \quad 0.6e^{-0.6t} = 0.4e^{-0.4t} \quad \Leftrightarrow$

$\frac{0.6}{0.4} = e^{-0.4t+0.6t} \quad \Leftrightarrow \quad \frac{3}{2} = e^{0.2t} \quad \Leftrightarrow \quad 0.2t = \ln\frac{3}{2} \quad \Leftrightarrow \quad t = 5\ln\frac{3}{2} \approx 2.027$ h. $\quad C(0) = 8(1 - 1) = 0$,

$C\left(5\ln\frac{3}{2}\right) = 8(e^{-2\ln 3/2} - e^{-3\ln 3/2}) = 8\left[\left(\frac{3}{2}\right)^{-2} - \left(\frac{3}{2}\right)^{-3}\right] = 8\left(\frac{4}{9} - \frac{8}{27}\right) = \frac{32}{27} \approx 1.185$, and

$C(12) = 8(e^{-4.8} - e^{-7.2}) \approx 0.060$. The maximum concentration of the antibiotic during the first 12 hours is $\frac{32}{27}$ μg/mL.

75. The density is defined as $\rho = \dfrac{\text{mass}}{\text{volume}} = \dfrac{1000}{V(T)}$ (in g/cm^3). But a critical point of ρ will also be a critical point of V

$\left[\text{since } \dfrac{d\rho}{dT} = -1000V^{-2}\dfrac{dV}{dT} \text{ and } V \text{ is never } 0\right]$, and V is easier to differentiate than ρ.

$V(T) = 999.87 - 0.06426T + 0.0085043T^2 - 0.0000679T^3 \quad \Rightarrow \quad V'(T) = -0.06426 + 0.0170086T - 0.0002037T^2$.

Setting this equal to 0 and using the quadratic formula to find T, we get

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$T = \dfrac{-0.0170086 \pm \sqrt{0.0170086^2 - 4 \cdot 0.0002037 \cdot 0.06426}}{2(-0.0002037)} \approx 3.9665°\text{C}$ or $79.5318°\text{C}$. Since we are only interested in the region $0°\text{C} \le T \le 30°\text{C}$, we check the density ρ at the endpoints and at $3.9665°\text{C}$: $\rho(0) \approx \dfrac{1000}{999.87} \approx 1.00013$; $\rho(30) \approx \dfrac{1000}{1003.7628} \approx 0.99625$; $\rho(3.9665) \approx \dfrac{1000}{999.7447} \approx 1.000255$. So water has its maximum density at about $3.9665°\text{C}$.

76. $F = \dfrac{\mu W}{\mu \sin\theta + \cos\theta} \quad \Rightarrow \quad \dfrac{dF}{d\theta} = \dfrac{(\mu\sin\theta + \cos\theta)(0) - \mu W(\mu\cos\theta - \sin\theta)}{(\mu\sin\theta + \cos\theta)^2} = \dfrac{-\mu W(\mu\cos\theta - \sin\theta)}{(\mu\sin\theta + \cos\theta)^2}$.

So $\dfrac{dF}{d\theta} = 0 \quad \Rightarrow \quad \mu\cos\theta - \sin\theta = 0 \quad \Rightarrow \quad \mu = \dfrac{\sin\theta}{\cos\theta} = \tan\theta$. Substituting $\tan\theta$ for μ in F gives us

$F = \dfrac{(\tan\theta)W}{(\tan\theta)\sin\theta + \cos\theta} = \dfrac{W\tan\theta}{\dfrac{\sin^2\theta}{\cos\theta} + \cos\theta} = \dfrac{W\tan\theta\,\cos\theta}{\sin^2\theta + \cos^2\theta} = \dfrac{W\sin\theta}{1} = W\sin\theta.$

If $\tan\theta = \mu$, then $\sin\theta = \dfrac{\mu}{\sqrt{\mu^2+1}}$ (see the figure), so $F = \dfrac{\mu}{\sqrt{\mu^2+1}}W$.

We compare this with the value of F at the endpoints: $F(0) = \mu W$ and $F\left(\frac{\pi}{2}\right) = W$.

Now because $\dfrac{\mu}{\sqrt{\mu^2+1}} \le 1$ and $\dfrac{\mu}{\sqrt{\mu^2+1}} \le \mu$, we have that $\dfrac{\mu}{\sqrt{\mu^2+1}}W$ is less than or equal to each of $F(0)$ and $F\left(\frac{\pi}{2}\right)$.

Hence, $\dfrac{\mu}{\sqrt{\mu^2+1}}W$ is the absolute minimum value of $F(\theta)$, and it occurs when $\tan\theta = \mu$.

77. $L(t) = 0.01441t^3 - 0.4177t^2 + 2.703t + 1060.1 \quad \Rightarrow \quad L'(t) = 0.04323t^2 - 0.8354t + 2.703$. Use the quadratic formula to solve $L'(t) = 0$. $\quad t = \dfrac{0.8354 \pm \sqrt{(0.8354)^2 - 4(0.04323)(2.703)}}{2(0.04323)} \approx 4.1$ or 15.2. For $0 \le t \le 12$, we have $L(0) = 1060.1$, $L(4.1) \approx 1065.2$, and $L(12) \approx 1057.3$. Thus, the water level was highest during 2012 about 4.1 months after January 1.

78. (a) The equation of the graph in the figure is

$v(t) = 0.00146t^3 - 0.11553t^2 + 24.98169t - 21.26872.$

(b) $a(t) = v'(t) = 0.00438t^2 - 0.23106t + 24.98169 \quad \Rightarrow$

$a'(t) = 0.00876t - 0.23106.$

$a'(t) = 0 \quad \Rightarrow \quad t_1 = \frac{0.23106}{0.00876} \approx 26.4$. $a(0) \approx 24.98$, $a(t_1) \approx 21.93$, and $a(125) \approx 64.54$.

The maximum acceleration is about $64.5\ \text{ft/s}^2$ and the minimum acceleration is about $21.93\ \text{ft/s}^2$.

79. (a) $v(r) = k(r_0 - r)r^2 = kr_0r^2 - kr^3 \quad \Rightarrow \quad v'(r) = 2kr_0r - 3kr^2$. $\quad v'(r) = 0 \quad \Rightarrow \quad kr(2r_0 - 3r) = 0 \quad \Rightarrow$ $r = 0$ or $\frac{2}{3}r_0$ (but 0 is not in the interval). Evaluating v at $\frac{1}{2}r_0$, $\frac{2}{3}r_0$, and r_0, we get $v\left(\frac{1}{2}r_0\right) = \frac{1}{8}kr_0^3$, $v\left(\frac{2}{3}r_0\right) = \frac{4}{27}kr_0^3$, and $v(r_0) = 0$. Since $\frac{4}{27} > \frac{1}{8}$, v attains its maximum value at $r = \frac{2}{3}r_0$. This supports the statement in the text.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) From part (a), the maximum value of v is $\frac{4}{27}kr_0^3$.

(c)

80. $f(x) = x^{101} + x^{51} + x + 1 \quad \Rightarrow \quad f'(x) = 101x^{100} + 51x^{50} + 1$. Since $f'(x) \geq 1$ for all x, $f'(x) = 0$ has no solution. Thus, f has no critical number, and the function f has neither a local maximum nor a local minimum.

81. (a) Suppose that f has a local minimum value at c, so $f(x) \geq f(c)$ for all x near c. Then $g(x) = -f(x) \leq -f(c) = g(c)$ for all x near c, so $g(x)$ has a local maximum value at c.

(b) If f has a local minimum at c, then $g(x) = -f(x)$ has a local maximum at c, so $g'(c) = 0$ by the case of Fermat's Theorem proved in the text. Thus, $f'(c) = -g'(c) = 0$.

82. (a) $f(x) = ax^3 + bx^2 + cx + d$, where $a \neq 0$. So $f'(x) = 3ax^2 + 2bx + c$ is a quadratic function and hence, the quadratic equation $f'(x) = 0$ has either 2, 1, or 0 real solutions. Thus, a cubic function can have two, one, or no critical number(s).

Case (i) [2 critical numbers]: $f(x) = x^3 - 3x \quad \Rightarrow$
$f'(x) = 3x^2 - 3 = 3(x^2 - 1)$, so $x = -1, 1$
are critical numbers.

Case (ii) [1 critical number]: $f(x) = x^3 \quad \Rightarrow$
$f'(x) = 3x^2$, so $x = 0$
is the only critical number.

Case (iii) [no critical number]: $f(x) = x^3 + 3x \quad \Rightarrow$
$f'(x) = 3x^2 + 3 = 3(x^2 + 1)$,
so there is no critical number.

(b) Since there are at most two critical numbers, a cubic function can have at most two local extreme values, and by (a)(i), this can occur. By (a)(ii) and (a)(iii), it can have no local extreme value. Thus, a cubic function can have zero or two local extreme values.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

APPLIED PROJECT The Calculus of Rainbows

1. From Snell's Law, we have $\sin\alpha = k\sin\beta \approx \frac{4}{3}\sin\beta \quad\Leftrightarrow\quad \beta \approx \arcsin\left(\frac{3}{4}\sin\alpha\right)$. We substitute this into $D(\alpha) = \pi + 2\alpha - 4\beta = \pi + 2\alpha - 4\arcsin\left(\frac{3}{4}\sin\alpha\right)$, and then differentiate to find the minimum:

$D'(\alpha) = 2 - 4\left[1 - \left(\frac{3}{4}\sin\alpha\right)^2\right]^{-1/2}\left(\frac{3}{4}\cos\alpha\right) = 2 - \dfrac{3\cos\alpha}{\sqrt{1 - \frac{9}{16}\sin^2\alpha}}$. This is 0 when $\dfrac{3\cos\alpha}{\sqrt{1 - \frac{9}{16}\sin^2\alpha}} = 2 \quad\Leftrightarrow$

$\frac{9}{4}\cos^2\alpha = 1 - \frac{9}{16}\sin^2\alpha \quad\Leftrightarrow\quad \frac{9}{4}\cos^2\alpha = 1 - \frac{9}{16}\left(1 - \cos^2\alpha\right) \quad\Leftrightarrow\quad \frac{27}{16}\cos^2\alpha = \frac{7}{16} \quad\Leftrightarrow\quad \cos\alpha = \sqrt{\frac{7}{27}} \quad\Leftrightarrow$

$\alpha = \arccos\sqrt{\frac{7}{27}} \approx 59.4^\circ$, and so the local minimum is $D(59.4^\circ) \approx 2.4$ radians $\approx 138^\circ$.

To see that this is an absolute minimum, we check the endpoints, which in this case are $\alpha = 0$ and $\alpha = \frac{\pi}{2}$:

$D(0) = \pi$ radians $= 180^\circ$, and $D\left(\frac{\pi}{2}\right) \approx 166^\circ$.

Another method: We first calculate $\dfrac{d\beta}{d\alpha}$: $\sin\alpha = \frac{4}{3}\sin\beta \quad\Leftrightarrow\quad \cos\alpha = \frac{4}{3}\cos\beta\,\dfrac{d\beta}{d\alpha} \quad\Leftrightarrow\quad \dfrac{d\beta}{d\alpha} = \dfrac{3}{4}\dfrac{\cos\alpha}{\cos\beta}$, so since

$D'(\alpha) = 2 - 4\dfrac{d\beta}{d\alpha} = 0 \quad\Leftrightarrow\quad \dfrac{d\beta}{d\alpha} = \dfrac{1}{2}$, the minimum occurs when $3\cos\alpha = 2\cos\beta$. Now we square both sides and substitute $\sin\alpha = \frac{4}{3}\sin\beta$, leading to the same result.

2. If we repeat Problem 1 with k in place of $\frac{4}{3}$, we get $D(\alpha) = \pi + 2\alpha - 4\arcsin\left(\dfrac{1}{k}\sin\alpha\right) \quad\Rightarrow$

$D'(\alpha) = 2 - \dfrac{4\cos\alpha}{k\sqrt{1 - \left(\dfrac{\sin\alpha}{k}\right)^2}}$, which is 0 when $\dfrac{2\cos\alpha}{k} = \sqrt{1 - \left(\dfrac{\sin\alpha}{k}\right)^2} \quad\Leftrightarrow\quad \left(\dfrac{2\cos\alpha}{k}\right)^2 = 1 - \left(\dfrac{\sin\alpha}{k}\right)^2 \quad\Leftrightarrow$

$4\cos^2\alpha = k^2 - \sin^2\alpha \quad\Leftrightarrow\quad 3\cos^2\alpha = k^2 - 1 \quad\Leftrightarrow\quad \alpha = \arccos\sqrt{\dfrac{k^2 - 1}{3}}$. So for $k \approx 1.3318$ (red light) the minimum occurs at $\alpha_1 \approx 1.038$ radians, and so the rainbow angle is about $\pi - D(\alpha_1) \approx 42.3^\circ$. For $k \approx 1.3435$ (violet light) the minimum occurs at $\alpha_2 \approx 1.026$ radians, and so the rainbow angle is about $\pi - D(\alpha_2) \approx 40.6^\circ$.

Another method: As in Problem 1, we can instead find $D'(\alpha)$ in terms of $\dfrac{d\beta}{d\alpha}$, and then substitute $\dfrac{d\beta}{d\alpha} = \dfrac{\cos\alpha}{k\cos\beta}$.

3. At each reflection or refraction, the light is bent in a counterclockwise direction: the bend at A is $\alpha - \beta$, the bend at B is $\pi - 2\beta$, the bend at C is again $\pi - 2\beta$, and the bend at D is $\alpha - \beta$. So the total bend is

$D(\alpha) = 2(\alpha - \beta) + 2(\pi - 2\beta) = 2\alpha - 6\beta + 2\pi$, as required. We substitute $\beta = \arcsin\left(\dfrac{\sin\alpha}{k}\right)$ and differentiate, to get

$D'(\alpha) = 2 - \dfrac{6\cos\alpha}{k\sqrt{1 - \left(\dfrac{\sin\alpha}{k}\right)^2}}$, which is 0 when $\dfrac{3\cos\alpha}{k} = \sqrt{1 - \left(\dfrac{\sin\alpha}{k}\right)^2} \quad\Leftrightarrow\quad 9\cos^2\alpha = k^2 - \sin^2\alpha \quad\Leftrightarrow$

$8\cos^2\alpha = k^2 - 1 \quad\Leftrightarrow\quad \cos\alpha = \sqrt{\frac{1}{8}(k^2 - 1)}$. If $k = \frac{4}{3}$, then the minimum occurs at

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\alpha_1 = \arccos\sqrt{\dfrac{(4/3)^2 - 1}{8}} \approx 1.254$ radians. Thus, the minimum *counterclockwise* rotation is $D(\alpha_1) \approx 231°$, which is equivalent to a *clockwise* rotation of $360° - 231° = 129°$ (see the figure). So the rainbow angle for the secondary rainbow is about $180° - 129° = 51°$, as required. In general, the rainbow angle for the secondary rainbow is $\pi - [2\pi - D(\alpha)] = D(\alpha) - \pi$.

4. In the primary rainbow, the rainbow angle gets smaller as k gets larger, as we found in Problem 2, so the colors appear from top to bottom in order of increasing k. But in the secondary rainbow, the rainbow angle gets larger as k gets larger. To see this, we find the minimum deviations for red light and for violet light in the secondary rainbow. For $k \approx 1.3318$ (red light) the minimum occurs at $\alpha_1 \approx \arccos\sqrt{\dfrac{1.3318^2 - 1}{8}} \approx 1.255$ radians, and so the rainbow angle is $D(\alpha_1) - \pi \approx 50.6°$. For $k \approx 1.3435$ (violet light) the minimum occurs at $\alpha_2 \approx \arccos\sqrt{\dfrac{1.3435^2 - 1}{8}} \approx 1.248$ radians, and so the rainbow angle is $D(\alpha_2) - \pi \approx 53.6°$. Consequently, the rainbow angle is larger for colors with higher indices of refraction, and the colors appear from bottom to top in order of increasing k, the reverse of their order in the primary rainbow.

Note that our calculations above also explain why the secondary rainbow is more spread out than the primary rainbow: in the primary rainbow, the difference between rainbow angles for red and violet light is about $1.7°$, whereas in the secondary rainbow it is about $3°$.

4.2 The Mean Value Theorem

1. **(1)** f is continuous on the closed interval $[0, 8]$.

(2) f is differentiable on the open interval $(0, 8)$.

(3) $f(0) = 3$ and $f(8) = 3$

Thus, f satisfies the hypotheses of Rolle's Theorem. The numbers $c = 1$ and $c = 5$ satisfy the conclusion of Rolle's Theorem since $f'(1) = f'(5) = 0$.

2. The possible graphs fall into two general categories: **(1)** Not continuous and therefore not differentiable, **(2)** Continuous, but not differentiable.

Not continuous

Not differentiable

In either case, there is no number c such that $f'(c) = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a) **(1)** g is continuous on the closed interval $[0, 8]$.

(2) g is differentiable on the open interval $(0, 8)$.

(b) $g'(c) = \dfrac{g(8) - g(0)}{8 - 0} = \dfrac{4 - 1}{8} = \dfrac{3}{8}$.

It appears that $g'(c) = \frac{3}{8}$ when $c \approx 2.2$ and 6.4.

(c) $g'(c) = \dfrac{g(6) - g(2)}{6 - 2} = \dfrac{1 - 3}{4} = -\dfrac{1}{2}$.

It appears that $g'(c) = -\frac{1}{2}$ when $c \approx 3.7$ and 5.5.

4.

The function shown in the figure is continuous on $[0, 8]$ [but not differentiable on $(0, 8)$] with $f(0) = 1$ and $f(8) = 4$. The line passing through the two points has slope $\frac{3}{8}$. There is no number c in $(0, 8)$ such that $f'(c) = \frac{3}{8}$.

5. **(1)** f is continuous on the closed interval $[0, 5]$.

(2) f is not differentiable on the open interval $(0, 5)$ since f is not differentiable at 3.

Thus, f does not satisfy the hypotheses of the Mean Value Theorem on the interval $[0, 5]$.

6. **(1)** f is continuous on the closed interval $[0, 5]$.

(2) f is differentiable on the open interval $(0, 5)$.

Thus, f satisfies the hypotheses of the Mean Value Theorem on the interval $[0, 5]$.

The line passing through $(0, f(0))$ and $(5, f(5))$ has slope 0. It appears that $f'(c) = 0$ for $c = 1$ and $c = 3$.

7. **(1)** f is continuous on the closed interval $[0, 5]$.

(2) f is differentiable on the open interval $(0, 5)$.

Thus, f satisfies the hypotheses of the Mean Value Theorem on the interval $[0, 5]$.

The line passing through $(0, f(0))$ and $(5, f(5))$ has slope

$\dfrac{f(5) - f(0)}{5 - 0} = \dfrac{3 - 1}{5} = \dfrac{2}{5}$. It appears that $f'(c) = \frac{2}{5}$ when $c \approx 3.8$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. **(1)** f is continuous on the closed interval $[0, 5]$.

(2) f is not differentiable on the open interval $(0, 5)$ since f is not differentiable at 4.

Thus, f does not satisfy the hypotheses of the Mean Value Theorem on the interval $[0, 5]$.

9. $f(x) = 2x^2 - 4x + 5,\ [-1, 3]$. f is a polynomial, so it's continuous and differentiable on $\mathbb{R}$, and hence, continuous on $[-1, 3]$ and differentiable on $(-1, 3)$. Since $f(-1) = 11$ and $f(3) = 11$, f satisfies all the hypotheses of Rolle's Theorem. $f'(c) = 4c - 4$ and $f'(c) = 0 \quad \Leftrightarrow \quad 4c - 4 = 0 \quad \Leftrightarrow \quad c = 1$. $c = 1$ is in the interval $(-1, 3)$, so 1 satisfies the conclusion of Rolle's Theorem.

10. $f(x) = x^3 - 2x^2 - 4x + 2,\ [-2, 2]$. f is a polynomial, so it's continuous and differentiable on $\mathbb{R}$, and hence, continuous on $[-2, 2]$ and differentiable on $(-2, 2)$. Since $f(-2) = -6$ and $f(2) = -6$, f satisfies all the hypotheses of Rolle's Theorem. $f'(c) = 3c^2 - 4c - 4$ and $f'(c) = 0 \quad \Leftrightarrow \quad (3c + 2)(c - 2) = 0 \quad \Leftrightarrow \quad c = -\frac{2}{3}$ or 2. $c = -\frac{2}{3}$ is in the open interval $(-2, 2)$ (but 2 *isn't*), so only $-\frac{2}{3}$ satisfies the conclusion of Rolle's Theorem.

11. $f(x) = \sin(x/2),\ [\pi/2, 3\pi/2]$. f, being the composite of the sine function and the polynomial $x/2$, is continuous and differentiable on $\mathbb{R}$, so it is continuous on $[\pi/2, 3\pi/2]$ and differentiable on $(\pi/2, 3\pi/2)$. Also, $f\left(\frac{\pi}{2}\right) = \frac{1}{2}\sqrt{2} = f\left(\frac{3\pi}{2}\right)$.
$f'(c) = 0 \quad \Leftrightarrow \quad \frac{1}{2}\cos(c/2) = 0 \quad \Leftrightarrow \quad \cos(c/2) = 0 \quad \Leftrightarrow \quad c/2 = \frac{\pi}{2} + n\pi$ [n an integer] $\quad \Leftrightarrow \quad c = \pi + 2n\pi$.
Only $c = \pi$ [when $n = 0$] is in $(\pi/2, 3\pi/2)$, so π satisfies the conclusion of Rolle's Theorem.

12. $f(x) = x + 1/x,\ \left[\frac{1}{2}, 2\right]$. $f'(x) = 1 - 1/x^2 = \dfrac{x^2 - 1}{x^2}$. f is a rational function that is continuous on its domain, $(-\infty, 0) \cup (0, \infty)$, so it is continuous on $\left[\frac{1}{2}, 2\right]$. f' has the same domain and is differentiable on $\left(\frac{1}{2}, 2\right)$. Also, $f\left(\frac{1}{2}\right) = \frac{5}{2} = f(2)$. $f'(c) = 0 \quad \Leftrightarrow \quad \dfrac{c^2 - 1}{c^2} = 0 \quad \Leftrightarrow \quad c^2 - 1 = 0 \quad \Leftrightarrow \quad c = \pm 1$. Only 1 is in $\left(\frac{1}{2}, 2\right)$, so 1 satisfies the conclusion of Rolle's Theorem.

13. $f(x) = 1 - x^{2/3}$. $f(-1) = 1 - (-1)^{2/3} = 1 - 1 = 0 = f(1)$. $f'(x) = -\frac{2}{3}x^{-1/3}$, so $f'(c) = 0$ has no solution. This does not contradict Rolle's Theorem, since $f'(0)$ does not exist, and so f is not differentiable on $(-1, 1)$.

14. $f(x) = \tan x$. $f(0) = \tan 0 = 0 = \tan \pi = f(\pi)$. $f'(x) = \sec^2 x \geq 1$, so $f'(c) = 0$ has no solution. This does not contradict Rolle's Theorem, since $f'\left(\frac{\pi}{2}\right)$ does not exist, and so f is not differentiable on $(0, \pi)$. (Also, f is not continuous on $[0, \pi]$.)

15. $f(x) = 2x^2 - 3x + 1,\ [0, 2]$. f is continuous on $[0, 2]$ and differentiable on $(0, 2)$ since polynomials are continuous and differentiable on $\mathbb{R}$. $f'(c) = \dfrac{f(b) - f(a)}{b - a} \quad \Leftrightarrow \quad 4c - 3 = \dfrac{f(2) - f(0)}{2 - 0} = \dfrac{3 - 1}{2} = 1 \quad \Leftrightarrow \quad 4c = 4 \quad \Leftrightarrow \quad c = 1$, which is in $(0, 2)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. $f(x) = x^3 - 3x + 2,\ [-2, 2]$. f is continuous on $[-2, 2]$ and differentiable on $(-2, 2)$ since polynomials are continuous and differentiable on $\mathbb{R}$. $f'(c) = \dfrac{f(b) - f(a)}{b - a} \quad \Leftrightarrow \quad 3c^2 - 3 = \dfrac{f(2) - f(-2)}{2 - (-2)} = \dfrac{4 - 0}{4} = 1 \quad \Leftrightarrow \quad 3c^2 = 4 \quad \Leftrightarrow$

$c^2 = \dfrac{4}{3} \quad \Leftrightarrow \quad c = \pm\dfrac{2}{\sqrt{3}} \approx \pm 1.15$, which are both in $(-2, 2)$.

17. $f(x) = \ln x,\ [1, 4]$. f is continuous and differentiable on $(0, \infty)$, so f is continuous on $[1, 4]$ and differentiable on $(1, 4)$.

$f'(c) = \dfrac{f(b) - f(a)}{b - a} \quad \Leftrightarrow \quad \dfrac{1}{c} = \dfrac{f(4) - f(1)}{4 - 1} = \dfrac{\ln 4 - 0}{3} = \dfrac{\ln 4}{3} \quad \Leftrightarrow \quad c = \dfrac{3}{\ln 4} \approx 2.16$, which is in $(1, 4)$.

18. $f(x) = 1/x,\ [1, 3]$. f is continuous and differentiable on $(-\infty, 0) \cup (0, \infty)$, so f is continous on $[1, 3]$ and differentiable on $(1, 3)$. $f'(c) = \dfrac{f(b) - f(a)}{b - a} \quad \Leftrightarrow \quad -\dfrac{1}{c^2} = \dfrac{f(3) - f(1)}{3 - 1} = \dfrac{\frac{1}{3} - 1}{2} = -\dfrac{1}{3} \quad \Leftrightarrow \quad c^2 = 3 \quad \Leftrightarrow \quad c = \pm\sqrt{3} \approx \pm 1.73$, but only $\sqrt{3}$ is in $(1, 3)$.

19. $f(x) = \sqrt{x},\ [0, 4]$. $f'(c) = \dfrac{f(4) - f(0)}{4 - 0} \quad \Leftrightarrow \quad \dfrac{1}{2\sqrt{c}} = \dfrac{2 - 0}{4} \quad \Leftrightarrow$

$\dfrac{1}{2\sqrt{c}} = \dfrac{1}{2} \quad \Leftrightarrow \quad \sqrt{c} = 1 \quad \Leftrightarrow \quad c = 1$. The secant line and the tangent line are parallel.

20. $f(x) = e^{-x}, [0, 2]$. $f'(c) = \dfrac{f(2) - f(0)}{2 - 0} \quad \Leftrightarrow \quad -e^{-c} = \dfrac{e^{-2} - 1}{2} \quad \Leftrightarrow$

$e^{-c} = \dfrac{1 - e^{-2}}{2} \quad \Leftrightarrow \quad -c = \ln\dfrac{1 - e^{-2}}{2} \quad \Leftrightarrow$

$c = -\ln\dfrac{1 - e^{-2}}{2} \approx 0.84$. The secant line and the tangent line are parallel.

21. $f(x) = (x - 3)^{-2} \quad \Rightarrow \quad f'(x) = -2(x - 3)^{-3}$. $f(4) - f(1) = f'(c)(4 - 1) \quad \Rightarrow \quad \dfrac{1}{1^2} - \dfrac{1}{(-2)^2} = \dfrac{-2}{(c - 3)^3} \cdot 3 \quad \Rightarrow$

$\dfrac{3}{4} = \dfrac{-6}{(c - 3)^3} \quad \Rightarrow \quad (c - 3)^3 = -8 \quad \Rightarrow \quad c - 3 = -2 \quad \Rightarrow \quad c = 1$, which is not in the *open* interval $(1, 4)$. This does not contradict the Mean Value Theorem since f is not continuous at $x = 3$, which is in the interval $[1, 4]$.

22. $f(x) = 2 - |2x - 1| = \begin{cases} 2 - (2x - 1) & \text{if } 2x - 1 \geq 0 \\ 2 - [-(2x - 1)] & \text{if } 2x - 1 < 0 \end{cases} = \begin{cases} 3 - 2x & \text{if } x \geq \frac{1}{2} \\ 1 + 2x & \text{if } x < \frac{1}{2} \end{cases} \quad \Rightarrow \quad f'(x) = \begin{cases} -2 & \text{if } x > \frac{1}{2} \\ 2 & \text{if } x < \frac{1}{2} \end{cases}$

$f(3) - f(0) = f'(c)(3 - 0) \quad \Rightarrow \quad -3 - 1 = f'(c) \cdot 3 \quad \Rightarrow \quad f'(c) = -\frac{4}{3}$ [not ± 2]. This does not contradict the Mean Value Theorem since f is not differentiable at $x = \frac{1}{2}$, which is in the interval $(0, 3)$.

23. Let $f(x) = 2x + \cos x$. Then $f(-\pi) = -2\pi - 1 < 0$ and $f(0) = 1 > 0$. Since f is the sum of the polynomial $2x$ and the trignometric function $\cos x$, f is continuous and differentiable for all x. By the Intermediate Value Theorem, there is a

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

number c in $(-\pi, 0)$ such that $f(c) = 0$. Thus, the given equation has at least one real solution. If the equation has distinct real solutions a and b with $a < b$, then $f(a) = f(b) = 0$. Since f is continuous on $[a, b]$ and differentiable on (a, b), Rolle's Theorem implies that there is a number r in (a, b) such that $f'(r) = 0$. But $f'(r) = 2 - \sin r > 0$ since $\sin r \leq 1$. This contradiction shows that the given equation can't have two distinct real solutions, so it has exactly one solution.

24. Let $f(x) = x^3 + e^x$. Then $f(-1) = -1 + 1/e < 0$ and $f(0) = 1 > 0$. Since f is the sum of a polynomial and the natural exponential function, f is continous and differentiable for all x. By the Intermediate Value Theorem, there is a number c in $(-1, 0)$ such that $f(c) = 0$. Thus, the given equation has at least one real solution. If the equation has distinct real solutions a and b with $a < b$, then $f(a) = f(b) = 0$. Since f is continuous on $[a, b]$ and differentiable on (a, b), Rolle's Theorem implies that there is a number r in (a, b) such that $f'(r) = 0$. But $f'(r) = 3r^2 + e^r > 0$. This contradiction shows that the given equation can't have two distinct real solutions, so it has exactly one solution.

25. Let $f(x) = x^3 - 15x + c$ for x in $[-2, 2]$. If f has two real solutions a and b in $[-2, 2]$, with $a < b$, then $f(a) = f(b) = 0$. Since the polynomial f is continuous on $[a, b]$ and differentiable on (a, b), Rolle's Theorem implies that there is a number r in (a, b) such that $f'(r) = 0$. Now $f'(r) = 3r^2 - 15$. Since r is in (a, b), which is contained in $[-2, 2]$, we have $|r| < 2$, so $r^2 < 4$. It follows that $3r^2 - 15 < 3 \cdot 4 - 15 = -3 < 0$. This contradicts $f'(r) = 0$, so the given equation, $x^3 - 15x + c = 0$, can't have two real solutions in $[-2, 2]$. Hence, it has at most one real solution in $[-2, 2]$.

26. $f(x) = x^4 + 4x + c$. Suppose that $f(x) = 0$ has three distinct real solutions a, b, d where $a < b < d$. Then $f(a) = f(b) = f(d) = 0$. By Rolle's Theorem there are numbers c_1 and c_2 with $a < c_1 < b$ and $b < c_2 < d$ and $0 = f'(c_1) = f'(c_2)$, so $f'(x) = 0$ must have at least two real solutions. However $0 = f'(x) = 4x^3 + 4 = 4(x^3 + 1) = 4(x + 1)(x^2 - x + 1)$ has as its only real solution $x = -1$. Thus, the given equation, $x^4 + 4x + c = 0$, can have at most two real solutions.

27. (a) Suppose that a cubic polynomial $P(x)$ has zeros $a_1 < a_2 < a_3 < a_4$, so $P(a_1) = P(a_2) = P(a_3) = P(a_4)$. By Rolle's Theorem there are numbers c_1, c_2, c_3 with $a_1 < c_1 < a_2$, $a_2 < c_2 < a_3$, and $a_3 < c_3 < a_4$ and $P'(c_1) = P'(c_2) = P'(c_3) = 0$. Thus, the second-degree polynomial $P'(x)$ has three distinct real zeros, which is impossible. This contradiction tells us that a polynomial of degree 3 has at most three real zeros.

(b) We prove by induction that a polynomial of degree n has at most n real zeros. This is certainly true for $n = 1$. Suppose that the result is true for all polynomials of degree n and let $P(x)$ be a polynomial of degree $n + 1$. Suppose that $P(x)$ has more than $n + 1$ real zeros, say $a_1 < a_2 < a_3 < \cdots < a_{n+1} < a_{n+2}$. Then $P(a_1) = P(a_2) = \cdots = P(a_{n+2}) = 0$. By Rolle's Theorem there are real numbers $c_1, \ldots, c_{n+1}$ with $a_1 < c_1 < a_2, \ldots, a_{n+1} < c_{n+1} < a_{n+2}$ and $P'(c_1) = \cdots = P'(c_{n+1}) = 0$. Thus, the nth-degree polynomial $P'(x)$ has at least $n + 1$ zeros. This contradiction shows that $P(x)$ has at most $n + 1$ real zeros and hence, a polynomial of degree n has at most n real zeros.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. (a) Suppose that $f(a) = f(b) = 0$ where $a < b$. By Rolle's Theorem applied to f on $[a, b]$, there is a number c such that $a < c < b$ and $f'(c) = 0$.

(b) Suppose that $f(a) = f(b) = f(c) = 0$ where $a < b < c$. By Rolle's Theorem applied to f on $[a, b]$ and $[b, c]$, there are numbers d and e such that $a < d < b$ and $b < e < c$, with $f'(d) = 0$ and $f'(e) = 0$. By Rolle's Theorem applied to f' on $[d, e]$, there is a number g such that $d < g < e$ and $f''(g) = 0$.

(c) Suppose that f is n times differentiable on $\mathbb{R}$ and has $n + 1$ distinct real zeros. Then $f^{(n)}$ has at least one real zero.

29. By the Mean Value Theorem, $f(4) - f(1) = f'(c)(4 - 1)$ for some $c \in (1, 4)$. (f is differentiable for all x, so, in particular, f is differentiable on $(1, 4)$ and continuous on $[1, 4]$. Thus, the hypotheses of the Mean Value Theorem are satisfied.) For every $c \in (1, 4)$, we have $f'(c) \geq 2$. Putting $f'(c) \geq 2$ into the above equation and substituting $f(1) = 10$, we get $f(4) = f(1) + f'(c)(4 - 1) = 10 + 3f'(c) \geq 10 + 3 \cdot 2 = 16$. So the smallest possible value of $f(4)$ is 16.

30. By the Mean Value Theorem, $f(8) - f(2) = f'(c)(8 - 2)$ for some $c \in (2, 8)$. (f is differentiable for all x, so, in particular, f is differentiable on $(2, 8)$ and continuous on $[2, 8]$. Thus, the hypotheses of the Mean Value Theorem are satisfied.) Since $f(8) - f(2) = 6f'(c)$ and $3 \leq f'(c) \leq 5$, it follows that $6 \cdot 3 \leq 6f'(c) \leq 6 \cdot 5$, so $18 \leq f(8) - f(2) \leq 30$.

31. Suppose that such a function f exists. By the Mean Value Theorem, there is a number c such that $0 < c < 2$ with $f'(c) = \dfrac{f(2) - f(0)}{2 - 0} = \dfrac{4 - (-1)}{2 - 0} = \dfrac{5}{2}$. This result, $f'(c) = \dfrac{5}{2}$, is impossible since $f'(x) \leq 2$ for all x, so no such function f exists.

32. Let $h = f - g$. Note that since $f(a) = g(a)$, $h(a) = f(a) - g(a) = 0$. Then since f and g are continuous on $[a, b]$ and differentiable on (a, b), so is h, and thus h satisfies the assumptions of the Mean Value Theorem. Therefore, there is a number c with $a < c < b$ such that $h(b) = h(b) - h(a) = h'(c)(b - a)$. Given $f'(x) < g'(x)$, we have $f' - g' < 0$ or, equivalently, $h' < 0$. Now since $h'(c) < 0$, $h'(c)(b - a) < 0$, so $h(b) = f(b) - g(b) < 0$ and hence $f(b) < g(b)$.

33. Consider the function $f(x) = \sin x$, which is continuous and differentiable on $\mathbb{R}$. Let a be a number such that $0 < a < 2\pi$. Then f is continuous on $[0, a]$ and differentiable on $(0, a)$. By the Mean Value Theorem, there is a number c in $(0, a)$ such that $f(a) - f(0) = f'(c)(a - 0)$; that is, $\sin a - 0 = (\cos c)(a)$. Now $\cos c < 1$ for $0 < c < 2\pi$, so $\sin a < 1 \cdot a = a$. We took a to be an arbitrary number in $(0, 2\pi)$, so $\sin x < x$ for all x satisfying $0 < x < 2\pi$.

34. f satisfies the conditions for the Mean Value Theorem, so we use this theorem on the interval $[-b, b]$: $f'(c) = \dfrac{f(b) - f(-b)}{b - (-b)}$ for some $c \in (-b, b)$. Since f is odd, $f(-b) = -f(b)$. Substituting this into the above equation, we get $f'(c) = \dfrac{f(b) + f(b)}{2b}$, or, equivalently, $f'(c) = \dfrac{f(b)}{b}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. Let $f(x) = \sin x$ on $[a, b]$. Then f is continuous on $[a, b]$ and differentiable on (a, b). By the Mean Value Theorem, there is a number $c \in (a, b)$ with $f(b) - f(a) = f'(c)(b - a)$ or, equivalently, $\sin b - \sin a = (\cos c)(b - a)$. Taking absolute values, $|\sin b - \sin a| \le |\cos c|\,|b - a|$ or, equivalently, $|\sin a - \sin b| \le 1\,|b - a|$. If $b < a$, then $|\sin a - \sin b| \le |a - b|$. If $a = b$, both sides of the inequality are 0, which proves the given inequality for all a and b.

36. Suppose that $f'(x) = c$. Let $g(x) = cx$, so $g'(x) = c$. Then, by Corollary 7, $f(x) = g(x) + d$, where d is a constant, so $f(x) = cx + d$.

37. For $x > 0$, $f(x) = g(x)$, so $f'(x) = g'(x)$. For $x < 0$, $f'(x) = (1/x)' = -1/x^2$ and $g'(x) = (1 + 1/x)' = -1/x^2$, so again $f'(x) = g'(x)$. However, the domain of $g(x)$ is not an interval [it is $(-\infty, 0) \cup (0, \infty)$], so we cannot conclude that $f - g$ is constant (in fact, it is not).

38. Let $f(x) = \arctan x + \arctan\left(\dfrac{1}{x}\right)$. Then $f'(x) = \dfrac{1}{1+x^2} + \dfrac{1}{1+\left(\dfrac{1}{x}\right)^2} \cdot \left(-\dfrac{1}{x^2}\right) = \dfrac{1}{1+x^2} - \dfrac{1}{x^2+1} = 0$ on the given domain, $(0, \infty)$. Therefore, $f(x) = C$ on $(0, \infty)$ by Theorem 5. To find C, we let $x = 1$ to get

$\arctan 1 + \arctan\left(\dfrac{1}{1}\right) = C \quad \Rightarrow \quad \dfrac{\pi}{4} + \dfrac{\pi}{4} = \dfrac{\pi}{2} = C$. Thus, $f(x) = \dfrac{\pi}{2}$; that is, $\arctan x + \arctan\left(\dfrac{1}{x}\right) = \dfrac{\pi}{2}$.

39. Let $f(x) = 2\sin^{-1} x - \cos^{-1}(1 - 2x^2)$. Then

$$\begin{aligned} f'(x) &= \frac{2}{\sqrt{1-x^2}} - \left(-\frac{-4x}{\sqrt{1-(1-2x^2)^2}}\right) = \frac{2}{\sqrt{1-x^2}} - \frac{4x}{\sqrt{4x^2-4x^4}} \\ &= \frac{2}{\sqrt{1-x^2}} - \frac{4x}{2x\sqrt{1-x^2}} \ \text{[since } x \ge 0] \ = 0 \end{aligned}$$

Thus, $f'(x) = 0$ for all $x \in (0, 1)$, and hence, $f(x) = C$ on $(0, 1)$. To find C, let $x = 0.5$ to get $f(0.5) = 2\,\sin^{-1}(0.5) - \cos^{-1}(0.5) = 2\left(\frac{\pi}{6}\right) - \frac{\pi}{3} = 0 = C$. We conclude that $f(x) = 0$ for x in $(0, 1)$. By continuity of f, $f(x) = 0$ on $[0, 1]$. Therefore, we see that $f(x) = 2\,\sin^{-1} x - \cos^{-1}(1 - 2x^2) = 0 \quad \Rightarrow \quad 2\,\sin^{-1} x = \cos^{-1}(1 - 2x^2)$.

40. Let $v(t)$ be the velocity of the car t hours after 2:00 PM. Then $\dfrac{v(1/6) - v(0)}{1/6 - 0} = \dfrac{50 - 30}{1/6} = 120$. By the Mean Value Theorem, there is a number c such that $0 < c < \frac{1}{6}$ with $v'(c) = 120$. Since $v'(t)$ is the acceleration at time t, the acceleration c hours after 2:00 PM is exactly $120\ \text{mi/h}^2$.

41. Let $g(t)$ and $h(t)$ be the position functions of the two runners and let $f(t) = g(t) - h(t)$. By hypothesis, where b is the finishing time, $f(0) = g(0) - h(0) = 0$ and $f(b) = g(b) - h(b) = 0$. Then by the Mean Value Theorem, there is a time c, with $0 < c < b$, such that $f'(c) = \dfrac{f(b) - f(0)}{b - 0} = \dfrac{0 - 0}{b} = 0$. Since $f'(c) = g'(c) - h'(c) = 0$, we have $g'(c) = h'(c)$ [the velocities are equal]. So at time c, both runners have the same speed.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. Assume that f is differentiable (and hence continuous) on $\mathbb{R}$ and that $f'(x) \neq 1$ for all x. Suppose f has more than one fixed point. Then there are numbers a and b such that $a < b$, $f(a) = a$, and $f(b) = b$. Applying the Mean Value Theorem to the function f on $[a, b]$, we find that there is a number c in (a, b) such that $f'(c) = \dfrac{f(b) - f(a)}{b - a}$. But then $f'(c) = \dfrac{b - a}{b - a} = 1$, contradicting our assumption that $f'(x) \neq 1$ for every real number x. This shows that our supposition was wrong, that is, that f cannot have more than one fixed point.

4.3 What Derivatives Tell Us about the Shape of a Graph

1. (a) f is increasing on $(1, 3)$ and $(4, 6)$.
(b) f is decreasing on $(0, 1)$ and $(3, 4)$.
(c) f is concave upward on $(0, 2)$.
(d) f is concave downward on $(2, 4)$ and $(4, 6)$.
(e) The point of inflection is $(2, 3)$.

2. (a) f is increasing on $(0, 1)$ and $(3, 7)$.
(b) f is decreasing on $(1, 3)$.
(c) f is concave upward on $(2, 4)$ and $(5, 7)$.
(d) f is concave downward on $(0, 2)$ and $(4, 5)$.
(e) The points of inflection are $(2, 2)$, $(4, 3)$, and $(5, 4)$.

3. (a) Use the Increasing/Decreasing (I/D) Test.
(b) Use the Concavity Test.
(c) At any value of x where the concavity changes, we have an inflection point at $(x, f(x))$.

4. (a) See the First Derivative Test.
(b) See the Second Derivative Test and the note that precedes Example 6.

5. (a) Since $f'(x) > 0$ on $(0, 1)$ and $(3, 5)$, f is increasing on these intervals. Since $f'(x) < 0$ on $(1, 3)$ and $(5, 6)$, f is decreasing on these intervals.
(b) Since $f'(x) = 0$ at $x = 1$ and $x = 5$, and f' changes from positive to negative at both values, f changes from increasing to decreasing and has local maxima at $x = 1$ and $x = 5$. Since $f'(x) = 0$ at $x = 3$, and f' changes from negative to positive there, f changes from decreasing to increasing and has a local minimum at $x = 3$.

6. (a) Since $f'(x) > 0$ on $(1, 4)$ and $(5, 6)$, f is increasing on these intervals. Since $f'(x) < 0$ on $(0, 1)$ and $(4, 5)$, f is decreasing on these intervals.
(b) Since $f'(x) = 0$ at $x = 4$, and f' changes from positive to negative there, f changes from increasing to decreasing and has a local maximum at $x = 4$. Since $f'(x) = 0$ at $x = 1$ and $x = 5$, and f' changes from negative to positive at both values, f changes from decreasing to increasing and has local minima at $x = 1$ and $x = 5$.

7. (a) There is an IP at $x = 3$ because the graph of f changes from CD to CU there. There is an IP at $x = 5$ because the graph of f changes from CU to CD there.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) There is an IP at $x = 2$ and at $x = 6$ because $f'(x)$ has a maximum value there, and so $f''(x)$ changes from positive to negative there. There is an IP at $x = 4$ because $f'(x)$ has a minimum value there and so $f''(x)$ changes from negative to positive there.

(c) There is an inflection point at $x = 1$ because $f''(x)$ changes from negative to positive there, and so the graph of f changes from concave downward to concave upward. There is an inflection point at $x = 7$ because $f''(x)$ changes from positive to negative there, and so the graph of f changes from concave upward to concave downward.

8. (a) f is increasing when f' is positive. This happens on the intervals $(0, 4)$ and $(6, 8)$.

(b) f has a local maximum where it changes from increasing to decreasing, that is, where f' changes from positive to negative (at $x = 4$ and $x = 8$). Similarly, f has a local minimum where f' changes from negative to positive (at $x = 6$).

(c) f is concave upward where f' is increasing (hence f'' is positive). This happens on $(0, 1)$, $(2, 3)$, and $(5, 7)$. Similarly, f is concave downward where f' is decreasing, that is, on $(1, 2)$, $(3, 5)$, and $(7, 9)$.

(d) f has an inflection point where the concavity changes. This happens at $x = 1$, 2, 3, 5, and 7.

9. $f(x) = 2x^3 - 15x^2 + 24x - 5 \quad \Rightarrow \quad f'(x) = 6x^2 - 30x + 24 = 6(x^2 - 5x + 4) = 6(x - 1)(x - 4)$.

Interval	$x - 1$	$x - 4$	$f'(x)$	f
$x < 1$	$-$	$-$	$+$	increasing on $(-\infty, 1)$
$1 < x < 4$	$+$	$-$	$-$	decreasing on $(1, 4)$
$x > 4$	$+$	$+$	$+$	increasing on $(4, \infty)$

f changes from increasing to decreasing at $x = 1$ and from decreasing to increasing at $x = 4$. Thus, $f(1) = 6$ is a local maximum value and $f(4) = -21$ is a local minimum value.

10. $f(x) = x^3 - 6x^2 - 135x \quad \Rightarrow \quad f'(x) = 3x^2 - 12x - 135 = 3(x^2 - 4x - 45) = 3(x + 5)(x - 9)$.

Interval	$x + 5$	$x - 9$	$f'(x)$	f
$x < -5$	$-$	$-$	$+$	increasing on $(-\infty, -5)$
$-5 < x < 9$	$+$	$-$	$-$	decreasing on $(-5, 9)$
$x > 9$	$+$	$+$	$+$	increasing on $(9, \infty)$

f changes from increasing to decreasing at $x = -5$ and from decreasing to increasing at $x = 9$. Thus, $f(-5) = 400$ is a local maximum value and $f(9) = -972$ is a local minimum value.

11. $f(x) = 6x^4 - 16x^3 + 1 \quad \Rightarrow \quad f'(x) = 24x^3 - 48x^2 = 24x^2(x - 2)$.

Interval	x^2	$x - 2$	$f'(x)$	f
$x < 0$	$+$	$-$	$-$	decreasing on $(-\infty, 0)$
$0 < x < 2$	$+$	$-$	$-$	decreasing on $(0, 2)$
$x > 2$	$+$	$+$	$+$	increasing on $(2, \infty)$

Note that f is differentiable and $f'(x) < 0$ on the interval $(-\infty, 2)$ except for the single number $x = 0$. By applying the result

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

of Exercise 4.3.97, we can say that f is decreasing on the entire interval $(-\infty, 2)$. f changes from decreasing to increasing at $x = 2$. Thus, $f(2) = -31$ is a local minimum value.

12. $f(x) = x^{2/3}(x-3) \quad \Rightarrow \quad f'(x) = x^{2/3}(1) + (x-3)\cdot\frac{2}{3}x^{-1/3} = \frac{1}{3}x^{-1/3}\left[3x + (x-3)\cdot 2\right] = \frac{1}{3}x^{-1/3}(5x-6)$.

Interval	$x^{-1/3}$	$5x-6$	$f'(x)$	f
$x < 0$	$-$	$-$	$+$	increasing on $(-\infty, 0)$
$0 < x < \frac{6}{5}$	$+$	$-$	$-$	decreasing on $\left(0, \frac{6}{5}\right)$
$x > \frac{6}{5}$	$+$	$+$	$+$	increasing on $\left(\frac{6}{5}, \infty\right)$

f changes from increasing to decreasing at $x = 0$ and from decreasing to increasing at $x = \frac{6}{5}$. Thus, $f(0) = 0$ is a local maximum value and $f\left(\frac{6}{5}\right) = \left(\frac{6}{5}\right)^{2/3}\left(-\frac{9}{5}\right) \approx -2.03$ is a local minimum value.

13. $f(x) = \dfrac{x^2-24}{x-5} \quad \Rightarrow$

$$f'(x) = \frac{(x-5)(2x) - (x^2-24)(1)}{(x-5)^2} = \frac{2x^2 - 10x - x^2 + 24}{(x-5)^2} = \frac{x^2 - 10x + 24}{(x-5)^2} = \frac{(x-4)(x-6)}{(x-5)^2}.$$

Interval	$x-4$	$x-6$	$(x-5)^2$	$f'(x)$	f
$x < 4$	$-$	$-$	$+$	$+$	increasing on $(-\infty, 4)$
$4 < x < 5$	$+$	$-$	$+$	$-$	decreasing on $(4, 5)$
$5 < x < 6$	$+$	$-$	$+$	$-$	decreasing on $(5, 6)$
$x > 6$	$+$	$+$	$+$	$+$	increasing on $(6, \infty)$

$x = 5$ is not in the domain of f. f changes from increasing to decreasing at $x = 4$ and from decreasing to increasing at $x = 6$. Thus, $f(4) = 8$ is a local maximum value and $f(6) = 12$ is a local minimum value.

14. $f(x) = x + \dfrac{4}{x^2} = x + 4x^{-2} \quad \Rightarrow \quad f'(x) = 1 - 8x^{-3} = 1 - \dfrac{8}{x^3} = \dfrac{x^3-8}{x^3} = \dfrac{(x-2)(x^2+2x+4)}{x^3}$. The factor $x^2 + 2x + 4$ is always positive and does not affect the sign of $f'(x)$.

Interval	x^3	$x-2$	$f'(x)$	f
$x < 0$	$-$	$-$	$+$	increasing on $(-\infty, 0)$
$0 < x < 2$	$+$	$-$	$-$	decreasing on $(0, 2)$
$x > 2$	$+$	$+$	$+$	increasing on $(2, \infty)$

$x = 0$ is not in the domain of f. f changes from decreasing to increasing at $x = 2$. Thus, $f(2) = 3$ is a local minimum value.

15. $f(x) = \sin x + \cos x,\ 0 \le x \le 2\pi. \quad f'(x) = \cos x - \sin x = 0 \quad \Rightarrow \quad \cos x = \sin x \quad \Rightarrow \quad 1 = \dfrac{\sin x}{\cos x} \quad \Rightarrow$

$\tan x = 1 \quad \Rightarrow \quad x = \frac{\pi}{4}$ or $\frac{5\pi}{4}$. Thus, $f'(x) > 0 \quad \Leftrightarrow \quad \cos x - \sin x > 0 \quad \Leftrightarrow \quad \cos x > \sin x \quad \Leftrightarrow \quad 0 < x < \frac{\pi}{4}$ or

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\frac{5\pi}{4} < x < 2\pi$ and $f'(x) < 0 \quad \Leftrightarrow \quad \cos x < \sin x \quad \Leftrightarrow \quad \frac{\pi}{4} < x < \frac{5\pi}{4}$. So f is increasing on $\left(0, \frac{\pi}{4}\right)$ and $\left(\frac{5\pi}{4}, 2\pi\right)$ and f is decreasing on $\left(\frac{\pi}{4}, \frac{5\pi}{4}\right)$.

f changes from increasing to decreasing at $x = \frac{\pi}{4}$ and from decreasing to increasing at $x = \frac{5\pi}{4}$. Thus, $f\left(\frac{\pi}{4}\right) = \sqrt{2}$ is a local maximum value and $f\left(\frac{5\pi}{4}\right) = -\sqrt{2}$ is a local minimum value.

16. $f(x) = x^4 e^{-x} \quad \Rightarrow \quad f'(x) = x^4(-e^{-x}) + e^{-x}(4x^3) = x^3 e^{-x}(-x+4)$. Thus, $f'(x) > 0$ if $0 < x < 4$ and $f'(x) < 0$ if $x < 0$ or $x > 4$. So f is increasing on $(0, 4)$ and decreasing on $(-\infty, 0)$ and $(4, \infty)$.

f changes from decreasing to increasing at $x = 0$ and from increasing to decreasing at $x = 4$. Thus, $f(0) = 0$ is a local minimum value and $f(4) = 256e^{-4}$ $[\approx 4.69]$ is a local maximum value.

17. $f(x) = x^3 - 3x^2 - 9x + 4 \quad \Rightarrow \quad f'(x) = 3x^2 - 6x - 9 \quad \Rightarrow \quad f''(x) = 6x - 6 = 6(x-1)$. $\quad f''(x) > 0 \quad \Leftrightarrow \quad x > 1$ and $f''(x) < 0 \quad \Leftrightarrow \quad x < 1$. Thus, f is concave upward on $(1, \infty)$ and concave downward on $(-\infty, 1)$. There is an inflection point at $(1, f(1)) = (1, -7)$.

18. $f(x) = 2x^3 - 9x^2 + 12x - 3 \quad \Rightarrow \quad f'(x) = 6x^2 - 18x + 12 \quad \Rightarrow \quad f''(x) = 12x - 18 = 12\left(x - \frac{3}{2}\right)$. $\quad f''(x) > 0 \quad \Leftrightarrow$ $x > \frac{3}{2}$ and $f''(x) < 0 \quad \Leftrightarrow \quad x < \frac{3}{2}$. Thus, f is concave upward on $\left(\frac{3}{2}, \infty\right)$ and concave downward on $\left(-\infty, \frac{3}{2}\right)$. There is an inflection point at $\left(\frac{3}{2}, \frac{3}{2}\right)$.

19. $f(x) = \sin^2 x - \cos 2x,\ 0 \le x \le \pi$. $\quad f'(x) = 2\sin x\,\cos x + 2\sin 2x = \sin 2x + 2\sin 2x = 3\sin 2x$ and $f''(x) = 6\cos 2x$. $\quad f''(x) > 0 \quad \Leftrightarrow \quad \cos 2x > 0 \quad \Leftrightarrow \quad 0 < x < \frac{\pi}{4}$ and $\frac{3\pi}{4} < x < \pi$ and $f''(x) < 0 \quad \Leftrightarrow$ $\cos 2x < 0 \quad \Leftrightarrow \quad \frac{\pi}{4} < x < \frac{3\pi}{4}$. Thus, f is concave upward on $\left(0, \frac{\pi}{4}\right)$ and $\left(\frac{3\pi}{4}, \pi\right)$ and concave downward on $\left(\frac{\pi}{4}, \frac{3\pi}{4}\right)$. There are inflection points at $\left(\frac{\pi}{4}, \frac{1}{2}\right)$ and $\left(\frac{3\pi}{4}, \frac{1}{2}\right)$.

20. $f(x) = \ln(2 + \sin x),\ 0 \le x \le 2\pi$. $\quad f'(x) = \dfrac{1}{2 + \sin x}(\cos x) = \dfrac{\cos x}{2 + \sin x}$ and

$$f''(x) = \frac{(2+\sin x)(-\sin x) - \cos x\,(\cos x)}{(2+\sin x)^2} = \frac{-2\sin x - \sin^2 x - \cos^2 x}{(2+\sin x)^2} = \frac{-2\sin x - (\sin^2 x + \cos^2 x)}{(2+\sin x)^2}$$
$$= \frac{-2\sin x - 1}{(2+\sin x)^2} = -\frac{1 + 2\sin x}{(2+\sin x)^2}.$$

$f''(x) > 0 \quad \Rightarrow \quad 1 + 2\sin x < 0 \quad \Leftrightarrow \quad \sin x < -\frac{1}{2} \quad \Leftrightarrow \quad \frac{7\pi}{6} < x < \frac{11\pi}{6}$ and $f''(x) < 0 \quad \Rightarrow \quad 1 + 2\sin x > 0 \quad \Leftrightarrow$ $\sin x > -\frac{1}{2} \quad \Leftrightarrow \quad 0 < x < \frac{7\pi}{6}$ or $\frac{11\pi}{6} < x < 2\pi$. Thus, f is concave upward on $\left(\frac{7\pi}{6}, \frac{11\pi}{6}\right)$ and concave downward on $\left(0, \frac{7\pi}{6}\right)$ and $\left(\frac{11\pi}{6}, 2\pi\right)$. There are inflection points at $\left(\frac{7\pi}{6}, \ln\frac{3}{2}\right)$ and $\left(\frac{11\pi}{6}, \ln\frac{3}{2}\right)$.

21. $f(x) = \ln(x^2 + 5) \quad \Rightarrow \quad f'(x) = \dfrac{1}{x^2+5}(2x) = \dfrac{2x}{x^2+5}$ and

$$f''(x) = \frac{(x^2+5)(2) - 2x(2x)}{(x^2+5)^2} = \frac{2x^2 + 10 - 4x^2}{(x^2+5)^2} = \frac{-2x^2 + 10}{(x^2+5)^2} = -\frac{2(x^2-5)}{(x^2+5)^2}.$$

$f''(x) > 0 \quad \Rightarrow \quad x^2 - 5 < 0 \quad \Rightarrow \quad x^2 < 5 \quad \Rightarrow \quad -\sqrt{5} < x < \sqrt{5}$ and $f''(x) < 0 \quad \Rightarrow \quad x^2 - 5 > 0 \quad \Rightarrow \quad x^2 > 5 \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x < -\sqrt{5}$ or $x > \sqrt{5}$. Thus, f is concave upward on $\left(-\sqrt{5}, \sqrt{5}\right)$ and concave downward on $\left(-\infty, -\sqrt{5}\right)$ and $\left(\sqrt{5}, \infty\right)$. There are inflection points at $\left(-\sqrt{5}, \ln 10\right)$ and $\left(\sqrt{5}, \ln 10\right)$.

22. $f(x) = \dfrac{e^x}{e^x+2} \quad \Rightarrow \quad f'(x) = \dfrac{(e^x+2)e^x - e^x(e^x)}{(e^x+2)^2} = \dfrac{e^x\left[(e^x+2)-e^x\right]}{(e^x+2)^2} = \dfrac{2e^x}{(e^x+2)^2}$.

$f''(x) = \dfrac{(e^x+2)^2 \cdot 2e^x - 2e^x \cdot 2(e^x+2)e^x}{[(e^x+2)^2]^2} = \dfrac{2e^x(e^x+2)[(e^x+2)-2e^x]}{(e^x+2)^4} = \dfrac{2e^x(2-e^x)}{(e^x+2)^3}$.

$f''(x) > 0 \quad \Leftrightarrow \quad 2 - e^x > 0 \quad \Leftrightarrow \quad e^x < 2 \quad \Leftrightarrow \quad x < \ln 2$ and $f''(x) < 0 \quad \Leftrightarrow \quad 2 - e^x < 0 \quad \Leftrightarrow \quad e^x > 2 \quad \Leftrightarrow$ $x > \ln 2$. Thus, f is concave upward on $(-\infty, \ln 2)$ and concave downward on $(\ln 2, \infty)$. There is an inflection point at $(\ln 2, f(\ln 2)) = \left(\ln 2, \dfrac{2}{2+2}\right) = \left(\ln 2, \frac{1}{2}\right)$.

23. (a) $f(x) = x^4 - 2x^2 + 3 \quad \Rightarrow \quad f'(x) = 4x^3 - 4x = 4x\left(x^2-1\right) = 4x(x+1)(x-1)$.

Interval	$x+1$	$4x$	$x-1$	$f'(x)$	f
$x < -1$	$-$	$-$	$-$	$-$	decreasing on $(-\infty, -1)$
$-1 < x < 0$	$+$	$-$	$-$	$+$	increasing on $(-1, 0)$
$0 < x < 1$	$+$	$+$	$-$	$-$	decreasing on $(0, 1)$
$x > 1$	$+$	$+$	$+$	$+$	increasing on $(1, \infty)$

(b) f changes from increasing to decreasing at $x = 0$ and from decreasing to increasing at $x = -1$ and $x = 1$. Thus, $f(0) = 3$ is a local maximum value and $f(\pm 1) = 2$ are local minimum values.

(c) $f''(x) = 12x^2 - 4 = 12\left(x^2 - \frac{1}{3}\right) = 12\left(x + 1/\sqrt{3}\right)\left(x - 1/\sqrt{3}\right)$. $\quad f''(x) > 0 \quad \Leftrightarrow \quad x < -1/\sqrt{3}$ or $x > 1/\sqrt{3}$ and $f''(x) < 0 \quad \Leftrightarrow \quad -1/\sqrt{3} < x < 1/\sqrt{3}$. Thus, f is concave upward on $\left(-\infty, -\sqrt{3}/3\right)$ and $\left(\sqrt{3}/3, \infty\right)$ and concave downward on $\left(-\sqrt{3}/3, \sqrt{3}/3\right)$. There are inflection points at $\left(\pm\sqrt{3}/3, \frac{22}{9}\right)$.

24. (a) $f(x) = \dfrac{x}{x^2+1} \quad \Rightarrow \quad f'(x) = \dfrac{(x^2+1)(1) - x(2x)}{(x^2+1)^2} = \dfrac{1-x^2}{(x^2+1)^2} = -\dfrac{(x+1)(x-1)}{(x^2+1)^2}$. Thus, $f'(x) > 0$ if $(x+1)(x-1) < 0 \quad \Leftrightarrow \quad -1 < x < 1$, and $f'(x) < 0$ if $x < -1$ or $x > 1$. So f is increasing on $(-1, 1)$ and f is decreasing on $(-\infty, -1)$ and $(1, \infty)$.

(b) f changes from decreasing to increasing at $x = -1$ and from increasing to decreasing at $x = 1$. Thus, $f(-1) = -\frac{1}{2}$ is a local minimum value and $f(1) = \frac{1}{2}$ is a local maximum value.

(c) $f''(x) = \dfrac{(x^2+1)^2(-2x) - (1-x^2)[2(x^2+1)(2x)]}{[(x^2+1)^2]^2} = \dfrac{(x^2+1)(-2x)[(x^2+1)+2(1-x^2)]}{(x^2+1)^4} = \dfrac{2x(x^2-3)}{(x^2+1)^3}$.

$f''(x) > 0 \quad \Leftrightarrow \quad -\sqrt{3} < x < 0$ or $x > \sqrt{3}$, and $f''(x) < 0 \quad \Leftrightarrow \quad x < -\sqrt{3}$ or $0 < x < \sqrt{3}$. Thus, f is concave upward on $\left(-\sqrt{3}, 0\right)$ and $\left(\sqrt{3}, \infty\right)$ and concave downward on $\left(-\infty, -\sqrt{3}\right)$ and $\left(0, \sqrt{3}\right)$. There are inflection points at $\left(-\sqrt{3}, -\sqrt{3}/4\right)$, $(0, 0)$, and $\left(\sqrt{3}, \sqrt{3}/4\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. (a) $f(x) = x^2 - x - \ln x \quad \Rightarrow \quad f'(x) = 2x - 1 - \dfrac{1}{x} = \dfrac{2x^2 - x - 1}{x} = \dfrac{(2x+1)(x-1)}{x}$. Thus, $f'(x) > 0$ if $x > 1$ [note that $x > 0$] and $f'(x) < 0$ if $0 < x < 1$. So f is increasing on $(1, \infty)$ and f is decreasing on $(0, 1)$.

(b) f changes from decreasing to increasing at $x = 1$. Thus, $f(1) = 0$ is a local minimum value.

(c) $f''(x) = 2 + 1/x^2 > 0$ for all x, so f is concave upward on $(0, \infty)$. There is no inflection point.

26. (a) $f(x) = x^2 \ln x \quad \Rightarrow \quad f'(x) = x^2(1/x) + (\ln x)(2x) = x + 2x \ln x = x(1 + 2\ln x)$. The domain of f is $(0, \infty)$, so the sign of f' is determined solely by the factor $1 + 2\ln x$. $f'(x) > 0 \quad \Leftrightarrow \quad \ln x > -\frac{1}{2} \quad \Leftrightarrow \quad x > e^{-1/2}$ $[\approx 0.61]$ and $f'(x) < 0 \quad \Leftrightarrow \quad 0 < x < e^{-1/2}$. So f is increasing on $(e^{-1/2}, \infty)$ and f is decreasing on $(0, e^{-1/2})$.

(b) f changes from decreasing to increasing at $x = e^{-1/2}$. Thus, $f(e^{-1/2}) = (e^{-1/2})^2 \ln(e^{-1/2}) = e^{-1}(-1/2) = -1/(2e)$ $[\approx -0.18]$ is a local minimum value.

(c) $f'(x) = x(1 + 2\ln x) \quad \Rightarrow \quad f''(x) = x(2/x) + (1 + 2\ln x) \cdot 1 = 2 + 1 + 2\ln x = 3 + 2\ln x$. $\quad f''(x) > 0 \quad \Leftrightarrow$ $3 + 2\ln x > 0 \quad \Leftrightarrow \quad \ln x > -3/2 \quad \Leftrightarrow \quad x > e^{-3/2}$ $[\approx 0.22]$. Thus, f is concave upward on $(e^{-3/2}, \infty)$ and f is concave downward on $(0, e^{-3/2})$. $f(e^{-3/2}) = (e^{-3/2})^2 \ln e^{-3/2} = e^{-3}(-3/2) = -3/(2e^3)$ $[\approx -0.07]$. There is a point of inflection at $\left(e^{-3/2}, f(e^{-3/2})\right) = \left(e^{-3/2}, -3/(2e^3)\right)$.

27. (a) $f(x) = xe^{2x} \quad \Rightarrow \quad f'(x) = x(2e^{2x}) + e^{2x}(1) = e^{2x}(2x + 1)$. Thus, $f'(x) > 0$ if $x > -\frac{1}{2}$ and $f'(x) < 0$ if $x < -\frac{1}{2}$. So f is increasing on $\left(-\frac{1}{2}, \infty\right)$ and f is decreasing on $\left(-\infty, -\frac{1}{2}\right)$.

(b) f changes from decreasing to increasing at $x = -\frac{1}{2}$. Thus, $f\left(-\frac{1}{2}\right) = -\frac{1}{2}e^{-1} = -1/(2e)$ $[\approx -0.18]$ is a local minimum value.

(c) $f''(x) = e^{2x}(2) + (2x + 1) \cdot 2e^{2x} = 2e^{2x}[1 + (2x + 1)] = 2e^{2x}(2x + 2) = 4e^{2x}(x + 1)$. $\quad f''(x) > 0 \quad \Leftrightarrow \quad x > -1$ and $f''(x) < 0 \quad \Leftrightarrow \quad x < -1$. Thus, f is concave upward on $(-1, \infty)$ and f is concave downward on $(-\infty, -1)$. There is an inflection point at $\left(-1, -e^{-2}\right)$, or $\left(-1, -1/e^2\right)$.

28. (a) $f(x) = \cos^2 x - 2\sin x, \quad 0 \le x \le 2\pi$. $\quad f'(x) = -2\cos x \sin x - 2\cos x = -2\cos x\,(1 + \sin x)$. Note that $1 + \sin x \ge 0$ [since $\sin x \ge -1$], with equality $\quad \Leftrightarrow \quad \sin x = -1 \quad \Leftrightarrow \quad x = \frac{3\pi}{2}$ [since $0 \le x \le 2\pi$] $\quad \Rightarrow$ $\cos x = 0$. Thus, $f'(x) > 0 \quad \Leftrightarrow \quad \cos x < 0 \quad \Leftrightarrow \quad \frac{\pi}{2} < x < \frac{3\pi}{2}$ and $f'(x) < 0 \quad \Leftrightarrow \quad \cos x > 0 \quad \Leftrightarrow \quad 0 < x < \frac{\pi}{2}$ or $\frac{3\pi}{2} < x < 2\pi$. Thus, f is increasing on $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$ and f is decreasing on $\left(0, \frac{\pi}{2}\right)$ and $\left(\frac{3\pi}{2}, 2\pi\right)$.

(b) f changes from decreasing to increasing at $x = \frac{\pi}{2}$ and from increasing to decreasing at $x = \frac{3\pi}{2}$. Thus, $f\left(\frac{\pi}{2}\right) = -2$ is a local minimum value and $f\left(\frac{3\pi}{2}\right) = 2$ is a local maximum value.

(c) $f''(x) = 2\sin x\,(1 + \sin x) - 2\cos^2 x = 2\sin x + 2\sin^2 x - 2(1 - \sin^2 x)$

$$= 4\sin^2 x + 2\sin x - 2 = 2(2\sin x - 1)(\sin x + 1)$$

so $f''(x) > 0 \quad \Leftrightarrow \quad \sin x > \frac{1}{2} \quad \Leftrightarrow \quad \frac{\pi}{6} < x < \frac{5\pi}{6}$, and $f''(x) < 0 \quad \Leftrightarrow \quad \sin x < \frac{1}{2}$ and $\sin x \ne -1 \quad \Leftrightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$0 < x < \frac{\pi}{6}$ or $\frac{5\pi}{6} < x < \frac{3\pi}{2}$ or $\frac{3\pi}{2} < x < 2\pi$. Thus, f is concave upward on $\left(\frac{\pi}{6}, \frac{5\pi}{6}\right)$ and concave downward on $\left(0, \frac{\pi}{6}\right)$, $\left(\frac{5\pi}{6}, \frac{3\pi}{2}\right)$, and $\left(\frac{3\pi}{2}, 2\pi\right)$. There are inflection points at $\left(\frac{\pi}{6}, -\frac{1}{4}\right)$ and $\left(\frac{5\pi}{6}, -\frac{1}{4}\right)$.

29. $f(x) = 1 + 3x^2 - 2x^3 \quad \Rightarrow \quad f'(x) = 6x - 6x^2 = 6x(1-x)$.

First Derivative Test: $f'(x) > 0 \quad \Rightarrow \quad 0 < x < 1$ and $f'(x) < 0 \quad \Rightarrow \quad x < 0$ or $x > 1$. Since f' changes from negative to positive at $x = 0$, $f(0) = 1$ is a local minimum value; and since f' changes from positive to negative at $x = 1$, $f(1) = 2$ is a local maximum value.

Second Derivative Test: $f''(x) = 6 - 12x$. $f'(x) = 0 \quad \Leftrightarrow \quad x = 0, 1$. $f''(0) = 6 > 0 \quad \Rightarrow \quad f(0) = 1$ is a local minimum value. $f''(1) = -6 < 0 \quad \Rightarrow \quad f(1) = 2$ is a local maximum value.

Preference: For this function, the two tests are equally easy.

30. $f(x) = \dfrac{x^2}{x-1} \quad \Rightarrow \quad f'(x) = \dfrac{(x-1)(2x) - x^2(1)}{(x-1)^2} = \dfrac{x^2 - 2x}{(x-1)^2} = \dfrac{x(x-2)}{(x-1)^2}$.

First Derivative Test: $f'(x) > 0 \quad \Rightarrow \quad x < 0$ or $x > 2$ and $f'(x) < 0 \quad \Rightarrow \quad 0 < x < 1$ or $1 < x < 2$. Since f' changes from positive to negative at $x = 0$, $f(0) = 0$ is a local maximum value; and since f' changes from negative to positive at $x = 2$, $f(2) = 4$ is a local minimum value.

Second Derivative Test:

$$f''(x) = \frac{(x-1)^2(2x-2) - (x^2-2x)2(x-1)}{[(x-1)^2]^2} = \frac{2(x-1)[(x-1)^2 - (x^2-2x)]}{(x-1)^4} = \frac{2}{(x-1)^3}.$$

$f'(x) = 0 \quad \Leftrightarrow \quad x = 0, 2$. $f''(0) = -2 < 0 \quad \Rightarrow \quad f(0) = 0$ is a local maximum value. $f''(2) = 2 > 0 \quad \Rightarrow \quad f(2) = 4$ is a local minimum value.

Preference: Since calculating the second derivative is fairly difficult, the First Derivative Test is easier to use for this function.

31. $f'(x) = (x-4)^2(x+3)^7(x-5)^8$. The factors $(x-4)^2$ and $(x-5)^8$ are nonnegative. Hence, the sign of f' is determined by the sign of $(x+3)^7$, which is positive for $x > -3$. Thus, f increases on the intervals $(-3, 4)$, $(4, 5)$, and $(5, \infty)$. Note that f is differentiable and $f'(x) > 0$ on the interval $(-3, \infty)$ except for the numbers $x = 4$ and $x = 5$. By applying the result of Exercise 4.3.97, we can say that f is increasing on the entire interval $(-3, \infty)$.

32. (a) $f(x) = x^4(x-1)^3 \quad \Rightarrow \quad f'(x) = x^4 \cdot 3(x-1)^2 + (x-1)^3 \cdot 4x^3 = x^3(x-1)^2\,[3x + 4(x-1)] = x^3(x-1)^2(7x-4)$

The critical numbers are 0, 1, and $\frac{4}{7}$.

(b) $f''(x) = 3x^2(x-1)^2(7x-4) + x^3 \cdot 2(x-1)(7x-4) + x^3(x-1)^2 \cdot 7$

$= x^2(x-1)\,[3(x-1)(7x-4) + 2x(7x-4) + 7x(x-1)]$

Now $f''(0) = f''(1) = 0$, so the Second Derivative Test gives no information for $x = 0$ or $x = 1$.

$f''\left(\frac{4}{7}\right) = \left(\frac{4}{7}\right)^2\left(\frac{4}{7} - 1\right)\left[0 + 0 + 7\left(\frac{4}{7}\right)\left(\frac{4}{7} - 1\right)\right] = \left(\frac{4}{7}\right)^2\left(-\frac{3}{7}\right)(4)\left(-\frac{3}{7}\right) > 0$, so there is a local minimum at $x = \frac{4}{7}$.

(c) f' is positive on $(-\infty, 0)$, negative on $\left(0, \frac{4}{7}\right)$, positive on $\left(\frac{4}{7}, 1\right)$, and positive on $(1, \infty)$. So f has a local maximum at $x = 0$, a local minimum at $x = \frac{4}{7}$, and no local maximum or minimum at $x = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. (a) By the Second Derivative Test, if $f'(2) = 0$ and $f''(2) = -5 < 0$, f has a local maximum at $x = 2$.

(b) If $f'(6) = 0$, we know that f has a horizontal tangent at $x = 6$. Knowing that $f''(6) = 0$ does not provide any additional information since the Second Derivative Test fails. For example, the first and second derivatives of $y = (x-6)^4$, $y = -(x-6)^4$, and $y = (x-6)^3$ all equal zero for $x = 6$, but the first has a local minimum at $x = 6$, the second has a local maximum at $x = 6$, and the third has an inflection point at $x = 6$.

34. (a) $f'(x) < 0$ and $f''(x) < 0$ for all x

The function must be always decreasing (since the first derivative is always negative) and concave downward (since the second derivative is always negative).

(b) $f'(x) > 0$ and $f''(x) > 0$ for all x

The function must be always increasing (since the first derivative is always positive) and concave upward (since the second derivative is always positive).

35. (a) $f'(x) > 0$ and $f''(x) < 0$ for all x

The function must be always increasing (since the first derivative is always positive) and concave downward (since the second derivative is always negative).

(b) $f'(x) < 0$ and $f''(x) > 0$ for all x

The function must be always decreasing (since the first derivative is always negative) and concave upward (since the second derivative is always positive).

36. Vertical asymptote $x = 0$

$f'(x) > 0$ if $x < -2 \quad \Rightarrow \quad f$ is increasing on $(-\infty, -2)$.

$f'(x) < 0$ if $x > -2$ $(x \neq 0) \quad \Rightarrow \quad f$ is decreasing on $(-2, 0)$ and $(0, \infty)$.

$f''(x) < 0$ if $x < 0 \quad \Rightarrow \quad f$ is concave downward on $(-\infty, 0)$.

$f''(x) > 0$ if $x > 0 \quad \Rightarrow \quad f$ is concave upward on $(0, \infty)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. $f'(0) = f'(2) = f'(4) = 0 \quad \Rightarrow \quad$ horizontal tangents at $x = 0, 2, 4$.

$f'(x) > 0$ if $x < 0$ or $2 < x < 4 \quad \Rightarrow \quad f$ is increasing on $(-\infty, 0)$ and $(2, 4)$.

$f'(x) < 0$ if $0 < x < 2$ or $x > 4 \quad \Rightarrow \quad f$ is decreasing on $(0, 2)$ and $(4, \infty)$.

$f''(x) > 0$ if $1 < x < 3 \quad \Rightarrow \quad f$ is concave upward on $(1, 3)$.

$f''(x) < 0$ if $x < 1$ or $x > 3 \quad \Rightarrow \quad f$ is concave downward on $(-\infty, 1)$ and $(3, \infty)$. There are inflection points when $x = 1$ and 3.

38. $f'(x) > 0$ for all $x \neq 1 \quad \Rightarrow \quad f$ is increasing on $(-\infty, 1)$ and $(1, \infty)$.

Vertical asymptote $x = 1$

$f''(x) > 0$ if $x < 1$ or $x > 3 \quad \Rightarrow \quad f$ is concave upward on $(-\infty, 1)$ and $(3, \infty)$.

$f''(x) < 0$ if $1 < x < 3 \quad \Rightarrow \quad f$ is concave downward on $(1, 3)$.

There is an inflection point at $x = 3$.

39. $f'(5) = 0 \quad \Rightarrow \quad$ horizontal tangent at $x = 5$.

$f'(x) < 0$ when $x < 5 \quad \Rightarrow \quad f$ is decreasing on $(-\infty, 5)$.

$f'(x) > 0$ when $x > 5 \quad \Rightarrow \quad f$ is increasing on $(5, \infty)$.

$f''(2) = 0, \quad f''(8) = 0, \quad f''(x) < 0$ when $x < 2$ or $x > 8$,

$f''(x) > 0$ for $2 < x < 8 \quad \Rightarrow \quad f$ is concave upward on $(2, 8)$ and concave downward on $(-\infty, 2)$ and $(8, \infty)$.

There are inflection points at $x = 2$ and $x = 8$.

$\lim\limits_{x \to \infty} f(x) = 3, \quad \lim\limits_{x \to -\infty} f(x) = 3 \quad \Rightarrow \quad y = 3$ is a horizontal asymptote.

40. $f'(0) = f'(4) = 0 \quad \Rightarrow \quad$ horizontal tangents at $x = 0$ and 4.

$f'(x) = 1$ if $x < -1 \quad \Rightarrow \quad f$ is a line with slope 1 on $(-\infty, -1)$.

$f'(x) > 0$ if $0 < x < 2 \quad \Rightarrow \quad f$ is increasing on $(0, 2)$.

$f'(x) < 0$ if $-1 < x < 0$ or $2 < x < 4$ or $x > 4 \quad \Rightarrow \quad f$ is decreasing on $(-1, 0)$, $(2, 4)$, and $(4, \infty)$.

$\lim\limits_{x \to 2^-} f'(x) = \infty \quad \Rightarrow \quad f'$ increases without bound as $x \to 2^-$.

$\lim\limits_{x \to 2^+} f'(x) = -\infty \quad \Rightarrow \quad f'$ decreases without bound as $x \to 2^+$.

$f''(x) > 0$ if $-1 < x < 2$ or $2 < x < 4 \quad \Rightarrow \quad f$ is concave upward on $(-1, 2)$ and $(2, 4)$.

$f''(x) < 0$ if $x > 4 \quad \Rightarrow \quad f$ is concave downward on $(4, \infty)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. $f'(x) > 0$ if $x \neq 2 \quad \Rightarrow \quad f$ is increasing on $(-\infty, 2)$ and $(2, \infty)$.

$f''(x) > 0$ if $x < 2 \quad \Rightarrow \quad f$ is concave upward on $(-\infty, 2)$.

$f''(x) < 0$ if $x > 2 \quad \Rightarrow \quad f$ is concave downward on $(2, \infty)$.

f has inflection point $(2, 5) \quad \Rightarrow \quad f$ changes concavity at the point $(2, 5)$.

$\lim\limits_{x \to \infty} f(x) = 8 \quad \Rightarrow \quad f$ has a horizontal asymptote of $y = 8$ as $x \to \infty$.

$\lim\limits_{x \to -\infty} f(x) = 0 \quad \Rightarrow \quad f$ has a horizontal asymptote of $y = 0$ as $x \to -\infty$.

42. (a) $\dfrac{dy}{dx} > 0$ (f is increasing) and $\dfrac{d^2y}{dx^2} > 0$ (f is concave upward) at point B.

(b) $\dfrac{dy}{dx} < 0$ (f is decreasing) and $\dfrac{d^2y}{dx^2} < 0$ (f is concave downward) at point E.

(c) $\dfrac{dy}{dx} < 0$ (f is decreasing) and $\dfrac{d^2y}{dx^2} > 0$ (f is concave upward) at point A.

Note: At C, $\dfrac{dy}{dx} > 0$ and $\dfrac{d^2y}{dx^2} < 0$. At D, $\dfrac{dy}{dx} = 0$ and $\dfrac{d^2y}{dx^2} \leq 0$.

43. (a) f is increasing where f' is positive, that is, on $(0, 2)$, $(4, 6)$, and $(8, \infty)$; and decreasing where f' is negative, that is, on $(2, 4)$ and $(6, 8)$.

(b) f has local maxima where f' changes from positive to negative, at $x = 2$ and at $x = 6$, and local minima where f' changes from negative to positive, at $x = 4$ and at $x = 8$.

(c) f is concave upward (CU) where f' is increasing, that is, on $(3, 6)$ and $(6, \infty)$, and concave downward (CD) where f' is decreasing, that is, on $(0, 3)$.

(d) There is a point of inflection where f changes from being CD to being CU, that is, at $x = 3$.

(e)

44. (a) f is increasing where f' is positive, on $(1, 6)$ and $(8, \infty)$, and decreasing where f' is negative, on $(0, 1)$ and $(6, 8)$.

(b) f has a local maximum where f' changes from positive to negative, at $x = 6$, and local minima where f' changes from negative to positive, at $x = 1$ and at $x = 8$.

(c) f is concave upward where f' is increasing, that is, on $(0, 2)$, $(3, 5)$, and $(7, \infty)$, and concave downward where f' is decreasing, that is, on $(2, 3)$ and $(5, 7)$.

(d) There are points of inflection where f changes its direction of concavity, at $x = 2$, $x = 3$, $x = 5$ and $x = 7$.

(e)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. (a) $f(x) = x^3 - 3x^2 + 4 \quad \Rightarrow \quad f'(x) = 3x^2 - 6x = 3x(x-2)$.

Interval	$3x$	$x-2$	$f'(x)$	f
$x < 0$	$-$	$-$	$+$	increasing on $(-\infty, 0)$
$0 < x < 2$	$+$	$-$	$-$	decreasing on $(0, 2)$
$x > 2$	$+$	$+$	$+$	increasing on $(2, \infty)$

(b) f changes from increasing to decreasing at $x = 0$ and from decreasing to increasing at $x = 2$. Thus, $f(0) = 4$ is a local maximum value and $f(2) = 0$ is a local minimum value.

(c) $f''(x) = 6x - 6 = 6(x-1)$. $\quad f''(x) = 0 \quad \Leftrightarrow \quad x = 1$. $\quad f''(x) > 0$ on $(1, \infty)$ and $f''(x) < 0$ on $(-\infty, 1)$. So f is concave upward on $(1, \infty)$ and f is concave downward on $(-\infty, 1)$. There is an inflection point at $(1, 2)$.

(d)

46. (a) $f(x) = 36x + 3x^2 - 2x^3 \quad \Rightarrow \quad f'(x) = 36 + 6x - 6x^2 = -6(x^2 - x - 6) = -6(x+2)(x-3)$. $f'(x) > 0 \quad \Leftrightarrow$ $-2 < x < 3$ and $f'(x) < 0 \quad \Leftrightarrow \quad x < -2$ or $x > 3$. So f is increasing on $(-2, 3)$ and f is decreasing on $(-\infty, -2)$ and $(3, \infty)$.

(b) f changes from increasing to decreasing at $x = 3$, so $f(3) = 81$ is a local maximum value. f changes from decreasing to increasing at $x = -2$, so $f(-2) = -44$ is a local minimum value.

(c) $f''(x) = 6 - 12x$. $\quad f''(x) = 0 \quad \Leftrightarrow \quad x = \frac{1}{2}$. $f''(x) > 0$ on $\left(-\infty, \frac{1}{2}\right)$ and $f''(x) < 0$ on $\left(\frac{1}{2}, \infty\right)$. So f is CU on $\left(-\infty, \frac{1}{2}\right)$ and f is CD on $\left(\frac{1}{2}, \infty\right)$. There is an inflection point at $\left(\frac{1}{2}, \frac{37}{2}\right)$.

(d)

47. (a) $f(x) = \frac{1}{2}x^4 - 4x^2 + 3 \quad \Rightarrow \quad f'(x) = 2x^3 - 8x = 2x(x^2 - 4) = 2x(x+2)(x-2)$. $\quad f'(x) > 0 \quad \Leftrightarrow \quad -2 < x < 0$ or $x > 2$, and $f'(x) < 0 \quad \Leftrightarrow \quad x < -2$ or $0 < x < 2$. So f is increasing on $(-2, 0)$ and $(2, \infty)$ and f is decreasing on $(-\infty, -2)$ and $(0, 2)$.

(b) f changes from increasing to decreasing at $x = 0$, so $f(0) = 3$ is a local maximum value.

f changes from decreasing to increasing at $x = \pm 2$, so $f(\pm 2) = -5$ is a local minimum value.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $f''(x) = 6x^2 - 8 = 6\left(x^2 - \frac{4}{3}\right) = 6\left(x + \frac{2}{\sqrt{3}}\right)\left(x - \frac{2}{\sqrt{3}}\right)$.

$f''(x) = 0 \quad \Leftrightarrow \quad x = \pm\frac{2}{\sqrt{3}}$. $f''(x) > 0$ on $\left(-\infty, -\frac{2}{\sqrt{3}}\right)$ and $\left(\frac{2}{\sqrt{3}}, \infty\right)$ and $f''(x) < 0$ on $\left(-\frac{2}{\sqrt{3}}, \frac{2}{\sqrt{3}}\right)$. So f is CU on $\left(-\infty, -\frac{2}{\sqrt{3}}\right)$ and $\left(\frac{2}{\sqrt{3}}, \infty\right)$, and f is CD on $\left(-\frac{2}{\sqrt{3}}, \frac{2}{\sqrt{3}}\right)$. There are inflection points at $\left(\pm\frac{2}{\sqrt{3}}, -\frac{13}{9}\right)$.

(d)

48. (a) $g(x) = 200 + 8x^3 + x^4 \quad \Rightarrow \quad g'(x) = 24x^2 + 4x^3 = 4x^2(6 + x) = 0$ when $x = -6$ and when $x = 0$.

$g'(x) > 0 \quad \Leftrightarrow \quad x > -6 \;\; [x \neq 0]$ and $g'(x) < 0 \quad \Leftrightarrow \quad x < -6$, so g is decreasing on $(-\infty, -6)$ and g is increasing on $(-6, \infty)$, with a horizontal tangent at $x = 0$.

(b) $g(-6) = -232$ is a local minimum value. There is no local maximum value.

(c) $g''(x) = 48x + 12x^2 = 12x(4 + x) = 0$ when $x = -4$ and when $x = 0$.

$g''(x) > 0 \quad \Leftrightarrow \quad x < -4$ or $x > 0$ and $g''(x) < 0 \quad \Leftrightarrow \quad -4 < x < 0$, so g is CU on $(-\infty, -4)$ and $(0, \infty)$, and g is CD on $(-4, 0)$. There are inflection points at $(-4, -56)$ and $(0, 200)$.

(d)

49. (a) $g(t) = 3t^4 - 8t^3 + 12 \quad \Rightarrow \quad g'(t) = 12t^3 - 24t^2 = 12t^2(t - 2)$.

Interval	$12t^2$	$t - 2$	$g'(t)$	g
$t < 0$	$+$	$-$	$-$	decreasing on $(-\infty, 0)$
$0 < t < 2$	$+$	$-$	$-$	decreasing on $(0, 2)$
$t > 2$	$+$	$+$	$+$	increasing on $(2, \infty)$

(b) g changes from decreasing to increasing at $x = 2$. Thus, $g(2) = -4$ is a local minimum value.

(c) $g''(t) = 36t^2 - 48t = 12t(3t - 4)$. $g''(t) = 0 \quad \Leftrightarrow \quad t = 0$ or $t = \frac{4}{3}$.

Interval	$12t$	$3t - 4$	$g''(t)$	g
$t < 0$	$-$	$-$	$+$	concave up on $(-\infty, 0)$
$0 < t < \frac{4}{3}$	$+$	$-$	$-$	concave down on $\left(0, \frac{4}{3}\right)$
$t > \frac{4}{3}$	$+$	$+$	$+$	concave up on $\left(\frac{4}{3}, \infty\right)$

There are inflection points at $(0, 12)$ and $\left(\frac{4}{3}, \frac{68}{27}\right)$.

(d)

50. (a) $h(x) = 5x^3 - 3x^5 \quad \Rightarrow \quad h'(x) = 15x^2 - 15x^4 = 15x^2(1 - x^2) = 15x^2(1 + x)(1 - x)$. $h'(x) > 0 \quad \Leftrightarrow$ $-1 < x < 0$ and $0 < x < 1$ [note that $h'(0) = 0$] and $h'(x) < 0 \quad \Leftrightarrow \quad x < -1$ or $x > 1$. So h is increasing on $(-1, 1)$ and h is decreasing on $(-\infty, -1)$ and $(1, \infty)$.

(b) h changes from decreasing to increasing at $x = -1$, so $h(-1) = -2$ is a local minimum value. h changes from increasing to decreasing at $x = 1$, so $h(1) = 2$ is a local maximum value.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $h''(x) = 30x - 60x^3 = 30x(1 - 2x^2)$. $h''(x) = 0 \;\Leftrightarrow\; x = 0$ or $1 - 2x^2 = 0 \;\Leftrightarrow\; x = 0$ or $x = \pm 1/\sqrt{2}$. $h''(x) > 0$ on $\left(-\infty, -1/\sqrt{2}\right)$ and $\left(0, 1/\sqrt{2}\right)$, and $h''(x) < 0$ on $\left(-1/\sqrt{2}, 0\right)$ and $\left(1/\sqrt{2}, \infty\right)$. So h is CU on $\left(-\infty, -1/\sqrt{2}\right)$ and $\left(0, 1/\sqrt{2}\right)$, and h is CD on $\left(-1/\sqrt{2}, 0\right)$ and $\left(1/\sqrt{2}, \infty\right)$. There are inflection points at $\left(-1/\sqrt{2}, -7/\left(4\sqrt{2}\right)\right)$, $(0, 0)$, and $\left(1/\sqrt{2}, 7/\left(4\sqrt{2}\right)\right)$.

(d)

51. (a) $f(z) = z^7 - 112z^2 \;\Rightarrow\; f'(z) = 7z^6 - 224z = 7z(z^5 - 32)$. $f'(z) = 0 \;\Rightarrow\; z = 0, 2$.

Interval	$7z$	$z^5 - 32$	$f'(z)$	f
$z < 0$	$-$	$-$	$+$	increasing on $(-\infty, 0)$
$0 < z < 2$	$+$	$-$	$-$	decreasing on $(0, 2)$
$z > 2$	$+$	$+$	$+$	increasing on $(2, \infty)$

(b) f changes from increasing to decreasing at $x = 0$ and from decreasing to increasing at $x = 2$. Thus, $f(0) = 0$ is a local maximum value and $f(2) = -320$ is a local minimum value.

(c) $f''(z) = 42z^5 - 224 = 14\left(3z^5 - 16\right)$. $f''(z) = 0 \;\Leftrightarrow\; 3z^5 = 16 \;\Leftrightarrow\; z^5 = \frac{16}{3} \;\Leftrightarrow\; z = \sqrt[5]{\frac{16}{3}}$ [call this value a]. $f''(z) > 0 \;\Leftrightarrow\; z > a$ and $f''(z) < 0 \;\Leftrightarrow\; z < a$. So, f is concave up on (a, ∞) and concave down on $(-\infty, a)$. There is an inflection point at $(a, f(a)) = \left(\sqrt[5]{\frac{16}{3}}, -\frac{320}{3}\sqrt[5]{\frac{256}{9}}\right) \approx (1.398, -208.4)$.

(d)

52. (a) $f(x) = \left(x^2 - 4\right)^3 \;\Rightarrow\; f'(x) = 3(x^2 - 4)^2(2x) = 6x(x^2 - 4)^2$. Since $(x^2 - 4)^2$ is nonnegative, the sign of $f'(x)$ is determined by the sign of $6x$. Thus, $f'(x) < 0 \;\Leftrightarrow\; x < 0$ $[x \neq -2]$ and $f'(x) > 0 \;\Leftrightarrow\; x > 0$ $[x \neq 2]$. So f is increasing on $(0, 2)$ and $(2, \infty)$, and f is decreasing on $(-\infty, -2)$ and $(-2, 0)$. By Exercise 4.3.97, we can say that f is increasing on $(0, \infty)$ and decreasing on $(-\infty, 0)$.

(b) f changes from decreasing to increasing at $x = 0$. Thus, $f(0) = -64$ is a local minimum value.

(c) $f''(x) = 6x \cdot 2(x^2 - 4)(2x) + (x^2 - 4)^2 \cdot 6 = 6(x^2 - 4)[4x^2 + (x^2 - 4)]$
$= 6(x^2 - 4)(5x^2 - 4) = 6(x + 2)(x - 2)\left(\sqrt{5}\,x + 2\right)\left(\sqrt{5}\,x - 2\right)$

Interval	$x + 2$	$\sqrt{5}\,x + 2$	$\sqrt{5}\,x - 2$	$x - 2$	$f''(x)$	f
$x < -2$	$-$	$-$	$-$	$-$	$+$	concave up on $(-\infty, -2)$
$-2 < x < -\frac{2}{\sqrt{5}}$	$+$	$-$	$-$	$-$	$-$	concave down on $\left(-2, -\frac{2}{\sqrt{5}}\right)$
$-\frac{2}{\sqrt{5}} < x < \frac{2}{\sqrt{5}}$	$+$	$+$	$-$	$-$	$+$	concave up on $\left(-\frac{2}{\sqrt{5}}, \frac{2}{\sqrt{5}}\right)$
$\frac{2}{\sqrt{5}} < x < 2$	$+$	$+$	$+$	$-$	$-$	concave down on $\left(\frac{2}{\sqrt{5}}, 2\right)$
$x > 2$	$+$	$+$	$+$	$+$	$+$	concave up on $(2, \infty)$

There are inflection points at $(-2, 0)$, $\left(-\dfrac{2}{\sqrt{5}}, -\dfrac{4096}{125}\right)$, $\left(\dfrac{2}{\sqrt{5}}, -\dfrac{4096}{125}\right)$, and $(2, 0)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d)

53. (a) $F(x) = x\sqrt{6-x} \quad \Rightarrow$

$$F'(x) = x \cdot \tfrac{1}{2}(6-x)^{-1/2}(-1) + (6-x)^{1/2}(1) = \tfrac{1}{2}(6-x)^{-1/2}[-x + 2(6-x)] = \frac{-3x+12}{2\sqrt{6-x}}.$$

$F'(x) > 0 \quad \Leftrightarrow \quad -3x + 12 > 0 \quad \Leftrightarrow \quad x < 4$ and $F'(x) < 0 \quad \Leftrightarrow \quad 4 < x < 6$. So F is increasing on $(-\infty, 4)$ and F is decreasing on $(4, 6)$.

(b) F changes from increasing to decreasing at $x = 4$, so $F(4) = 4\sqrt{2}$ is a local maximum value. There is no local minimum value.

(c) $F'(x) = -\frac{3}{2}(x-4)(6-x)^{-1/2} \quad \Rightarrow$

$$F''(x) = -\tfrac{3}{2}\left[(x-4)\left(-\tfrac{1}{2}(6-x)^{-3/2}(-1)\right) + (6-x)^{-1/2}(1)\right]$$

$$= -\frac{3}{2} \cdot \frac{1}{2}(6-x)^{-3/2}[(x-4) + 2(6-x)] = \frac{3(x-8)}{4(6-x)^{3/2}}$$

$F''(x) < 0$ on $(-\infty, 6)$, so F is CD on $(-\infty, 6)$. There is no inflection point.

(d)

54. (a) $G(x) = 5x^{2/3} - 2x^{5/3} \quad \Rightarrow \quad G'(x) = \frac{10}{3}x^{-1/3} - \frac{10}{3}x^{2/3} = \frac{10}{3}x^{-1/3}(1-x) = \dfrac{10(1-x)}{3x^{1/3}}$.

$G'(x) > 0 \quad \Leftrightarrow \quad 0 < x < 1$ and $G'(x) < 0 \quad \Leftrightarrow \quad x < 0$ or $x > 1$. So G is increasing on $(0, 1)$ and G is decreasing on $(-\infty, 0)$ and $(1, \infty)$.

(b) G changes from decreasing to increasing at $x = 0$, so $G(0) = 0$ is a local minimum value. G changes from increasing to decreasing at $x = 1$, so $G(1) = 3$ is a local maximum value. Note that the First Derivative Test applies at $x = 0$ even though G' is not defined at $x = 0$, since G is continuous at 0.

(c) $G''(x) = -\frac{10}{9}x^{-4/3} - \frac{20}{9}x^{-1/3} = -\frac{10}{9}x^{-4/3}(1+2x)$. $\quad G''(x) > 0 \quad \Leftrightarrow$ $x < -\frac{1}{2}$ and $G''(x) < 0 \quad \Leftrightarrow \quad -\frac{1}{2} < x < 0$ or $x > 0$. So G is CU on $\left(-\infty, -\frac{1}{2}\right)$ and G is CD on $\left(-\frac{1}{2}, 0\right)$ and $(0, \infty)$. The only change in concavity occurs at $x = -\frac{1}{2}$, so there is an inflection point at $\left(-\frac{1}{2}, 6/\sqrt[3]{4}\right)$.

(d)

55. (a) $C(x) = x^{1/3}(x+4) = x^{4/3} + 4x^{1/3} \quad \Rightarrow \quad C'(x) = \frac{4}{3}x^{1/3} + \frac{4}{3}x^{-2/3} = \frac{4}{3}x^{-2/3}(x+1) = \dfrac{4(x+1)}{3\sqrt[3]{x^2}}$. $C'(x) > 0$ if $-1 < x < 0$ or $x > 0$ and $C'(x) < 0$ for $x < -1$, so C is increasing on $(-1, \infty)$ and C is decreasing on $(-\infty, -1)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $C(-1) = -3$ is a local minimum value.

(c) $C''(x) = \frac{4}{9}x^{-2/3} - \frac{8}{9}x^{-5/3} = \frac{4}{9}x^{-5/3}(x-2) = \dfrac{4(x-2)}{9\sqrt[3]{x^5}}$.

$C''(x) < 0$ for $0 < x < 2$ and $C''(x) > 0$ for $x < 0$ and $x > 2$, so C is concave downward on $(0, 2)$ and concave upward on $(-\infty, 0)$ and $(2, \infty)$. There are inflection points at $(0, 0)$ and $\left(2, 6\sqrt[3]{2}\right) \approx (2, 7.56)$.

(d)

56. (a) $f(x) = \ln(x^2+9) \quad \Rightarrow \quad f'(x) = \dfrac{1}{x^2+9} \cdot 2x = \dfrac{2x}{x^2+9}$. $\quad f'(x) > 0 \quad \Leftrightarrow \quad 2x > 0 \quad \Leftrightarrow \quad x > 0$ and $f'(x) < 0 \quad \Leftrightarrow$ $x < 0$. So f is increasing on $(0, \infty)$ and f is decreasing on $(-\infty, 0)$.

(b) f changes from decreasing to increasing at $x = 0$, so $f(0) = \ln 9$ is a local minimum value. There is no local maximum value.

(c) $f''(x) = \dfrac{(x^2+9)\cdot 2 - 2x(2x)}{(x^2+9)^2} = \dfrac{18-2x^2}{(x^2+9)^2} = \dfrac{-2(x+3)(x-3)}{(x^2+9)^2}$.

$f''(x) = 0 \quad \Leftrightarrow \quad x = \pm 3$. $f''(x) > 0$ on $(-3, 3)$ and $f''(x) < 0$ on $(-\infty, -3)$ and $(3, \infty)$. So f is CU on $(-3, 3)$, and f is CD on $(-\infty, -3)$ and $(3, \infty)$. There are inflection points at $(\pm 3, \ln 18)$.

(d)

57. (a) $f(\theta) = 2\cos\theta + \cos^2\theta,\ 0 \le \theta \le 2\pi \quad \Rightarrow \quad f'(\theta) = -2\sin\theta + 2\cos\theta\,(-\sin\theta) = -2\sin\theta\,(1+\cos\theta)$.

$f'(\theta) = 0 \quad \Leftrightarrow \quad \theta = 0, \pi$, and 2π. $f'(\theta) > 0 \quad \Leftrightarrow \quad \pi < \theta < 2\pi$ and $f'(\theta) < 0 \quad \Leftrightarrow \quad 0 < \theta < \pi$. So f is increasing on $(\pi, 2\pi)$ and f is decreasing on $(0, \pi)$.

(b) $f(\pi) = -1$ is a local minimum value.

(c) $f'(\theta) = -2\sin\theta\,(1+\cos\theta) \quad \Rightarrow$

$$\begin{aligned} f''(\theta) &= -2\sin\theta\,(-\sin\theta) + (1+\cos\theta)(-2\cos\theta) = 2\sin^2\theta - 2\cos\theta - 2\cos^2\theta \\ &= 2(1-\cos^2\theta) - 2\cos\theta - 2\cos^2\theta = -4\cos^2\theta - 2\cos\theta + 2 \\ &= -2(2\cos^2\theta + \cos\theta - 1) = -2(2\cos\theta - 1)(\cos\theta + 1) \end{aligned}$$

Since $-2(\cos\theta + 1) < 0$ [for $\theta \ne \pi$], $f''(\theta) > 0 \quad \Rightarrow \quad 2\cos\theta - 1 < 0 \quad \Rightarrow \quad \cos\theta < \frac{1}{2} \quad \Rightarrow \quad \frac{\pi}{3} < \theta < \frac{5\pi}{3}$ and $f''(\theta) < 0 \quad \Rightarrow \quad \cos\theta > \frac{1}{2} \quad \Rightarrow \quad 0 < \theta < \frac{\pi}{3}$ or $\frac{5\pi}{3} < \theta < 2\pi$. So f is CU on $\left(\frac{\pi}{3}, \frac{5\pi}{3}\right)$ and f is CD on $\left(0, \frac{\pi}{3}\right)$ and $\left(\frac{5\pi}{3}, 2\pi\right)$. There are points of inflection at $\left(\frac{\pi}{3}, f\left(\frac{\pi}{3}\right)\right) = \left(\frac{\pi}{3}, \frac{5}{4}\right)$ and $\left(\frac{5\pi}{3}, f\left(\frac{5\pi}{3}\right)\right) = \left(\frac{5\pi}{3}, \frac{5}{4}\right)$.

(d)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

58. (a) $S(x) = x - \sin x$, $0 \le x \le 4\pi \quad \Rightarrow \quad S'(x) = 1 - \cos x$. $S'(x) = 0 \quad \Leftrightarrow \quad \cos x = 1 \quad \Leftrightarrow \quad x = 0, 2\pi$, and 4π.

$S'(x) > 0 \quad \Leftrightarrow \quad \cos x < 1$, which is true for all x except integer multiples of 2π, so S is increasing on $(0, 4\pi)$ since $S'(2\pi) = 0$.

(b) There is no local maximum or minimum.

(c) $S''(x) = \sin x$. $S''(x) > 0$ if $0 < x < \pi$ or $2\pi < x < 3\pi$, and $S''(x) < 0$ if $\pi < x < 2\pi$ or $3\pi < x < 4\pi$. So S is CU on $(0, \pi)$ and $(2\pi, 3\pi)$, and S is CD on $(\pi, 2\pi)$ and $(3\pi, 4\pi)$. There are inflection points at (π, π), $(2\pi, 2\pi)$, and $(3\pi, 3\pi)$.

(d)

59. $f(x) = 1 + \dfrac{1}{x} - \dfrac{1}{x^2}$ has domain $(-\infty, 0) \cup (0, \infty)$.

(a) $\displaystyle\lim_{x\to\pm\infty}\left(1 + \frac{1}{x} - \frac{1}{x^2}\right) = 1$, so $y = 1$ is a HA. $\displaystyle\lim_{x\to 0^+}\left(1 + \frac{1}{x} - \frac{1}{x^2}\right) = \lim_{x\to 0^+}\left(\frac{x^2 + x - 1}{x^2}\right) = -\infty$ since $(x^2 + x - 1) \to -1$ and $x^2 \to 0$ as $x \to 0^+$ [a similar argument can be made for $x \to 0^-$], so $x = 0$ is a VA.

(b) $f'(x) = -\dfrac{1}{x^2} + \dfrac{2}{x^3} = -\dfrac{1}{x^3}(x - 2)$. $f'(x) = 0 \quad \Leftrightarrow \quad x = 2$. $f'(x) > 0 \quad \Leftrightarrow \quad 0 < x < 2$ and $f'(x) < 0 \quad \Leftrightarrow \quad x < 0$ or $x > 2$. So f is increasing on $(0, 2)$ and f is decreasing on $(-\infty, 0)$ and $(2, \infty)$.

(c) f changes from increasing to decreasing at $x = 2$, so $f(2) = \frac{5}{4}$ is a local maximum value. There is no local minimum value.

(d) $f''(x) = \dfrac{2}{x^3} - \dfrac{6}{x^4} = \dfrac{2}{x^4}(x - 3)$. $f''(x) = 0 \quad \Leftrightarrow \quad x = 3$. $f''(x) > 0 \quad \Leftrightarrow$ $x > 3$ and $f''(x) < 0 \quad \Leftrightarrow \quad x < 0$ or $0 < x < 3$. So f is CU on $(3, \infty)$ and f is CD on $(-\infty, 0)$ and $(0, 3)$. There is an inflection point at $\left(3, \frac{11}{9}\right)$.

(e)

60. $f(x) = \dfrac{x^2 - 4}{x^2 + 4}$ has domain $\mathbb{R}$.

(a) $\displaystyle\lim_{x\to\pm\infty}\frac{x^2 - 4}{x^2 + 4} = \lim_{x\to\pm\infty}\frac{1 - 4/x^2}{1 + 4/x^2} = \frac{1}{1} = 1$, so $y = 1$ is a HA. There is no vertical asymptote.

(b) $f'(x) = \dfrac{(x^2 + 4)(2x) - (x^2 - 4)(2x)}{(x^2 + 4)^2} = \dfrac{2x[(x^2 + 4) - (x^2 - 4)]}{(x^2 + 4)^2} = \dfrac{16x}{(x^2 + 4)^2}$. $f'(x) > 0 \quad \Leftrightarrow \quad x > 0$ and $f'(x) < 0 \quad \Leftrightarrow \quad x < 0$. So f is increasing on $(0, \infty)$ and f is decreasing on $(-\infty, 0)$.

(c) f changes from decreasing to increasing at $x = 0$, so $f(0) = -1$ is a local minimum value.

(d) $f''(x) = \dfrac{(x^2 + 4)^2(16) - 16x \cdot 2(x^2 + 4)(2x)}{[(x^2 + 4)^2]^2} = \dfrac{16(x^2 + 4)[(x^2 + 4) - 4x^2]}{(x^2 + 4)^4} = \dfrac{16(4 - 3x^2)}{(x^2 + 4)^3}$.

$f''(x) = 0 \quad \Leftrightarrow \quad x = \pm 2/\sqrt{3}$. $f''(x) > 0 \quad \Leftrightarrow \quad -2/\sqrt{3} < x < 2/\sqrt{3}$ and $f''(x) < 0 \quad \Leftrightarrow \quad x < -2/\sqrt{3}$ or $x > 2/\sqrt{3}$. So f is CU on $\left(-2/\sqrt{3}, 2/\sqrt{3}\right)$ and f is CD on $\left(-\infty, -2/\sqrt{3}\right)$ and $\left(2/\sqrt{3}, \infty\right)$. There are inflection points at $\left(\pm 2/\sqrt{3}, -\frac{1}{2}\right)$.

(e)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. $f(x) = e^{-2/x}$ has domain $(-\infty, 0) \cup (0, \infty)$.

(a) $\lim\limits_{x\to\pm\infty} e^{-2/x} = \lim\limits_{x\to\pm\infty} \dfrac{1}{e^{2/x}} = \dfrac{1}{e^0} = 1$, so $y = 1$ is a HA. $\lim\limits_{x\to 0^-} e^{-2/x} = \infty$, so $x = 0$ is a VA.

(b) $f'(x) = e^{-2/x}\left(\dfrac{2}{x^2}\right) = \dfrac{2}{x^2 e^{2/x}}$. f is always positive on its domain, so f increases on $(-\infty, 0)$ and $(0, \infty)$.

(c) f has no local maximum or minimum value.

(d) $f''(x) = \dfrac{x^2 e^{2/x} \cdot 0 - 2 \cdot \left[x^2 \cdot e^{2/x}\left(-\dfrac{2}{x^2}\right) + e^{2/x} \cdot 2x\right]}{\left(x^2 e^{2/x}\right)^2} = \dfrac{4e^{2/x}(1-x)}{\left(x^2 e^{2/x}\right)^2} = \dfrac{4(1-x)}{x^4 e^{2/x}}$

$f''(x) > 0 \quad \Leftrightarrow \quad x < 1 \ [x \neq 0]$ and $f''(x) < 0 \quad \Leftrightarrow$ $x > 1$ [since x^4 and $e^{2/x}$ are positive for $x \neq 0$], so f is concave up on $(-\infty, 0)$ and $(0, 1)$, and f is concave down on $(1, \infty)$. There is an inflection point at $\left(1, e^{-2}\right) \approx (1, 0.14)$.

(e)

62. $f(x) = \dfrac{e^x}{1 - e^x}$ has domain $\{x \mid 1 - e^x \neq 0\} = \{x \mid e^x \neq 1\} = \{x \mid x \neq 0\}$.

(a) $\lim\limits_{x\to\infty} \dfrac{e^x}{1-e^x} = \lim\limits_{x\to\infty} \dfrac{e^x/e^x}{(1-e^x)/e^x} = \lim\limits_{x\to\infty} \dfrac{1}{1/e^x - 1} = \dfrac{1}{0-1} = -1$, so $y = -1$ is a HA.

$\lim\limits_{x\to-\infty} \dfrac{e^x}{1-e^x} = \dfrac{0}{1-0} = 0$, so $y = 0$ is a HA. $\lim\limits_{x\to 0^+} \dfrac{e^x}{1-e^x} = -\infty$ and $\lim\limits_{x\to 0^-} \dfrac{e^x}{1-e^x} = \infty$, so $x = 0$ is a VA.

(b) $f'(x) = \dfrac{(1-e^x)e^x - e^x(-e^x)}{(1-e^x)^2} = \dfrac{e^x[(1-e^x)+e^x]}{(1-e^x)^2} = \dfrac{e^x}{(1-e^x)^2}$. $f'(x) > 0$ for $x \neq 0$, so f is increasing on $(-\infty, 0)$ and $(0, \infty)$.

(c) There is no local maximum or minimum.

(d) $f''(x) = \dfrac{(1-e^x)^2 e^x - e^x \cdot 2(1-e^x)(-e^x)}{[(1-e^x)^2]^2}$

$= \dfrac{(1-e^x)e^x[(1-e^x)+2e^x]}{(1-e^x)^4} = \dfrac{e^x(e^x+1)}{(1-e^x)^3}$

$f''(x) > 0 \quad \Leftrightarrow \quad (1-e^x)^3 > 0 \quad \Leftrightarrow \quad e^x < 1 \quad \Leftrightarrow \quad x < 0$ and

$f''(x) < 0 \quad \Leftrightarrow \quad x > 0$. So f is CU on $(-\infty, 0)$ and f is CD on $(0, \infty)$.

There is no inflection point.

(e)

63. $f(x) = e^{-x^2}$ has domain $\mathbb{R}$.

(a) $\lim\limits_{x\to\pm\infty} e^{-x^2} = \lim\limits_{x\to\pm\infty} \dfrac{1}{e^{x^2}} = 0$, so $y = 0$ is a HA. There is no VA.

(b) $f(x) = e^{-x^2} \quad \Rightarrow \quad f'(x) = e^{-x^2}(-2x)$. $f'(x) = 0 \quad \Leftrightarrow \quad x = 0$. $f'(x) > 0 \quad \Leftrightarrow \quad x < 0$ and $f'(x) < 0 \quad \Leftrightarrow$ $x > 0$. So f is increasing on $(-\infty, 0)$ and f is decreasing on $(0, \infty)$.

(c) f changes from increasing to decreasing at $x = 0$, so $f(0) = 1$ is a local maximum value. There is no local minimum value.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $f''(x) = e^{-x^2}(-2) + (-2x)e^{-x^2}(-2x) = -2e^{-x^2}(1-2x^2)$.

$f''(x) = 0 \quad \Leftrightarrow \quad x^2 = \frac{1}{2} \quad \Leftrightarrow \quad x = \pm 1/\sqrt{2}$. $\quad f''(x) > 0 \quad \Leftrightarrow$

$x < -1/\sqrt{2}$ or $x > 1/\sqrt{2}$ and $f''(x) < 0 \quad \Leftrightarrow \quad -1/\sqrt{2} < x < 1/\sqrt{2}$. So

f is CU on $\left(-\infty, -1/\sqrt{2}\right)$ and $\left(1/\sqrt{2}, \infty\right)$, and f is CD on $\left(-1/\sqrt{2}, 1/\sqrt{2}\right)$.

There are inflection points at $\left(\pm 1/\sqrt{2}, e^{-1/2}\right)$.

(e)

64. $f(x) = x - \frac{1}{6}x^2 - \frac{2}{3}\ln x$ has domain $(0, \infty)$.

(a) $\lim\limits_{x \to 0^+} \left(x - \frac{1}{6}x^2 - \frac{2}{3}\ln x\right) = \infty$ since $\ln x \to -\infty$ as $x \to 0^+$, so $x = 0$ is a VA. There is no HA.

(b) $f'(x) = 1 - \dfrac{1}{3}x - \dfrac{2}{3x} = \dfrac{3x - x^2 - 2}{3x} = \dfrac{-(x^2 - 3x + 2)}{3x} = \dfrac{-(x-1)(x-2)}{3x}$. $\quad f'(x) > 0 \quad \Leftrightarrow$

$(x-1)(x-2) < 0 \quad \Leftrightarrow \quad 1 < x < 2$ and $f'(x) < 0 \quad \Leftrightarrow \quad 0 < x < 1$ or $x > 2$. So f is increasing on $(1, 2)$ and f is decreasing on $(0, 1)$ and $(2, \infty)$.

(c) f changes from decreasing to increasing at $x = 1$, so $f(1) = \frac{5}{6}$ is a local minimum value. f changes from increasing to decreasing at $x = 2$, so $f(2) = \frac{4}{3} - \frac{2}{3}\ln 2 \approx 0.87$ is a local maximum value.

(d) $f''(x) = -\dfrac{1}{3} + \dfrac{2}{3x^2} = \dfrac{2 - x^2}{3x^2}$. $\quad f''(x) > 0 \quad \Leftrightarrow \quad 0 < x < \sqrt{2}$ and

$f''(x) < 0 \quad \Leftrightarrow \quad x > \sqrt{2}$. So f is CU on $\left(0, \sqrt{2}\right)$ and f is CD on $\left(\sqrt{2}, \infty\right)$. There is an inflection point at $\left(\sqrt{2}, \sqrt{2} - \frac{1}{3} - \frac{1}{3}\ln 2\right)$.

(e)

65. $f(x) = \ln(1 - \ln x)$ is defined when $x > 0$ (so that $\ln x$ is defined) and $1 - \ln x > 0$ [so that $\ln(1 - \ln x)$ is defined]. The second condition is equivalent to $1 > \ln x \quad \Leftrightarrow \quad x < e$, so f has domain $(0, e)$.

(a) As $x \to 0^+$, $\ln x \to -\infty$, so $1 - \ln x \to \infty$ and $f(x) \to \infty$. As $x \to e^-$, $\ln x \to 1^-$, so $1 - \ln x \to 0^+$ and $f(x) \to -\infty$. Thus, $x = 0$ and $x = e$ are vertical asymptotes. There is no horizontal asymptote.

(b) $f'(x) = \dfrac{1}{1 - \ln x}\left(-\dfrac{1}{x}\right) = -\dfrac{1}{x(1 - \ln x)} < 0$ on $(0, e)$. Thus, f is decreasing on its domain, $(0, e)$.

(c) $f'(x) \neq 0$ on $(0, e)$, so f has no local maximum or minimum value.

(d) $f''(x) = -\dfrac{-\left[x(1 - \ln x)\right]'}{\left[x(1 - \ln x)\right]^2} = \dfrac{x(-1/x) + (1 - \ln x)}{x^2(1 - \ln x)^2}$

$= -\dfrac{\ln x}{x^2(1 - \ln x)^2}$

so $f''(x) > 0 \quad \Leftrightarrow \quad \ln x < 0 \quad \Leftrightarrow \quad 0 < x < 1$. Thus, f is CU on $(0, 1)$ and CD on $(1, e)$. There is an inflection point at $(1, 0)$.

(e)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

66. $f(x) = e^{\arctan x}$ has domain $\mathbb{R}$.

(a) $\lim\limits_{x\to\infty} \arctan x = \frac{\pi}{2}$, so $\lim\limits_{x\to\infty} e^{\arctan x} = e^{\pi/2}$ [≈ 4.81], so $y = e^{\pi/2}$ is a HA.

$\lim\limits_{x\to-\infty} e^{\arctan x} = e^{-\pi/2}$ [≈ 0.21], so $y = e^{-\pi/2}$ is a HA. No VA.

(b) $f(x) = e^{\arctan x} \quad\Rightarrow\quad f'(x) = e^{\arctan x} \cdot \dfrac{1}{1+x^2} > 0$ for all x. Thus, f is increasing on $\mathbb{R}$.

(c) There is no local maximum or minimum.

(d) $f''(x) = e^{\arctan x}\left[\dfrac{-2x}{(1+x^2)^2}\right] + \dfrac{1}{1+x^2} \cdot e^{\arctan x} \cdot \dfrac{1}{1+x^2}$

$= \dfrac{e^{\arctan x}}{(1+x^2)^2}(-2x+1)$

$f''(x) > 0 \quad\Leftrightarrow\quad -2x+1 > 0 \quad\Leftrightarrow\quad x < \frac{1}{2}$ and $f''(x) < 0 \quad\Leftrightarrow$ $x > \frac{1}{2}$, so f is CU on $\left(-\infty, \frac{1}{2}\right)$ and f is CD on $\left(\frac{1}{2}, \infty\right)$. There is an inflection point at $\left(\frac{1}{2}, f\left(\frac{1}{2}\right)\right) = \left(\frac{1}{2}, e^{\arctan(1/2)}\right) \approx \left(\frac{1}{2}, 1.59\right)$.

(e)

67. $f(x) = x^4 - cx$, $c > 0 \quad\Rightarrow\quad f'(x) = 4x^3 - c \quad\Rightarrow\quad f'(x) = 0 \quad\Leftrightarrow\quad x = \sqrt[3]{c/4}$. $f'(x) > 0 \quad\Leftrightarrow\quad x > \sqrt[3]{c/4}$ and $f'(x) < 0 \quad\Leftrightarrow\quad x < \sqrt[3]{c/4}$. Thus, f is increasing on $\left(\sqrt[3]{c/4}, \infty\right)$ and decreasing on $\left(-\infty, \sqrt[3]{c/4}\right)$. f changes from decreasing to increasing at $x = \sqrt[3]{c/4}$. Thus, $f\left(\sqrt[3]{c/4}\right)$ is a local minimum value. $f''(x) = 12x^2$ is positive except at $x = 0$, so f is concave up on $(-\infty, 0)$ and $(0, \infty)$. There are no inflection points.

The members of this family have one local minimum point with increasing x-coordinate and decreasing y-coordinate as c increases. The graphs are concave up on $(-\infty, 0)$ and $(0, \infty)$. Since the graphs are continuous at $x = 0$, and the graphs lie above their tangents, we can say that the graphs are concave up on $(-\infty, \infty)$. There is no inflection point.

68. $f(x) = x^3 - 3c^2x + 2c^3$, $c > 0 \quad\Rightarrow\quad f'(x) = 3x^2 - 3c^2 = 3(x^2 - c^2) \quad\Rightarrow\quad f'(x) > 0 \quad\Leftrightarrow\quad |x| > c$ and $f'(x) < 0 \quad\Leftrightarrow\quad |x| < c$. Thus, f is increasing on $(-\infty, -c)$ and (c, ∞), and f is decreasing on $(-c, c)$. f changes from increasing to decreasing at $x = -c$ and from decreasing to increasing at $x = c$. Thus, $f(-c) = 4c^3$ is a local maximum value and $f(c) = 0$ is a local minimum value.

$f''(x) = 6x$, so $f''(x) > 0 \quad\Leftrightarrow\quad x > 0$ and $f''(x) < 0 \quad\Leftrightarrow\quad x < 0$. Thus, f is concave up on $(0, \infty)$ and concave down on $(-\infty, 0)$. There is an inflection point at $\left(0, 2c^3\right)$.

The members of this family have a local maximum point that moves higher and to the left as c increases, and there is a local minimum point on the x-axis that moves to the right as c increases. The graphs are concave down on $(-\infty, 0)$ and concave up on $(0, \infty)$. There is an inflection point on the y-axis that moves higher as c increases.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

69. (a)

From the graph, we get an estimate of $f(1) \approx 1.41$ as a local maximum value, and no local minimum value.

$f(x) = \dfrac{x+1}{\sqrt{x^2+1}} \quad \Rightarrow \quad f'(x) = \dfrac{1-x}{(x^2+1)^{3/2}}$.

$f'(x) = 0 \quad \Leftrightarrow \quad x = 1$. $f(1) = \frac{2}{\sqrt{2}} = \sqrt{2}$ is the exact value.

(b) From the graph in part (a), f increases most rapidly somewhere between $x = -\frac{1}{2}$ and $x = -\frac{1}{4}$. To find the exact value, we need to find the maximum value of f', which we can do by finding the critical numbers of f'.

$f''(x) = \dfrac{2x^2 - 3x - 1}{(x^2+1)^{5/2}} = 0 \quad \Leftrightarrow \quad x = \dfrac{3 \pm \sqrt{17}}{4}$. By the First Derivative Test applied to f', $x = \dfrac{3+\sqrt{17}}{4}$ corresponds to the *minimum* value of f' and the maximum value of f' occurs at $x = \frac{3-\sqrt{17}}{4} \approx -0.28$.

70. (a)

Tracing the graph gives us estimates of $f(0) = 0$ for a local minimum value and $f(2) = 0.54$ for a local maximum value.

$f(x) = x^2 e^{-x} \quad \Rightarrow \quad f'(x) = xe^{-x}(2-x)$. $\quad f'(x) = 0 \quad \Leftrightarrow \quad x = 0$ or 2.

$f(0) = 0$ and $f(2) = 4e^{-2}$ are the exact values.

(b) From the graph in part (a), f increases most rapidly around $x = \frac{3}{4}$. To find the exact value, we need to find the maximum value of f', which we can do by finding the critical numbers of f'. $\quad f''(x) = e^{-x}\left(x^2 - 4x + 2\right) = 0 \quad \Rightarrow$ $x = 2 \pm \sqrt{2}$. By the First Derivative Test applied to f', $x = 2 + \sqrt{2}$ corresponds to the *minimum* value of f' and the maximum value of f' is at $\left(2-\sqrt{2}, \left(2-\sqrt{2}\right)^2 e^{-2+\sqrt{2}}\right) \approx (0.59, 0.19)$.

71. $f(x) = \sin 2x + \sin 4x \quad \Rightarrow \quad f'(x) = 2\cos 2x + 4\cos 4x \quad \Rightarrow \quad f''(x) = -4\sin 2x - 16\sin 4x$

(a) From the graph of f, it seems that f is CD on $(0, 0.8)$, CU on $(0.8, 1.6)$, CD on $(1.6, 2.3)$, and CU on $(2.3, \pi)$. The inflection points appear to be at $(0.8, 0.7)$, $(1.6, 0)$, and $(2.3, -0.7)$.

(b) From the graph of f'' (and zooming in near the zeros), it seems that f is CD on $(0, 0.85)$, CU on $(0.85, 1.57)$, CD on $(1.57, 2.29)$, and CU on $(2.29, \pi)$. Refined estimates of the inflection points are $(0.85, 0.74)$, $(1.57, 0)$, and $(2.29, -0.74)$.

72. $f(x) = (x-1)^2(x+1)^3 \quad \Rightarrow$

$f'(x) = (x-1)^2 3(x+1)^2 + (x+1)^3 2(x-1)$

$\quad = (x-1)(x+1)^2\left[3(x-1) + 2(x+1)\right] = (x-1)(x+1)^2(5x-1) \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\begin{aligned} f''(x) &= (1)(x+1)^2(5x-1) + (x-1)(2)(x+1)(5x-1) + (x-1)(x+1)^2(5) \\ &= (x+1)[(x+1)(5x-1) + 2(x-1)(5x-1) + 5(x-1)(x+1)] \\ &= (x+1)[5x^2 + 4x - 1 + 10x^2 - 12x + 2 + 5x^2 - 5] \\ &= (x+1)(20x^2 - 8x - 4) = 4(x+1)(5x^2 - 2x - 1) \end{aligned}$$

(a) From the graph of f, it seems that f is CD on $(-\infty, -1)$, CU on $(-1, -0.3)$, CD on $(-0.3, 0.7)$, and CU on $(0.7, \infty)$. The inflection points appear to be at $(-1, 0)$, $(-0.3, 0.6)$, and $(0.7, 0.5)$.

(b) From the graph of f'' (and zooming in near the zeros), it seems that f is CD on $(-1, 0)$, CU on $(-1, -0.29)$, CD on $(-0.29, 0.69)$, and CU on $(0.69, \infty)$. Refined estimates of the inflection points are $(-1, 0)$, $(-0.29, 0.60)$, and $(0.69, 0.46)$.

73. $f(x) = \dfrac{x^4 + x^3 + 1}{\sqrt{x^2 + x + 1}}$. In Maple, we define f and then use the command `plot(diff(diff(f,x),x),x=-2..2);`. In Mathematica, we define f and then use `Plot[Dt[Dt[f,x],x],{x,-2,2}]`. We see that $f'' > 0$ for $x < -0.6$ and $x > 0.0$ $[\approx 0.03]$ and $f'' < 0$ for $-0.6 < x < 0.0$. So f is CU on $(-\infty, -0.6)$ and $(0.0, \infty)$ and CD on $(-0.6, 0.0)$.

74. $f(x) = \dfrac{x^2 \tan^{-1} x}{1 + x^3}$. It appears that f'' is positive (and thus f is concave upward) on $(-\infty, -1)$, $(0, 0.7)$, and $(2.5, \infty)$; and f'' is negative (and thus f is concave downward) on $(-1, 0)$ and $(0.7, 2.5)$.

75. (a) The rate of increase of the population is initially very small, then gets larger until it reaches a maximum at about $t = 8$ hours, and decreases toward 0 as the population begins to level off.

(b) The rate of increase has its maximum value at $t = 8$ hours.

(c) The population function is concave upward on $(0, 8)$ and concave downward on $(8, 18)$.

(d) At $t = 8$, the population is about 350, so the inflection point is about $(8, 350)$.

76. If $S(t)$ is the average SAT score as a function of time t, then $S'(t) < 0$ (since the SAT scores are declining) and $S''(t) > 0$ (since the rate of decrease of the scores is increasing—becoming less negative).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

77. If $D(t)$ is the size of the national deficit as a function of time t, then at the time of the speech $D'(t) > 0$ (since the deficit is increasing), and $D''(t) < 0$ (since the rate of increase of the deficit is decreasing).

78. (a) I'm very unhappy. It's uncomfortably hot and $f'(3) = 2$ indicates that the temperature is increasing, and $f''(3) = 4$ indicates that the rate of increase is increasing. (The temperature is rapidly getting warmer.)

(b) I'm still unhappy, but not as unhappy as in part (a). It's uncomfortably hot and $f'(3) = 2$ indicates that the temperature is increasing, but $f''(3) = -4$ indicates that the rate of increase is decreasing. (The temperature is slowly getting warmer.)

(c) I'm somewhat happy. It's uncomfortably hot and $f'(3) = -2$ indicates that the temperature is decreasing, but $f''(3) = 4$ indicates that the rate of change is increasing. (The rate of change is negative but it's becoming less negative. The temperature is slowly getting cooler.)

(d) I'm very happy. It's uncomfortably hot and $f'(3) = -2$ indicates that the temperature is decreasing, and $f''(3) = -4$ indicates that the rate of change is decreasing, that is, becoming more negative. (The temperature is rapidly getting cooler.)

79. Most students learn more in the third hour of studying than in the eighth hour, so $K(3) - K(2)$ is larger than $K(8) - K(7)$. In other words, as you begin studying for a test, the rate of knowledge gain is large and then starts to taper off, so $K'(t)$ decreases and the graph of K is concave downward.

80. At first the depth increases slowly because the base of the mug is wide. But as the mug narrows, the coffee rises more quickly. Thus, the depth d increases at an increasing rate and its graph is concave upward. The rate of increase of d has a maximum where the mug is narrowest; that is, when the mug is half full. It is there that the inflection point (IP) occurs. Then the rate of increase of d starts to decrease as the mug widens and the graph becomes concave down.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

81. $S(t) = At^p e^{-kt}$ with $A = 0.01$, $p = 4$, and $k = 0.07$. We will find the zeros of f'' for $f(t) = t^p e^{-kt}$.

$$f'(t) = t^p(-ke^{-kt}) + e^{-kt}(pt^{p-1}) = e^{-kt}(-kt^p + pt^{p-1})$$

$$\begin{aligned} f''(t) &= e^{-kt}(-kpt^{p-1} + p(p-1)t^{p-2}) + (-kt^p + pt^{p-1})(-ke^{-kt}) \\ &= t^{p-2}e^{-kt}[-kpt + p(p-1) + k^2t^2 - kpt] \\ &= t^{p-2}e^{-kt}(k^2t^2 - 2kpt + p^2 - p) \end{aligned}$$

Using the given values of p and k gives us $f''(t) = t^2 e^{-0.07t}(0.0049t^2 - 0.56t + 12)$. So $S''(t) = 0.01f''(t)$ and its zeros are $t = 0$ and the solutions of $0.0049t^2 - 0.56t + 12 = 0$, which are $t_1 = \frac{200}{7} \approx 28.57$ and $t_2 = \frac{600}{7} \approx 85.71$.

At t_1 minutes, the rate of increase of the level of medication in the bloodstream is at its greatest and at t_2 minutes, the rate of decrease is the greatest.

82. (a) As $|x| \to \infty$, $t = -x^2/(2\sigma^2) \to -\infty$, and $e^t \to 0$. The HA is $y = 0$. Since t takes on its maximum value at $x = 0$, so does e^t. Showing this result using derivatives, we have $f(x) = e^{-x^2/(2\sigma^2)} \quad \Rightarrow \quad f'(x) = e^{-x^2/(2\sigma^2)}(-x/\sigma^2)$.

$f'(x) = 0 \quad \Leftrightarrow \quad x = 0$. Because f' changes from positive to negative at $x = 0$, $f(0) = 1$ is a local maximum. For inflection points, we find $f''(x) = -\dfrac{1}{\sigma^2}\left[e^{-x^2/(2\sigma^2)} \cdot 1 + xe^{-x^2/(2\sigma^2)}(-x/\sigma^2)\right] = \dfrac{-1}{\sigma^2}e^{-x^2/(2\sigma^2)}(1 - x^2/\sigma^2)$.

$f''(x) = 0 \quad \Leftrightarrow \quad x^2 = \sigma^2 \quad \Leftrightarrow \quad x = \pm\sigma$. $\quad f''(x) < 0 \quad \Leftrightarrow \quad x^2 < \sigma^2 \quad \Leftrightarrow \quad -\sigma < x < \sigma$.

So f is CD on $(-\sigma, \sigma)$ and CU on $(-\infty, -\sigma)$ and (σ, ∞). IP at $(\pm\sigma, e^{-1/2})$.

(b) Since we have IP at $x = \pm\sigma$, the inflection points move away from the y-axis as σ increases.

(c)

From the graph, we see that as σ increases, the graph tends to spread out and there is more area between the curve and the x-axis.

83. $f(x) = ax^3 + bx^2 + cx + d \quad \Rightarrow \quad f'(x) = 3ax^2 + 2bx + c$.

We are given that $f(1) = 0$ and $f(-2) = 3$, so $f(1) = a + b + c + d = 0$ and $f(-2) = -8a + 4b - 2c + d = 3$. Also $f'(1) = 3a + 2b + c = 0$ and $f'(-2) = 12a - 4b + c = 0$ by Fermat's Theorem. Solving these four equations, we get $a = \frac{2}{9}$, $b = \frac{1}{3}$, $c = -\frac{4}{3}$, $d = \frac{7}{9}$, so the function is $f(x) = \frac{1}{9}\left(2x^3 + 3x^2 - 12x + 7\right)$.

84. $f(x) = axe^{bx^2} \quad \Rightarrow \quad f'(x) = a\left[xe^{bx^2} \cdot 2bx + e^{bx^2} \cdot 1\right] = ae^{bx^2}(2bx^2 + 1)$. For $f(2) = 1$ to be a maximum value, we must have $f'(2) = 0$. $f(2) = 1 \quad \Rightarrow \quad 1 = 2ae^{4b}$ and $f'(2) = 0 \quad \Rightarrow \quad 0 = (8b + 1)ae^{4b}$. So $8b + 1 = 0 \quad [a \neq 0] \quad \Rightarrow$ $b = -\frac{1}{8}$ and now $1 = 2ae^{-1/2} \quad \Rightarrow \quad a = \sqrt{e}/2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

85. $y = \dfrac{1+x}{1+x^2} \quad \Rightarrow \quad y' = \dfrac{(1+x^2)(1)-(1+x)(2x)}{(1+x^2)^2} = \dfrac{1-2x-x^2}{(1+x^2)^2} \quad \Rightarrow$

$$y'' = \frac{(1+x^2)^2(-2-2x)-(1-2x-x^2)\cdot 2(1+x^2)(2x)}{[(1+x^2)^2]^2} = \frac{2(1+x^2)[(1+x^2)(-1-x)-(1-2x-x^2)(2x)]}{(1+x^2)^4}$$

$$= \frac{2(-1-x-x^2-x^3-2x+4x^2+2x^3)}{(1+x^2)^3} = \frac{2(x^3+3x^2-3x-1)}{(1+x^2)^3} = \frac{2(x-1)(x^2+4x+1)}{(1+x^2)^3}$$

So $y''=0 \quad \Rightarrow \quad x = 1, -2\pm\sqrt{3}$. Let $a=-2-\sqrt{3}$, $b=-2+\sqrt{3}$, and $c=1$. We can show that $f(a)=\frac{1}{4}\left(1-\sqrt{3}\right)$, $f(b)=\frac{1}{4}\left(1+\sqrt{3}\right)$, and $f(c)=1$. To show that these three points of inflection lie on one straight line, we'll show that the slopes m_{ac} and m_{bc} are equal.

$$m_{ac} = \frac{f(c)-f(a)}{c-a} = \frac{1-\frac{1}{4}\left(1-\sqrt{3}\right)}{1-\left(-2-\sqrt{3}\right)} = \frac{\frac{3}{4}+\frac{1}{4}\sqrt{3}}{3+\sqrt{3}} = \frac{1}{4}$$

$$m_{bc} = \frac{f(c)-f(b)}{c-b} = \frac{1-\frac{1}{4}\left(1+\sqrt{3}\right)}{1-\left(-2+\sqrt{3}\right)} = \frac{\frac{3}{4}-\frac{1}{4}\sqrt{3}}{3-\sqrt{3}} = \frac{1}{4}$$

86. $y = f(x) = e^{-x}\sin x \quad \Rightarrow \quad y' = e^{-x}\cos x + \sin x(-e^{-x}) = e^{-x}(\cos x - \sin x) \quad \Rightarrow$

$y'' = e^{-x}(-\sin x - \cos x) + (\cos x - \sin x)(-e^{-x}) = e^{-x}(-\sin x - \cos x - \cos x + \sin x) = e^{-x}(-2\cos x)$.

So $y''=0 \quad \Rightarrow \quad \cos x = 0 \quad \Rightarrow \quad x = \frac{\pi}{2}+n\pi$. At these values of x, f has points of inflection and since $\sin\left(\frac{\pi}{2}+n\pi\right) = \pm 1$, we get $y = \pm e^{-x}$, so f intersects the other curves at its inflection points.

Let $g(x) = e^{-x}$ and $h(x) = -e^{-x}$, so that $g'(x) = -e^{-x}$ and $h'(x) = e^{-x}$. Now $f'\left(\frac{\pi}{2}+n\pi\right) = e^{-(\pi/2+n\pi)}\left[\cos\left(\frac{\pi}{2}+n\pi\right) - \sin\left(\frac{\pi}{2}+n\pi\right)\right] = -e^{-(\pi/2+n\pi)}\sin\left(\frac{\pi}{2}+n\pi\right)$. If n is odd, then $f'\left(\frac{\pi}{2}+n\pi\right) = e^{-(\pi/2+n\pi)} = h'\left(\frac{\pi}{2}+n\pi\right)$. If n is even, then $f'\left(\frac{\pi}{2}+n\pi\right) = -e^{-(\pi/2+n\pi)} = g'\left(\frac{\pi}{2}+n\pi\right)$. Thus, at $x = \frac{\pi}{2}+n\pi$, f has the same slope as either g or h, and hence, g and h touch f at its inflection points.

87. $y = x\sin x \quad \Rightarrow \quad y' = x\cos x + \sin x \quad \Rightarrow \quad y'' = -x\sin x + 2\cos x$. $\quad y''=0 \quad \Rightarrow \quad 2\cos x = x\sin x$ [which is y] $\quad \Rightarrow$ $(2\cos x)^2 = (x\sin x)^2 \quad \Rightarrow \quad 4\cos^2 x = x^2\sin^2 x \quad \Rightarrow \quad 4\cos^2 x = x^2(1-\cos^2 x) \quad \Rightarrow \quad 4\cos^2 x + x^2\cos^2 x = x^2 \quad \Rightarrow$ $\cos^2 x(4+x^2) = x^2 \quad \Rightarrow \quad 4\cos^2 x(x^2+4) = 4x^2 \quad \Rightarrow \quad y^2(x^2+4) = 4x^2$ since $y = 2\cos x$ when $y''=0$.

88. (a) We will make use of the converse of the Concavity Test (along with the stated assumptions); that is, if f is concave upward on I, then $f''>0$ on I. If f and g are CU on I, then $f''>0$ and $g''>0$ on I, so $(f+g)'' = f''+g''>0$ on $I \quad \Rightarrow$ $f+g$ is CU on I.

(b) Since f is positive and CU on I, $f>0$ and $f''>0$ on I. So $g(x) = [f(x)]^2 \quad \Rightarrow \quad g' = 2ff' \quad \Rightarrow$ $g'' = 2f'f' + 2ff'' = 2(f')^2 + 2ff'' > 0 \quad \Rightarrow \quad g$ is CU on I.

89. (a) Since f and g are positive, increasing, and CU on I with f'' and g'' never equal to 0, we have $f>0$, $f'\geq 0$, $f''>0$, $g>0$, $g'\geq 0$, $g''>0$ on I. Then $(fg)' = f'g+fg' \quad \Rightarrow \quad (fg)'' = f''g+2f'g'+fg'' \geq f''g+fg''>0$ on $I \quad \Rightarrow$ fg is CU on I.

(b) In part (a), if f and g are both decreasing instead of increasing, then $f'\leq 0$ and $g'\leq 0$ on I, so we still have $2f'g'\geq 0$ on I. Thus, $(fg)'' = f''g+2f'g'+fg'' \geq f''g+fg''>0$ on $I \quad \Rightarrow \quad fg$ is CU on I as in part (a).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) Suppose f is increasing and g is decreasing [with f and g positive and CU]. Then $f' \geq 0$ and $g' \leq 0$ on I, so $2f'g' \leq 0$ on I and the argument in parts (a) and (b) fails.

Example 1. $I = (0, \infty)$, $f(x) = x^3$, $g(x) = 1/x$. Then $(fg)(x) = x^2$, so $(fg)'(x) = 2x$ and $(fg)''(x) = 2 > 0$ on I. Thus, fg is CU on I.

Example 2. $I = (0, \infty)$, $f(x) = 4x\sqrt{x}$, $g(x) = 1/x$. Then $(fg)(x) = 4\sqrt{x}$, so $(fg)'(x) = 2/\sqrt{x}$ and $(fg)''(x) = -1/\sqrt{x^3} < 0$ on I. Thus, fg is CD on I.

Example 3. $I = (0, \infty)$, $f(x) = x^2$, $g(x) = 1/x$. Thus, $(fg)(x) = x$, so fg is linear on I.

90. Since f and g are CU on $(-\infty, \infty)$, $f'' > 0$ and $g'' > 0$ on $(-\infty, \infty)$. $h(x) = f(g(x)) \Rightarrow$ $h'(x) = f'(g(x))\, g'(x) \Rightarrow h''(x) = f''(g(x))\, g'(x)\, g'(x) + f'(g(x))\, g''(x) = f''(g(x))[g'(x)]^2 + f'(g(x))\, g''(x) > 0$ if $f' > 0$. So h is CU if f is increasing.

91. Let the cubic function be $f(x) = ax^3 + bx^2 + cx + d \Rightarrow f'(x) = 3ax^2 + 2bx + c \Rightarrow f''(x) = 6ax + 2b$. So f is CU when $6ax + 2b > 0 \Leftrightarrow x > -b/(3a)$, CD when $x < -b/(3a)$, and so the only point of inflection occurs when $x = -b/(3a)$. If the graph has three x-intercepts x_1, x_2 and x_3, then the expression for $f(x)$ must factor as $f(x) = a(x - x_1)(x - x_2)(x - x_3)$. Multiplying these factors together gives us

$$f(x) = a[x^3 - (x_1 + x_2 + x_3)x^2 + (x_1x_2 + x_1x_3 + x_2x_3)x - x_1x_2x_3]$$

Equating the coefficients of the x^2-terms for the two forms of f gives us $b = -a(x_1 + x_2 + x_3)$. Hence, the x-coordinate of the point of inflection is $-\dfrac{b}{3a} = -\dfrac{-a(x_1 + x_2 + x_3)}{3a} = \dfrac{x_1 + x_2 + x_3}{3}$.

92. $P(x) = x^4 + cx^3 + x^2 \Rightarrow P'(x) = 4x^3 + 3cx^2 + 2x \Rightarrow P''(x) = 12x^2 + 6cx + 2$. The graph of $P''(x)$ is a parabola. If $P''(x)$ has two solutions, then it changes sign twice and so has two inflection points. This happens when the discriminant of $P''(x)$ is positive, that is, $(6c)^2 - 4 \cdot 12 \cdot 2 > 0 \Leftrightarrow 36c^2 - 96 > 0 \Leftrightarrow |c| > \frac{2\sqrt{6}}{3} \approx 1.63$. If $36c^2 - 96 = 0 \Leftrightarrow c = \pm\frac{2\sqrt{6}}{3}$, $P''(x)$ is 0 at one point, but there is still no inflection point since $P''(x)$ never changes sign, and if $36c^2 - 96 < 0 \Leftrightarrow |c| < \frac{2\sqrt{6}}{3}$, then $P''(x)$ never changes sign, and so there is no inflection point.

$c = 6$

$c = 3$

$c = 1.8$

$c = \frac{2\sqrt{6}}{3}$

$c = 0$

$c = -2$

For large positive c, the graph of f has two inflection points and a large dip to the left of the y-axis. As c decreases, the graph

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

of f becomes flatter for $x < 0$, and eventually the dip rises above the x-axis, and then disappears entirely, along with the inflection points. As c continues to decrease, the dip and the inflection points reappear, to the right of the origin.

93. By hypothesis $g = f'$ is differentiable on an open interval containing c. Since $(c, f(c))$ is a point of inflection, the concavity changes at $x = c$, so $f''(x)$ changes signs at $x = c$. Hence, by the First Derivative Test, f' has a local extremum at $x = c$. Thus, by Fermat's Theorem $f''(c) = 0$.

94. $f(x) = x^4 \quad \Rightarrow \quad f'(x) = 4x^3 \quad \Rightarrow \quad f''(x) = 12x^2 \quad \Rightarrow \quad f''(0) = 0$. For $x < 0$, $f''(x) > 0$, so f is CU on $(-\infty, 0)$; for $x > 0$, $f''(x) > 0$, so f is also CU on $(0, \infty)$. Since f does not change concavity at 0, $(0, 0)$ is not an inflection point.

95. Using the fact that $|x| = \sqrt{x^2}$, we have that $g(x) = x\,|x| = x\sqrt{x^2} \quad \Rightarrow \quad g'(x) = \sqrt{x^2} + \sqrt{x^2} = 2\sqrt{x^2} = 2\,|x| \quad \Rightarrow$ $g''(x) = 2x\big(x^2\big)^{-1/2} = \dfrac{2x}{|x|} < 0$ for $x < 0$ and $g''(x) > 0$ for $x > 0$, so $(0, 0)$ is an inflection point. But $g''(0)$ does not exist.

96. There must exist some interval containing c on which f''' is positive, since $f'''(c)$ is positive and f''' is continuous. On this interval, f'' is increasing (since f''' is positive), so $f'' = (f')'$ changes from negative to positive at c. So by the First Derivative Test, f' has a local minimum at $x = c$ and thus cannot change sign there, so f has no maximum or minimum at c. But since f'' changes from negative to positive at c, f has a point of inflection at c (it changes from concave down to concave up).

97. Suppose that f is differentiable on an interval I and $f'(x) > 0$ for all x in I except $x = c$. To show that f is increasing on I, let x_1, x_2 be two numbers in I with $x_1 < x_2$.

Case 1 $x_1 < x_2 < c$. Let J be the interval $\{x \in I \mid x < c\}$. By applying the Increasing/Decreasing Test to f on J, we see that f is increasing on J, so $f(x_1) < f(x_2)$.

Case 2 $c < x_1 < x_2$. Apply the Increasing/Decreasing Test to f on $K = \{x \in I \mid x > c\}$.

Case 3 $x_1 < x_2 = c$. Apply the proof of the Increasing/Decreasing Test, using the Mean Value Theorem (MVT) on the interval $[x_1, x_2]$ and noting that the MVT does not require f to be differentiable at the endpoints of $[x_1, x_2]$.

Case 4 $c = x_1 < x_2$. Same proof as in Case 3.

Case 5 $x_1 < c < x_2$. By Cases 3 and 4, f is increasing on $[x_1, c]$ and on $[c, x_2]$, so $f(x_1) < f(c) < f(x_2)$.

In all cases, we have shown that $f(x_1) < f(x_2)$. Since x_1, x_2 were any numbers in I with $x_1 < x_2$, we have shown that f is increasing on I.

98. $f(x) = cx + \dfrac{1}{x^2+3} \quad \Rightarrow \quad f'(x) = c - \dfrac{2x}{(x^2+3)^2}$. $f'(x) > 0 \quad \Leftrightarrow \quad c > \dfrac{2x}{(x^2+3)^2}$ [call this $g(x)$]. Now f' is positive (and hence f increasing) if $c > g$, so we'll find the maximum value of g.

$$g'(x) = \frac{(x^2+3)^2 \cdot 2 - 2x \cdot 2(x^2+3) \cdot 2x}{[(x^2+3)^2]^2} = \frac{2(x^2+3)[(x^2+3) - 4x^2]}{(x^2+3)^4} = \frac{2(3-3x^2)}{(x^2+3)^3} = \frac{6(1+x)(1-x)}{(x^2+3)^3}.$$

$g'(x) = 0 \quad \Leftrightarrow \quad x = \pm 1$. $g'(x) > 0$ on $(0, 1)$ and $g'(x) < 0$ on $(1, \infty)$, so g is increasing on $(0, 1)$ and decreasing on

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$(1, \infty)$, and hence g has a maximum value on $(0, \infty)$ of $g(1) = \frac{2}{16} = \frac{1}{8}$. Also since $g(x) \leq 0$ if $x \leq 0$, the maximum value of g on $(-\infty, \infty)$ is $\frac{1}{8}$. Thus, when $c > \frac{1}{8}$, f is increasing. When $c = \frac{1}{8}$, $f'(x) > 0$ on $(-\infty, 1)$ and $(1, \infty)$, and hence f is increasing on these intervals. Since f is continuous, we may conclude that f is also increasing on $(-\infty, \infty)$ if $c = \frac{1}{8}$. Therefore, f is increasing on $(-\infty, \infty)$ if $c \geq \frac{1}{8}$.

99. (a) $f(x) = x^4 \sin \dfrac{1}{x} \quad \Rightarrow \quad f'(x) = x^4 \cos \dfrac{1}{x} \left(-\dfrac{1}{x^2}\right) + \sin \dfrac{1}{x}\,(4x^3) = 4x^3 \sin \dfrac{1}{x} - x^2 \cos \dfrac{1}{x}$.

$g(x) = x^4 \left(2 + \sin \dfrac{1}{x}\right) = 2x^4 + f(x) \quad \Rightarrow \quad g'(x) = 8x^3 + f'(x)$.

$h(x) = x^4 \left(-2 + \sin \dfrac{1}{x}\right) = -2x^4 + f(x) \quad \Rightarrow \quad h'(x) = -8x^3 + f'(x)$.

It is given that $f(0) = 0$, so $f'(0) = \lim\limits_{x \to 0} \dfrac{f(x) - f(0)}{x - 0} = \lim\limits_{x \to 0} \dfrac{x^4 \sin \frac{1}{x} - 0}{x} = \lim\limits_{x \to 0} x^3 \sin \dfrac{1}{x}$. Since $-\left|x^3\right| \leq x^3 \sin \dfrac{1}{x} \leq \left|x^3\right|$ and $\lim\limits_{x \to 0} \left|x^3\right| = 0$, we see that $f'(0) = 0$ by the Squeeze Theorem. Also, $g'(0) = 8(0)^3 + f'(0) = 0$ and $h'(0) = -8(0)^3 + f'(0) = 0$, so 0 is a critical number of f, g, and h.

For $x_{2n} = \dfrac{1}{2n\pi}$ [n a nonzero integer], $\sin \dfrac{1}{x_{2n}} = \sin 2n\pi = 0$ and $\cos \dfrac{1}{x_{2n}} = \cos 2n\pi = 1$, so $f'(x_{2n}) = -x_{2n}^2 < 0$.

For $x_{2n+1} = \dfrac{1}{(2n+1)\pi}$, $\sin \dfrac{1}{x_{2n+1}} = \sin(2n+1)\pi = 0$ and $\cos \dfrac{1}{x_{2n+1}} = \cos(2n+1)\pi = -1$, so $f'(x_{2n+1}) = x_{2n+1}^2 > 0$. Thus, f' changes sign infinitely often on both sides of 0.

Next, $g'(x_{2n}) = 8x_{2n}^3 + f'(x_{2n}) = 8x_{2n}^3 - x_{2n}^2 = x_{2n}^2(8x_{2n} - 1) < 0$ for $x_{2n} < \frac{1}{8}$, but $g'(x_{2n+1}) = 8x_{2n+1}^3 + x_{2n+1}^2 = x_{2n+1}^2(8x_{2n+1} + 1) > 0$ for $x_{2n+1} > -\frac{1}{8}$, so g' changes sign infinitely often on both sides of 0.

Last, $h'(x_{2n}) = -8x_{2n}^3 + f'(x_{2n}) = -8x_{2n}^3 - x_{2n}^2 = -x_{2n}^2(8x_{2n} + 1) < 0$ for $x_{2n} > -\frac{1}{8}$ and $h'(x_{2n+1}) = -8x_{2n+1}^3 + x_{2n+1}^2 = x_{2n+1}^2(-8x_{2n+1} + 1) > 0$ for $x_{2n+1} < \frac{1}{8}$, so h' changes sign infinitely often on both sides of 0.

(b) $f(0) = 0$ and since $\sin \dfrac{1}{x}$ and hence $x^4 \sin \dfrac{1}{x}$ is both positive and negative inifinitely often on both sides of 0, and arbitrarily close to 0, f has neither a local maximum nor a local minimum at 0.

Since $2 + \sin \dfrac{1}{x} \geq 1$, $g(x) = x^4 \left(2 + \sin \dfrac{1}{x}\right) > 0$ for $x \neq 0$, so $g(0) = 0$ is a local minimum.

Since $-2 + \sin \dfrac{1}{x} \leq -1$, $h(x) = x^4 \left(-2 + \sin \dfrac{1}{x}\right) < 0$ for $x \neq 0$, so $h(0) = 0$ is a local maximum.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4.4 Indeterminate Forms and l'Hospital's Rule

Note: The use of l'Hospital's Rule is indicated by an H above the equal sign: $\overset{\text{H}}{=}$

1. (a) $\lim\limits_{x\to a} \dfrac{f(x)}{g(x)}$ is an indeterminate form of type $\dfrac{0}{0}$.

(b) $\lim\limits_{x\to a} \dfrac{f(x)}{p(x)} = 0$ because the numerator approaches 0 while the denominator becomes large.

(c) $\lim\limits_{x\to a} \dfrac{h(x)}{p(x)} = 0$ because the numerator approaches a finite number while the denominator becomes large.

(d) If $\lim\limits_{x\to a} p(x) = \infty$ and $f(x) \to 0$ through positive values, then $\lim\limits_{x\to a} \dfrac{p(x)}{f(x)} = \infty$. [For example, take $a = 0$, $p(x) = 1/x^2$, and $f(x) = x^2$.] If $f(x) \to 0$ through negative values, then $\lim\limits_{x\to a} \dfrac{p(x)}{f(x)} = -\infty$. [For example, take $a = 0$, $p(x) = 1/x^2$, and $f(x) = -x^2$.] If $f(x) \to 0$ through both positive and negative values, then the limit might not exist. [For example, take $a = 0$, $p(x) = 1/x^2$, and $f(x) = x$.]

(e) $\lim\limits_{x\to a} \dfrac{p(x)}{q(x)}$ is an indeterminate form of type $\dfrac{\infty}{\infty}$.

2. (a) $\lim\limits_{x\to a} [f(x)p(x)]$ is an indeterminate form of type $0 \cdot \infty$.

(b) When x is near a, $p(x)$ is large and $h(x)$ is near 1, so $h(x)p(x)$ is large. Thus, $\lim\limits_{x\to a} [h(x)p(x)] = \infty$.

(c) When x is near a, $p(x)$ and $q(x)$ are both large, so $p(x)q(x)$ is large. Thus, $\lim\limits_{x\to a} [p(x)q(x)] = \infty$.

3. (a) When x is near a, $f(x)$ is near 0 and $p(x)$ is large, so $f(x) - p(x)$ is large negative. Thus, $\lim\limits_{x\to a} [f(x) - p(x)] = -\infty$.

(b) $\lim\limits_{x\to a} [p(x) - q(x)]$ is an indeterminate form of type $\infty - \infty$.

(c) When x is near a, $p(x)$ and $q(x)$ are both large, so $p(x) + q(x)$ is large. Thus, $\lim\limits_{x\to a} [p(x) + q(x)] = \infty$.

4. (a) $\lim\limits_{x\to a} [f(x)]^{g(x)}$ is an indeterminate form of type 0^0.

(b) If $y = [f(x)]^{p(x)}$, then $\ln y = p(x) \ln f(x)$. When x is near a, $p(x) \to \infty$ and $\ln f(x) \to -\infty$, so $\ln y \to -\infty$. Therefore, $\lim\limits_{x\to a} [f(x)]^{p(x)} = \lim\limits_{x\to a} y = \lim\limits_{x\to a} e^{\ln y} = 0$, provided f^p is defined.

(c) $\lim\limits_{x\to a} [h(x)]^{p(x)}$ is an indeterminate form of type 1^∞.

(d) $\lim\limits_{x\to a} [p(x)]^{f(x)}$ is an indeterminate form of type ∞^0.

(e) If $y = [p(x)]^{q(x)}$, then $\ln y = q(x) \ln p(x)$. When x is near a, $q(x) \to \infty$ and $\ln p(x) \to \infty$, so $\ln y \to \infty$. Therefore, $\lim\limits_{x\to a} [p(x)]^{q(x)} = \lim\limits_{x\to a} y = \lim\limits_{x\to a} e^{\ln y} = \infty$.

(f) $\lim\limits_{x\to a} \sqrt[q(x)]{p(x)} = \lim\limits_{x\to a} [p(x)]^{1/q(x)}$ is an indeterminate form of type ∞^0.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. From the graphs of f and g, we see that $\lim_{x\to 2} f(x) = 0$ and $\lim_{x\to 2} g(x) = 0$, so l'Hospital's Rule applies.

$$\lim_{x\to 2}\frac{f(x)}{g(x)} = \lim_{x\to 2}\frac{f'(x)}{g'(x)} = \frac{\lim_{x\to 2} f'(x)}{\lim_{x\to 2} g'(x)} = \frac{f'(2)}{g'(2)} = \frac{1.8}{\frac{4}{5}} = \frac{9}{4}$$

6. From the graphs of f and g, we see that $\lim_{x\to 2} f(x) = 0$ and $\lim_{x\to 2} g(x) = 0$, so l'Hospital's Rule applies.

$$\lim_{x\to 2}\frac{f(x)}{g(x)} = \lim_{x\to 2}\frac{f'(x)}{g'(x)} = \frac{\lim_{x\to 2} f'(x)}{\lim_{x\to 2} g'(x)} = \frac{f'(2)}{g'(2)} = \frac{1.5}{-1} = -\frac{3}{2}$$

7. f and $g = e^x - 1$ are differentiable and $g' = e^x \neq 0$ on an open interval that contains 0. $\lim_{x\to 0} f(x) = 0$ and $\lim_{x\to 0} g(x) = 0$, so we have the indeterminate form $\frac{0}{0}$ and can apply l'Hospital's Rule.

$$\lim_{x\to 0}\frac{f(x)}{e^x - 1} \overset{\text{H}}{=} \lim_{x\to 0}\frac{f'(x)}{e^x} = \frac{1}{1} = 1$$

Note that $\lim_{x\to 0} f'(x) = 1$ since the graph of f has the same slope as the line $y = x$ at $x = 0$.

8. This limit has the form $\frac{0}{0}$. $\lim_{x\to 3}\frac{x-3}{x^2-9} = \lim_{x\to 3}\frac{x-3}{(x+3)(x-3)} = \lim_{x\to 3}\frac{1}{x+3} = \frac{1}{3+3} = \frac{1}{6}$

Note: Alternatively, we could apply l'Hospital's Rule.

9. This limit has the form $\frac{0}{0}$. $\lim_{x\to 4}\frac{x^2-2x-8}{x-4} = \lim_{x\to 4}\frac{(x-4)(x+2)}{x-4} = \lim_{x\to 4}(x+2) = 4+2 = 6$

Note: Alternatively, we could apply l'Hospital's Rule.

10. This limit has the form $\frac{0}{0}$. $\lim_{x\to -2}\frac{x^3+8}{x+2} \overset{\text{H}}{=} \lim_{x\to -2}\frac{3x^2}{1} = 3(-2)^2 = 12$

Note: Alternatively, we could factor and simplify.

11. This limit has the form $\frac{0}{0}$. $\lim_{x\to 1}\frac{x^7-1}{x^3-1} \overset{\text{H}}{=} \lim_{x\to 1}\frac{7x^6}{3x^2} = \frac{7}{3}$

Note: Alternatively, we could factor and simplify.

12. This limit has the form $\frac{0}{0}$.

$$\lim_{x\to 4}\frac{\sqrt{x}-2}{x-4} = \lim_{x\to 4}\frac{\sqrt{x}-2}{x-4}\cdot\frac{\sqrt{x}+2}{\sqrt{x}+2} = \lim_{x\to 4}\frac{x-4}{(x-4)\left(\sqrt{x}+2\right)} = \lim_{x\to 4}\frac{1}{\sqrt{x}+2} = \frac{1}{\sqrt{4}+2} = \frac{1}{4}$$

Note: Alternatively, we could apply l'Hospital's Rule.

13. This limit has the form $\frac{0}{0}$. $\lim_{x\to \pi/4}\frac{\sin x - \cos x}{\tan x - 1} \overset{\text{H}}{=} \lim_{x\to \pi/4}\frac{\cos x + \sin x}{\sec^2 x} = \lim_{x\to 4}\frac{\frac{\sqrt{2}}{2}+\frac{\sqrt{2}}{2}}{\left(\sqrt{2}\right)^2} = \frac{\sqrt{2}}{2}$

14. This limit has the form $\frac{0}{0}$. $\lim_{x\to 0}\frac{\tan 3x}{\sin 2x} \overset{\text{H}}{=} \lim_{x\to 0}\frac{3\sec^2 3x}{2\cos 2x} = \frac{3(1)^2}{2(1)} = \frac{3}{2}$

15. This limit has the form $\frac{0}{0}$. $\lim_{t\to 0}\frac{e^{2t}-1}{\sin t} \overset{\text{H}}{=} \lim_{t\to 0}\frac{2e^{2t}}{\cos t} = \frac{2(1)}{1} = 2$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. This limit has the form $\frac{0}{0}$. $\lim\limits_{x\to 0} \dfrac{x^2}{1-\cos x} \overset{\text{H}}{=} \lim\limits_{x\to 0} \dfrac{2x}{\sin x} = \lim\limits_{x\to 0} \dfrac{2}{(\sin x)/x} = \dfrac{2}{1} = 2$

17. This limit has the form $\frac{0}{0}$. $\lim\limits_{x\to 1} \dfrac{\sin(x-1)}{x^3+x-2} \overset{\text{H}}{=} \lim\limits_{x\to 1} \dfrac{\cos(x-1)}{3x^2+1} = \dfrac{\cos 0}{3(1)^2+1} = \dfrac{1}{4}$

18. The limit can be evaluated by substituting π for θ. $\lim\limits_{\theta\to\pi} \dfrac{1+\cos\theta}{1-\cos\theta} = \dfrac{1+(-1)}{1-(-1)} = \dfrac{0}{2} = 0$

19. This limit has the form $\frac{\infty}{\infty}$. $\lim\limits_{x\to\infty} \dfrac{\sqrt{x}}{1+e^x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{\frac{1}{2\sqrt{x}}}{e^x} = \lim\limits_{x\to\infty} \dfrac{1}{e^x \cdot 2\sqrt{x}} = 0$

20. This limit has the form $\frac{\infty}{\infty}$. $\lim\limits_{x\to\infty} \dfrac{x+x^2}{1-2x^2} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1+2x}{-4x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{2}{-4} = -\dfrac{1}{2}$.

A better method is to divide the numerator and the denominator by x^2: $\lim\limits_{x\to\infty} \dfrac{x+x^2}{1-2x^2} = \lim\limits_{x\to\infty} \dfrac{\frac{1}{x}+1}{\frac{1}{x^2}-2} = \dfrac{0+1}{0-2} = -\dfrac{1}{2}$.

21. $\lim\limits_{x\to 0^+} [(\ln x)/x] = -\infty$ since $\ln x \to -\infty$ as $x \to 0^+$ and dividing by small values of x just increases the magnitude of the quotient $(\ln x)/x$. L'Hospital's Rule does not apply.

22. This limit has the form $\frac{\infty}{\infty}$. $\lim\limits_{x\to\infty} \dfrac{\ln\sqrt{x}}{x^2} = \lim\limits_{x\to\infty} \dfrac{\frac{1}{2}\ln x}{x^2} \overset{\text{H}}{=} \lim\limits_{x\to\infty} = \dfrac{\frac{1}{2x}}{2x} = \lim\limits_{x\to\infty} \dfrac{1}{4x^2} = 0$

23. This limit has the form $\frac{0}{0}$. $\lim\limits_{x\to 3} \dfrac{\ln(x/3)}{3-x} \overset{\text{H}}{=} \lim\limits_{x\to 3} \dfrac{1/(x/3)\cdot(1/3)}{-1} = \lim\limits_{x\to 3} \left(-\dfrac{1}{x}\right) = -\dfrac{1}{3}$

24. This limit has the form $\frac{0}{0}$. $\lim\limits_{t\to 0} \dfrac{8^t-5^t}{t} \overset{\text{H}}{=} \lim\limits_{t\to 0} \dfrac{8^t\ln 8 - 5^t\ln 5}{1} = \ln 8 - \ln 5 = \ln\dfrac{8}{5}$

25. This limit has the form $\frac{0}{0}$.

$$\lim_{x\to 0} \frac{\sqrt{1+2x}-\sqrt{1-4x}}{x} \overset{\text{H}}{=} \lim_{x\to 0} \frac{\frac{1}{2}(1+2x)^{-1/2}\cdot 2 - \frac{1}{2}(1-4x)^{-1/2}(-4)}{1}$$

$$= \lim_{x\to 0}\left(\frac{1}{\sqrt{1+2x}} + \frac{2}{\sqrt{1-4x}}\right) = \frac{1}{\sqrt{1}} + \frac{2}{\sqrt{1}} = 3$$

26. This limit has the form $\frac{\infty}{\infty}$.

$$\lim_{u\to\infty} \frac{e^{u/10}}{u^3} \overset{\text{H}}{=} \lim_{u\to\infty} \frac{e^{u/10}\cdot\frac{1}{10}}{3u^2} \overset{\text{H}}{=} \frac{1}{30}\lim_{u\to\infty} \frac{e^{u/10}\cdot\frac{1}{10}}{2u} \overset{\text{H}}{=} \frac{1}{600}\lim_{u\to\infty} \frac{e^{u/10}\cdot\frac{1}{10}}{1} = \frac{1}{6000}\lim_{u\to\infty} e^{u/10} = \infty$$

27. This limit has the form $\frac{0}{0}$. $\lim\limits_{x\to 0} \dfrac{e^x+e^{-x}-2}{e^x-x-1} \overset{\text{H}}{=} \lim\limits_{x\to 0} \dfrac{e^x-e^{-x}}{e^x-1} \overset{\text{H}}{=} \lim\limits_{x\to 0} \dfrac{e^x+e^{-x}}{e^x} = \dfrac{1+1}{1} = 2$

28. This limit has the form $\frac{0}{0}$. $\lim\limits_{x\to 0} \dfrac{\sinh x - x}{x^3} \overset{\text{H}}{=} \lim\limits_{x\to 0} \dfrac{\cosh x - 1}{3x^2} \overset{\text{H}}{=} \lim\limits_{x\to 0} \dfrac{\sinh x}{6x} \overset{\text{H}}{=} \lim\limits_{x\to 0} \dfrac{\cosh x}{6} = \dfrac{1}{6}$

29. This limit has the form $\frac{0}{0}$. $\lim\limits_{x\to 0} \dfrac{\tanh x}{\tan x} \overset{\text{H}}{=} \lim\limits_{x\to 0} \dfrac{\operatorname{sech}^2 x}{\sec^2 x} = \dfrac{\operatorname{sech}^2 0}{\sec^2 0} = \dfrac{1}{1} = 1$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. This limit has the form $\frac{0}{0}$.

$$\lim_{x\to 0}\frac{x-\sin x}{x-\tan x}\overset{\text{H}}{=}\lim_{x\to 0}\frac{1-\cos x}{1-\sec^2 x}\overset{\text{H}}{=}\lim_{x\to 0}\frac{-(-\sin x)}{-2\sec x\,(\sec x\,\tan x)}=-\frac{1}{2}\lim_{x\to 0}\frac{\sin x\left(\dfrac{\cos x}{\sin x}\right)}{\sec^2 x}$$
$$=-\tfrac{1}{2}\lim_{x\to 0}\cos^3 x=-\tfrac{1}{2}(1)^3=-\tfrac{1}{2}$$

Another method is to write the limit as $\displaystyle\lim_{x\to 0}\frac{1-\dfrac{\sin x}{x}}{1-\dfrac{\tan x}{x}}$.

31. This limit has the form $\frac{0}{0}$. $\displaystyle\lim_{x\to 0}\frac{\sin^{-1}x}{x}\overset{\text{H}}{=}\lim_{x\to 0}\frac{1/\sqrt{1-x^2}}{1}=\lim_{x\to 0}\frac{1}{\sqrt{1-x^2}}=\frac{1}{1}=1$

32. This limit has the form $\frac{\infty}{\infty}$. $\displaystyle\lim_{x\to\infty}\frac{(\ln x)^2}{x}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{2(\ln x)(1/x)}{1}=2\lim_{x\to\infty}\frac{\ln x}{x}\overset{\text{H}}{=}2\lim_{x\to\infty}\frac{1/x}{1}=2(0)=0$

33. This limit has the form $\frac{0}{0}$. $\displaystyle\lim_{x\to 0}\frac{x3^x}{3^x-1}\overset{\text{H}}{=}\lim_{x\to 0}\frac{x3^x\ln 3+3^x}{3^x\ln 3}=\lim_{x\to 0}\frac{3^x(x\ln 3+1)}{3^x\ln 3}=\lim_{x\to 0}\frac{x\ln 3+1}{\ln 3}=\frac{1}{\ln 3}$

34. This limit has the form $\frac{0}{0}$.

$$\lim_{x\to 0}\frac{e^x+e^{-x}-2\cos x}{x\sin x}\overset{\text{H}}{=}\lim_{x\to 0}\frac{e^x-e^{-x}+2\sin x}{x\cos x+\sin x}\overset{\text{H}}{=}\lim_{x\to 0}\frac{e^x+e^{-x}+2\cos x}{-x\sin x+\cos x+\cos x}=\frac{1+1+2}{0+1+1}=\frac{4}{2}=2$$

35. This limit can be evaluated by substituting 0 for x. $\displaystyle\lim_{x\to 0}\frac{\ln(1+x)}{\cos x+e^x-1}=\frac{\ln 1}{1+1-1}=\frac{0}{1}=0$

36. This limit has the form $\frac{0}{0}$. $\displaystyle\lim_{x\to 1}\frac{x\sin(x-1)}{2x^2-x-1}\overset{\text{H}}{=}\lim_{x\to 1}\frac{x\cos(x-1)+\sin(x-1)}{4x-1}=\frac{\cos 0}{4-1}=\frac{1}{3}$

37. This limit has the form $\frac{0}{\infty}$, so l'Hospital's Rule doesn't apply. As $x\to 0^+$, $\arctan 2x\to 0$ and $\ln x\to-\infty$,

so $\displaystyle\lim_{x\to 0^+}\frac{\arctan 2x}{\ln x}=0$.

38. This limit has the form $\frac{0}{0}$.

$$\lim_{x\to 0}\frac{x^2\sin x}{\sin x-x}\overset{\text{H}}{=}\lim_{x\to 0}\frac{x^2\cos x+2x\sin x}{\cos x-1}\overset{\text{H}}{=}\lim_{x\to 0}\frac{-x^2\sin x+2x\cos x+2x\cos x+2\sin x}{-\sin x}$$
$$=\lim_{x\to 0}\frac{(2-x^2)\sin x+4x\cos x}{-\sin x}$$
$$\overset{\text{H}}{=}\lim_{x\to 0}\frac{(2-x^2)\cos x-2x\sin x-4x\sin x+4\cos x}{-\cos x}=\frac{2(1)-0-0+4(1)}{-1}=-6$$

39. This limit has the form $\frac{0}{0}$. $\displaystyle\lim_{x\to 1}\frac{x^a-1}{x^b-1}$ [for $b\neq 0$] $\displaystyle\overset{\text{H}}{=}\lim_{x\to 1}\frac{ax^{a-1}}{bx^{b-1}}=\frac{a(1)}{b(1)}=\frac{a}{b}$

40. This limit has the form $\frac{0}{0}$. $\displaystyle\lim_{x\to\infty}\frac{e^{-x}}{(\pi/2)-\tan^{-1}x}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{-e^{-x}}{-\dfrac{1}{1+x^2}}=\lim_{x\to\infty}\frac{1+x^2}{e^x}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{2x}{e^x}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{2}{e^x}=0$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. This limit has the form $\frac{0}{0}$.

$$\lim_{x\to 0}\frac{\cos x-1+\frac{1}{2}x^2}{x^4}\overset{\text{H}}{=}\lim_{x\to 0}\frac{-\sin x+x}{4x^3}\overset{\text{H}}{=}\lim_{x\to 0}\frac{-\cos x+1}{12x^2}\overset{\text{H}}{=}\lim_{x\to 0}\frac{\sin x}{24x}\overset{\text{H}}{=}\lim_{x\to 0}\frac{\cos x}{24}=\frac{1}{24}$$

42. This limit has the form $\frac{0}{0}$.

$$\lim_{x\to 0}\frac{x-\sin x}{x\sin(x^2)}\overset{\text{H}}{=}\lim_{x\to 0}\frac{1-\cos x}{x\cos(x^2)\cdot 2x+\sin(x^2)}=\lim_{x\to 0}\frac{1-\cos x}{2x^2\cos(x^2)+\sin(x^2)}$$

$$\overset{\text{H}}{=}\lim_{x\to 0}\frac{\sin x}{2x^2\left[-\sin(x^2)\cdot 2x\right]+\cos(x^2)\cdot 4x+\cos(x^2)\cdot 2x}=\lim_{x\to 0}\frac{\sin x}{6x\cos(x^2)-4x^3\sin(x^2)}$$

$$\overset{\text{H}}{=}\lim_{x\to 0}\frac{\cos x}{6x\left[-\sin(x^2)\cdot 2x\right]+\cos(x^2)\cdot 6-4x^3\cos(x^2)\cdot 2x-\sin(x^2)\cdot 12x^2}=\frac{1}{0+1\cdot 6-0-0}=\frac{1}{6}$$

43. This limit has the form $\infty\cdot 0$. We'll change it to the form $\frac{0}{0}$.

$$\lim_{x\to\infty}x\sin(\pi/x)=\lim_{x\to\infty}\frac{\sin(\pi/x)}{1/x}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{\cos(\pi/x)(-\pi/x^2)}{-1/x^2}=\pi\lim_{x\to\infty}\cos(\pi/x)=\pi(1)=\pi$$

44. This limit has the form $\infty\cdot 0$. We'll change it to the form $\frac{\infty}{\infty}$.

$$\lim_{x\to\infty}\sqrt{x}\,e^{-x/2}=\lim_{x\to\infty}\frac{\sqrt{x}}{e^{x/2}}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{\frac{1}{2}x^{-1/2}}{\frac{1}{2}e^{x/2}}=\lim_{x\to\infty}\frac{1}{\sqrt{x}\,e^{x/2}}=0$$

45. This limit has the form $0\cdot\infty$. We'll change it to the form $\frac{0}{0}$. $\displaystyle\lim_{x\to 0}\sin 5x\csc 3x=\lim_{x\to 0}\frac{\sin 5x}{\sin 3x}\overset{\text{H}}{=}\lim_{x\to 0}\frac{5\cos 5x}{3\cos 3x}=\frac{5\cdot 1}{3\cdot 1}=\frac{5}{3}$

46. This limit has the form $(-\infty)\cdot 0$.

$$\lim_{x\to-\infty}x\ln\left(1-\frac{1}{x}\right)=\lim_{x\to-\infty}\frac{\ln\left(1-\frac{1}{x}\right)}{\frac{1}{x}}\overset{\text{H}}{=}\lim_{x\to-\infty}\frac{\frac{1}{1-1/x}\cdot\frac{1}{x^2}}{-\frac{1}{x^2}}=\lim_{x\to-\infty}\frac{-1}{1-\frac{1}{x}}=\frac{-1}{1}=-1$$

47. This limit has the form $\infty\cdot 0$. $\displaystyle\lim_{x\to\infty}x^3e^{-x^2}=\lim_{x\to\infty}\frac{x^3}{e^{x^2}}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{3x^2}{2xe^{x^2}}=\lim_{x\to\infty}\frac{3x}{2e^{x^2}}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{3}{4xe^{x^2}}=0$

48. This limit has the form $\infty\cdot 0$.

$$\lim_{x\to\infty}x^{3/2}\sin(1/x)=\lim_{x\to\infty}x^{1/2}\cdot\frac{\sin(1/x)}{1/x}=\lim_{t\to 0^+}\frac{1}{\sqrt{t}}\frac{\sin t}{t}\quad[\text{where } t=1/x]\quad=\infty \text{ since as } t\to 0^+,\ \frac{1}{\sqrt{t}}\to\infty$$

and $\dfrac{\sin t}{t}\to 1$.

49. This limit has the form $0\cdot(-\infty)$.

$$\lim_{x\to 1^+}\ln x\,\tan(\pi x/2)=\lim_{x\to 1^+}\frac{\ln x}{\cot(\pi x/2)}\overset{\text{H}}{=}\lim_{x\to 1^+}\frac{1/x}{(-\pi/2)\csc^2(\pi x/2)}=\frac{1}{(-\pi/2)(1)^2}=-\frac{2}{\pi}$$

50. This limit has the form $0\cdot\infty$. $\displaystyle\lim_{x\to(\pi/2)^-}\cos x\,\sec 5x=\lim_{x\to(\pi/2)^-}\frac{\cos x}{\cos 5x}\overset{\text{H}}{=}\lim_{x\to(\pi/2)^-}\frac{-\sin x}{-5\sin 5x}=\frac{-1}{-5}=\frac{1}{5}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. This limit has the form $\infty - \infty$.

$$\lim_{x\to 1}\left(\frac{x}{x-1} - \frac{1}{\ln x}\right) = \lim_{x\to 1}\frac{x\ln x - (x-1)}{(x-1)\ln x} \overset{\text{H}}{=} \lim_{x\to 1}\frac{x(1/x) + \ln x - 1}{(x-1)(1/x) + \ln x} = \lim_{x\to 1}\frac{\ln x}{1 - (1/x) + \ln x}$$

$$\overset{\text{H}}{=} \lim_{x\to 1}\frac{1/x}{1/x^2 + 1/x}\cdot\frac{x^2}{x^2} = \lim_{x\to 1}\frac{x}{1+x} = \frac{1}{1+1} = \frac{1}{2}$$

52. This limit has the form $\infty - \infty$. $\lim_{x\to 0}(\csc x - \cot x) = \lim_{x\to 0}\left(\frac{1}{\sin x} - \frac{\cos x}{\sin x}\right) = \lim_{x\to 0}\frac{1-\cos x}{\sin x} \overset{\text{H}}{=} \lim_{x\to 0}\frac{\sin x}{\cos x} = 0$

53. This limit has the form $\infty - \infty$.

$$\lim_{x\to 0^+}\left(\frac{1}{x} - \frac{1}{e^x - 1}\right) = \lim_{x\to 0^+}\frac{e^x - 1 - x}{x(e^x - 1)} \overset{\text{H}}{=} \lim_{x\to 0^+}\frac{e^x - 1}{xe^x + e^x - 1} \overset{\text{H}}{=} \lim_{x\to 0^+}\frac{e^x}{xe^x + e^x + e^x} = \frac{1}{0+1+1} = \frac{1}{2}$$

54. This limit has the form $\infty - \infty$.

$$\lim_{x\to 0^+}\left(\frac{1}{x} - \frac{1}{\tan^{-1} x}\right) = \lim_{x\to 0^+}\frac{\tan^{-1} x - x}{x\tan^{-1} x} \overset{\text{H}}{=} \lim_{x\to 0^+}\frac{1/(1+x^2) - 1}{x/(1+x^2) + \tan^{-1} x} = \lim_{x\to 0^+}\frac{1-(1+x^2)}{x + (1+x^2)\tan^{-1} x}$$

$$= \lim_{x\to 0^+}\frac{-x^2}{x + (1+x^2)\tan^{-1} x} \overset{\text{H}}{=} \lim_{x\to 0^+}\frac{-2x}{1 + (1+x^2)(1/(1+x^2)) + (\tan^{-1} x)(2x)}$$

$$= \lim_{x\to 0^+}\frac{-2x}{2 + 2x\tan^{-1} x} = \frac{0}{2+0} = 0$$

55. This limit has the form $\infty - \infty$.

$$\lim_{x\to 0^+}\left(\frac{1}{x} - \frac{1}{\tan x}\right) = \lim_{x\to 0^+}\frac{\tan x - x}{x\tan x} \overset{\text{H}}{=} \lim_{x\to 0^+}\frac{\sec^2 x - 1}{x\sec^2 x + \tan x} \overset{\text{H}}{=} \lim_{x\to 0^+}\frac{2\sec x\cdot\sec x\tan x}{x\cdot 2\sec x\cdot\sec x\tan x + \sec^2 x + \sec^2 x}$$

$$= \frac{0}{0+1+1} = 0$$

56. The limit has the form $\infty - \infty$ and we will change the form to a product by factoring out x.

$$\lim_{x\to\infty}(x - \ln x) = \lim_{x\to\infty} x\left(1 - \frac{\ln x}{x}\right) = \infty \text{ since } \lim_{x\to\infty}\frac{\ln x}{x} \overset{\text{H}}{=} \lim_{x\to\infty}\frac{1/x}{1} = 0.$$

57. $y = x^{\sqrt{x}} \quad\Rightarrow\quad \ln y = \sqrt{x}\,\ln x$, so

$$\lim_{x\to 0^+}\ln y = \lim_{x\to 0^+}\sqrt{x}\,\ln x = \lim_{x\to 0^+}\frac{\ln x}{x^{-1/2}} \overset{\text{H}}{=} \lim_{x\to 0^+}\frac{1/x}{-\frac{1}{2}x^{-3/2}} = -2\lim_{x\to 0^+}\sqrt{x} = 0 \quad\Rightarrow$$

$$\lim_{x\to 0^+} x^{\sqrt{x}} = \lim_{x\to 0^+} e^{\ln y} = e^0 = 1.$$

58. $y = (\tan 2x)^x \quad\Rightarrow\quad \ln y = x\cdot\ln\tan 2x$, so

$$\lim_{x\to 0^+}\ln y = \lim_{x\to 0^+} x\cdot\ln\tan 2x = \lim_{x\to 0^+}\frac{\ln\tan 2x}{1/x} \overset{\text{H}}{=} \lim_{x\to 0^+}\frac{(1/\tan 2x)(2\sec^2 2x)}{-1/x^2} = \lim_{x\to 0^+}\frac{-2x^2\cos 2x}{\sin 2x\,\cos^2 2x}$$

$$= \lim_{x\to 0^+}\frac{2x}{\sin 2x}\cdot\lim_{x\to 0^+}\frac{-x}{\cos 2x} = 1\cdot 0 = 0 \quad\Rightarrow$$

$$\lim_{x\to 0^+}(\tan 2x)^x = \lim_{x\to 0^+} e^{\ln y} = e^0 = 1.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. $y = (1-2x)^{1/x} \quad \Rightarrow \quad \ln y = \dfrac{1}{x}\ln(1-2x)$, so $\displaystyle\lim_{x\to 0}\ln y = \lim_{x\to 0}\frac{\ln(1-2x)}{x} \overset{\text{H}}{=} \lim_{x\to 0}\frac{-2/(1-2x)}{1} = -2 \quad \Rightarrow$

$\displaystyle\lim_{x\to 0}(1-2x)^{1/x} = \lim_{x\to 0} e^{\ln y} = e^{-2}$.

60. $y = \left(1+\dfrac{a}{x}\right)^{bx} \quad \Rightarrow \quad \ln y = bx\ln\left(1+\dfrac{a}{x}\right)$, so

$$\lim_{x\to\infty}\ln y = \lim_{x\to\infty}\frac{b\ln(1+a/x)}{1/x} \overset{\text{H}}{=} \lim_{x\to\infty}\frac{b\left(\dfrac{1}{1+a/x}\right)\left(-\dfrac{a}{x^2}\right)}{-1/x^2} = \lim_{x\to\infty}\frac{ab}{1+a/x} = ab \quad \Rightarrow$$

$$\lim_{x\to\infty}\left(1+\frac{a}{x}\right)^{bx} = \lim_{x\to\infty} e^{\ln y} = e^{ab}.$$

61. $y = x^{1/(1-x)} \quad \Rightarrow \quad \ln y = \dfrac{1}{1-x}\ln x$, so $\displaystyle\lim_{x\to 1^+}\ln y = \lim_{x\to 1^+}\frac{1}{1-x}\ln x = \lim_{x\to 1^+}\frac{\ln x}{1-x} \overset{\text{H}}{=} \lim_{x\to 1^+}\frac{1/x}{-1} = -1 \quad \Rightarrow$

$\displaystyle\lim_{x\to 1^+} x^{1/(1-x)} = \lim_{x\to 1^+} e^{\ln y} = e^{-1} = \frac{1}{e}$.

62. $y = (e^x+10x)^{1/x} \quad \Rightarrow \quad \ln y = \dfrac{1}{x}\ln(e^x+10x)$, so

$$\lim_{x\to\infty}\ln y = \lim_{x\to\infty}\frac{1}{x}\ln(e^x+10x) = \lim_{x\to\infty}\frac{\ln(e^x+10x)}{x} \overset{\text{H}}{=} \lim_{x\to\infty}\frac{\dfrac{1}{e^x+10x}\cdot(e^x+10)}{1}$$

$$= \lim_{x\to\infty}\frac{e^x+10}{e^x+10x} \overset{\text{H}}{=} \lim_{x\to\infty}\frac{e^x}{e^x+10} \overset{\text{H}}{=} \lim_{x\to\infty}\frac{e^x}{e^x} = \lim_{x\to\infty}(1) = 1 \quad \Rightarrow$$

$\displaystyle\lim_{x\to\infty}(e^x+10x)^{1/x} = \lim_{x\to\infty} e^{\ln y} = e^1 = e$

63. $y = x^{1/x} \quad \Rightarrow \quad \ln y = (1/x)\ln x \quad \Rightarrow \quad \displaystyle\lim_{x\to\infty}\ln y = \lim_{x\to\infty}\frac{\ln x}{x} \overset{\text{H}}{=} \lim_{x\to\infty}\frac{1/x}{1} = 0 \quad \Rightarrow$

$\displaystyle\lim_{x\to\infty} x^{1/x} = \lim_{x\to\infty} e^{\ln y} = e^0 = 1$

64. $y = x^{e^{-x}} \quad \Rightarrow \quad \ln y = e^{-x}\ln x \quad \Rightarrow \quad \displaystyle\lim_{x\to\infty}\ln y = \lim_{x\to\infty}\frac{\ln x}{e^x} \overset{\text{H}}{=} \lim_{x\to\infty}\frac{1/x}{e^x} = \lim_{x\to\infty}\frac{1}{xe^x} = 0 \quad \Rightarrow$

$\displaystyle\lim_{x\to\infty} x^{e^{-x}} = \lim_{x\to\infty} e^{\ln y} = e^0 = 1$

65. $y = (4x+1)^{\cot x} \quad \Rightarrow \quad \ln y = \cot x\,\ln(4x+1)$, so $\displaystyle\lim_{x\to 0^+}\ln y = \lim_{x\to 0^+}\frac{\ln(4x+1)}{\tan x} \overset{\text{H}}{=} \lim_{x\to 0^+}\frac{\dfrac{4}{4x+1}}{\sec^2 x} = 4 \quad \Rightarrow$

$\displaystyle\lim_{x\to 0^+}(4x+1)^{\cot x} = \lim_{x\to 0^+} e^{\ln y} = e^4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

66. $y = (1-\cos x)^{\sin x} \quad \Rightarrow \quad \ln y = \sin x \ln(1-\cos x)$, so

$$\lim_{x\to 0^+} \ln y = \lim_{x\to 0^+} \sin x \ln(1-\cos x) = \lim_{x\to 0^+} \frac{\ln(1-\cos x)}{\csc x} \overset{\text{H}}{=} \lim_{x\to 0^+} \frac{\dfrac{1}{1-\cos x}\cdot \sin x}{-\csc x \cot x}$$

$$= -\lim_{x\to 0^+} \frac{\sin x}{(1-\cos x)\csc x \cot x} = -\lim_{x\to 0^+} \frac{\sin x}{\csc x \cot x - \cot^2 x}\cdot\left(\frac{\sin^2 x}{\sin^2 x}\right)$$

$$= -\lim_{x\to 0^+} \frac{\sin^3 x}{\cos x - \cos^2 x} = -\lim_{x\to 0^+} \frac{\sin^3 x}{(1-\cos x)\cos x}$$

$$\overset{\text{H}}{=} -\lim_{x\to 0^+} \frac{3\sin^2 x \cos x}{(1-\cos x)(-\sin x) + \cos x\,(\sin x)} = -\lim_{x\to 0^+} \frac{3\sin x \cos x}{(\cos x - 1) + \cos x} = -\frac{0}{0+1} = 0 \quad \Rightarrow$$

$$\lim_{x\to 0^+} (1-\cos x)^{\sin x} = \lim_{x\to 0^+} e^{\ln y} = e^0 = 1$$

67. $y = (1+\sin 3x)^{1/x} \quad \Rightarrow \quad \ln y = \dfrac{1}{x}\ln(1+\sin 3x) \quad \Rightarrow$

$$\lim_{x\to 0^+} \ln y = \lim_{x\to 0^+} \frac{\ln(1+\sin 3x)}{x} \overset{\text{H}}{=} \lim_{x\to 0^+} \frac{[1/(1+\sin 3x)]\cdot 3\cos 3x}{1} = \lim_{x\to 0^+} \frac{3\cos 3x}{1+\sin 3x} = \frac{3\cdot 1}{1+0} = 3 \quad \Rightarrow$$

$$\lim_{x\to 0^+} (1+\sin 3x)^{1/x} = \lim_{x\to 0^+} e^{\ln y} = e^3$$

68. $y = (\cos x)^{1/x^2} \quad \Rightarrow \quad \ln y = \dfrac{1}{x^2}\ln\cos x$, so

$$\lim_{x\to 0} \ln y = \lim_{x\to 0} \frac{1}{x^2}\ln\cos x = \lim_{x\to 0} \frac{\ln\cos x}{x^2} \overset{\text{H}}{=} \lim_{x\to 0} \frac{\dfrac{1}{\cos x}\cdot(-\sin x)}{2x} = \lim_{x\to 0} \frac{-\tan x}{2x} \overset{\text{H}}{=} \lim_{x\to 0} \frac{-\sec^2 x}{2} = -\frac{1}{2} \quad \Rightarrow$$

$$\lim_{x\to 0} (\cos x)^{1/x^2} = \lim_{x\to 0} e^{\ln y} = e^{-1/2} = \frac{1}{\sqrt{e}}$$

69. The given limit is $\displaystyle\lim_{x\to 0^+} \frac{x^x - 1}{\ln x + x - 1}$. Note that $y = x^x \quad \Rightarrow \quad \ln y = x\ln x$, so

$$\lim_{x\to 0^+} \ln y = \lim_{x\to 0^+} x\ln x = \lim_{x\to 0^+} \frac{\ln x}{\dfrac{1}{x}} \overset{\text{H}}{=} \lim_{x\to 0^+} \frac{\dfrac{1}{x}}{-\dfrac{1}{x^2}} = \lim_{x\to 0^+} (-x) = 0 \quad \Rightarrow \quad \lim_{x\to 0^+} x^x = \lim_{x\to 0^+} e^{\ln y} = e^0 = 1.$$

Therefore, the numerator of the given limit has limit $1 - 1 = 0$ as $x \to 0^+$. The denominator of the given limit $\to -\infty$ as $x \to 0^+$ since $\ln x \to -\infty$ as $x \to 0^+$. Thus, $\displaystyle\lim_{x\to 0^+} \frac{x^x - 1}{\ln x + x - 1} = 0$.

70. $y = \left(\dfrac{2x-3}{2x+5}\right)^{2x+1} \quad \Rightarrow \quad \ln y = (2x+1)\,\ln\left(\dfrac{2x-3}{2x+5}\right) \quad \Rightarrow$

$$\lim_{x\to\infty} \ln y = \lim_{x\to\infty} \frac{\ln(2x-3) - \ln(2x+5)}{1/(2x+1)} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{2/(2x-3) - 2/(2x+5)}{-2/(2x+1)^2} = \lim_{x\to\infty} \frac{-8(2x+1)^2}{(2x-3)(2x+5)}$$

$$= \lim_{x\to\infty} \frac{-8(2+1/x)^2}{(2-3/x)(2+5/x)} = -8 \quad \Rightarrow \quad \lim_{x\to\infty} \left(\frac{2x-3}{2x+5}\right)^{2x+1} = \lim_{x\to\infty} e^{\ln y} = e^{-8}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

71.

From the graph, if $x = 500$, $y \approx 7.36$. The limit has the form 1^∞.

Now $y = \left(1 + \dfrac{2}{x}\right)^x \quad \Rightarrow \quad \ln y = x \ln\left(1 + \dfrac{2}{x}\right) \quad \Rightarrow$

$$\lim_{x\to\infty} \ln y = \lim_{x\to\infty} \frac{\ln(1+2/x)}{1/x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{\dfrac{1}{1+2/x}\left(-\dfrac{2}{x^2}\right)}{-1/x^2}$$

$$= 2 \lim_{x\to\infty} \frac{1}{1+2/x} = 2(1) = 2 \quad \Rightarrow$$

$$\lim_{x\to\infty} \left(1 + \frac{2}{x}\right)^x = \lim_{x\to\infty} e^{\ln y} = e^2 \ \ [\approx 7.39]$$

72.

From the graph, as $x \to 0$, $y \approx 0.55$. The limit has the form $\frac{0}{0}$.

$$\lim_{x\to 0} \frac{5^x - 4^x}{3^x - 2^x} \overset{\text{H}}{=} \lim_{x\to 0} \frac{5^x \ln 5 - 4^x \ln 4}{3^x \ln 3 - 2^x \ln 2} = \frac{\ln 5 - \ln 4}{\ln 3 - \ln 2} = \frac{\ln \frac{5}{4}}{\ln \frac{3}{2}} \ \ [\approx 0.55]$$

73.

From the graph, it appears that $\displaystyle\lim_{x\to 0} \frac{f(x)}{g(x)} = \lim_{x\to 0} \frac{f'(x)}{g'(x)} = 0.25$.

We calculate $\displaystyle\lim_{x\to 0} \frac{f(x)}{g(x)} = \lim_{x\to 0} \frac{e^x - 1}{x^3 + 4x} \overset{\text{H}}{=} \lim_{x\to 0} \frac{e^x}{3x^2 + 4} = \frac{1}{4}$.

74.

From the graph, it appears that $\displaystyle\lim_{x\to 0} \frac{f(x)}{g(x)} = \lim_{x\to 0} \frac{f'(x)}{g'(x)} = 4$. We calculate

$$\lim_{x\to 0} \frac{f(x)}{g(x)} = \lim_{x\to 0} \frac{2x \sin x}{\sec x - 1} \overset{\text{H}}{=} \lim_{x\to 0} \frac{2(x \cos x + \sin x)}{\sec x \ \tan x}$$

$$\overset{\text{H}}{=} \lim_{x\to 0} \frac{2(-x \sin x + \cos x + \cos x)}{\sec x(\sec^2 x) + \tan x(\sec x \ \tan x)} = \frac{4}{1} = 4$$

75. $\displaystyle\lim_{x\to\infty} \frac{e^x}{x^n} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{e^x}{nx^{n-1}} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{e^x}{n(n-1)x^{n-2}} \overset{\text{H}}{=} \cdots \overset{\text{H}}{=} \lim_{x\to\infty} \frac{e^x}{n!} = \infty$

76. This limit has the form $\frac{\infty}{\infty}$. $\displaystyle\lim_{x\to\infty} \frac{\ln x}{x^p} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{1/x}{px^{p-1}} = \lim_{x\to\infty} \frac{1}{px^p} = 0$ since $p > 0$.

77. $\displaystyle\lim_{x\to\infty} \frac{x}{\sqrt{x^2+1}} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{1}{\frac{1}{2}(x^2+1)^{-1/2}(2x)} = \lim_{x\to\infty} \frac{\sqrt{x^2+1}}{x}$. Repeated applications of l'Hospital's Rule result in the original limit or the limit of the reciprocal of the function. Another method is to try dividing the numerator and denominator by x: $\displaystyle\lim_{x\to\infty} \frac{x}{\sqrt{x^2+1}} = \lim_{x\to\infty} \frac{x/x}{\sqrt{x^2/x^2 + 1/x^2}} = \lim_{x\to\infty} \frac{1}{\sqrt{1 + 1/x^2}} = \frac{1}{1} = 1$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

78. $\lim\limits_{x\to(\pi/2)^-} \dfrac{\sec x}{\tan x} \overset{\text{H}}{=} \lim\limits_{x\to(\pi/2)^-} \dfrac{\sec x\,\tan x}{\sec^2 x} = \lim\limits_{x\to(\pi/2)^-} \dfrac{\tan x}{\sec x}$. Repeated applications of l'Hospital's Rule result in the original limit or the limit of the reciprocal of the function. Another method is to simplify first:

$$\lim_{x\to(\pi/2)^-} \frac{\sec x}{\tan x} = \lim_{x\to(\pi/2)^-} \frac{1/\cos x}{\sin x/\cos x} = \lim_{x\to(\pi/2)^-} \frac{1}{\sin x} = \frac{1}{1} = 1$$

79. $f(x) = e^x - cx \;\Rightarrow\; f'(x) = e^x - c = 0 \;\Leftrightarrow\; e^x = c \;\Leftrightarrow\; x = \ln c,\ c > 0$. $f''(x) = e^x > 0$, so f is CU on $(-\infty, \infty)$. $\lim\limits_{x\to\infty}(e^x - cx) = \lim\limits_{x\to\infty}\left[x\left(\dfrac{e^x}{x} - c\right)\right] = L_1$. Now $\lim\limits_{x\to\infty} \dfrac{e^x}{x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{e^x}{1} = \infty$, so $L_1 = \infty$, regardless of the value of c. For $L = \lim\limits_{x\to-\infty}(e^x - cx)$, $e^x \to 0$, so L is determined by $-cx$. If $c > 0$, $-cx \to \infty$, and $L = \infty$. If $c < 0$, $-cx \to -\infty$, and $L = -\infty$. Thus, f has an absolute minimum for $c > 0$. As c increases, the minimum points $(\ln c, c - c\ln c)$, get farther away from the origin.

80. (a) $\lim\limits_{t\to\infty} v = \lim\limits_{t\to\infty} \dfrac{mg}{c}\left(1 - e^{-ct/m}\right) = \dfrac{mg}{c} \lim\limits_{t\to\infty}\left(1 - e^{-ct/m}\right) = \dfrac{mg}{c}(1 - 0)$ [because $-ct/m \to -\infty$ as $t \to \infty$]

$= \dfrac{mg}{c}$, which is the speed the object approaches as time goes on, the so-called limiting velocity.

(b) $\lim\limits_{c\to0^+} v = \lim\limits_{c\to0^+} \dfrac{mg}{c}(1 - e^{-ct/m}) = mg \lim\limits_{c\to0^+} \dfrac{1 - e^{-ct/m}}{c}$ [form is $\frac{0}{0}$]

$$\overset{\text{H}}{=} mg \lim_{c\to0^+} \frac{(-e^{-ct/m})\cdot(-t/m)}{1} = \frac{mgt}{m} \lim_{c\to0^+} e^{-ct/m} = gt(1) = gt$$

The velocity of a falling object in a vacuum is directly proportional to the amount of time it falls.

81. First we will find $\lim\limits_{n\to\infty}\left(1 + \dfrac{r}{n}\right)^{nt}$, which is of the form 1^∞. $y = \left(1 + \dfrac{r}{n}\right)^{nt} \;\Rightarrow\; \ln y = nt\ln\left(1 + \dfrac{r}{n}\right)$, so

$$\lim_{n\to\infty} \ln y = \lim_{n\to\infty} nt\ln\left(1 + \frac{r}{n}\right) = t\lim_{n\to\infty} \frac{\ln(1 + r/n)}{1/n} \overset{\text{H}}{=} t\lim_{n\to\infty} \frac{(-r/n^2)}{(1 + r/n)(-1/n^2)} = t\lim_{n\to\infty} \frac{r}{1 + i/n} = tr \;\Rightarrow$$

$\lim\limits_{n\to\infty} y = e^{rt}$. Thus, as $n \to \infty$, $A = A_0\left(1 + \dfrac{r}{n}\right)^{nt} \to A_0 e^{rt}$.

82. (a) $r = 3,\ \rho = 0.05 \;\Rightarrow\; P = \dfrac{1 - 10^{-\rho r^2}}{\rho r^2 \ln 10} = \dfrac{1 - 10^{-0.45}}{0.45\ln 10} \approx 0.62$, or about 62%.

(b) $r = 2,\ \rho = 0.05 \;\Rightarrow\; P = \dfrac{1 - 10^{-0.2}}{0.2\ln 10} \approx 0.80$, or about 80%.

Yes, it makes sense. Since measured brightness decreases with light entering farther from the center of the pupil, a smaller pupil radius means that the average brightness measurements are higher than when including light entering at larger radii.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\lim\limits_{r\to 0^+} P = \lim\limits_{r\to 0^+} \dfrac{1-10^{-\rho r^2}}{\rho r^2 \ln 10} \overset{\text{H}}{=} \lim\limits_{r\to 0^+} \dfrac{-10^{-\rho r^2}(\ln 10)(-2\rho r)}{2\rho r(\ln 10)} = \lim\limits_{r\to 0^+} \dfrac{1}{10^{\rho r^2}} = 1$, or 100%.

We might expect that 100% of the brightness is sensed at the very center of the pupil, so a limit of 1 would make sense in this context if the radius r could approach 0. This result isn't physically possible because there are limitations on how small the pupil can shrink.

83. (a) $\lim\limits_{t\to\infty} P(t) = \lim\limits_{t\to\infty} \dfrac{M}{1+Ae^{-kt}} = \dfrac{M}{1+A\cdot 0} = M$

It is to be expected that a population that is growing will eventually reach the maximum population size that can be supported.

(b) $\lim\limits_{M\to\infty} P(t) = \lim\limits_{M\to\infty} \dfrac{M}{1+\dfrac{M-P_0}{P_0}e^{-kt}} = \lim\limits_{M\to\infty} \dfrac{M}{1+\left(\dfrac{M}{P_0}-1\right)e^{-kt}} \overset{\text{H}}{=} \lim\limits_{M\to\infty} \dfrac{1}{\dfrac{1}{P_0}e^{-kt}} = P_0e^{kt}$

P_0e^{kt} is an exponential function.

84. (a) $\lim\limits_{R\to r^+} v = \lim\limits_{R\to r^+} \left[-c\left(\dfrac{r}{R}\right)^2 \ln\left(\dfrac{r}{R}\right)\right] = -cr^2 \lim\limits_{R\to r^+} \left[\left(\dfrac{1}{R}\right)^2 \ln\left(\dfrac{r}{R}\right)\right] = -cr^2\cdot\dfrac{1}{r^2}\cdot\ln 1 = -c\cdot 0 = 0$

As the insulation of a metal cable becomes thinner, the velocity of an electrical impulse in the cable approaches zero.

(b) $\lim\limits_{r\to 0^+} v = \lim\limits_{r\to 0^+} \left[-c\left(\dfrac{r}{R}\right)^2 \ln\left(\dfrac{r}{R}\right)\right] = -\dfrac{c}{R^2} \lim\limits_{r\to 0^+} \left[r^2 \ln\left(\dfrac{r}{R}\right)\right]$ [form is $0\cdot\infty$]

$= -\dfrac{c}{R^2} \lim\limits_{r\to 0^+} \dfrac{\ln\left(\dfrac{r}{R}\right)}{\dfrac{1}{r^2}}$ [form is ∞/∞] $\overset{\text{H}}{=} -\dfrac{c}{R^2} \lim\limits_{r\to 0^+} \dfrac{\dfrac{R}{r}\cdot\dfrac{1}{R}}{\dfrac{-2}{r^3}} = -\dfrac{c}{R^2} \lim\limits_{r\to 0^+} \left(-\dfrac{r^2}{2}\right) = 0$

As the radius of the metal cable approaches zero, the velocity of an electrical impulse in the cable approaches zero.

85. We see that both numerator and denominator approach 0, so we can use l'Hospital's Rule:

$$\lim_{x\to a} \frac{\sqrt{2a^3x-x^4}-a\sqrt[3]{aax}}{a-\sqrt[4]{ax^3}} \overset{\text{H}}{=} \lim_{x\to a} \frac{\frac{1}{2}(2a^3x-x^4)^{-1/2}(2a^3-4x^3)-a\left(\frac{1}{3}\right)(aax)^{-2/3}a^2}{-\frac{1}{4}(ax^3)^{-3/4}(3ax^2)}$$

$$= \frac{\frac{1}{2}(2a^3a-a^4)^{-1/2}(2a^3-4a^3)-\frac{1}{3}a^3(a^2a)^{-2/3}}{-\frac{1}{4}(aa^3)^{-3/4}(3aa^2)}$$

$$= \frac{(a^4)^{-1/2}(-a^3)-\frac{1}{3}a^3(a^3)^{-2/3}}{-\frac{3}{4}a^3(a^4)^{-3/4}} = \frac{-a-\frac{1}{3}a}{-\frac{3}{4}} = \frac{4}{3}\left(\frac{4}{3}a\right) = \frac{16}{9}a$$

86. Let the radius of the circle be r. We see that $A(\theta)$ is the area of the whole figure (a sector of the circle with radius 1), minus the area of $\triangle OPR$. But the area of the sector of the circle is $\frac{1}{2}r^2\theta$ (see Reference Page 1), and the area of the triangle is $\frac{1}{2}r\,|PQ| = \frac{1}{2}r(r\sin\theta) = \frac{1}{2}r^2\sin\theta$. So we have $A(\theta) = \frac{1}{2}r^2\theta - \frac{1}{2}r^2\sin\theta = \frac{1}{2}r^2(\theta-\sin\theta)$. Now by elementary

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

trigonometry, $B(\theta) = \frac{1}{2}\,|QR|\,|PQ| = \frac{1}{2}(r - |OQ|)\,|PQ| = \frac{1}{2}(r - r\cos\theta)(r\sin\theta) = \frac{1}{2}r^2(1-\cos\theta)\sin\theta$.

So the limit we want is

$$\begin{aligned}\lim_{\theta\to 0^+}\frac{A(\theta)}{B(\theta)} &= \lim_{\theta\to 0^+}\frac{\frac{1}{2}r^2(\theta-\sin\theta)}{\frac{1}{2}r^2(1-\cos\theta)\sin\theta} \overset{\text{H}}{=} \lim_{\theta\to 0^+}\frac{1-\cos\theta}{(1-\cos\theta)\cos\theta+\sin\theta\,(\sin\theta)}\\ &= \lim_{\theta\to 0^+}\frac{1-\cos\theta}{\cos\theta-\cos^2\theta+\sin^2\theta} \overset{\text{H}}{=} \lim_{\theta\to 0^+}\frac{\sin\theta}{-\sin\theta-2\cos\theta\,(-\sin\theta)+2\sin\theta\,(\cos\theta)}\\ &= \lim_{\theta\to 0^+}\frac{\sin\theta}{-\sin\theta+4\sin\theta\,\cos\theta} = \lim_{\theta\to 0^+}\frac{1}{-1+4\cos\theta} = \frac{1}{-1+4\cos 0} = \frac{1}{3}\end{aligned}$$

87. The limit, $L = \lim\limits_{x\to\infty}\left[x - x^2\ln\left(\dfrac{1+x}{x}\right)\right] = \lim\limits_{x\to\infty}\left[x - x^2\ln\left(\dfrac{1}{x}+1\right)\right]$. Let $t = 1/x$, so as $x\to\infty$, $t\to 0^+$.

$$L = \lim_{t\to 0^+}\left[\frac{1}{t}-\frac{1}{t^2}\ln(t+1)\right] = \lim_{t\to 0^+}\frac{t-\ln(t+1)}{t^2} \overset{\text{H}}{=} \lim_{t\to 0^+}\frac{1-\dfrac{1}{t+1}}{2t} = \lim_{t\to 0^+}\frac{t/(t+1)}{2t} = \lim_{t\to 0^+}\frac{1}{2\,(t+1)} = \frac{1}{2}$$

Note: Starting the solution by factoring out x or x^2 leads to a more complicated solution.

88. $y = [f(x)]^{g(x)} \;\Rightarrow\; \ln y = g(x)\ln f(x)$. Since f is a positive function, $\ln f(x)$ is defined. Now $\lim\limits_{x\to a}\ln y = \lim\limits_{x\to a} g(x)\ln f(x) = -\infty$ since $\lim\limits_{x\to a} g(x) = \infty$ and $\lim\limits_{x\to a} f(x) = 0 \;\Rightarrow\; \lim\limits_{x\to a}\ln f(x) = -\infty$. Thus, if $t = \ln y$, $\lim\limits_{x\to a} y = \lim\limits_{t\to -\infty} e^t = 0$. Note that the limit, $\lim\limits_{x\to a} g(x)\ln f(x)$, is *not* of the form $\infty\cdot 0$.

89. (a) We look for functions f and g whose individual limits are ∞ as $x\to 0$, but whose quotient has a limit of 7 as $x\to 0$.

One such pair of functions is $f(x) = \dfrac{7}{x^2}$ and $g(x) = \dfrac{1}{x^2}$. We have $\lim\limits_{x\to 0} f(x) = \lim\limits_{x\to 0} g(x) = \infty$, and

$$\lim_{x\to 0}\frac{f(x)}{g(x)} = \lim_{x\to 0}\frac{7/x^2}{1/x^2} = \lim_{x\to 0} 7 = 7.$$

(b) We look for functions f and g whose individual limits are ∞ as $x\to 0$, but whose difference has a limit of 7 as $x\to 0$.

One such pair of functions is $f(x) = \dfrac{1}{x^2}+7$ and $g(x) = \dfrac{1}{x^2}$. We have $\lim\limits_{x\to 0} f(x) = \lim\limits_{x\to 0} g(x) = \infty$, and

$$\lim_{x\to 0}[f(x)-g(x)] = \lim_{x\to 0}\left[\left(\frac{1}{x^2}+7\right)-\frac{1}{x^2}\right] = \lim_{x\to 0} 7 = 7.$$

90. $L = \lim\limits_{x\to 0}\left(\dfrac{\sin 2x}{x^3}+a+\dfrac{b}{x^2}\right) = \lim\limits_{x\to 0}\dfrac{\sin 2x + ax^3 + bx}{x^3} \overset{\text{H}}{=} \lim\limits_{x\to 0}\dfrac{2\cos 2x + 3ax^2 + b}{3x^2}$. As $x\to 0$, $3x^2\to 0$, and $(2\cos 2x + 3ax^2 + b)\to b+2$, so the last limit exists only if $b+2=0$, that is, $b=-2$. Thus,

$\lim\limits_{x\to 0}\dfrac{2\cos 2x + 3ax^2 - 2}{3x^2} \overset{\text{H}}{=} \lim\limits_{x\to 0}\dfrac{-4\sin 2x + 6ax}{6x} \overset{\text{H}}{=} \lim\limits_{x\to 0}\dfrac{-8\cos 2x + 6a}{6} = \dfrac{6a-8}{6}$, which is equal to 0 if and only if $a = \frac{4}{3}$. Hence, $L = 0$ if and only if $b = -2$ and $a = \frac{4}{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

91. (a) We show that $\lim\limits_{x\to 0}\dfrac{f(x)}{x^n}=0$ for every integer $n\ge 0$. Let $y=\dfrac{1}{x^2}$. Then

$$\lim_{x\to 0}\frac{f(x)}{x^{2n}}=\lim_{x\to 0}\frac{e^{-1/x^2}}{\left(x^2\right)^n}=\lim_{y\to\infty}\frac{y^n}{e^y}\overset{\text{H}}{=}\lim_{y\to\infty}\frac{ny^{n-1}}{e^y}\overset{\text{H}}{=}\cdots\overset{\text{H}}{=}\lim_{y\to\infty}\frac{n!}{e^y}=0\quad\Rightarrow$$

$$\lim_{x\to 0}\frac{f(x)}{x^n}=\lim_{x\to 0}x^n\frac{f(x)}{x^{2n}}=\lim_{x\to 0}x^n\lim_{x\to 0}\frac{f(x)}{x^{2n}}=0.\ \text{Thus, } f'(0)=\lim_{x\to 0}\frac{f(x)-f(0)}{x-0}=\lim_{x\to 0}\frac{f(x)}{x}=0.$$

(b) Using the Chain Rule and the Quotient Rule we see that $f^{(n)}(x)$ exists for $x\ne 0$. In fact, we prove by induction that for each $n\ge 0$, there is a polynomial p_n and a non-negative integer k_n with $f^{(n)}(x)=p_n(x)f(x)/x^{k_n}$ for $x\ne 0$. This is true for $n=0$; suppose it is true for the nth derivative. Then $f'(x)=f(x)(2/x^3)$, so

$$\begin{aligned}f^{(n+1)}(x)&=\left[x^{k_n}\left[p_n'(x)\,f(x)+p_n(x)\,f'(x)\right]-k_nx^{k_n-1}p_n(x)\,f(x)\right]x^{-2k_n}\\&=\left[x^{k_n}p_n'(x)+p_n(x)(2/x^3)-k_nx^{k_n-1}p_n(x)\right]f(x)x^{-2k_n}\\&=\left[x^{k_n+3}p_n'(x)+2p_n(x)-k_nx^{k_n+2}\,p_n(x)\right]f(x)x^{-(2k_n+3)}\end{aligned}$$

which has the desired form.

Now we show by induction that $f^{(n)}(0)=0$ for all n. By part (a), $f'(0)=0$. Suppose that $f^{(n)}(0)=0$. Then

$$\begin{aligned}f^{(n+1)}(0)&=\lim_{x\to 0}\frac{f^{(n)}(x)-f^{(n)}(0)}{x-0}=\lim_{x\to 0}\frac{f^{(n)}(x)}{x}=\lim_{x\to 0}\frac{p_n(x)\,f(x)/x^{k_n}}{x}=\lim_{x\to 0}\frac{p_n(x)\,f(x)}{x^{k_n+1}}\\&=\lim_{x\to 0}p_n(x)\lim_{x\to 0}\frac{f(x)}{x^{k_n+1}}=p_n(0)\cdot 0=0\end{aligned}$$

92. (a) For f to be continuous, we need $\lim\limits_{x\to 0}f(x)=f(0)=1$. We note that for $x\ne 0$, $\ln f(x)=\ln|x|^x=x\ln|x|$.

So $\lim\limits_{x\to 0}\ln f(x)=\lim\limits_{x\to 0}x\ln|x|=\lim\limits_{x\to 0}\dfrac{\ln|x|}{1/x}\overset{\text{H}}{=}\lim\limits_{x\to 0}\dfrac{1/x}{-1/x^2}=0$. Therefore, $\lim\limits_{x\to 0}f(x)=\lim\limits_{x\to 0}e^{\ln f(x)}=e^0=1$.

So f is continuous at 0.

(b) From the graphs, it appears that f is differentiable at 0.

(c) To find f', we use logarithmic differentiation: $\ln f(x)=x\ln|x|\quad\Rightarrow\quad\dfrac{f'(x)}{f(x)}=x\left(\dfrac{1}{x}\right)+\ln|x|\quad\Rightarrow$

$f'(x)=f(x)(1+\ln|x|)=|x|^x(1+\ln|x|)$, $x\ne 0$. Now $f'(x)\to-\infty$ as $x\to 0$ [since $|x|^x\to 1$ and $(1+\ln|x|)\to-\infty$], so the curve has a vertical tangent at $(0,1)$ and is therefore not differentiable there.

The fact cannot be seen in the graphs in part (b) because $\ln|x|\to-\infty$ very slowly as $x\to 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4.5 Summary of Curve Sketching

1. $y = f(x) = x^3 + 3x^2 = x^2(x+3)$ **A.** f is a polynomial, so $D = \mathbb{R}$.

B. y-intercept $= f(0) = 0$, x-intercepts are 0 and -3 **C.** No symmetry

D. No asympote **E.** $f'(x) = 3x^2 + 6x = 3x(x+2) > 0 \quad \Leftrightarrow \quad x < -2$ or $x > 0$, so f is increasing on $(-\infty, -2)$ and $(0, \infty)$, and decreasing on $(-2, 0)$.

F. Local maximum value $f(-2) = 4$, local minimum value $f(0) = 0$

G. $f''(x) = 6x + 6 = 6(x+1) > 0 \quad \Leftrightarrow \quad x > -1$, so f is CU on $(-1, \infty)$ and CD on $(-\infty, -1)$. IP at $(-1, 2)$

H.

2. $y = f(x) = 2x^3 - 12x^2 + 18x = 2x(x^2 - 6x + 9) = 2x(x-3)^2$

A. $D = \mathbb{R}$ **B**. x-intercepts are 0 and 3, y-intercept $f(0) = 0$ **C**. No symmetry

D. No asymptote

E. $f'(x) = 6x^2 - 24x + 18 = 6(x^2 - 4x + 3)$
$= 6(x-1)(x-3) > 0 \quad \Leftrightarrow \quad x < 1$ or $x > 3$

and $f'(x) < 0 \quad \Leftrightarrow \quad 1 < x < 3$, so f is increasing on $(-\infty, 1)$ and $(3, \infty)$, and decreasing on $(1, 3)$. **F**. Local maximum value $f(1) = 8$, local minimum value $f(3) = 0$ **G**. $f''(x) = 12x - 24 = 12(x-2) > 0 \quad \Leftrightarrow \quad x > 2$, so f is CU on $(2, \infty)$, and f is CD on $(-\infty, 2)$. IP at $(2, 4)$

H.

3. $y = f(x) = x^4 - 4x = x(x^3 - 4)$ **A.** $D = \mathbb{R}$ **B.** x-intercepts are 0 and $\sqrt[3]{4}$, y-intercept $= f(0) = 0$ **C.** No symmetry **D.** No asymptote

E. $f'(x) = 4x^3 - 4 = 4(x^3 - 1) = 4(x-1)(x^2 + x + 1) > 0 \quad \Leftrightarrow \quad x > 1$, so f is increasing on $(1, \infty)$ and decreasing on $(-\infty, 1)$. **F.** Local minimum value $f(1) = -3$, no local maximum **G.** $f''(x) = 12x^2 > 0$ for all x, so f is CU on $(-\infty, \infty)$. No IP

H.

4. $y = f(x) = x^4 - 8x^2 + 8$ **A.** $D = \mathbb{R}$ **B.** y-intercept $f(0) = 8$; x-intercepts: $f(x) = 0 \quad \Rightarrow$ [by the quadratic formula] $x = \pm\sqrt{4 \pm 2\sqrt{2}} \approx \pm 2.61, \pm 1.08$ **C.** $f(-x) = f(x)$, so f is even and symmetric about the y-axis **D.** No asymptote

E. $f'(x) = 4x^3 - 16x = 4x(x^2 - 4) = 4x(x+2)(x-2) > 0 \quad \Leftrightarrow \quad -2 < x < 0$ or $x > 2$, so f is increasing on $(-2, 0)$ and $(2, \infty)$, and f is decreasing on $(-\infty, -2)$ and $(0, 2)$.

F. Local maximum value $f(0) = 8$, local minimum values $f(\pm 2) = -8$

G. $f''(x) = 12x^2 - 16 = 4(3x^2 - 4) > 0 \quad \Rightarrow \quad |x| > 2/\sqrt{3}$ [≈ 1.15], so f is CU on $\left(-\infty, -2/\sqrt{3}\right)$ and $\left(2/\sqrt{3}, \infty\right)$, and f is CD on $\left(-2/\sqrt{3}, 2/\sqrt{3}\right)$.

IP at $\left(\pm 2/\sqrt{3}, -\frac{8}{9}\right)$

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $y = f(x) = x(x-4)^3$ **A.** $D = \mathbb{R}$ **B.** x-intercepts are 0 and 4, y-intercept $f(0) = 0$ **C.** No symmetry

D. No asymptote

H.

E. $f'(x) = x \cdot 3(x-4)^2 + (x-4)^3 \cdot 1 = (x-4)^2[3x + (x-4)]$

$= (x-4)^2(4x-4) = 4(x-1)(x-4)^2 > 0 \quad \Leftrightarrow$

$x > 1$, so f is increasing on $(1, \infty)$ and decreasing on $(-\infty, 1)$.

F. Local minimum value $f(1) = -27$, no local maximum value

G. $f''(x) = 4[(x-1) \cdot 2(x-4) + (x-4)^2 \cdot 1] = 4(x-4)[2(x-1) + (x-4)]$

$= 4(x-4)(3x-6) = 12(x-4)(x-2) < 0 \quad \Leftrightarrow$

$2 < x < 4$, so f is CD on $(2, 4)$ and CU on $(-\infty, 2)$ and $(4, \infty)$. IPs at $(2, -16)$ and $(4, 0)$

6. $y = f(x) = x^5 - 5x = x(x^4 - 5)$ **A.** $D = \mathbb{R}$ **B.** x-intercepts $\pm\sqrt[4]{5}$ and 0, y-intercept $= f(0) = 0$

C. $f(-x) = -f(x)$, so f is odd; the curve is symmetric about the origin. **D.** No asymptote

H.

E. $f'(x) = 5x^4 - 5 = 5(x^4 - 1) = 5(x^2 - 1)(x^2 + 1)$

$= 5(x+1)(x-1)(x^2+1) > 0 \quad \Leftrightarrow$

$x < -1$ or $x > 1$, so f is increasing on $(-\infty, -1)$ and $(1, \infty)$, and f is decreasing on $(-1, 1)$. **F.** Local maximum value $f(-1) = 4$, local minimum value $f(1) = -4$ **G.** $f''(x) = 20x^3 > 0 \quad \Leftrightarrow \quad x > 0$, so f is CU on $(0, \infty)$ and CD on $(-\infty, 0)$. IP at $(0, 0)$

7. $y = f(x) = \frac{1}{5}x^5 - \frac{8}{3}x^3 + 16x = x\left(\frac{1}{5}x^4 - \frac{8}{3}x^2 + 16\right)$ **A.** $D = \mathbb{R}$ **B.** x-intercept 0, y-intercept $= f(0) = 0$

C. $f(-x) = -f(x)$, so f is odd; the curve is symmetric about the origin. **D.** No asymptote

H.

E. $f'(x) = x^4 - 8x^2 + 16 = (x^2 - 4)^2 = (x+2)^2(x-2)^2 > 0$ for all x except ± 2, so f is increasing on $\mathbb{R}$. **F.** There is no local maximum or minimum value.

G. $f''(x) = 4x^3 - 16x = 4x(x^2 - 4) = 4x(x+2)(x-2) > 0 \quad \Leftrightarrow$

$-2 < x < 0$ or $x > 2$, so f is CU on $(-2, 0)$ and $(2, \infty)$, and f is CD on $(-\infty, -2)$ and $(0, 2)$. IP at $\left(-2, -\frac{256}{15}\right)$, $(0, 0)$, and $\left(2, \frac{256}{15}\right)$

8. $y = f(x) = (4 - x^2)^5$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 4^5 = 1024$; x-intercepts: ± 2 **C.** $f(-x) = f(x) \quad \Rightarrow$ f is even; the curve is symmetric about the y-axis. **D.** No asymptote **E.** $f'(x) = 5(4 - x^2)^4(-2x) = -10x(4 - x^2)^4$, so for $x \neq \pm 2$ we have $f'(x) > 0 \quad \Leftrightarrow \quad x < 0$ and $f'(x) < 0 \quad \Leftrightarrow \quad x > 0$. Thus, f is increasing on $(-\infty, 0)$ and decreasing on $(0, \infty)$. **F.** Local maximum value $f(0) = 1024$

H.

G. $f''(x) = -10x \cdot 4(4 - x^2)^3(-2x) + (4 - x^2)^4(-10)$

$= -10(4 - x^2)^3[-8x^2 + 4 - x^2] = -10(4 - x^2)^3(4 - 9x^2)$

so $f''(x) = 0 \quad \Leftrightarrow \quad x = \pm 2, \pm\frac{2}{3}$. $f''(x) > 0 \quad \Leftrightarrow \quad -2 < x < -\frac{2}{3}$ and $\frac{2}{3} < x < 2$ and $f''(x) < 0 \quad \Leftrightarrow \quad x < -2, -\frac{2}{3} < x < \frac{2}{3}$, and $x > 2$, so f is CU on $(-\infty, 2)$, $\left(-\frac{2}{3}, \frac{2}{3}\right)$, and $(2, \infty)$, and CD on $\left(-2, -\frac{2}{3}\right)$ and $\left(\frac{2}{3}, 2\right)$.

IP at $(\pm 2, 0)$ and $\left(\pm\frac{2}{3}, \left(\frac{32}{9}\right)^5\right) \approx (\pm 0.67, 568.25)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $y = f(x) = \dfrac{2x+3}{x+2}$ **A.** $D = \{x \mid x \neq -2\} = (-\infty, -2) \cup (-2, \infty)$ **B.** x-intercept $= -\frac{3}{2}$, y-intercept $= f(0) = \frac{3}{2}$

C. No symmetry **D.** $\displaystyle\lim_{x\to\pm\infty} \frac{2x+3}{x+2} = 2$, so $y = 2$ is a HA. $\displaystyle\lim_{x\to-2^-} \frac{2x+3}{x+2} = \infty$, $\displaystyle\lim_{x\to-2^+} \frac{2x+3}{x+2} = -\infty$, so $x = -2$ is a VA. **E.** $f'(x) = \dfrac{(x+2)\cdot 2 - (2x+3)\cdot 1}{(x+2)^2} = \dfrac{1}{(x+2)^2} > 0$ for $x \neq -2$, so f is increasing on $(-\infty, -2)$ and $(-2, \infty)$. **F.** No extreme values

G. $f''(x) = \dfrac{-2}{(x+2)^3} > 0 \quad \Leftrightarrow \quad x < -2$, so f is CU on $(-\infty, -2)$ and CD on $(-2, \infty)$. No IP

H.

10. $y = f(x) = \dfrac{x^2+5x}{25-x^2} = \dfrac{x(x+5)}{(5+x)(5-x)} = \dfrac{x}{5-x}$ for $x \neq -5$. There is a hole in the graph at $\left(-5, -\frac{1}{2}\right)$.

A. $D = \{x \mid x \neq \pm 5\} = (-\infty, -5) \cup (-5, 5) \cup (5, \infty)$ **B.** x-intercept $= 0$, y-intercept $= f(0) = 0$ **C.** No symmetry

D. $\displaystyle\lim_{x\to\pm\infty} \frac{x}{5-x} = -1$, so $y = -1$ is a HA. $\displaystyle\lim_{x\to5^-} \frac{x}{5-x} = \infty$, $\displaystyle\lim_{x\to5^+} \frac{x}{5-x} = -\infty$, so $x = 5$ is a VA.

E. $f'(x) = \dfrac{(5-x)(1) - x(-1)}{(5-x)^2} = \dfrac{5}{(5-x)^2} > 0$ for all x in D, so f is increasing on $(-\infty, -5)$, $(-5, 5)$, and $(5, \infty)$. **F.** No extrema

G. $f'(x) = 5(5-x)^{-2} \quad \Rightarrow$

$f''(x) = -10(5-x)^{-3}(-1) = \dfrac{10}{(5-x)^3} > 0 \quad \Leftrightarrow \quad x < 5$, so f is CU on $(-\infty, -5)$ and $(-5, 5)$, and f is CD on $(5, \infty)$. No IP

H.

11. $y = f(x) = \dfrac{x-x^2}{2-3x+x^2} = \dfrac{x(1-x)}{(1-x)(2-x)} = \dfrac{x}{2-x}$ for $x \neq 1$. There is a hole in the graph at $(1, 1)$.

A. $D = \{x \mid x \neq 1, 2\} = (-\infty, 1) \cup (1, 2) \cup (2, \infty)$ **B.** x-intercept $= 0$, y-intercept $= f(0) = 0$ **C.** No symmetry

D. $\displaystyle\lim_{x\to\pm\infty} \frac{x}{2-x} = -1$, so $y = -1$ is a HA. $\displaystyle\lim_{x\to2^-} \frac{x}{2-x} = \infty$, $\displaystyle\lim_{x\to2^+} \frac{x}{2-x} = -\infty$, so $x = 2$ is a VA.

E. $f'(x) = \dfrac{(2-x)(1) - x(-1)}{(2-x)^2} = \dfrac{2}{(2-x)^2} > 0$ $[x \neq 1, 2]$, so f is increasing on $(-\infty, 1)$, $(1, 2)$, and $(2, \infty)$. **F.** No extrema

G. $f'(x) = 2(2-x)^{-2} \quad \Rightarrow$

$f''(x) = -4(2-x)^{-3}(-1) = \dfrac{4}{(2-x)^3} > 0 \quad \Leftrightarrow \quad x < 2$, so f is CU on $(-\infty, 1)$ and $(1, 2)$, and f is CD on $(2, \infty)$. No IP

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $y = f(x) = 1 + \dfrac{1}{x} + \dfrac{1}{x^2} = \dfrac{x^2 + x + 1}{x^2}$ **A.** $D = (-\infty, 0) \cup (0, \infty)$ **B.** y-intercept: none $[x \neq 0]$;

x-intercepts: $f(x) = 0 \quad \Leftrightarrow \quad x^2 + x + 1 = 0$, there is no real solution, and hence, no x-intercept **C.** No symmetry

D. $\lim\limits_{x \to \pm\infty} \left(1 + \dfrac{1}{x} + \dfrac{1}{x^2}\right) = 1$, so $y = 1$ is a HA. $\lim\limits_{x \to 0} f(x) = \infty$, so $x = 0$ is a VA. **E.** $f'(x) = -\dfrac{1}{x^2} - \dfrac{2}{x^3} = \dfrac{-x-2}{x^3}$.

$f'(x) > 0 \quad \Leftrightarrow \quad -2 < x < 0$ and $f'(x) < 0 \quad \Leftrightarrow \quad x < -2$ or $x > 0$, so f is increasing on $(-2, 0)$ and decreasing on $(-\infty, -2)$ and $(0, \infty)$. **F.** Local minimum value $f(-2) = \frac{3}{4}$; no local maximum **G.** $f''(x) = \dfrac{2}{x^3} + \dfrac{6}{x^4} = \dfrac{2x+6}{x^4}$. $f''(x) < 0 \quad \Leftrightarrow \quad x < -3$ and $f''(x) > 0 \quad \Leftrightarrow \quad -3 < x < 0$ and $x > 0$, so f is CD on $(-\infty, -3)$ and CU on $(-3, 0)$ and $(0, \infty)$. IP at $\left(-3, \frac{7}{9}\right)$

H.

13. $y = f(x) = \dfrac{x}{x^2 - 4} = \dfrac{x}{(x+2)(x-2)}$ **A.** $D = (-\infty, -2) \cup (-2, 2) \cup (2, \infty)$ **B.** x-intercept $= 0$, y-intercept $= f(0) = 0$ **C.** $f(-x) = -f(x)$, so f is odd; the graph is symmetric about the origin.

D. $\lim\limits_{x \to 2^+} \dfrac{x}{x^2 - 4} = \infty$, $\lim\limits_{x \to 2^-} f(x) = -\infty$, $\lim\limits_{x \to -2^+} f(x) = \infty$, $\lim\limits_{x \to -2^-} f(x) = -\infty$, so $x = \pm 2$ are VAs.

$\lim\limits_{x \to \pm\infty} \dfrac{x}{x^2 - 4} = 0$, so $y = 0$ is a HA. **E.** $f'(x) = \dfrac{(x^2-4)(1) - x(2x)}{(x^2-4)^2} = -\dfrac{x^2+4}{(x^2-4)^2} < 0$ for all x in D, so f is decreasing on $(-\infty, -2)$, $(-2, 2)$, and $(2, \infty)$.

F. No local extrema

G.
$$\begin{aligned} f''(x) &= -\frac{(x^2-4)^2(2x) - (x^2+4)2(x^2-4)(2x)}{[(x^2-4)^2]^2} \\ &= -\frac{2x(x^2-4)[(x^2-4) - 2(x^2+4)]}{(x^2-4)^4} \\ &= -\frac{2x(-x^2-12)}{(x^2-4)^3} = \frac{2x(x^2+12)}{(x+2)^3(x-2)^3}. \end{aligned}$$

$f''(x) < 0$ if $x < -2$ or $0 < x < 2$, so f is CD on $(-\infty, -2)$ and $(0, 2)$, and CU on $(-2, 0)$ and $(2, \infty)$. IP at $(0, 0)$

H.

14. $y = f(x) = \dfrac{1}{x^2 - 4} = \dfrac{1}{(x+2)(x-2)}$ **A.** $D = (-\infty, -2) \cup (-2, 2) \cup (2, \infty)$ **B.** No x-intercept, y-intercept $= f(0) = -\frac{1}{4}$ **C.** $f(-x) = f(x)$, so f is even; the graph is symmetric about the y-axis.

D. $\lim\limits_{x \to 2^+} \dfrac{1}{x^2 - 4} = \infty$, $\lim\limits_{x \to 2^-} f(x) = -\infty$, $\lim\limits_{x \to -2^+} f(x) = -\infty$, $\lim\limits_{x \to -2^-} f(x) = \infty$, so $x = \pm 2$ are VAs. $\lim\limits_{x \to \pm\infty} f(x) = 0$, so $y = 0$ is a HA. **E.** $f'(x) = -\dfrac{2x}{(x^2-4)^2}$ [Reciprocal Rule] > 0 if $x < 0$ and x is in D, so f is increasing on $(-\infty, -2)$ and $(-2, 0)$. f is decreasing on $(0, 2)$ and $(2, \infty)$. **F.** Local maximum value $f(0) = -\frac{1}{4}$, no local minimum value

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

G. $f''(x) = \dfrac{(x^2-4)^2(-2) - (-2x)2(x^2-4)(2x)}{[(x^2-4)^2]^2}$

$= \dfrac{-2(x^2-4)\left[(x^2-4)-4x^2\right]}{(x^2-4)^4}$

$= \dfrac{-2(-3x^2-4)}{(x^2-4)^3} = \dfrac{2(3x^2+4)}{(x^2-4)^3}$

$f''(x) < 0 \quad \Leftrightarrow \quad -2 < x < 2$, so f is CD on $(-2, 2)$ and CU on $(-\infty, -2)$ and $(2, \infty)$. No IP

H.

15. $y = f(x) = \dfrac{x^2}{x^2+3} = \dfrac{(x^2+3)-3}{x^2+3} = 1 - \dfrac{3}{x^2+3}$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \quad \Leftrightarrow \quad x = 0$ **C.** $f(-x) = f(x)$, so f is even; the graph is symmetric about the y-axis.

D. $\displaystyle\lim_{x\to\pm\infty} \frac{x^2}{x^2+3} = 1$, so $y = 1$ is a HA. No VA. **E.** Using the Reciprocal Rule, $f'(x) = -3 \cdot \dfrac{-2x}{(x^2+3)^2} = \dfrac{6x}{(x^2+3)^2}$.

$f'(x) > 0 \quad \Leftrightarrow \quad x > 0$ and $f'(x) < 0 \quad \Leftrightarrow \quad x < 0$, so f is decreasing on $(-\infty, 0)$ and increasing on $(0, \infty)$.

F. Local minimum value $f(0) = 0$, no local maximum.

G. $f''(x) = \dfrac{(x^2+3)^2 \cdot 6 - 6x \cdot 2(x^2+3) \cdot 2x}{[(x^2+3)^2]^2}$

$= \dfrac{6(x^2+3)[(x^2+3)-4x^2]}{(x^2+3)^4} = \dfrac{6(3-3x^2)}{(x^2+3)^3} = \dfrac{-18(x+1)(x-1)}{(x^2+3)^3}$

$f''(x)$ is negative on $(-\infty, -1)$ and $(1, \infty)$ and positive on $(-1, 1)$, so f is CD on $(-\infty, -1)$ and $(1, \infty)$ and CU on $(-1, 1)$. IP at $\left(\pm 1, \frac{1}{4}\right)$

H.

16. $y = f(x) = \dfrac{(x-1)^2}{x^2+1} \geq 0$ with equality $\quad \Leftrightarrow \quad x = 1$. **A.** $D = \mathbb{R}$ **B.** y-intercept $= f(0) = 1$; x-intercept 1 **C.** No symmetry **D.** $\displaystyle\lim_{x\to\pm\infty} f(x) = \lim_{x\to\pm\infty} \frac{x^2-2x+1}{x^2+1} = \lim_{x\to\pm\infty} \frac{1-2/x+1/x^2}{1+1/x^2} = 1$, so $y = 1$ is a HA. No VA

E. $f'(x) = \dfrac{(x^2+1)2(x-1) - (x-1)^2(2x)}{(x^2+1)^2} = \dfrac{2(x-1)\left[(x^2+1)-x(x-1)\right]}{(x^2+1)^2} = \dfrac{2(x-1)(x+1)}{(x^2+1)^2} < 0 \quad \Leftrightarrow$

$-1 < x < 1$, so f is decreasing on $(-1, 1)$ and increasing on $(-\infty, -1)$ and $(1, \infty)$ **F.** Local maximum value $f(-1) = 2$, local minimum value $f(1) = 0$

G. $f''(x) = \dfrac{(x^2+1)^2(4x) - (2x^2-2)2(x^2+1)(2x)}{[(x^2+1)^2]^2} = \dfrac{4x(x^2+1)\left[(x^2+1)-(2x^2-2)\right]}{(x^2+1)^4} = \dfrac{4x(3-x^2)}{(x^2+1)^3}$.

$f''(x) > 0 \quad \Leftrightarrow \quad x < -\sqrt{3}$ or $0 < x < \sqrt{3}$, so f is CU on $\left(-\infty, -\sqrt{3}\right)$ and $\left(0, \sqrt{3}\right)$, and f is CD on $\left(-\sqrt{3}, 0\right)$ and $\left(\sqrt{3}, \infty\right)$.

$f\left(\pm\sqrt{3}\right) = \frac{1}{4}\left(\sqrt{3} \mp 1\right)^2 = \frac{1}{4}\left(4 \mp 2\sqrt{3}\right) = 1 \mp \frac{1}{2}\sqrt{3}\,[\approx 0.13, 1.87]$, so there are IPs at $\left(-\sqrt{3}, 1 + \frac{1}{2}\sqrt{3}\right)$, $(0, 1)$, and $\left(\sqrt{3}, 1 - \frac{1}{2}\sqrt{3}\right)$. Note that the graph is symmetric about the point $(0, 1)$.

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $y = f(x) = \dfrac{x-1}{x^2}$ **A.** $D = \{x \mid x \neq 0\} = (-\infty, 0) \cup (0, \infty)$ **B.** No y-intercept; x-intercept: $f(x) = 0 \quad \Leftrightarrow \quad x = 1$

C. No symmetry **D.** $\lim\limits_{x\to\pm\infty} \dfrac{x-1}{x^2} = 0$, so $y = 0$ is a HA. $\lim\limits_{x\to 0} \dfrac{x-1}{x^2} = -\infty$, so $x = 0$ is a VA.

E. $f'(x) = \dfrac{x^2 \cdot 1 - (x-1)\cdot 2x}{(x^2)^2} = \dfrac{-x^2+2x}{x^4} = \dfrac{-(x-2)}{x^3}$, so $f'(x) > 0 \quad \Leftrightarrow \quad 0 < x < 2$ and $f'(x) < 0 \quad \Leftrightarrow$ $x < 0$ or $x > 2$. Thus, f is increasing on $(0, 2)$ and decreasing on $(-\infty, 0)$ and $(2, \infty)$. **F.** No local minimum, local maximum value $f(2) = \frac{1}{4}$.

G. $f''(x) = \dfrac{x^3 \cdot (-1) - [-(x-2)]\cdot 3x^2}{(x^3)^2} = \dfrac{2x^3 - 6x^2}{x^6} = \dfrac{2(x-3)}{x^4}$.

$f''(x)$ is negative on $(-\infty, 0)$ and $(0, 3)$ and positive on $(3, \infty)$, so f is CD on $(-\infty, 0)$ and $(0, 3)$ and CU on $(3, \infty)$. IP at $\left(3, \frac{2}{9}\right)$

H.

18. $y = f(x) = \dfrac{x}{x^3 - 1}$ **A.** $D = (-\infty, 1) \cup (1, \infty)$ **B.** y-intercept: $f(0) = 0$; x-intercept: $f(x) = 0 \quad \Leftrightarrow \quad x = 0$

C. No symmetry **D.** $\lim\limits_{x\to\pm\infty} \dfrac{x}{x^3-1} = 0$, so $y = 0$ is a HA. $\lim\limits_{x\to 1^-} f(x) = -\infty$ and $\lim\limits_{x\to 1^+} f(x) = \infty$, so $x = 1$ is a VA.

E. $f'(x) = \dfrac{(x^3-1)(1) - x(3x^2)}{(x^3-1)^2} = \dfrac{-2x^3-1}{(x^3-1)^2}$. $f'(x) = 0 \quad \Rightarrow \quad x = -\sqrt[3]{1/2}$. $f'(x) > 0 \quad \Leftrightarrow \quad x < -\sqrt[3]{1/2}$ and $f'(x) < 0 \quad \Leftrightarrow \quad -\sqrt[3]{1/2} < x < 1$ and $x > 1$, so f is increasing on $\left(-\infty, -\sqrt[3]{1/2}\right)$ and decreasing on $\left(-\sqrt[3]{1/2}, 1\right)$ and $(1, \infty)$. **F.** Local maximum value $f\left(-\sqrt[3]{1/2}\right) = \frac{2}{3}\sqrt[3]{1/2}$; no local minimum

G. $f''(x) = \dfrac{(x^3-1)^2(-6x^2) - (-2x^3-1)2(x^3-1)(3x^2)}{[(x^3-1)^2]^2}$

$= \dfrac{-6x^2(x^3-1)[(x^3-1) - (2x^3+1)]}{(x^3-1)^4} = \dfrac{6x^2(x^3+2)}{(x^3-1)^3}$.

$f''(x) > 0 \quad \Leftrightarrow \quad x < -\sqrt[3]{2}$ and $x > 1$, $f''(x) < 0 \quad \Leftrightarrow \quad -\sqrt[3]{2} < x < 0$ and $0 < x < 1$, so f is CU on $\left(-\infty, -\sqrt[3]{2}\right)$ and $(1, \infty)$ and CD on $\left(-\sqrt[3]{2}, 1\right)$.

IP at $\left(-\sqrt[3]{2}, \frac{1}{3}\sqrt[3]{2}\right)$

H.

19. $y = f(x) = \dfrac{x^3}{x^3+1} = \dfrac{x^3}{(x+1)(x^2-x+1)}$ **A.** $D = (-\infty, -1) \cup (-1, \infty)$ **B.** y-intercept: $f(0) = 0$; x-intercept: $f(x) = 0 \quad \Leftrightarrow \quad x = 0$ **C.** No symmetry **D.** $\lim\limits_{x\to\pm\infty} \dfrac{x^3}{x^3+1} = \dfrac{1}{1+1/x^3} = 1$, so $y = 1$ is a HA. $\lim\limits_{x\to -1^-} f(x) = \infty$ and $\lim\limits_{x\to -1^+} f(x) = -\infty$, so $x = -1$ is a VA. **E.** $f'(x) = \dfrac{(x^3+1)(3x^2) - x^3(3x^2)}{(x^3+1)^2} = \dfrac{3x^2}{(x^3+1)^2}$. $f'(x) > 0$ for $x \neq -1$ (not in the domain) and $x \neq 0$ ($f' = 0$), so f is increasing on $(-\infty, -1)$, $(-1, 0)$, and $(0, \infty)$, and furthermore, by Exercise 4.3.97, f is increasing on $(-\infty, -1)$, and $(-1, \infty)$. **F.** No local extrema

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

G. $f''(x) = \dfrac{(x^3+1)^2(6x) - 3x^2[2(x^3+1)(3x^2)]}{[(x^3+1)^2]^2}$

$= \dfrac{(x^3+1)(6x)[(x^3+1)-3x^3]}{(x^3+1)^4} = \dfrac{6x(1-2x^3)}{(x^3+1)^3}$

$f''(x) > 0 \quad \Leftrightarrow \quad x < -1$ or $0 < x < \sqrt[3]{\frac{1}{2}}$ $[\approx 0.79]$, so f is CU on $(-\infty, -1)$ and $\left(0, \sqrt[3]{\frac{1}{2}}\right)$ and CD on $(-1, 0)$ and $\left(\sqrt[3]{\frac{1}{2}}, \infty\right)$. There are IPs at $(0, 0)$ and $\left(\sqrt[3]{\frac{1}{2}}, \frac{1}{3}\right)$.

H.

20. $y = f(x) = \dfrac{x^3}{x-2} = x^2 + 2x + 4 + \dfrac{8}{x-2}$ [by long division] **A.** $D = (-\infty, 2) \cup (2, \infty)$ **B.** x-intercept $= 0$, y-intercept $= f(0) = 0$ **C.** No symmetry **D.** $\lim\limits_{x \to 2^-} \dfrac{x^3}{x-2} = -\infty$ and $\lim\limits_{x \to 2^+} \dfrac{x^3}{x-2} = \infty$, so $x = 2$ is a VA.

There are no horizontal or slant asymptotes. *Note:* Since $\lim\limits_{x \to \pm\infty} \dfrac{8}{x-2} = 0$, the parabola $y = x^2 + 2x + 4$ is approached asymptotically as $x \to \pm\infty$.

E. $f'(x) = \dfrac{(x-2)(3x^2) - x^3(1)}{(x-2)^2} = \dfrac{x^2[3(x-2)-x]}{(x-2)^2} = \dfrac{x^2(2x-6)}{(x-2)^2} = \dfrac{2x^2(x-3)}{(x-2)^2} > 0 \quad \Leftrightarrow \quad x > 3$ and

$f'(x) < 0 \quad \Leftrightarrow \quad x < 0$ or $0 < x < 2$ or $2 < x < 3$, so f is increasing on $(3, \infty)$ and f is decreasing on $(-\infty, 2)$ and $(2, 3)$.

F. Local minimum value $f(3) = 27$, no local maximum value **G.** $f'(x) = 2\dfrac{x^3 - 3x^2}{(x-2)^2} \quad \Rightarrow$

$f''(x) = 2\dfrac{(x-2)^2(3x^2-6x) - (x^3-3x^2)2(x-2)}{[(x-2)^2]^2}$

$= 2\dfrac{(x-2)x[(x-2)(3x-6) - (x^2-3x)2]}{(x-2)^4}$

$= \dfrac{2x(3x^2 - 12x + 12 - 2x^2 + 6x)}{(x-2)^3}$

$= \dfrac{2x(x^2 - 6x + 12)}{(x-2)^3} > 0 \quad \Leftrightarrow$

$x < 0$ or $x > 2$, so f is CU on $(-\infty, 0)$ and $(2, \infty)$, and f is CD on $(0, 2)$. IP at $(0, 0)$

H.

21. $y = f(x) = (x-3)\sqrt{x} = x^{3/2} - 3x^{1/2}$ **A.** $D = [0, \infty)$ **B.** x-intercepts: $0, 3$; y-intercept $= f(0) = 0$ **C.** No symmetry **D.** No asymptote **E.** $f'(x) = \frac{3}{2}x^{1/2} - \frac{3}{2}x^{-1/2} = \frac{3}{2}x^{-1/2}(x-1) = \dfrac{3(x-1)}{2\sqrt{x}} > 0 \quad \Leftrightarrow \quad x > 1$,

so f is increasing on $(1, \infty)$ and decreasing on $(0, 1)$.

F. Local minimum value $f(1) = -2$, no local maximum value

G. $f''(x) = \frac{3}{4}x^{-1/2} + \frac{3}{4}x^{-3/2} = \frac{3}{4}x^{-3/2}(x+1) = \dfrac{3(x+1)}{4x^{3/2}} > 0$ for $x > 0$,

so f is CU on $(0, \infty)$. No IP

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $y = f(x) = (x-4)\sqrt[3]{x} = x^{4/3} - 4x^{1/3}$ **A.** $D = \mathbb{R}$ **B.** y-intercept $= f(0) = 0$; x-intercepts: 0 and 4

C. No symmetry **D.** No asymptote

H.

E. $f'(x) = \frac{4}{3}x^{1/3} - \frac{4}{3}x^{-2/3} = \frac{4}{3}x^{-2/3}(x-1) = \dfrac{4(x-1)}{3x^{2/3}}$. $f'(x) > 0 \iff$ $x > 1$, so f is increasing on $(1, \infty)$ and f is decreasing on $(-\infty, 1)$.

F. Local minimum value $f(1) = -3$

G. $f''(x) = \frac{4}{9}x^{-2/3} + \frac{8}{9}x^{-5/3} = \frac{4}{9}x^{-5/3}(x+2) = \dfrac{4(x+2)}{9x^{5/3}}$.

$f''(x) < 0 \iff -2 < x < 0$, so f is CD on $(-2, 0)$, and f is CU on $(-\infty, -2)$ and $(0, \infty)$. There are IPs at $\left(-2, 6\sqrt[3]{2}\right)$ and $(0, 0)$.

23. $y = f(x) = \sqrt{x^2 + x - 2} = \sqrt{(x+2)(x-1)}$ **A.** $D = \{x \mid (x+2)(x-1) \geq 0\} = (-\infty, -2] \cup [1, \infty)$

B. y-intercept: none; x-intercepts: -2 and 1 **C.** No symmetry **D.** No asymptote

E. $f'(x) = \frac{1}{2}(x^2 + x - 2)^{-1/2}(2x+1) = \dfrac{2x+1}{2\sqrt{x^2+x-2}}$, $f'(x) = 0$ if $x = -\frac{1}{2}$, but $-\frac{1}{2}$ is not in the domain.

$f'(x) > 0 \Rightarrow x > -\frac{1}{2}$ and $f'(x) < 0 \Rightarrow x < -\frac{1}{2}$, so (considering the domain) f is increasing on $(1, \infty)$ and f is decreasing on $(-\infty, -2)$. **F.** No local extrema

G.
$$f''(x) = \frac{2(x^2+x-2)^{1/2}(2) - (2x+1)\cdot 2 \cdot \frac{1}{2}(x^2+x-2)^{-1/2}(2x+1)}{\left(2\sqrt{x^2+x-2}\right)^2}$$
$$= \frac{(x^2+x-2)^{-1/2}\left[4(x^2+x-2) - (4x^2+4x+1)\right]}{4(x^2+x-2)}$$
$$= \frac{-9}{4(x^2+x-2)^{3/2}} < 0$$

so f is CD on $(-\infty, -2)$ and $(1, \infty)$. No IP

H.

24. $y = f(x) = \sqrt{x^2 + x} - x = \sqrt{x(x+1)} - x$ **A.** $D = (-\infty, -1] \cup [0, \infty)$ **B.** y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \Rightarrow \sqrt{x^2+x} = x \Rightarrow x^2 + x = x^2 \Rightarrow x = 0$ **C.** No symmetry

D.
$$\lim_{x\to\infty} f(x) = \lim_{x\to\infty}\left(\sqrt{x^2+x} - x\right)\frac{\sqrt{x^2+x}+x}{\sqrt{x^2+x}+x} = \lim_{x\to\infty}\frac{x^2+x-x^2}{\sqrt{x^2+x}+x}$$
$$= \lim_{x\to\infty}\frac{x/x}{\left(\sqrt{x^2+x}+x\right)/x} = \lim_{x\to\infty}\frac{1}{\sqrt{1+1/x}+1} = \frac{1}{2},$$ so $y = \frac{1}{2}$ is a HA. No VA

E. $f'(x) = \frac{1}{2}(x^2+x)^{-1/2}(2x+1) - 1 = \dfrac{2x+1}{2\sqrt{x^2+x}} - 1 > 0 \iff 2x+1 > 2\sqrt{x^2+x} \iff$

$x + \frac{1}{2} > \sqrt{\left(x+\frac{1}{2}\right)^2 - \frac{1}{4}}$. Keep in mind that the domain excludes the interval $(-1, 0)$. When $x + \frac{1}{2}$ is positive (for $x \geq 0$), the last inequality is *true* since the value of the radical is less than $x + \frac{1}{2}$. When $x + \frac{1}{2}$ is negative (for $x \leq -1$), the last inequality is *false* since the value of the radical is positive. So f is increasing on $(0, \infty)$ and decreasing on $(-\infty, -1)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

F. No local extrema

G. $f''(x) = \dfrac{2(x^2+x)^{1/2}(2) - (2x+1)\cdot 2\cdot \frac{1}{2}(x^2+x)^{-1/2}(2x+1)}{\left(2\sqrt{x^2+x}\,\right)^2}$

$= \dfrac{(x^2+x)^{-1/2}[4(x^2+x)-(2x+1)^2]}{4(x^2+x)} = \dfrac{-1}{4(x^2+x)^{3/2}}.$

$f''(x) < 0$ when it is defined, so f is CD on $(-\infty, -1)$ and $(0, \infty)$. No IP

H.

25. $y = f(x) = x/\sqrt{x^2+1}$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \;\Rightarrow\; x = 0$

C. $f(-x) = -f(x)$, so f is odd; the graph is symmetric about the origin.

D. $\displaystyle\lim_{x\to\infty} f(x) = \lim_{x\to\infty} \frac{x}{\sqrt{x^2+1}} = \lim_{x\to\infty} \frac{x/x}{\sqrt{x^2+1}/x} = \lim_{x\to\infty} \frac{x/x}{\sqrt{x^2+1}/\sqrt{x^2}} = \lim_{x\to\infty} \frac{1}{\sqrt{1+1/x^2}} = \frac{1}{\sqrt{1+0}} = 1$

and

$\displaystyle\lim_{x\to-\infty} f(x) = \lim_{x\to-\infty} \frac{x}{\sqrt{x^2+1}} = \lim_{x\to-\infty} \frac{x/x}{\sqrt{x^2+1}/x} = \lim_{x\to-\infty} \frac{x/x}{\sqrt{x^2+1}/\left(-\sqrt{x^2}\right)} = \lim_{x\to-\infty} \frac{1}{-\sqrt{1+1/x^2}}$

$= \dfrac{1}{-\sqrt{1+0}} = -1$ so $y = \pm 1$ are HA. No VA

E. $f'(x) = \dfrac{\sqrt{x^2+1} - x\cdot\dfrac{2x}{2\sqrt{x^2+1}}}{[(x^2+1)^{1/2}]^2} = \dfrac{x^2+1-x^2}{(x^2+1)^{3/2}} = \dfrac{1}{(x^2+1)^{3/2}} > 0$ for all x, so f is increasing on $\mathbb{R}$.

F. No extreme values

G. $f''(x) = -\frac{3}{2}(x^2+1)^{-5/2}\cdot 2x = \dfrac{-3x}{(x^2+1)^{5/2}}$, so $f''(x) > 0$ for $x < 0$

and $f''(x) < 0$ for $x > 0$. Thus, f is CU on $(-\infty, 0)$ and CD on $(0, \infty)$.

IP at $(0, 0)$

H.

26. $y = f(x) = x\sqrt{2-x^2}$ **A.** $D = \left[-\sqrt{2}, \sqrt{2}\,\right]$ **B.** y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \;\Rightarrow$

$x = 0, \pm\sqrt{2}$. **C.** $f(-x) = -f(x)$, so f is odd; the graph is symmetric about the origin. **D.** No asymptote

E. $f'(x) = x\cdot\dfrac{-x}{\sqrt{2-x^2}} + \sqrt{2-x^2} = \dfrac{-x^2+2-x^2}{\sqrt{2-x^2}} = \dfrac{2(1+x)(1-x)}{\sqrt{2-x^2}}$. $f'(x)$ is negative for $-\sqrt{2} < x < -1$

and $1 < x < \sqrt{2}$, and positive for $-1 < x < 1$, so f is decreasing on $\left(-\sqrt{2}, -1\right)$ and $\left(1, \sqrt{2}\,\right)$ and increasing on $(-1, 1)$.

F. Local minimum value $f(-1) = -1$, local maximum value $f(1) = 1$.

G. $f''(x) = \dfrac{\sqrt{2-x^2}(-4x) - (2-2x^2)\dfrac{-x}{\sqrt{2-x^2}}}{[(2-x^2)^{1/2}]^2}$

$= \dfrac{(2-x^2)(-4x) + (2-2x^2)x}{(2-x^2)^{3/2}} = \dfrac{2x^3-6x}{(2-x^2)^{3/2}} = \dfrac{2x(x^2-3)}{(2-x^2)^{3/2}}$

Since $x^2 - 3 < 0$ for x in $\left[-\sqrt{2}, \sqrt{2}\,\right]$, $f''(x) > 0$ for $-\sqrt{2} < x < 0$ and

$f''(x) < 0$ for $0 < x < \sqrt{2}$. Thus, f is CU on $\left(-\sqrt{2}, 0\right)$ and CD on $\left(0, \sqrt{2}\,\right)$.

The only IP is $(0, 0)$.

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. $y = f(x) = \sqrt{1-x^2}/x$ **A.** $D = \{x \mid |x| \le 1, x \ne 0\} = [-1, 0) \cup (0, 1]$ **B.** x-intercepts ± 1, no y-intercept

C. $f(-x) = -f(x)$, so the curve is symmetric about $(0, 0)$. **D.** $\lim\limits_{x \to 0^+} \dfrac{\sqrt{1-x^2}}{x} = \infty$, $\lim\limits_{x \to 0^-} \dfrac{\sqrt{1-x^2}}{x} = -\infty$,

so $x = 0$ is a VA. **E.** $f'(x) = \dfrac{\left(-x^2/\sqrt{1-x^2}\right) - \sqrt{1-x^2}}{x^2} = -\dfrac{1}{x^2\sqrt{1-x^2}} < 0$, so f is decreasing

on $(-1, 0)$ and $(0, 1)$. **F.** No extreme values

G. $f''(x) = \dfrac{2-3x^2}{x^3(1-x^2)^{3/2}} > 0 \quad \Leftrightarrow \quad -1 < x < -\sqrt{\tfrac{2}{3}}$ or $0 < x < \sqrt{\tfrac{2}{3}}$, so

f is CU on $\left(-1, -\sqrt{\tfrac{2}{3}}\right)$ and $\left(0, \sqrt{\tfrac{2}{3}}\right)$ and CD on $\left(-\sqrt{\tfrac{2}{3}}, 0\right)$ and $\left(\sqrt{\tfrac{2}{3}}, 1\right)$.

IP at $\left(\pm\sqrt{\tfrac{2}{3}}, \pm\tfrac{1}{\sqrt{2}}\right)$

H.

28. $y = f(x) = x/\sqrt{x^2-1}$ **A.** $D = (-\infty, -1) \cup (1, \infty)$ **B.** No intercepts **C.** $f(-x) = -f(x)$, so f is odd;

the graph is symmetric about the origin. **D.** $\lim\limits_{x \to \infty} \dfrac{x}{\sqrt{x^2-1}} = 1$ and $\lim\limits_{x \to -\infty} \dfrac{x}{\sqrt{x^2-1}} = -1$, so $y = \pm 1$ are HA.

$\lim\limits_{x \to 1^+} f(x) = +\infty$ and $\lim\limits_{x \to -1^-} f(x) = -\infty$, so $x = \pm 1$ are VA.

E. $f'(x) = \dfrac{\sqrt{x^2-1} - x \cdot \dfrac{x}{\sqrt{x^2-1}}}{[(x^2-1)^{1/2}]^2} = \dfrac{x^2-1-x^2}{(x^2-1)^{3/2}} = \dfrac{-1}{(x^2-1)^{3/2}} < 0$, so f is decreasing on $(-\infty, -1)$ and $(1, \infty)$.

F. No extreme values

G. $f''(x) = (-1)\left(-\tfrac{3}{2}\right)(x^2-1)^{-5/2} \cdot 2x = \dfrac{3x}{(x^2-1)^{5/2}}$.

$f''(x) < 0$ on $(-\infty, -1)$ and $f''(x) > 0$ on $(1, \infty)$, so f is CD on $(-\infty, -1)$ and CU on $(1, \infty)$. No IP

H.

29. $y = f(x) = x - 3x^{1/3}$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \quad \Rightarrow \quad x = 3x^{1/3} \quad \Rightarrow$

$x^3 = 27x \quad \Rightarrow \quad x^3 - 27x = 0 \quad \Rightarrow \quad x(x^2 - 27) = 0 \quad \Rightarrow \quad x = 0, \pm 3\sqrt{3}$ **C.** $f(-x) = -f(x)$, so f is odd;

the graph is symmetric about the origin. **D.** No asymptote **E.** $f'(x) = 1 - x^{-2/3} = 1 - \dfrac{1}{x^{2/3}} = \dfrac{x^{2/3}-1}{x^{2/3}}$.

$f'(x) > 0$ when $|x| > 1$ and $f'(x) < 0$ when $0 < |x| < 1$, so f is increasing on $(-\infty, -1)$ and $(1, \infty)$, and

decreasing on $(-1, 0)$ and $(0, 1)$ [hence decreasing on $(-1, 1)$ since f is continuous on $(-1, 1)$]. **F.** Local maximum value $f(-1) = 2$, local minimum value $f(1) = -2$ **G.** $f''(x) = \tfrac{2}{3}x^{-5/3} < 0$ when $x < 0$ and $f''(x) > 0$ when $x > 0$, so f is CD on $(-\infty, 0)$ and CU on $(0, \infty)$. IP at $(0, 0)$

H.

30. $y = f(x) = x^{5/3} - 5x^{2/3} = x^{2/3}(x-5)$ **A.** $D = \mathbb{R}$ **B.** x-intercepts 0, 5; y-intercept 0 **C.** No symmetry

D. $\lim\limits_{x \to \pm\infty} x^{2/3}(x-5) = \pm\infty$, so there is no asymptote. **E.** $f'(x) = \tfrac{5}{3}x^{2/3} - \tfrac{10}{3}x^{-1/3} = \tfrac{5}{3}x^{-1/3}(x-2) > 0 \quad \Leftrightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x < 0$ or $x > 2$, so f is increasing on $(-\infty, 0)$, $(2, \infty)$ and decreasing on $(0, 2)$.

F. Local maximum value $f(0) = 0$, local minimum value $f(2) = -3\sqrt[3]{4}$

G. $f''(x) = \frac{10}{9}x^{-1/3} + \frac{10}{9}x^{-4/3} = \frac{10}{9}x^{-4/3}(x+1) > 0 \quad \Leftrightarrow \quad x > -1$, so f is CU on $(-1, 0)$ and $(0, \infty)$, CD on $(-\infty, -1)$. IP at $(-1, -6)$

H.

31. $y = f(x) = \sqrt[3]{x^2 - 1}$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = -1$; x-intercepts: $f(x) = 0 \quad \Leftrightarrow \quad x^2 - 1 = 0 \quad \Leftrightarrow \quad x = \pm 1$ **C.** $f(-x) = f(x)$, so the curve is symmetric about the y-axis. **D.** No asymptote

E. $f'(x) = \frac{1}{3}(x^2 - 1)^{-2/3}(2x) = \dfrac{2x}{3\sqrt[3]{(x^2-1)^2}}$. $f'(x) > 0 \quad \Leftrightarrow \quad x > 0$ and $f'(x) < 0 \quad \Leftrightarrow \quad x < 0$, so f is increasing on $(0, \infty)$ and decreasing on $(-\infty, 0)$. **F.** Local minimum value $f(0) = -1$

G.
$$f''(x) = \frac{2}{3} \cdot \frac{(x^2-1)^{2/3}(1) - x \cdot \frac{2}{3}(x^2-1)^{-1/3}(2x)}{[(x^2-1)^{2/3}]^2}$$
$$= \frac{2}{9} \cdot \frac{(x^2-1)^{-1/3}[3(x^2-1) - 4x^2]}{(x^2-1)^{4/3}} = -\frac{2(x^2+3)}{9(x^2-1)^{5/3}}$$

$f''(x) > 0 \quad \Leftrightarrow \quad -1 < x < 1$ and $f''(x) < 0 \quad \Leftrightarrow \quad x < -1$ or $x > 1$, so f is CU on $(-1, 1)$ and f is CD on $(-\infty, -1)$ and $(1, \infty)$. IP at $(\pm 1, 0)$

H.

32. $y = f(x) = \sqrt[3]{x^3 + 1}$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 1$; x-intercept: $f(x) = 0 \quad \Leftrightarrow \quad x^3 + 1 = 0 \quad \Rightarrow \quad x = -1$

C. No symmetry **D.** No asymptote **E.** $f'(x) = \frac{1}{3}(x^3+1)^{-2/3}(3x^2) = \dfrac{x^2}{\sqrt[3]{(x^3+1)^2}}$. $f'(x) > 0$ if $x < -1$, $-1 < x < 0$, and $x > 0$, so f is increasing on $\mathbb{R}$. **F.** No local extrema

G.
$$f''(x) = \frac{(x^3+1)^{2/3}(2x) - x^2 \cdot \frac{2}{3}(x^3+1)^{-1/3}(3x^2)}{[(x^3+1)^{2/3}]^2}$$
$$= \frac{x(x^3+1)^{-1/3}[2(x^3+1) - 2x^3]}{(x^3+1)^{4/3}} = \frac{2x}{(x^3+1)^{5/3}}$$

$f''(x) > 0 \quad \Leftrightarrow \quad x < -1$ or $x > 0$ and $f''(x) < 0 \quad \Leftrightarrow \quad -1 < x < 0$, so f is CU on $(-\infty, -1)$ and $(0, \infty)$ and CD on $(-1, 0)$. IP at $(-1, 0)$ and $(0, 1)$

H.

33. $y = f(x) = \sin^3 x$ **A.** $D = \mathbb{R}$ **B.** x-intercepts: $f(x) = 0 \quad \Leftrightarrow \quad x = n\pi$, n an integer; y-intercept $= f(0) = 0$

C. $f(-x) = -f(x)$, so f is odd and the curve is symmetric about the origin. Also, $f(x + 2\pi) = f(x)$, so f is periodic with period 2π, and we determine **E–G** for $0 \le x \le \pi$. Since f is odd, we can reflect the graph of f on $[0, \pi]$ about the origin to obtain the graph of f on $[-\pi, \pi]$, and then since f has period 2π, we can extend the graph of f for all real numbers.

D. No asymptote **E.** $f'(x) = 3\sin^2 x \cos x > 0 \quad \Leftrightarrow \quad \cos x > 0$ and $\sin x \neq 0 \quad \Leftrightarrow \quad 0 < x < \frac{\pi}{2}$, so f is increasing on $\left(0, \frac{\pi}{2}\right)$ and f is decreasing on $\left(\frac{\pi}{2}, \pi\right)$. **F.** Local maximum value $f\left(\frac{\pi}{2}\right) = 1$ $\left[\text{local minimum value } f\left(-\frac{\pi}{2}\right) = -1\right]$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

G. $f''(x) = 3\sin^2 x\,(-\sin x) + 3\cos x\,(2\sin x\,\cos x) = 3\sin x\,(2\cos^2 x - \sin^2 x)$

$= 3\sin x\,[2(1-\sin^2 x) - \sin^2 x] = 3\sin x(2 - 3\sin^2 x) > 0 \quad \Leftrightarrow$

$\sin x > 0$ and $\sin^2 x < \frac{2}{3} \quad \Leftrightarrow \quad 0 < x < \pi$ and $0 < \sin x < \sqrt{\frac{2}{3}} \quad \Leftrightarrow \quad 0 < x < \sin^{-1}\sqrt{\frac{2}{3}}$ $\left[\text{let } \alpha = \sin^{-1}\sqrt{\frac{2}{3}}\right]$ or

$\pi - \alpha < x < \pi$, so f is CU on $(0, \alpha)$ and $(\pi - \alpha, \pi)$, and f is CD on $(\alpha, \pi - \alpha)$. There are inflection points at $x = 0$, π, α, and $x = \pi - \alpha$.

H.

34. $y = f(x) = x + \cos x$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 1$; the x-intercept is about -0.74 and can be found using Newton's method **C.** No symmetry **D.** No asymptote **E.** $f'(x) = 1 - \sin x > 0$ except for $x = \frac{\pi}{2} + 2n\pi$, so f is increasing on $\mathbb{R}$. **F.** No local extrema

G. $f''(x) = -\cos x$. $f''(x) > 0 \quad \Rightarrow \quad -\cos x > 0 \quad \Rightarrow \quad \cos x < 0 \quad \Rightarrow$ x is in $\left(\frac{\pi}{2} + 2n\pi, \frac{3\pi}{2} + 2n\pi\right)$ and $f''(x) < 0 \quad \Rightarrow$ x is in $\left(-\frac{\pi}{2} + 2n\pi, \frac{\pi}{2} + 2n\pi\right)$, so f is CU on $\left(\frac{\pi}{2} + 2n\pi, \frac{3\pi}{2} + 2n\pi\right)$ and CD on $\left(-\frac{\pi}{2} + 2n\pi, \frac{\pi}{2} + 2n\pi\right)$. IP at $\left(\frac{\pi}{2} + n\pi, f\left(\frac{\pi}{2} + n\pi\right)\right) = \left(\frac{\pi}{2} + n\pi, \frac{\pi}{2} + n\pi\right)$ [on the line $y = x$]

H.

35. $y = f(x) = x\tan x$, $-\frac{\pi}{2} < x < \frac{\pi}{2}$ **A.** $D = \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ **B.** Intercepts are 0 **C.** $f(-x) = f(x)$, so the curve is symmetric about the y-axis. **D.** $\lim\limits_{x \to (\pi/2)^-} x\tan x = \infty$ and $\lim\limits_{x \to -(\pi/2)^+} x\tan x = \infty$, so $x = \frac{\pi}{2}$ and $x = -\frac{\pi}{2}$ are VA.

E. $f'(x) = \tan x + x\sec^2 x > 0 \quad \Leftrightarrow \quad 0 < x < \frac{\pi}{2}$, so f increases on $\left(0, \frac{\pi}{2}\right)$ and decreases on $\left(-\frac{\pi}{2}, 0\right)$. **F.** Absolute and local minimum value $f(0) = 0$.

G. $y'' = 2\sec^2 x + 2x\tan x\,\sec^2 x > 0$ for $-\frac{\pi}{2} < x < \frac{\pi}{2}$, so f is CU on $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$. No IP

H.

36. $y = f(x) = 2x - \tan x$, $-\frac{\pi}{2} < x < \frac{\pi}{2}$ **A.** $D = \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ **B.** y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \quad \Leftrightarrow$ $2x = \tan x \quad \Leftrightarrow \quad x = 0$ or $x \approx \pm 1.17$ **C.** $f(-x) = -f(x)$, so f is odd; the graph is symmetric about the origin.

D. $\lim\limits_{x \to (-\pi/2)^+} (2x - \tan x) = \infty$ and $\lim\limits_{x \to (\pi/2)^-} (2x - \tan x) = -\infty$, so $x = \pm\frac{\pi}{2}$ are VA. No HA.

E. $f'(x) = 2 - \sec^2 x < 0 \quad \Leftrightarrow \quad |\sec x| > \sqrt{2}$ and $f'(x) > 0 \quad \Leftrightarrow \quad |\sec x| < \sqrt{2}$, so f is decreasing on $\left(-\frac{\pi}{2}, -\frac{\pi}{4}\right)$,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

increasing on $\left(-\frac{\pi}{4}, \frac{\pi}{4}\right)$, and decreasing again on $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$ **F.** Local maximum value $f\left(\frac{\pi}{4}\right) = \frac{\pi}{2} - 1$, local minimum value $f\left(-\frac{\pi}{4}\right) = -\frac{\pi}{2} + 1$

G. $f''(x) = -2\sec x \cdot \sec x \tan x = -2\tan x \sec^2 x = -2\tan x(\tan^2 x + 1)$ so $f''(x) > 0 \Leftrightarrow \tan x < 0 \Leftrightarrow -\frac{\pi}{2} < x < 0$, and $f''(x) < 0 \Leftrightarrow \tan x > 0 \Leftrightarrow 0 < x < \frac{\pi}{2}$. Thus, f is CU on $\left(-\frac{\pi}{2}, 0\right)$ and CD on $\left(0, \frac{\pi}{2}\right)$. IP at $(0, 0)$

H.

37. $y = f(x) = \sin x + \sqrt{3}\cos x$, $-2\pi \le x \le 2\pi$ **A.** $D = [-2\pi, 2\pi]$ **B.** y-intercept: $f(0) = \sqrt{3}$; x-intercepts: $f(x) = 0 \Leftrightarrow \sin x = -\sqrt{3}\cos x \Leftrightarrow \tan x = -\sqrt{3} \Leftrightarrow x = -\frac{4\pi}{3}, -\frac{\pi}{3}, \frac{2\pi}{3}$, or $\frac{5\pi}{3}$ **C.** f is periodic with period 2π. **D.** No asymptote **E.** $f'(x) = \cos x - \sqrt{3}\sin x$. $f'(x) = 0 \Leftrightarrow \cos x = \sqrt{3}\sin x \Leftrightarrow \tan x = \dfrac{1}{\sqrt{3}} \Leftrightarrow x = -\frac{11\pi}{6}, -\frac{5\pi}{6}, \frac{\pi}{6}$, or $\frac{7\pi}{6}$. $f'(x) < 0 \Leftrightarrow -\frac{11\pi}{6} < x < -\frac{5\pi}{6}$ or $\frac{\pi}{6} < x < \frac{7\pi}{6}$, so f is decreasing on $\left(-\frac{11\pi}{6}, -\frac{5\pi}{6}\right)$ and $\left(\frac{\pi}{6}, \frac{7\pi}{6}\right)$, and f is increasing on $\left(-2\pi, -\frac{11\pi}{6}\right)$, $\left(-\frac{5\pi}{6}, \frac{\pi}{6}\right)$, and $\left(\frac{7\pi}{6}, 2\pi\right)$. **F.** Local maximum value $f\left(-\frac{11\pi}{6}\right) = f\left(\frac{\pi}{6}\right) = \frac{1}{2} + \sqrt{3}\left(\frac{1}{2}\sqrt{3}\right) = 2$, local minimum value $f\left(-\frac{5\pi}{6}\right) = f\left(\frac{7\pi}{6}\right) = -\frac{1}{2} + \sqrt{3}\left(-\frac{1}{2}\sqrt{3}\right) = -2$

G. $f''(x) = -\sin x - \sqrt{3}\cos x$. $f''(x) = 0 \Leftrightarrow \sin x = -\sqrt{3}\cos x \Leftrightarrow \tan x = -\dfrac{1}{\sqrt{3}} \Leftrightarrow x = -\frac{4\pi}{3}, -\frac{\pi}{3}, \frac{2\pi}{3}$, or $\frac{5\pi}{3}$. $f''(x) > 0 \Leftrightarrow -\frac{4\pi}{3} < x < -\frac{\pi}{3}$ or $\frac{2\pi}{3} < x < \frac{5\pi}{3}$, so f is CU on $\left(-\frac{4\pi}{3}, -\frac{\pi}{3}\right)$ and $\left(\frac{2\pi}{3}, \frac{5\pi}{3}\right)$, and f is CD on $\left(-2\pi, -\frac{4\pi}{3}\right)$, $\left(-\frac{\pi}{3}, \frac{2\pi}{3}\right)$, and $\left(\frac{5\pi}{3}, 2\pi\right)$. There are IPs at $\left(-\frac{4\pi}{3}, 0\right)$, $\left(-\frac{\pi}{3}, 0\right)$, $\left(\frac{2\pi}{3}, 0\right)$, and $\left(\frac{5\pi}{3}, 0\right)$.

H.

38. $y = f(x) = \csc x - 2\sin x$, $0 < x < \pi$ **A.** $D = (0, \pi)$ **B.** No y-intercept; x-intercept: $f(x) = 0 \Leftrightarrow \csc x = 2\sin x \Leftrightarrow \frac{1}{2} = \sin^2 x \Leftrightarrow \sin x = \pm\frac{1}{2}\sqrt{2} \Leftrightarrow x = \frac{\pi}{4}$ or $\frac{3\pi}{4}$ **C.** No symmetry

D. $\lim\limits_{x \to 0^+} f(x) = \infty$ and $\lim\limits_{x \to \pi^-} f(x) = \infty$, so $x = 0$ and $x = \pi$ are VAs.

E. $f'(x) = -\csc x \cot x - 2\cos x = -\dfrac{\cos x}{\sin^2 x} - 2\cos x = -\cos x\left(\dfrac{1}{\sin^2 x} + 2\right)$. $f'(x) > 0$ when $-\cos x > 0 \Leftrightarrow \cos x < 0 \Leftrightarrow \frac{\pi}{2} < x < \pi$, so f' is increasing on $\left(\frac{\pi}{2}, \pi\right)$, and f is decreasing on $\left(0, \frac{\pi}{2}\right)$. **F.** Local minimum value $f\left(\frac{\pi}{2}\right) = -1$

G. $f''(x) = (-\csc x)(-\csc^2 x) + (\cot x)(\csc x \ \cot x) + 2\sin x$

$$= \frac{1 + \cos^2 x + 2\sin^4 x}{\sin^3 x}$$

f'' has the same sign as $\sin x$, which is positive on $(0, \pi)$, so f is CU on $(0, \pi)$. No IP

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. $y = f(x) = \dfrac{\sin x}{1+\cos x}$ $\left[\overset{\substack{\text{when}\\ \cos x \neq 1}}{=} \dfrac{\sin x}{1+\cos x}\cdot\dfrac{1-\cos x}{1-\cos x} = \dfrac{\sin x\,(1-\cos x)}{\sin^2 x} = \dfrac{1-\cos x}{\sin x} = \csc x - \cot x\right]$

A. The domain of f is the set of all real numbers except odd integer multiples of π; that is, all reals except $(2n+1)\pi$, where n is an integer. **B.** y-intercept: $f(0) = 0$; x-intercepts: $x = 2n\pi$, n an integer. **C.** $f(-x) = -f(x)$, so f is an odd function; the graph is symmetric about the origin and has period 2π. **D.** When n is an odd integer, $\lim\limits_{x\to(n\pi)^-} f(x) = \infty$ and $\lim\limits_{x\to(n\pi)^+} f(x) = -\infty$, so $x = n\pi$ is a VA for each odd integer n. No HA.

E. $f'(x) = \dfrac{(1+\cos x)\cdot\cos x - \sin x(-\sin x)}{(1+\cos x)^2} = \dfrac{1+\cos x}{(1+\cos x)^2} = \dfrac{1}{1+\cos x}$. $f'(x) > 0$ for all x except odd multiples of π, so f is increasing on $((2k-1)\pi, (2k+1)\pi)$ for each integer k. **F.** No extreme values

G. $f''(x) = \dfrac{\sin x}{(1+\cos x)^2} > 0 \;\Rightarrow\; \sin x > 0 \;\Rightarrow$ $x \in (2k\pi, (2k+1)\pi)$ and $f''(x) < 0$ on $((2k-1)\pi, 2k\pi)$ for each integer k. f is CU on $(2k\pi, (2k+1)\pi)$ and CD on $((2k-1)\pi, 2k\pi)$ for each integer k. f has IPs at $(2k\pi, 0)$ for each integer k.

H.

40. $y = f(x) = \dfrac{\sin x}{2+\cos x}$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \;\Leftrightarrow\; \sin x = 0 \;\Leftrightarrow\; x = n\pi$

C. $f(-x) = -f(x)$, so the curve is symmetric about the origin. f is periodic with period 2π, so we determine **E–G** for $0 \le x \le 2\pi$. **D.** No asymptote

E. $f'(x) = \dfrac{(2+\cos x)\cos x - \sin x(-\sin x)}{(2+\cos x)^2} = \dfrac{2\cos x + \cos^2 x + \sin^2 x}{(2+\cos x)^2} = \dfrac{2\cos x + 1}{(2+\cos x)^2}$.

$f'(x) > 0 \;\Leftrightarrow\; 2\cos x + 1 > 0 \;\Leftrightarrow\; \cos x > -\frac{1}{2} \;\Leftrightarrow\; x$ is in $\left(0, \frac{2\pi}{3}\right)$ or $\left(\frac{4\pi}{3}, 2\pi\right)$, so f is increasing on $\left(0, \frac{2\pi}{3}\right)$ and $\left(\frac{4\pi}{3}, 2\pi\right)$, and f is decreasing on $\left(\frac{2\pi}{3}, \frac{4\pi}{3}\right)$.

F. Local maximum value $f\left(\frac{2\pi}{3}\right) = \dfrac{\sqrt{3}/2}{2-(1/2)} = \dfrac{\sqrt{3}}{3}$ and local minimum value $f\left(\frac{4\pi}{3}\right) = \dfrac{-\sqrt{3}/2}{2-(1/2)} = -\dfrac{\sqrt{3}}{3}$

G. $f''(x) = \dfrac{(2+\cos x)^2(-2\sin x) - (2\cos x+1)2(2+\cos x)(-\sin x)}{[(2+\cos x)^2]^2}$

$$= \frac{-2\sin x\,(2+\cos x)[(2+\cos x) - (2\cos x+1)]}{(2+\cos x)^4} = \frac{-2\sin x\,(1-\cos x)}{(2+\cos x)^3}$$

$f''(x) > 0 \;\Leftrightarrow\; -2\sin x > 0 \;\Leftrightarrow\; \sin x < 0 \;\Leftrightarrow\; x$ is in $(\pi, 2\pi)$ [f is CU] and $f''(x) < 0 \;\Leftrightarrow$ x is in $(0, \pi)$ [f is CD]. The inflection points are $(0, 0)$, $(\pi, 0)$, and $(2\pi, 0)$.

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. $y = f(x) = \arctan(e^x)$ **A.** $D = \mathbb{R}$ **B.** y-intercept $= f(0) = \arctan 1 = \frac{\pi}{4}$. $f(x) > 0$ so there are no x-intercepts.

C. No symmetry **D.** $\lim\limits_{x\to-\infty} \arctan(e^x) = 0$ and $\lim\limits_{x\to\infty} \arctan(e^x) = \frac{\pi}{2}$, so $y = 0$ and $y = \frac{\pi}{2}$ are HAs. No VA

E. $f'(x) = \dfrac{1}{1+(e^x)^2}\dfrac{d}{dx}e^x = \dfrac{e^x}{1+e^{2x}} > 0$, so f is increasing on $(-\infty, \infty)$. **F.** No local extrema

G. $f''(x) = \dfrac{(1+e^{2x})e^x - e^x(2e^{2x})}{(1+e^{2x})^2} = \dfrac{e^x[(1+e^{2x}) - 2e^{2x}]}{(1+e^{2x})^2}$

$= \dfrac{e^x(1-e^{2x})}{(1+e^{2x})^2} > 0 \quad \Leftrightarrow$

$1 - e^{2x} > 0 \quad \Leftrightarrow \quad e^{2x} < 1 \quad \Leftrightarrow \quad 2x < 0 \quad \Leftrightarrow \quad x < 0$, so f is CU on $(-\infty, 0)$ and CD on $(0, \infty)$. IP at $\left(0, \frac{\pi}{4}\right)$

H.

42. $y = f(x) = (1-x)e^x$ **A.** $D = \mathbb{R}$ **B.** x-intercept 1, y-intercept $= f(0) = 1$ **C.** No symmetry

D. $\lim\limits_{x\to-\infty} \dfrac{1-x}{e^{-x}}$ $\left[\text{form } \frac{\infty}{\infty}\right] \overset{\text{H}}{=} \lim\limits_{x\to-\infty} \dfrac{-1}{-e^{-x}} = 0$, so $y = 0$ is a HA. No VA

E. $f'(x) = (1-x)e^x + e^x(-1) = e^x[(1-x) + (-1)] = -xe^x > 0 \quad \Leftrightarrow \quad x < 0$, so f is increasing on $(-\infty, 0)$ and decreasing on $(0, \infty)$.

F. Local maximum value $f(0) = 1$, no local minimum value

G. $f''(x) = -xe^x + e^x(-1) = e^x(-x-1) = -(x+1)e^x > 0 \quad \Leftrightarrow$

$x < -1$, so f is CU on $(-\infty, -1)$ and CD on $(-1, \infty)$. IP at $(-1, 2/e)$

H.

43. $y = 1/(1+e^{-x})$ **A.** $D = \mathbb{R}$ **B.** No x-intercept; y-intercept $= f(0) = \frac{1}{2}$. **C.** No symmetry

D. $\lim\limits_{x\to\infty} 1/(1+e^{-x}) = \frac{1}{1+0} = 1$ and $\lim\limits_{x\to-\infty} 1/(1+e^{-x}) = 0$ since $\lim\limits_{x\to-\infty} e^{-x} = \infty$, so f has horizontal asymptotes $y = 0$ and $y = 1$. **E.** $f'(x) = -(1+e^{-x})^{-2}(-e^{-x}) = e^{-x}/(1+e^{-x})^2$. This is positive for all x, so f is increasing on $\mathbb{R}$.

F. No extreme values **G.** $f''(x) = \dfrac{(1+e^{-x})^2(-e^{-x}) - e^{-x}(2)(1+e^{-x})(-e^{-x})}{(1+e^{-x})^4} = \dfrac{e^{-x}(e^{-x}-1)}{(1+e^{-x})^3}$

The second factor in the numerator is negative for $x > 0$ and positive for $x < 0$, and the other factors are always positive, so f is CU on $(-\infty, 0)$ and CD on $(0, \infty)$. IP at $\left(0, \frac{1}{2}\right)$

H.

44. $y = f(x) = e^{-x}\sin x,\ 0 \le x \le 2\pi$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \quad \Leftrightarrow \quad \sin x = 0 \quad \Leftrightarrow$ $x = 0, \pi$, and 2π. **C.** No symmetry **D.** No asymptote **E.** $f'(x) = e^{-x}\cos x + \sin x\,(-e^{-x}) = e^{-x}(\cos x - \sin x)$.

$f'(x) = 0 \quad \Leftrightarrow \quad \cos x = \sin x \quad \Leftrightarrow \quad x = \frac{\pi}{4}, \frac{5\pi}{4}$. $f'(x) > 0$ if x is in $\left(0, \frac{\pi}{4}\right)$ or $\left(\frac{5\pi}{4}, 2\pi\right)$ [f is increasing] and $f'(x) < 0$ if x is in $\left(\frac{\pi}{4}, \frac{5\pi}{4}\right)$ [f is decreasing]. **F.** Local maximum value $f\left(\frac{\pi}{4}\right)$ and local minimum value $f\left(\frac{5\pi}{4}\right)$

G. $f''(x) = e^{-x}(-\sin x - \cos x) + (\cos x - \sin x)(-e^{-x}) = e^{-x}(-2\cos x)$. $f''(x) > 0 \quad \Leftrightarrow \quad -2\cos x > 0 \quad \Leftrightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\cos x < 0 \quad \Rightarrow \quad x$ is in $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$ [f is CU] and $f''(x) < 0 \quad \Leftrightarrow$

$\cos x > 0 \quad \Rightarrow \quad x$ is in $\left(0, \frac{\pi}{2}\right)$ or $\left(\frac{3\pi}{2}, 2\pi\right)$ [f is CD].

IP at $\left(\frac{\pi}{2} + n\pi, f\left(\frac{\pi}{2} + n\pi\right)\right)$

H.

45. $y = f(x) = \dfrac{1}{x} + \ln x$ **A.** $D = (0, \infty)$ [same as $\ln x$] **B.** No y-intercept; no x-intercept [$1/x > |\ln x|$ on $(0, 1)$, and $1/x$ and $\ln x$ are both positive on $(1, \infty)$] **C.** No symmetry **D.** $\lim\limits_{x \to 0^+} f(x) = \infty$, so $x = 0$ is a VA.

E. $f'(x) = -\dfrac{1}{x^2} + \dfrac{1}{x} = \dfrac{x-1}{x^2}$. $f'(x) > 0$ for $x > 1$, so f is increasing on $(1, \infty)$ and f is decreasing on $(0, 1)$.

F. Local minimum value $f(1) = 1$ **G.** $f''(x) = \dfrac{2}{x^3} - \dfrac{1}{x^2} = \dfrac{2-x}{x^3}$.

$f''(x) > 0$ for $0 < x < 2$, so f is CU on $(0, 2)$, and f is CD on $(2, \infty)$.

IP at $\left(2, \frac{1}{2} + \ln 2\right)$

H.

46. $y = x(\ln x)^2$ **A.** $D = (0, \infty)$ [same as $\ln x$] **B.** No y-intercept; x-intercepts: $f(x) = 0 \quad \Rightarrow$ $(\ln x)^2 = 0$ [$x = 0$ not in domain] $\Rightarrow \quad \ln x = 0 \quad \Rightarrow \quad x = 1$ **C.** No symmetry **D.** No HA;

$$\lim_{x \to 0^+} x(\ln x)^2 = \lim_{x \to 0^+} \frac{(\ln x)^2}{1/x} \overset{\text{H}}{=} \lim_{x \to 0^+} \frac{2\ln x \cdot 1/x}{-1/x^2} = \lim_{x \to 0^+} \frac{2\ln x}{-1/x} \overset{\text{H}}{=} \lim_{x \to 0^+} \frac{2/x}{1/x^2} = \lim_{x \to 0^+} (2x) = 0, \text{ no VA.}$$

E. $f'(x) = x \cdot 2\ln x \cdot \left(\dfrac{1}{x}\right) + (\ln x)^2 \cdot 1 = 2\ln x + (\ln x)^2 = \ln x\,(2 + \ln x) = 0 \quad \Leftrightarrow \quad x = 1$ or $x = e^{-2}$;

$f'(x) > 0 \quad \Leftrightarrow \quad 0 < x < e^{-2}$ or $x > 1$ and $f'(x) < 0 \quad \Leftrightarrow \quad e^{-2} < x < 1$, so f is increasing on $(0, e^{-2})$ and $(1, \infty)$, and decreasing on $\left(e^{-2}, 1\right)$. **F.** Local maximum value $f(e^{-2}) = 4e^{-2}$, local minimum value $f(1) = 0$ **G.** $f''(x) = \dfrac{2}{x} + 2(\ln x) \cdot \dfrac{1}{x} = \dfrac{2(1 + \ln x)}{x}$, so

$f''(x) > 0 \quad \Leftrightarrow \quad 1 + \ln x > 0 \quad \Leftrightarrow \quad x > e^{-1}$ and $f''(x) < 0 \quad \Leftrightarrow$

$0 < x < e^{-1}$. Thus, f is CD on $(0, e^{-1})$ and CU on (e^{-1}, ∞). IP at (e^{-1}, e^{-1})

H.

47. $y = f(x) = (1 + e^x)^{-2} = \dfrac{1}{(1 + e^x)^2}$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = \frac{1}{4}$. x-intercepts: none [since $f(x) > 0$]

C. No symmetry **D.** $\lim\limits_{x \to \infty} f(x) = 0$ and $\lim\limits_{x \to -\infty} f(x) = 1$, so $y = 0$ and $y = 1$ are HA; no VA

E. $f'(x) = -2(1 + e^x)^{-3} e^x = \dfrac{-2e^x}{(1 + e^x)^3} < 0$, so f is decreasing on $\mathbb{R}$ **F.** No local extrema

G. $f''(x) = (1 + e^x)^{-3}(-2e^x) + (-2e^x)(-3)(1 + e^x)^{-4} e^x$

$$= -2e^x(1 + e^x)^{-4}[(1 + e^x) - 3e^x] = \frac{-2e^x(1 - 2e^x)}{(1 + e^x)^4}.$$

$f''(x) > 0 \quad \Leftrightarrow \quad 1 - 2e^x < 0 \quad \Leftrightarrow \quad e^x > \frac{1}{2} \quad \Leftrightarrow \quad x > \ln \frac{1}{2}$ and

$f''(x) < 0 \quad \Leftrightarrow \quad x < \ln \frac{1}{2}$, so f is CU on $\left(\ln \frac{1}{2}, \infty\right)$ and CD on $\left(-\infty, \ln \frac{1}{2}\right)$.

IP at $\left(\ln \frac{1}{2}, \frac{4}{9}\right)$

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. $y = f(x) = e^x/x^2$ **A.** $D = (-\infty, 0) \cup (0, \infty)$ **B.** No intercept **C.** No symmetry **D.** $\lim\limits_{x\to-\infty} \dfrac{e^x}{x^2} = 0$, so $y = 0$ is HA. $\lim\limits_{x\to 0} \dfrac{e^x}{x^2} = \infty$, so $x = 0$ is a VA. **E.** $f'(x) = \dfrac{x^2e^x - e^x(2x)}{(x^2)^2} = \dfrac{xe^x(x-2)}{x^4} = \dfrac{e^x(x-2)}{x^3} > 0 \quad \Leftrightarrow \quad x < 0$ or $x > 2$, so f is increasing on $(-\infty, 0)$ and $(2, \infty)$, and f is decreasing on $(0, 2)$.

H.

F. Local minimum value $f(2) = e^2/4 \approx 1.85$, no local maximum value

G. $f''(x) = \dfrac{x^3[e^x(1) + (x-2)e^x] - e^x(x-2)(3x^2)}{(x^3)^2}$

$$= \frac{x^2e^x[x(x-1) - 3(x-2)]}{x^6} = \frac{e^x(x^2 - 4x + 6)}{x^4} > 0$$

for all x in the domain of f; that is, f is CU on $(-\infty, 0)$ and $(0, \infty)$. No IP

49. $y = f(x) = \ln(\sin x)$

A. $D = \{x \text{ in } \mathbb{R} \mid \sin x > 0\} = \bigcup\limits_{n=-\infty}^{\infty} (2n\pi, (2n+1)\pi) = \cdots \cup (-4\pi, -3\pi) \cup (-2\pi, -\pi) \cup (0, \pi) \cup (2\pi, 3\pi) \cup \cdots$

B. No y-intercept; x-intercepts: $f(x) = 0 \quad \Leftrightarrow \quad \ln(\sin x) = 0 \quad \Leftrightarrow \quad \sin x = e^0 = 1 \quad \Leftrightarrow \quad x = 2n\pi + \frac{\pi}{2}$ for each integer n. **C.** f is periodic with period 2π. **D.** $\lim\limits_{x\to(2n\pi)^+} f(x) = -\infty$ and $\lim\limits_{x\to[(2n+1)\pi]^-} f(x) = -\infty$, so the lines $x = n\pi$ are VAs for all integers n. **E.** $f'(x) = \dfrac{\cos x}{\sin x} = \cot x$, so $f'(x) > 0$ when $2n\pi < x < 2n\pi + \frac{\pi}{2}$ for each integer n, and $f'(x) < 0$ when $2n\pi + \frac{\pi}{2} < x < (2n+1)\pi$. Thus, f is increasing on $\left(2n\pi, 2n\pi + \frac{\pi}{2}\right)$ and decreasing on $\left(2n\pi + \frac{\pi}{2}, (2n+1)\pi\right)$ for each integer n.

H.

F. Local maximum values $f\left(2n\pi + \frac{\pi}{2}\right) = 0$, no local minimum.

G. $f''(x) = -\csc^2 x < 0$, so f is CD on $(2n\pi, (2n+1)\pi)$ for each integer n. No IP

50. $y = f(x) = \ln(1 + x^3)$ **A.** $1 + x^3 > 0 \quad \Leftrightarrow \quad x^3 > -1 \quad \Leftrightarrow \quad x > -1$, so $D = (-1, \infty)$. **B.** y-intercept: $f(0) = \ln 1 = 0$; x-intercept: $f(x) = 0 \quad \Leftrightarrow \quad \ln(1 + x^3) = 0 \quad \Leftrightarrow \quad 1 + x^3 = e^0 \quad \Leftrightarrow \quad x^3 = 0 \quad \Leftrightarrow \quad x = 0$ **C.** No symmetry **D.** $\lim\limits_{x\to-1^+} f(x) = -\infty$, so $x = -1$ is a VA **E.** $f'(x) = \dfrac{3x^2}{1 + x^3}$. $f'(x) > 0$ on $(-1, 0)$ and $(0, \infty)$ [$f'(x) = 0$ at $x = 0$], so by Exercise 4.3.97, f is increasing on $(-1, \infty)$. **F.** No extreme values

G. $f''(x) = \dfrac{(1 + x^3)(6x) - 3x^2(3x^2)}{(1 + x^3)^2}$

$$= \frac{3x[2(1 + x^3) - 3x^3]}{(1 + x^3)^2} = \frac{3x(2 - x^3)}{(1 + x^3)^2}$$

$f''(x) > 0 \quad \Leftrightarrow \quad 0 < x < \sqrt[3]{2}$, so f is CU on $\left(0, \sqrt[3]{2}\right)$ and f is CD on $(-1, 0)$ and $\left(\sqrt[3]{2}, \infty\right)$. IP at $(0, 0)$ and $\left(\sqrt[3]{2}, \ln 3\right)$

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. $y = f(x) = xe^{-1/x}$ **A.** $D = (-\infty, 0) \cup (0, \infty)$ **B.** No intercept **C.** No symmetry

D. $\lim\limits_{x\to 0^-} \dfrac{e^{-1/x}}{1/x} \overset{\text{H}}{=} \lim\limits_{x\to 0^-} \dfrac{e^{-1/x}(1/x^2)}{-1/x^2} = -\lim\limits_{x\to 0^-} e^{-1/x} = -\infty$, so $x = 0$ is a VA. Also, $\lim\limits_{x\to 0^+} xe^{-1/x} = 0$, so the graph approaches the origin as $x \to 0^+$. **E.** $f'(x) = xe^{-1/x}\left(\dfrac{1}{x^2}\right) + e^{-1/x}(1) = e^{-1/x}\left(\dfrac{1}{x} + 1\right) = \dfrac{x+1}{xe^{1/x}} > 0 \quad \Leftrightarrow$ $x < -1$ or $x > 0$, so f is increasing on $(-\infty, -1)$ and $(0, \infty)$, and f is decreasing on $(-1, 0)$.

F. Local maximum value $f(-1) = -e$, no local minimum value

G. $f'(x) = e^{-1/x}\left(\dfrac{1}{x} + 1\right) \quad \Rightarrow$

$$f''(x) = e^{-1/x}\left(-\frac{1}{x^2}\right) + \left(\frac{1}{x} + 1\right)e^{-1/x}\left(\frac{1}{x^2}\right)$$
$$= \frac{1}{x^2}e^{-1/x}\left[-1 + \left(\frac{1}{x} + 1\right)\right] = \frac{1}{x^3 e^{1/x}} > 0 \quad \Leftrightarrow$$

$x > 0$, so f is CU on $(0, \infty)$ and CD on $(-\infty, 0)$. No IP

H.

52. $y = f(x) = \dfrac{\ln x}{x^2}$ **A.** $D = (0, \infty)$ **B.** y-intercept: none; x-intercept: $f(x) = 0 \quad \Leftrightarrow \quad \ln x = 0 \quad \Leftrightarrow \quad x = 1$

C. No symmetry **D.** $\lim\limits_{x\to 0^+} f(x) = -\infty$, so $x = 0$ is a VA; $\lim\limits_{x\to\infty} \dfrac{\ln x}{x^2} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1/x}{2x} = 0$, so $y = 0$ is a HA.

E. $f'(x) = \dfrac{x^2(1/x) - (\ln x)(2x)}{(x^2)^2} = \dfrac{x(1 - 2\ln x)}{x^4} = \dfrac{1 - 2\ln x}{x^3}$. $f'(x) > 0 \quad \Leftrightarrow \quad 1 - 2\ln x > 0 \quad \Leftrightarrow \quad \ln x < \frac{1}{2} \quad \Rightarrow$ $0 < x < e^{1/2}$ and $f'(x) < 0 \quad \Rightarrow \quad x > e^{1/2}$, so f is increasing on $\left(0, \sqrt{e}\right)$ and decreasing on $\left(\sqrt{e}, \infty\right)$.

F. Local maximum value $f(e^{1/2}) = \dfrac{1/2}{e} = \dfrac{1}{2e}$

G. $f''(x) = \dfrac{x^3(-2/x) - (1 - 2\ln x)(3x^2)}{(x^3)^2}$

$$= \frac{x^2[-2 - 3(1 - 2\ln x)]}{x^6} = \frac{-5 + 6\ln x}{x^4}$$

$f''(x) > 0 \quad \Leftrightarrow \quad -5 + 6\ln x > 0 \quad \Leftrightarrow \quad \ln x > \frac{5}{6} \quad \Rightarrow \quad x > e^{5/6}$ [f is CU]

and $f''(x) < 0 \quad \Leftrightarrow \quad 0 < x < e^{5/6}$ [f is CD]. IP at $(e^{5/6}, 5/(6e^{5/3}))$

H.

53. $y = f(x) = e^{\arctan x}$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = e^0 = 1$; no x-intercept since $e^{\arctan x}$ is positive for all x.

C. No symmetry **D.** $\lim\limits_{x\to -\infty} f(x) = e^{-\pi/2}$ [≈ 0.21], so $y = e^{-\pi/2}$ is a HA. $\lim\limits_{x\to\infty} f(x) = e^{\pi/2}$ [≈ 4.81], so $y = e^{\pi/2}$ is a HA. **E.** $f'(x) = e^{\arctan x}\left(\dfrac{1}{1 + x^2}\right)$. $f'(x) > 0$ for all x, so f is increasing on $\mathbb{R}$. **F.** No local extrema

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

G. $f''(x) = \dfrac{(1+x^2)e^{\arctan x}\left(\dfrac{1}{1+x^2}\right) - e^{\arctan x}(2x)}{(1+x^2)^2}$

$= \dfrac{e^{\arctan x}(1-2x)}{(1+x^2)^2}$

$f''(x) > 0$ for $x < \frac{1}{2}$, so f is CU on $\left(-\infty, \frac{1}{2}\right)$ and f is CD on $\left(\frac{1}{2}, \infty\right)$.

IP at $\left(\frac{1}{2}, e^{\arctan 1/2}\right) \approx (0.5, 1.59)$

H.

54. $y = f(x) = \tan^{-1}\left(\dfrac{x-1}{x+1}\right)$ **A.** $D = \{x \mid x \neq -1\}$ **B.** x-intercept $= 1$, y-intercept $= f(0) = \tan^{-1}(-1) = -\frac{\pi}{4}$

C. No symmetry **D.** $\lim\limits_{x\to\pm\infty} \tan^{-1}\left(\dfrac{x-1}{x+1}\right) = \lim\limits_{x\to\pm\infty} \tan^{-1}\left(\dfrac{1-1/x}{1+1/x}\right) = \tan^{-1} 1 = \frac{\pi}{4}$, so $y = \frac{\pi}{4}$ is a HA.

Also $\lim\limits_{x\to-1^+} \tan^{-1}\left(\dfrac{x-1}{x+1}\right) = -\dfrac{\pi}{2}$ and $\lim\limits_{x\to-1^-} \tan^{-1}\left(\dfrac{x-1}{x+1}\right) = \dfrac{\pi}{2}$.

E. $f'(x) = \dfrac{1}{1+[(x-1)/(x+1)]^2}\dfrac{(x+1)-(x-1)}{(x+1)^2} = \dfrac{2}{(x+1)^2+(x-1)^2} = \dfrac{1}{x^2+1} > 0$,

so f is increasing on $(-\infty, -1)$ and $(-1, \infty)$. **F.** No extreme values

G. $f''(x) = -2x/\left(x^2+1\right)^2 > 0 \quad \Leftrightarrow \quad x < 0$, so f is CU on $(-\infty, -1)$ and $(-1, 0)$, and CD on $(0, \infty)$. IP at $\left(0, -\frac{\pi}{4}\right)$

H.

55. $g(x) = \sqrt{f(x)}$

(a) The domain of g consists of all x such that $f(x) \geq 0$, so g has domain $(-\infty, 7]$. $g'(x) = \dfrac{1}{2\sqrt{f(x)}} \cdot f'(x) = \dfrac{f'(x)}{2\sqrt{f(x)}}$.

Since $f'(3)$ does not exist, $g'(3)$ does not exist. (Note that $f(7) = 0$, but 7 is an endpoint of the domain of g.) The domain of g' is $(-\infty, 3) \cup (3, 7)$.

(b) $g'(x) = 0 \quad \Rightarrow \quad f'(x) = 0$ on the domain of $g \quad \Rightarrow \quad x = 5$ [there is a horizontal tangent line there]. From part (a), $g'(3)$ does not exist. So the critical numbers of g are 3 and 5.

(c) From part (a), $g'(x) = \dfrac{f'(x)}{2\sqrt{f(x)}}$. $g'(6) = \dfrac{f'(6)}{2\sqrt{f(6)}} \approx \dfrac{-2}{2\sqrt{3}} = -\dfrac{1}{\sqrt{3}} \approx -0.58$.

(d) $\lim\limits_{x\to-\infty} g(x) = \lim\limits_{x\to-\infty} \sqrt{f(x)} = \sqrt{2}$, so $y = \sqrt{2}$ is a horizontal asymptote. No VA

56. $g(x) = \sqrt[3]{f(x)}$

(a) Since the cube-root function is defined for all reals, the domain of g equals the domain of f, $(-\infty, \infty)$.

$g'(x) = \dfrac{1}{3\left(\sqrt[3]{f(x)}\right)^2} \cdot f'(x) = \dfrac{f'(x)}{3\left(\sqrt[3]{f(x)}\right)^2}$. Since $f'(3)$ does not exist and $f(7) = 0$, $g'(3)$ and $g'(7)$ do not exist.

The domain of g' is $(-\infty, 3) \cup (3, 7) \cup (7, \infty)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $g'(x) = 0 \quad \Rightarrow \quad f'(x) = 0 \quad \Rightarrow \quad x = 5$ or $x = 9$ [there are horizontal tangent lines there]. From part (a), $g'(x)$ does not exist at $x = 3$ and $x = 7$. So the critical numbers of g are 3, 5, 7, and 9.

(c) From part (a), $g'(x) = \dfrac{f'(x)}{3\left(\sqrt[3]{f(x)}\right)^2}$. $g'(6) = \dfrac{f'(6)}{3\left(\sqrt[3]{f(6)}\right)^2} \approx \dfrac{-2}{3\left(\sqrt[3]{3}\right)^2} = -\dfrac{2}{3^{5/3}} \approx -0.32$.

(d) $\lim\limits_{x\to-\infty} g(x) = \lim\limits_{x\to-\infty} \sqrt[3]{f(x)} = \sqrt[3]{2}$ and $\lim\limits_{x\to\infty} g(x) = \lim\limits_{x\to\infty} \sqrt[3]{f(x)} = \sqrt[3]{-1} = -1$, so $y = \sqrt[3]{2}$ and $y = -1$ are horizontal asymptotes. No VA

57. $g(x) = |f(x)|$

(a) Since the absolute-value function is defined for all reals, the domain of g equals the domain of f, $(-\infty, \infty)$. The domain of g' equals the domain of f' except for any values of x such that both $f(x) = 0$ and $f'(x) \neq 0$. $f'(3)$ does not exist, $f(7) = 0$, and $f'(7) \neq 0$. Thus, the domain of g' is $(-\infty, 3) \cup (3, 7) \cup (7, \infty)$.

(b) $g'(x) = 0 \quad \Rightarrow \quad f'(x) = 0 \quad \Rightarrow \quad x = 5$ or $x = 9$ [there are horizontal tangent lines there]. From part (a), g' does not exist at $x = 3$ and $x = 7$. So the critical numbers of g are 3, 5, 7, and 9.

(c) Since f is positive near $x = 6$, $g(x) = |f(x)| = f(x)$ near 6, so $g'(6) = f'(6) \approx -2$.

(d) $\lim\limits_{x\to-\infty} g(x) = \lim\limits_{x\to-\infty} |f(x)| = |2| = 2$ and $\lim\limits_{x\to\infty} g(x) = \lim\limits_{x\to\infty} |f(x)| = |-1| = 1$, so $y = 2$ and $y = 1$ are horizontal asymptotes. No VA

58. $g(x) = 1/f(x)$

(a) The domain of g consists of all x such that $f(x) \neq 0$, so g has domain $(-\infty, 7) \cup (7, \infty)$. $g(x) = \dfrac{1}{f(x)} = [f(x)]^{-1} \quad \Rightarrow$ $g'(x) = -1[f(x)]^{-2} \cdot f'(x) = -\dfrac{f'(x)}{[f(x)]^2}$. The domain of g' will equal the domain of f except for any values of f such that $f(x) = 0$. $f'(3)$ does not exist, and $f(7) = 0$. Thus, the domain of g' is $(-\infty, 3) \cup (3, 7) \cup (7, \infty)$.

(b) $g'(x) = 0 \quad \Rightarrow \quad f'(x) = 0 \quad \Rightarrow \quad x = 5$ or $x = 9$ [there are horizontal tangent lines there]. From part (a), g' does not exist at $x = 3$ and $x = 7$. So the critical numbers of g are 3, 5, 7, and 9.

(c) From part (a), $g'(x) = -\dfrac{f'(x)}{[f(x)]^2}$. $g'(6) = -\dfrac{f'(6)}{[f(6)]^2} \approx -\left(\dfrac{-2}{3^2}\right) = \dfrac{2}{9}$.

(d) $\lim\limits_{x\to-\infty} g(x) = \lim\limits_{x\to-\infty} \dfrac{1}{f(x)} = \dfrac{1}{2}$ and $\lim\limits_{x\to\infty} g(x) = \lim\limits_{x\to\infty} \dfrac{1}{f(x)} = \dfrac{1}{-1} = -1$, so $y = \dfrac{1}{2}$ and $y = -1$ are horizontal asymptotes. $\lim\limits_{x\to7^-} \dfrac{1}{f(x)} = \dfrac{1}{\lim\limits_{x\to7^-} f(x)} = \infty$ and $\lim\limits_{x\to7^+} g(x) = \lim\limits_{x\to7^+} \dfrac{1}{f(x)} = \dfrac{1}{\lim\limits_{x\to7^+} f(x)} = -\infty$, so $x = 7$ is a vertical asymptote.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. $m = f(v) = \dfrac{m_0}{\sqrt{1 - v^2/c^2}}$. The m-intercept is $f(0) = m_0$. There are no v-intercepts. $\lim\limits_{v \to c^-} f(v) = \infty$, so $v = c$ is a VA.

$f'(v) = -\frac{1}{2} m_0 (1 - v^2/c^2)^{-3/2}(-2v/c^2) = \dfrac{m_0 v}{c^2(1 - v^2/c^2)^{3/2}} = \dfrac{m_0 v}{\dfrac{c^2(c^2 - v^2)^{3/2}}{c^3}} = \dfrac{m_0 c v}{(c^2 - v^2)^{3/2}} > 0$, so f is increasing on $(0, c)$. There are no local extreme values.

$$f''(v) = \frac{(c^2 - v^2)^{3/2}(m_0 c) - m_0 c v \cdot \frac{3}{2}(c^2 - v^2)^{1/2}(-2v)}{[(c^2 - v^2)^{3/2}]^2}$$

$$= \frac{m_0 c (c^2 - v^2)^{1/2}[(c^2 - v^2) + 3v^2]}{(c^2 - v^2)^3} = \frac{m_0 c (c^2 + 2v^2)}{(c^2 - v^2)^{5/2}} > 0,$$

so f is CU on $(0, c)$. There are no inflection points.

60. Let $a = m_0^2 c^4$ and $b = h^2 c^2$, so the equation can be written as $E = f(\lambda) = \sqrt{a + b/\lambda^2} = \sqrt{\dfrac{a\lambda^2 + b}{\lambda^2}} = \dfrac{\sqrt{a\lambda^2 + b}}{\lambda}$.

$\lim\limits_{\lambda \to 0^+} \dfrac{\sqrt{a\lambda^2 + b}}{\lambda} = \infty$, so $\lambda = 0$ is a VA.

$\lim\limits_{\lambda \to \infty} \dfrac{\sqrt{a\lambda^2 + b}}{\lambda} = \lim\limits_{\lambda \to \infty} \dfrac{\sqrt{a\lambda^2 + b}/\lambda}{\lambda/\lambda} = \lim\limits_{\lambda \to \infty} \dfrac{\sqrt{a + b/\lambda^2}}{1} = \sqrt{a}$, so $E = \sqrt{a} = m_0 c^2$ is a HA.

$$f'(\lambda) = \frac{\lambda \cdot \frac{1}{2}(a\lambda^2 + b)^{-1/2}(2a\lambda) - (a\lambda^2 + b)^{1/2}(1)}{\lambda^2} = \frac{(a\lambda^2 + b)^{-1/2}[a\lambda^2 - (a\lambda^2 + b)]}{\lambda^2} = \frac{-b}{\lambda^2 \sqrt{a\lambda^2 + b}} < 0,$$

so f is decreasing on $(0, \infty)$. Using the Reciprocal Rule,

$$f''(\lambda) = b \cdot \frac{\lambda^2 \cdot \frac{1}{2}(a\lambda^2 + b)^{-1/2}(2a\lambda) + (a\lambda^2 + b)^{1/2}(2\lambda)}{\left(\lambda^2 \sqrt{a\lambda^2 + b}\right)^2}$$

$$= \frac{b\lambda(a\lambda^2 + b)^{-1/2}[a\lambda^2 + 2(a\lambda^2 + b)]}{\left(\lambda^2 \sqrt{a\lambda^2 + b}\right)^2} = \frac{b(3a\lambda^2 + 2b)}{\lambda^3 (a\lambda^2 + b)^{3/2}} > 0,$$

so f is CU on $(0, \infty)$. There are no extrema or inflection points. The graph shows that as λ decreases, the energy increases and as λ increases, the energy decreases. For large wavelengths, the energy is very close to the energy at rest.

61. (a) $p(t) = \dfrac{1}{2} \;\Rightarrow\; \dfrac{1}{2} = \dfrac{1}{1 + ae^{-kt}} \;\Leftrightarrow\; 1 + ae^{-kt} = 2 \;\Leftrightarrow\; ae^{-kt} = 1 \;\Leftrightarrow\; e^{-kt} = \dfrac{1}{a} \;\Leftrightarrow$

$\ln e^{-kt} = \ln a^{-1} \;\Leftrightarrow\; -kt = -\ln a \;\Leftrightarrow\; t = \dfrac{\ln a}{k}$, which is when half the population will have heard the rumor.

(b) The rate of spread is given by $p'(t) = \dfrac{ake^{-kt}}{(1 + ae^{-kt})^2}$. To find the greatest rate of spread, we'll apply the First Derivative Test to $p'(t)$ [not $p(t)$].

$$[p'(t)]' = p''(t) = \frac{(1 + ae^{-kt})^2(-ak^2 e^{-kt}) - ake^{-kt} \cdot 2(1 + ae^{-kt})(-ake^{-kt})}{[(1 + ae^{-kt})^2]^2}$$

$$= \frac{(1 + ae^{-kt})(-ake^{-kt})[k(1 + ae^{-kt}) - 2ake^{-kt}]}{(1 + ae^{-kt})^4} = \frac{-ake^{-kt}(k)(1 - ae^{-kt})}{(1 + ae^{-kt})^3} = \frac{ak^2 e^{-kt}(ae^{-kt} - 1)}{(1 + ae^{-kt})^3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$p''(t) > 0 \quad \Leftrightarrow \quad ae^{-kt} > 1 \quad \Leftrightarrow \quad -kt > \ln a^{-1} \quad \Leftrightarrow \quad t < \dfrac{\ln a}{k}$, so $p'(t)$ is increasing for $t < \dfrac{\ln a}{k}$ and $p'(t)$ is decreasing for $t > \dfrac{\ln a}{k}$. Thus, $p'(t)$, the rate of spread of the rumor, is greatest at the same time, $\dfrac{\ln a}{k}$, as when half the population [by part (a)] has heard it.

(c) $p(0) = \dfrac{1}{1+a}$ and $\lim\limits_{t\to\infty} p(t) = 1$. The graph is shown with $a = 4$ and $k = \frac{1}{2}$.

62. $C(t) = K(e^{-at} - e^{-bt})$, where $K > 0$ and $b > a > 0$. $C(0) = K(1-1) = 0$ is the only intercept. $\lim\limits_{t\to\infty} C(t) = 0$, so $C = 0$ is a HA. $C'(t) = K(-ae^{-at} + be^{-bt}) > 0 \quad \Leftrightarrow \quad be^{-bt} > ae^{-at} \quad \Leftrightarrow \quad e^{at}e^{-bt} > \dfrac{a}{b} \quad \Leftrightarrow \quad e^{(a-b)t} > \dfrac{a}{b} \quad \Leftrightarrow$ $(a-b)t > \ln\dfrac{a}{b} \quad \Leftrightarrow \quad t > \dfrac{\ln(a/b)}{a-b}$ or $\dfrac{\ln(b/a)}{b-a}$ [call this value D]. C is increasing for $t < D$ and decreasing for $t > D$, so $C(D)$ is a local maximum [and absolute] value. $C''(t) = K(a^2e^{-at} - b^2e^{-bt}) > 0 \quad \Leftrightarrow \quad a^2e^{-at} > b^2e^{-bt} \quad \Leftrightarrow$ $e^{bt}e^{-at} > \dfrac{b^2}{a^2} \quad \Leftrightarrow \quad e^{(b-a)t} > \left(\dfrac{b}{a}\right)^2 \quad \Leftrightarrow \quad (b-a)t > \ln\left(\dfrac{b}{a}\right)^2 \quad \Leftrightarrow \quad t > \dfrac{2\ln(b/a)}{b-a} = 2D$, so C is CU on $(2D, \infty)$ and CD on $(0, 2D)$. The inflection point is $(2D, C(2D))$. For the graph shown, $K = 1$, $a = 1$, $b = 2$, $D = \ln 2$, $C(D) = \frac{1}{4}$, and $C(2D) = \frac{3}{16}$. The graph tells us that when the drug is injected into the bloodstream, its concentration rises rapidly to a maximum at time D, then falls, reaching its maximum rate of decrease at time $2D$, then continues to decrease more and more slowly, approaching 0 as $t \to \infty$.

63. $y = -\dfrac{W}{24EI}x^4 + \dfrac{WL}{12EI}x^3 - \dfrac{WL^2}{24EI}x^2 = -\dfrac{W}{24EI}x^2\left(x^2 - 2Lx + L^2\right)$

$= \dfrac{-W}{24EI}x^2(x-L)^2 = cx^2(x-L)^2$

where $c = -\dfrac{W}{24EI}$ is a negative constant and $0 \le x \le L$. We sketch

$f(x) = cx^2(x-L)^2$ for $c = -1$. $f(0) = f(L) = 0$.

$f'(x) = cx^2[2(x-L)] + (x-L)^2(2cx) = 2cx(x-L)[x + (x-L)] = 2cx(x-L)(2x-L)$. So for $0 < x < L$, $f'(x) > 0 \quad \Leftrightarrow \quad x(x-L)(2x-L) < 0$ [since $c < 0$] $\quad \Leftrightarrow \quad L/2 < x < L$ and $f'(x) < 0 \quad \Leftrightarrow \quad 0 < x < L/2$. Thus, f is increasing on $(L/2, L)$ and decreasing on $(0, L/2)$, and there is a local and absolute minimum at the point $(L/2, f(L/2)) = \left(L/2, cL^4/16\right)$. $f'(x) = 2c[x(x-L)(2x-L)] \quad \Rightarrow$ $f''(x) = 2c[1(x-L)(2x-L) + x(1)(2x-L) + x(x-L)(2)] = 2c(6x^2 - 6Lx + L^2) = 0 \quad \Leftrightarrow$ $x = \dfrac{6L \pm \sqrt{12L^2}}{12} = \frac{1}{2}L \pm \frac{\sqrt{3}}{6}L$, and these are the x-coordinates of the two inflection points.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. $F(x) = -\dfrac{k}{x^2} + \dfrac{k}{(x-2)^2}$, where $k > 0$ and $0 < x < 2$. For $0 < x < 2$, $x - 2 < 0$, so $F'(x) = \dfrac{2k}{x^3} - \dfrac{2k}{(x-2)^3} > 0$ and F is increasing. $\lim\limits_{x \to 0^+} F(x) = -\infty$ and $\lim\limits_{x \to 2^-} F(x) = \infty$, so $x = 0$ and $x = 2$ are vertical asymptotes. Notice that when the middle particle is at $x = 1$, the net force acting on it is 0. When $x > 1$, the net force is positive, meaning that it acts to the right. And if the particle approaches $x = 2$, the force on it rapidly becomes very large. When $x < 1$, the net force is negative, so it acts to the left. If the particle approaches 0, the force becomes very large to the left.

65. $y = \dfrac{x^2+1}{x+1}$. Long division gives us:

$$\begin{array}{r|l} & \quad x - 1 \\ \hline x+1 & x^2 \qquad + 1 \\ & x^2 + x \\ \hline & \quad -x + 1 \\ & \quad -x - 1 \\ \hline & \qquad\quad 2 \end{array}$$

Thus, $y = f(x) = \dfrac{x^2+1}{x+1} = x - 1 + \dfrac{2}{x+1}$ and $f(x) - (x-1) = \dfrac{2}{x+1} = \dfrac{\frac{2}{x}}{1 + \frac{1}{x}}$ [for $x \neq 0$] $\to 0$ as $x \to \pm\infty$.

So the line $y = x - 1$ is a slant asymptote (SA).

66. $y = \dfrac{4x^3 - 10x^2 - 11x + 1}{x^2 - 3x}$. Long division gives us:

$$\begin{array}{r|l} & \qquad\qquad 4x + 2 \\ \hline x^2 - 3x & 4x^3 - 10x^2 - 11x + 1 \\ & 4x^3 - 12x^2 \\ \hline & \qquad 2x^2 - 11x \\ & \qquad 2x^2 - \ \ 6x \\ \hline & \qquad\qquad -5x + 1 \end{array}$$

Thus, $y = f(x) = \dfrac{4x^3 - 10x^2 - 11x + 1}{x^2 - 3x} = 4x + 2 + \dfrac{-5x+1}{x^2 - 3x}$ and $f(x) - (4x+2) = \dfrac{-5x+1}{x^2-3x} = \dfrac{-\frac{5}{x} + \frac{1}{x^2}}{1 - \frac{3}{x}}$

[for $x \neq 0$] $\to \frac{0}{1} = 0$ as $x \to \pm\infty$. So the line $y = 4x + 2$ is a slant asymptote (SA).

67. $y = \dfrac{2x^3 - 5x^2 + 3x}{x^2 - x - 2}$. Long division gives us:

$$\begin{array}{r|l} & \qquad\qquad 2x - 3 \\ \hline x^2 - x - 2 & 2x^3 - 5x^2 + 3x \\ & 2x^3 - 2x^2 - 4x \\ \hline & \quad -3x^2 + 7x \\ & \quad -3x^2 + 3x + 6 \\ \hline & \qquad\qquad 4x - 6 \end{array}$$

Thus, $y = f(x) = \dfrac{2x^3 - 5x^2 + 3x}{x^2 - x - 2} = 2x - 3 + \dfrac{4x-6}{x^2 - x - 2}$ and $f(x) - (2x-3) = \dfrac{4x-6}{x^2-x-2} = \dfrac{\frac{4}{x} - \frac{6}{x^2}}{1 - \frac{1}{x} - \frac{1}{x^2}}$

[for $x \neq 0$] $\to \frac{0}{1} = 0$ as $x \to \pm\infty$. So the line $y = 2x - 3$ is a slant asymptote (SA).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

68. $y = \dfrac{-6x^4 + 2x^3 + 3}{2x^3 - x}$. Long division gives us:

$$\begin{array}{r|l} & \qquad\qquad\qquad\qquad -3x + 1 \\ \hline 2x^3 - x & -6x^4 + 2x^3 \qquad\qquad + 3 \\ & -6x^4 \qquad + 3x^2 \\ \hline & \qquad 2x^3 - 3x^2 \\ & \qquad 2x^3 \qquad - x \\ \hline & \qquad\qquad -3x^2 + x + 3 \end{array}$$

Thus, $y = f(x) = \dfrac{-6x^4 + 2x^3 + 3}{2x^3 - x} = -3x + 1 + \dfrac{-3x^2 + x + 3}{2x^3 - x}$ and

$f(x) - (-3x + 1) = \dfrac{-3x^2 + x + 3}{2x^3 - x} = \dfrac{-\dfrac{3}{x} + \dfrac{1}{x^2} + \dfrac{3}{x^3}}{2 - \dfrac{1}{x^2}}$ [for $x \neq 0$] $\to \frac{0}{2} = 0$ as $x \to \pm\infty$. So the line $y = -3x + 1$ is a slant asymptote (SA).

69. $y = f(x) = \dfrac{x^2}{x - 1} = x + 1 + \dfrac{1}{x - 1}$ **A.** $D = (-\infty, 1) \cup (1, \infty)$ **B.** x-intercept: $f(x) = 0 \iff x = 0$; y-intercept: $f(0) = 0$ **C.** No symmetry **D.** $\lim\limits_{x \to 1^-} f(x) = -\infty$ and $\lim\limits_{x \to 1^+} f(x) = \infty$, so $x = 1$ is a VA.

$\lim\limits_{x \to \pm\infty} [f(x) - (x + 1)] = \lim\limits_{x \to \pm\infty} \dfrac{1}{x - 1} = 0$, so the line $y = x + 1$ is a SA.

E. $f'(x) = 1 - \dfrac{1}{(x - 1)^2} = \dfrac{(x - 1)^2 - 1}{(x - 1)^2} = \dfrac{x^2 - 2x}{(x - 1)^2} = \dfrac{x(x - 2)}{(x - 1)^2} > 0$ for $x < 0$ or $x > 2$, so f is increasing on $(-\infty, 0)$ and $(2, \infty)$, and f is decreasing on $(0, 1)$ and $(1, 2)$. **F.** Local maximum value $f(0) = 0$, local minimum value $f(2) = 4$ **G.** $f''(x) = \dfrac{2}{(x - 1)^3} > 0$ for $x > 1$, so f is CU on $(1, \infty)$ and f is CD on $(-\infty, 1)$. No IP

H.

70. $y = f(x) = \dfrac{1 + 5x - 2x^2}{x - 2} = -2x + 1 + \dfrac{3}{x - 2}$ **A.** $D = (-\infty, 2) \cup (2, \infty)$ **B.** x-intercepts: $f(x) = 0 \iff$ $1 + 5x - 2x^2 = 0 \Rightarrow x = \dfrac{-5 \pm \sqrt{33}}{-4} \Rightarrow x \approx -0.19, 2.69$; y-intercept: $f(0) = -\frac{1}{2}$ **C.** No symmetry

D. $\lim\limits_{x \to 2^-} f(x) = -\infty$ and $\lim\limits_{x \to 2^+} f(x) = \infty$, so $x = 2$ is a VA. $\lim\limits_{x \to \pm\infty} [f(x) - (-2x + 1)] = \lim\limits_{x \to \pm\infty} \dfrac{3}{x - 2} = 0$, so $y = -2x + 1$ is a SA.

E. $f'(x) = -2 - \dfrac{3}{(x - 2)^2} = \dfrac{-2(x^2 - 4x + 4) - 3}{(x - 2)^2}$

$= \dfrac{-2x^2 + 8x - 11}{(x - 2)^2} < 0$

for $x \neq 2$, so f is decreasing on $(-\infty, 2)$ and $(2, \infty)$. **F.** No local extrema

G. $f''(x) = \dfrac{6}{(x - 2)^3} > 0$ for $x > 2$, so f is CU on $(2, \infty)$ and CD on $(-\infty, 2)$.

No IP

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

71. $y = f(x) = \dfrac{x^3+4}{x^2} = x + \dfrac{4}{x^2}$ **A.** $D = (-\infty, 0) \cup (0, \infty)$ **B.** x-intercept: $f(x) = 0 \iff x = -\sqrt[3]{4}$; no y-intercept

C. No symmetry **D.** $\lim\limits_{x\to 0} f(x) = \infty$, so $x = 0$ is a VA. $\lim\limits_{x\to\pm\infty}[f(x) - x] = \lim\limits_{x\to\pm\infty} \dfrac{4}{x^2} = 0$, so $y = x$ is a SA.

E. $f'(x) = 1 - \dfrac{8}{x^3} = \dfrac{x^3-8}{x^3} > 0$ for $x < 0$ or $x > 2$, so f is increasing on $(-\infty, 0)$ and $(2, \infty)$, and f is decreasing on $(0, 2)$. **F.** Local minimum value $f(2) = 3$, no local maximum value **G.** $f''(x) = \dfrac{24}{x^4} > 0$ for $x \neq 0$, so f is CU on $(-\infty, 0)$ and $(0, \infty)$. No IP

H.

72. $y = f(x) = \dfrac{x^3}{(x+1)^2} = x - 2 + \dfrac{3x+2}{(x+1)^2}$ **A.** $D = (-\infty, -1) \cup (-1, \infty)$ **B.** x-intercept: 0; y-intercept: $f(0) = 0$

C. No symmetry **D.** $\lim\limits_{x\to -1^-} f(x) = -\infty$ and $\lim\limits_{x\to -1^+} f(x) = -\infty$, so $x = -1$ is a VA.

$\lim\limits_{x\to\pm\infty}[f(x) - (x-2)] = \lim\limits_{x\to\pm\infty} \dfrac{3x+2}{(x+1)^2} = 0$, so $y = x - 2$ is a SA.

E. $f'(x) = \dfrac{(x+1)^2(3x^2) - x^3 \cdot 2(x+1)}{[(x+1)^2]^2} = \dfrac{x^2(x+1)[3(x+1) - 2x]}{(x+1)^4} = \dfrac{x^2(x+3)}{(x+1)^3} > 0 \iff x < -3$ or $x > -1$ $[x \neq 0]$, so f is increasing on $(-\infty, -3)$ and $(-1, \infty)$, and f is decreasing on $(-3, -1)$.

F. Local maximum value $f(-3) = -\frac{27}{4}$, no local minimum

G.
$$\begin{aligned} f''(x) &= \frac{(x+1)^3(3x^2+6x) - (x^3+3x^2)\cdot 3(x+1)^2}{[(x+1)^3]^2} \\ &= \frac{3x(x+1)^2[(x+1)(x+2) - (x^2+3x)]}{(x+1)^6} \\ &= \frac{3x(x^2+3x+2-x^2-3x)}{(x+1)^4} = \frac{6x}{(x+1)^4} > 0 \iff \end{aligned}$$
$x > 0$, so f is CU on $(0, \infty)$ and f is CD on $(-\infty, -1)$ and $(-1, 0)$. IP at $(0, 0)$

H.

73. $y = f(x) = 1 + \frac{1}{2}x + e^{-x}$ **A.** $D = \mathbb{R}$ **B.** y-intercept $= f(0) = 2$, no x-intercept [see part **F**] **C.** No symmetry

D. No VA or HA. $\lim\limits_{x\to\infty}\left[f(x) - \left(1 + \frac{1}{2}x\right)\right] = \lim\limits_{x\to\infty} e^{-x} = 0$, so $y = 1 + \frac{1}{2}x$ is a SA. **E.** $f'(x) = \frac{1}{2} - e^{-x} > 0 \iff \frac{1}{2} > e^{-x} \iff -x < \ln\frac{1}{2} \iff x > -\ln 2^{-1} \iff x > \ln 2$, so f is increasing on $(\ln 2, \infty)$ and decreasing on $(-\infty, \ln 2)$. **F.** Local and absolute minimum value

$$\begin{aligned} f(\ln 2) &= 1 + \tfrac{1}{2}\ln 2 + e^{-\ln 2} = 1 + \tfrac{1}{2}\ln 2 + (e^{\ln 2})^{-1} \\ &= 1 + \tfrac{1}{2}\ln 2 + \tfrac{1}{2} = \tfrac{3}{2} + \tfrac{1}{2}\ln 2 \approx 1.85, \end{aligned}$$

no local maximum value **G.** $f''(x) = e^{-x} > 0$ for all x, so f is CU on $(-\infty, \infty)$. No IP

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

74. $y = f(x) = 1 - x + e^{1+x/3}$ **A.** $D = \mathbb{R}$ **B.** y-intercept $= f(0) = 1 + e$, no x-intercept [see part **F**]

C. No symmetry **D.** No VA or HA $\lim\limits_{x\to-\infty}[f(x) - (1-x)] = \lim\limits_{x\to-\infty} e^{1+x/3} = 0$, so $y = 1 - x$ is a SA.

E. $f'(x) = -1 + \frac{1}{3}e^{1+x/3} > 0 \;\Leftrightarrow\; \frac{1}{3}e^{1+x/3} > 1 \;\Leftrightarrow\; e^{1+x/3} > 3 \;\Leftrightarrow\; 1 + \dfrac{x}{3} > \ln 3 \;\Leftrightarrow\; \dfrac{x}{3} > \ln 3 - 1 \;\Leftrightarrow$

$x > 3(\ln 3 - 1) \approx 0.3$, so f isincreasing on $(3\ln 3 - 3, \infty)$ and decreasing on $(-\infty, 3\ln 3 - 3)$. **F.** Local and absolute minimum value $f(3\ln 3 - 3) = 1 - (3\ln 3 - 3) + e^{1+\ln 3 - 1} = 4 - 3\ln 3 + 3 = 7 - 3\ln 3 \approx 3.7$, no local maximum value **G.** $f''(x) = \frac{1}{9}e^{1+x/3} > 0$ for all x, so f is CU on $(-\infty, \infty)$. No IP

H.

75. $y = f(x) = x - \tan^{-1} x$, $f'(x) = 1 - \dfrac{1}{1+x^2} = \dfrac{1+x^2-1}{1+x^2} = \dfrac{x^2}{1+x^2}$,

$f''(x) = \dfrac{(1+x^2)(2x) - x^2(2x)}{(1+x^2)^2} = \dfrac{2x(1+x^2-x^2)}{(1+x^2)^2} = \dfrac{2x}{(1+x^2)^2}$.

$\lim\limits_{x\to\infty}\left[f(x) - \left(x - \frac{\pi}{2}\right)\right] = \lim\limits_{x\to\infty}\left(\frac{\pi}{2} - \tan^{-1} x\right) = \frac{\pi}{2} - \frac{\pi}{2} = 0$, so $y = x - \frac{\pi}{2}$ is a SA.

Also, $\lim\limits_{x\to-\infty}\left[f(x) - \left(x + \frac{\pi}{2}\right)\right] = \lim\limits_{x\to-\infty}\left(-\frac{\pi}{2} - \tan^{-1} x\right) = -\frac{\pi}{2} - \left(-\frac{\pi}{2}\right) = 0$,

so $y = x + \frac{\pi}{2}$ is also a SA. $f'(x) \geq 0$ for all x, with equality $\Leftrightarrow$ $x = 0$, so f is increasing on $\mathbb{R}$. $f''(x)$ has the same sign as x, so f is CD on $(-\infty, 0)$ and CU on $(0, \infty)$. $f(-x) = -f(x)$, so f is an odd function; its graph is symmetric about the origin. f has no local extreme values. Its only IP is at $(0, 0)$.

76. $y = f(x) = \sqrt{x^2 + 4x} = \sqrt{x(x+4)}$. $x(x+4) \geq 0 \;\Leftrightarrow\; x \leq -4$ or $x \geq 0$, so $D = (-\infty, -4] \cup [0, \infty)$.

y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \;\Rightarrow\; x = -4, 0$.

$$\sqrt{x^2+4x} \mp (x+2) = \frac{\sqrt{x^2+4x} \mp (x+2)}{1} \cdot \frac{\sqrt{x^2+4x} \pm (x+2)}{\sqrt{x^2+4x} \pm (x+2)} = \frac{(x^2+4x) - (x^2+4x+4)}{\sqrt{x^2+4x} \pm (x+2)}$$
$$= \frac{-4}{\sqrt{x^2+4x} \pm (x+2)}$$

so $\lim\limits_{x\to\pm\infty}[f(x) \mp (x+2)] = 0$. Thus, the graph of f approaches the slant asymptote $y = x + 2$ as $x \to \infty$ and it approaches the slant asymptote $y = -(x+2)$ as $x \to -\infty$. $f'(x) = \dfrac{x+2}{\sqrt{x^2+4x}}$, so $f'(x) < 0$ for $x < -4$ and $f'(x) > 0$ for $x > 0$; that is, f is decreasing on $(-\infty, -4)$ and increasing on $(0, \infty)$. There are no local extreme values. $f'(x) = (x+2)(x^2+4x)^{-1/2} \;\Rightarrow$

$$\begin{aligned} f''(x) &= (x+2) \cdot \left(-\tfrac{1}{2}\right)(x^2+4x)^{-3/2} \cdot (2x+4) + (x^2+4x)^{-1/2} \\ &= (x^2+4x)^{-3/2}\left[-(x+2)^2 + (x^2+4x)\right] = -4(x^2+4x)^{-3/2} < 0 \text{ on } D \end{aligned}$$

so f is CD on $(-\infty, -4)$ and $(0, \infty)$. No IP

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

77. $\dfrac{x^2}{a^2} - \dfrac{y^2}{b^2} = 1 \quad \Rightarrow \quad y = \pm\dfrac{b}{a}\sqrt{x^2 - a^2}$. Now

$$\lim_{x\to\infty}\left[\frac{b}{a}\sqrt{x^2-a^2} - \frac{b}{a}x\right] = \frac{b}{a}\cdot\lim_{x\to\infty}\left(\sqrt{x^2-a^2}-x\right)\frac{\sqrt{x^2-a^2}+x}{\sqrt{x^2-a^2}+x} = \frac{b}{a}\cdot\lim_{x\to\infty}\frac{-a^2}{\sqrt{x^2-a^2}+x} = 0,$$

which shows that $y = \dfrac{b}{a}x$ is a slant asymptote. Similarly,

$$\lim_{x\to\infty}\left[-\frac{b}{a}\sqrt{x^2-a^2} - \left(-\frac{b}{a}x\right)\right] = -\frac{b}{a}\cdot\lim_{x\to\infty}\frac{-a^2}{\sqrt{x^2-a^2}+x} = 0, \text{ so } y = -\frac{b}{a}x \text{ is a slant asymptote.}$$

78. $f(x) - x^2 = \dfrac{x^3+1}{x} - x^2 = \dfrac{x^3+1-x^3}{x} = \dfrac{1}{x}$, and $\displaystyle\lim_{x\to\pm\infty}\frac{1}{x} = 0$. Therefore, $\displaystyle\lim_{x\to\pm\infty}[f(x)-x^2] = 0$,

and so the graph of f is asymptotic to that of $y = x^2$. For purposes of differentiation, we will use $f(x) = x^2 + 1/x$.

A. $D = \{x \mid x \neq 0\}$ **B.** No y-intercept; to find the x-intercept, we set $y = 0 \quad \Leftrightarrow \quad x = -1$.

C. No symmetry **D.** $\displaystyle\lim_{x\to 0^+}\frac{x^3+1}{x} = \infty$ and $\displaystyle\lim_{x\to 0^-}\frac{x^3+1}{x} = -\infty$, so $x = 0$ is a vertical asymptote. Also, the graph is asymptotic to the parabola $y = x^2$, as shown above. **E.** $f'(x) = 2x - 1/x^2 > 0 \quad \Leftrightarrow \quad x > \frac{1}{\sqrt[3]{2}}$, so f is increasing on $\left(\dfrac{1}{\sqrt[3]{2}}, \infty\right)$ and decreasing on $(-\infty, 0)$ and $\left(0, \dfrac{1}{\sqrt[3]{2}}\right)$.

F. Local minimum value $f\left(\dfrac{1}{\sqrt[3]{2}}\right) = \dfrac{3\sqrt[3]{3}}{2}$, no local maximum

G. $f''(x) = 2 + 2/x^3 > 0 \quad \Leftrightarrow \quad x < -1$ or $x > 0$, so f is CU on $(-\infty, -1)$ and $(0, \infty)$, and CD on $(-1, 0)$. IP at $(-1, 0)$

H.

79. $\displaystyle\lim_{x\to\pm\infty}\left[f(x) - x^3\right] = \lim_{x\to\pm\infty}\frac{x^4+1}{x} - \frac{x^4}{x} = \lim_{x\to\pm\infty}\frac{1}{x} = 0$, so the graph of f is asymptotic to that of $y = x^3$.

A. $D = \{x \mid x \neq 0\}$ **B.** No intercept **C.** f is symmetric about the origin. **D.** $\displaystyle\lim_{x\to 0^-}\left(x^3 + \frac{1}{x}\right) = -\infty$ and

$\displaystyle\lim_{x\to 0^+}\left(x^3 + \frac{1}{x}\right) = \infty$, so $x = 0$ is a vertical asymptote, and as shown above, the graph of f is asymptotic to that of $y = x^3$.

E. $f'(x) = 3x^2 - 1/x^2 > 0 \quad \Leftrightarrow \quad x^4 > \frac{1}{3} \quad \Leftrightarrow \quad |x| > \frac{1}{\sqrt[4]{3}}$, so f is increasing on $\left(-\infty, -\dfrac{1}{\sqrt[4]{3}}\right)$ and $\left(\dfrac{1}{\sqrt[4]{3}}, \infty\right)$ and decreasing on $\left(-\dfrac{1}{\sqrt[4]{3}}, 0\right)$ and $\left(0, \dfrac{1}{\sqrt[4]{3}}\right)$. **F.** Local maximum value $f\left(-\dfrac{1}{\sqrt[4]{3}}\right) = -4\cdot 3^{-5/4}$, local minimum value $f\left(\dfrac{1}{\sqrt[4]{3}}\right) = 4\cdot 3^{-5/4}$

G. $f''(x) = 6x + 2/x^3 > 0 \quad \Leftrightarrow \quad x > 0$, so f is CU on $(0, \infty)$ and CD on $(-\infty, 0)$. No IP

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

80. $f(x) = \sin x + e^{-x}$. $\lim\limits_{x\to\infty} [f(x) - \sin x] = \lim\limits_{x\to\infty} e^{-x} = 0$, so the graph of f is asymptotic to the graph of $\sin x$ as $x \to \infty$. $\lim\limits_{x\to-\infty} e^{-x} = \infty$, whereas $|\sin x| \le 1$, so for large negative x, the graph of f looks like the graph of e^{-x}.

4.6 Graphing with Calculus and Technology

1. $f(x) = x^5 - 5x^4 - x^3 + 28x^2 - 2x \;\Rightarrow\; f'(x) = 5x^4 - 20x^3 - 3x^2 + 56x - 2 \;\Rightarrow\; f''(x) = 20x^3 - 60x^2 - 6x + 56$. $f(x) = 0 \;\Leftrightarrow\; x = 0$ or $x \approx -2.09, 0.07$; $f'(x) = 0 \;\Leftrightarrow\; x \approx -1.50, 0.04, 2.62, 2.84$; $f''(x) = 0 \;\Leftrightarrow\; x \approx -0.89, 1.15, 2.74$.

From the graphs of f', we estimate that $f' < 0$ and that f is decreasing on $(-1.50, 0.04)$ and $(2.62, 2.84)$, and that $f' > 0$ and f is increasing on $(-\infty, -1.50)$, $(0.04, 2.62)$, and $(2.84, \infty)$ with local minimum values $f(0.04) \approx -0.04$ and $f(2.84) \approx 56.73$ and local maximum values $f(-1.50) \approx 36.47$ and $f(2.62) \approx 56.83$.

From the graph of f'', we estimate that $f'' > 0$ and that f is CU on $(-0.89, 1.15)$ and $(2.74, \infty)$, and that $f'' < 0$ and f is CD on $(-\infty, -0.89)$ and $(1.15, 2.74)$. There are inflection points at about $(-0.89, 20.90)$, $(1.15, 26.57)$, and $(2.74, 56.78)$.

2. $f(x) = -2x^6 + 5x^5 + 140x^3 - 110x^2 \;\Rightarrow\; f'(x) = -12x^5 + 25x^4 + 420x^2 - 220x \;\Rightarrow\; f''(x) = -60x^4 + 100x^3 + 840x - 220$. $f(x) = 0 \;\Leftrightarrow\; x = 0$ or $x \approx 0.77, 4.93$; $f'(x) = 0 \;\Leftrightarrow\; x = 0$ or

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x \approx 0.52, 3.99;\ f''(x) = 0 \quad \Leftrightarrow \quad x \approx 0.26, 3.05.$

From the graphs of f', we estimate that $f' > 0$ and that f is increasing on $(-\infty, 0)$ and $(0.52, 3.99)$, and that $f' < 0$ and that f is decreasing on $(0, 0.52)$ and $(3.99, \infty)$. f has local maximum values $f(0) = 0$ and $f(3.99) \approx 4128.20$, and f has local minimum value $f(0.52) \approx -9.91$. From the graph of f'', we estimate that $f'' > 0$ and f is CU on $(0.26, 3.05)$, and that $f'' < 0$ and f is CD on $(-\infty, 0.26)$ and $(3.05, \infty)$. There are inflection points at about $(0.26, -4.97)$ and $(3.05, 2649.46)$.

3. $f(x) = x^6 - 5x^5 + 25x^3 - 6x^2 - 48x \quad \Rightarrow$

$f'(x) = 6x^5 - 25x^4 + 75x^2 - 12x - 48 \quad \Rightarrow$

$f''(x) = 30x^4 - 100x^3 + 150x - 12.\quad f(x) = 0 \quad \Leftrightarrow \quad x = 0$ or $x \approx 3.20;$

$f'(x) = 0 \quad \Leftrightarrow \quad x \approx -1.31, -0.84, 1.06, 2.50, 2.75;\ f''(x) = 0 \quad \Leftrightarrow$

$x \approx -1.10, 0.08, 1.72, 2.64.$

From the graph of f', we estimate that f is decreasing on $(-\infty, -1.31)$, increasing on $(-1.31, -0.84)$, decreasing on $(-0.84, 1.06)$, increasing on $(1.06, 2.50)$, decreasing on $(2.50, 2.75)$, and increasing on $(2.75, \infty)$. f has local minimum values $f(-1.31) \approx 20.72$, $f(1.06) \approx -33.12$, and $f(2.75) \approx -11.33$. f has local maximum values $f(-0.84) \approx 23.71$ and $f(2.50) \approx -11.02$.

From the graph of f'', we estimate that f is CU on $(-\infty, -1.10)$, CD on $(-1.10, 0.08)$, CU on $(0.08, 1.72)$, CD on $(1.72, 2.64)$, and CU on $(2.64, \infty)$. There are inflection points at about $(-1.10, 22.09)$, $(0.08, -3.88)$, $(1.72, -22.53)$, and $(2.64, -11.18)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $f(x) = \dfrac{x^4 - x^3 - 8}{x^2 - x - 6} \quad \Rightarrow \quad f'(x) = \dfrac{2(x^5 - 2x^4 - 11x^3 + 9x^2 + 8x - 4)}{(x^2 - x - 6)^2} \quad \Rightarrow$

$f''(x) = \dfrac{2(x^6 - 3x^5 - 15x^4 + 41x^3 + 174x^2 - 84x - 56}{(x^2 - x - 6)^3}$. $\quad f(x) = 0 \quad \Leftrightarrow \quad x \approx -1.48$ or $x = 2$; $f'(x) = 0 \quad \Leftrightarrow$

$x \approx -2.74, -0.81, 0.41, 1.08, 4.06$; $f''(x) = 0 \quad \Leftrightarrow \quad x \approx -0.39, 0.79$. The VAs are $x = -2$ and $x = 3$.

From the graphs of f', we estimate that f is decreasing on $(-\infty, -2.74)$, increasing on $(-2.74, -2)$, increasing on $(-2, -0.81)$, decreasing on $(-0.81, 0.41)$, increasing on $(0.41, 1.08)$, decreasing on $(1.08, 3)$, decreasing on $(3, 4.06)$, and increasing on $(4.06, \infty)$. f has local minimum values $f(-2.74) \approx 16.23$, $f(0.41) \approx 1.29$, and $f(4.06) \approx 30.63$. f has local maximum values $f(-0.81) \approx 1.55$ and $f(1.08) \approx 1.34$.

From the graphs of f'', we estimate that f is CU on $(-\infty, -2)$, CD on $(-2, -0.39)$, CU on $(-0.39, 0.79)$, CD on $(0.79, 3)$, and CU on $(3, \infty)$. There are inflection points at about $(-0.39, 1.45)$ and $(0.79, 1.31)$.

5. $f(x) = \dfrac{x}{x^3 + x^2 + 1} \quad \Rightarrow \quad f'(x) = -\dfrac{2x^3 + x^2 - 1}{(x^3 + x^2 + 1)^2} \quad \Rightarrow \quad f''(x) = \dfrac{2x(3x^4 + 3x^3 + x^2 - 6x - 3)}{(x^3 + x^2 + 1)^3}$

From the graph of f, we see that there is a VA at $x \approx -1.47$. From the graph of f', we estimate that f is increasing on $(-\infty, -1.47)$, increasing on $(-1.47, 0.66)$, and decreasing on $(0.66, \infty)$, with local maximum value $f(0.66) \approx 0.38$.

From the graph of f'', we estimate that f is CU on $(-\infty, -1.47)$, CD on $(-1.47, -0.49)$, CU on $(-0.49, 0)$, CD on $(0, 1.10)$, and CU on $(1.10, \infty)$. There is an inflection point at $(0, 0)$ and at about $(-0.49, -0.44)$ and $(1.10, 0.31)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $f(x) = 6\sin x - x^2, -5 \le x \le 3 \quad \Rightarrow \quad f'(x) = 6\cos x - 2x \quad \Rightarrow \quad f''(x) = -6\sin x - 2$

From the graph of f', which has two negative zeros, we estimate that f is increasing on $(-5, -2.94)$, decreasing on $(-2.94, -2.66)$, increasing on $(-2.66, 1.17)$, and decreasing on $(1.17, 3)$, with local maximum values $f(-2.94) \approx -9.84$ and $f(1.17) \approx 4.16$, and local minimum value $f(-2.66) \approx -9.85$.

From the graph of f'', we estimate that f is CD on $(-5, -2.80)$, CU on $(-2.80, -0.34)$, and CD on $(-0.34, 3)$. There are inflection points at about $(-2.80, -9.85)$ and $(-0.34, -2.12)$.

7. $f(x) = 6\sin x + \cot x, -\pi \le x \le \pi \quad \Rightarrow \quad f'(x) = 6\cos x - \csc^2 x \quad \Rightarrow \quad f''(x) = -6\sin x + 2\csc^2 x \cot x$

From the graph of f, we see that there are VAs at $x = 0$ and $x = \pm\pi$. f is an odd function, so its graph is symmetric about the origin. From the graph of f', we estimate that f is decreasing on $(-\pi, -1.40)$, increasing on $(-1.40, -0.44)$, decreasing on $(-0.44, 0)$, decreasing on $(0, 0.44)$, increasing on $(0.44, 1.40)$, and decreasing on $(1.40, \pi)$, with local minimum values $f(-1.40) \approx -6.09$ and $f(0.44) \approx 4.68$, and local maximum values $f(-0.44) \approx -4.68$ and $f(1.40) \approx 6.09$.

From the graph of f'', we estimate that f is CU on $(-\pi, -0.77)$, CD on $(-0.77, 0)$, CU on $(0, 0.77)$, and CD on $(0.77, \pi)$. There are IPs at about $(-0.77, -5.22)$ and $(0.77, 5.22)$.

8. $f(x) = e^x - 0.186x^4 \quad \Rightarrow \quad f'(x) = e^x - 0.744x^3 \quad \Rightarrow \quad f''(x) = e^x - 2.232x^2$

From the graph of f', which has two positive zeros, we estimate that f is increasing on $(-\infty, 2.973)$, decreasing on $(2.973, 3.027)$, and increasing on $(3.027, \infty)$, with local maximum value $f(2.973) \approx 5.01958$ and local minimum value $f(3.027) \approx 5.01949$.

From the graph of f'', we estimate that f is CD on $(-\infty, -0.52)$, CU on $(-0.52, 1.25)$, CD on $(1.25, 3.00)$, and CU on $(3.00, \infty)$. There are inflection points at about $(-0.52, 0.58)$, $(1.25, 3.04)$ and $(3.00, 5.01954)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $f(x) = 1 + \dfrac{1}{x} + \dfrac{8}{x^2} + \dfrac{1}{x^3} \quad \Rightarrow \quad f'(x) = -\dfrac{1}{x^2} - \dfrac{16}{x^3} - \dfrac{3}{x^4} = -\dfrac{1}{x^4}(x^2 + 16x + 3) \quad \Rightarrow$

$f''(x) = \dfrac{2}{x^3} + \dfrac{48}{x^4} + \dfrac{12}{x^5} = \dfrac{2}{x^5}(x^2 + 24x + 6).$

From the graphs, it appears that f increases on $(-15.8, -0.2)$ and decreases on $(-\infty, -15.8)$, $(-0.2, 0)$, and $(0, \infty)$; that f has a local minimum value of $f(-15.8) \approx 0.97$ and a local maximum value of $f(-0.2) \approx 72$; that f is CD on $(-\infty, -24)$ and $(-0.25, 0)$ and is CU on $(-24, -0.25)$ and $(0, \infty)$; and that f has IPs at $(-24, 0.97)$ and $(-0.25, 60)$.

To find the exact values, note that $f' = 0 \quad \Rightarrow \quad x = \dfrac{-16 \pm \sqrt{256 - 12}}{2} = -8 \pm \sqrt{61} \quad [\approx -0.19 \text{ and } -15.81]$.

f' is positive (f is increasing) on $\left(-8 - \sqrt{61}, -8 + \sqrt{61}\right)$ and f' is negative (f is decreasing) on $\left(-\infty, -8 - \sqrt{61}\right)$, $\left(-8 + \sqrt{61}, 0\right)$, and $(0, \infty)$. $\quad f'' = 0 \quad \Rightarrow \quad x = \dfrac{-24 \pm \sqrt{576 - 24}}{2} = -12 \pm \sqrt{138} \quad [\approx -0.25 \text{ and } -23.75]$. $\quad f''$ is positive (f is CU) on $\left(-12 - \sqrt{138}, -12 + \sqrt{138}\right)$ and $(0, \infty)$ and f'' is negative (f is CD) on $\left(-\infty, -12 - \sqrt{138}\right)$ and $\left(-12 + \sqrt{138}, 0\right)$.

10. $f(x) = \dfrac{1}{x^8} - \dfrac{c}{x^4} \quad [c = 2 \times 10^8] \quad \Rightarrow$

$f'(x) = -\dfrac{8}{x^9} + \dfrac{4c}{x^5} = -\dfrac{4}{x^9}(2 - cx^4) \quad \Rightarrow$

$f''(x) = \dfrac{72}{x^{10}} - \dfrac{20c}{x^6} = \dfrac{4}{x^{10}}(18 - 5cx^4).$

From the graph, it appears that f increases on $(-0.01, 0)$ and $(0.01, \infty)$ and decreases on $(-\infty, -0.01)$ and $(0, 0.01)$; that f has a local minimum value of $f(\pm 0.01) = -10^{16}$; and that f is CU on $(-0.012, 0)$ and $(0, 0.012)$ and f is CD on $(-\infty, -0.012)$ and $(0.012, \infty)$.

To find the exact values, note that $f' = 0 \quad \Rightarrow \quad x^4 = \frac{2}{c} \quad \Rightarrow \quad x \pm \sqrt[4]{\frac{2}{c}} = \pm\frac{1}{100} \quad [c = 2 \times 10^8]$. $\quad f'$ is positive (f is increasing) on $(-0.01, 0)$ and $(0.01, \infty)$ and f' is negative (f is decreasing) on $(-\infty, -0.01)$ and $(0, 0.01)$.

$f'' = 0 \quad \Rightarrow \quad x^4 = \dfrac{18}{5c} \quad \Rightarrow \quad x = \pm\sqrt[4]{\dfrac{18}{5c}} = \pm\dfrac{1}{100}\sqrt[4]{1.8} \quad [\approx \pm 0.0116]$. $\quad f''$ is positive (f is CU) on $\left(-\frac{1}{100}\sqrt[4]{1.8}, 0\right)$ and $\left(0, \frac{1}{100}\sqrt[4]{1.8}\right)$ and f'' is negative (f is CD) on $\left(-\infty, -\frac{1}{100}\sqrt[4]{1.8}\right)$ and $\left(\frac{1}{100}\sqrt[4]{1.8}, \infty\right)$.

11. (a) $f(x) = x^2 \ln x$. The domain of f is $(0, \infty)$.

(b) $\displaystyle\lim_{x \to 0^+} x^2 \ln x = \lim_{x \to 0^+} \frac{\ln x}{1/x^2} \overset{\text{H}}{=} \lim_{x \to 0^+} \frac{1/x}{-2/x^3} = \lim_{x \to 0^+} \left(-\frac{x^2}{2}\right) = 0.$

There is a hole at $(0, 0)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) It appears that there is an IP at about $(0.2, -0.06)$ and a local minimum at $(0.6, -0.18)$. $f(x) = x^2 \ln x \quad \Rightarrow$ $f'(x) = x^2(1/x) + (\ln x)(2x) = x(2\ln x + 1) > 0 \quad \Leftrightarrow \quad \ln x > -\frac{1}{2} \quad \Leftrightarrow \quad x > e^{-1/2}$, so f is increasing on $\left(1/\sqrt{e}, \infty\right)$, decreasing on $\left(0, 1/\sqrt{e}\right)$. By the FDT, $f\left(1/\sqrt{e}\right) = -1/(2e)$ is a local minimum value. This point is approximately $(0.6065, -0.1839)$, which agrees with our estimate.

$f''(x) = x(2/x) + (2\ln x + 1) = 2\ln x + 3 > 0 \quad \Leftrightarrow \quad \ln x > -\frac{3}{2} \quad \Leftrightarrow \quad x > e^{-3/2}$, so f is CU on $(e^{-3/2}, \infty)$ and CD on $(0, e^{-3/2})$. IP is $(e^{-3/2}, -3/(2e^3)) \approx (0.2231, -0.0747)$.

12. (a) $f(x) = xe^{1/x}$. The domain of f is $(-\infty, 0) \cup (0, \infty)$.

(b) $\lim\limits_{x\to 0^+} xe^{1/x} = \lim\limits_{x\to 0^+} \dfrac{e^{1/x}}{1/x} \overset{\text{H}}{=} \lim\limits_{x\to 0^+} \dfrac{e^{1/x}(-1/x^2)}{-1/x^2} = \lim\limits_{x\to 0^+} e^{1/x} = \infty$,

so $x = 0$ is a VA.

Also $\lim\limits_{x\to 0^-} xe^{1/x} = 0$ since $1/x \to -\infty \quad \Rightarrow \quad e^{1/x} \to 0$.

(c) It appears that there is a local minimum at $(1, 2.7)$. There are no IP and f is CD on $(-\infty, 0)$ and CU on $(0, \infty)$.

$f(x) = xe^{1/x} \quad \Rightarrow \quad f'(x) = xe^{1/x}\left(-\dfrac{1}{x^2}\right) + e^{1/x} = e^{1/x}\left(1 - \dfrac{1}{x}\right) > 0 \quad \Leftrightarrow \quad \dfrac{1}{x} < 1 \quad \Leftrightarrow \quad x < 0$ or $x > 1$,

so f is increasing on $(-\infty, 0)$ and $(1, \infty)$, and decreasing on $(0, 1)$. By the FDT, $f(1) = e$ is a local minimum value, which agrees with our estimate.

$f''(x) = e^{1/x}(1/x^2) + (1 - 1/x)e^{1/x}(-1/x^2) = (e^{1/x}/x^2)(1 - 1 + 1/x) = e^{1/x}/x^3 > 0 \quad \Leftrightarrow \quad x > 0$, so f is CU on $(0, \infty)$ and CD on $(-\infty, 0)$. No IP

13.

$f(x) = \dfrac{(x+4)(x-3)^2}{x^4(x-1)}$ has VA at $x = 0$ and at $x = 1$ since $\lim\limits_{x\to 0} f(x) = -\infty$, $\lim\limits_{x\to 1^-} f(x) = -\infty$ and $\lim\limits_{x\to 1^+} f(x) = \infty$.

$$f(x) = \frac{\dfrac{x+4}{x} \cdot \dfrac{(x-3)^2}{x^2}}{\dfrac{x^4}{x^3} \cdot (x-1)} \quad \left[\begin{array}{c}\text{dividing numerator}\\ \text{and denominator by } x^3\end{array}\right] = \frac{(1+4/x)(1-3/x)^2}{x(x-1)} \to 0$$

as $x \to \pm\infty$, so f is asymptotic to the x-axis.

Since f is undefined at $x = 0$, it has no y-intercept. $f(x) = 0 \quad \Rightarrow \quad (x+4)(x-3)^2 = 0 \quad \Rightarrow \quad x = -4$ or $x = 3$, so f has x-intercepts -4 and 3. Note, however, that the graph of f is only tangent to the x-axis and does not cross it at $x = 3$, since f is positive as $x \to 3^-$ and as $x \to 3^+$.

From these graphs, it appears that f has three maximum values and one minimum value. The maximum values are

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

approximately $f(-5.6) = 0.0182$, $f(0.82) = -281.5$ and $f(5.2) = 0.0145$ and we know (since the graph is tangent to the x-axis at $x = 3$) that the minimum value is $f(3) = 0$.

14.

$f(x) = \dfrac{(2x+3)^2(x-2)^5}{x^3(x-5)^2}$ has VAs at $x = 0$ and $x = 5$ since $\lim\limits_{x\to 0^-} f(x) = \infty$, $\lim\limits_{x\to 0^+} f(x) = -\infty$, and $\lim\limits_{x\to 5} f(x) = \infty$. No HA since $\lim\limits_{x\to\pm\infty} f(x) = \infty$.

Since f is undefined at $x = 0$, it has no y-intercept.

$f(x) = 0 \quad \Leftrightarrow \quad (2x+3)^2\,(x-2)^5 = 0 \quad \Leftrightarrow \quad x = -\frac{3}{2}$ or $x = 2$, so f has x-intercepts at $-\frac{3}{2}$ and 2. Note, however, that the graph of f is only tangent to the x-axis and does not cross it at $x = -\frac{3}{2}$, since f is positive as $x \to \left(-\frac{3}{2}\right)^-$ and as $x \to \left(-\frac{3}{2}\right)^+$. There is a local minimum value of $f\left(-\frac{3}{2}\right) = 0$.

The only "mystery" feature is the local minimum to the right of the VA $x = 5$. From the graph, we see that $f(7.98) \approx 609$ is a local minimum value.

15. $f(x) = \dfrac{x^2(x+1)^3}{(x-2)^2(x-4)^4} \quad \Rightarrow \quad f'(x) = -\dfrac{x(x+1)^2(x^3+18x^2-44x-16)}{(x-2)^3(x-4)^5}$ [from CAS].

From the graphs of f', it seems that the critical points which indicate extrema occur at $x \approx -20$, -0.3, and 2.5, as estimated in Example 3. (There is another critical point at $x = -1$, but the sign of f' does not change there.) We differentiate again, obtaining $f''(x) = 2\dfrac{(x+1)(x^6+36x^5+6x^4-628x^3+684x^2+672x+64)}{(x-2)^4(x-4)^6}$.

From the graphs of f'', it appears that f is CU on $(-35.3, -5.0)$, $(-1, -0.5)$, $(-0.1, 2)$, $(2, 4)$, and $(4, \infty)$ and CD on $(-\infty, -35.3)$, $(-5.0, -1)$ and $(-0.5, -0.1)$. We check back on the graphs of f to find the y-coordinates of the inflection points, and find that these points are approximately $(-35.3, -0.015)$, $(-5.0, -0.005)$, $(-1, 0)$, $(-0.5, 0.00001)$, and $(-0.1, 0.0000066)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. From a CAS,
$$f'(x) = \frac{2(x-2)^4(2x+3)(2x^3-14x^2-10x-45)}{x^4(x-5)^3}$$
and
$$f''(x) = \frac{2(x-2)^3(4x^6-56x^5+216x^4+460x^3+805x^2+1710x+5400)}{x^5(x-5)^4}$$

From Exercise 14 and $f'(x)$ above, we know that the zeros of f' are -1.5, 2, and 7.98. From the graph of f', we conclude that f is decreasing on $(-\infty, -1.5)$, increasing on $(-1.5, 0)$ and $(0, 5)$, decreasing on $(5, 7.98)$, and increasing on $(7.98, \infty)$.

From $f''(x)$, we know that $x = 2$ is a zero, and the graph of f'' shows us that $x = 2$ is the only zero of f''. Thus, f is CU on $(-\infty, 0)$, CD on $(0, 2)$, CU on $(2, 5)$, and CU on $(5, \infty)$.

17. $f(x) = \dfrac{x^3+5x^2+1}{x^4+x^3-x^2+2}$. From a CAS, $f'(x) = \dfrac{-x(x^5+10x^4+6x^3+4x^2-3x-22)}{(x^4+x^3-x^2+2)^2}$ and
$$f''(x) = \frac{2(x^9+15x^8+18x^7+21x^6-9x^5-135x^4-76x^3+21x^2+6x+22)}{(x^4+x^3-x^2+2)^3}$$

The first graph of f shows that $y = 0$ is a HA. As $x \to \infty$, $f(x) \to 0$ through positive values. As $x \to -\infty$, it is not clear if $f(x) \to 0$ through positive or negative values. The second graph of f shows that f has an x-intercept near -5, and will have a local minimum and inflection point to the left of -5.

From the two graphs of f', we see that f' has four zeros. We conclude that f is decreasing on $(-\infty, -9.41)$, increasing on $(-9.41, -1.29)$, decreasing on $(-1.29, 0)$, increasing on $(0, 1.05)$, and decreasing on $(1.05, \infty)$. We have local minimum values $f(-9.41) \approx -0.056$ and $f(0) = 0.5$, and local maximum values $f(-1.29) \approx 7.49$ and $f(1.05) \approx 2.35$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

From the two graphs of f'', we see that f'' has five zeros. We conclude that f is CD on $(-\infty, -13.81)$, CU on $(-13.81, -1.55)$, CD on $(-1.55, -1.03)$, CU on $(-1.03, 0.60)$, CD on $(0.60, 1.48)$, and CU on $(1.48, \infty)$. There are five inflection points: $(-13.81, -0.05)$, $(-1.55, 5.64)$, $(-1.03, 5.39)$, $(0.60, 1.52)$, and $(1.48, 1.93)$.

18. $y = f(x) = \dfrac{x^{2/3}}{1 + x + x^4}$. From a CAS, $y' = -\dfrac{10x^4 + x - 2}{3x^{1/3}(x^4 + x + 1)^2}$ and $y'' = \dfrac{2(65x^8 - 14x^5 - 80x^4 + 2x^2 - 8x - 1)}{9x^{4/3}(x^4 + x + 1)^3}$

$f'(x)$ does not exist at $x = 0$ and $f'(x) = 0 \quad \Leftrightarrow \quad x \approx -0.72$ and 0.61, so f is increasing on $(-\infty, -0.72)$, decreasing on $(-0.72, 0)$, increasing on $(0, 0.61)$, and decreasing on $(0.61, \infty)$. There is a local maximum value of $f(-0.72) \approx 1.46$ and a local minimum value of $f(0.61) \approx 0.41$. $f''(x)$ does not exist at $x = 0$ and $f''(x) = 0 \quad \Leftrightarrow \quad x \approx -0.97, -0.46, -0.12$, and 1.11, so f is CU on $(-\infty, -0.97)$, CD on $(-0.97, -0.46)$, CU on $(-0.46, -0.12)$, CD on $(-0.12, 0)$, CD on $(0, 1.11)$, and CU on $(1.11, \infty)$. There are inflection points at $(-0.97, 1.08)$, $(-0.46, 1.01)$, $(-0.12, 0.28)$, and $(1.11, 0.29)$.

19. $y = f(x) = \sqrt{x + 5\sin x}$, $x \le 20$.

From a CAS, $y' = \dfrac{5\cos x + 1}{2\sqrt{x + 5\sin x}}$ and $y'' = -\dfrac{10\cos x + 25\sin^2 x + 10x\sin x + 26}{4(x + 5\sin x)^{3/2}}$.

We'll start with a graph of $g(x) = x + 5\sin x$. Note that $f(x) = \sqrt{g(x)}$ is only defined if $g(x) \ge 0$. $g(x) = 0 \quad \Leftrightarrow \quad x = 0$ or $x \approx -4.91, -4.10, 4.10$, and 4.91. Thus, the domain of f is $[-4.91, -4.10] \cup [0, 4.10] \cup [4.91, 20]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

From the expression for y', we see that $y' = 0 \quad \Leftrightarrow \quad 5\cos x + 1 = 0 \quad \Rightarrow \quad x_1 = \cos^{-1}\left(-\frac{1}{5}\right) \approx 1.77$ and $x_2 = 2\pi - x_1 \approx -4.51$ (not in the domain of f). The leftmost zero of f' is $x_1 - 2\pi \approx -4.51$. Moving to the right, the zeros of f' are x_1, $x_1 + 2\pi$, $x_2 + 2\pi$, $x_1 + 4\pi$, and $x_2 + 4\pi$. Thus, f is increasing on $(-4.91, -4.51)$, decreasing on $(-4.51, -4.10)$, increasing on $(0, 1.77)$, decreasing on $(1.77, 4.10)$, increasing on $(4.91, 8.06)$, decreasing on $(8.06, 10.79)$, increasing on $(10.79, 14.34)$, decreasing on $(14.34, 17.08)$, and increasing on $(17.08, 20)$. The local maximum values are $f(-4.51) \approx 0.62$, $f(1.77) \approx 2.58$, $f(8.06) \approx 3.60$, and $f(14.34) \approx 4.39$. The local minimum values are $f(10.79) \approx 2.43$ and $f(17.08) \approx 3.49$.

f is CD on $(-4.91, -4.10)$, $(0, 4.10)$, $(4.91, 9.60)$, CU on $(9.60, 12.25)$, CD on $(12.25, 15.81)$, CU on $(15.81, 18.65)$, and CD on $(18.65, 20)$. There are inflection points at $(9.60, 2.95)$, $(12.25, 3.27)$, $(15.81, 3.91)$, and $(18.65, 4.20)$.

20. $y = f(x) = x - \tan^{-1} x^2$. From a CAS, $y' = \dfrac{x^4 - 2x + 1}{x^4 + 1}$ and $y'' = \dfrac{2(3x^4 - 1)}{(x^4 + 1)^2}$. $\quad y' = 0 \quad \Leftrightarrow \quad x \approx 0.54$ or $x = 1$.

$y'' = 0 \quad \Leftrightarrow \quad x \approx \pm 0.76$.

From the graphs of f and f', we estimate that f is increasing on $(-\infty, 0.54)$, decreasing on $(0.54, 1)$, and increasing on $(1, \infty)$. $\quad f$ has local maximum value $f(0.54) \approx 0.26$ and local minimum value $f(1) \approx 0.21$.

From the graph of f'', we estimate that f is CU on $(-\infty, -0.76)$, CD on $(-0.76, 0.76)$, and CU on $(0.76, \infty)$. There are inflection points at about $(-0.76, -1.28)$ and $(0.76, 0.24)$.

21. $y = f(x) = \dfrac{1 - e^{1/x}}{1 + e^{1/x}}$. $\quad$ From a CAS, $y' = \dfrac{2e^{1/x}}{x^2(1 + e^{1/x})^2}$ and $y'' = \dfrac{-2e^{1/x}(1 - e^{1/x} + 2x + 2xe^{1/x})}{x^4(1 + e^{1/x})^3}$.

f is an odd function defined on $(-\infty, 0) \cup (0, \infty)$. Its graph has no x- or y-intercepts. Since $\lim\limits_{x \to \pm\infty} f(x) = 0$, the x-axis is a HA. $f'(x) > 0$ for $x \neq 0$, so f is increasing on $(-\infty, 0)$ and $(0, \infty)$. It has no local extreme values. $f''(x) = 0$ for $x \approx \pm 0.417$, so f is CU on $(-\infty, -0.417)$, CD on $(-0.417, 0)$, CU on $(0, 0.417)$, and CD on $(0.417, \infty)$. f has IPs at $(-0.417, 0.834)$ and $(0.417, -0.834)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $y = f(x) = \dfrac{3}{3 + 2\sin x}$. From a CAS, $y' = -\dfrac{6\cos x}{(3 + 2\sin x)^2}$ and $y'' = \dfrac{6(2\sin^2 x + 4\cos^2 x + 3\sin x)}{(3 + 2\sin x)^3}$. Since f is periodic with period 2π, we'll restrict our attention to the interval $[0, 2\pi)$. $y' = 0 \;\Leftrightarrow\; 6\cos x = 0 \;\Leftrightarrow\; x = \frac{\pi}{2}$ or $\frac{3\pi}{2}$. $y'' = 0 \;\Leftrightarrow\; x \approx 4.16$ or 5.27.

From the graphs of f and f', we conclude that f is decreasing on $\left(0, \frac{\pi}{2}\right)$, increasing on $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$, and decreasing on $\left(\frac{3\pi}{2}, 2\pi\right)$. f has local minimum value $f\left(\frac{\pi}{2}\right) = \frac{3}{5}$ and local maximum value $f\left(\frac{3\pi}{2}\right) = 3$.

From the graph of f'', we conclude that f is CU on $(0, 4.16)$, CD on $(4.16, 5.27)$, and CU on $(5.27, 2\pi)$. There are inflection points at about $(4.16, 2.31)$ and $(5.27, 2.31)$.

23. $f(x) = \dfrac{1 - \cos(x^4)}{x^8} \geq 0$. f is an even function, so its graph is symmetric with respect to the y-axis. The first graph shows that f levels off at $y = \frac{1}{2}$ for $|x| < 0.7$. It also shows that f then drops to the x-axis. Your graphing utility may show some severe oscillations near the origin, but there are none. See the discussion in Section 2.2 after Example 2, as well as "Lies My Calculator and Computer Told Me" on the website.

The second graph indicates that as $|x|$ increases, f has progressively smaller humps.

24. $f(x) = e^x + \ln|x - 4|$. The first graph shows the big picture of f but conceals hidden behavior.

The second graph shows that for large negative values of x, f looks like $g(x) = \ln|x|$. It also shows a minimum value and a point of inflection.

The third graph hints at the vertical asymptote that we know exists at $x = 4$ because $\lim\limits_{x \to 4} (e^x + \ln|x - 4|) = -\infty$.

A graphing calculator is unable to show much of the dip of the curve toward the vertical asymptote because of limited resolution. A computer can show more if we restrict ourselves to a narrow interval around $x = 4$. See the solution to Exercise 2.2.46 for a hand-drawn graph of this function.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. (a) $f(x) = x^{1/x}$

(b) Recall that $a^b = e^{b \ln a}$. $\lim\limits_{x \to 0^+} x^{1/x} = \lim\limits_{x \to 0^+} e^{(1/x) \ln x}$. As $x \to 0^+$, $\dfrac{\ln x}{x} \to -\infty$, so $x^{1/x} = e^{(1/x) \ln x} \to 0$. This indicates that there is a hole at $(0, 0)$. As $x \to \infty$, we have the indeterminate form ∞^0. $\lim\limits_{x \to \infty} x^{1/x} = \lim\limits_{x \to \infty} e^{(1/x) \ln x}$, but $\lim\limits_{x \to \infty} \dfrac{\ln x}{x} \overset{\text{H}}{=} \lim\limits_{x \to \infty} \dfrac{1/x}{1} = 0$, so $\lim\limits_{x \to \infty} x^{1/x} = e^0 = 1$. This indicates that $y = 1$ is a HA.

(c) Estimated maximum: $(2.72, 1.45)$. No estimated minimum. We use logarithmic differentiation to find any critical numbers. $y = x^{1/x} \;\Rightarrow\; \ln y = \dfrac{1}{x} \ln x \;\Rightarrow\; \dfrac{y'}{y} = \dfrac{1}{x} \cdot \dfrac{1}{x} + (\ln x)\left(-\dfrac{1}{x^2}\right) \;\Rightarrow\; y' = x^{1/x}\left(\dfrac{1 - \ln x}{x^2}\right) = 0 \;\Rightarrow$ $\ln x = 1 \;\Rightarrow\; x = e$. For $0 < x < e$, $y' > 0$ and for $x > e$, $y' < 0$, so $f(e) = e^{1/e}$ is a local maximum value. This point is approximately $(2.7183, 1.4447)$, which agrees with our estimate.

(d)

From the graph, we see that $f''(x) = 0$ at $x \approx 0.58$ and $x \approx 4.37$. Since f'' changes sign at these values, they are x-coordinates of inflection points.

26. (a) $f(x) = (\sin x)^{\sin x}$ is continuous where $\sin x > 0$, that is, on intervals of the form $(2n\pi, (2n+1)\pi)$, so we have graphed f on $(0, \pi)$.

(b) $y = (\sin x)^{\sin x} \;\Rightarrow\; \ln y = \sin x \ln \sin x$, so

$$\lim_{x \to 0^+} \ln y = \lim_{x \to 0^+} \sin x \ln \sin x = \lim_{x \to 0^+} \frac{\ln \sin x}{\csc x} \overset{\text{H}}{=} \lim_{x \to 0^+} \frac{\cot x}{-\csc x \cot x}$$
$$= \lim_{x \to 0^+} (-\sin x) = 0 \quad \Rightarrow \quad \lim_{x \to 0^+} y = e^0 = 1.$$

(c) It appears that we have a local maximum at $(1.57, 1)$ and local minima at $(0.38, 0.69)$ and $(2.76, 0.69)$.

$y = (\sin x)^{\sin x} \;\Rightarrow\; \ln y = \sin x \ln \sin x \;\Rightarrow\; \dfrac{y'}{y} = (\sin x)\left(\dfrac{\cos x}{\sin x}\right) + (\ln \sin x) \cos x = \cos x\,(1 + \ln \sin x) \;\Rightarrow$ $y' = (\sin x)^{\sin x}(\cos x)(1 + \ln \sin x)$. $y' = 0 \;\Rightarrow\; \cos x = 0$ or $\ln \sin x = -1 \;\Rightarrow\; x_2 = \frac{\pi}{2}$ or $\sin x = e^{-1}$. On $(0, \pi)$, $\sin x = e^{-1} \;\Rightarrow\; x_1 = \sin^{-1}(e^{-1})$ and $x_3 = \pi - \sin^{-1}(e^{-1})$. Approximating these points gives us $(x_1, f(x_1)) \approx (0.3767, 0.6922)$, $(x_2, f(x_2)) \approx (1.5708, 1)$, and $(x_3, f(x_3)) \approx (2.7649, 0.6922)$. The approximations confirm our estimates.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d)

From the graph, we see that $f''(x) = 0$ at $x \approx 0.94$ and $x \approx 2.20$. Since f'' changes sign at these values, they are x-coordinates of inflection points.

27.

From the graph of $f(x) = \sin(x + \sin 3x)$ in the viewing rectangle $[0, \pi]$ by $[-1.2, 1.2]$, it looks like f has two maxima and two minima. If we calculate and graph $f'(x) = [\cos(x + \sin 3x)]\,(1 + 3\cos 3x)$ on $[0, 2\pi]$, we see that the graph of f' appears to be almost tangent to the x-axis at about $x = 0.7$. The graph of

$$f'' = -\,[\sin(x + \sin 3x)]\,(1 + 3\cos 3x)^2 + \cos(x + \sin 3x)(-9\sin 3x)$$

is even more interesting near this x-value: it seems to just touch the x-axis.

If we zoom in on this place on the graph of f'', we see that f'' actually does cross the axis twice near $x = 0.65$, indicating a change in concavity for a very short interval. If we look at the graph of f' on the same interval, we see that it changes sign three times near $x = 0.65$, indicating that what we had thought was a broad extremum at about $x = 0.7$ actually consists of three extrema (two maxima and a minimum). These maximum values are roughly $f(0.59) = 1$ and $f(0.68) = 1$, and the minimum value is roughly $f(0.64) = 0.99996$. There are also a maximum value of about $f(1.96) = 1$ and minimum values of about $f(1.46) = 0.49$ and $f(2.73) = -0.51$. The points of inflection on $(0, \pi)$ are about $(0.61, 0.99998)$, $(0.66, 0.99998)$, $(1.17, 0.72)$, $(1.75, 0.77)$, and $(2.28, 0.34)$. On $(\pi, 2\pi)$, they are about $(4.01, -0.34)$, $(4.54, -0.77)$, $(5.11, -0.72)$, $(5.62, -0.99998)$, and $(5.67, -0.99998)$. There are also IP at $(0, 0)$ and $(\pi, 0)$. Note that the function is odd and periodic with period 2π, and it is also rotationally symmetric about all points of the form $((2n + 1)\pi, 0)$, n an integer.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $f(x) = x^3 + cx = x(x^2 + c) \quad \Rightarrow \quad f'(x) = 3x^2 + c \quad \Rightarrow \quad f''(x) = 6x$

$c = -6$ $\qquad$ $c = 0$ $\qquad$ $c = 6$

x-intercepts: When $c \geq 0$, 0 is the only x-intercept. When $c < 0$, the x-intercepts are 0 and $\pm\sqrt{-c}$.

y-intercept $= f(0) = 0$. f is odd, so the graph is symmetric with respect to the origin. $f''(x) < 0$ for $x < 0$ and $f''(x) > 0$ for $x > 0$, so f is CD on $(-\infty, 0)$ and CU on $(0, \infty)$. The origin is the only inflection point.

If $c > 0$, then $f'(x) > 0$ for all x, so f is increasing and has no local maximum or minimum.

If $c = 0$, then $f'(x) \geq 0$ with equality at $x = 0$, so again f is increasing and has no local maximum or minimum.

If $c < 0$, then $f'(x) = 3[x^2 - (-c/3)] = 3\left(x + \sqrt{-c/3}\right)\left(x - \sqrt{-c/3}\right)$, so $f'(x) > 0$ on $\left(-\infty, -\sqrt{-c/3}\right)$ and $\left(\sqrt{-c/3}, \infty\right)$; $f'(x) < 0$ on $\left(-\sqrt{-c/3}, \sqrt{-c/3}\right)$. It follows that $f\left(-\sqrt{-c/3}\right) = -\frac{2}{3}c\sqrt{-c/3}$ is a local maximum value and $f\left(\sqrt{-c/3}\right) = \frac{2}{3}c\sqrt{-c/3}$ is a local minimum value. As c decreases (toward more negative values), the local maximum and minimum move further apart.

There is no absolute maximum or minimum value. The only transitional value of c corresponding to a change in character of the graph is $c = 0$.

29. $f(x) = x^2 + 6x + c/x \quad \Rightarrow \quad f'(x) = 2x + 6 - c/x^2 \quad \Rightarrow \quad f''(x) = 2 + 2c/x^3$

$\boldsymbol{c = 0}$: The graph is the parabola $y = x^2 + 6x$, which has x-intercepts -6 and 0, vertex $(-3, -9)$, and opens upward.

$\boldsymbol{c \neq 0}$: The parabola $y = x^2 + 6x$ is an asymptote that the graph of f approaches as $x \to \pm\infty$. The y-axis is a vertical asymptote.

$\boldsymbol{c < 0}$**:** The x-intercepts are found by solving $f(x) = 0 \quad \Leftrightarrow \quad x^3 + 6x^2 + c = g(x) = 0$. Now $g'(x) = 0 \quad \Leftrightarrow \quad x = -4$ or 0, and g (not f) has a local maximum at $x = -4$. $\quad g(-4) = 32 + c$, so if $c < -32$, the maximum is negative and there are no negative x-intercepts; if $c = -32$, the maximum is 0 and there is one negative x-intercept; if $-32 < c < 0$, the maximum is positive and there are two negative x-intercepts. In all cases, there is one positive x-intercept.

As $c \to 0^-$, the local minimum point moves down and right, approaching $(-3, -9)$. [Note that since $f'(x) = \dfrac{2x^3 + 6x^2 - c}{x^2}$, Descartes' Rule of Signs implies that f' has no positive solutions and one negative solution when $c < 0$. $\quad f''(x) = \dfrac{2(x^3 + c)}{x^3} > 0$ at that negative solution, so that critical point yields a local minimum value. This tells us that there are no local maximums when $c < 0$.] $\quad f'(x) > 0$ for $x > 0$, so f is increasing on $(0, \infty)$. From

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$f''(x) = \dfrac{2(x^3+c)}{x^3}$, we see that f has an inflection point at $\left(\sqrt[3]{-c}, 6\sqrt[3]{-c}\right)$. This inflection point moves down and left, approaching the origin as $c \to 0^-$.

f is CU on $(-\infty, 0)$, CD on $\left(0, \sqrt[3]{-c}\right)$, and CU on $\left(\sqrt[3]{-c}, \infty\right)$.

$\boldsymbol{c > 0}$**:** The inflection point $\left(\sqrt[3]{-c}, 6\sqrt[3]{-c}\right)$ is now in the third quadrant and moves up and right, approaching the origin as $c \to 0^+$. f is CU on $\left(-\infty, \sqrt[3]{-c}\right)$, CD on $\left(\sqrt[3]{-c}, 0\right)$, and CU on $(0, \infty)$. f has a local minimum point in the first quadrant. It moves down and left, approaching the origin as $c \to 0^+$. $f'(x) = 0 \;\Leftrightarrow\; 2x^3 + 6x^2 - c = h(x) = 0$. Now $h'(x) = 0 \;\Leftrightarrow\; x = -2$ or 0, and h (*not* f) has a local maximum at $x = -2$. $h(-2) = 8 - c$, so $c = 8$ makes $h(x) = 0$, and hence, $f'(x) = 0$. When $c > 8$, $f'(x) < 0$ and f is decreasing on $(-\infty, 0)$. For $0 < c < 8$, there is a local minimum that moves toward $(-3, -9)$ and a local maximum that moves toward the origin as c decreases.

$\boldsymbol{c \le 0}$

$\boldsymbol{c \ge 0}$

30. With $c = 0$ in $y = f(x) = x\sqrt{c^2 - x^2}$, the graph of f is just the point $(0, 0)$. Since $(-c)^2 = c^2$, we only consider $c > 0$. Since $f(-x) = -f(x)$, the graph is symmetric about the origin. The domain of f is found by solving $c^2 - x^2 \ge 0 \;\Leftrightarrow\; x^2 \le c^2 \;\Leftrightarrow\; |x| \le c$, which gives us $[-c, c]$.

$$f'(x) = x \cdot \tfrac{1}{2}(c^2 - x^2)^{-1/2}(-2x) + (c^2 - x^2)^{1/2}(1) = (c^2 - x^2)^{-1/2}[-x^2 + (c^2 - x^2)] = \frac{c^2 - 2x^2}{\sqrt{c^2 - x^2}}.$$

$f'(x) > 0 \;\Leftrightarrow\; c^2 - 2x^2 > 0 \;\Leftrightarrow\; x^2 < c^2/2 \;\Leftrightarrow\; |x| < c/\sqrt{2}$, so f is increasing on $\left(-c/\sqrt{2}, c/\sqrt{2}\right)$ and decreasing on $\left(-c, -c/\sqrt{2}\right)$ and $\left(c/\sqrt{2}, c\right)$. There is a local minimum value of $f\left(-c/\sqrt{2}\right) = \left(-c/\sqrt{2}\right)\sqrt{c^2 - c^2/2} = \left(-c/\sqrt{2}\right)\left(c/\sqrt{2}\right) = -c^2/2$ and a local maximum value of $f\left(c/\sqrt{2}\right) = c^2/2$.

$$f''(x) = \frac{(c^2 - x^2)^{1/2}(-4x) - (c^2 - 2x^2)\frac{1}{2}\left(c^2 - x^2\right)^{-1/2}(-2x)}{[(c^2 - x^2)^{1/2}]^2}$$

$$= \frac{x(c^2 - x^2)^{-1/2}[(c^2 - x^2)(-4) + (c^2 - 2x^2)]}{(c^2 - x^2)^1} = \frac{2x\left(2x^2 - 3c^2\right)}{(c^2 - x^2)^{3/2}},$$

so $f''(x) = 0 \;\Leftrightarrow\; x = 0$ or $x = \pm\sqrt{\frac{3}{2}}\,c$, but only 0 is in the domain of f.

$f''(x) < 0$ for $0 < x < c$ and $f''(x) > 0$ for $-c < x < 0$, so f is CD on $(0, c)$ and CU on $(-c, 0)$. There is an IP at $(0, 0)$. So as $|c|$ gets larger, the maximum and minimum values increase in magnitude. The value of c does not affect the concavity of f.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. $f(x) = e^x + ce^{-x}$. $f = 0 \;\Rightarrow\; ce^{-x} = -e^x \;\Rightarrow\; c = -e^{2x} \;\Rightarrow\; 2x = \ln(-c) \;\Rightarrow\; x = \frac{1}{2}\ln(-c)$.

$f'(x) = e^x - ce^{-x}$. $f' = 0 \;\Rightarrow\; ce^{-x} = e^x \;\Rightarrow\; c = e^{2x} \;\Rightarrow\; 2x = \ln c \;\Rightarrow\; x = \frac{1}{2}\ln c$.

$f''(x) = e^x + ce^{-x} = f(x)$.

The only transitional value of c is 0. As c increases from $-\infty$ to 0, $\frac{1}{2}\ln(-c)$ is both the the x-intercept and inflection point, and this decreases from ∞ to $-\infty$. Also $f' > 0$, so f is increasing. When $c = 0$, $f(x) = f'(x) = f''(x) = e^x$, f is positive, increasing, and concave upward. As c increases from 0 to ∞, the absolute minimum occurs at $x = \frac{1}{2}\ln c$, which increases from $-\infty$ to ∞. Also, $f = f'' > 0$, so f is positive and concave upward. The value of the y-intercept is $f(0) = 1 + c$, and this increases as c increases from $-\infty$ to ∞.

Note: The minimum point $\left(\frac{1}{2}\ln c, 2\sqrt{c}\right)$ can be parameterized by $x = \frac{1}{2}\ln c$, $y = 2\sqrt{c}$, and after eliminating the parameter c, we see that the minimum point lies on the graph of $y = 2e^x$.

32. We see that if $c \le 0$, $f(x) = \ln(x^2 + c)$ is only defined for $x^2 > -c \;\Rightarrow\; |x| > \sqrt{-c}$, and $\lim\limits_{x\to\sqrt{-c}^+} f(x) = \lim\limits_{x\to-\sqrt{-c}^-} f(x) = -\infty$, since $\ln y \to -\infty$ as $y \to 0$. Thus, for $c < 0$, there are vertical asymptotes at $x = \pm\sqrt{c}$, and as c decreases (that is, $|c|$ increases), the asymptotes get further apart. For $c = 0$, $\lim\limits_{x\to 0} f(x) = -\infty$, so there is a vertical asymptote at $x = 0$. If $c > 0$, there are no asymptotes. To find the extrema and inflection points, we differentiate: $f(x) = \ln(x^2 + c) \;\Rightarrow\; f'(x) = \dfrac{1}{x^2 + c}(2x)$, so by the First Derivative Test there is a local and absolute minimum at $x = 0$. Differentiating again, we get $f''(x) = \dfrac{1}{x^2 + c}(2) + 2x\left[-(x^2 + c)^{-2}(2x)\right] = \dfrac{2(c - x^2)}{(x^2 + c)^2}$.

Now if $c \le 0$, f'' is always negative, so f is concave down on both of the intervals on which it is defined. If $c > 0$, then f'' changes sign when $c = x^2 \;\Leftrightarrow\; x = \pm\sqrt{c}$. So for $c > 0$ there are inflection points at $x = \pm\sqrt{c}$, and as c increases, the inflection points get further apart.

33. Note that $c = 0$ is a transitional value at which the graph consists of the x-axis. Also, we can see that if we substitute $-c$ for c, the function $f(x) = \dfrac{cx}{1 + c^2x^2}$ will be reflected in the x-axis, so we investigate only positive values of c (except $c = -1$, as a demonstration of this reflective property). Also, f is an odd function. $\lim\limits_{x\to\pm\infty} f(x) = 0$, so $y = 0$ is a horizontal asymptote for all c. We calculate $f'(x) = \dfrac{(1 + c^2x^2)c - cx(2c^2x)}{(1 + c^2x^2)^2} = -\dfrac{c(c^2x^2 - 1)}{(1 + c^2x^2)^2}$. $f'(x) = 0 \;\Leftrightarrow\; c^2x^2 - 1 = 0 \;\Leftrightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x = \pm 1/c$. So there is an absolute maximum value of $f(1/c) = \frac{1}{2}$ and an absolute minimum value of $f(-1/c) = -\frac{1}{2}$. These extrema have the same value regardless of c, but the maximum points move closer to the y-axis as c increases.

$$f''(x) = \frac{(-2c^3x)(1+c^2x^2)^2 - (-c^3x^2+c)[2(1+c^2x^2)(2c^2x)]}{(1+c^2x^2)^4}$$
$$= \frac{(-2c^3x)(1+c^2x^2) + (c^3x^2-c)(4c^2x)}{(1+c^2x^2)^3} = \frac{2c^3x(c^2x^2-3)}{(1+c^2x^2)^3}$$

$f''(x) = 0 \quad \Leftrightarrow \quad x = 0$ or $\pm\sqrt{3}/c$, so there are inflection points at $(0, 0)$ and at $\left(\pm\sqrt{3}/c, \pm\sqrt{3}/4\right)$. Again, the y-coordinate of the inflection points does not depend on c, but as c increases, both inflection points approach the y-axis.

34. $f(x) = \dfrac{\sin x}{c + \cos x} \quad \Rightarrow \quad f'(x) = \dfrac{1 + c\cos x}{\cos^2 x + 2c\cos x + c^2} \quad \Rightarrow \quad f''(x) = \dfrac{\sin x(c\cos x - c^2 + 2)}{\cos^3 x + 3c\cos^2 x + 3c^2\cos x + c^3}$. Notice that f is an odd function and has period 2π. We will graph f for $0 \le x \le 2\pi$.

$|\mathbf{c}| \le \mathbf{1}$**:** See the first figure. f has VAs when the denominator is zero, that is, at $x = \cos^{-1}(-c)$ and $x = 2\pi - \cos^{-1}(-c)$. So for $c = -1$, there are VAs at $x = 0$ and $x = 2\pi$, and as c increases, they move closer to $x = \pi$, which is the single VA when $c = 1$. Note that if $c = 0$, then $f(x) = \tan x$. There are no extreme points (on the entire domain) and inflection points occur at multiples of π.

$\mathbf{c} > \mathbf{1}$**:** See the second figure. $f'(x) = 0 \quad \Leftrightarrow \quad x = \cos^{-1}\left(\dfrac{-1}{c}\right)$ or $x = 2\pi - \cos^{-1}\left(\dfrac{-1}{c}\right)$. The VA disappears and there is now a local maximum and a local minimum. As $c \to 1^+$, the coordinates of the local maximum approach π and ∞, and the coordinates of the local minimum approach π and $-\infty$.

As $c \to \infty$, the graph of f looks like a graph of $y = \sin x$ that is vertically compressed, and the local maximum and local minimum approach $\left(\frac{\pi}{2}, 0\right)$ and $\left(\frac{3\pi}{2}, 0\right)$, respectively.

$f''(x) = 0 \quad \Leftrightarrow \quad \sin x = 0$ (IPs at $x = n\pi$) or $c\cos x - c^2 + 2 = 0$. The second condition is true if $\cos x = \dfrac{c^2 - 2}{c}$ $[c \ne 0]$. The last equation has two solutions if $-1 < \dfrac{c^2-2}{c} < 1 \quad \Rightarrow \quad -c < c^2 - 2 < c \quad \Rightarrow \quad -c < c^2 - 2$ and $c^2 - 2 < c \quad \Rightarrow \quad c^2 + c - 2 > 0$ and $c^2 - c - 2 < 0 \quad \Rightarrow \quad (c+2)(c-1) > 0$ and $(c-2)(c+1) < 0 \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$c-1>0$ [since $c>1$] and $c-2<0 \quad \Rightarrow \quad c>1$ and $c<2$. Thus, for $1<c<2$, we have 2 nontrivial IPs at $x=\cos^{-1}\left(\dfrac{c^2-2}{c}\right)$ and $x=2\pi-\cos^{-1}\left(\dfrac{c^2-2}{c}\right)$.

c < −1: See the third figure. The VAs for $c=-1$ at $x=0$ and $x=2\pi$ in the first figure disappear and we now have a local minimum and a local maximum. As $c \to -1^+$, the coordinates of the local minimum approach 0 and $-\infty$, and the coordinates of the local maximum approach 2π and ∞. As $c \to -\infty$, the graph of f looks like a graph of $y=\sin x$ that is vertically compressed, and the local minimum and local maximum approach $\left(\frac{\pi}{2},0\right)$ and $\left(\frac{3\pi}{2},0\right)$, respectively. As above, we have two nontrivial IPs for $-2<c<-1$.

35. $f(x)=cx+\sin x \quad \Rightarrow \quad f'(x)=c+\cos x \quad \Rightarrow \quad f''(x)=-\sin x$

$f(-x)=-f(x)$, so f is an odd function and its graph is symmetric with respect to the origin.

$f(x)=0 \quad \Leftrightarrow \quad \sin x=-cx$, so 0 is always an x-intercept.

$f'(x)=0 \quad \Leftrightarrow \quad \cos x=-c$, so there is no critical number when $|c|>1$. If $|c|\le 1$, then there are infinitely many critical numbers. If x_1 is the unique solution of $\cos x=-c$ in the interval $[0,\pi]$, then the critical numbers are $2n\pi \pm x_1$, where n ranges over the integers. (Special cases: When $c=-1$, $x_1=0$; when $c=0$, $x=\frac{\pi}{2}$; and when $c=1$, $x_1=\pi$.)

$f''(x)<0 \quad \Leftrightarrow \quad \sin x>0$, so f is CD on intervals of the form $(2n\pi,(2n+1)\pi)$. f is CU on intervals of the form $((2n-1)\pi,2n\pi)$. The inflection points of f are the points $(n\pi,n\pi c)$, where n is an integer.

If $c\ge 1$, then $f'(x)\ge 0$ for all x, so f is increasing and has no extremum. If $c\le -1$, then $f'(x)\le 0$ for all x, so f is decreasing and has no extremum. If $|c|<1$, then $f'(x)>0 \quad \Leftrightarrow \quad \cos x>-c \quad \Leftrightarrow \quad x$ is in an interval of the form $(2n\pi-x_1,2n\pi+x_1)$ for some integer n. These are the intervals on which f is increasing. Similarly, we find that f is decreasing on the intervals of the form $(2n\pi+x_1,2(n+1)\pi-x_1)$. Thus, f has local maxima at the points $2n\pi+x_1$, where f has the values $c(2n\pi+x_1)+\sin x_1=c(2n\pi+x_1)+\sqrt{1-c^2}$, and f has local minima at the points $2n\pi-x_1$, where we have $f(2n\pi-x_1)=c(2n\pi-x_1)-\sin x_1=c(2n\pi-x_1)-\sqrt{1-c^2}$.

The transitional values of c are -1 and 1. The inflection points move vertically, but not horizontally, when c changes.

When $|c|\ge 1$, there is no extremum. For $|c|<1$, the maxima are spaced 2π apart horizontally, as are the minima. The horizontal spacing between maxima and adjacent minima is regular (and equals π) when $c=0$, but the horizontal space between a local maximum and the nearest local minimum shrinks as $|c|$ approaches 1.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. For $f(t) = C(e^{-at} - e^{-bt})$, C affects only vertical stretching, so we let $C = 1$. From the first figure, we notice that the graphs all pass through the origin, approach the t-axis as t increases, and approach $-\infty$ as $t \to -\infty$. Next we let $a = 2$ and produce the second figure.

Here, as b increases, the slope of the tangent at the origin increases and the local maximum value increases.

$f(t) = e^{-2t} - e^{-bt} \quad \Rightarrow \quad f'(t) = be^{-bt} - 2e^{-2t}$. $f'(0) = b - 2$, which increases as b increases.

$f'(t) = 0 \quad \Rightarrow \quad be^{-bt} = 2e^{-2t} \quad \Rightarrow \quad \dfrac{b}{2} = e^{(b-2)t} \quad \Rightarrow \quad \ln \dfrac{b}{2} = (b-2)t \quad \Rightarrow \quad t = t_1 = \dfrac{\ln b - \ln 2}{b-2}$, which decreases as b increases (the maximum is getting closer to the y-axis). $f(t_1) = \dfrac{(b-2)2^{2/(b-2)}}{b^{1+2/(b-2)}}$. We can show that this value increases as b increases by considering it to be a function of b and graphing its derivative with respect to b, which is always positive.

37. If $c < 0$, then $\lim\limits_{x\to-\infty} f(x) = \lim\limits_{x\to-\infty} xe^{-cx} = \lim\limits_{x\to-\infty} \dfrac{x}{e^{cx}} \overset{\text{H}}{=} \lim\limits_{x\to-\infty} \dfrac{1}{ce^{cx}} = 0$, and $\lim\limits_{x\to\infty} f(x) = \infty$.

If $c > 0$, then $\lim\limits_{x\to-\infty} f(x) = -\infty$, and $\lim\limits_{x\to\infty} f(x) \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1}{ce^{cx}} = 0$.

If $c = 0$, then $f(x) = x$, so $\lim\limits_{x\to\pm\infty} f(x) = \pm\infty$, respectively.

So we see that $c = 0$ is a transitional value. We now exclude the case $c = 0$, since we know how the function behaves in that case. To find the maxima and minima of f, we differentiate: $f(x) = xe^{-cx} \quad \Rightarrow$ $f'(x) = x(-ce^{-cx}) + e^{-cx} = (1 - cx)e^{-cx}$. This is 0 when $1 - cx = 0 \quad \Leftrightarrow \quad x = 1/c$. If $c < 0$ then this represents a minimum value of $f(1/c) = 1/(ce)$, since $f'(x)$ changes from negative to positive at $x = 1/c$; and if $c > 0$, it represents a maximum value. As $|c|$ increases, the maximum or minimum point gets closer to the origin. To find the inflection points, we differentiate again: $f'(x) = e^{-cx}(1 - cx) \quad \Rightarrow$ $f''(x) = e^{-cx}(-c) + (1 - cx)(-ce^{-cx}) = (cx - 2)ce^{-cx}$. This changes sign when $cx - 2 = 0 \quad \Leftrightarrow \quad x = 2/c$. So as $|c|$ increases, the points of inflection get closer to the origin.

38. (a) $f(x) = cx^4 - 4x^2 + 1 \quad \Rightarrow \quad f'(x) = 4cx^3 - 8x = 4x(cx^2 - 2)$. If $c \le 0$, then the only real solution of $f'(x) = 0$ is $x = 0$. f' changes from positive to negative at $x = 0$, so f has only a maximum point in this case. If $c > 0$, then $f'(x) = 4x(cx^2 - 2) = 4x\left(\sqrt{c}\,x + \sqrt{2}\right)\left(\sqrt{c}\,x - \sqrt{2}\right)$, and f changes from negative to positive at $x = \pm\sqrt{2}/\sqrt{c}$. Thus, if $c > 0$, the curve has minimum points.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $f\left(\pm\frac{\sqrt{2}}{\sqrt{c}}\right) = c\left(\pm\frac{\sqrt{2}}{c}\right)^4 - 4\left(\pm\frac{\sqrt{2}}{\sqrt{c}}\right)^2 + 1 = \frac{4}{c} - \frac{8}{c} + 1 = -\frac{4}{c} + 1$. For $y = g(x) = -2x^2 + 1$, we have $g\left(\pm\frac{\sqrt{2}}{\sqrt{c}}\right) = -2\left(\pm\frac{\sqrt{2}}{c}\right)^2 + 1 = -\frac{4}{c} + 1$. Also, $f(0) = 1$ and $g(0) = 1$. Thus, the minimum points $\left(\pm\frac{\sqrt{2}}{\sqrt{c}}, -\frac{4}{c} + 1\right)$ and the maximum point $(0,1)$ of every curve in the family $f(x) = cx^4 - 4x^2 + 1$ lie on the parabola $y = -2x^2 + 1$.

39. For $c = 0$, there is no inflection point; the curve is CU everywhere. If c increases, the curve simply becomes steeper, and there are still no inflection points. If c starts at 0 and decreases, a slight upward bulge appears near $x = 0$, so that there are two inflection points for any $c < 0$. This can be seen algebraically by calculating the second derivative: $f(x) = x^4 + cx^2 + x \quad \Rightarrow \quad f'(x) = 4x^3 + 2cx + 1 \quad \Rightarrow \quad f''(x) = 12x^2 + 2c$. Thus, $f''(x) > 0$ when $c > 0$. For $c < 0$, there are inflection points when $x = \pm\sqrt{-\frac{1}{6}c}$. For $c = 0$, the graph has one critical number, at the absolute minimum somewhere around $x = -0.6$. As c increases, the number of critical points does not change. If c instead decreases from 0, we see that the graph eventually sprouts another local minimum, to the right of the origin, somewhere between $x = 1$ and $x = 2$. Consequently, there is also a maximum near $x = 0$.

After a bit of experimentation, we find that at $c = -1.5$, there appear to be two critical numbers: the absolute minimum at about $x = -1$, and a horizontal tangent with no extremum at about $x = 0.5$. For any c smaller than this there will be 3 critical points, as shown in the graphs with $c = -3$ and with $c = -5$.

To prove this algebraically, we calculate $f'(x) = 4x^3 + 2cx + 1$. Now if we substitute our value of $c = -1.5$, the formula for $f'(x)$ becomes $4x^3 - 3x + 1 = (x + 1)(2x - 1)^2$. This has a double solution at $x = \frac{1}{2}$, indicating that the function has two critical points: $x = -1$ and $x = \frac{1}{2}$, just as we had guessed from the graph.

40. (a) $f(x) = 2x^3 + cx^2 + 2x \quad \Rightarrow \quad f'(x) = 6x^2 + 2cx + 2 = 2(3x^2 + cx + 1)$. $f'(x) = 0 \quad \Leftrightarrow \quad x = \dfrac{-c \pm \sqrt{c^2 - 12}}{6}$.

So f has critical points $\quad \Leftrightarrow \quad c^2 - 12 \geq 0 \quad \Leftrightarrow \quad |c| \geq 2\sqrt{3}$. For $c = \pm 2\sqrt{3}$, $f'(x) \geq 0$ on $(-\infty, \infty)$, so f' does not change signs at $-c/6$, and there is no extremum. If $c^2 - 12 > 0$, then f' changes from positive to negative at $x = \dfrac{-c - \sqrt{c^2 - 12}}{6}$ and from negative to positive at $x = \dfrac{-c + \sqrt{c^2 - 12}}{6}$. So f has a local maximum at $x = \dfrac{-c - \sqrt{c^2 - 12}}{6}$ and a local minimum at $x = \dfrac{-c + \sqrt{c^2 - 12}}{6}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Let x_0 be a critical number for $f(x)$. Then $f'(x_0) = 0 \quad \Rightarrow$

$3x_0^2 + cx_0 + 1 = 0 \quad \Leftrightarrow \quad c = \dfrac{-1 - 3x_0^2}{x_0}$. Now

$$f(x_0) = 2x_0^3 + cx_0^2 + 2x_0 = 2x_0^3 + x_0^2\left(\frac{-1 - 3x_0^2}{x_0}\right) + 2x_0$$

$$= 2x_0^3 - x_0 - 3x_0^3 + 2x_0 = x_0 - x_0^3$$

So the point is $(x_0, y_0) = \left(x_0, x_0 - x_0^3\right)$; that is, the point lies on the curve $y = x - x^3$.

4.7 Optimization Problems

1. (a)

First Number	Second Number	Product
1	22	22
2	21	42
3	20	60
4	19	76
5	18	90
6	17	102
7	16	112
8	15	120
9	14	126
10	13	130
11	12	132

We needn't consider pairs where the first number is larger than the second, since we can just interchange the numbers in such cases. The answer appears to be 11 and 12, but we have considered only integers in the table.

(b) Call the two numbers x and y. Then $x + y = 23$, so $y = 23 - x$. Call the product P. Then $P = xy = x(23 - x) = 23x - x^2$, so we wish to maximize the function $P(x) = 23x - x^2$. Since $P'(x) = 23 - 2x$, we see that $P'(x) = 0 \quad \Leftrightarrow \quad x = \frac{23}{2} = 11.5$. Thus, the maximum value of P is $P(11.5) = (11.5)^2 = 132.25$ and it occurs when $x = y = 11.5$.

Or: Note that $P''(x) = -2 < 0$ for all x, so P is everywhere concave downward and the local maximum at $x = 11.5$ must be an absolute maximum.

2. The two numbers are $x + 100$ and x. Minimize $f(x) = (x + 100)x = x^2 + 100x$. $f'(x) = 2x + 100 = 0 \quad \Rightarrow \quad x = -50$. Since $f''(x) = 2 > 0$, there is an absolute minimum at $x = -50$. The two numbers are 50 and -50.

3. The two numbers are x and $\dfrac{100}{x}$, where $x > 0$. Minimize $f(x) = x + \dfrac{100}{x}$. $f'(x) = 1 - \dfrac{100}{x^2} = \dfrac{x^2 - 100}{x^2}$. The critical number is $x = 10$. Since $f'(x) < 0$ for $0 < x < 10$ and $f'(x) > 0$ for $x > 10$, there is an absolute minimum at $x = 10$. The numbers are 10 and 10.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. Call the two numbers x and y. Then $x + y = 16$, so $y = 16 - x$. Call the sum of their squares S. Then

$S = x^2 + y^2 = x^2 + (16 - x)^2 \quad \Rightarrow \quad S' = 2x + 2(16 - x)(-1) = 2x - 32 + 2x = 4x - 32.$ $S' = 0 \quad \Rightarrow \quad x = 8.$

Since $S'(x) < 0$ for $0 < x < 8$ and $S'(x) > 0$ for $x > 8$, there is an absolute minimum at $x = 8$. Thus, $y = 16 - 8 = 8$ and $S = 8^2 + 8^2 = 128$.

5. Let the vertical distance be given by $v(x) = (x + 2) - x^2$, $-1 \le x \le 2$.

$v'(x) = 1 - 2x = 0 \quad \Leftrightarrow \quad x = \frac{1}{2}$. $v(-1) = 0$, $v\left(\frac{1}{2}\right) = \frac{9}{4}$, and $v(2) = 0$, so there is an absolute maximum at $x = \frac{1}{2}$. The maximum distance is $v\left(\frac{1}{2}\right) = \frac{1}{2} + 2 - \frac{1}{4} = \frac{9}{4}$.

6. Let the vertical distance be given by

$v(x) = (x^2 + 1) - (x - x^2) = 2x^2 - x + 1$. $v'(x) = 4x - 1 = 0 \quad \Leftrightarrow$ $x = \frac{1}{4}$. $v'(x) < 0$ for $x < \frac{1}{4}$ and $v'(x) > 0$ for $x > \frac{1}{4}$, so there is an absolute minimum at $x = \frac{1}{4}$. The minimum distance is $v\left(\frac{1}{4}\right) = \frac{1}{8} - \frac{1}{4} + 1 = \frac{7}{8}$.

7. If the rectangle has dimensions x and y, then its perimeter is $2x + 2y = 100$ m, so $y = 50 - x$. Thus, the area is $A = xy = x(50 - x)$. We wish to maximize the function $A(x) = x(50 - x) = 50x - x^2$, where $0 < x < 50$. Since $A'(x) = 50 - 2x = -2(x - 25)$, $A'(x) > 0$ for $0 < x < 25$ and $A'(x) < 0$ for $25 < x < 50$. Thus, A has an absolute maximum at $x = 25$, and $A(25) = 25^2 = 625$ m^2. The dimensions of the rectangle that maximize its area are $x = y = 25$ m. (The rectangle is a square.)

8. If the rectangle has dimensions x and y, then its area is $xy = 1000$ m^2, so $y = 1000/x$. The perimeter $P = 2x + 2y = 2x + 2000/x$. We wish to minimize the function $P(x) = 2x + 2000/x$ for $x > 0$.

$P'(x) = 2 - 2000/x^2 = (2/x^2)(x^2 - 1000)$, so the only critical number in the domain of P is $x = \sqrt{1000}$.

$P''(x) = 4000/x^3 > 0$, so P is concave upward throughout its domain and $P\left(\sqrt{1000}\right) = 4\sqrt{1000}$ is an absolute minimum value. The dimensions of the rectangle with minimal perimeter are $x = y = \sqrt{1000} = 10\sqrt{10}$ m. (The rectangle is a square.)

9. We need to maximize Y for $N \ge 0$. $Y(N) = \dfrac{kN}{1 + N^2} \quad \Rightarrow$

$$Y'(N) = \frac{(1 + N^2)k - kN(2N)}{(1 + N^2)^2} = \frac{k(1 - N^2)}{(1 + N^2)^2} = \frac{k(1 + N)(1 - N)}{(1 + N^2)^2}.$$

$Y'(N) > 0$ for $0 < N < 1$ and $Y'(N) < 0$ for $N > 1$. Thus, Y has an absolute maximum of $Y(1) = \frac{1}{2}k$ at $N = 1$.

10. We need to maximize P for $I \ge 0$. $P(I) = \dfrac{100I}{I^2 + I + 4} \quad \Rightarrow$

$$P'(I) = \frac{(I^2 + I + 4)(100) - 100I(2I + 1)}{(I^2 + I + 4)^2} = \frac{100(I^2 + I + 4 - 2I^2 - I)}{(I^2 + I + 4)^2} = \frac{-100(I^2 - 4)}{(I^2 + I + 4)^2} = \frac{-100(I + 2)(I - 2)}{(I^2 + I + 4)^2}.$$

$P'(I) > 0$ for $0 < I < 2$ and $P'(I) < 0$ for $I > 2$. Thus, P has an absolute maximum of $P(2) = 20$ at $I = 2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. (a)

The areas of the three figures are 12,500, 12,500, and 9000 ft^2. There appears to be a maximum area of at least 12,500 ft^2.

(b) Let x denote the length of each of two sides and three dividers.

Let y denote the length of the other two sides.

(c) Area $A =$ length $\times$ width $= y \cdot x$

(d) Length of fencing $= 750 \quad \Rightarrow \quad 5x + 2y = 750$

(e) $5x + 2y = 750 \quad \Rightarrow \quad y = 375 - \frac{5}{2}x \quad \Rightarrow \quad A(x) = \left(375 - \frac{5}{2}x\right)x = 375x - \frac{5}{2}x^2$

(f) $A'(x) = 375 - 5x = 0 \quad \Rightarrow \quad x = 75$. Since $A''(x) = -5 < 0$ there is an absolute maximum when $x = 75$. Then $y = \frac{375}{2} = 187.5$. The largest area is $75\left(\frac{375}{2}\right) = 14{,}062.5$ ft^2. These values of x and y are between the values in the first and second figures in part (a). Our original estimate was low.

12. (a)

The volumes of the resulting boxes are 1, 1.6875, and 2 ft^3. There appears to be a maximum volume of at least 2 ft^3.

(b) Let x denote the length of the side of the square being cut out. Let y denote the length of the base.

(c) Volume $V =$ length $\times$ width $\times$ height $\quad \Rightarrow \quad V = y \cdot y \cdot x = xy^2$

(d) Length of cardboard $= 3 \quad \Rightarrow \quad x + y + x = 3 \quad \Rightarrow \quad y + 2x = 3$

(e) $y + 2x = 3 \quad \Rightarrow \quad y = 3 - 2x \quad \Rightarrow \quad V(x) = x(3 - 2x)^2$

(f) $V(x) = x(3 - 2x)^2 \quad \Rightarrow$

$V'(x) = x \cdot 2(3 - 2x)(-2) + (3 - 2x)^2 \cdot 1 = (3 - 2x)[-4x + (3 - 2x)] = (3 - 2x)(-6x + 3)$,

so the critical numbers are $x = \frac{3}{2}$ and $x = \frac{1}{2}$. Now $0 \le x \le \frac{3}{2}$ and $V(0) = V\left(\frac{3}{2}\right) = 0$, so the maximum is $V\left(\frac{1}{2}\right) = \left(\frac{1}{2}\right)(2)^2 = 2$ ft^3, which is the value found from our third figure in part (a).

13.

$xy = 1.5 \times 10^6$, so $y = 1.5 \times 10^6/x$. Minimize the amount of fencing, which is $3x + 2y = 3x + 2(1.5 \times 10^6/x) = 3x + 3 \times 10^6/x = F(x)$.

$F'(x) = 3 - 3 \times 10^6/x^2 = 3(x^2 - 10^6)/x^2$. The critical number is $x = 10^3$ and $F'(x) < 0$ for $0 < x < 10^3$ and $F'(x) > 0$ if $x > 10^3$, so the absolute minimum occurs when $x = 10^3$ and $y = 1.5 \times 10^3$.

The field should be 1000 feet by 1500 feet with the middle fence parallel to the short side of the field.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14.

See the figure. We have $4x + y = 1200 \quad \Rightarrow \quad y = 1200 - 4x$. The trapezoidal area A is $\frac{1}{2}(x + 3x)y = 2xy = 2x(1200 - 4x) = 2400x - 8x^2,\ 0 < x < 300$. $A'(x) = 2400 - 16x = 0 \quad \Leftrightarrow \quad x = 150$. Since $A'(x) > 0$ for $0 < x < 150$ and $A'(x) < 0$ for $150 < x < 300$, there is an absolute maximum when $x = 150$ by the First Derivative Test for Absolute Extreme Values. $x = 150 \quad \Rightarrow \quad y = 1200 - 4(150) = 600$. The maximum area is $2xy = 2(150)(600) = 180{,}000$ ft^2.

15.

See the figure. The fencing cost \$20 per linear foot to install and the cost of the fencing on the west side will be split with the neighbor, so the farmer's cost C will be $C = \frac{1}{2}(20x) + 20y + 20x = 20y + 30x$. The area A will be maximized when $C = 5000$, so $5000 = 20y + 30x \quad \Leftrightarrow \quad 20y = 5000 - 30x \quad \Leftrightarrow$ $y = 250 - \frac{3}{2}x$. Now $A = xy = x\left(250 - \frac{3}{2}x\right) = 250x - \frac{3}{2}x^2 \quad \Rightarrow \quad A' = 250 - 3x. \quad A' = 0 \quad \Leftrightarrow \quad x = \frac{250}{3}$ and since $A'' = -3 < 0$, we have a maximum for A when $x = \frac{250}{3}$ ft and $y = 250 - \frac{3}{2}\left(\frac{250}{3}\right) = 125$ ft. [The maximum area is $125\left(\frac{250}{3}\right) = 10{,}416.\overline{6}$ ft^2.]

16.

See the figure. The fencing cost \$20 per linear foot to install and the cost of the fencing on the west side will be split with the neighbor, so the farmer's cost C will be $C = \frac{1}{2}(20x) + 20y + 20x = 20y + 30x$. The area A to be enclosed is 8000 ft^2, so $A = xy = 8000 \quad \Rightarrow \quad y = \dfrac{8000}{x}$.

Now $C = 20y + 30x = 20\left(\dfrac{8000}{x}\right) + 30x = \dfrac{160{,}000}{x} + 30x \quad \Rightarrow \quad C' = -\dfrac{160{,}000}{x^2} + 30. \quad C' = 0 \quad \Leftrightarrow$

$30 = \dfrac{160{,}000}{x^2} \quad \Leftrightarrow \quad x^2 = \dfrac{16{,}000}{3} \quad \Rightarrow \quad x = \sqrt{\dfrac{16{,}000}{3}} = 40\sqrt{\dfrac{10}{3}} = \dfrac{40}{3}\sqrt{30}$. Since $C'' = \dfrac{320{,}000}{x^3} > 0$ [for $x > 0$],

we have a minimum for C when $x = \dfrac{40}{3}\sqrt{30}$ ft and $y = \dfrac{8000}{x} = \dfrac{8000}{40} \cdot \dfrac{3}{\sqrt{30}} \cdot \dfrac{\sqrt{30}}{\sqrt{30}} = 20\sqrt{30}$ ft. [The minimum cost is $20(20\sqrt{30}) + 30\left(\frac{40}{3}\sqrt{30}\right) = 800\sqrt{30} \approx \$4381.78$.]

17. (a) Let the rectangle have sides x and y and area A, so $A = xy$ or $y = A/x$. The problem is to minimize the perimeter $= 2x + 2y = 2x + 2A/x = P(x)$. Now $P'(x) = 2 - 2A/x^2 = 2\left(x^2 - A\right)/x^2$. So the critical number is $x = \sqrt{A}$. Since $P'(x) < 0$ for $0 < x < \sqrt{A}$ and $P'(x) > 0$ for $x > \sqrt{A}$, there is an absolute minimum at $x = \sqrt{A}$. The sides of the rectangle are $\sqrt{A}$ and $A/\sqrt{A} = \sqrt{A}$, so the rectangle is a square.

(b) Let p be the perimeter and x and y the lengths of the sides, so $p = 2x + 2y \quad \Rightarrow \quad 2y = p - 2x \quad \Rightarrow \quad y = \frac{1}{2}p - x$. The area is $A(x) = x\left(\frac{1}{2}p - x\right) = \frac{1}{2}px - x^2$. Now $A'(x) = 0 \quad \Rightarrow \quad \frac{1}{2}p - 2x = 0 \quad \Rightarrow \quad 2x = \frac{1}{2}p \quad \Rightarrow \quad x = \frac{1}{4}p$. Since $A''(x) = -2 < 0$, there is an absolute maximum for A when $x = \frac{1}{4}p$ by the Second Derivative Test. The sides of the rectangle are $\frac{1}{4}p$ and $\frac{1}{2}p - \frac{1}{4}p = \frac{1}{4}p$, so the rectangle is a square.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. Let b be the length of the base of the box and h the height. The volume is $32{,}000 = b^2h \quad \Rightarrow \quad h = 32{,}000/b^2$.

The surface area of the open box is $S = b^2 + 4hb = b^2 + 4(32{,}000/b^2)b = b^2 + 4(32{,}000)/b$.

So $S'(b) = 2b - 4(32{,}000)/b^2 = 2\left(b^3 - 64{,}000\right)/b^2 = 0 \quad \Leftrightarrow \quad b = \sqrt[3]{64{,}000} = 40$. This gives an absolute minimum since $S'(b) < 0$ if $0 < b < 40$ and $S'(b) > 0$ if $b > 40$. The box should be $40 \times 40 \times 20$.

19. Let b be the length of the base of the box and h the height. The surface area is $1200 = b^2 + 4hb \quad \Rightarrow \quad h = (1200 - b^2)/(4b)$.

The volume is $V = b^2h = b^2(1200 - b^2)/4b = 300b - b^3/4 \quad \Rightarrow \quad V'(b) = 300 - \frac{3}{4}b^2$.

$V'(b) = 0 \quad \Rightarrow \quad 300 = \frac{3}{4}b^2 \quad \Rightarrow \quad b^2 = 400 \quad \Rightarrow \quad b = \sqrt{400} = 20$. Since $V'(b) > 0$ for $0 < b < 20$ and $V'(b) < 0$ for $b > 20$, there is an absolute maximum when $b = 20$ by the First Derivative Test for Absolute Extreme Values. If $b = 20$, then $h = (1200 - 20^2)/(4 \cdot 20) = 10$, so the largest possible volume is $b^2h = (20)^2(10) = 4000 \text{ cm}^3$.

20.

$V = x(4 - 2x)(3 - 3x) = 12x - 18x^2 + 6x^3,\ 0 \le x \le 1$.

$V'(x) = 12 - 36x + 18x^2 = 6(2 - 6x + 3x^2)$.

$V'(x) = 0 \quad \Rightarrow \quad x = \dfrac{3 - \sqrt{3}}{3}$ by the quadratic formula (rejecting $x = \dfrac{3 + \sqrt{3}}{3}$ since it is greater than 1). $V''(x) = -36 + 36x \quad \Rightarrow \quad V''\left(\dfrac{3 - \sqrt{3}}{3}\right) = -36 + 36\left(\dfrac{3 - \sqrt{3}}{3}\right) = -36 + 36 - 12\sqrt{3} < 0$,

so there is an absolute maximum at $x = \dfrac{3 - \sqrt{3}}{3}$ since $V(0) = V(1) = 0$. The maximum volume is

$V\left(\dfrac{3 - \sqrt{3}}{3}\right) = \dfrac{4\sqrt{3}}{3} \approx 2.31 \text{ ft}^3$.

21.

$V = lwh \quad \Rightarrow \quad 10 = (2w)(w)h = 2w^2h$, so $h = 5/w^2$.

The cost is $10(2w^2) + 6[2(2wh) + 2(hw)] = 20w^2 + 36wh$, so

$C(w) = 20w^2 + 36w\left(5/w^2\right) = 20w^2 + 180/w$.

$C'(w) = 40w - 180/w^2 = (40w^3 - 180)/w^2 = 40\left(w^3 - \frac{9}{2}\right)/w^2 \quad \Rightarrow \quad w = \sqrt[3]{\frac{9}{2}}$ is the critical number. There is an absolute minimum for C when $w = \sqrt[3]{\frac{9}{2}}$ since $C'(w) < 0$ for $0 < w < \sqrt[3]{\frac{9}{2}}$ and $C'(w) > 0$ for $w > \sqrt[3]{\frac{9}{2}}$. The minimum cost is $C\left(\sqrt[3]{\frac{9}{2}}\right) = 20\left(\sqrt[3]{\frac{9}{2}}\right)^2 + \dfrac{180}{\sqrt[3]{9/2}} \approx \$163.54$.

22.

$V = lwh \quad \Rightarrow \quad 10 = (2w)(w)h = 2w^2h$, so $h = 5/w^2$.

The cost is $10(2w^2) + 6[2(2wh) + 2(hw)] + 6(2w^2) = 32w^2 + 36wh$, so

$C(w) = 32w^2 + 36w(5/w^2) = 32w^2 + 180/w$.

$C'(w) = 64w - 180/w^2 = (64w^3 - 180)/w^2 = 4(16w^3 - 45)/w^2 \quad \Rightarrow \quad w = \sqrt[3]{\frac{45}{16}}$ is the critical number. There is an absolute minimum for C when $w = \sqrt[3]{\frac{45}{16}}$ since $C'(w) < 0$ for $0 < w < \sqrt[3]{\frac{45}{16}}$ and $C'(w) > 0$ for $w > \sqrt[3]{\frac{45}{16}}$. The minimum cost is $C\left(\sqrt[3]{\frac{45}{16}}\right) = 32\left(\sqrt[3]{\frac{45}{16}}\right)^2 + \dfrac{180}{\sqrt[3]{45/16}} \approx \$191.28$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. Let $x > 0$ be the length of the package and $y > 0$ be the length of the sides of the square base. We have $x + 4y = 108 \quad \Rightarrow$ $x = 108 - 4y$. The volume is $V = xy^2 = (108 - 4y)y^2 = 108y^2 - 4y^3$ [for $0 < y < 27$].

$V'(y) = 216y - 12y^2 = 12y(18 - y) = 0 \quad \Rightarrow \quad y = 18$ [since $y \neq 0$]. Since $V'(y) > 0$ for $0 < y < 18$ and $V'(y) < 0$ for $y > 18$, there is an absolute maximum when $y = 18$ by the First Derivative Test for Absolute Extreme Values. If $y = 18$, then $x = 108 - 4(18) = 36$, so the dimensions that give the greatest volume are $18 \text{ in} \times 18 \text{ in} \times 36 \text{ in}$, giving a greatest possible volume of $11{,}664 \text{ in}^3$.

24. Let $x > 0$ be the length of the package and $r > 0$ be the radius of the circular base. We have $x + 2\pi r = 108 \quad \Rightarrow$ $x = 108 - 2\pi r$. The volume is $V = \pi r^2 x = \pi r^2(108 - 2\pi r) = \pi(108r^2 - 2\pi r^3)$ [for $0 < r < 54/\pi$].

$V'(r) = \pi(216r - 6\pi r^2) = 6\pi r(36 - \pi r) = 0 \quad \Rightarrow \quad r = \dfrac{36}{\pi}$ [since $r \neq 0$]. Since $V'(r) > 0$ for $0 < r < \dfrac{36}{\pi}$ and $V'(r) < 0$ for $r > \dfrac{36}{\pi}$, there is an absolute maximum when $r = \dfrac{36}{\pi}$ by the First Derivative Test for Absolute Extreme Values. If $r = \dfrac{36}{\pi}$, then $x = 108 - 2\pi\left(\dfrac{36}{\pi}\right) = 108 - 72 = 36$, so the dimensions that give the greatest volume are a length of 36 in. and a base radius of $\dfrac{36}{\pi} \approx 11.46$ in., giving a greatest possible volume of $\dfrac{46{,}656}{\pi}$ in^3 [$\approx 14{,}851.1 \text{ in}^3$].

25. The distance d from the origin $(0, 0)$ to a point $(x, 2x + 3)$ on the line is given by $d = \sqrt{(x - 0)^2 + (2x + 3 - 0)^2}$ and the square of the distance is $S = d^2 = x^2 + (2x + 3)^2$. $S' = 2x + 2(2x + 3)2 = 10x + 12$ and $S' = 0 \quad \Leftrightarrow \quad x = -\frac{6}{5}$. Now $S'' = 10 > 0$, so we know that S has a minimum at $x = -\frac{6}{5}$. Thus, the y-value is $2\left(-\frac{6}{5}\right) + 3 = \frac{3}{5}$ and the point is $\left(-\frac{6}{5}, \frac{3}{5}\right)$.

26. The distance d from the point $(3, 0)$ to a point $(x, \sqrt{x}\,)$ on the curve is given by $d = \sqrt{(x - 3)^2 + (\sqrt{x} - 0)^2}$ and the square of the distance is $S = d^2 = (x - 3)^2 + x$. $S' = 2(x - 3) + 1 = 2x - 5$ and $S' = 0 \quad \Leftrightarrow \quad x = \frac{5}{2}$. Now $S'' = 2 > 0$, so we know that S has a minimum at $x = \frac{5}{2}$. Thus, the y-value is $\sqrt{\frac{5}{2}}$ and the point is $\left(\frac{5}{2}, \sqrt{\frac{5}{2}}\right)$.

27.

From the figure, we see that there are two points that are farthest away from $A(1, 0)$. The distance d from A to an arbitrary point $P(x, y)$ on the ellipse is $d = \sqrt{(x - 1)^2 + (y - 0)^2}$ and the square of the distance is $S = d^2 = x^2 - 2x + 1 + y^2 = x^2 - 2x + 1 + (4 - 4x^2) = -3x^2 - 2x + 5$. $S' = -6x - 2$ and $S' = 0 \quad \Rightarrow \quad x = -\frac{1}{3}$. Now $S'' = -6 < 0$, so we know that S has a maximum at $x = -\frac{1}{3}$. Since $-1 \le x \le 1$, $S(-1) = 4$, $S\left(-\frac{1}{3}\right) = \frac{16}{3}$, and $S(1) = 0$, we see that the maximum distance is $\sqrt{\frac{16}{3}}$. The corresponding y-values are $y = \pm\sqrt{4 - 4\left(-\frac{1}{3}\right)^2} = \pm\sqrt{\frac{32}{9}} = \pm\frac{4}{3}\sqrt{2} \approx \pm 1.89$. The points are $\left(-\frac{1}{3}, \pm\frac{4}{3}\sqrt{2}\right)$.

28. The distance d from the point $(4, 2)$ to a point $(x, \sin x)$ on the curve is given by $d = \sqrt{(x - 4)^2 + (\sin x - 2)^2}$ and the square of the distance is $S = d^2 = (x - 4)^2 + (\sin x - 2)^2$. $\quad S' = 2(x - 4) + 2(\sin x - 2)\cos x$. Using a calculator, it is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

clear that S has a minimum between 0 and 5, and from a graph of S', we find that $S' = 0 \quad \Rightarrow \quad x \approx 2.65$, so the point is about $(2.65, 0.47)$.

29.

The area of the rectangle is $(2x)(2y) = 4xy$. Also $r^2 = x^2 + y^2$ so $y = \sqrt{r^2 - x^2}$, so the area is $A(x) = 4x\sqrt{r^2 - x^2}$. Now $A'(x) = 4\left(\sqrt{r^2 - x^2} - \dfrac{x^2}{\sqrt{r^2 - x^2}}\right) = 4\,\dfrac{r^2 - 2x^2}{\sqrt{r^2 - x^2}}$. The critical number is $x = \frac{1}{\sqrt{2}}r$. Clearly this gives a maximum.

$y = \sqrt{r^2 - \left(\frac{1}{\sqrt{2}}r\right)^2} = \sqrt{\frac{1}{2}r^2} = \frac{1}{\sqrt{2}}r = x$, which tells us that the rectangle is a square. The dimensions are $2x = \sqrt{2}\,r$ and $2y = \sqrt{2}\,r$.

30.

The area of the rectangle is $(2x)(2y) = 4xy$. Now $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1$ gives $y = \dfrac{b}{a}\sqrt{a^2 - x^2}$, so we maximize $A(x) = 4\dfrac{b}{a}x\sqrt{a^2 - x^2}$.

$$A'(x) = \frac{4b}{a}\left[x \cdot \tfrac{1}{2}(a^2 - x^2)^{-1/2}(-2x) + (a^2 - x^2)^{1/2} \cdot 1\right]$$
$$= \frac{4b}{a}\,(a^2 - x^{2-1/2}[-x^2 + a^2 - x^2] = \frac{4b}{a\sqrt{a^2 - x^2}}[a^2 - 2x^2]$$

So the critical number is $x = \frac{1}{\sqrt{2}}\,a$, and this clearly gives a maximum. Then $y = \frac{1}{\sqrt{2}}\,b$, so the maximum area is $4\left(\frac{1}{\sqrt{2}}\,a\right)\left(\frac{1}{\sqrt{2}}\,b\right) = 2ab$.

31.

The height h of the equilateral triangle with sides of length L is $\frac{\sqrt{3}}{2}\,L$, since $h^2 + (L/2)^2 = L^2 \quad \Rightarrow \quad h^2 = L^2 - \frac{1}{4}L^2 = \frac{3}{4}L^2 \quad \Rightarrow$ $h = \frac{\sqrt{3}}{2}L$. Using similar triangles, $\dfrac{\frac{\sqrt{3}}{2}L - y}{x} = \dfrac{\frac{\sqrt{3}}{2}L}{L/2} = \sqrt{3} \quad \Rightarrow$ $\sqrt{3}\,x = \frac{\sqrt{3}}{2}L - y \quad \Rightarrow \quad y = \frac{\sqrt{3}}{2}L - \sqrt{3}\,x \quad \Rightarrow \quad y = \frac{\sqrt{3}}{2}(L - 2x)$.

The area of the inscribed rectangle is $A(x) = (2x)y = \sqrt{3}\,x(L - 2x) = \sqrt{3}\,Lx - 2\sqrt{3}\,x^2$, where $0 \le x \le L/2$. Now $0 = A'(x) = \sqrt{3}\,L - 4\sqrt{3}\,x \quad \Rightarrow \quad x = \sqrt{3}\,L/\left(4\sqrt{3}\right) = L/4$. Since $A(0) = A(L/2) = 0$, the maximum occurs when $x = L/4$, and $y = \frac{\sqrt{3}}{2}L - \frac{\sqrt{3}}{4}L = \frac{\sqrt{3}}{4}L$, so the dimensions are $L/2$ and $\frac{\sqrt{3}}{4}L$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32.

The area A of a trapezoid is given by $A = \frac{1}{2}h(B + b)$. From the diagram, $h = y$, $B = 2$, and $b = 2x$, so $A = \frac{1}{2}y(2 + 2x) = y(1 + x)$. Since it's easier to substitute for y^2, we'll let $T = A^2 = y^2(1 + x)^2 = (1 - x^2)(1 + x)^2$. Now

$$T' = (1 - x^2)2(1 + x) + (1 + x)^2(-2x) = -2(1 + x)[-(1 - x^2) + (1 + x)x]$$
$$= -2(1 + x)(2x^2 + x - 1) = -2(1 + x)(2x - 1)(x + 1)$$

$T' = 0 \quad \Leftrightarrow \quad x = -1$ or $x = \frac{1}{2}$. $T' > 0$ if $x < \frac{1}{2}$ and $T' < 0$ if $x > \frac{1}{2}$, so we get a maximum at $x = \frac{1}{2}$ [$x = -1$ gives us $A = 0$]. Thus, $y = \sqrt{1 - \left(\frac{1}{2}\right)^2} = \frac{\sqrt{3}}{2}$ and the maximum area is $A = y(1 + x) = \frac{\sqrt{3}}{2}\left(1 + \frac{1}{2}\right) = \frac{3\sqrt{3}}{4}$.

33.

The area of the triangle is

$A(x) = \frac{1}{2}(2t)(r + x) = t(r + x) = \sqrt{r^2 - x^2}(r + x)$. Then

$$0 = A'(x) = r\frac{-2x}{2\sqrt{r^2 - x^2}} + \sqrt{r^2 - x^2} + x\frac{-2x}{2\sqrt{r^2 - x^2}}$$
$$= -\frac{x^2 + rx}{\sqrt{r^2 - x^2}} + \sqrt{r^2 - x^2} \quad \Rightarrow$$

$$\frac{x^2 + rx}{\sqrt{r^2 - x^2}} = \sqrt{r^2 - x^2} \quad \Rightarrow \quad x^2 + rx = r^2 - x^2 \quad \Rightarrow \quad 0 = 2x^2 + rx - r^2 = (2x - r)(x + r) \quad \Rightarrow$$

$x = \frac{1}{2}r$ or $x = -r$. Now $A(r) = 0 = A(-r) \quad \Rightarrow$ the maximum occurs where $x = \frac{1}{2}r$, so the triangle has height $r + \frac{1}{2}r = \frac{3}{2}r$ and base $2\sqrt{r^2 - \left(\frac{1}{2}r\right)^2} = 2\sqrt{\frac{3}{4}r^2} = \sqrt{3}\,r$.

34.

From the figure, we have $x^2 + h^2 = a^2 \quad \Rightarrow \quad h = \sqrt{a^2 - x^2}$. The area of the isosceles triangle is $A = \frac{1}{2}(2x)h = xh = x\sqrt{a^2 - x^2}$ with $0 \le x \le a$. Now

$$A' = x \cdot \tfrac{1}{2}(a^2 - x^2)^{-1/2}(-2x) + (a^2 - x^2)^{1/2}(1)$$
$$= (a^2 - x^2)^{-1/2}[-x^2 + (a^2 - x^2)] = \frac{a^2 - 2x^2}{\sqrt{a^2 - x^2}}$$

$A' = 0 \quad \Leftrightarrow \quad x^2 = \frac{1}{2}a^2 \quad \Rightarrow \quad x = a/\sqrt{2}$. Since $A(0) = 0$, $A(a) = 0$, and $A(a/\sqrt{2}\,) = (a/\sqrt{2}\,)\sqrt{a^2/2} = \frac{1}{2}a^2$, we see that $x = a/\sqrt{2}$ gives us the maximum area and the length of the base is $2x = 2(a/\sqrt{2}\,) = \sqrt{2}\,a$. Note that the triangle has sides a, a, and $\sqrt{2}\,a$, which form a *right* triangle, with the right angle between the two sides of equal length.

35. The area of the triangle is $A = \frac{1}{2}a(2a)\sin\theta$ for $0 < \theta < \pi$. $A'(\theta) = a^2\cos\theta = 0 \quad \Rightarrow \quad \cos\theta = 0 \quad \Rightarrow \quad \theta = \frac{\pi}{2}$. Since $A'(\theta) > 0$ for $0 < \theta < \frac{\pi}{2}$ and $A'(\theta) < 0$ for $\frac{\pi}{2} < \theta < \pi$, there is an absolute maximum when $\theta = \frac{\pi}{2}$ by the First Derivative Test for Absolute Extreme Values. (The maximum area of $\frac{1}{2}a(2a)\sin\frac{\pi}{2} = a^2$ results from the triangle being a right triangle.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36.

The area of the rectangle is $A = 2xy = 2x(4 - x^2) = 8x - 2x^3,\ 0 < x < 2$.

$A'(x) = 8 - 6x^2 = 2(4 - 3x^2) = 0 \quad \Rightarrow \quad x = \dfrac{2}{\sqrt{3}}$ [since $x > 0$]. Since $A'(x) > 0$ for $0 < x < \dfrac{2}{\sqrt{3}}$ and $A'(x) < 0$ for $\dfrac{2}{\sqrt{3}} < x < 2$, there is an absolute maximum when $x = \dfrac{2}{\sqrt{3}}$ by the First Derivative Test for Absolute Extreme Values. Thus, the largest possible area of the rectangle is $A = 2\left(\dfrac{2}{\sqrt{3}}\right)\left[4 - \left(\dfrac{2}{\sqrt{3}}\right)^2\right] = \left(\dfrac{4}{\sqrt{3}}\right)\left(\dfrac{8}{3}\right) = \dfrac{32}{3\sqrt{3}} \approx 6.16$.

37.

The cylinder has volume $V = \pi y^2(2x)$. Also $x^2 + y^2 = r^2 \quad \Rightarrow \quad y^2 = r^2 - x^2$, so $V(x) = \pi(r^2 - x^2)(2x) = 2\pi(r^2x - x^3)$, where $0 \le x \le r$.

$V'(x) = 2\pi\left(r^2 - 3x^2\right) = 0 \quad \Rightarrow \quad x = r/\sqrt{3}$. Now $V(0) = V(r) = 0$, so there is a maximum when $x = r/\sqrt{3}$ and $V\left(r/\sqrt{3}\right) = \pi(r^2 - r^2/3)\left(2r/\sqrt{3}\right) = 4\pi r^3/\left(3\sqrt{3}\right)$.

38.

By similar triangles, $y/x = h/r$, so $y = hx/r$. The volume of the cylinder is $\pi x^2(h - y) = \pi hx^2 - (\pi h/r)x^3 = V(x)$. Now

$V'(x) = 2\pi hx - (3\pi h/r)x^2 = \pi hx(2 - 3x/r)$.

So $V'(x) = 0 \quad \Rightarrow \quad x = 0$ or $x = \frac{2}{3}r$. The maximum clearly occurs when $x = \frac{2}{3}r$ and then the volume is

$\pi hx^2 - (\pi h/r)x^3 = \pi hx^2(1 - x/r) = \pi\left(\frac{2}{3}r\right)^2 h\left(1 - \frac{2}{3}\right) = \frac{4}{27}\pi r^2 h$.

39.

The cylinder has surface area

$$\begin{aligned} 2(\text{area of the base}) + (\text{lateral surface area}) &= 2\pi(\text{radius})^2 + 2\pi(\text{radius})(\text{height}) \\ &= 2\pi y^2 + 2\pi y(2x) \end{aligned}$$

Now $x^2 + y^2 = r^2 \quad \Rightarrow \quad y^2 = r^2 - x^2 \quad \Rightarrow \quad y = \sqrt{r^2 - x^2}$, so the surface area is

$$\begin{aligned} S(x) &= 2\pi(r^2 - x^2) + 4\pi x\sqrt{r^2 - x^2},\ \ 0 \le x \le r \\ &= 2\pi r^2 - 2\pi x^2 + 4\pi\left(x\sqrt{r^2 - x^2}\right) \end{aligned}$$

Thus,

$$\begin{aligned} S'(x) &= 0 - 4\pi x + 4\pi\left[x \cdot \tfrac{1}{2}(r^2 - x^2)^{-1/2}(-2x) + (r^2 - x^2)^{1/2} \cdot 1\right] \\ &= 4\pi\left[-x - \frac{x^2}{\sqrt{r^2 - x^2}} + \sqrt{r^2 - x^2}\right] = 4\pi \cdot \frac{-x\sqrt{r^2 - x^2} - x^2 + r^2 - x^2}{\sqrt{r^2 - x^2}} \end{aligned}$$

$S'(x) = 0 \quad \Rightarrow \quad x\sqrt{r^2 - x^2} = r^2 - 2x^2 \quad (\star) \quad \Rightarrow \quad \left(x\sqrt{r^2 - x^2}\right)^2 = (r^2 - 2x^2)^2 \quad \Rightarrow$

$x^2(r^2 - x^2) = r^4 - 4r^2x^2 + 4x^4 \quad \Rightarrow \quad r^2x^2 - x^4 = r^4 - 4r^2x^2 + 4x^4 \quad \Rightarrow \quad 5x^4 - 5r^2x^2 + r^4 = 0$.

This is a quadratic equation in x^2. By the quadratic formula, $x^2 = \frac{5 \pm \sqrt{5}}{10}r^2$, but we reject the solution with the + sign since it doesn't satisfy $(\star)$. [The right side is negative and the left side is positive.] So $x = \sqrt{\frac{5 - \sqrt{5}}{10}}\,r$. Since $S(0) = S(r) = 0$, the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

maximum surface area occurs at the critical number and $x^2 = \frac{5-\sqrt{5}}{10}r^2 \quad \Rightarrow \quad y^2 = r^2 - \frac{5-\sqrt{5}}{10}r^2 = \frac{5+\sqrt{5}}{10}r^2 \quad \Rightarrow$ the surface area is

$$2\pi\left(\tfrac{5+\sqrt{5}}{10}\right)r^2 + 4\pi\sqrt{\tfrac{5-\sqrt{5}}{10}}\sqrt{\tfrac{5+\sqrt{5}}{10}}r^2 = \pi r^2\left[2\cdot\tfrac{5+\sqrt{5}}{10} + 4\tfrac{\sqrt{(5-\sqrt{5})(5+\sqrt{5})}}{10}\right] = \pi r^2\left[\tfrac{5+\sqrt{5}}{5} + \tfrac{2\sqrt{20}}{5}\right]$$
$$= \pi r^2\left[\tfrac{5+\sqrt{5}+2\cdot 2\sqrt{5}}{5}\right] = \pi r^2\left[\tfrac{5+5\sqrt{5}}{5}\right] = \pi r^2\left(1+\sqrt{5}\right).$$

40.

Perimeter $= 30 \quad \Rightarrow \quad 2y + x + \pi\left(\dfrac{x}{2}\right) = 30 \quad \Rightarrow$

$y = \dfrac{1}{2}\left(30 - x - \dfrac{\pi x}{2}\right) = 15 - \dfrac{x}{2} - \dfrac{\pi x}{4}$. The area is the area of the rectangle plus the area of the semicircle, or $xy + \frac{1}{2}\pi\left(\dfrac{x}{2}\right)^2$, so $A(x) = x\left(15 - \dfrac{x}{2} - \dfrac{\pi x}{4}\right) + \frac{1}{8}\pi x^2 = 15x - \frac{1}{2}x^2 - \frac{\pi}{8}x^2$.

$A'(x) = 15 - \left(1 + \frac{\pi}{4}\right)x = 0 \quad \Rightarrow \quad x = \dfrac{15}{1+\pi/4} = \dfrac{60}{4+\pi}$. $\quad A''(x) = -\left(1 + \dfrac{\pi}{4}\right) < 0$, so this gives a maximum.

The dimensions are $x = \dfrac{60}{4+\pi}$ ft and $y = 15 - \dfrac{30}{4+\pi} - \dfrac{15\pi}{4+\pi} = \dfrac{60 + 15\pi - 30 - 15\pi}{4+\pi} = \dfrac{30}{4+\pi}$ ft, so the height of the rectangle is half the base.

41.

$xy = 384 \quad \Rightarrow \quad y = 384/x$. Total area is

$A(x) = (8+x)(12 + 384/x) = 12(40 + x + 256/x)$, so

$A'(x) = 12(1 - 256/x^2) = 0 \quad \Rightarrow \quad x = 16$. There is an absolute minimum when $x = 16$ since $A'(x) < 0$ for $0 < x < 16$ and $A'(x) > 0$ for $x > 16$.

When $x = 16$, $y = 384/16 = 24$, so the dimensions are 24 cm and 36 cm.

42.

$xy = 180$, so $y = 180/x$. The printed area is

$(x-2)(y-3) = (x-2)(180/x - 3) = 186 - 3x - 360/x = A(x)$.

$A'(x) = -3 + 360/x^2 = 0$ when $x^2 = 120 \quad \Rightarrow \quad x = 2\sqrt{30}$. This gives an absolute maximum since $A'(x) > 0$ for $0 < x < 2\sqrt{30}$ and $A'(x) < 0$ for $x > 2\sqrt{30}$. When $x = 2\sqrt{30}$, $y = 180/(2\sqrt{30})$, so the dimensions are $2\sqrt{30}$ in. and $90/\sqrt{30}$ in.

43.

Let x be the length of the wire used for the square. The total area is

$$A(x) = \left(\frac{x}{4}\right)^2 + \frac{1}{2}\left(\frac{10-x}{3}\right)\frac{\sqrt{3}}{2}\left(\frac{10-x}{3}\right)$$
$$= \tfrac{1}{16}x^2 + \tfrac{\sqrt{3}}{36}(10-x)^2,\ 0 \le x \le 10$$

$A'(x) = \frac{1}{8}x - \frac{\sqrt{3}}{18}(10-x) = 0 \quad \Leftrightarrow \quad \frac{9}{72}x + \frac{4\sqrt{3}}{72}x - \frac{40\sqrt{3}}{72} = 0 \quad \Leftrightarrow \quad x = \frac{40\sqrt{3}}{9+4\sqrt{3}}$.

Now $A(0) = \left(\frac{\sqrt{3}}{36}\right)100 \approx 4.81$, $A(10) = \frac{100}{16} = 6.25$ and $A\left(\frac{40\sqrt{3}}{9+4\sqrt{3}}\right) \approx 2.72$, so

(a) The maximum area occurs when $x = 10$ m, and all the wire is used for the square.

(b) The minimum area occurs when $x = \frac{40\sqrt{3}}{9+4\sqrt{3}} \approx 4.35$ m.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44.

Total area is $A(x) = \left(\frac{x}{4}\right)^2 + \pi\left(\frac{10-x}{2\pi}\right)^2 = \frac{x^2}{16} + \frac{(10-x)^2}{4\pi},\ 0 \le x \le 10.$

$A'(x) = \frac{x}{8} - \frac{10-x}{2\pi} = \left(\frac{1}{2\pi} + \frac{1}{8}\right)x - \frac{5}{\pi} = 0 \quad \Rightarrow \quad x = 40/(4+\pi).$

$A(0) = 25/\pi \approx 7.96$, $A(10) = 6.25$, and $A(40/(4+\pi)) \approx 3.5$, so the maximum occurs when $x = 0$ m and the minimum occurs when $x = 40/(4+\pi)$ m.

45.

From the figure, the perimeter of the slice is $2r + r\theta = 32$, so $\theta = \frac{32-2r}{r}$. The area A of the slice is $A = \frac{1}{2}r^2\theta = \frac{1}{2}r^2\left(\frac{32-2r}{r}\right) = r(16-r) = 16r - r^2$ for $0 \le r \le 16$. $A'(r) = 16 - 2r$, so $A' = 0$ when $r = 8$. Since $A(0) = 0$, $A(16) = 0$, and $A(8) = 64$ in.^{2}, the largest piece comes from a pizza with radius 8 in. and diameter 16 in. Note that $\theta = 2$ radians $\approx 114.6°$, which is about 32% of the whole pizza.

46.

$L = 8\csc\theta + 4\sec\theta$, $0 < \theta < \frac{\pi}{2}$, $\frac{dL}{d\theta} = -8\csc\theta\,\cot\theta + 4\sec\theta\,\tan\theta = 0$ when $\sec\theta\,\tan\theta = 2\csc\theta\,\cot\theta \quad \Leftrightarrow \quad \tan^3\theta = 2 \quad \Leftrightarrow \quad \tan\theta = \sqrt[3]{2} \quad \Leftrightarrow \quad \theta = \tan^{-1}\sqrt[3]{2}$. $dL/d\theta < 0$ when $0 < \theta < \tan^{-1}\sqrt[3]{2}$, $dL/d\theta > 0$ when $\tan^{-1}\sqrt[3]{2} < \theta < \frac{\pi}{2}$, so L has an absolute minimum when $\theta = \tan^{-1}\sqrt[3]{2}$, and the shortest ladder has length

$L = 8\frac{\sqrt{1+2^{2/3}}}{2^{1/3}} + 4\sqrt{1+2^{2/3}} \approx 16.65$ ft.

Another method: Minimize $L^2 = x^2 + (4+y)^2$, where $\frac{x}{4+y} = \frac{8}{y}$.

47.

$h^2 + r^2 = R^2 \quad \Rightarrow \quad V = \frac{\pi}{3}r^2h = \frac{\pi}{3}(R^2 - h^2)h = \frac{\pi}{3}(R^2h - h^3).$

$V'(h) = \frac{\pi}{3}(R^2 - 3h^2) = 0$ when $h = \frac{1}{\sqrt{3}}R$. This gives an absolute maximum, since $V'(h) > 0$ for $0 < h < \frac{1}{\sqrt{3}}R$ and $V'(h) < 0$ for $h > \frac{1}{\sqrt{3}}R$. The maximum volume is $V\left(\frac{1}{\sqrt{3}}R\right) = \frac{\pi}{3}\left(\frac{1}{\sqrt{3}}R^3 - \frac{1}{3\sqrt{3}}R^3\right) = \frac{2}{9\sqrt{3}}\pi R^3.$

48. The volume and surface area of a cone with radius r and height h are given by $V = \frac{1}{3}\pi r^2 h$ and $S = \pi r\sqrt{r^2+h^2}$.

We'll minimize $A = S^2$ subject to $V = 27$. $V = 27 \quad \Rightarrow \quad \frac{1}{3}\pi r^2 h = 27 \quad \Rightarrow \quad r^2 = \frac{81}{\pi h}$ **(1)**.

$A = \pi^2 r^2(r^2+h^2) = \pi^2\left(\frac{81}{\pi h}\right)\left(\frac{81}{\pi h} + h^2\right) = \frac{81^2}{h^2} + 81\pi h$, so $A' = 0 \quad \Rightarrow \quad \frac{-2\cdot 81^2}{h^3} + 81\pi = 0 \quad \Rightarrow$

$81\pi = \frac{2\cdot 81^2}{h^3} \quad \Rightarrow \quad h^3 = \frac{162}{\pi} \quad \Rightarrow \quad h = \sqrt[3]{\frac{162}{\pi}} = 3\sqrt[3]{\frac{6}{\pi}} \approx 3.722$. From **(1)**, $r^2 = \frac{81}{\pi h} = \frac{81}{\pi\cdot 3\sqrt[3]{6/\pi}} = \frac{27}{\sqrt[3]{6\pi^2}} \quad \Rightarrow$

$r = \frac{3\sqrt{3}}{\sqrt[6]{6\pi^2}} \approx 2.632$. $A'' = 6\cdot 81^2/h^4 > 0$, so A and hence S has an absolute minimum at these values of r and h.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49.

By similar triangles, $\dfrac{H}{R} = \dfrac{H-h}{r}$ **(1)**. The volume of the inner cone is $V = \frac{1}{3}\pi r^2 h$, so we'll solve **(1)** for h. $\dfrac{Hr}{R} = H - h \quad \Rightarrow$

$h = H - \dfrac{Hr}{R} = \dfrac{HR - Hr}{R} = \dfrac{H}{R}(R-r)$ **(2)**.

Thus, $V(r) = \dfrac{\pi}{3}r^2 \cdot \dfrac{H}{R}(R-r) = \dfrac{\pi H}{3R}(Rr^2 - r^3) \quad \Rightarrow$

$V'(r) = \dfrac{\pi H}{3R}(2Rr - 3r^2) = \dfrac{\pi H}{3R}r(2R-3r)$.

$V'(r) = 0 \quad \Rightarrow \quad r = 0$ or $2R = 3r \quad \Rightarrow \quad r = \frac{2}{3}R$ and from **(2)**, $h = \dfrac{H}{R}\left(R - \frac{2}{3}R\right) = \dfrac{H}{R}\left(\frac{1}{3}R\right) = \frac{1}{3}H$.

$V'(r)$ changes from positive to negative at $r = \frac{2}{3}R$, so the inner cone has a maximum volume of

$V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi\left(\frac{2}{3}R\right)^2\left(\frac{1}{3}H\right) = \frac{4}{27} \cdot \frac{1}{3}\pi R^2 H$, which is approximately 15% of the volume of the larger cone.

50. We need to minimize F for $0 \le \theta < \pi/2$. $F(\theta) = \dfrac{\mu W}{\mu \sin\theta + \cos\theta} \quad \Rightarrow \quad F'(\theta) = \dfrac{-\mu W\,(\mu\cos\theta - \sin\theta)}{(\mu\sin\theta + \cos\theta)^2}$ [by the Reciprocal Rule]. $F'(\theta) > 0 \quad \Rightarrow \quad \mu\cos\theta - \sin\theta < 0 \quad \Rightarrow \quad \mu\cos\theta < \sin\theta \quad \Rightarrow \quad \mu < \tan\theta \quad \Rightarrow \quad \theta > \tan^{-1}\mu$. So F is decreasing on $\left(0, \tan^{-1}\mu\right)$ and increasing on $\left(\tan^{-1}\mu, \frac{\pi}{2}\right)$. Thus, F attains its minimum value at $\theta = \tan^{-1}\mu$. This maximum value is $F(\tan^{-1}\mu) = \dfrac{\mu W}{\sqrt{\mu^2+1}}$.

51. $P(R) = \dfrac{E^2 R}{(R+r)^2} \quad \Rightarrow$

$$P'(R) = \frac{(R+r)^2 \cdot E^2 - E^2 R \cdot 2(R+r)}{[(R+r)^2]^2} = \frac{(R^2+2Rr+r^2)E^2 - 2E^2R^2 - 2E^2Rr}{(R+r)^4}$$

$$= \frac{E^2r^2 - E^2R^2}{(R+r)^4} = \frac{E^2(r^2-R^2)}{(R+r)^4} = \frac{E^2(r+R)(r-R)}{(R+r)^4} = \frac{E^2(r-R)}{(R+r)^3}$$

$P'(R) = 0 \quad \Rightarrow \quad R = r \quad \Rightarrow \quad P(r) = \dfrac{E^2 r}{(r+r)^2} = \dfrac{E^2 r}{4r^2} = \dfrac{E^2}{4r}$.

The expression for $P'(R)$ shows that $P'(R) > 0$ for $R < r$ and $P'(R) < 0$ for $R > r$. Thus, the maximum value of the power is $E^2/(4r)$, and this occurs when $R = r$.

52. (a) $E(v) = \dfrac{aLv^3}{v-u} \quad \Rightarrow \quad E'(v) = aL\dfrac{(v-u)3v^2 - v^3}{(v-u)^2} = 0$ when

$2v^3 = 3uv^2 \quad \Rightarrow \quad 2v = 3u \quad \Rightarrow \quad v = \frac{3}{2}u$.

The First Derivative Test shows that this value of v gives the minimum value of E.

(b)

53. $S = 6sh - \frac{3}{2}s^2\cot\theta + \left(\frac{3}{2}\sqrt{3}\,s^2\right)\csc\theta$

(a) $\dfrac{dS}{d\theta} = \frac{3}{2}s^2\csc^2\theta - \left(\frac{3}{2}\sqrt{3}\,s^2\right)\csc\theta\,\cot\theta$ or $\frac{3}{2}s^2\csc\theta\left(\csc\theta - \sqrt{3}\cot\theta\right)$.

(b) $\dfrac{dS}{d\theta} = 0$ when $\csc\theta - \sqrt{3}\cot\theta = 0 \quad \Rightarrow \quad \dfrac{1}{\sin\theta} - \sqrt{3}\,\dfrac{\cos\theta}{\sin\theta} = 0 \quad \Rightarrow \quad \cos\theta = \frac{1}{\sqrt{3}}$. The First Derivative Test shows that the minimum surface area occurs when $\theta = \cos^{-1}\left(\frac{1}{\sqrt{3}}\right) \approx 55°$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c)

If $\cos\theta = \frac{1}{\sqrt{3}}$, then $\cot\theta = \frac{1}{\sqrt{2}}$ and $\csc\theta = \frac{\sqrt{3}}{\sqrt{2}}$, so the surface area is

$$S = 6sh - \tfrac{3}{2}s^2\tfrac{1}{\sqrt{2}} + 3s^2\tfrac{\sqrt{3}}{2}\tfrac{\sqrt{3}}{\sqrt{2}} = 6sh - \tfrac{3}{2\sqrt{2}}s^2 + \tfrac{9}{2\sqrt{2}}s^2$$
$$= 6sh + \tfrac{6}{2\sqrt{2}}s^2 = 6s\left(h + \tfrac{1}{2\sqrt{2}}s\right)$$

54.

Let t be the time, in hours, after 2:00 PM. The position of the boat heading south at time t is $(0, -20t)$. The position of the boat heading east at time t is $(-15 + 15t, 0)$. If $D(t)$ is the distance between the boats at time t, we minimize $f(t) = [D(t)]^2 = 20^2t^2 + 15^2(t-1)^2$.

$f'(t) = 800t + 450(t-1) = 1250t - 450 = 0$ when $t = \frac{450}{1250} = 0.36$ h.

$0.36 \text{ h} \times \frac{60 \text{ min}}{\text{h}} = 21.6 \text{ min} = 21 \text{ min } 36 \text{ s}$. Since $f''(t) > 0$, this gives a minimum, so the boats are closest together at 2:21:36 PM.

55. Here $T(x) = \dfrac{\sqrt{x^2+25}}{6} + \dfrac{5-x}{8}$, $0 \le x \le 5 \;\Rightarrow\; T'(x) = \dfrac{x}{6\sqrt{x^2+25}} - \dfrac{1}{8} = 0 \;\Leftrightarrow\; 8x = 6\sqrt{x^2+25} \;\Leftrightarrow$ $16x^2 = 9(x^2+25) \;\Leftrightarrow\; x = \frac{15}{\sqrt{7}}$. But $\frac{15}{\sqrt{7}} > 5$, so T has no critical number. Since $T(0) \approx 1.46$ and $T(5) \approx 1.18$, she should row directly to B.

56.

In isosceles triangle AOB, $\angle O = 180° - \theta - \theta$, so $\angle BOC = 2\theta$. The distance rowed is $4\cos\theta$ while the distance walked is the length of arc $BC = 2(2\theta) = 4\theta$. The time taken is given by $T(\theta) = \dfrac{4\cos\theta}{2} + \dfrac{4\theta}{4} = 2\cos\theta + \theta$, $0 \le \theta \le \frac{\pi}{2}$.

$T'(\theta) = -2\sin\theta + 1 = 0 \;\Leftrightarrow\; \sin\theta = \frac{1}{2} \;\Rightarrow\; \theta = \frac{\pi}{6}$.

Check the value of T at $\theta = \frac{\pi}{6}$ and at the endpoints of the domain of T; that is, $\theta = 0$ and $\theta = \frac{\pi}{2}$.

$T(0) = 2$, $T\left(\frac{\pi}{6}\right) = \sqrt{3} + \frac{\pi}{6} \approx 2.26$, and $T\left(\frac{\pi}{2}\right) = \frac{\pi}{2} \approx 1.57$. Therefore, the minimum value of T is $\frac{\pi}{2}$ when $\theta = \frac{\pi}{2}$; that is, the woman should walk all the way. Note that $T''(\theta) = -2\cos\theta < 0$ for $0 \le \theta < \frac{\pi}{2}$, so $\theta = \frac{\pi}{6}$ gives a maximum time.

57. There are $(6-x)$ km over land and $\sqrt{x^2+4}$ km under the river.

We need to minimize the cost C (measured in \$100,000) of the pipeline.

$C(x) = (6-x)(4) + \left(\sqrt{x^2+4}\right)(8) \;\Rightarrow$

$C'(x) = -4 + 8 \cdot \frac{1}{2}(x^2+4)^{-1/2}(2x) = -4 + \dfrac{8x}{\sqrt{x^2+4}}$.

$C'(x) = 0 \;\Rightarrow\; 4 = \dfrac{8x}{\sqrt{x^2+4}} \;\Rightarrow\; \sqrt{x^2+4} = 2x \;\Rightarrow\; x^2 + 4 = 4x^2 \;\Rightarrow\; 4 = 3x^2 \;\Rightarrow\; x^2 = \frac{4}{3} \;\Rightarrow$

$x = 2/\sqrt{3}$ $[0 \le x \le 6]$. Compare the costs for $x = 0$, $2/\sqrt{3}$, and 6. $C(0) = 24 + 16 = 40$,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$C\big(2/\sqrt{3}\,\big) = 24 - 8/\sqrt{3} + 32/\sqrt{3} = 24 + 24/\sqrt{3} \approx 37.9$, and $C(6) = 0 + 8\sqrt{40} \approx 50.6$. So the minimum cost is about \$3.79 million when P is $6 - 2/\sqrt{3} \approx 4.85$ km east of the refinery.

58. The distance from the refinery to P is now $\sqrt{(6-x)^2 + 1^2} = \sqrt{x^2 - 12x + 37}$.

Thus, $C(x) = 4\sqrt{x^2 - 12x + 37} + 8\sqrt{x^2 + 4} \quad \Rightarrow$

$$C'(x) = 4 \cdot \tfrac{1}{2}(x^2 - 12x + 37)^{-1/2}(2x - 12) + 8 \cdot \tfrac{1}{2}(x^2 + 4)^{-1/2}(2x) = \frac{4(x-6)}{\sqrt{x^2 - 12x + 37}} + \frac{8x}{\sqrt{x^2 + 4}}.$$

$C'(x) = 0 \quad \Rightarrow \quad x \approx 1.12$ [from a graph of C' or a numerical rootfinder]. $C(0) \approx 40.3$, $C(1.12) \approx 38.3$, and $C(6) \approx 54.6$. So the minimum cost is slightly higher (than in the previous exercise) at about \$3.83 million when P is approximately 4.88 km from the point on the bank 1 km south of the refinery.

59.

The total illumination is $I(x) = \dfrac{3k}{x^2} + \dfrac{k}{(10-x)^2}$, $0 < x < 10$. Then

$$I'(x) = \frac{-6k}{x^3} + \frac{2k}{(10-x)^3} = 0 \quad \Rightarrow \quad 6k(10-x)^3 = 2kx^3 \quad \Rightarrow$$

$3(10-x)^3 = x^3 \quad \Rightarrow \quad \sqrt[3]{3}\,(10-x) = x \quad \Rightarrow \quad 10\sqrt[3]{3} - \sqrt[3]{3}\,x = x \quad \Rightarrow \quad 10\sqrt[3]{3} = x + \sqrt[3]{3}\,x \quad \Rightarrow$

$10\sqrt[3]{3} = \big(1 + \sqrt[3]{3}\,\big)x \quad \Rightarrow \quad x = \dfrac{10\sqrt[3]{3}}{1 + \sqrt[3]{3}} \approx 5.9$ ft. This gives a minimum since $I''(x) > 0$ for $0 < x < 10$.

60.

The line with slope m (where $m < 0$) through $(3, 5)$ has equation $y - 5 = m(x - 3)$ or $y = mx + (5 - 3m)$. The y-intercept is $5 - 3m$ and the x-intercept is $-5/m + 3$. So the triangle has area $A(m) = \frac{1}{2}(5 - 3m)(-5/m + 3) = 15 - 25/(2m) - \frac{9}{2}m$. Now

$$A'(m) = \frac{25}{2m^2} - \frac{9}{2} = 0 \quad \Leftrightarrow \quad m^2 = \tfrac{25}{9} \quad \Rightarrow \quad m = -\tfrac{5}{3} \text{ (since } m < 0).$$

$A''(m) = -\dfrac{25}{m^3} > 0$, so there is an absolute minimum when $m = -\frac{5}{3}$. Thus, an equation of the line is $y - 5 = -\frac{5}{3}(x - 3)$ or $y = -\frac{5}{3}x + 10$.

61.

Every line segment in the first quadrant passing through (a, b) with endpoints on the x- and y-axes satisfies an equation of the form $y - b = m(x - a)$, where $m < 0$. By setting $x = 0$ and then $y = 0$, we find its endpoints, $A(0, b - am)$ and $B\big(a - \frac{b}{m}, 0\big)$. The distance d from A to B is given by $d = \sqrt{[(a - \frac{b}{m}) - 0]^2 + [0 - (b - am)]^2}$.

It follows that the square of the length of the line segment, as a function of m, is given by

$S(m) = \left(a - \dfrac{b}{m}\right)^2 + (am - b)^2 = a^2 - \dfrac{2ab}{m} + \dfrac{b^2}{m^2} + a^2m^2 - 2abm + b^2$. Thus,

$$\begin{aligned} S'(m) &= \frac{2ab}{m^2} - \frac{2b^2}{m^3} + 2a^2m - 2ab = \frac{2}{m^3}(abm - b^2 + a^2m^4 - abm^3) \\ &= \frac{2}{m^3}[b(am - b) + am^3(am - b)] = \frac{2}{m^3}(am - b)(b + am^3) \end{aligned}$$

Thus, $S'(m) = 0 \quad \Leftrightarrow \quad m = b/a$ or $m = -\sqrt[3]{\frac{b}{a}}$. Since $b/a > 0$ and $m < 0$, m must equal $-\sqrt[3]{\frac{b}{a}}$. Since $\dfrac{2}{m^3} < 0$, we see

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

that $S'(m) < 0$ for $m < -\sqrt[3]{\frac{b}{a}}$ and $S'(m) > 0$ for $m > -\sqrt[3]{\frac{b}{a}}$. Thus, S has its absolute minimum value when $m = -\sqrt[3]{\frac{b}{a}}$. That value is

$$S\left(-\sqrt[3]{\tfrac{b}{a}}\right) = \left(a + b\sqrt[3]{\tfrac{a}{b}}\right)^2 + \left(-a\sqrt[3]{\tfrac{b}{a}} - b\right)^2 = \left(a + \sqrt[3]{ab^2}\right)^2 + \left(\sqrt[3]{a^2b} + b\right)^2$$

$$= a^2 + 2a^{4/3}b^{2/3} + a^{2/3}b^{4/3} + a^{4/3}b^{2/3} + 2a^{2/3}b^{4/3} + b^2 = a^2 + 3a^{4/3}b^{2/3} + 3a^{2/3}b^{4/3} + b^2$$

The last expression is of the form $x^3 + 3x^2y + 3xy^2 + y^3 \quad [= (x+y)^3]$ with $x = a^{2/3}$ and $y = b^{2/3}$, so we can write it as $(a^{2/3} + b^{2/3})^3$ and the shortest such line segment has length $\sqrt{S} = (a^{2/3} + b^{2/3})^{3/2}$.

62. $y = 1 + 40x^3 - 3x^5 \quad \Rightarrow \quad y' = 120x^2 - 15x^4$, so the tangent line to the curve at $x = a$ has slope $m(a) = 120a^2 - 15a^4$. Now $m'(a) = 240a - 60a^3 = -60a(a^2 - 4) = -60a(a+2)(a-2)$, so $m'(a) > 0$ for $a < -2$, and $0 < a < 2$, and $m'(a) < 0$ for $-2 < a < 0$ and $a > 2$. Thus, m is increasing on $(-\infty, -2)$, decreasing on $(-2, 0)$, increasing on $(0, 2)$, and decreasing on $(2, \infty)$. Clearly, $m(a) \to -\infty$ as $a \to \pm\infty$, so the maximum value of $m(a)$ must be one of the two local maxima, $m(-2)$ or $m(2)$. But both $m(-2)$ and $m(2)$ equal $120 \cdot 2^2 - 15 \cdot 2^4 = 480 - 240 = 240$. So 240 is the largest slope, and it occurs at the points $(-2, -223)$ and $(2, 225)$. *Note:* $a = 0$ corresponds to a local *minimum* of m.

63. $y = \dfrac{3}{x} \quad \Rightarrow \quad y' = -\dfrac{3}{x^2}$, so an equation of the tangent line at the point $(a, \frac{3}{a})$ is $y - \dfrac{3}{a} = -\dfrac{3}{a^2}(x - a)$, or $y = -\dfrac{3}{a^2}x + \dfrac{6}{a}$. The y-intercept $\quad [x = 0] \quad$ is $6/a$. The x-intercept $\quad [y = 0] \quad$ is $2a$. The distance d of the line segment that has endpoints at the intercepts is $d = \sqrt{(2a-0)^2 + (0 - 6/a)^2}$. Let $S = d^2$, so $S = 4a^2 + \dfrac{36}{a^2} \quad \Rightarrow$

$S' = 8a - \dfrac{72}{a^3}$. $S' = 0 \quad \Leftrightarrow \quad \dfrac{72}{a^3} = 8a \quad \Leftrightarrow \quad a^4 = 9 \quad \Leftrightarrow \quad a^2 = 3 \quad \Rightarrow \quad a = \sqrt{3}$.

$S'' = 8 + \dfrac{216}{a^4} > 0$, so there is an absolute minimum at $a = \sqrt{3}$. Thus, $S = 4(3) + \frac{36}{3} = 12 + 12 = 24$ and hence, $d = \sqrt{24} = 2\sqrt{6}$.

64. $y = 4 - x^2 \quad \Rightarrow \quad y' = -2x$, so an equation of the tangent line at $(a, 4 - a^2)$ is $y - (4 - a^2) = -2a(x - a)$, or $y = -2ax + a^2 + 4$. The y-intercept $\quad [x = 0]$ is $a^2 + 4$. The x-intercept $\quad [y = 0] \quad$ is $\dfrac{a^2 + 4}{2a}$. The area A of the triangle is

$A = \dfrac{1}{2}(\text{base})(\text{height}) = \dfrac{1}{2} \cdot \dfrac{a^2 + 4}{2a}(a^2 + 4) = \dfrac{1}{4}\dfrac{a^4 + 8a^2 + 16}{a} = \dfrac{1}{4}\left(a^3 + 8a + \dfrac{16}{a}\right)$.

$A' = 0 \quad \Rightarrow \quad \dfrac{1}{4}\left(3a^2 + 8 - \dfrac{16}{a^2}\right) = 0 \quad \Rightarrow \quad 3a^4 + 8a^2 - 16 = 0 \quad \Rightarrow$

$(3a^2 - 4)(a^2 + 4) = 0 \quad \Rightarrow \quad a^2 = \dfrac{4}{3} \quad \Rightarrow \quad a = \dfrac{2}{\sqrt{3}}$. $A'' = \dfrac{1}{4}\left(6a + \dfrac{32}{a^3}\right) > 0$, so there is an absolute minimum at $a = \dfrac{2}{\sqrt{3}}$. Thus, $A = \dfrac{1}{2} \cdot \dfrac{4/3 + 4}{2(2/\sqrt{3})}\left(\dfrac{4}{3} + 4\right) = \dfrac{1}{2} \cdot \dfrac{4\sqrt{3}}{3} \cdot \dfrac{16}{3} = \dfrac{32}{9}\sqrt{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

65. (a) If $c(x) = \dfrac{C(x)}{x}$, then, by the Quotient Rule, we have $c'(x) = \dfrac{xC'(x) - C(x)}{x^2}$. Now $c'(x) = 0$ when $xC'(x) - C(x) = 0$ and this gives $C'(x) = \dfrac{C(x)}{x} = c(x)$. Therefore, the marginal cost equals the average cost.

(b) (i) $C(x) = 16{,}000 + 200x + 4x^{3/2}$, $C(1000) = 16{,}000 + 200{,}000 + 40{,}000\sqrt{10} \approx 216{,}000 + 126{,}491$, so $C(1000) \approx \$342{,}491$. $c(x) = C(x)/x = \dfrac{16{,}000}{x} + 200 + 4x^{1/2}$, $c(1000) \approx \$342.49/\text{unit}$. $C'(x) = 200 + 6x^{1/2}$, $C'(1000) = 200 + 60\sqrt{10} \approx \$389.74/\text{unit}$.

(ii) We must have $C'(x) = c(x) \quad \Leftrightarrow \quad 200 + 6x^{1/2} = \dfrac{16{,}000}{x} + 200 + 4x^{1/2} \quad \Leftrightarrow \quad 2x^{3/2} = 16{,}000 \quad \Leftrightarrow$ $x = (8{,}000)^{2/3} = 400$ units. To check that this is a minimum, we calculate $c'(x) = \dfrac{-16{,}000}{x^2} + \dfrac{2}{\sqrt{x}} = \dfrac{2}{x^2}\left(x^{3/2} - 8000\right)$. This is negative for $x < (8000)^{2/3} = 400$, zero at $x = 400$, and positive for $x > 400$, so c is decreasing on $(0, 400)$ and increasing on $(400, \infty)$. Thus, c has an absolute minimum at $x = 400$. [*Note:* $c''(x)$ is *not* positive for all $x > 0$.]

(iii) The minimum average cost is $c(400) = 40 + 200 + 80 = \$320/\text{unit}$.

66. (a) The total profit is $P(x) = R(x) - C(x)$. In order to maximize profit we look for the critical numbers of P, that is, the numbers where the marginal profit is 0. But if $P'(x) = R'(x) - C'(x) = 0$, then $R'(x) = C'(x)$. Therefore, if the profit is a maximum, then the marginal revenue equals the marginal cost.

(b) $C(x) = 16{,}000 + 500x - 1.6x^2 + 0.004x^3$, $p(x) = 1700 - 7x$. Then $R(x) = xp(x) = 1700x - 7x^2$. If the profit is maximum, then $R'(x) = C'(x) \quad \Leftrightarrow \quad 1700 - 14x = 500 - 3.2x + 0.012x^2 \quad \Leftrightarrow \quad 0.012x^2 + 10.8x - 1200 = 0 \quad \Leftrightarrow$ $x^2 + 900x - 100{,}000 = 0 \quad \Leftrightarrow \quad (x + 1000)(x - 100) = 0 \quad \Leftrightarrow \quad x = 100$ (since $x > 0$). The profit is maximized if $P''(x) < 0$, but since $P''(x) = R''(x) - C''(x)$, we can just check the condition $R''(x) < C''(x)$. Now $R''(x) = -14 < -3.2 + 0.024x = C''(x)$ for $x > 0$, so there is a maximum at $x = 100$.

67. (a) We are given that the demand function p is linear and $p(27{,}000) = 10$, $p(33{,}000) = 8$, so the slope is $\frac{10 - 8}{27{,}000 - 33{,}000} = -\frac{1}{3000}$ and an equation of the line is $y - 10 = \left(-\frac{1}{3000}\right)(x - 27{,}000) \quad \Rightarrow$ $y = p(x) = -\frac{1}{3000}x + 19 = 19 - (x/3000)$.

(b) The revenue is $R(x) = xp(x) = 19x - (x^2/3000) \quad \Rightarrow \quad R'(x) = 19 - (x/1500) = 0$ when $x = 28{,}500$. Since $R''(x) = -1/1500 < 0$, the maximum revenue occurs when $x = 28{,}500 \quad \Rightarrow \quad$ the price is $p(28{,}500) = \$9.50$.

68. (a) Let $p(x)$ be the demand function. Then $p(x)$ is linear and $y = p(x)$ passes through $(20, 10)$ and $(18, 11)$, so the slope is $-\frac{1}{2}$ and an equation of the line is $y - 10 = -\frac{1}{2}(x - 20) \quad \Leftrightarrow \quad y = -\frac{1}{2}x + 20$. Thus, the demand is $p(x) = -\frac{1}{2}x + 20$ and the revenue is $R(x) = xp(x) = -\frac{1}{2}x^2 + 20x$.

(b) The cost is $C(x) = 6x$, so the profit is $P(x) = R(x) - C(x) = -\frac{1}{2}x^2 + 14x$. Then $0 = P'(x) = -x + 14 \quad \Rightarrow$ $x = 14$. Since $P''(x) = -1 < 0$, the selling price for maximum profit is $p(14) = -\frac{1}{2}(14) + 20 = \$13$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

69. (a) As in Example 6, we see that the demand function p is linear. We are given that $p(1200) = 350$ and deduce that $p(1280) = 340$, since a \$10 reduction in price increases sales by 80 per week. The slope for p is $\dfrac{340-350}{1280-1200} = -\dfrac{1}{8}$, so an equation is $p - 350 = -\frac{1}{8}(x - 1200)$ or $p(x) = -\frac{1}{8}x + 500$, where $x \geq 1200$.

(b) $R(x) = x\,p(x) = -\frac{1}{8}x^2 + 500x$. $R'(x) = -\frac{1}{4}x + 500 = 0$ when $x = 4(500) = 2000$. $p(2000) = 250$, so the price should be set at \$250 to maximize revenue.

(c) $C(x) = 35{,}000 + 120x \quad\Rightarrow\quad P(x) = R(x) - C(x) = -\frac{1}{8}x^2 + 500x - 35{,}000 - 120x = -\frac{1}{8}x^2 + 380x - 35{,}000$. $P'(x) = -\frac{1}{4}x + 380 = 0$ when $x = 4(380) = 1520$. $p(1520) = 310$, so the price should be set at \$310 to maximize profit.

70. Let w denote the number of operating wells. Then the amount of daily oil production for each well is $240 - 8(w - 16) = 368 - 8w$, where $w \geq 16$. The total daily oil production P for all wells is given by $P(w) = w(368 - 8w) = 368w - 8w^2$. Now $P'(w) = 368 - 16w$ and $P'(w) = 0 \quad\Leftrightarrow\quad w = \frac{368}{16} = 23$. $P''(w) = -16 < 0$, so the daily production is maximized when the company adds $23 - 16 = 7$ wells.

71.

Here $s^2 = h^2 + b^2/4$, so $h^2 = s^2 - b^2/4$. The area is $A = \frac{1}{2}b\sqrt{s^2 - b^2/4}$. Let the perimeter be p, so $2s + b = p$ or $s = (p - b)/2 \quad\Rightarrow$ $A(b) = \frac{1}{2}b\sqrt{(p-b)^2/4 - b^2/4} = b\sqrt{p^2 - 2pb}/4$. Now

$$A'(b) = \frac{\sqrt{p^2 - 2pb}}{4} - \frac{bp/4}{\sqrt{p^2 - 2pb}} = \frac{-3pb + p^2}{4\sqrt{p^2 - 2pb}}.$$

Therefore, $A'(b) = 0 \quad\Rightarrow\quad -3pb + p^2 = 0 \quad\Rightarrow\quad b = p/3$. Since $A'(b) > 0$ for $b < p/3$ and $A'(b) < 0$ for $b > p/3$, there is an absolute maximum when $b = p/3$. But then $2s + p/3 = p$, so $s = p/3 \quad\Rightarrow\quad s = b \quad\Rightarrow$ the triangle is equilateral.

72. From Exercise 57, with K replacing 8 for the "under river" cost (measured in \$100,000), we see that $C'(x) = 0 \quad\Leftrightarrow$ $4\sqrt{x^2+4} = Kx \quad\Leftrightarrow\quad 16x^2 + 64 = K^2x^2 \quad\Leftrightarrow\quad 64 = (K^2 - 16)x^2 \quad\Leftrightarrow\quad x = \dfrac{8}{\sqrt{K^2 - 16}}$. Also from Exercise 57, we have $C(x) = (6 - x)4 + \sqrt{x^2+4}\,K$. We now compare costs for using the minimum distance possible under the river $[x = 0]$ and using the critical number above. $C(0) = 24 + 2K$ and

$$\begin{aligned} C\left(\frac{8}{\sqrt{K^2-16}}\right) &= 24 - \frac{32}{\sqrt{K^2-16}} + \sqrt{\frac{64}{K^2-16} + 4}\,K = 24 - \frac{32}{\sqrt{K^2-16}} + \sqrt{\frac{4K^2}{K^2-16}}\,K \\ &= 24 - \frac{32}{\sqrt{K^2-16}} + \frac{2K^2}{\sqrt{K^2-16}} = 24 + \frac{2(K^2-16)}{\sqrt{K^2-16}} = 24 + 2\sqrt{K^2-16} \end{aligned}$$

Since $\sqrt{K^2-16} < K$, we see that $C\left(\dfrac{8}{\sqrt{K^2-16}}\right) < C(0)$ for any cost K, so the minimum distance possible for the "under river" portion of the pipeline should *never* be used.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

73. (a)

Using implicit differentiation, $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 \quad \Rightarrow \quad \dfrac{2x}{a^2} + \dfrac{2y\,y'}{b^2} = 0 \quad \Rightarrow$ $\dfrac{2y\,y'}{b^2} = -\dfrac{2x}{a^2} \quad \Rightarrow \quad y' = -\dfrac{b^2x}{a^2y}$. At (p, q), $y' = -\dfrac{b^2p}{a^2q}$, and an equation of the tangent line is $y - q = -\dfrac{b^2p}{a^2q}(x - p) \quad \Leftrightarrow \quad y = -\dfrac{b^2p}{a^2q}x + \dfrac{b^2p^2}{a^2q} + q \quad \Leftrightarrow$ $y = -\dfrac{b^2p}{a^2q}x + \dfrac{b^2p^2 + a^2q^2}{a^2q}$. The last term is the y-intercept, but not the term we want, namely b^2/q. Since (p, q) is on the ellipse, we know $\dfrac{p^2}{a^2} + \dfrac{q^2}{b^2} = 1$. To use that relationship we must divide b^2p^2 in the y-intercept by a^2b^2, so divide all terms by a^2b^2. $\dfrac{(b^2p^2 + a^2q^2)/a^2b^2}{(a^2q)/a^2b^2} = \dfrac{p^2/a^2 + q^2/b^2}{q/b^2} = \dfrac{1}{q/b^2} = \dfrac{b^2}{q}$. So the tangent line has equation $y = -\dfrac{b^2p}{a^2q}x + \dfrac{b^2}{q}$. Let $y = 0$ and solve for x to find that x-intercept: $\dfrac{b^2p}{a^2q}x = \dfrac{b^2}{q} \quad \Leftrightarrow$ $x = \dfrac{b^2a^2q}{qb^2p} = \dfrac{a^2}{p}$.

(b) The portion of the tangent line cut off by the coordinate axes is the distance between the intercepts, $\left(a^2/p, 0\right)$ and $\left(0, b^2/q\right)$: $\sqrt{\left(\dfrac{a^2}{p}\right)^2 + \left(-\dfrac{b^2}{q}\right)^2} = \sqrt{\dfrac{a^4}{p^2} + \dfrac{b^4}{q^2}}$. To eliminate p or q, we turn to the relationship $\dfrac{p^2}{a^2} + \dfrac{q^2}{b^2} = 1 \quad \Leftrightarrow$ $\dfrac{q^2}{b^2} = 1 - \dfrac{p^2}{a^2} \quad \Leftrightarrow \quad q^2 = b^2 - \dfrac{b^2p^2}{a^2} \quad \Leftrightarrow \quad q^2 = \dfrac{b^2(a^2 - p^2)}{a^2}$. Now substitute for q^2 and use the square S of the distance. $S(p) = \dfrac{a^4}{p^2} + \dfrac{b^4a^2}{b^2(a^2 - p^2)} = \dfrac{a^4}{p^2} + \dfrac{a^2b^2}{a^2 - p^2}$ for $0 < p < a$. Note that as $p \to 0$ or $p \to a$, $S(p) \to \infty$, so the minimum value of S must occur at a critical number. Now $S'(p) = -\dfrac{2a^4}{p^3} + \dfrac{2a^2b^2p}{(a^2 - p^2)^2}$ and $S'(p) = 0 \quad \Leftrightarrow$ $\dfrac{2a^4}{p^3} = \dfrac{2a^2b^2p}{(a^2 - p^2)^2} \quad \Leftrightarrow \quad a^2(a^2 - p^2)^2 = b^2p^4 \quad \Rightarrow \quad a(a^2 - p^2) = bp^2 \quad \Leftrightarrow \quad a^3 = (a + b)p^2 \quad \Leftrightarrow \quad p^2 = \dfrac{a^3}{a + b}$.

Substitute for p^2 in $S(p)$:

$$\frac{a^4}{\dfrac{a^3}{a+b}} + \frac{a^2b^2}{a^2 - \dfrac{a^3}{a+b}} = \frac{a^4(a+b)}{a^3} + \frac{a^2b^2(a+b)}{a^2(a+b) - a^3} = \frac{a(a+b)}{1} + \frac{a^2b^2(a+b)}{a^2b}$$
$$= a(a+b) + b(a+b) = (a+b)(a+b) = (a+b)^2$$

Taking the square root gives us the desired minimum length of $a + b$.

(c) The triangle formed by the tangent line and the coordinate axes has area $A = \dfrac{1}{2}\left(\dfrac{a^2}{p}\right)\left(\dfrac{b^2}{q}\right)$. As in part (b), we'll use the square of the area and substitute for q^2. $S = \dfrac{a^4b^4}{4p^2q^2} = \dfrac{a^4b^4a^2}{4p^2b^2(a^2 - p^2)} = \dfrac{a^6b^2}{4p^2(a^2 - p^2)}$. Minimizing S (and hence A) is equivalent to maximizing $p^2(a^2 - p^2)$. Let $f(p) = p^2(a^2 - p^2) = a^2p^2 - p^4$ for $0 < p < a$. As in part (b), the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

minimum value of S must occur at a critical number. Now $f'(p) = 2a^2p - 4p^3 = 2p(a^2 - 2p^2)$. $\quad f'(p) = 0 \quad \Rightarrow$

$p^2 = a^2/2 \quad \Rightarrow \quad p = a/\sqrt{2}$ $[p > 0]$. Substitute for p^2 in $S(p)$: $\dfrac{a^6b^2}{4\left(\dfrac{a^2}{2}\right)\left(a^2 - \dfrac{a^2}{2}\right)} = \dfrac{a^6b^2}{a^4} = a^2b^2 = (ab)^2$.

Taking the square root gives us the desired minimum area of ab.

74. See the figure. The area is given by

$$A(x) = \tfrac{1}{2}\left(2\sqrt{a^2 - x^2}\,\right)x + \tfrac{1}{2}\left(2\sqrt{a^2 - x^2}\,\right)\left(\sqrt{x^2 + b^2 - a^2}\,\right)$$
$$= \sqrt{a^2 - x^2}\left(x + \sqrt{x^2 + b^2 - a^2}\,\right)$$

for $0 \le x \le a$. Now

$$A'(x) = \sqrt{a^2 - x^2}\left(1 + \frac{x}{\sqrt{x^2 + b^2 - a^2}}\right) + \left(x + \sqrt{x^2 + b^2 - a^2}\,\right)\frac{-x}{\sqrt{a^2 - x^2}}$$
$$= 0 \quad \Leftrightarrow$$
$$\frac{x}{\sqrt{a^2 - x^2}}\left(x + \sqrt{x^2 + b^2 - a^2}\,\right) = \sqrt{a^2 - x^2}\left(\frac{x + \sqrt{x^2 + b^2 - a^2}}{\sqrt{x^2 + b^2 - a^2}}\right).$$

Except for the trivial case where $x = 0$, $a = b$ and $A(x) = 0$, we have $x + \sqrt{x^2 + b^2 - a^2} > 0$. Hence, cancelling this factor gives $\dfrac{x}{\sqrt{a^2 - x^2}} = \dfrac{\sqrt{a^2 - x^2}}{\sqrt{x^2 + b^2 - a^2}} \quad \Rightarrow \quad x\sqrt{x^2 + b^2 - a^2} = a^2 - x^2 \quad \Rightarrow$

$x^2(x^2 + b^2 - a^2) = a^4 - 2a^2x^2 + x^4 \quad \Rightarrow \quad x^2(b^2 - a^2) = a^4 - 2a^2x^2 \quad \Rightarrow \quad x^2(b^2 + a^2) = a^4 \quad \Rightarrow \quad x = \dfrac{a^2}{\sqrt{a^2 + b^2}}$.

Now we must check the value of A at this point as well as at the endpoints of the domain to see which gives the maximum value. $A(0) = a\sqrt{b^2 - a^2}$, $A(a) = 0$ and

$$A\left(\frac{a^2}{\sqrt{a^2 + b^2}}\right) = \sqrt{a^2 - \left(\frac{a^2}{\sqrt{a^2 + b^2}}\right)^2}\left[\frac{a^2}{\sqrt{a^2 + b^2}} + \sqrt{\left(\frac{a^2}{\sqrt{a^2 + b^2}}\right)^2 + b^2 - a^2}\,\right]$$
$$= \frac{ab}{\sqrt{a^2 + b^2}}\left[\frac{a^2}{\sqrt{a^2 + b^2}} + \frac{b^2}{\sqrt{a^2 + b^2}}\right] = \frac{ab(a^2 + b^2)}{a^2 + b^2} = ab$$

Since $b \ge \sqrt{b^2 - a^2}$, $A\left(a^2/\sqrt{a^2 + b^2}\,\right) \ge A\,(0)$. So there is an absolute maximum when $x = \dfrac{a^2}{\sqrt{a^2 + b^2}}$. In this case the horizontal piece should be $\dfrac{2ab}{\sqrt{a^2 + b^2}}$ and the vertical piece should be $\dfrac{a^2 + b^2}{\sqrt{a^2 + b^2}} = \sqrt{a^2 + b^2}$.

75. Note that $|AD| = |AP| + |PD| \quad \Rightarrow \quad 5 = x + |PD| \quad \Rightarrow \quad |PD| = 5 - x$.

Using the Pythagorean Theorem for ΔPDB and ΔPDC gives us

$$L(x) = |AP| + |BP| + |CP| = x + \sqrt{(5 - x)^2 + 2^2} + \sqrt{(5 - x)^2 + 3^2}$$
$$= x + \sqrt{x^2 - 10x + 29} + \sqrt{x^2 - 10x + 34} \quad \Rightarrow$$

$L'(x) = 1 + \dfrac{x - 5}{\sqrt{x^2 - 10x + 29}} + \dfrac{x - 5}{\sqrt{x^2 - 10x + 34}}$. From the graphs of L and L', it seems that the minimum value of L is about $L(3.59) = 9.35$ m.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

76. We note that since c is the consumption in gallons per hour, and v is the velocity in miles per hour, then

$\dfrac{c}{v} = \dfrac{\text{gallons/hour}}{\text{miles/hour}} = \dfrac{\text{gallons}}{\text{mile}}$ gives us the consumption in gallons per mile, that is, the quantity G. To find the minimum,

we calculate $\dfrac{dG}{dv} = \dfrac{d}{dv}\left(\dfrac{c}{v}\right) = \dfrac{v\dfrac{dc}{dv} - c\dfrac{dv}{dv}}{v^2} = \dfrac{v\dfrac{dc}{dv} - c}{v^2}$.

This is 0 when $v\dfrac{dc}{dv} - c = 0 \quad \Leftrightarrow \quad \dfrac{dc}{dv} = \dfrac{c}{v}$. This implies that the tangent line of $c(v)$ passes through the origin, and this occurs when $v \approx 53$ mi/h. Note that the slope of the secant line through the origin and a point $(v, c(v))$ on the graph is equal to $G(v)$, and it is intuitively clear that G is minimized in the case where the secant is in fact a tangent.

77.

The total time is

$$T(x) = (\text{time from } A \text{ to } C) + (\text{time from } C \text{ to } B)$$
$$= \frac{\sqrt{a^2+x^2}}{v_1} + \frac{\sqrt{b^2+(d-x)^2}}{v_2}, \ 0 < x < d$$

$$T'(x) = \frac{x}{v_1\sqrt{a^2+x^2}} - \frac{d-x}{v_2\sqrt{b^2+(d-x)^2}} = \frac{\sin\theta_1}{v_1} - \frac{\sin\theta_2}{v_2}$$

The minimum occurs when $T'(x) = 0 \quad \Rightarrow \quad \dfrac{\sin\theta_1}{v_1} = \dfrac{\sin\theta_2}{v_2}$, or,

equivalently, $\dfrac{\sin\theta_1}{\sin\theta_2} = \dfrac{v_1}{v_2}$. [*Note:* $T''(x) > 0$]

78.

If $d = |QT|$, we minimize $f(\theta_1) = |PR| + |RS| = a\csc\theta_1 + b\csc\theta_2$.

Differentiating with respect to θ_1, and setting $\dfrac{df}{d\theta_1}$ equal to 0, we get

$$\frac{df}{d\theta_1} = 0 = -a\csc\theta_1\,\cot\theta_1 - b\csc\theta_2\,\cot\theta_2\,\frac{d\theta_2}{d\theta_1}.$$

So we need to find an expression for $\dfrac{d\theta_2}{d\theta_1}$. We can do this by observing that $|QT| = \text{constant} = a\cot\theta_1 + b\cot\theta_2$.

Differentiating this equation implicitly with respect to θ_1, we get $-a\csc^2\theta_1 - b\csc^2\theta_2\,\dfrac{d\theta_2}{d\theta_1} = 0 \quad \Rightarrow$

$\dfrac{d\theta_2}{d\theta_1} = -\dfrac{a\csc^2\theta_1}{b\csc^2\theta_2}$. We substitute this into the expression for $\dfrac{df}{d\theta_1}$ to get

$$-a\csc\theta_1\,\cot\theta_1 - b\csc\theta_2\,\cot\theta_2\left(-\frac{a\csc^2\theta_1}{b\csc^2\theta_2}\right) = 0 \quad \Leftrightarrow \quad -a\csc\theta_1\,\cot\theta_1 + a\frac{\csc^2\theta_1\,\cot\theta_2}{\csc\theta_2} = 0 \quad \Leftrightarrow$$

$\cot\theta_1\,\csc\theta_2 = \csc\theta_1\,\cot\theta_2 \quad \Leftrightarrow \quad \dfrac{\cot\theta_1}{\csc\theta_1} = \dfrac{\cot\theta_2}{\csc\theta_2} \quad \Leftrightarrow \quad \cos\theta_1 = \cos\theta_2$. Since θ_1 and θ_2 are both acute, we have $\theta_1 = \theta_2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

79.

$y^2 = x^2 + z^2$, but triangles CDE and BCA are similar, so

$z/8 = x/\left(4\sqrt{x-4}\right) \quad \Rightarrow \quad z = 2x/\sqrt{x-4}$. Thus, we minimize

$f(x) = y^2 = x^2 + 4x^2/(x-4) = x^3/(x-4),\ 4 < x \le 8.$

$$f'(x) = \frac{(x-4)(3x^2) - x^3}{(x-4)^2} = \frac{x^2[3(x-4)-x]}{(x-4)^2} = \frac{2x^2(x-6)}{(x-4)^2} = 0$$

when $x = 6$. $f'(x) < 0$ when $x < 6$, $f'(x) > 0$ when $x > 6$, so the minimum occurs when $x = 6$ in.

80.

Paradoxically, we solve this maximum problem by solving a minimum problem. Let L be the length of the line ACB going from wall to wall touching the inner corner C. As $\theta \to 0$ or $\theta \to \frac{\pi}{2}$, we have $L \to \infty$ and there will be an angle that makes L a minimum. A pipe of this length will just fit around the corner.

From the diagram, $L = L_1 + L_2 = 9\csc\theta + 6\sec\theta \quad \Rightarrow \quad dL/d\theta = -9\csc\theta\,\cot\theta + 6\sec\theta\,\tan\theta = 0$ when

$6\sec\theta\,\tan\theta = 9\csc\theta\,\cot\theta \quad \Leftrightarrow \quad \tan^3\theta = \frac{9}{6} = 1.5 \quad \Leftrightarrow \quad \tan\theta = \sqrt[3]{1.5}$. Then $\sec^2\theta = 1 + \left(\frac{3}{2}\right)^{2/3}$ and

$\csc^2\theta = 1 + \left(\frac{3}{2}\right)^{-2/3}$, so the longest pipe has length $L = 9\left[1 + \left(\frac{3}{2}\right)^{-2/3}\right]^{1/2} + 6\left[1 + \left(\frac{3}{2}\right)^{2/3}\right]^{1/2} \approx 21.07$ ft.

Or, use $\theta = \tan^{-1}\left(\sqrt[3]{1.5}\right) \approx 0.853 \quad \Rightarrow \quad L = 9\csc\theta + 6\sec\theta \approx 21.07$ ft.

81.

$$\theta = (\theta + \psi) - \psi = \arctan\frac{3t}{1} - \arctan\frac{t}{1} \quad \Rightarrow \quad \theta' = \frac{3}{1+9t^2} - \frac{1}{1+t^2}.$$

$\theta' = 0 \quad \Rightarrow \quad \dfrac{3}{1+9t^2} = \dfrac{1}{1+t^2} \quad \Rightarrow \quad 3 + 3t^2 = 1 + 9t^2 \quad \Rightarrow \quad 2 = 6t^2 \quad \Rightarrow$

$t^2 = \frac{1}{3} \quad \Rightarrow \quad t = 1/\sqrt{3}$. Thus,

$\theta = \arctan 3/\sqrt{3} - \arctan 1/\sqrt{3} = \frac{\pi}{3} - \frac{\pi}{6} = \frac{\pi}{6}.$

82. We maximize the cross-sectional area

$$A(\theta) = 10h + 2\left(\tfrac{1}{2}dh\right) = 10h + dh = 10(10\sin\theta) + (10\cos\theta)(10\sin\theta)$$
$$= 100(\sin\theta + \sin\theta\,\cos\theta),\ 0 \le \theta \le \tfrac{\pi}{2}$$

$$A'(\theta) = 100(\cos\theta + \cos^2\theta - \sin^2\theta) = 100(\cos\theta + 2\cos^2\theta - 1)$$
$$= 100(2\cos\theta - 1)(\cos\theta + 1) = 0 \text{ when } \cos\theta = \tfrac{1}{2} \quad \Leftrightarrow \quad \theta = \tfrac{\pi}{3} \qquad [\cos\theta \ne -1 \text{ since } 0 \le \theta \le \tfrac{\pi}{2}.]$$

Now $A(0) = 0$, $A\left(\frac{\pi}{2}\right) = 100$ and $A\left(\frac{\pi}{3}\right) = 75\sqrt{3} \approx 129.9$, so the maximum occurs when $\theta = \frac{\pi}{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

83.

From the figure, $\tan\alpha = \dfrac{5}{x}$ and $\tan\beta = \dfrac{2}{3-x}$. Since

$$\alpha + \beta + \theta = 180^\circ = \pi,\ \theta = \pi - \tan^{-1}\left(\frac{5}{x}\right) - \tan^{-1}\left(\frac{2}{3-x}\right) \quad \Rightarrow$$

$$\frac{d\theta}{dx} = -\frac{1}{1+\left(\dfrac{5}{x}\right)^2}\left(-\frac{5}{x^2}\right) - \frac{1}{1+\left(\dfrac{2}{3-x}\right)^2}\left[\frac{2}{(3-x)^2}\right]$$

$$= \frac{x^2}{x^2+25}\cdot\frac{5}{x^2} - \frac{(3-x)^2}{(3-x)^2+4}\cdot\frac{2}{(3-x)^2}.$$

Now $\dfrac{d\theta}{dx} = 0 \quad\Rightarrow\quad \dfrac{5}{x^2+25} = \dfrac{2}{x^2-6x+13} \quad\Rightarrow\quad 2x^2+50 = 5x^2-30x+65 \quad\Rightarrow$

$3x^2-30x+15=0 \quad\Rightarrow\quad x^2-10x+5=0 \quad\Rightarrow\quad x = 5\pm 2\sqrt{5}$. We reject the solution with the $+$ sign, since it is larger than 3. $d\theta/dx > 0$ for $x < 5-2\sqrt{5}$ and $d\theta/dx < 0$ for $x > 5-2\sqrt{5}$, so θ is maximized when $|AP| = x = 5-2\sqrt{5} \approx 0.53$.

84. Let x be the distance from the observer to the wall. Then, from the text figure,

$$\theta = \tan^{-1}\left(\frac{h+d}{x}\right) - \tan^{-1}\left(\frac{d}{x}\right),\quad x>0 \quad\Rightarrow$$

$$\frac{d\theta}{dx} = \frac{1}{1+[(h+d)/x]^2}\left[-\frac{h+d}{x^2}\right] - \frac{1}{1+(d/x)^2}\left[-\frac{d}{x^2}\right] = -\frac{h+d}{x^2+(h+d)^2} + \frac{d}{x^2+d^2}$$

$$= \frac{d[x^2+(h+d)^2]-(h+d)(x^2+d^2)}{[x^2+(h+d)^2](x^2+d^2)} = \frac{h^2d+hd^2-hx^2}{[x^2+(h+d)^2](x^2+d^2)} = 0 \quad\Leftrightarrow$$

$hx^2 = h^2d+hd^2 \quad\Leftrightarrow\quad x^2 = hd+d^2 \quad\Leftrightarrow\quad x = \sqrt{d(h+d)}$. Since $d\theta/dx > 0$ for all $x < \sqrt{d(h+d)}$ and $d\theta/dx < 0$ for all $x > \sqrt{d(h+d)}$, the absolute maximum occurs when $x = \sqrt{d(h+d)}$.

85.

In the small triangle with sides a and c and hypotenuse W, $\sin\theta = \dfrac{a}{W}$ and $\cos\theta = \dfrac{c}{W}$. In the triangle with sides b and d and hypotenuse L, $\sin\theta = \dfrac{d}{L}$ and $\cos\theta = \dfrac{b}{L}$. Thus, $a = W\sin\theta$, $c = W\cos\theta$, $d = L\sin\theta$, and $b = L\cos\theta$, so the area of the circumscribed rectangle is

$$\begin{aligned} A(\theta) &= (a+b)(c+d) = (W\sin\theta + L\cos\theta)(W\cos\theta + L\sin\theta) \\ &= W^2\sin\theta\,\cos\theta + WL\sin^2\theta + LW\cos^2\theta + L^2\sin\theta\,\cos\theta \\ &= LW\sin^2\theta + LW\cos^2\theta + (L^2+W^2)\sin\theta\,\cos\theta \\ &= LW(\sin^2\theta+\cos^2\theta) + (L^2+W^2)\cdot\tfrac{1}{2}\cdot 2\sin\theta\,\cos\theta = LW + \tfrac{1}{2}(L^2+W^2)\sin 2\theta,\ 0\le\theta\le\tfrac{\pi}{2} \end{aligned}$$

This expression shows, without calculus, that the maximum value of $A(\theta)$ occurs when $\sin 2\theta = 1 \quad\Leftrightarrow\quad 2\theta = \frac{\pi}{2} \quad\Rightarrow$ $\theta = \frac{\pi}{4}$. So the maximum area is $A\left(\frac{\pi}{4}\right) = LW + \frac{1}{2}(L^2+W^2) = \frac{1}{2}(L^2+2LW+W^2) = \frac{1}{2}(L+W)^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

86. (a) Let D be the point such that $a = |AD|$. From the text figure, $\sin\theta = \dfrac{b}{|BC|} \quad \Rightarrow \quad |BC| = b\csc\theta$ and

$\cos\theta = \dfrac{|BD|}{|BC|} = \dfrac{a - |AB|}{|BC|} \quad \Rightarrow \quad |BC| = (a - |AB|)\sec\theta$. Eliminating $|BC|$ gives

$(a - |AB|)\sec\theta = b\csc\theta \quad \Rightarrow \quad b\cot\theta = a - |AB| \quad \Rightarrow \quad |AB| = a - b\cot\theta$. The total resistance is

$R(\theta) = C\dfrac{|AB|}{r_1^4} + C\dfrac{|BC|}{r_2^4} = C\left(\dfrac{a - b\cot\theta}{r_1^4} + \dfrac{b\csc\theta}{r_2^4}\right)$.

(b) $R'(\theta) = C\left(\dfrac{b\csc^2\theta}{r_1^4} - \dfrac{b\csc\theta\,\cot\theta}{r_2^4}\right) = bC\csc\theta\left(\dfrac{\csc\theta}{r_1^4} - \dfrac{\cot\theta}{r_2^4}\right)$.

$R'(\theta) = 0 \quad \Leftrightarrow \quad \dfrac{\csc\theta}{r_1^4} = \dfrac{\cot\theta}{r_2^4} \quad \Leftrightarrow \quad \dfrac{r_2^4}{r_1^4} = \dfrac{\cot\theta}{\csc\theta} = \cos\theta$.

$R'(\theta) > 0 \quad \Leftrightarrow \quad \dfrac{\csc\theta}{r_1^4} > \dfrac{\cot\theta}{r_2^4} \quad \Rightarrow \quad \cos\theta < \dfrac{r_2^4}{r_1^4}$ and $R'(\theta) < 0$ when $\cos\theta > \dfrac{r_2^4}{r_1^4}$, so there is an absolute minimum when $\cos\theta = r_2^4/r_1^4$.

(c) When $r_2 = \frac{2}{3}r_1$, we have $\cos\theta = \left(\frac{2}{3}\right)^4$, so $\theta = \cos^{-1}\left(\frac{2}{3}\right)^4 \approx 79°$.

87. (a)

If $k =$ energy/km over land, then energy/km over water $= 1.4k$.

So the total energy is $E = 1.4k\sqrt{25 + x^2} + k(13 - x),\ 0 \le x \le 13$,

and so $\dfrac{dE}{dx} = \dfrac{1.4kx}{(25 + x^2)^{1/2}} - k$.

Set $\dfrac{dE}{dx} = 0$: $1.4kx = k(25 + x^2)^{1/2} \quad \Rightarrow \quad 1.96x^2 = x^2 + 25 \quad \Rightarrow \quad 0.96x^2 = 25 \quad \Rightarrow \quad x = \frac{5}{\sqrt{0.96}} \approx 5.1$.

Testing against the value of E at the endpoints: $E(0) = 1.4k(5) + 13k = 20k$, $E(5.1) \approx 17.9k$, $E(13) \approx 19.5k$.

Thus, to minimize energy, the bird should fly to a point about 5.1 km from B.

(b) If W/L is large, the bird would fly to a point C that is closer to B than to D to minimize the energy used flying over water.

If W/L is small, the bird would fly to a point C that is closer to D than to B to minimize the distance of the flight.

$E = W\sqrt{25 + x^2} + L(13 - x) \quad \Rightarrow \quad \dfrac{dE}{dx} = \dfrac{Wx}{\sqrt{25 + x^2}} - L = 0$ when $\dfrac{W}{L} = \dfrac{\sqrt{25 + x^2}}{x}$. By the same sort of argument as in part (a), this ratio will give the minimal expenditure of energy if the bird heads for the point x km from B.

(c) For flight direct to D, $x = 13$, so from part (b), $W/L = \frac{\sqrt{25 + 13^2}}{13} \approx 1.07$. There is no value of W/L for which the bird should fly directly to B. But note that $\lim\limits_{x \to 0^+}(W/L) = \infty$, so if the point at which E is a minimum is close to B, then W/L is large.

(d) Assuming that the birds instinctively choose the path that minimizes the energy expenditure, we can use the equation for $dE/dx = 0$ from part (a) with $1.4k = c$, $x = 4$, and $k = 1$: $c(4) = 1 \cdot (25 + 4^2)^{1/2} \quad \Rightarrow \quad c = \sqrt{41}/4 \approx 1.6$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

88. (a) $I(x) \propto \dfrac{\text{strength of source}}{(\text{distance from source})^2}$. Adding the intensities from the left and right lightbulbs,

$$I(x) = \frac{k}{x^2+d^2} + \frac{k}{(10-x)^2+d^2} = \frac{k}{x^2+d^2} + \frac{k}{x^2-20x+100+d^2}.$$

(b) The magnitude of the constant k won't affect the location of the point of maximum intensity, so for convenience we take $k = 1$. $I'(x) = -\dfrac{2x}{(x^2+d^2)^2} - \dfrac{2(x-10)}{(x^2-20x+100+d^2)^2}$.

Substituting $d = 5$ into the equations for $I(x)$and $I'(x)$, we get

$$I_5(x) = \frac{1}{x^2+25} + \frac{1}{x^2-20x+125} \quad \text{and} \quad I_5'(x) = -\frac{2x}{(x^2+25)^2} - \frac{2(x-10)}{(x^2-20x+125)^2}$$

From the graphs, it appears that $I_5(x)$ has a minimum at $x = 5$ m.

(c) Substituting $d = 10$ into the equations for $I(x)$ and $I'(x)$ gives

$$I_{10}(x) = \frac{1}{x^2+100} + \frac{1}{x^2-20x+200} \quad \text{and} \quad I_{10}'(x) = -\frac{2x}{(x^2+100)^2} - \frac{2\,(x-10)}{(x^2-20x+200)^2}$$

From the graphs, it seems that for $d = 10$, the intensity is minimized at the endpoints, that is, $x = 0$ and $x = 10$. The midpoint is now the most brightly lit point!

(d) From the first figures in parts (b) and (c), we see that the minimal illumination changes from the midpoint ($x = 5$ with $d = 5$) to the endpoints ($x = 0$ and $x = 10$ with $d = 10$).

So we try $d = 6$ (see the first figure) and we see that the minimum value still occurs at $x = 5$. Next, we let $d = 8$ (see the second figure) and we see that the minimum value occurs at the endpoints. It appears that for some value of d between 6 and 8, we must have minima at both the midpoint and the endpoints, that is, $I(5)$ must equal $I(0)$. To find this value of d, we solve $I(0) = I(5)$ (with $k = 1$):

$$\frac{1}{d^2} + \frac{1}{100+d^2} = \frac{1}{25+d^2} + \frac{1}{25+d^2} = \frac{2}{25+d^2} \quad \Rightarrow \quad (25+d^2)(100+d^2) + d^2(25+d^2) = 2d^2(100+d^2) \quad \Rightarrow$$

$$2500 + 125d^2 + d^4 + 25d^2 + d^4 = 200d^2 + 2d^4 \quad \Rightarrow \quad 2500 = 50d^2 \quad \Rightarrow \quad d^2 = 50 \quad \Rightarrow \quad d = 5\sqrt{2} \approx 7.071$$

[for $0 \le d \le 10$]. The third figure, a graph of $I(0) - I(5)$ with d independent, confirms that $I(0) - I(5) = 0$, that is,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$I(0) = I(5)$, when $d = 5\sqrt{2}$. Thus, the point of minimal illumination changes abruptly from the midpoint to the endpoints when $d = 5\sqrt{2}$.

APPLIED PROJECT The Shape of a Can

1. In this case, the amount of metal used in the making of each top or bottom is $(2r)^2 = 4r^2$. So the quantity we want to minimize is $A = 2\pi rh + 2(4r^2)$. But $V = \pi r^2 h \quad \Leftrightarrow \quad h = V/\pi r^2$. Substituting this expression for h in A gives $A = 2V/r + 8r^2$. Differentiating A with respect to r, we get $dA/dr = -2V/r^2 + 16r = 0 \quad \Rightarrow$ $16r^3 = 2V = 2\pi r^2 h \quad \Leftrightarrow \quad \dfrac{h}{r} = \dfrac{8}{\pi} \approx 2.55$. This gives a minimum because $\dfrac{d^2A}{dr^2} = 16 + \dfrac{4V}{r^3} > 0$.

2.

We need to find the area of metal used up by each end, that is, the area of each hexagon. We subdivide the hexagon into six congruent triangles, each sharing one side (s in the diagram) with the hexagon. We calculate the length of $s = 2r\tan\frac{\pi}{6} = \frac{2}{\sqrt{3}}r$, so the area of each triangle is $\frac{1}{2}sr = \frac{1}{\sqrt{3}}r^2$, and the total area of the hexagon is $6 \cdot \frac{1}{\sqrt{3}}r^2 = 2\sqrt{3}\,r^2$. So the quantity we want to minimize is $A = 2\pi rh + 2 \cdot 2\sqrt{3}\,r^2$. Substituting for h as in Problem 1 and differentiating, we get $\dfrac{dA}{dr} = -\dfrac{2V}{r^2} + 8\sqrt{3}\,r$. Setting this equal to 0, we get $8\sqrt{3}\,r^3 = 2V = 2\pi r^2 h \quad \Rightarrow \quad \dfrac{h}{r} = \dfrac{4\sqrt{3}}{\pi} \approx 2.21$. Again this minimizes A because $\dfrac{d^2A}{dr^2} = 8\sqrt{3} + \dfrac{4V}{r^3} > 0$.

3. Let $C = 4\sqrt{3}\,r^2 + 2\pi rh + k\,(4\pi r + h) = 4\sqrt{3}\,r^2 + 2\pi r\left(\dfrac{V}{\pi r^2}\right) + k\left(4\pi r + \dfrac{V}{\pi r^2}\right)$. Then $\dfrac{dC}{dr} = 8\sqrt{3}\,r - \dfrac{2V}{r^2} + 4k\pi - \dfrac{2kV}{\pi r^3}$. Setting this equal to 0, dividing by 2 and substituting $\dfrac{V}{r^2} = \pi h$ and $\dfrac{V}{\pi r^3} = \dfrac{h}{r}$ in the second and fourth terms respectively, we get $0 = 4\sqrt{3}\,r - \pi h + 2k\pi - \dfrac{kh}{r} \quad \Leftrightarrow$ $k\left(2\pi - \dfrac{h}{r}\right) = \pi h - 4\sqrt{3}\,r \quad \Rightarrow \quad \dfrac{k}{r}\dfrac{2\pi - h/r}{\pi h/r - 4\sqrt{3}} = 1$. We now multiply by $\dfrac{\sqrt[3]{V}}{k}$, noting that $\dfrac{\sqrt[3]{V}}{k}\dfrac{k}{r} = \sqrt[3]{\dfrac{V}{r^3}} = \sqrt[3]{\dfrac{\pi h}{r}}$, and get $\dfrac{\sqrt[3]{V}}{k} = \sqrt[3]{\dfrac{\pi h}{r}} \cdot \dfrac{2\pi - h/r}{\pi h/r - 4\sqrt{3}}$.

4.

Let $\sqrt[3]{V}/k = T$ and $h/r = x$ so that $T(x) = \sqrt[3]{\pi x} \cdot \dfrac{2\pi - x}{\pi x - 4\sqrt{3}}$. We see from the graph of T that when the ratio $\sqrt[3]{V}/k$ is large; that is, either the volume of the can is large or the cost of joining (proportional to k) is small, the optimum value of h/r is about 2.21, but when $\sqrt[3]{V}/k$ is small, indicating small volume or expensive joining, the optimum value of h/r is larger. (The part of the graph for $\sqrt[3]{V}/k < 0$ has no physical meaning, but confirms the location of the asymptote.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. Our conclusion is usually true in practice. But there are exceptions, such as cans of tuna, which may have to do with the shape of a reasonable slice of tuna. And for a comfortable grip on a soda or beer can, the geometry of the human hand is a restriction on the radius. Other possible considerations are packaging, transportation and stocking constraints, aesthetic appeal and other marketing concerns. Also, there may be better models than ours which prescribe a differently shaped can in special circumstances.

APPLIED PROJECT Planes and Birds: Minimizing Energy

1. $P(v) = Av^3 + \dfrac{BL^2}{v} \quad \Rightarrow \quad P'(v) = 3Av^2 - \dfrac{BL^2}{v^2}$. $\quad P'(v) = 0 \quad \Leftrightarrow \quad 3Av^2 = \dfrac{BL^2}{v^2} \quad \Leftrightarrow \quad v^4 = \dfrac{BL^2}{3A} \quad \Rightarrow$

$v = \sqrt[4]{\dfrac{BL^2}{3A}}$. $\quad P''(v) = 6Av + \dfrac{2BL^2}{v^3} > 0$, so the speed that minimizes the required power is $v_P = \left(\dfrac{BL^2}{3A}\right)^{1/4}$.

2. $E(v) = \dfrac{P(v)}{v} = Av^2 + \dfrac{BL^2}{v^2} \quad \Rightarrow \quad E'(v) = 2Av - \dfrac{2BL^2}{v^3}$. $\quad E'(v) = 0 \quad \Leftrightarrow \quad 2Av = \dfrac{2BL^2}{v^3} \quad \Leftrightarrow \quad v^4 = \dfrac{BL^2}{A} \quad \Rightarrow$

$v = \sqrt[4]{\dfrac{BL^2}{A}}$. $\quad E''(v) = 2A + \dfrac{6BL^2}{v^4} > 0$, so the speed that minimizes the energy needed to propel the plane is

$v_E = \left(\dfrac{BL^2}{A}\right)^{1/4}$.

3. $\dfrac{v_E}{v_P} = \dfrac{\left(\dfrac{BL^2}{A}\right)^{1/4}}{\left(\dfrac{BL^2}{3A}\right)^{1/4}} = \left(\dfrac{\dfrac{BL^2}{A}}{\dfrac{BL^2}{3A}}\right)^{1/4} = 3^{1/4} \approx 1.316$. Thus, $v_E \approx 1.316\, v_P$, so the speed for minimum energy is about 31.6% greater (faster) than the speed for minimum power.

4. Since x is the fraction of flying time spent in flapping mode, $1 - x$ is the fraction of time spent in folded mode. The average power $\overline{P}$ is the weighted average of P_{flap} and P_{fold}, so

$$\begin{aligned}\overline{P} &= xP_{\text{flap}} + (1-x)P_{\text{fold}} = x\left[(A_b + A_w)\,v^3 + \frac{B(mg/x)^2}{v}\right] + (1-x)A_b v^3 \\ &= xA_b v^3 + xA_w v^3 + x\,\frac{Bm^2g^2}{x^2 v} + A_b v^3 - xA_b v^3 = A_b v^3 + xA_w v^3 + \frac{Bm^2g^2}{xv}\end{aligned}$$

5. $\overline{P}(x) = A_b v^3 + xA_w v^3 + \dfrac{Bm^2g^2}{xv} \quad \Rightarrow \quad \overline{P}'(x) = A_w v^3 - \dfrac{Bm^2g^2}{x^2 v}$. $\quad \overline{P}'(x) = 0 \quad \Leftrightarrow \quad A_w v^3 = \dfrac{Bm^2g^2}{x^2 v} \quad \Leftrightarrow$

$x^2 = \dfrac{Bm^2g^2}{A_w v^4} \quad \Rightarrow \quad x = \dfrac{mg}{v^2}\sqrt{\dfrac{B}{A_w}}$. Since $\overline{P}''(x) = \dfrac{2Bm^2g^2}{x^3 v} > 0$, this critical number, call it $x_{\overline{P}}$, gives an absolute minimum for the average power. If the bird flies slowly, then v is smaller and $x_{\overline{P}}$ increases, and the bird spends a larger fraction of its flying time flapping. If the bird flies faster and faster, then v is larger and $x_{\overline{P}}$ decreases, and the bird spends a smaller fraction of its flying time flapping, while still minimizing average power.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $\overline{E}(x) = \dfrac{\overline{P}(x)}{v} \quad \Rightarrow \quad \overline{E}'(x) = \dfrac{1}{v}\overline{P}'(x)$, so $\overline{E}'(x) = 0 \quad \Leftrightarrow \quad \overline{P}'(x) = 0$. The value of x that minimizes $\overline{E}$ is the same value of x that minimizes $\overline{P}$, namely $x_{\overline{P}} = \dfrac{mg}{v^2}\sqrt{\dfrac{B}{A_w}}$.

4.8 Newton's Method

1. (a)

The tangent line at $x_1 = 6$ intersects the x-axis at $x \approx 7.3$, so $x_2 = 7.3$. The tangent line at $x = 7.3$ intersects the x-axis at $x \approx 6.8$, so $x_3 \approx 6.8$.

(b) $x_1 = 8$ would be a better first approximation because the tangent line at $x = 8$ intersects the x-axis closer to s than does the first approximation $x_1 = 6$.

2.

The tangent line at $x_1 = 1$ intersects the x-axis at $x \approx 3.5$, so $x_2 = 3.5$. The tangent line at $x = 3.5$ intersects the x-axis at $x \approx 2.8$, so $x_3 = 2.8$.

3. Since the tangent line $y = 9 - 2x$ is tangent to the curve $y = f(x)$ at the point $(2, 5)$, we have $x_1 = 2$, $f(x_1) = 5$, and $f'(x_1) = -2$ [the slope of the tangent line]. Thus, by Equation 2,

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)} = 2 - \frac{5}{-2} = \frac{9}{2}$$

Note that geometrically $\frac{9}{2}$ represents the x-intercept of the tangent line $y = 9 - 2x$.

4. (a)

If $x_1 = 0$, then x_2 is negative, and x_3 is even more negative. The sequence of approximations does not converge, that is, Newton's method fails.

(b)

If $x_1 = 1$, the tangent line is horizontal and Newton's method fails.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c)

If $x_1 = 3$, then $x_2 = 1$ and we have the same situation as in part (b). Newton's method fails again.

(d)

If $x_1 = 4$, the tangent line is horizontal and Newton's method fails.

(e)

If $x_1 = 5$, then x_2 is greater than 6, x_3 gets closer to 6, and the sequence of approximations converges to 6. Newton's method succeeds!

5. The initial approximations $x_1 = a, b$, and c will work, resulting in a second approximation closer to the origin, and lead to the solution of the equation $f(x) = 0$, namely, $x = 0$. The initial approximation $x_1 = d$ will not work because it will result in successive approximations farther and farther from the origin.

6. $f(x) = 2x^3 - 3x^2 + 2 \quad \Rightarrow \quad f'(x) = 6x^2 - 6x$, so $x_{n+1} = x_n - \dfrac{2x_n^3 - 3x_n^2 + 2}{6x_n^2 - 6x_n}$. Now $x_1 = -1 \quad \Rightarrow$

$$x_2 = -1 - \frac{2(-1)^3 - 3(-1)^2 + 2}{6(-1)^2 - 6(-1)} = -1 - \frac{-3}{12} = -\frac{3}{4} \quad \Rightarrow$$

$$x_3 = -\frac{3}{4} - \frac{2\left(-\frac{3}{4}\right)^3 - 3\left(-\frac{3}{4}\right)^2 + 2}{6\left(-\frac{3}{4}\right)^2 - 6\left(-\frac{3}{4}\right)} = -\frac{3}{4} - \frac{-17/32}{63/8} = -\frac{43}{63} \approx -0.6825.$$

7. $f(x) = \dfrac{2}{x} - x^2 + 1 \quad \Rightarrow \quad f'(x) = -\dfrac{2}{x^2} - 2x$, so $x_{n+1} = x_n - \dfrac{2/x_n - x_n^2 + 1}{-2/x_n^2 - 2x_n}$. Now $x_1 = 2 \quad \Rightarrow$

$$x_2 = 2 - \frac{1 - 4 + 1}{-1/2 - 4} = 2 - \frac{-2}{-9/2} = \frac{14}{9} \quad \Rightarrow \quad x_3 = \frac{14}{9} - \frac{2/(14/9) - (14/9)^2 + 1}{-2(14/9)^2 - 2(14/9)} \approx 1.5215.$$

8. Solving $x^5 = x^2 + 1$ is the same as solving $f(x) = x^5 - x^2 - 1 = 0$. $f'(x) = 5x^4 - 2x$, so $x_{n+1} = x_n - \dfrac{x_n^5 - x_n^2 - 1}{5x_n^4 - 2x_n}$.

Now, $x_1 = 1 \quad \Rightarrow \quad x_2 = 1 - \dfrac{1^5 - 1^2 - 1}{5(1)^4 - 2(1)} = 1 - \dfrac{-1}{3} = \dfrac{4}{3} \quad \Rightarrow \quad x_3 = \dfrac{4}{3} - \dfrac{(4/3)^5 - (4/3)^2 - 1}{5(4/3)^4 - 2(4/3)} \approx 1.2240.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $f(x) = x^3 + x + 3 \quad \Rightarrow \quad f'(x) = 3x^2 + 1$, so $x_{n+1} = x_n - \dfrac{x_n^3 + x_n + 3}{3x_n^2 + 1}$.

Now $x_1 = -1 \quad \Rightarrow$

$$x_2 = -1 - \frac{(-1)^3 + (-1) + 3}{3(-1)^2 + 1} = -1 - \frac{-1 - 1 + 3}{3 + 1} = -1 - \frac{1}{4} = -1.25.$$

Newton's method follows the tangent line at $(-1, 1)$ down to its intersection with the x-axis at $(-1.25, 0)$, giving the second approximation $x_2 = -1.25$.

10. $f(x) = x^4 - x - 1 \quad \Rightarrow \quad f'(x) = 4x^3 - 1$, so $x_{n+1} = x_n - \dfrac{x_n^4 - x_n - 1}{4x_n^3 - 1}$.

Now $x_1 = 1 \quad \Rightarrow \quad x_2 = 1 - \dfrac{1^4 - 1 - 1}{4 \cdot 1^3 - 1} = 1 - \dfrac{-1}{3} = \dfrac{4}{3}$. Newton's method follows the tangent line at $(1, -1)$ up to its intersection with the x-axis at $\left(\frac{4}{3}, 0\right)$, giving the second approximation $x_2 = \frac{4}{3}$.

11. To approximate $x = \sqrt[4]{75}$ (so that $x^4 = 75$), we can take $f(x) = x^4 - 75$. So $f'(x) = 4x^3$, and thus, $x_{n+1} = x_n - \dfrac{x_n^4 - 75}{4x_n^3}$. Since $\sqrt[4]{81} = 3$ and 81 is reasonably close to 75, we'll use $x_1 = 3$. We need to find approximations until they agree to eight decimal places. $x_1 = 3 \quad \Rightarrow \quad x_2 = 2.9\overline{4}$, $x_3 \approx 2.94283228$, $x_4 \approx 2.94283096 \approx x_5$. So $\sqrt[4]{75} \approx 2.94283096$, to eight decimal places.

To use Newton's method on a calculator, assign f to Y_1 and f' to Y_2. Then store x_1 in X and enter $\mathrm{X} - \mathrm{Y}_1/\mathrm{Y}_2 \to \mathrm{X}$ to get x_2 and further approximations (repeatedly press ENTER).

12. $f(x) = x^8 - 500 \quad \Rightarrow \quad f'(x) = 8x^7$, so $x_{n+1} = x_n - \dfrac{x_n^8 - 500}{8x_n^7}$. Since $\sqrt[8]{256} = 2$ and 256 is reasonably close to 500, we'll use $x_1 = 2$. We need to find approximations until they agree to eight decimal places. $x_1 = 2 \quad \Rightarrow \quad x_2 \approx 2.23828125$, $x_3 \approx 2.18055972$, $x_4 \approx 2.17461675$, $x_5 \approx 2.17455928 \approx x_6$. So $\sqrt[8]{500} \approx 2.17455928$, to eight decimal places.

13. (a) Let $f(x) = 3x^4 - 8x^3 + 2$. The polynomial f is continuous on $[2, 3]$, $f(2) = -14 < 0$, and $f(3) = 29 > 0$, so by the Intermediate Value Theorem, there is a number c in $(2, 3)$ such that $f(c) = 0$. In other words, the equation $3x^4 - 8x^3 + 2 = 0$ has a solution in $[2, 3]$.

(b) $f'(x) = 12x^3 - 24x^2 \quad \Rightarrow \quad x_{n+1} = x_n - \dfrac{3x_n^4 - 8x_n^3 + 2}{12x_n^3 - 24x_n^2}$. Taking $x_1 = 2.5$, we get $x_2 = 2.655$, $x_3 \approx 2.630725$, $x_4 \approx 2.630021$, $x_5 \approx 2.630020 \approx x_6$. To six decimal places, the solution is 2.630020. Note that taking $x_1 = 2$ is not allowed since $f'(2) = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. (a) Let $f(x) = -2x^5 + 9x^4 - 7x^3 - 11x$. The polynomial f is continuous on $[3, 4]$, $f(3) = 21 > 0$, and $f(4) = -236 < 0$, so by the Intermediate Value Theorem, there is a number c in $(3, 4)$ such that $f(c) = 0$. In other words, the equation $-2x^5 + 9x^4 - 7x^3 - 11x = 0$ has a solution in $[3, 4]$.

(b) $f'(x) = -10x^4 + 36x^3 - 21x^2 - 11$. $\quad x_{n+1} = x_n - \dfrac{-2x_n^5 + 9x_n^4 - 7x_n^3 - 11x_n}{-10x_n^4 + 36x_n^3 - 21x_n^2 - 11}$. Taking $x_1 = 3.5$, we get $x_2 \approx 3.329174$, $x_3 = 3.278706$, $x_4 \approx 3.274501$, and $x_5 \approx 3.274473 \approx x_6$. To six decimal places, the solution is 3.274473.

15.

From the graph, we see that the negative solution of $\cos x = x^2 - 4$ is near $x = -2$. Solving $\cos x = x^2 - 4$ is the same as solving $f(x) = \cos x - x^2 + 4 = 0$.

$f'(x) = -\sin x - 2x$, so $x_{n+1} = x_n - \dfrac{\cos x_n - x_n^2 + 4}{-\sin x_n - 2x_n}$. $x_1 = -2 \quad \Rightarrow$

$x_2 \approx -1.915233$, $x_3 \approx -1.914021 \approx x_4$. Thus, the negative solution is -1.914021, to six decimal places.

16.

From the graph, we see that the positive solution of $e^{2x} = x + 3$ is near $x = 1$. Solving $e^{2x} = x + 3$ is the same as solving $f(x) = e^{2x} - x - 3 = 0$.

$f'(x) = 2e^{2x} - 1$, so $x_{n+1} = x_n - \dfrac{e^{2x_n} - x_n - 3}{2e^{2x_n} - 1}$. $\quad x_1 = 1 \quad \Rightarrow$

$x_2 \approx 0.754026$, $x_3 \approx 0.658965$, $x_4 \approx 0.647110$, $x_5 \approx 0.646945 \approx x_6$. Thus, the positive solution is 0.646945, to six decimal places.

17.

From the graph, we see that there appears to be a point of intersection near $x = 2$. Solving $\sin x = x - 1$ is the same as solving $f(x) = \sin x - x + 1 = 0$.

$f'(x) = \cos x - 1$, so $x_{n+1} = x_n - \dfrac{\sin x_n - x_n + 1}{\cos x_n - 1}$. $\quad x_1 = 2 \quad \Rightarrow$

$x_2 \approx 1.935951$, $x_3 \approx 1.934564$, $x_4 \approx 1.934563 \approx x_5$. Thus, the solution is 1.934563, to six decimal places.

18.

From the graph, we see that there appears to be a point of intersection near $x = 0.5$. Solving $\cos 2x = x^3$ is the same as solving $f(x) = \cos 2x - x^3 = 0$.

$f'(x) = -2\sin 2x - 3x^2$, so $x_{n+1} = x_n - \dfrac{\cos 2x_n - x_n^3}{-2\sin 2x_n - 3x_n^2}$. $\quad x_1 = 0.5 \quad \Rightarrow$

$x_2 \approx 0.670700$, $x_3 \approx 0.648160$, $x_4 \approx 0.647766$, $x_5 \approx 0.647765 \approx x_6$. Thus, the solution 0.647765, to six decimal places.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19.

From the figure, we see that the graphs intersect between -2 and -1 and between 0 and 1. Solving $2^x = 2 - x^2$ is the same as solving $f(x) = 2^x - 2 + x^2 = 0$. $f'(x) = 2^x \ln 2 + 2x$, so

$$x_{n+1} = x_n - \frac{2^{x_n} - 2 + x_n^2}{2^{x_n} \ln 2 + 2x_n}.$$

$x_1 = -1$	$x_1 = 1$
$x_2 \approx -1.302402$	$x_2 \approx 0.704692$
$x_3 \approx -1.258636$	$x_3 \approx 0.654915$
$x_4 \approx -1.257692$	$x_4 \approx 0.653484$
$x_5 \approx -1.257691 \approx x_6$	$x_5 \approx 0.653483 \approx x_6$

To six decimal places, the solutions of the equation are -1.257691 and 0.653483.

20.

From the figure, we see that the graphs intersect between 0 and 1 and between 3 and 4. Solving $\ln x = \dfrac{1}{x-3}$ is the same as solving $f(x) = \ln x - \dfrac{1}{x-3} = 0$.

$f'(x) = \dfrac{1}{x} + \dfrac{1}{(x-3)^2}$, so $x_{n+1} = x_n - \dfrac{\ln x_n - 1/(x_n - 3)}{(1/x_n) + 1/(x_n - 3)^2}$.

$x_1 = 1$	$x_1 = 4$
$x_2 \approx 0.6$	$x_2 \approx 3.690965$
$x_3 \approx 0.651166$	$x_3 \approx 3.750726$
$x_4 \approx 0.653057$	$x_4 \approx 3.755672$
$x_5 \approx 0.653060 \approx x_6$	$x_5 \approx 3.755701 \approx x_6$

To six decimal places, the solutions of the equation are 0.653060 and 3.755701.

21.

From the figure, we see that there appear to be points of intersection near $x = -1$ and $x = 2$. Solving $\arctan x = x^2 - 3$ is the same as solving $f(x) = \arctan x - x^2 + 3 = 0$. $\quad f'(x) = \dfrac{1}{1+x^2} - 2x$, so

$$x_{n+1} = x_n - \frac{\arctan x_n - x_n^2 + 3}{\dfrac{1}{1+x_n^2} - 2x_n}.$$

$x_1 = -1$	$x_1 = 2$
$x_2 \approx -1.485841$	$x_2 \approx 2.028197$
$x_3 \approx -1.429153$	$x_3 \approx 2.027975 \approx x_4$
$x_4 \approx -1.428293 \approx x_5$	

To six decimal places, the solutions of the equation are -1.428293 and 2.027975.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22.

From the figure, we see that there appear to be points of intersection near $x = -2$, $x = 1$, and $x = 2$. Solving $x^3 = 5x - 3$ is the same as solving $f(x) = x^3 - 5x + 3 = 0$. $f'(x) = 3x^2 - 5$, so

$$x_{n+1} = x_n - \frac{x_n^3 - 5x_n + 3}{3x_n^2 - 5}.$$

$x_1 = -2$	$x_1 = 1$	$x_1 = 2$
$x_2 \approx -2.714286$	$x_2 = 0.5$	$x_2 \approx 1.857143$
$x_3 \approx -2.513979$	$x_3 \approx 0.647059$	$x_3 \approx 1.834787$
$x_4 \approx -2.491151$	$x_4 \approx 0.656573$	$x_4 \approx 1.834244$
$x_5 \approx -2.490864 \approx x_6$	$x_5 \approx 0.656620 \approx x_6$	$x_5 \approx 1.834243 \approx x_6$

To six decimal places, the solutions of the equation are -2.490864, 0.656620, and 1.834243.

23.

$f(x) = -2x^7 - 5x^4 + 9x^3 + 5 \quad \Rightarrow \quad f'(x) = -14x^6 - 20x^3 + 27x^2 \quad \Rightarrow$

$$x_{n+1} = x_n - \frac{-2x_n^7 - 5x_n^4 + 9x_n^3 + 5}{-14x_n^6 - 20x_n^3 + 27x_n^2}.$$

From the graph of f, there appear to be solutions near -1.7, -0.7, and 1.3.

$x_1 = -1.7$	$x_1 = -0.7$	$x_1 = 1.3$
$x_2 = -1.693255$	$x_2 \approx -0.74756345$	$x_2 = 1.268776$
$x_3 \approx -1.69312035$	$x_3 \approx -0.74467752$	$x_3 \approx 1.26589387$
$x_4 \approx -1.69312029 \approx x_5$	$x_4 \approx -0.74466668 \approx x_5$	$x_4 \approx 1.26587094 \approx x_5$

To eight decimal places, the solutions of the equation are -1.69312029, -0.74466668, and 1.26587094.

24.

$f(x) = x^5 - 3x^4 + x^3 - x^2 - x + 6 \quad \Rightarrow$

$f'(x) = 5x^4 - 12x^3 + 3x^2 - 2x - 1 \quad \Rightarrow$

$x_{n+1} = x_n - \dfrac{x_n^5 - 3x_n^4 + x_n^3 - x_n^2 - x_n + 6}{5x_n^4 - 12x_n^3 + 3x_n^2 - 2x_n - 1}$. From the graph of f, there appear to be solutions near -1, 1.3, and 2.7.

$x_1 = -1$	$x_1 = 1.3$	$x_1 = 2.7$
$x_2 \approx -1.04761905$	$x_2 \approx 1.33313045$	$x_2 \approx 2.70556135$
$x_3 \approx -1.04451724$	$x_3 \approx 1.33258330$	$x_3 \approx 2.70551210$
$x_4 \approx -1.04450307 \approx x_5$	$x_4 \approx 1.33258316 \approx x_5$	$x_4 \approx 2.70551209 \approx x_5$

To eight decimal places, the solutions of the equation are -1.04450307, 1.33258316, and 2.70551209.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25.

Solving $\dfrac{x}{x^2+1} = \sqrt{1-x}$ is the same as solving

$f(x) = \dfrac{x}{x^2+1} - \sqrt{1-x} = 0.$ $f'(x) = \dfrac{1-x^2}{(x^2+1)^2} + \dfrac{1}{2\sqrt{1-x}}$ $\Rightarrow$

$$x_{n+1} = x_n - \frac{\dfrac{x_n}{x_n^2+1} - \sqrt{1-x_n}}{\dfrac{1-x_n^2}{(x_n^2+1)^2} + \dfrac{1}{2\sqrt{1-x_n}}}.$$

From the graph, we see that the curves intersect at about 0.8. $x_1 = 0.8 \Rightarrow x_2 \approx 0.76757581$, $x_3 \approx 0.76682610$, $x_4 \approx 0.76682579 \approx x_5$. To eight decimal places, the solution of the equation is 0.76682579.

26.

Solving $\cos(x^2 - x) = x^4$ is the same as solving

$f(x) = \cos(x^2 - x) - x^4 = 0.$ $f'(x) = -(2x-1)\sin(x^2 - x) - 4x^3 \Rightarrow$

$x_{n+1} = x_n - \dfrac{\cos(x_n^2 - x_n) - x_n^4}{-(2x_n - 1)\sin(x_n^2 - x_n) - 4x_n^3}$. From the equations

$y = \cos(x^2 - x)$ and $y = x^4$ and the graph, we deduce that one solution of the equation $\cos(x^2 - x) = x^4$ is $x = 1$. We also see that the graphs intersect at approximately $x = -0.7$. $x_1 = -0.7 \Rightarrow x_2 \approx -0.73654354$, $x_3 \approx -0.73486274$, $x_4 \approx -0.73485910 \approx x_5$. To eight decimal places, one solution of the equation is -0.73485910; the other solution is 1.

27.

From the graph, we see that the curves intersect at approximately $x = -0.8$ and $x = 0.8$. Solving $\sqrt{4-x^3} = e^{x^2}$ is the same as solving

$f(x) = \sqrt{4-x^3} - e^{x^2} = 0.$

$$f'(x) = \frac{1}{2\sqrt{4-x^3}} \cdot (-3x^2) - e^{x^2} \cdot 2x = -\frac{3x^2}{2\sqrt{4-x^3}} - 2xe^{x^2} \quad \Rightarrow \quad x_{n+1} = x_n - \frac{\sqrt{4-x_n^3} - e^{x_n^2}}{-\dfrac{3x_n^2}{2\sqrt{4-x_n^3}} - 2x_n e^{x_n^2}}.$$

$x_1 = -0.8$	$x_1 = 0.8$
$x_2 = -0.88815983$	$x_2 \approx 0.79186616$
$x_3 \approx -0.87842996$	$x_3 \approx 0.79177078$
$x_4 \approx -0.87828296$	$x_4 \approx 0.79177077 \approx x_5$
$x_5 \approx -0.87828292 \approx x_6$	

To eight decimal places, the solutions of the equation are -0.87828292 and 0.79177077.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28.

Solving $\ln(x^2+2) = \dfrac{3x}{\sqrt{x^2+1}}$ is the same as solving

$$f(x) = \ln(x^2+2) - \frac{3x}{\sqrt{x^2+1}} = 0.$$

$$\begin{aligned} f'(x) &= \frac{2x}{x^2+2} - \frac{(x^2+1)^{1/2}(3) - (3x)\frac{1}{2}(x^2+1)^{-1/2}(2x)}{[(x^2+1)^{1/2}]^2} \\ &= \frac{2x}{x^2+2} - \frac{(x^2+1)^{-1/2}[3(x^2+1)-3x^2]}{(x^2+1)^1} \\ &= \frac{2x}{x^2+2} - \frac{3}{(x^2+1)^{3/2}} \quad \Rightarrow \quad x_{n+1} = x_n - \frac{\ln(x_n^2+2) - \dfrac{3x_n}{\sqrt{x_n^2+1}}}{\dfrac{2x_n}{x_n^2+2} - \dfrac{3}{(x_n^2+1)^{3/2}}}. \end{aligned}$$

From the figure, we see that the graphs intersect at approximately $x = 0.2$ and $x = 4$.

$x_1 = 0.2$	$x_1 = 4$
$x_2 \approx 0.24733161$	$x_2 \approx 4.04993412$
$x_3 \approx 0.24852333$	$x_3 \approx 4.05010983$
$x_4 \approx 0.24852414 \approx x_5$	$x_4 \approx 4.05010984 \approx x_5$

To eight decimal places, the solutions of the equation are 0.24852414 and 4.05010984.

29. (a) $f(x) = x^2 - a \quad \Rightarrow \quad f'(x) = 2x$, so Newton's method gives

$$x_{n+1} = x_n - \frac{x_n^2 - a}{2x_n} = x_n - \frac{1}{2}x_n + \frac{a}{2x_n} = \frac{1}{2}x_n + \frac{a}{2x_n} = \frac{1}{2}\left(x_n + \frac{a}{x_n}\right).$$

(b) Using (a) with $a = 1000$ and $x_1 = \sqrt{900} = 30$, we get $x_2 \approx 31.666667$, $x_3 \approx 31.622807$, and $x_4 \approx 31.622777 \approx x_5$.

So $\sqrt{1000} \approx 31.622777$.

30. (a) $f(x) = \dfrac{1}{x} - a \quad \Rightarrow \quad f'(x) = -\dfrac{1}{x^2}$, so $x_{n+1} = x_n - \dfrac{1/x_n - a}{-1/x_n^2} = x_n + x_n - ax_n^2 = 2x_n - ax_n^2$.

(b) Using (a) with $a = 1.6894$ and $x_1 = \frac{1}{2} = 0.5$, we get $x_2 = 0.5754$, $x_3 \approx 0.588485$, and $x_4 \approx 0.588789 \approx x_5$.

So $1/1.6984 \approx 0.588789$.

31. $f(x) = x^3 - 3x + 6 \quad \Rightarrow \quad f'(x) = 3x^2 - 3$. If $x_1 = 1$, then $f'(x_1) = 0$ and the tangent line used for approximating x_2 is horizontal. Attempting to find x_2 results in trying to divide by zero.

32. $x^3 - x = 1 \quad \Leftrightarrow \quad x^3 - x - 1 = 0$. $f(x) = x^3 - x - 1 \quad \Rightarrow \quad f'(x) = 3x^2 - 1$, so $x_{n+1} = x_n - \dfrac{x_n^3 - x_n - 1}{3x_n^2 - 1}$.

(a) $x_1 = 1$, $x_2 = 1.5$, $x_3 \approx 1.347826$, $x_4 \approx 1.325200$, $x_5 \approx 1.324718 \approx x_6$

(b) $x_1 = 0.6$, $x_2 = 17.9$, $x_3 \approx 11.946802$, $x_4 \approx 7.985520$, $x_5 \approx 5.356909$, $x_6 \approx 3.624996$, $x_7 \approx 2.505589$, $x_8 \approx 1.820129$, $x_9 \approx 1.461044$, $x_{10} \approx 1.339323$, $x_{11} \approx 1.324913$, $x_{12} \approx 1.324718 \approx x_{13}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $x_1 = 0.57$, $x_2 \approx -54.165455$, $x_3 \approx -36.114293$, $x_4 \approx -24.082094$, $x_5 \approx -16.063387$, $x_6 \approx -10.721483$, $x_7 \approx -7.165534$, $x_8 \approx -4.801704$, $x_9 \approx -3.233425$, $x_{10} \approx -2.193674$, $x_{11} \approx -1.496867$, $x_{12} \approx -0.997546$, $x_{13} \approx -0.496305$, $x_{14} \approx -2.894162$, $x_{15} \approx -1.967962$, $x_{16} \approx -1.341355$, $x_{17} \approx -0.870187$, $x_{18} \approx -0.249949$, $x_{19} \approx -1.192219$, $x_{20} \approx -0.731952$, $x_{21} \approx 0.355213$, $x_{22} \approx -1.753322$, $x_{23} \approx -1.189420$, $x_{24} \approx -0.729123$, $x_{25} \approx 0.377844$, $x_{26} \approx -1.937872$, $x_{27} \approx -1.320350$, $x_{28} \approx -0.851919$, $x_{29} \approx -0.200959$, $x_{30} \approx -1.119386$, $x_{31} \approx -0.654291$, $x_{32} \approx 1.547010$, $x_{33} \approx 1.360051$, $x_{34} \approx 1.325828$, $x_{35} \approx 1.324719$, $x_{36} \approx 1.324718 \approx x_{37}$.

(d)

From the figure, we see that the tangent line corresponding to $x_1 = 1$ results in a sequence of approximations that converges quite quickly ($x_5 \approx x_6$). The tangent line corresponding to $x_1 = 0.6$ is close to being horizontal, so x_2 is quite far from the solution. But the sequence still converges — just a little more slowly ($x_{12} \approx x_{13}$). Lastly, the tangent line corresponding to $x_1 = 0.57$ is very nearly horizontal, x_2 is farther away from the solution, and the sequence takes more iterations to converge ($x_{36} \approx x_{37}$).

33. For $f(x) = x^{1/3}$, $f'(x) = \frac{1}{3}x^{-2/3}$ and

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^{1/3}}{\frac{1}{3}x_n^{-2/3}} = x_n - 3x_n = -2x_n.$$

Therefore, each successive approximation becomes twice as large as the previous one in absolute value, so the sequence of approximations fails to converge to the solution, which is 0. In the figure, we have $x_1 = 0.5$, $x_2 = -2(0.5) = -1$, and $x_3 = -2(-1) = 2$.

34. According to Newton's Method, for $x_n > 0$,

$$x_{n+1} = x_n - \frac{\sqrt{x_n}}{1/\left(2\sqrt{x_n}\right)} = x_n - 2x_n = -x_n \text{ and for } x_n < 0,$$

$$x_{n+1} = x_n - \frac{-\sqrt{-x_n}}{1/\left(2\sqrt{-x_n}\right)} = x_n - [-2(-x_n)] = -x_n. \text{ So we can see that}$$

after choosing any value x_1 the subsequent values will alternate between $-x_1$ and x_1 and never approach the solution.

35. (a) $f(x) = x^6 - x^4 + 3x^3 - 2x \quad \Rightarrow \quad f'(x) = 6x^5 - 4x^3 + 9x^2 - 2 \quad \Rightarrow$ $f''(x) = 30x^4 - 12x^2 + 18x$. To find the critical numbers of f, we'll find the zeros of f'. From the graph of f', it appears there are zeros at approximately $x = -1.3$, -0.4, and 0.5. Try $x_1 = -1.3 \quad \Rightarrow$

$$x_2 = x_1 - \frac{f'(x_1)}{f''(x_1)} \approx -1.293344 \quad \Rightarrow \quad x_3 \approx -1.293227 \approx x_4.$$

Now try $x_1 = -0.4 \quad \Rightarrow \quad x_2 \approx -0.443755 \quad \Rightarrow \quad x_3 \approx -0.441735 \quad \Rightarrow \quad x_4 \approx -0.441731 \approx x_5$. Finally try

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x_1 = 0.5 \quad \Rightarrow \quad x_2 \approx 0.507937 \quad \Rightarrow \quad x_3 \approx 0.507854 \approx x_4$. Therefore, $x = -1.293227$, -0.441731, and 0.507854 are all the critical numbers correct to six decimal places.

(b) There are two critical numbers where f' changes from negative to positive, so f changes from decreasing to increasing. $f(-1.293227) \approx -2.0212$ and $f(0.507854) \approx -0.6721$, so -2.0212 is the absolute minimum value of f correct to four decimal places.

36. $f(x) = x\cos x \quad \Rightarrow \quad f'(x) = \cos x - x\sin x$. $f'(x)$ exists for all x, so to find the maximum of f, we can examine the zeros of f'. From the graph of f', we see that a good choice for x_1 is $x_1 = 0.9$. Use $g(x) = \cos x - x\sin x$ and $g'(x) = -2\sin x - x\cos x$ to obtain $x_2 \approx 0.860781$, $x_3 \approx 0.860334 \approx x_4$. Now we have $f(0) = 0$, $f(\pi) = -\pi$, and $f(0.860334) \approx 0.561096$, so 0.561096 is the absolute maximum value of f correct to six decimal places.

37.

$$y = x^2\sin x \quad \Rightarrow \quad y' = x^2\cos x + (\sin x)(2x) \quad \Rightarrow$$
$$y'' = x^2(-\sin x) + (\cos x)(2x) + (\sin x)(2) + 2x\cos x$$
$$= -x^2\sin x + 4x\cos x + 2\sin x \quad \Rightarrow$$
$$y''' = -x^2\cos x + (\sin x)(-2x) + 4x(-\sin x) + (\cos x)(4) + 2\cos x$$
$$= -x^2\cos x - 6x\sin x + 6\cos x.$$

From the graph of $y = x^2\sin x$, we see that $x = 1.5$ is a reasonable guess for the x-coordinate of the inflection point. Using Newton's method with $g(x) = y''$ and $g'(x) = y'''$, we get $x_1 = 1.5 \quad \Rightarrow \quad x_2 \approx 1.520092$, $x_3 \approx 1.519855 \approx x_4$. The inflection point is about $(1.519855, 2.306964)$.

38.

$f(x) = -\sin x \quad \Rightarrow \quad f'(x) = -\cos x$. At $x = a$, the slope of the tangent line is $f'(a) = -\cos a$. The line through the origin and $(a, f(a))$ is $y = \dfrac{-\sin a - 0}{a - 0}x$. If this line is to be tangent to f at $x = a$, then its slope must equal $f'(a)$. Thus, $\dfrac{-\sin a}{a} = -\cos a \quad \Rightarrow \quad \tan a = a$.

To solve this equation using Newton's method, let $g(x) = \tan x - x$, $g'(x) = \sec^2 x - 1$, and $x_{n+1} = x_n - \dfrac{\tan x_n - x_n}{\sec^2 x_n - 1}$ with $x_1 = 4.5$ (estimated from the figure). $x_2 \approx 4.493614$, $\quad x_3 \approx 4.493410$, $\quad x_4 \approx 4.493409 \approx x_5$. Thus, the slope of the line that has the largest slope is $f'(x_5) \approx 0.217234$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. We need to minimize the distance from $(0, 0)$ to an arbitrary point (x, y) on the curve $y = (x-1)^2$. $d = \sqrt{x^2+y^2} \quad \Rightarrow$

$d(x) = \sqrt{x^2 + [(x-1)^2]^2} = \sqrt{x^2 + (x-1)^4}$. When $d' = 0$, d will be minimized and equivalently, $s = d^2$ will be minimized, so we will use Newton's method with $f = s'$ and $f' = s''$.

$f(x) = 2x + 4(x-1)^3 \quad \Rightarrow \quad f'(x) = 2 + 12(x-1)^2$, so $x_{n+1} = x_n - \dfrac{2x_n + 4(x_n-1)^3}{2 + 12(x_n-1)^2}$. Try $x_1 = 0.5 \quad \Rightarrow$

$x_2 = 0.4$, $x_3 \approx 0.410127$, $x_4 \approx 0.410245 \approx x_5$. Now $d(0.410245) \approx 0.537841$ is the minimum distance and the point on the parabola is $(0.410245, 0.347810)$, correct to six decimal places.

40. Let the radius of the circle be r. Using $s = r\theta$, we have $5 = r\theta$ and so $r = 5/\theta$. From the Law of Cosines we get $4^2 = r^2 + r^2 - 2 \cdot r \cdot r \cdot \cos\theta \quad \Leftrightarrow \quad 16 = 2r^2(1-\cos\theta) = 2(5/\theta)^2(1-\cos\theta)$. Multiplying by θ^2 gives $16\theta^2 = 50(1-\cos\theta)$, so we take $f(\theta) = 16\theta^2 + 50\cos\theta - 50$ and $f'(\theta) = 32\theta - 50\sin\theta$. The formula for Newton's method is $\theta_{n+1} = \theta_n - \dfrac{16\theta_n^2 + 50\cos\theta_n - 50}{32\theta_n - 50\sin\theta_n}$. From the graph of f, we can use $\theta_1 = 2.2$, giving us $\theta_2 \approx 2.2662$, $\theta_3 \approx 2.2622 \approx \theta_4$. So correct to four decimal places, the angle is 2.2622 radians $\approx 130°$.

41. In this case, $A = 18{,}000$, $R = 375$, and $n = 5(12) = 60$. So the formula $A = \dfrac{R}{i}[1-(1+i)^{-n}]$ becomes

$18{,}000 = \dfrac{375}{x}[1-(1+x)^{-60}] \quad \Leftrightarrow \quad 48x = 1-(1+x)^{-60}$ [multiply each term by $(1+x)^{60}$] $\Leftrightarrow$

$48x(1+x)^{60} - (1+x)^{60} + 1 = 0$. Let the LHS be called $f(x)$, so that

$$\begin{aligned} f'(x) &= 48x(60)(1+x)^{59} + 48(1+x)^{60} - 60(1+x)^{59} \\ &= 12(1+x)^{59}[4x(60) + 4(1+x) - 5] = 12(1+x)^{59}(244x - 1) \end{aligned}$$

$x_{n+1} = x_n - \dfrac{48x_n(1+x_n)^{60} - (1+x_n)^{60} + 1}{12(1+x_n)^{59}(244x_n - 1)}$. An interest rate of 1% per month seems like a reasonable estimate for $x = i$. So let $x_1 = 1\% = 0.01$, and we get $x_2 \approx 0.0082202$, $x_3 \approx 0.0076802$, $x_4 \approx 0.0076291$, $x_5 \approx 0.0076286 \approx x_6$. Thus, the dealer is charging a monthly interest rate of 0.76286% (or 9.55% per year, compounded monthly).

42. (a) $p(x) = x^5 - (2+r)x^4 + (1+2r)x^3 - (1-r)x^2 + 2(1-r)x + r - 1 \quad \Rightarrow$

$p'(x) = 5x^4 - 4(2+r)x^3 + 3(1+2r)x^2 - 2(1-r)x + 2(1-r)$. So we use

$$x_{n+1} = x_n - \frac{x_n^5 - (2+r)x_n^4 + (1+2r)x_n^3 - (1-r)x_n^2 + 2(1-r)x_n + r - 1}{5x_n^4 - 4(2+r)x_n^3 + 3(1+2r)x_n^2 - 2(1-r)x_n + 2(1-r)}.$$

We substitute in the value $r \approx 3.04042 \times 10^{-6}$ in order to evaluate the approximations numerically. The libration point L_1 is slightly less than 1 AU from the sun, so we take $x_1 = 0.95$ as our first approximation, and get $x_2 \approx 0.96682$,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x_3 \approx 0.97770$, $x_4 \approx 0.98451$, $x_5 \approx 0.98830$, $x_6 \approx 0.98976$, $x_7 \approx 0.98998$, $x_8 \approx 0.98999 \approx x_9$. So, to five decimal places, L_1 is located 0.98999 AU from the sun (or 0.01001 AU from the earth).

(b) In this case we use Newton's method with the function

$p(x) - 2rx^2 = x^5 - (2+r)x^4 + (1+2r)x^3 - (1+r)x^2 + 2(1-r)x + r - 1 \quad \Rightarrow$

$\left[p(x) - 2rx^2\right]' = 5x^4 - 4(2+r)x^3 + 3(1+2r)x^2 - 2(1+r)x + 2(1-r)$. So

$x_{n+1} = x_n - \dfrac{x_n^5 - (2+r)x_n^4 + (1+2r)x_n^3 - (1+r)x_n^2 + 2(1-r)x_n + r - 1}{5x_n^4 - 4(2+r)x_n^3 + 3(1+2r)x_n^2 - 2(1+r)x_n + 2(1-r)}$. Again, we substitute

$r \approx 3.04042 \times 10^{-6}$. L_2 is slightly more than 1 AU from the sun and, judging from the result of part (a), probably less than 0.02 AU from earth. So we take $x_1 = 1.02$ and get $x_2 \approx 1.01422$, $x_3 \approx 1.01118$, $x_4 \approx 1.01018$, $x_5 \approx 1.01008 \approx x_6$. So, to five decimal places, L_2 is located 1.01008 AU from the sun (or 0.01008 AU from the earth).

4.9 Antiderivatives

1. (a) $f(x) = 6 \quad \Rightarrow \quad F(x) = 6x$ is an antiderivative.

(b) $g(t) = 3t^2 \quad \Rightarrow \quad G(t) = 3\,\dfrac{t^{2+1}}{2+1} = t^3$ is an antiderivative.

2. (a) $f(x) = 2x = 2x^1 \quad \Rightarrow \quad F(x) = 2\,\dfrac{x^{1+1}}{1+1} = x^2$ is an antiderivative.

(b) $g(x) = -1/x^2 = -x^{-2} \quad \Rightarrow \quad G(x) = -\dfrac{x^{-2+1}}{-2+1} = x^{-1} = 1/x$ is an antiderivative.

3. (a) $h(q) = \cos q \quad \Rightarrow \quad H(q) = \sin q$ is an antiderivative.

(b) $f(x) = e^x \quad \Rightarrow \quad F(x) = e^x$ is an antiderivative.

4. (a) $g(t) = 1/t \quad \Rightarrow \quad G(t) = \ln|\,t\,|$ is an antiderivative.

(b) $r(\theta) = \sec^2\theta \quad \Rightarrow \quad R(\theta) = \tan\theta$ is an antiderivative.

5. $f(x) = 4x + 7 = 4x^1 + 7 \quad \Rightarrow \quad F(x) = 4\dfrac{x^{1+1}}{1+1} + 7x + C = 2x^2 + 7x + C$

Check: $F'(x) = 2(2x) + 7 + 0 = 4x + 7 = f(x)$

6. $f(x) = x^2 - 3x + 2 \quad \Rightarrow \quad F(x) = \dfrac{x^3}{3} - 3\dfrac{x^2}{2} + 2x + C = \frac{1}{3}x^3 - \frac{3}{2}x^2 + 2x + C$

Check: $F'(x) = \frac{1}{3}(3x^2) - \frac{3}{2}(2x) + 2 + 0 = x^2 - 3x + 2 = f(x)$

7. $f(x) = 2x^3 - \frac{2}{3}x^2 + 5x \quad \Rightarrow \quad F(x) = 2\,\dfrac{x^{3+1}}{3+1} - \dfrac{2}{3}\dfrac{x^{2+1}}{2+1} + 5\,\dfrac{x^{1+1}}{1+1} = \frac{1}{2}x^4 - \frac{2}{9}x^3 + \frac{5}{2}x^2 + C$

Check: $F'(x) = \frac{1}{2}(4x^3) - \frac{2}{9}(3x^2) + \frac{5}{2}(2x) + 0 = 2x^3 - \frac{2}{3}x^2 + 5x = f(x)$

8. $f(x) = 6x^5 - 8x^4 - 9x^2 \quad \Rightarrow \quad F(x) = 6\,\dfrac{x^6}{6} - 8\,\dfrac{x^5}{5} - 9\,\dfrac{x^3}{3} + C = x^6 - \frac{8}{5}x^5 - 3x^3 + C$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $f(x) = x(12x+8) = 12x^2 + 8x \quad \Rightarrow \quad F(x) = 12\,\dfrac{x^3}{3} + 8\,\dfrac{x^2}{2} + C = 4x^3 + 4x^2 + C$

10. $f(x) = (x-5)^2 = x^2 - 10x + 25 \quad \Rightarrow \quad F(x) = \dfrac{x^3}{3} - 10\dfrac{x^2}{2} + 25x + C = \frac{1}{3}x^3 - 5x^2 + 25x + C$

11. $g(x) = 4x^{-2/3} - 2x^{5/3} \quad \Rightarrow \quad G(x) = 4(3x^{1/3}) - 2\left(\frac{3}{8}x^{8/3}\right) + C = 12x^{1/3} - \frac{3}{4}x^{8/3} + C$

12. $h(z) = 3z^{0.8} + z^{-2.5} \quad \Rightarrow \quad H(z) = 3\,\dfrac{z^{1.8}}{1.8} + \dfrac{z^{-1.5}}{-1.5} = \dfrac{5}{3}\,z^{1.8} - \dfrac{2}{3}\,z^{-1.5} + C$

13. $f(x) = 3\sqrt{x} - 2\sqrt[3]{x} = 3x^{1/2} - 2x^{1/3} \quad \Rightarrow \quad F(x) = 3\left(\frac{2}{3}x^{3/2}\right) - 2\left(\frac{3}{4}x^{4/3}\right) + C = 2x^{3/2} - \frac{3}{2}x^{4/3} + C$

14. $g(x) = \sqrt{x}\,(2 - x + 6x^2) = 2x^{1/2} - x^{3/2} + 6x^{5/2} \quad \Rightarrow$

$G(x) = 2\,\dfrac{x^{3/2}}{3/2} - \dfrac{x^{5/2}}{5/2} + 6\,\dfrac{x^{7/2}}{7/2} + C = \dfrac{4}{3}\,x^{3/2} - \dfrac{2}{5}\,x^{5/2} + \dfrac{12}{7}\,x^{7/2} + C$

15. $f(t) = \dfrac{2t - 4 + 3\sqrt{t}}{\sqrt{t}} = 2t^{1/2} - 4t^{-1/2} + 3 \quad \Rightarrow \quad F(t) = 2\,\dfrac{t^{3/2}}{3/2} - 4\,\dfrac{t^{1/2}}{1/2} + 3t + C = \dfrac{4}{3}\,t^{3/2} - 8t^{1/2} + 3t + C$

16. $f(x) = \sqrt[4]{5} + \sqrt[4]{x} = \sqrt[4]{5} + x^{1/4} \quad \Rightarrow \quad F(x) = \sqrt[4]{5}\,x + \dfrac{x^{5/4}}{5/4} + C = \sqrt[4]{5}\,x + \dfrac{4}{5}\,x^{5/4} + C$

17. $f(x) = \dfrac{2}{5x} - \dfrac{3}{x^2} = \dfrac{2}{5}\left(\dfrac{1}{x}\right) - 3x^{-2}$ has domain $(-\infty, 0) \cup (0, \infty)$, so $F(x) = \begin{cases} \dfrac{2}{5}\ln(-x) + \dfrac{3}{x} + C_2 & \text{if } x < 0 \\ \dfrac{2}{5}\ln x + \dfrac{3}{x} + C_1 & \text{if } x > 0 \end{cases}$

See Example 1(b) for a similar problem.

18. $f(x) = \dfrac{5x^2 - 6x + 4}{x^2},\ x > 0 \quad \Rightarrow \quad f(x) = 5 - 6x^{-1} + 4x^{-2},\ x > 0 \quad \Rightarrow$

$F(x) = 5x - 6\ln x + 4(-x^{-1}) + C = 5x - 6\ln x - \dfrac{4}{x} + C,\ x > 0$

19. $g(t) = 7e^t - e^3 \quad \Rightarrow \quad G(t) = 7e^t - e^3 t + C$

20. $f(x) = \dfrac{10}{x^6} - 2e^x + 3$ has domain $(-\infty, 0) \cup (0, \infty)$, so $F(x) = \begin{cases} -\dfrac{2}{x^5} - 2e^x + 3x + C_1 & \text{if } x < 0 \\ -\dfrac{2}{x^5} - 2e^x + 3x + C_2 & \text{if } x > 0 \end{cases}$

21. $f(\theta) = 2\sin\theta - 3\sec\theta\tan\theta \quad \Rightarrow \quad F(\theta) = -2\cos\theta - 3\sec\theta + C_n$ on the interval $\left(n\pi - \dfrac{\pi}{2}, n\pi + \dfrac{\pi}{2}\right)$.

22. $h(x) = \sec^2 x + \pi\cos x \quad \Rightarrow \quad H(x) = \tan x + \pi\sin x + C_n$ on the interval $\left(n\pi - \dfrac{\pi}{2}, n\pi + \dfrac{\pi}{2}\right)$.

23. $f(r) = \dfrac{4}{1+r^2} - \sqrt[5]{r^4} = 4\left(\dfrac{1}{1+r^2}\right) - r^{4/5} \quad \Rightarrow \quad F(r) = 4\arctan r - \dfrac{5}{9}r^{9/5} + C$

24. $g(v) = 2\cos v - \dfrac{3}{\sqrt{1-v^2}} \quad \Rightarrow \quad G(v) = 2\sin v - 3\sin^{-1} v + C$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $f(x) = 2^x + 4\sinh x \quad \Rightarrow \quad F(x) = \dfrac{2^x}{\ln 2} + 4\cosh x + C$

26. $f(x) = \dfrac{2x^2+5}{x^2+1} = \dfrac{2(x^2+1)+3}{x^2+1} = 2 + \dfrac{3}{x^2+1} \quad \Rightarrow \quad F(x) = 2x + 3\tan^{-1} x + C$

27. $f(x) = 2e^x - 6x \quad \Rightarrow \quad F(x) = 2e^x - 6 \cdot \dfrac{x^2}{2} + C = 2e^x - 3x^2 + C.$

$F(0) = 1 \quad \Rightarrow \quad 2e^0 - 3(0)^2 + C = 1 \quad \Rightarrow \quad C = -1$, so $F(x) = 2e^x - 3x^2 - 1$.

The graph confirms our answer since $f(x) = 0$ when F has a local maximum and when F has a local minimum, f is positive when F is increasing, and f is negative when F is decreasing.

28. $f(x) = 4 - 3\left(1+x^2\right)^{-1} = 4 - \dfrac{3}{1+x^2} \quad \Rightarrow \quad F(x) = 4x - 3\tan^{-1} x + C.$

$F(1) = 0 \quad \Rightarrow \quad 4 - 3\left(\frac{\pi}{4}\right) + C = 0 \quad \Rightarrow \quad C = \frac{3\pi}{4} - 4$, so

$F(x) = 4x - 3\tan^{-1} x + \frac{3\pi}{4} - 4$. Note that f is positive and F is increasing on $\mathbb{R}$.

Also, f has smaller values where the slopes of the tangent lines of F are smaller.

29. $f''(x) = 24x \quad \Rightarrow \quad f'(x) = 24\left(\dfrac{x^2}{2}\right) + C = 12x^2 + C \quad \Rightarrow \quad f(x) = 12\left(\dfrac{x^3}{3}\right) + Cx + D = 4x^3 + Cx + D$

30. $f''(t) = t^2 - 4 \quad \Rightarrow \quad f'(t) = \dfrac{t^3}{3} - 4t + C \quad \Rightarrow \quad f(t) = \dfrac{1}{3}\left(\dfrac{t^4}{4}\right) - 4\left(\dfrac{t^2}{2}\right) + Ct + D = \dfrac{1}{12}t^4 - 2t^2 + Ct + D$

31. $f''(x) = 4x^3 + 24x - 1 \quad \Rightarrow \quad f'(x) = 4\left(\dfrac{x^4}{4}\right) + 24\left(\dfrac{x^2}{2}\right) - x + C = x^4 + 12x^2 - x + C \quad \Rightarrow$

$f(x) = \dfrac{x^5}{5} + 12\left(\dfrac{x^3}{3}\right) - \dfrac{x^2}{2} + Cx + D = \dfrac{1}{5}x^5 + 4x^3 - \dfrac{1}{2}x^2 + Cx + D$

32. $f''(x) = 6x - x^4 + 3x^5 \quad \Rightarrow \quad f'(x) = 6\left(\dfrac{x^2}{2}\right) - \dfrac{x^5}{5} + 3\left(\dfrac{x^6}{6}\right) + C = 3x^2 - \dfrac{1}{5}x^5 + \dfrac{1}{2}x^6 + C \quad \Rightarrow$

$f(x) = 3\left(\dfrac{x^3}{3}\right) - \dfrac{1}{5}\left(\dfrac{x^6}{6}\right) + \dfrac{1}{2}\left(\dfrac{x^7}{7}\right) + Cx + D = x^3 - \dfrac{1}{30}x^6 + \dfrac{1}{14}x^7 + Cx + D$

33. $f''(x) = 2x + 3e^x \quad \Rightarrow \quad f'(x) = x^2 + 3e^x + C \quad \Rightarrow \quad f(x) = \frac{1}{3}x^3 + 3e^x + Cx + D$

34. $f''(x) = 1/x^2 = x^{-2} \quad \Rightarrow \quad f'(x) = \begin{cases} -1/x + C_1 & \text{if } x < 0 \\ -1/x + C_2 & \text{if } x > 0 \end{cases} \quad \Rightarrow \quad f(x) = \begin{cases} -\ln(-x) + C_1x + D_1 & \text{if } x < 0 \\ -\ln x + C_2x + D_2 & \text{if } x > 0 \end{cases}$

35. $f'''(t) = 12 + \sin t \quad \Rightarrow \quad f''(t) = 12t - \cos t + C_1 \quad \Rightarrow \quad f'(t) = 6t^2 - \sin t + C_1t + D \quad \Rightarrow$

$f(t) = 2t^3 + \cos t + Ct^2 + Dt + E$, where $C = \frac{1}{2}C_1$.

36. $f'''(t) = \sqrt{t} - 2\cos t = t^{1/2} - 2\cos t \quad \Rightarrow \quad f''(t) = \frac{2}{3}t^{3/2} - 2\sin t + C_1 \quad \Rightarrow \quad f'(t) = \frac{4}{15}t^{5/2} + 2\cos t + C_1t + D \quad \Rightarrow$

$f(t) = \frac{8}{105}t^{7/2} + 2\sin t + Ct^2 + Dt + E$, where $C = \frac{1}{2}C_1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. $f'(x) = 8x^3 + \dfrac{1}{x}$, $x > 0 \quad \Rightarrow \quad f(x) = 2x^4 + \ln x + C$. $\quad f(1) = 2(1)^4 + \ln 1 + C = 2 + C$ and $f(1) = -3 \quad \Rightarrow$

$C = -5$, so $f(x) = 2x^4 + \ln x - 5$.

38. $f'(x) = \sqrt{x} - 2 \quad \Rightarrow \quad f(x) = \frac{2}{3}x^{3/2} - 2x + C$.

$f(9) = \frac{2}{3}(9)^{3/2} - 2(9) + C = 18 - 18 + C = C$ and $f(9) = 4 \quad \Rightarrow \quad C = 4$, so $f(x) = \frac{2}{3}x^{3/2} - 2x + 4$.

39. $f'(t) = \dfrac{4}{1+t^2} \quad \Rightarrow \quad f(t) = 4\arctan t + C$. $\quad f(1) = 4\left(\dfrac{\pi}{4}\right) + C$ and $f(1) = 0 \quad \Rightarrow \quad \pi + C = 0 \quad \Rightarrow \quad C = -\pi$,

so $f(t) = 4\arctan t - \pi$.

40. $f'(t) = t + \dfrac{1}{t^3}$, $t > 0 \quad \Rightarrow \quad f(t) = \dfrac{1}{2}t^2 - \dfrac{1}{2t^2} + C$. $\quad f(1) = \dfrac{1}{2} - \dfrac{1}{2} + C$ and $f(1) = 6 \quad \Rightarrow \quad C = 6$, so

$f(t) = \dfrac{1}{2}t^2 - \dfrac{1}{2t^2} + 6$.

41. $f'(x) = 5x^{2/3} \quad \Rightarrow \quad f(x) = 5\left(\frac{3}{5}x^{5/3}\right) + C = 3x^{5/3} + C$.

$f(8) = 3 \cdot 32 + C$ and $f(8) = 21 \quad \Rightarrow \quad 96 + C = 21 \quad \Rightarrow \quad C = -75$, so $f(x) = 3x^{5/3} - 75$.

42. $f'(x) = \dfrac{x+1}{\sqrt{x}} = x^{1/2} + x^{-1/2} \quad \Rightarrow \quad f(x) = \frac{2}{3}x^{3/2} + 2x^{1/2} + C$.

$f(1) = \frac{2}{3} + 2 + C = \frac{8}{3} + C$ and $f(1) = 5 \quad \Rightarrow \quad C = 5 - \frac{8}{3} = \frac{7}{3}$, so $f(x) = \frac{2}{3}x^{3/2} + 2\sqrt{x} + \frac{7}{3}$.

43. $f'(t) = \sec t\,(\sec t + \tan t) = \sec^2 t + \sec t\,\tan t$, $-\frac{\pi}{2} < t < \frac{\pi}{2} \quad \Rightarrow \quad f(t) = \tan t + \sec t + C$. $\quad f\left(\frac{\pi}{4}\right) = 1 + \sqrt{2} + C$

and $f\left(\frac{\pi}{4}\right) = -1 \quad \Rightarrow \quad 1 + \sqrt{2} + C = -1 \quad \Rightarrow \quad C = -2 - \sqrt{2}$, so $f(t) = \tan t + \sec t - 2 - \sqrt{2}$.

Note: The fact that f is defined and continuous on $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ means that we have only one constant of integration.

44. $f'(t) = 3^t - \dfrac{3}{t} \quad \Rightarrow \quad f(t) = \begin{cases} 3^t/\ln 3 - 3\ln(-t) + C & \text{if} \quad t < 0 \\ 3^t/\ln 3 - 3\ln t + D & \text{if} \quad t > 0 \end{cases}$

$f(-1) = \dfrac{1}{3\ln 3} - 3\ln 1 + C$ and $f(-1) = 1 \quad \Rightarrow \quad C = 1 - \dfrac{1}{3\ln 3}$.

$f(1) = \dfrac{3}{\ln 3} - 3\ln 1 + D$ and $f(1) = 2 \quad \Rightarrow \quad D = 2 - \dfrac{3}{\ln 3}$.

Thus, $f(t) = \begin{cases} 3^t/\ln 3 - 3\ln(-t) + 1 - 1/(3\ln 3) & \text{if} \quad t < 0 \\ 3^t/\ln 3 - 3\ln t + 2 - 3/\ln 3 & \text{if} \quad t > 0 \end{cases}$

45. $f''(x) = -2 + 12x - 12x^2 \quad \Rightarrow \quad f'(x) = -2x + 6x^2 - 4x^3 + C$. $f'(0) = C$ and $f'(0) = 12 \quad \Rightarrow \quad C = 12$, so

$f'(x) = -2x + 6x^2 - 4x^3 + 12$ and hence, $f(x) = -x^2 + 2x^3 - x^4 + 12x + D$. $f(0) = D$ and $f(0) = 4 \quad \Rightarrow \quad D = 4$,

so $f(x) = -x^2 + 2x^3 - x^4 + 12x + 4$.

46. $f''(x) = 8x^3 + 5 \quad \Rightarrow \quad f'(x) = 2x^4 + 5x + C$. $f'(1) = 2 + 5 + C$ and $f'(1) = 8 \quad \Rightarrow \quad C = 1$, so

$f'(x) = 2x^4 + 5x + 1$. $f(x) = \frac{2}{5}x^5 + \frac{5}{2}x^2 + x + D$. $f(1) = \frac{2}{5} + \frac{5}{2} + 1 + D = D + \frac{39}{10}$ and $f(1) = 0 \quad \Rightarrow \quad D = -\frac{39}{10}$,

so $f(x) = \frac{2}{5}x^5 + \frac{5}{2}x^2 + x - \frac{39}{10}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $f''(\theta) = \sin\theta + \cos\theta \quad \Rightarrow \quad f'(\theta) = -\cos\theta + \sin\theta + C$. $f'(0) = -1 + C$ and $f'(0) = 4 \quad \Rightarrow \quad C = 5$, so $f'(\theta) = -\cos\theta + \sin\theta + 5$ and hence, $f(\theta) = -\sin\theta - \cos\theta + 5\theta + D$. $f(0) = -1 + D$ and $f(0) = 3 \quad \Rightarrow \quad D = 4$, so $f(\theta) = -\sin\theta - \cos\theta + 5\theta + 4$.

48. $f''(t) = t^2 + \dfrac{1}{t^2} = t^2 + t^{-2},\ t > 0 \quad \Rightarrow \quad f'(t) = \frac{1}{3}t^3 - \dfrac{1}{t} + C$. $f'(1) = \frac{1}{3} - 1 + C$ and $f'(1) = 2 \quad \Rightarrow$ $C - \frac{2}{3} = 2 \quad \Rightarrow \quad C = \frac{8}{3}$, so $f'(t) = \frac{1}{3}t^3 - \dfrac{1}{t} + \frac{8}{3}$ and hence, $f(t) = \frac{1}{12}t^4 - \ln t + \frac{8}{3}t + D$. $f(2) = \frac{4}{3} - \ln 2 + \frac{16}{3} + D$ and $f(2) = 3 \quad \Rightarrow \quad \frac{20}{3} - \ln 2 + D = 3 \quad \Rightarrow \quad D = \ln 2 - \frac{11}{3}$, so $f(t) = \frac{1}{12}t^4 - \ln t + \frac{8}{3}t + \ln 2 - \frac{11}{3}$.

49. $f''(x) = 4 + 6x + 24x^2 \quad \Rightarrow \quad f'(x) = 4x + 3x^2 + 8x^3 + C \quad \Rightarrow \quad f(x) = 2x^2 + x^3 + 2x^4 + Cx + D$. $f(0) = D$ and $f(0) = 3 \quad \Rightarrow \quad D = 3$, so $f(x) = 2x^2 + x^3 + 2x^4 + Cx + 3$. $f(1) = 8 + C$ and $f(1) = 10 \quad \Rightarrow \quad C = 2$, so $f(x) = 2x^2 + x^3 + 2x^4 + 2x + 3$.

50. $f''(x) = x^3 + \sinh x \quad \Rightarrow \quad f'(x) = \frac{1}{4}x^4 + \cosh x + C \quad \Rightarrow \quad f(x) = \frac{1}{20}x^5 + \sinh x + Cx + D$. $f(0) = D$ and $f(0) = 1 \quad \Rightarrow \quad D = 1$, so $f(x) = \frac{1}{20}x^5 + \sinh x + Cx + 1$. $f(2) = \frac{32}{20} + \sinh 2 + 2C + 1$ and $f(2) = 2.6 \quad \Rightarrow$ $\sinh 2 + 2C = 0 \quad \Rightarrow \quad C = -\frac{1}{2}\sinh 2$, so $f(x) = \frac{1}{20}x^5 + \sinh x - \frac{1}{2}(\sinh 2)x + 1$.

51. $f''(x) = e^x - 2\sin x \quad \Rightarrow \quad f'(x) = e^x + 2\cos x + C \quad \Rightarrow \quad f(x) = e^x + 2\sin x + Cx + D$. $f(0) = 1 + 0 + D$ and $f(0) = 3 \quad \Rightarrow \quad D = 2$, so $f(x) = e^x + 2\sin x + Cx + 2$. $f\left(\frac{\pi}{2}\right) = e^{\pi/2} + 2 + \frac{\pi}{2}C + 2$ and $f\left(\frac{\pi}{2}\right) = 0 \quad \Rightarrow \quad e^{\pi/2} + 4 + \frac{\pi}{2}C = 0 \quad \Rightarrow \quad \frac{\pi}{2}C = -e^{\pi/2} - 4 \quad \Rightarrow \quad C = -\frac{2}{\pi}(e^{\pi/2} + 4)$, so $f(x) = e^x + 2\sin x - \frac{2}{\pi}(e^{\pi/2} + 4)x + 2$.

52. $f''(t) = \sqrt[3]{t} - \cos t = t^{1/3} - \cos t \quad \Rightarrow \quad f'(t) = \frac{3}{4}t^{4/3} - \sin t + C \quad \Rightarrow \quad f(t) = \frac{9}{28}t^{7/3} + \cos t + Ct + D$. $f(0) = 0 + 1 + 0 + D$ and $f(0) = 2 \quad \Rightarrow \quad D = 1$, so $f(t) = \frac{9}{28}t^{7/3} + \cos t + Ct + 1$. $f(1) = \frac{9}{28} + \cos 1 + C + 1$ and $f(1) = 2 \quad \Rightarrow \quad C = 2 - \frac{9}{28} - \cos 1 - 1 = \frac{19}{28} - \cos 1$, so $f(t) = \frac{9}{28}t^{7/3} + \cos t + \left(\frac{19}{28} - \cos 1\right)t + 1$.

53. $f''(x) = x^{-2}, x > 0 \quad \Rightarrow \quad f'(x) = -1/x + C \quad \Rightarrow \quad f(x) = -\ln|x| + Cx + D = -\ln x + Cx + D$ [since $x > 0$]. $f(1) = 0 \quad \Rightarrow \quad C + D = 0$ and $f(2) = 0 \quad \Rightarrow \quad -\ln 2 + 2C + D = 0 \quad \Rightarrow \quad -\ln 2 + 2C - C = 0$ [since $D = -C$] $\Rightarrow$ $-\ln 2 + C = 0 \quad \Rightarrow \quad C = \ln 2$ and $D = -\ln 2$. So $f(x) = -\ln x + (\ln 2)x - \ln 2$.

54. $f'''(x) = \cos x \quad \Rightarrow \quad f''(x) = \sin x + C$. $f''(0) = C$ and $f''(0) = 3 \quad \Rightarrow \quad C = 3$. $f''(x) = \sin x + 3 \quad \Rightarrow$ $f'(x) = -\cos x + 3x + D$. $f'(0) = -1 + D$ and $f'(0) = 2 \quad \Rightarrow \quad D = 3$. $f'(x) = -\cos x + 3x + 3 \quad \Rightarrow$ $f(x) = -\sin x + \frac{3}{2}x^2 + 3x + E$. $f(0) = E$ and $f(0) = 1 \quad \Rightarrow \quad E = 1$. Thus, $f(x) = -\sin x + \frac{3}{2}x^2 + 3x + 1$.

55. "The slope of its tangent line at $(x, f(x))$ is $3 - 4x$" means that $f'(x) = 3 - 4x$, so $f(x) = 3x - 2x^2 + C$. "The graph of f passes through the point $(2, 5)$" means that $f(2) = 5$, but $f(2) = 3(2) - 2(2)^2 + C$, so $5 = 6 - 8 + C \quad \Rightarrow$ $C = 7$. Thus, $f(x) = 3x - 2x^2 + 7$ and $f(1) = 3 - 2 + 7 = 8$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

56. $f'(x) = x^3 \quad \Rightarrow \quad f(x) = \frac{1}{4}x^4 + C$. $x + y = 0 \quad \Rightarrow \quad y = -x \quad \Rightarrow \quad m = -1$. Now $m = f'(x) \quad \Rightarrow \quad -1 = x^3 \quad \Rightarrow$ $x = -1 \quad \Rightarrow \quad y = 1$ (from the equation of the tangent line), so $(-1, 1)$ is a point on the graph of f. From f, $1 = \frac{1}{4}(-1)^4 + C \quad \Rightarrow \quad C = \frac{3}{4}$. Therefore, the function is $f(x) = \frac{1}{4}x^4 + \frac{3}{4}$.

57. b is the antiderivative of f. For small x, f is negative, so the graph of its antiderivative must be decreasing. But both a and c are increasing for small x, so only b can be f's antiderivative. Also, f is positive where b is increasing, which supports our conclusion.

58. We know right away that c cannot be f's antiderivative, since the slope of c is not zero at the x-value where $f = 0$. Now f is positive when a is increasing and negative when a is decreasing, so a is the antiderivative of f.

59.

The graph of F must start at $(0, 1)$. Where the given graph, $y = f(x)$, has a local minimum or maximum, the graph of F will have an inflection point.

Where f is negative (positive), F is decreasing (increasing).

Where f changes from negative to positive, F will have a minimum.

Where f changes from positive to negative, F will have a maximum.

Where f is decreasing (increasing), F is concave downward (upward).

60.

Where v is positive (negative), s is increasing (decreasing).

Where v is increasing (decreasing), s is concave upward (downward).

Where v is horizontal (a steady velocity), s is linear.

61.

$$f'(x) = \begin{cases} 2 & \text{if } 0 \le x < 1 \\ 1 & \text{if } 1 < x < 2 \\ -1 & \text{if } 2 < x < 3 \end{cases} \quad \Rightarrow \quad f(x) = \begin{cases} 2x + C & \text{if } 0 \le x < 1 \\ x + D & \text{if } 1 < x < 2 \\ -x + E & \text{if } 2 < x < 3 \end{cases}$$

$f(0) = -1 \quad \Rightarrow \quad 2(0) + C = -1 \quad \Rightarrow \quad C = -1$. Starting at the point $(0, -1)$ and moving to the right on a line with slope 2 gets us to the point $(1, 1)$.

The slope for $1 < x < 2$ is 1, so we get to the point $(2, 2)$. Here we have used the fact that f is continuous. We can include the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

point $x = 1$ on either the first or the second part of f. The line connecting $(1, 1)$ to $(2, 2)$ is $y = x$, so $D = 0$. The slope for $2 < x < 3$ is -1, so we get to $(3, 1)$. $f(2) = 2 \quad \Rightarrow \quad -2 + E = 2 \quad \Rightarrow \quad E = 4$. Thus,

$$f(x) = \begin{cases} 2x - 1 & \text{if } 0 \le x \le 1 \\ x & \text{if } 1 < x < 2 \\ -x + 4 & \text{if } 2 \le x < 3 \end{cases}$$

Note that $f'(x)$ does not exist at $x = 1$, 2, or 3.

62. (a)

(b) Since $F(0) = 1$, we can start our graph at $(0, 1)$. f has a minimum at about $x = 0.5$, so its derivative is zero there. f is decreasing on $(0, 0.5)$, so its derivative is negative and hence, F is CD on $(0, 0.5)$ and has an IP at $x \approx 0.5$. On $(0.5, 2.2)$, f is negative and increasing (f' is positive), so F is decreasing and CU. On $(2.2, \infty)$, f is positive and increasing, so F is increasing and CU.

(c) $f(x) = 2x - 3\sqrt{x} \quad \Rightarrow \quad F(x) = x^2 - 3 \cdot \frac{2}{3}x^{3/2} + C$.

$F(0) = C$ and $F(0) = 1 \quad \Rightarrow \quad C = 1$, so $F(x) = x^2 - 2x^{3/2} + 1$.

(d)

63. $f(x) = \dfrac{\sin x}{1 + x^2}, \ -2\pi \le x \le 2\pi$

Note that the graph of f is one of an odd function, so the graph of F will be one of an even function.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. $f(x) = \sqrt{x^4 - 2x^2 + 2} - 2,\ -3 \le x \le 3$

Note that the graph of f is one of an even function, so the graph of F will be one of an odd function.

65. $v(t) = s'(t) = 2\cos t + 4\sin t \quad\Rightarrow\quad s(t) = 2\sin t - 4\cos t + C.\quad s(0) = -4 + C$ and $s(0) = 3 \quad\Rightarrow\quad C = 7$, so $s(t) = 2\sin t - 4\cos t + 7$.

66. $v(t) = s'(t) = t^2 - 3\sqrt{t} = t^2 - 3t^{1/2} \quad\Rightarrow\quad s(t) = \frac{1}{3}t^3 - 2t^{3/2} + C.\quad s(4) = \frac{64}{3} - 16 + C$ and $s(4) = 8 \quad\Rightarrow$ $C = 8 - \frac{64}{3} + 16 = \frac{8}{3}$, so $s(t) = \frac{1}{3}t^3 - 2t^{3/2} + \frac{8}{3}$.

67. $a(t) = v'(t) = 2t + 1 \quad\Rightarrow\quad v(t) = t^2 + t + C.\quad v(0) = C$ and $v(0) = -2 \quad\Rightarrow\quad C = -2$, so $v(t) = t^2 + t - 2$ and $s(t) = \frac{1}{3}t^3 + \frac{1}{2}t^2 - 2t + D.\quad s(0) = D$ and $s(0) = 3 \quad\Rightarrow\quad D = 3$, so $s(t) = \frac{1}{3}t^3 + \frac{1}{2}t^2 - 2t + 3$.

68. $a(t) = v'(t) = 3\cos t - 2\sin t \quad\Rightarrow\quad v(t) = 3\sin t + 2\cos t + C.\quad v(0) = 2 + C$ and $v(0) = 4 \quad\Rightarrow\quad C = 2$, so $v(t) = 3\sin t + 2\cos t + 2$ and $s(t) = -3\cos t + 2\sin t + 2t + D.\quad s(0) = -3 + D$ and $s(0) = 0 \quad\Rightarrow\quad D = 3$, so $s(t) = -3\cos t + 2\sin t + 2t + 3$.

69. $a(t) = v'(t) = \sin t - \cos t \quad\Rightarrow\quad v(t) = s'(t) = -\cos t - \sin t + C \quad\Rightarrow\quad s(t) = -\sin t + \cos t + Ct + D$.

$s(0) = 1 + D = 0$ and $s(\pi) = -1 + C\pi + D = 6 \quad\Rightarrow\quad D = -1$ and $C = \dfrac{8}{\pi}$. Thus, $s(t) = -\sin t + \cos t + \dfrac{8}{\pi}t - 1$.

70. $a(t) = t^2 - 4t + 6 \quad\Rightarrow\quad v(t) = \frac{1}{3}t^3 - 2t^2 + 6t + C \quad\Rightarrow\quad s(t) = \frac{1}{12}t^4 - \frac{2}{3}t^3 + 3t^2 + Ct + D.\ s(0) = D$ and $s(0) = 0 \quad\Rightarrow\quad D = 0.\ s(1) = \frac{29}{12} + C$ and $s(1) = 20 \quad\Rightarrow\quad C = \frac{211}{12}$. Thus, $s(t) = \frac{1}{12}t^4 - \frac{2}{3}t^3 + 3t^2 + \frac{211}{12}t$.

71. (a) We first observe that since the stone is dropped 450 m above the ground, $v(0) = 0$ and $s(0) = 450$.

$v'(t) = a(t) = -9.8 \quad\Rightarrow\quad v(t) = -9.8t + C$. Now $v(0) = 0 \quad\Rightarrow\quad C = 0$, so $v(t) = -9.8t \quad\Rightarrow$ $s(t) = -4.9t^2 + D$. Last, $s(0) = 450 \quad\Rightarrow\quad D = 450 \quad\Rightarrow\quad s(t) = 450 - 4.9t^2$.

(b) The stone reaches the ground when $s(t) = 0$. $\ 450 - 4.9t^2 = 0 \quad\Rightarrow\quad t^2 = 450/4.9 \quad\Rightarrow\quad t_1 = \sqrt{450/4.9} \approx 9.58$ s.

(c) The velocity with which the stone strikes the ground is $v(t_1) = -9.8\sqrt{450/4.9} \approx -93.9$ m/s.

(d) This is just reworking parts (a) and (b) with $v(0) = -5$. Using $v(t) = -9.8t + C$, $v(0) = -5 \quad\Rightarrow\quad 0 + C = -5 \quad\Rightarrow$ $v(t) = -9.8t - 5$. So $s(t) = -4.9t^2 - 5t + D$ and $s(0) = 450 \quad\Rightarrow\quad D = 450 \quad\Rightarrow\quad s(t) = -4.9t^2 - 5t + 450$.

Solving $s(t) = 0$ by using the quadratic formula gives us $t = \left(5 \pm \sqrt{8845}\right)/(-9.8) \quad\Rightarrow\quad t_1 \approx 9.09$ s.

72. $v'(t) = a(t) = a \quad\Rightarrow\quad v(t) = at + C$ and $v_0 = v(0) = C \quad\Rightarrow\quad v(t) = at + v_0 \quad\Rightarrow$ $s(t) = \frac{1}{2}at^2 + v_0t + D \quad\Rightarrow\quad s_0 = s(0) = D \quad\Rightarrow\quad s(t) = \frac{1}{2}at^2 + v_0t + s_0$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

73. By Exercise 72 with $a = -9.8$, $s(t) = -4.9t^2 + v_0 t + s_0$ and $v(t) = s'(t) = -9.8t + v_0$. So

$[v(t)]^2 = (-9.8t + v_0)^2 = (9.8)^2 t^2 - 19.6v_0 t + v_0^2 = v_0^2 + 96.04t^2 - 19.6v_0 t = v_0^2 - 19.6\left(-4.9t^2 + v_0 t\right)$.

But $-4.9t^2 + v_0 t$ is just $s(t)$ without the s_0 term; that is, $s(t) - s_0$. Thus, $[v(t)]^2 = v_0^2 - 19.6\,[s(t) - s_0]$.

74. For the first ball, $s_1(t) = -16t^2 + 48t + 432$ from Example 7. For the second ball, $a(t) = -32 \quad\Rightarrow\quad v(t) = -32t + C$, but

$v(1) = -32(1) + C = 24 \quad\Rightarrow\quad C = 56$, so $v(t) = -32t + 56 \quad\Rightarrow\quad s(t) = -16t^2 + 56t + D$, but

$s(1) = -16(1)^2 + 56(1) + D = 432 \quad\Rightarrow\quad D = 392$, and $s_2(t) = -16t^2 + 56t + 392$. The balls pass each other

when $s_1(t) = s_2(t) \quad\Rightarrow\quad -16t^2 + 48t + 432 = -16t^2 + 56t + 392 \quad\Leftrightarrow\quad 8t = 40 \quad\Leftrightarrow\quad t = 5$ s.

Another solution: From Exercise 72, we have $s_1(t) = -16t^2 + 48t + 432$ and $s_2(t) = -16t^2 + 24t + 432$.

We now want to solve $s_1(t) = s_2(t-1) \quad\Rightarrow\quad -16t^2 + 48t + 432 = -16(t-1)^2 + 24(t-1) + 432 \quad\Rightarrow$

$48t = 32t - 16 + 24t - 24 \quad\Rightarrow\quad 40 = 8t \quad\Rightarrow\quad t = 5$ s.

75. Using Exercise 72 with $a = -32$, $v_0 = 0$, and $s_0 = h$ (the height of the cliff), we know that the height at time t is

$s(t) = -16t^2 + h$. $v(t) = s'(t) = -32t$ and $v(t) = -120 \quad\Rightarrow\quad -32t = -120 \quad\Rightarrow\quad t = 3.75$, so

$0 = s(3.75) = -16(3.75)^2 + h \quad\Rightarrow\quad h = 16(3.75)^2 = 225$ ft.

76. (a) $EIy'' = mg(L-x) + \frac{1}{2}\rho g(L-x)^2 \quad\Rightarrow\quad EIy' = -\frac{1}{2}mg(L-x)^2 - \frac{1}{6}\rho g(L-x)^3 + C \quad\Rightarrow$

$EIy = \frac{1}{6}mg(L-x)^3 + \frac{1}{24}\rho g(L-x)^4 + Cx + D$. Since the left end of the board is fixed, we must have $y = y' = 0$

when $x = 0$. Thus, $0 = -\frac{1}{2}mgL^2 - \frac{1}{6}\rho gL^3 + C$ and $0 = \frac{1}{6}mgL^3 + \frac{1}{24}\rho gL^4 + D$. It follows that

$EIy = \frac{1}{6}mg(L-x)^3 + \frac{1}{24}\rho g(L-x)^4 + \left(\frac{1}{2}mgL^2 + \frac{1}{6}\rho gL^3\right)x - \left(\frac{1}{6}mgL^3 + \frac{1}{24}\rho gL^4\right)$ and

$f(x) = y = \dfrac{1}{EI}\left[\frac{1}{6}mg(L-x)^3 + \frac{1}{24}\rho g(L-x)^4 + \left(\frac{1}{2}mgL^2 + \frac{1}{6}\rho gL^3\right)x - \left(\frac{1}{6}mgL^3 + \frac{1}{24}\rho gL^4\right)\right]$

(b) $f(L) < 0$, so the end of the board is a *distance* approximately $-f(L)$ below the horizontal. From our result in (a), we calculate

$$-f(L) = \frac{-1}{EI}\left[\tfrac{1}{2}mgL^3 + \tfrac{1}{6}\rho gL^4 - \tfrac{1}{6}mgL^3 - \tfrac{1}{24}\rho gL^4\right] = \frac{-1}{EI}\left(\tfrac{1}{3}mgL^3 + \tfrac{1}{8}\rho gL^4\right) = -\frac{gL^3}{EI}\left(\frac{m}{3} + \frac{\rho L}{8}\right)$$

Note: This is positive because g is negative.

77. Marginal cost $= 1.92 - 0.002x = C'(x) \quad\Rightarrow\quad C(x) = 1.92x - 0.001x^2 + K$. But $C(1) = 1.92 - 0.001 + K = 562 \quad\Rightarrow$

$K = 560.081$. Therefore, $C(x) = 1.92x - 0.001x^2 + 560.081 \quad\Rightarrow\quad C(100) = 742.081$, so the cost of producing

100 items is \$742.08.

78. Let the mass, measured from one end, be $m(x)$. Then $m(0) = 0$ and $\rho = \dfrac{dm}{dx} = x^{-1/2} \quad\Rightarrow\quad m(x) = 2x^{1/2} + C$ and

$m(0) = C = 0$, so $m(x) = 2\sqrt{x}$. Thus, the mass of the 100-centimeter rod is $m(100) = 2\sqrt{100} = 20$ g.

79. Taking the upward direction to be positive we have that for $0 \le t \le 10$ (using the subscript 1 to refer to $0 \le t \le 10$),

$a_1(t) = -(9 - 0.9t) = v_1'(t) \quad\Rightarrow\quad v_1(t) = -9t + 0.45t^2 + v_0$, but $v_1(0) = v_0 = -10 \quad\Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$v_1(t) = -9t + 0.45t^2 - 10 = s_1'(t) \quad \Rightarrow \quad s_1(t) = -\frac{9}{2}t^2 + 0.15t^3 - 10t + s_0$. But $s_1(0) = 500 = s_0 \quad \Rightarrow$ $s_1(t) = -\frac{9}{2}t^2 + 0.15t^3 - 10t + 500$. $s_1(10) = -450 + 150 - 100 + 500 = 100$, so it takes more than 10 seconds for the raindrop to fall. Now for $t > 10$, $a(t) = 0 = v'(t) \quad \Rightarrow$ $v(t) = \text{constant} = v_1(10) = -9(10) + 0.45(10)^2 - 10 = -55 \quad \Rightarrow \quad v(t) = -55$.

At 55 m/s, it will take $100/55 \approx 1.8$ s to fall the last 100 m. Hence, the total time is $10 + \frac{100}{55} = \frac{130}{11} \approx 11.8$ s.

80. $v'(t) = a(t) = -22$. The initial velocity is 50 mi/h $= \frac{50 \cdot 5280}{3600} = \frac{220}{3}$ ft/s, so $v(t) = -22t + \frac{220}{3}$.

The car stops when $v(t) = 0 \quad \Leftrightarrow \quad t = \frac{220}{3 \cdot 22} = \frac{10}{3}$. Since $s(t) = -11t^2 + \frac{220}{3}t$, the distance covered is $s\left(\frac{10}{3}\right) = -11\left(\frac{10}{3}\right)^2 + \frac{220}{3} \cdot \frac{10}{3} = \frac{1100}{9} = 122.\overline{2}$ ft.

81. $a(t) = k$, the initial velocity is 30 mi/h $= 30 \cdot \frac{5280}{3600} = 44$ ft/s, and the final velocity (after 5 seconds) is 50 mi/h $= 50 \cdot \frac{5280}{3600} = \frac{220}{3}$ ft/s. So $v(t) = kt + C$ and $v(0) = 44 \quad \Rightarrow \quad C = 44$. Thus, $v(t) = kt + 44 \quad \Rightarrow$ $v(5) = 5k + 44$. But $v(5) = \frac{220}{3}$, so $5k + 44 = \frac{220}{3} \quad \Rightarrow \quad 5k = \frac{88}{3} \quad \Rightarrow \quad k = \frac{88}{15} \approx 5.87$ ft/s^2.

82. $a(t) = -16 \quad \Rightarrow \quad v(t) = -16t + v_0$ where v_0 is the car's speed (in ft/s) when the brakes were applied. The car stops when $-16t + v_0 = 0 \quad \Leftrightarrow \quad t = \frac{1}{16}v_0$. Now $s(t) = \frac{1}{2}(-16)t^2 + v_0 t = -8t^2 + v_0 t$. The car travels 200 ft in the time that it takes to stop, so $s\left(\frac{1}{16}v_0\right) = 200 \quad \Rightarrow \quad 200 = -8\left(\frac{1}{16}v_0\right)^2 + v_0\left(\frac{1}{16}v_0\right) = \frac{1}{32}v_0^2 \quad \Rightarrow \quad v_0^2 = 32 \cdot 200 = 6400 \quad \Rightarrow$ $v_0 = 80$ ft/s $[54.\overline{54}$ mi/h$]$.

83. Let the acceleration be $a(t) = k$ km/h^2. We have $v(0) = 100$ km/h and we can take the initial position $s(0)$ to be 0. We want the time t_f for which $v(t) = 0$ to satisfy $s(t) < 0.08$ km. In general, $v'(t) = a(t) = k$, so $v(t) = kt + C$, where $C = v(0) = 100$. Now $s'(t) = v(t) = kt + 100$, so $s(t) = \frac{1}{2}kt^2 + 100t + D$, where $D = s(0) = 0$. Thus, $s(t) = \frac{1}{2}kt^2 + 100t$. Since $v(t_f) = 0$, we have $kt_f + 100 = 0$ or $t_f = -100/k$, so

$s(t_f) = \dfrac{1}{2}k\left(-\dfrac{100}{k}\right)^2 + 100\left(-\dfrac{100}{k}\right) = 10{,}000\left(\dfrac{1}{2k} - \dfrac{1}{k}\right) = -\dfrac{5{,}000}{k}$. The condition $s(t_f)$ must satisfy is

$-\dfrac{5{,}000}{k} < 0.08 \quad \Rightarrow \quad -\dfrac{5{,}000}{0.08} > k \quad [k \text{ is negative}] \quad \Rightarrow \quad k < -62{,}500$ km/h^2, or equivalently, $k < -\frac{3125}{648} \approx -4.82$ m/s^2.

84. (a) For $0 \le t \le 3$ we have $a(t) = 60t \quad \Rightarrow \quad v(t) = 30t^2 + C \quad \Rightarrow \quad v(0) = 0 = C \quad \Rightarrow \quad v(t) = 30t^2$, so $s(t) = 10t^3 + C \quad \Rightarrow \quad s(0) = 0 = C \quad \Rightarrow \quad s(t) = 10t^3$. Note that $v(3) = 270$ and $s(3) = 270$.

For $3 < t \le 17$: $a(t) = -g = -32$ ft/s $\quad \Rightarrow \quad v(t) = -32(t-3) + C \quad \Rightarrow \quad v(3) = 270 = C \quad \Rightarrow$ $v(t) = -32(t-3) + 270 \quad \Rightarrow \quad s(t) = -16(t-3)^2 + 270(t-3) + C \quad \Rightarrow \quad s(3) = 270 = C \quad \Rightarrow$ $s(t) = -16(t-3)^2 + 270(t-3) + 270$. Note that $v(17) = -178$ and $s(17) = 914$.

For $17 < t \le 22$: The velocity increases linearly from -178 ft/s to -18 ft/s during this period, so

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\dfrac{\Delta v}{\Delta t} = \dfrac{-18-(-178)}{22-17} = \dfrac{160}{5} = 32$. Thus, $v(t) = 32(t-17) - 178 \quad \Rightarrow$

$s(t) = 16(t-17)^2 - 178(t-17) + 914$ and $s(22) = 424$ ft.

For $t > 22$: $v(t) = -18 \quad \Rightarrow \quad s(t) = -18(t-22) + C$. But $s(22) = 424 = C \quad \Rightarrow \quad s(t) = -18(t-22) + 424$.

Therefore, until the rocket lands, we have

$$v(t) = \begin{cases} 30t^2 & \text{if } 0 \le t \le 3 \\ -32\,(t-3) + 270 & \text{if } 3 < t \le 17 \\ 32(t-17) - 178 & \text{if } 17 < t \le 22 \\ -18 & \text{if } t > 22 \end{cases}$$

and

$$s(t) = \begin{cases} 10t^3 & \text{if } 0 \le t \le 3 \\ -16(t-3)^2 + 270(t-3) + 270 & \text{if } 3 < t \le 17 \\ 16(t-17)^2 - 178\,(t-17) + 914 & \text{if } 17 < t \le 22 \\ -18(t-22) + 424 & \text{if } t > 22 \end{cases}$$

(b) To find the maximum height, set $v(t)$ on $3 < t \le 17$ equal to 0. $-32(t-3) + 270 = 0 \quad \Rightarrow \quad t_1 = 11.4375$ s and the maximum height is $s(t_1) = -16(t_1-3)^2 + 270(t_1-3) + 270 = 1409.0625$ ft.

(c) To find the time to land, set $s(t) = -18(t-22) + 424 = 0$. Then $t - 22 = \frac{424}{18} = 23.\overline{5}$, so $t \approx 45.6$ s.

85. (a) First note that $180 \text{ mi/h} = 180 \times \dfrac{5280}{3600} \text{ ft/s} = 264 \text{ ft/s}$. Then $a(t) = 2.4 \text{ ft/s}^2 \quad \Rightarrow \quad v(t) = 2.4t + C$, but $v(0) = 0 \quad \Rightarrow \quad C = 0$. Now $2.4t = 264$ when $t = \dfrac{264}{2.4} = 110$ s, so it takes 110 s to reach 264 ft/s. Therefore, taking $s(0) = 0$, we have $s(t) = 1.2t^2$, $0 \le t \le 110$. So $s(110) = 14{,}520$ ft. 20 minutes $= 20(60) = 1200$ s, so for $110 \le t \le 1310$ we have $v(t) = 264 \text{ ft/s} \quad \Rightarrow \quad s(1310) = 264(1200) + 14{,}520 = 331{,}320 \text{ ft} = 62.75$ mi.

(b) As in part (a), the train accelerates for 110 s and travels 14,520 ft while doing so. Similarly, it decelerates for 110 s and travels 14,520 ft at the end of its trip. During the remaining $1200 - 2(110) = 980$ s it travels at 264 ft/s, so the distance traveled is $264 \cdot 980 = 258{,}720$ ft. Thus, the total distance is $14{,}520 + 258{,}720 + 14{,}520 = 287{,}760 \text{ ft} = 54.5$ mi.

(c) $60 \text{ mi} = 60(5280) = 316{,}800$ ft. Subtract $2(14{,}520)$ to take care of the speeding up and slowing down, and we have 287,760 ft at 264 ft/s for a trip of $287{,}760/264 = 1090$ s at 180 mi/h. The total time is $1090 + 2(110) = 1310 \text{ s} = 21 \text{ min } 50 \text{ s} = 21.8\overline{3}$ min.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $37.5(60) = 2250$ s. Then $2250 - 2(110) = 2030$ s at maximum speed. $2030(264) + 2(14{,}520) = 564{,}960$ total feet or $564{,}960/5280 = 107$ mi.

4 Review

TRUE-FALSE QUIZ

1. False. For example, take $f(x) = x^3$, then $f'(x) = 3x^2$ and $f'(0) = 0$, but $f(0) = 0$ is not a maximum or minimum; $(0, 0)$ is an inflection point.

2. False. For example, $f(x) = |x|$ has an absolute minimum at 0, but $f'(0)$ does not exist.

3. False. For example, $f(x) = x$ is continuous on $(0, 1)$ but attains neither a maximum nor a minimum value on $(0, 1)$. Don't confuse this with f being continuous on the *closed* interval $[a, b]$, which would make the statement true.

4. True. By the Mean Value Theorem, $f'(c) = \dfrac{f(1) - f(-1)}{1 - (-1)} = \dfrac{0}{2} = 0$. Note that $|c| < 1 \quad \Leftrightarrow \quad c \in (-1, 1)$.

5. True. This is an example of part (b) of the Increasing/Decreasing Test.

6. False. For example, the curve $y = f(x) = 1$ has no inflection points but $f''(c) = 0$ for all c.

7. False. $f'(x) = g'(x) \quad \Rightarrow \quad f(x) = g(x) + C$. For example, if $f(x) = x + 2$ and $g(x) = x + 1$, then $f'(x) = g'(x) = 1$, but $f(x) \neq g(x)$.

8. False. Assume there is a function f such that $f(1) = -2$ and $f(3) = 0$. Then by the Mean Value Theorem there exists a number $c \in (1, 3)$ such that $f'(c) = \dfrac{f(3) - f(1)}{3 - 1} = \dfrac{0 - (-2)}{2} = 1$. But $f'(x) > 1$ for all x, a contradiction.

9. True. The graph of one such function is sketched.

10. False. At any point $(a, f(a))$, we know that $f'(a) < 0$. So since the tangent line at $(a, f(a))$ is not horizontal, it must cross the x-axis—at $x = b$, say. But since $f''(x) > 0$ for all x, the graph of f must lie above all of its tangents; in particular, $f(b) > 0$. But this is a contradiction, since we are given that $f(x) < 0$ for all x.

11. True. Let $x_1 < x_2$ where $x_1, x_2 \in I$. Then $f(x_1) < f(x_2)$ and $g(x_1) < g(x_2)$ [since f and g are increasing on I], so $(f + g)(x_1) = f(x_1) + g(x_1) < f(x_2) + g(x_2) = (f + g)(x_2)$.

12. False. $f(x) = x$ and $g(x) = 2x$ are both increasing on $(0, 1)$, but $f(x) - g(x) = -x$ is not increasing on $(0, 1)$.

13. False. Take $f(x) = x$ and $g(x) = x - 1$. Then both f and g are increasing on $(0, 1)$. But $f(x)\,g(x) = x(x - 1)$ is not increasing on $(0, 1)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. True. Let $x_1 < x_2$ where $x_1, x_2 \in I$. Then $0 < f(x_1) < f(x_2)$ and $0 < g(x_1) < g(x_2)$ [since f and g are both positive and increasing]. Hence, $f(x_1)\,g(x_1) < f(x_2)\,g(x_1) < f(x_2)\,g(x_2)$. So fg is increasing on I.

15. True. Let $x_1, x_2 \in I$ and $x_1 < x_2$. Then $f(x_1) < f(x_2)$ [f is increasing] $\Rightarrow$ $\dfrac{1}{f(x_1)} > \dfrac{1}{f(x_2)}$ [f is positive] $\Rightarrow$ $g(x_1) > g(x_2)$ $\Rightarrow$ $g(x) = 1/f(x)$ is decreasing on I.

16. False. If f is even, then $f(x) = f(-x)$. Using the Chain Rule to differentiate this equation, we get $f'(x) = f'(-x)\dfrac{d}{dx}(-x) = -f'(-x)$. Thus, $f'(-x) = -f'(x)$, so f' is odd.

17. True. If f is periodic, then there is a number p such that $f(x+p) = f(p)$ for all x. Differentiating gives $f'(x) = f'(x+p)\cdot(x+p)' = f'(x+p)\cdot 1 = f'(x+p)$, so f' is periodic.

18. False. The most general antiderivative of $f(x) = x^{-2}$ is $F(x) = -1/x + C_1$ for $x < 0$ and $F(x) = -1/x + C_2$ for $x > 0$ [see Example 4.9.1(b)].

19. True. By the Mean Value Theorem, there exists a number c in $(0, 1)$ such that $f(1) - f(0) = f'(c)(1-0) = f'(c)$. Since $f'(c)$ is nonzero, $f(1) - f(0) \neq 0$, so $f(1) \neq f(0)$.

20. False. Let $f(x) = 1 + \dfrac{1}{x}$ and $g(x) = x$. Then $\lim\limits_{x\to\infty} f(x) = 1$ and $\lim\limits_{x\to\infty} g(x) = \infty$, but

$$\lim_{x\to\infty}[f(x)]^{g(x)} = \lim_{x\to\infty}\left(1+\frac{1}{x}\right)^x = e, \text{ not } 1.$$

21. False. $\displaystyle\lim_{x\to 0}\frac{x}{e^x} = \frac{\lim\limits_{x\to 0} x}{\lim\limits_{x\to 0} e^x} = \frac{0}{1} = 0$, not 1.

EXERCISES

1. $f(x) = x^3 - 9x^2 + 24x - 2$, $[0, 5]$. $f'(x) = 3x^2 - 18x + 24 = 3(x^2 - 6x + 8) = 3(x-2)(x-4)$. $f'(x) = 0 \Leftrightarrow x = 2$ or $x = 4$. $f'(x) > 0$ for $0 < x < 2$, $f'(x) < 0$ for $2 < x < 4$, and $f'(x) > 0$ for $4 < x < 5$, so $f(2) = 18$ is a local maximum value and $f(4) = 14$ is a local minimum value. Checking the endpoints, we find $f(0) = -2$ and $f(5) = 18$. Thus, $f(0) = -2$ is the absolute minimum value and $f(2) = f(5) = 18$ is the absolute maximum value.

2. $f(x) = x\sqrt{1-x}$, $[-1, 1]$. $f'(x) = x\cdot\frac{1}{2}(1-x)^{-1/2}(-1) + (1-x)^{1/2}(1) = (1-x)^{-1/2}\left[-\frac{1}{2}x + (1-x)\right] = \dfrac{1-\frac{3}{2}x}{\sqrt{1-x}}$. $f'(x) = 0 \Rightarrow x = \frac{2}{3}$. $f'(x)$ does not exist $\Leftrightarrow x = 1$. $f'(x) > 0$ for $-1 < x < \frac{2}{3}$ and $f'(x) < 0$ for $\frac{2}{3} < x < 1$, so $f\left(\frac{2}{3}\right) = \frac{2}{3}\sqrt{\frac{1}{3}} = \frac{2}{9}\sqrt{3}$ [≈ 0.38] is a local maximum value. Checking the endpoints, we find $f(-1) = -\sqrt{2}$ and $f(1) = 0$. Thus, $f(-1) = -\sqrt{2}$ is the absolute minimum value and $f\left(\frac{2}{3}\right) = \frac{2}{9}\sqrt{3}$ is the absolute maximum value.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. $f(x) = \dfrac{3x-4}{x^2+1}$, $[-2, 2]$. $f'(x) = \dfrac{(x^2+1)(3)-(3x-4)(2x)}{(x^2+1)^2} = \dfrac{-(3x^2-8x-3)}{(x^2+1)^2} = \dfrac{-(3x+1)(x-3)}{(x^2+1)^2}$.

$f'(x) = 0 \;\Rightarrow\; x = -\frac{1}{3}$ or $x = 3$, but 3 is not in the interval. $f'(x) > 0$ for $-\frac{1}{3} < x < 2$ and $f'(x) < 0$ for $-2 < x < -\frac{1}{3}$, so $f\left(-\frac{1}{3}\right) = \frac{-5}{10/9} = -\frac{9}{2}$ is a local minimum value. Checking the endpoints, we find $f(-2) = -2$ and $f(2) = \frac{2}{5}$. Thus, $f\left(-\frac{1}{3}\right) = -\frac{9}{2}$ is the absolute minimum value and $f(2) = \frac{2}{5}$ is the absolute maximum value.

4. $f(x) = \sqrt{x^2+x+1}$, $[-2, 1]$. $f'(x) = \frac{1}{2}(x^2+x+1)^{-1/2}(2x+1) = \dfrac{2x+1}{2\sqrt{x^2+x+1}}$. $f'(x) = 0 \;\Rightarrow\; x = -\dfrac{1}{2}$.

$f'(x) > 0$ for $-\frac{1}{2} < x < 1$ and $f'(x) < 0$ for $-2 < x < -\frac{1}{2}$, so $f\left(-\frac{1}{2}\right) = \sqrt{3}/2$ is a local minimum value. Checking the endpoints, we find $f(-2) = f(1) = \sqrt{3}$. Thus, $f\left(-\frac{1}{2}\right) = \sqrt{3}/2$ is the absolute minimum value and $f(-2) = f(1) = \sqrt{3}$ is the absolute maximum value.

5. $f(x) = x + 2\cos x$, $[-\pi, \pi]$. $f'(x) = 1 - 2\sin x$. $f'(x) = 0 \;\Rightarrow\; \sin x = \frac{1}{2} \;\Rightarrow\; x = \frac{\pi}{6}, \frac{5\pi}{6}$. $f'(x) > 0$ for $\left(-\pi, \frac{\pi}{6}\right)$ and $\left(\frac{5\pi}{6}, \pi\right)$, and $f'(x) < 0$ for $\left(\frac{\pi}{6}, \frac{5\pi}{6}\right)$, so $f\left(\frac{\pi}{6}\right) = \frac{\pi}{6} + \sqrt{3} \approx 2.26$ is a local maximum value and $f\left(\frac{5\pi}{6}\right) = \frac{5\pi}{6} - \sqrt{3} \approx 0.89$ is a local minimum value. Checking the endpoints, we find $f(-\pi) = -\pi - 2 \approx -5.14$ and $f(\pi) = \pi - 2 \approx 1.14$. Thus, $f(-\pi) = -\pi - 2$ is the absolute minimum value and $f\left(\frac{\pi}{6}\right) = \frac{\pi}{6} + \sqrt{3}$ is the absolute maximum value.

6. $f(x) = x^2e^{-x}$, $[-1, 3]$. $f'(x) = x^2(-e^{-x}) + e^{-x}(2x) = xe^{-x}(-x+2)$. $f'(x) = 0 \;\Rightarrow\; x = 0$ or $x = 2$.

$f'(x) > 0$ for $0 < x < 2$ and $f'(x) < 0$ for $-1 < x < 0$ and $2 < x < 3$, so $f(0) = 0$ is a local minimum value and $f(2) = 4e^{-2} \approx 0.54$ is a local maximum value. Checking the endpoints, we find $f(-1) = e \approx 2.72$ and $f(3) = 9e^{-3} \approx 0.45$. Thus, $f(0) = 0$ is the absolute minimum value and $f(-1) = e$ is the absolute maximum value.

7. This limit has the form $\frac{0}{0}$. $\displaystyle\lim_{x\to 0} \frac{e^x - 1}{\tan x} \overset{\text{H}}{=} \lim_{x\to 0} \frac{e^x}{\sec^2 x} = \frac{1}{1} = 1$

8. This limit has the form $\frac{0}{0}$. $\displaystyle\lim_{x\to 0} \frac{\tan 4x}{x + \sin 2x} \overset{\text{H}}{=} \lim_{x\to 0} \frac{4\sec^2 4x}{1 + 2\cos 2x} = \frac{4(1)}{1+2(1)} = \frac{4}{3}$

9. This limit has the form $\frac{0}{0}$. $\displaystyle\lim_{x\to 0} \frac{e^{2x} - e^{-2x}}{\ln(x+1)} \overset{\text{H}}{=} \lim_{x\to 0} \frac{2e^{2x} + 2e^{-2x}}{1/(x+1)} = \frac{2+2}{1} = 4$

10. This limit has the form $\frac{\infty}{\infty}$. $\displaystyle\lim_{x\to\infty} \frac{e^{2x} - e^{-2x}}{\ln(x+1)} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{2e^{2x} + 2e^{-2x}}{1/(x+1)} = \lim_{x\to\infty} 2(x+1)(e^{2x} + e^{-2x}) = \infty$

since $2(x+1) \to \infty$ and $(e^{2x} + e^{-2x}) \to \infty$ as $x \to \infty$.

11. This limit has the form $\infty \cdot 0$.

$$\lim_{x\to-\infty} (x^2 - x^3)e^{2x} = \lim_{x\to-\infty} \frac{x^2 - x^3}{e^{-2x}} \quad \left[\tfrac{\infty}{\infty} \text{ form}\right] \quad \overset{\text{H}}{=} \lim_{x\to-\infty} \frac{2x - 3x^2}{-2e^{-2x}} \quad \left[\tfrac{\infty}{\infty} \text{ form}\right]$$
$$\overset{\text{H}}{=} \lim_{x\to-\infty} \frac{2 - 6x}{4e^{-2x}} \quad \left[\tfrac{\infty}{\infty} \text{ form}\right] \quad \overset{\text{H}}{=} \lim_{x\to-\infty} \frac{-6}{-8e^{-2x}} = 0$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. This limit has the form $0 \cdot \infty$. $\displaystyle\lim_{x\to\pi^-}(x-\pi)\csc x = \lim_{x\to\pi^-}\frac{x-\pi}{\sin x}$ $\left[\frac{0}{0}\text{ form}\right] \overset{\text{H}}{=} \displaystyle\lim_{x\to\pi^-}\frac{1}{\cos x} = \frac{1}{-1} = -1$

13. This limit has the form $\infty - \infty$.

$$\lim_{x\to1^+}\left(\frac{x}{x-1}-\frac{1}{\ln x}\right) = \lim_{x\to1^+}\left(\frac{x\ln x - x + 1}{(x-1)\ln x}\right) \overset{\text{H}}{=} \lim_{x\to1^+}\frac{x\cdot(1/x)+\ln x - 1}{(x-1)\cdot(1/x)+\ln x} = \lim_{x\to1^+}\frac{\ln x}{1-1/x+\ln x}$$

$$\overset{\text{H}}{=} \lim_{x\to1^+}\frac{1/x}{1/x^2+1/x} = \frac{1}{1+1} = \frac{1}{2}$$

14. $y = (\tan x)^{\cos x} \quad\Rightarrow\quad \ln y = \cos x\,\ln\tan x$, so

$$\lim_{x\to(\pi/2)^-}\ln y = \lim_{x\to(\pi/2)^-}\frac{\ln\tan x}{\sec x} \overset{\text{H}}{=} \lim_{x\to(\pi/2)^-}\frac{(1/\tan x)\sec^2 x}{\sec x\,\tan x} = \lim_{x\to(\pi/2)^-}\frac{\sec x}{\tan^2 x} = \lim_{x\to(\pi/2)^-}\frac{\cos x}{\sin^2 x} = \frac{0}{1^2} = 0,$$

so $\displaystyle\lim_{x\to(\pi/2)^-}(\tan x)^{\cos x} = \lim_{x\to(\pi/2)^-}e^{\ln y} = e^0 = 1$.

15. $f(0)=0$, $f'(-2)=f'(1)=f'(9)=0$, $\displaystyle\lim_{x\to\infty}f(x)=0$, $\displaystyle\lim_{x\to6}f(x)=-\infty$,

$f'(x)<0$ on $(-\infty,-2)$, $(1,6)$, and $(9,\infty)$, $f'(x)>0$ on $(-2,1)$ and $(6,9)$,

$f''(x)>0$ on $(-\infty,0)$ and $(12,\infty)$, $f''(x)<0$ on $(0,6)$ and $(6,12)$

16. For $0<x<1$, $f'(x)=2x$, so $f(x)=x^2+C$. Since $f(0)=0$,

$f(x)=x^2$ on $[0,1]$. For $1<x<3$, $f'(x)=-1$, so $f(x)=-x+D$.

$1=f(1)=-1+D \quad\Rightarrow\quad D=2$, so $f(x)=2-x$. For $x>3$, $f'(x)=1$,

so $f(x)=x+E$. $-1=f(3)=3+E \quad\Rightarrow\quad E=-4$, so $f(x)=x-4$.

Since f is even, its graph is symmetric about the y-axis.

17. f is odd, $f'(x)<0$ for $0<x<2$, $\quad f'(x)>0$ for $x>2$,

$f''(x)>0$ for $0<x<3$, $\quad f''(x)<0$ for $x>3$,

$\lim_{x\to\infty}f(x)=-2$

18. (a) Since $f'(x)>0$ on $(1,4)$ and $(4,7)$, f is increasing on these intervals. Since $f'(x)<0$ on $(0,1)$ and $(7,8)$, f is decreasing on these intervals.

(b) Since $f'(7)=0$ and f' changes from positive to negative there, f changes from increasing to decreasing and has a local maximum at $x=7$. Since $f'(x)=0$ at $x=1$ and f' changes from negative to positive there, f changes from decreasing to increasing and has a local minimum at $x=1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c)

(d)

19. $y = f(x) = 2 - 2x - x^3$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 2$. The x-intercept (approximately 0.770917) can be found using Newton's Method. **C.** No symmetry **D.** No asymptote

E. $f'(x) = -2 - 3x^2 = -(3x^2 + 2) < 0$, so f is decreasing on $\mathbb{R}$.

F. No extreme value **G.** $f''(x) = -6x < 0$ on $(0, \infty)$ and $f''(x) > 0$ on $(-\infty, 0)$, so f is CD on $(0, \infty)$ and CU on $(-\infty, 0)$. There is an IP at $(0, 2)$.

H.

20. $y = f(x) = -2x^3 - 3x^2 + 12x + 5$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 5$; x-intercept: $f(x) = 0 \Leftrightarrow x \approx -3.15, -0.39, 2.04$ **C.** No symmetry **D.** No asymptote

E. $f'(x) = -6x^2 - 6x + 12 = -6(x^2 + x - 2) = -6(x + 2)(x - 1)$. $f'(x) > 0$ for $-2 < x < 1$, so f is increasing on $(-2, 1)$ and decreasing on $(-\infty, -2)$ and $(1, \infty)$. **F.** Local minimum value $f(-2) = -15$, local maximum value $f(1) = 12$ **G.** $f''(x) = -12x - 6 = -12\left(x + \frac{1}{2}\right)$. $f''(x) > 0$ for $x < -\frac{1}{2}$, so f is CU on $\left(-\infty, -\frac{1}{2}\right)$ and CD on $\left(-\frac{1}{2}, \infty\right)$. There is an IP at $\left(-\frac{1}{2}, -\frac{3}{2}\right)$.

H.

21. $y = f(x) = 3x^4 - 4x^3 + 2$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 2$; no x-intercept **C.** No symmetry **D.** No asymptote

E. $f'(x) = 12x^3 - 12x^2 = 12x^2(x - 1)$. $f'(x) > 0$ for $x > 1$, so f is increasing on $(1, \infty)$ and decreasing on $(-\infty, 1)$. **F.** $f'(x)$ does not change sign at $x = 0$, so there is no local extremum there. $f(1) = 1$ is a local minimum value. **G.** $f''(x) = 36x^2 - 24x = 12x(3x - 2)$. $f''(x) < 0$ for $0 < x < \frac{2}{3}$, so f is CD on $\left(0, \frac{2}{3}\right)$ and f is CU on $(-\infty, 0)$ and $\left(\frac{2}{3}, \infty\right)$. There are inflection points at $(0, 2)$ and $\left(\frac{2}{3}, \frac{38}{27}\right)$.

H.

22. $y = f(x) = \dfrac{x}{1 - x^2}$ **A.** $D = (-\infty, -1) \cup (-1, 1) \cup (1, \infty)$ **B.** y-intercept: $f(0) = 0$; x-intercept: 0

C. $f(-x) = -f(x)$, so f is odd and the graph is symmetric about the origin. **D.** $\displaystyle\lim_{x \to \pm\infty} \frac{x}{1 - x^2} = 0$, so $y = 0$ is a HA.

$\displaystyle\lim_{x \to -1^-} \frac{x}{1 - x^2} = \infty$ and $\displaystyle\lim_{x \to -1^+} \frac{x}{1 - x^2} = -\infty$, so $x = -1$ is a VA. Similarly, $\displaystyle\lim_{x \to 1^-} \frac{x}{1 - x^2} = \infty$ and

$\displaystyle\lim_{x \to 1^+} \frac{x}{1 - x^2} = -\infty$, so $x = 1$ is a VA. **E.** $f'(x) = \dfrac{(1 - x^2)(1) - x(-2x)}{(1 - x^2)^2} = \dfrac{1 + x^2}{(1 - x^2)^2} > 0$ for $x \neq \pm 1$, so f is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

increasing on $(-\infty, -1)$, $(-1, 1)$, and $(1, \infty)$. **F.** No local extrema

G. $f''(x) = \dfrac{(1-x^2)^2(2x) - (1+x^2)2(1-x^2)(-2x)}{[(1-x^2)^2]^2}$

$= \dfrac{2x(1-x^2)[(1-x^2)+2(1+x^2)]}{(1-x^2)^4} = \dfrac{2x(3+x^2)}{(1-x^2)^3}$

$f''(x) > 0$ for $x < -1$ and $0 < x < 1$, and $f''(x) < 0$ for $-1 < x < 0$ and $x > 1$, so f is CU on $(-\infty, -1)$ and $(0, 1)$, and f is CD on $(-1, 0)$ and $(1, \infty)$. $(0, 0)$ is an IP.

H.

23. $y = f(x) = \dfrac{1}{x(x-3)^2}$ **A.** $D = \{x \mid x \neq 0, 3\} = (-\infty, 0) \cup (0, 3) \cup (3, \infty)$ **B.** No intercepts. **C.** No symmetry.

D. $\lim\limits_{x\to\pm\infty} \dfrac{1}{x(x-3)^2} = 0$, so $y = 0$ is a HA. $\lim\limits_{x\to 0^+} \dfrac{1}{x(x-3)^2} = \infty$, $\lim\limits_{x\to 0^-} \dfrac{1}{x(x-3)^2} = -\infty$, $\lim\limits_{x\to 3} \dfrac{1}{x(x-3)^2} = \infty$, so $x = 0$ and $x = 3$ are VA. **E.** $f'(x) = -\dfrac{(x-3)^2 + 2x(x-3)}{x^2(x-3)^4} = \dfrac{3(1-x)}{x^2(x-3)^3} \Rightarrow f'(x) > 0 \Leftrightarrow 1 < x < 3$, so f is increasing on $(1, 3)$ and decreasing on $(-\infty, 0)$, $(0, 1)$, and $(3, \infty)$.

F. Local minimum value $f(1) = \frac{1}{4}$ **G.** $f''(x) = \dfrac{6(2x^2 - 4x + 3)}{x^3(x-3)^4}$.

Note that $2x^2 - 4x + 3 > 0$ for all x since it has negative discriminant. So $f''(x) > 0 \Leftrightarrow x > 0 \Rightarrow f$ is CU on $(0, 3)$ and $(3, \infty)$ and CD on $(-\infty, 0)$. No IP

H.

24. $y = f(x) = \dfrac{1}{x^2} - \dfrac{1}{(x-2)^2}$ **A.** $D = \{x \mid x \neq 0, 2\}$ **B.** y-intercept: none; x-intercept: $f(x) = 0 \Rightarrow$

$\dfrac{1}{x^2} = \dfrac{1}{(x-2)^2} \Leftrightarrow (x-2)^2 = x^2 \Leftrightarrow x^2 - 4x + 4 = x^2 \Leftrightarrow 4x = 4 \Leftrightarrow x = 1$ **C.** No symmetry

D. $\lim\limits_{x\to 0} f(x) = \infty$ and $\lim\limits_{x\to 2} f(x) = -\infty$, so $x = 0$ and $x = 2$ are VA; $\lim\limits_{x\to\pm\infty} f(x) = 0$, so $y = 0$ is a HA

E. $f'(x) = -\dfrac{2}{x^3} + \dfrac{2}{(x-2)^3} > 0 \Rightarrow \dfrac{-(x-2)^3 + x^3}{x^3(x-2)^3} > 0 \Leftrightarrow \dfrac{-x^3 + 6x^2 - 12x + 8 + x^3}{x^3(x-2)^3} > 0 \Leftrightarrow$

$\dfrac{2(3x^2 - 6x + 4)}{x^3(x-2)^3} > 0$. The numerator is positive (the discriminant of the quadratic is negative), so $f'(x) > 0$ if $x < 0$ or $x > 2$, and hence, f is increasing on $(-\infty, 0)$ and $(2, \infty)$ and decreasing on $(0, 2)$.

F. No local extreme values **G.** $f''(x) = \dfrac{6}{x^4} - \dfrac{6}{(x-2)^4} > 0 \Rightarrow$

$\dfrac{(x-2)^4 - x^4}{x^4(x-2)^4} > 0 \Leftrightarrow \dfrac{x^4 - 8x^3 + 24x^2 - 32x + 16 - x^4}{x^4(x-2)^4} > 0 \Leftrightarrow$

$\dfrac{-8(x^3 - 3x^2 + 4x - 2)}{x^4(x-2)^4} > 0 \Leftrightarrow \dfrac{-8(x-1)(x^2 - 2x + 2)}{x^4(x-2)^4} > 0$. So f'' is positive for $x < 1$ $[x \neq 0]$ and negative for $x > 1$ $[x \neq 2]$. Thus, f is CU on $(-\infty, 0)$ and $(0, 1)$ and f is CD on $(1, 2)$ and $(2, \infty)$. IP at $(1, 0)$

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $y = f(x) = \dfrac{(x-1)^3}{x^2} = \dfrac{x^3 - 3x^2 + 3x - 1}{x^2} = x - 3 + \dfrac{3x-1}{x^2}$ **A.** $D = \{x \mid x \neq 0\} = (-\infty, 0) \cup (0, \infty)$

B. y-intercept: none; x-intercept: $f(x) = 0 \quad \Leftrightarrow \quad x = 1$ **C.** No symmetry **D.** $\displaystyle\lim_{x \to 0^-} \frac{(x-1)^3}{x^2} = -\infty$ and

$\displaystyle\lim_{x \to 0^+} f(x) = -\infty$, so $x = 0$ is a VA. $f(x) - (x-3) = \dfrac{3x-1}{x^2} \to 0$ as $x \to \pm\infty$, so $y = x - 3$ is a SA.

E. $f'(x) = \dfrac{x^2 \cdot 3(x-1)^2 - (x-1)^3(2x)}{(x^2)^2} = \dfrac{x(x-1)^2[3x - 2(x-1)]}{x^4} = \dfrac{(x-1)^2(x+2)}{x^3}$. $f'(x) < 0$ for $-2 < x < 0$,

so f is increasing on $(-\infty, -2)$, decreasing on $(-2, 0)$, and increasing on $(0, \infty)$.

H.

F. Local maximum value $f(-2) = -\frac{27}{4}$ **G.** $f(x) = x - 3 + \dfrac{3}{x} - \dfrac{1}{x^2} \quad \Rightarrow$

$f'(x) = 1 - \dfrac{3}{x^2} + \dfrac{2}{x^3} \quad \Rightarrow \quad f''(x) = \dfrac{6}{x^3} - \dfrac{6}{x^4} = \dfrac{6x-6}{x^4} = \dfrac{6(x-1)}{x^4}$.

$f''(x) > 0$ for $x > 1$, so f is CD on $(-\infty, 0)$ and $(0, 1)$, and f is CU on $(1, \infty)$.

There is an inflection point at $(1, 0)$.

26. $y = f(x) = \sqrt{1-x} + \sqrt{1+x}$ **A.** $1 - x \geq 0$ and $1 + x \geq 0 \quad \Rightarrow \quad x \leq 1$ and $x \geq -1$, so $D = [-1, 1]$.

B. y-intercept: $f(0) = 1 + 1 = 2$; no x-intercept because $f(x) > 0$ for all x.

C. $f(-x) = f(x)$, so the curve is symmetric about the y-axis. **D.** No asymptote

E. $f'(x) = \frac{1}{2}(1-x)^{-1/2}(-1) + \frac{1}{2}(1+x)^{-1/2} = \dfrac{-1}{2\sqrt{1-x}} + \dfrac{1}{2\sqrt{1+x}} = \dfrac{-\sqrt{1+x} + \sqrt{1-x}}{2\sqrt{1-x}\sqrt{1+x}} > 0 \quad \Rightarrow$

$-\sqrt{1+x} + \sqrt{1-x} > 0 \quad \Rightarrow \quad \sqrt{1-x} > \sqrt{1+x} \quad \Rightarrow \quad 1 - x > 1 + x \quad \Rightarrow \quad -2x > 0 \quad \Rightarrow \quad x < 0$, so $f'(x) > 0$ for

$-1 < x < 0$ and $f'(x) < 0$ for $0 < x < 1$. Thus, f is increasing on $(-1, 0)$

and decreasing on $(0, 1)$. **F.** Local maximum value $f(0) = 2$

H.

G. $f''(x) = -\frac{1}{2}\left(-\frac{1}{2}\right)(1-x)^{-3/2}(-1) + \frac{1}{2}\left(-\frac{1}{2}\right)(1+x)^{-3/2}$

$$= \frac{-1}{4(1-x)^{3/2}} + \frac{-1}{4(1+x)^{3/2}} < 0$$

for all x in the domain, so f is CD on $(-1, 1)$. No IP

27. $y = f(x) = x\sqrt{2+x}$ **A.** $D = [-2, \infty)$ **B.** y-intercept: $f(0) = 0$; x-intercepts: -2 and 0 **C.** No symmetry

D. No asymptote **E.** $f'(x) = \dfrac{x}{2\sqrt{2+x}} + \sqrt{2+x} = \dfrac{1}{2\sqrt{2+x}}[x + 2(2+x)] = \dfrac{3x+4}{2\sqrt{2+x}} = 0$ when $x = -\frac{4}{3}$, so f is

decreasing on $\left(-2, -\frac{4}{3}\right)$ and increasing on $\left(-\frac{4}{3}, \infty\right)$. **F.** Local minimum value $f\left(-\frac{4}{3}\right) = -\frac{4}{3}\sqrt{\frac{2}{3}} = -\frac{4\sqrt{6}}{9} \approx -1.09$,

no local maximum

H.

G. $f''(x) = \dfrac{2\sqrt{2+x} \cdot 3 - (3x+4)\dfrac{1}{\sqrt{2+x}}}{4(2+x)} = \dfrac{6(2+x) - (3x+4)}{4(2+x)^{3/2}}$

$$= \frac{3x+8}{4(2+x)^{3/2}}$$

$f''(x) > 0$ for $x > -2$, so f is CU on $(-2, \infty)$. No IP

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $y = f(x) = x^{2/3}(x-3)^2$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \quad \Leftrightarrow \quad x = 0, 3$

C. No symmetry **D.** No asymptote

E. $f'(x) = x^{2/3} \cdot 2(x-3) + (x-3)^2 \cdot \frac{2}{3}x^{-1/3} = \frac{2}{3}x^{-1/3}(x-3)[3x + (x-3)] = \frac{2}{3}x^{-1/3}(x-3)(4x-3)$.

$f'(x) > 0 \quad \Leftrightarrow \quad 0 < x < \frac{3}{4}$ or $x > 3$, so f is decreasing on $(-\infty, 0)$, increasing on $\left(0, \frac{3}{4}\right)$, decreasing on $\left(\frac{3}{4}, 3\right)$, and increasing on $(3, \infty)$. **F.** Local minimum value $f(0) = f(3) = 0$; local maximum value

$f\left(\frac{3}{4}\right) = \left(\frac{3}{4}\right)^{2/3}\left(-\frac{9}{4}\right)^2 = \frac{81}{16}\sqrt[3]{\frac{9}{16}} = \frac{81}{32}\sqrt[3]{\frac{9}{2}}$ $[\approx 4.18]$

G. $f'(x) = \left(\frac{2}{3}x^{-1/3}\right)(4x^2 - 15x + 9) \quad \Rightarrow$

$$\begin{aligned} f''(x) &= \left(\tfrac{2}{3}x^{-1/3}\right)(8x-15) + (4x^2 - 15x + 9)\left(-\tfrac{2}{9}x^{-4/3}\right) \\ &= \tfrac{2}{9}x^{-4/3}[3x(8x-15) - (4x^2 - 15x + 9)] \\ &= \tfrac{2}{9}x^{-4/3}(20x^2 - 30x - 9) \end{aligned}$$

$f''(x) = 0 \quad \Leftrightarrow \quad x \approx -0.26$ or 1.76. $f''(x)$ does not exist at $x = 0$.

f is CU on $(-\infty, -0.26)$, CD on $(-0.26, 0)$, CD on $(0, 1.76)$, and CU on $(1.76, \infty)$. There are inflection points at $(-0.26, 4.28)$ and $(1.76, 2.25)$.

H.

29. $y = f(x) = e^x \sin x, -\pi \le x \le \pi$ **A.** $D = [-\pi, \pi]$ **B.** y-intercept: $f(0) = 0$; $f(x) = 0 \quad \Leftrightarrow \quad \sin x = 0 \quad \Rightarrow$ $x = -\pi, 0, \pi$. **C.** No symmetry **D.** No asymptote **E.** $f'(x) = e^x \cos x + \sin x \cdot e^x = e^x(\cos x + \sin x)$.

$f'(x) = 0 \quad \Leftrightarrow \quad -\cos x = \sin x \quad \Leftrightarrow \quad -1 = \tan x \quad \Rightarrow \quad x = -\frac{\pi}{4}, \frac{3\pi}{4}$. $f'(x) > 0$ for $-\frac{\pi}{4} < x < \frac{3\pi}{4}$ and $f'(x) < 0$ for $-\pi < x < -\frac{\pi}{4}$ and $\frac{3\pi}{4} < x < \pi$, so f is increasing on $\left(-\frac{\pi}{4}, \frac{3\pi}{4}\right)$ and f is decreasing on $\left(-\pi, -\frac{\pi}{4}\right)$ and $\left(\frac{3\pi}{4}, \pi\right)$.

F. Local minimum value $f\left(-\frac{\pi}{4}\right) = (-\sqrt{2}/2)e^{-\pi/4} \approx -0.32$ and local maximum value $f\left(\frac{3\pi}{4}\right) = \left(\sqrt{2}/2\right)e^{3\pi/4} \approx 7.46$

G. $f''(x) = e^x(-\sin x + \cos x) + (\cos x + \sin x)e^x = e^x(2\cos x) > 0 \quad \Rightarrow$ $-\frac{\pi}{2} < x < \frac{\pi}{2}$ and $f''(x) < 0 \quad \Rightarrow \quad -\pi < x < -\frac{\pi}{2}$ and $\frac{\pi}{2} < x < \pi$, so f is CU on $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, and f is CD on $\left(-\pi, -\frac{\pi}{2}\right)$ and $\left(\frac{\pi}{2}, \pi\right)$. There are inflection points at $\left(-\frac{\pi}{2}, -e^{-\pi/2}\right)$ and $\left(\frac{\pi}{2}, e^{\pi/2}\right)$.

H.

30. $y = f(x) = 4x - \tan x, -\frac{\pi}{2} < x < \frac{\pi}{2}$ **A.** $D = \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$. **B.** y-intercept $= f(0) = 0$ **C.** $f(-x) = -f(x)$, so the curve is symmetric about $(0, 0)$. **D.** $\lim\limits_{x \to \pi/2^-}(4x - \tan x) = -\infty$, $\lim\limits_{x \to -\pi/2^+}(4x - \tan x) = \infty$, so $x = \frac{\pi}{2}$ and $x = -\frac{\pi}{2}$ are VA. **E.** $f'(x) = 4 - \sec^2 x > 0 \quad \Leftrightarrow \quad \sec x < 2 \quad \Leftrightarrow \quad \cos x > \frac{1}{2} \quad \Leftrightarrow \quad -\frac{\pi}{3} < x < \frac{\pi}{3}$, so f is increasing on $\left(-\frac{\pi}{3}, \frac{\pi}{3}\right)$ and decreasing on $\left(-\frac{\pi}{2}, -\frac{\pi}{3}\right)$ and $\left(\frac{\pi}{3}, \frac{\pi}{2}\right)$. **F.** $f\left(\frac{\pi}{3}\right) = \frac{4\pi}{3} - \sqrt{3}$ is a local maximum value, $f\left(-\frac{\pi}{3}\right) = \sqrt{3} - \frac{4\pi}{3}$ is a local minimum value.

G. $f''(x) = -2\sec^2 x \tan x > 0 \quad \Leftrightarrow \quad \tan x < 0 \quad \Leftrightarrow \quad -\frac{\pi}{2} < x < 0$, so f is CU on $\left(-\frac{\pi}{2}, 0\right)$ and CD on $\left(0, \frac{\pi}{2}\right)$. IP at $(0, 0)$

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. $y = f(x) = \sin^{-1}(1/x)$ **A.** $D = \{x \mid -1 \le 1/x \le 1\} = (-\infty, -1] \cup [1, \infty)$. **B.** No intercept

C. $f(-x) = -f(x)$, symmetric about the origin **D.** $\lim\limits_{x \to \pm\infty} \sin^{-1}(1/x) = \sin^{-1}(0) = 0$, so $y = 0$ is a HA.

E. $f'(x) = \dfrac{1}{\sqrt{1-(1/x)^2}}\left(-\dfrac{1}{x^2}\right) = \dfrac{-1}{\sqrt{x^4 - x^2}} < 0$, so f is decreasing on $(-\infty, -1)$ and $(1, \infty)$.

F. No local extreme value, but $f(1) = \frac{\pi}{2}$ is the absolute maximum value and $f(-1) = -\frac{\pi}{2}$ is the absolute minimum value.

H.

G. $f''(x) = \dfrac{4x^3 - 2x}{2(x^4 - x^2)^{3/2}} = \dfrac{x(2x^2 - 1)}{(x^4 - x^2)^{3/2}} > 0$ for $x > 1$ and $f''(x) < 0$ for $x < -1$, so f is CU on $(1, \infty)$ and CD on $(-\infty, -1)$. No IP

32. $y = f(x) = e^{2x - x^2}$ **A.** $D = \mathbb{R}$ **B.** y-intercept 1; no x-intercept **C.** No symmetry **D.** $\lim\limits_{x \to \pm\infty} e^{2x - x^2} = 0$, so $y = 0$ is a HA. **E.** $y = f(x) = e^{2x - x^2} \Rightarrow f'(x) = 2(1 - x)e^{2x - x^2} > 0 \Leftrightarrow x < 1$, so f is increasing on $(-\infty, 1)$ and decreasing on $(1, \infty)$. **F.** $f(1) = e$ is a local and absolute maximum value.

G. $f''(x) = 2(2x^2 - 4x + 1)e^{2x - x^2} = 0 \Leftrightarrow x = 1 \pm \frac{\sqrt{2}}{2}$.

$f''(x) > 0 \Leftrightarrow x < 1 - \frac{\sqrt{2}}{2}$ or $x > 1 + \frac{\sqrt{2}}{2}$, so f is CU on $\left(-\infty, 1 - \frac{\sqrt{2}}{2}\right)$ and $\left(1 + \frac{\sqrt{2}}{2}, \infty\right)$, and CD on $\left(1 - \frac{\sqrt{2}}{2}, 1 + \frac{\sqrt{2}}{2}\right)$. IP at $\left(1 \pm \frac{\sqrt{2}}{2}, \sqrt{e}\right)$

H.

33. $y = f(x) = (x - 2)e^{-x}$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = -2$; x-intercept: $f(x) = 0 \Leftrightarrow x = 2$

C. No symmetry **D.** $\lim\limits_{x \to \infty} \dfrac{x - 2}{e^x} \overset{\text{H}}{=} \lim\limits_{x \to \infty} \dfrac{1}{e^x} = 0$, so $y = 0$ is a HA. No VA

E. $f'(x) = (x - 2)(-e^{-x}) + e^{-x}(1) = e^{-x}[-(x - 2) + 1] = (3 - x)e^{-x}$.

$f'(x) > 0$ for $x < 3$, so f is increasing on $(-\infty, 3)$ and decreasing on $(3, \infty)$.

F. Local maximum value $f(3) = e^{-3}$, no local minimum value

G. $f''(x) = (3 - x)(-e^{-x}) + e^{-x}(-1) = e^{-x}[-(3 - x) + (-1)]$
$= (x - 4)e^{-x} > 0$

for $x > 4$, so f is CU on $(4, \infty)$ and CD on $(-\infty, 4)$. IP at $(4, 2e^{-4})$

H.

34. $y = f(x) = x + \ln(x^2 + 1)$ **A.** $D = \mathbb{R}$ **B.** y-intercept: $f(0) = 0 + \ln 1 = 0$; x-intercept: $f(x) = 0 \Leftrightarrow \ln(x^2 + 1) = -x \Leftrightarrow x^2 + 1 = e^{-x} \Rightarrow x = 0$ since the graphs of $y = x^2 + 1$ and $y = e^{-x}$ intersect only at $x = 0$.

C. No symmetry **D.** No asymptote **E.** $f'(x) = 1 + \dfrac{2x}{x^2 + 1} = \dfrac{x^2 + 2x + 1}{x^2 + 1} = \dfrac{(x + 1)^2}{x^2 + 1}$. $f'(x) > 0$ if $x \neq -1$ and f is increasing on $\mathbb{R}$. **F.** No local extreme values

G. $f''(x) = \dfrac{(x^2 + 1)2 - 2x(2x)}{(x^2 + 1)^2} = \dfrac{2[(x^2 + 1) - 2x^2]}{(x^2 + 1)^2} = \dfrac{2(1 - x^2)}{(x^2 + 1)^2}$.

$f''(x) > 0 \Leftrightarrow -1 < x < 1$ and $f''(x) < 0 \Leftrightarrow x < -1$ or $x > 1$, so f is CU on $(-1, 1)$ and f is CD on $(-\infty, -1)$ and $(1, \infty)$. IP at $(-1, -1 + \ln 2)$ and $(1, 1 + \ln 2)$

H.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. $f(x) = \dfrac{x^2-1}{x^3} \quad \Rightarrow \quad f'(x) = \dfrac{x^3(2x) - (x^2-1)3x^2}{x^6} = \dfrac{3-x^2}{x^4} \quad \Rightarrow$

$f''(x) = \dfrac{x^4(-2x) - (3-x^2)4x^3}{x^8} = \dfrac{2x^2-12}{x^5}$

Estimates: From the graphs of f' and f'', it appears that f is increasing on $(-1.73, 0)$ and $(0, 1.73)$ and decreasing on $(-\infty, -1.73)$ and $(1.73, \infty)$; f has a local maximum of about $f(1.73) = 0.38$ and a local minimum of about $f(-1.7) = -0.38$; f is CU on $(-2.45, 0)$ and $(2.45, \infty)$, and CD on $(-\infty, -2.45)$ and $(0, 2.45)$; and f has inflection points at about $(-2.45, -0.34)$ and $(2.45, 0.34)$.

Exact: Now $f'(x) = \dfrac{3-x^2}{x^4}$ is positive for $0 < x^2 < 3$, that is, f is increasing on $\left(-\sqrt{3}, 0\right)$ and $\left(0, \sqrt{3}\right)$; and $f'(x)$ is negative (and so f is decreasing) on $\left(-\infty, -\sqrt{3}\right)$ and $\left(\sqrt{3}, \infty\right)$. $f'(x) = 0$ when $x = \pm\sqrt{3}$.

f' goes from positive to negative at $x = \sqrt{3}$, so f has a local maximum of $f\left(\sqrt{3}\right) = \frac{\left(\sqrt{3}\right)^2 - 1}{\left(\sqrt{3}\right)^3} = \frac{2\sqrt{3}}{9}$; and since f is odd, we know that maxima on the interval $(0, \infty)$ correspond to minima on $(-\infty, 0)$, so f has a local minimum of $f\left(-\sqrt{3}\right) = -\frac{2\sqrt{3}}{9}$. Also, $f''(x) = \dfrac{2x^2-12}{x^5}$ is positive (so f is CU) on $\left(-\sqrt{6}, 0\right)$ and $\left(\sqrt{6}, \infty\right)$, and negative (so f is CD) on $\left(-\infty, -\sqrt{6}\right)$ and $\left(0, \sqrt{6}\right)$. There are IP at $\left(\sqrt{6}, \frac{5\sqrt{6}}{36}\right)$ and $\left(-\sqrt{6}, -\frac{5\sqrt{6}}{36}\right)$.

36. $f(x) = \dfrac{x^3+1}{x^6+1} \quad \Rightarrow \quad f'(x) = -\dfrac{3x^2(x^6+2x^3-1)}{(x^6+1)^2} \quad \Rightarrow \quad f''(x) = \dfrac{6x(2x^{12}+7x^9-9x^6-5x^3+1)}{(x^6+1)^3}$.

$f(x) = 0 \quad \Leftrightarrow \quad x = -1$. $f'(x) = 0 \quad \Leftrightarrow \quad x = 0$ or $x \approx -1.34, 0.75$. $f''(x) = 0 \quad \Leftrightarrow \quad x = 0$ or $x \approx -1.64, -0.82, 0.54, 1.09$. From the graphs of f and f', it appears that f is decreasing on $(-\infty, -1.34)$, increasing on $(-1.34, 0.75)$, and decreasing on $(0.75, \infty)$. f has a local minimum value of $f(-1.34) \approx -0.21$ and a local maximum value of $f(0.75) \approx 1.21$. From the graphs of f and f'', it appears that f is CD on $(-\infty, -1.64)$, CU on $(-1.64, -0.82)$, CD on $(-0.82, 0)$, CU on $(0, 0.54)$, CD on $(0.54, 1.09)$ and CU on $(1.09, \infty)$. There are inflection points at about $(-1.64, -0.17)$, $(-0.82, 0.34)$, $(0.54, 1.13)$, $(1.09, 0.86)$, and at $(0, 1)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. $f(x) = 3x^6 - 5x^5 + x^4 - 5x^3 - 2x^2 + 2 \quad \Rightarrow \quad f'(x) = 18x^5 - 25x^4 + 4x^3 - 15x^2 - 4x \quad \Rightarrow$
$f''(x) = 90x^4 - 100x^3 + 12x^2 - 30x - 4$

From the graphs of f' and f'', it appears that f is increasing on $(-0.23, 0)$ and $(1.62, \infty)$ and decreasing on $(-\infty, -0.23)$ and $(0, 1.62)$; f has a local maximum of $f(0) = 2$ and local minima of about $f(-0.23) = 1.96$ and $f(1.62) = -19.2$; f is CU on $(-\infty, -0.12)$ and $(1.24, \infty)$ and CD on $(-0.12, 1.24)$; and f has inflection points at about $(-0.12, 1.98)$ and $(1.24, -12.1)$.

38. $f(x) = x^2 + 6.5\sin x,\ -5 \le x \le 5 \quad \Rightarrow \quad f'(x) = 2x + 6.5\cos x \quad \Rightarrow \quad f''(x) = 2 - 6.5\sin x.$ $\quad f(x) = 0 \quad \Leftrightarrow$
$x \approx -2.25$ and $x = 0$; $\quad f'(x) = 0 \quad \Leftrightarrow \quad x \approx -1.19, 2.40, 3.24$; $\quad f''(x) = 0 \quad \Leftrightarrow \quad x \approx -3.45, 0.31, 2.83.$

From the graphs of f' and f'', it appears that f is decreasing on $(-5, -1.19)$ and $(2.40, 3.24)$ and increasing on $(-1.19, 2.40)$ and $(3.24, 5)$; f has a local maximum of about $f(2.40) = 10.15$ and local minima of about $f(-1.19) = -4.62$ and $f(3.24) = 9.86$; f is CU on $(-3.45, 0.31)$ and $(2.83, 5)$ and CD on $(-5, -3.45)$ and $(0.31, 2.83)$; and f has inflection points at about $(-3.45, 13.93)$, $(0.31, 2.10)$, and $(2.83, 10.00)$.

39.

From the graph, we estimate the points of inflection to be about $(\pm 0.82, 0.22)$.

$f(x) = e^{-1/x^2} \quad \Rightarrow \quad f'(x) = 2x^{-3}e^{-1/x^2} \quad \Rightarrow$

$f''(x) = 2[x^{-3}(2x^{-3})e^{-1/x^2} + e^{-1/x^2}(-3x^{-4})] = 2x^{-6}e^{-1/x^2}\left(2 - 3x^2\right).$

This is 0 when $2 - 3x^2 = 0 \quad \Leftrightarrow \quad x = \pm\sqrt{\frac{2}{3}}$, so the inflection points are $\left(\pm\sqrt{\frac{2}{3}}, e^{-3/2}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. (a)

(b) $f(x) = \dfrac{1}{1+e^{1/x}}$.

$\lim\limits_{x\to\infty} f(x) = \dfrac{1}{1+1} = \dfrac{1}{2}$, $\lim\limits_{x\to-\infty} f(x) = \dfrac{1}{1+1} = \dfrac{1}{2}$,

as $x \to 0^+$, $1/x \to \infty$, so $e^{1/x} \to \infty \quad \Rightarrow \quad \lim\limits_{x\to 0^+} f(x) = 0$,

as $x \to 0^-$, $1/x \to -\infty$, so $e^{1/x} \to 0 \quad \Rightarrow \quad \lim\limits_{x\to 0^-} f(x) = \dfrac{1}{1+0} = 1$

(c) From the graph of f, estimates for the IP are $(-0.4, 0.9)$ and $(0.4, 0.08)$.

(d) $f''(x) = -\dfrac{e^{1/x}[e^{1/x}(2x-1)+2x+1]}{x^4(e^{1/x}+1)^3}$

(e) From the graph, we see that f'' changes sign at $x = \pm 0.417$ ($x = 0$ is not in the domain of f). IP are approximately $(0.417, 0.083)$ and $(-0.417, 0.917)$.

41. $f(x) = \dfrac{\cos^2 x}{\sqrt{x^2+x+1}}$, $-\pi \le x \le \pi \quad \Rightarrow \quad f'(x) = -\dfrac{\cos x\,[(2x+1)\cos x + 4(x^2+x+1)\sin x]}{2(x^2+x+1)^{3/2}} \quad \Rightarrow$

$$f''(x) = -\frac{(8x^4+16x^3+16x^2+8x+9)\cos^2 x - 8(x^2+x+1)(2x+1)\sin x\,\cos x - 8(x^2+x+1)^2\sin^2 x}{4(x^2+x+1)^{5/2}}$$

$f(x) = 0 \quad \Leftrightarrow \quad x = \pm\frac{\pi}{2}$; $\quad f'(x) = 0 \quad \Leftrightarrow \quad x \approx -2.96, -1.57, -0.18, 1.57, 3.01$;

$f''(x) = 0 \quad \Leftrightarrow \quad x \approx -2.16, -0.75, 0.46$, and 2.21.

The x-coordinates of the maximum points are the values at which f' changes from positive to negative, that is, -2.96, -0.18, and 3.01. The x-coordinates of the minimum points are the values at which f' changes from negative to positive, that is, -1.57 and 1.57. The x-coordinates of the inflection points are the values at which f'' changes sign, that is, -2.16, -0.75, 0.46, and 2.21.

42. $f(x) = e^{-0.1x}\ln(x^2-1) \quad \Rightarrow \quad f'(x) = \dfrac{e^{-0.1x}\left[(x^2-1)\,\ln(x^2-1) - 20x\right]}{10(1-x^2)} \quad \Rightarrow$

$$f''(x) = \frac{e^{-0.1x}\left[(x^2-1)^2\ln(x^2-1) - 40(x^3+5x^2-x+5)\right]}{100(x^2-1)^2}.$$

The domain of f is $(-\infty,-1)\cup(1,\infty)$. $f(x) = 0 \quad \Leftrightarrow \quad x = \pm\sqrt{2}$; $\quad f'(x) = 0 \quad \Leftrightarrow \quad x \approx 5.87$;

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$f''(x) = 0 \quad \Leftrightarrow \quad x \approx -4.31$ and 11.74.

f' changes from positive to negative at $x \approx 5.87$, so 5.87 is the x-coordinate of the maximum point. There is no minimum point. The x-coordinates of the inflection points are the values at which f'' changes sign, that is, -4.31 and 11.74.

43. The family of functions $f(x) = \ln(\sin x + c)$ all have the same period and all have maximum values at $x = \frac{\pi}{2} + 2\pi n$. Since the domain of ln is $(0, \infty)$, f has a graph only if $\sin x + c > 0$ somewhere. Since $-1 \le \sin x \le 1$, this happens if $c > -1$, that is, f has no graph if $c \le -1$. Similarly, if $c > 1$, then $\sin x + c > 0$ and f is continuous on $(-\infty, \infty)$. As c increases, the graph of f is shifted vertically upward and flattens out. If $-1 < c \le 1$, f is defined where $\sin x + c > 0 \quad \Leftrightarrow \quad \sin x > -c \quad \Leftrightarrow$ $\sin^{-1}(-c) < x < \pi - \sin^{-1}(-c)$. Since the period is 2π, the domain of f is $\big(2n\pi + \sin^{-1}(-c), (2n+1)\pi - \sin^{-1}(-c)\big)$, n an integer.

44. We exclude the case $c = 0$, since in that case $f(x) = 0$ for all x. To find the maxima and minima, we differentiate:

$$f(x) = cxe^{-cx^2} \quad \Rightarrow \quad f'(x) = c\Big[xe^{-cx^2}(-2cx) + e^{-cx^2}(1)\Big] = ce^{-cx^2}(-2cx^2 + 1)$$

This is 0 where $-2cx^2 + 1 = 0 \quad \Leftrightarrow \quad x = \pm 1/\sqrt{2c}$. So if $c > 0$, there are two maxima or minima, whose x-coordinates approach 0 as c increases. The negative solution gives a minimum and the positive solution gives a maximum, by the First Derivative Test. By substituting back into the equation, we see that $f\big(\pm 1/\sqrt{2c}\big) = c\big(\pm 1/\sqrt{2c}\big)\, e^{-c\left(\pm 1/\sqrt{2c}\right)^2} = \pm\sqrt{c/2e}$. So as c increases, the extreme points become more pronounced. Note that if $c > 0$, then $\lim\limits_{x \to \pm\infty} f(x) = 0$. If $c < 0$, then there are no extreme values, and $\lim\limits_{x \to \pm\infty} f(x) = \mp\infty$.

To find the points of inflection, we differentiate again: $f'(x) = ce^{-cx^2}\big(-2cx^2 + 1\big) \quad \Rightarrow$ $f''(x) = c\Big[e^{-cx^2}(-4cx) + (-2cx^2 + 1)(-2cxe^{-cx^2})\Big] = -2c^2xe^{-cx^2}(3 - 2cx^2)$. This is 0 at $x = 0$ and where $3 - 2cx^2 = 0 \quad \Leftrightarrow \quad x = \pm\sqrt{3/(2c)} \quad \Rightarrow$ IP at $\Big(\pm\sqrt{3/(2c)}, \pm\sqrt{3c/2}\, e^{-3/2}\Big)$. If $c > 0$ there are three inflection points, and as c increases, the x-coordinates of the nonzero inflection points approach 0. If $c < 0$, there is only one inflection point, the origin.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. Let $f(x) = 3x + 2\cos x + 5$. Then $f(0) = 7 > 0$ and $f(-\pi) = -3\pi - 2 + 5 = -3\pi + 3 = -3(\pi - 1) < 0$, and since f is continuous on $\mathbb{R}$ (hence on $[-\pi, 0]$), the Intermediate Value Theorem assures us that there is at least one solution of f in $[-\pi, 0]$. Now $f'(x) = 3 - 2\sin x > 0$ implies that f is increasing on $\mathbb{R}$, so there is exactly one solution of f, and hence, exactly one real solution of the equation $3x + 2\cos x + 5 = 0$.

46. By the Mean Value Theorem, $f'(c) = \dfrac{f(4) - f(0)}{4 - 0} \quad \Leftrightarrow \quad 4f'(c) = f(4) - 1$ for some c with $0 < c < 4$. Since $2 \le f'(c) \le 5$, we have $4(2) \le 4f'(c) \le 4(5) \quad \Leftrightarrow \quad 4(2) \le f(4) - 1 \le 4(5) \quad \Leftrightarrow \quad 8 \le f(4) - 1 \le 20 \quad \Leftrightarrow$ $9 \le f(4) \le 21$.

47. Since f is continuous on $[32, 33]$ and differentiable on $(32, 33)$, then by the Mean Value Theorem there exists a number c in $(32, 33)$ such that $f'(c) = \frac{1}{5}c^{-4/5} = \dfrac{\sqrt[5]{33} - \sqrt[5]{32}}{33 - 32} = \sqrt[5]{33} - 2$, but $\frac{1}{5}c^{-4/5} > 0 \quad \Rightarrow \quad \sqrt[5]{33} - 2 > 0 \quad \Rightarrow \quad \sqrt[5]{33} > 2$. Also f' is decreasing, so that $f'(c) < f'(32) = \frac{1}{5}(32)^{-4/5} = 0.0125 \quad \Rightarrow \quad 0.0125 > f'(c) = \sqrt[5]{33} - 2 \quad \Rightarrow \quad \sqrt[5]{33} < 2.0125$. Therefore, $2 < \sqrt[5]{33} < 2.0125$.

48. Since the point $(1, 3)$ is on the curve $y = ax^3 + bx^2$, we have $3 = a(1)^3 + b(1)^2 \quad \Rightarrow \quad 3 = a + b$ **(1)**.

$y' = 3ax^2 + 2bx \quad \Rightarrow \quad y'' = 6ax + 2b$. $\quad y'' = 0$ [for inflection points] $\quad \Leftrightarrow \quad x = \dfrac{-2b}{6a} = -\dfrac{b}{3a}$. Since we want $x = 1$, $1 = -\dfrac{b}{3a} \quad \Rightarrow \quad b = -3a$. Combining with **(1)** gives us $3 = a - 3a \quad \Leftrightarrow \quad 3 = -2a \quad \Leftrightarrow \quad a = -\frac{3}{2}$. Hence, $b = -3\left(-\frac{3}{2}\right) = \frac{9}{2}$ and the curve is $y = -\frac{3}{2}x^3 + \frac{9}{2}x^2$.

49. (a) $g(x) = f(x^2) \quad \Rightarrow \quad g'(x) = 2xf'(x^2)$ by the Chain Rule. Since $f'(x) > 0$ for all $x \neq 0$, we must have $f'(x^2) > 0$ for $x \neq 0$, so $g'(x) = 0 \quad \Leftrightarrow \quad x = 0$. Now $g'(x)$ changes sign (from negative to positive) at $x = 0$, since one of its factors, $f'(x^2)$, is positive for all x, and its other factor, $2x$, changes from negative to positive at this point, so by the First Derivative Test, f has a local and absolute minimum at $x = 0$.

(b) $g'(x) = 2xf'(x^2) \quad \Rightarrow \quad g''(x) = 2[xf''(x^2)(2x) + f'(x^2)] = 4x^2 f''(x^2) + 2f'(x^2)$ by the Product Rule and the Chain Rule. But $x^2 > 0$ for all $x \neq 0$, $f''(x^2) > 0$ [since f is CU for $x > 0$], and $f'(x^2) > 0$ for all $x \neq 0$, so since all of its factors are positive, $g''(x) > 0$ for $x \neq 0$. Whether $g''(0)$ is positive or 0 doesn't matter [since the sign of g'' does not change there]; g is concave upward on $\mathbb{R}$.

50. Call the two integers x and y. Then $x + 4y = 1000$, so $x = 1000 - 4y$. Their product is $P = xy = (1000 - 4y)y$, so our problem is to maximize the function $P(y) = 1000y - 4y^2$, where $0 < y < 250$ and y is an integer. $P'(y) = 1000 - 8y$, so $P'(y) = 0 \quad \Leftrightarrow \quad y = 125$. $\quad P''(y) = -8 < 0$, so $P(125) = 62{,}500$ is an absolute maximum. Since the optimal y turned out to be an integer, we have found the desired pair of numbers, namely $x = 1000 - 4(125) = 500$ and $y = 125$.

51. If $B = 0$, the line is vertical and the distance from $x = -\dfrac{C}{A}$ to (x_1, y_1) is $\left|x_1 + \dfrac{C}{A}\right| = \dfrac{|Ax_1 + By_1 + C|}{\sqrt{A^2 + B^2}}$, so assume $B \neq 0$. The square of the distance from (x_1, y_1) to the line is $f(x) = (x - x_1)^2 + (y - y_1)^2$ where $Ax + By + C = 0$, so

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

we minimize $f(x) = (x - x_1)^2 + \left(-\dfrac{A}{B}x - \dfrac{C}{B} - y_1\right)^2 \quad \Rightarrow \quad f'(x) = 2\,(x - x_1) + 2\left(-\dfrac{A}{B}x - \dfrac{C}{B} - y_1\right)\left(-\dfrac{A}{B}\right)$.

$f'(x) = 0 \quad \Rightarrow \quad x = \dfrac{B^2x_1 - ABy_1 - AC}{A^2 + B^2}$ and this gives a minimum since $f''(x) = 2\left(1 + \dfrac{A^2}{B^2}\right) > 0$. Substituting this value of x into $f(x)$ and simplifying gives $f(x) = \dfrac{(Ax_1 + By_1 + C)^2}{A^2 + B^2}$, so the minimum distance is $\sqrt{f(x)} = \dfrac{|Ax_1 + By_1 + C|}{\sqrt{A^2 + B^2}}$.

52. On the hyperbola $xy = 8$, if $d(x)$ is the distance from the point $(x, y) = (x, 8/x)$ to the point $(3, 0)$, then $[d(x)]^2 = (x - 3)^2 + 64/x^2 = f(x)$. $f'(x) = 2(x - 3) - 128/x^3 = 0 \quad \Rightarrow \quad x^4 - 3x^3 - 64 = 0 \quad \Rightarrow$ $(x - 4)\left(x^3 + x^2 + 4x + 16\right) = 0 \quad \Rightarrow \quad x = 4$ since the solution must have $x > 0$. Then $y = \frac{8}{4} = 2$, so the point is $(4, 2)$.

53.

By similar triangles, $\dfrac{y}{x} = \dfrac{r}{\sqrt{x^2 - 2rx}}$, so the area of the triangle is

$$A(x) = \tfrac{1}{2}(2y)x = xy = \frac{rx^2}{\sqrt{x^2 - 2rx}} \quad \Rightarrow$$

$$A'(x) = \frac{2rx\sqrt{x^2 - 2rx} - rx^2(x - r)/\sqrt{x^2 - 2rx}}{x^2 - 2rx} = \frac{rx^2\,(x - 3r)}{(x^2 - 2rx)^{3/2}} = 0$$

when $x = 3r$.

$A'(x) < 0$ when $2r < x < 3r$, $A'(x) > 0$ when $x > 3r$. So $x = 3r$ gives a minimum and $A(3r) = \dfrac{r(9r^2)}{\sqrt{3}\,r} = 3\sqrt{3}\,r^2$.

54.

The volume of the cone is $V = \frac{1}{3}\pi y^2(r + x) = \frac{1}{3}\pi(r^2 - x^2)(r + x),\ -r \le x \le r$.

$$V'(x) = \tfrac{\pi}{3}[(r^2 - x^2)\,(1) + (r + x)(-2x)] = \tfrac{\pi}{3}[(r + x)(r - x - 2x)]$$
$$= \tfrac{\pi}{3}(r + x)(r - 3x) = 0 \text{ when } x = -r \text{ or } x = r/3.$$

Now $V(r) = 0 = V(-r)$, so the maximum occurs at $x = r/3$ and the volume is

$$V\left(\frac{r}{3}\right) = \frac{\pi}{3}\left(r^2 - \frac{r^2}{9}\right)\left(\frac{4r}{3}\right) = \frac{32\pi r^3}{81}.$$

55.

We minimize $L(x) = |PA| + |PB| + |PC| = 2\sqrt{x^2 + 16} + (5 - x)$, $0 \le x \le 5$. $L'(x) = 2x/\sqrt{x^2 + 16} - 1 = 0 \quad \Leftrightarrow \quad 2x = \sqrt{x^2 + 16} \quad \Leftrightarrow$ $4x^2 = x^2 + 16 \quad \Leftrightarrow \quad x = \frac{4}{\sqrt{3}}$. $L(0) = 13$, $L\left(\frac{4}{\sqrt{3}}\right) \approx 11.9$, $L(5) \approx 12.8$, so the minimum occurs when $x = \frac{4}{\sqrt{3}} \approx 2.3$.

56.

If $|CD| = 2$, the last part of $L(x)$ changes from $(5 - x)$ to $(2 - x)$ with $0 \le x \le 2$. But we still get $L'(x) = 0 \quad \Leftrightarrow \quad x = \frac{4}{\sqrt{3}}$, which isn't in the interval $[0, 2]$. Now $L(0) = 10$ and $L(2) = 2\sqrt{20} = 4\sqrt{5} \approx 8.9$. The minimum occurs when $P = C$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. $v = K\sqrt{\dfrac{L}{C} + \dfrac{C}{L}} \quad \Rightarrow \quad \dfrac{dv}{dL} = \dfrac{K}{2\sqrt{(L/C) + (C/L)}}\left(\dfrac{1}{C} - \dfrac{C}{L^2}\right) = 0 \quad \Leftrightarrow \quad \dfrac{1}{C} = \dfrac{C}{L^2} \quad \Leftrightarrow \quad L^2 = C^2 \quad \Leftrightarrow \quad L = C.$

This gives the minimum velocity since $v' < 0$ for $0 < L < C$ and $v' > 0$ for $L > C$.

58.

We minimize the surface area $S = \pi r^2 + 2\pi rh + \frac{1}{2}(4\pi r^2) = 3\pi r^2 + 2\pi rh$.

Solving $V = \pi r^2 h + \frac{2}{3}\pi r^3$ for h, we get $h = \dfrac{V - \frac{2}{3}\pi r^3}{\pi r^2} = \dfrac{V}{\pi r^2} - \frac{2}{3}r$, so

$S(r) = 3\pi r^2 + 2\pi r\left[\dfrac{V}{\pi r^2} - \frac{2}{3}r\right] = \frac{5}{3}\pi r^2 + \dfrac{2V}{r}$.

$S'(r) = -\dfrac{2V}{r^2} + \frac{10}{3}\pi r = \dfrac{\frac{10}{3}\pi r^3 - 2V}{r^2} = 0 \quad \Leftrightarrow \quad \frac{10}{3}\pi r^3 = 2V \quad \Leftrightarrow \quad r^3 = \dfrac{3V}{5\pi} \quad \Leftrightarrow \quad r = \sqrt[3]{\dfrac{3V}{5\pi}}$.

This gives an absolute minimum since $S'(r) < 0$ for $0 < r < \sqrt[3]{\dfrac{3V}{5\pi}}$ and $S'(r) > 0$ for $r > \sqrt[3]{\dfrac{3V}{5\pi}}$. Thus,

$$h = \frac{V - \frac{2}{3}\pi \cdot \dfrac{3V}{5\pi}}{\pi\sqrt[3]{\dfrac{(3V)^2}{(5\pi)^2}}} = \frac{\left(V - \frac{2}{5}V\right)\sqrt[3]{(5\pi)^2}}{\pi\sqrt[3]{(3V)^2}} = \frac{3V\sqrt[3]{(5\pi)^2}}{5\pi\sqrt[3]{(3V)^2}} = \sqrt[3]{\frac{3V}{5\pi}} = r$$

59. Let x denote the number of \$1 decreases in ticket price. Then the ticket price is $\$12 - \$1(x)$, and the average attendance is $11{,}000 + 1000(x)$. Now the revenue per game is

$$\begin{aligned} R(x) &= (\text{price per person}) \times (\text{number of people per game}) \\ &= (12 - x)(11{,}000 + 1000x) = -1000x^2 + 1000x + 132{,}000 \end{aligned}$$

for $0 \le x \le 4$ [since the seating capacity is 15,000] $\Rightarrow$ $R'(x) = -2000x + 1000 = 0 \quad \Leftrightarrow \quad x = 0.5$. This is a maximum since $R''(x) = -2000 < 0$ for all x. Now we must check the value of $R(x) = (12 - x)(11{,}000 + 1000x)$ at $x = 0.5$ and at the endpoints of the domain to see which value of x gives the maximum value of R.

$R(0) = (12)(11{,}000) = 132{,}000$, $R(0.5) = (11.5)(11{,}500) = 132{,}250$, and $R(4) = (8)(15{,}000) = 120{,}000$. Thus, the maximum revenue of \$132,250 per game occurs when the average attendance is 11,500 and the ticket price is \$11.50.

60. (a) $C(x) = 1800 + 25x - 0.2x^2 + 0.001x^3$ and

$R(x) = xp(x) = 48.2x - 0.03x^2$.

The profit is maximized when $C'(x) = R'(x)$.

From the figure, we estimate that the tangents are parallel when $x \approx 160$.

(b) $C'(x) = 25 - 0.4x + 0.003x^2$ and $R'(x) = 48.2 - 0.06x$. $C'(x) = R'(x) \quad \Rightarrow \quad 0.003x^2 - 0.34x - 23.2 = 0 \quad \Rightarrow$ $x_1 \approx 161.3$ $(x > 0)$. $R''(x) = -0.06$ and $C''(x) = -0.4 + 0.006x$, so $R''(x_1) = -0.06 < C''(x_1) \approx 0.57 \quad \Rightarrow$ profit is maximized by producing 161 units.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $c(x) = \dfrac{C(x)}{x} = \dfrac{1800}{x} + 25 - 0.2x + 0.001x^2$ is the average cost. Since the average cost is minimized when the marginal cost equals the average cost, we graph $c(x)$ and $C'(x)$ and estimate the point of intersection. From the figure, $C'(x) = c(x) \quad \Leftrightarrow \quad x \approx 144$.

61. $f(x) = x^5 - x^4 + 3x^2 - 3x - 2 \quad \Rightarrow \quad f'(x) = 5x^4 - 4x^3 + 6x - 3$, so $x_{n+1} = x_n - \dfrac{x_n^5 - x_n^4 + 3x_n^2 - 3x_n - 2}{5x_n^4 - 4x_n^3 + 6x_n - 3}$.

Now $x_1 = 1 \quad \Rightarrow \quad x_2 = 1.5 \quad \Rightarrow \quad x_3 \approx 1.343860 \quad \Rightarrow \quad x_4 \approx 1.300320 \quad \Rightarrow \quad x_5 \approx 1.297396 \quad \Rightarrow$ $x_6 \approx 1.297383 \approx x_7$, so the solution in $[1, 2]$ is 1.297383, to six decimal places.

62. Graphing $y = \sin x$ and $y = x^2 - 3x + 1$ shows that there are two solutions, one about 0.3 and the other about 2.8.

$f(x) = \sin x - x^2 + 3x - 1 \quad \Rightarrow \quad f'(x) = \cos x - 2x + 3 \quad \Rightarrow$

$x_{n+1} = x_n - \dfrac{\sin x_n - x_n^2 + 3x_n - 1}{\cos x_n - 2x_n + 3}$. Now $x_1 = 0.3 \quad \Rightarrow$

$x_2 \approx 0.268552 \quad \Rightarrow \quad x_3 \approx 0.268881 \approx x_4$ and $x_1 = 2.8 \quad \Rightarrow$

$x_2 \approx 2.770354 \quad \Rightarrow \quad x_3 \approx 2.770058 \approx x_4$, so to six decimal places, the solutions are 0.268881 and 2.770058.

63. $f(t) = \cos t + t - t^2 \quad \Rightarrow \quad f'(t) = -\sin t + 1 - 2t$. $f'(t)$ exists for all t, so to find the maximum of f, we can examine the zeros of f'.

From the graph of f', we see that a good choice for t_1 is $t_1 = 0.3$.

Use $g(t) = -\sin t + 1 - 2t$ and $g'(t) = -\cos t - 2$ to obtain $t_2 \approx 0.33535293$, $t_3 \approx 0.33541803 \approx t_4$. Since $f''(t) = -\cos t - 2 < 0$ for all t, $f(0.33541803) \approx 1.16718557$ is the absolute maximum.

64. $y = f(x) = x \sin x, \quad 0 \le x \le 2\pi$. **A.** $D = [0, 2\pi]$ **B.** y-intercept: $f(0) = 0$; x-intercepts: $f(x) = 0 \quad \Leftrightarrow \quad x = 0$ or $\sin x = 0 \quad \Leftrightarrow \quad x = 0, \pi$, or 2π. **C.** There is no symmetry on D, but if f is defined for all real numbers x, then f is an even function. **D.** No asymptote **E.** $f'(x) = x \cos x + \sin x$. To find critical numbers in $(0, 2\pi)$, we graph f' and see that there are two critical numbers, about 2 and 4.9. To find them more precisely, we use Newton's method, setting

$g(x) = f'(x) = x \cos x + \sin x$, so that $g'(x) = f''(x) = 2 \cos x - x \sin x$ and $x_{n+1} = x_n - \dfrac{x_n \cos x_n + \sin x_n}{2 \cos x_n - x_n \sin x_n}$.

$x_1 = 2 \quad \Rightarrow \quad x_2 \approx 2.029048$, $x_3 \approx 2.028758 \approx x_4$ and $x_1 = 4.9 \quad \Rightarrow \quad x_2 \approx 4.913214$, $x_3 \approx 4.913180 \approx x_4$, so the critical numbers, to six decimal places, are $r_1 = 2.028758$ and $r_2 = 4.913180$. By checking sample values of f' in $(0, r_1)$, (r_1, r_2), and $(r_2, 2\pi)$, we see that f is increasing on $(0, r_1)$, decreasing on (r_1, r_2), and increasing on $(r_2, 2\pi)$. **F.** Local maximum value $f(r_1) \approx 1.819706$, local minimum value $f(r_2) \approx -4.814470$. **G.** $f''(x) = 2 \cos x - x \sin x$. To find

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

points where $f''(x) = 0$, we graph f'' and find that $f''(x) = 0$ at about 1 and 3.6. To find the values more precisely, we use Newton's method. Set $h(x) = f''(x) = 2\cos x - x\sin x$. Then $h'(x) = -3\sin x - x\cos x$, so

$x_{n+1} = x_n - \dfrac{2\cos x_n - x_n \sin x_n}{-3\sin x_n - x_n \cos x_n}$. $x_1 = 1 \Rightarrow x_2 \approx 1.078028$, $x_3 \approx 1.076874 \approx x_4$ and $x_1 = 3.6 \Rightarrow$ $x_2 \approx 3.643996$, $x_3 \approx 3.643597 \approx x_4$, so the zeros of f'', to six decimal places, are $r_3 = 1.076874$ and $r_4 = 3.643597$.

By checking sample values of f'' in $(0, r_3)$, (r_3, r_4), and $(r_4, 2\pi)$, we see that f is CU on $(0, r_3)$, CD on (r_3, r_4), and CU on $(r_4, 2\pi)$. f has inflection points at $(r_3, f(r_3) \approx 0.948166)$ and $(r_4, f(r_4) \approx -1.753240)$.

H.

65. $f(x) = 4\sqrt{x} - 6x^2 + 3 = 4x^{1/2} - 6x^2 + 3 \Rightarrow F(x) = 4\left(\frac{2}{3}x^{3/2}\right) - 6\left(\frac{1}{3}x^3\right) + 3x + C = \frac{8}{3}x^{3/2} - 2x^3 + 3x + C$

66. $g(x) = \dfrac{1}{x} + \dfrac{1}{x^2+1} \Rightarrow G(x) = \begin{cases} \ln x + \tan^{-1} x + C_1 & \text{if } x > 0 \\ \ln(-x) + \tan^{-1} x + C_2 & \text{if } x < 0 \end{cases}$

67. $f(t) = 2\sin t - 3e^t \Rightarrow F(t) = -2\cos t - 3e^t + C$

68. $f(x) = x^{-3} + \cosh x \Rightarrow F(x) = \begin{cases} -1/(2x^2) + \sinh x + C_1 & \text{if } x > 0 \\ -1/(2x^2) + \sinh x + C_2 & \text{if } x < 0 \end{cases}$

69. $f'(t) = 2t - 3\sin t \Rightarrow f(t) = t^2 + 3\cos t + C$.

$f(0) = 3 + C$ and $f(0) = 5 \Rightarrow C = 2$, so $f(t) = t^2 + 3\cos t + 2$.

70. $f'(u) = \dfrac{u^2 + \sqrt{u}}{u} = u + u^{-1/2} \Rightarrow f(u) = \frac{1}{2}u^2 + 2u^{1/2} + C$.

$f(1) = \frac{1}{2} + 2 + C$ and $f(1) = 3 \Rightarrow C = \frac{1}{2}$, so $f(u) = \frac{1}{2}u^2 + 2\sqrt{u} + \frac{1}{2}$.

71. $f''(x) = 1 - 6x + 48x^2 \Rightarrow f'(x) = x - 3x^2 + 16x^3 + C$. $f'(0) = C$ and $f'(0) = 2 \Rightarrow C = 2$, so $f'(x) = x - 3x^2 + 16x^3 + 2$ and hence, $f(x) = \frac{1}{2}x^2 - x^3 + 4x^4 + 2x + D$.

$f(0) = D$ and $f(0) = 1 \Rightarrow D = 1$, so $f(x) = \frac{1}{2}x^2 - x^3 + 4x^4 + 2x + 1$.

72. $f''(x) = 5x^3 + 6x^2 + 2 \Rightarrow f'(x) = \frac{5}{4}x^4 + 2x^3 + 2x + C \Rightarrow f(x) = \frac{1}{4}x^5 + \frac{1}{2}x^4 + x^2 + Cx + D$. Now $f(0) = D$ and $f(0) = 3$, so $D = 3$. Also, $f(1) = \frac{1}{4} + \frac{1}{2} + 1 + C + 3 = C + \frac{19}{4}$ and $f(1) = -2$, so $C + \frac{19}{4} = -2 \Rightarrow C = -\frac{27}{4}$. Thus, $f(x) = \frac{1}{4}x^5 + \frac{1}{2}x^4 + x^2 - \frac{27}{4}x + 3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

73. $v(t) = s'(t) = 2t - \dfrac{1}{1+t^2} \quad \Rightarrow \quad s(t) = t^2 - \tan^{-1} t + C.$

$s(0) = 0 - 0 + C = C$ and $s(0) = 1 \quad \Rightarrow \quad C = 1$, so $s(t) = t^2 - \tan^{-1} t + 1.$

74. $a(t) = v'(t) = \sin t + 3\cos t \quad \Rightarrow \quad v(t) = -\cos t + 3\sin t + C.$

$v(0) = -1 + 0 + C$ and $v(0) = 2 \quad \Rightarrow \quad C = 3$, so $v(t) = -\cos t + 3\sin t + 3$ and $s(t) = -\sin t - 3\cos t + 3t + D.$

$s(0) = -3 + D$ and $s(0) = 0 \quad \Rightarrow \quad D = 3$, and $s(t) = -\sin t - 3\cos t + 3t + 3.$

75. (a) Since f is 0 just to the left of the y-axis, we must have a minimum of F at the same place since we are increasing through $(0, 0)$ on F. There must be a local maximum to the left of $x = -3$, since f changes from positive to negative there.

(b) $f(x) = 0.1e^x + \sin x \quad \Rightarrow$

$F(x) = 0.1e^x - \cos x + C$. $F(0) = 0 \quad \Rightarrow$

$0.1 - 1 + C = 0 \quad \Rightarrow \quad C = 0.9$, so

$F(x) = 0.1e^x - \cos x + 0.9.$

(c)

76. $f(x) = x^4 + x^3 + cx^2 \quad \Rightarrow \quad f'(x) = 4x^3 + 3x^2 + 2cx$. This is 0 when $x\big(4x^2 + 3x + 2c\big) = 0 \quad \Leftrightarrow \quad x = 0$ or $4x^2 + 3x + 2c = 0$. Using the quadratic formula, we find that the solutions of this last equation are $x = \dfrac{-3 \pm \sqrt{9 - 32c}}{8}$.

Now if $9 - 32c < 0 \quad \Leftrightarrow \quad c > \frac{9}{32}$, then $(0, 0)$ is the only critical point, a minimum. If $c = \frac{9}{32}$, then there are two critical points (a minimum at $x = 0$, and a horizontal tangent with no maximum or minimum at $x = -\frac{3}{8}$) and if $c < \frac{9}{32}$, then there are three critical points except when $c = 0$, in which case the solution with the $+$ sign coincides with the critical point at $x = 0$.

For $0 < c < \frac{9}{32}$, there is a minimum at $x = -\dfrac{3}{8} - \dfrac{\sqrt{9 - 32c}}{8}$, a maximum at $x = -\dfrac{3}{8} + \dfrac{\sqrt{9 - 32c}}{8}$, and a minimum at $x = 0$. For $c = 0$, there is a minimum at $x = -\frac{3}{4}$ and a horizontal tangent with no extremum at $x = 0$, and for $c < 0$, there is a maximum at $x = 0$, and there are minima at $x = -\dfrac{3}{8} \pm \dfrac{\sqrt{9 - 32c}}{8}$. Now we calculate $f''(x) = 12x^2 + 6x + 2c$.

The solutions of this equation are $x = \dfrac{-6 \pm \sqrt{36 - 4 \cdot 12 \cdot 2c}}{24}$. So if $36 - 96c \le 0 \quad \Leftrightarrow \quad c \ge \frac{3}{8}$, then there is no inflection point. If $c < \frac{3}{8}$, then there are two inflection points at $x = -\dfrac{1}{4} \pm \dfrac{\sqrt{9 - 24c}}{12}$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Value of c	No. of CP	No. of IP
$c < 0$	3	2
$c = 0$	2	2
$0 < c < \frac{9}{32}$	3	2
$c = \frac{9}{32}$	2	2
$\frac{9}{32} < c < \frac{3}{8}$	1	2
$c \geq \frac{3}{8}$	1	0

77. Choosing the positive direction to be upward, we have $a(t) = -9.8 \;\Rightarrow\; v(t) = -9.8t + v_0$, but $v(0) = 0 = v_0 \;\Rightarrow\; v(t) = -9.8t = s'(t) \;\Rightarrow\; s(t) = -4.9t^2 + s_0$, but $s(0) = s_0 = 500 \;\Rightarrow\; s(t) = -4.9t^2 + 500$. When $s = 0$, $-4.9t^2 + 500 = 0 \;\Rightarrow\; t_1 = \sqrt{\frac{500}{4.9}} \approx 10.1 \;\Rightarrow\; v(t_1) = -9.8\sqrt{\frac{500}{4.9}} \approx -98.995$ m/s. Since the canister has been designed to withstand an impact velocity of 100 m/s, the canister will *not burst*.

78. Let $s_A(t)$ and $s_B(t)$ be the position functions for cars A and B and let $f(t) = s_A(t) - s(t)$. Since A passed B twice, there must be three values of t such that $f(t) = 0$. Then by three applications of Rolle's Theorem (see Exercise 4.2.26), there is a number c such that $f''(c) = 0$. So $s_A''(c) = s_B''(c)$; that is, A and B had equal accelerations at $t = c$. We assume that f is continuous on $[0, T]$ and twice differentiable on $(0, T)$, where T is the total time of the race.

79. (a)

The cross-sectional area of the rectangular beam is

$A = 2x \cdot 2y = 4xy = 4x\sqrt{100 - x^2}$, $0 \leq x \leq 10$, so

$$\frac{dA}{dx} = 4x\left(\tfrac{1}{2}\right)(100 - x^2)^{-1/2}(-2x) + (100 - x^2)^{1/2} \cdot 4$$

$$= \frac{-4x^2}{(100 - x^2)^{1/2}} + 4(100 - x^2)^{1/2} = \frac{4[-x^2 + (100 - x^2)]}{(100 - x^2)^{1/2}}.$$

$\dfrac{dA}{dx} = 0$ when $-x^2 + (100 - x^2) = 0 \;\Rightarrow\; x^2 = 50 \;\Rightarrow\; x = \sqrt{50} \approx 7.07 \;\Rightarrow\; y = \sqrt{100 - \left(\sqrt{50}\right)^2} = \sqrt{50}$.

Since $A(0) = A(10) = 0$, the rectangle of maximum area is a square.

(b)

The cross-sectional area of each rectangular plank (shaded in the figure) is

$A = 2x\left(y - \sqrt{50}\right) = 2x\left[\sqrt{100 - x^2} - \sqrt{50}\right]$, $0 \leq x \leq \sqrt{50}$, so

$$\frac{dA}{dx} = 2\left(\sqrt{100 - x^2} - \sqrt{50}\right) + 2x\left(\tfrac{1}{2}\right)(100 - x^2)^{-1/2}(-2x)$$

$$= 2(100 - x^2)^{1/2} - 2\sqrt{50} - \frac{2x^2}{(100 - x^2)^{1/2}}$$

Set $\dfrac{dA}{dx} = 0$: $(100 - x^2) - \sqrt{50}\,(100 - x^2)^{1/2} - x^2 = 0 \;\Rightarrow\; 100 - 2x^2 = \sqrt{50}\,(100 - x^2)^{1/2} \;\Rightarrow$

$10{,}000 - 400x^2 + 4x^4 = 50(100 - x^2) \;\Rightarrow\; 4x^4 - 350x^2 + 5000 = 0 \;\Rightarrow\; 2x^4 - 175x^2 + 2500 = 0 \;\Rightarrow$

$x^2 = \dfrac{175 \pm \sqrt{10{,}625}}{4} \approx 69.52$ or $17.98 \;\Rightarrow\; x \approx 8.34$ or 4.24. But $8.34 > \sqrt{50}$, so $x_1 \approx 4.24 \;\Rightarrow$

$y - \sqrt{50} = \sqrt{100 - x_1^2} - \sqrt{50} \approx 1.99$. Each plank should have dimensions about $8\frac{1}{2}$ inches by 2 inches.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) From the figure in part (a), the width is $2x$ and the depth is $2y$, so the strength is

$S = k(2x)(2y)^2 = 8kxy^2 = 8kx(100 - x^2) = 800kx - 8kx^3,\ 0 \le x \le 10.\ dS/dx = 800k - 24kx^2 = 0$ when

$24kx^2 = 800k \quad \Rightarrow \quad x^2 = \frac{100}{3} \quad \Rightarrow \quad x = \frac{10}{\sqrt{3}} \quad \Rightarrow \quad y = \sqrt{\frac{200}{3}} = \frac{10\sqrt{2}}{\sqrt{3}} = \sqrt{2}\,x$. Since $S(0) = S(10) = 0$, the maximum strength occurs when $x = \frac{10}{\sqrt{3}}$. The dimensions should be $\frac{20}{\sqrt{3}} \approx 11.55$ inches by $\frac{20\sqrt{2}}{\sqrt{3}} \approx 16.33$ inches.

80. (a)

$y = (\tan\theta)x - \dfrac{g}{2v^2\cos^2\theta}x^2$. The parabola intersects the line when

$(\tan\alpha)x = (\tan\theta)x - \dfrac{g}{2v^2\cos^2\theta}x^2 \quad \Rightarrow$

$x = \dfrac{(\tan\theta - \tan\alpha)2v^2\cos^2\theta}{g} \quad \Rightarrow$

$$R(\theta) = \frac{x}{\cos\alpha} = \left(\frac{\sin\theta}{\cos\theta} - \frac{\sin\alpha}{\cos\alpha}\right)\frac{2v^2\cos^2\theta}{g\cos\alpha} = \left(\frac{\sin\theta}{\cos\theta} - \frac{\sin\alpha}{\cos\alpha}\right)(\cos\theta\,\cos\alpha)\,\frac{2v^2\cos\theta}{g\cos^2\alpha}$$

$$= (\sin\theta\,\cos\alpha - \sin\alpha\,\cos\theta)\,\frac{2v^2\cos\theta}{g\cos^2\alpha} = \sin(\theta - \alpha)\,\frac{2v^2\cos\theta}{g\cos^2\alpha}$$

(b) $R'(\theta) = \dfrac{2v^2}{g\cos^2\alpha}\,[\cos\theta\cdot\cos(\theta-\alpha) + \sin(\theta-\alpha)(-\sin\theta)] = \dfrac{2v^2}{g\cos^2\alpha}\cos[\theta + (\theta - \alpha)]$

$= \dfrac{2v^2}{g\cos^2\alpha}\cos(2\theta - \alpha) = 0$

when $\cos(2\theta - \alpha) = 0 \quad \Rightarrow \quad 2\theta - \alpha = \frac{\pi}{2} \quad \Rightarrow \quad \theta = \dfrac{\pi/2 + \alpha}{2} = \dfrac{\pi}{4} + \dfrac{\alpha}{2}$. The First Derivative Test shows that this gives a maximum value for $R(\theta)$. [This could be done without calculus by applying the formula for $\sin x\,\cos y$ to $R(\theta)$.]

(c)

Replacing α by $-\alpha$ in part (a), we get $R(\theta) = \dfrac{2v^2\cos\theta\,\sin(\theta+\alpha)}{g\cos^2\alpha}$.

Proceeding as in part (b), or simply by replacing α by $-\alpha$ in the result of part (b), we see that $R(\theta)$ is maximized when $\theta = \dfrac{\pi}{4} - \dfrac{\alpha}{2}$.

81. $\displaystyle\lim_{E\to 0^+} P(E) = \lim_{E\to 0^+}\left(\frac{e^E + e^{-E}}{e^E - e^{-E}} - \frac{1}{E}\right)$

$$= \lim_{E\to 0^+}\frac{E\big(e^E + e^{-E}\big) - 1\big(e^E - e^{-E}\big)}{\big(e^E - e^{-E}\big)E} = \lim_{E\to 0^+}\frac{Ee^E + Ee^{-E} - e^E + e^{-E}}{Ee^E - Ee^{-E}} \qquad \left[\text{form is } \tfrac{0}{0}\right]$$

$$\overset{\text{H}}{=} \lim_{E\to 0^+}\frac{Ee^E + e^E\cdot 1 + E\big(-e^{-E}\big) + e^{-E}\cdot 1 - e^E + \big(-e^{-E}\big)}{Ee^E + e^E\cdot 1 - [E(-e^{-E}) + e^{-E}\cdot 1]}$$

$$= \lim_{E\to 0^+}\frac{Ee^E - Ee^{-E}}{Ee^E + e^E + Ee^{-E} - e^{-E}} = \lim_{E\to 0^+}\frac{e^E - e^{-E}}{e^E + \dfrac{e^E}{E} + e^{-E} - \dfrac{e^{-E}}{E}} \qquad [\text{divide by } E]$$

$$= \frac{0}{2 + L}, \quad \text{where } L = \lim_{E\to 0^+}\frac{e^E - e^{-E}}{E} \quad \left[\text{form is } \tfrac{0}{0}\right] \quad \overset{\text{H}}{=} \lim_{E\to 0^+}\frac{e^E + e^{-E}}{1} = \frac{1+1}{1} = 2$$

Thus, $\displaystyle\lim_{E\to 0^+} P(E) = \frac{0}{2+2} = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

82. $\lim\limits_{c\to 0^+} s(t) = \lim\limits_{c\to 0^+}\left(\dfrac{m}{c}\ln\cosh\sqrt{\dfrac{gc}{mt}}\right) = m\lim\limits_{c\to 0^+}\dfrac{\ln\cosh\sqrt{ac}}{c}$ [let $a = g/(mt)$]

$$\overset{\text{H}}{=} m\lim_{c\to 0^+}\frac{\dfrac{1}{\cosh\sqrt{ac}}\left(\sinh\sqrt{ac}\right)\left(\dfrac{\sqrt{a}}{2\sqrt{c}}\right)}{1} = \frac{m\sqrt{a}}{2}\lim_{c\to 0^+}\frac{\tanh\sqrt{ac}}{\sqrt{c}}$$

$$\overset{\text{H}}{=} \frac{m\sqrt{a}}{2}\lim_{c\to 0^+}\frac{\operatorname{sech}^2\sqrt{ac}\left[\sqrt{a}/\left(2\sqrt{c}\right)\right]}{1/\left(2\sqrt{c}\right)} = \frac{ma}{2}\lim_{c\to 0^+}\operatorname{sech}^2\sqrt{ac} = \frac{ma}{2}(1)^2 = \frac{mg}{2mt} = \frac{g}{2t}$$

83. We first show that $\dfrac{x}{1+x^2} < \tan^{-1}x$ for $x > 0$. Let $f(x) = \tan^{-1}x - \dfrac{x}{1+x^2}$. Then

$f'(x) = \dfrac{1}{1+x^2} - \dfrac{1(1+x^2)-x(2x)}{(1+x^2)^2} = \dfrac{(1+x^2)-(1-x^2)}{(1+x^2)^2} = \dfrac{2x^2}{(1+x^2)^2} > 0$ for $x > 0$. So $f(x)$ is increasing on $(0,\infty)$. Hence, $0 < x \;\Rightarrow\; 0 = f(0) < f(x) = \tan^{-1}x - \dfrac{x}{1+x^2}$. So $\dfrac{x}{1+x^2} < \tan^{-1}x$ for $0 < x$. We next show that $\tan^{-1}x < x$ for $x > 0$. Let $h(x) = x - \tan^{-1}x$. Then $h'(x) = 1 - \dfrac{1}{1+x^2} = \dfrac{x^2}{1+x^2} > 0$. Hence, $h(x)$ is increasing on $(0,\infty)$. So for $0 < x$, $0 = h(0) < h(x) = x - \tan^{-1}x$. Hence, $\tan^{-1}x < x$ for $x > 0$, and we conclude that $\dfrac{x}{1+x^2} < \tan^{-1}x < x$ for $x > 0$.

84. If $f'(x) < 0$ for all x, $f''(x) > 0$ for $|x| > 1$, $f''(x) < 0$ for $|x| < 1$, and $\lim\limits_{x\to\pm\infty}[f(x)+x] = 0$, then f is decreasing everywhere, concave up on $(-\infty,-1)$ and $(1,\infty)$, concave down on $(-1,1)$, and approaches the line $y = -x$ as $x \to \pm\infty$. An example of such a graph is sketched.

85. Let c be the diameter of the semicircle. Then, from the given figure and the Law of Cosines, $c^2 = a^2 + a^2 - 2a\cdot a\cdot\cos\theta$. The radius of the semicircle is $\frac{1}{2}c$, or $\frac{1}{2}\sqrt{2a^2 - 2a^2\cos\theta}$. The area of the figure is given by

$$\begin{aligned} A(\theta) &= \text{area of triangle} + \text{area of circle} \\ &= \tfrac{1}{2}a\cdot a\cdot\sin\theta + \tfrac{1}{2}\pi\left(\tfrac{1}{2}\sqrt{2a^2-2a^2\cos\theta}\right)^2 = \tfrac{1}{2}a^2\sin\theta + \tfrac{1}{8}\pi(2a^2 - 2a^2\cos\theta) \end{aligned}$$

$A'(\theta) = \frac{1}{2}a^2\cos\theta + \frac{1}{4}\pi a^2\sin\theta = 0 \;\Rightarrow\; \frac{1}{4}\pi a^2\sin\theta = -\frac{1}{2}a^2\cos\theta \;\Rightarrow\; \tan\theta = -\dfrac{2}{\pi} \;\Rightarrow$

$\theta = \tan^{-1}\left(-\dfrac{2}{\pi}\right) + n\pi$ (n an integer). We let $n = 1$ so that $0 \le \theta \le \pi$, giving $\theta = \tan^{-1}\left(-\dfrac{2}{\pi}\right) + \pi \approx 147.5^\circ$.

$A''(\theta) = -\frac{1}{2}a^2\sin\theta + \frac{1}{4}\pi a^2\cos\theta < 0$ [θ. is a second-quadrant angle, so $\sin\theta > 0$ and $\cos\theta < 0$], so this value of θ gives a maximum.

86. (a) $V'(t)$ is the rate of change of the volume of the water with respect to time. $H'(t)$ is the rate of change of the height of the water with respect to time. Since the volume and the height are increasing, $V'(t)$ and $H'(t)$ are positive.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $V'(t)$ is constant, so $V''(t)$ is zero (the slope of a constant function is 0).

(c) At first, the height H of the water increases quickly because the tank is narrow. But as the sphere widens, the rate of increase of the height slows down, reaching a minimum at $t = t_2$. Thus, the height is increasing at a decreasing rate on $(0, t_2)$, so its graph is concave downward and $H''(t_1) < 0$. As the sphere narrows for $t > t_2$, the rate of increase of the height begins to increase, and the graph of H is concave upward. Therefore, $H''(t_2) = 0$ and $H''(t_3) > 0$.

87. (a) $I = \dfrac{k\cos\theta}{d^2} = \dfrac{k(h/d)}{d^2} = k\dfrac{h}{d^3} = k\,\dfrac{h}{\left(\sqrt{40^2+h^2}\,\right)^3} = k\,\dfrac{h}{(1600+h^2)^{3/2}} \quad \Rightarrow$

$$\frac{dI}{dh} = k\,\frac{(1600+h^2)^{3/2} - h\frac{3}{2}(1600+h^2)^{1/2}\cdot 2h}{[(1600+h^2)^{3/2}]^2} = \frac{k(1600+h^2)^{1/2}(1600+h^2-3h^2)}{(1600+h^2)^3}$$

$$= \frac{k(1600-2h^2)}{(1600+h^2)^{5/2}} \qquad [k \text{ is the constant of proportionality}]$$

Set $dI/dh = 0$: $1600 - 2h^2 = 0 \quad \Rightarrow \quad h^2 = 800 \quad \Rightarrow \quad h = \sqrt{800} = 20\sqrt{2}$. By the First Derivative Test, I has a local maximum at $h = 20\sqrt{2} \approx 28$ ft.

(b)

$\dfrac{dx}{dt} = 4$ ft/s

$$I = \frac{k\cos\theta}{d^2} = \frac{k[(h-4)/d]}{d^2} = \frac{k(h-4)}{d^3}$$

$$= \frac{k(h-4)}{[(h-4)^2+x^2]^{3/2}} = k(h-4)\left[(h-4)^2+x^2\right]^{-3/2}$$

$$\frac{dI}{dt} = \frac{dI}{dx}\cdot\frac{dx}{dt} = k(h-4)\left(-\tfrac{3}{2}\right)\left[(h-4)^2+x^2\right]^{-5/2}\cdot 2x\cdot\frac{dx}{dt}$$

$$= k(h-4)(-3x)\left[(h-4)^2+x^2\right]^{-5/2}\cdot 4 = \frac{-12xk(h-4)}{[(h-4)^2+x^2]^{5/2}}$$

$$\left.\frac{dI}{dt}\right|_{x=40} = -\frac{480k(h-4)}{[(h-4)^2+1600]^{5/2}}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1. Let $y = f(x) = e^{-x^2}$. The area of the rectangle under the curve from $-x$ to x is $A(x) = 2xe^{-x^2}$ where $x \geq 0$. We maximize $A(x)$: $A'(x) = 2e^{-x^2} - 4x^2e^{-x^2} = 2e^{-x^2}\left(1 - 2x^2\right) = 0 \quad \Rightarrow \quad x = \frac{1}{\sqrt{2}}$. This gives a maximum since $A'(x) > 0$ for $0 \leq x < \frac{1}{\sqrt{2}}$ and $A'(x) < 0$ for $x > \frac{1}{\sqrt{2}}$. We next determine the points of inflection of $f(x)$. Notice that $f'(x) = -2xe^{-x^2} = -A(x)$. So $f''(x) = -A'(x)$ and hence, $f''(x) < 0$ for $-\frac{1}{\sqrt{2}} < x < \frac{1}{\sqrt{2}}$ and $f''(x) > 0$ for $x < -\frac{1}{\sqrt{2}}$ and $x > \frac{1}{\sqrt{2}}$. So $f(x)$ changes concavity at $x = \pm\frac{1}{\sqrt{2}}$, and the two vertices of the rectangle of largest area are at the inflection points.

2. Let $f(x) = \sin x - \cos x$ on $[0, 2\pi]$ since f has period 2π. $f'(x) = \cos x + \sin x = 0 \quad \Leftrightarrow \quad \cos x = -\sin x \quad \Leftrightarrow$ $\tan x = -1 \quad \Leftrightarrow \quad x = \frac{3\pi}{4}$ or $\frac{7\pi}{4}$. Evaluating f at its critical numbers and endpoints, we get $f(0) = -1$, $f\left(\frac{3\pi}{4}\right) = \sqrt{2}$, $f\left(\frac{7\pi}{4}\right) = -\sqrt{2}$, and $f(2\pi) = -1$. So f has absolute maximum value $\sqrt{2}$ and absolute minimum value $-\sqrt{2}$. Thus, $-\sqrt{2} \leq \sin x - \cos x \leq \sqrt{2} \quad \Rightarrow \quad |\sin x - \cos x| \leq \sqrt{2}$.

3. $f(x)$ has the form $e^{g(x)}$, so it will have an absolute maximum (minimum) where g has an absolute maximum (minimum).

$$g(x) = 10|x-2| - x^2 = \begin{cases} 10(x-2) - x^2 & \text{if } x - 2 > 0 \\ 10[-(x-2)] - x^2 & \text{if } x - 2 < 0 \end{cases} = \begin{cases} -x^2 + 10x - 20 & \text{if } x > 2 \\ -x^2 - 10x + 20 & \text{if } x < 2 \end{cases} \quad \Rightarrow$$

$$g'(x) = \begin{cases} -2x + 10 & \text{if } x > 2 \\ -2x - 10 & \text{if } x < 2 \end{cases}$$

$g'(x) = 0$ if $x = -5$ or $x = 5$, and $g'(2)$ does not exist, so the critical numbers of g are -5, 2, and 5. Since $g''(x) = -2$ for all $x \neq 2$, g is concave downward on $(-\infty, 2)$ and $(2, \infty)$, and g will attain its absolute maximum at one of the critical numbers. Since $g(-5) = 45$, $g(2) = -4$, and $g(5) = 5$, we see that $f(-5) = e^{45}$ is the absolute maximum value of f. Also, $\lim\limits_{x\to\infty} g(x) = -\infty$, so $\lim\limits_{x\to\infty} f(x) = \lim\limits_{x\to\infty} e^{g(x)} = 0$. But $f(x) > 0$ for all x, so there is no absolute minimum value of f.

4. $x^2y^2\left(4 - x^2\right)\left(4 - y^2\right) = x^2\left(4 - x^2\right)y^2\left(4 - y^2\right) = f(x)f(y)$, where $f(t) = t^2\left(4 - t^2\right)$. We will show that $0 \leq f(t) \leq 4$ for $|t| \leq 2$, which gives $0 \leq f(x)f(y) \leq 16$ for $|x| \leq 2$ and $|y| \leq 2$.

$f(t) = 4t^2 - t^4 \quad \Rightarrow \quad f'(t) = 8t - 4t^3 = 4t\left(2 - t^2\right) = 0 \quad \Rightarrow \quad t = 0$ or $\pm\sqrt{2}$.

$f(0) = 0$, $f\left(\pm\sqrt{2}\right) = 2(4-2) = 4$, and $f(2) = 0$. So 0 is the absolute minimum value of $f(t)$ on $[-2, 2]$ and 4 is the absolute maximum value of $f(t)$ on $[-2, 2]$. We conclude that $0 \leq f(t) \leq 4$ for $|t| \leq 2$ and hence, $0 \leq f(x)f(y) \leq 4^2$ or $0 \leq x^2\left(4 - x^2\right)y^2\left(4 - y^2\right) \leq 16$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $y = \dfrac{\sin x}{x} \quad \Rightarrow \quad y' = \dfrac{x\cos x - \sin x}{x^2} \quad \Rightarrow \quad y'' = \dfrac{-x^2\sin x - 2x\cos x + 2\sin x}{x^3}$. If (x, y) is an inflection point, then $y'' = 0 \quad \Rightarrow \quad (2 - x^2)\sin x = 2x\cos x \quad \Rightarrow \quad (2 - x^2)^2\sin^2 x = 4x^2\cos^2 x \quad \Rightarrow$

$(2 - x^2)^2\sin^2 x = 4x^2(1 - \sin^2 x) \quad \Rightarrow \quad (4 - 4x^2 + x^4)\sin^2 x = 4x^2 - 4x^2\sin^2 x \quad \Rightarrow$

$(4 + x^4)\sin^2 x = 4x^2 \quad \Rightarrow \quad (x^4 + 4)\dfrac{\sin^2 x}{x^2} = 4 \quad \Rightarrow \quad y^2(x^4 + 4) = 4$ since $y = \dfrac{\sin x}{x}$.

6. Let $P\big(a, 1 - a^2\big)$ be the point of contact. The equation of the tangent line at P is $y - \big(1 - a^2\big) = (-2a)(x - a) \quad \Rightarrow$

$y - 1 + a^2 = -2ax + 2a^2 \quad \Rightarrow \quad y = -2ax + a^2 + 1$. To find the x-intercept, put $y = 0$: $2ax = a^2 + 1 \quad \Rightarrow$

$x = \dfrac{a^2 + 1}{2a}$. To find the y-intercept, put $x = 0$: $y = a^2 + 1$. Therefore, the area of the triangle is

$\dfrac{1}{2}\left(\dfrac{a^2 + 1}{2a}\right)\big(a^2 + 1\big) = \dfrac{\big(a^2 + 1\big)^2}{4a}$. Therefore, we minimize the function $A(a) = \dfrac{\big(a^2 + 1\big)^2}{4a}$, $a > 0$.

$A'(a) = \dfrac{(4a)2\big(a^2 + 1\big)\,(2a) - \big(a^2 + 1\big)^2(4)}{16a^2} = \dfrac{\big(a^2 + 1\big)[4a^2 - \big(a^2 + 1\big)]}{4a^2} = \dfrac{\big(a^2 + 1\big)\big(3a^2 - 1\big)}{4a^2}$.

$A'(a) = 0$ when $3a^2 - 1 = 0 \quad \Rightarrow \quad a = \frac{1}{\sqrt{3}}$. $A'(a) < 0$ for $a < \frac{1}{\sqrt{3}}$, $A'(a) > 0$ for $a > \frac{1}{\sqrt{3}}$. So by the First Derivative Test, there is an absolute minimum when $a = \frac{1}{\sqrt{3}}$. The required point is $\left(\frac{1}{\sqrt{3}}, \frac{2}{3}\right)$ and the corresponding minimum area is $A\left(\frac{1}{\sqrt{3}}\right) = \frac{4\sqrt{3}}{9}$.

7. Let $L = \lim\limits_{x \to 0} \dfrac{ax^2 + \sin bx + \sin cx + \sin dx}{3x^2 + 5x^4 + 7x^6}$. Now L has the indeterminate form of type $\frac{0}{0}$, so we can apply l'Hospital's Rule. $L = \lim\limits_{x \to 0} \dfrac{2ax + b\cos bx + c\cos cx + d\cos dx}{6x + 20x^3 + 42x^5}$. The denominator approaches 0 as $x \to 0$, so the numerator must also approach 0 (because the limit exists). But the numerator approaches $0 + b + c + d$, so $b + c + d = 0$. Apply l'Hospital's Rule again. $L = \lim\limits_{x \to 0} \dfrac{2a - b^2\sin bx - c^2\sin cx - d^2\sin dx}{6 + 60x^2 + 210x^4} = \dfrac{2a - 0}{6 + 0} = \dfrac{2a}{6}$, which must equal 8.

$\dfrac{2a}{6} = 8 \quad \Rightarrow \quad a = 24$. Thus, $a + b + c + d = a + (b + c + d) = 24 + 0 = 24$.

8. We first present some preliminary results that we will invoke when calculating the limit.

(1) If $y = (1 + ax)^x$, then $\ln y = x\ln(1 + ax)$, and $\lim\limits_{x \to 0^+} \ln y = \lim\limits_{x \to 0^+} x\ln(1 + ax) = 0$. Thus, $\lim\limits_{x \to 0^+} (1 + ax)^x = e^0 = 1$.

(2) If $y = (1 + ax)^x$, then $\ln y = x\ln(1 + ax)$, and implicitly differentiating gives us $\dfrac{y'}{y} = x \cdot \dfrac{a}{1 + ax} + \ln(1 + ax) \quad \Rightarrow$

$y' = y\left[\dfrac{ax}{1 + ax} + \ln(1 + ax)\right]$. Thus, $y = (1 + ax)^x \quad \Rightarrow \quad y' = (1 + ax)^x\left[\dfrac{ax}{1 + ax} + \ln(1 + ax)\right]$.

(3) If $y = \dfrac{ax}{1 + ax}$, then $y' = \dfrac{(1 + ax)a - ax(a)}{(1 + ax)^2} = \dfrac{a + a^2x - a^2x}{(1 + ax)^2} = \dfrac{a}{(1 + ax)^2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\lim_{x\to\infty}\frac{(x+2)^{1/x}-x^{1/x}}{(x+3)^{1/x}-x^{1/x}}=\lim_{x\to\infty}\frac{x^{1/x}[(1+2/x)^{1/x}-1]}{x^{1/x}[(1+3/x)^{1/x}-1]} \qquad \text{[factor out } x^{1/x}\text{]}$$

$$=\lim_{x\to\infty}\frac{(1+2/x)^{1/x}-1}{(1+3/x)^{1/x}-1}$$

$$=\lim_{t\to0^+}\frac{(1+2t)^t-1}{(1+3t)^t-1} \qquad \text{[let } t=1/x\text{, form } 0/0 \text{ by } \textbf{(1)}\text{]}$$

$$\overset{\text{H}}{=}\lim_{t\to0^+}\frac{(1+2t)^t\left[\dfrac{2t}{1+2t}+\ln(1+2t)\right]}{(1+3t)^t\left[\dfrac{3t}{1+3t}+\ln(1+3t)\right]} \qquad \text{[by } \textbf{(2)}\text{]}$$

$$=\lim_{t\to0^+}\frac{(1+2t)^t}{(1+3t)^t}\cdot\lim_{t\to0^+}\frac{\dfrac{2t}{1+2t}+\ln(1+2t)}{\dfrac{3t}{1+3t}+\ln(1+3t)}$$

$$=\frac{1}{1}\cdot\lim_{t\to0^+}\frac{\dfrac{2t}{1+2t}+\ln(1+2t)}{\dfrac{3t}{1+3t}+\ln(1+3t)} \qquad \text{[by } \textbf{(1)}\text{, now form } 0/0\text{]}$$

$$\overset{\text{H}}{=}\lim_{t\to0^+}\frac{\dfrac{2}{(1+2t)^2}+\dfrac{2}{1+2t}}{\dfrac{3}{(1+3t)^2}+\dfrac{3}{1+3t}} \qquad \text{[by } \textbf{(3)}\text{]}$$

$$=\frac{2+2}{3+3}=\frac{4}{6}=\frac{2}{3}$$

9. Differentiating $x^2+xy+y^2=12$ implicitly with respect to x gives $2x+y+x\dfrac{dy}{dx}+2y\dfrac{dy}{dx}=0$, so $\dfrac{dy}{dx}=-\dfrac{2x+y}{x+2y}$.

At a highest or lowest point, $\dfrac{dy}{dx}=0 \quad\Leftrightarrow\quad y=-2x$. Substituting $-2x$ for y in the original equation gives $x^2+x(-2x)+(-2x)^2=12$, so $3x^2=12$ and $x=\pm2$. If $x=2$, then $y=-2x=-4$, and if $x=-2$ then $y=4$. Thus, the highest and lowest points are $(-2,4)$ and $(2,-4)$.

10. Since

$$\frac{f(x+n)-f(x)}{n}=f'(x) \qquad \textbf{(1)}$$

holds for all real numbers x and all positive integers n, we have

$$\frac{f(x+n)-f(x)}{n}=\frac{f(x+2n)-f(x)}{2n}$$

for every real number x. It follows that

$$f(x+2n)-2f(x+n)=-f(x) \qquad \textbf{(2)}$$

Now, again from **(1)**, we can write

$$nf'(x)=f(x+n)-f(x)$$

The right-hand side of this equation is differentiable by hypothesis, so the left-hand side is also differentiable. Differentiating

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and then using **(1)** again — twice this time — we get

$$\begin{aligned} nf''(x) &= f'(x+n) - f'(x) \\ &= \frac{f(x+2n) - f(x+n)}{n} - \frac{f(x+n) - f(x)}{n} = \frac{f(x+2n) - 2f(x+n) + f(x)}{n} \end{aligned}$$

Rearranging this last equation and simplifying using **(2)**, we get

$$n^2 f''(x) = f(x+2n) - 2f(x+n) + f(x) = -f(x) + f(x) = 0$$

Thus, $f''(x) = 0$ for all x, so f is a linear function.

11. (a) $y = x^2 \quad \Rightarrow \quad y' = 2x$, so the slope of the tangent line at $P(a, a^2)$ is $2a$ and the slope of the normal line is $-\dfrac{1}{2a}$ for $a \neq 0$. An equation of the normal line is $y - a^2 = -\dfrac{1}{2a}(x - a)$. Substitute x^2 for y to find the x-coordinates of the two points of intersection of the parabola and the normal line. $x^2 - a^2 = -\dfrac{x}{2a} + \dfrac{1}{2} \quad \Leftrightarrow \quad x^2 + \left(\dfrac{1}{2a}\right)x - \dfrac{1}{2} - a^2 = 0$. We know that a is a root of this quadratic equation, so $x - a$ is a factor, and we have $(x - a)\left(x + \dfrac{1}{2a} + a\right) = 0$, and hence, $x = -a - \dfrac{1}{2a}$ is the x-coordinate of the point Q. We want to minimize the y-coordinate of Q, which is $\left(-a - \dfrac{1}{2a}\right)^2 = a^2 + 1 + \dfrac{1}{4a^2} = y(a)$. Now $y'(a) = 2a - \dfrac{1}{2a^3} = \dfrac{4a^4 - 1}{2a^3} = \dfrac{(2a^2+1)(2a^2-1)}{2a^3} = 0 \quad \Rightarrow$ $a = \dfrac{1}{\sqrt{2}}$ for $a > 0$. Since $y''(a) = 2 + \dfrac{3}{2a^4} > 0$, we see that $a = \dfrac{1}{\sqrt{2}}$ gives us the minimum value of the y-coordinate of Q.

(b) The square S of the distance from $P(a, a^2)$ to $Q\left(-a - \dfrac{1}{2a}, \left(-a - \dfrac{1}{2a}\right)^2\right)$ is given by

$$\begin{aligned} S &= \left(-a - \frac{1}{2a} - a\right)^2 + \left[\left(-a - \frac{1}{2a}\right)^2 - a^2\right]^2 = \left(-2a - \frac{1}{2a}\right)^2 + \left[\left(a^2 + 1 + \frac{1}{4a^2}\right) - a^2\right]^2 \\ &= \left(4a^2 + 2 + \frac{1}{4a^2}\right) + \left(1 + \frac{1}{4a^2}\right)^2 = \left(4a^2 + 2 + \frac{1}{4a^2}\right) + 1 + \frac{2}{4a^2} + \frac{1}{16a^4} \\ &= 4a^2 + 3 + \frac{3}{4a^2} + \frac{1}{16a^4} \end{aligned}$$

$S' = 8a - \dfrac{6}{4a^3} - \dfrac{4}{16a^5} = 8a - \dfrac{3}{2a^3} - \dfrac{1}{4a^5} = \dfrac{32a^6 - 6a^2 - 1}{4a^5} = \dfrac{(2a^2-1)(4a^2+1)^2}{4a^5}$. The only real positive zero of the equation $S' = 0$ is $a = \dfrac{1}{\sqrt{2}}$. Since $S'' = 8 + \dfrac{9}{2a^4} + \dfrac{5}{4a^6} > 0$, $a = \dfrac{1}{\sqrt{2}}$ corresponds to the shortest possible length of the line segment PQ.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12.

$y = cx^3 + e^x \quad \Rightarrow \quad y' = 3cx^2 + e^x \quad \Rightarrow \quad y'' = 6cx + e^x$. The curve will have inflection points when y'' changes sign. $y'' = 0 \quad \Rightarrow \quad -6cx = e^x$, so y'' will change sign when the line $y = -6cx$ intersects the curve $y = e^x$ (but is not tangent to it). Note that if $c = 0$, the curve is just $y = e^x$, which has no inflection point.

The first figure shows that for $c > 0$, $y = -6cx$ will intersect $y = e^x$ once, so $y = cx^3 + e^x$ will have one inflection point.

The second figure shows that for $c < 0$, the line $y = -6cx$ can intersect the curve $y = e^x$ in two points (two inflection points), be tangent to it (no inflection point), or not intersect it (no inflection point). The tangent line at (a, e^a) has slope e^a, but from the diagram we see that the slope is $\dfrac{e^a}{a}$. So $\dfrac{e^a}{a} = e^a \quad \Rightarrow \quad a = 1$. Thus, the slope is e. The line $y = -6cx$ must have slope greater than e, so $-6c > e \quad \Rightarrow \quad c < -e/6$.

Therefore, the curve $y = cx^3 + e^x$ will have one inflection point if $c > 0$ and two inflection points if $c < -e/6$.

13.

$\overline{AC}$ is tangent to the unit circle at D. To find the slope of $\overline{AC}$ at D, use implicit differentiation. $x^2 + y^2 = 1 \quad \Rightarrow \quad 2x + 2y\,y' = 0 \quad \Rightarrow \quad y\,y' = -x \quad \Rightarrow \quad y' = -\dfrac{x}{y}$. Thus, the tangent line at $D(b, c)$ has equation $y = -\dfrac{b}{c}\,x + a$. At D, $x = b$ and $y = c$, so $c = -\dfrac{b}{c}(b) + a \quad \Rightarrow \quad a = c + \dfrac{b^2}{c} = \dfrac{c^2 + b^2}{c} = \dfrac{1}{c}$, and hence $c = \dfrac{1}{a}$.

Since $b^2 + c^2 = 1$, $b = \sqrt{1 - c^2} = \sqrt{1 - 1/a^2} = \sqrt{\dfrac{a^2 - 1}{a^2}} = \dfrac{\sqrt{a^2 - 1}}{a}$, and now we have both b and c in terms of a. At C, $y = -1$, so $-1 = -\dfrac{b}{c}\,x + a \quad \Rightarrow \quad \dfrac{b}{c}\,x = a + 1 \quad \Rightarrow$

$x = \dfrac{c}{b}(a + 1) = \dfrac{1/a}{\sqrt{a^2 - 1}/a}(a + 1) = \dfrac{a + 1}{\sqrt{(a + 1)(a - 1)}} = \sqrt{\dfrac{a + 1}{a - 1}}$, and C has coordinates $\left(\sqrt{\dfrac{a + 1}{a - 1}}, -1\right)$. Let S be the square of the distance from A to C. Then $S(a) = \left(0 - \sqrt{\dfrac{a + 1}{a - 1}}\right)^2 + (a + 1)^2 = \dfrac{a + 1}{a - 1} + (a + 1)^2 \quad \Rightarrow$

$$S'(a) = \frac{(a - 1)(1) - (a + 1)(1)}{(a - 1)^2} + 2(a + 1) = \frac{-2 + 2(a + 1)(a - 1)^2}{(a - 1)^2}$$

$$= \frac{-2 + 2(a^3 - a^2 - a + 1)}{(a - 1)^2} = \frac{2a^3 - 2a^2 - 2a}{(a - 1)^2} = \frac{2a(a^2 - a - 1)}{(a - 1)^2}$$

Using the quadratic formula, we find that the solutions of $a^2 - a - 1 = 0$ are $a = \dfrac{1 \pm \sqrt{5}}{2}$, so $a_1 = \dfrac{1 + \sqrt{5}}{2}$ (the "golden mean") since $a > 0$. For $1 < a < a_1$, $S'(a) < 0$, and for $a > a_1$, $S'(a) > 0$, so a_1 minimizes S.

Note: The minimum length of the equal sides is $\sqrt{S(a_1)} = \cdots = \sqrt{\dfrac{11 + 5\sqrt{5}}{2}} \approx 3.33$ and the corresponding length of the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

third side is $2\sqrt{\dfrac{a_1+1}{a_1-1}} = \cdots = 2\sqrt{2+\sqrt{5}} \approx 4.12$, so the triangle is *not* equilateral.

Another method: In ΔABC, $\cos\theta = \dfrac{a+1}{\overline{AC}}$, so $\overline{AC} = \dfrac{a+1}{\cos\theta}$. In ΔADO, $\sin\theta = \dfrac{1}{a}$, so $\cos\theta = \sqrt{1-\sin^2\theta} = \sqrt{1-1/a^2} = \dfrac{1}{a}\sqrt{a^2-1}$. Thus $\overline{AC} = \dfrac{a+1}{(1/a)\sqrt{a^2-1}} = \dfrac{a(a+1)}{\sqrt{a^2-1}} = f(a)$. Now find the minimum of f.

14. To sketch the region $\{(x,y) \mid 2xy \le |x-y| \le x^2+y^2\}$, we consider two cases.

Case 1: $x \ge y$ This is the case in which (x,y) lies on or below the line $y=x$. The double inequality becomes $2xy \le x-y \le x^2+y^2$. The right-hand inequality holds if and only if $x^2-x+y^2+y \ge 0 \quad \Leftrightarrow$ $\left(x-\frac{1}{2}\right)^2+\left(y+\frac{1}{2}\right)^2 \ge \frac{1}{2} \quad \Leftrightarrow \quad (x,y)$ lies on or outside the circle with radius $\frac{1}{\sqrt{2}}$ centered at $\left(\frac{1}{2},-\frac{1}{2}\right)$. The left-hand inequality holds if and only if $2xy-x+y \le 0 \quad \Leftrightarrow \quad xy-\frac{1}{2}x+\frac{1}{2}y \le 0 \quad \Leftrightarrow$ $\left(x+\frac{1}{2}\right)\left(y-\frac{1}{2}\right) \le -\frac{1}{4} \quad \Leftrightarrow \quad (x,y)$ lies on or below the hyperbola $\left(x+\frac{1}{2}\right)\left(y-\frac{1}{2}\right) = -\frac{1}{4}$, which passes through the origin and approaches the lines $y=\frac{1}{2}$ and $x=-\frac{1}{2}$ asymptotically.

Case 2: $y \ge x$ This is the case in which (x,y) lies on or above the line $y=x$. The double inequality becomes $2xy \le y-x \le x^2+y^2$. The right-hand inequality holds if and only if $x^2+x+y^2-y \ge 0 \quad \Leftrightarrow$ $\left(x+\frac{1}{2}\right)^2+\left(y-\frac{1}{2}\right)^2 \ge \frac{1}{2} \quad \Leftrightarrow \quad (x,y)$ lies on or outside the circle of radius $\frac{1}{\sqrt{2}}$ centered at $\left(-\frac{1}{2},\frac{1}{2}\right)$. The left-hand inequality holds if and only if $2xy+x-y \le 0 \quad \Leftrightarrow \quad xy+\frac{1}{2}x-\frac{1}{2}y \le 0 \quad \Leftrightarrow \quad \left(x-\frac{1}{2}\right)\left(y+\frac{1}{2}\right) \le -\frac{1}{4} \quad \Leftrightarrow \quad (x,y)$ lies on or above the left-hand branch of the hyperbola $\left(x-\frac{1}{2}\right)\left(y+\frac{1}{2}\right) = -\frac{1}{4}$, which passes through the origin and approaches the lines $y=-\frac{1}{2}$ and $x=\frac{1}{2}$ asymptotically. Therefore, the region of interest consists of the points on or above the left branch of the hyperbola $\left(x-\frac{1}{2}\right)\left(y+\frac{1}{2}\right) = -\frac{1}{4}$ that are on or outside the circle $\left(x+\frac{1}{2}\right)^2+\left(y-\frac{1}{2}\right)^2 = \frac{1}{2}$, together with the points on or below the right branch of the hyperbola $\left(x+\frac{1}{2}\right)\left(y-\frac{1}{2}\right) = -\frac{1}{4}$ that are on or outside the circle $\left(x-\frac{1}{2}\right)^2+\left(y+\frac{1}{2}\right)^2 = \frac{1}{2}$. Note that the inequalities are unchanged when x and y are interchanged, so the region is symmetric about the line $y=x$. So we need only have analyzed case 1 and then reflected that region about the line $y=x$, instead of considering case 2.

15. $A=\left(x_1,x_1^2\right)$ and $B=\left(x_2,x_2^2\right)$, where x_1 and x_2 are the solutions of the quadratic equation $x^2=mx+b$. Let $P=\left(x,x^2\right)$ and set $A_1=(x_1,0)$, $B_1=(x_2,0)$, and $P_1=(x,0)$. Let $f(x)$ denote the area of triangle PAB. Then $f(x)$ can be expressed in terms of the areas of three trapezoids as follows:

$$\begin{aligned} f(x) &= \text{area}\,(A_1ABB_1) - \text{area}\,(A_1APP_1) - \text{area}\,(B_1BPP_1) \\ &= \tfrac{1}{2}\left(x_1^2+x_2^2\right)(x_2-x_1) - \tfrac{1}{2}\left(x_1^2+x^2\right)(x-x_1) - \tfrac{1}{2}\left(x^2+x_2^2\right)(x_2-x) \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

After expanding and canceling terms, we get

$f(x) = \frac{1}{2}\left(x_2x_1^2 - x_1x_2^2 - xx_1^2 + x_1x^2 - x_2x^2 + xx_2^2\right) = \frac{1}{2}\left[x_1^2(x_2 - x) + x_2^2(x - x_1) + x^2(x_1 - x_2)\right]$

$f'(x) = \frac{1}{2}\left[-x_1^2 + x_2^2 + 2x(x_1 - x_2)\right]$. $\quad f''(x) = \frac{1}{2}[2(x_1 - x_2)] = x_1 - x_2 < 0$ since $x_2 > x_1$.

$f'(x) = 0 \quad \Rightarrow \quad 2x(x_1 - x_2) = x_1^2 - x_2^2 \quad \Rightarrow \quad x_P = \frac{1}{2}(x_1 + x_2)$.

$$\begin{aligned} f(x_P) &= \tfrac{1}{2}\left(x_1^2\left[\tfrac{1}{2}(x_2 - x_1)\right] + x_2^2\left[\tfrac{1}{2}(x_2 - x_1)\right] + \tfrac{1}{4}(x_1 + x_2)^2(x_1 - x_2)\right) \\ &= \tfrac{1}{2}\left[\tfrac{1}{2}(x_2 - x_1)\left(x_1^2 + x_2^2\right) - \tfrac{1}{4}(x_2 - x_1)(x_1 + x_2)^2\right] = \tfrac{1}{8}(x_2 - x_1)\left[2\left(x_1^2 + x_2^2\right) - \left(x_1^2 + 2x_1x_2 + x_2^2\right)\right] \\ &= \tfrac{1}{8}(x_2 - x_1)\left(x_1^2 - 2x_1x_2 + x_2^2\right) = \tfrac{1}{8}(x_2 - x_1)(x_1 - x_2)^2 = \tfrac{1}{8}(x_2 - x_1)(x_2 - x_1)^2 = \tfrac{1}{8}(x_2 - x_1)^3 \end{aligned}$$

To put this in terms of m and b, we solve the system $y = x_1^2$ and $y = mx_1 + b$, giving us $x_1^2 - mx_1 - b = 0 \quad \Rightarrow$ $x_1 = \frac{1}{2}\left(m - \sqrt{m^2 + 4b}\right)$. Similarly, $x_2 = \frac{1}{2}\left(m + \sqrt{m^2 + 4b}\right)$. The area is then $\frac{1}{8}(x_2 - x_1)^3 = \frac{1}{8}\left(\sqrt{m^2 + 4b}\right)^3$, and is attained at the point $P\left(x_P, x_P^2\right) = P\left(\frac{1}{2}m, \frac{1}{4}m^2\right)$.

Note: Another way to get an expression for $f(x)$ is to use the formula for an area of a triangle in terms of the coordinates of the vertices: $f(x) = \frac{1}{2}\left[\left(x_2x_1^2 - x_1x_2^2\right) + \left(x_1x^2 - xx_1^2\right) + \left(xx_2^2 - x_2x^2\right)\right]$.

16. Let $x = |AE|$, $y = |AF|$ as shown. The area $\mathcal{A}$ of the ΔAEF is $\mathcal{A} = \frac{1}{2}xy$. We need to find a relationship between x and y, so that we can take the derivative $d\mathcal{A}/dx$ and then find the maximum and minimum areas. Now let A' be the point on which A ends up after the fold has been performed, and let P be the intersection of AA' and EF. Note that AA' is perpendicular to EF since we are reflecting A through the line EF to get to A', and that $|AP| = |PA'|$ for the same reason.

But $|AA'| = 1$, since AA' is a radius of the circle. Since $|AP| + |PA'| = |AA'|$, we have $|AP| = \frac{1}{2}$. Another way to express the area of the triangle is $\mathcal{A} = \frac{1}{2}|EF|\,|AP| = \frac{1}{2}\sqrt{x^2 + y^2}\left(\frac{1}{2}\right) = \frac{1}{4}\sqrt{x^2 + y^2}$. Equating the two expressions for $\mathcal{A}$, we get $\frac{1}{2}xy = \frac{1}{4}\sqrt{x^2 + y^2} \quad \Rightarrow \quad 4x^2y^2 = x^2 + y^2 \quad \Rightarrow \quad y^2\left(4x^2 - 1\right) = x^2 \quad \Rightarrow \quad y = x/\sqrt{4x^2 - 1}$.

(Note that we could also have derived this result from the similarity of $\triangle A'PE$ and $\triangle A'FE$; that is,

$\dfrac{|A'P|}{|PE|} = \dfrac{|A'F|}{|A'E|} \quad \Rightarrow \quad \dfrac{\frac{1}{2}}{\sqrt{x^2 - \left(\frac{1}{2}\right)^2}} = \dfrac{y}{x} \quad \Rightarrow \quad y = \dfrac{\frac{1}{2}x}{\sqrt{4x^2 - 1}/2} = \dfrac{x}{\sqrt{4x^2 - 1}}$.) Now we can substitute for y and calculate $\dfrac{d\mathcal{A}}{dx}$: $\mathcal{A} = \dfrac{1}{2}\dfrac{x^2}{\sqrt{4x^2 - 1}} \quad \Rightarrow \quad \dfrac{d\mathcal{A}}{dx} = \dfrac{1}{2}\left[\dfrac{\sqrt{4x^2 - 1}\,(2x) - x^2\left(\frac{1}{2}\right)\left(4x^2 - 1\right)^{-1/2}(8x)}{4x^2 - 1}\right]$. This is 0 when

$2x\sqrt{4x^2 - 1} - 4x^3\left(4x^2 - 1\right)^{-1/2} = 0 \quad \Leftrightarrow \quad 2x\left(4x^2 - 1\right)^{-1/2}\left[\left(4x^2 - 1\right) - 2x^2\right] = 0 \quad \Rightarrow \quad \left(4x^2 - 1\right) - 2x^2 = 0$ $(x > 0) \quad \Leftrightarrow \quad 2x^2 = 1 \quad \Rightarrow \quad x = \frac{1}{\sqrt{2}}$. So this is one possible value for an extremum. We must also test the endpoints of the interval over which x ranges. The largest value that x can attain is 1, and the smallest value of x occurs when $y = 1 \quad \Leftrightarrow$ $1 = x/\sqrt{4x^2 - 1} \quad \Leftrightarrow \quad x^2 = 4x^2 - 1 \quad \Leftrightarrow \quad 3x^2 = 1 \quad \Leftrightarrow \quad x = \frac{1}{\sqrt{3}}$. This will give the same value of $\mathcal{A}$ as will $x = 1$, since the geometric situation is the same (reflected through the line $y = x$). We calculate

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathcal{A}\left(\frac{1}{\sqrt{2}}\right) = \dfrac{1}{2}\dfrac{\left(1/\sqrt{2}\right)^2}{\sqrt{4\left(1/\sqrt{2}\right)^2 - 1}} = \dfrac{1}{4} = 0.25$, and $\mathcal{A}(1) = \dfrac{1}{2}\dfrac{1^2}{\sqrt{4(1)^2 - 1}} = \dfrac{1}{2\sqrt{3}} \approx 0.29$. So the maximum area is $\mathcal{A}(1) = \mathcal{A}\left(\frac{1}{\sqrt{3}}\right) = \frac{1}{2\sqrt{3}}$ and the minimum area is $\mathcal{A}\left(\frac{1}{\sqrt{2}}\right) = \frac{1}{4}$.

Another method: Use the angle θ (see diagram above) as a variable:

$\mathcal{A} = \frac{1}{2}xy = \frac{1}{2}\left(\frac{1}{2}\sec\theta\right)\left(\frac{1}{2}\csc\theta\right) = \dfrac{1}{8\sin\theta\,\cos\theta} = \dfrac{1}{4\sin 2\theta}$. $\mathcal{A}$ is minimized when $\sin 2\theta$ is maximal, that is, when $\sin\,2\theta = 1 \;\Rightarrow\; 2\theta = \frac{\pi}{2} \;\Rightarrow\; \theta = \frac{\pi}{4}$. Also note that $A'E = x = \frac{1}{2}\sec\theta \le 1 \;\Rightarrow\; \sec\theta \le 2 \;\Rightarrow$ $\cos\theta \ge \frac{1}{2} \;\Rightarrow\; \theta \le \frac{\pi}{3}$, and similarly, $A'F = y = \frac{1}{2}\csc\theta \le 1 \;\Rightarrow\; \csc\theta \le 2 \;\Rightarrow\; \sin\theta \le \frac{1}{2} \;\Rightarrow\; \theta \ge \frac{\pi}{6}$.

As above, we find that $\mathcal{A}$ is maximized at these endpoints: $\mathcal{A}\left(\frac{\pi}{6}\right) = \dfrac{1}{4\sin\frac{\pi}{3}} = \dfrac{1}{2\sqrt{3}} = \dfrac{1}{4\sin\frac{2\pi}{3}} = \mathcal{A}\left(\frac{\pi}{3}\right)$;

and minimized at $\theta = \frac{\pi}{4}$: $\mathcal{A}\left(\frac{\pi}{4}\right) = \dfrac{1}{4\sin\frac{\pi}{2}} = \dfrac{1}{4}$.

17. Suppose that the curve $y = a^x$ intersects the line $y = x$. Then $a^{x_0} = x_0$ for some $x_0 > 0$, and hence $a = x_0^{1/x_0}$. We find the maximum value of $g(x) = x^{1/x}$, $x > 0$, because if a is larger than the maximum value of this function, then the curve $y = a^x$ does not intersect the line $y = x$. $g'(x) = e^{(1/x)\ln x}\left(-\dfrac{1}{x^2}\ln x + \dfrac{1}{x}\cdot\dfrac{1}{x}\right) = x^{1/x}\left(\dfrac{1}{x^2}\right)(1 - \ln x)$. This is 0 only where $x = e$, and for $0 < x < e$, $f'(x) > 0$, while for $x > e$, $f'(x) < 0$, so g has an absolute maximum of $g(e) = e^{1/e}$. So if $y = a^x$ intersects $y = x$, we must have $0 < a \le e^{1/e}$. Conversely, suppose that $0 < a \le e^{1/e}$. Then $a^e \le e$, so the graph of $y = a^x$ lies below or touches the graph of $y = x$ at $x = e$. Also $a^0 = 1 > 0$, so the graph of $y = a^x$ lies above that of $y = x$ at $x = 0$. Therefore, by the Intermediate Value Theorem, the graphs of $y = a^x$ and $y = x$ must intersect somewhere between $x = 0$ and $x = e$.

18. If $L = \lim\limits_{x\to\infty}\left(\dfrac{x+a}{x-a}\right)^x$, then L has the indeterminate form 1^∞, so

$$\ln L = \lim_{x\to\infty}\ln\left(\frac{x+a}{x-a}\right)^x = \lim_{x\to\infty} x\ln\left(\frac{x+a}{x-a}\right) = \lim_{x\to\infty}\frac{\ln(x+a) - \ln(x-a)}{1/x} \overset{\text{H}}{=} \lim_{x\to\infty}\frac{\dfrac{1}{x+a} - \dfrac{1}{x-a}}{-1/x^2}$$

$$= \lim_{x\to\infty}\left[\frac{(x-a)-(x+a)}{(x+a)(x-a)}\cdot\frac{-x^2}{1}\right] = \lim_{x\to\infty}\frac{2ax^2}{x^2 - a^2} = \lim_{x\to\infty}\frac{2a}{1 - a^2/x^2} = 2a$$

Hence, $\ln L = 2a$, so $L = e^{2a}$. From the original equation, we want $L = e^1 \;\Rightarrow\; 2a = 1 \;\Rightarrow\; a = \frac{1}{2}$.

19. Note that $f(0) = 0$, so for $x \ne 0$, $\left|\dfrac{f(x) - f(0)}{x - 0}\right| = \left|\dfrac{f(x)}{x}\right| = \dfrac{|f(x)|}{|x|} \le \dfrac{|\sin x|}{|x|} = \dfrac{\sin x}{x}$.

Therefore, $|f'(0)| = \left|\lim\limits_{x\to 0}\dfrac{f(x) - f(0)}{x - 0}\right| = \lim\limits_{x\to 0}\left|\dfrac{f(x) - f(0)}{x - 0}\right| \le \lim\limits_{x\to 0}\dfrac{\sin x}{x} = 1$.

But $f(x) = a_1\sin x + a_2\sin 2x + \cdots + a_n\sin nx \;\Rightarrow\; f'(x) = a_1\cos x + 2a_2\cos 2x + \cdots + na_n\cos nx$, so $|f'(0)| = |a_1 + 2a_2 + \cdots + na_n| \le 1$.

Another solution: We are given that $\left|\sum_{k=1}^n a_k\sin kx\right| \le |\sin x|$. So for x close to 0, and $x \ne 0$, we have

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\left|\sum_{k=1}^{n} a_k \dfrac{\sin kx}{\sin x}\right| \le 1 \quad\Rightarrow\quad \lim_{x\to 0}\left|\sum_{k=1}^{n} a_k \dfrac{\sin kx}{\sin x}\right| \le 1 \quad\Rightarrow\quad \left|\sum_{k=1}^{n} a_k \lim_{x\to 0}\dfrac{\sin kx}{\sin x}\right| \le 1$. But by l'Hospital's Rule,

$\lim_{x\to 0}\dfrac{\sin kx}{\sin x} = \lim_{x\to 0}\dfrac{k\cos kx}{\cos x} = k$, so $\left|\sum_{k=1}^{n} ka_k\right| \le 1$.

20. Let the circle have radius r, so $|OP| = |OQ| = r$, where O is the center of the circle. Now $\angle POR$ has measure $\frac{1}{2}\theta$, and $\angle OPR$ is a right angle, so $\tan\frac{1}{2}\theta = \dfrac{|PR|}{r}$ and the area of $\triangle OPR$ is $\frac{1}{2}|OP|\,|PR| = \frac{1}{2}r^2\tan\frac{1}{2}\theta$. The area of the sector cut by OP and OR is $\frac{1}{2}r^2\left(\frac{1}{2}\theta\right) = \frac{1}{4}r^2\theta$. Let S be the intersection of PQ and OR. Then $\sin\frac{1}{2}\theta = \dfrac{|PS|}{r}$ and $\cos\frac{1}{2}\theta = \dfrac{|OS|}{r}$, and the area of $\triangle OSP$ is $\frac{1}{2}|OS|\,|PS| = \frac{1}{2}\left(r\cos\frac{1}{2}\theta\right)\left(r\sin\frac{1}{2}\theta\right) = \frac{1}{2}r^2\sin\frac{1}{2}\theta\,\cos\frac{1}{2}\theta = \frac{1}{4}r^2\sin\theta$.

So $B(\theta) = 2\left(\frac{1}{2}r^2\tan\frac{1}{2}\theta - \frac{1}{4}r^2\theta\right) = r^2\left(\tan\frac{1}{2}\theta - \frac{1}{2}\theta\right)$ and $A(\theta) = 2\left(\frac{1}{4}r^2\theta - \frac{1}{4}r^2\sin\theta\right) = \frac{1}{2}r^2(\theta - \sin\theta)$. Thus,

$$\begin{aligned}
\lim_{\theta\to 0^+}\frac{A(\theta)}{B(\theta)} &= \lim_{\theta\to 0^+}\frac{\frac{1}{2}r^2(\theta-\sin\theta)}{r^2\left(\tan\frac{1}{2}\theta - \frac{1}{2}\theta\right)} = \lim_{\theta\to 0^+}\frac{\theta-\sin\theta}{2\left(\tan\frac{1}{2}\theta-\frac{1}{2}\theta\right)} \overset{\text{H}}{=} \lim_{\theta\to 0^+}\frac{1-\cos\theta}{2\left(\frac{1}{2}\sec^2\frac{1}{2}\theta - \frac{1}{2}\right)} \\
&= \lim_{\theta\to 0^+}\frac{1-\cos\theta}{\sec^2\frac{1}{2}\theta - 1} = \lim_{\theta\to 0^+}\frac{1-\cos\theta}{\tan^2\frac{1}{2}\theta} \overset{\text{H}}{=} \lim_{\theta\to 0^+}\frac{\sin\theta}{2\left(\tan\frac{1}{2}\theta\right)\left(\sec^2\frac{1}{2}\theta\right)\frac{1}{2}} \\
&= \lim_{\theta\to 0^+}\frac{\sin\theta\cos^3\frac{1}{2}\theta}{\sin\frac{1}{2}\theta} = \lim_{\theta\to 0^+}\frac{\left(2\sin\frac{1}{2}\theta\,\cos\frac{1}{2}\theta\right)\cos^3\frac{1}{2}\theta}{\sin\frac{1}{2}\theta} = 2\lim_{\theta\to 0^+}\cos^4\left(\tfrac{1}{2}\theta\right) = 2(1)^4 = 2
\end{aligned}$$

21. (a) Distance = rate × time, so time = distance/rate. $T_1 = \dfrac{D}{c_1}$, $T_2 = \dfrac{2\,|PR|}{c_1} + \dfrac{|RS|}{c_2} = \dfrac{2h\sec\theta}{c_1} + \dfrac{D - 2h\tan\theta}{c_2}$,

$T_3 = \dfrac{2\sqrt{h^2 + D^2/4}}{c_1} = \dfrac{\sqrt{4h^2 + D^2}}{c_1}$.

(b) $\dfrac{dT_2}{d\theta} = \dfrac{2h}{c_1}\cdot\sec\theta\,\tan\theta - \dfrac{2h}{c_2}\sec^2\theta = 0$ when $2h\sec\theta\left(\dfrac{1}{c_1}\tan\theta - \dfrac{1}{c_2}\sec\theta\right) = 0 \quad\Rightarrow$

$\dfrac{1}{c_1}\dfrac{\sin\theta}{\cos\theta} - \dfrac{1}{c_2}\dfrac{1}{\cos\theta} = 0 \quad\Rightarrow\quad \dfrac{\sin\theta}{c_1\cos\theta} = \dfrac{1}{c_2\cos\theta} \quad\Rightarrow\quad \sin\theta = \dfrac{c_1}{c_2}$. The First Derivative Test shows that this gives a minimum.

(c) Using part (a) with $D = 1$ and $T_1 = 0.26$, we have $T_1 = \dfrac{D}{c_1} \quad\Rightarrow\quad c_1 = \frac{1}{0.26} \approx 3.85$ km/s. $T_3 = \dfrac{\sqrt{4h^2 + D^2}}{c_1} \quad\Rightarrow$

$4h^2 + D^2 = T_3^2c_1^2 \quad\Rightarrow\quad h = \frac{1}{2}\sqrt{T_3^2c_1^2 - D^2} = \frac{1}{2}\sqrt{(0.34)^2(1/0.26)^2 - 1^2} \approx 0.42$ km. To find c_2, we use $\sin\theta = \dfrac{c_1}{c_2}$ from part (b) and $T_2 = \dfrac{2h\sec\theta}{c_1} + \dfrac{D - 2h\tan\theta}{c_2}$ from part (a). From the figure,

$\sin\theta = \dfrac{c_1}{c_2} \quad\Rightarrow\quad \sec\theta = \dfrac{c_2}{\sqrt{c_2^2 - c_1^2}}$ and $\tan\theta = \dfrac{c_1}{\sqrt{c_2^2 - c_1^2}}$, so

$T_2 = \dfrac{2hc_2}{c_1\sqrt{c_2^2 - c_1^2}} + \dfrac{D\sqrt{c_2^2 - c_1^2} - 2hc_1}{c_2\sqrt{c_2^2 - c_1^2}}$. Using the values for T_2 [given as 0.32], h, c_1, and D, we can graph $\text{Y}_1 = T_2$ and $\text{Y}_2 = \dfrac{2hc_2}{c_1\sqrt{c_2^2 - c_1^2}} + \dfrac{D\sqrt{c_2^2 - c_1^2} - 2hc_1}{c_2\sqrt{c_2^2 - c_1^2}}$ and find their intersection

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

points. Doing so gives us $c_2 \approx 4.10$ and 7.66, but if $c_2 = 4.10$, then $\theta = \arcsin(c_1/c_2) \approx 69.6°$, which implies that point S is to the left of point R in the diagram. So $c_2 = 7.66$ km/s.

22. A straight line intersects the curve $y = f(x) = x^4 + cx^3 + 12x^2 - 5x + 2$ in four distinct points if and only if the graph of f has two inflection points. $f'(x) = 4x^3 + 3cx^2 + 24x - 5$ and $f''(x) = 12x^2 + 6cx + 24$.

$f''(x) = 0 \quad \Leftrightarrow \quad x = \dfrac{-6c \pm \sqrt{(6c)^2 - 4(12)(24)}}{2(12)}$. There are two distinct roots for $f''(x) = 0$ (and hence two inflection points) if and only if the discriminant is positive; that is, $36c^2 - 1152 > 0 \quad \Leftrightarrow \quad c^2 > 32 \quad \Leftrightarrow \quad |c| > \sqrt{32}$. Thus, the desired values of c are $c < -4\sqrt{2}$ or $c > 4\sqrt{2}$.

23.

Let $a = |EF|$ and $b = |BF|$ as shown in the figure.

Since $\ell = |BF| + |FD|$, $|FD| = \ell - b$. Now

$$\begin{aligned} |ED| &= |EF| + |FD| = a + \ell - b \\ &\sqrt{r^2 - x^2} + \ell - \sqrt{(d-x)^2 + a^2} \\ &= \sqrt{r^2 - x^2} + \ell - \sqrt{(d-x)^2 + \left(\sqrt{r^2 - x^2}\right)^2} \\ &= \sqrt{r^2 - x^2} + \ell - \sqrt{d^2 - 2dx + x^2 + r^2 - x^2} \end{aligned}$$

Let $f(x) = \sqrt{r^2 - x^2} + \ell - \sqrt{d^2 + r^2 - 2dx}$.

$f'(x) = \frac{1}{2}\left(r^2 - x^2\right)^{-1/2}(-2x) - \frac{1}{2}\left(d^2 + r^2 - 2dx\right)^{-1/2}(-2d) = \dfrac{-x}{\sqrt{r^2 - x^2}} + \dfrac{d}{\sqrt{d^2 + r^2 - 2dx}}$.

$f'(x) = 0 \quad \Rightarrow \quad \dfrac{x}{\sqrt{r^2 - x^2}} = \dfrac{d}{\sqrt{d^2 + r^2 - 2dx}} \quad \Rightarrow \quad \dfrac{x^2}{r^2 - x^2} = \dfrac{d^2}{d^2 + r^2 - 2dx} \quad \Rightarrow$

$d^2x^2 + r^2x^2 - 2dx^3 = d^2r^2 - d^2x^2 \quad \Rightarrow \quad 0 = 2dx^3 - 2d^2x^2 - r^2x^2 + d^2r^2 \quad \Rightarrow$

$0 = 2dx^2(x - d) - r^2\left(x^2 - d^2\right) \quad \Rightarrow \quad 0 = 2dx^2(x - d) - r^2(x + d)(x - d) \quad \Rightarrow \quad 0 = (x - d)\left[2dx^2 - r^2(x + d)\right]$

But $d > r > x$, so $x \neq d$. Thus, we solve $2dx^2 - r^2x - dr^2 = 0$ for x:

$x = \dfrac{-\left(-r^2\right) \pm \sqrt{\left(-r^2\right)^2 - 4(2d)(-dr^2)}}{2(2d)} = \dfrac{r^2 \pm \sqrt{r^4 + 8d^2r^2}}{4d}$. Because $\sqrt{r^4 + 8d^2r^2} > r^2$, the "negative" can be discarded. Thus, $x = \dfrac{r^2 + \sqrt{r^2}\sqrt{r^2 + 8d^2}}{4d} = \dfrac{r^2 + r\sqrt{r^2 + 8d^2}}{4d} \quad [r > 0] \quad = \dfrac{r}{4d}\left(r + \sqrt{r^2 + 8d^2}\right)$. The maximum value of $|ED|$ occurs at this value of x.

24.

Let $a = \overline{CD}$ denote the distance from the center C of the base to the midpoint D of a side of the base.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Since ΔPQR is similar to ΔDCR, $\dfrac{a}{h} = \dfrac{r}{\sqrt{h(h-2r)}} \quad \Rightarrow \quad a = \dfrac{rh}{\sqrt{h(h-2r)}} = r\dfrac{\sqrt{h}}{\sqrt{h-2r}}$.

Let b denote one-half the length of a side of the base. The area A of the base is

$A = 8(\text{area of } \Delta CDE) = 8\left(\frac{1}{2}ab\right) = 4a\left(a\tan\frac{\pi}{4}\right) = 4a^2$.

The volume of the pyramid is $V = \frac{1}{3}Ah = \frac{1}{3}\left(4a^2\right)h = \dfrac{4}{3}\left(r\dfrac{\sqrt{h}}{\sqrt{h-2r}}\right)^2 h = \frac{4}{3}r^2\dfrac{h^2}{h-2r}$, with domain $h > 2r$.

Now $\dfrac{dV}{dh} = \dfrac{4}{3}r^2 \cdot \dfrac{(h-2r)(2h) - h^2(1)}{(h-2r)^2} = \dfrac{4}{3}r^2\dfrac{h^2-4hr}{(h-2r)^2} = \dfrac{4}{3}r^2\dfrac{h(h-4r)}{(h-2r)^2}$

and
$$\begin{aligned}\frac{d^2V}{dh^2} &= \frac{4}{3}r^2 \cdot \frac{(h-2r)^2(2h-4r) - \left(h^2-4hr\right)(2)(h-2r)(1)}{\left[(h-2r)^2\right]^2}\\ &= \frac{4}{3}r^2 \cdot \frac{2(h-2r)\left[\left(h^2-4hr+4r^2\right) - \left(h^2-4hr\right)\right]}{(h-2r)^2}\\ &= \frac{8}{3}r^2 \cdot \frac{4r^2}{(h-2r)^3} = \frac{32}{3}r^4 \cdot \frac{1}{(h-2r)^3}.\end{aligned}$$

The first derivative is equal to zero for $h = 4r$ and the second derivative is positive for $h > 2r$, so the volume of the pyramid is minimized when $h = 4r$.

To extend our solution to a regular n-gon, we make the following changes:

(1) the number of sides of the base is n

(2) the number of triangles in the base is $2n$

(3) $\angle DCE = \dfrac{\pi}{n}$

(4) $b = a\tan\dfrac{\pi}{n}$

We then obtain the following results: $A = na^2\tan\dfrac{\pi}{n}$, $V = \dfrac{nr^2}{3}\cdot\tan\left(\dfrac{\pi}{n}\right)\cdot\dfrac{h^2}{h-2r}$, $\dfrac{dV}{dh} = \dfrac{nr^2}{3}\cdot\tan\left(\dfrac{\pi}{n}\right)\cdot\dfrac{h(h-4r)}{(h-2r)^2}$, and $\dfrac{d^2V}{dh^2} = \dfrac{8nr^4}{3}\cdot\tan\left(\dfrac{\pi}{n}\right)\cdot\dfrac{1}{(h-2r)^3}$. Notice that the answer, $h = 4r$, is independent of the number of sides of the base of the polygon!

25. $V = \frac{4}{3}\pi r^3 \quad \Rightarrow \quad \dfrac{dV}{dt} = 4\pi r^2\dfrac{dr}{dt}$. But $\dfrac{dV}{dt}$ is proportional to the surface area, so $\dfrac{dV}{dt} = k\cdot 4\pi r^2$ for some constant k.

Therefore, $4\pi r^2\dfrac{dr}{dt} = k\cdot 4\pi r^2 \quad \Leftrightarrow \quad \dfrac{dr}{dt} = k =$ constant. An antiderivative of k with respect to t is kt, so $r = kt + C$.

When $t = 0$, the radius r must equal the original radius r_0, so $C = r_0$, and $r = kt + r_0$. To find k we use the fact that when $t = 3$, $r = 3k + r_0$ and $V = \frac{1}{2}V_0 \quad \Rightarrow \quad \frac{4}{3}\pi(3k+r_0)^3 = \frac{1}{2}\cdot\frac{4}{3}\pi r_0^3 \quad \Rightarrow \quad (3k+r_0)^3 = \frac{1}{2}r_0^3 \quad \Rightarrow$

$3k + r_0 = \frac{1}{\sqrt[3]{2}}r_0 \quad \Rightarrow \quad k = \frac{1}{3}r_0\left(\frac{1}{\sqrt[3]{2}} - 1\right)$. Since $r = kt + r_0$, $r = \frac{1}{3}r_0\left(\frac{1}{\sqrt[3]{2}} - 1\right)t + r_0$. When the snowball

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

has melted completely we have $r = 0 \quad \Rightarrow \quad \frac{1}{3}r_0\left(\frac{1}{\sqrt[3]{2}} - 1\right)t + r_0 = 0$ which gives $t = \dfrac{3\sqrt[3]{2}}{\sqrt[3]{2} - 1}$. Hence, it takes

$\dfrac{3\sqrt[3]{2}}{\sqrt[3]{2} - 1} - 3 = \dfrac{3}{\sqrt[3]{2} - 1} \approx 11$ h 33 min longer.

26. By ignoring the bottom hemisphere of the initial spherical bubble, we can rephrase the problem as follows: Prove that the maximum height of a stack of n hemispherical bubbles is $\sqrt{n}$ if the radius of the bottom hemisphere is 1. We proceed by induction. The case $n = 1$ is obvious since $\sqrt{1}$ is the height of the first hemisphere. Suppose the assertion is true for $n = k$ and let's suppose we have $k + 1$ hemispherical bubbles forming a stack of maximum height. Suppose the second hemisphere (counting from the bottom) has radius r. Then by our induction hypothesis (scaled to the setting of a bottom hemisphere of radius r), the height of the stack formed by the top k bubbles is $\sqrt{k}\,r$. (If it were shorter, then the total stack of $k + 1$ bubbles wouldn't have maximum height.)

The height of the whole stack is $H(r) = \sqrt{k}\,r + \sqrt{1 - r^2}$. (See the figure.) We want to choose r so as to maximize $H(r)$. Note that $0 < r < 1$.

We calculate $H'(r) = \sqrt{k} - \dfrac{r}{\sqrt{1 - r^2}}$ and $H''(r) = \dfrac{-1}{(1 - r^2)^{3/2}}$.

$H'(r) = 0 \quad \Leftrightarrow \quad r^2 = k(1 - r^2) \quad \Leftrightarrow \quad (k + 1)r^2 = k \quad \Leftrightarrow \quad r = \sqrt{\dfrac{k}{k + 1}}$.

This is the only critical number in $(0, 1)$ and it represents a local maximum (hence an absolute maximum) since $H''(r) < 0$ on $(0, 1)$. When $r = \sqrt{\dfrac{k}{k + 1}}$,

$H(r) = \sqrt{k}\dfrac{\sqrt{k}}{\sqrt{k + 1}} + \sqrt{1 - \dfrac{k}{k + 1}} = \dfrac{k}{\sqrt{k + 1}} + \dfrac{1}{\sqrt{k + 1}} = \sqrt{k + 1}$. Thus, the assertion is true for $n = k + 1$ when it is true for $n = k$. By induction, it is true for all positive integers n.

Note: In general, a maximally tall stack of n hemispherical bubbles consists of bubbles with radii

$$1, \sqrt{\frac{n-1}{n}}, \sqrt{\frac{n-2}{n}}, \ldots, \sqrt{\frac{2}{n}}, \sqrt{\frac{1}{n}}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5 ☐ INTEGRALS

5.1 The Area and Distance Problems

1. (a) Since f is *decreasing*, we can obtain a *lower* estimate by using *right* endpoints. We are instructed to use five rectangles, so $n = 5$.

$$R_5 = \sum_{i=1}^{5} f(x_i)\,\Delta x \qquad \left[\Delta x = \frac{b-a}{n} = \frac{10-0}{5} = 2\right]$$
$$= f(x_1)\cdot 2 + f(x_2)\cdot 2 + f(x_3)\cdot 2 + f(x_4)\cdot 2 + f(x_5)\cdot 2$$
$$= 2[f(2) + f(4) + f(6) + f(8) + f(10)]$$
$$\approx 2(3.2 + 1.8 + 0.8 + 0.2 + 0)$$
$$= 2(6) = 12$$

Since f is *decreasing*, we can obtain an *upper* estimate by using *left* endpoints.

$$L_5 = \sum_{i=1}^{5} f(x_{i-1})\,\Delta x$$
$$= f(x_0)\cdot 2 + f(x_1)\cdot 2 + f(x_2)\cdot 2 + f(x_3)\cdot 2 + f(x_4)\cdot 2$$
$$= 2[f(0) + f(2) + f(4) + f(6) + f(8)]$$
$$\approx 2(5 + 3.2 + 1.8 + 0.8 + 0.2)$$
$$= 2(11) = 22$$

(b) $R_{10} = \sum_{i=1}^{10} f(x_i)\,\Delta x \qquad \left[\Delta x = \frac{10-0}{10} = 1\right]$

$$= 1[f(x_1) + f(x_2) + \cdots + f(x_{10})]$$
$$= f(1) + f(2) + \cdots + f(10)$$
$$\approx 4 + 3.2 + 2.5 + 1.8 + 1.3 + 0.8 + 0.5 + 0.2 + 0.1 + 0$$
$$= 14.4$$

$$L_{10} = \sum_{i=1}^{10} f(x_{i-1})\,\Delta x$$
$$= f(0) + f(1) + \cdots + f(9)$$
$$= R_{10} + 1\cdot f(0) - 1\cdot f(10) \qquad \left[\begin{array}{c}\text{add leftmost upper rectangle,}\\ \text{subtract rightmost lower rectangle}\end{array}\right]$$
$$= 14.4 + 5 - 0$$
$$= 19.4$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. (a) (i) $L_6 = \sum_{i=1}^{6} f(x_{i-1})\,\Delta x \qquad [\Delta x = \frac{12-0}{6} = 2]$

$$= 2[f(x_0) + f(x_1) + f(x_2) + f(x_3) + f(x_4) + f(x_5)]$$
$$= 2[f(0) + f(2) + f(4) + f(6) + f(8) + f(10)]$$
$$\approx 2(9 + 8.8 + 8.2 + 7.3 + 5.9 + 4.1)$$
$$= 2(43.3) = 86.6$$

(ii) $R_6 = L_6 + 2 \cdot f(12) - 2 \cdot f(0)$

$$\approx 86.6 + 2(1) - 2(9) = 70.6$$

[Add area of rightmost lower rectangle and subtract area of leftmost upper rectangle.]

(iii) $M_6 = \sum_{i=1}^{6} f(x_i)\,\Delta x$

$$= 2[f(1) + f(3) + f(5) + f(7) + f(9) + f(11)]$$
$$\approx 2(8.9 + 8.5 + 7.8 + 6.6 + 5.1 + 2.8)$$
$$= 2(39.7) = 79.4$$

(b) Since f is *decreasing*, we obtain an *overestimate* by using *left* endpoints; that is, L_6.

(c) Since f is *decreasing*, we obtain an *underestimate* by using *right* endpoints; that is, R_6.

(d) M_6 gives the best estimate, since the area of each rectangle appears to be closer to the true area than the overestimates and underestimates in L_6 and R_6.

3. (a) $R_4 = \sum_{i=1}^{4} f(x_i)\,\Delta x \qquad \left[\Delta x = \frac{2-1}{4} = \frac{1}{4}\right] = \left[\sum_{i=1}^{4} f(x_i)\right]\Delta x$

$$= [f(x_1) + f(x_2) + f(x_3) + f(x_4)]\ \Delta x$$
$$= \left[\frac{1}{5/4} + \frac{1}{6/4} + \frac{1}{7/4} + \frac{1}{8/4}\right]\frac{1}{4} = \left[\tfrac{4}{5} + \tfrac{2}{3} + \tfrac{4}{7} + \tfrac{1}{2}\right]\tfrac{1}{4} \approx 0.6345$$

Since f is *decreasing* on $[1, 2]$, an *underestimate* is obtained by using the *right* endpoint approximation, R_4.

(b) $L_4 = \sum_{i=1}^{4} f(x_{i-1})\,\Delta x = \left[\sum_{i=1}^{4} f(x_{i-1})\right]\Delta x$

$$= [f(x_0) + f(x_1) + f(x_2) + f(x_3)]\ \Delta x$$
$$= \left[\frac{1}{1} + \frac{1}{5/4} + \frac{1}{6/4} + \frac{1}{7/4}\right]\frac{1}{4} = \left[1 + \tfrac{4}{5} + \tfrac{2}{3} + \tfrac{4}{7}\right]\tfrac{1}{4} \approx 0.7595$$

L_4 is an overestimate. Alternatively, we could just add the area of the leftmost upper rectangle and subtract the area of the rightmost lower rectangle; that is, $L_4 = R_4 + f(1) \cdot \frac{1}{4} - f(2) \cdot \frac{1}{4}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a) $R_4 = \sum_{i=1}^{4} f(x_i)\ \Delta x \qquad \left[\Delta x = \dfrac{\pi/2 - 0}{4} = \dfrac{\pi}{8}\right] = \left[\sum_{i=1}^{4} f(x_i)\right] \Delta x$

$= [f(x_1) + f(x_2) + f(x_3) + f(x_4)]\ \Delta x$

$= \left[\sin \frac{\pi}{8} + \sin \frac{2\pi}{8} + \sin \frac{3\pi}{8} + \sin \frac{4\pi}{8}\right] \frac{\pi}{8}$

≈ 1.1835

Since f is *increasing* on $\left[0, \frac{\pi}{2}\right]$, R_4 is an *overestimate*.

(b) $L_4 = \sum_{i=1}^{4} f(x_{i-1})\ \Delta x = \left[\sum_{i=1}^{4} f(x_{i-1})\right] \Delta x$

$= [f(x_0) + f(x_1) + f(x_2) + f(x_3)]\ \Delta x$

$= \left[\sin 0 + \sin \frac{\pi}{8} + \sin \frac{2\pi}{8} + \sin \frac{3\pi}{8}\right] \frac{\pi}{8}$

≈ 0.7908

Since f is *increasing* on $\left[0, \frac{\pi}{2}\right]$, L_4 is an *underestimate*.

5. (a) $f(x) = 1 + x^2$ and $\Delta x = \dfrac{2-(-1)}{3} = 1 \quad \Rightarrow$

$R_3 = 1 \cdot f(0) + 1 \cdot f(1) + 1 \cdot f(2) = 1 \cdot 1 + 1 \cdot 2 + 1 \cdot 5 = 8.$

$\Delta x = \dfrac{2-(-1)}{6} = 0.5 \quad \Rightarrow$

$R_6 = 0.5[f(-0.5) + f(0) + f(0.5) + f(1) + f(1.5) + f(2)]$

$= 0.5(1.25 + 1 + 1.25 + 2 + 3.25 + 5)$

$= 0.5(13.75) = 6.875$

(b) $L_3 = 1 \cdot f(-1) + 1 \cdot f(0) + 1 \cdot f(1) = 1 \cdot 2 + 1 \cdot 1 + 1 \cdot 2 = 5$

$L_6 = 0.5[f(-1) + f(-0.5) + f(0) + f(0.5) + f(1) + f(1.5)]$

$= 0.5(2 + 1.25 + 1 + 1.25 + 2 + 3.25)$

$= 0.5(10.75) = 5.375$

(c) $M_3 = 1 \cdot f(-0.5) + 1 \cdot f(0.5) + 1 \cdot f(1.5)$

$= 1 \cdot 1.25 + 1 \cdot 1.25 + 1 \cdot 3.25 = 5.75$

$M_6 = 0.5[f(-0.75) + f(-0.25) + f(0.25)$

$+\ f(0.75) + f(1.25) + f(1.75)]$

$= 0.5(1.5625 + 1.0625 + 1.0625 + 1.5625 + 2.5625 + 4.0625)$

$= 0.5(11.875) = 5.9375$

(d) M_6 appears to be the best estimate.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. (a) $f(x) = e^{x-x^2},\ 0 \le x \le 2$

(b) $f(x) = e^{x-x^2}$ and $\Delta x = \dfrac{2-0}{4} = \dfrac{1}{2} \quad \Rightarrow$

(i) $R_4 = \frac{1}{2} \cdot f\left(\frac{1}{2}\right) + \frac{1}{2} \cdot f(1) + \frac{1}{2} \cdot f\left(\frac{3}{2}\right) + \frac{1}{2} \cdot f(2)$

$= \frac{1}{2}\left[f\left(\frac{1}{2}\right) + f(1) + f\left(\frac{3}{2}\right) + f(2)\right]$

$= \frac{1}{2}\left[e^{1/2-(1/2)^2} + e^{1-1^2} + e^{3/2-(3/2)^2} + e^{2-2^2}\right]$

≈ 1.446

(ii) $M_4 = \frac{1}{2} \cdot f\left(\frac{1}{4}\right) + \frac{1}{2} \cdot f\left(\frac{3}{4}\right) + \frac{1}{2} \cdot f\left(\frac{5}{4}\right) + \frac{1}{2} \cdot f\left(\frac{7}{4}\right)$

$= \frac{1}{2}\left[f\left(\frac{1}{4}\right) + f\left(\frac{3}{4}\right) + f\left(\frac{5}{4}\right) + f\left(\frac{7}{4}\right)\right]$

$= \frac{1}{2}\left[e^{1/4-(1/4)^2} + e^{3/4-(3/4)^2} + e^{5/4-(5/4)^2} + e^{7/4-(7/4)^2}\right]$

≈ 1.707

(c) $f(x) = e^{x-x^2}$ and $\Delta x = \dfrac{2-0}{8} = \dfrac{1}{4} \quad \Rightarrow$

(i) $R_8 = \frac{1}{4}\left[f\left(\frac{1}{4}\right) + f\left(\frac{1}{2}\right) + \cdots + f(2)\right]$

$= \frac{1}{4}\left[e^{1/4-(1/4)^2} + e^{1/2-(1/2)^2} + \cdots + e^{2-2^2}\right]$

≈ 1.576

(ii) $M_8 = \frac{1}{4}\left[f\left(\frac{1}{8}\right) + f\left(\frac{3}{8}\right) + \cdots + f\left(\frac{15}{8}\right)\right]$

$= \frac{1}{4}\left[e^{1/8-(1/8)^2} + e^{3/8-(3/8)^2} + \cdots + e^{15/8-(15/8)^2}\right]$

≈ 1.695

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. $f(x) = 6 - x^2,\ -2 \le x \le 2,\ \Delta x = \dfrac{2-(-2)}{n} = \dfrac{4}{n}.$

$\boldsymbol{n = 2}$**:**

The maximum values of f on both subintervals occur at $x = 0$, so

upper sum $= f(0) \cdot 2 + f(0) \cdot 2 = 6 \cdot 2 + 6 \cdot 2 = 24.$

The minimum values of f on the subintervals occur at $x = -2$ and $x = 2$, so

lower sum $= f(-2) \cdot 2 + f(2) \cdot 2 = 2 \cdot 2 + 2 \cdot 2 = 8.$

$\boldsymbol{n = 4}$**:**

$$\begin{aligned}\text{upper sum} &= [f(-1) + f(0) + f(0) + f(1)](1)\\ &= [5 + 6 + 6 + 5](1)\\ &= 22\end{aligned}$$

$$\begin{aligned}\text{lower sum} &= [f(-2) + f(-1) + f(1) + f(2)](1)\\ &= [2 + 5 + 5 + 2](1)\\ &= 14\end{aligned}$$

$\boldsymbol{n = 8}$**:**

$$\begin{aligned}\text{upper sum} &= [f(-1.5) + f(-1) + f(-0.5) + f(0)\\ &\qquad + f(0) + f(0.5) + f(1) + f(1.5)](0.5)\\ &= 20.5\end{aligned}$$

$$\begin{aligned}\text{lower sum} &= [f(-2) + f(-1.5) + f(-1) + f(-0.5)\\ &\qquad + f(0.5) + f(1) + f(1.5) + f(2)](0.5)\\ &= 16.5\end{aligned}$$

8. $f(x) = 1 + \cos(x/2),\ -\pi \le x \le \pi,\ \Delta x = \dfrac{2\pi}{n}.$

$\boldsymbol{n = 3}$**:**

$$\begin{aligned}\text{upper sum} &= \left[f\left(-\tfrac{\pi}{3}\right) + f(0) + f\left(\tfrac{\pi}{3}\right)\right]\left(\tfrac{2\pi}{3}\right)\\ &\approx 12.005\end{aligned}$$

$$\begin{aligned}\text{lower sum} &= \left[f(-\pi) + f\left(\pm\tfrac{\pi}{3}\right) + f(\pi)\right]\left(\tfrac{2\pi}{3}\right)\\ &\approx 8.097\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\boldsymbol{n = 4}$**:**

$$\text{upper sum} = \left[f\left(-\tfrac{\pi}{2}\right) + f(0) + f(0) + f\left(\tfrac{\pi}{2}\right)\right]\left(\tfrac{\pi}{2}\right) \approx 11.646$$

$$\text{lower sum} = \left[f(-\pi) + f\left(-\tfrac{\pi}{2}\right) + f\left(\tfrac{\pi}{2}\right) + f(\pi)\right]\left(\tfrac{\pi}{2}\right) \approx 8.505$$

$\boldsymbol{n = 6}$**:**

$$\text{upper sum} = \left[f\left(-\tfrac{2\pi}{3}\right) + f\left(-\tfrac{\pi}{3}\right) + f(0) + f(0) + f\left(\tfrac{\pi}{3}\right) + f\left(\tfrac{2\pi}{3}\right)\right]\left(\tfrac{\pi}{3}\right) \approx 11.239$$

$$\text{lower sum} = \left[f(-\pi) + f\left(-\tfrac{2\pi}{3}\right) + f\left(-\tfrac{\pi}{3}\right) + f\left(\tfrac{\pi}{3}\right) + f\left(\tfrac{2\pi}{3}\right) + f(\pi)\right]\left(\tfrac{\pi}{3}\right) \approx 9.144$$

9. Since v is an increasing function, L_6 will give us a lower estimate and R_6 will give us an upper estimate.

$$L_6 = (0 \text{ ft/s})(0.5 \text{ s}) + (6.2)(0.5) + (10.8)(0.5) + (14.9)(0.5) + (18.1)(0.5) + (19.4)(0.5) = 0.5(69.4) = 34.7 \text{ ft}$$

$$R_6 = 0.5(6.2 + 10.8 + 14.9 + 18.1 + 19.4 + 20.2) = 0.5(89.6) = 44.8 \text{ ft}$$

10. (a) The velocities are given with units mi/h, so we must convert the 10-second intervals to hours:

$$10 \text{ seconds} = \frac{10 \text{ seconds}}{3600 \text{ seconds/h}} = \frac{1}{360} \text{ h}$$

$$\begin{aligned}\text{distance} \approx L_6 &= (182.9 \text{ mi/h})\left(\tfrac{1}{360} \text{ h}\right) + (168.0)\left(\tfrac{1}{360}\right) + (106.6)\left(\tfrac{1}{360}\right) + (99.8)\left(\tfrac{1}{360}\right) \\ &\quad + (124.5)\left(\tfrac{1}{360}\right) + (176.1)\left(\tfrac{1}{360}\right) \\ &= \frac{857.9}{360} \approx 2.383 \text{ miles}\end{aligned}$$

(b) $\text{Distance} \approx R_6 = \left(\tfrac{1}{360}\right)(168.0 + 106.6 + 99.8 + 124.5 + 176.1 + 175.6) = \dfrac{850.6}{360} \approx 2.363 \text{ miles}$

(c) The velocity is neither increasing nor decreasing on the given interval, so the estimates in parts (a) and (b) are neither upper nor lower estimates.

11. Lower estimate for oil leakage: $R_5 = (7.6 + 6.8 + 6.2 + 5.7 + 5.3)(2) = (31.6)(2) = 63.2$ L.

Upper estimate for oil leakage: $L_5 = (8.7 + 7.6 + 6.8 + 6.2 + 5.7)(2) = (35)(2) = 70$ L.

12. We can find an upper estimate by using the final velocity for each time interval. Thus, the distance d traveled after 62 seconds can be approximated by

$$d = \sum_{i=1}^{6} v(t_i)\,\Delta t_i = (185 \text{ ft/s})(10 \text{ s}) + 319 \cdot 5 + 447 \cdot 5 + 742 \cdot 12 + 1325 \cdot 27 + 1445 \cdot 3 = 54{,}694 \text{ ft}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. For a decreasing function, using left endpoints gives us an overestimate and using right endpoints results in an underestimate. We will use M_6 to get an estimate. $\Delta t = 1$, so

$$M_6 = 1[v(0.5) + v(1.5) + v(2.5) + v(3.5) + v(4.5) + v(5.5)] \approx 55 + 40 + 28 + 18 + 10 + 4 = 155 \text{ ft}$$

For a very rough check on the above calculation, we can draw a line from $(0, 70)$ to $(6, 0)$ and calculate the area of the triangle: $\frac{1}{2}(70)(6) = 210$. This is clearly an overestimate, so our midpoint estimate of 155 is reasonable.

14. For an increasing function, using left endpoints gives us an underestimate and using right endpoints results in an overestimate. We will use M_6 to get an estimate. $\Delta t = \frac{30-0}{6} = 5 \text{ s} = \frac{5}{3600} \text{ h} = \frac{1}{720} \text{ h}$.

$$\begin{aligned} M_6 &= \tfrac{1}{720}[v(2.5) + v(7.5) + v(12.5) + v(17.5) + v(22.5) + v(27.5)] \\ &= \tfrac{1}{720}(31.25 + 66 + 88 + 103.5 + 113.75 + 119.25) = \tfrac{1}{720}(521.75) \approx 0.725 \text{ km} \end{aligned}$$

For a very rough check on the above calculation, we can draw a line from $(0, 0)$ to $(30, 120)$ and calculate the area of the triangle: $\frac{1}{2}(30)(120) = 1800$. Divide by 3600 to get 0.5, which is clearly an underestimate, making our midpoint estimate of 0.725 seem reasonable. Of course, answers will vary due to different readings of the graph.

15. $f(t) = -t(t-21)(t+1)$ and $\Delta t = \frac{12-0}{6} = 2$

$$\begin{aligned} M_6 &= 2 \cdot f(1) + 2 \cdot f(3) + 2 \cdot f(5) + 2 \cdot f(7) + 2 \cdot f(9) + 2 \cdot f(11) \\ &= 2 \cdot 40 + 2 \cdot 216 + 2 \cdot 480 + 2 \cdot 784 + 2 \cdot 1080 + 2 \cdot 1320 \\ &= 7840 \text{ (infected cells/mL)} \cdot \text{days} \end{aligned}$$

Thus, the total amount of infection needed to develop symptoms of measles is about 7840 infected cells per mL of blood plasma.

16. $f(x) = x^2 e^x,\ 0 \le x \le 4.\quad \Delta x = (4-0)/n = 4/n$ and $x_i = 0 + i\,\Delta x = 4i/n$.

$$A = \lim_{n\to\infty} R_n = \lim_{n\to\infty} \sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty} \sum_{i=1}^{n} (4i/n)^2 e^{4i/n} \cdot \frac{4}{n}.$$

17. $f(x) = 2 + \sin^2 x,\ 0 \le x \le \pi.\quad \Delta x = (\pi - 0)/n = \pi/n$ and $x_i = 0 + i\,\Delta x = \pi i/n$.

$$A = \lim_{n\to\infty} R_n = \lim_{n\to\infty} \sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty} \sum_{i=1}^{n} \left[2 + \sin^2(\pi i/n)\right] \cdot \frac{\pi}{n}.$$

18. $f(x) = x + \ln x,\ 3 \le x \le 8.\quad \Delta x = (8-3)/n = 5/n$ and $x_i = 3 + i\,\Delta x = 3 + 5i/n$.

$$A = \lim_{n\to\infty} R_n = \lim_{n\to\infty} \sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty} \sum_{i=1}^{n} [(3 + 5i/n) + \ln(3 + 5i/n)] \cdot \frac{5}{n}.$$

19. $f(x) = x\sqrt{x^3 + 8},\ 1 \le x \le 5.\quad \Delta x = (5-1)/n = 4/n$ and $x_i = 1 + i\,\Delta x = 1 + 4i/n$.

$$A = \lim_{n\to\infty} R_n = \lim_{n\to\infty} \sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty} \sum_{i=1}^{n} \left[(1 + 4i/n)\sqrt{(1 + 4i/n)^3 + 8}\right] \cdot \frac{4}{n}.$$

20. $\displaystyle\lim_{n\to\infty} \sum_{i=1}^{n} \frac{1}{n}\left(\frac{i}{n}\right)^3$ can be interpreted as the area of the region lying under the graph of $y = x^3$ on the interval $[0, 1]$, since for $y = x^3$ on $[0, 1]$ with $\Delta x = \dfrac{1-0}{n} = \dfrac{1}{n}$, $x_i = 0 + i\,\Delta x = \dfrac{1i}{n} = \dfrac{i}{n}$, and $x_i^* = x_i$, the expression for area is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$A = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} f(x_i^*)\,\Delta x = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \left(\dfrac{i}{n}\right)^3 \cdot \dfrac{1}{n}$. Note that this answer is not unique. We could use $y = (x-1)^3$ on $[1, 2]$ or, in general, $y = (x-n)^3$ on $[n, n+1]$, where n is any real number.

21. $\lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \dfrac{2}{n} \dfrac{1}{1+(2i/n)}$ can be interpreted as the area of the region lying under the graph of $y = \dfrac{1}{1+x}$ on the interval $[0, 2]$, since for $y = \dfrac{1}{1+x}$ on $[0, 2]$ with $\Delta x = \dfrac{2-0}{n} = \dfrac{2}{n}$, $x_i = 0 + i\,\Delta x = \dfrac{2i}{n}$, and $x_i^* = x_i$, the expression for area is $A = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} f(x_i^*)\,\Delta x = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \left(\dfrac{1}{1+(2i/n)}\right) \cdot \dfrac{2}{n}$. Note that this answer is not unique. We could use $y = \dfrac{1}{x}$ on $[1, 3]$ or, in general, $y = \dfrac{1}{x-n}$ on $[n+1, n+3]$, where n is any real number. The given answer results from the general case with $n = -1$.

22. $\lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \dfrac{3}{n} \sqrt{1 + \dfrac{3i}{n}}$ can be interpreted as the area of the region lying under the graph of $y = \sqrt{1+x}$ on the interval $[0, 3]$, since for $y = \sqrt{1+x}$ on $[0, 3]$ with $\Delta x = \dfrac{3-0}{n} = \dfrac{3}{n}$, $x_i = 0 + i\,\Delta x = \dfrac{3i}{n}$, and $x_i^* = x_i$, the expression for the area is $A = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} f(x_i^*)\,\Delta x = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \sqrt{1 + \dfrac{3i}{n}}\,\dfrac{3}{n}$. Note that this answer is not unique. We could use $y = \sqrt{x}$ on $[1, 4]$ or, in general, $y = \sqrt{x-n}$ on $[n+1, n+4]$, where n is any real number.

23. $\lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \dfrac{\pi}{4n} \tan \dfrac{i\pi}{4n}$ can be interpreted as the area of the region lying under the graph of $y = \tan x$ on the interval $\left[0, \frac{\pi}{4}\right]$, since for $y = \tan x$ on $\left[0, \frac{\pi}{4}\right]$ with $\Delta x = \dfrac{\pi/4 - 0}{n} = \dfrac{\pi}{4n}$, $x_i = 0 + i\,\Delta x = \dfrac{i\pi}{4n}$, and $x_i^* = x_i$, the expression for the area is $A = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} f(x_i^*)\,\Delta x = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \tan\left(\dfrac{i\pi}{4n}\right) \dfrac{\pi}{4n}$. Note that this answer is not unique, since the expression for the area is the same for the function $y = \tan(x - k\pi)$ on the interval $\left[k\pi, k\pi + \frac{\pi}{4}\right]$, where k is any integer.

24. (a) $\Delta x = \dfrac{1-0}{n} = \dfrac{1}{n}$ and $x_i = 0 + i\,\Delta x = \dfrac{i}{n}$. $\quad A = \lim\limits_{n\to\infty} R_n = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} f(x_i)\,\Delta x = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \left(\dfrac{i}{n}\right)^3 \cdot \dfrac{1}{n}$.

(b) $\lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \dfrac{i^3}{n^3} \cdot \dfrac{1}{n} = \lim\limits_{n\to\infty} \dfrac{1}{n^4} \sum\limits_{i=1}^{n} i^3 = \lim\limits_{n\to\infty} \dfrac{1}{n^4} \left[\dfrac{n(n+1)}{2}\right]^2 = \lim\limits_{n\to\infty} \dfrac{(n+1)^2}{4n^2} = \dfrac{1}{4} \lim\limits_{n\to\infty} \left(1 + \dfrac{1}{n}\right)^2 = \dfrac{1}{4}$

25. (a) Since f is an increasing function, L_n is an underestimate of A [lower sum] and R_n is an overestimate of A [upper sum]. Thus, A, L_n, and R_n are related by the inequality $L_n < A < R_n$.

(b)
$$\begin{aligned} R_n &= f(x_1)\Delta x + f(x_2)\Delta x + \cdots + f(x_n)\Delta x \\ L_n &= f(x_0)\Delta x + f(x_1)\Delta x + \cdots + f(x_{n-1})\Delta x \\ R_n - L_n &= f(x_n)\Delta x - f(x_0)\Delta x \\ &= \Delta x[f(x_n) - f(x_0)] \\ &= \frac{b-a}{n}[f(b) - f(a)] \end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

In the diagram, $R_n - L_n$ is the sum of the areas of the shaded rectangles. By sliding the shaded rectangles to the left so that they stack on top of the leftmost shaded rectangle, we form a rectangle of height $f(b) - f(a)$ and width $\dfrac{b-a}{n}$.

(c) $A > L_n$, so $R_n - A < R_n - L_n$; that is, $R_n - A < \dfrac{b-a}{n}[f(b) - f(a)]$.

26. From Exercise 25, we have $R_n - A < \dfrac{b-a}{n}[f(b) - f(a)] = \dfrac{3-1}{n}[f(3) - f(1)] = \dfrac{2}{n}(e^3 - e)$.

Solving $\dfrac{2}{n}(e^3 - e) < 0.0001$ for n gives us $2(e^3 - e) < 0.0001n \quad \Rightarrow \quad n > \dfrac{2(e^3 - e)}{0.0001} \quad \Rightarrow \quad n > 347{,}345.1$.

Thus, a value of n that assures us that $R_n - A < 0.0001$ is $n = 347{,}346$. [This is not the *least* value of n.]

27. Here is one possible algorithm (ordered sequence of operations) for calculating the sums:

1 Let SUM = 0, X_MIN = 0, X_MAX = 1, N = 10 (depending on which sum we are calculating), DELTA_X = (X_MAX - X_MIN)/N, and RIGHT_ENDPOINT = X_MIN + DELTA_X.

2 Repeat steps 2a, 2b in sequence until RIGHT_ENDPOINT > X_MAX.

2a Add (RIGHT_ENDPOINT)^4 to SUM.

2b Add DELTA_X to RIGHT_ENDPOINT.

At the end of this procedure, (DELTA_X)·(SUM) is equal to the answer we are looking for. We find that $R_{10} = \dfrac{1}{10}\sum_{i=1}^{10}\left(\dfrac{i}{10}\right)^4 \approx 0.2533$, $R_{30} = \dfrac{1}{30}\sum_{i=1}^{30}\left(\dfrac{i}{30}\right)^4 \approx 0.2170$, $R_{50} = \dfrac{1}{50}\sum_{i=1}^{50}\left(\dfrac{i}{50}\right)^4 \approx 0.2101$, and $R_{100} = \dfrac{1}{100}\sum_{i=1}^{100}\left(\dfrac{i}{100}\right)^4 \approx 0.2050$. It appears that the exact area is 0.2. The following display shows the program SUMRIGHT and its output from a TI-83/4 Plus calculator. To generalize the program, we have input (rather than assign) values for Xmin, Xmax, and N. Also, the function, x^4, is assigned to Y_1, enabling us to evaluate any right sum merely by changing Y_1 and running the program.

```
PROGRAM:SUMRIGHT
:0→S
:Prompt Xmin
:Prompt Xmax
:Prompt N
:(Xmax-Xmin)/N→D
:Xmin+D→R
:For(I,1,N)
:S+Y1(R)→S
:R+D→R
:End
:D*S→Z
:Disp Z
```

```
prgmSUMRIGHT
Xmin=?0
Xmax=?1
N=?10
              .25333
                Done
```

28. We can use the algorithm from Exercise 27 with X_MIN = 0, X_MAX = $\pi/2$, and cos(RIGHT_ENDPOINT) instead of (RIGHT_ENDPOINT)^4 in step 2a. We find that $R_{10} = \dfrac{\pi/2}{10}\sum_{i=1}^{10}\cos\left(\dfrac{i\pi}{20}\right) \approx 0.9194$, $R_{30} = \dfrac{\pi/2}{30}\sum_{i=1}^{30}\cos\left(\dfrac{i\pi}{60}\right) \approx 0.9736$, and $R_{50} = \dfrac{\pi/2}{50}\sum_{i=1}^{50}\cos\left(\dfrac{i\pi}{100}\right) \approx 0.9842$, and $R_{100} = \dfrac{\pi/2}{100}\sum_{i=1}^{100}\cos\left(\dfrac{i\pi}{200}\right) \approx 0.9921$. It appears that the exact area is 1.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. In Maple, we have to perform a number of steps before getting a numerical answer. After loading the student package [command: `with(student);`] we use the command `left_sum:=leftsum(1/(x^2+1),x=0..1,10` [or `30`, or `50`]`);` which gives us the expression in summation notation. To get a numerical approximation to the sum, we use `evalf(left_sum);`. Mathematica does not have a special command for these sums, so we must type them in manually. For example, the first left sum is given by `(1/10)*Sum[1/(((i-1)/10)^2+1)],{i,1,10}]`, and we use the `N` command on the resulting output to get a numerical approximation.

(a) With $f(x) = \dfrac{1}{x^2+1}$, $0 \le x \le 1$, the left sums are of the form $L_n = \dfrac{1}{n}\sum\limits_{i=1}^{n} \dfrac{1}{\left(\frac{i-1}{n}\right)^2+1}$. Specifically, $L_{10} \approx 0.8100$, $L_{30} \approx 0.7937$, and $L_{50} \approx 0.7904$. The right sums are of the form $R_n = \dfrac{1}{n}\sum\limits_{i=1}^{n} \dfrac{1}{\left(\frac{i}{n}\right)^2+1}$. Specifically, $R_{10} \approx 0.7600$, $R_{30} \approx 0.7770$, and $R_{50} \approx 0.7804$.

(b) In Maple, we use the `leftbox` (with the same arguments as `left_sum`) and `rightbox` commands to generate the graphs.

left endpoints, $n = 10$ left endpoints, $n = 30$ left endpoints, $n = 50$

right endpoints, $n = 10$ right endpoints, $n = 30$ right endpoints, $n = 50$

(c) We know that since $y = 1/(x^2+1)$ is a decreasing function on $(0, 1)$, all of the left sums are larger than the actual area, and all of the right sums are smaller than the actual area. Since the left sum with $n = 50$ is about $0.7904 < 0.791$ and the right sum with $n = 50$ is about $0.7804 > 0.780$, we conclude that $0.780 < R_{50} <$ exact area $< L_{50} < 0.791$, so the exact area is between 0.780 and 0.791.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. See the solution to Exercise 29 for the CAS commands for evaluating the sums.

(a) With $f(x) = \ln x$, $1 \le x \le 4$, the left sums are of the form $L_n = \dfrac{3}{n}\sum\limits_{i=1}^{n} \ln\left(1 + \dfrac{3(i-1)}{n}\right)$. In particular, $L_{10} \approx 2.3316$, $L_{30} \approx 2.4752$, and $L_{50} \approx 2.5034$. The right sums are of the form $R_n = \dfrac{3}{n}\sum\limits_{i=1}^{n} \ln\left(1 + \dfrac{3i}{n}\right)$. In particular, $R_{10} \approx 2.7475$, $R_{30} \approx 2.6139$, and $R_{50} \approx 2.5865$.

(b) In Maple, we use the `leftbox` (with the same arguments as `left_sum`) and `rightbox` commands to generate the graphs.

left endpoints, $n = 10$ left endpoints, $n = 30$ left endpoints, $n = 50$

right endpoints, $n = 10$ right endpoints, $n = 30$ right endpoints, $n = 50$

(c) We know that since $y = \ln x$ is an increasing function on $(1, 4)$, all of the left sums are smaller than the actual area, and all of the right sums are larger than the actual area. Since the left sum with $n = 50$ is about $2.503 > 2.50$ and the right sum with $n = 50$ is about $2.587 < 2.59$, we conclude that $2.50 < L_{50} <$ exact area $< R_{50} < 2.59$, so the exact area is between 2.50 and 2.59.

31. (a) $y = f(x) = x^5$. $\Delta x = \dfrac{2-0}{n} = \dfrac{2}{n}$ and $x_i = 0 + i\,\Delta x = \dfrac{2i}{n}$.

$$A = \lim_{n\to\infty} R_n = \lim_{n\to\infty}\sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty}\sum_{i=1}^{n}\left(\frac{2i}{n}\right)^5 \cdot \frac{2}{n} = \lim_{n\to\infty}\sum_{i=1}^{n}\frac{32i^5}{n^5}\cdot\frac{2}{n} = \lim_{n\to\infty}\frac{64}{n^6}\sum_{i=1}^{n} i^5.$$

(b) $\sum\limits_{i=1}^{n} i^5 \overset{\text{CAS}}{=} \dfrac{n^2(n+1)^2\left(2n^2+2n-1\right)}{12}$

(c) $$\lim_{n\to\infty}\frac{64}{n^6}\cdot\frac{n^2(n+1)^2\left(2n^2+2n-1\right)}{12} = \frac{64}{12}\lim_{n\to\infty}\frac{\left(n^2+2n+1\right)\left(2n^2+2n-1\right)}{n^2\cdot n^2}$$
$$= \frac{16}{3}\lim_{n\to\infty}\left(1+\frac{2}{n}+\frac{1}{n^2}\right)\left(2+\frac{2}{n}-\frac{1}{n^2}\right) = \tfrac{16}{3}\cdot 1\cdot 2 = \tfrac{32}{3}$$

32. From Example 3(a), we have $A = \lim\limits_{n\to\infty}\dfrac{2}{n}\sum\limits_{i=1}^{n} e^{-2i/n}$. Using a CAS, $\sum\limits_{i=1}^{n} e^{-2i/n} = \dfrac{e^{-2}\left(e^2-1\right)}{e^{2/n}-1}$ and

$\lim\limits_{n\to\infty}\dfrac{2}{n}\cdot\dfrac{e^{-2}\left(e^2-1\right)}{e^{2/n}-1} = e^{-2}\left(e^2-1\right) \approx 0.8647$, whereas the estimate from Example 3(b) using M_{10} was 0.8632.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. $y = f(x) = \cos x$. $\Delta x = \dfrac{b-0}{n} = \dfrac{b}{n}$ and $x_i = 0 + i\,\Delta x = \dfrac{bi}{n}$.

$$A = \lim_{n\to\infty} R_n = \lim_{n\to\infty} \sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty} \sum_{i=1}^{n} \cos\left(\frac{bi}{n}\right)\cdot\frac{b}{n}$$

$$\overset{\text{CAS}}{=} \lim_{n\to\infty}\left[\frac{b\sin\left(b\left(\frac{1}{2n}+1\right)\right)}{2n\sin\left(\frac{b}{2n}\right)} - \frac{b}{2n}\right] \overset{\text{CAS}}{=} \sin b$$

If $b = \frac{\pi}{2}$, then $A = \sin\frac{\pi}{2} = 1$.

34. (a)

The diagram shows one of the n congruent triangles, $\triangle AOB$, with central angle $2\pi/n$. O is the center of the circle and AB is one of the sides of the polygon. Radius OC is drawn so as to bisect $\angle AOB$. It follows that OC intersects AB at right angles and bisects AB. Thus, $\triangle AOB$ is divided into two right triangles with legs of length $\frac{1}{2}(AB) = r\sin(\pi/n)$ and $r\cos(\pi/n)$. $\triangle AOB$ has area $2\cdot\frac{1}{2}[r\sin(\pi/n)][r\cos(\pi/n)] = r^2\sin(\pi/n)\,\cos(\pi/n) = \frac{1}{2}r^2\sin(2\pi/n)$, so $A_n = n\cdot\text{area}(\triangle AOB) = \frac{1}{2}nr^2\sin(2\pi/n)$.

(b) To use Equation 3.3.5, $\lim\limits_{\theta\to 0}\dfrac{\sin\theta}{\theta} = 1$, we need to have the same expression in the denominator as we have in the argument of the sine function—in this case, $2\pi/n$.

$$\lim_{n\to\infty} A_n = \lim_{n\to\infty}\tfrac{1}{2}nr^2\sin(2\pi/n) = \lim_{n\to\infty}\tfrac{1}{2}nr^2\frac{\sin(2\pi/n)}{2\pi/n}\cdot\frac{2\pi}{n} = \lim_{n\to\infty}\frac{\sin(2\pi/n)}{2\pi/n}\pi r^2. \text{ Let } \theta = \frac{2\pi}{n}.$$

Then as $n\to\infty$, $\theta\to 0$, so $\lim\limits_{n\to\infty} A_n = \lim\limits_{n\to\infty}\dfrac{\sin(2\pi/n)}{2\pi/n}\pi r^2 = \lim\limits_{\theta\to 0}\dfrac{\sin\theta}{\theta}\pi r^2 = (1)\,\pi r^2 = \pi r^2$.

5.2 The Definite Integral

1. $f(x) = x - 1$, $-6 \le x \le 4$. $\Delta x = \dfrac{b-a}{n} = \dfrac{4-(-6)}{5} = 2$.

Since we are using right endpoints, $x_i^* = x_i$.

$$\begin{aligned} R_5 &= \sum_{i=1}^{5} f(x_i)\,\Delta x \\ &= (\Delta x)[f(x_1)+f(x_2)+f(x_3)+f(x_4)+f(x_5)+f(x_6)] \\ &= 2[f(-4)+f(-2)+f(0)+f(2)+f(4)] \\ &= 2[-5+(-3)+(-1)+1+3] \\ &= 2(-5) = -10 \end{aligned}$$

The Riemann sum represents the sum of the areas of the two rectangles above the x-axis minus the sum of the areas of the three rectangles below the x-axis; that is, the *net area* of the rectangles with respect to the x-axis.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. $f(x) = \cos x$, $0 \le x \le \frac{3\pi}{4}$. $\Delta x = \dfrac{b-a}{n} = \dfrac{3\pi/4 - 0}{6} = \dfrac{\pi}{8}$.

Since we are using left endpoints, $x_i^* = x_{i-1}$.

$$\begin{aligned} L_6 &= \sum_{i=1}^{6} f(x_{i-1})\,\Delta x \\ &= (\Delta x)[f(x_0) + f(x_1) + f(x_2) + f(x_3) + f(x_4) + f(x_5)] \\ &= \tfrac{\pi}{8}[f(0) + f\left(\tfrac{\pi}{8}\right) + f\left(\tfrac{2\pi}{8}\right) + f\left(\tfrac{3\pi}{8}\right) + f\left(\tfrac{4\pi}{8}\right) + f\left(\tfrac{5\pi}{8}\right)] \\ &\approx 1.033186 \end{aligned}$$

The Riemann sum represents the sum of the areas of the four rectangles above the x-axis minus the area of the rectangle below the x-axis; that is, the *net area* of the rectangles with respect to the x-axis. A sixth rectangle is degenerate, with height 0, and has no area.

3. $f(x) = x^2 - 4$, $0 \le x \le 3$. $\Delta x = \dfrac{b-a}{n} = \dfrac{3-0}{6} = \dfrac{1}{2}$.

Since we are using midpoints, $x_i^* = \overline{x}_i = \frac{1}{2}(x_{i-1} + x_i)$.

$$\begin{aligned} M_6 &= \sum_{i=1}^{6} f(\overline{x}_i)\,\Delta x \\ &= (\Delta x)[f(\overline{x}_1) + f(\overline{x}_2) + f(\overline{x}_3) + f(\overline{x}_4) + f(\overline{x}_5) + f(\overline{x}_6)] \\ &= \tfrac{1}{2}\left[f\left(\tfrac{1}{4}\right) + f\left(\tfrac{3}{4}\right) + f\left(\tfrac{5}{4}\right) + f\left(\tfrac{7}{4}\right) + f\left(\tfrac{9}{4}\right) + f\left(\tfrac{11}{4}\right)\right] \\ &= \tfrac{1}{2}\left(-\tfrac{63}{16} - \tfrac{55}{16} - \tfrac{39}{16} - \tfrac{15}{16} + \tfrac{17}{16} + \tfrac{57}{16}\right) = \tfrac{1}{2}\left(-\tfrac{98}{16}\right) = -\tfrac{49}{16} \end{aligned}$$

The Riemann sum represents the sum of the areas of the two rectangles above the x-axis minus the sum of the areas of the four rectangles below the x-axis; that is, the *net area* of the rectangles with respect to the x-axis.

4. (a) $f(x) = \dfrac{1}{x}$, $1 \le x \le 2$. $\Delta x = \dfrac{b-a}{n} = \dfrac{2-1}{4} = \dfrac{1}{4}$.

Since we are using right endpoints, $x_i^* = x_i$.

$$\begin{aligned} R_4 &= \sum_{i=1}^{4} f(x_i)\,\Delta x \\ &= (\Delta x)[f(x_1) + f(x_2) + f(x_3) + f(x_4)] \\ &= \tfrac{1}{4}[f\left(\tfrac{5}{4}\right) + f\left(\tfrac{6}{4}\right) + f\left(\tfrac{7}{4}\right) + f\left(\tfrac{8}{4}\right)] \\ &= \tfrac{1}{4}\left[\tfrac{4}{5} + \tfrac{2}{3} + \tfrac{4}{7} + \tfrac{1}{2}\right] \\ &\approx 0.634524 \end{aligned}$$

The Riemann sum represents the sum of the areas of the four rectangles.

(b) Since we are using midpoints, $x_i^* = \overline{x}_i = \frac{1}{2}(x_{i-1} + x_i)$.

$$\begin{aligned} M_4 &= \sum_{i=1}^{4} f(\overline{x}_i)\,\Delta x \\ &= (\Delta x)[f(\overline{x}_1) + f(\overline{x}_2) + f(\overline{x}_3) + f(\overline{x}_4)] \\ &= \tfrac{1}{4}\left[f\left(\tfrac{9}{8}\right) + f\left(\tfrac{11}{8}\right) + f\left(\tfrac{13}{8}\right) + f\left(\tfrac{15}{8}\right)\right] \\ &= \tfrac{1}{4}\left(\tfrac{8}{9} + \tfrac{8}{11} + \tfrac{8}{13} + \tfrac{8}{15}\right) \approx 0.691220 \end{aligned}$$

The Riemann sum represents the sum of the areas of the four rectangles.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. (a) $\int_0^{10} f(x)\,dx \approx R_5 = [f(2)+f(4)+f(6)+f(8)+f(10)]\,\Delta x$

$= [-1+1+3+(-1)+0](2) = 2(2) = 4$

(b) $\int_0^{10} f(x)\,dx \approx L_5 = [f(0)+f(2)+f(4)+f(6)+f(8)]\,\Delta x$

$= [-1+(-1)+1+3+(-1)](2) = 1(2) = 2$

(c) $\int_0^{10} f(x)\,dx \approx M_5 = [f(1)+f(3)+f(5)+f(7)+f(9)]\,\Delta x$

$= [2+0+2+1+(-2)](2) = 3(2) = 6$

6. (a) $\int_{-2}^{4} g(x)\,dx \approx R_6 = [\,g(-1)+g(0)+g(1)+g(2)+g(3)+g(4)]\,\Delta x$

$= \left[\frac{3}{2}+0+\left(-\frac{3}{2}\right)+\frac{1}{2}+(-1)+\frac{1}{2}\right](1) = 0(1) = 0$

(b) $\int_{-2}^{4} g(x)\,dx \approx L_6 = [g(-2)+g(-1)+g(0)+g(1)+g(2)+g(3)]\,\Delta x$

$= \left[0+\frac{3}{2}+0+\left(-\frac{3}{2}\right)+\frac{1}{2}+(-1)\right](1) = -\frac{1}{2}(1) = -\frac{1}{2}$

(c) $\int_{-2}^{4} g(x)\,dx \approx M_6 = \left[g\left(-\frac{3}{2}\right)+g\left(-\frac{1}{2}\right)+g\left(\frac{1}{2}\right)+g\left(\frac{3}{2}\right)+g\left(\frac{5}{2}\right)+g\left(\frac{7}{2}\right)\right]\Delta x$

$= \left[1+1+(-1)+\left(-\frac{1}{2}\right)+0+\left(-\frac{1}{2}\right)\right](1) = 0(1) = 0$

7. Since f is increasing, $L_5 \le \int_{10}^{30} f(x)\,dx \le R_5$.

Lower estimate $= L_5 = \sum_{i=1}^{5} f(x_{i-1})\Delta x = 4[f(10)+f(14)+f(18)+f(22)+f(26)]$

$= 4[-12+(-6)+(-2)+1+3] = 4(-16) = -64$

Upper estimate $= R_5 = \sum_{i=1}^{5} f(x)\Delta x = 4[f(14)+f(18)+f(22)+f(26)+f(30)]$

$= 4[-6+(-2)+1+3+8] = 4(4) = 16$

8. (a) Using the right endpoints to approximate $\int_3^9 f(x)\,dx$, we have

$\sum_{i=1}^{3} f(x_i)\,\Delta x = 2[f(5)+f(7)+f(9)] = 2\,(-0.6+0.9+1.8) = 4.2.$

Since f is *increasing*, using *right* endpoints gives an *overestimate*.

(b) Using the left endpoints to approximate $\int_3^9 f(x)\,dx$, we have

$\sum_{i=1}^{3} f(x_{i-1})\,\Delta x = 2[f(3)+f(5)+f(7)] = 2\,(-3.4-0.6+0.9) = -6.2.$

Since f is *increasing*, using *left* endpoints gives an *underestimate*.

(c) Using the midpoint of each interval to approximate $\int_3^9 f(x)\,dx$, we have

$\sum_{i=1}^{3} f(\overline{x}_i)\,\Delta x = 2[f(4)+f(6)+f(8)] = 2\,(-2.1+0.3+1.4) = -0.8.$

We cannot say anything about the midpoint estimate compared to the exact value of the integral.

9. $\Delta x = (8-0)/4 = 2$, so the endpoints are 0, 2, 4, 6, and 8, and the midpoints are 1, 3, 5, and 7. The Midpoint Rule gives $\int_0^8 x^2\,dx \approx \sum_{i=1}^{4} f(\overline{x}_i)\,\Delta x = 2(1^2+3^2+5^2+7^2) = 2(84) = 168.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $\Delta x = (2-0)/4 = \frac{1}{2}$, so the endpoints are 0, $\frac{1}{2}$, 1, $\frac{3}{2}$, and 2, and the midpoints are $\frac{1}{4}$, $\frac{3}{4}$, $\frac{5}{4}$, and $\frac{7}{4}$. The Midpoint Rule gives

$$\int_0^2 (8x+3)\,dx \approx \sum_{i=1}^{4} f(\overline{x}_i)\,\Delta x = \left[\left(8\cdot\tfrac{1}{4}+3\right)+\left(8\cdot\tfrac{3}{4}+3\right)+\left(8\cdot\tfrac{5}{4}+3\right)+\left(8\cdot\tfrac{7}{4}+3\right)\right]\left(\tfrac{1}{2}\right) = (44)\left(\tfrac{1}{2}\right) = 22.$$

11. $\Delta x = (3-0)/6 = \frac{1}{2}$, so the endpoints are 0, $\frac{1}{2}$, 1, $\frac{3}{2}$, 2, $\frac{5}{2}$, and 3, and the midpoints are $\frac{1}{4}$, $\frac{3}{4}$, $\frac{5}{4}$, $\frac{7}{4}$, $\frac{9}{4}$, and $\frac{11}{4}$. The Midpoint Rule gives $\displaystyle\int_0^3 e^{\sqrt{x}}\,dx \approx \sum_{i=1}^{6} f(\overline{x}_i)\,\Delta x = \left(e^{\sqrt{1/4}}+e^{\sqrt{3/4}}+e^{\sqrt{5/4}}+e^{\sqrt{7/4}}+e^{\sqrt{9/4}}+e^{\sqrt{11/4}}\right)\left(\tfrac{1}{2}\right) \approx 10.2857.$

12. $\Delta x = (1-0)/5 = \frac{1}{5}$, so the endpoints are 0, $\frac{1}{5}$, $\frac{2}{5}$, $\frac{3}{5}$, $\frac{4}{5}$, and 1, and the midpoints are $\frac{1}{10}$, $\frac{3}{10}$, $\frac{5}{10}$, $\frac{7}{10}$, and $\frac{9}{10}$. The Midpoint Rule gives

$$\int_0^1 \sqrt{x^3+1}\,dx \approx \sum_{i=1}^{5} f(\overline{x}_i)\,\Delta x = \tfrac{1}{5}\left(\sqrt{\left(\tfrac{1}{10}\right)^3+1}+\sqrt{\left(\tfrac{3}{10}\right)^3+1}+\sqrt{\left(\tfrac{5}{10}\right)^3+1}+\sqrt{\left(\tfrac{7}{10}\right)^3+1}+\sqrt{\left(\tfrac{9}{10}\right)^3+1}\right)$$
$$\approx 1.1097$$

13. $\Delta x = (3-1)/5 = \frac{2}{5}$, so the endpoints are 1, $\frac{7}{5}$, $\frac{9}{5}$, $\frac{11}{5}$, $\frac{13}{5}$, and 3, and the midpoints are $\frac{6}{5}$, $\frac{8}{5}$, 2, $\frac{12}{5}$, and $\frac{14}{5}$. The Midpoint Rule gives

$$\int_1^3 \frac{x}{x^2+8}\,dx \approx \sum_{i=1}^{5} f(\overline{x}_i)\,\Delta x = \left(\frac{6/5}{(6/5)^2+8}+\frac{8/5}{(8/5)^2+8}+\frac{2}{2^2+8}+\frac{12/5}{(12/5)^2+8}+\frac{14/5}{(14/5)^2+8}\right)\left(\frac{2}{5}\right)$$
$$\approx 0.3186$$

14. $\Delta x = (\pi-0)/4 = \frac{\pi}{4}$, so the endpoints are $\frac{\pi}{4}$, $\frac{2\pi}{4}$, $\frac{3\pi}{4}$, and $\frac{4\pi}{4}$, and the midpoints are $\frac{\pi}{8}$, $\frac{3\pi}{8}$, $\frac{5\pi}{8}$, and $\frac{7\pi}{8}$. The Midpoint Rule gives

$$\int_0^\pi x\sin^2 x\,dx \approx \sum_{i=1}^{5} f(\overline{x}_i)\,\Delta x = \frac{\pi}{4}\left(\frac{\pi}{8}\sin^2\frac{\pi}{8}+\frac{3\pi}{8}\sin^2\frac{3\pi}{8}+\frac{5\pi}{8}\sin^2\frac{5\pi}{8}+\frac{7\pi}{8}\sin^2\frac{7\pi}{8}\right) \approx 2.4674$$

15. Using Mathematica and the Riemann Sum notebook from MathWorld, we obtain the following for $f(x) = x/(x^2+8)$:

$M_5 \approx 0.318595$

$M_{10} \approx 0.318144$

$M_{20} \approx 0.318032$

16. For $f(x) = x/(x+1)$ on $[0,2]$, we calculate $L_{100} \approx 0.89469$ and $R_{100} \approx 0.90802$. Since f is increasing on $[0,2]$, L_{100} is an underestimate of $\displaystyle\int_0^2 \frac{x}{x+1}\,dx$ and R_{100} is an overestimate. Thus, $0.8946 < \displaystyle\int_0^2 \frac{x}{x+1}\,dx < 0.9081$.

17. We'll create the table of values to approximate $\int_0^\pi \sin x\,dx$ by using the program in the solution to Exercise 5.1.27 with $\mathrm{Y}_1 = \sin x$, $\mathrm{Xmin} = 0$, $\mathrm{Xmax} = \pi$, and $n = 5$, 10, 50, and 100.

The values of R_n appear to be approaching 2.

n	R_n
5	1.933766
10	1.983524
50	1.999342
100	1.999836

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $\int_0^2 e^{-x^2}\,dx$ with $n = 5$, 10, 50, and 100.

n	L_n	R_n
5	1.077467	0.684794
10	0.980007	0.783670
50	0.901705	0.862438
100	0.891896	0.872262

The value of the integral lies between 0.872 and 0.892. Note that $f(x) = e^{-x^2}$ is decreasing on $(0, 2)$. We cannot make a similar statement for $\int_{-1}^2 e^{-x^2}\,dx$ since f is increasing on $(-1, 0)$.

19. On $[0, 1]$, $\displaystyle\lim_{n\to\infty} \sum_{i=1}^{n} \frac{e^{x_i}}{1 + x_i}\,\Delta x = \int_0^1 \frac{e^x}{1 + x}\,dx$.

20. On $[2, 5]$, $\displaystyle\lim_{n\to\infty} \sum_{i=1}^{n} x_i\sqrt{1 + x_i^3}\,\Delta x = \int_2^5 x\sqrt{1 + x^3}\,dx$.

21. On $[2, 7]$, $\displaystyle\lim_{n\to\infty} \sum_{i=1}^{n} [5(x_i^*)^3 - 4x_i^*]\,\Delta x = \int_2^7 (5x^3 - 4x)\,dx$.

22. On $[1, 3]$, $\displaystyle\lim_{n\to\infty} \sum_{i=1}^{n} \frac{x_i^*}{(x_i^*)^2 + 4}\,\Delta x = \int_1^3 \frac{x}{x^2 + 4}\,dx$.

23. For $\displaystyle\int_0^4 (x - x^2)\,dx$, $\Delta x = \dfrac{4 - 0}{n} = \dfrac{4}{n}$, and $x_i = 0 + i\,\Delta x = \dfrac{4i}{n}$. Then

$$\int_0^4 (x - x^2)\,dx = \lim_{n\to\infty} \sum_{i=1}^{n} f\left(\frac{4i}{n}\right)\frac{4}{n} = \lim_{n\to\infty} \sum_{i=1}^{n} \left[\left(\frac{4i}{n}\right) - \left(\frac{4i}{n}\right)^2\right]\frac{4}{n} = \lim_{n\to\infty} \frac{4}{n}\sum_{i=1}^{n}\left[\frac{4i}{n} - \frac{16i^2}{n^2}\right] = \lim_{n\to\infty} R_n.$$

$$\begin{aligned}
\lim_{n\to\infty} \frac{4}{n}\sum_{i=1}^{n}\left[\frac{4i}{n} - \frac{16i^2}{n^2}\right] &= \lim_{n\to\infty} \frac{4}{n}\left[\frac{4}{n}\sum_{i=1}^{n} i - \frac{16}{n^2}\sum_{i=1}^{n} i^2\right] = \lim_{n\to\infty}\left[\frac{16}{n^2}\frac{n(n+1)}{2} - \frac{64}{n^3}\frac{n(n+1)(2n+1)}{6}\right] \\
&= \lim_{n\to\infty}\left[\frac{8}{n}(n+1) - \frac{32}{3n^2}(n+1)(2n+1)\right] \\
&= \lim_{n\to\infty}\left[8\left(1 + \frac{1}{n}\right) - \frac{32}{3}\left(1 + \frac{1}{n}\right)\left(2 + \frac{1}{n}\right)\right] = 8(1) - \frac{32}{3}(1)(2) = -\frac{40}{3}
\end{aligned}$$

24. For $\displaystyle\int_1^3 (x^3 + 5x^2)\,dx$, $\Delta x = \dfrac{3 - 1}{n} = \dfrac{2}{n}$, and $x_i = 1 + i\,\Delta x = 1 + \dfrac{2i}{n}$. Then

$$\begin{aligned}
\int_1^3 (x^3 + 5x^2)\,dx &= \lim_{n\to\infty} \sum_{i=1}^{n} f\left(1 + \frac{2i}{n}\right)\frac{2}{n} = \lim_{n\to\infty} \sum_{i=1}^{n}\left[\left(1 + \frac{2i}{n}\right)^3 + 5\left(1 + \frac{2i}{n}\right)^2\right]\frac{2}{n} \\
&= \lim_{n\to\infty} \frac{2}{n}\sum_{i=1}^{n}\left[\left(1 + \frac{6i}{n} + \frac{12i^2}{n^2} + \frac{8i^3}{n^3}\right) + \left(5 + \frac{20i}{n} + \frac{20i^2}{n^2}\right)\right] \\
&= \lim_{n\to\infty} \frac{2}{n}\sum_{i=1}^{n}\left[6 + \frac{26i}{n} + \frac{32i^2}{n^2} + \frac{8i^3}{n^3}\right] = \lim_{n\to\infty} R_n
\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\lim_{n\to\infty}\frac{2}{n}\sum_{i=1}^{n}\left[6+\frac{26i}{n}+\frac{32i^2}{n^2}+\frac{8i^3}{n^3}\right]=\lim_{n\to\infty}\frac{2}{n}\left[\sum_{i=1}^{n}6+\frac{26}{n}\sum_{i=1}^{n}i+\frac{32}{n^2}\sum_{i=1}^{n}i^2+\frac{8}{n^3}\sum_{i=1}^{n}i^3\right]$$

$$=\lim_{n\to\infty}\left\{\frac{2}{n}\cdot n(6)+\frac{52}{n^2}\frac{n(n+1)}{2}+\frac{64}{n^3}\frac{n(n+1)(2n+1)}{6}+\frac{16}{n^4}\left[\frac{n(n+1)}{2}\right]^2\right\}$$

$$=\lim_{n\to\infty}\left[2(6)+\frac{26}{n}(n+1)+\frac{32}{3n^2}(n+1)(2n+1)+\frac{4}{n^2}(n+1)^2\right]$$

$$=\lim_{n\to\infty}\left[12+26\left(1+\frac{1}{n}\right)+\frac{32}{3}\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right)+4\left(1+\frac{1}{n}\right)^2\right]$$

$$=12+26(1)+\frac{32}{3}(1)(2)+4(1)^2=\frac{190}{3}$$

25. $f(x)=\sqrt{4+x^2}$, $a=1$, $b=3$, and $\Delta x=\dfrac{3-1}{n}=\dfrac{2}{n}$. Using Theorem 4, we get $x_i^*=x_i=1+i\,\Delta x=1+\dfrac{2i}{n}$, so

$$\int_1^3\sqrt{4+x^2}\,dx=\lim_{n\to\infty}R_n=\lim_{n\to\infty}\sum_{i=1}^{n}\sqrt{4+\left(1+\frac{2i}{n}\right)^2}\cdot\frac{2}{n}.$$

26. $f(x)=x^2+\dfrac{1}{x}$, $a=2$, $b=5$, and $\Delta x=\dfrac{5-2}{n}=\dfrac{3}{n}$. Using Theorem 4, we get $x_i^*=x_i=2+i\,\Delta x=2+\dfrac{3i}{n}$, so

$$\int_2^5\left(x^2+\frac{1}{x}\right)dx=\lim_{n\to\infty}R_n=\lim_{n\to\infty}\sum_{i=1}^{n}\left[\left(2+\frac{3i}{n}\right)^2+\frac{1}{2+\frac{3i}{n}}\right]\cdot\frac{3}{n}.$$

27. Note that $\Delta x=\dfrac{2-0}{n}=\dfrac{2}{n}$ and $x_i=0+i\,\Delta x=\dfrac{2i}{n}$.

$$\int_0^2 3x\,dx=\lim_{n\to\infty}\sum_{i=1}^{n}f(x_i)\,\Delta x=\lim_{n\to\infty}\sum_{i=1}^{n}f\left(\frac{2i}{n}\right)\frac{2}{n}=\lim_{n\to\infty}\frac{2}{n}\sum_{i=1}^{n}3\left(\frac{2i}{n}\right)=\lim_{n\to\infty}\frac{2}{n}\sum_{i=1}^{n}\frac{6i}{n}$$

$$=\lim_{n\to\infty}\frac{2}{n}\left(\frac{6}{n}\right)\sum_{i=1}^{n}i=\lim_{n\to\infty}\frac{2}{n}\left(\frac{6}{n}\right)\left[\frac{n(n+1)}{2}\right]=\lim_{n\to\infty}6\left(\frac{n+1}{n}\right)$$

$$=\lim_{n\to\infty}6\left(1+\frac{1}{n}\right)=6(1)=6$$

28. Note that $\Delta x=\dfrac{3-0}{n}=\dfrac{3}{n}$ and $x_i=0+i\,\Delta x=\dfrac{3i}{n}$.

$$\int_0^3 x^2\,dx=\lim_{n\to\infty}\sum_{i=1}^{n}f(x_i)\,\Delta x=\lim_{n\to\infty}\sum_{i=1}^{n}f\left(\frac{3i}{n}\right)\frac{3}{n}=\lim_{n\to\infty}\frac{3}{n}\sum_{i=1}^{n}\left(\frac{3i}{n}\right)^2=\lim_{n\to\infty}\frac{3}{n}\sum_{i=1}^{n}\frac{9i^2}{n^2}$$

$$=\lim_{n\to\infty}\frac{3}{n}\left(\frac{9}{n^2}\right)\sum_{i=1}^{n}i^2=\lim_{n\to\infty}\frac{3}{n}\left(\frac{9}{n^2}\right)\left[\frac{n(n+1)(2n+1)}{6}\right]$$

$$=\frac{9}{2}\lim_{n\to\infty}\frac{(n+1)(2n+1)}{n^2}=\frac{9}{2}\lim_{n\to\infty}\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right)$$

$$=\frac{9}{2}(1)(2)=9$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. Note that $\Delta x = \dfrac{3-0}{n} = \dfrac{3}{n}$ and $x_i = 0 + i\,\Delta x = \dfrac{3i}{n}$.

$$\begin{aligned}
\int_0^3 (5x+2)\,dx &= \lim_{n\to\infty} \sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty} \sum_{i=1}^{n} \left[5\left(\frac{3i}{n}\right)+2\right]\frac{3}{n} = \lim_{n\to\infty} \frac{3}{n}\sum_{i=1}^{n}\left(\frac{15i}{n}+2\right) \\
&= \lim_{n\to\infty} \frac{3}{n}\left(\frac{15}{n}\sum_{i=1}^{n} i + \sum_{i=1}^{n} 2\right) = \lim_{n\to\infty}\left[\frac{45}{n^2}\frac{n(n+1)}{2} + \frac{3}{n}\cdot n(2)\right] \\
&= \lim_{n\to\infty}\left[\frac{45}{2n}(n+1)+3(2)\right] = \lim_{n\to\infty}\left[\frac{45}{2}\left(1+\frac{1}{n}\right)+6\right] = \frac{45}{2}(1)+6 = \frac{57}{2}
\end{aligned}$$

30. Note that $\Delta x = \dfrac{4-0}{n} = \dfrac{4}{n}$ and $x_i = 0 + i\,\Delta x = \dfrac{4i}{n}$.

$$\begin{aligned}
\int_0^4 (6-x^2)\,dx &= \lim_{n\to\infty} \sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty} \sum_{i=1}^{n} f\left(\frac{4i}{n}\right)\frac{4}{n} = \lim_{n\to\infty} \frac{4}{n}\sum_{i=1}^{n}\left[6-\left(\frac{4i}{n}\right)^2\right] \\
&= \lim_{n\to\infty} \frac{4}{n}\sum_{i=1}^{n}\left(6-\frac{16i^2}{n^2}\right) = \lim_{n\to\infty}\left(\frac{4}{n}\sum_{i=1}^{n} 6 - \frac{64}{n^3}\sum_{i=1}^{n} i^2\right) \\
&= \lim_{n\to\infty}\left[\frac{4}{n}\cdot n(6) - \frac{64}{n^3}\frac{n(n+1)(2n+1)}{6}\right] = \lim_{n\to\infty}\left[4(6) - \frac{32}{3n^2}(n+1)(2n+1)\right] \\
&= \lim_{n\to\infty}\left[24 - \frac{32}{3}\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right)\right] = 24 - \frac{32}{3}(1)(2) = \frac{8}{3}
\end{aligned}$$

31. Note that $\Delta x = \dfrac{5-1}{n} = \dfrac{4}{n}$ and $x_i = 1 + i\,\Delta x = 1 + \dfrac{4i}{n}$.

$$\begin{aligned}
\int_1^5 (3x^2+7x)\,dx &= \lim_{n\to\infty} \sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty} \sum_{i=1}^{n} f\left(1+\frac{4i}{n}\right)\frac{4}{n} \\
&= \lim_{n\to\infty} \frac{4}{n}\sum_{i=1}^{n}\left[3\left(1+\frac{4i}{n}\right)^2 + 7\left(1+\frac{4i}{n}\right)\right] \\
&= \lim_{n\to\infty} \frac{4}{n}\sum_{i=1}^{n}\left[3\left(1+\frac{8i}{n}+\frac{16i^2}{n^2}\right) + 7\left(1+\frac{4i}{n}\right)\right] \\
&= \lim_{n\to\infty} \frac{4}{n}\sum_{i=1}^{n}\left[3+\frac{24i}{n}+\frac{48i^2}{n^2}+7+\frac{28i}{n}\right] = \lim_{n\to\infty} \frac{4}{n}\sum_{i=1}^{n}\left[10+\frac{52i}{n}+\frac{48i^2}{n^2}\right] \\
&= \lim_{n\to\infty} \frac{4}{n}\left[\sum_{i=1}^{n} 10 + \frac{52}{n}\sum_{i=1}^{n} i + \frac{48}{n^2}\sum_{i=1}^{n} i^2\right] \\
&= \lim_{n\to\infty}\left[\frac{4}{n}\cdot n(10) + \frac{208}{n^2}\frac{n(n+1)}{2} + \frac{192}{n^3}\frac{n(n+1)(2n+1)}{6}\right] \\
&= \lim_{n\to\infty}\left[4\,(10) + \frac{104}{n}(n+1) + \frac{32}{n^2}(n+1)(2n+1)\right] \\
&= \lim_{n\to\infty}\left[40 + 104\left(1+\frac{1}{n}\right) + 32\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right)\right] \\
&= 40 + 104(1) + 32(1)(2) = 208
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. Note that $\Delta x = \dfrac{2-(-1)}{n} = \dfrac{3}{n}$ and $x_i = -1 + i\,\Delta x = -1 + \dfrac{3i}{n}$.

$$\begin{aligned}
\int_{-1}^{2}(4x^2+x+2)\,dx &= \lim_{n\to\infty}\sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty}\sum_{i=1}^{n} f\left(-1+\frac{3i}{n}\right)\frac{3}{n} \\
&= \lim_{n\to\infty}\frac{3}{n}\sum_{i=1}^{n}\left[4\left(-1+\frac{3i}{n}\right)^2+\left(-1+\frac{3i}{n}\right)+2\right] \\
&= \lim_{n\to\infty}\frac{3}{n}\sum_{i=1}^{n}\left[4\left(1-\frac{6i}{n}+\frac{9i^2}{n^2}\right)+\left(-1+\frac{3i}{n}\right)+2\right] \\
&= \lim_{n\to\infty}\frac{3}{n}\sum_{i=1}^{n}\left[5-\frac{21i}{n}+\frac{36i^2}{n^2}\right] = \lim_{n\to\infty}\frac{3}{n}\left[\sum_{i=1}^{n}5-\frac{21}{n}\sum_{i=1}^{n}i+\frac{36}{n^2}\sum_{i=1}^{n}i^2\right] \\
&= \lim_{n\to\infty}\left[\frac{3}{n}\cdot n(5)-\frac{63}{n^2}\frac{n(n+1)}{2}+\frac{108}{n^3}\frac{n(n+1)(2n+1)}{6}\right] \\
&= \lim_{n\to\infty}\left[3(5)-\frac{63}{2n}(n+1)+\frac{18}{n^2}(n+1)(2n+1)\right] \\
&= \lim_{n\to\infty}\left[15-\frac{63}{2}\left(1+\frac{1}{n}\right)+18\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right)\right] = 15-\frac{63}{2}(1)+18(1)(2) = \frac{39}{2}
\end{aligned}$$

33. Note that $\Delta x = \dfrac{1-0}{n} = \dfrac{1}{n}$ and $x_i = 0 + i\,\Delta x = \dfrac{i}{n}$.

$$\begin{aligned}
\int_0^1 (x^3-3x^2)\,dx &= \lim_{n\to\infty}\sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty}\sum_{i=1}^{n} f\left(\frac{i}{n}\right)\Delta x = \lim_{n\to\infty}\sum_{i=1}^{n}\left[\left(\frac{i}{n}\right)^3-3\left(\frac{i}{n}\right)^2\right]\frac{1}{n} \\
&= \lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^{n}\left[\frac{i^3}{n^3}-\frac{3i^2}{n^2}\right] = \lim_{n\to\infty}\frac{1}{n}\left[\frac{1}{n^3}\sum_{i=1}^{n}i^3-\frac{3}{n^2}\sum_{i=1}^{n}i^2\right] \\
&= \lim_{n\to\infty}\left\{\frac{1}{n^4}\left[\frac{n(n+1)}{2}\right]^2-\frac{3}{n^3}\,\frac{n(n+1)(2n+1)}{6}\right\} = \lim_{n\to\infty}\left[\frac{1}{4}\,\frac{n+1}{n}\,\frac{n+1}{n}-\frac{1}{2}\,\frac{n+1}{n}\,\frac{2n+1}{n}\right] \\
&= \lim_{n\to\infty}\left[\frac{1}{4}\left(1+\frac{1}{n}\right)\left(1+\frac{1}{n}\right)-\frac{1}{2}\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right)\right] = \frac{1}{4}(1)(1)-\frac{1}{2}(1)(2) = -\frac{3}{4}
\end{aligned}$$

34. Note that $\Delta x = \dfrac{2-0}{n} = \dfrac{2}{n}$ and $x_i = 0 + i\,\Delta x = \dfrac{2i}{n}$.

$$\begin{aligned}
\int_0^2 (2x-x^3)\,dx &= \lim_{n\to\infty}\sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty}\sum_{i=1}^{n} f\left(\frac{2i}{n}\right)\frac{2}{n} = \lim_{n\to\infty}\frac{2}{n}\sum_{i=1}^{n}\left[2\left(\frac{2i}{n}\right)-\left(\frac{2i}{n}\right)^3\right] \\
&= \lim_{n\to\infty}\frac{2}{n}\sum_{i=1}^{n}\left[\frac{4i}{n}-\frac{8i^3}{n^3}\right] = \lim_{n\to\infty}\frac{2}{n}\left[\frac{4}{n}\sum_{i=1}^{n}i-\frac{8}{n^3}\sum_{i=1}^{n}i^3\right] \\
&= \lim_{n\to\infty}\left\{\frac{8}{n^2}\,\frac{n(n+1)}{2}-\frac{16}{n^4}\left[\frac{n(n+1)}{2}\right]^2\right\} = \lim_{n\to\infty}\left[4\frac{n+1}{n}-4\,\frac{(n+1)^2}{n^2}\right] \\
&= \lim_{n\to\infty}\left[4\left(1+\frac{1}{n}\right)-4\,\frac{n+1}{n}\,\frac{n+1}{n}\right] = \lim_{n\to\infty}\left[4\left(1+\frac{1}{n}\right)-4\left(1+\frac{1}{n}\right)\left(1+\frac{1}{n}\right)\right] \\
&= 4(1)-4(1)(1) = 0
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. (a) Think of $\int_0^2 f(x)\,dx$ as the area of a trapezoid with bases 1 and 3 and height 2. The area of a trapezoid is $A = \frac{1}{2}(b+B)h$, so $\int_0^2 f(x)\,dx = \frac{1}{2}(1+3)2 = 4$.

(b) $\int_0^5 f(x)\,dx = \underbrace{\int_0^2 f(x)\,dx}_{\text{trapezoid}} + \underbrace{\int_2^3 f(x)\,dx}_{\text{rectangle}} + \underbrace{\int_3^5 f(x)\,dx}_{\text{triangle}}$

$$= \tfrac{1}{2}(1+3)2 + 3\cdot 1 + \tfrac{1}{2}\cdot 2\cdot 3 = 4+3+3 = 10$$

(c) $\int_5^7 f(x)\,dx$ is the negative of the area of the triangle with base 2 and height 3. $\int_5^7 f(x)\,dx = -\frac{1}{2}\cdot 2\cdot 3 = -3$.

(d) $\int_3^7 f(x)\,dx = \int_3^5 f(x)\,dx + \int_5^7 f(x)\,dx$. $\int_3^5 f(x)\,dx$ is the area of the triangle with base 2 and height 3. $\int_3^5 f(x)\,dx = \frac{1}{2}\cdot 2\cdot 3 = 3$. From part (c), $\int_5^7 f(x)\,dx = -3$. Thus, $\int_3^7 f(x)\,dx = 3 + (-3) = 0$.

Or: Since $\int_3^5 f(x)\,dx$ is the same figure as in part (c), but with opposite sign, it has value 3. Thus, $\int_3^7 f(x)\,dx = 3 + (-3) = 0$.

(e) $\int_3^7 |f(x)|\,dx = \int_3^5 |f(x)|\,dx + \int_5^7 |f(x)|\,dx = \int_3^5 f(x)\,dx + \int_5^7 [-f(x)]\,dx$. From part (d), $\int_3^5 f(x)\,dx = 3$. From part (c), $\int_5^7 f(x)\,dx = -3$, so $\int_5^7 [-f(x)]\,dx = -(-3) = 3$. Thus, $\int_3^7 |f(x)|\,dx = 3+3 = 6$.

(f) $\int_2^0 f(x)\,dx = -\int_0^2 f(x)\,dx$. From part (a), $\int_0^2 f(x)\,dx = 4$, so $\int_2^0 f(x)\,dx = -4$.

36. (a) $\int_0^2 g(x)\,dx = \frac{1}{2}\cdot 4\cdot 2 = 4$ [area of a triangle]

(b) $\int_2^6 g(x)\,dx = -\frac{1}{2}\pi(2)^2 = -2\pi$ [negative of the area of a semicircle]

(c) $\int_6^7 g(x)\,dx = \frac{1}{2}\cdot 1\cdot 1 = \frac{1}{2}$ [area of a triangle]

$$\int_0^7 g(x)\,dx = \int_0^2 g(x)\,dx + \int_2^6 g(x)\,dx + \int_6^7 g(x)\,dx = 4 - 2\pi + \tfrac{1}{2} = 4.5 - 2\pi$$

37. (a) Note that $\Delta x = \dfrac{3-0}{n} = \dfrac{3}{n}$ and $x_i = 0 + i\,\Delta x = \dfrac{3i}{n}$.

$$\int_0^3 4x\,dx = \lim_{n\to\infty}\sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty}\sum_{i=1}^{n} f\left(\frac{3i}{n}\right)\frac{3}{n} = \lim_{n\to\infty}\frac{3}{n}\sum_{i=1}^{n} 4\left(\frac{3i}{n}\right)$$

$$= \lim_{n\to\infty}\frac{3}{n}\sum_{i=1}^{n}\frac{12i}{n} = \lim_{n\to\infty}\frac{3}{n}\left(\frac{12}{n}\right)\sum_{i=1}^{n} i = \lim_{n\to\infty}\frac{36}{n^2}\left[\frac{n\,(n+1)}{2}\right]$$

$$= \lim_{n\to\infty} 18\left(\frac{n+1}{n}\right) = \lim_{n\to\infty} 18\left(1+\frac{1}{n}\right) = 18(1) = 18$$

(b) $\int_0^3 4x\,dx$ can be interpreted as the area of the shaded triangle; that is, $\frac{1}{2}(3)(12) = 18$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. (a) Note that $\Delta x = \dfrac{4-(-1)}{n} = \dfrac{5}{n}$ and $x_i = -1 + i\,\Delta x = -1 + \dfrac{5i}{n}$.

$$\int_{-1}^{4}\left(2-\frac{1}{2}x\right)dx = \lim_{n\to\infty}\sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty}\sum_{i=1}^{n} f\left(-1+\frac{5i}{n}\right)\frac{5}{n} = \lim_{n\to\infty}\frac{5}{n}\sum_{i=1}^{n}\left[2-\frac{1}{2}\left(-1+\frac{5i}{n}\right)\right]$$

$$= \lim_{n\to\infty}\frac{5}{n}\sum_{i=1}^{n}\left[2+\frac{1}{2}-\frac{5i}{2n}\right] = \lim_{n\to\infty}\frac{5}{n}\sum_{i=1}^{n}\left[\frac{5}{2}-\frac{5i}{2n}\right] = \lim_{n\to\infty}\left[\frac{5}{n}\sum_{i=1}^{n}\frac{5}{2}-\frac{25}{2n^2}\sum_{i=1}^{n} i\right]$$

$$= \lim_{n\to\infty}\left[\frac{5}{n}\cdot n\left(\frac{5}{2}\right)-\frac{25}{2n^2}\frac{n(n+1)}{2}\right] = \lim_{n\to\infty}\left[\frac{25}{2}-\frac{25}{4n}(n+1)\right]$$

$$= \lim_{n\to\infty}\left[\frac{25}{2}-\frac{25}{4}\left(1+\frac{1}{n}\right)\right] = \frac{25}{2}-\frac{25}{4}(1) = \frac{25}{4}$$

(b) $\displaystyle\int_{-1}^{4}\left(2-\frac{1}{2}x\right)dx$ can be interpreted as the area of the shaded triangle; that is, $\dfrac{1}{2}(5)\left(\dfrac{5}{2}\right) = \dfrac{25}{4}$.

39. (a) $\Delta x = (8-0)/8 = 1$ and $x_i^* = x_i = 0 + 1i = i$.

$$\int_0^8 (3-2x)\,dx$$
$$\approx \sum_{i=1}^{8} f(x_i^*)\,\Delta x$$
$$= 1\{[3-2(1)]+[3-2(2)]+\cdots+[3-2(8)]\}$$
$$= 1[1+(-1)+(-3)+(-5)+(-7)+(-9)+(-11)+(-13)]$$
$$= -48$$

(b)

(c) Note that $\Delta x = \dfrac{8-0}{n} = \dfrac{8}{n}$ and $x_i = 0 + i\,\Delta x = \dfrac{8i}{n}$.

$$\int_0^8 (3-2x)\,dx = \lim_{n\to\infty}\sum_{i=1}^{n} f(x_i)\,\Delta x = \lim_{n\to\infty}\sum_{i=1}^{n}\left[3-2\left(\frac{8i}{n}\right)\right]\left(\frac{8}{n}\right) = \lim_{n\to\infty}\frac{8}{n}\sum_{i=1}^{n}\left[3-\frac{16i}{n}\right]$$

$$= \lim_{n\to\infty}\left[\frac{8}{n}\sum_{i=1}^{n} 3-\frac{128}{n^2}\sum_{i=1}^{n} i\right] = \lim_{n\to\infty}\left[\frac{8}{n}\cdot n(3)-\frac{128}{n^2}\frac{n(n+1)}{2}\right]$$

$$= \lim_{n\to\infty}\left[8(3)-\frac{64}{n}(n+1)\right] = \lim_{n\to\infty}\left[24-64\left(1+\frac{1}{n}\right)\right]$$

$$= 24-64(1) = -40$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $\int_0^8 (3-2x)\,dx = A_1 - A_2$, where A_1 is the area marked $+$ and A_2 is the area marked $-$.

40. (a) $\Delta x = (4-0)/8 = 0.5$ and $x_i^* = x_i = 0.5i$.

$$\int_0^4 (x^2-3x)\,dx \approx \sum_{i=1}^{8} f(x_i^*)\,\Delta x$$

$$= 0.5\{[0.5^2 - 3(0.5)] + [1.0^2 - 3(1.0)] + \cdots + [3.5^2 - 3(3.5)] + [4.0^2 - 3(4.0)]\}$$

$$= \tfrac{1}{2}\left(-\tfrac{5}{4} - 2 - \tfrac{9}{4} - 2 - \tfrac{5}{4} + 0 + \tfrac{7}{4} + 4\right) = -1.5$$

(b)

(c) $$\int_0^4 (x^2-3x)\,dx = \lim_{n\to\infty} \sum_{i=1}^{n} \left[\left(\frac{4i}{n}\right)^2 - 3\left(\frac{4i}{n}\right)\right]\left(\frac{4}{n}\right)$$

$$= \lim_{n\to\infty} \frac{4}{n}\left[\frac{16}{n^2}\sum_{i=1}^{n} i^2 - \frac{12}{n}\sum_{i=1}^{n} i\right] = \lim_{n\to\infty}\left[\frac{64}{n^3}\cdot\frac{n(n+1)(2n+1)}{6} - \frac{48}{n^2}\cdot\frac{n(n+1)}{2}\right]$$

$$= \lim_{n\to\infty}\left[\frac{32}{3}\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right) - 24\left(1+\frac{1}{n}\right)\right]$$

$$= \tfrac{32}{3}\cdot 2 - 24 = -\tfrac{8}{3}$$

(d) $\int_0^4 (x^2-3x)\,dx = A_1 - A_2$, where A_1 is the area marked $+$ and A_2 is the area marked $-$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. $\int_{-2}^{5}(10-5x)\,dx$ can be interpreted as the difference of the areas of the two shaded triangles; that is, $\frac{1}{2}(4)(20)-\frac{1}{2}(3)(15)=40-\frac{45}{2}=\frac{35}{2}$.

42. $\int_{-1}^{3}(2x-1)\,dx$ can be interpreted as the difference of the areas of the two shaded triangles; that is, $-\frac{1}{2}\left(\frac{3}{2}\right)(3)+\frac{1}{2}\left(\frac{5}{2}\right)(5)=-\frac{9}{4}+\frac{25}{4}=\frac{16}{4}=4$.

43. $\int_{-4}^{3}\left|\frac{1}{2}x\right|\,dx$ can be interpreted as the sum of the areas of the two shaded triangles; that is, $\frac{1}{2}(4)(2)+\frac{1}{2}(3)\left(\frac{3}{2}\right)=4+\frac{9}{4}=\frac{25}{4}$.

44. $\int_{0}^{1}|2x-1|\,dx$ can be interpreted as the sum of the areas of the two shaded triangles; that is, $2\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)(1)=\frac{1}{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. $\int_{-3}^{0}\left(1+\sqrt{9-x^2}\right)dx$ can be interpreted as the area under the graph of $f(x)=1+\sqrt{9-x^2}$ between $x=-3$ and $x=0$. This is equal to one-quarter the area of the circle with radius 3, plus the area of the rectangle, so $\int_{-3}^{0}\left(1+\sqrt{9-x^2}\right)dx=\frac{1}{4}\pi\cdot 3^2+1\cdot 3=3+\frac{9}{4}\pi$.

46. $\int_{-4}^{4}\left(2x-\sqrt{16-x^2}\right)dx=\int_{-4}^{4}2x\,dx-\int_{-4}^{4}\sqrt{16-x^2}\,dx$. By symmetry, the value of the first integral is 0 since the shaded area above the x-axis equals the shaded area below the x-axis. The second integral can be interpreted as one-half the area of a circle with radius 4; that is, $\frac{1}{2}\pi(4)^2=8\pi$. Thus, the value of the original integral is $0-8\pi=-8\pi$.

47.
$$\int_a^b x\,dx=\lim_{n\to\infty}\frac{b-a}{n}\sum_{i=1}^{n}\left[a+\frac{b-a}{n}i\right]=\lim_{n\to\infty}\left[\frac{a(b-a)}{n}\sum_{i=1}^{n}1+\frac{(b-a)^2}{n^2}\sum_{i=1}^{n}i\right]$$
$$=\lim_{n\to\infty}\left[\frac{a(b-a)}{n}n+\frac{(b-a)^2}{n^2}\cdot\frac{n(n+1)}{2}\right]=a\,(b-a)+\lim_{n\to\infty}\frac{(b-a)^2}{2}\left(1+\frac{1}{n}\right)$$
$$=a(b-a)+\tfrac{1}{2}(b-a)^2=(b-a)\left(a+\tfrac{1}{2}b-\tfrac{1}{2}a\right)=(b-a)\tfrac{1}{2}(b+a)=\tfrac{1}{2}\left(b^2-a^2\right)$$

48.
$$\int_a^b x^2\,dx=\lim_{n\to\infty}\frac{b-a}{n}\sum_{i=1}^{n}\left[a+\frac{b-a}{n}\,i\right]^2=\lim_{n\to\infty}\frac{b-a}{n}\sum_{i=1}^{n}\left[a^2+2a\frac{b-a}{n}i+\frac{(b-a)^2}{n^2}\,i^2\right]$$
$$=\lim_{n\to\infty}\left[\frac{(b-a)^3}{n^3}\sum_{i=1}^{n}i^2+\frac{2a\,(b-a)^2}{n^2}\sum_{i=1}^{n}i+\frac{a^2\,(b-a)}{n}\sum_{i=1}^{n}1\right]$$
$$=\lim_{n\to\infty}\left[\frac{(b-a)^3}{n^3}\frac{n\,(n+1)\,(2n+1)}{6}+\frac{2a\,(b-a)^2}{n^2}\frac{n\,(n+1)}{2}+\frac{a^2\,(b-a)}{n}n\right]$$
$$=\lim_{n\to\infty}\left[\frac{(b-a)^3}{6}\cdot 1\cdot\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right)+a\,(b-a)^2\cdot 1\cdot\left(1+\frac{1}{n}\right)+a^2\,(b-a)\right]$$
$$=\frac{(b-a)^3}{3}+a\,(b-a)^2+a^2\,(b-a)=\frac{b^3-3ab^2+3a^2b-a^3}{3}+ab^2-2a^2b+a^3+a^2b-a^3$$
$$=\frac{b^3}{3}-\frac{a^3}{3}-ab^2+a^2b+ab^2-a^2b=\frac{b^3-a^3}{3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. $\Delta x = (\pi - 0)/n = \pi/n$ and $x_i^* = x_i = \pi i/n$.

$$\int_0^{\pi} \sin 5x\,dx = \lim_{n\to\infty} \sum_{i=1}^{n} (\sin 5x_i)\left(\frac{\pi}{n}\right) = \lim_{n\to\infty} \sum_{i=1}^{n} \left(\sin \frac{5\pi i}{n}\right)\frac{\pi}{n} \overset{\text{CAS}}{=} \pi \lim_{n\to\infty} \frac{1}{n} \cot\left(\frac{5\pi}{2n}\right) \overset{\text{CAS}}{=} \pi\left(\frac{2}{5\pi}\right) = \frac{2}{5}$$

50. $\Delta x = (10 - 2)/n = 8/n$ and $x_i^* = x_i = 2 + 8i/n$.

$$\begin{aligned}\int_2^{10} x^6\,dx &= \lim_{n\to\infty} \sum_{i=1}^{n} \left(2 + \frac{8i}{n}\right)^6 \left(\frac{8}{n}\right) = 8 \lim_{n\to\infty} \frac{1}{n} \sum_{i=1}^{n} \left(2 + \frac{8i}{n}\right)^6 \\ &\overset{\text{CAS}}{=} 8 \lim_{n\to\infty} \frac{1}{n} \cdot \frac{64\left(58{,}593n^6 + 164{,}052n^5 + 131{,}208n^4 - 27{,}776n^2 + 2048\right)}{21n^5} \\ &\overset{\text{CAS}}{=} 8\left(\frac{1{,}249{,}984}{7}\right) = \frac{9{,}999{,}872}{7} \approx 1{,}428{,}553.1\end{aligned}$$

51. $\int_1^1 \sqrt{1 + x^4}\,dx = 0$ since the limits of integration are equal.

52. $\int_\pi^0 \sin^4\theta\,d\theta = -\int_0^\pi \sin^4\theta\,d\theta$ [because we reversed the limits of integration]

$= -\int_0^\pi \sin^4 x\,dx$ [we can use any letter without changing the value of the integral]

$= -\frac{3}{8}\pi$ [given value]

53. $\int_0^1 (5 - 6x^2)\,dx = \int_0^1 5\,dx - 6\int_0^1 x^2\,dx = 5(1 - 0) - 6\left(\frac{1}{3}\right) = 5 - 2 = 3$

54. $\int_1^3 (2e^x - 1)\,dx = 2\int_1^3 e^x\,dx - \int_1^3 1\,dx = 2(e^3 - e) - 1(3 - 1) = 2e^3 - 2e - 2$

55. $\int_1^3 e^{x+2}\,dx = \int_1^3 e^x \cdot e^2\,dx = e^2 \int_1^3 e^x\,dx = e^2(e^3 - e) = e^5 - e^3$

56. $\int_0^{\pi/2} (2\cos x - 5x)\,dx = \int_0^{\pi/2} 2\cos x\,dx - \int_0^{\pi/2} 5x\,dx = 2\int_0^{\pi/2} \cos x\,dx - 5\int_0^{\pi/2} x\,dx$

$$= 2(1) - 5\frac{(\pi/2)^2 - 0^2}{2} = 2 - \frac{5\pi^2}{8}$$

57. $\int_{-2}^2 f(x)\,dx + \int_2^5 f(x)\,dx - \int_{-2}^{-1} f(x)\,dx = \int_{-2}^5 f(x)\,dx + \int_{-1}^{-2} f(x)\,dx$ [by Property 5 and reversing limits]

$= \int_{-1}^5 f(x)\,dx$ [Property 5]

58. $\int_2^4 f(x)\,dx + \int_4^8 f(x)\,dx = \int_2^8 f(x)\,dx$, so $\int_4^8 f(x)\,dx = \int_2^8 f(x)\,dx - \int_2^4 f(x)\,dx = 7.3 - 5.9 = 1.4$.

59. $\int_0^9 [2f(x) + 3g(x)]\,dx = 2\int_0^9 f(x)\,dx + 3\int_0^9 g(x)\,dx = 2(37) + 3(16) = 122$

60. If $f(x) = \begin{cases} 3 & \text{for } x < 3 \\ x & \text{for } x \geq 3 \end{cases}$, then $\int_0^5 f(x)\,dx$ can be interpreted as the area of the shaded region, which consists of a 5-by-3 rectangle surmounted by an isosceles right triangle whose legs have length 2. Thus, $\int_0^5 f(x)\,dx = 5(3) + \frac{1}{2}(2)(2) = 17$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. $\int_0^3 f(x)\,dx$ is clearly less than -1 and has the smallest value. The slope of the tangent line of f at $x=1$, $f'(1)$, has a value between -1 and 0, so it has the next smallest value. The largest value is $\int_3^8 f(x)\,dx$, followed by $\int_4^8 f(x)\,dx$, which has a value about 1 unit less than $\int_3^8 f(x)\,dx$. Still positive, but with a smaller value than $\int_4^8 f(x)\,dx$, is $\int_0^8 f(x)\,dx$. Ordering these quantities from smallest to largest gives us

$$\int_0^3 f(x)\,dx < f'(1) < \int_0^8 f(x)\,dx < \int_4^8 f(x)\,dx < \int_3^8 f(x)\,dx \text{ or } \mathrm{B} < \mathrm{E} < \mathrm{A} < \mathrm{D} < \mathrm{C}$$

62. $F(0) = \int_2^0 f(t)\,dt = -\int_0^2 f(t)\,dt$, so $F(0)$ is negative, and similarly, so is $F(1)$. $F(3)$ and $F(4)$ are negative since they represent negatives of areas below the x-axis. Since $F(2) = \int_2^2 f(t)\,dt = 0$ is the only non-negative value, choice C is the largest.

63. $I = \int_{-4}^2 [f(x) + 2x + 5]\,dx = \int_{-4}^2 f(x)\,dx + 2\int_{-4}^2 x\,dx + \int_{-4}^2 5\,dx = I_1 + 2I_2 + I_3$

$I_1 = -3$ [area below x-axis] $+3 - 3 = -3$

$I_2 = -\frac{1}{2}(4)(4)$ [area of triangle, see figure] $+\frac{1}{2}(2)(2)$
$= -8 + 2 = -6$

$I_3 = 5[2 - (-4)] = 5(6) = 30$

Thus, $I = -3 + 2(-6) + 30 = 15$.

64. Using Integral Comparison Property 8, $m \le f(x) \le M \quad\Rightarrow\quad m(2-0) \le \int_0^2 f(x)\,dx \le M(2-0) \quad\Rightarrow$ $2m \le \int_0^2 f(x)\,dx \le 2M$.

65. $x^2 - 4x + 4 = (x-2)^2 \ge 0$ on $[0,4]$, so $\int_0^4 (x^2 - 4x + 4)\,dx \ge 0$ [Property 6].

66. $x^2 \le x$ on $[0,1]$, so $\sqrt{1+x^2} \le \sqrt{1+x}$ on $[0,1]$. Hence, $\int_0^1 \sqrt{1+x^2}\,dx \le \int_0^1 \sqrt{1+x}\,dx$ [Property 7].

67. If $-1 \le x \le 1$, then $0 \le x^2 \le 1$ and $1 \le 1 + x^2 \le 2$, so $1 \le \sqrt{1+x^2} \le \sqrt{2}$ and $1[1-(-1)] \le \int_{-1}^1 \sqrt{1+x^2}\,dx \le \sqrt{2}\,[1-(-1)]$ [Property 8]; that is, $2 \le \int_{-1}^1 \sqrt{1+x^2}\,dx \le 2\sqrt{2}$.

68. If $\dfrac{\pi}{6} \le x \le \dfrac{\pi}{3}$, then $\dfrac{1}{2} \le \sin x \le \dfrac{\sqrt{3}}{2}$ $\left(\sin x \text{ is increasing on } \left[\frac{\pi}{6}, \frac{\pi}{3}\right]\right)$, so

$\dfrac{1}{2}\left(\dfrac{\pi}{3} - \dfrac{\pi}{6}\right) \le \displaystyle\int_{\pi/6}^{\pi/3} \sin x\,dx \le \dfrac{\sqrt{3}}{2}\left(\dfrac{\pi}{3} - \dfrac{\pi}{6}\right)$ [Property 8]; that is, $\dfrac{\pi}{12} \le \displaystyle\int_{\pi/6}^{\pi/3} \sin x\,dx \le \dfrac{\sqrt{3}\,\pi}{12}$.

69. If $0 \le x \le 1$, then $0 \le x^3 \le 1$, so $0(1-0) \le \int_0^1 x^3\,dx \le 1(1-0)$ [Property 8]; that is, $0 \le \int_0^1 x^3\,dx \le 1$.

70. If $0 \le x \le 3$, then $4 \le x + 4 \le 7$ and $\dfrac{1}{7} \le \dfrac{1}{x+4} \le \dfrac{1}{4}$, so $\dfrac{1}{7}(3-0) \le \displaystyle\int_0^3 \dfrac{1}{x+4}\,dx \le \dfrac{1}{4}(3-0)$ [Property 8]; that is,

$\dfrac{3}{7} \le \displaystyle\int_0^3 \dfrac{1}{x+4}\,dx \le \dfrac{3}{4}$.

71. If $\frac{\pi}{4} \le x \le \frac{\pi}{3}$, then $1 \le \tan x \le \sqrt{3}$, so $1\left(\frac{\pi}{3} - \frac{\pi}{4}\right) \le \int_{\pi/4}^{\pi/3} \tan x\,dx \le \sqrt{3}\left(\frac{\pi}{3} - \frac{\pi}{4}\right)$ or $\frac{\pi}{12} \le \int_{\pi/4}^{\pi/3} \tan x\,dx \le \frac{\pi}{12}\sqrt{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

72. Let $f(x) = x^3 - 3x + 3$ for $0 \le x \le 2$. Then $f'(x) = 3x^2 - 3 = 3(x+1)(x-1)$, so f is decreasing on $(0,1)$ and increasing on $(1,2)$. f has the absolute minimum value $f(1) = 1$. Since $f(0) = 3$ and $f(2) = 5$, the absolute maximum value of f is $f(2) = 5$. Thus, $1 \le x^3 - 3x + 3 \le 5$ for x in $[0,2]$. It follows from Property 8 that $1 \cdot (2-0) \le \int_0^2 \left(x^3 - 3x + 3\right) dx \le 5 \cdot (2-0)$; that is, $2 \le \int_0^2 \left(x^3 - 3x + 3\right) dx \le 10$.

73. The only critical number of $f(x) = xe^{-x}$ on $[0,2]$ is $x = 1$. Since $f(0) = 0$, $f(1) = e^{-1} \approx 0.368$, and $f(2) = 2e^{-2} \approx 0.271$, we know that the absolute minimum value of f on $[0,2]$ is 0, and the absolute maximum is e^{-1}. By Property 8, $0 \le xe^{-x} \le e^{-1}$ for $0 \le x \le 2 \quad \Rightarrow \quad 0(2-0) \le \int_0^2 xe^{-x}\,dx \le e^{-1}(2-0) \quad \Rightarrow \quad 0 \le \int_0^2 xe^{-x}\,dx \le 2/e$.

74. Let $f(x) = x - 2\sin x$ for $\pi \le x \le 2\pi$. Then $f'(x) = 1 - 2\cos x$ and $f'(x) = 0 \quad \Rightarrow \quad \cos x = \frac{1}{2} \quad \Rightarrow \quad x = \frac{5\pi}{3}$. f has the absolute maximum value $f\left(\frac{5\pi}{3}\right) = \frac{5\pi}{3} - 2\sin\frac{5\pi}{3} = \frac{5\pi}{3} + \sqrt{3} \approx 6.97$ since $f(\pi) = \pi$ and $f(2\pi) = 2\pi$ are both smaller than 6.97. Thus, $\pi \le f(x) \le \frac{5\pi}{3} + \sqrt{3} \quad \Rightarrow \quad \pi(2\pi - \pi) \le \int_\pi^{2\pi} f(x)\,dx \le \left(\frac{5\pi}{3} + \sqrt{3}\right)(2\pi - \pi)$; that is, $\pi^2 \le \int_\pi^{2\pi} (x - 2\sin x)\,dx \le \frac{5}{3}\pi^2 + \sqrt{3}\,\pi$.

75. $\sqrt{x^4+1} \ge \sqrt{x^4} = x^2$, so $\int_1^3 \sqrt{x^4+1}\,dx \ge \int_1^3 x^2\,dx = \frac{1}{3}\left(3^3 - 1^3\right) = \frac{26}{3}$.

76. $0 \le \sin x \le 1$ for $0 \le x \le \frac{\pi}{2}$, so $x \sin x \le x \quad \Rightarrow \quad \int_0^{\pi/2} x\sin x\,dx \le \int_0^{\pi/2} x\,dx = \frac{1}{2}\left[\left(\frac{\pi}{2}\right)^2 - 0^2\right] = \frac{\pi^2}{8}$.

77. $\sin x < \sqrt{x} < x$ for $1 \le x \le 2$ and $\arctan$ is an increasing function, so $\arctan(\sin x) < \arctan\sqrt{x} < \arctan x$, and hence, $\int_1^2 \arctan(\sin x)\,dx < \int_1^2 \arctan\sqrt{x}\,dx < \int_1^2 \arctan x\,dx$. Thus, $\int_1^2 \arctan x\,dx$ has the largest value.

78. $x^2 < \sqrt{x}$ for $0 < x \le 0.5$ and cosine is a decreasing function on $[0, 0.5]$, so $\cos(x^2) > \cos\sqrt{x}$, and hence, $\int_0^{0.5} \cos(x^2)\,dx > \int_0^{0.5} \cos\sqrt{x}\,dx$. Thus, $\int_0^{0.5} \cos(x^2)\,dx$ is larger.

79. Using right endpoints as in the proof of Property 2, we calculate

$$\int_a^b cf(x)\,dx = \lim_{n\to\infty} \sum_{i=1}^n cf(x_i)\,\Delta x = \lim_{n\to\infty} c\sum_{i=1}^n f(x_i)\,\Delta x = c\lim_{n\to\infty}\sum_{i=1}^n f(x_i)\,\Delta x = c\int_a^b f(x)\,dx.$$

80. (a) See the figure.

$$\begin{aligned}\left|\int_a^b f(x)\,dx\right| &= |A_1 - A_2 + A_3 - A_4 + A_5| \\ &\le |A_1 + A_2 + A_3 + A_4 + A_5| \\ &= \int_a^b |f(x)|\,dx\end{aligned}$$

(b) Since $-|f(x)| \le f(x) \le |f(x)|$, it follows from Property 7 that

$$-\int_a^b |f(x)|\,dx \le \int_a^b f(x)\,dx \le \int_a^b |f(x)|\,dx \quad \Rightarrow \quad \left|\int_a^b f(x)\,dx\right| \le \int_a^b |f(x)|\,dx$$

Note that the definite integral is a real number, and so the following property applies: $-a \le b \le a \quad \Rightarrow \quad |b| \le a$ for all real numbers b and nonnegative numbers a.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\left|\int_a^b f(x)\sin 2x\,dx\right| \le \int_a^b |f(x)\sin 2x|\ dx$ [by part (b)] $= \int_a^b |f(x)|\,|\sin 2x|\,dx \le \int_a^b |f(x)|\ dx$ by Property 7 since $|\sin 2x| \le 1 \quad \Rightarrow \quad |f(x)|\,|\sin 2x| \le |f(x)|$.

81. Suppose that f is integrable on $[0,1]$, that is, $\lim\limits_{n\to\infty}\sum\limits_{i=1}^{n} f(x_i^*)\,\Delta x$ exists for any choice of x_i^* in $[x_{i-1}, x_i]$. Let n denote a positive integer and divide the interval $[0,1]$ into n equal subintervals $\left[0,\frac{1}{n}\right], \left[\frac{1}{n},\frac{2}{n}\right], \ldots, \left[\frac{n-1}{n},1\right]$. If we choose x_i^* to be a rational number in the ith subinterval, then we obtain the Riemann sum $\sum\limits_{i=1}^{n} f(x_i^*)\cdot\frac{1}{n} = 0$, so $\lim\limits_{n\to\infty}\sum\limits_{i=1}^{n} f(x_i^*)\cdot\frac{1}{n} = \lim\limits_{n\to\infty} 0 = 0$. Now suppose we choose x_i^* to be an irrational number. Then we get $\sum\limits_{i=1}^{n} f(x_i^*)\cdot\frac{1}{n} = \sum\limits_{i=1}^{n} 1\cdot\frac{1}{n} = n\cdot\frac{1}{n} = 1$ for each n, so $\lim\limits_{n\to\infty}\sum\limits_{i=1}^{n} f(x_i^*)\cdot\frac{1}{n} = \lim\limits_{n\to\infty} 1 = 1$. Since the value of $\lim\limits_{n\to\infty}\sum\limits_{i=1}^{n} f(x_i^*)\,\Delta x$ depends on the choice of the sample points x_i^*, the limit does not exist, and f is not integrable on $[0,1]$.

82. Partition the interval $[0,1]$ into n equal subintervals and choose $x_1^* = \frac{1}{n^2}$. Then with $f(x) = \frac{1}{x}$, $\sum\limits_{i=1}^{n} f(x_i^*)\,\Delta x \ge f(x_1^*)\Delta x = \frac{1}{1/n^2}\cdot\frac{1}{n} = n$. Thus, $\sum\limits_{i=1}^{n} f(x_i^*)\,\Delta x$ can be made arbitrarily large and hence, f is not integrable on $[0,1]$.

83. $\lim\limits_{n\to\infty}\sum\limits_{i=1}^{n}\frac{i^4}{n^5} = \lim\limits_{n\to\infty}\sum\limits_{i=1}^{n}\frac{i^4}{n^4}\cdot\frac{1}{n} = \lim\limits_{n\to\infty}\sum\limits_{i=1}^{n}\left(\frac{i}{n}\right)^4\frac{1}{n}$. At this point, we need to recognize the limit as being of the form $\lim\limits_{n\to\infty}\sum\limits_{i=1}^{n} f(x_i)\,\Delta x$, where $\Delta x = (1-0)/n = 1/n$, $x_i = 0 + i\,\Delta x = i/n$, and $f(x) = x^4$. Thus, the definite integral is $\int_0^1 x^4\,dx$.

84. $\lim\limits_{n\to\infty}\frac{1}{n}\sum\limits_{i=1}^{n}\frac{1}{1+(i/n)^2} = \lim\limits_{n\to\infty}\sum\limits_{i=1}^{n}\frac{1}{1+(i/n)^2}\cdot\frac{1}{n} = \lim\limits_{n\to\infty}\sum\limits_{i=1}^{n} f(x_i)\,\Delta x$, where $\Delta x = (1-0)/n = 1/n$, $x_i = 0 + i\,\Delta x = i/n$, and $f(x) = \frac{1}{1+x^2}$. Thus, the definite integral is $\displaystyle\int_0^1 \frac{dx}{1+x^2}$.

85. Choose $x_i = 1 + \frac{i}{n}$ and $x_i^* = \sqrt{x_{i-1}x_i} = \sqrt{\left(1+\frac{i-1}{n}\right)\left(1+\frac{i}{n}\right)}$. Then

$$\begin{aligned}
\int_1^2 x^{-2}\,dx &= \lim_{n\to\infty}\frac{1}{n}\sum_{i=1}^{n}\frac{1}{\left(1+\frac{i-1}{n}\right)\left(1+\frac{i}{n}\right)} = \lim_{n\to\infty} n\sum_{i=1}^{n}\frac{1}{(n+i-1)(n+i)} \\
&= \lim_{n\to\infty} n\sum_{i=1}^{n}\left(\frac{1}{n+i-1} - \frac{1}{n+i}\right) \quad \text{[by the hint]} \quad = \lim_{n\to\infty} n\left(\sum_{i=0}^{n-1}\frac{1}{n+i} - \sum_{i=1}^{n}\frac{1}{n+i}\right) \\
&= \lim_{n\to\infty} n\left(\left[\frac{1}{n} + \frac{1}{n+1} + \cdots + \frac{1}{2n-1}\right] - \left[\frac{1}{n+1} + \cdots + \frac{1}{2n-1} + \frac{1}{2n}\right]\right) \\
&= \lim_{n\to\infty} n\left(\frac{1}{n} - \frac{1}{2n}\right) = \lim_{n\to\infty}\left(1 - \tfrac{1}{2}\right) = \tfrac{1}{2}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

DISCOVERY PROJECT Area Functions

1. (a)

Area of trapezoid $= \frac{1}{2}(b_1 + b_2)h = \frac{1}{2}(3+7)2$
$= 10$ square units

Or:

Area of rectangle + area of triangle
$= b_r h_r + \frac{1}{2} b_t h_t = (2)(3) + \frac{1}{2}(2)(4) = 10$ square units

(b)

As in part (a),

$A(x) = \frac{1}{2}[3 + (2x+1)](x-1) = \frac{1}{2}(2x+4)(x-1)$
$= (x+2)(x-1) = x^2 + x - 2$ square units

(c) $A'(x) = 2x + 1$. This is the y-coordinate of the point $(x, 2x+1)$ on the given line.

2. (a)

(b) $A(x) = \int_{-1}^{x} (1+t^2)\,dt = \int_{-1}^{x} 1\,dt + \int_{-1}^{x} t^2\,dt$ [Property 2]

$= 1[x - (-1)] + \dfrac{x^3 - (-1)^3}{3}$ [Property 1 and Exercise 5.2.48]

$= x + 1 + \frac{1}{3}x^3 + \frac{1}{3}$

$= \frac{1}{3}x^3 + x + \frac{4}{3}$

(c) $A'(x) = x^2 + 1$. This is the y-coordinate of the point $(x, 1+x^2)$ on the given curve.

(d)

$A(x+h) - A(x)$ is the area under the curve $y = 1 + t^2$ from $t = x$ to $t = x + h$.

(e)

An approximating rectangle is shown in the figure.

It has height $1 + x^2$, width h, and area $h(1+x^2)$, so

$$A(x+h) - A(x) \approx h(1+x^2) \quad \Rightarrow \quad \frac{A(x+h) - A(x)}{h} \approx 1 + x^2.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(f) Part (e) says that the average rate of change of A is approximately $1 + x^2$. As h approaches 0, the quotient approaches the instantaneous rate of change—namely, $A'(x)$. So the result of part (c), $A'(x) = x^2 + 1$, is geometrically plausible.

3. (a) $f(x) = \cos(x^2)$

(b) $g(x)$ starts to decrease at that value of x where $\cos(t^2)$ changes from positive to negative; that is, at about $x = 1.25$.

(c) $g(x) = \int_0^x \cos(t^2)\,dt$. Using an integration command, we find that

$g(0) = 0$, $g(0.2) \approx 0.200$, $g(0.4) \approx 0.399$, $g(0.6) \approx 0.592$,

$g(0.8) \approx 0.768$, $g(1.0) \approx 0.905$, $g(1.2) \approx 0.974$, $g(1.4) \approx 0.950$,

$g(1.6) \approx 0.826$, $g(1.8) \approx 0.635$, and $g(2.0) \approx 0.461$.

(d) We sketch the graph of g' using the method of Example 1 in Section 2.8.

The graphs of $g'(x)$ and $f(x)$ look alike, so we guess that $g'(x) = f(x)$.

4. In Problems 1 and 2, we showed that if $g(x) = \int_a^x f(t)\,dt$, then $g'(x) = f(x)$, for the functions $f(t) = 2t + 1$ and $f(t) = 1 + t^2$. In Problem 3 we guessed that the same is true for $f(t) = \cos(t^2)$, based on visual evidence. So we conjecture that $g'(x) = f(x)$ for any continuous function f. This turns out to be true and is proved in Section 5.3 (the Fundamental Theorem of Calculus).

5.3 The Fundamental Theorem of Calculus

1. One process undoes what the other one does. The precise version of this statement is given by the Fundamental Theorem of Calculus. See the statement of this theorem and the paragraph that follows it.

2. (a) $g(x) = \int_0^x f(t)\,dt$, so $g(0) = \int_0^0 f(t)\,dt = 0$.

$$\begin{aligned}
g(1) &= \textstyle\int_0^1 f(t)\,dt = \frac{1}{2}\cdot 1\cdot 1 \quad \text{[area of triangle]} \quad = \frac{1}{2}.\\
g(2) &= \textstyle\int_0^2 f(t)\,dt = \int_0^1 f(t)\,dt + \int_1^2 f(t)\,dt \quad \text{[below the } t\text{-axis]}\\
&= \tfrac{1}{2} - \tfrac{1}{2}\cdot 1\cdot 1 = 0.\\
g(3) &= g(2) + \textstyle\int_2^3 f(t)\,dt = 0 - \frac{1}{2}\cdot 1\cdot 1 = -\frac{1}{2}.\\
g(4) &= g(3) + \textstyle\int_3^4 f(t)\,dt = -\frac{1}{2} + \frac{1}{2}\cdot 1\cdot 1 = 0.\\
g(5) &= g(4) + \textstyle\int_4^5 f(t)\,dt = 0 + 1.5 = 1.5.\\
g(6) &= g(5) + \textstyle\int_5^6 f(t)\,dt = 1.5 + 2.5 = 4.
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $g(7) = g(6) + \int_6^7 f(t)\,dt \approx 4 + 2.2$ [estimate from the graph] $= 6.2$.

(c) The answers from part (a) and part (b) indicate that g has a minimum at $x = 3$ and a maximum at $x = 7$. This makes sense from the graph of f since we are subtracting area on $1 < x < 3$ and adding area on $3 < x < 7$.

(d)

3. (a) $g(x) = \int_0^x f(t)\,dt$.

$g(0) = \int_0^0 f(t)\,dt = 0$

$g(1) = \int_0^1 f(t)\,dt = 1 \cdot 2 = 2$ [rectangle],

$g(2) = \int_0^2 f(t)\,dt = \int_0^1 f(t)\,dt + \int_1^2 f(t)\,dt = g(1) + \int_1^2 f(t)\,dt$
$= 2 + 1 \cdot 2 + \frac{1}{2} \cdot 1 \cdot 2 = 5$ [rectangle plus triangle],

$g(3) = \int_0^3 f(t)\,dt = g(2) + \int_2^3 f(t)\,dt = 5 + \frac{1}{2} \cdot 1 \cdot 4 = 7$,

$g(6) = g(3) + \int_3^6 f(t)\,dt$ [the integral is negative since f lies under the t-axis]
$= 7 + \left[-\left(\frac{1}{2} \cdot 2 \cdot 2 + 1 \cdot 2\right)\right] = 7 - 4 = 3$

(b) g is increasing on $(0, 3)$ because as x increases from 0 to 3, we keep adding more area.

(c) g has a maximum value when we start subtracting area; that is, at $x = 3$.

(d)

4. (a) $g(x) = \int_0^x f(t)\,dt \quad \Rightarrow \quad g'(x) = f(x)$ by FTC1.

Thus, the graph of g' is the graph of f.

(b) $g(3) = \int_0^3 f(t)\,dt = \frac{1}{2} \cdot 2 \cdot 5 + \frac{1}{2}(5 + 3) \cdot 1 = 9$ [triangle plus trapezoid]

$g'(3) = f(3) = 3$ by FTC1.

From part (a), $g''(3) = f'(3)$, which is the slope of the given graph at $x = 3$.

Thus, $g''(3) = \dfrac{1 - 5}{4 - 2} = \dfrac{-4}{2} = -2$ [slope of line between $(2, 5)$ and $(4, 1)$].

(c) g has a local maximum at $x = 6$ because $f = g'$ changes from positive to negative there, so g changes from increasing to decreasing there. (We start subtracting area at $x = 6$).

(d) g has a local minimum at $x = 9$ because $f = g'$ changes from negative to positive there, so g changes from decreasing to increasing there. (We start adding area at $x = 9$.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. (a) $g(x) = \int_0^x f(t)\,dt = 3x$ [area of a rectangle with base x and height 3]

(b) $g(x) = 3x \quad \Rightarrow \quad g'(x) = 3 = f(x)$, so g is an antiderivative of f. Since $f(t) = 3$, 3 is the integrand in part (a), and 3 is the integrand evaluated at upper limit of integration x (since 3 is constant), verifying FTC1.

6. (a) $g(x) = \int_0^x f(t)\,dt = \frac{1}{2} \cdot x \cdot 3x = \frac{3}{2}x^2$ [area of a triangle with base x and height $3x$]

(b) $g(x) = \frac{3}{2}x^2 \quad \Rightarrow \quad g'(x) = \frac{3}{2} \cdot 2x = 3x = f(x)$, so g is an antiderivative of f. Since $f(t) = 3t$, $3t$ is the integrand in part (a), and $3x$ is the integrand evaluated at upper limit of integration x, verifying FTC1.

7.

(a) By FTC1 with $f(t) = t^2$ and $a = 1$, $g(x) = \int_1^x t^2\,dt \quad \Rightarrow$

$g'(x) = f(x) = x^2$.

(b) Using FTC2, $g(x) = \int_1^x t^2\,dt = \left[\frac{1}{3}t^3\right]_1^x = \frac{1}{3}x^3 - \frac{1}{3} \quad \Rightarrow \quad g'(x) = x^2$.

8.

(a) By FTC1 with $f(t) = 2 + \sin t$ and $a = 0$, $g(x) = \int_0^x (2 + \sin t)\,dt \quad \Rightarrow$

$g'(x) = f(x) = 2 + \sin x$.

(b) Using FTC2,

$$g(x) = \int_0^x (2 + \sin t)\,dt = [2t - \cos t]_0^x = (2x - \cos x) - (0 - 1)$$
$$= 2x - \cos x + 1 \quad \Rightarrow$$
$$g'(x) = 2 - (-\sin x) + 0 = 2 + \sin x$$

9. $f(t) = \sqrt{t + t^3}$ and $g(x) = \int_0^x \sqrt{t + t^3}\,dt$, so by FTC1, $g'(x) = f(x) = \sqrt{x + x^3}$.

10. $f(t) = \ln(1 + t^2)$ and $g(x) = \int_1^x \ln(1 + t^2)\,dt$, so by FTC1, $g'(x) = f(x) = \ln(1 + x^2)$.

11. $f(t) = \sin(1 + t^3)$ and $g(w) = \int_0^w \sin(1 + t^3)\,dt$, so by FTC1, $g'(w) = f(w) = \sin(1 + w^3)$.

12. $f(t) = \dfrac{\sqrt{t}}{t+1}$ and $h(u) = \displaystyle\int_0^u \frac{\sqrt{t}}{t+1}\,dt$, so by FTC1, $h'(u) = f(u) = \dfrac{\sqrt{u}}{u+1}$.

13. $F(x) = \displaystyle\int_x^0 \sqrt{1 + \sec t}\,dt = -\int_0^x \sqrt{1 + \sec t}\,dt \overset{\text{FTC1}}{\Rightarrow} F'(x) = -\frac{d}{dx}\int_0^x \sqrt{1 + \sec t}\,dt = -\sqrt{1 + \sec x}$

14. $A(w) = \int_w^{-1} e^{t+t^2}\,dt = -\int_{-1}^w e^{t+t^2}\,dt \overset{\text{FTC1}}{\Rightarrow} A'(w) = -\dfrac{d}{dw}\int_{-1}^w e^{t+t^2} = -e^{w+w^2}$

15. Let $u = e^x$. Then $\dfrac{du}{dx} = e^x$. Also, $\dfrac{dh}{dx} = \dfrac{dh}{du}\dfrac{du}{dx}$, so by FTC1,

$$h'(x) = \frac{d}{dx}\int_1^{e^x} \ln t\,dt = \frac{d}{du}\int_1^u \ln t\,dt \cdot \frac{du}{dx} = \ln u \frac{du}{dx} = (\ln e^x) \cdot e^x = xe^x.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. Let $u = \sqrt{x}$. Then $\dfrac{du}{dx} = \dfrac{1}{2\sqrt{x}}$. Also, $\dfrac{dh}{dx} = \dfrac{dh}{du}\dfrac{du}{dx}$, so by FTC1,

$$h'(x) = \frac{d}{dx}\int_1^{\sqrt{x}} \frac{z^2}{z^4+1}\,dz = \frac{d}{du}\int_1^{u} \frac{z^2}{z^4+1}\,dz \cdot \frac{du}{dx} = \frac{u^2}{u^4+1}\,\frac{du}{dx} = \frac{x}{x^2+1}\,\frac{1}{2\sqrt{x}} = \frac{\sqrt{x}}{2(x^2+1)}.$$

17. Let $u = 3x+2$. Then $\dfrac{du}{dx} = 3$. Also, $\dfrac{dy}{dx} = \dfrac{dy}{du}\dfrac{du}{dx}$, so by FTC1,

$$y' = \frac{d}{dx}\int_1^{3x+2} \frac{t}{1+t^3}\,dt = \frac{d}{du}\int_1^{u} \frac{t}{1+t^3}\,dt \cdot \frac{du}{dx} = \frac{u}{1+u^3}\,\frac{du}{dx} = \frac{3x+2}{1+(3x+2)^3}\cdot 3 = \frac{3(3x+2)}{1+(3x+2)^3}$$

18. Let $u = \tan x$. Then $\dfrac{du}{dx} = \sec^2 x$. Also, $\dfrac{dy}{dx} = \dfrac{dy}{du}\dfrac{du}{dx}$, so by FTC1,

$$y' = \frac{d}{dx}\int_0^{\tan x} e^{-t^2}\,dt = \frac{d}{du}\int_0^{u} e^{-t^2}\,dt \cdot \frac{du}{dx} = e^{-u^2}\cdot \sec^2 x = e^{-\tan^2 x}\cdot \sec^2 x$$

19. Let $u = \sqrt{x}$. Then $\dfrac{du}{dx} = \dfrac{1}{2\sqrt{x}}$. Also, $\dfrac{dy}{dx} = \dfrac{dy}{du}\dfrac{du}{dx}$, so by FTC1,

$$y' = \frac{d}{dx}\int_{\sqrt{x}}^{\pi/4} \theta\tan\theta\,d\theta = -\frac{d}{du}\int_{\pi/4}^{\sqrt{x}} \theta\tan\theta\,d\theta \cdot \frac{du}{dx} = -u\tan u\,\frac{du}{dx} = -\sqrt{x}\tan\sqrt{x}\cdot\frac{1}{2\sqrt{x}} = -\frac{1}{2}\tan\sqrt{x}$$

20. $y = \displaystyle\int_{1/x}^{4} \sqrt{1+\frac{1}{t}}\,dt = -\int_4^{1/x} \sqrt{1+\frac{1}{t}}\,dt$. Let $u = \dfrac{1}{x}$. Then $\dfrac{du}{dx} = -\dfrac{1}{x^2}$. Also, $\dfrac{dy}{dx} = \dfrac{dy}{du}\dfrac{du}{dx}$, so by FTC1,

$$\begin{aligned} y' &= \frac{d}{dx}\left[-\int_4^{1/x} \sqrt{1+\frac{1}{t}}\,dt\right] = -\frac{d}{du}\int_4^{u} \sqrt{1+\frac{1}{t}}\,dt \cdot \frac{du}{dx} \\ &= -\sqrt{1+\frac{1}{u}}\cdot\left(-\frac{1}{x^2}\right) = -\sqrt{1+\frac{1}{1/x}}\cdot\left(-\frac{1}{x^2}\right) = \frac{1}{x^2}\sqrt{1+x} \end{aligned}$$

21. $\int_{-1}^{2} x^3\,dx = \left[\frac{1}{4}x^4\right]_{-1}^{2} = 4 - \frac{1}{4} = \frac{15}{4} = 3.75$

22. $\int_0^4 (x^2 - 4x)\,dx = \left[\frac{1}{3}x^3 - 2x^2\right]_0^4$

$= \left(\frac{64}{3} - 32\right) - 0 = -\frac{32}{3}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. $\int_{\pi/2}^{2\pi}(2\sin x)\,dx = \Big[-2\cos x\Big]_{\pi/2}^{2\pi}$

$= (-2\cos 2\pi) - \left(-2\cos\frac{\pi}{2}\right)$

$= -2(1) + 2(0) = -2$

24. $\int_{-1}^{2}(e^x + 2)\,dx = \Big[e^x + 2x\Big]_{-1}^{2}$

$= [e^2 + 2(2)] - \left[e^{-1} + 2(-1)\right]$

$= e^2 - e^{-1} + 6$

25. $\int_1^3 (x^2 + 2x - 4)\,dx = \left[\frac{1}{3}x^3 + x^2 - 4x\right]_1^3 = (9 + 9 - 12) - \left(\frac{1}{3} + 1 - 4\right) = 6 + \frac{8}{3} = \frac{26}{3}$

26. $\int_{-1}^{1} x^{100}\,dx = \left[\frac{1}{101}x^{101}\right]_{-1}^{1} = \frac{1}{101} - \left(-\frac{1}{101}\right) = \frac{2}{101}$

27. $\int_0^2 \left(\frac{4}{5}t^3 - \frac{3}{4}t^2 + \frac{2}{5}t\right)dt = \left[\frac{1}{5}t^4 - \frac{1}{4}t^3 + \frac{1}{5}t^2\right]_0^2 = \left(\frac{16}{5} - 2 + \frac{4}{5}\right) - 0 = 2$

28. $\int_0^1 (1 - 8v^3 + 16v^7)\,dv = \left[v - 2v^4 + 2v^8\right]_0^1 = (1 - 2 + 2) - 0 = 1$

29. $\displaystyle\int_1^9 \sqrt{x}\,dx = \int_1^9 x^{1/2}\,dx = \left[\frac{x^{3/2}}{3/2}\right]_1^9 = \frac{2}{3}\left[x^{3/2}\right]_1^9 = \frac{2}{3}(9^{3/2} - 1^{3/2}) = \frac{2}{3}(27 - 1) = \frac{52}{3}$

30. $\displaystyle\int_1^8 x^{-2/3}\,dx = \left[\frac{x^{1/3}}{1/3}\right]_1^8 = 3\left[x^{1/3}\right]_1^8 = 3(8^{1/3} - 1^{1/3}) = 3(2 - 1) = 3$

31. $\displaystyle\int_0^4 (t^2 + t^{3/2})\,dt = \left[\frac{1}{3}t^3 + \frac{2}{5}t^{5/2}\right]_0^4 = \left(\frac{64}{3} + \frac{64}{5}\right) - 0 = \frac{512}{15}$

32. $\displaystyle\int_1^3 \left(\frac{1}{z^2} + \frac{1}{z^3}\right)dz = \left[-\frac{1}{z} - \frac{1}{2z^2}\right]_1^3 = \left(-\frac{1}{3} - \frac{1}{18}\right) - \left(-1 - \frac{1}{2}\right) = \frac{10}{9}$

33. $\int_{\pi/2}^{0} \cos\theta\,d\theta = \Big[\sin\theta\Big]_{\pi/2}^{0} = \sin 0 - \sin\dfrac{\pi}{2} = 0 - 1 = -1$

34. $\int_{-5}^{5} e\,dx = \Big[ex\Big]_{-5}^{5} = 5e - (-5e) = 10e$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. $\displaystyle\int_0^1 (u+2)(u-3)\,du = \int_0^1 (u^2-u-6)\,du = \left[\tfrac{1}{3}u^3 - \tfrac{1}{2}u^2 - 6u\right]_0^1 = \left(\tfrac{1}{3}-\tfrac{1}{2}-6\right) - 0 = -\tfrac{37}{6}$

36. $\displaystyle\int_0^4 (4-t)\sqrt{t}\,dt = \int_0^4 (4-t)t^{1/2}\,dt = \int_0^4 (4t^{1/2} - t^{3/2})\,dt = \left[\tfrac{8}{3}t^{3/2} - \tfrac{2}{5}t^{5/2}\right]_0^4 = \tfrac{8}{3}(8) - \tfrac{2}{5}(32) = \tfrac{320-192}{15} = \tfrac{128}{15}$

37. $\displaystyle\int_1^4 \frac{2+x^2}{\sqrt{x}}\,dx = \int_1^4 \left(\frac{2}{\sqrt{x}} + \frac{x^2}{\sqrt{x}}\right) dx = \int_1^4 (2x^{-1/2} + x^{3/2})\,dx$

$\displaystyle = \left[4x^{1/2} + \tfrac{2}{5}x^{5/2}\right]_1^4 = \left[4(2) + \tfrac{2}{5}(32)\right] - \left(4 + \tfrac{2}{5}\right) = 8 + \tfrac{64}{5} - 4 - \tfrac{2}{5} = \tfrac{82}{5}$

38. $\int_{-1}^2 (3u-2)(u+1)\,du = \int_{-1}^2 (3u^2+u-2)\,du = \left[u^3 + \tfrac{1}{2}u^2 - 2u\right]_{-1}^2 = (8+2-4) - \left(-1+\tfrac{1}{2}+2\right) = 6 - \tfrac{3}{2} = \tfrac{9}{2}$

39. $\displaystyle\int_1^3 \left(2x + \frac{1}{x}\right) dx = \left[x^2 + \ln|x|\right]_1^3 = (9+\ln 3) - (1+\ln 1) = 8 + \ln 3$

40. $\int_5^5 \sqrt{t^2+\sin t}\,dt = F(5) - F(5)$ [where F is any antiderivative of $\sqrt{t^2-5t}$] $= 0$

41. $\int_0^{\pi/3} \sec\theta\,\tan\theta\,d\theta = \left[\sec\theta\right]_0^{\pi/3} = \sec\dfrac{\pi}{3} - \sec 0 = 2 - 1 = 1$

42. $\displaystyle\int_1^3 \frac{y^3-2y^2-y}{y^2}\,dy = \int_1^3 \left(y - 2 - \frac{1}{y}\right) dy = \left[\tfrac{1}{2}y^2 - 2y - \ln|y|\right]_1^3 = \left(\tfrac{9}{2} - 6 - \ln 3\right) - \left(\tfrac{1}{2} - 2 - 0\right) = -\ln 3$

43. $\int_0^1 (1+r)^3\,dr = \int_0^1 (1+3r+3r^2+r^3)\,dr = \left[r + \tfrac{3}{2}r^2 + r^3 + \tfrac{1}{4}r^4\right]_0^1 = \left(1 + \tfrac{3}{2} + 1 + \tfrac{1}{4}\right) - 0 = \tfrac{15}{4}$

44. $\displaystyle\int_0^3 (2\sin x - e^x)\,dx = \left[-2\cos x - e^x\right]_0^3 = (-2\cos 3 - e^3) - (-2-1) = 3 - 2\cos 3 - e^3$

45. $\displaystyle\int_1^2 \frac{v^3+3v^6}{v^4} = \int_1^2 \left(\frac{1}{v} + 3v^2\right) dv = \left[\ln|v| + v^3\right]_1^2 = (\ln 2 + 8) - (\ln 1 + 1) = \ln 2 + 7$

46. $\displaystyle\int_1^{18} \sqrt{\frac{3}{z}}\,dz = \int_1^{18} \sqrt{3}z^{-1/2}\,dz = \sqrt{3}\left[2z^{1/2}\right]_1^{18} = 2\sqrt{3}(18^{1/2} - 1^{1/2}) = 2\sqrt{3}(3\sqrt{2} - 1)$

47. $\displaystyle\int_0^1 (x^e + e^x)\,dx = \left[\frac{x^{e+1}}{e+1} + e^x\right]_0^1 = \left(\frac{1}{e+1} + e\right) - (0+1) = \frac{1}{e+1} + e - 1$

48. $\int_0^1 \cosh t\,dt = \left[\sinh t\right]_0^1 = \sinh 1 - \sinh 0 = \sinh 1 \quad \left[\text{or } \tfrac{1}{2}(e - e^{-1})\right]$

49. $\displaystyle\int_{1/\sqrt{3}}^{\sqrt{3}} \frac{8}{1+x^2}\,dx = \left[8\arctan x\right]_{1/\sqrt{3}}^{\sqrt{3}} = 8\left(\frac{\pi}{3} - \frac{\pi}{6}\right) = 8\left(\frac{\pi}{6}\right) = \frac{4\pi}{3}$

50. $\displaystyle\int_1^3 \frac{(3x+1)^2}{x^3}\,dx = \int_1^3 \frac{9x^2+6x+1}{x^3}\,dx = \int_1^3 \left(\frac{9}{x} + \frac{6}{x^2} + \frac{1}{x^3}\right) dx = \left[9\ln|x| - \frac{6}{x} - \frac{1}{2x^2}\right]_1^3$

$\displaystyle = \left(9\ln 3 - 2 - \tfrac{1}{18}\right) - \left(9\ln 1 - 6 - \tfrac{1}{2}\right) = 9\ln 3 + \tfrac{40}{9}$

51. $\displaystyle\int_0^4 2^s\,ds = \left[\frac{1}{\ln 2}2^s\right]_0^4 = \frac{16}{\ln 2} - \frac{1}{\ln 2} = \frac{15}{\ln 2}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

52. $\displaystyle\int_{1/2}^{1/\sqrt{2}} \frac{4}{\sqrt{1-x^2}}\,dx = \Big[4\arcsin x\Big]_{1/2}^{1/\sqrt{2}} = 4\Big(\frac{\pi}{4}-\frac{\pi}{6}\Big) = 4\Big(\frac{\pi}{12}\Big) = \frac{\pi}{3}$

53. If $f(x) = \begin{cases} \sin x & \text{if } 0 \le x < \pi/2 \\ \cos x & \text{if } \pi/2 \le x \le \pi \end{cases}$ then

$$\int_0^{\pi} f(x)\,dx = \int_0^{\pi/2} \sin x\,dx + \int_{\pi/2}^{\pi} \cos x\,dx = \big[-\cos x\big]_0^{\pi/2} + \big[\sin x\big]_{\pi/2}^{\pi} = -\cos\tfrac{\pi}{2} + \cos 0 + \sin\pi - \sin\tfrac{\pi}{2}$$
$$= -0 + 1 + 0 - 1 = 0$$

Note that f is integrable by Theorem 3 in Section 5.2.

54. If $f(x) = \begin{cases} 2 & \text{if } -2 \le x \le 0 \\ 4-x^2 & \text{if } 0 < x \le 2 \end{cases}$ then

$$\int_{-2}^{2} f(x)\,dx = \int_{-2}^{0} 2\,dx + \int_0^2 (4-x^2)\,dx = \big[2x\big]_{-2}^{0} + \big[4x - \tfrac{1}{3}x^3\big]_0^2 = [0-(-4)] + \big(\tfrac{16}{3} - 0\big) = \tfrac{28}{3}$$

Note that f is integrable by Theorem 3 in Section 5.2.

55. Area $= \displaystyle\int_0^4 \sqrt{x}\,dx = \int_0^4 x^{1/2}\,dx = \Big[\tfrac{2}{3}x^{3/2}\Big]_0^4 = \tfrac{2}{3}(8) - 0 = \tfrac{16}{3}$

56. Area $= \displaystyle\int_0^1 x^3\,dx = \big[\tfrac{1}{4}x^4\big]_0^1 = \tfrac{1}{4} - 0 = \tfrac{1}{4}$

57. Area $= \displaystyle\int_{-2}^{2} (4-x^2)\,dx = \big[4x - \tfrac{1}{3}x^3\big]_{-2}^{2} = \big(8 - \tfrac{8}{3}\big) - \big(-8 + \tfrac{8}{3}\big) = \tfrac{32}{3}$

58. Area $= \displaystyle\int_0^2 (2x - x^2)\,dx = \big[x^2 - \tfrac{1}{3}x^3\big]_0^2 = \big(4 - \tfrac{8}{3}\big) - 0 = \tfrac{4}{3}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. From the graph, it appears that the area is about 60. The actual area is $\int_0^{27} x^{1/3}dx = \left[\frac{3}{4}x^{4/3}\right]_0^{27} = \frac{3}{4}\cdot 81 - 0 = \frac{243}{4} = 60.75$. This is $\frac{3}{4}$ of the area of the viewing rectangle.

60. From the graph, it appears that the area is about $\frac{1}{3}$. The actual area is

$$\int_1^6 x^{-4}\,dx = \left[\frac{x^{-3}}{-3}\right]_1^6 = \left[\frac{-1}{3x^3}\right]_1^6 = -\frac{1}{3\cdot 216} + \frac{1}{3} = \frac{215}{648} \approx 0.3318.$$

61. It appears that the area under the graph is about $\frac{2}{3}$ of the area of the viewing rectangle, or about $\frac{2}{3}\pi \approx 2.1$. The actual area is $\int_0^{\pi} \sin x\,dx = [-\cos x]_0^{\pi} = (-\cos\pi) - (-\cos 0) = -(-1) + 1 = 2$.

62. Splitting up the region as shown, we estimate that the area under the graph is $\frac{\pi}{3} + \frac{1}{4}\left(3\cdot\frac{\pi}{3}\right) \approx 1.8$. The actual area is $\int_0^{\pi/3} \sec^2 x\,dx = [\tan x]_0^{\pi/3} = \sqrt{3} - 0 = \sqrt{3} \approx 1.73$.

63. $f(x) = x^{-4}$ is not continuous on the interval $[-2, 1]$, so FTC2 cannot be applied. In fact, f has an infinite discontinuity at $x = 0$, so $\int_{-2}^{1} x^{-4}\,dx$ does not exist.

64. $f(x) = \dfrac{4}{x^3}$ is not continuous on the interval $[-1, 2]$, so FTC2 cannot be applied. In fact, f has an infinite discontinuity at $x = 0$, so $\displaystyle\int_{-1}^{2} \frac{4}{x^3}\,dx$ does not exist.

65. $f(\theta) = \sec\theta\,\tan\theta$ is not continuous on the interval $[\pi/3, \pi]$, so FTC2 cannot be applied. In fact, f has an infinite discontinuity at $x = \pi/2$, so $\int_{\pi/3}^{\pi} \sec\theta\,\tan\theta\,d\theta$ does not exist.

66. $f(x) = \sec^2 x$ is not continuous on the interval $[0, \pi]$, so FTC2 cannot be applied. In fact, f has an infinite discontinuity at $x = \pi/2$, so $\int_0^{\pi} \sec^2 x\,dx$ does not exist.

67. $$g(x) = \int_{2x}^{3x} \frac{u^2-1}{u^2+1}\,du = \int_{2x}^{0} \frac{u^2-1}{u^2+1}\,du + \int_0^{3x} \frac{u^2-1}{u^2+1}\,du = -\int_0^{2x} \frac{u^2-1}{u^2+1}\,du + \int_0^{3x} \frac{u^2-1}{u^2+1}\,du \quad \Rightarrow$$

$$g'(x) = -\frac{(2x)^2-1}{(2x)^2+1}\cdot\frac{d}{dx}(2x) + \frac{(3x)^2-1}{(3x)^2+1}\cdot\frac{d}{dx}(3x) = -2\cdot\frac{4x^2-1}{4x^2+1} + 3\cdot\frac{9x^2-1}{9x^2+1}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

68. $g(x) = \int_{1-2x}^{1+2x} t\sin t\,dt = \int_{1-2x}^{0} t\sin t\,dt + \int_{0}^{1+2x} t\sin t\,dt = -\int_{0}^{1-2x} t\sin t\,dt + \int_{0}^{1+2x} t\sin t\,dt \quad \Rightarrow$

$$\begin{aligned} g'(x) &= -(1-2x)\sin(1-2x)\cdot\frac{d}{dx}(1-2x) + (1+2x)\sin(1+2x)\cdot\frac{d}{dx}(1+2x) \\ &= 2(1-2x)\sin(1-2x) + 2(1+2x)\sin(1+2x) \end{aligned}$$

69. $F(x) = \int_{x}^{x^2} e^{t^2}\,dt = \int_{x}^{0} e^{t^2}\,dt + \int_{0}^{x^2} e^{t^2}\,dt = -\int_{0}^{x} e^{t^2}\,dt + \int_{0}^{x^2} e^{t^2}\,dt \quad \Rightarrow$

$F'(x) = -e^{x^2} + e^{(x^2)^2}\cdot\frac{d}{dx}(x^2) = -e^{x^2} + 2xe^{x^4}$

70. $F(x) = \int_{\sqrt{x}}^{2x} \arctan t\,dt = \int_{\sqrt{x}}^{0} \arctan t\,dt + \int_{0}^{2x} \arctan t\,dt = -\int_{0}^{\sqrt{x}} \arctan t\,dt + \int_{0}^{2x} \arctan t\,dt \quad \Rightarrow$

$F'(x) = -\arctan\sqrt{x}\cdot\frac{d}{dx}(\sqrt{x}) + \arctan 2x\cdot\frac{d}{dx}(2x) = -\frac{1}{2\sqrt{x}}\arctan\sqrt{x} + 2\arctan 2x$

71. $$\begin{aligned} y &= \int_{\cos x}^{\sin x} \ln(1+2v)\,dv = \int_{\cos x}^{0} \ln(1+2v)\,dv + \int_{0}^{\sin x} \ln(1+2v)\,dv \\ &= -\int_{0}^{\cos x} \ln(1+2v)\,dv + \int_{0}^{\sin x} \ln(1+2v)\,dv \quad \Rightarrow \end{aligned}$$

$y' = -\ln(1+2\cos x)\cdot\frac{d}{dx}\cos x + \ln(1+2\sin x)\cdot\frac{d}{dx}\sin x = \sin x\,\ln(1+2\cos x) + \cos x\,\ln(1+2\sin x)$

72. $f(x) = \int_{0}^{x} (1-t^2)e^{t^2}\,dt$ is increasing when $f'(x) = (1-x^2)e^{x^2}$ is positive.

Since $e^{x^2} > 0$, $f'(x) > 0 \quad \Leftrightarrow \quad 1 - x^2 > 0 \quad \Leftrightarrow \quad |x| < 1$, so f is increasing on $(-1, 1)$.

73. $y = \int_{0}^{x} \frac{t^2}{t^2+t+2}\,dt \quad \Rightarrow \quad y' = \frac{x^2}{x^2+x+2} \quad \Rightarrow$

$$y'' = \frac{(x^2+x+2)(2x) - x^2(2x+1)}{(x^2+x+2)^2} = \frac{2x^3+2x^2+4x-2x^3-x^2}{(x^2+x+2)^2} = \frac{x^2+4x}{(x^2+x+2)^2} = \frac{x(x+4)}{(x^2+x+2)^2}.$$

The curve y is concave downward when $y'' < 0$; that is, on the interval $(-4, 0)$.

74. If $F(x) = \int_1^x f(t)\,dt$, then by FTC1, $F'(x) = f(x)$, and also, $F''(x) = f'(x)$. F is concave downward where F'' is negative; that is, where f' is negative. The given graph shows that f is decreasing ($f' < 0$) on the interval $(-1, 1)$.

75. $F(x) = \int_2^x e^{t^2}\,dt \quad \Rightarrow \quad F'(x) = e^{x^2}$, so the slope at $x = 2$ is $e^{2^2} = e^4$. The y-coordinate of the point on F at $x = 2$ is $F(2) = \int_2^2 e^{t^2}\,dt = 0$ since the limits are equal. An equation of the tangent line is $y - 0 = e^4(x-2)$, or $y = e^4x - 2e^4$.

76. $g(y) = \int_3^y f(x)\,dx \quad \Rightarrow \quad g'(y) = f(y)$. Since $f(x) = \int_0^{\sin x} \sqrt{1+t^2}\,dt$, $g''(y) = f'(y) = \sqrt{1+\sin^2 y}\cdot\cos y$, so $g''(\frac{\pi}{6}) = \sqrt{1+\sin^2(\frac{\pi}{6})}\cdot\cos\frac{\pi}{6} = \sqrt{1+(\frac{1}{2})^2}\cdot\frac{\sqrt{3}}{2} = \frac{\sqrt{5}}{2}\cdot\frac{\sqrt{3}}{2} = \frac{\sqrt{15}}{4}$.

77. $\lim_{x\to 0}\frac{1}{x^2}\int_0^x \frac{2t}{\sqrt{t^3+1}}\,dt = \lim_{x\to 0}\frac{\int_0^x \frac{2t}{\sqrt{t^3+1}}\,dt}{x^2}$ $\left[\text{form } \frac{0}{0}\right] \overset{\text{H}}{=} \lim_{x\to 0}\frac{\frac{2x}{\sqrt{x^3+1}}}{2x} = \lim_{x\to 0}\frac{1}{\sqrt{x^3+1}} = 1$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

78. $\displaystyle\lim_{x\to\infty}\frac{1}{x^2}\int_0^x \ln(1+e^t)\,dt = \lim_{x\to\infty}\frac{\int_0^x \ln(1+e^t)\,dt}{x^2}$ $\left[\text{form } \frac{\infty}{\infty}\right]$ $\overset{\text{H}}{=} \displaystyle\lim_{x\to\infty}\frac{\ln(1+e^x)}{2x} \overset{\text{H}}{=} \lim_{x\to\infty}\frac{\frac{e^x}{1+e^x}}{2}$

$$= \lim_{x\to\infty}\frac{e^x}{2(1+e^x)} = \lim_{x\to\infty}\frac{e^x/e^x}{2(1+e^x)/e^x} = \lim_{x\to\infty}\frac{1}{2\left(\frac{1}{e^x}+1\right)} = \frac{1}{2(0+1)} = \frac{1}{2}$$

79. By FTC2, $\int_1^4 f'(x)\,dx = f(4)-f(1)$, so $17 = f(4)-12 \;\Rightarrow\; f(4) = 17+12 = 29$.

80. (a) $\operatorname{erf}(x) = \dfrac{2}{\sqrt{\pi}}\displaystyle\int_0^x e^{-t^2}\,dt \;\Rightarrow\; \int_0^x e^{-t^2}\,dt = \frac{\sqrt{\pi}}{2}\operatorname{erf}(x)$. By Property 5 of definite integrals in Section 5.2,

$\int_0^b e^{-t^2}\,dt = \int_0^a e^{-t^2}\,dt + \int_a^b e^{-t^2}\,dt$, so

$$\int_a^b e^{-t^2}\,dt = \int_0^b e^{-t^2}\,dt - \int_0^a e^{-t^2}\,dt = \frac{\sqrt{\pi}}{2}\operatorname{erf}(b) - \frac{\sqrt{\pi}}{2}\operatorname{erf}(a) = \tfrac{1}{2}\sqrt{\pi}\,[\operatorname{erf}(b)-\operatorname{erf}(a)].$$

(b) $y = e^{x^2}\operatorname{erf}(x) \;\Rightarrow\; y' = 2xe^{x^2}\operatorname{erf}(x) + e^{x^2}\operatorname{erf}'(x) = 2xy + e^{x^2}\cdot\dfrac{2}{\sqrt{\pi}}e^{-x^2}$ [by FTC1] $= 2xy + \dfrac{2}{\sqrt{\pi}}$.

81. (a) The Fresnel function $S(x) = \int_0^x \sin\left(\frac{\pi}{2}t^2\right)dt$ has local maximum values where $0 = S'(x) = \sin\left(\frac{\pi}{2}t^2\right)$ and S' changes from positive to negative. For $x > 0$, this happens when $\frac{\pi}{2}x^2 = (2n-1)\pi$ [odd multiples of π] $\Leftrightarrow$ $x^2 = 2(2n-1) \;\Leftrightarrow\; x = \sqrt{4n-2}$, n any positive integer. For $x < 0$, S' changes from positive to negative where $\frac{\pi}{2}x^2 = 2n\pi$ [even multiples of π] $\Leftrightarrow$ $x^2 = 4n \;\Leftrightarrow\; x = -2\sqrt{n}$. S' does not change sign at $x = 0$.

(b) S is concave upward on those intervals where $S''(x) > 0$. Differentiating our expression for $S'(x)$, we get $S''(x) = \cos\left(\frac{\pi}{2}x^2\right)\left(2\frac{\pi}{2}x\right) = \pi x\cos\left(\frac{\pi}{2}x^2\right)$. For $x > 0$, $S''(x) > 0$ where $\cos(\frac{\pi}{2}x^2) > 0 \;\Leftrightarrow\; 0 < \frac{\pi}{2}x^2 < \frac{\pi}{2}$ or $\left(2n-\frac{1}{2}\right)\pi < \frac{\pi}{2}x^2 < \left(2n+\frac{1}{2}\right)\pi$, n any integer $\Leftrightarrow$ $0 < x < 1$ or $\sqrt{4n-1} < x < \sqrt{4n+1}$, n any positive integer. For $x < 0$, $S''(x) > 0$ where $\cos(\frac{\pi}{2}x^2) < 0 \;\Leftrightarrow\; \left(2n-\frac{3}{2}\right)\pi < \frac{\pi}{2}x^2 < \left(2n-\frac{1}{2}\right)\pi$, n any integer $\Leftrightarrow$ $4n-3 < x^2 < 4n-1 \;\Leftrightarrow\; \sqrt{4n-3} < |x| < \sqrt{4n-1} \;\Rightarrow\; \sqrt{4n-3} < -x < \sqrt{4n-1} \;\Rightarrow$ $-\sqrt{4n-3} > x > -\sqrt{4n-1}$, so the intervals of upward concavity for $x < 0$ are $\left(-\sqrt{4n-1}, -\sqrt{4n-3}\right)$, n any positive integer. To summarize: S is concave upward on the intervals $(0,1)$, $\left(-\sqrt{3},-1\right)$, $\left(\sqrt{3},\sqrt{5}\right)$, $\left(-\sqrt{7},-\sqrt{5}\right)$, $\left(\sqrt{7},3\right), \ldots$.

(c) In Maple, we use `plot({int(sin(Pi*t^2/2),t=0..x),0.2},x=0..2);`. Note that Maple recognizes the Fresnel function, calling it `FresnelS(x)`. In Mathematica, we use `Plot[{Integrate[Sin[Pi*t^2/2],{t,0,x}],0.2},{x,0,2}]`. In Derive, we load the utility file `FRESNEL` and plot `FRESNEL_SIN(x)`. From the graphs, we see that $\int_0^x \sin\left(\frac{\pi}{2}t^2\right)dt = 0.2$ at $x \approx 0.74$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

82. (a) In Maple, we should start by setting `si:=int(sin(t)/t,t=0..x);`. In Mathematica, the command is `si=Integrate[Sin[t]/t,{t,0,x}]`. Note that both systems recognize this function; Maple calls it `Si(x)` and Mathematica calls it `SinIntegral[x]`. In Maple, the command to generate the graph is `plot(si,x=-4*Pi..4*Pi);`. In Mathematica, it is `Plot[si,{x,-4*Pi,4*Pi}]`. In Derive, we load the utility file `EXP_INT` and plot `SI(x)`.

(b) $\mathrm{Si}(x)$ has local maximum values where $\mathrm{Si}'(x)$ changes from positive to negative, passing through 0. From the Fundamental Theorem we know that $\mathrm{Si}'(x) = \dfrac{d}{dx}\displaystyle\int_0^x \frac{\sin t}{t}\,dt = \frac{\sin x}{x}$, so we must have $\sin x = 0$ for a maximum, and for $x > 0$ we must have $x = (2n-1)\pi$, n any positive integer, for Si' to be changing from positive to negative at x. For $x < 0$, we must have $x = 2n\pi$, n any positive integer, for a maximum, since the denominator of $\mathrm{Si}'(x)$ is negative for $x < 0$. Thus, the local maxima occur at $x = \pi, -2\pi, 3\pi, -4\pi, 5\pi, -6\pi, \ldots$.

(c) To find the first inflection point, we solve $\mathrm{Si}''(x) = \dfrac{\cos x}{x} - \dfrac{\sin x}{x^2} = 0$. We can see from the graph that the first inflection point lies somewhere between $x = 3$ and $x = 5$. Using a rootfinder gives the value $x \approx 4.4934$. To find the y-coordinate of the inflection point, we evaluate $\mathrm{Si}(4.4934) \approx 1.6556$. So the coordinates of the first inflection point to the right of the origin are about $(4.4934, 1.6556)$. Alternatively, we could graph $S''(x)$ and estimate the first positive x-value at which it changes sign.

(d) It seems from the graph that the function has horizontal asymptotes at $y \approx 1.5$, with $\lim\limits_{x\to\pm\infty} \mathrm{Si}(x) \approx \pm 1.5$ respectively. Using the limit command, we get $\lim\limits_{x\to\infty} \mathrm{Si}(x) = \frac{\pi}{2}$. Since $\mathrm{Si}(x)$ is an odd function, $\lim\limits_{x\to-\infty} \mathrm{Si}(x) = -\frac{\pi}{2}$. So $\mathrm{Si}(x)$ has the horizontal asymptotes $y = \pm\frac{\pi}{2}$.

(e) We use the `fsolve` command in Maple (or `FindRoot` in Mathematica) to find that the solution is $x \approx 1.1$. Or, as in Exercise 81(c), we graph $y = \mathrm{Si}(x)$ and $y = 1$ on the same screen to see where they intersect.

83. (a) By FTC1, $g'(x) = f(x)$. So $g'(x) = f(x) = 0$ at $x = 1, 3, 5, 7$, and 9. g has local maxima at $x = 1$ and 5 (since $f = g'$ changes from positive to negative there) and local minima at $x = 3$ and 7. There is no local maximum or minimum at $x = 9$, since f is not defined for $x > 9$.

(b) We can see from the graph that $\left|\int_0^1 f\,dt\right| < \left|\int_1^3 f\,dt\right| < \left|\int_3^5 f\,dt\right| < \left|\int_5^7 f\,dt\right| < \left|\int_7^9 f\,dt\right|$. So $g(1) = \left|\int_0^1 f\,dt\right|$, $g(5) = \int_0^5 f\,dt = g(1) - \left|\int_1^3 f\,dt\right| + \left|\int_3^5 f\,dt\right|$, and $g(9) = \int_0^9 f\,dt = g(5) - \left|\int_5^7 f\,dt\right| + \left|\int_7^9 f\,dt\right|$. Thus, $g(1) < g(5) < g(9)$, and so the absolute maximum of $g(x)$ occurs at $x = 9$.

(c) g is concave downward on those intervals where $g'' < 0$. But $g'(x) = f(x)$, so $g''(x) = f'(x)$, which is negative on (approximately) $\left(\frac{1}{2}, 2\right)$, $(4, 6)$ and $(8, 9)$. So g is concave downward on these intervals.

(d)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

84. (a) By FTC1, $g'(x) = f(x)$. So $g'(x) = f(x) = 0$ at $x = 2, 4, 6, 8$, and 10. g has local maxima at $x = 2$ and 6 (since $f = g'$ changes from positive to negative there) and local minima at $x = 4$ and 8. There is no local maximum or minimum at $x = 10$, since f is not defined for $x > 10$.

(b) We can see from the graph that $\left|\int_0^2 f\,dt\right| > \left|\int_2^4 f\,dt\right| > \left|\int_4^6 f\,dt\right| > \left|\int_6^8 f\,dt\right| > \left|\int_8^{10} f\,dt\right|$. So $g(2) = \left|\int_0^2 f\,dt\right|$, $g(6) = \int_0^6 f\,dt = g(2) - \left|\int_2^4 f\,dt\right| + \left|\int_4^6 f\,dt\right|$, and $g(10) = \int_0^{10} f\,dt = g(6) - \left|\int_6^8 f\,dt\right| + \left|\int_8^{10} f\,dt\right|$. Thus, $g(2) > g(6) > g(10)$, and so the absolute maximum of $g(x)$ occurs at $x = 2$.

(c) g is concave downward on those intervals where $g'' < 0$. But $g'(x) = f(x)$, so $g''(x) = f'(x)$, which is negative on $(1, 3)$, $(5, 7)$ and $(9, 10)$. So g is concave downward on these intervals.

(d)

85. $$\lim_{n\to\infty} \sum_{i=1}^{n} \left(\frac{i^4}{n^5} + \frac{i}{n^2}\right) = \lim_{n\to\infty} \sum_{i=1}^{n} \left(\frac{i^4}{n^4} + \frac{i}{n}\right)\frac{1}{n} = \lim_{n\to\infty} \frac{1-0}{n} \sum_{i=1}^{n} \left[\left(\frac{i}{n}\right)^4 + \frac{i}{n}\right] = \int_0^1 (x^4 + x)\,dx$$
$$= \left[\tfrac{1}{5}x^5 + \tfrac{1}{2}x^2\right]_0^1 = \left(\tfrac{1}{5} + \tfrac{1}{2}\right) - 0 = \tfrac{7}{10}$$

86. $$\lim_{n\to\infty} \frac{1}{n}\left(\sqrt{\frac{1}{n}} + \sqrt{\frac{2}{n}} + \cdots + \sqrt{\frac{n}{n}}\right) = \lim_{n\to\infty} \frac{1-0}{n} \sum_{i=1}^{n} \sqrt{\frac{i}{n}} = \int_0^1 \sqrt{x}\,dx = \left[\frac{2x^{3/2}}{3}\right]_0^1 = \frac{2}{3} - 0 = \frac{2}{3}$$

87. Suppose $h < 0$. Since f is continuous on $[x + h, x]$, the Extreme Value Theorem says that there are numbers u and v in $[x + h, x]$ such that $f(u) = m$ and $f(v) = M$, where m and M are the absolute minimum and maximum values of f on $[x + h, x]$. By Integral Property 5.2.8, $m(-h) \le \int_{x+h}^{x} f(t)\,dt \le M(-h)$; that is, $f(u)(-h) \le -\int_x^{x+h} f(t)\,dt \le f(v)(-h)$. Since $-h > 0$, we can divide this inequality by $-h$: $f(u) \le \dfrac{1}{h}\displaystyle\int_x^{x+h} f(t)\,dt \le f(v)$. By Equation 2, $\dfrac{g(x+h) - g(x)}{h} = \dfrac{1}{h}\displaystyle\int_x^{x+h} f(t)\,dt$ for $h \ne 0$, and hence $f(u) \le \dfrac{g(x+h) - g(x)}{h} \le f(v)$, which is Equation 3 in the case where $h < 0$.

88. $$\frac{d}{dx}\int_{g(x)}^{h(x)} f(t)\,dt = \frac{d}{dx}\left[\int_{g(x)}^{0} f(t)\,dt + \int_0^{h(x)} f(t)\,dt\right] = \frac{d}{dx}\int_0^{h(x)} f(t)\,dt - \frac{d}{dx}\int_0^{g(x)} f(t)\,dt.$$
Using the Chain Rule in conjunction with FTC1 twice gives us $f(h(x))\,h'(x) - f(g(x))\,g'(x)$.

89. (a) Let $f(x) = \sqrt{x} \;\Rightarrow\; f'(x) = 1/(2\sqrt{x}) > 0$ for $x > 0 \;\Rightarrow\; f$ is increasing on $(0, \infty)$. If $x \ge 0$, then $x^3 \ge 0$, so $1 + x^3 \ge 1$ and since f is increasing, this means that $f(1 + x^3) \ge f(1) \;\Rightarrow\; \sqrt{1 + x^3} \ge 1$ for $x \ge 0$. Next let $g(t) = t^2 - t \;\Rightarrow\; g'(t) = 2t - 1 \;\Rightarrow\; g'(t) > 0$ when $t \ge 1$. Thus, g is increasing on $(1, \infty)$. And since $g(1) = 0$, $g(t) \ge 0$ when $t \ge 1$. Now let $t = \sqrt{1 + x^3}$, where $x \ge 0$. $\sqrt{1 + x^3} \ge 1$ (from above) $\;\Rightarrow\; t \ge 1 \;\Rightarrow\; g(t) \ge 0 \;\Rightarrow\; (1 + x^3) - \sqrt{1 + x^3} \ge 0$ for $x \ge 0$. Therefore, $1 \le \sqrt{1 + x^3} \le 1 + x^3$ for $x \ge 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) From part (a) and Integral Property 5.2.7: $\int_0^1 1\,dx \le \int_0^1 \sqrt{1+x^3}\,dx \le \int_0^1 (1+x^3)\,dx \quad \Leftrightarrow$

$\left[x\right]_0^1 \le \int_0^1 \sqrt{1+x^3}\,dx \le \left[x+\frac{1}{4}x^4\right]_0^1 \quad \Leftrightarrow \quad 1 \le \int_0^1 \sqrt{1+x^3}\,dx \le 1+\frac{1}{4} = 1.25.$

90. (a) For $0 \le x \le 1$, we have $x^2 \le x$. Since $f(x) = \cos x$ is a decreasing function on $[0,1]$, $\cos(x^2) \ge \cos x$.

(b) $\pi/6 < 1$, so by part (a), $\cos(x^2) \ge \cos x$ on $[0, \pi/6]$. Thus,

$\int_0^{\pi/6} \cos(x^2)\,dx \ge \int_0^{\pi/6} \cos x\,dx = \left[\sin x\right]_0^{\pi/6} = \sin(\pi/6) - \sin 0 = \frac{1}{2} - 0 = \frac{1}{2}.$

91. $0 < \dfrac{x^2}{x^4+x^2+1} < \dfrac{x^2}{x^4} = \dfrac{1}{x^2}$ on $[5, 10]$, so

$$0 \le \int_5^{10} \frac{x^2}{x^4+x^2+1}\,dx < \int_5^{10} \frac{1}{x^2}\,dx = \left[-\frac{1}{x}\right]_5^{10} = -\frac{1}{10} - \left(-\frac{1}{5}\right) = \frac{1}{10} = 0.1.$$

92. (a) If $x < 0$, then $g(x) = \int_0^x f(t)\,dt = \int_0^x 0\,dt = 0$.

If $0 \le x \le 1$, then $g(x) = \int_0^x f(t)\,dt = \int_0^x t\,dt = \left[\frac{1}{2}t^2\right]_0^x = \frac{1}{2}x^2$.

If $1 < x \le 2$, then

$g(x) = \int_0^x f(t)\,dt = \int_0^1 f(t)\,dt + \int_1^x f(t)\,dt = g(1) + \int_1^x (2-t)\,dt$

$= \frac{1}{2}(1)^2 + \left[2t - \frac{1}{2}t^2\right]_1^x = \frac{1}{2} + \left(2x - \frac{1}{2}x^2\right) - \left(2 - \frac{1}{2}\right) = 2x - \frac{1}{2}x^2 - 1.$

If $x > 2$, then $g(x) = \int_0^x f(t)\,dt = g(2) + \int_2^x 0\,dt = 1 + 0 = 1$. So

$$g(x) = \begin{cases} 0 & \text{if } x < 0 \\ \frac{1}{2}x^2 & \text{if } 0 \le x \le 1 \\ 2x - \frac{1}{2}x^2 - 1 & \text{if } 1 < x \le 2 \\ 1 & \text{if } x > 2 \end{cases}$$

(b)

(c) f is not differentiable at its corners at $x = 0$, 1, and 2. f is differentiable on $(-\infty, 0)$, $(0,1)$, $(1,2)$ and $(2, \infty)$.

g is differentiable on $(-\infty, \infty)$.

93. Using FTC1, we differentiate both sides of $6 + \displaystyle\int_a^x \frac{f(t)}{t^2}\,dt = 2\sqrt{x}$ to get $\dfrac{f(x)}{x^2} = 2\,\dfrac{1}{2\sqrt{x}} \quad \Rightarrow \quad f(x) = x^{3/2}$.

To find a, we substitute $x = a$ in the original equation to obtain $6 + \displaystyle\int_a^a \frac{f(t)}{t^2}\,dt = 2\sqrt{a} \quad \Rightarrow \quad 6 + 0 = 2\sqrt{a} \quad \Rightarrow$

$3 = \sqrt{a} \quad \Rightarrow \quad a = 9.$

94. $B = 3A \quad \Rightarrow \quad \int_0^b e^x\,dx = 3\int_0^a e^x\,dx \quad \Rightarrow \quad [e^x]_0^b = 3\,[e^x]_0^a \quad \Rightarrow \quad e^b - 1 = 3(e^a - 1) \quad \Rightarrow \quad e^b = 3e^a - 2 \quad \Rightarrow$

$b = \ln(3e^a - 2)$

95. (a) Let $F(t) = \int_0^t f(s)\,ds$. Then, by FTC1, $F'(t) = f(t) =$ rate of depreciation, so $F(t)$ represents the loss in value over the interval $[0, t]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $C(t) = \dfrac{1}{t}\left[A + \displaystyle\int_0^t f(s)\,ds\right] = \dfrac{A + F(t)}{t}$ represents the average expenditure per unit of t during the interval $[0, t]$, assuming that there has been only one overhaul during that time period. The company wants to minimize average expenditure.

(c) $C(t) = \dfrac{1}{t}\left[A + \displaystyle\int_0^t f(s)\,ds\right]$. Using FTC1, we have $C'(t) = -\dfrac{1}{t^2}\left[A + \displaystyle\int_0^t f(s)\,ds\right] + \dfrac{1}{t}f(t)$.

$$C'(t) = 0 \quad\Rightarrow\quad t\,f(t) = A + \int_0^t f(s)\,ds \quad\Rightarrow\quad f(t) = \frac{1}{t}\left[A + \int_0^t f(s)\,ds\right] = C(t).$$

5.4 Indefinite Integrals and the Net Change Theorem

1. $\dfrac{d}{dx}\left[x\ln x - x + C\right] = x\cdot\dfrac{1}{x} + (\ln x)\cdot 1 - 1 + 0 = 1 + \ln x - 1 = \ln x$

2. $\dfrac{d}{dx}\left(\tan x - x + C\right) = \sec^2 x - 1 + 0 = \tan^2 x$

3. $$\frac{d}{dx}\left[-\frac{\sqrt{1+x^2}}{x} + C\right] = \frac{d}{dx}\left[-\frac{(1+x^2)^{1/2}}{x} + C\right] = -\frac{x\cdot\frac{1}{2}(1+x^2)^{-1/2}(2x) - (1+x^2)^{1/2}\cdot 1}{(x)^2} + 0$$
$$= -\frac{(1+x^2)^{-1/2}\left[x^2 - (1+x^2)\right]}{x^2} = -\frac{-1}{(1+x^2)^{1/2}x^2} = \frac{1}{x^2\sqrt{1+x^2}}$$

4. $$\frac{d}{dx}\left[\frac{2}{15b^2}(3bx - 2a)(a+bx)^{3/2} + C\right] = \frac{2}{15b^2}\left[(3bx-2a)\tfrac{3}{2}(a+bx)^{1/2}(b) + (a+bx)^{3/2}(3b) + 0\right]$$
$$= \frac{2}{15b^2}(3b)(a+bx)^{1/2}\left[(3bx-2a)\tfrac{1}{2} + (a+bx)\right]$$
$$= \frac{2}{5b}(a+bx)^{1/2}\left(\frac{5}{2}bx\right) = x\sqrt{a+bx}$$

5. $\int(3x^2 + 4x + 1)\,dx = 3\cdot\frac{1}{3}x^3 + 4\cdot\frac{1}{2}x^2 + x + C = x^3 + 2x^2 + x + C$

6. $\int(5 + 2\sqrt{x}\,)\,dx = \int(5 + 2x^{1/2})\,dx = 5x + 2\cdot\frac{2}{3}x^{3/2} + C = 5x + \frac{4}{3}x^{3/2} + C$

7. $\int(x + \cos x)\,dx = \frac{1}{2}x^2 + \sin x + C$

8. $\displaystyle\int\left(\sqrt[3]{x} + \frac{1}{\sqrt[3]{x}}\right)dx = \int(x^{1/3} + x^{-1/3})\,dx = \frac{3}{4}x^{4/3} + \frac{3}{2}x^{2/3} + C$

9. $\displaystyle\int(x^{1.3} + 7x^{2.5})\,dx = \frac{1}{2.3}x^{2.3} + \frac{7}{3.5}x^{3.5} + C = \frac{1}{2.3}x^{2.3} + 2x^{3.5} + C$

10. $\int\sqrt[4]{x^5}\,dx = \int x^{5/4}\,dx = \frac{4}{9}x^{9/4} + C$

11. $\int\left(5 + \frac{2}{3}x^2 + \frac{3}{4}x^3\right)\,dx = 5x + \frac{2}{3}\cdot\frac{1}{3}x^3 + \frac{3}{4}\cdot\frac{1}{4}x^4 + C = 5x + \frac{2}{9}x^3 + \frac{3}{16}x^4 + C$

12. $\int(u^6 - 2u^5 - u^3 + \frac{2}{7})\,du = \frac{1}{7}u^7 - 2\cdot\frac{1}{6}u^6 - \frac{1}{4}u^4 + \frac{2}{7}u + C = \frac{1}{7}u^7 - \frac{1}{3}u^6 - \frac{1}{4}u^4 + \frac{2}{7}u + C$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $\displaystyle\int (u+4)(2u+1)\,du = \int (2u^2+9u+4)\,du = 2\,\frac{u^3}{3} + 9\,\frac{u^2}{2} + 4u + C = \frac{2}{3}u^3 + \frac{9}{2}u^2 + 4u + C$

14. $\displaystyle\int \sqrt{t}(t^2+3t+2)\,dt = \int t^{1/2}(t^2+3t+2)\,dt = \int (t^{5/2}+3t^{3/2}+2t^{1/2})\,dt$

$$= \tfrac{2}{7}t^{7/2} + 3\cdot\tfrac{2}{5}t^{5/2} + 2\cdot\tfrac{2}{3}t^{3/2} + C = \tfrac{2}{7}t^{7/2} + \tfrac{6}{5}t^{5/2} + \tfrac{4}{3}t^{3/2} + C$$

15. $\displaystyle\int \frac{1+\sqrt{x}+x}{x}\,dx = \int \left(\frac{1}{x} + \frac{\sqrt{x}}{x} + \frac{x}{x}\right) dx = \int \left(\frac{1}{x} + x^{-1/2} + 1\right) dx$

$$= \ln|x| + 2x^{1/2} + x + C = \ln|x| + 2\sqrt{x} + x + C$$

16. $\displaystyle\int \left(x^2 + 1 + \frac{1}{x^2+1}\right) dx = \frac{x^3}{3} + x + \tan^{-1} x + C$

17. $\displaystyle\int \left(e^x + \frac{1}{x}\right) dx = e^x + \ln|x| + C$

18. $\displaystyle\int (2+3^x)\,dx = 2x + \frac{3^x}{\ln 3} + C$

19. $\displaystyle\int (\sin x + \sinh x)\,dx = -\cos x + \cosh x + C$

20. $\displaystyle\int \left(\frac{1+r}{r}\right)^2 dr = \int \frac{1+2r+r^2}{r^2}\,dr = \int (r^{-2}+2r^{-1}+1)\,dr = -r^{-1} + 2\ln|r| + r + C = -\frac{1}{r} + 2\ln|r| + r + C$

21. $\displaystyle\int (2+\tan^2\theta)\,d\theta = \int [2+(\sec^2\theta - 1)]\,d\theta = \int (1+\sec^2\theta)\,d\theta = \theta + \tan\theta + C$

22. $\displaystyle\int \sec t\,(\sec t + \tan t)\,dt = \int (\sec^2 t + \sec t\,\tan t)\,dt = \tan t + \sec t + C$

23. $\displaystyle\int 3\csc^2 t\,dt = -3\cot t + C$

24. $\displaystyle\int \frac{\sin 2x}{\sin x}\,dx = \int \frac{2\sin x\,\cos x}{\sin x}\,dx = \int 2\cos x\,dx = 2\sin x + C$

25. $\displaystyle\int \left(\cos x + \tfrac{1}{2}x\right) dx = \sin x + \tfrac{1}{4}x^2 + C$. The members of the family in the figure correspond to $C = -5$, 0, 5, and 10.

26. $\displaystyle\int (e^x - 2x^2)\,dx = e^x - \tfrac{2}{3}x^3 + C$. The members of the family in the figure correspond to $C = -5$, 0, 2, and 5.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. $\int_{-2}^{3}(x^2-3)\,dx = \left[\frac{1}{3}x^3-3x\right]_{-2}^{3} = (9-9)-\left(-\frac{8}{3}+6\right) = \frac{8}{3}-\frac{18}{3} = -\frac{10}{3}$

28. $\int_{1}^{2}(4x^3-3x^2+2x)\,dx = \left[x^4-x^3+x^2\right]_{1}^{2} = (16-8+4)-(1-1+1) = 12-1 = 11$

29. $\displaystyle\int_1^4 (8t^3-6t^{-2})\,dt = \left[2t^4+\frac{6}{t}\right]_1^4 = \left(2\cdot 4^4+\frac{6}{4}\right)-\left(2\cdot 1^4+\frac{6}{1}\right) = \left(512+\frac{3}{2}\right)-(2+6) = \frac{1011}{2} = 505.5$

30. $\int_0^8 \left(\frac{1}{8}+\frac{1}{2}w+\frac{1}{3}w^{1/3}\right)dw = \left[\frac{1}{8}w+\frac{1}{4}w^2+\frac{1}{4}w^{4/3}\right]_0^8 = (1+16+4)-0 = 21$

31. $\int_0^2(2x-3)(4x^2+1)\,dx = \int_0^2(8x^3-12x^2+2x-3)\,dx = \left[2x^4-4x^3+x^2-3x\right]_0^2 = (32-32+4-6)-0 = -2$

32. $\displaystyle\int_1^2\left(\frac{1}{x^2}-\frac{4}{x^3}\right)dx = \int_1^2(x^{-2}-4x^{-3})\,dx = \left[\frac{x^{-1}}{-1}-\frac{4x^{-2}}{-2}\right]_1^2 = \left[-\frac{1}{x}+\frac{2}{x^2}\right]_1^2 = \left(-\frac{1}{2}+\frac{1}{2}\right)-(-1+2) = -1$

33. $\displaystyle\int_1^3\left(\frac{3x^2+4x+1}{x}\right)dx = \int_1^3\left(3x+4+\frac{1}{x}\right)dx = \left[\frac{3}{2}x^2+4x+\ln|x|\right]_1^3 = \left(\frac{27}{2}+12+\ln 3\right)-\left(\frac{3}{2}+4+\ln 1\right)$

$\displaystyle = 20+\ln 3$

34. $\int_{-1}^{1}t(1-t)^2\,dt = \int_{-1}^{1}t(1-2t+t^2)\,dt = \int_{-1}^{1}(t-2t^2+t^3)\,dt = \left[\frac{1}{2}t^2-\frac{2}{3}t^3+\frac{1}{4}t^4\right]_{-1}^{1}$

$= \left(\frac{1}{2}-\frac{2}{3}+\frac{1}{4}\right)-\left(\frac{1}{2}+\frac{2}{3}+\frac{1}{4}\right) = -\frac{4}{3}$

35. $\displaystyle\int_1^4\left(\frac{4+6u}{\sqrt{u}}\right)du = \int_1^4\left(\frac{4}{\sqrt{u}}+\frac{6u}{\sqrt{u}}\right)du = \int_1^4(4u^{-1/2}+6u^{1/2})\,du = \left[8u^{1/2}+4u^{3/2}\right]_1^4 = (16+32)-(8+4) = 36$

36. $\displaystyle\int_0^1\frac{4}{1+p^2}\,dp = \left[4\arctan p\right]_0^1 = 4\arctan 1-4\arctan 0 = 4\left(\frac{\pi}{4}\right)-4(0) = \pi$

37. $\displaystyle\int_{\pi/6}^{\pi/3}(4\sec^2 y)\,dy = \left[4\tan y\right]_{\pi/6}^{\pi/3} = 4\tan\frac{\pi}{3}-4\tan\frac{\pi}{6} = 4\cdot\sqrt{3}-4\cdot\frac{1}{\sqrt{3}} = 4\sqrt{3}-\frac{4\sqrt{3}}{3} = \frac{8\sqrt{3}}{3}$ or $\dfrac{8}{\sqrt{3}}$

38. $\displaystyle\int_0^{\pi/2}\left(\sqrt{t}-3\cos t\right)dt = \int_0^{\pi/2}\left(t^{1/2}-3\cos t\right)dt = \left[\frac{2}{3}t^{3/2}-3\sin t\right]_0^{\pi/2} = \left[\frac{2}{3}\left(\frac{\pi}{2}\right)^{3/2}-3\sin\frac{\pi}{2}\right]-0$

$\displaystyle = \frac{2}{3}\left(\frac{\pi}{2}\right)^{3/2}-3\cdot 1 = \frac{2}{3}\left(\frac{\pi}{2}\right)^{3/2}-3$

39. $\int_0^1 x\left(\sqrt[3]{x}+\sqrt[4]{x}\right)\,dx = \int_0^1(x^{4/3}+x^{5/4})\,dx = \left[\frac{3}{7}x^{7/3}+\frac{4}{9}x^{9/4}\right]_0^1 = \left(\frac{3}{7}+\frac{4}{9}\right)-0 = \frac{55}{63}$

40. $\displaystyle\int_1^4\frac{\sqrt{y}-y}{y^2}\,dy = \int_1^4\left(\frac{\sqrt{y}}{y^2}-\frac{y}{y^2}\right)dy = \int_1^4(y^{-3/2}-y^{-1})\,dy = \left[-2y^{-1/2}-\ln|y|\right]_1^4 = \left[-\frac{2}{\sqrt{y}}-\ln|y|\right]_1^4$

$= (-1-\ln 4)-(-2-\ln 1) = 1-\ln 4$

41. $\displaystyle\int_1^2\left(\frac{x}{2}-\frac{2}{x}\right)dx = \left[\frac{1}{4}x^2-2\ln|x|\right]_1^2 = (1-2\ln 2)-\left(\frac{1}{4}-2\ln 1\right) = \frac{3}{4}-2\ln 2$

42. $\displaystyle\int_0^1(5x-5^x)\,dx = \left[\frac{5}{2}x^2-\frac{5^x}{\ln 5}\right]_0^1 = \left(\frac{5}{2}-\frac{5}{\ln 5}\right)-\left(0-\frac{1}{\ln 5}\right) = \frac{5}{2}-\frac{4}{\ln 5}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. $\int_{-2}^{2}(\sinh x+\cosh x)\,dx=\Big[\cosh x+\sinh x\Big]_{-2}^{2}=(\cosh 2+\sinh 2)-(\cosh(-2)+\sinh(-2))$

$=(\cosh 2+\sinh 2)-(\cosh 2-\sinh 2)=2\sinh 2$

44. $\int_0^{\pi/4}(3e^x-4\sec x\,\tan x)\,dx=\Big[3e^x-4\sec x\Big]_0^{\pi/4}=\Big(3e^{\pi/4}-4\sec\dfrac{\pi}{4}\Big)-(3e^0-4\sec 0)$

$=\Big(3e^{\pi/4}-4\cdot\sqrt{2}\Big)-(3\cdot 1-4\cdot 1)=3e^{\pi/4}-4\sqrt{2}+1$

45. $\displaystyle\int_0^{\pi/4}\frac{1+\cos^2\theta}{\cos^2\theta}\,d\theta=\int_0^{\pi/4}\left(\frac{1}{\cos^2\theta}+\frac{\cos^2\theta}{\cos^2\theta}\right)d\theta=\int_0^{\pi/4}(\sec^2\theta+1)\,d\theta$

$=\big[\tan\theta+\theta\big]_0^{\pi/4}=\big(\tan\frac{\pi}{4}+\frac{\pi}{4}\big)-(0+0)=1+\frac{\pi}{4}$

46. $\displaystyle\int_0^{\pi/3}\frac{\sin\theta+\sin\theta\,\tan^2\theta}{\sec^2\theta}\,d\theta=\int_0^{\pi/3}\frac{\sin\theta\,(1+\tan^2\theta)}{\sec^2\theta}\,d\theta=\int_0^{\pi/3}\frac{\sin\theta\,\sec^2\theta}{\sec^2\theta}\,d\theta=\int_0^{\pi/3}\sin\theta\,d\theta$

$=\big[-\cos\theta\big]_0^{\pi/3}=-\frac{1}{2}-(-1)=\frac{1}{2}$

47. $\displaystyle\int_3^4\sqrt{\frac{3}{x}}\,dx=\int_3^4\sqrt{3}x^{-1/2}\,dx=\sqrt{3}\Big[2x^{1/2}\Big]_3^4=2\sqrt{3}\Big[2-\sqrt{3}\Big]=4\sqrt{3}-6$

48. $\displaystyle\int_{-10}^{10}\frac{2e^x}{\sinh x+\cosh x}\,dx=\int_{-10}^{10}\frac{2e^x}{\dfrac{e^x-e^{-x}}{2}+\dfrac{e^x+e^{-x}}{2}}\,dx=\int_{-10}^{10}\frac{2e^x}{e^x}\,dx=\int_{-10}^{10}2\,dx=\big[2x\big]_{-10}^{10}$

$=20-(-20)=40$

49. $\displaystyle\int_0^{\sqrt{3}/2}\frac{dr}{\sqrt{1-r^2}}=\Big[\arcsin r\Big]_0^{\sqrt{3}/2}=\arcsin\Big(\sqrt{3}/2\Big)-\arcsin 0=\frac{\pi}{3}-0=\frac{\pi}{3}$

50. $\displaystyle\int_{\pi/6}^{\pi/2}\csc t\,\cot t\,dt=\Big[-\csc t\Big]_{\pi/6}^{\pi/2}=\big(-\csc\tfrac{\pi}{2}\big)-\big(-\csc\tfrac{\pi}{6}\big)=-1-(-2)=1$

51. $\displaystyle\int_0^{1/\sqrt{3}}\frac{t^2-1}{t^4-1}\,dt=\int_0^{1/\sqrt{3}}\frac{t^2-1}{(t^2+1)(t^2-1)}\,dt=\int_0^{1/\sqrt{3}}\frac{1}{t^2+1}\,dt=\big[\arctan t\big]_0^{1/\sqrt{3}}=\arctan\Big(1/\sqrt{3}\Big)-\arctan 0$

$=\frac{\pi}{6}-0=\frac{\pi}{6}$

52. $|2x-1|=\begin{cases}2x-1 & \text{if } 2x-1\ge 0\\ -(2x-1) & \text{if } 2x-1<0\end{cases}=\begin{cases}2x-1 & \text{if } x\ge\frac{1}{2}\\ 1-2x & \text{if } x<\frac{1}{2}\end{cases}$

Thus, $\int_0^2|2x-1|\,dx=\int_0^{1/2}(1-2x)\,dx+\int_{1/2}^2(2x-1)\,dx=\big[x-x^2\big]_0^{1/2}+\big[x^2-x\big]_{1/2}^2$

$=(\frac{1}{2}-\frac{1}{4})-0+(4-2)-(\frac{1}{4}-\frac{1}{2})=\frac{1}{4}+2-(-\frac{1}{4})=\frac{5}{2}$

53. $\int_{-1}^2(x-2\,|x|)\,dx=\int_{-1}^0[x-2(-x)]\,dx+\int_0^2[x-2(x)]\,dx=\int_{-1}^0 3x\,dx+\int_0^2(-x)\,dx=3\big[\frac{1}{2}x^2\big]_{-1}^0-\big[\frac{1}{2}x^2\big]_0^2$

$=3\big(0-\frac{1}{2}\big)-(2-0)=-\frac{7}{2}=-3.5$

54. $\int_0^{3\pi/2}|\sin x|\;dx=\int_0^{\pi}\sin x\,dx+\int_{\pi}^{3\pi/2}(-\sin x)\,dx=\big[-\cos x\big]_0^{\pi}+\big[\cos x\big]_{\pi}^{3\pi/2}$

$=[1-(-1)]+[0-(-1)]=2+1=3$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

55. The graph shows that $y = 1 - 2x - 5x^4$ has x-intercepts at $x = a \approx -0.86$ and at $x = b \approx 0.42$. So the area of the region that lies under the curve and above the x-axis is

$$\int_a^b (1 - 2x - 5x^4)\,dx = \left[x - x^2 - x^5\right]_a^b$$
$$= (b - b^2 - b^5) - (a - a^2 - a^5) \approx 1.36$$

56. The graph shows that $y = (x^2 + 1)^{-1} - x^4$ has x-intercepts at $x = a \approx -0.87$ and at $x = b \approx 0.87$. So the area of the region that lies under the curve and above the x-axis is

$$\int_a^b \left[(x^2 + 1)^{-1} - x^4\right]\,dx = \left[\tan^{-1} x - \tfrac{1}{5}x^5\right]_a^b$$
$$= \left(\tan^{-1} b - \tfrac{1}{5}b^5\right) - \left(\tan^{-1} a - \tfrac{1}{5}a^5\right)$$
$$\approx 1.23$$

57. $A = \int_0^2 \left(2y - y^2\right)\,dy = \left[y^2 - \frac{1}{3}y^3\right]_0^2 = \left(4 - \frac{8}{3}\right) - 0 = \frac{4}{3}$

58. $y = \sqrt[4]{x} \quad \Rightarrow \quad x = y^4$, so $A = \int_0^1 y^4\,dy = \left[\frac{1}{5}y^5\right]_0^1 = \frac{1}{5}$.

59. If $w'(t)$ is the rate of change of weight in pounds per year, then $w(t)$ represents the weight in pounds of the child at age t. We know from the Net Change Theorem that $\int_5^{10} w'(t)\,dt = w(10) - w(5)$, so the integral represents the increase in the child's weight (in pounds) between the ages of 5 and 10.

60. $\int_a^b I(t)\,dt = \int_a^b Q'(t)\,dt = Q(b) - Q(a)$ by the Net Change Theorem, so it represents the change in the charge Q from time $t = a$ to $t = b$.

61. Since $r(t)$ is the rate at which oil leaks, we can write $r(t) = -V'(t)$, where $V(t)$ is the volume of oil at time t. [Note that the minus sign is needed because V is decreasing, so $V'(t)$ is negative, but $r(t)$ is positive.] Thus, by the Net Change Theorem, $\int_0^{120} r(t)\,dt = -\int_0^{120} V'(t)\,dt = -\left[V(120) - V(0)\right] = V(0) - V(120)$, which is the number of gallons of oil that leaked from the tank in the first two hours (120 minutes).

62. By the Net Change Theorem, $\int_0^{15} n'(t)\,dt = n(15) - n(0) = n(15) - 100$ represents the increase in the bee population in 15 weeks. So $100 + \int_0^{15} n'(t)\,dt = n(15)$ represents the total bee population after 15 weeks.

63. By the Net Change Theorem, $\int_{1000}^{5000} R'(x)\,dx = R(5000) - R(1000)$, so it represents the increase in revenue when production is increased from 1000 units to 5000 units.

64. The slope of the trail is the rate of change of the elevation E, so $f(x) = E'(x)$. By the Net Change Theorem, $\int_3^5 f(x)\,dx = \int_3^5 E'(x)\,dx = E(5) - E(3)$ is the change in the elevation E between $x = 3$ miles and $x = 5$ miles from the start of the trail.

65. The function h gives the rate of change in the total number of heartbeats H, so $h(t) = H'(t)$. By the Net Change Theorem, $\int_0^{30} h(t)\,dt = H(30) - H(0) = H(30) - 0 = H(30)$ represents the total number of heartbeats during the first 30 minutes of an exercise session.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

66. The units for $a(x)$ are pounds per foot and the units for x are feet, so the units for da/dx are pounds per foot per foot, denoted (lb/ft)/ft. The unit of measurement for $\int_2^8 a(x)\,dx$ is the product of pounds per foot and feet; that is, pounds.

67. In general, the unit of measurement for $\int_a^b f(x)\,dx$ is the product of the unit for $f(x)$ and the unit for x. Since $f(x)$ is measured in newtons and x is measured in meters, the units for $\int_0^{100} f(x)\,dx$ are newton-meters (or joules). (A newton-meter is abbreviated N·m.)

68. (a) At $t=2$, the distance from the charging station (the displacement) is given by $\int_0^2 v(t)\,dt = \frac{1}{2}\cdot 2\cdot 5$ [triangle] $= 5$ m.

At $t=4$, the displacement is $\int_0^4 v(t)\,dt = 5 + \int_2^4 v(t)\,dt = 5 + 2\cdot 2$ [square] $+ \frac{1}{2}\cdot 1\cdot 3$ [triangle] $= \frac{21}{2}$ m.

At $t=6$, the displacement is $\frac{21}{2} + \int_4^6 v(t)\,dt = \frac{21}{2} + \frac{1}{2}\cdot 2\cdot 2$ [triangle] $= \frac{25}{2}$ m.

At $t=8$, the displacement is $\frac{25}{2} + \int_6^8 v(t)\,dt = \frac{25}{2} + 0 = \frac{25}{2}$ m.

At $t=10$, the displacement is $\frac{25}{2} + \int_8^{10} v(t)\,dt = \frac{25}{2} - \frac{1}{2}\cdot 2\cdot 4$ [triangle] $= \frac{17}{2}$ m.

At $t=12$, the displacement is $\frac{17}{2} + \int_{10}^{12} v(t)\,dt = \frac{17}{2} - \frac{1}{2}\cdot 2\cdot 4$ [triangle] $= \frac{9}{2}$ m.

(b) The vehicle is farthest from the charging station when the absolute value of the displacement is greatest. The displacement of this autonomous vehicle is never negative, so we look for the largest values of the displacement integral (or, equivalently, the places where the area above the t-axis is the greatest). This occurs from $t=6$ to $t=8$, so the vehicle is farthest from the charging station then.

(c) The total distance traveled is given by $\int_0^{12} |v(t)|\,dt = \int_0^8 v(t)\,dt + \int_8^{12}[-v(t)]\,dt = \frac{25}{2} + \frac{1}{2}\cdot 4\cdot 4 = \frac{41}{2}$ m.

69. (a) Displacement $= \int_0^3 (3t-5)\,dt = \left[\frac{3}{2}t^2 - 5t\right]_0^3 = \frac{27}{2} - 15 = -\frac{3}{2}$ m

(b) Distance traveled $= \int_0^3 |3t-5|\,dt = \int_0^{5/3}(5-3t)\,dt + \int_{5/3}^3 (3t-5)\,dt$

$$= \left[5t - \tfrac{3}{2}t^2\right]_0^{5/3} + \left[\tfrac{3}{2}t^2 - 5t\right]_{5/3}^3 = \tfrac{25}{3} - \tfrac{3}{2}\cdot\tfrac{25}{9} + \tfrac{27}{2} - 15 - \left(\tfrac{3}{2}\cdot\tfrac{25}{9} - \tfrac{25}{3}\right) = \tfrac{41}{6}\text{ m}$$

70. (a) Displacement $= \int_2^4 (t^2-2t-3)\,dt = \left[\frac{1}{3}t^3 - t^2 - 3t\right]_2^4 = \left(\frac{64}{3} - 16 - 12\right) - \left(\frac{8}{3} - 4 - 6\right) = \frac{2}{3}$ m

(b) $v(t) = t^2 - 2t - 3 = (t+1)(t-3)$, so $v(t) < 0$ for $-1 < t < 3$, but on the interval $[2,4]$, $v(t) < 0$ for $2 \le t < 3$.

$$\begin{aligned}\text{Distance traveled} &= \int_2^4 \left|t^2-2t-3\right|\,dt = \int_2^3 -(t^2-2t-3)\,dt + \int_3^4 (t^2-2t-3)\,dt \\ &= \left[-\tfrac{1}{3}t^3 + t^2 + 3t\right]_2^3 + \left[\tfrac{1}{3}t^3 - t^2 - 3t\right]_3^4 \\ &= (-9+9+9) - \left(-\tfrac{8}{3}+4+6\right) + \left(\tfrac{64}{3} - 16 - 12\right) - (9-9-9) = 4\text{ m}\end{aligned}$$

71. (a) $v'(t) = a(t) = t+4 \;\Rightarrow\; v(t) = \frac{1}{2}t^2 + 4t + C \;\Rightarrow\; v(0) = C = 5 \;\Rightarrow\; v(t) = \frac{1}{2}t^2 + 4t + 5$ m/s

(b) Distance traveled $= \int_0^{10} |v(t)|\,dt = \int_0^{10} \left|\frac{1}{2}t^2 + 4t + 5\right|\,dt = \int_0^{10}\left(\frac{1}{2}t^2 + 4t + 5\right)dt = \left[\frac{1}{6}t^3 + 2t^2 + 5t\right]_0^{10}$

$$= \tfrac{500}{3} + 200 + 50 = 416\tfrac{2}{3}\text{ m}$$

72. (a) $v'(t) = a(t) = 2t+3 \;\Rightarrow\; v(t) = t^2 + 3t + C \;\Rightarrow\; v(0) = C = -4 \;\Rightarrow\; v(t) = t^2 + 3t - 4$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Distance traveled $= \int_0^3 \left|t^2+3t-4\right| dt = \int_0^3 |(t+4)(t-1)|\, dt = \int_0^1 \left(-t^2-3t+4\right) dt + \int_1^3 \left(t^2+3t-4\right) dt$

$$= \left[-\tfrac{1}{3}t^3 - \tfrac{3}{2}t^2 + 4t\right]_0^1 + \left[\tfrac{1}{3}t^3 + \tfrac{3}{2}t^2 - 4t\right]_1^3$$

$$= \left(-\tfrac{1}{3} - \tfrac{3}{2} + 4\right) + \left(9 + \tfrac{27}{2} - 12\right) - \left(\tfrac{1}{3} + \tfrac{3}{2} - 4\right) = \tfrac{89}{6} \text{ m}$$

73. Since $m'(x) = \rho(x)$, $m = \int_0^4 \rho(x)\, dx = \int_0^4 \left(9 + 2\sqrt{x}\right) dx = \left[9x + \tfrac{4}{3}x^{3/2}\right]_0^4 = 36 + \tfrac{32}{3} - 0 = \tfrac{140}{3} = 46\tfrac{2}{3}$ kg.

74. By the Net Change Theorem, the amount of water that flows from the tank during the first 10 minutes is

$\int_0^{10} r(t)\, dt = \int_0^{10} (200 - 4t)\, dt = \left[200t - 2t^2\right]_0^{10} = (2000 - 200) - 0 = 1800$ liters.

75. Let s be the position of the car. We know from Equation 2 that $s(100) - s(0) = \int_0^{100} v(t)\, dt$. We use the Midpoint Rule for $0 \le t \le 100$ with $n = 5$. Note that the length of each of the five time intervals is 20 seconds $= \frac{20}{3600}$ hour $= \frac{1}{180}$ hour. So the distance traveled is

$\int_0^{100} v(t)\, dt \approx \frac{1}{180}[v(10) + v(30) + v(50) + v(70) + v(90)] = \frac{1}{180}(38 + 58 + 51 + 53 + 47) = \frac{247}{180} \approx 1.37$ miles.

76. (a) By the Net Change Theorem, the total amount spewed into the atmosphere is $Q(6) - Q(0) = \int_0^6 r(t)\, dt = Q(6)$ since $Q(0) = 0$. The rate $r(t)$ is positive, so Q is an increasing function. Thus, an upper estimate for $Q(6)$ is R_6 and a lower estimate for $Q(6)$ is L_6. $\Delta t = \dfrac{b-a}{n} = \dfrac{6-0}{6} = 1$.

$R_6 = \sum_{i=1}^{6} r(t_i)\, \Delta t = 10 + 24 + 36 + 46 + 54 + 60 = 230$ tonnes.

$L_6 = \sum_{i=1}^{6} r(t_{i-1})\, \Delta t = R_6 + r(0) - r(6) = 230 + 2 - 60 = 172$ tonnes.

(b) $\Delta t = \dfrac{b-a}{n} = \dfrac{6-0}{3} = 2$. $Q(6) \approx M_3 = 2[r(1) + r(3) + r(5)] = 2(10 + 36 + 54) = 2(100) = 200$ tonnes.

77. From the Net Change Theorem, the increase in cost if the production level is raised from 2000 yards to 4000 yards is

$C(4000) - C(2000) = \int_{2000}^{4000} C'(x)\, dx$.

$$\int_{2000}^{4000} C'(x)\, dx = \int_{2000}^{4000} (3 - 0.01x + 0.000006x^2)\, dx = \left[3x - 0.005x^2 + 0.000002x^3\right]_{2000}^{4000}$$

$$= 60{,}000 - 2{,}000 = \$58{,}000$$

78. By the Net Change Theorem, the amount of water after four days is

$$25{,}000 + \int_0^4 r(t)\, dt \approx 25{,}000 + M_4 = 25{,}000 + \tfrac{4-0}{4}\left[r(0.5) + r(1.5) + r(2.5) + r(3.5)\right]$$

$$\approx 25{,}000 + [1500 + 1770 + 740 + (-690)] = 28{,}320 \text{ liters}$$

79. To use the Midpoint Rule, we'll use the midpoint of each of three 2-second intervals.

$v(6) - v(0) = \int_0^6 a(t)\, dt \approx [a(1) + a(3) + a(5)]\, \dfrac{6-0}{3} \approx (0.6 + 10 + 9.3)(2) = 39.8$ ft/s

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

80. Use the midpoint of each of four 2-day intervals. Let $t = 0$ correspond to July 18 and note that the inflow rate, $r(t)$, is in ft^3/s.

Amount of water $= \int_0^8 r(t)\,dt \approx [r(1) + r(3) + r(5) + r(7)]\,\dfrac{8-0}{4} \approx [6401 + 4249 + 3821 + 2628](2) = 34{,}198.$

Now multiply by the number of seconds in a day, $24 \cdot 60^2$, to get 2,954,707,200 ft^3.

81. Let $P(t)$ denote the bacteria population at time t (in hours). By the Net Change Theorem,

$$P(1) - P(0) = \int_0^1 P'(t)\,dt = \int_0^1 (1000 \cdot 2^t)\,dt = \left[1000\,\frac{2^t}{\ln 2}\right]_0^1 = \frac{1000}{\ln 2}\,(2^1 - 2^0) = \frac{1000}{\ln 2} \approx 1443.$$

Thus, the population after one hour is $4000 + 1443 = 5443$.

82. Let $M(t)$ denote the number of megabits transmitted at time t (in hours) [note that $D(t)$ is measured in megabits/second]. By the Net Change Theorem and the Midpoint Rule,

$$\begin{aligned} M(8) - M(0) &= \textstyle\int_0^8 3600D(t)\,dt \approx 3600 \cdot \frac{8-0}{4}[D(1) + D(3) + D(5) + D(7)] \\ &\approx 7200(0.32 + 0.50 + 0.56 + 0.83) = 7200(2.21) = 15{,}912 \text{ megabits} \end{aligned}$$

83. Power is the rate of change of energy with respect to time; that is, $P(t) = E'(t)$. By the Net Change Theorem and the Midpoint Rule, the electric energy used on that day is

$$\begin{aligned} E(24) - E(0) &= \int_0^{24} P(t)\,dt \approx \frac{24-0}{12}[P(1) + P(3) + P(5) + \cdots + P(21) + P(23)] \\ &\approx 2(11 + 10.5 + 11 + 14 + 15.1 + 15.5 + 15.1 + 15 + 15 + 16.1 + 15 + 13) \\ &= 2(166.3) = 332.6 \text{ gigawatt-hours} \end{aligned}$$

84. (a) From Exercise 4.1.78(a), $v(t) = 0.00146t^3 - 0.11553t^2 + 24.98169t - 21.26872$.

(b) $h(125) - h(0) = \int_0^{125} v(t)\,dt = \left[0.000365t^4 - 0.03851t^3 + 12.490845t^2 - 21.26872t\right]_0^{125} \approx 206{,}407$ ft

5.5 The Substitution Rule

1. Let $u = 2x$. Then $du = 2\,dx$ and $dx = \frac{1}{2}\,du$, so $\int \cos 2x\,dx = \int \cos u\,\left(\frac{1}{2}\,du\right) = \frac{1}{2}\sin u + C = \frac{1}{2}\sin 2x + C$.

2. Let $u = -x^2$. Then $du = -2x\,dx$ and $x\,dx = -\frac{1}{2}\,du$, so $\int xe^{-x^2}\,dx = \int e^u\,\left(-\frac{1}{2}\,du\right) = -\frac{1}{2}e^u + C = -\frac{1}{2}e^{-x^2} + C$.

3. Let $u = x^3 + 1$. Then $du = 3x^2\,dx$ and $x^2\,dx = \frac{1}{3}\,du$, so

$$\int x^2\sqrt{x^3+1}\,dx = \int \sqrt{u}\,\left(\tfrac{1}{3}\,du\right) = \frac{1}{3}\frac{u^{3/2}}{3/2} + C = \frac{1}{3} \cdot \frac{2}{3}u^{3/2} + C = \tfrac{2}{9}(x^3+1)^{3/2} + C.$$

4. Let $u = \sin\theta$. Then $du = \cos\theta\,d\theta$, so $\int \sin^2\theta\,\cos\theta\,d\theta = \int u^2\,du = \frac{1}{3}u^3 + C = \frac{1}{3}\sin^3\theta + C$.

5. Let $u = x^4 - 5$. Then $du = 4x^3\,dx$ and $x^3\,dx = \frac{1}{4}\,du$, so

$$\int \frac{x^3}{x^4-5}\,dx = \int \frac{1}{u}\left(\frac{1}{4}\,du\right) = \frac{1}{4}\ln|u| + C = \frac{1}{4}\ln\left|x^4 - 5\right| + C.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. Let $u = 1 + \dfrac{1}{x}$. Then $du = -\dfrac{1}{x^2}\,dx$ and $\dfrac{1}{x^2}\,dx = -du$, so

$$\int \frac{1}{x^2}\sqrt{1+\frac{1}{x}}\,dx = \int \sqrt{u}\,(-du) = -\frac{2}{3}u^{3/2} + C = -\frac{2}{3}\left(1+\frac{1}{x}\right)^{3/2} + C.$$

7. Let $u = \sqrt{t}$. Then $du = \dfrac{1}{2\sqrt{t}}\,dt$ and $\dfrac{1}{\sqrt{t}}\,dt = 2\,du$, so $\displaystyle\int \frac{\cos\sqrt{t}}{\sqrt{t}}\,dt = \int \cos u\ (2\,du) = 2\sin u + C = 2\sin\sqrt{t} + C$.

8. Let $u = z - 1$. Then $z = u + 1$ and $du = dz$, so

$$\begin{aligned}\textstyle\int z\sqrt{z-1}\,dz = \int (u+1)\sqrt{u}\,du = \int (u^{3/2} + u^{1/2})\,du &= \tfrac{2}{5}u^{5/2} + \tfrac{2}{3}u^{3/2} + C \\ &= \tfrac{2}{5}(z-1)^{5/2} + \tfrac{2}{3}(z-1)^{3/2} + C\end{aligned}$$

9. Let $u = 1 - x^2$. Then $du = -2x\,dx$ and $x\,dx = -\frac{1}{2}\,du$, so

$\int x\sqrt{1-x^2}\,dx = \int \sqrt{u}\,\left(-\frac{1}{2}\,du\right) = -\frac{1}{2}\cdot\frac{2}{3}u^{3/2} + C = -\frac{1}{3}(1-x^2)^{3/2} + C.$

10. Let $u = 5 - 3x$. Then $du = -3\,dx$ and $dx = -\frac{1}{3}\,du$, so

$\int (5-3x)^{10}\,dx = \int u^{10}\,\left(-\frac{1}{3}\,du\right) = -\frac{1}{3}\cdot\frac{1}{11}u^{11} + C = -\frac{1}{33}(5-3x)^{11} + C.$

11. Let $u = -t^4$. Then $du = -4t^3\,dt$ and $t^3\,dt = -\frac{1}{4}\,du$, so $\int t^3 e^{-t^4}\,dt = \int e^u\,\left(-\frac{1}{4}\,du\right) = -\frac{1}{4}e^u + C = -\frac{1}{4}e^{-t^4} + C.$

12. Let $u = 1 + \cos t$. Then $du = -\sin t\,dt$ and $\sin t\,dt = -du$, so

$\int \sin t\,\sqrt{1+\cos t}\,dt = \int \sqrt{u}\,(-du) = -\frac{2}{3}u^{3/2} + C = -\frac{2}{3}(1+\cos t)^{3/2} + C.$

13. Let $u = \dfrac{\pi}{3}t$. Then $du = \dfrac{\pi}{3}\,dt$ and $dt = \dfrac{3}{\pi}\,du$, so

$$\int \sin\left(\frac{\pi t}{3}\right)dt = \int \sin u\,\left(\frac{3}{\pi}\,du\right) = \frac{3}{\pi}\cdot(-\cos u) + C = -\frac{3}{\pi}\cos\left(\frac{\pi}{3}t\right) + C.$$

14. Let $u = 2\theta$. Then $du = 2\,d\theta$ and $d\theta = \frac{1}{2}\,du$, so $\int \sec^2 2\theta\,d\theta = \int \sec^2 u\,\left(\frac{1}{2}\,du\right) = \frac{1}{2}\tan u + C = \frac{1}{2}\tan 2\theta + C.$

15. Let $u = 4x + 7$. Then $du = 4\,dx$ and $dx = \dfrac{1}{4}\,du$, so $\displaystyle\int \frac{dx}{4x+7} = \int \frac{1}{u}\left(\frac{1}{4}\,du\right) = \frac{1}{4}\ln|u| + C = \frac{1}{4}\ln|4x+7| + C.$

16. Let $u = 4 - y^3$. Then $du = -3y^2\,dy$ and $y^2\,dy = -\frac{1}{3}\,du$, so

$\int y^2(4-y^3)^{2/3}\,dy = \int u^{2/3}\,\left(-\frac{1}{3}\,du\right) = -\frac{1}{3}\cdot\frac{3}{5}u^{5/3} + C = -\frac{1}{5}(4-y^3)^{5/3} + C.$

17. Let $u = 1 + \sin\theta$. Then $du = \cos\theta\,d\theta$, so

$\displaystyle\int \frac{\cos\theta}{1+\sin\theta}\,d\theta = \int \frac{1}{u}\,du = \ln|u| + C = \ln|1+\sin\theta| + C = \ln(1+\sin\theta) + C$ [since $1 + \sin\theta$ is nonnegative].

18. Let $u = z^3 + 1$. Then $du = 3z^2\,dz$ and $\frac{1}{3}\,du = z^2\,dz$, so

$$\int \frac{z^2}{z^3+1}\,dz = \int \frac{1}{u}\left(\frac{1}{3}\,du\right) = \frac{1}{3}\ln|u| + C = \frac{1}{3}\ln|z^3+1| + C.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. Let $u = \cos\theta$. Then $du = -\sin\theta\, d\theta$ and $\sin\theta\, d\theta = -du$, so

$\int \cos^3\theta\, \sin\theta\, d\theta = \int u^3(-du) = -\frac{1}{4}u^4 + C = -\frac{1}{4}\cos^4\theta + C.$

20. Let $u = -5r$. Then $du = -5\, dr$ and $dr = -\frac{1}{5}\, du$, so $\int e^{-5r}\, dr = \int e^u\left(-\frac{1}{5}\, du\right) = -\frac{1}{5}e^u + C = -\frac{1}{5}e^{-5r} + C.$

21. Let $x = 1 - e^u$. Then $dx = -e^u\, du$ and $e^u\, du = -dx$, so

$$\int \frac{e^u}{(1-e^u)^2}\, du = \int \frac{1}{x^2}(-dx) = -\int x^{-2}\, dx = -(-x^{-1}) + C = \frac{1}{x} + C = \frac{1}{1-e^u} + C.$$

22. Let $u = 1/x$. Then $du = -\dfrac{1}{x^2}\, dx$ and $\dfrac{1}{x^2}\, dx = -du$, so $\displaystyle\int \frac{\sin(1/x)}{x^2}\, dx = \int \sin u\,(-du) = \cos u + C = \cos(1/x) + C.$

23. Let $u = 3ax + bx^3$. Then $du = (3a + 3bx^2)\, dx = 3(a + bx^2)\, dx$, so

$$\int \frac{a + bx^2}{\sqrt{3ax + bx^3}}\, dx = \int \frac{\frac{1}{3}\, du}{u^{1/2}} = \frac{1}{3}\int u^{-1/2}\, du = \tfrac{1}{3}\cdot 2u^{1/2} + C = \tfrac{2}{3}\sqrt{3ax + bx^3} + C.$$

24. Let $u = 3t^2 + 6t - 5$. Then $du = (6t + 6)\, dt = 6(t + 1)\, dt$ and $(t + 1)\, dt = \frac{1}{6}\, du$, so

$$\int \frac{t+1}{3t^2 + 6t - 5}\, dt = \int \frac{1}{u}\left(\frac{1}{6}\, du\right) = \frac{1}{6}\ln|u| + C = \frac{1}{6}\ln\left|3t^2 + 6t - 5\right| + C.$$

25. Let $u = \ln x$. Then $du = \dfrac{dx}{x}$, so $\displaystyle\int \frac{(\ln x)^2}{x}\, dx = \int u^2\, du = \tfrac{1}{3}u^3 + C = \tfrac{1}{3}(\ln x)^3 + C.$

26. Let $u = \cos x$. Then $du = -\sin x\, dx$ and $-du = \sin x\, dx$, so

$\int \sin x\, \sin(\cos x)\, dx = \int \sin u\,(-du) = (-\cos u)(-1) + C = \cos(\cos x) + C.$

27. Let $u = \tan\theta$. Then $du = \sec^2\theta\, d\theta$, so $\int \sec^2\theta\, \tan^3\theta\, d\theta = \int u^3\, du = \frac{1}{4}u^4 + C = \frac{1}{4}\tan^4\theta + C.$

28. Let $u = x + 2$. Then $du = dx$ and $x = u - 2$, so

$\int x\sqrt{x+2}\, dx = \int (u-2)\sqrt{u}\, du = \int (u^{3/2} - 2u^{1/2})\, du = \frac{2}{5}u^{5/2} - 2\cdot\frac{2}{3}u^{3/2} + C = \frac{2}{5}(x+2)^{5/2} - \frac{4}{3}(x+2)^{3/2} + C.$

29. Let $u = x^2 + \dfrac{2}{x}$. Then $du = \left(2x - \dfrac{2}{x^2}\right) dx = 2\left(x - \dfrac{1}{x^2}\right) dx$ and $\left(x - \dfrac{1}{x^2}\right) dx = \dfrac{1}{2}\, du$, so

$$\int \left(x - \frac{1}{x^2}\right)\left(x^2 + \frac{2}{x}\right)^5 dx = \int u^5\left(\frac{1}{2}\, du\right) = \frac{1}{2}\cdot\frac{1}{6}u^6 + C = \frac{1}{12}\left(x^2 + \frac{2}{x}\right)^6 + C.$$

30. Let $u = ax + b$. Then $du = a\, dx$ and $dx = (1/a)\, du$, so

$$\int \frac{dx}{ax+b} = \int \frac{(1/a)\, du}{u} = \frac{1}{a}\int \frac{1}{u}\, du = \frac{1}{a}\ln|u| + C = \frac{1}{a}\ln|ax+b| + C.$$

31. Let $u = 2 + 3e^r$. Then $du = 3e^r\, dr$ and $e^r\, dr = \frac{1}{3}\, du$, so

$\int e^r(2 + 3e^r)^{3/2}\, dr = \int u^{3/2}\left(\frac{1}{3}\, du\right) = \frac{1}{3}\cdot\frac{2}{5}u^{5/2} + C = \frac{2}{15}(2 + 3e^r)^{5/2} + C.$

32. Let $u = \arcsin x$. Then $du = \dfrac{1}{\sqrt{1-x^2}}\, dx$, so $\displaystyle\int \frac{e^{\arcsin x}}{\sqrt{1-x^2}}\, dx = \int e^u\, du = e^u + C = e^{\arcsin x} + C.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. Let $u = \tan\theta$. Then $du = \sec^2\theta\, d\theta$, so $\displaystyle\int \frac{\sec^2\theta}{\tan\theta}\, d\theta = \int \frac{1}{u}\, du = \ln|u| + C = \ln|\tan\theta| + C$.

34. Let $u = \tan x$. Then $du = \sec^2 x\, dx$, so

$$\int \frac{\sec^2 x}{\tan^2 x}\, dx = \int \frac{1}{u^2}\, du = \int u^{-2}\, du = -1u^{-1} + C = -\frac{1}{\tan x} + C = -\cot x + C.$$

Or: $\displaystyle\int \frac{\sec^2 x}{\tan^2 x}\, dx = \int \left(\frac{1}{\cos^2 x} \cdot \frac{\cos^2 x}{\sin^2 x}\right) dx = \int \csc^2 x\, dx = -\cot x + C$

35. Let $u = \arctan x$. Then $du = \dfrac{1}{x^2+1}\, dx$, so $\displaystyle\int \frac{(\arctan x)^2}{x^2+1}\, dx = \int u^2\, du = \tfrac{1}{3}u^3 + C = \tfrac{1}{3}(\arctan x)^3 + C$.

36. Let $u = \arctan x$. Then $du = \dfrac{1}{1+x^2}\, dx = \dfrac{1}{x^2+1}\, dx$, so

$$\int \frac{1}{(x^2+1)\arctan x}\, dx = \int \frac{1}{u}\, du = \ln|u| + C = \ln|\arctan x| + C.$$

37. Let $u = 5^t$. Then $du = 5^t \ln 5\, dt$ and $5^t\, dt = \dfrac{1}{\ln 5}\, du$, so

$$\int 5^t \sin(5^t)\, dt = \int \sin u \left(\frac{1}{\ln 5}\, du\right) = -\frac{1}{\ln 5}\cos u + C = -\frac{1}{\ln 5}\cos(5^t) + C.$$

38. Let $u = 1 + \sin^2\theta$. Then $du = 2\sin\theta\cos\theta\, d\theta$ and $\sin\theta\cos\theta\, d\theta = \frac{1}{2}\, du$, so

$$\int \frac{\sin\theta\cos\theta}{1+\sin^2\theta}\, d\theta = \int \frac{1}{u}\left(\frac{1}{2}\, du\right) = \frac{1}{2}\ln|u| + C = \frac{1}{2}\ln\left|1+\sin^2\theta\right| + C$$

$$= \tfrac{1}{2}\ln(1+\sin^2\theta) + C \qquad \text{[since } 1+\sin^2\theta \text{ is positive]}$$

39. Let $u = 1 + 5t$. Then $du = 5\, dt$ and $dt = \frac{1}{5}\, du$, so

$\int \cos(1+5t)\, dt = \int \cos u\left(\frac{1}{5}\, du\right) = \frac{1}{5}\sin u + C = \frac{1}{5}\sin(1+5t) + C$.

40. Let $u = \dfrac{\pi}{x}$. Then $du = -\dfrac{\pi}{x^2}\, dx$ and $\dfrac{1}{x^2}\, dx = -\dfrac{1}{\pi}\, du$, so

$$\int \frac{\cos(\pi/x)}{x^2}\, dx = \int \cos u\left(-\frac{1}{\pi}\, du\right) = -\frac{1}{\pi}\sin u + C = -\frac{1}{\pi}\sin\frac{\pi}{x} + C$$

41. Let $u = \cot x$. Then $du = -\csc^2 x\, dx$ and $\csc^2 x\, dx = -du$, so

$$\int \sqrt{\cot x}\csc^2 x\, dx = \int \sqrt{u}\,(-du) = -\frac{u^{3/2}}{3/2} + C = -\tfrac{2}{3}(\cot x)^{3/2} + C.$$

42. Let $u = 2^t + 3$. Then $du = 2^t \ln 2\, dt$ and $2^t\, dt = \dfrac{1}{\ln 2}\, du$, so

$$\int \frac{2^t}{2^t+3}\, dt = \int \frac{1}{u}\left(\frac{1}{\ln 2}\, du\right) = \frac{1}{\ln 2}\ln|u| + C = \frac{1}{\ln 2}\ln(2^t+3) + C.$$

43. Let $u = \sinh x$. Then $du = \cosh x\, dx$, so $\int \sinh^2 x\, \cosh x\, dx = \int u^2\, du = \frac{1}{3}u^3 + C = \frac{1}{3}\sinh^3 x + C$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. Let $u = 1 + \tan t$. Then $du = \sec^2 t\,dt$, so

$$\int \frac{dt}{\cos^2 t\sqrt{1+\tan t}} = \int \frac{\sec^2 t\,dt}{\sqrt{1+\tan t}} = \int \frac{du}{\sqrt{u}} = \int u^{-1/2}\,du = \frac{u^{1/2}}{1/2} + C = 2\sqrt{1+\tan t} + C.$$

45. $\displaystyle\int \frac{\sin 2x}{1+\cos^2 x}\,dx = 2\int \frac{\sin x\cos x}{1+\cos^2 x}\,dx = 2I$. Let $u = \cos x$. Then $du = -\sin x\,dx$, so

$$2I = -2\int \frac{u\,du}{1+u^2} = -2\cdot\tfrac{1}{2}\ln(1+u^2) + C = -\ln(1+u^2) + C = -\ln(1+\cos^2 x) + C.$$

Or: Let $u = 1 + \cos^2 x$.

46. Let $u = \cos x$. Then $du = -\sin x\,dx$ and $\sin x\,dx = -du$, so

$$\int \frac{\sin x}{1+\cos^2 x}\,dx = \int \frac{-du}{1+u^2} = -\tan^{-1} u + C = -\tan^{-1}(\cos x) + C.$$

47. $\displaystyle\int \cot x\,dx = \int \frac{\cos x}{\sin x}\,dx$. Let $u = \sin x$. Then $du = \cos x\,dx$, so $\displaystyle\int \cot x\,dx = \int \frac{1}{u}\,du = \ln|u| + C = \ln|\sin x| + C$.

48. Let $u = \ln t$. Then $du = \dfrac{1}{t}\,dt$, so $\displaystyle\int \frac{\cos(\ln t)}{t}\,dt = \int \cos u\,du = \sin u + C = \sin(\ln t) + C$.

49. Let $u = \sin^{-1} x$. Then $du = \dfrac{1}{\sqrt{1-x^2}}\,dx$, so $\displaystyle\int \frac{dx}{\sqrt{1-x^2}\sin^{-1} x} = \int \frac{1}{u}\,du = \ln|u| + C = \ln\left|\sin^{-1} x\right| + C$.

50. Let $u = x^2$. Then $du = 2x\,dx$, so $\displaystyle\int \frac{x}{1+x^4}\,dx = \int \frac{\frac{1}{2}\,du}{1+u^2} = \tfrac{1}{2}\tan^{-1} u + C = \tfrac{1}{2}\tan^{-1}(x^2) + C$.

51. Let $u = 1 + x^2$. Then $du = 2x\,dx$, so

$$\begin{aligned}\int \frac{1+x}{1+x^2}\,dx &= \int \frac{1}{1+x^2}\,dx + \int \frac{x}{1+x^2}\,dx = \tan^{-1} x + \int \frac{\frac{1}{2}\,du}{u} = \tan^{-1} x + \tfrac{1}{2}\ln|u| + C \\ &= \tan^{-1} x + \tfrac{1}{2}\ln\left|1+x^2\right| + C = \tan^{-1} x + \tfrac{1}{2}\ln\left(1+x^2\right) + C \quad [\text{since } 1+x^2 > 0].\end{aligned}$$

52. Let $u = 2 + x$. Then $du = dx$, $x = u - 2$, and $x^2 = (u-2)^2$, so

$$\begin{aligned}\textstyle\int x^2\sqrt{2+x}\,dx &= \textstyle\int (u-2)^2\sqrt{u}\,du = \int (u^2 - 4u + 4)u^{1/2}\,du = \int (u^{5/2} - 4u^{3/2} + 4u^{1/2})\,du \\ &= \tfrac{2}{7}u^{7/2} - \tfrac{8}{5}u^{5/2} + \tfrac{8}{3}u^{3/2} + C = \tfrac{2}{7}(2+x)^{7/2} - \tfrac{8}{5}(2+x)^{5/2} + \tfrac{8}{3}(2+x)^{3/2} + C\end{aligned}$$

53. Let $u = 2x + 5$. Then $du = 2\,dx$ and $x = \frac{1}{2}(u-5)$, so

$$\begin{aligned}\textstyle\int x(2x+5)^8\,dx &= \textstyle\int \frac{1}{2}(u-5)u^8\left(\frac{1}{2}\,du\right) = \frac{1}{4}\int (u^9 - 5u^8)\,du \\ &= \tfrac{1}{4}\left(\tfrac{1}{10}u^{10} - \tfrac{5}{9}u^9\right) + C = \tfrac{1}{40}(2x+5)^{10} - \tfrac{5}{36}(2x+5)^9 + C\end{aligned}$$

54. Let $u = x^2 + 1$ [so $x^2 = u - 1$]. Then $du = 2x\,dx$ and $x\,dx = \frac{1}{2}\,du$, so

$$\begin{aligned}\textstyle\int x^3\sqrt{x^2+1}\,dx &= \textstyle\int x^2\sqrt{x^2+1}\,x\,dx = \int (u-1)\sqrt{u}\left(\frac{1}{2}\,du\right) = \frac{1}{2}\int (u^{3/2} - u^{1/2})\,du \\ &= \tfrac{1}{2}\left(\tfrac{2}{5}u^{5/2} - \tfrac{2}{3}u^{3/2}\right) + C = \tfrac{1}{5}(x^2+1)^{5/2} - \tfrac{1}{3}(x^2+1)^{3/2} + C\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Or: Let $u = \sqrt{x^2+1}$. Then $u^2 = x^2+1 \quad \Rightarrow \quad 2u\,du = 2x\,dx \quad \Rightarrow \quad u\,du = x\,dx$, so

$$\int x^3\sqrt{x^2+1}\,dx = \int x^2\sqrt{x^2+1}\,x\,dx = \int (u^2-1)\,u\cdot u\,du = \int (u^4-u^2)\,du$$
$$= \tfrac{1}{5}u^5 - \tfrac{1}{3}u^3 + C = \tfrac{1}{5}(x^2+1)^{5/2} - \tfrac{1}{3}(x^2+1)^{3/2} + C$$

Note: This answer can be written as $\frac{1}{15}\sqrt{x^2+1}(3x^4+x^2-2)+C$.

55. $f(x) = x(x^2-1)^3$. $\quad u = x^2-1 \quad \Rightarrow \quad du = 2x\,dx$, so

$$\int x(x^2-1)^3\,dx = \int u^3\left(\tfrac{1}{2}\,du\right) = \tfrac{1}{8}u^4 + C = \tfrac{1}{8}(x^2-1)^4 + C$$

Where f is positive (negative), F is increasing (decreasing). Where f changes from negative to positive (positive to negative), F has a local minimum (maximum).

56. $f(\theta) = \tan^2\theta\,\sec^2\theta$. $u = \tan\theta \quad \Rightarrow \quad du = \sec^2\theta\,d\theta$, so

$$\int \tan^2\theta\,\sec^2\theta\,d\theta = \int u^2\,du = \tfrac{1}{3}u^3 + C = \tfrac{1}{3}\tan^3\theta + C$$

Note that f is positive and F is increasing. At $x=0$, $f=0$ and F has a horizontal tangent.

57. $f(x) = e^{\cos x}\sin x$. $u = \cos x \quad \Rightarrow \quad du = -\sin x\,dx$, so

$$\int e^{\cos x}\sin x\,dx = \int e^u\,(-du) = -e^u + C = -e^{\cos x} + C$$

Note that at $x=\pi$, f changes from positive to negative and F has a local maximum. Also, both f and F are periodic with period 2π, so at $x=0$ and at $x=2\pi$, f changes from negative to positive and F has a local minimum.

58. $f(x) = \sin x\cos^4 x$. $u = \cos x \quad \Rightarrow \quad du = -\sin x\,dx$, so

$$\int \sin x\cos^4 x\,dx = \int u^4\,(-du) = -\tfrac{1}{5}u^5 + C = -\tfrac{1}{5}\cos^5 x + C$$

Note that at $x=\pi$, f changes from positive to negative and F has a local maximum. Also, both f and F are periodic with period 2π, so at $x=0$ and at $x=2\pi$, f changes from negative to positive and F has a local minimum.

59. Let $u = \frac{\pi}{2}t$, so $du = \frac{\pi}{2}\,dt$. When $t=0$, $u=0$; when $t=1$, $u=\frac{\pi}{2}$. Thus,

$$\int_0^1 \cos(\pi t/2)\,dt = \int_0^{\pi/2}\cos u\,\left(\tfrac{2}{\pi}\,du\right) = \tfrac{2}{\pi}\left[\sin u\right]_0^{\pi/2} = \tfrac{2}{\pi}\left(\sin\tfrac{\pi}{2} - \sin 0\right) = \tfrac{2}{\pi}(1-0) = \tfrac{2}{\pi}$$

60. Let $u = 3t-1$, so $du = 3\,dt$. When $t=0$, $u=-1$; when $t=1$, $u=2$. Thus,

$$\int_0^1 (3t-1)^{50}\,dt = \int_{-1}^2 u^{50}\,\left(\tfrac{1}{3}\,du\right) = \tfrac{1}{3}\left[\tfrac{1}{51}u^{51}\right]_{-1}^2 = \tfrac{1}{153}\left[2^{51} - (-1)^{51}\right] = \tfrac{1}{153}(2^{51}+1)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. Let $u = 1 + 7x$, so $du = 7\,dx$. When $x = 0$, $u = 1$; when $x = 1$, $u = 8$. Thus,

$$\int_0^1 \sqrt[3]{1+7x}\,dx = \int_1^8 u^{1/3}\left(\tfrac{1}{7}\,du\right) = \tfrac{1}{7}\left[\tfrac{3}{4}u^{4/3}\right]_1^8 = \tfrac{3}{28}\left(8^{4/3} - 1^{4/3}\right) = \tfrac{3}{28}(16-1) = \tfrac{45}{28}$$

62. Let $u = \frac{1}{2}t$, so $du = \frac{1}{2}\,dt$. When $t = \frac{\pi}{3}$, $u = \frac{\pi}{6}$; when $t = \frac{2\pi}{3}$, $u = \frac{\pi}{3}$. Thus,

$$\int_{\pi/3}^{2\pi/3} \csc^2\left(\tfrac{1}{2}t\right)dt = \int_{\pi/6}^{\pi/3} \csc^2 u\,(2\,du) = 2\left[-\cot u\right]_{\pi/6}^{\pi/3} = -2\left(\cot\frac{\pi}{3} - \cot\frac{\pi}{6}\right)$$

$$= -2\left(\frac{1}{\sqrt{3}} - \sqrt{3}\right) = -2\left(\tfrac{1}{3}\sqrt{3} - \sqrt{3}\right) = \tfrac{4}{3}\sqrt{3}$$

63. Let $u = \cos t$, so $du = -\sin t\,dt$. When $t = 0$, $u = 1$; when $t = \frac{\pi}{6}$, $u = \sqrt{3}/2$. Thus,

$$\int_0^{\pi/6} \frac{\sin t}{\cos^2 t}\,dt = \int_1^{\sqrt{3}/2} \frac{1}{u^2}\,(-du) = \left[\frac{1}{u}\right]_1^{\sqrt{3}/2} = \frac{2}{\sqrt{3}} - 1.$$

64. Let $u = 2 + \sqrt{x}$. Then $du = \dfrac{1}{2\sqrt{x}}\,dx$ and $\dfrac{1}{\sqrt{x}}\,dx = 2\,du$. When $x = 1$, $u = 3$; when $x = 4$, $u = 4$. Thus,

$$\int_1^4 \frac{\sqrt{2+\sqrt{x}}}{\sqrt{x}}\,dx = \int_3^4 \sqrt{u}\,(2\,du) = \left[\frac{4}{3}u^{3/2}\right]_3^4 = \frac{4}{3}\left(4^{3/2} - 3^{3/2}\right) = \frac{4}{3}\left(8 - 3\sqrt{3}\right) \text{ or } \frac{32}{3} - 4\sqrt{3}.$$

65. Let $u = 1/x$, so $du = -1/x^2\,dx$. When $x = 1$, $u = 1$; when $x = 2$, $u = \frac{1}{2}$. Thus,

$$\int_1^2 \frac{e^{1/x}}{x^2}\,dx = \int_1^{1/2} e^u\,(-du) = -\left[e^u\right]_1^{1/2} = -\left(e^{1/2} - e\right) = e - \sqrt{e}.$$

66. Let $u = e^x$. Then $du = e^x\,dx$. When $x = 0$, $u = 1$; when $x = 1$, $u = e$. Thus,

$$\int_0^1 \frac{e^x}{1+e^{2x}}\,dx = \int_1^e \frac{1}{1+u^2}\,du = \Big[\arctan u\Big]_1^e = \arctan e - \arctan 1 = \arctan e - \frac{\pi}{4}.$$

67. $\int_{-\pi/4}^{\pi/4}(x^3 + x^4\tan x)\,dx = 0$ by Theorem 7(b), since $f(x) = x^3 + x^4\tan x$ is an odd function.

68. Let $u = \sin x$, so $du = \cos x\,dx$. When $x = 0$, $u = 0$; when $x = \frac{\pi}{2}$, $u = 1$. Thus,

$\int_0^{\pi/2}\cos x\,\sin(\sin x)\,dx = \int_0^1 \sin u\,du = \big[-\cos u\big]_0^1 = -(\cos 1 - 1) = 1 - \cos 1.$

69. Let $u = 1 + 2x$, so $du = 2\,dx$. When $x = 0$, $u = 1$; when $x = 13$, $u = 27$. Thus,

$$\int_0^{13} \frac{dx}{\sqrt[3]{(1+2x)^2}} = \int_1^{27} u^{-2/3}\left(\tfrac{1}{2}\,du\right) = \left[\tfrac{1}{2}\cdot 3u^{1/3}\right]_1^{27} = \tfrac{3}{2}(3-1) = 3.$$

70. Assume $a > 0$. Let $u = a^2 - x^2$, so $du = -2x\,dx$. When $x = 0$, $u = a^2$; when $x = a$, $u = 0$. Thus,

$\int_0^a x\sqrt{a^2 - x^2}\,dx = \int_{a^2}^0 u^{1/2}\left(-\tfrac{1}{2}\,du\right) = \tfrac{1}{2}\int_0^{a^2} u^{1/2}\,du = \tfrac{1}{2}\cdot\left[\tfrac{2}{3}u^{3/2}\right]_0^{a^2} = \tfrac{1}{3}a^3.$

71. Let $u = x^2 + a^2$, so $du = 2x\,dx$ and $x\,dx = \frac{1}{2}\,du$. When $x = 0$, $u = a^2$; when $x = a$, $u = 2a^2$. Thus,

$$\int_0^a x\sqrt{x^2+a^2}\,dx = \int_{a^2}^{2a^2} u^{1/2}\left(\tfrac{1}{2}\,du\right) = \tfrac{1}{2}\left[\tfrac{2}{3}u^{3/2}\right]_{a^2}^{2a^2} = \left[\tfrac{1}{3}u^{3/2}\right]_{a^2}^{2a^2} = \tfrac{1}{3}\left[(2a^2)^{3/2} - (a^2)^{3/2}\right] = \tfrac{1}{3}\left(2\sqrt{2} - 1\right)a^3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

72. $\int_{-\pi/3}^{\pi/3} x^4 \sin x\,dx = 0$ by Theorem 7(b), since $f(x) = x^4 \sin x$ is an odd function.

73. Let $u = x - 1$, so $u + 1 = x$ and $du = dx$. When $x = 1$, $u = 0$; when $x = 2$, $u = 1$. Thus,

$$\int_1^2 x\sqrt{x-1}\,dx = \int_0^1 (u+1)\sqrt{u}\,du = \int_0^1 (u^{3/2} + u^{1/2})\,du = \left[\tfrac{2}{5}u^{5/2} + \tfrac{2}{3}u^{3/2}\right]_0^1 = \tfrac{2}{5} + \tfrac{2}{3} = \tfrac{16}{15}.$$

74. Let $u = 1 + 2x$, so $x = \frac{1}{2}(u-1)$ and $du = 2\,dx$. When $x = 0$, $u = 1$; when $x = 4$, $u = 9$. Thus,

$$\int_0^4 \frac{x\,dx}{\sqrt{1+2x}} = \int_1^9 \frac{\frac{1}{2}(u-1)}{\sqrt{u}}\frac{du}{2} = \tfrac{1}{4}\int_1^9 (u^{1/2} - u^{-1/2})\,du = \tfrac{1}{4}\left[\tfrac{2}{3}u^{3/2} - 2u^{1/2}\right]_1^9 = \tfrac{1}{4}\cdot\tfrac{2}{3}\left[u^{3/2} - 3u^{1/2}\right]_1^9$$
$$= \tfrac{1}{6}[(27-9) - (1-3)] = \tfrac{20}{6} = \tfrac{10}{3}$$

75. Let $u = \ln x$, so $du = \dfrac{dx}{x}$. When $x = e$, $u = 1$; when $x = e^4$; $u = 4$. Thus,

$$\int_e^{e^4} \frac{dx}{x\sqrt{\ln x}} = \int_1^4 u^{-1/2}\,du = 2\left[u^{1/2}\right]_1^4 = 2(2-1) = 2.$$

76. Let $u = (x-1)^2$, so $du = 2(x-1)\,dx$. When $x = 0$, $u = 1$; when $x = 2$, $u = 1$. Thus,

$$\int_0^2 (x-1)e^{(x-1)^2}\,dx = \int_1^1 e^u\left(\tfrac{1}{2}\,du\right) = 0 \text{ since the limits are equal.}$$

77. Let $u = e^z + z$, so $du = (e^z + 1)\,dz$. When $z = 0$, $u = 1$; when $z = 1$, $u = e + 1$. Thus,

$$\int_0^1 \frac{e^z+1}{e^z+z}\,dz = \int_1^{e+1} \frac{1}{u}\,du = \Big[\ln|u|\Big]_1^{e+1} = \ln|e+1| - \ln|1| = \ln(e+1).$$

78. Let $u = \sqrt{x}$. Then $u^2 = x$, $2u\,du = dx$, $du = \dfrac{1}{2\sqrt{x}}\,dx$, and $\dfrac{1}{\sqrt{x}}\,dx = 2\,du$. When $x = 1$, $u = 1$; when $x = 4$, $u = 2$.

Thus,

$$\int_1^4 \frac{1}{(x+1)\sqrt{x}}\,dx = \int_1^2 \frac{1}{u^2+1}\,(2\,du) = 2\Big[\arctan u\Big]_1^2 = 2(\arctan 2 - \arctan 1)$$
$$= 2\left(\arctan 2 - \frac{\pi}{4}\right) = 2\arctan 2 - \frac{\pi}{2}$$

79. Let $u = 1 + \sqrt{x}$, so $du = \dfrac{1}{2\sqrt{x}}\,dx \;\Rightarrow\; 2\sqrt{x}\,du = dx \;\Rightarrow\; 2(u-1)\,du = dx$. When $x = 0$, $u = 1$; when $x = 1$, $u = 2$. Thus,

$$\int_0^1 \frac{dx}{(1+\sqrt{x})^4} = \int_1^2 \frac{1}{u^4}\cdot[2(u-1)\,du] = 2\int_1^2 \left(\frac{1}{u^3} - \frac{1}{u^4}\right)du = 2\left[-\frac{1}{2u^2} + \frac{1}{3u^3}\right]_1^2$$
$$= 2\left[\left(-\tfrac{1}{8} + \tfrac{1}{24}\right) - \left(-\tfrac{1}{2} + \tfrac{1}{3}\right)\right] = 2\left(\tfrac{1}{12}\right) = \tfrac{1}{6}$$

80. Let $u = 1 + x^{3/4}$. Then $x^{3/4} = u - 1$, $du = \frac{3}{4}x^{-1/4}\,dx$, and $x^{-1/4}\,dx = \frac{4}{3}\,du$. When $x = 1$, $u = 2$; when $x = 16$, $u = 9$. Thus,

$$\int_1^{16} \frac{x^{1/2}}{1+x^{3/4}}\,dx = \int_1^{16} \frac{x^{3/4}\cdot x^{-1/4}}{1+x^{3/4}}\,dx = \int_2^9 \frac{u-1}{u}\left(\frac{4}{3}\,du\right) = \frac{4}{3}\int_2^9 \left(1 - \frac{1}{u}\right)du = \frac{4}{3}\Big[u - \ln|u|\Big]_2^9$$
$$= \tfrac{4}{3}[(9 - \ln 9) - (2 - \ln 2)] = \tfrac{4}{3}(7 - \ln 9 + \ln 2) = \tfrac{4}{3}\left(7 + \ln\tfrac{2}{9}\right)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

81. From the graph, it appears that the area under the curve is about $1 + \left(\text{a little more than } \frac{1}{2} \cdot 1 \cdot 0.7\right)$, or about 1.4. The exact area is given by $A = \int_0^1 \sqrt{2x+1}\,dx$. Let $u = 2x+1$, so $du = 2\,dx$. The limits change to $2 \cdot 0 + 1 = 1$ and $2 \cdot 1 + 1 = 3$, and

$A = \int_1^3 \sqrt{u}\left(\frac{1}{2}\,du\right) = \frac{1}{2}\left[\frac{2}{3}u^{3/2}\right]_1^3 = \frac{1}{3}\left(3\sqrt{3} - 1\right) = \sqrt{3} - \frac{1}{3} \approx 1.399.$

82. From the graph, it appears that the area under the curve is almost $\frac{1}{2} \cdot \pi \cdot 2.6$, or about 4. The exact area is given by

$$A = \int_0^\pi (2\sin x - \sin 2x)\,dx = -2\left[\cos x\right]_0^\pi - \int_0^\pi \sin 2x\,dx$$
$$= -2(-1-1) - 0 = 4$$

Note: $\int_0^\pi \sin 2x\,dx = 0$ since it is clear from the graph of $y = \sin 2x$ that $\int_{\pi/2}^\pi \sin 2x\,dx = -\int_0^{\pi/2} \sin 2x\,dx$.

83. First write the integral as a sum of two integrals:

$I = \int_{-2}^2 (x+3)\sqrt{4-x^2}\,dx = I_1 + I_2 = \int_{-2}^2 x\sqrt{4-x^2}\,dx + \int_{-2}^2 3\sqrt{4-x^2}\,dx$. $I_1 = 0$ by Theorem 7(b), since $f(x) = x\sqrt{4-x^2}$ is an odd function and we are integrating from $x = -2$ to $x = 2$. We interpret I_2 as three times the area of a semicircle with radius 2, so $I = 0 + 3 \cdot \frac{1}{2}\left(\pi \cdot 2^2\right) = 6\pi$.

84. Let $u = x^2$. Then $du = 2x\,dx$ and the limits are unchanged ($0^2 = 0$ and $1^2 = 1$), so

$I = \int_0^1 x\sqrt{1-x^4}\,dx = \frac{1}{2}\int_0^1 \sqrt{1-u^2}\,du$. But this integral can be interpreted as the area of a quarter-circle with radius 1. So $I = \frac{1}{2} \cdot \frac{1}{4}\left(\pi \cdot 1^2\right) = \frac{1}{8}\pi$.

85. **First Figure** Let $u = \sqrt{x}$, so $x = u^2$ and $dx = 2u\,du$. When $x = 0$, $u = 0$; when $x = 1$, $u = 1$. Thus,
$A_1 = \int_0^1 e^{\sqrt{x}}\,dx = \int_0^1 e^u(2u\,du) = 2\int_0^1 ue^u\,du.$

Second Figure $A_2 = \int_0^1 2xe^x\,dx = 2\int_0^1 ue^u\,du.$

Third Figure Let $u = \sin x$, so $du = \cos x\,dx$. When $x = 0$, $u = 0$; when $x = \frac{\pi}{2}$, $u = 1$. Thus,
$A_3 = \int_0^{\pi/2} e^{\sin x}\sin 2x\,dx = \int_0^{\pi/2} e^{\sin x}(2\sin x\,\cos x)\,dx = \int_0^1 e^u(2u\,du) = 2\int_0^1 ue^u\,du.$

Since $A_1 = A_2 = A_3$, all three areas are equal.

86. Let $u = \dfrac{\pi t}{12}$. Then $du = \dfrac{\pi}{12}\,dt$ and

$$\int_0^{24} R(t)\,dt = \int_0^{24}\left[85 - 0.18\cos\left(\frac{\pi t}{12}\right)\right]dt = \int_0^{2\pi}(85 - 0.18\cos u)\left(\frac{12}{\pi}\,du\right) = \frac{12}{\pi}\left[85u - 0.18\sin u\right]_0^{2\pi}$$
$$= \frac{12}{\pi}\left[(85 \cdot 2\pi - 0) - (0 - 0)\right] = 2040 \text{ kcal}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

87. The rate is measured in liters per minute. Integrating from $t = 0$ minutes to $t = 60$ minutes will give us the total amount of oil that leaks out (in liters) during the first hour.

$$\int_0^{60} r(t)\,dt = \int_0^{60} 100e^{-0.01t}\,dt \qquad [u = -0.01t, du = -0.01dt]$$
$$= 100\int_0^{-0.6} e^u(-100\,du) = -10{,}000\left[e^u\right]_0^{-0.6} = -10{,}000(e^{-0.6} - 1) \approx 4511.9 \approx 4512 \text{ liters}$$

88. Let $r(t) = ae^{bt}$ with $a = 450.268$ and $b = 1.12567$, and $n(t) =$ population after t hours. Since $r(t) = n'(t)$, $\int_0^3 r(t)\,dt = n(3) - n(0)$ is the total change in the population after three hours. Since we start with 400 bacteria, the population will be

$$n(3) = 400 + \int_0^3 r(t)\,dt = 400 + \int_0^3 ae^{bt}\,dt = 400 + \frac{a}{b}\left[e^{bt}\right]_0^3 = 400 + \frac{a}{b}\left(e^{3b} - 1\right)$$
$$\approx 400 + 11{,}313 = 11{,}713 \text{ bacteria}$$

89. The volume of inhaled air in the lungs at time t is

$$V(t) = \int_0^t f(u)\,du = \int_0^t \frac{1}{2}\sin\left(\frac{2\pi}{5}u\right)du = \int_0^{2\pi t/5} \frac{1}{2}\sin v\left(\frac{5}{2\pi}\,dv\right) \qquad \left[\text{substitute } v = \tfrac{2\pi}{5}u,\ dv = \tfrac{2\pi}{5}\,du\right]$$
$$= \frac{5}{4\pi}\left[-\cos v\right]_0^{2\pi t/5} = \frac{5}{4\pi}\left[-\cos\left(\frac{2\pi}{5}t\right) + 1\right] = \frac{5}{4\pi}\left[1 - \cos\left(\frac{2\pi}{5}t\right)\right] \text{ liters}$$

90. The rate G is measured in kilograms per year. Integrating from $t = 0$ years (2000) to $t = 20$ years (2020) will give us the net change in biomass from 2000 to 2020.

$$\int_0^{20} \frac{60{,}000e^{-0.6t}}{(1 + 5e^{-0.6t})^2}\,dt = \int_6^{1+5e^{-12}} \frac{60{,}000}{u^2}\left(-\tfrac{1}{3}\,du\right) \qquad \begin{bmatrix} u = 1 + 5e^{-0.6t}, \\ du = -3e^{-0.6t}dt \end{bmatrix}$$
$$= \left[\frac{20{,}000}{u}\right]_6^{1+5e^{-12}} = \frac{20{,}000}{1 + 5e^{-12}} - \frac{20{,}000}{6} \approx 16{,}666$$

Thus, the predicted biomass for the year 2020 is approximately $25{,}000 + 16{,}666 = 41{,}666$ kg.

91. $$\int_0^{30} u(t)\,dt = \int_0^{30} \frac{r}{V}C_0e^{-rt/V}\,dt = C_0\int_1^{e^{-30r/V}} (-dx) \qquad \begin{bmatrix} x = e^{-rt/V}, \\ dx = -\frac{r}{V}e^{-rt/V}dt \end{bmatrix}$$
$$= C_0\Big[-x\Big]_1^{e^{-30r/V}} = C_0(-e^{-30r/V} + 1)$$

The integral $\int_0^{30} u(t)\,dt$ represents the total amount of urea removed from the blood in the first 30 minutes of dialysis.

92. Number of calculators $= x(4) - x(2) = \int_2^4 5000\left[1 - 100(t + 10)^{-2}\right]dt$
$$= 5000\left[t + 100(t + 10)^{-1}\right]_2^4 = 5000\left[\left(4 + \tfrac{100}{14}\right) - \left(2 + \tfrac{100}{12}\right)\right] \approx 4048$$

93. Let $u = 2x$. Then $du = 2\,dx$, so $\int_0^2 f(2x)\,dx = \int_0^4 f(u)\left(\frac{1}{2}\,du\right) = \frac{1}{2}\int_0^4 f(u)\,du = \frac{1}{2}(10) = 5$.

94. Let $u = x^2$. Then $du = 2x\,dx$, so $\int_0^3 xf(x^2)\,dx = \int_0^9 f(u)\left(\frac{1}{2}\,du\right) = \frac{1}{2}\int_0^9 f(u)\,du = \frac{1}{2}(4) = 2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

95. Let $u = -x$. Then $du = -dx$, so

$$\int_a^b f(-x)\,dx = \int_{-a}^{-b} f(u)(-du) = \int_{-b}^{-a} f(u)\,du = \int_{-b}^{-a} f(x)\,dx$$

From the diagram, we see that the equality follows from the fact that we are reflecting the graph of f, and the limits of integration, about the y-axis.

96. Let $u = x + c$. Then $du = dx$, so

$$\int_a^b f(x+c)\,dx = \int_{a+c}^{b+c} f(u)\,du = \int_{a+c}^{b+c} f(x)\,dx$$

From the diagram, we see that the equality follows from the fact that we are translating the graph of f, and the limits of integration, by a distance c.

97. Let $u = 1 - x$. Then $x = 1 - u$ and $dx = -du$, so

$$\int_0^1 x^a(1-x)^b\,dx = \int_1^0 (1-u)^a\,u^b(-du) = \int_0^1 u^b(1-u)^a\,du = \int_0^1 x^b(1-x)^a\,dx.$$

98. Let $u = \pi - x$. Then $du = -dx$. When $x = \pi$, $u = 0$ and when $x = 0$, $u = \pi$. So

$$\begin{aligned}\int_0^\pi x f(\sin x)\,dx &= -\int_\pi^0 (\pi - u)\,f(\sin(\pi - u))\,du = \int_0^\pi (\pi - u)\,f(\sin u)\,du \\ &= \pi\int_0^\pi f(\sin u)\,du - \int_0^\pi u\,f(\sin u)\,du = \pi\int_0^\pi f(\sin x)\,dx - \int_0^\pi x\,f(\sin x)\,dx \quad \Rightarrow\end{aligned}$$

$$2\int_0^\pi x\,f(\sin x)\,dx = \pi\int_0^\pi f(\sin x)\,dx \quad \Rightarrow \quad \int_0^\pi x\,f(\sin x)\,dx = \tfrac{\pi}{2}\int_0^\pi f(\sin x)\,dx.$$

99. $\dfrac{x\sin x}{1+\cos^2 x} = x \cdot \dfrac{\sin x}{2 - \sin^2 x} = x\,f(\sin x)$, where $f(t) = \dfrac{t}{2-t^2}$. By Exercise 98,

$$\int_0^\pi \frac{x\sin x}{1+\cos^2 x}\,dx = \int_0^\pi x\,f(\sin x)\,dx = \frac{\pi}{2}\int_0^\pi f(\sin x)\,dx = \frac{\pi}{2}\int_0^\pi \frac{\sin x}{1+\cos^2 x}\,dx$$

Let $u = \cos x$. Then $du = -\sin x\,dx$. When $x = \pi$, $u = -1$ and when $x = 0$, $u = 1$. So

$$\begin{aligned}\frac{\pi}{2}\int_0^\pi \frac{\sin x}{1+\cos^2 x}\,dx &= -\frac{\pi}{2}\int_1^{-1} \frac{du}{1+u^2} = \frac{\pi}{2}\int_{-1}^1 \frac{du}{1+u^2} = \frac{\pi}{2}\left[\tan^{-1} u\right]_{-1}^1 \\ &= \frac{\pi}{2}[\tan^{-1} 1 - \tan^{-1}(-1)] = \frac{\pi}{2}\left[\frac{\pi}{4} - \left(-\frac{\pi}{4}\right)\right] = \frac{\pi^2}{4}\end{aligned}$$

100. (a) $\int_0^{\pi/2} f(\cos x)\,dx = \int_0^{\pi/2} f\left[\sin\left(\frac{\pi}{2} - x\right)\right]dx \qquad \left[u = \frac{\pi}{2} - x,\ du = -dx\right]$

$$= \int_{\pi/2}^0 f(\sin u)(-du) = \int_0^{\pi/2} f(\sin u)\,du = \int_0^{\pi/2} f(\sin x)\,dx$$

Continuity of f is needed in order to apply the substitution rule for definite integrals.

(b) In part (a), take $f(x) = x^2$, so $\int_0^{\pi/2} \cos^2 x\,dx = \int_0^{\pi/2} \sin^2 x\,dx$. Now

$$\int_0^{\pi/2} \cos^2 x\,dx + \int_0^{\pi/2} \sin^2 x\,dx = \int_0^{\pi/2} (\cos^2 x + \sin^2 x)\,dx = \int_0^{\pi/2} 1\,dx = \left[x\right]_0^{\pi/2} = \tfrac{\pi}{2},$$

so $2\int_0^{\pi/2} \cos^2 x\,dx = \frac{\pi}{2} \quad \Rightarrow \quad \int_0^{\pi/2} \cos^2 x\,dx = \frac{\pi}{4} \quad \left[= \int_0^{\pi/2} \sin^2 x\,dx\right].$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5 Review

TRUE-FALSE QUIZ

1. True by Property 2 of the Integral in Section 5.2.

2. False. Try $a = 0$, $b = 2$, $f(x) = g(x) = 1$ as a counterexample.

3. True by Property 3 of the Integral in Section 5.2.

4. False. You can't take a variable outside the integral sign. For example, using $f(x) = 1$ on $[0, 1]$, $\int_0^1 x\, f(x)\, dx = \int_0^1 x\, dx = \left[\frac{1}{2}x^2\right]_0^1 = \frac{1}{2}$ (a constant) while $x \int_0^1 1\, dx = x\,[x]_0^1 = x \cdot 1 = x$ (a variable).

5. False. For example, let $f(x) = x^2$. Then $\int_0^1 \sqrt{x^2}\, dx = \int_0^1 x\, dx = \frac{1}{2}$, but $\sqrt{\int_0^1 x^2\, dx} = \sqrt{\frac{1}{3}} = \frac{1}{\sqrt{3}}$.

6. True. The definite integral is a number; it does not depend on the choice of variable name. See Note 2 of Section 5.2.

7. True by the Net Change Theorem.

8. False. For example, let $v(t) = 1 - t$. On $0 \le t \le 2$, the distance traveled is $\int_0^2 |v(t)|\, dt = \int_0^1 (1 - t)\, dt + \int_1^2 (t - 1)\, dt = \frac{1}{2} + \frac{1}{2} = 1$, but $\int_0^2 v(t)\, dt = \int_0^2 (1 - t)\, dt = 0$. The given integral represents net displacement.

9. False. $\int_a^b f'(x)\,[f(x)]^4\, dx$ is a definite integral and, thus, is a number; $\frac{1}{5}[f(x)]^5 + C$ is a family of functions. The statement would be true without the limits of integration.

10. False. For example, let $a = 0$, $b = 1$, $f(x) = 3$, $g(x) = x$. $f(x) > g(x)$ for each x in $(0, 1)$, but $f'(x) = 0 < 1 = g'(x)$ for $x \in (0, 1)$.

11. True by Comparison Property 7 of the Integral in Section 5.2.

12. True.
$$\begin{aligned}\int_{-5}^{5} \left(ax^2 + bx + c\right) dx &= \int_{-5}^{5} \left(ax^2 + c\right) dx + \int_{-5}^{5} bx\, dx \\ &= 2\int_0^5 \left(ax^2 + c\right) dx + 0 \quad \text{[because } ax^2 + c \text{ is even and } bx \text{ is odd]}\end{aligned}$$

13. False. For example, the function $y = |x|$ is continuous on $\mathbb{R}$, but has no derivative at $x = 0$.

14. True by FTC1.

15. True by Property 5 of Integrals.

16. False. For example, $\int_0^1 \left(x - \frac{1}{2}\right) dx = \left[\frac{1}{2}x^2 - \frac{1}{2}x\right]_0^1 = \left(\frac{1}{2} - \frac{1}{2}\right) - (0 - 0) = 0$, but $f(x) = x - \frac{1}{2} \neq 0$.

17. False. $\int_a^b f(x)\, dx$ is a constant, so $\dfrac{d}{dx}\left(\int_a^b f(x)\, dx\right) = 0$, not $f(x)$ [unless $f(x) = 0$]. Compare the given statement carefully with FTC1, in which the upper limit in the integral is x.

18. False. See the paragraph before Figure 4 in Section 5.2, and notice that $y = x - x^3 < 0$ for $1 < x \le 2$.

19. False. The function $f(x) = 1/x^4$ is not bounded on the interval $[-2, 1]$. It has an infinite discontinuity at $x = 0$, so it is not integrable on the interval. (If the integral were to exist, a positive value would be expected, by Comparison Property 5.2.6 of Integrals.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. False. For example, if $f(x) = \begin{cases} 1 & \text{if } 0 \le x \le 1 \\ 0 & \text{if } -1 \le x < 0 \end{cases}$ then f has a jump discontinuity at 0, but $\int_{-1}^{1} f(x)\,dx$ exists and is equal to 1.

EXERCISES

1. (a)

$$L_6 = \sum_{i=1}^{6} f(x_{i-1})\,\Delta x \quad [\Delta x = \tfrac{6-0}{6} = 1]$$

$$= f(x_0)\cdot 1 + f(x_1)\cdot 1 + f(x_2)\cdot 1 + f(x_3)\cdot 1 + f(x_4)\cdot 1 + f(x_5)\cdot 1$$

$$\approx 2 + 3.5 + 4 + 2 + (-1) + (-2.5) = 8$$

The Riemann sum represents the sum of the areas of the four rectangles above the x-axis minus the sum of the areas of the two rectangles below the x-axis.

(b)

$$M_6 = \sum_{i=1}^{6} f(\overline{x}_i)\,\Delta x \quad [\Delta x = \tfrac{6-0}{6} = 1]$$

$$= f(\overline{x}_1)\cdot 1 + f(\overline{x}_2)\cdot 1 + f(\overline{x}_3)\cdot 1 + f(\overline{x}_4)\cdot 1 + f(\overline{x}_5)\cdot 1 + f(\overline{x}_6)\cdot 1$$

$$= f(0.5) + f(1.5) + f(2.5) + f(3.5) + f(4.5) + f(5.5)$$

$$\approx 3 + 3.9 + 3.4 + 0.3 + (-2) + (-2.9) = 5.7$$

The Riemann sum represents the sum of the areas of the four rectangles above the x-axis minus the sum of the areas of the two rectangles below the x-axis.

2. (a)

$f(x) = x^2 - x$ and $\Delta x = \frac{2-0}{4} = 0.5 \quad \Rightarrow$

$$R_4 = 0.5f(0.5) + 0.5f(1) + 0.5f(1.5) + 0.5f(2)$$

$$= 0.5(-0.25 + 0 + 0.75 + 2) = 1.25$$

The Riemann sum represents the sum of the areas of the two rectangles above the x-axis minus the area of the rectangle below the x-axis. (The second rectangle vanishes.)

(b) $\int_0^2 \left(x^2 - x\right) dx = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} f(x_i)\ \Delta x \qquad [\Delta x = 2/n \text{ and } x_i = 2i/n]$

$$= \lim_{n\to\infty} \sum_{i=1}^{n} \left(\frac{4i^2}{n^2} - \frac{2i}{n}\right)\left(\frac{2}{n}\right) = \lim_{n\to\infty} \frac{2}{n}\left[\frac{4}{n^2}\sum_{i=1}^{n} i^2 - \frac{2}{n}\sum_{i=1}^{n} i\right]$$

$$= \lim_{n\to\infty} \left[\frac{8}{n^3}\cdot\frac{n(n+1)(2n+1)}{6} - \frac{4}{n^2}\cdot\frac{n(n+1)}{2}\right] = \lim_{n\to\infty}\left[\frac{4}{3}\cdot\frac{n+1}{n}\cdot\frac{2n+1}{n} - 2\cdot\frac{n+1}{n}\right]$$

$$= \lim_{n\to\infty}\left[\frac{4}{3}\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right) - 2\left(1+\frac{1}{n}\right)\right] = \tfrac{4}{3}\cdot 1\cdot 2 - 2\cdot 1 = \tfrac{2}{3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\int_0^2 \left(x^2 - x\right) dx = \left[\frac{1}{3}x^3 - \frac{1}{2}x^2\right]_0^2 = \left(\frac{8}{3} - 2\right) = \frac{2}{3}$

(d)

$\int_0^2 \left(x^2 - x\right) dx = A_1 - A_2$, where A_1 and A_2 are the areas shown in the diagram.

3. $\int_0^1 \left(x + \sqrt{1 - x^2}\right) dx = \int_0^1 x\, dx + \int_0^1 \sqrt{1 - x^2}\, dx = I_1 + I_2$.

I_1 can be interpreted as the area of the triangle shown in the figure and I_2 can be interpreted as the area of the quarter-circle.

Area $= \frac{1}{2}(1)(1) + \frac{1}{4}(\pi)(1)^2 = \frac{1}{2} + \frac{\pi}{4}$.

4. On $[0, \pi]$, $\lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \sin x_i\, \Delta x = \int_0^\pi \sin x\, dx = [-\cos x]_0^\pi = -(-1) - (-1) = 2$.

5. $\int_0^6 f(x)\, dx = \int_0^4 f(x)\, dx + \int_4^6 f(x)\, dx \quad \Rightarrow \quad 10 = 7 + \int_4^6 f(x)\, dx \quad \Rightarrow \quad \int_4^6 f(x)\, dx = 10 - 7 = 3$

6. (a) $\int_1^5 (x + 2x^5)\, dx = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} f(x_i)\, \Delta x \qquad \left[\Delta x = \dfrac{5-1}{n} = \dfrac{4}{n},\ x_i = 1 + \dfrac{4i}{n}\right]$

$$= \lim_{n\to\infty} \sum_{i=1}^{n} \left[\left(1 + \frac{4i}{n}\right) + 2\left(1 + \frac{4i}{n}\right)^5\right] \cdot \frac{4}{n} \overset{\text{CAS}}{=} \lim_{n\to\infty} \frac{1305n^4 + 3126n^3 + 2080n^2 - 256}{n^3} \cdot \frac{4}{n}$$

$$= 5220$$

(b) $\int_1^5 (x + 2x^5)\, dx = \left[\frac{1}{2}x^2 + \frac{2}{6}x^6\right]_1^5 = \left(\frac{25}{2} + \frac{15{,}625}{3}\right) - \left(\frac{1}{2} + \frac{1}{3}\right) = 12 + 5208 = 5220$

7. First note that either a or b must be the graph of $\int_0^x f(t)\, dt$, since $\int_0^0 f(t)\, dt = 0$, and $c(0) \neq 0$. Now notice that $b > 0$ when c is increasing, and that $c > 0$ when a is increasing. It follows that c is the graph of $f(x)$, b is the graph of $f'(x)$, and a is the graph of $\int_0^x f(t)\, dt$.

8. (a) By the Net Change Theorem (FTC2), $\displaystyle\int_0^1 \frac{d}{dx}\left(e^{\arctan x}\right) dx = \left[e^{\arctan x}\right]_0^1 = e^{\pi/4} - 1$

(b) $\displaystyle\frac{d}{dx}\int_0^1 e^{\arctan x}\, dx = 0$ since this is the derivative of a constant.

(c) By FTC1, $\displaystyle\frac{d}{dx}\int_0^x e^{\arctan t}\, dt = e^{\arctan x}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $g(4) = \int_0^4 f(t)\,dt = \int_0^1 f(t)\,dt + \int_1^2 f(t)\,dt + \int_2^3 f(t)\,dt + \int_3^4 f(t)\,dt$

$= -\frac{1}{2}\cdot 1\cdot 2 \quad \begin{bmatrix}\text{area of triangle,}\\ \textit{below } t\text{-axis}\end{bmatrix} \quad + \frac{1}{2}\cdot 1\cdot 2 + 1\cdot 2 + \frac{1}{2}\cdot 1\cdot 2 = 3$

By FTC1, $g'(x) = f(x)$, so $g'(4) = f(4) = 0$.

10. $g(x) = \int_0^x f(t)\,dt \quad\Rightarrow\quad g'(x) = f(x)$ [by FTC1] $\quad\Rightarrow\quad g''(x) = f'(x)$, so $g''(4) = f'(4) = -2$, which is the slope of the line segment at $x = 4$.

11. $\int_{-1}^0 (x^2 + 5x)\,dx = \left[\frac{1}{3}x^3 + \frac{5}{2}x^2\right]_{-1}^0 = 0 - \left(-\frac{1}{3} + \frac{5}{2}\right) = -\frac{13}{6}$

12. $\int_0^T (x^4 - 8x + 7)\,dx = \left[\frac{1}{5}x^5 - 4x^2 + 7x\right]_0^T = \left(\frac{1}{5}T^5 - 4T^2 + 7T\right) - 0 = \frac{1}{5}T^5 - 4T^2 + 7T$

13. $\int_0^1 \left(1 - x^9\right)dx = \left[x - \frac{1}{10}x^{10}\right]_0^1 = \left(1 - \frac{1}{10}\right) - 0 = \frac{9}{10}$

14. Let $u = 1 - x$, so $du = -dx$ and $dx = -du$. When $x = 0$, $u = 1$; when $x = 1$, $u = 0$. Thus,

$\int_0^1 (1 - x)^9\,dx = \int_1^0 u^9(-du) = \int_0^1 u^9\,du = \frac{1}{10}\left[u^{10}\right]_0^1 = \frac{1}{10}(1 - 0) = \frac{1}{10}$.

15. $\displaystyle\int_1^9 \frac{\sqrt{u} - 2u^2}{u}\,du = \int_1^9 (u^{-1/2} - 2u)\,du = \left[2u^{1/2} - u^2\right]_1^9 = (6 - 81) - (2 - 1) = -76$

16. $\int_0^1 \left(\sqrt[4]{u} + 1\right)^2\,du = \int_0^1 (u^{1/2} + 2u^{1/4} + 1)\,du = \left[\frac{2}{3}u^{3/2} + \frac{8}{5}u^{5/4} + u\right]_0^1 = \left(\frac{2}{3} + \frac{8}{5} + 1\right) - 0 = \frac{49}{15}$

17. Let $u = y^2 + 1$, so $du = 2y\,dy$ and $y\,dy = \frac{1}{2}\,du$. When $y = 0$, $u = 1$; when $y = 1$, $u = 2$. Thus,

$\int_0^1 y(y^2 + 1)^5\,dy = \int_1^2 u^5\left(\frac{1}{2}\,du\right) = \frac{1}{2}\left[\frac{1}{6}u^6\right]_1^2 = \frac{1}{12}(64 - 1) = \frac{63}{12} = \frac{21}{4}$.

18. Let $u = 1 + y^3$, so $du = 3y^2\,dy$ and $y^2\,dy = \frac{1}{3}\,du$. When $y = 0$, $u = 1$; when $y = 2$, $u = 9$. Thus,

$\int_0^2 y^2\sqrt{1 + y^3}\,dy = \int_1^9 u^{1/2}\left(\frac{1}{3}\,du\right) = \frac{1}{3}\left[\frac{2}{3}u^{3/2}\right]_1^9 = \frac{2}{9}(27 - 1) = \frac{52}{9}$.

19. $\displaystyle\int_1^5 \frac{dt}{(t - 4)^2}$ does not exist because the function $f(t) = \dfrac{1}{(t - 4)^2}$ has an infinite discontinuity at $t = 4$; that is, f is discontinuous on the interval $[1, 5]$.

20. Let $u = 3\pi t$, so $du = 3\pi\,dt$. When $t = 0$, $u = 1$; when $t = 1$, $u = 3\pi$. Thus,

$$\int_0^1 \sin(3\pi t)\,dt = \int_0^{3\pi} \sin u\left(\frac{1}{3\pi}\,du\right) = \frac{1}{3\pi}\left[-\cos u\right]_0^{3\pi} = -\frac{1}{3\pi}(-1 - 1) = \frac{2}{3\pi}.$$

21. Let $u = v^3$, so $du = 3v^2\,dv$. When $v = 0$, $u = 0$; when $v = 1$, $u = 1$. Thus,

$\int_0^1 v^2\cos(v^3)\,dv = \int_0^1 \cos u\left(\frac{1}{3}\,du\right) = \frac{1}{3}\left[\sin u\right]_0^1 = \frac{1}{3}(\sin 1 - 0) = \frac{1}{3}\sin 1$.

22. $\displaystyle\int_{-1}^1 \frac{\sin x}{1 + x^2}\,dx = 0$ by Theorem 5.5.7(b), since $f(x) = \dfrac{\sin x}{1 + x^2}$ is an odd function.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. $\displaystyle\int_{-\pi/4}^{\pi/4} \frac{t^4 \tan t}{2+\cos t}\,dt = 0$ by Theorem 5.5.7(b), since $f(t) = \dfrac{t^4 \tan t}{2+\cos t}$ is an odd function.

24. $\displaystyle\int_{-2}^{-1} \frac{z^2+1}{z}\,dz = \int_{-2}^{-1} \left(z + \frac{1}{z}\right) dz = \left[\frac{1}{2}z^2 + \ln|z|\right]_{-2}^{-1} = \left(\frac{1}{2} + \ln|-1|\right) - \left(2 + \ln|-2|\right) = -\frac{3}{2} - \ln 2$

25. Let $u = x^2 + 1$. Then $du = 2x\,dx$ and $x\,dx = \dfrac{1}{2}\,du$, so

$$\int \frac{x}{x^2+1}\,dx = \int \frac{1}{u}\left(\frac{1}{2}\,du\right) = \frac{1}{2}\ln|u| + C = \frac{1}{2}\ln\left|x^2+1\right| + C = \frac{1}{2}\ln(x^2+1) + C \quad \text{[since } x^2+1 \text{ is positive]}.$$

26. $\displaystyle\int \frac{dx}{x^2+1} = \int \frac{1}{1+x^2}\,dx = \arctan x + C$

27. Let $u = x^2 + 4x$. Then $du = (2x+4)\,dx = 2(x+2)\,dx$, so

$$\int \frac{x+2}{\sqrt{x^2+4x}}\,dx = \int u^{-1/2}\left(\tfrac{1}{2}\,du\right) = \tfrac{1}{2}\cdot 2u^{1/2} + C = \sqrt{u} + C = \sqrt{x^2+4x} + C.$$

28. Let $u = 1 + \cot x$. Then $du = -\csc^2 x\,dx$, so $\displaystyle\int \frac{\csc^2 x}{1+\cot x}\,dx = \int \frac{1}{u}\,(-du) = -\ln|u| + C = -\ln|1+\cot x| + C$.

29. Let $u = \sin \pi t$. Then $du = \pi\,\cos \pi t\,dt$, so $\int \sin \pi t\,\cos \pi t\,dt = \int u\left(\frac{1}{\pi}\,du\right) = \frac{1}{\pi}\cdot\frac{1}{2}u^2 + C = \frac{1}{2\pi}(\sin \pi t)^2 + C$.

30. Let $u = \cos x$. Then $du = -\sin x\,dx$, so $\int \sin x\,\cos(\cos x)\,dx = -\int \cos u\,du = -\sin u + C = -\sin(\cos x) + C$.

31. Let $u = \sqrt{x}$. Then $du = \dfrac{dx}{2\sqrt{x}}$, so $\displaystyle\int \frac{e^{\sqrt{x}}}{\sqrt{x}}\,dx = 2\int e^u\,du = 2e^u + C = 2e^{\sqrt{x}} + C$.

32. Let $u = \ln x$. Then $du = \dfrac{1}{x}\,dx$, so $\displaystyle\int \frac{\sin(\ln x)}{x}\,dx = \int \sin u\,du = -\cos u + C = -\cos(\ln x) + C$.

33. Let $u = \ln(\cos x)$. Then $du = \dfrac{-\sin x}{\cos x}\,dx = -\tan x\,dx$, so

$\int \tan x\,\ln(\cos x)\,dx = -\int u\,du = -\frac{1}{2}u^2 + C = -\frac{1}{2}[\ln(\cos x)]^2 + C$.

34. Let $u = x^2$. Then $du = 2x\,dx$, so $\displaystyle\int \frac{x}{\sqrt{1-x^4}}\,dx = \frac{1}{2}\int \frac{du}{\sqrt{1-u^2}} = \frac{1}{2}\sin^{-1}u + C = \frac{1}{2}\sin^{-1}\left(x^2\right) + C$.

35. Let $u = 1 + x^4$. Then $du = 4x^3\,dx$, so $\displaystyle\int \frac{x^3}{1+x^4}\,dx = \frac{1}{4}\int \frac{1}{u}\,du = \frac{1}{4}\ln|u| + C = \frac{1}{4}\ln\left(1+x^4\right) + C$.

36. Let $u = 1 + 4x$. Then $du = 4\,dx$, so $\int \sinh(1+4x)\,dx = \frac{1}{4}\int \sinh u\,du = \frac{1}{4}\cosh u + C = \frac{1}{4}\cosh(1+4x) + C$.

37. Let $u = 1 + \sec\theta$. Then $du = \sec\theta\,\tan\theta\,d\theta$, so

$$\int \frac{\sec\theta\,\tan\theta}{1+\sec\theta}\,d\theta = \int \frac{1}{1+\sec\theta}\,(\sec\theta\,\tan\theta\,d\theta) = \int \frac{1}{u}\,du = \ln|u| + C = \ln|1+\sec\theta| + C.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. Let $u = 1 + \tan t$, so $du = \sec^2 t\,dt$. When $t = 0$, $u = 1$; when $t = \frac{\pi}{4}$, $u = 2$. Thus,

$\int_0^{\pi/4} (1 + \tan t)^3 \sec^2 t\,dt = \int_1^2 u^3\,du = \left[\frac{1}{4}u^4\right]_1^2 = \frac{1}{4}\left(2^4 - 1^4\right) = \frac{1}{4}(16 - 1) = \frac{15}{4}$.

39. Let $u = 1 - x$. Then $x = 1 - u$, $du = -dx$, and $dx = -du$, so

$$\begin{aligned}\int x(1-x)^{2/3}\,dx &= \int (1-u)\cdot u^{2/3}\,(-du) = \int (u^{5/3} - u^{2/3})\,du = \tfrac{3}{8}u^{8/3} - \tfrac{3}{5}u^{5/3} + C \\ &= \tfrac{3}{8}(1-x)^{8/3} - \tfrac{3}{5}(1-x)^{5/3} + C\end{aligned}$$

40. Let $u = x - 3$. Then $x = u + 3$ and $du = dx$, so

$$\int \frac{x}{x-3}\,dx = \int \frac{u+3}{u}\,du = \int \left(1 + \frac{3}{u}\right) du = u + 3\ln|u| + C_1 = x - 3 + 3\ln|x-3| + C_1 = x + 3\ln|x-3| + C$$

[where $C = C_1 - 3$].

41. Since $x^2 - 4 < 0$ for $0 \le x < 2$ and $x^2 - 4 > 0$ for $2 < x \le 3$, we have $\left|x^2 - 4\right| = -(x^2 - 4) = 4 - x^2$ for $0 \le x < 2$ and $\left|x^2 - 4\right| = x^2 - 4$ for $2 < x \le 3$. Thus,

$$\begin{aligned}\int_0^3 \left|x^2 - 4\right| dx &= \int_0^2 (4 - x^2)\,dx + \int_2^3 (x^2 - 4)\,dx = \left[4x - \frac{x^3}{3}\right]_0^2 + \left[\frac{x^3}{3} - 4x\right]_2^3 \\ &= \left(8 - \tfrac{8}{3}\right) - 0 + (9 - 12) - \left(\tfrac{8}{3} - 8\right) = \tfrac{16}{3} - 3 + \tfrac{16}{3} = \tfrac{32}{3} - \tfrac{9}{3} = \tfrac{23}{3}\end{aligned}$$

42. Since $\sqrt{x} - 1 < 0$ for $0 \le x < 1$ and $\sqrt{x} - 1 > 0$ for $1 < x \le 4$, we have $\left|\sqrt{x} - 1\right| = -\left(\sqrt{x} - 1\right) = 1 - \sqrt{x}$ for $0 \le x < 1$ and $\left|\sqrt{x} - 1\right| = \sqrt{x} - 1$ for $1 < x \le 4$. Thus,

$$\begin{aligned}\int_0^4 \left|\sqrt{x} - 1\right| dx &= \int_0^1 \left(1 - \sqrt{x}\right) dx + \int_1^4 \left(\sqrt{x} - 1\right) dx = \left[x - \tfrac{2}{3}x^{3/2}\right]_0^1 + \left[\tfrac{2}{3}x^{3/2} - x\right]_1^4 \\ &= \left(1 - \tfrac{2}{3}\right) - 0 + \left(\tfrac{16}{3} - 4\right) - \left(\tfrac{2}{3} - 1\right) = \tfrac{1}{3} + \tfrac{16}{3} - 4 + \tfrac{1}{3} = 6 - 4 = 2\end{aligned}$$

43. Let $u = 1 + \sin x$. Then $du = \cos x\,dx$, so

$$\int \frac{\cos x\,dx}{\sqrt{1 + \sin x}} = \int u^{-1/2}\,du = 2u^{1/2} + C = 2\sqrt{1 + \sin x} + C.$$

44. Let $u = x^2 + 1$. Then $x^2 = u - 1$ and $x\,dx = \frac{1}{2}\,du$, so

$$\begin{aligned}\int \frac{x^3}{\sqrt{x^2+1}}\,dx &= \int \frac{(u-1)}{\sqrt{u}}\left(\tfrac{1}{2}\,du\right) = \frac{1}{2}\int \left(u^{1/2} - u^{-1/2}\right) du \\ &= \tfrac{1}{2}\left(\tfrac{2}{3}u^{3/2} - 2u^{1/2}\right) + C = \tfrac{1}{3}(x^2+1)^{3/2} - (x^2+1)^{1/2} + C \\ &= \tfrac{1}{3}(x^2+1)^{1/2}\left[(x^2+1) - 3\right] + C = \tfrac{1}{3}\sqrt{x^2+1}\,(x^2 - 2) + C\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. From the graph, it appears that the area under the curve $y = x\sqrt{x}$ between $x = 0$ and $x = 4$ is somewhat less than half the area of an 8×4 rectangle, so perhaps about 13 or 14. To find the exact value, we evaluate

$\int_0^4 x\sqrt{x}\,dx = \int_0^4 x^{3/2}\,dx = \left[\frac{2}{5}x^{5/2}\right]_0^4 = \frac{2}{5}(4)^{5/2} = \frac{64}{5} = 12.8.$

46. From the graph, it seems as though $\int_0^{2\pi} \cos^2 x\,\sin x\,dx$ is equal to 0. To evaluate the integral, let $u = \cos x \quad \Rightarrow \quad du = -\sin x\,dx$. Thus, $I = \int_1^1 u^2\,(-du) = 0$.

47. $\text{Area} = \int_0^4 (x^2+5)\,dx = \left[\frac{1}{3}x^3 + 5x\right]_0^4 = \left(\frac{64}{3} + 20\right) - 0 = \frac{124}{3}$

48. $\text{Area} = \int_0^{\pi/2} \sin x\,dx = \Big[-\cos x\Big]_0^{\pi/2} = -\cos\dfrac{\pi}{2} - (-\cos 0) = 0 + 1 = 1.$

49. (a) $\int_1^5 f(x)\,dx = \int_1^3 f(x)\,dx + \int_3^4 f(x)\,dx + \int_4^5 f(x)\,dx = 3 - 2 + 1 = 2$

(b) $\int_1^5 |f(x)|\,dx = \int_1^3 f(x)\,dx + \int_3^4 [-f(x)]\,dx + \int_4^5 f(x)\,dx = 3 + 2 + 1 = 6$

50. (a) $\int_1^4 f(x)\,dx + \int_3^5 f(x)\,dx = \int_1^3 f(x)\,dx + \int_3^4 f(x)\,dx + \int_3^4 f(x)\,dx + \int_4^5 f(x)\,dx = 3 - 2 - 2 + 1 = 0$

(b) $\int_1^3 2f(x)\,dx + \int_3^5 6f(x)\,dx = 2\left[\int_1^3 f(x)\,dx\right] + 6\left[\int_3^5 f(x)\,dx\right]$

$= 2\left[\int_1^3 f(x)\,dx\right] + 6\left[\int_3^4 f(x)\,dx + \int_4^5 f(x)\,dx\right]$

$= 2(3) + 6(-2+1) = 6 + 6(-1) = 0$

51. $F(x) = \displaystyle\int_0^x \frac{t^2}{1+t^3}\,dt \quad \Rightarrow \quad F'(x) = \frac{d}{dx}\int_0^x \frac{t^2}{1+t^3}\,dt = \frac{x^2}{1+x^3}$

52. $F(x) = \displaystyle\int_x^1 \sqrt{t+\sin t}\,dt = -\int_1^x \sqrt{t+\sin t}\,dt \quad \Rightarrow \quad F'(x) = -\frac{d}{dx}\int_1^x \sqrt{t+\sin t}\,dt = -\sqrt{x+\sin x}$

53. Let $u = x^4$. Then $\dfrac{du}{dx} = 4x^3$. Also, $\dfrac{dg}{dx} = \dfrac{dg}{du}\dfrac{du}{dx}$, so

$$g'(x) = \frac{d}{dx}\int_0^{x^4} \cos(t^2)\,dt = \frac{d}{du}\int_0^u \cos(t^2)\,dt \cdot \frac{du}{dx} = \cos(u^2)\,\frac{du}{dx} = 4x^3\cos(x^8).$$

54. Let $u = \sin x$. Then $\dfrac{du}{dx} = \cos x$. Also, $\dfrac{dg}{dx} = \dfrac{dg}{du}\dfrac{du}{dx}$, so

$$g'(x) = \frac{d}{dx}\int_1^{\sin x} \frac{1-t^2}{1+t^4}\,dt = \frac{d}{du}\int_1^u \frac{1-t^2}{1+t^4}\,dt \cdot \frac{du}{dx} = \frac{1-u^2}{1+u^4}\cdot\frac{du}{dx} = \frac{1-\sin^2 x}{1+\sin^4 x}\cdot\cos x = \frac{\cos^3 x}{1+\sin^4 x}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

55. $y = \int_{\sqrt{x}}^{x} \frac{e^t}{t}\,dt = \int_{\sqrt{x}}^{1} \frac{e^t}{t}\,dt + \int_{1}^{x} \frac{e^t}{t}\,dt = -\int_{1}^{\sqrt{x}} \frac{e^t}{t}\,dt + \int_{1}^{x} \frac{e^t}{t}\,dt \quad \Rightarrow$

$\frac{dy}{dx} = -\frac{d}{dx}\left(\int_{1}^{\sqrt{x}} \frac{e^t}{t}\,dt\right) + \frac{d}{dx}\left(\int_{1}^{x} \frac{e^t}{t}\,dt\right)$. Let $u = \sqrt{x}$. Then

$$\frac{d}{dx}\int_{1}^{\sqrt{x}} \frac{e^t}{t}\,dt = \frac{d}{dx}\int_{1}^{u} \frac{e^t}{t}\,dt = \frac{d}{du}\left(\int_{1}^{u} \frac{e^t}{t}\,dt\right)\frac{du}{dx} = \frac{e^u}{u}\cdot\frac{1}{2\sqrt{x}} = \frac{e^{\sqrt{x}}}{\sqrt{x}}\cdot\frac{1}{2\sqrt{x}} = \frac{e^{\sqrt{x}}}{2x},$$

so $\frac{dy}{dx} = -\frac{e^{\sqrt{x}}}{2x} + \frac{e^x}{x} = \frac{2e^x - e^{\sqrt{x}}}{2x}$.

56. $y = \int_{2x}^{3x+1} \sin(t^4)\,dt = \int_{2x}^{0} \sin(t^4)\,dt + \int_{0}^{3x+1} \sin(t^4)\,dt = \int_{0}^{3x+1} \sin(t^4)\,dt - \int_{0}^{2x} \sin(t^4)\,dt \quad \Rightarrow$

$y' = \sin\left[(3x+1)^4\right]\cdot\frac{d}{dx}(3x+1) - \sin\left[(2x)^4\right]\cdot\frac{d}{dx}(2x) = 3\sin\left[(3x+1)^4\right] - 2\sin\left[(2x)^4\right]$

57. If $1 \le x \le 3$, then $\sqrt{1^2+3} \le \sqrt{x^2+3} \le \sqrt{3^2+3} \quad \Rightarrow \quad 2 \le \sqrt{x^2+3} \le 2\sqrt{3}$, so

$2(3-1) \le \int_1^3 \sqrt{x^2+3}\,dx \le 2\sqrt{3}(3-1)$; that is, $4 \le \int_1^3 \sqrt{x^2+3}\,dx \le 4\sqrt{3}$.

58. If $2 \le x \le 4$, then $2^3 + 2 \le x^3 + 2 \le 4^3 + 2 \quad \Rightarrow \quad 10 \le x^3 + 2 \le 66$ and $\frac{1}{66} \le \frac{1}{x^3+2} \le \frac{1}{10}$, so

$\frac{1}{66}(4-2) \le \int_2^4 \frac{1}{x^3+2}\,dx \le \frac{1}{10}(4-2)$; that is, $\frac{1}{33} \le \int_2^4 \frac{1}{x^3+2}\,dx \le \frac{1}{5}$.

59. $0 \le x \le 1 \quad \Rightarrow \quad 0 \le \cos x \le 1 \quad \Rightarrow \quad x^2\cos x \le x^2 \quad \Rightarrow \quad \int_0^1 x^2\cos x\,dx \le \int_0^1 x^2\,dx = \frac{1}{3}\left[x^3\right]_0^1 = \frac{1}{3}$ [Property 5.2.7].

60. From a graph we see that $\frac{\sin x}{x}$ is decreasing on the interval $\left[\frac{\pi}{4}, \frac{\pi}{2}\right]$. Therefore, the largest value of $\frac{\sin x}{x}$ on $\left[\frac{\pi}{4}, \frac{\pi}{2}\right]$ is

$\frac{\sin(\pi/4)}{\pi/4} = \frac{\sqrt{2}/2}{\pi/4} = \frac{2\sqrt{2}}{\pi}$. (Alternatively we could apply the Closed Interval Method to find the maximum value.) By

Property 5.2.8 with $M = \frac{2\sqrt{2}}{\pi}$, we have $\int_{\pi/4}^{\pi/2} \frac{\sin x}{x}\,dx \le \frac{2\sqrt{2}}{\pi}\left(\frac{\pi}{2} - \frac{\pi}{4}\right) = \frac{\sqrt{2}}{2}$.

61. $\cos x \le 1 \quad \Rightarrow \quad e^x\cos x \le e^x \quad \Rightarrow \quad \int_0^1 e^x\cos x\,dx \le \int_0^1 e^x\,dx = [e^x]_0^1 = e - 1$

62. For $0 \le x \le 1$, $0 \le \sin^{-1} x \le \frac{\pi}{2}$, so $\int_0^1 x\sin^{-1} x\,dx \le \int_0^1 x\left(\frac{\pi}{2}\right)dx = \left[\frac{\pi}{4}x^2\right]_0^1 = \frac{\pi}{4}$.

63. $\Delta x = (3-0)/6 = \frac{1}{2}$, so the endpoints are 0, $\frac{1}{2}$, 1, $\frac{3}{2}$, 2, $\frac{5}{2}$, and 3, and the midpoints are $\frac{1}{4}$, $\frac{3}{4}$, $\frac{5}{4}$, $\frac{7}{4}$, $\frac{9}{4}$, and $\frac{11}{4}$.

The Midpoint Rule gives

$\int_0^3 \sin(x^3)\,dx \approx \sum_{i=1}^{6} f(\overline{x}_i)\,\Delta x = \frac{1}{2}\left[\sin\left(\frac{1}{4}\right)^3 + \sin\left(\frac{3}{4}\right)^3 + \sin\left(\frac{5}{4}\right)^3 + \sin\left(\frac{7}{4}\right)^3 + \sin\left(\frac{9}{4}\right)^3 + \sin\left(\frac{11}{4}\right)^3\right] \approx 0.2810.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. (a) Displacement $= \int_0^5 \left(t^2 - t\right) dt = \left[\frac{1}{3}t^3 - \frac{1}{2}t^2\right]_0^5 = \frac{125}{3} - \frac{25}{2} = \frac{175}{6} = 29.1\overline{6}$ meters

(b) Distance traveled $= \int_0^5 \left|t^2 - t\right| dt = \int_0^5 |t(t-1)|\, dt = \int_0^1 \left(t - t^2\right) dt + \int_1^5 \left(t^2 - t\right) dt$

$$= \left[\tfrac{1}{2}t^2 - \tfrac{1}{3}t^3\right]_0^1 + \left[\tfrac{1}{3}t^3 - \tfrac{1}{2}t^2\right]_1^5 = \tfrac{1}{2} - \tfrac{1}{3} - 0 + \left(\tfrac{125}{3} - \tfrac{25}{2}\right) - \left(\tfrac{1}{3} - \tfrac{1}{2}\right) = \tfrac{177}{6} = 29.5 \text{ meters}$$

65. Note that $r(t) = b'(t)$, where $b(t) =$ the number of barrels of oil consumed up to time t. So, by the Net Change Theorem, $\int_{15}^{20} r(t)\, dt = b(20) - b(15)$ represents the number of barrels of oil consumed from Jan. 1, 2015, through Jan. 1, 2020.

66. Distance covered $= \int_0^{5.0} v(t)\, dt \approx M_5 = \frac{5.0-0}{5}[v(0.5) + v(1.5) + v(2.5) + v(3.5) + v(4.5)]$

$$= 1(4.67 + 8.86 + 10.22 + 10.67 + 10.81) = 45.23 \text{ m}$$

67. We use the Midpoint Rule with $n = 6$ and $\Delta t = \frac{24-0}{6} = 4$. The increase in the bee population was

$$\int_0^{24} r(t)\, dt \approx M_6 = 4[r(2) + r(6) + r(10) + r(14) + r(18) + r(22)]$$
$$\approx 4[50 + 1000 + 7000 + 8550 + 1350 + 150] = 4(18{,}100) = 72{,}400$$

68. $A_1 = \frac{1}{2}bh = \frac{1}{2}(2)(2) = 2$, $A_2 = \frac{1}{2}bh = \frac{1}{2}(1)(1) = \frac{1}{2}$, and since $y = -\sqrt{1-x^2}$ for $0 \le x \le 1$ represents a quarter-circle with radius 1, $A_3 = \frac{1}{4}\pi r^2 = \frac{1}{4}\pi(1)^2 = \frac{\pi}{4}$. So

$$\int_{-3}^{1} f(x)\, dx = A_1 - A_2 - A_3 = 2 - \tfrac{1}{2} - \tfrac{\pi}{4} = \tfrac{1}{4}(6 - \pi)$$

69. Let $u = 2\sin\theta$. Then $du = 2\cos\theta\, d\theta$ and when $\theta = 0$, $u = 0$; when $\theta = \frac{\pi}{2}$, $u = 2$. Thus,

$$\int_0^{\pi/2} f(2\sin\theta)\cos\theta\, d\theta = \int_0^2 f(u)\left(\tfrac{1}{2}\, du\right) = \tfrac{1}{2}\int_0^2 f(u)\, du = \tfrac{1}{2}\int_0^2 f(x)\, dx = \tfrac{1}{2}(6) = 3.$$

70. (a) C is increasing on those intervals where C' is positive. By the Fundamental Theorem of Calculus, $C'(x) = \dfrac{d}{dx}\left[\int_0^x \cos\left(\frac{\pi}{2}t^2\right)dt\right] = \cos\left(\frac{\pi}{2}x^2\right)$. This is positive when $\frac{\pi}{2}x^2$ is in the interval $\left(\left(2n - \frac{1}{2}\right)\pi, \left(2n + \frac{1}{2}\right)\pi\right)$, n any integer. This implies that $\left(2n - \frac{1}{2}\right)\pi < \frac{\pi}{2}x^2 < \left(2n + \frac{1}{2}\right)\pi \quad \Leftrightarrow \quad 0 \le |x| < 1$ or $\sqrt{4n-1} < |x| < \sqrt{4n+1}$, n any positive integer. So C is increasing on the intervals $(-1, 1)$, $\left(\sqrt{3}, \sqrt{5}\right)$, $\left(-\sqrt{5}, -\sqrt{3}\right)$, $\left(\sqrt{7}, 3\right)$, $\left(-3, -\sqrt{7}\right), \ldots$.

(b) C is concave upward on those intervals where $C'' > 0$. We differentiate C' to find C'': $C'(x) = \cos\left(\frac{\pi}{2}x^2\right) \quad \Rightarrow$ $C''(x) = -\sin\left(\frac{\pi}{2}x^2\right)\left(\frac{\pi}{2} \cdot 2x\right) = -\pi x \sin\left(\frac{\pi}{2}x^2\right)$. For $x > 0$, this is positive where $(2n-1)\pi < \frac{\pi}{2}x^2 < 2n\pi$, n any positive integer $\quad \Leftrightarrow \quad \sqrt{2(2n-1)} < x < 2\sqrt{n}$, n any positive integer. Since there is a factor of $-x$ in C'', the intervals of upward concavity for $x < 0$ are $\left(-\sqrt{2(2n+1)}, -2\sqrt{n}\right)$, n any nonnegative integer. That is, C is concave upward on $\left(-\sqrt{2}, 0\right)$, $\left(\sqrt{2}, 2\right)$, $\left(-\sqrt{6}, -2\right)$, $\left(\sqrt{6}, 2\sqrt{2}\right), \ldots$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c)

From the graphs, we can determine that $\int_0^x \cos\left(\frac{\pi}{2}t^2\right)\,dt = 0.7$ at $x \approx 0.76$ and $x \approx 1.22$.

(d)

The graphs of $S(x)$ and $C(x)$ have similar shapes, except that S's flattens out near the origin, while C's does not. Note that for $x > 0$, C is increasing where S is concave up, and C is decreasing where S is concave down. Similarly, S is increasing where C is concave down, and S is decreasing where C is concave up. For $x < 0$, these relationships are reversed; that is, C is increasing where S is concave down, and S is increasing where C is concave up. See Example 5.3.3 and Exercise 5.3.81 for a discussion of $S(x)$.

71. Area under the curve $y = \sinh cx$ between $x = 0$ and $x = 1$ is equal to 1 $\Rightarrow$ $\int_0^1 \sinh cx\,dx = 1 \;\Rightarrow\; \frac{1}{c}\left[\cosh cx\right]_0^1 = 1 \;\Rightarrow\; \frac{1}{c}(\cosh c - 1) = 1 \;\Rightarrow$ $\cosh c - 1 = c \;\Rightarrow\; \cosh c = c + 1$. From the graph, we get $c = 0$ and $c \approx 1.6161$, but $c = 0$ isn't a solution for this problem since the curve $y = \sinh cx$ becomes $y = 0$ and the area under it is 0. Thus, $c \approx 1.6161$.

72. Both numerator and denominator approach 0 as $a \to 0$, so we use l'Hospital's Rule. (Note that we are differentiating *with respect to* a, since that is the quantity which is changing.) We also use FTC1:

$$\lim_{a\to 0} T(x,t) = \lim_{a\to 0} \frac{C\int_0^a e^{-(x-u)^2/(4kt)}du}{a\sqrt{4\pi kt}} \overset{\text{H}}{=} \lim_{a\to 0} \frac{Ce^{-(x-a)^2/(4kt)}}{\sqrt{4\pi kt}} = \frac{Ce^{-x^2/(4kt)}}{\sqrt{4\pi kt}}$$

73. Using FTC1, we differentiate both sides of the given equation, $\int_1^x f(t)\,dt = (x-1)e^{2x} + \int_1^x e^{-t}f(t)\,dt$, and get

$$f(x) = e^{2x} + 2(x-1)e^{2x} + e^{-x}f(x) \;\Rightarrow\; f(x)(1-e^{-x}) = e^{2x} + 2(x-1)e^{2x} \;\Rightarrow\; f(x) = \frac{e^{2x}(2x-1)}{1-e^{-x}}.$$

74. Note that $h'(u)$ is an antiderivative of $h''(u)$, so by FTC2, $\int_1^2 h''(u)\,du = \int_1^2 (h')'(u)\,du = h'(2) - h'(1) = 5 - 2 = 3$. The other information is unnecessary.

75. Let $u = f(x)$ and $du = f'(x)\,dx$. So $2\int_a^b f(x)f'(x)\,dx = 2\int_{f(a)}^{f(b)} u\,du = \left[u^2\right]_{f(a)}^{f(b)} = [f(b)]^2 - [f(a)]^2$.

76. Let $F(x) = \displaystyle\int_2^x \sqrt{1+t^3}\,dt$. Then $F'(2) = \displaystyle\lim_{h\to 0}\frac{F(2+h)-F(2)}{h} = \lim_{h\to 0}\frac{1}{h}\int_2^{2+h}\sqrt{1+t^3}\,dt$, and $F'(x) = \sqrt{1+x^3}$, so

$$\lim_{h\to 0}\frac{1}{h}\int_2^{2+h}\sqrt{1+t^3}\,dt = F'(2) = \sqrt{1+2^3} = \sqrt{9} = 3.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

77. Let $u = 1 - x$. Then $du = -dx$, so $\int_0^1 f(1-x)\,dx = \int_1^0 f(u)(-du) = \int_0^1 f(u)\,du = \int_0^1 f(x)\,dx$.

78. $\lim\limits_{n\to\infty} \dfrac{1}{n}\left[\left(\dfrac{1}{n}\right)^9 + \left(\dfrac{2}{n}\right)^9 + \left(\dfrac{3}{n}\right)^9 + \cdots + \left(\dfrac{n}{n}\right)^9\right] = \lim\limits_{n\to\infty} \dfrac{1-0}{n} \displaystyle\sum_{i=1}^{n} \left(\dfrac{i}{n}\right)^9 = \int_0^1 x^9\,dx = \left[\dfrac{x^{10}}{10}\right]_0^1 = \dfrac{1}{10}$

The limit is based on Riemann sums using right endpoints and subintervals of equal length.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1. Differentiating both sides of the equation $x \sin \pi x = \int_0^{x^2} f(t)\,dt$ (using FTC1 and the Chain Rule for the right side) gives $\sin \pi x + \pi x \cos \pi x = 2x f(x^2)$. Letting $x = 2$ so that $f(x^2) = f(4)$, we obtain $\sin 2\pi + 2\pi \cos 2\pi = 4f(4)$, so $f(4) = \frac{1}{4}(0 + 2\pi \cdot 1) = \frac{\pi}{2}$.

2. The shaded region has area $\int_0^1 f(x)\,dx = \frac{1}{3}$. The integral $\int_0^1 f^{-1}(y)\,dy$ gives the area of the unshaded region, which we know to be $1 - \frac{1}{3} = \frac{2}{3}$. So $\int_0^1 f^{-1}(y)\,dy = \frac{2}{3}$.

3. For $I = \int_0^4 x e^{(x-2)^4}\,dx$, let $u = x - 2$ so that $x = u + 2$ and $dx = du$. Then

$I = \int_{-2}^{2} (u+2) e^{u^4}\,du = \int_{-2}^{2} u e^{u^4}\,du + \int_{-2}^{2} 2e^{u^4}\,du = 0$ [by 5.5.7(b)] $+ 2\int_0^4 e^{(x-2)^4}\,dx = 2k$.

4. (a)

From the graph of $f(x) = \dfrac{2cx - x^2}{c^3}$, it appears that the areas are equal; that is, the area enclosed is independent of c.

(b) We first find the x-intercepts of the curve, to determine the limits of integration: $y = 0 \iff 2cx - x^2 = 0 \iff x = 0$ or $x = 2c$. Now we integrate the function between these limits to find the enclosed area:

$$A = \int_0^{2c} \frac{2cx - x^2}{c^3}\,dx = \frac{1}{c^3}\left[cx^2 - \tfrac{1}{3}x^3\right]_0^{2c} = \frac{1}{c^3}\left[c(2c)^2 - \tfrac{1}{3}(2c)^3\right] = \frac{1}{c^3}\left[4c^3 - \tfrac{8}{3}c^3\right] = \tfrac{4}{3}, \text{ a constant.}$$

(c)

The vertices of the family of parabolas seem to determine a branch of a hyperbola.

(d) For a particular c, the vertex is the point where the maximum occurs. We have seen that the x-intercepts are 0 and $2c$, so by symmetry, the maximum occurs at $x = c$, and its value is $\dfrac{2c(c) - c^2}{c^3} = \dfrac{1}{c}$. So we are interested in the curve consisting of all points of the form $\left(c, \dfrac{1}{c}\right)$, $c > 0$. This is the part of the hyperbola $y = 1/x$ lying in the first quadrant.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $f(x) = \displaystyle\int_0^{g(x)} \frac{1}{\sqrt{1+t^3}}\,dt$, where $g(x) = \displaystyle\int_0^{\cos x} [1+\sin(t^2)]\,dt$. Using FTC1 and the Chain Rule (twice) we have

$f'(x) = \dfrac{1}{\sqrt{1+[g(x)]^3}}\,g'(x) = \dfrac{1}{\sqrt{1+[g(x)]^3}}\,[1+\sin(\cos^2 x)](-\sin x)$. Now $g\left(\frac{\pi}{2}\right) = \displaystyle\int_0^0 [1+\sin(t^2)]\,dt = 0$, so

$f'\left(\frac{\pi}{2}\right) = \dfrac{1}{\sqrt{1+0}}\,(1+\sin 0)(-1) = 1\cdot 1\cdot(-1) = -1$.

6. If $f(x) = \int_0^x x^2\sin(t^2)\,dt = x^2\int_0^x \sin(t^2)\,dt$, then $f'(x) = x^2\sin(x^2) + 2x\int_0^x \sin(t^2)\,dt$, by the Product Rule and FTC1.

7. As in Exercise 5.3.82, assume that the integrand is defined at $t=0$ so that it is continuous there. By l'Hospital's Rule and the Fundamental Theorem, using the notation $\exp(y) = e^y$,

$$\lim_{x\to 0}\frac{\int_0^x (1-\tan 2t)^{1/t}\,dt}{x} \overset{\text{H}}{=} \lim_{x\to 0}\frac{(1-\tan 2x)^{1/x}}{1} = \exp\left(\lim_{x\to 0}\frac{\ln(1-\tan 2x)}{x}\right)$$

$$\overset{\text{H}}{=} \exp\left(\lim_{x\to 0}\frac{-2\sec^2 2x}{1-\tan 2x}\right) = \exp\left(\frac{-2\cdot 1^2}{1-0}\right) = e^{-2}$$

8. The area $A(t) = \int_0^t \sin(x^2)\,dx$, and the area $B(t) = \frac{1}{2}t\sin(t^2)$. Since $\lim\limits_{t\to 0^+} A(t) = 0 = \lim\limits_{t\to 0^+} B(t)$, we can use l'Hospital's Rule:

$$\lim_{t\to 0^+}\frac{A(t)}{B(t)} \overset{\text{H}}{=} \lim_{t\to 0^+}\frac{\sin(t^2)}{\frac{1}{2}\sin(t^2)+\frac{1}{2}t[2t\cos(t^2)]} \qquad \text{[by FTC1 and the Product Rule]}$$

$$\overset{\text{H}}{=} \lim_{t\to 0^+}\frac{2t\cos(t^2)}{t\cos(t^2)-2t^3\sin(t^2)+2t\cos(t^2)} = \lim_{t\to 0^+}\frac{2\cos(t^2)}{3\cos(t^2)-2t^2\sin(t^2)} = \frac{2}{3-0} = \frac{2}{3}$$

9. $f(x) = 2+x-x^2 = (-x+2)(x+1) = 0 \quad\Leftrightarrow\quad x=2$ or $x=-1$. $f(x)\ge 0$ for $x\in[-1,2]$ and $f(x)<0$ everywhere else. The integral $\int_a^b (2+x-x^2)\,dx$ has a maximum on the interval where the integrand is positive, which is $[-1,2]$. So $a=-1$, $b=2$. (Any larger interval gives a smaller integral since $f(x)<0$ outside $[-1,2]$. Any smaller interval also gives a smaller integral since $f(x)\ge 0$ in $[-1,2]$.)

10. This sum can be interpreted as a Riemann sum, with the right endpoints of the subintervals as sample points and with $a=0$, $b=10{,}000$, and $f(x)=\sqrt{x}$. So we approximate

$\sum\limits_{i=1}^{10{,}000}\sqrt{i} \approx \lim\limits_{n\to\infty}\frac{10{,}000}{n}\sum\limits_{i=1}^{n}\sqrt{\frac{10{,}000i}{n}} = \int_0^{10{,}000}\sqrt{x}\,dx = \left[\frac{2}{3}x^{3/2}\right]_0^{10{,}000} = \frac{2}{3}(1{,}000{,}000) \approx 666{,}667$.

Alternate method: We can use graphical methods as follows:

From the figure we see that $\int_{i-1}^{i}\sqrt{x}\,dx < \sqrt{i} < \int_i^{i+1}\sqrt{x}\,dx$, so

$\int_0^{10{,}000}\sqrt{x}\,dx < \sum\limits_{i=1}^{10{,}000}\sqrt{i} < \int_1^{10{,}001}\sqrt{x}\,dx$. Since

$\int\sqrt{x}\,dx = \frac{2}{3}x^{3/2}+C$, we get $\int_0^{10{,}000}\sqrt{x}\,dx = 666{,}666.\overline{6}$ and

$\int_1^{10{,}001}\sqrt{x}\,dx = \frac{2}{3}[(10{,}001)^{3/2}-1] \approx 666{,}766$.

Hence, $666{,}666.\overline{6} < \sum\limits_{i=1}^{10{,}000}\sqrt{i} < 666{,}766$. We can estimate the sum by averaging these bounds:

$\sum\limits_{i=1}^{10{,}000} \approx \dfrac{666{,}666.\overline{6}+666{,}766}{2} \approx 666{,}716$. The actual value is about 666,716.46.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. (a) We can split the integral $\int_0^n [\![x]\!]\,dx$ into the sum $\sum\limits_{i=1}^{n}\left[\int_{i-1}^{i}[\![x]\!]\,dx\right]$. But on each of the intervals $[i-1, i)$ of integration, $[\![x]\!]$ is a constant function, namely $i-1$. So the ith integral in the sum is equal to $(i-1)[i-(i-1)] = (i-1)$. So the original integral is equal to $\sum\limits_{i=1}^{n}(i-1) = \sum\limits_{i=1}^{n-1} i = \dfrac{(n-1)n}{2}$.

(b) We can write $\int_a^b [\![x]\!]\,dx = \int_0^b [\![x]\!]\,dx - \int_0^a [\![x]\!]\,dx$.

Now $\int_0^b [\![x]\!]\,dx = \int_0^{[\![b]\!]} [\![x]\!]\,dx + \int_{[\![b]\!]}^b [\![x]\!]\,dx$. The first of these integrals is equal to $\frac{1}{2}([\![b]\!]-1)\,[\![b]\!]$, by part (a), and since $[\![x]\!] = [\![b]\!]$ on $[[\![b]\!], b]$, the second integral is just $[\![b]\!]\,(b-[\![b]\!])$. So

$\int_0^b [\![x]\!]\,dx = \frac{1}{2}([\![b]\!]-1)\,[\![b]\!] + [\![b]\!]\,(b-[\![b]\!]) = \frac{1}{2}\,[\![b]\!]\,(2b-[\![b]\!]-1)$ and similarly $\int_0^a [\![x]\!]\,dx = \frac{1}{2}\,[\![a]\!]\,(2a-[\![a]\!]-1)$.

Therefore, $\int_a^b [\![x]\!]\,dx = \frac{1}{2}\,[\![b]\!]\,(2b-[\![b]\!]-1) - \frac{1}{2}\,[\![a]\!]\,(2a-[\![a]\!]-1)$.

12. By FTC1, $\dfrac{d}{dx}\displaystyle\int_0^x\left(\int_1^{\sin t}\sqrt{1+u^4}\,du\right)dt = \int_1^{\sin x}\sqrt{1+u^4}\,du$. Again using FTC1,

$$\frac{d^2}{dx^2}\int_0^x\left(\int_1^{\sin t}\sqrt{1+u^4}\,du\right)dt = \frac{d}{dx}\int_1^{\sin x}\sqrt{1+u^4}\,du = \sqrt{1+\sin^4 x}\cos x.$$

13. Let $Q(x) = \displaystyle\int_0^x P(t)\,dt = \left[at + \frac{b}{2}t^2 + \frac{c}{3}t^3 + \frac{d}{4}t^4\right]_0^x = ax + \frac{b}{2}x^2 + \frac{c}{3}x^3 + \frac{d}{4}x^4$. Then $Q(0) = 0$, and $Q(1) = 0$ by the given condition, $a + \dfrac{b}{2} + \dfrac{c}{3} + \dfrac{d}{4} = 0$. Also, $Q'(x) = P(x) = a + bx + cx^2 + dx^3$ by FTC1. By Rolle's Theorem, applied to Q on $[0, 1]$, there is a number r in $(0, 1)$ such that $Q'(r) = 0$, that is, such that $P(r) = 0$. Thus, the equation $P(x) = 0$ has a solution between 0 and 1.

More generally, if $P(x) = a_0 + a_1x + a_2x^2 + \cdots + a_nx^n$ and if $a_0 + \dfrac{a_1}{2} + \dfrac{a_2}{3} + \cdots + \dfrac{a_n}{n+1} = 0$, then the equation $P(x) = 0$ has a solution between 0 and 1. The proof is the same as before:

Let $Q(x) = \displaystyle\int_0^x P(t)\,dt = a_0x + \frac{a_1}{2}x^2 + \frac{a_2}{3}x^3 + \cdots + \frac{a_n}{n+1}x^n$. Then $Q(0) = Q(1) = 0$ and $Q'(x) = P(x)$. By Rolle's Theorem applied to Q on $[0, 1]$, there is a number r in $(0, 1)$ such that $Q'(r) = 0$, that is, such that $P(r) = 0$.

14.

Let x be the distance between the center of the disk and the surface of the liquid. The wetted circular region has area $\pi r^2 - \pi x^2$ while the unexposed wetted region (shaded in the diagram) has area $2\int_x^r \sqrt{r^2-t^2}\,dt$, so the exposed wetted region has area $A(x) = \pi r^2 - \pi x^2 - 2\int_x^r \sqrt{r^2-t^2}\,dt$, $0 \le x \le r$. By FTC1, we have $A'(x) = -2\pi x + 2\sqrt{r^2-x^2}$.

Now $A'(x) > 0 \;\Rightarrow\; -2\pi x + 2\sqrt{r^2-x^2} > 0 \;\Rightarrow\; \sqrt{r^2-x^2} > \pi x \;\Rightarrow\; r^2 - x^2 > \pi^2x^2 \;\Rightarrow$

$r^2 > \pi^2x^2 + x^2 \;\Rightarrow\; r^2 > x^2(\pi^2+1) \;\Rightarrow\; x^2 < \dfrac{r^2}{\pi^2+1} \;\Rightarrow\; x < \dfrac{r}{\sqrt{\pi^2+1}}$, and we'll call this value x^*.

Since $A'(x) > 0$ for $0 < x < x^*$ and $A'(x) < 0$ for $x^* < x < r$, we have an absolute maximum when $x = x^*$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. Note that $\dfrac{d}{dx}\left(\displaystyle\int_0^x \left[\int_0^u f(t)\,dt\right] du\right) = \displaystyle\int_0^x f(t)\,dt$ by FTC1, while

$$\frac{d}{dx}\left[\int_0^x f(u)(x-u)\,du\right] = \frac{d}{dx}\left[x\int_0^x f(u)\,du\right] - \frac{d}{dx}\left[\int_0^x f(u)u\,du\right]$$
$$= \textstyle\int_0^x f(u)\,du + xf(x) - f(x)x = \int_0^x f(u)\,du$$

Hence, $\int_0^x f(u)(x-u)\,du = \int_0^x \left[\int_0^u f(t)\,dt\right] du + C$. Setting $x = 0$ gives $C = 0$.

16. The parabola $y = 4 - x^2$ and the line $y = x + 2$ intersect when $4 - x^2 = x + 2 \;\Leftrightarrow\; x^2 + x - 2 = 0 \;\Leftrightarrow\; (x+2)(x-1) = 0 \;\Leftrightarrow$ $x = -2$ or 1. So the point A is $(-2, 0)$ and B is $(1, 3)$. The slope of the line $y = x + 2$ is 1 and the slope of the parabola $y = 4 - x^2$ at x-coordinate x is $-2x$. These slopes are equal when $x = -\frac{1}{2}$, so the point C is $\left(-\frac{1}{2}, \frac{15}{4}\right)$.

The area $\mathcal{A}_1$ of the parabolic segment is the area under the parabola from $x = -2$ to $x = 1$, minus the area under the line $y = x + 2$ from -2 to 1. Thus,

$$\mathcal{A}_1 = \textstyle\int_{-2}^1 (4 - x^2)\,dx - \int_{-2}^1 (x+2)\,dx = \left[4x - \frac{1}{3}x^3\right]_{-2}^1 - \left[\frac{1}{2}x^2 + 2x\right]_{-2}^1$$
$$= \left[\left(4 - \tfrac{1}{3}\right) - \left(-8 + \tfrac{8}{3}\right)\right] - \left[\left(\tfrac{1}{2} + 2\right) - (2 - 4)\right] = 9 - \tfrac{9}{2} = \tfrac{9}{2}.$$

The area $\mathcal{A}_2$ of the inscribed triangle is the area under the line segment AC plus the area under the line segment CB minus the area under the line segment AB. The line through A and C has slope $\dfrac{15/4 - 0}{-1/2 + 2} = \dfrac{5}{2}$ and equation $y - 0 = \frac{5}{2}(x + 2)$, or $y = \frac{5}{2}x + 5$. The line through C and B has slope $\dfrac{3 - 15/4}{1 + 1/2} = -\dfrac{1}{2}$ and equation $y - 3 = -\frac{1}{2}(x - 1)$, or $y = -\frac{1}{2}x + \frac{7}{2}$. Thus,

$$\mathcal{A}_2 = \int_{-2}^{-1/2} \left(\tfrac{5}{2}x + 5\right) dx + \int_{-1/2}^{1} \left(-\tfrac{1}{2}x + \tfrac{7}{2}\right) dx - \int_{-2}^{1} (x+2)\,dx = \left[\tfrac{5}{4}x^2 + 5x\right]_{-2}^{-1/2} + \left[-\tfrac{1}{4}x^2 + \tfrac{7}{2}x\right]_{-1/2}^{1} - \tfrac{9}{2}$$
$$= \left[\left(\tfrac{5}{16} - \tfrac{5}{2}\right) - (5 - 10)\right] + \left[\left(-\tfrac{1}{4} + \tfrac{7}{2}\right) - \left(-\tfrac{1}{16} - \tfrac{7}{4}\right)\right] - \tfrac{9}{2} = \tfrac{45}{16} + \tfrac{81}{16} - \tfrac{72}{16} = \tfrac{54}{16} = \tfrac{27}{8}$$

Archimedes' result states that $\mathcal{A}_1 = \frac{4}{3}\mathcal{A}_2$, which is verified in this case since $\frac{4}{3} \cdot \frac{27}{8} = \frac{9}{2}$.

17.

Let c be the nonzero x-intercept so that the parabola has equation $f(x) = kx(x - c)$, or $y = kx^2 - ckx$, where $k < 0$. The area A under the parabola is

$$A = \textstyle\int_0^c kx(x-c)\,dx = k\int_0^c (x^2 - cx)\,dx = k\left[\frac{1}{3}x^3 - \frac{1}{2}cx^2\right]_0^c$$
$$= k\left(\tfrac{1}{3}c^3 - \tfrac{1}{2}c^3\right) = -\tfrac{1}{6}kc^3$$

The point (a, b) is on the parabola, so $f(a) = b \;\Rightarrow\; b = ka(a - c) \;\Rightarrow$ $k = \dfrac{b}{a(a-c)}$. Substituting for k in A gives $A(c) = -\dfrac{b}{6a} \cdot \dfrac{c^3}{a - c} \;\Rightarrow$

$$A'(c) = -\frac{b}{6a} \cdot \frac{(a-c)(3c^2) - c^3(-1)}{(a-c)^2} = -\frac{b}{6a} \cdot \frac{c^2[3(a-c) + c]}{(a-c)^2} = -\frac{bc^2(3a - 2c)}{6a(a-c)^2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Now $A' = 0 \quad \Rightarrow \quad c = \frac{3}{2}a$. Since $A'(c) < 0$ for $a < c < \frac{3}{2}a$ and $A'(c) > 0$ for $c > \frac{3}{2}a$, so A has an absolute minimum when $c = \frac{3}{2}a$. Substituting for c in k gives us $k = \dfrac{b}{a\left(a - \frac{3}{2}a\right)} = -\dfrac{2b}{a^2}$, so $f(x) = -\dfrac{2b}{a^2}\,x\left(x - \frac{3}{2}a\right)$, or $f(x) = -\dfrac{2b}{a^2}\,x^2 + \dfrac{3b}{a}\,x$. Note that the vertex of the parabola is $\left(\frac{3}{4}a, \frac{9}{8}b\right)$ and the minimal area under the parabola is $A\left(\frac{3}{2}a\right) = \frac{9}{8}ab$.

18.

We restrict our attention to the triangle shown. A point in this triangle is closer to the side shown than to any other side, so if we find the area of the region R consisting of all points in the triangle that are closer to the center than to that side, we can multiply this area by 4 to find the total area. We find the equation of the set of points which are equidistant from the center and the side: the distance of the point (x, y) from the side is $1 - y$, and its distance from the center is $\sqrt{x^2 + y^2}$.

So the distances are equal if $\sqrt{x^2 + y^2} = 1 - y \quad \Leftrightarrow \quad x^2 + y^2 = 1 - 2y + y^2 \quad \Leftrightarrow \quad y = \frac{1}{2}(1 - x^2)$. Note that the area we are interested in is equal to the area of a triangle plus a crescent-shaped area. To find these areas, we have to find the y-coordinate h of the horizontal line separating them. From the diagram, $1 - h = \sqrt{2}\,h \quad \Leftrightarrow \quad h = \frac{1}{1 + \sqrt{2}} = \sqrt{2} - 1$. We calculate the areas in terms of h, and substitute afterward.

The area of the triangle is $\frac{1}{2}(2h)(h) = h^2$, and the area of the crescent-shaped section is

$$\int_{-h}^{h} \left[\tfrac{1}{2}(1 - x^2) - h\right] dx = 2\int_0^h \left(\tfrac{1}{2} - h - \tfrac{1}{2}x^2\right) dx = 2\left[\left(\tfrac{1}{2} - h\right)x - \tfrac{1}{6}x^3\right]_0^h = h - 2h^2 - \tfrac{1}{3}h^3.$$

So the area of the whole region is

$$\begin{aligned} 4\left[\left(h - 2h^2 - \tfrac{1}{3}h^3\right) + h^2\right] &= 4h\left(1 - h - \tfrac{1}{3}h^2\right) = 4\left(\sqrt{2} - 1\right)\left[1 - \left(\sqrt{2} - 1\right) - \tfrac{1}{3}\left(\sqrt{2} - 1\right)^2\right] \\ &= 4\left(\sqrt{2} - 1\right)\left(1 - \tfrac{1}{3}\sqrt{2}\right) = \tfrac{4}{3}\left(4\sqrt{2} - 5\right) \end{aligned}$$

19. $\displaystyle \lim_{n\to\infty}\left(\frac{1}{\sqrt{n}\,\sqrt{n+1}} + \frac{1}{\sqrt{n}\,\sqrt{n+2}} + \cdots + \frac{1}{\sqrt{n}\,\sqrt{n+n}}\right)$

$$\begin{aligned} &= \lim_{n\to\infty} \frac{1}{n}\left(\sqrt{\frac{n}{n+1}} + \sqrt{\frac{n}{n+2}} + \cdots + \sqrt{\frac{n}{n+n}}\right) \\ &= \lim_{n\to\infty} \frac{1}{n}\left(\frac{1}{\sqrt{1+1/n}} + \frac{1}{\sqrt{1+2/n}} + \cdots + \frac{1}{\sqrt{1+1}}\right) \\ &= \lim_{n\to\infty} \frac{1}{n}\sum_{i=1}^{n} f\left(\frac{i}{n}\right) \qquad \left[\text{where } f(x) = \frac{1}{\sqrt{1+x}}\right] \\ &= \int_0^1 \frac{1}{\sqrt{1+x}}\,dx = \left[2\sqrt{1+x}\,\right]_0^1 = 2\left(\sqrt{2} - 1\right) \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. Note that the graphs of $(x-c)^2$ and $[(x-c)-2]^2$ intersect when $|x-c| = |x-c-2| \quad \Leftrightarrow$ $c - x = x - c - 2 \quad \Leftrightarrow \quad x = c+1$. The integration will proceed differently depending on the value of c.

Case 1: $-2 \le c < -1$

In this case, $f_c(x) = (x-c-2)^2$ for $x \in [0,1]$, so

$$g(c) = \int_0^1 (x-c-2)^2\,dx = \tfrac{1}{3}\left[(x-c-2)^3\right]_0^1 = \tfrac{1}{3}\left[(-c-1)^3 - (-c-2)^3\right]$$
$$= \tfrac{1}{3}(3c^2+9c+7) = c^2+3c+\tfrac{7}{3} = \left(c+\tfrac{3}{2}\right)^2 + \tfrac{1}{12}$$

This is a parabola; its maximum value for $-2 \le c < -1$ is $g(-2) = \frac{1}{3}$, and its minimum value is $g\left(-\frac{3}{2}\right) = \frac{1}{12}$.

Case 2: $-1 \le c < 0$

In this case, $f_c(x) = \begin{cases} (x-c)^2 & \text{if } 0 \le x \le c+1 \\ (x-c-2)^2 & \text{if } c+1 < x \le 1 \end{cases}$

Therefore,

$$g(c) = \int_0^1 f_c(x)\,dx = \int_0^{c+1} (x-c)^2\,dx + \int_{c+1}^1 (x-c-2)^2\,dx$$
$$= \tfrac{1}{3}\left[(x-c)^3\right]_0^{c+1} + \tfrac{1}{3}\left[(x-c-2)^3\right]_{c+1}^1 = \tfrac{1}{3}\left[1 + c^3 + (-c-1)^3 - (-1)\right]$$
$$= -c^2 - c + \tfrac{1}{3} = -\left(c+\tfrac{1}{2}\right)^2 + \tfrac{7}{12}$$

Again, this is a parabola, whose maximum value for $-1 \le c < 0$ is $g\left(-\frac{1}{2}\right) = \frac{7}{12}$, and whose minimum value on this c-interval is $g(-1) = \frac{1}{3}$.

Case 3: $0 \le c \le 2$

In this case, $f_c(x) = (x-c)^2$ for $x \in [0,1]$, so

$$g(c) = \int_0^1 (x-c)^2\,dx = \tfrac{1}{3}\left[(x-c)^3\right]_0^1 = \tfrac{1}{3}\left[(1-c)^3 - (-c)^3\right]$$
$$= c^2 - c + \tfrac{1}{3} = \left(c-\tfrac{1}{2}\right)^2 + \tfrac{1}{12}$$

This parabola has a maximum value of $g(2) = \frac{7}{3}$ and a minimum value of $g\left(\frac{1}{2}\right) = \frac{1}{12}$.

We conclude that $g(c)$ has an absolute maximum value of $g(2) = \frac{7}{3}$, and absolute minimum values of $g\left(-\frac{3}{2}\right) = g\left(\frac{1}{2}\right) = \frac{1}{12}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6 ☐ APPLICATIONS OF INTEGRATION

6.1 Areas Between Curves

1. (a) $A = \int_{x=0}^{x=2} (y_T - y_B)\,dx = \int_0^2 [(3x - x^2) - x]\,dx = \int_0^2 (2x - x^2)\,dx$

(b) $\int_0^2 (2x - x^2)\,dx = \left[x^2 - \frac{1}{3}x^3\right]_0^2 = \left(4 - \frac{8}{3}\right) - 0 = \frac{4}{3}$

2. (a) $A = \int_{x=0}^{x=1} (y_T - y_B)\,dx = \int_0^1 (e^x - x^2)\,dx$

(b) $\int_0^1 (e^x - x^2)\,dx = \left[e^x - \frac{1}{3}x^3\right]_0^1 = \left(e - \frac{1}{3}\right) - (1 - 0) = e - \frac{4}{3}$

3. (a) $A = \int_{y=-1}^{y=1} (x_R - x_L)\,dy = \int_{-1}^1 [e^y - (y^2 - 2)]\,dy = \int_{-1}^1 (e^y - y^2 + 2)\,dy$

(b) $\int_{-1}^1 (e^y - y^2 + 2)\,dy = \left[e^y - \frac{1}{3}y^3 + 2y\right]_{-1}^1 = \left(e^1 - \frac{1}{3} + 2\right) - \left(e^{-1} + \frac{1}{3} - 2\right) = e - \frac{1}{e} + \frac{10}{3}$

4. (a) $A = \int_{y=0}^{y=3} (x_R - x_L)\,dy - \int_0^3 [(2y - y^2) - (y^2 - 4y)]\,dy = \int_0^3 (-2y^2 + 6y)\,dy$

(b) $\int_0^3 (-2y^2 + 6y)\,dy = \left[-\frac{2}{3}y^3 + 3y^2\right]_0^3 = (-18 + 27) - 0 = 9$

5. $A = \int_{-2}^0 [(x^3 - 3x) - x]\,dx + \int_0^2 [x - (x^3 - 3x)]\,dx \overset{\text{symmetry}}{=} 2\int_0^2 [x - (x^3 - 3x)]\,dx$

$= 2\int_0^2 (4x - x^3)\,dx = 2\left[2x^2 - \frac{1}{4}x^4\right]_0^2 = 2[(8 - 4) - 0] = 8$

6. $A = \int_{-2}^1 \left[\left(\frac{2}{3}x + \frac{16}{3}\right) - x^2\right] dx + \int_1^2 [(-2x + 8) - x^2]\,dx$

$= \left[\frac{1}{3}x^2 + \frac{16}{3}x - \frac{1}{3}x^3\right]_{-2}^1 + \left[-x^2 + 8x - \frac{1}{3}x^3\right]_1^2$

$= \left[\left(\frac{1}{3} + \frac{16}{3} - \frac{1}{3}\right) - \left(\frac{4}{3} - \frac{32}{3} + \frac{8}{3}\right)\right] + \left[\left(-4 + 16 - \frac{8}{3}\right) - \left(-1 + 8 - \frac{1}{3}\right)\right] = \frac{44}{3}$

7. By inspection, we see that the curves intersect at $x = 0$.

$A = \int_0^1 (3^x - 2^x)\,dx$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. By inspection, we see that the curves intersect at $x = 1$.

$$A = \int_1^2 [\ln(x^2) - \ln x]\, dx$$

9. The curves intersect when $2 - x = 2x - x^2 \quad \Leftrightarrow$

$x^2 - 3x + 2 = 0 \quad \Leftrightarrow \quad (x-2)(x-1) = 0 \quad \Leftrightarrow$

$x = 1$ or $x = 2$.

$$A = \int_1^2 [(2x - x^2) - (2 - x)]\, dx$$

10. The curves intersect when $y^4 = 2 - y^2 \quad \Leftrightarrow$

$y^4 + y^2 - 2 = 0 \quad \Leftrightarrow \quad (y^2+2)(y^2-1) = 0 \quad \Leftrightarrow$

$y = \pm 1$.

$$A = \int_{-1}^1 [(2 - y^2) - y^4]\, dy$$

11. $A = \displaystyle\int_0^1 [(x^2+2) - (-x-1)]\, dx$

$= \displaystyle\int_0^1 (x^2 + x + 3)\, dx$

$= \left[\dfrac{1}{3}x^3 + \dfrac{1}{2}x^2 + 3x\right]_0^1 = \left(\dfrac{1}{3} + \dfrac{1}{2} + 3\right) - 0$

$= \dfrac{23}{6}$

12. $A = \displaystyle\int_{-1}^0 [(2 - x) - (1 + x^3)]\, dx$

$= \displaystyle\int_{-1}^0 (1 - x - x^3)\, dx$

$= \left[x - \dfrac{1}{2}x^2 - \dfrac{1}{4}x^4\right]_{-1}^0$

$= \left[0 - \left(-1 - \dfrac{1}{2} - \dfrac{1}{4}\right)\right]$

$= \dfrac{7}{4}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $A = \int_1^2 \left(\frac{1}{x} - \frac{1}{x^2}\right) dx = \left[\ln|x| + \frac{1}{x}\right]_1^2$

$= \left(\ln 2 + \frac{1}{2}\right) - (\ln 1 + 1)$

$= \ln 2 - \frac{1}{2} \approx 0.19$

14. By inspection, we see that the curves intersect at $x = 0$.

$A = \int_0^{\pi/2} (e^x - \cos x)\, dx = \left[e^x - \sin x\right]_0^{\pi/2}$

$= \left(e^{\pi/2} - \sin\frac{\pi}{2}\right) - (e^0 - \sin 0)$

$= [(e^{\pi/2} - 1) - (1 - 0)] = e^{\pi/2} - 2$

15. The curves intersect when $(x-2)^2 = x \quad \Leftrightarrow \quad x^2 - 4x + 4 = x \quad \Leftrightarrow \quad x^2 - 5x + 4 = 0 \quad \Leftrightarrow$ $(x-1)(x-4) = 0 \quad \Leftrightarrow \quad x = 1$ or 4.

$A = \int_1^4 [x - (x-2)^2]\, dx = \int_1^4 (-x^2 + 5x - 4)\, dx$

$= \left[-\frac{1}{3}x^3 + \frac{5}{2}x^2 - 4x\right]_1^4$

$= \left(-\frac{64}{3} + 40 - 16\right) - \left(-\frac{1}{3} + \frac{5}{2} - 4\right)$

$= \frac{9}{2}$

16. The curves intesect when $x^2 - 4x = 2x \quad \Rightarrow \quad x^2 - 6x = 0 \quad \Rightarrow \quad x(x-6) = 0 \quad \Rightarrow \quad x = 0$ or 6.

$A = \int_0^6 [2x - (x^2 - 4x)]\, dx$

$= \int_0^6 (6x - x^2)\, dx = \left[3x^2 - \frac{1}{3}x^3\right]_0^6$

$= \left[3(6)^2 - \frac{1}{3}(6)^3\right] - (0 - 0)$

$= 108 - 72 = 36$

17. The curves intersect when $1 - y^2 = y^2 - 1 \quad \Leftrightarrow \quad 2 = 2y^2 \quad \Leftrightarrow \quad y^2 = 1 \quad \Leftrightarrow \quad y = \pm 1$.

$A = \int_{-1}^1 \left[(1 - y^2) - (y^2 - 1)\right] dy$

$= \int_{-1}^1 2(1 - y^2)\, dy = 2 \cdot 2 \int_0^1 (1 - y^2)\, dy$

$= 4\left[y - \frac{1}{3}y^3\right]_0^1 = 4\left(1 - \frac{1}{3}\right) = \frac{8}{3}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $4x + y^2 = 12$ and $x = y \quad \Rightarrow \quad 4x + x^2 = 12 \quad \Leftrightarrow \quad (x+6)(x-2) = 0 \quad \Leftrightarrow \quad x = -6$ or $x = 2$, so $y = -6$ or $y = 2$ and

$$A = \int_{-6}^{2} \left[\left(-\tfrac{1}{4}y^2 + 3\right) - y\right] dy$$
$$= \left[-\tfrac{1}{12}y^3 - \tfrac{1}{2}y^2 + 3y\right]_{-6}^{2}$$
$$= \left(-\tfrac{2}{3} - 2 + 6\right) - (18 - 18 - 18)$$
$$= 22 - \tfrac{2}{3} = \tfrac{64}{3}$$

19. $12 - x^2 = x^2 - 6 \quad \Leftrightarrow \quad 2x^2 = 18 \quad \Leftrightarrow$
$x^2 = 9 \quad \Leftrightarrow \quad x = \pm 3$, so

$$A = \int_{-3}^{3} \left[(12 - x^2) - (x^2 - 6)\right] dx$$
$$= 2\int_{0}^{3} \left(18 - 2x^2\right) dx \qquad \text{[by symmetry]}$$
$$= 2\left[18x - \tfrac{2}{3}x^3\right]_0^3 = 2\left[(54 - 18) - 0\right]$$
$$= 2(36) = 72$$

20. $x^2 = 4x - x^2 \quad \Leftrightarrow \quad 2x^2 - 4x = 0 \quad \Leftrightarrow$
$2x(x-2) = 0 \quad \Leftrightarrow \quad x = 0$ or 2, so

$$A = \int_{0}^{2} \left[(4x - x^2) - x^2\right] dx = \int_{0}^{2} (4x - 2x^2)\, dx$$
$$= \left[2x^2 - \tfrac{2}{3}x^3\right]_0^2 = 8 - \tfrac{16}{3} = \tfrac{8}{3}$$

21. $2y^2 = 4 + y^2 \quad \Leftrightarrow \quad y^2 = 4 \quad \Leftrightarrow \quad y = \pm 2$, so

$$A = \int_{-2}^{2} \left[(4 + y^2) - 2y^2\right] dy$$
$$= 2\int_{0}^{2} (4 - y^2)\, dy \qquad \text{[by symmetry]}$$
$$= 2\left[4y - \tfrac{1}{3}y^3\right]_0^2 = 2\left(8 - \tfrac{8}{3}\right) = \tfrac{32}{3}$$

22. The curves intersect when $\sqrt{x-1} = x - 1 \quad \Rightarrow$
$x - 1 = x^2 - 2x + 1 \quad \Leftrightarrow \quad 0 = x^2 - 3x + 2 \quad \Leftrightarrow$
$0 = (x-1)(x-2) \quad \Leftrightarrow \quad x = 1$ or 2.

$$A = \int_{1}^{2} \left[\sqrt{x-1} - (x-1)\right] dx$$
$$= \left[\tfrac{2}{3}(x-1)^{3/2} - \tfrac{1}{2}(x-1)^2\right]_1^2 = \left(\tfrac{2}{3} - \tfrac{1}{2}\right) - (0 - 0) = \tfrac{1}{6}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. $\sqrt[3]{2x} = \frac{1}{2}x \quad \Leftrightarrow \quad 2x = \left(\frac{1}{2}x\right)^3 = \frac{1}{8}x^3$

$\Leftrightarrow \quad 16x = x^3 \quad \Leftrightarrow \quad x^3 - 16x = 0$

$\Leftrightarrow \quad x(x^2 - 16) = 0 \quad \Leftrightarrow \quad x = -4, 0, \text{ and } 4$

By symmetry,

$A = 2\int_0^4 \left(\sqrt[3]{2x} - \frac{1}{2}x\right) dx = 2\left[\frac{3}{8}(2x)^{4/3} - \frac{1}{4}x^2\right]_0^4$

$= 2[(6-4) - 0] = 4$

24. The curves intersect when $x^3 = x \quad \Leftrightarrow \quad x^3 - x = 0 \quad \Leftrightarrow$

$x(x^2 - 1) = 0 \quad \Leftrightarrow \quad x(x+1)(x-1) = 0 \quad \Leftrightarrow$

$x = 0$ or $x = \pm 1$.

$A = 2\int_0^1 (x - x^3)\, dx$ [by symmetry]

$= 2\left[\frac{1}{2}x^2 - \frac{1}{4}x^4\right]_0^1 = 2\left(\frac{1}{2} - \frac{1}{4}\right) = \frac{1}{2}$

25. $\sqrt{x} = \frac{1}{3}x \quad \Rightarrow \quad x = \frac{1}{9}x^2 \quad \Leftrightarrow \quad \frac{1}{9}x^2 - x = 0 \quad \Leftrightarrow$

$\frac{1}{9}x(x-9) = 0 \quad \Leftrightarrow \quad x = 0$ or $x = 9$

$A = \int_0^9 \left(\sqrt{x} - \frac{1}{3}x\right) dx + \int_9^{16} \left(\frac{1}{3}x - \sqrt{x}\right) dx$

$= \left[\frac{2}{3}x^{3/2} - \frac{1}{6}x^2\right]_0^9 + \left[\frac{1}{6}x^2 - \frac{2}{3}x^{3/2}\right]_9^{16}$

$= \left[\left(18 - \frac{27}{2}\right) - 0\right] + \left[\left(\frac{128}{3} - \frac{128}{3}\right) - \left(\frac{27}{2} - 18\right)\right] = 9$

26. $A = \int_0^{2\pi} [(2 - \cos x) - \cos x]\, dx$

$= \int_0^{2\pi} (2 - 2\cos x)\, dx$

$= \Big[2x - 2\sin x\Big]_0^{2\pi}$

$= (4\pi - 0) - 0 = 4\pi$

27. $\cos x = \sin 2x = 2\sin x \cos x \quad \Leftrightarrow \quad 2\sin x\cos x - \cos x = 0 \quad \Leftrightarrow \quad \cos x(2\sin x - 1) = 0 \quad \Leftrightarrow$

$\cos x = 0$ or $\sin x = \frac{1}{2} \quad \Leftrightarrow \quad x = \frac{\pi}{2}$ or $x = \frac{\pi}{6}$ on $\left[0, \frac{\pi}{2}\right]$

$A = \int_0^{\pi/6} (\cos x - \sin 2x)\, dx + \int_{\pi/6}^{\pi/2} (\sin 2x - \cos x)\, dx$

$= \left[\sin x + \frac{1}{2}\cos 2x\right]_0^{\pi/6} + \left[-\frac{1}{2}\cos 2x - \sin x\right]_{\pi/6}^{\pi/2}$

$= \left[\left(\sin\frac{\pi}{6} + \frac{1}{2}\cos\frac{\pi}{3}\right) - \left(\sin 0 + \frac{1}{2}\cos 0\right)\right]$

$\quad + \left[\left(-\frac{1}{2}\cos\pi - \sin\frac{\pi}{2}\right) - \left(-\frac{1}{2}\cos\frac{\pi}{3} - \sin\frac{\pi}{6}\right)\right]$

$= \left[\left(\frac{1}{2} + \frac{1}{4}\right) - \left(0 + \frac{1}{2}\right)\right] + \left[\left(\frac{1}{2} - 1\right) - \left(-\frac{1}{4} - \frac{1}{2}\right)\right] = \frac{1}{2}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. The curves intersect when $\cos x = 1 - \cos x$ (on $[0, \pi]$) $\Leftrightarrow$ $2\cos x = 1$ $\Leftrightarrow$ $\cos x = \frac{1}{2}$ $\Leftrightarrow$ $x = \frac{\pi}{3}$.

$$A = \int_0^{\pi/3} [\cos x - (1 - \cos x)]\,dx + \int_{\pi/3}^{\pi} [(1 - \cos x) - \cos x]\,dx$$
$$= \int_0^{\pi/3} (2\cos x - 1)\,dx + \int_{\pi/3}^{\pi} (1 - 2\cos x)\,dx$$
$$= \Big[2\sin x - x\Big]_0^{\pi/3} + \Big[x - 2\sin x\Big]_{\pi/3}^{\pi}$$
$$= \left(\sqrt{3} - \frac{\pi}{3}\right) - 0 + (\pi - 0) - \left(\frac{\pi}{3} - \sqrt{3}\right)$$
$$= 2\sqrt{3} + \frac{\pi}{3}$$

29. The curves intersect when $8\cos x = \sec^2 x$ $\Rightarrow$ $8\cos^3 x = 1$ $\Rightarrow$ $\cos^3 x = \frac{1}{8}$ $\Rightarrow$ $\cos x = \frac{1}{2}$ $\Rightarrow$ $x = \frac{\pi}{3}$ for $0 < x < \frac{\pi}{2}$. By symmetry,

$$A = 2\int_0^{\pi/3} (8\cos x - \sec^2 x)\,dx$$
$$= 2\left[8\sin x - \tan x\right]_0^{\pi/3}$$
$$= 2\left(8 \cdot \tfrac{\sqrt{3}}{2} - \sqrt{3}\right) = 2\left(3\sqrt{3}\right)$$
$$= 6\sqrt{3}$$

30. $x^4 - 3x^2 = x^2$ $\Leftrightarrow$ $x^4 - 4x^2 = 0$ $\Leftrightarrow$ $x^2(x^2 - 4) = 0$ $\Leftrightarrow$ $x = 0, x = \pm 2$

$$A = 2\int_0^2 [x^2 - (x^4 - 3x^2)]\,dx = 2\int_0^2 (4x^2 - x^4)\,dx$$
$$= 2\left[\frac{4}{3}x^3 - \frac{1}{5}x^5\right]_0^2 = 2\left[\left(\frac{32}{3} - \frac{32}{5}\right) - 0\right]$$
$$= \frac{128}{15}$$

31. By inspection, we see that the curves intersect at $x = \pm 1$ and that the area of the region enclosed by the curves is twice the area enclosed in the first quadrant.

$$A = 2\int_0^1 [(2 - x) - x^4]\,dx = 2\left[2x - \tfrac{1}{2}x^2 - \tfrac{1}{5}x^5\right]_0^1$$
$$= 2\left[\left(2 - \tfrac{1}{2} - \tfrac{1}{5}\right) - 0\right] = 2\left(\tfrac{13}{10}\right) = \tfrac{13}{5}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. $x^2 = \dfrac{32}{x^2+4} \quad \Leftrightarrow \quad x^2(x^2+4) = 32 \quad \Leftrightarrow$

$x^4 + 4x^2 - 32 = 0 \quad \Leftrightarrow \quad (x^2+8)(x^2-4) = 0 \quad \Leftrightarrow \quad x = \pm 2$

By symmetry,

$$A = 2\int_0^2 \left(\frac{32}{x^2+4} - x^2\right) dx$$

$$= 2\left[16\arctan\left(\frac{x}{2}\right) - \frac{1}{3}x^3\right]_0^2 = 2\left[\left(16\arctan 1 - \frac{8}{3}\right) - 0\right]$$

$$= 2\left(16 \cdot \frac{\pi}{4} - \frac{8}{3}\right) = 8\pi - \frac{16}{3}$$

33. By inspection, we see that the curves intersect at $x = -1$, 0, and 1.

By symmetry,

$$A = 2\int_0^1 \left[\sin\left(\frac{\pi x}{2}\right) - x^3\right] dx = 2\left[-\frac{2}{\pi}\cos\left(\frac{\pi x}{2}\right) - \frac{1}{4}x^4\right]_0^1$$

$$= 2\left[\left(-\frac{2}{\pi}\cos\frac{\pi}{2} - \frac{1}{4}\right) - \left(-\frac{2}{\pi}\cos 0 - 0\right)\right]$$

$$= 2\left[\left(0 - \frac{1}{4}\right) - \left(-\frac{2}{\pi} - 0\right)\right] = 2\left(\frac{2}{\pi} - \frac{1}{4}\right) = \frac{4}{\pi} - \frac{1}{2}$$

34. $4 - 2\cosh x = \dfrac{1}{2}\sinh x \quad \Leftrightarrow \quad 4 - 2\left(\dfrac{e^x + e^{-x}}{2}\right) = \dfrac{1}{2}\left(\dfrac{e^x - e^{-x}}{2}\right) \quad \Leftrightarrow \quad 4 - e^x - e^{-x} = \dfrac{1}{4}e^x - \dfrac{1}{4}e^{-x} \quad \Leftrightarrow$

$4e^x(4 - e^x - e^{-x}) = 4e^x\left(\dfrac{1}{4}e^x - \dfrac{1}{4}e^{-x}\right) \quad \Leftrightarrow \quad 16e^x - 4e^{2x} - 4 = e^{2x} - 1 \quad \Leftrightarrow \quad 5e^{2x} - 16e^x + 3 = 0 \quad \Leftrightarrow$

$(5e^x - 1)(e^x - 3) = 0 \quad \Leftrightarrow \quad e^x = \frac{1}{5}$ or $e^x = 3 \quad \Leftrightarrow \quad x = \ln\frac{1}{5} = -\ln 5$ or $x = \ln 3$

$$A = \int_{-\ln 5}^{\ln 3} \left[(4 - 2\cosh x) - \frac{1}{2}\sinh x\right] dx = \left[4x - 2\sinh x - \frac{1}{2}\cosh x\right]_{-\ln 5}^{\ln 3}$$

$$= \left[4\ln 3 - 2\left(\frac{e^{\ln 3} - e^{-\ln 3}}{2}\right) - \frac{1}{2}\left(\frac{e^{\ln 3} + e^{-\ln 3}}{2}\right)\right] - \left[-4\ln 5 - 2\left(\frac{e^{-\ln 5} - e^{\ln 5}}{2}\right) - \frac{1}{2}\left(\frac{e^{-\ln 5} + e^{\ln 5}}{2}\right)\right]$$

$$= \left[4\ln 3 - \left(3 - \frac{1}{3}\right) - \frac{1}{4}\left(3 + \frac{1}{3}\right)\right] - \left[-4\ln 5 - \left(\frac{1}{5} - 5\right) - \frac{1}{4}\left(\frac{1}{5} + 5\right)\right]$$

$$= 4\ln 15 - 7$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. $1/x = x \quad \Leftrightarrow \quad 1 = x^2 \quad \Leftrightarrow \quad x = \pm 1$ and $1/x = \frac{1}{4}x \quad \Leftrightarrow$

$4 = x^2 \quad \Leftrightarrow \quad x = \pm 2$, so for $x > 0$,

$$A = \int_0^1 \left(x - \frac{1}{4}x\right) dx + \int_1^2 \left(\frac{1}{x} - \frac{1}{4}x\right) dx$$

$$= \int_0^1 \left(\frac{3}{4}x\right) dx + \int_1^2 \left(\frac{1}{x} - \frac{1}{4}x\right) dx$$

$$= \left[\tfrac{3}{8}x^2\right]_0^1 + \left[\ln|x| - \tfrac{1}{8}x^2\right]_1^2$$

$$= \tfrac{3}{8} + \left(\ln 2 - \tfrac{1}{2}\right) - \left(0 - \tfrac{1}{8}\right) = \ln 2$$

36. $\frac{1}{4}x^2 = -x + 3 \quad \Leftrightarrow \quad x^2 + 4x - 12 = 0 \quad \Leftrightarrow \quad (x+6)(x-2) = 0 \quad \Leftrightarrow \quad x = -6$ or 2 and $2x^2 = -x + 3 \quad \Leftrightarrow$

$2x^2 + x - 3 = 0 \quad \Leftrightarrow \quad (2x+3)(x-1) = 0 \quad \Leftrightarrow \quad x = -\frac{3}{2}$ or 1, so for $x \geq 0$,

$$A = \int_0^1 \left(2x^2 - \tfrac{1}{4}x^2\right) dx + \int_1^2 \left[(-x+3) - \tfrac{1}{4}x^2\right] dx$$

$$= \int_0^1 \tfrac{7}{4}x^2\, dx + \int_1^2 \left(-\tfrac{1}{4}x^2 - x + 3\right) dx$$

$$= \left[\tfrac{7}{12}x^3\right]_0^1 + \left[-\tfrac{1}{12}x^3 - \tfrac{1}{2}x^2 + 3x\right]_1^2$$

$$= \tfrac{7}{12} + \left(-\tfrac{2}{3} - 2 + 6\right) - \left(-\tfrac{1}{12} - \tfrac{1}{2} + 3\right) = \tfrac{3}{2}$$

37. (a) Total area $= 12 + 27 = 39$.

(b) $f(x) \leq g(x)$ for $0 \leq x \leq 2$ and $f(x) \geq g(x)$ for $2 \leq x \leq 5$, so

$$\int_0^5 [f(x) - g(x)]\, dx = \int_0^2 [f(x) - g(x)]\, dx + \int_2^5 [f(x) - g(x)]\, dx$$

$$= -\int_0^2 [g(x) - f(x)]\, dx + \int_2^5 [f(x) - g(x)]\, dx$$

$$= -(12) + 27 = 15$$

38. $\dfrac{x}{\sqrt{1+x^2}} = \dfrac{x}{\sqrt{9-x^2}} \quad \Leftrightarrow \quad x = 0$ or $\sqrt{1+x^2} = \sqrt{9-x^2} \quad \Rightarrow$

$1 + x^2 = 9 - x^2 \quad \Rightarrow \quad 2x^2 = 8 \quad \Rightarrow \quad x^2 = 4 \quad \Rightarrow \quad x = 2 \ (x \geq 0)$.

$$A = \int_0^2 \left(\frac{x}{\sqrt{1+x^2}} - \frac{x}{\sqrt{9-x^2}}\right) dx = \left[\sqrt{1+x^2} + \sqrt{9-x^2}\right]_0^2$$

$$= (\sqrt{5} + \sqrt{5}) - (1 + 3) = 2\sqrt{5} - 4$$

39. $\dfrac{x}{1+x^2} = \dfrac{x^2}{1+x^3} \quad \Leftrightarrow \quad x + x^4 = x^2 + x^4 \quad \Leftrightarrow \quad x = x^2 \quad \Leftrightarrow$

$0 = x^2 - x \quad \Leftrightarrow \quad 0 = x(x-1) \quad \Leftrightarrow \quad x = 0$ or $x = 1$.

$$A = \int_0^1 \left(\frac{x}{1+x^2} - \frac{x^2}{1+x^3}\right) dx = \left[\tfrac{1}{2}\ln(1+x^2) - \tfrac{1}{3}\ln(1+x^3)\right]_0^1$$

$$= \left(\tfrac{1}{2}\ln 2 - \tfrac{1}{3}\ln 2\right) - (0 - 0) = \tfrac{1}{6}\ln 2$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $\dfrac{\ln x}{x} = \dfrac{(\ln x)^2}{x} \quad \Leftrightarrow \quad \ln x = (\ln x)^2 \quad \Leftrightarrow \quad 0 = (\ln x)^2 - \ln x \quad \Leftrightarrow$

$0 = \ln x(\ln x - 1) \quad \Leftrightarrow \quad \ln x = 0 \text{ or } 1 \quad \Leftrightarrow \quad x = e^0 \text{ or } e^1 \ \ [1 \text{ or } e]$

$$A = \int_1^e \left[\frac{\ln x}{x} - \frac{(\ln x)^2}{x}\right] dx = \left[\tfrac{1}{2}(\ln x)^2 - \tfrac{1}{3}(\ln x)^3\right]_1^e$$

$$= \left(\tfrac{1}{2} - \tfrac{1}{3}\right) - (0 - 0) = \tfrac{1}{6}$$

41. An equation of the line through $(0, 0)$ and $(3, 1)$ is $y = \frac{1}{3}x$; through $(0, 0)$ and $(1, 2)$ is $y = 2x$; through $(3, 1)$ and $(1, 2)$ is $y = -\frac{1}{2}x + \frac{5}{2}$.

$$A = \int_0^1 \left(2x - \tfrac{1}{3}x\right) dx + \int_1^3 \left[\left(-\tfrac{1}{2}x + \tfrac{5}{2}\right) - \tfrac{1}{3}x\right] dx$$

$$= \int_0^1 \tfrac{5}{3}x\, dx + \int_1^3 \left(-\tfrac{5}{6}x + \tfrac{5}{2}\right) dx = \left[\tfrac{5}{6}x^2\right]_0^1 + \left[-\tfrac{5}{12}x^2 + \tfrac{5}{2}x\right]_1^3$$

$$= \tfrac{5}{6} + \left(-\tfrac{15}{4} + \tfrac{15}{2}\right) - \left(-\tfrac{5}{12} + \tfrac{5}{2}\right) = \tfrac{5}{2}$$

42. An equation of the line through $(2, 0)$ and $(0, 2)$ is $y = -x + 2$; through $(2, 0)$ and $(-1, 1)$ is $y = -\frac{1}{3}x + \frac{2}{3}$; through $(0, 2)$ and $(-1, 1)$ is $y = x + 2$.

$$A = \int_{-1}^0 \left[(x + 2) - \left(-\tfrac{1}{3}x + \tfrac{2}{3}\right)\right] dx + \int_0^2 \left[(-x + 2) - \left(-\tfrac{1}{3}x + \tfrac{2}{3}\right)\right] dx$$

$$= \int_{-1}^0 \left(\tfrac{4}{3}x + \tfrac{4}{3}\right) dx + \int_0^2 \left(-\tfrac{2}{3}x + \tfrac{4}{3}\right) dx$$

$$= \left[\tfrac{2}{3}x^2 + \tfrac{4}{3}x\right]_{-1}^0 + \left[-\tfrac{1}{3}x^2 + \tfrac{4}{3}x\right]_0^2$$

$$= 0 - \left(\tfrac{2}{3} - \tfrac{4}{3}\right) + \left(-\tfrac{4}{3} + \tfrac{8}{3}\right) - 0 = 2$$

43. The curves intersect when $\sin x = \cos 2x$ (on $[0, \pi/2]$) $\Leftrightarrow$ $\sin x = 1 - 2\sin^2 x$ $\Leftrightarrow$ $2\sin^2 x + \sin x - 1 = 0$ $\Leftrightarrow$ $(2\sin x - 1)(\sin x + 1) = 0$ $\Rightarrow$ $\sin x = \frac{1}{2}$ $\Rightarrow$ $x = \frac{\pi}{6}$.

$$A = \int_0^{\pi/2} |\sin x - \cos 2x|\, dx$$

$$= \int_0^{\pi/6} (\cos 2x - \sin x)\, dx + \int_{\pi/6}^{\pi/2} (\sin x - \cos 2x)\, dx$$

$$= \left[\tfrac{1}{2}\sin 2x + \cos x\right]_0^{\pi/6} + \left[-\cos x - \tfrac{1}{2}\sin 2x\right]_{\pi/6}^{\pi/2}$$

$$= \left(\tfrac{1}{4}\sqrt{3} + \tfrac{1}{2}\sqrt{3}\right) - (0 + 1) + (0 - 0) - \left(-\tfrac{1}{2}\sqrt{3} - \tfrac{1}{4}\sqrt{3}\right)$$

$$= \tfrac{3}{2}\sqrt{3} - 1$$

44. $$A = \int_{-1}^1 |3^x - 2^x|\, dx = \int_{-1}^0 (2^x - 3^x)\, dx + \int_0^1 (3^x - 2^x)\, dx$$

$$= \left[\frac{2^x}{\ln 2} - \frac{3^x}{\ln 3}\right]_{-1}^0 + \left[\frac{3^x}{\ln 3} - \frac{2^x}{\ln 2}\right]_0^1$$

$$= \left(\frac{1}{\ln 2} - \frac{1}{\ln 3}\right) - \left(\frac{1}{2\ln 2} - \frac{1}{3\ln 3}\right) + \left(\frac{3}{\ln 3} - \frac{2}{\ln 2}\right) - \left(\frac{1}{\ln 3} - \frac{1}{\ln 2}\right)$$

$$= \frac{2 - 1 - 4 + 2}{2\ln 2} + \frac{-3 + 1 + 9 - 3}{3\ln 3} = \frac{4}{3\ln 3} - \frac{1}{2\ln 2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45.

From the graph, we see that the curves intersect at $x = 0$ and $x = a \approx 0.896$, with $x\sin(x^2) > x^4$ on $(0, a)$. So the area A of the region bounded by the curves is

$$A = \int_0^a \left[x\sin(x^2) - x^4\right] dx = \left[-\tfrac{1}{2}\cos(x^2) - \tfrac{1}{5}x^5\right]_0^a$$
$$= -\tfrac{1}{2}\cos(a^2) - \tfrac{1}{5}a^5 + \tfrac{1}{2} \approx 0.037$$

46. From the graph, we see that the curves intersect (with $x \geq 0$) at $x = 0$ and $x = a$, where $a \approx 1.052$, with $x/(x^2+1)^2 > x^5 - x$ on $(0, a)$. The area A of the region bounded by the curves is

$$A = \int_0^a \left[\frac{x}{(x^2+1)^2} - (x^5 - x)\right] dx = \left[-\frac{1}{2}\cdot\frac{1}{x^2+1} - \frac{1}{6}x^6 + \frac{1}{2}x^2\right]_0^a$$
$$\approx 0.59$$

47.

From the graph, we see that the curves intersect at $x = a \approx -1.11$, $x = b \approx 1.25$, and $x = c \approx 2.86$, with $x^3 - 3x + 4 > 3x^2 - 2x$ on (a, b) and $3x^2 - 2x > x^3 - 3x + 4$ on (b, c). So the area of the region bounded by the curves is

$$A = \int_a^b \left[(x^3 - 3x + 4) - (3x^2 - 2x)\right] dx + \int_b^c \left[(3x^2 - 2x) - (x^3 - 3x + 4)\right] dx$$
$$= \int_a^b (x^3 - 3x^2 - x + 4)\, dx + \int_b^c (-x^3 + 3x^2 + x - 4)\, dx$$
$$= \left[\tfrac{1}{4}x^4 - x^3 - \tfrac{1}{2}x^2 + 4x\right]_a^b + \left[-\tfrac{1}{4}x^4 + x^3 + \tfrac{1}{2}x^2 - 4x\right]_b^c \approx 8.38$$

48.

From the graph, we see that the curves intersect at $x = a \approx 0.29$ and $x = b \approx 6.08$. $y = 2\sqrt{x}$ is the upper curve, so the area of the region bounded by the curves is

$$A \approx \int_a^b \left(2\sqrt{x} - 1.3^x\right) dx = \left[\tfrac{4}{3}x^{3/2} - \frac{1}{\ln 1.3}\, 1.3^x\right]_a^b \approx 5.11$$

49.

Graph `Y1=2/(1+x^4)` and `Y2=x^2`. We see that `Y1` > `Y2` on $(-1, 1)$, so the area is given by $\displaystyle\int_{-1}^{1} \left(\frac{2}{1+x^4} - x^2\right) dx$. Evaluate the integral with a command such as `fnInt(Y1-Y2,x,-1,1)` to get 2.80123 to five decimal places.

Another method: Graph $f(x) =$ `Y1=2/(1+x^4)-x^2` and from the graph evaluate $\int f(x)\, dx$ from -1 to 1.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50.

The curves intersect at $x = \pm 1$.

$$A = \int_{-1}^{1} (e^{1-x^2} - x^4)\, dx \approx 3.66016$$

51.

The curves intersect at $x = 0$ and $x = a \approx 0.749363$.

$$A = \int_0^a \left(\sqrt{x} - \tan^2 x\right)\, dx \approx 0.25142$$

52.

The curves intersect at $x = a \approx -1.911917$, $x = b \approx -1.223676$, and $x = c \approx 0.607946$.

$$A = \int_a^b \left[(x + 2\sin^4 x) - \cos x\right] dx + \int_b^c \left[\cos x - (x + 2\sin^4 x)\right] dx$$
$$\approx 1.70413$$

53. As the figure illustrates, the curves $y = x$ and $y = x^5 - 6x^3 + 4x$ enclose a four-part region symmetric about the origin (since $x^5 - 6x^3 + 4x$ and x are odd functions of x). The curves intersect at values of x where $x^5 - 6x^3 + 4x = x$; that is, where $x(x^4 - 6x^2 + 3) = 0$. That happens at $x = 0$ and where

$x^2 = \dfrac{6 \pm \sqrt{36 - 12}}{2} = 3 \pm \sqrt{6}$; that is, at $x = -\sqrt{3+\sqrt{6}}, -\sqrt{3-\sqrt{6}}, 0, \sqrt{3-\sqrt{6}}$, and $\sqrt{3+\sqrt{6}}$. The exact area is

$$\begin{aligned} 2\int_0^{\sqrt{3+\sqrt{6}}} \left|(x^5 - 6x^3 + 4x) - x\right| dx &= 2\int_0^{\sqrt{3+\sqrt{6}}} \left|x^5 - 6x^3 + 3x\right| dx \\ &= 2\int_0^{\sqrt{3-\sqrt{6}}} (x^5 - 6x^3 + 3x)\, dx + 2\int_{\sqrt{3-\sqrt{6}}}^{\sqrt{3+\sqrt{6}}} (-x^5 + 6x^3 - 3x)\, dx \\ &\overset{\text{CAS}}{=} 12\sqrt{6} - 9 \end{aligned}$$

54. The inequality $x \geq 2y^2$ describes the region that lies on, or to the right of, the parabola $x = 2y^2$. The inequality $x \leq 1 - |y|$ describes the region that lies on, or to the left of, the curve $x = 1 - |y| = \begin{cases} 1 - y & \text{if } y \geq 0 \\ 1 + y & \text{if } y < 0 \end{cases}$.

So the given region is the shaded region that lies between the curves.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The graphs of $x = 1 - y$ and $x = 2y^2$ intersect when $1 - y = 2y^2 \quad \Leftrightarrow$

$2y^2 + y - 1 = 0 \quad \Leftrightarrow \quad (2y-1)(y+1) = 0 \quad \Rightarrow \quad y = \frac{1}{2}$ [for $y \geq 0$]. By symmetry,

$A = 2\int_0^{1/2} \left[(1-y) - 2y^2\right] dy = 2\left[-\frac{2}{3}y^3 - \frac{1}{2}y^2 + y\right]_0^{1/2} = 2\left[\left(-\frac{1}{12} - \frac{1}{8} + \frac{1}{2}\right) - 0\right] = 2\left(\frac{7}{24}\right) = \frac{7}{12}.$

55. 1 second $= \frac{1}{3600}$ hour, so 10 s $= \frac{1}{360}$ h. With the given data, we can take $n = 5$ to use the Midpoint Rule.

$\Delta t = \frac{1/360 - 0}{5} = \frac{1}{1800}$, so

$$\begin{aligned}
\text{distance}_{\text{Kelly}} - \text{distance}_{\text{Chris}} &= \int_0^{1/360} v_K\, dt - \int_0^{1/360} v_C\, dt = \int_0^{1/360} (v_K - v_C)\, dt \\
&\approx M_5 = \frac{1}{1800}\left[(v_K - v_C)(1) + (v_K - v_C)(3) + (v_K - v_C)(5)\right. \\
&\qquad \left. + (v_K - v_C)(7) + (v_K - v_C)(9)\right] \\
&= \frac{1}{1800}[(22-20) + (52-46) + (71-62) + (86-75) + (98-86)] \\
&= \frac{1}{1800}(2 + 6 + 9 + 11 + 12) = \frac{1}{1800}(40) = \frac{1}{45} \text{ mile, or } 117\tfrac{1}{3} \text{ feet}
\end{aligned}$$

56. If $x =$ distance from left end of pool and $w = w(x) =$ width at x, then the Midpoint Rule with $n = 4$ and

$\Delta x = \dfrac{b-a}{n} = \dfrac{8 \cdot 2 - 0}{4} = 4$ gives Area $= \int_0^{16} w\, dx \approx 4(6.2 + 6.8 + 5.0 + 4.8) = 4(22.8) = 91.2 \text{ m}^2$.

57. Let $h(x)$ denote the height of the wing at x cm from the left end.

$$\begin{aligned}
A \approx M_5 &= \frac{200-0}{5}\left[h(20) + h(60) + h(100) + h(140) + h(180)\right] \\
&= 40(20.3 + 29.0 + 27.3 + 20.5 + 8.7) = 40(105.8) = 4232 \text{ cm}^2
\end{aligned}$$

58. For $0 \leq t \leq 10$, $b(t) > d(t)$, so the area between the curves is given by

$$\begin{aligned}
\int_0^{10} [b(t) - d(t)]\, dt &= \int_0^{10} (2200e^{0.024t} - 1460e^{0.018t})\, dt = \left[\frac{2200}{0.024}e^{0.024t} - \frac{1460}{0.018}e^{0.018t}\right]_0^{10} \\
&= \left(\frac{275{,}000}{3}e^{0.24} - \frac{730{,}000}{9}e^{0.18}\right) - \left(\frac{275{,}000}{3} - \frac{730{,}000}{9}\right) \approx 8868 \text{ people}
\end{aligned}$$

This area A represents the increase in population over a 10-year period.

59. (a) From Example 8(a), the infectiousness concentration is 1210 cells/mL. $g(t) = 1210 \quad \Leftrightarrow \quad 0.9f(t) = 1210 \quad \Leftrightarrow$ $0.9(-t)(t-21)(t+1) = 1210$. Using a calculator to solve the last equation for $t > 0$ gives us two solutions with the lesser being $t = t_3 \approx 11.26$ days, or the 12th day.

(b) From Example 8(b), the slope of the line through P_1 and P_2 is -23. From part (a), $P_3 = (t_3, 1210)$. An equation of the line through P_3 that is parallel to $\overline{P_1P_2}$ is $N - 1210 = -23(t - t_3)$, or $N = -23t + 23t_3 + 1210$. Using a calculator, we find that this line intersects g at $t = t_4 \approx 17.18$, or the 18th day. So in the patient with some immunity, the infection lasts about 2 days less than in the patient without immunity.

(c) The level of infectiousness for this patient is the area between the graph of g and the line in part (b). This area is

$$\begin{aligned}
\int_{t_3}^{t_4} \left[g(t) - (-23t + 23t_3 + 1210)\right] dt &\approx \int_{11.26}^{17.18} (-0.9t^3 + 18t^2 + 41.9t - 1468.94)\, dt \\
&= \left[-0.225t^4 + 6t^3 + 20.95t^2 - 1468.94t\right]_{11.26}^{17.18} \approx 706 \text{ (cells/mL)} \cdot \text{days}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. From the figure, $g(t) > f(t)$ for $0 \le t \le 2$. The area between the curves is given by

$$\begin{aligned}\int_0^2 [g(t) - f(t)]\,dt &= \int_0^2 [(0.17t^2 - 0.5t + 1.1) - (0.73t^3 - 2t^2 + t + 0.6)]\,dt \\ &= \int_0^2 (-0.73t^3 + 2.17t^2 - 1.5t + 0.5)\,dt \\ &= \left[-\frac{0.73}{4}t^4 + \frac{2.17}{3}t^3 - 0.75t^2 + 0.5t\right]_0^2 \\ &= -2.92 + \frac{17.36}{3} - 3 + 1 - 0 = 0.8\overline{6} \approx 0.87\end{aligned}$$

Thus, about 0.87 more inches of rain fell at the second location than at the first during the first two hours of the storm.

61. We know that the area under curve A between $t = 0$ and $t = x$ is $\int_0^x v_A(t)\,dt = s_A(x)$, where $v_A(t)$ is the velocity of car A and s_{A} is its displacement. Similarly, the area under curve B between $t = 0$ and $t = x$ is $\int_0^x v_{\mathrm{B}}(t)\,dt = s_{\mathrm{B}}(x)$.

(a) After one minute, the area under curve A is greater than the area under curve B. So car A is ahead after one minute.

(b) The area of the shaded region has numerical value $s_{\mathrm{A}}(1) - s_{\mathrm{B}}(1)$, which is the distance by which A is ahead of B after 1 minute.

(c) After two minutes, car B is traveling faster than car A and has gained some ground, but the area under curve A from $t = 0$ to $t = 2$ is still greater than the corresponding area for curve B, so car A is still ahead.

(d) From the graph, it appears that the area between curves A and B for $0 \le t \le 1$ (when car A is going faster), which corresponds to the distance by which car A is ahead, seems to be about 3 squares. Therefore, the cars will be side by side at the time x where the area between the curves for $1 \le t \le x$ (when car B is going faster) is the same as the area for $0 \le t \le 1$. From the graph, it appears that this time is $x \approx 2.2$. So the cars are side by side when $t \approx 2.2$ minutes.

62. The area under $R'(x)$ from $x = 50$ to $x = 100$ represents the change in revenue, and the area under $C'(x)$ from $x = 50$ to $x = 100$ represents the change in cost. The shaded region represents the difference between these two values; that is, the increase in profit as the production level increases from 50 units to 100 units. We use the Midpoint Rule with $n = 5$ and $\Delta x = 10$:

$$\begin{aligned}M_5 &= \Delta x\{[R'(55) - C'(55)] + [R'(65) - C'(65)] + [R'(75) - C'(75)] + [R'(85) - C'(85)] + [R'(95) - C'(95)]\} \\ &\approx 10(2.40 - 0.85 + 2.20 - 0.90 + 2.00 - 1.00 + 1.80 - 1.10 + 1.70 - 1.20) \\ &= 10(5.05) = 50.5 \text{ thousand dollars}\end{aligned}$$

Using M_1 would give us $50(2 - 1) = 50$ thousand dollars.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

63.

To graph this function, we must first express it as a combination of explicit functions of y; namely, $y = \pm x\sqrt{x+3}$. We can see from the graph that the loop extends from $x = -3$ to $x = 0$, and that by symmetry, the area we seek is just twice the area under the top half of the curve on this interval, the equation of the top half being $y = -x\sqrt{x+3}$. So the area is $A = 2\int_{-3}^{0}\left(-x\sqrt{x+3}\right)dx$. We substitute $u = x+3$, so $du = dx$ and the limits change to 0 and 3, and we get

$$A = -2\int_0^3 [(u-3)\sqrt{u}\,]\,du = -2\int_0^3 (u^{3/2} - 3u^{1/2})\,du$$
$$= -2\left[\tfrac{2}{5}u^{5/2} - 2u^{3/2}\right]_0^3 = -2\left[\tfrac{2}{5}\left(3^2\sqrt{3}\right) - 2\left(3\sqrt{3}\right)\right] = \tfrac{24}{5}\sqrt{3}$$

64.

We start by finding the equation of the tangent line to $y = x^2$ at the point $(1, 1)$: $y' = 2x$, so the slope of the tangent is $2(1) = 2$, and its equation is $y - 1 = 2(x-1)$, or $y = 2x - 1$. We would need two integrals to integrate with respect to x, but only one to integrate with respect to y.

$$A = \int_0^1 \left[\tfrac{1}{2}(y+1) - \sqrt{y}\,\right]dy = \left[\tfrac{1}{4}y^2 + \tfrac{1}{2}y - \tfrac{2}{3}y^{3/2}\right]_0^1$$
$$= \tfrac{1}{4} + \tfrac{1}{2} - \tfrac{2}{3} = \tfrac{1}{12}$$

65.

By the symmetry of the problem, we consider only the first quadrant, where $y = x^2 \;\Rightarrow\; x = \sqrt{y}$. We are looking for a number b such that

$$\int_0^b \sqrt{y}\,dy = \int_b^4 \sqrt{y}\,dy \;\Rightarrow\; \tfrac{2}{3}\left[y^{3/2}\right]_0^b = \tfrac{2}{3}\left[y^{3/2}\right]_b^4 \;\Rightarrow$$
$$b^{3/2} = 4^{3/2} - b^{3/2} \;\Rightarrow\; 2b^{3/2} = 8 \;\Rightarrow\; b^{3/2} = 4 \;\Rightarrow\; b = 4^{2/3} \approx 2.52.$$

66. (a) We want to choose a so that

$$\int_1^a \frac{1}{x^2}\,dx = \int_a^4 \frac{1}{x^2}\,dx \;\Rightarrow\; \left[\frac{-1}{x}\right]_1^a = \left[\frac{-1}{x}\right]_a^4 \;\Rightarrow\; -\frac{1}{a} + 1 = -\frac{1}{4} + \frac{1}{a} \;\Rightarrow\; \frac{5}{4} = \frac{2}{a} \;\Rightarrow\; a = \frac{8}{5}.$$

(b) The area under the curve $y = 1/x^2$ from $x = 1$ to $x = 4$ is $\frac{3}{4}$ [take $a = 4$ in the first integral in part (a)]. Now the line $y = b$ must intersect the curve $x = 1/\sqrt{y}$ and not the line $x = 4$, since the area under the line $y = 1/4^2$ from $x = 1$ to $x = 4$ is only $\frac{3}{16}$, which is less than half of $\frac{3}{4}$. We want to choose b so that the upper area in the diagram is half of the total area under the curve $y = 1/x^2$ from $x = 1$ to $x = 4$. This implies that

$\int_b^1 \left(1/\sqrt{y} - 1\right)dy = \frac{1}{2}\cdot\frac{3}{4} \;\Rightarrow\; \left[2\sqrt{y} - y\right]_b^1 = \frac{3}{8} \;\Rightarrow\; 1 - 2\sqrt{b} + b = \frac{3}{8} \;\Rightarrow$ $b - 2\sqrt{b} + \frac{5}{8} = 0$. Letting $c = \sqrt{b}$, we get $c^2 - 2c + \frac{5}{8} = 0 \;\Rightarrow$ $8c^2 - 16c + 5 = 0$. Thus, $c = \frac{16 \pm \sqrt{256 - 160}}{16} = 1 \pm \frac{\sqrt{6}}{4}$. But $c = \sqrt{b} < 1 \;\Rightarrow$ $c = 1 - \frac{\sqrt{6}}{4} \;\Rightarrow\; b = c^2 = 1 + \frac{3}{8} - \frac{\sqrt{6}}{2} = \frac{1}{8}\left(11 - 4\sqrt{6}\right) \approx 0.1503.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. We first assume that $c > 0$, since c can be replaced by $-c$ in both equations without changing the graphs, and if $c = 0$ the curves do not enclose a region. We see from the graph that the enclosed area A lies between $x = -c$ and $x = c$, and by symmetry, it is equal to four times the area in the first quadrant. The enclosed area is

$A = 4\int_0^c (c^2 - x^2)\,dx = 4\left[c^2x - \frac{1}{3}x^3\right]_0^c = 4\left(c^3 - \frac{1}{3}c^3\right) = 4\left(\frac{2}{3}c^3\right) = \frac{8}{3}c^3$

So $A = 576 \;\Leftrightarrow\; \frac{8}{3}c^3 = 576 \;\Leftrightarrow\; c^3 = 216 \;\Leftrightarrow\; c = \sqrt[3]{216} = 6.$

Note that $c = -6$ is another solution, since the graphs are the same.

68.

It appears from the diagram that the curves $y = \cos x$ and $y = \cos(x - c)$ intersect halfway between 0 and c, namely, when $x = c/2$. We can verify that this is indeed true by noting that $\cos(c/2 - c) = \cos(-c/2) = \cos(c/2)$. The point where $\cos(x - c)$ crosses the x-axis is $x = \frac{\pi}{2} + c$. So we require that $\int_0^{c/2} [\cos x - \cos(x - c)]\,dx = -\int_{\pi/2+c}^{\pi} \cos(x - c)\,dx$ [the negative sign on the RHS is needed since the second area is beneath the x-axis] $\Leftrightarrow$ $[\sin x - \sin(x - c)]_0^{c/2} = -[\sin(x - c)]_{\pi/2+c}^{\pi}$ $\Rightarrow$ $[\sin(c/2) - \sin(-c/2)] - [-\sin(-c)] = -\sin(\pi - c) + \sin\left[\left(\frac{\pi}{2} + c\right) - c\right]$ $\Leftrightarrow$ $2\sin(c/2) - \sin c = -\sin c + 1$. [Here we have used the oddness of the sine function, and the fact that $\sin(\pi - c) = \sin c$]. So $2\sin(c/2) = 1$ $\Leftrightarrow$ $\sin(c/2) = \frac{1}{2}$ $\Leftrightarrow$ $c/2 = \frac{\pi}{6}$ $\Leftrightarrow$ $c = \frac{\pi}{3}$.

69. Let a and b be the x-coordinates of the points where the line intersects the curve. From the figure, $R_1 = R_2 \;\Rightarrow$

$$\int_0^a \left[c - \left(8x - 27x^3\right)\right] dx = \int_a^b \left[\left(8x - 27x^3\right) - c\right] dx$$

$$\left[cx - 4x^2 + \tfrac{27}{4}x^4\right]_0^a = \left[4x^2 - \tfrac{27}{4}x^4 - cx\right]_a^b$$

$$ac - 4a^2 + \tfrac{27}{4}a^4 = \left(4b^2 - \tfrac{27}{4}b^4 - bc\right) - \left(4a^2 - \tfrac{27}{4}a^4 - ac\right)$$

$$0 = 4b^2 - \tfrac{27}{4}b^4 - bc = 4b^2 - \tfrac{27}{4}b^4 - b\left(8b - 27b^3\right)$$

$$= 4b^2 - \tfrac{27}{4}b^4 - 8b^2 + 27b^4 = \tfrac{81}{4}b^4 - 4b^2$$

$$= b^2\left(\tfrac{81}{4}b^2 - 4\right)$$

So for $b > 0$, $b^2 = \frac{16}{81} \;\Rightarrow\; b = \frac{4}{9}$. Thus, $c = 8b - 27b^3 = 8\left(\frac{4}{9}\right) - 27\left(\frac{64}{729}\right) = \frac{32}{9} - \frac{64}{27} = \frac{32}{27}$.

70. The curve and the line will determine a region when they intersect at two or more points. So we solve the equation $x/(x^2 + 1) = mx \;\Rightarrow$

$x = x(mx^2 + m) \;\Rightarrow\; x(mx^2 + m) - x = 0 \;\Rightarrow$

$x(mx^2 + m - 1) = 0 \;\Rightarrow\; x = 0$ or $mx^2 + m - 1 = 0 \;\Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x = 0$ or $x^2 = \dfrac{1-m}{m} \quad \Rightarrow \quad x = 0$ or $x = \pm\sqrt{\dfrac{1}{m} - 1}$. Note that if $m = 1$, this has only the solution $x = 0$, and no region is determined. But if $1/m - 1 > 0 \quad \Leftrightarrow \quad 1/m > 1 \quad \Leftrightarrow \quad 0 < m < 1$, then there are two solutions. [Another way of seeing this is to observe that the slope of the tangent to $y = x/(x^2+1)$ at the origin is $y'(0) = 1$ and therefore we must have $0 < m < 1$.] Note that we cannot just integrate between the positive and negative roots, since the curve and the line cross at the origin. Since mx and $x/(x^2+1)$ are both odd functions, the total area is twice the area between the curves on the interval $\left[0, \sqrt{1/m - 1}\right]$. So the total area enclosed is

$$2\int_0^{\sqrt{1/m-1}} \left[\frac{x}{x^2+1} - mx\right] dx = 2\left[\tfrac{1}{2}\ln(x^2+1) - \tfrac{1}{2}mx^2\right]_0^{\sqrt{1/m-1}} = [\ln(1/m - 1 + 1) - m(1/m - 1)] - (\ln 1 - 0)$$
$$= \ln(1/m) - 1 + m = m - \ln m - 1$$

APPLIED PROJECT The Gini Index

1. (a) $G = \dfrac{\text{area between } L \text{ and } y = x}{\text{area under } y = x} = \dfrac{\int_0^1 [x - L(x)]\,dx}{\frac{1}{2}} = 2\displaystyle\int_0^1 [x - L(x)]\,dx$

(b) For a perfectly egalitarian society, $L(x) = x$, so $G = 2\int_0^1 [x - x]\,dx = 0$. For a perfectly totalitarian society,

$$L(x) = \begin{cases} 1 & \text{if } x = 1 \\ 0 & \text{if } 0 \le x < 1 \end{cases} \quad \text{so } G = 2\int_0^1 (x - 0)\,dx = 2\left[\tfrac{1}{2}x^2\right]_0^1 = 2\left(\tfrac{1}{2}\right) = 1.$$

2. (a) The richest 20% of the population in 2016 received $1 - L(0.8) = 1 - 0.485 = 0.515$, or 51.5%, of the total US income.

(b) A quadratic model has the form $Q(x) = ax^2 + bx + c$. Rounding to six decimal places, we get $a = 1.341\,071$, $b = -0.411\,929$, and $c = 0.028\,571$. The quadratic model appears to be a reasonable fit, but note that $Q(0) \neq 0$ and Q is both decreasing and increasing.

(c) $G = 2\displaystyle\int_0^1 [x - Q(x)]\,dx \approx 0.4607$

3.

Year	$Q(x) = ax^2 + bx + c$			Gini
	a	b	c	
1980	1.149 554	−0.189 696	0.016 179	0.3910
1990	1.214 732	−0.265 589	0.020 393	0.4150
2000	1.280 804	−0.345 232	0.025 821	0.4397
2010	1.312 946	−0.378 518	0.026 679	0.4499

The Gini index has risen steadily from 1980 to 2016. The trend is toward a less egalitarian society.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. Using TI's `PwrReg` command and omitting the point $(0, 0)$ gives us $P(x) = 0.839\,312x^{2.103\,154}$ and a Gini index of $G = 2\int_0^1 [x - P(x)]\,dx \approx 0.4591$. Note that the power function is nearly quadratic.

6.2 Volumes

1. (a)

(b) A cross-section is a disk with radius $x^2 + 5$, so its area is

$A(x) = \pi(x^2 + 5)^2 = \pi(x^4 + 10x^2 + 25)$.

$$V = \int_0^3 A(x)\,dx = \int_0^3 \pi(x^4 + 10x^2 + 25)\,dx$$

(c) $\displaystyle\int_0^3 \pi(x^4 + 10x^2 + 25)\,dx = \pi\left[\frac{1}{5}x^5 + \frac{10}{3}x^3 + 25x\right]_0^3 = \pi\left(\frac{243}{5} + 90 + 75\right) = \frac{1068}{5}\pi$

2. (a)

(b) A cross-section is a washer (annulus) with inner radius $\frac{1}{2}x$ and outer radius $\sqrt{x}$, so its area is

$A(x) = \pi\left[(\sqrt{x}\,)^2 - \left(\frac{1}{2}x\right)^2\right] = \pi\left(x - \frac{1}{4}x^2\right)$.

$$V = \int_0^4 A(x)\,dx = \int_0^4 \pi\left(x - \tfrac{1}{4}x^2\right)dx$$

(c) $\displaystyle\int_0^4 \pi\left(x - \tfrac{1}{4}x^2\right)dx = \pi\left[\tfrac{1}{2}x^2 - \tfrac{1}{12}x^3\right]_0^4 = \pi\left(8 - \frac{16}{3}\right) = \frac{8}{3}\pi$

3. (a)

(b) $y = x^3 + 1 \;\Rightarrow\; y - 1 = x^3 \;\Rightarrow\; x = \sqrt[3]{y - 1}$. Therefore, a cross-section is a disk with radius $\sqrt[3]{y - 1}$, so its area is

$A(y) = \pi\left(\sqrt[3]{y - 1}\,\right)^2 = \pi(y - 1)^{2/3}$.

$$V = \int_1^9 A(y)\,dy = \int_1^9 \pi(y - 1)^{2/3}\,dy$$

(c) $\displaystyle\int_1^9 \pi(y - 1)^{2/3}\,dy = \pi\left[\frac{3}{5}(y - 1)^{5/3}\right]_1^9 = \frac{3}{5}\pi(32 - 0) = \frac{96}{5}\pi$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a)

(b) $y = x/2 \quad \Rightarrow \quad x = 2y$; $y = 2/x \quad \Rightarrow \quad x = 2/y$. A cross-section is a washer (annulus) with inner radius $2/y$ and outer radius $2y$, so its area is $A(y) = \pi\left[(2y)^2 - \left(\dfrac{2}{y}\right)^2\right] = \pi\left(4y^2 - \dfrac{4}{y^2}\right)$.

$$V = \int_1^2 A(y)\,dy = \int_1^2 \pi\left(4y^2 - \frac{4}{y^2}\right)dy$$

(c) $\displaystyle\int_1^2 \pi\left(4y^2 - \frac{4}{y^2}\right)dy = \pi\left[\frac{4}{3}y^3 + \frac{4}{y}\right]_1^2 = \pi\left[\left(\frac{32}{3} + 2\right) - \left(\frac{4}{3} + 4\right)\right] = \frac{22}{3}\pi$

5. $V = \displaystyle\int_1^3 \pi(\ln x)^2\,dx$

6. $V = \displaystyle\int_0^5 \pi\left(\sqrt{5-y}\right)^2 dy$

7. $8y = x^2 \quad \Rightarrow \quad x = \sqrt{8y}$ for $x \geq 0$; $y = \sqrt{x} \quad \Rightarrow \quad x = y^2$ for $y \geq 0$.

$y^2 = \sqrt{8y} \quad \Rightarrow \quad y^4 = 8y \quad \Leftrightarrow \quad y^4 - 8y = 0 \quad \Leftrightarrow \quad y(y^3 - 8) = 0 \quad \Leftrightarrow$

$y = 0$ or $y = 2$.

$$V = \int_0^2 \pi\left[\left(\sqrt{8y}\right)^2 - (y^2)^2\right] dy$$

8. $(x-2)^2 = x + 10 \quad \Rightarrow \quad x^2 - 4x + 4 = x + 10 \quad \Rightarrow$

$x^2 - 5x - 6 = 0 \quad \Rightarrow \quad (x+1)(x-6) = 0 \quad \Rightarrow$

$x = -1$ or $x = 6$.

$$V = \int_{-1}^6 \pi\{(x+10)^2 - [(x-2)^2]^2\}\,dx$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $V = \pi \int_0^{\pi} \left\{ [\sin x - (-2)]^2 - [0 - (-2)]^2 \right\} dx$

$= \pi \int_0^{\pi} \left[(\sin x + 2)^2 - 2^2 \right] dx$

10. $y = \sqrt{x} \quad \Rightarrow \quad x = y^2$ for $y \geq 0$.

$V = \int_0^2 \pi[(6 - y^2)^2 - (6 - 4)^2]\, dy$

11. A cross-section is a disk with radius $x + 1$, so its area is

$A(x) = \pi(x + 1)^2 = \pi(x^2 + 2x + 1)$.

$V = \int_0^2 A(x)\, dx = \int_0^2 \pi(x^2 + 2x + 1)\, dx$

$= \pi\left[\frac{1}{3}x^3 + x^2 + x\right]_0^2$

$= \pi\left(\frac{8}{3} + 4 + 2\right) = \frac{26\pi}{3}$

12. A cross-section is a disk with radius $\dfrac{1}{x}$, so its area is $A(x) = \pi\left(\dfrac{1}{x}\right)^2 = \pi x^{-2}$.

$V = \int_1^4 A(x)\, dx = \int_1^4 \pi x^{-2}\, dx$

$= \pi\Big[-x^{-1}\Big]_1^4 = \pi\left(-\frac{1}{4} + 1\right)$

$= \frac{3\pi}{4}$

13. A cross-section is a disk with radius $\sqrt{x - 1}$, so its area is $A(x) = \pi\left(\sqrt{x - 1}\right)^2 = \pi(x - 1)$.

$$V = \int_1^5 A(x)\, dx = \int_1^5 \pi(x - 1)\, dx = \pi\left[\tfrac{1}{2}x^2 - x\right]_1^5 = \pi\left[\left(\tfrac{25}{2} - 5\right) - \left(\tfrac{1}{2} - 1\right)\right] = 8\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. A cross-section is a disk with radius e^x, so its area is $A(x) = \pi(e^x)^2 = \pi e^{2x}$.

$$V = \int_{-1}^{1} A(x)\,dx = \int_{-1}^{1} \pi e^{2x}\,dx$$
$$= \pi\left[\tfrac{1}{2}e^{2x}\right]_{-1}^{1} = \tfrac{\pi}{2}(e^2 - e^{-2})$$

15. A cross-section is a disk with radius $2\sqrt{y}$, so its area is $A(y) = \pi\left(2\sqrt{y}\right)^2$.

$$V = \int_0^9 A(y)\,dy = \int_0^9 \pi\left(2\sqrt{y}\right)^2 dy = 4\pi\int_0^9 y\,dy$$
$$= 4\pi\left[\tfrac{1}{2}y^2\right]_0^9 = 2\pi(81) = 162\pi$$

16. A cross-section is a disk with radius $\frac{1}{2}y^2$, so its area is $A(y) = \pi\left(\frac{1}{2}y^2\right)^2 = \frac{1}{4}\pi y^4$.

$$V = \int_0^4 A(y)\,dy = \int_0^4 \pi\left(\tfrac{1}{4}y^4\right) dy$$
$$= \tfrac{\pi}{4}\left[\tfrac{1}{5}y^5\right]_0^4 = \tfrac{\pi}{20}(4^5) = \tfrac{256\pi}{5}$$

17. A cross-section is a washer with inner radius $\frac{1}{2}y$ and outer radius $\sqrt{y}$, so its area is

$$A(y) = \pi\left[\left(\sqrt{y}\right)^2 - \left(\tfrac{1}{2}y\right)^2\right] = \pi\left(y - \tfrac{1}{4}y^2\right).$$

$$V = \int_0^4 \pi\left(y - \tfrac{1}{4}y^2\right) dy = \pi\left[\tfrac{1}{2}y^2 - \tfrac{1}{12}y^3\right]_0^4$$
$$= \pi\left[\left(8 - \tfrac{16}{3}\right) - 0\right] = \tfrac{8}{3}\pi$$

18. A cross-section is a washer (annulus) with inner radius 2 and outer radius $6 - x^2$, so its area is

$A(x) = \pi[(6 - x^2)^2 - 2^2] = \pi(x^4 - 12x^2 + 32)$.

$$V = \int_{-2}^{2} A(x)\,dx = 2\int_0^2 \pi(x^4 - 12x^2 + 32)\,dx$$
$$= 2\pi\left[\tfrac{1}{5}x^5 - 4x^3 + 32x\right]_0^2$$
$$= 2\pi\left(\tfrac{32}{5} - 32 + 64\right) = 2\pi\left(\tfrac{192}{5}\right) = \tfrac{384\pi}{5}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. A cross-section is a washer with inner radius x^3 and outer radius $\sqrt{x}$, so its area is

$$A(x) = \pi\left[\left(\sqrt{x}\right)^2 - (x^3)^2\right] = \pi(x - x^6).$$

$$V = \int_0^1 \pi(x - x^6)\,dx = \pi\left[\tfrac{1}{2}x^2 - \tfrac{1}{7}x^7\right]_0^1$$
$$= \pi\left[\left(\tfrac{1}{2} - \tfrac{1}{7}\right) - 0\right] = \tfrac{5}{14}\pi$$

20. A cross-section is a washer with inner radius y^4 and outer radius $2 - y^2$, so its area is

$$A(y) = \pi(2-y^2)^2 - \pi(y^4)^2 = \pi(4 - 4y^2 + y^4 - y^8).$$

$$V = \int_{-1}^{1} A(y)\,dy = 2\int_0^1 \pi(4 - 4y^2 + y^4 - y^8)\,dy$$
$$= 2\pi\left[4y - \tfrac{4}{3}y^3 + \tfrac{1}{5}y^5 - \tfrac{1}{9}y^8\right]_0^1$$
$$= 2\pi\left(4 - \tfrac{4}{3} + \tfrac{1}{5} - \tfrac{1}{9}\right) = 2\pi\left(\tfrac{124}{45}\right) = \tfrac{248\pi}{45}$$

21. A cross-section is a washer with inner radius $1 - \sqrt{x}$ and outer radius $1 - x^2$, so its area is

$$A(x) = \pi\left[(1 - x^2)^2 - (1 - \sqrt{x}\,)^2\right]$$
$$= \pi\left[(1 - 2x^2 + x^4) - (1 - 2\sqrt{x} + x)\right]$$
$$= \pi\left(x^4 - 2x^2 + 2\sqrt{x} - x\right).$$

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi(x^4 - 2x^2 + 2x^{1/2} - x)\,dx$$
$$= \pi\left[\tfrac{1}{5}x^5 - \tfrac{2}{3}x^3 + \tfrac{4}{3}x^{3/2} - \tfrac{1}{2}x^2\right]_0^1$$
$$= \pi\left(\tfrac{1}{5} - \tfrac{2}{3} + \tfrac{4}{3} - \tfrac{1}{2}\right) = \tfrac{11}{30}\pi$$

22. A cross-section is a washer with inner radius $1 - (-3) = 4$ and outer radius $x^3 - (-3) = x^3 + 3$, so its area is

$$A(x) = \pi(x^3 + 3)^2 - \pi(4)^2 = \pi(x^6 + 6x^3 - 7).$$

$$V = \int_1^2 A(x)\,dx = \int_1^2 \pi(x^6 + 6x^3 - 7)\,dx$$
$$= \pi\left[\tfrac{1}{7}x^7 + \tfrac{3}{2}x^4 - 7x\right]_1^2$$
$$= \pi\left[\left(\tfrac{128}{7} + 24 - 14\right) - \left(\tfrac{1}{7} + \tfrac{3}{2} - 7\right)\right] = \tfrac{471\pi}{14}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. A cross-section is a washer with inner radius $(1+\sec x)-1=\sec x$ and outer radius $3-1=2$, so its area is

$$A(x)=\pi\left[2^2-(\sec x)^2\right]=\pi(4-\sec^2 x).$$

$$V=\int_{-\pi/3}^{\pi/3} A(x)\,dx=\int_{-\pi/3}^{\pi/3}\pi(4-\sec^2 x)\,dx$$

$$=2\pi\int_0^{\pi/3}(4-\sec^2 x)\,dx \qquad \text{[by symmetry]}$$

$$=2\pi\Big[4x-\tan x\Big]_0^{\pi/3}=2\pi\left[\left(\tfrac{4\pi}{3}-\sqrt{3}\right)-0\right]$$

$$=2\pi\left(\tfrac{4\pi}{3}-\sqrt{3}\right)$$

24. A cross-section is a washer with inner radius $\sin x-(-1)$ and outer radius $\cos x-(-1)$, so its area is

$$A(x)=\pi\left[(\cos x+1)^2-(\sin x+1)^2\right]$$

$$=\pi(\cos^2 x+2\cos x-\sin^2 x-2\sin x)$$

$$=\pi(\cos 2x+2\cos x-2\sin x).$$

$$V=\int_0^{\pi/4}A(x)\,dx=\int_0^{\pi/4}\pi(\cos 2x+2\cos x-2\sin x)\,dx$$

$$=\pi\left[\tfrac{1}{2}\sin 2x+2\sin x+2\cos x\right]_0^{\pi/4}$$

$$=\pi\left[\left(\tfrac{1}{2}+\sqrt{2}+\sqrt{2}\right)-(0+0+2)\right]=\left(2\sqrt{2}-\tfrac{3}{2}\right)\pi$$

25. A cross-section is a washer with inner radius $2-1$ and outer radius $2-\sqrt[3]{y}$, so its area is

$$A(y)=\pi\left[(2-\sqrt[3]{y})^2-(2-1)^2\right]=\pi\left[4-4\sqrt[3]{y}+\sqrt[3]{y^2}-1\right].$$

$$V=\int_0^1 A(y)\,dy=\int_0^1\pi(3-4y^{1/3}+y^{2/3})\,dy=\pi\left[3y-3y^{4/3}+\tfrac{3}{5}y^{2/3}\right]_0^1=\pi\left(3-3+\tfrac{3}{5}\right)=\tfrac{3}{5}\pi.$$

26. For $0\le y<\frac{1}{2}$, a cross-section is a washer with inner radius $1-(-1)$ and outer radius $2-(-1)$, so its area is $A(y)=\pi(3^2-2^2)=5\pi$. For $\frac{1}{2}\le y\le 1$, a cross-section is a washer with inner radius $1-(-1)$ and outer radius $1/y-(-1)$, so its area is $A(y)=\pi\left[(1/y+1)^2-(2)^2\right]=\pi(1/y^2+2/y+1-4)$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$V = \int_0^{1/2} 5\pi\,dy + \int_{1/2}^1 \pi\left(\frac{1}{y^2} + \frac{2}{y} - 3\right) dy = 5\pi\left[y\right]_0^{1/2} + \pi\left[-\frac{1}{y} + 2\ln y - 3y\right]_{1/2}^1$$
$$= 5\pi\left(\tfrac{1}{2} - 0\right) + \pi\left[(-1 + 0 - 3) - \left(-2 + 2\ln\tfrac{1}{2} - \tfrac{3}{2}\right)\right] = \tfrac{5}{2}\pi + \pi\left(-\tfrac{1}{2} + 2\ln 2\right)$$
$$= (2 + 2\ln 2)\pi = 2\pi(1 + \ln 2)$$

27. From the symmetry of the curves, we see they intersect at $x = \frac{1}{2}$ and so $y^2 = \frac{1}{2} \quad \Leftrightarrow \quad y = \pm\sqrt{\frac{1}{2}}$. A cross-section is a washer with inner radius $3 - (1 - y^2)$ and outer radius $3 - y^2$, so its area is

$$A(y) = \pi\left[(3 - y^2)^2 - (2 + y^2)^2\right]$$
$$= \pi\left[(9 - 6y^2 + y^4) - (4 + 4y^2 + y^4)\right]$$
$$= \pi(5 - 10y^2).$$

$$V = \int_{-\sqrt{1/2}}^{\sqrt{1/2}} A(y)\,dy$$
$$= 2\int_0^{\sqrt{1/2}} 5\pi(1 - 2y^2)\,dy \qquad \text{[by symmetry]}$$
$$= 10\pi\left[y - \tfrac{2}{3}y^3\right]_0^{\sqrt{2}/2} = 10\pi\left(\tfrac{\sqrt{2}}{2} - \tfrac{\sqrt{2}}{6}\right)$$
$$= 10\pi\left(\tfrac{\sqrt{2}}{3}\right) = \tfrac{10}{3}\sqrt{2}\,\pi$$

28. For $0 \le y < 2$, a cross-section is an annulus with inner radius $2 - 1$ and outer radius $4 - 1$, the area of which is $A_1(y) = \pi(4 - 1)^2 - \pi(2 - 1)^2$. For $2 \le y \le 4$, a cross-section is an annulus with inner radius $y - 1$ and outer radius $4 - 1$, the area of which is $A_2(y) = \pi(4 - 1)^2 - \pi(y - 1)^2$.

$$V = \int_0^4 A(y)\,dy = \pi\int_0^2 \left[(4 - 1)^2 - (2 - 1)^2\right] dy + \pi\int_2^4 \left[(4 - 1)^2 - (y - 1)^2\right] dy$$
$$= \pi\left[8y\right]_0^2 + \pi\int_2^4 (8 + 2y - y^2)\,dy$$
$$= 16\pi + \pi\left[8y + y^2 - \tfrac{1}{3}y^3\right]_2^4$$
$$= 16\pi + \pi\left[\left(32 + 16 - \tfrac{64}{3}\right) - \left(16 + 4 - \tfrac{8}{3}\right)\right]$$
$$= \tfrac{76}{3}\pi$$

29. $\mathcal{R}_1$ about OA (the line $y = 0$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi(x)^2\,dx = \pi\left[\tfrac{1}{3}x^3\right]_0^1 = \tfrac{1}{3}\pi$$

30. $\mathcal{R}_1$ about OC (the line $x = 0$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi(1^2 - y^2)\,dy = \pi\left[y - \tfrac{1}{3}y^3\right]_0^1 = \pi\left(1 - \tfrac{1}{3}\right) = \tfrac{2}{3}\pi$$

31. $\mathcal{R}_1$ about AB (the line $x = 1$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi(1 - y)^2\,dy = \pi\int_0^1 (1 - 2y + y^2)\,dy = \pi\left[y - y^2 + \tfrac{1}{3}y^3\right]_0^1 = \tfrac{1}{3}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. $\mathcal{R}_1$ about BC (the line $y = 1$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi[(1-0)^2 - (1-x)^2]\,dx = \pi \int_0^1 [1 - (1 - 2x + x^2)]\,dx$$
$$= \pi \int_0^1 (-x^2 + 2x)\,dx = \pi\left[-\tfrac{1}{3}x^3 + x^2\right]_0^1 = \pi\left(-\tfrac{1}{3} + 1\right) = \tfrac{2}{3}\pi$$

33. $\mathcal{R}_2$ about OA (the line $y = 0$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi\left[1^2 - \left(\sqrt[4]{x}\right)^2\right]dx = \pi \int_0^1 (1 - x^{1/2})\,dx = \pi\left[x - \tfrac{2}{3}x^{3/2}\right]_0^1 = \pi\left(1 - \tfrac{2}{3}\right) = \tfrac{1}{3}\pi$$

34. $\mathcal{R}_2$ about OC (the line $x = 0$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi[(y^4)^2]\,dy = \pi \int_0^1 y^8\,dy = \pi\left[\tfrac{1}{9}y^9\right]_0^1 = \tfrac{1}{9}\pi$$

35. $\mathcal{R}_2$ about AB (the line $x = 1$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi[1^2 - (1 - y^4)^2]\,dy = \pi \int_0^1 [1 - (1 - 2y^4 + y^8)]\,dy$$
$$= \pi \int_0^1 (2y^4 - y^8)\,dy = \pi\left[\tfrac{2}{5}y^5 - \tfrac{1}{9}y^9\right]_0^1 = \pi\left(\tfrac{2}{5} - \tfrac{1}{9}\right) = \tfrac{13}{45}\pi$$

36. $\mathcal{R}_2$ about BC (the line $y = 1$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi(1 - \sqrt[4]{x})^2\,dx = \pi \int_0^1 (1 - 2x^{1/4} + x^{1/2})\,dx$$
$$= \pi\left[x - \tfrac{8}{5}x^{5/4} + \tfrac{2}{3}x^{3/2}\right]_0^1 = \pi\left(1 - \tfrac{8}{5} + \tfrac{2}{3}\right) = \tfrac{1}{15}\pi$$

37. $\mathcal{R}_3$ about OA (the line $y = 0$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi\left[\left(\sqrt[4]{x}\right)^2 - x^2\right]dx = \pi \int_0^1 (x^{1/2} - x^2)\,dx = \pi\left[\tfrac{2}{3}x^{3/2} - \tfrac{1}{3}x^3\right]_0^1 = \pi\left(\tfrac{2}{3} - \tfrac{1}{3}\right) = \tfrac{1}{3}\pi$$

Note: Let $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2 \cup \mathcal{R}_3$. If we rotate $\mathcal{R}$ about any of the segments OA, OC, AB, or BC, we obtain a right circular cylinder of height 1 and radius 1. Its volume is $\pi r^2 h = \pi(1)^2 \cdot 1 = \pi$. As a check for Exercises 29, 33, and 37, we can add the answers, and that sum must equal π. Thus, $\frac{1}{3}\pi + \frac{1}{3}\pi + \frac{1}{3}\pi = \pi$.

38. $\mathcal{R}_3$ about OC (the line $x = 0$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi[y^2 - (y^4)^2]\,dy = \pi \int_0^1 (y^2 - y^8)\,dy = \pi\left[\tfrac{1}{3}y^3 - \tfrac{1}{9}y^9\right]_0^1 = \pi\left(\tfrac{1}{3} - \tfrac{1}{9}\right) = \tfrac{2}{9}\pi$$

Note: See the note in the solution to Exercise 37. For Exercises 30, 34, and 38, we have $\frac{2}{3}\pi + \frac{1}{9}\pi + \frac{2}{9}\pi = \pi$.

39. $\mathcal{R}_3$ about AB (the line $x = 1$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi[(1 - y^4)^2 - (1 - y)^2]\,dy = \pi \int_0^1 [(1 - 2y^4 + y^8) - (1 - 2y + y^2)]\,dy$$
$$= \pi \int_0^1 (y^8 - 2y^4 - y^2 + 2y)\,dy = \pi\left[\tfrac{1}{9}y^9 - \tfrac{2}{5}y^5 - \tfrac{1}{3}y^3 + y^2\right]_0^1 = \pi\left(\tfrac{1}{9} - \tfrac{2}{5} - \tfrac{1}{3} + 1\right) = \tfrac{17}{45}\pi$$

Note: See the note in the solution to Exercise 37. For Exercises 31, 35, and 39, we have $\frac{1}{3}\pi + \frac{13}{45}\pi + \frac{17}{45}\pi = \pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $\mathcal{R}_3$ about BC (the line $y = 1$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi[(1-x)^2 - \left(1-\sqrt[4]{x}\right)^2]\,dx = \pi\int_0^1 [(1-2x+x^2) - (1-2x^{1/4}+x^{1/2})]\,dx$$

$$= \pi\int_0^1 (x^2 - 2x - x^{1/2} + 2x^{1/4})\,dx = \pi\left[\tfrac{1}{3}x^3 - x^2 - \tfrac{2}{3}x^{3/2} + \tfrac{8}{5}x^{5/4}\right]_0^1 = \pi\left(\tfrac{1}{3} - 1 - \tfrac{2}{3} + \tfrac{8}{5}\right) = \tfrac{4}{15}\pi$$

Note: See the note in the solution to Exercise 37. For Exercises 32, 36, and 40, we have $\frac{2}{3}\pi + \frac{1}{15}\pi + \frac{4}{15}\pi = \pi$.

41. (a) About the x-axis:

$$V = \int_{-1}^1 \pi(e^{-x^2})^2\,dx = 2\pi\int_0^1 e^{-2x^2}\,dx \quad \text{[by symmetry]}$$

$$\approx 3.75825$$

(b) About $y = -1$:

$$V = \int_{-1}^1 \pi\left\{[e^{-x^2} - (-1)]^2 - [0-(-1)]^2\right\}\,dx$$

$$= 2\pi\int_0^1 [(e^{-x^2}+1)^2 - 1]\,dx = 2\pi\int_0^1 (e^{-2x^2} + 2e^{-x^2})\,dx$$

$$\approx 13.14312$$

42. (a) About the x-axis:

$$V = \int_{-\pi/2}^{\pi/2} \pi(\cos^2 x)^2\,dx = 2\pi\int_0^{\pi/2} \cos^4 x\,dx \quad \text{[by symmetry]}$$

$$\approx 3.70110$$

(b) About $y = 1$:

$$V = \int_{-\pi/2}^{\pi/2} \pi[(1-0)^2 - (1-\cos^2 x)^2]\,dx$$

$$= 2\pi\int_0^{\pi/2} [1 - (1 - 2\cos^2 x + \cos^4 x)]\,dx$$

$$= 2\pi\int_0^{\pi/2} (2\cos^2 x - \cos^4 x)\,dx \approx 6.16850$$

43. (a) About $y = 2$:

$$x^2 + 4y^2 = 4 \quad\Rightarrow\quad 4y^2 = 4 - x^2 \quad\Rightarrow\quad y^2 = 1 - x^2/4 \quad\Rightarrow$$

$$y = \pm\sqrt{1 - x^2/4}$$

$$V = \int_{-2}^2 \pi\left\{\left[2 - \left(-\sqrt{1-x^2/4}\right)\right]^2 - \left(2 - \sqrt{1-x^2/4}\right)^2\right\}dx$$

$$= 2\pi\int_0^2 8\sqrt{1-x^2/4}\,dx \approx 78.95684$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) About $x = 2$:

$$x^2 + 4y^2 = 4 \quad \Rightarrow \quad x^2 = 4 - 4y^2 \quad \Rightarrow \quad x = \pm\sqrt{4 - 4y^2}$$

$$V = \int_{-1}^{1} \pi \left\{ \left[2 - \left(-\sqrt{4 - 4y^2}\right)\right]^2 - \left(2 - \sqrt{4 - 4y^2}\right)^2 \right\} dy$$

$$= 2\pi \int_0^1 8\sqrt{4 - 4y^2}\, dy \approx 78.95684$$

[Notice that this is the same approximation as in part (a). This can be explained by Pappus's Theorem in Section 8.3.]

44. (a) About the x-axis:

$y = x^2$ and $x^2 + y^2 = 1 \quad \Rightarrow \quad x^2 + x^4 = 1 \quad \Rightarrow \quad x^4 + x^2 - 1 = 0 \quad \Rightarrow$

$$x^2 = \frac{-1 + \sqrt{5}}{2} \approx 0.618 \quad \Rightarrow \quad x = \pm a = \pm\sqrt{\frac{-1 + \sqrt{5}}{2}} \approx \pm 0.786.$$

$$V = \int_{-a}^{a} \pi \left[\left(\sqrt{1 - x^2}\right)^2 - (x^2)^2 \right] dx = 2\pi \int_0^a (1 - x^2 - x^4)\, dx$$

$$\approx 3.54459$$

(b) About the y-axis:

$$V = \int_0^{a^2} \pi \left(\sqrt{y}\right)^2 dy + \int_{a^2}^{1} \pi \left(\sqrt{1 - y^2}\right)^2 dy$$

$$= \pi \int_0^{a^2} y\, dy + \pi \int_{a^2}^{1} (1 - y^2)\, dy \approx 0.99998$$

45. $y = \ln(x^6 + 2)$ and $y = \sqrt{3 - x^3}$ intersect at $x = a \approx -4.091$, $x = b \approx -1.467$, and $x = c \approx 1.091$.

$$V = \pi \int_a^b \left\{ \left[\ln(x^6 + 2)\right]^2 - \left(\sqrt{3 - x^3}\right)^2 \right\} dx + \pi \int_b^c \left\{ \left(\sqrt{3 - x^3}\right)^2 - \left[\ln(x^6 + 2)\right]^2 \right\} dx \approx 89.023$$

46. $y = 1 + xe^{-x^3}$ and $y = \arctan x^2$ intersect at $x = a \approx -0.570$ and $x = b \approx 1.391$.

$$V = \pi \int_a^b \left[\left(1 + xe^{-x^3}\right)^2 - \left(\arctan x^2\right)^2 \right] dx \approx 6.923$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $V = \pi \int_0^{\pi} \left\{ \left[\sin^2 x - (-1)\right]^2 - \left[0 - (-1)\right]^2 \right\} dx$

$\overset{\text{CAS}}{=} \frac{11}{8}\pi^2$

48. $V = \pi \int_0^2 \left[(3-x)^2 - (3 - xe^{1-x/2})^2\right] dx$

$\overset{\text{CAS}}{=} \pi\left(-2e^2 + 24e - \frac{142}{3}\right)$

49. $\pi \int_0^{\pi/2} \sin^2 x\, dx = \pi \int_0^{\pi/2} (\sin x)^2\, dx$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \{(x, y) \mid 0 \le x \le \pi/2, 0 \le y \le \sin x\}$ of the xy-plane about the x-axis.

50. $\pi \int_0^{\ln 2} e^{2x}\, dx = \pi \int_0^{\ln 2} (e^x)^2\, dx$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \{(x, y) \mid 0 \le x \le \ln 2, 0 \le y \le e^x\}$ of the xy-plane about the x-axis.

51. $\pi \int_0^1 (x^4 - x^6)\, dx = \pi \int_0^1 [(x^2)^2 - (x^3)^2]\, dx$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \{(x, y) \mid 0 \le x \le 1, x^3 \le y \le x^2\}$ of the xy-plane about the x-axis.

52. $\pi \int_{-1}^1 (1 - y^2)^2\, dy$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \{(x, y) \mid -1 \le y \le 1, 0 \le x \le 1 - y^2\}$ of the xy-plane about the y-axis.

53. $\pi \int_0^4 y\, dy = \pi \int_0^4 \left(\sqrt{y}\right)^2 dy$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \left\{(x, y) \mid 0 \le y \le 4, 0 \le x \le \sqrt{y}\right\}$ of the xy-plane about the y-axis.

54. $\pi \int_1^4 \left[3^2 - \left(3 - \sqrt{x}\right)^2\right] dx$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \left\{(x, y) \mid 1 \le x \le 4, 3 - \sqrt{x} \le y \le 3\right\}$ of the xy-plane about the x-axis.

55. There are 10 subintervals over the 15-cm length, so we'll use $n = 10/2 = 5$ for the Midpoint Rule.

$$V = \int_0^{15} A(x)\, dx \approx M_5 = \tfrac{15-0}{5}[A(1.5) + A(4.5) + A(7.5) + A(10.5) + A(13.5)]$$
$$= 3(18 + 79 + 106 + 128 + 39) = 3 \cdot 370 = 1110 \text{ cm}^3$$

56. $V = \int_0^{10} A(x)\, dx \approx M_5 = \frac{10-0}{5}[A(1) + A(3) + A(5) + A(7) + A(9)]$

$$= 2(0.65 + 0.61 + 0.59 + 0.55 + 0.50) = 2(2.90) = 5.80 \text{ m}^3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. (a) $V = \int_2^{10} \pi\,[f(x)]^2\,dx \approx \pi\frac{10-2}{4}\left\{[f(3)]^2 + [f(5)]^2 + [f(7)]^2 + [f(9)]^2\right\}$

$\approx 2\pi\left[(1.5)^2 + (2.2)^2 + (3.8)^2 + (3.1)^2\right] \approx 196 \text{ units}^3$

(b) $V = \int_0^4 \pi\left[(\text{outer radius})^2 - (\text{inner radius})^2\right] dy$

$\approx \pi\frac{4-0}{4}\left\{\left[(9.9)^2 - (2.2)^2\right] + \left[(9.7)^2 - (3.0)^2\right] + \left[(9.3)^2 - (5.6)^2\right] + \left[(8.7)^2 - (6.5)^2\right]\right\}$

$\approx 838 \text{ units}^3$

58. (a) $V = \displaystyle\int_{-1}^{1} \pi\left[\left(ax^3 + bx^2 + cx + d\right)\sqrt{1-x^2}\,\right]^2 dx \overset{\text{CAS}}{=} \frac{4\left\{5a^2 + 18ac + 3\left[3b^2 + 14bd + 7(c^2 + 5d^2)\right]\right\}\pi}{315}$

(b) $y = (-0.06x^3 + 0.04x^2 + 0.1x + 0.54)\sqrt{1-x^2}$ is graphed in the figure. Substitute $a = -0.06$, $b = 0.04$, $c = 0.1$, and $d = 0.54$ in the answer for part (a) to get $V \overset{\text{CAS}}{=} \dfrac{3769\pi}{9375} \approx 1.263$.

59. We'll form a right circular cone with height h and base radius r by revolving the line $y = \frac{r}{h}x$ about the x-axis.

$$V = \pi\int_0^h \left(\frac{r}{h}\,x\right)^2 dx = \pi\int_0^h \frac{r^2}{h^2}\,x^2 dx = \pi\,\frac{r^2}{h^2}\left[\frac{1}{3}x^3\right]_0^h$$

$$= \pi\,\frac{r^2}{h^2}\left(\frac{1}{3}h^3\right) = \frac{1}{3}\pi r^2 h$$

Another solution: Revolve $x = -\dfrac{r}{h}\,y + r$ about the y-axis.

$$V = \pi\int_0^h \left(-\frac{r}{h}y + r\right)^2 dy \overset{*}{=} \pi\int_0^h \left[\frac{r^2}{h^2}y^2 - \frac{2r^2}{h}y + r^2\right] dy$$

$$= \pi\left[\frac{r^2}{3h^2}y^3 - \frac{r^2}{h}y^2 + r^2 y\right]_0^h = \pi\left(\tfrac{1}{3}r^2 h - r^2 h + r^2 h\right) = \tfrac{1}{3}\pi r^2 h$$

* Or use substitution with $u = r - \dfrac{r}{h}\,y$ and $du = -\dfrac{r}{h}\,dy$ to get

$$\pi\int_r^0 u^2\left(-\frac{h}{r}\,du\right) = -\pi\,\frac{h}{r}\left[\frac{1}{3}u^3\right]_r^0 = -\pi\,\frac{h}{r}\left(-\frac{1}{3}r^3\right) = \frac{1}{3}\pi r^2 h.$$

60. $V = \pi\displaystyle\int_0^h \left(R - \frac{R-r}{h}\,y\right)^2 dy$

$$= \pi\int_0^h \left[R^2 - \frac{2R(R-r)}{h}\,y + \left(\frac{R-r}{h}\right)^2 y^2\right] dy$$

$$= \pi\left[R^2 y - \frac{R(R-r)}{h}\,y^2 + \frac{1}{3}\left(\frac{R-r}{h}\right)^2 y^3\right]_0^h$$

$$= \pi\left[R^2 h - R(R-r)h + \tfrac{1}{3}(R-r)^2 h\right]$$

$$= \tfrac{1}{3}\pi h\left[3Rr + (R^2 - 2Rr + r^2)\right] = \tfrac{1}{3}\pi h(R^2 + Rr + r^2)$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Another solution: $\dfrac{H}{R} = \dfrac{H-h}{r}$ by similar triangles. Therefore, $Hr = HR - hR \;\Rightarrow\; hR = H(R-r) \;\Rightarrow$

$H = \dfrac{hR}{R-r}$. Now

$$\begin{aligned}
V &= \tfrac{1}{3}\pi R^2 H - \tfrac{1}{3}\pi r^2 (H-h) \qquad \text{[by Exercise 59]} \\
&= \tfrac{1}{3}\pi R^2 \frac{hR}{R-r} - \tfrac{1}{3}\pi r^2 \frac{rh}{R-r} \qquad \left[H - h = \frac{rH}{R} = \frac{rhR}{R(R-r)}\right] \\
&= \frac{1}{3}\pi h \frac{R^3 - r^3}{R-r} = \tfrac{1}{3}\pi h\left(R^2 + Rr + r^2\right) \\
&= \tfrac{1}{3}\left[\pi R^2 + \pi r^2 + \sqrt{(\pi R^2)(\pi r^2)}\,\right] h = \tfrac{1}{3}\left(A_1 + A_2 + \sqrt{A_1 A_2}\,\right) h
\end{aligned}$$

where A_1 and A_2 are the areas of the bases of the frustum. (See Exercise 62 for a related result.)

61. $x^2 + y^2 = r^2 \quad\Leftrightarrow\quad x^2 = r^2 - y^2$

$$\begin{aligned}
V &= \pi \int_{r-h}^{r} \left(r^2 - y^2\right) dy = \pi\left[r^2 y - \frac{y^3}{3}\right]_{r-h}^{r} = \pi\left\{\left[r^3 - \frac{r^3}{3}\right] - \left[r^2(r-h) - \frac{(r-h)^3}{3}\right]\right\} \\
&= \pi\left\{\tfrac{2}{3}r^3 - \tfrac{1}{3}(r-h)\left[3r^2 - (r-h)^2\right]\right\} \\
&= \tfrac{1}{3}\pi\left\{2r^3 - (r-h)\left[3r^2 - \left(r^2 - 2rh + h^2\right)\right]\right\} \\
&= \tfrac{1}{3}\pi\left\{2r^3 - (r-h)\left[2r^2 + 2rh - h^2\right]\right\} \\
&= \tfrac{1}{3}\pi\left(2r^3 - 2r^3 - 2r^2 h + rh^2 + 2r^2 h + 2rh^2 - h^3\right) \\
&= \tfrac{1}{3}\pi\left(3rh^2 - h^3\right) = \tfrac{1}{3}\pi h^2 (3r - h), \text{ or, equivalently, } \pi h^2\left(r - \frac{h}{3}\right)
\end{aligned}$$

62. An equation of the line is $x = \dfrac{\Delta x}{\Delta y}\, y + (x\text{-intercept}) = \dfrac{a/2 - b/2}{h - 0}\, y + \dfrac{b}{2} = \dfrac{a-b}{2h}\, y + \dfrac{b}{2}$.

$$\begin{aligned}
V &= \int_0^h A(y)\, dy = \int_0^h (2x)^2\, dy \\
&= \int_0^h \left[2\left(\frac{a-b}{2h}y + \frac{b}{2}\right)\right]^2 dy = \int_0^h \left[\frac{a-b}{h}y + b\right]^2 dy \\
&= \int_0^h \left[\frac{(a-b)^2}{h^2}y^2 + \frac{2b(a-b)}{h}y + b^2\right] dy \\
&= \left[\frac{(a-b)^2}{3h^2}y^3 + \frac{b(a-b)}{h}y^2 + b^2 y\right]_0^h \\
&= \tfrac{1}{3}(a-b)^2 h + b(a-b)h + b^2 h = \tfrac{1}{3}\left(a^2 - 2ab + b^2 + 3ab\right)h \\
&= \tfrac{1}{3}\left(a^2 + ab + b^2\right)h
\end{aligned}$$

[Note that this can be written as $\tfrac{1}{3}\left(A_1 + A_2 + \sqrt{A_1 A_2}\,\right)h$, as in Exercise 60.]

If $a = b$, we get a rectangular solid with volume $b^2 h$. If $a = 0$, we get a square pyramid with volume $\tfrac{1}{3}b^2 h$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

63. For a cross-section at height y, we see from similar triangles that $\dfrac{\alpha/2}{b/2} = \dfrac{h-y}{h}$, so $\alpha = b\left(1 - \dfrac{y}{h}\right)$.

Similarly, for cross-sections having $2b$ as their base and β replacing α, $\beta = 2b\left(1 - \dfrac{y}{h}\right)$. So

$$V = \int_0^h A(y)\,dy = \int_0^h \left[b\left(1-\frac{y}{h}\right)\right]\left[2b\left(1-\frac{y}{h}\right)\right]dy$$

$$= \int_0^h 2b^2\left(1-\frac{y}{h}\right)^2 dy = 2b^2\int_0^h \left(1 - \frac{2y}{h} + \frac{y^2}{h^2}\right)dy$$

$$= 2b^2\left[y - \frac{y^2}{h} + \frac{y^3}{3h^2}\right]_0^h = 2b^2\left[h - h + \tfrac{1}{3}h\right]$$

$$= \tfrac{2}{3}b^2h \quad \left[= \tfrac{1}{3}Bh \text{ where } B \text{ is the area of the base, as with any pyramid.}\right]$$

64. Consider the triangle consisting of two vertices of the base and the center of the base. This triangle is similar to the corresponding triangle at a height y, so $a/b = \alpha/\beta \quad \Rightarrow \quad \alpha = a\beta/b$. Also by similar triangles, $b/h = \beta/(h-y) \Rightarrow \beta = b(h-y)/h$. These two equations imply that $\alpha = a(1 - y/h)$, and since the cross-section is an equilateral triangle, it has area

$$A(y) = \frac{1}{2}\cdot\alpha\cdot\frac{\sqrt{3}}{2}\alpha = \frac{a^2(1-y/h)^2}{4}\sqrt{3}, \text{ so}$$

$$V = \int_0^h A(y)\,dy = \frac{a^2\sqrt{3}}{4}\int_0^h \left(1-\frac{y}{h}\right)^2 dy$$

$$= \frac{a^2\sqrt{3}}{4}\left[-\frac{h}{3}\left(1-\frac{y}{h}\right)^3\right]_0^h = -\frac{\sqrt{3}}{12}a^2h(-1) = \frac{\sqrt{3}}{12}a^2h$$

65. A cross-section at height z is a triangle similar to the base, so we'll multiply the legs of the base triangle, 3 and 4, by a proportionality factor of $(5-z)/5$. Thus, the triangle at height z has area

$$A(z) = \frac{1}{2}\cdot 3\left(\frac{5-z}{5}\right)\cdot 4\left(\frac{5-z}{5}\right) = 6\left(1-\frac{z}{5}\right)^2, \text{ so}$$

$$V = \int_0^5 A(z)\,dz = 6\int_0^5\left(1-\frac{z}{5}\right)^2 dz = 6\int_1^0 u^2(-5\,du) \qquad \left[\begin{array}{l} u = 1 - z/5, \\ du = -\frac{1}{5}\,dz \end{array}\right]$$

$$= -30\left[\tfrac{1}{3}u^3\right]_1^0 = -30\left(-\tfrac{1}{3}\right) = 10 \text{ cm}^3$$

66. A cross-section is shaded in the diagram.

$$A(x) = (2y)^2 = \left(2\sqrt{r^2 - x^2}\right)^2, \text{ so}$$

$$V = \int_{-r}^{r} A(x)\,dx = 2\int_0^r 4(r^2 - x^2)\,dx$$

$$= 8\left[r^2x - \tfrac{1}{3}x^3\right]_0^r = 8\left(\tfrac{2}{3}r^3\right) = \tfrac{16}{3}r^3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. If l is a leg of the isosceles right triangle and $2y$ is the hypotenuse,

then $l^2 + l^2 = (2y)^2 \quad \Rightarrow \quad 2l^2 = 4y^2 \quad \Rightarrow \quad l^2 = 2y^2$.

$V = \int_{-2}^{2} A(x)\,dx = 2\int_0^2 A(x)\,dx = 2\int_0^2 \frac{1}{2}(l)(l)\,dx = 2\int_0^2 y^2\,dx$

$= 2\int_0^2 \frac{1}{4}(36 - 9x^2)\,dx = \frac{9}{2}\int_0^2 (4 - x^2)\,dx$

$= \frac{9}{2}\left[4x - \frac{1}{3}x^3\right]_0^2 = \frac{9}{2}\left(8 - \frac{8}{3}\right) = 24$

68. The cross-section of the base corresponding to the coordinate y has length $x = 1 - y$. The corresponding equilateral triangle with side s has area $A(y) = s^2\left(\frac{\sqrt{3}}{4}\right) = (1-y)^2\left(\frac{\sqrt{3}}{4}\right)$. Therefore,

$$V = \int_0^1 A(y)\,dy = \int_0^1 (1-y)^2\left(\frac{\sqrt{3}}{4}\right)dy$$

$$= \frac{\sqrt{3}}{4}\int_0^1 (1 - 2y + y^2)\,dy = \frac{\sqrt{3}}{4}\left[y - y^2 + \frac{1}{3}y^3\right]_0^1$$

$$= \frac{\sqrt{3}}{4}\left(\frac{1}{3}\right) = \frac{\sqrt{3}}{12}$$

Or: $\displaystyle\int_0^1 (1-y)^2\left(\frac{\sqrt{3}}{4}\right)dy = \frac{\sqrt{3}}{4}\int_1^0 u^2(-du) \quad [u = 1 - y] \; = \frac{\sqrt{3}}{4}\left[\frac{1}{3}u^3\right]_0^1 = \frac{\sqrt{3}}{12}$

69. The cross-section of the base corresponding to the coordinate x has length $y = 1 - x$. The corresponding square with side s has area $A(x) = s^2 = (1-x)^2 = 1 - 2x + x^2$. Therefore,

$$V = \int_0^1 A(x)\,dx = \int_0^1 (1 - 2x + x^2)\,dx$$

$$= \left[x - x^2 + \frac{1}{3}x^3\right]_0^1 = \left(1 - 1 + \frac{1}{3}\right) - 0 = \frac{1}{3}$$

Or: $\displaystyle\int_0^1 (1-x)^2\,dx = \int_1^0 u^2(-du) \quad [u = 1 - x] \; = \left[\frac{1}{3}u^3\right]_0^1 = \frac{1}{3}$

70. The cross-section of the base corresponding to the coordinate y has length $2x = 2\sqrt{1-y}$. $\left[y = 1 - x^2 \quad \Leftrightarrow \quad x = \pm\sqrt{1-y}\,\right]$ The corresponding square with side s has area $A(x) = s^2 = \left(2\sqrt{1-y}\right)^2 = 4(1-y)$. Therefore,

$V = \int_0^1 A(y)\,dy = \int_0^1 4(1-y)\,dy = 4\left[y - \frac{1}{2}y^2\right]_0^1 = 4\left[\left(1 - \frac{1}{2}\right) - 0\right] = 2.$

71. The cross-section of the base b corresponding to the coordinate x has length $1 - x^2$. The height h also has length $1 - x^2$, so the corresponding isosceles triangle has area $A(x) = \frac{1}{2}bh = \frac{1}{2}(1 - x^2)^2$. Therefore,

$$V = \int_{-1}^{1} A(x)\,dx = \int_{-1}^{1} \frac{1}{2}(1 - x^2)^2\,dx$$

$$= 2 \cdot \frac{1}{2}\int_0^1 (1 - 2x^2 + x^4)\,dx \qquad \text{[by symmetry]}$$

$$= \left[x - \frac{2}{3}x^3 + \frac{1}{5}x^5\right]_0^1 = \left(1 - \frac{2}{3} + \frac{1}{5}\right) - 0 = \frac{8}{15}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

72. The cross-section of the base corresponding to the coordinate y has length $2x = 2\sqrt{2-y}$. $[y = 2 - x^2 \quad \Leftrightarrow$ $x = \pm\sqrt{2-y}\,]$ The corresponding cross-section of the solid S is a quarter-circle with radius $2\sqrt{2-y}$ and area $A(y) = \frac{1}{4}\pi(2\sqrt{2-y}\,)^2 = \pi(2-y)$. Therefore,

$$V = \int_0^2 A(y)\,dy = \int_0^2 \pi(2-y)\,dy$$
$$= \pi\left[2y - \tfrac{1}{2}y^2\right]_0^2 = \pi(4-2) = 2\pi$$

73. The cross-section of S at coordinate x, $-1 \le x \le 1$, is a circle centered at the point $\left(x, \frac{1}{2}(1-x^2)\right)$ with radius $\frac{1}{2}(1-x^2)$. The area of the cross-section is

$$A(x) = \pi\left[\tfrac{1}{2}(1-x^2)\right]^2 = \tfrac{\pi}{4}(1 - 2x^2 + x^4)$$

The volume of S is

$$V = \int_{-1}^{1} A(x)\,dx = 2\int_0^1 \tfrac{\pi}{4}(1-2x^2+x^4)\,dx = \tfrac{\pi}{2}\left[x - \tfrac{2}{3}x^3 + \tfrac{1}{5}x^5\right]_0^1 = \tfrac{\pi}{2}\left(1 - \tfrac{2}{3} + \tfrac{1}{5}\right) = \tfrac{\pi}{2}\left(\tfrac{8}{15}\right) = \tfrac{4\pi}{15}$$

74. The cross-section of S at coordinate x, $0 \le x \le 4$, is a circle centered at the point $\left(x, \frac{1}{2}\left(\frac{1}{2}\sqrt{x} + \sqrt{x}\right)\right)$ with radius $\frac{1}{2}\left(\sqrt{x} - \frac{1}{2}\sqrt{x}\right)$. The area of the cross-section is $A(x) = \pi\left[\frac{1}{2}\left(\sqrt{x} - \frac{1}{2}\sqrt{x}\right)\right]^2 = \pi \cdot \frac{1}{4} \cdot \left(\frac{1}{2}\sqrt{x}\right)^2 = \dfrac{\pi x}{16}$. The volume of S is $V = \displaystyle\int_0^4 A(x)\,dx = \int_0^4 \frac{\pi x}{16}\,dx = \frac{\pi}{32}\left[x^2\right]_0^4 = \frac{\pi}{32}(16-0) = \frac{\pi}{2}$.

75. (a) The torus is obtained by rotating the circle $(x-R)^2 + y^2 = r^2$ about the y-axis. Solving for x, we see that the right half of the circle is given by $x = R + \sqrt{r^2 - y^2} = f(y)$ and the left half by $x = R - \sqrt{r^2 - y^2} = g(y)$. So

$$V = \pi\int_{-r}^{r}\left\{[f(y)]^2 - [g(y)]^2\right\}dy$$
$$= 2\pi\int_0^r\left[\left(R^2 + 2R\sqrt{r^2-y^2} + r^2 - y^2\right) - \left(R^2 - 2R\sqrt{r^2-y^2} + r^2 - y^2\right)\right]dy$$
$$= 2\pi\int_0^r 4R\sqrt{r^2-y^2}\,dy = 8\pi R\int_0^r \sqrt{r^2-y^2}\,dy$$

(b) Observe that the integral represents a quarter of the area of a circle with radius r, so $8\pi R\int_0^r \sqrt{r^2-y^2}\,dy = 8\pi R \cdot \frac{1}{4}\pi r^2 = 2\pi^2 r^2 R$.

76. (a) $V = \int_{-r}^{r} A(x)\,dx = 2\int_0^r A(x)\,dx = 2\int_0^r \frac{1}{2}h\left(2\sqrt{r^2-x^2}\,\right)dx = 2h\int_0^r \sqrt{r^2-x^2}\,dx$

(b) Observe that the integral represents one quarter of the area of a circle of radius r, so $V = 2h \cdot \frac{1}{4}\pi r^2 = \frac{1}{2}\pi h r^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

77. The cross-sections perpendicular to the y-axis in Figure 17 are rectangles. The rectangle corresponding to the coordinate y has a base of length $2\sqrt{16-y^2}$ in the xy-plane and a height of $\frac{1}{\sqrt{3}}\,y$, since $\angle BAC = 30^\circ$ and $|BC| = \frac{1}{\sqrt{3}}\,|AB|$. Thus, $A(y) = \frac{2}{\sqrt{3}}\,y\sqrt{16-y^2}$ and

$$V = \int_0^4 A(y)\,dy = \tfrac{2}{\sqrt{3}}\int_0^4 \sqrt{16-y^2}\,y\,dy = \tfrac{2}{\sqrt{3}}\int_{16}^0 u^{1/2}\left(-\tfrac{1}{2}\,du\right) \qquad [\text{Put } u = 16-y^2, \text{ so } du = -2y\,dy]$$

$$= \tfrac{1}{\sqrt{3}}\int_0^{16} u^{1/2}\,du = \tfrac{1}{\sqrt{3}}\,\tfrac{2}{3}\left[u^{3/2}\right]_0^{16} = \tfrac{2}{3\sqrt{3}}\,(64) = \tfrac{128}{3\sqrt{3}}$$

78. (a) $\text{Volume}(S_1) = \int_0^h A(z)\,dz = \text{Volume}(S_2)$ since the cross-sectional area $A(z)$ at height z is the same for both solids.

(b) By Cavalieri's Principle, the volume of the cylinder in the figure is the same as that of a right circular cylinder with radius r and height h, that is, $\pi r^2 h$.

79.

By similar triangles, the radius x_1 of the cross-section at height y of the cone removed from the cylinder satisfies $\dfrac{x_1}{y} = \dfrac{r}{r} \quad\Rightarrow\quad x_1 = y$. Thus, the area of the annular cross-section at height y remaining once the cone is removed from the cylinder is $\pi(r^2-y^2)$.

The radius x_2 of the cross-section at height y of the hemisphere satisfies $x_2^2 + y^2 = r^2 \quad\Rightarrow\quad x_2 = \sqrt{r^2-y^2}$.

The area of the circular cross-section at height y is then $\pi\left(\sqrt{r^2-y^2}\right)^2 = \pi(r^2-y^2)$.

Each cross-section at height y of the cylinder with cone removed has area equal to that of the corresponding cross-section at height y of the hemisphere. By Cavalieri's Principle, the volumes of the solids are then equal.

80. Each cross-section of the solid S in a plane perpendicular to the x-axis is a square (since the edges of the cut lie on the cylinders, which are perpendicular). One-quarter of this square and one-eighth of S are shown. The area of this quarter-square is $|PQ|^2 = r^2 - x^2$. Therefore, $A(x) = 4(r^2-x^2)$ and the volume of S is

$$V = \int_{-r}^{r} A(x)\,dx = 4\int_{-r}^{r}(r^2-x^2)\,dx$$

$$= 8(r^2-x^2)\,dx = 8\left[r^2x - \tfrac{1}{3}x^3\right]_0^r = \tfrac{16}{3}r^3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

81. The volume is obtained by rotating the area common to two circles of radius r, as shown. The volume of the right half is

$$V_{\text{right}} = \pi \int_0^{r/2} y^2\,dx = \pi \int_0^{r/2} \left[r^2 - \left(\tfrac{1}{2}r + x\right)^2\right] dx$$

$$= \pi\left[r^2x - \tfrac{1}{3}\left(\tfrac{1}{2}r + x\right)^3\right]_0^{r/2} = \pi\left[\left(\tfrac{1}{2}r^3 - \tfrac{1}{3}r^3\right) - \left(0 - \tfrac{1}{24}r^3\right)\right] = \tfrac{5}{24}\pi r^3$$

So by symmetry, the total volume is twice this, or $\frac{5}{12}\pi r^3$.

Another solution: We observe that the volume is the twice the volume of a cap of a sphere, so we can use the formula from Exercise 61 with $h = \frac{1}{2}r$: $V = 2 \cdot \frac{1}{3}\pi h^2(3r - h) = \frac{2}{3}\pi\left(\frac{1}{2}r\right)^2\left(3r - \frac{1}{2}r\right) = \frac{5}{12}\pi r^3$.

82. We consider two cases: one in which the ball is not completely submerged and the other in which it is.

Case 1: $0 \le h \le 10$ The ball will not be completely submerged, and so a cross-section of the water parallel to the surface will be the shaded area shown in the first diagram. We can find the area of the cross-section at height x above the bottom of the bowl by using the Pythagorean Theorem: $R^2 = 15^2 - (15 - x)^2$ and $r^2 = 5^2 - (x - 5)^2$, so $A(x) = \pi\left(R^2 - r^2\right) = 20\pi x$. The volume of water when it has depth h is then $V(h) = \int_0^h A(x)\,dx = \int_0^h 20\pi x\,dx = \left[10\pi x^2\right]_0^h = 10\pi h^2$ cm^3, $0 \le h \le 10$.

Case 2: $10 < h \le 15$ In this case we can find the volume by simply subtracting the volume displaced by the ball from the total volume inside the bowl underneath the surface of the water. The total volume underneath the surface is just the volume of a cap of the bowl, so we use the formula from Exercise 61: $V_{\text{cap}}(h) = \frac{1}{3}\pi h^2(45 - h)$. The volume of the small sphere is $V_{\text{ball}} = \frac{4}{3}\pi(5)^3 = \frac{500}{3}\pi$, so the total volume is $V_{\text{cap}} - V_{\text{ball}} = \frac{1}{3}\pi(45h^2 - h^3 - 500)$ cm^3.

83. Take the x-axis to be the axis of the cylindrical hole of radius r. A quarter of the cross-section through y, perpendicular to the y-axis, is the rectangle shown. Using the Pythagorean Theorem twice, we see that the dimensions of this rectangle are $x = \sqrt{R^2 - y^2}$ and $z = \sqrt{r^2 - y^2}$, so $\frac{1}{4}A(y) = xz = \sqrt{r^2 - y^2}\,\sqrt{R^2 - y^2}$, and

$$V = \int_{-r}^{r} A(y)\,dy = \int_{-r}^{r} 4\sqrt{r^2 - y^2}\,\sqrt{R^2 - y^2}\,dy = 8\int_0^r \sqrt{r^2 - y^2}\,\sqrt{R^2 - y^2}\,dy$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

84. The line $y = r$ intersects the semicircle $y = \sqrt{R^2 - x^2}$ when $r = \sqrt{R^2 - x^2} \quad \Rightarrow \quad r^2 = R^2 - x^2 \quad \Rightarrow$ $x^2 = R^2 - r^2 \quad \Rightarrow \quad x = \pm\sqrt{R^2 - r^2}$. Rotating the shaded region about the x-axis gives us

$$V = \int_{-\sqrt{R^2-r^2}}^{\sqrt{R^2-r^2}} \pi\left[\left(\sqrt{R^2 - x^2}\right)^2 - r^2\right] dx = 2\pi \int_0^{\sqrt{R^2-r^2}} (R^2 - x^2 - r^2)\, dx \qquad \text{[by symmetry]}$$

$$= 2\pi \int_0^{\sqrt{R^2-r^2}} \left[(R^2 - r^2) - x^2\right] dx = 2\pi\left[(R^2 - r^2)x - \tfrac{1}{3}x^3\right]_0^{\sqrt{R^2-r^2}}$$

$$= 2\pi\left[(R^2 - r^2)^{3/2} - \tfrac{1}{3}(R^2 - r^2)^{3/2}\right] = 2\pi \cdot \tfrac{2}{3}(R^2 - r^2)^{3/2} = \tfrac{4\pi}{3}(R^2 - r^2)^{3/2}$$

Our answer makes sense in limiting cases. As $r \to 0$, $V \to \frac{4}{3}\pi R^3$, which is the volume of the full sphere. As $r \to R$, $V \to 0$, which makes sense because the hole's radius is approaching that of the sphere.

85. (a) The radius of the barrel is the same at each end by symmetry, since the function $y = R - cx^2$ is even. Since the barrel is obtained by rotating the graph of the function y about the x-axis, this radius is equal to the value of y at $x = \frac{1}{2}h$, which is $R - c\left(\frac{1}{2}h\right)^2 = R - d = r$.

(b) The barrel is symmetric about the y-axis, so its volume is twice the volume of that part of the barrel for $x > 0$. Also, the barrel is a volume of rotation, so

$$V = 2\int_0^{h/2} \pi y^2\, dx = 2\pi \int_0^{h/2} \left(R - cx^2\right)^2 dx = 2\pi\left[R^2 x - \tfrac{2}{3}Rcx^3 + \tfrac{1}{5}c^2x^5\right]_0^{h/2}$$

$$= 2\pi\left(\tfrac{1}{2}R^2 h - \tfrac{1}{12}Rch^3 + \tfrac{1}{160}c^2h^5\right)$$

Trying to make this look more like the expression we want, we rewrite it as $V = \frac{1}{3}\pi h\left[2R^2 + \left(R^2 - \frac{1}{2}Rch^2 + \frac{3}{80}c^2h^4\right)\right]$.

But $R^2 - \frac{1}{2}Rch^2 + \frac{3}{80}c^2h^4 = \left(R - \frac{1}{4}ch^2\right)^2 - \frac{1}{40}c^2h^4 = (R - d)^2 - \frac{2}{5}\left(\frac{1}{4}ch^2\right)^2 = r^2 - \frac{2}{5}d^2$.

Substituting this back into V, we see that $V = \frac{1}{3}\pi h\left(2R^2 + r^2 - \frac{2}{5}d^2\right)$, as required.

86. It suffices to consider the case where $\mathcal{R}$ is bounded by the curves $y = f(x)$ and $y = g(x)$ for $a \le x \le b$, where $g(x) \le f(x)$ for all x in $[a, b]$, since other regions can be decomposed into subregions of this type. We are concerned with the volume obtained when $\mathcal{R}$ is rotated about the line $y = -k$, which is equal to

$$V_2 = \pi \int_a^b \left([f(x) + k]^2 - [g(x) + k]^2\right) dx$$

$$= \pi \int_a^b \left([f(x)]^2 - [g(x)]^2\right) dx + 2\pi k \int_a^b [f(x) - g(x)]\, dx = V_1 + 2\pi kA$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

87. (a) $y = x^3 \quad \Rightarrow \quad x = \sqrt[3]{y}$. $V_1 = \int_1^8 \pi \left(\sqrt[3]{y}\right)^2 \, dy = \int_1^8 \pi y^{2/3} \, dy = \frac{3}{5}\pi \left[y^{5/3}\right]_1^8 = \frac{3}{5}\pi(32-1) = \frac{93}{5}\pi$

(b) If each y is replaced with cy, then $y = 1$ will be mapped to $y = 1c = c$, and $y = 8$ will be mapped to $y = 8c$, each as shown. If each x is replaced with cx, then $y = (cx)^3/c^2 = c^3x^3/c^2 = cx^3$, so y has again been mapped to cy. A dilation with scaling factor c therefore transforms the region R_1 into R_2.

(c) $y = x^3/c^2 \quad \Rightarrow \quad c^2y = x^3 \quad \Rightarrow \quad x = \sqrt[3]{c^2y}$. Then

$$V_2 = \int_c^{8c} \pi\left(\sqrt[3]{c^2y}\right)^2 \, dy = \int_c^{8c} \pi \cdot c^{4/3}y^{2/3} \, dy = \tfrac{3}{5}\pi \cdot c^{4/3}\left[y^{5/3}\right]_c^{8c} = \tfrac{3}{5}\pi \cdot c^{4/3}(32c^{5/3} - c^{5/3})$$
$$= \tfrac{3}{5}\pi \cdot c^{4/3} \cdot 31c^{5/3} = \tfrac{3}{5}\pi \cdot 31c^3 = \tfrac{93}{5}\pi c^3 = c^3V_1$$

(d) $V_2 = 5 \text{ L} \quad \Rightarrow \quad \dfrac{93}{5}\pi c^3 = 5000 \text{ cm}^3 \quad \Rightarrow \quad c^3 = \dfrac{25{,}000}{93\pi} \quad \Rightarrow \quad c = \sqrt[3]{\dfrac{25{,}000}{93\pi}} \approx 4.41$

6.3 Volumes by Cylindrical Shells

1.

If we were to use the "washer" method, we would first have to locate the local maximum point (a, b) of $y = x(x-1)^2$ using the methods of Chapter 4. Then we would have to solve the equation $y = x(x-1)^2$ for x in terms of y to obtain the functions $x = g_1(y)$ and $x = g_2(y)$ shown in the first figure. This step would be difficult because it involves the cubic formula. Finally we would find the volume using $V = \pi \int_0^b \left\{[g_1(y)]^2 - [g_2(y)]^2\right\} dy$.

Using shells, we find that a typical approximating shell has radius x, so its circumference is $2\pi x$. Its height is y, that is, $x(x-1)^2$. So the total volume is

$$V = \int_0^1 2\pi x\left[x(x-1)^2\right] dx = 2\pi \int_0^1 \left(x^4 - 2x^3 + x^2\right) dx = 2\pi\left[\frac{x^5}{5} - 2\frac{x^4}{4} + \frac{x^3}{3}\right]_0^1 = \frac{\pi}{15}$$

2.

A typical cylindrical shell has circumference $2\pi x$ and height $\sin(x^2)$. $V = \int_0^{\sqrt{\pi}} 2\pi x \sin(x^2) \, dx$. Let $u = x^2$. Then $du = 2x \, dx$, so $V = \pi \int_0^{\pi} \sin u \, du = \pi[-\cos u]_0^{\pi} = \pi[1 - (-1)] = 2\pi$. For washers, we would first have to locate the local maximum point (a, b) of $y = \sin(x^2)$ using the methods of Chapter 4. Then we would have to solve the equation $y = \sin(x^2)$ for x in terms of y to obtain the functions $x = g_1(y)$ and $x = g_2(y)$ shown in the second figure. Finally we would find the volume using $V = \pi \int_0^b \left\{[g_1(y)]^2 - [g_2(y)]^2\right\} dy$. Using shells is definitely preferable to using washers.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a)

$$V = \int_0^{\sqrt{\pi/2}} 2\pi x \cos(x^2)\,dx$$

(b) $V = \displaystyle\int_0^{\sqrt{\pi/2}} 2\pi x\cos(x^2)\,dx = 2\pi\int_0^{\sqrt{\pi/2}} x\cos(x^2)\,dx = 2\pi\cdot\tfrac{1}{2}\Big[\sin(x^2)\Big]_0^{\sqrt{\pi/2}} = \pi\Big[\sin\Big(\frac{\pi}{2}\Big) - \sin 0\Big]$

$= \pi(1-0) = \pi$

4. (a)

$y = 2 - x \quad\Rightarrow\quad x = 2 - y;$

$y = \sqrt{x} \quad\Rightarrow\quad x = y^2$

$$V = \int_0^1 2\pi y[(2-y) - y^2]\,dy$$

(b) $V = \int_0^1 2\pi y[(2-y)-y^2]\,dy = 2\pi\int_0^1 (2y - y^2 - y^3)\,dy = 2\pi\big[y^2 - \frac{1}{3}y^3 - \frac{1}{4}y^4\big]_0^1$

$2\pi\big[\big(1 - \frac{1}{3} - \frac{1}{4}\big) - 0\big] = 2\pi\big(\frac{5}{12}\big) = \frac{5}{6}\pi$

5. $V = \int_1^2 2\pi x \ln x\,dx$

6. $y = x^3 \quad\Rightarrow\quad x = \sqrt[3]{y}$

$V = \int_0^8 2\pi y\sqrt[3]{y}\,dy$

7. $y = \sin^{-1} x \quad\Rightarrow\quad x = \sin y$

$V = \int_0^{\pi/2} 2\pi(3-y)\sin y\,dy$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $V = \int_0^3 2\pi(7-x)[(4x-x^2)-x]\,dx$

9. The shell has radius x, circumference $2\pi x$, and height $\sqrt{x}$, so

$$V = \int_0^4 2\pi x\sqrt{x}\,dx = \int_0^4 2\pi x^{3/2}\,dx = 2\pi\left[\tfrac{2}{5}x^{5/2}\right]_0^4 = 2\pi\cdot\tfrac{2}{5}(32-0) = \tfrac{128}{5}\pi.$$

10. The shell has radius x, circumference $2\pi x$, and height x^3, so

$$V = \int_1^2 2\pi x\cdot x^3\,dx = 2\pi\int_1^2 x^4\,dx$$
$$= 2\pi\left[\tfrac{1}{5}x^5\right]_1^2 = 2\pi\left(\tfrac{32}{5}-\tfrac{1}{5}\right) = \tfrac{62}{5}\pi$$

11. The shell has radius x, circumference $2\pi x$, and height $1/x$, so

$$V = \int_1^4 2\pi x\left(\frac{1}{x}\right)dx = \int_1^4 2\pi\,dx = 2\pi\Big[x\Big]_1^4 = 2\pi(4-1) = 6\pi.$$

12. The shell has radius x, circumference $2\pi x$, and height e^{-x^2}, so $V = \int_0^1 2\pi x e^{-x^2}\,dx$.

Let $u = x^2$. Thus, $du = 2x\,dx$, so

$$V = \pi\int_0^1 e^{-u}\,du = \pi\left[-e^{-u}\right]_0^1 = \pi(1-1/e).$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. The shell has radius x, circumference $2\pi x$, and height $\sqrt{5+x^2}$, so

$$V = \int_0^2 2\pi x\sqrt{5+x^2}\,dx \quad [\,u = 5+x^2,\, du = 2x\,dx\,]$$
$$= \int_5^9 2\pi \cdot \frac{1}{2}u^{1/2}\,du = \pi \cdot \frac{2}{3}\left[u^{3/2}\right]_5^9$$
$$= \frac{2}{3}\pi(27 - 5^{3/2})$$

14. $4x - x^2 = x \quad \Leftrightarrow \quad 0 = x^2 - 3x \quad \Leftrightarrow \quad 0 = x(x-3) \quad \Leftrightarrow \quad x = 0$ or 3.

The shell has radius x, circumference $2\pi x$, and height $[(4x - x^2) - x]$, so

$$V = \int_0^3 2\pi x[(4x - x^2) - x]\,dx$$
$$= 2\pi \int_0^3 (-x^3 + 3x^2)\,dx$$
$$= 2\pi\left[-\tfrac{1}{4}x^4 + x^3\right]_0^3$$
$$= 2\pi\left(-\tfrac{81}{4} + 27\right) = 2\pi\left(\tfrac{27}{4}\right) = \tfrac{27}{2}\pi$$

15. $xy = 1 \quad \Rightarrow \quad x = \frac{1}{y}$. The shell has radius y, circumference $2\pi y$, and height $1/y$, so

$$V = \int_1^3 2\pi y\left(\frac{1}{y}\right)dy$$
$$= 2\pi \int_1^3 dy = 2\pi\left[y\right]_1^3$$
$$= 2\pi(3-1) = 4\pi$$

16. $y = \sqrt{x} \quad \Rightarrow \quad x = y^2$. The shell has radius y, circumference $2\pi y$, and height y^2, so

$$V = \int_0^2 2\pi y(y^2)\,dy = 2\pi \int_0^2 y^3\,dy$$
$$= 2\pi\left[\tfrac{1}{4}y^4\right]_0^2$$
$$= 2\pi(4) = 8\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $y = x^{3/2} \quad \Rightarrow \quad x = y^{2/3}$. The shell has radius y, circumference $2\pi y$, and height $y^{2/3}$, so

$$V = \int_0^8 2\pi y(y^{2/3})\,dy = 2\pi \int_0^8 y^{5/3}\,dy$$
$$= 2\pi\left[\tfrac{3}{8}y^{8/3}\right]_0^8$$
$$= 2\pi \cdot \tfrac{3}{8} \cdot 256 = 192\pi$$

18. The shell has radius y, circumference $2\pi y$, and height $-3y^2 + 12y - 9$, so

$$V = \int_1^3 2\pi y(-3y^2 + 12y - 9)\,dy$$
$$= 2\pi \int_1^3 (-3y^3 + 12y^2 - 9y)\,dy$$
$$= -6\pi \int_1^3 (y^3 - 4y^2 + 3y)\,dy$$
$$= -6\pi\left[\tfrac{1}{4}y^4 - \tfrac{4}{3}y^3 + \tfrac{3}{2}y^2\right]_1^3$$
$$= -6\pi\left[\left(\tfrac{81}{4} - 36 + \tfrac{27}{2}\right) - \left(\tfrac{1}{4} - \tfrac{4}{3} + \tfrac{3}{2}\right)\right]$$
$$= -6\pi\left(-\tfrac{8}{3}\right) = 16\pi$$

19. The shell has radius y, circumference $2\pi y$, and height

$$2 - \left[1 + (y-2)^2\right] = 1 - (y-2)^2 = 1 - \left(y^2 - 4y + 4\right)$$
$$= -y^2 + 4y - 3, \text{ so}$$

$$V = \int_1^3 2\pi y(-y^2 + 4y - 3)\,dy$$
$$= 2\pi \int_1^3 (-y^3 + 4y^2 - 3y)\,dy$$
$$= 2\pi\left[-\tfrac{1}{4}y^4 + \tfrac{4}{3}y^3 - \tfrac{3}{2}y^2\right]_1^3$$
$$= 2\pi\left[\left(-\tfrac{81}{4} + 36 - \tfrac{27}{2}\right) - \left(-\tfrac{1}{4} + \tfrac{4}{3} - \tfrac{3}{2}\right)\right]$$
$$= 2\pi\left(\tfrac{8}{3}\right) = \tfrac{16}{3}\pi$$

20. The curves intersect when $4 - y = y^2 - 4y + 4 \quad \Leftrightarrow$

$0 = y^2 - 3y \quad \Leftrightarrow \quad 0 = y(y-3) \quad \Leftrightarrow \quad y = 0$ or 3.

The shell has radius y, circumference $2\pi y$, and height

$(4 - y) - (y^2 - 4y + 4) = -y^2 + 3y$, so

$$V = \int_0^3 2\pi y(-y^2 + 3y)\,dy = 2\pi \int_0^3 (3y^2 - y^3)\,dy$$
$$= 2\pi\left[y^3 - \tfrac{1}{4}y^4\right]_0^3 = 2\pi\left(27 - \tfrac{81}{4}\right) = 2\pi\left(\tfrac{27}{4}\right) = \tfrac{27\pi}{2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. $x^2 = 8\sqrt{x} \quad \Rightarrow \quad x^4 = 64x \quad \Rightarrow \quad x^4 - 64x = 0 \quad \Rightarrow$

$x(x^3 - 64) = 0 \quad \Rightarrow \quad x = 0$ or $x = 4$

(a) By cylindrical shells:

$$V = \int_0^4 2\pi x \left(8\sqrt{x} - x^2\right) dx$$
$$= \int_0^4 2\pi(8x^{3/2} - x^3)\, dx$$
$$= 2\pi\left[\tfrac{16}{5}x^{5/2} - \tfrac{1}{4}x^4\right]_0^4$$
$$= 2\pi\left[\left(\tfrac{512}{5} - 64\right) - 0\right] = \tfrac{384}{5}\pi$$

(b) By washers:

$$V = \int_0^{16} \pi\left[\left(\sqrt{y}\right)^2 - \left(\tfrac{1}{64}y^2\right)^2\right] dy$$
$$= \int_0^{16} \pi\left(y - \tfrac{1}{4096}y^4\right) dy = \pi\left[\tfrac{1}{2}y^2 - \tfrac{1}{20{,}480}y^5\right]_0^{16}$$
$$= \pi\left(128 - \tfrac{256}{5}\right) = \tfrac{384}{5}\pi$$

22. $x^3 = 4x^2 \quad \Rightarrow \quad x^3 - 4x^2 = 0 \quad \Rightarrow \quad x^2(x - 4) = 0 \quad \Rightarrow$

$x = 0$ or $x = 4$

(a) By washers:

$$V = \int_0^4 \pi[(4x^2)^2 - (x^3)^2]\, dx = \int_0^4 \pi(16x^4 - x^6)\, dx$$
$$= \pi\left[\frac{16}{5}x^5 - \frac{1}{7}x^7\right]_0^4 = \pi\left(\frac{16{,}384}{5} - \frac{16{,}384}{7}\right)$$
$$= \frac{32{,}768}{35}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) By cylindrical shells:

$$V = \int_0^{64} 2\pi y\left(\sqrt[3]{y} - \tfrac{1}{2}\sqrt{y}\right) dy = \int_0^{64} 2\pi\left(y^{4/3} - \tfrac{1}{2}y^{3/2}\right) dy$$

$$= 2\pi\left[\frac{3}{7}y^{7/3} - \frac{1}{2}\cdot\frac{2}{5}y^{5/2}\right]_0^{64}$$

$$= 2\pi\left(\frac{49{,}152}{7} - \frac{32{,}768}{5}\right) = \frac{32{,}768}{35}\pi$$

23. (a) The shell has radius $x - (-2) = x + 2$,

circumference $2\pi(x+2)$, and height $4x - x^2$.

(b) $V = \displaystyle\int_0^4 2\pi(x+2)(4x - x^2)\,dx$

(c) $V = \displaystyle\int_0^4 2\pi(x+2)(4x - x^2)\,dx$

$$= \int_0^4 2\pi(2x^2 - x^3 + 8x)\,dx = 2\pi\left[\frac{2}{3}x^3 - \frac{1}{4}x^4 + 4x^2\right]_0^4$$

$$= 2\pi\left[\left(\frac{128}{3} - 64 + 64\right) - 0\right] = \frac{256}{3}\pi$$

24. (a) $y = \sqrt{x} \;\Rightarrow\; x = y^2$; $y = x^3 \;\Rightarrow\; x = \sqrt[3]{y}$.

The shell has radius $y - (-1) = y + 1$,

circumference $2\pi(y+1)$, and height $\sqrt[3]{y} - y^2$.

(b) $V = \displaystyle\int_0^1 2\pi(y+1)\left(\sqrt[3]{y} - y^2\right) dy$

(c) $V = \displaystyle\int_0^1 2\pi(y+1)\left(\sqrt[3]{y} - y^2\right) dy$

$$= \int_0^1 2\pi(y^{4/3} - y^3 + y^{1/3} - y^2)\,dy$$

$$= 2\pi\left[\tfrac{3}{7}y^{7/3} - \tfrac{1}{4}y^4 + \tfrac{3}{4}y^{4/3} - \tfrac{1}{3}y^3\right]_0^1 = 2\pi\left[\left(\tfrac{3}{7} - \tfrac{1}{4} + \tfrac{3}{4} - \tfrac{1}{3}\right) - 0\right]$$

$$= \tfrac{25}{21}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. The shell has radius $3 - x$, circumference $2\pi(3 - x)$, and height $8 - x^3$.

$$V = \int_0^2 2\pi(3 - x)(8 - x^3)\,dx$$
$$= 2\pi \int_0^2 (x^4 - 3x^3 - 8x + 24)\,dx$$
$$= 2\pi \left[\tfrac{1}{5}x^5 - \tfrac{3}{4}x^4 - 4x^2 + 24x\right]_0^2$$
$$= 2\pi\left(\tfrac{32}{5} - 12 - 16 + 48\right) = 2\pi\left(\tfrac{132}{5}\right) = \tfrac{264\pi}{5}$$

26. The shell has radius $x - (-1) = x + 1$, circumference $2\pi(x + 1)$, and height $4 - 2x$.

$$V = \int_0^2 2\pi(x + 1)(4 - 2x)\,dx$$
$$= 4\pi \int_0^2 (x + 1)(2 - x)\,dx$$
$$= 4\pi \int_0^2 (-x^2 + x + 2)\,dx$$
$$= 4\pi \left[-\tfrac{1}{3}x^3 + \tfrac{1}{2}x^2 + 2x\right]_0^2$$
$$= 4\pi\left(-\tfrac{8}{3} + 2 + 4\right) = 4\pi\left(\tfrac{10}{3}\right) = \tfrac{40\pi}{3}$$

27. The shell has radius $x - 1$, circumference $2\pi(x - 1)$, and height $(4x - x^2) - 3 = -x^2 + 4x - 3$.

$$V = \int_1^3 2\pi(x - 1)(-x^2 + 4x - 3)\,dx$$
$$= 2\pi \int_1^3 (-x^3 + 5x^2 - 7x + 3)\,dx$$
$$= 2\pi\left[-\tfrac{1}{4}x^4 + \tfrac{5}{3}x^3 - \tfrac{7}{2}x^2 + 3x\right]_1^3$$
$$= 2\pi\left[\left(-\tfrac{81}{4} + 45 - \tfrac{63}{2} + 9\right) - \left(-\tfrac{1}{4} + \tfrac{5}{3} - \tfrac{7}{2} + 3\right)\right]$$
$$= 2\pi\left(\tfrac{4}{3}\right) = \tfrac{8}{3}\pi$$

28. The shell has radius $5 - x$, circumference $2\pi(5 - x)$, and height $\sqrt{x} - \frac{1}{2}x$.

$$V = \int_0^4 2\pi(5 - x)\left(\sqrt{x} - \tfrac{1}{2}x\right)\,dx = 2\pi \int_0^4 (5x^{1/2} - \tfrac{5}{2}x - x^{3/2} + \tfrac{1}{2}x^2)\,dx$$
$$= 2\pi\left[\tfrac{10}{3}x^{3/2} - \tfrac{5}{4}x^2 - \tfrac{2}{5}x^{5/2} + \tfrac{1}{6}x^3\right]_0^4 = 2\pi\left(\tfrac{80}{3} - 20 - \tfrac{64}{5} + \tfrac{32}{3}\right)$$
$$= 2\pi\left(\tfrac{68}{15}\right) = \tfrac{136\pi}{15}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. The shell has radius $2-y$, circumference $2\pi(2-y)$, and height $2-2y^2$.

$$\begin{aligned} V &= \int_0^1 2\pi(2-y)(2-2y^2)\,dy \\ &= 4\pi\int_0^1 (2-y)(1-y^2)\,dy \\ &= 4\pi\int_0^1 (y^3-2y^2-y+2)\,dy \\ &= 4\pi\left[\tfrac{1}{4}y^4-\tfrac{2}{3}y^3-\tfrac{1}{2}y^2+2y\right]_0^1 \\ &= 4\pi\left(\tfrac{1}{4}-\tfrac{2}{3}-\tfrac{1}{2}+2\right) \\ &= 4\pi\left(\tfrac{13}{12}\right) = \tfrac{13\pi}{3} \end{aligned}$$

30. The shell has radius $y-(-2)=y+2$, circumference $2\pi(y+2)$, and height $(y^2+1)-2y^2=1-y^2$.

$$\begin{aligned} V &= \int_{-1}^1 2\pi(y+2)(1-y^2)\,dy \\ &= 2\pi\int_{-1}^1 (-y^3-2y^2+y+2)\,dy \\ &= 4\pi\int_0^1 (-2y^2+2)\,dy \qquad \text{[by Theorem 5.5.7]} \\ &= 8\pi\int_0^1 (1-y^2)\,dy = 8\pi\left[y-\tfrac{1}{3}y^3\right]_0^1 \\ &= 8\pi\left(1-\tfrac{1}{3}\right) = 8\pi\left(\tfrac{2}{3}\right) = \tfrac{16\pi}{3} \end{aligned}$$

31. (a) $V = 2\pi\displaystyle\int_0^2 x(xe^{-x})\,dx = 2\pi\int_0^2 x^2e^{-x}\,dx$

(b) $V \approx 4.06300$

32. (a) $V = 2\pi\displaystyle\int_0^{\pi/4}\left(\frac{\pi}{2}-x\right)\tan x\,dx$

(b) $V \approx 2.25323$

33. (a) $V = 2\pi\displaystyle\int_{-\pi/2}^{\pi/2}(\pi-x)[\cos^4 x-(-\cos^4 x)]\,dx$

$$= 4\pi\int_{-\pi/2}^{\pi/2}(\pi-x)\,\cos^4 x\,dx$$

[or $8\pi^2\int_0^{\pi/2}\cos^4 x\,dx$ using Theorem 5.5.7]

(b) $V \approx 46.50942$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. (a) $x = \dfrac{2x}{1+x^3} \quad \Rightarrow \quad x + x^4 = 2x \quad \Rightarrow \quad x^4 - x = 0 \quad \Rightarrow$

$x(x^3 - 1) = 0 \quad \Rightarrow \quad x(x-1)(x^2+x+1) = 0 \quad \Rightarrow \quad x = 0 \text{ or } 1$

$V = 2\pi \displaystyle\int_0^1 [x - (-1)] \left(\frac{2x}{1+x^3} - x \right) dx$

(b) $V \approx 2.36164$

35. (a) $V = \int_0^{\pi} 2\pi(4-y)\sqrt{\sin y}\, dy$ (b) $V \approx 36.57476$

36. (a) $V = \int_{-3}^{3} 2\pi(5-y)\left(4 - \sqrt{y^2+7}\right) dy$ (b) $V \approx 163.02712$

37. $V = \int_0^1 2\pi x\sqrt{1+x^3}\, dx$. Let $f(x) = x\sqrt{1+x^3}$.

Then the Midpoint Rule with $n = 5$ gives

$\int_0^1 f(x)\, dx \approx \frac{1-0}{5}\,[f(0.1) + f(0.3) + f(0.5) + f(0.7) + f(0.9)]$

$\approx 0.2(2.9290)$

Multiplying by 2π gives $V \approx 3.68$.

38. $V = \int_0^{10} 2\pi x f(x)\, dx$. Let $g(x) = xf(x)$, where the values of f are obtained from the graph.

Using the Midpoint Rule with $n = 5$ gives

$$\int_0^{10} g(x)\, dx \approx \tfrac{10-0}{5}[g(1) + g(3) + g(5) + g(7) + g(9)] = 2[1f(1) + 3f(3) + 5f(5) + 7f(7) + 9f(9)]$$

$$= 2[1(4-2) + 3(5-1) + 5(4-1) + 7(4-2) + 9(4-2)]$$

$$= 2(2 + 12 + 15 + 14 + 18) = 2(61) = 122$$

Multiplying by 2π gives $V \approx 244\pi \approx 766.5$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. $\int_0^3 2\pi x^5\,dx = 2\pi\int_0^3 x(x^4)\,dx$. By the method of cylindrical shells, this integral represents the volume of the solid obtained by rotating the region $0 \le y \le x^4$, $0 \le x \le 3$ about the y-axis ($x = 0$).

40. $\int_1^3 2\pi y \ln y\,dy$. By the method of cylindrical shells, this integral represents the volume of the solid obtained by rotating the region $0 \le x \le \ln y$, $1 \le y \le 3$ about the x-axis.

41. $2\pi\displaystyle\int_1^4 \frac{y+2}{y^2}\,dy = 2\pi\int_1^4 (y+2)\left(\frac{1}{y^2}\right)dy$. By the method of cylindrical shells, this integral represents the volume of the solid obtained by rotating the region $0 \le x \le 1/y^2$, $1 \le y \le 4$ about the line $y = -2$.

42. $\int_0^1 2\pi(2-x)(3^x - 2^x)\,dx$. By the method of cylindrical shells, this integral represents the volume of the solid obtained by rotating the region $2^x \le y \le 3^x$, $0 \le x \le 1$ about the line $x = 2$.

43. From the graph, the curves intersect at $x = 0$ and $x = a \approx 2.175$, with $\dfrac{x}{x^2+1} > x^2 - 2x$ on the interval $(0, a)$. So the volume of the solid obtained by rotating the region about the y-axis is

$$V = 2\pi\int_0^a x\left[\frac{x}{x^2+1} - (x^2 - 2x)\right]dx \approx 14.450$$

44. From the graph, the curves intersect at $x = a \approx 0.906$ and $x = b \approx 2.715$, with $e^{\sin x} > x^2 - 4x + 5$ on the interval (a, b). So the volume of the solid obtained by rotating the region about the y-axis is

$$V = 2\pi\int_a^b x\left[e^{\sin x} - (x^2 - 4x + 5)\right]dx \approx 21.253$$

45. $V = 2\pi\displaystyle\int_0^{\pi/2} \left[\left(\tfrac{\pi}{2} - x\right)\left(\sin^2 x - \sin^4 x\right)\right]dx$

$\overset{\text{CAS}}{=} \frac{1}{32}\pi^3$

46. $V = 2\pi\int_0^\pi \left\{[x - (-1)](x^3 \sin x)\right\}dx$

$\overset{\text{CAS}}{=} 2\pi(\pi^4 + \pi^3 - 12\pi^2 - 6\pi + 48)$

$= 2\pi^5 + 2\pi^4 - 24\pi^3 - 12\pi^2 + 96\pi$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. (a) Use shells. Each shell has radius x, circumference $2\pi x$, and height $\dfrac{1}{1+x^2} - \dfrac{x}{2}$.

$$V = \int_0^1 2\pi x\left(\frac{1}{1+x^2} - \frac{x}{2}\right) dx.$$

(b) $$V = \int_0^1 2\pi x\left(\frac{1}{1+x^2} - \frac{x}{2}\right) dx = \int_0^1 2\pi\left(\frac{x}{1+x^2} - \frac{1}{2}x^2\right) dx = 2\pi\left[\frac{1}{2}\ln\left|1+x^2\right| - \frac{1}{6}x^3\right]_0^1$$

$$= 2\pi\left[\left(\frac{1}{2}\ln 2 - \frac{1}{6}\right) - 0\right] = \pi\left(\ln 2 - \frac{1}{3}\right)$$

48. (a) Use washers. $2 - x^2 = x^2 \quad\Rightarrow\quad 2x^2 = 2 \quad\Rightarrow\quad x = 1 \;\; [x > 0]$

$V = \int_0^1 \pi[(2-x^2)^2 - (x^2)^2]\,dx$

(b) $V = \int_0^1 \pi[(2-x^2)^2 - (x^2)^2]\,dx = \int_0^1 \pi(4-4x^2)\,dx = \pi\left[4x - 4\cdot\frac{1}{3}x^3\right]_0^1 = 4\pi\left[\left(1-\frac{1}{3}\right) - 0\right] = \frac{8}{3}\pi$

49. (a) Use disks. $V = \int_0^\pi \pi\left(\sqrt{\sin x}\right)^2\,dx$

(b) $V = \int_0^\pi \pi\left(\sqrt{\sin x}\right)^2 dx = \int_0^\pi \pi\sin x\,dx = \pi\Big[-\cos x\Big]_0^\pi = \pi[-\cos\pi - (-\cos 0)] = \pi(1+1) = 2\pi$

50. (a) Use shells. Each shell has radius x, circumference $2\pi x$, and height $[(4x - x^2) - x]$.

$4x - x^2 = x \quad\Rightarrow\quad x^2 - 3x = 0 \quad\Rightarrow\quad x(x-3) = 0 \quad\Rightarrow\quad x = 0$ or $x = 3$

$V = \int_0^3 2\pi x[(4x-x^2) - x]\,dx$

(b) $V = \int_0^3 2\pi x[(4x-x^2) - x]\,dx = \int_0^3 2\pi(3x^2 - x^3)\,dx = 2\pi\left[x^3 - \frac{1}{4}x^4\right]_0^3 = 2\pi\left[\left(27 - \frac{81}{4}\right) - 0\right] = \frac{27}{2}\pi$

51. (a) Use shells. Each shell has radius $x - (-2) = x + 2$, circumference $2\pi(x+2)$, and height $x^2 - x^3$.

$V = \int_0^{1/2} 2\pi(x+2)(x^2 - x^3)\,dx$

(b) $V = \int_0^{1/2} 2\pi(x+2)(x^2-x^3)\,dx = \int_0^{1/2} 2\pi(-x^3 - x^4 + 2x^2)\,dx = 2\pi\left[-\frac{1}{4}x^4 - \frac{1}{5}x^5 + \frac{2}{3}x^3\right]_0^{1/2}$

$= 2\pi\left[\left(-\frac{1}{64} - \frac{1}{160} + \frac{1}{12}\right) - 0\right] = \frac{59}{480}\pi$

52. (a) Use shells. Each shell has radius $3 - y$, circumference $2\pi(3-y)$, and height $[(3y - y^2) - 2]$.

$3y - y^2 = 2 \quad\Rightarrow\quad y^2 - 3y + 2 = 0 \quad\Rightarrow\quad (y-1)(y-2) = 0 \quad\Rightarrow\quad y = 1$ or $y = 2$

$V = \int_1^2 2\pi(3-y)[(3y-y^2) - 2]\,dy$

(b) $V = \int_1^2 2\pi(3-y)[(3y-y^2)-2]\,dy = \int_1^2 2\pi(y^3 - 6y^2 + 11y - 6)\,dy$

$= 2\pi\left[\frac{1}{4}y^4 - 2y^3 + \frac{11}{2}y^2 - 6y\right]_1^2 = 2\pi\left[(4 - 16 + 22 - 12) - \left(\frac{1}{4} - 2 + \frac{11}{2} - 6\right)\right] = \frac{1}{2}\pi$

53. Use shells:

$V = \int_2^4 2\pi x(-x^2 + 6x - 8)\,dx = 2\pi\int_2^4(-x^3 + 6x^2 - 8x)\,dx$

$= 2\pi\left[-\frac{1}{4}x^4 + 2x^3 - 4x^2\right]_2^4$

$= 2\pi[(-64 + 128 - 64) - (-4 + 16 - 16)]$

$= 2\pi(4) = 8\pi$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. Use disks:

$$V = \int_2^4 \pi(-x^2+6x-8)^2\,dx$$
$$= \pi\int_2^4 (x^4-12x^3+52x^2-96x+64)\,dx$$
$$= \pi\left[\tfrac{1}{5}x^5-3x^4+\tfrac{52}{3}x^3-48x^2+64x\right]_2^4$$
$$= \pi\left(\tfrac{512}{15}-\tfrac{496}{15}\right) = \tfrac{16}{15}\pi$$

55. Use washers: $y^2 - x^2 = 1 \quad\Rightarrow\quad y = \pm\sqrt{x^2 \pm 1}$

$$V = \int_{-\sqrt{3}}^{\sqrt{3}} \pi\left[(2-0)^2 - \left(\sqrt{x^2+1}-0\right)^2\right] dx$$
$$= 2\pi\int_0^{\sqrt{3}} [4-(x^2+1)]\,dx \qquad \text{[by symmetry]}$$
$$= 2\pi\int_0^{\sqrt{3}} (3-x^2)\,dx = 2\pi\left[3x-\tfrac{1}{3}x^3\right]_0^{\sqrt{3}}$$
$$= 2\pi\left(3\sqrt{3}-\sqrt{3}\right) = 4\sqrt{3}\,\pi$$

56. Use disks: $y^2 - x^2 = 1 \quad\Rightarrow\quad x = \pm\sqrt{y^2-1}$

$$V = \pi\int_1^2 \left(\sqrt{y^2-1}\right)^2 dy = \pi\int_1^2 (y^2-1)\,dy$$
$$= \pi\left[\tfrac{1}{3}y^3-y\right]_1^2 = \pi\left[\left(\tfrac{8}{3}-2\right)-\left(\tfrac{1}{3}-1\right)\right] = \tfrac{4}{3}\pi$$

57. Use disks: $x^2 + (y-1)^2 = 1 \quad\Leftrightarrow\quad x = \pm\sqrt{1-(y-1)^2}$

$$V = \pi\int_0^2 \left[\sqrt{1-(y-1)^2}\right]^2 dy = \pi\int_0^2 (2y-y^2)\,dy$$
$$= \pi\left[y^2-\tfrac{1}{3}y^3\right]_0^2 = \pi\left(4-\tfrac{8}{3}\right) = \tfrac{4}{3}\pi$$

58. Use shells:

$$V = \int_1^5 2\pi(y-1)[4-(y-3)^2]\,dy$$
$$= 2\pi\int_1^5 (y-1)(-y^2+6y-5)\,dy$$
$$= 2\pi\int_1^5 (-y^3+7y^2-11y+5)\,dy$$
$$= 2\pi\left[-\tfrac{1}{4}y^4+\tfrac{7}{3}y^3-\tfrac{11}{2}y^2+5y\right]_1^5$$
$$= 2\pi\left(\tfrac{275}{12}-\tfrac{19}{12}\right) = \tfrac{128}{3}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. $y+1=(y-1)^2 \quad \Leftrightarrow \quad y+1=y^2-2y+1 \quad \Leftrightarrow \quad 0=y^2-3y \quad \Leftrightarrow$

$0=y(y-3) \quad \Leftrightarrow \quad y=0$ or 3.

Use washers:

$$V=\pi\int_0^3 \left\{[(y+1)-(-1)]^2-[(y-1)^2-(-1)]^2\right\}dy$$

$$=\pi\int_0^3 [(y+2)^2-(y^2-2y+2)^2]\,dy$$

$$=\pi\int_0^3 [(y^2+4y+4)-(y^4-4y^3+8y^2-8y+4)]\,dy=\pi\int_0^3(-y^4+4y^3-7y^2+12y)\,dy$$

$$=\pi\left[-\tfrac{1}{5}y^5+y^4-\tfrac{7}{3}y^3+6y^2\right]_0^3=\pi\left(-\tfrac{243}{5}+81-63+54\right)=\tfrac{117}{5}\pi$$

60. Use cylindrical shells to find the volume V.

$$V=\int_0^1 2\pi(a-x)(2x)\,dx=4\pi\int_0^1(ax-x^2)\,dx$$

$$=4\pi\left[\tfrac{1}{2}ax^2-\tfrac{1}{3}x^3\right]_0^1=4\pi\left(\tfrac{1}{2}a-\tfrac{1}{3}\right)$$

Now solve for a in terms of V:

$$V=4\pi\left(\tfrac{1}{2}a-\tfrac{1}{3}\right) \quad \Leftrightarrow \quad \frac{V}{4\pi}=\frac{1}{2}a-\frac{1}{3} \quad \Leftrightarrow \quad \frac{1}{2}a=\frac{V}{4\pi}+\frac{1}{3} \quad \Leftrightarrow$$

$$a=\frac{V}{2\pi}+\frac{2}{3}$$

61. Use shells:

$$V=2\int_0^r 2\pi x\sqrt{r^2-x^2}\,dx=-2\pi\int_0^r(r^2-x^2)^{1/2}(-2x)\,dx$$

$$=\left[-2\pi\cdot\tfrac{2}{3}(r^2-x^2)^{3/2}\right]_0^r=-\tfrac{4}{3}\pi(0-r^3)=\tfrac{4}{3}\pi r^3$$

62. $V=\int_{R-r}^{R+r} 2\pi x\cdot 2\sqrt{r^2-(x-R)^2}\,dx$

$$=\int_{-r}^{r} 4\pi(u+R)\sqrt{r^2-u^2}\,du \qquad [\text{let } u=x-R]$$

$$=4\pi R\int_{-r}^{r}\sqrt{r^2-u^2}\,du+4\pi\int_{-r}^{r}u\sqrt{r^2-u^2}\,du$$

The first integral is the area of a semicircle of radius r, that is, $\frac{1}{2}\pi r^2$, and the second is zero since the integrand is an odd function. Thus, $V=4\pi R\left(\frac{1}{2}\pi r^2\right)+4\pi\cdot 0=2\pi^2Rr^2$.

63. $V=2\pi\displaystyle\int_0^r x\left(-\frac{h}{r}x+h\right)dx=2\pi h\int_0^r\left(-\frac{x^2}{r}+x\right)dx$

$$=2\pi h\left[-\frac{x^3}{3r}+\frac{x^2}{2}\right]_0^r=2\pi h\,\frac{r^2}{6}=\frac{\pi r^2 h}{3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. By symmetry, the volume of a napkin ring obtained by drilling a hole of radius r through a sphere with radius R is twice the volume obtained by rotating the area above the x-axis and below the curve $y = \sqrt{R^2 - x^2}$ (the equation of the top half of the cross-section of the sphere), between $x = r$ and $x = R$, about the y-axis. This volume is equal to

$$2\int_{\text{inner radius}}^{\text{outer radius}} 2\pi r h\,dx = 2\cdot 2\pi\int_r^R x\sqrt{R^2-x^2}\,dx = 4\pi\left[-\tfrac{1}{3}\left(R^2-x^2\right)^{3/2}\right]_r^R = \tfrac{4}{3}\pi(R^2-r^2)^{3/2}$$

But by the Pythagorean Theorem, $R^2 - r^2 = \left(\frac{1}{2}h\right)^2$, so the volume of the napkin ring is $\frac{4}{3}\pi\left(\frac{1}{2}h\right)^3 = \frac{1}{6}\pi h^3$, which is independent of both R and r; that is, the amount of wood in a napkin ring of height h is the same regardless of the size of the sphere used. Note that most of this calculation has been done already, but with more difficulty, in Exercise 6.2.84.

Another solution: The height of the missing cap is the radius of the sphere minus half the height of the cut-out cylinder, that is, $R - \frac{1}{2}h$. Using Exercise 6.2.61,

$$V_{\text{napkin ring}} = V_{\text{sphere}} - V_{\text{cylinder}} - 2V_{\text{cap}} = \tfrac{4}{3}\pi R^3 - \pi r^2 h - 2\cdot\tfrac{\pi}{3}\left(R-\tfrac{1}{2}h\right)^2\left[3R-\left(R-\tfrac{1}{2}h\right)\right] = \tfrac{1}{6}\pi h^3$$

6.4 Work

1. The force exerted by the weight lifter is $F = mg = (200\text{ kg})(9.8\text{ m/s}^2) = 1960\text{ N}$. The work done by the weight lifter in lifting the weight from 1.5 m to 2.0 m above the ground is then

$$W = Fd = (1960\text{ N})(2.0\text{ m} - 1.5\text{ m}) = (1960\text{ N})(0.5\text{ m}) = 980\text{ N-m} = 980\text{ J}$$

2. $W = Fd = (1100\text{ lb})(35\text{ ft}) = 38{,}500$ ft-lb.

3. $W = \int_a^b f(x)\,dx = \int_1^{10} 5x^{-2}\,dx = 5\left[-x^{-1}\right]_1^{10} = 5\left(-\frac{1}{10}+1\right) = 4.5$ ft-lb

4. $W = \displaystyle\int_a^b f(x)\,dx = \int_4^{16} 4\sqrt{x}\,dx = 4\left[\tfrac{2}{3}x^{3/2}\right]_4^{16} = \tfrac{8}{3}(64-8) = \tfrac{448}{3}\text{ N-m} = \tfrac{448}{3}\text{ J}.$

5. The force function is given by $F(x)$ (in newtons) and the work (in joules) is the area under the curve, given by $\int_0^8 F(x)\,dx = \int_0^4 F(x)\,dx + \int_4^8 F(x)\,dx = \frac{1}{2}(4)(30) + (4)(30) = 180$ J.

6. $W = \int_4^{20} f(x)\,dx \approx M_4 = \Delta x[f(6)+f(10)+f(14)+f(18)] = \frac{20-4}{4}[5.8+8.8+8.2+5.2] = 4(28) = 112$ J

7. According to Hooke's Law, the force required to maintain a spring stretched x units beyond its natural length (or compressed x units less than its natural length) is proportional to x, that is, $f(x) = kx$. Here, the amount stretched is 4 in. $= \frac{1}{3}$ ft and the force is 10 lb. Thus, $10 = k\left(\frac{1}{3}\right) \quad\Rightarrow\quad k = 30$ lb/ft, and $f(x) = 30x$. The work done in stretching the spring from its natural length to 6 in. $= \frac{1}{2}$ ft beyond its natural length is $W = \int_0^{1/2} 30x\,dx = \left[15x^2\right]_0^{1/2} = \frac{15}{4}$ ft-lb.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. According to Hooke's Law, the force required to maintain a spring stretched x units beyond its natural length (or compressed x units less than its natural length) is proportional to x, that is, $f(x) = kx$. Here, the amount compressed is $40 - 30 = 10$ cm $= 0.1$ m and the force is 60 N. Thus, $60 = k(0.1) \quad \Rightarrow \quad k = 600$ N/m, and $f(x) = 600x$. The work required to compress the spring 0.1 m is $W = \int_0^{0.1} 600x\,dx = \left[300x^2\right]_0^{0.1} = 300(0.01) = 3$ N-m (or J). The work required to compress the spring $40 - 25 = 15$ cm $= 0.15$ m is $W = \int_0^{0.15} 600x\,dx = \left[300x^2\right]_0^{0.15} = 300(0.0225) = 6.75$ J.

9. (a) If $\int_0^{0.12} kx\,dx = 2$ J, then $2 = \left[\frac{1}{2}kx^2\right]_0^{0.12} = \frac{1}{2}k(0.0144) = 0.0072k$ and $k = \frac{2}{0.0072} = \frac{2500}{9} \approx 277.78$ N/m.

Thus, the work needed to stretch the spring from 35 cm to 40 cm is

$\int_{0.05}^{0.10} \frac{2500}{9}x\,dx = \left[\frac{1250}{9}x^2\right]_{1/20}^{1/10} = \frac{1250}{9}\left(\frac{1}{100} - \frac{1}{400}\right) = \frac{25}{24} \approx 1.04$ J.

(b) $f(x) = kx$, so $30 = \frac{2500}{9}x$ and $x = \frac{270}{2500}$ m $= 10.8$ cm

10. If $12 = \int_0^1 kx\,dx = \left[\frac{1}{2}kx^2\right]_0^1 = \frac{1}{2}k$, then $k = 24$ lb/ft and the work required is

$\int_0^{3/4} 24x\,dx = \left[12x^2\right]_0^{3/4} = 12 \cdot \frac{9}{16} = \frac{27}{4} = 6.75$ ft-lb.

11. The distance from 20 cm to 30 cm is 0.1 m, so with $f(x) = kx$, we get $W_1 = \int_0^{0.1} kx\,dx = k\left[\frac{1}{2}x^2\right]_0^{0.1} = \frac{1}{200}k$.

Now $W_2 = \int_{0.1}^{0.2} kx\,dx = k\left[\frac{1}{2}x^2\right]_{0.1}^{0.2} = k\left(\frac{4}{200} - \frac{1}{200}\right) = \frac{3}{200}k$. Thus, $W_2 = 3W_1$.

12. Let L be the natural length of the spring in meters. Then

$6 = \int_{0.10-L}^{0.12-L} kx\,dx = \left[\frac{1}{2}kx^2\right]_{0.10-L}^{0.12-L} = \frac{1}{2}k\left[(0.12 - L)^2 - (0.10 - L)^2\right]$ and

$10 = \int_{0.12-L}^{0.14-L} kx\,dx = \left[\frac{1}{2}kx^2\right]_{0.12-L}^{0.14-L} = \frac{1}{2}k\left[(0.14 - L)^2 - (0.12 - L)^2\right]$.

Simplifying gives us $12 = k(0.0044 - 0.04L)$ and $20 = k(0.0052 - 0.04L)$. Subtracting the first equation from the second gives $8 = 0.0008k$, so $k = 10{,}000$. Now the second equation becomes $20 = 52 - 400L$, so $L = \frac{32}{400}$ m $= 8$ cm.

In Exercises 13–22, n is the number of subintervals of length Δx, and x_i^* is a sample point in the ith subinterval $[x_{i-1}, x_i]$.

13. (a) The portion of the rope from x ft to $(x + \Delta x)$ ft below the top of the building weighs $\frac{1}{2}\,\Delta x$ lb and must be lifted x_i^* ft, so its contribution to the total work is $\frac{1}{2}x_i^*\,\Delta x$ ft-lb. The total work is

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} \frac{1}{2}x_i^*\,\Delta x = \int_0^{50} \frac{1}{2}x\,dx = \left[\frac{1}{4}x^2\right]_0^{50} = \frac{2500}{4} = 625 \text{ ft-lb}$$

Notice that the exact height of the building does not matter (as long as it is more than 50 ft).

(b) When half the rope is pulled to the top of the building, the work to lift the top half of the rope is

$W_1 = \int_0^{25} \frac{1}{2}x\,dx = \left[\frac{1}{4}x^2\right]_0^{25} = \frac{625}{4}$ ft-lb. The bottom half of the rope is lifted 25 ft and the work needed to accomplish that is $W_2 = \int_{25}^{50} \frac{1}{2} \cdot 25\,dx = \frac{25}{2}\left[x\right]_{25}^{50} = \frac{625}{2}$ ft-lb. The total work done in pulling half the rope to the top of the building is $W = W_1 + W_2 = \frac{625}{2} + \frac{625}{4} = \frac{3}{4} \cdot 625 = \frac{1875}{4}$ ft-lb.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. (a) The 60 ft cable weighs 180 lb, or 3 lb/ft. If we divide the cable into n equal parts of length $\Delta x = 60/n$ ft, then for large n, all points in the ith part are lifted by approximately the same amount. Choose a representative distance from the winch in the ith part of the cable, say x_i^*. If $x_i^* < 25$ ft, then the ith part has to be lifted roughly x_i^* ft. If $x_i^* \geq 25$ ft, then the ith part has to be lifted 25 ft. The ith part weighs $(3 \text{ lb/ft})(\Delta x \text{ ft}) = 3\,\Delta x$ lb, so the work done in lifting it is $(3\,\Delta x)x_i^*$ if $x_i^* < 25$ ft and $(3\,\Delta x)(25) = 75\,\Delta x$ if $x_i^* \geq 25$ ft. The work of lifting the top 25 ft of the cable is

$W_1 = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n_1} 3x_i^*\,\Delta x = \int_0^{25} 3x\,dx = \left[\frac{3}{2}x^2\right]_0^{25} = \frac{3}{2}(625) = 937.5$ ft-lb. Here n_1 represents the number of parts of the cable in the top 25 ft. The work of lifting the bottom 35 ft of the cable is

$W_2 = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n_2} 75\,\Delta x = \int_{25}^{60} 75\,dx = 75(60-25) = 2625$ ft-lb, where n_2 represents the number of small parts in the bottom 35 feet of the cable. The total work done is $W = W_1 + W_2 = 937.5 + 2625 = 3562.5$ ft-lb.

(b) Once x feet of cable have been wound up by the winch, there is $(60 - x)$ ft of cable still hanging from the winch. That portion of the cable weighs $3(60 - x)$ lb. Lifting it Δx feet requires $3(60 - x)\,\Delta x$ ft-lb of work. Thus, the total work needed to lift the cable 25 ft is $W = \int_0^{25} 3(60 - x)\,dx = \left[180x - \frac{3}{2}x^2\right]_0^{25} = 4500 - 937.5 = 3562.5$ ft-lb.

15. The work needed to lift the cable is $\lim\limits_{n\to\infty} \sum_{i=1}^{n} 2x_i^*\,\Delta x = \int_0^{500} 2x\,dx = \left[x^2\right]_0^{500} = 250{,}000$ ft-lb. The work needed to lift the coal is 800 lb · 500 ft = 400,000 ft-lb. Thus, the total work required is 250,000 + 400,000 = 650,000 ft-lb.

16. *Assumptions*:

1. After lifting, the chain is L-shaped, with 4 m of the chain lying along the ground.
2. The chain slides effortlessly and without friction along the ground while its end is lifted.
3. The weight density of the chain is constant throughout its length and therefore equals $(8 \text{ kg/m})(9.8 \text{ m/s}^2) = 78.4$ N/m.

The part of the chain x m from the lifted end is raised $6 - x$ m if $0 \leq x \leq 6$ m, and it is lifted 0 m if $x > 6$ m.

Thus, the work needed is

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} (6 - x_i^*) \cdot 78.4\,\Delta x = \int_0^6 (6 - x)78.4\,dx = 78.4\left[6x - \tfrac{1}{2}x^2\right]_0^6 = (78.4)(18) = 1411.2 \text{ J}$$

17. The chain's weight density is $\dfrac{25 \text{ lb}}{10 \text{ ft}} = 2.5$ lb/ft. The part of the chain x ft below the ceiling (for $5 \leq x \leq 10$) has to be lifted $2(x - 5)$ ft, so the work needed to lift the ith subinterval of the chain is $2(x_i^* - 5)(2.5\,\Delta x)$. The total work needed is

$$\begin{aligned} W &= \lim_{n\to\infty} \sum_{i=1}^{n} 2(x_i^* - 5)(2.5)\,\Delta x = \int_5^{10} [2(x-5)(2.5)]\,dx = 5\int_5^{10} (x-5)\,dx \\ &= 5\left[\tfrac{1}{2}x^2 - 5x\right]_5^{10} = 5\left[(50-50) - \left(\tfrac{25}{2} - 25\right)\right] = 5\left(\tfrac{25}{2}\right) = 62.5 \text{ ft-lb} \end{aligned}$$

18. The work needed to lift the model rocket itself is $Fd = (mg)d = (0.4 \text{ kg})(9.8 \text{ m/s}^2)(20 \text{ m}) = 78.4$ N-m $= 78.4$ J. At time t (in seconds) the rocket is $x_i^* = 4t$ m above the ground, but it now holds only $(0.75 - 0.15t)$ kg of rocket fuel. In terms

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

of distance, the rocket holds $\left[0.75 - 0.15\left(\frac{1}{4}x_i^*\right)\right]$ kg of rocket fuel when it is x_i^* m above the ground. Moving this mass of rocket fuel a distance of Δx m requires $(9.8)(0.75 - 0.0375x_i^*)\,\Delta x$ J of work. Thus, the work needed to lift the rocket fuel is

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} 9.8(0.75 - 0.0375x_i^*)\,\Delta x = \int_0^{20} 9.8(0.75 - 0.0375x)\,dx$$
$$= 9.8\Big[0.75x - 0.01875x^2\Big]_0^{20} = 9.8[(15 - 7.5) - 0] = 73.5 \text{ J}$$

Adding the work of lifting the model rocket itself gives a total of 151.9 J of work.

19. At a height of x meters ($0 \le x \le 12$), the mass of the rope is $(0.8 \text{ kg/m})(12 - x \text{ m}) = (9.6 - 0.8x)$ kg and the mass of the water is $\left(\frac{36}{12} \text{ kg/m}\right)(12 - x \text{ m}) = (36 - 3x)$ kg. The mass of the bucket is 10 kg, so the total mass is $(9.6 - 0.8x) + (36 - 3x) + 10 = (55.6 - 3.8x)$ kg, and hence, the total force is $9.8(55.6 - 3.8x)$ N. The work needed to lift the bucket Δx m through the ith subinterval of $[0, 12]$ is $9.8(55.6 - 3.8x_i^*)\Delta x$, so the total work is

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} 9.8(55.6 - 3.8x_i^*)\,\Delta x = \int_0^{12} (9.8)(55.6 - 3.8x)\,dx = 9.8\Big[55.6x - 1.9x^2\Big]_0^{12} = 9.8(393.6) \approx 3857 \text{ J}$$

20. A horizontal cylindrical slice of water Δx ft thick has a volume of $\pi r^2 h = \pi \cdot 12^2 \cdot \Delta x$ ft^3 and weighs about $\left(62.5 \text{ lb/ft}^3\right)\left(144\pi\,\Delta x \text{ ft}^3\right) = 9000\pi\,\Delta x$ lb. If the slice lies x_i^* ft below the edge of the pool (where $1 \le x_i^* \le 5$), then the work needed to pump it out is about $9000\pi x_i^*\,\Delta x$. Thus,

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} 9000\pi x_i^*\,\Delta x = \int_1^5 9000\pi x\,dx = \left[4500\pi x^2\right]_1^5 = 4500\pi(25 - 1) = 108{,}000\pi \text{ ft-lb}$$

21. A "slice" of water Δx m thick and lying at a depth of x_i^* m (where $0 \le x_i^* \le \frac{1}{2}$) has volume $(2 \times 1 \times \Delta x)$ m^3, a mass of $2000\,\Delta x$ kg, weighs about $(9.8)(2000\,\Delta x) = 19{,}600\,\Delta x$ N, and thus requires about $19{,}600x_i^*\,\Delta x$ J of work for its removal. So $W = \lim_{n\to\infty} \sum_{i=1}^{n} 19{,}600x_i^*\,\Delta x = \int_0^{1/2} 19{,}600x\,dx = \left[9800x^2\right]_0^{1/2} = 2450$ J.

22. We use a vertical coodinate x measured from the center of the water tank. The top and bottom of the tank have coordinates $x = -12$ ft and $x = 12$ ft, respectively.

A thin horizontal slice of water at coordinate x is a disk of radius $\sqrt{12^2 - x^2}$ as shown in the figure. The disk has area $\pi r^2 = \pi(12^2 - x^2)$, so if the slice has thickness Δx, the slice has volume $\pi(12^2 - x^2)\,\Delta x$ and weight $62.5\pi(12^2 - x^2)\,\Delta x$. The work needed to raise this water from ground level (coordinate 72) to coordinate x, a distance of $(72 - x)$ ft, is $62.5\pi(12^2 - x^2)(72 - x)\,\Delta x$ ft-lb. The total work needed to fill the tank is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

approximated by a Riemann sum $\sum_{i=1}^{n} 62.5\pi[(12^2-(x_i^*)^2)](72-x_i^*)\,\Delta x$. Thus, the total work is

$$\begin{aligned}
W &= \lim_{n\to\infty}\sum_{i=1}^{n} 62.5\pi[(12^2-(x_i^*)^2)](72-x_i^*)\,\Delta x = \int_{-12}^{12} 62.5\pi(12^2-x^2)(72-x)\,dx \\
&= 62.5\pi\int_{-12}^{12}[\underbrace{72(12^2-x^2)}_{\text{even function}} - \underbrace{x(12^2-x^2)}_{\text{odd function}}]\,dx = 62.5\pi(2)\int_0^{12} 72(12^2-x^2)\,dx \qquad \text{[by Theorem 5.5.7]} \\
&= 125\pi(72)\left[12^2x-\tfrac{1}{3}x^3\right]_0^{12} = 9000\pi\left(12^3-\tfrac{1}{3}\cdot 12^3\right) = 9000\pi\left(\tfrac{2}{3}\cdot 12^3\right) \\
&= 10{,}368{,}000\pi \text{ ft-lb}
\end{aligned}$$

The 1.5 horsepower pump does $1.5(550) = 825$ ft-lb of work per second. To fill the tank, it will take

$\dfrac{10{,}368{,}000\pi \text{ ft-lb}}{825 \text{ ft-lb/s}} \approx 39{,}481 \text{ s} \approx 10.97$ hours.

23. A rectangular "slice" of water Δx m thick and lying x m above the bottom has width x m and volume $8x\,\Delta x$ m^3. It weighs about $(9.8\times 1000)(8x\,\Delta x)$ N, and must be lifted $(5-x)$ m by the pump, so the work needed is about $(9.8\times 10^3)(5-x)(8x\,\Delta x)$ J. The total work required is

$$\begin{aligned}
W &\approx \int_0^3 (9.8\times 10^3)(5-x)8x\,dx = (9.8\times 10^3)\int_0^3 (40x-8x^2)\,dx = (9.8\times 10^3)\left[20x^2-\tfrac{8}{3}x^3\right]_0^3 \\
&= (9.8\times 10^3)(180-72) = (9.8\times 10^3)(108) = 1058.4\times 10^3 \approx 1.06\times 10^6 \text{ J}
\end{aligned}$$

24. Let y measure depth (in meters) below the center of the spherical tank, so that $y=-3$ at the top of the tank and $y=-4$ at the spigot. A horizontal disk-shaped "slice" of water Δy m thick and lying at coordinate y has radius $\sqrt{9-y^2}$ m and volume $\pi r^2\Delta y = \pi(9-y^2)\,\Delta y$ m^3. It weighs about $(9.8\times 1000)\pi(9-y^2)\,\Delta y$ N and must be lifted $(y+4)$ m by the pump, so the work needed to pump it out is about $(9.8\times 10^3)(y+4)\pi(9-y^2)\,\Delta y$ J. The total work required is

$$\begin{aligned}
W &\approx \int_{-3}^{3}(9.8\times 10^3)(y+4)\pi(9-y^2)\,dy = (9.8\times 10^3)\pi\int_{-3}^{3}[y(9-y^2)+4(9-y^2)]\,dy \\
&= (9.8\times 10^3)\pi(2)(4)\int_0^3 (9-y^2)\,dy \qquad \text{[by Theorem 5.5.7]} \\
&= (78.4\times 10^3)\pi\left[9y-\tfrac{1}{3}y^3\right]_0^3 = (78.4\times 10^3)\pi(18) = 1{,}411{,}200\pi \approx 4.43\times 10^6 \text{ J}
\end{aligned}$$

25. Let x measure depth (in feet) below the spout at the top of the tank. A horizontal disk-shaped "slice" of water Δx ft thick and lying at coordinate x has radius $\frac{3}{8}(16-x)$ ft $(\star)$ and volume $\pi r^2\Delta x = \pi\cdot\frac{9}{64}(16-x)^2\,\Delta x$ ft^3. It weighs about $(62.5)\frac{9\pi}{64}(16-x)^2\,\Delta x$ lb and must be lifted x ft by the pump, so the work needed to pump it out is about $(62.5)x\,\frac{9\pi}{64}(16-x)^2\,\Delta x$ ft-lb. The total work required is

$$\begin{aligned}
W &\approx \int_0^8 (62.5)x\,\tfrac{9\pi}{64}(16-x)^2\,dx = (62.5)\tfrac{9\pi}{64}\int_0^8 x(256-32x+x^2)\,dx \\
&= (62.5)\tfrac{9\pi}{64}\int_0^8 (256x-32x^2+x^3)\,dx = (62.5)\tfrac{9\pi}{64}\left[128x^2-\tfrac{32}{3}x^3+\tfrac{1}{4}x^4\right]_0^8 \\
&= (62.5)\frac{9\pi}{64}\left(\frac{11{,}264}{3}\right) = 33{,}000\pi \approx 1.04\times 10^5 \text{ ft-lb}
\end{aligned}$$

$(\star)$ From similar triangles, $\dfrac{d}{8-x} = \dfrac{3}{8}$.

So $r = 3 + d = 3 + \frac{3}{8}(8-x)$

$= \dfrac{3(8)}{8} + \dfrac{3}{8}(8-x)$

$= \frac{3}{8}(16-x)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. Let x measure the distance (in feet) above the bottom of the tank. A horizontal "slice" of water Δx ft thick and lying at coordinate x has volume $10(2x)\,\Delta x$ ft^3. It weighs about $(62.5)20x\,\Delta x$ lb and must be lifted $(6-x)$ ft by the pump, so the work needed to pump it out is about $(62.5)(6-x)20x\,\Delta x$ ft-lb. The total work required is

$W \approx \int_0^6 (62.5)(6-x)20x\,dx = 1250\int_0^6 (6x-x^2)\,dx = 1250\big[3x^2 - \frac{1}{3}x^3\big]_0^6 = 1250(36) = 45{,}000$ ft-lb.

27. If only 4.7×10^5 J of work is done, then only the water above a certain level (call it h) will be pumped out. So we use the same formula as in Exercise 23, except that the work is fixed, and we are trying to find the lower limit of integration:

$4.7\times 10^5 \approx \int_h^3 (9.8\times 10^3)(5-x)8x\,dx = \big(9.8\times 10^3\big)\big[20x^2 - \frac{8}{3}x^3\big]_h^3 \quad \Leftrightarrow$

$\frac{4.7}{9.8}\times 10^2 \approx 48 = \big(20\cdot 3^2 - \frac{8}{3}\cdot 3^3\big) - \big(20h^2 - \frac{8}{3}h^3\big) \quad \Leftrightarrow$

$2h^3 - 15h^2 + 45 = 0$. To find the solution of this equation, we plot $2h^3 - 15h^2 + 45$ between $h = 0$ and $h = 3$. We see that the equation is satisfied for $h \approx 2.0$. So the depth of water remaining in the tank is about 2.0 m.

28. The only changes needed in the solution for Exercise 24 are: (1) change the lower limit from -3 to 0 and (2) change 1000 to 900.

$$\begin{aligned} W &\approx \int_0^3 (9.8\times 900)(y+4)\pi(9-y^2)\,dy = (9.8\times 900)\,\pi\int_0^3 (9y - y^3 + 36 - 4y^2)\,dy \\ &= (9.8\times 900)\pi\left[\tfrac{9}{2}y^2 - \tfrac{1}{4}y^4 + 36y - \tfrac{4}{3}y^3\right]_0^3 = (9.8\times 900)\pi(92.25) = 813{,}645\pi \\ &\approx 2.56\times 10^6 \text{ J} \quad \text{[about 58\% of the work in Exercise 24]} \end{aligned}$$

29. $V = \pi r^2 x$, so V is a function of x and P can also be regarded as a function of x. If $V_1 = \pi r^2 x_1$ and $V_2 = \pi r^2 x_2$, then

$$\begin{aligned} W &= \int_{x_1}^{x_2} F(x)\,dx = \int_{x_1}^{x_2} \pi r^2 P(V(x))\,dx = \int_{x_1}^{x_2} P(V(x))\,dV(x) \qquad \text{[Let } V(x) = \pi r^2 x\text{, so } dV(x) = \pi r^2\,dx.\text{]} \\ &= \int_{V_1}^{V_2} P(V)\,dV \quad \text{by the Substitution Rule.} \end{aligned}$$

30. 160 lb/in^2 $= 160\cdot 144$ lb/ft^2, 100 in^3 $= \frac{100}{1728}$ ft^3, and 800 in^3 $= \frac{800}{1728}$ ft^3.

$k = PV^{1.4} = (160\cdot 144)\big(\frac{100}{1728}\big)^{1.4} = 23{,}040\big(\frac{25}{432}\big)^{1.4} \approx 426.5$. Therefore, $P \approx 426.5V^{-1.4}$ and

$$W = \int_{100/1728}^{800/1728} 426.5V^{-1.4}\,dV = 426.5\left[\frac{1}{-0.4}V^{-0.4}\right]_{25/432}^{25/54} = (426.5)(2.5)\left[\big(\tfrac{432}{25}\big)^{0.4} - \big(\tfrac{54}{25}\big)^{0.4}\right] \approx 1.88\times 10^3 \text{ ft-lb.}$$

31.
$$\begin{aligned} W &= \int_{x_1}^{x_2} f(x)\,dx = \int_{t_1}^{t_2} f(s(t))\,v(t)\,dt \qquad \begin{bmatrix} x = s(t), \\ dx = v(t)\,dt \end{bmatrix} \\ &= \int_{t_1}^{t_2} m\,a(t)\,v(t)\,dt = \int_{v_1}^{v_2} m\,u\,du \qquad \begin{bmatrix} u = v(t), \\ du = a(t)\,dt \end{bmatrix} \\ &= \big[\tfrac{1}{2}mu^2\big]_{v_1}^{v_2} = \tfrac{1}{2}mv_2^2 - \tfrac{1}{2}mv_1^2 \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. The mass of the bowling ball is $\dfrac{12\text{ lb}}{32\text{ ft/s}^2} = \dfrac{3}{8}$ slug. Converting 20 mi/h to ft/s^2 gives us

$\dfrac{20\text{ mi}}{\text{h}} \cdot \dfrac{5280\text{ ft}}{1\text{ mi}} \cdot \dfrac{1\text{ h}}{3600\text{ s}^2} = \dfrac{88}{3}$ ft/s^2. From part (a) with $v_1 = 0$ and $v_2 = \frac{88}{3}$, the work required to hurl the bowling ball is

$W = \frac{1}{2} \cdot \frac{3}{8}\left(\frac{88}{3}\right)^2 - \frac{1}{2} \cdot \frac{3}{8}(0)^2 = \frac{484}{3} = 161.\overline{3}$ ft-lb.

33. The work required to move the 800-kg roller coaster car is

$W = \int_0^{60} (5.7x^2 + 1.5x)\,dx = \left[1.9x^3 + 0.75x^2\right]_0^{60} = 410{,}400 + 2700 = 413{,}100$ J.

Using Exercise 31 with $v_1 = 0$, we get $W = \frac{1}{2}mv_2^2 \quad \Rightarrow \quad v_2 = \sqrt{\dfrac{2W}{m}} = \sqrt{\dfrac{2(413{,}100)}{800}} \approx 32.14$ m/s.

34. $W = \int_1^2 \cos\left(\frac{1}{3}\pi x\right) dx = \frac{3}{\pi}\left[\sin\left(\frac{1}{3}\pi x\right)\right]_1^2 = \frac{3}{\pi}\left(\frac{\sqrt{3}}{2} - \frac{\sqrt{3}}{2}\right) = 0$ N·m $= 0$ J.

Interpretation: From $x = 1$ to $x = \frac{3}{2}$, the force does work equal to $\int_1^{3/2} \cos\left(\frac{1}{3}\pi x\right) dx = \frac{3}{\pi}\left(1 - \frac{\sqrt{3}}{2}\right)$ J in accelerating the particle and increasing its kinetic energy. From $x = \frac{3}{2}$ to $x = 2$, the force opposes the motion of the particle, decreasing its kinetic energy. This is negative work, equal in magnitude but opposite in sign to the work done from $x = 1$ to $x = \frac{3}{2}$.

35. (a) $W = \displaystyle\int_a^b F(r)\,dr = \int_a^b G\frac{m_1 m_2}{r^2}\,dr = Gm_1m_2\left[\frac{-1}{r}\right]_a^b = Gm_1m_2\left(\frac{1}{a} - \frac{1}{b}\right)$

(b) By part (a), $W = GMm\left(\dfrac{1}{R} - \dfrac{1}{R + 1{,}000{,}000}\right)$ where $M =$ mass of the earth in kg, $R =$ radius of the earth in m, and $m =$ mass of satellite in kg. (Note that 1000 km $= 1{,}000{,}000$ m.) Thus,

$$W = (6.67 \times 10^{-11})(5.98 \times 10^{24})(1000) \times \left(\frac{1}{6.37 \times 10^6} - \frac{1}{7.37 \times 10^6}\right) \approx 8.50 \times 10^9 \text{ J}$$

36. (a) Assume the pyramid has smooth sides. From the figure for $0 \le x \le 378$, an equation for the side is $y = \frac{-481}{378}x + 481 \quad \Leftrightarrow$ $x = -\frac{378}{481}(y - 481)$. The horizontal length of a cross-section is $2x$ and the area of a cross-section is $A = (2x)^2 = 4x^2 = 4\,\dfrac{378^2}{481^2}(y - 481)^2$. A slice of thickness Δy at height y has volume $\Delta V = A\,\Delta y$ ft^3 and weight $150\,\Delta V$ lb, so the work needed to build the pyramid was

$$W_1 = \int_0^{481} 150y \cdot 4\,\frac{378^2}{481^2}(y - 481)^2\,dy = 600\,\frac{378^2}{481^2}\int_0^{481} (y^3 - 2 \cdot 481y^2 + 481^2 y)\,dy$$

$$= 600\,\frac{378^2}{481^2}\left[\frac{1}{4}y^4 - \frac{2 \cdot 481}{3}y^3 + \frac{481^2}{2}y^2\right]_0^{481} = 600\,\frac{378^2}{481^2}\left(\frac{481^4}{4} - \frac{2 \cdot 481^4}{3} + \frac{481^4}{2}\right)$$

$$= 600\,\frac{378^2}{481^2}\,\frac{481^4}{12} = 50 \cdot 378^2 \cdot 481^2 \approx 1.653 \times 10^{12} \text{ ft-lb}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Work done $= W_2 = \dfrac{10\text{ h}}{\text{day}} \cdot \dfrac{340\text{ days}}{\text{year}} \cdot \dfrac{20\text{ yr}}{1\text{ laborer}} \cdot \dfrac{200\text{ ft-lb}}{\text{hour}} = 1.36 \times 10^7 \dfrac{\text{ft-lb}}{\text{laborer}}$. Dividing W_1 by W_2 gives us about 121,536 laborers.

6.5 Average Value of a Function

1. $f_{\text{avg}} = \frac{1}{b-a}\int_a^b f(x)\,dx = \frac{1}{2-(-1)}\int_{-1}^{2}(3x^2+8x)\,dx = \frac{1}{3}[x^3+4x^2]_{-1}^{2} = \frac{1}{3}[(8+16)-(-1+4)] = 7$

2. $f_{\text{avg}} = \frac{1}{b-a}\int_a^b f(x)\,dx = \frac{1}{4-0}\int_0^4 \sqrt{x}\,dx = \frac{1}{4}\left[\frac{2}{3}x^{3/2}\right]_0^4 = \frac{1}{4}\left(\frac{2}{3}\cdot 8\right) = \frac{4}{3}$

3. $g_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b g(x)\,dx = \dfrac{1}{\pi/2-(-\pi/2)}\int_{-\pi/2}^{\pi/2} 3\cos x\,dx = \dfrac{3\cdot 2}{\pi}\int_0^{\pi/2}\cos x\,dx$ [by Theorem 5.5.7(a)]

$= \frac{6}{\pi}\Big[\sin x\Big]_0^{\pi/2} = \frac{6}{\pi}(1-0) = \frac{6}{\pi}$

4. $f_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b f(z)\,dz = \dfrac{1}{4-1}\int_1^4 \dfrac{e^{1/z}}{z^2}\,dz = \dfrac{1}{3}\int_1^{1/4} e^u\,(-du)$ $\left[\begin{array}{l} u = 1/z, \\ du = -1/z^2\,dz \end{array}\right]$

$= -\dfrac{1}{3}\Big[e^u\Big]_1^{1/4} = -\dfrac{1}{3}(e^{1/4}-e) = \dfrac{1}{3}(e-e^{1/4})$

5. $g_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b g(t)\,dt = \dfrac{1}{2-0}\int_0^2 \dfrac{9}{1+t^2}\,dt = \dfrac{1}{2}\Big[9\arctan t\Big]_0^2 = \dfrac{9}{2}(\arctan 2 - \arctan 0) = \dfrac{9}{2}\arctan 2$

6. $f_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b f(x)\,dx = \dfrac{1}{1-(-1)}\int_{-1}^{1}\dfrac{x^2}{(x^3+3)^2}\,dx = \dfrac{1}{2}\int_2^4 \dfrac{1}{u^2}\left(\dfrac{1}{3}\,du\right)$ $\left[\begin{array}{l} u = x^3+3, \\ du = 3x^2\,dx \end{array}\right]$

$= \dfrac{1}{6}\left[-\dfrac{1}{u}\right]_2^4 = \dfrac{1}{6}\left(-\dfrac{1}{4}+\dfrac{1}{2}\right) = \dfrac{1}{24}$

7. $h_{\text{avg}} = \frac{1}{b-a}\int_a^b h(x)\,dx = \frac{1}{\pi-0}\int_0^\pi \cos^4 x\,\sin x\,dx = \frac{1}{\pi}\int_1^{-1} u^4(-du)$ $[u = \cos x, du = -\sin x\,dx]$

$= \frac{1}{\pi}\int_{-1}^{1} u^4\,du = \frac{1}{\pi}\cdot 2\int_0^1 u^4\,du$ [by Theorem 5.5.7(a)] $= \frac{2}{\pi}\left[\frac{1}{5}u^5\right]_0^1 = \frac{2}{5\pi}$

8. $h_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b h(u)\,du = \dfrac{1}{5-1}\int_1^5 \dfrac{\ln u}{u}\,du = \dfrac{1}{4}\int_0^{\ln 5} y\,dy$ $\left[\begin{array}{l} y = \ln u, \\ dy = 1/u\,du \end{array}\right]$

$= \frac{1}{4}\left[\frac{1}{2}y^2\right]_0^{\ln 5} = \frac{1}{8}(\ln 5)^2$

9. (a) $f_{\text{avg}} = \dfrac{1}{3-1}\displaystyle\int_1^3 \dfrac{1}{t^2}\,dt = \dfrac{1}{2}\left[-\dfrac{1}{t}\right]_1^3$

$= \dfrac{1}{2}\left[-\dfrac{1}{3}-(-1)\right] = \dfrac{1}{3}$

(b) $f(c) = f_{\text{avg}} \;\Leftrightarrow\; \dfrac{1}{c^2} = \dfrac{1}{3} \;\Leftrightarrow\; c^2 = 3 \;\Rightarrow$

$c = \sqrt{3}$ since $-\sqrt{3}$ is not in $[1, 3]$.

(c)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. (a) $g_{\text{avg}} = \dfrac{1}{2-0}\displaystyle\int_0^2 (x+1)^3\,dx = \dfrac{1}{2}\left[\dfrac{1}{4}(x+1)^4\right]_0^2$

$= \dfrac{1}{8}(3^4 - 1^4) = \dfrac{1}{8}\cdot 80 = 10$

(b) $g(c) = g_{\text{avg}} \;\Leftrightarrow\; (c+1)^3 = 10 \;\Leftrightarrow\; c+1 = \sqrt[3]{10}$

$\Leftrightarrow\; c = -1 + \sqrt[3]{10} \approx 1.154$

(c)

11. (a) $f_{\text{avg}} = \dfrac{1}{\pi - 0}\displaystyle\int_0^{\pi} (2\sin x - \sin 2x)\,dx$

$= \frac{1}{\pi}\left[-2\cos x + \frac{1}{2}\cos 2x\right]_0^{\pi}$

$= \frac{1}{\pi}\left[\left(2 + \frac{1}{2}\right) - \left(-2 + \frac{1}{2}\right)\right] = \frac{4}{\pi}$

(b) $f(c) = f_{\text{avg}} \;\Leftrightarrow\; 2\sin c - \sin 2c = \frac{4}{\pi} \;\Leftrightarrow$

$c = c_1 \approx 1.238$ or $c = c_2 \approx 2.808$

(c)

12. (a) $f_{\text{avg}} = \frac{1}{2-0}\int_0^2 2xe^{-x^2}\,dx$

$= \frac{1}{2}\left[-e^{-x^2}\right]_0^2 = \frac{1}{2}(-e^{-4} + 1)$

(b) $f(c) = f_{\text{avg}} \;\Leftrightarrow\; 2ce^{-c^2} = \frac{1}{2}(1 - e^{-4}) \;\Leftrightarrow$

$c = c_1 \approx 0.263$ or $c = c_2 \approx 1.287$

(c)

13. f is continuous on $[1, 3]$, so by the Mean Value Theorem for Integrals there exists a number c in $[1, 3]$ such that $\int_1^3 f(x)\,dx = f(c)(3-1) \;\Rightarrow\; 8 = 2f(c)$; that is, there is a number c such that $f(c) = \frac{8}{2} = 4$.

14. The requirement is that $\dfrac{1}{b-0}\displaystyle\int_0^b f(x)\,dx = 3$. The LHS of this equation is equal to

$\dfrac{1}{b}\displaystyle\int_0^b \left(2 + 6x - 3x^2\right)dx = \dfrac{1}{b}\left[2x + 3x^2 - x^3\right]_0^b = 2 + 3b - b^2$, so we solve the equation $2 + 3b - b^2 = 3 \;\Leftrightarrow$

$b^2 - 3b + 1 = 0 \;\Leftrightarrow\; b = \dfrac{3 \pm \sqrt{(-3)^2 - 4\cdot 1\cdot 1}}{2\cdot 1} = \dfrac{3 \pm \sqrt{5}}{2}$. Both roots are valid since they are positive.

15. Use geometric interpretations to find the values of the integrals.

$\int_0^8 f(x)\,dx = \int_0^1 f(x)\,dx + \int_1^2 f(x)\,dx + \int_2^3 f(x)\,dx + \int_3^4 f(x)\,dx + \int_4^6 f(x)\,dx + \int_6^7 f(x)\,dx + \int_7^8 f(x)\,dx$

$= -\frac{1}{2} + \frac{1}{2} + \frac{1}{2} + 1 + 4 + \frac{3}{2} + 2 = 9$

Thus, the average value of f on $[0, 8] = f_{\text{avg}} = \frac{1}{8-0}\int_0^8 f(x)\,dx = \frac{1}{8}(9) = \frac{9}{8}$.

16. (a) $v_{\text{avg}} = \frac{1}{12-0}\int_0^{12} v(t)\,dt = \frac{1}{12}I$. Use the Midpoint Rule with $n = 3$ and $\Delta t = \frac{12-0}{3} = 4$ to estimate I.

$I \approx M_3 = 4[v(2) + v(6) + v(10)] = 4[21 + 50 + 66] = 4(137) = 548$. Thus, $v_{\text{avg}} \approx \frac{1}{12}(548) = 45\frac{2}{3}$ km/h.

(b) Estimating from the graph, $v(t) = 45\frac{2}{3}$ when $t \approx 5.2$ s.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. Let $t = 0$ and $t = 12$ correspond to 9 AM and 9 PM, respectively.

$$T_{\text{avg}} = \tfrac{1}{12-0}\int_0^{12}\left[50 + 14\sin\tfrac{1}{12}\pi t\right]dt = \tfrac{1}{12}\left[50t - 14\cdot\tfrac{12}{\pi}\cos\tfrac{1}{12}\pi t\right]_0^{12}$$
$$= \tfrac{1}{12}\left[50\cdot 12 + 14\cdot\tfrac{12}{\pi} + 14\cdot\tfrac{12}{\pi}\right] = 50 + \tfrac{28}{\pi} \approx 59\,^\circ\text{F}$$

18. East Coast: $T_{E_\text{avg}} = \frac{1}{24-0}\int_0^{24} T_E(t)\,dt = \frac{1}{24}E$; West Coast: $T_{W_\text{avg}} = \frac{1}{24-0}\int_0^{24} T_W(t)\,dt = \frac{1}{24}W$

Use the Midpoint Rule with $n = 12$ and $\Delta t = \frac{24-0}{12} = 2$ to estimate E and W.

$$E \approx M_{12} = 2[T_E(1) + T_E(3) + \cdots + T_E(23)] = 2(60+60+65+65+75+70+65+70+55+65+60+55)$$
$$= 2(765) = 1530$$

$$W \approx M_{12} = 2[T_W(1) + T_W(3) + \cdots + T_W(23)] = 2(65+55+50+55+60+65+80+65+60+55+50+50)$$
$$= 2(710) = 1420$$

Thus, $T_{E_\text{avg}} = \frac{1}{24}(1530) = 63\frac{3}{4}\,^\circ\text{F}$, and $T_{W_\text{avg}} = \frac{1}{24}(1420) = 59\frac{1}{6}\,^\circ\text{F}$.

While the West Coast city had the highest temperature that day, 80 °F, the East Coast city was "warmer" overall, because the East Coast city had the higher average temperature.

19. $\rho_{\text{avg}} = \dfrac{1}{8}\displaystyle\int_0^8 \frac{12}{\sqrt{x+1}}\,dx = \frac{3}{2}\int_0^8 (x+1)^{-1/2}\,dx = \left[3\sqrt{x+1}\right]_0^8 = 9 - 3 = 6\ \text{kg/m}$

20. $v_{\text{avg}} = \dfrac{1}{R-0}\displaystyle\int_0^R v(r)\,dr = \frac{1}{R}\int_0^R \frac{P}{4\eta l}(R^2 - r^2)\,dr = \frac{P}{4\eta l R}\left[R^2 r - \tfrac{1}{3}r^3\right]_0^R = \frac{P}{4\eta l R}\left(\tfrac{2}{3}\right)R^3 = \frac{PR^2}{6\eta l}.$

Since $v(r)$ is decreasing on $(0, R]$, $v_{\text{max}} = v(0) = \dfrac{PR^2}{4\eta l}$. Thus, $v_{\text{avg}} = \frac{2}{3}v_{\text{max}}$.

21. $P_{\text{avg}} = \frac{1}{50-0}\int_0^{50} P(t)\,dt = \frac{1}{50}\int_0^{50} 2560e^{bt}\,dt \qquad$ [with $b = 0.017185$]

$$= \frac{2560}{50}\left[\frac{1}{b}e^{bt}\right]_0^{50} = \frac{2560}{50b}(e^{50b} - 1) \approx 4056 \text{ million, or about 4 billion people}$$

22. (a) Similar to Example 3.8.3, we have $T_s = 20^\circ\text{C}$ and hence $\dfrac{dT}{dt} = c(T - 20)$. Let $y = T - 20$, so that

$y(0) = T(0) - 20 = 95 - 20 = 75$. Now y satisfies (3.8.2), so $y = 75e^{ct}$. We are given that $T(30) = 61$, so

$y(30) = 61 - 20 = 41$ and $41 = 75e^{c(30)} \;\Rightarrow\; \frac{41}{75} = e^{30c} \;\Rightarrow\; 30c = \ln\frac{41}{75} \;\Rightarrow\; c = \frac{1}{30}\ln\frac{41}{75} \approx -0.020131.$

Thus, $T(t) = 20 + 75e^{-kt}$, where $k = -c \approx 0.02$.

(b) $T_{\text{avg}} = \frac{1}{30-0}\int_0^{30} T(t)\,dt = \frac{1}{30}\int_0^{30}(20 + 75e^{-kt})\,dt = \frac{1}{30}\left[20t - \frac{75}{k}e^{-kt}\right]_0^{30} = \frac{1}{30}\left[\left(600 - \frac{75}{k}e^{-30k}\right) - \left(0 - \frac{75}{k}\right)\right]$

$$= \tfrac{1}{30}\left(600 - \tfrac{75}{k}\cdot\tfrac{41}{75} + \tfrac{75}{k}\right) = \tfrac{1}{30}\left(600 + \tfrac{34}{k}\right) = 20 + \tfrac{34}{30k} \approx 76.3^\circ\text{C}$$

23. $V_{\text{avg}} = \frac{1}{5}\int_0^5 V(t)\,dt = \frac{1}{5}\int_0^5 \frac{5}{4\pi}\left[1 - \cos\left(\frac{2}{5}\pi t\right)\right]dt = \frac{1}{4\pi}\int_0^5\left[1 - \cos\left(\frac{2}{5}\pi t\right)\right]dt$

$$= \tfrac{1}{4\pi}\left[t - \tfrac{5}{2\pi}\sin\left(\tfrac{2}{5}\pi t\right)\right]_0^5 = \tfrac{1}{4\pi}\left[(5-0) - 0\right] = \tfrac{5}{4\pi} \approx 0.4\ \text{L}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $s = \frac{1}{2}gt^2 \quad \Rightarrow \quad t = \sqrt{2s/g}$ [since $t \geq 0$]. Now $v = ds/dt = gt = g\sqrt{2s/g} = \sqrt{2gs} \quad \Rightarrow \quad v^2 = 2gs \quad \Rightarrow \quad s = \dfrac{v^2}{2g}$.

We see that v can be regarded as a function of t or of s: $v = F(t) = gt$ and $v = G(s) = \sqrt{2gs}$. Note that $v_T = F(T) = gT$.

Displacement can be viewed as a function of t: $s = s(t) = \frac{1}{2}gt^2$; also $s(t) = \dfrac{v^2}{2g} = \dfrac{[F(t)]^2}{2g}$. When $t = T$, these two formulas for $s(t)$ imply that

$$\sqrt{2gs(T)} = F(T) = v_T = gT = 2\left(\tfrac{1}{2}gT^2\right)/T = 2s(T)/T \qquad (\star)$$

The average of the velocities with respect to time t during the interval $[0, T]$ is

$$v_{t\text{-avg}} = f_{\text{avg}} = \frac{1}{T-0}\int_0^T F(t)\,dt = \frac{1}{T}\left[s(T) - s(0)\right] \quad \text{[by FTC]} \quad = \frac{s(T)}{T} \quad \text{[since } s(0) = 0\text{]} \quad = \frac{1}{2}v_T \quad \text{[by } (\star)\text{]}$$

But the average of the velocities with respect to displacement s during the corresponding displacement interval $[s(0), s(T)] = [0, s(T)]$ is

$$v_{s\text{-avg}} = g_{\text{avg}} = \frac{1}{s(T)-0}\int_0^{s(T)} G(s)\,ds = \frac{1}{s(T)}\int_0^{s(T)} \sqrt{2gs}\,ds = \frac{\sqrt{2g}}{s(T)}\int_0^{s(T)} s^{1/2}\,ds$$

$$= \frac{\sqrt{2g}}{s(T)}\cdot\frac{2}{3}\left[s^{3/2}\right]_0^{s(T)} = \frac{2}{3}\cdot\frac{\sqrt{2g}}{s(T)}\cdot\left[s(T)\right]^{3/2} = \frac{2}{3}\sqrt{2gs(T)} = \frac{2}{3}v_T \quad \text{[by } (\star)\text{]}$$

25. $f_{\text{avg}} = \frac{1}{b-a}\int_a^b f(x)\,dx$

$> \frac{1}{b-a}$ (area of trapezoid $ABDF$)

$= \frac{1}{b-a}$ (area of rectangle $ACEF$)

$= \frac{1}{b-a}\left[f\left(\frac{a+b}{2}\right)\cdot(b-a)\right]$

$= f\left(\frac{a+b}{2}\right)$

26. $f_{\text{avg}}[a, b] = \dfrac{1}{b-a}\displaystyle\int_a^b f(x)\,dx = \dfrac{1}{b-a}\int_a^c f(x)\,dx + \dfrac{1}{b-a}\int_c^b f(x)\,dx$

$$= \frac{c-a}{b-a}\left[\frac{1}{c-a}\int_a^c f(x)\,dx\right] + \frac{b-c}{b-a}\left[\frac{1}{b-c}\int_c^b f(x)\,dx\right] = \frac{c-a}{b-a}f_{\text{avg}}[a, c] + \frac{b-c}{b-a}f_{\text{avg}}[c, b]$$

27. Since f is continuous, we can apply l'Hospital's Rule.

$$\lim_{t\to a^+} f_{\text{avg}}[a, t] = \lim_{t\to a^+}\frac{1}{t-a}\int_a^t f(x)\,dx = \lim_{t\to a^+}\frac{\displaystyle\int_a^t f(x)\,dx}{t-a} \quad \left[\text{form } \tfrac{0}{0}\right] \overset{\text{H}}{=} \lim_{t\to a^+}\frac{f(t)}{1} = f(a)$$

28. Let $F(x) = \int_a^x f(t)\,dt$ for x in $[a, b]$. Then F is continuous on $[a, b]$ and differentiable on (a, b), so by the Mean Value Theorem there is a number c in (a, b) such that $F(b) - F(a) = F'(c)(b-a)$. But $F'(x) = f(x)$ by the Fundamental Theorem of Calculus. Therefore, $\int_a^b f(t)\,dt - 0 = f(c)(b-a)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

APPLIED PROJECT Calculus and Baseball

1. (a) $F = ma = m\dfrac{dv}{dt}$, so by the Substitution Rule we have

$$\int_{t_0}^{t_1} F(t)\,dt = \int_{t_0}^{t_1} m\left(\frac{dv}{dt}\right)dt = m\int_{v_0}^{v_1} dv = \big[mv\big]_{v_0}^{v_1} = mv_1 - mv_0 = p(t_1) - p(t_0)$$

(b) (i) We have $v_1 = 110$ mi/h $= \frac{110(5280)}{3600}$ ft/s $= 161.\overline{3}$ ft/s, $v_0 = -90$ mi/h $= -132$ ft/s, and the mass of the baseball is $m = \dfrac{w}{g} = \frac{5/16}{32} = \frac{5}{512}$. So the change in momentum is

$p(t_1) - p(t_0) = mv_1 - mv_0 = \frac{5}{512}\,[161.\overline{3} - (-132)] \approx 2.86$ slug-ft/s.

(ii) From part (a) and part (b)(i), we have $\int_0^{0.001} F(t)\,dt = p(0.001) - p(0) \approx 2.86$, so the average force over the interval $[0, 0.001]$ is $\frac{1}{0.001}\int_0^{0.001} F(t)\,dt \approx \frac{1}{0.001}(2.86) = 2860$ lb.

2. (a) $W = \displaystyle\int_{s_0}^{s_1} F(s)\,ds$, where $F(s) = m\dfrac{dv}{dt} = m\dfrac{dv}{ds}\dfrac{ds}{dt} = mv\dfrac{dv}{ds}$ and so, by the Substitution Rule,

$$W = \int_{s_0}^{s_1} F(s)\,ds = \int_{s_0}^{s_1} mv\frac{dv}{ds}\,ds = \int_{v(s_0)}^{v(s_1)} mv\,dv = \big[\tfrac{1}{2}mv^2\big]_{v_0}^{v_1} = \tfrac{1}{2}mv_1^2 - \tfrac{1}{2}mv_0^2$$

(b) From part (b)(i), 90 mi/h $= 132$ ft/s. Assume $v_0 = v(s_0) = 0$ and $v_1 = v(s_1) = 132$ ft/s [note that s_1 is the point of release of the baseball]. $m = \frac{5}{512}$, so the work done is $W = \frac{1}{2}mv_1^2 - \frac{1}{2}mv_0^2 = \frac{1}{2}\cdot\frac{5}{512}\cdot(132)^2 \approx 85$ ft-lb.

3. (a) Here we have a differential equation of the form $dv/dt = kv$, so by Theorem 3.8.2, the solution is $v(t) = v(0)e^{kt}$. In this case $k = -\frac{1}{10}$ and $v(0) = 100$ ft/s, so $v(t) = 100e^{-t/10}$. We are interested in the time t that the ball takes to travel 280 ft, so we find the distance function

$$s(t) = \int_0^t v(x)\,dx = \int_0^t 100e^{-x/10}\,dx = 100\Big[-10e^{-x/10}\Big]_0^t = -1000(e^{-t/10} - 1) = 1000(1 - e^{-t/10})$$

Now we set $s(t) = 280$ and solve for t: $280 = 1000(1 - e^{-t/10}) \;\Rightarrow\; 1 - e^{-t/10} = \frac{7}{25} \;\Rightarrow$ $-\frac{1}{10}t = \ln\big(1 - \frac{7}{25}\big) \;\Rightarrow\; t \approx 3.285$ seconds.

(b) Let x be the distance of the shortstop from home plate. We calculate the time for the ball to reach home plate as a function of x, then differentiate with respect to x to find the value of x which corresponds to the minimum time. The total time that it takes the ball to reach home is the sum of the times of the two throws, plus the relay time $\big(\frac{1}{2}\text{ s}\big)$. The distance from the fielder to the shortstop is $280 - x$, so to find the time t_1 taken by the first throw, we solve the equation

$s_1(t_1) = 280 - x \;\Leftrightarrow\; 1 - e^{-t_1/10} = \dfrac{280 - x}{1000} \;\Leftrightarrow\; t_1 = -10\ln\dfrac{720 + x}{1000}$. We find the time t_2 taken by the second throw if the shortstop throws with velocity w, since we see that this velocity varies in the rest of the problem. We use

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$v = we^{-t/10}$ and isolate t_2 in the equation $s(t_2) = 10w(1 - e^{-t_2/10}) = x \quad \Leftrightarrow \quad e^{-t_2/10} = 1 - \dfrac{x}{10w} \quad \Leftrightarrow$

$t_2 = -10 \ln \dfrac{10w - x}{10w}$, so the total time is $t_w(x) = \dfrac{1}{2} - 10\left[\ln \dfrac{720 + x}{1000} + \ln \dfrac{10w - x}{10w}\right]$.

To find the minimum, we differentiate: $\dfrac{dt_w}{dx} = -10\left[\dfrac{1}{720 + x} - \dfrac{1}{10w - x}\right]$, which changes from negative to positive when $720 + x = 10w - x \quad \Leftrightarrow \quad x = 5w - 360$. By the First Derivative Test, t_w has a minimum at this distance from the shortstop to home plate. So if the shortstop throws at $w = 105$ ft/s from a point $x = 5(105) - 360 = 165$ ft from home plate, the minimum time is $t_{105}(165) = \frac{1}{2} - 10\left(\ln \frac{720 + 165}{1000} + \ln \frac{1050 - 165}{1050}\right) \approx 3.431$ seconds. This is longer than the time taken in part (a), so in this case the manager should encourage a direct throw. If $w = 115$ ft/s, then $x = 215$ ft from home, and the minimum time is $t_{115}(215) = \frac{1}{2} - 10\left(\ln \frac{720 + 215}{1000} + \ln \frac{1150 - 215}{1150}\right) \approx 3.242$ seconds. This is less than the time taken in part (a), so in this case, the manager should encourage a relayed throw.

(c) In general, the minimum time is $t_w(5w - 360) = \dfrac{1}{2} - 10\left[\ln \dfrac{360 + 5w}{1000} + \ln \dfrac{360 + 5w}{10w}\right] = \dfrac{1}{2} - 10 \ln \dfrac{(w + 72)^2}{400w}$.

We want to find out when this is about 3.285 seconds, the same time as the direct throw. From the graph, we estimate that this is the case for $w \approx 112.8$ ft/s. So if the shortstop can throw the ball with this velocity, then a relayed throw takes the same time as a direct throw.

APPLIED PROJECT Where to Sit at the Movies

1. $|VP| = 9 + x\cos\alpha$, $|PT| = 35 - (4 + x\sin\alpha) = 31 - x\sin\alpha$, and $|PB| = (4 + x\sin\alpha) - 10 = x\sin\alpha - 6$. So using the Pythagorean Theorem, we have $|VT| = \sqrt{|VP|^2 + |PT|^2} = \sqrt{(9 + x\cos\alpha)^2 + (31 - x\sin\alpha)^2} = a$, and $|VB| = \sqrt{|VP|^2 + |PB|^2} = \sqrt{(9 + x\cos\alpha)^2 + (x\sin\alpha - 6)^2} = b$.

Using the Law of Cosines on $\triangle VBT$, we get $25^2 = a^2 + b^2 - 2ab\cos\theta \quad \Leftrightarrow \quad \cos\theta = \dfrac{a^2 + b^2 - 625}{2ab} \quad \Leftrightarrow$

$\theta = \arccos\left(\dfrac{a^2 + b^2 - 625}{2ab}\right)$, as required.

2. From the graph of θ, it appears that the value of x which maximizes θ is $x \approx 8.25$ ft. Assuming that the first row is at $x = 0$, the row closest to this value of x is the fourth row, at $x = 9$ ft, and from the graph, the viewing angle in this row seems to be about 0.85 radians, or about $49°$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. With a CAS, we type in the definition of θ, substitute in the proper values of a and b in terms of x and $\alpha = 20^\circ = \frac{\pi}{9}$ radians, and then use the differentiation command to find the derivative. We use a numerical rootfinder and find that the root of the equation $d\theta/dx = 0$ is $x \approx 8.253062$, as approximated in Problem 2.

4. From the graph in Problem 2, it seems that the average value of the function on the interval $[0, 60]$ is about 0.6. We can use a CAS to approximate $\frac{1}{60}\int_0^{60} \theta(x)\,dx \approx 0.625 \approx 36^\circ$. (The calculation is much faster if we reduce the number of digits of accuracy required.) The minimum value is $\theta(60) \approx 0.38$ and, from Problem 2, the maximum value is about 0.85.

6 Review

TRUE-FALSE QUIZ

1. False. For example, let $f(x) = x$, $g(x) = 2x$, $a = 1$, and $b = 2$. The area between the curves for $a \le x \le b$ is $A = \int_a^b [g(x) - f(x)]\,dx$. The given integral represents area when $f(x) \ge g(x)$ for $a \le x \le b$.

2. False. In a solid of revolution, cross-sections perpendicular to the axis of rotation are circular. A cube has no circular cross-sections.

3. False. Cross-sections perpendicular to the x-axis are washers, and we find cross-sectional area by subtracting the area of the inner circle from the area of the outer circle. Thus, $A(x) = \pi\left(\sqrt{x}\right)^2 - \pi(x)^2 = \pi\left[\left(\sqrt{x}\right)^2 - x^2\right]$, and the volume of the resulting solid is $V = \displaystyle\int_0^1 A(x)\,dx = \int_0^1 \pi\left[\left(\sqrt{x}\right)^2 - x^2\right]\,dx$.

4. True. See "The Method of Cylindrical Shells" in Section 6.3.

5. True. See "Volumes of Solids of Revolution" in Section 6.2.

6. True. See "Volumes of Solids of Revolution" in Section 6.2.

7. False. Cross-sections perpendicular to the y-axis are washers.

8. False. Using the method of cylindrical shells, a typical shell of the solid obtained by revolving $\mathcal{R}$ about the y-axis has radius x, and the volume of the solid is $V = \int_a^b 2\pi x\, f(x)\,dx$. Again using the method of cylindrical shells, a typical shell of the solid obtained by revolving about the line $x = -2$ has radius $x - (-2) = x + 2$, and the volume of the solid is $V = \int_a^b 2\pi(x+2)\, f(x)\,dx$.

9. True. A cross-section of S perpendicular to the x-axis is a square with side length $f(x)$, so each cross-section has area $A(x) = [f(x)]^2$ and volume $V = \int_a^b A(x)\,dx = \int_a^b [f(x)]^2\,dx$.

10. False. The work done to pull up the top half of the cable will be more than half of the work required to pull up the entire cable. When the top half of the cable is being pulled up, the bottom half is still attached, and extra work is required to pull that bottom half up as the top half is pulled up. There is no such extra work required as the remaining bottom half is pulled up.

11. True. By defnition of the average value of f on the interval $[a, b]$, the average value of f on $[2, 5]$ is $\frac{1}{5-2}\int_2^5 f(x)\,dx = \frac{1}{3}(12) = 4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

EXERCISES

1. The curves intersect when $x^2 = 8x - x^2 \quad \Leftrightarrow \quad 2x^2 - 8x = 0 \quad \Leftrightarrow$ $2x(x-4) = 0 \quad \Leftrightarrow \quad x = 0$ or 4.

$$A = \int_0^4 \left[(8x - x^2) - x^2\right] dx = \int_0^4 (8x - 2x^2)\, dx$$

$$= \left[4x^2 - \tfrac{2}{3}x^3\right]_0^4 = \left[\left(64 - \tfrac{128}{3}\right) - 0\right] = \tfrac{64}{3}$$

2. The line $y = x - 2$ intersects the curve $y = \sqrt{x}$ at $(4, 2)$ and it intersects the curve $y = -\sqrt[3]{x}$ at $(1, -1)$.

$$A = \int_0^1 [\sqrt{x} - (-\sqrt[3]{x})]\, dx + \int_1^4 [\sqrt{x} - (x-2)]\, dx$$

$$= \left[\tfrac{2}{3}x^{3/2} + \tfrac{3}{4}x^{4/3}\right]_0^1 + \left[\tfrac{2}{3}x^{3/2} - \tfrac{1}{2}x^2 + 2x\right]_1^4$$

$$= \left(\tfrac{2}{3} + \tfrac{3}{4}\right) - 0 + \left(\tfrac{16}{3} - 8 + 8\right) - \left(\tfrac{2}{3} - \tfrac{1}{2} + 2\right)$$

$$= \tfrac{16}{3} + \tfrac{3}{4} - \tfrac{3}{2} = \tfrac{55}{12}$$

Or, integrating with respect to y: $A = \int_{-1}^0 [(y+2) - (-y^3)]\, dy + \int_0^2 [(y+2) - y^2]\, dy$

3. If $x \geq 0$, then $|\,x\,| = x$, and the graphs intersect when $x = 1 - 2x^2 \quad \Leftrightarrow \quad 2x^2 + x - 1 = 0 \quad \Leftrightarrow \quad (2x-1)(x+1) = 0 \quad \Leftrightarrow$ $x = \frac{1}{2}$ or -1, but $-1 < 0$. By symmetry, we can double the area from $x = 0$ to $x = \frac{1}{2}$.

$$A = 2\int_0^{1/2} \left[(1 - 2x^2) - x\right] dx = 2\int_0^{1/2} (-2x^2 - x + 1)\, dx$$

$$= 2\left[-\tfrac{2}{3}x^3 - \tfrac{1}{2}x^2 + x\right]_0^{1/2} = 2\left[\left(-\tfrac{1}{12} - \tfrac{1}{8} + \tfrac{1}{2}\right) - 0\right]$$

$$= 2\left(\tfrac{7}{24}\right) = \tfrac{7}{12}$$

4. $y^2 + 3y = -y \quad \Leftrightarrow \quad y^2 + 4y = 0 \quad \Leftrightarrow \quad y(y+4) = 0 \quad \Leftrightarrow$ $y = 0$ or -4.

$$A = \int_{-4}^0 \left[-y - (y^2 + 3y)\right] dy = \int_{-4}^0 (-y^2 - 4y)\, dy$$

$$= \left[-\tfrac{1}{3}y^3 - 2y^2\right]_{-4}^0 = 0 - \left(\tfrac{64}{3} - 32\right) = \tfrac{32}{3}$$

5. $$A = \int_0^2 \left[\sin\left(\frac{\pi x}{2}\right) - (x^2 - 2x)\right] dx$$

$$= \left[-\frac{2}{\pi}\cos\left(\frac{\pi x}{2}\right) - \frac{1}{3}x^3 + x^2\right]_0^2$$

$$= \left(\tfrac{2}{\pi} - \tfrac{8}{3} + 4\right) - \left(-\tfrac{2}{\pi} - 0 + 0\right) = \tfrac{4}{3} + \tfrac{4}{\pi}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $A = \int_0^1 \left(\sqrt{x} - x^2\right) dx + \int_1^2 \left(x^2 - \sqrt{x}\right) dx$

$= \left[\frac{2}{3}x^{3/2} - \frac{1}{3}x^3\right]_0^1 + \left[\frac{1}{3}x^3 - \frac{2}{3}x^{3/2}\right]_1^2$

$= \left[\left(\frac{2}{3} - \frac{1}{3}\right) - 0\right] + \left[\left(\frac{8}{3} - \frac{4}{3}\sqrt{2}\right) - \left(\frac{1}{3} - \frac{2}{3}\right)\right]$

$= \frac{10}{3} - \frac{4}{3}\sqrt{2}$

7. Using washers with inner radius x^2 and outer radius $2x$, we have

$V = \pi \int_0^2 \left[(2x)^2 - (x^2)^2\right] dx = \pi \int_0^2 (4x^2 - x^4)\, dx$

$= \pi\left[\frac{4}{3}x^3 - \frac{1}{5}x^5\right]_0^2 = \pi\left(\frac{32}{3} - \frac{32}{5}\right)$

$= 32\pi \cdot \frac{2}{15} = \frac{64}{15}\pi$

8. $1 + y^2 = y + 3 \quad \Leftrightarrow \quad y^2 - y - 2 = 0 \quad \Leftrightarrow \quad (y-2)(y+1) = 0 \quad \Leftrightarrow$

$y = 2$ or -1.

$V = \pi \int_{-1}^2 \left[(y+3)^2 - (1+y^2)^2\right] dy = \pi \int_{-1}^2 (y^2 + 6y + 9 - 1 - 2y^2 - y^4)\, dy$

$= \pi \int_{-1}^2 (8 + 6y - y^2 - y^4)\, dy = \pi\left[8y + 3y^2 - \frac{1}{3}y^3 - \frac{1}{5}y^5\right]_{-1}^2$

$= \pi\left[\left(16 + 12 - \frac{8}{3} - \frac{32}{5}\right) - \left(-8 + 3 + \frac{1}{3} + \frac{1}{5}\right)\right] = \pi\left(33 - \frac{9}{3} - \frac{33}{5}\right) = \frac{117}{5}\pi$

9. $V = \pi \int_{-3}^3 \left\{\left[(9 - y^2) - (-1)\right]^2 - [0 - (-1)]^2\right\} dy$

$= 2\pi \int_0^3 \left[(10 - y^2)^2 - 1\right] dy = 2\pi \int_0^3 (100 - 20y^2 + y^4 - 1)\, dy$

$= 2\pi \int_0^3 (99 - 20y^2 + y^4)\, dy = 2\pi\left[99y - \frac{20}{3}y^3 + \frac{1}{5}y^5\right]_0^3$

$= 2\pi\left(297 - 180 + \frac{243}{5}\right) = \frac{1656}{5}\pi$

10. $V = \pi \int_{-2}^2 \left\{\left[(9 - x^2) - (-1)\right]^2 - \left[(x^2 + 1) - (-1)\right]^2\right\} dx$

$= \pi \int_{-2}^2 \left[(10 - x^2)^2 - (x^2 + 2)^2\right] dx$

$= 2\pi \int_0^2 (96 - 24x^2)\, dx = 48\pi \int_0^2 (4 - x^2)\, dx$

$= 48\pi\left[4x - \frac{1}{3}x^3\right]_0^2 = 48\pi\left(8 - \frac{8}{3}\right) = 256\pi$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. The graph of $x^2 - y^2 = a^2$ is a hyperbola with right and left branches.

Solving for y gives us $y^2 = x^2 - a^2 \quad \Rightarrow \quad y = \pm\sqrt{x^2 - a^2}$.

We'll use shells and the height of each shell is

$\sqrt{x^2 - a^2} - \left(-\sqrt{x^2 - a^2}\right) = 2\sqrt{x^2 - a^2}$.

The volume is $V = \int_a^{a+h} 2\pi x \cdot 2\sqrt{x^2 - a^2}\, dx$. To evaluate, let $u = x^2 - a^2$, so $du = 2x\, dx$ and $x\, dx = \frac{1}{2}\, du$. When $x = a$, $u = 0$, and when $x = a + h$, $u = (a+h)^2 - a^2 = a^2 + 2ah + h^2 - a^2 = 2ah + h^2$.

Thus, $V = 4\pi \displaystyle\int_0^{2ah+h^2} \sqrt{u}\left(\frac{1}{2}\, du\right) = 2\pi\left[\frac{2}{3}u^{3/2}\right]_0^{2ah+h^2} = \frac{4}{3}\pi\left(2ah + h^2\right)^{3/2}$.

12. A shell has radius x, circumference $2\pi x$, and height $\tan x - x$.

$V = \int_0^{\pi/3} 2\pi x\,(\tan x - x)\, dx$

13. A shell has radius $\frac{\pi}{2} - x$, circumference $2\pi\left(\frac{\pi}{2} - x\right)$, and height $\cos^2 x - \frac{1}{4}$.

$y = \cos^2 x$ intersects $y = \frac{1}{4}$ when $\cos^2 x = \frac{1}{4} \quad \Leftrightarrow$

$\cos x = \pm\frac{1}{2} \quad [\,|x| \le \pi/2] \quad \Leftrightarrow \quad x = \pm\frac{\pi}{3}$.

$$V = \int_{-\pi/3}^{\pi/3} 2\pi\left(\frac{\pi}{2} - x\right)\left(\cos^2 x - \frac{1}{4}\right) dx$$

14. A shell has radius $x - (-1) = x + 1$,

circumference $2\pi(x + 1)$, and height $\ln x$.

$V = \int_1^4 2\pi(x + 1)\, \ln x\, dx$

15. $3x^2 = x^3 \quad \Leftrightarrow \quad x^3 - 3x^2 = 0 \quad \Leftrightarrow$

$x^2(x - 3) = 0 \quad \Leftrightarrow \quad x = 0$ or $x = 3$

$y = 3x^2 \quad \Rightarrow \quad x = \sqrt{\dfrac{y}{3}}\ (x > 0)$

$y = x^3 \quad \Rightarrow \quad x = \sqrt[3]{y}$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(a) With x as the variable of integration, we use the method of cylindrical shells.

$$V = \int_0^3 2\pi(x+1)(3x^2 - x^3)\,dx = \int_0^3 2\pi(2x^3 - x^4 + 3x^2)\,dx = 2\pi\left[\tfrac{1}{2}x^4 - \tfrac{1}{5}x^5 + x^3\right]_0^3$$

$$= 2\pi\left[\left(\frac{81}{2} - \frac{243}{5} + 27\right) - 0\right] = 2\pi \cdot \frac{189}{10} = \frac{189}{5}\pi$$

(b) With y as the variable of integration, we use washers with inner radius $\sqrt{\dfrac{y}{3}} + 1$ and outer radius $\sqrt[3]{y} + 1$.

The area of a cross-section is

$$\pi\left(\sqrt[3]{y} + 1\right)^2 - \pi\left(\sqrt{\frac{y}{3}} + 1\right)^2 = \pi\left[(y^{2/3} + 2y^{1/3} + 1) - \left(\frac{y}{3} + \frac{2}{\sqrt{3}}\,y^{1/2} + 1)\right)\right]$$

$$= \pi\left(y^{2/3} + 2y^{1/3} - \frac{y}{3} - \frac{2}{\sqrt{3}}\,y^{1/2}\right)$$

$$V = \int_0^{27} \pi\left(y^{2/3} + 2y^{1/3} - \frac{y}{3} - \frac{2}{\sqrt{3}}\,y^{1/2}\right) dy = \pi\left[\frac{3}{5}y^{5/3} + \frac{3}{2}y^{4/3} - \frac{1}{6}y^2 - \frac{4}{3\sqrt{3}}\,y^{3/2}\right]_0^{27}$$

$$= \pi\left[\left(\frac{729}{5} + \frac{243}{2} - \frac{243}{2} - 108\right) - 0\right] = \frac{189}{5}\pi$$

16. $\sqrt{x} = x^2 \;\Rightarrow\; x = x^4 \;\Leftrightarrow\; x^4 - x = 0 \;\Leftrightarrow$

$x(x^3 - 1) = 0 \;\Leftrightarrow\; x = 0$ or $x = 1$

$y = \sqrt{x} \;\Rightarrow\; x = y^2\ (x > 0)$

$y = x^2\ (x > 0) \;\Rightarrow\; x = \sqrt{y}$

(a) With x as the variable of integration, we use washers with inner radius $3 - \sqrt{x}$ and outer radius $3 - x^2$. The area of a cross-section is $\pi(3 - x^2)^2 - \pi\left(3 - \sqrt{x}\right)^2 = \pi[(9 - 6x^2 + x^4) - (9 - 6x^{1/2} + x)] = \pi(-6x^2 + x^4 + 6x^{1/2} - x)$.

$$V = \int_0^1 \pi(-6x^2 + x^4 + 6x^{1/2} - x)\,dx = \pi\left[-2x^3 + \tfrac{1}{5}x^5 + 4x^{3/2} - \tfrac{1}{2}x^2\right]_0^1 = \pi\left[\left(-2 + \tfrac{1}{5} + 4 - \tfrac{1}{2}\right) - 0\right] = \tfrac{17}{10}\pi$$

(b) With y as the variable of integration, we use the method of cylindrical shells.

$$V = \int_0^1 2\pi(3 - y)\left(\sqrt{y} - y^2\right) dy = \int_0^1 2\pi(3y^{1/2} - 3y^2 - y^{3/2} + y^3)\,dy$$

$$= 2\pi\left[2y^{3/2} - y^3 - \tfrac{2}{5}y^{5/2} + \tfrac{1}{4}y^4\right]_0^1 = 2\pi\left[\left(2 - 1 - \tfrac{2}{5} + \tfrac{1}{4}\right) - 0\right] = 2\pi \cdot \tfrac{17}{20} = \tfrac{17}{10}\pi$$

17. (a) A cross-section is a washer with inner radius x^2 and outer radius x.

$$V = \int_0^1 \pi\left[(x)^2 - (x^2)^2\right] dx = \int_0^1 \pi(x^2 - x^4)\,dx = \pi\left[\tfrac{1}{3}x^3 - \tfrac{1}{5}x^5\right]_0^1 = \pi\left[\tfrac{1}{3} - \tfrac{1}{5}\right] = \tfrac{2}{15}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) A cross-section is a washer with inner radius y and outer radius $\sqrt{y}$.

$$V = \int_0^1 \pi\left[\left(\sqrt{y}\right)^2 - y^2\right] dy = \int_0^1 \pi(y - y^2)\, dy = \pi\left[\tfrac{1}{2}y^2 - \tfrac{1}{3}y^3\right]_0^1 = \pi\left[\tfrac{1}{2} - \tfrac{1}{3}\right] = \tfrac{\pi}{6}$$

(c) A cross-section is a washer with inner radius $2 - x$ and outer radius $2 - x^2$.

$$V = \int_0^1 \pi\left[(2 - x^2)^2 - (2 - x)^2\right] dx = \int_0^1 \pi(x^4 - 5x^2 + 4x)\, dx = \pi\left[\tfrac{1}{5}x^5 - \tfrac{5}{3}x^3 + 2x^2\right]_0^1 = \pi\left[\tfrac{1}{5} - \tfrac{5}{3} + 2\right] = \tfrac{8}{15}\pi$$

18. (a) $A = \int_0^1 (2x - x^2 - x^3)\, dx = \left[x^2 - \tfrac{1}{3}x^3 - \tfrac{1}{4}x^4\right]_0^1 = 1 - \tfrac{1}{3} - \tfrac{1}{4} = \tfrac{5}{12}$

(b) A cross-section is a washer with inner radius x^3 and outer radius $2x - x^2$, so its area is $\pi(2x - x^2)^2 - \pi(x^3)^2$.

$$\begin{aligned} V &= \int_0^1 A(x)\, dx = \int_0^1 \pi[(2x - x^2)^2 - (x^3)^2]\, dx = \int_0^1 \pi(4x^2 - 4x^3 + x^4 - x^6)\, dx \\ &= \pi\left[\tfrac{4}{3}x^3 - x^4 + \tfrac{1}{5}x^5 - \tfrac{1}{7}x^7\right]_0^1 = \pi\left(\tfrac{4}{3} - 1 + \tfrac{1}{5} - \tfrac{1}{7}\right) = \tfrac{41}{105}\pi \end{aligned}$$

(c) Using the method of cylindrical shells,

$$V = \int_0^1 2\pi x(2x - x^2 - x^3)\, dx = \int_0^1 2\pi(2x^2 - x^3 - x^4)\, dx = 2\pi\left[\tfrac{2}{3}x^3 - \tfrac{1}{4}x^4 - \tfrac{1}{5}x^5\right]_0^1 = 2\pi\left(\tfrac{2}{3} - \tfrac{1}{4} - \tfrac{1}{5}\right) = \tfrac{13}{30}\pi.$$

19. (a) Using the Midpoint Rule on $[0, 1]$ with $f(x) = \tan(x^2)$ and $n = 4$, we estimate

$$A = \int_0^1 \tan(x^2)\, dx \approx \tfrac{1}{4}\left[\tan\left(\left(\tfrac{1}{8}\right)^2\right) + \tan\left(\left(\tfrac{3}{8}\right)^2\right) + \tan\left(\left(\tfrac{5}{8}\right)^2\right) + \tan\left(\left(\tfrac{7}{8}\right)^2\right)\right] \approx \tfrac{1}{4}(1.53) \approx 0.38$$

(b) Using the Midpoint Rule on $[0, 1]$ with $f(x) = \pi \tan^2(x^2)$ (for disks) and $n = 4$, we estimate

$$V = \int_0^1 f(x)\, dx \approx \tfrac{1}{4}\pi\left[\tan^2\left(\left(\tfrac{1}{8}\right)^2\right) + \tan^2\left(\left(\tfrac{3}{8}\right)^2\right) + \tan^2\left(\left(\tfrac{5}{8}\right)^2\right) + \tan^2\left(\left(\tfrac{7}{8}\right)^2\right)\right] \approx \tfrac{\pi}{4}(1.114) \approx 0.87$$

20. (a)

From the graph, we see that the curves intersect at $x = 0$ and at $x = a \approx 0.75$, with $1 - x^2 > x^6 - x + 1$ on $(0, a)$.

(b) The area of $\mathcal{R}$ is $A = \int_0^a \left[(1 - x^2) - (x^6 - x + 1)\right] dx = \left[-\tfrac{1}{3}x^3 - \tfrac{1}{7}x^7 + \tfrac{1}{2}x^2\right]_0^a \approx 0.12$.

(c) Using washers, the volume generated when $\mathcal{R}$ is rotated about the x-axis is

$$\begin{aligned} V &= \pi \int_0^a [(1 - x^2)^2 - (x^6 - x + 1)^2]\, dx = \pi \int_0^a (-x^{12} + 2x^7 - 2x^6 + x^4 - 3x^2 + 2x)\, dx \\ &= \pi\left[-\tfrac{1}{13}x^{13} + \tfrac{1}{4}x^8 - \tfrac{2}{7}x^7 + \tfrac{1}{5}x^5 - x^3 + x^2\right]_0^a \approx 0.54 \end{aligned}$$

(d) Using shells, the volume generated when $\mathcal{R}$ is rotated about the y-axis is

$$V = \int_0^a 2\pi x[(1 - x^2) - (x^6 - x + 1)]\, dx = 2\pi \int_0^a (-x^3 - x^7 + x^2)\, dx = 2\pi\left[-\tfrac{1}{4}x^4 - \tfrac{1}{8}x^8 + \tfrac{1}{3}x^3\right]_0^a \approx 0.31.$$

21. $\int_0^{\pi/2} 2\pi x \cos x\, dx = \int_0^{\pi/2} (2\pi x) \cos x\, dx$

The solid is obtained by rotating the region $\mathcal{R} = \left\{(x, y) \mid 0 \le x \le \tfrac{\pi}{2}, 0 \le y \le \cos x\right\}$ about the y-axis.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $\int_0^{\pi/2} 2\pi \cos^2 x\, dx = \int_0^{\pi/2} \pi\left(\sqrt{2}\cos x\right)^2\, dx$

The solid is obtained by rotating the region $\mathcal{R} = \left\{(x, y) \mid 0 \le x \le \frac{\pi}{2}, 0 \le y \le \sqrt{2}\,\cos x\right\}$ about the x-axis.

23. $\int_0^{\pi} \pi(2 - \sin x)^2\, dx$

The solid is obtained by rotating the region $\mathcal{R} = \{(x, y) \mid 0 \le x \le \pi, 0 \le y \le 2 - \sin x\}$ about the x-axis.

24. $\int_0^4 2\pi(6 - y)(4y - y^2)\, dy$

The solid is obtained by rotating the region $\mathcal{R} = \left\{(x, y) \mid 0 \le x \le 4y - y^2, 0 \le y \le 4\right\}$ about the line $y = 6$.

25. Take the base to be the disk $x^2 + y^2 \le 9$. Then $V = \int_{-3}^{3} A(x)\, dx$, where $A(x_0)$ is the area of the isosceles right triangle whose hypotenuse lies along the line $x = x_0$ in the xy-plane. The length of the hypotenuse is $2\sqrt{9 - x^2}$ and the length of each leg is $\sqrt{2}\sqrt{9 - x^2}$. $A(x) = \frac{1}{2}\left(\sqrt{2}\sqrt{9 - x^2}\right)^2 = 9 - x^2$, so

$$V = 2\int_0^3 A(x)\, dx = 2\int_0^3 (9 - x^2)\, dx = 2\left[9x - \tfrac{1}{3}x^3\right]_0^3 = 2(27 - 9) = 36$$

26. $V = \int_{-1}^{1} A(x)\, dx = 2\int_0^1 A(x)\, dx = 2\int_0^1 \left[(2 - x^2) - x^2\right]^2\, dx = 2\int_0^1 \left[2(1 - x^2)\right]^2\, dx$

$= 8\int_0^1 (1 - 2x^2 + x^4)\, dx = 8\left[x - \frac{2}{3}x^3 + \frac{1}{5}x^5\right]_0^1 = 8\left(1 - \frac{2}{3} + \frac{1}{5}\right) = \frac{64}{15}$

27. Equilateral triangles with sides measuring $\frac{1}{4}x$ meters have height $\frac{1}{4}x\sin 60^\circ = \frac{\sqrt{3}}{8}x$. Therefore,

$A(x) = \frac{1}{2}\cdot\frac{1}{4}x\cdot\frac{\sqrt{3}}{8}x = \frac{\sqrt{3}}{64}x^2$. $\quad V = \int_0^{20} A(x)\, dx = \frac{\sqrt{3}}{64}\int_0^{20} x^2\, dx = \frac{\sqrt{3}}{64}\left[\frac{1}{3}x^3\right]_0^{20} = \frac{8000\sqrt{3}}{64\cdot 3} = \frac{125\sqrt{3}}{3}\ \text{m}^3$.

28. (a) By the symmetry of the problem, we consider only the solid to the right of the origin. The semicircular cross-sections perpendicular to the x-axis have radius $1 - x$, so $A(x) = \frac{1}{2}\pi(1 - x)^2$. Now we can calculate

$V = 2\int_0^1 A(x)\, dx = 2\int_0^1 \frac{1}{2}\pi(1 - x)^2\, dx = \int_0^1 \pi(1 - x)^2\, dx = -\frac{\pi}{3}\left[(1 - x)^3\right]_0^1 = \frac{\pi}{3}$.

(b) Cut the solid with a plane perpendicular to the x-axis and passing through the y-axis. Fold the half of the solid in the region $x \le 0$ under the xy-plane so that the point $(-1, 0)$ comes around and touches the point $(1, 0)$. The resulting solid is a right circular cone of radius 1 with vertex at $(x, y, z) = (1, 0, 0)$ and with its base in the yz-plane, centered at the origin. The volume of this cone is $\frac{1}{3}\pi r^2 h = \frac{1}{3}\pi\cdot 1^2\cdot 1 = \frac{\pi}{3}$.

29. $f(x) = kx \;\Rightarrow\; 30\ \text{N} = k(15 - 12)\ \text{cm} \;\Rightarrow\; k = 10\ \text{N/cm} = 1000\ \text{N/m}$. $\;20\ \text{cm} - 12\ \text{cm} = 0.08\ \text{m} \;\Rightarrow$

$W = \int_0^{0.08} kx\, dx = 1000\int_0^{0.08} x\, dx = 500\left[x^2\right]_0^{0.08} = 500(0.08)^2 = 3.2\ \text{N·m} = 3.2\ \text{J}$.

30. The work needed to raise the elevator alone is $1600\ \text{lb} \times 30\ \text{ft} = 48{,}000$ ft-lb. The work needed to raise the bottom 170 ft of cable is $170\ \text{ft} \times 10\ \text{lb/ft} \times 30\ \text{ft} = 51{,}000$ ft-lb. The work needed to raise the top 30 ft of cable is $\int_0^{30} 10x\, dx = \left[5x^2\right]_0^{30} = 5\cdot 900 = 4500$ ft-lb. Adding these, we see that the total work needed is $48{,}000 + 51{,}000 + 4{,}500 = 103{,}500$ ft-lb.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. (a) The parabola has equation $y = ax^2$ with vertex at the origin and passing through $(4, 4)$. $4 = a \cdot 4^2 \;\Rightarrow\; a = \frac{1}{4} \;\Rightarrow\; y = \frac{1}{4}x^2 \;\Rightarrow\; x^2 = 4y \;\Rightarrow$ $x = 2\sqrt{y}$. Each circular disk has radius $2\sqrt{y}$ and is moved $4 - y$ ft.

$$W = \int_0^4 \pi \left(2\sqrt{y}\right)^2 62.5(4-y)\,dy = 250\pi \int_0^4 y(4-y)\,dy$$

$$= 250\pi\left[2y^2 - \tfrac{1}{3}y^3\right]_0^4 = 250\pi\left(32 - \tfrac{64}{3}\right) = \tfrac{8000\pi}{3} \approx 8378 \text{ ft-lb}$$

(b) In part (a) we knew the final water level (0) but not the amount of work done. Here we use the same equation, except with the work fixed, and the lower limit of integration (that is, the final water level — call it h) unknown: $W = 4000 \;\Leftrightarrow$ $250\pi\left[2y^2 - \frac{1}{3}y^3\right]_h^4 = 4000 \;\Leftrightarrow\; \frac{16}{\pi} = \left[\left(32 - \frac{64}{3}\right) - \left(2h^2 - \frac{1}{3}h^3\right)\right] \;\Leftrightarrow$ $h^3 - 6h^2 + 32 - \frac{48}{\pi} = 0$. We graph the function $f(h) = h^3 - 6h^2 + 32 - \frac{48}{\pi}$ on the interval $[0, 4]$ to see where it is 0. From the graph, $f(h) = 0$ for $h \approx 2.1$. So the depth of water remaining is about 2.1 ft.

32. A horizontal slice of cooking oil Δx m thick has a volume of $\pi r^2 h = \pi \cdot 2^2 \cdot \Delta x \text{ m}^3$, a mass of $920(4\pi\,\Delta x)$ kg, weighs about $(9.8)(3680\pi\,\Delta x) = 36{,}064\pi\Delta x$ N, and thus requires about $36{,}064\pi x_i^* \Delta x$ J of work for its removal (where $3 \le x_i^* \le 6$). The total work needed to empty the tank is

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} 36{,}064\pi x_i^*\,\Delta x = \int_3^6 36{,}064\pi x\,dx = 36{,}064\pi\left[\tfrac{1}{2}x^2\right]_3^6 = 18{,}032\pi(36-9) = 486{,}864\pi \approx 1.53 \times 10^6 \text{ J}.$$

33. $f_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b f(t)\,dt = \dfrac{1}{\pi/4 - 0}\int_0^{\pi/4} \sec^2 t\,dt = \dfrac{4}{\pi}\left[\tan t\right]_0^{\pi/4} = \dfrac{4}{\pi}(1-0) = \dfrac{4}{\pi}$

34. (a) $f_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b f(x)\,dx = \dfrac{1}{4-1}\int_1^4 \dfrac{1}{\sqrt{x}}\,dx$

$$= \frac{1}{3}\int_1^4 x^{-1/2}\,dx = \frac{1}{3}\left[2\sqrt{x}\right]_1^4$$

$$= \tfrac{2}{3}(2-1) = \tfrac{2}{3}$$

(b) $f(c) = f_{\text{avg}} \;\Leftrightarrow\; \dfrac{1}{\sqrt{c}} = \dfrac{2}{3} \;\Leftrightarrow\; \sqrt{c} = \dfrac{3}{2} \;\Leftrightarrow\; c = \dfrac{9}{4}$

(c)

35. (a) The regions $\mathcal{R}_1$ and $\mathcal{R}_2$ are shown in the figure.

The area of $\mathcal{R}_1$ is $A_1 = \int_0^b x^2\,dx = \left[\frac{1}{3}x^3\right]_0^b = \frac{1}{3}b^3$, and the area of $\mathcal{R}_2$ is $A_2 = \int_0^{b^2} \sqrt{y}\,dy = \left[\frac{2}{3}y^{3/2}\right]_0^{b^2} = \frac{2}{3}b^3$. So there is no solution to $A_1 = A_2$ for $b \neq 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Using disks, we calculate the volume of rotation of $\mathcal{R}_1$ about the x-axis to be $V_{1,x} = \pi \int_0^b (x^2)^2\, dx = \frac{1}{5}\pi b^5$.

Using cylindrical shells, we calculate the volume of rotation of $\mathcal{R}_1$ about the y-axis to be

$V_{1,y} = 2\pi \int_0^b x(x^2)\, dx = 2\pi \left[\frac{1}{4}x^4\right]_0^b = \frac{1}{2}\pi b^4$. So $V_{1,x} = V_{1,y} \quad \Leftrightarrow \quad \frac{1}{5}\pi b^5 = \frac{1}{2}\pi b^4 \quad \Leftrightarrow \quad 2b = 5 \quad \Leftrightarrow \quad b = \frac{5}{2}$.

So the volumes of rotation about the x- and y-axes are the same for $b = \frac{5}{2}$.

(c) We use cylindrical shells to calculate the volume of rotation of $\mathcal{R}_2$ about the x-axis:

$\mathcal{R}_{2,x} = 2\pi \displaystyle\int_0^{b^2} y\left(\sqrt{y}\right) dy = 2\pi \left[\frac{2}{5}y^{5/2}\right]_0^{b^2} = \frac{4}{5}\pi b^5$. We already know the volume of rotation of $\mathcal{R}_1$ about the x-axis from part (b), and $\mathcal{R}_{1,x} = \mathcal{R}_{2,x} \quad \Leftrightarrow \quad \frac{1}{5}\pi b^5 = \frac{4}{5}\pi b^5$, which has no solution for $b \neq 0$.

(d) We use disks to calculate the volume of rotation of $\mathcal{R}_2$ about the y-axis: $\mathcal{R}_{2,y} = \pi \int_0^{b^2} \left(\sqrt{y}\right)^2 dy = \pi \left[\frac{1}{2}y^2\right]_0^{b^2} = \frac{1}{2}\pi b^4$.

We know the volume of rotation of $\mathcal{R}_1$ about the y-axis from part (b), and $\mathcal{R}_{1,y} = \mathcal{R}_{2,y} \quad \Leftrightarrow \quad \frac{1}{2}\pi b^4 = \frac{1}{2}\pi b^4$. But this equation is true for all b, so the volumes of rotation about the y-axis are equal for all values of b.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1. The volume generated from $x = 0$ to $x = b$ is $\int_0^b \pi[f(x)]^2\ dx$. Hence, we are given that $b^2 = \int_0^b \pi[f(x)]^2\ dx$ for all $b > 0$. Differentiating both sides of this equation with respect to b using the Fundamental Theorem of Calculus gives $2b = \pi[f(b)]^2 \quad \Rightarrow \quad f(b) = \sqrt{2b/\pi}$, since f is positive. Therefore, $f(x) = \sqrt{2x/\pi}$.

2. The total area of the region bounded by the parabola $y = x - x^2 = x(1-x)$ and the x-axis is $\int_0^1 (x - x^2)\,dx = \left[\frac{1}{2}x^2 - \frac{1}{3}x^3\right]_0^1 = \frac{1}{6}$. Let the slope of the line we are looking for be m. Then the area above this line but below the parabola is $\int_0^a \left[(x - x^2) - mx\right] dx$, where a is the x-coordinate of the point of intersection of the line and the parabola. We find the point of intersection by solving the equation $x - x^2 = mx \quad \Leftrightarrow \quad 1 - x = m \quad \Leftrightarrow \quad x = 1 - m$. So the value of a is $1 - m$, and

$$\begin{aligned}\int_0^{1-m} \left[(x - x^2) - mx\right] dx &= \int_0^{1-m} \left[(1-m)x - x^2\right] dx = \left[\tfrac{1}{2}(1-m)x^2 - \tfrac{1}{3}x^3\right]_0^{1-m} \\ &= \tfrac{1}{2}(1-m)(1-m)^2 - \tfrac{1}{3}(1-m)^3 = \tfrac{1}{6}(1-m)^3\end{aligned}$$

We want this to be half of $\frac{1}{6}$, so $\frac{1}{6}(1-m)^3 = \frac{1}{12} \quad \Leftrightarrow \quad (1-m)^3 = \frac{6}{12} \quad \Leftrightarrow \quad 1 - m = \sqrt[3]{\frac{1}{2}} \quad \Leftrightarrow \quad m = 1 - \frac{1}{\sqrt[3]{2}}$. So the slope of the required line is $1 - \frac{1}{\sqrt[3]{2}} \approx 0.206$.

3. We must find expressions for the areas A and B, and then set them equal and see what this says about the curve C. If $P = \left(a, 2a^2\right)$, then area A is just $\int_0^a (2x^2 - x^2)\,dx = \int_0^a x^2\,dx = \frac{1}{3}a^3$. To find area B, we use y as the variable of integration. So we find the equation of the middle curve as a function of y: $y = 2x^2 \quad \Leftrightarrow \quad x = \sqrt{y/2}$, since we are concerned with the first quadrant only. We can express area B as

$$\int_0^{2a^2} \left[\sqrt{y/2} - C(y)\right] dy = \left[\frac{4}{3}(y/2)^{3/2}\right]_0^{2a^2} - \int_0^{2a^2} C(y)\,dy = \frac{4}{3}a^3 - \int_0^{2a^2} C(y)\,dy$$

where $C(y)$ is the function with graph C. Setting $A = B$, we get $\frac{1}{3}a^3 = \frac{4}{3}a^3 - \int_0^{2a^2} C(y)\,dy \quad \Leftrightarrow \quad \int_0^{2a^2} C(y)\,dy = a^3$. Now we differentiate this equation with respect to a using the Chain Rule and the Fundamental Theorem: $C(2a^2)(4a) = 3a^2 \quad \Rightarrow \quad C(y) = \frac{3}{4}\sqrt{y/2}$, where $y = 2a^2$. Now we can solve for y: $x = \frac{3}{4}\sqrt{y/2} \quad \Rightarrow$ $x^2 = \frac{9}{16}(y/2) \quad \Rightarrow \quad y = \frac{32}{9}x^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a) Take slices perpendicular to the line through the center C of the bottom of the glass and the point P where the top surface of the water meets the bottom of the glass.

A typical rectangular cross-section y units above the axis of the glass has width $2\,|QR| = 2\sqrt{r^2-y^2}$ and length $h = |QS| = \dfrac{L}{2r}\,(r-y)$. [Triangles PQS and PAB are similar, so $\dfrac{h}{L} = \dfrac{|PQ|}{|PA|} = \dfrac{r-y}{2r}$.] Thus,

$$V = \int_{-r}^{r} 2\sqrt{r^2-y^2}\cdot\frac{L}{2r}\,(r-y)\,dy = L\int_{-r}^{r}\left(1-\frac{y}{r}\right)\sqrt{r^2-y^2}\,dy$$

$$= L\int_{-r}^{r}\sqrt{r^2-y^2}\,dy - \frac{L}{r}\int_{-r}^{r} y\sqrt{r^2-y^2}\,dy$$

$$= L\cdot\frac{\pi r^2}{2} - \frac{L}{r}\cdot 0 \quad \left[\begin{array}{c}\text{the first integral is the area of a semicircle of radius } r,\\ \text{and the second has an odd integrand}\end{array}\right] \quad = \frac{\pi r^2 L}{2}$$

(b) Slice parallel to the plane through the axis of the glass and the point of contact P. (This is the plane determined by P, B, and C in the figure.) $STUV$ is a typical trapezoidal slice. With respect to an x-axis with origin at C as shown, if S and V have x-coordinate x, then $|SV| = 2\sqrt{r^2-x^2}$. Projecting the trapezoid $STUV$ onto the plane of the triangle PAB (call the projection $S'T'U'V'$), we see that $|AP| = 2r$, $|SV| = 2\sqrt{r^2-x^2}$, and $|S'P| = |V'A| = \frac{1}{2}(|AP| - |SV|) = r - \sqrt{r^2-x^2}$.

By similar triangles, $\dfrac{|ST|}{|S'P|} = \dfrac{|AB|}{|AP|}$, so $|ST| = \left(r-\sqrt{r^2-x^2}\right)\cdot\dfrac{L}{2r}$. In the same way, we find that $\dfrac{|VU|}{|V'P|} = \dfrac{|AB|}{|AP|}$, so $|VU| = |V'P|\cdot\dfrac{L}{2r} = (|AP| - |V'A|)\cdot\dfrac{L}{2r} = \left(r+\sqrt{r^2-x^2}\right)\cdot\dfrac{L}{2r}$. The area $A(x)$ of the trapezoid $STUV$ is $\frac{1}{2}|SV|\cdot(|ST| + |VU|)$; that is,

$A(x) = \frac{1}{2}\cdot 2\sqrt{r^2-x^2}\cdot\left[\left(r-\sqrt{r^2-x^2}\right)\cdot\dfrac{L}{2r} + \left(r+\sqrt{r^2-x^2}\right)\cdot\dfrac{L}{2r}\right] = L\sqrt{r^2-x^2}$. Thus,

$$V = \int_{-r}^{r} A(x)\,dx = L\int_{-r}^{r}\sqrt{r^2-x^2}\,dx = L\cdot\frac{\pi r^2}{2} = \frac{\pi r^2 L}{2}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) See the computation of V in part (a) or part (b).

(d) The volume of the water is exactly half the volume of the cylindrical glass, so $V = \frac{1}{2}\pi r^2 L$.

(e) Choose x-, y-, and z-axes as shown in the figure. Then slices perpendicular to the x-axis are triangular, slices perpendicular to the y-axis are rectangular, and slices perpendicular to the z-axis are segments of circles. Using triangular slices, we find that the area $A(x)$ of a typical slice DEF, where D has x-coordinate x, is given by

$A(x) = \frac{1}{2}|DE| \cdot |EF| = \frac{1}{2}|DE| \cdot \left(\dfrac{L}{r}\,|DE|\right) = \dfrac{L}{2r}\,|DE|^2 = \dfrac{L}{2r}\left(r^2 - x^2\right)$. Thus,

$$V = \int_{-r}^{r} A(x)\,dx = \frac{L}{2r}\int_{-r}^{r}(r^2 - x^2)\,dx = \frac{L}{r}\int_{-r}^{r}(r^2 - x^2)\,dx = \frac{L}{r}\left[r^2 x - \frac{x^3}{3}\right]_0^r$$

$$= \frac{L}{r}\left(r^3 - \frac{r^3}{3}\right) = \frac{L}{r}\cdot\tfrac{2}{3}r^3 = \tfrac{2}{3}r^2 L \qquad \text{[This is } 2/(3\pi) \approx 0.21 \text{ of the volume of the glass.]}$$

5. We are given that the rate of change of the volume of water is $\dfrac{dV}{dt} = -kA(x)$, where k is some positive constant and $A(x)$ is the area of the surface when the water has depth x. Now we are concerned with the rate of change of the depth of the water with respect to time, that is, $\dfrac{dx}{dt}$. But by the Chain Rule, $\dfrac{dV}{dt} = \dfrac{dV}{dx}\dfrac{dx}{dt}$, so the first equation can be written $\dfrac{dV}{dx}\dfrac{dx}{dt} = -kA(x)$ $(\star)$. Also, we know that the total volume of water up to a depth x is $V(x) = \int_0^x A(s)\,ds$, where $A(s)$ is the area of a cross-section of the water at a depth s. Differentiating this equation with respect to x, we get $dV/dx = A(x)$. Substituting this into equation $\star$, we get $A(x)(dx/dt) = -kA(x) \quad\Rightarrow\quad dx/dt = -k$, a constant.

6. (a) The volume above the surface is $\int_0^{L-h} A(y)\,dy = \int_{-h}^{L-h} A(y)\,dy - \int_{-h}^{0} A(y)\,dy$. So the proportion of volume above the surface is $\dfrac{\int_0^{L-h} A(y)\,dy}{\int_{-h}^{L-h} A(y)\,dy} = \dfrac{\int_{-h}^{L-h} A(y)\,dy - \int_{-h}^{0} A(y)\,dy}{\int_{-h}^{L-h} A(y)\,dy}$. Now by Archimedes' Principle, we have $F = W \quad\Rightarrow$ $\rho_f g \int_{-h}^{0} A(y)\,dy = \rho_0 g \int_{-h}^{L-h} A(y)\,dy$, so $\int_{-h}^{0} A(y)\,dy = \left(\rho_0/\rho_f\right)\int_{-h}^{L-h} A(y)\,dy$. Therefore,

$$\frac{\int_0^{L-h} A(y)\,dy}{\int_{-h}^{L-h} A(y)\,dy} = \frac{\int_{-h}^{L-h} A(y)\,dy - \left(\rho_0/\rho_f\right)\int_{-h}^{L-h} A(y)\,dy}{\int_{-h}^{L-h} A(y)\,dy} = \frac{\rho_f - \rho_0}{\rho_f},$$

so the percentage of volume above the surface is $100\left(\dfrac{\rho_f - \rho_0}{\rho_f}\right)\%$.

(b) For an iceberg, the percentage of volume above the surface is $100\left(\frac{1030 - 917}{1030}\right)\% \approx 11\%$.

(c) No, the water does not overflow. Let V_i be the volume of the ice cube, and let V_w be the volume of the water which results from the melting. Then by the formula derived in part (a), the volume of ice above the surface of the water is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\left[(\rho_f - \rho_0)/\rho_f\right]V_i$, so the volume below the surface is $V_i - \left[(\rho_f - \rho_0)/\rho_f\right]V_i = (\rho_0/\rho_f)V_i$. Now the mass of the ice cube is the same as the mass of the water which is created when it melts, namely $m = \rho_0 V_i = \rho_f V_w \quad \Rightarrow$ $V_w = (\rho_0/\rho_f)V_i$. So when the ice cube melts, the volume of the resulting water is the same as the underwater volume of the ice cube, and so the water does not overflow.

(d) The figure shows the instant when the height of the exposed part of the ball is y. Using the volume formula from Exercise 6.2.61, $V = \frac{1}{3}\pi h^2(3r - h)$, with $r = 0.4$ and $h = 0.8 - y$, we see that the volume of the submerged part of the sphere is $\frac{1}{3}\pi(0.8 - y)^2[1.2 - (0.8 - y)]$, so its weight is $1000g \cdot \frac{1}{3}\pi s^2(1.2 - s)$, where $s = 0.8 - y$. Then the work done to submerge the sphere is

$$W = \int_0^{0.8} g\tfrac{1000}{3}\pi s^2(1.2 - s)\,ds = g\tfrac{1000}{3}\pi \int_0^{0.8}(1.2s^2 - s^3)\,ds$$
$$= g\tfrac{1000}{3}\pi\left[0.4s^3 - \tfrac{1}{4}s^4\right]_0^{0.8} = g\tfrac{1000}{3}\pi(0.2048 - 0.1024) = 9.8\,\tfrac{1000}{3}\,\pi(0.1024) \approx 1.05 \times 10^3 \text{ J}$$

7. A typical sphere of radius r is shown in the figure. We wish to maximize the shaded volume V, which can be thought of as the volume of a hemisphere of radius r minus the volume of the spherical cap with height $h = 1 - \sqrt{1 - r^2}$ and radius 1.

$$V = \tfrac{1}{2}\cdot\tfrac{4}{3}\pi r^3 - \tfrac{1}{3}\pi\left(1 - \sqrt{1 - r^2}\right)^2\left[3(1) - \left(1 - \sqrt{1 - r^2}\right)\right] \quad \text{[by Exercise 6.2.61]}$$
$$= \tfrac{1}{3}\pi\left[2r^3 - \left(2 - 2\sqrt{1 - r^2} - r^2\right)\left(2 + \sqrt{1 - r^2}\right)\right]$$
$$= \tfrac{1}{3}\pi\left[2r^3 - 2 + \left(r^2 + 2\right)\sqrt{1 - r^2}\right]$$
$$V' = \tfrac{1}{3}\pi\left[6r^2 + \frac{\left(r^2 + 2\right)(-r)}{\sqrt{1 - r^2}} + \sqrt{1 - r^2}(2r)\right] = \tfrac{1}{3}\pi\left[\frac{6r^2\sqrt{1 - r^2} - r\left(r^2 + 2\right) + 2r\left(1 - r^2\right)}{\sqrt{1 - r^2}}\right]$$
$$= \tfrac{1}{3}\pi\left(\frac{6r^2\sqrt{1 - r^2} - 3r^3}{\sqrt{1 - r^2}}\right) = \frac{\pi r^2\left(2\sqrt{1 - r^2} - r\right)}{\sqrt{1 - r^2}}$$

$V'(r) = 0 \quad \Leftrightarrow \quad 2\sqrt{1 - r^2} = r \quad \Leftrightarrow \quad 4 - 4r^2 = r^2 \quad \Leftrightarrow \quad r^2 = \frac{4}{5} \quad \Leftrightarrow \quad r = \frac{2}{\sqrt{5}} \approx 0.89.$

Since $V'(r) > 0$ for $0 < r < \frac{2}{\sqrt{5}}$ and $V'(r) < 0$ for $\frac{2}{\sqrt{5}} < r < 1$, we know that V attains a maximum at $r = \frac{2}{\sqrt{5}}$.

8. We want to find the volume of that part of the sphere which is below the surface of the water. As we can see from the diagram, this region is a cap of a sphere with radius r and height $r + d$. If we can find an expression for d in terms of h, r and θ, then we can determine the volume of the region [see Exercise 6.2.61], and then differentiate with respect to r to find the maximum. We see that

$$\sin\theta = \frac{r}{h - d} \quad \Leftrightarrow \quad h - d = \frac{r}{\sin\theta} \quad \Leftrightarrow \quad d = h - r\csc\theta.$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Now we can use the formula from Exercise 6.2.61 to find the volume of water displaced:

$$V = \tfrac{1}{3}\pi h^2(3r-h) = \tfrac{1}{3}\pi(r+d)^2\,[3r-(r+d)] = \tfrac{1}{3}\pi(r+h-r\csc\theta)^2(2r-h+r\csc\theta)$$
$$= \tfrac{\pi}{3}[r(1-\csc\theta)+h]^2[r(2+\csc\theta)-h]$$

Now we differentiate with respect to r:

$$dV/dr = \tfrac{\pi}{3}\big([r(1-\csc\theta)+h]^2(2+\csc\theta)+2[r(1-\csc\theta)+h](1-\csc\theta)[r(2+\csc\theta)-h]\big)$$
$$= \tfrac{\pi}{3}[r(1-\csc\theta)+h]([r(1-\csc\theta)+h](2+\csc\theta)+2(1-\csc\theta)[r(2+\csc\theta)-h])$$
$$= \tfrac{\pi}{3}[r(1-\csc\theta)+h](3(2+\csc\theta)(1-\csc\theta)r+[(2+\csc\theta)-2(1-\csc\theta)]h)$$
$$= \tfrac{\pi}{3}[r(1-\csc\theta)+h][3(2+\csc\theta)(1-\csc\theta)r+3h\csc\theta]$$

This is 0 when $r = \dfrac{h}{\csc\theta-1}$ and when $r = \dfrac{h\csc\theta}{(\csc\theta+2)(\csc\theta-1)}$. Now since $V\left(\dfrac{h}{\csc\theta-1}\right) = 0$ (the first factor vanishes; this corresponds to $d=-r$), the maximum volume of water is displaced when $r = \dfrac{h\csc\theta}{(\csc\theta-1)(\csc\theta+2)}$. (Our intuition tells us that a maximum value does exist, and it must occur at a critical number.) Multiplying numerator and denominator by $\sin^2\theta$, we get an alternative form of the answer: $r = \dfrac{h\sin\theta}{\sin\theta+\cos 2\theta}$.

9. (a) Stacking disks along the y-axis gives us $V = \int_0^h \pi\,[f(y)]^2\,dy$.

(b) Using the Chain Rule, $\dfrac{dV}{dt} = \dfrac{dV}{dh}\cdot\dfrac{dh}{dt} = \pi\,[f(h)]^2\,\dfrac{dh}{dt}$.

(c) $kA\sqrt{h} = \pi[f(h)]^2\,\dfrac{dh}{dt}$. Set $\dfrac{dh}{dt} = C$: $\pi[f(h)]^2\,C = kA\sqrt{h} \;\Rightarrow\; [f(h)]^2 = \dfrac{kA}{\pi C}\sqrt{h} \;\Rightarrow\; f(h) = \sqrt{\dfrac{kA}{\pi C}}\,h^{1/4}$; that is, $f(y) = \sqrt{\dfrac{kA}{\pi C}}\,y^{1/4}$. The advantage of having $\dfrac{dh}{dt} = C$ is that the markings on the container are equally spaced.

10. (a) We first use the cylindrical shell method to express the volume V in terms of h, r, and ω:

$$V = \int_0^r 2\pi xy\,dx = \int_0^r 2\pi x\left[h+\frac{\omega^2x^2}{2g}\right]dx = 2\pi\int_0^r\left(hx+\frac{\omega^2x^3}{2g}\right)dx$$
$$= 2\pi\left[\frac{hx^2}{2}+\frac{\omega^2x^4}{8g}\right]_0^r = 2\pi\left[\frac{hr^2}{2}+\frac{\omega^2r^4}{8g}\right] = \pi hr^2+\frac{\pi\omega^2r^4}{4g} \quad\Rightarrow$$

$h = \dfrac{V-\big(\pi\omega^2r^4\big)/(4g)}{\pi r^2} = \dfrac{4gV-\pi\omega^2r^4}{4\pi gr^2}$.

(b) The surface touches the bottom when $h=0 \;\Rightarrow\; 4gV-\pi\omega^2r^4=0 \;\Rightarrow\; \omega^2 = \dfrac{4gV}{\pi r^4} \;\Rightarrow\; \omega = \dfrac{2\sqrt{gV}}{\sqrt{\pi}r^2}$.

To spill over the top, $y(r) > L \;\Leftrightarrow$

$$L < h+\frac{\omega^2r^2}{2g} = \frac{4gV-\pi\omega^2r^4}{4\pi gr^2}+\frac{\omega^2r^2}{2g} = \frac{4gV}{4\pi gr^2}-\frac{\pi\omega^2r^2}{4\pi gr^2}+\frac{\omega^2r^2}{2g}$$
$$= \frac{V}{\pi r^2}-\frac{\omega^2r^2}{4g}+\frac{\omega^2r^2}{2g} = \frac{V}{\pi r^2}+\frac{\omega^2r^2}{4g} \quad\Leftrightarrow$$

$\dfrac{\omega^2r^2}{4g} > L-\dfrac{V}{\pi r^2} = \dfrac{\pi r^2L-V}{\pi r^2} \;\Leftrightarrow\; \omega^2 > \dfrac{4g\big(\pi r^2L-V\big)}{\pi r^4}$. So for spillage, the angular speed should be $\omega > \dfrac{2\sqrt{g(\pi r^2L-V)}}{r^2\sqrt{\pi}}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) (i) Here we have $r = 2$, $L = 7$, $h = 7 - 5 = 2$. When $x = 1$, $y = 7 - 4 = 3$. Therefore, $3 = 2 + \dfrac{\omega^2 \cdot 1^2}{2 \cdot 32} \quad \Rightarrow$

$1 = \dfrac{\omega^2}{2 \cdot 32} \quad \Rightarrow \quad \omega^2 = 64 \quad \Rightarrow \quad \omega = 8$ rad/s. $V = \pi(2)(2)^2 + \dfrac{\pi \cdot 8^2 \cdot 2^4}{4g} = 8\pi + 8\pi = 16\pi$ ft^2.

(ii) At the wall, $x = 2$, so $y = 2 + \dfrac{8^2 \cdot 2^2}{2 \cdot 32} = 6$ and the surface is $7 - 6 = 1$ ft below the top of the tank.

11. The cubic polynomial passes through the origin, so let its equation be $y = px^3 + qx^2 + rx$. The curves intersect when $px^3 + qx^2 + rx = x^2 \quad \Leftrightarrow$ $px^3 + (q-1)x^2 + rx = 0$. Call the left side $f(x)$. Since $f(a) = f(b) = 0$, another form of f is

$$\begin{aligned} f(x) &= px(x-a)(x-b) = px[x^2 - (a+b)x + ab] \\ &= p[x^3 - (a+b)x^2 + abx] \end{aligned}$$

Since the two areas are equal, we must have $\int_0^a f(x)\,dx = -\int_a^b f(x)\,dx \quad \Rightarrow$ $[F(x)]_0^a = [F(x)]_b^a \quad \Rightarrow \quad F(a) - F(0) = F(a) - F(b) \quad \Rightarrow \quad F(0) = F(b)$, where F is an antiderivative of f. Now $F(x) = \int f(x)\,dx = \int p[x^3 - (a+b)x^2 + abx]\,dx = p\left[\frac{1}{4}x^4 - \frac{1}{3}(a+b)x^3 + \frac{1}{2}abx^2\right] + C$, so $F(0) = F(b) \quad \Rightarrow \quad C = p\left[\frac{1}{4}b^4 - \frac{1}{3}(a+b)b^3 + \frac{1}{2}ab^3\right] + C \quad \Rightarrow \quad 0 = p\left[\frac{1}{4}b^4 - \frac{1}{3}(a+b)b^3 + \frac{1}{2}ab^3\right] \quad \Rightarrow$ $0 = 3b - 4(a+b) + 6a$ [multiply by $12/(pb^3)$, $b \neq 0$] $\Rightarrow \quad 0 = 3b - 4a - 4b + 6a \quad \Rightarrow \quad b = 2a$. Hence, b is twice the value of a.

12. (a) Place the round flat tortilla on an xy-coordinate system as shown in the first figure. An equation of the circle is $x^2 + y^2 = 4^2$ and the height of a cross-section is $2\sqrt{16 - x^2}$.

Now look at a cross-section with central angle θ_x as shown in the second figure (r is the radius of the circular cylinder). The filled area $A(x)$ is equal to the area $A_1(x)$ of the sector minus the area $A_2(x)$ of the triangle.

$$\begin{aligned} A(x) = A_1(x) - A_2(x) &= \tfrac{1}{2}r^2\theta_x - \tfrac{1}{2}r^2 \sin\theta_x \qquad \text{[area formulas from trigonometry]} \\ &= \tfrac{1}{2}r(r\theta_x) - \tfrac{1}{2}r^2 \sin\left(\frac{s}{r}\right) \qquad [\text{arc length } s = r\theta_x \quad \Rightarrow \quad \theta_x = s/r] \\ &= \tfrac{1}{2}r \cdot 2\sqrt{16 - x^2} - \tfrac{1}{2}r^2 \sin\left(\frac{2\sqrt{16 - x^2}}{r}\right) \qquad [s = 2\sqrt{16 - x^2}\,] \\ &= r\sqrt{16 - x^2} - \tfrac{1}{2}r^2 \sin\left(\frac{2}{r}\sqrt{16 - x^2}\right) \qquad (\star) \end{aligned}$$

Note that the central angle θ_x will be small near the ends of the tortilla; that is, when $|x| \approx 4$. But near the center of

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

the tortilla (when $|x| \approx 0$), the central angle θ_x may exceed $180°$. Thus, the sine of θ_x will be negative and the second term in ($\star$) will be positive (actually adding area to the area of the sector). The volume of the taco can be found by integrating the cross-sectional areas from $x = -4$ to $x = 4$. Thus,

$$V(x) = \int_{-4}^{4} A(x)\,dx = \int_{-4}^{4} \left[r\sqrt{16 - x^2} - \tfrac{1}{2}r^2 \sin\left(\frac{2}{r}\sqrt{16 - x^2}\right)\right] dx$$

(b) To find the value of r that maximizes the volume of the taco, we can define the function

$$V(r) = \int_{-4}^{4} \left[r\sqrt{16 - x^2} - \tfrac{1}{2}r^2 \sin\left(\frac{2}{r}\sqrt{16 - x^2}\right)\right] dx$$

The figure shows a graph of $y = V(r)$ and $y = V'(r)$. The maximum volume of about 52.94 occurs when $r \approx 2.2912$.

13. We assume that P lies in the region of positive x. Since $y = x^3$ is an odd function, this assumption will not affect the result of the calculation. Let $P = (a, a^3)$. The slope of the tangent to the curve $y = x^3$ at P is $3a^2$, and so the equation of the tangent is $y - a^3 = 3a^2(x - a) \quad \Leftrightarrow \quad y = 3a^2x - 2a^3$.

We solve this simultaneously with $y = x^3$ to find the other point of intersection: $x^3 = 3a^2x - 2a^3 \quad \Leftrightarrow$ $(x - a)^2(x + 2a) = 0$. So $Q = (-2a, -8a^3)$ is the other point of intersection. The equation of the tangent at Q is $y - (-8a^3) = 12a^2[x - (-2a)] \quad \Leftrightarrow \quad y = 12a^2x + 16a^3$. By symmetry, this tangent will intersect the curve again at $x = -2(-2a) = 4a$. The curve lies above the first tangent, and below the second, so we are looking for a relationship between $A = \int_{-2a}^{a} \left[x^3 - (3a^2x - 2a^3)\right] dx$ and $B = \int_{-2a}^{4a} \left[(12a^2x + 16a^3) - x^3\right] dx$. We calculate $A = \left[\frac{1}{4}x^4 - \frac{3}{2}a^2x^2 + 2a^3x\right]_{-2a}^{a} = \frac{3}{4}a^4 - (-6a^4) = \frac{27}{4}a^4$, and $B = \left[6a^2x^2 + 16a^3x - \frac{1}{4}x^4\right]_{-2a}^{4a} = 96a^4 - (-12a^4) = 108a^4$. We see that $B = 16A = 2^4A$. This is because our calculation of area B was essentially the same as that of area A, with a replaced by $-2a$, so if we replace a with $-2a$ in our expression for A, we get $\frac{27}{4}(-2a)^4 = 108a^4 = B$.

14. From the solution to Problem 11 in Problems Plus following Chapter 4, an equation of the normal line through P is $y - a^2 = -\dfrac{1}{2a}(x - a) \quad \Rightarrow \quad y = -\dfrac{1}{2a}x + \dfrac{1}{2} + a^2$, and the x-coordinate of Q is $x = -a - \dfrac{1}{2a}$. The area of $\mathcal{R}$ is given by

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$A(a) = \int_{-a-1/(2a)}^{a} \left[\left(-\frac{1}{2a}x + \frac{1}{2} + a^2\right) - x^2\right] dx = \left[-\frac{1}{4a}x^2 + \frac{1}{2}x + a^2x - \frac{1}{3}x^3\right]_{-a-1/(2a)}^{a}$$

$$= \left(-\frac{1}{4}a + \frac{1}{2}a + a^3 - \frac{1}{3}a^3\right) - \left[-\frac{1}{4a}\left(-a - \frac{1}{2a}\right)^2 + \frac{1}{2}\left(-a - \frac{1}{2a}\right) + a^2\left(-a - \frac{1}{2a}\right) - \frac{1}{3}\left(-a - \frac{1}{2a}\right)^3\right]$$

$$= \left(\frac{1}{4}a + \frac{2}{3}a^3\right) - \left[-\frac{1}{4a}\left(a^2 + 1 + \frac{1}{4a^2}\right) - \frac{1}{2}a - \frac{1}{4a} - a^3 - \frac{1}{2}a - \frac{1}{3}\left(-a^3 - \frac{3}{2}a - \frac{3}{4a} - \frac{1}{8a^3}\right)\right]$$

$$= \left(\frac{1}{4}a + \frac{2}{3}a^3\right) - \left(-\frac{2}{3}a^3 - \frac{1}{48a^3} - \frac{3}{4}a - \frac{1}{4a}\right)$$

$$= \frac{4}{3}a^3 + \frac{1}{48a^3} + a + \frac{1}{4a} = \frac{64a^6 + 48a^4 + 12a^2 + 1}{48a^3} = \frac{(4a^2+1)^3}{48a^3}$$

$$A'(a) = \frac{48a^3 \cdot 3(4a^2+1)^2 \cdot 8a - (4a^2+1)^3 \cdot 144a^2}{(48a^3)^2} = \frac{48 \cdot 3a^2(4a^2+1)^2[a \cdot 8a - (4a^2+1)]}{48 \cdot 48 \cdot a^3 \cdot a^3}$$

$$= \frac{(4a^2+1)^2(4a^2-1)}{16a^4} = 0 \quad \Rightarrow \quad a^2 = \frac{1}{4} \quad \Rightarrow \quad a = \frac{1}{2} \ (a > 0)$$

Since $A'(a) < 0$ for $0 < a < \frac{1}{2}$ and $A'(a) > 0$ for $a > \frac{1}{2}$, there is an absolute minimum when $a = \frac{1}{2}$ by the First Derivative Test for Absolute Extreme Values.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7 ☐ TECHNIQUES OF INTEGRATION

7.1 Integration by Parts

1. Let $u = x$, $dv = e^{2x}\,dx \quad \Rightarrow \quad du = dx$, $v = \frac{1}{2}e^{2x}$. Then by Equation 2,

$\int xe^{2x}\,dx = \frac{1}{2}xe^{2x} - \int \frac{1}{2}e^{2x}\,dx = \frac{1}{2}xe^{2x} - \frac{1}{4}e^{2x} + C$.

2. Let $u = \ln x$, $dv = \sqrt{x}\,dx \quad \Rightarrow \quad du = \dfrac{1}{x}\,dx$, $v = \frac{2}{3}x^{3/2}$. Then by Equation 2,

$$\int \sqrt{x}\ln x\,dx = \tfrac{2}{3}x^{3/2}\ln x - \int \tfrac{2}{3}x^{3/2}\cdot\frac{1}{x}\,dx = \tfrac{2}{3}x^{3/2}\ln x - \int \tfrac{2}{3}x^{1/2}\,dx = \tfrac{2}{3}x^{3/2}\ln x - \tfrac{4}{9}x^{3/2} + C.$$

3. Let $u = x$, $dv = \cos 4x\,dx \quad \Rightarrow \quad du = dx$, $v = \frac{1}{4}\sin 4x$. Then by Equation 2,

$\int x\cos 4x\,dx = \frac{1}{4}x\sin 4x - \int \frac{1}{4}\sin 4x\,dx = \frac{1}{4}x\sin 4x + \frac{1}{16}\cos 4x + C$.

4. Let $u = \sin^{-1}x$, $dv = dx \quad \Rightarrow \quad du = \dfrac{1}{\sqrt{1-x^2}}\,dx$, $v = x$. Then by Equation 2,

$$\begin{aligned}\int \sin^{-1}x\,dx &= x\sin^{-1}x - \int \frac{x}{\sqrt{1-x^2}}\,dx = x\sin^{-1}x - \int \frac{1}{\sqrt{t}}\left(-\frac{1}{2}\,dt\right) \qquad \begin{bmatrix} t = 1-x^2, \\ dt = -2x\,dx \end{bmatrix} \\ &= x\sin^{-1}x + \tfrac{1}{2}\int t^{-1/2}\,dt = x\sin^{-1}x + \tfrac{1}{2}\cdot 2t^{1/2} + C = x\sin^{-1}x + \sqrt{1-x^2} + C\end{aligned}$$

Note: A mnemonic device which is helpful for selecting u when using integration by parts is the LIATE principle of precedence for u:

Logarithmic
Inverse trigonometric
Algebraic
Trigonometric
Exponential

If the integrand has several factors, then we try to choose among them a u which appears as high as possible on the list. For example, in $\int xe^{2x}\,dx$ the integrand is xe^{2x}, which is the product of an algebraic function (x) and an exponential function (e^{2x}). Since Algebraic appears before Exponential, we choose $u = x$. Sometimes the integration turns out to be similar regardless of the selection of u and dv, but it is advisable to refer to LIATE when in doubt.

5. Let $u = t$, $dv = e^{2t}\,dt \quad \Rightarrow \quad du = dt$, $v = \frac{1}{2}e^{2t}$. Then by Equation 2,

$\int te^{2t}\,dt = \frac{1}{2}te^{2t} - \int \frac{1}{2}e^{2t}\,dt = \frac{1}{2}te^{2t} - \frac{1}{4}e^{2t} + C$.

6. Let $u = y$, $dv = e^{-y}\,dy \quad \Rightarrow \quad du = dy$, $v = -e^{-y}$. Then by Equation 2,

$\int ye^{-y}\,dy = -ye^{-y} - \int -e^{-y}\,dy = -ye^{-y} + \int e^{-y}\,dy = -ye^{-y} - e^{-y} + C$.

7. Let $u = x$, $dv = \sin 10x\,dx \quad \Rightarrow \quad du = dx$, $v = -\frac{1}{10}\cos 10x$. Then by Equation 2,

$$\begin{aligned}\int x\sin 10x\,dx &= -\tfrac{1}{10}x\cos 10x - \int -\tfrac{1}{10}\cos 10x\,dx = -\tfrac{1}{10}x\cos 10x + \tfrac{1}{10}\int \cos 10x\,dx \\ &= -\tfrac{1}{10}x\cos 10x + \tfrac{1}{100}\sin 10x + C\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. Let $u = \pi - x$, $dv = \cos \pi x\, dx \quad \Rightarrow \quad du = -dx$, $v = \dfrac{1}{\pi} \sin \pi x$. Then by Equation 2,

$$\int (\pi - x) \cos \pi x\, dx = \frac{1}{\pi}(\pi - x) \sin \pi x - \int -\frac{1}{\pi} \sin \pi x\, dx = \frac{1}{\pi}(\pi - x) \sin \pi x - \frac{1}{\pi^2} \cos \pi x + C.$$

9. Let $u = \ln w$, $dv = w\, dw \quad \Rightarrow \quad du = \dfrac{1}{w}\, dw$, $v = \dfrac{1}{2} w^2$. Then by Equation 2,

$$\int w \ln w\, dw = \frac{1}{2} w^2 \ln w - \int \frac{1}{2} w^2 \cdot \frac{1}{w}\, dw = \frac{1}{2} w^2 \ln w - \frac{1}{2} \int w\, dw = \frac{1}{2} w^2 \ln w - \frac{1}{4} w^2 + C.$$

10. Let $u = \ln x$, $dv = \dfrac{1}{x^2}\, dx = x^{-2}\, dx \quad \Rightarrow \quad du = \dfrac{1}{x}\, dx = x^{-1}\, dx$, $v = -x^{-1}$. Then by Equation 2,

$$\int \frac{\ln x}{x^2}\, dx = -\frac{\ln x}{x} - \int -x^{-1} \cdot x^{-1}\, dx = -\frac{\ln x}{x} + \int x^{-2}\, dx = -\frac{\ln x}{x} - x^{-1} + C = -\frac{\ln x}{x} - \frac{1}{x} + C.$$

11. First let $u = x^2 + 2x$, $dv = \cos x\, dx \quad \Rightarrow \quad du = (2x + 2)\, dx$, $v = \sin x$. Then by Equation 2,

$I = \int (x^2 + 2x) \cos x\, dx = (x^2 + 2x) \sin x - \int (2x + 2) \sin x\, dx$. Next let $U = 2x + 2$, $dV = \sin x\, dx \quad \Rightarrow$

$dU = 2\, dx$, $V = -\cos x$, so $\int (2x + 2) \sin x\, dx = -(2x + 2) \cos x - \int -2 \cos x\, dx = -(2x + 2) \cos x + 2 \sin x$.

Thus, $I = (x^2 + 2x) \sin x + (2x + 2) \cos x - 2 \sin x + C$.

12. First let $u = t^2$, $dv = \sin \beta t\, dt \quad \Rightarrow \quad du = 2t\, dt$, $v = -\frac{1}{\beta} \cos \beta t$. Then by Equation 2,

$I = \int t^2 \sin \beta t\, dt = -\frac{1}{\beta} t^2 \cos \beta t - \int -\frac{2}{\beta} t \cos \beta t\, dt$. Next let $U = t$, $dV = \cos \beta t\, dt \quad \Rightarrow \quad dU = dt$,

$V = \dfrac{1}{\beta} \sin \beta t$, so $\int t \cos \beta t\, dt = \dfrac{1}{\beta} t \sin \beta t - \displaystyle\int \frac{1}{\beta} \sin \beta t\, dt = \frac{1}{\beta} t \sin \beta t + \frac{1}{\beta^2} \cos \beta t$. Thus,

$$I = -\frac{1}{\beta} t^2 \cos \beta t + \frac{2}{\beta}\left(\frac{1}{\beta} t \sin \beta t + \frac{1}{\beta^2} \cos \beta t\right) + C = -\frac{1}{\beta} t^2 \cos \beta t + \frac{2}{\beta^2} t \sin \beta t + \frac{2}{\beta^3} \cos \beta t + C.$$

13. Let $u = \cos^{-1} x$, $dv = dx \quad \Rightarrow \quad du = \dfrac{-1}{\sqrt{1 - x^2}}\, dx$, $v = x$. Then by Equation 2,

$$\begin{aligned} \int \cos^{-1} x\, dx &= x \cos^{-1} x - \int \frac{-x}{\sqrt{1 - x^2}}\, dx = x \cos^{-1} x - \int \frac{1}{\sqrt{t}} \left(\frac{1}{2}\, dt\right) \qquad \left[\begin{smallmatrix} t = 1 - x^2, \\ dt = -2x\, dx \end{smallmatrix}\right] \\ &= x \cos^{-1} x - \tfrac{1}{2} \cdot 2t^{1/2} + C = x \cos^{-1} x - \sqrt{1 - x^2} + C \end{aligned}$$

14. Let $u = \ln \sqrt{x}$, $dv = dx \quad \Rightarrow \quad du = \dfrac{1}{\sqrt{x}} \cdot \dfrac{1}{2\sqrt{x}}\, dx = \dfrac{1}{2x}\, dx$, $v = x$. Then by Equation 2,

$$\int \ln \sqrt{x}\, dx = x \ln \sqrt{x} - \int x \cdot \frac{1}{2x}\, dx = x \ln \sqrt{x} - \int \frac{1}{2}\, dx = x \ln \sqrt{x} - \frac{1}{2} x + C.$$

Note: We could start by using $\ln \sqrt{x} = \frac{1}{2} \ln x$.

15. Let $u = \ln t$, $dv = t^4\, dt \quad \Rightarrow \quad du = \dfrac{1}{t}\, dt$, $v = \dfrac{1}{5} t^5$. Then by Equation 2,

$$\int t^4 \ln t\, dt = \frac{1}{5} t^5 \ln t - \int \frac{1}{5} t^5 \cdot \frac{1}{t}\, dt = \frac{1}{5} t^5 \ln t - \int \frac{1}{5} t^4\, dt = \frac{1}{5} t^5 \ln t - \frac{1}{25} t^5 + C.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. Let $u = \tan^{-1}(2y)$, $dv = dy \quad \Rightarrow \quad du = \dfrac{2}{1+4y^2}\,dy$, $v = y$. Then by Equation 2,

$$\int \tan^{-1}(2y)\,dy = y\tan^{-1}(2y) - \int \frac{2y}{1+4y^2}\,dy = y\tan^{-1}(2y) - \int \frac{1}{t}\left(\frac{1}{4}\,dt\right) \qquad \begin{bmatrix} t = 1+4y^2, \\ dt = 8y\,dy \end{bmatrix}$$

$$= y\tan^{-1}(2y) - \tfrac{1}{4}\ln|t| + C = y\tan^{-1}(2y) - \tfrac{1}{4}\ln(1+4y^2) + C$$

17. Let $u = t$, $dv = \csc^2 t\,dt \quad \Rightarrow \quad du = dt$, $v = -\cot t$. Then by Equation 2,

$$\int t\csc^2 t\,dt = -t\cot t - \int -\cot t\,dt = -t\cot t + \int \frac{\cos t}{\sin t}\,dt = -t\cot t + \int \frac{1}{z}\,dz \qquad \begin{bmatrix} z = \sin t, \\ dz = \cos t\,dt \end{bmatrix}$$

$$= -t\cot t + \ln|z| + C = -t\cot t + \ln|\sin t| + C$$

18. Let $u = x$, $dv = \cosh ax\,dx \quad \Rightarrow \quad du = dx$, $v = \dfrac{1}{a}\sinh ax$. Then by Equation 2,

$$\int x\cosh ax\,dx = \frac{1}{a}x\sinh ax - \int \frac{1}{a}\sinh ax\,dx = \frac{1}{a}x\sinh ax - \frac{1}{a^2}\cosh ax + C.$$

19. First let $u = (\ln x)^2$, $dv = dx \quad \Rightarrow \quad du = 2\ln x \cdot \frac{1}{x}\,dx$, $v = x$. Then by Equation 2,

$I = \int (\ln x)^2\,dx = x(\ln x)^2 - 2\int x\ln x \cdot \frac{1}{x}\,dx = x(\ln x)^2 - 2\int \ln x\,dx$. Next let $U = \ln x$, $dV = dx \quad \Rightarrow$

$dU = 1/x\,dx$, $V = x$ to get $\int \ln x\,dx = x\ln x - \int x\cdot(1/x)\,dx = x\ln x - \int dx = x\ln x - x + C_1$. Thus,

$I = x(\ln x)^2 - 2(x\ln x - x + C_1) = x(\ln x)^2 - 2x\ln x + 2x + C$, where $C = -2C_1$.

20. $\displaystyle\int \frac{z}{10^z}\,dz = \int z\,10^{-z}\,dz$. Let $u = z$, $dv = 10^{-z}\,dz \quad \Rightarrow \quad du = dz$, $v = \dfrac{-10^{-z}}{\ln 10}$. Then by Equation 2,

$$\int z\,10^{-z}\,dz = \frac{-z\,10^{-z}}{\ln 10} - \int \frac{-10^{-z}}{\ln 10}\,dz = \frac{-z}{10^z\ln 10} - \frac{10^{-z}}{(\ln 10)(\ln 10)} + C = -\frac{z}{10^z\ln 10} - \frac{1}{10^z(\ln 10)^2} + C.$$

21. First let $u = e^{3x}$, $dv = \cos x\,dx \quad \Rightarrow \quad du = 3e^{3x}\,dx$, $v = \sin x$. Then

$I = \int e^{3x}\cos x\,dx = e^{3x}\sin x - 3\int e^{3x}\sin x\,dx$. Next, let $U = e^{3x}$, $dV = \sin x\,dx \quad \Rightarrow \quad dU = 3e^{3x}\,dx$, $V = -\cos x$,

so $\int e^{3x}\sin x\,dx = -e^{3x}\cos x + 3\int e^{3x}\cos x\,dx$. Substituting in the previous formula gives

$I = e^{3x}\sin x - 3(-e^{3x}\cos x + 3I) = e^{3x}\sin x + 3e^{3x}\cos x - 9I \quad \Rightarrow \quad 10I = e^{3x}\sin x + 3e^{3x}\cos x + C_1 \quad \Rightarrow$

$I = \frac{1}{10}e^{3x}\sin x + \frac{3}{10}e^{3x}\cos x + C$, where $C = \frac{1}{10}C_1$.

22. First let $u = e^x$, $dv = \sin \pi x\,dx \quad \Rightarrow \quad du = e^x\,dx$, $v = -\dfrac{1}{\pi}\cos \pi x$. Then

$I = \displaystyle\int e^x\sin \pi x\,dx = -\frac{1}{\pi}e^x\cos \pi x + \frac{1}{\pi}\int e^x\cos \pi x\,dx$. Next, let $U = e^x$, $dV = \cos \pi x\,dx \quad \Rightarrow$

$dU = e^x\,dx$, $V = \dfrac{1}{\pi}\sin \pi x$, so $\displaystyle\int e^x\cos \pi x\,dx = \frac{1}{\pi}e^x\sin \pi x - \frac{1}{\pi}\int e^x\sin \pi x\,dx$. Substituting in the previous formula

gives $I = -\dfrac{1}{\pi}e^x\cos \pi x + \dfrac{1}{\pi}\left(\dfrac{1}{\pi}e^x\sin \pi x - \dfrac{1}{\pi}I\right) = -\dfrac{1}{\pi}e^x\cos \pi x + \dfrac{1}{\pi^2}e^x\sin \pi x - \dfrac{1}{\pi^2}I \quad \Rightarrow$

$\left(1 + \dfrac{1}{\pi^2}\right)I = -\dfrac{1}{\pi}e^x\cos \pi x + \dfrac{1}{\pi^2}e^x\sin \pi x + C_1 \quad \Rightarrow \quad I = -\dfrac{\pi}{\pi^2+1}\,e^x\cos \pi x + \dfrac{1}{\pi^2+1}\,e^x\sin \pi x + C,$

where $C = \dfrac{\pi^2}{\pi^2+1}\,C_1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. First let $u = \sin 3\theta$, $dv = e^{2\theta}\,d\theta \;\Rightarrow\; du = 3\cos 3\theta\,d\theta$, $v = \frac{1}{2}e^{2\theta}$. Then

$I = \int e^{2\theta}\sin 3\theta\,d\theta = \frac{1}{2}e^{2\theta}\sin 3\theta - \frac{3}{2}\int e^{2\theta}\cos 3\theta\,d\theta$. Next let $U = \cos 3\theta$, $dV = e^{2\theta}\,d\theta \;\Rightarrow\; dU = -3\sin 3\theta\,d\theta$,

$V = \frac{1}{2}e^{2\theta}$ to get $\int e^{2\theta}\cos 3\theta\,d\theta = \frac{1}{2}e^{2\theta}\cos 3\theta + \frac{3}{2}\int e^{2\theta}\sin 3\theta\,d\theta$. Substituting in the previous formula gives

$I = \frac{1}{2}e^{2\theta}\sin 3\theta - \frac{3}{4}e^{2\theta}\cos 3\theta - \frac{9}{4}\int e^{2\theta}\sin 3\theta\,d\theta = \frac{1}{2}e^{2\theta}\sin 3\theta - \frac{3}{4}e^{2\theta}\cos 3\theta - \frac{9}{4}I \;\Rightarrow$

$\frac{13}{4}I = \frac{1}{2}e^{2\theta}\sin 3\theta - \frac{3}{4}e^{2\theta}\cos 3\theta + C_1$. Hence, $I = \frac{1}{13}e^{2\theta}(2\sin 3\theta - 3\cos 3\theta) + C$, where $C = \frac{4}{13}C_1$.

24. First let $u = e^{-\theta}$, $dv = \cos 2\theta\,d\theta \;\Rightarrow\; du = -e^{-\theta}\,d\theta$, $v = \frac{1}{2}\sin 2\theta$. Then

$I = \int e^{-\theta}\cos 2\theta\,d\theta = \frac{1}{2}e^{-\theta}\sin 2\theta - \int \frac{1}{2}\sin 2\theta\left(-e^{-\theta}\,d\theta\right) = \frac{1}{2}e^{-\theta}\sin 2\theta + \frac{1}{2}\int e^{-\theta}\sin 2\theta\,d\theta$.

Next let $U = e^{-\theta}$, $dV = \sin 2\theta\,d\theta \;\Rightarrow\; dU = -e^{-\theta}\,d\theta$, $V = -\frac{1}{2}\cos 2\theta$, so

$\int e^{-\theta}\sin 2\theta\,d\theta = -\frac{1}{2}e^{-\theta}\cos 2\theta - \int\left(-\frac{1}{2}\right)\cos 2\theta\left(-e^{-\theta}\,d\theta\right) = -\frac{1}{2}e^{-\theta}\cos 2\theta - \frac{1}{2}\int e^{-\theta}\cos 2\theta\,d\theta$.

So $I = \frac{1}{2}e^{-\theta}\sin 2\theta + \frac{1}{2}\left[\left(-\frac{1}{2}e^{-\theta}\cos 2\theta\right) - \frac{1}{2}I\right] = \frac{1}{2}e^{-\theta}\sin 2\theta - \frac{1}{4}e^{-\theta}\cos 2\theta - \frac{1}{4}I \;\Rightarrow$

$\frac{5}{4}I = \frac{1}{2}e^{-\theta}\sin 2\theta - \frac{1}{4}e^{-\theta}\cos 2\theta + C_1 \;\Rightarrow\; I = \frac{4}{5}\left(\frac{1}{2}e^{-\theta}\sin 2\theta - \frac{1}{4}e^{-\theta}\cos 2\theta + C_1\right) = \frac{2}{5}e^{-\theta}\sin 2\theta - \frac{1}{5}e^{-\theta}\cos 2\theta + C$.

25. First let $u = z^3$, $dv = e^z dz \;\Rightarrow\; du = 3z^2 dz$, $v = e^z$. Then $I_1 = \int z^3 e^z dz = z^3 e^z - 3\int z^2 e^z dz$. Next let $u_1 = z^2$,

$dv_1 = e^z dz \;\Rightarrow\; du_1 = 2z\,dz$, $v_1 = e^z$. Then $I_2 = z^2 e^z - 2\int z e^z dz$. Finally, let $u_2 = z$, $dv_2 = e^z dz \;\Rightarrow\; du_2 = dz$,

$v_2 = e^z$. Then $\int z e^z dz = z e^z - \int e^z dz = z e^z - e^z + C_1$. Substituting in the expression for I_2, we get

$I_2 = z^2 e^z - 2(z e^z - e^z + C_1) = z^2 e^z - 2z e^z + 2e^z - 2C_1$. Substituting the last expression for I_2 into I_1 gives

$I_1 = z^3 e^z - 3(z^2 e^z - 2z e^z + 2e^z - 2C_1) = z^3 e^z - 3z^2 e^z + 6z e^z - 6e^z + C$, where $C = 6C_1$.

26. First let $u = (\arcsin x)^2$, $dv = dx \;\Rightarrow\; du = 2\arcsin x \cdot \dfrac{1}{\sqrt{1-x^2}}\,dx$, $v = x$. Then

$I = \displaystyle\int (\arcsin x)^2 dx = x(\arcsin x)^2 - 2\int \frac{x\arcsin x}{\sqrt{1-x^2}}\,dx$. To simplify the last integral, let $t = \arcsin x$ $[x = \sin t]$, so

$dt = \dfrac{1}{\sqrt{1-x^2}}\,dx$, and $\displaystyle\int \frac{x\arcsin x}{\sqrt{1-x^2}}\,dx = \int t\sin t\,dt$. To evaluate just the last integral, now let $U = t$, $dV = \sin t\,dt \;\Rightarrow$

$dU = dt$, $V = -\cos t$. Thus,

$$\int t\sin t\,dt = -t\cos t + \int \cos t\,dt = -t\cos t + \sin t + C$$

$$= -\arcsin x \cdot \frac{\sqrt{1-x^2}}{1} + x + C_1 \quad \text{[refer to the figure]}$$

Returning to I, we get $I = x(\arcsin x)^2 + 2\sqrt{1-x^2}\,\arcsin x - 2x + C$,

where $C = -2C_1$.

27. First let $u = 1 + x^2$, $dv = e^{3x}\,dx \;\Rightarrow\; du = 2x\,dx$, $v = \frac{1}{3}e^{3x}$. Then

$I = \int (1+x^2)e^{3x}\,dx = \frac{1}{3}e^{3x}(1+x^2) - \frac{2}{3}\int x e^{3x}\,dx$. Next, let $U = x$, $dV = e^{3x}\,dx \;\Rightarrow\; dU = dx$, $V = \frac{1}{3}e^{3x}$, so

$\int x e^{3x}\,dx = \frac{1}{3}x e^{3x} - \frac{1}{3}\int e^{3x}\,dx = \frac{1}{3}x e^{3x} - \frac{1}{9}e^{3x} + C_1$. Substituting in the previous formula gives

$$I = \tfrac{1}{3}e^{3x}(1+x^2) - \tfrac{2}{3}\left(\tfrac{1}{3}x e^{3x} - \tfrac{1}{9}e^{3x} + C_1\right) = \tfrac{1}{3}e^{3x} + \tfrac{1}{3}x^2 e^{3x} - \tfrac{2}{9}x e^{3x} + \tfrac{2}{27}e^{3x} - \tfrac{2}{3}C_1$$

$$= \tfrac{11}{27}e^{3x} - \tfrac{2}{9}x e^{3x} + \tfrac{1}{3}x^2 e^{3x} + C, \text{ where } C = -\tfrac{2}{3}C_1$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. Let $u = \theta$, $dv = \sin 3\pi\theta\, d\theta \quad \Rightarrow \quad du = d\theta$, $v = -\dfrac{1}{3\pi}\cos 3\pi\theta$. By (6),

$$\int_0^{1/2} \theta \sin 3\pi\theta\, d\theta = \left[-\frac{1}{3\pi}\theta\cos 3\pi\theta\right]_0^{1/2} + \frac{1}{3\pi}\int_0^{1/2} \cos 3\pi\theta\, d\theta = (0+0) + \frac{1}{9\pi^2}\left[\sin 3\pi\theta\right]_0^{1/2}$$
$$= \frac{1}{9\pi^2}(-1-0) = -\frac{1}{9\pi^2}$$

29. Let $u = x$, $dv = 3^x\, dx \quad \Rightarrow \quad du = dx$, $v = \dfrac{1}{\ln 3}3^x$. By (6),

$$\int_0^1 x3^x\, dx = \left[\frac{1}{\ln 3}\, x\, 3^x\right]_0^1 - \frac{1}{\ln 3}\int_0^1 3^x\, dx = \left(\frac{3}{\ln 3} - 0\right) - \frac{1}{\ln 3}\left[\frac{1}{\ln 3}\, 3^x\right]_0^1 = \frac{3}{\ln 3} - \frac{1}{(\ln 3)^2}(3-1)$$
$$= \frac{3}{\ln 3} - \frac{2}{(\ln 3)^2}$$

30. Let $u = xe^x$, $dv = \dfrac{1}{(1+x)^2}\, dx \quad \Rightarrow \quad du = (xe^x + e^x)\, dx = e^x(x+1)\, dx$, $v = -\dfrac{1}{1+x}$. By (6),

$$\int_0^1 \frac{xe^x}{(1+x)^2}\, dx = \left[-\frac{xe^x}{1+x}\right]_0^1 - \int_0^1 \left(-\frac{1}{1+x}\right) e^x(1+x)\, dx = \left(-\frac{e}{2} + 0\right) + \int_0^1 e^x\, dx = -\frac{1}{2}e + \left[e^x\right]_0^1$$
$$= -\tfrac{1}{2}e + e - 1 = \tfrac{1}{2}e - 1$$

31. Let $u = y$, $dv = \sinh y\, dy \quad \Rightarrow \quad du = dy$, $v = \cosh y$. By (6),

$$\int_0^2 y\sinh y\, dy = \Big[y\cosh y\Big]_0^2 - \int_0^2 \cosh y\, dy = 2\cosh 2 - 0 - \Big[\sinh y\Big]_0^2 = 2\cosh 2 - \sinh 2.$$

32. Let $u = \ln w$, $dv = w^2\, dw \quad \Rightarrow \quad du = \dfrac{1}{w}\, dw$, $v = \frac{1}{3}w^3$. By (6),

$$\textstyle\int_1^2 w^2 \ln w\, dw = \left[\frac{1}{3}w^3 \ln w\right]_1^2 - \int_1^2 \frac{1}{3}w^2\, dw = \frac{8}{3}\ln 2 - 0 - \left[\frac{1}{9}w^3\right]_1^2 = \frac{8}{3}\ln 2 - \left(\frac{8}{9} - \frac{1}{9}\right) = \frac{8}{3}\ln 2 - \frac{7}{9}.$$

33. Let $u = \ln R$, $dv = \dfrac{1}{R^2}\, dR \quad \Rightarrow \quad du = \dfrac{1}{R}\, dR$, $v = -\dfrac{1}{R}$. By (6),

$$\int_1^5 \frac{\ln R}{R^2}\, dR = \left[-\frac{1}{R}\ln R\right]_1^5 - \int_1^5 -\frac{1}{R^2}\, dR = -\tfrac{1}{5}\ln 5 - 0 - \left[\frac{1}{R}\right]_1^5 = -\tfrac{1}{5}\ln 5 - \left(\tfrac{1}{5} - 1\right) = \tfrac{4}{5} - \tfrac{1}{5}\ln 5.$$

34. First let $u = t^2$, $dv = \sin 2t\, dt \quad \Rightarrow \quad du = 2t\, dt$, $v = -\frac{1}{2}\cos 2t$. By (6),

$\int_0^{2\pi} t^2 \sin 2t\, dt = \left[-\frac{1}{2}t^2\cos 2t\right]_0^{2\pi} + \int_0^{2\pi} t\cos 2t\, dt = -2\pi^2 + \int_0^{2\pi} t\cos 2t\, dt.$

Next let $U = t$, $dV = \cos 2t\, dt \quad \Rightarrow \quad dU = dt$, $V = \frac{1}{2}\sin 2t$. By (6) again,

$\int_0^{2\pi} t\cos 2t\, dt = \left[\frac{1}{2}t\sin 2t\right]_0^{2\pi} - \int_0^{2\pi} \frac{1}{2}\sin 2t\, dt = 0 - \left[-\frac{1}{4}\cos 2t\right]_0^{2\pi} = \frac{1}{4} - \frac{1}{4} = 0$. Thus, $\int_0^{2\pi} t^2 \sin 2t\, dt = -2\pi^2$.

35. $\sin 2x = 2\sin x\, \cos x$, so $\int_0^{\pi} x\, \sin x\, \cos x\, dx = \frac{1}{2}\int_0^{\pi} x\, \sin 2x\, dx$. Let $u = x$, $dv = \sin 2x\, dx \quad \Rightarrow$

$du = dx$, $v = -\frac{1}{2}\cos 2x$. By (6),

$$\tfrac{1}{2}\textstyle\int_0^{\pi} x\, \sin 2x\, dx = \frac{1}{2}\left[-\frac{1}{2}x\cos 2x\right]_0^{\pi} - \frac{1}{2}\int_0^{\pi} -\frac{1}{2}\cos 2x\, dx = -\frac{1}{4}\pi - 0 + \frac{1}{4}\left[\frac{1}{2}\sin 2x\right]_0^{\pi} = -\frac{\pi}{4}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. Let $u = \arctan(1/x)$, $dv = dx \quad \Rightarrow \quad du = \dfrac{1}{1+(1/x)^2} \cdot \dfrac{-1}{x^2}\,dx = \dfrac{-dx}{x^2+1}$, $v = x$. By (6),

$$\int_1^{\sqrt{3}} \arctan\left(\frac{1}{x}\right)\,dx = \left[x\arctan\left(\frac{1}{x}\right)\right]_1^{\sqrt{3}} + \int_1^{\sqrt{3}} \frac{x\,dx}{x^2+1} = \sqrt{3}\,\frac{\pi}{6} - 1\cdot\frac{\pi}{4} + \frac{1}{2}\Big[\ln(x^2+1)\Big]_1^{\sqrt{3}}$$
$$= \frac{\pi\sqrt{3}}{6} - \frac{\pi}{4} + \frac{1}{2}(\ln 4 - \ln 2) = \frac{\pi\sqrt{3}}{6} - \frac{\pi}{4} + \frac{1}{2}\ln\frac{4}{2} = \frac{\pi\sqrt{3}}{6} - \frac{\pi}{4} + \frac{1}{2}\ln 2$$

37. Let $u = M$, $dv = e^{-M}\,dM \quad \Rightarrow \quad du = dM$, $v = -e^{-M}$. By (6),

$$\int_1^5 \frac{M}{e^M}\,dM = \int_1^5 Me^{-M}\,dM = \Big[-Me^{-M}\Big]_1^5 - \int_1^5 -e^{-M}\,dM = -5e^{-5} + e^{-1} - \Big[e^{-M}\Big]_1^5$$
$$= -5e^{-5} + e^{-1} - (e^{-5} - e^{-1}) = 2e^{-1} - 6e^{-5}$$

38. Let $u = (\ln x)^2$, $dv = x^{-3}\,dx \quad \Rightarrow \quad du = \dfrac{2\ln x}{x}\,dx$, $v = -\frac{1}{2}x^{-2}$. By (6),

$I = \displaystyle\int_1^2 \frac{(\ln x)^2}{x^3}\,dx = \left[-\frac{(\ln x)^2}{2x^2}\right]_1^2 + \int_1^2 \frac{\ln x}{x^3}\,dx$. Now let $U = \ln x$, $dV = x^{-3}\,dx \quad \Rightarrow \quad dU = \dfrac{1}{x}\,dx$, $V = -\frac{1}{2}x^{-2}$.

Then

$$\int_1^2 \frac{\ln x}{x^3}\,dx = \left[-\frac{\ln x}{2x^2}\right]_1^2 + \tfrac{1}{2}\int_1^2 x^{-3}dx = -\tfrac{1}{8}\ln 2 + 0 + \tfrac{1}{2}\left[-\tfrac{1}{2x^2}\right]_1^2 = -\tfrac{1}{8}\ln 2 + \tfrac{1}{2}\left(-\tfrac{1}{8} + \tfrac{1}{2}\right) = \tfrac{3}{16} - \tfrac{1}{8}\ln 2.$$

Thus $I = \left(-\frac{1}{8}(\ln 2)^2 + 0\right) + \left(\frac{3}{16} - \frac{1}{8}\ln 2\right) = -\frac{1}{8}(\ln 2)^2 - \frac{1}{8}\ln 2 + \frac{3}{16}$.

39. Let $u = \ln(\cos x)$, $dv = \sin x\,dx \quad \Rightarrow \quad du = \dfrac{1}{\cos x}(-\sin x)\,dx$, $v = -\cos x$. By (6),

$$\int_0^{\pi/3} \sin x\,\ln(\cos x)\,dx = \Big[-\cos x\,\ln(\cos x)\Big]_0^{\pi/3} - \int_0^{\pi/3} \sin x\,dx = -\tfrac{1}{2}\ln\tfrac{1}{2} - 0 - \Big[-\cos x\Big]_0^{\pi/3}$$
$$= -\tfrac{1}{2}\ln\tfrac{1}{2} + \left(\tfrac{1}{2} - 1\right) = \tfrac{1}{2}\ln 2 - \tfrac{1}{2}$$

40. Let $u = r^2$, $dv = \dfrac{r}{\sqrt{4+r^2}}\,dr \quad \Rightarrow \quad du = 2r\,dr$, $v = \sqrt{4+r^2}$. By (6),

$$\int_0^1 \frac{r^3}{\sqrt{4+r^2}}\,dr = \Big[r^2\sqrt{4+r^2}\Big]_0^1 - 2\int_0^1 r\sqrt{4+r^2}\,dr = \sqrt{5} - \tfrac{2}{3}\Big[(4+r^2)^{3/2}\Big]_0^1$$
$$= \sqrt{5} - \tfrac{2}{3}(5)^{3/2} + \tfrac{2}{3}(8) = \sqrt{5}\left(1 - \tfrac{10}{3}\right) + \tfrac{16}{3} = \tfrac{16}{3} - \tfrac{7}{3}\sqrt{5}$$

41. Let $u = \cos x$, $dv = \sinh x\,dx \quad \Rightarrow \quad du = -\sin x\,dx$, $v = \cosh x$. By (6),

$I = \displaystyle\int_0^{\pi} \cos x\sinh x\,dx = \Big[\cos x\cosh x\Big]_0^{\pi} - \int_0^{\pi} -\sin x\cosh x\,dx = -\cosh\pi - 1 + \int_0^{\pi} \sin x\cosh x\,dx.$

Now let $U = \sin x$, $dV = \cosh x\,dx \quad \Rightarrow \quad dU = \cos x\,dx$, $V = \sinh x$. Then

$$\int_0^{\pi} \sin x\cosh x\,dx = \Big[\sin x\sinh x\Big]_0^{\pi} - \int_0^{\pi} \cos x\sinh x\,dx = (0-0) - \int_0^{\pi} \cos x\sinh x\,dx = -I.$$

Substituting in the previous formula gives $I = -\cosh\pi - 1 - I \quad \Rightarrow \quad 2I = -(\cosh\pi + 1) \quad \Rightarrow \quad I = -\dfrac{\cosh\pi + 1}{2}$.

[We could also write the answer as $I = -\frac{1}{4}(2 + e^{\pi} + e^{-\pi})$.]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. Let $u = \sin(t-s)$, $dv = e^s\,ds \;\Rightarrow\; du = -\cos(t-s)\,ds$, $v = e^s$. Then

$I = \int_0^t e^s \sin(t-s)\,ds = \Big[e^s \sin(t-s)\Big]_0^t + \int_0^t e^s \cos(t-s)\,ds = e^t \sin 0 - e^0 \sin t + I_1$. For I_1, let $U = \cos(t-s)$,

$dV = e^s\,ds \;\Rightarrow\; dU = \sin(t-s)\,ds$, $V = e^s$. So $I_1 = \big[e^s \cos(t-s)\big]_0^t - \int_0^t e^s \sin(t-s)\,ds = e^t \cos 0 - e^0 \cos t - I$.

Thus, $I = -\sin t + e^t - \cos t - I \;\Rightarrow\; 2I = e^t - \cos t - \sin t \;\Rightarrow\; I = \frac{1}{2}(e^t - \cos t - \sin t)$.

43. Let $t = \sqrt{x}$, so that $t^2 = x$ and $2t\,dt = dx$. Thus, $\int e^{\sqrt{x}}\,dx = \int e^t(2t)\,dt$. Now use parts with $u = t$, $dv = e^t\,dt$, $du = dt$, and $v = e^t$ to get $2\int te^t\,dt = 2t\,e^t - 2\int e^t\,dt = 2t\,e^t - 2e^t + C = 2\sqrt{x}\,e^{\sqrt{x}} - 2e^{\sqrt{x}} + C$.

44. Let $t = \ln x$, so that $e^t = x$ and $e^t\,dt = dx$. Thus, $\int \cos(\ln x)\,dx = \int \cos t \cdot e^t\,dt = I$. Now use parts with $u = \cos t$, $dv = e^t\,dt$, $du = -\sin t\,dt$, and $v = e^t$ to get $\int e^t \cos t\,dt = e^t \cos t - \int -e^t \sin t\,dt = e^t \cos t + \int e^t \sin t\,dt$. Now use parts with $U = \sin t$, $dV = e^t\,dt$, $dU = \cos t\,dt$, and $V = e^t$ to get

$\int e^t \sin t\,dt = e^t \sin t - \int e^t \cos t\,dt$. Thus, $I = e^t \cos t + e^t \sin t - I \;\Rightarrow\; 2I = e^t \cos t + e^t \sin t \;\Rightarrow$

$I = \frac{1}{2}e^t \cos t + \frac{1}{2}e^t \sin t + C = \frac{1}{2}x\cos(\ln x) + \frac{1}{2}x\sin(\ln x) + C$.

45. Let $x = \theta^2$, so that $dx = 2\theta\,d\theta$. Thus, $\displaystyle\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos\big(\theta^2\big)\;d\theta = \int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^2 \cos\big(\theta^2\big) \cdot \tfrac{1}{2}(2\theta\,d\theta) = \tfrac{1}{2}\int_{\pi/2}^{\pi} x\cos x\,dx$. Now use parts with $u = x$, $dv = \cos x\,dx$, $du = dx$, $v = \sin x$ to get

$$\begin{aligned}\tfrac{1}{2}\int_{\pi/2}^{\pi} x\cos x\,dx &= \tfrac{1}{2}\left(\big[x\sin x\big]_{\pi/2}^{\pi} - \int_{\pi/2}^{\pi} \sin x\,dx\right) = \tfrac{1}{2}\big[x\sin x + \cos x\big]_{\pi/2}^{\pi} \\ &= \tfrac{1}{2}(\pi\sin\pi + \cos\pi) - \tfrac{1}{2}\big(\tfrac{\pi}{2}\sin\tfrac{\pi}{2} + \cos\tfrac{\pi}{2}\big) = \tfrac{1}{2}(\pi\cdot 0 - 1) - \tfrac{1}{2}\big(\tfrac{\pi}{2}\cdot 1 + 0\big) = -\tfrac{1}{2} - \tfrac{\pi}{4}\end{aligned}$$

46. Let $x = \cos t$, so that $dx = -\sin t\,dt$. Thus,

$\int_0^{\pi} e^{\cos t} \sin 2t\,dt = \int_0^{\pi} e^{\cos t}(2\sin t\,\cos t)\,dt = \int_1^{-1} e^x \cdot 2x\,(-dx) = 2\int_{-1}^{1} xe^x\,dx$. Now use parts with $u = x$, $dv = e^x\,dx$, $du = dx$, $v = e^x$ to get

$2\int_{-1}^{1} xe^x\,dx = 2\left(\big[xe^x\big]_{-1}^{1} - \int_{-1}^{1} e^x\,dx\right) = 2\left(e^1 + e^{-1} - \big[\,e^x\,\big]_{-1}^{1}\right) = 2(e + e^{-1} - [e^1 - e^{-1}]) = 2(2e^{-1}) = 4/e$.

47. Let $y = 1 + x$, so that $dy = dx$. Thus, $\int x\ln(1+x)\,dx = \int (y-1)\ln y\,dy$. Now use parts with $u = \ln y$, $dv = (y-1)\,dy$, $du = \frac{1}{y}\,dy$, $v = \frac{1}{2}y^2 - y$ to get

$$\begin{aligned}\int (y-1)\ln y\,dy &= \big(\tfrac{1}{2}y^2 - y\big)\ln y - \int \big(\tfrac{1}{2}y - 1\big)\,dy = \tfrac{1}{2}y(y-2)\ln y - \tfrac{1}{4}y^2 + y + C \\ &= \tfrac{1}{2}(1+x)(x-1)\ln(1+x) - \tfrac{1}{4}(1+x)^2 + 1 + x + C,\end{aligned}$$

which can be written as $\frac{1}{2}(x^2-1)\ln(1+x) - \frac{1}{4}x^2 + \frac{1}{2}x + \frac{3}{4} + C$.

48. Let $y = \ln x$, so that $dy = \dfrac{1}{x}\,dx$. Thus, $\displaystyle\int \frac{\arcsin(\ln x)}{x}\,dx = \int \arcsin y\,dy$. Now use parts with $u = \arcsin y$, $dv = dy$, $du = \dfrac{1}{\sqrt{1-y^2}}\,dy$, and $v = y$ to get

$$\int \arcsin y\,dy = y\arcsin y - \int \frac{y}{\sqrt{1-y^2}}\,dy = y\arcsin y + \sqrt{1-y^2} + C = (\ln x)\arcsin(\ln x) + \sqrt{1-(\ln x)^2} + C.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. Let $u = x$, $dv = e^{-2x}\,dx \quad\Rightarrow\quad du = dx$, $v = -\frac{1}{2}e^{-2x}$. Then

$\int xe^{-2x}\,dx = -\frac{1}{2}xe^{-2x} + \int \frac{1}{2}e^{-2x}\,dx = -\frac{1}{2}xe^{-2x} - \frac{1}{4}e^{-2x} + C$. We see from the graph that this is reasonable, since F has a minimum where f changes from negative to positive. Also, F increases where f is positive and F decreases where f is negative.

50. Let $u = \ln x$, $dv = x^{3/2}\,dx \quad\Rightarrow\quad du = \frac{1}{x}\,dx$, $v = \frac{2}{5}x^{5/2}$. Then

$$\int x^{3/2}\ln x\,dx = \tfrac{2}{5}x^{5/2}\ln x - \tfrac{2}{5}\int x^{3/2}\,dx = \tfrac{2}{5}x^{5/2}\ln x - \left(\tfrac{2}{5}\right)^2 x^{5/2} + C$$
$$= \tfrac{2}{5}x^{5/2}\ln x - \tfrac{4}{25}x^{5/2} + C$$

We see from the graph that this is reasonable, since F has a minimum where f changes from negative to positive.

51. Let $u = \frac{1}{2}x^2$, $dv = 2x\sqrt{1+x^2}\,dx \quad\Rightarrow\quad du = x\,dx$, $v = \frac{2}{3}(1+x^2)^{3/2}$. Then

$$\int x^3\sqrt{1+x^2}\,dx = \tfrac{1}{2}x^2\left[\tfrac{2}{3}(1+x^2)^{3/2}\right] - \tfrac{2}{3}\int x(1+x^2)^{3/2}dx$$
$$= \tfrac{1}{3}x^2(1+x^2)^{3/2} - \tfrac{2}{3}\cdot\tfrac{2}{5}\cdot\tfrac{1}{2}(1+x^2)^{5/2} + C$$
$$= \tfrac{1}{3}x^2(1+x^2)^{3/2} - \tfrac{2}{15}(1+x^2)^{5/2} + C$$

We see from the graph that this is reasonable, since F increases where f is positive and F decreases where f is negative. Note also that f is an odd function and F is an even function.

Another method: Use substitution with $u = 1 + x^2$ to get $\frac{1}{5}(1+x^2)^{5/2} - \frac{1}{3}(1+x^2)^{3/2} + C$.

52. First let $u = x^2$, $dv = \sin 2x\,dx \quad\Rightarrow\quad du = 2x\,dx$, $v = -\frac{1}{2}\cos 2x$.

Then $I = \int x^2\sin 2x\,dx = -\frac{1}{2}x^2\cos 2x + \int x\cos 2x\,dx$.

Next let $U = x$, $dV = \cos 2x\,dx \quad\Rightarrow\quad dU = dx$, $V = \frac{1}{2}\sin 2x$, so

$\int x\cos 2x\,dx = \frac{1}{2}x\sin 2x - \int \frac{1}{2}\sin 2x\,dx = \frac{1}{2}x\sin 2x + \frac{1}{4}\cos 2x + C$.

Thus, $I = -\frac{1}{2}x^2\cos 2x + \frac{1}{2}x\sin 2x + \frac{1}{4}\cos 2x + C$.

We see from the graph that this is reasonable, since F increases where f is positive and F decreases where f is negative. Note also that f is an odd function and F is an even function.

53. (a) Take $n = 2$ in Example 6 to get $\displaystyle\int \sin^2 x\,dx = -\frac{1}{2}\cos x\sin x + \frac{1}{2}\int 1\,dx = \frac{x}{2} - \frac{\sin 2x}{4} + C$.

(b) $\int \sin^4 x\,dx = -\frac{1}{4}\cos x\sin^3 x + \frac{3}{4}\int \sin^2 x\,dx = -\frac{1}{4}\cos x\sin^3 x + \frac{3}{8}x - \frac{3}{16}\sin 2x + C$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. (a) Let $u = \cos^{n-1} x$, $dv = \cos x\,dx \quad \Rightarrow \quad du = -(n-1)\cos^{n-2} x\,\sin x\,dx$, $v = \sin x$ in (2):

$$\begin{aligned}\int \cos^n x\,dx &= \cos^{n-1} x\,\sin x + (n-1)\int \cos^{n-2} x\,\sin^2 x\,dx \\ &= \cos^{n-1} x\,\sin x + (n-1)\int \cos^{n-2} x\,(1-\cos^2 x)dx \\ &= \cos^{n-1} x\,\sin x + (n-1)\int \cos^{n-2} x\,dx - (n-1)\int \cos^n x\,dx\end{aligned}$$

Rearranging terms gives $n\int \cos^n x\,dx = \cos^{n-1} x\,\sin x + (n-1)\int \cos^{n-2} x\,dx$ or

$$\int \cos^n x\,dx = \frac{1}{n}\cos^{n-1} x\,\sin x + \frac{n-1}{n}\int \cos^{n-2} x\,dx$$

(b) Take $n = 2$ in part (a) to get $\int \cos^2 x\,dx = \frac{1}{2}\cos x\,\sin x + \frac{1}{2}\int 1\,dx = \dfrac{x}{2} + \dfrac{\sin 2x}{4} + C$.

(c) $\int \cos^4 x\,dx = \frac{1}{4}\cos^3 x\,\sin x + \frac{3}{4}\int \cos^2 x\,dx = \frac{1}{4}\cos^3 x\,\sin x + \frac{3}{8}x + \frac{3}{16}\sin 2x + C$

55. (a) From Example 6, $\displaystyle\int \sin^n x\,dx = -\frac{1}{n}\cos x\,\sin^{n-1} x + \frac{n-1}{n}\int \sin^{n-2} x\,dx$. Using (6),

$$\begin{aligned}\int_0^{\pi/2} \sin^n x\,dx &= \left[-\frac{\cos x\,\sin^{n-1} x}{n}\right]_0^{\pi/2} + \frac{n-1}{n}\int_0^{\pi/2} \sin^{n-2} x\,dx \\ &= (0-0) + \frac{n-1}{n}\int_0^{\pi/2} \sin^{n-2} x\,dx = \frac{n-1}{n}\int_0^{\pi/2} \sin^{n-2} x\,dx\end{aligned}$$

(b) Using $n = 3$ in part (a), we have $\int_0^{\pi/2} \sin^3 x\,dx = \frac{2}{3}\int_0^{\pi/2} \sin x\,dx = \left[-\frac{2}{3}\cos x\right]_0^{\pi/2} = \frac{2}{3}$.

Using $n = 5$ in part (a), we have $\int_0^{\pi/2} \sin^5 x\,dx = \frac{4}{5}\int_0^{\pi/2} \sin^3 x\,dx = \frac{4}{5}\cdot\frac{2}{3} = \frac{8}{15}$.

(c) The formula holds for $n = 1$ (that is, $2n+1 = 3$) by (b). Assume it holds for some $k \geq 1$. Then

$\displaystyle\int_0^{\pi/2} \sin^{2k+1} x\,dx = \frac{2\cdot 4\cdot 6\cdot\cdots\cdot(2k)}{3\cdot 5\cdot 7\cdot\cdots\cdot(2k+1)}$. By Example 6,

$$\begin{aligned}\int_0^{\pi/2} \sin^{2k+3} x\,dx &= \frac{2k+2}{2k+3}\int_0^{\pi/2} \sin^{2k+1} x\,dx = \frac{2k+2}{2k+3}\cdot\frac{2\cdot 4\cdot 6\cdot\cdots\cdot(2k)}{3\cdot 5\cdot 7\cdot\cdots\cdot(2k+1)} \\ &= \frac{2\cdot 4\cdot 6\cdot\cdots\cdot(2k)[2(k+1)]}{3\cdot 5\cdot 7\cdot\cdots\cdot(2k+1)[2(k+1)+1]},\end{aligned}$$

so the formula holds for $n = k+1$. By induction, the formula holds for all $n \geq 1$.

56. Using Exercise 53(a), we see that the formula holds for $n = 1$, because $\int_0^{\pi/2} \sin^2 x\,dx = \frac{1}{2}\int_0^{\pi/2} 1\,dx = \frac{1}{2}\left[x\right]_0^{\pi/2} = \frac{1}{2}\cdot\frac{\pi}{2}$.

Now assume it holds for some $k \geq 1$. Then $\displaystyle\int_0^{\pi/2} \sin^{2k} x\,dx = \frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2k-1)}{2\cdot 4\cdot 6\cdot\cdots\cdot(2k)}\frac{\pi}{2}$. By Exercise 53(a),

$$\begin{aligned}\int_0^{\pi/2} \sin^{2(k+1)} x\,dx &= \frac{2k+1}{2k+2}\int_0^{\pi/2} \sin^{2k} x\,dx = \frac{2k+1}{2k+2}\cdot\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2k-1)}{2\cdot 4\cdot 6\cdot\cdots\cdot(2k)}\frac{\pi}{2} \\ &= \frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2k-1)(2k+1)}{2\cdot 4\cdot 6\cdot\cdots\cdot(2k)(2k+2)}\cdot\frac{\pi}{2},\end{aligned}$$

so the formula holds for $n = k+1$. By induction, the formula holds for all $n \geq 1$.

57. Let $u = (\ln x)^n$, $dv = dx \quad \Rightarrow \quad du = n(\ln x)^{n-1}(dx/x)$, $v = x$. By Equation 2,

$\int (\ln x)^n\,dx = x(\ln x)^n - \int nx(\ln x)^{n-1}(dx/x) = x(\ln x)^n - n\int (\ln x)^{n-1}\,dx$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

58. Let $u = x^n$, $dv = e^x\,dx \quad\Rightarrow\quad du = nx^{n-1}\,dx$, $v = e^x$. By Equation 2, $\int x^n e^x\,dx = x^n e^x - n\int x^{n-1}e^x\,dx$.

59. $\int \tan^n x\,dx = \int \tan^{n-2} x\,\tan^2 x\,dx = \int \tan^{n-2} x\,(\sec^2 x - 1)\,dx = \int \tan^{n-2} x\,\sec^2 x\,dx - \int \tan^{n-2} x\,dx$

$= I - \int \tan^{n-2} x\,dx.$

Let $u = \tan^{n-2} x$, $dv = \sec^2 x\,dx \quad\Rightarrow\quad du = (n-2)\,\tan^{n-3} x\,\sec^2 x\,dx$, $v = \tan x$. Then, by Equation 2,

$$\begin{aligned} I &= \tan^{n-1} x - (n-2)\int \tan^{n-2} x\,\sec^2 x\,dx \\ 1I &= \tan^{n-1} x - (n-2)I \\ (n-1)I &= \tan^{n-1} x \\ I &= \frac{\tan^{n-1} x}{n-1} \end{aligned}$$

Returning to the original integral, $\int \tan^n x\,dx = \dfrac{\tan^{n-1} x}{n-1} - \int \tan^{n-2} x\,dx$.

60. Let $u = \sec^{n-2} x$, $dv = \sec^2 x\,dx \quad\Rightarrow\quad du = (n-2)\,\sec^{n-3} x\,\sec x\,\tan x\,dx$, $v = \tan x$. Then, by Equation 2,

$$\begin{aligned} \int \sec^n x\,dx &= \tan x\,\sec^{n-2} x - (n-2)\int \sec^{n-2} x\,\tan^2 x\,dx \\ &= \tan x\,\sec^{n-2} x - (n-2)\int \sec^{n-2} x\,(\sec^2 x - 1)\,dx \\ &= \tan x\,\sec^{n-2} x - (n-2)\int \sec^n x\,dx + (n-2)\int \sec^{n-2} x\,dx \end{aligned}$$

so $(n-1)\int \sec^n x\,dx = \tan x\,\sec^{n-2} x + (n-2)\int \sec^{n-2} x\,dx$. If $n - 1 \neq 0$, then

$$\int \sec^n x\,dx = \frac{\tan x\,\sec^{n-2} x}{n-1} + \frac{n-2}{n-1}\int \sec^{n-2} x\,dx.$$

61. By repeated applications of the reduction formula in Exercise 57,

$$\begin{aligned} \int (\ln x)^3\,dx &= x\,(\ln x)^3 - 3\int (\ln x)^2\,dx = x(\ln x)^3 - 3\big[x(\ln x)^2 - 2\int (\ln x)^1\,dx\big] \\ &= x\,(\ln x)^3 - 3x(\ln x)^2 + 6\big[x(\ln x)^1 - 1\int (\ln x)^0\,dx\big] \\ &= x\,(\ln x)^3 - 3x(\ln x)^2 + 6x\ln x - 6\int 1\,dx = x\,(\ln x)^3 - 3x(\ln x)^2 + 6x\ln x - 6x + C \end{aligned}$$

62. By repeated applications of the reduction formula in Exercise 58,

$$\begin{aligned} \int x^4 e^x\,dx &= x^4 e^x - 4\int x^3 e^x\,dx = x^4 e^x - 4\big(x^3 e^x - 3\int x^2 e^x\,dx\big) \\ &= x^4 e^x - 4x^3 e^x + 12\big(x^2 e^x - 2\int x^1 e^x\,dx\big) = x^4 e^x - 4x^3 e^x + 12x^2 e^x - 24\big(x^1 e^x - \int x^0 e^x\,dx\big) \\ &= x^4 e^x - 4x^3 e^x + 12x^2 e^x - 24xe^x + 24e^x + C \quad \big[\text{or } e^x(x^4 - 4x^3 + 12x^2 - 24x + 24) + C\big] \end{aligned}$$

63. The curves $y = x^2 \ln x$ and $y = 4\ln x$ intersect when $x^2 \ln x = 4\ln x \quad\Leftrightarrow$

$x^2 \ln x - 4\ln x = 0 \quad\Leftrightarrow\quad (x^2 - 4)\ln x = 0 \quad\Leftrightarrow$

$x = 1$ or 2 [since $x > 0$]. For $1 < x < 2$, $4\ln x > x^2 \ln x$. Thus,

area $= \int_1^2 (4\ln x - x^2 \ln x)\,dx = \int_1^2 [(4 - x^2)\ln x]\,dx$. Let $u = \ln x$,

$dv = (4 - x^2)\,dx \quad\Rightarrow\quad du = \frac{1}{x}\,dx$, $v = 4x - \frac{1}{3}x^3$. Then

$$\text{area} = \big[(\ln x)\big(4x - \tfrac{1}{3}x^3\big)\big]_1^2 - \int_1^2 \left[\big(4x - \tfrac{1}{3}x^3\big)\frac{1}{x}\right] dx = (\ln 2)\big(\tfrac{16}{3}\big) - 0 - \int_1^2 \big(4 - \tfrac{1}{3}x^2\big)\,dx$$

$$= \tfrac{16}{3}\ln 2 - \big[4x - \tfrac{1}{9}x^3\big]_1^2 = \tfrac{16}{3}\ln 2 - \big(\tfrac{64}{9} - \tfrac{35}{9}\big) = \tfrac{16}{3}\ln 2 - \tfrac{29}{9}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. The curves $y = x^2e^{-x}$ and $y = xe^{-x}$ intersect when $x^2e^{-x} = xe^{-x} \quad \Leftrightarrow$

$x^2 - x = 0 \quad \Leftrightarrow \quad x(x-1) = 0 \quad \Leftrightarrow \quad x = 0$ or 1.

For $0 < x < 1$, $xe^{-x} > x^2e^{-x}$. Thus,

area $= \int_0^1 (xe^{-x} - x^2e^{-x})\,dx = \int_0^1 (x - x^2)e^{-x}\,dx$. Let $u = x - x^2$,

$dv = e^{-x}dx \quad \Rightarrow \quad du = (1-2x)\,dx$, $v = -e^{-x}$. Then

area $= \left[(x-x^2)(-e^{-x})\right]_0^1 - \int_0^1 [-e^{-x}(1-2x)]\,dx = 0 + \int_0^1 (1-2x)e^{-x}\,dx$.

Now let $U = 1 - 2x$, $dV = e^{-x}dx \quad \Rightarrow \quad dU = -2\,dx$, $V = -e^{-x}$. Now

area $= \left[(1-2x)(-e^{-x})\right]_0^1 - \int_0^1 2e^{-x}\,dx = e^{-1} + 1 - \left[-2e^{-x}\right]_0^1 = e^{-1} + 1 + 2(e^{-1} - 1) = 3e^{-1} - 1$.

65. The curves $y = \arcsin\left(\frac{1}{2}x\right)$ and $y = 2 - x^2$ intersect at $x = a \approx -1.75119$ and $x = b \approx 1.17210$. From the figure, the area bounded by the curves is given by

$A = \int_a^b [(2 - x^2) - \arcsin\left(\frac{1}{2}x\right)]\,dx = \left[2x - \frac{1}{3}x^3\right]_a^b - \int_a^b \arcsin\left(\frac{1}{2}x\right)dx$.

Let $u = \arcsin\left(\frac{1}{2}x\right)$, $dv = dx \quad \Rightarrow \quad du = \dfrac{1}{\sqrt{1 - \left(\frac{1}{2}x\right)^2}} \cdot \dfrac{1}{2}\,dx$, $v = x$.

Then

$$A = \left[2x - \frac{1}{3}x^3\right]_a^b - \left\{\left[x\arcsin\left(\frac{1}{2}x\right)\right]_a^b - \int_a^b \frac{x}{2\sqrt{1 - \frac{1}{4}x^2}}\,dx\right\}$$

$$= \left[2x - \tfrac{1}{3}x^3 - x\arcsin\left(\tfrac{1}{2}x\right) - 2\sqrt{1 - \tfrac{1}{4}x^2}\,\right]_a^b \approx 3.99926$$

66. The curves $y = x\ln(x+1)$ and $y = 3x - x^2$ intersect at $x = 0$ and $x = a \approx 1.92627$. From the figure, the area bounded by the curves is given by

$A = \int_0^a [(3x - x^2) - x\ln(x+1)]\,dx = \left[\frac{3}{2}x^2 - \frac{1}{3}x^3\right]_0^a - \int_0^a x\ln(x+1)\,dx$.

Let $u = \ln(x+1)$, $dv = x\,dx \quad \Rightarrow \quad du = \dfrac{1}{x+1}\,dx$, $v = \frac{1}{2}x^2$. Then

$$A = \left[\frac{3}{2}x^2 - \frac{1}{3}x^3\right]_0^a - \left\{\left[\frac{1}{2}x^2\ln(x+1)\right]_0^a - \frac{1}{2}\int_0^a \frac{x^2}{x+1}\,dx\right\}$$

$$= \left[\frac{3}{2}x^2 - \frac{1}{3}x^3\right]_0^a - \left[\frac{1}{2}x^2\ln(x+1)\right]_0^a + \frac{1}{2}\int_0^a \left(x - 1 + \frac{1}{x+1}\right)dx$$

$$= \left[\tfrac{3}{2}x^2 - \tfrac{1}{3}x^3 - \tfrac{1}{2}x^2\ln(x+1) + \tfrac{1}{4}x^2 - \tfrac{1}{2}x + \tfrac{1}{2}\ln|x+1|\right]_0^a \approx 1.69260$$

67. Volume $= \int_0^1 2\pi x\cos(\pi x/2)\,dx$. Let $u = x$, $dv = \cos(\pi x/2)\,dx \quad \Rightarrow \quad du = dx$, $v = \frac{2}{\pi}\sin(\pi x/2)$.

$$V = 2\pi\left[\frac{2}{\pi}x\sin\left(\frac{\pi x}{2}\right)\right]_0^1 - 2\pi\cdot\frac{2}{\pi}\int_0^1 \sin\left(\frac{\pi x}{2}\right)dx = 2\pi\left(\frac{2}{\pi} - 0\right) - 4\left[-\frac{2}{\pi}\cos\left(\frac{\pi x}{2}\right)\right]_0^1 = 4 + \frac{8}{\pi}(0-1) = 4 - \frac{8}{\pi}.$$

68. Volume $= \int_0^1 2\pi x(e^x - e^{-x})\,dx = 2\pi\int_0^1 (xe^x - xe^{-x})\,dx = 2\pi\left[\int_0^1 xe^x\,dx - \int_0^1 xe^{-x}\,dx\right]$ [both integrals by parts]

$$= 2\pi\left[(xe^x - e^x) - \left(-xe^{-x} - e^{-x}\right)\right]_0^1 = 2\pi[2/e - 0] = 4\pi/e$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

69. Volume $= \int_{-1}^{0} 2\pi(1-x)e^{-x}\,dx$. Let $u = 1-x$, $dv = e^{-x}\,dx \;\Rightarrow\; du = -\,dx$, $v = -e^{-x}$.

$$V = 2\pi\left[(1-x)(-e^{-x})\right]_{-1}^{0} - 2\pi\int_{-1}^{0} e^{-x}\,dx = 2\pi\left[(x-1)(e^{-x}) + e^{-x}\right]_{-1}^{0} = 2\pi\left[xe^{-x}\right]_{-1}^{0} = 2\pi(0+e) = 2\pi e.$$

70. $y = e^x \;\Leftrightarrow\; x = \ln y$. Volume $= \int_1^3 2\pi y\,\ln y\,dy$. Let $u = \ln y$, $dv = y\,dy \;\Rightarrow\; du = \dfrac{1}{y}\,dy$, $v = \frac{1}{2}y^2$.

$$V = 2\pi\left[\tfrac{1}{2}y^2\ln y\right]_1^3 - 2\pi\int_1^3 \tfrac{1}{2}y\,dy = 2\pi\left[\tfrac{1}{2}y^2\ln y - \tfrac{1}{4}y^2\right]_1^3$$
$$= 2\pi\left[\left(\tfrac{9}{2}\ln 3 - \tfrac{9}{4}\right) - \left(0 - \tfrac{1}{4}\right)\right] = 2\pi\left(\tfrac{9}{2}\ln 3 - 2\right) = (9\ln 3 - 4)\,\pi$$

71. (a) Use shells about the y-axis:

$$V = \int_1^2 2\pi x\ln x\,dx \qquad \begin{bmatrix} u = \ln x, & dv = x\,dx \\ du = \frac{1}{x}\,dx, & v = \frac{1}{2}x^2 \end{bmatrix}$$
$$= 2\pi\left\{\left[\tfrac{1}{2}x^2\ln x\right]_1^2 - \int_1^2 \tfrac{1}{2}x\,dx\right\} = 2\pi\left\{(2\ln 2 - 0) - \left[\tfrac{1}{4}x^2\right]_1^2\right\} = 2\pi\left(2\ln 2 - \tfrac{3}{4}\right)$$

(b) Use disks about the x-axis:

$$V = \int_1^2 \pi(\ln x)^2\,dx \qquad \begin{bmatrix} u = (\ln x)^2, & dv = dx \\ du = 2\ln x\cdot\frac{1}{x}\,dx, & v = x \end{bmatrix}$$
$$= \pi\left\{\left[x(\ln x)^2\right]_1^2 - \int_1^2 2\ln x\,dx\right\} \qquad \begin{bmatrix} u = \ln x, & dv = dx \\ du = \frac{1}{x}\,dx, & v = x \end{bmatrix}$$
$$= \pi\left\{2(\ln 2)^2 - 2\left(\Big[x\ln x\Big]_1^2 - \int_1^2 dx\right)\right\} = \pi\left\{2(\ln 2)^2 - 4\ln 2 + 2\Big[x\Big]_1^2\right\}$$
$$= \pi[2(\ln 2)^2 - 4\ln 2 + 2] = 2\pi[(\ln 2)^2 - 2\ln 2 + 1]$$

72. $$f_{\text{ave}} = \frac{1}{b-a}\int_a^b f(x)\,dx = \frac{1}{\pi/4 - 0}\int_0^{\pi/4} x\sec^2 x\,dx \qquad \begin{bmatrix} u = x, & dv = \sec^2 x\,dx \\ du = dx, & v = \tan x \end{bmatrix}$$
$$= \frac{4}{\pi}\left\{\Big[x\tan x\Big]_0^{\pi/4} - \int_0^{\pi/4}\tan x\,dx\right\} = \frac{4}{\pi}\left\{\frac{\pi}{4} - \Big[\ln|\sec x|\Big]_0^{\pi/4}\right\} = \frac{4}{\pi}\left(\frac{\pi}{4} - \ln\sqrt{2}\right)$$
$$= 1 - \tfrac{4}{\pi}\ln\sqrt{2} \;\text{ or }\; 1 - \tfrac{2}{\pi}\ln 2$$

73. $S(x) = \displaystyle\int_0^x \sin\left(\tfrac{1}{2}\pi t^2\right)dt \;\Rightarrow\; \int S(x)\,dx = \int\left[\int_0^x \sin\left(\tfrac{1}{2}\pi t^2\right)dt\right]dx.$

Let $u = \displaystyle\int_0^x \sin\left(\tfrac{1}{2}\pi t^2\right)dt = S(x)$, $dv = dx \;\Rightarrow\; du = \sin\left(\tfrac{1}{2}\pi x^2\right)dx$, $v = x$. Thus,

$$\int S(x)\,dx = xS(x) - \int x\sin\left(\tfrac{1}{2}\pi x^2\right)dx = xS(x) - \int \sin y\left(\tfrac{1}{\pi}\,dy\right) \qquad \begin{bmatrix} u = \frac{1}{2}\pi x^2, \\ du = \pi x\,dx \end{bmatrix}$$
$$= xS(x) + \tfrac{1}{\pi}\cos y + C = xS(x) + \tfrac{1}{\pi}\cos\left(\tfrac{1}{2}\pi x^2\right) + C$$

74. (a) The rocket will have height $H = \int_0^T v(t)\,dt$ after T seconds.

$$H = \int_0^T\left[-gt - v_e\ln\left(\frac{m-rt}{m}\right)\right]dt = -g\left[\tfrac{1}{2}t^2\right]_0^T - v_e\left[\int_0^T \ln(m-rt)\,dt - \int_0^T \ln m\,dt\right]$$
$$= -\tfrac{1}{2}gT^2 + v_e(\ln m)T - v_e\int_0^T \ln(m-rt)\,dt$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Let $u = \ln(m - rt)$, $dv = dt \quad \Rightarrow \quad du = \dfrac{1}{m - rt}(-r)\,dt$, $v = t$. Then

$$\int_0^T \ln(m - rt)\,dt = \Big[t\ln(m - rt)\Big]_0^T + \int_0^T \frac{rt}{m - rt}\,dt = T\ln(m - rT) + \int_0^T \left(-1 + \frac{m}{m - rt}\right)dt$$

$$= T\ln(m - rT) + \left[-t - \frac{m}{r}\ln(m - rt)\right]_0^T$$

$$= T\ln(m - rT) - T - \frac{m}{r}\ln(m - rT) + \frac{m}{r}\ln m$$

So $H = -\frac{1}{2}gT^2 + v_e(\ln m)T - v_eT\ln(m - rT) + v_eT + \dfrac{m}{r}v_e\ln(m - rT) - \dfrac{m}{r}v_e\ln m$. Substituting $T = 60$, $g = 9.8$, $m = 30{,}000$, $r = 160$, and $v_e = 3000$ gives us $H \approx 14{,}844$ m.

(b) The time taken to consume 6000 kg of fuel is $T = \dfrac{6000}{r} = \dfrac{6000}{160} = 37.5$ s. The rocket will have height $H = \int_0^{37.5} v(t)\,dt$ after 37.5 seconds. Evaluating this integral using the results of part (a) with $T = 37.5$, $g = 9.8$, $m = 30{,}000$, $r = 160$, and $v_e = 3000$ gives us $H \approx 5195$ m.

75. Since $v(t) > 0$ for all t, the desired distance is $s(t) = \int_0^t v(w)\,dw = \int_0^t w^2e^{-w}\,dw$.

First let $u = w^2$, $dv = e^{-w}\,dw \quad \Rightarrow \quad du = 2w\,dw$, $v = -e^{-w}$. Then $s(t) = \left[-w^2e^{-w}\right]_0^t + 2\int_0^t we^{-w}\,dw$.

Next let $U = w$, $dV = e^{-w}\,dw \quad \Rightarrow \quad dU = dw$, $V = -e^{-w}$. Then

$$s(t) = -t^2e^{-t} + 2\left(\left[-we^{-w}\right]_0^t + \int_0^t e^{-w}\,dw\right) = -t^2e^{-t} + 2\left(-te^{-t} + 0 + \left[-e^{-w}\right]_0^t\right)$$

$$= -t^2e^{-t} + 2(-te^{-t} - e^{-t} + 1) = -t^2e^{-t} - 2te^{-t} - 2e^{-t} + 2 = 2 - e^{-t}(t^2 + 2t + 2) \text{ meters}$$

76. Suppose $f(0) = g(0) = 0$ and let $u = f(x)$, $dv = g''(x)\,dx \quad \Rightarrow \quad du = f'(x)\,dx$, $v = g'(x)$.

Then $\int_0^a f(x)\,g''(x)\,dx = \Big[f(x)\,g'(x)\Big]_0^a - \int_0^a f'(x)\,g'(x)\,dx = f(a)\,g'(a) - \int_0^a f'(x)\,g'(x)\,dx$.

Now let $U = f'(x)$, $dV = g'(x)\,dx \quad \Rightarrow \quad dU = f''(x)\,dx$ and $V = g(x)$, so

$\int_0^a f'(x)\,g'(x)\,dx = \Big[f'(x)\,g(x)\Big]_0^a - \int_0^a f''(x)\,g(x)\,dx = f'(a)\,g(a) - \int_0^a f''(x)\,g(x)\,dx$.

Combining the two results, we get $\int_0^a f(x)\,g''(x)\,dx = f(a)\,g'(a) - f'(a)\,g(a) + \int_0^a f''(x)\,g(x)\,dx$.

77. For $I = \int_1^4 xf''(x)\,dx$, let $u = x$, $dv = f''(x)\,dx \quad \Rightarrow \quad du = dx$, $v = f'(x)$. Then

$I = \left[xf'(x)\right]_1^4 - \int_1^4 f'(x)\,dx = 4f'(4) - 1\cdot f'(1) - [f(4) - f(1)] = 4\cdot 3 - 1\cdot 5 - (7 - 2) = 12 - 5 - 5 = 2$.

We used the fact that f'' is continuous to guarantee that I exists.

78. (a) Take $g(x) = x$ and $g'(x) = 1$ in Equation 1.

(b) By part (a), $\int_a^b f(x)\,dx = bf(b) - a\,f(a) - \int_a^b x\,f'(x)\,dx$. Now let $y = f(x)$, so that $x = g(y)$ and $dy = f'(x)\,dx$.

Then $\int_a^b x\,f'(x)\,dx = \int_{f(a)}^{f(b)} g(y)\,dy$. The result follows.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) Part (b) says that the area of region $ABFC$ is

$$\begin{aligned} &= bf(b) - af(a) - \textstyle\int_{f(a)}^{f(b)} g(y)\,dy \\ &= (\text{area of rectangle } OBFE) - (\text{area of rectangle } OACD) - (\text{area of region } DCFE) \end{aligned}$$

(d) We have $f(x) = \ln x$, so $f^{-1}(x) = e^x$, and since $g = f^{-1}$, we have $g(y) = e^y$. By part (b),

$$\int_1^e \ln x\,dx = e\ln e - 1\ln 1 - \int_{\ln 1}^{\ln e} e^y\,dy = e - \int_0^1 e^y\,dy = e - \left[e^y\right]_0^1 = e - (e-1) = 1.$$

79. (a) Assuming $f(x)$ and $g(x)$ are differentiable functions, the Quotient Rule for differentiation states

$\dfrac{d}{dx}\left[\dfrac{f(x)}{g(x)}\right] = \dfrac{g(x)\,f'(x) - f(x)\,g'(x)}{[g(x)]^2}$. Writing in integral form gives

$\dfrac{f(x)}{g(x)} = \displaystyle\int \frac{g(x)\,f'(x) - f(x)\,g'(x)}{[g(x)]^2}\,dx = \int \frac{1}{g(x)}\,f'(x)\,dx - \int \frac{f(x)}{[g(x)]^2}\,g'(x)\,dx$. Now let $u = f(x)$ and $v = g(x)$ so

that $du = f'(x)\,dx$ and $dv = g'(x)\,dx$. Substituting into the above equation gives $\dfrac{u}{v} = \displaystyle\int \frac{1}{v}\,du - \int \frac{u}{v^2}\,dv \quad \Rightarrow$

$\displaystyle\int \frac{u}{v^2}\,dv = -\frac{u}{v} + \int \frac{1}{v}\,du.$

(b) Let $u = \ln x$, $v = x \quad \Rightarrow \quad du = \dfrac{1}{x}\,dx$. Then, using the formula from part (a), we get

$$\int \frac{\ln x}{x^2}\,dx = -\frac{\ln x}{x} + \int \frac{1}{x}\left(\frac{1}{x}\,dx\right) = -\frac{\ln x}{x} + \int \frac{1}{x^2}\,dx = -\frac{\ln x}{x} - \frac{1}{x} + C.$$

80. (a) We note that for $0 \le x \le \frac{\pi}{2}$, $0 \le \sin x \le 1$, so $\sin^{2n+2} x \le \sin^{2n+1} x \le \sin^{2n} x$. So by the second Comparison Property of the Integral, $I_{2n+2} \le I_{2n+1} \le I_{2n}$.

(b) Substituting directly into the result from Exercise 56, we get

$$\frac{I_{2n+2}}{I_{2n}} = \frac{\dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot[2(n+1)-1]}{2\cdot 4\cdot 6\cdot\cdots\cdot[2(n+1)]}\dfrac{\pi}{2}}{\dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)}\dfrac{\pi}{2}} = \frac{2(n+1)-1}{2(n+1)} = \frac{2n+1}{2n+2}$$

(c) We divide the result from part (a) by I_{2n}. The inequalities are preserved since I_{2n} is positive: $\dfrac{I_{2n+2}}{I_{2n}} \le \dfrac{I_{2n+1}}{I_{2n}} \le \dfrac{I_{2n}}{I_{2n}}$.

Now from part (b), the left term is equal to $\dfrac{2n+1}{2n+2}$, so the expression becomes $\dfrac{2n+1}{2n+2} \le \dfrac{I_{2n+1}}{I_{2n}} \le 1$. Now

$\displaystyle\lim_{n\to\infty} \frac{2n+1}{2n+2} = \lim_{n\to\infty} 1 = 1$, so by the Squeeze Theorem, $\displaystyle\lim_{n\to\infty} \frac{I_{2n+1}}{I_{2n}} = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) We substitute the results from Exercises 55 and 56 into the result from part (c):

$$1 = \lim_{n\to\infty} \frac{I_{2n+1}}{I_{2n}} = \lim_{n\to\infty} \frac{\dfrac{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)}{3\cdot 5\cdot 7\cdot\cdots\cdot(2n+1)}}{\dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)}\dfrac{\pi}{2}} = \lim_{n\to\infty}\left[\frac{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)}{3\cdot 5\cdot 7\cdot\cdots\cdot(2n+1)}\right]\left[\frac{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)}{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}\left(\frac{2}{\pi}\right)\right]$$

$$= \lim_{n\to\infty} \frac{2}{1}\cdot\frac{2}{3}\cdot\frac{4}{3}\cdot\frac{4}{5}\cdot\frac{6}{5}\cdot\frac{6}{7}\cdot\cdots\cdot\frac{2n}{2n-1}\cdot\frac{2n}{2n+1}\cdot\frac{2}{\pi} \qquad \text{[rearrange terms]}$$

Multiplying both sides by $\frac{\pi}{2}$ gives us the *Wallis product*:

$$\frac{\pi}{2} = \frac{2}{1}\cdot\frac{2}{3}\cdot\frac{4}{3}\cdot\frac{4}{5}\cdot\frac{6}{5}\cdot\frac{6}{7}\cdot\cdots$$

(e) The area of the kth rectangle is k. At the $2n$th step, the area is increased from $2n-1$ to $2n$ by multiplying the width by $\dfrac{2n}{2n-1}$, and at the $(2n+1)$th step, the area is increased from $2n$ to $2n+1$ by multiplying the height by $\dfrac{2n+1}{2n}$. These two steps multiply the ratio of width to height by $\dfrac{2n}{2n-1}$ and $\dfrac{1}{(2n+1)/(2n)} = \dfrac{2n}{2n+1}$ respectively. So, by part (d), the limiting ratio is $\dfrac{2}{1}\cdot\dfrac{2}{3}\cdot\dfrac{4}{3}\cdot\dfrac{4}{5}\cdot\dfrac{6}{5}\cdot\dfrac{6}{7}\cdot\cdots = \dfrac{\pi}{2}$.

81. Using the formula for volumes of rotation and the figure, we see that

Volume $= \int_0^d \pi b^2\,dy - \int_0^c \pi a^2\,dy - \int_c^d \pi[g(y)]^2\,dy = \pi b^2 d - \pi a^2 c - \int_c^d \pi[g(y)]^2\,dy$. Let $y = f(x)$,

which gives $dy = f'(x)\,dx$ and $g(y) = x$, so that $V = \pi b^2 d - \pi a^2 c - \pi\int_a^b x^2 f'(x)\,dx$.

Now integrate by parts with $u = x^2$, and $dv = f'(x)\,dx \;\Rightarrow\; du = 2x\,dx$, $v = f(x)$, and

$\int_a^b x^2 f'(x)\,dx = \left[x^2 f(x)\right]_a^b - \int_a^b 2x\,f(x)\,dx = b^2 f(b) - a^2 f(a) - \int_a^b 2x\,f(x)\,dx$, but $f(a) = c$ and $f(b) = d \;\Rightarrow$

$V = \pi b^2 d - \pi a^2 c - \pi\left[b^2 d - a^2 c - \int_a^b 2x\,f(x)\,dx\right] = \int_a^b 2\pi x\,f(x)\,dx$.

7.2 Trigonometric Integrals

The symbols $\overset{s}{=}$ and $\overset{c}{=}$ indicate the use of the substitutions $\{u = \sin x, du = \cos x\,dx\}$ and $\{u = \cos x, du = -\sin x\,dx\}$, respectively.

1. $\int \sin^3 x\cos^2 x\,dx = \int \sin^2 x\cos^2 x\sin x\,dx = \int (1-\cos^2 x)\cos^2 x\sin x\,dx \overset{c}{=} \int (1-u^2)u^2\,(-du)$

$$= \int (u^4 - u^2)\,du = \tfrac{1}{5}u^5 - \tfrac{1}{3}u^3 + C = \tfrac{1}{5}\cos^5 x - \tfrac{1}{3}\cos^3 x + C$$

2. $\int \cos^6 y\sin^3 y\,dy = \int \cos^6 y\sin^2 y\sin y\,dy = \int \cos^6 y(1-\cos^2 y)\sin y\,dy \overset{c}{=} \int u^6(1-u^2)\,(-du)$

$$= \int (u^8 - u^6)\,du = \tfrac{1}{9}u^9 - \tfrac{1}{7}u^7 + C = \tfrac{1}{9}\cos^9 y - \tfrac{1}{7}\cos^7 y + C$$

3. $\displaystyle\int_0^{\pi/2} \cos^9 x\sin^5 x\,dx = \int_0^{\pi/2} \cos^9 x\,(\sin^2 x)^2\sin x\,dx = \int_0^{\pi/2} \cos^9 x\,(1-\cos^2 x)^2\sin x\,dx$

$$\overset{c}{=} \int_1^0 u^9(1-u^2)^2\,(-du) = \int_0^1 u^9(1-2u^2+u^4)\,du = \int_0^1 (u^9 - 2u^{11} + u^{13})\,du$$

$$= \left[\tfrac{1}{10}u^{10} - \tfrac{1}{6}u^{12} + \tfrac{1}{14}u^{14}\right]_0^1 = \left(\tfrac{1}{10} - \tfrac{1}{6} + \tfrac{1}{14}\right) - 0 = \tfrac{1}{210}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $$\int_0^{\pi/4} \sin^5 x\,dx = \int_0^{\pi/4} (\sin^2 x)^2 \sin x\,dx = \int_0^{\pi/4} (1-\cos^2 x)^2 \sin x\,dx \overset{c}{=} \int_1^{1/\sqrt{2}} (1-u^2)^2\,(-du)$$
$$= \int_{1/\sqrt{2}}^{1} (1-2u^2+u^4)\,du = \left[u - \frac{2}{3}u^3 + \frac{1}{5}u^5\right]_{1/\sqrt{2}}^{1} = \left(1 - \frac{2}{3} + \frac{1}{5}\right) - \left(\frac{1}{\sqrt{2}} - \frac{2}{3\sqrt{8}} + \frac{1}{5\sqrt{32}}\right)$$
$$= \frac{8}{15} - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{6} + \frac{\sqrt{2}}{40}\right) = \frac{8}{15} - \frac{43\sqrt{2}}{120}$$

5. $$\int \sin^5(2t)\cos^2(2t)\,dt = \int \sin^4(2t)\,\cos^2(2t)\,\sin(2t)\,dt = \int [1-\cos^2(2t)]^2 \cos^2(2t)\,\sin(2t)\,dt$$
$$= \int (1-u^2)^2\,u^2\,\left(-\tfrac{1}{2}\,du\right) \qquad [u = \cos(2t),\, du = -2\sin(2t)\,dt]$$
$$= -\tfrac{1}{2}\int (u^4 - 2u^2 + 1)u^2\,du = -\tfrac{1}{2}\int (u^6 - 2u^4 + u^2)\,du$$
$$= -\tfrac{1}{2}\left(\tfrac{1}{7}u^7 - \tfrac{2}{5}u^5 + \tfrac{1}{3}u^3\right) + C = -\tfrac{1}{14}\cos^7(2t) + \tfrac{1}{5}\cos^5(2t) - \tfrac{1}{6}\cos^3(2t) + C$$

6. $$\int \cos^3\left(\frac{t}{2}\right)\sin^2\left(\frac{t}{2}\right)dt = \int \cos^2\left(\frac{t}{2}\right)\sin^2\left(\frac{t}{2}\right)\cos\left(\frac{t}{2}\right)dt = \int \left(1-\sin^2\left(\frac{t}{2}\right)\right)\sin^2\left(\frac{t}{2}\right)\cos\left(\frac{t}{2}\right)dt$$
$$= \int (1-u^2)u^2(2\,du) \qquad \left[u = \sin\left(\frac{t}{2}\right),\, du = \frac{1}{2}\cos\left(\frac{t}{2}\right)dt\right]$$
$$= 2\int (u^2 - u^4)\,du = 2\left(\frac{1}{3}u^3 - \frac{1}{5}u^5\right) + C = \frac{2}{3}\sin^3\left(\frac{t}{2}\right) - \frac{2}{5}\sin^5\left(\frac{t}{2}\right) + C$$

7. $$\int_0^{\pi/2} \cos^2\theta\,d\theta = \int_0^{\pi/2} \tfrac{1}{2}(1+\cos 2\theta)\,d\theta \qquad \text{[half-angle identity]}$$
$$= \tfrac{1}{2}\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/2} = \tfrac{1}{2}\left[\left(\tfrac{\pi}{2}+0\right) - (0+0)\right] = \tfrac{\pi}{4}$$

8. $$\int_0^{\pi/4} \sin^2(2\theta)\,d\theta = \int_0^{\pi/4} \frac{1}{2}(1-\cos 4\theta)\,d\theta \qquad \text{[half-angle identity]}$$
$$= \frac{1}{2}\left[\theta - \frac{1}{4}\sin 4\theta\right]_0^{\pi/4} = \frac{1}{2}\left[\left(\frac{\pi}{4} - 0\right) - 0\right] = \frac{\pi}{8}$$

9. $$\int_0^{\pi} \cos^4(2t)\,dt = \int_0^{\pi} [\cos^2(2t)]^2\,dt = \int_0^{\pi} \left[\tfrac{1}{2}(1+\cos(2\cdot 2t))\right]^2 dt \qquad \text{[half-angle identity]}$$
$$= \tfrac{1}{4}\int_0^{\pi} [1 + 2\cos 4t + \cos^2(4t)]\,dt = \tfrac{1}{4}\int_0^{\pi} [1 + 2\cos 4t + \tfrac{1}{2}(1+\cos 8t)]\,dt$$
$$= \tfrac{1}{4}\int_0^{\pi} \left(\tfrac{3}{2} + 2\cos 4t + \tfrac{1}{2}\cos 8t\right)\,dt = \tfrac{1}{4}\left[\tfrac{3}{2}t + \tfrac{1}{2}\sin 4t + \tfrac{1}{16}\sin 8t\right]_0^{\pi} = \tfrac{1}{4}\left[\left(\tfrac{3}{2}\pi + 0 + 0\right) - 0\right] = \tfrac{3}{8}\pi$$

10. $$\int_0^{\pi} \sin^2 t\,\cos^4 t\,dt = \tfrac{1}{4}\int_0^{\pi} (4\sin^2 t\,\cos^2 t)\cos^2 t\,dt = \tfrac{1}{4}\int_0^{\pi} (2\sin t\,\cos t)^2\,\tfrac{1}{2}(1+\cos 2t)\,dt$$
$$= \tfrac{1}{8}\int_0^{\pi} (\sin 2t)^2(1+\cos 2t)\,dt = \tfrac{1}{8}\int_0^{\pi} (\sin^2 2t + \sin^2 2t\,\cos 2t)\,dt$$
$$= \tfrac{1}{8}\int_0^{\pi} \sin^2 2t\,dt + \tfrac{1}{8}\int_0^{\pi} \sin^2 2t\,\cos 2t\,dt = \tfrac{1}{8}\int_0^{\pi} \tfrac{1}{2}(1-\cos 4t)\,dt + \tfrac{1}{8}\left[\tfrac{1}{3}\cdot\tfrac{1}{2}\sin^3 2t\right]_0^{\pi}$$
$$= \tfrac{1}{16}\left[t - \tfrac{1}{4}\sin 4t\right]_0^{\pi} + \tfrac{1}{8}(0-0) = \tfrac{1}{16}\left[(\pi - 0) - 0\right] = \tfrac{\pi}{16}$$

11. $$\int_0^{\pi/2} \sin^2 x\,\cos^2 x\,dx = \int_0^{\pi/2} \tfrac{1}{4}(4\sin^2 x\,\cos^2 x)\,dx = \int_0^{\pi/2} \tfrac{1}{4}(2\sin x\,\cos x)^2 dx = \tfrac{1}{4}\int_0^{\pi/2} \sin^2 2x\,dx$$
$$= \tfrac{1}{4}\int_0^{\pi/2} \tfrac{1}{2}(1-\cos 4x)\,dx = \tfrac{1}{8}\int_0^{\pi/2} (1-\cos 4x)\,dx = \tfrac{1}{8}\left[x - \tfrac{1}{4}\sin 4x\right]_0^{\pi/2} = \tfrac{1}{8}\left(\tfrac{\pi}{2}\right) = \tfrac{\pi}{16}$$

12. $$\int_0^{\pi/2} (2-\sin\theta)^2\,d\theta = \int_0^{\pi/2} (4 - 4\sin\theta + \sin^2\theta)\,d\theta = \int_0^{\pi/2} \left[4 - 4\sin\theta + \tfrac{1}{2}(1-\cos 2\theta)\right]d\theta$$
$$= \int_0^{\pi/2} \left(\tfrac{9}{2} - 4\sin\theta - \tfrac{1}{2}\cos 2\theta\right)d\theta = \left[\tfrac{9}{2}\theta + 4\cos\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{\pi/2}$$
$$= \left(\tfrac{9\pi}{4} + 0 - 0\right) - (0 + 4 - 0) = \tfrac{9}{4}\pi - 4$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $\int \sqrt{\cos\theta}\sin^3\theta\,d\theta = \int \sqrt{\cos\theta}\sin^2\theta\,\sin\theta\,d\theta = \int (\cos\theta)^{1/2}(1-\cos^2\theta)\sin\theta\,d\theta$

$$\overset{c}{=} \int u^{1/2}(1-u^2)\,(-du) = \int (u^{5/2}-u^{1/2})\,du$$
$$= \tfrac{2}{7}u^{7/2} - \tfrac{2}{3}u^{3/2} + C = \tfrac{2}{7}(\cos\theta)^{7/2} - \tfrac{2}{3}(\cos\theta)^{3/2} + C$$

14. $\int \left(1+\sqrt[3]{\sin t}\right)\cos^3 t\,dt = \int \left(1+(\sin t)^{1/3}\right)\cos^2 t\cos t\,dt = \int \left(1+(\sin t)^{1/3}\right)(1-\sin^2 t)\cos t\,dt$

$$\overset{s}{=} \int (1+u^{1/3})(1-u^2)\,du = \int (1-u^2+u^{1/3}-u^{7/3})\,du$$
$$= u - \tfrac{1}{3}u^3 + \tfrac{3}{4}u^{4/3} - \tfrac{3}{10}u^{10/3} + C$$
$$= \sin t - \tfrac{1}{3}\sin^3 t + \tfrac{3}{4}\sqrt[3]{\sin^4 t} - \tfrac{3}{10}\sqrt[3]{\sin^{10} t} + C$$

15. $\int \sin x\,\sec^5 x\,dx = \int \frac{\sin x}{\cos^5 x}\,dx \overset{c}{=} \int \frac{1}{u^5}\,(-du) = \frac{1}{4u^4} + C = \frac{1}{4\cos^4 x} + C = \frac{1}{4}\sec^4 x + C$

16. $\int \csc^5\theta\,\cos^3\theta\,d\theta = \int \frac{\cos^2\theta}{\sin^5\theta}\cos\theta\,d\theta = \int \frac{1-\sin^2\theta}{\sin^5\theta}\cos\theta\,d\theta \overset{s}{=} \int \frac{1-u^2}{u^5}\,du = \int (u^{-5}-u^{-3})\,du$

$$= -\frac{1}{4}u^{-4} + \frac{1}{2}u^{-2} + C = -\frac{1}{4\sin^4\theta} + \frac{1}{2\sin^2\theta} + C$$

Alternate solution:

$$\int \csc^5\theta\cos^3\theta\,d\theta = \int \left(\frac{\cos^3\theta}{\sin^3\theta}\right)\left(\frac{1}{\sin^2\theta}\right)d\theta = \int \cot^3\theta\,\csc^2\theta\,d\theta$$
$$= \int u^3\,(-du) \qquad [u=\cot\theta,\ du=-\csc^2\theta\,d\theta]$$
$$= -\int u^3\,du = -\tfrac{1}{4}u^4 + C = -\tfrac{1}{4}\cot^4\theta + C$$

17. $\int \cot x\,\cos^2 x\,dx = \int \frac{\cos x}{\sin x}\,(1-\sin^2 x)\,dx$

$$\overset{s}{=} \int \frac{1-u^2}{u}\,du = \int \left(\frac{1}{u}-u\right)du = \ln|u| - \tfrac{1}{2}u^2 + C = \ln|\sin x| - \tfrac{1}{2}\sin^2 x + C$$

18. $\int \tan^2 x\,\cos^3 x\,dx = \int \frac{\sin^2 x}{\cos^2 x}\cos^3 x\,dx = \int \sin^2 x\,\cos x\,dx \overset{s}{=} \int u^2\,du = \tfrac{1}{3}u^3 + C = \tfrac{1}{3}\sin^3 x + C$

19. $\int \sin^2 x\,\sin 2x\,dx = \int \sin^2 x\,(2\sin x\,\cos x)\,dx \overset{s}{=} \int 2u^3\,du = \tfrac{1}{2}u^4 + C = \tfrac{1}{2}\sin^4 x + C$

20. $\int \sin x\,\cos\left(\tfrac{1}{2}x\right)dx = \int \sin\left(2\cdot\tfrac{1}{2}x\right)\cos\left(\tfrac{1}{2}x\right)dx = \int 2\sin\left(\tfrac{1}{2}x\right)\cos^2\left(\tfrac{1}{2}x\right)dx$

$$= \int 2u^2\,(-2\,du) \qquad [u=\cos\left(\tfrac{1}{2}x\right),\ du=-\tfrac{1}{2}\sin\left(\tfrac{1}{2}x\right)dx]$$
$$= -\tfrac{4}{3}u^3 + C = -\tfrac{4}{3}\cos^3\left(\tfrac{1}{2}x\right) + C$$

21. $\int \tan x\,\sec^3 x\,dx = \int \tan x\,\sec x\,\sec^2 x\,dx = \int u^2\,du \qquad [u=\sec x,\ du=\sec x\,\tan x\,dx]$

$$= \tfrac{1}{3}u^3 + C = \tfrac{1}{3}\sec^3 x + C$$

22. $\int \tan^2\theta\,\sec^4\theta\,d\theta = \int \tan^2\theta\,\sec^2\theta\,\sec^2\theta\,d\theta = \int \tan^2\theta\,(\tan^2\theta+1)\,\sec^2\theta\,d\theta$

$$= \int u^2(u^2+1)\,du \qquad [u=\tan\theta,\ du=\sec^2\theta\,d\theta]$$
$$= \int (u^4+u^2)\,du = \tfrac{1}{5}u^5 + \tfrac{1}{3}u^3 + C = \tfrac{1}{5}\tan^5\theta + \tfrac{1}{3}\tan^3\theta + C$$

23. $\int \tan^2 x\,dx = \int (\sec^2 x - 1)\,dx = \tan x - x + C$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $\int(\tan^2 x+\tan^4 x)\,dx=\int \tan^2 x\,(1+\tan^2 x)\,dx=\int \tan^2 x\,\sec^2 x\,dx=\int u^2\,du \qquad [u=\tan x,\ du=\sec^2 x\,dx]$

$$=\tfrac{1}{3}u^3+C=\tfrac{1}{3}\tan^3 x+C$$

25. Let $u=\tan x$. Then $du=\sec^2 x\,dx$, so

$$\int \tan^4 x\sec^6 x\,dx=\int \tan^4 x\sec^4 x\,(\sec^2 x\,dx)=\int \tan^4 x(1+\tan^2 x)^2\,(\sec^2 x\,dx)$$
$$=\int u^4(1+u^2)^2\,du=\int(u^8+2u^6+u^4)\,du$$
$$=\tfrac{1}{9}u^9+\tfrac{2}{7}u^7+\tfrac{1}{5}u^5+C=\tfrac{1}{9}\tan^9 x+\tfrac{2}{7}\tan^7 x+\tfrac{1}{5}\tan^5 x+C$$

26. $\int_0^{\pi/4}\sec^6\theta\,\tan^6\theta\,d\theta=\int_0^{\pi/4}\tan^6\theta\,\sec^4\theta\,\sec^2\theta\,d\theta=\int_0^{\pi/4}\tan^6\theta(1+\tan^2\theta)^2\sec^2\theta\,d\theta$

$$=\int_0^1 u^6(1+u^2)^2\,du \qquad \begin{bmatrix} u=\tan\theta, \\ du=\sec^2\theta\,d\theta \end{bmatrix}$$
$$=\int_0^1 u^6(u^4+2u^2+1)\,du=\int_0^1(u^{10}+2u^8+u^6)\,du$$
$$=\left[\tfrac{1}{11}u^{11}+\tfrac{2}{9}u^9+\tfrac{1}{7}u^7\right]_0^1=\tfrac{1}{11}+\tfrac{2}{9}+\tfrac{1}{7}=\tfrac{63+154+99}{693}=\tfrac{316}{693}$$

27. $\int \tan^3 x\,\sec x\,dx=\int \tan^2 x\,\sec x\,\tan x\,dx=\int(\sec^2 x-1)\,\sec x\,\tan x\,dx$

$$=\int(u^2-1)\,du \quad \left[u=\sec x, du=\sec x\,\tan x\,dx\right] \quad =\tfrac{1}{3}u^3-u+C=\tfrac{1}{3}\sec^3 x-\sec x+C$$

28. Let $u=\sec x$, so $du=\sec x\tan x\,dx$. Thus,

$$\int \tan^5 x\sec^3 x\,dx=\int \tan^4 x\sec^2 x\,(\sec x\tan x)\,dx=\int(\sec^2 x-1)^2\sec^2 x\,(\sec x\tan x\,dx)$$
$$=\int(u^2-1)^2u^2\,du=\int(u^6-2u^4+u^2)\,du$$
$$=\tfrac{1}{7}u^7-\tfrac{2}{5}u^5+\tfrac{1}{3}u^3+C=\tfrac{1}{7}\sec^7 x-\tfrac{2}{5}\sec^5 x+\tfrac{1}{3}\sec^3 x+C$$

29. $\int \tan^3 x\,\sec^6 x\,dx=\int \tan^3 x\,\sec^4 x\,\sec^2 x\,dx=\int \tan^3 x\,(1+\tan^2 x)^2\sec^2 x\,dx$

$$=\int u^3(1+u^2)^2\,du \qquad \begin{bmatrix} u=\tan x, \\ du=\sec^2 x\,dx \end{bmatrix}$$
$$=\int u^3(u^4+2u^2+1)\,du=\int(u^7+2u^5+u^3)\,du$$
$$=\tfrac{1}{8}u^8+\tfrac{1}{3}u^6+\tfrac{1}{4}u^4+C=\tfrac{1}{8}\tan^8 x+\tfrac{1}{3}\tan^6 x+\tfrac{1}{4}\tan^4 x+C$$

30. $\int_0^{\pi/4}\tan^4 t\,dt=\int_0^{\pi/4}\tan^2 t\,(\sec^2 t-1)\,dt=\int_0^{\pi/4}\tan^2 t\,\sec^2 t\,dt-\int_0^{\pi/4}\tan^2 t\,dt$

$$=\int_0^1 u^2\,du\ \ [u=\tan t]\ \ -\int_0^{\pi/4}(\sec^2 t-1)\,dt=\left[\tfrac{1}{3}u^3\right]_0^1-\Big[\tan t-t\Big]_0^{\pi/4}$$
$$=\tfrac{1}{3}-\left[\left(1-\tfrac{\pi}{4}\right)-0\right]=\tfrac{\pi}{4}-\tfrac{2}{3}$$

31. $\int \tan^5 x\,dx=\int(\sec^2 x-1)^2\,\tan x\,dx=\int \sec^4 x\,\tan x\,dx-2\int \sec^2 x\,\tan x\,dx+\int \tan x\,dx$

$$=\int \sec^3 x\,\sec x\,\tan x\,dx-2\int \tan x\,\sec^2 x\,dx+\int \tan x\,dx$$
$$=\tfrac{1}{4}\sec^4 x-\tan^2 x+\ln|\sec x|+C \quad \left[\text{or } \tfrac{1}{4}\sec^4 x-\sec^2 x+\ln|\sec x|+C\right]$$

32. $\int \tan^2 x\,\sec x\,dx=\int(\sec^2 x-1)\,\sec x\,dx=\int \sec^3 x\,dx-\int \sec x\,dx$

$$=\tfrac{1}{2}(\sec x\,\tan x+\ln|\sec x+\tan x|)-\ln|\sec x+\tan x|+C \qquad \text{[by Example 8 and (1)]}$$
$$=\tfrac{1}{2}(\sec x\,\tan x-\ln|\sec x+\tan x|)+C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. $\displaystyle\int \frac{1-\tan^2 x}{\sec^2 x}\,dx = \int \left(\cos^2 x - \sin^2 x\right) dx = \int \cos 2x\,dx = \tfrac{1}{2}\sin 2x + C$

34. $\displaystyle\int \frac{\tan x \sec^2 x}{\cos x}\,dx = \int \sec^2 x \tan x \sec x\,dx = \int u^2\,du \qquad [u = \sec x,\ du = \sec x\ \tan x\,dx]$

$\displaystyle = \tfrac{1}{3}u^3 + C = \tfrac{1}{3}\sec^3 x + C$

35. $\displaystyle\int_0^{\pi/4} \frac{\sin^3 x}{\cos x}\,dx = \int_0^{\pi/4} \frac{\sin^2 x}{\cos x}\sin x\,dx = \int_0^{\pi/4} \frac{1-\cos^2 x}{\cos x}\sin x\,dx \overset{c}{=} \int_1^{1/\sqrt{2}} \frac{1-u^2}{u}\,(-du)$

$\displaystyle = \int_{1/\sqrt{2}}^{1} \left(\frac{1}{u} - u\right) du = \left[\ln|u| - \frac{1}{2}u^2\right]_{1/\sqrt{2}}^{1} = \left(\ln 1 - \frac{1}{2}\right) - \left(\ln\frac{1}{\sqrt{2}} - \frac{1}{4}\right) = -\frac{1}{4} - \ln\frac{\sqrt{2}}{2}$

36. $\displaystyle\int \frac{\sin\theta + \tan\theta}{\cos^3\theta}\,d\theta = \int \frac{\sin\theta}{\cos^3\theta}\,d\theta + \int \frac{\tan\theta}{\cos^3\theta}\,d\theta = \int \frac{\sin\theta}{\cos^3\theta}\,d\theta + \int \sec^2\theta\tan\theta\sec\theta\,d\theta$

$\displaystyle = -\int \frac{1}{u^3}\,du + \int v^2\,dv \qquad \begin{bmatrix} u = \cos\theta,\ du = -\sin\theta\,d\theta \\ v = \sec\theta,\ dv = \sec\theta\tan\theta\,d\theta \end{bmatrix}$

$\displaystyle = \frac{1}{2u^2} + \frac{1}{3}v^3 + C = \frac{1}{2\cos^2\theta} + \frac{1}{3}\sec^3\theta + C = \frac{1}{2}\sec^2\theta + \frac{1}{3}\sec^3\theta + C$

37. $\int_{\pi/6}^{\pi/2} \cot^2 x\,dx = \int_{\pi/6}^{\pi/2} (\csc^2 x - 1)\,dx = \left[-\cot x - x\right]_{\pi/6}^{\pi/2} = \left(0 - \frac{\pi}{2}\right) - \left(-\sqrt{3} - \frac{\pi}{6}\right) = \sqrt{3} - \frac{\pi}{3}$

38. $\displaystyle\int_{\pi/4}^{\pi/2} \cot^3 x\,dx = \int_{\pi/4}^{\pi/2} \cot x\,(\csc^2 x - 1)\,dx = \int_{\pi/4}^{\pi/2} \cot x\ \csc^2 x\,dx - \int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin x}\,dx$

$\displaystyle = \left[-\tfrac{1}{2}\cot^2 x - \ln|\sin x|\right]_{\pi/4}^{\pi/2} = (0 - \ln 1) - \left[-\tfrac{1}{2} - \ln\tfrac{1}{\sqrt{2}}\right] = \tfrac{1}{2} + \ln\tfrac{1}{\sqrt{2}} = \tfrac{1}{2}(1 - \ln 2)$

39. $\int_{\pi/4}^{\pi/2} \cot^5\phi\ \csc^3\phi\,d\phi = \int_{\pi/4}^{\pi/2} \cot^4\phi\ \csc^2\phi\ \csc\phi\ \cot\phi\,d\phi = \int_{\pi/4}^{\pi/2} (\csc^2\phi - 1)^2\ \csc^2\phi\ \csc\phi\ \cot\phi\,d\phi$

$\displaystyle = \int_{\sqrt{2}}^{1} (u^2 - 1)^2 u^2\,(-du) \qquad [u = \csc\phi,\ du = -\csc\phi\ \cot\phi\,d\phi]$

$\displaystyle = \int_1^{\sqrt{2}} (u^6 - 2u^4 + u^2)\,du = \left[\tfrac{1}{7}u^7 - \tfrac{2}{5}u^5 + \tfrac{1}{3}u^3\right]_1^{\sqrt{2}} = \left(\tfrac{8}{7}\sqrt{2} - \tfrac{8}{5}\sqrt{2} + \tfrac{2}{3}\sqrt{2}\right) - \left(\tfrac{1}{7} - \tfrac{2}{5} + \tfrac{1}{3}\right)$

$\displaystyle = \frac{120 - 168 + 70}{105}\sqrt{2} - \frac{15 - 42 + 35}{105} = \frac{22}{105}\sqrt{2} - \frac{8}{105}$

40. $\int_{\pi/4}^{\pi/2} \csc^4\theta\ \cot^4\theta\,d\theta = \int_{\pi/4}^{\pi/2} \cot^4\theta\ \csc^2\theta\ \csc^2\theta\,d\theta = \int_{\pi/4}^{\pi/2} \cot^4\theta\,(\cot^2\theta + 1)\ \csc^2\theta\,d\theta$

$\displaystyle = \int_1^0 u^4(u^2 + 1)\,(-du) \qquad \begin{bmatrix} u = \cot\theta, \\ du = -\csc^2\theta\,d\theta \end{bmatrix}$

$\displaystyle = \int_0^1 (u^6 + u^4)\,du$

$\displaystyle = \left[\tfrac{1}{7}u^7 + \tfrac{1}{5}u^5\right]_0^1 = \tfrac{1}{7} + \tfrac{1}{5} = \tfrac{12}{35}$

41. $I = \displaystyle\int \csc x\,dx = \int \frac{\csc x\,(\csc x - \cot x)}{\csc x - \cot x}\,dx = \int \frac{-\csc x\ \cot x + \csc^2 x}{\csc x - \cot x}\,dx$. Let $u = \csc x - \cot x \quad \Rightarrow$

$du = (-\csc x\ \cot x + \csc^2 x)\,dx$. Then $I = \int du/u = \ln|u| = \ln|\csc x - \cot x| + C$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. Let $u = \csc x$, $dv = \csc^2 x\,dx$. Then $du = -\csc x\,\cot x\,dx$, $v = -\cot x \quad \Rightarrow$

$$\int \csc^3 x\,dx = -\csc x\,\cot x - \int \csc x\,\cot^2 x\,dx = -\csc x\,\cot x - \int \csc x\,(\csc^2 x - 1)\,dx$$
$$= -\csc x\,\cot x + \int \csc x\,dx - \int \csc^3 x\,dx$$

Solving for $\int \csc^3 x\,dx$ and using Exercise 41, we get

$\int \csc^3 x\,dx = -\frac{1}{2}\csc x\,\cot x + \frac{1}{2}\int \csc x\,dx = -\frac{1}{2}\csc x\,\cot x + \frac{1}{2}\ln|\csc x - \cot x| + C$. Thus,

$$\int_{\pi/6}^{\pi/3} \csc^3 x\,dx = \left[-\tfrac{1}{2}\csc x\,\cot x + \tfrac{1}{2}\ln|\csc x - \cot x|\right]_{\pi/6}^{\pi/3}$$
$$= -\tfrac{1}{2}\cdot\tfrac{2}{\sqrt{3}}\cdot\tfrac{1}{\sqrt{3}} + \tfrac{1}{2}\ln\left|\tfrac{2}{\sqrt{3}} - \tfrac{1}{\sqrt{3}}\right| + \tfrac{1}{2}\cdot 2\cdot\sqrt{3} - \tfrac{1}{2}\ln\left|2 - \sqrt{3}\,\right|$$
$$= -\tfrac{1}{3} + \sqrt{3} + \tfrac{1}{2}\ln\tfrac{1}{\sqrt{3}} - \tfrac{1}{2}\ln\left(2 - \sqrt{3}\,\right) \approx 1.7825$$

43. $\int \sin 8x\,\cos 5x\,dx \overset{2a}{=} \int \frac{1}{2}[\sin(8x - 5x) + \sin(8x + 5x)]\,dx = \frac{1}{2}\int(\sin 3x + \sin 13x)\,dx$

$$= \tfrac{1}{2}\left(-\tfrac{1}{3}\cos 3x - \tfrac{1}{13}\cos 13x\right) + C = -\tfrac{1}{6}\cos 3x - \tfrac{1}{26}\cos 13x + C$$

44. $\int \sin 2\theta\,\sin 6\theta\,d\theta \overset{2b}{=} \int \frac{1}{2}[\cos(2\theta - 6\theta) - \cos(2\theta + 6\theta)]\,d\theta$

$$= \tfrac{1}{2}\int[\cos(-4\theta) - \cos 8\theta]\,d\theta = \tfrac{1}{2}\int(\cos 4\theta - \cos 8\theta)\,d\theta$$
$$= \tfrac{1}{2}\left(\tfrac{1}{4}\sin 4\theta - \tfrac{1}{8}\sin 8\theta\right) + C = \tfrac{1}{8}\sin 4\theta - \tfrac{1}{16}\sin 8\theta + C$$

45. $\int_0^{\pi/2} \cos 5t\,\cos 10t\,dt \overset{2c}{=} \int_0^{\pi/2} \frac{1}{2}[\cos(5t - 10t) + \cos(5t + 10t)]\,dt$

$$= \tfrac{1}{2}\int_0^{\pi/2}[\cos(-5t) + \cos 15t]\,dt = \tfrac{1}{2}\int_0^{\pi/2}(\cos 5t + \cos 15t)\,dt$$
$$= \tfrac{1}{2}\left[\tfrac{1}{5}\sin 5t + \tfrac{1}{15}\sin 15t\right]_0^{\pi/2} = \tfrac{1}{2}\left(\tfrac{1}{5} - \tfrac{1}{15}\right) = \tfrac{1}{15}$$

46. $\int t\cos^5(t^2)\,dt = \int t\cos^4(t^2)\,\cos(t^2)\,dt = \int t[1 - \sin^2(t^2)]^2\cos(t^2)\,dt$

$$= \int \tfrac{1}{2}(1 - u^2)^2\,du \qquad \left[u = \sin(t^2),\ du = 2t\cos(t^2)\,dt\right]$$
$$= \tfrac{1}{2}\int(u^4 - 2u^2 + 1)\,du = \tfrac{1}{2}\left(\tfrac{1}{5}u^5 - \tfrac{2}{3}u^3 + u\right) + C = \tfrac{1}{10}\sin^5(t^2) - \tfrac{1}{3}\sin^3(t^2) + \tfrac{1}{2}\sin(t^2) + C$$

47. $$\int \frac{\sin^2(1/t)}{t^2}\,dt = \int \sin^2 u\,(-du) \qquad \left[u = \tfrac{1}{t},\ du = -\tfrac{1}{t^2}\,dt\right]$$
$$= -\int \frac{1}{2}(1 - \cos 2u)\,du = -\frac{1}{2}\left(u - \frac{1}{2}\sin 2u\right) + C = -\frac{1}{2t} + \frac{1}{4}\sin\left(\frac{2}{t}\right) + C$$

48. $\int \sec^2 y\cos^3(\tan y)\,dy = \int \cos^3 u\,du \qquad \left[u = \tan y,\ du = \sec^2 y\,dy\right]$

$$= \sin\,u - \tfrac{1}{3}\sin^3 u + C \qquad [\text{by Example 1}]$$
$$= \sin(\tan y) - \tfrac{1}{3}\sin^3(\tan y) + C$$

49. $\int_0^{\pi/6} \sqrt{1 + \cos 2x}\,dx = \int_0^{\pi/6} \sqrt{1 + (2\cos^2 x - 1)}\,dx = \int_0^{\pi/6} \sqrt{2\cos^2 x}\,dx = \sqrt{2}\int_0^{\pi/6} \sqrt{\cos^2 x}\,dx$

$$= \sqrt{2}\int_0^{\pi/6} |\cos x|\,dx = \sqrt{2}\int_0^{\pi/6} \cos x\,dx \qquad [\text{since } \cos x > 0 \text{ for } 0 \le x \le \pi/6]$$
$$= \sqrt{2}\left[\sin x\right]_0^{\pi/6} = \sqrt{2}\left(\tfrac{1}{2} - 0\right) = \tfrac{1}{2}\sqrt{2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50. $\int_0^{\pi/4} \sqrt{1-\cos 4\theta}\, d\theta = \int_0^{\pi/4} \sqrt{1-(1-2\sin^2(2\theta))}\, d\theta = \int_0^{\pi/4} \sqrt{2\sin^2(2\theta)}\, d\theta = \sqrt{2}\int_0^{\pi/4} \sqrt{\sin^2(2\theta)}\, d\theta$

$$= \sqrt{2}\int_0^{\pi/4} |\sin 2\theta|\, d\theta = \sqrt{2}\int_0^{\pi/4} \sin 2\theta\, d\theta \qquad \text{[since } \sin 2\theta \geq 0 \text{ for } 0 \leq \theta \leq \pi/4\text{]}$$

$$= \sqrt{2}\left[-\tfrac{1}{2}\cos 2\theta\right]_0^{\pi/4} = -\tfrac{1}{2}\sqrt{2}\,(0-1) = \tfrac{1}{2}\sqrt{2}$$

51. $\int t\sin^2 t\, dt = \int t\left[\tfrac{1}{2}(1-\cos 2t)\right]\, dt = \tfrac{1}{2}\int (t - t\cos 2t)\, dt = \tfrac{1}{2}\int t\, dt - \tfrac{1}{2}\int t\cos 2t\, dt$

$$= \tfrac{1}{2}\left(\tfrac{1}{2}t^2\right) - \tfrac{1}{2}\left(\tfrac{1}{2}t\sin 2t - \int \tfrac{1}{2}\sin 2t\, dt\right) \qquad \begin{bmatrix} u = t, & dv = \cos 2t\, dt \\ du = dt, & v = \tfrac{1}{2}\sin 2t \end{bmatrix}$$

$$= \tfrac{1}{4}t^2 - \tfrac{1}{4}t\sin 2t + \tfrac{1}{2}\left(-\tfrac{1}{4}\cos 2t\right) + C = \tfrac{1}{4}t^2 - \tfrac{1}{4}t\sin 2t - \tfrac{1}{8}\cos 2t + C$$

52. Let $u = x, dv = \sec x\, \tan x\, dx \quad \Rightarrow \quad du = dx, v = \sec x$. Then

$\int x\, \sec x\, \tan x\, dx = x\sec x - \int \sec x\, dx = x\sec x - \ln|\sec x + \tan x| + C.$

53. $\int x\tan^2 x\, dx = \int x(\sec^2 x - 1)\, dx = \int x\sec^2 x\, dx - \int x\, dx$

$$= x\tan x - \int \tan x\, dx - \tfrac{1}{2}x^2 \qquad \begin{bmatrix} u = x, & dv = \sec^2 x\, dx \\ du = dx, & v = \tan x \end{bmatrix}$$

$$= x\tan x - \ln|\sec x| - \tfrac{1}{2}x^2 + C$$

54. $I = \int x\sin^3 x\, dx$. First, evaluate

$$\int \sin^3 x\, dx = \int (1-\cos^2 x)\, \sin x\, dx \overset{c}{=} \int (1-u^2)(-du) = \int (u^2-1)\, du$$

$$= \tfrac{1}{3}u^3 - u + C_1 = \tfrac{1}{3}\cos^3 x - \cos x + C_1.$$

Now for I, let $u = x$, $dv = \sin^3 x \quad \Rightarrow \quad du = dx$, $v = \tfrac{1}{3}\cos^3 x - \cos x$, so

$$I = \tfrac{1}{3}x\cos^3 x - x\cos x - \int \left(\tfrac{1}{3}\cos^3 x - \cos x\right) dx = \tfrac{1}{3}x\cos^3 x - x\cos x - \tfrac{1}{3}\int \cos^3 x\, dx + \sin x$$

$$= \tfrac{1}{3}x\cos^3 x - x\cos x - \tfrac{1}{3}(\sin x - \tfrac{1}{3}\sin^3 x) + \sin x + C \qquad \text{[by Example 1]}$$

$$= \tfrac{1}{3}x\cos^3 x - x\cos x + \tfrac{2}{3}\sin x + \tfrac{1}{9}\sin^3 x + C$$

55. $\displaystyle\int \frac{dx}{\cos x - 1} = \int \frac{1}{\cos x - 1}\cdot\frac{\cos x + 1}{\cos x + 1}\, dx = \int \frac{\cos x + 1}{\cos^2 x - 1}\, dx = \int \frac{\cos x + 1}{-\sin^2 x}\, dx$

$$= \int \left(-\cot x\, \csc x - \csc^2 x\right) dx = \csc x + \cot x + C$$

56. $\displaystyle\int \frac{1}{\sec\theta + 1}\, d\theta = \int \frac{1}{\sec\theta + 1}\cdot\frac{\sec\theta - 1}{\sec\theta - 1}\, d\theta = \int \frac{\sec\theta - 1}{\sec^2\theta - 1}\, d\theta = \int \frac{\sec\theta - 1}{\tan^2\theta}\, d\theta$

$$= \int \frac{\cos\theta}{\sin^2\theta}\, d\theta - \int \frac{\cos^2\theta}{\sin^2\theta}\, d\theta = \int \frac{\cos\theta\, d\theta}{\sin^2\theta} - \int \frac{1-\sin^2\theta}{\sin^2\theta}\, d\theta \overset{s}{=} \int \frac{1}{u^2}\, du - \int \csc^2\theta\, d\theta + \int d\theta$$

$$= -\frac{1}{\sin\theta} + \cot\theta + \theta + C$$

Alternate solution:

$$\int \frac{1}{\sec\theta + 1}\, d\theta = \int \frac{\cos\theta}{1+\cos\theta}\, d\theta = \int \frac{2\cos^2\left(\dfrac{\theta}{2}\right) - 1}{2\cos^2\left(\dfrac{\theta}{2}\right)}\, d\theta \qquad \text{[double-angle identities]}$$

$$= \int 1\, d\theta - \int \frac{1}{2}\sec^2\left(\frac{\theta}{2}\right)\, d\theta = \theta - \tan\left(\frac{\theta}{2}\right) + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

In Exercises 57–60, let $f(x)$ denote the integrand and $F(x)$ its antiderivative (with $C = 0$).

57. Let $u = x^2$, so that $du = 2x\,dx$. Then

$$\begin{aligned}\int x\sin^2(x^2)\,dx &= \int \sin^2 u\left(\tfrac{1}{2}\,du\right) = \tfrac{1}{2}\int \tfrac{1}{2}(1-\cos 2u)\,du \\ &= \tfrac{1}{4}\left(u - \tfrac{1}{2}\sin 2u\right) + C = \tfrac{1}{4}u - \tfrac{1}{4}\left(\tfrac{1}{2}\cdot 2\sin u\,\cos u\right) + C \\ &= \tfrac{1}{4}x^2 - \tfrac{1}{4}\sin(x^2)\,\cos(x^2) + C\end{aligned}$$

We see from the graph that this is reasonable, since F increases where f is positive and F decreases where f is negative. Note also that f is an odd function and F is an even function.

58.
$$\begin{aligned}\int \sin^5 x\,\cos^3 x\,dx &= \int \sin^5 x\,\cos^2 x\,\cos x\,dx \\ &= \int \sin^5 x\,(1-\sin^2 x)\,\cos x\,dx \\ &\overset{s}{=} \int u^5(1-u^2)\,du = \int (u^5 - u^7)\,du \\ &= \tfrac{1}{6}\sin^6 x - \tfrac{1}{8}\sin^8 x + C\end{aligned}$$

We see from the graph that this is reasonable, since F increases where f is positive and F decreases where f is negative. Note also that f is an odd function and F is an even function.

59.
$$\begin{aligned}\int \sin 3x\,\sin 6x\,dx &= \int \tfrac{1}{2}[\cos(3x-6x) - \cos(3x+6x)]\,dx \\ &= \tfrac{1}{2}\int(\cos 3x - \cos 9x)\,dx \\ &= \tfrac{1}{6}\sin 3x - \tfrac{1}{18}\sin 9x + C\end{aligned}$$

Notice that $f(x) = 0$ whenever F has a horizontal tangent.

60.
$$\begin{aligned}\int \sec^4\left(\tfrac{1}{2}x\right)\,dx &= \int \left(\tan^2\tfrac{x}{2} + 1\right)\sec^2\tfrac{x}{2}\,dx \\ &= \int (u^2+1)\,2\,du \quad \left[u = \tan\tfrac{x}{2},\ du = \tfrac{1}{2}\sec^2\tfrac{x}{2}\,dx\right] \\ &= \tfrac{2}{3}u^3 + 2u + C = \tfrac{2}{3}\tan^3\tfrac{x}{2} + 2\tan\tfrac{x}{2} + C\end{aligned}$$

Notice that F is increasing and f is positive on the intervals on which they are defined. Also, F has no horizontal tangent and f is never zero.

61. Let $u = \tan^7 x$, $dv = \sec x\,\tan x\,dx \quad\Rightarrow\quad du = 7\tan^6 x\,\sec^2 x\,dx$, $v = \sec x$. Then

$$\begin{aligned}\int \tan^8 x\,\sec x\,dx &= \int \tan^7 x\cdot\sec x\,\tan x\,dx = \tan^7 x\,\sec x - \int 7\tan^6 x\sec^2 x\sec x\,dx \\ &= \tan^7 x\,\sec x - 7\int \tan^6 x\,(\tan^2 x + 1)\,\sec x\,dx \\ &= \tan^7 x\,\sec x - 7\int \tan^8 x\sec x\,dx - 7\int \tan^6 x\sec x\,dx\end{aligned}$$

Thus, $8\int \tan^8 x\sec x\,dx = \tan^7 x\,\sec x - 7\int \tan^6 x\sec x\,dx$ and

$$\int_0^{\pi/4} \tan^8 x\sec x\,dx = \frac{1}{8}\left[\tan^7 x\,\sec x\right]_0^{\pi/4} - \frac{7}{8}\int_0^{\pi/4} \tan^6 x\,\sec x\,dx = \frac{\sqrt{2}}{8} - \frac{7}{8}I.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

62. (a) $\int \tan^{2n} x\,dx = \int \tan^{2n-2} x \tan^2 x\,dx = \int \tan^{2n-2} x\,(\sec^2 x - 1)\,dx$

$$= \int \tan^{2n-2} x \sec^2 x\,dx - \int \tan^{2n-2} x\,dx$$

$$= \int u^{2n-2}\,du - \int \tan^{2n-2} x\,dx \qquad [u = \tan x, du = \sec^2 x\,dx]$$

$$= \frac{u^{2n-1}}{2n-1} - \int \tan^{2n-2} x\,dx = \frac{\tan^{2n-1} x}{2n-1} - \int \tan^{2n-2} x\,dx$$

(b) Starting with $n = 4$, repeated applications of the reduction formula in part (a) gives

$$\int \tan^8 x\,dx = \frac{\tan^7 x}{7} - \int \tan^6 x\,dx = \frac{\tan^7 x}{7} - \left(\frac{\tan^5 x}{5} - \int \tan^4 x\,dx\right)$$

$$= \frac{\tan^7 x}{7} - \frac{\tan^5 x}{5} + \left(\frac{\tan^3 x}{3} - \int \tan^2 x\,dx\right)$$

$$= \frac{\tan^7 x}{7} - \frac{\tan^5 x}{5} + \frac{\tan^3 x}{3} - \left(\frac{\tan x}{1} - \int 1\,dx\right)$$

$$= \frac{\tan^7 x}{7} - \frac{\tan^5 x}{5} + \frac{\tan^3 x}{3} - \tan x + x + C$$

63. $f_{\text{avg}} = \frac{1}{2\pi}\int_{-\pi}^{\pi} \sin^2 x\,\cos^3 x\,dx = \frac{1}{2\pi}\int_{-\pi}^{\pi} \sin^2 x\,(1 - \sin^2 x)\,\cos x\,dx$

$= \frac{1}{2\pi}\int_0^0 u^2(1-u^2)\,du$ [where $u = \sin x$] $= 0$

64. (a) Let $u = \cos x$. Then $du = -\sin x\,dx \;\Rightarrow\; \int \sin x\,\cos x\,dx = \int u(-du) = -\frac{1}{2}u^2 + C = -\frac{1}{2}\cos^2 x + C_1$.

(b) Let $u = \sin x$. Then $du = \cos x\,dx \;\Rightarrow\; \int \sin x\,\cos x\,dx = \int u\,du = \frac{1}{2}u^2 + C = \frac{1}{2}\sin^2 x + C_2$.

(c) $\int \sin x\,\cos x\,dx = \int \frac{1}{2}\sin 2x\,dx = -\frac{1}{4}\cos 2x + C_3$

(d) Let $u = \sin x$, $dv = \cos x\,dx$. Then $du = \cos x\,dx$, $v = \sin x$, so $\int \sin x\,\cos x\,dx = \sin^2 x - \int \sin x\,\cos x\,dx$, by Equation 7.1.2, so $\int \sin x\,\cos x\,dx = \frac{1}{2}\sin^2 x + C_4$.

Using $\cos^2 x = 1 - \sin^2 x$ and $\cos 2x = 1 - 2\sin^2 x$, we see that the answers differ only by a constant.

65. $A = \int_0^{\pi} (\sin^2 x - \sin^3 x)\,dx = \int_0^{\pi} \left[\frac{1}{2}(1 - \cos 2x) - \sin x\,(1 - \cos^2 x)\right]\,dx$

$= \int_0^{\pi} \left(\frac{1}{2} - \frac{1}{2}\cos 2x\right)\,dx + \int_1^{-1}(1-u^2)\,du \qquad \begin{bmatrix} u = \cos x, \\ du = -\sin x\,dx \end{bmatrix}$

$= \left[\frac{1}{2}x - \frac{1}{4}\sin 2x\right]_0^{\pi} + 2\int_0^1 (u^2 - 1)\,du$

$= \left(\frac{1}{2}\pi - 0\right) - (0 - 0) + 2\left[\frac{1}{3}u^3 - u\right]_0^1$

$= \frac{1}{2}\pi + 2\left(\frac{1}{3} - 1\right) = \frac{1}{2}\pi - \frac{4}{3}$

66. $A = \int_0^{\pi/4} (\tan x - \tan^2 x)\,dx = \int_0^{\pi/4} (\tan x - \sec^2 x + 1)\,dx$

$= \Big[\ln|\sec x| - \tan x + x\Big]_0^{\pi/4} = \left(\ln\sqrt{2} - 1 + \frac{\pi}{4}\right) - (\ln 1 - 0 + 0)$

$= \ln\sqrt{2} - 1 + \frac{\pi}{4}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67.

It seems from the graph that $\int_0^{2\pi} \cos^3 x\,dx = 0$, since the area below the x-axis and above the graph looks about equal to the area above the axis and below the graph. By Example 1, the integral is $\left[\sin x - \frac{1}{3}\sin^3 x\right]_0^{2\pi} = 0$. Note that due to symmetry, the integral of any odd power of $\sin x$ or $\cos x$ between limits which differ by $2n\pi$ (n any integer) is 0.

68.

It seems from the graph that $\int_0^2 \sin 2\pi x\,\cos 5\pi x\,dx = 0$, since each bulge above the x-axis seems to have a corresponding depression below the x-axis. To evaluate the integral, we use a trigonometric identity:

$$\begin{aligned}\int_0^1 \sin 2\pi x\,\cos 5\pi x\,dx &= \tfrac{1}{2}\int_0^2 [\sin(2\pi x - 5\pi x) + \sin(2\pi x + 5\pi x)]\,dx \\ &= \tfrac{1}{2}\int_0^2 [\sin(-3\pi x) + \sin 7\pi x]\,dx \\ &= \tfrac{1}{2}\left[\tfrac{1}{3\pi}\cos(-3\pi x) - \tfrac{1}{7\pi}\cos 7\pi x\right]_0^2 \\ &= \tfrac{1}{2}\left[\tfrac{1}{3\pi}(1-1) - \tfrac{1}{7\pi}(1-1)\right] = 0\end{aligned}$$

69. Using disks, $V = \int_{\pi/2}^{\pi} \pi \sin^2 x\,dx = \pi\int_{\pi/2}^{\pi} \frac{1}{2}(1-\cos 2x)\,dx = \pi\left[\frac{1}{2}x - \frac{1}{4}\sin 2x\right]_{\pi/2}^{\pi} = \pi\left(\frac{\pi}{2} - 0 - \frac{\pi}{4} + 0\right) = \frac{\pi^2}{4}$

70. Using disks,

$$\begin{aligned}V &= \int_0^{\pi} \pi(\sin^2 x)^2\,dx = 2\pi\int_0^{\pi/2}\left[\tfrac{1}{2}(1-\cos 2x)\right]^2 dx \\ &= \tfrac{\pi}{2}\int_0^{\pi/2}(1 - 2\cos 2x + \cos^2 2x)\,dx \\ &= \tfrac{\pi}{2}\int_0^{\pi/2}\left[1 - 2\cos 2x + \tfrac{1}{2}(1-\cos 4x)\right]dx \\ &= \tfrac{\pi}{2}\int_0^{\pi/2}\left(\tfrac{3}{2} - 2\cos 2x - \tfrac{1}{2}\cos 4x\right)dx = \tfrac{\pi}{2}\left[\tfrac{3}{2}x - \sin 2x + \tfrac{1}{8}\sin 4x\right]_0^{\pi/2} \\ &= \tfrac{\pi}{2}\left[\left(\tfrac{3\pi}{4} - 0 + 0\right) - (0 - 0 + 0)\right] = \tfrac{3}{8}\pi^2\end{aligned}$$

71. Using washers,

$$\begin{aligned}V &= \int_0^{\pi/4} \pi\left[(1-\sin x)^2 - (1-\cos x)^2\right]dx \\ &= \pi\int_0^{\pi/4}\left[(1 - 2\sin x + \sin^2 x) - (1 - 2\cos x + \cos^2 x)\right]dx \\ &= \pi\int_0^{\pi/4}(2\cos x - 2\sin x + \sin^2 x - \cos^2 x)\,dx \\ &= \pi\int_0^{\pi/4}(2\cos x - 2\sin x - \cos 2x)\,dx = \pi\left[2\sin x + 2\cos x - \tfrac{1}{2}\sin 2x\right]_0^{\pi/4} \\ &= \pi\left[\left(\sqrt{2} + \sqrt{2} - \tfrac{1}{2}\right) - (0 + 2 - 0)\right] = \pi\left(2\sqrt{2} - \tfrac{5}{2}\right)\end{aligned}$$

72. Using washers,

$$\begin{aligned}V &= \int_0^{\pi/3} \pi\left\{[\sec x - (-1)]^2 - [\cos x - (-1)]^2\right\}dx \\ &= \pi\int_0^{\pi/3}[(\sec^2 x + 2\sec x + 1) - (\cos^2 x + 2\cos x + 1)]\,dx \\ &= \pi\int_0^{\pi/3}\left[\sec^2 x + 2\sec x - \tfrac{1}{2}(1+\cos 2x) - 2\cos x\right]dx \\ &= \pi\left[\tan x + 2\ln|\sec x + \tan x| - \tfrac{1}{2}x - \tfrac{1}{4}\sin 2x - 2\sin x\right]_0^{\pi/3} \\ &= \pi\left[\left(\sqrt{3} + 2\ln\left(2+\sqrt{3}\right) - \tfrac{\pi}{6} - \tfrac{1}{8}\sqrt{3} - \sqrt{3}\right) - 0\right] \\ &= 2\pi\ln\left(2+\sqrt{3}\right) - \tfrac{1}{6}\pi^2 - \tfrac{1}{8}\pi\sqrt{3}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

73. $s = f(t) = \int_0^t \sin \omega u \, \cos^2 \omega u \, du$. Let $y = \cos \omega u \quad \Rightarrow \quad dy = -\omega \sin \omega u \, du$. Then

$s = -\frac{1}{\omega} \int_1^{\cos \omega t} y^2 dy = -\frac{1}{\omega} \left[\frac{1}{3} y^3\right]_1^{\cos \omega t} = \frac{1}{3\omega}(1 - \cos^3 \omega t)$.

74. (a) We want to calculate the square root of the average value of $[E(t)]^2 = [155 \sin(120\pi t)]^2 = 155^2 \sin^2(120\pi t)$. First, we calculate the average value itself, by integrating $[E(t)]^2$ over one cycle (between $t = 0$ and $t = \frac{1}{60}$, since there are 60 cycles per second) and dividing by $\left(\frac{1}{60} - 0\right)$:

$$[E(t)]^2_{\text{avg}} = \frac{1}{1/60} \int_0^{1/60} [155^2 \sin^2(120\pi t)] \, dt = 60 \cdot 155^2 \int_0^{1/60} \tfrac{1}{2}[1 - \cos(240\pi t)] \, dt$$

$$= 60 \cdot 155^2 \left(\tfrac{1}{2}\right) \left[t - \tfrac{1}{240\pi} \sin(240\pi t)\right]_0^{1/60} = 60 \cdot 155^2 \left(\tfrac{1}{2}\right) \left[\left(\tfrac{1}{60} - 0\right) - (0 - 0)\right] = \tfrac{155^2}{2}$$

The RMS value is just the square root of this quantity, which is $\frac{155}{\sqrt{2}} \approx 110$ V.

(b) $220 = \sqrt{[E(t)]^2_{\text{avg}}} \quad \Rightarrow$

$$220^2 = [E(t)]^2_{\text{avg}} = \frac{1}{1/60} \int_0^{1/60} A^2 \sin^2(120\pi t) \, dt = 60A^2 \int_0^{1/60} \tfrac{1}{2}[1 - \cos(240\pi t)] \, dt$$

$$= 30A^2 \left[t - \tfrac{1}{240\pi} \sin(240\pi t)\right]_0^{1/60} = 30A^2 \left[\left(\tfrac{1}{60} - 0\right) - (0 - 0)\right] = \tfrac{1}{2} A^2$$

Thus, $220^2 = \frac{1}{2} A^2 \quad \Rightarrow \quad A = 220\sqrt{2} \approx 311$ V.

75. Just note that the integrand is odd $[f(-x) = -f(x)]$.

Or: If $m \neq n$, calculate

$$\int_{-\pi}^{\pi} \sin mx \, \cos nx \, dx = \int_{-\pi}^{\pi} \tfrac{1}{2}[\sin(m-n)x + \sin(m+n)x] \, dx = \tfrac{1}{2}\left[-\frac{\cos(m-n)x}{m-n} - \frac{\cos(m+n)x}{m+n}\right]_{-\pi}^{\pi} = 0$$

If $m = n$, then the first term in each set of brackets is zero.

76. $\int_{-\pi}^{\pi} \sin mx \, \sin nx \, dx = \int_{-\pi}^{\pi} \frac{1}{2}[\cos(m-n)x - \cos(m+n)x] \, dx$.

If $m \neq n$, this is equal to $\dfrac{1}{2}\left[\dfrac{\sin(m-n)x}{m-n} - \dfrac{\sin(m+n)x}{m+n}\right]_{-\pi}^{\pi} = 0$.

If $m = n$, we get $\int_{-\pi}^{\pi} \frac{1}{2}[1 - \cos(m+n)x] \, dx = \left[\frac{1}{2}x\right]_{-\pi}^{\pi} - \left[\dfrac{\sin(m+n)x}{2(m+n)}\right]_{-\pi}^{\pi} = \pi - 0 = \pi$.

77. $\int_{-\pi}^{\pi} \cos mx \, \cos nx \, dx = \int_{-\pi}^{\pi} \frac{1}{2}[\cos(m-n)x + \cos(m+n)x] \, dx$.

If $m \neq n$, this is equal to $\dfrac{1}{2}\left[\dfrac{\sin(m-n)x}{m-n} + \dfrac{\sin(m+n)x}{m+n}\right]_{-\pi}^{\pi} = 0$.

If $m = n$, we get $\int_{-\pi}^{\pi} \frac{1}{2}[1 + \cos(m+n)x] \, dx = \left[\frac{1}{2}x\right]_{-\pi}^{\pi} + \left[\dfrac{\sin(m+n)x}{2(m+n)}\right]_{-\pi}^{\pi} = \pi + 0 = \pi$.

78. $\dfrac{1}{\pi} \displaystyle\int_{-\pi}^{\pi} f(x) \, \sin mx \, dx = \dfrac{1}{\pi} \int_{-\pi}^{\pi} \left[\left(\sum_{n=1}^{m} a_n \sin nx\right) \sin mx\right] dx = \sum_{n=1}^{m} \dfrac{a_n}{\pi} \int_{-\pi}^{\pi} \sin mx \, \sin nx \, dx$. By Exercise 76, every term is zero except the mth one, and that term is $\dfrac{a_m}{\pi} \cdot \pi = a_m$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7.3 Trigonometric Substitution

1. (a) Use $x = \tan\theta$, where $-\pi/2 < \theta < \pi/2$, since the integrand contains the expression $\sqrt{1+x^2}$.

(b) $x = \tan\theta \quad\Rightarrow\quad dx = \sec^2\theta\, d\theta$ and $\sqrt{1+x^2} = \sqrt{1+\tan^2\theta} = \sqrt{\sec^2\theta} = |\sec\theta| = \sec\theta$.

Then $\displaystyle\int \frac{x^3}{\sqrt{1+x^2}}\,dx = \int \frac{\tan^3\theta}{\sec\theta}\sec^2\theta\,d\theta = \int \tan^3\theta\sec\theta\,d\theta.$

2. (a) Use $x = 3\sin\theta$, where $-\pi/2 \le \theta \le \pi/2$, since the integrand contains the expression $\sqrt{3^2 - x^2}$.

(b) $x = 3\sin\theta \quad\Rightarrow\quad dx = 3\cos\theta\,d\theta$ and

$$\begin{aligned}\sqrt{9-x^2} &= \sqrt{9-9\sin^2\theta} = \sqrt{9(1-\sin^2\theta)} = \sqrt{9\cos^2\theta}\\ &= 3\,|\cos\theta| = 3\cos\theta\end{aligned}$$

Then $\displaystyle\int \frac{x^3}{\sqrt{9-x^2}}\,dx = \int \frac{27\sin^3\theta}{3\cos\theta}\,3\cos\theta\,d\theta = \int 27\sin^3\theta\,d\theta.$

3. (a) Use $x = \sqrt{2}\sec\theta$, where $0 < \theta < \pi/2$ or $\pi < \theta < 3\pi/2$, since the integrand contains the expression $\sqrt{x^2-2}$.

(b) $x = \sqrt{2}\sec\theta \quad\Rightarrow\quad dx = \sqrt{2}\sec\theta\tan\theta\,d\theta$ and

$\sqrt{x^2-2} = \sqrt{2\sec^2\theta - 2} = \sqrt{2(\sec^2\theta - 1)} = \sqrt{2\tan^2\theta} = \sqrt{2}\,|\tan\theta| = \sqrt{2}\tan\theta.$

Then $\displaystyle\int \frac{x^2}{\sqrt{x^2-2}}\,dx = \int \frac{2\sec^2\theta}{\sqrt{2}\tan\theta}\sqrt{2}\sec\theta\tan\theta\,d\theta = \int 2\sec^3\theta\,d\theta.$

4. (a) Use $x = \frac{3}{2}\sin\theta$, where $-\pi/2 \le \theta \le \pi/2$, since the integrand contains the expression

$(9-4x^2)^{3/2} = 4^{3/2}\left(\frac{9}{4} - x^2\right)^{3/2}.$

(b) $x = \frac{3}{2}\sin\theta \quad\Rightarrow\quad dx = \frac{3}{2}\cos\theta\,d\theta$ and

$(9-4x^2)^{3/2} = \left(\sqrt{9-9\sin^2\theta}\right)^3 = \left(\sqrt{9(1-\sin^2\theta)}\right)^3 = \left(\sqrt{9\cos^2\theta}\right)^3 = \left(3\,|\cos\theta|\right)^3 = 27\cos^3\theta.$

Then $\displaystyle\int \frac{x^3}{(9-4x^2)^{3/2}}\,dx = \int \frac{\frac{27}{8}\sin^3\theta}{27\cos^3\theta}\left(\frac{3}{2}\cos\theta\,d\theta\right) = \int \frac{3}{16}\sin^3\theta\sec^2\theta\,d\theta.$

5. Let $x = \sin\theta$, where $-\pi/2 \le \theta \le \pi/2$. Then $dx = \cos\theta\,d\theta$ and

$\sqrt{1-x^2} = \sqrt{1-\sin^2\theta} = \sqrt{\cos^2\theta} = |\cos\theta| = \cos\theta$. Thus,

$$\begin{aligned}\int \frac{x^3}{\sqrt{1-x^2}}\,dx &= \int \frac{\sin^3\theta}{\cos\theta}\cos\theta\;d\theta = \int (1-\cos^2\theta)\sin\theta\,d\theta\\ &\overset{c}{=} \int (1-u^2)\,(-du) = \int (-1+u^2)\,du = -u + \frac{1}{3}u^3 + C\\ &= -\cos\theta + \tfrac{1}{3}\cos^3\theta + C = -\sqrt{1-x^2} + \tfrac{1}{3}\left(\sqrt{1-x^2}\right)^3 + C\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. Let $x = 3\tan\theta$, where $-\pi/2 < \theta < \pi/2$. Then $dx = 3\sec^2\theta\, d\theta$ and

$\sqrt{9+x^2} = \sqrt{9+9\tan^2\theta} = \sqrt{9\left(1+\tan^2\theta\right)} = \sqrt{9\sec^2\theta} = 3\,|\sec\theta| = 3\sec\theta$. Thus,

$$\begin{aligned}
\int \frac{x^3}{\sqrt{9+x^2}}\,dx &= \int \frac{27\tan^3\theta}{3\sec\theta}\left(3\sec^2\theta\, d\theta\right) = \int 27\tan^3\theta\sec\theta\, d\theta \\
&= 27\int\left(\sec^2\theta - 1\right)\tan\theta\sec\theta\, d\theta \\
&= 27\int(u^2-1)\,du \qquad \left[u = \sec\theta,\ du = \sec\theta\tan\theta\, d\theta\right] \\
&= 27\left(\tfrac{1}{3}u^3 - u\right) + C = 9\sec^3\theta - 27\sec\theta + C \\
&= 9\left(\frac{\sqrt{x^2+9}}{3}\right)^3 - \frac{27\sqrt{x^2+9}}{3} + C = \frac{1}{3}(x^2+9)^{3/2} - 9\sqrt{x^2+9} + C
\end{aligned}$$

7. Let $x = \frac{5}{2}\sec\theta$, where $0 \le \theta \le \frac{\pi}{2}$ or $\pi \le \theta < \frac{3\pi}{2}$. Then $dx = \frac{5}{2}\sec\theta\tan\theta\, d\theta$ and $\sqrt{4x^2-25} = \sqrt{25\sec^2\theta - 25} = \sqrt{25\tan^2\theta} = 5\,|\tan\theta| = 5\tan\theta$ for the relevant values of θ, so

$$\begin{aligned}
\int \frac{\sqrt{4x^2-25}}{x}\,dx &= \int \frac{5\tan\theta}{\frac{5}{2}\sec\theta}\left(\frac{5}{2}\sec\theta\tan\theta\, d\theta\right) = 5\int \tan^2\theta\, d\theta \\
&= 5(\tan\theta - \theta) + C \qquad \left[\text{by Exercise 7.1.59 or integration by parts}\right] \\
&= 5\left(\frac{\sqrt{4x^2-25}}{5} - \sec^{-1}\left(\frac{2x}{5}\right)\right) + C = \sqrt{4x^2-25} - 5\sec^{-1}\left(\frac{2}{5}x\right) + C
\end{aligned}$$

8. Let $x = \sqrt{2}\,\sin\theta$, where $-\pi/2 \le \theta \le \pi/2$. Then $dx = \sqrt{2}\cos\theta\, d\theta$ and

$\sqrt{2-x^2} = \sqrt{2-2\sin^2\theta} = \sqrt{2\cos^2\theta} = \sqrt{2}\,|\cos\theta| = \sqrt{2}\cos\theta$. Thus,

$$\begin{aligned}
\int \frac{\sqrt{2-x^2}}{x^2}\,dx &= \int \frac{\sqrt{2}\cos\theta}{2\sin^2\theta}\sqrt{2}\,\cos\theta\, d\theta = \int \frac{\cos^2\theta}{\sin^2\theta}d\theta = \int \cot^2\theta\, d\theta \\
&= \int(\csc^2\theta - 1)\,d\theta = -\cot\theta - \theta + C \\
&= -\frac{\sqrt{2-x^2}}{x} - \sin^{-1}\left(\frac{x}{\sqrt{2}}\right) + C
\end{aligned}$$

9. Let $x = 4\tan\theta$, where $-\pi/2 < \theta < \pi/2$. Then $dx = 4\sec^2\theta\, d\theta$ and

$\sqrt{16+x^2} = \sqrt{16+16\tan^2\theta} = \sqrt{16\sec^2\theta} = 4\,|\sec\theta| = 4\sec\theta$. Thus,

$$\begin{aligned}
\int x^3\sqrt{16+x^2}\,dx &= \int 64\tan^3\theta\,(4\sec\theta)(4\sec^2\theta\, d\theta) = 1024\int \tan^3\theta\sec^3\theta\, d\theta \\
&= 1024\int \tan^2\theta\sec^2\theta\sec\theta\tan\theta\, d\theta = 1024\int(\sec^2\theta - 1)\,\sec^2\theta\sec\theta\,\tan\theta\, d\theta \\
&= 1024\int(u^2-1)u^2\,du \qquad \left[u = \sec\theta,\ du = \sec\theta\,\tan\theta\, d\theta\right] \\
&= 1024\left(\frac{1}{5}u^5 - \frac{1}{3}u^3\right) + C = \frac{1024}{5}\left(\frac{\sqrt{16+x^2}}{4}\right)^5 - \frac{1024}{3}\left(\frac{\sqrt{16+x^2}}{4}\right)^3 + C \\
&= \left(16+x^2\right)^{3/2}\left(\frac{1}{5}\left(16+x^2\right) - \frac{16}{3}\right) + C = \left(16+x^2\right)^{3/2}\left(\frac{1}{5}x^2 - \frac{32}{15}\right) + C
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. Let $x = 3\sin\theta$, where $-\pi/2 \le \theta \le \pi/2$. Then $dx = 3\cos\theta\, d\theta$ and $\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3\,|\cos\theta| = 3\cos\theta$.

$$\int \frac{x^2}{\sqrt{9-x^2}}\,dx = \int \frac{9\sin^2\theta}{3\cos\theta}\,3\cos\theta\,d\theta = 9\int \sin^2\theta\,d\theta$$

$$= 9\int \tfrac{1}{2}(1-\cos 2\theta)\,d\theta = \tfrac{9}{2}\left(\theta - \tfrac{1}{2}\sin 2\theta\right) + C = \tfrac{9}{2}\theta - \tfrac{9}{4}(2\sin\theta\,\cos\theta) + C$$

$$= \frac{9}{2}\sin^{-1}\left(\frac{x}{3}\right) - \frac{9}{2}\cdot\frac{x}{3}\cdot\frac{\sqrt{9-x^2}}{3} + C = \frac{9}{2}\sin^{-1}\left(\frac{x}{3}\right) - \frac{1}{2}x\sqrt{9-x^2} + C$$

11. Let $x = \sec\theta$, where $0 \le \theta \le \frac{\pi}{2}$ or $\pi \le \theta < \frac{3\pi}{2}$. Then $dx = \sec\theta\,\tan\theta\,d\theta$ and $\sqrt{x^2-1} = \sqrt{\sec^2\theta - 1} = \sqrt{\tan^2\theta} = |\tan\theta| = \tan\theta$ for the relevant values of θ, so

$$\int \frac{\sqrt{x^2-1}}{x^4}\,dx = \int \frac{\tan\theta}{\sec^4\theta}\,\sec\theta\,\tan\theta\,d\theta = \int \tan^2\theta\,\cos^3\theta\,d\theta$$

$$= \int \sin^2\theta\,\cos\theta\,d\theta \overset{s}{=} \int u^2\,du = \tfrac{1}{3}u^3 + C = \tfrac{1}{3}\sin^3\theta + C$$

$$= \frac{1}{3}\left(\frac{\sqrt{x^2-1}}{x}\right)^3 + C = \frac{1}{3}\frac{(x^2-1)^{3/2}}{x^3} + C$$

12. Let $u = 36 - x^2$, so $du = -2x\,dx$. When $x = 0$, $u = 36$; when $x = 3$, $u = 27$. Thus,

$$\int_0^3 \frac{x}{\sqrt{36-x^2}}\,dx = \int_{36}^{27} \frac{1}{\sqrt{u}}\left(-\frac{1}{2}\,du\right) = -\frac{1}{2}\Big[2\sqrt{u}\Big]_{36}^{27} = -\left(\sqrt{27} - \sqrt{36}\right) = 6 - 3\sqrt{3}$$

Another method: Let $x = 6\sin\theta$, so $dx = 6\cos\theta\,d\theta$, $x = 0 \;\Rightarrow\; \theta = 0$, and $x = 3 \;\Rightarrow\; \theta = \frac{\pi}{6}$. Then

$$\int_0^3 \frac{x}{\sqrt{36-x^2}}\,dx = \int_0^{\pi/6} \frac{6\sin\theta}{\sqrt{36(1-\sin^2\theta)}}\,6\cos\theta\,d\theta = \int_0^{\pi/6} \frac{6\sin\theta}{6\cos\theta}\,6\cos\theta\,d\theta = 6\int_0^{\pi/6} \sin\theta\,d\theta$$

$$= 6\Big[-\cos\theta\Big]_0^{\pi/6} = 6\left(-\tfrac{\sqrt{3}}{2} + 1\right) = 6 - 3\sqrt{3}$$

13. Let $x = a\tan\theta$, where $a > 0$ and $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$. Then $dx = a\,\sec^2\theta\,d\theta$, $x = 0 \;\Rightarrow\; \theta = 0$, and $x = a \;\Rightarrow\; \theta = \frac{\pi}{4}$. Thus,

$$\int_0^a \frac{dx}{(a^2+x^2)^{3/2}} = \int_0^{\pi/4} \frac{a\sec^2\theta\,d\theta}{[a^2(1+\tan^2\theta)]^{3/2}} = \int_0^{\pi/4} \frac{a\sec^2\theta\,d\theta}{a^3\sec^3\theta} = \frac{1}{a^2}\int_0^{\pi/4} \cos\theta\,d\theta$$

$$= \frac{1}{a^2}\Big[\sin\theta\Big]_0^{\pi/4} = \frac{1}{a^2}\left(\frac{\sqrt{2}}{2} - 0\right) = \frac{1}{\sqrt{2}\,a^2}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. Let $t = 4\sec\theta$, where $0 \le \theta < \frac{\pi}{2}$ or $\pi \le \theta < \frac{3\pi}{2}$. Then $dt = 4\sec\theta\,\tan\theta\,d\theta$ and $\sqrt{t^2-16} = \sqrt{16\sec^2\theta - 16} = \sqrt{16\tan^2\theta} = 4\tan\theta$ for the relevant values of θ, so

$$\int \frac{dt}{t^2\sqrt{t^2-16}} = \int \frac{4\sec\theta\,\tan\theta\,d\theta}{16\sec^2\theta\cdot 4\tan\theta} = \frac{1}{16}\int \frac{1}{\sec\theta}\,d\theta = \frac{1}{16}\int \cos\theta\,d\theta$$

$$= \frac{1}{16}\sin\theta + C = \frac{1}{16}\frac{\sqrt{t^2-16}}{t} + C = \frac{\sqrt{t^2-16}}{16t} + C$$

15. Let $x = \sec\theta$, so $dx = \sec\theta\,\tan\theta\,d\theta$, $x = 2 \;\Rightarrow\; \theta = \frac{\pi}{3}$, and $x = 3 \;\Rightarrow\; \theta = \sec^{-1}3$. Then

$$\int_2^3 \frac{dx}{(x^2-1)^{3/2}} = \int_{\pi/3}^{\sec^{-1}3} \frac{\sec\theta\,\tan\theta\,d\theta}{\tan^3\theta} = \int_{\pi/3}^{\sec^{-1}3} \frac{\cos\theta}{\sin^2\theta}\,d\theta$$

$$\overset{s}{=} \int_{\sqrt{3}/2}^{\sqrt{8}/3} \frac{1}{u^2}\,du = \left[-\frac{1}{u}\right]_{\sqrt{3}/2}^{\sqrt{8}/3} = \frac{-3}{\sqrt{8}} + \frac{2}{\sqrt{3}} = -\frac{3}{4}\sqrt{2} + \frac{2}{3}\sqrt{3}$$

16. Let $x = \frac{2}{3}\sin\theta$, so $dx = \frac{2}{3}\cos\theta\,d\theta$, $x = 0 \;\Rightarrow\; \theta = 0$, and $x = \frac{2}{3} \;\Rightarrow\; \theta = \frac{\pi}{2}$. Thus,

$$\int_0^{2/3} \sqrt{4-9x^2}\,dx = \int_0^{\pi/2} \sqrt{4 - 9\cdot\frac{4}{9}\sin^2\theta}\;\frac{2}{3}\cos\theta\,d\theta$$

$$= \int_0^{\pi/2} 2\cos\theta\cdot\frac{2}{3}\cos\theta\,d\theta = \frac{4}{3}\int_0^{\pi/2}\cos^2\theta\,d\theta$$

$$= \frac{4}{3}\int_0^{\pi/2}\frac{1}{2}(1+\cos 2\theta)\,d\theta = \frac{2}{3}\left[\theta + \frac{1}{2}\sin 2\theta\right]_0^{\pi/2} = \frac{2}{3}\left[\left(\frac{\pi}{2}+0\right) - (0+0)\right] = \frac{\pi}{3}$$

17. $$\int_0^{1/2} x\sqrt{1-4x^2}\,dx = \int_1^0 u^{1/2}\left(-\frac{1}{8}\,du\right) \qquad \begin{bmatrix} u = 1-4x^2, \\ du = -8x\,dx \end{bmatrix}$$

$$= \tfrac{1}{8}\left[\tfrac{2}{3}u^{3/2}\right]_0^1 = \tfrac{1}{12}(1-0) = \tfrac{1}{12}$$

18. Let $t = 2\tan\theta$, so $dt = 2\sec^2\theta\,d\theta$, $t = 0 \;\Rightarrow\; \theta = 0$, and $t = 2 \;\Rightarrow\; \theta = \frac{\pi}{4}$. Thus,

$$\int_0^2 \frac{dt}{\sqrt{4+t^2}} = \int_0^{\pi/4} \frac{2\sec^2\theta\,d\theta}{\sqrt{4+4\tan^2\theta}} = \int_0^{\pi/4} \frac{2\sec^2\theta\,d\theta}{2\sec\theta} = \int_0^{\pi/4}\sec\theta\,d\theta = \Big[\ln|\sec\theta + \tan\theta|\Big]_0^{\pi/4}$$

$$= \ln\left|\sqrt{2}+1\right| - \ln|1+0| = \ln(\sqrt{2}+1)$$

19. Let $x = 3\sec\theta$, where $0 \le \theta < \frac{\pi}{2}$ or $\pi \le \theta < \frac{3\pi}{2}$. Then $dx = 3\sec\theta\,\tan\theta\,d\theta$ and $\sqrt{x^2-9} = 3\tan\theta$, so

$$\int \frac{\sqrt{x^2-9}}{x^3}\,dx = \int \frac{3\tan\theta}{27\sec^3\theta}\,3\sec\theta\,\tan\theta\,d\theta = \frac{1}{3}\int \frac{\tan^2\theta}{\sec^2\theta}\,d\theta$$

$$= \tfrac{1}{3}\int \sin^2\theta\,d\theta = \tfrac{1}{3}\int \tfrac{1}{2}(1-\cos 2\theta)\,d\theta = \tfrac{1}{6}\theta - \tfrac{1}{12}\sin 2\theta + C = \tfrac{1}{6}\theta - \tfrac{1}{6}\sin\theta\,\cos\theta + C$$

$$= \frac{1}{6}\sec^{-1}\left(\frac{x}{3}\right) - \frac{1}{6}\frac{\sqrt{x^2-9}}{x}\,\frac{3}{x} + C = \frac{1}{6}\sec^{-1}\left(\frac{x}{3}\right) - \frac{\sqrt{x^2-9}}{2x^2} + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. Let $x = \tan\theta$, so $dx = \sec^2\theta\, d\theta$, $x = 0 \;\Rightarrow\; \theta = 0$, and $x = 1 \;\Rightarrow\; \theta = \frac{\pi}{4}$. Then

$$\begin{aligned}\int_0^1 \frac{dx}{(x^2+1)^2} &= \int_0^{\pi/4} \frac{\sec^2\theta\, d\theta}{(\tan^2\theta+1)^2} = \int_0^{\pi/4} \frac{\sec^2\theta\, d\theta}{(\sec^2\theta)^2} \\ &= \int_0^{\pi/4} \cos^2\theta\, d\theta = \int_0^{\pi/4} \frac{1}{2}\left(1+\cos 2\theta\right)\, d\theta \\ &= \tfrac{1}{2}\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/4} = \tfrac{1}{2}\left[\left(\tfrac{\pi}{4}+\tfrac{1}{2}\right) - 0\right] = \tfrac{\pi}{8} + \tfrac{1}{4}\end{aligned}$$

21. Let $x = a\sin\theta$, $dx = a\cos\theta\, d\theta$, $x = 0 \;\Rightarrow\; \theta = 0$ and $x = a \;\Rightarrow\; \theta = \frac{\pi}{2}$. Then

$$\begin{aligned}\int_0^a x^2\sqrt{a^2-x^2}\, dx &= \int_0^{\pi/2} a^2\sin^2\theta\,(a\cos\theta)\, a\cos\theta\, d\theta = a^4\int_0^{\pi/2} \sin^2\theta\,\cos^2\theta\, d\theta \\ &= a^4\int_0^{\pi/2} \left[\tfrac{1}{2}(2\sin\theta\,\cos\theta)\right]^2\, d\theta = \frac{a^4}{4}\int_0^{\pi/2} \sin^2 2\theta\, d\theta = \frac{a^4}{4}\int_0^{\pi/2} \tfrac{1}{2}(1-\cos 4\theta)\, d\theta \\ &= \frac{a^4}{8}\left[\theta - \tfrac{1}{4}\sin 4\theta\right]_0^{\pi/2} = \frac{a^4}{8}\left[\left(\frac{\pi}{2}-0\right)-0\right] = \frac{\pi}{16}a^4\end{aligned}$$

22. Let $x = \dfrac{1}{2}\sin\theta$, so $dx = \dfrac{1}{2}\cos\theta\, d\theta$, $x = \dfrac{1}{4} \;\Rightarrow\; \theta = \dfrac{\pi}{6}$, and $x = \dfrac{\sqrt{3}}{4} \;\Rightarrow\; \theta = \dfrac{\pi}{3}$. Then

$$\begin{aligned}\int_{1/4}^{\sqrt{3}/4} \sqrt{1-4x^2}\, dx &= \int_{\pi/6}^{\pi/3} \sqrt{1-\sin^2\theta}\left(\frac{1}{2}\cos\theta\, d\theta\right) = \frac{1}{2}\int_{\pi/6}^{\pi/3} \cos^2\theta\, d\theta = \frac{1}{2}\int_{\pi/6}^{\pi/3} \frac{1}{2}(1+\cos 2\theta)\, d\theta \\ &= \frac{1}{4}\left[\theta + \frac{1}{2}\sin 2\theta\right]_{\pi/6}^{\pi/3} = \frac{1}{4}\left[\left(\frac{\pi}{3}+\frac{1}{2}\sin\frac{2\pi}{3}\right) - \left(\frac{\pi}{6}+\frac{1}{2}\sin\frac{\pi}{3}\right)\right] \\ &= \frac{1}{4}\left[\left(\frac{\pi}{3}+\frac{\sqrt{3}}{4}\right) - \left(\frac{\pi}{6}+\frac{\sqrt{3}}{4}\right)\right] = \frac{\pi}{24}\end{aligned}$$

23. Let $u = x^2 - 7$, so $du = 2x\, dx$. Then $\displaystyle\int \frac{x}{\sqrt{x^2-7}}\, dx = \frac{1}{2}\int \frac{1}{\sqrt{u}}\, du = \tfrac{1}{2}\cdot 2\sqrt{u} + C = \sqrt{x^2-7} + C$.

24. Let $u = 1 + x^2$, so $du = 2x\, dx$. Then

$$\int \frac{x}{\sqrt{1+x^2}}\, dx = \int \frac{1}{\sqrt{u}}\left(\frac{1}{2}\, du\right) = \frac{1}{2}\int u^{-1/2}\, du = \frac{1}{2}\cdot 2u^{1/2} + C = \sqrt{1+x^2} + C$$

25. Let $x = \tan\theta$, where $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$. Then $dx = \sec^2\theta\, d\theta$ and $\sqrt{1+x^2} = \sec\theta$, so

$$\begin{aligned}\int \frac{\sqrt{1+x^2}}{x}\, dx &= \int \frac{\sec\theta}{\tan\theta}\sec^2\theta\, d\theta = \int \frac{\sec\theta}{\tan\theta}\left(1+\tan^2\theta\right) d\theta \\ &= \int (\csc\theta + \sec\theta\,\tan\theta)\, d\theta \\ &= \ln|\csc\theta - \cot\theta| + \sec\theta + C \qquad \text{[by Exercise 7.2.41]} \\ &= \ln\left|\frac{\sqrt{1+x^2}}{x} - \frac{1}{x}\right| + \frac{\sqrt{1+x^2}}{1} + C = \ln\left|\frac{\sqrt{1+x^2}-1}{x}\right| + \sqrt{1+x^2} + C\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. Let $x = \frac{3}{5}\sin\theta$, so $dx = \frac{3}{5}\cos\theta\,d\theta$, $x = 0 \;\Rightarrow\; \theta = 0$, and $x = 0.3 \;\Rightarrow\; \theta = \frac{\pi}{6}$. Then

$$\int_0^{0.3} \frac{x}{(9-25x^2)^{3/2}}\,dx = \int_0^{\pi/6} \frac{\frac{3}{5}\sin\theta}{\left(\sqrt{9-9\sin^2\theta}\right)^3}\left(\tfrac{3}{5}\cos\theta\,d\theta\right)$$

$$= \frac{9}{25}\int_0^{\pi/6} \frac{\sin\theta}{(3\cos\theta)^3}\cos\theta\,d\theta = \frac{1}{75}\int_0^{\pi/6} \frac{\sin\theta}{\cos^2\theta}\,d\theta \overset{c}{=} -\frac{1}{75}\int_1^{\sqrt{3}/2} \frac{1}{u^2}\,du$$

$$= -\frac{1}{75}\left[-\frac{1}{u}\right]_1^{\sqrt{3}/2} = \frac{1}{75}\left(\frac{2}{\sqrt{3}} - 1\right) = \frac{2\sqrt{3}}{225} - \frac{1}{75}$$

27. Let $x = \frac{3}{5}\sin\theta$, so $dx = \frac{3}{5}\cos\theta\,d\theta$, $x = 0 \;\Rightarrow\; \theta = 0$, and $x = 0.6 \;\Rightarrow\; \theta = \frac{\pi}{2}$. Then

$$\int_0^{0.6} \frac{x^2}{\sqrt{9-25x^2}}\,dx = \int_0^{\pi/2} \frac{\left(\frac{3}{5}\right)^2\sin^2\theta}{3\cos\theta}\left(\tfrac{3}{5}\cos\theta\,d\theta\right) = \tfrac{9}{125}\int_0^{\pi/2} \sin^2\theta\,d\theta$$

$$= \tfrac{9}{125}\int_0^{\pi/2} \tfrac{1}{2}(1-\cos 2\theta)\,d\theta = \tfrac{9}{250}\left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/2}$$

$$= \tfrac{9}{250}\left[\left(\tfrac{\pi}{2} - 0\right) - 0\right] = \tfrac{9}{500}\pi$$

28. Let $x = \tan\theta$, where $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$. Then $dx = \sec^2\theta\,d\theta$,

$\sqrt{x^2+1} = \sec\theta$ and $x = 0 \;\Rightarrow\; \theta = 0$, $x = 1 \;\Rightarrow\; \theta = \frac{\pi}{4}$, so

$$\int_0^1 \sqrt{x^2+1}\,dx = \int_0^{\pi/4} \sec\theta\,\sec^2\theta\,d\theta = \int_0^{\pi/4} \sec^3\theta\,d\theta$$

$$= \tfrac{1}{2}\Big[\sec\theta\,\tan\theta + \ln|\sec\theta + \tan\theta|\Big]_0^{\pi/4} \qquad \text{[by Example 7.2.8]}$$

$$= \tfrac{1}{2}\left[\sqrt{2}\cdot 1 + \ln\left(1+\sqrt{2}\right) - 0 - \ln(1+0)\right] = \tfrac{1}{2}\left[\sqrt{2} + \ln\left(1+\sqrt{2}\right)\right]$$

29. $$\int \frac{dx}{\sqrt{x^2+2x+5}} = \int \frac{dx}{\sqrt{(x+1)^2+4}} = \int \frac{2\sec^2\theta\,d\theta}{\sqrt{4\tan^2\theta+4}} \qquad \left[\begin{array}{l} x+1 = 2\tan\theta, \\ dx = 2\sec^2\theta\,d\theta \end{array}\right]$$

$$= \int \frac{2\sec^2\theta\,d\theta}{2\sec\theta} = \int \sec\theta\,d\theta = \ln|\sec\theta + \tan\theta| + C_1$$

$$= \ln\left|\frac{\sqrt{x^2+2x+5}}{2} + \frac{x+1}{2}\right| + C_1,$$

or $\ln\left|\sqrt{x^2+2x+5} + x + 1\right| + C$, where $C = C_1 - \ln 2$.

30. $$\int_0^1 \sqrt{x-x^2}\,dx = \int_0^1 \sqrt{\tfrac{1}{4} - \left(x^2 - x + \tfrac{1}{4}\right)}\,dx = \int_0^1 \sqrt{\tfrac{1}{4} - \left(x - \tfrac{1}{2}\right)^2}\,dx$$

$$= \int_{-\pi/2}^{\pi/2} \sqrt{\tfrac{1}{4} - \tfrac{1}{4}\sin^2\theta}\;\tfrac{1}{2}\cos\theta\,d\theta \qquad \left[\begin{array}{l} x - \frac{1}{2} = \frac{1}{2}\sin\theta, \\ dx = \frac{1}{2}\cos\theta\,d\theta \end{array}\right]$$

$$= 2\int_0^{\pi/2} \tfrac{1}{2}\cos\theta\,\tfrac{1}{2}\cos\theta\,d\theta = \tfrac{1}{2}\int_0^{\pi/2} \cos^2\theta\,d\theta = \tfrac{1}{2}\int_0^{\pi/2} \tfrac{1}{2}(1+\cos 2\theta)\,d\theta$$

$$= \tfrac{1}{4}\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/2} = \tfrac{1}{4}\left(\tfrac{\pi}{2}\right) = \tfrac{\pi}{8}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. $\int x^2\sqrt{3+2x-x^2}\,dx = \int x^2\sqrt{4-(x^2+2x+1)}\,dx = \int x^2\sqrt{2^2-(x-1)^2}\,dx$

$$= \int (1+2\sin\theta)^2\sqrt{4\cos^2\theta}\,2\cos\theta\,d\theta \qquad \begin{bmatrix} x-1 = 2\sin\theta, \\ dx = 2\cos\theta\,d\theta \end{bmatrix}$$

$$= \int (1+4\sin\theta+4\sin^2\theta)\,4\cos^2\theta\,d\theta$$

$$= 4\int(\cos^2\theta + 4\sin\theta\,\cos^2\theta + 4\sin^2\theta\,\cos^2\theta)\,d\theta$$

$$= 4\int \tfrac{1}{2}(1+\cos 2\theta)\,d\theta + 4\int 4\sin\theta\,\cos^2\theta\,d\theta + 4\int(2\sin\theta\,\cos\theta)^2\,d\theta$$

$$= 2\int(1+\cos 2\theta)\,d\theta + 16\int\sin\theta\,\cos^2\theta\,d\theta + 4\int\sin^2 2\theta\,d\theta$$

$$= 2\big(\theta + \tfrac{1}{2}\sin 2\theta\big) + 16\big(-\tfrac{1}{3}\cos^3\theta\big) + 4\int\tfrac{1}{2}(1-\cos 4\theta)\,d\theta$$

$$= 2\theta + \sin 2\theta - \tfrac{16}{3}\cos^3\theta + 2\big(\theta - \tfrac{1}{4}\sin 4\theta\big) + C$$

$$= 4\theta - \tfrac{1}{2}\sin 4\theta + \sin 2\theta - \tfrac{16}{3}\cos^3\theta + C$$

$$= 4\theta - \tfrac{1}{2}(2\sin 2\theta\,\cos 2\theta) + \sin 2\theta - \tfrac{16}{3}\cos^3\theta + C$$

$$= 4\theta + \sin 2\theta(1-\cos 2\theta) - \tfrac{16}{3}\cos^3\theta + C$$

$$= 4\theta + (2\sin\theta\,\cos\theta)(2\sin^2\theta) - \tfrac{16}{3}\cos^3\theta + C$$

$$= 4\theta + 4\sin^3\theta\,\cos\theta - \tfrac{16}{3}\cos^3\theta + C$$

$$= 4\sin^{-1}\left(\frac{x-1}{2}\right) + 4\left(\frac{x-1}{2}\right)^3\frac{\sqrt{3+2x-x^2}}{2} - \frac{16}{3}\frac{(3+2x-x^2)^{3/2}}{2^3} + C$$

$$= 4\sin^{-1}\left(\frac{x-1}{2}\right) + \frac{1}{4}(x-1)^3\sqrt{3+2x-x^2} - \frac{2}{3}(3+2x-x^2)^{3/2} + C$$

32. $3+4x-4x^2 = -(4x^2-4x+1)+4 = 2^2-(2x-1)^2$.

Let $2x-1 = 2\sin\theta$, so $2\,dx = 2\cos\theta\,d\theta$ and $\sqrt{3+4x-4x^2} = 2\cos\theta$.

Then

$$\int\frac{x^2}{(3+4x-4x^2)^{3/2}}\,dx = \int\frac{\left[\frac{1}{2}(1+2\sin\theta)\right]^2}{(2\cos\theta)^3}\cos\theta\,d\theta$$

$$= \frac{1}{32}\int\frac{1+4\sin\theta+4\sin^2\theta}{\cos^2\theta}\,d\theta = \frac{1}{32}\int(\sec^2\theta + 4\tan\theta\,\sec\theta + 4\tan^2\theta)\,d\theta$$

$$= \tfrac{1}{32}\int[\sec^2\theta + 4\tan\theta\,\sec\theta + 4(\sec^2\theta - 1)]\,d\theta$$

$$= \tfrac{1}{32}\int(5\sec^2\theta + 4\tan\theta\,\sec\theta - 4)\,d\theta = \tfrac{1}{32}(5\tan\theta + 4\sec\theta - 4\theta) + C$$

$$= \frac{1}{32}\left[5\cdot\frac{2x-1}{\sqrt{3+4x-4x^2}} + 4\cdot\frac{2}{\sqrt{3+4x-4x^2}} - 4\cdot\sin^{-1}\left(\frac{2x-1}{2}\right)\right] + C$$

$$= \frac{10x+3}{32\sqrt{3+4x-4x^2}} - \frac{1}{8}\sin^{-1}\left(\frac{2x-1}{2}\right) + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. $x^2+2x=(x^2+2x+1)-1=(x+1)^2-1$. Let $x+1=1\sec\theta$, so $dx=\sec\theta\,\tan\theta\,d\theta$ and $\sqrt{x^2+2x}=\tan\theta$. Then

$$\int\sqrt{x^2+2x}\,dx=\int\tan\theta\,(\sec\theta\,\tan\theta\,d\theta)=\int\tan^2\theta\,\sec\theta\,d\theta$$

$$=\int(\sec^2\theta-1)\,\sec\theta\,d\theta=\int\sec^3\theta\,d\theta-\int\sec\theta\,d\theta$$

$$=\tfrac12\sec\theta\,\tan\theta+\tfrac12\ln|\sec\theta+\tan\theta|-\ln|\sec\theta+\tan\theta|+C$$

$$=\tfrac12\sec\theta\,\tan\theta-\tfrac12\ln|\sec\theta+\tan\theta|+C=\tfrac12(x+1)\sqrt{x^2+2x}-\tfrac12\ln\left|x+1+\sqrt{x^2+2x}\,\right|+C$$

34. $x^2-2x+2=(x^2-2x+1)+1=(x-1)^2+1$. Let $x-1=1\tan\theta$, so $dx=\sec^2\theta\,d\theta$ and $\sqrt{x^2-2x+2}=\sec\theta$. Then

$$\int\frac{x^2+1}{(x^2-2x+2)^2}\,dx=\int\frac{(\tan\theta+1)^2+1}{\sec^4\theta}\,\sec^2\theta\,d\theta$$

$$=\int\frac{\tan^2\theta+2\tan\theta+2}{\sec^2\theta}\,d\theta$$

$$=\int(\sin^2\theta+2\sin\theta\,\cos\theta+2\cos^2\theta)\,d\theta=\int(1+2\sin\theta\,\cos\theta+\cos^2\theta)\,d\theta$$

$$=\int\left[1+2\sin\theta\,\cos\theta+\tfrac12(1+\cos2\theta)\right]\,d\theta=\int\left(\tfrac32+2\sin\theta\,\cos\theta+\tfrac12\cos2\theta\right)\,d\theta$$

$$=\tfrac32\theta+\sin^2\theta+\tfrac14\sin2\theta+C=\tfrac32\theta+\sin^2\theta+\tfrac12\sin\theta\,\cos\theta+C$$

$$=\frac32\tan^{-1}\left(\frac{x-1}{1}\right)+\frac{(x-1)^2}{x^2-2x+2}+\frac12\frac{x-1}{\sqrt{x^2-2x+2}}\frac{1}{\sqrt{x^2-2x+2}}+C$$

$$=\frac32\tan^{-1}(x-1)+\frac{2(x^2-2x+1)+x-1}{2(x^2-2x+2)}+C=\frac32\tan^{-1}(x-1)+\frac{2x^2-3x+1}{2(x^2-2x+2)}+C$$

We can write the answer as

$$\frac32\tan^{-1}(x-1)+\frac{(2x^2-4x+4)+x-3}{2(x^2-2x+2)}+C=\frac32\tan^{-1}(x-1)+1+\frac{x-3}{2(x^2-2x+2)}+C$$

$$=\frac32\tan^{-1}(x-1)+\frac{x-3}{2(x^2-2x+2)}+C_1,\ \text{where } C_1=1+C$$

35. Let $u=x^2$, $du=2x\,dx$. Then

$$\int x\sqrt{1-x^4}\,dx=\int\sqrt{1-u^2}\left(\tfrac12\,du\right)=\tfrac12\int\cos\theta\cdot\cos\theta\,d\theta\qquad\left[\begin{array}{c}\text{where } u=\sin\theta,\ du=\cos\theta\,d\theta,\\ \text{and } \sqrt{1-u^2}=\cos\theta\end{array}\right]$$

$$=\tfrac12\int\tfrac12(1+\cos2\theta)\,d\theta=\tfrac14\theta+\tfrac18\sin2\theta+C=\tfrac14\theta+\tfrac14\sin\theta\,\cos\theta+C$$

$$=\tfrac14\sin^{-1}u+\tfrac14u\sqrt{1-u^2}+C=\tfrac14\sin^{-1}(x^2)+\tfrac14x^2\sqrt{1-x^4}+C$$

36. Let $u=\sin t$, $du=\cos t\,dt$. Then

$$\int_0^{\pi/2}\frac{\cos t}{\sqrt{1+\sin^2t}}\,dt=\int_0^1\frac{1}{\sqrt{1+u^2}}\,du=\int_0^{\pi/4}\frac{1}{\sec\theta}\,\sec^2\theta\,d\theta\qquad\left[\begin{array}{c}\text{where } u=\tan\theta\,,\ du=\sec^2\theta\,d\theta,\\ \text{and } \sqrt{1+u^2}=\sec\theta\end{array}\right]$$

$$=\int_0^{\pi/4}\sec\theta\,d\theta=\Big[\ln|\sec\theta+\tan\theta|\Big]_0^{\pi/4}\qquad\text{[by (1) in Section 7.2]}$$

$$=\ln\left(\sqrt2+1\right)-\ln(1+0)=\ln\left(\sqrt2+1\right)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. (a) Let $x = a\tan\theta$, where $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$. Then $\sqrt{x^2+a^2} = a\sec\theta$ and

$$\int \frac{dx}{\sqrt{x^2+a^2}} = \int \frac{a\sec^2\theta\,d\theta}{a\sec\theta} = \int \sec\theta\,d\theta = \ln|\sec\theta+\tan\theta| + C_1 = \ln\left|\frac{\sqrt{x^2+a^2}}{a} + \frac{x}{a}\right| + C_1$$

$$= \ln\left(x+\sqrt{x^2+a^2}\right) + C \quad \text{where } C = C_1 - \ln|a|$$

(b) Let $x = a\sinh t$, so that $dx = a\cosh t\,dt$ and $\sqrt{x^2+a^2} = a\cosh t$. Then

$$\int \frac{dx}{\sqrt{x^2+a^2}} = \int \frac{a\cosh t\,dt}{a\cosh t} = t + C = \sinh^{-1}\frac{x}{a} + C.$$

38. (a) Let $x = a\tan\theta$, $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$. Then

$$I = \int \frac{x^2}{(x^2+a^2)^{3/2}}\,dx = \int \frac{a^2\tan^2\theta}{a^3\sec^3\theta}\,a\sec^2\theta\,d\theta = \int \frac{\tan^2\theta}{\sec\theta}\,d\theta = \int \frac{\sec^2\theta - 1}{\sec\theta}\,d\theta$$

$$= \int (\sec\theta - \cos\theta)\,d\theta = \ln|\sec\theta+\tan\theta| - \sin\theta + C$$

$$= \ln\left|\frac{\sqrt{x^2+a^2}}{a} + \frac{x}{a}\right| - \frac{x}{\sqrt{x^2+a^2}} + C = \ln\left(x+\sqrt{x^2+a^2}\right) - \frac{x}{\sqrt{x^2+a^2}} + C_1$$

(b) Let $x = a\sinh t$. Then

$$I = \int \frac{a^2\sinh^2 t}{a^3\cosh^3 t}\,a\cosh t\,dt = \int \tanh^2 t\,dt = \int (1-\operatorname{sech}^2 t)\,dt = t - \tanh t + C$$

$$= \sinh^{-1}\left(\frac{x}{a}\right) - \frac{x}{\sqrt{a^2+x^2}} + C$$

39. The average value of $f(x) = \sqrt{x^2-1}/x$ on the interval $[1, 7]$ is

$$\frac{1}{7-1}\int_1^7 \frac{\sqrt{x^2-1}}{x}\,dx = \frac{1}{6}\int_0^\alpha \frac{\tan\theta}{\sec\theta}\cdot\sec\theta\tan\theta\,d\theta \qquad \left[\begin{array}{c}\text{where } x = \sec\theta,\ dx = \sec\theta\tan\theta\,d\theta,\\ \sqrt{x^2-1} = \tan\theta, \text{ and } \alpha = \sec^{-1}7\end{array}\right]$$

$$= \tfrac{1}{6}\int_0^\alpha \tan^2\theta\,d\theta = \tfrac{1}{6}\int_0^\alpha (\sec^2\theta - 1)\,d\theta = \tfrac{1}{6}\left[\tan\theta - \theta\right]_0^\alpha$$

$$= \tfrac{1}{6}(\tan\alpha - \alpha) = \tfrac{1}{6}\left(\sqrt{48} - \sec^{-1}7\right)$$

40. $9x^2 - 4y^2 = 36 \quad\Rightarrow\quad y = \pm\frac{3}{2}\sqrt{x^2-4} \quad\Rightarrow$

$$\text{area} = 2\int_2^3 \tfrac{3}{2}\sqrt{x^2-4}\,dx = 3\int_2^3 \sqrt{x^2-4}\,dx$$

$$= 3\int_0^\alpha 2\tan\theta\,2\sec\theta\,\tan\theta\,d\theta \qquad \left[\begin{array}{c}\text{where } x = 2\sec\theta,\\ dx = 2\sec\theta\,\tan\theta\,d\theta,\\ \alpha = \sec^{-1}\left(\frac{3}{2}\right)\end{array}\right]$$

$$= 12\int_0^\alpha \left(\sec^2\theta - 1\right)\sec\theta\,d\theta = 12\int_0^\alpha \left(\sec^3\theta - \sec\theta\right)d\theta$$

$$= 12\left[\tfrac{1}{2}(\sec\theta\,\tan\theta + \ln|\sec\theta+\tan\theta|) - \ln|\sec\theta+\tan\theta|\right]_0^\alpha$$

$$= 6\Big[\sec\theta\,\tan\theta - \ln|\sec\theta+\tan\theta|\Big]_0^\alpha = 6\left[\tfrac{3\sqrt{5}}{4} - \ln\left(\tfrac{3}{2} + \tfrac{\sqrt{5}}{2}\right)\right] = \tfrac{9\sqrt{5}}{2} - 6\ln\left(\tfrac{3+\sqrt{5}}{2}\right)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. Area of $\triangle POQ = \frac{1}{2}(r\cos\theta)(r\sin\theta) = \frac{1}{2}r^2\sin\theta\,\cos\theta$. Area of region $PQR = \int_{r\cos\theta}^{r}\sqrt{r^2-x^2}\,dx$.

Let $x = r\cos u \quad\Rightarrow\quad dx = -r\sin u\,du$ for $\theta \le u \le \frac{\pi}{2}$. Then we obtain

$$\int\sqrt{r^2-x^2}\,dx = \int r\sin u\,(-r\sin u)\,du = -r^2\int\sin^2 u\,du = -\tfrac{1}{2}r^2(u-\sin u\,\cos u)+C$$
$$= -\tfrac{1}{2}r^2\cos^{-1}(x/r) + \tfrac{1}{2}x\sqrt{r^2-x^2}+C$$

so
$$\text{area of region } PQR = \tfrac{1}{2}\left[-r^2\cos^{-1}(x/r) + x\sqrt{r^2-x^2}\,\right]_{r\cos\theta}^{r}$$
$$= \tfrac{1}{2}\left[0-(-r^2\theta + r\cos\theta\, r\sin\theta)\right] = \tfrac{1}{2}r^2\theta - \tfrac{1}{2}r^2\sin\theta\,\cos\theta$$

and thus, (area of sector POR) = (area of $\triangle POQ$) + (area of region PQR) $= \frac{1}{2}r^2\theta$.

42. Let $x = \sqrt{2}\sec\theta$, where $0 \le \theta < \frac{\pi}{2}$ or $\pi \le \theta < \frac{3\pi}{2}$, so $dx = \sqrt{2}\,\sec\theta\tan\theta\,d\theta$. Then

$$\int\frac{dx}{x^4\sqrt{x^2-2}} = \int\frac{\sqrt{2}\sec\theta\,\tan\theta\,d\theta}{4\sec^4\theta\sqrt{2}\,\tan\theta}$$
$$= \tfrac{1}{4}\int\cos^3\theta\,d\theta = \tfrac{1}{4}\int\left(1-\sin^2\theta\right)\cos\theta\,d\theta$$
$$= \tfrac{1}{4}\left[\sin\theta - \tfrac{1}{3}\sin^3\theta\right]+C \qquad [\text{substitute } u = \sin\theta]$$
$$= \frac{1}{4}\left[\frac{\sqrt{x^2-2}}{x} - \frac{\left(x^2-2\right)^{3/2}}{3x^3}\right]+C$$

From the graph, it appears that our answer is reasonable. [Notice that $f(x)$ is large when F increases rapidly and small when F levels out.]

43. Use disks about the x-axis:

$$V = \int_0^3 \pi\left(\frac{9}{x^2+9}\right)^2 dx = 81\pi\int_0^3\frac{1}{(x^2+9)^2}\,dx$$

Let $x = 3\tan\theta$, so $dx = 3\sec^2\theta\,d\theta$, $x = 0 \quad\Rightarrow\quad \theta = 0$ and $x = 3 \quad\Rightarrow\quad \theta = \frac{\pi}{4}$. Thus,

$$V = 81\pi\int_0^{\pi/4}\frac{1}{(9\sec^2\theta)^2}\,3\sec^2\theta\,d\theta = 3\pi\int_0^{\pi/4}\cos^2\theta\,d\theta = 3\pi\int_0^{\pi/4}\frac{1}{2}(1+\cos 2\theta)\,d\theta$$
$$= \tfrac{3\pi}{2}\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/4} = \tfrac{3\pi}{2}\left[\left(\tfrac{\pi}{4}+\tfrac{1}{2}\right)-0\right] = \tfrac{3}{8}\pi^2 + \tfrac{3}{4}\pi$$

44. Use shells about $x = 1$:

$$V = \int_0^1 2\pi(1-x)\,x\sqrt{1-x^2}\,dx$$
$$= 2\pi\int_0^1 x\sqrt{1-x^2}\,dx - 2\pi\int_0^1 x^2\sqrt{1-x^2}\,dx = 2\pi V_1 - 2\pi V_2$$

For V_1, let $u = 1-x^2$, so $du = -2x\,dx$, and

$$V_1 = \int_1^0\sqrt{u}\,\left(-\tfrac{1}{2}\,du\right) = \tfrac{1}{2}\int_0^1 u^{1/2}\,du = \tfrac{1}{2}\left[\tfrac{2}{3}u^{3/2}\right]_0^1 = \tfrac{1}{2}\left(\tfrac{2}{3}\right) = \tfrac{1}{3}.$$

For V_2, let $x = \sin\theta$, so $dx = \cos\theta\,d\theta$, and

$$V_2 = \int_0^{\pi/2}\sin^2\theta\sqrt{\cos^2\theta}\,\cos\theta\,d\theta = \int_0^{\pi/2}\sin^2\theta\,\cos^2\theta\,d\theta = \int_0^{\pi/2}\tfrac{1}{4}(2\sin\theta\,\cos\theta)^2\,d\theta$$
$$= \tfrac{1}{4}\int_0^{\pi/2}\sin^2 2\theta\,d\theta = \tfrac{1}{4}\int_0^{\pi/2}\tfrac{1}{2}(1-\cos 2\theta)\,d\theta = \tfrac{1}{8}\left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/2} = \tfrac{1}{8}\left(\tfrac{\pi}{2}\right) = \tfrac{\pi}{16}$$

Thus, $V = 2\pi\left(\frac{1}{3}\right) - 2\pi\left(\frac{\pi}{16}\right) = \frac{2}{3}\pi - \frac{1}{8}\pi^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. (a) Let $t = a\sin\theta$, $dt = a\cos\theta\,d\theta$, $t = 0 \;\Rightarrow\; \theta = 0$ and $t = x \;\Rightarrow$ $\theta = \sin^{-1}(x/a)$. Then

$$\int_0^x \sqrt{a^2-t^2}\,dt = \int_0^{\sin^{-1}(x/a)} a\cos\theta\,(a\cos\theta\,d\theta) = a^2\int_0^{\sin^{-1}(x/a)} \cos^2\theta\,d\theta$$

$$= \frac{a^2}{2}\int_0^{\sin^{-1}(x/a)} (1+\cos 2\theta)\,d\theta = \frac{a^2}{2}\Big[\theta + \tfrac{1}{2}\sin 2\theta\Big]_0^{\sin^{-1}(x/a)} = \frac{a^2}{2}\Big[\theta + \sin\theta\,\cos\theta\Big]_0^{\sin^{-1}(x/a)}$$

$$= \frac{a^2}{2}\left[\left(\sin^{-1}\left(\frac{x}{a}\right) + \frac{x}{a}\cdot\frac{\sqrt{a^2-x^2}}{a}\right) - 0\right] = \tfrac{1}{2}a^2\sin^{-1}(x/a) + \tfrac{1}{2}x\sqrt{a^2-x^2}$$

(b) The integral $\int_0^x \sqrt{a^2-t^2}\,dt$ represents the area under the curve $y = \sqrt{a^2-t^2}$ between the vertical lines $t = 0$ and $t = x$. The figure shows that this area consists of a triangular region and a sector of the circle $t^2 + y^2 = a^2$. The triangular region has base x and height $\sqrt{a^2-x^2}$, so its area is $\frac{1}{2}x\sqrt{a^2-x^2}$. The sector has area $\frac{1}{2}a^2\theta = \frac{1}{2}a^2\sin^{-1}(x/a)$.

46. The curves intersect when $x^2 + \left(\frac{1}{2}x^2\right)^2 = 8 \;\Leftrightarrow\; x^2 + \frac{1}{4}x^4 = 8 \;\Leftrightarrow\; x^4 + 4x^2 - 32 = 0 \;\Leftrightarrow$ $(x^2+8)(x^2-4) = 0 \;\Leftrightarrow\; x = \pm 2$. The area inside the circle and above the parabola is given by

$$A_1 = \int_{-2}^{2}\left(\sqrt{8-x^2} - \tfrac{1}{2}x^2\right)dx = 2\int_0^2\sqrt{8-x^2}\,dx - 2\int_0^2 \tfrac{1}{2}x^2\,dx$$

$$= 2\left[\tfrac{1}{2}(8)\sin^{-1}\left(\tfrac{2}{\sqrt{8}}\right) + \tfrac{1}{2}(2)\sqrt{8-2^2} - \tfrac{1}{2}\left[\tfrac{1}{3}x^3\right]_0^2\right] \qquad \text{[by Exercise 45]}$$

$$= 8\sin^{-1}\left(\tfrac{1}{\sqrt{2}}\right) + 2\sqrt{4} - \tfrac{8}{3} = 8\left(\tfrac{\pi}{4}\right) + 4 - \tfrac{8}{3} = 2\pi + \tfrac{4}{3}$$

Since the area of the disk is $\pi\left(\sqrt{8}\right)^2 = 8\pi$, the area inside the circle and below the parabola ia $A_2 = 8\pi - \left(2\pi + \frac{4}{3}\right) = 6\pi - \frac{4}{3}$.

47. We use cylindrical shells and assume that $R > r$. $x^2 = r^2 - (y-R)^2 \;\Rightarrow\; x = \pm\sqrt{r^2-(y-R)^2}$, so $g(y) = 2\sqrt{r^2-(y-R)^2}$ and

$$V = \int_{R-r}^{R+r} 2\pi y\cdot 2\sqrt{r^2-(y-R)^2}\,dy = \int_{-r}^{r} 4\pi(u+R)\sqrt{r^2-u^2}\,du \qquad [\text{where } u = y-R]$$

$$= 4\pi\int_{-r}^{r} u\sqrt{r^2-u^2}\,du + 4\pi R\int_{-r}^{r}\sqrt{r^2-u^2}\,du \qquad \left[\begin{array}{c}\text{where } u = r\sin\theta\,,\ du = r\cos\theta\,d\theta \\ \text{in the second integral}\end{array}\right]$$

$$= 4\pi\left[-\tfrac{1}{3}(r^2-u^2)^{3/2}\right]_{-r}^{r} + 4\pi R\int_{-\pi/2}^{\pi/2} r^2\cos^2\theta\,d\theta = -\tfrac{4\pi}{3}(0-0) + 4\pi R r^2\int_{-\pi/2}^{\pi/2}\cos^2\theta\,d\theta$$

$$= 2\pi R r^2\int_{-\pi/2}^{\pi/2}(1+\cos 2\theta)\,d\theta = 2\pi R r^2\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_{-\pi/2}^{\pi/2} = 2\pi^2 R r^2$$

Another method: Use washers instead of shells, so $V = 8\pi R\int_0^r\sqrt{r^2-y^2}\,dy$ as in Exercise 6.2.75(a), but evaluate the integral using $y = r\sin\theta$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. Let $x = b\tan\theta$, so that $dx = b\sec^2\theta\,d\theta$ and $\sqrt{x^2+b^2} = b\sec\theta$.

$$E(P) = \int_{-a}^{L-a} \frac{\lambda b}{4\pi\varepsilon_0(x^2+b^2)^{3/2}}\,dx = \frac{\lambda b}{4\pi\varepsilon_0}\int_{\theta_1}^{\theta_2} \frac{1}{(b\sec\theta)^3}\,b\sec^2\theta\,d\theta$$

$$= \frac{\lambda}{4\pi\varepsilon_0 b}\int_{\theta_1}^{\theta_2} \frac{1}{\sec\theta}\,d\theta = \frac{\lambda}{4\pi\varepsilon_0 b}\int_{\theta_1}^{\theta_2} \cos\theta\,d\theta = \frac{\lambda}{4\pi\varepsilon_0 b}\Big[\sin\theta\Big]_{\theta_1}^{\theta_2}$$

$$= \frac{\lambda}{4\pi\varepsilon_0 b}\left[\frac{x}{\sqrt{x^2+b^2}}\right]_{-a}^{L-a} = \frac{\lambda}{4\pi\varepsilon_0 b}\left(\frac{L-a}{\sqrt{(L-a)^2+b^2}} + \frac{a}{\sqrt{a^2+b^2}}\right)$$

49. Let the equation of the large circle be $x^2+y^2=R^2$. Then the equation of the small circle is $x^2+(y-b)^2=r^2$, where $b=\sqrt{R^2-r^2}$ is the distance between the centers of the circles. The desired area is

$$A = \int_{-r}^{r}\left[\left(b+\sqrt{r^2-x^2}\right) - \sqrt{R^2-x^2}\,\right]dx$$
$$= 2\int_0^r \left(b+\sqrt{r^2-x^2}-\sqrt{R^2-x^2}\right)dx$$
$$= 2\int_0^r b\,dx + 2\int_0^r \sqrt{r^2-x^2}\,dx - 2\int_0^r \sqrt{R^2-x^2}\,dx$$

The first integral is just $2br = 2r\sqrt{R^2-r^2}$. The second integral represents the area of a quarter-circle of radius r, so its value is $\frac{1}{4}\pi r^2$. To evaluate the other integral, note that

$$\int \sqrt{a^2-x^2}\,dx = \int a^2\cos^2\theta\,d\theta \quad [x = a\sin\theta,\ dx = a\cos\theta\,d\theta] \ = \left(\tfrac{1}{2}a^2\right)\int(1+\cos 2\theta)\,d\theta$$
$$= \tfrac{1}{2}a^2\left(\theta + \tfrac{1}{2}\sin 2\theta\right) + C = \tfrac{1}{2}a^2(\theta + \sin\theta\,\cos\theta) + C$$
$$= \frac{a^2}{2}\arcsin\left(\frac{x}{a}\right) + \frac{a^2}{2}\left(\frac{x}{a}\right)\frac{\sqrt{a^2-x^2}}{a} + C = \frac{a^2}{2}\arcsin\left(\frac{x}{a}\right) + \frac{x}{2}\sqrt{a^2-x^2} + C$$

Thus, the desired area is

$$A = 2r\sqrt{R^2-r^2} + 2\left(\tfrac{1}{4}\pi r^2\right) - \left[R^2\arcsin(x/R) + x\sqrt{R^2-x^2}\,\right]_0^r$$
$$= 2r\sqrt{R^2-r^2} + \tfrac{1}{2}\pi r^2 - \left[R^2\arcsin(r/R) + r\sqrt{R^2-r^2}\,\right] = r\sqrt{R^2-r^2} + \tfrac{\pi}{2}r^2 - R^2\arcsin(r/R)$$

50. Note that the circular cross-sections of the tank are the same everywhere, so the percentage of the total capacity that is being used is equal to the percentage of any cross-section that is under water. The underwater area is

$$A = 2\int_{-5}^{2}\sqrt{25-y^2}\,dy$$
$$= \Big[25\arcsin(y/5) + y\sqrt{25-y^2}\,\Big]_{-5}^{2} \qquad [\text{substitute } y = 5\sin\theta]$$
$$= 25\arcsin\tfrac{2}{5} + 2\sqrt{21} + \tfrac{25}{2}\pi \approx 58.72\ \text{ft}^2$$

so the fraction of the total capacity in use is $\dfrac{A}{\pi(5)^2} \approx \dfrac{58.72}{25\pi} \approx 0.748$ or 74.8%.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7.4 Integration of Rational Functions by Partial Fractions

1. (a) $\dfrac{1}{(x-3)(x+5)} = \dfrac{A}{x-3} + \dfrac{B}{x+5}$

(b) $\dfrac{2x+5}{(x-2)^2(x^2+2)} = \dfrac{A}{x-2} + \dfrac{B}{(x-2)^2} + \dfrac{Cx+D}{x^2+2}$

2. (a) $\dfrac{x-6}{x^2+x-6} = \dfrac{x-6}{(x+3)(x-2)} = \dfrac{A}{x+3} + \dfrac{B}{x-2}$

(b) $\dfrac{1}{x^2+x^4} = \dfrac{1}{x^2(1+x^2)} = \dfrac{A}{x} + \dfrac{B}{x^2} + \dfrac{Cx+D}{1+x^2}$

3. (a) $\dfrac{x^2+4}{x^3-3x^2+2x} = \dfrac{x^2+4}{x(x^2-3x+2)} = \dfrac{x^2+4}{x(x-1)(x-2)} = \dfrac{A}{x} + \dfrac{B}{x-1} + \dfrac{C}{x-2}$

(b) $\dfrac{x^3+x}{x(2x-1)^2(x^2+3)^2} = \dfrac{A}{x} + \dfrac{B}{2x-1} + \dfrac{C}{(2x-1)^2} + \dfrac{Dx+E}{x^2+3} + \dfrac{Fx+G}{(x^2+3)^2}$

4. (a) $\dfrac{5}{x^4-1} = \dfrac{5}{(x^2+1)(x^2-1)} = \dfrac{5}{(x^2+1)(x+1)(x-1)} = \dfrac{A}{x+1} + \dfrac{B}{x-1} + \dfrac{Cx+D}{x^2+1}$

(b) $$\begin{aligned}\frac{x^4+x+1}{(x^3-1)(x^2-1)} &= \frac{x^4+x+1}{(x-1)(x^2+x+1)(x+1)(x-1)} = \frac{x^4+x+1}{(x+1)(x-1)^2(x^2+x+1)} \\ &= \frac{A}{x+1} + \frac{B}{x-1} + \frac{C}{(x-1)^2} + \frac{Dx+E}{x^2+x+1}\end{aligned}$$

5. (a) $\dfrac{x^5+1}{(x^2-x)(x^4+2x^2+1)} = \dfrac{x^5+1}{x(x-1)(x^2+1)^2} = \dfrac{A}{x} + \dfrac{B}{x-1} + \dfrac{Cx+D}{x^2+1} + \dfrac{Ex+F}{(x^2+1)^2}$

(b) $\dfrac{x^2}{x^2+x-6} = 1 + \dfrac{-x+6}{x^2+x-6} = 1 + \dfrac{-x+6}{(x-2)(x+3)} = 1 + \dfrac{A}{x-2} + \dfrac{B}{x+3}$

6. (a) $$\begin{aligned}\frac{x^6}{x^2-4} &= x^4 + 4x^2 + 16 + \frac{64}{(x+2)(x-2)} \quad \text{[by long division]} \\ &= x^4 + 4x^2 + 16 + \frac{A}{x+2} + \frac{B}{x-2}\end{aligned}$$

(b) $\dfrac{x^4}{(x^2-x+1)(x^2+2)^2} = \dfrac{Ax+B}{x^2-x+1} + \dfrac{Cx+D}{x^2+2} + \dfrac{Ex+F}{(x^2+2)^2}$

7. $\dfrac{5}{(x-1)(x+4)} = \dfrac{A}{x-1} + \dfrac{B}{x+4}$. Multiply both sides by $(x-1)(x+4)$ to get $5 = A(x+4) + B(x-1) \quad \Rightarrow$ $5 = (A+B)x + (4A-B)$. The coefficients of x must be equal and the constant terms are also equal, so $A+B=0$ and $4A-B=5$. Adding the equations together gives $5A = 5 \quad \Leftrightarrow \quad A = 1$, and hence $B = -1$. Thus,

$$\int \frac{5}{(x-1)(x+4)}\,dx = \int \left(\frac{1}{x-1} - \frac{1}{x+4}\right) dx = \ln|x-1| - \ln|x+4| + C.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $\dfrac{x-12}{x^2-4x} = \dfrac{x-12}{x(x-4)} = \dfrac{A}{x} + \dfrac{B}{x-4}$. Multiply both sides by $x(x-4)$ to get $x-12 = A(x-4)+Bx \quad \Rightarrow$

$x-12 = (A+B)x + (-4A)$. The coefficients of x must be equal and the constant terms are also equal, so $A+B=1$ and $-4A=-12$. The second equation gives $A=3$, which after substituting in the first equation gives $B=-2$. Thus,

$$\int \frac{x-12}{x^2-4x}\,dx = \int \left(\frac{3}{x} - \frac{2}{x-4}\right) dx = 3\ln|x| - 2\ln|x-4| + C.$$

9. $\dfrac{5x+1}{(2x+1)(x-1)} = \dfrac{A}{2x+1} + \dfrac{B}{x-1}$. Multiply both sides by $(2x+1)(x-1)$ to get $5x+1 = A(x-1)+B(2x+1) \quad \Rightarrow$

$5x+1 = Ax - A + 2Bx + B \quad \Rightarrow \quad 5x+1 = (A+2B)x + (-A+B)$.

The coefficients of x must be equal and the constant terms are also equal, so $A+2B=5$ and $-A+B=1$. Adding these equations gives us $3B=6 \quad \Leftrightarrow \quad B=2$, and hence, $A=1$. Thus,

$$\int \frac{5x+1}{(2x+1)(x-1)}\,dx = \int \left(\frac{1}{2x+1} + \frac{2}{x-1}\right) dx = \tfrac{1}{2}\ln|2x+1| + 2\ln|x-1| + C.$$

Another method: Substituting 1 for x in the equation $5x+1 = A(x-1)+B(2x+1)$ gives $6=3B \quad \Leftrightarrow \quad B=2$. Substituting $-\frac{1}{2}$ for x gives $-\frac{3}{2} = -\frac{3}{2}A \quad \Leftrightarrow \quad A=1$.

10. $\dfrac{y}{(y+4)(2y-1)} = \dfrac{A}{y+4} + \dfrac{B}{2y-1}$. Multiply both sides by $(y+4)(2y-1)$ to get $y = A(2y-1)+B(y+4) \quad \Rightarrow$

$y = 2Ay - A + By + 4B \quad \Rightarrow \quad y = (2A+B)y + (-A+4B)$. The coefficients of y must be equal and the constant terms are also equal, so $2A+B=1$ and $-A+4B=0$. Adding 2 times the second equation and the first equation gives us $9B=1 \quad \Leftrightarrow \quad B=\frac{1}{9}$ and hence, $A=\frac{4}{9}$. Thus,

$$\begin{aligned}\int \frac{y}{(y+4)(2y-1)}\,dy &= \int \left(\frac{\frac{4}{9}}{y+4} + \frac{\frac{1}{9}}{2y-1}\right) dy = \frac{4}{9}\ln|y+4| + \frac{1}{9}\cdot\frac{1}{2}\ln|2y-1| + C \\ &= \tfrac{4}{9}\ln|y+4| + \tfrac{1}{18}\ln|2y-1| + C\end{aligned}$$

Another method: Substituting $\frac{1}{2}$ for y in the equation $y = A(2y-1)+B(y+4)$ gives $\frac{1}{2} = \frac{9}{2}B \quad \Leftrightarrow \quad B=\frac{1}{9}$. Substituting -4 for y gives $-4=-9A \quad \Leftrightarrow \quad A=\frac{4}{9}$.

11. $\dfrac{2}{2x^2+3x+1} = \dfrac{2}{(2x+1)(x+1)} = \dfrac{A}{2x+1} + \dfrac{B}{x+1}$. Multiply both sides by $(2x+1)(x+1)$ to get

$2 = A(x+1)+B(2x+1)$. The coefficients of x must be equal and the constant terms are also equal, so $A+2B=0$ and $A+B=2$. Subtracting the second equation from the first gives $B=-2$, and hence, $A=4$. Thus,

$$\int_0^1 \frac{2}{2x^2+3x+1}\,dx = \int_0^1 \left(\frac{4}{2x+1} - \frac{2}{x+1}\right) dx = \left[\frac{4}{2}\ln|2x+1| - 2\ln|x+1|\right]_0^1 = (2\ln 3 - 2\ln 2) - 0 = 2\ln\frac{3}{2}.$$

Another method: Substituting -1 for x in the equation $2 = A(x+1)+B(2x+1)$ gives $2=-B \quad \Leftrightarrow \quad B=-2$. Substituting $-\frac{1}{2}$ for x gives $2=\frac{1}{2}A \quad \Leftrightarrow \quad A=4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $\dfrac{x-4}{x^2-5x+6} = \dfrac{A}{x-2} + \dfrac{B}{x-3}$. Multiply both sides by $(x-2)(x-3)$ to get $x-4 = A(x-3)+B(x-2) \quad \Rightarrow$

$x-4 = Ax-3A+Bx-2B \quad \Rightarrow \quad x-4 = (A+B)x+(-3A-2B)$.

The coefficients of x must be equal and the constant terms are also equal, so $A+B=1$ and $-3A-2B=-4$.

Adding twice the first equation to the second gives us $-A=-2 \quad \Leftrightarrow \quad A=2$, and hence, $B=-1$.Thus,

$$\int_0^1 \frac{x-4}{x^2-5x+6}\,dx = \int_0^1 \left(\frac{2}{x-2} - \frac{1}{x-3}\right) dx = \left[2\ln|x-2| - \ln|x-3|\right]_0^1$$
$$= (0-\ln 2) - (2\ln 2 - \ln 3) = -3\ln 2 + \ln 3 \ \ [\text{or } \ln\tfrac{3}{8}]$$

Another method: Substituting 3 for x in the equation $x-4 = A(x-3)+B(x-2)$ gives $-1=B$. Substituting 2 for x gives $-2=-A \quad \Leftrightarrow \quad A=2$.

13. $\dfrac{1}{x(x-a)} = \dfrac{A}{x} + \dfrac{B}{x-a}$. Multiply both sides by $x(x-a)$ to get $1 = A(x-a)+Bx \quad \Rightarrow \quad 1 = (A+B)x+(-aA)$.

The coefficients of x must be equal and the constant terms are also equal, so $A+B=0$ and $-aA=1$. The second equation gives $A=-1/a$, which after substituting in the first equation gives $B=1/a$. Thus,

$$\int \frac{1}{x(x-a)}\,dx = \int \left(-\frac{1/a}{x} + \frac{1/a}{x-a}\right) dx = -\frac{1}{a}\ln|x| + \frac{1}{a}\ln|x-a| + C.$$

14. If $a \neq b$, $\dfrac{1}{(x+a)(x+b)} = \dfrac{1}{b-a}\left(\dfrac{1}{x+a} - \dfrac{1}{x+b}\right)$, so if $a \neq b$, then

$$\int \frac{1}{(x+a)(x+b)}\,dx = \frac{1}{b-a}\left(\ln|x+a| - \ln|x+b|\right) + C = \frac{1}{b-a}\ln\left|\frac{x+a}{x+b}\right| + C$$

If $a=b$, then $\displaystyle\int \frac{1}{(x+a)^2}\,dx = -\frac{1}{x+a} + C$.

15. $\dfrac{x^2}{x-1} = \dfrac{(x^2-1)+1}{x-1} = \dfrac{(x+1)(x-1)+1}{x-1} = x+1+\dfrac{1}{x-1}$. [This result can also be obtained using long division.]

Thus, $\displaystyle\int \frac{x^2}{x-1}\,dx = \int \left(x+1+\frac{1}{x-1}\right) dx = \frac{1}{2}x^2 + x + \ln|x-1| + C$.

16. $\dfrac{3t-2}{t+1} = \dfrac{3t+3-5}{t+1} = \dfrac{3(t+1)-5}{t+1} = 3 - \dfrac{5}{t+1}$. Thus, $\displaystyle\int \frac{3t-2}{t+1}\,dt = \int \left(3 - \frac{5}{t+1}\right) dt = 3t - 5\ln|t+1| + C$.

17. $\dfrac{4y^2-7y-12}{y(y+2)(y-3)} = \dfrac{A}{y} + \dfrac{B}{y+2} + \dfrac{C}{y-3} \quad \Rightarrow \quad 4y^2-7y-12 = A(y+2)(y-3)+By(y-3)+Cy(y+2)$. Setting $y=0$ gives $-12=-6A$, so $A=2$. Setting $y=-2$ gives $18=10B$, so $B=\frac{9}{5}$. Setting $y=3$ gives $3=15C$, so $C=\frac{1}{5}$.

Now

$$\int_1^2 \frac{4y^2-7y-12}{y(y+2)(y-3)}\,dy = \int_1^2 \left(\frac{2}{y} + \frac{9/5}{y+2} + \frac{1/5}{y-3}\right) dy = \left[2\ln|y| + \tfrac{9}{5}\ln|y+2| + \tfrac{1}{5}\ln|y-3|\right]_1^2$$
$$= 2\ln 2 + \tfrac{9}{5}\ln 4 + \tfrac{1}{5}\ln 1 - 2\ln 1 - \tfrac{9}{5}\ln 3 - \tfrac{1}{5}\ln 2$$
$$= 2\ln 2 + \tfrac{18}{5}\ln 2 - \tfrac{1}{5}\ln 2 - \tfrac{9}{5}\ln 3 = \tfrac{27}{5}\ln 2 - \tfrac{9}{5}\ln 3 = \tfrac{9}{5}(3\ln 2 - \ln 3) = \tfrac{9}{5}\ln\tfrac{8}{3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $\dfrac{3x^2+6x+2}{x^2+3x+2} = 3 + \dfrac{-3x-4}{(x+1)(x+2)}$. Write $\dfrac{-3x-4}{(x+1)(x+2)} = \dfrac{A}{x+1} + \dfrac{B}{x+2}$. Multiplying both sides by $(x+1)(x+2)$ gives $-3x-4 = A(x+2)+B(x+1)$. Substituting -2 for x gives $2=-B \quad\Leftrightarrow\quad B=-2$. Substituting -1 for x gives $-1=A$. Thus,

$$\int_1^2 \frac{3x^2+6x+2}{x^2+3x+2}\,dx = \int_1^2 \left(3 - \frac{1}{x+1} - \frac{2}{x+2}\right) dx = \Big[3x - \ln|x+1| - 2\ln|x+2|\Big]_1^2$$
$$= (6 - \ln 3 - 2\ln 4) - (3 - \ln 2 - 2\ln 3) = 3 + \ln 2 + \ln 3 - 2\ln 4, \text{ or } 3 + \ln\tfrac{3}{8}$$

19. $\dfrac{x^2+x+1}{(x+1)^2(x+2)} = \dfrac{A}{x+1} + \dfrac{B}{(x+1)^2} + \dfrac{C}{x+2}$. Multiplying both sides by $(x+1)^2(x+2)$ gives $x^2+x+1 = A(x+1)(x+2)+B(x+2)+C(x+1)^2$. Substituting -1 for x gives $1=B$. Substituting -2 for x gives $3=C$. Equating coefficients of x^2 gives $1=A+C=A+3$, so $A=-2$. Thus,

$$\int_0^1 \frac{x^2+x+1}{(x+1)^2(x+2)}\,dx = \int_0^1 \left(\frac{-2}{x+1} + \frac{1}{(x+1)^2} + \frac{3}{x+2}\right) dx = \left[-2\ln|x+1| - \frac{1}{x+1} + 3\ln|x+2|\right]_0^1$$
$$= \left(-2\ln 2 - \tfrac{1}{2} + 3\ln 3\right) - (0 - 1 + 3\ln 2) = \tfrac{1}{2} - 5\ln 2 + 3\ln 3, \text{ or } \tfrac{1}{2} + \ln\tfrac{27}{32}$$

20. $\dfrac{x(3-5x)}{(3x-1)(x-1)^2} = \dfrac{A}{3x-1} + \dfrac{B}{x-1} + \dfrac{C}{(x-1)^2}$. Multiplying both sides by $(3x-1)(x-1)^2$ gives $x(3-5x) = A(x-1)^2 + B(x-1)(3x-1) + C(3x-1)$. Substituting 1 for x gives $-2=2C \quad\Leftrightarrow\quad C=-1$. Substituting $\frac{1}{3}$ for x gives $\frac{4}{9} = \frac{4}{9}A \quad\Leftrightarrow\quad A=1$. Substituting 0 for x gives $0 = A+B-C = 1+B+1$, so $B=-2$. Thus,

$$\int_2^3 \frac{x(3-5x)}{(3x-1)(x-1)^2}\,dx = \int_2^3 \left[\frac{1}{3x-1} - \frac{2}{x-1} - \frac{1}{(x-1)^2}\right] dx = \left[\frac{1}{3}\ln|3x-1| - 2\ln|x-1| + \frac{1}{x-1}\right]_2^3$$
$$= \left(\tfrac{1}{3}\ln 8 - 2\ln 2 + \tfrac{1}{2}\right) - \left(\tfrac{1}{3}\ln 5 - 0 + 1\right) = -\ln 2 - \tfrac{1}{3}\ln 5 - \tfrac{1}{2}$$

21. $\dfrac{1}{(t^2-1)^2} = \dfrac{1}{(t+1)^2(t-1)^2} = \dfrac{A}{t+1} + \dfrac{B}{(t+1)^2} + \dfrac{C}{t-1} + \dfrac{D}{(t-1)^2}$. Multiplying both sides by $(t+1)^2(t-1)^2$ gives $1 = A(t+1)(t-1)^2 + B(t-1)^2 + C(t-1)(t+1)^2 + D(t+1)^2$. Substituting 1 for t gives $1=4D \quad\Leftrightarrow\quad D=\frac{1}{4}$. Substituting -1 for t gives $1=4B \quad\Leftrightarrow\quad B=\frac{1}{4}$. Substituting 0 for t gives $1 = A+B-C+D = A+\frac{1}{4}-C+\frac{1}{4}$, so $\frac{1}{2} = A-C$. Equating coefficients of t^3 gives $0=A+C$. Adding the last two equations gives $2A=\frac{1}{2} \quad\Leftrightarrow\quad A=\frac{1}{4}$, and so $C=-\frac{1}{4}$. Thus,

$$\int \frac{dt}{(t^2-1)^2} = \int \left[\frac{1/4}{t+1} + \frac{1/4}{(t+1)^2} - \frac{1/4}{t-1} + \frac{1/4}{(t-1)^2}\right] dt$$
$$= \frac{1}{4}\left[\ln|t+1| - \frac{1}{t+1} - \ln|t-1| - \frac{1}{t-1}\right] + C, \text{ or } \frac{1}{4}\left(\ln\left|\frac{t+1}{t-1}\right| + \frac{2t}{1-t^2}\right) + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $\dfrac{3x^2+12x-20}{x^4-8x^2+16} = \dfrac{3x^2+12x-20}{(x^2-4)^2} = \dfrac{3x^2+12x-20}{(x-2)^2(x+2)^2} = \dfrac{A}{x-2} + \dfrac{B}{(x-2)^2} + \dfrac{C}{x+2} + \dfrac{D}{(x+2)^2}$. Multiply both sides by $(x-2)^2(x+2)^2$ to get $3x^2+12x-20 = A(x-2)(x+2)^2 + B(x+2)^2 + C(x-2)^2(x+2) + D(x-2)^2$. Setting $x=2$ gives $16=16B$, so $B=1$, and setting $x=-2$ gives $-32=16D$, so $D=-2$. Now, using these values of B and D and setting $x=0$ gives $-20=-8A+4+8C-8 \quad\Leftrightarrow\quad -2=-A+C$ **(1)**. Also, setting $x=1$ gives $-5=-9A+9+3C-2 \quad\Leftrightarrow\quad -4=-3A+C$ **(2)**. Subtracting **(2)** from **(1)** gives $2=2A \quad\Leftrightarrow\quad A=1$, and hence $C=-1$. Thus,

$$\int \frac{3x^2+12x-20}{x^4-8x^2+16}\,dx = \int\left(\frac{1}{x-2}+\frac{1}{(x-2)^2}-\frac{1}{x+2}-\frac{2}{(x+2)^2}\right)dx$$
$$= \ln|x-2| - \frac{1}{x-2} - \ln|x+2| + \frac{2}{x+2} + C$$

23. $\dfrac{10}{(x-1)(x^2+9)} = \dfrac{A}{x-1} + \dfrac{Bx+C}{x^2+9}$. Multiply both sides by $(x-1)\big(x^2+9\big)$ to get $10 = A\big(x^2+9\big) + (Bx+C)(x-1)$ $(\star)$. Substituting 1 for x gives $10=10A \quad\Leftrightarrow\quad A=1$. Substituting 0 for x gives $10=9A-C \quad\Rightarrow\quad C=9(1)-10=-1$. The coefficients of the x^2-terms in $(\star)$ must be equal, so $0=A+B \quad\Rightarrow\quad B=-1$. Thus,

$$\int \frac{10}{(x-1)(x^2+9)}\,dx = \int\left(\frac{1}{x-1}+\frac{-x-1}{x^2+9}\right)dx = \int\left(\frac{1}{x-1}-\frac{x}{x^2+9}-\frac{1}{x^2+9}\right)dx$$
$$= \ln|x-1| - \tfrac{1}{2}\ln(x^2+9) - \tfrac{1}{3}\tan^{-1}\left(\tfrac{x}{3}\right) + C$$

In the second term we used the substitution $u=x^2+9$ and in the last term we used Formula 10.

24. $\dfrac{3x^2-x+8}{x^3+4x} = \dfrac{3x^2-x+8}{x(x^2+4)} = \dfrac{A}{x} + \dfrac{Bx+C}{x^2+4}$. Multiply both sides by $x(x^2+4)$ to get $3x^2-x+8 = A(x^2+4)+x(Bx+C) \quad\Rightarrow\quad 3x^2-x+8=(A+B)x^2+Cx+4A$. Equating constant terms, we get $4A=8 \quad\Leftrightarrow\quad A=2$. Equating coefficients of x gives $C=-1$. Now equating coefficients of x^2 gives $A+B=3$, so $B=1$. Thus,

$$\int \frac{3x^2-x+8}{x^3+4x}\,dx = \int\left(\frac{2}{x}+\frac{x-1}{x^2+4}\right)dx = \int \frac{2}{x}+\frac{x}{x^2+4}-\frac{1}{x^2+4}dx$$
$$= 2\,\ln(x) + \tfrac{1}{2}\ln(x^2+4) - \tfrac{1}{2}\tan^{-1}\left(\tfrac{x}{2}\right) + C$$

25. $\dfrac{x^3-4x+1}{x^2-3x+2} = x+3+\dfrac{3x-5}{(x-1)(x-2)}$. Write $\dfrac{3x-5}{(x-1)(x-2)} = \dfrac{A}{x-1}+\dfrac{B}{x-2}$. Multiplying both sides by $(x-1)(x-2)$ gives $3x-5=A(x-2)+B(x-1)$. Substituting 2 for x gives $1=B$. Substituting 1 for x gives $-2=-A \quad\Leftrightarrow\quad A=2$. Thus,

$$\int_{-1}^{0} \frac{x^3-4x+1}{x^2-3x+2}\,dx = \int_{-1}^{0}\left(x+3+\frac{2}{x-1}+\frac{1}{x-2}\right)dx = \left[\tfrac{1}{2}x^2+3x+2\ln|x-1|+\ln|x-2|\right]_{-1}^{0}$$
$$= (0+0+0+\ln 2) - \left(\tfrac{1}{2}-3+2\ln 2+\ln 3\right) = \tfrac{5}{2}-\ln 2-\ln 3, \text{ or } \tfrac{5}{2}-\ln 6$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $\dfrac{x^3+4x^2+x-1}{x^3+x^2} = 1 + \dfrac{3x^2+x-1}{x^2(x+1)}$. Write $\dfrac{3x^2+x-1}{x^2(x+1)} = \dfrac{A}{x} + \dfrac{B}{x^2} + \dfrac{C}{x+1}$. Multiplying both sides by $x^2(x+1)$ gives $3x^2+x-1 = Ax(x+1)+B(x+1)+Cx^2$. Substituting 0 for x gives $-1=B$. Substituting -1 for x gives $1=C$. Equating coefficients of x^2 gives $3 = A+C = A+1$, so $A=2$. Thus,

$$\int_1^2 \frac{x^3+4x^2+x-1}{x^3+x^2}\,dx = \int_1^2 \left(1+\frac{2}{x}-\frac{1}{x^2}+\frac{1}{x+1}\right)dx = \left[x+2\ln|x|+\frac{1}{x}+\ln|x+1|\right]_1^2$$
$$= \left(2+2\ln 2+\tfrac{1}{2}+\ln 3\right)-(1+0+1+\ln 2) = \tfrac{1}{2}+\ln 2+\ln 3, \text{ or } \tfrac{1}{2}+\ln 6.$$

27. $\dfrac{4x}{x^3+x^2+x+1} = \dfrac{4x}{x^2(x+1)+1(x+1)} = \dfrac{4x}{(x+1)(x^2+1)} = \dfrac{A}{x+1} + \dfrac{Bx+C}{x^2+1}$. Multiply both sides by $(x+1)(x^2+1)$ to get $4x = A(x^2+1)+(Bx+C)(x+1) \quad\Leftrightarrow\quad 4x = Ax^2+A+Bx^2+Bx+Cx+C \quad\Leftrightarrow$ $4x = (A+B)x^2+(B+C)x+(A+C)$. Comparing coefficients gives us the following system of equations:

$$A+B=0 \quad \textbf{(1)} \qquad B+C=4 \quad \textbf{(2)} \qquad A+C=0 \quad \textbf{(3)}$$

Subtracting equation **(1)** from equation **(2)** gives us $-A+C=4$, and adding that equation to equation **(3)** gives us $2C=4 \quad\Leftrightarrow\quad C=2$, and hence $A=-2$ and $B=2$. Thus,

$$\int \frac{4x}{x^3+x^2+x+1}\,dx = \int\left(\frac{-2}{x+1}+\frac{2x+2}{x^2+1}\right)dx = \int\left(\frac{-2}{x+1}+\frac{2x}{x^2+1}+\frac{2}{x^2+1}\right)dx$$
$$= -2\ln|x+1|+\ln(x^2+1)+2\tan^{-1}x+C$$

28. $\displaystyle\int \frac{x^2+x+1}{(x^2+1)^2}\,dx = \int \frac{x^2+1}{(x^2+1)^2}\,dx + \int \frac{x}{(x^2+1)^2}\,dx = \int \frac{1}{x^2+1}\,dx + \frac{1}{2}\int \frac{1}{u^2}\,du \quad \left[u=x^2+1,\ du=2x\,dx\right]$

$$= \tan^{-1}x + \frac{1}{2}\left(-\frac{1}{u}\right)+C = \tan^{-1}x - \frac{1}{2(x^2+1)}+C$$

29. $\dfrac{x^3+4x+3}{x^4+5x^2+4} = \dfrac{x^3+4x+3}{(x^2+1)(x^2+4)} = \dfrac{Ax+B}{x^2+1} + \dfrac{Cx+D}{x^2+4}$. Multiply both sides by $(x^2+1)(x^2+4)$ to get $x^3+4x+3 = (Ax+B)(x^2+4)+(Cx+D)(x^2+1) \quad\Leftrightarrow$

$x^3+4x+3 = Ax^3+Bx^2+4Ax+4B+Cx^3+Dx^2+Cx+D \quad\Leftrightarrow$

$x^3+4x+3 = (A+C)x^3+(B+D)x^2+(4A+C)x+(4B+D)$. Comparing coefficients gives us the following system of equations:

$$A+C=1 \quad \textbf{(1)} \qquad B+D=0 \quad \textbf{(2)} \qquad 4A+C=4 \quad \textbf{(3)} \qquad 4B+D=3 \quad \textbf{(4)}$$

Subtracting equation **(1)** from equation **(3)** gives us $A=1$ and hence, $C=0$. Subtracting equation **(2)** from equation **(4)** gives us $B=1$ and hence, $D=-1$. Thus,

$$\int \frac{x^3+4x+3}{x^4+5x^2+4}\,dx = \int\left(\frac{x+1}{x^2+1}+\frac{-1}{x^2+4}\right)dx = \int\left(\frac{x}{x^2+1}+\frac{1}{x^2+1}-\frac{1}{x^2+4}\right)dx$$
$$= \frac{1}{2}\ln(x^2+1)+\tan^{-1}x-\frac{1}{2}\tan^{-1}\left(\frac{x}{2}\right)+C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $\dfrac{x^3+6x-2}{x^4+6x^2} = \dfrac{x^3+6x-2}{x^2(x^2+6)} = \dfrac{A}{x} + \dfrac{B}{x^2} + \dfrac{Cx+D}{x^2+6}$. Multiply both sides by $x^2(x^2+6)$ to get

$x^3+6x-2 = Ax(x^2+6) + B(x^2+6) + (Cx+D)x^2 \quad \Leftrightarrow$

$x^3+6x-2 = Ax^3+6Ax+Bx^2+6B+Cx^3+Dx^2 \quad \Leftrightarrow \quad x^3+6x-2 = (A+C)x^3+(B+D)x^2+6Ax+6B.$

Substituting 0 for x gives $-2 = 6B \quad \Leftrightarrow \quad B = -\frac{1}{3}$. Equating coefficients of x^2 gives $0 = B + D$, so $D = \frac{1}{3}$. Equating coefficients of x gives $6 = 6A \quad \Leftrightarrow \quad A = 1$. Equating coefficients of x^3 gives $1 = A + C$, so $C = 0$. Thus,

$$\int \frac{x^3+6x-2}{x^4+6x^2}\,dx = \int \left(\frac{1}{x} + \frac{-1/3}{x^2} + \frac{1/3}{x^2+6}\right) dx = \ln|x| + \frac{1}{3x} + \frac{1}{3\sqrt{6}}\tan^{-1}\left(\frac{x}{\sqrt{6}}\right) + C.$$

31.
$$\begin{aligned}
\int \frac{x+4}{x^2+2x+5}\,dx &= \int \frac{x+1}{x^2+2x+5}\,dx + \int \frac{3}{x^2+2x+5}\,dx = \frac{1}{2}\int \frac{(2x+2)\,dx}{x^2+2x+5} + \int \frac{3\,dx}{(x+1)^2+4} \\
&= \frac{1}{2}\ln\left|x^2+2x+5\right| + 3\int \frac{2\,du}{4(u^2+1)} \qquad \left[\begin{array}{c}\text{where } x+1 = 2u, \\ \text{and } dx = 2\,du\end{array}\right] \\
&= \frac{1}{2}\ln(x^2+2x+5) + \frac{3}{2}\tan^{-1}u + C = \frac{1}{2}\ln(x^2+2x+5) + \frac{3}{2}\tan^{-1}\left(\frac{x+1}{2}\right) + C
\end{aligned}$$

32.
$$\begin{aligned}
\int_0^1 \frac{x}{x^2+4x+13}\,dx &= \int_0^1 \frac{\frac{1}{2}(2x+4)}{x^2+4x+13}\,dx - 2\int_0^1 \frac{dx}{(x+2)^2+9} \\
&= \frac{1}{2}\int_{13}^{18} \frac{dy}{y} - 2\int_{2/3}^{1} \frac{3\,du}{9u^2+9} \qquad \left[\begin{array}{c}\text{where } y = x^2+4x+13,\ dy = (2x+4)\,dx, \\ x+2 = 3u, \text{ and } dx = 3\,du\end{array}\right] \\
&= \tfrac{1}{2}\left[\ln y\right]_{13}^{18} - \tfrac{2}{3}\left[\tan^{-1}u\right]_{2/3}^{1} = \tfrac{1}{2}\ln\tfrac{18}{13} - \tfrac{2}{3}\left(\tfrac{\pi}{4} - \tan^{-1}\left(\tfrac{2}{3}\right)\right) \\
&= \tfrac{1}{2}\ln\tfrac{18}{13} - \tfrac{\pi}{6} + \tfrac{2}{3}\tan^{-1}\left(\tfrac{2}{3}\right)
\end{aligned}$$

33. $\dfrac{1}{x^3-1} = \dfrac{1}{(x-1)(x^2+x+1)} = \dfrac{A}{x-1} + \dfrac{Bx+C}{x^2+x+1} \quad \Rightarrow \quad 1 = A\left(x^2+x+1\right) + (Bx+C)(x-1).$

Take $x = 1$ to get $A = \frac{1}{3}$. Equating coefficients of x^2 and then comparing the constant terms, we get $0 = \frac{1}{3} + B$, $1 = \frac{1}{3} - C$, so $B = -\frac{1}{3}$, $C = -\frac{2}{3} \quad \Rightarrow$

$$\begin{aligned}
\int \frac{1}{x^3-1}dx &= \int \frac{\frac{1}{3}}{x-1}\,dx + \int \frac{-\frac{1}{3}x - \frac{2}{3}}{x^2+x+1}\,dx = \tfrac{1}{3}\ln|x-1| - \frac{1}{3}\int \frac{x+2}{x^2+x+1}dx \\
&= \tfrac{1}{3}\ln|x-1| - \frac{1}{3}\int \frac{x+1/2}{x^2+x+1}dx - \frac{1}{3}\int \frac{(3/2)\,dx}{(x+1/2)^2+3/4} \\
&= \tfrac{1}{3}\ln|x-1| - \tfrac{1}{6}\ln\left(x^2+x+1\right) - \tfrac{1}{2}\left(\frac{2}{\sqrt{3}}\right)\tan^{-1}\left(\frac{x+\frac{1}{2}}{\sqrt{3}/2}\right) + K \\
&= \tfrac{1}{3}\ln|x-1| - \tfrac{1}{6}\ln(x^2+x+1) - \tfrac{1}{\sqrt{3}}\tan^{-1}\left(\tfrac{1}{\sqrt{3}}(2x+1)\right) + K
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. $\dfrac{x^3-2x^2+2x-5}{x^4+4x^2+3} = \dfrac{x^3-2x^2+2x-5}{(x^2+1)(x^2+3)} = \dfrac{Ax+B}{x^2+1} + \dfrac{Cx+D}{x^2+3}$. Multiply both sides by $(x^2+1)(x^2+3)$ to get

$x^3-2x^2+2x-5 = (Ax+B)(x^2+3)+(Cx+D)(x^2+1) \quad \Leftrightarrow$

$x^3-2x^2+2x-5 = Ax^3+Bx^2+3Ax+3B+Cx^3+Dx^2+Cx+D \quad \Leftrightarrow$

$x^3-2x^2+2x-5 = (A+C)x^3+(B+D)x^2+(3A+C)x+(3B+D)$. Comparing coefficients gives us the following system of equations:

$A+C=1$ **(1)** $\qquad B+D=-2$ **(2)** $\qquad 3A+C=2$ **(3)** $\qquad 3B+D=-5$ **(4)**

Subtracting equation **(1)** from equation **(3)** gives us $2A=1 \quad \Leftrightarrow \quad A=\frac{1}{2}$, and hence, $C=\frac{1}{2}$. Subtracting equation **(2)** from equation **(4)** gives us $2B=-3 \quad \Leftrightarrow \quad B=-\frac{3}{2}$, and hence, $D=-\frac{1}{2}$.

Thus,

$$\int \frac{x^3-2x^2+2x-5}{x^4+4x^2+3}\,dx = \int\left(\frac{\frac{1}{2}x-\frac{3}{2}}{x^2+1}+\frac{\frac{1}{2}x-\frac{1}{2}}{x^2+3}\right)dx = \int\left(\frac{\frac{1}{2}x}{x^2+1}-\frac{\frac{3}{2}}{x^2+1}+\frac{\frac{1}{2}x}{x^2+3}-\frac{\frac{1}{2}}{x^2+3}\right)dx$$

$$= \tfrac{1}{4}\ln(x^2+1)-\tfrac{3}{2}\tan^{-1}x+\tfrac{1}{4}\ln(x^2+3)-\frac{1}{2\sqrt{3}}\tan^{-1}\left(\frac{x}{\sqrt{3}}\right)+C$$

35. Let $u=x^4+4x^2+3$, so that $du=(4x^3+8x)\,dx=4(x^3+2x)\,dx$, $x=0 \quad \Rightarrow \quad u=3$, and $x=1 \quad \Rightarrow \quad u=8$.

Then $\displaystyle\int_0^1 \frac{x^3+2x}{x^4+4x^2+3}\,dx = \int_3^8 \frac{1}{u}\left(\frac{1}{4}\,du\right) = \frac{1}{4}\big[\ln|u|\,\big]_3^8 = \frac{1}{4}(\ln 8-\ln 3) = \frac{1}{4}\ln\frac{8}{3}$.

36. $\dfrac{x^5+x-1}{x^3+1} = x^2+\dfrac{-x^2+x-1}{x^3+1} = x^2+\dfrac{-x^2+x-1}{(x+1)(x^2-x+1)} = x^2+\dfrac{-1}{x+1}$, so

$$\int \frac{x^5+x-1}{x^3+1}\,dx = \int\left(x^2-\frac{1}{x+1}\right)dx = \frac{1}{3}x^3-\ln|x+1|+C$$

37. $\dfrac{5x^4+7x^2+x+2}{x(x^2+1)^2} = \dfrac{A}{x}+\dfrac{Bx+C}{x^2+1}+\dfrac{Dx+E}{(x^2+1)^2}$. Multiply by $x(x^2+1)^2$ to get

$5x^4+7x^2+x+2 = A(x^2+1)^2+(Bx+C)x(x^2+1)+(Dx+E)x \quad \Leftrightarrow$

$5x^4+7x^2+x+2 = A(x^4+2x^2+1)+(Bx^2+Cx)(x^2+1)+Dx^2+Ex \quad \Leftrightarrow$

$5x^4+7x^2+x+2 = Ax^4+2Ax^2+A+Bx^4+Cx^3+Bx^2+Cx+Dx^2+Ex \quad \Leftrightarrow$

$5x^4+7x^2+x+2 = (A+B)x^4+Cx^3+(2A+B+D)x^2+(C+E)x+A$. Equating coefficients gives us $C=0$, $A=2$, $A+B=5 \quad \Rightarrow \quad B=3$, $C+E=1 \quad \Rightarrow \quad E=1$, and $2A+B+D=7 \quad \Rightarrow \quad D=0$. Thus,

$\displaystyle\int \frac{5x^4+7x^2+x+2}{x(x^2+1)^2}\,dx = \int\left[\frac{2}{x}+\frac{3x}{x^2+1}+\frac{1}{(x^2+1)^2}\right]dx = I$. Now

$$\int \frac{dx}{(x^2+1)^2} = \int \frac{\sec^2\theta\,d\theta}{(\tan^2\theta+1)^2} \qquad \left[\begin{array}{l} x=\tan\theta, \\ dx=\sec^2\theta\,d\theta \end{array}\right]$$

$$= \int \frac{\sec^2\theta}{\sec^4\theta}\,d\theta = \int \cos^2\theta\,d\theta = \int \tfrac{1}{2}(1+\cos 2\theta)\,d\theta$$

$$= \tfrac{1}{2}\theta+\tfrac{1}{4}\sin 2\theta+C = \tfrac{1}{2}\theta+\tfrac{1}{2}\sin\theta\,\cos\theta+C$$

$$= \frac{1}{2}\tan^{-1}x+\frac{1}{2}\frac{x}{\sqrt{x^2+1}}\frac{1}{\sqrt{x^2+1}}+C$$

Therefore, $I = 2\ln|x|+\frac{3}{2}\ln(x^2+1)+\frac{1}{2}\tan^{-1}x+\dfrac{x}{2(x^2+1)}+C$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. Let $u = x^5 + 5x^3 + 5x$, so that $du = (5x^4 + 15x^2 + 5)dx = 5(x^4 + 3x^2 + 1)dx$. Then

$$\int \frac{x^4 + 3x^2 + 1}{x^5 + 5x^3 + 5x}\,dx = \int \frac{1}{u}\left(\frac{1}{5}\,du\right) = \frac{1}{5}\ln|u| + C = \frac{1}{5}\ln\left|x^5 + 5x^3 + 5x\right| + C$$

39. $\dfrac{x^2 - 3x + 7}{(x^2 - 4x + 6)^2} = \dfrac{Ax + B}{x^2 - 4x + 6} + \dfrac{Cx + D}{(x^2 - 4x + 6)^2} \quad \Rightarrow \quad x^2 - 3x + 7 = (Ax + B)(x^2 - 4x + 6) + Cx + D \quad \Rightarrow$

$x^2 - 3x + 7 = Ax^3 + (-4A + B)x^2 + (6A - 4B + C)x + (6B + D)$. So $A = 0$, $-4A + B = 1 \quad \Rightarrow \quad B = 1$,

$6A - 4B + C = -3 \quad \Rightarrow \quad C = 1, 6B + D = 7 \quad \Rightarrow \quad D = 1$. Thus,

$$\begin{aligned}
I &= \int \frac{x^2 - 3x + 7}{(x^2 - 4x + 6)^2}\,dx = \int \left(\frac{1}{x^2 - 4x + 6} + \frac{x + 1}{(x^2 - 4x + 6)^2}\right)dx \\
&= \int \frac{1}{(x - 2)^2 + 2}\,dx + \int \frac{x - 2}{(x^2 - 4x + 6)^2}\,dx + \int \frac{3}{(x^2 - 4x + 6)^2}\,dx \\
&= I_1 + I_2 + I_3.
\end{aligned}$$

$$I_1 = \int \frac{1}{(x - 2)^2 + \left(\sqrt{2}\right)^2}\,dx = \frac{1}{\sqrt{2}}\tan^{-1}\left(\frac{x - 2}{\sqrt{2}}\right) + C_1$$

$$I_2 = \frac{1}{2}\int \frac{2x - 4}{(x^2 - 4x + 6)^2}\,dx = \tfrac{1}{2}\int \frac{1}{u^2}\,du = \tfrac{1}{2}\left(-\frac{1}{u}\right) + C_2 = -\frac{1}{2(x^2 - 4x + 6)} + C_2$$

$$\begin{aligned}
I_3 &= 3\int \frac{1}{\left[(x - 2)^2 + \left(\sqrt{2}\right)^2\right]^2}\,dx = 3\int \frac{1}{[2(\tan^2\theta + 1)]^2}\sqrt{2}\sec^2\theta\,d\theta \qquad \begin{bmatrix} x - 2 = \sqrt{2}\tan\theta, \\ dx = \sqrt{2}\sec^2\theta\,d\theta \end{bmatrix} \\
&= \frac{3\sqrt{2}}{4}\int \frac{\sec^2\theta}{\sec^4\theta}\,d\theta = \frac{3\sqrt{2}}{4}\int \cos^2\theta\,d\theta = \frac{3\sqrt{2}}{4}\int \tfrac{1}{2}(1 + \cos 2\theta)\,d\theta \\
&= \frac{3\sqrt{2}}{8}\left(\theta + \tfrac{1}{2}\sin 2\theta\right) + C_3 = \frac{3\sqrt{2}}{8}\tan^{-1}\left(\frac{x - 2}{\sqrt{2}}\right) + \frac{3\sqrt{2}}{8}\left(\tfrac{1}{2}\cdot 2\sin\theta\,\cos\theta\right) + C_3 \\
&= \frac{3\sqrt{2}}{8}\tan^{-1}\left(\frac{x - 2}{\sqrt{2}}\right) + \frac{3\sqrt{2}}{8}\cdot\frac{x - 2}{\sqrt{x^2 - 4x + 6}}\cdot\frac{\sqrt{2}}{\sqrt{x^2 - 4x + 6}} + C_3 \\
&= \frac{3\sqrt{2}}{8}\tan^{-1}\left(\frac{x - 2}{\sqrt{2}}\right) + \frac{3(x - 2)}{4(x^2 - 4x + 6)} + C_3
\end{aligned}$$

So $\quad I = I_1 + I_2 + I_3 \qquad [C = C_1 + C_2 + C_3]$

$$\begin{aligned}
&= \frac{1}{\sqrt{2}}\tan^{-1}\left(\frac{x - 2}{\sqrt{2}}\right) + \frac{-1}{2(x^2 - 4x + 6)} + \frac{3\sqrt{2}}{8}\tan^{-1}\left(\frac{x - 2}{\sqrt{2}}\right) + \frac{3(x - 2)}{4(x^2 - 4x + 6)} + C \\
&= \left(\frac{4\sqrt{2}}{8} + \frac{3\sqrt{2}}{8}\right)\tan^{-1}\left(\frac{x - 2}{\sqrt{2}}\right) + \frac{3(x - 2) - 2}{4(x^2 - 4x + 6)} + C = \frac{7\sqrt{2}}{8}\tan^{-1}\left(\frac{x - 2}{\sqrt{2}}\right) + \frac{3x - 8}{4(x^2 - 4x + 6)} + C
\end{aligned}$$

40. $\dfrac{x^3 + 2x^2 + 3x - 2}{(x^2 + 2x + 2)^2} = \dfrac{Ax + B}{x^2 + 2x + 2} + \dfrac{Cx + D}{(x^2 + 2x + 2)^2} \quad \Rightarrow$

$x^3 + 2x^2 + 3x - 2 = (Ax + B)(x^2 + 2x + 2) + Cx + D \quad \Rightarrow$

$x^3 + 2x^2 + 3x - 2 = Ax^3 + (2A + B)x^2 + (2A + 2B + C)x + 2B + D.$

So $A = 1, 2A + B = 2 \quad \Rightarrow \quad B = 0, 2A + 2B + C = 3 \quad \Rightarrow \quad C = 1$, and $2B + D = -2 \quad \Rightarrow \quad D = -2$. Thus,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$I = \int \frac{x^3+2x^2+3x-2}{(x^2+2x+2)^2}\,dx = \int \left(\frac{x}{x^2+2x+2} + \frac{x-2}{(x^2+2x+2)^2}\right) dx$$

$$= \int \frac{x+1}{x^2+2x+2}\,dx + \int \frac{-1}{x^2+2x+2}\,dx + \int \frac{x+1}{(x^2+2x+2)^2}\,dx + \int \frac{-3}{(x^2+2x+2)^2}\,dx$$

$$= I_1 + I_2 + I_3 + I_4.$$

$$I_1 = \int \frac{x+1}{x^2+2x+2}\,dx = \int \frac{1}{u}\left(\frac{1}{2}\,du\right) \quad \begin{bmatrix} u = x^2+2x+2, \\ du = 2(x+1)\,dx \end{bmatrix} \quad = \frac{1}{2}\ln\left|x^2+2x+2\right| + C_1$$

$$I_2 = -\int \frac{1}{(x+1)^2+1}\,dx = -\frac{1}{1}\tan^{-1}\left(\frac{x+1}{1}\right) + C_2 = -\tan^{-1}(x+1) + C_2$$

$$I_3 = \int \frac{x+1}{(x^2+2x+2)^2}\,dx = \int \frac{1}{u^2}\left(\frac{1}{2}\,du\right) = -\frac{1}{2u} + C_3 = -\frac{1}{2(x^2+2x+2)} + C_3$$

$$I_4 = -3\int \frac{1}{[(x+1)^2+1]^2}\,dx = -3\int \frac{1}{(\tan^2\theta+1)^2}\sec^2\theta\,d\theta \quad \begin{bmatrix} x+1 = 1\tan\theta, \\ dx = \sec^2\theta\,d\theta \end{bmatrix}$$

$$= -3\int \frac{1}{\sec^2\theta}\,d\theta = -3\int \cos^2\theta\,d\theta = -\frac{3}{2}\int (1+\cos 2\theta)\,d\theta$$

$$= -\tfrac{3}{2}\left(\theta + \tfrac{1}{2}\sin 2\theta\right) + C_4 = -\tfrac{3}{2}\theta - \tfrac{3}{2}\left(\tfrac{1}{2}\cdot 2\sin\theta\,\cos\theta\right) + C_4$$

$$= -\frac{3}{2}\tan^{-1}\left(\frac{x+1}{1}\right) - \frac{3}{2}\cdot\frac{x+1}{\sqrt{x^2+2x+2}}\cdot\frac{1}{\sqrt{x^2+2x+2}} + C_4$$

$$= -\frac{3}{2}\tan^{-1}(x+1) - \frac{3(x+1)}{2(x^2+2x+2)} + C_4$$

So $\quad I = I_1 + I_2 + I_3 + I_4 \qquad [C = C_1 + C_2 + C_3 + C_4]$

$$= \frac{1}{2}\ln(x^2+2x+2) - \tan^{-1}(x+1) - \frac{1}{2(x^2+2x+2)} - \frac{3}{2}\tan^{-1}(x+1) - \frac{3(x+1)}{2(x^2+2x+2)} + C$$

$$= \frac{1}{2}\ln(x^2+2x+2) - \frac{5}{2}\tan^{-1}(x+1) - \frac{3x+4}{2(x^2+2x+2)} + C$$

41. $$\int \frac{dx}{x\sqrt{x-1}} = \int \frac{2u}{u(u^2+1)}\,du \qquad \begin{bmatrix} u = \sqrt{x-1}, & x = u^2+1 \\ u^2 = x-1, & dx = 2u\,du \end{bmatrix}$$

$$= 2\int \frac{1}{u^2+1}\,du = 2\tan^{-1}u + C = 2\tan^{-1}\sqrt{x-1} + C$$

42. Let $u = \sqrt{x+3}$, so $u^2 = x+3$ and $2u\,du = dx$. Then

$$\int \frac{dx}{2\sqrt{x+3}+x} = \int \frac{2u\,du}{2u+(u^2-3)} = \int \frac{2u}{u^2+2u-3}\,du = \int \frac{2u}{(u+3)(u-1)}\,du.$$ Now

$$\frac{2u}{(u+3)(u-1)} = \frac{A}{u+3} + \frac{B}{u-1} \quad \Rightarrow \quad 2u = A(u-1) + B(u+3).$$ Setting $u = 1$ gives $2 = 4B$, so $B = \frac{1}{2}$.

Setting $u = -3$ gives $-6 = -4A$, so $A = \frac{3}{2}$. Thus,

$$\int \frac{2u}{(u+3)(u-1)}\,du = \int \left(\frac{\frac{3}{2}}{u+3} + \frac{\frac{1}{2}}{u-1}\,du\right)$$

$$= \tfrac{3}{2}\ln|u+3| + \tfrac{1}{2}\ln|u-1| + C = \tfrac{3}{2}\ln\left(\sqrt{x+3}+3\right) + \tfrac{1}{2}\ln\left|\sqrt{x+3}-1\right| + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. Let $u=\sqrt{x}$, so $u^2=x$ and $2u\,du=dx$. Then $\displaystyle\int\frac{dx}{x^2+x\sqrt{x}}=\int\frac{2u\,du}{u^4+u^3}=\int\frac{2\,du}{u^3+u^2}=\int\frac{2\,du}{u^2(u+1)}$.

$\dfrac{2}{u^2(u+1)}=\dfrac{A}{u}+\dfrac{B}{u^2}+\dfrac{C}{u+1}\quad\Rightarrow\quad 2=Au(u+1)+B(u+1)+Cu^2$. Setting $u=0$ gives $B=2$. Setting $u=-1$ gives $C=2$. Equating coefficients of u^2, we get $0=A+C$, so $A=-2$. Thus,

$$\int\frac{2\,du}{u^2(u+1)}=\int\left(\frac{-2}{u}+\frac{2}{u^2}+\frac{2}{u+1}\right)du=-2\ln|u|-\frac{2}{u}+2\ln|u+1|+C=-2\ln\sqrt{x}-\frac{2}{\sqrt{x}}+2\ln\left(\sqrt{x}+1\right)+C.$$

44. Let $u=\sqrt[3]{x}$. Then $x=u^3$, $dx=3u^2\,du\quad\Rightarrow$

$$\int_0^1\frac{1}{1+\sqrt[3]{x}}\,dx=\int_0^1\frac{3u^2\,du}{1+u}=\int_0^1\left(3u-3+\frac{3}{1+u}\right)du=\left[\tfrac{3}{2}u^2-3u+3\ln(1+u)\right]_0^1=3\left(\ln 2-\tfrac{1}{2}\right).$$

45. Let $u=\sqrt[3]{x^2+1}$. Then $x^2=u^3-1$, $2x\,dx=3u^2\,du\quad\Rightarrow$

$$\int\frac{x^3}{\sqrt[3]{x^2+1}}\,dx=\int\frac{(u^3-1)\frac{3}{2}u^2\,du}{u}=\frac{3}{2}\int(u^4-u)\,du$$
$$=\tfrac{3}{10}u^5-\tfrac{3}{4}u^2+C=\tfrac{3}{10}(x^2+1)^{5/3}-\tfrac{3}{4}(x^2+1)^{2/3}+C$$

46. $$\int\frac{dx}{(1+\sqrt{x}\,)^2}=\int\frac{2(u-1)\,du}{u^2}\qquad\left[\begin{matrix}u=1+\sqrt{x},\\ x=(u-1)^2,\ dx=2(u-1)\,du\end{matrix}\right]$$
$$=2\int\left(\frac{1}{u}-\frac{1}{u^2}\right)du=2\ln|u|+\frac{2}{u}+C=2\ln(1+\sqrt{x}\,)+\frac{2}{1+\sqrt{x}}+C$$

47. If we were to substitute $u=\sqrt{x}$, then the square root would disappear but a cube root would remain. On the other hand, the substitution $u=\sqrt[3]{x}$ would eliminate the cube root but leave a square root. We can eliminate both roots by means of the substitution $u=\sqrt[6]{x}$. (Note that 6 is the least common multiple of 2 and 3.)

Let $u=\sqrt[6]{x}$. Then $x=u^6$, so $dx=6u^5\,du$ and $\sqrt{x}=u^3$, $\sqrt[3]{x}=u^2$. Thus,

$$\int\frac{1}{\sqrt{x}-\sqrt[3]{x}}\,dx=\int\frac{6u^5\,du}{u^3-u^2}=6\int\frac{u^5}{u^2(u-1)}\,du=6\int\frac{u^3}{u-1}\,du$$
$$=6\int\left(u^2+u+1+\frac{1}{u-1}\right)du\qquad\text{[by long division]}$$
$$=6\left(\tfrac{1}{3}u^3+\tfrac{1}{2}u^2+u+\ln|u-1|\right)+C=2\sqrt{x}+3\sqrt[3]{x}+6\sqrt[6]{x}+6\ln\left|\sqrt[6]{x}-1\right|+C$$

48. Let $u=x^{1/5}\quad\Rightarrow\quad x=u^5$, so $dx=5u^4\,du$. This substitution gives

$$I=\int\frac{1}{x-x^{1/5}}\,dx=\int\frac{5u^4}{u^5-u}\,du=\int\frac{5u^3}{u^4-1}\,du=\int\frac{5u^3}{(u^2-1)(u^2+1)}\,du=\int\frac{5u^3}{(u-1)(u+1)(u^2+1)}\,du.$$

Now $\dfrac{5u^3}{(u-1)(u+1)(u^2+1)}=\dfrac{A}{u-1}+\dfrac{B}{u+1}+\dfrac{Cu+D}{u^2+1}$. Multiply both sides by $(u-1)(u+1)(u^2+1)$ to get

$5u^3=A(u+1)(u^2+1)+B(u-1)(u^2+1)+(Cu+D)(u-1)(u+1)\quad\Leftrightarrow$

$5u^3=A(u^3+u^2+u+1)+B(u^3-u^2+u-1)+Cu(u^2-1)+D(u^2-1)\quad\Leftrightarrow$

$5u^3=(A+B+C)u^3+(A-B+D)u^2+(A+B-C)u+(A-B-D)$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Setting $u = 1$ gives $5 = 4A$, so $A = \frac{5}{4}$. Now, comparing coefficients gives us the following system of equations:

$$A + B + C = 5 \quad \textbf{(1)} \qquad A - B + D = 0 \quad \textbf{(2)}$$

$$A + B - C = 0 \quad \textbf{(3)} \qquad A - B - D = 0 \quad \textbf{(4)}$$

Adding equations **(1)** and **(3)** gives $2A + 2B = 5$, so $B = \frac{5}{4}$. Subtracting equation **(4)** from equation **(2)** gives $D = 0$. Finally, substituting the value of A and B in equation **(3)** gives $C = \frac{5}{2}$. Thus,

$$\begin{aligned} I &= \int \left(\frac{5/4}{u-1} + \frac{5/4}{u+1} + \frac{5u/2}{u^2+1} \right) du = \frac{5}{4} \ln|u-1| + \frac{5}{4} \ln|u+1| + \frac{5}{4} \ln(u^2+1) + C \\ &= \tfrac{5}{4} \ln\left|(u^2-1)(u^2+1)\right| + C = \tfrac{5}{4} \ln\left|u^4 - 1\right| + C = \tfrac{5}{4} \ln\left|x^{4/5} - 1\right| + C \end{aligned}$$

49. Let $u = \sqrt{x} \quad \Rightarrow \quad x = u^2$, so $dx = 2u\,du$. This substitution gives $I = \displaystyle\int \frac{1}{x - 3\sqrt{x} + 2}\,dx = \int \frac{2u}{u^2 - 3u + 2}\,du$. Now $\dfrac{2u}{u^2 - 3u + 2} = \dfrac{2u}{(u-2)(u-1)} = \dfrac{A}{u-2} + \dfrac{B}{u-1}$. Multiply both sides by $(u-2)(u-1)$ to get $2u = A(u-1) + B(u-2)$. Setting $u = 1$ gives $2 = -B$ or $B = -2$, and setting $u = 2$ gives $A = 4$. Thus,

$$I = \int \left(\frac{4}{u-2} - \frac{2}{u-1} \right) du = 4 \ln|u-2| - 2 \ln|u-1| + C = 4 \ln\left|\sqrt{x} - 2\right| - 2 \ln\left|\sqrt{x} - 1\right| + C.$$

50. Let $u = \sqrt{1 + \sqrt{x}}$, so that $u^2 = 1 + \sqrt{x}$, $x = (u^2 - 1)^2$, and $dx = 2(u^2 - 1) \cdot 2u\,du = 4u(u^2 - 1)\,du$. Then $\displaystyle\int \frac{\sqrt{1+\sqrt{x}}}{x}\,dx = \int \frac{u}{(u^2-1)^2} \cdot 4u(u^2-1)\,du = \int \frac{4u^2}{u^2-1}\,du = \int \left(4 + \frac{4}{u^2-1}\right) du$. Now $\dfrac{4}{u^2-1} = \dfrac{A}{u+1} + \dfrac{B}{u-1} \quad \Rightarrow \quad 4 = A(u-1) + B(u+1)$. Setting $u = 1$ gives $4 = 2B$, so $B = 2$. Setting $u = -1$ gives $4 = -2A$, so $A = -2$. Thus,

$$\begin{aligned} \int \left(4 + \frac{4}{u^2-1}\right) du &= \int \left(4 - \frac{2}{u+1} + \frac{2}{u-1}\right) du = 4u - 2\ln|u+1| + 2\ln|u-1| + C \\ &= 4\sqrt{1+\sqrt{x}} - 2\ln\left(\sqrt{1+\sqrt{x}} + 1\right) + 2\ln\left(\sqrt{1+\sqrt{x}} - 1\right) + C \end{aligned}$$

51. Let $u = e^x$. Then $x = \ln u$, $dx = \dfrac{du}{u} \quad \Rightarrow$

$$\begin{aligned} \int \frac{e^{2x}\,dx}{e^{2x} + 3e^x + 2} &= \int \frac{u^2\,(du/u)}{u^2 + 3u + 2} = \int \frac{u\,du}{(u+1)(u+2)} = \int \left[\frac{-1}{u+1} + \frac{2}{u+2}\right] du \\ &= 2\ln|u+2| - \ln|u+1| + C = \ln\frac{(e^x+2)^2}{e^x+1} + C \end{aligned}$$

52. Let $u = \cos x$, so that $du = -\sin x\,dx$. Then $\displaystyle\int \frac{\sin x}{\cos^2 x - 3\cos x}\,dx = \int \frac{1}{u^2 - 3u}\,(-du) = \int \frac{-1}{u(u-3)}\,du$. $\dfrac{-1}{u(u-3)} = \dfrac{A}{u} + \dfrac{B}{u-3} \quad \Rightarrow \quad -1 = A(u-3) + Bu$. Setting $u = 3$ gives $B = -\frac{1}{3}$. Setting $u = 0$ gives $A = \frac{1}{3}$. Thus, $\displaystyle\int \frac{-1}{u(u-3)}\,du = \int \left(\frac{\frac{1}{3}}{u} - \frac{\frac{1}{3}}{u-3}\right) du = \frac{1}{3}\ln|u| - \frac{1}{3}\ln|u-3| + C = \frac{1}{3}\ln|\cos x| - \frac{1}{3}\ln|\cos x - 3| + C$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

53. Let $u = \tan t$, so that $du = \sec^2 t\,dt$. Then $\displaystyle\int \frac{\sec^2 t}{\tan^2 t + 3\tan t + 2}\,dt = \int \frac{1}{u^2+3u+2}\,du = \int \frac{1}{(u+1)(u+2)}\,du$.

Now $\dfrac{1}{(u+1)(u+2)} = \dfrac{A}{u+1} + \dfrac{B}{u+2} \quad\Rightarrow\quad 1 = A(u+2) + B(u+1)$.

Setting $u = -2$ gives $1 = -B$, so $B = -1$. Setting $u = -1$ gives $1 = A$.

Thus, $\displaystyle\int \frac{1}{(u+1)(u+2)}\,du = \int \left(\frac{1}{u+1} - \frac{1}{u+2}\right) du = \ln|u+1| - \ln|u+2| + C = \ln|\tan t + 1| - \ln|\tan t + 2| + C$.

54. Let $u = e^x$, so that $du = e^x\,dx$. Then $\displaystyle\int \frac{e^x}{(e^x-2)(e^{2x}+1)}\,dx = \int \frac{1}{(u-2)(u^2+1)}\,du$. Now

$\dfrac{1}{(u-2)(u^2+1)} = \dfrac{A}{u-2} + \dfrac{Bu+C}{u^2+1} \quad\Rightarrow\quad 1 = A(u^2+1) + (Bu+C)(u-2)$. Setting $u = 2$ gives $1 = 5A$, so $A = \frac{1}{5}$.

Setting $u = 0$ gives $1 = \frac{1}{5} - 2C$, so $C = -\frac{2}{5}$. Comparing coefficients of u^2 gives $0 = \frac{1}{5} + B$, so $B = -\frac{1}{5}$. Thus,

$$\begin{aligned}\int \frac{1}{(u-2)(u^2+1)}\,du &= \int \left(\frac{\frac{1}{5}}{u-2} + \frac{-\frac{1}{5}u - \frac{2}{5}}{u^2+1}\right) du = \tfrac{1}{5}\int \frac{1}{u-2}\,du - \tfrac{1}{5}\int \frac{u}{u^2+1}\,du - \tfrac{2}{5}\int \frac{1}{u^2+1}\,du \\ &= \tfrac{1}{5}\ln|u-2| - \tfrac{1}{5}\cdot\tfrac{1}{2}\ln\left|u^2+1\right| - \tfrac{2}{5}\tan^{-1} u + C \\ &= \tfrac{1}{5}\ln|e^x-2| - \tfrac{1}{10}\ln(e^{2x}+1) - \tfrac{2}{5}\tan^{-1} e^x + C\end{aligned}$$

55. Let $u = e^x$, so that $du = e^x\,dx$ and $dx = \dfrac{du}{u}$. Then $\displaystyle\int \frac{dx}{1+e^x} = \int \frac{du}{(1+u)u}$. $\dfrac{1}{u(u+1)} = \dfrac{A}{u} + \dfrac{B}{u+1} \quad\Rightarrow$

$1 = A(u+1) + Bu$. Setting $u = -1$ gives $B = -1$. Setting $u = 0$ gives $A = 1$. Thus,

$\displaystyle\int \frac{du}{u(u+1)} = \int \left(\frac{1}{u} - \frac{1}{u+1}\right) du = \ln|u| - \ln|u+1| + C = \ln e^x - \ln(e^x+1) + C = x - \ln(e^x+1) + C$.

56. Let $u = \sinh t$, so that $du = \cosh t\,dt$. Then $\displaystyle\int \frac{\cosh t}{\sinh^2 t + \sinh^4 t}\,dt = \int \frac{1}{u^2+u^4}\,du = \int \frac{1}{u^2(u^2+1)}\,du$.

$\dfrac{1}{u^2(u^2+1)} = \dfrac{A}{u} + \dfrac{B}{u^2} + \dfrac{Cu+D}{u^2+1} \quad\Rightarrow\quad 1 = Au(u^2+1) + B(u^2+1) + (Cu+D)u^2$. Setting $u = 0$ gives $B = 1$.

Comparing coefficients of u^2, we get $0 = B + D$, so $D = -1$. Comparing coefficients of u, we get $0 = A$. Comparing coefficients of u^3, we get $0 = A + C$, so $C = 0$. Thus,

$$\begin{aligned}\int \frac{1}{u^2(u^2+1)}\,du &= \int \left(\frac{1}{u^2} - \frac{1}{u^2+1}\right) du = -\frac{1}{u} - \tan^{-1} u + C = -\frac{1}{\sinh t} - \tan^{-1}(\sinh t) + C \\ &= -\operatorname{csch} t - \tan^{-1}(\sinh t) + C\end{aligned}$$

57. Let $u = \ln(x^2 - x + 2)$, $dv = dx$. Then $du = \dfrac{2x-1}{x^2-x+2}\,dx$, $v = x$, and (by integration by parts)

$$\begin{aligned}\int \ln(x^2-x+2)\,dx &= x\ln(x^2-x+2) - \int \frac{2x^2-x}{x^2-x+2}\,dx = x\ln(x^2-x+2) - \int \left(2 + \frac{x-4}{x^2-x+2}\right) dx \\ &= x\ln(x^2-x+2) - 2x - \int \frac{\frac{1}{2}(2x-1)}{x^2-x+2}\,dx + \frac{7}{2}\int \frac{dx}{(x-\frac{1}{2})^2 + \frac{7}{4}}\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$= x\ln(x^2 - x + 2) - 2x - \frac{1}{2}\ln(x^2 - x + 2) + \frac{7}{2}\int \frac{\frac{\sqrt{7}}{2}\,du}{\frac{7}{4}(u^2+1)} \qquad \left[\begin{array}{c}\text{where } x - \frac{1}{2} = \frac{\sqrt{7}}{2}u,\\ dx = \frac{\sqrt{7}}{2}\,du,\\ (x-\frac{1}{2})^2 + \frac{7}{4} = \frac{7}{4}(u^2+1)\end{array}\right]$$

$$= (x - \tfrac{1}{2})\ln(x^2 - x + 2) - 2x + \sqrt{7}\tan^{-1} u + C$$

$$= (x - \tfrac{1}{2})\ln(x^2 - x + 2) - 2x + \sqrt{7}\tan^{-1}\frac{2x-1}{\sqrt{7}} + C$$

58. Let $u = \tan^{-1} x$, $dv = x\,dx \quad\Rightarrow\quad du = dx/(1+x^2)$, $v = \frac{1}{2}x^2$.

Then $\displaystyle\int x\tan^{-1} x\,dx = \tfrac{1}{2}x^2\tan^{-1} x - \frac{1}{2}\int \frac{x^2}{1+x^2}\,dx$. To evaluate the last integral, use long division or observe that $\displaystyle\int \frac{x^2}{1+x^2}\,dx = \int \frac{(1+x^2)-1}{1+x^2}\,dx = \int 1\,dx - \int \frac{1}{1+x^2}\,dx = x - \tan^{-1} x + C_1$. So

$\int x\tan^{-1} x\,dx = \frac{1}{2}x^2\tan^{-1} x - \frac{1}{2}(x - \tan^{-1} x + C_1) = \frac{1}{2}(x^2\tan^{-1} x + \tan^{-1} x - x) + C$.

59.

From the graph, we see that the integral will be negative, and we guess that the area is about the same as that of a rectangle with width 2 and height 0.3, so we estimate the integral to be $-(2\cdot 0.3) = -0.6$. Now

$$\frac{1}{x^2 - 2x - 3} = \frac{1}{(x-3)(x+1)} = \frac{A}{x-3} + \frac{B}{x+1} \quad\Leftrightarrow$$

$1 = (A+B)x + A - 3B$, so $A = -B$ and $A - 3B = 1 \quad\Leftrightarrow\quad A = \frac{1}{4}$

and $B = -\frac{1}{4}$, so the integral becomes

$$\int_0^2 \frac{dx}{x^2 - 2x - 3} = \frac{1}{4}\int_0^2 \frac{dx}{x-3} - \frac{1}{4}\int_0^2 \frac{dx}{x+1} = \frac{1}{4}\Big[\ln|x-3| - \ln|x+1|\Big]_0^2 = \frac{1}{4}\left[\ln\left|\frac{x-3}{x+1}\right|\right]_0^2$$

$$= \tfrac{1}{4}\big(\ln\tfrac{1}{3} - \ln 3\big) = -\tfrac{1}{2}\ln 3 \approx -0.55$$

60. $\boldsymbol{k = 0}$: $\displaystyle\int \frac{dx}{x^2+k} = \int \frac{dx}{x^2} = -\frac{1}{x} + C$

$\boldsymbol{k > 0}$: $\displaystyle\int \frac{dx}{x^2+k} = \int \frac{dx}{x^2 + (\sqrt{k}\,)^2} = \frac{1}{\sqrt{k}}\tan^{-1}\left(\frac{x}{\sqrt{k}}\right) + C$

$\boldsymbol{k < 0}$: $\displaystyle\int \frac{dx}{x^2+k} = \int \frac{dx}{x^2 - (-k)} = \int \frac{dx}{x^2 - \left(\sqrt{-k}\right)^2} = \frac{1}{2\sqrt{-k}}\ln\left|\frac{x-\sqrt{-k}}{x+\sqrt{-k}}\right| + C$ [by Example 3]

61. $\displaystyle\int \frac{dx}{x^2 - 2x} = \int \frac{dx}{(x-1)^2 - 1} = \int \frac{du}{u^2 - 1}$ [put $u = x - 1$]

$\displaystyle = \frac{1}{2}\ln\left|\frac{u-1}{u+1}\right| + C$ [by Equation 6] $\displaystyle = \frac{1}{2}\ln\left|\frac{x-2}{x}\right| + C$

62. $\displaystyle\int \frac{(2x+1)\,dx}{4x^2 + 12x - 7} = \frac{1}{4}\int \frac{(8x+12)\,dx}{4x^2 + 12x - 7} - \int \frac{2\,dx}{(2x+3)^2 - 16}$

$\displaystyle = \tfrac{1}{4}\ln\left|4x^2 + 12x - 7\right| - \int \frac{du}{u^2 - 16}$ [put $u = 2x + 3$]

$= \frac{1}{4}\ln\left|4x^2 + 12x - 7\right| - \frac{1}{8}\ln|(u-4)/(u+4)| + C$ [by Equation 6]

$= \frac{1}{4}\ln\left|4x^2 + 12x - 7\right| - \frac{1}{8}\ln|(2x-1)/(2x+7)| + C$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

63. (a) If $t = \tan\left(\frac{x}{2}\right)$, then $\frac{x}{2} = \tan^{-1} t$. The figure gives

$$\cos\left(\frac{x}{2}\right) = \frac{1}{\sqrt{1+t^2}} \text{ and } \sin\left(\frac{x}{2}\right) = \frac{t}{\sqrt{1+t^2}}.$$

(b) $\cos x = \cos\left(2 \cdot \frac{x}{2}\right) = 2\cos^2\left(\frac{x}{2}\right) - 1$

$$= 2\left(\frac{1}{\sqrt{1+t^2}}\right)^2 - 1 = \frac{2}{1+t^2} - 1 = \frac{1-t^2}{1+t^2}$$

$$\sin x = \sin\left(2 \cdot \frac{x}{2}\right) = 2\sin\left(\frac{x}{2}\right)\cos\left(\frac{x}{2}\right) = 2\frac{t}{\sqrt{t^2+1}} \cdot \frac{1}{\sqrt{t^2+1}} = \frac{2t}{t^2+1}$$

(c) $\frac{x}{2} = \arctan t \quad \Rightarrow \quad x = 2\arctan t \quad \Rightarrow \quad dx = \frac{2}{1+t^2}\,dt$

64. Let $t = \tan(x/2)$. Then, by using the expressions in Exercise 63, we have

$$\int \frac{dx}{1-\cos x} = \int \frac{2\,dt/(1+t^2)}{1-(1-t^2)/(1+t^2)} = \int \frac{2\,dt}{(1+t^2)-(1-t^2)} = \int \frac{2\,dt}{2t^2} = \int \frac{1}{t^2}\,dt$$

$$= -\frac{1}{t} + C = -\frac{1}{\tan(x/2)} + C = -\cot(x/2) + C$$

Another method: $\displaystyle\int \frac{dx}{1-\cos x} = \int \left(\frac{1}{1-\cos x} \cdot \frac{1+\cos x}{1+\cos x}\right) dx = \int \frac{1+\cos x}{1-\cos^2 x}\,dx$

$$= \int \frac{1+\cos x}{\sin^2 x}\,dx = \int \left(\frac{1}{\sin^2 x} + \frac{\cos x}{\sin^2 x}\right) dx$$

$$= \int (\csc^2 x + \csc x \cot x)\,dx = -\cot x - \csc x + C$$

65. Let $t = \tan(x/2)$. Then, using the expressions in Exercise 63, we have

$$\int \frac{1}{3\sin x - 4\cos x}\,dx = \int \frac{1}{3\left(\frac{2t}{1+t^2}\right) - 4\left(\frac{1-t^2}{1+t^2}\right)} \frac{2\,dt}{1+t^2} = 2\int \frac{dt}{3(2t) - 4(1-t^2)} = \int \frac{dt}{2t^2+3t-2}$$

$$= \int \frac{dt}{(2t-1)(t+2)} = \int \left[\frac{2}{5}\frac{1}{2t-1} - \frac{1}{5}\frac{1}{t+2}\right] dt \qquad \text{[using partial fractions]}$$

$$= \tfrac{1}{5}\Big[\ln|2t-1| - \ln|t+2|\Big] + C = \frac{1}{5}\ln\left|\frac{2t-1}{t+2}\right| + C = \frac{1}{5}\ln\left|\frac{2\tan(x/2)-1}{\tan(x/2)+2}\right| + C$$

66. Let $t = \tan(x/2)$. Then, by Exercise 63,

$$\int_{\pi/3}^{\pi/2} \frac{dx}{1+\sin x - \cos x} = \int_{1/\sqrt{3}}^{1} \frac{2\,dt/(1+t^2)}{1+2t/(1+t^2)-(1-t^2)/(1+t^2)} = \int_{1/\sqrt{3}}^{1} \frac{2\,dt}{1+t^2+2t-1+t^2}$$

$$= \int_{1/\sqrt{3}}^{1} \left[\frac{1}{t} - \frac{1}{t+1}\right] dt = \Big[\ln t - \ln(t+1)\Big]_{1/\sqrt{3}}^{1}$$

$$= \ln\frac{1}{2} - \ln\frac{1}{\sqrt{3}+1} = \ln\frac{\sqrt{3}+1}{2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. Let $t = \tan(x/2)$. Then, by Exercise 63,

$$\int_0^{\pi/2} \frac{\sin 2x}{2+\cos x}\,dx = \int_0^{\pi/2} \frac{2\sin x\,\cos x}{2+\cos x}\,dx = \int_0^1 \frac{2\cdot\dfrac{2t}{1+t^2}\cdot\dfrac{1-t^2}{1+t^2}}{2+\dfrac{1-t^2}{1+t^2}}\,\frac{2}{1+t^2}\,dt = \int_0^1 \frac{\dfrac{8t(1-t^2)}{(1+t^2)^2}}{2(1+t^2)+(1-t^2)}\,dt$$

$$= \int_0^1 8t\cdot\frac{1-t^2}{(t^2+3)(t^2+1)^2}\,dt = I$$

If we now let $u = t^2$, then $\dfrac{1-t^2}{(t^2+3)(t^2+1)^2} = \dfrac{1-u}{(u+3)(u+1)^2} = \dfrac{A}{u+3} + \dfrac{B}{u+1} + \dfrac{C}{(u+1)^2} \quad\Rightarrow$

$1-u = A(u+1)^2 + B(u+3)(u+1) + C(u+3)$. Set $u=-1$ to get $2 = 2C$, so $C = 1$. Set $u = -3$ to get $4 = 4A$, so $A = 1$. Set $u = 0$ to get $1 = 1 + 3B + 3$, so $B = -1$. So

$$I = \int_0^1 \left[\frac{8t}{t^2+3} - \frac{8t}{t^2+1} + \frac{8t}{(t^2+1)^2}\right]dt = \left[4\ln(t^2+3) - 4\ln(t^2+1) - \frac{4}{t^2+1}\right]_0^1$$

$$= (4\ln 4 - 4\ln 2 - 2) - (4\ln 3 - 0 - 4) = 8\ln 2 - 4\ln 2 - 4\ln 3 + 2 = 4\ln\tfrac{2}{3} + 2$$

68. $\dfrac{1}{x^3+x} = \dfrac{1}{x(x^2+1)} = \dfrac{A}{x} + \dfrac{Bx+C}{x^2+1} \quad\Rightarrow\quad 1 = A(x^2+1) + (Bx+C)x$. Set $x = 0$ to get $1 = A$. So

$1 = (1+B)x^2 + Cx + 1 \quad\Rightarrow\quad B + 1 = 0 \quad [B = -1]$ and $C = 0$. Thus, the area is

$$\int_1^2 \frac{1}{x^3+x}\,dx = \int_1^2 \left(\frac{1}{x} - \frac{x}{x^2+1}\right)dx = \left[\ln|x| - \tfrac{1}{2}\ln\left|x^2+1\right|\right]_1^2 = \left(\ln 2 - \tfrac{1}{2}\ln 5\right) - \left(0 - \tfrac{1}{2}\ln 2\right)$$

$$= \tfrac{3}{2}\ln 2 - \tfrac{1}{2}\ln 5 \quad \left[\text{or } \tfrac{1}{2}\ln\tfrac{8}{5}\right]$$

69. By long division, $\dfrac{x^2+1}{3x-x^2} = -1 + \dfrac{3x+1}{3x-x^2}$. Now

$\dfrac{3x+1}{3x-x^2} = \dfrac{3x+1}{x(3-x)} = \dfrac{A}{x} + \dfrac{B}{3-x} \quad\Rightarrow\quad 3x+1 = A(3-x) + Bx$. Set $x = 3$ to get $10 = 3B$, so $B = \frac{10}{3}$. Set $x = 0$ to get $1 = 3A$, so $A = \frac{1}{3}$. Thus, the area is

$$\int_1^2 \frac{x^2+1}{3x-x^2}\,dx = \int_1^2 \left(-1 + \frac{\frac{1}{3}}{x} + \frac{\frac{10}{3}}{3-x}\right)dx = \left[-x + \tfrac{1}{3}\ln|x| - \tfrac{10}{3}\ln|3-x|\right]_1^2$$

$$= \left(-2 + \tfrac{1}{3}\ln 2 - 0\right) - \left(-1 + 0 - \tfrac{10}{3}\ln 2\right) = -1 + \tfrac{11}{3}\ln 2$$

70. (a) We use disks, so the volume is $V = \pi\displaystyle\int_0^1 \left[\frac{1}{x^2+3x+2}\right]^2 dx = \pi\int_0^1 \frac{dx}{(x+1)^2(x+2)^2}$. To evaluate the integral,

we use partial fractions: $\dfrac{1}{(x+1)^2(x+2)^2} = \dfrac{A}{x+1} + \dfrac{B}{(x+1)^2} + \dfrac{C}{x+2} + \dfrac{D}{(x+2)^2} \quad\Rightarrow$

$1 = A(x+1)(x+2)^2 + B(x+2)^2 + C(x+1)^2(x+2) + D(x+1)^2$. We set $x = -1$, giving $B = 1$, then set $x = -2$, giving $D = 1$. Now equating coefficients of x^3 gives $A = -C$, and then equating constants gives $1 = 4A + 4 + 2(-A) + 1 \quad\Rightarrow\quad A = -2 \quad\Rightarrow\quad C = 2$. So the expression becomes

$$V = \pi\int_0^1 \left[\frac{-2}{x+1} + \frac{1}{(x+1)^2} + \frac{2}{(x+2)} + \frac{1}{(x+2)^2}\right]dx = \pi\left[2\ln\left|\frac{x+2}{x+1}\right| - \frac{1}{x+1} - \frac{1}{x+2}\right]_0^1$$

$$= \pi\left[\left(2\ln\tfrac{3}{2} - \tfrac{1}{2} - \tfrac{1}{3}\right) - \left(2\ln 2 - 1 - \tfrac{1}{2}\right)\right] = \pi\left(2\ln\tfrac{3/2}{2} + \tfrac{2}{3}\right) = \pi\left(\tfrac{2}{3} + \ln\tfrac{9}{16}\right)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) In this case, we use cylindrical shells, so the volume is $V = 2\pi \int_0^1 \frac{x\,dx}{x^2+3x+2} = 2\pi \int_0^1 \frac{x\,dx}{(x+1)(x+2)}$. We use partial fractions to simplify the integrand: $\frac{x}{(x+1)(x+2)} = \frac{A}{x+1} + \frac{B}{x+2} \quad \Rightarrow \quad x = (A+B)x + 2A + B$. So $A + B = 1$ and $2A + B = 0 \quad \Rightarrow \quad A = -1$ and $B = 2$. So the volume is

$$2\pi \int_0^1 \left[\frac{-1}{x+1} + \frac{2}{x+2}\right] dx = 2\pi \Big[-\ln|x+1| + 2\,\ln|x+2|\Big]_0^1$$
$$= 2\pi(-\ln 2 + 2\ln 3 + \ln 1 - 2\ln 2) = 2\pi(2\ln 3 - 3\ln 2) = 2\pi \ln \tfrac{9}{8}$$

71. $t = \int \frac{P+S}{P[(r-1)P-S]}\,dP = \int \frac{P+S}{P(0.1P-S)}\,dP \quad [r = 1.1]$. Now $\frac{P+S}{P(0.1P-S)} = \frac{A}{P} + \frac{B}{0.1P-S} \quad \Rightarrow$ $P + S = A(0.1P - S) + BP$. Substituting 0 for P gives $S = -AS \quad \Rightarrow \quad A = -1$. Substituting $10S$ for P gives $11S = 10BS \quad \Rightarrow \quad B = \frac{11}{10}$. Thus, $t = \int \left(\frac{-1}{P} + \frac{11/10}{0.1P-S}\right) dP \quad \Rightarrow \quad t = -\ln P + 11\ln(0.1P - S) + C$.

When $t = 0$, $P = 10{,}000$ and $S = 900$, so $0 = -\ln 10{,}000 + 11\ln(1000 - 900) + C \quad \Rightarrow$ $C = \ln 10{,}000 - 11\ln 100 \quad [= \ln 10^{-18} \approx -41.45]$.

Therefore, $t = -\ln P + 11\ln\left(\frac{1}{10}P - 900\right) + \ln 10{,}000 - 11\ln 100 \quad \Rightarrow \quad t = \ln\frac{10{,}000}{P} + 11\ln\frac{P-9000}{1000}$.

72. If we add and subtract $2x^2$ (because $2x^2$ completes the square for x^4+1), we get

$$x^4 + 1 = x^4 + 2x^2 + 1 - 2x^2 = \left(x^2+1\right)^2 - 2x^2 = \left(x^2+1\right)^2 - \left(\sqrt{2}\,x\right)^2$$
$$= \left[(x^2+1) - \sqrt{2}\,x\right]\left[\left(x^2+1\right) + \sqrt{2}\,x\right] = \left(x^2 - \sqrt{2}\,x + 1\right)\left(x^2 + \sqrt{2}\,x + 1\right)$$

So we can decompose $\frac{1}{x^4+1} = \frac{Ax+B}{x^2+\sqrt{2}\,x+1} + \frac{Cx+D}{x^2-\sqrt{2}\,x+1} \quad \Rightarrow$

$1 = (Ax+B)\left(x^2 - \sqrt{2}\,x + 1\right) + (Cx+D)\left(x^2 + \sqrt{2}\,x + 1\right)$. Setting the constant terms equal gives $B + D = 1$, then from the coefficients of x^3 we get $A + C = 0$. Now from the coefficients of x we get $A + C + (B - D)\sqrt{2} = 0 \quad \Leftrightarrow$ $[(1-D) - D]\sqrt{2} = 0 \quad \Rightarrow \quad D = \frac{1}{2} \quad \Rightarrow \quad B = \frac{1}{2}$, and finally, from the coefficients of x^2 we get $\sqrt{2}\,(C - A) + B + D = 0 \quad \Rightarrow \quad C - A = -\frac{1}{\sqrt{2}} \quad \Rightarrow \quad C = -\frac{\sqrt{2}}{4}$ and $A = \frac{\sqrt{2}}{4}$. So we rewrite the integrand, splitting the terms into forms which we know how to integrate:

$$\frac{1}{x^4+1} = \frac{\frac{\sqrt{2}}{4}x + \frac{1}{2}}{x^2+\sqrt{2}\,x+1} + \frac{-\frac{\sqrt{2}}{4}x + \frac{1}{2}}{x^2-\sqrt{2}\,x+1} = \frac{1}{4\sqrt{2}}\left[\frac{2x+2\sqrt{2}}{x^2+\sqrt{2}\,x+1} - \frac{2x-2\sqrt{2}}{x^2-\sqrt{2}\,x+1}\right]$$
$$= \frac{\sqrt{2}}{8}\left[\frac{2x+\sqrt{2}}{x^2+\sqrt{2}\,x+1} - \frac{2x-\sqrt{2}}{x^2-\sqrt{2}\,x+1}\right] + \frac{1}{4}\left[\frac{1}{\left(x+\frac{1}{\sqrt{2}}\right)^2 + \frac{1}{2}} + \frac{1}{\left(x-\frac{1}{\sqrt{2}}\right)^2 + \frac{1}{2}}\right]$$

Now we integrate:

$$\int \frac{dx}{x^4+1} = \frac{\sqrt{2}}{8}\ln\left(\frac{x^2+\sqrt{2}\,x+1}{x^2-\sqrt{2}\,x+1}\right) + \frac{\sqrt{2}}{4}\left[\tan^{-1}\left(\sqrt{2}\,x+1\right) + \tan^{-1}\left(\sqrt{2}\,x-1\right)\right] + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

73. (a) In Maple, we define $f(x)$, and then use `convert(f,parfrac,x);` to obtain

$$f(x) = \frac{24{,}110/4879}{5x+2} - \frac{668/323}{2x+1} - \frac{9438/80{,}155}{3x-7} + \frac{(22{,}098x + 48{,}935)/260{,}015}{x^2+x+5}$$

In Mathematica, we use the command `Apart`.

(b)
$$\begin{aligned}\int f(x)\,dx &= \tfrac{24{,}110}{4879}\cdot\tfrac{1}{5}\ln|5x+2| - \tfrac{668}{323}\cdot\tfrac{1}{2}\ln|2x+1| - \tfrac{9438}{80{,}155}\cdot\tfrac{1}{3}\ln|3x-7| \\ &\quad + \frac{1}{260{,}015}\int \frac{22{,}098\left(x+\frac{1}{2}\right)+37{,}886}{\left(x+\frac{1}{2}\right)^2+\frac{19}{4}}\,dx + C \\ &= \tfrac{24{,}110}{4879}\cdot\tfrac{1}{5}\ln|5x+2| - \tfrac{668}{323}\cdot\tfrac{1}{2}\ln|2x+1| - \tfrac{9438}{80{,}155}\cdot\tfrac{1}{3}\ln|3x-7| \\ &\quad + \tfrac{1}{260{,}015}\left[22{,}098\cdot\tfrac{1}{2}\ln\left(x^2+x+5\right) + 37{,}886\cdot\sqrt{\tfrac{4}{19}}\tan^{-1}\left(\tfrac{1}{\sqrt{19/4}}\left(x+\tfrac{1}{2}\right)\right)\right] + C \\ &= \tfrac{4822}{4879}\ln|5x+2| - \tfrac{334}{323}\ln|2x+1| - \tfrac{3146}{80{,}155}\ln|3x-7| + \tfrac{11{,}049}{260{,}015}\ln\left(x^2+x+5\right) \\ &\quad + \tfrac{75{,}772}{260{,}015\sqrt{19}}\tan^{-1}\left[\tfrac{1}{\sqrt{19}}(2x+1)\right] + C\end{aligned}$$

Using a CAS, we get

$$\begin{aligned}&\frac{4822\ln(5x+2)}{4879} - \frac{334\ln(2x+1)}{323} - \frac{3146\ln(3x-7)}{80{,}155} \\ &\qquad + \frac{11{,}049\ln(x^2+x+5)}{260{,}015} + \frac{3988\sqrt{19}}{260{,}015}\tan^{-1}\left[\frac{\sqrt{19}}{19}(2x+1)\right]\end{aligned}$$

The main difference in this answer is that the absolute value signs and the constant of integration have been omitted. Also, the fractions have been reduced and the denominators rationalized.

74. (a) In Maple, we define $f(x)$, and then use `convert(f,parfrac,x);` to get

$$f(x) = \frac{5828/1815}{(5x-2)^2} - \frac{59{,}096/19{,}965}{5x-2} + \frac{2(2843x+816)/3993}{2x^2+1} + \frac{(313x-251)/363}{(2x^2+1)^2}.$$

In Mathematica, we use the command `Apart`, and in Derive, we use `Expand`.

(b) As we saw in Exercise 73, computer algebra systems omit the absolute value signs in $\int (1/y)\,dy = \ln|y|$. So we use the CAS to integrate the expression in part (a) and add the necessary absolute value signs and constant of integration to get

$$\begin{aligned}\int f(x)\,dx &= -\frac{5828}{9075(5x-2)} - \frac{59{,}096\ln|5x-2|}{99{,}825} + \frac{2843\ln\left(2x^2+1\right)}{7986} \\ &\quad + \frac{503}{15{,}972}\sqrt{2}\tan^{-1}\left(\sqrt{2}\,x\right) - \frac{1}{2904}\,\frac{1004x+626}{2x^2+1} + C\end{aligned}$$

(c) From the graph, we see that f goes from negative to positive at $x \approx -0.78$, then back to negative at $x \approx 0.8$, and finally back to positive at $x = 1$. Also, $\lim_{x\to 0.4} f(x) = \infty$. So we see (by the First Derivative Test) that $\int f(x)\,dx$ has minima at $x \approx -0.78$ and $x = 1$, and a maximum at $x \approx 0.80$, and that $\int f(x)\,dx$ is unbounded as $x \to 0.4$. Note also that

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

just to the right of $x = 0.4$, f has large values, so $\int f(x)\,dx$ increases rapidly, but slows down as f drops toward 0. $\int f(x)\,dx$ decreases from about 0.8 to 1, then increases slowly since f stays small and positive.

75. There are only finitely many values of x where $Q(x) = 0$ (assuming that Q is not the zero polynomial). At all other values of x, $F(x)/Q(x) = G(x)/Q(x)$, so $F(x) = G(x)$. In other words, the values of F and G agree at all except perhaps finitely many values of x. By continuity of F and G, the polynomials F and G must agree at those values of x too.

More explicitly: if a is a value of x such that $Q(a) = 0$, then $Q(x) \neq 0$ for all x sufficiently close to a. Thus,

$$\begin{aligned} F(a) &= \lim_{x\to a} F(x) && \text{[by continuity of } F\text{]} \\ &= \lim_{x\to a} G(x) && \text{[whenever } Q(x) \neq 0\text{]} \\ &= G(a) && \text{[by continuity of } G\text{]} \end{aligned}$$

76. (a) Let $u = (x^2 + a^2)^{-n}$, $dv = dx \quad\Rightarrow\quad du = -n(x^2 + a^2)^{-n-1}\,2x\,dx$, $v = x$.

$$\begin{aligned} I_n = \int \frac{dx}{(x^2+a^2)^n} &= \frac{x}{(x^2+a^2)^n} - \int \frac{-2nx^2}{(x^2+a^2)^{n+1}}\,dx \qquad \text{[by parts]} \\ &= \frac{x}{(x^2+a^2)^n} + 2n\int \frac{(x^2+a^2)-a^2}{(x^2+a^2)^{n+1}}\,dx \\ &= \frac{x}{(x^2+a^2)^n} + 2n\int \frac{dx}{(x^2+a^2)^n} - 2na^2\int \frac{dx}{(x^2+a^2)^{n+1}} \end{aligned}$$

Recognizing the last two integrals as I_n and I_{n+1}, we can solve for I_{n+1} in terms of I_n.

$$2na^2 I_{n+1} = \frac{x}{(x^2+a^2)^n} + 2nI_n - I_n \quad\Rightarrow\quad I_{n+1} = \frac{x}{2a^2n(x^2+a^2)^n} + \frac{2n-1}{2a^2n}I_n \quad\Rightarrow$$

$$I_n = \frac{x}{2a^2(n-1)(x^2+a^2)^{n-1}} + \frac{2n-3}{2a^2(n-1)}I_{n-1}$$ [decrease n-values by 1], which is the desired result.

(b) Using part (a) with $a = 1$ and $n = 2$, we get

$$\int \frac{dx}{(x^2+1)^2} = \frac{x}{2(x^2+1)} + \frac{1}{2}\int \frac{dx}{x^2+1} = \frac{x}{2(x^2+1)} + \frac{1}{2}\tan^{-1}x + C$$

Using part (a) with $a = 1$ and $n = 3$, we get

$$\begin{aligned} \int \frac{dx}{(x^2+1)^3} &= \frac{x}{2(2)(x^2+1)^2} + \frac{3}{2(2)}\int \frac{dx}{(x^2+1)^2} = \frac{x}{4(x^2+1)^2} + \frac{3}{4}\left[\frac{x}{2(x^2+1)} + \frac{1}{2}\tan^{-1}x\right] + C \\ &= \frac{x}{4(x^2+1)^2} + \frac{3x}{8(x^2+1)} + \frac{3}{8}\tan^{-1}x + C \end{aligned}$$

77. If $a \neq 0$ and n is a positive integer, then $f(x) = \dfrac{1}{x^n(x-a)} = \dfrac{A_1}{x} + \dfrac{A_2}{x^2} + \cdots + \dfrac{A_n}{x^n} + \dfrac{B}{x-a}$. Multiply both sides by $x^n(x-a)$ to get $1 = A_1x^{n-1}(x-a) + A_2x^{n-2}(x-a) + \cdots + A_n(x-a) + Bx^n$. Let $x = a$ in the last equation to

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

get $1 = Ba^n \quad \Rightarrow \quad B = 1/a^n$. So

$$\begin{aligned} f(x) - \frac{B}{x-a} &= \frac{1}{x^n(x-a)} - \frac{1}{a^n(x-a)} = \frac{a^n - x^n}{x^n a^n (x-a)} = -\frac{x^n - a^n}{a^n x^n (x-a)} \\ &= -\frac{(x-a)(x^{n-1} + x^{n-2}a + x^{n-3}a^2 + \cdots + xa^{n-2} + a^{n-1})}{a^n x^n (x-a)} \\ &= -\left(\frac{x^{n-1}}{a^n x^n} + \frac{x^{n-2}a}{a^n x^n} + \frac{x^{n-3}a^2}{a^n x^n} + \cdots + \frac{xa^{n-2}}{a^n x^n} + \frac{a^{n-1}}{a^n x^n}\right) \\ &= -\frac{1}{a^n x} - \frac{1}{a^{n-1}x^2} - \frac{1}{a^{n-2}x^3} - \cdots - \frac{1}{a^2 x^{n-1}} - \frac{1}{ax^n} \end{aligned}$$

Thus, $f(x) = \dfrac{1}{x^n(x-a)} = -\dfrac{1}{a^n x} - \dfrac{1}{a^{n-1}x^2} - \cdots - \dfrac{1}{ax^n} + \dfrac{1}{a^n(x-a)}$.

78. Let $f(x) = ax^2 + bx + c$. We calculate the partial fraction decomposition of $\dfrac{f(x)}{x^2(x+1)^3}$. Since $f(0) = 1$, we must have $c = 1$, so $\dfrac{f(x)}{x^2(x+1)^3} = \dfrac{ax^2+bx+1}{x^2(x+1)^3} = \dfrac{A}{x} + \dfrac{B}{x^2} + \dfrac{C}{x+1} + \dfrac{D}{(x+1)^2} + \dfrac{E}{(x+1)^3}$. Now in order for the integral not to contain any logarithms (that is, in order for it to be a rational function), we must have $A = C = 0$, so $ax^2 + bx + 1 = B(x+1)^3 + Dx^2(x+1) + Ex^2$. Equating constant terms gives $B = 1$, then equating coefficients of x gives $3B = b \quad \Rightarrow \quad b = 3$. This is the quantity we are looking for, since $f'(0) = b$.

7.5 Strategy for Integration

1. (a) Let $u = 1 + x^2$, so that $du = 2x\,dx \quad \Rightarrow \quad \frac{1}{2}\,du = x\,dx$. Then,

$$\int \frac{x}{1+x^2}\,dx = \frac{1}{2}\int \frac{1}{u}\,du = \frac{1}{2}\ln|u| + C = \frac{1}{2}\ln(1+x^2) + C$$

Note the absolute value has been omitted in the last step since $1 + x^2 > 0$ for all $x \in \mathbb{R}$.

(b) $\displaystyle\int \frac{1}{1+x^2}\,dx = \tan^{-1}x + C$

(c) $$\begin{aligned} \int \frac{1}{1-x^2}\,dx &= \int \frac{1}{(1+x)(1-x)}\,dx = \frac{1}{2}\int\left(\frac{1}{1+x} + \frac{1}{1-x}\right)dx \qquad [\text{by partial fractions}] \\ &= \tfrac{1}{2}\ln|1+x| - \tfrac{1}{2}\ln|1-x| + C \end{aligned}$$

2. (a) Let $u = x^2 - 1$, so that $du = 2x\,dx \quad \Rightarrow \quad \frac{1}{2}\,du = x\,dx$. Then

$$\int x\sqrt{x^2-1}\,dx = \frac{1}{2}\int u^{1/2}\,du = \frac{1}{3}u^{3/2} + C = \frac{1}{3}(x^2-1)^{3/2} + C.$$

(b) Let $x = \sec\theta$ where $0 \le \theta < \pi/2$. Then $dx = \sec\theta\tan\theta\,d\theta$ and $\sqrt{x^2-1} = \sqrt{\sec^2\theta - 1} = \sqrt{\tan^2\theta} = |\tan\theta| = \tan\theta$.

Thus, $\displaystyle\int \frac{1}{x\sqrt{x^2-1}}\,dx = \int \frac{\sec\theta\tan\theta}{\sec\theta\tan\theta}\,d\theta = \int d\theta = \theta + C = \sec^{-1}x + C.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) Let $x = \sec\theta$ where $0 \le \theta < \pi/2$. Then $dx = \sec\theta\tan\theta\,d\theta$ and

$\sqrt{x^2-1} = \sqrt{\sec^2\theta - 1} = \sqrt{\tan^2\theta} = |\tan\theta| = \tan\theta$. Thus,

$$\int \frac{\sqrt{x^2-1}}{x}\,dx = \int \frac{\tan\theta}{\sec\theta}\sec\theta\tan\theta\,d\theta = \int \tan^2\theta\,d\theta = \int (\sec^2\theta - 1)\,d\theta$$
$$= \tan\theta - \theta + C = \sqrt{x^2-1} - \sec^{-1}x + C$$

3. (a) Let $u = \ln x$, so that $du = \dfrac{1}{x}\,dx$. Then $\displaystyle\int \frac{\ln x}{x}dx = \int u\,du = \frac{1}{2}u^2 + C = \frac{1}{2}(\ln x)^2 + C$.

(b) Use integration by parts with $u = \ln(2x)$, $dv = dx \;\Rightarrow\; du = \dfrac{1}{2x}(2)\,dx = \dfrac{1}{x}\,dx$, $v = x$. Then

$$\int \ln(2x)\,dx = x\ln(2x) - \int x\left(\frac{1}{x}\,dx\right) = x\ln(2x) - \int dx = x\ln(2x) - x + C.$$

(c) Use integration by parts with $u = \ln x$, $dv = x\,dx \;\Rightarrow\; du = \dfrac{1}{x}\,dx$, $v = \dfrac{1}{2}x^2$. Then

$$\int x\ln x\,dx = \frac{1}{2}x^2\ln x - \int \frac{1}{2}x^2\left(\frac{1}{x}\,dx\right) = \frac{1}{2}x^2\ln x - \frac{1}{2}\int x\,dx = \frac{1}{2}x^2\ln x - \frac{1}{4}x^2 + C.$$

4. (a) $\displaystyle\int \sin^2 x\,dx = \int \tfrac{1}{2}(1-\cos 2x)\,dx = \tfrac{1}{2}\left(x - \tfrac{1}{2}\sin 2x\right) + C = \tfrac{1}{2}x - \tfrac{1}{4}\sin 2x + C$

(b) $\displaystyle\int \sin^3 x\,dx = \int \sin^2 x\sin x\,dx = \int \left(1-\cos^2 x\right)\sin x\,dx \overset{c}{=} \int (1-u^2)\,(-du)$

$$= -\left(u - \tfrac{1}{3}u^3\right) + C = -\cos x + \tfrac{1}{3}\cos^3 x + C$$

(c) Let $u = 2x$ so that $du = 2\,dx$. Then $\int \sin 2x\,dx = \int \sin u\left(\frac{1}{2}\,du\right) = \frac{1}{2}(-\cos u) + C = -\frac{1}{2}\cos 2x + C$.

5. (a) $\dfrac{1}{x^2-4x+3} = \dfrac{1}{(x-3)(x-1)} = \dfrac{A}{x-3} + \dfrac{B}{x-1}$. Multiply both sides by $(x-3)(x-1)$ to get

$1 = A(x-1) + B(x-3)$. Setting $x = 3$ gives $1 = 2A$, so $A = \frac{1}{2}$. Now setting $x = 1$ gives $1 = -2B$, so $B = -\frac{1}{2}$.

Thus, $\displaystyle\int \frac{1}{x^2-4x+3}\,dx = \frac{1}{2}\int \frac{1}{x-3} - \frac{1}{x-1}\,dx = \tfrac{1}{2}\ln|x-3| - \tfrac{1}{2}\ln|x-1| + C$.

(b) $\dfrac{1}{x^2-4x+4} = \dfrac{1}{(x-2)^2}$. Let $u = x-2$, so that $du = dx$. Thus,

$$\int \frac{1}{x^2-4x+4}\,dx = \int \frac{1}{(x-2)^2}\,dx = \int \frac{1}{u^2}\,du = \int u^{-2}\,du = -u^{-1} + C = -\frac{1}{x-2} + C.$$

(c) $x^2 - 4x + 5$ is an irreducible quadratic, so it cannot be factored. Completing the square gives

$x^2 - 4x + 5 = (x^2 - 4x + 4) - 4 + 5 = (x-2)^2 + 1$. Now, use the substitution $u = x - 2$, so that $du = dx$. Thus,

$$\int \frac{1}{x^2-4x+5}\,dx = \int \frac{1}{(x-2)^2+1}\,dx = \int \frac{1}{u^2+1}\,du = \tan^{-1}u + C = \tan^{-1}(x-2) + C.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. (a) Let $u = x^2$, so that $du = 2x\,dx \quad \Rightarrow \quad \frac{1}{2}\,du = x\,dx$. Thus,

$$\int x\cos x^2\,dx = \frac{1}{2}\int \cos u\,du = \frac{1}{2}\sin u + C = \frac{1}{2}\sin x^2 + C.$$

(b) $\int x\cos^2 x\,dx = \frac{1}{2}\int x(1+\cos 2x)\,dx = \frac{1}{2}\int x\,dx + \frac{1}{2}\int x\cos 2x\,dx = \frac{1}{4}x^2 + \frac{1}{2}\int x\cos 2x\,dx$.

The remaining integral can be evaluated using integration by parts with $u = x$, $dv = \cos 2x\,dx \quad \Rightarrow$ $du = dx$, $v = \frac{1}{2}\sin 2x$. Thus,

$$\begin{aligned}\int x\cos^2 x\,dx &= \tfrac{1}{4}x^2 + \tfrac{1}{2}\left(\tfrac{1}{2}x\sin 2x - \int \tfrac{1}{2}\sin 2x\,dx\right) = \tfrac{1}{4}x^2 + \tfrac{1}{2}\left(\tfrac{1}{2}x\sin 2x + \tfrac{1}{4}\cos 2x\right) + C\\ &= \tfrac{1}{4}x^2 + \tfrac{1}{4}x\sin 2x + \tfrac{1}{8}\cos 2x + C\end{aligned}$$

(c) First, use integration by parts with $u = x^2$, $dv = \cos x\,dx \quad \Rightarrow \quad du = 2x\,dx$, $v = \sin x$. This gives $I = \int x^2\cos x\,dx = x^2\sin x - \int 2x\sin x\,dx$. Next, use integration by parts for the remaining integral with $U = 2x$, $dV = \sin x\,dx \quad \Rightarrow \quad dU = 2\,dx$, $V = -\cos x$. Thus, $I = x^2\sin x - \left(-2x\cos x + \int 2\cos x\,dx\right) = x^2\sin x + 2x\cos x - 2\sin x + C$.

7. (a) Let $u = x^3$, so that $du = 3x^2\,dx \quad \Rightarrow \quad \frac{1}{3}\,du = x^2\,dx$. Thus, $\displaystyle\int x^2 e^{x^3}\,dx = \frac{1}{3}\int e^u\,du = \frac{1}{3}e^u + C = \frac{1}{3}e^{x^3} + C$.

(b) First, use integration by parts with $u = x^2$, $dv = e^x\,dx \quad \Rightarrow \quad du = 2x\,dx$, $v = e^x$. This gives $I = \int x^2 e^x\,dx = x^2e^x - \int 2xe^x\,dx$. Next, use integration by parts for the remaining integral with $U = 2x$, $dV = e^x\,dx \quad \Rightarrow \quad dU = 2\,dx$, $V = e^x$. Thus, $I = x^2e^x - \left(2xe^x - \int 2e^x\,dx\right) = x^2e^x - 2xe^x + 2e^x + C$.

(c) Let $y = x^2$, so that $dy = 2x\,dx$. Thus, $\int x^3 e^{x^2}\,dx = \int x^2 e^{x^2} x\,dx = \frac{1}{2}\int ye^y\,dy$. Now use integration by parts with $u = y$, $dv = e^y \quad \Rightarrow \quad du = dy$, $v = e^y$. This gives $\int x^3 e^{x^2}\,dx = \frac{1}{2}\left(ye^y - \int e^y dy\right) = \frac{1}{2}ye^y - \frac{1}{2}e^y + C = \frac{1}{2}x^2e^{x^2} - \frac{1}{2}e^{x^2} + C$.

8. (a) Let $u = e^x - 1$, so that $du = e^x\,dx$. Thus, $\int e^x\sqrt{e^x - 1}\,dx = \int u^{1/2}\,du = \frac{2}{3}u^{3/2} + C = \frac{2}{3}(e^x - 1)^{3/2} + C$.

(b) Let $u = e^x$, so that $du = e^x\,dx$. Thus, $\displaystyle\int \frac{e^x}{\sqrt{1 - e^{2x}}}\,dx = \int \frac{1}{\sqrt{1 - u^2}}\,du = \sin^{-1} u + C = \sin^{-1}(e^x) + C$.

(c) Let $u = \sqrt{e^x - 1}$, so that $u^2 = e^x - 1 \quad \Rightarrow \quad 2u\,du = e^x\,dx$, and $\dfrac{2u\,du}{u^2 + 1} = dx$. Then

$$\int \frac{1}{\sqrt{e^x - 1}}\,dx = \int \frac{1}{u}\frac{2u\,du}{u^2 + 1} = 2\int \frac{1}{u^2 + 1}\,du = 2\tan^{-1} u + C = 2\tan^{-1}\sqrt{e^x - 1} + C.$$

9. Let $u = 1 - \sin x$. Then $du = -\cos x\,dx \quad \Rightarrow$

$$\int \frac{\cos x}{1 - \sin x}\,dx = \int \frac{1}{u}(-du) = -\ln|u| + C = -\ln|1 - \sin x| + C = -\ln(1 - \sin x) + C.$$

10. Let $u = 3x + 1$. Then $du = 3\,dx \quad \Rightarrow$

$$\int_0^1 (3x + 1)^{\sqrt{2}}\,dx = \int_1^4 u^{\sqrt{2}}\left(\frac{1}{3}\,du\right) = \frac{1}{3}\left[\frac{1}{\sqrt{2} + 1}\,u^{\sqrt{2}+1}\right]_1^4 = \frac{1}{3(\sqrt{2} + 1)}\left(4^{\sqrt{2}+1} - 1\right).$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. Let $u = \ln y$, $dv = \sqrt{y}\,dy \quad \Rightarrow \quad du = \dfrac{1}{y}\,dy$, $v = \dfrac{2}{3}y^{3/2}$. Then

$$\int_1^4 \sqrt{y}\ln y\,dy = \left[\frac{2}{3}y^{3/2}\ln y\right]_1^4 - \int_1^4 \frac{2}{3}y^{1/2}\,dy = \frac{2}{3}\cdot 8\ln 4 - 0 - \left[\frac{4}{9}y^{3/2}\right]_1^4$$
$$= \tfrac{16}{3}(2\ln 2) - \left(\tfrac{4}{9}\cdot 8 - \tfrac{4}{9}\right) = \tfrac{32}{3}\ln 2 - \tfrac{28}{9}$$

12. Let $u = \arcsin x$, so that $du = \dfrac{1}{\sqrt{1-x^2}}\,dx$. Thus, $\displaystyle\int \frac{e^{\arcsin x}}{\sqrt{1-x^2}}\,dx = \int e^u\,du = e^u + C = e^{\arcsin x} + C$.

13. Let $x = \ln y$, so that $dx = \dfrac{1}{y}\,dy$. Thus, $I = \displaystyle\int \frac{\ln(\ln y)}{y}dy = \int \ln x\,dx$. Now use integration by parts with

$u = \ln x$, $dv = dx \quad \Rightarrow \quad du = dx/x$, $v = x$. This gives

$$I = x\ln x - \int \frac{x}{x}\,dx = x\ln x - \int dx = x\ln x - x + C = \ln y\,[\ln(\ln y)] - \ln y + C.$$

14. Let $u = 2x+1$. Then $du = 2\,dx \quad \Rightarrow$

$$\int_0^1 \frac{x}{(2x+1)^3}\,dx = \int_1^3 \frac{(u-1)/2}{u^3}\left(\frac{1}{2}\,du\right) = \frac{1}{4}\int_1^3\left(\frac{1}{u^2} - \frac{1}{u^3}\right)du = \frac{1}{4}\left[-\frac{1}{u} + \frac{1}{2u^2}\right]_1^3$$
$$= \tfrac{1}{4}\left[\left(-\tfrac{1}{3} + \tfrac{1}{18}\right) - \left(-1 + \tfrac{1}{2}\right)\right] = \tfrac{1}{4}\left(\tfrac{2}{9}\right) = \tfrac{1}{18}$$

15. Let $u = x^2$, so that $du = 2x\,dx$. Thus,

$$\int \frac{x}{x^4+9}\,dx = \frac{1}{2}\int \frac{1}{u^2+9}\,du = \frac{1}{2}\int \frac{(1/3)^2}{(u/3)^2+1}\,du = \frac{1}{2}\left(\frac{1}{3}\tan^{-1}\left(\frac{u}{3}\right)\right) + C = \frac{1}{6}\tan^{-1}\left(\frac{x^2}{3}\right) + C.$$

16. $\int t\sin t\,\cos t\,dt = \int t\cdot\frac{1}{2}(2\sin t\,\cos t)\,dt = \frac{1}{2}\int t\sin 2t\,dt$

$$= \tfrac{1}{2}\left(-\tfrac{1}{2}t\cos 2t - \int -\tfrac{1}{2}\cos 2t\,dt\right) \qquad \begin{bmatrix} u = t, & dv = \sin 2t\,dt \\ du = dt, & v = -\frac{1}{2}\cos 2t \end{bmatrix}$$
$$= -\tfrac{1}{4}t\cos 2t + \tfrac{1}{4}\int \cos 2t\,dt = -\tfrac{1}{4}t\cos 2t + \tfrac{1}{8}\sin 2t + C$$

17. $\dfrac{x+2}{x^2+3x-4} = \dfrac{x+2}{(x+4)(x-1)} = \dfrac{A}{x+4} + \dfrac{B}{x-1}$. Multiply by $(x+4)(x-1)$ to get $x+2 = A(x-1) + B(x+4)$.

Substituting 1 for x gives $3 = 5B \quad \Leftrightarrow \quad B = \frac{3}{5}$. Substituting -4 for x gives $-2 = -5A \quad \Leftrightarrow \quad A = \frac{2}{5}$. Thus,

$$\int_2^4 \frac{x+2}{x^2+3x-4}\,dx = \int_2^4\left(\frac{2/5}{x+4} + \frac{3/5}{x-1}\right)dx = \left[\frac{2}{5}\ln|x+4| + \frac{3}{5}\ln|x-1|\right]_2^4$$
$$= \left(\tfrac{2}{5}\ln 8 + \tfrac{3}{5}\ln 3\right) - \left(\tfrac{2}{5}\ln 6 + 0\right) = \tfrac{2}{5}(3\ln 2) + \tfrac{3}{5}\ln 3 - \tfrac{2}{5}(\ln 2 + \ln 3)$$
$$= \tfrac{4}{5}\ln 2 + \tfrac{1}{5}\ln 3, \text{ or } \tfrac{1}{5}\ln 48$$

18. Let $u = \dfrac{1}{x}$, $dv = \dfrac{\cos(1/x)}{x^2} \quad \Rightarrow \quad du = -\dfrac{1}{x^2}\,dx$, $v = -\sin\left(\dfrac{1}{x}\right)$. Then

$$\int \frac{\cos(1/x)}{x^3}\,dx = -\frac{1}{x}\sin\left(\frac{1}{x}\right) - \int \frac{1}{x^2}\sin\left(\frac{1}{x}\right)dx = -\frac{1}{x}\sin\left(\frac{1}{x}\right) - \cos\left(\frac{1}{x}\right) + C.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. Let $x = \sec\theta$, where $0 \le \theta \le \frac{\pi}{2}$ or $\pi \le \theta < \frac{3\pi}{2}$. Then $dx = \sec\theta\,\tan\theta\,d\theta$ and

$\sqrt{x^2-1} = \sqrt{\sec^2\theta - 1} = \sqrt{\tan^2\theta} = |\tan\theta| = \tan\theta$ for the relevant values of θ, so

$$\int \frac{1}{x^3\sqrt{x^2-1}}\,dx = \int \frac{\sec\theta\,\tan\theta}{\sec^3\theta\,\tan\theta}\,d\theta = \int \cos^2\theta\,d\theta = \int \tfrac{1}{2}(1+\cos 2\theta)\,d\theta$$

$$= \tfrac{1}{2}\theta + \tfrac{1}{4}\sin 2\theta + C = \tfrac{1}{2}\theta + \tfrac{1}{2}\sin\theta\,\cos\theta + C$$

$$= \frac{1}{2}\sec^{-1}x + \frac{1}{2}\frac{\sqrt{x^2-1}}{x}\,\frac{1}{x} + C = \frac{1}{2}\sec^{-1}x + \frac{\sqrt{x^2-1}}{2x^2} + C$$

20. $\dfrac{2x-3}{x^3+3x} = \dfrac{2x-3}{x(x^2+3)} = \dfrac{A}{x} + \dfrac{Bx+C}{x^2+3}$. Multiply by $x(x^2+3)$ to get $2x-3 = A(x^2+3) + (Bx+C)x \quad \Leftrightarrow$

$2x-3 = (A+B)x^2 + Cx + 3A$. Equating coefficients gives us $C = 2$, $3A = -3 \quad \Leftrightarrow \quad A = -1$, and $A + B = 0$, so $B = 1$. Thus,

$$\int \frac{2x-3}{x^3+3x}\,dx = \int \left(\frac{-1}{x} + \frac{x+2}{x^2+3}\right)dx = \int \left(-\frac{1}{x} + \frac{x}{x^2+3} + \frac{2}{x^2+3}\right)dx$$

$$= -\ln|x| + \frac{1}{2}\ln(x^2+3) + \frac{2}{\sqrt{3}}\tan^{-1}\left(\frac{x}{\sqrt{3}}\right) + C$$

21. $\displaystyle\int \frac{\cos^3 x}{\csc x}\,dx = \int \cos^3 x\sin x\,dx \overset{c}{=} \int u^3\,(-du) = -\frac{1}{4}u^4 + C = -\frac{1}{4}\cos^4 x + C$

22. Let $u = \ln(1+x^2)$, $dv = dx \quad \Rightarrow \quad du = \dfrac{2x}{1+x^2}\,dx$, $v = x$. Then

$$\int \ln(1+x^2)\,dx = x\ln(1+x^2) - \int \frac{2x^2}{1+x^2}\,dx = x\ln(1+x^2) - 2\int \frac{(x^2+1)-1}{1+x^2}\,dx$$

$$= x\ln(1+x^2) - 2\int \left(1 - \frac{1}{1+x^2}\right)dx = x\ln(1+x^2) - 2x + 2\tan^{-1}x + C$$

23. Let $u = x$, $dv = \sec x\,\tan x\,dx \quad \Rightarrow \quad du = dx$, $v = \sec x$. Then

$\int x\sec x\,\tan x\,dx = x\sec x - \int \sec x\,dx = x\sec x - \ln|\sec x + \tan x| + C$.

24. $\displaystyle\int_0^{\sqrt{2}/2} \frac{x^2}{\sqrt{1-x^2}}\,dx = \int_0^{\pi/4} \frac{\sin^2\theta}{\cos\theta}\cos\theta\,d\theta \qquad \begin{bmatrix} u = \sin\theta, \\ du = \cos\theta\,d\theta \end{bmatrix}$

$$= \int_0^{\pi/4} \tfrac{1}{2}(1-\cos 2\theta)\,d\theta = \tfrac{1}{2}\left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/4} = \tfrac{1}{2}\left[\left(\tfrac{\pi}{4} - \tfrac{1}{2}\right) - (0-0)\right] = \tfrac{\pi}{8} - \tfrac{1}{4}$$

25. $\int_0^{\pi} t\,\cos^2 t\,dt = \int_0^{\pi} t\left[\frac{1}{2}(1+\cos 2t)\right]dt = \frac{1}{2}\int_0^{\pi} t\,dt + \frac{1}{2}\int_0^{\pi} t\cos 2t\,dt$

$$= \tfrac{1}{2}\left[\tfrac{1}{2}t^2\right]_0^{\pi} + \tfrac{1}{2}\left[\tfrac{1}{2}t\sin 2t\right]_0^{\pi} - \tfrac{1}{2}\int_0^{\pi} \tfrac{1}{2}\sin 2t\,dt \qquad \begin{bmatrix} u = t, & dv = \cos 2t\,dt \\ du = dt, & v = \frac{1}{2}\sin 2t \end{bmatrix}$$

$$= \tfrac{1}{4}\pi^2 + 0 - \tfrac{1}{4}\left[-\tfrac{1}{2}\cos 2t\right]_0^{\pi} = \tfrac{1}{4}\pi^2 + \tfrac{1}{8}(1-1) = \tfrac{1}{4}\pi^2$$

26. Let $u = \sqrt{t}$. Then $du = \dfrac{1}{2\sqrt{t}}\,dt \quad \Rightarrow \quad \displaystyle\int_1^4 \frac{e^{\sqrt{t}}}{\sqrt{t}}\,dt = \int_1^2 e^u\,(2\,du) = 2\left[e^u\right]_1^2 = 2(e^2 - e)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. Let $u = e^x$. Then $\int e^{x+e^x}\,dx = \int e^{e^x} e^x\,dx = \int e^u\,du = e^u + C = e^{e^x} + C$.

28. $\displaystyle\int \frac{e^x}{1+e^{2a}}\,dx = \frac{1}{1+e^{2a}}\int e^x\,dx = \frac{e^x}{1+e^{2a}} + C$

29. Let $t = \sqrt{x}$, so that $t^2 = x$ and $2t\,dt = dx$. Then $\int \arctan\sqrt{x}\,dx = \int \arctan t\,(2t\,dt) = I$. Now use parts with

$u = \arctan t$, $dv = 2t\,dt \quad\Rightarrow\quad du = \dfrac{1}{1+t^2}\,dt$, $v = t^2$. Thus,

$$I = t^2\arctan t - \int \frac{t^2}{1+t^2}\,dt = t^2\arctan t - \int\left(1 - \frac{1}{1+t^2}\right)dt = t^2\arctan t - t + \arctan t + C$$

$$= x\arctan\sqrt{x} - \sqrt{x} + \arctan\sqrt{x} + C \quad \left[\text{or } (x+1)\arctan\sqrt{x} - \sqrt{x} + C\right]$$

30. Let $u = 1 + (\ln x)^2$, so that $du = \dfrac{2\ln x}{x}\,dx$. Then

$$\int \frac{\ln x}{x\sqrt{1+(\ln x)^2}}\,dx = \frac{1}{2}\int \frac{1}{\sqrt{u}}\,du = \frac{1}{2}\left(2\sqrt{u}\right) + C = \sqrt{1+(\ln x)^2} + C.$$

31. Let $u = 1 + \sqrt{x}$. Then $x = (u-1)^2$, $dx = 2(u-1)\,du \quad\Rightarrow$

$$\int_0^1 \left(1+\sqrt{x}\right)^8 dx = \int_1^2 u^8 \cdot 2(u-1)\,du = 2\int_1^2 (u^9 - u^8)\,du$$

$$= \left[\tfrac{1}{5}u^{10} - 2\cdot\tfrac{1}{9}u^9\right]_1^2 = \tfrac{1024}{5} - \tfrac{1024}{9} - \tfrac{1}{5} + \tfrac{2}{9} = \tfrac{4097}{45}$$

32. $\int (1+\tan x)^2 \sec x\,dx = \int (1 + 2\tan x + \tan^2 x)\sec x\,dx$

$$= \int [\sec x + 2\sec x\,\tan x + (\sec^2 x - 1)\sec x]\,dx = \int (2\sec x\,\tan x + \sec^3 x)\,dx$$

$$= 2\sec x + \tfrac{1}{2}(\sec x\,\tan x + \ln|\sec x + \tan x|) + C \quad \text{[by Example 7.2.8]}$$

33. $\displaystyle\int_0^1 \frac{1+12t}{1+3t}\,dt = \int_0^1 \frac{(12t+4)-3}{3t+1}\,dt = \int_0^1 \left(4 - \frac{3}{3t+1}\right)dt = \Big[4t - \ln|3t+1|\Big]_0^1$

$$= (4 - \ln 4) - (0 - 0) = 4 - \ln 4$$

34. $\dfrac{3x^2+1}{x^3+x^2+x+1} = \dfrac{3x^2+1}{(x^2+1)(x+1)} = \dfrac{A}{x+1} + \dfrac{Bx+C}{x^2+1}$. Multiply by $(x+1)(x^2+1)$ to get

$3x^2 + 1 = A(x^2+1) + (Bx+C)(x+1) \quad\Leftrightarrow\quad 3x^2 + 1 = (A+B)x^2 + (B+C)x + (A+C)$. Substituting -1 for x gives $4 = 2A \quad\Leftrightarrow\quad A = 2$. Equating coefficients of x^2 gives $3 = A + B = 2 + B \quad\Leftrightarrow\quad B = 1$. Equating coefficients of x gives $0 = B + C = 1 + C \quad\Leftrightarrow\quad C = -1$. Thus,

$$\int_0^1 \frac{3x^2+1}{x^3+x^2+x+1}\,dx = \int_0^1 \left(\frac{2}{x+1} + \frac{x-1}{x^2+1}\right)dx = \int_0^1 \left(\frac{2}{x+1} + \frac{x}{x^2+1} - \frac{1}{x^2+1}\right)dx$$

$$= \left[2\ln|x+1| + \tfrac{1}{2}\ln(x^2+1) - \tan^{-1}x\right]_0^1 = (2\ln 2 + \tfrac{1}{2}\ln 2 - \tfrac{\pi}{4}) - (0 + 0 - 0)$$

$$= \tfrac{5}{2}\ln 2 - \tfrac{\pi}{4}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. Let $u = 1 + e^x$, so that $du = e^x\,dx = (u-1)\,dx$. Then $\displaystyle\int \frac{1}{1+e^x}\,dx = \int \frac{1}{u}\cdot\frac{du}{u-1} = \int \frac{1}{u(u-1)}\,du = I$. Now

$\displaystyle\frac{1}{u(u-1)} = \frac{A}{u} + \frac{B}{u-1} \quad\Rightarrow\quad 1 = A(u-1) + Bu$. Set $u = 1$ to get $1 = B$. Set $u = 0$ to get $1 = -A$, so $A = -1$.

Thus, $\displaystyle I = \int\left(\frac{-1}{u} + \frac{1}{u-1}\right)du = -\ln|u| + \ln|u-1| + C = -\ln(1+e^x) + \ln e^x + C = x - \ln(1+e^x) + C$.

Another method: Multiply numerator and denominator by e^{-x} and let $u = e^{-x} + 1$. This gives the answer in the form $-\ln(e^{-x}+1) + C$.

36. $\displaystyle\int \sin\sqrt{at}\,dt = \int \sin u \cdot \tfrac{2}{a}u\,du \quad [u = \sqrt{at},\ u^2 = at,\ 2u\,du = a\,dt] \quad = \tfrac{2}{a}\int u\sin u\,du$

$\displaystyle = \tfrac{2}{a}[-u\cos u + \sin u] + C$ [integration by parts] $\displaystyle = -\tfrac{2}{a}\sqrt{at}\,\cos\sqrt{at} + \tfrac{2}{a}\,\sin\sqrt{at} + C$

$\displaystyle = -2\sqrt{\tfrac{t}{a}}\,\cos\sqrt{at} + \tfrac{2}{a}\,\sin\sqrt{at} + C$

37. Use integration by parts with $u = \ln\left(x + \sqrt{x^2-1}\right)$, $dv = dx \quad\Rightarrow$

$\displaystyle du = \frac{1}{x+\sqrt{x^2-1}}\left(1 + \frac{x}{\sqrt{x^2-1}}\right)dx = \frac{1}{x+\sqrt{x^2-1}}\left(\frac{\sqrt{x^2-1}+x}{\sqrt{x^2-1}}\right)dx = \frac{1}{\sqrt{x^2-1}}\,dx$, $v = x$. Then

$\displaystyle\int \ln\left(x+\sqrt{x^2-1}\right)dx = x\ln\left(x+\sqrt{x^2-1}\right) - \int\frac{x}{\sqrt{x^2-1}}\,dx = x\ln\left(x+\sqrt{x^2-1}\right) - \sqrt{x^2-1} + C$.

38. $\displaystyle|e^x - 1| = \begin{cases} e^x - 1 & \text{if } e^x - 1 \ge 0 \\ -(e^x-1) & \text{if } e^x - 1 < 0 \end{cases} = \begin{cases} e^x - 1 & \text{if } x \ge 0 \\ 1 - e^x & \text{if } x < 0 \end{cases}$

Thus, $\displaystyle\int_{-1}^{2}|e^x-1|\,dx = \int_{-1}^{0}(1-e^x)\,dx + \int_0^2(e^x-1)\,dx = \Big[x - e^x\Big]_{-1}^{0} + \Big[e^x - x\Big]_0^2$

$\displaystyle = (0-1) - (-1-e^{-1}) + (e^2-2) - (1-0) = e^2 + e^{-1} - 3$

39. As in Example 5,

$$\int\sqrt{\frac{1+x}{1-x}}\,dx = \int\frac{\sqrt{1+x}}{\sqrt{1-x}}\cdot\frac{\sqrt{1+x}}{\sqrt{1+x}}\,dx = \int\frac{1+x}{\sqrt{1-x^2}}\,dx = \int\frac{dx}{\sqrt{1-x^2}} + \int\frac{x\,dx}{\sqrt{1-x^2}} = \sin^{-1}x - \sqrt{1-x^2} + C.$$

Another method: Substitute $u = \sqrt{(1+x)/(1-x)}$.

40. $\displaystyle\int_1^3 \frac{e^{3/x}}{x^2}\,dx = \int_3^1 e^u\left(-\tfrac{1}{3}\,du\right) \quad \begin{bmatrix} u = 3/x, \\ du = -3/x^2\,dx \end{bmatrix} \quad = -\tfrac{1}{3}\Big[e^u\Big]_3^1 = -\tfrac{1}{3}(e - e^3) = \tfrac{1}{3}(e^3 - e)$

41. $3 - 2x - x^2 = -(x^2+2x+1) + 4 = 4 - (x+1)^2$. Let $x + 1 = 2\sin\theta$, where $-\frac{\pi}{2} \le \theta \le \frac{\pi}{2}$. Then $dx = 2\cos\theta\,d\theta$ and

$\displaystyle\int\sqrt{3-2x-x^2}\,dx = \int\sqrt{4-(x+1)^2}\,dx = \int\sqrt{4-4\sin^2\theta}\,2\cos\theta\,d\theta$

$\displaystyle = 4\int\cos^2\theta\,d\theta = 2\int(1+\cos 2\theta)\,d\theta$

$\displaystyle = 2\theta + \sin 2\theta + C = 2\theta + 2\sin\theta\cos\theta + C$

$\displaystyle = 2\sin^{-1}\left(\frac{x+1}{2}\right) + 2\cdot\frac{x+1}{2}\cdot\frac{\sqrt{3-2x-x^2}}{2} + C$

$\displaystyle = 2\sin^{-1}\left(\frac{x+1}{2}\right) + \frac{x+1}{2}\sqrt{3-2x-x^2} + C$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. $\displaystyle\int_{\pi/4}^{\pi/2} \frac{1+4\cot x}{4-\cot x}\,dx = \int_{\pi/4}^{\pi/2} \left[\frac{(1+4\cos x/\sin x)}{(4-\cos x/\sin x)}\cdot\frac{\sin x}{\sin x}\right] dx = \int_{\pi/4}^{\pi/2} \frac{\sin x + 4\cos x}{4\sin x - \cos x}\,dx$

$$= \int_{3/\sqrt{2}}^{4} \frac{1}{u}\,du \quad \left[\begin{array}{l} u = 4\sin x - \cos x, \\ du = (4\cos x + \sin x)\,dx \end{array}\right]$$

$$= \Big[\ln|u|\Big]_{3/\sqrt{2}}^{4} = \ln 4 - \ln\frac{3}{\sqrt{2}} = \ln\frac{4}{3/\sqrt{2}} = \ln\left(\frac{4}{3}\sqrt{2}\right)$$

43. The integrand is an odd function, so $\displaystyle\int_{-\pi/2}^{\pi/2} \frac{x}{1+\cos^2 x}\,dx = 0$ [by 5.5.7].

44. $\displaystyle\int \frac{1+\sin x}{1+\cos x}\,dx = \int \frac{(1+\sin x)(1-\cos x)}{(1+\cos x)(1-\cos x)}\,dx = \int \frac{1-\cos x+\sin x-\sin x\,\cos x}{\sin^2 x}\,dx$

$$= \int \left(\csc^2 x - \frac{\cos x}{\sin^2 x} + \csc x - \frac{\cos x}{\sin x}\right) dx$$

$$\overset{\text{s}}{=} -\cot x + \frac{1}{\sin x} + \ln|\csc x - \cot x| - \ln|\sin x| + C \quad \text{[by Exercise 7.2.41]}$$

The answer can be written as $\dfrac{1-\cos x}{\sin x} - \ln(1+\cos x) + C$.

45. Let $u = \tan\theta$. Then $du = \sec^2\theta\,d\theta \;\Rightarrow\; \int_0^{\pi/4} \tan^3\theta\,\sec^2\theta\,d\theta = \int_0^1 u^3\,du = \left[\frac{1}{4}u^4\right]_0^1 = \frac{1}{4}$.

46. $\displaystyle\int_{\pi/6}^{\pi/3} \frac{\sin\theta\,\cot\theta}{\sec\theta}\,d\theta = \int_{\pi/6}^{\pi/3} \cos^2\theta\,d\theta = \frac{1}{2}\int_{\pi/6}^{\pi/3} (1+\cos 2\theta)\,d\theta = \frac{1}{2}\left[\theta + \frac{1}{2}\sin 2\theta\right]_{\pi/6}^{\pi/3}$

$$= \frac{1}{2}\left[\left(\frac{\pi}{3}+\frac{\sqrt{3}}{4}\right) - \left(\frac{\pi}{6}+\frac{\sqrt{3}}{4}\right)\right] = \frac{1}{2}\left(\frac{\pi}{6}\right) = \frac{\pi}{12}$$

47. Let $u = \sec\theta$, so that $du = \sec\theta\,\tan\theta\,d\theta$. Then $\displaystyle\int \frac{\sec\theta\,\tan\theta}{\sec^2\theta - \sec\theta}\,d\theta = \int \frac{1}{u^2-u}\,du = \int \frac{1}{u(u-1)}\,du = I$. Now

$\dfrac{1}{u(u-1)} = \dfrac{A}{u} + \dfrac{B}{u-1} \;\Rightarrow\; 1 = A(u-1) + Bu$. Set $u = 1$ to get $1 = B$. Set $u = 0$ to get $1 = -A$, so $A = -1$.

Thus, $I = \displaystyle\int \left(\frac{-1}{u} + \frac{1}{u-1}\right) du = -\ln|u| + \ln|u-1| + C = \ln|\sec\theta - 1| - \ln|\sec\theta| + C$ [or $\ln|1-\cos\theta| + C$].

48. Using product formula 2(a) in Section 7.2, $\sin 6x\,\cos 3x = \frac{1}{2}[\sin(6x-3x) + \sin(6x+3x)] = \frac{1}{2}(\sin 3x + \sin 9x)$. Thus,

$$\int_0^\pi \sin 6x\,\cos 3x\,dx = \int_0^\pi \tfrac{1}{2}(\sin 3x + \sin 9x)\,dx = \tfrac{1}{2}\left[-\tfrac{1}{3}\cos 3x - \tfrac{1}{9}\cos 9x\right]_0^\pi$$

$$= \tfrac{1}{2}\left[\left(\tfrac{1}{3}+\tfrac{1}{9}\right) - \left(-\tfrac{1}{3}-\tfrac{1}{9}\right)\right] = \tfrac{1}{2}\left(\tfrac{4}{9}+\tfrac{4}{9}\right) = \tfrac{4}{9}$$

49. Let $u = \theta$, $dv = \tan^2\theta\,d\theta = \left(\sec^2\theta - 1\right)d\theta \;\Rightarrow\; du = d\theta$ and $v = \tan\theta - \theta$. So

$$\int \theta\tan^2\theta\,d\theta = \theta(\tan\theta - \theta) - \int(\tan\theta - \theta)\,d\theta = \theta\tan\theta - \theta^2 - \ln|\sec\theta| + \tfrac{1}{2}\theta^2 + C$$

$$= \theta\tan\theta - \tfrac{1}{2}\theta^2 - \ln|\sec\theta| + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50. Let $u = \sqrt{x-1}$, so that $x = u^2 + 1$ and $dx = 2u\,du$. Thus,

$$\int \frac{1}{x\sqrt{x-1}}\,dx = \int \frac{2u}{(u^2+1)u}\,du = 2\int \frac{1}{u^2+1}\,du = 2\tan^{-1}u + C = 2\tan^{-1}\left(\sqrt{x-1}\right) + C.$$

51. Let $u = \sqrt{x}$, so that $du = \dfrac{1}{2\sqrt{x}}\,dx$. Then

$$\int \frac{\sqrt{x}}{1+x^3}\,dx = \int \frac{u}{1+u^6}\,(2u\,du) = 2\int \frac{u^2}{1+(u^3)^2}\,du = 2\int \frac{1}{1+t^2}\left(\frac{1}{3}\,dt\right) \qquad \left[\begin{array}{l} t = u^3 \\ dt = 3u^2\,du \end{array}\right]$$

$$= \tfrac{2}{3}\tan^{-1}t + C = \tfrac{2}{3}\tan^{-1}u^3 + C = \tfrac{2}{3}\tan^{-1}(x^{3/2}) + C$$

Another method: Let $u = x^{3/2}$ so that $u^2 = x^3$ and $du = \frac{3}{2}x^{1/2}\,dx \quad\Rightarrow\quad \sqrt{x}\,dx = \frac{2}{3}\,du$. Then

$$\int \frac{\sqrt{x}}{1+x^3}\,dx = \int \frac{\frac{2}{3}}{1+u^2}\,du = \frac{2}{3}\tan^{-1}u + C = \frac{2}{3}\tan^{-1}(x^{3/2}) + C.$$

52. Let $u = \sqrt{1+e^x}$. Then $u^2 = 1 + e^x$, $2u\,du = e^x\,dx = (u^2-1)\,dx$, and $dx = \dfrac{2u}{u^2-1}du$, so

$$\int \sqrt{1+e^x}\,dx = \int u\cdot\frac{2u}{u^2-1}\,du = \int \frac{2u^2}{u^2-1}\,du = \int\left(2 + \frac{2}{u^2-1}\right)du = \int\left(2 + \frac{1}{u-1} - \frac{1}{u+1}\right)du$$

$$= 2u + \ln|u-1| - \ln|u+1| + C = 2\sqrt{1+e^x} + \ln\left(\sqrt{1+e^x}-1\right) - \ln\left(\sqrt{1+e^x}+1\right) + C$$

53. Let $u = \sqrt{x}$, so that $x = u^2$ and $dx = 2u\,du$. Thus,

$$\int \frac{x}{1+\sqrt{x}}\,dx = \int \frac{u^2}{1+u}\,(2u\,du) = 2\int \frac{u^3}{1+u}\,du = 2\int \frac{\left(u^3+1\right)-1}{u+1}\,du \qquad \text{[or use long division]}$$

$$= 2\int \frac{(u+1)(u^2-u+1)-1}{u+1}\,du = 2\int\left(u^2 - u + 1 - \frac{1}{u+1}\right)du$$

$$= 2\left(\tfrac{1}{3}u^3 - \tfrac{1}{2}u^2 + u - \ln|u+1|\right) + C = \tfrac{2}{3}u^3 - u^2 + 2u - 2\ln|u+1| + C$$

$$= \tfrac{2}{3}x^{3/2} - x + 2\sqrt{x} - 2\ln\left(\sqrt{x}+1\right) + C$$

54. Use integration by parts with $u = (x-1)e^x$, $dv = \dfrac{1}{x^2}\,dx \quad\Rightarrow\quad du = [(x-1)e^x + e^x]\,dx = xe^x\,dx$, $v = -\dfrac{1}{x}$. Then

$$\int \frac{(x-1)e^x}{x^2}\,dx = (x-1)e^x\left(-\frac{1}{x}\right) - \int -e^x\,dx = -e^x + \frac{e^x}{x} + e^x + C = \frac{e^x}{x} + C.$$

55. Let $u = x - 1$, so that $du = dx$. Then

$$\int x^3(x-1)^{-4}\,dx = \int (u+1)^3u^{-4}\,du = \int (u^3+3u^2+3u+1)u^{-4}\,du = \int (u^{-1}+3u^{-2}+3u^{-3}+u^{-4})\,du$$

$$= \ln|u| - 3u^{-1} - \tfrac{3}{2}u^{-2} - \tfrac{1}{3}u^{-3} + C = \ln|x-1| - 3(x-1)^{-1} - \tfrac{3}{2}(x-1)^{-2} - \tfrac{1}{3}(x-1)^{-3} + C$$

56. Let $u = \sqrt{1-x^2}$, so $u^2 = 1 - x^2$, and $2u\,du = -2x\,dx$. Then $\int_0^1 x\sqrt{2-\sqrt{1-x^2}}\,dx = \int_1^0 \sqrt{2-u}\,(-u\,du)$.

Now let $v = \sqrt{2-u}$, so $v^2 = 2 - u$, and $2v\,dv = -du$. Thus,

$$\int_1^0 \sqrt{2-u}\,(-u\,du) = \int_1^{\sqrt{2}} v(2-v^2)\,(2v\,dv) = \int_1^{\sqrt{2}} (4v^2 - 2v^4)\,dv = \left[\tfrac{4}{3}v^3 - \tfrac{2}{5}v^5\right]_1^{\sqrt{2}}$$

$$= \left(\tfrac{8}{3}\sqrt{2} - \tfrac{8}{5}\sqrt{2}\right) - \left(\tfrac{4}{3} - \tfrac{2}{5}\right) = \tfrac{16}{15}\sqrt{2} - \tfrac{14}{15}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. Let $u=\sqrt{4x+1} \quad\Rightarrow\quad u^2=4x+1 \quad\Rightarrow\quad 2u\,du=4\,dx \quad\Rightarrow\quad dx=\frac{1}{2}u\,du$. So

$$\int \frac{1}{x\sqrt{4x+1}}\,dx=\int \frac{\frac{1}{2}u\,du}{\frac{1}{4}(u^2-1)\,u}=2\int \frac{du}{u^2-1}=2\left(\tfrac{1}{2}\right)\ln\left|\frac{u-1}{u+1}\right|+C \qquad \text{[by Formula 19]}$$

$$=\ln\left|\frac{\sqrt{4x+1}-1}{\sqrt{4x+1}+1}\right|+C$$

58. As in Exercise 57, let $u=\sqrt{4x+1}$. Then $\displaystyle\int \frac{dx}{x^2\sqrt{4x+1}}=\int \frac{\frac{1}{2}u\,du}{\left[\frac{1}{4}(u^2-1)\right]^2 u}=8\int \frac{du}{(u^2-1)^2}$. Now

$$\frac{1}{(u^2-1)^2}=\frac{1}{(u+1)^2(u-1)^2}=\frac{A}{u+1}+\frac{B}{(u+1)^2}+\frac{C}{u-1}+\frac{D}{(u-1)^2} \quad\Rightarrow$$

$1=A(u+1)(u-1)^2+B(u-1)^2+C(u-1)(u+1)^2+D(u+1)^2$. $u=1 \quad\Rightarrow\quad D=\frac{1}{4}$, $u=-1 \quad\Rightarrow\quad B=\frac{1}{4}$.

Equating coefficients of u^3 gives $A+C=0$, and equating coefficients of 1 gives $1=A+B-C+D \quad\Rightarrow$

$1=A+\frac{1}{4}-C+\frac{1}{4} \quad\Rightarrow\quad \frac{1}{2}=A-C$. So $A=\frac{1}{4}$ and $C=-\frac{1}{4}$. Therefore,

$$\int \frac{dx}{x^2\sqrt{4x+1}}=8\int\left[\frac{1/4}{u+1}+\frac{1/4}{(u+1)^2}+\frac{-1/4}{u-1}+\frac{1/4}{(u-1)^2}\right]du$$

$$=\int\left[\frac{2}{u+1}+2(u+1)^{-2}-\frac{2}{u-1}+2(u-1)^{-2}\right]du$$

$$=2\ln|u+1|-\frac{2}{u+1}-2\ln|u-1|-\frac{2}{u-1}+C$$

$$=2\ln\left(\sqrt{4x+1}+1\right)-\frac{2}{\sqrt{4x+1}+1}-2\ln\left|\sqrt{4x+1}-1\right|-\frac{2}{\sqrt{4x+1}-1}+C$$

59. Let $2x=\tan\theta \quad\Rightarrow\quad x=\frac{1}{2}\tan\theta$, $dx=\frac{1}{2}\sec^2\theta\,d\theta$, $\sqrt{4x^2+1}=\sec\theta$, so

$$\int \frac{dx}{x\sqrt{4x^2+1}}=\int \frac{\frac{1}{2}\sec^2\theta\,d\theta}{\frac{1}{2}\tan\theta\,\sec\theta}=\int \frac{\sec\theta}{\tan\theta}\,d\theta=\int \csc\theta\,d\theta$$

$$=-\ln|\csc\theta+\cot\theta|+C \qquad [\text{or } \ln|\csc\theta-\cot\theta|+C]$$

$$=-\ln\left|\frac{\sqrt{4x^2+1}}{2x}+\frac{1}{2x}\right|+C \qquad \left[\text{or } \ln\left|\frac{\sqrt{4x^2+1}}{2x}-\frac{1}{2x}\right|+C\right]$$

60. Let $u=x^2$. Then $du=2x\,dx \quad\Rightarrow$

$$\int \frac{dx}{x(x^4+1)}=\int \frac{x\,dx}{x^2(x^4+1)}=\frac{1}{2}\int \frac{du}{u(u^2+1)}=\frac{1}{2}\int\left[\frac{1}{u}-\frac{u}{u^2+1}\right]du=\tfrac{1}{2}\ln|u|-\tfrac{1}{4}\ln(u^2+1)+C$$

$$=\tfrac{1}{2}\ln(x^2)-\tfrac{1}{4}\ln(x^4+1)+C=\tfrac{1}{4}\left[\ln(x^4)-\ln(x^4+1)\right]+C=\frac{1}{4}\ln\left(\frac{x^4}{x^4+1}\right)+C$$

Or: Write $I=\displaystyle\int \frac{x^3\,dx}{x^4(x^4+1)}$ and let $u=x^4$.

61. $\displaystyle\int x^2\sinh mx\,dx=\frac{1}{m}x^2\cosh mx-\frac{2}{m}\int x\cosh mx\,dx \qquad \begin{bmatrix} u=x^2, & dv=\sinh mx\,dx, \\ du=2x\,dx & v=\frac{1}{m}\cosh mx \end{bmatrix}$

$$=\frac{1}{m}x^2\cosh\,mx-\frac{2}{m}\left(\frac{1}{m}x\sinh mx-\frac{1}{m}\int \sinh mx\,dx\right) \qquad \begin{bmatrix} U=x, & dV=\cosh mx\,dx, \\ dU=dx & V=\frac{1}{m}\sinh mx \end{bmatrix}$$

$$=\frac{1}{m}x^2\cosh mx-\frac{2}{m^2}x\sinh mx+\frac{2}{m^3}\cosh mx+C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

62. $\int (x+\sin x)^2\,dx = \int \left(x^2+2x\sin x+\sin^2 x\right)dx = \frac{1}{3}x^3+2(\sin x - x\cos x)+\frac{1}{2}(x-\sin x\cos x)+C$
$= \frac{1}{3}x^3+\frac{1}{2}x+2\sin x-\frac{1}{2}\sin x\cos x-2x\cos x+C$

63. Let $u=\sqrt{x}$, so that $x=u^2$ and $dx=2u\,du$. Then $\displaystyle\int \frac{dx}{x+x\sqrt{x}} = \int \frac{2u\,du}{u^2+u^2\cdot u} = \int \frac{2}{u(1+u)}\,du = I$.

Now $\dfrac{2}{u(1+u)} = \dfrac{A}{u}+\dfrac{B}{1+u} \quad\Rightarrow\quad 2=A(1+u)+Bu$. Set $u=-1$ to get $2=-B$, so $B=-2$. Set $u=0$ to get $2=A$.

Thus, $I=\displaystyle\int\left(\frac{2}{u}-\frac{2}{1+u}\right)du = 2\ln|u|-2\ln|1+u|+C = 2\ln\sqrt{x}-2\ln\left(1+\sqrt{x}\right)+C$.

64. Let $u=\sqrt{x}$, so that $x=u^2$ and $dx=2u\,du$. Then

$$\int \frac{dx}{\sqrt{x}+x\sqrt{x}} = \int \frac{2u\,du}{u+u^2\cdot u} = \int \frac{2}{1+u^2}\,du = 2\tan^{-1}u+C = 2\tan^{-1}\sqrt{x}+C.$$

65. Let $u=\sqrt[3]{x+c}$. Then $x=u^3-c \quad\Rightarrow$

$\int x\sqrt[3]{x+c}\,dx = \int (u^3-c)u\cdot 3u^2\,du = 3\int(u^6-cu^3)\,du = \frac{3}{7}u^7-\frac{3}{4}cu^4+C = \frac{3}{7}(x+c)^{7/3}-\frac{3}{4}c(x+c)^{4/3}+C$

66. Let $t=\sqrt{x^2-1}$. Then $dt=\left(x/\sqrt{x^2-1}\right)dx$, $x^2-1=t^2$, $x=\sqrt{t^2+1}$, so

$I=\displaystyle\int \frac{x\ln x}{\sqrt{x^2-1}}\,dx = \int \ln\sqrt{t^2+1}\,dt = \frac{1}{2}\int \ln(t^2+1)\,dt$. Now use parts with $u=\ln(t^2+1)$, $dv=dt$:

$$\begin{aligned} I &= \tfrac{1}{2}t\ln\left(t^2+1\right) - \int \frac{t^2}{t^2+1}\,dt = \tfrac{1}{2}t\ln\left(t^2+1\right) - \int\left[1-\frac{1}{t^2+1}\right]dt \\ &= \tfrac{1}{2}t\ln\left(t^2+1\right) - t + \tan^{-1}t + C = \sqrt{x^2-1}\,\ln x - \sqrt{x^2-1} + \tan^{-1}\sqrt{x^2-1} + C \end{aligned}$$

Another method: First integrate by parts with $u=\ln x$, $dv=\left(x/\sqrt{x^2-1}\right)dx$ and then use substitution $\left(x=\sec\theta \text{ or } u=\sqrt{x^2-1}\right)$.

67. $\dfrac{1}{x^4-16} = \dfrac{1}{(x^2-4)(x^2+4)} = \dfrac{1}{(x-2)(x+2)(x^2+4)} = \dfrac{A}{x-2}+\dfrac{B}{x+2}+\dfrac{Cx+D}{x^2+4}$. Multiply by $(x-2)(x+2)(x^2+4)$ to get $1=A(x+2)(x^2+4)+B(x-2)(x^2+4)+(Cx+D)(x-2)(x+2)$. Substituting 2 for x gives $1=32A \quad\Leftrightarrow\quad A=\frac{1}{32}$. Substituting -2 for x gives $1=-32B \quad\Leftrightarrow\quad B=-\frac{1}{32}$. Equating coefficients of x^3 gives $0=A+B+C=\frac{1}{32}-\frac{1}{32}+C$, so $C=0$. Equating constant terms gives $1=8A-8B-4D=\frac{1}{4}+\frac{1}{4}-4D$, so $\frac{1}{2}=-4D \quad\Leftrightarrow\quad D=-\frac{1}{8}$. Thus,

$$\begin{aligned} \int \frac{dx}{x^4-16} &= \int\left(\frac{1/32}{x-2}-\frac{1/32}{x+2}-\frac{1/8}{x^2+4}\right)dx = \frac{1}{32}\ln|x-2|-\frac{1}{32}\ln|x+2|-\frac{1}{8}\cdot\frac{1}{2}\tan^{-1}\left(\frac{x}{2}\right)+C \\ &= \frac{1}{32}\ln\left|\frac{x-2}{x+2}\right| - \frac{1}{16}\tan^{-1}\left(\frac{x}{2}\right)+C \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

68. Let $2x = \sec\theta$, so that $2\,dx = \sec\theta\,\tan\theta\,d\theta$. Then

$$\int \frac{dx}{x^2\sqrt{4x^2-1}} = \int \frac{\frac{1}{2}\sec\theta\,\tan\theta\,d\theta}{\frac{1}{4}\sec^2\theta\sqrt{\sec^2\theta-1}} = \int \frac{2\tan\theta\,d\theta}{\sec\theta\,\tan\theta}$$

$$= 2\int \cos\theta\,d\theta = 2\sin\theta + C$$

$$= 2\cdot\frac{\sqrt{4x^2-1}}{2x} + C = \frac{\sqrt{4x^2-1}}{x} + C$$

69. $$\int \frac{d\theta}{1+\cos\theta} = \int\left(\frac{1}{1+\cos\theta}\cdot\frac{1-\cos\theta}{1-\cos\theta}\right)d\theta = \int\frac{1-\cos\theta}{1-\cos^2\theta}\,d\theta = \int\frac{1-\cos\theta}{\sin^2\theta}\,d\theta = \int\left(\frac{1}{\sin^2\theta}-\frac{\cos\theta}{\sin^2\theta}\right)d\theta$$

$$= \int(\csc^2\theta - \cot\theta\,\csc\theta)\,d\theta = -\cot\theta + \csc\theta + C$$

Another method: Use the substitutions in Exercise 7.4.63.

$$\int\frac{d\theta}{1+\cos\theta} = \int\frac{2/(1+t^2)\,dt}{1+(1-t^2)/(1+t^2)} = \int\frac{2\,dt}{(1+t^2)+(1-t^2)} = \int dt = t + C = \tan\left(\frac{\theta}{2}\right) + C$$

70. $$\int\frac{d\theta}{1+\cos^2\theta} = \int\frac{(1/\cos^2\theta)\,d\theta}{(1+\cos^2\theta)/\cos^2\theta} = \int\frac{\sec^2\theta}{\sec^2\theta+1}\,d\theta = \int\frac{\sec^2\theta}{\tan^2\theta+2}\,d\theta = \int\frac{1}{u^2+2}\,du \quad \begin{bmatrix} u=\tan\theta, \\ du=\sec^2\theta\,d\theta \end{bmatrix}$$

$$= \int\frac{1}{u^2+\left(\sqrt{2}\right)^2}\,du = \frac{1}{\sqrt{2}}\tan^{-1}\left(\frac{u}{\sqrt{2}}\right) + C = \frac{1}{\sqrt{2}}\tan^{-1}\left(\frac{\tan\theta}{\sqrt{2}}\right) + C$$

71. Let $y = \sqrt{x}$ so that $dy = \dfrac{1}{2\sqrt{x}}\,dx \quad\Rightarrow\quad dx = 2\sqrt{x}\,dy = 2y\,dy$. Then

$$\int\sqrt{x}\,e^{\sqrt{x}}\,dx = \int ye^y(2y\,dy) = \int 2y^2e^y\,dy \qquad \begin{bmatrix} u=2y^2, & dv=e^y\,dy, \\ du=4y\,dy & v=e^y \end{bmatrix}$$

$$= 2y^2e^y - \int 4ye^y\,dy \qquad \begin{bmatrix} U=4y, & dV=e^y\,dy, \\ dU=4\,dy & V=e^y \end{bmatrix}$$

$$= 2y^2e^y - \left(4ye^y - \int 4e^y\,dy\right) = 2y^2e^y - 4ye^y + 4e^y + C$$

$$= 2(y^2-2y+2)e^y + C = 2\left(x - 2\sqrt{x} + 2\right)e^{\sqrt{x}} + C$$

72. Let $u = \sqrt{x}+1$, so that $x = (u-1)^2$ and $dx = 2(u-1)\,du$. Then

$$\int\frac{1}{\sqrt{\sqrt{x}+1}}\,dx = \int\frac{2(u-1)\,du}{\sqrt{u}} = \int(2u^{1/2}-2u^{-1/2})\,du = \frac{4}{3}u^{3/2} - 4u^{1/2} + C = \frac{4}{3}\left(\sqrt{x}+1\right)^{3/2} - 4\sqrt{\sqrt{x}+1} + C.$$

73. Let $u = \cos^2 x$, so that $du = 2\cos x\,(-\sin x)\,dx$. Then

$$\int\frac{\sin 2x}{1+\cos^4 x}\,dx = \int\frac{2\sin x\,\cos x}{1+(\cos^2 x)^2}\,dx = \int\frac{1}{1+u^2}\,(-du) = -\tan^{-1}u + C = -\tan^{-1}(\cos^2 x) + C.$$

74. Let $u = \tan x$. Then

$$\int_{\pi/4}^{\pi/3}\frac{\ln(\tan x)\,dx}{\sin x\,\cos x} = \int_{\pi/4}^{\pi/3}\frac{\ln(\tan x)}{\tan x}\,\sec^2 x\,dx = \int_1^{\sqrt{3}}\frac{\ln u}{u}\,du = \left[\tfrac{1}{2}(\ln u)^2\right]_1^{\sqrt{3}} = \tfrac{1}{2}\left(\ln\sqrt{3}\right)^2 = \tfrac{1}{8}(\ln 3)^2.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

75. $$\int \frac{1}{\sqrt{x+1}+\sqrt{x}}\,dx = \int \left(\frac{1}{\sqrt{x+1}+\sqrt{x}} \cdot \frac{\sqrt{x+1}-\sqrt{x\sqrt{x}}}{\sqrt{x+1}-\sqrt{x}}\right) dx = \int \left(\sqrt{x+1}-\sqrt{x}\right) dx$$
$$= \tfrac{2}{3}\left[(x+1)^{3/2} - x^{3/2}\right] + C$$

76. $$\int \frac{x^2}{x^6+3x^3+2}\,dx = \int \frac{x^2\,dx}{(x^3+1)(x^3+2)} = \int \frac{\frac{1}{3}\,du}{(u+1)(u+2)} \quad \left[\begin{matrix} u = x^3, \\ du = 3x^2\,dx \end{matrix}\right].$$

Now $\dfrac{1}{(u+1)(u+2)} = \dfrac{A}{u+1} + \dfrac{B}{u+2} \quad \Rightarrow \quad 1 = A(u+2) + B(u+1)$. Setting $u = -2$ gives $B = -1$. Setting $u = -1$ gives $A = 1$. Thus,

$$\frac{1}{3}\int \frac{du}{(u+1)(u+2)} = \frac{1}{3}\int \left(\frac{1}{u+1} - \frac{1}{u+2}\right) du = \frac{1}{3}\ln|u+1| - \frac{1}{3}\ln|u+2| + C$$
$$= \tfrac{1}{3}\ln\left|x^3+1\right| - \tfrac{1}{3}\ln\left|x^3+2\right| + C$$

77. Let $x = \tan\theta$, so that $dx = \sec^2\theta\,d\theta$, $x = \sqrt{3} \quad \Rightarrow \quad \theta = \frac{\pi}{3}$, and $x = 1 \quad \Rightarrow \quad \theta = \frac{\pi}{4}$. Then

$$\int_1^{\sqrt{3}} \frac{\sqrt{1+x^2}}{x^2}\,dx = \int_{\pi/4}^{\pi/3} \frac{\sec\theta}{\tan^2\theta}\,\sec^2\theta\,d\theta = \int_{\pi/4}^{\pi/3} \frac{\sec\theta\,(\tan^2\theta+1)}{\tan^2\theta}\,d\theta = \int_{\pi/4}^{\pi/3} \left(\frac{\sec\theta\,\tan^2\theta}{\tan^2\theta} + \frac{\sec\theta}{\tan^2\theta}\right) d\theta$$
$$= \int_{\pi/4}^{\pi/3} (\sec\theta + \csc\theta\,\cot\theta)\,d\theta = \Big[\ln|\sec\theta + \tan\theta| - \csc\theta\Big]_{\pi/4}^{\pi/3}$$
$$= \left(\ln\left|2+\sqrt{3}\right| - \tfrac{2}{\sqrt{3}}\right) - \left(\ln\left|\sqrt{2}+1\right| - \sqrt{2}\right) = \sqrt{2} - \tfrac{2}{\sqrt{3}} + \ln\left(2+\sqrt{3}\right) - \ln\left(1+\sqrt{2}\right)$$

78. Let $u = e^x$. Then $x = \ln u$, $dx = du/u \quad \Rightarrow$

$$\int \frac{1}{1+2e^x-e^{-x}}\,dx = \int \frac{du/u}{1+2u-1/u} = \int \frac{du}{2u^2+u-1} = \int \left[\frac{2/3}{2u-1} - \frac{1/3}{u+1}\right] du$$
$$= \tfrac{1}{3}\ln|2u-1| - \tfrac{1}{3}\ln|u+1| + C = \tfrac{1}{3}\ln|(2e^x-1)/(e^x+1)| + C$$

79. Let $u = e^x$. Then $x = \ln u$, $dx = du/u \quad \Rightarrow$

$$\int \frac{e^{2x}}{1+e^x}\,dx = \int \frac{u^2}{1+u}\frac{du}{u} = \int \frac{u}{1+u}\,du = \int \left(1 - \frac{1}{1+u}\right) du = u - \ln|1+u| + C = e^x - \ln(1+e^x) + C.$$

80. Use parts with $u = \ln(x+1)$, $dv = dx/x^2$:

$$\int \frac{\ln(x+1)}{x^2}\,dx = -\frac{1}{x}\ln(x+1) + \int \frac{dx}{x(x+1)} = -\frac{1}{x}\ln(x+1) + \int \left[\frac{1}{x} - \frac{1}{x+1}\right] dx$$
$$= -\frac{1}{x}\ln(x+1) + \ln|x| - \ln(x+1) + C = -\left(1+\frac{1}{x}\right)\ln(x+1) + \ln|x| + C$$

81. Let $\theta = \arcsin x$, so that $d\theta = \dfrac{1}{\sqrt{1-x^2}}\,dx$ and $x = \sin\theta$. Then

$$\int \frac{x+\arcsin x}{\sqrt{1-x^2}}\,dx = \int (\sin\theta + \theta)\,d\theta = -\cos\theta + \tfrac{1}{2}\theta^2 + C$$
$$= -\sqrt{1-x^2} + \tfrac{1}{2}(\arcsin x)^2 + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

82. $\displaystyle\int \frac{4^x+10^x}{2^x}\,dx = \int\left(\frac{4^x}{2^x}+\frac{10^x}{2^x}\right)dx = \int(2^x+5^x)\,dx = \frac{2^x}{\ln 2}+\frac{5^x}{\ln 5}+C$

83. $\displaystyle\int \frac{dx}{x\ln x - x} = \int \frac{dx}{x(\ln x-1)} = \int \frac{du}{u} \qquad \begin{bmatrix} u=\ln x-1, \\ du=(1/x)\,dx \end{bmatrix}$

$\displaystyle = \ln|u|+C = \ln|\ln x-1|+C$

84. $\displaystyle\int \frac{x^2}{\sqrt{x^2+1}}\,dx = \int \frac{\tan^2\theta}{\sec\theta}\sec^2\theta\,d\theta \qquad \begin{bmatrix} x=\tan\theta, \\ dx=\sec^2\theta\,d\theta \end{bmatrix}$

$\displaystyle = \int \tan^2\theta\,\sec\theta\,d\theta = \int(\sec^2\theta-1)\sec\theta\,d\theta$

$\displaystyle = \int(\sec^3\theta-\sec\theta)\,d\theta$

$\displaystyle = \tfrac{1}{2}(\sec\theta\,\tan\theta+\ln|\sec\theta+\tan\theta|)-\ln|\sec\theta+\tan\theta|+C$ [by (7.2.1) and Example 7.2.8]

$\displaystyle = \tfrac{1}{2}(\sec\theta\,\tan\theta-\ln|\sec\theta+\tan\theta|)+C = \tfrac{1}{2}\left[x\sqrt{x^2+1}-\ln(\sqrt{x^2+1}+x)\right]+C$

85. Let $y=\sqrt{1+e^x}$, so that $y^2=1+e^x$, $2y\,dy=e^x\,dx$, $e^x=y^2-1$, and $x=\ln(y^2-1)$. Then

$\displaystyle\int \frac{xe^x}{\sqrt{1+e^x}}\,dx = \int \frac{\ln(y^2-1)}{y}(2y\,dy) = 2\int[\ln(y+1)+\ln(y-1)]\,dy$

$\displaystyle = 2[(y+1)\ln(y+1)-(y+1)+(y-1)\ln(y-1)-(y-1)]+C$ [by Example 7.1.2]

$\displaystyle = 2[y\ln(y+1)+\ln(y+1)-y-1+y\ln(y-1)-\ln(y-1)-y+1]+C$

$\displaystyle = 2[y(\ln(y+1)+\ln(y-1))+\ln(y+1)-\ln(y-1)-2y]+C$

$\displaystyle = 2\left[y\ln(y^2-1)+\ln\frac{y+1}{y-1}-2y\right]+C = 2\left[\sqrt{1+e^x}\,\ln(e^x)+\ln\frac{\sqrt{1+e^x}+1}{\sqrt{1+e^x}-1}-2\sqrt{1+e^x}\right]+C$

$\displaystyle = 2x\sqrt{1+e^x}+2\ln\frac{\sqrt{1+e^x}+1}{\sqrt{1+e^x}-1}-4\sqrt{1+e^x}+C = 2(x-2)\sqrt{1+e^x}+2\ln\frac{\sqrt{1+e^x}+1}{\sqrt{1+e^x}-1}+C$

86. $\displaystyle\frac{1+\sin x}{1-\sin x} = \frac{1+\sin x}{1-\sin x}\cdot\frac{1+\sin x}{1+\sin x} = \frac{1+2\sin x+\sin^2 x}{1-\sin^2 x} = \frac{1+2\sin x+\sin^2 x}{\cos^2 x} = \frac{1}{\cos^2 x}+\frac{2\sin x}{\cos^2 x}+\frac{\sin^2 x}{\cos^2 x}$

$\displaystyle = \sec^2 x+2\sec x\,\tan x+\tan^2 x = \sec^2 x+2\sec x\,\tan x+\sec^2 x-1 = 2\sec^2 x+2\sec x\,\tan x-1$

Thus, $\displaystyle\int \frac{1+\sin x}{1-\sin x}\,dx = \int(2\sec^2 x+2\sec x\,\tan x-1)\,dx = 2\tan x+2\sec x-x+C$

87. Let $u=x$, $dv=\sin^2 x\cos x\,dx \;\Rightarrow\; du=dx$, $v=\frac{1}{3}\sin^3 x$. Then

$\displaystyle\int x\sin^2 x\,\cos x\,dx = \tfrac{1}{3}x\sin^3 x-\int\tfrac{1}{3}\sin^3 x\,dx = \tfrac{1}{3}x\sin^3 x-\tfrac{1}{3}\int(1-\cos^2 x)\sin x\,dx$

$\displaystyle = \frac{1}{3}x\sin^3 x+\frac{1}{3}\int(1-y^2)\,dy \qquad \begin{bmatrix} u=\cos x, \\ du=-\sin x\,dx \end{bmatrix}$

$\displaystyle = \tfrac{1}{3}x\sin^3 x+\tfrac{1}{3}y-\tfrac{1}{9}y^3+C = \tfrac{1}{3}x\sin^3 x+\tfrac{1}{3}\cos x-\tfrac{1}{9}\cos^3 x+C$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

88. $$\int \frac{\sec x\,\cos 2x}{\sin x+\sec x}\,dx = \int \frac{\sec x\,\cos 2x}{\sin x+\sec x}\cdot\frac{2\cos x}{2\cos x}\,dx = \int \frac{2\cos 2x}{2\sin x\cos x+2}\,dx$$

$$= \int \frac{2\cos 2x}{\sin 2x+2}\,dx = \int \frac{1}{u}\,du \qquad \begin{bmatrix} u=\sin 2x+2, \\ du = 2\cos 2x\,dx \end{bmatrix}$$

$$= \ln|u| + C = \ln|\sin 2x+2| + C = \ln(\sin 2x+2) + C$$

89. $$\int \sqrt{1-\sin x}\,dx = \int \sqrt{\frac{1-\sin x}{1}\cdot\frac{1+\sin x}{1+\sin x}}\,dx = \int \sqrt{\frac{1-\sin^2 x}{1+\sin x}}\,dx$$

$$= \int \sqrt{\frac{\cos^2 x}{1+\sin x}}\,dx = \int \frac{\cos x\,dx}{\sqrt{1+\sin x}} \qquad [\text{assume } \cos x > 0]$$

$$= \int \frac{du}{\sqrt{u}} \qquad \begin{bmatrix} u = 1+\sin x, \\ du = \cos x\,dx \end{bmatrix}$$

$$= 2\sqrt{u} + C = 2\sqrt{1+\sin x} + C$$

Another method: Let $u = \sin x$ so that $du = \cos x\,dx = \sqrt{1-\sin^2 x}\,dx = \sqrt{1-u^2}\,dx$. Then

$$\int \sqrt{1-\sin x}\,dx = \int \sqrt{1-u}\left(\frac{du}{\sqrt{1-u^2}}\right) = \int \frac{1}{\sqrt{1+u}}\,du = 2\sqrt{1+u} + C = 2\sqrt{1+\sin x} + C.$$

90. $$\int \frac{\sin x\,\cos x}{\sin^4 x+\cos^4 x}\,dx = \int \frac{\sin x\,\cos x}{(\sin^2 x)^2+(\cos^2 x)^2}\,dx = \int \frac{\sin x\,\cos x}{(\sin^2 x)^2+(1-\sin^2 x)^2}\,dx$$

$$= \int \frac{1}{u^2+(1-u)^2}\left(\frac{1}{2}\,du\right) \qquad \begin{bmatrix} u = \sin^2 x, \\ du = 2\sin x\,\cos x\,dx \end{bmatrix}$$

$$= \int \frac{1}{4u^2-4u+2}\,du = \int \frac{1}{(4u^2-4u+1)+1}\,du$$

$$= \int \frac{1}{(2u-1)^2+1}\,du = \frac{1}{2}\int \frac{1}{y^2+1}\,dy \qquad \begin{bmatrix} y = 2u-1, \\ dy = 2\,du \end{bmatrix}$$

$$= \tfrac{1}{2}\tan^{-1} y + C = \tfrac{1}{2}\tan^{-1}(2u-1) + C = \tfrac{1}{2}\tan^{-1}(2\sin^2 x - 1) + C$$

Another solution:

$$\int \frac{\sin x\,\cos x}{\sin^4 x+\cos^4 x}\,dx = \int \frac{(\sin x\,\cos x)/\cos^4 x}{(\sin^4 x+\cos^4 x)/\cos^4 x}\,dx = \int \frac{\tan x\,\sec^2 x}{\tan^4 x+1}\,dx$$

$$= \int \frac{1}{u^2+1}\left(\frac{1}{2}\,du\right) \qquad \begin{bmatrix} u = \tan^2 x, \\ du = 2\tan x\,\sec^2 x\,dx \end{bmatrix}$$

$$= \tfrac{1}{2}\tan^{-1} u + C = \tfrac{1}{2}\tan^{-1}(\tan^2 x) + C$$

91. $$\int_1^3 \left(\sqrt{\frac{9-x}{x}} - \sqrt{\frac{x}{9-x}}\right)dx = \int_1^3 \left(\frac{\sqrt{9-x}}{\sqrt{x}} - \frac{\sqrt{x}}{\sqrt{9-x}}\right)dx = \int_1^3 \left(\frac{9-x-x}{\sqrt{x}\sqrt{9-x}}\right)dx = \int_1^3 \left(\frac{9-2x}{\sqrt{9x-x^2}}\right)dx$$

$$= \int_8^{18} \left(\frac{1}{\sqrt{u}}\right)du \qquad \begin{bmatrix} u = 9x-x^2 \\ du = (9-2x)\,dx \end{bmatrix} \qquad = \int_8^{18} u^{-1/2}\,du = \left[2u^{1/2}\right]_8^{18}$$

$$= 2\sqrt{18} - 2\sqrt{8} = 6\sqrt{2} - 4\sqrt{2} = 2\sqrt{2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

92. $$\int \frac{1}{(\sin x+\cos x)^2}\,dx = \int \frac{1}{\sin^2 x + 2\sin x\cos x + \cos^2 x}\,dx = \int \frac{1}{\cos^2 x\left(\dfrac{\sin^2 x}{\cos^2 x} + \dfrac{2\sin x}{\cos x} + 1\right)}\,dx$$

$$= \int \frac{1}{\cos^2 x\,(\tan^2 x + 2\tan x + 1)}\,dx = \int \frac{1}{\cos^2 x\,(\tan x+1)^2}\,dx = \int \frac{\sec^2 x}{(\tan x+1)^2}\,dx$$

$$= \int \frac{1}{u^2}\,du \quad \begin{bmatrix} u = \tan x + 1 \\ du = \sec^2 x\,dx \end{bmatrix} \quad = -\frac{1}{u} + C = -\frac{1}{\tan x + 1} + C$$

93. $$\int_0^{\pi/6} \sqrt{1+\sin 2\theta}\,d\theta = \int_0^{\pi/6} \sqrt{\left(\sin^2\theta + \cos^2\theta\right) + 2\sin\theta\cos\theta}\,d\theta = \int_0^{\pi/6} \sqrt{(\sin\theta+\cos\theta)^2}\,d\theta$$

$$= \int_0^{\pi/6} |\sin\theta + \cos\theta|\,d\theta = \int_0^{\pi/6} (\sin\theta+\cos\theta)\,d\theta \qquad \begin{bmatrix} \text{since integrand is} \\ \text{positive on } [0,\,\pi/6] \end{bmatrix}$$

$$= \Big[-\cos\theta + \sin\theta\Big]_0^{\pi/6} = \left(-\frac{\sqrt{3}}{2} + \frac{1}{2}\right) - (-1+0) = \frac{3-\sqrt{3}}{2}$$

Alternate solution:

$$\int_0^{\pi/6} \sqrt{1+\sin 2\theta}\,d\theta = \int_0^{\pi/6} \sqrt{1+\sin 2\theta}\cdot\frac{\sqrt{1-\sin 2\theta}}{\sqrt{1-\sin 2\theta}}\,d\theta = \int_0^{\pi/6} \frac{\sqrt{1-\sin^2 2\theta}}{\sqrt{1-\sin 2\theta}}\,d\theta$$

$$= \int_0^{\pi/6} \frac{\sqrt{\cos^2 2\theta}}{\sqrt{1-\sin 2\theta}}\,d\theta = \int_0^{\pi/6} \frac{|\cos 2\theta|}{\sqrt{1-\sin 2\theta}}\,d\theta = \int_0^{\pi/6} \frac{\cos 2\theta}{\sqrt{1-\sin 2\theta}}\,d\theta$$

$$= -\frac{1}{2}\int_1^{1-\sqrt{3}/2} u^{-1/2}\,du \qquad \left[u = 1-\sin 2\theta,\, du = -2\cos 2\theta\,d\theta\right]$$

$$= -\tfrac{1}{2}\Big[2u^{1/2}\Big]_1^{1-\sqrt{3}/2} = 1 - \sqrt{1-\left(\sqrt{3}/2\right)}$$

94. (a) $$\int_1^2 \frac{e^x}{x}\,dx = \int_0^{\ln 2} \frac{e^{e^t}}{e^t}\,e^t\,dt \quad \begin{bmatrix} x = e^t, \\ dx = e^t dt \end{bmatrix} \quad = \int_0^{\ln 2} e^{e^t}\,dt = F(\ln 2)$$

(b) $$\int_2^3 \frac{1}{\ln x}\,dx = \int_{\ln 2}^{\ln 3} \frac{1}{u}\left(e^u\,du\right) \quad \begin{bmatrix} u = \ln x, \\ du = \dfrac{1}{x}\,dx \end{bmatrix} \quad = \int_{\ln\ln 2}^{\ln\ln 3} \frac{e^{e^v}}{e^v}\,e^v\,dv \quad \begin{bmatrix} u = e^v, \\ du = e^v\,dv \end{bmatrix}$$

$$= \int_{\ln\ln 2}^{0} e^{e^v}\,dv + \int_0^{\ln\ln 3} e^{e^v}\,dv \quad \left[\text{note that } \ln\ln 2 < 0\right]$$

$$= \int_0^{\ln\ln 3} e^{e^v}\,dv - \int_0^{\ln\ln 2} e^{e^v}\,dv = F(\ln\ln 3) - F(\ln\ln 2)$$

Another method: Substitute $x = e^{e^t}$ in the original integral.

95. The function $y = 2xe^{x^2}$ *does* have an elementary antiderivative, so we'll use this fact to help evaluate the integral.

$$\int (2x^2+1)e^{x^2}\,dx = \int 2x^2 e^{x^2}\,dx + \int e^{x^2}\,dx = \int x\left(2xe^{x^2}\right)dx + \int e^{x^2}\,dx$$

$$= xe^{x^2} - \int e^{x^2}\,dx + \int e^{x^2}\,dx \quad \begin{bmatrix} u = x, & dv = 2xe^{x^2}\,dx, \\ du = dx & v = e^{x^2} \end{bmatrix} \quad = xe^{x^2} + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7.6 Integration Using Tables and Technology

Keep in mind that there are several ways to approach many of these exercises, and different methods can lead to different forms of the answer.

1. $$\int_0^{\pi/2} \cos 5x \cos 2x\,dx \overset{80}{=} \left[\frac{\sin(5-2)x}{2(5-2)} + \frac{\sin(5+2)x}{2(5+2)}\right]_0^{\pi/2} \qquad \begin{bmatrix} a=5, \\ b=2 \end{bmatrix}$$
$$= \left[\frac{\sin 3x}{6} + \frac{\sin 7x}{14}\right]_0^{\pi/2} = \left(-\frac{1}{6} - \frac{1}{14}\right) - 0 = \frac{-7-3}{42} = -\frac{5}{21}$$

2. $$\int_0^1 \sqrt{x-x^2}\,dx = \int_0^1 \sqrt{2\left(\tfrac{1}{2}\right)x - x^2}\,dx \overset{113}{=} \left[\frac{x-\frac{1}{2}}{2}\sqrt{2\left(\tfrac{1}{2}\right)x-x^2} + \frac{\left(\frac{1}{2}\right)^2}{2}\cos^{-1}\left(\frac{\frac{1}{2}-x}{\frac{1}{2}}\right)\right]_0^1$$
$$= \left[\frac{2x-1}{4}\sqrt{x-x^2} + \frac{1}{8}\cos^{-1}(1-2x)\right]_0^1 = \left(0 + \frac{1}{8}\cdot\pi\right) - \left(0 + \frac{1}{8}\cdot 0\right) = \frac{1}{8}\pi$$

3. Let $u = x^2$, so that $du = 2x\,dx$. Thus,
$$\int x \arcsin(x^2)\,dx = \tfrac{1}{2}\int \sin^{-1} u\,du \overset{87}{=} \tfrac{1}{2}\left[u\sin^{-1}u + \sqrt{1-u^2}\right] + C = \tfrac{1}{2}x^2\sin^{-1}(x^2) + \tfrac{1}{2}\sqrt{1-x^4} + C.$$

4. $$\int \frac{\tan\theta}{\sqrt{2+\cos\theta}}\,d\theta = \int \frac{\sin\theta}{\cos\theta\sqrt{2+\cos\theta}}\,d\theta \overset{c}{=} \int \frac{1}{u\sqrt{2+u}}(-du)$$
$$\overset{57}{=} -\frac{1}{\sqrt{2}}\ln\left|\frac{\sqrt{2+u}-\sqrt{2}}{\sqrt{2+u}+\sqrt{2}}\right| + C = -\frac{1}{\sqrt{2}}\ln\left|\frac{\sqrt{2+\cos\theta}-\sqrt{2}}{\sqrt{2+\cos\theta}+\sqrt{2}}\right| + C$$

5. Let $u = y^2$, so that $du = 2y\,dy$. Then,
$$\int \frac{y^5}{\sqrt{4+y^4}}\,dy = \int \frac{y^4}{\sqrt{4+y^4}}\,y\,dy = \frac{1}{2}\int \frac{u^2}{\sqrt{4+u^2}}\,du \overset{26}{=} \frac{1}{2}\left[\frac{u}{2}\sqrt{4+u^2} - \frac{4}{2}\ln\left(u + \sqrt{4+u^2}\right)\right] + C$$
$$= \frac{y^2}{4}\sqrt{4+y^4} - \ln\left(y^2 + \sqrt{4+y^4}\right) + C$$

6. Let $u = t^3$, so that $du = 3t^2\,dt$. Thus,
$$\int \frac{\sqrt{t^6-5}}{t}\,dt = \int \frac{\sqrt{t^6-5}}{3t^3}\,3t^2\,dt = \frac{1}{3}\int \frac{\sqrt{u^2-5}}{u}\,du \overset{41}{=} \frac{1}{3}\left[\sqrt{u^2-5} - \sqrt{5}\cos^{-1}\left(\frac{\sqrt{5}}{|u|}\right)\right] + C$$
$$= \frac{1}{3}\sqrt{t^6-5} - \frac{\sqrt{5}}{3}\cos^{-1}\left(\frac{\sqrt{5}}{|t^3|}\right) + C$$

7. $$\int_0^{\pi/8} \arctan 2x\,dx = \tfrac{1}{2}\int_0^{\pi/4} \arctan u\,du \qquad \left[u = 2x,\ du = 2\,dx\right]$$
$$\overset{89}{=} \frac{1}{2}\left[u\arctan u - \frac{1}{2}\ln(1+u^2)\right]_0^{\pi/4} = \frac{1}{2}\left\{\left[\frac{\pi}{4}\arctan\frac{\pi}{4} - \frac{1}{2}\ln\left(1+\frac{\pi^2}{16}\right)\right] - 0\right\}$$
$$= \frac{\pi}{8}\arctan\frac{\pi}{4} - \frac{1}{4}\ln\left(1+\frac{\pi^2}{16}\right)$$

8. $$\int_0^2 x^2\sqrt{4-x^2}\,dx \overset{31}{=} \left[\frac{x}{8}(2x^2-4)\sqrt{4-x^2} + \frac{16}{8}\sin^{-1}\left(\frac{x}{2}\right)\right]_0^2 = \left(0 + 2\cdot\frac{\pi}{2}\right) - 0 = \pi$$

9. $$\int \frac{\cos x}{\sin^2 x - 9}\,dx = \int \frac{1}{u^2-9}\,du \quad \begin{bmatrix} u = \sin x, \\ du = \cos x\,dx \end{bmatrix} \quad \overset{20}{=} \frac{1}{2(3)}\ln\left|\frac{u-3}{u+3}\right| + C = \frac{1}{6}\ln\left|\frac{\sin x - 3}{\sin x + 3}\right| + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $\displaystyle\int \frac{e^x}{4-e^{2x}}\,dx = \int \frac{1}{4-u^2}\,du \quad \begin{bmatrix} u = e^x, \\ du = e^x\,dx \end{bmatrix} \quad \overset{19}{=} \frac{1}{2(2)} \ln\left|\frac{u+2}{u-2}\right| + C = \frac{1}{4}\ln\left|\frac{e^x+2}{e^x-2}\right| + C$

11. $\displaystyle\int \frac{\sqrt{9x^2+4}}{x^2}\,dx = \int \frac{\sqrt{u^2+4}}{u^2/9}\left(\frac{1}{3}du\right) \quad \begin{bmatrix} u = 3x, \\ du = 3\,dx \end{bmatrix}$

$$= 3\int \frac{\sqrt{4+u^2}}{u^2}\,du \overset{24}{=} 3\left[-\frac{\sqrt{4+u^2}}{u} + \ln(u+\sqrt{4+u^2})\right] + C$$

$$= -\frac{3\sqrt{4+9x^2}}{3x} + 3\ln(3x+\sqrt{4+9x^2}) + C = -\frac{\sqrt{9x^2+4}}{x} + 3\ln(3x+\sqrt{9x^2+4}) + C$$

12. Let $u = \sqrt{2}\,y$ and $a = \sqrt{3}$. Then $du = \sqrt{2}\,dy$ and

$$\int \frac{\sqrt{2y^2-3}}{y^2}\,dy = \int \frac{\sqrt{u^2-a^2}}{\frac{1}{2}u^2}\,\frac{du}{\sqrt{2}} = \sqrt{2}\int \frac{\sqrt{u^2-a^2}}{u^2}\,du$$

$$\overset{42}{=} \sqrt{2}\left(-\frac{\sqrt{u^2-a^2}}{u} + \ln\left|u+\sqrt{u^2-a^2}\right|\right) + C$$

$$= \sqrt{2}\left(-\frac{\sqrt{2y^2-3}}{\sqrt{2}\,y} + \ln\left|\sqrt{2}\,y+\sqrt{2y^2-3}\right|\right) + C$$

$$= -\frac{\sqrt{2y^2-3}}{y} + \sqrt{2}\ln\left|\sqrt{2}\,y+\sqrt{2y^2-3}\right| + C$$

13. $\int_0^\pi \cos^6\theta\,d\theta \overset{74}{=} \left[\frac{1}{6}\cos^5\theta\,\sin\theta\right]_0^\pi + \frac{5}{6}\int_0^\pi \cos^4\theta\,d\theta \overset{74}{=} 0 + \frac{5}{6}\left\{\left[\frac{1}{4}\cos^3\theta\,\sin\theta\right]_0^\pi + \frac{3}{4}\int_0^\pi \cos^2\theta\,d\theta\right\}$

$$\overset{64}{=} \frac{5}{6}\left\{0 + \frac{3}{4}\left[\frac{1}{2}\theta + \frac{1}{4}\sin 2\theta\right]_0^\pi\right\} = \frac{5}{6}\cdot\frac{3}{4}\cdot\frac{\pi}{2} = \frac{5\pi}{16}$$

14. $\int x\sqrt{2+x^4}\,dx = \int \sqrt{2+u^2}\,(\frac{1}{2}\,du) \quad \begin{bmatrix} u = x^2, \\ du = 2x\,dx \end{bmatrix}$

$$\overset{21}{=} \frac{1}{2}\left[\frac{u}{2}\sqrt{2+u^2} + \frac{2}{2}\ln\left(u+\sqrt{2+u^2}\right)\right] + C = \frac{x^2}{4}\sqrt{2+x^4} + \frac{1}{2}\ln(x^2+\sqrt{2+x^4}) + C$$

15. $\displaystyle\int \frac{\arctan\sqrt{x}}{\sqrt{x}}\,dx = \int \arctan u\,(2\,du) \quad \begin{bmatrix} u = \sqrt{x}, \\ du = 1/(2\sqrt{x})\,dx \end{bmatrix}$

$$\overset{89}{=} 2\left[u\arctan u - \tfrac{1}{2}\ln(1+u^2)\right] + C = 2\sqrt{x}\arctan\sqrt{x} - \ln(1+x) + C$$

16. $\int_0^\pi x^3\sin x\,dx \overset{84}{=} \left[-x^3\cos x\right]_0^\pi + 3\int_0^\pi x^2\cos x\,dx \overset{85}{=} -\pi^3(-1) + 3\left\{\left[x^2\sin x\right]_0^\pi - 2\int_0^\pi x\sin x\,dx\right\}$

$$= \pi^3 - 6\int_0^\pi x\sin x\,dx \overset{84}{=} \pi^3 - 6\left\{\left[-x\cos x\right]_0^\pi + \int_0^\pi \cos x\,dx\right\} = \pi^3 - 6[\pi] - 6\left[\sin x\right]_0^\pi$$

$$= \pi^3 - 6\pi$$

17. $\displaystyle\int \frac{\coth(1/y)}{y^2}\,dy = \int \coth u\,(-du) \quad \begin{bmatrix} u = 1/y, \\ du = -1/y^2\,dy \end{bmatrix}$

$$\overset{106}{=} -\ln|\sinh u| + C = -\ln|\sinh(1/y)| + C$$

18. $\displaystyle\int \frac{e^{3t}}{\sqrt{e^{2t}-1}}\,dt = \int \frac{e^{2t}}{\sqrt{e^{2t}-1}}\,(e^t\,dt) = \int \frac{u^2}{\sqrt{u^2-1}}\,du \quad \begin{bmatrix} u = e^t, \\ du = e^t\,dt \end{bmatrix}$

$$\overset{44}{=} \frac{u}{2}\sqrt{u^2-1} + \tfrac{1}{2}\ln\left|u+\sqrt{u^2-1}\right| + C = \tfrac{1}{2}e^t\sqrt{e^{2t}-1} + \tfrac{1}{2}\ln\left(e^t+\sqrt{e^{2t}-1}\right) + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. Let $z = 6 + 4y - 4y^2 = 6 - (4y^2 - 4y + 1) + 1 = 7 - (2y-1)^2$, $u = 2y - 1$, and $a = \sqrt{7}$.

Then $z = a^2 - u^2$, $du = 2\,dy$, and

$$\begin{aligned}
\int y\sqrt{6+4y-4y^2}\,dy &= \int y\sqrt{z}\,dy = \int \tfrac{1}{2}(u+1)\sqrt{a^2-u^2}\,\tfrac{1}{2}\,du = \tfrac{1}{4}\int u\sqrt{a^2-u^2}\,du + \tfrac{1}{4}\int \sqrt{a^2-u^2}\,du \\
&= \tfrac{1}{4}\int \sqrt{a^2-u^2}\,du - \tfrac{1}{8}\int(-2u)\sqrt{a^2-u^2}\,du \\
&\overset{30}{=} \frac{u}{8}\sqrt{a^2-u^2} + \frac{a^2}{8}\sin^{-1}\left(\frac{u}{a}\right) - \frac{1}{8}\int \sqrt{w}\,dw \qquad \left[\begin{array}{l} w = a^2 - u^2, \\ dw = -2u\,du \end{array}\right] \\
&= \frac{2y-1}{8}\sqrt{6+4y-4y^2} + \frac{7}{8}\sin^{-1}\frac{2y-1}{\sqrt{7}} - \frac{1}{8}\cdot\frac{2}{3}w^{3/2} + C \\
&= \frac{2y-1}{8}\sqrt{6+4y-4y^2} + \frac{7}{8}\sin^{-1}\frac{2y-1}{\sqrt{7}} - \frac{1}{12}(6+4y-4y^2)^{3/2} + C
\end{aligned}$$

This can be rewritten as

$$\begin{aligned}
&\sqrt{6+4y-4y^2}\left[\frac{1}{8}(2y-1) - \frac{1}{12}(6+4y-4y^2)\right] + \frac{7}{8}\sin^{-1}\frac{2y-1}{\sqrt{7}} + C \\
&\qquad = \left(\frac{1}{3}y^2 - \frac{1}{12}y - \frac{5}{8}\right)\sqrt{6+4y-4y^2} + \frac{7}{8}\sin^{-1}\left(\frac{2y-1}{\sqrt{7}}\right) + C \\
&\qquad = \frac{1}{24}(8y^2 - 2y - 15)\sqrt{6+4y-4y^2} + \frac{7}{8}\sin^{-1}\left(\frac{2y-1}{\sqrt{7}}\right) + C
\end{aligned}$$

20. $$\int \frac{dx}{2x^3 - 3x^2} = \int \frac{dx}{x^2(-3+2x)} \overset{50}{=} -\frac{1}{-3x} + \frac{2}{(-3)^2}\ln\left|\frac{-3+2x}{x}\right| + C = \frac{1}{3x} + \tfrac{2}{9}\ln\left|\frac{2x-3}{x}\right| + C$$

21. Let $u = \sin x$. Then $du = \cos x\,dx$, so

$$\begin{aligned}
\int \sin^2 x\,\cos x\,\ln(\sin x)\,dx &= \int u^2 \ln u\,du \overset{101}{=} \frac{u^{2+1}}{(2+1)^2}\left[(2+1)\ln u - 1\right] + C = \tfrac{1}{9}u^3(3\ln u - 1) + C \\
&= \tfrac{1}{9}\sin^3 x\,[3\ln(\sin x) - 1] + C
\end{aligned}$$

22. Let $u = \sin\theta$, so that $du = \cos\theta\,d\theta$. Then

$$\begin{aligned}
\int \frac{\sin 2\theta}{\sqrt{5-\sin\theta}}\,d\theta &= \int \frac{2\sin\theta\,\cos\theta}{\sqrt{5-\sin\theta}}\,d\theta = 2\int \frac{u}{\sqrt{5-u}}\,du \overset{55}{=} 2\cdot\frac{2}{3(-1)^2}\left[-1u - 2(5)\right]\sqrt{5-u} + C \\
&= \tfrac{4}{3}(-u-10)\sqrt{5-u} + C = -\tfrac{4}{3}(\sin\theta + 10)\sqrt{5-\sin\theta} + C
\end{aligned}$$

23. $$\begin{aligned}
\int \frac{\sin 2\theta}{\sqrt{\cos^4\theta + 4}}\,d\theta &= \int \frac{2\sin\theta\cos\theta}{\sqrt{\cos^4\theta+4}}\,d\theta = -\int \frac{1}{\sqrt{u^2+4}}\,du \qquad \left[\begin{array}{l} u = \cos^2\theta \\ du = -2\sin\theta\cos\theta\,d\theta \end{array}\right] \\
&\overset{25}{=} \ln\left(u + \sqrt{u^2+4}\right) + C = -\ln\left(\cos^2\theta + \sqrt{\cos^4\theta + 4}\right) + C
\end{aligned}$$

24. Let $u = x^2$ and $a = 2$. Then $du = 2x\,dx$ and

$$\begin{aligned}
\int_0^2 x^3\sqrt{4x^2 - x^4}\,dx &= \tfrac{1}{2}\int_0^2 x^2\sqrt{2\cdot 2\cdot x^2 - (x^2)^2}\cdot 2x\,dx = \tfrac{1}{2}\int_0^4 u\sqrt{2au - u^2}\,du \\
&\overset{114}{=} \left[\frac{2u^2 - au - 3a^2}{12}\sqrt{2au - u^2} + \frac{a^3}{4}\cos^{-1}\left(\frac{a-u}{a}\right)\right]_0^4 \\
&= \left[\frac{2u^2 - 2u - 12}{12}\sqrt{4u - u^2} + \frac{8}{4}\cos^{-1}\left(\frac{2-u}{2}\right)\right]_0^4 \\
&= \left[\frac{u^2 - u - 6}{6}\sqrt{4u - u^2} + 2\cos^{-1}\left(\frac{2-u}{2}\right)\right]_0^4 \\
&= [0 + 2\cos^{-1}(-1)] - (0 + 2\cos^{-1} 1) = 2\cdot\pi - 2\cdot 0 = 2\pi
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $\int x^3 e^{2x}\,dx \overset{97}{=} \frac{1}{2}x^3e^{2x} - \frac{3}{2}\int x^2e^{2x}\,dx \overset{97}{=} \frac{1}{2}x^3e^{2x} - \frac{3}{2}\left(\frac{1}{2}x^2e^{2x} - \int xe^{2x}\,dx\right)$

$$\overset{96}{=} \tfrac{1}{2}x^3e^{2x} - \tfrac{3}{2}\left(\tfrac{1}{2}x^2e^{2x} - \tfrac{1}{4}(2x-1)e^{2x}\right) + C$$
$$= \tfrac{1}{2}x^3e^{2x} - \tfrac{3}{4}x^2e^{2x} + \tfrac{3}{4}xe^{2x} - \tfrac{3}{8}e^{2x} + C$$
$$= \tfrac{1}{2}e^{2x}\left(x^3 - \tfrac{3}{2}x^2 + \tfrac{3}{2}x - \tfrac{3}{4}\right) + C$$

26. $\int x^3 \arcsin(x^2)\,dx = \int u \arcsin u \left(\frac{1}{2}du\right) \qquad \begin{bmatrix} u = x^2, \\ du = 2x\,dx \end{bmatrix}$

$$\overset{90}{=} \frac{1}{2}\left[\frac{2u^2-1}{4}\arcsin u + \frac{u\sqrt{1-u^2}}{4}\right] + C = \frac{2x^4-1}{8}\arcsin(x^2) + \frac{x^2\sqrt{1-x^4}}{8} + C$$

27. $\int \cos^5 y\,dy \overset{74}{=} \frac{1}{5}\cos^4 y \sin y + \frac{4}{5}\int \cos^3 y\,dy \overset{68}{=} \frac{1}{5}\cos^4 y \sin y + \frac{4}{5}\left[\frac{1}{3}(2+\cos^2 y)\sin y\right] + C$

$$= \tfrac{1}{5}\cos^4 y \sin y + \tfrac{8}{15}\sin y + \tfrac{4}{15}\cos^2 y \sin y + C = \tfrac{1}{5}\sin y\left(\cos^4 y + \tfrac{4}{3}\cos^2 y + \tfrac{8}{3}\right) + C$$

28. Let $u = \ln x$, so that $du = (1/x)\,dx$. Thus,

$$\int \frac{\sqrt{(\ln x)^2 - 9}}{x \ln x}\,dx = \int \frac{\sqrt{u^2-9}}{u}\,du \overset{41}{=} \sqrt{u^2-9} - 3\cos^{-1}\left(\frac{3}{|u|}\right) + C$$
$$= \sqrt{(\ln x)^2 - 9} - 3\cos^{-1}\left(\frac{3}{|\ln x|}\right) + C$$

29. $\displaystyle\int \frac{\cos^{-1}(x^{-2})}{x^3}\,dx = -\frac{1}{2}\int \cos^{-1} u\,du \qquad \begin{bmatrix} u = x^{-2}, \\ du = -2x^{-3}\,dx \end{bmatrix}$

$$\overset{88}{=} -\tfrac{1}{2}\left(u\cos^{-1} u - \sqrt{1-u^2}\right) + C = -\tfrac{1}{2}x^{-2}\cos^{-1}(x^{-2}) + \tfrac{1}{2}\sqrt{1-x^{-4}} + C$$

30. $\displaystyle\int \frac{dx}{\sqrt{1-e^{2x}}} = \int \frac{1}{\sqrt{1-u^2}}\left(\frac{du}{u}\right) \qquad \begin{bmatrix} u = e^x, \\ du = e^x\,dx,\ dx = du/u \end{bmatrix}$

$$\overset{35}{=} -\frac{1}{1}\ln\left|\frac{1+\sqrt{1-u^2}}{u}\right| + C = -\ln\left|\frac{1+\sqrt{1-e^{2x}}}{e^x}\right| + C = -\ln\left(\frac{1+\sqrt{1-e^{2x}}}{e^x}\right) + C$$

31. Let $u = e^x$. Then $x = \ln u$, $dx = du/u$, so

$$\int \sqrt{e^{2x}-1}\,dx = \int \frac{\sqrt{u^2-1}}{u}\,du \overset{41}{=} \sqrt{u^2-1} - \cos^{-1}(1/u) + C = \sqrt{e^{2x}-1} - \cos^{-1}(e^{-x}) + C.$$

32. $\displaystyle\int \sin 2\theta \arctan(\sin\theta)\,d\theta = \int 2\sin\theta\cos\theta\tan^{-1}(\sin\theta)\,d\theta \overset{s}{=} 2\int u\tan^{-1} u\,du$

$$\overset{92}{=} 2\left(\frac{u^2+1}{2}\tan^{-1} u - \frac{u}{2}\right) + C = (\sin^2\theta + 1)\tan^{-1}(\sin\theta) - \sin\theta + C$$

33. $\displaystyle\int \frac{x^4}{\sqrt{x^{10}-2}}\,dx = \int \frac{x^4}{\sqrt{(x^5)^2-2}}\,dx = \frac{1}{5}\int \frac{1}{\sqrt{u^2-2}}\,du \qquad \begin{bmatrix} u = x^5, \\ du = 5x^4\,dx \end{bmatrix}$

$$\overset{43}{=} \tfrac{1}{5}\ln\left|u + \sqrt{u^2-2}\right| + C = \tfrac{1}{5}\ln\left|x^5 + \sqrt{x^{10}-2}\right| + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. Let $u = \tan\theta$ and $a = 3$. Then $du = \sec^2\theta\, d\theta$ and

$$\int \frac{\sec^2\theta\,\tan^2\theta}{\sqrt{9-\tan^2\theta}}\,d\theta = \int \frac{u^2}{\sqrt{a^2-u^2}}\,du \overset{34}{=} -\frac{u}{2}\sqrt{a^2-u^2} + \frac{a^2}{2}\sin^{-1}\left(\frac{u}{a}\right) + C$$
$$= -\frac{1}{2}\tan\theta\sqrt{9-\tan^2\theta} + \frac{9}{2}\sin^{-1}\left(\frac{\tan\theta}{3}\right) + C$$

35. Use disks about the x-axis:

$$V = \int_0^\pi \pi(\sin^2 x)^2\,dx = \pi\int_0^\pi \sin^4 x\,dx \overset{73}{=} \pi\left\{\left[-\tfrac{1}{4}\sin^3 x\,\cos x\right]_0^\pi + \tfrac{3}{4}\int_0^\pi \sin^2 x\,dx\right\}$$
$$\overset{63}{=} \pi\left\{0 + \tfrac{3}{4}\left[\tfrac{1}{2}x - \tfrac{1}{4}\sin 2x\right]_0^\pi\right\} = \pi\left[\tfrac{3}{4}\left(\tfrac{1}{2}\pi - 0\right)\right] = \tfrac{3}{8}\pi^2$$

36. Use shells about the y-axis:

$$V = \int_0^1 2\pi x\arcsin x\,dx \overset{90}{=} 2\pi\left[\frac{2x^2-1}{4}\sin^{-1}x + \frac{x\sqrt{1-x^2}}{4}\right]_0^1 = 2\pi\left[\left(\frac{1}{4}\cdot\frac{\pi}{2}+0\right)-0\right] = \frac{1}{4}\pi^2$$

37. (a) $$\frac{d}{du}\left[\frac{1}{b^3}\left(a+bu-\frac{a^2}{a+bu}-2a\ln|a+bu|\right)+C\right] = \frac{1}{b^3}\left[b+\frac{ba^2}{(a+bu)^2}-\frac{2ab}{(a+bu)}\right]$$
$$= \frac{1}{b^3}\left[\frac{b(a+bu)^2+ba^2-(a+bu)2ab}{(a+bu)^2}\right]$$
$$= \frac{1}{b^3}\left[\frac{b^3u^2}{(a+bu)^2}\right] = \frac{u^2}{(a+bu)^2}$$

(b) Let $t = a + bu \quad\Rightarrow\quad dt = b\,du$. Note that $u = \dfrac{t-a}{b}$ and $du = \dfrac{1}{b}\,dt$.

$$\int \frac{u^2\,du}{(a+bu)^2} = \frac{1}{b^3}\int\frac{(t-a)^2}{t^2}\,dt = \frac{1}{b^3}\int\frac{t^2-2at+a^2}{t^2}\,dt = \frac{1}{b^3}\int\left(1-\frac{2a}{t}+\frac{a^2}{t^2}\right)dt$$
$$= \frac{1}{b^3}\left(t-2a\ln|t|-\frac{a^2}{t}\right)+C = \frac{1}{b^3}\left(a+bu-\frac{a^2}{a+bu}-2a\ln|a+bu|\right)+C$$

38. (a) $$\frac{d}{du}\left[\frac{u}{8}(2u^2-a^2)\sqrt{a^2-u^2}+\frac{a^4}{8}\sin^{-1}\frac{u}{a}+C\right]$$
$$= \frac{u}{8}(2u^2-a^2)\frac{-u}{\sqrt{a^2-u^2}} + \sqrt{a^2-u^2}\left[\frac{u}{8}(4u)+(2u^2-a^2)\tfrac{1}{8}\right] + \frac{a^4}{8}\frac{1/a}{\sqrt{1-u^2/a^2}}$$
$$= -\frac{u^2(2u^2-a^2)}{8\sqrt{a^2-u^2}} + \sqrt{a^2-u^2}\left[\frac{u^2}{2}+\frac{2u^2-a^2}{8}\right] + \frac{a^4}{8\sqrt{a^2-u^2}}$$
$$= \tfrac{1}{2}(a^2-u^2)^{-1/2}\left[-\frac{u^2}{4}(2u^2-a^2)+u^2(a^2-u^2)+\tfrac{1}{4}(a^2-u^2)(2u^2-a^2)+\frac{a^4}{4}\right]$$
$$= \tfrac{1}{2}(a^2-u^2)^{-1/2}[2u^2a^2-2u^4]$$
$$= \frac{u^2(a^2-u^2)}{\sqrt{a^2-u^2}} = u^2\sqrt{a^2-u^2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Let $u = a\sin\theta \quad\Rightarrow\quad du = a\cos\theta\,d\theta$. Then

$$\begin{aligned}
\int u^2\sqrt{a^2-u^2}\,du &= \int a^2\sin^2\theta\, a\sqrt{1-\sin^2\theta}\, a\cos\theta\,d\theta = a^4\int \sin^2\theta\,\cos^2\theta\,d\theta \\
&= a^4\int \tfrac{1}{2}(1+\cos 2\theta)\tfrac{1}{2}(1-\cos 2\theta)\,d\theta = \tfrac{1}{4}a^4\int(1-\cos^2 2\theta)\,d\theta \\
&= \tfrac{1}{4}a^4\int\left[1-\tfrac{1}{2}(1+\cos 4\theta)\right]d\theta = \tfrac{1}{4}a^4\left(\tfrac{1}{2}\theta-\tfrac{1}{8}\sin 4\theta\right)+C \\
&= \tfrac{1}{4}a^4\left(\tfrac{1}{2}\theta-\tfrac{1}{8}\cdot 2\sin 2\theta\,\cos 2\theta\right)+C = \tfrac{1}{4}a^4\left[\tfrac{1}{2}\theta-\tfrac{1}{2}\sin\theta\,\cos\theta(1-2\sin^2\theta)\right]+C \\
&= \frac{a^4}{8}\left[\sin^{-1}\frac{u}{a}-\frac{u}{a}\frac{\sqrt{a^2-u^2}}{a}\left(1-\frac{2u^2}{a^2}\right)\right]+C = \frac{a^4}{8}\left[\sin^{-1}\frac{u}{a}-\frac{u}{a}\frac{\sqrt{a^2-u^2}}{a}\frac{a^2-2u^2}{a^2}\right]+C \\
&= \frac{u}{8}(2u^2-a^2)\sqrt{a^2-u^2}+\frac{a^4}{8}\sin^{-1}\frac{u}{a}+C
\end{aligned}$$

39. Maple and Mathematica both give $\int \sec^4 x\,dx = \frac{2}{3}\tan x + \frac{1}{3}\tan x\,\sec^2 x$. Using Formula 77, we get $\int \sec^4 x\,dx = \frac{1}{3}\tan x\,\sec^2 x + \frac{2}{3}\int \sec^2 x\,dx = \frac{1}{3}\tan x\,\sec^2 x + \frac{2}{3}\tan x + C$.

40. Maple gives $I = \displaystyle\int \csc^5 x\,dx = -\frac{1}{4}\frac{\cos x}{\sin^4 x} - \frac{3}{8}\frac{\cos x}{\sin^2 x} + \frac{3}{8}\ln(\csc x - \cot x)$. Mathematica gives

$$\begin{aligned}
I &= -\frac{3}{32}\csc^2\frac{x}{2} - \frac{1}{64}\csc^4\frac{x}{2} - \frac{3}{8}\log\cos\frac{x}{2} + \frac{3}{8}\log\sin\frac{x}{2} + \frac{3}{32}\sec^2\frac{x}{2} + \frac{1}{64}\sec^4\frac{x}{2} \\
&= \frac{3}{8}\left(\log\sin\frac{x}{2} - \log\cos\frac{x}{2}\right) + \frac{3}{32}\left(\sec^2\frac{x}{2} - \csc^2\frac{x}{2}\right) + \frac{1}{64}\left(\sec^4\frac{x}{2} - \csc^4\frac{x}{2}\right) \\
&= \frac{3}{8}\log\frac{\sin(x/2)}{\cos(x/2)} + \frac{3}{32}\left[\frac{1}{\cos^2(x/2)} - \frac{1}{\sin^2(x/2)}\right] + \frac{1}{64}\left[\frac{1}{\cos^4(x/2)} - \frac{1}{\sin^4(x/2)}\right] \\
&= \frac{3}{8}\log\tan\frac{x}{2} + \frac{3}{32}\left[\frac{\sin^2(x/2)-\cos^2(x/2)}{\cos^2(x/2)\,\sin^2(x/2)}\right] + \frac{1}{64}\left[\frac{\sin^4(x/2)-\cos^4(x/2)}{\cos^4(x/2)\,\sin^4(x/2)}\right]
\end{aligned}$$

Now
$$\frac{\sin^2(x/2)-\cos^2(x/2)}{\cos^2(x/2)\,\sin^2(x/2)} = \frac{\dfrac{1-\cos x}{2} - \dfrac{1+\cos x}{2}}{\dfrac{1+\cos x}{2}\cdot\dfrac{1-\cos x}{2}} = \frac{-\dfrac{2\cos x}{2}}{\dfrac{1-\cos^2 x}{4}} = \frac{-4\cos x}{\sin^2 x}$$

and
$$\begin{aligned}
\frac{\sin^4(x/2)-\cos^4(x/2)}{\cos^4(x/2)\,\sin^4(x/2)} &= \frac{\sin^2(x/2)-\cos^2(x/2)}{\cos^2(x/2)\,\sin^2(x/2)}\,\frac{\sin^2(x/2)+\cos^2(x/2)}{\cos^2(x/2)\,\sin^2(x/2)} \\
&= \frac{-4\cos x}{\sin^2 x}\,\frac{1}{\dfrac{1+\cos x}{2}\cdot\dfrac{1-\cos x}{2}} = -\frac{4\cos x}{\sin^2 x}\,\frac{4}{1-\cos^2 x} = -\frac{16\cos x}{\sin^4 x}
\end{aligned}$$

Returning to the expression for I, we get

$$I = \frac{3}{8}\log\tan\frac{x}{2} + \frac{3}{32}\left(\frac{-4\cos x}{\sin^2 x}\right) + \frac{1}{64}\left(\frac{-16\cos x}{\sin^4 x}\right) = \frac{3}{8}\log\tan\frac{x}{2} - \frac{3}{8}\frac{\cos x}{\sin^2 x} - \frac{1}{4}\frac{\cos x}{\sin^4 x},$$

so all are equivalent.

Now use Formula 78 to get

$$\begin{aligned}
\int \csc^5 x\,dx &= \frac{-1}{4}\cot x\,\csc^3 x + \frac{3}{4}\int \csc^3 x\,dx = -\frac{1}{4}\frac{\cos x}{\sin x}\frac{1}{\sin^3 x} + \frac{3}{4}\left(\frac{-1}{2}\cot x\csc x + \frac{1}{2}\int \csc x\,dx\right) \\
&= -\frac{1}{4}\frac{\cos x}{\sin^4 x} - \frac{3}{8}\frac{\cos x}{\sin x}\frac{1}{\sin x} + \frac{3}{8}\int \csc x\,dx = -\frac{1}{4}\frac{\cos x}{\sin^4 x} - \frac{3}{8}\frac{\cos x}{\sin^2 x} + \frac{3}{8}\ln|\csc x - \cot x| + C
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. Maple gives $\int x^2\sqrt{2^2+x^2}\,dx = \frac{1}{4}x(x^2+4)^{3/2} - \frac{1}{2}x\sqrt{x^2+4} - 2\operatorname{arcsinh}\left(\frac{1}{2}x\right)$. Applying the command `convert(%,ln);` yields

$$\begin{aligned}\tfrac{1}{4}x(x^2+4)^{3/2} - \tfrac{1}{2}x\sqrt{x^2+4} - 2\ln\left(\tfrac{1}{2}x + \tfrac{1}{2}\sqrt{x^2+4}\right) &= \tfrac{1}{4}x(x^2+4)^{1/2}\left[(x^2+4)-2\right] - 2\ln\left[\left(x+\sqrt{x^2+4}\right)/2\right]\\ &= \tfrac{1}{4}x(x^2+2)\sqrt{x^2+4} - 2\ln\left(\sqrt{x^2+4}+x\right) + 2\ln 2\end{aligned}$$

Mathematica gives $\frac{1}{4}x(2+x^2)\sqrt{3+x^2} - 2\operatorname{arcsinh}(x/2)$. Applying the `TrigToExp` and `Simplify` commands gives $\frac{1}{4}\left[x(2+x^2)\sqrt{4+x^2} - 8\log\left(\frac{1}{2}\left(x+\sqrt{4+x^2}\right)\right)\right] = \frac{1}{4}x(x^2+2)\sqrt{x^2+4} - 2\ln\left(x+\sqrt{4+x^2}\right) + 2\ln 2$, so all are equivalent (without constant).

Now use Formula 22 to get

$$\begin{aligned}\int x^2\sqrt{2^2+x^2}\,dx &= \frac{x}{8}(2^2+2x^2)\sqrt{2^2+x^2} - \frac{2^4}{8}\ln\left(x+\sqrt{2^2+x^2}\right) + C\\ &= \frac{x}{8}(2)(2+x^2)\sqrt{4+x^2} - 2\ln\left(x+\sqrt{4+x^2}\right) + C\\ &= \tfrac{1}{4}x(x^2+2)\sqrt{x^2+4} - 2\ln\left(\sqrt{x^2+4}+x\right) + C\end{aligned}$$

42. Maple gives $\int \frac{1}{e^x(3e^x+2)}\,dx = \frac{3}{4}\ln(3e^x+2) - \frac{1}{2e^x} - \frac{3}{4}\ln(e^x)$, whereas Mathematica gives

$$-\frac{e^{-x}}{2} + \frac{3}{4}\log(3+2e^{-x}) = -\frac{e^{-x}}{2} + \frac{3}{4}\log\left(\frac{3e^x+2}{e^x}\right) = -\frac{e^{-x}}{2} + \frac{3}{4}\frac{\ln(3e^x+2)}{\ln e^x} = -\frac{e^{-x}}{2} + \frac{3}{4}\ln(3e^x+2) - \frac{3}{4}x,$$

so both are equivalent. Now let $u = e^x$, so $du = e^x\,dx$ and $dx = du/u$. Then

$$\begin{aligned}\int \frac{1}{e^x(3e^x+2)}\,dx &= \int \frac{1}{u(3u+2)}\frac{du}{u} = \int \frac{1}{u^2(2+3u)}\,du \overset{50}{=} -\frac{1}{2u} + \frac{3}{2^2}\ln\left|\frac{2+3u}{u}\right| + C\\ &= -\frac{1}{2e^x} + \frac{3}{4}\ln(2+3e^x) - \frac{3}{4}\ln e^x + C = -\frac{1}{2e^x} + \frac{3}{4}\ln(3e^x+2) - \frac{3}{4}x + C\end{aligned}$$

43. Maple gives $\int \cos^4 x\,dx = \frac{\sin x\,\cos^3 x}{4} + \frac{3\sin x\,\cos x}{8} + \frac{3x}{8}$, whereas Mathematica gives

$$\begin{aligned}\frac{3x}{8} + \frac{1}{4}\sin(2x) + \frac{1}{32}\sin(4x) &= \frac{3x}{8} + \frac{1}{4}(2\sin x\,\cos x) + \frac{1}{32}(2\sin 2x\,\cos 2x)\\ &= \frac{3x}{8} + \frac{1}{2}\sin x\,\cos x + \frac{1}{16}[2\sin x\,\cos x\,(2\cos^2 x - 1)]\\ &= \frac{3x}{8} + \frac{1}{2}\sin x\,\cos x + \frac{1}{4}\sin x\,\cos^3 x - \frac{1}{8}\sin x\,\cos x,\end{aligned}$$

so both are equivalent.

Using tables,

$$\begin{aligned}\int \cos^4 x\,dx &\overset{74}{=} \tfrac{1}{4}\cos^3 x\,\sin x + \tfrac{3}{4}\int \cos^2 x\,dx \overset{64}{=} \tfrac{1}{4}\cos^3 x\,\sin x + \tfrac{3}{4}\left(\tfrac{1}{2}x + \tfrac{1}{4}\sin 2x\right) + C\\ &= \tfrac{1}{4}\cos^3 x\,\sin x + \tfrac{3}{8}x + \tfrac{3}{16}(2\sin x\,\cos x) + C = \tfrac{1}{4}\cos^3 x\,\sin x + \tfrac{3}{8}x + \tfrac{3}{8}\sin x\,\cos x + C\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. Maple gives

$$\int x^2\sqrt{1-x^2}\,dx = -\frac{x}{4}(1-x^2)^{3/2} + \frac{x}{8}\sqrt{1-x^2} + \frac{1}{8}\arcsin x = \frac{x}{8}(1-x^2)^{1/2}[-2(1-x^2)+1] + \frac{1}{8}\arcsin x$$
$$= \frac{x}{8}(1-x^2)^{1/2}(2x^2-1) + \frac{1}{8}\arcsin x,$$

and Mathematica gives $\frac{1}{8}\left(x\sqrt{1-x^2}(-1+2x^2) + \arcsin x\right)$, so both are equivalent.

Now use Formula 31 to get

$$\int x^2\sqrt{1-x^2}\,dx = \frac{x}{8}(2x^2-1)\sqrt{1-x^2} + \frac{1}{8}\sin^{-1}x + C$$

45. Maple gives $\int \tan^5 x\,dx = \frac{1}{4}\tan^4 x - \frac{1}{2}\tan^2 x + \frac{1}{2}\ln(1+\tan^2 x)$, and Mathematica gives $\int \tan^5 x\,dx = \frac{1}{4}[-1-2\cos(2x)]\sec^4 x - \ln(\cos x)$. These expressions are equivalent, and neither includes absolute value bars or a constant of integration. Note that Mathematica's expression suggests that the integral is undefined where $\cos x < 0$, which is not the case. Using Formula 75, $\int \tan^5 x\,dx = \frac{1}{5-1}\tan^{5-1}x - \int \tan^{5-2}x\,dx = \frac{1}{4}\tan^4 x - \int \tan^3 x\,dx$. Using Formula 69, $\int \tan^3 x\,dx = \frac{1}{2}\tan^2 x + \ln|\cos x| + C$, so $\int \tan^5 x\,dx = \frac{1}{4}\tan^4 x - \frac{1}{2}\tan^2 x - \ln|\cos x| + C$.

46. Maple and Mathematica both give $\displaystyle\int \frac{1}{\sqrt{1+\sqrt[3]{x}}}\,dx = \frac{2}{5}\sqrt{\sqrt[3]{x}+1}\left(3\sqrt[3]{x^2} - 4\sqrt[3]{x} + 8\right)$. [Maple adds a constant of $-\frac{16}{5}$.] We'll change the form of the integral by letting $u = \sqrt[3]{x}$, so that $u^3 = x$ and $3u^2\,du = dx$. Then

$$\int \frac{1}{\sqrt{1+\sqrt[3]{x}}}\,dx = \int \frac{3u^2\,du}{\sqrt{1+u}} \overset{56}{=} 3\left[\frac{2}{15(1)^3}\left(8(1)^2 + 3(1)^2u^2 - 4(1)(1)u\right)\sqrt{1+u}\right] + C$$
$$= \tfrac{2}{5}(8+3u^2-4u)\sqrt{1+u} + C = \tfrac{2}{5}\left(8 + 3\sqrt[3]{x^2} - 4\sqrt[3]{x}\right)\sqrt{1+\sqrt[3]{x}} + C$$

47. (a) $F(x) = \displaystyle\int f(x)\,dx = \int \frac{1}{x\sqrt{1-x^2}}\,dx \overset{35}{=} -\frac{1}{1}\ln\left|\frac{1+\sqrt{1-x^2}}{x}\right| + C = -\ln\left|\frac{1+\sqrt{1-x^2}}{x}\right| + C.$

f has domain $\{x \mid x \neq 0, 1-x^2 > 0\} = \{x \mid x \neq 0,\ |x| < 1\} = (-1,0)\cup(0,1)$. F has the same domain.

(b) Mathematica gives $F(x) = \ln x - \ln\left(1+\sqrt{1-x^2}\right)$. Maple gives $F(x) = -\operatorname{arctanh}\left(1/\sqrt{1-x^2}\right)$. This function has domain $\left\{x \;\middle|\; |x| < 1, -1 < 1/\sqrt{1-x^2} < 1\right\} = \left\{x \;\middle|\; |x| < 1, 1/\sqrt{1-x^2} < 1\right\} = \left\{x \;\middle|\; |x| < 1, \sqrt{1-x^2} > 1\right\} = \emptyset$, the empty set! If we apply the command `convert(%,ln);` to Maple's answer, we get

$-\dfrac{1}{2}\ln\left(\dfrac{1}{\sqrt{1-x^2}}+1\right) + \dfrac{1}{2}\ln\left(1 - \dfrac{1}{\sqrt{1-x^2}}\right)$, which has the same domain, $\emptyset$.

48. Neither Maple nor Mathematica is able to evaluate $\int (1+\ln x)\sqrt{1+(x\ln x)^2}\,dx$. However, if we let $u = x\ln x$, then $du = (1+\ln x)\,dx$ and the integral is simply $\int \sqrt{1+u^2}\,du$, which any CAS can evaluate. The antiderivative is $\frac{1}{2}\ln\left(x\ln x + \sqrt{1+(x\ln x)^2}\right) + \frac{1}{2}x\ln x\sqrt{1+(x\ln x)^2} + C$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

DISCOVERY PROJECT Patterns in Integrals

1. (a) The CAS results are listed. Note that the absolute value symbols are missing, as is the familiar "$+C$".

(i) $\displaystyle\int \frac{1}{(x+2)(x+3)}\,dx = \ln(x+2) - \ln(x+3)$ (ii) $\displaystyle\int \frac{1}{(x+1)(x+5)}\,dx = \frac{\ln(x+1)}{4} - \frac{\ln(x+5)}{4}$

(iii) $\displaystyle\int \frac{1}{(x+2)(x-5)}\,dx = \frac{\ln(x-5)}{7} - \frac{\ln(x+2)}{7}$ (iv) $\displaystyle\int \frac{1}{(x+2)^2}\,dx = -\frac{1}{x+2}$

(b) If $a \neq b$, it appears that $\ln(x+a)$ is divided by $b-a$ and $\ln(x+b)$ is divided by $a-b$, so we guess that $\displaystyle\int \frac{1}{(x+a)(x+b)}\,dx = \frac{\ln(x+a)}{b-a} + \frac{\ln(x+b)}{a-b} + C$. If $a=b$, as in part (a)(iv), it appears that $\displaystyle\int \frac{1}{(x+a)^2}\,dx = -\frac{1}{x+a} + C$.

(c) The CAS verifies our guesses. Now $\dfrac{1}{(x+a)(x+b)} = \dfrac{A}{x+a} + \dfrac{B}{x+b} \quad\Rightarrow\quad 1 = A(x+b) + B(x+a)$.

Setting $x=-b$ gives $B = 1/(a-b)$ and setting $x=-a$ gives $A = 1/(b-a)$. So

$$\int \frac{1}{(x+a)(x+b)}\,dx = \int \left[\frac{1/(b-a)}{x+a} + \frac{1/(a-b)}{x+b}\right] dx = \frac{\ln|x+a|}{b-a} + \frac{\ln|x+b|}{a-b} + C$$

and our guess for $a \neq b$ is correct. If $a=b$, then $\dfrac{1}{(x+a)(x+b)} = \dfrac{1}{(x+a)^2} = (x+a)^{-2}$. Letting $u = x+a \quad\Rightarrow$ $du = dx$, we have $\displaystyle\int (x+a)^{-2}\,dx = \int u^{-2}\,du = -\frac{1}{u} + C = -\frac{1}{x+a} + C$, and our guess for $a=b$ is also correct.

2. (a) (i) $\displaystyle\int \sin x\,\cos 2x\,dx = \frac{\cos x}{2} - \frac{\cos 3x}{6}$ (ii) $\displaystyle\int \sin 3x\,\cos 7x\,dx = \frac{\cos 4x}{8} - \frac{\cos 10x}{20}$

(iii) $\displaystyle\int \sin 8x\,\cos 3x\,dx = -\frac{\cos 11x}{22} - \frac{\cos 5x}{10}$

(b) Looking at the sums and differences of a and b in part (a), we guess that

$$\int \sin ax\,\cos bx\,dx = \frac{\cos((a-b)x)}{2(b-a)} - \frac{\cos((a+b)x)}{2(a+b)} + C$$

Note that $\cos((a-b)x) = \cos((b-a)x)$.

(c) The CAS verifies our guess. Again, we can prove that the guess is correct by differentiating:

$$\begin{aligned}
\frac{d}{dx}\left[\frac{\cos((a-b)x)}{2(b-a)} - \frac{\cos((a+b)x)}{2(a+b)}\right] &= \frac{1}{2(b-a)}\,[-\sin((a-b)x)](a-b) - \frac{1}{2(a+b)}\,[-\sin((a+b)x)](a+b) \\
&= \tfrac{1}{2}\sin(ax-bx) + \tfrac{1}{2}\sin(ax+bx) \\
&= \tfrac{1}{2}(\sin ax\,\cos bx - \cos ax\,\sin bx) + \tfrac{1}{2}(\sin ax\,\cos bx + \cos ax\,\sin bx) \\
&= \sin ax\,\cos bx
\end{aligned}$$

Our formula is valid for $a \neq b$.

3. (a) (i) $\int \ln x\,dx = x\ln x - x$ (ii) $\int x\ln x\,dx = \frac{1}{2}x^2\ln x - \frac{1}{4}x^2$

(iii) $\int x^2\ln x\,dx = \frac{1}{3}x^3\ln x - \frac{1}{9}x^3$ (iv) $\int x^3\ln x\,dx = \frac{1}{4}x^4\ln x - \frac{1}{16}x^4$

(v) $\int x^7\ln x\,dx = \frac{1}{8}x^8\ln x - \frac{1}{64}x^8$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) We guess that $\int x^n \ln x\,dx = \dfrac{1}{n+1}x^{n+1}\ln x - \dfrac{1}{(n+1)^2}x^{n+1}$.

(c) Let $u = \ln x$, $dv = x^n\,dx \quad\Rightarrow\quad du = \dfrac{dx}{x}$, $v = \dfrac{1}{n+1}x^{n+1}$. Then

$$\int x^n \ln x\,dx = \frac{1}{n+1}x^{n+1}\ln x - \frac{1}{n+1}\int x^n\,dx = \frac{1}{n+1}x^{n+1}\ln x - \frac{1}{n+1}\cdot\frac{1}{n+1}x^{n+1},$$

which verifies our guess. We must have $n+1 \neq 0 \quad\Leftrightarrow\quad n \neq -1$.

4. (a) (i) $\int xe^x\,dx = e^x(x-1)$ (ii) $\int x^2e^x\,dx = e^x(x^2-2x+2)$

(iii) $\int x^3e^x\,dx = e^x(x^3-3x^2+6x-6)$ (iv) $\int x^4e^x\,dx = e^x(x^4-4x^3+12x^2-24x+24)$

(v) $\int x^5e^x\,dx = e^x\left(x^5-5x^4+20x^3-60x^2+120x-120\right)$

(b) Notice from part (a) that we can write

$$\int x^4e^x\,dx = e^x(x^4-4x^3+4\cdot 3x^2-4\cdot 3\cdot 2x+4\cdot 3\cdot 2\cdot 1)$$

and $$\int x^5e^x\,dx = e^x(x^5-5x^4+5\cdot 4x^3-5\cdot 4\cdot 3x^2+5\cdot 4\cdot 3\cdot 2x-5\cdot 4\cdot 3\cdot 2\cdot 1)$$

So we guess that

$$\begin{aligned}\int x^6e^x\,dx &= e^x(x^6-6x^5+6\cdot 5x^4-6\cdot 5\cdot 4x^3+6\cdot 5\cdot 4\cdot 3x^2-6\cdot 5\cdot 4\cdot 3\cdot 2x+6\cdot 5\cdot 4\cdot 3\cdot 2\cdot 1)\\ &= e^x(x^6-6x^5+30x^4-120x^3+360x^2-720x+720)\end{aligned}$$

The CAS verifies our guess.

(c) From the results in part (a), as well as our prediction in part (b), we speculate that

$$\int x^ne^x\,dx = e^x\left[x^n - nx^{n-1}+n(n-1)x^{n-2}-n(n-1)(n-2)x^{n-3}+\cdots\pm n!x\mp n!\right] = e^x\sum_{i=0}^{n}(-1)^{n-i}\frac{n!}{i!}\,x^i.$$

(We have reversed the order of the polynomial's terms.)

(d) Let S_n be the statement that $\int x^ne^x\,dx = e^x\sum\limits_{i=0}^{n}(-1)^{n-i}\dfrac{n!}{i!}\,x^i$.

S_1 is true by part (a)(i). Suppose S_k is true for some k, and consider S_{k+1}. Integrating by parts with $u = x^{k+1}$, $dv = e^x\,dx \quad\Rightarrow\quad du = (k+1)x^k\,dx$, $v = e^x$, we get

$$\begin{aligned}\int x^{k+1}e^x\,dx &= x^{k+1}e^x - (k+1)\int x^ke^x\,dx = x^{k+1}e^x - (k+1)\left[e^x\sum_{i=0}^{k}(-1)^{k-i}\frac{k!}{i!}\,x^i\right]\\ &= e^x\left[x^{k+1}-(k+1)\sum_{i=0}^{k}(-1)^{k-i}\frac{k!}{i!}\,x^i\right] = e^x\left[x^{k+1}+\sum_{i=0}^{k}(-1)^{k-i+1}\frac{(k+1)k!}{i!}\,x^i\right]\\ &= e^x\sum_{i=0}^{k+1}(-1)^{(k+1)-i}\frac{(k+1)!}{i!}\,x^i\end{aligned}$$

This verifies S_n for $n = k+1$. Thus, by mathematical induction, S_n is true for all n, where n is a positive integer.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7.7 Approximate Integration

1. (a) $\Delta x = (b-a)/n = (4-0)/2 = 2$

$$L_2 = \sum_{i=1}^{2} f(x_{i-1})\,\Delta x = f(x_0)\cdot 2 + f(x_1)\cdot 2 = 2\,[f(0)+f(2)] = 2(0.5+2.5) = 6$$

$$R_2 = \sum_{i=1}^{2} f(x_i)\,\Delta x = f(x_1)\cdot 2 + f(x_2)\cdot 2 = 2\,[f(2)+f(4)] = 2(2.5+3.5) = 12$$

$$M_2 = \sum_{i=1}^{2} f(\overline{x}_i)\Delta x = f(\overline{x}_1)\cdot 2 + f(\overline{x}_2)\cdot 2 = 2\,[f(1)+f(3)] \approx 2(1.6+3.2) = 9.6$$

(b)

L_2 is an underestimate, since the area under the small rectangles is less than the area under the curve, and R_2 is an overestimate, since the area under the large rectangles is greater than the area under the curve. It appears that M_2 is an overestimate, though it is fairly close to I. See the solution to Exercise 47 for a proof of the fact that if f is concave down on $[a,b]$, then the Midpoint Rule is an overestimate of $\int_a^b f(x)\,dx$.

(c) $T_2 = \left(\frac{1}{2}\,\Delta x\right)[f(x_0)+2f(x_1)+f(x_2)] = \frac{2}{2}[f(0)+2f(2)+f(4)] = 0.5+2(2.5)+3.5 = 9.$

This approximation is an underestimate, since the graph is concave down. Thus, $T_2 = 9 < I$. See the solution to Exercise 47 for a general proof of this conclusion.

(d) For any n, we will have $L_n < T_n < I < M_n < R_n$.

2.

The diagram shows that $L_4 > T_4 > \int_0^2 f(x)\,dx > R_4$, and it appears that M_4 is a bit less than $\int_0^2 f(x)\,dx$. In fact, for any function that is concave upward, it can be shown that $L_n > T_n > \int_0^2 f(x)\,dx > M_n > R_n$.

(a) Since $0.9540 > 0.8675 > 0.8632 > 0.7811$, it follows that $L_n = 0.9540$, $T_n = 0.8675$, $M_n = 0.8632$, and $R_n = 0.7811$.

(b) Since $M_n < \int_0^2 f(x)\,dx < T_n$, we have $0.8632 < \int_0^2 f(x)\,dx < 0.8675$.

3. $f(x) = \cos\left(x^2\right)$, $\Delta x = \frac{1-0}{4} = \frac{1}{4}$

(a) $T_4 = \frac{1}{4\cdot 2}\left[f(0)+2f\left(\frac{1}{4}\right)+2f\left(\frac{2}{4}\right)+2f\left(\frac{3}{4}\right)+f(1)\right] \approx 0.895759$

(b) $M_4 = \frac{1}{4}\left[f\left(\frac{1}{8}\right)+f\left(\frac{3}{8}\right)+f\left(\frac{5}{8}\right)+f\left(\frac{7}{8}\right)\right] \approx 0.908907$

The graph shows that f is concave down on $[0,1]$. So T_4 is an underestimate and M_4 is an overestimate. We can conclude that $0.895759 < \int_0^1 \cos\left(x^2\right)\,dx < 0.908907$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4.

(a) $f(x) = \sin\left(\frac{1}{2}x^2\right)$. Since f is increasing on $[0, 1]$, L_2 will underestimate I (since the area of the darkest rectangle is less than the area under the curve), and R_2 will overestimate I. Since f is concave upward on $[0, 1]$, M_2 will underestimate I and T_2 will overestimate I (the area under the straight line segments is greater than the area under the curve).

(b) For any n, we will have $L_n < M_n < I < T_n < R_n$.

(c) $L_5 = \sum_{i=1}^{5} f(x_{i-1})\ \Delta x = \frac{1}{5}[f(0.0) + f(0.2) + f(0.4) + f(0.6) + f(0.8)] \approx 0.1187$

$R_5 = \sum_{i=1}^{5} f(x_i)\ \Delta x = \frac{1}{5}[f(0.2) + f(0.4) + f(0.6) + f(0.8) + f(1)] \approx 0.2146$

$M_5 = \sum_{i=1}^{5} f(\overline{x}_i)\ \Delta x = \frac{1}{5}[f(0.1) + f(0.3) + f(0.5) + f(0.7) + f(0.9)] \approx 0.1622$

$T_5 = \left(\frac{1}{2}\,\Delta x\right)[f(0) + 2f(0.2) + 2f(0.4) + 2f(0.6) + 2f(0.8) + f(1)] \approx 0.1666$

From the graph, it appears that the Midpoint Rule gives the best approximation. (This is in fact the case, since $I \approx 0.16371405$.)

5. (a) $f(x) = x \sin x,\ \Delta x = \dfrac{b-a}{n} = \dfrac{\pi - 0}{6} = \dfrac{\pi}{6}$

$$M_6 = \frac{\pi}{6}\left[f\left(\frac{\pi}{12}\right) + f\left(\frac{3\pi}{12}\right) + f\left(\frac{5\pi}{12}\right) + f\left(\frac{7\pi}{12}\right) + f\left(\frac{9\pi}{12}\right) + f\left(\frac{11\pi}{12}\right)\right] \approx 3.177769$$

(b) $S_6 = \dfrac{\pi}{6 \cdot 3}\left[f(0) + 4f\left(\dfrac{\pi}{6}\right) + 2f\left(\dfrac{2\pi}{6}\right) + 4f\left(\dfrac{3\pi}{6}\right) + 2f\left(\dfrac{4\pi}{6}\right) + 4f\left(\dfrac{5\pi}{6}\right) + f\left(\dfrac{6\pi}{6}\right)\right] \approx 3.142949$

Actual: $I = \int_0^{\pi} x \sin x\, dx = \Big[-x\cos x + \sin x\Big]_0^{\pi}$ $\quad$ [use parts with $u = x$ and $dv = \sin x\, dx$]

$= (-\pi(-1) - 0) - (0 + 0) = \pi \approx 3.141593$

Errors: $E_M = \text{actual} - M_6 \approx 3.141593 - 3.177769 \approx -0.036176$

$Es = \text{actual} - S_6 \approx 3.141593 - 3.142949 \approx -0.001356$

6. (a) $f(x) = \dfrac{x}{\sqrt{1+x^2}},\ \Delta x = \dfrac{b-a}{n} = \dfrac{2-0}{8} = \dfrac{1}{4}$

$M_8 = \frac{1}{4}\left[f\left(\frac{1}{8}\right) + f\left(\frac{3}{8}\right) + f\left(\frac{5}{8}\right) + f\left(\frac{7}{8}\right) + f\left(\frac{9}{8}\right) + f\left(\frac{11}{8}\right) + f\left(\frac{13}{8}\right) + f\left(\frac{15}{8}\right)\right] \approx 1.238455$

(b) $S_8 = \dfrac{1}{4 \cdot 3}\left[f(0) + 4f\left(\dfrac{1}{4}\right) + 2f\left(\dfrac{2}{4}\right) + 4f\left(\dfrac{3}{4}\right) + 2f\left(\dfrac{4}{4}\right) + 4f\left(\dfrac{5}{4}\right) + 2f\left(\dfrac{6}{4}\right) + 4f\left(\dfrac{7}{4}\right) + f\left(\dfrac{8}{4}\right)\right]$

≈ 1.236147

Actual: $I = \displaystyle\int_0^2 \frac{x}{\sqrt{1+x^2}}\, dx = \Big[\sqrt{1+x^2}\Big]_0^2$ $\quad$ [$u = 1 + x^2$, $du = 2x\, dx$]

$= \sqrt{1+4} - \sqrt{1} = \sqrt{5} - 1 \approx 1.236068$

Errors: $E_M = \text{actual} - M_8 \approx 1.236068 - 1.238455 \approx -0.002387$

$Es = \text{actual} - S_8 \approx 1.236068 - 1.236147 \approx -0.000079$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. $f(x) = \sqrt{1+x^3}$, $\Delta x = \dfrac{b-a}{n} = \dfrac{1-0}{4} = \dfrac{1}{4}$

(a) $T_4 = \frac{1}{4\cdot 2}[f(0) + 2f(0.25) + 2f(0.5) + 2f(0.75) + f(1)] \approx 1.116993$

(b) $M_4 = \frac{1}{4}[f(0.125) + f(0.375) + f(0.625) + f(0.875)] \approx 1.108667$

(c) $S_4 = \frac{1}{4\cdot 3}[f(0) + 4f(0.25) + 2f(0.5) + 4f(0.75) + f(1)] \approx 1.111363$

8. $f(x) = \sin\sqrt{x}$, $\Delta x = \dfrac{b-a}{n} = \dfrac{4-1}{6} = \dfrac{3}{6} = \dfrac{1}{2}$

(a) $T_6 = \frac{1}{2\cdot 2}[f(1) + 2f(1.5) + 2f(2) + 2f(2.5) + 2f(3) + 2f(3.5) + f(4)] \approx 2.873085$

(b) $M_6 = \frac{1}{2}[f(1.25) + f(1.75) + f(2.25) + f(2.75) + f(3.25) + f(3.75)] \approx 2.884712$

(c) $S_6 = \frac{1}{2\cdot 3}[f(1) + 4f(1.5) + 2f(2) + 4f(2.5) + 2f(3) + 4f(3.5) + f(4)] \approx 2.880721$

9. $f(x) = \sqrt{e^x - 1}$, $\Delta x = \dfrac{b-a}{n} = \dfrac{1-0}{10} = \dfrac{1}{10}$

(a) $T_{10} = \frac{1}{10\cdot 2}[f(0) + 2f(0.1) + 2f(0.2) + 2f(0.3) + 2f(0.4) + 2f(0.5) + 2f(0.6)$
$+ 2f(0.7) + 2f(0.8) + 2f(0.9) + f(1)]$
≈ 0.777722

(b) $M_{10} = \frac{1}{10}[f(0.05) + f(0.15) + f(0.25) + f(0.35) + f(0.45) + f(0.55)$
$+ f(0.65) + f(0.75) + f(0.85) + f(0.95)]$
≈ 0.784958

(c) $S_{10} = \frac{1}{10\cdot 3}[f(0) + 4f(0.1) + 2f(0.2) + 4f(0.3) + 2f(0.4) + 4f(0.5) + 2f(0.6)$
$+ 4f(0.7) + 2f(0.8) + 4f(0.9) + f(1)]$
≈ 0.780895

10. $f(x) = \sqrt[3]{1-x^2}$, $\Delta x = \dfrac{b-a}{n} = \dfrac{2-0}{10} = \dfrac{2}{10} = \dfrac{1}{5}$

(a) $T_{10} = \frac{1}{5\cdot 2}[f(0) + 2f(0.2) + 2f(0.4) + 2f(0.6) + 2f(0.8) + 2f(1)$
$+ 2f(1.2) + 2f(1.4) + 2f(1.6) + 2f(1.8) + f(2)]$
≈ -0.186646

(b) $M_{10} = \frac{1}{5}[f(0.1) + f(0.3) + f(0.5) + f(0.7) + f(0.9) + f(1.1) + f(1.3) + f(1.5) + f(1.7) + f(1.9)]$
≈ -0.184073

(c) $S_{10} = \frac{1}{5\cdot 3}[f(0) + 4f(0.2) + 2f(0.4) + 4f(0.6) + 2f(0.8) + 4f(1) + 2f(1.2)$
$+ 4f(1.4) + 2f(1.6) + 4f(1.8) + f(2)]$
≈ -0.183984

11. $f(x) = e^{x+\cos x}$, $\Delta x = \frac{2-(-1)}{6} = \frac{1}{2}$

(a) $T_6 = \frac{1}{2}\,[f(-1.0) + 2f(-0.5) + 2f(0) + 2f(0.5) + 2f(1) + 2f(1.5) + f(2.0)] \approx 10.185560$

(b) $M_6 = \frac{1}{2}\,[f(-0.75) + f(-0.25) + f(0.25) + f(0.75) + f(1.25) + f(1.75)] \approx 10.208618$

(c) $S_6 = \frac{1}{2\cdot 3}\,[f(-1.0) + 4f(-0.5) + 2f(0) + 4f(0.5) + 2f(1.0) + 4f(1.5) + f(2.0)] \approx 10.201790$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $f(x) = e^{1/x}$, $\Delta x = \frac{3-1}{8} = \frac{1}{4}$

(a) $T_8 = \frac{1}{4 \cdot 2}\left[f(1) + 2f\left(\frac{5}{4}\right) + 2f\left(\frac{3}{2}\right) + 2f\left(\frac{7}{4}\right) + 2f(2) + 2f\left(\frac{9}{4}\right) + 2f\left(\frac{5}{2}\right) + 2f\left(\frac{11}{4}\right) + f(3)\right] \approx 3.534934$

(b) $M_8 = \frac{1}{4}\left[f\left(\frac{9}{8}\right) + f\left(\frac{11}{8}\right) + f\left(\frac{13}{8}\right) + f\left(\frac{15}{8}\right) + f\left(\frac{17}{8}\right) + f\left(\frac{19}{8}\right) + f\left(\frac{21}{8}\right) + f\left(\frac{23}{8}\right)\right] \approx 3.515248$

(c) $S_8 = \frac{1}{4 \cdot 3}\left[f(1) + 4f\left(\frac{5}{4}\right) + 2f\left(\frac{3}{2}\right) + 4f\left(\frac{7}{4}\right) + 2f(2) + 4f\left(\frac{9}{4}\right) + 2f\left(\frac{5}{2}\right) + 4f\left(\frac{11}{4}\right) + f(3)\right] \approx 3.522375$

13. $f(y) = \sqrt{y}\cos y$, $\Delta y = \frac{4-0}{8} = \frac{1}{2}$

(a) $T_8 = \frac{1}{2 \cdot 2}\left[f(0) + 2f\left(\frac{1}{2}\right) + 2f(1) + 2f\left(\frac{3}{2}\right) + 2f(2) + 2f\left(\frac{5}{2}\right) + 2f(3) + 2f\left(\frac{7}{2}\right) + f(4)\right] \approx -2.364034$

(b) $M_8 = \frac{1}{2}\left[f\left(\frac{1}{4}\right) + f\left(\frac{3}{4}\right) + f\left(\frac{5}{4}\right) + f\left(\frac{7}{4}\right) + f\left(\frac{9}{4}\right) + f\left(\frac{11}{4}\right) + f\left(\frac{13}{4}\right) + f\left(\frac{15}{4}\right)\right] \approx -2.310690$

(c) $S_8 = \frac{1}{2 \cdot 3}\left[f(0) + 4f\left(\frac{1}{2}\right) + 2f(1) + 4f\left(\frac{3}{2}\right) + 2f(2) + 4f\left(\frac{5}{2}\right) + 2f(3) + 4f\left(\frac{7}{2}\right) + f(4)\right] \approx -2.346520$

14. $f(t) = \dfrac{1}{\ln t}$, $\Delta t = \dfrac{3-2}{10} = \dfrac{1}{10}$

(a) $T_{10} = \frac{1}{10 \cdot 2}\{f(2) + 2[f(2.1) + f(2.2) + \cdots + f(2.9)] + f(3)\} \approx 1.119061$

(b) $M_{10} = \frac{1}{10}[f(2.05) + f(2.15) + \cdots + f(2.85) + f(2.95)] \approx 1.118107$

(c) $S_{10} = \frac{1}{10 \cdot 3}[f(2) + 4f(2.1) + 2f(2.2) + 4f(2.3) + 2f(2.4) + 4f(2.5) + 2f(2.6)$
$+ 4f(2.7) + 2f(2.8) + 4f(2.9) + f(3)] \approx 1.118428$

15. $f(x) = \dfrac{x^2}{1+x^4}$, $\Delta x = \dfrac{1-0}{10} = \dfrac{1}{10}$

(a) $T_{10} = \frac{1}{10 \cdot 2}\{f(0) + 2[f(0.1 + f(0.2) + \cdots + f(0.9)] + f(1)\} \approx 0.243747$

(b) $M_{10} = \frac{1}{10}[f(0.05) + f(0.15) + \cdots + f(0.85) + f(0.95)] \approx 0.243748$

(c) $S_{10} = \frac{1}{10 \cdot 3}[f(0) + 4f(0.1) + 2f(0.2) + 4f(0.3) + 2f(0.4) + 4f(0.5) + 2f(0.6)$
$+ 4f(0.7) + 2f(0.8) + 4f(0.9) + f(1)] \approx 0.243751$

Note: $\int_0^1 f(x)\,dx \approx 0.24374775$. This is a rare case where the Trapezoidal and Midpoint Rules give better approximations than Simpson's Rule.

16. $f(t) = \dfrac{\sin t}{t}$, $\Delta t = \dfrac{3-1}{4} = \dfrac{1}{2}$

(a) $T_4 = \frac{1}{2 \cdot 2}[f(1) + 2f(1.5) + 2f(2) + 2f(2.5) + f(3)] \approx 0.901645$

(b) $M_4 = \frac{1}{2}[f(1.25) + f(1.75) + f(2.25) + f(2.75)] \approx 0.903031$

(c) $S_4 = \frac{1}{2 \cdot 3}[f(1) + 4f(1.5) + 2f(2) + 4f(2.5) + f(3)] \approx 0.902558$

17. $f(x) = \ln(1 + e^x)$, $\Delta x = \frac{4-0}{8} = \frac{1}{2}$

(a) $T_8 = \frac{1}{2 \cdot 2}\{f(0) + 2[f(0.5) + f(1) + \cdots + f(3) + f(3.5)] + f(4)\} \approx 8.814278$

(b) $M_8 = \frac{1}{2}[f(0.25) + f(0.75) + \cdots + f(3.25) + f(3.75)] \approx 8.799212$

(c) $S_8 = \frac{1}{2 \cdot 3}[f(0) + 4f(0.5) + 2f(1) + 4f(1.5) + 2f(2) + 4f(2.5) + 2f(3) + 4f(3.5) + f(4)] \approx 8.804229$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $f(x) = \sqrt{x + x^3}$, $\Delta x = \frac{1-0}{10} = \frac{1}{10}$

(a) $T_{10} = \frac{1}{2 \cdot 2}\{f(0) + 2[f(0.1) + f(0.2) + \cdots + f(0.8) + f(0.9)] + f(1)\} \approx 0.787092$

(b) $M_{10} = \frac{1}{2}[f(0.05) + f(0.15) + \cdots + f(0.85) + f(0.95)] \approx 0.793821$

(c) $S_{10} = \frac{1}{2 \cdot 3}[f(0) + 4f(0.1) + 2f(0.2) + 4f(0.3) + 2f(0.4) + 4f(0.5) + 2f(0.6)$
$+ 4f(0.7) + 2f(0.8) + 4f(0.9) + f(1)]$
≈ 0.789915

19. $f(x) = \cos(x^2)$, $\Delta x = \frac{1-0}{8} = \frac{1}{8}$

(a) $T_8 = \frac{1}{8 \cdot 2}\left\{f(0) + 2\left[f\left(\frac{1}{8}\right) + f\left(\frac{2}{8}\right) + \cdots + f\left(\frac{7}{8}\right)\right] + f(1)\right\} \approx 0.902333$

$M_8 = \frac{1}{8}\left[f\left(\frac{1}{16}\right) + f\left(\frac{3}{16}\right) + f\left(\frac{5}{16}\right) + \cdots + f\left(\frac{15}{16}\right)\right] = 0.905620$

(b) $f(x) = \cos(x^2)$, $f'(x) = -2x\sin(x^2)$, $f''(x) = -2\sin(x^2) - 4x^2\cos(x^2)$. For $0 \le x \le 1$, sin and cos are positive, so $|f''(x)| = 2\sin(x^2) + 4x^2\cos(x^2) \le 2 \cdot 1 + 4 \cdot 1 \cdot 1 = 6$ since $\sin(x^2) \le 1$ and $\cos\left(x^2\right) \le 1$ for all x, and $x^2 \le 1$ for $0 \le x \le 1$. So for $n = 8$, we take $K = 6$, $a = 0$, and $b = 1$ in Theorem 3, to get $|E_T| \le 6 \cdot 1^3/(12 \cdot 8^2) = \frac{1}{128} = 0.0078125$ and $|E_M| \le \frac{1}{256} = 0.00390625$. [A better estimate is obtained by noting from a graph of f'' that $|f''(x)| \le 4$ for $0 \le x \le 1$.]

(c) Take $K = 6$ [as in part (b)] in Theorem 3. $|E_T| \le \dfrac{K(b-a)^3}{12n^2} \le 0.0001 \quad \Leftrightarrow \quad \dfrac{6(1-0)^3}{12n^2} \le 10^{-4} \quad \Leftrightarrow$

$\dfrac{1}{2n^2} \le \dfrac{1}{10^4} \quad \Leftrightarrow \quad 2n^2 \ge 10^4 \quad \Leftrightarrow \quad n^2 \ge 5000 \quad \Leftrightarrow \quad n \ge 71$. Take $n = 71$ for T_n. For E_M, again take $K = 6$ in Theorem 3 to get $|E_M| \le 10^{-4} \quad \Leftrightarrow \quad 4n^2 \ge 10^4 \quad \Leftrightarrow \quad n^2 \ge 2500 \quad \Leftrightarrow \quad n \ge 50$. Take $n = 50$ for M_n.

20. $f(x) = e^{1/x}$, $\Delta x = \frac{2-1}{10} = \frac{1}{10}$

(a) $T_{10} = \frac{1}{10 \cdot 2}[f(1) + 2f(1.1) + 2f(1.2) + \cdots + 2f(1.9) + f(2)] \approx 2.021976$

$M_{10} = \frac{1}{10}\,[f(1.05) + f(1.15) + f(1.25) + \cdots + f(1.95)] \approx 2.019102$

(b) $f(x) = e^{1/x}$, $f'(x) = -\dfrac{1}{x^2}e^{1/x}$, $f''(x) = \dfrac{2x+1}{x^4}\,e^{1/x}$. Now f'' is decreasing on $[1, 2]$, so let $x = 1$ to take $K = 3e$.

$|E_T| \le \dfrac{3e(2-1)^3}{12(10)^2} = \dfrac{e}{400} \approx 0.006796. \quad |E_M| \le \dfrac{|E_T|}{2} = \dfrac{e}{800} \approx 0.003398.$

(c) Take $K = 3e$ [as in part (b)] in Theorem 3. $|E_T| \le \dfrac{K(b-a)^3}{12n^2} \le 0.0001 \quad \Leftrightarrow \quad \dfrac{3e(2-1)^3}{12n^2} \le 10^{-4} \quad \Leftrightarrow$

$\dfrac{e}{4n^2} \le \dfrac{1}{10^4} \quad \Leftrightarrow \quad n^2 \ge \dfrac{10^4 e}{4} \quad \Leftrightarrow \quad n \ge 83$. Take $n = 83$ for T_n. For E_M, again take $K = 3e$ in Theorem 3 to get $|E_M| \le 10^{-4} \quad \Leftrightarrow \quad n^2 \ge \dfrac{10^4 e}{8} \quad \Leftrightarrow \quad n \ge 59$. Take $n = 59$ for M_n.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. $f(x) = \sin x$, $\Delta x = \frac{\pi - 0}{10} = \frac{\pi}{10}$

(a) $T_{10} = \frac{\pi}{10 \cdot 2}\left[f(0) + 2f\left(\frac{\pi}{10}\right) + 2f\left(\frac{2\pi}{10}\right) + \cdots + 2f\left(\frac{9\pi}{10}\right) + f(\pi)\right] \approx 1.983524$

$M_{10} = \frac{\pi}{10}\left[f\left(\frac{\pi}{20}\right) + f\left(\frac{3\pi}{20}\right) + f\left(\frac{5\pi}{20}\right) + \cdots + f\left(\frac{19\pi}{20}\right)\right] \approx 2.008248$

$S_{10} = \frac{\pi}{10 \cdot 3}\left[f(0) + 4f\left(\frac{\pi}{10}\right) + 2f\left(\frac{2\pi}{10}\right) + 4f\left(\frac{3\pi}{10}\right) + \cdots + 4f\left(\frac{9\pi}{10}\right) + f(\pi)\right] \approx 2.000110$

Since $I = \int_0^\pi \sin x\,dx = \left[-\cos x\right]_0^\pi = 1 - (-1) = 2$, $E_T = I - T_{10} \approx 0.016476$, $E_M = I - M_{10} \approx -0.008248$, and $E_S = I - S_{10} \approx -0.000110$.

(b) $f(x) = \sin x \quad \Rightarrow \quad \left|f^{(n)}(x)\right| \le 1$, so take $K = 1$ for all error estimates.

$$|E_T| \le \frac{K(b-a)^3}{12n^2} = \frac{1(\pi - 0)^3}{12(10)^2} = \frac{\pi^3}{1200} \approx 0.025839. \qquad |E_M| \le \frac{|E_T|}{2} = \frac{\pi^3}{2400} \approx 0.012919.$$

$$|E_S| \le \frac{K(b-a)^5}{180n^4} = \frac{1(\pi - 0)^5}{180(10)^4} = \frac{\pi^5}{1{,}800{,}000} \approx 0.000170.$$

The actual error is about 64% of the error estimate in all three cases.

(c) $|E_T| \le 0.00001 \quad \Leftrightarrow \quad \dfrac{\pi^3}{12n^2} \le \dfrac{1}{10^5} \quad \Leftrightarrow \quad n^2 \ge \dfrac{10^5\pi^3}{12} \quad \Rightarrow \quad n \ge 508.3$. Take $n = 509$ for T_n.

$|E_M| \le 0.00001 \quad \Leftrightarrow \quad \dfrac{\pi^3}{24n^2} \le \dfrac{1}{10^5} \quad \Leftrightarrow \quad n^2 \ge \dfrac{10^5\pi^3}{24} \quad \Rightarrow \quad n \ge 359.4$. Take $n = 360$ for M_n.

$|E_S| \le 0.00001 \quad \Leftrightarrow \quad \dfrac{\pi^5}{180n^4} \le \dfrac{1}{10^5} \quad \Leftrightarrow \quad n^4 \ge \dfrac{10^5\pi^5}{180} \quad \Rightarrow \quad n \ge 20.3.$

Take $n = 22$ for S_n (since n must be even).

22. From Example 7(b), we take $K = 76e$ to get $|E_S| \le \dfrac{76e(1)^5}{180n^4} \le 0.00001 \quad \Rightarrow \quad n^4 \ge \dfrac{76e}{180(0.00001)} \quad \Rightarrow \quad n \ge 18.4.$

Take $n = 20$ (since n must be even).

23. (a) Using a CAS, we differentiate $f(x) = e^{\cos x}$ twice, and find that $f''(x) = e^{\cos x}(\sin^2 x - \cos x)$. From the graph, we see that the maximum value of $|f''(x)|$ occurs at the endpoints of the interval $[0, 2\pi]$. Since $f''(0) = -e$, we can use $K = e$ or $K = 2.8$.

(b) A CAS gives $M_{10} \approx 7.954926518$. (In Maple, use `Student[Calculus1][RiemannSum]` or `Student[Calculus1][ApproximateInt]`.)

(c) Using Theorem 3 for the Midpoint Rule, with $K = e$, we get $|E_M| \le \dfrac{e(2\pi - 0)^3}{24 \cdot 10^2} \approx 0.280945995$.

With $K = 2.8$, we get $|E_M| \le \dfrac{2.8(2\pi - 0)^3}{24 \cdot 10^2} = 0.289391916$.

(d) A CAS gives $I \approx 7.954926521$.

(e) The actual error is only about 3×10^{-9}, much less than the estimate in part (c).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(f) We use the CAS to differentiate twice more, and then graph

$f^{(4)}(x) = e^{\cos x}(\sin^4 x - 6\sin^2 x\,\cos x + 3 - 7\sin^2 x + \cos x)$.

From the graph, we see that the maximum value of $\left|f^{(4)}(x)\right|$ occurs at the endpoints of the interval $[0, 2\pi]$. Since $f^{(4)}(0) = 4e$, we can use $K = 4e$ or $K = 10.9$.

(g) A CAS gives $S_{10} \approx 7.953789422$. (In Maple, use `Student[Calculus1][ApproximateInt]`.)

(h) Using Theorem 4 with $K = 4e$, we get $|E_S| \le \dfrac{4e(2\pi - 0)^5}{180 \cdot 10^4} \approx 0.059153618$.

With $K = 10.9$, we get $|E_S| \le \dfrac{10.9(2\pi - 0)^5}{180 \cdot 10^4} \approx 0.059299814$.

(i) The actual error is about $7.954926521 - 7.953789422 \approx 0.00114$. This is quite a bit smaller than the estimate in part (h), though the difference is not nearly as great as it was in the case of the Midpoint Rule.

(j) To ensure that $|E_S| \le 0.0001$, we use Theorem 4: $|E_S| \le \dfrac{4e(2\pi)^5}{180 \cdot n^4} \le 0.0001 \quad \Rightarrow \quad \dfrac{4e(2\pi)^5}{180 \cdot 0.0001} \le n^4 \quad \Rightarrow$ $n^4 \ge 5{,}915{,}362 \quad \Leftrightarrow \quad n \ge 49.3$. So we must take $n \ge 50$ to ensure that $|I - S_n| \le 0.0001$. ($K = 10.9$ leads to the same value of n.)

24. (a) Using the CAS, we differentiate $f(x) = \sqrt{4 - x^3}$ twice, and find that $f''(x) = -\dfrac{9x^4}{4(4 - x^3)^{3/2}} - \dfrac{3x}{(4 - x^3)^{1/2}}$.

From the graph, we see that $|f''(x)| < 2.2$ on $[-1, 1]$.

(b) A CAS gives $M_{10} \approx 3.995804152$. (In Maple, use `Student[Calculus1][RiemannSum]` or `Student[Calculus1][ApproximateInt]`.)

(c) Using Theorem 3 for the Midpoint Rule, with $K = 2.2$, we get $|E_M| \le \dfrac{2.2\,[1 - (-1)]^3}{24 \cdot 10^2} \approx 0.00733$.

(d) A CAS gives $I \approx 3.995487677$.

(e) The actual error is about -0.0003165, much less than the estimate in part (c).

(f) We use the CAS to differentiate twice more, and then graph

$f^{(4)}(x) = \dfrac{9}{16}\dfrac{x^2(x^6 - 224x^3 - 1280)}{(4 - x^3)^{7/2}}$.

From the graph, we see that $\left|f^{(4)}(x)\right| < 18.1$ on $[-1, 1]$.

(g) A CAS gives $S_{10} \approx 3.995449790$. (In Maple, use `Student[Calculus1][ApproximateInt]`.)

(h) Using Theorem 4 with $K = 18.1$, we get $|E_S| \le \dfrac{18.1\,[1 - (-1)]^5}{180 \cdot 10^4} \approx 0.000322$.

(i) The actual error is about $3.995487677 - 3.995449790 \approx 0.0000379$. This is quite a bit smaller than the estimate in part (h).

(j) To ensure that $|E_S| \le 0.0001$, we use Theorem 4: $|E_S| \le \dfrac{18.1(2)^5}{180 \cdot n^4} \le 0.0001 \quad \Rightarrow \quad \dfrac{18.1(2)^5}{180 \cdot 0.0001} \le n^4 \quad \Rightarrow$ $n^4 \ge 32{,}178 \quad \Rightarrow n \ge 13.4$. So we must take $n \ge 14$ to ensure that $|I - S_n| \le 0.0001$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $I = \int_0^1 xe^x\,dx = [(x-1)e^x]_0^1$ [parts or Formula 96] $= 0-(-1) = 1$, $f(x) = xe^x$, $\Delta x = 1/n$

$n = 5$:
$$L_5 = \tfrac{1}{5}[f(0)+f(0.2)+f(0.4)+f(0.6)+f(0.8)] \approx 0.742943$$
$$R_5 = \tfrac{1}{5}[f(0.2)+f(0.4)+f(0.6)+f(0.8)+f(1)] \approx 1.286599$$
$$T_5 = \tfrac{1}{5\cdot 2}[f(0)+2f(0.2)+2f(0.4)+2f(0.6)+2f(0.8)+f(1)] \approx 1.014771$$
$$M_5 = \tfrac{1}{5}[f(0.1)+f(0.3)+f(0.5)+f(0.7)+f(0.9)] \approx 0.992621$$
$$E_L = I - L_5 \approx 1 - 0.742943 = 0.257057$$
$$E_R \approx 1 - 1.286599 = -0.286599$$
$$E_T \approx 1 - 1.014771 = -0.014771$$
$$E_M \approx 1 - 0.992621 = 0.007379$$

$n = 10$:
$$L_{10} = \tfrac{1}{10}[f(0)+f(0.1)+f(0.2)+\cdots+f(0.9)] \approx 0.867782$$
$$R_{10} = \tfrac{1}{10}[f(0.1)+f(0.2)+\cdots+f(0.9)+f(1)] \approx 1.139610$$
$$T_{10} = \tfrac{1}{10\cdot 2}\{f(0)+2[f(0.1)+f(0.2)+\cdots+f(0.9)]+f(1)\} \approx 1.003696$$
$$M_{10} = \tfrac{1}{10}[f(0.05)+f(0.15)+\cdots+f(0.85)+f(0.95)] \approx 0.998152$$
$$E_L = I - L_{10} \approx 1 - 0.867782 = 0.132218$$
$$E_R \approx 1 - 1.139610 = -0.139610$$
$$E_T \approx 1 - 1.003696 = -0.003696$$
$$E_M \approx 1 - 0.998152 = 0.001848$$

$n = 20$:
$$L_{20} = \tfrac{1}{20}[f(0)+f(0.05)+f(0.10)+\cdots+f(0.95)] \approx 0.932967$$
$$R_{20} = \tfrac{1}{20}[f(0.05)+f(0.10)+\cdots+f(0.95)+f(1)] \approx 1.068881$$
$$T_{20} = \tfrac{1}{20\cdot 2}\{f(0)+2[f(0.05)+f(0.10)+\cdots+f(0.95)]+f(1)\} \approx 1.000924$$
$$M_{20} = \tfrac{1}{20}[f(0.025)+f(0.075)+f(0.125)+\cdots+f(0.975)] \approx 0.999538$$
$$E_L = I - L_{20} \approx 1 - 0.932967 = 0.067033$$
$$E_R \approx 1 - 1.068881 = -0.068881$$
$$E_T \approx 1 - 1.000924 = -0.000924$$
$$E_M \approx 1 - 0.999538 = 0.000462$$

n	L_n	R_n	T_n	M_n
5	0.742943	1.286599	1.014771	0.992621
10	0.867782	1.139610	1.003696	0.998152
20	0.932967	1.068881	1.000924	0.999538

n	E_L	E_R	E_T	E_M
5	0.257057	−0.286599	−0.014771	0.007379
10	0.132218	−0.139610	−0.003696	0.001848
20	0.067033	−0.068881	−0.000924	0.000462

Observations:

1. E_L and E_R are always opposite in sign, as are E_T and E_M.
2. As n is doubled, E_L and E_R are decreased by about a factor of 2, and E_T and E_M are decreased by a factor of about 4.
3. The Midpoint approximation is about twice as accurate as the Trapezoidal approximation.
4. All the approximations become more accurate as the value of n increases.
5. The Midpoint and Trapezoidal approximations are much more accurate than the endpoint approximations.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $I = \int_1^2 \frac{1}{x^2}\,dx = \left[-\frac{1}{x}\right]_1^2 = -\frac{1}{2} - (-1) = \frac{1}{2}, f(x) = \frac{1}{x^2}, \Delta x = \frac{1}{n}$

$n = 5$: $L_5 = \frac{1}{5}[f(1) + f(1.2) + f(1.4) + f(1.6) + f(1.8)] \approx 0.580783$

$R_5 = \frac{1}{5}[f(1.2) + f(1.4) + f(1.6) + f(1.8) + f(2)] \approx 0.430783$

$T_5 = \frac{1}{5 \cdot 2}[f(1) + 2f(1.2) + 2f(1.4) + 2f(1.6) + 2f(1.8) + f(2)] \approx 0.505783$

$M_5 = \frac{1}{5}[f(1.1) + f(1.3) + f(1.5) + f(1.7) + f(1.9)] \approx 0.497127$

$E_L = I - L_5 \approx \frac{1}{2} - 0.580783 = -0.080783$

$E_R \approx \frac{1}{2} - 0.430783 = 0.069217$

$E_T \approx \frac{1}{2} - 0.505783 = -0.005783$

$E_M \approx \frac{1}{2} - 0.497127 = 0.002873$

$n = 10$: $L_{10} = \frac{1}{10}[f(1) + f(1.1) + f(1.2) + \cdots + f(1.9)] \approx 0.538955$

$R_{10} = \frac{1}{10}[f(1.1) + f(1.2) + \cdots + f(1.9) + f(2)] \approx 0.463955$

$T_{10} = \frac{1}{10 \cdot 2}\{f(1) + 2[f(1.1) + f(1.2) + \cdots + f(1.9)] + f(2)\} \approx 0.501455$

$M_{10} = \frac{1}{10}[f(1.05) + f(1.15) + \cdots + f(1.85) + f(1.95)] \approx 0.499274$

$E_L = I - L_{10} \approx \frac{1}{2} - 0.538955 = -0.038955$

$E_R \approx \frac{1}{2} - 0.463955 = 0.036049$

$E_T \approx \frac{1}{2} - 0.501455 = -0.001455$

$E_M \approx \frac{1}{2} - 0.499274 = 0.000726$

$n = 20$: $L_{20} = \frac{1}{20}[f(1) + f(1.05) + f(1.10) + \cdots + f(1.95)] \approx 0.519114$

$R_{20} = \frac{1}{20}[f(1.05) + f(1.10) + \cdots + f(1.95) + f(2)] \approx 0.481614$

$T_{20} = \frac{1}{20 \cdot 2}\{f(1) + 2[f(1.05) + f(1.10) + \cdots + f(1.95)] + f(2)\} \approx 0.500364$

$M_{20} = \frac{1}{20}[f(1.025) + f(1.075) + f(1.125) + \cdots + f(1.975)] \approx 0.499818$

$E_L = I - L_{20} \approx \frac{1}{2} - 0.519114 = -0.019114$

$E_R \approx \frac{1}{2} - 0.481614 = 0.018386$

$E_T \approx \frac{1}{2} - 0.500364 = -0.000364$

$E_M \approx \frac{1}{2} - 0.499818 = 0.000182$

n	L_n	R_n	T_n	M_n
5	0.580783	0.430783	0.505783	0.497127
10	0.538955	0.463955	0.501455	0.499274
20	0.519114	0.481614	0.500364	0.499818

n	E_L	E_R	E_T	E_M
5	−0.080783	0.069217	−0.005783	0.002873
10	−0.038955	0.036049	−0.001455	0.000726
20	−0.019114	0.018386	−0.000364	0.000182

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Observations:

1. E_L and E_R are always opposite in sign, as are E_T and E_M.
2. As n is doubled, E_L and E_R are decreased by about a factor of 2, and E_T and E_M are decreased by a factor of about 4.
3. The Midpoint approximation is about twice as accurate as the Trapezoidal approximation.
4. All the approximations become more accurate as the value of n increases.
5. The Midpoint and Trapezoidal approximations are much more accurate than the endpoint approximations.

27. $I = \int_0^2 x^4\,dx = \left[\frac{1}{5}x^5\right]_0^2 = \frac{32}{5} - 0 = 6.4$, $f(x) = x^4$, $\Delta x = \frac{2-0}{n} = \frac{2}{n}$

$n = 6$:
$$\begin{aligned}
T_6 &= \tfrac{2}{6\cdot 2}\left\{f(0) + 2\left[f\left(\tfrac{1}{3}\right) + f\left(\tfrac{2}{3}\right) + f\left(\tfrac{3}{3}\right) + f\left(\tfrac{4}{3}\right) + f\left(\tfrac{5}{3}\right)\right] + f(2)\right\} \approx 6.695473\\
M_6 &= \tfrac{2}{6}\left[f\left(\tfrac{1}{6}\right) + f\left(\tfrac{3}{6}\right) + f\left(\tfrac{5}{6}\right) + f\left(\tfrac{7}{6}\right) + f\left(\tfrac{9}{6}\right) + f\left(\tfrac{11}{6}\right)\right] \approx 6.252572\\
S_6 &= \tfrac{2}{6\cdot 3}\left[f(0) + 4f\left(\tfrac{1}{3}\right) + 2f\left(\tfrac{2}{3}\right) + 4f\left(\tfrac{3}{3}\right) + 2f\left(\tfrac{4}{3}\right) + 4f\left(\tfrac{5}{3}\right) + f(2)\right] \approx 6.403292\\
E_T &= I - T_6 \approx 6.4 - 6.695473 = -0.295473\\
E_M &\approx 6.4 - 6.252572 = 0.147428\\
E_S &\approx 6.4 - 6.403292 = -0.003292
\end{aligned}$$

$n = 12$:
$$\begin{aligned}
T_{12} &= \tfrac{2}{12\cdot 2}\left\{f(0) + 2\left[f\left(\tfrac{1}{6}\right) + f\left(\tfrac{2}{6}\right) + f\left(\tfrac{3}{6}\right) + \cdots + f\left(\tfrac{11}{6}\right)\right] + f(2)\right\} \approx 6.474023\\
M_6 &= \tfrac{2}{12}\left[f\left(\tfrac{1}{12}\right) + f\left(\tfrac{3}{12}\right) + f\left(\tfrac{5}{12}\right) + \cdots + f\left(\tfrac{23}{12}\right)\right] \approx 6.363008\\
S_6 &= \tfrac{2}{12\cdot 3}\left[f(0) + 4f\left(\tfrac{1}{6}\right) + 2f\left(\tfrac{2}{6}\right) + 4f\left(\tfrac{3}{6}\right) + 2f\left(\tfrac{4}{6}\right) + \cdots + 4f\left(\tfrac{11}{6}\right) + f(2)\right] \approx 6.400206\\
E_T &= I - T_{12} \approx 6.4 - 6.474023 = -0.074023\\
E_M &\approx 6.4 - 6.363008 = 0.036992\\
E_S &\approx 6.4 - 6.400206 = -0.000206
\end{aligned}$$

n	T_n	M_n	S_n
6	6.695473	6.252572	6.403292
12	6.474023	6.363008	6.400206

n	E_T	E_M	E_S
6	−0.295473	0.147428	−0.003292
12	−0.074023	0.036992	−0.000206

Observations:

1. E_T and E_M are opposite in sign and decrease by a factor of about 4 as n is doubled.
2. The Simpson's approximation is much more accurate than the Midpoint and Trapezoidal approximations, and E_S seems to decrease by a factor of about 16 as n is doubled.

28. $I = \displaystyle\int_1^4 \frac{1}{\sqrt{x}}\,dx = \left[2\sqrt{x}\right]_1^4 = 4 - 2 = 2$, $f(x) = \dfrac{1}{\sqrt{x}}$, $\Delta x = \dfrac{4-1}{n} = \dfrac{3}{n}$

$n = 6$:
$$\begin{aligned}
T_6 &= \tfrac{3}{6\cdot 2}\left\{f(1) + 2\left[f\left(\tfrac{3}{2}\right) + f\left(\tfrac{4}{2}\right) + f\left(\tfrac{5}{2}\right) + f\left(\tfrac{6}{2}\right) + f\left(\tfrac{7}{2}\right)\right] + f(4)\right\} \approx 2.008966\\
M_6 &= \tfrac{3}{6}\left[f\left(\tfrac{5}{4}\right) + f\left(\tfrac{7}{4}\right) + f\left(\tfrac{9}{4}\right) + f\left(\tfrac{11}{4}\right) + f\left(\tfrac{13}{4}\right) + f\left(\tfrac{15}{4}\right)\right] \approx 1.995572\\
S_6 &= \tfrac{3}{6\cdot 3}\left[f(1) + 4f\left(\tfrac{3}{2}\right) + 2f\left(\tfrac{4}{2}\right) + 4f\left(\tfrac{5}{2}\right) + 2f\left(\tfrac{6}{2}\right) + 4f\left(\tfrac{7}{2}\right) + f(4)\right] \approx 2.000469\\
E_T &= I - T_6 \approx 2 - 2.008966 = -0.008966,\\
E_M &\approx 2 - 1.995572 = 0.004428,\\
E_S &\approx 2 - 2.000469 = -0.000469
\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$n = 12$: $T_{12} = \frac{3}{12 \cdot 2}\left\{f(1) + 2\left[f\left(\frac{5}{4}\right) + f\left(\frac{6}{4}\right) + f\left(\frac{7}{4}\right) + \cdots + f\left(\frac{15}{4}\right)\right] + f(4)\right\} \approx 2.002269$

$M_{12} = \frac{3}{12}\left[f\left(\frac{9}{8}\right) + f\left(\frac{11}{8}\right) + f\left(\frac{13}{8}\right) + \cdots + f\left(\frac{31}{8}\right)\right] \approx 1.998869$

$S_{12} = \frac{3}{12 \cdot 3}\left[f(1) + 4f\left(\frac{5}{4}\right) + 2f\left(\frac{6}{4}\right) + 4f\left(\frac{7}{4}\right) + 2f\left(\frac{8}{4}\right) + \cdots + 4f\left(\frac{15}{4}\right) + f(4)\right] \approx 2.000036$

$E_T = I - T_{12} \approx 2 - 2.002269 = -0.002269$

$E_M \approx 2 - 1.998869 = 0.001131$

$E_S \approx 2 - 2.000036 = -0.000036$

n	T_n	M_n	S_n
6	2.008966	1.995572	2.000469
12	2.002269	1.998869	2.000036

n	E_T	E_M	E_S
6	−0.008966	0.004428	−0.000469
12	−0.002269	0.001131	−0.000036

Observations:

1. E_T and E_M are opposite in sign and decrease by a factor of about 4 as n is doubled.
2. The Simpson's approximation is much more accurate than the Midpoint and Trapezoidal approximations, and E_S seems to decrease by a factor of about 16 as n is doubled.

29. (a) $\Delta x = (b - a)/n = (6 - 0)/6 = 1$

$T_6 = \frac{1}{2}[f(0) + 2f(1) + 2f(2) + 2f(3) + 2f(4) + 2f(5) + f(6)]$

$\approx \frac{1}{2}[2 + 2(1) + 2(3) + 2(5) + 2(4) + 2(3) + 4] = \frac{1}{2}(38) = 19$

(b) $M_6 = 1[f(0.5) + f(1.5) + f(2.5) + f(3.5) + f(4.5) + f(5.5)] \approx 1.3 + 1.5 + 4.6 + 4.7 + 3.3 + 3.2 = 18.6$

(c) $S_6 = \frac{1}{3}[f(0) + 4f(1) + 2f(2) + 4f(3) + 2f(4) + 4f(5) + f(6)]$

$\approx \frac{1}{3}[2 + 4(1) + 2(3) + 4(5) + 2(4) + 4(3) + 4] = \frac{1}{3}(56) = 18.\overline{6}$

30. If $x =$ distance from left end of pool and $w = w(x) =$ width at x, then Simpson's Rule with $n = 8$ and $\Delta x = 2$ gives

Area $= \int_0^{16} w\,dx \approx \frac{2}{3}[0 + 4(6.2) + 2(7.2) + 4(6.8) + 2(5.6) + 4(5.0) + 2(4.8) + 4(4.8) + 0] \approx 84\ \text{m}^2$.

31. (a) $\int_1^5 f(x)\,dx \approx M_4 = \frac{5-1}{4}[f(1.5) + f(2.5) + f(3.5) + f(4.5)] = 1(2.9 + 3.6 + 4.0 + 3.9) = 14.4$

(b) $-2 \le f''(x) \le 3 \quad \Rightarrow \quad |f''(x)| \le 3 \quad \Rightarrow \quad K = 3$, since $|f''(x)| \le K$. The error estimate for the Midpoint Rule is

$$|E_M| \le \frac{K(b-a)^3}{24n^2} = \frac{3(5-1)^3}{24(4)^2} = \frac{1}{2}.$$

32. (a) $\int_0^{1.6} g(x)\,dx \approx S_8 = \frac{1.6-0}{8 \cdot 3}[g(0) + 4g(0.2) + 2g(0.4) + 4g(0.6) + 2g(0.8) + 4g(1.0) + 2g(1.2) + 4g(1.4) + g(1.6)]$

$= \frac{1}{15}[12.1 + 4(11.6) + 2(11.3) + 4(11.1) + 2(11.7) + 4(12.2) + 2(12.6) + 4(13.0) + 13.2]$

$= \frac{1}{15}(288.1) = \frac{2881}{150} \approx 19.2$

(b) $-5 \le g^{(4)}(x) \le 2 \quad \Rightarrow \quad \left|g^{(4)}(x)\right| \le 5 \quad \Rightarrow \quad K = 5$, since $\left|g^{(4)}(x)\right| \le K$. The error estimate for Simpson's Rule is

$$|E_S| \le \frac{K(b-a)^5}{180n^4} = \frac{5(1.6-0)^5}{180(8)^4} = \frac{2}{28{,}125} = 7.\overline{1} \times 10^{-5}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. We use Simpson's Rule with $n = 12$ and $\Delta t = \frac{24-0}{12} = 2$.

$$\begin{aligned} S_{12} &= \tfrac{2}{3}[\,T(0) + 4T(2) + 2T(4) + 4T(6) + 2T(8) + 4T(10) + 2T(12) \\ &\qquad + 4T(14) + 2T(16) + 4T(18) + 2T(20) + 4T(22) + T(24)] \\ &\approx \tfrac{2}{3}[\,66.6 + 4(65.4) + 2(64.4) + 4(61.7) + 2(67.3) + 4(72.1) + 2(74.9) \\ &\qquad + 4(77.4) + 2(79.1) + 4(75.4) + 2(75.6) + 4(71.4) + 67.5] = \tfrac{2}{3}(2550.3) = 1700.2. \end{aligned}$$

Thus, $\int_0^{24} T(t)\,dt \approx S_{12}$ and $T_{\text{ave}} = \frac{1}{24-0}\int_0^{24} T(t)\,dt \approx 70.84°\text{F}$.

34. We use Simpson's Rule with $n = 10$ and $\Delta x = \frac{1}{2}$:

$$\begin{aligned} \text{distance} = \int_0^5 v(t)\,dt \approx S_{10} &= \tfrac{1}{2\cdot 3}[f(0) + 4f(0.5) + 2f(1) + \cdots + 4f(4.5) + f(5)] \\ &= \tfrac{1}{6}[0 + 4(4.67) + 2(7.34) + 4(8.86) + 2(9.73) + 4(10.22) \\ &\qquad + 2(10.51) + 4(10.67) + 2(10.76) + 4(10.81) + 10.81] \\ &= \tfrac{1}{6}(268.41) = 44.735 \text{ m} \end{aligned}$$

35. By the Net Change Theorem, the increase in velocity is equal to $\int_0^6 a(t)\,dt$. We use Simpson's Rule with $n = 6$ and $\Delta t = (6-0)/6 = 1$ to estimate this integral:

$$\begin{aligned} \int_0^6 a(t)\,dt \approx S_6 &= \tfrac{1}{3}[a(0) + 4a(1) + 2a(2) + 4a(3) + 2a(4) + 4a(5) + a(6)] \\ &\approx \tfrac{1}{3}[0 + 4(0.5) + 2(4.1) + 4(9.8) + 2(12.9) + 4(9.5) + 0] = \tfrac{1}{3}(113.2) = 37.7\overline{3} \text{ ft/s} \end{aligned}$$

36. By the Net Change Theorem, the total amount of water that leaked out during the first six hours is equal to $\int_0^6 r(t)\,dt$. We use Simpson's Rule with $n = 6$ and $\Delta t = \frac{6-0}{6} = 1$ to estimate this integral:

$$\begin{aligned} \int_0^6 r(t)\,dt \approx S_6 &= \tfrac{1}{3}[r(0) + 4r(1) + 2r(2) + 4r(3) + 2r(4) + 4r(5) + r(6)] \\ &\approx \tfrac{1}{3}[4 + 4(3) + 2(2.4) + 4(1.9) + 2(1.4) + 4(1.1) + 1] = \tfrac{1}{3}(36.6) = 12.2 \text{ liters} \end{aligned}$$

37. By the Net Change Theorem, the energy used is equal to $\int_0^6 P(t)\,dt$. We use Simpson's Rule with $n = 12$ and $\Delta t = \frac{6-0}{12} = \frac{1}{2}$ to estimate this integral:

$$\begin{aligned} \int_0^6 P(t)\,dt \approx S_{12} &= \tfrac{1/2}{3}[P(0) + 4P(0.5) + 2P(1) + 4P(1.5) + 2P(2) + 4P(2.5) + 2P(3) \\ &\qquad + 4P(3.5) + 2P(4) + 4P(4.5) + 2P(5) + 4P(5.5) + P(6)] \\ &= \tfrac{1}{6}[1814 + 4(1735) + 2(1686) + 4(1646) + 2(1637) + 4(1609) + 2(1604) \\ &\qquad + 4(1611) + 2(1621) + 4(1666) + 2(1745) + 4(1886) + 2052] \\ &= \tfrac{1}{6}(61{,}064) = 10{,}177.\overline{3} \text{ megawatt-hours} \end{aligned}$$

38. By the Net Change Theorem, the total amount of data transmitted is equal to $\int_0^8 D(t)\,dt \times 3600$ [since $D(t)$ is measured in megabits per second and t is in hours]. We use Simpson's Rule with $n = 8$ and $\Delta t = (8-0)/8 = 1$ to estimate this integral:

$$\begin{aligned} \int_0^8 D(t)\,dt \approx S_8 &= \tfrac{1}{3}[D(0) + 4D(1) + 2D(2) + 4D(3) + 2D(4) + 4D(5) + 2D(6) + 4D(7) + D(8)] \\ &\approx \tfrac{1}{3}[0.35 + 4(0.32) + 2(0.41) + 4(0.50) + 2(0.51) + 4(0.56) + 2(0.56) + 4(0.83) + 0.88] \\ &= \tfrac{1}{3}(13.03) = 4.34\overline{3} \end{aligned}$$

Now multiply by 3600 to obtain 15,636 megabits.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. (a) Let $y = f(x)$ denote the curve. Using disks, $V = \int_2^{10} \pi[f(x)]^2\,dx = \pi \int_2^{10} g(x)\,dx = \pi I_1$.

Now use Simpson's Rule to approximate I_1:

$$\begin{aligned} I_1 \approx S_8 &= \tfrac{10-2}{3(8)}[g(2) + 4g(3) + 2g(4) + 4g(5) + 2g(6) + 4g(7) + g(8)] \\ &\approx \tfrac{1}{3}[0^2 + 4(1.5)^2 + 2(1.9)^2 + 4(2.2)^2 + 2(3.0)^2 + 4(3.8)^2 + 2(4.0)^2 + 4(3.1)^2 + 0^2] \\ &= \tfrac{1}{3}(181.78) \end{aligned}$$

Thus, $V \approx \pi \cdot \frac{1}{3}(181.78) \approx 190.4$ or 190 cubic units.

(b) Using cylindrical shells, $V = \int_2^{10} 2\pi x f(x)\,dx = 2\pi \int_2^{10} x f(x)\,dx = 2\pi I_1$.

Now use Simpson's Rule to approximate I_1:

$$\begin{aligned} I_1 \approx S_8 &= \tfrac{10-2}{3(8)}[2f(2) + 4\cdot 3f(3) + 2\cdot 4f(4) + 4\cdot 5f(5) + 2\cdot 6f(6) \\ &\qquad + 4\cdot 7f(7) + 2\cdot 8f(8) + 4\cdot 9f(9) + 10f(10)] \\ &\approx \tfrac{1}{3}[2(0) + 12(1.5) + 8(1.9) + 20(2.2) + 12(3.0) + 28(3.8) + 16(4.0) + 36(3.1) + 10(0)] \\ &= \tfrac{1}{3}(395.2) \end{aligned}$$

Thus, $V \approx 2\pi \cdot \frac{1}{3}(395.2) \approx 827.7$ or 828 cubic units.

40. $$\begin{aligned} \text{Work} = \int_0^{18} f(x)\,dx \approx S_6 &= \tfrac{18-0}{6\cdot 3}\,[f(0) + 4f(3) + 2f(6) + 4f(9) + 2f(12) + 4f(15) + f(18)] \\ &= 1\cdot[9.8 + 4(9.1) + 2(8.5) + 4(8.0) + 2(7.7) + 4(7.5) + 7.4] = 148 \text{ joules} \end{aligned}$$

41. The curve is $y = f(x) = 1/(1 + e^{-x})$. Using disks, $V = \int_0^{10} \pi[f(x)]^2\,dx = \pi \int_0^{10} g(x)\,dx = \pi I_1$. Now use Simpson's Rule to approximate I_1:

$$\begin{aligned} I_1 \approx S_{10} &= \tfrac{10-0}{10\cdot 3}[g(0) + 4g(1) + 2g(2) + 4g(3) + 2g(4) + 4g(5) + 2g(6) + 4g(7) + 2g(8) + 4g(9) + g(10)] \\ &\approx 8.80825 \end{aligned}$$

Thus, $V \approx \pi I_1 \approx 27.7$ or 28 cubic units.

42. Using Simpson's Rule with $n = 10$, $\Delta x = \frac{\pi/2}{10}$, $L = 1$, $\theta_0 = \frac{42\pi}{180}$ radians, $g = 9.8\ \text{m/s}^2$, $k^2 = \sin^2\left(\frac{1}{2}\theta_0\right)$, and $f(x) = 1/\sqrt{1 - k^2\sin^2 x}$, we get

$$\begin{aligned} T &= 4\sqrt{\frac{L}{g}}\int_0^{\pi/2} \frac{dx}{\sqrt{1 - k^2\sin^2 x}} \approx 4\sqrt{\frac{L}{g}}\,S_{10} \\ &= 4\sqrt{\tfrac{1}{9.8}}\left(\tfrac{\pi/2}{10\cdot 3}\right)\left[f(0) + 4f\left(\tfrac{\pi}{20}\right) + 2f\left(\tfrac{2\pi}{20}\right) + \cdots + 4f\left(\tfrac{9\pi}{20}\right) + f\left(\tfrac{\pi}{2}\right)\right] \approx 2.07665 \end{aligned}$$

43. $I(\theta) = \dfrac{N^2\sin^2 k}{k^2}$, where $k = \dfrac{\pi N d\sin\theta}{\lambda}$, $N = 10{,}000$, $d = 10^{-4}$, and $\lambda = 632.8 \times 10^{-9}$. So $I(\theta) = \dfrac{(10^4)^2\sin^2 k}{k^2}$, where $k = \dfrac{\pi(10^4)(10^{-4})\,\sin\theta}{632.8 \times 10^{-9}}$. Now $n = 10$ and $\Delta\theta = \dfrac{10^{-6} - (-10^{-6})}{10} = 2 \times 10^{-7}$, so

$M_{10} = 2 \times 10^{-7}[I(-0.0000009) + I(-0.0000007) + \cdots + I(0.0000009)] \approx 59.4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. $f(x) = \cos(\pi x)$, $\Delta x = \frac{20-0}{10} = 2 \quad \Rightarrow$

$$T_{10} = \tfrac{2}{2}\{f(0) + 2[f(2) + f(4) + \cdots + f(18)] + f(20)\} = 1[\cos 0 + 2(\cos 2\pi + \cos 4\pi + \cdots + \cos 18\pi) + \cos 20\pi]$$
$$= 1 + 2(1+1+1+1+1+1+1+1+1+1) + 1 = 20$$

The actual value is $\int_0^{20} \cos(\pi x)\,dx = \frac{1}{\pi}\big[\sin \pi x\big]_0^{20} = \frac{1}{\pi}(\sin 20\pi - \sin 0) = 0$. The discrepancy is due to the fact that the function is sampled only at points of the form $2n$, where its value is $f(2n) = \cos(2n\pi) = 1$.

45. Consider the function f whose graph is shown. The area $\int_0^2 f(x)\,dx$ is close to 2. The Trapezoidal Rule gives $T_2 = \frac{2-0}{2\cdot 2}[f(0) + 2f(1) + f(2)] = \frac{1}{2}[1 + 2\cdot 1 + 1] = 2$. The Midpoint Rule gives $M_2 = \frac{2-0}{2}[f(0.5) + f(1.5)] = 1[0+0] = 0$, so the Trapezoidal Rule is more accurate.

46. Consider the function $f(x) = |x-1|$, $0 \le x \le 2$. The area $\int_0^2 f(x)\,dx$ is exactly 1. So is the right endpoint approximation: $R_2 = f(1)\,\Delta x + f(2)\,\Delta x = 0\cdot 1 + 1\cdot 1 = 1$. But Simpson's Rule approximates f with the parabola $y = (x-1)^2$, shown dashed, and

$$S_2 = \frac{\Delta x}{3}[f(0) + 4f(1) + f(2)] = \frac{1}{3}[1 + 4\cdot 0 + 1] = \frac{2}{3}.$$

47. Since the Trapezoidal and Midpoint approximations on the interval $[a, b]$ are the sums of the Trapezoidal and Midpoint approximations on the subintervals $[x_{i-1}, x_i]$, $i = 1, 2, \ldots, n$, we can focus our attention on one such interval. The condition $f''(x) < 0$ for $a \le x \le b$ means that the graph of f is concave down as in Figure 5. In that figure, T_n is the area of the trapezoid $AQRD$, $\int_a^b f(x)\,dx$ is the area of the region $AQPRD$, and M_n is the area of the trapezoid $ABCD$, so $T_n < \int_a^b f(x)\,dx < M_n$. In general, the condition $f'' < 0$ implies that the graph of f on $[a, b]$ lies above the chord joining the points $(a, f(a))$ and $(b, f(b))$. Thus, $\int_a^b f(x)\,dx > T_n$. Since M_n is the area under a tangent to the graph, and since $f'' < 0$ implies that the tangent lies above the graph, we also have $M_n > \int_a^b f(x)\,dx$. Thus, $T_n < \int_a^b f(x)\,dx < M_n$.

48. (a) Let f be a polynomial of degree ≤ 3; say $f(x) = Ax^3 + Bx^2 + Cx + D$. It will suffice to show that Simpson's estimate is exact when there are two subintervals ($n = 2$), because for a larger even number of subintervals the sum of exact estimates is exact. As in the derivation of Simpson's Rule, we can assume that $x_0 = -h$, $x_1 = 0$, and $x_2 = h$. Then Simpson's approximation is

$$\int_{-h}^{h} f(x)\,dx \approx \tfrac{1}{3}\cdot\tfrac{2h}{2}[f(-h) + 4f(0) + f(h)] = \tfrac{1}{3}h\big[\big(-Ah^3 + Bh^2 - Ch + D\big) + 4D + \big(Ah^3 + Bh^2 + Ch + D\big)\big]$$
$$= \tfrac{1}{3}h[2Bh^2 + 6D] = \tfrac{2}{3}Bh^3 + 2Dh$$

The exact value of the integral is

$$\int_{-h}^{h}(Ax^3 + Bx^2 + Cx + D)\,dx = 2\int_0^h (Bx^2 + D)\,dx \qquad \text{[by Theorem 5.5.7]}$$
$$= 2\big[\tfrac{1}{3}Bx^3 + Dx\big]_0^h = \tfrac{2}{3}Bh^3 + 2Dh$$

Thus, Simpson's Rule is exact.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Using Simpson's Rule with $n = 4$, $\Delta x = \frac{b-a}{n} = \frac{8-0}{4} = 2$, and $f(x) = x^3 - 6x^2 + 4x$, we get

$S_4 = \frac{2}{3}[f(0) + 4f(2) + 2f(4) + 4f(6) + f(8)] = \frac{2}{3}(192) = 128$. The exact value of the integral is

$\int_0^8 (x^3 - 6x^2 + 4x)\,dx = \left[\frac{1}{4}x^4 - 2x^3 + 2x^2\right]_0^8 = (1024 - 1024 + 128) - 0 = 128$.

Thus, $S_4 = \int_0^8 (x^3 - 6x^2 + 4x)\,dx$.

(c) $f(x) = Ax^3 + Bx^2 + Cx + D \;\Rightarrow\; f'(x) = 3Ax^2 + 2Bx + C \;\Rightarrow\; f''(x) = 6Ax + 2B \;\Rightarrow\; f'''(x) = 6A \;\Rightarrow$

$f^{(4)}(x) = 0$. Since $\left|f^{(4)}(x)\right| = 0$ for all x, the error bound in (4) gives $|E_S| \le \dfrac{(0)(b-a)^5}{180n^4} = 0$, indicating the error in using Simpson's Rule is zero. Hence, Simpson's Rule gives the exact value of the integral for a polynomial of degree 3 or lower.

49. $T_n = \frac{1}{2}\,\Delta x\,[f(x_0) + 2f(x_1) + \cdots + 2f(x_{n-1}) + f(x_n)]$ and

$M_n = \Delta x\,[f(\overline{x}_1) + f(\overline{x}_2) + \cdots + f(\overline{x}_{n-1}) + f(\overline{x}_n)]$, where $\overline{x}_i = \frac{1}{2}(x_{i-1} + x_i)$. Now

$T_{2n} = \frac{1}{2}\left(\frac{1}{2}\Delta x\right)[f(x_0) + 2f(\overline{x}_1) + 2f(x_1) + 2f(\overline{x}_2) + 2f(x_2) + \cdots + 2f(\overline{x}_{n-1}) + 2f(x_{n-1}) + 2f(\overline{x}_n) + f(x_n)]$, so

$$\begin{aligned}
\tfrac{1}{2}(T_n + M_n) &= \tfrac{1}{2}T_n + \tfrac{1}{2}M_n \\
&= \tfrac{1}{4}\Delta x[f(x_0) + 2f(x_1) + \cdots + 2f(x_{n-1}) + f(x_n)] \\
&\qquad + \tfrac{1}{4}\Delta x[2f(\overline{x}_1) + 2f(\overline{x}_2) + \cdots + 2f(\overline{x}_{n-1}) + 2f(\overline{x}_n)] \\
&= T_{2n}
\end{aligned}$$

50. $T_n = \dfrac{\Delta x}{2}\left[f(x_0) + 2\sum_{i=1}^{n-1} f(x_i) + f(x_n)\right]$ and $M_n = \Delta x \sum_{i=1}^{n} f\left(x_i - \dfrac{\Delta x}{2}\right)$, so

$$\tfrac{1}{3}T_n + \tfrac{2}{3}M_n = \tfrac{1}{3}(T_n + 2M_n) = \frac{\Delta x}{3 \cdot 2}\left[f(x_0) + 2\sum_{i=1}^{n-1} f(x_i) + f(x_n) + 4\sum_{i=1}^{n} f\left(x_i - \frac{\Delta x}{2}\right)\right]$$

where $\Delta x = \dfrac{b-a}{n}$. Let $\delta x = \dfrac{b-a}{2n}$. Then $\Delta x = 2\delta x$, so

$$\begin{aligned}
\tfrac{1}{3}T_n + \tfrac{2}{3}M_n &= \frac{\delta x}{3}\left[f(x_0) + 2\sum_{i=1}^{n-1} f(x_i) + f(x_n) + 4\sum_{i=1}^{n} f(x_i - \delta x)\right] \\
&= \tfrac{1}{3}\delta x[f(x_0) + 4f(x_1 - \delta x) + 2f(x_1) + 4f(x_2 - \delta x) \\
&\qquad + 2f(x_2) + \cdots + 2f(x_{n-1}) + 4f(x_n - \delta x) + f(x_n)]
\end{aligned}$$

Since $x_0, x_1 - \delta x, x_1, x_2 - \delta x, x_2, \ldots, x_{n-1}, x_n - \delta x, x_n$ are the subinterval endpoints for S_{2n}, and since $\delta x = \dfrac{b-a}{2n}$ is the width of the subintervals for S_{2n}, the last expression for $\frac{1}{3}T_n + \frac{2}{3}M_n$ is the usual expression for S_{2n}. Therefore, $\frac{1}{3}T_n + \frac{2}{3}M_n = S_{2n}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7.8 Improper Integrals

1. (a) Since $y = \dfrac{1}{x-3}$ has an infinite discontinuity at $x = 3$, $\displaystyle\int_1^4 \frac{dx}{x-3}$ is a Type 2 improper integral.

(b) Since $\displaystyle\int_3^\infty \frac{dx}{x^2-4}$ has an infinite interval of integration, it is an improper integral of Type 1.

(c) Since $y = \tan \pi x$ has an infinite discontinuity at $x = \frac{1}{2}$, $\int_0^1 \tan \pi x \, dx$ is a Type 2 improper integral.

(d) Since $\displaystyle\int_{-\infty}^{-1} \frac{e^x}{x}\, dx$ has an infinite interval of integration, it is an improper integral of Type 1.

2. (a) Since $y = \sec x$ has an infinite discontinuity at $x = \frac{\pi}{2}$, $\int_0^\pi \sec x \, dx$ is a Type 2 improper integral.

(b) Since $y = \dfrac{1}{x-5}$ is defined and continuous on the interval $[0, 4]$, $\displaystyle\int_0^4 \frac{dx}{x-5}$ is a proper integral.

(c) Since $y = \dfrac{1}{x+x^3} = \dfrac{1}{x\left(1+x^2\right)}$ has an infinite discontinuity at $x = 0$, $\displaystyle\int_{-1}^3 \frac{1}{x+x^3}\, dx$ is a Type 2 improper integral.

(d) Since $\displaystyle\int_1^\infty \frac{1}{x+x^3}\, dx$ has an infinite interval of integration, it is an improper integral of Type 1.

3. The area under the graph of $y = 1/x^3 = x^{-3}$ between $x = 1$ and $x = t$ is

$A(t) = \int_1^t x^{-3}\, dx = \left[-\frac{1}{2}x^{-2}\right]_1^t = -\frac{1}{2}t^{-2} - \left(-\frac{1}{2}\right) = \frac{1}{2} - 1/\left(2t^2\right)$. So the area for $1 \le x \le 10$ is $A(10) = 0.5 - 0.005 = 0.495$, the area for $1 \le x \le 100$ is $A(100) = 0.5 - 0.00005 = 0.49995$, and the area for $1 \le x \le 1000$ is $A(1000) = 0.5 - 0.0000005 = 0.4999995$. The total area under the curve for $x \ge 1$ is $\displaystyle\lim_{t\to\infty} A(t) = \lim_{t\to\infty} \left[\frac{1}{2} - 1/(2t^2)\right] = \frac{1}{2}$.

4. (a)

(b) The area under the graph of f from $x = 1$ to $x = t$ is

$$F(t) = \int_1^t f(x)\, dx = \int_1^t x^{-1.1}\, dx = \left[-\frac{1}{0.1}x^{-0.1}\right]_1^t$$
$$= -10(t^{-0.1} - 1) = 10(1 - t^{-0.1})$$

and the area under the graph of g is

$$G(t) = \int_1^t g(x)\, dx = \int_1^t x^{-0.9}\, dx = \left[\frac{1}{0.1}x^{0.1}\right]_1^t = 10(t^{0.1} - 1).$$

t	$F(t)$	$G(t)$
10	2.06	2.59
100	3.69	5.85
10^4	6.02	15.12
10^6	7.49	29.81
10^{10}	9	90
10^{20}	9.9	990

(c) The total area under the graph of f is $\displaystyle\lim_{t\to\infty} F(t) = \lim_{t\to\infty} 10(1 - t^{-0.1}) = 10$.

The total area under the graph of g does not exist, since $\displaystyle\lim_{t\to\infty} G(t) = \lim_{t\to\infty} 10(t^{0.1} - 1) = \infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $\int_1^\infty 2x^{-3}\,dx = \lim\limits_{t\to\infty}\int_1^t 2x^{-3}\,dx = \lim\limits_{t\to\infty}\left[-x^{-2}\right]_1^t = \lim\limits_{t\to\infty}\left[-t^{-2}+1\right] = 0+1 = 1.$ Convergent

6. $\displaystyle\int_{-\infty}^{-1}\frac{1}{\sqrt[3]{x}}\,dx = \lim_{t\to-\infty}\int_t^{-1} x^{-1/3}\,dx = \lim_{t\to-\infty}\left[\tfrac{3}{2}x^{2/3}\right]_t^{-1} = \lim_{t\to-\infty}\left[\tfrac{3}{2}-\tfrac{3}{2}t^{2/3}\right] = -\infty.$ Divergent

7. $\int_0^\infty e^{-2x}\,dx = \lim\limits_{t\to\infty}\int_0^t e^{-2x}\,dx = \lim\limits_{t\to\infty}\left[-\tfrac{1}{2}e^{-2x}\right]_0^t = \lim\limits_{t\to\infty}\left[-\tfrac{1}{2}e^{-2t}+\tfrac{1}{2}\right] = 0+\tfrac{1}{2} = \tfrac{1}{2}.$ Convergent

8. $\displaystyle\int_1^\infty\left(\frac{1}{3}\right)^x dx = \lim_{t\to\infty}\int_1^t 3^{-x}\,dx = \lim_{t\to\infty}\left[-\frac{3^{-x}}{\ln 3}\right]_1^t = \lim_{t\to\infty}\left[-\frac{3^{-t}}{\ln 3}+\frac{3^{-1}}{\ln 3}\right] = 0+\frac{1}{3\ln 3} = \frac{1}{\ln 27}.$ Convergent

9. $\displaystyle\int_{-2}^\infty\frac{1}{x+4}\,dx = \lim_{t\to\infty}\int_{-2}^t\frac{1}{x+4}\,dx = \lim_{t\to\infty}\Big[\ln|x+4|\,\Big]_{-2}^t = \lim_{t\to\infty}\Big[\ln|t+4|-\ln 2\Big] = \infty$ since $\lim\limits_{t\to\infty}\ln|x+4| = \infty.$

Divergent

10. $\displaystyle\int_1^\infty\frac{1}{x^2+4}\,dx = \lim_{t\to\infty}\int_1^t\frac{1}{x^2+4}\,dx = \lim_{t\to\infty}\left[\frac{1}{2}\tan^{-1}\left(\frac{x}{2}\right)\right]_1^t = \lim_{t\to\infty}\left[\frac{1}{2}\tan^{-1}\left(\frac{t}{2}\right)-\frac{1}{2}\tan^{-1}\left(\frac{1}{2}\right)\right]$

$\displaystyle = \frac{\pi}{4}-\frac{1}{2}\tan^{-1}\left(\frac{1}{2}\right).$ Convergent

11. $\displaystyle\int_3^\infty\frac{1}{(x-2)^{3/2}}\,dx = \lim_{t\to\infty}\int_3^t(x-2)^{-3/2}\,dx = \lim_{t\to\infty}\Big[-2\,(x-2)^{-1/2}\Big]_3^t$ $[u = x-2,\, du = dx]$

$\displaystyle = \lim_{t\to\infty}\left(\frac{-2}{\sqrt{t-2}}+\frac{2}{\sqrt{1}}\right) = 0+2 = 2.$ Convergent

12. $\displaystyle\int_0^\infty\frac{1}{\sqrt[4]{1+x}}\,dx = \lim_{t\to\infty}\int_0^t(1+x)^{-1/4}\,dx = \lim_{t\to\infty}\left[\tfrac{4}{3}(1+x)^{3/4}\right]_0^t$ $[u = 1+x,\, du = dx]$

$\displaystyle = \lim_{t\to\infty}\left[\tfrac{4}{3}(1+t)^{3/4}-\tfrac{4}{3}\right] = \infty.$ Divergent

13. $\displaystyle\int_{-\infty}^0\frac{x}{(x^2+1)^3}\,dx = \lim_{t\to-\infty}\int_t^0\frac{x}{(x^2+1)^3}\,dx = \lim_{t\to-\infty}\left[-\frac{1}{4}(x^2+1)^{-2}\right]_t^0 = \lim_{t\to-\infty}\left[-\frac{1}{4\,(x^2+1)^2}\right]_t^0$

$\displaystyle = \lim_{t\to-\infty}\left[-\frac{1}{4}+\frac{1}{4(t^2+1)^2}\right] = -\frac{1}{4}+0 = -\frac{1}{4}.$ Convergent

14. $\displaystyle\int_{-\infty}^{-3}\frac{x}{4-x^2}\,dx = \lim_{t\to-\infty}\int_t^{-3}\frac{x}{4-x^2}\,dx = \lim_{t\to-\infty}\left[-\frac{1}{2}\ln\left|4-x^2\right|\right]_t^{-3}$

$= \lim\limits_{t\to-\infty}\left[-\tfrac{1}{2}\ln 5+\tfrac{1}{2}\ln\left|4-t^2\right|\right] = \infty$ since $\lim\limits_{t\to-\infty}\ln\left|4-t^2\right| = \infty.$ Divergent

15. $\displaystyle\int_1^\infty\frac{x^2+x+1}{x^4}\,dx = \lim_{t\to\infty}\int_1^t(x^{-2}+x^{-3}+x^{-4})\,dx$

$\displaystyle = \lim_{t\to\infty}\left[-x^{-1}-\frac{1}{2}x^{-2}-\frac{1}{3}x^{-3}\right]_1^t = \lim_{t\to\infty}\left[-\frac{1}{x}-\frac{1}{2x^2}-\frac{1}{3x^3}\right]_1^t$

$\displaystyle = \lim_{t\to\infty}\left[\left(-\frac{1}{t}-\frac{1}{2t^2}-\frac{1}{3t^3}\right)-\left(-1-\frac{1}{2}-\frac{1}{3}\right)\right] = 0+\frac{11}{6} = \frac{11}{6}.$ Convergent

16. $\displaystyle\int_2^\infty\frac{x}{\sqrt{x^2-1}}\,dx = \lim_{t\to\infty}\int_2^t\frac{x}{\sqrt{x^2-1}}\,dx = \lim_{t\to\infty}\Big[\sqrt{x^2-1}\,\Big]_2^t = \lim_{t\to\infty}\left[\sqrt{t^2-1}-\sqrt{3}\right] = \infty.$ Divergent

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\displaystyle\int_0^\infty \frac{e^x}{(1+e^x)^2}\,dx = \lim_{t\to\infty}\int_0^t \frac{e^x}{(1+e^x)^2}\,dx = \lim_{t\to\infty}\Big[-(1+e^x)^{-1}\Big]_0^t = \lim_{t\to\infty}\left[-\frac{1}{1+e^x}\right]_0^t$

$\displaystyle = \lim_{t\to\infty}\left[-\frac{1}{1+e^t}+\frac{1}{2}\right] = 0+\frac{1}{2} = \frac{1}{2}.$ Convergent

18. $\displaystyle I = \int_{-\infty}^{-1}\frac{x^2+x}{x^3}\,dx = \int_{-\infty}^{-1}\frac{1}{x}\,dx + \int_{-\infty}^{-1}\frac{1}{x^2}\,dx = I_1 + I_2.$

Now, $\displaystyle\int_{-\infty}^{-1}\frac{1}{x}\,dx = \lim_{t\to-\infty}\int_t^{-1}\frac{1}{x}\,dx = \lim_{t\to-\infty}\Big[\ln|x|\Big]_t^{-1} = \lim_{t\to-\infty}\Big[\ln 1 - \ln|t|\Big] = -\infty.$

Since I_1 is divergent, I is divergent. Divergent

19. $\int_{-\infty}^{\infty} xe^{-x^2}\,dx = \int_{-\infty}^{0} xe^{-x^2}\,dx + \int_0^\infty xe^{-x^2}\,dx.$

$\int_{-\infty}^{0} xe^{-x^2}\,dx = \lim_{t\to-\infty}\left(-\frac{1}{2}\right)\Big[e^{-x^2}\Big]_t^0 = \lim_{t\to-\infty}\left(-\frac{1}{2}\right)\Big(1-e^{-t^2}\Big) = -\frac{1}{2}\cdot(1-0) = -\frac{1}{2}$, and

$\int_0^\infty xe^{-x^2}\,dx = \lim_{t\to\infty}\left(-\frac{1}{2}\right)\Big[e^{-x^2}\Big]_0^t = \lim_{t\to\infty}\left(-\frac{1}{2}\right)\Big(e^{-t^2}-1\Big) = -\frac{1}{2}\cdot(0-1) = \frac{1}{2}.$

Therefore, $\int_{-\infty}^{\infty} xe^{-x^2}\,dx = -\frac{1}{2}+\frac{1}{2} = 0.$ Convergent

20. $\displaystyle I = \int_{-\infty}^{\infty}\frac{x}{x^2+1}\,dx = \int_{-\infty}^{0}\frac{x}{x^2+1}\,dx + \int_0^\infty\frac{x}{x^2+1}\,dx = I_1 + I_2$, but

$\displaystyle I_2 = \lim_{t\to\infty}\int_0^t\frac{x}{x^2+1}\,dx = \lim_{t\to\infty}\int_1^{t^2+1}\frac{1/2}{u}\,du \quad \begin{bmatrix} u = x^2+1, \\ du = 2x\,dx \end{bmatrix} \quad = \frac{1}{2}\lim_{t\to\infty}\Big[\ln|u|\Big]_1^{t^2+1}$

$\displaystyle = \frac{1}{2}\lim_{t\to\infty}\Big[\ln\left|t^2+1\right| - 0\Big] = \infty$

Since I_2 is divergent, I is divergent, and there is no need to evaluate I_1. Divergent

21. $I = \int_{-\infty}^{\infty}\cos 2t\,dt = \int_{-\infty}^{0}\cos 2t\,dt + \int_0^\infty \cos 2t\,dt = I_1 + I_2$, but $I_1 = \displaystyle\lim_{s\to-\infty}\Big[\tfrac{1}{2}\sin 2t\Big]_s^0 = \lim_{s\to-\infty}\left(-\tfrac{1}{2}\sin 2s\right)$, and this limit does not exist. Since I_1 is divergent, I is divergent, and there is no need to evaluate I_2. Divergent

22. $\displaystyle\int_1^\infty\frac{e^{-1/x}}{x^2}\,dx = \lim_{t\to\infty}\int_1^t\frac{e^{-1/x}}{x^2}\,dx = \lim_{t\to\infty}\Big[e^{-1/x}\Big]_1^t = \lim_{t\to\infty}(e^{-1/t}-e^{-1}) = 1-\frac{1}{e}.$ Convergent

23. $\int_0^\infty \sin^2\alpha\,d\alpha = \lim_{t\to\infty}\int_0^t \frac{1}{2}(1-\cos 2\alpha)\,d\alpha = \lim_{t\to\infty}\left[\frac{1}{2}\left(\alpha-\frac{1}{2}\sin 2\alpha\right)\right]_0^t = \lim_{t\to\infty}\left[\frac{1}{2}\left(t-\frac{1}{2}\sin 2t\right)-0\right] = \infty.$

Divergent

24. $\int_0^\infty \sin\theta\, e^{\cos\theta}\,d\theta = \lim_{t\to\infty}\int_0^t \sin\theta\, e^{\cos\theta}\,d\theta = \lim_{t\to\infty}\Big[-e^{\cos\theta}\Big]_0^t = \lim_{t\to\infty}(-e^{\cos t}+e)$

This limit does not exist since $\cos t$ oscillates in value between -1 and 1, so $e^{\cos t}$ oscillates in value between e^{-1} and e^1. Divergent

25. $\displaystyle\int_1^\infty\frac{1}{x^2+x}\,dx = \lim_{t\to\infty}\int_1^t\frac{1}{x(x+1)}\,dx = \lim_{t\to\infty}\int_1^t\left(\frac{1}{x}-\frac{1}{x+1}\right)dx$ [partial fractions]

$\displaystyle = \lim_{t\to\infty}\Big[\ln|x| - \ln|x+1|\Big]_1^t = \lim_{t\to\infty}\left[\ln\left|\frac{x}{x+1}\right|\right]_1^t = \lim_{t\to\infty}\left(\ln\frac{t}{t+1} - \ln\frac{1}{2}\right) = 0 - \ln\frac{1}{2} = \ln 2.$

Convergent

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $\displaystyle\int_2^\infty \frac{dv}{v^2+2v-3} = \lim_{t\to\infty}\int_2^t \frac{dv}{(v+3)(v-1)} = \lim_{t\to\infty}\int_2^t \left(\frac{-\frac{1}{4}}{v+3}+\frac{\frac{1}{4}}{v-1}\right)dv = \lim_{t\to\infty}\left[-\tfrac{1}{4}\ln|v+3|+\tfrac{1}{4}\ln|v-1|\right]_2^t$

$$= \frac{1}{4}\lim_{t\to\infty}\left[\ln\frac{v-1}{v+3}\right]_2^t = \frac{1}{4}\lim_{t\to\infty}\left(\ln\frac{t-1}{t+3}-\ln\frac{1}{5}\right) = \frac{1}{4}(0+\ln 5) = \frac{1}{4}\ln 5. \quad \text{Convergent}$$

27. $\displaystyle\int_{-\infty}^0 ze^{2z}\,dz = \lim_{t\to-\infty}\int_t^0 ze^{2z}\,dz = \lim_{t\to-\infty}\left[\tfrac{1}{2}ze^{2z}-\tfrac{1}{4}e^{2z}\right]_t^0$ $\left[\begin{array}{c}\text{integration by parts with}\\ u=z,\ dv=e^{2z}\,dz\end{array}\right]$

$$= \lim_{t\to-\infty}\left[\left(0-\tfrac{1}{4}\right)-\left(\tfrac{1}{2}te^{2t}-\tfrac{1}{4}e^{2t}\right)\right] = -\tfrac{1}{4}-0+0 \quad \text{[by l'Hospital's Rule]} \quad = -\tfrac{1}{4}. \quad \text{Convergent}$$

28. $\displaystyle\int_2^\infty ye^{-3y}\,dy = \lim_{t\to\infty}\int_2^t ye^{-3y}\,dy = \lim_{t\to\infty}\left[-\tfrac{1}{3}ye^{-3y}-\tfrac{1}{9}e^{-3y}\right]_2^t$ $\left[\begin{array}{c}\text{integration by parts with}\\ u=y,\ dv=e^{-3y}\,dy\end{array}\right]$

$$= \lim_{t\to\infty}\left[\left(-\tfrac{1}{3}te^{-3t}-\tfrac{1}{9}e^{-3t}\right)-\left(-\tfrac{2}{3}e^{-6}-\tfrac{1}{9}e^{-6}\right)\right] = 0-0+\tfrac{7}{9}e^{-6} \quad \text{[by l'Hospital's Rule]} \quad = \tfrac{7}{9}e^{-6}.$$

Convergent

29. $\displaystyle\int_1^\infty \frac{\ln x}{x}\,dx = \lim_{t\to\infty}\left[\frac{(\ln x)^2}{2}\right]_1^t$ $\left[\begin{array}{c}\text{by substitution with}\\ u=\ln x,\ du=dx/x\end{array}\right]$ $\displaystyle = \lim_{t\to\infty}\frac{(\ln t)^2}{2} = \infty.$ Divergent

30. $\displaystyle\int_1^\infty \frac{\ln x}{x^2}\,dx = \lim_{t\to\infty}\int_1^t \frac{\ln x}{x^2}\,dx = \lim_{t\to\infty}\left[-\frac{\ln x}{x}-\frac{1}{x}\right]_1^t$ $\left[\begin{array}{c}\text{integration by parts with}\\ u=\ln x,\ dv=(1/x^2)\,dx\end{array}\right]$

$$= \lim_{t\to\infty}\left(-\frac{\ln t}{t}-\frac{1}{t}+1\right) \overset{\text{H}}{=} \lim_{t\to\infty}\left(-\frac{1/t}{1}\right)-\lim_{t\to\infty}\frac{1}{t}+\lim_{t\to\infty}1 = 0-0+1 = 1. \quad \text{Convergent}$$

31. $\displaystyle\int_{-\infty}^0 \frac{z}{z^4+4}\,dz = \lim_{t\to-\infty}\int_t^0 \frac{z}{z^4+4}\,dz = \lim_{t\to-\infty}\frac{1}{2}\left[\frac{1}{2}\tan^{-1}\left(\frac{z^2}{2}\right)\right]_t^0$ $\left[\begin{array}{c}u=z^2,\\ du=2z\,dz\end{array}\right]$

$$= \lim_{t\to-\infty}\left[0-\frac{1}{4}\tan^{-1}\left(\frac{t^2}{2}\right)\right] = -\frac{1}{4}\left(\frac{\pi}{2}\right) = -\frac{\pi}{8}. \quad \text{Convergent}$$

32. $\displaystyle\int_e^\infty \frac{1}{x(\ln x)^2}\,dx = \lim_{t\to\infty}\int_e^t \frac{1}{x(\ln x)^2}\,dx = \lim_{t\to\infty}\left[-\frac{1}{\ln x}\right]_e^t$ $\left[\begin{array}{c}u=\ln x,\\ du=(1/x)\,dx\end{array}\right]$

$$= \lim_{t\to\infty}\left(-\frac{1}{\ln t}+1\right) = 0+1 = 1. \quad \text{Convergent}$$

33. $\displaystyle\int_0^\infty e^{-\sqrt{y}}\,dy = \lim_{t\to\infty}\int_0^t e^{-\sqrt{y}}\,dy = \lim_{t\to\infty}\int_0^{\sqrt{t}} e^{-x}\,(2x\,dx)$ $\left[\begin{array}{c}x=\sqrt{y},\\ dx=1/(2\sqrt{y}\,)\,dy\end{array}\right]$

$$= \lim_{t\to\infty}\left\{\left[-2xe^{-x}\right]_0^{\sqrt{t}}+\int_0^{\sqrt{t}} 2e^{-x}\,dx\right\} \qquad \left[\begin{array}{ll}u=2x, & dv=e^{-x}\,dx\\ du=2\,dx, & v=-e^{-x}\end{array}\right]$$

$$= \lim_{t\to\infty}\left(-2\sqrt{t}\,e^{-\sqrt{t}}+\left[-2e^{-x}\right]_0^{\sqrt{t}}\right) = \lim_{t\to\infty}\left(\frac{-2\sqrt{t}}{e^{\sqrt{t}}}-\frac{2}{e^{\sqrt{t}}}+2\right) = 0-0+2 = 2.$$

Convergent

Note: $\displaystyle\lim_{t\to\infty}\frac{\sqrt{t}}{e^{\sqrt{t}}} \overset{\text{H}}{=} \lim_{t\to\infty}\frac{2\sqrt{t}}{2\sqrt{t}\,e^{\sqrt{t}}} = \lim_{t\to\infty}\frac{1}{e^{\sqrt{t}}} = 0$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. $\displaystyle\int_1^\infty \frac{dx}{\sqrt{x}+x\sqrt{x}}\,dx = \lim_{t\to\infty}\int_1^t \frac{dx}{\sqrt{x}\,(1+x)} = \lim_{t\to\infty}\int_1^{\sqrt{t}} \frac{1}{u(1+u^2)}\,(2u\,du) \qquad \left[\begin{array}{l} u=\sqrt{x}, \\ du = 1/(2\sqrt{x}\,)\,dx \end{array}\right]$

$\displaystyle = \lim_{t\to\infty}\int_1^{\sqrt{t}} \frac{2}{1+u^2}\,du = \lim_{t\to\infty}\left[2\tan^{-1}u\right]_1^{\sqrt{t}} = \lim_{t\to\infty} 2(\tan^{-1}\sqrt{t}-\tan^{-1}1)$

$= 2(\frac{\pi}{2}-\frac{\pi}{4}) = \frac{\pi}{2}.$ Convergent

35. $\displaystyle\int_0^1 \frac{1}{x}\,dx = \lim_{t\to0^+}\int_t^1 \frac{1}{x}\,dx = \lim_{t\to0^+}\Big[\ln|x|\Big]_t^1 = \lim_{t\to0^+}(-\ln t) = \infty.$ Divergent

36. $\displaystyle\int_0^5 \frac{1}{\sqrt[3]{5-x}}\,dx = \lim_{t\to5^-}\int_0^t (5-x)^{-1/3}\,dx = \lim_{t\to5^-}\left[-\tfrac{3}{2}(5-x)^{2/3}\right]_0^t = \lim_{t\to5^-}\left\{-\tfrac{3}{2}[(5-t)^{2/3}-5^{2/3}]\right\}$

$= \frac{3}{2}5^{2/3}.$ Convergent

37. $\displaystyle\int_{-2}^{14} \frac{dx}{\sqrt[4]{x+2}} = \lim_{t\to-2^+}\int_t^{14}(x+2)^{-1/4}\,dx = \lim_{t\to-2^+}\left[\frac{4}{3}(x+2)^{3/4}\right]_t^{14} = \frac{4}{3}\lim_{t\to-2^+}\left[16^{3/4}-(t+2)^{3/4}\right]$

$= \frac{4}{3}(8-0) = \frac{32}{3}.$ Convergent

38. $\displaystyle\int_{-1}^2 \frac{x}{(x+1)^2}\,dx = \lim_{t\to-1^+}\int_t^2 \frac{x}{(x+1)^2}\,dx = \lim_{t\to-1^+}\int_t^2\left[\frac{1}{x+1}-\frac{1}{(x+1)^2}\right]dx$ [partial fractions]

$\displaystyle = \lim_{t\to-1^+}\left[\ln|x+1|+\frac{1}{x+1}\right]_t^2 = \lim_{t\to-1^+}\left[\ln3+\frac{1}{3}-\left(\ln(t+1)+\frac{1}{t+1}\right)\right] = -\infty.$ Divergent

Note: To justify the last step, $\displaystyle\lim_{t\to-1^+}\left[\ln(t+1)+\frac{1}{t+1}\right] = \lim_{x\to0^+}\left(\ln x+\frac{1}{x}\right) \quad \left[\begin{array}{c}\text{substitute}\\ x \text{ for } t+1\end{array}\right] = \lim_{x\to0^+}\frac{x\ln x+1}{x} = \infty$

since $\displaystyle\lim_{x\to0^+}(x\ln x) = \lim_{x\to0^+}\frac{\ln x}{1/x} \overset{\text{H}}{=} \lim_{x\to0^+}\frac{1/x}{-1/x^2} = \lim_{x\to0^+}(-x) = 0.$

39. $\displaystyle\int_{-2}^3 \frac{1}{x^4}\,dx = \int_{-2}^0 \frac{dx}{x^4}+\int_0^3\frac{dx}{x^4}$, but $\displaystyle\int_{-2}^0\frac{dx}{x^4} = \lim_{t\to0^-}\left[-\frac{x^{-3}}{3}\right]_{-2}^t = \lim_{t\to0^-}\left[-\frac{1}{3t^3}-\frac{1}{24}\right] = \infty.$ Divergent

40. $\displaystyle\int_0^1 \frac{dx}{\sqrt{1-x^2}} = \lim_{t\to1^-}\int_0^t\frac{dx}{\sqrt{1-x^2}} = \lim_{t\to1^-}\left[\sin^{-1}x\right]_0^t = \lim_{t\to1^-}\sin^{-1}t = \frac{\pi}{2}.$ Convergent

41. There is an infinite discontinuity at $x=1$. $\displaystyle\int_0^9 \frac{1}{\sqrt[3]{x-1}}\,dx = \int_0^1(x-1)^{-1/3}\,dx+\int_1^9(x-1)^{-1/3}\,dx.$

Here $\int_0^1(x-1)^{-1/3}\,dx = \lim_{t\to1^-}\int_0^t(x-1)^{-1/3}\,dx = \lim_{t\to1^-}\left[\frac{3}{2}(x-1)^{2/3}\right]_0^t = \lim_{t\to1^-}\left[\frac{3}{2}(t-1)^{2/3}-\frac{3}{2}\right] = -\frac{3}{2}$

and $\int_1^9(x-1)^{-1/3}\,dx = \lim_{t\to1^+}\int_t^9(x-1)^{-1/3}\,dx = \lim_{t\to1^+}\left[\frac{3}{2}(x-1)^{2/3}\right]_t^9 = \lim_{t\to1^+}\left[6-\frac{3}{2}(t-1)^{2/3}\right] = 6$. Thus,

$\displaystyle\int_0^9\frac{1}{\sqrt[3]{x-1}}\,dx = -\frac{3}{2}+6 = \frac{9}{2}.$ Convergent

42. There is an infinite discontinuity at $w=2$.

$\displaystyle\int_0^2\frac{w}{w-2}\,dw = \lim_{t\to2^-}\int_0^t\left(1+\frac{2}{w-2}\right)dw = \lim_{t\to2^-}\Big[w+2\ln|w-2|\Big]_0^t = \lim_{t\to2^-}(t+2\ln|t-2|-2\ln2) = -\infty$, so

$\displaystyle\int_0^2\frac{w}{w-2}\,dw$ diverges, and hence, $\displaystyle\int_0^5\frac{w}{w-2}\,dw$ diverges. Divergent

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. $\int_0^{\pi/2} \tan^2\theta\, d\theta = \lim\limits_{t\to(\pi/2)^-} \int_0^t \tan^2\theta\, d\theta = \lim\limits_{t\to(\pi/2)^-} \int_0^t (\sec^2\theta - 1)\, d\theta = \lim\limits_{t\to(\pi/2)^-} \Big[\tan\theta - \theta\Big]_0^t$

$= \lim\limits_{t\to(\pi/2)^-} (\tan t - t) = \infty$ since $\tan t \to \infty$ as $t \to \frac{\pi}{2}^-$. Divergent

44. $\displaystyle\int_0^4 \frac{dx}{x^2 - x - 2} = \int_0^4 \frac{dx}{(x-2)(x+1)} = \int_0^2 \frac{dx}{(x-2)(x+1)} + \int_2^4 \frac{dx}{(x-2)(x+1)}$

Considering only $\displaystyle\int_0^2 \frac{dx}{(x-2)(x+1)}$ and using partial fractions, we have

$$\int_0^2 \frac{dx}{(x-2)(x+1)} = \lim_{t\to 2^-} \int_0^t \left(\frac{\frac{1}{3}}{x-2} - \frac{\frac{1}{3}}{x+1}\right) dx = \lim_{t\to 2^-} \Big[\tfrac{1}{3}\ln|x-2| - \tfrac{1}{3}\ln|x+1|\Big]_0^t$$

$$= \lim_{t\to 2^-} \big[\tfrac{1}{3}\ln|t-2| - \tfrac{1}{3}\ln|t+1| - \tfrac{1}{3}\ln 2 + 0\big] = -\infty \text{ since } \ln|t-2| \to -\infty \text{ as } t \to 2^-.$$

Thus, $\displaystyle\int_0^2 \frac{dx}{x^2 - x - 2}$ is divergent, and hence, $\displaystyle\int_0^4 \frac{dx}{x^2 - x - 2}$ is divergent as well.

45. $\int_0^1 r\,\ln r\, dr = \lim\limits_{t\to 0^+} \int_t^1 r\,\ln r\, dr = \lim\limits_{t\to 0^+} \Big[\frac{1}{2}r^2\ln r - \frac{1}{4}r^2\Big]_t^1$ $\quad \begin{bmatrix} u = \ln r, & dv = r\,dr \\ du = (1/r)\,dr, & v = \frac{1}{2}r^2 \end{bmatrix}$

$= \lim\limits_{t\to 0^+} \big[(0 - \frac{1}{4}) - (\frac{1}{2}t^2\ln t - \frac{1}{4}t^2)\big] = -\frac{1}{4} - 0 = -\frac{1}{4}$

since $\lim\limits_{t\to 0^+} t^2\ln t = \lim\limits_{t\to 0^+} \dfrac{\ln t}{1/t^2} \overset{\text{H}}{=} \lim\limits_{t\to 0^+} \dfrac{1/t}{-2/t^3} = \lim\limits_{t\to 0^+} (-\frac{1}{2}t^2) = 0.$ Convergent

46. $\displaystyle\int_0^{\pi/2} \frac{\cos\theta}{\sqrt{\sin\theta}}\, d\theta = \lim_{t\to 0^+} \int_t^{\pi/2} \frac{\cos\theta}{\sqrt{\sin\theta}}\, d\theta = \lim_{t\to 0^+} \Big[2\sqrt{\sin\theta}\Big]_t^{\pi/2}$ $\quad \begin{bmatrix} u = \sin\theta, \\ du = \cos\theta\, d\theta \end{bmatrix}$

$= \lim\limits_{t\to 0^+} (2 - 2\sqrt{\sin t}\,) = 2 - 0 = 2.$ Convergent

47. $\displaystyle\int_{-1}^0 \frac{e^{1/x}}{x^3}\, dx = \lim_{t\to 0^-} \int_{-1}^t \frac{1}{x}e^{1/x} \cdot \frac{1}{x^2}\, dx = \lim_{t\to 0^-} \int_{-1}^{1/t} ue^u\,(-du)$ $\quad \begin{bmatrix} u = 1/x, \\ du = -dx/x^2 \end{bmatrix}$

$= \lim\limits_{t\to 0^-} \big[(u-1)e^u\big]_{1/t}^{-1}$ $\begin{bmatrix} \text{use parts} \\ \text{or Formula 96} \end{bmatrix}$ $= \lim\limits_{t\to 0^-} \left[-2e^{-1} - \left(\dfrac{1}{t} - 1\right)e^{1/t}\right]$

$= -\dfrac{2}{e} - \lim\limits_{s\to -\infty} (s-1)e^s$ $\quad [s = 1/t]$ $\quad = -\dfrac{2}{e} - \lim\limits_{s\to -\infty} \dfrac{s-1}{e^{-s}} \overset{\text{H}}{=} -\dfrac{2}{e} - \lim\limits_{s\to -\infty} \dfrac{1}{-e^{-s}}$

$= -\dfrac{2}{e} - 0 = -\dfrac{2}{e}.$ Convergent

48. $\displaystyle\int_0^1 \frac{e^{1/x}}{x^3}\, dx = \lim_{t\to 0^+} \int_t^1 \frac{1}{x}e^{1/x} \cdot \frac{1}{x^2}\, dx = \lim_{t\to 0^+} \int_{1/t}^1 ue^u\,(-du)$ $\quad \begin{bmatrix} u = 1/x, \\ du = -dx/x^2 \end{bmatrix}$

$= \lim\limits_{t\to 0^+} \big[(u-1)e^u\big]_1^{1/t}$ $\begin{bmatrix} \text{use parts} \\ \text{or Formula 96} \end{bmatrix}$ $= \lim\limits_{t\to 0^+} \left[\left(\dfrac{1}{t} - 1\right)e^{1/t} - 0\right]$

$= \lim\limits_{s\to\infty} (s-1)e^s$ $\quad [s = 1/t]$ $\quad = \infty.$ Divergent

49.

$$\text{Area} = \int_1^\infty e^{-x}\, dx = \lim_{t\to\infty} \int_1^t e^{-x}\, dx = \lim_{t\to\infty} \Big[-e^{-x}\Big]_1^t$$

$$= \lim_{t\to\infty} (-e^{-t} + e^{-1}) = 0 + e^{-1} = 1/e$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50.

$$\text{Area} = \int_{-\infty}^{0} e^x\,dx = \lim_{t\to-\infty}\int_t^0 e^x\,dx = \lim_{t\to-\infty}\Big[e^x\Big]_t^0$$
$$= \lim_{t\to-\infty}(e^0 - e^t) = 1 - 0 = 1$$

51.

$$\text{Area} = \int_1^{\infty} \frac{1}{x^3+x}\,dx = \lim_{t\to\infty}\int_1^t \frac{1}{x(x^2+1)}\,dx$$
$$= \lim_{t\to\infty}\int_1^t \left(\frac{1}{x} - \frac{x}{x^2+1}\right)dx \qquad \text{[partial fractions]}$$
$$= \lim_{t\to\infty}\left[\ln|x| - \frac{1}{2}\ln\left|x^2+1\right|\right]_1^t = \lim_{t\to\infty}\left[\ln\frac{x}{\sqrt{x^2+1}}\right]_1^t$$
$$= \lim_{t\to\infty}\left(\ln\frac{t}{\sqrt{t^2+1}} - \ln\frac{1}{\sqrt{2}}\right) = \ln 1 - \ln 2^{-1/2} = \frac{1}{2}\ln 2$$

52.

$$\text{Area} = \int_0^{\infty} xe^{-x}\,dx = \lim_{t\to\infty}\int_0^t xe^{-x}\,dx$$
$$= \lim_{t\to\infty}\Big[-xe^{-x} - e^{-x}\Big]_0^t \qquad \text{[use parts wtih } u = x \text{ and } dv = e^{-x}\,dx\text{]}$$
$$= \lim_{t\to\infty}\left[(-te^{-t} - e^{-t}) - (-1)\right]$$
$$= 0 \quad \text{[use l'Hospital's Rule]} \quad -0 + 1 = 1$$

53.

$$\text{Area} = \int_0^{\pi/2} \sec^2 x\,dx = \lim_{t\to(\pi/2)^-}\int_0^t \sec^2 x\,dx = \lim_{t\to(\pi/2)^-}\Big[\tan x\Big]_0^t$$
$$= \lim_{t\to(\pi/2)^-}(\tan t - 0) = \infty$$

Infinite area

54.

$$\text{Area} = \int_{-2}^{0} \frac{1}{\sqrt{x+2}}\,dx = \lim_{t\to-2^+}\int_t^0 \frac{1}{\sqrt{x+2}}\,dx = \lim_{t\to-2^+}\left[2\sqrt{x+2}\right]_t^0$$
$$= \lim_{t\to-2^+}\left(2\sqrt{2} - 2\sqrt{t+2}\right) = 2\sqrt{2} - 0 = 2\sqrt{2}$$

55. (a)

t	$\int_1^t g(x)\,dx$
2	0.447453
5	0.577101
10	0.621306
100	0.668479
1000	0.672957
10,000	0.673407

$$g(x) = \frac{\sin^2 x}{x^2}.$$

It appears that the integral is convergent.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $-1 \le \sin x \le 1 \quad \Rightarrow \quad 0 \le \sin^2 x \le 1 \quad \Rightarrow \quad 0 \le \dfrac{\sin^2 x}{x^2} \le \dfrac{1}{x^2}$. Since $\displaystyle\int_1^\infty \frac{1}{x^2}\,dx$ is convergent [Theorem 2 with $p = 2 > 1$], $\displaystyle\int_1^\infty \frac{\sin^2 x}{x^2}\,dx$ is convergent by the Comparison Theorem.

(c)

Since $\int_1^\infty f(x)\,dx$ is finite and the area under $g(x)$ is less than the area under $f(x)$ on any interval $[1, t]$, $\int_1^\infty g(x)\,dx$ must be finite; that is, the integral is convergent.

56. (a)

t	$\int_2^t g(x)dx$
5	3.830327
10	6.801200
100	23.328769
1000	69.023361
10,000	208.124560

$g(x) = \dfrac{1}{\sqrt{x} - 1}$.

It appears that the integral is divergent.

(b) For $x \ge 2$, $\sqrt{x} > \sqrt{x} - 1 \quad \Rightarrow \quad \dfrac{1}{\sqrt{x}} < \dfrac{1}{\sqrt{x} - 1}$. Since $\displaystyle\int_2^\infty \frac{1}{\sqrt{x}}\,dx$ is divergent [Theorem 2 with $p = \frac{1}{2} \le 1$], $\displaystyle\int_2^\infty \frac{1}{\sqrt{x} - 1}\,dx$ is divergent by the Comparison Theorem.

(c)

Since $\int_2^\infty f(x)\,dx$ is infinite and the area under $g(x)$ is greater than the area under $f(x)$ on any interval $[2, t]$, $\int_2^\infty g(x)\,dx$ must be infinite; that is, the integral is divergent.

57. For $x > 0$, $\dfrac{x}{x^3 + 1} < \dfrac{x}{x^3} = \dfrac{1}{x^2}$. $\displaystyle\int_1^\infty \frac{1}{x^2}\,dx$ is convergent by Theorem 2 with $p = 2 > 1$, so $\displaystyle\int_1^\infty \frac{x}{x^3 + 1}\,dx$ is convergent by the Comparison Theorem. $\displaystyle\int_0^1 \frac{x}{x^3 + 1}\,dx$ is a constant, so $\displaystyle\int_0^\infty \frac{x}{x^3 + 1}\,dx = \int_0^1 \frac{x}{x^3 + 1}\,dx + \int_1^\infty \frac{x}{x^3 + 1}\,dx$ is also convergent.

58. For $x \ge 1$, $\dfrac{1 + \sin^2 x}{\sqrt{x}} \ge \dfrac{1}{\sqrt{x}}$. $\displaystyle\int_1^\infty \frac{1}{\sqrt{x}}\,dx$ is divergent by Theorem 2 with $p = \frac{1}{2} \le 1$, so $\displaystyle\int_1^\infty \frac{1 + \sin^2 x}{\sqrt{x}}\,dx$ is divergent by the Comparison Theorem.

59. For $x \ge 1$, $\dfrac{1}{x - \ln x} \ge \dfrac{1}{x}$. $\displaystyle\int_2^\infty \frac{1}{x}\,dx$ is divergent by Equation 2 with $p = 1 \le 1$, so $\displaystyle\int_2^\infty \frac{1}{x - \ln x}\,dx$ is divergent by the Comparison Theorem.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. For $x \geq 0$, $\arctan x < \dfrac{\pi}{2} < 2$, so $\dfrac{\arctan x}{2+e^x} < \dfrac{2}{2+e^x} < \dfrac{2}{e^x} = 2e^{-x}$. Now

$I = \displaystyle\int_0^\infty 2e^{-x}\,dx = \lim_{t\to\infty}\int_0^t 2e^{-x}\,dx = \lim_{t\to\infty}\left[-2e^{-x}\right]_0^t = \lim_{t\to\infty}\left(-\frac{2}{e^t}+2\right) = 2$, so I is convergent, and by comparison,

$\displaystyle\int_0^\infty \frac{\arctan x}{2+e^x}\,dx$ is convergent.

61. For $x > 1$, $f(x) = \dfrac{x+1}{\sqrt{x^4-x}} > \dfrac{x+1}{\sqrt{x^4}} > \dfrac{x}{x^2} = \dfrac{1}{x}$, so $\displaystyle\int_2^\infty f(x)\,dx$ diverges by comparison with $\displaystyle\int_2^\infty \frac{1}{x}\,dx$, which diverges by Theorem 2 with $p = 1 \leq 1$. Thus, $\int_1^\infty f(x)\,dx = \int_1^2 f(x)\,dx + \int_2^\infty f(x)\,dx$ also diverges.

62. For $x > 1$, $\dfrac{2+\cos x}{\sqrt{x^4+x^2}} \leq \dfrac{2+1}{\sqrt{x^4+x^2}} < \dfrac{3}{\sqrt{x^4}} = \dfrac{3}{x^2}$. $\displaystyle\int_1^\infty \frac{3}{x^2}\,dx = 3\int_1^\infty \frac{1}{x^2}\,dx$ is convergent by Equation 2 with $p = 2 > 1$, so $\displaystyle\int_1^\infty \frac{2+\cos x}{\sqrt{x^4+x^2}}\,dx$ is convergent by the Comparison Theorem.

63. For $0 < x \leq 1$, $\dfrac{\sec^2 x}{x\sqrt{x}} > \dfrac{1}{x^{3/2}}$. Now

$I = \displaystyle\int_0^1 x^{-3/2}\,dx = \lim_{t\to 0^+}\int_t^1 x^{-3/2}\,dx = \lim_{t\to 0^+}\left[-2x^{-1/2}\right]_t^1 = \lim_{t\to 0^+}\left(-2+\frac{2}{\sqrt{t}}\right) = \infty$, so I is divergent, and by comparison, $\displaystyle\int_0^1 \frac{\sec^2 x}{x\sqrt{x}}$ is divergent.

64. For $0 < x \leq 1$, $\dfrac{\sin^2 x}{\sqrt{x}} \leq \dfrac{1}{\sqrt{x}}$. Now

$I = \displaystyle\int_0^\pi \frac{1}{\sqrt{x}}\,dx = \lim_{t\to 0^+}\int_t^\pi x^{-1/2}\,dx = \lim_{t\to 0^+}\left[2x^{1/2}\right]_t^\pi = \lim_{t\to 0^+}\left(2\pi - 2\sqrt{t}\right) = 2\pi - 0 = 2\pi$, so I is convergent, and by comparison, $\displaystyle\int_0^\pi \frac{\sin^2 x}{\sqrt{x}}\,dx$ is convergent.

65. $I = \displaystyle\int_0^\infty \frac{1}{x^2}\,dx = \int_0^1 \frac{1}{x^2}\,dx + \int_1^\infty \frac{1}{x^2}\,dx = I_1 + I_2$. Now,

$I_1 = \displaystyle\lim_{t\to 0^+}\int_t^1 \frac{1}{x^2}\,dx = \lim_{t\to 0^+}\left[-x^{-1}\right]_t^1 = \lim_{t\to 0^+}\left[-1+\frac{1}{t}\right] = \infty$. Since I_1 is divergent, I is divergent, and there is no need to evaluate I_2.

66. $I = \displaystyle\int_0^\infty \frac{1}{\sqrt{x}}\,dx = \int_0^1 \frac{1}{x^{1/2}}\,dx + \int_1^\infty \frac{1}{x^{1/2}}\,dx = I_1 + I_2$. Since I_2 is divergent $\left[\text{Equation 2 with } p = \frac{1}{2} \leq 1\right]$, I is divergent, and there is no need to evaluate I_1.

67. $\displaystyle\int_0^\infty \frac{1}{\sqrt{x}\,(1+x)}\,dx = \int_0^1 \frac{dx}{\sqrt{x}\,(1+x)} + \int_1^\infty \frac{dx}{\sqrt{x}\,(1+x)} = \lim_{t\to 0^+}\int_t^1 \frac{dx}{\sqrt{x}\,(1+x)} + \lim_{t\to\infty}\int_1^t \frac{dx}{\sqrt{x}\,(1+x)}$. Now

$\displaystyle\int \frac{dx}{\sqrt{x}\,(1+x)} = \int \frac{2u\,du}{u(1+u^2)} \quad \left[\begin{matrix} u=\sqrt{x},\ x=u^2, \\ dx = 2u\,du \end{matrix}\right] \quad = 2\int \frac{du}{1+u^2} = 2\tan^{-1} u + C = 2\tan^{-1}\sqrt{x} + C$, so

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\int_0^\infty \frac{dx}{\sqrt{x}\,(1+x)} = \lim_{t\to 0^+}\left[2\tan^{-1}\sqrt{x}\,\right]_t^1 + \lim_{t\to\infty}\left[2\tan^{-1}\sqrt{x}\,\right]_1^t$$

$$= \lim_{t\to 0^+}\left[2\left(\tfrac{\pi}{4}\right) - 2\tan^{-1}\sqrt{t}\,\right] + \lim_{t\to\infty}\left[2\tan^{-1}\sqrt{t} - 2\left(\tfrac{\pi}{4}\right)\right] = \tfrac{\pi}{2} - 0 + 2\left(\tfrac{\pi}{2}\right) - \tfrac{\pi}{2} = \pi.$$

68. $\displaystyle\int_2^\infty \frac{1}{x\sqrt{x^2-4}}\,dx = \int_2^3 \frac{dx}{x\sqrt{x^2-4}} + \int_3^\infty \frac{dx}{x\sqrt{x^2-4}} = \lim_{t\to 2^+}\int_t^3 \frac{dx}{x\sqrt{x^2-4}} + \lim_{t\to\infty}\int_3^t \frac{dx}{x\sqrt{x^2-4}}$. Now

$$\int \frac{dx}{x\sqrt{x^2-4}} = \int \frac{2\sec\theta\,\tan\theta\,d\theta}{2\sec\theta\,2\tan\theta} \quad \left[\begin{array}{c} x = 2\sec\theta, \text{ where} \\ 0 \le \theta < \pi/2 \text{ or } \pi \le \theta < 3\pi/2 \end{array}\right] \quad = \tfrac{1}{2}\theta + C = \tfrac{1}{2}\sec^{-1}\left(\tfrac{1}{2}x\right) + C, \text{ so}$$

$$\int_2^\infty \frac{dx}{x\sqrt{x^2-4}} = \lim_{t\to 2^+}\left[\tfrac{1}{2}\sec^{-1}\left(\tfrac{1}{2}x\right)\right]_t^3 + \lim_{t\to\infty}\left[\tfrac{1}{2}\sec^{-1}\left(\tfrac{1}{2}x\right)\right]_3^t = \tfrac{1}{2}\sec^{-1}\left(\tfrac{3}{2}\right) - 0 + \tfrac{1}{2}\left(\tfrac{\pi}{2}\right) - \tfrac{1}{2}\sec^{-1}\left(\tfrac{3}{2}\right) = \tfrac{\pi}{4}.$$

69. If $p = 1$, then $\displaystyle\int_0^1 \frac{1}{x^p}\,dx = \lim_{t\to 0^+}\int_t^1 \frac{dx}{x} = \lim_{t\to 0^+}\left[\ln x\right]_t^1 = \infty.$ Divergent

If $p \neq 1$, then $\displaystyle\int_0^1 \frac{dx}{x^p} = \lim_{t\to 0^+}\int_t^1 \frac{dx}{x^p}$ [note that the integral is not improper if $p < 0$]

$$= \lim_{t\to 0^+}\left[\frac{x^{-p+1}}{-p+1}\right]_t^1 = \lim_{t\to 0^+}\frac{1}{1-p}\left[1 - \frac{1}{t^{p-1}}\right]$$

If $p > 1$, then $p - 1 > 0$, so $\dfrac{1}{t^{p-1}} \to \infty$ as $t \to 0^+$, and the integral diverges.

If $p < 1$, then $p - 1 < 0$, so $\dfrac{1}{t^{p-1}} \to 0$ as $t \to 0^+$ and $\displaystyle\int_0^1 \frac{dx}{x^p} = \frac{1}{1-p}\left[\lim_{t\to 0^+}\left(1 - t^{1-p}\right)\right] = \frac{1}{1-p}.$

Thus, the integral converges if and only if $p < 1$, and in that case its value is $\dfrac{1}{1-p}$.

70. Let $u = \ln x$. Then $du = dx/x \;\Rightarrow\; \displaystyle\int_e^\infty \frac{1}{x\,(\ln x)^p}\,dx = \int_1^\infty \frac{du}{u^p}$. By Example 4, this converges to $\dfrac{1}{p-1}$ if $p > 1$ and diverges otherwise.

71. First suppose $p = -1$. Then

$\displaystyle\int_0^1 x^p \ln x\,dx = \int_0^1 \frac{\ln x}{x}\,dx = \lim_{t\to 0^+}\int_t^1 \frac{\ln x}{x}\,dx = \lim_{t\to 0^+}\left[\tfrac{1}{2}(\ln x)^2\right]_t^1 = -\tfrac{1}{2}\lim_{t\to 0^+}(\ln t)^2 = -\infty$, so the integral diverges. Now suppose $p \neq -1$. Then integration by parts gives

$\displaystyle\int x^p \ln x\,dx = \frac{x^{p+1}}{p+1}\ln x - \int \frac{x^p}{p+1}\,dx = \frac{x^{p+1}}{p+1}\ln x - \frac{x^{p+1}}{(p+1)^2} + C$. If $p < -1$, then $p + 1 < 0$, so

$$\int_0^1 x^p \ln x\,dx = \lim_{t\to 0^+}\left[\frac{x^{p+1}}{p+1}\ln x - \frac{x^{p+1}}{(p+1)^2}\right]_t^1 = \frac{-1}{(p+1)^2} - \left(\frac{1}{p+1}\right)\lim_{t\to 0^+}\left[t^{p+1}\left(\ln t - \frac{1}{p+1}\right)\right] = \infty.$$

If $p > -1$, then $p + 1 > 0$ and

$$\int_0^1 x^p \ln x\,dx = \frac{-1}{(p+1)^2} - \left(\frac{1}{p+1}\right)\lim_{t\to 0^+}\frac{\ln t - 1/(p+1)}{t^{-(p+1)}} \overset{\text{H}}{=} \frac{-1}{(p+1)^2} - \left(\frac{1}{p+1}\right)\lim_{t\to 0^+}\frac{1/t}{-(p+1)t^{-(p+2)}}$$

$$= \frac{-1}{(p+1)^2} + \frac{1}{(p+1)^2}\lim_{t\to 0^+} t^{p+1} = \frac{-1}{(p+1)^2}$$

Thus, the integral converges to $-\dfrac{1}{(p+1)^2}$ if $p > -1$ and diverges otherwise.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

72. (a) $\boldsymbol{n=0}$: $\displaystyle\int_0^\infty x^n e^{-x}\,dx = \lim_{t\to\infty}\int_0^t e^{-x}\,dx = \lim_{t\to\infty}\left[-e^{-x}\right]_0^t = \lim_{t\to\infty}\left[-e^{-t}+1\right] = 0+1 = 1$

$\boldsymbol{n=1}$: $\displaystyle\int_0^\infty x^n e^{-x}\,dx = \lim_{t\to\infty}\int_0^t xe^{-x}\,dx$. To evaluate $\displaystyle\int xe^{-x}\,dx$, we'll use integration by parts with $u=x$, $dv=e^{-x}\,dx \;\Rightarrow\; du=dx$, $v=-e^{-x}$.

So $\displaystyle\int xe^{-x}\,dx = -xe^{-x} - \int -e^{-x}\,dx = -xe^{-x} - e^{-x} + C = (-x-1)e^{-x} + C$ and

$$\lim_{t\to\infty}\int_0^t xe^{-x}\,dx = \lim_{t\to\infty}\left[(-x-1)e^{-x}\right]_0^t = \lim_{t\to\infty}\left[(-t-1)e^{-t}+1\right] = \lim_{t\to\infty}\left[-te^{-t}-e^{-t}+1\right]$$
$$= 0-0+1 \quad [\text{use l'Hospital's Rule}] \quad = 1$$

$\boldsymbol{n=2}$: $\displaystyle\int_0^\infty x^n e^{-x}\,dx = \lim_{t\to\infty}\int_0^t x^2e^{-x}\,dx$. To evaluate $\displaystyle\int x^2e^{-x}\,dx$, we could use integration by parts again or Formula 97. Thus,

$$\lim_{t\to\infty}\int_0^t x^2e^{-x}\,dx = \lim_{t\to\infty}\left[-x^2e^{-x}\right]_0^t + 2\lim_{t\to\infty}\int_0^t xe^{-x}\,dx$$
$$= 0+0+2(1) \quad [\text{use l'Hospital's Rule and the result for } n=1] \quad = 2$$

$\boldsymbol{n=3}$: $\displaystyle\int_0^\infty x^n e^{-x}dx = \lim_{t\to\infty}\int_0^t x^3e^{-x}\,dx \overset{97}{=} \lim_{t\to\infty}\left[-x^3e^{-x}\right]_0^t + 3\lim_{t\to\infty}\int_0^t x^2e^{-x}\,dx$

$$= 0+0+3(2) \quad [\text{use l'Hospital's Rule and the result for } n=2] \quad = 6$$

(b) For $n=1$, 2, and 3, we have $\int_0^\infty x^ne^{-x}\,dx = 1$, 2, and 6. The values for the integral are equal to the factorials for n, so we guess $\int_0^\infty x^ne^{-x}\,dx = n!$.

(c) Suppose that $\int_0^\infty x^ke^{-x}\,dx = k!$ for some positive integer k. Then $\int_0^\infty x^{k+1}e^{-x}\,dx = \lim\limits_{t\to\infty}\int_0^t x^{k+1}e^{-x}\,dx$.

To evaluate $\int x^{k+1}e^{-x}\,dx$, we use parts with $u=x^{k+1}$, $dv=e^{-x}\,dx \;\Rightarrow\; du=(k+1)x^k\,dx$, $v=-e^{-x}$.

So $\int x^{k+1}e^{-x}dx = -x^{k+1}e^{-x} - \int -(k+1)x^ke^{-x}\,dx = -x^{k+1}e^{-x} + (k+1)\int x^ke^{-x}\,dx$ and

$$\lim_{t\to\infty}\int_0^t x^{k+1}e^{-x}\,dx = \lim_{t\to\infty}\left[-x^{k+1}e^{-x}\right]_0^t + (k+1)\lim_{t\to\infty}\int_0^t x^ke^{-x}\,dx$$
$$= \lim_{t\to\infty}\left[-t^{k+1}e^{-t}+0\right] + (k+1)k! = 0+0+(k+1)! = (k+1)!,$$

so the formula holds for $k+1$. By induction, the formula holds for all positive integers. (Since $0! = 1$, the formula holds for $n=0$, too.)

73. $\displaystyle I = \int_{-\infty}^\infty x\,dx = \int_{-\infty}^0 x\,dx + \int_0^\infty x\,dx$ and $\displaystyle\int_0^\infty x\,dx = \lim_{t\to\infty}\int_0^t x\,dx = \lim_{t\to\infty}\left[\tfrac{1}{2}x^2\right]_0^t = \lim_{t\to\infty}\left[\tfrac{1}{2}t^2-0\right] = \infty$,

so I is divergent. The Cauchy principal value of I is given by

$\displaystyle\lim_{t\to\infty}\int_{-t}^t x\,dx = \lim_{t\to\infty}\left[\tfrac{1}{2}x^2\right]_{-t}^t = \lim_{t\to\infty}\left[\tfrac{1}{2}t^2-\tfrac{1}{2}(-t)^2\right] = \lim_{t\to\infty}[\,0\,] = 0$. Hence, I is divergent, but its Cauchy principal value is 0.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

74. Let $k = \dfrac{M}{2RT}$ so that $\overline{v} = \dfrac{4}{\sqrt{\pi}}\, k^{3/2} \displaystyle\int_0^\infty v^3 e^{-kv^2}\, dv$. Let I denote the integral and use parts to integrate I. Let $\alpha = v^2$, $d\beta = ve^{-kv^2}\, dv \quad \Rightarrow \quad d\alpha = 2v\, dv$, $\beta = -\dfrac{1}{2k} e^{-kv^2}$:

$$I = \lim_{t\to\infty} \left[-\frac{1}{2k} v^2 e^{-kv^2}\right]_0^t + \frac{1}{k}\int_0^\infty ve^{-kv^2}\, dv_0^t = -\frac{1}{2k}\lim_{t\to\infty}\left(t^2 e^{-kt^2}\right) + \frac{1}{k}\lim_{t\to\infty}\left[-\frac{1}{2k}e^{-kv^2}\right]$$

$$\overset{\text{H}}{=} -\frac{1}{2k}\cdot 0 - \frac{1}{2k^2}(0-1) = \frac{1}{2k^2}$$

Thus, $\overline{v} = \dfrac{4}{\sqrt{\pi}} k^{3/2} \cdot \dfrac{1}{2k^2} = \dfrac{2}{(k\pi)^{1/2}} = \dfrac{2}{[\pi M/(2RT)]^{1/2}} = \dfrac{2\sqrt{2}\sqrt{RT}}{\sqrt{\pi M}} = \sqrt{\dfrac{8RT}{\pi M}}$.

75. Volume $= \displaystyle\int_1^\infty \pi\left(\frac{1}{x}\right)^2 dx = \pi \lim_{t\to\infty}\int_1^t \frac{dx}{x^2} = \pi\lim_{t\to\infty}\left[-\frac{1}{x}\right]_1^t = \pi\lim_{t\to\infty}\left(1-\frac{1}{t}\right) = \pi < \infty$.

76. Work $= \displaystyle\int_R^\infty \frac{GMm}{r^2}\, dr = \lim_{t\to\infty}\int_R^t \frac{GMm}{r^2}\, dr = \lim_{t\to\infty} GMm\left[\frac{-1}{r}\right]_R^t = GMm\lim_{t\to\infty}\left(\frac{-1}{t}+\frac{1}{R}\right) = \frac{GMm}{R}$, where $M =$ mass of the earth $= 5.98\times 10^{24}$ kg, $m =$ mass of satellite $= 10^3$ kg, $R =$ radius of the earth $= 6.37\times 10^6$ m, and $G =$ gravitational constant $= 6.67\times 10^{-11}$ N·m^2/kg.

Therefore, Work $= \dfrac{6.67\times 10^{-11}\cdot 5.98\times 10^{24}\cdot 10^3}{6.37\times 10^6} \approx 6.26\times 10^{10}$ J.

77. Work $= \displaystyle\int_R^\infty F\, dr = \lim_{t\to\infty}\int_R^t \frac{GmM}{r^2}\, dr = \lim_{t\to\infty} GmM\left(\frac{1}{R}-\frac{1}{t}\right) = \frac{GmM}{R}$. The initial kinetic energy provides the work, so $\frac{1}{2}mv_0^2 = \dfrac{GmM}{R} \quad \Rightarrow \quad v_0 = \sqrt{\dfrac{2GM}{R}}$.

78. $y(s) = \displaystyle\int_s^R \frac{2r}{\sqrt{r^2-s^2}}\, x(r)\, dr$ and $x(r) = \frac{1}{2}(R-r)^2 \quad \Rightarrow$

$$y(s) = \lim_{t\to s^+}\int_t^R \frac{r(R-r)^2}{\sqrt{r^2-s^2}}\, dr = \lim_{t\to s^+}\int_t^R \frac{r^3-2Rr^2+R^2r}{\sqrt{r^2-s^2}}\, dr$$

$$= \lim_{t\to s^+}\left[\int_t^R \frac{r^3\, dr}{\sqrt{r^2-s^2}} - 2R\int_t^R \frac{r^2\, dr}{\sqrt{r^2-s^2}} + R^2\int_t^R \frac{r\, dr}{\sqrt{r^2-s^2}}\right] = \lim_{t\to s^+}\left(I_1 - 2RI_2 + R^2 I_3\right) = L$$

For I_1: Let $u = \sqrt{r^2-s^2} \quad \Rightarrow \quad u^2 = r^2 - s^2$, $r^2 = u^2 + s^2$, $2r\, dr = 2u\, du$, so, omitting limits and constant of integration,

$$I_1 = \int \frac{(u^2+s^2)u}{u}\, du = \int (u^2+s^2)\, du = \tfrac{1}{3}u^3 + s^2 u = \tfrac{1}{3}u(u^2+3s^2)$$
$$= \tfrac{1}{3}\sqrt{r^2-s^2}\,(r^2-s^2+3s^2) = \tfrac{1}{3}\sqrt{r^2-s^2}\,(r^2+2s^2)$$

For I_2: Using Formula 44, $I_2 = \dfrac{r}{2}\sqrt{r^2-s^2} + \dfrac{s^2}{2}\ln\left|r+\sqrt{r^2-s^2}\right|$.

For I_3: Let $u = r^2 - s^2 \quad \Rightarrow \quad du = 2r\, dr$. Then $I_3 = \dfrac{1}{2}\displaystyle\int \frac{du}{\sqrt{u}} = \frac{1}{2}\cdot 2\sqrt{u} = \sqrt{r^2-s^2}$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Thus,

$$\begin{aligned}
L &= \lim_{t\to s^+}\left[\tfrac{1}{3}\sqrt{r^2-s^2}\,(r^2+2s^2)-2R\left(\frac{r}{2}\sqrt{r^2-s^2}+\frac{s^2}{2}\ln\left|r+\sqrt{r^2-s^2}\,\right|\right)+R^2\sqrt{r^2-s^2}\right]_t^R \\
&= \lim_{t\to s^+}\left[\tfrac{1}{3}\sqrt{R^2-s^2}(R^2+2s^2)-2R\left(\frac{R}{2}\sqrt{R^2-s^2}+\frac{s^2}{2}\ln\left|R+\sqrt{R^2-s^2}\,\right|\right)+R^2\sqrt{R^2-s^2}\right] \\
&\qquad -\lim_{t\to s^+}\left[\tfrac{1}{3}\sqrt{t^2-s^2}\,(t^2+2s^2)-2R\left(\frac{t}{2}\sqrt{t^2-s^2}+\frac{s^2}{2}\ln\left|t+\sqrt{t^2-s^2}\,\right|\right)+R^2\sqrt{t^2-s^2}\right] \\
&= \left[\tfrac{1}{3}\sqrt{R^2-s^2}\,(R^2+2s^2)-Rs^2\ln\left|R+\sqrt{R^2-s^2}\,\right|\right]-\left[-Rs^2\ln|s|\right] \\
&= \tfrac{1}{3}\sqrt{R^2-s^2}\,(R^2+2s^2)-Rs^2\ln\left(\frac{R+\sqrt{R^2-s^2}}{s}\right)
\end{aligned}$$

79. We would expect a small percentage of bulbs to burn out in the first few hundred hours, most of the bulbs to burn out after close to 700 hours, and a few overachievers to burn on and on.

(a)

(b) $r(t)=F'(t)$ is the rate at which the fraction $F(t)$ of burnt-out bulbs increases as t increases. This could be interpreted as a fractional burnout rate.

(c) $\int_0^\infty r(t)\,dt=\lim\limits_{x\to\infty}F(x)=1$, since all of the bulbs will eventually burn out.

80. $I=\displaystyle\int_0^\infty te^{kt}\,dt=\lim_{s\to\infty}\left[\frac{1}{k^2}(kt-1)\,e^{kt}\right]_0^s$ [Formula 96, or parts] $=\displaystyle\lim_{s\to\infty}\left[\left(\frac{1}{k}se^{ks}-\frac{1}{k^2}e^{ks}\right)-\left(-\frac{1}{k^2}\right)\right]$.

Since $k<0$ the first two terms approach 0 (you can verify that the first term does so with l'Hospital's Rule), so the limit is equal to $1/k^2$. Thus, $M=-kI=-k\big(1/k^2\big)=-1/k=-1/(-0.000121)\approx 8264.5$ years.

81. $\gamma=\displaystyle\int_0^\infty\frac{cN(1-e^{-kt})}{k}\,e^{-\lambda t}\,dt=\frac{cN}{k}\lim_{x\to\infty}\int_0^x\left[e^{-\lambda t}-e^{(-k-\lambda)t}\right]dt$

$$\begin{aligned}
&= \frac{cN}{k}\lim_{x\to\infty}\left[\frac{1}{-\lambda}\,e^{-\lambda t}-\frac{1}{-k-\lambda}\,e^{(-k-\lambda)t}\right]_0^x=\frac{cN}{k}\lim_{x\to\infty}\left[\frac{1}{-\lambda e^{\lambda x}}+\frac{1}{(k+\lambda)e^{(k+\lambda)x}}-\left(\frac{1}{-\lambda}+\frac{1}{k+\lambda}\right)\right] \\
&= \frac{cN}{k}\left(\frac{1}{\lambda}-\frac{1}{k+\lambda}\right)=\frac{cN}{k}\left(\frac{k+\lambda-\lambda}{\lambda(k+\lambda)}\right)=\frac{cN}{\lambda(k+\lambda)}
\end{aligned}$$

82. $\displaystyle\int_0^\infty u(t)\,dt=\lim_{x\to\infty}\int_0^x\frac{r}{V}\,C_0\,e^{-rt/V}\,dt=\frac{r}{V}\,C_0\lim_{x\to\infty}\left[\frac{e^{-rt/V}}{-r/V}\right]_0^x=\frac{r}{V}\,C_0\left(-\frac{V}{r}\right)\lim_{x\to\infty}\left(e^{-rx/V}-1\right)$

$$= -C_0(0-1)=C_0.$$

$\int_0^\infty u(t)\,dt$ represents the total amount of urea removed from the blood if dialysis is continued indefinitely. The fact that $\int_0^\infty u(t)\,dt=C_0$ means that, in the limit, as $t\to\infty$, all the urea in the blood at time $t=0$ is removed. The calculation says nothing about how rapidly that limit is approached.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

83. $I = \displaystyle\int_a^\infty \frac{1}{x^2+1}\,dx = \lim_{t\to\infty}\int_a^t \frac{1}{x^2+1}\,dx = \lim_{t\to\infty}\left[\tan^{-1}x\right]_a^t = \lim_{t\to\infty}\left(\tan^{-1}t - \tan^{-1}a\right) = \frac{\pi}{2} - \tan^{-1}a.$

$I < 0.001 \;\Rightarrow\; \frac{\pi}{2} - \tan^{-1}a < 0.001 \;\Rightarrow\; \tan^{-1}a > \frac{\pi}{2} - 0.001 \;\Rightarrow\; a > \tan\left(\frac{\pi}{2} - 0.001\right) \approx 1000.$

84. $f(x) = e^{-x^2}$ and $\Delta x = \frac{4-0}{8} = \frac{1}{2}$.

$$\textstyle\int_0^4 f(x)\,dx \approx S_8 = \frac{1}{2\cdot 3}[f(0) + 4f(0.5) + 2f(1) + \cdots + 2f(3) + 4f(3.5) + f(4)] \approx \frac{1}{6}(5.31717808) \approx 0.8862$$

Now $x > 4 \;\Rightarrow\; -x\cdot x < -x\cdot 4 \;\Rightarrow\; e^{-x^2} < e^{-4x} \;\Rightarrow\; \int_4^\infty e^{-x^2}\,dx < \int_4^\infty e^{-4x}\,dx.$

$\int_4^\infty e^{-4x}\,dx = \lim_{t\to\infty}\left[-\frac{1}{4}e^{-4x}\right]_4^t = -\frac{1}{4}\left(0 - e^{-16}\right) = 1/(4e^{16}) \approx 0.0000000281 < 0.0000001$, as desired.

85. (a) $F(s) = \displaystyle\int_0^\infty f(t)e^{-st}\,dt = \int_0^\infty e^{-st}\,dt = \lim_{n\to\infty}\left[-\frac{e^{-st}}{s}\right]_0^n = \lim_{n\to\infty}\left(\frac{e^{-sn}}{-s} + \frac{1}{s}\right)$. This converges to $\dfrac{1}{s}$ only if $s > 0$.

Therefore $F(s) = \dfrac{1}{s}$ with domain $\{s \mid s > 0\}$.

(b) $F(s) = \displaystyle\int_0^\infty f(t)e^{-st}\,dt = \int_0^\infty e^t e^{-st}\,dt = \lim_{n\to\infty}\int_0^n e^{t(1-s)}\,dt = \lim_{n\to\infty}\left[\frac{1}{1-s}e^{t(1-s)}\right]_0^n$

$$= \lim_{n\to\infty}\left(\frac{e^{(1-s)n}}{1-s} - \frac{1}{1-s}\right)$$

This converges only if $1 - s < 0 \;\Rightarrow\; s > 1$, in which case $F(s) = \dfrac{1}{s-1}$ with domain $\{s \mid s > 1\}$.

(c) $F(s) = \int_0^\infty f(t)e^{-st}\,dt = \lim_{n\to\infty}\int_0^n te^{-st}\,dt$. Use integration by parts: let $u = t$, $dv = e^{-st}\,dt \;\Rightarrow\; du = dt$,

$v = -\dfrac{e^{-st}}{s}$. Then $F(s) = \displaystyle\lim_{n\to\infty}\left[-\frac{t}{s}e^{-st} - \frac{1}{s^2}e^{-st}\right]_0^n = \lim_{n\to\infty}\left(\frac{-n}{se^{sn}} - \frac{1}{s^2e^{sn}} + 0 + \frac{1}{s^2}\right) = \frac{1}{s^2}$ only if $s > 0$.

Therefore, $F(s) = \dfrac{1}{s^2}$ and the domain of F is $\{s \mid s > 0\}$.

86. $0 \le f(t) \le Me^{at} \;\Rightarrow\; 0 \le f(t)e^{-st} \le Me^{at}e^{-st}$ for $t \ge 0$. Now use the Comparison Theorem:

$$\int_0^\infty Me^{at}e^{-st}\,dt = \lim_{n\to\infty} M\int_0^n e^{t(a-s)}\,dt = M\cdot\lim_{n\to\infty}\left[\frac{1}{a-s}e^{t(a-s)}\right]_0^n = M\cdot\lim_{n\to\infty}\frac{1}{a-s}\left[e^{n(a-s)} - 1\right]$$

This is convergent only when $a - s < 0 \;\Rightarrow\; s > a$. Therefore, by the Comparison Theorem, $F(s) = \int_0^\infty f(t)\,e^{-st}\,dt$ is also convergent for $s > a$.

87. $G(s) = \int_0^\infty f'(t)e^{-st}\,dt$. Integrate by parts with $u = e^{-st}$, $dv = f'(t)\,dt \;\Rightarrow\; du = -se^{-st}$, $v = f(t)$:

$$G(s) = \lim_{n\to\infty}\left[f(t)e^{-st}\right]_0^n + s\int_0^\infty f(t)e^{-st}\,dt = \lim_{n\to\infty} f(n)e^{-sn} - f(0) + sF(s)$$

But $0 \le f(t) \le Me^{at} \;\Rightarrow\; 0 \le f(t)e^{-st} \le Me^{at}e^{-st}$ and $\lim_{t\to\infty} Me^{t(a-s)} = 0$ for $s > a$. So by the Squeeze Theorem,

$\lim_{t\to\infty} f(t)e^{-st} = 0$ for $s > a \;\Rightarrow\; G(s) = 0 - f(0) + sF(s) = sF(s) - f(0)$ for $s > a$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

88. Assume without loss of generality that $a < b$. Then

$$\begin{aligned}
\int_{-\infty}^{a} f(x)\,dx + \int_{a}^{\infty} f(x)\,dx &= \lim_{t\to-\infty} \int_t^a f(x)\,dx + \lim_{u\to\infty} \int_a^u f(x)\,dx \\
&= \lim_{t\to-\infty} \int_t^a f(x)\,dx + \lim_{u\to\infty} \left[\int_a^b f(x)\,dx + \int_b^u f(x)\,dx\right] \\
&= \lim_{t\to-\infty} \int_t^a f(x)\,dx + \int_a^b f(x)\,dx + \lim_{u\to\infty} \int_b^u f(x)\,dx \\
&= \lim_{t\to-\infty} \left[\int_t^a f(x)\,dx + \int_a^b f(x)\,dx\right] + \int_b^\infty f(x)\,dx \\
&= \lim_{t\to-\infty} \int_t^b f(x)\,dx + \int_b^\infty f(x)\,dx = \int_{-\infty}^{b} f(x)\,dx + \int_b^\infty f(x)\,dx
\end{aligned}$$

89. We use integration by parts: let $u = x$, $dv = xe^{-x^2}\,dx \quad\Rightarrow\quad du = dx$, $v = -\frac{1}{2}e^{-x^2}$. So

$$\int_0^\infty x^2 e^{-x^2}\,dx = \lim_{t\to\infty}\left[-\frac{1}{2}xe^{-x^2}\right]_0^t + \frac{1}{2}\int_0^\infty e^{-x^2}\,dx = \lim_{t\to\infty}\left[-\frac{t}{2e^{t^2}}\right] + \frac{1}{2}\int_0^\infty e^{-x^2}\,dx = \frac{1}{2}\int_0^\infty e^{-x^2}\,dx$$

(The limit is 0 by l'Hospital's Rule.)

90. $\int_0^\infty e^{-x^2}\,dx$ is the area under the curve $y = e^{-x^2}$ for $0 \le x < \infty$ and $0 < y \le 1$. Solving $y = e^{-x^2}$ for x, we get $y = e^{-x^2} \quad\Rightarrow\quad \ln y = -x^2 \quad\Rightarrow\quad -\ln y = x^2 \quad\Rightarrow\quad x = \pm\sqrt{-\ln y}$. Since x is positive, choose $x = \sqrt{-\ln y}$, and the area is represented by $\int_0^1 \sqrt{-\ln y}\,dy$. Therefore, each integral represents the same area, so the integrals are equal.

91. For the first part of the integral, let $x = 2\tan\theta \quad\Rightarrow\quad dx = 2\sec^2\theta\,d\theta$.

$$\int \frac{1}{\sqrt{x^2+4}}\,dx = \int \frac{2\sec^2\theta}{2\sec\theta}\,d\theta = \int \sec\theta\,d\theta = \ln|\sec\theta + \tan\theta|.$$

From the figure, $\tan\theta = \dfrac{x}{2}$, and $\sec\theta = \dfrac{\sqrt{x^2+4}}{2}$. So

$$\begin{aligned}
I = \int_0^\infty \left(\frac{1}{\sqrt{x^2+4}} - \frac{C}{x+2}\right)dx &= \lim_{t\to\infty}\left[\ln\left|\frac{\sqrt{x^2+4}}{2} + \frac{x}{2}\right| - C\ln|x+2|\right]_0^t \\
&= \lim_{t\to\infty}\left[\ln\frac{\sqrt{t^2+4}+t}{2} - C\ln(t+2) - (\ln 1 - C\ln 2)\right] \\
&= \lim_{t\to\infty}\left[\ln\left(\frac{\sqrt{t^2+4}+t}{2\,(t+2)^C}\right) + \ln 2^C\right] = \ln\left(\lim_{t\to\infty}\frac{t+\sqrt{t^2+4}}{(t+2)^C}\right) + \ln 2^{C-1}
\end{aligned}$$

Now $L = \displaystyle\lim_{t\to\infty}\frac{t+\sqrt{t^2+4}}{(t+2)^C} \overset{\text{H}}{=} \lim_{t\to\infty}\frac{1+t/\sqrt{t^2+4}}{C\,(t+2)^{C-1}} = \frac{2}{C\lim\limits_{t\to\infty}(t+2)^{C-1}}$.

If $C < 1$, $L = \infty$ and I diverges.

If $C = 1$, $L = 2$ and I converges to $\ln 2 + \ln 2^0 = \ln 2$.

If $C > 1$, $L = 0$ and I diverges to $-\infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

92. $I = \int_0^\infty \left(\frac{x}{x^2+1} - \frac{C}{3x+1} \right) dx = \lim_{t\to\infty} \left[\frac{1}{2}\ln(x^2+1) - \frac{1}{3}C\ln(3x+1) \right]_0^t = \lim_{t\to\infty} \left[\ln(t^2+1)^{1/2} - \ln(3t+1)^{C/3} \right]$

$$= \lim_{t\to\infty} \left(\ln \frac{(t^2+1)^{1/2}}{(3t+1)^{C/3}} \right) = \ln\left(\lim_{t\to\infty} \frac{\sqrt{t^2+1}}{(3t+1)^{C/3}} \right)$$

For $C \le 0$, the integral diverges. For $C > 0$, we have

$$L = \lim_{t\to\infty} \frac{\sqrt{t^2+1}}{(3t+1)^{C/3}} \overset{\text{H}}{=} \lim_{t\to\infty} \frac{t/\sqrt{t^2+1}}{C(3t+1)^{(C/3)-1}} = \frac{1}{C} \lim_{t\to\infty} \frac{1}{(3t+1)^{(C/3)-1}}$$

For $C/3 < 1 \quad \Leftrightarrow \quad C < 3$, $L = \infty$ and I diverges.

For $C = 3$, $L = \frac{1}{3}$ and $I = \ln \frac{1}{3}$.

For $C > 3$, $L = 0$ and I diverges to $-\infty$.

93. No, $I = \int_0^\infty f(x)\,dx$ must be *divergent*. Since $\lim_{x\to\infty} f(x) = 1$, there must exist an N such that if $x \ge N$, then $f(x) \ge \frac{1}{2}$.

Thus, $I = I_1 + I_2 = \int_0^N f(x)\,dx + \int_N^\infty f(x)\,dx$, where I_1 is an ordinary definite integral that has a finite value, and I_2 is improper and diverges by comparison with the divergent integral $\int_N^\infty \frac{1}{2}\,dx$.

94. As in Exercises 65–68, we let $I = \int_0^\infty \frac{x^a}{1+x^b}\,dx = I_1 + I_2$, where $I_1 = \int_0^1 \frac{x^a}{1+x^b}\,dx$ and $I_2 = \int_1^\infty \frac{x^a}{1+x^b}\,dx$. We will show that I_1 converges for $a > -1$ and I_2 converges for $b > a+1$, so that I converges when $a > -1$ and $b > a+1$.

I_1 is improper only when $a < 0$. When $0 \le x \le 1$, we have $\frac{1}{1+x^b} \le 1 \quad \Rightarrow \quad \frac{1}{x^{-a}(1+x^b)} \le \frac{1}{x^{-a}}$. The integral $\int_0^1 \frac{1}{x^{-a}}\,dx$ converges for $-a < 1$ [or $a > -1$] by Exercise 69, so by the Comparison Theorem, $\int_0^1 \frac{1}{x^{-a}(1+x^b)}\,dx$ converges for $-1 < a < 0$. I_1 is not improper when $a \ge 0$, so it has a finite real value in that case. Therefore, I_1 has a finite real value (converges) when $a > -1$.

I_2 is always improper. When $x \ge 1$, $\frac{x^a}{1+x^b} = \frac{1}{x^{-a}(1+x^b)} = \frac{1}{x^{-a}+x^{b-a}} < \frac{1}{x^{b-a}}$. By (2), $\int_1^\infty \frac{1}{x^{b-a}}\,dx$ converges for $b-a > 1$ (or $b > a+1$), so by the Comparison Theorem, $\int_1^\infty \frac{x^a}{1+x^b}\,dx$ converges for $b > a+1$.

Thus, I converges if $a > -1$ and $b > a+1$.

7 Review

TRUE-FALSE QUIZ

1. True. See Example 5 in Section 7.1.

2. True. Integration by parts can be used to show that $\int x^n e^x\,dx = x^n e^x - n\int x^{n-1}e^x\,dx$, so that the power of x in the new integrand is reduced by 1. Hence, when $n = 5$, repeatedly applying integration by parts five times will reduce the final integral to $\int e^x\,dx$, which evaluates to e^x.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. False. Substituting $x = 5\sin\theta$ into $\sqrt{25+x^2}$ gives $\sqrt{25+25\sin^2\theta}$. This expression cannot be further simplified using a trigonometric identity. A more useful substitiution would be $x = 5\tan\theta$.

4. False. To use entry 25, we need to first write $\displaystyle\int \frac{dx}{\sqrt{9+e^{2x}}}$ in the form $\displaystyle\int \frac{du}{\sqrt{9+u^2}}$, which suggests making the substitution $u = e^x$, so that $du = e^x\,dx$, or $du/u = dx$. Thus, $\displaystyle\int \frac{dx}{\sqrt{9+e^{2x}}} = \int \frac{du}{u\sqrt{9+u^2}}$, however, entry 25 cannot be used to evaluate this new integral. Instead, entry 27 would be needed.

5. False. Since the numerator has a higher degree than the denominator, $\dfrac{x(x^2+4)}{x^2-4} = x + \dfrac{8x}{x^2-4} = x + \dfrac{A}{x+2} + \dfrac{B}{x-2}$.

6. True. $\dfrac{x^2+4}{x(x^2-4)} = \dfrac{-1}{x} + \dfrac{1}{x+2} + \dfrac{1}{x-2}$

7. False. $\dfrac{x^2+4}{x^2(x-4)}$ can be put in the form $\dfrac{A}{x} + \dfrac{B}{x^2} + \dfrac{C}{x-4}$.

8. False. $\dfrac{x^2-4}{x(x^2+4)}$ can be put into the form $\dfrac{A}{x} + \dfrac{Bx+C}{x^2+4}$.

9. False. This is an improper integral, since the denominator vanishes at $x = 1$.

$$\int_0^4 \frac{x}{x^2-1}\,dx = \int_0^1 \frac{x}{x^2-1}\,dx + \int_1^4 \frac{x}{x^2-1}\,dx \text{ and}$$

$$\int_0^1 \frac{x}{x^2-1}\,dx = \lim_{t\to 1^-} \int_0^t \frac{x}{x^2-1}\,dx = \lim_{t\to 1^-} \left[\tfrac{1}{2}\ln\left|x^2-1\right|\right]_0^t = \lim_{t\to 1^-} \tfrac{1}{2}\ln\left|t^2-1\right| = \infty$$

So the integral diverges.

10. True by Theorem 7.8.2 with $p = \sqrt{2} > 1$.

11. True. $\displaystyle\int_{-\infty}^{\infty} f(x)\,dx = \int_{-\infty}^{0} f(x)\,dx + \int_0^{\infty} f(x)\,dx = I_1 + I_2$. If $\displaystyle\int_{-\infty}^{\infty} f(x)\,dx$ is convergent, it follows that both I_1 and I_2 must be convergent.

12. False. For example, with $n = 1$ the Trapezoidal Rule is much more accurate than the Midpoint Rule for the function in the diagram.

13. (a) True. See the end of Section 7.5.

(b) False. Examples include the functions $f(x) = e^{x^2}$, $g(x) = \sin(x^2)$, and $h(x) = \dfrac{\sin x}{x}$.

14. True. If f is continuous on $[0,\infty)$, then $\int_0^1 f(x)\,dx$ is finite. Since $\int_1^\infty f(x)\,dx$ is finite, so is $\int_0^\infty f(x)\,dx = \int_0^1 f(x)\,dx + \int_1^\infty f(x)\,dx$.

15. False. If $f(x) = 1/x$, then f is continuous and decreasing on $[1,\infty)$ with $\lim\limits_{x\to\infty} f(x) = 0$, but $\int_1^\infty f(x)\,dx$ is divergent.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. True. $\int_a^\infty [f(x)+g(x)]\,dx = \lim_{t\to\infty}\int_a^t [f(x)+g(x)]\,dx = \lim_{t\to\infty}\left(\int_a^t f(x)\,dx + \int_a^t g(x)\,dx\right)$

$$= \lim_{t\to\infty}\int_a^t f(x)\,dx + \lim_{t\to\infty}\int_a^t g(x)\,dx \quad \left[\begin{array}{c}\text{since both limits}\\ \text{in the sum exist}\end{array}\right]$$

$$= \int_a^\infty f(x)\,dx + \int_a^\infty g(x)\,dx$$

Since the two integrals are finite, so is their sum.

17. False. Take $f(x)=1$ for all x and $g(x)=-1$ for all x. Then $\int_a^\infty f(x)\,dx=\infty$ [divergent] and $\int_a^\infty g(x)\,dx=-\infty$ [divergent], but $\int_a^\infty [f(x)+g(x)]\,dx=0$ [convergent].

18. False. $\int_0^\infty f(x)\,dx$ could converge or diverge. For example, if $g(x)=1$, then $\int_0^\infty f(x)\,dx$ diverges if $f(x)=1$ and converges if $f(x)=0$.

EXERCISES

1. $$\int_1^2 \frac{(x+1)^2}{x}\,dx = \int_1^2 \frac{x^2+2x+1}{x}\,dx = \int_1^2\left(x+2+\frac{1}{x}\right)dx = \left[\frac{1}{2}x^2+2x+\ln|x|\right]_1^2$$
$$= (2+4+\ln 2)-\left(\tfrac{1}{2}+2+0\right) = \tfrac{7}{2}+\ln 2$$

2. $$\int_1^2 \frac{x}{(x+1)^2}\,dx = \int_2^3 \frac{u-1}{u^2}\,du \qquad \left[\begin{array}{l} u=x+1,\\ du=dx\end{array}\right]$$
$$= \int_2^3\left(\frac{1}{u}-\frac{1}{u^2}\right)du = \left[\ln|u|+\frac{1}{u}\right]_2^3 = \left(\ln 3+\frac{1}{3}\right)-\left(\ln 2+\frac{1}{2}\right) = \ln\frac{3}{2}-\frac{1}{6}$$

3. $$\int \frac{e^{\sin x}}{\sec x}\,dx = \int \cos x\, e^{\sin x}\,dx = \int e^u\,du \qquad \left[\begin{array}{l} u=\sin x,\\ du=\cos x\,dx\end{array}\right]$$
$$= e^u + C = e^{\sin x}+C$$

4. $$\int_0^{\pi/6} t\sin 2t\,dt = \left[-\tfrac{1}{2}t\cos 2t\right]_0^{\pi/6} - \int_0^{\pi/6}\left(-\tfrac{1}{2}\cos 2t\right)dt \qquad \left[\begin{array}{ll} u=t, & dv=\sin 2t\\ du=dt, & v=-\frac{1}{2}\cos 2t\end{array}\right]$$
$$= \left(-\tfrac{\pi}{12}\cdot\tfrac{1}{2}\right)-(0)+\left[\tfrac{1}{4}\sin 2t\right]_0^{\pi/6} = -\tfrac{\pi}{24}+\tfrac{1}{8}\sqrt{3}$$

5. $$\int \frac{dt}{2t^2+3t+1} = \int \frac{1}{(2t+1)(t+1)}\,dt = \int\left(\frac{2}{2t+1}-\frac{1}{t+1}\right)dt \quad \text{[partial fractions]} \quad = \ln|2t+1|-\ln|t+1|+C$$

6. $$\int_1^2 x^5\ln x\,dx = \left[\tfrac{1}{6}x^6\ln x\right]_1^2 - \int_1^2 \tfrac{1}{6}x^5\,dx \qquad \left[\begin{array}{ll} u=\ln x, & dv=x^5\,dx\\ du=\frac{1}{x}\,dx, & v=\frac{1}{6}x^6\end{array}\right]$$
$$= \tfrac{64}{6}\ln 2 - 0 - \left[\tfrac{1}{36}x^6\right]_1^2 = \tfrac{32}{3}\ln 2 - \left(\tfrac{64}{36}-\tfrac{1}{36}\right) = \tfrac{32}{3}\ln 2 - \tfrac{7}{4}$$

7. $$\int_0^{\pi/2}\sin^3\theta\,\cos^2\theta\,d\theta = \int_0^{\pi/2}(1-\cos^2\theta)\cos^2\theta\,\sin\theta\,d\theta = \int_1^0(1-u^2)u^2\,(-du) \qquad \left[\begin{array}{l} u=\cos\theta,\\ du=-\sin\theta\,d\theta\end{array}\right]$$
$$= \int_0^1(u^2-u^4)\,du = \left[\tfrac{1}{3}u^3-\tfrac{1}{5}u^5\right]_0^1 = \left(\tfrac{1}{3}-\tfrac{1}{5}\right)-0 = \tfrac{2}{15}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. Let $x = 4\sin\theta$, where $-\pi/2 \le \theta \le \pi/2$. Then $dx = 4\cos\theta\, d\theta$ and

$\sqrt{16-x^2} = \sqrt{16-16\sin^2\theta} = \sqrt{16\cos^2\theta} = 4\,|\cos\theta\,| = 4\cos\theta$. Thus,

$$\int \frac{dx}{x^2\sqrt{16-x^2}} = \int \frac{4\cos\theta}{16\sin^2\theta\,(4\cos\theta)}\,d\theta = \frac{1}{16}\int \csc^2\theta\,d\theta$$

$$= -\frac{1}{16}\cot\theta + C = -\frac{\sqrt{16-x^2}}{16x} + C$$

9. Let $u = \ln t$, $du = dt/t$. Then $\displaystyle\int \frac{\sin(\ln t)}{t}\,dt = \int \sin u\,du = -\cos u + C = -\cos(\ln t) + C$.

10. Let $u = \arctan x$, $du = dx/(1+x^2)$. Then

$$\int_0^1 \frac{\sqrt{\arctan x}}{1+x^2}\,dx = \int_0^{\pi/4} \sqrt{u}\,du = \frac{2}{3}\Big[u^{3/2}\Big]_0^{\pi/4} = \frac{2}{3}\left[\frac{\pi^{3/2}}{4^{3/2}} - 0\right] = \frac{2}{3}\cdot\frac{1}{8}\pi^{3/2} = \frac{1}{12}\pi^{3/2}.$$

11. First let $u = (\ln x)^2$, $dv = x\,dx \;\Rightarrow\; du = \dfrac{2\ln x}{x}\,dx$, $v = \dfrac{1}{2}x^2$. Then $I = \displaystyle\int x\,(\ln x)^2\,dx = \frac{1}{2}x^2\,(\ln x)^2 - \int x\ln x\,dx$.

Next, let $U = \ln x$, $dV = x\,dx \;\Rightarrow\; dU = \dfrac{1}{x}\,dx$, $V = \dfrac{1}{2}x^2$, so $\displaystyle\int x\ln x\,dx = \frac{1}{2}x^2\ln x - \frac{1}{2}\int x\,dx = \frac{1}{2}x^2\ln x - \frac{1}{4}x^2$.

Substituting in the previous formula gives $I = \frac{1}{2}x^2\,(\ln x)^2 - \left(\frac{1}{2}x^2\ln x - \frac{1}{4}x^2\right) + C = \frac{1}{4}x^2[2(\ln x)^2 - 2\ln x + 1] + C$.

12. Let $t = \cos x$, so that $dt = -\sin x\,dx$. Then

$$\int \sin x\cos x\ln(\cos x)\,dx = \int t\ln t\,(-dt) = -\tfrac{1}{2}t^2\ln t + \int \tfrac{1}{2}t\,dt \qquad \begin{bmatrix} u = \ln t,\ dv = -t\,dt \\ du = \frac{1}{t}\,dt,\ v = -\frac{1}{2}t^2 \end{bmatrix}$$

$$= -\tfrac{1}{2}t^2\ln t + \tfrac{1}{4}t^2 + C$$

$$= -\tfrac{1}{2}\cos^2 x\ln(\cos x) + \tfrac{1}{4}\cos^2 x + C$$

13. Let $x = \sec\theta$. Then

$$\int_1^2 \frac{\sqrt{x^2-1}}{x}\,dx = \int_0^{\pi/3} \frac{\tan\theta}{\sec\theta}\,\sec\theta\,\tan\theta\,d\theta = \int_0^{\pi/3} \tan^2\theta\,d\theta = \int_0^{\pi/3} (\sec^2\theta - 1)\,d\theta = \big[\tan\theta - \theta\big]_0^{\pi/3} = \sqrt{3} - \tfrac{\pi}{3}.$$

14. $$\int \frac{e^{2x}}{1+e^{4x}}\,dx = \int \frac{1}{1+u^2}\left(\tfrac{1}{2}\,du\right) \qquad \begin{bmatrix} u = e^{2x}, \\ du = 2e^{2x}\,dx \end{bmatrix}$$

$$= \tfrac{1}{2}\tan^{-1} u + C = \tfrac{1}{2}\tan^{-1} e^{2x} + C$$

15. Let $w = \sqrt[3]{x}$. Then $w^3 = x$ and $3w^2\,dw = dx$, so $\int e^{\sqrt[3]{x}}\,dx = \int e^w \cdot 3w^2\,dw = 3I$. To evaluate I, let $u = w^2$, $dv = e^w\,dw \;\Rightarrow\; du = 2w\,dw$, $v = e^w$, so $I = \int w^2e^w\,dw = w^2e^w - \int 2we^w\,dw$. Now let $U = w$, $dV = e^w\,dw \;\Rightarrow\; dU = dw$, $V = e^w$. Thus, $I = w^2e^w - 2\big[we^w - \int e^w\,dw\big] = w^2e^w - 2we^w + 2e^w + C_1$, and hence

$3I = 3e^w(w^2 - 2w + 2) + C = 3e^{\sqrt[3]{x}}(x^{2/3} - 2x^{1/3} + 2) + C$.

16. $$\int \frac{x^2+2}{x+2}\,dx = \int \left(x - 2 + \frac{6}{x+2}\right)dx = \tfrac{1}{2}x^2 - 2x + 6\ln|x+2| + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. Integrate by parts with $u = \tan^{-1} x$, $dv = x^2\,dx$, so that $du = \dfrac{1}{1+x^2}\,dx$, $v = \dfrac{1}{3}x^3$. Then

$$\int x^2 \tan^{-1}x\,dx = \frac{1}{3}x^3\tan^{-1}x - \frac{1}{3}\int \frac{x^3}{1+x^2}\,dx = \frac{1}{3}x^3\tan^{-1}x - \frac{1}{3}\cdot\frac{1}{2}\int \frac{y-1}{y}\,dy \qquad \begin{bmatrix} y = 1+x^2, \\ dy = 2x\,dx \end{bmatrix}$$

$$= \frac{1}{3}x^3\tan^{-1}x - \frac{1}{6}\int\left(1-\frac{1}{y}\right)dy = \frac{1}{3}x^3\tan^{-1}x - \frac{1}{6}\Big(y - \ln|y|\Big) + C_1$$

$$= \tfrac{1}{3}x^3\tan^{-1}x - \tfrac{1}{6}\big[1 + x^2 - \ln(1+x^2)\big] + C_1$$

$$= \tfrac{1}{6}[2x^3\tan^{-1}x - x^2 + \ln(1+x^2)] + C, \text{ where } C = C_1 - \tfrac{1}{6}$$

18. Let $u = x+1$ so that $u+1 = x+2$ and $du = dx$. Thus,

$$\int (x+2)^2(x+1)^{20}\,dx = \int (u+1)^2u^{20}\,du = \int (u^2+2u+1)\,u^{20}\,du = \int (u^{22}+2u^{21}+u^{20})\,du$$

$$= \tfrac{1}{23}u^{23} + \tfrac{2}{22}u^{22} + \tfrac{1}{21}u^{21} + C = \tfrac{1}{23}(x+1)^{23} + \tfrac{1}{11}(x+1)^{22} + \tfrac{1}{21}(x+1)^{21} + C$$

19. $\dfrac{x-1}{x^2+2x} = \dfrac{x-1}{x(x+2)} = \dfrac{A}{x} + \dfrac{B}{x+2} \quad\Rightarrow\quad x-1 = A(x+2) + Bx$. Set $x = -2$ to get $-3 = -2B$, so $B = \frac{3}{2}$. Set $x = 0$ to get $-1 = 2A$, so $A = -\frac{1}{2}$. Thus, $\displaystyle\int \frac{x-1}{x^2+2x}\,dx = \int\left(\frac{-\frac{1}{2}}{x} + \frac{\frac{3}{2}}{x+2}\right)dx = -\tfrac{1}{2}\ln|x| + \tfrac{3}{2}\ln|x+2| + C$.

20. $$\int \frac{\sec^6\theta}{\tan^2\theta}\,d\theta = \int \frac{(\tan^2\theta+1)^2\sec^2\theta}{\tan^2\theta}\,d\theta \quad \begin{bmatrix} u = \tan\theta, \\ du = -\sec^2\theta\,d\theta \end{bmatrix} \quad = \int \frac{(u^2+1)^2}{u^2}\,du = \int \frac{u^4+2u^2+1}{u^2}\,du$$

$$= \int\left(u^2 + 2 + \frac{1}{u^2}\right)du = \frac{u^3}{3} + 2u - \frac{1}{u} + C = \tfrac{1}{3}\tan^3\theta + 2\tan\theta - \cot\theta + C$$

21. $$\int x\cosh x\,dx = x\sinh x - \int \sinh x\,dx \qquad \begin{bmatrix} u = x, & dv = \cosh x\,dx \\ du = dx, & v = \sinh x \end{bmatrix}$$

$$= x\sinh x - \cosh x + C$$

22. $\dfrac{x^2+8x-3}{x^3+3x^2} = \dfrac{x^2+8x-3}{x^2(x+3)} = \dfrac{A}{x} + \dfrac{B}{x^2} + \dfrac{C}{x+3} \quad\Rightarrow\quad x^2+8x-3 = Ax(x+3) + B(x+3) + Cx^2$.

Taking $x = 0$, we get $-3 = 3B$, so $B = -1$. Taking $x = -3$, we get $-18 = 9C$, so $C = -2$.

Taking $x = 1$, we get $6 = 4A + 4B + C = 4A - 4 - 2$, so $4A = 12$ and $A = 3$. Now

$$\int \frac{x^2+8x-3}{x^3+3x^2}\,dx = \int\left(\frac{3}{x} - \frac{1}{x^2} - \frac{2}{x+3}\right)dx = 3\ln|x| + \frac{1}{x} - 2\ln|x+3| + C.$$

23. $$\int \frac{dx}{\sqrt{x^2-4x}} = \int \frac{dx}{\sqrt{(x^2-4x+4)-4}} = \int \frac{dx}{\sqrt{(x-2)^2-2^2}}$$

$$= \int \frac{2\sec\theta\,\tan\theta\,d\theta}{2\tan\theta} \qquad \begin{bmatrix} x-2 = 2\sec\theta, \\ dx = 2\sec\theta\,\tan\theta\,d\theta \end{bmatrix}$$

$$= \int \sec\theta\,d\theta = \ln|\sec\theta + \tan\theta| + C_1$$

$$= \ln\left|\frac{x-2}{2} + \frac{\sqrt{x^2-4x}}{2}\right| + C_1$$

$$= \ln\left|x - 2 + \sqrt{x^2-4x}\,\right| + C, \text{ where } C = C_1 - \ln 2$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $\displaystyle\int \frac{2^{\sqrt{x}}}{\sqrt{x}}\,dx = \int 2^u\,(2\,du) \quad \begin{bmatrix} u=\sqrt{x}, \\ du = 1/(2\sqrt{x}\,)\,dx \end{bmatrix} \quad = 2\cdot\frac{2^u}{\ln 2} + C = \frac{2^{\sqrt{x}+1}}{\ln 2} + C$

25. $\displaystyle\int \frac{x+1}{9x^2+6x+5}\,dx = \int \frac{x+1}{(9x^2+6x+1)+4}\,dx = \int \frac{x+1}{(3x+1)^2+4}\,dx \qquad \begin{bmatrix} u=3x+1, \\ du=3\,dx \end{bmatrix}$

$$= \int \frac{\left[\frac{1}{3}(u-1)\right]+1}{u^2+4}\left(\frac{1}{3}\,du\right) = \frac{1}{3}\cdot\frac{1}{3}\int\frac{(u-1)+3}{u^2+4}\,du$$

$$= \frac{1}{9}\int\frac{u}{u^2+4}\,du + \frac{1}{9}\int\frac{2}{u^2+2^2}\,du = \frac{1}{9}\cdot\frac{1}{2}\ln(u^2+4) + \frac{2}{9}\cdot\frac{1}{2}\tan^{-1}\left(\frac{1}{2}u\right) + C$$

$$= \tfrac{1}{18}\ln(9x^2+6x+5) + \tfrac{1}{9}\tan^{-1}\left[\tfrac{1}{2}(3x+1)\right] + C$$

26. $\int \tan^5\theta\,\sec^3\theta\,d\theta = \int \tan^4\theta\,\sec^2\theta\,\sec\theta\,\tan\theta\,d\theta = \int(\sec^2\theta-1)^2\sec^2\theta\,\sec\theta\,\tan\theta\,d\theta \qquad \begin{bmatrix} u=\sec\theta, \\ du=\sec\theta\,\tan\theta\,d\theta \end{bmatrix}$

$$= \int(u^2-1)^2u^2\,du = \int(u^6-2u^4+u^2)\,du$$

$$= \tfrac{1}{7}u^7 - \tfrac{2}{5}u^5 + \tfrac{1}{3}u^3 + C = \tfrac{1}{7}\sec^7\theta - \tfrac{2}{5}\sec^5\theta + \tfrac{1}{3}\sec^3\theta + C$$

27. $\sqrt{x^2-2x+2} = \sqrt{x^2-2x+1+1} = \sqrt{(x-1)^2+1}$. Since this is a sum of squares, we try the substitution $x-1=\tan\theta$, where $-\pi/2<\theta<\pi/2$. Then $dx=\sec^2\theta\,d\theta$ and $\sqrt{(x-1)^2+1} = \sqrt{\tan^2\theta+1} = \sqrt{\sec^2\theta} = |\sec\theta| = \sec\theta$. Also, $x=0 \;\Rightarrow\; \theta=-\pi/4$ and $x=2 \;\Rightarrow\; \theta=\pi/4$. Thus,

$$\int_0^2 \sqrt{x^2-2x+2}\,dx = \int_{-\pi/4}^{\pi/4}\sec\theta\,(\sec^2\theta\,d\theta) = \int_{-\pi/4}^{\pi/4}\sec^3\theta\,d\theta$$

$$= \frac{1}{2}\Big[\sec\theta\,\tan\theta + \ln|\sec\theta+\tan\theta|\,\Big]_{-\pi/4}^{\pi/4} \qquad \text{[by Example 8 in Section 7.2]}$$

$$= \frac{1}{2}\left[\left(\sqrt{2}+\ln\left(\sqrt{2}+1\right)\right) - \left(-\sqrt{2}+\ln\left(\sqrt{2}-1\right)\right)\right] = \frac{1}{2}\left[2\sqrt{2}+\ln\left(\frac{\sqrt{2}+1}{\sqrt{2}-1}\right)\right]$$

$$= \frac{1}{2}\left[2\sqrt{2}+\ln\left(\frac{\sqrt{2}+1}{\sqrt{2}-1}\cdot\frac{\sqrt{2}+1}{\sqrt{2}+1}\right)\right] = \frac{1}{2}\left[2\sqrt{2}+\ln\left(\sqrt{2}+1\right)^2\right]$$

$$= \tfrac{1}{2}\left[2\sqrt{2}+2\ln\left(\sqrt{2}+1\right)\right] = \sqrt{2}+\ln\left(\sqrt{2}+1\right)$$

28. $\int \cos\sqrt{t}\,dt = \int 2x\,\cos x\,dx \qquad \begin{bmatrix} x=\sqrt{t}, \\ x^2=t, \quad 2x\,dx=dt \end{bmatrix}$

$$= 2x\,\sin x - \int 2\sin x\,dx \qquad \begin{bmatrix} u=x, & dv=\cos x\,dx \\ du=dx, & v=\sin x \end{bmatrix}$$

$$= 2x\,\sin x + 2\cos x + C = 2\sqrt{t}\,\sin\sqrt{t} + 2\cos\sqrt{t} + C$$

29. Let $x=\tan\theta$, so that $dx=\sec^2\theta\,d\theta$. Then

$$\int\frac{dx}{x\sqrt{x^2+1}} = \int\frac{\sec^2\theta\,d\theta}{\tan\theta\,\sec\theta} = \int\frac{\sec\theta}{\tan\theta}\,d\theta$$

$$= \int\csc\theta\,d\theta = \ln|\csc\theta-\cot\theta| + C$$

$$= \ln\left|\frac{\sqrt{x^2+1}}{x} - \frac{1}{x}\right| + C = \ln\left|\frac{\sqrt{x^2+1}-1}{x}\right| + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. Let $u = \cos x$, $dv = e^x\,dx \quad\Rightarrow\quad du = -\sin x\,dx$, $v = e^x$: $(*)$ $I = \int e^x \cos x\,dx = e^x \cos x + \int e^x \sin x\,dx$.

To integrate $\int e^x \sin x\,dx$, let $U = \sin x$, $dV = e^x\,dx \quad\Rightarrow\quad dU = \cos x\,dx$, $V = e^x$. Then

$\int e^x \sin x\,dx = e^x \sin x - \int e^x \cos x\,dx = e^x \sin x - I$. By substitution in $(*)$, $I = e^x \cos x + e^x \sin x - I \quad\Rightarrow$

$2I = e^x(\cos x + \sin x) \quad\Rightarrow\quad I = \frac{1}{2}e^x(\cos x + \sin x) + C$.

31. Let $u = \sqrt{1+x^2}$, so that $du = \dfrac{x}{\sqrt{1+x^2}}\,dx$. Thus,

$$\int \frac{x\sin\left(\sqrt{1+x^2}\right)}{\sqrt{1+x^2}}\,dx = \int \sin u\,du = -\cos u + C = -\cos\left(\sqrt{1+x^2}\right) + C.$$

32. Let $u = x^{1/4} \quad\Rightarrow\quad x = u^4$, so that $dx = 4u^3\,du$. Thus,

$$\int \frac{1}{x^{1/2}+x^{1/4}}\,dx = \int \frac{4u^3}{u^2+u}\,du = 4\int \frac{u^2}{u+1}\,du = 4\int \left(u - 1 + \frac{1}{u+1}\right)du \qquad \text{[using long division]}$$

$$= 4\left(\tfrac{1}{2}u^2 - u + \ln|u+1|\right) + C = 2x^{1/2} - 4x^{1/4} + 4\ln(x^{1/4}+1) + C$$

33. $\dfrac{3x^3 - x^2 + 6x - 4}{(x^2+1)(x^2+2)} = \dfrac{Ax+B}{x^2+1} + \dfrac{Cx+D}{x^2+2} \quad\Rightarrow\quad 3x^3 - x^2 + 6x - 4 = (Ax+B)\left(x^2+2\right) + (Cx+D)\left(x^2+1\right)$.

Equating the coefficients gives $A + C = 3$, $B + D = -1$, $2A + C = 6$, and $2B + D = -4 \quad\Rightarrow$

$A = 3$, $C = 0$, $B = -3$, and $D = 2$. Now

$$\int \frac{3x^3 - x^2 + 6x - 4}{(x^2+1)(x^2+2)}\,dx = 3\int \frac{x-1}{x^2+1}\,dx + 2\int \frac{dx}{x^2+2} = \frac{3}{2}\ln\left(x^2+1\right) - 3\tan^{-1}x + \sqrt{2}\tan^{-1}\left(\frac{x}{\sqrt{2}}\right) + C.$$

34. $\int x\sin x\,\cos x\,dx = \int \frac{1}{2}x\sin 2x\,dx \qquad \begin{bmatrix} u = \frac{1}{2}x, & dv = \sin 2x\,dx, \\ du = \frac{1}{2}\,dx & v = -\frac{1}{2}\cos 2x \end{bmatrix}$

$$= -\tfrac{1}{4}x\cos 2x + \int \tfrac{1}{4}\cos 2x\,dx = -\tfrac{1}{4}x\cos 2x + \tfrac{1}{8}\sin 2x + C$$

35. $\int_0^{\pi/2} \cos^3 x\,\sin 2x\,dx = \int_0^{\pi/2} \cos^3 x\,(2\sin x\,\cos x)\,dx = \int_0^{\pi/2} 2\cos^4 x\,\sin x\,dx = \left[-\frac{2}{5}\cos^5 x\right]_0^{\pi/2} = \frac{2}{5}$

36. Let $u = \sqrt[3]{x}$. Then $x = u^3$, $dx = 3u^2\,du \quad\Rightarrow$

$$\int \frac{\sqrt[3]{x}+1}{\sqrt[3]{x}-1}\,dx = \int \frac{u+1}{u-1}\,3u^2 du = 3\int \left(u^2 + 2u + 2 + \frac{2}{u-1}\right)du$$

$$= u^3 + 3u^2 + 6u + 6\ln|u-1| + C = x + 3x^{2/3} + 6\sqrt[3]{x} + 6\ln|\sqrt[3]{x} - 1| + C$$

37. The integrand is an odd function, so $\displaystyle\int_{-3}^{3} \frac{x}{1+|x|}\,dx = 0$ [by 5.5.7(b)].

38. Let $u = e^{-x}$, $du = -e^{-x}\,dx$. Then

$$\int \frac{dx}{e^x\sqrt{1-e^{-2x}}} = \int \frac{e^{-x}\,dx}{\sqrt{1-(e^{-x})^2}} = \int \frac{-du}{\sqrt{1-u^2}} = -\sin^{-1}u + C = -\sin^{-1}(e^{-x}) + C.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. Let $u = \sqrt{e^x - 1}$. Then $u^2 = e^x - 1$ and $2u\,du = e^x\,dx$. Also, $e^x + 8 = u^2 + 9$. Thus,

$$\int_0^{\ln 10} \frac{e^x\sqrt{e^x-1}}{e^x+8}\,dx = \int_0^3 \frac{u\cdot 2u\,du}{u^2+9} = 2\int_0^3 \frac{u^2}{u^2+9}\,du = 2\int_0^3 \left(1 - \frac{9}{u^2+9}\right)du$$

$$= 2\left[u - \frac{9}{3}\tan^{-1}\left(\frac{u}{3}\right)\right]_0^3 = 2\left[(3 - 3\tan^{-1}1) - 0\right] = 2\left(3 - 3\cdot\frac{\pi}{4}\right) = 6 - \frac{3\pi}{2}$$

40. $$\int_0^{\pi/4} \frac{x\sin x}{\cos^3 x}\,dx = \int_0^{\pi/4} x\tan x\,\sec^2 x\,dx \qquad \begin{bmatrix} u = x, & dv = \tan x\,\sec^2 x\,dx, \\ du = dx & v = \frac{1}{2}\tan^2 x \end{bmatrix}$$

$$= \left[\frac{x}{2}\tan^2 x\right]_0^{\pi/4} - \frac{1}{2}\int_0^{\pi/4} \tan^2 x\,dx = \frac{\pi}{8}\cdot 1^2 - 0 - \frac{1}{2}\int_0^{\pi/4}(\sec^2 x - 1)\,dx$$

$$= \frac{\pi}{8} - \frac{1}{2}\left[\tan x - x\right]_0^{\pi/4} = \frac{\pi}{8} - \frac{1}{2}\left(1 - \frac{\pi}{4}\right) = \frac{\pi}{4} - \frac{1}{2}$$

41. Let $x = 2\sin\theta \;\Rightarrow\; \left(4 - x^2\right)^{3/2} = (2\cos\theta)^3$, $dx = 2\cos\theta\,d\theta$, so

$$\int \frac{x^2}{\left(4-x^2\right)^{3/2}}\,dx = \int \frac{4\sin^2\theta}{8\cos^3\theta}\,2\cos\theta\,d\theta = \int \tan^2\theta\,d\theta = \int \left(\sec^2\theta - 1\right)d\theta$$

$$= \tan\theta - \theta + C = \frac{x}{\sqrt{4-x^2}} - \sin^{-1}\left(\frac{x}{2}\right) + C$$

42. Integrate by parts twice, first with $u = (\arcsin x)^2$, $dv = dx$:

$$I = \int (\arcsin x)^2\,dx = x(\arcsin x)^2 - \int 2x\arcsin x\left(\frac{dx}{\sqrt{1-x^2}}\right)$$

Now let $U = \arcsin x$, $dV = \dfrac{x}{\sqrt{1-x^2}}\,dx \;\Rightarrow\; dU = \dfrac{1}{\sqrt{1-x^2}}\,dx$, $V = -\sqrt{1-x^2}$. So

$$I = x(\arcsin x)^2 - 2\left[\arcsin x\left(-\sqrt{1-x^2}\right) + \int dx\right] = x(\arcsin x)^2 + 2\sqrt{1-x^2}\arcsin x - 2x + C$$

43. $$\int \frac{1}{\sqrt{x + x^{3/2}}}\,dx = \int \frac{dx}{\sqrt{x\left(1+\sqrt{x}\right)}} = \int \frac{dx}{\sqrt{x}\sqrt{1+\sqrt{x}}} \qquad \begin{bmatrix} u = 1 + \sqrt{x}, \\ du = \frac{dx}{2\sqrt{x}} \end{bmatrix} \quad = \int \frac{2\,du}{\sqrt{u}} = \int 2u^{-1/2}\,du$$

$$= 4\sqrt{u} + C = 4\sqrt{1+\sqrt{x}} + C$$

44. $$\int \frac{1-\tan\theta}{1+\tan\theta}\,d\theta = \int \frac{\dfrac{\cos\theta}{\cos\theta} - \dfrac{\sin\theta}{\cos\theta}}{\dfrac{\cos\theta}{\cos\theta} + \dfrac{\sin\theta}{\cos\theta}}\,d\theta = \int \frac{\cos\theta - \sin\theta}{\cos\theta + \sin\theta}\,d\theta = \ln|\cos\theta + \sin\theta| + C$$

45. $$\int (\cos x + \sin x)^2\cos 2x\,dx = \int \left(\cos^2 x + 2\sin x\cos x + \sin^2 x\right)\cos 2x\,dx = \int (1 + \sin 2x)\cos 2x\,dx$$

$$= \int \cos 2x\,dx + \tfrac{1}{2}\int \sin 4x\,dx = \tfrac{1}{2}\sin 2x - \tfrac{1}{8}\cos 4x + C$$

Or: $$\int (\cos x + \sin x)^2\cos 2x\,dx = \int (\cos x + \sin x)^2(\cos^2 x - \sin^2 x)\,dx$$

$$= \int (\cos x + \sin x)^3(\cos x - \sin x)\,dx = \tfrac{1}{4}(\cos x + \sin x)^4 + C_1$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. Let $u = \sin(x^2)$, so that $du = 2x\cos(x^2)\,dx$ and $\cos^2(x^2) = 1 - u^2$. Thus,

$$\begin{aligned}\int x\cos^3(x^2)\sqrt{\sin(x^2)}\,dx &= \int x\cos(x^2)\cos^2(x^2)\sqrt{\sin(x^2)}\,dx = \frac{1}{2}\int \left(1-u^2\right)\sqrt{u}\,du \\ &= \frac{1}{2}\int (u^{1/2} - u^{5/2})\,du = \frac{1}{2}\left(\frac{2}{3}u^{3/2} - \frac{2}{7}u^{7/2}\right) + C \\ &= \tfrac{1}{3}\sqrt{\sin^3(x^2)} - \tfrac{1}{7}\sqrt{\sin^7(x^2)} + C\end{aligned}$$

47. We'll integrate $I = \displaystyle\int \frac{xe^{2x}}{(1+2x)^2}\,dx$ by parts with $u = xe^{2x}$ and $dv = \dfrac{dx}{(1+2x)^2}$. Then $du = (x\cdot 2e^{2x} + e^{2x}\cdot 1)\,dx$ and $v = -\dfrac{1}{2}\cdot\dfrac{1}{1+2x}$, so

$$I = -\frac{1}{2}\cdot\frac{xe^{2x}}{1+2x} - \int\left[-\frac{1}{2}\cdot\frac{e^{2x}(2x+1)}{1+2x}\right]dx = -\frac{xe^{2x}}{4x+2} + \frac{1}{2}\cdot\frac{1}{2}e^{2x} + C = e^{2x}\left(\frac{1}{4} - \frac{x}{4x+2}\right) + C$$

Thus, $\displaystyle\int_0^{1/2} \frac{xe^{2x}}{(1+2x)^2}\,dx = \left[e^{2x}\left(\frac{1}{4} - \frac{x}{4x+2}\right)\right]_0^{1/2} = e\left(\frac{1}{4} - \frac{1}{8}\right) - 1\left(\frac{1}{4} - 0\right) = \frac{1}{8}e - \frac{1}{4}.$

48. $$\begin{aligned}\int_{\pi/4}^{\pi/3} \frac{\sqrt{\tan\theta}}{\sin 2\theta}\,d\theta &= \int_{\pi/4}^{\pi/3} \frac{\sqrt{\frac{\sin\theta}{\cos\theta}}}{2\sin\theta\cos\theta}\,d\theta = \int_{\pi/4}^{\pi/3} \frac{1}{2}(\sin\theta)^{-1/2}(\cos\theta)^{-3/2}\,d\theta = \int_{\pi/4}^{\pi/3} \frac{1}{2}\left(\frac{\sin\theta}{\cos\theta}\right)^{-1/2}(\cos\theta)^{-2}d\theta \\ &= \int_{\pi/4}^{\pi/3} \frac{1}{2}(\tan\theta)^{-1/2}\sec^2\theta\,d\theta = \left[\sqrt{\tan\theta}\right]_{\pi/4}^{\pi/3} = \sqrt{\sqrt{3}} - \sqrt{1} = \sqrt[4]{3} - 1\end{aligned}$$

49. Let $u = \sqrt{e^x - 4}$, so that $e^x = u^2 + 4$ and $2u\,du = e^x\,dx = (u^2+4)\,dx \quad\Leftrightarrow\quad \dfrac{2u}{u^2+4}\,du = dx$. Thus,

$$\int \frac{1}{\sqrt{e^x-4}}\,dx = \int \frac{1}{u}\cdot\frac{2u}{u^2+4}\,du = 2\int\frac{1}{u^2+4}\,du = 2\left(\frac{1}{2}\tan^{-1}\left(\frac{u}{2}\right)\right) + C = \tan^{-1}\left(\frac{\sqrt{e^x-4}}{2}\right) + C.$$

50. Let $y = \sqrt{1+x^2}$, so that $y^2 = 1 + x^2$ and $2y\,dy = 2x\,dx \quad\Rightarrow\quad dx = \dfrac{y}{x}\,dy$. Thus,

$$\begin{aligned}\int x\sin\left(\sqrt{1+x^2}\right)dx &= \int y\sin y\,dy \qquad \begin{bmatrix} u = y,\ dv = \sin y\,dy \\ du = dy,\ v = -\cos y\end{bmatrix} \\ &= -y\cos y + \int \cos y\,dy = -y\cos y + \sin y + C \\ &= -\sqrt{1+x^2}\cos\sqrt{1+x^2} + \sin\sqrt{1+x^2} + C\end{aligned}$$

51. $$\begin{aligned}\int_1^\infty \frac{1}{(2x+1)^3}\,dx &= \lim_{t\to\infty}\int_1^t \frac{1}{(2x+1)^3}\,dx = \lim_{t\to\infty}\int_1^t \tfrac{1}{2}(2x+1)^{-3}\,2\,dx = \lim_{t\to\infty}\left[-\frac{1}{4(2x+1)^2}\right]_1^t \\ &= -\frac{1}{4}\lim_{t\to\infty}\left[\frac{1}{(2t+1)^2} - \frac{1}{9}\right] = -\frac{1}{4}\left(0 - \frac{1}{9}\right) = \frac{1}{36}\end{aligned}$$

52. $$\begin{aligned}\int_1^\infty \frac{\ln x}{x^4}\,dx &= \lim_{t\to\infty}\int_1^t \frac{\ln x}{x^4}\,dx \qquad \begin{bmatrix} u = \ln x, & dv = dx/x^4, \\ du = dx/x & v = -1/(3x^3)\end{bmatrix} \\ &= \lim_{t\to\infty}\left[-\frac{\ln x}{3x^3}\right]_1^t + \int_1^t \frac{1}{3x^4}\,dx = \lim_{t\to\infty}\left(-\frac{\ln t}{3t^3} + 0 + \left[\frac{-1}{9x^3}\right]_1^t\right) \overset{\text{H}}{=} \lim_{t\to\infty}\left(-\frac{1}{9t^3} + \left[\frac{-1}{9t^3} + \frac{1}{9}\right]\right) \\ &= 0 + 0 + \tfrac{1}{9} = \tfrac{1}{9}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

53. $\displaystyle\int \frac{dx}{x\ln x} \quad \begin{bmatrix} u=\ln x, \\ du = dx/x \end{bmatrix} \quad = \int \frac{du}{u} = \ln|u| + C = \ln|\ln x| + C$, so

$$\int_2^\infty \frac{dx}{x\ln x} = \lim_{t\to\infty}\int_2^t \frac{dx}{x\ln x} = \lim_{t\to\infty}\Big[\ln|\ln x|\Big]_2^t = \lim_{t\to\infty}[\ln(\ln t) - \ln(\ln 2)] = \infty,$$ so the integral is divergent.

54. Let $u = \sqrt{y-2}$. Then $y = u^2 + 2$ and $dy = 2u\,du$, so

$$\int \frac{y\,dy}{\sqrt{y-2}} = \int \frac{(u^2+2)2u\,du}{u} = 2\int (u^2+2)\,du = 2\big[\tfrac{1}{3}u^3 + 2u\big] + C$$

Thus, $\displaystyle\int_2^6 \frac{y}{\sqrt{y-2}}\,dy = \lim_{t\to 2^+}\int_t^6 \frac{y}{\sqrt{y-2}}\,dy = \lim_{t\to 2^+}\Big[\tfrac{2}{3}(y-2)^{3/2} + 4\sqrt{y-2}\Big]_t^6$

$$= \lim_{t\to 2^+}\Big[\tfrac{16}{3} + 8 - \tfrac{2}{3}(t-2)^{3/2} - 4\sqrt{t-2}\Big] = \tfrac{40}{3}.$$

55. $\displaystyle\int_0^4 \frac{\ln x}{\sqrt{x}}\,dx = \lim_{t\to 0^+}\int_t^4 \frac{\ln x}{\sqrt{x}}\,dx \overset{\star}{=} \lim_{t\to 0^+}\Big[2\sqrt{x}\,\ln x - 4\sqrt{x}\Big]_t^4$

$$= \lim_{t\to 0^+}\big[(2\cdot 2\ln 4 - 4\cdot 2) - \big(2\sqrt{t}\,\ln t - 4\sqrt{t}\big)\big] \overset{\star\star}{=} (4\ln 4 - 8) - (0-0) = 4\ln 4 - 8$$

($\star$) Let $u = \ln x$, $dv = \dfrac{1}{\sqrt{x}}\,dx \quad\Rightarrow\quad du = \dfrac{1}{x}\,dx$, $v = 2\sqrt{x}$. Then

$$\int \frac{\ln x}{\sqrt{x}}\,dx = 2\sqrt{x}\,\ln x - 2\int \frac{dx}{\sqrt{x}} = 2\sqrt{x}\,\ln x - 4\sqrt{x} + C$$

($\star\star$) $$\lim_{t\to 0^+}\big(2\sqrt{t}\,\ln t\big) = \lim_{t\to 0^+}\frac{2\ln t}{t^{-1/2}} \overset{\text{H}}{=} \lim_{t\to 0^+}\frac{2/t}{-\frac{1}{2}t^{-3/2}} = \lim_{t\to 0^+}\big(-4\sqrt{t}\big) = 0$$

56. Note that $f(x) = 1/(2-3x)$ has an infinite discontinuity at $x = \frac{2}{3}$. Now

$$\int_0^{2/3} \frac{1}{2-3x}\,dx = \lim_{t\to(2/3)^-}\int_0^t \frac{1}{2-3x}\,dx = \lim_{t\to(2/3)^-}\Big[-\tfrac{1}{3}\ln|2-3x|\Big]_0^t = -\tfrac{1}{3}\lim_{t\to(2/3)^-}\Big[\ln|2-3t| - \ln 2\Big] = \infty$$

Since $\displaystyle\int_0^{2/3} \frac{1}{2-3x}\,dx$ diverges, so does $\displaystyle\int_0^1 \frac{1}{2-3x}\,dx$.

57. $\displaystyle\int_0^1 \frac{x-1}{\sqrt{x}}\,dx = \lim_{t\to 0^+}\int_t^1 \left(\frac{x}{\sqrt{x}} - \frac{1}{\sqrt{x}}\right)dx = \lim_{t\to 0^+}\int_t^1 (x^{1/2} - x^{-1/2})\,dx = \lim_{t\to 0^+}\Big[\tfrac{2}{3}x^{3/2} - 2x^{1/2}\Big]_t^1$

$$= \lim_{t\to 0^+}\Big[\big(\tfrac{2}{3} - 2\big) - \big(\tfrac{2}{3}t^{3/2} - 2t^{1/2}\big)\Big] = -\tfrac{4}{3} - 0 = -\tfrac{4}{3}$$

58. $\displaystyle I = \int_{-1}^1 \frac{dx}{x^2-2x} = \int_{-1}^1 \frac{dx}{x(x-2)} = \int_{-1}^0 \frac{dx}{x(x-2)} + \int_0^1 \frac{dx}{x(x-2)} = I_1 + I_2$. Now

$\dfrac{1}{x(x-2)} = \dfrac{A}{x} + \dfrac{B}{x-2} \quad\Rightarrow\quad 1 = A(x-2) + Bx$. Set $x = 2$ to get $1 = 2B$, so $B = \frac{1}{2}$. Set $x = 0$ to get $1 = -2A$, $A = -\frac{1}{2}$. Thus,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$I_2 = \lim_{t\to 0^+} \int_t^1 \left(\frac{-\frac{1}{2}}{x} + \frac{\frac{1}{2}}{x-2} \right) dx = \lim_{t\to 0^+} \left[-\tfrac{1}{2}\ln|x| + \tfrac{1}{2}\ln|x-2| \right]_t^1 = \lim_{t\to 0^+} \left[(0+0) - \left(-\tfrac{1}{2}\ln t + \tfrac{1}{2}\ln|t-2| \right) \right]$$
$$= -\tfrac{1}{2}\ln 2 + \tfrac{1}{2}\lim_{t\to 0^+} \ln t = -\infty.$$

Since I_2 diverges, I is divergent.

59. Let $u = 2x+1$. Then

$$\int_{-\infty}^{\infty} \frac{dx}{4x^2+4x+5} = \int_{-\infty}^{\infty} \frac{\frac{1}{2}\,du}{u^2+4} = \tfrac{1}{2}\int_{-\infty}^{0} \frac{du}{u^2+4} + \tfrac{1}{2}\int_0^{\infty} \frac{du}{u^2+4}$$
$$= \tfrac{1}{2}\lim_{t\to-\infty} \left[\tfrac{1}{2}\tan^{-1}\left(\tfrac{1}{2}u\right) \right]_t^0 + \tfrac{1}{2}\lim_{t\to\infty} \left[\tfrac{1}{2}\tan^{-1}\left(\tfrac{1}{2}u\right) \right]_0^t = \tfrac{1}{4}\left[0 - \left(-\tfrac{\pi}{2}\right)\right] + \tfrac{1}{4}\left[\tfrac{\pi}{2} - 0\right] = \tfrac{\pi}{4}.$$

60. $\displaystyle\int_1^{\infty} \frac{\tan^{-1} x}{x^2}\,dx = \lim_{t\to\infty} \int_1^t \frac{\tan^{-1} x}{x^2}\,dx$. Integrate by parts:

$$\int \frac{\tan^{-1} x}{x^2}\,dx = \frac{-\tan^{-1} x}{x} + \int \frac{1}{x}\frac{dx}{1+x^2} = \frac{-\tan^{-1} x}{x} + \int \left[\frac{1}{x} - \frac{x}{x^2+1} \right] dx$$
$$= \frac{-\tan^{-1} x}{x} + \ln|x| - \tfrac{1}{2}\ln(x^2+1) + C = \frac{-\tan^{-1} x}{x} + \frac{1}{2}\ln\frac{x^2}{x^2+1} + C$$

Thus,

$$\int_1^{\infty} \frac{\tan^{-1} x}{x^2}\,dx = \lim_{t\to\infty} \left[-\frac{\tan^{-1} x}{x} + \frac{1}{2}\ln\frac{x^2}{x^2+1} \right]_1^t = \lim_{t\to\infty} \left[-\frac{\tan^{-1} t}{t} + \frac{1}{2}\ln\frac{t^2}{t^2+1} + \frac{\pi}{4} - \frac{1}{2}\ln\frac{1}{2} \right]$$
$$= 0 + \tfrac{1}{2}\ln 1 + \tfrac{\pi}{4} + \tfrac{1}{2}\ln 2 = \tfrac{\pi}{4} + \tfrac{1}{2}\ln 2$$

61. We first make the substitution $t = x+1$, so $\ln(x^2+2x+2) = \ln\left[(x+1)^2+1\right] = \ln(t^2+1)$. Then we use parts with $u = \ln(t^2+1)$, $dv = dt$:

$$\int \ln(t^2+1)\,dt = t\,\ln(t^2+1) - \int \frac{t(2t)\,dt}{t^2+1} = t\,\ln(t^2+1) - 2\int \frac{t^2\,dt}{t^2+1} = t\,\ln(t^2+1) - 2\int \left(1 - \frac{1}{t^2+1}\right) dt$$
$$= t\,\ln(t^2+1) - 2t + 2\arctan t + C$$
$$= (x+1)\,\ln(x^2+2x+2) - 2x + 2\arctan(x+1) + K, \text{ where } K = C - 2$$

[Alternatively, we could have integrated by parts immediately with $u = \ln(x^2+2x+2)$.] Notice from the graph that $f = 0$ where F has a horizontal tangent. Also, F is always increasing, and $f \geq 0$.

62. Let $u = x^2+1$. Then $x^2 = u-1$ and $x\,dx = \frac{1}{2}\,du$, so

$$\int \frac{x^3}{\sqrt{x^2+1}}\,dx = \int \frac{(u-1)}{\sqrt{u}}\left(\tfrac{1}{2}\,du\right) = \frac{1}{2}\int \left(u^{1/2} - u^{-1/2}\right) du$$
$$= \tfrac{1}{2}\left(\tfrac{2}{3}u^{3/2} - 2u^{1/2}\right) + C = \tfrac{1}{3}(x^2+1)^{3/2} - (x^2+1)^{1/2} + C$$
$$= \tfrac{1}{3}(x^2+1)^{1/2}\left[(x^2+1) - 3\right] + C = \tfrac{1}{3}\sqrt{x^2+1}\,(x^2-2) + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

63. From the graph, it seems as though $\int_0^{2\pi} \cos^2 x \, \sin^3 x \, dx$ is equal to 0.

To evaluate the integral, we write the integral as

$I = \int_0^{2\pi} \cos^2 x \, (1 - \cos^2 x) \, \sin x \, dx$ and let $u = \cos x \quad \Rightarrow$

$du = -\sin x \, dx$. Thus, $I = \int_1^1 u^2 (1 - u^2)(-du) = 0$.

64. (a) To evaluate $\int x^5 e^{-2x} \, dx$ by hand, we would integrate by parts repeatedly, always taking $dv = e^{-2x}$ and starting with $u = x^5$. Each time we would reduce the degree of the x-factor by 1.

(b) To evaluate the integral using tables, we would use Formula 97 (which is proved using integration by parts) until the exponent of x was reduced to 1, and then we would use Formula 96.

(c) $\int x^5 e^{-2x} \, dx = -\frac{1}{8} e^{-2x} \left(4x^5 + 10x^4 + 20x^3 + 30x^2 + 30x + 15\right) + C$

(d)

65. $\int \sqrt{4x^2 - 4x - 3} \, dx = \int \sqrt{(2x-1)^2 - 4} \, dx \quad \begin{bmatrix} u = 2x - 1, \\ du = 2\,dx \end{bmatrix} \quad = \int \sqrt{u^2 - 2^2} \left(\frac{1}{2} \, du\right)$

$$\overset{39}{=} \frac{1}{2}\left(\frac{u}{2}\sqrt{u^2 - 2^2} - \frac{2^2}{2} \ln\left|u + \sqrt{u^2 - 2^2}\right|\right) + C = \tfrac{1}{4} u \sqrt{u^2 - 4} - \ln\left|u + \sqrt{u^2 - 4}\right| + C$$

$$= \tfrac{1}{4}(2x - 1)\sqrt{4x^2 - 4x - 3} - \ln\left|2x - 1 + \sqrt{4x^2 - 4x - 3}\right| + C$$

66. $\int \csc^5 t \, dt \overset{78}{=} -\frac{1}{4} \cot t \, \csc^3 t + \frac{3}{4} \int \csc^3 t \, dt \overset{72}{=} -\frac{1}{4} \cot t \, \csc^3 t + \frac{3}{4}\left[-\frac{1}{2} \csc t \, \cot t + \frac{1}{2} \ln|\csc t - \cot t|\right] + C$

$$= -\tfrac{1}{4} \cot t \, \csc^3 t - \tfrac{3}{8} \csc t \, \cot t + \tfrac{3}{8} \ln|\csc t - \cot t| + C$$

67. Let $u = \sin x$, so that $du = \cos x \, dx$. Then

$$\int \cos x \sqrt{4 + \sin^2 x} \, dx = \int \sqrt{2^2 + u^2} \, du \overset{21}{=} \frac{u}{2}\sqrt{2^2 + u^2} + \frac{2^2}{2} \ln\left(u + \sqrt{2^2 + u^2}\right) + C$$

$$= \tfrac{1}{2} \sin x \sqrt{4 + \sin^2 x} + 2 \ln\left(\sin x + \sqrt{4 + \sin^2 x}\right) + C$$

68. Let $u = \sin x$. Then $du = \cos x \, dx$, so

$$\int \frac{\cot x \, dx}{\sqrt{1 + 2\sin x}} = \int \frac{du}{u\sqrt{1 + 2u}} \overset{\substack{57 \text{ with} \\ a=1,\, b=2}}{=} \ln\left|\frac{\sqrt{1 + 2u} - 1}{\sqrt{1 + 2u} + 1}\right| + C = \ln\left|\frac{\sqrt{1 + 2\sin x} - 1}{\sqrt{1 + 2\sin x} + 1}\right| + C$$

69. (a) $\dfrac{d}{du}\left[-\dfrac{1}{u}\sqrt{a^2 - u^2} - \sin^{-1}\left(\dfrac{u}{a}\right) + C\right] = \dfrac{1}{u^2}\sqrt{a^2 - u^2} + \dfrac{1}{\sqrt{a^2 - u^2}} - \dfrac{1}{\sqrt{1 - u^2/a^2}} \cdot \dfrac{1}{a}$

$$= \left(a^2 - u^2\right)^{-1/2}\left[\frac{1}{u^2}\left(a^2 - u^2\right) + 1 - 1\right] = \frac{\sqrt{a^2 - u^2}}{u^2}$$

(b) Let $u = a \sin\theta \quad \Rightarrow \quad du = a\cos\theta \, d\theta$, $a^2 - u^2 = a^2\left(1 - \sin^2\theta\right) = a^2 \cos^2\theta$.

$$\int \frac{\sqrt{a^2 - u^2}}{u^2} \, du = \int \frac{a^2 \cos^2\theta}{a^2 \sin^2\theta} \, d\theta = \int \frac{1 - \sin^2\theta}{\sin^2\theta} \, d\theta = \int \left(\csc^2\theta - 1\right) d\theta = -\cot\theta - \theta + C$$

$$= -\frac{\sqrt{a^2 - u^2}}{u} - \sin^{-1}\left(\frac{u}{a}\right) + C$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

70. Work backward, and use integration by parts with $U = u^{-(n-1)}$ and $dV = (a+bu)^{-1/2}\,du \quad \Rightarrow$

$dU = \dfrac{-(n-1)\,du}{u^n}$ and $V = \dfrac{2}{b}\sqrt{a+bu}$, to get

$$\int \frac{du}{u^{n-1}\sqrt{a+bu}} = \int U\,dV = UV - \int V\,dU = \frac{2\sqrt{a+bu}}{bu^{n-1}} + \frac{2(n-1)}{b}\int \frac{\sqrt{a+bu}}{u^n}\,du$$

$$= \frac{2\sqrt{a+bu}}{bu^{n-1}} + \frac{2(n-1)}{b}\int \frac{a+bu}{u^n\sqrt{a+bu}}\,du$$

$$= \frac{2\sqrt{a+bu}}{bu^{n-1}} + 2(n-1)\int \frac{du}{u^{n-1}\sqrt{a+bu}} + \frac{2a(n-1)}{b}\int \frac{du}{u^n\sqrt{a+bu}}$$

Rearranging the equation gives $\dfrac{2a(n-1)}{b}\displaystyle\int \frac{du}{u^n\sqrt{a+bu}} = -\frac{2\sqrt{a+bu}}{bu^{n-1}} - (2n-3)\int \frac{du}{u^{n-1}\sqrt{a+bu}} \quad \Rightarrow$

$$\int \frac{du}{u^n\sqrt{a+bu}} = \frac{-\sqrt{a+bu}}{a(n-1)u^{n-1}} - \frac{b(2n-3)}{2a(n-1)}\int \frac{du}{u^{n-1}\sqrt{a+bu}}$$

71. For $n \geq 0$, $\int_0^\infty x^n\,dx = \lim\limits_{t\to\infty}\left[x^{n+1}/(n+1)\right]_0^t = \infty$. For $n < 0$, $\int_0^\infty x^n\,dx = \int_0^1 x^n\,dx + \int_1^\infty x^n\,dx$. Both integrals are improper. By (7.8.2), the second integral diverges if $-1 \leq n < 0$. By Exercise 7.8.69, the first integral diverges if $n \leq -1$. Thus, $\int_0^\infty x^n\,dx$ is divergent for all values of n.

72. $I = \displaystyle\int_0^\infty e^{ax}\cos x\,dx = \lim_{t\to\infty}\int_0^t e^{ax}\cos x\,dx \overset{\substack{99\text{ with}\\ b=1}}{=} \lim_{t\to\infty}\left[\frac{e^{ax}}{a^2+1}(a\cos x + \sin x)\right]_0^t$

$$= \lim_{t\to\infty}\left[\frac{e^{at}}{a^2+1}(a\cos t + \sin t) - \frac{1}{a^2+1}(a)\right] = \frac{1}{a^2+1}\lim_{t\to\infty}\left[e^{at}(a\cos t + \sin t) - a\right].$$

For $a \geq 0$, the limit does not exist due to oscillation. For $a < 0$, $\lim\limits_{t\to\infty}\left[e^{at}(a\cos t + \sin t)\right] = 0$ by the Squeeze Theorem, because $\left|e^{at}(a\cos t + \sin t)\right| \leq e^{at}(|a|+1)$, so $I = \dfrac{1}{a^2+1}(-a) = -\dfrac{a}{a^2+1}$.

73. $f(x) = \dfrac{1}{\ln x}$, $\Delta x = \dfrac{b-a}{n} = \dfrac{4-2}{10} = \dfrac{1}{5}$

(a) $T_{10} = \frac{1}{5\cdot 2}\{f(2) + 2[f(2.2) + f(2.4) + \cdots + f(3.8)] + f(4)\} \approx 1.925444$

(b) $M_{10} = \frac{1}{5}[f(2.1) + f(2.3) + f(2.5) + \cdots + f(3.9)] \approx 1.920915$

(c) $S_{10} = \frac{1}{5\cdot 3}[f(2) + 4f(2.2) + 2f(2.4) + \cdots + 2f(3.6) + 4f(3.8) + f(4)] \approx 1.922470$

74. $f(x) = \sqrt{x}\cos x$, $\Delta x = \dfrac{b-a}{n} = \dfrac{4-1}{10} = \dfrac{3}{10}$

(a) $T_{10} = \frac{3}{10\cdot 2}\{f(1) + 2[f(1.3) + f(1.6) + \cdots + f(3.7)] + f(4)\} \approx -2.835151$

(b) $M_{10} = \frac{3}{10}[f(1.15) + f(1.45) + f(1.75) + \cdots + f(3.85)] \approx -2.856809$

(c) $S_{10} = \frac{3}{10\cdot 3}[f(1) + 4f(1.3) + 2f(1.6) + \cdots + 2f(3.4) + 4f(3.7) + f(4)] \approx -2.849672$

75. $f(x) = \dfrac{1}{\ln x} \quad \Rightarrow \quad f'(x) = -\dfrac{1}{x(\ln x)^2} \quad \Rightarrow \quad f''(x) = \dfrac{2+\ln x}{x^2(\ln x)^3} = \dfrac{2}{x^2(\ln x)^3} + \dfrac{1}{x^2(\ln x)^2}$. Note that each term of $f''(x)$ decreases on $[2,4]$, so we'll take $K = f''(2) \approx 2.022$. $|E_T| \leq \dfrac{K(b-a)^3}{12n^2} \approx \dfrac{2.022(4-2)^3}{12(10)^2} = 0.01348$ and

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$|E_M| \le \dfrac{K(b-a)^3}{24n^2} = 0.00674$. $|E_T| \le 0.00001 \quad \Leftrightarrow \quad \dfrac{2.022(8)}{12n^2} \le \dfrac{1}{10^5} \quad \Leftrightarrow \quad n^2 \ge \dfrac{10^5(2.022)(8)}{12} \quad \Rightarrow \quad n \ge 367.2.$

Take $n = 368$ for T_n. $|E_M| \le 0.00001 \quad \Leftrightarrow \quad n^2 \ge \dfrac{10^5(2.022)(8)}{24} \quad \Rightarrow \quad n \ge 259.6$. Take $n = 260$ for M_n.

76. $\displaystyle\int_1^4 \frac{e^x}{x}\,dx \approx S_6 = \frac{(4-1)/6}{3}\,[f(1) + 4f(1.5) + 2f(2) + 4f(2.5) + 2f(3) + 4f(3.5) + f(4)] \approx 17.739438$

77. $\Delta t = \left(\frac{10}{60} - 0\right)/\,10 = \frac{1}{60}$.

$$\begin{aligned}\text{Distance traveled } &= \textstyle\int_0^{10} v\,dt \approx S_{10}\\ &= \tfrac{1}{60\cdot 3}[40 + 4(42) + 2(45) + 4(49) + 2(52) + 4(54) + 2(56) + 4(57) + 2(57) + 4(55) + 56]\\ &= \tfrac{1}{180}(1544) = 8.5\overline{7}\text{ mi}\end{aligned}$$

78. We use Simpson's Rule with $n = 6$ and $\Delta t = \frac{24-0}{6} = 4$:

$$\begin{aligned}\text{Increase in bee population } &= \textstyle\int_0^{24} r(t)\,dt \approx S_6\\ &= \tfrac{4}{3}[r(0) + 4r(4) + 2r(8) + 4r(12) + 2r(16) + 4r(20) + r(24)]\\ &= \tfrac{4}{3}[0 + 4(300) + 2(3000) + 4(11{,}000) + 2(4000) + 4(400) + 0]\\ &= \tfrac{4}{3}(60{,}800) \approx 81{,}067\text{ bees}\end{aligned}$$

79. (a) $f(x) = \sin(\sin x)$. A CAS gives

$$\begin{aligned}f^{(4)}(x) = \sin(\sin x)[\cos^4 x + 7\cos^2 x - 3]\\ +\ \cos(\sin x)\left[6\cos^2 x \sin x + \sin x\right]\end{aligned}$$

From the graph, we see that $\left|f^{(4)}(x)\right| < 3.8$ for $x \in [0, \pi]$.

(b) We use Simpson's Rule with $f(x) = \sin(\sin x)$ and $\Delta x = \frac{\pi}{10}$:

$$\textstyle\int_0^{\pi} f(x)\,dx \approx \frac{\pi}{10\cdot 3}\left[f(0) + 4f\left(\frac{\pi}{10}\right) + 2f\left(\frac{2\pi}{10}\right) + \cdots + 4f\left(\frac{9\pi}{10}\right) + f(\pi)\right] \approx 1.786721$$

From part (a), we know that $\left|f^{(4)}(x)\right| < 3.8$ on $[0, \pi]$, so we use Theorem 7.7.4 with $K = 3.8$, and estimate the error as $|E_S| \le \dfrac{3.8(\pi - 0)^5}{180(10)^4} \approx 0.000646$.

(c) If we want the error to be less than 0.00001, we must have $|E_S| \le \dfrac{3.8\pi^5}{180n^4} \le 0.00001$,

so $n^4 \ge \dfrac{3.8\pi^5}{180(0.00001)} \approx 646{,}041.6 \quad \Rightarrow \quad n \ge 28.35$. Since n must be even for Simpson's Rule, we must have $n \ge 30$ to ensure the desired accuracy.

80. With an x-axis in the normal position, at $x = 7$ we have $C = 2\pi r = 45 \quad \Rightarrow \quad r(7) = \frac{2\pi}{45}$.

Using Simpson's Rule with $n = 4$ and $\Delta x = 7$, we have

$V = \int_0^{28} \pi[r(x)]^2\,dx \approx S_4 = \frac{7}{3}\left[0 + 4\pi\left(\frac{45}{2\pi}\right)^2 + 2\pi\left(\frac{53}{2\pi}\right)^2 + 4\pi\left(\frac{45}{2\pi}\right)^2 + 0\right] = \frac{7}{3}\left(\frac{21{,}818}{4\pi}\right) \approx 4051\text{ cm}^3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

81. (a) $\dfrac{2+\sin x}{\sqrt{x}} \geq \dfrac{1}{\sqrt{x}}$ for x in $[1,\infty)$. $\displaystyle\int_1^\infty \frac{1}{\sqrt{x}}\,dx$ is divergent by (7.8.2) with $p=\dfrac{1}{2}\leq 1$. Therefore, $\displaystyle\int_1^\infty \frac{2+\sin x}{\sqrt{x}}\,dx$ is divergent by the Comparison Theorem.

(b) $\dfrac{1}{\sqrt{1+x^4}} < \dfrac{1}{\sqrt{x^4}} = \dfrac{1}{x^2}$ for x in $[1,\infty)$. $\displaystyle\int_1^\infty \frac{1}{x^2}\,dx$ is convergent by (7.8.2) with $p=2>1$. Therefore, $\displaystyle\int_1^\infty \frac{1}{\sqrt{1+x^4}}\,dx$ is convergent by the Comparison Theorem.

82. The line $y=3$ intersects the hyperbola $y^2-x^2=1$ at two points on its upper branch, namely $\left(-2\sqrt{2},3\right)$ and $\left(2\sqrt{2},3\right)$. The desired area is

$$A=\int_{-2\sqrt{2}}^{2\sqrt{2}}\left(3-\sqrt{x^2+1}\right)dx = 2\int_0^{2\sqrt{2}}\left(3-\sqrt{x^2+1}\right)dx \overset{21}{=} 2\left[3x-\tfrac{1}{2}x\sqrt{x^2+1}-\tfrac{1}{2}\ln\left(x+\sqrt{x^2+1}\right)\right]_0^{2\sqrt{2}}$$
$$=\left[6x-x\sqrt{x^2+1}-\ln\left(x+\sqrt{x^2+1}\right)\right]_0^{2\sqrt{2}} = 12\sqrt{2}-2\sqrt{2}\cdot 3-\ln\left(2\sqrt{2}+3\right)=6\sqrt{2}-\ln\left(3+2\sqrt{2}\right)$$

Another method: $A=2\int_1^3\sqrt{y^2-1}\,dy$ and use Formula 39.

83. For x in $\left[0,\frac{\pi}{2}\right]$, $0\leq\cos^2 x\leq\cos x$. For x in $\left[\frac{\pi}{2},\pi\right]$, $\cos x\leq 0\leq\cos^2 x$. Thus,

$$\text{area}=\int_0^{\pi/2}(\cos x-\cos^2 x)\,dx+\int_{\pi/2}^{\pi}(\cos^2 x-\cos x)\,dx$$
$$=\left[\sin x-\tfrac{1}{2}x-\tfrac{1}{4}\sin 2x\right]_0^{\pi/2}+\left[\tfrac{1}{2}x+\tfrac{1}{4}\sin 2x-\sin x\right]_{\pi/2}^{\pi}=\left[\left(1-\tfrac{\pi}{4}\right)-0\right]+\left[\tfrac{\pi}{2}-\left(\tfrac{\pi}{4}-1\right)\right]=2$$

84. The curves $y=\dfrac{1}{2\pm\sqrt{x}}$ are defined for $x\geq 0$. For $x>0$, $\dfrac{1}{2-\sqrt{x}}>\dfrac{1}{2+\sqrt{x}}$. Thus, the required area is

$$\int_0^1\left(\frac{1}{2-\sqrt{x}}-\frac{1}{2+\sqrt{x}}\right)dx=\int_0^1\left(\frac{1}{2-u}-\frac{1}{2+u}\right)2u\,du \quad \left[u=\sqrt{x}\right] \quad =2\int_0^1\left(-\frac{u}{u-2}-\frac{u}{u+2}\right)du$$
$$=2\int_0^1\left(-1-\frac{2}{u-2}-1+\frac{2}{u+2}\right)du=2\left[2\ln\left|\frac{u+2}{u-2}\right|-2u\right]_0^1=4\ln 3-4.$$

85. Using the formula for disks, the volume is

$$V=\int_0^{\pi/2}\pi\,[f(x)]^2\,dx=\pi\int_0^{\pi/2}(\cos^2 x)^2\,dx=\pi\int_0^{\pi/2}\left[\tfrac{1}{2}(1+\cos 2x)\right]^2\,dx$$
$$=\tfrac{\pi}{4}\int_0^{\pi/2}(1+\cos^2 2x+2\cos 2x)\,dx=\tfrac{\pi}{4}\int_0^{\pi/2}\left[1+\tfrac{1}{2}(1+\cos 4x)+2\cos 2x\right]dx$$
$$=\tfrac{\pi}{4}\left[\tfrac{3}{2}x+\tfrac{1}{2}\left(\tfrac{1}{4}\sin 4x\right)+2\left(\tfrac{1}{2}\sin 2x\right)\right]_0^{\pi/2}=\tfrac{\pi}{4}\left[\left(\tfrac{3\pi}{4}+\tfrac{1}{8}\cdot 0+0\right)-0\right]=\tfrac{3}{16}\pi^2$$

86. Using the formula for cylindrical shells, the volume is

$$V=\int_0^{\pi/2}2\pi x f(x)\,dx=2\pi\int_0^{\pi/2}x\cos^2 x\,dx=2\pi\int_0^{\pi/2}x\left[\tfrac{1}{2}(1+\cos 2x)\right]dx=2\left(\tfrac{1}{2}\right)\pi\int_0^{\pi/2}(x+x\cos 2x)\,dx$$
$$=\pi\left(\left[\tfrac{1}{2}x^2\right]_0^{\pi/2}+\left[x\left(\tfrac{1}{2}\sin 2x\right)\right]_0^{\pi/2}-\int_0^{\pi/2}\tfrac{1}{2}\sin 2x\,dx\right) \quad \left[\begin{array}{l}\text{parts with } u=x,\\ dv=\cos 2x\,dx\end{array}\right]$$
$$=\pi\left[\tfrac{1}{2}\left(\tfrac{\pi}{2}\right)^2+0-\tfrac{1}{2}\left[-\tfrac{1}{2}\cos 2x\right]_0^{\pi/2}\right]=\tfrac{\pi^3}{8}+\tfrac{\pi}{4}(-1-1)=\tfrac{1}{8}(\pi^3-4\pi)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

87. By the Fundamental Theorem of Calculus,

$$\int_0^\infty f'(x)\,dx = \lim_{t\to\infty}\int_0^t f'(x)\,dx = \lim_{t\to\infty}[f(t)-f(0)] = \lim_{t\to\infty} f(t) - f(0) = 0 - f(0) = -f(0).$$

88. (a) $(\tan^{-1}x)_{\text{avg}} = \lim\limits_{t\to\infty}\dfrac{1}{t-0}\displaystyle\int_0^t \tan^{-1}x\,dx \overset{89}{=} \lim_{t\to\infty}\left\{\frac{1}{t}\left[x\tan^{-1}x - \tfrac{1}{2}\ln(1+x^2)\right]_0^t\right\}$

$$= \lim_{t\to\infty}\left[\frac{1}{t}\left(t\,\tan^{-1}t - \tfrac{1}{2}\ln(1+t^2)\right)\right] = \lim_{t\to\infty}\left[\tan^{-1}t - \frac{\ln\left(1+t^2\right)}{2t}\right]$$

$$\overset{\text{H}}{=} \frac{\pi}{2} - \lim_{t\to\infty}\frac{2t/(1+t^2)}{2} = \frac{\pi}{2} - 0 = \frac{\pi}{2}$$

(b) $f(x) \geq 0$ and $\int_a^\infty f(x)\,dx$ is divergent $\Rightarrow$ $\lim\limits_{t\to\infty}\int_a^t f(x)\,dx = \infty$.

$$f_{\text{avg}} = \lim_{t\to\infty}\frac{\int_a^t f(x)\,dx}{t-a}\,dx \overset{\text{H}}{=} \lim_{t\to\infty}\frac{f(t)}{1} \quad \text{[by FTC1]} \quad = \lim_{x\to\infty} f(x), \text{ if this limit exists.}$$

(c) Suppose $\int_a^\infty f(x)\,dx$ converges; that is, $\lim\limits_{t\to\infty}\int_a^t f(x)\,dx = L < \infty$. Then

$$f_{\text{avg}} = \lim_{t\to\infty}\left[\frac{1}{t-a}\int_a^t f(x)\,dx\right] = \lim_{t\to\infty}\frac{1}{t-a}\cdot\lim_{t\to\infty}\int_a^t f(x)\,dx = 0\cdot L = 0.$$

(d) $(\sin x)_{\text{avg}} = \lim\limits_{t\to\infty}\dfrac{1}{t}\displaystyle\int_0^t \sin x\,dx = \lim_{t\to\infty}\left(\frac{1}{t}\left[-\cos x\right]_0^t\right) = \lim_{t\to\infty}\left(-\frac{\cos t}{t} + \frac{1}{t}\right) = \lim_{t\to\infty}\frac{1-\cos t}{t} = 0$

89. Let $u = 1/x \;\Rightarrow\; x = 1/u \;\Rightarrow\; dx = -(1/u^2)\,du$.

$$\int_0^\infty \frac{\ln x}{1+x^2}\,dx = \int_\infty^0 \frac{\ln(1/u)}{1+1/u^2}\left(-\frac{du}{u^2}\right) = \int_\infty^0 \frac{-\ln u}{u^2+1}(-du) = \int_\infty^0 \frac{\ln u}{1+u^2}\,du = -\int_0^\infty \frac{\ln u}{1+u^2}\,du$$

Therefore, $\displaystyle\int_0^\infty \frac{\ln x}{1+x^2}\,dx = -\int_0^\infty \frac{\ln x}{1+x^2}\,dx = 0.$

90. If the distance between P and the point charge is d, then the potential V at P is

$$V = W = \int_\infty^d F\,dr = \int_\infty^d \frac{q}{4\pi\varepsilon_0 r^2}\,dr = \lim_{t\to\infty}\frac{q}{4\pi\varepsilon_0}\left[-\frac{1}{r}\right]_t^d = \frac{q}{4\pi\varepsilon_0}\lim_{t\to\infty}\left(-\frac{1}{d}+\frac{1}{t}\right) = -\frac{q}{4\pi\varepsilon_0 d}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1.

By symmetry, the problem can be reduced to finding the line $x = c$ such that the shaded area is one-third of the area of the quarter-circle. An equation of the semicircle is $y = \sqrt{49 - x^2}$, so we require that $\int_0^c \sqrt{49 - x^2}\,dx = \frac{1}{3} \cdot \frac{1}{4}\pi(7)^2 \quad \Leftrightarrow$ $\left[\frac{1}{2}x\sqrt{49 - x^2} + \frac{49}{2}\sin^{-1}(x/7)\right]_0^c = \frac{49}{12}\pi$ [by Formula 30] $\Leftrightarrow$ $\frac{1}{2}c\sqrt{49 - c^2} + \frac{49}{2}\sin^{-1}(c/7) = \frac{49}{12}\pi$.

This equation would be difficult to solve exactly, so we plot the left-hand side as a function of c, and find that the equation holds for $c \approx 1.85$. So the cuts should be made at distances of about 1.85 inches from the center of the pizza.

2.
$$\int \frac{1}{x^7 - x}\,dx = \int \frac{dx}{x(x^6 - 1)} = \int \frac{x^5}{x^6(x^6 - 1)}\,dx = \frac{1}{6}\int \frac{1}{u(u-1)}\,du \qquad \begin{bmatrix} u = x^6, \\ du = 6x^5\,dx \end{bmatrix}$$
$$= \frac{1}{6}\int \left(\frac{1}{u-1} - \frac{1}{u}\right) du = \frac{1}{6}(\ln|u-1| - \ln|u|) + C$$
$$= \frac{1}{6}\ln\left|\frac{u-1}{u}\right| + C = \frac{1}{6}\ln\left|\frac{x^6 - 1}{x^6}\right| + C$$

Alternate method:
$$\int \frac{1}{x^7 - x}\,dx = \int \frac{x^{-7}}{1 - x^{-6}}\,dx \qquad \begin{bmatrix} u = 1 - x^{-6}, \\ du = 6x^{-7}\,dx \end{bmatrix}$$
$$= \tfrac{1}{6}\int du/u = \tfrac{1}{6}\ln|u| + C = \tfrac{1}{6}\ln\left|1 - x^{-6}\right| + C$$

Other methods: Substitute $u = x^3$ or $x^3 = \sec\theta$.

3. The given integral represents the difference of the shaded areas, which appears to be 0. It can be calculated by integrating with respect to either x or y, so we find x in terms of y for each curve: $y = \sqrt[3]{1 - x^7} \quad \Rightarrow \quad x = \sqrt[7]{1 - y^3}$ and $y = \sqrt[7]{1 - x^3} \quad \Rightarrow \quad x = \sqrt[3]{1 - y^7}$, so $\int_0^1 \left(\sqrt[3]{1 - y^7} - \sqrt[7]{1 - y^3}\right) dy = \int_0^1 \left(\sqrt[7]{1 - x^3} - \sqrt[3]{1 - x^7}\right) dx$. But this equation is of the form $z = -z$. So $\int_0^1 \left(\sqrt[3]{1 - x^7} - \sqrt[7]{1 - x^3}\right) dx = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. First note that since f is increasing, it is one-to-one and hence has an inverse. Now

$$\begin{aligned}\int_0^1 f(x) + f^{-1}(x)\,dx &= \int_0^1 f(x)\,dx + \int_0^1 f^{-1}(x)\,dx \\ &= \int_0^1 f(x)\,dx + \int_0^1 f^{-1}(y)\,dy \\ &= 1\end{aligned}$$

The last equality is true because, viewing f^{-1} as a function of y and using the interpretation of the integral as the area under a graph, we see from the figure that the integral gives the area of the unit square, which is 1.

5. $I = \displaystyle\int_{-r}^{r} \frac{f(x)}{1+a^x}\,dx = \int_{-r}^{0} \frac{f(x)}{1+a^x}\,dx + \int_0^r \frac{f(x)}{1+a^x}\,dx = I_1 + I_2$

Using the substitution $u = -x$, $du = -dx$ to evaluate I_1 gives

$$\begin{aligned}\int_{-r}^{0} \frac{f(x)}{1+a^x}\,dx &= \int_r^0 \frac{f(-u)}{1+a^{-u}}\,(-du) = \int_0^r f(-u)\left[\frac{1}{1+a^{-u}}\right] du \\ &= \int_0^r f(u)\left[\frac{1}{1+a^{-u}}\right] du \quad \text{[since } f(x) \text{ is even]} \\ &= \int_0^r f(u)\left[1 - \frac{1}{1+a^u}\right] du \qquad \text{[using the provided hint]} \\ &= \int_0^r f(u)\,du - \int_0^r \frac{f(u)}{1+a^u}\,du\end{aligned}$$

Thus, $I = I_1 + I_2 = \left(\displaystyle\int_0^r f(u)\,du - \int_0^r \frac{f(u)}{1+a^u}\,du\right) + \int_0^r \frac{f(x)}{1+a^x}\,dx = \int_0^r f(u)\,du.$

6. The area of each circle is $\pi(1)^2 = \pi$. By symmetry, the area of the union of the two disks is $A = \pi + \pi - 4I$.

$$\begin{aligned}I &= \textstyle\int_{1/2}^1 \sqrt{1-x^2}\,dx \\ &\overset{30}{=} \left[\tfrac{x}{2}\sqrt{1-x^2} + \tfrac{1}{2}\sin^{-1}\left(\tfrac{x}{1}\right)\right]_{1/2}^1 \quad \text{[or substitute } x = \sin\theta\text{]} \\ &= \left(0 + \tfrac{\pi}{4}\right) - \left(\tfrac{1}{4}\tfrac{\sqrt{3}}{2} + \tfrac{1}{2}\tfrac{\pi}{6}\right) = \tfrac{\pi}{4} - \tfrac{\sqrt{3}}{8} - \tfrac{\pi}{12} = \tfrac{\pi}{6} - \tfrac{\sqrt{3}}{8}\end{aligned}$$

Thus, $A = 2\pi - 4\left(\frac{\pi}{6} - \frac{\sqrt{3}}{8}\right) = 2\pi - \frac{2\pi}{3} + \frac{\sqrt{3}}{2} = \frac{4\pi}{3} + \frac{\sqrt{3}}{2}$.

Alternate solution (no calculus): The area of the sector, with central angle at the origin, containing I is $\frac{1}{2}r^2\theta = \frac{1}{2}(1)^2\left(\frac{\pi}{3}\right) = \frac{\pi}{6}$. The area of the triangle with hypotenuse OP is $\frac{1}{2}\left(\frac{1}{2}\right)\left(\frac{\sqrt{3}}{2}\right) = \frac{\sqrt{3}}{8}$.

Thus, the area of I is $\frac{\pi}{6} - \frac{\sqrt{3}}{8}$, as calculated above.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. The area A of the remaining part of the circle is given by

$$A = 4I = 4\int_0^a \left(\sqrt{a^2 - x^2} - \frac{b}{a}\sqrt{a^2 - x^2}\right) dx = 4\left(1 - \frac{b}{a}\right)\int_0^a \sqrt{a^2 - x^2}\, dx$$

$$\overset{30}{=} \frac{4}{a}(a-b)\left[\frac{x}{2}\sqrt{a^2 - x^2} + \frac{a^2}{2}\sin^{-1}\frac{x}{a}\right]_0^a$$

$$= \frac{4}{a}(a-b)\left[\left(0 + \frac{a^2}{2}\frac{\pi}{2}\right) - 0\right] = \frac{4}{a}(a-b)\left(\frac{a^2\pi}{4}\right) = \pi a(a-b),$$

which is the area of an ellipse with semiaxes a and $a - b$.

Alternate solution: Subtracting the area of the ellipse from the area of the circle gives us $\pi a^2 - \pi ab = \pi a\,(a-b)$, as calculated above. (The formula for the area of an ellipse was derived in Example 2 in Section 7.3.)

8. (a) The tangent to the curve $y = f(x)$ at $x = x_0$ has the equation $y - f(x_0) = f'(x_0)(x - x_0)$. The y-intercept of this tangent line is $f(x_0) - f'(x_0)x_0$. Thus, L is the distance from the point $(0, f(x_0) - f'(x_0)x_0)$ to the point $(x_0, f(x_0))$; that is, $L^2 = x_0^2 + [f'(x_0)]^2 x_0^2$, so $[f'(x_0)]^2 = \dfrac{L^2 - x_0^2}{x_0^2}$ and $f'(x_0) = -\dfrac{\sqrt{L^2 - x_0^2}}{x_0}$ for $0 < x_0 < L$.

(b) $\dfrac{dy}{dx} = -\dfrac{\sqrt{L^2 - x^2}}{x} \quad \Rightarrow \quad y = \displaystyle\int \left(-\frac{\sqrt{L^2 - x^2}}{x}\right) dx.$

Let $x = L\sin\theta$. Then $dx = L\cos\theta\, d\theta$ and

$$y = \int \frac{-L\cos\theta\, L\cos\theta\, d\theta}{L\sin\theta} = L\int \frac{\sin^2\theta - 1}{\sin\theta}\, d\theta = L\int (\sin\theta - \csc\theta)\, d\theta$$

$$= -L\cos\theta - L\ln|\csc\theta - \cot\theta| + C = -\sqrt{L^2 - x^2} - L\ln\left(\frac{L}{x} - \frac{\sqrt{L^2 - x^2}}{x}\right) + C$$

When $x = L$, $y = 0$, and $0 = -0 - L\ln(1 - 0) + C$, so $C = 0$. Therefore, $y = -\sqrt{L^2 - x^2} - L\ln\left(\dfrac{L - \sqrt{L^2 - x^2}}{x}\right)$.

9. Recall that $\cos A\cos B = \frac{1}{2}[\cos(A+B) + \cos(A-B)]$. So

$$f(x) = \int_0^\pi \cos t\cos(x-t)\, dt = \tfrac{1}{2}\int_0^\pi [\cos(t + x - t) + \cos(t - x + t)]\, dt = \tfrac{1}{2}\int_0^\pi [\cos x + \cos(2t - x)]\, dt$$

$$= \tfrac{1}{2}\left[t\cos x + \tfrac{1}{2}\sin(2t - x)\right]_0^\pi = \tfrac{\pi}{2}\cos x + \tfrac{1}{4}\sin(2\pi - x) - \tfrac{1}{4}\sin(-x)$$

$$= \tfrac{\pi}{2}\cos x + \tfrac{1}{4}\sin(-x) - \tfrac{1}{4}\sin(-x) = \tfrac{\pi}{2}\cos x$$

The minimum of $\cos x$ on this domain is -1, so the minimum value of $f(x)$ is $f(\pi) = -\frac{\pi}{2}$.

10. n is a positive integer, so

$$\int (\ln x)^n\, dx = x(\ln x)^n - \int x \cdot n(\ln x)^{n-1}\,(dx/x) \quad \text{[by parts]} \; = x(\ln x)^n - n\int (\ln x)^{n-1}\, dx$$

Thus,

$$\int_0^1 (\ln x)^n\, dx = \lim_{t\to 0^+}\int_t^1 (\ln x)^n\, dx = \lim_{t\to 0^+}\left[x(\ln x)^n\right]_t^1 - n\lim_{t\to 0^+}\int_t^1 (\ln x)^{n-1}\, dx$$

$$= -\lim_{t\to 0^+}\frac{(\ln t)^n}{1/t} - n\int_0^1 (\ln x)^{n-1}\, dx = -n\int_0^1 (\ln x)^{n-1}\, dx$$

by repeated application of l'Hospital's Rule. We want to prove that $\int_0^1 (\ln x)^n\, dx = (-1)^n n!$ for every positive integer n. For

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$n = 1$, we have

$$\int_0^1 (\ln x)^1\,dx = (-1)\int_0^1 (\ln x)^0\,dx = -\int_0^1 dx = -1 \qquad \left[\text{or } \int_0^1 \ln x\,dx = \lim_{t\to 0^+}\left[x\ln x - x\right]_t^1 = -1\right]$$

Assuming that the formula holds for n, we find that

$$\int_0^1 (\ln x)^{n+1}\,dx = -(n+1)\int_0^1 (\ln x)^n\,dx = -(n+1)(-1)^n\,n! = (-1)^{n+1}(n+1)!$$

This is the formula for $n+1$. Thus, the formula holds for all positive integers n by induction.

11. In accordance with the hint, we let $I_k = \int_0^1 (1-x^2)^k\,dx$, and we find an expression for I_{k+1} in terms of I_k. We integrate I_{k+1} by parts with $u = (1-x^2)^{k+1} \;\Rightarrow\; du = (k+1)(1-x^2)^k(-2x)$, $dv = dx \;\Rightarrow\; v = x$, and then split the remaining integral into identifiable quantities:

$$\begin{aligned} I_{k+1} &= x(1-x^2)^{k+1}\big|_0^1 + 2(k+1)\int_0^1 x^2(1-x^2)^k\,dx = (2k+2)\int_0^1 (1-x^2)^k[1-(1-x^2)]\,dx \\ &= (2k+2)(I_k - I_{k+1}) \end{aligned}$$

So $I_{k+1}[1+(2k+2)] = (2k+2)I_k \;\Rightarrow\; I_{k+1} = \dfrac{2k+2}{2k+3}I_k$. Now to complete the proof, we use induction:

$I_0 = 1 = \dfrac{2^0(0!)^2}{1!}$, so the formula holds for $n = 0$. Now suppose it holds for $n = k$. Then

$$\begin{aligned} I_{k+1} &= \frac{2k+2}{2k+3}I_k = \frac{2k+2}{2k+3}\left[\frac{2^{2k}(k!)^2}{(2k+1)!}\right] = \frac{2(k+1)2^{2k}(k!)^2}{(2k+3)(2k+1)!} = \frac{2(k+1)}{2k+2}\cdot\frac{2(k+1)2^{2k}(k!)^2}{(2k+3)(2k+1)!} \\ &= \frac{[2(k+1)]^2\,2^{2k}(k!)^2}{(2k+3)(2k+2)(2k+1)!} = \frac{2^{2(k+1)}\,[(k+1)!]^2}{[2(k+1)+1]!} \end{aligned}$$

So by induction, the formula holds for all integers $n \geq 0$.

12. (a) Since $-1 \leq \sin \leq 1$, we have $-f(x) \leq f(x)\sin nx \leq f(x)$, and the graph of $y = f(x)\,\sin nx$ oscillates between $f(x)$ and $-f(x)$. (The diagram shows the case $f(x) = e^x$ and $n = 10$.) As $n \to \infty$, the graph oscillates more and more frequently; see the graphs in part (b).

(b) From the graphs of the integrand, it seems that $\lim_{n\to\infty}\int_0^1 f(x)\,\sin nx\,dx = 0$, since as n increases, the integrand oscillates more and more rapidly, and thus (since f' is continuous) it makes sense that the areas above the x-axis and below it during each oscillation approach equality.

$n = 100$

$n = 200$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) We integrate by parts with $u = f(x) \quad \Rightarrow \quad du = f'(x)\,dx$, $dv = \sin nx\,dx \quad \Rightarrow \quad v = -\dfrac{\cos nx}{n}$:

$$\int_0^1 f(x)\,\sin nx\,dx = \left[-\frac{f(x)\cos nx}{n}\right]_0^1 + \int_0^1 \frac{\cos nx}{n}\,f'(x)\,dx = \frac{1}{n}\left(\int_0^1 \cos nx f'(x)\,dx - \big[f(x)\,\cos nx\big]_0^1\right)$$
$$= \tfrac{1}{n}\left[\textstyle\int_0^1 \cos nx f'(x)\,dx + f(0) - f(1)\,\cos n\right]$$

Taking absolute values of the first and last terms in this equality, and using the facts that $|\alpha \pm \beta| \le |\alpha| + |\beta|$, $\int_0^1 f(x)\,dx \le \int_0^1 |f(x)|\,dx$, $|f(0)| = f(0)$ [f is positive], $|f'(x)| \le M$ for $0 \le x \le 1$, and $|\cos nx| \le 1$,

$$\left|\textstyle\int_0^1 f(x)\,\sin nx\,dx\right| \le \tfrac{1}{n}\left[\left|\textstyle\int_0^1 M\,dx\right| + |f(0)| + |f(1)|\right] = \tfrac{1}{n}\big[M + |f(0)| + |f(1)|\big]$$

which approaches 0 as $n \to \infty$. The result follows by the Squeeze Theorem.

13. $0 < a < b$. Now

$$\int_0^1 [bx + a(1-x)]^t\,dx = \int_a^b \frac{u^t}{(b-a)}\,du \quad [u = bx + a(1-x)] \quad = \left[\frac{u^{t+1}}{(t+1)(b-a)}\right]_a^b = \frac{b^{t+1} - a^{t+1}}{(t+1)(b-a)}.$$

Now let $y = \lim\limits_{t\to 0}\left[\dfrac{b^{t+1} - a^{t+1}}{(t+1)(b-a)}\right]^{1/t}$. Then $\ln y = \lim\limits_{t\to 0}\left[\dfrac{1}{t}\ln\dfrac{b^{t+1} - a^{t+1}}{(t+1)(b-a)}\right]$. This limit is of the form $0/0$, so we can apply l'Hospital's Rule to get

$$\ln y = \lim_{t\to 0}\left[\frac{b^{t+1}\ln b - a^{t+1}\ln a}{b^{t+1} - a^{t+1}} - \frac{1}{t+1}\right] = \frac{b\ln b - a\ln a}{b-a} - 1 = \frac{b\ln b}{b-a} - \frac{a\ln a}{b-a} - \ln e = \ln\frac{b^{b/(b-a)}}{ea^{a/(b-a)}}.$$

Therefore, $y = e^{-1}\left(\dfrac{b^b}{a^a}\right)^{1/(b-a)}$.

14.

From the graph, it appears that the area under the graph of $f(x) = \sin(e^x)$ on the interval $[t, t+1]$ is greatest when $t \approx -0.2$. To find the exact value, we write the integral as $I = \int_t^{t+1} f(x)\,dx = \int_0^{t+1} f(x)\,dx - \int_0^t f(x)\,dx$, and use FTC1 to find $dI/dt = f(t+1) - f(t) = \sin\big(e^{t+1}\big) - \sin\big(e^t\big) = 0$ when $\sin\big(e^{t+1}\big) = \sin\big(e^t\big)$.

Now we have $\sin x = \sin y$ whenever $x - y = 2k\pi$ and also whenever x and y are the same distance from $\big(k + \frac{1}{2}\big)\pi$, k any integer, since $\sin x$ is symmetric about the line $x = \big(k + \frac{1}{2}\big)\pi$. The first possibility is the more obvious one, but if we calculate $e^{t+1} - e^t = 2k\pi$, we get $t = \ln(2k\pi/(e-1))$, which is about 1.3 for $k = 1$ (the least possible value of k). From the graph, this looks unlikely to give the maximum we are looking for. So instead we set $e^{t+1} - \big(k + \frac{1}{2}\big)\pi = \big(k + \frac{1}{2}\big)\pi - e^t \quad \Leftrightarrow$ $e^{t+1} + e^t = (2k+1)\pi \quad \Leftrightarrow \quad e^t(e+1) = (2k+1)\pi \quad \Leftrightarrow \quad t = \ln((2k+1)\pi/(e+1))$. Now $k = 0 \quad \Rightarrow$ $t = \ln(\pi/(e+1)) \approx -0.16853$, which does give the maximum value, as we have seen from the graph of f.

15. Write $I = \displaystyle\int \frac{x^8}{(1+x^6)^2}\,dx = \int x^3 \cdot \frac{x^5}{(1+x^6)^2}\,dx$. Integrate by parts with $u = x^3$, $dv = \dfrac{x^5}{(1+x^6)^2}\,dx$. Then $du = 3x^2\,dx$, $v = -\dfrac{1}{6(1+x^6)} \quad \Rightarrow \quad I = -\dfrac{x^3}{6(1+x^6)} + \dfrac{1}{2}\displaystyle\int \frac{x^2}{1+x^6}\,dx$. Substitute $t = x^3$ in this latter

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

integral. $\displaystyle\int \frac{x^2}{1+x^6}\,dx = \frac{1}{3}\int \frac{dt}{1+t^2} = \frac{1}{3}\tan^{-1} t + C = \frac{1}{3}\tan^{-1}(x^3) + C$. Therefore

$I = -\dfrac{x^3}{6(1+x^6)} + \dfrac{1}{6}\tan^{-1}(x^3) + C$. Returning to the improper integral,

$$\begin{aligned}\int_{-1}^{\infty}\left(\frac{x^4}{1+x^6}\right)^2 dx &= \lim_{t\to\infty}\int_{-1}^{t}\frac{x^8}{(1+x^6)^2}\,dx = \lim_{t\to\infty}\left[-\frac{x^3}{6(1+x^6)} + \frac{1}{6}\tan^{-1}(x^3)\right]_{-1}^{t}\\ &= \lim_{t\to\infty}\left(-\frac{t^3}{6(1+t^6)} + \frac{1}{6}\tan^{-1}(t^3) + \frac{-1}{6(1+1)} - \frac{1}{6}\tan^{-1}(-1)\right)\\ &= 0 + \frac{1}{6}\left(\frac{\pi}{2}\right) - \frac{1}{12} - \frac{1}{6}\left(-\frac{\pi}{4}\right) = \frac{\pi}{12} - \frac{1}{12} + \frac{\pi}{24} = \frac{\pi}{8} - \frac{1}{12}\end{aligned}$$

16. $\displaystyle\int \sqrt{\tan x}\,dx = \int u\left(\frac{2u}{u^4+1}\,du\right)$ $\quad\left[\begin{array}{ll} u = \sqrt{\tan x}, & u^2 = \tan x \\ 2u\,du = \sec^2 x\,dx, & (\tan^2 x + 1)\,dx = (u^4+1)\,dx\end{array}\right]$

Factoring the denominator, we get

$u^4 + 1 = u^4 + 2u^2 + 1 - 2u^2 = (u^2+1)^2 - \left(\sqrt{2}\,u\right)^2 = \left(u^2 + \sqrt{2}\,u + 1\right)\left(u^2 - \sqrt{2}\,u + 1\right)$. So

$\dfrac{2u^2}{u^4+1} = \dfrac{Au+B}{u^2+\sqrt{2}\,u+1} + \dfrac{Cu+D}{u^2-\sqrt{2}\,u+1} \quad\Rightarrow\quad 2u^2 = (Au+B)\left(u^2-\sqrt{2}\,u+1\right) + (Cu+D)\left(u^2+\sqrt{2}\,u+1\right)$.

Equating coefficients of powers of u, we get $A + C = 0$ (u^3), $B - \sqrt{2}\,A + D + \sqrt{2}\,C = 2$ (u^2),

$A - \sqrt{2}\,B + C + \sqrt{2}\,D = 0$ (u), $B + D = 0$ (constants). Substituting $-A$ for C and $-B$ for D in the u-equation leads to

$B = 0$ and $D = 0$, and then substituting those values in the u^2-equation gives us $A = -1/\sqrt{2}$ and $C = 1/\sqrt{2}$. Thus,

$$\begin{aligned}\int \frac{2u^2}{u^4+1}\,du &= -\frac{1}{\sqrt{2}}\int \frac{u}{u^2+\sqrt{2}\,u+1}\,du + \frac{1}{\sqrt{2}}\int \frac{u}{u^2-\sqrt{2}\,u+1}\,du\\ &= \frac{1}{\sqrt{2}}\int \frac{\frac{1}{2}\left(2u-\sqrt{2}\right) + 1/\sqrt{2}}{u^2-\sqrt{2}\,u+1}\,du - \frac{1}{\sqrt{2}}\int \frac{\frac{1}{2}\left(2u+\sqrt{2}\right) - 1/\sqrt{2}}{u^2+\sqrt{2}\,u+1}\,du\\ &= \frac{1}{2\sqrt{2}}\int \frac{2u-\sqrt{2}}{u^2-\sqrt{2}\,u+1}\,du + \frac{1}{2}\int \frac{du}{u^2-\sqrt{2}\,u+1} - \frac{1}{2\sqrt{2}}\int \frac{2u+\sqrt{2}}{u^2+\sqrt{2}\,u+1}\,du + \frac{1}{2}\int \frac{du}{u^2+\sqrt{2}\,u+1}\\ &= \frac{1}{2\sqrt{2}}\ln\left(u^2-\sqrt{2}\,u+1\right) + \frac{1}{2}\int \frac{du}{\left(u-1/\sqrt{2}\right)^2+\frac{1}{2}} - \frac{1}{2\sqrt{2}}\ln\left(u^2+\sqrt{2}\,u+1\right) + \frac{1}{2}\int \frac{du}{\left(u+1/\sqrt{2}\right)^2+\frac{1}{2}}\\ &= \frac{1}{2\sqrt{2}}\ln\frac{u^2-\sqrt{2}\,u+1}{u^2+\sqrt{2}\,u+1} + \frac{1}{2}\frac{1}{1/\sqrt{2}}\tan^{-1}\frac{u-1/\sqrt{2}}{1/\sqrt{2}} + \frac{1}{2}\frac{1}{1/\sqrt{2}}\tan^{-1}\frac{u+1/\sqrt{2}}{1/\sqrt{2}} + C\\ &= \frac{\sqrt{2}}{4}\ln\frac{\tan x - \sqrt{2\tan x}+1}{\tan x + \sqrt{2\tan x}+1} + \frac{\sqrt{2}}{2}\tan^{-1}\left(\sqrt{2\tan x}-1\right) + \frac{\sqrt{2}}{2}\tan^{-1}\left(\sqrt{2\tan x}+1\right) + C\end{aligned}$$

17.

An equation of the circle with center $(0, c)$ and radius 1 is $x^2 + (y-c)^2 = 1^2$, so an equation of the lower semicircle is $y = c - \sqrt{1-x^2}$. At the points of tangency, the slopes of the line and semicircle must be equal. For $x \geq 0$, we must have

$y' = 2 \;\Rightarrow\; \dfrac{x}{\sqrt{1-x^2}} = 2 \;\Rightarrow\; x = 2\sqrt{1-x^2} \;\Rightarrow\; x^2 = 4(1-x^2) \;\Rightarrow$

$5x^2 = 4 \;\Rightarrow\; x^2 = \frac{4}{5} \;\Rightarrow\; x = \frac{2}{5}\sqrt{5}$ and so $y = 2\left(\frac{2}{5}\sqrt{5}\right) = \frac{4}{5}\sqrt{5}$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The slope of the perpendicular line segment is $-\frac{1}{2}$, so an equation of the line segment is $y - \frac{4}{5}\sqrt{5} = -\frac{1}{2}\left(x - \frac{2}{5}\sqrt{5}\right) \quad \Leftrightarrow$ $y = -\frac{1}{2}x + \frac{1}{5}\sqrt{5} + \frac{4}{5}\sqrt{5} \quad \Leftrightarrow \quad y = -\frac{1}{2}x + \sqrt{5}$, so $c = \sqrt{5}$ and an equation of the lower semicircle is $y = \sqrt{5} - \sqrt{1-x^2}$. Thus, the shaded area is

$$\begin{aligned} 2\int_0^{(2/5)\sqrt{5}} \left[\left(\sqrt{5} - \sqrt{1-x^2}\right) - 2x\right] dx &\overset{30}{=} 2\left[\sqrt{5}\,x - \frac{x}{2}\sqrt{1-x^2} - \frac{1}{2}\sin^{-1} x - x^2\right]_0^{(2/5)\sqrt{5}} \\ &= 2\left[2 - \frac{\sqrt{5}}{5}\cdot\frac{1}{\sqrt{5}} - \frac{1}{2}\sin^{-1}\left(\frac{2}{\sqrt{5}}\right) - \frac{4}{5}\right] - 2(0) \\ &= 2\left[1 - \frac{1}{2}\sin^{-1}\left(\frac{2}{\sqrt{5}}\right)\right] = 2 - \sin^{-1}\left(\frac{2}{\sqrt{5}}\right) \end{aligned}$$

18. (a) $M\dfrac{dv}{dt} - ub = -Mg \quad \Rightarrow \quad (M_0 - bt)\dfrac{dv}{dt} = ub - (M_0 - bt)g \quad \Rightarrow \quad \dfrac{dv}{dt} = \dfrac{ub}{M_0 - bt} - g \quad \Rightarrow$

$v(t) = -u\ln(M_0 - bt) - gt + C$. Now $0 = v(0) = -u\ln M_0 + C$, so $C = u\ln M_0$. Thus

$v(t) = u\ln M_0 - u\ln(M_0 - bt) - gt = u\ln\dfrac{M_0}{M_0 - bt} - gt$.

(b) Burnout velocity $= v\left(\dfrac{M_2}{b}\right) = u\ln\dfrac{M_0}{M_0 - M_2} - g\dfrac{M_2}{b} = u\ln\dfrac{M_0}{M_1} - g\dfrac{M_2}{b}$.

Note: The reason for the term "burnout velocity" is that M_2 kilograms of fuel is used in M_2/b seconds, so $v(M_2/b)$ is the rocket's velocity when the fuel is used up.

(c) Height at burnout time $= y\left(\dfrac{M_2}{b}\right)$. Now $\dfrac{dy}{dt} = v(t) = u\ln M_0 - gt - u\ln(M_0 - bt)$, so

$y(t) = (u\ln M_0)t - \dfrac{gt^2}{2} - \dfrac{u}{b}(M_0 - bt)\ln(M_0 - bt) + ut + C$. Since $0 = y(0) = \dfrac{u}{b}M_0\ln M_0 + C$, we get

$C = -\dfrac{u}{b}M_0\ln M_0$ and $y(t) = u(1 + \ln M_0)t - \dfrac{gt^2}{2} + \dfrac{u}{b}(M_0 - bt)\ln(M_0 - bt) - \dfrac{u}{b}M_0\ln M_0$.

Therefore, the height at burnout is

$$\begin{aligned} y\left(\frac{M_2}{b}\right) &= u(1 + \ln M_0)\frac{M_2}{b} - \frac{g}{2}\left(\frac{M_2}{b}\right)^2 + \frac{u}{b}M_1\ln M_1 - \frac{u}{b}M_0\ln M_0 \\ &= \frac{u}{b}M_2 - \frac{u}{b}M_1\ln M_0 + \frac{u}{b}M_1\ln M_1 - \frac{g}{2}\left(\frac{M_2}{b}\right)^2 = \frac{u}{b}M_2 + \frac{u}{b}M_1\ln\frac{M_1}{M_0} - \frac{g}{2}\left(\frac{M_2}{b}\right)^2 \end{aligned}$$

[In the calculation of $y(M_2/b)$, repeated use was made of the relation $M_0 = M_1 + M_2$. In particular, $t = M_2/b \quad \Rightarrow \quad M_0 - bt = M_1$.]

(d) The formula for $y(t)$ in part (c) holds while there is still fuel. Once the fuel is used up, gravity is the only force acting on the rocket. $-M_1 g = M_1\dfrac{dv}{dt} \quad \Rightarrow \quad \dfrac{dv}{dt} = -g \quad \Rightarrow \quad v(t) = -gt + c_1$, where $c_1 = v\left(\dfrac{M_2}{b}\right) + \dfrac{gM_2}{b} \quad \Rightarrow$

$v(t) = v\left(\dfrac{M_2}{b}\right) - g\left(t - \dfrac{M_2}{b}\right) \quad \Rightarrow \quad y(t) = v\left(\dfrac{M_2}{b}\right)\left(t - \dfrac{M_2}{b}\right) - \dfrac{g}{2}\left(t - \dfrac{M_2}{b}\right)^2 + c_2$, where $c_2 = y\left(\dfrac{M_2}{b}\right)$,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

so $y(t) = y\left(\frac{M_2}{b}\right) + v\left(\frac{M_2}{b}\right)\left(t - \frac{M_2}{b}\right) - \frac{g}{2}\left(t - \frac{M_2}{b}\right)^2, t \geq \frac{M_2}{b}$.

To summarize: For $0 \leq t \leq \frac{M_2}{b}$, $y(t) = u(1 + \ln M_0)t - \frac{gt^2}{2} + \frac{u}{b}(M_0 - bt)\ln(M_0 - bt) - \frac{u}{b}M_0 \ln M_0$

[from part (c)], and for $t \geq \frac{M_2}{b}$, $y(t) = y\left(\frac{M_2}{b}\right) + v\left(\frac{M_2}{b}\right)\left(t - \frac{M_2}{b}\right) - \frac{g}{2}\left(t - \frac{M_2}{b}\right)^2$ [from above].

$y\left(\frac{M_2}{b}\right)$ and $v\left(\frac{M_2}{b}\right)$ are given in parts (c) and (b), respectively.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8 ☐ FURTHER APPLICATIONS OF INTEGRATION

8.1 Arc Length

1. $y = 3 - 2x \quad \Rightarrow \quad L = \int_{-1}^{3} \sqrt{1 + (dy/dx)^2}\, dx = \int_{-1}^{3} \sqrt{1 + (-2)^2}\, dx = \sqrt{5}\,\big[\,x\,\big]_{-1}^{3} = \sqrt{5}\,[3 - (-1)] = 4\sqrt{5}.$

The arc length can be calculated using the distance formula, since the curve is a line segment, so

$L = [\text{distance from } (-1, 5) \text{ to } (3, -3)] = \sqrt{[3 - (-1)]^2 + (-3 - 5)^2} = \sqrt{80} = 4\sqrt{5}.$

2. Using the arc length formula with $y = \sqrt{4 - x^2} \quad \Rightarrow \quad \dfrac{dy}{dx} = -\dfrac{x}{\sqrt{4 - x^2}}$, we get

$$L = \int_0^2 \sqrt{1 + \left(\frac{dy}{dx}\right)^2}\, dx = \int_0^2 \sqrt{1 + \frac{x^2}{4 - x^2}}\, dx = \int_0^2 \frac{\sqrt{4}}{\sqrt{4 - x^2}}\, dx = 2\int_0^2 \frac{dx}{\sqrt{4 - x^2}}$$
$$= 2\left[\sin^{-1}\left(\frac{x}{2}\right)\right]_0^2 = 2(\sin^{-1} 1 - \sin^{-1} 0) = 2\left(\frac{\pi}{2} - 0\right) = \pi$$

The curve is one-quarter of a circle with radius 2, so the length of the arc is $\frac{1}{4}(2\pi \cdot 2) = \pi$.

3. $y = x^3 \quad \Rightarrow \quad dy/dx = 3x^2 \quad \Rightarrow \quad 1 + (dy/dx)^2 = 1 + (3x^2)^2$. So $L = \int_0^2 \sqrt{1 + 9x^4}\, dx$.

4. $y = e^x \quad \Rightarrow \quad dy/dx = e^x \quad \Rightarrow \quad 1 + (dy/dx)^2 = 1 + (e^x)^2 = 1 + e^{2x}$. So $L = \int_1^3 \sqrt{1 + e^{2x}}\, dx$.

5. $y = x - \ln x \quad \Rightarrow \quad dy/dx = 1 - 1/x \quad \Rightarrow \quad 1 + (dy/dx)^2 = 1 + (1 - 1/x)^2$. So $L = \int_1^4 \sqrt{1 + (1 - 1/x)^2}\, dx$.

6. $x = y^2 + y \quad \Rightarrow \quad dx/dy = 2y + 1 \quad \Rightarrow \quad 1 + (dx/dy)^2 = 1 + (2y + 1)^2$. So $L = \int_0^3 \sqrt{1 + (2y + 1)^2}\, dy$.

7. $x = \sin y \quad \Rightarrow \quad dx/dy = \cos y \quad \Rightarrow \quad 1 + (dx/dy)^2 = 1 + \cos^2 y$. So $L = \int_0^{\pi/2} \sqrt{1 + \cos^2 y}\, dy$.

8. $y^2 = \ln x \quad \Leftrightarrow \quad x = e^{y^2} \quad \Rightarrow \quad dx/dy = 2ye^{y^2} \quad \Rightarrow \quad 1 + (dx/dy)^2 = 1 + 4y^2 e^{2y^2}$. So $L = \int_{-1}^1 \sqrt{1 + 4y^2 e^{2y^2}}\, dy$.

9. $y = \frac{2}{3}x^{3/2} \quad \Rightarrow \quad dy/dx = x^{1/2} \quad \Rightarrow \quad 1 + (dy/dx)^2 = 1 + x$. So

$L = \int_0^2 \sqrt{1 + x}\, dx = \int_0^2 (1 + x)^{1/2}\, dx = \left[\frac{2}{3}(1 + x)^{3/2}\right]_0^2 = \frac{2}{3}(3^{3/2} - 1^{3/2}) = \frac{2}{3}\left(3\sqrt{3} - 1\right) = 2\sqrt{3} - \frac{2}{3}.$

10. $y = (x + 4)^{3/2} \quad \Rightarrow \quad dy/dx = \frac{3}{2}(x + 4)^{1/2} \quad \Rightarrow \quad 1 + (dy/dx)^2 = 1 + \frac{9}{4}(x + 4)$. So

$L = \int_0^4 \sqrt{1 + \frac{9}{4}(x + 4)}\, dx = \int_0^4 \left(10 + \frac{9}{4}x\right)^{1/2} dx = \left[\frac{8}{27}\left(10 + \frac{9}{4}x\right)^{3/2}\right]_0^4 = \frac{8}{27}(19^{3/2} - 10^{3/2}).$

11. $y = \frac{2}{3}(1 + x^2)^{3/2} \quad \Rightarrow \quad dy/dx = 2x(1 + x^2)^{1/2} \quad \Rightarrow \quad 1 + (dy/dx)^2 = 1 + 4x^2(1 + x^2)$. So

$$L = \int_0^1 \sqrt{1 + 4x^2(1 + x^2)}\, dx = \int_0^1 \sqrt{4x^4 + 4x^2 + 1}\, dx = \int_0^1 \sqrt{(2x^2 + 1)^2}\, dx = \int_0^1 \left|2x^2 + 1\right| dx$$
$$= \int_0^1 (2x^2 + 1)\, dx = \left[\tfrac{2}{3}x^3 + x\right]_0^1 = \left(\tfrac{2}{3} + 1\right) - 0 = \tfrac{5}{3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $36y^2 = (x^2-4)^3,\ y \ge 0 \quad\Rightarrow\quad y = \frac{1}{6}(x^2-4)^{3/2} \quad\Rightarrow\quad dy/dx = \frac{1}{6}\cdot\frac{3}{2}(x^2-4)^{1/2}(2x) = \frac{1}{2}x(x^2-4)^{1/2} \quad\Rightarrow$

$1+(dy/dx)^2 = 1+\frac{1}{4}x^2(x^2-4) = \frac{1}{4}x^4 - x^2 + 1 = \frac{1}{4}(x^4-4x^2+4) = \left[\frac{1}{2}(x^2-2)\right]^2$. So

$L = \int_2^3 \sqrt{\left[\frac{1}{2}(x^2-2)\right]^2}\,dx = \int_2^3 \frac{1}{2}(x^2-2)\,dx = \frac{1}{2}\left[\frac{1}{3}x^3-2x\right]_2^3 = \frac{1}{2}\left[(9-6)-\left(\frac{8}{3}-4\right)\right] = \frac{1}{2}\left(\frac{13}{3}\right) = \frac{13}{6}$.

13. $y = \dfrac{x^3}{3} + \dfrac{1}{4x} \quad\Rightarrow\quad \dfrac{dy}{dx} = x^2 - \dfrac{1}{4x^2} \quad\Rightarrow$

$1+\left(\dfrac{dy}{dx}\right)^2 = 1+\left(x^4 - \dfrac{1}{2} + \dfrac{1}{16x^4}\right) = x^4 + \dfrac{1}{2} + \dfrac{1}{16x^4} = \left(x^2 + \dfrac{1}{4x^2}\right)^2$. So

$$L = \int_1^2 \sqrt{1+\left(\frac{dy}{dx}\right)^2}\,dx = \int_1^2 \left|x^2+\frac{1}{4x^2}\right|\,dx = \int_1^2 \left(x^2+\frac{1}{4x^2}\right)\,dx$$

$$= \left[\frac{1}{3}x^3 - \frac{1}{4x}\right]_1^2 = \left(\frac{8}{3}-\frac{1}{8}\right) - \left(\frac{1}{3}-\frac{1}{4}\right) = \frac{7}{3}+\frac{1}{8} = \frac{59}{24}$$

14. $x = \dfrac{y^4}{8} + \dfrac{1}{4y^2} \quad\Rightarrow\quad \dfrac{dx}{dy} = \dfrac{1}{2}y^3 - \dfrac{1}{2}y^{-3} \quad\Rightarrow$

$1+(dx/dy)^2 = 1+\frac{1}{4}y^6 - \frac{1}{2} + \frac{1}{4}y^{-6} = \frac{1}{4}y^6 + \frac{1}{2} + \frac{1}{4}y^{-6} = \left(\frac{1}{2}y^3+\frac{1}{2}y^{-3}\right)^2$. So

$L = \int_1^2 \sqrt{\left(\frac{1}{2}y^3+\frac{1}{2}y^{-3}\right)^2}\,dy = \int_1^2 \left(\frac{1}{2}y^3+\frac{1}{2}y^{-3}\right)dy = \left[\frac{1}{8}y^4 - \frac{1}{4}y^{-2}\right]_1^2 = \left(2-\frac{1}{16}\right) - \left(\frac{1}{8}-\frac{1}{4}\right)$

$= 2 + \frac{1}{16} = \frac{33}{16}$

15. $y = \frac{1}{2}\ln(\sin 2x) \quad\Rightarrow\quad \dfrac{dy}{dx} = \dfrac{\cos 2x}{\sin 2x} = \cot 2x \quad\Rightarrow\quad 1+\left(\dfrac{dy}{dx}\right)^2 = 1+\cot^2 2x = \csc^2 2x$. So

$$L = \int_{\pi/8}^{\pi/6} \sqrt{\csc^2 2x}\,dx = \int_{\pi/8}^{\pi/6} |\csc 2x|\,dx = \int_{\pi/8}^{\pi/6} \csc 2x\,dx = \frac{1}{2}\int_{\pi/4}^{\pi/3} \csc u\,du \qquad \begin{bmatrix} u = 2x \\ du = 2\,dx \end{bmatrix}$$

$$= \frac{1}{2}\Big[\ln|\csc u - \cot u|\Big]_{\pi/4}^{\pi/3} = \frac{1}{2}\left[\ln\left(\frac{2}{\sqrt{3}} - \frac{1}{\sqrt{3}}\right) - \ln\left(\sqrt{2}-1\right)\right] = \frac{1}{2}\left[\ln\frac{1}{\sqrt{3}} - \ln\left(\sqrt{2}-1\right)\right]$$

16. $y = \ln(\cos x) \quad\Rightarrow\quad dy/dx = -\tan x \quad\Rightarrow\quad 1+(dy/dx)^2 = 1+\tan^2 x = \sec^2 x$. So

$L = \int_0^{\pi/3} \sqrt{\sec^2 x}\,dx = \int_0^{\pi/3} \sec x\,dx = \Big[\ln|\sec x + \tan x|\Big]_0^{\pi/3} = \ln\left(2+\sqrt{3}\right) - \ln(1+0) = \ln\left(2+\sqrt{3}\right)$.

17. $y = \ln(\sec x) \quad\Rightarrow\quad \dfrac{dy}{dx} = \dfrac{\sec x\,\tan x}{\sec x} = \tan x \quad\Rightarrow\quad 1+\left(\dfrac{dy}{dx}\right)^2 = 1+\tan^2 x = \sec^2 x$. So

$L = \int_0^{\pi/4} \sqrt{\sec^2 x}\,dx = \int_0^{\pi/4} |\sec x|\,dx = \int_0^{\pi/4} \sec x\,dx = \Big[\ln|\sec x + \tan x|\Big]_0^{\pi/4}$

$= \ln\left(\sqrt{2}+1\right) - \ln(1+0) = \ln\left(\sqrt{2}+1\right)$

18. $x = e^y + \frac{1}{4}e^{-y} \quad\Rightarrow\quad dx/dy = e^y - \frac{1}{4}e^{-y} \quad\Rightarrow$

$1+(dx/dy)^2 = 1+\left(e^y - \frac{1}{4}e^{-y}\right)^2 = 1+\left(e^{2y} - \frac{1}{2} + \frac{1}{16}e^{-2y}\right) = e^{2y} + \frac{1}{2} + \frac{1}{16}e^{-2y} = \left(e^y + \frac{1}{4}e^{-y}\right)^2$. So

$$L = \int_0^1 \sqrt{\left(e^y+\frac{1}{4}e^{-y}\right)^2}\,dy = \int_0^1 \left|e^y+\frac{1}{4}e^{-y}\right|\,dy = \int_0^1 \left(e^y+\frac{1}{4}e^{-y}\right)\,dy = \left[e^y - \frac{1}{4}e^{-y}\right]_0^1$$

$$= e - \frac{1}{4}e^{-1} - \left(1-\frac{1}{4}\right) = e - \frac{1}{4e} - \frac{3}{4}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $x = \frac{1}{3}\sqrt{y}\,(y-3) = \frac{1}{3}y^{3/2} - y^{1/2} \quad \Rightarrow \quad dx/dy = \frac{1}{2}y^{1/2} - \frac{1}{2}y^{-1/2} \quad \Rightarrow$

$1 + (dx/dy)^2 = 1 + \frac{1}{4}y - \frac{1}{2} + \frac{1}{4}y^{-1} = \frac{1}{4}y + \frac{1}{2} + \frac{1}{4}y^{-1} = \left(\frac{1}{2}y^{1/2} + \frac{1}{2}y^{-1/2}\right)^2$. So

$L = \int_1^9 \left(\frac{1}{2}y^{1/2} + \frac{1}{2}y^{-1/2}\right) dy = \frac{1}{2}\left[\frac{2}{3}y^{3/2} + 2y^{1/2}\right]_1^9 = \frac{1}{2}\left[\left(\frac{2}{3}\cdot 27 + 2\cdot 3\right) - \left(\frac{2}{3}\cdot 1 + 2\cdot 1\right)\right]$

$= \frac{1}{2}\left(24 - \frac{8}{3}\right) = \frac{1}{2}\left(\frac{64}{3}\right) = \frac{32}{3}$.

20. $y = 3 + \frac{1}{2}\cosh 2x \quad \Rightarrow \quad dy/dx = \sinh 2x \quad \Rightarrow \quad 1 + (dy/dx)^2 = 1 + \sinh^2(2x) = \cosh^2(2x)$. So

$L = \int_0^1 \sqrt{\cosh^2(2x)}\,dx = \int_0^1 \cosh 2x\,dx = \left[\frac{1}{2}\sinh 2x\right]_0^1 = \frac{1}{2}\sinh 2 - 0 = \frac{1}{2}\sinh 2$.

21. $y = \dfrac{1}{4}x^2 - \dfrac{1}{2}\ln x \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{1}{2}x - \dfrac{1}{2x} \quad \Rightarrow$

$1 + \left(\dfrac{dy}{dx}\right)^2 = 1 + \left(\dfrac{1}{4}x^2 - \dfrac{1}{2} + \dfrac{1}{4x^2}\right) = \dfrac{1}{4}x^2 + \dfrac{1}{2} + \dfrac{1}{4x^2} = \left(\dfrac{1}{2}x + \dfrac{1}{2x}\right)^2$. So

$$L = \int_1^2 \sqrt{1 + \left(\frac{dy}{dx}\right)^2}\,dx = \int_1^2 \left|\frac{1}{2}x + \frac{1}{2x}\right| dx = \int_1^2 \left(\frac{1}{2}x + \frac{1}{2x}\right) dx$$

$$= \left[\frac{1}{4}x^2 + \frac{1}{2}\ln|x|\right]_1^2 = \left(1 + \frac{1}{2}\ln 2\right) - \left(\frac{1}{4} + 0\right) = \frac{3}{4} + \frac{1}{2}\ln 2$$

22. $y = \sqrt{x - x^2} + \sin^{-1}\left(\sqrt{x}\right) \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{1-2x}{2\sqrt{x-x^2}} + \dfrac{1}{2\sqrt{x}\sqrt{1-x}} = \dfrac{2-2x}{2\sqrt{x}\sqrt{1-x}} = \sqrt{\dfrac{1-x}{x}} \quad \Rightarrow$

$1 + \left(\dfrac{dy}{dx}\right)^2 = 1 + \dfrac{1-x}{x} = \dfrac{1}{x}$. The curve has endpoints $(0,0)$ and $\left(1, \frac{\pi}{2}\right)$,

so $L = \int_0^1 \sqrt{1/x}\,dx = \lim\limits_{t\to 0^+} \int_t^1 \sqrt{1/x}\,dx = \lim\limits_{t\to 0^+} \left[2\sqrt{x}\right]_t^1 = \lim\limits_{t\to 0^+} \left[2\sqrt{1} - 2\sqrt{t}\right] = 2 - 0 = 2$.

23. $y = \ln(1 - x^2) \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{1}{1-x^2}\cdot(-2x) \quad \Rightarrow$

$1 + \left(\dfrac{dy}{dx}\right)^2 = 1 + \dfrac{4x^2}{(1-x^2)^2} = \dfrac{1 - 2x^2 + x^4 + 4x^2}{(1-x^2)^2} = \dfrac{1 + 2x^2 + x^4}{(1-x^2)^2} = \dfrac{(1+x^2)^2}{(1-x^2)^2} \quad \Rightarrow$

$\sqrt{1 + \left(\dfrac{dy}{dx}\right)^2} = \sqrt{\left(\dfrac{1+x^2}{1-x^2}\right)^2} = \dfrac{1+x^2}{1-x^2} = -1 + \dfrac{2}{1-x^2}$ [by division] $= -1 + \dfrac{1}{1+x} + \dfrac{1}{1-x}$ [partial fractions].

So $L = \displaystyle\int_0^{1/2} \left(-1 + \frac{1}{1+x} + \frac{1}{1-x}\right) dx = \left[-x + \ln|1+x| - \ln|1-x|\right]_0^{1/2} = \left(-\frac{1}{2} + \ln\frac{3}{2} - \ln\frac{1}{2}\right) - 0 = \ln 3 - \frac{1}{2}$.

24. $y = 1 - e^{-x} \quad \Rightarrow \quad dy/dx = -(-e^{-x}) = e^{-x} \quad \Rightarrow \quad 1 + (dy/dx)^2 = 1 + e^{-2x}$. So

$$L = \int_0^2 \sqrt{1 + e^{-2x}}\,dx = \int_1^{e^{-2}} \sqrt{1+u^2}\left(-\frac{1}{u}\,du\right) \qquad [u = e^{-x}]$$

$$\overset{23}{=} \left[\ln\left|\frac{1 + \sqrt{1+u^2}}{u}\right| - \sqrt{1+u^2}\right]_1^{e^{-2}} \qquad [\text{or substitute } u = \tan\theta]$$

$$= \ln\left|\frac{1 + \sqrt{1+e^{-4}}}{e^{-2}}\right| - \sqrt{1+e^{-4}} - \ln\left|\frac{1+\sqrt{2}}{1}\right| + \sqrt{2}$$

$$= \ln\left(1 + \sqrt{1+e^{-4}}\right) - \ln e^{-2} - \sqrt{1+e^{-4}} - \ln\left(1+\sqrt{2}\right) + \sqrt{2}$$

$$= \ln\left(1 + \sqrt{1+e^{-4}}\right) + 2 - \sqrt{1+e^{-4}} - \ln\left(1+\sqrt{2}\right) + \sqrt{2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $y = \frac{1}{2}x^2 \quad \Rightarrow \quad dy/dx = x \quad \Rightarrow \quad 1 + (dy/dx)^2 = 1 + x^2$. So

$$L = \int_{-1}^{1} \sqrt{1+x^2}\,dx = 2\int_0^1 \sqrt{1+x^2}\,dx \quad \text{[by symmetry]} \quad \overset{21}{=} 2\left[\tfrac{x}{2}\sqrt{1+x^2} + \tfrac{1}{2}\ln\left(x+\sqrt{1+x^2}\right)\right]_0^1 \quad \begin{bmatrix}\text{or substitute}\\ x = \tan\theta\end{bmatrix}$$

$$= 2\left[\left(\tfrac{1}{2}\sqrt{2} + \tfrac{1}{2}\ln\left(1+\sqrt{2}\right)\right) - \left(0 + \tfrac{1}{2}\ln 1\right)\right] = \sqrt{2} + \ln\left(1+\sqrt{2}\right)$$

26. $x^2 = (y-4)^3 \quad \Rightarrow \quad x = (y-4)^{3/2} \quad [\text{for } x > 0] \quad \Rightarrow \quad dx/dy = \frac{3}{2}(y-4)^{1/2} \quad \Rightarrow$

$1 + (dx/dy)^2 = 1 + \frac{9}{4}(y-4) = \frac{9}{4}y - 8$. So

$$L = \int_5^8 \sqrt{\tfrac{9}{4}y - 8}\,dy = \int_{13/4}^{10} \sqrt{u}\,\left(\tfrac{4}{9}\,du\right) \quad \begin{bmatrix} u = \frac{9}{4}y - 8, \\ du = \frac{9}{4}\,dy \end{bmatrix} \quad = \tfrac{4}{9}\left[\tfrac{2}{3}u^{3/2}\right]_{13/4}^{10}$$

$$= \tfrac{8}{27}\left[10^{3/2} - \left(\tfrac{13}{4}\right)^{3/2}\right] \quad \left[\text{or } \tfrac{1}{27}\left(80\sqrt{10} - 13\sqrt{13}\right)\right]$$

27.

From the figure, the length of the curve is slightly larger than the hypotenuse of the triangle formed by the points $(1, 2)$, $(1, 12)$, and $(2, 12)$. This length is about $\sqrt{10^2 + 1^2} \approx 10$, so we might estimate the length to be 10. $y = x^2 + x^3 \quad \Rightarrow \quad y' = 2x + 3x^2 \quad \Rightarrow \quad 1 + (y')^2 = 1 + (2x + 3x^2)^2$. So $L = \int_1^2 \sqrt{1 + (2x + 3x^2)^2}\,dx \approx 10.0556$.

28.

From the figure, the length of the curve is slightly larger than the hypotenuse of the triangle formed by the points $(1, 1)$, $\left(\frac{\pi}{2}, 1\right)$, and $\left(\frac{\pi}{2}, \frac{\pi}{2}\right)$. This length is about $\sqrt{\left(\frac{\pi}{2}\right)^2 + \left(\frac{\pi}{2} - 1\right)^2} \approx 1.7$, so we might estimate the length to be 1.7. $y = x + \cos x \quad \Rightarrow \quad y' = 1 - \sin x \quad \Rightarrow$ $1 + (y')^2 = 1 + (1 - \sin x)^2$. So $L = \int_0^{\pi/2} \sqrt{1 + (1 - \sin x)^2}\,dx \approx 1.7294$.

29.

From the figure, the length of the curve is slightly larger than the line segment joining the points $(1, 1)$ and $\left(4, \sqrt[3]{4}\right)$. This length is $\sqrt{(4-1)^2 + \left(\sqrt[3]{4} - 1\right)^2} \approx 3.057$, so we might estimate the length of the curve to be 3.06. $y = \sqrt[3]{x} = x^{1/3} \quad \Rightarrow \quad y' = \frac{1}{3}x^{-2/3} \quad \Rightarrow$ $1 + (y')^2 = 1 + \frac{1}{9}x^{-4/3}$. So $L = \int_1^4 \sqrt{1 + \frac{1}{9}x^{-4/3}}\,dx \approx 3.0609$.

30.

From the figure, the length of the curve is slightly larger than the line segment joining the points $(0, 0)$ and $(1, \tan 1)$. This length is $\sqrt{(1-0)^2 + (\tan 1 - 0)^2} \approx 1.851$, so we might estimate the length of the curve to be 1.9. $y = x\tan x \quad \Rightarrow$ $y' = x\sec^2 x + \tan x \quad \Rightarrow \quad 1 + (y')^2 = 1 + (x\sec^2 x + \tan x)^2$. So $L = \int_0^1 \sqrt{1 + (x\sec^2 x + \tan x)^2}\,dx \approx 1.9799$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31.

From the figure, the length of the curve is slightly larger than the line segment joining the points $\left(1, e^{-1}\right)$ and $\left(2, 2e^{-2}\right)$. This length is $\sqrt{(2-1)^2 + (2e^{-2} - e^{-1})^2} \approx 1.0047$, so we might estimate the length of the curve to be 1.005. $y = xe^{-x} \quad \Rightarrow \quad y' = -xe^{-x} + e^{-x} \quad \Rightarrow$ $1 + (y')^2 = 1 + (e^{-x} - xe^{-x})^2$. So $L = \int_1^2 \sqrt{1 + (e^{-x} - xe^{-x})^2}\, dx \approx 1.0054$.

32.

From the figure, the curve is slightly larger than the sum of the lengths of the line segments joining the points $(-2, \ln 8)$ to $(0, \ln 4)$, and $(0, \ln 4)$ to $(2, \ln 8)$. These line segments each have length $\sqrt{(0+2)^2 + (\ln 4 - \ln 8)^2} \approx 2.1167$, so we might estimate the length of the curve to be $2(2.1167) = 4.2334$ or 4.25.

$$y = \ln(x^2 + 4) \quad \Rightarrow \quad y' = \frac{2x}{x^2 + 4} \quad \Rightarrow \quad 1 + (y')^2 = 1 + \frac{4x^2}{(x^2+4)^2}.$$

So $L = \displaystyle\int_{-2}^{2} \sqrt{1 + \frac{4x^2}{(x^2+4)^2}}\, dx \approx 4.2726$.

33. $y = x \sin x \quad \Rightarrow \quad dy/dx = x\cos x + (\sin x)(1) \quad \Rightarrow \quad 1 + (dy/dx)^2 = 1 + (x\cos x + \sin x)^2$. Let $f(x) = \sqrt{1 + (dy/dx)^2} = \sqrt{1 + (x\cos x + \sin x)^2}$. Then $L = \int_0^{2\pi} f(x)\, dx$. Since $n = 10$, $\Delta x = \frac{2\pi - 0}{10} = \frac{\pi}{5}$. Now

$$\begin{aligned} L \approx S_{10} &= \tfrac{\pi/5}{3}\left[f(0) + 4f\left(\tfrac{\pi}{5}\right) + 2f\left(\tfrac{2\pi}{5}\right) + 4f\left(\tfrac{3\pi}{5}\right) + 2f\left(\tfrac{4\pi}{5}\right) + 4f\left(\tfrac{5\pi}{5}\right) + 2f\left(\tfrac{6\pi}{5}\right)\right. \\ &\qquad \left. + 4f\left(\tfrac{7\pi}{5}\right) + 2f\left(\tfrac{8\pi}{5}\right) + 4f\left(\tfrac{9\pi}{5}\right) + f(2\pi)\right] \\ &\approx 15.498085 \end{aligned}$$

The value of the integral produced by a calculator is 15.374568 (to six decimal places).

34. $y = e^{-x^2} \quad \Rightarrow \quad dy/dx = e^{-x^2}(-2x) \quad \Rightarrow \quad L = \int_0^2 f(x)\, dx$, where $f(x) = \sqrt{1 + 4x^2 e^{-2x^2}}$.

Since $n = 10$, $\Delta x = \frac{2-0}{10} = \frac{1}{5}$. Now

$$\begin{aligned} L \approx S_{10} &= \tfrac{1/5}{3}[f(0) + 4f(0.2) + 2f(0.4) + 4f(0.6) + 2f(0.8) + 4f(1) + 2f(1.2) \\ &\qquad + 4f(1.4) + 2f(1.6) + 4f(1.8) + f(2)] \\ &\approx 2.280559 \end{aligned}$$

The value of the integral produced by a calculator is 2.280526 (to six decimal places).

35. (a) Let $f(x) = y = x\sqrt[3]{4 - x}$ with $0 \le x \le 4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The polygon with one side is just the line segment joining the points $(0, f(0)) = (0, 0)$ and $(4, f(4)) = (4, 0)$, and its length $L_1 = 4$.

The polygon with two sides joins the points $(0, 0)$, $(2, f(2)) = \left(2, 2\sqrt[3]{2}\right)$ and $(4, 0)$. Its length

$$L_2 = \sqrt{(2-0)^2 + \left(2\sqrt[3]{2} - 0\right)^2} + \sqrt{(4-2)^2 + \left(0 - 2\sqrt[3]{2}\right)^2} = 2\sqrt{4 + 2^{8/3}} \approx 6.43$$

Similarly, the inscribed polygon with four sides joins the points $(0, 0)$, $\left(1, \sqrt[3]{3}\right)$, $\left(2, 2\sqrt[3]{2}\right)$, $(3, 3)$, and $(4, 0)$, so its length

$$L_4 = \sqrt{1 + \left(\sqrt[3]{3}\right)^2} + \sqrt{1 + \left(2\sqrt[3]{2} - \sqrt[3]{3}\right)^2} + \sqrt{1 + \left(3 - 2\sqrt[3]{2}\right)^2} + \sqrt{1+9} \approx 7.50$$

(c) Using the arc length formula with $\dfrac{dy}{dx} = x\left[\frac{1}{3}(4-x)^{-2/3}(-1)\right] + \sqrt[3]{4-x} = \dfrac{12-4x}{3(4-x)^{2/3}}$, the length of the curve is

$$L = \int_0^4 \sqrt{1 + \left(\frac{dy}{dx}\right)^2}\, dx = \int_0^4 \sqrt{1 + \left[\frac{12-4x}{3(4-x)^{2/3}}\right]^2}\, dx.$$

(d) According to a calculator, the length of the curve is $L \approx 7.7988$. The actual value is larger than any of the approximations in part (b). This is always true, since any approximating straight line between two points on the curve is shorter than the length of the curve between the two points.

36. (a) Let $f(x) = y = x + \sin x$ with $0 \le x \le 2\pi$.

(b) The polygon with one side is just the line segment joining the points $(0, f(0)) = (0, 0)$ and $(2\pi, f(2\pi)) = (2\pi, 2\pi)$, and its length is $\sqrt{(2\pi - 0)^2 + (2\pi - 0)^2} = 2\sqrt{2}\,\pi \approx 8.9$.

The polygon with two sides joins the points $(0, 0)$, $(\pi, f(\pi)) = (\pi, \pi)$, and $(2\pi, 2\pi)$. Its length is

$$\sqrt{(\pi - 0)^2 + (\pi - 0)^2} + \sqrt{(2\pi - \pi)^2 + (2\pi - \pi)^2} = \sqrt{2}\,\pi + \sqrt{2}\,\pi$$
$$= 2\sqrt{2}\,\pi \approx 8.9$$

Note from the diagram that the two approximations are the same because the sides of the two-sided polygon are in fact on the same line, since $f(\pi) = \pi = \frac{1}{2} f(2\pi)$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The four-sided polygon joins the points $(0,0)$, $\left(\frac{\pi}{2}, \frac{\pi}{2}+1\right)$, (π,π), $\left(\frac{3\pi}{2}, \frac{3\pi}{2}-1\right)$, and $(2\pi, 2\pi)$, so its length is

$$\sqrt{\left(\tfrac{\pi}{2}\right)^2+\left(\tfrac{\pi}{2}+1\right)^2}+\sqrt{\left(\tfrac{\pi}{2}\right)^2+\left(\tfrac{\pi}{2}-1\right)^2}+\sqrt{\left(\tfrac{\pi}{2}\right)^2+\left(\tfrac{\pi}{2}-1\right)^2}+\sqrt{\left(\tfrac{\pi}{2}\right)^2+\left(\tfrac{\pi}{2}+1\right)^2}\approx 9.4$$

(c) Using the arc length formula with $dy/dx = 1+\cos x$, the length of the curve is

$$L=\int_0^{2\pi}\sqrt{1+(1+\cos x)^2}\,dx=\int_0^{2\pi}\sqrt{2+2\cos x+\cos^2 x}\,dx$$

(d) The calculator approximates the integral as 9.5076. The actual length is larger than the approximations in part (b).

37. $y=e^x \;\Rightarrow\; dy/dx=e^x \;\Rightarrow\; 1+(dy/dx)^2 \;\Rightarrow\; 1+e^{2x} \;\Rightarrow$

$$L=\int_0^2\sqrt{1+e^{2x}}\,dx=\int_1^{e^2}\sqrt{1+u^2}\left(\frac{1}{u}\,du\right) \qquad \begin{bmatrix} u=e^x, \\ du=e^x\,dx \end{bmatrix}$$

$$\overset{23}{=}\left[\sqrt{1+u^2}-\ln\left|\frac{1+\sqrt{1+u^2}}{u}\right|\right]_1^{e^2}=\left(\sqrt{1+e^4}-\ln\frac{1+\sqrt{1+e^4}}{e^2}\right)-\left(\sqrt{2}-\ln\frac{1+\sqrt{2}}{1}\right)$$

$$=\sqrt{1+e^4}-\ln(1+\sqrt{1+e^4})+2-\sqrt{2}+\ln(1+\sqrt{2})$$

An equivalent answer from a CAS is

$$-\sqrt{2}+\operatorname{arctanh}(\sqrt{2}/2)+\sqrt{e^4+1}-\operatorname{arctanh}(1/\sqrt{e^4+1}).$$

38. $y=x^{4/3} \;\Rightarrow\; dy/dx=\frac{4}{3}x^{1/3} \;\Rightarrow\; 1+(dy/dx)^2=1+\frac{16}{9}x^{2/3} \;\Rightarrow$

$$L=\int_0^1\sqrt{1+\tfrac{16}{9}x^{2/3}}\,dx=\int_0^{4/3}\sqrt{1+u^2}\,\tfrac{81}{64}u^2\,du \qquad \begin{bmatrix} u=\frac{4}{3}x^{1/3},\ du=\frac{4}{9}x^{-2/3}\,dx, \\ dx=\frac{9}{4}x^{2/3}\,du=\frac{9}{4}\cdot\frac{9}{16}u^2\,du=\frac{81}{64}u^2\,du \end{bmatrix}$$

$$\overset{22}{=}\tfrac{81}{64}\left[\tfrac{1}{8}u(1+2u^2)\sqrt{1+u^2}-\tfrac{1}{8}\ln\left(u+\sqrt{1+u^2}\right)\right]_0^{4/3}=\tfrac{81}{64}\left[\tfrac{1}{6}\left(1+\tfrac{32}{9}\right)\sqrt{\tfrac{25}{9}}-\tfrac{1}{8}\ln\left(\tfrac{4}{3}+\sqrt{\tfrac{25}{9}}\right)\right]$$

$$=\tfrac{81}{64}\left(\tfrac{1}{6}\cdot\tfrac{41}{9}\cdot\tfrac{5}{3}-\tfrac{1}{8}\ln 3\right)=\tfrac{205}{128}-\tfrac{81}{512}\ln 3\approx 1.4277586$$

39. The astroid $x^{2/3}+y^{2/3}=1$ has an equal length of arc in each quadrant. Thus, we can find the length of the curve in the first quadrant and then multiply by 4. The top half of the astroid has equation $y=(1-x^{2/3})^{3/2}$. Then

$$dy/dx=-x^{-1/3}(1-x^{2/3})^{1/2} \;\Rightarrow\; 1+(dy/dx)^2=1+\left[-x^{-1/3}(1-x^{2/3})^{1/2}\right]^2=1+x^{-2/3}(1-x^{2/3})=x^{-2/3}.$$

So the portion of the astroid in quadrant 1 has length $L=\int_0^1\sqrt{x^{-2/3}}\,dx=\int_0^1 x^{-1/3}\,dx=\left[\frac{3}{2}x^{2/3}\right]_0^1=\frac{3}{2}-0=\frac{3}{2}$. Thus, the astroid has length $4\left(\frac{3}{2}\right)=6$.

40. (a) Graph of $y^3=x^2$:

(b) $y=x^{2/3} \;\Rightarrow\; 1+\left(\dfrac{dy}{dx}\right)^2=1+\left(\frac{2}{3}x^{-1/3}\right)^2=1+\frac{4}{9}x^{-2/3}$. So $L=\int_0^1\sqrt{1+\frac{4}{9}x^{-2/3}}\,dx$ [an improper integral].

$x=y^{3/2} \;\Rightarrow\; 1+\left(\dfrac{dx}{dy}\right)^2=1+\left(\frac{3}{2}y^{1/2}\right)^2=1+\frac{9}{4}y$. So $L=\int_0^1\sqrt{1+\frac{9}{4}y}\,dy$.

The second integral equals $\frac{4}{9}\cdot\frac{2}{3}\left[\left(1+\frac{9}{4}y\right)^{3/2}\right]_0^1=\frac{8}{27}\left(\frac{13\sqrt{13}}{8}-1\right)=\frac{13\sqrt{13}-8}{27}$. [continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The first integral can be evaluated as follows:

$$\int_0^1 \sqrt{1+\tfrac{4}{9}x^{-2/3}}\,dx = \lim_{t\to 0^+}\int_t^1 \frac{\sqrt{9x^{2/3}+4}}{3x^{1/3}}\,dx = \lim_{t\to 0^+}\int_{9t^{2/3}}^9 \frac{\sqrt{u+4}}{18}\,du \qquad \left[\begin{matrix} u = 9x^{2/3}, \\ du = 6x^{-1/3}\,dx \end{matrix}\right]$$

$$= \int_0^9 \frac{\sqrt{u+4}}{18}\,du = \frac{1}{18}\cdot\left[\frac{2}{3}(u+4)^{3/2}\right]_0^9 = \frac{1}{27}(13^{3/2}-4^{3/2}) = \frac{13\sqrt{13}-8}{27}$$

(c) L = length of the arc of this curve from $(-1,1)$ to $(8,4)$

$$= \int_0^1 \sqrt{1+\frac{9}{4}y}\,dy + \int_0^4 \sqrt{1+\frac{9}{4}y}\,dy = \frac{13\sqrt{13}-8}{27} + \frac{8}{27}\left[\left(1+\frac{9}{4}y\right)^{3/2}\right]_0^4 \qquad \text{[from part (b)]}$$

$$= \frac{13\sqrt{13}-8}{27} + \frac{8}{27}\left(10\sqrt{10}-1\right) = \frac{13\sqrt{13}+80\sqrt{10}-16}{27}$$

41. $y = 2x^{3/2} \;\Rightarrow\; y' = 3x^{1/2} \;\Rightarrow\; 1+(y')^2 = 1+9x$. The arc length function with starting point $P_0(1,2)$ is $s(x) = \int_1^x \sqrt{1+9t}\,dt = \left[\frac{2}{27}(1+9t)^{3/2}\right]_1^x = \frac{2}{27}\left[(1+9x)^{3/2} - 10\sqrt{10}\right]$.

42. (a) $y = f(x) = \ln(\sin x) \;\Rightarrow\; y' = \dfrac{1}{\sin x}\cdot\cos x = \cot x \;\Rightarrow\; 1+(y')^2 = 1+\cot^2 x = \csc^2 x \;\Rightarrow$ $\sqrt{1+(y')^2} = \sqrt{\csc^2 x} = |\csc x|$. Therefore,

$$s(x) = \int_{\pi/2}^x \sqrt{1+[f'(t)]^2}\,dt = \int_{\pi/2}^x \csc t\,dt = \Big[\ln|\csc t - \cos t|\Big]_{\pi/2}^x$$

$$= \ln|\csc x - \cot x| - \ln|1-0| = \ln(\csc x - \cot x)$$

(b) Note that s is increasing on $(0,\pi)$ and that $x=0$ and $x=\pi$ are vertical asymptotes for both f and s.

The arc length function $s(x)$ uses $x = \pi/2$ as a starting point, so when $x < \pi/2$, it gives a length of arc moving in the negative x direction. Thus, $s(x)$ is negative when $x < \pi/2$.

43. $y = \sin^{-1}x + \sqrt{1-x^2} \;\Rightarrow\; y' = \dfrac{1}{\sqrt{1-x^2}} - \dfrac{x}{\sqrt{1-x^2}} = \dfrac{1-x}{\sqrt{1-x^2}} \;\Rightarrow$

$1+(y')^2 = 1+\dfrac{(1-x)^2}{1-x^2} = \dfrac{1-x^2+1-2x+x^2}{1-x^2} = \dfrac{2-2x}{1-x^2} = \dfrac{2(1-x)}{(1+x)(1-x)} = \dfrac{2}{1+x} \;\Rightarrow$

$\sqrt{1+(y')^2} = \sqrt{\dfrac{2}{1+x}}$. Thus, the arc length function with starting point $(0,1)$ is given by

$$s(x) = \int_0^x \sqrt{1+[f'(t)]^2}\,dt = \int_0^x \sqrt{\frac{2}{1+t}}\,dt = \sqrt{2}\left[2\sqrt{1+t}\,\right]_0^x = 2\sqrt{2}\left(\sqrt{1+x}-1\right).$$

44. (a) $s(x) = \int_a^x \sqrt{1+[f'(t)]^2}\,dt$ and $s(x) = \int_0^x \sqrt{3t+5}\,dt \;\Rightarrow\; 1+[f'(t)]^2 = 3t+5 \;\Rightarrow\; [f'(t)]^2 = 3t+4 \;\Rightarrow$ $f'(t) = \sqrt{3t+4}$ [since f is increasing]. So $f(t) = \int(3t+4)^{1/2}\,dt = \frac{2}{3}\cdot\frac{1}{3}(3t+4)^{3/2} + C$ and since f has y-intercept 2, $f(0) = \frac{2}{9}\cdot 8 + C$ and $f(0) = 2 \;\Rightarrow\; C = 2 - \frac{16}{9} = \frac{2}{9}$. Thus, $f(t) = \frac{2}{9}(3t+4)^{3/2} + \frac{2}{9}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $s(x) = \int_0^x \sqrt{3t+5}\,dt = \left[\frac{2}{9}(3t+5)^{3/2}\right]_0^x = \frac{2}{9}(3x+5)^{3/2} - \frac{2}{9}(5)^{3/2}$.

$s(x) = 3 \;\Leftrightarrow\; \frac{2}{9}(3x+5)^{3/2} = 3 + \frac{2}{9}(5\sqrt{5}) \;\Leftrightarrow\; (3x+5)^{3/2} = \frac{27}{2} + 5\sqrt{5} \;\Leftrightarrow\; 3x+5 = \left(\frac{27}{2} + 5\sqrt{5}\right)^{2/3} \;\Rightarrow$

$x_1 = \frac{1}{3}\left[\left(\frac{27}{2} + 5\sqrt{5}\right)^{2/3} - 5\right]$. Thus, the point on the graph of f that is 3 units along the curve from the y-intercept is $(x_1, f(x_1)) \approx (1.159, 4.765)$.

45. The prey hits the ground when $y = 0 \;\Leftrightarrow\; 180 - \frac{1}{45}x^2 = 0 \;\Leftrightarrow\; x^2 = 45 \cdot 180 \;\Rightarrow\; x = \sqrt{8100} = 90$, since x must be positive. $y' = -\frac{2}{45}x \;\Rightarrow\; 1 + (y')^2 = 1 + \frac{4}{45^2}x^2$, so the distance traveled by the prey is

$$L = \int_0^{90} \sqrt{1 + \frac{4}{45^2}x^2}\,dx = \int_0^4 \sqrt{1+u^2}\left(\tfrac{45}{2}\,du\right) \qquad \begin{bmatrix} u = \frac{2}{45}x, \\ du = \frac{2}{45}\,dx \end{bmatrix}$$

$$\overset{21}{=} \tfrac{45}{2}\left[\tfrac{1}{2}u\sqrt{1+u^2} + \tfrac{1}{2}\ln\left(u + \sqrt{1+u^2}\right)\right]_0^4 = \tfrac{45}{2}\left[2\sqrt{17} + \tfrac{1}{2}\ln\left(4+\sqrt{17}\right)\right] = 45\sqrt{17} + \tfrac{45}{4}\ln\left(4+\sqrt{17}\right) \approx 209.1 \text{ m}$$

46. $y = 150 - \frac{1}{40}(x-50)^2 \;\Rightarrow\; y' = -\frac{1}{20}(x-50) \;\Rightarrow\; 1 + (y')^2 = 1 + \frac{1}{20^2}(x-50)^2$, so the distance traveled by the kite is

$$L = \int_0^{80} \sqrt{1 + \frac{1}{20^2}(x-50)^2}\,dx = \int_{-5/2}^{3/2} \sqrt{1+u^2}\,(20\,du) \qquad \begin{bmatrix} u = \frac{1}{20}(x-50), \\ du = \frac{1}{20}\,dx \end{bmatrix}$$

$$\overset{21}{=} 20\left[\tfrac{1}{2}u\sqrt{1+u^2} + \tfrac{1}{2}\ln\left(u+\sqrt{1+u^2}\right)\right]_{-5/2}^{3/2} = 10\left[\tfrac{3}{2}\sqrt{\tfrac{13}{4}} + \ln\left(\tfrac{3}{2} + \sqrt{\tfrac{13}{4}}\right) + \tfrac{5}{2}\sqrt{\tfrac{29}{4}} - \ln\left(-\tfrac{5}{2} + \sqrt{\tfrac{29}{4}}\right)\right]$$

$$= \tfrac{15}{2}\sqrt{13} + \tfrac{25}{2}\sqrt{29} + 10\ln\left(\tfrac{3+\sqrt{13}}{-5+\sqrt{29}}\right) \approx 122.8 \text{ ft}$$

47. The sine wave has amplitude 1 and period 14, since it goes through two periods in a distance of 28 in., so its equation is $y = 1\sin\left(\frac{2\pi}{14}x\right) = \sin\left(\frac{\pi}{7}x\right)$. The width w of the flat metal sheet needed to make the panel is the arc length of the sine curve from $x = 0$ to $x = 28$. We set up the integral to evaluate w using the arc length formula with $\frac{dy}{dx} = \frac{\pi}{7}\cos\left(\frac{\pi}{7}x\right)$:

$L = \int_0^{28} \sqrt{1 + \left[\frac{\pi}{7}\cos\left(\frac{\pi}{7}x\right)\right]^2}\,dx = 2\int_0^{14} \sqrt{1 + \left[\frac{\pi}{7}\cos\left(\frac{\pi}{7}x\right)\right]^2}\,dx$. This integral would be very difficult to evaluate exactly, so we use a CAS, and find that $L \approx 29.36$ inches.

48. (a) $y = a\cosh\left(\frac{x}{a}\right) \;\Rightarrow\; \frac{dy}{dx} = \sinh\left(\frac{x}{a}\right) \;\Rightarrow\; 1 + \left(\frac{dy}{dx}\right)^2 = 1 + \sinh^2\left(\frac{x}{a}\right) = \cosh^2\left(\frac{x}{a}\right)$. So

$$L = \int_c^d \sqrt{\cosh^2\left(\frac{x}{a}\right)}\,dx = \int_c^d \cosh\left(\frac{x}{a}\right)dx = \left[a\sinh\left(\frac{x}{a}\right)\right]_c^d = a\left[\sinh\left(\frac{d}{a}\right) - \sinh\left(\frac{c}{a}\right)\right].$$

(b) $A = \int_c^d a\,\cosh\left(\frac{x}{a}\right)dx = \left[a^2\sinh\left(\frac{x}{a}\right)\right]_c^d = a^2\left[\sinh\left(\frac{d}{a}\right) - \sinh\left(\frac{c}{a}\right)\right]$. The ratio of the area under the catenary to its arc length is $\dfrac{A}{L} = \dfrac{a^2\left[\sinh\left(\frac{d}{a}\right) - \sinh\left(\frac{c}{a}\right)\right]}{a\left[\sinh\left(\frac{d}{a}\right) - \sinh\left(\frac{c}{a}\right)\right]} = a$. Thus, the ratio of the area under the catenary to its arc length over any interval is a, a constant, that does not depend on the interval.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. $y = c + a\cosh\left(\dfrac{x}{a}\right) \quad\Rightarrow\quad \dfrac{dy}{dx} = \sinh\left(\dfrac{x}{a}\right) \quad\Rightarrow\quad 1 + \left(\dfrac{dy}{dx}\right)^2 = 1 + \sinh^2\left(\dfrac{x}{a}\right) = \cosh^2\left(\dfrac{x}{a}\right)$. So

$L = \displaystyle\int_{-25}^{25} \sqrt{\cosh^2\left(\frac{x}{a}\right)}\,dx = 2\int_0^{25} \cosh\left(\frac{x}{a}\right)dx = 2\left[a\sinh\left(\frac{x}{a}\right)\right]_0^{25} = 2a\sinh\left(\frac{25}{a}\right)$. Now, $L = 51 \quad\Rightarrow$

$51 = 2a\sinh\left(\dfrac{25}{a}\right)$. From the figure, we see that $y = 51$ intersects

$y = 2x\sinh\left(\dfrac{25}{x}\right)$ at $x \approx 72.3843$ for $x > 0$. So $a \approx 72.3843$. At $x = 0$,

$y = c + a$, so $c + a = 20 \quad\Rightarrow\quad c = 20 - a$. Thus, the wire should be attached

at a distance of $y = c + a\cosh\left(\dfrac{25}{a}\right) = 20 - a + a\cosh\left(\dfrac{25}{a}\right) \approx 24.36$ ft

above the ground.

50. Let $y = a - b\cosh cx$, where $a = 211.49$, $b = 20.96$, and $c = 0.03291765$. Then $y' = -bc\sinh cx \quad\Rightarrow$

$1 + (y')^2 = 1 + b^2c^2\sinh^2(cx)$. So $L = \int_{-91.2}^{91.2} \sqrt{1 + b^2c^2\sinh^2(cx)}\,dx \approx 451.137 \approx 451$, to the nearest meter.

51. $f(x) = \frac{1}{4}e^x + e^{-x} \quad\Rightarrow\quad f'(x) = \frac{1}{4}e^x - e^{-x} \quad\Rightarrow$

$1 + [f'(x)]^2 = 1 + \left(\frac{1}{4}e^x - e^{-x}\right)^2 = 1 + \frac{1}{16}e^{2x} - \frac{1}{2} + e^{-2x} = \frac{1}{16}e^{2x} + \frac{1}{2} + e^{-2x} = \left(\frac{1}{4}e^x + e^{-x}\right)^2 = [f(x)]^2$. The arc length of the curve $y = f(x)$ on the interval $[a, b]$ is $L = \int_a^b \sqrt{1 + [f'(x)]^2}\,dx = \int_a^b \sqrt{[f(x)]^2}\,dx = \int_a^b f(x)\,dx$, which is the area under the curve $y = f(x)$ on the interval $[a, b]$.

52. By symmetry, the length of the curve in each quadrant is the same,

so we'll find the length in the first quadrant and multiply by 4.

$x^{2k} + y^{2k} = 1 \quad\Rightarrow\quad y^{2k} = 1 - x^{2k} \quad\Rightarrow\quad y = (1 - x^{2k})^{1/(2k)}$

(in the first quadrant), so we use the arc length formula with

$\dfrac{dy}{dx} = \dfrac{1}{2k}(1 - x^{2k})^{1/(2k)-1}(-2kx^{2k-1}) = -x^{2k-1}(1 - x^{2k})^{1/(2k)-1}$

The total length is therefore

$$L_{2k} = 4\int_0^1 \sqrt{1 + [-x^{2k-1}(1 - x^{2k})^{1/(2k)-1}]^2}\,dx = 4\int_0^1 \sqrt{1 + x^{2(2k-1)}(1 - x^{2k})^{1/k-2}}\,dx$$

Now from the graph, we see that as k increases, the "corners" of these fat circles get closer to the points $(\pm1, \pm1)$ and $(\pm1, \mp1)$, and the "edges" of the fat circles approach the lines joining these four points. It seems plausible that as $k \to \infty$, the total length of the fat circle with $n = 2k$ will approach the length of the perimeter of the square with sides of length 2. This is supported by taking the limit as $k \to \infty$ of the equation of the fat circle in the first quadrant: $\lim\limits_{k\to\infty}(1 - x^{2k})^{1/(2k)} = 1$ for $0 \le x < 1$. So we guess that $\lim\limits_{k\to\infty} L_{2k} = 4 \cdot 2 = 8$.

53. $y = \int_1^x \sqrt{t^3 - 1}\,dt \quad\Rightarrow\quad dy/dx = \sqrt{x^3 - 1}$ [by FTC1] $\quad\Rightarrow\quad 1 + (dy/dx)^2 = 1 + \left(\sqrt{x^3 - 1}\right)^2 = x^3 \quad\Rightarrow$

$L = \int_1^4 \sqrt{x^3}\,dx = \int_1^4 x^{3/2}\,dx = \frac{2}{5}\left[x^{5/2}\right]_1^4 = \frac{2}{5}(32 - 1) = \frac{62}{5} = 12.4$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

DISCOVERY PROJECT Arc Length Contest

For advice on how to run the contest and a list of student entries, see the article "Arc Length Contest" by Larry Riddle in *The College Mathematics Journal*, Volume 29, No. 4, September 1998, pages 314–320.

8.2 Area of a Surface of Revolution

1. (a) $y = \sqrt[3]{x} = x^{1/3} \quad \Rightarrow \quad dy/dx = \frac{1}{3}x^{-2/3} \quad \Rightarrow \quad ds = \sqrt{1+(dy/dx)^2}\,dx = \sqrt{1+\frac{1}{9}x^{-4/3}}\,dx.$

By (7), $S = \int 2\pi y\,ds = \int_1^8 2\pi\sqrt[3]{x}\sqrt{1+\frac{1}{9}x^{-4/3}}\,dx.$

(b) $y = \sqrt[3]{x} \quad \Rightarrow \quad x = y^3 \quad \Rightarrow \quad dx/dy = 3y^2 \quad \Rightarrow \quad ds = \sqrt{1+(dx/dy)^2}\,dy = \sqrt{1+9y^4}\,dy.$

When $x = 1$, $y = 1$ and when $x = 8$, $y = 2$. Thus, $S = \int 2\pi y\,ds = \int_1^2 2\pi y\sqrt{1+9y^4}\,dy.$

2. (a) $x^2 = e^y \quad \Rightarrow \quad y = \ln x^2 = 2\ln x$ [for $x > 0$] $\quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{2}{x} \quad \Rightarrow \quad ds = \sqrt{1+\left(\dfrac{dy}{dx}\right)^2}\,dx = \sqrt{1+\dfrac{4}{x^2}}\,dx.$

By (7), $S = \displaystyle\int 2\pi y\,ds = \int_1^e 4\pi\ln x\sqrt{1+\frac{4}{x^2}}\,dx.$

(b) $x^2 = e^y \quad \Rightarrow \quad x = \sqrt{e^y} \quad \Rightarrow \quad e^{y/2}$ [positive since $x > 0$] $\quad \Rightarrow \quad dx/dy = \frac{1}{2}e^{y/2} \quad \Rightarrow$

$ds = \sqrt{1+(dx/dy)^2}\,dy = \sqrt{1+\frac{1}{4}e^y}\,dy.$ When $x = 1$, $y = 0$ and when $x = e$, $y = 2$. Thus,

$S = \int 2\pi y\,ds = \int_0^2 2\pi y\sqrt{1+\frac{1}{4}e^y}\,dy.$

3. (a) $x = \ln(2y+1) \quad \Rightarrow \quad e^x = 2y+1 \quad \Rightarrow \quad y = \frac{1}{2}e^x - \frac{1}{2} \quad \Rightarrow \quad dy/dx = \frac{1}{2}e^x \quad \Rightarrow$

$ds = \sqrt{1+(dy/dx)^2}\,dx = \sqrt{1+\frac{1}{4}e^{2x}}\,dx.$ When $y = 0$, $x = 0$ and when $y = 1$, $x = \ln 3$. Thus, by (7),

$S = \int 2\pi y\,ds = \int_0^{\ln 3} 2\pi\left(\frac{1}{2}e^x - \frac{1}{2}\right)\sqrt{1+\frac{1}{4}e^{2x}}\,dx = \int_0^{\ln 3}\pi(e^x-1)\sqrt{1+\frac{1}{4}e^{2x}}\,dx.$

(b) $x = \ln(2y+1) \quad \Rightarrow \quad dx/dy = \dfrac{2}{2y+1} \quad \Rightarrow \quad ds = \sqrt{1+(dx/dy)^2}\,dy = \sqrt{1+4/(2y+1)^2}\,dy.$ By (7),

$S = \int 2\pi y\,ds = \int_0^1 2\pi y\sqrt{1+4/(2y+1)^2}\,dy.$

4. (a) $y = \tan^{-1}x \quad \Rightarrow \quad dy/dx = 1/(1+x^2) \quad \Rightarrow \quad ds = \sqrt{1+(dy/dx)^2}\,dx = \sqrt{1+1/(1+x^2)^2}\,dx.$ By (7),

$S = \int 2\pi y\,ds = \int_0^1 2\pi\tan^{-1}x\sqrt{1+1/(1+x^2)^2}\,dx.$

(b) $y = \tan^{-1}x \quad \Rightarrow \quad x = \tan y \quad \Rightarrow \quad dx/dy = \sec^2 y \quad \Rightarrow \quad ds = \sqrt{1+(dx/dy)^2}\,dy = \sqrt{1+\sec^4 y}\,dy.$ When $x = 0$,

$y = 0$ and when $x = 1$, $y = \frac{\pi}{4}$. Thus $S = \int 2\pi y\,ds = \int_0^{\pi/4} 2\pi y\sqrt{1+\sec^4 y}\,dy.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. (a) $xy = 4 \quad \Rightarrow \quad y = \dfrac{4}{x} = 4x^{-1} \quad \Rightarrow \quad dy/dx = -4x^{-2} \quad \Rightarrow \quad ds = \sqrt{1 + (dy/dx)^2}\,dx = \sqrt{1 + 16/x^4}\,dx.$

By (8), $S = \int 2\pi x\,ds = \int_1^8 2\pi x\sqrt{1 + 16/x^4}\,dx.$

(b) $xy = 4 \quad \Rightarrow \quad x = \dfrac{4}{y} = 4y^{-1} \quad \Rightarrow \quad dx/dy = -4y^{-2} \quad \Rightarrow \quad ds = \sqrt{1 + (dx/dy)^2}\,dy = \sqrt{1 + 16/y^4}\,dy.$

When $x = 1$, $y = 4$ and when $x = 8$, $y = \frac{1}{2}$. By (8), $S = \displaystyle\int 2\pi x\,ds = \int_{1/2}^{4} \frac{8\pi}{y}\sqrt{1 + 16/y^4}\,dy.$

6. (a) $y = (x+1)^4 \quad \Rightarrow \quad dy/dx = 4(x+1)^3 \quad \Rightarrow \quad ds = \sqrt{1 + (dy/dx)^2}\,dx = \sqrt{1 + 16(x+1)^6}\,dx$. By (8),

$S = \int 2\pi x\,ds = \int_0^2 2\pi x\sqrt{1 + 16(x+1)^6}\,dx.$

(b) $y = (x+1)^4 \quad \Rightarrow \quad x = y^{1/4} - 1 \quad \Rightarrow \quad \dfrac{dx}{dy} = \dfrac{1}{4}y^{-3/4} \quad \Rightarrow \quad ds = \sqrt{1 + (dx/dy)^2}\,dy = \sqrt{1 + \frac{1}{16}y^{-3/2}}\,dy.$

When $x = 0$, $y = 1$ and when $x = 2$, $y = 81$. By (8), $S = \int 2\pi x\,ds = \int_1^{81} 2\pi(y^{1/4} - 1)\sqrt{1 + \frac{1}{16}y^{-3/2}}\,dy.$

7. (a) $y = 1 + \sin x \quad \Rightarrow \quad dy/dx = \cos x \quad \Rightarrow \quad ds = \sqrt{1 + (dy/dx)^2}\,dx = \sqrt{1 + \cos^2 x}\,dx.$

By (8), $S = \int 2\pi x\,ds = \int_0^{\pi/2} 2\pi x\sqrt{1 + \cos^2 x}\,dx.$

(b) $y = 1 + \sin x \quad \Rightarrow \quad x = \sin^{-1}(y-1) \quad \Rightarrow \quad \dfrac{dx}{dy} = \dfrac{1}{\sqrt{1 - (y-1)^2}} \quad \Rightarrow$

$ds = \sqrt{1 + \left(\dfrac{dx}{dy}\right)^2}\,dy = \sqrt{1 + \dfrac{1}{1 - (y-1)^2}}\,dy$. When $x = 0$, $y = 1$ and when $x = \pi/2$, $y = 2$.

By (8), $S = \displaystyle\int 2\pi x\,ds = \int_1^2 2\pi \sin^{-1}(y-1)\sqrt{1 + \frac{1}{1 - (y-1)^2}}\,dy$ or $\displaystyle\int_1^2 2\pi \sin^{-1}(y-1)\sqrt{1 + \frac{1}{2y - y^2}}\,dy.$

8. (a) $x = e^{2y} \quad \Rightarrow \quad y = \dfrac{1}{2}\ln x \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{1}{2x} \quad \Rightarrow \quad ds = \sqrt{1 + \left(\dfrac{dy}{dx}\right)^2}\,dx = \sqrt{1 + \dfrac{1}{4x^2}}\,dx.$

When $y = 0$, $x = 1$ and when $y = 2$, $x = e^4$. By (8), $S = \displaystyle\int 2\pi x\,ds = \int_1^{e^4} 2\pi x\sqrt{1 + \frac{1}{4x^2}}\,dx.$

(b) $x = e^{2y} \quad \Rightarrow \quad dx/dy = 2e^{2y} \quad \Rightarrow \quad ds = \sqrt{1 + (dx/dy)^2}\,dy = \sqrt{1 + 4e^{4y}}\,dy.$

By (8), $S = \int 2\pi x\,ds = \int_0^2 2\pi e^{2y}\sqrt{1 + 4e^{4y}}\,dy.$

9. $y = x^3 \quad \Rightarrow \quad y' = 3x^2$. So

$$S = \int_0^2 2\pi y\,\sqrt{1 + (y')^2}\,dx = 2\pi\int_0^2 x^3\,\sqrt{1 + 9x^4}\,dx$$

$$= \tfrac{2\pi}{36}\int_1^{145}\sqrt{u}\,du \qquad [u = 1 + 9x^4,\ du = 36x^3\,dx]$$

$$= \tfrac{\pi}{18}\left[\tfrac{2}{3}u^{3/2}\right]_1^{145} = \tfrac{\pi}{27}\left(145\sqrt{145} - 1\right)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $y=\sqrt{5-x} \quad\Rightarrow\quad y'=\frac{1}{2}(5-x)^{-1/2}(-1)=-1/(2\sqrt{5-x}\,)$. So

$$S=\int_3^5 2\pi y\sqrt{1+(y')^2}\,dx=\int_3^5 2\pi\sqrt{5-x}\sqrt{1+\frac{1}{4(5-x)}}\,dx=2\pi\int_3^5\sqrt{5-x+\tfrac{1}{4}}\,dx$$

$$=2\pi\int_3^5\sqrt{\tfrac{21}{4}-x}\,dx=2\pi\int_{9/4}^{1/4}\sqrt{u}\,(-du)\qquad\begin{bmatrix}u=\frac{21}{4}-x,\\ du=-dx\end{bmatrix}$$

$$=2\pi\int_{1/4}^{9/4}u^{1/2}\,du=2\pi\left[\tfrac{2}{3}u^{3/2}\right]_{1/4}^{9/4}=\tfrac{4\pi}{3}\left(\tfrac{27}{8}-\tfrac{1}{8}\right)=\tfrac{13\pi}{3}$$

11. $y^2=x+1 \quad\Rightarrow\quad y=\sqrt{x+1}$ (for $0\le x\le 3$ and $1\le y\le 2$) $\quad\Rightarrow\quad y'=1/(2\sqrt{x+1}\,)$. So

$$S=\int_0^3 2\pi y\sqrt{1+(y')^2}\,dx=2\pi\int_0^3\sqrt{x+1}\sqrt{1+\frac{1}{4(x+1)}}\,dx=2\pi\int_0^3\sqrt{x+1+\tfrac{1}{4}}\,dx$$

$$=2\pi\int_0^3\sqrt{x+\tfrac{5}{4}}\,dx=2\pi\int_{5/4}^{17/4}\sqrt{u}\,du\qquad\begin{bmatrix}u=x+\frac{5}{4},\\ du=dx\end{bmatrix}$$

$$=2\pi\left[\tfrac{2}{3}u^{3/2}\right]_{5/4}^{17/4}=2\pi\cdot\frac{2}{3}\left(\frac{17^{3/2}}{8}-\frac{5^{3/2}}{8}\right)=\frac{\pi}{6}(17\sqrt{17}-5\sqrt{5}).$$

12. $y=\sqrt{1+e^x} \quad\Rightarrow\quad y'=\frac{1}{2}(1+e^x)^{-1/2}(e^x)=\dfrac{e^x}{2\sqrt{1+e^x}} \quad\Rightarrow$

$$\sqrt{1+(y')^2}=\sqrt{1+\frac{e^{2x}}{4(1+e^x)}}=\sqrt{\frac{4+4e^x+e^{2x}}{4(1+e^x)}}=\sqrt{\frac{(e^x+2)^2}{4(1+e^x)}}=\frac{e^x+2}{2\sqrt{1+e^x}}.\text{ So}$$

$$S=\int_0^1 2\pi y\sqrt{1+(y')^2}\,dx=2\pi\int_0^1\sqrt{1+e^x}\,\frac{e^x+2}{2\sqrt{1+e^x}}\,dx=\pi\int_0^1(e^x+2)\,dx$$

$$=\pi\left[e^x+2x\right]_0^1=\pi[(e+2)-(1+0)]=\pi(e+1)$$

13. $y=\cos\left(\frac{1}{2}x\right) \quad\Rightarrow\quad y'=-\frac{1}{2}\sin\left(\frac{1}{2}x\right)$. So

$$S=\int_0^\pi 2\pi y\sqrt{1+(y')^2}\,dx=2\pi\int_0^\pi\cos\left(\tfrac{1}{2}x\right)\sqrt{1+\tfrac{1}{4}\sin^2\left(\tfrac{1}{2}x\right)}\,dx$$

$$=2\pi\int_0^1\sqrt{1+\tfrac{1}{4}u^2}\,(2\,du)\qquad\begin{bmatrix}u=\sin\left(\frac{1}{2}x\right),\\ du=\frac{1}{2}\cos\left(\frac{1}{2}x\right)dx\end{bmatrix}$$

$$=2\pi\int_0^1\sqrt{4+u^2}\,du\overset{21}{=}2\pi\left[\frac{u}{2}\sqrt{4+u^2}+2\ln\left(u+\sqrt{4+u^2}\right)\right]_0^1$$

$$=2\pi\left[\left(\tfrac{1}{2}\sqrt{5}+2\ln(1+\sqrt{5}\,)\right)-(0+2\ln 2)\right]=\pi\sqrt{5}+4\pi\ln\left(\frac{1+\sqrt{5}}{2}\right)$$

14. $y=\dfrac{x^3}{6}+\dfrac{1}{2x} \quad\Rightarrow\quad \dfrac{dy}{dx}=\dfrac{x^2}{2}-\dfrac{1}{2x^2} \quad\Rightarrow\quad \sqrt{1+\left(\dfrac{dy}{dx}\right)^2}=\sqrt{\dfrac{x^4}{4}+\dfrac{1}{2}+\dfrac{1}{4x^4}}=\sqrt{\left(\dfrac{x^2}{2}+\dfrac{1}{2x^2}\right)^2}=\dfrac{x^2}{2}+\dfrac{1}{2x^2}$. So

$$S=\int_{1/2}^1 2\pi\left(\frac{x^3}{6}+\frac{1}{2x}\right)\left(\frac{x^2}{2}+\frac{1}{2x^2}\right)dx=2\pi\int_{1/2}^1\left(\frac{x^5}{12}+\frac{x}{12}+\frac{x}{4}+\frac{1}{4x^3}\right)dx$$

$$=2\pi\int_{1/2}^1\left(\frac{x^5}{12}+\frac{x}{3}+\frac{x^{-3}}{4}\right)dx=2\pi\left[\frac{x^6}{72}+\frac{x^2}{6}-\frac{x^{-2}}{8}\right]_{1/2}^1$$

$$=2\pi\left[\left(\tfrac{1}{72}+\tfrac{1}{6}-\tfrac{1}{8}\right)-\left(\tfrac{1}{64\cdot 72}+\tfrac{1}{24}-\tfrac{1}{2}\right)\right]=2\pi\left(\tfrac{263}{512}\right)=\tfrac{263}{256}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $x = \frac{1}{3}(y^2+2)^{3/2} \quad \Rightarrow \quad dx/dy = \frac{1}{2}(y^2+2)^{1/2}(2y) = y\sqrt{y^2+2} \quad \Rightarrow \quad 1+(dx/dy)^2 = 1+y^2(y^2+2) = (y^2+1)^2.$

So $S = 2\pi\int_1^2 y(y^2+1)\,dy = 2\pi\left[\frac{1}{4}y^4+\frac{1}{2}y^2\right]_1^2 = 2\pi\left(4+2-\frac{1}{4}-\frac{1}{2}\right) = \frac{21\pi}{2}.$

16. $x = 1+2y^2 \quad \Rightarrow \quad 1+(dx/dy)^2 = 1+(4y)^2 = 1+16y^2.$

So $S = 2\pi\int_1^2 y\sqrt{1+16y^2}\,dy = \frac{\pi}{16}\int_1^2(16y^2+1)^{1/2}32y\,dy = \frac{\pi}{16}\left[\frac{2}{3}(16y^2+1)^{3/2}\right]_1^2 = \frac{\pi}{24}\left(65\sqrt{65}-17\sqrt{17}\right).$

17. $y = \frac{1}{3}x^{3/2} \quad \Rightarrow \quad y' = \frac{1}{2}x^{1/2} \quad \Rightarrow \quad 1+(y')^2 = 1+\frac{1}{4}x$. So

$$\begin{aligned}
S &= \int_0^{12} 2\pi x\sqrt{1+(y')^2}\,dx = 2\pi\int_0^{12} x\sqrt{1+\tfrac{1}{4}x}\,dx = 2\pi\int_0^{12} x\,\tfrac{1}{2}\sqrt{4+x}\,dx \\
&= \pi\int_4^{16}(u-4)\sqrt{u}\,du \qquad \begin{bmatrix} u = x+4, \\ du = dx \end{bmatrix} \\
&= \pi\int_4^{16}(u^{3/2}-4u^{1/2})\,du = \pi\left[\tfrac{2}{5}u^{5/2}-\tfrac{8}{3}u^{3/2}\right]_4^{16} = \pi\left[\left(\tfrac{2}{5}\cdot 1024-\tfrac{8}{3}\cdot 64\right)-\left(\tfrac{2}{5}\cdot 32-\tfrac{8}{3}\cdot 8\right)\right] \\
&= \pi\left(\tfrac{2}{5}\cdot 992-\tfrac{8}{3}\cdot 56\right) = \pi\left(\tfrac{5952-2240}{15}\right) = \tfrac{3712\pi}{15}
\end{aligned}$$

18. $x^{2/3}+y^{2/3} = 1,\ 0 \le y \le 1$. The curve is symmetric about the y-axis from $x=-1$ to $x=1$, so we'll use the portion of the curve from $x=0$ to $x=1$. $\quad y^{2/3} = 1-x^{2/3} \quad \Rightarrow \quad y = (1-x^{2/3})^{3/2} \quad \Rightarrow$

$y' = \frac{3}{2}(1-x^{2/3})^{1/2}\left(-\frac{2}{3}x^{-1/3}\right) = -\dfrac{\sqrt{1-x^{2/3}}}{x^{1/3}} \quad \Rightarrow \quad 1+(y')^2 = 1+\dfrac{1-x^{2/3}}{x^{2/3}} = \dfrac{x^{2/3}+1-x^{2/3}}{x^{2/3}} = x^{-2/3}$. So

$S = \int_0^1 2\pi x\sqrt{1+(y')^2}\,dx = 2\pi\int_0^1 x(x^{-1/3})\,dx = 2\pi\int_0^1 x^{2/3}\,dx = 2\pi\left[\frac{3}{5}x^{5/3}\right]_0^1 = 2\pi\left(\frac{3}{5}\right) = \frac{6\pi}{5}.$

19. $x = \sqrt{a^2-y^2} \quad \Rightarrow \quad dx/dy = \frac{1}{2}(a^2-y^2)^{-1/2}(-2y) = -y/\sqrt{a^2-y^2} \quad \Rightarrow$

$1+\left(\dfrac{dx}{dy}\right)^2 = 1+\dfrac{y^2}{a^2-y^2} = \dfrac{a^2-y^2}{a^2-y^2}+\dfrac{y^2}{a^2-y^2} = \dfrac{a^2}{a^2-y^2}$. So

$S = \displaystyle\int_0^{a/2} 2\pi\sqrt{a^2-y^2}\,\frac{a}{\sqrt{a^2-y^2}}\,dy = 2\pi\int_0^{a/2} a\,dy = 2\pi a\left[y\right]_0^{a/2} = 2\pi a\left(\frac{a}{2}-0\right) = \pi a^2.$

Note that this is $\frac{1}{4}$ the surface area of a sphere of radius a, and the length of the interval $y=0$ to $y=a/2$ is $\frac{1}{4}$ the length of the interval $y=-a$ to $y=a$.

20. $y = \frac{1}{4}x^2-\frac{1}{2}\ln x \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{x}{2}-\dfrac{1}{2x} \quad \Rightarrow \quad 1+\left(\dfrac{dy}{dx}\right)^2 = 1+\dfrac{x^2}{4}-\dfrac{1}{2}+\dfrac{1}{4x^2} = \dfrac{x^2}{4}+\dfrac{1}{2}+\dfrac{1}{4x^2} = \left(\dfrac{x}{2}+\dfrac{1}{2x}\right)^2$. So

$$\begin{aligned}
S &= \int_1^2 2\pi x\sqrt{\left(\frac{x}{2}+\frac{1}{2x}\right)^2}\,dx = 2\pi\int_1^2 x\left(\frac{x}{2}+\frac{1}{2x}\right)dx = \pi\int_1^2(x^2+1)\,dx = \pi\left[\tfrac{1}{3}x^3+x\right]_1^2 \\
&= \pi\left[\left(\tfrac{8}{3}+2\right)-\left(\tfrac{1}{3}+1\right)\right] = \tfrac{10}{3}\pi
\end{aligned}$$

21. $y = e^{-x^2} \quad \Rightarrow \quad dy/dx = -2xe^{-x^2} \quad \Rightarrow \quad ds = \sqrt{1+(dy/dx)^2}\,dx = \sqrt{1+4x^2e^{-2x^2}}\,dx$. By (7),

$S = \int 2\pi y\,ds = \int_{-1}^1 2\pi e^{-x^2}\sqrt{1+4x^2e^{-2x^2}}\,dx \approx 11.0753.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $xy = y^2 - 1 \quad \Rightarrow \quad x = y - y^{-1} \quad \Rightarrow \quad dx/dy = 1 + y^{-2} \quad \Rightarrow \quad ds = \sqrt{1 + (dx/dy)^2}\,dy = \sqrt{1 + (1 + y^{-2})^2}\,dy.$

By(7), $S = \int 2\pi y\,ds = \int_1^3 2\pi y\sqrt{1 + (1 + y^{-2})^2}\,dy \approx 40.8099.$

23. $x = y + y^3 \quad \Rightarrow \quad dx/dy = 1 + 3y^2 \quad \Rightarrow \quad ds = \sqrt{1 + (dx/dy)^2}\,dy = \sqrt{1 + (1 + 3y^2)^2}\,dy.$

By (8), $S = \int 2\pi x\,ds = \int_0^1 2\pi(y + y^3)\sqrt{1 + (1 + 3y^2)^2}\,dy \approx 13.5134.$

24. $y = x + \sin x \quad \Rightarrow \quad dy/dx = 1 + \cos x \quad \Rightarrow \quad ds = \sqrt{1 + (dy/dx)^2}\,dx = \sqrt{1 + (1 + \cos x)^2}\,dx.$

By (8), $S = \int 2\pi x\,ds = \int_0^{2\pi/3} 2\pi x\sqrt{1 + (1 + \cos x)^2}\,dx \approx 21.2980.$

25. $\ln y = x - y^2 \quad \Rightarrow \quad x = \ln y + y^2 \quad \Rightarrow \quad \dfrac{dx}{dy} = \dfrac{1}{y} + 2y \quad \Rightarrow \quad ds = \sqrt{1 + (dx/dy)^2}\,dy = \sqrt{1 + (y^{-1} + 2y)^2}\,dy.$

By (7), $S = \int 2\pi y\,ds = \int_1^4 2\pi y\sqrt{1 + (y^{-1} + 2y)^2}\,dy \approx 286.9239.$

26. $x = \cos^2 y \quad \Rightarrow \quad dx/dy = 2\cos y\,(-\sin y) \quad \Rightarrow \quad ds = \sqrt{1 + (dx/dy)^2}\,dy = \sqrt{1 + 4\sin^2 y\cos^2 y}\,dy.$

By (8), $S = \int 2\pi x\,ds = \int_0^{\pi/2} 2\pi\cos^2 y\sqrt{1 + 4\sin^2 y\cos^2 y}\,dy \approx 6.0008.$

27. $y = 1/x \quad \Rightarrow \quad ds = \sqrt{1 + (dy/dx)^2}\,dx = \sqrt{1 + (-1/x^2)^2}\,dx = \sqrt{1 + 1/x^4}\,dx \quad \Rightarrow$

$$S = \int_1^2 2\pi \cdot \frac{1}{x}\sqrt{1 + \frac{1}{x^4}}\,dx = 2\pi\int_1^2 \frac{\sqrt{x^4 + 1}}{x^3}\,dx = 2\pi\int_1^4 \frac{\sqrt{u^2 + 1}}{u^2}\left(\tfrac{1}{2}\,du\right) \qquad [u = x^2,\ du = 2x\,dx]$$

$$= \pi\int_1^4 \frac{\sqrt{1 + u^2}}{u^2}\,du \overset{24}{=} \pi\left[-\frac{\sqrt{1 + u^2}}{u} + \ln\left(u + \sqrt{1 + u^2}\right)\right]_1^4$$

$$= \pi\left[-\tfrac{\sqrt{17}}{4} + \ln\left(4 + \sqrt{17}\right) + \tfrac{\sqrt{2}}{1} - \ln\left(1 + \sqrt{2}\right)\right] = \frac{\pi}{4}\left[4\ln\left(\sqrt{17} + 4\right) - 4\ln\left(\sqrt{2} + 1\right) - \sqrt{17} + 4\sqrt{2}\right]$$

28. $y = \sqrt{x^2 + 1} \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{x}{\sqrt{x^2 + 1}} \quad \Rightarrow \quad ds = \sqrt{1 + \left(\dfrac{dy}{dx}\right)^2}\,dx = \sqrt{1 + \dfrac{x^2}{x^2 + 1}}\,dx \quad \Rightarrow$

$$S = \int_0^3 2\pi\sqrt{x^2 + 1}\sqrt{1 + \frac{x^2}{x^2 + 1}}\,dx = 2\pi\int_0^3 \sqrt{2x^2 + 1}\,dx = 2\sqrt{2}\,\pi\int_0^3 \sqrt{x^2 + \left(\tfrac{1}{\sqrt{2}}\right)^2}\,dx$$

$$\overset{21}{=} 2\sqrt{2}\pi\left[\tfrac{1}{2}x\sqrt{x^2 + \tfrac{1}{2}} + \tfrac{1}{4}\ln\left(x + \sqrt{x^2 + \tfrac{1}{2}}\right)\right]_0^3 = 2\sqrt{2}\pi\left[\tfrac{3}{2}\sqrt{9 + \tfrac{1}{2}} + \tfrac{1}{4}\ln\left(3 + \sqrt{9 + \tfrac{1}{2}}\right) - \tfrac{1}{4}\ln\tfrac{1}{\sqrt{2}}\right]$$

$$= 2\sqrt{2}\pi\left[\tfrac{3}{2}\sqrt{\tfrac{19}{2}} + \tfrac{1}{4}\ln\left(3 + \sqrt{\tfrac{19}{2}}\right) + \tfrac{1}{4}\ln\sqrt{2}\right] = 2\sqrt{2}\pi\left[\tfrac{3}{2}\tfrac{\sqrt{19}}{\sqrt{2}} + \tfrac{1}{4}\ln\left(3\sqrt{2} + \sqrt{19}\right)\right]$$

$$= 3\sqrt{19}\pi + \tfrac{\pi}{\sqrt{2}}\ln\left(3\sqrt{2} + \sqrt{19}\right)$$

29. $y = x^3$ and $0 \le y \le 1 \quad \Rightarrow \quad y' = 3x^2$ and $0 \le x \le 1$.

$$S = \int_0^1 2\pi x\sqrt{1 + (3x^2)^2}\,dx = 2\pi\int_0^3 \sqrt{1 + u^2}\,\tfrac{1}{6}\,du \qquad \begin{bmatrix} u = 3x^2, \\ du = 6x\,dx \end{bmatrix}$$

$$= \tfrac{\pi}{3}\int_0^3 \sqrt{1 + u^2}\,du \overset{21}{=} \text{[or use CAS]}\ \tfrac{\pi}{3}\left[\tfrac{1}{2}u\sqrt{1 + u^2} + \tfrac{1}{2}\ln\left(u + \sqrt{1 + u^2}\right)\right]_0^3$$

$$= \tfrac{\pi}{3}\left[\tfrac{3}{2}\sqrt{10} + \tfrac{1}{2}\ln\left(3 + \sqrt{10}\right)\right] = \tfrac{\pi}{6}\left[3\sqrt{10} + \ln\left(3 + \sqrt{10}\right)\right]$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $y = \ln(x+1)$, $0 \le x \le 1$. $ds = \sqrt{1 + \left(\dfrac{dy}{dx}\right)^2}\,dx = \sqrt{1 + \left(\dfrac{1}{x+1}\right)^2}\,dx$, so

$$S = \int_0^1 2\pi x \sqrt{1 + \frac{1}{(x+1)^2}}\,dx = \int_1^2 2\pi(u-1)\sqrt{1 + \frac{1}{u^2}}\,du \qquad [u = x+1,\ du = dx]$$

$$= 2\pi \int_1^2 u\,\frac{\sqrt{1+u^2}}{u}\,du - 2\pi \int_1^2 \frac{\sqrt{1+u^2}}{u}\,du = 2\pi \int_1^2 \sqrt{1+u^2}\,du - 2\pi \int_1^2 \frac{\sqrt{1+u^2}}{u}\,du$$

$$\overset{21,\,23}{=} \text{[or use CAS]} \quad 2\pi\left[\tfrac{1}{2}u\sqrt{1+u^2} + \tfrac{1}{2}\ln\left(u + \sqrt{1+u^2}\right)\right]_1^2 - 2\pi\left[\sqrt{1+u^2} - \ln\left(\frac{1+\sqrt{1+u^2}}{u}\right)\right]_1^2$$

$$= 2\pi\left[\sqrt{5} + \tfrac{1}{2}\ln\left(2+\sqrt{5}\right) - \tfrac{1}{2}\sqrt{2} - \tfrac{1}{2}\ln\left(1+\sqrt{2}\right)\right] - 2\pi\left[\sqrt{5} - \ln\left(\tfrac{1+\sqrt{5}}{2}\right) - \sqrt{2} + \ln\left(1+\sqrt{2}\right)\right]$$

$$= 2\pi\left[\tfrac{1}{2}\ln\left(2+\sqrt{5}\right) + \ln\left(\tfrac{1+\sqrt{5}}{2}\right) + \tfrac{\sqrt{2}}{2} - \tfrac{3}{2}\ln\left(1+\sqrt{2}\right)\right]$$

31. $y = \frac{1}{5}x^5 \;\Rightarrow\; dy/dx = x^4 \;\Rightarrow\; 1 + (dy/dx)^2 = 1 + x^8 \;\Rightarrow\; S = \int_0^5 2\pi\left(\frac{1}{5}x^5\right)\sqrt{1+x^8}\,dx$.

Let $f(x) = \frac{2}{5}\pi x^5\sqrt{1+x^8}$. Since $n = 10$, $\Delta x = \frac{5-0}{10} = \frac{1}{2}$. Then

$$\begin{aligned} S \approx S_{10} &= \tfrac{1/2}{3}[f(0) + 4f(0.5) + 2f(1) + 4f(1.5) + 2f(2) + 4f(2.5) + 2f(3) \\ &\qquad + 4f(3.5) + 2f(4) + 4f(4.5) + f(5)] \\ &\approx 1{,}230{,}507 \end{aligned}$$

The value of the integral produced by a calculator is approximately 1,227,192.

32. $y = x\ln x \;\Rightarrow\; dy/dx = x \cdot \dfrac{1}{x} + \ln x = 1 + \ln x \;\Rightarrow\; 1 + (dy/dx)^2 = 1 + (1+\ln x)^2 \;\Rightarrow$

$S = \int_1^2 2\pi x \ln x\,\sqrt{1 + (1+\ln x)^2}\,dx$. Let $f(x) = 2\pi x \ln x\,\sqrt{1+(1+\ln x)^2}$. Since $n = 10$, $\Delta x = \frac{2-1}{10} = \frac{1}{10}$. Then

$$\begin{aligned} S \approx S_{10} &= \tfrac{1/10}{3}[f(1) + 4f(1.1) + 2f(1.2) + 4f(1.3) + 2f(1.4) + 4f(1.5) + 2f(1.6) \\ &\qquad + 4f(1.7) + 2f(1.8) + 4f(1.9) + f(2)] \\ &\approx 7.248933 \end{aligned}$$

The value of the integral produced by a calculator is 7.248934 (to six decimal places).

33. $S = 2\pi \displaystyle\int_1^\infty y\sqrt{1 + \left(\frac{dy}{dx}\right)^2}\,dx = 2\pi \int_1^\infty \frac{1}{x}\sqrt{1 + \frac{1}{x^4}}\,dx = 2\pi \int_1^\infty \frac{\sqrt{x^4+1}}{x^3}\,dx$. Rather than trying to evaluate this integral, note that $\sqrt{x^4+1} > \sqrt{x^4} = x^2$ for $x > 0$. Thus, if the area is finite,

$S = 2\pi \displaystyle\int_1^\infty \frac{\sqrt{x^4+1}}{x^3}\,dx > 2\pi \int_1^\infty \frac{x^2}{x^3}\,dx = 2\pi \int_1^\infty \frac{1}{x}\,dx$. But we know that this integral diverges, so the area S is infinite.

34. $S = \int_0^\infty 2\pi y\,\sqrt{1 + (dy/dx)^2}\,dx = 2\pi \int_0^\infty e^{-x}\sqrt{1 + (-e^{-x})^2}\,dx \qquad [y = e^{-x},\ y' = -e^{-x}]$.

Evaluate $I = \int e^{-x}\sqrt{1 + (-e^{-x})^2}\,dx$ by using the substitution $u = -e^{-x}$, $du = e^{-x}\,dx$:

$I = \int \sqrt{1+u^2}\,du \overset{21}{=} \frac{1}{2}u\sqrt{1+u^2} + \frac{1}{2}\ln\left(u + \sqrt{1+u^2}\right) + C = \frac{1}{2}(-e^{-x})\sqrt{1+e^{-2x}} + \frac{1}{2}\ln\left(-e^{-x} + \sqrt{1+e^{-2x}}\right) + C$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Returning to the surface area integral, we have

$$S = 2\pi \lim_{t\to\infty} \int_0^t e^{-x}\sqrt{1+(-e^{-x})^2}\,dx = 2\pi \lim_{t\to\infty} \left[\tfrac{1}{2}(-e^{-x})\sqrt{1+e^{-2x}} + \tfrac{1}{2}\ln\left(-e^{-x}+\sqrt{1+e^{-2x}}\right)\right]_0^t$$

$$= 2\pi \lim_{t\to\infty} \left\{\left[\tfrac{1}{2}(-e^{-t})\sqrt{1+e^{-2t}} + \tfrac{1}{2}\ln\left(-e^{-t}+\sqrt{1+e^{-2t}}\right)\right] - \left[\tfrac{1}{2}(-1)\sqrt{1+1} + \tfrac{1}{2}\ln\left(-1+\sqrt{1+1}\right)\right]\right\}$$

$$= 2\pi\left\{\left[\tfrac{1}{2}(0)\sqrt{1} + \tfrac{1}{2}\ln\left(0+\sqrt{1}\right)\right] - \left[-\tfrac{1}{2}\sqrt{2} + \tfrac{1}{2}\ln\left(-1+\sqrt{2}\right)\right]\right\}$$

$$= 2\pi\left\{[0] + \tfrac{1}{2}\left[\sqrt{2} - \ln\left(\sqrt{2}-1\right)\right]\right\} = \pi\left[\sqrt{2} - \ln\left(\sqrt{2}-1\right)\right]$$

35. As seen in the exercise figure, the loop of the curve $3ay^2 = x(a-x)^2$ extends from $x=0$ to $x=a$. The top half of the loop is given by $y = \sqrt{\dfrac{1}{3a}x(a-x)^2} = \dfrac{1}{\sqrt{3a}}\sqrt{x}\,|a-x| = \dfrac{1}{\sqrt{3a}}\sqrt{x}(a-x)$, since $x \le a$. Now,

$$\frac{dy}{dx} = \frac{1}{\sqrt{3a}}\left[\sqrt{x}\,(-1) + \frac{1}{2\sqrt{x}}(a-x)\right] = \frac{1}{\sqrt{3a}}\left[-\frac{2x}{2\sqrt{x}} + \frac{a-x}{2\sqrt{x}}\right] = \frac{1}{\sqrt{3a}}\left[\frac{a-3x}{2\sqrt{x}}\right] \quad \Rightarrow$$

$$1 + \left(\frac{dy}{dx}\right)^2 = 1 + \frac{(a-3x)^2}{12ax} = \frac{12ax}{12ax} + \frac{a^2-6ax+9x^2}{12ax} = \frac{a^2+6ax+9x^2}{12ax} = \frac{(a+3x)^2}{12ax}.$$

(a) $$S = \int 2\pi y\,ds = 2\pi\int_0^a \frac{\sqrt{x}(a-x)}{\sqrt{3a}}\sqrt{\frac{(a+3x)^2}{12ax}}\,dx = 2\pi\int_0^a \frac{(a-x)(a+3x)}{6a}\,dx$$

$$= \frac{\pi}{3a}\int_0^a (a^2+2ax-3x^2)\,dx = \frac{\pi}{3a}\left[a^2x + ax^2 - x^3\right]_0^a = \frac{\pi}{3a}(a^3+a^3-a^3)$$

$$= \frac{\pi}{3a}\cdot a^3 = \frac{\pi a^2}{3}$$

Note that the top half of the loop has been rotated about the x-axis, producing the full surface.

(b) We must rotate the full loop about the y-axis, so we get double the area obtained by rotating the top half of the loop:

$$S = 2\int_{x=0}^a 2\pi x\,ds = 4\pi\int_0^a x\,\frac{a+3x}{\sqrt{12ax}}\,dx = \frac{4\pi}{2\sqrt{3a}}\int_0^a x^{1/2}(a+3x)\,dx = \frac{2\pi}{\sqrt{3a}}\int_0^a (ax^{1/2}+3x^{3/2})\,dx$$

$$= \frac{2\pi}{\sqrt{3a}}\left[\frac{2}{3}ax^{3/2} + \frac{6}{5}x^{5/2}\right]_0^a = \frac{2\pi\sqrt{3}}{3\sqrt{a}}\left(\frac{2}{3}a^{5/2} + \frac{6}{5}a^{5/2}\right) = \frac{2\pi\sqrt{3}}{3}\left(\frac{2}{3}+\frac{6}{5}\right)a^2 = \frac{2\pi\sqrt{3}}{3}\left(\frac{28}{15}\right)a^2$$

$$= \frac{56\pi\sqrt{3}a^2}{45}$$

36. In general, if the parabola $y = ax^2$, $-c \le x \le c$, is rotated about the y-axis, the surface area it generates is

$$S = \int 2\pi x\,ds = 2\pi\int_0^c x\sqrt{1+(2ax)^2}\,dx = 2\pi\int_0^{2ac} \frac{u}{2a}\sqrt{1+u^2}\,\frac{1}{2a}\,du \quad \begin{bmatrix} u = 2ax, \\ du = 2a\,dx \end{bmatrix}$$

$$= \frac{\pi}{4a^2}\int_0^{2ac} \left(1+u^2\right)^{1/2} 2u\,du = \frac{\pi}{4a^2}\left[\tfrac{2}{3}\left(1+u^2\right)^{3/2}\right]_0^{2ac} = \frac{\pi}{6a^2}\left[\left(1+4a^2c^2\right)^{3/2} - 1\right]$$

Here $2c = 10$ ft and $ac^2 = 2$ ft, so $c = 5$ and $a = \frac{2}{25}$. Thus, the surface area is

$$S = \tfrac{\pi}{6}\tfrac{625}{4}\left[\left(1+4\cdot\tfrac{4}{625}\cdot 25\right)^{3/2} - 1\right] = \tfrac{625\pi}{24}\left[\left(1+\tfrac{16}{25}\right)^{3/2} - 1\right]$$

$$= \tfrac{625\pi}{24}\left(\tfrac{41\sqrt{41}}{125} - 1\right) = \tfrac{5\pi}{24}\left(41\sqrt{41} - 125\right) \approx 90.01 \text{ ft}^2$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. (a) $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1 \quad\Rightarrow\quad \dfrac{y\,(dy/dx)}{b^2}=-\dfrac{x}{a^2} \quad\Rightarrow\quad \dfrac{dy}{dx}=-\dfrac{b^2x}{a^2y} \quad\Rightarrow$

$$1+\left(\frac{dy}{dx}\right)^2=1+\frac{b^4x^2}{a^4y^2}=\frac{b^4x^2+a^4y^2}{a^4y^2}=\frac{b^4x^2+a^4b^2\left(1-x^2/a^2\right)}{a^4b^2\left(1-x^2/a^2\right)}=\frac{a^4b^2+b^4x^2-a^2b^2x^2}{a^4b^2-a^2b^2x^2}$$

$$=\frac{a^4+b^2x^2-a^2x^2}{a^4-a^2x^2}=\frac{a^4-\left(a^2-b^2\right)x^2}{a^2(a^2-x^2)}$$

The ellipsoid's surface area is twice the area generated by rotating the first-quadrant portion of the ellipse about the x-axis. Thus,

$$S=2\int_0^a 2\pi y\sqrt{1+\left(\frac{dy}{dx}\right)^2}\,dx=4\pi\int_0^a \frac{b}{a}\sqrt{a^2-x^2}\,\frac{\sqrt{a^4-(a^2-b^2)x^2}}{a\sqrt{a^2-x^2}}\,dx=\frac{4\pi b}{a^2}\int_0^a \sqrt{a^4-(a^2-b^2)x^2}\,dx$$

$$=\frac{4\pi b}{a^2}\int_0^{a\sqrt{a^2-b^2}}\sqrt{a^4-u^2}\,\frac{du}{\sqrt{a^2-b^2}}\quad\left[u=\sqrt{a^2-b^2}\,x\right]\overset{30}{=}\frac{4\pi b}{a^2\sqrt{a^2-b^2}}\left[\frac{u}{2}\sqrt{a^4-u^2}+\frac{a^4}{2}\sin^{-1}\left(\frac{u}{a^2}\right)\right]_0^{a\sqrt{a^2-b^2}}$$

$$=\frac{4\pi b}{a^2\sqrt{a^2-b^2}}\left[\frac{a\sqrt{a^2-b^2}}{2}\sqrt{a^4-a^2(a^2-b^2)}+\frac{a^4}{2}\sin^{-1}\frac{\sqrt{a^2-b^2}}{a}\right]=2\pi\left[b^2+\frac{a^2b\sin^{-1}\dfrac{\sqrt{a^2-b^2}}{a}}{\sqrt{a^2-b^2}}\right]$$

(b) $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1 \quad\Rightarrow\quad \dfrac{x\,(dx/dy)}{a^2}=-\dfrac{y}{b^2} \quad\Rightarrow\quad \dfrac{dx}{dy}=-\dfrac{a^2y}{b^2x} \quad\Rightarrow$

$$1+\left(\frac{dx}{dy}\right)^2=1+\frac{a^4y^2}{b^4x^2}=\frac{b^4x^2+a^4y^2}{b^4x^2}=\frac{b^4a^2(1-y^2/b^2)+a^4y^2}{b^4a^2(1-y^2/b^2)}=\frac{a^2b^4-a^2b^2y^2+a^4y^2}{a^2b^4-a^2b^2y^2}$$

$$=\frac{b^4-b^2y^2+a^2y^2}{b^4-b^2y^2}=\frac{b^4-(b^2-a^2)y^2}{b^2(b^2-y^2)}$$

The oblate spheroid's surface area is twice the area generated by rotating the first-quadrant portion of the ellipse about the y-axis. Thus,

$$S=2\int_0^b 2\pi\,x\sqrt{1+\left(\frac{dx}{dy}\right)^2}\,dy=4\pi\int_0^b \frac{a}{b}\sqrt{b^2-y^2}\,\frac{\sqrt{b^4-(b^2-a^2)y^2}}{b\sqrt{b^2-y^2}}\,dy$$

$$=\frac{4\pi a}{b^2}\int_0^b \sqrt{b^4-\left(b^2-a^2\right)y^2}\,dy=\frac{4\pi a}{b^2}\int_0^b \sqrt{b^4+\left(a^2-b^2\right)\,y^2}\,dy \qquad \left[\text{since } a>b\right]$$

$$=\frac{4\pi a}{b^2}\int_0^{b\sqrt{a^2-b^2}}\sqrt{b^4+u^2}\,\frac{du}{\sqrt{a^2-b^2}} \qquad \left[u=\sqrt{a^2-b^2}\,y\right]$$

$$\overset{21}{=}\frac{4\pi a}{b^2\sqrt{a^2-b^2}}\left[\frac{u}{2}\sqrt{b^4+u^2}+\frac{b^4}{2}\ln\left(u+\sqrt{b^4+u^2}\right)\right]_0^{b\sqrt{a^2-b^2}}$$

$$=\frac{4\pi a}{b^2\sqrt{a^2-b^2}}\left\{\left[\frac{b\sqrt{a^2-b^2}}{2}(ab)+\frac{b^4}{2}\ln\left(b\sqrt{a^2-b^2}+ab\right)\right]-\left[0+\frac{b^4}{2}\ln(b^2)\right]\right\}$$

$$=\frac{4\pi a}{b^2\sqrt{a^2-b^2}}\left[\frac{ab^2\sqrt{a^2-b^2}}{2}+\frac{b^4}{2}\ln\frac{b\sqrt{a^2-b^2}+ab}{b^2}\right]=2\pi a^2+\frac{2\pi ab^2}{\sqrt{a^2-b^2}}\ln\frac{\sqrt{a^2-b^2}+a}{b}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. The upper half of the torus is generated by rotating the curve $(x-R)^2+y^2=r^2$, $y>0$, about the y-axis.

$$y\,\frac{dy}{dx}=-(x-R)\quad\Rightarrow\quad 1+\left(\frac{dy}{dx}\right)^2=1+\frac{(x-R)^2}{y^2}=\frac{y^2+(x-R)^2}{y^2}=\frac{r^2}{r^2-(x-R)^2}.\text{ Thus,}$$

$$S=2\int_{R-r}^{R+r}2\pi x\sqrt{1+\left(\frac{dy}{dx}\right)^2}\,dx=4\pi\int_{R-r}^{R+r}\frac{rx}{\sqrt{r^2-(x-R)^2}}\,dx=4\pi r\int_{-r}^{r}\frac{u+R}{\sqrt{r^2-u^2}}\,du\qquad\left[u=x-R\right]$$

$$=4\pi r\int_{-r}^{r}\frac{u\,du}{\sqrt{r^2-u^2}}+4\pi Rr\int_{-r}^{r}\frac{du}{\sqrt{r^2-u^2}}=4\pi r\cdot 0+8\pi Rr\int_0^r\frac{du}{\sqrt{r^2-u^2}}\qquad\left[\begin{array}{c}\text{since the first integrand is odd}\\ \text{and the second is even}\end{array}\right]$$

$$=8\pi Rr\left[\sin^{-1}(u/r)\right]_0^r=8\pi Rr\left(\tfrac{\pi}{2}\right)=4\pi^2Rr$$

39. (a) The analogue of $f(x_i^*)$ in the derivation of (4) is now $c-f(x_i^*)$, so

$$S=\lim_{n\to\infty}\sum_{i=1}^{n}2\pi[c-f(x_i^*)]\sqrt{1+[f'(x_i^*)]^2}\,\Delta x=\int_a^b 2\pi[c-f(x)]\sqrt{1+[f'(x)]^2}\,dx.$$

(b) $y=x^{1/2}\quad\Rightarrow\quad y'=\frac{1}{2}x^{-1/2}\quad\Rightarrow\quad 1+(y')^2=1+1/4x$, so by part (a), $S=\int_0^4 2\pi\left(4-\sqrt{x}\right)\sqrt{1+1/(4x)}\,dx$.

Using a CAS, we get $S=2\pi\ln\left(\sqrt{17}+4\right)+\frac{\pi}{6}\left(31\sqrt{17}+1\right)\approx 80.6095$.

40. $y=x^3\quad\Rightarrow\quad dy/dx=3x^2\quad\Rightarrow\quad ds=\sqrt{1+(dy/dx)^2}\,dx=\sqrt{1+9x^4}\,dx$. Also, $x=y^{1/3}\quad\Rightarrow$

$dx/dy=\frac{1}{3}y^{-2/3}\quad\Rightarrow\quad ds=\sqrt{1+(dx/dy)^2}\,dy=\sqrt{1+\frac{1}{9}y^{-4/3}}\,dy$. When $x=1$, $y=1$ and when $x=2$, $y=8$.

(a) Since $x>-1$ over the x-interval $[1,2]$, the area of the surface obtained by rotating the curve about $x=-1$ is given by

$S=\int 2\pi(x+1)\,ds=\int_1^2 2\pi(x+1)\sqrt{1+9x^4}\,dx\approx 115.91$.

Alternative method: $S=\int 2\pi(x+1)\,ds=\int_1^8 2\pi(y^{1/3}+1)\sqrt{1+\frac{1}{9}y^{-4/3}}\,dy\approx 115.91$.

(b) Since $x<4$ over the x-interval $[1,2]$, the area of the surface obtained by rotating the curve about $x=4$ is given by

$S=\int 2\pi(4-x)\,ds=\int_1^2 2\pi(4-x)\sqrt{1+9x^4}\,dx\approx 106.60$.

Alternative method: $S=\int 2\pi(4-x)\,ds=\int_1^8 2\pi(4-y^{1/3})\sqrt{1+\frac{1}{9}y^{-4/3}}\,dy\approx 106.60$.

(c) Since $y>\frac{1}{2}$ over the y-interval $[1,8]$, the area of the surface obtained by rotating the curve about $y=\frac{1}{2}$ is given by

$S=\int 2\pi\left(y-\frac{1}{2}\right)ds=\int_1^2 2\pi\left(x^3-\frac{1}{2}\right)\sqrt{1+9x^4}\,dx\approx 177.23$.

Alternative method: $S=\int 2\pi\left(y-\frac{1}{2}\right)ds=\int_1^8 2\pi\left(y-\frac{1}{2}\right)\sqrt{1+\frac{1}{9}y^{-4/3}}\,dy\approx 177.23$.

(d) Since $y<10$ over the y-interval $[1,8]$, the area of the surface obtained by rotating the curve about $y=10$ is given by

$S=\int 2\pi(10-y)\,ds=\int_1^2 2\pi(10-x^3)\sqrt{1+9x^4}\,dx\approx 245.52$.

Alternative method: $S=\int 2\pi(10-y)\,ds=\int_1^8 2\pi(10-y)\sqrt{1+\frac{1}{9}y^{-4/3}}\,dy\approx 245.52$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. For the upper semicircle, $y = \sqrt{r^2 - x^2}$, $dy/dx = -x/\sqrt{r^2 - x^2}$. The surface area generated is

$$S_1 = \int_{-r}^{r} 2\pi(r - y)\,ds = \int_{-r}^{r} 2\pi\left(r - \sqrt{r^2 - x^2}\right)\sqrt{1 + \frac{x^2}{r^2 - x^2}}\,dx$$

$$= 4\pi \int_0^r \left(r - \sqrt{r^2 - x^2}\right)\frac{r}{\sqrt{r^2 - x^2}}\,dx = 4\pi \int_0^r \left(\frac{r^2}{\sqrt{r^2 - x^2}} - r\right)dx$$

For the lower semicircle, $y = -\sqrt{r^2 - x^2}$ and $\dfrac{dy}{dx} = \dfrac{x}{\sqrt{r^2 - x^2}}$, so $S_2 = 4\pi \displaystyle\int_0^r \left(\frac{r^2}{\sqrt{r^2 - x^2}} + r\right)dx$.

Thus, the total area is $S = S_1 + S_2 = 8\pi \displaystyle\int_0^r \left(\frac{r^2}{\sqrt{r^2 - x^2}}\right)dx = 8\pi\left[r^2 \sin^{-1}\left(\frac{x}{r}\right)\right]_0^r = 8\pi r^2\left(\frac{\pi}{2}\right) = 4\pi^2 r^2$.

42. Rotate $y = \sqrt{R^2 - x^2}$ with $a \le x \le a + h$ about the x-axis to generate a zone of a sphere. $y = \sqrt{R^2 - x^2} \quad \Rightarrow$

$y' = \frac{1}{2}(R^2 - x^2)^{-1/2}(-2x) \quad \Rightarrow \quad ds = \sqrt{1 + \left(\dfrac{-x}{\sqrt{R^2 - x^2}}\right)^2}\,dx$. The surface area is

$$S = \int_a^{a+h} 2\pi y\,ds = 2\pi \int_a^{a+h} \sqrt{R^2 - x^2}\sqrt{1 + \frac{x^2}{R^2 - x^2}}\,dx$$

$$= 2\pi \int_a^{a+h} \sqrt{R^2 - x^2 + x^2}\,dx = 2\pi R\left[\,x\,\right]_a^{a+h}$$

$$= 2\pi R(a + h - a) = 2\pi Rh$$

43. Rotate $y = R$ with $0 \le x \le h$ about the x-axis to generate a zone of a cylinder. $y = R \quad \Rightarrow \quad y' = 0 \quad \Rightarrow$

$ds = \sqrt{1 + 0^2}\,dx = dx$. The surface area is $S = \int_0^h 2\pi y\,ds = 2\pi \int_0^h R\,dx = 2\pi R\left[\,x\,\right]_0^h = 2\pi Rh$.

44. Since $g(x) = f(x) + c$, we have $g'(x) = f'(x)$. Thus, the surface area generated by rotating the curve $g(x) = f(x) + c$ about the x-axis

$$S_g = \int_a^b 2\pi g(x)\sqrt{1 + [g'(x)]^2}\,dx = \int_a^b 2\pi[f(x) + c]\sqrt{1 + [f'(x)]^2}\,dx$$

$$= \int_a^b 2\pi f(x)\sqrt{1 + [f'(x)]^2}\,dx + 2\pi c\int_a^b \sqrt{1 + [f'(x)]^2}\,dx = S_f + 2\pi cL$$

45. $y = e^{x/2} + e^{-x/2} \quad \Rightarrow \quad y' = \frac{1}{2}e^{x/2} - \frac{1}{2}e^{-x/2} \quad \Rightarrow$

$1 + (y')^2 = 1 + \left(\frac{1}{2}e^{x/2} - \frac{1}{2}e^{-x/2}\right)^2 = 1 + \frac{1}{4}e^x - \frac{1}{2} + \frac{1}{4}e^{-x} = \frac{1}{4}e^x + \frac{1}{2} + \frac{1}{4}e^{-x} = \left(\frac{1}{2}e^{x/2} + \frac{1}{2}e^{-x/2}\right)^2$.

If we rotate the curve about the x-axis on the interval $a \le x \le b$, the resulting surface area is

$S = \int_a^b 2\pi y\sqrt{1 + (y')^2}\,dx = 2\pi \int_a^b (e^{x/2} + e^{-x/2})\left(\frac{1}{2}e^{x/2} + \frac{1}{2}e^{-x/2}\right)dx = \pi \int_a^b (e^{x/2} + e^{-x/2})^2\,dx$, which is the same as the volume obtained by rotating the curve y about the x-axis on the interval $a \le x \le b$, namely, $V = \pi \int_a^b y^2\,dx$.

46. In the derivation of (4), we computed a typical contribution to the surface area to be $2\pi\,\dfrac{y_{i-1} + y_i}{2}\,|P_{i-1}P_i|$, the area of a frustum of a cone. When $f(x)$ is not necessarily positive, the approximations $y_i = f(x_i) \approx f(x_i^*)$ and $y_{i-1} = f(x_{i-1}) \approx f(x_i^*)$ must be replaced by $y_i = |f(x_i)| \approx |f(x_i^*)|$ and $y_{i-1} = |f(x_{i-1})| \approx |f(x_i^*)|$. Thus, $2\pi\,\dfrac{y_{i-1} + y_i}{2}\,|P_{i-1}P_i| \approx 2\pi\,|f(x_i^*)|\sqrt{1 + [f'(x_i^*)]^2}\,\Delta x$. Continuing with the rest of the derivation as before, we obtain $S = \int_a^b 2\pi\,|f(x)|\sqrt{1 + [f'(x)]^2}\,dx$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

DISCOVERY PROJECT Rotating on a Slant

1.

In the figure, the segment a lying above the interval $[x_i - \Delta x, x_i]$ along the tangent to C has length $\Delta x \sec\alpha = \Delta x\sqrt{1+\tan^2\alpha} = \sqrt{1+[f'(x_i)]^2}\,\Delta x$. The segment from $(x_i, f(x_i))$ drawn perpendicular to the line $y = mx + b$ has length

$$g(x_i) = [f(x_i) - mx_i - b]\cos\beta = \frac{f(x_i) - mx_i - b}{\sec\beta} = \frac{f(x_i) - mx_i - b}{\sqrt{1+\tan^2\beta}} = \frac{f(x_i) - mx_i - b}{\sqrt{1+m^2}}$$

Also, $\cos(\beta - \alpha) = \dfrac{\Delta u}{\Delta x\ \sec\alpha} \quad \Rightarrow$

$$\Delta u = \Delta x\ \sec\alpha\ \cos(\beta-\alpha) = \Delta x\,\frac{\cos\beta\ \cos\alpha + \sin\beta\ \sin\alpha}{\cos\alpha} = \Delta x(\cos\beta + \sin\beta\ \tan\alpha)$$

$$= \Delta x\left[\frac{1}{\sqrt{1+m^2}} + \frac{m}{\sqrt{1+m^2}}f'(x_i)\right] = \frac{1+mf'(x_i)}{\sqrt{1+m^2}}\,\Delta x$$

Thus,
$$\text{Area}(\mathcal{R}) = \lim_{n\to\infty}\sum_{i=1}^{n} g(x_i)\,\Delta u = \lim_{n\to\infty}\sum_{i=1}^{n}\frac{f(x_i) - mx_i - b}{\sqrt{1+m^2}}\cdot\frac{1+mf'(x_i)}{\sqrt{1+m^2}}\,\Delta x$$

$$= \frac{1}{1+m^2}\int_p^q [f(x) - mx - b][1 + mf'(x)]\,dx$$

2. From Problem 1 with $m = 1$, $f(x) = x + \sin x$, $mx + b = x - 2$, $p = 0$, and $q = 2\pi$,

$$\text{Area} = \frac{1}{1\ +\ 1^2}\int_0^{2\pi}[x + \sin x - (x-2)]\,[1 + 1(1+\cos x)]\,dx = \tfrac{1}{2}\int_0^{2\pi}(\sin x + 2)(2 + \cos x)\,dx$$

$$= \tfrac{1}{2}\int_0^{2\pi}(2\sin x + \sin x\ \cos x + 4 + 2\cos x)\,dx = \tfrac{1}{2}\left[-2\cos x + \tfrac{1}{2}\sin^2 x + 4x + 2\sin x\right]_0^{2\pi}$$

$$= \tfrac{1}{2}[(-2 + 0 + 8\pi + 0) - (-2 + 0 + 0 + 0)] = \tfrac{1}{2}(8\pi) = 4\pi$$

3. $V = \displaystyle\lim_{n\to\infty}\sum_{i=1}^{n}\pi[g(x_i)]^2\,\Delta u = \lim_{n\to\infty}\sum_{i=1}^{n}\pi\left[\frac{f(x_i) - mx_i - b}{\sqrt{1+m^2}}\right]^2\frac{1+mf'(x_i)}{\sqrt{1+m^2}}\,\Delta x$

$$= \frac{\pi}{(1+m^2)^{3/2}}\int_p^q [f(x) - mx - b]^2[1 + mf'(x)]\,dx$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $V = \dfrac{\pi}{(1+1^2)^{3/2}} \displaystyle\int_0^{2\pi} (x + \sin x - x + 2)^2 (1 + 1 + \cos x)\, dx$

$$= \frac{\pi}{2\sqrt{2}} \int_0^{2\pi} (\sin x + 2)^2 (\cos x + 2)\, dx = \frac{\pi}{2\sqrt{2}} \int_0^{2\pi} \left(\sin^2 x + 4\sin x + 4\right)(\cos x + 2)\ dx$$

$$= \frac{\pi}{2\sqrt{2}} \int_0^{2\pi} \left(\sin^2 x\ \cos x + 4\sin x\ \cos x + 4\cos x + 2\sin^2 x + 8\sin x + 8\right) dx$$

$$= \frac{\pi}{2\sqrt{2}} \left[\tfrac{1}{3}\sin^3 x + 2\sin^2 x + 4\sin x + x - \tfrac{1}{2}\sin 2x - 8\cos x + 8x\right]_0^{2\pi} \qquad [\text{since } 2\sin^2 x = 1 - \cos 2x]$$

$$= \frac{\pi}{2\sqrt{2}} [(2\pi - 8 + 16\pi) - (-8)] = \frac{9\sqrt{2}}{2}\pi^2$$

5. $S = \displaystyle\int_p^q 2\pi g(x)\sqrt{1 + [f'(x)]^2}\, dx = \frac{2\pi}{\sqrt{1+m^2}} \int_p^q [f(x) - mx - b]\sqrt{1 + [f'(x)]^2}\, dx$

6. From Problem 5 with $f(x) = \sqrt{x}$, $p = 0$, $q = 4$, $m = \frac{1}{2}$, and $b = 0$,

$$S = \frac{2\pi}{\sqrt{1 + \left(\frac{1}{2}\right)^2}} \int_0^4 \left(\sqrt{x} - \tfrac{1}{2}x\right)\sqrt{1 + \left(\frac{1}{2\sqrt{x}}\right)^2}\, dx \overset{\text{CAS}}{=} \frac{\pi}{\sqrt{5}}\left[\frac{\ln\left(\sqrt{17}+4\right)}{32} + \frac{37\sqrt{17}}{24} - \frac{1}{3}\right] \approx 8.554$$

8.3 Applications to Physics and Engineering

1. The weight density of water is $\delta = 62.5\ \text{lb/ft}^3$.

(a) $P = \delta d \approx (62.5\ \text{lb/ft}^3)(3\ \text{ft}) = 187.5\ \text{lb/ft}^2$

(b) $F = PA \approx (187.5\ \text{lb/ft}^2)(5\ \text{ft})(2\ \text{ft}) = 1875\ \text{lb}$. ($A$ is the area of the bottom of the tank.)

(c) By reasoning as in Example 1, the area of the ith strip is $2\,(\Delta x)$ and the pressure is $\delta d = \delta x_i$. Thus,

$F = \int_0^3 \delta x \cdot 2\, dx \approx (62.5)(2)\int_0^3 x\, dx = 125\left[\frac{1}{2}x^2\right]_0^3 = 125\left(\frac{9}{2}\right) = 562.5\ \text{lb}$.

2. (a) $P = \rho g d = (820\ \text{kg/m}^3)(9.8\ \text{m/s}^2)(1.5\ \text{m}) = 12{,}054\ \text{Pa} \approx 12\ \text{kPa}$

(b) $F = PA = (12{,}054\ \text{Pa})(8\ \text{m})(4\ \text{m}) \approx 3.86 \times 10^5\ \text{N}$ (A is the area at the bottom of the tank.)

(c) The area of the ith strip is $4(\Delta x)$ and the pressure is $\rho g d = \rho g\, x_i$. Thus,

$F = \int_0^{1.5} \rho g x \cdot 4\, dx = (820)(9.8)\cdot 4\int_0^{1.5} x\, dx = 32{,}144\left[\frac{1}{2}x^2\right]_0^{3/2} = 16{,}072\left(\frac{9}{4}\right) \approx 3.62 \times 10^4\ \text{N}$.

In Exercises 3–9, n is the number of subintervals of length Δx and x_i^* is a sample point in the ith subinterval $[x_{i-1}, x_i]$.

3. Set up a vertical x-axis as shown, with $x = 0$ at the water's surface and x increasing in the downward direction. Then the area of the ith rectangular strip is $2\,\Delta x$ and the pressure on the strip is δx_i^* (where $\delta \approx 62.5\ \text{lb/ft}^3$). Thus, the hydrostatic force on the strip is $\delta x_i^* \cdot 2\,\Delta x$ and the total hydrostatic force $\approx \sum_{i=1}^{n} \delta x_i^* \cdot 2\,\Delta x$. The total force is

$$F = \lim_{n\to\infty} \sum_{i=1}^{n} \delta x_i^* \cdot 2\,\Delta x = \int_3^{11} \delta x \cdot 2\, dx = 2\delta\int_3^{11} x\, dx = 2\delta\left[\frac{1}{2}x^2\right]_3^{11} = \delta(121 - 9)$$

$$= 112\delta \approx 7000\ \text{lb}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. Set up a vertical axis as shown. Then the area of the ith rectangular strip is $2(x_i^* - 2)\,\Delta x$. $\left[\text{By similar triangles, } \dfrac{w_i}{x_i^* - 2} = \dfrac{10}{5}, \text{ so } w_i = 2(x_i^* - 2).\right]$

The pressure on the strip is δx_i^*, so the hydrostatic force on the strip is $\delta x_i^* \cdot 2(x_i^* - 2)\,\Delta x$ and the total hydrostatic force on the plate $\approx \sum\limits_{i=1}^{n} \delta x_i^* \cdot 2(x_i^* - 2)\,\Delta x$. The total force is

$$F = \lim_{n\to\infty} \sum_{i=1}^{n} \delta x_i^* \cdot 2(x_i^* - 2)\,\Delta x = \int_2^7 \delta x \cdot 2(x-2)\,dx = 2\delta \int_2^7 (x^2 - 2x)\,dx$$

$$= 2\delta\left[\tfrac{1}{3}x^3 - x^2\right]_2^7 = 2\delta\left[\left(\tfrac{343}{3} - 49\right) - \left(\tfrac{8}{3} - 4\right)\right] = 2\delta\left(\tfrac{200}{3}\right) = \tfrac{400}{3}\delta \approx \tfrac{400}{3}(62.5) = 8333.\overline{3} \text{ lb.}$$

5. Set up a coordinate system as shown. Then the area of the ith rectangular strip is $2\sqrt{8^2 - (y_i^*)^2}\,\Delta y$. The pressure on the strip is $\delta d_i = \rho g(12 - y_i^*)$, so the hydrostatic force on the strip is $\rho g(12 - y_i^*)\,2\sqrt{64 - (y_i^*)^2}\,\Delta y$ and the total hydrostatic force on the plate $\approx \sum\limits_{i=1}^{n} \rho g(12 - y_i^*)\,2\sqrt{64 - (y_i^*)^2}\,\Delta y$.

The total force $F = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \rho g(12 - y_i^*)\,2\sqrt{64 - (y_i^*)^2}\,\Delta y = \int_{-8}^{8} \rho g(12 - y)\,2\sqrt{64 - y^2}\,dy$

$$= 2\rho g \cdot 12 \int_{-8}^{8} \sqrt{64 - y^2}\,dy - 2\rho g \int_{-8}^{8} y\sqrt{64 - y^2}\,dy.$$

The second integral is 0 because the integrand is an odd function. The first integral is the area of a semicircular disk with radius 8. Thus, $F = 24\rho g\left(\frac{1}{2}\pi(8)^2\right) = 768\pi\rho g \approx 768\pi(1000)(9.8) \approx 2.36 \times 10^7$ N.

6. Set up a coordinate system as shown. Then the area of the ith rectangular strip is $2\sqrt{6^2 - (y_i^*)^2}\,\Delta y$. The pressure on the strip is $\delta d_i = \rho g(4 - y_i^*)$, so the hydrostatic force on the strip is $\rho g(4 - y_i^*)\,2\sqrt{36 - (y_i^*)^2}\,\Delta y$ and the hydrostatic force on the plate $\approx \sum\limits_{i=1}^{n} \rho g(4 - y_i^*)\,2\sqrt{36 - (y_i^*)^2}\,\Delta y$. The total force $F = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n} \rho g(4 - y_i^*)\,2\sqrt{36 - (y_i^*)^2}\,\Delta y = \int_0^4 \rho g(4 - y)\,2\sqrt{36 - y^2}\,dy = 8\rho g I_1 - 2\rho g I_2$.

$$I_1 = \int_0^4 \sqrt{36 - y^2}\,dy = \int_0^\alpha \sqrt{36 - 36\sin^2\theta}\,(6\cos\theta\,d\theta) \qquad \begin{bmatrix} y = 6\sin\theta, \\ dy = 6\cos\theta\,d\theta \\ \alpha = \sin^{-1}(2/3) \end{bmatrix}$$

$$= \int_0^\alpha 36\cos^2\theta\,d\theta = \int_0^\alpha 36 \cdot \tfrac{1}{2}(1 + \cos 2\theta)\,d\theta = 18\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_0^\alpha$$

$$= 18\left(\alpha + \tfrac{1}{2}\sin 2\alpha\right) = 18(\alpha + \sin\alpha\,\cos\alpha).$$

$$I_2 = \int_0^4 y\sqrt{36 - y^2}\,dy = \int_{36}^{20} \sqrt{u}\,(-\tfrac{1}{2}du) \qquad \begin{bmatrix} u = 36 - y^2, \\ du = -2y\,dy \end{bmatrix}$$

$$= -\tfrac{1}{2}\left[\tfrac{2}{3}u^{3/2}\right]_{36}^{20} = -\tfrac{1}{3}(20^{3/2} - 216) = 72 - \tfrac{40}{3}\sqrt{5}.$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Thus,

$$F = 8\rho g \cdot 18(\alpha + \sin\alpha\,\cos\alpha) - 2\rho g\big(72 - \tfrac{40}{3}\sqrt{5}\big) = 144\rho g\Big(\sin^{-1}\tfrac{2}{3} + \tfrac{2}{3}\tfrac{\sqrt{5}}{3}\Big) - 2\rho g\big(72 - \tfrac{40}{3}\sqrt{5}\big)$$
$$= \rho g\big(144\sin^{-1}\tfrac{2}{3} + \tfrac{176}{3}\sqrt{5} - 144\big) \approx 9.04 \times 10^5 \text{ N} \quad [\rho = 1000,\ g \approx 9.8].$$

7. Set up a vertical x-axis as shown. By similar triangles, $w_i/4 = x_i^*/6$, so $w_i = \frac{2}{3}x_i^*$, and the area of the ith rectangular strip is $\frac{2}{3}x_i^*\,\Delta x$. The pressure on the ith strip is $\rho g x_i^*$, so the hydrostatic force on the strip is $\rho g x_i^* \cdot \frac{2}{3}x_i^*\,\Delta x$, and the hydrostatic force on the plate is $\approx \sum_{i=1}^{n} \rho g x_i^* \cdot \frac{2}{3}x_i^*\,\Delta x$. The total force is

$$F = \lim_{n\to\infty} \sum_{i=1}^{n} \rho g x_i^* \cdot \tfrac{2}{3}x_i^*\,\Delta x = \tfrac{2}{3}\rho g \int_0^6 x^2\,dx = \tfrac{2}{3}\rho g\big[\tfrac{1}{3}x^3\big]_0^6 = \tfrac{2}{9}\rho g(216 - 0)$$
$$= 48\rho g = 48(1000)(9.8) = 470{,}400 \text{ N}$$

8. Set up a vertical x-axis as shown. The height of the triangle is $\sqrt{5^2 - (6/2)^2} = 4$. By similar triangles, $\dfrac{w_i}{6} = \dfrac{4 - x_i^*}{4}$, so $w_i = 6 - \frac{3}{2}x_i^*$, and the area of the ith rectangular strip is $\big(6 - \frac{3}{2}x_i^*\big)\,\Delta x$. The pressure on the ith strip is $\delta(3 + x_i^*)$, so the hydrostatic force on the strip is $\delta(3 + x_i^*)\big(6 - \frac{3}{2}x_i^*\big)\,\Delta x$, and the hydrostatic force on the plate is $\approx \sum_{i=1}^{n} \delta(3 + x_i^*)\big(6 - \frac{3}{2}x_i^*\big)\,\Delta x$. The total force is

$$F = \lim_{n\to\infty} \sum_{i=1}^{n} \delta(3 + x_i^*)\big(6 - \tfrac{3}{2}x_i^*\big)\ \Delta x = \delta \int_0^4 (3 + x)\,\big(6 - \tfrac{3}{2}x\big)\,dx = \delta \int_0^4 \big(18 + \tfrac{3}{2}x - \tfrac{3}{2}x^2\big)\,dx$$
$$= \delta\big[18x + \tfrac{3}{4}x^2 - \tfrac{1}{2}x^3\big]_0^4 = \delta(52 - 0) = 52\delta \approx 3250 \text{ lb} \quad [\delta \approx 62.5]$$

9. Set up a vertical x-axis as shown. Then the area of the ith rectangular strip is $w_i\,\Delta x = \big(4 + 2 \cdot \frac{2}{3}x_i^*\big)\,\Delta x$. The pressure on the strip is $\delta(x_i^* - 1)$, so the hydrostatic force on the strip is $\delta(x_i^* - 1)\big(4 + \frac{4}{3}x_i^*\big)\,\Delta x$ and the hydrostatic force on the plate $\approx \sum_{i=1}^{n} \delta(x_i^* - 1)\big(4 + \frac{4}{3}x_i^*\big)\,\Delta x$. The total force is

$$F = \lim_{n\to\infty} \sum_{i=1}^{n} \delta(x_i^* - 1)\big(4 + \tfrac{4}{3}x_i^*\big)\,\Delta x = \int_1^3 \delta(x - 1)\big(4 + \tfrac{4}{3}x\big)\,dx$$
$$= \delta \int_1^3 \big(\tfrac{4}{3}x^2 + \tfrac{8}{3}x - 4\big)\,dx = \delta\big[\tfrac{4}{9}x^3 + \tfrac{4}{3}x^2 - 4x\big]_1^3$$
$$= \delta\big[(12 + 12 - 12) - \big(\tfrac{4}{9} + \tfrac{4}{3} - 4\big)\big] = \delta\big(\tfrac{128}{9}\big)$$
$$\approx 889 \text{ lb} \quad [\delta \approx 62.5]$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. Set up coordinate axes as shown in the figure. For the *top half,* the length of the ith strip is $2\left(a/\sqrt{2}-y_i^*\right)$ and its area is $2\left(a/\sqrt{2}-y_i^*\right)\Delta y$. The pressure on this strip is approximately $\delta d_i=\delta\left(a/\sqrt{2}-y_i^*\right)$ and so the force on the strip is approximately $2\delta\left(a/\sqrt{2}-y_i^*\right)^2\Delta y$. The total force is

$$F_1=\lim_{n\to\infty}\sum_{i=1}^{n}2\delta\left(\frac{a}{\sqrt{2}}-y_i^*\right)^2\Delta y=2\delta\int_0^{a/\sqrt{2}}\left(\frac{a}{\sqrt{2}}-y\right)^2dy$$

$$=2\delta\left[-\frac{1}{3}\left(\frac{a}{\sqrt{2}}-y\right)^3\right]_0^{a/\sqrt{2}}=-\frac{2}{3}\delta\left[0-\left(\frac{a}{\sqrt{2}}\right)^3\right]=\frac{2\delta}{3}\frac{a^3}{2\sqrt{2}}=\frac{\sqrt{2}\,a^3\delta}{6}$$

For the *bottom half,* the length is $2\left(a/\sqrt{2}+y_i^*\right)$ and the total force is

$$F_2=\lim_{n\to\infty}\sum_{i=1}^{n}2\delta\left(\frac{a}{\sqrt{2}}+y_i^*\right)\left(\frac{a}{\sqrt{2}}-y_i^*\right)\Delta y=2\delta\int_{-a/\sqrt{2}}^{0}\left(\frac{a^2}{2}-y^2\right)dy=2\delta\left[\tfrac{1}{2}a^2y-\tfrac{1}{3}y^3\right]_{-a/\sqrt{2}}^{0}$$

$$=2\delta\left[0-\left(-\frac{\sqrt{2}\,a^3}{4}+\frac{\sqrt{2}\,a^3}{12}\right)\right]=2\delta\left(\frac{\sqrt{2}\,a^3}{6}\right)=\frac{2\sqrt{2}\,a^3\delta}{6}\qquad[F_2=2F_1]$$

Thus, the total force $F=F_1+F_2=\dfrac{3\sqrt{2}\,a^3\delta}{6}=\dfrac{\sqrt{2}\,a^3\delta}{2}$.

11. Set up a vertical x-axis as shown. Then the area of the ith rectangular strip is $\dfrac{a}{h}(2h-x_i^*)\,\Delta x$. $\left[\text{By similar triangles, }\dfrac{w_i}{2h-x_i^*}=\dfrac{2a}{2h}\text{, so }w_i=\dfrac{a}{h}(2h-x_i^*).\right]$ The pressure on the strip is δx_i^*, so the hydrostatic force on the plate $\approx\sum_{i=1}^{n}\delta x_i^*\,\dfrac{a}{h}(2h-x_i^*)\,\Delta x$. The total force is

$$F=\lim_{n\to\infty}\sum_{i=1}^{n}\delta x_i^*\,\frac{a}{h}(2h-x_i^*)\,\Delta x=\delta\,\frac{a}{h}\int_0^h x(2h-x)\,dx=\frac{a\delta}{h}\int_0^h\left(2hx-x^2\right)dx$$

$$=\frac{a\delta}{h}\left[hx^2-\tfrac{1}{3}x^3\right]_0^h=\frac{a\delta}{h}\left(h^3-\tfrac{1}{3}h^3\right)=\frac{a\delta}{h}\left(\frac{2h^3}{3}\right)=\tfrac{2}{3}\delta ah^2$$

12. $F=\int_0^2\rho g(10-x)2\sqrt{4-x^2}\,dx$

$$=20\rho g\int_0^2\sqrt{4-x^2}\,dx-\rho g\int_0^2\sqrt{4-x^2}\,2x\,dx$$

$$=20\rho g\tfrac{1}{4}\pi(2^2)-\rho g\int_0^4 u^{1/2}\,du\qquad[u=4-x^2,\,du=-2x\,dx]$$

$$=20\pi\rho g-\tfrac{2}{3}\rho g\left[u^{3/2}\right]_0^4=20\pi\rho g-\tfrac{16}{3}\rho g=\rho g\left(20\pi-\tfrac{16}{3}\right)$$

$$=(1000)(9.8)\left(20\pi-\tfrac{16}{3}\right)\approx5.63\times10^5\text{ N}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. The solution is similar to the solution for Example 2. The pressure on a strip is approximately $\delta d_i = 47(4 - y_i^*)$ and the area of the ith rectangular strip is $2\sqrt{16 - (y_i^*)^2}\,\Delta y$, so the total force is

$$F = \lim_{n\to\infty} \sum_{i=1}^{n} 47(4 - y_i^*)\,2\sqrt{16 - (y_i^*)^2}\,\Delta y = 94\int_{-4}^{4}(4 - y)\sqrt{16 - y^2}\,dy$$

$$= 94 \cdot 4\int_{-4}^{4}\sqrt{16 - y^2}\,dy - 94\int_{-4}^{4} y\sqrt{16 - y^2}\,dy$$

$$= 376 \cdot \tfrac{1}{2}\pi(4)^2 - 0 \qquad \left[\begin{array}{c}\text{The first integral is the area of a semicircular disk with radius 4}\\ \text{and the second integral is 0 because the integrand is an odd function.}\end{array}\right]$$

$$= 3008\pi \approx 9450 \text{ lb}$$

14. (a) Set up a coordinate system as shown. The slanted side of the trough has equation $y = 3x - 3$ or $x = \frac{1}{3}y + 1$, so the area of the ith rectangular strip is $\left(\frac{1}{3}y_i^* + 1\right)\Delta y$. The pressure on the strip is $\rho g d_i = \rho g(3 - y_i^*)$, so the hydrostatic force on the strip is $\rho g(3 - y_i^*)\left(\frac{1}{3}y_i^* + 1\right)\Delta y$ and the total force on the end of the trough is

$$F = \lim_{n\to\infty} \sum_{i=1}^{n} \rho g(3 - y_i^*)\left(\tfrac{1}{3}y_i^* + 1\right)\Delta y = \rho g\int_0^3 (3 - y)\left(\tfrac{1}{3}y + 1\right)dy$$

$$= \rho g\int_0^3 \left(3 - \tfrac{1}{3}y^2\right)dy = \rho g\left[3y - \tfrac{1}{9}y^3\right]_0^3 = \rho g(9 - 3)$$

$$\approx (925)(9.8)(6) = 54{,}390 \text{ N}$$

(b) When filled to a depth of 1.2 m, the pressure on the ith rectangular strip is $\rho g(1.2 - y_i^*)$, so the hydrostatic force on the strip is $\rho g(1.2 - y_i^*)\left(\frac{1}{3}y_i^* + 1\right)\Delta y$ and the total force on the end of the trough is

$$F = \lim_{n\to\infty} \sum_{i=1}^{n} \rho g(1.2 - y_i^*)\left(\tfrac{1}{3}y_i^* + 1\right)\Delta y = \rho g\int_0^{1.2}(1.2 - y)\left(\tfrac{1}{3}y + 1\right)dy = \rho g\int_0^{1.2}\left(1.2 - 0.6y - \tfrac{1}{3}y^2\right)dy$$

$$= \rho g\left[1.2y - 0.3y^2 - \tfrac{1}{9}y^3\right]_0^{1.2} = (925)(9.8)(1.44 - 0.432 - 0.192) = 7397.04 \text{ N}$$

Note that this is about 14% of the force for the completely filled trough.

15. (a) The top of the cube has depth $d = 1 \text{ m} - 20 \text{ cm} = 80 \text{ cm} = 0.8 \text{ m}$.

$$F = \rho g d A \approx (1000)(9.8)(0.8)(0.2)^2 = 313.6 \approx 314 \text{ N}$$

(b) The area of a strip is $0.2\,\Delta x$ and the pressure on it is $\rho g x_i^*$.

$$F = \int_{0.8}^{1} \rho g x(0.2)\,dx = 0.2\rho g\left[\tfrac{1}{2}x^2\right]_{0.8}^{1} = (0.2\rho g)(0.18) = 0.036\rho g = 0.036(1000)(9.8) = 352.8 \approx 353 \text{ N}$$

16. The height of the dam is $h = \sqrt{70^2 - 25^2}\,\cos 30^\circ = 15\sqrt{19}\left(\frac{\sqrt{3}}{2}\right)$.

The width of the trapezoid is $w = 50 + 2a$.

By similar triangles, $\dfrac{25}{h} = \dfrac{a}{h - x} \quad \Rightarrow \quad a = \dfrac{25}{h}(h - x)$. Thus,

$$w = 50 + 2 \cdot \frac{25}{h}(h - x) = 50 + \frac{50}{h} \cdot h - \frac{50}{h} \cdot x = 50 + 50 - \frac{50x}{h} = 100 - \frac{50x}{h}.$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

From the small triangle in the second figure, $\cos 30^\circ = \dfrac{\Delta x}{z} \quad \Rightarrow$

$z = \Delta x \sec 30^\circ = 2\,\Delta x/\sqrt{3}$.

$$F = \int_0^h \delta x\left(100 - \frac{50x}{h}\right)\frac{2}{\sqrt{3}}\,dx = \frac{200\delta}{\sqrt{3}}\int_0^h x\,dx - \frac{100\delta}{h\sqrt{3}}\int_0^h x^2\,dx$$

$$= \frac{200\delta}{\sqrt{3}}\,\frac{h^2}{2} - \frac{100\delta}{h\sqrt{3}}\,\frac{h^3}{3} = \frac{200\delta h^2}{3\sqrt{3}} = \frac{200(62.5)}{3\sqrt{3}}\cdot\frac{12{,}825}{4} \approx 7.71\times 10^6 \text{ lb}$$

17. (a) *Shallow end:* The area of a strip is $20\,\Delta x$ and the pressure on it is δx_i.

$$F = \int_0^3 \delta x 20\,dx = 20\delta\left[\tfrac{1}{2}x^2\right]_0^3 = 20\delta\cdot\tfrac{9}{2} = 90\delta$$

$$= 90(62.5) = 5625 \text{ lb} \approx 5.63\times 10^3 \text{ lb}$$

Deep end: $F = \int_0^9 \delta x 20\,dx = 20\delta\left[\tfrac{1}{2}x^2\right]_0^9 = 20\delta\cdot\tfrac{81}{2} = 810\delta = 810(62.5) = 50{,}625 \text{ lb} \approx 5.06\times 10^4$ lb.

Sides: For the first 3 ft, the length of a side is constant at 40 ft. For $3 < x \le 9$, we can use similar triangles to find the length a: $\dfrac{a}{40} = \dfrac{9-x}{6} \quad\Rightarrow\quad a = 40\cdot\dfrac{9-x}{6}$.

$$F = \int_0^3 \delta x 40\,dx + \int_3^9 \delta x(40)\,\frac{9-x}{6}\,dx = 40\delta\left[\tfrac{1}{2}x^2\right]_0^3 + \tfrac{20}{3}\delta\int_3^9 (9x - x^2)\,dx = 180\delta + \tfrac{20}{3}\delta\left[\tfrac{9}{2}x^2 - \tfrac{1}{3}x^3\right]_3^9$$

$$= 180\delta + \tfrac{20}{3}\delta\left[\left(\tfrac{729}{2} - 243\right) - \left(\tfrac{81}{2} - 9\right)\right] = 180\delta + 600\delta = 780\delta = 780(62.5) = 48{,}750 \text{ lb} \approx 4.88\times 10^4 \text{ lb}$$

(b) For any right triangle with hypotenuse on the bottom,

$\sin\theta = \dfrac{\Delta x}{\text{hypotenuse}} \quad\Rightarrow$

$\text{hypotenuse} = \Delta x \csc\theta = \Delta x\,\dfrac{\sqrt{40^2 + 6^2}}{6} = \dfrac{\sqrt{409}}{3}\,\Delta x$.

$$F = \int_3^9 \delta x 20\,\tfrac{\sqrt{409}}{3}\,dx = \tfrac{1}{3}\left(20\sqrt{409}\right)\delta\left[\tfrac{1}{2}x^2\right]_3^9$$

$$= \tfrac{1}{3}\cdot 10\sqrt{409}\,\delta(81 - 9) \approx 303{,}356 \text{ lb} \approx 3.03\times 10^5 \text{ lb}$$

18. Partition the interval $[a, b]$ by points x_i as usual and choose $x_i^* \in [x_{i-1}, x_i]$ for each i. The ith horizontal strip of the immersed plate is approximated by a rectangle of height Δx_i and width $w(x_i^*)$, so its area is $A_i \approx w(x_i^*)\,\Delta x_i$. For small Δx_i, the pressure P_i on the ith strip is almost constant and $P_i \approx \rho g x_i^*$ by Equation 1. The hydrostatic force F_i acting on the ith strip is $F_i = P_i A_i \approx \rho g x_i^* w(x_i^*)\,\Delta x_i$. Adding these forces and taking the limit as $n \to \infty$, we obtain the hydrostatic force on the immersed plate:

$$F = \lim_{n\to\infty}\sum_{i=1}^{n} F_i = \lim_{n\to\infty}\sum_{i=1}^{n} \rho g x_i^* w(x_i^*)\,\Delta x_i = \int_a^b \rho g x w(x)\,dx$$

19. From Exercise 18, we have $F = \int_a^b \rho g x w(x)\,dx = \int_{7.0}^{9.4} 64 x w(x)\,dx$. From the table, we see that $\Delta x = 0.4$, so using Simpson's Rule to estimate F, we get

$$F \approx 64\,\tfrac{0.4}{3}[7.0w(7.0) + 4(7.4)w(7.4) + 2(7.8)w(7.8) + 4(8.2)w(8.2) + 2(8.6)w(8.6) + 4(9.0)w(9.0) + 9.4w(9.4)]$$

$$= \tfrac{25.6}{3}[7(1.2) + 29.6(1.8) + 15.6(2.9) + 32.8(3.8) + 17.2(3.6) + 36(4.2) + 9.4(4.4)]$$

$$= \tfrac{25.6}{3}(486.04) \approx 4148 \text{ lb}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. (a) From Equation 8, $\overline{x} = \frac{1}{A}\int_a^b xw(x)\,dx \quad\Rightarrow\quad A\overline{x} = \int_a^b xw(x)\,dx \quad\Rightarrow\quad \rho g A\overline{x} = \rho g\int_a^b xw(x)\,dx \quad\Rightarrow$

$(\rho g\overline{x})A = \int_a^b \rho g\, xw(x)\,dx = F$ by Exercise 18.

(b) For the figure in Exercise 10, let the coordinates of the centroid $(\overline{x},\overline{y}) = \left(a/\sqrt{2}, 0\right)$.

$$F = (\rho g\overline{x})A = \rho g\,\frac{a}{\sqrt{2}}\,a^2 = \delta\,\frac{\sqrt{2}\,a}{2}\,a^2 = \frac{\sqrt{2}\,a^3\delta}{2}.$$

21. The moment M of the system about the origin is $M = \sum_{i=1}^{2} m_i x_i = m_1x_1 + m_2x_2 = 6\cdot 10 + 9\cdot 30 = 330$.

The mass m of the system is $m = \sum_{i=1}^{2} m_i = m_1 + m_2 = 6 + 9 = 15$.

The center of mass of the system is $\overline{x} = M/m = \frac{330}{15} = 22$.

22. The moment M is $m_1x_1 + m_2x_2 + m_3x_3 = 12(-3) + 15(2) + 20(8) = 154$. The mass m is $m_1 + m_2 + m_3 = 12 + 15 + 20 = 47$. The center of mass is $\overline{x} = M/m = \frac{154}{47}$.

23. The mass is $m = \sum_{i=1}^{3} m_i = 5 + 8 + 7 = 20$. The moment about the x-axis is $M_x = \sum_{i=1}^{3} m_i y_i = 5(1) + 8(4) + 7(-2) = 23$.

The moment about the y-axis is $M_y = \sum_{i=1}^{3} m_i x_i = 5(3) + 8(0) + 7(-5) = -20$. The center of mass is

$$(\overline{x},\overline{y}) = \left(\frac{M_y}{m}, \frac{M_x}{m}\right) = \left(-\frac{20}{20}, \frac{23}{20}\right) = (-1, 1.15).$$

24. The mass is $m = \sum_{i=1}^{4} m_i = 4 + 3 + 6 + 3 = 16$.

The moment about the x-axis is $M_x = \sum_{i=1}^{4} m_i y_i = 4(1) + 3(-1) + 6(2) + 3(-5) = -2$.

The moment about the y-axis is $M_y = \sum_{i=1}^{4} m_i x_i = 4(6) + 3(3) + 6(-2) + 3(-2) = 15$.

The center of mass is $(\overline{x},\overline{y}) = \left(\frac{M_y}{m}, \frac{M_x}{m}\right) = \left(\frac{15}{16}, -\frac{2}{16}\right) = \left(\frac{15}{16}, -\frac{1}{8}\right)$.

25. The region in the figure is "left-heavy" and "bottom-heavy," so we know that $\overline{x} < 1$ and $\overline{y} < 2$, and we might guess that $\overline{x} = 0.7$ and $\overline{y} = 1.3$. The line $2x + y = 4$ can be expressed as $y = 4 - 2x$, so $A = \int_0^2 (4 - 2x)\,dx = \left[4x - x^2\right]_0^2 = 8 - 4 = 4$.

$$\overline{x} = \frac{1}{A}\int_0^2 x(4 - 2x)\,dx = \tfrac{1}{4}\int_0^2 (4x - 2x^2)\,dx = \tfrac{1}{4}\left[2x^2 - \tfrac{2}{3}x^3\right]_0^2 = \tfrac{1}{4}\left(8 - \tfrac{16}{3}\right) = \tfrac{2}{3}.$$

$$\overline{y} = \frac{1}{A}\int_0^2 \tfrac{1}{2}(4 - 2x)^2 dx = \tfrac{1}{4}\int_0^2 \tfrac{1}{2}\cdot 4(2 - x)^2 dx = \tfrac{1}{2}\left[-\tfrac{1}{3}(2 - x)^3\right]_0^2 = -\tfrac{1}{6}(0 - 8) = \tfrac{4}{3}.$$

Thus, the centroid is $(\overline{x},\overline{y}) = \left(\frac{2}{3}, \frac{4}{3}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. The region in the figure is "right-heavy" and "bottom-heavy," so we know that $\overline{x} > 0$ and $\overline{y} < 1$, and we might guess that $\overline{x} = 0.3$ and $\overline{y} = 0.8$.

$A = \int_{-1}^{1} e^x\,dx = \left[e^x\right]_{-1}^{1} = e - e^{-1}$.

$$\overline{x} = \frac{1}{A}\int_{-1}^{1} xe^x\,dx = \frac{1}{e-e^{-1}}\left[xe^x - e^x\right]_{-1}^{1} \quad \text{[by parts]}$$

$$= \frac{1}{e-e^{-1}}\left[(e-e) - (-e^{-1} - e^{-1})\right] = \frac{2e^{-1}}{e-e^{-1}} \cdot \frac{e}{e} = \frac{2}{e^2-1}$$

$$\overline{y} = \frac{1}{A}\int_{-1}^{1} \frac{1}{2}(e^x)^2\,dx = \frac{1}{2(e-e^{-1})}\int_{-1}^{1} e^{2x}\,dx = \frac{1}{2(e-e^{-1})} \cdot \frac{1}{2}\left[e^{2x}\right]_{-1}^{1} = \frac{e^2-e^{-2}}{4(e-e^{-1})} \cdot \frac{e^2}{e^2} = \frac{e^4-1}{4e(e^2-1)}$$

$$= \frac{(e^2+1)(e^2-1)}{4e(e^2-1)} = \frac{e^2+1}{4e}$$

Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\dfrac{2}{e^2-1}, \dfrac{e^2+1}{4e}\right) \approx (0.31, 0.77)$.

27. The region in the figure is "right-heavy" and "bottom-heavy," so we know that $\overline{x} > 1$ and $\overline{y} < 1$, and we might guess that $\overline{x} = 1.5$ and $\overline{y} = 0.5$.

$A = \int_0^2 \frac{1}{2}x^2\,dx = \left[\frac{1}{6}x^3\right]_0^2 = \frac{8}{6} - 0 = \frac{4}{3}$.

$$\overline{x} = \frac{1}{A}\int_0^2 x \cdot \tfrac{1}{2}x^2 dx = \tfrac{3}{4}\int_0^2 \tfrac{1}{2}x^3 dx = \tfrac{3}{8}\int_0^2 x^3 dx = \tfrac{3}{8}\left[\tfrac{1}{4}x^4\right]_0^2 = \tfrac{3}{32}(16-0) = \tfrac{3}{2}.$$

$$\overline{y} = \frac{1}{A}\int_0^2 \tfrac{1}{2}\left(\tfrac{1}{2}x^2\right)^2 dx = \tfrac{3}{4}\int_0^2 \tfrac{1}{8}x^4\,dx = \tfrac{3}{32}\left[\tfrac{1}{5}x^5\right]_0^2 = \tfrac{3}{160}(32-0) = \tfrac{3}{5}.$$

Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\frac{3}{2}, \frac{3}{5}\right)$.

28. The region in the figure is "left-heavy" and "bottom-heavy," so we know $\overline{x} < 1.5$ and $\overline{y} < 0.5$, and we might guess that $\overline{x} = 1.4$ and $\overline{y} = 0.4$.

$A = \int_1^2 \frac{1}{x}\,dx = [\ln x]_1^2 = \ln 2$. $\overline{x} = \frac{1}{A}\int_1^2 x \cdot \frac{1}{x}\,dx = \frac{1}{A}[x]_1^2 = \frac{1}{A} = \frac{1}{\ln 2}$.

$\overline{y} = \frac{1}{A}\int_1^2 \frac{1}{2}\left(\frac{1}{x}\right)^2\,dx = \frac{1}{2A}\int_1^2 x^{-2}\,dx = \frac{1}{2A}\left[-\frac{1}{x}\right]_1^2 = \frac{1}{2\ln 2}\left(-\frac{1}{2}+1\right) = \frac{1}{4\ln 2}$.

Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\frac{1}{\ln 2}, \frac{1}{4\ln 2}\right) \approx (1.44, 0.36)$.

29. $A = \int_0^1 (x^{1/2} - x^2)\,dx = \left[\frac{2}{3}x^{3/2} - \frac{1}{3}x^3\right]_0^1 = \left(\frac{2}{3} - \frac{1}{3}\right) - 0 = \frac{1}{3}$.

$$\overline{x} = \tfrac{1}{A}\int_0^1 x(x^{1/2} - x^2)\,dx = 3\int_0^1 (x^{3/2} - x^3)\,dx = 3\left[\tfrac{2}{5}x^{5/2} - \tfrac{1}{4}x^4\right]_0^1$$

$$= 3\left(\tfrac{2}{5} - \tfrac{1}{4}\right) = 3\left(\tfrac{3}{20}\right) = \tfrac{9}{20}.$$

$$\overline{y} = \tfrac{1}{A}\int_0^1 \tfrac{1}{2}\left[(x^{1/2})^2 - (x^2)^2\right] dx = 3\left(\tfrac{1}{2}\right)\int_0^1 (x - x^4)\,dx$$

$$= \tfrac{3}{2}\left[\tfrac{1}{2}x^2 - \tfrac{1}{5}x^5\right]_0^1 = \tfrac{3}{2}\left(\tfrac{1}{2} - \tfrac{1}{5}\right) = \tfrac{3}{2}\left(\tfrac{3}{10}\right) = \tfrac{9}{20}.$$

Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\frac{9}{20}, \frac{9}{20}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. The curves intersect when $2 - x^2 = x \quad \Leftrightarrow \quad 0 = x^2 + x - 2 \quad \Leftrightarrow$ $0 = (x+2)(x-1) \quad \Leftrightarrow \quad x = -2$ or $x = 1$.

$A = \int_{-2}^{1}(2 - x^2 - x)\,dx = \left[2x - \frac{1}{3}x^3 - \frac{1}{2}x^2\right]_{-2}^{1} = \frac{7}{6} - \left(-\frac{10}{3}\right) = \frac{9}{2}$.

$\overline{x} = \frac{1}{A}\int_{-2}^{1} x(2 - x^2 - x)\,dx = \frac{2}{9}\int_{-2}^{1}(2x - x^3 - x^2)\,dx$

$= \frac{2}{9}\left[x^2 - \frac{1}{4}x^4 - \frac{1}{3}x^3\right]_{-2}^{1} = \frac{2}{9}\left(\frac{5}{12} - \frac{8}{3}\right) = -\frac{1}{2}$.

$\overline{y} = \frac{1}{A}\int_{-2}^{1} \frac{1}{2}[(2 - x^2)^2 - x^2]\,dx = \frac{2}{9}\cdot\frac{1}{2}\int_{-2}^{1}(4 - 5x^2 + x^4)\,dx$

$= \frac{1}{9}\left[4x - \frac{5}{3}x^3 + \frac{1}{5}x^5\right]_{-2}^{1} = \frac{1}{9}\left[\frac{38}{15} - \left(-\frac{16}{15}\right)\right] = \frac{2}{5}$.

Thus, the centroid is $(\overline{x}, \overline{y}) = (-\frac{1}{2}, \frac{2}{5})$.

31. $A = \int_0^{\pi/3}(\sin 2x - \sin x)\,dx = \left[-\frac{1}{2}\cos 2x + \cos x\right]_0^{\pi/3} = \left(\frac{1}{4} + \frac{1}{2}\right) - \left(-\frac{1}{2} + 1\right) = \frac{1}{4}$.

$$\overline{x} = \frac{1}{A}\int_0^{\pi/3} x(\sin 2x - \sin x)\,dx = 4\left[x\left(-\frac{1}{2}\cos 2x + \cos x\right) - \left(-\frac{1}{4}\sin 2x + \sin x\right)\right]_0^{\pi/3} \qquad \left[\begin{array}{l}\text{by parts with } u = x \text{ and}\\ dv = (\sin 2x - \sin x)\,dx\end{array}\right]$$

$$= 4\left[\frac{\pi}{3}\left(\frac{1}{4} + \frac{1}{2}\right) - \left(-\frac{\sqrt{3}}{8} + \frac{\sqrt{3}}{2}\right)\right] = \pi - \frac{3\sqrt{3}}{2}.$$

$$\overline{y} = \frac{1}{A}\int_0^{\pi/3} \frac{1}{2}(\sin^2 2x - \sin^2 x)\,dx$$

$$= 4\int_0^{\pi/3} \frac{1}{2}(4\sin^2 x\cos^2 x - \sin^2 x)\,dx \qquad \left[\begin{array}{c}\text{double-angle}\\ \text{identity}\end{array}\right]$$

$$= 2\int_0^{\pi/3}[4\sin^2 x\,(1 - \sin^2 x) - \sin^2 x]\,dx$$

$$= 6\int_0^{\pi/3}\sin^2 x\,dx - 8\int_0^{\pi/3}\sin^4 x\,dx$$

Now, $\displaystyle\int_0^{\pi/3}\sin^2 x\,dx \overset{63}{=} \left[\frac{1}{2}x - \frac{1}{4}\sin 2x\right]_0^{\pi/3} = \frac{\pi}{6} - \frac{\sqrt{3}}{8}$ and

$$\int_0^{\pi/3}\sin^4 x\,dx = \frac{1}{4}\left[\frac{3}{2}x - \sin 2x + \frac{1}{8}\sin 4x\right]_0^{\pi/3} \qquad [\text{by Example 7.2.4}]$$

$$= \frac{1}{4}\left[\frac{\pi}{2} - \frac{\sqrt{3}}{2} - \frac{\sqrt{3}}{16}\right] = \frac{\pi}{8} - \frac{9\sqrt{3}}{64}$$

So $\overline{y} = 6\left(\dfrac{\pi}{6} - \dfrac{\sqrt{3}}{8}\right) - 8\left(\dfrac{\pi}{8} - \dfrac{9\sqrt{3}}{64}\right) = \dfrac{3\sqrt{3}}{8}$. Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\pi - \dfrac{3\sqrt{3}}{2}, \dfrac{3\sqrt{3}}{8}\right) \approx (0.54, 0.65)$.

32. $A = \int_0^1 x^3\,dx + \int_1^2(2 - x)\,dx = \left[\frac{1}{4}x^4\right]_0^1 + \left[2x - \frac{1}{2}x^2\right]_1^2$

$= \frac{1}{4} + (4 - 2) - \left(2 - \frac{1}{2}\right) = \frac{3}{4}$.

$\overline{x} = \frac{1}{A}\left[\int_0^1 x(x^3)\,dx + \int_1^2 x(2 - x)\,dx\right] = \frac{4}{3}\left[\int_0^1 x^4\,dx + \int_1^2(2x - x^2)\,dx\right]$

$= \frac{4}{3}\left\{\left[\frac{1}{5}x^5\right]_0^1 + \left[x^2 - \frac{1}{3}x^3\right]_1^2\right\} = \frac{4}{3}\left[\frac{1}{5} + \left(4 - \frac{8}{3}\right) - \left(1 - \frac{1}{3}\right)\right]$

$= \frac{4}{3}\left(\frac{13}{15}\right) = \frac{52}{45}$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\overline{y} = \frac{1}{A}\left[\int_0^1 \frac{1}{2}(x^3)^2\,dx + \int_1^2 \frac{1}{2}(2-x)^2\,dx\right] = \frac{2}{3}\left[\int_0^1 x^6 dx + \int_1^2 (x-2)^2\,dx\right] = \frac{2}{3}\left\{\left[\frac{1}{7}x^7\right]_0^1 + \left[\frac{1}{3}(x-2)^3\right]_1^2\right\}$

$= \frac{2}{3}\left(\frac{1}{7} - 0 + 0 + \frac{1}{3}\right) = \frac{2}{3}\left(\frac{10}{21}\right) = \frac{20}{63}.$

Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\frac{52}{45}, \frac{20}{63}\right)$.

33. The curves intersect when $2 - y = y^2 \quad \Leftrightarrow \quad 0 = y^2 + y - 2 \quad \Leftrightarrow$

$0 = (y+2)(y-1) \quad \Leftrightarrow \quad y = -2$ or $y = 1$.

$A = \int_{-2}^1 (2 - y - y^2)\,dy = \left[2y - \frac{1}{2}y^2 - \frac{1}{3}y^3\right]_{-2}^1 = \frac{7}{6} - \left(-\frac{10}{3}\right) = \frac{9}{2}.$

$\overline{x} = \frac{1}{A}\int_{-2}^1 \frac{1}{2}[(2-y)^2 - (y^2)^2]\,dy = \frac{2}{9}\cdot\frac{1}{2}\int_{-2}^1 (4 - 4y + y^2 - y^4)\,dy$

$= \frac{1}{9}\left[4y - 2y^2 + \frac{1}{3}y^3 - \frac{1}{5}y^5\right]_{-2}^1 = \frac{1}{9}\left[\frac{32}{15} - \left(-\frac{184}{15}\right)\right] = \frac{8}{5}.$

$\overline{y} = \frac{1}{A}\int_{-2}^1 y(2 - y - y^2)\,dy = \frac{2}{9}\int_{-2}^1 (2y - y^2 - y^3)\,dy$

$= \frac{2}{9}\left[y^2 - \frac{1}{3}y^3 - \frac{1}{4}y^4\right]_{-2}^1 = \frac{2}{9}\left(\frac{5}{12} - \frac{8}{3}\right) = -\frac{1}{2}.$

Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\frac{8}{5}, -\frac{1}{2}\right)$.

34. An equation of the line is $y = -\frac{3}{2}x + 3$. $A = \frac{1}{2}(2)(3) = 3$, so $m = \rho A = 4(3) = 12$.

$M_x = \rho\int_0^2 \frac{1}{2}\left(-\frac{3}{2}x + 3\right)^2\,dx = \frac{1}{2}\rho\int_0^2 \left(\frac{9}{4}x^2 - 9x + 9\right)dx = \frac{1}{2}(4)\left[\frac{3}{4}x^3 - \frac{9}{2}x^2 + 9x\right]_0^2 = 2(6 - 18 + 18) = 12.$

$M_y = \rho\int_0^2 x\left(-\frac{3}{2}x + 3\right)dx = \rho\int_0^2 \left(-\frac{3}{2}x^2 + 3x\right)dx = 4\left[-\frac{1}{2}x^3 + \frac{3}{2}x^2\right]_0^2 = 4(-4 + 6) = 8.$

$\overline{x} = \dfrac{M_y}{m} = \dfrac{8}{12} = \dfrac{2}{3}$ and $\overline{y} = \dfrac{M_x}{m} = \dfrac{12}{12} = 1$. Thus, the center of mass is $(\overline{x}, \overline{y}) = \left(\frac{2}{3}, 1\right)$. Since ρ is constant, the center of mass is also the centroid.

35. The quarter-circle has equation $y = \sqrt{4^2 - x^2}$ for $0 \le x \le 4$ and the line has equation $y = -2$.

$A = \frac{1}{4}\pi(4)^2 + 2(4) = 4\pi + 8 = 4(\pi + 2)$, so $m = \rho A = 6\cdot 4(\pi + 2) = 24(\pi + 2)$.

$M_x = \rho\int_0^4 \frac{1}{2}\left[\left(\sqrt{16 - x^2}\right)^2 - (-2)^2\right]dx = \frac{1}{2}\rho\int_0^4 (16 - x^2 - 4)\,dx = \frac{1}{2}(6)\left[12x - \frac{1}{3}x^3\right]_0^4 = 3\left(48 - \frac{64}{3}\right) = 80.$

$M_y = \rho\int_0^4 x\left[\sqrt{16 - x^2} - (-2)\right]dx = \rho\int_0^4 x\sqrt{16 - x^2}\,dx + \rho\int_0^4 2x\,dx = 6\left[-\frac{1}{3}(16 - x^2)^{3/2}\right]_0^4 + 6\left[x^2\right]_0^4$

$= 6\left(0 + \frac{64}{3}\right) + 6(16) = 224.$

$\overline{x} = \dfrac{M_y}{m} = \dfrac{224}{24(\pi + 2)} = \dfrac{28}{3(\pi + 2)}$ and $\overline{y} = \dfrac{M_x}{m} = \dfrac{80}{24(\pi + 2)} = \dfrac{10}{3(\pi + 2)}$.

Thus, the center of mass is $\left(\dfrac{28}{3(\pi + 2)}, \dfrac{10}{3(\pi + 2)}\right) \approx (1.82, 0.65)$.

36. We'll use $n = 8$, so $\Delta x = \frac{b - a}{n} = \frac{8 - 0}{8} = 1$.

$A = \int_0^8 f(x)\,dx \approx S_{10} = \frac{1}{3}[f(0) + 4f(1) + 2f(2) + 4f(3) + 2f(4) + 4f(5) + 2f(6) + 4f(7) + f(8)]$

$\approx \frac{1}{3}[0 + 4(2.0) + 2(2.6) + 4(2.3) + 2(2.2) + 4(3.3) + 2(4.0) + 4(3.2) + 0]$

$= \frac{1}{3}(60.8) = 20.2\overline{6} \quad \left[\text{or } \frac{304}{15}\right]$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Now $\int_0^8 x\,f(x)\,dx \approx \frac{1}{3}[0\cdot f(0) + 4\cdot 1\cdot f(1) + 2\cdot 2\cdot f(2) + 4\cdot 3\cdot f(3)$
$+ 2\cdot 4\cdot f(4) + 4\cdot 5\cdot f(5) + 2\cdot 6\cdot f(6) + 4\cdot 7\cdot f(7) + 8\cdot f(8)]$
$\approx \frac{1}{3}[0 + 8 + 10.4 + 27.6 + 17.6 + 66 + 48 + 89.6 + 0]$
$= \frac{1}{3}(267.2) = 89.0\overline{6}$ $\left[\text{or } \frac{1336}{15}\right]$, so $\overline{x} = \frac{1}{A}\int_0^8 xf(x)\,dx \approx 4.39$.

Also, $\int_0^8 [f(x)]^2\,dx \approx \frac{1}{3}[0^2 + 4(2.0)^2 + 2(2.6)^2 + 4(2.3)^2 + 2(2.2)^2 + 4(3.3)^2 + 2(4.0)^2 + 4(3.2)^2 + 0^2]$
$= \frac{1}{3}(176.88) = 58.96$, so $\overline{y} = \frac{1}{A}\int_0^8 \frac{1}{2}[f(x)]^2\,dx \approx 1.45$.

Thus, the centroid is $(\overline{x}, \overline{y}) \approx (4.4, 1.5)$.

37. $A = \int_{-1}^1 [(x^3 - x) - (x^2 - 1)]\,dx = \int_{-1}^1 (1 - x^2)\,dx$ $\left[\begin{array}{c}\text{odd-degree terms}\\ \text{drop out}\end{array}\right]$

$= 2\int_0^1 (1 - x^2)\,dx = 2\left[x - \frac{1}{3}x^3\right]_0^1 = 2\left(\frac{2}{3}\right) = \frac{4}{3}$.

$\overline{x} = \frac{1}{A}\int_{-1}^1 x(x^3 - x - x^2 + 1)\,dx = \frac{3}{4}\int_{-1}^1 (x^4 - x^2 - x^3 + x)\,dx$

$= \frac{3}{4}\int_{-1}^1 (x^4 - x^2)\,dx = \frac{3}{4}\cdot 2\int_0^1 (x^4 - x^2)\,dx$

$= \frac{3}{2}\left[\frac{1}{5}x^5 - \frac{1}{3}x^3\right]_0^1 = \frac{3}{2}\left(-\frac{2}{15}\right) = -\frac{1}{5}$.

$\overline{y} = \frac{1}{A}\int_{-1}^1 \frac{1}{2}[(x^3 - x)^2 - (x^2 - 1)^2]\,dx = \frac{3}{4}\cdot\frac{1}{2}\int_{-1}^1 (x^6 - 2x^4 + x^2 - x^4 + 2x^2 - 1)\,dx$

$= \frac{3}{8}\cdot 2\int_0^1 (x^6 - 3x^4 + 3x^2 - 1)\,dx = \frac{3}{4}\left[\frac{1}{7}x^7 - \frac{3}{5}x^5 + x^3 - x\right]_0^1 = \frac{3}{4}\left(-\frac{16}{35}\right) = -\frac{12}{35}$.

Thus, the centroid is $(\overline{x}, \overline{y}) = \left(-\frac{1}{5}, -\frac{12}{35}\right)$.

38. The curves intersect at $x = a \approx -1.315974$ and $x = b \approx 0.53727445$.

$A = \int_a^b [(2 - x^2) - e^x]\,dx = \left[2x - \frac{1}{3}x^3 - e^x\right]_a^b \approx 1.452014$.

$\overline{x} = \frac{1}{A}\int_a^b x(2 - x^2 - e^x)\,dx = \frac{1}{A}\left[x^2 - \frac{1}{4}x^4 - xe^x + e^x\right]_a^b$
≈ -0.374293

$\overline{y} = \frac{1}{A}\int_a^b \frac{1}{2}[(2 - x^2)^2 - (e^x)^2]\,dx = \frac{1}{2A}\int_a^b (4 - 4x^2 + x^4 - e^{2x})\,dx$
$= \frac{1}{2A}\left[4x - \frac{4}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{2}e^{2x}\right]_a^b \approx 1.218131$

Thus, the centroid is $(\overline{x}, \overline{y}) \approx (-0.37, 1.22)$.

39. Choose x- and y-axes so that the base (one side of the triangle) lies along the x-axis with the other vertex along the positive y-axis as shown. From geometry, we know the medians intersect at a point $\frac{2}{3}$ of the way from each vertex (along the median) to the opposite side. The median from B goes to the midpoint $\left(\frac{1}{2}(a + c), 0\right)$ of side AC, so the point of intersection of the medians is $\left(\frac{2}{3}\cdot\frac{1}{2}(a + c), \frac{1}{3}b\right) = \left(\frac{1}{3}(a + c), \frac{1}{3}b\right)$.

This can also be verified by finding the equations of two medians, and solving them simultaneously to find their point of intersection. Now let us compute the location of the centroid of the triangle. The area is $A = \frac{1}{2}(c - a)b$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\overline{x} = \frac{1}{A}\left[\int_a^0 x \cdot \frac{b}{a}(a-x)\,dx + \int_0^c x \cdot \frac{b}{c}(c-x)\,dx\right] = \frac{1}{A}\left[\frac{b}{a}\int_a^0 (ax - x^2)\,dx + \frac{b}{c}\int_0^c \left(cx - x^2\right)dx\right]$$

$$= \frac{b}{Aa}\left[\frac{1}{2}ax^2 - \frac{1}{3}x^3\right]_a^0 + \frac{b}{Ac}\left[\frac{1}{2}cx^2 - \frac{1}{3}x^3\right]_0^c = \frac{b}{Aa}\left[-\frac{1}{2}a^3 + \frac{1}{3}a^3\right] + \frac{b}{Ac}\left[\frac{1}{2}c^3 - \frac{1}{3}c^3\right]$$

$$= \frac{2}{a\,(c\,-\,a)} \cdot \frac{-a^3}{6} + \frac{2}{c\,(c\,-\,a)} \cdot \frac{c^3}{6} = \frac{1}{3\,(c\,-\,a)}(c^2 - a^2) = \frac{a\,+\,c}{3}$$

and $$\overline{y} = \frac{1}{A}\left[\int_a^0 \frac{1}{2}\left(\frac{b}{a}(a-x)\right)^2 dx + \int_0^c \frac{1}{2}\left(\frac{b}{c}(c-x)\right)^2 dx\right]$$

$$= \frac{1}{A}\left[\frac{b^2}{2a^2}\int_a^0 (a^2 - 2ax + x^2)\,dx + \frac{b^2}{2c^2}\int_0^c (c^2 - 2cx + x^2)\,dx\right]$$

$$= \frac{1}{A}\left[\frac{b^2}{2a^2}\left[a^2x - ax^2 + \tfrac{1}{3}x^3\right]_a^0 + \frac{b^2}{2c^2}\left[c^2x - cx^2 + \tfrac{1}{3}x^3\right]_0^c\right]$$

$$= \frac{1}{A}\left[\frac{b^2}{2a^2}\left(-a^3 + a^3 - \tfrac{1}{3}a^3\right) + \frac{b^2}{2c^2}\left(c^3 - c^3 + \tfrac{1}{3}c^3\right)\right] = \frac{1}{A}\left[\frac{b^2}{6}\,(-a + c)\right] = \frac{2}{(c\,-\,a)\,b} \cdot \frac{(c-a)b^2}{6} = \frac{b}{3}$$

Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\dfrac{a+c}{3}, \dfrac{b}{3}\right)$, as claimed.

Remarks: Actually the computation of $\overline{y}$ is all that is needed. By considering each side of the triangle in turn to be the base, we see that the centroid is $\frac{1}{3}$ of the way from each side to the opposite vertex and must therefore be the intersection of the medians.

The computation of $\overline{y}$ in this problem (and many others) can be simplified by using horizontal rather than vertical approximating rectangles. If the length of a thin rectangle at coordinate y is $\ell(y)$, then its area is $\ell(y)\,\Delta y$, its mass is $\rho\ell(y)\,\Delta y$, and its moment about the x-axis is $\Delta M_x = \rho y \ell(y)\,\Delta y$. Thus,

$$M_x = \int \rho y \ell(y)\,dy \qquad \text{and} \qquad \overline{y} = \frac{\int \rho y \ell(y)\,dy}{\rho A} = \frac{1}{A}\int y\ell(y)\,dy$$

In this problem, $\ell(y) = \dfrac{c-a}{b}\,(b-y)$ by similar triangles, so

$$\overline{y} = \frac{1}{A}\int_0^b \frac{c-a}{b}\,y(b-y)\,dy = \frac{2}{b^2}\int_0^b (by - y^2)\,dy = \frac{2}{b^2}\left[\tfrac{1}{2}by^2 - \tfrac{1}{3}y^3\right]_0^b = \frac{2}{b^2} \cdot \frac{b^3}{6} = \frac{b}{3}$$

Notice that only one integral is needed when this method is used.

40. The rectangle to the left of the y-axis has centroid $\left(-\frac{1}{2}, 1\right)$ and area 2. The triangle to the right of the y-axis has area 2 and centroid $\left(\frac{2}{3}, \frac{2}{3}\right)$ [by Exercise 39, the centroid is two-thirds of the way from the vertex $(0, 0)$ to the point $(1, 1)$].

$$\overline{x} = \frac{M_y}{m} = \frac{1}{m}\sum_{i=1}^{2} m_i x_i = \frac{1}{2+2}\left[2\left(-\tfrac{1}{2}\right) + 2\left(\tfrac{2}{3}\right)\right] = \tfrac{1}{4}\left(\tfrac{1}{3}\right) = \tfrac{1}{12}.$$

$$\overline{y} = \frac{M_x}{m} = \frac{1}{m}\sum_{i=1}^{2} m_i y_i = \frac{1}{2+2}\left[2(1) + 2\left(\tfrac{2}{3}\right)\right] = \tfrac{1}{4}\left(\tfrac{10}{3}\right) = \tfrac{5}{6}.$$ Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\tfrac{1}{12}, \tfrac{5}{6}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. Divide the lamina into two triangles and one rectangle with respective masses of 2, 2 and 4, so that the total mass is 8. Using the result of Exercise 39, the triangles have centroids $\left(-1, \frac{2}{3}\right)$ and $\left(1, \frac{2}{3}\right)$. The centroid of the rectangle (its center) is $\left(0, -\frac{1}{2}\right)$.

So, using Formulas 6 and 7, we have $\overline{y} = \dfrac{M_x}{m} = \dfrac{1}{m}\sum\limits_{i=1}^{3} m_i\, y_i = \frac{1}{8}\left[2\left(\frac{2}{3}\right) + 2\left(\frac{2}{3}\right) + 4\left(-\frac{1}{2}\right)\right] = \frac{1}{8}\left(\frac{2}{3}\right) = \frac{1}{12}$, and $\overline{x} = 0$, since the lamina is symmetric about the line $x = 0$. Thus, the centroid is $(\overline{x}, \overline{y}) = \left(0, \frac{1}{12}\right)$.

42. The parabola has equation $y = kx^2$ and passes through (a, b),

so $b = ka^2 \quad \Rightarrow \quad k = \dfrac{b}{a^2}$ and hence, $y = \dfrac{b}{a^2}x^2$.

$\mathcal{R}_1$ has area $A_1 = \displaystyle\int_0^a \frac{b}{a^2}x^2\,dx = \frac{b}{a^2}\left[\frac{1}{3}x^3\right]_0^a = \frac{b}{a^2}\left(\frac{a^3}{3}\right) = \frac{1}{3}ab.$

Since $\mathcal{R}$ has area ab, $\mathcal{R}_2$ has area $A_2 = ab - \frac{1}{3}ab = \frac{2}{3}ab$.

For $\mathcal{R}_1$:

$$\overline{x}_1 = \frac{1}{A_1}\int_0^a x\left(\frac{b}{a^2}x^2\right)dx = \frac{3}{ab}\,\frac{b}{a^2}\int_0^a x^3\,dx = \frac{3}{a^3}\left[\frac{1}{4}x^4\right]_0^a = \frac{3}{a^3}\left(\frac{1}{4}a^4\right) = \frac{3}{4}a$$

$$\overline{y}_1 = \frac{1}{A_1}\int_0^a \frac{1}{2}\left(\frac{b}{a^2}x^2\right)^2 dx = \frac{3}{ab}\,\frac{b^2}{2a^4}\int_0^a x^4\,dx = \frac{3b}{2a^5}\left[\frac{1}{5}x^5\right]_0^a = \frac{3b}{2a^5}\left(\frac{1}{5}a^5\right) = \frac{3}{10}b$$

Thus, the centroid for $\mathcal{R}_1$ is $(\overline{x}_1, \overline{y}_1) = \left(\frac{3}{4}a, \frac{3}{10}b\right)$.

For $\mathcal{R}_2$:

$$\overline{x}_2 = \frac{1}{A_2}\int_0^a x\left(b - \frac{b}{a^2}x^2\right)dx = \frac{3}{2ab}\int_0^a b\left(x - \frac{1}{a^2}x^3\right)dx = \frac{3}{2a}\left[\frac{1}{2}x^2 - \frac{1}{4a^2}x^4\right]_0^a$$

$$= \frac{3}{2a}\left(\frac{a^2}{2} - \frac{a^2}{4}\right) = \frac{3}{2a}\left(\frac{a^2}{4}\right) = \frac{3}{8}a$$

$$\overline{y}_2 = \frac{1}{A_2}\int_0^a \frac{1}{2}\left[(b)^2 - \left(\frac{b}{a^2}x^2\right)^2\right]dx = \frac{3}{2ab}\,\frac{1}{2}\int_0^a b^2\left(1 - \frac{1}{a^4}x^4\right)dx = \frac{3b}{4a}\left[x - \frac{1}{5a^4}x^5\right]_0^a$$

$$= \frac{3b}{4a}\left(a - \frac{1}{5}a\right) = \frac{3b}{4a}\left(\frac{4a}{5}\right) = \frac{3}{5}b$$

Thus, the centroid for $\mathcal{R}_2$ is $(\overline{x}_2, \overline{y}_2) = \left(\frac{3}{8}a, \frac{3}{5}b\right)$. Note the relationships: $A_2 = 2A_1, \overline{x}_1 = 2\,\overline{x}_2, \overline{y}_2 = 2\,\overline{y}_1$.

43. $\int_a^b (cx + d)\, f(x)\,dx = \int_a^b cx\, f(x)\,dx + \int_a^b d f(x)\,dx = c\int_a^b x\, f(x)\,dx + d\int_a^b f(x)\,dx = c\overline{x}A + d\int_a^b f(x)\,dx$ [by (8)]

$$= c\overline{x}\int_a^b f(x)\,dx + d\int_a^b f(x)\,dx = (c\overline{x} + d)\int_a^b f(x)\,dx$$

44. A sphere can be generated by rotating a semicircle about its diameter. The center of mass travels a distance

$2\pi\overline{y} = 2\pi\left(\dfrac{4r}{3\pi}\right)$ [from Example 4] $= \dfrac{8r}{3}$, so by the Theorem of Pappus, the volume of the sphere is

$V = Ad = \dfrac{\pi r^2}{2}\cdot\dfrac{8r}{3} = \dfrac{4}{3}\pi r^3.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. A cone of height h and radius r can be generated by rotating a right triangle about one of its legs as shown. By Exercise 39, $\overline{x} = \frac{1}{3}r$, so by the Theorem of Pappus, the volume of the cone is

$$V = Ad = \left(\tfrac{1}{2} \cdot \text{base} \cdot \text{height}\right) \cdot (2\pi\overline{x}) = \tfrac{1}{2}rh \cdot 2\pi\left(\tfrac{1}{3}r\right) = \tfrac{1}{3}\pi r^2 h.$$

46. From the symmetry in the figure, $\overline{y} = 4$. So the distance traveled by the centroid when rotating the triangle about the x-axis is $d = 2\pi \cdot 4 = 8\pi$. The area of the triangle is $A = \frac{1}{2}bh = \frac{1}{2}(2)(3) = 3$. By the Theorem of Pappus, the volume of the resulting solid is $Ad = 3(8\pi) = 24\pi$.

47. The curve C is the quarter-circle $y = \sqrt{16 - x^2},\ 0 \le x \le 4$. Its length L is $\frac{1}{4}(2\pi \cdot 4) = 2\pi$.

Now $y' = \frac{1}{2}(16 - x^2)^{-1/2}(-2x) = \dfrac{-x}{\sqrt{16 - x^2}} \quad \Rightarrow \quad 1 + (y')^2 = 1 + \dfrac{x^2}{16 - x^2} = \dfrac{16}{16 - x^2} \quad \Rightarrow$

$ds = \sqrt{1 + (y')^2}\,dx = \dfrac{4}{\sqrt{16 - x^2}}\,dx$, so

$\overline{x} = \dfrac{1}{L}\displaystyle\int x\,ds = \dfrac{1}{2\pi}\int_0^4 4x(16 - x^2)^{-1/2}\,dx = \dfrac{4}{2\pi}\Big[-(16 - x^2)^{1/2}\Big]_0^4 = \dfrac{2}{\pi}(0 + 4) = \dfrac{8}{\pi}$ and

$\overline{y} = \dfrac{1}{L}\displaystyle\int y\,ds = \dfrac{1}{2\pi}\int_0^4 \sqrt{16 - x^2} \cdot \dfrac{4}{\sqrt{16 - x^2}}\,dx = \dfrac{4}{2\pi}\int_0^4 dx = \dfrac{2}{\pi}\Big[x\Big]_0^4 = \dfrac{2}{\pi}(4 - 0) = \dfrac{8}{\pi}$. Thus, the centroid is $\left(\dfrac{8}{\pi}, \dfrac{8}{\pi}\right)$. Note that the centroid does not lie on the curve, but does lie on the line $y = x$, as expected, due to the symmetry of the curve.

48. (a) From Exercise 47, we have $\overline{y} = (1/L)\int y\,ds \quad \Leftrightarrow \quad \overline{y}L = \int y\,ds$. The surface area is

$S = \int 2\pi y\,ds = 2\pi\int y\,ds = 2\pi(\overline{y}L) = L(2\pi\overline{y})$, which is the product of the arc length of C and the distance traveled by the centroid of C.

(b) From Exercise 47, $L = 2\pi$ and $\overline{y} = \frac{8}{\pi}$. By the Second Theorem of Pappus, the surface area is

$S = L(2\pi\overline{y}) = 2\pi(2\pi \cdot \frac{8}{\pi}) = 32\pi$.

A geometric formula for the surface area of a half-sphere is $S = 2\pi r^2$. With $r = 4$, we get $S = 32\pi$, which agrees with our first answer.

49. The circle has arc length (circumference) $L = 2\pi r$. As in Example 7, the distance traveled by the centroid during a rotation is $d = 2\pi R$. Therefore, by the Second Theorem of Pappus, the surface area is

$$S = Ld = (2\pi r)(2\pi R) = 4\pi^2 rR$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50. (a) Let $0 \le x \le 1$. If $n < m$, then $x^n > x^m$; that is, raising x to a larger power produces a smaller number.

(b) Using Formulas 9 and the fact that the area of $\mathscr{R}$ is

$A = \int_0^1 (x^n - x^m)\,dx = \dfrac{1}{n+1} - \dfrac{1}{m+1} = \dfrac{m-n}{(n+1)(m+1)}$, we get

$$\overline{x} = \frac{(n+1)(m+1)}{m-n}\int_0^1 x[x^n - x^m]\,dx = \frac{(n+1)(m+1)}{m-n}\int_0^1 \left(x^{n+1} - x^{m+1}\right)dx$$

$$= \frac{(n+1)(m+1)}{m-n}\left[\frac{1}{n+2} - \frac{1}{m+2}\right] = \frac{(n+1)(m+1)}{(n+2)(m+2)}$$

and
$$\overline{y} = \frac{(n+1)(m+1)}{m-n}\int_0^1 \tfrac{1}{2}\left[(x^n)^2 - (x^m)^2\right]dx = \frac{(n+1)(m+1)}{2(m-n)}\int_0^1 \left(x^{2n} - x^{2m}\right)dx$$

$$= \frac{(n+1)(m+1)}{2(m-n)}\left[\frac{1}{2n+1} - \frac{1}{2m+1}\right] = \frac{(n+1)(m+1)}{(2n+1)(2m+1)}$$

(c) If we take $n = 3$ and $m = 4$, then

$$(\overline{x}, \overline{y}) = \left(\frac{4\cdot 5}{5\cdot 6}, \frac{4\cdot 5}{7\cdot 9}\right) = \left(\frac{2}{3}, \frac{20}{63}\right)$$

which lies outside $\mathscr{R}$ since $\left(\frac{2}{3}\right)^3 = \frac{8}{27} < \frac{20}{63}$. This is the simplest of many possibilities.

51. Suppose the region lies between two curves $y = f(x)$ and $y = g(x)$ where $f(x) \ge g(x)$, as illustrated in Figure 13. Choose points x_i with $a = x_0 < x_1 < \cdots < x_n = b$ and choose x_i^* to be the midpoint of the ith subinterval; that is, $x_i^* = \overline{x}_i = \frac{1}{2}(x_{i-1} + x_i)$. Then the centroid of the ith approximating rectangle R_i is its center $C_i = \left(\overline{x}_i, \frac{1}{2}[f(\overline{x}_i) + g(\overline{x}_i)]\right)$. Its area is $[f(\overline{x}_i) - g(\overline{x}_i)]\,\Delta x$, so its mass is $\rho[f(\overline{x}_i) - g(\overline{x}_i)]\,\Delta x$. Thus, $M_y(R_i) = \rho[f(\overline{x}_i) - g(\overline{x}_i)]\,\Delta x \cdot \overline{x}_i = \rho\overline{x}_i\,[f(\overline{x}_i) - g(\overline{x}_i)]\,\Delta x$ and $M_x(R_i) = \rho[f(\overline{x}_i) - g(\overline{x}_i)]\,\Delta x \cdot \frac{1}{2}[f(\overline{x}_i) + g(\overline{x}_i)] = \rho \cdot \frac{1}{2}\left[f(\overline{x}_i)^2 - g(\overline{x}_i)^2\right]\Delta x$. Summing over i and taking the limit as $n \to \infty$, we get $M_y = \lim\limits_{n\to\infty} \sum_i \rho\overline{x}_i\,[f(\overline{x}_i) - g(\overline{x}_i)]\,\Delta x = \rho\int_a^b x[f(x) - g(x)]\,dx$ and $M_x = \lim\limits_{n\to\infty} \sum_i \rho \cdot \frac{1}{2}\left[f(\overline{x}_i)^2 - g(\overline{x}_i)^2\right]\Delta x = \rho\int_a^b \frac{1}{2}\left[f(x)^2 - g(x)^2\right]dx$.

Thus, $\overline{x} = \dfrac{M_y}{m} = \dfrac{M_y}{\rho A} = \dfrac{1}{A}\displaystyle\int_a^b x[f(x) - g(x)]\,dx$ and $\overline{y} = \dfrac{M_x}{m} = \dfrac{M_x}{\rho A} = \dfrac{1}{A}\displaystyle\int_a^b \frac{1}{2}\left[f(x)^2 - g(x)^2\right]dx$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

DISCOVERY PROJECT Complementary Coffee Cups

1. Cup A has volume $V_A = \int_0^h \pi[f(y)]^2\,dy$ and cup B has volume

$$V_B = \int_0^h \pi[k - f(y)]^2\,dy = \int_0^h \pi\{k^2 - 2kf(y) + [f(y)]^2\}\,dy$$
$$= \left[\pi k^2 y\right]_0^h - 2\pi k \int_0^h f(y)\,dy + \int_0^h \pi[f(y)]^2\,dy = \pi k^2 h - 2\pi k A_1 + V_A$$

Thus, $V_A = V_B \quad\Leftrightarrow\quad \pi k(kh - 2A_1) = 0 \quad\Leftrightarrow\quad k = 2(A_1/h)$; that is, k is twice the average value of f on the interval $[0, h]$.

2. From Problem 1, $V_A = V_B \quad\Leftrightarrow\quad kh = 2A_1 \quad\Leftrightarrow\quad A_1 + A_2 = 2A_1 \quad\Leftrightarrow\quad A_2 = A_1$.

3. Let $\overline{x}_1$ and $\overline{x}_2$ denote the x-coordinates of the centroids of A_1 and A_2, respectively. By Pappus's Theorem,

$V_A = 2\pi\overline{x}_1 A_1$ and $V_B = 2\pi(k - \overline{x}_2)A_2$, so $V_A = V_B \quad\Leftrightarrow\quad \overline{x}_1 A_1 = kA_2 - \overline{x}_2 A_2 \quad\Leftrightarrow\quad kA_2 = \overline{x}_1 A_1 + \overline{x}_2 A_2 \overset{(\star)}{\Leftrightarrow}$ $kA_2 = \frac{1}{2}k\,(A_1 + A_2) \quad\Leftrightarrow\quad \frac{1}{2}kA_2 = \frac{1}{2}kA_1 \quad\Leftrightarrow\quad A_2 = A_1$, as shown in Problem 2. [$(\star)$ The sum of the moments of the regions of areas A_1 and A_2 about the y-axis equals the moment of the entire k-by-h rectangle about the y-axis.]

So, since $A_1 + A_2 = kh$, we have $V_A = V_B \quad\Leftrightarrow\quad A_1 = A_2 \quad\Leftrightarrow\quad A_1 = \frac{1}{2}(A_1 + A_2) \quad\Leftrightarrow\quad A_1 = \frac{1}{2}(kh) \quad\Leftrightarrow$ $k = 2\,(A_1/h)$, as shown in Problem 1.

4. We'll use a cup that is $h = 8$ cm high with a diameter of 6 cm on the top and the bottom and symmetrically bulging to a diameter of 8 cm in the middle (all inside dimensions).

For an equation, we'll use a parabola with a vertex at $(4, 4)$; that is, $x = a(y-4)^2 + 4$. To find a, use the point $(3, 0)$: $3 = a(0-4)^2 + 4 \quad\Rightarrow\quad -1 = 16a \quad\Rightarrow\quad a = -\frac{1}{16}$. To find k, we'll use the relationship in Problem 1, so we need A_1.

$$A_1 = \int_0^8 \left[-\tfrac{1}{16}(y-4)^2 + 4\right] dy = \int_{-4}^4 \left(-\tfrac{1}{16}u^2 + 4\right) du \quad [u = y - 4]$$
$$= 2\int_0^4 \left(-\tfrac{1}{16}u^2 + 4\right) du = 2\left[-\tfrac{1}{48}u^3 + 4u\right]_0^4 = 2\left(-\tfrac{4}{3} + 16\right) = \tfrac{88}{3}.$$

Thus, $k = 2(A_1/h) = 2\left(\frac{88/3}{8}\right) = \frac{22}{3}$.

So with $h = 8$ and curve $x = -\frac{1}{16}(y-4)^2 + 4$, we have

$$V_A = \int_0^8 \pi\left[-\tfrac{1}{16}(y-4)^2 + 4\right]^2 dy = \pi\int_{-4}^4 \left(-\tfrac{1}{16}u^2 + 4\right)^2 du \quad [u = y - 4] \quad = 2\pi\int_0^4 \left(\tfrac{1}{256}u^4 - \tfrac{1}{2}u^2 + 16\right) du$$
$$= 2\pi\left[\tfrac{1}{1280}u^5 - \tfrac{1}{6}u^3 + 16u\right]_0^4 = 2\pi\left(\tfrac{4}{5} - \tfrac{32}{3} + 64\right) = 2\pi\left(\tfrac{812}{15}\right) = \tfrac{1624}{15}\pi$$

This is approximately 340 cm^3 or 11.5 fl. oz. And with $k = \frac{22}{3}$, we know from Problem 1 that cup B holds the same amount.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8.4 Applications to Economics and Biology

1. By the Net Change Theorem, $C(4000) - C(0) = \int_0^{4000} C'(x)\,dx \quad \Rightarrow$

$$C(4000) = 18{,}000 + \int_0^{4000} (0.82 - 0.000\,03x + 0.000\,000\,003x^2)\,dx$$
$$= 18{,}000 + \left[0.82x - 0.000\,015x^2 + 0.000\,000\,001x^3\right]_0^{4000} = 18{,}000 + 3104 = \$21{,}104$$

2. By the Net Change Theorem,

$$R(10{,}000) - R(5000) = \int_{5000}^{10{,}000} R'(x)\,dx = \int_{5000}^{10{,}000} (48 - 0.0012x)\,dx = \left[48x - 0.0006x^2\right]_{5000}^{10{,}000}$$
$$= 420{,}000 - 225{,}000 = \$195{,}000$$

3. By the Net Change Theorem, $C(50) - C(0) = \int_0^{50} (0.6 + 0.008x)\,dx \quad \Rightarrow$

$C(50) = 100 + \left[0.6x + 0.004x^2\right]_0^{50} = 100 + (40 - 0) = 140$, or \$140,000.

Similarly, $C(100) - C(50) = \left[0.6x + 0.004x^2\right]_{50}^{100} = 100 - 40 = 60$, or \$60,000.

4. Consumer surplus $= \int_0^{400} [p(x) - p(400)]\,dx = \int_0^{400} [(2000 - 46\sqrt{x}\,) - 1080]\,dx$

$$= \int_0^{400} (920 - 46\sqrt{x}\,)\,dx = 46\int_0^{400} (20 - x^{1/2})\,dx$$
$$= 46\left[20x - \tfrac{2}{3}x^{3/2}\right]_0^{400} = 46\left(8000 - \tfrac{2}{3}\cdot 8000\right)$$
$$= 46 \cdot \tfrac{1}{3} \cdot 8000 \approx \$122{,}666.67$$

5. Consumer surplus $= \int_0^{45} [p(x) - p(45)]\,dx$

$$= \int_0^{45} (870e^{-0.03x} - 870e^{-0.03(45)})\,dx$$
$$= 870\left[-\frac{1}{0.03}e^{-0.03x} - e^{-1.35}x\right]_0^{45}$$
$$= 870\left(-\frac{1}{0.03}e^{-1.35} - 45e^{-1.35} + \frac{1}{0.03}\right) \approx \$11{,}332.78$$

6. $p = 2.80 \quad \Rightarrow \quad 6 - \dfrac{x}{3500} = 2.80 \quad \Rightarrow \quad \dfrac{x}{3500} = 3.2 \quad \Rightarrow \quad x = 11{,}200$

$$\text{Consumer surplus} = \int_0^{11{,}200} [p(x) - 2.80]\,dx = \int_0^{11{,}200} \left(6 - \frac{x}{3500} - 2.80\right) dx$$
$$= \int_0^{11{,}200} \left(3.2 - \frac{x}{3500}\right) dx = \left[3.2x - \frac{x^2}{7000}\right]_0^{11{,}200} = 35{,}840 - 17{,}920 = \$17{,}920$$

7. Since the demand increases by 30 for each dollar the price is lowered, the demand function, $p(x)$, is linear with slope $-\frac{1}{30}$. Also, $p(210) = 18$, so an equation for the demand is $p - 18 = -\frac{1}{30}(x - 210)$ or $p = -\frac{1}{30}x + 25$. A selling price of \$15 implies that $15 = -\frac{1}{30}x + 25 \quad \Rightarrow \quad \frac{1}{30}x = 10 \quad \Rightarrow \quad x = 300$.

Consumer surplus $= \int_0^{300} \left(-\frac{1}{30}x + 25 - 15\right) dx = \int_0^{300} \left(-\frac{1}{30}x + 10\right) dx = \left[-\frac{1}{60}x^2 + 10x\right]_0^{300} = \$1500$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $p(x) = \dfrac{800{,}000e^{-x/5000}}{x + 20{,}000} = 16 \quad \Rightarrow \quad x = x_1 \approx 3727.04.$

Consumer surplus $= \int_0^{x_1} [p(x) - 16]\, dx \approx \$37{,}753$

9. $p_S(x) = 3 + 0.01x^2. \quad P = p_S(10) = 3 + 1 = 4.$

$$\text{Producer surplus} = \int_0^{10} [P - p_S(x)]\, dx = \int_0^{10} \left[4 - 3 - 0.01x^2\right] dx$$
$$= \left[x - \tfrac{0.01}{3}x^3\right]_0^{10} \approx 10 - 3.33 = \$6.67$$

10. $P = p_S(x) \quad \Rightarrow \quad 625 = 125 + 0.002x^2 \quad \Rightarrow \quad 500 = \frac{1}{500}x^2 \quad \Rightarrow \quad x^2 = 500^2 \quad \Rightarrow \quad x = 500.$

$$\text{Producer surplus} = \int_0^{500} [P - p_S(x)]\, dx = \int_0^{500} [625 - (125 + 0.002x^2)]\, dx = \int_0^{500} \left(500 - \tfrac{1}{500}x^2\right) dx$$
$$= \left[500x - \tfrac{1}{1500}x^3\right]_0^{500} = 500^2 - \tfrac{1}{1500}(500^3) \approx \$166{,}666.67$$

11. $p = \sqrt{30 + 0.01xe^{0.001x}} = 30$ when $x \approx 3278.5$ (using a graphing calculator or other computing device). Then the producer surplus is approximately $\int_0^{3278.5} \left[30 - \sqrt{30 + 0.01xe^{0.001x}}\,\right] dx \approx \$55{,}735.$

12. (a) Demand curve $p_D(x) =$ supply curve $p_S(x) \quad \Leftrightarrow \quad 50 - \frac{1}{20}x = 20 + \frac{1}{10}x \quad \Leftrightarrow \quad 30 = \frac{3}{20}x \quad \Leftrightarrow \quad x = 200.$

$p_D(200) = 50 - \frac{1}{20}(200) = 40$, so the market for this good is in equilibrium when the quantity is 200 and the price is \$40.

(b) At equilibrium, the

$$\text{Consumer surplus} = \int_0^{200} [p_D(x) - 40]\, dx = \int_0^{200} \left(50 - \tfrac{1}{20}x - 40\right) dx$$
$$= \left[10x - \tfrac{1}{40}x^2\right]_0^{200} = \$1000$$

and the

$$\text{Producer surplus} = \int_0^{200} [40 - p_S(x)]\, dx = \int_0^{200} \left(40 - 20 - \tfrac{1}{10}x\right) dx$$
$$= \left[20x - \tfrac{1}{20}x^2\right]_0^{200} = \$2000$$

13. (a) Demand function $p(x) =$ supply function $p_S(x) \quad \Leftrightarrow \quad 228.4 - 18x = 27x + 57.4 \quad \Leftrightarrow \quad 171 = 45x \quad \Leftrightarrow$ $x = \frac{19}{5}$ [3.8 thousand]. $p(3.8) = 228.4 - 18(3.8) = 160$. The market for the stereos is in equilibrium when the quantity is 3800 and the price is \$160.

(b) $$\text{Consumer surplus} = \int_0^{3.8} [p(x) - 160]\, dx = \int_0^{3.8} (228.4 - 18x - 160)\, dx = \int_0^{3.8} (68.4 - 18x)\, dx$$
$$= \Big[68.4x - 9x^2\Big]_0^{3.8} = 68.4(3.8) - 9(3.8)^2 = 129.96$$

$$\text{Producer surplus} = \int_0^{3.8} [160 - p_S(x)]\, dx = \int_0^{3.8} [160 - (27x + 57.4)]\, dx = \int_0^{3.8} (102.6 - 27x)\, dx$$
$$= \Big[102.6x - 13.5x^2\Big]_0^{3.8} = 102.6(3.8) - 13.5(3.8)^2 = 194.94$$

Thus, the maximum total surplus for the stereos is $129.96 + 194.94 = 324.9$, or \$324,900.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $p(x) = p_S(x) \quad \Leftrightarrow \quad 312e^{-0.14x} = 26e^{0.2x} \quad \Leftrightarrow \quad \dfrac{312}{26} = \dfrac{e^{0.2x}}{e^{-0.14x}} \quad \Leftrightarrow \quad 12 = e^{0.34x} \quad \Leftrightarrow \quad \ln 12 = 0.34x \quad \Leftrightarrow$

$x = X = \dfrac{\ln 12}{0.34}$. $\quad X \approx 7.3085$ (in thousands) and $p(X) \approx 112.1465$.

Consumer surplus $= \int_0^X [p(x) - p(X)]\,dx \approx \int_0^{7.3085} (312e^{-0.14x} - 112.1465)\,dx \approx 607.896$

Producer surplus $= \int_0^X [p_S(X) - p_S(x)]\,dx \approx \int_0^{7.3085} (112.1465 - 26e^{0.2x})\,dx \approx 388.896$

Maximum total surplus $\approx 607.896 + 388.896 = 996.792$, or \$996,792.

Note: Since $p(X) = p_S(X)$, the maximum total surplus could be found by calculating $\int_0^X [p(x) - p_S(x)]\,dx$.

15. $f(8) - f(4) = \int_4^8 f'(t)\,dt = \int_4^8 \sqrt{t}\,dt = \left[\frac{2}{3}t^{3/2}\right]_4^8 = \frac{2}{3}\left(16\sqrt{2} - 8\right) \approx \$9.75$ million

16. The total revenue R obtained in the first four years is

$$R = \int_0^4 f(t)\,dt = \int_0^4 9000\sqrt{1+2t}\,dt = \int_1^9 9000u^{1/2}\left(\tfrac{1}{2}\,du\right) \qquad [u = 1 + 2t,\ du = 2\,dt]$$

$$= 4500\left[\tfrac{2}{3}u^{3/2}\right]_1^9 = 3000(27 - 1) = \$78{,}000$$

17. Future value $= \int_0^T f(t)\,e^{r(T-t)}\,dt = \int_0^6 8000e^{0.04t}\,e^{0.062(6-t)}\,dt = 8000\int_0^6 e^{0.04t}\,e^{0.372-0.062t}\,dt$

$$= 8000\int_0^6 e^{0.372-0.022t}\,dt = 8000e^{0.372}\int_0^6 e^{-0.022t}\,dt = 8000e^{0.372}\left[\frac{e^{-0.022t}}{-0.022}\right]_0^6$$

$$= \frac{8000e^{0.372}}{-0.022}(e^{-0.132} - 1) \approx \$65{,}230.48$$

18. Present value $= \int_0^T f(t)\,e^{-rt}\,dt = \int_0^6 8000e^{0.04t}\,e^{-0.062t}\,dt = 8000\int_0^6 e^{-0.022t}\,dt = 8000\left[\dfrac{e^{-0.022t}}{-0.022}\right]_0^6$

$$= \frac{8000}{-0.022}\left(e^{-0.132} - 1\right) \approx \$44{,}966.91$$

19. $N = \displaystyle\int_a^b Ax^{-k}\,dx = A\left[\frac{x^{-k+1}}{-k+1}\right]_a^b = \frac{A}{1-k}\left(b^{1-k} - a^{1-k}\right)$.

Similarly, $\displaystyle\int_a^b Ax^{1-k}\,dx = A\left[\frac{x^{2-k}}{2-k}\right]_a^b = \frac{A}{2-k}\left(b^{2-k} - a^{2-k}\right)$.

Thus, $\overline{x} = \dfrac{1}{N}\displaystyle\int_a^b Ax^{1-k}\,dx = \frac{[A/(2-k)](b^{2-k} - a^{2-k})}{[A/(1-k)](b^{1-k} - a^{1-k})} = \frac{(1-k)(b^{2-k} - a^{2-k})}{(2-k)(b^{1-k} - a^{1-k})}$.

20. $n(9) - n(5) = \displaystyle\int_5^9 (2200 + 10e^{0.8t})\,dt = \left[2200t + \frac{10e^{0.8t}}{0.8}\right]_5^9 = \left[2200t\right]_5^9 + \tfrac{25}{2}\left[e^{0.8t}\right]_5^9$

$$= 2200(9 - 5) + 12.5(e^{7.2} - e^4) \approx 24{,}860$$

21. $F = \dfrac{\pi PR^4}{8\eta l} = \dfrac{\pi(4000)(0.008)^4}{8(0.027)(2)} \approx 1.19 \times 10^{-4}\ \text{cm}^3/\text{s}$

22. If the flux remains constant, then $\dfrac{\pi P_0 R_0^4}{8\eta l} = \dfrac{\pi PR^4}{8\eta l} \quad \Rightarrow \quad P_0R_0^4 = PR^4 \quad \Rightarrow \quad \dfrac{P}{P_0} = \left(\dfrac{R_0}{R}\right)^4$.

$R = \frac{3}{4}R_0 \quad \Rightarrow \quad \dfrac{P}{P_0} = \left(\dfrac{R_0}{\frac{3}{4}R_0}\right)^4 \quad \Rightarrow \quad P = P_0\left(\frac{4}{3}\right)^4 \approx 3.1605P_0 > 3P_0$; that is, the blood pressure is more than tripled.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. From (3), $F = \dfrac{A}{\int_0^T c\,(t)\ dt} = \dfrac{6}{20I}$, where

$$I = \int_0^{10} te^{-0.6t}\,dt = \left[\frac{1}{(-0.6)^2}\,(-0.6t-1)\,e^{-0.6t}\right]_0^{10} \left[\begin{array}{c}\text{integrating}\\ \text{by parts}\end{array}\right] = \tfrac{1}{0.36}(-7e^{-6}+1)$$

Thus, $F = \dfrac{6(0.36)}{20(1-7e^{-6})} = \dfrac{0.108}{1-7e^{-6}} \approx 0.1099$ L/s or 6.594 L/min.

24. As in Example 2, we will estimate the cardiac output using Simpson's Rule with $\Delta t = (16-0)/8 = 2$.

$$\begin{aligned}\textstyle\int_0^{16} c(t)\,dt &\approx \tfrac{2}{3}[c(0)+4c(2)+2c(4)+4c(6)+2c(8)+4c(10)+2c(12)+4c(14)+c(16)]\\ &= \tfrac{2}{3}[0+4(4.1)+2(8.9)+4(8.5)+2(6.7)+4(4.3)+2(2.5)+4(1.2)+0.2]\\ &= \tfrac{2}{3}(108.8) = 72.5\overline{3}\ \text{mg}\cdot\text{s/L}\end{aligned}$$

Therefore, $F \approx \dfrac{A}{72.5\overline{3}} = \dfrac{5.5}{72.5\overline{3}} \approx 0.0758$ L/s or 4.55 L/min.

25. As in Example 2, we will estimate the cardiac output using Simpson's Rule with $\Delta t = (16-0)/8 = 2$.

$$\begin{aligned}\textstyle\int_0^{16} c(t)\,dt &\approx \tfrac{2}{3}[c(0)+4c(2)+2c(4)+4c(6)+2c(8)+4c(10)+2c(12)+4c(14)+c(16)]\\ &\approx \tfrac{2}{3}[0+4(6.1)+2(7.4)+4(6.7)+2(5.4)+4(4.1)+2(3.0)+4(2.1)+1.5]\\ &= \tfrac{2}{3}(109.1) = 72.7\overline{3}\ \text{mg}\cdot\text{s/L}\end{aligned}$$

Therefore, $F \approx \dfrac{A}{72.7\overline{3}} = \dfrac{7}{72.7\overline{3}} \approx 0.0962$ L/s or 5.77 L/min.

8.5 Probability

1. (a) $\int_{30,000}^{40,000} f(x)\,dx$ is the probability that a randomly chosen tire will have a lifetime between 30,000 and 40,000 miles.

(b) $\int_{25,000}^{\infty} f(x)\,dx$ is the probability that a randomly chosen tire will have a lifetime of at least 25,000 miles.

2. (a) The probability that you drive to school in less than 15 minutes is $\int_0^{15} f(t)\,dt$.

(b) The probability that it takes you more than half an hour to get to school is $\int_{30}^{\infty} f(t)\,dt$.

3. (a) In general, we must satisfy the two conditions that are mentioned before Example 1—namely, **(1)** $f(x) \geq 0$ for all x, and **(2)** $\int_{-\infty}^{\infty} f(x)\,dx = 1$. For $0 \leq x \leq 1$, $f(x) = 30x^2(1-x)^2 \geq 0$ and $f(x) = 0$ for all other values of x, so $f(x) \geq 0$ for all x. Also,

$$\begin{aligned}\textstyle\int_{-\infty}^{\infty} f(x)\,dx &= \int_0^1 30x^2(1-x)^2\,dx = \int_0^1 30x^2(1-2x+x^2)\,dx = \int_0^1 (30x^2-60x^3+30x^4)\,dx\\ &= \left[10x^3-15x^4+6x^5\right]_0^1 = 10-15+6 = 1\end{aligned}$$

Therefore, f is a probability density function.

(b) $P\left(X \leq \frac{1}{3}\right) = \int_{-\infty}^{1/3} f(x)\,dx = \int_0^{1/3} 30x^2(1-x)^2\,dx = \left[10x^3-15x^4+6x^5\right]_0^{1/3} = \frac{10}{27} - \frac{15}{81} + \frac{6}{243} = \frac{17}{81}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a) In general, we must satisfy the two conditions that are mentioned before Example 1 — namely, **(1)** $f(x) \geq 0$ for all x, and **(2)** $\int_{-\infty}^{\infty} f(x)\,dx = 1$. For $f(x) = \dfrac{e^{3-x}}{(1+e^{3-x})^2}$, the numerator and denominator are both positive, so $f(x) \geq 0$ for all x.

Also,

$$\int_{-\infty}^{\infty} f(x)\,dx = \int_{-\infty}^{0} f(x)\,dx + \int_{0}^{\infty} f(x)\,dx = \lim_{t\to-\infty}\int_t^0 \frac{e^{3-x}}{(1+e^{3-x})^2}\,dx + \lim_{s\to\infty}\int_0^s \frac{e^{3-x}}{(1+e^{3-x})^2}\,dx$$

$$= \lim_{t\to-\infty}\int_{x=t}^{0} \frac{-du}{u^2} + \lim_{s\to\infty}\int_{x=0}^{s} \frac{-du}{u^2} \qquad \begin{bmatrix} u = 1+e^{3-x}, \\ du = -e^{3-x}\,dx \end{bmatrix}$$

$$= \lim_{t\to-\infty}\left[\frac{1}{u}\right]_{x=t}^{0} + \lim_{s\to\infty}\left[\frac{1}{u}\right]_{x=0}^{s} = \lim_{t\to-\infty}\left[\frac{1}{1+e^{3-x}}\right]_t^0 + \lim_{s\to\infty}\left[\frac{1}{1+e^{3-x}}\right]_0^s$$

$$= \lim_{t\to-\infty}\left(\frac{1}{1+e^3} - \frac{1}{1+e^{3-t}}\right) + \lim_{s\to\infty}\left(\frac{1}{1+e^{3-s}} - \frac{1}{1+e^3}\right) = \frac{1}{1+e^3} - 0 + 1 - \frac{1}{1+e^3} = 1.$$

Therefore, f is a probability density function.

(b) $P(3 \leq X \leq 4) = \displaystyle\int_3^4 f(x)\,dx = \left[\frac{1}{1+e^{3-x}}\right]_3^4$ [from part (a)] $= \dfrac{1}{1+e^{-1}} - \dfrac{1}{1+1} \approx 0.231$

(c)

The graph of f appears to be symmetric about the line $x = 3$, so the mean appears to be 3. Similarly, half the area under the graph of f appears to lie to the right of $x = 3$, so the median also appears to be 3.

5. (a) In general, we must satisfy the two conditions that are mentioned before Example 1—namely, **(1)** $f(x) \geq 0$ for all x, and **(2)** $\int_{-\infty}^{\infty} f(x)\,dx = 1$. If $c \geq 0$, then $f(x) \geq 0$, so condition (1) is satisfied. For condition (2), we see that

$$\int_{-\infty}^{\infty} f(x)\,dx = \int_{-\infty}^{\infty} \frac{c}{1+x^2}\,dx \text{ and}$$

$$\int_0^{\infty} \frac{c}{1+x^2}\,dx = \lim_{t\to\infty}\int_0^t \frac{c}{1+x^2}\,dx = c\lim_{t\to\infty}\left[\tan^{-1}x\right]_0^t = c\lim_{t\to\infty}\tan^{-1}t = c\left(\frac{\pi}{2}\right)$$

Similarly, $\displaystyle\int_{-\infty}^{0} \frac{c}{1+x^2}\,dx = c\left(\frac{\pi}{2}\right)$, so $\displaystyle\int_{-\infty}^{\infty} \frac{c}{1+x^2}\,dx = 2c\left(\frac{\pi}{2}\right) = c\pi.$

Since $c\pi$ must equal 1, we must have $c = 1/\pi$ so that f is a probability density function.

(b) $P(-1 < X < 1) = \displaystyle\int_{-1}^{1} \frac{1/\pi}{1+x^2}\,dx = \frac{2}{\pi}\int_0^1 \frac{1}{1+x^2}\,dx = \frac{2}{\pi}\left[\tan^{-1}x\right]_0^1 = \frac{2}{\pi}\left(\frac{\pi}{4} - 0\right) = \frac{1}{2}$

6. (a) For $0 \leq x \leq 3$, we have $f(x) = k(3x - x^2)$, which is nonnegative if and only if $k \geq 0$. Also, $\int_{-\infty}^{\infty} f(x)\,dx = \int_0^3 k(3x - x^2)\,dx = k\left[\frac{3}{2}x^2 - \frac{1}{3}x^3\right]_0^3 = k\left(\frac{27}{2} - 9\right) = \frac{9}{2}k$. Now $\frac{9}{2}k = 1 \quad\Rightarrow\quad k = \frac{2}{9}$. Therefore, f is a probability density function if and only if $k = \frac{2}{9}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Let $k = \frac{2}{9}$.

$P(X > 1) = \int_1^\infty f(x)\,dx = \int_1^3 \frac{2}{9}(3x - x^2)\,dx = \frac{2}{9}\left[\frac{3}{2}x^2 - \frac{1}{3}x^3\right]_1^3 = \frac{2}{9}\left[\left(\frac{27}{2} - 9\right) - \left(\frac{3}{2} - \frac{1}{3}\right)\right] = \frac{2}{9}\left(\frac{10}{3}\right) = \frac{20}{27}$.

(c) The mean $\mu = \int_{-\infty}^\infty xf(x)\,dx = \int_0^3 x\left[\frac{2}{9}(3x - x^2)\right]dx = \frac{2}{9}\int_0^3 (3x^2 - x^3)\,dx$

$= \frac{2}{9}\left[x^3 - \frac{1}{4}x^4\right]_0^3 = \frac{2}{9}\left(27 - \frac{81}{4}\right) = \frac{2}{9}\left(\frac{27}{4}\right) = \frac{3}{2}$.

7. (a) In general, we must satisfy the two conditions that are mentioned before Example 1—namely, **(1)** $f(x) \geq 0$ for all x, and **(2)** $\int_{-\infty}^\infty f(x)\,dx = 1$. Since $f(x) = 0$ or $f(x) = 0.1$, condition (1) is satisfied. For condition (2), we see that $\int_{-\infty}^\infty f(x)\,dx = \int_0^{10} 0.1\,dx = \left[\frac{1}{10}x\right]_0^{10} = 1$. Thus, $f(x)$ is a probability density function for the spinner's values.

(b) Since all the numbers between 0 and 10 are equally likely to be selected, we expect the mean to be halfway between the endpoints of the interval; that is, $x = 5$.

$$\mu = \int_{-\infty}^\infty xf(x)\,dx = \int_0^{10} x(0.1)\,dx = \left[\frac{1}{20}x^2\right]_0^{10} = \frac{100}{20} = 5, \quad \text{as expected.}$$

8. (a) As in the preceding exercise, **(1)** $f(x) \geq 0$ and **(2)** $\int_{-\infty}^\infty f(x)\,dx = \int_0^{10} f(x)\,dx = \frac{1}{2}(10)(0.2)$ [area of a triangle] $= 1$. So $f(x)$ is a probability density function.

(b) (i) $P(X < 3) = \int_0^3 f(x)\,dx = \frac{1}{2}(3)(0.1) = \frac{3}{20} = 0.15$

(ii) We first compute $P(X > 8)$ and then subtract that value and our answer in (i) from 1 (the total probability).

$P(X > 8) = \int_8^{10} f(x)\,dx = \frac{1}{2}(2)(0.1) = \frac{2}{20} = 0.10$. So $P(3 \leq X \leq 8) = 1 - 0.15 - 0.10 = 0.75$.

(c) We find equations of the lines from $(0, 0)$ to $(6, 0.2)$ and from $(6, 0.2)$ to $(10, 0)$, and find that

$$f(x) = \begin{cases} \frac{1}{30}x & \text{if } 0 \leq x < 6 \\ -\frac{1}{20}x + \frac{1}{2} & \text{if } 6 \leq x < 10 \\ 0 & \text{otherwise} \end{cases}$$

$$\mu = \int_{-\infty}^\infty xf(x)\,dx = \int_0^6 x\left(\frac{1}{30}x\right)dx + \int_6^{10} x\left(-\frac{1}{20}x + \frac{1}{2}\right)dx = \left[\frac{1}{90}x^3\right]_0^6 + \left[-\frac{1}{60}x^3 + \frac{1}{4}x^2\right]_6^{10}$$
$$= \frac{216}{90} + \left(-\frac{1000}{60} + \frac{100}{4}\right) - \left(-\frac{216}{60} + \frac{36}{4}\right) = \frac{16}{3} = 5.\overline{3}$$

9. We need to find m so that $\int_m^\infty f(t)\,dt = \frac{1}{2} \;\Rightarrow\; \lim_{x\to\infty} \int_m^x \frac{1}{5}e^{-t/5}\,dt = \frac{1}{2} \;\Rightarrow\; \lim_{x\to\infty} \left[\frac{1}{5}(-5)e^{-t/5}\right]_m^x = \frac{1}{2} \;\Rightarrow$ $(-1)(0 - e^{-m/5}) = \frac{1}{2} \;\Rightarrow\; e^{-m/5} = \frac{1}{2} \;\Rightarrow\; -m/5 = \ln\frac{1}{2} \;\Rightarrow\; m = -5\ln\frac{1}{2} = 5\ln 2 \approx 3.47$ min.

10. (a) $\mu = 1000 \;\Rightarrow\; f(t) = \begin{cases} 0 & \text{if } t < 0 \\ \frac{1}{1000}e^{-t/1000} & \text{if } t \geq 0 \end{cases}$

(i) $P(0 \leq X \leq 200) = \int_0^{200} \frac{1}{1000}e^{-t/1000}\,dt = \left[-e^{-t/1000}\right]_0^{200} = -e^{-1/5} + 1 \approx 0.181$

(ii) $P(X > 800) = \int_{800}^\infty \frac{1}{1000}e^{-t/1000}\,dt = \lim_{x\to\infty}\left[-e^{-t/1000}\right]_{800}^x = 0 + e^{-4/5} \approx 0.449$

(b) We need to find m so that $\int_m^\infty f(t)\,dt = \frac{1}{2} \;\Rightarrow\; \lim_{x\to\infty} \int_m^x \frac{1}{1000}e^{-t/1000}\,dt = \frac{1}{2} \;\Rightarrow\; \lim_{x\to\infty}\left[-e^{-t/1000}\right]_m^x = \frac{1}{2} \;\Rightarrow$ $0 + e^{-m/1000} = \frac{1}{2} \;\Rightarrow\; -m/1000 = \ln\frac{1}{2} \;\Rightarrow\; m = -1000\ln\frac{1}{2} = 1000\ln 2 \approx 693.1$ h.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. (a) An exponential density function with $\mu = 1.6$ is $f(t) = \begin{cases} 0 & \text{if } t < 0 \\ \frac{1}{1.6}e^{-t/1.6} & \text{if } t \geq 0 \end{cases}$.

The probability that a customer waits less than a second is

$P(X < 1) = \int_0^1 f(t)\,dt = \int_0^1 \frac{1}{1.6}e^{-t/1.6}\,dt = \left[-e^{-t/1.6}\right]_0^1 = -e^{-1/1.6} + 1 \approx 0.465.$

(b) The probability that a customer waits more than 3 seconds is

$P(X > 3) = \int_3^\infty f(t)\,dt = \lim_{s\to\infty} \int_3^s f(t)\,dt = \lim_{s\to\infty} \left[-e^{-t/1.6}\right]_3^s = \lim_{s\to\infty} (-e^{-s/1.6} + e^{-3/1.6}) = e^{-3/1.6} \approx 0.153.$

Or: Calculate $1 - \int_0^3 f(t)\,dt$.

(c) We want to find b such that $P(X > b) = 0.05$. From part (b), $P(X > b) = e^{-b/1.6}$. Solving $e^{-b/1.6} = 0.05$ gives us

$-\frac{b}{1.6} = \ln 0.05 \quad \Rightarrow \quad b = -1.6 \ln 0.05 \approx 4.79$ seconds.

Or: Solve $\int_0^b f(t)\,dt = 0.95$ for b.

12. (a) We first find an antiderivative of $g(t) = t^2\,e^{at}$.

$$\begin{aligned}
\int t^2\,e^{at}\,dt &= \frac{1}{a}\,t^2\,e^{at} - \int \frac{2}{a}\,t\,e^{at}\,dt \qquad \begin{bmatrix} u = t^2, & dv = e^{at}\,dt \\ du = 2t\,dt, & v = \frac{1}{a}e^{at} \end{bmatrix} \\
&= \frac{1}{a}\,t^2\,e^{at} - \frac{2}{a}\left[\frac{1}{a}\,t\,e^{at} - \int \frac{1}{a}\,e^{at}\,dt\right] \qquad \begin{bmatrix} u = t, & dv = e^{at}\,dt \\ du = dt, & v = \frac{1}{a}e^{at} \end{bmatrix} \\
&= \frac{1}{a}\,t^2\,e^{at} - \frac{2}{a^2}\,t\,e^{at} + \frac{2}{a^3}\,e^{at} + C = \frac{1}{a}\,e^{at}\left(t^2 - \frac{2}{a}t + \frac{2}{a^2}\right) + C \\
&= -20e^{-0.05t}(t^2 + 40t + 800) + C \qquad [\text{with } a = -0.05]
\end{aligned}$$

$$\begin{aligned}
P(0 \leq X \leq 48) &= \int_0^{48} f(t)\,dt = \frac{1}{15{,}676}\int_0^{48} g(t)\,dt = \frac{1}{15{,}676}\left[-20e^{-0.05t}(t^2 + 40t + 800)\right]_0^{48} \\
&= \frac{-20}{15{,}676}(5024e^{-2.4} - 800) \approx 0.439.
\end{aligned}$$

(b) $$\begin{aligned}
P(X > 36) = P(36 < X \leq 150) &= \frac{1}{15{,}676}\int_{36}^{150} g(t)\,dt = \frac{1}{15{,}676}\left[-20e^{-0.05t}(t^2 + 40t + 800)\right]_{36}^{150} \\
&= \frac{-20}{15{,}676}(29{,}300e^{-7.5} - 3536e^{-1.8}) \approx 0.725
\end{aligned}$$

13. (a) $f(t) = \begin{cases} \frac{1}{1600}t & \text{if } 0 \leq t \leq 40 \\ \frac{1}{20} - \frac{1}{1600}t & \text{if } 40 < t \leq 80 \\ 0 & \text{otherwise} \end{cases}$

$$\begin{aligned}
P(30 \leq T \leq 60) &= \int_{30}^{60} f(t)\,dt = \int_{30}^{40} \frac{t}{1600}\,dt + \int_{40}^{60}\left(\frac{1}{20} - \frac{t}{1600}\right)dt = \left[\frac{t^2}{3200}\right]_{30}^{40} + \left[\frac{t}{20} - \frac{t^2}{3200}\right]_{40}^{60} \\
&= \left(\frac{1600}{3200} - \frac{900}{3200}\right) + \left(\frac{60}{20} - \frac{3600}{3200}\right) - \left(\frac{40}{20} - \frac{1600}{3200}\right) = -\frac{1300}{3200} + 1 = \frac{19}{32}
\end{aligned}$$

The probability that the amount of REM sleep is between 30 and 60 minutes is $\frac{19}{32} \approx 59.4\%$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\mu = \int_{-\infty}^{\infty} t\, f(t)\, dt = \int_0^{40} t\left(\frac{t}{1600}\right) dt + \int_{40}^{80} t\left(\frac{1}{20} - \frac{t}{1600}\right) dt = \left[\frac{t^3}{4800}\right]_0^{40} + \left[\frac{t^2}{40} - \frac{t^3}{4800}\right]_{40}^{80}$

$$= \frac{64{,}000}{4800} + \left(\frac{6400}{40} - \frac{512{,}000}{4800}\right) - \left(\frac{1600}{40} - \frac{64{,}000}{4800}\right) = -\frac{384{,}000}{4800} + 120 = 40$$

The mean amount of REM sleep is 40 minutes.

14. (a) With $\mu = 69$ and $\sigma = 2.8$, we have $P(65 \le X \le 73) = \int_{65}^{73} \frac{1}{2.8\sqrt{2\pi}} \exp\left(-\frac{(x-69)^2}{2 \cdot 2.8^2}\right) dx \approx 0.847$ (using a calculator or computer to estimate the integral).

(b) $P(X > 6 \text{ feet}) = P(X > 72 \text{ inches}) = 1 - P(0 \le X \le 72) \approx 1 - 0.858 = 0.142$, so 14.2% of the adult male population is more than 6 feet tall.

15. $P(X \ge 10) = \int_{10}^{\infty} \frac{1}{4.2\sqrt{2\pi}} \exp\left(-\frac{(x-9.4)^2}{2 \cdot 4.2^2}\right) dx$. To avoid the improper integral we approximate it by the integral from 10 to 100. Thus, $P(X \ge 10) \approx \int_{10}^{100} \frac{1}{4.2\sqrt{2\pi}} \exp\left(-\frac{(x-9.4)^2}{2 \cdot 4.2^2}\right) dx \approx 0.443$ (using a calculator or computer to estimate the integral), so about 44 percent of the households throw out at least 10 lb of paper a week.

Note: We can't evaluate $1 - P(0 \le X \le 10)$ for this problem since a significant amount of area lies to the left of $X = 0$.

16. (a) $P(0 \le X \le 480) = \int_0^{480} \frac{1}{12\sqrt{2\pi}} \exp\left(-\frac{(x-500)^2}{2 \cdot 12^2}\right) dx \approx 0.0478$ (using a calculator or computer to estimate the integral), so there is about a 4.78% chance that a particular box contains less than 480 g of cereal.

(b) We need to find μ so that $P(0 \le X < 500) = 0.05$. Using our calculator or computer to find $P(0 \le X \le 500)$ for various values of μ, we find that if $\mu = 519.73$, $P = 0.05007$; and if $\mu = 519.74$, $P = 0.04998$. So a good target weight is at least 519.74 g.

17. (a) $P(0 \le X \le 100) = \int_0^{100} \frac{1}{8\sqrt{2\pi}} \exp\left(-\frac{(x-112)^2}{2 \cdot 8^2}\right) dx \approx 0.0668$ (using a calculator or computer to estimate the integral), so there is about a 6.68% chance that a randomly chosen vehicle is traveling at a legal speed.

(b) $P(X \ge 125) = \int_{125}^{\infty} \frac{1}{8\sqrt{2\pi}} \exp\left(-\frac{(x-112)^2}{2 \cdot 8^2}\right) dx = \int_{125}^{\infty} f(x)\, dx$. In this case, we could use a calculator or computer to estimate either $\int_{125}^{300} f(x)\, dx$ or $1 - \int_0^{125} f(x)\, dx$. Both are approximately 0.0521, so about 5.21% of the motorists are targeted.

18. $f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)} \quad \Rightarrow \quad f'(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)} \frac{-2(x-\mu)}{2\sigma^2} = \frac{-1}{\sigma^3\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)}(x-\mu) \quad \Rightarrow$

$$f''(x) = \frac{-1}{\sigma^3\sqrt{2\pi}} \left[e^{-(x-\mu)^2/(2\sigma^2)} \cdot 1 + (x-\mu)e^{-(x-\mu)^2/(2\sigma^2)} \frac{-2(x-\mu)}{2\sigma^2}\right]$$

$$= \frac{-1}{\sigma^3\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)} \left[1 - \frac{(x-\mu)^2}{\sigma^2}\right] = \frac{1}{\sigma^5\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)} \left[(x-\mu)^2 - \sigma^2\right]$$

$f''(x) < 0 \quad \Rightarrow \quad (x-\mu)^2 - \sigma^2 < 0 \quad \Rightarrow \quad |x-\mu| < \sigma \quad \Rightarrow \quad -\sigma < x - \mu < \sigma \quad \Rightarrow \quad \mu - \sigma < x < \mu + \sigma$ and similarly, $f''(x) > 0 \quad \Rightarrow \quad x < \mu - \sigma$ or $x > \mu + \sigma$. Thus, f changes concavity and has inflection points at $x = \mu \pm \sigma$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $P(\mu - 2\sigma \le X \le \mu + 2\sigma) = \displaystyle\int_{\mu-2\sigma}^{\mu+2\sigma} \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) dx$. Substituting $t = \dfrac{x-\mu}{\sigma}$ and $dt = \dfrac{1}{\sigma}\,dx$ gives us

$$\int_{-2}^{2} \frac{1}{\sigma\sqrt{2\pi}} e^{-t^2/2}(\sigma\,dt) = \frac{1}{\sqrt{2\pi}} \int_{-2}^{2} e^{-t^2/2}\,dt \approx 0.9545.$$

20. Let $f(x) = \begin{cases} 0 & \text{if } x < 0 \\ ce^{-cx} & \text{if } x \ge 0 \end{cases}$ where $c = 1/\mu$. By using parts, tables, or a CAS, we find that

(1): $\int xe^{bx}\,dx = (e^{bx}/b^2)(bx-1)$

(2): $\int x^2e^{bx}\,dx = (e^{bx}/b^3)(b^2x^2 - 2bx + 2)$

Now
$$\begin{aligned}\sigma^2 &= \textstyle\int_{-\infty}^{\infty}(x-\mu)^2 f(x)\,dx = \int_{-\infty}^{0}(x-\mu)^2 f(x)\,dx + \int_0^{\infty}(x-\mu)^2 f(x)\,dx \\ &= 0 + \lim_{t\to\infty} c\textstyle\int_0^t (x-\mu)^2 e^{-cx}\,dx = c\cdot \lim_{t\to\infty}\int_0^t \left(x^2e^{-cx} - 2x\mu e^{-cx} + \mu^2 e^{-cx}\right) dx\end{aligned}$$

Next we use (2) and (1) with $b = -c$ to get

$$\sigma^2 = c\lim_{t\to\infty}\left[-\frac{e^{-cx}}{c^3}\left(c^2x^2 + 2cx + 2\right) - 2\mu\frac{e^{-cx}}{c^2}(-cx-1) + \mu^2\frac{e^{-cx}}{-c}\right]_0^t$$

Using l'Hospital's Rule several times, along with the fact that $\mu = 1/c$, we get

$$\sigma^2 = c\left[0 - \left(-\frac{2}{c^3} + \frac{2}{c}\cdot\frac{1}{c^2} + \frac{1}{c^2}\cdot\frac{1}{-c}\right)\right] = c\left(\frac{1}{c^3}\right) = \frac{1}{c^2} \quad\Rightarrow\quad \sigma = \frac{1}{c} = \mu$$

21. (a) First $p(r) = \dfrac{4}{a_0^3} r^2 e^{-2r/a_0} \ge 0$ for $r \ge 0$. Next,

$$\int_{-\infty}^{\infty} p(r)\,dr = \int_0^{\infty} \frac{4}{a_0^3} r^2 e^{-2r/a_0}\,dr = \frac{4}{a_0^3}\lim_{t\to\infty}\int_0^t r^2 e^{-2r/a_0}\,dr$$

By using parts, tables, or a CAS [or as in Exercise 20] , we find that $\int x^2e^{bx}\,dx = (e^{bx}/b^3)(b^2x^2 - 2bx + 2)$. $(\star)$

Next, we use $(\star)$ (with $b = -2/a_0$) and l'Hospital's Rule to get $\dfrac{4}{a_0^3}\left[\dfrac{a_0^3}{-8}(-2)\right] = 1$. This satisfies the second condition for a function to be a probability density function.

(b) Using l'Hospital's Rule, $\dfrac{4}{a_0^3}\displaystyle\lim_{r\to\infty}\frac{r^2}{e^{2r/a_0}} = \frac{4}{a_0^3}\lim_{r\to\infty}\frac{2r}{(2/a_0)e^{2r/a_0}} = \frac{2}{a_0^2}\lim_{r\to\infty}\frac{2}{(2/a_0)e^{2r/a_0}} = 0.$

To find the maximum of p, we differentiate:

$$p'(r) = \frac{4}{a_0^3}\left[r^2e^{-2r/a_0}\left(-\frac{2}{a_0}\right) + e^{-2r/a_0}(2r)\right] = \frac{4}{a_0^3}e^{-2r/a_0}(2r)\left(-\frac{r}{a_0} + 1\right)$$

$p'(r) = 0 \quad\Leftrightarrow\quad r = 0$ or $1 = \dfrac{r}{a_0} \quad\Leftrightarrow\quad r = a_0 \quad [a_0 \approx 5.59\times 10^{-11}$ m].

$p'(r)$ changes from positive to negative at $r = a_0$, so $p(r)$ has its maximum value at $r = a_0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) It is fairly difficult to find a viewing rectangle, but knowing the maximum value from part (b) helps.

$$p(a_0) = \frac{4}{a_0^3}\, a_0^2 e^{-2a_0/a_0} = \frac{4}{a_0}\, e^{-2} \approx 9{,}684{,}098{,}979$$

With a maximum of nearly 10 billion and a total area under the curve of 1, we know that the "hump" in the graph must be extremely narrow.

(d) $P(r) = \displaystyle\int_0^r \frac{4}{a_0^3} s^2 e^{-2s/a_0}\, ds \quad \Rightarrow \quad P(4a_0) = \int_0^{4a_0} \frac{4}{a_0^3} s^2 e^{-2s/a_0}\, ds$. Using ($\star$) from part (a) [with $b = -2/a_0$],

$$P(4a_0) = \frac{4}{a_0^3}\left[\frac{e^{-2s/a_0}}{-8/a_0^3}\left(\frac{4}{a_0^2}s^2 + \frac{4}{a_0}s + 2\right)\right]_0^{4a_0} = \frac{4}{a_0^3}\left(\frac{a_0^3}{-8}\right)[e^{-8}(64+16+2) - 1(2)] = -\tfrac{1}{2}(82e^{-8} - 2)$$
$$= 1 - 41e^{-8} \approx 0.986$$

(e) $\mu = \displaystyle\int_{-\infty}^{\infty} rp(r)\, dr = \frac{4}{a_0^3} \lim_{t\to\infty} \int_0^t r^3 e^{-2r/a_0}\, dr$. Integrating by parts three times or using a CAS, we find that

$\displaystyle\int x^3 e^{bx}\, dx = \frac{e^{bx}}{b^4}\left(b^3x^3 - 3b^2x^2 + 6bx - 6\right)$. So with $b = -\dfrac{2}{a_0}$, we use l'Hospital's Rule, and get

$\mu = \dfrac{4}{a_0^3}\left[-\dfrac{a_0^4}{16}(-6)\right] = \tfrac{3}{2}a_0$.

8 Review

TRUE-FALSE QUIZ

1. True. The graph of $y = f(x) + c$ is obtained by vertically translating $y = f(x)$ by c units. The arc length over the interval $a \le x \le b$ will be unchanged by this transformation.

2. False. Suppose $g(x) = 1$ and $h(x) = g(x) + 2 = 3$. Rotating the graph of $g(x)$ about the x-axis over the interval $[0, 2]$ will produce an open cylinder with radius 1 and height 2, so its surface area will be $2\pi(1)(2) = 4\pi$. Similarly, rotating $h(x)$ in the same way will generate an open cylinder with radius 3 and height 2, so its surface area will be $2\pi(3)(2) = 12\pi$.

3. False. Suppose $f(x) = x$ and $g(x) = 1$ so that $f(x) \le g(x)$ in the interval $[0, 1]$. $f(x)$ is a straight line so its arc length over $[0, 1]$ is the distance between its endpoints and is given by $\sqrt{(1-0)^2 + [f(1) - f(0)]^2} = \sqrt{2}$. Similarly, the arc length of $g(x)$ over $[0, 1]$ is 1, which is less than $\sqrt{2}$.

4. False. $y = x^3 \quad \Rightarrow \quad dy/dx = 3x^2 \quad \Rightarrow \quad L = \displaystyle\int_0^1 \sqrt{1 + (dy/dx)^2}\, dx = \int_0^1 \sqrt{1 + (3x^2)^2}\, dx = \int_0^1 \sqrt{1 + 9x^4}\, dx$

5. True. The smallest possible length of arc between the points $(0, 0)$ and $(3, 4)$ is the length of a straight line segment connecting the two points. This length is $\sqrt{(3-0)^2 + (4-0)^2} = \sqrt{25} = 5$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. True. By Equation 8.3.9, the centroid of a lamina depends only on its area and the curves $y = f(x)$ and $y = g(x)$, which define its shape.

7. True. The hydrostatic pressure depends only on the depth of the fluid, d, and the fluid's weight density, δ, as given by $P = \delta d$. See margin note next to Equation 8.3.1.

8. True. The total probability must be 1. See Equation 8.5.2 and the preceding discussion.

EXERCISES

1. $y = 4(x-1)^{3/2} \quad \Rightarrow \quad \dfrac{dy}{dx} = 6(x-1)^{1/2} \quad \Rightarrow \quad 1 + \left(\dfrac{dy}{dx}\right)^2 = 1 + 36(x-1) = 36x - 35$. Thus,

$$L = \int_1^4 \sqrt{36x - 35}\,dx = \int_1^{109} \sqrt{u}\,(\frac{1}{36}\,du) \qquad \begin{bmatrix} u = 36x - 35, \\ du = 36\,dx \end{bmatrix}$$

$$= \tfrac{1}{36}\left[\tfrac{2}{3}u^{3/2}\right]_1^{109} = \tfrac{1}{54}(109\sqrt{109} - 1)$$

2. $y = 2\ln\left(\sin \tfrac{1}{2}x\right) \quad \Rightarrow \quad \dfrac{dy}{dx} = 2 \cdot \dfrac{1}{\sin\left(\frac{1}{2}x\right)} \cdot \cos\left(\tfrac{1}{2}x\right) \cdot \tfrac{1}{2} = \cot\left(\tfrac{1}{2}x\right) \quad \Rightarrow \quad 1 + \left(\dfrac{dy}{dx}\right)^2 = 1 + \cot^2\left(\tfrac{1}{2}x\right) = \csc^2\left(\tfrac{1}{2}x\right)$.

Thus,

$$L = \int_{\pi/3}^{\pi} \sqrt{\csc^2\left(\tfrac{1}{2}x\right)}\,dx = \int_{\pi/3}^{\pi} \left|\csc\left(\tfrac{1}{2}x\right)\right|\,dx = \int_{\pi/3}^{\pi} \csc\left(\tfrac{1}{2}x\right)\,dx = \int_{\pi/6}^{\pi/2} \csc u\,(2\,du) \qquad \begin{bmatrix} u = \frac{1}{2}x, \\ du = \frac{1}{2}\,dx \end{bmatrix}$$

$$= 2\Big[\ln|\csc u - \cot u|\Big]_{\pi/6}^{\pi/2} = 2\left[\ln\left|\csc\tfrac{\pi}{2} - \cot\tfrac{\pi}{2}\right| - \ln\left|\csc\tfrac{\pi}{6} - \cot\tfrac{\pi}{6}\right|\right]$$

$$= 2\Big[\ln|1 - 0| - \ln\left|2 - \sqrt{3}\right|\Big] = -2\ln\left(2 - \sqrt{3}\right) \approx 2.63$$

3. $12x = 4y^3 + 3y^{-1} \quad \Rightarrow \quad x = \tfrac{1}{3}y^3 + \tfrac{1}{4}y^{-1} \quad \Rightarrow \quad \dfrac{dx}{dy} = y^2 - \tfrac{1}{4}y^{-2} \quad \Rightarrow$

$1 + \left(\dfrac{dx}{dy}\right)^2 = 1 + y^4 - \tfrac{1}{2} + \tfrac{1}{16}y^{-4} = y^4 + \tfrac{1}{2} + \tfrac{1}{16}y^{-4} = (y^2 + \tfrac{1}{4}y^{-2})^2$. Thus,

$$L = \int_1^3 \sqrt{\left(y^2 + \tfrac{1}{4}y^{-2}\right)^2}\,dy = \int_1^3 \left|y^2 + \tfrac{1}{4}y^{-2}\right|\,dy = \int_1^3 \left(y^2 + \tfrac{1}{4}y^{-2}\right)\,dy = \left[\tfrac{1}{3}y^3 - \tfrac{1}{4}y^{-1}\right]_1^3$$

$$= (9 - \tfrac{1}{12}) - (\tfrac{1}{3} - \tfrac{1}{4}) = \tfrac{106}{12} = \tfrac{53}{6}$$

4. (a) $y = \dfrac{x^4}{16} + \dfrac{1}{2x^2} = \dfrac{1}{16}x^4 + \dfrac{1}{2}x^{-2} \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{1}{4}x^3 - x^{-3} \quad \Rightarrow$

$1 + (dy/dx)^2 = 1 + \left(\tfrac{1}{4}x^3 - x^{-3}\right)^2 = 1 + \tfrac{1}{16}x^6 - \tfrac{1}{2} + x^{-6} = \tfrac{1}{16}x^6 + \tfrac{1}{2} + x^{-6} = \left(\tfrac{1}{4}x^3 + x^{-3}\right)^2$.

Thus, $L = \int_1^2 \left(\tfrac{1}{4}x^3 + x^{-3}\right)\,dx = \left[\tfrac{1}{16}x^4 - \tfrac{1}{2}x^{-2}\right]_1^2 = \left(1 - \tfrac{1}{8}\right) - \left(\tfrac{1}{16} - \tfrac{1}{2}\right) = \tfrac{21}{16}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $S = \int_1^2 2\pi x\,ds = \int_1^2 2\pi x\sqrt{1+(dy/dx)^2}\,dx = \int_1^2 2\pi x\left(\frac{1}{4}x^3 + x^{-3}\right)dx = 2\pi\int_1^2\left(\frac{1}{4}x^4 + x^{-2}\right)dx$

$= 2\pi\left[\frac{1}{20}x^5 - \frac{1}{x}\right]_1^2 = 2\pi\left[\left(\frac{32}{20} - \frac{1}{2}\right) - \left(\frac{1}{20} - 1\right)\right] = 2\pi\left(\frac{8}{5} - \frac{1}{2} - \frac{1}{20} + 1\right) = 2\pi\left(\frac{41}{20}\right) = \frac{41}{10}\pi$

5. (a) $y = \dfrac{2}{x+1} \quad\Rightarrow\quad y' = \dfrac{-2}{(x+1)^2} \quad\Rightarrow\quad 1 + (y')^2 = 1 + \dfrac{4}{(x+1)^4}.$

For $0 \le x \le 3$, $L = \int_0^3 \sqrt{1+(y')^2}\,dx = \int_0^3 \sqrt{1+4/(x+1)^4}\,dx \approx 3.5121.$

(b) The area of the surface obtained by rotating C about the x-axis is

$$S = \int_0^3 2\pi y\,ds = 2\pi\int_0^3 \frac{2}{x+1}\sqrt{1+4/(x+1)^4}\,dx \approx 22.1391.$$

(c) The area of the surface obtained by rotating C about the y-axis is

$S = \int_0^3 2\pi x\,ds = 2\pi\int_0^3 x\sqrt{1+4/(x+1)^4}\,dx \approx 29.8522.$

6. (a) $y = x^2 \quad\Rightarrow\quad 1 + (y')^2 = 1 + 4x^2$. Rotate about the y-axis for $0 \le x \le 1$:

$$S = \int_0^1 2\pi x\sqrt{1+4x^2}\,dx = \int_1^5 \frac{\pi}{4}\sqrt{u}\,du \quad [u = 1+4x^2] \;=\; \frac{\pi}{6}\left[u^{3/2}\right]_1^5 = \frac{\pi}{6}(5^{3/2} - 1)$$

(b) $y = x^2 \quad\Rightarrow\quad 1 + (y')^2 = 1 + 4x^2$. Rotate about the x-axis for $0 \le x \le 1$:

$$S = 2\pi\int_0^1 x^2\sqrt{1+4x^2}\,dx = 2\pi\int_0^2 \frac{1}{4}u^2\sqrt{1+u^2}\,\frac{1}{2}\,du \quad [u = 2x] \quad = \frac{\pi}{4}\int_0^2 u^2\sqrt{1+u^2}\,du$$

$$= \frac{\pi}{4}\left[\frac{1}{8}u(1+2u^2)\sqrt{1+u^2} - \frac{1}{8}\ln\left|u + \sqrt{1+u^2}\,\right|\right]_0^2 \qquad [u = \tan\theta \text{ or use Formula 22}]$$

$$= \frac{\pi}{4}\left[\frac{1}{4}(9)\sqrt{5} - \frac{1}{8}\ln\left(2+\sqrt{5}\right) - 0\right] = \frac{\pi}{32}\left[18\sqrt{5} - \ln\left(2+\sqrt{5}\right)\right]$$

7. $y = \sin x \quad\Rightarrow\quad y' = \cos x \quad\Rightarrow\quad 1 + (y')^2 = 1 + \cos^2 x$. Let $f(x) = \sqrt{1+\cos^2 x}$. Then

$$L = \int_0^\pi f(x)\,dx \approx S_{10}$$

$$= \frac{(\pi - 0)/10}{3}\left[f(0) + 4f\left(\frac{\pi}{10}\right) + 2f\left(\frac{2\pi}{10}\right) + 4f\left(\frac{3\pi}{10}\right) + 2f\left(\frac{4\pi}{10}\right)\right.$$

$$\left. + 4f\left(\frac{5\pi}{10}\right) + 2f\left(\frac{6\pi}{10}\right) + 4f\left(\frac{7\pi}{10}\right) + 2f\left(\frac{8\pi}{10}\right) + 4f\left(\frac{9\pi}{10}\right) + f(\pi)\right]$$

$$\approx 3.8202$$

8. (a) $y = \sin x \quad\Rightarrow\quad y' = \cos x \quad\Rightarrow\quad 1 + (y')^2 = 1 + \cos^2 x.$

$S = \int_0^\pi 2\pi y\,ds = \int_0^\pi 2\pi\sin x\sqrt{1+\cos^2 x}\,dx.$

(b) $S \approx 14.4260$

9. $y = \int_1^x \sqrt{\sqrt{t} - 1}\,dt \quad\Rightarrow\quad dy/dx = \sqrt{\sqrt{x} - 1} \quad\Rightarrow\quad 1 + (dy/dx)^2 = 1 + \left(\sqrt{x} - 1\right) = \sqrt{x}.$

Thus, $L = \int_1^{16} \sqrt{\sqrt{x}}\,dx = \int_1^{16} x^{1/4}\,dx = \frac{4}{5}\left[x^{5/4}\right]_1^{16} = \frac{4}{5}(32 - 1) = \frac{124}{5}.$

10. $S = \int_1^{16} 2\pi x\,ds = 2\pi\int_1^{16} x\cdot x^{1/4}\,dx = 2\pi\int_1^{16} x^{5/4}\,dx = 2\pi\cdot\frac{4}{9}\left[x^{9/4}\right]_1^{16} = \frac{8\pi}{9}(512 - 1) = \frac{4088}{9}\pi$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. By reasoning as in Example 8.3.1, $\dfrac{a}{2-x} = \dfrac{1}{2} \quad \Rightarrow \quad 2a = 2 - x$ and $w = 2(1.5 + a) = 3 + 2a = 3 + 2 - x = 5 - x$.

Thus, $F = \int_0^2 \delta x(5-x)\,dx = \delta\left[\frac{5}{2}x^2 - \frac{1}{3}x^3\right]_0^2 = \delta\left(10 - \frac{8}{3}\right) = \frac{22}{3}\delta \approx 458$ lb $[\delta \approx 62.5\ \text{lb/ft}^3]$.

12. The parabola has equation $y = ax^2$. Since the point $(4, 4)$ is on the graph, $4 = a(4)^2 \quad \Rightarrow \quad a = \frac{1}{4}$. Thus, an equation is $y = \frac{1}{4}x^2$ or $x = \pm 2\sqrt{y}$.

$$F = \int_0^4 \delta(4-y)2\left(2\sqrt{y}\right)dy = 4\delta\int_0^4 (4y^{1/2} - y^{3/2})\,dy$$
$$= 4\delta\left[\tfrac{8}{3}y^{3/2} - \tfrac{2}{5}y^{5/2}\right]_0^4 = 4\delta\left(\tfrac{64}{3} - \tfrac{64}{5}\right) = 256\delta\left(\tfrac{1}{3} - \tfrac{1}{5}\right)$$
$$= \tfrac{512}{15}\delta \approx 2133.3\ \text{lb} \qquad [\delta \approx 62.5\ \text{lb/ft}^3]$$

13. The area of the triangular region is $A = \frac{1}{2}(2)(4) = 4$. An equation of the line is $y = \frac{1}{2}x$ or $x = 2y$.

$$\overline{x} = \frac{1}{A}\int_0^2 \frac{1}{2}[f(y)]^2\,dy = \frac{1}{4}\int_0^2 \frac{1}{2}(2y)^2\,dy = \frac{1}{8}\int_0^2 4y^2\,dy = \frac{1}{8}\left[\frac{4}{3}y^3\right]_0^2 = \frac{1}{6}(8) = \frac{4}{3}$$

$$\overline{y} = \frac{1}{A}\int_0^2 y\,f(y)\,dy = \frac{1}{4}\int_0^2 y(2y)\,dy = \frac{1}{2}\int_0^2 y^2\,dy = \frac{1}{2}\left[\frac{1}{3}y^3\right]_0^2 = \frac{1}{6}(8) = \frac{4}{3}$$

The centroid of the region is $\left(\frac{4}{3}, \frac{4}{3}\right)$.

14. An equation of the line is $y = 8 - x$. An equation of the quarter-circle is $y = -\sqrt{8^2 - x^2}$ with $0 \le x \le 8$. The area of the region is $A = \frac{1}{2}(8)(8) + \frac{1}{4}\pi(8)^2 = 32 + 16\pi = 16(2+\pi)$.

$$\overline{x} = \frac{1}{A}\int_0^8 x[f(x) - g(x)]\,dx = \frac{1}{A}\int_0^8 x\left[(8-x) + \sqrt{64 - x^2}\right]dx$$
$$= \frac{1}{A}\int_0^8 \left[8x - x^2 + x(64 - x^2)^{1/2}\right]dx = \frac{1}{A}\left[4x^2 - \frac{1}{3}x^3 - \frac{1}{3}(64 - x^2)^{3/2}\right]_0^8$$
$$= \frac{1}{A}\left[\left(256 - \frac{512}{3} - 0\right) - \left(0 - 0 - \frac{512}{3}\right)\right] = \frac{256}{16(2+\pi)} = \frac{16}{2+\pi}$$

$$\overline{y} = \frac{1}{A}\int_0^8 \frac{1}{2}\{[f(x)]^2 - [g(x)]^2\}\,dx = \frac{1}{2A}\int_0^8 \left[(8-x)^2 - (-\sqrt{64 - x^2}\,)^2\right]dx$$
$$= \frac{1}{2A}\int_0^8 \left[64 - 16x + x^2 - (64 - x^2)\right]dx = \frac{1}{2A}\int_0^8 (2x^2 - 16x)\,dx$$
$$= \frac{1}{A}\int_0^8 (x^2 - 8x)\,dx = \frac{1}{A}\left[\frac{1}{3}x^3 - 4x^2\right]_0^8 = \frac{1}{A}\left(\frac{512}{3} - 256\right)$$
$$= \frac{1}{16(2+\pi)}\left(-\frac{256}{3}\right) = -\frac{16}{3(2+\pi)}$$

The centroid of the region is $\left(\dfrac{16}{2+\pi}, -\dfrac{16}{3(2+\pi)}\right) \approx (3.11, -1.04)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $A = \int_0^4 \left(\sqrt{x} - \frac{1}{2}x\right) dx = \left[\frac{2}{3}x^{3/2} - \frac{1}{4}x^2\right]_0^4 = \frac{16}{3} - 4 = \frac{4}{3}$

$\overline{x} = \frac{1}{A}\int_0^4 x\left(\sqrt{x} - \frac{1}{2}x\right) dx = \frac{3}{4}\int_0^4 \left(x^{3/2} - \frac{1}{2}x^2\right) dx$

$= \frac{3}{4}\left[\frac{2}{5}x^{5/2} - \frac{1}{6}x^3\right]_0^4 = \frac{3}{4}\left(\frac{64}{5} - \frac{64}{6}\right) = \frac{3}{4}\left(\frac{64}{30}\right) = \frac{8}{5}$

$\overline{y} = \frac{1}{A}\int_0^4 \frac{1}{2}\left[\left(\sqrt{x}\right)^2 - \left(\frac{1}{2}x\right)^2\right] dx = \frac{3}{4}\int_0^4 \frac{1}{2}\left(x - \frac{1}{4}x^2\right) dx = \frac{3}{8}\left[\frac{1}{2}x^2 - \frac{1}{12}x^3\right]_0^4 = \frac{3}{8}\left(8 - \frac{16}{3}\right) = \frac{3}{8}\left(\frac{8}{3}\right) = 1$

Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\frac{8}{5}, 1\right)$.

16. From the symmetry of the region, $\overline{x} = \frac{\pi}{2}$. $A = \int_{\pi/4}^{3\pi/4} \sin x\, dx = \left[-\cos x\right]_{\pi/4}^{3\pi/4} = \frac{1}{\sqrt{2}} - \left(-\frac{1}{\sqrt{2}}\right) = \sqrt{2}$

$$\overline{y} = \frac{1}{A}\int_{\pi/4}^{3\pi/4} \frac{1}{2}\sin^2 x\, dx = \frac{1}{A}\int_{\pi/4}^{3\pi/4} \frac{1}{4}\left(1 - \cos 2x\right) dx$$

$$= \frac{1}{4\sqrt{2}}\left[x - \frac{1}{2}\sin 2x\right]_{\pi/4}^{3\pi/4}$$

$$= \frac{1}{4\sqrt{2}}\left[\frac{3\pi}{4} - \frac{1}{2}(-1) - \frac{\pi}{4} + \frac{1}{2}\cdot 1\right] = \frac{1}{4\sqrt{2}}\left(\frac{\pi}{2} + 1\right)$$

Thus, the centroid is $(\overline{x}, \overline{y}) = \left(\frac{\pi}{2}, \frac{1}{4\sqrt{2}}\left(\frac{\pi}{2} + 1\right)\right) \approx (1.57, 0.45)$.

17. The centroid of this circle, $(1, 0)$, travels a distance $2\pi(1)$ when the lamina is rotated about the y-axis. The area of the circle is $\pi(1)^2$. So by the Theorem of Pappus, $V = A(2\pi\overline{x}) = \pi(1)^2 2\pi(1) = 2\pi^2$.

18. The semicircular region has an area of $\frac{1}{2}\pi r^2$, and sweeps out a sphere of radius r when rotated about the x-axis. $\overline{x} = 0$ because of symmetry about the line $x = 0$. And by the Theorem of Pappus, $V = A(2\pi\overline{y}) \quad \Rightarrow$ $\frac{4}{3}\pi r^3 = \frac{1}{2}\pi r^2(2\pi\overline{y}) \quad \Rightarrow \quad \overline{y} = \frac{4}{3\pi}r$. Thus, the centroid is $(\overline{x}, \overline{y}) = \left(0, \frac{4}{3\pi}r\right)$.

19. $x = 100 \quad \Rightarrow \quad P = 2000 - 0.1(100) - 0.01(100)^2 = 1890$

$$\text{Consumer surplus} = \int_0^{100}[p(x) - P]\, dx = \int_0^{100}\left(2000 - 0.1x - 0.01x^2 - 1890\right) dx$$

$$= \left[110x - 0.05x^2 - \frac{0.01}{3}x^3\right]_0^{100} = 11{,}000 - 500 - \frac{10{,}000}{3} \approx \$7166.67$$

20. $\int_0^{24} c(t)\, dt \approx S_{12} = \frac{24-0}{12\cdot 3}[1(0) + 4(1.9) + 2(3.3) + 4(5.1) + 2(7.6) + 4(7.1) + 2(5.8)$
$+ 4(4.7) + 2(3.3) + 4(2.1) + 2(1.1) + 4(0.5) + 1(0)]$

$= \frac{2}{3}(127.8) = 85.2 \text{ mg}\cdot\text{s/L}$

Therefore, $F \approx A/85.2 = 6/85.2 \approx 0.0704$ L/s or 4.225 L/min.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. $f(x) = \begin{cases} \frac{\pi}{20}\sin\left(\frac{\pi}{10}x\right) & \text{if } 0 \le x \le 10 \\ 0 & \text{if } x < 0 \text{ or } x > 10 \end{cases}$

(a) $f(x) \ge 0$ for all real numbers x and

$$\int_{-\infty}^{\infty} f(x)\,dx = \int_0^{10} \tfrac{\pi}{20}\sin\left(\tfrac{\pi}{10}x\right)dx = \tfrac{\pi}{20}\cdot\tfrac{10}{\pi}\left[-\cos\left(\tfrac{\pi}{10}x\right)\right]_0^{10} = \tfrac{1}{2}(-\cos\pi + \cos 0) = \tfrac{1}{2}(1+1) = 1$$

Therefore, f is a probability density function.

(b) $P(X < 4) = \int_{-\infty}^{4} f(x)\,dx = \int_0^4 \frac{\pi}{20}\sin\left(\frac{\pi}{10}x\right)dx = \frac{1}{2}\left[-\cos\left(\frac{\pi}{10}x\right)\right]_0^4 = \frac{1}{2}\left(-\cos\frac{2\pi}{5} + \cos 0\right)$

$\approx \frac{1}{2}(-0.309017 + 1) \approx 0.3455$

(c) $\mu = \int_{-\infty}^{\infty} xf(x)\,dx = \int_0^{10} \frac{\pi}{20}x\sin\left(\frac{\pi}{10}x\right)dx$

$= \int_0^{\pi} \frac{\pi}{20}\cdot\frac{10}{\pi}u(\sin u)\left(\frac{10}{\pi}\right)du \qquad \left[u = \frac{\pi}{10}x,\ du = \frac{\pi}{10}\,dx\right]$

$= \frac{5}{\pi}\int_0^{\pi} u\sin u\,du \overset{82}{=} \frac{5}{\pi}\left[\sin u - u\cos u\right]_0^{\pi} = \frac{5}{\pi}[0 - \pi(-1)] = 5$

This answer is expected because the graph of f is symmetric about the line $x = 5$.

22. $P(250 \le X \le 280) = \displaystyle\int_{250}^{280} \frac{1}{\sigma\sqrt{2\pi}}\,e^{-(x-\mu)^2/(2\sigma^2)}\,dx = \int_{250}^{280} \frac{1}{15\sqrt{2\pi}}\exp\left(\frac{-(x-268)^2}{2\cdot 15^2}\right)dx \approx 0.673.$

Thus, the percentage of pregnancies that last between 250 and 280 days is about 67.3%.

23. (a) The probability density function is $f(t) = \begin{cases} 0 & \text{if } t < 0 \\ \frac{1}{8}e^{-t/8} & \text{if } t \ge 0 \end{cases}$

$$P(0 \le X \le 3) = \int_0^3 \tfrac{1}{8}e^{-t/8}\,dt = \left[-e^{-t/8}\right]_0^3 = -e^{-3/8} + 1 \approx 0.3127$$

(b) $P(X > 10) = \int_{10}^{\infty} \frac{1}{8}e^{-t/8}\,dt = \lim\limits_{x\to\infty}\left[-e^{-t/8}\right]_{10}^{x} = \lim\limits_{x\to\infty}(-e^{-x/8} + e^{-10/8}) = 0 + e^{-5/4} \approx 0.2865$

(c) We need to find m such that $P(X \ge m) = \frac{1}{2} \;\Rightarrow\; \int_m^{\infty} \frac{1}{8}e^{-t/8}\,dt = \frac{1}{2} \;\Rightarrow\; \lim\limits_{x\to\infty}\left[-e^{-t/8}\right]_m^x = \frac{1}{2} \;\Rightarrow$

$\lim\limits_{x\to\infty}(-e^{-x/8} + e^{-m/8}) = \frac{1}{2} \;\Rightarrow\; e^{-m/8} = \frac{1}{2} \;\Rightarrow\; -m/8 = \ln\frac{1}{2} \;\Rightarrow\; m = -8\ln\frac{1}{2} = 8\ln 2 \approx 5.55$ minutes.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

PROBLEMS PLUS

1. $x^2+y^2 \le 4y \quad \Leftrightarrow \quad x^2+(y-2)^2 \le 4$, so S is part of a circle, as shown in the diagram. The area of S is

$$\int_0^1 \sqrt{4y-y^2}\,dy \overset{113}{=} \left[\tfrac{y-2}{2}\sqrt{4y-y^2}+2\cos^{-1}\left(\tfrac{2-y}{2}\right)\right]_0^1 \quad [a=2]$$
$$= -\tfrac{1}{2}\sqrt{3}+2\cos^{-1}\left(\tfrac{1}{2}\right)-2\cos^{-1}1$$
$$= -\tfrac{\sqrt{3}}{2}+2\left(\tfrac{\pi}{3}\right)-2(0)=\tfrac{2\pi}{3}-\tfrac{\sqrt{3}}{2}$$

Another method (without calculus): Note that $\theta=\angle CAB=\frac{\pi}{3}$, so the area is

$$(\text{area of sector } OAB)-(\text{area of } \triangle ABC)=\tfrac{1}{2}\left(2^2\right)\tfrac{\pi}{3}-\tfrac{1}{2}(1)\sqrt{3}=\tfrac{2\pi}{3}-\tfrac{\sqrt{3}}{2}$$

2. $y=\pm\sqrt{x^3-x^4} \quad \Rightarrow$ The loop of the curve is symmetric about $y=0$, and therefore $\overline{y}=0$. At each point x where $0\le x\le 1$, the lamina has a vertical length of $\sqrt{x^3-x^4}-\left(-\sqrt{x^3-x^4}\right)=2\sqrt{x^3-x^4}$. Therefore,

$\overline{x}=\dfrac{\int_0^1 x\cdot 2\sqrt{x^3-x^4}\,dx}{\int_0^1 2\sqrt{x^3-x^4}\,dx}=\dfrac{\int_0^1 x\sqrt{x^3-x^4}\,dx}{\int_0^1 \sqrt{x^3-x^4}\,dx}$. We evaluate the integrals separately:

$$\int_0^1 x\sqrt{x^3-x^4}\,dx=\int_0^1 x^{5/2}\sqrt{1-x}\,dx$$
$$=\int_0^{\pi/2}2\sin^6\theta\cos\theta\sqrt{1-\sin^2\theta}\,d\theta \qquad \left[\begin{matrix}\sin\theta=\sqrt{x},\ \cos\theta\,d\theta=dx/(2\sqrt{x}),\\ 2\sin\theta\,\cos\theta\,d\theta=dx\end{matrix}\right]$$
$$=\int_0^{\pi/2}2\sin^6\theta\cos^2\theta\,d\theta=\int_0^{\pi/2}2\left[\tfrac{1}{2}(1-\cos 2\theta)\right]^3\tfrac{1}{2}(1+\cos 2\theta)d\theta$$
$$=\int_0^{\pi/2}\tfrac{1}{8}(1-2\cos 2\theta+2\cos^3 2\theta-\cos^4 2\theta)\,d\theta$$
$$=\int_0^{\pi/2}\tfrac{1}{8}\left[1-2\cos 2\theta+2\cos 2\theta(1-\sin^2 2\theta)-\tfrac{1}{4}(1+\cos 4\theta)^2\right]d\theta$$
$$=\tfrac{1}{8}\left[\theta-\tfrac{1}{3}\sin^3 2\theta\right]_0^{\pi/2}-\tfrac{1}{32}\int_0^{\pi/2}\left(1+2\cos 4\theta+\cos^2 4\theta\right)d\theta$$
$$=\tfrac{\pi}{16}-\tfrac{1}{32}\left[\theta+\tfrac{1}{2}\sin 4\theta\right]_0^{\pi/2}-\tfrac{1}{64}\int_0^{\pi/2}(1+\cos 8\theta)\,d\theta$$
$$=\tfrac{\pi}{16}-\tfrac{1}{32}\left[\theta+\tfrac{1}{2}\sin 4\theta\right]_0^{\pi/2}-\tfrac{1}{64}\int_0^{\pi/2}(1+\cos 8\theta)\,d\theta$$
$$=\tfrac{3\pi}{64}-\tfrac{1}{64}\left[\theta+\tfrac{1}{8}\sin 8\theta\right]_0^{\pi/2}=\tfrac{5\pi}{128}$$

$$\int_0^1\sqrt{x^3-x^4}\,dx=\int_0^1 x^{3/2}\sqrt{1-x}\,dx=\int_0^{\pi/2}2\sin^4\theta\cos\theta\sqrt{1-\sin^2\theta}\,d\theta \qquad [\sin\theta=\sqrt{x}\,]$$
$$=\int_0^{\pi/2}2\sin^4\theta\cos^2\theta\,d\theta=\int_0^{\pi/2}2\cdot\tfrac{1}{4}(1-\cos 2\theta)^2\cdot\tfrac{1}{2}(1+\cos 2\theta)\,d\theta$$
$$=\int_0^{\pi/2}\tfrac{1}{4}(1-\cos 2\theta-\cos^2 2\theta+\cos^3 2\theta)\,d\theta$$
$$=\int_0^{\pi/2}\tfrac{1}{4}\left[1-\cos 2\theta-\tfrac{1}{2}(1+\cos 4\theta)+\cos 2\theta(1-\sin^2 2\theta)\right]d\theta$$
$$=\tfrac{1}{4}\left[\tfrac{\theta}{2}-\tfrac{1}{8}\sin 4\theta-\tfrac{1}{6}\sin^3 2\theta\right]_0^{\pi/2}=\tfrac{\pi}{16}$$

Therefore, $\overline{x}=\dfrac{5\pi/128}{\pi/16}=\dfrac{5}{8}$, and $(\overline{x},\overline{y})=\left(\frac{5}{8},0\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a) The two spherical zones, whose surface areas we will call S_1 and S_2, are generated by rotation about the y-axis of circular arcs, as indicated in the figure. The arcs are the upper and lower portions of the circle $x^2 + y^2 = r^2$ that are obtained when the circle is cut with the line $y = d$. The portion of the upper arc in the first quadrant is sufficient to generate the upper spherical zone. That portion of the arc can be described by the relation $x = \sqrt{r^2 - y^2}$ for $d \le y \le r$. Thus, $dx/dy = -y/\sqrt{r^2 - y^2}$ and

$$ds = \sqrt{1 + \left(\frac{dx}{dy}\right)^2}\, dy = \sqrt{1 + \frac{y^2}{r^2 - y^2}}\, dy = \sqrt{\frac{r^2}{r^2 - y^2}}\, dy = \frac{r\, dy}{\sqrt{r^2 - y^2}}$$

From Formula 8.2.8 we have

$$S_1 = \int_d^r 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2}\, dy = \int_d^r 2\pi \sqrt{r^2 - y^2} \frac{r\, dy}{\sqrt{r^2 - y^2}} = \int_d^r 2\pi r\, dy = 2\pi r(r - d)$$

Similarly, we can compute $S_2 = \int_{-r}^{d} 2\pi x \sqrt{1 + (dx/dy)^2}\, dy = \int_{-r}^{d} 2\pi r\, dy = 2\pi r(r + d)$. Note that $S_1 + S_2 = 4\pi r^2$, the surface area of the entire sphere.

(b) $r = 3960$ mi and $d = r\,(\sin 75^\circ) \approx 3825$ mi, so the surface area of the Arctic Ocean is about $2\pi r(r - d) \approx 2\pi(3960)(135) \approx 3.36 \times 10^6$ mi^2.

(c) The area on the sphere lies between planes $y = y_1$ and $y = y_2$, where $y_2 - y_1 = h$. Thus, we compute the surface area on the sphere to be $S = \displaystyle\int_{y_1}^{y_2} 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2}\, dy = \int_{y_1}^{y_2} 2\pi r\, dy = 2\pi r(y_2 - y_1) = 2\pi r h.$

This equals the lateral area of a cylinder of radius r and height h, since such a cylinder is obtained by rotating the line $x = r$ about the y-axis, so the surface area of the cylinder between the planes $y = y_1$ and $y = y_2$ is

$$A = \int_{y_1}^{y_2} 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2}\, dy = \int_{y_1}^{y_2} 2\pi r \sqrt{1 + 0^2}\, dy$$
$$= 2\pi r y \Big|_{y=y_1}^{y_2} = 2\pi r(y_2 - y_1) = 2\pi r h$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $h = 2r \sin 23.45° \approx 3152$ mi, so the surface area of the Torrid Zone is $2\pi rh \approx 2\pi(3960)(3152) \approx 7.84 \times 10^7$ mi^2.

4. (a) Since the right triangles OAT and OTB are similar, we have $\dfrac{r+H}{r} = \dfrac{r}{a} \quad \Rightarrow$

$a = \dfrac{r^2}{r+H}$. The surface area visible from B is $S = \int_a^r 2\pi x \sqrt{1 + (dx/dy)^2}\, dy$.

From $x^2 + y^2 = r^2$, we get $\dfrac{d}{dy}(x^2+y^2) = \dfrac{d}{dy}(r^2) \quad \Rightarrow \quad 2x\dfrac{dx}{dy} + 2y = 0 \quad \Rightarrow$

$\dfrac{dx}{dy} = -\dfrac{y}{x}$ and $1 + \left(\dfrac{dx}{dy}\right)^2 = \dfrac{x^2+y^2}{x^2} = \dfrac{r^2}{x^2}$. Thus,

$$S = \int_a^r 2\pi x \cdot \frac{r}{x}\, dy = 2\pi r(r-a) = 2\pi r\left(r - \frac{r^2}{r+H}\right) = 2\pi r^2\left(1 - \frac{r}{r+H}\right) = 2\pi r^2 \cdot \frac{H}{r+H} = \frac{2\pi r^2 H}{r+H}.$$

(b) Assume $R \geq r$. If a light is placed at point L, at a distance x from the center of the sphere of radius r, then from part (a) we find that the total illuminated area A on the two spheres is [with $r + H = x$ and $r + H = d - x$].

$$A(x) = \frac{2\pi r^2(x-r)}{x} + \frac{2\pi R^2(d-x-R)}{d-x} \quad [r \leq x \leq d-R]. \quad \frac{A(x)}{2\pi} = r^2\left(1 - \frac{r}{x}\right) + R^2\left(1 - \frac{R}{d-x}\right),$$

so $A'(x) = 0 \quad \Leftrightarrow \quad 0 = r^2 \cdot \dfrac{r}{x^2} + R^2 \cdot \dfrac{-R}{(d-x)^2} \quad \Leftrightarrow \quad \dfrac{r^3}{x^2} = \dfrac{R^3}{(d-x)^2} \quad \Leftrightarrow \quad \dfrac{(d-x)^2}{x^2} = \dfrac{R^3}{r^3} \quad \Leftrightarrow$

$$\left(\frac{d}{x} - 1\right)^2 = \left(\frac{R}{r}\right)^3 \quad \Rightarrow \quad \frac{d}{x} - 1 = \left(\frac{R}{r}\right)^{3/2} \quad \Leftrightarrow \quad \frac{d}{x} = 1 + \left(\frac{R}{r}\right)^{3/2} \quad \Leftrightarrow \quad x = x^* = \frac{d}{1 + (R/r)^{3/2}}.$$

Now $A'(x) = 2\pi\left(\dfrac{r^3}{x^2} - \dfrac{R^3}{(d-x)^2}\right) \quad \Rightarrow \quad A''(x) = 2\pi\left(-\dfrac{2r^3}{x^3} - \dfrac{2R^3}{(d-x)^3}\right)$ and $A''(x^*) < 0$, so we have a local maximum at $x = x^*$.

However, x^* may not be an allowable value of x — we must show that x^* is between r and $d - R$.

(1) $x^* \geq r \quad \Leftrightarrow \quad \dfrac{d}{1 + (R/r)^{3/2}} \geq r \quad \Leftrightarrow \quad d \geq r + R\sqrt{R/r}$

(2) $x^* \leq d - R \quad \Leftrightarrow \quad \dfrac{d}{1 + (R/r)^{3/2}} \leq d - R \quad \Leftrightarrow \quad d \leq d - R + d\left(\dfrac{R}{r}\right)^{3/2} - R\left(\dfrac{R}{r}\right)^{3/2} \quad \Leftrightarrow$

$R + R\left(\dfrac{R}{r}\right)^{3/2} \leq d\left(\dfrac{R}{r}\right)^{3/2} \quad \Leftrightarrow \quad d \geq \dfrac{R}{(R/r)^{3/2}} + R = R + r\sqrt{r/R}$, but

$R + r\sqrt{r/R} \leq R + r$, and since $d > r + R$ [given], we conclude that $x^* \leq d - R$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Thus, from (1) and (2), x^* is not an allowable value of x if $d < r + R\sqrt{R/r}$.

So A may have a maximum at $x = r$, x^*, or $d - R$.

$$A(r) = \frac{2\pi R^2(d-r-R)}{d-r} \qquad \text{and} \qquad A(d-R) = \frac{2\pi r^2(d-r-R)}{d-R}$$

$A(r) > A(d-R) \;\Leftrightarrow\; \dfrac{R^2}{d-r} > \dfrac{r^2}{d-R} \;\Leftrightarrow\; R^2(d-R) > r^2(d-r) \;\Leftrightarrow\; R^2d - R^3 > r^2d - r^3 \;\Leftrightarrow$ $R^2d - r^2d > R^3 - r^3 \;\Leftrightarrow\; d(R-r)(R+r) > (R-r)(R^2+Rr+r^2) \;\Leftrightarrow\; d > (R^2+Rr+r^2)/(R+r) \;\Leftrightarrow$ $d > [(R+r)^2 - Rr]/(R+r) \;\Leftrightarrow\; d > R + r - Rr/(R+r)$. Now $R + r - Rr/(R+r) < R + r$, and we know that $d > R + r$, so we conclude that $A(r) > A(d-R)$.

In conclusion, A has an absolute maximum at $x = x^*$ provided $d \geq r + R\sqrt{R/r}$; otherwise, A has its maximum at $x = r$.

5. (a) Choose a vertical x-axis pointing downward with its origin at the surface. In order to calculate the pressure at depth z, consider n subintervals of the interval $[0, z]$ by points x_i and choose a point $x_i^* \in [x_{i-1}, x_i]$ for each i. The thin layer of water lying between depth x_{i-1} and depth x_i has a density of approximately $\rho(x_i^*)$, so the weight of a piece of that layer with unit cross-sectional area is $\rho(x_i^*)g\,\Delta x$. The total weight of a column of water extending from the surface to depth z (with unit cross-sectional area) would be approximately $\sum_{i=1}^{n} \rho(x_i^*)g\,\Delta x$. The estimate becomes exact if we take the limit as $n \to \infty$; weight (or force) per unit area at depth z is $W = \lim_{n\to\infty} \sum_{i=1}^{n} \rho(x_i^*)g\,\Delta x$. In other words, $P(z) = \int_0^z \rho(x)g\,dx$. More generally, if we make no assumptions about the location of the origin, then $P(z) = P_0 + \int_0^z \rho(x)g\,dx$, where P_0 is the pressure at $x = 0$. Differentiating, we get $dP/dz = \rho(z)g$.

(b)

$$\begin{aligned}
F &= \textstyle\int_{-r}^{r} P(L+x)\cdot 2\sqrt{r^2-x^2}\,dx \\
&= \textstyle\int_{-r}^{r}\Big(P_0 + \int_0^{L+x} \rho_0 e^{z/H} g\,dz\Big)\cdot 2\sqrt{r^2-x^2}\,dx \\
&= P_0\textstyle\int_{-r}^{r} 2\sqrt{r^2-x^2}\,dx + \rho_0 gH\int_{-r}^{r}\Big(e^{(L+x)/H}-1\Big)\cdot 2\sqrt{r^2-x^2}\,dx \\
&= (P_0-\rho_0 gH)\textstyle\int_{-r}^{r} 2\sqrt{r^2-x^2}\,dx + \rho_0 gH\int_{-r}^{r} e^{(L+x)/H}\cdot 2\sqrt{r^2-x^2}\,dx \\
&= (P_0-\rho_0 gH)\big(\pi r^2\big) + \rho_0 gHe^{L/H}\textstyle\int_{-r}^{r} e^{x/H}\cdot 2\sqrt{r^2-x^2}\,dx
\end{aligned}$$

6. The problem can be reduced to finding the line which minimizes the shaded area in the diagram. An equation of the circle in the first quadrant is $x = \sqrt{1-y^2}$. So the shaded area is

$$\begin{aligned}
A(h) &= \int_0^h \Big(1-\sqrt{1-y^2}\Big)\,dy + \int_h^1 \sqrt{1-y^2}\,dy \\
&= \int_0^h \Big(1-\sqrt{1-y^2}\Big)\,dy - \int_1^h \sqrt{1-y^2}\,dy
\end{aligned}$$

$$A'(h) = 1 - \sqrt{1-h^2} - \sqrt{1-h^2} \quad \text{[by FTC]} \; = 1 - 2\sqrt{1-h^2}$$

$A' = 0 \;\Leftrightarrow\; \sqrt{1-h^2} = \frac{1}{2} \;\Rightarrow\; 1 - h^2 = \frac{1}{4} \;\Rightarrow\; h^2 = \frac{3}{4} \;\Rightarrow\; h = \frac{\sqrt{3}}{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$A''(h) = -2 \cdot \frac{1}{2}(1-h^2)^{-1/2}(-2h) = \dfrac{2h}{\sqrt{1-h^2}} > 0$, so $h = \dfrac{\sqrt{3}}{2}$ gives a minimum value of A.

Note: Another strategy is to use the angle θ as the variable (see the diagram above) and show that $A = \theta + \cos\theta - \frac{\pi}{4} - \frac{1}{2}\sin 2\theta$, which is minimized when $\theta = \frac{\pi}{6}$.

7. To find the height of the pyramid, we use similar triangles. The first figure shows a cross-section of the pyramid passing through the top and through two opposite corners of the square base. Now $|BD| = b$, since it is a radius of the sphere, which has diameter $2b$ since it is tangent to the opposite sides of the square base. Also, $|AD| = b$ since $\triangle ADB$ is isosceles. So the height is $|AB| = \sqrt{b^2+b^2} = \sqrt{2}\,b$.

We first observe that the shared volume is equal to half the volume of the sphere, minus the sum of the four equal volumes (caps of the sphere) cut off by the triangular faces of the pyramid. See Exercise 6.2.49 for a derivation of the formula for the volume of a cap of a sphere. To use the formula, we need to find the perpendicular distance h of each triangular face from the surface of the sphere. We first find the distance d from the center of the sphere to one of the triangular faces. The third figure shows a cross-section of the pyramid through the top and through the midpoints of opposite sides of the square base. From similar triangles we find that

$$\frac{d}{b} = \frac{|AB|}{|AC|} = \frac{\sqrt{2}\,b}{\sqrt{b^2 + \left(\sqrt{2}\,b\right)^2}} \quad \Rightarrow \quad d = \frac{\sqrt{2}\,b^2}{\sqrt{3b^2}} = \frac{\sqrt{6}}{3}\,b$$

So $h = b - d = b - \frac{\sqrt{6}}{3}b = \frac{3-\sqrt{6}}{3}b$. So, using the formula $V = \pi h^2(r - h/3)$ from Exercise 6.2.49 with $r = b$, we find that the volume of each of the caps is $\pi\left(\frac{3-\sqrt{6}}{3}b\right)^2\left(b - \frac{3-\sqrt{6}}{3\cdot 3}b\right) = \frac{15-6\sqrt{6}}{9} \cdot \frac{6+\sqrt{6}}{9}\pi b^3 = \left(\frac{2}{3} - \frac{7}{27}\sqrt{6}\right)\pi b^3$. So, using our first observation, the shared volume is $V = \frac{1}{2}\left(\frac{4}{3}\pi b^3\right) - 4\left(\frac{2}{3} - \frac{7}{27}\sqrt{6}\right)\pi b^3 = \left(\frac{28}{27}\sqrt{6} - 2\right)\pi b^3$.

8. Orient the positive x-axis as in the figure.

Suppose that the plate has height h and is symmetric about the x-axis. At depth x below the water ($2 \le x \le 2+h$), let the width of the plate be $2f(x)$. Now each of the n horizontal strips has height h/n and the ith strip ($1 \le i \le n$) goes from $x = 2 + \left(\dfrac{i-1}{n}\right)h$ to $x = 2 + \left(\dfrac{i}{n}\right)h$. The hydrostatic force on the ith strip is $F(i) = \displaystyle\int_{2+[(i-1)/n]h}^{2+(i/n)h} 62.5x[2f(x)]\,dx$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

If we now let $x[2f(x)] = k$ (a constant) so that $f(x) = k/(2x)$, then

$$F(i) = \int_{2+[(i-1)/n]h}^{2+(i/n)h} 62.5k\,dx = 62.5k\,\big[\,x\,\big]_{2+[(i-1)/n]h}^{2+(i/n)h} = 62.5k\left[\left(2+\frac{i}{n}h\right)-\left(2+\frac{i-1}{n}\,h\right)\right] = 62.5k\left(\frac{h}{n}\right)$$

So the hydrostatic force on the ith strip is independent of i, that is, the force on each strip is the same. So the plate can be shaped as shown in the figure. (In fact, the required condition is satisfied whenever the plate has width C/x at depth x, for some constant C. Many shapes are possible.)

9. We can assume that the cut is made along a vertical line $x = b > 0$, that the disk's boundary is the circle $x^2 + y^2 = 1$, and that the center of mass of the smaller piece (to the right of $x = b$) is $\left(\frac{1}{2}, 0\right)$. We wish to find b to two decimal places. We have $\dfrac{1}{2} = \overline{x} = \dfrac{\int_b^1 x \cdot 2\sqrt{1-x^2}\,dx}{\int_b^1 2\sqrt{1-x^2}\,dx}$. Evaluating the numerator gives us $-\int_b^1 (1-x^2)^{1/2}(-2x)\,dx = -\frac{2}{3}\left[\left(1-x^2\right)^{3/2}\right]_b^1 = -\frac{2}{3}\left[0-\left(1-b^2\right)^{3/2}\right] = \frac{2}{3}(1-b^2)^{3/2}$.

Using Formula 30 in the table of integrals, we find that the denominator is

$\left[x\sqrt{1-x^2} + \sin^{-1}x\right]_b^1 = \left(0 + \frac{\pi}{2}\right) - \left(b\sqrt{1-b^2} + \sin^{-1}b\right)$. Thus, we have $\dfrac{1}{2} = \overline{x} = \dfrac{\frac{2}{3}(1-b^2)^{3/2}}{\frac{\pi}{2} - b\sqrt{1-b^2} - \sin^{-1}b}$, or, equivalently, $\frac{2}{3}(1-b^2)^{3/2} = \frac{\pi}{4} - \frac{1}{2}b\sqrt{1-b^2} - \frac{1}{2}\sin^{-1}b$. Solving this equation numerically with a calculator or CAS, we obtain $b \approx 0.138173$, or $b = 0.14$ m to two decimal places.

10. $A_1 = 30 \;\Rightarrow\; \frac{1}{2}bh = 30 \;\Rightarrow\; bh = 60.$

$\overline{x} = 6 \;\Rightarrow\; \dfrac{1}{A_2}\displaystyle\int_0^{10} x f(x)\,dx = 6 \;\Rightarrow$

$\displaystyle\int_0^b x\left(\frac{h}{b}x + 10 - h\right)dx + \int_b^{10} x(10)\,dx = 6(70) \;\Rightarrow$

$\displaystyle\int_0^b \left(\frac{h}{b}x^2 + 10x - hx\right)dx + 10\cdot\tfrac{1}{2}\left[x^2\right]_b^{10} = 420 \;\Rightarrow$

$\left[\dfrac{h}{3b}x^3 + 5x^2 - \dfrac{h}{2}x^2\right]_0^b + 5(100 - b^2) = 420 \;\Rightarrow\; \frac{1}{3}hb^2 + 5b^2 - \frac{1}{2}hb^2 + 500 - 5b^2 = 420 \;\Rightarrow\; 80 = \frac{1}{6}hb^2 \;\Rightarrow$

$480 = (hb)b \;\Rightarrow\; 480 = 60b \;\Rightarrow\; b = 8$. So $h = \frac{60}{8} = \frac{15}{2}$ and an equation of the line is

$y = \dfrac{15/2}{8}x + \left(10 - \dfrac{15}{2}\right) = \dfrac{15}{16}x + \dfrac{5}{2}$. Now

$$\begin{aligned}\overline{y} &= \frac{1}{A_2}\int_0^{10} \tfrac{1}{2}[f(x)]^2\,dx = \frac{1}{70\cdot 2}\left[\int_0^8 \left(\frac{15}{16}x + \frac{5}{2}\right)^2 dx + \int_8^{10}(10)^2\,dx\right]\\ &= \tfrac{1}{140}\left[\int_0^8 \left(\tfrac{225}{256}x^2 + \tfrac{75}{16}x + \tfrac{25}{4}\right)dx + 100(10-8)\right] = \tfrac{1}{140}\left(\left[\tfrac{225}{768}x^3 + \tfrac{75}{32}x^2 + \tfrac{25}{4}x\right]_0^8 + 200\right)\\ &= \tfrac{1}{140}(150 + 150 + 50 + 200) = \tfrac{550}{140} = \tfrac{55}{14}\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Another solution: Assume that the right triangle cut from the square has legs a cm and b cm long as shown. The triangle has area 30 cm^2, so $\frac{1}{2}ab = 30$ and $ab = 60$. We place the square in the first quadrant of the xy-plane as shown, and we let T, R, and S denote the triangle, the remaining portion of the square, and the full square, respectively. By symmetry, the centroid of S is $(5, 5)$. By Exercise 8.3.39, the centroid of T is $\left(\dfrac{b}{3}, 10 - \dfrac{a}{3}\right)$.

We are given that the centroid of R is $(6, c)$, where c is to be determined. We take the density of the square to be 1, so that areas can be used as masses. Then T has mass $m_T = 30$, S has mass $m_S = 100$, and R has mass $m_R = m_S - m_T = 70$. By reasoning as in Exercises 40 and 41 of Section 8.3, we view S as consisting of a mass m_T at the centroid $(\overline{x}_T, \overline{y}_T)$ of T and a mass R at the centroid $(\overline{x}_R, \overline{y}_R)$ of R. Then $\overline{x}_S = \dfrac{m_T\overline{x}_T + m_R\overline{x}_R}{m_T + m_R}$ and $\overline{y}_S = \dfrac{m_T\overline{y}_T + m_R\overline{y}_R}{m_T + m_R}$; that is,

$5 = \dfrac{30(b/3) + 70(6)}{100}$ and $5 = \dfrac{30(10 - a/3) + 70c}{100}$.

Solving the first equation for b, we get $b = 8$ cm. Since $ab = 60$ cm^2, it follows that $a = \frac{60}{8} = 7.5$ cm. Now the second equation says that $70c = 200 + 10a$, so $7c = 20 + a = \frac{55}{2}$ and $c = \frac{55}{14} = 3.9\overline{285714}$ cm. The solution is depicted in the figure.

11. If $h = L$, then $P = \dfrac{\text{area under } y = L\sin\theta}{\text{area of rectangle}} = \dfrac{\int_0^\pi L\sin\theta\,d\theta}{\pi L} = \dfrac{[-\cos\theta]_0^\pi}{\pi} = \dfrac{-(-1)+1}{\pi} = \dfrac{2}{\pi}$.

If $h = \frac{1}{2}L$, we replace L with $\frac{1}{2}L$ in the above calculation to get $P = \frac{1}{2}\left(\frac{2}{\pi}\right) = \frac{1}{\pi}$.

12. (a) The total set of possibilities can be identified with the rectangular region $\mathcal{R} = \{(\theta, y) \mid 0 \le y < L, 0 \le \theta < \pi\}$. Even when $h > L$, the needle intersects at least one line if and only if $y \le h\sin\theta$. Let $\mathcal{R}_1 = \{(\theta, y) \mid 0 \le y \le h\sin\theta, 0 \le \theta < \pi\}$. When $h \le L$, $\mathcal{R}_1$ is contained in $\mathcal{R}$, but that is no longer true when $h > L$. Thus, the probability that the needle intersects a line becomes

$$P = \frac{\text{area}(\mathcal{R} \cap \mathcal{R}_1)}{\text{area}(\mathcal{R})} = \frac{\text{area}(\mathcal{R} \cap \mathcal{R}_1)}{\pi L}$$

When $h > L$, the curve $y = h\sin\theta$ intersects the line $y = L$ twice — at $\left(\sin^{-1}(L/h), L\right)$ and at $\left(\pi - \sin^{-1}(L/h), L\right)$. Set $\theta_1 = \sin^{-1}(L/h)$ and $\theta_2 = \pi - \theta_1$. Then

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\begin{aligned}
\text{area}(\mathcal{R}\cap\mathcal{R}_1) &= \int_0^{\theta_1} h\sin\theta\,d\theta + \int_{\theta_1}^{\theta_2} L\,d\theta + \int_{\theta_2}^{\pi} h\sin\theta\,d\theta \\
&= 2\int_0^{\theta_1} h\sin\theta\,d\theta + L(\theta_2-\theta_1) = 2h\left[-\cos\theta\right]_0^{\theta_1} + L(\pi-2\theta_1) \\
&= 2h(1-\cos\theta_1) + L(\pi-2\theta_1) \\
&= 2h\left(1-\frac{\sqrt{h^2-L^2}}{h}\right) + L\left[\pi - 2\sin^{-1}\left(\frac{L}{h}\right)\right] \\
&= 2h - 2\sqrt{h^2-L^2} + \pi L - 2L\sin^{-1}\left(\frac{L}{h}\right)
\end{aligned}$$

We are told that $L=4$ and $h=7$, so $\text{area}(\mathcal{R}\cap\mathcal{R}_1) = 14 - 2\sqrt{33} + 4\pi - 8\sin^{-1}\left(\frac{4}{7}\right) \approx 10.21128$ and $P = \frac{1}{4\pi}\,\text{area}(\mathcal{R}\cap\mathcal{R}_1) \approx 0.812588$. (By comparison, $P = \frac{2}{\pi} \approx 0.636620$ when $h=L$, as shown in the solution to Problem 11.)

(b) The needle intersects at least two lines when $y + L \le h\sin\theta$; that is, when $y \le h\sin\theta - L$. Set $\mathcal{R}_2 = \{(\theta, y) \mid 0 \le y \le h\sin\theta - L, 0 \le \theta < \pi\}$. Then the probability that the needle intersects at least two lines is

$$P_2 = \frac{\text{area}(\mathcal{R}\cap\mathcal{R}_2)}{\text{area}(\mathcal{R})} = \frac{\text{area}(\mathcal{R}\cap\mathcal{R}_2)}{\pi L}$$

When $L=4$ and $h=7$, $\mathcal{R}_2$ is contained in $\mathcal{R}$ (see the figure). Thus,

$$\begin{aligned}
P_2 &= \frac{1}{4\pi}\,\text{area}(\mathcal{R}_2) = \frac{1}{4\pi}\int_{\sin^{-1}(4/7)}^{\pi-\sin^{-1}(4/7)} (7\sin\theta - 4)\,d\theta = \tfrac{1}{4\pi}\cdot 2\int_{\sin^{-1}(4/7)}^{\pi/2} (7\sin\theta-4)\,d\theta \\
&= \frac{1}{2\pi}\left[-7\cos\theta - 4\theta\right]_{\sin^{-1}(4/7)}^{\pi/2} = \frac{1}{2\pi}\left[0 - 2\pi + 7\frac{\sqrt{33}}{7} + 4\sin^{-1}\left(\tfrac{4}{7}\right)\right] = \frac{\sqrt{33} + 4\sin^{-1}\left(\frac{4}{7}\right) - 2\pi}{2\pi} \\
&\approx 0.301497
\end{aligned}$$

(c) The needle intersects at least three lines when $y + 2L \le h\sin\theta$: that is, when $y \le h\sin\theta - 2L$. Set $\mathcal{R}_3 = \{(\theta, y) \mid 0 \le y \le h\sin\theta - 2L, 0 \le \theta < \pi\}$. Then the probability that the needle intersects at least three lines is $P_3 = \dfrac{\text{area}(\mathcal{R}\cap\mathcal{R}_3)}{\text{area}(\mathcal{R})} = \dfrac{\text{area}(\mathcal{R}\cap\mathcal{R}_3)}{\pi L}$. (At this point, the generalization to P_n, n any positive integer, should be clear.)

Under the given assumption,

$$\begin{aligned}
P_3 &= \frac{1}{\pi L}\,\text{area}(\mathcal{R}_3) = \frac{1}{\pi L}\int_{\sin^{-1}(2L/h)}^{\pi-\sin^{-1}(2L/h)} (h\sin\theta - 2L)\,d\theta = \frac{2}{\pi L}\int_{\sin^{-1}(2L/h)}^{\pi/2} (h\sin\theta - 2L)\,d\theta \\
&= \frac{2}{\pi L}\left[-h\cos\theta - 2L\theta\right]_{\sin^{-1}(2L/h)}^{\pi/2} = \frac{2}{\pi L}\left[-\pi L + \sqrt{h^2-4L^2} + 2L\sin^{-1}(2L/h)\right]
\end{aligned}$$

Note that the probability that a needle touches exactly one line is $P_1 - P_2$, the probability that it touches exactly two lines is $P_2 - P_3$, and so on.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. Solve for y: $x^2 + (x+y+1)^2 = 1 \quad \Rightarrow \quad (x+y+1)^2 = 1 - x^2 \quad \Rightarrow \quad x+y+1 = \pm\sqrt{1-x^2} \quad \Rightarrow$

$y = -x - 1 \pm \sqrt{1-x^2}$.

$$A = \int_{-1}^{1} \left[\left(-x-1+\sqrt{1-x^2}\right) - \left(-x-1-\sqrt{1-x^2}\right)\right] dx$$

$$= \int_{-1}^{1} 2\sqrt{1-x^2}\, dx = 2\left(\tfrac{\pi}{2}\right) \quad \left[\begin{array}{c}\text{area of}\\ \text{semicircle}\end{array}\right] \quad = \pi$$

$$\overline{x} = \frac{1}{A}\int_{-1}^{1} x \cdot 2\sqrt{1-x^2}\, dx = 0 \quad [\text{odd integrand}]$$

$$\overline{y} = \frac{1}{A}\int_{-1}^{1} \frac{1}{2}\left[\left(-x-1+\sqrt{1-x^2}\right)^2 - \left(-x-1-\sqrt{1-x^2}\right)^2\right] dx = \frac{1}{\pi}\int_{-1}^{1} \frac{1}{2}\left(-4x\sqrt{1-x^2} - 4\sqrt{1-x^2}\right) dx$$

$$= -\frac{2}{\pi}\int_{-1}^{1} \left(x\sqrt{1-x^2} + \sqrt{1-x^2}\right) dx = -\frac{2}{\pi}\int_{-1}^{1} x\sqrt{1-x^2}\, dx - \frac{2}{\pi}\int_{-1}^{1} \sqrt{1-x^2}\, dx$$

$$= -\frac{2}{\pi}(0) \quad [\text{odd integrand}] \quad -\frac{2}{\pi}\left(\frac{\pi}{2}\right) \quad \left[\begin{array}{c}\text{area of}\\ \text{semicircle}\end{array}\right] \quad = -1$$

Thus, as expected, the centroid is $(\overline{x}, \overline{y}) = (0, -1)$. We might expect this result since the centroid of an ellipse is located at its center.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9 ☐ DIFFERENTIAL EQUATIONS

9.1 Modeling with Differential Equations

1. $\dfrac{dr}{dt} = \dfrac{k}{r}$, where k is a proportionality constant.

2. $\dfrac{dv}{dt} = k$, where k is a constant.

3. $\dfrac{dv}{dt} = k\,(M - v)$, where k is a proportionality constant.

4. $\dfrac{dy}{dt} = ky\,(N - y)$, where k is a proportionality constant.

5. The number of individuals who have *not* heard about the product is $N - y$. Thus, $\dfrac{dy}{dt} = k\,(N - y)$, where k is a proportionality constant.

6. $y = \sin x - \cos x \quad \Rightarrow \quad y' = \cos x + \sin x$. To determine whether y is a solution of the differential equation, we will substitute the expressions for y and y' in the left-hand side of the equation and see if the left-hand side is equal to the right-hand side. $\text{LHS} = y' + y = (\cos x + \sin x) + (\sin x - \cos x) = 2\sin x = \text{RHS}$, so $y = \sin x - \cos x$ **is** a solution of the differential equation.

7. $y = \frac{2}{3}e^x + e^{-2x} \quad \Rightarrow \quad y' = \frac{2}{3}e^x - 2e^{-2x}$. To determine whether y is a solution of the differential equation, we will substitute the expressions for y and y' in the left-hand side of the equation and see if the left-hand side is equal to the right-hand side.

$$\text{LHS} = y' + 2y = \tfrac{2}{3}e^x - 2e^{-2x} + 2\left(\tfrac{2}{3}e^x + e^{-2x}\right) = \tfrac{2}{3}e^x - 2e^{-2x} + \tfrac{4}{3}e^x + 2e^{-2x} = \tfrac{6}{3}e^x = 2e^x = \text{RHS}$$

so $y = \frac{2}{3}e^x + e^{-2x}$ **is** a solution of the differential equation.

8. $y = \tan x \quad \Rightarrow \quad y' = \sec^2 x$.

$\text{LHS} = y' - y^2 = \sec^2 x - \tan^2 x = 1 \left[\begin{array}{c}\text{using the identity}\\ 1 + \tan^2 x = \sec^2 x\end{array}\right] = \text{RHS}$, so $y = \tan x$ **is** a solution of the differential equation.

9. $y = \sqrt{x} = x^{1/2} \quad \Rightarrow \quad y' = \frac{1}{2}x^{-1/2}$.

$\text{LHS} = xy' - y = x\left(\frac{1}{2}x^{-1/2}\right) - x^{1/2} = \frac{1}{2}x^{1/2} - x^{1/2} = -\frac{1}{2}x^{1/2} \neq 0$, so $y = \sqrt{x}$ **is not** a solution of the differential equation.

10. $y = \sqrt{1 - x^2} = \left(1 - x^2\right)^{1/2} \quad \Rightarrow \quad y' = \dfrac{1}{2}(1 - x^2)^{-1/2}(-2x) = -\dfrac{x}{\sqrt{1 - x^2}}$.

$\text{LHS} = y\,y' - x = \sqrt{1 - x^2}\left(-\dfrac{x}{\sqrt{1 - x^2}}\right) - x = -x - x = -2x \neq 0$, so $y = \sqrt{1 - x^2}$ **is not** a solution of the differential equation.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. $y = x^3 \quad \Rightarrow \quad y' = 3x^2 \quad \Rightarrow \quad y'' = 6x.$

LHS $= x^2y'' - 6y = x^2 \cdot 6x - 6 \cdot x^3 = 6x^3 - 6x^3 = 0 =$ RHS, so $y = x^3$ **is** a solution of the differential equation.

12. $y = \ln x \quad \Rightarrow \quad y' = 1/x \quad \Rightarrow \quad y'' = -1/x^2.$

LHS $= x\,y'' - y' = x\left(-\dfrac{1}{x^2}\right) - \dfrac{1}{x} = -\dfrac{1}{x} - \dfrac{1}{x} = -\dfrac{2}{x} \neq 0$, so $y = \ln x$ **is not** a solution of the differential equation.

13. $y = -t\cos t - t \quad \Rightarrow \quad dy/dt = -t(-\sin t) + \cos t(-1) - 1 = t\sin t - \cos t - 1.$

LHS $= t\,\frac{dy}{dt} = t(t\sin t - \cos t - 1) = t^2\sin t - t\cos t - t = t^2\sin t + (-t\cos t - t) = t^2\sin t + y =$ RHS,

so y **is** a solution of the differential equation. Also, $y(\pi) = -\pi\cos\pi - \pi = -\pi(-1) - \pi = \pi - \pi = 0$, so the initial condition, $y(\pi) = 0$, is satisfied.

14. $y = 5e^{2x} + x \quad \Rightarrow \quad dy/dx = 10e^{2x} + 1.$

LHS $= \frac{dy}{dx} - 2y = 10e^{2x} + 1 - 2(5e^{2x} + x) = 1 - 2x =$ RHS, so y **is** a solution of the differential equation. Also,

$y(0) = 5e^{2(0)} + 0 = 5$, so the initial condition, $y(0) = 5$, is satisfied.

15. (a) $y = e^{rx} \quad \Rightarrow \quad y' = re^{rx} \quad \Rightarrow \quad y'' = r^2e^{rx}$. Substituting these expressions into the differential equation

$2y'' + y' - y = 0$, we get $2r^2e^{rx} + re^{rx} - e^{rx} = 0 \quad \Rightarrow \quad (2r^2 + r - 1)e^{rx} = 0 \quad \Rightarrow$

$(2r - 1)(r + 1) = 0$ [since e^{rx} is never zero] $\quad \Rightarrow \quad r = \frac{1}{2}$ or -1.

(b) Let $r_1 = \frac{1}{2}$ and $r_2 = -1$, so we need to show that every member of the family of functions $y = ae^{x/2} + be^{-x}$ is a solution of the differential equation $2y'' + y' - y = 0$.

$y = ae^{x/2} + be^{-x} \quad \Rightarrow \quad y' = \frac{1}{2}ae^{x/2} - be^{-x} \quad \Rightarrow \quad y'' = \frac{1}{4}ae^{x/2} + be^{-x}.$

$$\begin{aligned}
\text{LHS} = 2y'' + y' - y &= 2\left(\tfrac{1}{4}ae^{x/2} + be^{-x}\right) + \left(\tfrac{1}{2}ae^{x/2} - be^{-x}\right) - (ae^{x/2} + be^{-x}) \\
&= \tfrac{1}{2}ae^{x/2} + 2be^{-x} + \tfrac{1}{2}ae^{x/2} - be^{-x} - ae^{x/2} - be^{-x} \\
&= \left(\tfrac{1}{2}a + \tfrac{1}{2}a - a\right)e^{x/2} + (2b - b - b)e^{-x} \\
&= 0 = \text{RHS}
\end{aligned}$$

16. (a) $y = \cos kt \quad \Rightarrow \quad y' = -k\sin kt \quad \Rightarrow \quad y'' = -k^2\cos kt$. Substituting these expressions into the differential equation

$4y'' = -25y$, we get $4(-k^2\cos kt) = -25(\cos kt) \quad \Rightarrow \quad (25 - 4k^2)\cos kt = 0$ [for all t] $\quad \Rightarrow \quad 25 - 4k^2 = 0 \quad \Rightarrow$

$k^2 = \frac{25}{4} \quad \Rightarrow \quad k = \pm\frac{5}{2}$.

(b) $y = A\sin kt + B\cos kt \quad \Rightarrow \quad y' = Ak\cos kt - Bk\sin kt \quad \Rightarrow \quad y'' = -Ak^2\sin kt - Bk^2\cos kt.$

The given differential equation $4y'' = -25y$ is equivalent to $4y'' + 25y = 0$. Thus,

$$\begin{aligned}
\text{LHS} = 4y'' + 25y &= 4(-Ak^2\sin kt - Bk^2\cos kt) + 25(A\sin kt + B\cos kt) \\
&= -4Ak^2\sin kt - 4Bk^2\cos kt + 25A\sin kt + 25B\cos kt \\
&= (25 - 4k^2)A\sin kt + (25 - 4k^2)B\cos kt \\
&= 0 \quad \text{since } k^2 = \tfrac{25}{4}.
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. (a) $y = \sin x \quad \Rightarrow \quad y' = \cos x \quad \Rightarrow \quad y'' = -\sin x.$

LHS $= y'' + y = -\sin x + \sin x = 0 \neq \sin x$, so $y = \sin x$ **is not** a solution of the differential equation.

(b) $y = \cos x \quad \Rightarrow \quad y' = -\sin x \quad \Rightarrow \quad y'' = -\cos x.$

LHS $= y'' + y = -\cos x + \cos x = 0 \neq \sin x$, so $y = \cos x$ **is not** a solution of the differential equation.

(c) $y = \frac{1}{2}x\sin x \quad \Rightarrow \quad y' = \frac{1}{2}(x\cos x + \sin x) \quad \Rightarrow \quad y'' = \frac{1}{2}(-x\sin x + \cos x + \cos x).$

LHS $= y'' + y = \frac{1}{2}(-x\sin x + 2\cos x) + \frac{1}{2}x\sin x = \cos x \neq \sin x$, so $y = \frac{1}{2}x\sin x$ **is not** a solution of the differential equation.

(d) $y = -\frac{1}{2}x\cos x \quad \Rightarrow \quad y' = -\frac{1}{2}(-x\sin x + \cos x) \quad \Rightarrow \quad y'' = -\frac{1}{2}(-x\cos x - \sin x - \sin x).$

LHS $= y'' + y = -\frac{1}{2}(-x\cos x - 2\sin x) + \left(-\frac{1}{2}x\cos x\right) = \sin x =$ RHS, so $y = -\frac{1}{2}x\cos x$ **is** a solution of the differential equation.

18. (a) $y = \dfrac{\ln x + C}{x} \quad \Rightarrow \quad y' = \dfrac{x \cdot (1/x) - (\ln x + C)}{x^2} = \dfrac{1 - \ln x - C}{x^2}.$

$$\text{LHS} = x^2y' + xy = x^2 \cdot \frac{1 - \ln x - C}{x^2} + x \cdot \frac{\ln x + C}{x}$$
$$= 1 - \ln x - C + \ln x + C = 1 = \text{RHS}, \text{ so } y \text{ is a solution of the differential equation.}$$

(b)

A few notes about the graph of $y = (\ln x + C)/x$:

(1) There is a vertical asymptote of $x = 0$.

(2) There is a horizontal asymptote of $y = 0$.

(3) $y = 0 \quad \Rightarrow \quad \ln x + C = 0 \quad \Rightarrow \quad x = e^{-C}$, so there is an x-intercept at e^{-C}.

(4) $y' = 0 \quad \Rightarrow \quad \ln x = 1 - C \quad \Rightarrow \quad x = e^{1-C}$, so there is a local maximum at $x = e^{1-C}$.

(c) $y(1) = 2 \quad \Rightarrow \quad 2 = \dfrac{\ln 1 + C}{1} \quad \Rightarrow \quad 2 = C$, so the solution is $y = \dfrac{\ln x + 2}{x}$ [shown in part (b)].

(d) $y(2) = 1 \quad \Rightarrow \quad 1 = \dfrac{\ln 2 + C}{2} \quad \Rightarrow \quad 2 + \ln 2 + C \quad \Rightarrow \quad C = 2 - \ln 2$, so the solution is $y = \dfrac{\ln x + 2 - \ln 2}{x}$ [shown in part (b)].

19. (a) Since the derivative $y' = -y^2$ is always negative (or 0, if $y = 0$), the function y must be decreasing (or equal to 0) on any interval on which it is defined.

(b) $y = \dfrac{1}{x + C} \quad \Rightarrow \quad y' = -\dfrac{1}{(x + C)^2}.$ LHS $= y' = -\dfrac{1}{(x + C)^2} = -\left(\dfrac{1}{x + C}\right)^2 = -y^2 =$ RHS

(c) $y = 0$ is a solution of $y' = -y^2$ that is not a member of the family in part (b).

(d) If $y(x) = \dfrac{1}{x + C}$, then $y(0) = \dfrac{1}{0 + C} = \dfrac{1}{C}$. Since $y(0) = 0.5$, $\dfrac{1}{C} = \dfrac{1}{2} \quad \Rightarrow \quad C = 2$, so $y = \dfrac{1}{x + 2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. (a) If x is close to 0, then xy^3 is close to 0, and hence, y' is close to 0. Thus, the graph of y must have a tangent line that is nearly horizontal. If x is large, then xy^3 is large, and the graph of y must have a tangent line that is nearly vertical. (In both cases, we assume reasonable values for y.)

(b) $y = (c - x^2)^{-1/2} \quad \Rightarrow \quad y' = x(c - x^2)^{-3/2}$. $\text{RHS} = xy^3 = x[(c - x^2)^{-1/2}]^3 = x(c - x^2)^{-3/2} = y' = \text{LHS}$

(c)

When x is close to 0, y' is also close to 0.

As x gets larger, so does $|y'|$.

(d) $y(0) = (c - 0)^{-1/2} = 1/\sqrt{c}$ and $y(0) = 2 \quad \Rightarrow \quad \sqrt{c} = \frac{1}{2} \quad \Rightarrow \quad c = \frac{1}{4}$, so $y = \left(\frac{1}{4} - x^2\right)^{-1/2}$.

21. (a) $\dfrac{dP}{dt} = 1.2P\left(1 - \dfrac{P}{4200}\right)$. Now $\dfrac{dP}{dt} > 0 \quad \Rightarrow \quad 1 - \dfrac{P}{4200} > 0$ [assuming that $P > 0$] $\quad \Rightarrow \quad \dfrac{P}{4200} < 1 \quad \Rightarrow$

$P < 4200 \quad \Rightarrow$ the population is increasing for $0 < P < 4200$.

(b) $\dfrac{dP}{dt} < 0 \quad \Rightarrow \quad P > 4200$

(c) $\dfrac{dP}{dt} = 0 \quad \Rightarrow \quad P = 4200$ or $P = 0$

22. (a) $\dfrac{dv}{dt} = -v[v^2 - (1 + a)v + a] = -v(v - a)(v - 1)$, so $\dfrac{dv}{dt} = 0 \quad \Leftrightarrow \quad v = 0, a$, or 1.

(b) With $0 < a < 1$, $dv/dt = -v(v - a)(v - 1) > 0 \quad \Leftrightarrow \quad v < 0$ or $a < v < 1$, so v is increasing on $(-\infty, 0)$ and $(a, 1)$.

(c) With $0 < a < 1$, $dv/dt = -v(v - a)(v - 1) < 0 \quad \Leftrightarrow \quad 0 < v < a$ or $v > 1$, so v is decreasing on $(0, a)$ and $(1, \infty)$.

23. (a) This function is increasing *and* also decreasing. But $dy/dt = e^t(y - 1)^2 \geq 0$ for all t, implying that the graph of the solution of the differential equation cannot be decreasing on any interval.

(b) When $y = 1$, $dy/dt = 0$, but the graph does not have a horizontal tangent line.

24. The graph for this exercise is shown in the figure at the right.

A. $y' = 1 + xy > 1$ for points in the first quadrant, but we can see that $y' < 0$ for some points in the first quadrant.

B. $y' = -2xy = 0$ when $x = 0$, but we can see that $y' > 0$ for $x = 0$. Thus, equations A and B are incorrect, so the correct equation is C.

C. $y' = 1 - 2xy$ seems reasonable since:

(1) When $x = 0$, y' could be 1.

(2) When $x < 0$, y' could be greater than 1.

(3) Solving $y' = 1 - 2xy$ for y gives us $y = \dfrac{1 - y'}{2x}$. If y' takes on small negative values, then as $x \to \infty$, $y \to 0^+$, as shown in the figure.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. (a) $y' = 1 + x^2 + y^2 \geq 1$ and $y' \to \infty$ as $x \to \infty$. The only curve satisfying these conditions is labeled III.

(b) $y' = xe^{-x^2-y^2} > 0$ if $x > 0$ and $y' < 0$ if $x < 0$. The only curve with negative tangent slopes when $x < 0$ and positive tangent slopes when $x > 0$ is labeled I.

(c) $y' = \dfrac{1}{1 + e^{x^2+y^2}} > 0$ and $y' \to 0$ as $x \to \infty$. The only curve satisfying these conditions is labeled IV.

(d) $y' = \sin(xy)\ \cos(xy) = 0$ if $y = 0$, which is the solution graph labeled II.

26. (a) The coffee cools most quickly as soon as it is removed from the heat source. The rate of cooling decreases toward 0 since the coffee approaches room temperature.

(b) $\dfrac{dy}{dt} = k(y - R)$, where k is a proportionality constant, y is the temperature of the coffee, and R is the room temperature. The initial condition is $y(0) = 95°\text{C}$. The answer and the model support each other because as y approaches R, dy/dt approaches 0, so the model seems appropriate.

(c)

27. (a) P increases most rapidly at the beginning, since there are usually many simple, easily-learned sub-skills associated with learning a skill. As t increases, we would expect dP/dt to remain positive, but decrease. This is because as time progresses, the only points left to learn are the more difficult ones.

(b) $\dfrac{dP}{dt} = k(M - P)$ is always positive, so the level of performance P is increasing. As P gets close to M, dP/dt gets close to 0; that is, the performance levels off, as explained in part (a).

(c)

28. (a) $\dfrac{dL}{dt} = k(L_\infty - L)$. Assuming $L_\infty > L$, we have $k > 0$ and $dL/dt > 0$ for all t.

(b)

29. If $c(t) = c_s\left(1 - e^{-\alpha t^{1-b}}\right) = c_s - c_s e^{-\alpha t^{1-b}}$ for $t > 0$, where $k > 0$, $c_s > 0$, $0 < b < 1$, and $\alpha = k/(1-b)$, then $\dfrac{dc}{dt} = c_s\left[0 - e^{-\alpha t^{1-b}} \cdot \dfrac{d}{dt}\left(-\alpha t^{1-b}\right)\right] = -c_s e^{-\alpha t^{1-b}} \cdot (-\alpha)(1-b)t^{-b} = \dfrac{\alpha(1-b)}{t^b}\, c_s\, e^{-\alpha t^{1-b}} = \dfrac{k}{t^b}(c_s - c)$. The equation for c indicates that as t increases, c approaches c_s. The differential equation indicates that as t increases, the rate of increase of c decreases steadily and approaches 0 as c approaches c_s.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9.2 Direction Fields and Euler's Method

1. (a)

(b) It appears that the constant functions $y = 0.5$ and $y = 1.5$ are equilibrium solutions. Note that these two values of y satisfy the given differential equation $y' = x \cos \pi y$.

2. (a)

(b) It appears that the constant functions $y = 0$, $y = 2$, and $y = 4$ are equilibrium solutions. Note that these three values of y satisfy the given differential equation $y' = \tan\left(\frac{1}{2}\pi y\right)$.

3. $y' = 2 - y$. The slopes at each point are independent of x, so the slopes are the same along each line parallel to the x-axis. Thus, III is the direction field for this equation. Note that for $y = 2$, $y' = 0$.

4. $y' = x(2 - y) = 0$ on the lines $x = 0$ and $y = 2$. Direction field I satisfies these conditions.

5. $y' = x + y - 1 = 0$ on the line $y = -x + 1$. Direction field IV satisfies this condition. Notice also that on the line $y = -x$ we have $y' = -1$, which is true in IV.

6. $y' = \sin x \sin y = 0$ on the lines $x = 0$ and $y = 0$, and $y' > 0$ for $0 < x < \pi$, $0 < y < \pi$. Direction field II satisfies these conditions.

7.

8.

9.

x	y	$y' = \frac{1}{2}y$
0	0	0
0	1	0.5
0	2	1
0	−3	−1.5
0	−2	−1

Note that for $y = 0$, $y' = 0$. The three solution curves sketched go through $(0, 0)$, $(0, 1)$, and $(0, -1)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10.

x	y	$y' = x - y + 1$
-1	0	0
-1	-1	1
0	0	1
0	1	0
0	2	-1
0	-1	2
0	-2	3
1	0	2
1	1	1

Note that $y' = 0$ for $y = x + 1$ and that $y' = 1$ for $y = x$. For any constant value of x, y' decreases as y increases and y' increases as y decreases. The three solution curves sketched go through $(0, 0)$, $(0, 1)$, and $(0, -1)$.

11.

x	y	$y' = y - 2x$
-2	-2	2
-2	2	6
2	2	-2
2	-2	-6

Note that $y' = 0$ for any point on the line $y = 2x$. The slopes are positive to the left of the line and negative to the right of the line. The solution curve in the graph passes through $(1, 0)$.

12.

x	y	$y' = xy - x^2$
2	3	2
-2	-3	2
± 2	0	-4
0	0	0
2	2	0

$y' = xy - x^2 = x(y - x)$, so $y' = 0$ for $x = 0$ and $y = x$. The slopes are positive only in the regions in quadrants I and III that are bounded by $x = 0$ and $y = x$. The solution curve in the graph passes through $(0, 1)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13.

x	y	$y' = y + xy$
0	± 2	± 2
1	± 2	± 4
-3	± 2	∓ 4

Note that $y' = y(x+1) = 0$ for any point on $y = 0$ or on $x = -1$. The slopes are positive when the factors y and $x+1$ have the same sign and negative when they have opposite signs. The solution curve in the graph passes through $(0, 1)$.

14.

x	y	$y' = x + y^2$
-2	± 1	-1
-2	± 2	2
2	± 1	3
0	± 2	4
0	0	0

Note that $y' = x + y^2 = 0$ only on the parabola $x = -y^2$. The slopes are positive "outside" $x = -y^2$ and negative "inside" $x = -y^2$. The solution curve in the graph passes through $(0, 0)$.

15. $y' = x^2 y - \frac{1}{2} y^2$ and $y(0) = 1$.

In Maple, use the following commands to obtain a similar figure.

```
with(DETools):
ODE:=diff(y(x),x)=x^2*y(x)-(1/2)*y(x)^2;
ivs:=[y(0)=1];
DEplot({ODE},y(x),x=-3..2,y=0..4,ivs,linecolor=black);
```

16. $y' = \cos(x + y)$ and $y(0) = 1$.

In Maple, use the following commands to obtain a similar figure.

```
with(DETools):
ODE:=diff(y(x),x)=cos(x+y(x));
ivs:=[y(0)=1];
DEplot({ODE},y(x),x=-1.5*Pi..1.5*Pi,y=-1.5*Pi..1.5*Pi,
ivs,linecolor=black);
```

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17.

The direction field is for the differential equation $y' = y^3 - 4y$.

$L = \lim_{t\to\infty} y(t)$ exists for $-2 \le c \le 2$;

$L = \pm 2$ for $c = \pm 2$ and $L = 0$ for $-2 < c < 2$.

For other values of c, L does not exist.

18.

Note that when $f(y) = 0$ on the graph in the text, we have $y' = f(y) = 0$; so we get horizontal segments at $y = \pm 1, \pm 2$. We get segments with negative slopes only for $1 < |y| < 2$. All other segments have positive slope. For the limiting behavior of solutions:

- If $y(0) > 2$, then $\lim_{t\to\infty} y = \infty$ and $\lim_{t\to-\infty} y = 2$.
- If $1 < y(0) < 2$, then $\lim_{t\to\infty} y = 1$ and $\lim_{t\to-\infty} y = 2$.
- If $-1 < y(0) < 1$, then $\lim_{t\to\infty} y = 1$ and $\lim_{t\to-\infty} y = -1$.
- If $-2 < y(0) < -1$, then $\lim_{t\to\infty} y = -2$ and $\lim_{t\to-\infty} y = -1$.
- If $y < -2$, then $\lim_{t\to\infty} y = -2$ and $\lim_{t\to-\infty} y = -\infty$.

19. (a) $y' = F(x, y) = y$ and $y(0) = 1 \quad \Rightarrow \quad x_0 = 0, y_0 = 1$.

(i) $h = 0.4$ and $y_1 = y_0 + hF(x_0, y_0) \quad \Rightarrow \quad y_1 = 1 + 0.4 \cdot 1 = 1.4$. $x_1 = x_0 + h = 0 + 0.4 = 0.4$,

so $y_1 = y\,(0.4) = 1.4$.

(ii) $h = 0.2 \quad \Rightarrow \quad x_1 = 0.2$ and $x_2 = 0.4$, so we need to find y_2.

$y_1 = y_0 + hF(x_0, y_0) = 1 + 0.2y_0 = 1 + 0.2 \cdot 1 = 1.2$,

$y_2 = y_1 + hF(x_1, y_1) = 1.2 + 0.2y_1 = 1.2 + 0.2 \cdot 1.2 = 1.44$.

(iii) $h = 0.1 \quad \Rightarrow \quad x_4 = 0.4$, so we need to find y_4. $y_1 = y_0 + hF(x_0, y_0) = 1 + 0.1y_0 = 1 + 0.1 \cdot 1 = 1.1$,

$y_2 = y_1 + hF(x_1, y_1) = 1.1 + 0.1y_1 = 1.1 + 0.1 \cdot 1.1 = 1.21$,

$y_3 = y_2 + hF(x_2, y_2) = 1.21 + 0.1y_2 = 1.21 + 0.1 \cdot 1.21 = 1.331$,

$y_4 = y_3 + hF(x_3, y_3) = 1.331 + 0.1y_3 = 1.331 + 0.1 \cdot 1.331 = 1.4641$.

(b)

We see that the estimates are underestimates since they are all below the graph of $y = e^x$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) (i) For $h = 0.4$: (exact value) $-$ (approximate value) $= e^{0.4} - 1.4 \approx 0.0918$

(ii) For $h = 0.2$: (exact value) $-$ (approximate value) $= e^{0.4} - 1.44 \approx 0.0518$

(iii) For $h = 0.1$: (exact value) $-$ (approximate value) $= e^{0.4} - 1.4641 \approx 0.0277$

Each time the step size is halved, the error estimate also appears to be halved (approximately).

20.

As x increases, the slopes decrease and all of the estimates are above the true values. Thus, all of the estimates are overestimates.

21. $h = 0.5$, $x_0 = 1$, $y_0 = 0$, and $F(x, y) = y - 2x$.

Note that $x_1 = x_0 + h = 1 + 0.5 = 1.5$, $x_2 = 2$, and $x_3 = 2.5$.

$y_1 = y_0 + hF(x_0, y_0) = 0 + 0.5F(1, 0) = 0.5[0 - 2(1)] = -1$.

$y_2 = y_1 + hF(x_1, y_1) = -1 + 0.5F(1.5, -1) = -1 + 0.5[-1 - 2(1.5)] = -3$.

$y_3 = y_2 + hF(x_2, y_2) = -3 + 0.5F(2, -3) = -3 + 0.5[-3 - 2(2)] = -6.5$.

$y_4 = y_3 + hF(x_3, y_3) = -6.5 + 0.5F(2.5, -6.5) = -6.5 + 0.5[-6.5 - 2(2.5)] = -12.25$.

22. $h = 0.2$, $x_0 = 0$, $y_0 = 1$, and $F(x, y) = x^2y - \frac{1}{2}y^2$. Note that $x_1 = x_0 + h = 0 + 0.2 = 0.2$, $x_2 = 0.4$, $x_3 = 0.6$, $x_4 = 0.8$, and $x_5 = 1$.

$y_1 = y_0 + hF(x_0, y_0) = 1 + 0.2F(0, 1) = 1 + 0.2\left[0^2(1) - \frac{1}{2}(1)^2\right] = 1 + 0.2\left(-\frac{1}{2}\right) = 0.9$.

$y_2 = y_1 + hF(x_1, y_1) = 0.9 + 0.2F(0.2, 0.9) = 0.9 + 0.2\left[(0.2)^2(0.9) - \frac{1}{2}(0.9)^2\right] = 0.8262$.

$y_3 = y_2 + hF(x_2, y_2) = 0.8262 + 0.2F(0.4, 0.8262) = 0.8262 + 0.2\left[(0.4)^2(0.8262) - \frac{1}{2}(0.8262)^2\right] = 0.784377756$.

$y_4 = y_3 + hF(x_3, y_3) = 0.784377756 + 0.2F(0.6, 0.784377756) \approx 0.779328108$.

$y_5 = y_4 + hF(x_4, y_4) \approx 0.779328108 + 0.2F(0.8, 0.779328108) \approx 0.818346876$.

Thus, $y(1) \approx 0.8183$.

23. $h = 0.1$, $x_0 = 0$, $y_0 = 1$, and $F(x, y) = y + xy$.

Note that $x_1 = x_0 + h = 0 + 0.1 = 0.1$, $x_2 = 0.2$, $x_3 = 0.3$, and $x_4 = 0.4$.

$y_1 = y_0 + hF(x_0, y_0) = 1 + 0.1F(0, 1) = 1 + 0.1[1 + (0)(1)] = 1.1$.

$y_2 = y_1 + hF(x_1, y_1) = 1.1 + 0.1F(0.1, 1.1) = 1.1 + 0.1[1.1 + (0.1)(1.1)] = 1.221$.

$y_3 = y_2 + hF(x_2, y_2) = 1.221 + 0.1F(0.2, 1.221) = 1.221 + 0.1[1.221 + (0.2)(1.221)] = 1.36752$.

$$\begin{aligned} y_4 = y_3 + hF(x_3, y_3) &= 1.36752 + 0.1F(0.3, 1.36752) = 1.36752 + 0.1[1.36752 + (0.3)(1.36752)] \\ &= 1.5452976. \end{aligned}$$

$$\begin{aligned} y_5 = y_4 + hF(x_4, y_4) &= 1.5452976 + 0.1F(0.4, 1.5452976) \\ &= 1.5452976 + 0.1[1.5452976 + (0.4)(1.5452976)] = 1.761639264. \end{aligned}$$

Thus, $y(0.5) \approx 1.7616$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. (a) $h = 0.2$, $x_0 = 0$, $y_0 = 0$, and $F(x, y) = \cos(x + y)$. Note that $x_1 = x_0 + h = 0 + 0.2 = 0.2$, $x_2 = 0.4$, and $x_3 = 0.6$.

$y_1 = y_0 + hF(x_0, y_0) = 0 + 0.2F(0, 0) = 0.2\cos(0 + 0) = 0.2(1) = 0.2.$

$y_2 = y_1 + hF(x_1, y_1) = 0.2 + 0.2F(0.2, 0.2) = 0.2 + 0.2\cos(0.4) \approx 0.3842121988.$

$y_3 = y_2 + hF(x_2, y_2) \approx 0.3842 + 0.2F(0.4, 0.3842) \approx 0.5258011763.$

Thus, $y(0.6) \approx 0.5258$.

(b) Now use $h = 0.1$. For $1 \le n \le 6$, $x_n = 0.n$.

$y_1 = y_0 + hF(x_0, y_0) = 0 + 0.1\cos(0 + 0) = 0.1(1) = 0.1.$

$y_2 = y_1 + hF(x_1, y_1) = 0.1 + 0.1\cos(0.2) \approx 0.1980.$

$y_3 = y_2 + hF(x_2, y_2) \approx 0.1980 + 0.1\cos(0.3980) \approx 0.2902.$

$y_4 = y_3 + hF(x_3, y_3) \approx 0.2902 + 0.1\cos(0.5902) \approx 0.3733.$

$y_5 = y_4 + hF(x_4, y_4) \approx 0.3733 + 0.1\cos(0.7733) \approx 0.4448.$

$y_6 = y_5 + hF(x_5, y_5) \approx 0.4448 + 0.1\cos(0.9448) \approx 0.5034.$

Thus, $y(0.6) \approx 0.5034$.

25. (a) $dy/dx + 3x^2y = 6x^2 \quad \Rightarrow \quad y' = 6x^2 - 3x^2y$. Store this expression in Y_1 and use the following simple program to evaluate $y(1)$ for each part, using $\mathrm{H} = h = 1$ and $\mathrm{N} = 1$ for part (i), $\mathrm{H} = 0.1$ and $\mathrm{N} = 10$ for part (ii), and so forth.

```
h → H: 0 → X: 3 → Y:
For(I, 1, N): Y + H × Y1 → Y: X + H → X:
End(loop):
Display Y.   [To see all iterations, include this statement in the loop.]
```

(i) $\mathrm{H} = 1, \mathrm{N} = 1 \quad \Rightarrow \quad y(1) = 3$

(ii) $\mathrm{H} = 0.1, \mathrm{N} = 10 \quad \Rightarrow \quad y(1) \approx 2.3928$

(iii) $\mathrm{H} = 0.01, \mathrm{N} = 100 \quad \Rightarrow \quad y(1) \approx 2.3701$

(iv) $\mathrm{H} = 0.001, \mathrm{N} = 1000 \quad \Rightarrow \quad y(1) \approx 2.3681$

(b) $y = 2 + e^{-x^3} \quad \Rightarrow \quad y' = -3x^2e^{-x^3}$

$$\text{LHS} = y' + 3x^2y = -3x^2e^{-x^3} + 3x^2\left(2 + e^{-x^3}\right) = -3x^2e^{-x^3} + 6x^2 + 3x^2e^{-x^3} = 6x^2 = \text{RHS}$$

$y(0) = 2 + e^{-0} = 2 + 1 = 3$

(c) The exact value of $y(1)$ is $2 + e^{-1^3} = 2 + e^{-1}$.

(i) For $h = 1$: (exact value) $-$ (approximate value) $= 2 + e^{-1} - 3 \approx -0.6321$

(ii) For $h = 0.1$: (exact value) $-$ (approximate value) $= 2 + e^{-1} - 2.3928 \approx -0.0249$

(iii) For $h = 0.01$: (exact value) $-$ (approximate value) $= 2 + e^{-1} - 2.3701 \approx -0.0022$

(iv) For $h = 0.001$: (exact value) $-$ (approximate value) $= 2 + e^{-1} - 2.3681 \approx -0.0002$

In (ii)–(iv), it seems that when the step size is divided by 10, the error estimate is also divided by 10 (approximately).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. (a) We use the program from the solution to Exercise 25 with $Y_1 = x^3 - y^3$, H $= 0.01$, and N $= \frac{2-0}{0.01} = 200$. With $(x_0, y_0) = (0, 1)$, we get $y(2) \approx 1.9000$.

(b)

Notice from the graph that $y(2) \approx 1.9$, which serves as a check on our calculation in part (a).

27. (a) $R\dfrac{dQ}{dt} + \dfrac{1}{C}Q = E(t)$ becomes $5Q' + \dfrac{1}{0.05}Q = 60$ or $Q' + 4Q = 12$.

(b) From the graph, it appears that the limiting value of the charge Q is about 3.

(c) If $Q' = 0$, then $4Q = 12 \quad \Rightarrow \quad Q = 3$ is an equilibrium solution.

(d)

(e) $Q' + 4Q = 12 \quad \Rightarrow \quad Q' = 12 - 4Q$. Now $Q(0) = 0$, so $t_0 = 0$ and $Q_0 = 0$.

$$Q_1 = Q_0 + hF(t_0, Q_0) = 0 + 0.1(12 - 4 \cdot 0) = 1.2$$
$$Q_2 = Q_1 + hF(t_1, Q_1) = 1.2 + 0.1(12 - 4 \cdot 1.2) = 1.92$$
$$Q_3 = Q_2 + hF(t_2, Q_2) = 1.92 + 0.1(12 - 4 \cdot 1.92) = 2.352$$
$$Q_4 = Q_3 + hF(t_3, Q_3) = 2.352 + 0.1(12 - 4 \cdot 2.352) = 2.6112$$
$$Q_5 = Q_4 + hF(t_4, Q_4) = 2.6112 + 0.1(12 - 4 \cdot 2.6112) = 2.76672$$

Thus, $Q_5 = Q(0.5) \approx 2.77$ C.

28. (a) From the solution to Exercise 9.1.26, we have $dy/dt = k(y - R)$. We are given that $R = 20°$C and $dy/dt = -1°$C/min when $y = 70°$C. Thus, $-1 = k(70 - 20) \quad \Rightarrow \quad k = -\frac{1}{50}$ and the differential equation becomes $dy/dt = -\frac{1}{50}(y - 20)$.

(b)

The limiting value of the temperature is 20°C; that is, the temperature of the room.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) From part (a), $dy/dt = -\frac{1}{50}(y-20)$. With $t_0 = 0$, $y_0 = 95$, and $h = 2$ min, we get

$$\begin{aligned}
y_1 &= y_0 + hF(t_0, y_0) = 95 + 2\left[-\tfrac{1}{50}(95-20)\right] = 92 \\
y_2 &= y_1 + hF(t_1, y_1) = 92 + 2\left[-\tfrac{1}{50}(92-20)\right] = 89.12 \\
y_3 &= y_2 + hF(t_2, y_2) = 89.12 + 2\left[-\tfrac{1}{50}(89.12-20)\right] = 86.3552 \\
y_4 &= y_3 + hF(t_3, y_3) = 86.3552 + 2\left[-\tfrac{1}{50}(86.3552-20)\right] = 83.700992 \\
y_5 &= y_4 + hF(t_4, y_4) = 83.700992 + 2\left[-\tfrac{1}{50}(83.700992-20)\right] = 81.15295232
\end{aligned}$$

Thus, $y(10) \approx 81.15°\text{C}$.

9.3 Separable Equations

1. $\dfrac{dy}{dx} = 3x^2y^2 \;\Rightarrow\; \dfrac{dy}{y^2} = 3x^2\,dx \quad [y \neq 0] \;\Rightarrow\; \int y^{-2}\,dy = \int 3x^2\,dx \;\Rightarrow\; -y^{-1} = x^3 + C \;\Rightarrow$

$\dfrac{-1}{y} = x^3 + C \;\Rightarrow\; y = \dfrac{-1}{x^3+C}$. $y = 0$ is also a solution.

2. $\dfrac{dy}{dx} = \dfrac{x}{y^4} \;\Rightarrow\; y^4\,dy = x\,dx \;\Rightarrow\; \int y^4\,dy = \int x\,dx \;\Rightarrow\; \frac{1}{5}y^5 = \frac{1}{2}x^2 + K \;\Rightarrow\; y^5 = \frac{5}{2}x^2 + 5K \;\Rightarrow$

$y = \sqrt[5]{\frac{5}{2}x^2 + C}$, where $C = 5K$.

3. $\dfrac{dy}{dx} = x\sqrt{y} \;\Rightarrow\; \dfrac{dy}{\sqrt{y}} = x\,dx \quad [y \neq 0] \;\Rightarrow\; \int y^{-1/2}\,dy = \int x\,dx \;\Rightarrow\; 2y^{1/2} = \frac{1}{2}x^2 + K \;\Rightarrow$

$\sqrt{y} = \frac{1}{4}x^2 + \frac{1}{2}K \;\Rightarrow\; y = \left(\frac{1}{4}x^2 + C\right)^2$, where $C = \frac{1}{2}K$. $y = 0$ is also a solution.

4. $x\,y' = y + 3 \;\Rightarrow\; x\dfrac{dy}{dx} = y + 3 \;\Rightarrow\; \dfrac{dy}{y+3} = \dfrac{dx}{x} \quad [x \neq 0, y \neq -3] \;\Rightarrow\; \displaystyle\int \frac{1}{y+3}\,dy = \int \frac{1}{x}\,dx \;\Rightarrow$

$\ln|y+3| = \ln|x| + C \;\Rightarrow\; |y+3| = e^{\ln|x|+C} = e^{\ln|x|}\,e^C = e^C\,|x| \;\Rightarrow\; y + 3 = kx$, where $k = \pm e^C$. Thus, $y = kx - 3$. (In our derivation, k was nonzero, but we can restore the excluded case $y = -3$ by allowing k to be zero.)

5. $xy\,y' = x^2 + 1 \;\Rightarrow\; xy\dfrac{dy}{dx} = x^2 + 1 \;\Rightarrow\; y\,dy = \dfrac{x^2+1}{x}\,dx \quad [x \neq 0] \;\Rightarrow\; \displaystyle\int y\,dy = \int \left(x + \frac{1}{x}\right)dx \;\Rightarrow$

$\frac{1}{2}y^2 = \frac{1}{2}x^2 + \ln|x| + K \;\Rightarrow\; y^2 = x^2 + 2\ln|x| + 2K \;\Rightarrow\; y = \pm\sqrt{x^2 + 2\ln|x| + C}$, where $C = 2K$.

6. $y' + xe^y = 0 \;\Rightarrow\; \dfrac{dy}{dx} = -xe^y \;\Rightarrow\; e^{-y}\,dy = -x\,dx \;\Rightarrow\; \int e^{-y}\,dy = \int -x\,dx \;\Rightarrow\; -e^{-y} = -\frac{1}{2}x^2 + C \;\Rightarrow$

$e^{-y} = \frac{1}{2}x^2 - C \;\Rightarrow\; -y = \ln\left(\frac{1}{2}x^2 - C\right) \;\Rightarrow\; y = -\ln\left(\frac{1}{2}x^2 - C\right)$

7. $(e^y - 1)\,y' = 2 + \cos x \;\Rightarrow\; (e^y - 1)\dfrac{dy}{dx} = 2 + \cos x \;\Rightarrow\; (e^y - 1)\,dy = (2 + \cos x)\,dx \;\Rightarrow$

$\int (e^y - 1)\,dy = \int (2 + \cos x)\,dx \;\Rightarrow\; e^y - y = 2x + \sin x + C$. We cannot solve explicitly for y.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $\dfrac{dy}{dx} = 2x(y^2+1) \quad \Rightarrow \quad \dfrac{dy}{y^2+1} = 2x\,dx \quad \Rightarrow \quad \displaystyle\int \frac{1}{y^2+1}dy = \int 2x\,dx \quad \Rightarrow \quad \tan^{-1}y = x^2 + C \quad \Rightarrow$

$y = \tan(x^2 + C)$

9. $\dfrac{dp}{dt} = t^2p - p + t^2 - 1 = p(t^2-1) + 1(t^2-1) = (p+1)(t^2-1) \quad \Rightarrow \quad \dfrac{1}{p+1}\,dp = (t^2-1)\,dt \quad \Rightarrow$

$\displaystyle\int \frac{1}{p+1}\,dp = \int (t^2-1)\,dt \quad \Rightarrow \quad \ln|p+1| = \tfrac{1}{3}t^3 - t + C \quad \Rightarrow \quad |p+1| = e^{t^3/3-t+C} \quad \Rightarrow \quad p+1 = \pm e^C e^{t^3/3-t} \quad \Rightarrow$

$p = Ke^{t^3/3-t} - 1$, where $K = \pm e^C$. Since $p = -1$ is also a solution, K can equal 0, and hence, K can be any real number.

10. $\dfrac{dz}{dt} + e^{t+z} = 0 \quad \Rightarrow \quad \dfrac{dz}{dt} = -e^te^z \quad \Rightarrow \quad \int e^{-z}\,dz = -\int e^t\,dt \quad \Rightarrow \quad -e^{-z} = -e^t + C \quad \Rightarrow \quad e^{-z} = e^t - C \quad \Rightarrow$

$\dfrac{1}{e^z} = e^t - C \quad \Rightarrow \quad e^z = \dfrac{1}{e^t - C} \quad \Rightarrow \quad z = \ln\left(\dfrac{1}{e^t - C}\right) \quad \Rightarrow \quad z = -\ln\left(e^t - C\right)$

11. $\dfrac{d\theta}{dt} = \dfrac{t\sec\theta}{\theta e^{t^2}} \quad \Rightarrow \quad \theta\cos\theta\,d\theta = te^{-t^2}\,dt \quad \Rightarrow \quad \int \theta\cos\theta\,d\theta = \int te^{-t^2}\,dt \quad \Rightarrow$

$\theta\sin\theta + \cos\theta = -\frac{1}{2}e^{-t^2} + C$ [by parts]. We cannot solve explicitly for θ.

12. $\dfrac{dH}{dR} = \dfrac{RH^2\sqrt{1+R^2}}{\ln H} \quad \Rightarrow \quad \dfrac{\ln H}{H^2}\,dH = R\sqrt{1+R^2}\,dR \quad \Rightarrow \quad \displaystyle\int \frac{\ln H}{H^2}\,dH = \int R(1+R^2)^{1/2}\,dR \quad \Rightarrow$

$-\dfrac{\ln H}{H} - \dfrac{1}{H} = \dfrac{1}{3}(1+R^2)^{3/2} + C$ [by parts]. We cannot solve explicitly for H.

13. $\dfrac{dy}{dx} = xe^y \quad \Rightarrow \quad e^{-y}\,dy = x\,dx \quad \Rightarrow \quad \int e^{-y}\,dy = \int x\,dx \quad \Rightarrow \quad -e^{-y} = \frac{1}{2}x^2 + C.$

$y(0) = 0 \quad \Rightarrow \quad -e^{-0} = \frac{1}{2}(0)^2 + C \quad \Rightarrow \quad C = -1$, so $-e^{-y} = \frac{1}{2}x^2 - 1 \quad \Rightarrow \quad e^{-y} = -\frac{1}{2}x^2 + 1 \quad \Rightarrow$

$-y = \ln\left(1 - \frac{1}{2}x^2\right) \quad \Rightarrow \quad y = -\ln\left(1 - \frac{1}{2}x^2\right).$

14. $\dfrac{dP}{dt} = \sqrt{Pt} \quad \Rightarrow \quad dP/\sqrt{P} = \sqrt{t}\,dt \quad \Rightarrow \quad \int P^{-1/2}\,dP = \int t^{1/2}\,dt \quad \Rightarrow \quad 2P^{1/2} = \frac{2}{3}t^{3/2} + C.$

$P(1) = 2 \quad \Rightarrow \quad 2\sqrt{2} = \frac{2}{3} + C \quad \Rightarrow \quad C = 2\sqrt{2} - \frac{2}{3}$, so $2P^{1/2} = \frac{2}{3}t^{3/2} + 2\sqrt{2} - \frac{2}{3} \quad \Rightarrow \quad \sqrt{P} = \frac{1}{3}t^{3/2} + \sqrt{2} - \frac{1}{3} \quad \Rightarrow$

$P = \left(\frac{1}{3}t^{3/2} + \sqrt{2} - \frac{1}{3}\right)^2.$

15. $\dfrac{dA}{dr} = Ab^2\cos br \quad \Rightarrow \quad \dfrac{dA}{A} = b^2\cos br\,dr \quad \Rightarrow \quad \displaystyle\int \frac{1}{A}\,dA = \int b^2\cos br\,dr \quad \Rightarrow \quad \ln|A| = b\sin br + C.$

$A(0) = b^3 \quad \Rightarrow \quad \ln\left|b^3\right| = b\sin 0 + C \quad \Rightarrow \quad C = \ln\left|b^3\right|$, so $\ln|A| = b\sin br + \ln\left|b^3\right| \quad \Rightarrow$

$|A| = e^{b\sin br + \ln\left|b^3\right|} = e^{b\sin br}e^{\ln\left|b^3\right|} = \left|b^3\right|e^{b\sin br} \quad \Rightarrow \quad A = \pm b^3e^{b\sin br}$. Since $A(0) = b^3$, the solution is

$A = b^3e^{b\sin br}$.

16. $x^2\,y' = k\sec y \quad \Rightarrow \quad x^2\,\dfrac{dy}{dx} = \dfrac{k}{\cos y} \quad \Rightarrow \quad \cos y\,dy = k\,\dfrac{dx}{x^2} \quad \Rightarrow \quad \int \cos y\,dy = \int kx^{-2}dx \quad \Rightarrow \quad \sin y = -kx^{-1} + C.$

$y(1) = \frac{\pi}{6} \quad \Rightarrow \quad \sin\frac{\pi}{6} = -k(1)^{-1} + C \quad \Rightarrow \quad \frac{1}{2} = -k + C \quad \Rightarrow \quad C = \frac{1}{2} + k$, so

$\sin y = -\dfrac{k}{x} + \dfrac{1}{2} + k \quad \Rightarrow \quad y = \sin^{-1}\left(-\dfrac{k}{x} + \dfrac{1}{2} + k\right).$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\dfrac{du}{dt} = \dfrac{2t + \sec^2 t}{2u}$, $u(0) = -5$. $\int 2u\,du = \int \left(2t + \sec^2 t\right) dt \;\Rightarrow\; u^2 = t^2 + \tan t + C$,

where $[u(0)]^2 = 0^2 + \tan 0 + C \;\Rightarrow\; C = (-5)^2 = 25$. Therefore, $u^2 = t^2 + \tan t + 25$, so $u = \pm\sqrt{t^2 + \tan t + 25}$.

Since $u(0) = -5 < 0$, we must have $u = -\sqrt{t^2 + \tan t + 25}$.

18. $x + 3y^2\sqrt{x^2+1}\,\dfrac{dy}{dx} = 0 \;\Rightarrow\; 3y^2\sqrt{x^2+1}\,\dfrac{dy}{dx} = -x \;\Rightarrow\; 3y^2\,dy = \dfrac{-x}{\sqrt{x^2+1}}\,dx \;\Rightarrow$

$\int 3y^2\,dy = \int -x(x^2+1)^{-1/2}\,dx \;\Rightarrow\; y^3 = -(x^2+1)^{1/2} + C$. $y(0) = 1 \;\Rightarrow\; 1^3 = -(0^2+1)^{1/2} + C \;\Rightarrow$

$C = 2$, so $y^3 = -(x^2+1)^{1/2} + 2 \;\Rightarrow\; y = (2 - \sqrt{x^2+1})^{1/3}$.

19. $x\ln x = y\left(1 + \sqrt{3+y^2}\right)y'$, $y(1) = 1$. $\int x\ln x\,dx = \int \left(y + y\sqrt{3+y^2}\right)\,dy \;\Rightarrow\; \frac{1}{2}x^2\ln x - \int \frac{1}{2}x\,dx$

[use parts with $u = \ln x$, $dv = x\,dx$] $= \frac{1}{2}y^2 + \frac{1}{3}(3+y^2)^{3/2} \;\Rightarrow\; \frac{1}{2}x^2\ln x - \frac{1}{4}x^2 + C = \frac{1}{2}y^2 + \frac{1}{3}(3+y^2)^{3/2}$.

Now $y(1) = 1 \;\Rightarrow\; 0 - \frac{1}{4} + C = \frac{1}{2} + \frac{1}{3}(4)^{3/2} \;\Rightarrow\; C = \frac{1}{2} + \frac{8}{3} + \frac{1}{4} = \frac{41}{12}$, so

$\frac{1}{2}x^2\ln x - \frac{1}{4}x^2 + \frac{41}{12} = \frac{1}{2}y^2 + \frac{1}{3}(3+y^2)^{3/2}$. We do not solve explicitly for y.

20. $\dfrac{dy}{dx} = \dfrac{x\sin x}{y} \;\Rightarrow\; y\,dy = x\sin x\,dx \;\Rightarrow\; \int y\,dy = \int x\sin x\,dx \;\Rightarrow\; \frac{1}{2}y^2 = -x\cos x + \sin x + C$ [by parts].

$y(0) = -1 \;\Rightarrow\; \frac{1}{2}(-1)^2 = -0\cos 0 + \sin 0 + C \;\Rightarrow\; C = \frac{1}{2}$, so $\frac{1}{2}y^2 = -x\cos x + \sin x + \frac{1}{2} \;\Rightarrow$

$y^2 = -2x\cos x + 2\sin x + 1 \;\Rightarrow\; y = -\sqrt{-2x\cos x + 2\sin x + 1}$ since $y(0) = -1 < 0$.

21. $\dfrac{dy}{dx} = \dfrac{x}{y} \;\Rightarrow\; y\,dy = x\,dx \;\Rightarrow\; \int y\,dy = \int x\,dx \;\Rightarrow\; \frac{1}{2}y^2 = \frac{1}{2}x^2 + C$. $y(0) = 2 \;\Rightarrow\; \frac{1}{2}(2)^2 = \frac{1}{2}(0)^2 + C \;\Rightarrow$

$C = 2$, so $\frac{1}{2}y^2 = \frac{1}{2}x^2 + 2 \;\Rightarrow\; y^2 = x^2 + 4 \;\Rightarrow\; y = \sqrt{x^2+4}$ since $y(0) = 2 > 0$.

22. $f'(x) = x\,f(x) - x \;\Rightarrow\; \dfrac{dy}{dx} = xy - x \;\Rightarrow\; \dfrac{dy}{dx} = x(y-1) \;\Rightarrow\; \dfrac{dy}{y-1} = x\,dx \quad [y \neq 1] \;\Rightarrow$

$\displaystyle\int \frac{dy}{y-1} = \int x\,dx \;\Rightarrow\; \ln|y-1| = \frac{1}{2}x^2 + C$. $f(0) = 2 \;\Rightarrow\; \ln|2-1| = \frac{1}{2}(0)^2 + C \;\Rightarrow\; C = 0$, so

$\ln|y-1| = \frac{1}{2}x^2 \;\Rightarrow\; |y-1| = e^{x^2/2} \;\Rightarrow\; y - 1 = e^{x^2/2}$ [since $f(0) = 2$] $\;\Rightarrow\; y = e^{x^2/2} + 1$.

23. $u = x + y \;\Rightarrow\; \dfrac{d}{dx}(u) = \dfrac{d}{dx}(x+y) \;\Rightarrow\; \dfrac{du}{dx} = 1 + \dfrac{dy}{dx}$, but $\dfrac{dy}{dx} = x + y = u$, so $\dfrac{du}{dx} = 1 + u \;\Rightarrow$

$\dfrac{du}{1+u} = dx \quad [u \neq -1] \;\Rightarrow\; \displaystyle\int \frac{du}{1+u} = \int dx \;\Rightarrow\; \ln|1+u| = x + C \;\Rightarrow\; |1+u| = e^{x+C} \;\Rightarrow$

$1 + u = \pm e^C e^x \;\Rightarrow\; u = \pm e^C e^x - 1 \;\Rightarrow\; x + y = \pm e^C e^x - 1 \;\Rightarrow\; y = Ke^x - x - 1$, where $K = \pm e^C \neq 0$.

If $u = -1$, then $-1 = x + y \;\Rightarrow\; y = -x - 1$, which is just $y = Ke^x - x - 1$ with $K = 0$. Thus, the general solution is $y = Ke^x - x - 1$, where $K \in \mathbb{R}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $xy' = y + xe^{y/x} \quad \Rightarrow \quad y' = y/x + e^{y/x} \quad \Rightarrow \quad \dfrac{dy}{dx} = v + e^v$. Also, $v = y/x \quad \Rightarrow \quad xv = y \quad \Rightarrow \quad \dfrac{dy}{dx} = x\dfrac{dv}{dx} + v$,

so $v + e^v = x\dfrac{dv}{dx} + v \quad \Rightarrow \quad \dfrac{dv}{e^v} = \dfrac{dx}{x} \quad [x \neq 0] \quad \Rightarrow \quad \displaystyle\int \frac{dv}{e^v} = \int \frac{dx}{x} \quad \Rightarrow \quad -e^{-v} = \ln|x| + C \quad \Rightarrow$

$e^{-v} = -\ln|x| - C \quad \Rightarrow \quad -v = \ln(-\ln|x| - C) \quad \Rightarrow \quad y/x = -\ln(-\ln|x| - C) \quad \Rightarrow \quad y = -x\ln(-\ln|x| - C)$.

25. (a) $y' = 2x\sqrt{1-y^2} \quad \Rightarrow \quad \dfrac{dy}{dx} = 2x\sqrt{1-y^2} \quad \Rightarrow \quad \dfrac{dy}{\sqrt{1-y^2}} = 2x\,dx \quad \Rightarrow \quad \displaystyle\int \frac{dy}{\sqrt{1-y^2}} = \int 2x\,dx \quad \Rightarrow$

$\sin^{-1} y = x^2 + C$ for $-\frac{\pi}{2} \leq x^2 + C \leq \frac{\pi}{2}$.

(b) $y(0) = 0 \quad \Rightarrow \quad \sin^{-1} 0 = 0^2 + C \quad \Rightarrow \quad C = 0$,

so $\sin^{-1} y = x^2$ and $y = \sin(x^2)$ for $-\sqrt{\pi/2} \leq x \leq \sqrt{\pi/2}$.

(c) For $\sqrt{1-y^2}$ to be a real number, we must have $-1 \leq y \leq 1$; that is, $-1 \leq y(0) \leq 1$. Thus, the initial-value problem $y' = 2x\sqrt{1-y^2}$, $y(0) = 2$ does *not* have a solution.

26. $e^{-y}y' + \cos x = 0 \quad \Leftrightarrow \quad \int e^{-y}\,dy = -\int \cos x\,dx \quad \Leftrightarrow \quad -e^{-y} = -\sin x + C_1 \quad \Leftrightarrow \quad y = -\ln(\sin x + C)$. The solution is periodic, with period 2π. Note that for $C > 1$, the domain of the solution is $\mathbb{R}$, but for $-1 < C \leq 1$ it is only defined on the intervals where $\sin x + C > 0$, and it is meaningless for $C \leq -1$, since then $\sin x + C \leq 0$, and the logarithm is undefined.

For $-1 < C < 1$, the solution curve consists of concave-up pieces separated by intervals on which the solution is not defined (where $\sin x + C \leq 0$). For $C = 1$, the solution curve consists of concave-up pieces separated by vertical asymptotes at the points where $\sin x + C = 0 \quad \Leftrightarrow \quad \sin x = -1$. For $C > 1$, the curve is continuous, and as C increases, the graph moves downward, and the amplitude of the oscillations decreases.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. $\dfrac{dy}{dx} = \dfrac{\sin x}{\sin y}$, $y(0) = \dfrac{\pi}{2}$. So $\int \sin y\,dy = \int \sin x\,dx \quad \Leftrightarrow \quad -\cos y = -\cos x + C \quad \Leftrightarrow \quad \cos y = \cos x - C$. From the initial condition, we need $\cos\frac{\pi}{2} = \cos 0 - C \quad \Rightarrow \quad 0 = 1 - C \quad \Rightarrow \quad C = 1$, so the solution is $\cos y = \cos x - 1$. Note that we cannot take $\cos^{-1}$ of both sides, since that would unnecessarily restrict the solution to the case where $-1 \le \cos x - 1 \quad \Leftrightarrow$ $0 \le \cos x$, as $\cos^{-1}$ is defined only on $[-1, 1]$. Instead we plot the graph using Maple's `plots[implicitplot]` or Mathematica's `Plot[Evaluate[···]]`.

28. $\dfrac{dy}{dx} = \dfrac{x\sqrt{x^2+1}}{ye^y} \quad \Leftrightarrow \quad \int ye^y\,dy = \int x\sqrt{x^2+1}\,dx$. We use parts on the LHS with $u = y$, $dv = e^y\,dy$, and on the RHS we use the substitution $z = x^2 + 1$, so $dz = 2x\,dx$. The equation becomes $ye^y - \int e^y\,dy = \frac{1}{2}\int \sqrt{z}\,dz \quad \Leftrightarrow$ $e^y(y-1) = \frac{1}{3}(x^2+1)^{3/2} + C$, so we see that the curves are symmetric about the y-axis. Every point (x, y) in the plane lies on one of the curves, namely the one for which $C = (y-1)e^y - \frac{1}{3}(x^2+1)^{3/2}$. For example, along the y-axis, $C = (y-1)e^y - \frac{1}{3}$, so the origin lies on the curve with $C = -\frac{4}{3}$. We use Maple's `plots[implicitplot]` command or `Plot[Evaluate[···]]` in Mathematica to plot the solution curves for various values of C.

It seems that the transitional values of C are $-\frac{4}{3}$ and $-\frac{1}{3}$. For $C < -\frac{4}{3}$, the graph consists of left and right branches. At $C = -\frac{4}{3}$, the two branches become connected at the origin, and as C increases, the graph splits into top and bottom branches. At $C = -\frac{1}{3}$, the bottom half disappears. As C increases further, the graph moves upward, but doesn't change shape much.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. (a), (c)

(b) $y' = y^2 \quad \Rightarrow \quad \dfrac{dy}{dx} = y^2 \quad \Rightarrow \quad \displaystyle\int y^{-2}\,dy = \int dx \quad \Rightarrow$

$-y^{-1} = x + C \quad \Rightarrow \quad \dfrac{1}{y} = -x - C \quad \Rightarrow$

$y = \dfrac{1}{K - x}$, where $K = -C$. $y = 0$ is also a solution.

30. (a), (c)

(b) $y' = xy \quad \Rightarrow \quad \dfrac{dy}{dx} = xy \quad \Rightarrow \quad \displaystyle\int \frac{dy}{y} = \int x\,dx \quad \Rightarrow$

$\ln|y| = \frac{1}{2}x^2 + C \quad \Rightarrow \quad |y| = e^{x^2/2 + C} = e^{x^2/2}e^C \quad \Rightarrow$

$y = Ke^{x^2/2}$, where $K = \pm e^C$. Taking $K = 0$ gives us the solution $y = 0$.

31. The curves $x^2 + 2y^2 = k^2$ form a family of ellipses with major axis on the x-axis. Differentiating gives

$\dfrac{d}{dx}(x^2 + 2y^2) = \dfrac{d}{dx}(k^2) \quad \Rightarrow \quad 2x + 4yy' = 0 \quad \Rightarrow \quad 4yy' = -2x \quad \Rightarrow \quad y' = \dfrac{-x}{2y}$. Thus, the slope of the tangent line at any point (x, y) on one of the ellipses is $y' = \dfrac{-x}{2y}$, so the orthogonal trajectories must satisfy $y' = \dfrac{2y}{x} \quad \Leftrightarrow \quad \dfrac{dy}{dx} = \dfrac{2y}{x} \quad \Leftrightarrow \quad \dfrac{dy}{y} = 2 = \dfrac{dx}{x} \quad \Leftrightarrow$

$\displaystyle\int \frac{dy}{y} = 2\int \frac{dx}{x} \quad \Leftrightarrow \quad \ln|y| = 2\ln|x| + C_1 \quad \Leftrightarrow \quad \ln|y| = \ln|x|^2 + C_1 \quad \Leftrightarrow$

$|y| = e^{\ln x^2 + C_1} \quad \Leftrightarrow \quad y = \pm x^2 \cdot e^{C_1} = Cx^2$. This is a family of parabolas.

32. The curves $y^2 = kx^3$ form a family of power functions. Differentiating gives $\dfrac{d}{dx}(y^2) = \dfrac{d}{dx}(kx^3) \quad \Rightarrow \quad 2yy' = 3kx^2 \quad \Rightarrow$

$y' = \dfrac{3kx^2}{2y} = \dfrac{3(y^2/x^3)x^2}{2y} = \dfrac{3y}{2x}$, the slope of the tangent line at (x, y) on one of the curves. Thus, the orthogonal trajectories must satisfy $y' = -\dfrac{2x}{3y} \quad \Leftrightarrow \quad \dfrac{dy}{dx} = -\dfrac{2x}{3y} \quad \Leftrightarrow$

$3y\,dy = -2x\,dx \quad \Leftrightarrow \quad \int 3y\,dy = \int -2x\,dx \quad \Leftrightarrow \quad \frac{3}{2}y^2 = -x^2 + C_1 \quad \Leftrightarrow$

$3y^2 = -2x^2 + C_2 \quad \Leftrightarrow \quad 2x^2 + 3y^2 = C$. This is a family of ellipses.

33. The curves $y = k/x$ form a family of hyperbolas with asymptotes $x = 0$ and $y = 0$. Differentiating gives

$\dfrac{d}{dx}(y) = \dfrac{d}{dx}\left(\dfrac{k}{x}\right) \quad \Rightarrow \quad y' = -\dfrac{k}{x^2} \quad \Rightarrow \quad y' = -\dfrac{xy}{x^2}$ [since $y = k/x \quad \Rightarrow \quad xy = k$] $\Rightarrow \quad y' = -\dfrac{y}{x}$. Thus, the slope

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

of the tangent line at any point (x, y) on one of the hyperbolas is $y' = -y/x$, so the orthogonal trajectories must satisfy $y' = x/y \quad \Leftrightarrow \quad \dfrac{dy}{dx} = \dfrac{x}{y} \quad \Leftrightarrow$ $y\,dy = x\,dx \quad \Leftrightarrow \quad \int y\,dy = \int x\,dx \quad \Leftrightarrow \quad \frac{1}{2}y^2 = \frac{1}{2}x^2 + C_1 \quad \Leftrightarrow$ $y^2 = x^2 + C_2 \quad \Leftrightarrow \quad x^2 - y^2 = C$. This is a family of hyperbolas with asymptotes $y = \pm x$.

34. The curves $y = 1/(x + k)$ form a family of hyperbolas with asymptotes $x = -k$ and $y = 0$. Differentiating gives $\dfrac{d}{dx}(y) = \dfrac{d}{dx}\left(\dfrac{1}{x+k}\right) \quad \Rightarrow \quad y' = -\dfrac{1}{(x+k)^2} \quad \Rightarrow \quad y' = -y^2$ [since $y = 1/(x + k)$]. Thus, the slope of the tangent line at any point (x, y) on one of the hyperbolas is $y' = -y^2$, so the orthogonal trajectories must satisfy $y' = 1/y^2 \quad \Leftrightarrow \quad \dfrac{dy}{dx} = \dfrac{1}{y^2} \quad \Leftrightarrow$ $y^2\,dy = dx \quad \Leftrightarrow \quad \int y^2\,dy = \int dx \quad \Leftrightarrow \quad \frac{1}{3}y^3 = x + C_1 \quad \Leftrightarrow$ $y^3 = 3x + C \quad \Leftrightarrow \quad y = (3x + C)^{1/3}$. This is a family of cube root functions with vertical tangents on the x-axis $\quad [y = 0]$.

35. $y(x) = 2 + \displaystyle\int_2^x [t - ty(t)]\,dt \quad \Rightarrow \quad y'(x) = x - xy(x)$ [by FTC 1] $\quad \Rightarrow \quad \dfrac{dy}{dx} = x(1 - y) \quad \Rightarrow$

$\displaystyle\int \frac{dy}{1-y} = \int x\,dx \quad \Rightarrow \quad -\ln|1 - y| = \tfrac{1}{2}x^2 + C$. Letting $x = 2$ in the original integral equation gives us $y(2) = 2 + 0 = 2$. Thus, $-\ln|1 - 2| = \frac{1}{2}(2)^2 + C \quad \Rightarrow \quad 0 = 2 + C \quad \Rightarrow \quad C = -2$.

Thus, $-\ln|1 - y| = \frac{1}{2}x^2 - 2 \quad \Rightarrow \quad \ln|1 - y| = 2 - \frac{1}{2}x^2 \quad \Rightarrow \quad |1 - y| = e^{2 - x^2/2} \quad \Rightarrow$

$1 - y = \pm e^{2 - x^2/2} \quad \Rightarrow \quad y = 1 + e^{2 - x^2/2}$ $[y(2) = 2]$.

36. $y(x) = 2 + \displaystyle\int_1^x \frac{dt}{ty(t)}, x > 0 \quad \Rightarrow \quad y'(x) = \frac{1}{xy(x)} \quad \Rightarrow \quad \frac{dy}{dx} = \frac{1}{xy} \quad \Rightarrow \quad \int y\,dy = \int \frac{1}{x}\,dx \quad \Rightarrow$

$\frac{1}{2}y^2 = \ln x + C$ $[x > 0]$. Letting $x = 1$ in the original integral equation gives us $y(1) = 2 + 0 = 2$.

Thus, $\frac{1}{2}(2)^2 = \ln 1 + C \quad \Rightarrow \quad C = 2$. $\frac{1}{2}y^2 = \ln x + 2 \quad \Rightarrow \quad y^2 = 2\ln x + 4$ $[> 0] \quad \Rightarrow \quad y = \sqrt{2\ln x + 4}$.

37. $y(x) = 4 + \displaystyle\int_0^x 2t\sqrt{y(t)}\,dt \quad \Rightarrow \quad y'(x) = 2x\sqrt{y(x)} \quad \Rightarrow \quad \frac{dy}{dx} = 2x\sqrt{y} \quad \Rightarrow \quad \int \frac{dy}{\sqrt{y}} = \int 2x\,dx \quad \Rightarrow$

$2\sqrt{y} = x^2 + C$. Letting $x = 0$ in the original integral equation gives us $y(0) = 4 + 0 = 4$.

Thus, $2\sqrt{4} = 0^2 + C \quad \Rightarrow \quad C = 4$. $2\sqrt{y} = x^2 + 4 \quad \Rightarrow \quad \sqrt{y} = \frac{1}{2}x^2 + 2 \quad \Rightarrow \quad y = \left(\frac{1}{2}x^2 + 2\right)^2$.

38. $(t^2 + 1)f'(t) + [f(t)]^2 + 1 = 0 \quad \Rightarrow \quad (t^2 + 1)\dfrac{dy}{dt} + y^2 + 1 = 0 \quad \Rightarrow \quad \dfrac{dy}{dt} = \dfrac{-y^2 - 1}{t^2 + 1} \quad \Rightarrow$

$\displaystyle\int \frac{dy}{y^2 + 1} = -\int \frac{dt}{t^2 + 1} \quad \Rightarrow \quad \arctan y = -\arctan t + C \quad \Rightarrow \quad \arctan t + \arctan y = C \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\tan(\arctan t + \arctan y) = \tan C \quad \Rightarrow \quad \dfrac{\tan(\arctan t) + \tan(\arctan y)}{1 - \tan(\arctan t)\,\tan(\arctan y)} = \tan C \quad \Rightarrow \quad \dfrac{t+y}{1-ty} = \tan C = k \quad \Rightarrow$

$t + y = k - kty \quad \Rightarrow \quad y + kty = k - t \quad \Rightarrow \quad y(1+kt) = k - t \quad \Rightarrow \quad f(t) = y = \dfrac{k-t}{1+kt}$.

Since $f(3) = 2 = \dfrac{k-3}{1+3k} \quad \Rightarrow \quad 2 + 6k = k - 3 \quad \Rightarrow \quad 5k = -5 \quad \Rightarrow \quad k = -1$, we have $y = \dfrac{-1-t}{1+(-1)t} = \dfrac{t+1}{t-1}$.

39. From Exercise 9.2.27, $\dfrac{dQ}{dt} = 12 - 4Q \quad \Leftrightarrow \quad \displaystyle\int \frac{dQ}{12-4Q} = \int dt \quad \Leftrightarrow \quad -\tfrac{1}{4}\ln|12-4Q| = t + C \quad \Leftrightarrow$

$\ln|12-4Q| = -4t - 4C \quad \Leftrightarrow \quad |12-4Q| = e^{-4t-4C} \quad \Leftrightarrow \quad 12 - 4Q = Ke^{-4t} \quad [K = \pm e^{-4C}] \quad \Leftrightarrow$

$4Q = 12 - Ke^{-4t} \quad \Leftrightarrow \quad Q = 3 - Ae^{-4t} \quad [A = K/4]$. $Q(0) = 0 \quad \Leftrightarrow \quad 0 = 3 - A \quad \Leftrightarrow \quad A = 3 \quad \Leftrightarrow$

$Q(t) = 3 - 3e^{-4t}$. As $t \to \infty$, $Q(t) \to 3 - 0 = 3$ (the limiting value).

40. From Exercise 9.2.28, $\dfrac{dy}{dt} = -\frac{1}{50}(y-20) \quad \Leftrightarrow \quad \displaystyle\int \frac{dy}{y-20} = \int \left(-\tfrac{1}{50}\right) dt \quad \Leftrightarrow \quad \ln|y-20| = -\tfrac{1}{50}t + C \quad \Leftrightarrow$

$y - 20 = Ke^{-t/50} \quad \Leftrightarrow \quad y(t) = Ke^{-t/50} + 20$. $y(0) = 95 \quad \Leftrightarrow \quad 95 = K + 20 \quad \Leftrightarrow \quad K = 75 \quad \Leftrightarrow$

$y(t) = 75e^{-t/50} + 20$.

41. $\dfrac{dP}{dt} = k(M-P) \quad \Leftrightarrow \quad \displaystyle\int \frac{dP}{P-M} = \int (-k)\,dt \quad \Leftrightarrow \quad \ln|P-M| = -kt + C \quad \Leftrightarrow \quad |P-M| = e^{-kt+C} \quad \Leftrightarrow$

$P - M = Ae^{-kt} \quad [A = \pm e^C] \quad \Leftrightarrow \quad P = M + Ae^{-kt}$. If we assume that performance is at level 0 when $t = 0$, then

$P(0) = 0 \quad \Leftrightarrow \quad 0 = M + A \quad \Leftrightarrow \quad A = -M \quad \Leftrightarrow \quad P(t) = M - Me^{-kt}$. $\displaystyle\lim_{t\to\infty} P(t) = M - M \cdot 0 = M$.

42. (a) $\dfrac{dx}{dt} = k(a-x)(b-x),\ a \neq b$. Using partial fractions, $\dfrac{1}{(a-x)(b-x)} = \dfrac{1/(b-a)}{a-x} - \dfrac{1/(b-a)}{b-x}$, so

$$\int \frac{dx}{(a-x)(b-x)} = \int k\,dt \quad \Rightarrow \quad \frac{1}{b-a}\left(-\ln|a-x| + \ln|b-x|\right) = kt + C \quad \Rightarrow \quad \ln\left|\frac{b-x}{a-x}\right| = (b-a)(kt+C).$$

The concentrations [A] $= a - x$ and [B] $= b - x$ cannot be negative, so $\dfrac{b-x}{a-x} \geq 0$ and $\left|\dfrac{b-x}{a-x}\right| = \dfrac{b-x}{a-x}$.

We now have $\ln\left(\dfrac{b-x}{a-x}\right) = (b-a)(kt+C)$. Since $x(0) = 0$, we get $\ln\left(\dfrac{b}{a}\right) = (b-a)C$. Hence,

$$\ln\left(\frac{b-x}{a-x}\right) = (b-a)kt + \ln\left(\frac{b}{a}\right) \Rightarrow \frac{b-x}{a-x} = \frac{b}{a}e^{(b-a)kt} \Rightarrow x = \frac{b[e^{(b-a)kt}-1]}{be^{(b-a)kt}/a - 1} = \frac{ab[e^{(b-a)kt}-1]}{be^{(b-a)kt} - a}\ \frac{\text{moles}}{\text{L}}.$$

(b) If $b = a$, then $\dfrac{dx}{dt} = k(a-x)^2$, so $\displaystyle\int \frac{dx}{(a-x)^2} = \int k\,dt$ and $\dfrac{1}{a-x} = kt + C$. Since $x(0) = 0$, we get $C = \dfrac{1}{a}$.

Thus, $a - x = \dfrac{1}{kt + 1/a}$ and $x = a - \dfrac{a}{akt+1} = \dfrac{a^2kt}{akt+1}\,\dfrac{\text{moles}}{\text{L}}$. Suppose $x = [\text{C}] = \frac{1}{2}a$ when $t = 20$. Then

$x(20) = \dfrac{a}{2} \quad \Rightarrow \quad \dfrac{a}{2} = \dfrac{20a^2k}{20ak+1} \quad \Rightarrow \quad 40a^2k = 20a^2k + a \quad \Rightarrow \quad 20a^2k = a \quad \Rightarrow \quad k = \dfrac{1}{20a}$, so

$x = \dfrac{a^2t/(20a)}{1 + at/(20a)} = \dfrac{at/20}{1 + t/20} = \dfrac{at}{t+20}\,\dfrac{\text{moles}}{\text{L}}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. (a) If $a = b$, then $\dfrac{dx}{dt} = k(a-x)(b-x)^{1/2}$ becomes $\dfrac{dx}{dt} = k(a-x)^{3/2} \;\Rightarrow\; (a-x)^{-3/2}\,dx = k\,dt \;\Rightarrow$

$\int (a-x)^{-3/2}\,dx = \int k\,dt \;\Rightarrow\; 2(a-x)^{-1/2} = kt + C$ [by substitution] $\;\Rightarrow\; \dfrac{2}{kt+C} = \sqrt{a-x} \;\Rightarrow$

$\left(\dfrac{2}{kt+C}\right)^2 = a - x \;\Rightarrow\; x(t) = a - \dfrac{4}{(kt+C)^2}$. The initial concentration of HBr is 0, so $x(0) = 0 \;\Rightarrow$

$0 = a - \dfrac{4}{C^2} \;\Rightarrow\; \dfrac{4}{C^2} = a \;\Rightarrow\; C^2 = \dfrac{4}{a} \;\Rightarrow\; C = 2/\sqrt{a}$ [C is positive since $kt + C = 2(a-x)^{-1/2} > 0$].

Thus, $x(t) = a - \dfrac{4}{\left(kt + 2/\sqrt{a}\,\right)^2}$.

(b) $\dfrac{dx}{dt} = k(a-x)(b-x)^{1/2} \;\Rightarrow\; \dfrac{dx}{(a-x)\sqrt{b-x}} = k\,dt \;\Rightarrow\; \displaystyle\int \frac{dx}{(a-x)\sqrt{b-x}} = \int k\,dt \quad (\star)$.

From the hint, $u = \sqrt{b-x} \;\Rightarrow\; u^2 = b - x \;\Rightarrow\; 2u\,du = -dx$, so

$$\int \frac{dx}{(a-x)\sqrt{b-x}} = \int \frac{-2u\,du}{[a-(b-u^2)]u} = -2\int \frac{du}{a-b+u^2} = -2\int \frac{du}{\left(\sqrt{a-b}\,\right)^2 + u^2}$$

$$\overset{17}{=} -2\left(\frac{1}{\sqrt{a-b}}\tan^{-1}\frac{u}{\sqrt{a-b}}\right)$$

So $(\star)$ becomes $\dfrac{-2}{\sqrt{a-b}}\tan^{-1}\dfrac{\sqrt{b-x}}{\sqrt{a-b}} = kt + C$. Now $x(0) = 0 \;\Rightarrow\; C = \dfrac{-2}{\sqrt{a-b}}\tan^{-1}\dfrac{\sqrt{b}}{\sqrt{a-b}}$ and we have

$\dfrac{-2}{\sqrt{a-b}}\tan^{-1}\dfrac{\sqrt{b-x}}{\sqrt{a-b}} = kt - \dfrac{2}{\sqrt{a-b}}\tan^{-1}\dfrac{\sqrt{b}}{\sqrt{a-b}} \;\Rightarrow\; \dfrac{2}{\sqrt{a-b}}\left(\tan^{-1}\sqrt{\dfrac{b}{a-b}} - \tan^{-1}\sqrt{\dfrac{b-x}{a-b}}\right) = kt \;\Rightarrow$

$t(x) = \dfrac{2}{k\sqrt{a-b}}\left(\tan^{-1}\sqrt{\dfrac{b}{a-b}} - \tan^{-1}\sqrt{\dfrac{b-x}{a-b}}\right)$.

44. If $S = \dfrac{dT}{dr}$, then $\dfrac{dS}{dr} = \dfrac{d^2T}{dr^2}$. The differential equation $\dfrac{d^2T}{dr^2} + \dfrac{2}{r}\dfrac{dT}{dr} = 0$ can be written as $\dfrac{dS}{dr} + \dfrac{2}{r}S = 0$. Thus,

$\dfrac{dS}{dr} = \dfrac{-2S}{r} \;\Rightarrow\; \dfrac{dS}{S} = -\dfrac{2}{r}\,dr \;\Rightarrow\; \displaystyle\int \frac{1}{S}\,dS = \int -\frac{2}{r}\,dr \;\Rightarrow\; \ln|S| = -2\ln|r| + C$. Assuming $S = dT/dr > 0$

and $r > 0$, we have $S = e^{-2\ln r + C} = e^{\ln r^{-2}}e^C = r^{-2}k \;\; [k = e^C] \;\Rightarrow\; S = \dfrac{1}{r^2}k \;\Rightarrow\; \dfrac{dT}{dr} = \dfrac{1}{r^2}k \;\Rightarrow$

$dT = \dfrac{1}{r^2}k\,dr \;\Rightarrow\; \displaystyle\int dT = \int \frac{1}{r^2}k\,dr \;\Rightarrow\; T(r) = -\frac{k}{r} + A$.

$T(1) = 15 \;\Rightarrow\; 15 = -k + A$ **(1)** and $T(2) = 25 \;\Rightarrow\; 25 = -\frac{1}{2}k + A$ **(2)**.

Now solve for k and A: -2**(2)** $+$ **(1)** $\;\Rightarrow\; -35 = -A$, so $A = 35$ and $k = 20$, and $T(r) = -20/r + 35$.

45. (a) $\dfrac{dC}{dt} = r - kC \;\Rightarrow\; \dfrac{dC}{dt} = -(kC - r) \;\Rightarrow\; \displaystyle\int \frac{dC}{kC-r} = \int -dt \;\Rightarrow\; (1/k)\ln|kC - r| = -t + M_1 \;\Rightarrow$

$\ln|kC - r| = -kt + M_2 \;\Rightarrow\; |kC - r| = e^{-kt+M_2} \;\Rightarrow\; kC - r = M_3e^{-kt} \;\Rightarrow\; kC = M_3e^{-kt} + r \;\Rightarrow$

$C(t) = M_4e^{-kt} + r/k$. $C(0) = C_0 \;\Rightarrow\; C_0 = M_4 + r/k \;\Rightarrow\; M_4 = C_0 - r/k \;\Rightarrow$

$C(t) = (C_0 - r/k)e^{-kt} + r/k$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) If $C_0 < r/k$, then $C_0 - r/k < 0$ and the formula for $C(t)$ shows that $C(t)$ increases and $\lim_{t\to\infty} C(t) = r/k$.

As t increases, the formula for $C(t)$ shows how the role of C_0 steadily diminishes as that of r/k increases.

46. (a) Use 1 billion dollars as the x-unit and 1 day as the t-unit. Initially, there is \$10 billion of old currency in circulation, so all of the \$50 million returned to the banks is old. At time t, the amount of new currency is $x(t)$ billion dollars, so $10 - x(t)$ billion dollars of currency is old. The fraction of circulating money that is old is $[10 - x(t)]/10$, and the amount of old currency being returned to the banks each day is $\dfrac{10 - x(t)}{10}\,0.05$ billion dollars. This amount of new currency per day is introduced into circulation, so $\dfrac{dx}{dt} = \dfrac{10 - x}{10} \cdot 0.05 = 0.005(10 - x)$ billion dollars per day.

(b) $\dfrac{dx}{10 - x} = 0.005\,dt \quad\Rightarrow\quad \dfrac{-dx}{10 - x} = -0.005\,dt \quad\Rightarrow\quad \ln(10 - x) = -0.005t + c \quad\Rightarrow\quad 10 - x = Ce^{-0.005t}$,

where $C = e^c \quad\Rightarrow\quad x(t) = 10 - Ce^{-0.005t}$. From $x(0) = 0$, we get $C = 10$, so $x(t) = 10(1 - e^{-0.005t})$.

(c) The new bills make up 90% of the circulating currency when $x(t) = 0.9 \cdot 10 = 9$ billion dollars.

$9 = 10(1 - e^{-0.005t}) \quad\Rightarrow\quad 0.9 = 1 - e^{-0.005t} \quad\Rightarrow\quad e^{-0.005t} = 0.1 \quad\Rightarrow\quad -0.005t = -\ln 10 \quad\Rightarrow$

$t = 200 \ln 10 \approx 460.517$ days ≈ 1.26 years.

47. (a) Let $y(t)$ be the amount of salt (in kg) after t minutes. Then $y(0) = 15$. The amount of liquid in the tank is 1000 L at all times, so the concentration at time t (in minutes) is $y(t)/1000$ kg/L and $\dfrac{dy}{dt} = -\left[\dfrac{y(t)}{1000}\,\dfrac{\text{kg}}{\text{L}}\right]\left(10\,\dfrac{\text{L}}{\text{min}}\right) = -\dfrac{y(t)}{100}\,\dfrac{\text{kg}}{\text{min}}$.

$\displaystyle\int \frac{dy}{y} = -\frac{1}{100}\int dt \quad\Rightarrow\quad \ln y = -\frac{t}{100} + C$, and $y(0) = 15 \quad\Rightarrow\quad \ln 15 = C$, so $\ln y = \ln 15 - \dfrac{t}{100}$.

It follows that $\ln\left(\dfrac{y}{15}\right) = -\dfrac{t}{100}$ and $\dfrac{y}{15} = e^{-t/100}$, so $y = 15e^{-t/100}$ kg.

(b) After 20 minutes, $y = 15e^{-20/100} = 15e^{-0.2} \approx 12.3$ kg.

48. Let $y(t)$ be the amount of carbon dioxide in the room after t minutes. Then $y(0) = 0.0015(180) = 0.27\ \text{m}^3$. The amount of air in the room is $180\ \text{m}^3$ at all times, so the percentage at time t (in mimutes) is $y(t)/180 \times 100$, and the change in the amount of carbon dioxide with respect to time is

$$\frac{dy}{dt} = (0.0005)\left(2\,\frac{\text{m}^3}{\text{min}}\right) - \frac{y(t)}{180}\left(2\,\frac{\text{m}^3}{\text{min}}\right) = 0.001 - \frac{y}{90} = \frac{9 - 100y}{9000}\,\frac{\text{m}^3}{\text{min}}$$

Hence, $\displaystyle\int \frac{dy}{9 - 100y} = \int \frac{dt}{9000}$ and $-\dfrac{1}{100}\ln|9 - 100y| = \dfrac{1}{9000}t + C$. Because $y(0) = 0.27$, we have

$-\frac{1}{100}\ln 18 = C$, so $-\frac{1}{100}\ln|9 - 100y| = \frac{1}{9000}t - \frac{1}{100}\ln 18 \quad\Rightarrow\quad \ln|9 - 100y| = -\frac{1}{90}t + \ln 18 \quad\Rightarrow$

$\ln|9 - 100y| = \ln e^{-t/90} + \ln 18 \quad\Rightarrow\quad \ln|9 - 100y| = \ln(18e^{-t/90})$, and $|9 - 100y| = 18e^{-t/90}$. Since y is continuous, $y(0) = 0.27$, and the right-hand side is never zero, we deduce that $9 - 100y$ is always negative. Thus, $|9 - 100y| = 100y - 9$ and we have $100y - 9 = 18e^{-t/90} \quad\Rightarrow\quad 100y = 9 + 18e^{-t/90} \quad\Rightarrow\quad y = 0.09 + 0.18e^{-t/90}$. The percentage of carbon

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

dioxide in the room is

$$p(t) = \frac{y}{180} \times 100 = \frac{0.09 + 0.18e^{-t/90}}{180} \times 100 = (0.0005 + 0.001e^{-t/90}) \times 100 = 0.05 + 0.1e^{-t/90}$$

In the long run, we have $\lim_{t\to\infty} p(t) = 0.05 + 0.1(0) = 0.05$; that is, the amount of carbon dioxide approaches 0.05% as time goes on.

49. Let $y(t)$ be the amount of alcohol in the vat after t minutes. Then $y(0) = 0.04(500) = 20$ gal. The amount of beer in the vat is 500 gallons at all times, so the percentage at time t (in minutes) is $y(t)/500 \times 100$, and the change in the amount of alcohol with respect to time t is $\dfrac{dy}{dt} = \text{rate in} - \text{rate out} = 0.06\left(5\ \dfrac{\text{gal}}{\text{min}}\right) - \dfrac{y(t)}{500}\left(5\ \dfrac{\text{gal}}{\text{min}}\right) = 0.3 - \dfrac{y}{100} = \dfrac{30-y}{100}\ \dfrac{\text{gal}}{\text{min}}$.

Hence, $\displaystyle\int \frac{dy}{30-y} = \int \frac{dt}{100}$ and $-\ln|30-y| = \frac{1}{100}t + C$. Because $y(0) = 20$, we have $-\ln 10 = C$, so

$-\ln|30-y| = \frac{1}{100}t - \ln 10 \quad\Rightarrow\quad \ln|30-y| = -t/100 + \ln 10 \quad\Rightarrow\quad \ln|30-y| = \ln e^{-t/100} + \ln 10 \quad\Rightarrow$

$\ln|30-y| = \ln(10e^{-t/100}) \quad\Rightarrow\quad |30-y| = 10e^{-t/100}$. Since y is continuous, $y(0) = 20$, and the right-hand side is never zero, we deduce that $30 - y$ is always positive. Thus, $30 - y = 10e^{-t/100} \quad\Rightarrow\quad y = 30 - 10e^{-t/100}$. The percentage of alcohol is $p(t) = y(t)/500 \times 100 = y(t)/5 = 6 - 2e^{-t/100}$. The percentage of alcohol after one hour is $p(60) = 6 - 2e^{-60/100} \approx 4.9$.

50. (a) If $y(t)$ is the amount of salt (in kg) after t minutes, then $y(0) = 0$ and the total amount of liquid in the tank remains constant at 1000 L.

$$\begin{aligned}\frac{dy}{dt} &= \left(0.05\ \frac{\text{kg}}{\text{L}}\right)\left(5\ \frac{\text{L}}{\text{min}}\right) + \left(0.04\ \frac{\text{kg}}{\text{L}}\right)\left(10\ \frac{\text{L}}{\text{min}}\right) - \left(\frac{y(t)}{1000}\ \frac{\text{kg}}{\text{L}}\right)\left(15\ \frac{\text{L}}{\text{min}}\right)\\ &= 0.25 + 0.40 - 0.015y = 0.65 - 0.015y = \frac{130-3y}{200}\ \frac{\text{kg}}{\text{min}}\end{aligned}$$

Hence, $\displaystyle\int \frac{dy}{130-3y} = \int \frac{dt}{200}$ and $-\frac{1}{3}\ln|130-3y| = \frac{1}{200}t + C$. Because $y(0) = 0$, we have $-\frac{1}{3}\ln 130 = C$, so $-\frac{1}{3}\ln|130-3y| = \frac{1}{200}t - \frac{1}{3}\ln 130 \quad\Rightarrow\quad \ln|130-3y| = -\frac{3}{200}t + \ln 130 = \ln(130e^{-3t/200})$, and $|130-3y| = 130e^{-3t/200}$. Since y is continuous, $y(0) = 0$, and the right-hand side is never zero, we deduce that $130 - 3y$ is always positive. Thus, $130 - 3y = 130e^{-3t/200}$ and $y = \frac{130}{3}(1 - e^{-3t/200})$ kg.

(b) After one hour, $y = \frac{130}{3}(1 - e^{-3\cdot 60/200}) = \frac{130}{3}(1 - e^{-0.9}) \approx 25.7$ kg.

Note: As $t \to \infty$, $y(t) \to \frac{130}{3} = 43\frac{1}{3}$ kg.

51. Assume that the raindrop begins at rest, so that $v(0) = 0$. $dm/dt = km$ and $(mv)' = gm \quad\Rightarrow\quad mv' + vm' = gm \quad\Rightarrow$

$mv' + v(km) = gm \quad\Rightarrow\quad v' + vk = g \quad\Rightarrow\quad \dfrac{dv}{dt} = g - kv \quad\Rightarrow\quad \displaystyle\int \frac{dv}{g-kv} = \int dt \quad\Rightarrow$

$-(1/k)\ln|g-kv| = t + C \quad\Rightarrow\quad \ln|g-kv| = -kt - kC \quad\Rightarrow\quad g - kv = Ae^{-kt}$. $v(0) = 0 \quad\Rightarrow\quad A = g$.

So $kv = g - ge^{-kt} \quad\Rightarrow\quad v = (g/k)(1 - e^{-kt})$. Since $k > 0$, as $t \to \infty$, $e^{-kt} \to 0$ and therefore, $\lim_{t\to\infty} v(t) = g/k$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

52. (a) $m\dfrac{dv}{dt} = -kv \quad\Rightarrow\quad \dfrac{dv}{v} = -\dfrac{k}{m}\,dt \quad\Rightarrow\quad \ln|v| = -\dfrac{k}{m}t + C$. Since $v(0) = v_0$, $\ln|v_0| = C$. Therefore,

$\ln\left|\dfrac{v}{v_0}\right| = -\dfrac{k}{m}t \quad\Rightarrow\quad \left|\dfrac{v}{v_0}\right| = e^{-kt/m} \quad\Rightarrow\quad v(t) = \pm v_0 e^{-kt/m}$. The sign is $+$ when $t = 0$, and we assume v is continuous, so that the sign is $+$ for all t. Thus, $v(t) = v_0 e^{-kt/m}$. $ds/dt = v_0 e^{-kt/m} \quad\Rightarrow$ $s(t) = -\dfrac{mv_0}{k}e^{-kt/m} + C'$.

From $s(0) = s_0$, we get $s_0 = -\dfrac{mv_0}{k} + C'$, so $C' = s_0 + \dfrac{mv_0}{k}$ and $s(t) = s_0 + \dfrac{mv_0}{k}(1 - e^{-kt/m})$.

The distance traveled from time 0 to time t is $s(t) - s_0$, so the total distance traveled is $\lim\limits_{t\to\infty}[s(t) - s_0] = \dfrac{mv_0}{k}$.

Note: In finding the limit, we use the fact that $k > 0$ to conclude that $\lim\limits_{t\to\infty} e^{-kt/m} = 0$.

(b) $m\dfrac{dv}{dt} = -kv^2 \quad\Rightarrow\quad \dfrac{dv}{v^2} = -\dfrac{k}{m}\,dt \quad\Rightarrow\quad \dfrac{-1}{v} = -\dfrac{kt}{m} + C \quad\Rightarrow\quad \dfrac{1}{v} = \dfrac{kt}{m} - C$. Since $v(0) = v_0$,

$C = -\dfrac{1}{v_0}$ and $\dfrac{1}{v} = \dfrac{kt}{m} + \dfrac{1}{v_0}$. Therefore, $v(t) = \dfrac{1}{kt/m + 1/v_0} = \dfrac{mv_0}{kv_0t + m}$. $\dfrac{ds}{dt} = \dfrac{mv_0}{kv_0t + m} \quad\Rightarrow$

$s(t) = \dfrac{m}{k}\displaystyle\int \dfrac{kv_0\,dt}{kv_0t + m} = \dfrac{m}{k}\ln|kv_0t + m| + C'$. Since $s(0) = s_0$, we get $s_0 = \dfrac{m}{k}\ln m + C' \quad\Rightarrow$

$C' = s_0 - \dfrac{m}{k}\ln m \quad\Rightarrow\quad s(t) = s_0 + \dfrac{m}{k}(\ln|kv_0t + m| - \ln m) = s_0 + \dfrac{m}{k}\ln\left|\dfrac{kv_0t + m}{m}\right|$.

We can rewrite the formulas for $v(t)$ and $s(t)$ as $v(t) = \dfrac{v_0}{1 + (kv_0/m)t}$ and $s(t) = s_0 + \dfrac{m}{k}\ln\left|1 + \dfrac{kv_0}{m}t\right|$.

Remarks: This model of horizontal motion through a resistive medium was designed to handle the case in which $v_0 > 0$. Then the term $-kv^2$ representing the resisting force causes the object to decelerate. The absolute value in the expression for $s(t)$ is unnecessary (since k, v_0, and m are all positive), and $\lim\limits_{t\to\infty} s(t) = \infty$. In other words, the object travels infinitely far. However, $\lim\limits_{t\to\infty} v(t) = 0$. When $v_0 < 0$, the term $-kv^2$ increases the magnitude of the object's negative velocity. According to the formula for $s(t)$, the position of the object approaches $-\infty$ as t approaches $m/k(-v_0)$: $\lim\limits_{t\to -m/(kv_0)} s(t) = -\infty$. Again the object travels infinitely far, but this time the feat is accomplished in a finite amount of time. Notice also that $\lim\limits_{t\to -m/(kv_0)} v(t) = -\infty$ when $v_0 < 0$, showing that the speed of the object increases without limit.

53. (a) $\dfrac{1}{L_1}\dfrac{dL_1}{dt} = k\dfrac{1}{L_2}\dfrac{dL_2}{dt} \quad\Rightarrow\quad \dfrac{d}{dt}(\ln L_1) = \dfrac{d}{dt}(k\ln L_2) \quad\Rightarrow\quad \displaystyle\int \dfrac{d}{dt}(\ln L_1)\,dt = \int \dfrac{d}{dt}(\ln L_2^k)\,dt \quad\Rightarrow$

$\ln L_1 = \ln L_2^k + C \quad\Rightarrow\quad L_1 = e^{\ln L_2^k + C} = e^{\ln L_2^k}e^C \quad\Rightarrow\quad L_1 = KL_2^k$, where $K = e^C$.

(b) From part (a) with $L_1 = B$, $L_2 = V$, and $k = 0.0794$, we have $B = KV^{0.0794}$.

54. (a) $\dfrac{dV}{dt} = a(\ln b - \ln V)V \quad\Rightarrow\quad \dfrac{dV}{dt} = -aV(\ln V - \ln b) \quad\Rightarrow\quad \dfrac{dV}{V\ln(V/b)} = -a\,dt \quad\Rightarrow$

$\displaystyle\int \dfrac{dV}{V\ln(V/b)} = \int -a\,dt \quad\Rightarrow\quad \int \dfrac{1}{u}\,du = \int -a\,dt \qquad \begin{bmatrix} u = \ln(V/b), \\ du = (1/V)\,dV \end{bmatrix} \quad\Rightarrow\quad \ln|u| = -at + k \quad\Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$|u| = e^{-at}e^k \quad \Rightarrow \quad u = Ce^{-at}$ [where $C = \pm e^k$] $\Rightarrow \quad \ln(V/b) = Ce^{-at} \quad \Rightarrow \quad \dfrac{V}{b} = e^{Ce^{-at}} \quad \Rightarrow$

$V = be^{Ce^{-at}}$ with $C \neq 0$.

(b) $V(0) = 1 \quad \Rightarrow \quad 1 = be^{Ce^{-a(0)}} \quad \Rightarrow \quad 1 = be^C \quad \Rightarrow \quad b = e^{-C}$, so $V = e^{-C}e^{Ce^{-at}} = e^{Ce^{-at}-C} = e^{C(e^{-at}-1)}$.

55. (a) The rate of growth of the area is jointly proportional to $\sqrt{A(t)}$ and $M - A(t)$; that is, the rate is proportional to the product of those two quantities. So for some constant k, $dA/dt = k\sqrt{A}\,(M - A)$. We are interested in the maximum of the function dA/dt (when the tissue grows the fastest), so we differentiate, using the Chain Rule and then substituting for dA/dt from the differential equation:

$$\frac{d}{dt}\left(\frac{dA}{dt}\right) = k\left[\sqrt{A}\,(-1)\frac{dA}{dt} + (M - A)\cdot \tfrac{1}{2}A^{-1/2}\frac{dA}{dt}\right] = \tfrac{1}{2}kA^{-1/2}\frac{dA}{dt}\left[-2A + (M - A)\right]$$

$$= \tfrac{1}{2}kA^{-1/2}\left[k\sqrt{A}(M - A)\right][M - 3A] = \tfrac{1}{2}k^2(M - A)(M - 3A)$$

This is 0 when $M - A = 0$ [this situation never actually occurs, since the graph of $A(t)$ is asymptotic to the line $y = M$, as in the logistic model] and when $M - 3A = 0 \quad \Leftrightarrow \quad A(t) = M/3$. This represents a maximum by the First Derivative Test, since $\dfrac{d}{dt}\left(\dfrac{dA}{dt}\right)$ goes from positive to negative when $A(t) = M/3$.

(b) From the CAS, we get $A(t) = M\left(\dfrac{Ce^{\sqrt{M}kt} - 1}{Ce^{\sqrt{M}kt} + 1}\right)^2$. To get C in terms of the initial area A_0 and the maximum area M, we substitute $t = 0$ and $A = A_0 = A(0)$: $A_0 = M\left(\dfrac{C - 1}{C + 1}\right)^2 \quad \Leftrightarrow \quad (C + 1)\sqrt{A_0} = (C - 1)\sqrt{M} \quad \Leftrightarrow$

$C\sqrt{A_0} + \sqrt{A_0} = C\sqrt{M} - \sqrt{M} \quad \Leftrightarrow \quad \sqrt{M} + \sqrt{A_0} = C\sqrt{M} - C\sqrt{A_0} \quad \Leftrightarrow$

$\sqrt{M} + \sqrt{A_0} = C\left(\sqrt{M} - \sqrt{A_0}\right) \quad \Leftrightarrow \quad C = \dfrac{\sqrt{M} + \sqrt{A_0}}{\sqrt{M} - \sqrt{A_0}}$. [Notice that if $A_0 = 0$, then $C = 1$.]

56. (a) The volume of the newly frozen water layer is $\Delta V = A\,\Delta h$ and since the mass density $D = \Delta M/\Delta V$, the mass of this layer is $\Delta M = D\,\Delta V = D \cdot A\,\Delta h$. The heat loss required to freeze ΔM kg of water is $\Delta Q = L\,\Delta M = L \cdot AD\,\Delta h$, which after rearranging gives $\Delta h = (1/LAD)\,\Delta Q$ or $\Delta h/\Delta Q = 1/LAD$. Taking the limit as $\Delta Q \to 0$ then gives $dh/dQ = 1/LAD$.

(b) $\dfrac{dh}{dt} = \dfrac{dh}{dQ}\dfrac{dQ}{dt} = \dfrac{1}{LAD}\cdot\dfrac{kA}{h}(T_w - T_a) = \dfrac{k(T_w - T_a)}{LDh}$

The growth rate of the ice, dh/dt, is inversely proportional to the thickness of the ice, h, so as the ice gets thicker its rate of growth decreases until some equilibrium is reached. Hence, thin ice grows more rapidly than thick ice.

(c) $\dfrac{dh}{dt} = \dfrac{k(T_w - T_a)}{LDh} \quad \Rightarrow \quad h\,dh = \dfrac{k(T_w - T_a)}{LD}\,dt \quad \Rightarrow \quad \displaystyle\int h\,dh = \int \frac{k(T_w - T_a)}{LD}\,dt \quad \Rightarrow$

$\dfrac{1}{2}h^2 = \dfrac{k(T_w - T_a)}{LD}t + C$. $h(0) = h_0 \quad \Rightarrow \quad \dfrac{1}{2}h_0^2 = 0 + C = C$, so $\dfrac{1}{2}h^2 = \dfrac{k(T_w - T_a)}{LD}t + \dfrac{1}{2}h_0^2 \quad \Rightarrow$

$h = \sqrt{\dfrac{2k(T_w - T_a)}{LD}t + h_0^2}$ [only take the positive root since $h \geq 0$].

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. (a) According to the hint we use the Chain Rule: $m\dfrac{dv}{dt} = m\dfrac{dv}{dx}\cdot\dfrac{dx}{dt} = mv\dfrac{dv}{dx} = -\dfrac{mgR^2}{(x+R)^2} \quad\Rightarrow$

$\displaystyle\int v\,dv = \int \frac{-gR^2\,dx}{(x+R)^2} \quad\Rightarrow\quad \frac{v^2}{2} = \frac{gR^2}{x+R} + C$. When $x=0$, $v=v_0$, so $\dfrac{v_0^2}{2} = \dfrac{gR^2}{0+R} + C \quad\Rightarrow$

$C = \frac{1}{2}v_0^2 - gR \quad\Rightarrow\quad \frac{1}{2}v^2 - \frac{1}{2}v_0^2 = \dfrac{gR^2}{x+R} - gR$. Now at the top of its flight, the rocket's velocity will be 0, and its height will be $x=h$. Solving for v_0: $-\frac{1}{2}v_0^2 = \dfrac{gR^2}{h+R} - gR \quad\Rightarrow\quad \dfrac{v_0^2}{2} = g\left[-\dfrac{R^2}{R+h} + \dfrac{R(R+h)}{R+h}\right] = \dfrac{gRh}{R+h} \quad\Rightarrow$

$v_0 = \sqrt{\dfrac{2gRh}{R+h}}$.

(b) $v_e = \lim\limits_{h\to\infty} v_0 = \lim\limits_{h\to\infty}\sqrt{\dfrac{2gRh}{R+h}} = \lim\limits_{h\to\infty}\sqrt{\dfrac{2gR}{(R/h)+1}} = \sqrt{2gR}$

(c) $v_e = \sqrt{2\cdot 32\text{ ft/s}^2\cdot 3960\text{ mi}\cdot 5280\text{ ft/mi}} \approx 36{,}581\text{ ft/s} \approx 6.93\text{ mi/s}$

APPLIED PROJECT How Fast Does a Tank Drain?

1. (a) $V = \pi r^2 h \quad\Rightarrow\quad \dfrac{dV}{dt} = \pi r^2\dfrac{dh}{dt}$ [implicit differentiation] $\Rightarrow$

$$\frac{dh}{dt} = \frac{1}{\pi r^2}\frac{dV}{dt} = \frac{1}{\pi r^2}\left(-a\sqrt{2gh}\right) = \frac{1}{\pi 2^2}\left[-\pi\left(\tfrac{1}{12}\right)^2\sqrt{2\cdot 32}\,\sqrt{h}\right] = -\tfrac{1}{72}\sqrt{h}$$

(b) $\dfrac{dh}{dt} = -\frac{1}{72}\sqrt{h} \quad\Rightarrow\quad h^{-1/2}\,dh = -\frac{1}{72}\,dt \quad\Rightarrow\quad 2\sqrt{h} = -\frac{1}{72}t + C$.

$h(0) = 6 \quad\Rightarrow\quad 2\sqrt{6} = 0 + C \quad\Rightarrow\quad C = 2\sqrt{6} \quad\Rightarrow\quad h(t) = \left(-\frac{1}{144}t + \sqrt{6}\right)^2$.

(c) We want to find t when $h=0$, so we set $h = 0 = \left(-\frac{1}{144}t + \sqrt{6}\right)^2 \quad\Rightarrow\quad t = 144\sqrt{6} \approx 5$ min 53 s.

2. (a) $\dfrac{dh}{dt} = k\sqrt{h} \quad\Rightarrow\quad h^{-1/2}\,dh = k\,dt \quad [h\neq 0] \quad\Rightarrow\quad 2\sqrt{h} = kt + C \quad\Rightarrow$

$h(t) = \frac{1}{4}(kt+C)^2$. Since $h(0) = 10$ cm, the relation $2\sqrt{h(t)} = kt + C$ gives us $2\sqrt{10} = C$. Also, $h(68) = 3$ cm, so $2\sqrt{3} = 68k + 2\sqrt{10}$ and $k = -\dfrac{\sqrt{10}-\sqrt{3}}{34}$. Thus,

$h(t) = \dfrac{1}{4}\left(2\sqrt{10} - \dfrac{\sqrt{10}-\sqrt{3}}{34}\,t\right)^2 \approx 10 - 0.133t + 0.00044t^2$.

Here is a table of values of $h(t)$ correct to one decimal place.

t (in s)	$h(t)$ (in cm)
10	8.7
20	7.5
30	6.4
40	5.4
50	4.5
60	3.6

(b) The answers to this part are to be obtained experimentally. See the article by Tom Farmer and Fred Gass, *Physical Demonstrations in the Calculus Classroom,* College Mathematics Journal 1992, pp. 146–148.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. $V(t) = \pi r^2 h(t) = 100\pi h(t) \quad \Rightarrow \quad \dfrac{dV}{dh} = 100\pi$ and $\dfrac{dV}{dt} = \dfrac{dV}{dh}\dfrac{dh}{dt} = 100\pi\dfrac{dh}{dt}$.

Diameter $= 2.5$ inches $\Rightarrow$ radius $= 1.25$ inches $= \frac{5}{4}\cdot\frac{1}{12}$ foot $= \frac{5}{48}$ foot. Thus, $\dfrac{dV}{dt} = -a\sqrt{2gh} \quad \Rightarrow$

$100\pi\dfrac{dh}{dt} = -\pi\left(\frac{5}{48}\right)^2\sqrt{2\cdot 32h} = -\dfrac{25\pi}{288}\sqrt{h} \quad \Rightarrow \quad \dfrac{dh}{dt} = -\dfrac{\sqrt{h}}{1152} \quad \Rightarrow \quad \int h^{-1/2}\,dh = \int -\frac{1}{1152}\,dt \quad \Rightarrow$

$2\sqrt{h} = -\frac{1}{1152}t + C \quad \Rightarrow \quad \sqrt{h} = -\frac{1}{2304}t + k \quad \Rightarrow \quad h(t) = \left(-\frac{1}{2304}t + k\right)^2$. The water pressure after t seconds is $62.5h(t)$ lb/ft^2, so the condition that the pressure be at least 2160 lb/ft^2 for 10 minutes (600 seconds) is the condition $62.5\cdot h(600) \geq 2160$; that is, $\left(k - \frac{600}{2304}\right)^2 \geq \frac{2160}{62.5} \quad \Rightarrow \quad \left|k - \frac{25}{96}\right| \geq \sqrt{34.56} \quad \Rightarrow \quad k \geq \frac{25}{96} + \sqrt{34.56}$. Now $h(0) = k^2$, so the height of the tank should be at least $\left(\frac{25}{96} + \sqrt{34.56}\right)^2 \approx 37.69$ ft.

4. (a) If the radius of the circular cross-section at height h is r, then the Pythagorean Theorem gives $r^2 = 2^2 - (2-h)^2$ since the radius of the tank is 2 m. So $A(h) = \pi r^2 = \pi[4 - (2-h)^2] = \pi(4h - h^2)$. Thus, $A(h)\dfrac{dh}{dt} = -a\sqrt{2gh} \quad \Rightarrow$

$\pi(4h - h^2)\dfrac{dh}{dt} = -\pi(0.01)^2\sqrt{2\cdot 10h} \quad \Rightarrow \quad (4h - h^2)\dfrac{dh}{dt} = -0.0001\sqrt{20h}$.

(b) From part (a) we have $(4h^{1/2} - h^{3/2})\,dh = \left(-0.0001\sqrt{20}\right)dt \;\Rightarrow\; \frac{8}{3}h^{3/2} - \frac{2}{5}h^{5/2} = \left(-0.0001\sqrt{20}\right)t + C$.

$h(0) = 2 \quad \Rightarrow \quad \frac{8}{3}(2)^{3/2} - \frac{2}{5}(2)^{5/2} = C \quad \Rightarrow \quad C = \left(\frac{16}{3} - \frac{8}{5}\right)\sqrt{2} = \frac{56}{15}\sqrt{2}$. To find out how long it will take to drain all the water we evaluate t when $h = 0$: $0 = \left(-0.0001\sqrt{20}\right)t + C \quad \Rightarrow$

$$t = \frac{C}{0.0001\sqrt{20}} = \frac{56\sqrt{2}/15}{0.0001\sqrt{20}} = \frac{11{,}200\sqrt{10}}{3} \approx 11{,}806 \text{ s} \approx 3 \text{ h } 17 \text{ min}$$

9.4 Models for Population Growth

1. (a) Comparing the given equation, $\dfrac{dP}{dt} = 0.04P\left(1 - \dfrac{P}{1200}\right)$, to Equation 4, $\dfrac{dP}{dt} = kP\left(1 - \dfrac{P}{M}\right)$, we see that the carrying capacity is $M = 1200$ and the value of k is 0.04.

(b) By Equation 7, the solution of the equation is $P(t) = \dfrac{M}{1 + Ae^{-kt}}$, where $A = \dfrac{M - P_0}{P_0}$. Since $P(0) = P_0 = 60$, we have $A = \dfrac{1200 - 60}{60} = 19$, and hence, $P(t) = \dfrac{1200}{1 + 19e^{-0.04t}}$.

(c) The population after 10 weeks is $P(10) = \dfrac{1200}{1 + 19e^{-0.04(10)}} \approx 87$.

2. (a) $dP/dt = 0.02P - 0.0004P^2 = 0.02P(1 - 0.02P) = 0.02P(1 - P/50)$. Comparing to Equation 4, $dP/dt = kP(1 - P/M)$, we see that the carrying capacity is $M = 50$ and the value of k is 0.02.

(b) By Equation 7, the solution of the equation is $P(t) = \dfrac{M}{1 + Ae^{-kt}}$, where $A = \dfrac{M - P_0}{P_0}$. Since $P(0) = P_0 = 40$, we have $A = \dfrac{50 - 40}{40} = 0.25$, and hence, $P(t) = \dfrac{50}{1 + 0.25e^{-0.02t}}$.

(c) The population after 10 weeks is $P(10) = \dfrac{50}{1 + 0.25e^{-0.02(10)}} \approx 42$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a) $dP/dt = 0.05P - 0.0005P^2 = 0.05P(1 - 0.01P) = 0.05P(1 - P/100)$. Comparing to Equation 4, $dP/dt = kP(1 - P/M)$, we see that the carrying capacity is $M = 100$ and the value of k is 0.05.

(b) The slopes close to 0 occur where P is near 0 or 100. The largest slopes appear to be on the line $P = 50$. The solutions are increasing for $0 < P_0 < 100$ and decreasing for $P_0 > 100$.

(c)

All of the solutions approach $P = 100$ as t increases. As in part (b), the solutions differ since for $0 < P_0 < 100$ they are increasing, and for $P_0 > 100$ they are decreasing. Also, some have an IP and some don't. It appears that the solutions which have $P_0 = 20$ and $P_0 = 40$ have inflection points at $P = 50$.

(d) The equilibrium solutions are $P = 0$ (trivial solution) and $P = 100$. The increasing solutions move away from $P = 0$ and all nonzero solutions approach $P = 100$ as $t \to \infty$.

4. (a) $M = 6000$ and $k = 0.0015 \quad \Rightarrow \quad dP/dt = 0.0015P(1 - P/6000)$.

(b)

All of the solution curves approach 6000 as $t \to \infty$.

(c)

The curves with $P_0 = 1000$ and $P_0 = 2000$ appear to be concave upward at first and then concave downward. The curve with $P_0 = 4000$ appears to be concave downward everywhere. The curve with $P_0 = 8000$ appears to be concave upward everywhere. The inflection points are where the population grows the fastest.

(d) See the solution to Exercise 9.2.25 for a possible program to calculate $P(50)$. [In this case, we use X = 0, H = 1, N = 50, $\text{Y}_1 = 0.0015y(1 - y/6000)$, and Y = 1000.] We find that $P(50) \approx 1064$.

(e) Using Equation 7 with $M = 6000$, $k = 0.0015$, and $P_0 = 1000$, we have $P(t) = \dfrac{M}{1 + Ae^{-kt}} = \dfrac{6000}{1 + Ae^{-0.0015t}}$, where $A = \dfrac{M - P_0}{P_0} = \dfrac{6000 - 1000}{1000} = 5$. Thus, $P(50) = \dfrac{6000}{1 + 5e^{-0.0015(50)}} \approx 1064.1$, which is extremely close to the estimate obtained in part (d).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(f)

The curves are very similar.

5. (a) $\dfrac{dy}{dt} = ky\left(1 - \dfrac{y}{M}\right) \quad \Rightarrow \quad y(t) = \dfrac{M}{1 + Ae^{-kt}}$ with $A = \dfrac{M - y(0)}{y(0)}$. With $M = 8 \times 10^7$, $k = 0.71$, and $y(0) = 2 \times 10^7$, we get the model $y(t) = \dfrac{8 \times 10^7}{1 + 3e^{-0.71t}}$, so $y(1) = \dfrac{8 \times 10^7}{1 + 3e^{-0.71}} \approx 3.23 \times 10^7$ kg.

(b) $y(t) = 4 \times 10^7 \quad \Rightarrow \quad \dfrac{8 \times 10^7}{1 + 3e^{-0.71t}} = 4 \times 10^7 \quad \Rightarrow \quad 2 = 1 + 3e^{-0.71t} \quad \Rightarrow \quad e^{-0.71t} = \frac{1}{3} \quad \Rightarrow$

$-0.71t = \ln \frac{1}{3} \quad \Rightarrow \quad t = \dfrac{\ln 3}{0.71} \approx 1.55$ years

6. (a) $\dfrac{dP}{dt} = 0.4P - 0.001P^2 = 0.4P(1 - 0.0025P)$ $\left[\frac{0.001}{0.4} = 0.0025\right]$ $= 0.4P\left(1 - \dfrac{P}{400}\right)$ $\left[0.0025^{-1} = 400\right]$

Thus, by Equation 4, $k = 0.4$ and the carrying capacity is 400.

(b) Using the fact that $P(0) = 50$ and the formula for dP/dt, we get

$P'(0) = \left.\dfrac{dP}{dt}\right|_{t=0} = 0.4(50) - 0.001(50)^2 = 20 - 2.5 = 17.5.$

(c) From Equation 7, $A = \dfrac{M - P_0}{P_0} = \dfrac{400 - 50}{50} = 7$, so $P = \dfrac{400}{1 + 7e^{-0.4t}}$. The population reaches 50% of the carrying capacity, 200, when $200 = \dfrac{400}{1 + 7e^{-0.4t}} \quad \Rightarrow \quad 1 + 7e^{-0.4t} = 2 \quad \Rightarrow \quad e^{-0.4t} = \frac{1}{7} \quad \Rightarrow \quad -0.4t = \ln \frac{1}{7} \quad \Rightarrow$

$t = \left(\ln \frac{1}{7}\right)/(-0.4) \approx 4.86$ years.

7. Using Equation 7, $A = \dfrac{M - P_0}{P_0} = \dfrac{10{,}000 - 1000}{1000} = 9$, so $P(t) = \dfrac{10{,}000}{1 + 9e^{-kt}}$. $P(1) = 2500 \quad \Rightarrow$

$2500 = \dfrac{10{,}000}{1 + 9e^{-k(1)}} \quad \Rightarrow \quad 1 + 9e^{-k} = 4 \quad \Rightarrow \quad 9e^{-k} = 3 \quad \Rightarrow \quad e^{-k} = \frac{1}{3} \quad \Rightarrow \quad -k = \ln \frac{1}{3} \quad \Rightarrow \quad k = \ln 3$. After another three years, $t = 4$, and $P(4) = \dfrac{10{,}000}{1 + 9e^{-(\ln 3)4}} = \dfrac{10{,}000}{1 + 9\left(e^{\ln 3}\right)^{-4}} = \dfrac{10{,}000}{1 + 9(3)^{-4}} = \dfrac{10{,}000}{1 + \frac{1}{9}} = \dfrac{10{,}000}{\frac{10}{9}} = 9000.$

8. (a)

From the graph, we estimate the carrying capacity M for the yeast population to be 680.

(b) An estimate of the initial relative growth rate is $\dfrac{1}{P_0}\dfrac{dP}{dt} = \dfrac{1}{18} \cdot \dfrac{39 - 18}{2 - 0} = \dfrac{7}{12} = 0.58\overline{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) An exponential model is $P(t) = 18e^{7t/12}$. A logistic model is $P(t) = \dfrac{680}{1 + Ae^{-7t/12}}$, where $A = \frac{680 - 18}{18} = \frac{331}{9}$.

(d)

Time in Hours	Observed Values	Exponential Model	Logistic Model
0	18	18	18
2	39	58	55
4	80	186	149
6	171	596	322
8	336	1914	505
10	509	6147	614
12	597	19,739	658
14	640	63,389	673
16	664	203,558	678
18	672	653,679	679

The exponential model is a poor fit for anything beyond the first two observed values. The logistic model varies more for the middle values than it does for the values at either end, but provides a good general fit, as shown in the figure.

(e) $P(7) = \dfrac{680}{1 + \frac{331}{9}e^{-7(7/12)}} \approx 420$ yeast cells

9. (a) We will assume that the difference in birth and death rates is 20 million/year. Let $t = 0$ correspond to the year 2000. Thus,

$k \approx \dfrac{1}{P}\dfrac{dP}{dt} = \dfrac{1}{6.1 \text{ billion}}\left(\dfrac{20 \text{ million}}{\text{year}}\right) = \dfrac{1}{305}$, and $\dfrac{dP}{dt} = kP\left(1 - \dfrac{P}{M}\right) = \dfrac{1}{305}P\left(1 - \dfrac{P}{20}\right)$ with P in billions.

(b) $A = \dfrac{M - P_0}{P_0} = \dfrac{20 - 6.1}{6.1} = \dfrac{139}{61} \approx 2.2787$. $P(t) = \dfrac{M}{1 + Ae^{-kt}} = \dfrac{20}{1 + \frac{139}{61}e^{-t/305}}$, so

$P(10) = \dfrac{20}{1 + \frac{139}{61}e^{-10/305}} \approx 6.24$ billion, which underestimates the actual 2010 population of 6.9 billion.

(c) The years 2100 and 2500 correspond to $t = 100$ and $t = 500$, respectively. $P(100) = \dfrac{20}{1 + \frac{139}{61}e^{-100/305}} \approx 7.57$ billion

and $P(500) = \dfrac{20}{1 + \frac{139}{61}e^{-500/305}} \approx 13.87$ billion.

10. (a) Let $t = 0$ correspond to the year 2000. $A = \dfrac{M - P_0}{P_0} = \dfrac{800 - 282}{282} = \dfrac{259}{141} \approx 1.8369$.

$P(t) = \dfrac{M}{1 + Ae^{-kt}} = \dfrac{800}{1 + \frac{259}{141}e^{-kt}}$ with P in millions.

(b) $P(10) = 309 \quad \Leftrightarrow \quad \dfrac{800}{1 + \frac{259}{141}e^{-10k}} = 309 \quad \Leftrightarrow \quad \dfrac{800}{309} = 1 + \dfrac{259}{141}e^{-10k} \quad \Leftrightarrow \quad \dfrac{491}{309} = \dfrac{259}{141}e^{-10k} \quad \Leftrightarrow$

$\dfrac{491 \cdot 141}{309 \cdot 259} = e^{-10k} \quad \Leftrightarrow \quad -10k = \ln \dfrac{491 \cdot 47}{103 \cdot 259} \quad \Leftrightarrow \quad k = -\dfrac{1}{10}\ln \dfrac{23{,}077}{26{,}677} \approx 0.0145$.

(c) The years 2100 and 2200 correspond to $t = 100$ and $t = 200$, respectively. $P(100) = \dfrac{800}{1 + \frac{259}{141}e^{-100k}} \approx 559$ million and

$P(200) = \dfrac{800}{1 + \frac{259}{141}e^{-200k}} \approx 727$ million.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $P(t) = 500 \quad \Leftrightarrow \quad \dfrac{800}{1+\frac{259}{141}e^{-kt}} = 500 \quad \Leftrightarrow \quad \dfrac{800}{500} = 1 + \dfrac{259}{141}e^{-kt} \quad \Leftrightarrow \quad \dfrac{3}{5} = \dfrac{259}{141}e^{-kt} \quad \Leftrightarrow \quad \dfrac{3\cdot 141}{5\cdot 259} = e^{-kt} \quad \Leftrightarrow$

$-kt = \ln\dfrac{423}{1295} \quad \Leftrightarrow \quad t = 10\,\dfrac{\ln(423/1295)}{\ln(23{,}077/26{,}677)} \approx 77.18$ years. Our logistic model predicts that the US population will exceed 500 million in 77.18 years; that is, in the year 2077.

11. (a) Our assumption is that $\dfrac{dy}{dt} = ky(1-y)$, where y is the fraction of the population that has heard the rumor.

(b) Using the logistic equation (4), $\dfrac{dP}{dt} = kP\left(1 - \dfrac{P}{M}\right)$, we substitute $y = \dfrac{P}{M}$, $P = My$, and $\dfrac{dP}{dt} = M\,\dfrac{dy}{dt}$,

to obtain $M\,\dfrac{dy}{dt} = k(My)(1-y) \quad \Leftrightarrow \quad \dfrac{dy}{dt} = ky(1-y)$, our equation in part (a).

Now the solution to (4) is $P(t) = \dfrac{M}{1+Ae^{-kt}}$, where $A = \dfrac{M-P_0}{P_0}$.

We use the same substitution to obtain $My = \dfrac{M}{1+\dfrac{M-My_0}{My_0}e^{-kt}} \quad \Rightarrow \quad y = \dfrac{y_0}{y_0+(1-y_0)e^{-kt}}$.

Alternatively, we could use the same steps as outlined in the solution of Equation 4.

(c) Let t be the number of hours since 8 AM. Then $y_0 = y(0) = \frac{80}{1000} = 0.08$ and $y(4) = \frac{1}{2}$, so

$\dfrac{1}{2} = y(4) = \dfrac{0.08}{0.08+0.92e^{-4k}}$. Thus, $0.08 + 0.92e^{-4k} = 0.16$, $e^{-4k} = \frac{0.08}{0.92} = \frac{2}{23}$, and $e^{-k} = \left(\frac{2}{23}\right)^{1/4}$,

so $y = \dfrac{0.08}{0.08+0.92(2/23)^{t/4}} = \dfrac{2}{2+23(2/23)^{t/4}}$. Solving this equation for t, we get

$2y + 23y\left(\dfrac{2}{23}\right)^{t/4} = 2 \quad \Rightarrow \quad \left(\dfrac{2}{23}\right)^{t/4} = \dfrac{2-2y}{23y} \quad \Rightarrow \quad \left(\dfrac{2}{23}\right)^{t/4} = \dfrac{2}{23}\cdot\dfrac{1-y}{y} \quad \Rightarrow \quad \left(\dfrac{2}{23}\right)^{t/4-1} = \dfrac{1-y}{y}$.

It follows that $\dfrac{t}{4} - 1 = \dfrac{\ln[(1-y)/y]}{\ln\frac{2}{23}}$, so $t = 4\left[1 + \dfrac{\ln((1-y)/y)}{\ln\frac{2}{23}}\right]$.

When $y = 0.9$, $\dfrac{1-y}{y} = \frac{1}{9}$, so $t = 4\left(1 - \dfrac{\ln 9}{\ln\frac{2}{23}}\right) \approx 7.6$ h or 7 h 36 min. Thus, 90% of the population will have heard the rumor by 3:36 PM.

12. (a) $P(0) = P_0 = 400$, $P(1) = 1200$ and $M = 10{,}000$. From the solution to the logistic differential equation

$P(t) = \dfrac{P_0M}{P_0+(M-P_0)e^{-kt}}$, we get $P = \dfrac{400\,(10{,}000)}{400+(9600)e^{-kt}} = \dfrac{10{,}000}{1+24e^{-kt}}$. $\quad P(1) = 1200 \quad \Rightarrow$

$1 + 24e^{-k} = \frac{100}{12} \quad \Rightarrow \quad e^k = \frac{288}{88} \quad \Rightarrow \quad k = \ln\frac{36}{11}$. So $P = \dfrac{10{,}000}{1+24e^{-t\ln(36/11)}} = \dfrac{10{,}000}{1+24\cdot(11/36)^t}$.

(b) $5000 = \dfrac{10{,}000}{1+24(11/36)^t} \quad \Rightarrow \quad 24\left(\frac{11}{36}\right)^t = 1 \quad \Rightarrow \quad t\ln\frac{11}{36} = \ln\frac{1}{24} \quad \Rightarrow \quad t \approx 2.68$ years.

13. (a) $\dfrac{dP}{dt} = kP\left(1-\dfrac{P}{M}\right) \quad \Rightarrow \quad \dfrac{d^2P}{dt^2} = k\left[P\left(-\dfrac{1}{M}\dfrac{dP}{dt}\right) + \left(1-\dfrac{P}{M}\right)\dfrac{dP}{dt}\right] = k\,\dfrac{dP}{dt}\left(-\dfrac{P}{M} + 1 - \dfrac{P}{M}\right)$

$$= k\left[kP\left(1-\frac{P}{M}\right)\right]\left(1-\frac{2P}{M}\right) = k^2P\left(1-\frac{P}{M}\right)\left(1-\frac{2P}{M}\right)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) P grows fastest when P' has a maximum, that is, when $P'' = 0$. From part (a), $P'' = 0 \quad \Leftrightarrow \quad P = 0, P = M$, or $P = M/2$. Since $0 < P < M$, we see that $P'' = 0 \quad \Leftrightarrow \quad P = M/2$.

14. First we keep k constant (at 0.1, say) and change P_0 in the function $P = \dfrac{10P_0}{P_0 + (10 - P_0)e^{-0.1t}}$. (Notice that P_0 is the P-intercept.) If $P_0 = 0$, the function is 0 everywhere. For $0 < P_0 < 5$, the curve has an inflection point, which moves to the right as P_0 decreases. If $5 < P_0 < 10$, the graph is concave down everywhere. (We are considering only $t \geq 0$.) If $P_0 = 10$, the function is the constant function $P = 10$, and if $P_0 > 10$, the function decreases. For all $P_0 \neq 0$, $\lim\limits_{t \to \infty} P = 10$.

Now we instead keep P_0 constant (at $P_0 = 1$) and change k in the function $P = \dfrac{10}{1 + 9e^{-kt}}$. It seems that as k increases, the graph approaches the line $P = 10$ more and more quickly. (Note that the only difference in the shape of the curves is in the horizontal scaling; if we choose suitable x-scales, the graphs all look the same.)

15. (a)

(b) After subtracting 900 from each value of P, we get the logistic model $f(t) = \dfrac{345.5899}{1 + 7.9977e^{-0.2482t}}$.

(c) $P(t) = 900 + \dfrac{345.5899}{1 + 7.9977e^{-0.2482t}}$

The model fits the data reasonably well, even though the data is decreasing for the last five values and a logistic function is never decreasing.

(d) As $t \to \infty$, $P(t) \to 900 + 345.6 = 1245.6$, so the population approaches 1.246 million.

16. Let $N(t)$ represent the number of active Twitter users (in millions) t years since January 1, 2010. After using a calculator to fit the models to the data, we find the exponential function is $N(t) = 53.1390(1.4039)^t$ and the logistic function is $N_L(t) = \dfrac{325.9251}{1 + 8.7349e^{-0.8906t}}$. N_L is a reasonably accurate model, whereas N is not, since the exponential model grows at a much faster rate than the data as t increases.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. (a) $\dfrac{dP}{dt} = kP - m = k\left(P - \dfrac{m}{k}\right)$. Let $y = P - \dfrac{m}{k}$, so $\dfrac{dy}{dt} = \dfrac{dP}{dt}$ and the differential equation becomes $\dfrac{dy}{dt} = ky$.

The solution is $y = y_0 e^{kt} \;\Rightarrow\; P - \dfrac{m}{k} = \left(P_0 - \dfrac{m}{k}\right)e^{kt} \;\Rightarrow\; P(t) = \dfrac{m}{k} + \left(P_0 - \dfrac{m}{k}\right)e^{kt}$.

(b) Since $k > 0$, there will be an exponential expansion $\;\Leftrightarrow\; P_0 - \dfrac{m}{k} > 0 \;\Leftrightarrow\; m < kP_0$.

(c) The population will be constant if $P_0 - \dfrac{m}{k} = 0 \;\Leftrightarrow\; m = kP_0$. It will decline if $P_0 - \dfrac{m}{k} < 0 \;\Leftrightarrow\; m > kP_0$.

(d) $P_0 = 8{,}000{,}000$, $k = \alpha - \beta = 0.016$, $m = 210{,}000 \;\Rightarrow\; m > kP_0$ $(= 128{,}000)$, so by part (c), the population was declining.

18. (a) $\dfrac{dy}{dt} = ky^{1+c} \;\Rightarrow\; y^{-1-c}\,dy = k\,dt \;\Rightarrow\; \dfrac{y^{-c}}{-c} = kt + C$. Since $y(0) = y_0$, we have $C = \dfrac{y_0^{-c}}{-c}$. Thus,

$\dfrac{y^{-c}}{-c} = kt + \dfrac{y_0^{-c}}{-c}$, or $y^{-c} = y_0^{-c} - ckt$. So $y^c = \dfrac{1}{y_0^{-c} - ckt} = \dfrac{y_0^c}{1 - cy_0^c kt}$ and $y(t) = \dfrac{y_0}{(1 - cy_0^c kt)^{1/c}}$.

(b) $y(t) \to \infty$ as $1 - cy_0^c kt \to 0$, that is, as $t \to \dfrac{1}{cy_0^c k}$. Define $T = \dfrac{1}{cy_0^c k}$. Then $\lim\limits_{t \to T^-} y(t) = \infty$.

(c) According to the data given, we have $c = 0.01$, $y(0) = 2$, and $y(3) = 16$, where the time t is given in months. Thus,

$y_0 = 2$ and $16 = y(3) = \dfrac{y_0}{(1 - cy_0^c k \cdot 3)^{1/c}}$. Since $T = \dfrac{1}{cy_0^c k}$, we will solve for $cy_0^c k$. $\;16 = \dfrac{2}{(1 - 3cy_0^c k)^{100}} \;\Rightarrow$

$1 - 3cy_0^c k = \left(\frac{1}{8}\right)^{0.01} = 8^{-0.01} \;\Rightarrow\; cy_0^c k = \frac{1}{3}\left(1 - 8^{-0.01}\right)$. Thus, doomsday occurs when

$t = T = \dfrac{1}{cy_0^c k} = \dfrac{3}{1 - 8^{-0.01}} \approx 145.77$ months or 12.15 years.

19. (a) The term -15 represents a harvesting of fish at a constant rate—in this case, 15 fish/week. This is the rate at which fish are caught.

(b)

(c) From the graph in part (b), it appears that $P(t) = 250$ and $P(t) = 750$ are the equilibrium solutions. We confirm this analytically by solving the equation $dP/dt = 0$ as follows: $0.08P(1 - P/1000) - 15 = 0 \;\Rightarrow$

$0.08P - 0.00008P^2 - 15 = 0 \;\Rightarrow$

$-0.00008(P^2 - 1000P + 187{,}500) = 0 \;\Rightarrow$

$(P - 250)(P - 750) = 0 \;\Rightarrow\; P = 250$ or 750.

(d)

For $0 < P_0 < 250$, $P(t)$ decreases to 0. For $P_0 = 250$, $P(t)$ remains constant. For $250 < P_0 < 750$, $P(t)$ increases and approaches 750. For $P_0 = 750$, $P(t)$ remains constant. For $P_0 > 750$, $P(t)$ decreases and approaches 750.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(e) $\dfrac{dP}{dt} = 0.08P\left(1 - \dfrac{P}{1000}\right) - 15 \quad \Leftrightarrow \quad -\dfrac{100{,}000}{8} \cdot \dfrac{dP}{dt} = (0.08P - 0.00008P^2 - 15) \cdot \left(-\dfrac{100{,}000}{8}\right) \quad \Leftrightarrow$

$-12{,}500\,\dfrac{dP}{dt} = P^2 - 1000P + 187{,}500 \quad \Leftrightarrow \quad \dfrac{dP}{(P-250)(P-750)} = -\dfrac{1}{12{,}500}\,dt \quad \Leftrightarrow$

$\displaystyle\int\left(\frac{-1/500}{P-250} + \frac{1/500}{P-750}\right)dP = -\frac{1}{12{,}500}\,dt \quad \Leftrightarrow \quad \int\left(\frac{1}{P-250} - \frac{1}{P-750}\right)dP = \tfrac{1}{25}\,dt \quad \Leftrightarrow$

$\ln|P-250| - \ln|P-750| = \frac{1}{25}t + C \quad \Leftrightarrow \quad \ln\left|\dfrac{P-250}{P-750}\right| = \frac{1}{25}t + C \quad \Leftrightarrow \quad \left|\dfrac{P-250}{P-750}\right| = e^{t/25+C} = ke^{t/25} \quad \Leftrightarrow$

$\dfrac{P-250}{P-750} = ke^{t/25} \quad \Leftrightarrow \quad P - 250 = Pke^{t/25} - 750ke^{t/25} \quad \Leftrightarrow \quad P - Pke^{t/25} = 250 - 750ke^{t/25} \quad \Leftrightarrow$

$P(t) = \dfrac{250 - 750ke^{t/25}}{1 - ke^{t/25}}$. If $t = 0$ and $P = 200$, then $200 = \dfrac{250 - 750k}{1-k} \quad \Leftrightarrow \quad 200 - 200k = 250 - 750k \quad \Leftrightarrow$

$550k = 50 \quad \Leftrightarrow \quad k = \frac{1}{11}$. Similarly, if $t = 0$ and $P = 300$, then $k = -\frac{1}{9}$. Simplifying P with these two values of k gives us

$P(t) = \dfrac{250(3e^{t/25} - 11)}{e^{t/25} - 11}$ and $P(t) = \dfrac{750(e^{t/25} + 3)}{e^{t/25} + 9}$.

20. (a)

$c = 0$

$c = 10$

$c = 20$

$c = 21$

$c = 25$

$c = 30$

(b) For $0 \le c \le 20$, there is at least one equilibrium solution. For $c > 20$, the population always dies out.

(c) $\dfrac{dP}{dt} = 0.08P - 0.00008P^2 - c$. $\dfrac{dP}{dt} = 0 \quad \Leftrightarrow \quad P = \dfrac{-0.08 \pm \sqrt{(0.08)^2 - 4(-0.00008)(-c)}}{2(-0.00008)}$, which has at least one solution when the discriminant is nonnegative $\Rightarrow \quad 0.0064 - 0.00032c \ge 0 \quad \Leftrightarrow \quad c \le 20$. For $0 \le c \le 20$, there is at least one value of P such that $dP/dt = 0$ and hence, at least one equilibrium solution. For $c > 20$, $dP/dt < 0$ and the population always dies out.

(d) The weekly catch should be less than 20 fish per week.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. (a) $\dfrac{dP}{dt} = (kP)\left(1 - \dfrac{P}{M}\right)\left(1 - \dfrac{m}{P}\right)$. If $m < P < M$, then $dP/dt = (+)(+)(+) = +$ $\Rightarrow$ P is increasing.

If $0 < P < m$, then $dP/dt = (+)(+)(-) = -$ $\Rightarrow$ P is decreasing.

(b)

$k = 0.08$, $M = 1000$, and $m = 200$ $\Rightarrow$

$$\frac{dP}{dt} = 0.08P\left(1 - \frac{P}{1000}\right)\left(1 - \frac{200}{P}\right)$$

For $0 < P_0 < 200$, the population dies out. For $P_0 = 200$, the population is steady. For $200 < P_0 < 1000$, the population increases and approaches 1000. For $P_0 > 1000$, the population decreases and approaches 1000.

The equilibrium solutions are $P(t) = 200$ and $P(t) = 1000$.

(c) $\dfrac{dP}{dt} = kP\left(1 - \dfrac{P}{M}\right)\left(1 - \dfrac{m}{P}\right) = kP\left(\dfrac{M-P}{M}\right)\left(\dfrac{P-m}{P}\right) = \dfrac{k}{M}(M-P)(P-m)$ $\Leftrightarrow$

$\displaystyle\int \frac{dP}{(M-P)(P-m)} = \int \frac{k}{M}\,dt$. By partial fractions, $\dfrac{1}{(M-P)(P-m)} = \dfrac{A}{M-P} + \dfrac{B}{P-m}$, so

$A(P-m) + B(M-P) = 1$.

If $P = m$, $B = \dfrac{1}{M-m}$; if $P = M$, $A = \dfrac{1}{M-m}$, so $\displaystyle\frac{1}{M-m}\int\left(\frac{1}{M-P} + \frac{1}{P-m}\right)dP = \int \frac{k}{M}\,dt$ $\Rightarrow$

$\dfrac{1}{M-m}\left(-\ln|M-P| + \ln|P-m|\right) = \dfrac{k}{M}t + C$ $\Rightarrow$ $\dfrac{1}{M-m}\ln\left|\dfrac{P-m}{M-P}\right| = \dfrac{k}{M}t + C$ $\Rightarrow$

$\ln\left|\dfrac{P-m}{M-P}\right| = (M-m)\dfrac{k}{M}t + C_1$ $\Leftrightarrow$ $\dfrac{P-m}{M-P} = De^{(M-m)(k/M)t}$ $\quad [D = \pm e^{C_1}]$.

Let $t = 0$: $\dfrac{P_0 - m}{M - P_0} = D$. So $\dfrac{P-m}{M-P} = \dfrac{P_0 - m}{M - P_0}e^{(M-m)(k/M)t}$.

Solving for P, we get $P(t) = \dfrac{m(M-P_0) + M(P_0-m)e^{(M-m)(k/M)t}}{M - P_0 + (P_0 - m)e^{(M-m)(k/M)t}}$.

(d) If $P_0 < m$, then $P_0 - m < 0$. Let $N(t)$ be the numerator of the expression for $P(t)$ in part (c). Then

$N(0) = P_0(M-m) > 0$, and $P_0 - m < 0$ $\Leftrightarrow$ $\displaystyle\lim_{t\to\infty} M(P_0-m)e^{(M-m)(k/M)t} = -\infty$ $\Rightarrow$ $\displaystyle\lim_{t\to\infty} N(t) = -\infty$.

Since N is continuous, there is a number t such that $N(t) = 0$ and thus $P(t) = 0$. So the species will become extinct.

22. (a) $\dfrac{dP}{dt} = c\ln\left(\dfrac{M}{P}\right)P$ $\Rightarrow$ $\displaystyle\int \frac{dP}{P\ln(M/P)} = \int c\,dt$. Let $u = \ln\left(\dfrac{M}{P}\right) = \ln M - \ln P$ $\Rightarrow$ $du = -\dfrac{dP}{P}$ $\Rightarrow$

$\displaystyle\int -\frac{du}{u} = ct + D$ $\Rightarrow$ $\ln|u| = -ct - D$ $\Rightarrow$ $|u| = e^{-(ct+D)}$ $\Rightarrow$ $|\ln(M/P)| = e^{-(ct+D)}$ $\Rightarrow$

$\ln(M/P) = \pm e^{-(ct+D)}$. Letting $t = 0$, we get $\ln(M/P_0) = \pm e^{-D}$, so

$\ln(M/P) = \pm e^{-ct-D} = \pm e^{-ct}e^{-D} = \ln(M/P_0)e^{-ct}$ $\Rightarrow$ $M/P = e^{\ln(M/P_0)e^{-ct}}$ $\Rightarrow$

$P(t) = Me^{-\ln(M/P_0)e^{-ct}}$, $c \neq 0$.

(b) $\displaystyle\lim_{t\to\infty} P(t) = \lim_{t\to\infty} Me^{-\ln(M/P_0)e^{-ct}} = Me^{-\ln(M/P_0)\cdot 0} = Me^0 = M$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c)

The graphs look very similar. For the Gompertz function, $P(40) \approx 732$, nearly the same as the logistic function. The Gompertz function reaches $P = 900$ at $t \approx 61.7$ and its value at $t = 80$ is about 959, so it doesn't increase quite as fast as the logistic curve.

(d) $\dfrac{dP}{dt} = c\ln\left(\dfrac{M}{P}\right)P = cP(\ln M - \ln P) \quad \Rightarrow$

$$\frac{d^2P}{dt^2} = c\left[P\left(-\frac{1}{P}\frac{dP}{dt}\right) + (\ln M - \ln P)\frac{dP}{dt}\right] = c\,\frac{dP}{dt}\left[-1 + \ln\left(\frac{M}{P}\right)\right]$$
$$= c[c\ln(M/P)P][\ln(M/P) - 1] = c^2P\ln(M/P)\,[\ln(M/P) - 1]$$

Since $0 < P < M$, $P'' = 0 \quad \Leftrightarrow \quad \ln(M/P) = 1 \quad \Leftrightarrow \quad M/P = e \quad \Leftrightarrow \quad P = M/e$. $\quad P'' > 0$ for $0 < P < M/e$ and $P'' < 0$ for $M/e < P < M$, so P' is a maximum (and P grows fastest) when $P = M/e$.

Note: If $P > M$, then $\ln(M/P) < 0$, so $P''(t) > 0$.

23. (a) $dP/dt = kP\cos(rt - \phi) \quad \Rightarrow \quad (dP)/P = k\cos(rt - \phi)\,dt \quad \Rightarrow \quad \int (dP)/P = k\int \cos(rt - \phi)\,dt \quad \Rightarrow$ $\ln P = (k/r)\sin(rt - \phi) + C$. (Since this is a growth model, $P > 0$ and we can write $\ln P$ instead of $\ln|P|$.) Since $P(0) = P_0$, we obtain $\ln P_0 = (k/r)\sin(-\phi) + C = -(k/r)\sin\phi + C \quad \Rightarrow \quad C = \ln P_0 + (k/r)\sin\phi$. Thus, $\ln P = (k/r)\sin(rt - \phi) + \ln P_0 + (k/r)\sin\phi$, which we can rewrite as $\ln(P/P_0) = (k/r)[\sin(rt - \phi) + \sin\phi]$ or, after exponentiation, $P(t) = P_0e^{(k/r)[\sin(rt-\phi)+\sin\phi]}$.

(b) As k increases, the amplitude increases, but the minimum value stays the same.

Comparing values of k with $P_0 = 1$, $r = 2$, and $\phi = \pi/2$

As r increases, the amplitude and the period decrease.

Comparing values of r with $P_0 = 1$, $k = 1$, and $\phi = \pi/2$

A change in ϕ produces slight adjustments in the phase shift and amplitude.

Comparing values of ϕ with $P_0 = 1$, $k = 1$, and $r = 2$

$P(t)$ oscillates between $P_0e^{(k/r)(1+\sin\phi)}$ and $P_0e^{(k/r)(-1+\sin\phi)}$ (the extreme values are attained when $rt - \phi$ is an odd multiple of $\frac{\pi}{2}$), so $\lim\limits_{t\to\infty} P(t)$ does not exist.

24. (a) $dP/dt = kP\cos^2(rt - \phi) \quad \Rightarrow \quad (dP)/P = k\cos^2(rt - \phi)\,dt \quad \Rightarrow \quad \int (dP)/P = k\int \cos^2(rt - \phi)\,dt \quad \Rightarrow$

$\ln P = k\displaystyle\int \frac{1 + \cos(2(rt - \phi))}{2}\,dt = \frac{k}{2}t + \frac{k}{4r}\sin(2(rt - \phi)) + C$. From $P(0) = P_0$, we get

$\ln P_0 = \dfrac{k}{4r}\sin(-2\phi) + C = C - \dfrac{k}{4r}\sin 2\phi$, so $C = \ln P_0 + \dfrac{k}{4r}\sin 2\phi$ and

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\ln P = \dfrac{k}{2}\,t + \dfrac{k}{4r}\,\sin(2(rt-\phi)) + \ln P_0 + \dfrac{k}{4r}\,\sin 2\phi$. Simplifying, we get

$\ln \dfrac{P}{P_0} = \dfrac{k}{2}\,t + \dfrac{k}{4r}\,[\sin(2(rt-\phi)) + \sin 2\phi] = f(t)$, or $P(t) = P_0 e^{f(t)}$.

(b) An increase in k stretches the graph of P vertically while maintaining $P(0) = P_0$.

Comparing values of k with $P_0 = 1$, $r = 2$, and $\phi = \pi/2$

An increase in r compresses the graph of P horizontally—similar to changing the period in Exercise 23.

Comparing values of r with $P_0 = 1$, $k = 0.5$, and $\phi = \pi/2$

As in Exercise 23, a change in ϕ only makes slight adjustments in the growth of P, as shown in the figure.

Comparing values of ϕ with $P_0 = 1$, $k = 0.5$, and $r = 2$

$f'(t) = k/2 + [k/(4r)][2r\cos(2(rt-\phi))] = (k/2)[1+\cos(2(rt-\phi))] \geq 0$. Since $P(t) = P_0 e^{f(t)}$, we have $P'(t) = P_0 f'(t) e^{f(t)} \geq 0$, with equality only when $\cos(2(rt-\phi)) = -1$; that is, when $rt - \phi$ is an odd multiple of $\frac{\pi}{2}$. Therefore, $P(t)$ is an increasing function on $(0, \infty)$. P can also be written as $P(t) = P_0 e^{kt/2} e^{(k/4r)[\sin(2(rt-\phi))+\sin 2\phi]}$. The second exponential oscillates between $e^{(k/4r)(1+\sin 2\phi)}$ and $e^{(k/4r)(-1+\sin 2\phi)}$, while the first one, $e^{kt/2}$, grows without bound. So $\lim\limits_{t\to\infty} P(t) = \infty$.

25. By Equation 7, $P(t) = \dfrac{K}{1+Ae^{-kt}}$. By comparison, if $c = (\ln A)/k$ and $u = \frac{1}{2}k(t-c)$, then

$$1 + \tanh u = 1 + \frac{e^u - e^{-u}}{e^u + e^{-u}} = \frac{e^u + e^{-u}}{e^u + e^{-u}} + \frac{e^u - e^{-u}}{e^u + e^{-u}} = \frac{2e^u}{e^u + e^{-u}} \cdot \frac{e^{-u}}{e^{-u}} = \frac{2}{1+e^{-2u}}$$

and $e^{-2u} = e^{-k(t-c)} = e^{kc}e^{-kt} = e^{\ln A}e^{-kt} = Ae^{-kt}$, so

$$\tfrac{1}{2}K\big[1 + \tanh\big(\tfrac{1}{2}k(t-c)\big)\big] = \frac{K}{2}[1+\tanh u] = \frac{K}{2} \cdot \frac{2}{1+e^{-2u}} = \frac{K}{1+e^{-2u}} = \frac{K}{1+Ae^{-kt}} = P(t).$$

9.5 Linear Equations

1. $y' + x\sqrt{y} = x^2$ is not linear since it cannot be put into the standard form (1), $y' + P(x)\,y = Q(x)$.

2. $y' - x = y\tan x \quad\Leftrightarrow\quad y' + (-\tan x)y = x$ is linear since it can be put into the standard form (1), $y' + P(x)\,y = Q(x)$.

3. $ue^t = t + \sqrt{t}\,\dfrac{du}{dt} \quad\Leftrightarrow\quad \sqrt{t}\,u' - e^t u = -t \quad\Leftrightarrow\quad u' - \dfrac{e^t}{\sqrt{t}}\,u = -\sqrt{t}$ is linear since it can be put into the standard form, $u' + P(t)\,u = Q(t)$.

4. $\dfrac{dR}{dt} + t\cos R = e^{-t} \quad\Leftrightarrow\quad R' + t\cos R = e^{-t}$ is not linear since it cannot be put into the standard form $R' + P(t)\,R = Q(t)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. Comparing the given equation, $y' + y = 1$, with the general form, $y' + P(x)\,y = Q(x)$, we see that $P(x) = 1$ and the integrating factor is $I(x) = e^{\int P(x)\,dx} = e^{\int 1\,dx} = e^x$. Multiplying the differential equation by $I(x)$ gives

$e^x y' + e^x y = e^x \quad \Rightarrow \quad (e^x y)' = e^x \quad \Rightarrow \quad e^x y = \int e^x\,dx \quad \Rightarrow \quad e^x y = e^x + C \quad \Rightarrow \quad \dfrac{e^x y}{e^x} = \dfrac{e^x}{e^x} + \dfrac{C}{e^x} \quad \Rightarrow$

$y = 1 + Ce^{-x}$.

6. $y' - y = e^x \quad \Leftrightarrow \quad y' + (-1)y = e^x \quad \Rightarrow \quad P(x) = -1. \quad I(x) = e^{\int P(x)\,dx} = e^{\int -1\,dx} = e^{-x}$. Multiplying the original differential equation by $I(x)$ gives $e^{-x}y' - e^{-x}y = e^0 \quad \Rightarrow \quad (e^{-x}y)' = 1 \quad \Rightarrow \quad e^{-x}y = \int 1\,dx \quad \Rightarrow$

$e^{-x}y = x + C \quad \Rightarrow \quad y = \dfrac{x + C}{e^{-x}} \quad \Rightarrow \quad y = xe^x + Ce^x$.

7. $y' = x - y \quad \Rightarrow \quad y' + y = x \ (\star)$. $I(x) = e^{\int P(x)\,dx} = e^{\int 1\,dx} = e^x$. Multiplying the differential equation $(\star)$ by $I(x)$ gives

$e^x y' + e^x y = xe^x \quad \Rightarrow \quad (e^x y)' = xe^x \quad \Rightarrow \quad e^x y = \int xe^x\,dx \quad \Rightarrow \quad e^x y = xe^x - e^x + C$ [by parts] $\Rightarrow$

$y = x - 1 + Ce^{-x}$ [divide by e^x].

8. $4x^3 y + x^4 y' = \sin^3 x \quad \Rightarrow \quad (x^4 y)' = \sin^3 x \quad \Rightarrow \quad x^4 y = \int \sin^3 x\,dx \quad \Rightarrow$

$x^4 y = \int \sin x\,(1 - \cos^2 x)\,dx = \int (1 - u^2)(-du) \quad \begin{bmatrix} u = \cos x, \\ du = -\sin x\,dx \end{bmatrix}$

$\quad = \int (u^2 - 1)\,du = \frac{1}{3}u^3 - u + C = \frac{1}{3}u(u^2 - 3) + C = \frac{1}{3}\cos x\,(\cos^2 x - 3) + C \quad \Rightarrow$

$y = \dfrac{1}{3x^4}\cos x\,(\cos^2 x - 3) + \dfrac{C}{x^4}$

9. Since $P(x)$ is the derivative of the coefficient of y' $[P(x) = 1$ and the coefficient is $x]$, we can write the differential equation $xy' + y = \sqrt{x}$ in the easily integrable form $(xy)' = \sqrt{x} \quad \Rightarrow \quad xy = \frac{2}{3}x^{3/2} + C \quad \Rightarrow \quad y = \frac{2}{3}\sqrt{x} + C/x$.

10. $2xy' + y = 2\sqrt{x} \quad \Rightarrow \quad y' + \dfrac{1}{2x}\,y = \dfrac{1}{\sqrt{x}} \quad [x > 0] \quad \Rightarrow \quad P(x) = \dfrac{1}{2x}$.

$I(x) = e^{\int P(x)\,dx} = e^{\int 1/(2x)\,dx} = e^{(1/2)\ln|x|} = (e^{\ln x})^{1/2} = \sqrt{x}$. Multiplying the differential equation by $I(x)$ gives

$\sqrt{x}\,y' + \dfrac{1}{2\sqrt{x}}\,y = 1 \quad \Rightarrow \quad (\sqrt{x}\,y)' = 1 \quad \Rightarrow \quad \sqrt{x}\,y = \int 1\,dx \quad \Rightarrow \quad \sqrt{x}\,y = x + C \quad \Rightarrow \quad y = \dfrac{x + C}{\sqrt{x}}$.

11. $xy' - 2y = x^2 \quad \Rightarrow \quad y' - \dfrac{2}{x}\,y = x \quad \Rightarrow \quad P(x) = -\dfrac{2}{x}$.

$I(x) = e^{\int P(x)\,dx} = e^{\int -2/x\,dx} = e^{-2\ln x} \quad [x > 0] \quad = x^{-2} = \dfrac{1}{x^2}$. Multiplying the differential equation by $I(x)$ gives

$\dfrac{1}{x^2}\,y' - \dfrac{2}{x^3}\,y = \dfrac{1}{x} \quad \Rightarrow \quad \left(\dfrac{1}{x^2}\,y\right)' = \dfrac{1}{x} \quad \Rightarrow \quad \dfrac{1}{x^2}\,y = \displaystyle\int \dfrac{1}{x}\,dx \quad \Rightarrow \quad \dfrac{1}{x^2}\,y = \ln x + C \quad \Rightarrow \quad y = x^2(\ln x + C)$.

12. $y' - 3x^2 y = x^2 \quad \Rightarrow \quad P(x) = -3x^2$. $I(x) = e^{\int P(x)\,dx} = e^{\int -3x^2\,dx} = e^{-x^3}$. Multiplying the differential equation by $I(x)$ gives $e^{-x^3}y' - 3x^2 e^{-x^3}y = x^2 e^{-x^3} \quad \Rightarrow \quad \left(e^{-x^3}y\right)' = x^2 e^{-x^3} \quad \Rightarrow \quad e^{-x^3}y = \int x^2 e^{-x^3}\,dx \quad \Rightarrow$

$e^{-x^3}y = -\frac{1}{3}e^{-x^3} + C \quad \begin{bmatrix} \text{by substitution} \\ \text{with } u = -x^3 \end{bmatrix} \quad \Rightarrow \quad y = -\frac{1}{3} + Ce^{x^3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $t^2\dfrac{dy}{dt}+3ty=\sqrt{1+t^2}\quad\Rightarrow\quad y'+\dfrac{3}{t}\,y=\dfrac{\sqrt{1+t^2}}{t^2}\quad\Rightarrow\quad P(t)=\dfrac{3}{t}.$

$I(t)=e^{\int P(t)\,dt}=e^{\int 3/t\,dt}=e^{3\ln t}\quad[t>0]\quad=t^3$. Multiplying by t^3 gives $t^3y'+3t^2y=t\sqrt{1+t^2}\quad\Rightarrow$

$(t^3y)'=t\sqrt{1+t^2}\quad\Rightarrow\quad t^3y=\int t\sqrt{1+t^2}\,dt\quad\Rightarrow\quad t^3y=\frac{1}{3}(1+t^2)^{3/2}+C\quad\Rightarrow\quad y=\frac{1}{3}t^{-3}(1+t^2)^{3/2}+Ct^{-3}.$

14. $t\ln t\,\dfrac{dr}{dt}+r=te^t\quad\Rightarrow\quad\dfrac{dr}{dt}+\dfrac{1}{t\ln t}\,r=\dfrac{e^t}{\ln t}.\quad I(t)=e^{\int dt/(t\ln t)}=e^{\ln(\ln t)}=\ln t$. Multiplying by $\ln t$ gives

$\ln t\,\dfrac{dr}{dt}+\dfrac{1}{t}\,r=e^t\quad\Rightarrow\quad[(\ln t)r]'=e^t\quad\Rightarrow\quad(\ln t)r=e^t+C\quad\Rightarrow\quad r=\dfrac{e^t+C}{\ln t}.$

15. $y'+y\cos x=x\quad\Rightarrow\quad P(x)=\cos x.\ I(x)=e^{\int P(x)\,dx}=e^{\int\cos x\,dx}=e^{\sin x}$. Multiplying the differential equation by $I(x)$ gives $e^{\sin x}y'+e^{\sin x}\cos x\cdot y=xe^{\sin x}\quad\Rightarrow\quad(e^{\sin x}y)'=xe^{\sin x}\quad\Rightarrow\quad e^{\sin x}y=\int xe^{\sin x}\,dx+C\quad\Rightarrow$

$y=e^{-\sin x}\int xe^{\sin x}\,dx+Ce^{-\sin x}$. *Note:* $f(x)=xe^{\sin x}$ has an antiderivative F that is *not* an elementary function [see Section 7.5].

16. $y'+2xy=x^3\,e^{x^2}\quad\Rightarrow\quad P(x)=2x.\ I(x)=e^{\int P(x)\,dx}=e^{\int 2x\,dx}=e^{x^2}$. Multiplying the differential equation by $I(x)$ gives $e^{x^2}y'+2xe^{x^2}y=x^3e^{2x^2}\quad\Rightarrow\quad\left(e^{x^2}y\right)'=x^3e^{2x^2}\quad\Rightarrow\quad e^{x^2}y=\int x^3e^{2x^2}\,dx\quad\Rightarrow$

$e^{x^2}y=\frac{1}{4}x^2e^{2x^2}-\int\frac{1}{2}xe^{2x^2}\,dx\quad\begin{bmatrix}u=\frac{1}{4}x^2, & dv=4xe^{2x^2}\,dx\\ du=\frac{1}{2}x\,dx, & v=e^{2x^2}\end{bmatrix}\quad\Rightarrow$

$e^{x^2}y=\frac{1}{4}x^2e^{2x^2}-\frac{1}{2}\int e^z\left(\frac{1}{4}\,dz\right)\quad[z=2x^2,dz=4x\,dx]\quad\Rightarrow\quad e^{x^2}y=\frac{1}{4}x^2e^{2x^2}-\frac{1}{8}e^{2x^2}+C\quad\Rightarrow$

$y=\frac{1}{4}x^2e^{x^2}-\frac{1}{8}e^{x^2}+Ce^{-x^2}.$

17. $xy'+y=3x^2\quad\Rightarrow\quad(xy)'=3x^2\quad\Rightarrow\quad xy=\int 3x^2dx\quad\Rightarrow\quad xy=x^3+C\quad\Rightarrow\quad y=x^2+\dfrac{C}{x}$. Since $y(1)=4$,

$4=1^2+\dfrac{C}{1}\quad\Rightarrow\quad C=3$, so $y=x^2+\dfrac{3}{x}$.

18. $xy'-2y=2x\quad\Rightarrow\quad y'-\dfrac{2}{x}y=2\ (\star).\quad I(x)=e^{-2\int 1/x\,dx}=e^{-2\ln|x|}=e^{\ln|x|^{-2}}=|x|^{-2}=x^{-2}$. Multiplying $(\star)$ by

$I(x)$ gives $x^{-2}\,y'-\dfrac{2x^{-2}}{x}\,y=2x^{-2}\quad\Rightarrow\quad(x^{-2}y)'=2x^{-2}\quad\Rightarrow\quad x^{-2}y=\int 2x^{-2}\,dx\quad\Rightarrow\quad x^{-2}y=-2x^{-1}+C\quad\Rightarrow$

$y=-2x+Cx^2$. Since $y(2)=0$, $0=-2(2)+C(2)^2\quad\Rightarrow\quad C=1$, so $y=x^2-2x$.

19. $x^2y'+2xy=\ln x\quad\Rightarrow\quad(x^2y)'=\ln x\quad\Rightarrow\quad x^2y=\int\ln x\,dx\quad\Rightarrow\quad x^2y=x\ln x-x+C$ [by parts]. Since $y(1)=2$,

$1^2(2)=1\ln 1-1+C\quad\Rightarrow\quad 2=-1+C\quad\Rightarrow\quad C=3$, so $x^2y=x\ln x-x+3$, or $y=\dfrac{1}{x}\ln x-\dfrac{1}{x}+\dfrac{3}{x^2}$.

20. $t^3\dfrac{dy}{dt}+3t^2y=\cos t\quad\Rightarrow\quad(t^3y)'=\cos t\quad\Rightarrow\quad t^3y=\int\cos t\,dt\quad\Rightarrow\quad t^3y=\sin t+C$. Since $y(\pi)=0$,

$\pi^3(0)=\sin\pi+C\quad\Rightarrow\quad C=0$, so $t^3y=\sin t$, or $y=\dfrac{\sin t}{t^3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. $t\,\dfrac{du}{dt} = t^2 + 3u \;\Rightarrow\; u' - \dfrac{3}{t}u = t \;(\star)$. $I(t) = e^{\int -3/t\,dt} = e^{-3\ln|t|} = (e^{\ln|t|})^{-3} = t^{-3} \quad [t>0] \;=\; \dfrac{1}{t^3}$. Multiplying $(\star)$ by $I(t)$ gives $\dfrac{1}{t^3}\,u' - \dfrac{3}{t^4}u = \dfrac{1}{t^2} \;\Rightarrow\; \left(\dfrac{1}{t^3}u\right)' = \dfrac{1}{t^2} \;\Rightarrow\; \dfrac{1}{t^3}\,u = \displaystyle\int \dfrac{1}{t^2}\,dt \;\Rightarrow\; \dfrac{1}{t^3}\,u = -\dfrac{1}{t} + C$. Since $u(2) = 4$, $\dfrac{1}{2^3}(4) = -\dfrac{1}{2} + C \;\Rightarrow\; C = 1$, so $\dfrac{1}{t^3}\,u = -\dfrac{1}{t} + 1$, or $u = -t^2 + t^3$.

22. $xy' + y = x\ln x \;\Rightarrow\; (x\,y)' = x\ln x \;\Rightarrow\; xy = \int x\ln x\,dx \;\Rightarrow\; xy = \frac{1}{2}x^2\ln x - \frac{1}{4}x^2 + C \;\left[\begin{array}{c}\text{by parts}\\ \text{with } u = \ln x\end{array}\right] \;\Rightarrow$ $y = \dfrac{1}{2}x\ln x - \dfrac{1}{4}x + \dfrac{C}{x}$. $y(1) = 0 \;\Rightarrow\; 0 = 0 - \dfrac{1}{4} + C \;\Rightarrow\; C = \dfrac{1}{4}$, so $y = \dfrac{1}{2}x\ln x - \dfrac{1}{4}x + \dfrac{1}{4x}$.

23. $xy' = y + x^2\sin x \;\Rightarrow\; y' - \dfrac{1}{x}\,y = x\sin x$. $I(x) = e^{\int(-1/x)\,dx} = e^{-\ln x} = e^{\ln x^{-1}} = \dfrac{1}{x}$.

Multiplying by $\dfrac{1}{x}$ gives $\dfrac{1}{x}\,y' - \dfrac{1}{x^2}\,y = \sin x \;\Rightarrow\; \left(\dfrac{1}{x}\,y\right)' = \sin x \;\Rightarrow\; \dfrac{1}{x}\,y = -\cos x + C \;\Rightarrow\; y = -x\cos x + Cx$.

$y(\pi) = 0 \;\Rightarrow\; -\pi\cdot(-1) + C\pi = 0 \;\Rightarrow\; C = -1$, so $y = -x\cos x - x$.

24. $(x^2+1)\,\dfrac{dy}{dx} + 3x(y-1) = 0 \;\Rightarrow\; (x^2+1)\,y' + 3xy = 3x \;\Rightarrow\; y' + \dfrac{3x}{x^2+1}\,y = \dfrac{3x}{x^2+1}$.

$I(x) = e^{\int 3x/(x^2+1)\,dx} = e^{(3/2)\ln|x^2+1|} = \left(e^{\ln(x^2+1)}\right)^{3/2} = (x^2+1)^{3/2}$. Multiplying by $(x^2+1)^{3/2}$ gives

$(x^2+1)^{3/2}\,y' + 3x(x^2+1)^{1/2}\,y = 3x(x^2+1)^{1/2} \;\Rightarrow\; \left[(x^2+1)^{3/2}y\right]' = 3x(x^2+1)^{1/2} \;\Rightarrow$

$(x^2+1)^{3/2}\,y = \int 3x(x^2+1)^{1/2}\,dx = (x^2+1)^{3/2} + C \;\Rightarrow\; y = 1 + C(x^2+1)^{-3/2}$. Since $y(0) = 2$, we have $2 = 1 + C(1) \;\Rightarrow\; C = 1$ and hence, $y = 1 + (x^2+1)^{-3/2}$.

25. $xy' + 2y = e^x \;\Rightarrow\; y' + \dfrac{2}{x}\,y = \dfrac{e^x}{x}$.

$I(x) = e^{\int(2/x)\,dx} = e^{2\ln|x|} = \left(e^{\ln|x|}\right)^2 = |x|^2 = x^2$.

Multiplying by $I(x)$ gives $x^2\,y' + 2xy = xe^x \;\Rightarrow\; (x^2y)' = xe^x \;\Rightarrow$

$x^2y = \int xe^x\,dx = (x-1)e^x + C$ [by parts] $\Rightarrow$

$y = [(x-1)e^x + C]/x^2$. The graphs for $C = -5, -3, -1, 1, 3, 5$, and 7 are shown. $C = 1$ is a transitional value. For $C < 1$, there is an inflection point and for $C > 1$, there is a local minimum. As $|C|$ gets larger, the "branches" get further from the origin.

26. $xy' = x^2 + 2y \;\Leftrightarrow\; xy' - 2y = x^2 \;\Leftrightarrow\; y' - \dfrac{2}{x}\,y = x$.

$I(x) = e^{\int -2/x\,dx} = e^{-2\ln|x|} = (e^{\ln|x|})^{-2} = |x|^{-2} = \dfrac{1}{x^2}$. Multiplying by $I(x)$ gives $\dfrac{1}{x^2}\,y' - \dfrac{2}{x^3}\,y = \dfrac{1}{x} \;\Rightarrow\; \left(\dfrac{1}{x^2}\,y\right)' = \dfrac{1}{x} \;\Rightarrow\; \dfrac{1}{x^2}\,y = \displaystyle\int \dfrac{1}{x}\,dx \;\Rightarrow$

$\dfrac{1}{x^2}\,y = \ln|x| + C \;\Rightarrow\; y = (\ln|x| + C)x^2$. For all values of C, as $|x| \to 0$,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$y \to 0$, and as $|x| \to \infty$, $y \to \infty$. As $|x|$ increases from 0, the function decreases and attains an absolute minimum. The inflection points, absolute minimums, and x-intercepts all move farther from the origin as C decreases.

27. Setting $u = y^{1-n}$, $\dfrac{du}{dx} = (1-n)\,y^{-n}\dfrac{dy}{dx}$ or $\dfrac{dy}{dx} = \dfrac{y^n}{1-n}\dfrac{du}{dx} = \dfrac{u^{n/(1-n)}}{1-n}\dfrac{du}{dx}$. Then the Bernoulli differential equation becomes $\dfrac{u^{n/(1-n)}}{1-n}\dfrac{du}{dx} + P(x)u^{1/(1-n)} = Q(x)u^{n/(1-n)}$ or $\dfrac{du}{dx} + (1-n)P(x)u = Q(x)(1-n)$.

28. Here $xy' + y = -xy^2 \quad\Rightarrow\quad y' + \dfrac{y}{x} = -y^2$, so $n = 2$, $P(x) = \dfrac{1}{x}$ and $Q(x) = -1$. Setting $u = y^{-1}$, u satisfies $u' - \dfrac{1}{x}u = 1$. Then $I(x) = e^{\int(-1/x)dx} = \dfrac{1}{x}$ (for $x > 0$) and $u = x\left(\displaystyle\int \frac{1}{x}\,dx + C\right) = x(\ln|x| + C)$. Thus, $y = \dfrac{1}{x(C + \ln|x|)}$.

29. Here $y' + \dfrac{2}{x}\,y = \dfrac{y^3}{x^2}$, so $n = 3$, $P(x) = \dfrac{2}{x}$ and $Q(x) = \dfrac{1}{x^2}$. Setting $u = y^{-2}$, u satisfies $u' - \dfrac{4u}{x} = -\dfrac{2}{x^2}$.

Then $I(x) = e^{\int(-4/x)\,dx} = x^{-4}$ and $u = x^4\left(\displaystyle\int -\frac{2}{x^6}\,dx + C\right) = x^4\left(\frac{2}{5x^5} + C\right) = Cx^4 + \dfrac{2}{5x}$.

Thus, $y = \pm\left(Cx^4 + \dfrac{2}{5x}\right)^{-1/2}$.

30. $xy'' + 2y' = 12x^2$ and $u = y' \quad\Rightarrow\quad xu' + 2u = 12x^2 \quad\Rightarrow\quad u' + \dfrac{2}{x}\,u = 12x$.

$I(x) = e^{\int(2/x)\,dx} = e^{2\ln|x|} = \left(e^{\ln|x|}\right)^2 = |x|^2 = x^2$. Multiplying the last differential equation by x^2 gives $x^2u' + 2xu = 12x^3 \quad\Rightarrow\quad (x^2u)' = 12x^3 \quad\Rightarrow\quad x^2u = \int 12x^3\,dx = 3x^4 + C \quad\Rightarrow\quad u = 3x^2 + C/x^2 \quad\Rightarrow$ $y' = 3x^2 + C/x^2 \quad\Rightarrow\quad y = x^3 - C/x + D$.

31. (a) $2\dfrac{dI}{dt} + 10I = 40$ or $\dfrac{dI}{dt} + 5I = 20$. Then the integrating factor is $e^{\int 5\,dt} = e^{5t}$. Multiplying the differential equation by the integrating factor gives $e^{5t}\dfrac{dI}{dt} + 5Ie^{5t} = 20e^{5t} \quad\Rightarrow\quad (e^{5t}I)' = 20e^{5t} \quad\Rightarrow$ $I(t) = e^{-5t}\left[\int 20e^{5t}\,dt + C\right] = 4 + Ce^{-5t}$. But $0 = I(0) = 4 + C$, so $I(t) = 4 - 4e^{-5t}$.

(b) $I(0.1) = 4 - 4e^{-0.5} \approx 1.57$ A

32. (a) $\dfrac{dI}{dt} + 20I = 40\sin 60t$, so the integrating factor is e^{20t}. Multiplying the differential equation by the integrating factor gives $e^{20t}\dfrac{dI}{dt} + 20Ie^{20t} = 40e^{20t}\sin 60t \quad\Rightarrow\quad (e^{20t}I)' = 40e^{20t}\sin 60t \quad\Rightarrow$

$$\begin{aligned} I(t) &= e^{-20t}\left[\int 40e^{20t}\sin 60t\,dt + C\right] = e^{-20t}\left[40e^{20t}\left(\tfrac{1}{4000}\right)(20\sin 60t - 60\cos 60t)\right] + Ce^{-20t} \\ &= \frac{\sin 60t - 3\cos 60t}{5} + Ce^{-20t} \end{aligned}$$

But $1 = I(0) = -\frac{3}{5} + C$, so $I(t) = \dfrac{\sin 60t - 3\cos 60t\ + 8e^{-20t}}{5}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $I(0.1) = \dfrac{\sin 6 - 3\cos 6 + 8e^{-2}}{5} \approx -0.42$ A

(c)

33. $5\dfrac{dQ}{dt} + 20Q = 60$ with $Q(0) = 0$ C. Then the integrating factor is $e^{\int 4\,dt} = e^{4t}$, and multiplying the differential equation by the integrating factor gives $e^{4t}\dfrac{dQ}{dt} + 4e^{4t}Q = 12e^{4t} \;\Rightarrow\; (e^{4t}Q)' = 12e^{4t} \;\Rightarrow$ $Q(t) = e^{-4t}\left[\int 12e^{4t}\,dt + C\right] = 3 + Ce^{-4t}$. But $0 = Q(0) = 3 + C$ so $Q(t) = 3(1 - e^{-4t})$ is the charge at time t and $I = dQ/dt = 12e^{-4t}$ is the current at time t.

34. $2\dfrac{dQ}{dt} + 100Q = 10\sin 60t$ or $\dfrac{dQ}{dt} + 50Q = 5\sin 60t$. Then the integrating factor is $e^{\int 50\,dt} = e^{50t}$, and multiplying the differential equation by the integrating factor gives $e^{50t}\dfrac{dQ}{dt} + 50e^{50t}Q = 5e^{50t}\sin 60t \;\Rightarrow\; (e^{50t}Q)' = 5e^{50t}\sin 60t \;\Rightarrow$

$$\begin{aligned} Q(t) &= e^{-50t}\left[\int 5e^{50t}\sin 60t\,dt + C\right] = e^{-50t}\left[5e^{50t}\left(\tfrac{1}{6100}\right)(50\sin 60t - 60\cos 60t)\right] + Ce^{-50t} \\ &= \tfrac{1}{122}(5\sin 60t - 6\cos 60t) + Ce^{-50t} \end{aligned}$$

But $0 = Q(0) = -\frac{6}{122} + C$ so $C = \frac{3}{61}$ and $Q(t) = \dfrac{5\sin 60t - 6\cos 60t}{122} + \dfrac{3e^{-50t}}{61}$ is the charge at time t, while the current is $I(t) = \dfrac{dQ}{dt} = \dfrac{150\cos 60t + 180\sin 60t - 150e^{-50t}}{61}$.

35. $\dfrac{dP}{dt} = k[M - P(t)] \;\Rightarrow\; \dfrac{dP}{dt} + kP = kM$ $(\star)$, so $I(t) = e^{\int k\,dt} = e^{kt}$.

Multiplying $(\star)$ by $I(t)$ gives $e^{kt}\dfrac{dP}{dt} + kPe^{kt} = kMe^{kt} \;\Rightarrow$

$(e^{kt}P)' = kMe^{kt} \;\Rightarrow\; P(t) = e^{-kt}\left(\int kMe^{kt}dt + C\right) = M + Ce^{-kt}$, $k > 0$.

Furthermore, it is reasonable to assume that $0 \le P(0) \le M$, so $-M \le C \le 0$.

36. Since $P(0) = 0$, we have $P(t) = M(1 - e^{-kt})$. If $P_1(t)$ is Jim's learning curve, then $P_1(1) = 25$ and $P_1(2) = 45$. Hence, $25 = M_1(1 - e^{-k})$ and $45 = M_1(1 - e^{-2k})$, so $1 - 25/M_1 = e^{-k}$ or $k = -\ln\left(1 - \dfrac{25}{M_1}\right) = \ln\left(\dfrac{M_1}{M_1 - 25}\right)$. But $45 = M_1(1 - e^{-2k})$ so $45 = M_1\left[1 - \left(\dfrac{M_1 - 25}{M_1}\right)^2\right]$ or $45 = \dfrac{50M_1 - 625}{M_1}$. Thus, $M_1 = 125$ is the maximum number of units per hour Jim is capable of processing. Similarly, if $P_2(t)$ is Mark's learning curve, then $P_2(1) = 35$ and $P_2(2) = 50$. So $k = \ln\left(\dfrac{M_2}{M_2 - 35}\right)$ and $50 = M_2\left[1 - \left(\dfrac{M_2 - 35}{M_2}\right)^2\right]$ or $M_2 = 61.25$. Hence the maximum number of units per hour for Mark is approximately 61. Another approach would be to use the midpoints of the intervals so that $P_1(0.5) = 25$ and $P_1(1.5) = 45$. Doing so gives us $M_1 \approx 52.6$ and $M_2 \approx 51.8$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. $y(0) = 0$ kg. Salt is added at a rate of $\left(0.4\ \dfrac{\text{kg}}{\text{L}}\right)\left(5\ \dfrac{\text{L}}{\text{min}}\right) = 2\ \dfrac{\text{kg}}{\text{min}}$. Since solution is drained from the tank at a rate of 3 L/min, but salt solution is added at a rate of 5 L/min, the tank, which starts out with 100 L of water, contains $(100 + 2t)$ L of liquid after t min. Thus, the salt concentration at time t is $\dfrac{y(t)}{100+2t}\ \dfrac{\text{kg}}{\text{L}}$. Salt therefore leaves the tank at a rate of $\left(\dfrac{y(t)}{100+2t}\ \dfrac{\text{kg}}{\text{L}}\right)\left(3\ \dfrac{\text{L}}{\text{min}}\right) = \dfrac{3y}{100+2t}\ \dfrac{\text{kg}}{\text{min}}$. Combining the rates at which salt enters and leaves the tank, we get $\dfrac{dy}{dt} = 2 - \dfrac{3y}{100+2t}$. Rewriting this equation as $\dfrac{dy}{dt} + \left(\dfrac{3}{100+2t}\right) y = 2$, we see that it is linear.

$$I(t) = \exp\left(\int \frac{3\,dt}{100+2t}\right) = \exp\left(\tfrac{3}{2}\ln(100+2t)\right) = (100+2t)^{3/2}$$

Multiplying the differential equation by $I(t)$ gives $(100+2t)^{3/2}\dfrac{dy}{dt} + 3(100+2t)^{1/2}y = 2(100+2t)^{3/2} \quad \Rightarrow$ $[(100+2t)^{3/2}y]' = 2(100+2t)^{3/2} \quad \Rightarrow \quad (100+2t)^{3/2}y = \frac{2}{5}(100+2t)^{5/2} + C \quad \Rightarrow$ $y = \frac{2}{5}(100+2t) + C(100+2t)^{-3/2}$. Now $0 = y(0) = \frac{2}{5}(100) + C \cdot 100^{-3/2} = 40 + \frac{1}{1000}C \quad \Rightarrow \quad C = -40{,}000$, so $y = \left[\frac{2}{5}(100+2t) - 40{,}000(100+2t)^{-3/2}\right]$ kg. From this solution (no pun intended), we calculate the salt concentration at time t to be $C(t) = \dfrac{y(t)}{100+2t} = \left[\dfrac{-40{,}000}{(100+2t)^{5/2}} + \dfrac{2}{5}\right]\ \dfrac{\text{kg}}{\text{L}}$. In particular, $C(20) = \dfrac{-40{,}000}{140^{5/2}} + \dfrac{2}{5} \approx 0.2275\ \dfrac{\text{kg}}{\text{L}}$ and $y(20) = \frac{2}{5}(140) - 40{,}000(140)^{-3/2} \approx 31.85$ kg.

38. Let $y(t)$ denote the amount of chlorine in the tank at time t (in seconds). $y(0) = (0.05\ \text{g/L})\,(400\ \text{L}) = 20$ g. The amount of liquid in the tank at time t is $(400 - 6t)$ L since 4 L of water enters the tank each second and 10 L of liquid leaves the tank each second. Thus, the concentration of chlorine at time t is $\dfrac{y(t)}{400-6t}\ \dfrac{\text{g}}{\text{L}}$. Chlorine doesn't enter the tank, but it leaves at a rate of $\left[\dfrac{y(t)}{400-6t}\ \dfrac{\text{g}}{\text{L}}\right]\left[10\ \dfrac{\text{L}}{\text{s}}\right] = \dfrac{10\,y(t)}{400-6t}\ \dfrac{\text{g}}{\text{s}} = \dfrac{5\,y(t)}{200-3t}\ \dfrac{\text{g}}{\text{s}}$. Therefore, $\dfrac{dy}{dt} = -\dfrac{5y}{200-3t} \quad \Rightarrow \quad \displaystyle\int \frac{dy}{y} = \int \frac{-5\,dt}{200-3t} \quad \Rightarrow$ $\ln y = \frac{5}{3}\ln(200-3t) + C \quad \Rightarrow \quad y = \exp\left(\frac{5}{3}\ln(200-3t) + C\right) = e^C(200-3t)^{5/3}$. Now $20 = y(0) = e^C \cdot 200^{5/3} \quad \Rightarrow$ $e^C = \dfrac{20}{200^{5/3}}$, so $y(t) = 20\dfrac{(200-3t)^{5/3}}{200^{5/3}} = 20(1-0.015t)^{5/3}$ g for $0 \le t \le 66\frac{2}{3}$ s, at which time the tank is empty.

39. (a) $m\dfrac{dv}{dt} = mg - cv \quad \Rightarrow \quad \dfrac{dv}{dt} + \dfrac{c}{m}v = g$ and $I(t) = e^{\int (c/m)\,dt} = e^{(c/m)t}$, and multiplying the last differential equation by $I(t)$ gives $e^{(c/m)t}\dfrac{dv}{dt} + \dfrac{vce^{(c/m)t}}{m} = ge^{(c/m)t} \quad \Rightarrow \quad \left[e^{(c/m)t}v\right]' = ge^{(c/m)t}$. Hence, $v(t) = e^{-(c/m)t}\left[\int ge^{(c/m)t}\,dt + K\right] = mg/c + Ke^{-(c/m)t}$. But the object is dropped from rest, so $v(0) = 0$ and $K = -mg/c$. Thus, the velocity at time t is $v(t) = (mg/c)[1 - e^{-(c/m)t}]$.

(b) $\displaystyle\lim_{t\to\infty} v(t) = mg/c$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $s(t) = \int v(t)\,dt = (mg/c)[t + (m/c)e^{-(c/m)t}] + c_1$ where $c_1 = s(0) - m^2g/c^2$.

$s(0)$ is the initial position, so $s(0) = 0$ and $s(t) = (mg/c)[t + (m/c)e^{-(c/m)t}] - m^2g/c^2$.

40. $v = (mg/c)(1 - e^{-ct/m}) \quad \Rightarrow$

$$\frac{dv}{dm} = \frac{mg}{c}\left(0 - e^{-ct/m} \cdot \frac{ct}{m^2}\right) + \frac{g}{c}\,(1 - e^{-ct/m}) \cdot 1 = -\frac{gt}{m}\,e^{-ct/m} + \frac{g}{c} - \frac{g}{c}\,e^{-ct/m}$$

$$= \frac{g}{c}\left(1 - e^{-ct/m} - \frac{ct}{m}\,e^{-ct/m}\right) \quad \Rightarrow$$

$\dfrac{c}{g}\dfrac{dv}{dm} = 1 - \left(1 + \dfrac{ct}{m}\right)e^{-ct/m} = 1 - \dfrac{1 + ct/m}{e^{ct/m}} = 1 - \dfrac{1+Q}{e^Q}$, where $Q = \dfrac{ct}{m} \geq 0$. Since $e^Q > 1 + Q$ for all $Q > 0$, it follows that $dv/dm > 0$ for $t > 0$. In other words, for all $t > 0$, v increases as m increases.

41. (a) $z = \dfrac{1}{P} \quad \Rightarrow \quad P = \dfrac{1}{z} \quad \Rightarrow \quad P' = -\dfrac{z'}{z^2}$. Substituting into $P' = kP(1 - P/M)$ gives us $-\dfrac{z'}{z^2} = k\,\dfrac{1}{z}\left(1 - \dfrac{1}{zM}\right) \quad \Rightarrow$

$z' = -kz\left(1 - \dfrac{1}{zM}\right) \quad \Rightarrow \quad z' = -kz + \dfrac{k}{M} \quad \Rightarrow \quad z' + kz = \dfrac{k}{M}$ $(\star)$.

(b) The integrating factor is $e^{\int k\,dt} = e^{kt}$. Multiplying $(\star)$ by e^{kt} gives $e^{kt}z' + ke^{kt}z = \dfrac{ke^{kt}}{M} \quad \Rightarrow \quad (e^{kt}z)' = \dfrac{k}{M}\,e^{kt} \quad \Rightarrow$

$e^{kt}z = \displaystyle\int \frac{k}{M}\,e^{kt}\,dt \quad \Rightarrow \quad e^{kt}z = \frac{1}{M}\,e^{kt} + C \quad \Rightarrow \quad z = \frac{1}{M} + Ce^{-kt}$. Since $P = \dfrac{1}{z}$, we have

$P = \dfrac{1}{\dfrac{1}{M} + Ce^{-kt}} \quad \Rightarrow \quad P = \dfrac{M}{1 + MCe^{-kt}}$, which agrees with Equation 9.4.7, $P = \dfrac{M}{1 + Ae^{-kt}}$, when $MC = A$.

42. (a) $z = \dfrac{1}{P} \quad \Rightarrow \quad P = \dfrac{1}{z} \quad \Rightarrow \quad P' = \dfrac{-z'}{z^2}$. Substituting into $\dfrac{dP}{dt} = k(t)P\left(1 - \dfrac{P}{M(t)}\right)$ gives us

$-\dfrac{z'}{z^2} = \dfrac{k(t)}{z}\left(1 - \dfrac{1}{M(t)z}\right) \quad \Rightarrow \quad z' = -k(t)z\left(1 - \dfrac{1}{M(t)z}\right) \quad \Rightarrow \quad z' = -k(t)z + \dfrac{k(t)}{M(t)} \quad \Rightarrow$

$\dfrac{dz}{dt} + k(t)z = \dfrac{k(t)}{M(t)}$ $(\star)$.

(b) The integrating factor is $e^{K(t)}$, where $K(t) = \int_0^t k(s)\,ds$, so that $K'(t) = k(t)$. Multiplying $(\star)$ by

$e^{K(t)}$ gives $e^{K(t)}\dfrac{dz}{dt} + e^{K(t)}k(t)z = \dfrac{e^{K(t)}k(t)}{M(t)} \quad \Rightarrow \quad (e^{K(t)}z)' = \dfrac{K'(t)e^{K(t)}}{M(t)} \quad \Rightarrow$

$e^{K(t)}z = \displaystyle\int_0^t \frac{K'(s)e^{K(s)}}{M(s)}\,ds + C$, so $P = \dfrac{1}{z} = \dfrac{e^{K(t)}}{\displaystyle\int_0^t \frac{K'(s)e^{K(s)}}{M(s)}\,ds + C}$. Now suppose that M is a constant. Then

$P(t) = \dfrac{Me^{K(t)}}{\int_0^t K'(s)e^{K(s)}\,ds + CM} = \dfrac{Me^{K(t)}}{e^{K(t)} + CM} = \dfrac{M}{1 + CMe^{-K(t)}}$. If $\int_0^\infty k(t)\,dt = \infty$, then $\lim\limits_{t\to\infty} K(t) = \infty$, so

$\lim\limits_{t\to\infty} P(t) = \dfrac{M}{1 + CM \lim\limits_{t\to\infty} e^{-K(t)}} = \dfrac{M}{1 + CM \cdot 0} = M$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) If k is constant, but M varies, then $K(t) = kt$ and we get $e^{kt}z = \int_0^t \frac{ke^{ks}}{M(s)}\,ds + C \quad \Rightarrow$

$z(t) = \dfrac{\int_0^t \frac{ke^{ks}}{M(s)}\,ds + C}{e^{kt}} \quad \Rightarrow \quad z(t) = e^{-kt}\int_0^t \frac{ke^{ks}}{M(s)}\,ds + Ce^{-kt}$. Suppose $M(t)$ has a limit as $t \to \infty$, say $\lim\limits_{t\to\infty} M(t) = L$. Then

$$\lim_{t\to\infty} P(t) = \lim_{t\to\infty} \frac{1}{z(t)} = \lim_{t\to\infty} \frac{e^{kt}}{\int_0^t \frac{ke^{ks}}{M(s)}\,ds + C} = \lim_{t\to\infty} \frac{ke^{kt}}{\frac{ke^{kt}}{M(t)} + 0} \begin{bmatrix}\text{l'Hospital's}\\ \text{and FTC 1}\end{bmatrix} = \lim_{t\to\infty} M(t) = L.$$

APPLIED PROJECT Which Is Faster, Going Up or Coming Down?

1. $mv' = -pv - mg \quad \Rightarrow \quad m\dfrac{dv}{dt} = -(pv + mg) \quad \Rightarrow \quad \displaystyle\int \frac{dv}{pv + mg} = \int -\frac{1}{m}\,dt \quad \Rightarrow$

$\dfrac{1}{p}\ln(pv + mg) = -\dfrac{1}{m}t + C \quad [pv + mg > 0]$. At $t = 0$, $v = v_0$, so $C = \dfrac{1}{p}\ln(pv_0 + mg)$.

Thus, $\dfrac{1}{p}\ln(pv + mg) = -\dfrac{1}{m}t + \dfrac{1}{p}\ln(pv_0 + mg) \quad \Rightarrow \quad \ln(pv + mg) = -\dfrac{p}{m}t + \ln(pv_0 + mg) \quad \Rightarrow$

$pv + mg = e^{-pt/m}(pv_0 + mg) \quad \Rightarrow \quad pv = (pv_0 + mg)e^{-pt/m} - mg \quad \Rightarrow \quad v(t) = \left(v_0 + \dfrac{mg}{p}\right)e^{-pt/m} - \dfrac{mg}{p}$.

2. $y(t) = \int v(t)\,dt = \displaystyle\int \left[\left(v_0 + \frac{mg}{p}\right)e^{-pt/m} - \frac{mg}{p}\right] dt = \left(v_0 + \frac{mg}{p}\right)e^{-pt/m}\left(-\frac{m}{p}\right) - \frac{mg}{p}t + C$.

At $t = 0$, $y = 0$, so $C = \left(v_0 + \dfrac{mg}{p}\right)\dfrac{m}{p}$. Thus,

$$y(t) = \left(v_0 + \frac{mg}{p}\right)\frac{m}{p} - \left(v_0 + \frac{mg}{p}\right)\frac{m}{p}e^{-pt/m} - \frac{mgt}{p} = \left(v_0 + \frac{mg}{p}\right)\frac{m}{p}\left(1 - e^{-pt/m}\right) - \frac{mgt}{p}$$

3. $v(t) = 0 \quad \Rightarrow \quad \dfrac{mg}{p} = \left(v_0 + \dfrac{mg}{p}\right)e^{-pt/m} \quad \Rightarrow \quad e^{pt/m} = \dfrac{pv_0}{mg} + 1 \quad \Rightarrow \quad \dfrac{pt}{m} = \ln\left(\dfrac{pv_0}{mg} + 1\right) \quad \Rightarrow$

$t_1 = \dfrac{m}{p}\ln\left(\dfrac{mg + pv_0}{mg}\right)$. With $m = 1$, $v_0 = 20$, $p = \frac{1}{10}$, and $g = 9.8$, we have $t_1 = 10\ln\left(\frac{11.8}{9.8}\right) \approx 1.86$ s.

4.

The figure shows the graph of $y = 1180(1 - e^{-0.1t}) - 98t$. The zeros are at $t = 0$ and $t_2 \approx 3.84$. Thus, $t_1 - 0 \approx 1.86$ and $t_2 - t_1 \approx 1.98$. So the time it takes to come down is about 0.12 s longer than the time it takes to go up; hence, going up is faster.

5. $y(2t_1) = \left(v_0 + \dfrac{mg}{p}\right)\dfrac{m}{p}(1 - e^{-2pt_1/m}) - \dfrac{mg}{p}\cdot 2t_1$

$$= \left(\frac{pv_0 + mg}{p}\right)\frac{m}{p}\left[1 - (e^{pt_1/m})^{-2}\right] - \frac{mg}{p}\cdot 2\frac{m}{p}\ln\left(\frac{pv_0 + mg}{mg}\right)$$

Substituting $x = e^{pt_1/m} = \dfrac{pv_0}{mg} + 1 = \dfrac{pv_0 + mg}{mg}$ (from Problem 3), we get

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$y(2t_1) = \left(x \cdot \dfrac{mg}{p}\right)\dfrac{m}{p}(1 - x^{-2}) - \dfrac{m^2 g}{p^2} \cdot 2\ln x = \dfrac{m^2 g}{p^2}\left(x - \dfrac{1}{x} - 2\ln x\right)$. Now $p > 0$, $m > 0$, $t_1 > 0 \quad \Rightarrow$

$x = e^{pt_1/m} > e^0 = 1. \quad f(x) = x - \dfrac{1}{x} - 2\ln x \quad \Rightarrow \quad f'(x) = 1 + \dfrac{1}{x^2} - \dfrac{2}{x} = \dfrac{x^2 - 2x + 1}{x^2} = \dfrac{(x-1)^2}{x^2} > 0$

for $x > 1 \quad \Rightarrow \quad f(x)$ is increasing for $x > 1$. Since $f(1) = 0$, it follows that $f(x) > 0$ for every $x > 1$. Therefore, $y(2t_1) = \dfrac{m^2 g}{p^2} f(x)$ is positive, which means that the ball has not yet reached the ground at time $2t_1$. This tells us that the time spent going up is always less than the time spent coming down, so *ascent is faster*.

9.6 Predator-Prey Systems

1. (a) $dx/dt = -0.05x + 0.0001xy$. If $y = 0$, we have $dx/dt = -0.05x$, which indicates that in the absence of y, x declines at a rate proportional to itself. So x represents the predator population and y represents the prey population. The growth of the prey population, $0.1y$ (from $dy/dt = 0.1y - 0.005xy$), is restricted only by encounters with predators (the term $-0.005xy$). The predator population increases only through the term $0.0001xy$; that is, by encounters with the prey and not through additional food sources.

(b) $dy/dt = -0.015y + 0.00008xy$. If $x = 0$, we have $dy/dt = -0.015y$, which indicates that in the absence of x, y would decline at a rate proportional to itself. So y represents the predator population and x represents the prey population. The growth of the prey population, $0.2x$ (from $dx/dt = 0.2x - 0.0002x^2 - 0.006xy = 0.2x(1 - 0.001x) - 0.006xy$), is restricted by a carrying capacity of 1000 [from the term $1 - 0.001x = 1 - x/1000$] and by encounters with predators (the term $-0.006xy$). The predator population increases only through the term $0.00008xy$; that is, by encounters with the prey and not through additional food sources.

2. (a) $dx/dt = 0.12x - 0.0006x^2 + 0.00001xy$. $dy/dt = 0.08y + 0.00004xy$.

The xy terms represent encounters between the two species x and y. An increase in y makes dx/dt (the growth rate of x) larger due to the positive term $0.00001xy$. An increase in x makes dy/dt (the growth rate of y) larger due to the positive term $0.00004xy$. Hence, the system describes a cooperation model.

(b) $dx/dt = 0.15x - 0.0002x^2 - 0.0006xy = 0.15x(1 - x/750) - 0.0006xy$.

$dy/dt = 0.2y - 0.00008y^2 - 0.0002xy = 0.2y(1 - y/2500) - 0.0002xy$.

The system shows that x and y have carrying capacities of 750 and 2500. An increase in x reduces the growth rate of y due to the negative term $-0.0002xy$. An increase in y reduces the growth rate of x due to the negative term $-0.0006xy$. Hence, the system describes a competition model.

3. (a) $dx/dt = 0.5x - 0.004x^2 - 0.001xy = 0.5x(1 - x/125) - 0.001xy$.

$dy/dt = 0.4y - 0.001y^2 - 0.002xy = 0.4y(1 - y/400) - 0.002xy$.

The system shows that x and y have carrying capacities of 125 and 400. An increase in x reduces the growth rate of y due to the negative term $-0.002xy$. An increase in y reduces the growth rate of x due to the negative term $-0.001xy$. Hence the system describes a competition model.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $dx/dt = 0 \quad \Rightarrow \quad x(0.5 - 0.004x - 0.001y) = 0 \quad \Rightarrow \quad x(500 - 4x - y) = 0$ **(1)** and $dy/dt = 0 \quad \Rightarrow$ $y(0.4 - 0.001y - 0.002x) = 0 \quad \Rightarrow \quad y(400 - y - 2x) = 0$ **(2).**

From **(1)** and **(2)**, we get four equilibrium solutions.

(i) $x = 0$ and $y = 0$: If the populations are zero, there is no change.

(ii) $x = 0$ and $400 - y - 2x = 0 \quad \Rightarrow \quad x = 0$ and $y = 400$: In the absence of an x-population, the y-population stabilizes at 400.

(iii) $500 - 4x - y = 0$ and $y = 0 \quad \Rightarrow \quad x = 125$ and $y = 0$: In the absence of y-population, the x-population stabilizes at 125.

(iv) $500 - 4x - y = 0$ and $400 - y - 2x = 0 \quad \Rightarrow \quad y = 500 - 4x$ and $y = 400 - 2x \quad \Rightarrow \quad 500 - 4x = 400 - 2x \quad \Rightarrow$ $100 = 2x \quad \Rightarrow \quad x = 50$ and $y = 300$: A y-population of 300 is just enough to support a constant x-population of 50.

4. Let $L(t)$, $H(t)$, and $W(t)$ represent the populations of lynx, hares, and willows at time t. Let the k_i's and the c_i's denote positive constants, so that a plus sign means an increase and a minus sign means a decrease in the corresponding growth rate. "In the absence of hares, the willow population will grow exponentially and the lynx population will decay exponentially" gives us $dW/dt = +k_1W$ and $dL/dt = -k_2L$. "In the absence of lynx and willow, the hare population will decay exponentially" gives us $dH/dt = -k_3H$. "Lynx eat snowshoe hares and snowshoe hares eat woody plants like willows" gives us encounters that lynx win, hares lose and win, and willows lose. In terms of growth rates, this means that $dL/dt = +c_1LH$, $dH/dt = -c_2LH + c_3HW$, and $dW/dt = -c_4HW$. Putting this information together gives us the following system of differential equations.

$$\begin{aligned} dL/dt &= -k_2L + c_1LH \\ dH/dt &= -k_3H - c_2LH + c_3HW \\ dW/dt &= +k_1W - c_4HW \end{aligned}$$

5. (a) At $t = 0$, there are about 300 rabbits and 100 foxes. At $t = t_1$, the number of foxes reaches a minimum of about 20 while the number of rabbits is about 1000. At $t = t_2$, the number of rabbits reaches a maximum of about 2400, while the number of foxes rebounds to 100. At $t = t_3$, the number of rabbits decreases to about 1000 and the number of foxes reaches a maximum of about 315. As t increases, the number of foxes decreases greatly to 100, and the number of rabbits decreases to 300 (the initial populations), and the cycle starts again.

(b)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. (a) At $t = 0$, there are about 600 rabbits and 160 foxes. At $t = t_1$, the number of rabbits reaches a minimum of about 80 and the number of foxes is also 80. At $t = t_2$, the number of foxes reaches a minimum of about 25 while the number of rabbits rebounds to 1000. At $t = t_3$, the number of foxes has increased to 40 and the rabbit population has reached a maximum of about 1750. The curve ends at $t = t_4$, where the number of foxes has increased to 65 and the number of rabbits has decreased to about 950.

(b)

7.

8.

9. (a) $\dfrac{dW}{dR} = \dfrac{-0.02W + 0.00002RW}{0.08R - 0.001RW} \quad \Leftrightarrow \quad (0.08 - 0.001W)R\,dW = (-0.02 + 0.00002R)W\,dR \quad \Leftrightarrow$

$\dfrac{0.08 - 0.001W}{W}\,dW = \dfrac{-0.02 + 0.00002R}{R}\,dR \quad \Leftrightarrow \quad \displaystyle\int\left(\frac{0.08}{W} - 0.001\right)dW = \int\left(-\frac{0.02}{R} + 0.00002\right)dR \quad \Leftrightarrow$

$0.08\,\ln|W| - 0.001W = -0.02\,\ln|R| + 0.00002R + K \quad \Leftrightarrow$

$0.08\,\ln W + 0.02\,\ln R = 0.001W + 0.00002R + K \quad \Leftrightarrow \quad \ln\left(W^{0.08}R^{0.02}\right) = 0.00002R + 0.001W + K \quad \Leftrightarrow$

$W^{0.08}R^{0.02} = e^{0.00002R + 0.001W + K} \quad \Leftrightarrow \quad R^{0.02}W^{0.08} = Ce^{0.00002R}e^{0.001W} \quad \Leftrightarrow \quad \dfrac{R^{0.02}W^{0.08}}{e^{0.00002R}e^{0.001W}} = C.$

In general, if $\dfrac{dy}{dx} = \dfrac{-ry + bxy}{kx - axy}$, then $C = \dfrac{x^r y^k}{e^{bx}e^{ay}}$.

(b)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. (a) A and L are constant $\Rightarrow$ $A' = 0$ and $L' = 0$ $\Rightarrow$ $\left.\begin{cases} 0 = 2A - 0.01AL \\ 0 = -0.5L + 0.0001AL \end{cases}\right\}$ $\Rightarrow$ $\begin{cases} 0 = A(2 - 0.01L) \\ 0 = L(-0.5 + 0.0001A) \end{cases}$

So either $A = L = 0$ or $L = \frac{2}{0.01} = 200$ and $A = \frac{0.5}{0.0001} = 5000$. The trivial solution $A = L = 0$ just says that if there aren't any aphids or ladybugs, then the populations will not change. The non-trivial solution, $L = 200$ and $A = 5000$, indicates the population sizes needed so that there are no changes in either the number of aphids or the number of ladybugs.

(b) $\dfrac{dL}{dA} = \dfrac{dL/dt}{dA/dt} = \dfrac{-0.5L + 0.0001AL}{2A - 0.01AL}$

(c) The solution curves (phase trajectories) are all closed curves that have the equilibrium point $(5000, 200)$ inside them.

(d)

At $P_0(1000, 200)$, $dA/dt = 0$ and $dL/dt = -80 < 0$, so the number of ladybugs is decreasing and hence, we are proceeding in a counterclockwise direction. At P_0, there aren't enough aphids to support the ladybug population, so the number of ladybugs decreases and the number of aphids begins to increase. The ladybug population reaches a minimum at $P_1(5000, 100)$ while the aphid population increases in a dramatic way, reaching its maximum at $P_2(14,250, 200)$.

Meanwhile, the ladybug population is increasing from P_1 to $P_3(5000, 355)$, and as we pass through P_2, the increasing number of ladybugs starts to deplete the aphid population. At P_3 the ladybugs reach a maximum population, and start to decrease due to the reduced aphid population. Both populations then decrease until P_0, where the cycle starts over again.

(e) Both graphs have the same period and the graph of L peaks about a quarter of a cycle after the graph of A.

11. (a) Letting $W = 0$ gives us $dR/dt = 0.08R(1 - 0.0002R)$. $dR/dt = 0$ $\Leftrightarrow$ $R = 0$ or 5000. Since $dR/dt > 0$ for $0 < R < 5000$, we would expect the rabbit population to *increase* to 5000 for these values of R. Since $dR/dt < 0$ for $R > 5000$, we would expect the rabbit population to *decrease* to 5000 for these values of R. Hence, in the absence of wolves, we would expect the rabbit population to stabilize at 5000.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) R and W are constant $\Rightarrow$ $R' = 0$ and $W' = 0$ $\Rightarrow$

$$\left.\begin{cases} 0 = 0.08R(1 - 0.0002R) - 0.001RW \\ 0 = -0.02W + 0.00002RW \end{cases}\right\} \Rightarrow \begin{cases} 0 = R[0.08(1 - 0.0002R) - 0.001W] \\ 0 = W(-0.02 + 0.00002R) \end{cases}$$

The second equation is true if $W = 0$ or $R = \frac{0.02}{0.00002} = 1000$. If $W = 0$ in the first equation, then either $R = 0$ or $R = \frac{1}{0.0002} = 5000$ [as in part (a)]. If $R = 1000$, then $0 = 1000[0.08(1 - 0.0002 \cdot 1000) - 0.001W] \Leftrightarrow$ $0 = 80(1 - 0.2) - W \Leftrightarrow W = 64$.

Case (i): $W = 0$, $R = 0$: both populations are zero

Case (ii): $W = 0$, $R = 5000$: see part (a)

Case (iii): $R = 1000$, $W = 64$: the predator/prey interaction balances and the populations are stable.

(c) The populations of wolves and rabbits fluctuate around 64 and 1000, respectively, and eventually stabilize at those values.

(d)

12. (a) If $L = 0$, $dA/dt = 2A(1 - 0.0001A)$, so $dA/dt = 0 \Leftrightarrow A = 0$ or $A = \frac{1}{0.0001} = 10{,}000$. Since $dA/dt > 0$ for $0 < A < 10{,}000$, we expect the aphid population to *increase* to 10,000 for these values of A. Since $dA/dt < 0$ for $A > 10{,}000$, we expect the aphid population to *decrease* to 10,000 for these values of A. Hence, in the absence of ladybugs we expect the aphid population to stabilize at 10,000.

(b) A and L are constant $\Rightarrow$ $A' = 0$ and $L' = 0$ $\Rightarrow$

$$\left.\begin{cases} 0 = 2A(1 - 0.0001A) - 0.01AL \\ 0 = -0.5L + 0.0001AL \end{cases}\right\} \Rightarrow \begin{cases} 0 = A[2(1 - 0.0001A) - 0.01L] \\ 0 = L(-0.5 + 0.0001A) \end{cases}$$

The second equation is true if $L = 0$ or $A = \frac{0.5}{0.0001} = 5000$. If $L = 0$ in the first equation, then either $A = 0$ or $A = \frac{1}{0.0001} = 10{,}000$. If $A = 5000$, then $0 = 5000[2(1 - 0.0001 \cdot 5000) - 0.01L] \Leftrightarrow$ $0 = 10{,}000(1 - 0.5) - 50L \Leftrightarrow 50L = 5000 \Leftrightarrow L = 100$.

The equilibrium solutions are: (i) $L = 0$, $A = 0$ (ii) $L = 0$, $A = 10{,}000$ (iii) $A = 5000$, $L = 100$

(c) $\dfrac{dL}{dA} = \dfrac{dL/dt}{dA/dt} = \dfrac{-0.5L + 0.0001AL}{2A(1 - 0.0001A) - 0.01AL}$

(d)

All of the phase trajectories spiral tightly around the equilibrium solution $(5000, 100)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(e)

At $t = 0$, the ladybug population decreases rapidly and the aphid population decreases slightly before beginning to increase. As the aphid population continues to increase, the ladybug population reaches a minimum at about $(5000, 75)$. The ladybug population starts to increase and quickly stabilizes at 100, while the aphid population stabilizes at 5000.

(f)

The graph of A peaks just after the graph of L has a minimum.

9 Review

TRUE-FALSE QUIZ

1. True. Since $y^4 \geq 0$, $y' = -1 - y^4 < 0$ and the solutions are decreasing functions.

2. True. $f(x) = y = \dfrac{\ln x}{x} \quad \Rightarrow \quad y' = \dfrac{1 - \ln x}{x^2}$.

LHS $= x^2 y' + xy = x^2 \cdot \dfrac{1 - \ln x}{x^2} + x \cdot \dfrac{\ln x}{x} = (1 - \ln x) + \ln x = 1 =$ RHS, so $y = \dfrac{\ln x}{x}$ is a solution of $x^2 y' + xy = 1$.

3. False. $y = 3e^{2x} - 1 \quad \Rightarrow \quad y' = 6e^{2x}$. LHS $= y' - 2y = 6e^{2x} - 2(3e^{2x} - 1) = 6e^{2x} - 6e^{2x} + 2 = 2 \neq 1$, so $y = 3e^{2x} - 1$ is not a solution to the initial-value problem.

4. False. $x + y$ cannot be written in the form $g(x)f(y)$.

5. True. $y' = 3y - 2x + 6xy - 1 = 6xy - 2x + 3y - 1 = 2x(3y - 1) + 1(3y - 1) = (2x + 1)(3y - 1)$, so y' can be written in the form $g(x)f(y)$, and hence, is separable.

6. True. $e^x y' = y \quad \Rightarrow \quad y' = e^{-x} y \quad \Rightarrow \quad y' + (-e^{-x})y = 0$, which is of the form $y' + P(x)\,y = Q(x)$, so the equation is linear.

7. False. $y' + xy = e^y$ cannot be put in the form $y' + P(x)\,y = Q(x)$, so it is not linear.

8. True. Substituting $x = 3$, $y = 1$, and $y' = 1$ into the differential equation $(2x - y)\,y' = x + 2y$ gives $(2 \cdot 3 - 1)(1) = 3 + 2(1) \quad \Rightarrow \quad 5 = 5$. Hence, the statement is correct.

9. True. By comparing $\dfrac{dy}{dt} = 2y\left(1 - \dfrac{y}{5}\right)$ with the logistic differential equation (9.4.4), we see that the carrying capacity is 5; that is, $\lim\limits_{t \to \infty} y = 5$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

EXERCISES

1. (a) $y' = y(y-2)(y-4)$

(b) $\lim_{t\to\infty} y(t)$ appears to be finite for $0 \le c \le 4$. In fact $\lim_{t\to\infty} y(t) = 4$ for $c = 4$, $\lim_{t\to\infty} y(t) = 2$ for $0 < c < 4$, and $\lim_{t\to\infty} y(t) = 0$ for $c = 0$. The equilibrium solutions are $y(t) = 0$, $y(t) = 2$, and $y(t) = 4$.

2. (a)

We sketch the direction field and four solution curves, as shown. Note that the slope $y' = x/y$ is not defined on the line $y = 0$.

(b) $y' = x/y \quad \Leftrightarrow \quad y\,dy = x\,dx \quad \Leftrightarrow \quad y^2 = x^2 + C$. For $C = 0$, this is the pair of lines $y = \pm x$. For $C \neq 0$, it is the hyperbola $x^2 - y^2 = -C$.

3. (a) $y' = x^2 - y^2$

We estimate that when $x = 0.3$, $y = 0.8$, so $y(0.3) \approx 0.8$.

(b) $h = 0.1$, $x_0 = 0$, $y_0 = 1$ and $F(x, y) = x^2 - y^2$. So $y_n = y_{n-1} + 0.1\left(x_{n-1}^2 - y_{n-1}^2\right)$. Thus,

$y_1 = 1 + 0.1\left(0^2 - 1^2\right) = 0.9$, $y_2 = 0.9 + 0.1\left(0.1^2 - 0.9^2\right) = 0.82$, $y_3 = 0.82 + 0.1\left(0.2^2 - 0.82^2\right) = 0.75676$.

This is close to our graphical estimate of $y(0.3) \approx 0.8$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) The centers of the horizontal line segments of the direction field are located on the lines $y = x$ and $y = -x$.

When a solution curve crosses one of these lines, it has a local maximum or minimum.

4. (a) $h = 0.2$, $x_0 = 0$, $y_0 = 1$ and $F(x, y) = 2xy^2$. We need y_2.

$y_1 = 1 + 0.2(2 \cdot 0 \cdot 1^2) = 1$, $y_2 = 1 + 0.2(2 \cdot 0.2 \cdot 1^2) = 1.08 \approx y(0.4)$.

(b) $h = 0.1$ now, so $y_1 = 1 + 0.1(2 \cdot 0 \cdot 1^2) = 1$, $y_2 = 1 + 0.1(2 \cdot 0.1 \cdot 1^2) = 1.02$,

$y_3 = 1.02 + 0.1(2 \cdot 0.2 \cdot 1.02^2) \approx 1.06162$, $y_4 = 1.06162 + 0.1(2 \cdot 0.3 \cdot 1.06162^2) \approx 1.1292 \approx y(0.4)$.

(c) The equation is separable, so we write $\dfrac{dy}{y^2} = 2x\,dx \;\Rightarrow\; \displaystyle\int \frac{dy}{y^2} = \int 2x\,dx \;\Leftrightarrow\; -\frac{1}{y} = x^2 + C$, but $y(0) = 1$, so

$C = -1$ and $y(x) = \dfrac{1}{1 - x^2} \;\Leftrightarrow\; y(0.4) = \dfrac{1}{1 - 0.16} \approx 1.1905$. From this we see that the approximation was greatly improved by increasing the number of steps, but the approximations were still far off.

5. $y' = xe^{-\sin x} - y\cos x \;\Rightarrow\; y' + (\cos x)\,y = xe^{-\sin x}$ $(\star)$. This is a linear equation and the integrating factor is $I(x) = e^{\int \cos x\,dx} = e^{\sin x}$. Multiplying $(\star)$ by $e^{\sin x}$ gives $e^{\sin x}\,y' + e^{\sin x}(\cos x)\,y = x \;\Rightarrow\; (e^{\sin x}\,y)' = x \;\Rightarrow$ $e^{\sin x}\,y = \frac{1}{2}x^2 + C \;\Rightarrow\; y = \left(\frac{1}{2}x^2 + C\right)e^{-\sin x}$.

6. $\dfrac{dx}{dt} = 1 - t + x - tx = 1(1 - t) + x(1 - t) = (1 + x)(1 - t) \;\Rightarrow\; \dfrac{dx}{1 + x} = (1 - t)\,dt \;\Rightarrow$

$\displaystyle\int \frac{dx}{1 + x} = \int (1 - t)\,dt \;\Rightarrow\; \ln|1 + x| = t - \tfrac{1}{2}t^2 + C \;\Rightarrow\; |1 + x| = e^{t - t^2/2 + C} \;\Rightarrow$

$1 + x = \pm e^{t - t^2/2} \cdot e^C \;\Rightarrow\; x = -1 + Ke^{t - t^2/2}$, where K is any nonzero constant.

7. $2ye^{y^2}y' = 2x + 3\sqrt{x} \;\Rightarrow\; 2ye^{y^2}\dfrac{dy}{dx} = 2x + 3\sqrt{x} \;\Rightarrow\; 2ye^{y^2}\,dy = \left(2x + 3\sqrt{x}\right)dx \;\Rightarrow$

$\int 2ye^{y^2}\,dy = \int \left(2x + 3\sqrt{x}\right)dx \;\Rightarrow\; e^{y^2} = x^2 + 2x^{3/2} + C \;\Rightarrow\; y^2 = \ln(x^2 + 2x^{3/2} + C) \;\Rightarrow$

$y = \pm\sqrt{\ln(x^2 + 2x^{3/2} + C)}$

8. $x^2y' - y = 2x^3e^{-1/x} \;\Rightarrow\; y' - \dfrac{1}{x^2}\,y = 2xe^{-1/x}$ $(\star)$. This is a linear equation and the integrating factor is

$I(x) = e^{\int(-1/x^2)\,dx} = e^{1/x}$. Multiplying $(\star)$ by $e^{1/x}$ gives $e^{1/x}\,y' - e^{1/x} \cdot \dfrac{1}{x^2}\,y = 2x \;\Rightarrow\; (e^{1/x}\,y)' = 2x \;\Rightarrow$

$e^{1/x}\,y = x^2 + C \;\Rightarrow\; y = e^{-1/x}(x^2 + C)$.

9. $\dfrac{dr}{dt} + 2tr = r \;\Rightarrow\; \dfrac{dr}{dt} = r - 2tr = r(1 - 2t) \;\Rightarrow\; \displaystyle\int \frac{dr}{r} = \int (1 - 2t)\,dt \;\Rightarrow\; \ln|r| = t - t^2 + C \;\Rightarrow$

$|r| = e^{t - t^2 + C} = ke^{t - t^2}$. Since $r(0) = 5$, $5 = ke^0 = k$. Thus, $r(t) = 5e^{t - t^2}$.

10. $(1 + \cos x)y' = (1 + e^{-y})\sin x \;\Rightarrow\; \dfrac{dy}{1 + e^{-y}} = \dfrac{\sin x\,dx}{1 + \cos x} \;\Rightarrow\; \displaystyle\int \frac{dy}{1 + 1/e^y} = \int \frac{\sin x\,dx}{1 + \cos x} \;\Rightarrow$

$\displaystyle\int \frac{e^y\,dy}{1 + e^y} = \int \frac{\sin x\,dx}{1 + \cos x} \;\Rightarrow\; \ln|1 + e^y| = -\ln|1 + \cos x| + C \;\Rightarrow\; \ln(1 + e^y) = -\ln(1 + \cos x) + C \;\Rightarrow$

$1 + e^y = e^{-\ln(1 + \cos x)} \cdot e^C \;\Rightarrow\; e^y = ke^{-\ln(1 + \cos x)} - 1 \;\Rightarrow\; y = \ln[ke^{-\ln(1 + \cos x)} - 1]$. Since $y(0) = 0$,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$0 = \ln[ke^{-\ln 2} - 1] \;\Rightarrow\; e^0 = k\left(\frac{1}{2}\right) - 1 \;\Rightarrow\; k = 4$. Thus, $y(x) = \ln[4e^{-\ln(1+\cos x)} - 1]$. An equivalent form is $y(x) = \ln \dfrac{3 - \cos x}{1 + \cos x}$.

11. $xy' - y = x \ln x \;\Rightarrow\; y' - \dfrac{1}{x}\, y = \ln x$. $I(x) = e^{\int(-1/x)\,dx} = e^{-\ln|x|} = \left(e^{\ln|x|}\right)^{-1} = |x|^{-1} = 1/x$ since the condition $y(1) = 2$ implies that we want a solution with $x > 0$. Multiplying the last differential equation by $I(x)$ gives

$\dfrac{1}{x}\, y' - \dfrac{1}{x^2}\, y = \dfrac{1}{x} \ln x \;\Rightarrow\; \left(\dfrac{1}{x}\, y\right)' = \dfrac{1}{x} \ln x \;\Rightarrow\; \dfrac{1}{x}\, y = \displaystyle\int \frac{\ln x}{x}\, dx \;\Rightarrow\; \dfrac{1}{x}\, y = \tfrac{1}{2}(\ln x)^2 + C \;\Rightarrow$

$y = \frac{1}{2}x(\ln x)^2 + Cx$. Now $y(1) = 2 \;\Rightarrow\; 2 = 0 + C \;\Rightarrow\; C = 2$, so $y = \frac{1}{2}x(\ln x)^2 + 2x$.

12. $y' = 3x^2 e^y \;\Rightarrow\; \dfrac{dy}{dx} = 3x^2 e^y \;\Rightarrow\; e^{-y}\, dy = 3x^2\, dx \;\Rightarrow$

$\int e^{-y}\, dy = \int 3x^2\, dx \;\Rightarrow\; -e^{-y} = x^3 + C$. Now $y(0) = 1 \;\Rightarrow$

$-e^{-1} = C$, so $-e^{-y} = x^3 - e^{-1} \;\Rightarrow\; e^{-y} = -x^3 + e^{-1} \;\Rightarrow$

$-y = \ln(-x^3 + e^{-1}) \;\Rightarrow\; y = -\ln(-x^3 + e^{-1})$. To find the domain, solve $-x^3 + e^{-1} > 0 \;\Rightarrow\; x^3 < e^{-1} \;\Rightarrow\; x < e^{-1/3}$, so the domain is $(-\infty, e^{-1/3})$ and $x = e^{-1/3}$ $[\approx 0.72]$ is a vertical asymptote.

13. $\dfrac{d}{dx}(y) = \dfrac{d}{dx}(ke^x) \;\Rightarrow\; y' = ke^x = y$, so the orthogonal trajectories must have $y' = -\dfrac{1}{y} \;\Rightarrow\; \dfrac{dy}{dx} = -\dfrac{1}{y} \;\Rightarrow$

$y\, dy = -dx \;\Rightarrow\; \int y\, dy = -\int dx \;\Rightarrow\; \frac{1}{2}y^2 = -x + C \;\Rightarrow\; x = C - \frac{1}{2}y^2$, which are parabolas with a horizontal axis.

14. $\dfrac{d}{dx}(y) = \dfrac{d}{dx}(e^{kx}) \;\Rightarrow\; y' = ke^{kx} = ky = \dfrac{\ln y}{x} \cdot y$, so the orthogonal trajectories must have $y' = -\dfrac{x}{y \ln y} \;\Rightarrow$

$\dfrac{dy}{dx} = -\dfrac{x}{y \ln y} \;\Rightarrow\; y \ln y\, dy = -x\, dx \;\Rightarrow\; \int y \ln y\, dy = -\int x\, dx \;\Rightarrow\; \frac{1}{2}y^2 \ln y - \frac{1}{4}y^2$ [parts with $u = \ln y$, $dv = y\, dy$] $= -\frac{1}{2}x^2 + C_1 \;\Rightarrow\; 2y^2 \ln y - y^2 = C - 2x^2$.

15. (a) Using (4) and (7) in Section 9.4, we see that for $\dfrac{dP}{dt} = 0.1P\left(1 - \dfrac{P}{2000}\right)$ with $P(0) = 100$, we have $k = 0.1$, $M = 2000$, $P_0 = 100$, and $A = \dfrac{2000 - 100}{100} = 19$. Thus, the solution of the initial-value problem is

$P(t) = \dfrac{2000}{1 + 19e^{-0.1t}}$ and $P(20) = \dfrac{2000}{1 + 19e^{-2}} \approx 560$.

(b) $P = 1200 \;\Leftrightarrow\; 1200 = \dfrac{2000}{1 + 19e^{-0.1t}} \;\Leftrightarrow\; 1 + 19e^{-0.1t} = \dfrac{2000}{1200} \;\Leftrightarrow\; 19e^{-0.1t} = \dfrac{5}{3} - 1 \;\Leftrightarrow$

$e^{-0.1t} = \left(\frac{2}{3}\right)/19 \;\Leftrightarrow\; -0.1t = \ln \frac{2}{57} \;\Leftrightarrow\; t = -10 \ln \frac{2}{57} \approx 33.5$.

16. (a) Let $t = 0$ correspond to the year 2000. An exponential model is $P(t) = ae^{kt}$. $P(0) = 6.08$, so $P(t) = 6.08e^{kt}$.

$P(15) = 6.08e^{15k}$ and $P(15) = 7.35$, so $6.08e^{15k} = 7.35 \;\Rightarrow\; e^{15k} = \dfrac{7.35}{6.08} \;\Rightarrow\; 15k = \ln \dfrac{735}{608} \;\Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$k = \dfrac{1}{15}\ln\dfrac{735}{608} \approx 0.0126$. Thus, $P(t) = 6.08e^{kt}$ and $P(30) = 6.08e^{30k} \approx 8.89$. Our model predicts that the world population in the year 2030 will be 8.89 billion.

(b) $P(t) = 10 \;\Rightarrow\; 6.08e^{kt} = 10 \;\Rightarrow\; e^{kt} = \dfrac{10}{6.08} \;\Rightarrow\; kt = \ln\dfrac{10}{6.08} \;\Rightarrow\; t = 15\,\dfrac{\ln(10/6.08)}{\ln(735/608)} \approx 39.35$ years. Our exponential model predicts that the world population will exceed 10 billion in about 39.35 years; that is, in the year 2039.

(c) $A = \dfrac{M - P_0}{P_0} = \dfrac{20 - 6.08}{6.08} = \dfrac{87}{38}$, so $P(t) = \dfrac{M}{1 + Ae^{-kt}} = \dfrac{20}{1 + \frac{87}{38}e^{-kt}}$. $P(15) = 7.35 \;\Rightarrow$

$\dfrac{20}{1 + \frac{87}{38}e^{-15k}} = 7.35 \;\Rightarrow\; \dfrac{20}{7.35} = 1 + \dfrac{87}{38}e^{-15k} \;\Rightarrow\; \dfrac{253}{147} = \dfrac{87}{38}e^{-15k} \;\Rightarrow\; \dfrac{9614}{12{,}789} = e^{-15k} \;\Rightarrow$

$k = -\dfrac{1}{15}\ln\dfrac{9614}{12{,}789} \approx 0.01902$. Thus, $P(30) = \dfrac{20}{1 + \frac{87}{38}e^{-30k}} \approx 8.72$ billion, which is less than our prediction of 8.89 billion from the exponential model in part (a).

(d) $P(t) = 10 \;\Rightarrow\; \dfrac{20}{1 + \frac{87}{38}e^{-kt}} = 10 \;\Rightarrow\; \dfrac{20}{10} = 1 + \dfrac{87}{38}e^{-kt} \;\Rightarrow\; 1 = \dfrac{87}{38}e^{-kt} \;\Rightarrow\; \dfrac{38}{87} = e^{-kt} \;\Rightarrow$

$\ln\dfrac{38}{87} = -kt \;\Rightarrow\; t = 15\,\dfrac{\ln(38/87)}{\ln(9614/12{,}789)} \approx 43.54$ years; that is, in the year 2043, which is four years later than our prediction of 2039 from the exponential model in part (b).

17. (a) $\dfrac{dL}{dt} \propto L_\infty - L \;\Rightarrow\; \dfrac{dL}{dt} = k(L_\infty - L) \;\Rightarrow\; \displaystyle\int \frac{dL}{L_\infty - L} = \int k\,dt \;\Rightarrow\; -\ln|L_\infty - L| = kt + C \;\Rightarrow$

$\ln|L_\infty - L| = -kt - C \;\Rightarrow\; |L_\infty - L| = e^{-kt-C} \;\Rightarrow\; L_\infty - L = Ae^{-kt} \;\Rightarrow\; L = L_\infty - Ae^{-kt}$.

At $t = 0$, $L = L(0) = L_\infty - A \;\Rightarrow\; A = L_\infty - L(0) \;\Rightarrow\; L(t) = L_\infty - [L_\infty - L(0)]e^{-kt}$.

(b) $L_\infty = 53$ cm, $L(0) = 10$ cm, and $k = 0.2 \;\Rightarrow\; L(t) = 53 - (53 - 10)e^{-0.2t} = 53 - 43e^{-0.2t}$.

18. Denote the amount of salt in the tank (in kg) by y. $y(0) = 0$ since initially there is only water in the tank.

The rate at which y increases is equal to the rate at which salt flows into the tank minus the rate at which it flows out.

That rate is $\dfrac{dy}{dt} = 0.1\dfrac{\text{kg}}{\text{L}} \times 10\dfrac{\text{L}}{\text{min}} - \dfrac{y}{100}\dfrac{\text{kg}}{\text{L}} \times 10\dfrac{\text{L}}{\text{min}} = 1 - \dfrac{y}{10}\dfrac{\text{kg}}{\text{min}} \;\Rightarrow\; \displaystyle\int \frac{dy}{10 - y} = \int \frac{1}{10}\,dt \;\Rightarrow$

$-\ln|10 - y| = \frac{1}{10}t + C \;\Rightarrow\; 10 - y = Ae^{-t/10}$. $y(0) = 0 \;\Rightarrow\; 10 = A \;\Rightarrow\; y = 10(1 - e^{-t/10})$.

At $t = 6$ minutes, $y = 10(1 - e^{-6/10}) \approx 4.512$ kg.

19. Let P represent the population and I the number of infected people. The rate of spread dI/dt is jointly proportional to I and to $P - I$, so for some constant k, $\dfrac{dI}{dt} = kI(P - I) \;\Rightarrow\; I(t) = \dfrac{I_0 P}{I_0 + (P - I_0)e^{-kPt}}$ [from the discussion of logistic growth in Section 9.4].

Now, measuring t in days, we substitute $t = 7$, $P = 5000$, $I_0 = 160$ and $I(7) = 1200$ to find k:

$1200 = \dfrac{160 \cdot 5000}{160 + (5000 - 160)e^{-5000 \cdot 7 \cdot k}} \;\Leftrightarrow\; 3 = \dfrac{2000}{160 + 4840e^{-35{,}000k}} \;\Leftrightarrow\; 480 + 14{,}520e^{-35{,}000k} = 2000 \;\Leftrightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$e^{-35{,}000k} = \dfrac{2000-480}{14{,}520} \quad \Leftrightarrow \quad -35{,}000k = \ln\dfrac{38}{363} \quad \Leftrightarrow \quad k = \dfrac{-1}{35{,}000}\ln\dfrac{38}{363} \approx 0.00006448$. Next, let

$I = 5000 \times 80\% = 4000$, and solve for t: $4000 = \dfrac{160 \cdot 5000}{160 + (5000-160)e^{-k\cdot 5000\cdot t}} \quad \Leftrightarrow \quad 1 = \dfrac{200}{160 + 4840e^{-5000kt}} \quad \Leftrightarrow$

$160 + 4840e^{-5000kt} = 200 \quad \Leftrightarrow \quad e^{-5000kt} = \dfrac{200-160}{4840} \quad \Leftrightarrow \quad -5000kt = \ln\dfrac{1}{121} \quad \Leftrightarrow$

$t = \dfrac{-1}{5000k}\ln\dfrac{1}{121} = \dfrac{1}{\frac{1}{7}\ln\frac{38}{363}} \cdot \ln\dfrac{1}{121} = 7 \cdot \dfrac{\ln 121}{\ln\frac{363}{38}} \approx 14.875$. So it takes about 15 days for 80% of the population to be infected.

20. $\dfrac{1}{R}\dfrac{dR}{dt} = \dfrac{k}{S}\dfrac{dS}{dt} \quad \Rightarrow \quad \dfrac{d}{dt}(\ln R) = \dfrac{d}{dt}(k\ln S) \quad \Rightarrow \quad \ln R = k\ln S + C \quad \Rightarrow$

$R = e^{k\ln S + C} = e^C\left(e^{\ln S}\right)^k \quad \Rightarrow \quad R = AS^k$, where $A = e^C$ is a positive constant.

21. $\dfrac{dh}{dt} = -\dfrac{R}{V}\left(\dfrac{h}{k+h}\right) \quad \Rightarrow \quad \displaystyle\int \frac{k+h}{h}\,dh = \int\left(-\frac{R}{V}\right)dt \quad \Rightarrow \quad \int\left(1+\frac{k}{h}\right)dh = -\frac{R}{V}\int 1\,dt \quad \Rightarrow$

$h + k\ln h = -\dfrac{R}{V}\,t + C$. This equation gives a relationship between h and t, but it is not possible to isolate h and express it in terms of t.

22. $dx/dt = 0.4x - 0.002xy$, $dy/dt = -0.2y + 0.000008xy$

(a) The xy terms represent encounters between the birds and the insects. Since the y-population increases from these terms and the x-population decreases, we expect y to represent the birds and x the insects.

(b) x and y are constant $\Rightarrow$ $x' = 0$ and $y' = 0$ $\Rightarrow$

$$\left\{\begin{aligned} 0 &= 0.4x - 0.002xy \\ 0 &= -0.2y + 0.000008xy \end{aligned}\right\} \quad \Rightarrow \quad \left\{\begin{aligned} 0 &= 0.4x(1-0.005y) \\ 0 &= -0.2y(1-0.00004x) \end{aligned}\right. \quad \Rightarrow \quad y = 0 \text{ and } x = 0 \text{ (zero populations)}$$

or $y = \frac{1}{0.005} = 200$ and $x = \frac{1}{0.00004} = 25{,}000$. The non-trivial solution represents the population sizes needed so that there are no changes in either the number of birds or the number of insects.

(c) $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{-0.2y + 0.000008xy}{0.4x - 0.002xy}$

(d)

At $(x, y) = (40{,}000, 100)$, $dx/dt = 8000 > 0$, so as t increases we are proceeding in a counterclockwise direction. The populations increase to approximately $(59{,}646, 200)$, at which point the insect population starts to decrease. The birds attain a maximum population of about 380 when the insect population is 25,000. The populations decrease to about $(7370, 200)$, at which point the insect population starts to increase. The birds attain a minimum population of about 88 when the insect population is 25,000, and then the cycle repeats.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(e)

Both graphs have the same period and the bird population peaks about a quarter-cycle after the insect population.

23. (a) $dx/dt = 0.4x(1 - 0.000005x) - 0.002xy$, $dy/dt = -0.2y + 0.000008xy$. If $y = 0$, then $dx/dt = 0.4x(1 - 0.000005x)$, so $dx/dt = 0 \quad \Leftrightarrow \quad x = 0$ or $x = 200{,}000$, which shows that the insect population increases logistically with a carrying capacity of 200,000. Since $dx/dt > 0$ for $0 < x < 200{,}000$ and $dx/dt < 0$ for $x > 200{,}000$, we expect the insect population to stabilize at 200,000.

(b) x and y are constant $\quad \Rightarrow \quad x' = 0$ and $y' = 0 \quad \Rightarrow$

$$\left.\begin{cases} 0 = 0.4x(1 - 0.000005x) - 0.002xy \\ 0 = -0.2y + 0.000008xy \end{cases}\right\} \quad \Rightarrow \quad \begin{cases} 0 = 0.4x[(1 - 0.000005x) - 0.005y] \\ 0 = y(-0.2 + 0.000008x) \end{cases}$$

The second equation is true if $y = 0$ or $x = \frac{0.2}{0.000008} = 25{,}000$. If $y = 0$ in the first equation, then either $x = 0$ or $x = \frac{1}{0.000005} = 200{,}000$. If $x = 25{,}000$, then $0 = 0.4(25{,}000)[(1 - 0.000005 \cdot 25{,}000) - 0.005y] \quad \Rightarrow$ $0 = 10{,}000[(1 - 0.125) - 0.005y] \quad \Rightarrow \quad 0 = 8750 - 50y \quad \Rightarrow \quad y = 175$.

Case (i): $y = 0$, $x = 0$: Zero populations

Case (ii): $y = 0$, $x = 200{,}000$: In the absence of birds, the insect population is always 200,000.

Case (iii): $x = 25{,}000$, $y = 175$: The predator/prey interaction balances and the populations are stable.

(c) The populations of the birds and insects fluctuate around 175 and 25,000, respectively, and eventually stabilize at those values.

(d)

24. First note that, in this question, "weighs" is used in the informal sense, so what we really require is Brett's mass m in kg as a function of t. Brett's net intake of calories per day at time t (measured in days) is $c(t) = 2200 - 1200 - 15m(t) = 1000 - 15m(t)$, where $m(t)$ is his mass at time t. We are given that $m(0) = 85$ kg and $\dfrac{dm}{dt} = \dfrac{c(t)}{10{,}000}$, so $\dfrac{dm}{dt} = \dfrac{1000 - 15m}{10{,}000} = \dfrac{200 - 3m}{2000}$ with $m(0) = 85$. From $\displaystyle\int \frac{dm}{200 - 3m} = \int \frac{dt}{2000}$, we get $-\frac{1}{3}\ln|200 - 3m| = \frac{1}{2000}t + C$. Since $m(0) = 85$, $C = -\frac{1}{3}\ln 55$. Now, $-\frac{1}{3}\ln|200 - 3m| = \frac{1}{2000}t - \frac{1}{3}\ln 55 \quad \Rightarrow$ $\ln|200 - 3m| = -\frac{3}{2000}t + \ln 55 \quad \Rightarrow \quad |200 - 3m| = e^{-3t/2000 + \ln 55} = 55e^{-3t/2000}$. The quantity $200 - 3m$ is continuous, initially negative, and the right-hand side is never zero. Thus, $200 - 3m$ is negative for all t, so $200 - 3m = -55e^{-3t/2000} \quad \Rightarrow \quad m(t) = \frac{200}{3} + \frac{55}{3}e^{-3t/2000}$ kg. As $t \to \infty$, $m(t) \to \frac{200}{3} \approx 66.7$ kg. Thus, Brett's mass gradually settles down to 66.7 kg.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1. We use the Fundamental Theorem of Calculus to differentiate the given equation:

$[f(x)]^2 = 100 + \int_0^x \left\{[f(t)]^2 + [f'(t)]^2\right\} dt \quad \Rightarrow \quad 2f(x)f'(x) = [f(x)]^2 + [f'(x)]^2 \quad \Rightarrow$

$[f(x)]^2 + [f'(x)]^2 - 2f(x)f'(x) = 0 \quad \Rightarrow \quad [f(x) - f'(x)]^2 = 0 \quad \Leftrightarrow \quad f(x) = f'(x)$. We can solve this as a separable equation, or else use Theorem 9.4.2 with $k = 1$, which says that the solutions are $f(x) = Ce^x$. Now $[f(0)]^2 = 100$, so $f(0) = C = \pm 10$, and hence $f(x) = \pm 10e^x$ are the only functions satisfying the given equation.

2. $(fg)' = f'g'$, where $f(x) = e^{x^2} \quad \Rightarrow \quad \left(e^{x^2} g\right)' = 2xe^{x^2} g'$. Since the student's mistake did not affect the answer,

$\left(e^{x^2} g\right)' = e^{x^2} g' + 2xe^{x^2} g = 2xe^{x^2} g'$. So $(2x-1)g' = 2xg$, or $\dfrac{g'}{g} = \dfrac{2x}{2x-1} = 1 + \dfrac{1}{2x-1} \quad \Rightarrow$

$\ln|g(x)| = x + \frac{1}{2}\ln(2x-1) + C \quad \Rightarrow \quad g(x) = Ae^x\sqrt{2x-1}$.

3. $f'(x) = \lim\limits_{h\to 0} \dfrac{f(x+h) - f(x)}{h} = \lim\limits_{h\to 0} \dfrac{f(x)\,[f(h) - 1]}{h}$ [since $f(x+h) = f(x)f(h)$]

$$= f(x) \lim_{h\to 0} \frac{f(h) - 1}{h} = f(x) \lim_{h\to 0} \frac{f(h) - f(0)}{h - 0} = f(x)f'(0) = f(x) \cdot 1 = f(x)$$

Therefore, $f'(x) = f(x)$ for all x and from Theorem 9.4.2 we get $f(x) = Ae^x$.

Now $f(0) = 1 \quad \Rightarrow \quad A = 1 \quad \Rightarrow \quad f(x) = e^x$.

4. $\left(\int f(x)\,dx\right)\left(\int \dfrac{1}{f(x)}\,dx\right) = -1 \quad \Rightarrow \quad \int \dfrac{1}{f(x)}\,dx = \dfrac{-1}{\int f(x)\,dx} \quad \Rightarrow$

$\dfrac{1}{f(x)} = \dfrac{f(x)}{\left[\int f(x)\,dx\right]^2}$ [after differentiating] $\quad \Rightarrow \quad \int f(x)\,dx = \pm f(x)$ [after taking square roots] $\quad \Rightarrow$

$f(x) = \pm f'(x)$ [after differentiating again] $\quad \Rightarrow \quad y = Ae^x$ or $y = Ae^{-x}$ by Theorem 9.4.2. Therefore, $f(x) = Ae^x$ or $f(x) = Ae^{-x}$, for all nonzero constants A, are the functions satisfying the original equation.

5. "The area under the graph of f from 0 to x is proportional to the $(n+1)$st power of $f(x)$" translates to

$\int_0^x f(t)\,dt = k[f(x)]^{n+1}$ for some constant k. By FTC1, $\dfrac{d}{dx}\displaystyle\int_0^x f(t)\,dt = \dfrac{d}{dx}\left\{k[f(x)]^{n+1}\right\} \quad \Rightarrow$

$f(x) = k(n+1)[f(x)]^n f'(x) \quad \Rightarrow \quad 1 = k(n+1)[f(x)]^{n-1} f'(x) \quad \Rightarrow \quad 1 = k(n+1)y^{n-1}\dfrac{dy}{dx} \quad \Rightarrow$

$k(n+1)y^{n-1}\,dy = dx \quad \Rightarrow \quad \int k(n+1)y^{n-1}\,dy = \int dx \quad \Rightarrow \quad k(n+1)\dfrac{1}{n}y^n = x + C$.

Now $f(0) = 0 \quad \Rightarrow \quad 0 = 0 + C \quad \Rightarrow \quad C = 0$ and then $f(1) = 1 \quad \Rightarrow \quad k(n+1)\dfrac{1}{n} = 1 \quad \Rightarrow \quad k = \dfrac{n}{n+1}$,

so $y^n = x$ and $y = f(x) = x^{1/n}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. Let $y = f(x)$ be a curve that passes through the point $(c, 1)$ and whose subtangents all have length c. The tangent line at $x = a$ has equation $y - f(a) = f'(a)(x - a)$. Assuming $f(a) \neq 0$ and $f'(a) \neq 0$, it has x-intercept $a - \dfrac{f(a)}{f'(a)}$ [let $y = 0$ and solve for x]. Thus, the length of the subtangent is c, so $\left|a - \left(a - \dfrac{f(a)}{f'(a)}\right)\right| = \left|\dfrac{f(a)}{f'(a)}\right| = c \;\Rightarrow\; \dfrac{f'(a)}{f(a)} = \pm\dfrac{1}{c}$.

Now $\dfrac{f'(x)}{f(x)} = \pm\dfrac{1}{c} \;\Rightarrow\; f'(x) = \pm\dfrac{1}{c}\,f(x) \;\Rightarrow\; \dfrac{dy}{dx} = \pm\dfrac{1}{c}\,y \;\Rightarrow\; \dfrac{dy}{y} = \pm\dfrac{1}{c}\,dx \;\Rightarrow\; \displaystyle\int \frac{1}{y}\,dy = \pm\frac{1}{c}\int dx \;\Rightarrow$ $\ln|y| = \pm\dfrac{1}{c}\,x + K$. Since $f(c) = 1$, $\ln 1 = \pm 1 + K \;\Rightarrow\; K = \mp 1$. Thus, $y = e^{\pm x/c \mp 1}$, or $y = e^{\pm(x/c - 1)}$. One curve is an increasing exponential (as shown in the figure) and the other curve is its reflection about the line $x = c$.

7. Let $y(t)$ denote the temperature of the peach pie t minutes after 5:00 PM and R the temperature of the room. Newton's Law of Cooling gives us $dy/dt = k(y - R)$. Solving for y we get $\dfrac{dy}{y - R} = k\,dt \;\Rightarrow\; \ln|y - R| = kt + C \;\Rightarrow$ $|y - R| = e^{kt + C} \;\Rightarrow\; y - R = \pm e^{kt} \cdot e^{C} \;\Rightarrow\; y = Me^{kt} + R$, where M is a nonzero constant. We are given temperatures at three times.

$$\begin{aligned} y(0) &= 100 \quad\Rightarrow\quad 100 = M + R \quad\Rightarrow\quad R = 100 - M \\ y(10) &= 80 \quad\Rightarrow\quad 80 = Me^{10k} + R \qquad\qquad (\mathbf{1}) \\ y(20) &= 65 \quad\Rightarrow\quad 65 = Me^{20k} + R \qquad\qquad (\mathbf{2}) \end{aligned}$$

Substituting $100 - M$ for R in **(1)** and **(2)** gives us

$$-20 = Me^{10k} - M \;\; \textbf{(3)} \quad \text{and} \quad -35 = Me^{20k} - M \;\; \textbf{(4)}$$

Dividing **(3)** by **(4)** gives us $\dfrac{-20}{-35} = \dfrac{M\left(e^{10k} - 1\right)}{M\left(e^{20k} - 1\right)} \;\Rightarrow\; \dfrac{4}{7} = \dfrac{e^{10k} - 1}{e^{20k} - 1} \;\Rightarrow\; 4e^{20k} - 4 = 7e^{10k} - 7 \;\Rightarrow$ $4e^{20k} - 7e^{10k} + 3 = 0$. This is a quadratic equation in e^{10k}. $\left(4e^{10k} - 3\right)\left(e^{10k} - 1\right) = 0 \;\Rightarrow\; e^{10k} = \frac{3}{4}$ or $1 \;\Rightarrow$ $10k = \ln\frac{3}{4}$ or $\ln 1 \;\Rightarrow\; k = \frac{1}{10}\ln\frac{3}{4}$ since k is a nonzero constant of proportionality. Substituting $\frac{3}{4}$ for e^{10k} in **(3)** gives us $-20 = M \cdot \frac{3}{4} - M \;\Rightarrow\; -20 = -\frac{1}{4}M \;\Rightarrow\; M = 80$. Now $R = 100 - M$ so $R = 20°\text{C}$.

8. Let b be the number of hours before noon that it began to snow, t the time measured in hours after noon, and $x = x(t) =$ distance traveled by the plow at time t. Then $dx/dt =$ speed of plow. Since the snow falls steadily, the height at time t is $h(t) = k(t + b)$, where k is a constant. We are given that the rate of removal is constant, say R (in m^3/h). If the width of the path is w, then $R =$ height $\times$ width $\times$ speed $= h(t) \times w \times \dfrac{dx}{dt} = k(t + b)w\,\dfrac{dx}{dt}$. Thus, $\dfrac{dx}{dt} = \dfrac{C}{t + b}$, where $C = \dfrac{R}{kw}$ is a constant. This is a separable equation. $\displaystyle\int dx = C\int \frac{dt}{t + b} \;\Rightarrow\; x(t) = C\ln(t + b) + K$.

Put $t = 0$: $0 = C\ln b + K \;\Rightarrow\; K = -C\ln b$, so $x(t) = C\ln(t + b) - C\ln b = C\ln(1 + t/b)$.

Put $t = 1$: $6000 = C\ln(1 + 1/b)$ $\quad[x = 6 \text{ km}]$.

Put $t = 2$: $9000 = C\ln(1 + 2/b)$ $\quad[x = (6 + 3) \text{ km}]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Solve for b: $\dfrac{\ln(1+1/b)}{6000} = \dfrac{\ln(1+2/b)}{9000} \;\Rightarrow\; 3\ln\left(1+\dfrac{1}{b}\right) = 2\ln\left(1+\dfrac{2}{b}\right) \;\Rightarrow\; \left(1+\dfrac{1}{b}\right)^3 = \left(1+\dfrac{2}{b}\right)^2 \;\Rightarrow$

$1+\dfrac{3}{b}+\dfrac{3}{b^2}+\dfrac{1}{b^3} = 1+\dfrac{4}{b}+\dfrac{4}{b^2} \;\Rightarrow\; \dfrac{1}{b}+\dfrac{1}{b^2}-\dfrac{1}{b^3} = 0 \;\Rightarrow\; b^2+b-1=0 \;\Rightarrow\; b = \frac{-1\pm\sqrt{5}}{2}$.

But $b>0$, so $b = \frac{-1+\sqrt{5}}{2} \approx 0.618$ h ≈ 37 min. The snow began to fall $\frac{\sqrt{5}-1}{2}$ hours before noon; that is, at about 11:23 AM.

9. (a) While running from $(L,0)$ to (x,y), the dog travels a distance

$s = \int_x^L \sqrt{1+(dy/dx)^2}\,dx = -\int_L^x \sqrt{1+(dy/dx)^2}\,dx$, so

$\dfrac{ds}{dx} = -\sqrt{1+(dy/dx)^2}$. The dog and rabbit run at the same speed, so the rabbit's position when the dog has traveled a distance s is $(0,s)$. Since the dog runs straight for the rabbit, $\dfrac{dy}{dx} = \dfrac{s-y}{0-x}$ (see the figure).

Thus, $s = y - x\dfrac{dy}{dx} \;\Rightarrow\; \dfrac{ds}{dx} = \dfrac{dy}{dx} - \left(x\dfrac{d^2y}{dx^2} + 1\dfrac{dy}{dx}\right) = -x\dfrac{d^2y}{dx^2}$. Equating the two expressions for $\dfrac{ds}{dx}$ gives us $x\dfrac{d^2y}{dx^2} = \sqrt{1+\left(\dfrac{dy}{dx}\right)^2}$, as claimed.

(b) Letting $z = \dfrac{dy}{dx}$, we obtain the differential equation $x\dfrac{dz}{dx} = \sqrt{1+z^2}$, or $\dfrac{dz}{\sqrt{1+z^2}} = \dfrac{dx}{x}$. Integrating:

$\ln x = \displaystyle\int \frac{dz}{\sqrt{1+z^2}} \overset{25}{=} \ln\left(z+\sqrt{1+z^2}\right) + C$. When $x=L$, $z = dy/dx = 0$, so $\ln L = \ln 1 + C$. Therefore,

$C = \ln L$, so $\ln x = \ln\left(\sqrt{1+z^2}+z\right) + \ln L = \ln\left[L\left(\sqrt{1+z^2}+z\right)\right] \;\Rightarrow\; x = L\left(\sqrt{1+z^2}+z\right) \;\Rightarrow$

$\sqrt{1+z^2} = \dfrac{x}{L} - z \;\Rightarrow\; 1+z^2 = \left(\dfrac{x}{L}\right)^2 - \dfrac{2xz}{L} + z^2 \;\Rightarrow\; \left(\dfrac{x}{L}\right)^2 - 2z\left(\dfrac{x}{L}\right) - 1 = 0 \;\Rightarrow$

$z = \dfrac{(x/L)^2-1}{2(x/L)} = \dfrac{x^2-L^2}{2Lx} = \dfrac{x}{2L} - \dfrac{L}{2}\dfrac{1}{x}$ [for $x>0$]. Since $z = \dfrac{dy}{dx}$, $y = \dfrac{x^2}{4L} - \dfrac{L}{2}\ln x + C_1$.

Since $y=0$ when $x=L$, $0 = \dfrac{L}{4} - \dfrac{L}{2}\ln L + C_1 \;\Rightarrow\; C_1 = \dfrac{L}{2}\ln L - \dfrac{L}{4}$. Thus,

$y = \dfrac{x^2}{4L} - \dfrac{L}{2}\ln x + \dfrac{L}{2}\ln L - \dfrac{L}{4} = \dfrac{x^2-L^2}{4L} - \dfrac{L}{2}\ln\left(\dfrac{x}{L}\right)$.

(c) As $x \to 0^+$, $y \to \infty$, so the dog never catches the rabbit.

10. (a) If the dog runs twice as fast as the rabbit, then the rabbit's position when the dog has traveled a distance s is $(0, s/2)$.

Since the dog runs straight toward the rabbit, the tangent line to the dog's path has slope $\dfrac{dy}{dx} = \dfrac{s/2-y}{0-x}$.

Thus, $s = 2y - 2x\dfrac{dy}{dx} \;\Rightarrow\; \dfrac{ds}{dx} = 2\dfrac{dy}{dx} - \left(2x\dfrac{d^2y}{dx^2} + 2\dfrac{dy}{dx}\right) = -2x\dfrac{d^2y}{dx^2}$.

From Problem 9(a), $\dfrac{ds}{dx} = -\sqrt{1+\left(\dfrac{dy}{dx}\right)^2}$, so $2x\dfrac{d^2y}{dx^2} = \sqrt{1+\left(\dfrac{dy}{dx}\right)^2}$.

Letting $z = \dfrac{dy}{dx}$, we obtain the differential equation $2x\dfrac{dz}{dx} = \sqrt{1+z^2}$, or $\dfrac{2\,dz}{\sqrt{1+z^2}} = \dfrac{dx}{x}$. Integrating, we get

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\ln x = \int \frac{2\,dz}{\sqrt{1+z^2}} = 2\ln\left(\sqrt{1+z^2}+z\right) + C$. [See Problem 9(b).] When $x = L$, $z = dy/dx = 0$, so $\ln L = 2\ln 1 + C = C$. Thus,

$$\ln x = 2\ln\left(\sqrt{1+z^2}+z\right) + \ln L = \ln\left(L\left(\sqrt{1+z^2}+z\right)^2\right) \quad\Rightarrow\quad x = L\left(\sqrt{1+z^2}+z\right)^2 \quad\Rightarrow$$

$$\sqrt{1+z^2} = \sqrt{\frac{x}{L}} - z \quad\Rightarrow\quad 1+z^2 = \frac{x}{L} - 2\sqrt{\frac{x}{L}}\,z + z^2 \quad\Rightarrow\quad 2\sqrt{\frac{x}{L}}\,z = \frac{x}{L} - 1 \quad\Rightarrow$$

$$\frac{dy}{dx} = z = \frac{1}{2}\sqrt{\frac{x}{L}} - \frac{1}{2\sqrt{x/L}} = \frac{1}{2\sqrt{L}}\,x^{1/2} - \frac{\sqrt{L}}{2}\,x^{-1/2} \quad\Rightarrow\quad y = \frac{1}{3\sqrt{L}}\,x^{3/2} - \sqrt{L}\,x^{1/2} + C_1.$$

When $x = L$, $y = 0$, so $0 = \frac{1}{3\sqrt{L}}\,L^{3/2} - \sqrt{L}\,L^{1/2} + C_1 = \frac{L}{3} - L + C_1 = C_1 - \frac{2}{3}L$. Therefore, $C_1 = \frac{2}{3}L$ and

$y = \frac{x^{3/2}}{3\sqrt{L}} - \sqrt{L}\,x^{1/2} + \frac{2}{3}L$. As $x \to 0$, $y \to \frac{2}{3}L$, so the dog catches the rabbit when the rabbit is at $\left(0, \frac{2}{3}L\right)$.

(At that point, the dog has traveled a distance of $\frac{4}{3}L$, twice as far as the rabbit has run.)

(b) As in the solutions to part (a) and Problem 9, we get $z = \frac{dy}{dx} = \frac{x^2}{2L^2} - \frac{L^2}{2x^2}$ and hence $y = \frac{x^3}{6L^2} + \frac{L^2}{2x} - \frac{2}{3}L$.

We want to minimize the distance D from the dog at (x, y) to the rabbit at $(0, 2s)$. Now $s = \frac{1}{2}y - \frac{1}{2}x\,\frac{dy}{dx} \quad\Rightarrow$

$2s = y - xz \quad\Rightarrow\quad y - 2s = xz = x\left(\frac{x^2}{2L^2} - \frac{L^2}{2x^2}\right) = \frac{x^3}{2L^2} - \frac{L^2}{2x}$, so

$$D = \sqrt{(x-0)^2 + (y-2s)^2} = \sqrt{x^2 + \left(\frac{x^3}{2L^2} - \frac{L^2}{2x}\right)^2} = \sqrt{\frac{x^6}{4L^4} + \frac{x^2}{2} + \frac{L^4}{4x^2}} = \sqrt{\left(\frac{x^3}{2L^2} + \frac{L^2}{2x}\right)^2}$$
$$= \frac{x^3}{2L^2} + \frac{L^2}{2x}$$

$$D' = 0 \quad\Leftrightarrow\quad \frac{3x^2}{2L^2} - \frac{L^2}{2x^2} = 0 \quad\Leftrightarrow\quad \frac{3x^2}{2L^2} = \frac{L^2}{2x^2} \quad\Leftrightarrow\quad x^4 = \frac{L^4}{3} \quad\Leftrightarrow\quad x = \frac{L}{\sqrt[4]{3}},\ x > 0,\ L > 0.$$

Since $D''(x) = \frac{3x}{L^2} + \frac{L^2}{x^3} > 0$ for all $x > 0$, we know that $D\left(\frac{L}{\sqrt[4]{3}}\right) = \frac{(L \cdot 3^{-1/4})^3}{2L^2} + \frac{L^2}{2L \cdot 3^{-1/4})} = \frac{2L}{3^{3/4}}$ is the minimum value of D, that is, the closest the dog gets to the rabbit. The positions at this distance are

$$\text{Dog: } (x, y) = \left(\frac{L}{\sqrt[4]{3}}, \left(\frac{5}{3^{7/4}} - \frac{2}{3}\right)L\right) = \left(\frac{L}{\sqrt[4]{3}}, \frac{5\sqrt[4]{3} - 6}{9}L\right)$$

$$\text{Rabbit: } (0, 2s) = \left(0, \frac{8\sqrt[4]{3}L}{9} - \frac{2L}{3}\right) = \left(0, \frac{8\sqrt[4]{3} - 6}{9}L\right)$$

11. (a) We are given that $V = \frac{1}{3}\pi r^2 h$, $dV/dt = 60{,}000\pi$ ft^3/h, and $r = 1.5h = \frac{3}{2}h$. So $V = \frac{1}{3}\pi\left(\frac{3}{2}h\right)^2 h = \frac{3}{4}\pi h^3 \quad\Rightarrow$

$\frac{dV}{dt} = \frac{3}{4}\pi \cdot 3h^2\,\frac{dh}{dt} = \frac{9}{4}\pi h^2\,\frac{dh}{dt}$. Therefore, $\frac{dh}{dt} = \frac{4(dV/dt)}{9\pi h^2} = \frac{240{,}000\pi}{9\pi h^2} = \frac{80{,}000}{3h^2}$ $(\star) \quad\Rightarrow$

$\int 3h^2\,dh = \int 80{,}000\,dt \quad\Rightarrow\quad h^3 = 80{,}000t + C$. When $t = 0$, $h = 60$. Thus, $C = 60^3 = 216{,}000$, so

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$h^3 = 80{,}000t + 216{,}000$. Let $h = 100$. Then $100^3 = 1{,}000{,}000 = 80{,}000t + 216{,}000 \quad \Rightarrow$ $80{,}000t = 784{,}000 \quad \Rightarrow \quad t = 9.8$, so the time required is 9.8 hours.

(b) The floor area of the silo is $F = \pi \cdot 200^2 = 40{,}000\pi$ ft^2, and the area of the base of the pile is $A = \pi r^2 = \pi\left(\frac{3}{2}h\right)^2 = \frac{9\pi}{4}h^2$. So the area of the floor which is not covered when $h = 60$ is $F - A = 40{,}000\pi - 8100\pi = 31{,}900\pi \approx 100{,}217$ ft^2. Now $A = \frac{9\pi}{4}h^2 \quad \Rightarrow \quad dA/dt = \frac{9\pi}{4} \cdot 2h\,(dh/dt)$, and from ($\star$) in part (a) we know that when $h = 60$, $dh/dt = \frac{80{,}000}{3(60)^2} = \frac{200}{27}$ ft/h. Therefore, $dA/dt = \frac{9\pi}{4}(2)(60)\left(\frac{200}{27}\right) = 2000\pi \approx 6283$ ft^2/h.

(c) At $h = 90$ ft, $dV/dt = 60{,}000\pi - 20{,}000\pi = 40{,}000\pi$ ft^3/h. From ($\star$) in part (a),

$$\frac{dh}{dt} = \frac{4(dV/dt)}{9\pi h^2} = \frac{4(40{,}000\pi)}{9\pi h^2} = \frac{160{,}000}{9h^2} \quad \Rightarrow \quad \int 9h^2\,dh = \int 160{,}000\,dt \quad \Rightarrow \quad 3h^3 = 160{,}000t + C.$$

When $t = 0$, $h = 90$; therefore, $C = 3 \cdot 729{,}000 = 2{,}187{,}000$. So $3h^3 = 160{,}000t + 2{,}187{,}000$. At the top, $h = 100 \quad \Rightarrow$ $3(100)^3 = 160{,}000t + 2{,}187{,}000 \quad \Rightarrow \quad t = \frac{813{,}000}{160{,}000} \approx 5.1$. The pile reaches the top after about 5.1 h.

12. Let $P(a, b)$ be any first-quadrant point on the curve $y = f(x)$. The tangent line at P has equation $y - b = f'(a)(x - a)$, or equivalently, $y = mx + b - ma$, where $m = f'(a)$. If $Q(0, c)$ is the y-intercept, then $c = b - am$. If $R(k, 0)$ is the x-intercept, then $k = \dfrac{am - b}{m} = a - \dfrac{b}{m}$. Since the tangent line is bisected at P, we know that $|PQ| = |PR|$; that is, $\sqrt{(a-0)^2 + [b - (b - am)]^2} = \sqrt{[a - (a - b/m)]^2 + (b - 0)^2}$. Squaring and simplifying gives us $a^2 + a^2m^2 = b^2/m^2 + b^2 \quad \Rightarrow \quad a^2m^2 + a^2m^4 = b^2 + b^2m^2 \quad \Rightarrow \quad a^2m^4 + \left(a^2 - b^2\right)m^2 - b^2 = 0 \quad \Rightarrow$ $\left(a^2m^2 - b^2\right)\left(m^2 + 1\right) = 0 \quad \Rightarrow \quad m^2 = b^2/a^2$. Since m is the slope of the line from a positive y-intercept to a positive x-intercept, m must be negative. Since a and b are positive, we have $m = -b/a$, so we will solve the equivalent differential equation $\dfrac{dy}{dx} = -\dfrac{y}{x} \quad \Rightarrow \quad \dfrac{dy}{y} = -\dfrac{dx}{x} \quad \Rightarrow \quad \displaystyle\int \frac{dy}{y} = -\int \frac{dx}{x} \quad \Rightarrow \quad \ln y = -\ln x + C \;\; [x, y > 0] \quad \Rightarrow$ $y = e^{-\ln x + C} = e^{\ln x^{-1}} \cdot e^C = x^{-1} \cdot A \quad \Rightarrow \quad y = A/x$. Since the point $(3, 2)$ is on the curve, $3 = A/2 \quad \Rightarrow \quad A = 6$ and the curve is $y = 6/x$ with $x > 0$.

13. Let $P(a, b)$ be any point on the curve. If m is the slope of the tangent line at P, then $m = y'(a)$, and an equation of the normal line at P is $y - b = -\dfrac{1}{m}(x - a)$, or equivalently, $y = -\dfrac{1}{m}x + b + \dfrac{a}{m}$. The y-intercept is always 6, so $b + \dfrac{a}{m} = 6 \quad \Rightarrow \quad \dfrac{a}{m} = 6 - b \quad \Rightarrow \quad m = \dfrac{a}{6 - b}$. We will solve the equivalent differential equation $\dfrac{dy}{dx} = \dfrac{x}{6 - y} \quad \Rightarrow$

$$(6 - y)\,dy = x\,dx \quad \Rightarrow \quad \int (6 - y)\,dy = \int x\,dx \quad \Rightarrow \quad 6y - \tfrac{1}{2}y^2 = \tfrac{1}{2}x^2 + C \quad \Rightarrow \quad 12y - y^2 = x^2 + K.$$

Since $(3, 2)$ is on the curve, $12(2) - 2^2 = 3^2 + K \quad \Rightarrow \quad K = 11$. So the curve is given by $12y - y^2 = x^2 + 11 \quad \Rightarrow$ $x^2 + y^2 - 12y + 36 = -11 + 36 \quad \Rightarrow \quad x^2 + (y - 6)^2 = 25$, a circle with center $(0, 6)$ and radius 5.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. Let $P(x_0, y_0)$ be a point on the curve. Since the midpoint of the line segment determined by the normal line from (x_0, y_0) to its intersection with the x-axis has x-coordinate 0, the x-coordinate of the point of intersection with the x-axis must be $-x_0$. Hence, the normal line has slope $\dfrac{y_0 - 0}{x_0 - (-x_0)} = \dfrac{y_0}{2x_0}$. So the tangent line has slope $-\dfrac{2x_0}{y_0}$. This gives the differential equation $y' = -\dfrac{2x}{y} \quad \Rightarrow \quad y\,dy = -2x\,dx \quad \Rightarrow \quad \int y\,dy = \int(-2x)\,dx \quad \Rightarrow \quad \frac{1}{2}y^2 = -x^2 + C \quad \Rightarrow \quad x^2 + \frac{1}{2}y^2 = C$ $[C > 0]$. This is a family of ellipses.

15. From the figure, slope $OA = \dfrac{y}{x}$. If triangle OAB is isosceles, then slope AB must be $-\dfrac{y}{x}$, the negative of slope OA. This slope is also equal to $f'(x)$, so we have $\dfrac{dy}{dx} = -\dfrac{y}{x} \quad \Rightarrow \quad \displaystyle\int \frac{dy}{y} = -\int \frac{dx}{x} \quad \Rightarrow$

$\ln|y| = -\ln|x| + C \quad \Rightarrow \quad |y| = e^{-\ln|x| + C} \quad \Rightarrow$

$|y| = (e^{\ln|x|})^{-1}e^C \quad \Rightarrow \quad |y| = \dfrac{1}{|x|}e^C \quad \Rightarrow \quad y = \dfrac{K}{x}, K \neq 0.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10 ☐ PARAMETRIC EQUATIONS AND POLAR COORDINATES

10.1 Curves Defined by Parametric Equations

1. $x = t^2 + t, \quad y = 3^{t+1}, \quad t = -2, -1, 0, 1, 2$

t	-2	-1	0	1	2
x	2	0	0	2	6
y	$\frac{1}{3}$	1	3	9	27

Therefore, the coordinates are $\left(2, \frac{1}{3}\right)$, $(0, 1)$, $(0, 3)$, $(2, 9)$, and $(6, 27)$.

2. $x = \ln(t^2 + 1), \quad y = t/(t + 4), \quad t = -2, -1, 0, 1, 2$

t	-2	-1	0	1	2
x	$\ln 5$	$\ln 2$	0	$\ln 2$	$\ln 5$
y	-1	$-\frac{1}{3}$	0	$\frac{1}{5}$	$\frac{1}{3}$

Therefore, the coordinates are $(\ln 5, -1)$, $\left(\ln 2, -\frac{1}{3}\right)$, $(0, 0)$, $\left(\ln 2, \frac{1}{5}\right)$, and $\left(\ln 5, \frac{1}{3}\right)$.

3. $x = 1 - t^2, \quad y = 2t - t^2, \quad -1 \le t \le 2$

t	-1	0	1	2
x	0	1	0	-3
y	-3	0	1	0

4. $x = t^3 + t, \quad y = t^2 + 2, \quad -2 \le t \le 2$

t	-2	-1	0	1	2
x	-10	-2	0	2	10
y	6	3	2	3	6

5. $x = 2^t - t, \quad y = 2^{-t} + t, \quad -3 \le t \le 3$

t	-3	-2	-1	0	1	2	3
x	3.125	2.25	1.5	1	1	2	5
y	5	2	1	1	1.5	2.25	3.125

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $x = \cos^2 t, \quad y = 1 + \cos t, \quad 0 \le t \le \pi$

t	0	$\pi/4$	$\pi/2$	$3\pi/4$	π
x	1	0.5	0	0.5	1
y	2	1.707	1	0.293	0

7. $x = 2t - 1, \quad y = \frac{1}{2}t + 1$

(a)

t	-4	-2	0	2	4
x	-9	-5	-1	3	7
y	-1	0	1	2	3

(b) $x = 2t - 1 \quad \Rightarrow \quad 2t = x + 1 \quad \Rightarrow \quad t = \frac{1}{2}x + \frac{1}{2}$, so

$y = \frac{1}{2}t + 1 = \frac{1}{2}\left(\frac{1}{2}x + \frac{1}{2}\right) + 1 = \frac{1}{4}x + \frac{1}{4} + 1 \quad \Rightarrow \quad y = \frac{1}{4}x + \frac{5}{4}$

8. $x = 3t + 2, \quad y = 2t + 3$

(a)

t	-4	-2	0	2	4
x	-10	-4	2	8	14
y	-5	-1	3	7	11

(b) $x = 3t + 2 \quad \Rightarrow \quad 3t = x - 2 \quad \Rightarrow \quad t = \frac{1}{3}x - \frac{2}{3}$, so

$y = 2t + 3 = 2\left(\frac{1}{3}x - \frac{2}{3}\right) + 3 = \frac{2}{3}x - \frac{4}{3} + 3 \quad \Rightarrow \quad y = \frac{2}{3}x + \frac{5}{3}$

9. $x = t^2 - 3, \quad y = t + 2, \quad -3 \le t \le 3$

(a)

t	-3	-1	1	3
x	6	-2	-2	6
y	-1	1	3	5

(b) $y = t + 2 \quad \Rightarrow \quad t = y - 2$, so

$x = t^2 - 3 = (y - 2)^2 - 3 = y^2 - 4y + 4 - 3 \quad \Rightarrow$

$x = y^2 - 4y + 1, -1 \le y \le 5$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $x = \sin t, \quad y = 1 - \cos t, \quad 0 \le t \le 2\pi$

(a)

t	0	$\pi/2$	π	$3\pi/2$	2π
x	0	1	0	-1	0
y	0	1	2	1	0

(b) $x = \sin t, y = 1 - \cos t \quad [\text{or } y - 1 = -\cos t] \quad \Rightarrow$

$x^2 + (y-1)^2 = (\sin t)^2 + (-\cos t)^2 \quad \Rightarrow \quad x^2 + (y-1)^2 = 1.$

As t varies from 0 to 2π, the circle with center $(0, 1)$ and radius 1 is traced out.

11. $x = \sqrt{t}, \; y = 1 - t$

(a)

t	0	1	2	3	4
x	0	1	1.414	1.732	2
y	1	0	-1	-2	-3

(b) $x = \sqrt{t} \quad \Rightarrow \quad t = x^2 \quad \Rightarrow \quad y = 1 - t = 1 - x^2$. Since $t \ge 0$, $x \ge 0$.

So the curve is the right half of the parabola $y = 1 - x^2$.

12. $x = t^2, \quad y = t^3$

(a)

t	-2	-1	0	1	2
x	4	1	0	1	4
y	-8	-1	0	1	8

(b) $y = t^3 \quad \Rightarrow \quad t = \sqrt[3]{y} \quad \Rightarrow \quad x = t^2 = \left(\sqrt[3]{y}\right)^2 = y^{2/3}. \quad t \in \mathbb{R}, y \in \mathbb{R}, x \ge 0.$

13. (a) $x = 3\cos t, \quad y = 3\sin t, \quad 0 \le t \le \pi$

$x^2 + y^2 = 9\cos^2 t + 9\sin^2 t = 9(\cos^2 t + \sin^2 t) = 9$, which is the equation of a circle with radius 3. For $0 \le t \le \pi/2$, we have $3 \ge x \ge 0$ and $0 \le y \le 3$. For $\pi/2 < t \le \pi$, we have $0 > x \ge -3$ and $3 > y \ge 0$. Thus, the curve is the top half of the circle $x^2 + y^2 = 9$ traced counterclockwise.

(b)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. (a) $x = \sin 4\theta, \quad y = \cos 4\theta, \quad 0 \le \theta \le \pi/2$

$x^2 + y^2 = \sin^2 4\theta + \cos^2 4\theta = 1$, which is the equation of a circle with radius 1. When $\theta = 0$, we have $x = 0$ and $y = 1$. For $0 \le \theta \le \pi/4$, we have $x \ge 0$. For $\pi/4 < \theta \le \pi/2$, we have $x \le 0$. Thus, the curve is the circle $x^2 + y^2 = 1$ traced clockwise starting at $(0, 1)$.

(b)

15. (a) $x = \cos\theta, \quad y = \sec^2\theta, \quad 0 \le \theta < \pi/2.$

$y = \sec^2\theta = \dfrac{1}{\cos^2\theta} = \dfrac{1}{x^2}$. For $0 \le \theta < \pi/2$, we have $1 \ge x > 0$ and $1 \le y$.

(b)

16. (a) $x = \csc t, y = \cot t, 0 < t < \pi$

$y^2 - x^2 = \cot^2 t - \csc^2 t = 1$. For $0 < t < \pi$, we have $x > 1$. Thus, the curve is the right branch of the hyperbola $y^2 - x^2 = 1$.

(b)

17. (a) $y = e^t = 1/e^{-t} = 1/x$ for $x > 0$ since $x = e^{-t}$. Thus, the curve is the portion of the hyperbola $y = 1/x$ with $x > 0$.

(b)

18. (a) $x = t + 2 \quad \Rightarrow \quad t = x - 2$. $y = 1/t = 1/(x - 2)$. For $t > 0$, we have $x > 2$ and $y > 0$. Thus, the curve is the portion of the hyperbola $y = 1/(x - 2)$ with $x > 2$.

(b)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. (a) $x = \ln t$, $y = \sqrt{t}$, $t \geq 1$.

$x = \ln t \quad \Rightarrow \quad t = e^x \quad \Rightarrow \quad y = \sqrt{t} = e^{x/2}$, $x \geq 0$.

(b)

20. (a) $x = |t|$, $y = \Big|1 - |t|\Big| = |1 - x|$. For all t, we have $x \geq 0$ and $y \geq 0$. Thus, the curve is the portion of the absolute value function $y = |1 - x|$ with $x \geq 0$.

(b)

21. (a) $x = \sin^2 t$, $y = \cos^2 t$. $x + y = \sin^2 t + \cos^2 t = 1$. For all t, we have $0 \leq x \leq 1$ and $0 \leq y \leq 1$. Thus, the curve is the portion of the line $x + y = 1$ or $y = -x + 1$ in the first quadrant.

(b)

22. (a) $x = \sinh t$, $y = \cosh t \quad \Rightarrow \quad y^2 - x^2 = \cosh^2 t - \sinh^2 t = 1$. Since $y = \cosh t \geq 1$, we have the upper branch of the hyperbola $y^2 - x^2 = 1$.

(b)

23. The parametric equations $x = 5\cos t$ and $y = -5\sin t$ both have period 2π. When $t = 0$, we have $x = 5$ and $y = 0$. When $t = \pi/2$, we have $x = 0$ and $y = -5$. This is one-fourth of a circle. Thus, the object completes one revolution in $4 \cdot \frac{\pi}{2} = 2\pi$ seconds following a clockwise path.

24. The parametric equations $x = 3\sin\left(\dfrac{\pi}{4}t\right)$ and $y = 3\cos\left(\dfrac{\pi}{4}t\right)$ both have period $\dfrac{2\pi}{\pi/4} = 8$. When $t = 0$, we have $x = 0$ and $y = 3$. When $t = 2$, we have $x = 3$ and $y = 0$. This is one-fourth of a circle. Thus, the object completes one revolution in $4 \cdot 2 = 8$ seconds following a clockwise path.

25. $x = 5 + 2\cos \pi t$, $y = 3 + 2\sin \pi t \quad \Rightarrow \quad \cos \pi t = \dfrac{x-5}{2}$, $\sin \pi t = \dfrac{y-3}{2}$. $\cos^2(\pi t) + \sin^2(\pi t) = 1 \quad \Rightarrow$ $\left(\dfrac{x-5}{2}\right)^2 + \left(\dfrac{y-3}{2}\right)^2 = 1$. The motion of the particle takes place on a circle centered at $(5, 3)$ with a radius 2. As t goes from 1 to 2, the particle starts at the point $(3, 3)$ and moves counterclockwise along the circle $\left(\dfrac{x-5}{2}\right)^2 + \left(\dfrac{y-3}{2}\right)^2 = 1$ to $(7, 3)$ [one-half of a circle].

26. $x = 2 + \sin t$, $y = 1 + 3\cos t \quad \Rightarrow \quad \sin t = x - 2$, $\cos t = \dfrac{y-1}{3}$. $\sin^2 t + \cos^2 t = 1 \quad \Rightarrow \quad (x-2)^2 + \left(\dfrac{y-1}{3}\right)^2 = 1$. The motion of the particle takes place on an ellipse centered at $(2, 1)$. As t goes from $\pi/2$ to 2π, the particle starts at the point $(3, 1)$ and moves counterclockwise three-fourths of the way around the ellipse to $(2, 4)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. $x = 5\sin t,\ y = 2\cos t \quad \Rightarrow \quad \sin t = \dfrac{x}{5},\ \cos t = \dfrac{y}{2}. \quad \sin^2 t + \cos^2 t = 1 \quad \Rightarrow \quad \left(\dfrac{x}{5}\right)^2 + \left(\dfrac{y}{2}\right)^2 = 1.$ The motion of the particle takes place on an ellipse centered at $(0, 0)$. As t goes from $-\pi$ to 5π, the particle starts at the point $(0, -2)$ and moves clockwise around the ellipse 3 times.

28. $y = \cos^2 t = 1 - \sin^2 t = 1 - x^2$. The motion of the particle takes place on the parabola $y = 1 - x^2$. As t goes from -2π to $-\pi$, the particle starts at the point $(0, 1)$, moves to $(1, 0)$, and goes back to $(0, 1)$. As t goes from $-\pi$ to 0, the particle moves to $(-1, 0)$ and goes back to $(0, 1)$. The particle repeats this motion as t goes from 0 to 2π.

29. We must have $1 \le x \le 4$ and $2 \le y \le 3$. So the graph of the curve must be contained in the rectangle $[1, 4]$ by $[2, 3]$.

30. (a) From the first graph, we have $1 \le x \le 2$. From the second graph, we have $-1 \le y \le 1$. The only choice that satisfies either of those conditions is III.

(b) From the first graph, the values of x cycle through the values from -2 to 2 four times. From the second graph, the values of y cycle through the values from -2 to 2 six times. Choice I satisfies these conditions.

(c) From the first graph, the values of x cycle through the values from -2 to 2 three times. From the second graph, we have $0 \le y \le 2$. Choice IV satisfies these conditions.

(d) From the first graph, the values of x cycle through the values from -2 to 2 two times. From the second graph, the values of y do the same thing. Choice II satisfies these conditions.

31. When $t = -1$, $(x, y) = (1, 1)$. As t increases to 0, x and y both decrease to 0. As t increases from 0 to 1, x increases from 0 to 1 and y decreases from 0 to -1. As t increases beyond 1, x continues to increase and y continues to decrease. For $t < -1$, x and y are both positive and decreasing. We could achieve greater accuracy by estimating x- and y-values for selected values of t from the given graphs and plotting the corresponding points.

32. When $t = -1$, $(x, y) = (0, 0)$. As t increases to 0, x increases from 0 to 1, while y first decreases to -1 and then increases to 0. As t increases from 0 to 1, x decreases from 1 to 0, while y first increases to 1 and then decreases to 0. We could achieve greater accuracy by estimating x- and y-values for selected values of t from the given graphs and plotting the corresponding points.

33. When $t = -1$, $(x, y) = (0, 1)$. As t increases to 0, x increases from 0 to 1 and y decreases from 1 to 0. As t increases from 0 to 1, the curve is retraced in the opposite direction with x decreasing from 1 to 0 and y increasing from 0 to 1. We could achieve greater accuracy by estimating x- and y-values for selected values of t from the given graphs and plotting the corresponding points.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. (a) $x = t^4 - t + 1 = (t^4 + 1) - t > 0$ [think of the graphs of $y = t^4 + 1$ and $y = t$] and $y = t^2 \geq 0$, so these equations are matched with graph V.

(b) $y = \sqrt{t} \geq 0$. $x = t^2 - 2t = t(t-2)$ is negative for $0 < t < 2$, so these equations are matched with graph I.

(c) $x = t^3 - 2t = t(t^2 - 2) = t\left(t + \sqrt{2}\right)\left(t - \sqrt{2}\right)$, $y = t^2 - t = t(t-1)$. The equation $x = 0$ has three solutions and the equation $y = 0$ has two solutions. Thus, the curve has three y-intercepts and two x-intercepts, which matches graph II.

Alternate method: $x = t^3 - 2t$, $y = t^2 - t = \left(t^2 - t + \frac{1}{4}\right) - \frac{1}{4} = \left(t - \frac{1}{2}\right)^2 - \frac{1}{4}$ so $y \geq -\frac{1}{4}$ on this curve, whereas x is unbounded. These equations are matched with graph II.

(d) $x = \cos 5t$ has period $2\pi/5$ and $y = \sin 2t$ has period π, so x will take on the values -1 to 1, and then 1 to -1, before y takes on the values -1 to 1. Note that when $t = 0$, $(x, y) = (1, 0)$. These equations are matched with graph VI.

(e) $x = t + \sin 4t$, $y = t^2 + \cos 3t$. As t becomes large, t and t^2 become the dominant terms in the expressions for x and y, so the graph will look like the graph of $y = x^2$, but with oscillations. These equations are matched with graph IV.

(f) $x = t + \sin 2t$, $y = t + \sin 3t$. As t becomes large, t becomes the dominant term in the expressions for both x and y, so the graph will look like the graph of $y = x$, but with oscillations. These equations are matched with graph III.

35. Use $y = t$ and $x = t - 2\sin \pi t$ with a t-interval of $[-\pi, \pi]$.

36. Use $x_1 = t$, $y_1 = t^3 - 4t$ and $x_2 = t^3 - 4t$, $y_2 = t$ with a t-interval of $[-3, 3]$. There are 9 points of intersection; $(0, 0)$ is fairly obvious. The point in quadrant I is approximately $(2.2, 2.2)$, and by symmetry, the point in quadrant III is approximately $(-2.2, -2.2)$. The other six points are approximately $(\mp 1.9, \pm 0.5)$, $(\mp 1.7, \pm 1.7)$, and $(\mp 0.5, \pm 1.9)$.

37. (a) $x = x_1 + (x_2 - x_1)t$, $y = y_1 + (y_2 - y_1)t$, $0 \leq t \leq 1$. Clearly the curve passes through $P_1(x_1, y_1)$ when $t = 0$ and through $P_2(x_2, y_2)$ when $t = 1$. For $0 < t < 1$, x is strictly between x_1 and x_2 and y is strictly between y_1 and y_2. For every value of t, x and y satisfy the relation $y - y_1 = \dfrac{y_2 - y_1}{x_2 - x_1}(x - x_1)$, which is the equation of the line through $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$.

Finally, any point (x, y) on that line satisfies $\dfrac{y - y_1}{y_2 - y_1} = \dfrac{x - x_1}{x_2 - x_1}$; if we call that common value t, then the given parametric equations yield the point (x, y); and any (x, y) on the line between $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$ yields a value of t in $[0, 1]$. So the given parametric equations exactly specify the line segment from $P_1(x_1, y_1)$ to $P_2(x_2, y_2)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $x = -2 + [3 - (-2)]t = -2 + 5t$ and $y = 7 + (-1 - 7)t = 7 - 8t$ for $0 \le t \le 1$.

38. For the side of the triangle from A to B, use $(x_1, y_1) = (1, 1)$ and $(x_2, y_2) = (4, 2)$. Hence, the equations are

$$x = x_1 + (x_2 - x_1)\,t = 1 + (4 - 1)\,t = 1 + 3t,$$
$$y = y_1 + (y_2 - y_1)\,t = 1 + (2 - 1)\,t = 1 + t.$$

Graphing $x = 1 + 3t$ and $y = 1 + t$ with $0 \le t \le 1$ gives us the side of the triangle from A to B. Similarly, for the side BC we use $x = 4 - 3t$ and $y = 2 + 3t$, and for the side AC we use $x = 1$ and $y = 1 + 4t$.

39. The result in Example 4 indicates the parametric equations have the form $x = h + r\sin bt$ and $y = k + r\cos bt$ where (h, k) is the center of the circle with radius r and $b = 2\pi/\text{period}$. (The use of positive sine in the x-equation and positive cosine in the y-equation results in a clockwise motion.) With $h = 0$, $k = 0$ and $b = 2\pi/4\pi = 1/2$, we have $x = 5\sin\left(\frac{1}{2}t\right)$, $y = 5\cos\left(\frac{1}{2}t\right)$.

40. As in Example 4, we use parametric equations of the form $x = h + r\cos bt$ and $y = k + r\sin bt$ where $(h, k) = (1, 3)$ is the center of the circle with radius $r = 1$ and $b = 2\pi/\text{period} = 2\pi/3$. (The use of positive cosine in the x-equation and positive sine in the y-equation results in a counterclockwise motion.) Thus, $x = 1 + \cos\left(\frac{2\pi}{3}t\right)$, $y = 3 + \sin\left(\frac{2\pi}{3}t\right)$.

41. The circle $x^2 + (y - 1)^2 = 4$ has center $(0, 1)$ and radius 2, so by Example 4 it can be represented by $x = 2\cos t$, $y = 1 + 2\sin t$, $0 \le t \le 2\pi$. This representation gives us the circle with a counterclockwise orientation starting at $(2, 1)$.

(a) To get a clockwise orientation, we could change the equations to $x = 2\cos t$, $y = 1 - 2\sin t$, $0 \le t \le 2\pi$.

(b) To get three times around in the counterclockwise direction, we use the original equations $x = 2\cos t$, $y = 1 + 2\sin t$ with the domain expanded to $0 \le t \le 6\pi$.

(c) To start at $(0, 3)$ using the original equations, we must have $x_1 = 0$; that is, $2\cos t = 0$. Hence, $t = \frac{\pi}{2}$. So we use $x = 2\cos t$, $y = 1 + 2\sin t$, $\frac{\pi}{2} \le t \le \frac{3\pi}{2}$.

Alternatively, if we want t to start at 0, we could change the equations of the curve. For example, we could use $x = -2\sin t$, $y = 1 + 2\cos t$, $0 \le t \le \pi$.

42. (a) Let $x^2/a^2 = \sin^2 t$ and $y^2/b^2 = \cos^2 t$ to obtain $x = a\sin t$ and $y = b\cos t$ with $0 \le t \le 2\pi$ as possible parametric equations for the ellipse $x^2/a^2 + y^2/b^2 = 1$.

(b) The equations are $x = 3\sin t$ and $y = b\cos t$ for $b \in \{1, 2, 4, 8\}$.

(c) As b increases, the ellipse stretches vertically.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. *Big circle:* It's centered at $(2, 2)$ with a radius of 2, so by Example 4, parametric equations are

$$x = 2 + 2\cos t, \qquad y = 2 + 2\sin t, \qquad 0 \le t \le 2\pi$$

Small circles: They are centered at $(1, 3)$ and $(3, 3)$ with a radius of 0.1. By Example 4, parametric equations are

$$\textit{(left)} \qquad x = 1 + 0.1\cos t, \qquad y = 3 + 0.1\sin t, \qquad 0 \le t \le 2\pi$$

and

$$\textit{(right)} \qquad x = 3 + 0.1\cos t, \qquad y = 3 + 0.1\sin t, \qquad 0 \le t \le 2\pi$$

Semicircle: It's the lower half of a circle centered at $(2, 2)$ with radius 1. By Example 4, parametric equations are

$$x = 2 + 1\cos t, \qquad y = 2 + 1\sin t, \qquad \pi \le t \le 2\pi$$

To get all four graphs on the same screen with a typical graphing calculator, we need to change the last t-interval to$[0, 2\pi]$ in order to match the others. We can do this by changing t to $0.5t$. This change gives us the upper half. There are several ways to get the lower half—one is to change the "+" to a "−" in the y-assignment, giving us

$$x = 2 + 1\cos(0.5t), \qquad y = 2 - 1\sin(0.5t), \qquad 0 \le t \le 2\pi$$

44. If you are using a calculator or computer that can overlay graphs (using multiple t-intervals), the following is appropriate.

Left side: $x = 1$ and y goes from 1.5 to 4, so use

$$x = 1, \qquad y = t, \qquad 1.5 \le t \le 4$$

Right side: $x = 10$ and y goes from 1.5 to 4, so use

$$x = 10, \qquad y = t, \qquad 1.5 \le t \le 4$$

Bottom: x goes from 1 to 10 and $y = 1.5$, so use

$$x = t, \qquad y = 1.5, \qquad 1 \le t \le 10$$

Handle: It starts at $(10, 4)$ and ends at $(13, 7)$, so use

$$x = 10 + t, \qquad y = 4 + t, \qquad 0 \le t \le 3$$

Left wheel: It's centered at $(3, 1)$, has a radius of 1, and appears to go about $30°$ above the horizontal, so use

$$x = 3 + 1\cos t, \qquad y = 1 + 1\sin t, \qquad \tfrac{5\pi}{6} \le t \le \tfrac{13\pi}{6}$$

Right wheel: Similar to the left wheel with center $(8, 1)$, so use

$$x = 8 + 1\cos t, \qquad y = 1 + 1\sin t, \qquad \tfrac{5\pi}{6} \le t \le \tfrac{13\pi}{6}$$

If you are using a calculator or computer that cannot overlay graphs (using one t-interval), the following is appropriate. We'll start by picking the t-interval $[0, 2.5]$ since it easily matches the t-values for the two sides. We now need to find parametric equations for all graphs with $0 \le t \le 2.5$.

Left side: $x = 1$ and y goes from 1.5 to 4, so use

$$x = 1, \qquad y = 1.5 + t, \qquad 0 \le t \le 2.5$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Right side: $x = 10$ and y goes from 1.5 to 4, so use

$$x = 10, \qquad y = 1.5 + t, \qquad 0 \le t \le 2.5$$

Bottom: x goes from 1 to 10 and $y = 1.5$, so use

$$x = 1 + 3.6t, \qquad y = 1.5, \qquad 0 \le t \le 2.5$$

To get the x-assignment, think of creating a linear function such that when $t = 0$, $x = 1$ and when $t = 2.5$, $x = 10$. We can use the point-slope form of a line with $(t_1, x_1) = (0, 1)$ and $(t_2, x_2) = (2.5, 10)$.

$x - 1 = \dfrac{10 - 1}{2.5 - 0}(t - 0) \quad \Rightarrow \quad x = 1 + 3.6t.$

Handle: It starts at $(10, 4)$ and ends at $(13, 7)$, so use

$$x = 10 + 1.2t, \qquad y = 4 + 1.2t, \qquad 0 \le t \le 2.5$$

$(t_1, x_1) = (0, 10)$ and $(t_2, x_2) = (2.5, 13)$ gives us $x - 10 = \dfrac{13 - 10}{2.5 - 0}(t - 0) \quad \Rightarrow \quad x = 10 + 1.2t.$

$(t_1, y_1) = (0, 4)$ and $(t_2, y_2) = (2.5, 7)$ gives us $y - 4 = \dfrac{7 - 4}{2.5 - 0}(t - 0) \quad \Rightarrow \quad y = 4 + 1.2t.$

Left wheel: It's centered at $(3, 1)$, has a radius of 1, and appears to go about 30° above the horizontal, so use

$$x = 3 + 1\cos\left(\tfrac{8\pi}{15}t + \tfrac{5\pi}{6}\right), \qquad y = 1 + 1\sin\left(\tfrac{8\pi}{15}t + \tfrac{5\pi}{6}\right), \qquad 0 \le t \le 2.5$$

$(t_1, \theta_1) = \left(0, \tfrac{5\pi}{6}\right)$ and $(t_2, \theta_2) = \left(\tfrac{5}{2}, \tfrac{13\pi}{6}\right)$ gives us $\theta - \tfrac{5\pi}{6} = \dfrac{\frac{13\pi}{6} - \frac{5\pi}{6}}{\frac{5}{2} - 0}(t - 0) \quad \Rightarrow \quad \theta = \tfrac{5\pi}{6} + \tfrac{8\pi}{15}t.$

Right wheel: Similar to the left wheel with center $(8, 1)$, so use

$$x = 8 + 1\cos\left(\tfrac{8\pi}{15}t + \tfrac{5\pi}{6}\right), \qquad y = 1 + 1\sin\left(\tfrac{8\pi}{15}t + \tfrac{5\pi}{6}\right), \qquad 0 \le t \le 2.5$$

45. (a) (i) $x = t^2, y = t \quad \Rightarrow \quad y^2 = t^2 = x$ (ii) $x = t, y = \sqrt{t} \quad \Rightarrow \quad y^2 = t = x$

(iii) $x = \cos^2 t, y = \cos t \quad \Rightarrow \quad y^2 = \cos^2 t = x$ (iv) $x = 3^{2t}, y = 3^t \quad \Rightarrow \quad y^2 = (3^t)^2 = 3^{2t} = x.$

Thus, the points on all four of the given parametric curves satisfy the Cartesian equation $y^2 = x$.

(b) The graph of $y^2 = x$ is a right-opening parabola with vertex at the origin. For curve (i), $x \ge 0$ and y is unbounded so the graph contains the entire parabola. For (ii), $y = \sqrt{t}$ requires that $t \ge 0$, so that both $x \ge 0$ and $y \ge 0$, which captures the upper half of the parabola, including the origin. For (iii), $-1 \le \cos t \le 1$ so the graph is the portion of the parabola contained in the intervals $0 \le x \le 1$ and $-1 \le y \le 1$. For (iv), $x > 0$ and $y > 0$, which captures the upper half of the parabola excluding the origin.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. (a) $x = t$, so $y = t^{-2} = x^{-2}$. We get the entire curve $y = 1/x^2$ traversed in a left-to-right direction.

(b) $x = \cos t$, $y = \sec^2 t = \dfrac{1}{\cos^2 t} = \dfrac{1}{x^2}$. Since $\sec t \geq 1$, we only get the parts of the curve $y = 1/x^2$ with $y \geq 1$. We get the first quadrant portion of the curve when $x > 0$, that is, $\cos t > 0$, and we get the second quadrant portion of the curve when $x < 0$, that is, $\cos t < 0$.

(c) $x = e^t$, $y = e^{-2t} = (e^t)^{-2} = x^{-2}$. Since e^t and e^{-2t} are both positive, we only get the first quadrant portion of the curve $y = 1/x^2$.

47. (a) $x = t^3 \quad \Rightarrow \quad t = x^{1/3}$, so $y = t^2 = x^{2/3}$.

We get the entire curve $y = x^{2/3}$ traversed in a left to right direction.

(b) $x = t^6 \quad \Rightarrow \quad t = x^{1/6}$, so $y = t^4 = x^{4/6} = x^{2/3}$.

Since $x = t^6 \geq 0$, we only get the right half of the curve $y = x^{2/3}$.

(c) $x = e^{-3t} = (e^{-t})^3 \quad [\text{so } e^{-t} = x^{1/3}]$,

$y = e^{-2t} = (e^{-t})^2 = (x^{1/3})^2 = x^{2/3}$.

If $t < 0$, then x and y are both larger than 1. If $t > 0$, then x and y are between 0 and 1. Since $x > 0$ and $y > 0$, the curve never quite reaches the origin.

48. The case $\frac{\pi}{2} < \theta < \pi$ is illustrated. C has coordinates $(r\theta, r)$ as in Example 7, and Q has coordinates $(r\theta, r + r\cos(\pi - \theta)) = (r\theta, r(1 - \cos\theta))$ [since $\cos(\pi - \alpha) = \cos\pi\cos\alpha + \sin\pi\sin\alpha = -\cos\alpha$], so P has coordinates $(r\theta - r\sin(\pi - \theta), r(1 - \cos\theta)) = (r(\theta - \sin\theta), r(1 - \cos\theta))$ [since $\sin(\pi - \alpha) = \sin\pi\cos\alpha - \cos\pi\sin\alpha = \sin\alpha$]. Again we have the parametric equations $x = r(\theta - \sin\theta)$, $y = r(1 - \cos\theta)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. The first two diagrams depict the case $\pi < \theta < \frac{3\pi}{2}$, $d < r$. As in Example 7, C has coordinates $(r\theta, r)$. Now Q (in the second diagram) has coordinates $(r\theta, r + d\cos(\theta - \pi)) = (r\theta, r - d\cos\theta)$, so a typical point P of the trochoid has coordinates $(r\theta + d\sin(\theta - \pi), r - d\cos\theta)$. That is, P has coordinates (x, y), where $x = r\theta - d\sin\theta$ and $y = r - d\cos\theta$. When $d = r$, these equations agree with those of the cycloid.

50. In polar coordinates, an equation for the circle is $r = 2a\sin\theta$. Thus, the coordinates of Q are $x = r\cos\theta = 2a\sin\theta\,\cos\theta$ and $y = r\sin\theta = 2a\sin^2\theta$. The coordinates of R are $x = 2a\cot\theta$ and $y = 2a$. Since P is the midpoint of QR, we use the midpoint formula to get $x = a(\sin\theta\,\cos\theta + \cot\theta)$ and $y = a(1 + \sin^2\theta)$.

51. It is apparent that $x = |OQ|$ and $y = |QP| = |ST|$. From the diagram, $x = |OQ| = a\cos\theta$ and $y = |ST| = b\sin\theta$. Thus, the parametric equations are $x = a\cos\theta$ and $y = b\sin\theta$. To eliminate θ we rearrange: $\sin\theta = y/b \;\Rightarrow\; \sin^2\theta = (y/b)^2$ and $\cos\theta = x/a \;\Rightarrow\; \cos^2\theta = (x/a)^2$. Adding the two equations: $\sin^2\theta + \cos^2\theta = 1 = x^2/a^2 + y^2/b^2$. Thus, we have an ellipse.

52. A has coordinates $(a\cos\theta, a\sin\theta)$. Since OA is perpendicular to AB, ΔOAB is a right triangle and B has coordinates $(a\sec\theta, 0)$. It follows that P has coordinates $(a\sec\theta, b\sin\,\theta)$. Thus, the parametric equations are $x = a\sec\theta$, $y = b\sin\theta$.

53. $C = (2a\cot\theta, 2a)$, so the x-coordinate of P is $x = 2a\cot\theta$. Let $B = (0, 2a)$. Then $\angle OAB$ is a right angle and $\angle OBA = \theta$, so $|OA| = 2a\sin\theta$ and $A = ((2a\sin\theta)\cos\theta, (2a\sin\theta)\sin\theta)$. Thus, the y-coordinate of P is $y = 2a\sin^2\theta$.

54. (a) Let θ be the angle of inclination of segment OP. Then $|OB| = \dfrac{2a}{\cos\theta}$.

Let $C = (2a, 0)$. Then by use of right triangle OAC we see that $|OA| = 2a\cos\theta$.

Now

$$|OP| = |AB| = |OB| - |OA|$$
$$= 2a\left(\frac{1}{\cos\theta} - \cos\theta\right) = 2a\,\frac{1 - \cos^2\theta}{\cos\theta} = 2a\,\frac{\sin^2\theta}{\cos\theta} = 2a\sin\theta\,\tan\theta$$

So P has coordinates $x = 2a\sin\theta\,\tan\theta\cdot\cos\theta = 2a\sin^2\theta$ and $y = 2a\sin\theta\,\tan\theta\cdot\sin\theta = 2a\sin^2\theta\,\tan\theta$.

(b)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

55. (a) *Red particle*: $x = t + 5$, $y = t^2 + 4t + 6$

Blue particle: $x = 2t + 1$, $y = 2t + 6$

Substituting $x = 1$ and $y = 6$ into the parametric equations for the red particle gives $1 = t + 5$ and $6 = t^2 + 4t + 6$, which are both satisfied when $t = -4$. Making the same substitution for the blue particle gives $1 = 2t + 1$ and $6 = 2t + 6$, which are both satisfied when $t = 0$. Repeating the process for $x = 6$ and $y = 11$, the red particle's equations become $6 = t + 5$ and $11 = t^2 + 4t + 6$, which are both satisfied when $t = 1$. Similarly, the blue particle's equations become $6 = 2t + 1$ and $11 = 2t + 6$, which are both satisfied when $t = 2.5$. Thus, $(1, 6)$ and $(6, 11)$ are both intersection points, but they are not collision points, since the particles reach each of these points at different times.

(b) *Blue particle*: $x = 2t + 1 \quad \Rightarrow \quad t = \frac{1}{2}(x - 1)$.

Substituting into the equation for y gives $y = 2t + 6 = 2\left[\frac{1}{2}(x - 1)\right] + 6 = x + 5$.

Green particle: $x = 2t + 4 \quad \Rightarrow \quad t = \frac{1}{2}(x - 4)$.

Substituting into the equation for y gives $y = 2t + 9 = 2\left[\frac{1}{2}(x - 4)\right] + 9 = x + 5$.

Thus, the green and blue particles both move along the line $y = x + 5$.

Now, the red and green particles will collide if there is a time t when both particles are at the same point. Equating the x parametric equations, we find $t + 5 = 2t + 4$, which is satisfied when $t = 1$, and gives $x = 1 + 5 = 6$. Substituting $t = 1$ into the red and green particles' y equations gives $y = (1)^2 + 4(1) + 6 = 11$ and $y = 2(1) + 9 = 11$, respectively. Thus, the red and green particles collide at the point $(6, 11)$ when $t = 1$.

56. (a) $x = 3\sin t,\ y = 2\cos t,\ 0 \le t \le 2\pi$;

$x = -3 + \cos t,\ y = 1 + \sin t,\ 0 \le t \le 2\pi$

There are 2 points of intersection:

$(-3, 0)$ and approximately $(-2.1, 1.4)$.

(b) A collision point occurs when $x_1 = x_2$ and $y_1 = y_2$ for the same t. So solve the equations:

$$3\sin t = -3 + \cos t \quad \textbf{(1)}$$

$$2\cos t = 1 + \sin t \quad \textbf{(2)}$$

From **(2)**, $\sin t = 2\cos t - 1$. Substituting into **(1)**, we get $3(2\cos t - 1) = -3 + \cos t \quad \Rightarrow \quad 5\cos t = 0 \quad (\star) \quad \Rightarrow$ $\cos t = 0 \quad \Rightarrow \quad t = \frac{\pi}{2}$ or $\frac{3\pi}{2}$. We check that $t = \frac{3\pi}{2}$ satisfies **(1)** and **(2)** but $t = \frac{\pi}{2}$ does not. So the only collision point occurs when $t = \frac{3\pi}{2}$, and this gives the point $(-3, 0)$. [We could check our work by graphing x_1 and x_2 together as functions of t and, on another plot, y_1 and y_2 as functions of t. If we do so, we see that the only value of t for which *both* pairs of graphs intersect is $t = \frac{3\pi}{2}$.]

(c) The circle is centered at $(3, 1)$ instead of $(-3, 1)$. There are still 2 intersection points: $(3, 0)$ and $(2.1, 1.4)$, but there are no collision points, since $(\star)$ in part (b) becomes $5\cos t = 6 \quad \Rightarrow \quad \cos t = \frac{6}{5} > 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. (a) $x = 1 - t^2$, $y = t - t^3$. The curve intersects itself if there are two distinct times $t = a$ and $t = b$ (with $a < b$) such that $x(a) = x(b)$ and $y(a) = y(b)$. The equation $x(a) = x(b)$ gives $1 - a^2 = 1 - b^2$ so that $a^2 = b^2$. Since $a \neq b$ by assumption, we must have $a = -b$. Substituting into the equation for y gives $y(-b) = y(b) \quad \Rightarrow$ $-b - (-b)^3 = b - b^3 \quad \Rightarrow \quad 2b^3 - 2b = 0 \quad \Rightarrow \quad 2b(b-1)(b+1) = 0 \quad \Rightarrow \quad b = -1, 0, 1$. Since $a < b$, the only valid solution is $b = 1$, which corresponds to $a = -1$ and results in the coordinates $x = 0$ and $y = 0$. Thus, the curve intersects itself at $(0, 0)$ when $t = -1$ and $t = 1$.

(b) $x = 2t - t^3$, $y = t - t^2$. Similar to part (a), we try to find the times $t = a$ and $t = b$ with $a < b$ such that $x(a) = x(b)$ and $y(a) = y(b)$. The equation $y(a) = y(b)$ gives $a - a^2 = b - b^2 \quad \Rightarrow \quad 0 = a^2 - a + (b - b^2)$. Using the quadratic formula to solve for a, we get

$a = \dfrac{1 \pm \sqrt{1 - 4(b - b^2)}}{2} = \dfrac{1 \pm \sqrt{4b^2 - 4b + 1}}{2} = \dfrac{1 \pm \sqrt{(2b-1)^2}}{2} = \dfrac{1 \pm (2b-1)}{2} \quad \Rightarrow \quad a = b$ or $a = 1 - b$. Since $a < b$ by assumption, we reject the first solution and substitute $a = 1 - b$ into $x(a) = x(b) \quad \Rightarrow \quad x(1-b) = x(b) \quad \Rightarrow$ $2(1-b) - (1-b)^3 = 2b - b^3$. Expanding and simplifying gives $2b^3 - 3b^2 - b + 1 = 0$. By graphing the equation, we see that $b = \frac{1}{2}$ is a zero, so $2b - 1$ is a factor, and by long division $b^2 - b - 1$ is another factor. Hence, the solutions are $b = \frac{1}{2}$ and $b = \frac{1}{2} \pm \frac{1}{2}\sqrt{5}$ (found using the quadratic formula). Since $a = 1 - b$ and we require $a < b$, the only valid solution is $b = \frac{1}{2} + \frac{1}{2}\sqrt{5}$, which corresponds to $a = \frac{1}{2} - \frac{1}{2}\sqrt{5}$ and results in the coordinates $x = 2\left(\frac{1}{2} - \frac{1}{2}\sqrt{5}\right) - \left(\frac{1}{2} - \frac{1}{2}\sqrt{5}\right)^3 = -1$ and $y = \frac{1}{2} - \frac{1}{2}\sqrt{5} - \left(\frac{1}{2} - \frac{1}{2}\sqrt{5}\right)^2 = -1$. Thus, the curve intersects itself at $(-1, -1)$ when $t = \frac{1}{2} - \frac{1}{2}\sqrt{5}$ and $t = \frac{1}{2} + \frac{1}{2}\sqrt{5}$.

58. (a) If $\alpha = 30°$ and $v_0 = 500$ m/s, then the equations become $x = (500 \cos 30°)t = 250\sqrt{3}t$ and $y = (500 \sin 30°)t - \frac{1}{2}(9.8)t^2 = 250t - 4.9t^2$. $y = 0$ when $t = 0$ (when the gun is fired) and again when $t = \frac{250}{4.9} \approx 51$ s. Then $x = \left(250\sqrt{3}\right)\left(\frac{250}{4.9}\right) \approx 22{,}092$ m, so the bullet hits the ground about 22 km from the gun. The formula for y is quadratic in t. To find the maximum y-value, we will complete the square:

$$y = -4.9\left(t^2 - \tfrac{250}{4.9}t\right) = -4.9\left[t^2 - \tfrac{250}{4.9}t + \left(\tfrac{125}{4.9}\right)^2\right] + \tfrac{125^2}{4.9} = -4.9\left(t - \tfrac{125}{4.9}\right)^2 + \tfrac{125^2}{4.9} \leq \tfrac{125^2}{4.9}$$

with equality when $t = \frac{125}{4.9}$ s, so the maximum height attained is $\frac{125^2}{4.9} \approx 3189$ m.

(b)

As α ($0° < \alpha < 90°$) increases up to $45°$, the projectile attains a greater height and a greater range. As α increases past $45°$, the projectile attains a greater height, but its range decreases.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $x = (v_0 \cos\alpha)t \quad \Rightarrow \quad t = \dfrac{x}{v_0 \cos\alpha}$.

$$y = (v_0 \sin\alpha)t - \tfrac{1}{2}gt^2 \quad \Rightarrow \quad y = (v_0 \sin\alpha)\,\frac{x}{v_0 \cos\alpha} - \frac{g}{2}\left(\frac{x}{v_0 \cos\alpha}\right)^2 = (\tan\alpha)x - \left(\frac{g}{2v_0^2 \cos^2\alpha}\right)x^2,$$

which is the equation of a parabola (quadratic in x).

59. $x = t^2, y = t^3 - ct$. We use a graphing device to produce the graphs for various values of c with $-\pi \le t \le \pi$. Note that all the members of the family are symmetric about the x-axis. For $c < 0$, the graph does not cross itself, but for $c = 0$ it has a cusp at $(0, 0)$ and for $c > 0$ the graph crosses itself at $x = c$, so the loop grows larger as c increases.

60. $x = 2ct - 4t^3, y = -ct^2 + 3t^4$. We use a graphing device to produce the graphs for various values of c with $-\pi \le t \le \pi$. Note that all the members of the family are symmetric about the y-axis. When $c < 0$, the graph resembles that of a polynomial of even degree, but when $c = 0$ there is a corner at the origin, and when $c > 0$, the graph crosses itself at the origin, and has two cusps below the x-axis. The size of the "swallowtail" increases as c increases.

61. $x = t + a\cos t, y = t + a\sin t, a > 0$. From the first figure, we see that curves roughly follow the line $y = x$, and they start having loops when a is between 1.4 and 1.6. The loops increase in size as a increases.

While not required, the following is a solution to determine the *exact* values for which the curve has a loop, that is, we seek the values of a for which there exist parameter values t and u such that $t < u$ and $(t + a\cos t, t + a\sin t) = (u + a\cos u, u + a\sin u)$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

In the diagram at the left, T denotes the point (t, t), U the point (u, u), and P the point $(t + a\cos t, t + a\sin t) = (u + a\cos u, u + a\sin u)$. Since $\overline{PT} = \overline{PU} = a$, the triangle PTU is isosceles. Therefore its base angles, $\alpha = \angle PTU$ and $\beta = \angle PUT$ are equal. Since $\alpha = t - \frac{\pi}{4}$ and $\beta = 2\pi - \frac{3\pi}{4} - u = \frac{5\pi}{4} - u$, the relation $\alpha = \beta$ implies that $u + t = \frac{3\pi}{2}$ **(1)**.

Since $\overline{TU} = \text{distance}((t,t),(u,u)) = \sqrt{2(u-t)^2} = \sqrt{2}\,(u-t)$, we see that $\cos\alpha = \dfrac{\frac{1}{2}\overline{TU}}{\overline{PT}} = \dfrac{(u-t)/\sqrt{2}}{a}$, so $u - t = \sqrt{2}\,a\cos\alpha$, that is, $u - t = \sqrt{2}\,a\cos\left(t - \frac{\pi}{4}\right)$ **(2)**. Now $\cos\left(t - \frac{\pi}{4}\right) = \sin\left[\frac{\pi}{2} - \left(t - \frac{\pi}{4}\right)\right] = \sin\left(\frac{3\pi}{4} - t\right)$, so we can rewrite **(2)** as $u - t = \sqrt{2}\,a\sin\left(\frac{3\pi}{4} - t\right)$ **(2′)**. Subtracting **(2′)** from **(1)** and dividing by 2, we obtain $t = \frac{3\pi}{4} - \frac{\sqrt{2}}{2}a\sin\left(\frac{3\pi}{4} - t\right)$, or $\frac{3\pi}{4} - t = \frac{a}{\sqrt{2}}\sin\left(\frac{3\pi}{4} - t\right)$ **(3)**. Since $a > 0$ and $t < u$, it follows from **(2′)** that $\sin\left(\frac{3\pi}{4} - t\right) > 0$. Thus from **(3)** we see that $t < \frac{3\pi}{4}$. [We have implicitly assumed that $0 < t < \pi$ by the way we drew our diagram, but we lost no generality by doing so since replacing t by $t + 2\pi$ merely increases x and y by 2π. The curve's basic shape repeats every time we change t by 2π.] Solving for a in **(3)**, we get $a = \dfrac{\sqrt{2}\left(\frac{3\pi}{4} - t\right)}{\sin\left(\frac{3\pi}{4} - t\right)}$. Write $z = \frac{3\pi}{4} - t$. Then $a = \dfrac{\sqrt{2}\,z}{\sin z}$, where $z > 0$. Now $\sin z < z$ for $z > 0$, so $a > \sqrt{2}$. $\left[\text{As } z \to 0^+, \text{ that is, as } t \to \left(\frac{3\pi}{4}\right)^-, a \to \sqrt{2}\right]$.

62. Consider the curves $x = \sin t + \sin nt$, $y = \cos t + \cos nt$, where n is a positive integer. For $n = 1$, we get a circle of radius 2 centered at the origin. For $n > 1$, we get a curve lying on or inside that circle that traces out $n - 1$ loops as t ranges from 0 to 2π.

Note:
$$\begin{aligned} x^2 + y^2 &= (\sin t + \sin nt)^2 + (\cos t + \cos nt)^2 \\ &= \sin^2 t + 2\sin t\,\sin nt + \sin^2 nt + \cos^2 t + 2\cos t\,\cos nt + \cos^2 nt \\ &= (\sin^2 t + \cos^2 t) + (\sin^2 nt + \cos^2 nt) + 2(\cos t\,\cos nt + \sin t\,\sin nt) \\ &= 1 + 1 + 2\cos(t - nt) = 2 + 2\cos((1-n)t) \le 4 = 2^2, \end{aligned}$$

with equality for $n = 1$. This shows that each curve lies on or inside the curve for $n = 1$, which is a circle of radius 2 centered at the origin.

$n = 1$ $n = 2$ $n = 3$ $n = 5$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

63. Note that all the Lissajous figures are symmetric about the x-axis. The parameters a and b simply stretch the graph in the x- and y-directions respectively. For $a = b = n = 1$ the graph is simply a circle with radius 1. For $n = 2$ the graph crosses itself at the origin and there are loops above and below the x-axis. In general, the figures have $n - 1$ points of intersection, all of which are on the y-axis, and a total of n closed loops.

$a = b = 1$ $\qquad$ $n = 2$ $\qquad$ $n = 3$

64. $x = \cos t$, $y = \sin t - \sin ct$. If $c = 1$, then $y = 0$, and the curve is simply the line segment from $(-1, 0)$ to $(1, 0)$. The graphs are shown for $c = 2, 3, 4$ and 5.

It is easy to see that all the curves lie in the rectangle $[-1, 1]$ by $[-2, 2]$. When c is an integer, $x(t + 2\pi) = x(t)$ and $y(t + 2\pi) = y(t)$, so the curve is closed. When c is a positive integer greater than 1, the curve intersects the x-axis $c + 1$ times and has c loops (one of which degenerates to a tangency at the origin when c is an odd integer of the form $4k + 1$).

As c increases, the curve's loops become thinner, but stay in the region bounded by the semicircles $y = \pm\left(1 + \sqrt{1 - x^2}\right)$ and the line segments from $(-1, -1)$ to $(-1, 1)$ and from $(1, -1)$ to $(1, 1)$. This is true because $|y| = |\sin t - \sin ct| \le |\sin t| + |\sin ct| \le \sqrt{1 - x^2} + 1$. This curve appears to fill the entire region when c is very large, as shown in the figure for $c = 1000$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

When c is a fraction, we get a variety of shapes with multiple loops, but always within the same region. For some fractional values, such as $c = 2.359$, the curve again appears to fill the region.

DISCOVERY PROJECT Running Circles Around Circles

1. The center Q of the smaller circle has coordinates $((a-b)\cos\theta, (a-b)\sin\theta)$. Arc PS on circle C has length $a\theta$ since it is equal in length to arc AS (the smaller circle rolls without slipping against the larger.)

Thus, $\angle PQS = \dfrac{a}{b}\theta$ and $\angle PQT = \dfrac{a}{b}\theta - \theta$, so P has coordinates

$$x = (a-b)\cos\theta + b\cos(\angle PQT) = (a-b)\cos\theta + b\cos\left(\frac{a-b}{b}\theta\right)$$

and $$y = (a-b)\sin\theta - b\sin(\angle PQT) = (a-b)\sin\theta - b\sin\left(\frac{a-b}{b}\theta\right).$$

2. With $b = 1$ and a a positive integer greater than 2, we obtain a hypocycloid of a cusps. Shown in the figure is the graph for $a = 4$. Let $a = 4$ and $b = 1$. Using the sum identities to expand $\cos 3\theta$ and $\sin 3\theta$, we obtain

$$x = 3\cos\theta + \cos 3\theta = 3\cos\theta + \left(4\cos^3\theta - 3\cos\theta\right) = 4\cos^3\theta$$

and $$y = 3\sin\theta - \sin 3\theta = 3\sin\theta - \left(3\sin\theta - 4\sin^3\theta\right) = 4\sin^3\theta.$$

3. The graphs at the right are obtained with $b = 1$ and $a = \frac{1}{2}, \frac{1}{3}, \frac{1}{4}$, and $\frac{1}{10}$ with $-2\pi \le \theta \le 2\pi$. We conclude that as the denominator d increases, the graph gets smaller, but maintains the basic shape shown.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Letting $d = 2$ and $n = 3$, 5, and 7 with $-2\pi \le \theta \le 2\pi$ gives us the following:

So if d is held constant and n varies, we get a graph with n cusps (assuming n/d is in lowest form). When $n = d + 1$, we obtain a hypocycloid of n cusps. As n increases, we must expand the range of θ in order to get a closed curve. The following graphs have $a = \frac{3}{2}$, $\frac{5}{4}$, and $\frac{11}{10}$.

4. If $b = 1$, the equations for the hypocycloid are

$$x = (a-1)\cos\theta + \cos((a-1)\theta) \qquad y = (a-1)\sin\theta - \sin((a-1)\theta)$$

which is a hypocycloid of a cusps (from Problem 2). In general, if $a > 1$, we get a figure with cusps on the "outside ring" and if $a < 1$, the cusps are on the "inside ring". In any case, as the values of θ get larger, we get a figure that looks more and more like a washer. If we were to graph the hypocycloid for all values of θ, every point on the washer would eventually be arbitrarily close to a point on the curve.

$a = \sqrt{2}, \quad -10\pi \le \theta \le 10\pi$

$a = e - 2, \quad 0 \le \theta \le 446$

5. The center Q of the smaller circle has coordinates $((a+b)\cos\theta, (a+b)\sin\theta)$.

Arc PS has length $a\theta$ (as in Problem 1), so that $\angle PQS = \dfrac{a\theta}{b}$, $\angle PQR = \pi - \dfrac{a\theta}{b}$,

and $\angle PQT = \pi - \dfrac{a\theta}{b} - \theta = \pi - \left(\dfrac{a+b}{b}\right)\theta$ since $\angle RQT = \theta$.

Thus, the coordinates of P are

$$x = (a+b)\cos\theta + b\cos\left(\pi - \frac{a+b}{b}\theta\right) = (a+b)\cos\theta - b\cos\left(\frac{a+b}{b}\theta\right)$$

and $$y = (a+b)\sin\theta - b\sin\left(\pi - \frac{a+b}{b}\theta\right) = (a+b)\sin\theta - b\sin\left(\frac{a+b}{b}\theta\right).$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. Let $b = 1$ and the equations become

$$x = (a+1)\cos\theta - \cos((a+1)\theta) \qquad y = (a+1)\sin\theta - \sin((a+1)\theta)$$

If $a = 1$, we have a cardioid. If a is a positive integer greater than 1, we get the graph of an "a-leafed clover", with cusps that are a units from the origin. (Some of the pairs of figures are not to scale.)

$a = 3$, $-2\pi \le \theta \le 2\pi$

$a = 10$, $-2\pi \le \theta \le 2\pi$

If $a = n/d$ with $n = 1$, we obtain a figure that does not increase in size and requires $-d\pi \le \theta \le d\pi$ to be a closed curve traced exactly once.

$a = \frac{1}{4}$, $-4\pi \le \theta \le 4\pi$

$a = \frac{1}{7}$, $-7\pi \le \theta \le 7\pi$

Next, we keep d constant and let n vary. As n increases, so does the size of the figure. There is an n-pointed star in the middle.

$a = \frac{2}{5}$, $-5\pi \le \theta \le 5\pi$

$a = \frac{7}{5}$, $-5\pi \le \theta \le 5\pi$

Now if $n = d + 1$ we obtain figures similar to the previous ones, but the size of the figure does not increase.

$a = \frac{4}{3}$, $-3\pi \le \theta \le 3\pi$

$a = \frac{7}{6}$, $-6\pi \le \theta \le 6\pi$

If a is irrational, we get washers that increase in size as a increases.

$a = \sqrt{2}$, $0 \le \theta \le 200$

$a = e - 2$, $0 \le \theta \le 446$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10.2 Calculus with Parametric Curves

1. $x = 2t^3 + 3t,\ y = 4t - 5t^2 \quad \Rightarrow \quad \dfrac{dx}{dt} = 6t^2 + 3,\ \dfrac{dy}{dt} = 4 - 10t$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{4 - 10t}{6t^2 + 3}$.

2. $x = t - \ln t,\ y = t^2 - t^{-2} \quad \Rightarrow \quad \dfrac{dx}{dt} = 1 - t^{-1},\ \dfrac{dy}{dt} = 2t + 2t^{-3}$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t + 2t^{-3}}{1 - t^{-1}} \cdot \dfrac{t^3}{t^3} = \dfrac{2t^4 + 2}{t^3 - t^2}$.

3. $x = te^t,\ y = t + \sin t \quad \Rightarrow \quad \dfrac{dx}{dt} = te^t + e^t = e^t(t + 1),\ \dfrac{dy}{dt} = 1 + \cos t$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{1 + \cos t}{e^t(t + 1)}$.

4. $x = t + \sin(t^2 + 2),\ y = \tan(t^2 + 2) \quad \Rightarrow \quad \dfrac{dx}{dt} = 1 + 2t\cos(t^2 + 2),\ \dfrac{dy}{dt} = 2t\sec^2(t^2 + 2)$, and

$\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t\sec^2(t^2 + 2)}{1 + 2t\cos(t^2 + 2)}$.

5. $x = t^2 + 2t,\ y = 2^t - 2t;\ (15, 2)$. $\dfrac{dy}{dt} = 2^t \ln 2 - 2,\ \dfrac{dx}{dt} = 2t + 2$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2^t \ln 2 - 2}{2t + 2}$.

At $(15, 2)$, $x = t^2 + 2t = 15 \quad \Rightarrow \quad t^2 + 2t - 15 = 0 \quad \Rightarrow \quad (t + 5)(t - 3) = 0 \quad \Rightarrow \quad t = -5$ or $t = 3$. Only $t = 3$ gives $y = 2$. With $t = 3$, $\dfrac{dy}{dx} = \dfrac{2^3 \ln 2 - 2}{2(3) + 2} = \dfrac{4\ln 2 - 1}{4} = \ln 2 - \dfrac{1}{4} \approx 0.44$.

6. $x = t + \cos \pi t,\ y = -t + \sin \pi t;\ (3, -2)$. $\dfrac{dy}{dt} = -1 + \pi \cos \pi t,\ \dfrac{dx}{dt} = 1 - \pi \sin \pi t$, and

$\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{-1 + \pi \cos \pi t}{1 - \pi \sin \pi t}$. When $x = 3$, we have $t + \cos \pi t = 3 \quad \Rightarrow \quad \cos^2 \pi t = (3 - t)^2$ **(1)**. When $y = -2$, we have $-t + \sin \pi t = -2 \quad \Rightarrow \quad \sin^2 \pi t = (t - 2)^2$ **(2)**. Adding **(1)** and **(2)** gives $\sin^2 \pi t + \cos^2 \pi t = (t - 2)^2 + (3 - t)^2 \quad \Rightarrow$ $1 = t^2 - 4t + 4 + 9 - 6t + t^2 \quad \Rightarrow \quad 0 = 2t^2 - 10t + 12 \quad \Rightarrow \quad 0 = 2(t - 2)(t - 3) \quad \Rightarrow \quad t = 2$ or $t = 3$. Only $t = 2$ gives $y = -2$. With $t = 2$, $\dfrac{dy}{dx} = \dfrac{-1 + \pi \cos 2\pi}{1 - \pi \sin 2\pi} = \dfrac{-1 + \pi}{1 - 0} = \pi - 1 \approx 2.14$.

7. $x = t^3 + 1,\ y = t^4 + t;\quad t = -1$. $\dfrac{dy}{dt} = 4t^3 + 1,\ \dfrac{dx}{dt} = 3t^2$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{4t^3 + 1}{3t^2}$. When $t = -1$, $(x, y) = (0, 0)$ and $dy/dx = -3/3 = -1$, so an equation of the tangent to the curve at the point corresponding to $t = -1$ is $y - 0 = -1(x - 0)$, or $y = -x$.

8. $x = \sqrt{t},\ y = t^2 - 2t;\quad t = 4$. $\dfrac{dy}{dt} = 2t - 2,\ \dfrac{dx}{dt} = \dfrac{1}{2\sqrt{t}}$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = (2t - 2)2\sqrt{t} = 4(t - 1)\sqrt{t}$. When $t = 4$, $(x, y) = (2, 8)$ and $dy/dx = 4(3)(2) = 24$, so an equation of the tangent to the curve at the point corresponding to $t = 4$ is $y - 8 = 24(x - 2)$, or $y = 24x - 40$.

9. $x = \sin 2t + \cos t,\ y = \cos 2t - \sin t;\ t = \pi$. $\dfrac{dy}{dt} = -2\sin 2t - \cos t,\ \dfrac{dx}{dt} = 2\cos 2t - \sin t$, and

$\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{-2\sin 2t - \cos t}{2\cos 2t - \sin t}$. When $t = \pi$, $(x, y) = (-1, 1)$, and $\dfrac{dy}{dx} = \dfrac{1}{2}$, so an equation of the tangent to the curve at the point corresponding to $t = \pi$ is $y - 1 = \frac{1}{2}[x - (-1)]$, or $y = \frac{1}{2}x + \frac{3}{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $x = e^t \sin \pi t,\ y = e^{2t};\quad t = 0.\ \dfrac{dy}{dt} = 2e^{2t},\ \dfrac{dx}{dt} = e^t(\pi \cos \pi t) + (\sin \pi t)e^t = e^t(\pi \cos \pi t + \sin \pi t)$, and

$\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2e^{2t}}{e^t(\pi \cos \pi t + \sin \pi t)} = \dfrac{2e^t}{\pi \cos \pi t + \sin \pi t}$. When $t = 0$, $(x, y) = (0, 1)$ and $\dfrac{dy}{dx} = \dfrac{2}{\pi}$, so an equation of the tangent to the curve at the point corresponding to $t = 0$ is $y - 1 = \frac{2}{\pi}(x - 0)$, or $y = \frac{2}{\pi}x + 1$.

11. (a) $x = \sin t,\ y = \cos^2 t;\ \left(\frac{1}{2}, \frac{3}{4}\right).\quad \dfrac{dy}{dt} = 2\cos t(-\sin t),\ \dfrac{dx}{dt} = \cos t$, and $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{-2\sin t \cos t}{\cos t} = -2\sin t$.

At $\left(\frac{1}{2}, \frac{3}{4}\right)$, $x = \sin t = \frac{1}{2} \quad\Rightarrow\quad t = \frac{\pi}{6}$, so $dy/dx = -2\sin\frac{\pi}{6} = -2\left(\frac{1}{2}\right) = -1$, and an equation of the tangent is $y - \frac{3}{4} = -1\left(x - \frac{1}{2}\right)$, or $y = -x + \frac{5}{4}$.

(b) $x = \sin t \quad\Rightarrow\quad x^2 = \sin^2 t = 1 - \cos^2 t = 1 - y$, so $y = 1 - x^2$, and $y' = -2x$. At $\left(\frac{1}{2}, \frac{3}{4}\right)$, $y' = -2 \cdot \frac{1}{2} = -1$, so an equation of the tangent is $y - \frac{3}{4} = -1\left(x - \frac{1}{2}\right)$, or $y = -x + \frac{5}{4}$.

12. (a) $x = \sqrt{t+4},\ y = 1/(t+4);\ \left(2, \frac{1}{4}\right).\quad \dfrac{dy}{dt} = -\dfrac{1}{(t+4)^2},\ \dfrac{dx}{dt} = \dfrac{1}{2\sqrt{t+4}}$, and

$\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{-1/(t+4)^2}{1/\left(2\sqrt{t+4}\right)} = -2(t+4)^{-3/2}$. At $\left(2, \frac{1}{4}\right)$, $x = \sqrt{t+4} = 2 \quad\Rightarrow\quad t + 4 = 4 \quad\Rightarrow\quad t = 0$ and $dy/dx = -2(4)^{-3/2} = -\frac{1}{4}$, so an equation of the tangent is $y - \frac{1}{4} = -\frac{1}{4}(x - 2)$, or $y = -\frac{1}{4}x + \frac{3}{4}$.

(b) $x = \sqrt{t+4} \quad\Rightarrow\quad x^2 = t + 4 \quad\Rightarrow\quad t = x^2 - 4$, so $y = \dfrac{1}{t+4} = \dfrac{1}{x^2 - 4 + 4} = \dfrac{1}{x^2}$, and $y' = -\dfrac{2}{x^3}$. At $\left(2, \frac{1}{4}\right)$, $y' = -2/2^3 = -1/4$, so an equation of the tangent is $y - \frac{1}{4} = -\frac{1}{4}(x - 2)$, or $y = -\frac{1}{4}x + \frac{3}{4}$.

13. $x = t^2 - t,\ y = t^2 + t + 1;\ (0, 3).\quad \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t+1}{2t-1}$. To find the value of t corresponding to the point $(0, 3)$, solve $x = 0 \quad\Rightarrow$ $t^2 - t = 0 \quad\Rightarrow\quad t(t-1) = 0 \quad\Rightarrow\quad t = 0$ or $t = 1$. Only $t = 1$ gives $y = 3$. With $t = 1$, $dy/dx = 3$, and an equation of the tangent is $y - 3 = 3(x - 0)$, or $y = 3x + 3$.

14. $x = \sin \pi t,\ y = t^2 + t;\ (0, 2).\quad \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t+1}{\pi \cos \pi t}$. To find the value of t corresponding to the point $(0, 2)$, solve $y = 2 \quad\Rightarrow$ $t^2 + t - 2 = 0 \quad\Rightarrow\quad (t+2)(t-1) = 0 \quad\Rightarrow\quad t = -2$ or $t = 1$. Either value gives $dy/dx = -3/\pi$, so an equation of the tangent is $y - 2 = -\frac{3}{\pi}(x - 0)$, or $y = -\frac{3}{\pi}x + 2$.

15. $x = t^2 + 1,\ y = t^2 + t \quad\Rightarrow\quad \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t+1}{2t} = 1 + \dfrac{1}{2t} \quad\Rightarrow\quad \dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{-1/(2t^2)}{2t} = -\dfrac{1}{4t^3}$.

The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $t < 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. $x = t^3 + 1,\ y = t^2 - t \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t-1}{3t^2} = \dfrac{2}{3t} - \dfrac{1}{3t^2} \quad \Rightarrow$

$\dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{-\dfrac{2}{3t^2} + \dfrac{2}{3t^3}}{3t^2} = \dfrac{\dfrac{2-2t}{3t^3}}{3t^2} = \dfrac{2(1-t)}{9t^5}$. The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $0 < t < 1$.

17. $x = e^t,\ y = te^{-t} \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{-te^{-t} + e^{-t}}{e^t} = \dfrac{e^{-t}(1-t)}{e^t} = e^{-2t}(1-t) \quad \Rightarrow$

$\dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{e^{-2t}(-1) + (1-t)(-2e^{-2t})}{e^t} = \dfrac{e^{-2t}(-1-2+2t)}{e^t} = e^{-3t}(2t-3)$. The curve is CU when

$\dfrac{d^2y}{dx^2} > 0$, that is, when $t > \frac{3}{2}$.

18. $x = t^2 + 1,\ y = e^t - 1 \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{e^t}{2t} \quad \Rightarrow \quad \dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{\dfrac{2te^t - e^t \cdot 2}{(2t)^2}}{2t} = \dfrac{2e^t(t-1)}{(2t)^3} = \dfrac{e^t(t-1)}{4t^3}$.

The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $t < 0$ or $t > 1$.

19. $x = t - \ln t,\ y = t + \ln t$ [note that $t > 0$] $\Rightarrow \quad \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{1 + 1/t}{1 - 1/t} = \dfrac{t+1}{t-1} \quad \Rightarrow$

$\dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{\dfrac{(t-1)(1) - (t+1)(1)}{(t-1)^2}}{(t-1)/t} = \dfrac{-2t}{(t-1)^3}$. The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $0 < t < 1$.

20. $x = \cos t,\ y = \sin 2t,\ 0 < t < \pi \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2\cos 2t}{-\sin t} \quad \Rightarrow$

$$\frac{d^2y}{dx^2} = \frac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \frac{\dfrac{(-\sin t)(-4\sin 2t) - (2\cos 2t)(-\cos t)}{(-\sin t)^2}}{-\sin t} = \frac{(\sin t)(8\sin t\,\cos t) + [2(1 - 2\sin^2 t)](\cos t)}{(-\sin t)\sin^2 t}$$

$$= \frac{(\cos t)(8\sin^2 t + 2 - 4\sin^2 t)}{(-\sin t)\sin^2 t} = -\frac{\cos t}{\sin t} \cdot \frac{4\sin^2 t + 2}{\sin^2 t} \quad [\,(-\cot t) \cdot \text{positive expression}]$$

The curve is CU when $\dfrac{d^2y}{dx^2} > 0$, that is, when $-\cot t > 0 \quad \Leftrightarrow \quad \cot t < 0 \quad \Leftrightarrow \quad \frac{\pi}{2} < t < \pi$.

21. $x = t^3 - 3t,\ y = t^2 - 3.\quad \dfrac{dy}{dt} = 2t$, so $\dfrac{dy}{dt} = 0 \quad \Leftrightarrow \quad t = 0 \quad \Leftrightarrow$

$(x, y) = (0, -3).\quad \dfrac{dx}{dt} = 3t^2 - 3 = 3(t+1)(t-1)$, so $\dfrac{dx}{dt} = 0 \quad \Leftrightarrow$

$t = -1$ or $1 \quad \Leftrightarrow \quad (x, y) = (2, -2)$ or $(-2, -2)$. The curve has a horizontal tangent at $(0, -3)$ and vertical tangents at $(2, -2)$ and $(-2, -2)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $x = t^3 - 3t,\ y = t^3 - 3t^2.$ $\dfrac{dy}{dt} = 3t^2 - 6t = 3t(t-2)$, so $\dfrac{dy}{dt} = 0 \quad \Leftrightarrow$ $t = 0$ or $2 \quad \Leftrightarrow \quad (x, y) = (0, 0)$ or $(2, -4)$. $\dfrac{dx}{dt} = 3t^2 - 3 = 3(t+1)(t-1)$, so $\dfrac{dx}{dt} = 0 \quad \Leftrightarrow \quad t = -1$ or $1 \quad \Leftrightarrow \quad (x, y) = (2, -4)$ or $(-2, -2)$. The curve has horizontal tangents at $(0, 0)$ and $(2, -4)$, and vertical tangents at $(2, -4)$ and $(-2, -2)$.

23. $x = \cos\theta,\ y = \cos 3\theta$. The whole curve is traced out for $0 \le \theta \le \pi$.

$\dfrac{dy}{d\theta} = -3\sin 3\theta$, so $\dfrac{dy}{d\theta} = 0 \quad \Leftrightarrow \quad \sin 3\theta = 0 \quad \Leftrightarrow \quad 3\theta = 0, \pi, 2\pi$, or $3\pi \quad \Leftrightarrow$ $\theta = 0, \frac{\pi}{3}, \frac{2\pi}{3}$, or $\pi \quad \Leftrightarrow \quad (x, y) = (1, 1), \left(\frac{1}{2}, -1\right), \left(-\frac{1}{2}, 1\right)$, or $(-1, -1)$.

$\dfrac{dx}{d\theta} = -\sin\theta$, so $\dfrac{dx}{d\theta} = 0 \quad \Leftrightarrow \quad \sin\theta = 0 \quad \Leftrightarrow \quad \theta = 0$ or $\pi \quad \Leftrightarrow$ $(x, y) = (1, 1)$ or $(-1, -1)$. Both $\dfrac{dy}{d\theta}$ and $\dfrac{dx}{d\theta}$ equal 0 when $\theta = 0$ and π.

To find the slope when $\theta = 0$, we find $\displaystyle\lim_{\theta\to 0} \frac{dy}{dx} = \lim_{\theta\to 0} \frac{-3\sin 3\theta}{-\sin\theta} \overset{\text{H}}{=} \lim_{\theta\to 0} \frac{-9\cos 3\theta}{-\cos\theta} = 9$, which is the same slope when $\theta = \pi$. Thus, the curve has horizontal tangents at $\left(\frac{1}{2}, -1\right)$ and $\left(-\frac{1}{2}, 1\right)$, and there are no vertical tangents.

24. $x = e^{\sin\theta}$, $y = e^{\cos\theta}$. The whole curve is traced out for $0 \le \theta < 2\pi$.

$\dfrac{dy}{d\theta} = -\sin\theta\, e^{\cos\theta}$, so $\dfrac{dy}{d\theta} = \theta \quad \Leftrightarrow \quad \sin\theta = 0 \quad \Leftrightarrow \quad \theta = 0$ or $\pi \quad \Leftrightarrow$ $(x, y) = (1, e)$ or $(1, 1/e)$. $\dfrac{dx}{d\theta} = \cos\theta\, e^{\sin\theta}$, so $\dfrac{dx}{d\theta} = 0 \quad \Leftrightarrow \quad \cos\theta = 0 \quad \Leftrightarrow$ $\theta = \frac{\pi}{2}$ or $\frac{3\pi}{2} \quad \Leftrightarrow \quad (x, y) = (e, 1)$ or $(1/e, 1)$. The curve has horizontal tangents at $(1, e)$ and $(1, 1/e)$, and vertical tangents at $(e, 1)$ and $(1/e, 1)$.

25. From the graph, it appears that the rightmost point on the curve $x = t - t^6$, $y = e^t$ is about $(0.6, 2)$. To find the exact coordinates, we find the value of t for which the graph has a vertical tangent, that is, $0 = dx/dt = 1 - 6t^5 \quad \Leftrightarrow \quad t = 1/\sqrt[5]{6}$. Hence, the rightmost point is

$\left(1/\sqrt[5]{6} - 1/\left(6\sqrt[5]{6}\right), e^{1/\sqrt[5]{6}}\right) = \left(5 \cdot 6^{-6/5}, e^{6^{-1/5}}\right) \approx (0.58, 2.01)$.

26. From the graph, it appears that the lowest point and the leftmost point on the curve $x = t^4 - 2t$, $y = t + t^4$ are $(1.5, -0.5)$ and $(-1.2, 1.2)$, respectively. To find the exact coordinates, we solve $dy/dt = 0$ (horizontal tangents) and $dx/dt = 0$ (vertical tangents).

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\dfrac{dy}{dt} = 0 \quad \Leftrightarrow \quad 1 + 4t^3 = 0 \quad \Leftrightarrow \quad t = -\dfrac{1}{\sqrt[3]{4}}$, so the lowest point is

$$\left(\frac{1}{\sqrt[3]{256}} + \frac{2}{\sqrt[3]{4}}, -\frac{1}{\sqrt[3]{4}} + \frac{1}{\sqrt[3]{256}}\right) = \left(\frac{9}{\sqrt[3]{256}}, -\frac{3}{\sqrt[3]{256}}\right) \approx (1.42, -0.47)$$

$\dfrac{dx}{dt} = 0 \quad \Leftrightarrow \quad 4t^3 - 2 = 0 \quad \Leftrightarrow \quad t = \dfrac{1}{\sqrt[3]{2}}$, so the leftmost point is

$$\left(\frac{1}{\sqrt[3]{16}} - \frac{2}{\sqrt[3]{2}}, \frac{1}{\sqrt[3]{2}} + \frac{1}{\sqrt[3]{16}}\right) = \left(-\frac{3}{\sqrt[3]{16}}, \frac{3}{\sqrt[3]{16}}\right) \approx (-1.19, 1.19)$$

27. We graph the curve $x = t^4 - 2t^3 - 2t^2$, $y = t^3 - t$ in the viewing rectangle $[-2, 1.1]$ by $[-0.5, 0.5]$. This rectangle corresponds approximately to $t \in [-1, 0.8]$.

We estimate that the curve has horizontal tangents at about $(-1, -0.4)$ and $(-0.17, 0.39)$ and vertical tangents at about $(0, 0)$ and $(-0.19, 0.37)$. We calculate $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{3t^2 - 1}{4t^3 - 6t^2 - 4t}$. The horizontal tangents occur when $dy/dt = 3t^2 - 1 = 0 \quad \Leftrightarrow \quad t = \pm\frac{1}{\sqrt{3}}$, so both horizontal tangents are shown in our graph. The vertical tangents occur when $dx/dt = 2t(2t^2 - 3t - 2) = 0 \quad \Leftrightarrow \quad 2t(2t + 1)(t - 2) = 0 \quad \Leftrightarrow \quad t = 0, -\frac{1}{2}$ or 2. It seems that we have missed one vertical tangent, and indeed if we plot the curve on the t-interval $[-1.2, 2.2]$ we see that there is another vertical tangent at $(-8, 6)$.

28. We graph the curve $x = t^4 + 4t^3 - 8t^2$, $y = 2t^2 - t$ in the viewing rectangle $[-3.7, 0.2]$ by $[-0.2, 1.4]$. It appears that there is a horizontal tangent at about $(-0.4, -0.1)$, and vertical tangents at about $(-3, 1)$ and $(0, 0)$.

We calculate $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{4t - 1}{4t^3 + 12t^2 - 16t}$, so there is a horizontal tangent where $dy/dt = 4t - 1 = 0 \quad \Leftrightarrow \quad t = \frac{1}{4}$. This point (the lowest point) is shown in the first graph. There are vertical tangents where $dx/dt = 4t^3 + 12t^2 - 16t = 0 \quad \Leftrightarrow$ $4t(t^2 + 3t - 4) = 0 \quad \Leftrightarrow \quad 4t(t + 4)(t - 1) = 0$. We have missed one vertical tangent corresponding to $t = -4$, and if we plot the graph for $t \in [-5, 3]$, we see that the curve has another vertical tangent line at approximately $(-128, 36)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. $x = \cos t,\ y = \sin t \cos t.\quad dx/dt = -\sin t,$

$dy/dt = -\sin^2 t + \cos^2 t = \cos 2t.\ (x, y) = (0, 0) \quad \Leftrightarrow \quad \cos t = 0 \quad \Leftrightarrow \quad t$ is an odd multiple of $\frac{\pi}{2}$. When $t = \frac{\pi}{2}$, $dx/dt = -1$ and $dy/dt = -1$, so $dy/dx = 1$. When $t = \frac{3\pi}{2}$, $dx/dt = 1$ and $dy/dt = -1$. So $dy/dx = -1$. Thus, $y = x$ and $y = -x$ are both tangent to the curve at $(0, 0)$.

30. $x = -2\cos t,\ y = \sin t + \sin 2t$. From the graph, it appears that the curve crosses itself at the point $(1, 0)$. If this is true, then $x = 1 \quad \Leftrightarrow$ $-2\cos t = 1 \quad \Leftrightarrow \quad \cos t = -\frac{1}{2} \quad \Leftrightarrow \quad t = \frac{2\pi}{3}$ or $\frac{4\pi}{3}$ for $0 \le t \le 2\pi$. Substituting either value of t into y gives $y = 0$, confirming that $(1, 0)$ is the point where the curve crosses itself. $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{\cos t + 2\cos 2t}{2\sin t}$.

When $t = \dfrac{2\pi}{3}$, $\dfrac{dy}{dx} = \dfrac{-1/2 + 2(-1/2)}{2(\sqrt{3}/2)} = \dfrac{-3/2}{\sqrt{3}} = -\dfrac{\sqrt{3}}{2}$, so an equation of the tangent line is $y - 0 = -\dfrac{\sqrt{3}}{2}(x - 1)$, or $y = -\dfrac{\sqrt{3}}{2}x + \dfrac{\sqrt{3}}{2}$. Similarly, when $t = \dfrac{4\pi}{3}$, an equation of the tangent line is $y = \dfrac{\sqrt{3}}{2}x - \dfrac{\sqrt{3}}{2}$.

31. $x = r\theta - d\sin\theta,\ y = r - d\cos\theta.$

(a) $\dfrac{dx}{d\theta} = r - d\cos\theta$, $\dfrac{dy}{d\theta} = d\sin\theta$, so $\dfrac{dy}{dx} = \dfrac{d\sin\theta}{r - d\cos\theta}$.

(b) If $0 < d < r$, then $|d\cos\theta| \le d < r$, so $r - d\cos\theta \ge r - d > 0$. This shows that $dx/d\theta$ never vanishes, so the trochoid can have no vertical tangent if $d < r$.

32. $x = a\cos^3\theta,\ y = a\sin^3\theta.$

(a) $\dfrac{dx}{d\theta} = -3a\cos^2\theta\sin\theta$, $\dfrac{dy}{d\theta} = 3a\sin^2\theta\cos\theta$, so $\dfrac{dy}{dx} = -\dfrac{\sin\theta}{\cos\theta} = -\tan\theta$.

(b) The tangent is horizontal $\quad\Leftrightarrow\quad dy/dx = 0 \quad\Leftrightarrow\quad \tan\theta = 0 \quad\Leftrightarrow\quad \theta = n\pi \quad\Leftrightarrow\quad (x, y) = (\pm a, 0)$.

The tangent is vertical $\quad\Leftrightarrow\quad \cos\theta = 0 \quad\Leftrightarrow\quad \theta$ is an odd multiple of $\frac{\pi}{2} \quad\Leftrightarrow\quad (x, y) = (0, \pm a)$.

(c) $dy/dx = \pm 1 \quad\Leftrightarrow\quad \tan\theta = \pm 1 \quad\Leftrightarrow\quad \theta$ is an odd multiple of $\frac{\pi}{4} \quad\Leftrightarrow\quad (x, y) = \left(\pm\frac{\sqrt{2}}{4}a, \pm\frac{\sqrt{2}}{4}a\right)$

[All sign choices are valid.]

33. $x = 3t^2 + 1,\ y = t^3 - 1 \quad\Rightarrow\quad \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{3t^2}{6t} = \dfrac{t}{2}$. The tangent line has slope $\dfrac{1}{2}$ when $\dfrac{t}{2} = \dfrac{1}{2} \quad\Leftrightarrow\quad t = 1$, so the point is $(4, 0)$.

34. $x = 3t^2 + 1$, $y = 2t^3 + 1$, $\dfrac{dx}{dt} = 6t$, $\dfrac{dy}{dt} = 6t^2$, so $\dfrac{dy}{dx} = \dfrac{6t^2}{6t} = t$ [even where $t = 0$].

So at the point corresponding to parameter value t, an equation of the tangent line is $y - (2t^3 + 1) = t[x - (3t^2 + 1)]$. If this line is to pass through $(4, 3)$, we must have $3 - (2t^3 + 1) = t[4 - (3t^2 + 1)] \quad\Leftrightarrow\quad 2t^3 - 2 = 3t^3 - 3t \quad\Leftrightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$t^3 - 3t + 2 = 0 \quad \Leftrightarrow \quad (t-1)^2(t+2) = 0 \quad \Leftrightarrow \quad t = 1$ or -2. Hence, the desired equations are $y - 3 = x - 4$, or $y = x - 1$, tangent to the curve at $(4, 3)$, and $y - (-15) = -2(x - 13)$, or $y = -2x + 11$, tangent to the curve at $(13, -15)$.

35. The curve $x = t^3 + 1$, $y = 2t - t^2 = t(2 - t)$ intersects the x-axis when $y = 0$, that is, when $t = 0$ and $t = 2$. The corresponding values of x are 1 and 9. The shaded area is given by

$$\int_{x=1}^{x=9} (y_T - y_B)\,dx = \int_{t=0}^{t=2} [y(t) - 0]\,x'(t)\,dt = \int_0^2 (2t - t^2)(3t^2)\,dt$$
$$= 3\textstyle\int_0^2 (2t^3 - t^4)\,dt = 3\left[\frac{1}{2}t^4 - \frac{1}{5}t^5\right]_0^2 = 3\left(8 - \frac{32}{5}\right) = \frac{24}{5}$$

36. The curve $x = \sin t$, $y = \sin t \cos t$, $0 \le t \le \pi/2$ intersects the x-axis when $y = 0$, that is, when $t = 0$ and $t = \pi/2$. The corresponding values of x are 0 and 1, so the area enclosed by the curve and the x-axis is given by

$$\int_{x=0}^{x=1} y\,dx = \int_{t=0}^{t=\pi/2} y(t)\,x'(t)\,dt = \int_0^{\pi/2} \sin t \cos t\,(\cos t)\,dt \overset{c}{=} -\int_1^0 u^2\,du = \left[\tfrac{1}{3}u^3\right]_0^1 = \tfrac{1}{3}$$

37. The curve $x = \sin^2 t$, $y = \cos t$ intersects the y-axis when $x = 0$, that is, when $t = 0$ and $t = \pi$. (Any integer multiple of π will result in $x = 0$, though we choose two values of t over which the curve is traced out once.) The corresponding values of y are 1 and -1, so the area enclosed by the curve and the y-axis is given by

$$\int_{y=-1}^{y=1} x\,dy = \int_{t=\pi}^{t=0} x(t)\,y'(t)\,dt = \int_\pi^0 \sin^2 t\,(-\sin t)\,dt = \int_0^\pi \sin^3 t\,dt \overset{67}{=} \left[-\frac{1}{3}(2 + \sin^2 t)\cos t\right]_0^\pi$$
$$= \tfrac{2}{3} - \left(-\tfrac{2}{3}\right) = \tfrac{4}{3}$$

38. The curve $x = t^2 - 2t = t(t - 2)$, $y = \sqrt{t}$ intersects the y-axis when $x = 0$, that is, when $t = 0$ and $t = 2$. The corresponding values of y are 0 and $\sqrt{2}$. The shaded area is given by

$$\int_{y=0}^{y=\sqrt{2}} (x_R - x_L)\,dy = \int_{t=0}^{t=2} [0 - x(t)]\,y'(t)\,dt = -\int_0^2 (t^2 - 2t)\left(\frac{1}{2\sqrt{t}}\,dt\right)$$
$$= -\textstyle\int_0^2 \left(\frac{1}{2}t^{3/2} - t^{1/2}\right) dt = -\left[\frac{1}{5}t^{5/2} - \frac{2}{3}t^{3/2}\right]_0^2$$
$$= -\left(\tfrac{1}{5}\cdot 2^{5/2} - \tfrac{2}{3}\cdot 2^{3/2}\right) = -2^{1/2}\left(\tfrac{4}{5} - \tfrac{4}{3}\right)$$
$$= -\sqrt{2}\left(-\tfrac{8}{15}\right) = \tfrac{8}{15}\sqrt{2}$$

39. $x = a\cos\theta$, $y = b\sin\theta$, $0 \le \theta \le 2\pi$. By symmetry of the ellipse about the x- and y-axes,

$$A = 4\int_{x=0}^{x=a} y\,dx = 4\int_{\theta=\pi/2}^{\theta=0} b\sin\theta\,(-a\sin\theta)\,d\theta = 4ab\textstyle\int_0^{\pi/2} \sin^2\theta\,d\theta = 4ab\int_0^{\pi/2} \frac{1}{2}(1 - \cos 2\theta)\,d\theta$$
$$= 2ab\left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/2} = 2ab\left(\tfrac{\pi}{2}\right) = \pi ab$$

40. By symmetry, the area of the shaded region is twice the area of the shaded portion above the x-axis. The top half of the loop is described by $x = 1 - t^2$, $y = t - t^3 = t(1 - t)(1 + t)$, $0 \le t \le 1$ with x-intercepts 0 and 1 corresponding to $t = 1$ and $t = 0$, respectively. Thus, the area of the shaded region is

$$2\textstyle\int_0^1 y\,dx = 2\int_1^0 y(t)\,x'(t)\,dt = 2\int_1^0 (t - t^3)(-2t)\,dt = 4\int_0^1 (t^2 - t^4)\,dt = 4\left[\frac{1}{3}t^3 - \frac{1}{5}t^5\right]_0^1 = 4\left(\frac{1}{3} - \frac{1}{5}\right) = \frac{8}{15}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. $x = r\theta - d\sin\theta,\ y = r - d\cos\theta.$

$$A = \int_0^{2\pi r} y\,dx = \int_0^{2\pi}(r - d\cos\theta)(r - d\cos\theta)\,d\theta = \int_0^{2\pi}(r^2 - 2dr\cos\theta + d^2\cos^2\theta)\,d\theta$$
$$= \left[r^2\theta - 2dr\sin\theta + \tfrac{1}{2}d^2\left(\theta + \tfrac{1}{2}\sin 2\theta\right)\right]_0^{2\pi} = 2\pi r^2 + \pi d^2$$

42. (a) By symmetry, the area of $\mathcal{R}$ is twice the area inside $\mathcal{R}$ above the x-axis. The top half of the loop is described by $x = t^2$, $y = t^3 - 3t$, $-\sqrt{3} \le t \le 0$, so, using the Substitution Rule with $y = t^3 - 3t$ and $dx = 2t\,dt$, we find that

$$\text{area} = 2\int_0^3 y\,dx = 2\int_0^{-\sqrt{3}}(t^3 - 3t)2t\,dt = 2\int_0^{-\sqrt{3}}(2t^4 - 6t^2)\,dt = 2\left[\tfrac{2}{5}t^5 - 2t^3\right]_0^{-\sqrt{3}}$$
$$= 2\left[\tfrac{2}{5}(-3^{1/2})^5 - 2(-3^{1/2})^3\right] = 2\left[\tfrac{2}{5}\left(-9\sqrt{3}\right) - 2\left(-3\sqrt{3}\right)\right] = \tfrac{24}{5}\sqrt{3}$$

(b) Here we use the formula for disks and use the Substitution Rule as in part (a):

$$\text{volume} = \pi\int_0^3 y^2\,dx = \pi\int_0^{-\sqrt{3}}(t^3 - 3t)^2 2t\,dt = 2\pi\int_0^{-\sqrt{3}}(t^6 - 6t^4 + 9t^2)t\,dt = 2\pi\left[\tfrac{1}{8}t^8 - t^6 + \tfrac{9}{4}t^4\right]_0^{-\sqrt{3}}$$
$$= 2\pi\left[\tfrac{1}{8}(-3^{1/2})^8 - (-3^{1/2})^6 + \tfrac{9}{4}(-3^{1/2})^4\right] = 2\pi\left[\tfrac{81}{8} - 27 + \tfrac{81}{4}\right] = \tfrac{27}{4}\pi$$

(c) By symmetry, the y-coordinate of the centroid is 0. To find the x-coordinate, we note that it is the same as the x-coordinate of the centroid of the top half of $\mathcal{R}$, the area of which is $\frac{1}{2}\cdot\frac{24}{5}\sqrt{3} = \frac{12}{5}\sqrt{3}$. So, using Formula 8.3.8 with $A = \frac{12}{5}\sqrt{3}$, we get

$$\overline{x} = \tfrac{5}{12\sqrt{3}}\int_0^3 xy\,dx = \tfrac{5}{12\sqrt{3}}\int_0^{-\sqrt{3}} t^2(t^3 - 3t)2t\,dt = \tfrac{5}{6\sqrt{3}}\left[\tfrac{1}{7}t^7 - \tfrac{3}{5}t^5\right]_0^{-\sqrt{3}}$$
$$= \tfrac{5}{6\sqrt{3}}\left[\tfrac{1}{7}(-3^{1/2})^7 - \tfrac{3}{5}(-3^{1/2})^5\right] = \tfrac{5}{6\sqrt{3}}\left[-\tfrac{27}{7}\sqrt{3} + \tfrac{27}{5}\sqrt{3}\right] = \tfrac{9}{7}$$

So the coordinates of the centroid of $\mathcal{R}$ are $(x, y) = \left(\frac{9}{7}, 0\right)$.

43. $x = 3t^2 - t^3$, $y = t^2 - 2t$. $dx/dt = 6t - 3t^2$ and $dy/dt = 2t - 2$, so $(dx/dt)^2 + (dy/dt)^2 = (6t - 3t^2)^2 + (2t - 2)^2 = 36t^2 - 36t^3 + 9t^4 + 4t^2 - 8t + 4 = 9t^4 - 36t^3 + 40t^2 - 8t + 4$. The endpoints of the curve both have $y = 3$, so the value of t at these points must satisfy $t^2 - 2t = 3 \;\Rightarrow\; t^2 - 2t - 3 = 0 \;\Rightarrow\; (t + 1)(t - 3) = 0 \;\Rightarrow\; t = -1$ or $t = 3$. Thus,

$$L = \int_a^b \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_{-1}^3 \sqrt{9t^4 - 36t^3 + 40t^2 - 8t + 4}\,dt \approx 15.2092$$

44. $x = t + e^{-t}$, $y = t^2 + t$. $dx/dt = 1 - e^{-t}$ and $dy/dt = 2t + 1$, so $(dx/dt)^2 + (dy/dt)^2 = (1 - e^{-t})^2 + (2t + 1)^2 = 1 - 2e^{-t} + e^{-2t} + 4t^2 + 4t + 1$. One endpoint of the curve has $y = 2$, so the value of t must satisfy $t^2 + t = 2 \;\Rightarrow\; t^2 + t - 2 = 0 \;\Rightarrow\; (t + 2)(t - 1) = 0 \;\Rightarrow\; t = -2$. (The solution $t = 1$ corresponds to $x \approx 1.37$, which is not an endpoint.) The other endoint has $y = 6 \;\Rightarrow\; t^2 + t = 6 \;\Rightarrow\; t^2 + t - 6 = 0 \;\Rightarrow\; (t + 3)(t - 2) = 0 \;\Rightarrow\; t = 2$. (The solution $t = -3$ corresponds to $x \approx 17.1$, which is not a point on the graph.) Thus,

$$L = \int_a^b \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_{-2}^2 \sqrt{2 - 2e^{-t} + e^{-2t} + 4t^2 + 4t}\,dt \approx 11.2485$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. $x = t - 2\sin t,\ y = 1 - 2\cos t,\ 0 \le t \le 4\pi.$ $dx/dt = 1 - 2\cos t$ and $dy/dt = 2\sin t$, so $(dx/dt)^2 + (dy/dt)^2 = (1 - 2\cos t)^2 + (2\sin t)^2 = 1 - 4\cos t + 4\cos^2 t + 4\sin^2 t = 5 - 4\cos t$. Thus, $L = \int_a^b \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_0^{4\pi} \sqrt{5 - 4\cos t}\,dt \approx 26.7298$.

46. $x = t\cos t,\ y = t - 5\sin t.$ $dx/dt = \cos t - t\sin t$ and $dy/dt = 1 - 5\cos t$, so $(dx/dt)^2 + (dy/dt)^2 = (\cos t - t\sin t)^2 + (1 - 5\cos t)^2$. Observe that when $t = -\pi$, $(x, y) = (\pi, -\pi)$ and when $t = \pi$, $(x, y) = (-\pi, \pi)$. Thus, $L = \int_a^b \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_{-\pi}^{\pi} \sqrt{(\cos t - t\sin t)^2 + (1 - 5\cos t)^2}\,dt \approx 22.8546$.

47. $x = \frac{2}{3}t^3,\ y = t^2 - 2,\ 0 \le t \le 3.$ $dx/dt = 2t^2$ and $dy/dt = 2t$, so $(dx/dt)^2 + (dy/dt)^2 = 4t^4 + 4t^2 = 4t^2(t^2 + 1)$. Thus,

$$L = \int_a^b \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_0^3 \sqrt{4t^2(t^2+1)}\,dt = \int_0^3 2t\sqrt{t^2+1}\,dt$$

$$= \int_1^{10} \sqrt{u}\,du \quad [\,u = t^2 + 1,\ du = 2t\,dt\,] \quad = \left[\tfrac{2}{3}u^{3/2}\right]_1^{10} = \tfrac{2}{3}(10^{3/2} - 1) = \tfrac{2}{3}\left(10\sqrt{10} - 1\right)$$

48. $x = e^t - t,\ y = 4e^{t/2},\ 0 \le t \le 2.$ $dx/dt = e^t - 1$ and $dy/dt = 2e^{t/2}$, so $(dx/dt)^2 + (dy/dt)^2 = (e^t - 1)^2 + (2e^{t/2})^2 = e^{2t} - 2e^t + 1 + 4e^t = e^{2t} + 2e^t + 1 = (e^t + 1)^2$. Thus,

$$L = \int_0^2 \sqrt{(e^t+1)^2}\,dt = \int_0^2 \left|e^t + 1\right|\,dt = \int_0^2 (e^t + 1)\,dt = \left[e^t + t\right]_0^2 = (e^2 + 2) - (1 + 0) = e^2 + 1.$$

49. $x = t\sin t,\ y = t\cos t,\ 0 \le t \le 1.$ $\dfrac{dx}{dt} = t\cos t + \sin t$ and $\dfrac{dy}{dt} = -t\sin t + \cos t$, so

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = t^2\cos^2 t + 2t\sin t\,\cos t + \sin^2 t + t^2\sin^2 t - 2t\sin t\,\cos t + \cos^2 t$$

$$= t^2(\cos^2 t + \sin^2 t) + \sin^2 t + \cos^2 t = t^2 + 1.$$

Thus, $L = \int_0^1 \sqrt{t^2+1}\,dt \overset{21}{=} \left[\frac{1}{2}t\sqrt{t^2+1} + \frac{1}{2}\ln\left(t + \sqrt{t^2+1}\right)\right]_0^1 = \frac{1}{2}\sqrt{2} + \frac{1}{2}\ln\left(1 + \sqrt{2}\right)$.

50. $x = 3\cos t - \cos 3t,\ y = 3\sin t - \sin 3t,\ 0 \le t \le \pi.$ $\dfrac{dx}{dt} = -3\sin t + 3\sin 3t$ and $\dfrac{dy}{dt} = 3\cos t - 3\cos 3t$, so

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = 9\sin^2 t - 18\sin t\,\sin 3t + 9\sin^2 3t + 9\cos^2 t - 18\cos t\,\cos 3t + 9\cos^2 3t$$

$$= 9(\cos^2 t + \sin^2 t) - 18(\cos t\,\cos 3t + \sin t\,\sin 3t) + 9(\cos^2 3t + \sin^2 3t)$$

$$= 9(1) - 18\cos(t - 3t) + 9(1) = 18 - 18\cos(-2t) = 18(1 - \cos 2t)$$

$$= 18[1 - (1 - 2\sin^2 t)] = 36\sin^2 t.$$

Thus, $L = \int_0^\pi \sqrt{36\sin^2 t}\,dt = 6\int_0^\pi |\sin t|\,dt = 6\int_0^\pi \sin t\,dt = -6\left[\cos t\right]_0^\pi = -6\,(-1 - 1) = 12$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51.

$x = e^t \cos t,\ y = e^t \sin t,\ 0 \le t \le \pi.$

$$\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2 = [e^t(\cos t - \sin t)]^2 + [e^t(\sin t + \cos t)]^2$$
$$= (e^t)^2(\cos^2 t - 2\cos t\ \sin t + \sin^2 t)$$
$$+\ (e^t)^2(\sin^2 t + 2\sin t\ \cos t + \cos^2 t$$
$$= e^{2t}(2\cos^2 t + 2\sin^2 t) = 2e^{2t}$$

Thus, $L = \int_0^\pi \sqrt{2e^{2t}}\,dt = \int_0^\pi \sqrt{2}\,e^t\,dt = \sqrt{2}\ \left[\,e^t\,\right]_0^\pi = \sqrt{2}\,(e^\pi - 1).$

52. $x = \cos t + \ln(\tan \frac{1}{2}t), \quad y = \sin t, \quad \pi/4 \le t \le 3\pi/4.$

$\dfrac{dx}{dt} = -\sin t + \dfrac{\frac{1}{2}\sec^2(t/2)}{\tan(t/2)} = -\sin t + \dfrac{1}{2\sin(t/2)\ \cos(t/2)} = -\sin t + \dfrac{1}{\sin t}$ and $\dfrac{dy}{dt} = \cos t$, so

$\left(\dfrac{dx}{dt}\right)^2 + \left(\dfrac{dy}{dt}\right)^2 = \sin^2 t - 2 + \dfrac{1}{\sin^2 t} + \cos^2 t = 1 - 2 + \csc^2 t = \cot^2 t$. Thus,

$$L = \int_{\pi/4}^{3\pi/4} |\cot t|\ dt = 2\int_{\pi/4}^{\pi/2} \cot t\,dt$$
$$= 2\Big[\ln|\sin t|\Big]_{\pi/4}^{\pi/2} = 2\left(\ln 1 - \ln\frac{1}{\sqrt{2}}\right)$$
$$= 2\big(0 + \ln\sqrt{2}\big) = 2\big(\tfrac{1}{2}\ln 2\big) = \ln 2.$$

53.

The figure shows the curve $x = \sin t + \sin 1.5t$, $y = \cos t$ for $0 \le t \le 4\pi$.

$dx/dt = \cos t + 1.5\cos 1.5t$ and $dy/dt = -\sin t$, so

$(dx/dt)^2 + (dy/dt)^2 = \cos^2 t + 3\cos t\ \cos 1.5t + 2.25\cos^2 1.5t + \sin^2 t.$

Thus, $L = \int_0^{4\pi} \sqrt{1 + 3\cos t\ \cos 1.5t + 2.25\cos^2 1.5t}\,dt \approx 16.7102.$

54. $x = 3t - t^3$, $y = 3t^2$. $\quad dx/dt = 3 - 3t^2$ and $dy/dt = 6t$, so

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (3 - 3t^2)^2 + (6t)^2 = (3 + 3t^2)^2$$

and the length of the loop is given by

$$L = \int_{-\sqrt{3}}^{\sqrt{3}} (3 + 3t^2)\,dt = 2\int_0^{\sqrt{3}} (3 + 3t^2)\,dt = 2\left[3t + t^3\right]_0^{\sqrt{3}}$$
$$= 2\big(3\sqrt{3} + 3\sqrt{3}\big) = 12\sqrt{3}$$

55. $x = \sin^2 t,\ y = \cos^2 t,\ 0 \le t \le 3\pi.$

$(dx/dt)^2 + (dy/dt)^2 = (2\sin t\ \cos t)^2 + (-2\cos t\sin t)^2 = 8\sin^2 t\cos^2 t = 2\sin^2 2t \quad \Rightarrow$

Distance $= \int_0^{3\pi} \sqrt{2}\,|\sin 2t|\ dt = 6\sqrt{2}\int_0^{\pi/2} \sin 2t\,dt$ [by symmetry] $= -3\sqrt{2}\left[\cos 2t\right]_0^{\pi/2} = -3\sqrt{2}\,(-1 - 1) = 6\sqrt{2}.$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The full curve is traversed as t goes from 0 to $\frac{\pi}{2}$, because the curve is the segment of $x + y = 1$ that lies in the first quadrant (since $x, y \geq 0$), and this segment is completely traversed as t goes from 0 to $\frac{\pi}{2}$. Thus, $L = \int_0^{\pi/2} \sin 2t\, dt = \sqrt{2}$, as above.

56. $x = \cos^2 t,\ y = \cos t,\ 0 \leq t \leq 4\pi.$ $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (-2\cos t \sin t)^2 + (-\sin t)^2 = \sin^2 t\,(4\cos^2 t + 1)$

$$\begin{aligned}
\text{Distance } &= \int_0^{4\pi} |\sin t| \sqrt{4\cos^2 t + 1}\, dt = 4\int_0^{\pi} \sin t \sqrt{4\cos^2 t + 1}\, dt \\
&= -4\int_1^{-1} \sqrt{4u^2 + 1}\, du \quad [u = \cos t,\ du = -\sin t\, dt] \quad = 4\int_{-1}^{1} \sqrt{4u^2 + 1}\, du \\
&= 8\int_0^1 \sqrt{4u^2 + 1}\, du = 8\int_0^{\tan^{-1} 2} \sec\theta \cdot \tfrac{1}{2}\sec^2\theta\, d\theta \qquad [2u = \tan\theta,\ 2\,du = \sec^2\theta\, d\theta] \\
&= 4\int_0^{\tan^{-1} 2} \sec^3\theta\, d\theta \overset{71}{=} \Big[2\sec\theta\tan\theta + 2\ln|\sec\theta + \tan\theta|\Big]_0^{\tan^{-1} 2} = 4\sqrt{5} + 2\ln\big(\sqrt{5} + 2\big)
\end{aligned}$$

Thus, $L = \int_0^{\pi} |\sin t| \sqrt{4\cos^2 t + 1}\, dt = \sqrt{5} + \frac{1}{2}\ln\big(\sqrt{5} + 2\big)$.

57. $x = 2t - 3,\ y = 2t^2 - 3t + 6.$ $dx/dt = 2$ and $dy/dt = 4t - 3$, so $v(t) = s'(t) = \sqrt{2^2 + (4t - 3)^2}$. Thus, the speed of the particle at $t = 5$ is $v(5) = \sqrt{4 + (4 \cdot 5 - 3)^2} = \sqrt{293} \approx 17.12$ m/s.

58. $x = 2 + 5\cos\left(\frac{\pi}{3}t\right),\ y = -2 + 7\sin\left(\frac{\pi}{3}t\right).$ $dx/dt = -\frac{5\pi}{3}\sin\left(\frac{\pi}{3}t\right)$ and $dy/dt = \frac{7\pi}{3}\cos\left(\frac{\pi}{3}t\right)$, so $v(t) = s'(t) = \sqrt{\left[-\frac{5\pi}{3}\sin\left(\frac{\pi}{3}t\right)\right]^2 + \left[\frac{7\pi}{3}\cos\left(\frac{\pi}{3}t\right)\right]^2}$. Thus, the speed of the particle at $t = 3$ is

$$v(3) = \sqrt{\frac{25\pi^2}{9}\sin^2\pi + \frac{49\pi^2}{9}\cos^2\pi} = \sqrt{\frac{49\pi^2}{9}} = \frac{7\pi}{3} \approx 7.33 \text{ m/s}.$$

59. $x = e^t,\ y = te^t.$ $dx/dt = e^t$ and $dy/dt = te^t + e^t$, so $v(t) = s'(t) = \sqrt{(e^t)^2 + (te^t + e^t)^2}$. At (e, e), $x = e^t = e \Rightarrow t = 1$. Thus, the speed of the particle at (e, e) is $v(1) = \sqrt{e^2 + (e + e)^2} = \sqrt{5e^2} = \sqrt{5}\,e \approx 6.08$ m/s.

60. $x = t^2 + 1,\ y = t^4 + 2t^2 + 1.$ $dx/dt = 2t$ and $dy/dt = 4t^3 + 4t$, so $v(t) = s'(t) = \sqrt{4t^2 + (4t^3 + 4t)^2}$. At $(2, 4)$, $x = t^2 + 1 = 2 \Rightarrow t^2 = 1 \Rightarrow t = -1$ or $t = 1$. Both values result in the same speed since the function v is even. Thus, the speed of the particle at $(2, 4)$ is $v(1) = \sqrt{4 \cdot 1^2 + (4 \cdot 1^3 + 4 \cdot 1)^2} = \sqrt{68} = 2\sqrt{17} \approx 8.25$ m/s.

61. $x = (v_0\cos\alpha)t,\ y = (v_0\sin\alpha)t - \frac{1}{2}gt^2.$ $dx/dt = v_0\cos\alpha$ and $dy/dt = v_0\sin\alpha - gt$, so speed $= v(t) = s'(t) = \sqrt{(dx/dt)^2 + (dy/dt)^2} = \sqrt{v_0^2\cos^2\alpha + (v_0\sin\alpha - gt)^2}$.

(a) The projectile hits the ground when $y = 0 \Rightarrow (v_0\sin\alpha)t - \frac{1}{2}gt^2 = 0 \Rightarrow t\big(v_0\sin\alpha - \frac{1}{2}gt\big) = 0 \Rightarrow t = 0$ or $t = \dfrac{2v_0\sin\alpha}{g}$. The second solution gives the time at which the projectile hits the ground, and at this time it will have a speed of

$$\begin{aligned}
v\left(\frac{2v_0\sin\alpha}{g}\right) &= \sqrt{v_0^2\cos^2\alpha + \left[v_0\sin\alpha - g\left(\frac{2v_0\sin\alpha}{g}\right)\right]^2} = \sqrt{v_0^2\cos^2\alpha + (-v_0\sin\alpha)^2} \\
&= \sqrt{v_0^2\cos^2\alpha + v_0^2\sin^2\alpha} = \sqrt{v_0^2(\cos^2\alpha + \sin^2\alpha)} = \sqrt{v_0^2} = v_0 \text{ m/s}.
\end{aligned}$$

Thus, the projectile hits the ground with the same speed at which it was fired.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The projectile is at its highest point (maximum height) when $dy/dt = 0$. Thus, the speed of the projectile at this time is

$v(t) = \sqrt{(dx/dt)^2 + (dy/dt)^2} = \sqrt{v_0^2 \cos^2\alpha + (0)^2} = v_0 \cos\alpha$ m/s.

62. $x = a\sin\theta,\ y = b\cos\theta,\ 0 \le \theta \le 2\pi.$

$$\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2 = (a\cos\theta)^2 + (-b\sin\theta)^2 = a^2\cos^2\theta + b^2\sin^2\theta = a^2(1-\sin^2\theta) + b^2\sin^2\theta$$
$$= a^2 - (a^2 - b^2)\sin^2\theta = a^2 - c^2\sin^2\theta = a^2\left(1 - \frac{c^2}{a^2}\sin^2\theta\right) = a^2(1 - e^2\sin^2\theta)$$

So $L = 4\int_0^{\pi/2} \sqrt{a^2\left(1 - e^2\sin^2\theta\right)}\, d\theta$ [by symmetry] $= 4a\int_0^{\pi/2} \sqrt{1 - e^2\sin^2\theta}\, d\theta.$

63. (a) $x = 11\cos t - 4\cos(11t/2),\ y = 11\sin t - 4\sin(11t/2).$

Notice that $0 \le t \le 2\pi$ does not give the complete curve because $x(0) \ne x(2\pi)$. In fact, we must take $t \in [0, 4\pi]$ in order to obtain the complete curve, since the first term in each of the parametric equations has period 2π and the second has period $\frac{2\pi}{11/2} = \frac{4\pi}{11}$, and the least common integer multiple of these two numbers is 4π.

(b) We use the CAS to find the derivatives dx/dt and dy/dt, and then use Theorem 5 to find the arc length. Recent versions of Maple express the integral $\int_0^{4\pi} \sqrt{(dx/dt)^2 + (dy/dt)^2}\, dt$ as $88E\left(2\sqrt{2}\, i\right)$, where $E(x)$ is the elliptic integral $\displaystyle\int_0^1 \frac{\sqrt{1 - x^2t^2}}{\sqrt{1 - t^2}}\, dt$ and i is the imaginary number $\sqrt{-1}$.

Some earlier versions of Maple (as well as Mathematica) cannot do the integral exactly, so we use the command `evalf(Int(sqrt(diff(x,t)^2+diff(y,t)^2),t=0..4*Pi));` to estimate the length, and find that the arc length is approximately 294.03.

64. (a) It appears that as $t \to \infty$, $(x, y) \to \left(\frac{1}{2}, \frac{1}{2}\right)$, and as $t \to -\infty$, $(x, y) \to \left(-\frac{1}{2}, -\frac{1}{2}\right)$.

(b) By the Fundamental Theorem of Calculus, $x = C(t) = \int_0^t \cos(\pi u^2/2)\, du \quad \Rightarrow$ $dx/dt = \cos\left(\frac{\pi}{2}t^2\right)$ and $y = S(t) = \int_0^t \sin(\pi u^2/2)\, du \quad \Rightarrow \quad dy/dt = \sin\left(\frac{\pi}{2}t^2\right)$, so by Theorem 5, the length of the curve from the origin to the point with parameter value t is

$$L = \int_0^t \sqrt{\left(\tfrac{dx}{du}\right)^2 + \left(\tfrac{dy}{du}\right)^2}\, du = \int_0^t \sqrt{\cos^2\left(\tfrac{\pi}{2}u^2\right) + \sin^2\left(\tfrac{\pi}{2}u^2\right)}\, du$$
$$= \int_0^t 1\, du = t \qquad [\text{or } -t \text{ if } t < 0]$$

We have used u as the dummy variable so as not to confuse it with the upper limit of integration.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

65. $x = a\cos^3\theta,\ y = a\sin^3\theta$. By symmetry,

$A = 4\int_0^a y\,dx = 4\int_{\pi/2}^0 a\sin^3\theta(-3a\cos^2\theta\,\sin\theta)\,d\theta = 12a^2\int_0^{\pi/2}\sin^4\theta\,\cos^2\theta\,d\theta$. Now

$$\int \sin^4\theta\,\cos^2\theta\,d\theta = \int \sin^2\theta\left(\tfrac{1}{4}\sin^2 2\theta\right)d\theta = \tfrac{1}{8}\int(1-\cos 2\theta)\sin^2 2\theta\,d\theta$$
$$= \tfrac{1}{8}\int\left[\tfrac{1}{2}(1-\cos 4\theta) - \sin^2 2\theta\,\cos 2\theta\right]d\theta = \tfrac{1}{16}\theta - \tfrac{1}{64}\sin 4\theta - \tfrac{1}{48}\sin^3 2\theta + C$$

so $\int_0^{\pi/2}\sin^4\theta\,\cos^2\theta\,d\theta = \left[\tfrac{1}{16}\theta - \tfrac{1}{64}\sin 4\theta - \tfrac{1}{48}\sin^3 2\theta\right]_0^{\pi/2} = \tfrac{\pi}{32}$. Thus, $A = 12a^2\left(\tfrac{\pi}{32}\right) = \tfrac{3}{8}\pi a^2$.

66. By symmetry, the perimeter P of the astroid $x = a\cos^3\theta$, $y = a\sin^3\theta$ is given by

$$P = 4\int_0^{\pi/2}\sqrt{(dx/d\theta)^2 + (dy/d\theta)^2}\,d\theta = 4\int_0^{\pi/2}\sqrt{[3a\cos^2\theta(-\sin\theta)]^2 + [3a\sin^2\theta(\cos\theta)]^2}\,d\theta$$
$$= 4\int_0^{\pi/2}\sqrt{9a^2\sin^2\theta\cos^4\theta + 9a^2\sin^4\theta\cos^2\theta}\,d\theta = 4\int_0^{\pi/2}3a\,|\sin\theta\cos\theta|\,\sqrt{\cos^2\theta + \sin^2\theta}\,d\theta$$
$$= 6a\int_0^{\pi/2}2\sin\theta\cos\theta\sqrt{1}\,d\theta = 6a\int_0^{\pi/2}\sin 2\theta\,d\theta = 6a\left[-\tfrac{1}{2}\cos 2\theta\right]_0^{\pi/2} = -3a(-1-1) = 6a$$

67. $x = t\sin t,\ y = t\cos t,\ 0 \le t \le \pi/2$. $dx/dt = t\cos t + \sin t$ and $dy/dt = -t\sin t + \cos t$, so

$$(dx/dt)^2 + (dy/dt)^2 = t^2\cos^2 t + 2t\sin t\,\cos t + \sin^2 t + t^2\sin^2 t - 2t\sin t\,\cos t + \cos^2 t$$
$$= t^2(\cos^2 t + \sin^2 t) + \sin^2 t + \cos^2 t = t^2 + 1$$

$S = \int 2\pi y\,ds = \int_0^{\pi/2}2\pi t\cos t\sqrt{t^2+1}\,dt \approx 4.7394$.

68. $x = \sin t,\ y = \sin 2t,\ 0 \le t \le \pi/2$. $dx/dt = \cos t$ and $dy/dt = 2\cos 2t$, so $(dx/dt)^2 + (dy/dt)^2 = \cos^2 t + 4\cos^2 2t$.

$S = \int 2\pi y\,ds = \int_0^{\pi/2}2\pi\sin 2t\sqrt{\cos^2 t + 4\cos^2 2t}\,dt \approx 8.0285$.

69. $x = t + e^t,\ y = e^{-t},\ 0 \le t \le 1$.

$dx/dt = 1 + e^t$ and $dy/dt = -e^{-t}$, so $(dx/dt)^2 + (dy/dt)^2 = (1+e^t)^2 + (-e^{-t})^2 = 1 + 2e^t + e^{2t} + e^{-2t}$.

$S = \int 2\pi y\,ds = \int_0^1 2\pi e^{-t}\sqrt{1 + 2e^t + e^{2t} + e^{-2t}}\,dt \approx 10.6705$.

70. $x = t^2 - t^3,\ y = t + t^4,\ 0 \le t \le 1$.

$(dx/dt)^2 + (dy/dt)^2 = (2t - 3t^2)^2 + (1 + 4t^3)^2 = 4t^2 - 12t^3 + 9t^4 + 1 + 8t^3 + 16t^6$, so

$S = \int 2\pi y\,ds = \int_0^1 2\pi(t + t^4)\sqrt{16t^6 + 9t^4 - 4t^3 + 4t^2 + 1}\,dt \approx 12.7176$.

71. $x = t^3,\ y = t^2,\ 0 \le t \le 1$. $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = \left(3t^2\right)^2 + (2t)^2 = 9t^4 + 4t^2$.

$$S = \int_0^1 2\pi y\sqrt{\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2}\,dt = \int_0^1 2\pi t^2\sqrt{9t^4 + 4t^2}\,dt = 2\pi\int_0^1 t^2\sqrt{t^2(9t^2+4)}\,dt$$
$$= 2\pi\int_4^{13}\left(\frac{u-4}{9}\right)\sqrt{u}\left(\tfrac{1}{18}\,du\right) \quad \left[\begin{array}{l} u = 9t^2 + 4,\ t^2 = (u-4)/9, \\ du = 18t\,dt, \text{ so } t\,dt = \frac{1}{18}\,du \end{array}\right] \quad = \frac{2\pi}{9\cdot 18}\int_4^{13}(u^{3/2} - 4u^{1/2})\,du$$
$$= \tfrac{\pi}{81}\left[\tfrac{2}{5}u^{5/2} - \tfrac{8}{3}u^{3/2}\right]_4^{13} = \tfrac{\pi}{81}\cdot\tfrac{2}{15}\left[3u^{5/2} - 20u^{3/2}\right]_4^{13}$$
$$= \tfrac{2\pi}{1215}\left[\left(3\cdot 13^2\sqrt{13} - 20\cdot 13\sqrt{13}\right) - (3\cdot 32 - 20\cdot 8)\right] = \tfrac{2\pi}{1215}\left(247\sqrt{13} + 64\right)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

72. $x = 2t^2 + 1/t,\ y = 8\sqrt{t},\ 1 \le t \le 3.$

$$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = \left(4t - \frac{1}{t^2}\right)^2 + \left(\frac{4}{\sqrt{t}}\right)^2 = 16t^2 - \frac{8}{t} + \frac{1}{t^4} + \frac{16}{t} = 16t^2 + \frac{8}{t} + \frac{1}{t^4} = \left(4t + \frac{1}{t^2}\right)^2.$$

$$S = \int_1^3 2\pi y\sqrt{\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2}\,dt = \int_1^3 2\pi\left(8\sqrt{t}\right)\sqrt{\left(4t + \frac{1}{t^2}\right)^2}\,dt = 16\pi\int_1^3 t^{1/2}(4t + t^{-2})\,dt$$

$$= 16\pi\int_1^3 (4t^{3/2} + t^{-3/2})\,dt = 16\pi\left[\tfrac{8}{5}t^{5/2} - 2t^{-1/2}\right]_1^3 = 16\pi\left[\left(\tfrac{72}{5}\sqrt{3} - \tfrac{2}{3}\sqrt{3}\right) - \left(\tfrac{8}{5} - 2\right)\right]$$

$$= 16\pi\left(\tfrac{206}{15}\sqrt{3} + \tfrac{6}{15}\right) = \tfrac{32\pi}{15}\left(103\sqrt{3} + 3\right)$$

73. $x = a\cos^3\theta,\ y = a\sin^3\theta,\ 0 \le \theta \le \frac{\pi}{2}.\quad \left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2 = (-3a\cos^2\theta\,\sin\theta)^2 + (3a\sin^2\theta\,\cos\theta)^2 = 9a^2\sin^2\theta\,\cos^2\theta.$

$S = \int_0^{\pi/2} 2\pi\cdot a\sin^3\theta\cdot 3a\sin\theta\,\cos\theta\,d\theta = 6\pi a^2\int_0^{\pi/2}\sin^4\theta\,\cos\theta\,d\theta = \frac{6}{5}\pi a^2\left[\sin^5\theta\right]_0^{\pi/2} = \frac{6}{5}\pi a^2$

74. $x = 2\cos\theta - \cos 2\theta,\ y = 2\sin\theta - \sin 2\theta \quad\Rightarrow$

$$\left(\tfrac{dx}{d\theta}\right)^2 + \left(\tfrac{dy}{d\theta}\right)^2 = (-2\sin\theta + 2\sin 2\theta)^2 + (2\cos\theta - 2\cos 2\theta)^2$$

$$= 4[(\sin^2\theta - 2\sin\theta\,\sin 2\theta + \sin^2 2\theta) + (\cos^2\theta - 2\cos\theta\,\cos 2\theta + \cos^2 2\theta)]$$

$$= 4[1 + 1 - 2(\cos 2\theta\cos\theta + \sin 2\theta\sin\theta)] = 8[1 - \cos(2\theta - \theta)] = 8(1 - \cos\theta)$$

We plot the graph with parameter interval $[0, 2\pi]$, and see that we should only integrate between 0 and π. (If the interval $[0, 2\pi]$ were taken, the surface of revolution would be generated twice.) Also note that $y = 2\sin\theta - \sin 2\theta = 2\sin\theta(1 - \cos\theta)$. So

$S = \int_0^{\pi} 2\pi\cdot 2\sin\theta(1 - \cos\theta)\,2\sqrt{2}\sqrt{1 - \cos\theta}\,d\theta$

$= 8\sqrt{2}\pi\int_0^{\pi}(1 - \cos\theta)^{3/2}\sin\theta\,d\theta = 8\sqrt{2}\pi\int_0^2\sqrt{u^3}\,du \quad \begin{bmatrix} u = 1 - \cos\theta, \\ du = \sin\theta\,d\theta \end{bmatrix}$

$= 8\sqrt{2}\pi\left[\left(\frac{2}{5}\right)u^{5/2}\right]_0^2 = \frac{16}{5}\sqrt{2}\pi(2^{5/2}) = \frac{128}{5}\pi$

75. $x = 3t^2,\ y = 2t^3,\ 0 \le t \le 5 \quad\Rightarrow\quad \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (6t)^2 + (6t^2)^2 = 36t^2(1 + t^2) \quad\Rightarrow$

$$S = \int_0^5 2\pi x\sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_0^5 2\pi(3t^2)6t\sqrt{1 + t^2}\,dt = 18\pi\int_0^5 t^2\sqrt{1 + t^2}\,2t\,dt$$

$$= 18\pi\int_1^{26}(u - 1)\sqrt{u}\,du \quad \begin{bmatrix} u = 1 + t^2, \\ du = 2t\,dt \end{bmatrix} \quad = 18\pi\int_1^{26}(u^{3/2} - u^{1/2})\,du = 18\pi\left[\tfrac{2}{5}u^{5/2} - \tfrac{2}{3}u^{3/2}\right]_1^{26}$$

$$= 18\pi\left[\left(\tfrac{2}{5}\cdot 676\sqrt{26} - \tfrac{2}{3}\cdot 26\sqrt{26}\right) - \left(\tfrac{2}{5} - \tfrac{2}{3}\right)\right] = \tfrac{24}{5}\pi\left(949\sqrt{26} + 1\right)$$

76. $x = e^t - t, y = 4e^{t/2}, 0 \le t \le 1.\ \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (e^t - 1)^2 + (2e^{t/2})^2 = e^{2t} + 2e^t + 1 = (e^t + 1)^2.$

$$S = \int_0^1 2\pi(e^t - t)\sqrt{(e^t - 1)^2 + (2e^{t/2})^2}\,dt = \int_0^1 2\pi(e^t - t)(e^t + 1)d$$

$$= 2\pi\left[\tfrac{1}{2}e^{2t} + e^t - (t - 1)e^t - \tfrac{1}{2}t^2\right]_0^1 = \pi(e^2 + 2e - 6)$$

77. If f' is continuous and $f'(t) \ne 0$ for $a \le t \le b$, then either $f'(t) > 0$ for all t in $[a, b]$ or $f'(t) < 0$ for all t in $[a, b]$. Thus, f is monotonic (in fact, strictly increasing or strictly decreasing) on $[a, b]$. It follows that f has an inverse. Set $F = g \circ f^{-1}$, that is, define F by $F(x) = g(f^{-1}(x))$. Then $x = f(t) \quad\Rightarrow\quad f^{-1}(x) = t$, so $y = g(t) = g(f^{-1}(x)) = F(x)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

78. By Formula 8.2.5 with $y = F(x)$, $S = \int_a^b 2\pi F(x)\sqrt{1+[F'(x)]^2}\,dx$. But by Formula 10.2.1,

$1+[F'(x)]^2 = 1+\left(\dfrac{dy}{dx}\right)^2 = 1+\left(\dfrac{dy/dt}{dx/dt}\right)^2 = \dfrac{(dx/dt)^2+(dy/dt)^2}{(dx/dt)^2}$. Using the Substitution Rule with $x = x(t)$,

where $a = x(\alpha)$ and $b = x(\beta)$, we have $\left[\text{since } dx = \dfrac{dx}{dt}\,dt\right]$

$$S = \int_\alpha^\beta 2\pi\, F(x(t))\sqrt{\frac{(dx/dt)^2+(dy/dt)^2}{(dx/dt)^2}}\,\frac{dx}{dt}\,dt = \int_\alpha^\beta 2\pi y\sqrt{\left(\frac{dx}{dt}\right)^2+\left(\frac{dy}{dt}\right)^2}\,dt,$$ which is Formula 10.2.9.

79. $\phi = \tan^{-1}\left(\dfrac{dy}{dx}\right) \;\Rightarrow\; \dfrac{d\phi}{dt} = \dfrac{d}{dt}\tan^{-1}\left(\dfrac{dy}{dx}\right) = \dfrac{1}{1+(dy/dx)^2}\left[\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)\right]$. But $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{\dot y}{\dot x} \;\Rightarrow$

$\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right) = \dfrac{d}{dt}\left(\dfrac{\dot y}{\dot x}\right) = \dfrac{\ddot y\dot x - \ddot x\dot y}{\dot x^2} \;\Rightarrow\; \dfrac{d\phi}{dt} = \dfrac{1}{1+(\dot y/\dot x)^2}\left(\dfrac{\ddot y\dot x - \ddot x\dot y}{\dot x^2}\right) = \dfrac{\dot x\ddot y - \ddot x\dot y}{\dot x^2+\dot y^2}$. Using the Chain Rule, and the fact

that $s = \displaystyle\int_0^t \sqrt{\left(\tfrac{dx}{dt}\right)^2+\left(\tfrac{dy}{dt}\right)^2}\,dt \;\Rightarrow\; \tfrac{ds}{dt} = \sqrt{\left(\tfrac{dx}{dt}\right)^2+\left(\tfrac{dy}{dt}\right)^2} = \left(\dot x^2+\dot y^2\right)^{1/2}$, we have that

$\dfrac{d\phi}{ds} = \dfrac{d\phi/dt}{ds/dt} = \left(\dfrac{\dot x\ddot y - \ddot x\dot y}{\dot x^2+\dot y^2}\right)\dfrac{1}{(\dot x^2+\dot y^2)^{1/2}} = \dfrac{\dot x\ddot y - \ddot x\dot y}{(\dot x^2+\dot y^2)^{3/2}}$. So $\kappa = \left|\dfrac{d\phi}{ds}\right| = \left|\dfrac{\dot x\ddot y - \ddot x\dot y}{(\dot x^2+\dot y^2)^{3/2}}\right| = \dfrac{|\dot x\ddot y - \ddot x\dot y|}{(\dot x^2+\dot y^2)^{3/2}}$.

80. $x = x$ and $y = f(x) \;\Rightarrow\; \dot x = 1$, $\ddot x = 0$ and $\dot y = \dfrac{dy}{dx}$, $\ddot y = \dfrac{d^2y}{dx^2}$.

So $\kappa = \dfrac{\left|1\cdot(d^2y/dx^2) - 0\cdot(dy/dx)\right|}{[1+(dy/dx)^2]^{3/2}} = \dfrac{\left|d^2y/dx^2\right|}{[1+(dy/dx)^2]^{3/2}}$.

81. $x = \theta - \sin\theta \;\Rightarrow\; \dot x = 1-\cos\theta \;\Rightarrow\; \ddot x = \sin\theta$, and $y = 1-\cos\theta \;\Rightarrow\; \dot y = \sin\theta \;\Rightarrow\; \ddot y = \cos\theta$. Therefore,

$\kappa = \dfrac{\left|\cos\theta - \cos^2\theta - \sin^2\theta\right|}{[(1-\cos\theta)^2+\sin^2\theta]^{3/2}} = \dfrac{\left|\cos\theta - (\cos^2\theta+\sin^2\theta)\right|}{(1-2\cos\theta+\cos^2\theta+\sin^2\theta)^{3/2}} = \dfrac{|\cos\theta - 1|}{(2-2\cos\theta)^{3/2}}$. The top of the arch is

characterized by a horizontal tangent, and from Example 2(b) in Section 10.2, the tangent is horizontal when $\theta = (2n-1)\pi$,

so take $n = 1$ and substitute $\theta = \pi$ into the expression for κ: $\kappa = \dfrac{|\cos\pi - 1|}{(2-2\cos\pi)^{3/2}} = \dfrac{|-1-1|}{[2-2(-1)]^{3/2}} = \dfrac{1}{4}$.

82. (a) $y = x^2 \;\Rightarrow\; \dfrac{dy}{dx} = 2x \;\Rightarrow\; \dfrac{d^2y}{dx^2} = 2$. So $\kappa = \dfrac{\left|d^2y/dx^2\right|}{[1+(dy/dx)^2]^{3/2}} = \dfrac{2}{(1+4x^2)^{3/2}}$, and at $(1,1)$,

$\kappa = \dfrac{2}{5^{3/2}} = \dfrac{2}{5\sqrt{5}}$.

(b) $\kappa' = \dfrac{d\kappa}{dx} = -3(1+4x^2)^{-5/2}(8x) = 0 \;\Leftrightarrow\; x = 0 \;\Rightarrow\; y = 0$. This is a maximum since $\kappa' > 0$ for $x < 0$ and

$\kappa' < 0$ for $x > 0$. So the parabola $y = x^2$ has maximum curvature at the origin.

83. (a) Every straight line has parametrizations of the form $x = a + vt$, $y = b + wt$, where a, b are arbitrary and v, $w \neq 0$.

For example, a straight line passing through distinct points (a, b) and (c, d) can be described as the parametrized curve

$x = a + (c-a)t$, $y = b + (d-b)t$. Starting with $x = a + vt$, $y = b + wt$, we compute $\dot x = v$, $\dot y = w$, $\ddot x = \ddot y = 0$,

and $\kappa = \dfrac{|v\cdot 0 - w\cdot 0|}{(v^2+w^2)^{3/2}} = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Parametric equations for a circle of radius r are $x = r\cos\theta$ and $y = r\sin\theta$. We can take the center to be the origin. So $\dot{x} = -r\sin\theta \;\Rightarrow\; \ddot{x} = -r\cos\theta$ and $\dot{y} = r\cos\theta \;\Rightarrow\; \ddot{y} = -r\sin\theta$. Therefore,

$\kappa = \dfrac{\left|r^2\sin^2\theta + r^2\cos^2\theta\right|}{(r^2\sin^2\theta + r^2\cos^2\theta)^{3/2}} = \dfrac{r^2}{r^3} = \dfrac{1}{r}$. And so for any θ (and thus any point), $\kappa = \dfrac{1}{r}$.

84. If the cow walks with the rope taut, it traces out the portion of the involute in Exercise 85 corresponding to the range $0 \le \theta \le \pi$, arriving at the point $(-r, \pi r)$ when $\theta = \pi$. With the rope now fully extended, the cow walks in a semicircle of radius πr, arriving at $(-r, -\pi r)$. Finally, the cow traces out another portion of the involute, namely the reflection about the x-axis of the initial involute path. (This corresponds to the range $-\pi \le \theta \le 0$.) Referring to the figure, we see that the total grazing area is $2(A_1 + A_3)$. A_3 is one-quarter of the area of a circle of radius πr, so $A_3 = \frac{1}{4}\pi(\pi r)^2 = \frac{1}{4}\pi^3 r^2$. We will compute $A_1 + A_2$ and then subtract $A_2 = \frac{1}{2}\pi r^2$ to obtain A_1.

To find $A_1 + A_2$, first note that the rightmost point of the involute is $\left(\frac{\pi}{2}r, r\right)$. [To see this, note that $dx/d\theta = 0$ when $\theta = 0$ or $\frac{\pi}{2}$. $\theta = 0$ corresponds to the cusp at $(r, 0)$ and $\theta = \frac{\pi}{2}$ corresponds to $\left(\frac{\pi}{2}r, r\right)$.] The leftmost point of the involute is $(-r, \pi r)$. Thus, $A_1 + A_2 = \int_{\theta=\pi}^{\pi/2} y\,dx - \int_{\theta=0}^{\pi/2} y\,dx = \int_{\theta=\pi}^{0} y\,dx$.

Now $y\,dx = r(\sin\theta - \theta\cos\theta)\,r\theta\cos\theta\,d\theta = r^2(\theta\sin\theta\,\cos\theta - \theta^2\cos^2\theta)d\theta$. Integrate:

$(1/r^2)\int y\,dx = -\theta\cos^2\theta - \frac{1}{2}\left(\theta^2 - 1\right)\sin\theta\,\cos\theta - \frac{1}{6}\theta^3 + \frac{1}{2}\theta + C$. This enables us to compute

$$A_1 + A_2 = r^2\left[-\theta\cos^2\theta - \tfrac{1}{2}(\theta^2 - 1)\,\sin\theta\,\cos\theta - \tfrac{1}{6}\theta^3 + \tfrac{1}{2}\theta\right]_{\pi}^{0} = r^2\left[0 - \left(-\pi - \frac{\pi^3}{6} + \frac{\pi}{2}\right)\right] = r^2\left(\frac{\pi}{2} + \frac{\pi^3}{6}\right)$$

Therefore, $A_1 = (A_1 + A_2) - A_2 = \frac{1}{6}\pi^3 r^2$, so the grazing area is $2(A_1 + A_3) = 2\left(\frac{1}{6}\pi^3 r^2 + \frac{1}{4}\pi^3 r^2\right) = \frac{5}{6}\pi^3 r^2$.

85. The coordinates of T are $(r\cos\theta, r\sin\theta)$. Since TP was unwound from arc TA, TP has length $r\theta$. Also $\angle PTQ = \angle PTR - \angle QTR = \frac{1}{2}\pi - \theta$, so P has coordinates $x = r\cos\theta + r\theta\cos\left(\frac{1}{2}\pi - \theta\right) = r(\cos\theta + \theta\sin\theta)$, $y = r\sin\theta - r\theta\sin\left(\frac{1}{2}\pi - \theta\right) = r(\sin\theta - \theta\cos\theta)$.

DISCOVERY PROJECT Bézier Curves

1. The parametric equations for a cubic Bézier curve are

$$x = x_0(1-t)^3 + 3x_1t(1-t)^2 + 3x_2t^2(1-t) + x_3t^3$$
$$y = y_0(1-t)^3 + 3y_1t(1-t)^2 + 3y_2t^2(1-t) + y_3t^3$$

where $0 \le t \le 1$. We are given the points $P_0(x_0, y_0) = (4, 1)$, $P_1(x_1, y_1) = (28, 48)$, $P_2(x_2, y_2) = (50, 42)$, and

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$P_3(x_3, y_3) = (40, 5)$. The curve is then given by

$$x(t) = 4(1-t)^3 + 3 \cdot 28t(1-t)^2 + 3 \cdot 50t^2(1-t) + 40t^3$$
$$y(t) = 1(1-t)^3 + 3 \cdot 48t(1-t)^2 + 3 \cdot 42t^2(1-t) + 5t^3$$

where $0 \le t \le 1$. The line segments are of the form $x = x_0 + (x_1 - x_0)t$, $y = y_0 + (y_1 - y_0)t$:

P_0P_1	$x = 4 + 24t,$	$y = 1 + 47t$
P_1P_2	$x = 28 + 22t,$	$y = 48 - 6t$
P_2P_3	$x = 50 - 10t,$	$y = 42 - 37t$

2. It suffices to show that the slope of the tangent at P_0 is the same as that of line segment P_0P_1, namely $\dfrac{y_1 - y_0}{x_1 - x_0}$.

We calculate the slope of the tangent to the Bézier curve:

$$\frac{dy/dt}{dx/dt} = \frac{-3y_0(1-t)^2 + 3y_1\left[-2t(1-t) + (1-t)^2\right] + 3y_2\left[-t^2 + (2t)(1-t)\right] + 3y_3t^2}{-3x_0^2(1-t) + 3x_1[-2t(1-t) + (1-t)^2] + 3x_2[-t^2 + (2t)(1-t)] + 3x_3t^2}$$

At point P_0, $t = 0$, so the slope of the tangent is $\dfrac{-3y_0 + 3y_1}{-3x_0 + 3x_1} = \dfrac{y_1 - y_0}{x_1 - x_0}$. So the tangent to the curve at P_0 passes through P_1. Similarly, the slope of the tangent at point P_3 [where $t = 1$] is $\dfrac{-3y_2 + 3y_3}{-3x_2 + 3x_3} = \dfrac{y_3 - y_2}{x_3 - x_2}$, which is also the slope of line P_2P_3.

3. It seems that if P_1 were to the right of P_2, a loop would appear.

We try setting $P_1 = (110, 30)$, and the resulting curve does indeed have a loop.

4. Based on the behavior of the Bézier curve in Problems 1–3, we suspect that the four control points should be in an exaggerated C shape. We try $P_0(10, 12)$, $P_1(4, 15)$, $P_2(4, 5)$, and $P_3(10, 8)$, and these produce a decent C. If you are using a CAS, it may be necessary to instruct it to make the x- and y-scales the same so as not to distort the figure (this is called a "constrained projection" in Maple.)

5. We use the same P_0 and P_1 as in Problem 4, and use part of our C as the top of an S. To prevent the center line from slanting up too much, we move P_2 up to $(4, 6)$ and P_3 down and to the left, to $(8, 7)$. In order to have a smooth joint between the top and bottom halves of the S (and a symmetric S), we determine points P_4, P_5, and P_6 by rotating points P_2, P_1, and P_0 about the center of the letter (point P_3). The points are therefore $P_4(12, 8)$, $P_5(12, -1)$, and $P_6(6, 2)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10.3 Polar Coordinates

1. (a) $\left(1, \frac{\pi}{4}\right)$

By adding 2π to $\frac{\pi}{4}$, we obtain the point $\left(1, \frac{9\pi}{4}\right)$, which satisfies the $r > 0$ requirement. The direction opposite $\frac{\pi}{4}$ is $\frac{5\pi}{4}$, so $\left(-1, \frac{5\pi}{4}\right)$ is a point that satisfies the $r < 0$ requirement.

(b) $\left(-2, \frac{3\pi}{2}\right)$

$\boldsymbol{r > 0}$**:** $\left(-(-2), \frac{3\pi}{2} - \pi\right) = \left(2, \frac{\pi}{2}\right)$

$\boldsymbol{r < 0}$**:** $\left(-2, \frac{3\pi}{2} + 2\pi\right) = \left(-2, \frac{7\pi}{2}\right)$

(c) $\left(3, -\frac{\pi}{3}\right)$

$\boldsymbol{r > 0}$**:** $\left(3, -\frac{\pi}{3} + 2\pi\right) = \left(3, \frac{5\pi}{3}\right)$

$\boldsymbol{r < 0}$**:** $\left(-3, -\frac{\pi}{3} + \pi\right) = \left(-3, \frac{2\pi}{3}\right)$

2. (a) $\left(2, \frac{5\pi}{6}\right)$

$\boldsymbol{r > 0}$**:** $\left(2, \frac{5\pi}{6} + 2\pi\right) = \left(2, \frac{17\pi}{6}\right)$

$\boldsymbol{r < 0}$**:** $\left(-2, \frac{5\pi}{6} - \pi\right) = \left(-2, -\frac{\pi}{6}\right)$

(b) $\left(1, -\frac{2\pi}{3}\right)$

$\boldsymbol{r > 0}$**:** $\left(1, -\frac{2\pi}{3} + 2\pi\right) = \left(1, \frac{4\pi}{3}\right)$

$\boldsymbol{r < 0}$**:** $\left(-1, -\frac{2\pi}{3} + \pi\right) = \left(-1, \frac{\pi}{3}\right)$

(c) $\left(-1, \frac{5\pi}{4}\right)$

$\boldsymbol{r > 0}$**:** $\left(-(-1), \frac{5\pi}{4} - \pi\right) = \left(1, \frac{\pi}{4}\right)$

$\boldsymbol{r < 0}$**:** $\left(-1, \frac{5\pi}{4} - 2\pi\right) = \left(-1, -\frac{3\pi}{4}\right)$

3. (a)

$x = 2\cos\frac{3\pi}{2} = 2(0) = 0$ and $y = 2\sin\frac{3\pi}{2} = 2(-1) = -2$ give us the Cartesian coordinates $(0, -2)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b)

$x = \sqrt{2}\cos\dfrac{\pi}{4} = \sqrt{2}\left(\dfrac{1}{\sqrt{2}}\right) = 1$ and $y = \sqrt{2}\sin\dfrac{\pi}{4} = \sqrt{2}\left(\dfrac{1}{\sqrt{2}}\right) = 1$ give us the Cartesian coordinates $(1, 1)$.

(c)

$x = -1\cos\left(-\dfrac{\pi}{6}\right) = -1\left(\dfrac{\sqrt{3}}{2}\right) = -\dfrac{\sqrt{3}}{2}$ and

$y = -1\sin\left(-\dfrac{\pi}{6}\right) = -1\left(-\dfrac{1}{2}\right) = \dfrac{1}{2}$ give us the Cartesian coordinates $\left(-\dfrac{\sqrt{3}}{2}, \dfrac{1}{2}\right)$.

4. (a)

$x = 4\cos\dfrac{4\pi}{3} = 4\left(-\dfrac{1}{2}\right) = -2$ and

$y = 4\sin\dfrac{4\pi}{3} = 4\left(-\dfrac{\sqrt{3}}{2}\right) = -2\sqrt{3}$ give us the Cartesian coordinates $\left(-2, -2\sqrt{3}\right)$.

(b)

$x = -2\cos\dfrac{3\pi}{4} = -2\left(-\dfrac{\sqrt{2}}{2}\right) = \sqrt{2}$ and

$y = -2\sin\dfrac{3\pi}{4} = -2\left(\dfrac{\sqrt{2}}{2}\right) = -\sqrt{2}$ give us the Cartesian coordinates $\left(\sqrt{2}, -\sqrt{2}\right)$.

(c)

$x = -3\cos\left(-\dfrac{\pi}{3}\right) = -3\left(\dfrac{1}{2}\right) = -\dfrac{3}{2}$ and

$y = -3\sin\left(-\dfrac{\pi}{3}\right) = -3\left(-\dfrac{\sqrt{3}}{2}\right) = \dfrac{3\sqrt{3}}{2}$ give us the Cartesian coordinates $\left(-\dfrac{3}{2}, \dfrac{3\sqrt{3}}{2}\right)$.

5. (a) $x = -4$ and $y = 4 \quad \Rightarrow \quad r = \sqrt{(-4)^2 + 4^2} = 4\sqrt{2}$ and $\tan\theta = \frac{4}{-4} = -1 \quad [\theta = -\frac{\pi}{4} + n\pi]$. Since $(-4, 4)$ is in the second quadrant, the polar coordinates are (i) $\left(4\sqrt{2}, \frac{3\pi}{4}\right)$ and (ii) $\left(-4\sqrt{2}, \frac{7\pi}{4}\right)$.

(b) $x = 3$ and $y = 3\sqrt{3} \quad \Rightarrow \quad r = \sqrt{3^2 + \left(3\sqrt{3}\right)^2} = \sqrt{9 + 27} = 6$ and $\tan\theta = \dfrac{3\sqrt{3}}{3} = \sqrt{3} \quad [\theta = \frac{\pi}{3} + n\pi]$.

Since $\left(3, 3\sqrt{3}\right)$ is in the first quadrant, the polar coordinates are (i) $\left(6, \frac{\pi}{3}\right)$ and (ii) $\left(-6, \frac{4\pi}{3}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. (a) $x = \sqrt{3}$ and $y = -1 \quad \Rightarrow \quad r = \sqrt{\left(\sqrt{3}\right)^2 + (-1)^2} = 2$ and $\tan\theta = \dfrac{-1}{\sqrt{3}} \quad [\theta = -\frac{\pi}{6} + n\pi]$. Since $\left(\sqrt{3}, -1\right)$ is in the fourth quadrant, the polar coordinates are (i) $\left(2, \frac{11\pi}{6}\right)$ and (ii) $\left(-2, \frac{5\pi}{6}\right)$.

(b) $x = -6$ and $y = 0 \quad \Rightarrow \quad r = \sqrt{(-6)^2 + 0^2} = 6$ and $\tan\theta = \frac{0}{-6} = 0 \quad [\theta = n\pi]$. Since $(-6, 0)$ is on the negative x-axis, the polar coordinates are (i) $(6, \pi)$ and (ii) $(-6, 0)$.

7. $1 < r \le 3$. The curves $r = 1$ and $r = 3$ represent circles centered at O with radius 1 and 3, respectively. So $1 < r \le 3$ represents the region outside the radius 1 circle and on or inside the radius 3 circle. Note that θ can take on any value.

8. $r \ge 2,\ 0 \le \theta \le \pi$. This is the region on or outside the circle $r = 2$ in the first and second quadrants.

9. $0 \le r \le 1,\ -\pi/2 \le \theta \le \pi/2$. This is the region on or inside the circle $r = 1$ in the first and fourth quadrants.

10. $3 < r < 5,\ 2\pi/3 \le \theta \le 4\pi/3$

11. $2 \le r < 4, 3\pi/4 \le \theta \le 7\pi/4$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $r \geq 0,\ \pi \leq \theta \leq 5\pi/2$. This is the region in the third, fourth, and first quadrants including the origin and points on the negative x-axis and positive y-axis.

13. Converting the polar coordinates $\left(4, \frac{4\pi}{3}\right)$ and $\left(6, \frac{5\pi}{3}\right)$ to Cartesian coordinates gives us $\left(4\cos\frac{4\pi}{3}, 4\sin\frac{4\pi}{3}\right) = \left(-2, -2\sqrt{3}\right)$ and $\left(6\cos\frac{5\pi}{3}, 6\sin\frac{5\pi}{3}\right) = \left(3, -3\sqrt{3}\right)$. Now use the distance formula

$$\begin{aligned} d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{[3 - (-2)]^2 + \left[-3\sqrt{3} - \left(-2\sqrt{3}\right)\right]^2} \\ &= \sqrt{5^2 + \left(-\sqrt{3}\right)^2} = \sqrt{25 + 3} = \sqrt{28} = 2\sqrt{7} \end{aligned}$$

14. The points (r_1, θ_1) and (r_2, θ_2) in Cartesian coordinates are $(r_1\cos\theta_1, r_1\sin\theta_1)$ and $(r_2\cos\theta_2, r_2\sin\theta_2)$, respectively. The *square* of the distance between them is

$$\begin{aligned} &(r_2\cos\theta_2 - r_1\cos\theta_1)^2 + (r_2\sin\theta_2 - r_1\sin\theta_1)^2 \\ &\qquad = \left(r_2^2\cos^2\theta_2 - 2r_1r_2\cos\theta_1\cos\theta_2 + r_1^2\cos^2\theta_1\right) + \left(r_2^2\sin^2\theta_2 - 2r_1r_2\sin\theta_1\sin\theta_2 + r_1^2\sin^2\theta_1\right) \\ &\qquad = r_1^2\left(\sin^2\theta_1 + \cos^2\theta_1\right) + r_2^2\left(\sin^2\theta_2 + \cos^2\theta_2\right) - 2r_1r_2(\cos\theta_1\cos\theta_2 + \sin\theta_1\sin\theta_2) \\ &\qquad = r_1^2 - 2r_1r_2\cos(\theta_1 - \theta_2) + r_2^2, \end{aligned}$$

so the distance between them is $\sqrt{r_1^2 - 2r_1r_2\cos(\theta_1 - \theta_2) + r_2^2}$.

15. $r^2 = 5 \quad \Leftrightarrow \quad x^2 + y^2 = 5$, a circle of radius $\sqrt{5}$ centered at the origin.

16. $r = 4\sec\theta \quad \Leftrightarrow \quad \dfrac{r}{\sec\theta} = 4 \quad \Leftrightarrow \quad r\cos\theta = 4 \quad \Leftrightarrow \quad x = 4$, a vertical line.

17. $r = 5\cos\theta \quad \Rightarrow \quad r^2 = 5r\cos\theta \quad \Leftrightarrow \quad x^2 + y^2 = 5x \quad \Leftrightarrow \quad x^2 - 5x + \frac{25}{4} + y^2 = \frac{25}{4} \quad \Leftrightarrow \quad \left(x - \frac{5}{2}\right)^2 + y^2 = \frac{25}{4}$, a circle of radius $\frac{5}{2}$ centered at $\left(\frac{5}{2}, 0\right)$. The first two equations are actually equivalent since $r^2 = 5r\cos\theta \quad \Rightarrow$ $r(r - 5\cos\theta) = 0 \quad \Rightarrow \quad r = 0$ or $r = 5\cos\theta$. But $r = 5\cos\theta$ gives the point $r = 0$ (the pole) when $\theta = 0$. Thus, the equation $r = 5\cos\theta$ is equivalent to the compound condition ($r = 0$ or $r = 5\cos\theta$).

18. $\theta = \dfrac{\pi}{3} \quad \Rightarrow \quad \tan\theta = \tan\dfrac{\pi}{3} \quad \Rightarrow \quad \dfrac{y}{x} = \sqrt{3} \quad \Leftrightarrow \quad y = \sqrt{3}\,x$, a line through the origin.

19. $r^2\cos 2\theta = 1 \quad \Leftrightarrow \quad r^2(\cos^2\theta - \sin^2\theta) = 1 \quad \Leftrightarrow \quad (r\cos\theta)^2 - (r\sin\theta)^2 = 1 \quad \Leftrightarrow \quad x^2 - y^2 = 1$, a hyperbola centered at the origin with foci on the x-axis.

20. $r^2\sin 2\theta = 1 \quad \Leftrightarrow \quad r^2(2\sin\theta\,\cos\theta) = 1 \quad \Leftrightarrow \quad 2(r\cos\theta)(r\sin\theta) = 1 \quad \Leftrightarrow \quad 2xy = 1 \quad \Leftrightarrow \quad xy = \frac{1}{2}$, a hyperbola centered at the origin with foci on the line $y = x$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. $x^2 + y^2 = 7 \quad \Rightarrow \quad (r\cos\theta)^2 + (r\sin\theta)^2 = 7 \quad \Rightarrow \quad r^2(\cos^2\theta + \sin^2\theta) = 7 \quad \Rightarrow \quad r^2 = 7 \quad \Rightarrow \quad r = \sqrt{7}.$
Note that $r = -\sqrt{7}$ produces the same curve as $r = \sqrt{7}$.

22. $x = -1 \quad \Rightarrow \quad r\cos\theta = -1 \quad \Rightarrow \quad r = -\dfrac{1}{\cos\theta} = -\sec\theta$

23. $y = \sqrt{3}\,x \quad \Rightarrow \quad \dfrac{y}{x} = \sqrt{3} \quad [x \neq 0] \quad \Rightarrow \quad \tan\theta = \sqrt{3} \quad \Rightarrow \quad \theta = \dfrac{\pi}{3}$ or $\dfrac{4\pi}{3}$ [either incudes the pole]

24. $y = -2x^2 \quad \Rightarrow \quad r\sin\theta = -2(r\cos\theta)^2 \quad \Rightarrow \quad r\sin\theta + 2r^2\cos^2\theta = 0 \quad \Rightarrow \quad r(\sin\theta + 2r\cos^2\theta) = 0 \quad \Rightarrow$
$r = 0$ or $r = -\dfrac{\sin\theta}{2\cos^2\theta} = -\dfrac{1}{2}\tan\theta\sec\theta$. $\quad r = 0$ is included in $r = -\dfrac{1}{2}\tan\theta\sec\theta$ when $\theta = 0$, so the curve is represented by the single equation $r = -\frac{1}{2}\tan\theta\sec\theta$.

25. $x^2 + y^2 = 4y \quad \Rightarrow \quad r^2 = 4r\sin\theta \quad \Rightarrow \quad r^2 - 4r\sin\theta = 0 \quad \Rightarrow \quad r(r - 4\sin\theta) = 0 \quad \Rightarrow \quad r = 0$ or $r = 4\sin\theta$.
$r = 0$ is included in $r = 4\sin\theta$ when $\theta = 0$, so the curve is represented by the single equation $r = 4\sin\theta$.

26. $x^2 - y^2 = 4 \quad \Leftrightarrow \quad (r\cos\theta)^2 - (r\sin\theta)^2 = 4 \quad \Leftrightarrow \quad r^2\cos^2\theta - r^2\sin^2\theta = 4 \quad \Leftrightarrow \quad r^2(\cos^2\theta - \sin^2\theta) = 4 \quad \Leftrightarrow$
$r^2\cos 2\theta = 4$

27. (a) The description leads immediately to the polar equation $\theta = \frac{\pi}{6}$, and the Cartesian equation $y = \tan\left(\frac{\pi}{6}\right)x = \frac{1}{\sqrt{3}}\,x$ is slightly more difficult to derive.

(b) The easier description here is the Cartesian equation $x = 3$.

28. (a) Because its center is not at the origin, it is more easily described by its Cartesian equation, $(x-2)^2 + (y-3)^2 = 5^2$.

(b) This circle is more easily given in polar coordinates: $r = 4$. The Cartesian equation is also simple: $x^2 + y^2 = 16$.

29. For $\theta = 0$, π, and 2π, r has its minimum value of about 0.5. For $\theta = \frac{\pi}{2}$ and $\frac{3\pi}{2}$, r attains its maximum value of 2. We see that the graph has a similar shape for $0 \le \theta \le \pi$ and $\pi \le \theta \le 2\pi$.

30. For $\theta = 0$, π, and 2π, r has its maximum value of 2. For $\theta = \frac{\pi}{2}$ and $\frac{3\pi}{2}$, r has its minimum value of about -0.7.
Also, $r = 0$ for what appears to be $\theta = \frac{\pi}{3}$, $\frac{2\pi}{3}$, $\frac{4\pi}{3}$, and $\frac{5\pi}{3}$, so the curve passes through the pole at these angles. r is negative in the intervals $\left[\frac{\pi}{3}, \frac{2\pi}{3}\right]$ and $\left[\frac{4\pi}{3}, \frac{5\pi}{3}\right]$, so the curve will lie on the opposite side of the pole. The graph has a similar shape for $0 \le \theta \le \pi$ and $\pi \le \theta \le 2\pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. r has a maximum value of approximately 1 slightly before $\theta = \frac{\pi}{4}$ and slightly after $\theta = \frac{7\pi}{4}$. r has a minimum value of -2 when $\theta = \pi$. The graph touches the pole ($r = 0$) when $\theta = 0, \frac{\pi}{2}, \frac{3\pi}{2}$, and 2π. Since r is positive in the θ-intervals $\left(0, \frac{\pi}{2}\right)$ and $\left(\frac{3\pi}{2}, 2\pi\right)$, and negative in the interval $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$, the graph lies entirely in the first and fourth quadrants.

32. r increases from 0 to 1 (local max) in the interval $\left[0, \frac{\pi}{2}\right]$. It then decreases slightly, after which r increases to a maximum of 2 at $\theta = \pi$. The graph is symmetric about $\theta = \pi$, so the polar curve is symmetric about the polar axis.

33. $r = -2\sin\theta$

34. $r = 1 - \cos\theta$

35. $r = 2(1 + \cos\theta)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. $r = 1 + 2\cos\theta$

37. $r = \theta, \quad \theta \geq 0$

38. $r = \theta^2, \ -2\pi \leq \theta \leq 2\pi$

39. $r = 3\cos 3\theta$

40. $r = -\sin 5\theta$

41. $r = 2\cos 4\theta$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. $r = 2\sin 6\theta$

43. $r = 1 + 3\cos\theta$

44. $r = 1 + 5\sin\theta$

45. $r^2 = 9\sin 2\theta$

46. $r^2 = \cos 4\theta$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $r = 2 + \sin 3\theta$

48. $r^2\theta = 1 \quad \Leftrightarrow \quad r = \pm 1/\sqrt{\theta}$ for $\theta > 0$

49. $r = \sin(\theta/2)$

50. $r = \cos(\theta/3)$

51. $x = r\cos\theta = (4 + 2\sec\theta)\cos\theta = 4\cos\theta + 2$. Now, $r \to \infty \quad \Rightarrow$ $(4 + 2\sec\theta) \to \infty \quad \Rightarrow \quad \theta \to \left(\frac{\pi}{2}\right)^-$ or $\theta \to \left(\frac{3\pi}{2}\right)^+$ [since we need only consider $0 \le \theta < 2\pi$], so $\lim\limits_{r\to\infty} x = \lim\limits_{\theta\to\pi/2^-}(4\cos\theta + 2) = 2$. Also, $r \to -\infty \quad \Rightarrow \quad (4 + 2\sec\theta) \to -\infty \quad \Rightarrow \quad \theta \to \left(\frac{\pi}{2}\right)^+$ or $\theta \to \left(\frac{3\pi}{2}\right)^-$, so $\lim\limits_{r\to-\infty} x = \lim\limits_{\theta\to\pi/2^+}(4\cos\theta + 2) = 2$. Therefore, $\lim\limits_{r\to\pm\infty} x = 2 \quad \Rightarrow \quad x = 2$ is a vertical asymptote.

52. $y = r\sin\theta = (2 - \csc\theta)\cdot\sin\theta = 2\sin\theta - 1$. $r \to \infty \quad \Rightarrow \quad (2 - \csc\theta) \to \infty \quad \Rightarrow \quad \csc\theta \to -\infty \quad \Rightarrow \quad \theta \to \pi^+$ [since we need only consider $0 \le \theta < 2\pi$] and so $\lim\limits_{r\to\infty} y = \lim\limits_{\theta\to\pi^+} 2\sin\theta - 1 = -1$. Also $r \to -\infty \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$(2 - \csc\theta) \to -\infty \quad \Rightarrow \quad \csc\theta \to \infty \quad \Rightarrow \quad \theta \to \pi^-$ and so $\lim\limits_{r\to-\infty} x = \lim\limits_{\theta\to\pi^-} 2\sin\theta - 1 = -1$. Therefore $\lim\limits_{r\to\pm\infty} y = -1 \quad \Rightarrow \quad y = -1$ is a horizontal asymptote.

53. To show that $x = 1$ is an asymptote, we must prove $\lim\limits_{r\to\pm\infty} x = 1$.

$x = (r)\cos\theta = (\sin\theta\,\tan\theta)\cos\theta = \sin^2\theta$. Now, $r \to \infty \quad \Rightarrow \quad \sin\theta\,\tan\theta \to \infty \quad \Rightarrow$ $\theta \to \left(\frac{\pi}{2}\right)^-$, so $\lim\limits_{r\to\infty} x = \lim\limits_{\theta\to\pi/2^-} \sin^2\theta = 1$. Also, $r \to -\infty \quad \Rightarrow \quad \sin\theta\,\tan\theta \to -\infty \quad \Rightarrow$ $\theta \to \left(\frac{\pi}{2}\right)^+$, so $\lim\limits_{r\to-\infty} x = \lim\limits_{\theta\to\pi/2^+} \sin^2\theta = 1$. Therefore, $\lim\limits_{r\to\pm\infty} x = 1 \quad \Rightarrow \quad x = 1$ is a vertical asymptote. Also notice that $x = \sin^2\theta \geq 0$ for all θ, and $x = \sin^2\theta \leq 1$ for all θ. And $x \neq 1$, since the curve is not defined at odd multiples of $\frac{\pi}{2}$. Therefore, the curve lies entirely within the vertical strip $0 \leq x < 1$.

54. The equation is $(x^2 + y^2)^3 = 4x^2y^2$, but using polar coordinates we know that $x^2 + y^2 = r^2$ and $x = r\cos\theta$ and $y = r\sin\theta$. Substituting into the given equation: $r^6 = 4r^2\cos^2\theta\, r^2\sin^2\theta \quad \Rightarrow \quad r^2 = 4\cos^2\theta\,\sin^2\theta \quad \Rightarrow$ $r = \pm 2\cos\theta\,\sin\theta = \pm\sin 2\theta$. $r = \pm\sin 2\theta$ is sketched at right.

55. (a) We see that the curve $r = 1 + c\sin\theta$ crosses itself at the origin, where $r = 0$ (in fact the inner loop corresponds to negative r-values,) so we solve the equation of the limaçon for $r = 0 \quad \Leftrightarrow \quad c\sin\theta = -1 \quad \Leftrightarrow \quad \sin\theta = -1/c$. Now if $|c| < 1$, then this equation has no solution and hence there is no inner loop. But if $c < -1$, then on the interval $(0, 2\pi)$ the equation has the two solutions $\theta = \sin^{-1}(-1/c)$ and $\theta = \pi - \sin^{-1}(-1/c)$, and if $c > 1$, the solutions are $\theta = \pi + \sin^{-1}(1/c)$ and $\theta = 2\pi - \sin^{-1}(1/c)$. In each case, $r < 0$ for θ between the two solutions, indicating a loop.

(b) For $0 < c < 1$, the dimple (if it exists) is characterized by the fact that y has a local maximum at $\theta = \frac{3\pi}{2}$. So we determine for what c-values $\dfrac{d^2y}{d\theta^2}$ is negative at $\theta = \frac{3\pi}{2}$, since by the Second Derivative Test this indicates a maximum:

$$y = r\sin\theta = \sin\theta + c\sin^2\theta \quad \Rightarrow \quad \frac{dy}{d\theta} = \cos\theta + 2c\sin\theta\,\cos\theta = \cos\theta + c\sin 2\theta \quad \Rightarrow \quad \frac{d^2y}{d\theta^2} = -\sin\theta + 2c\cos 2\theta.$$

At $\theta = \frac{3\pi}{2}$, this is equal to $-(-1) + 2c(-1) = 1 - 2c$, which is negative only for $c > \frac{1}{2}$. A similar argument shows that for $-1 < c < 0$, y only has a local minimum at $\theta = \frac{\pi}{2}$ (indicating a dimple) for $c < -\frac{1}{2}$.

56. (a) The graph of $r = \cos 3\theta$ is a three-leaved rose, which is graph II.

(b) $r = \ln\theta,\ 1 \leq \theta \leq 6\pi$. r increases as θ increases and there are almost three full revolutions. The graph must be either III or VI. As θ increases, r grows slowly in VI and quickly in III. Since $r = \ln\theta$ grows slowly, its graph must be VI.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $r = \cos(\theta/2)$. For $\theta = 0$, $r = 1$, and as θ increases to π, r decreases to 0. Only graph V satisfies those values.

(d) $r = \cos(\theta/3)$. For $\theta = 0$, $r = 1$ and as θ increases to $3\pi/2$, $\theta/3 \to \pi/2$, and r decreases to 0. Only graph IX satisfies those values.

(e) $r = \sec(\theta/3)$. The secant function is never equal to zero, so this graph never intersects the pole. As $\theta \to 3\pi/2$, $\theta/3 \to \pi/2$, and $r \to \infty$. Only graph VII has these properties.

(f) $r = \sec\theta \quad \Rightarrow \quad r\cos\theta = 1 \quad \Rightarrow \quad x = 1$. This is the graph of a vertical line, so it must be graph VIII.

(g) $r = \theta^2,\ 0 \le \theta \le 8\pi$. See part (b). This is graph III.

(h) Since $-1 \le \cos 3\theta \le 1$, $1 \le 2 + \cos 3\theta \le 3$, so $r = 2 + \cos 3\theta$ is never 0; that is, the curve never intersects the pole. The graph must be I or IV. For $0 \le \theta \le 2\pi$, the graph assumes its minimum r-value of 1 three times, at $\theta = \frac{\pi}{3}$, π, and $\frac{5\pi}{3}$, so it must be graph IV.

(i) $r = 2 + \cos(3\theta/2)$. As in part (h), this graph never intersects the pole, so it must be graph I.

57. $r = a\sin\theta + b\cos\theta \quad \Rightarrow \quad r^2 = ar\sin\theta + br\cos\theta \quad \Rightarrow \quad x^2 + y^2 = ay + bx \quad \Rightarrow$
$x^2 - bx + \left(\frac{1}{2}b\right)^2 + y^2 - ay + \left(\frac{1}{2}a\right)^2 = \left(\frac{1}{2}b\right)^2 + \left(\frac{1}{2}a\right)^2 \quad \Rightarrow \quad \left(x - \frac{1}{2}b\right)^2 + \left(y - \frac{1}{2}a\right)^2 = \frac{1}{4}(a^2 + b^2)$, and this is a circle with center $\left(\frac{1}{2}b, \frac{1}{2}a\right)$ and radius $\frac{1}{2}\sqrt{a^2 + b^2}$.

58. These curves are circles which intersect at the origin and at $\left(\frac{1}{\sqrt{2}}a, \frac{\pi}{4}\right)$. At the origin, the first circle has a horizontal tangent and the second a vertical one, so the tangents are perpendicular here. For the first circle $[r = a\sin\theta]$, $dy/d\theta = a\cos\theta\,\sin\theta + a\sin\theta\,\cos\theta = a\sin 2\theta = a$ at $\theta = \frac{\pi}{4}$ and $dx/d\theta = a\cos^2\theta - a\sin^2\theta = a\cos 2\theta = 0$ at $\theta = \frac{\pi}{4}$, so the tangent here is vertical. Similarly, for the second circle $[r = a\cos\theta]$, $dy/d\theta = a\cos 2\theta = 0$ and $dx/d\theta = -a\sin 2\theta = -a$ at $\theta = \frac{\pi}{4}$, so the tangent is horizontal, and again the tangents are perpendicular.

59. $r = 1 + 2\sin(\theta/2)$. The parameter interval is $[0, 4\pi]$.

60. $r = \sqrt{1 - 0.8\sin^2\theta}$. The parameter interval is $[0, 2\pi]$.

61. $r = e^{\sin\theta} - 2\cos(4\theta)$.

The parameter interval is $[0, 2\pi]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

62. $r = |\tan\theta|^{|\cot\theta|}$. The parameter interval $[0, \pi]$ produces the heart-shaped valentine curve shown in the first window.

The complete curve, including the reflected heart, is produced by the parameter interval $[0, 2\pi]$, but perhaps you'll agree that the first curve is more appropriate.

63. $r = 1 + \cos^{999}\theta$. The parameter interval is $[0, 2\pi]$.

64. $r = 2 + \cos(9\theta/4)$. The parameter interval is $[0, 8\pi]$.

65. It appears that the graph of $r = 1 + \sin\left(\theta - \frac{\pi}{6}\right)$ is the same shape as the graph of $r = 1 + \sin\theta$, but rotated counterclockwise about the origin by $\frac{\pi}{6}$. Similarly, the graph of $r = 1 + \sin\left(\theta - \frac{\pi}{3}\right)$ is rotated by $\frac{\pi}{3}$. In general, the graph of $r = f(\theta - \alpha)$ is the same shape as that of $r = f(\theta)$, but rotated counterclockwise through α about the origin. That is, for any point (r_0, θ_0) on the curve $r = f(\theta)$, the point $(r_0, \theta_0 + \alpha)$ is on the curve $r = f(\theta - \alpha)$, since $r_0 = f(\theta_0) = f((\theta_0 + \alpha) - \alpha)$.

66.

From the graph, the highest points seem to have $y \approx 0.77$. To find the exact value, we solve $dy/d\theta = 0$. $y = r\sin\theta = \sin\theta\,\sin 2\theta \quad \Rightarrow$

$$\begin{aligned} dy/d\theta &= 2\sin\theta\,\cos 2\theta + \cos\theta\,\sin 2\theta \\ &= 2\sin\theta\,(2\cos^2\theta - 1) + \cos\theta\,(2\sin\theta\,\cos\theta) \\ &= 2\sin\theta\,(3\cos^2\theta - 1) \end{aligned}$$

In the first quadrant, this is 0 when $\cos\theta = \frac{1}{\sqrt{3}} \quad \Leftrightarrow \quad \sin\theta = \sqrt{\frac{2}{3}} \quad \Leftrightarrow$ $y = 2\sin^2\theta\cos\theta = 2 \cdot \frac{2}{3} \cdot \frac{1}{\sqrt{3}} = \frac{4}{9}\sqrt{3} \approx 0.77$.

67. Consider curves with polar equation $r = 1 + c\cos\theta$, where c is a real number. If $c = 0$, we get a circle of radius 1 centered at the pole. For $0 < c \le 0.5$, the curve gets slightly larger, moves right, and flattens out a bit on the left side. For $0.5 < c < 1$, the left side has a dimple shape. For $c = 1$, the dimple becomes a cusp. For $c > 1$, there is an internal loop. For $c \ge 0$, the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

rightmost point on the curve is $(1+c, 0)$. For $c < 0$, the curves are reflections through the vertical axis of the curves with $c > 0$.

$c = 0.25$ $c = 0.75$ $c = 1$ $c = 2$

68. Consider the polar curves $r = 1 + \cos^n \theta$, where n is a positive integer. First, let **n be an even positive integer**. The first figure shows that the curve has a peanut shape for $n = 2$, but as n increases, the ends are squeezed. As n becomes large, the curves look more and more like the unit circle, but with spikes to the points $(2, 0)$ and $(2, \pi)$.

The second figure shows r as a function of θ in Cartesian coordinates for the same values of n. We can see that for large n, the graph is similar to the graph of $y = 1$, but with spikes to $y = 2$ for $x = 0$, π, and 2π. (Note that when $0 < \cos\theta < 1$, $\cos^{1000}\theta$ is very small.)

Next, let **n be an odd positive integer**. The third figure shows that the curve is a cardioid for $n = 1$, but as n increases, the heart shape becomes more pronounced. As n becomes large, the curves again look more like the unit circle, but with an outward spike to $(2, 0)$ and an inward spike to $(0, \pi)$.

The fourth figure shows r as a function of θ in Cartesian coordinates for the same values of n. We can see that for large n, the graph is similar to the graph of $y = 1$, but spikes to $y = 2$ for $x = 0$ and π, and to $y = 0$ for $x = \pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

DISCOVERY PROJECT Families of Polar Curves

1. (a) $r = \sin n\theta$.

$n = 2$ $n = 3$ $n = 4$ $n = 5$

From the graphs, it seems that when n is even, the number of loops in the curve (called a rose) is $2n$, and when n is odd, the number of loops is simply n. This is because in the case of n odd, every point on the graph is traversed twice, due to the fact that

$$r(\theta + \pi) = \sin[n(\theta + \pi)] = \sin n\theta \cos n\pi + \cos n\theta \sin n\pi = \begin{cases} \sin n\theta & \text{if } n \text{ is even} \\ -\sin n\theta & \text{if } n \text{ is odd} \end{cases}$$

(b) The graph of $r = |\sin n\theta|$ has $2n$ loops whether n is odd or even, since $r(\theta + \pi) = r(\theta)$.

$n = 2$ $n = 3$ $n = 4$ $n = 5$

2. $r = 1 + c \sin n\theta$. We vary n while keeping c constant at 2. As n changes, the curves change in the same way as those in Exercise 1: the number of loops increases. Note that if n is even, the smaller loops are outside the larger ones; if n is odd, they are inside.

$\boldsymbol{c = 2}$

$n = 2$ $n = 3$ $n = 4$ $n = 5$

Now we vary c while keeping $n = 3$. As c increases toward 0, the entire graph gets smaller (the graphs below are not to scale) and the smaller loops shrink in relation to the large ones. At $c = -1$, the small loops disappear entirely, and for $-1 < c < 1$, the graph is a simple, closed curve (at $c = 0$ it is a circle). As c continues to increase, the same changes are seen, but in reverse order, since $1 + (-c)\sin n\theta = 1 + c \sin n(\theta + \pi)$, so the graph for $c = c_0$ is the same as that for $c = -c_0$, with a rotation

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

through π. As $c \to \infty$, the smaller loops get relatively closer in size to the large ones. Note that the distance between the outermost points of corresponding inner and outer loops is always 2. Maple's `animate` command (or Mathematica's `Animate`) is very useful for seeing the changes that occur as c varies.

$\boldsymbol{n = 3}$

3. $r = \dfrac{1 - a\cos\theta}{1 + a\cos\theta}$. We start with $a = 0$, since in this case the curve is simply the circle $r = 1$.

As a increases, the graph moves to the left, and its right side becomes flattened. As a increases through about 0.4, the right side seems to grow a dimple, which upon closer investigation (with narrower θ-ranges) seems to appear at $a \approx 0.42$ [the actual value is $\sqrt{2} - 1$]. As $a \to 1$, this dimple becomes more pronounced, and the curve begins to stretch out horizontally, until at $a = 1$ the denominator vanishes at $\theta = \pi$, and the dimple becomes an actual cusp. For $a > 1$ we must choose our parameter interval carefully, since $r \to \infty$ as $1 + a\cos\theta \to 0 \quad \Leftrightarrow \quad \theta \to \pm\cos^{-1}(-1/a)$. As a increases from 1, the curve splits into two parts. The left part has a loop, which grows larger as a increases, and the right part grows broader vertically, and its left tip develops a dimple when $a \approx 2.42$ [actually, $\sqrt{2} + 1$]. As a increases, the dimple grows more and more pronounced. If $a < 0$, we get the same graph as we do for the corresponding positive a-value, but with a rotation through π about the pole, as happened when c was replaced with $-c$ in Exercise 2.

$a = 0$ $\qquad$ $a = 0.3$ $\qquad$ $a = 0.41, |\theta| \le 0.5$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$a = 0.42, |\theta| \le 0.5$ $\quad$ $a = 0.9, |\theta| \le 0.5$ $\quad$ $a = 1, |\theta| \le 0.1$

$a = 2$ $\quad$ $a = 2.41, |\theta - \pi| \le 0.2$ $\quad$ $a = 4$

$a = 2.42, |\theta - \pi| \le 0.2$

4. Most graphing devices cannot plot implicit polar equations, so we must first find an explicit expression (or expressions) for r in terms of θ, a, and c. We note that the given equation, $r^4 - 2c^2r^2\cos 2\theta + c^4 - a^4 = 0$, is a quadratic in r^2, so we use the quadratic formula and find that

$$r^2 = \frac{2c^2\cos 2\theta \pm \sqrt{4c^4\cos^2 2\theta - 4(c^4 - a^4)}}{2} = c^2\cos 2\theta \pm \sqrt{a^4 - c^4\sin^2 2\theta}$$

so $r = \pm\sqrt{c^2\cos 2\theta \pm \sqrt{a^4 - c^4\sin^2 2\theta}}$. So for each graph, we must plot four curves to be sure of plotting all the points which satisfy the given equation. Note that all four functions have period π.

We start with the case $a = c = 1$, and the resulting curve resembles the symbol for infinity. If we let a decrease, the curve splits into two symmetric parts, and as a decreases further, the parts become smaller, further apart, and rounder. If instead we let a increase from 1, the two lobes of the curve join together, and as a increases further they continue to merge, until at $a \approx 1.4$, the graph no longer has dimples, and has an oval shape. As $a \to \infty$, the oval becomes larger and rounder, since the c^2 and c^4 terms lose their significance. Note that the shape of the graph seems to depend only on the ratio c/a, while the size of the graph varies as c and a jointly increase.

$(a, c) = (1, 1)$ $\quad$ $(a, c) = (0.99, 1)$ $\quad$ $(a, c) = (0.9, 1)$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$(a, c) = (0.6, 1)$ $(a, c) = (1.01, 1)$ $(a, c) = (4.04, 4)$

$(a, c) = (1.3, 1)$ $(a, c) = (1.5, 1)$ $(a, c) = (2, 1)$ $(a, c) = (4, 1)$

10.4 Calculus in Polar Coordinates

1. $r = \sqrt{2\theta},\ 0 \le \theta \le \pi/2.$

$$A = \int_0^{\pi/2} \frac{1}{2} r^2\, d\theta = \int_0^{\pi/2} \frac{1}{2}\left(\sqrt{2\theta}\right)^2 d\theta = \int_0^{\pi/2} \theta\, d\theta = \left[\frac{1}{2}\theta^2\right]_0^{\pi/2} = \frac{1}{2}\left(\frac{\pi}{2}\right)^2 = \frac{\pi^2}{8}$$

2. $r = e^\theta,\ 3\pi/4 \le \theta \le 3\pi/2.$

$$A = \int_{3\pi/4}^{3\pi/2} \tfrac{1}{2} r^2\, d\theta = \int_{3\pi/4}^{3\pi/2} \tfrac{1}{2}(e^\theta)^2\, d\theta = \int_{3\pi/4}^{3\pi/2} \tfrac{1}{2} e^{2\theta}\, d\theta = \tfrac{1}{2}\left[\tfrac{1}{2}e^{2\theta}\right]_{3\pi/4}^{3\pi/2} = \tfrac{1}{4}(e^{3\pi} - e^{3\pi/2})$$

3. $r = \sin\theta + \cos\theta,\ 0 \le \theta \le \pi.$

$$A = \int_0^\pi \frac{1}{2} r^2\, d\theta = \int_0^\pi \frac{1}{2}(\sin\theta + \cos\theta)^2\, d\theta = \int_0^\pi \frac{1}{2}(\sin^2\theta + 2\sin\theta\,\cos\theta + \cos^2\theta)\, d\theta = \int_0^\pi \frac{1}{2}(1 + \sin 2\theta)\, d\theta$$

$$= \tfrac{1}{2}\left[\theta - \tfrac{1}{2}\cos 2\theta\right]_0^\pi = \tfrac{1}{2}\left[\left(\pi - \tfrac{1}{2}\right) - \left(0 - \tfrac{1}{2}\right)\right] = \tfrac{\pi}{2}$$

4. $r = 1/\theta,\ \pi/2 \le \theta \le 2\pi.$

$$A = \int_{\pi/2}^{2\pi} \frac{1}{2} r^2\, d\theta = \int_{\pi/2}^{2\pi} \frac{1}{2}\left(\frac{1}{\theta}\right)^2 d\theta = \int_{\pi/2}^{2\pi} \frac{1}{2}\theta^{-2}\, d\theta = \frac{1}{2}\left[-\frac{1}{\theta}\right]_{\pi/2}^{2\pi}$$

$$= \tfrac{1}{2}\left(-\tfrac{1}{2\pi} + \tfrac{2}{\pi}\right) = \tfrac{1}{2}\left(-\tfrac{1}{2\pi} + \tfrac{4}{2\pi}\right) = \tfrac{3}{4\pi}$$

5. $r^2 = \sin 2\theta,\ 0 \le \theta \le \pi/2.$

$$A = \int_0^{\pi/2} \tfrac{1}{2} r^2\, d\theta = \int_0^{\pi/2} \tfrac{1}{2}\sin 2\theta\, d\theta = \left[-\tfrac{1}{4}\cos 2\theta\right]_0^{\pi/2} = -\tfrac{1}{4}(\cos\pi - \cos 0) = -\tfrac{1}{4}(-1 - 1) = \tfrac{1}{2}$$

6. $r = 2 + \cos\theta,\ \pi/2 \le \theta \le \pi.$

$$A = \int_{\pi/2}^{\pi} \tfrac{1}{2} r^2\, d\theta = \int_{\pi/2}^{\pi} \tfrac{1}{2}(2 + \cos\theta)^2\, d\theta = \int_{\pi/2}^{\pi} \tfrac{1}{2}(4 + 4\cos\theta + \cos^2\theta)\, d\theta = \int_{\pi/2}^{\pi} \tfrac{1}{2}[4 + 4\cos\theta + \tfrac{1}{2}(1 + \cos 2\theta)]\, d\theta$$

$$= \int_{\pi/2}^{\pi} \left(\tfrac{9}{4} + 2\cos\theta + \tfrac{1}{4}\cos 2\theta\right) d\theta = \left[\tfrac{9}{4}\theta + 2\sin\theta + \tfrac{1}{8}\sin 2\theta\right]_{\pi/2}^{\pi} = \left(\tfrac{9\pi}{4} + 0 + 0\right) - \left(\tfrac{9\pi}{8} + 2 + 0\right) = \tfrac{9\pi}{8} - 2$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. $r = 4 + 3\sin\theta,\ -\frac{\pi}{2} \le \theta \le \frac{\pi}{2}$.

$$\begin{aligned} A &= \int_{-\pi/2}^{\pi/2} \tfrac{1}{2}(4 + 3\sin\theta)^2\, d\theta = \tfrac{1}{2}\int_{-\pi/2}^{\pi/2} (16 + 24\sin\theta + 9\sin^2\theta)\, d\theta \\ &= \tfrac{1}{2}\int_{-\pi/2}^{\pi/2} (16 + 9\sin^2\theta)\, d\theta \qquad \text{[by Theorem 5.5.7]} \\ &= \tfrac{1}{2}\cdot 2\int_0^{\pi/2} \left[16 + 9\cdot\tfrac{1}{2}(1 - \cos 2\theta)\right] d\theta \qquad \text{[by Theorem 5.5.7]} \\ &= \int_0^{\pi/2} \left(\tfrac{41}{2} - \tfrac{9}{2}\cos 2\theta\right) d\theta = \left[\tfrac{41}{2}\theta - \tfrac{9}{4}\sin 2\theta\right]_0^{\pi/2} = \left(\tfrac{41\pi}{4} - 0\right) - (0 - 0) = \tfrac{41\pi}{4} \end{aligned}$$

8. $r = \sqrt{\ln\theta},\ 1 \le \theta \le 2\pi$.

$$A = \int_1^{2\pi} \tfrac{1}{2}\left(\sqrt{\ln\theta}\right)^2 d\theta = \int_1^{2\pi} \tfrac{1}{2}\ln\theta\, d\theta = \left[\tfrac{1}{2}\theta\ln\theta\right]_1^{2\pi} - \int_1^{2\pi} \tfrac{1}{2}\, d\theta \qquad \begin{bmatrix} u = \ln\theta, & dv = \frac{1}{2}\, d\theta \\ du = (1/\theta)\, d\theta, & v = \frac{1}{2}\theta \end{bmatrix}$$

$$= [\pi\ln(2\pi) - 0] - \left[\tfrac{1}{2}\theta\right]_1^{2\pi} = \pi\ln(2\pi) - \pi + \tfrac{1}{2}$$

9. The area is bounded by $r = 4\cos\theta$ for $\theta = 0$ to $\theta = \pi$.

$$A = \int_0^{\pi} \tfrac{1}{2}r^2\, d\theta = \int_0^{\pi} \tfrac{1}{2}(4\cos\theta)^2 d\theta = \int_0^{\pi} 8\cos^2\theta\, d\theta$$

$$= 8\int_0^{\pi} \tfrac{1}{2}(1 + \cos 2\theta)\, d\theta = 4\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{\pi} = 4\pi$$

Also, note that this is a circle with radius 2, so its area is $\pi(2)^2 = 4\pi$.

10. $A = \int_0^{2\pi} \tfrac{1}{2}r^2\, d\theta = \int_0^{2\pi} \tfrac{1}{2}(2 + 2\cos\theta)^2 d\theta$

$$= \int_0^{2\pi} \tfrac{1}{2}(4 + 8\cos\theta + 4\cos^2\theta)\, d\theta$$

$$= \int_0^{2\pi} \tfrac{1}{2}\left[4 + 8\cos\theta + 4\cdot\tfrac{1}{2}(1 + \cos 2\theta)\right] d\theta$$

$$= \int_0^{2\pi} (3 + 4\cos\theta + \cos 2\theta)\, d\theta = \left[3\theta + 4\sin\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{2\pi} = 6\pi$$

11. $A = \int_0^{2\pi} \tfrac{1}{2}r^2\, d\theta = \int_0^{2\pi} \tfrac{1}{2}(3 - 2\sin\theta)^2 d\theta$

$$= \tfrac{1}{2}\int_0^{2\pi} (9 - 12\sin\theta + 4\sin^2\theta)\, d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi} \left[9 - 12\sin\theta + 4\cdot\tfrac{1}{2}(1 - \cos 2\theta)\right] d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi} (11 - 12\sin\theta - 2\cos 2\theta)\, d\theta = \tfrac{1}{2}\Big[11\theta + 12\cos\theta - \sin 2\theta\Big]_0^{2\pi}$$

$$= \tfrac{1}{2}[(22\pi + 12) - 12] = 11\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. The area is bounded by $r = 2\sin 3\theta$ for $\theta = 0$ to $\theta = \pi$.

$$A = \int_0^{\pi} \tfrac{1}{2} r^2\, d\theta = \int_0^{\pi} \tfrac{1}{2}(2\sin 3\theta)^2 d\theta = \int_0^{\pi} 2\sin^2 3\theta\, d\theta$$

$$= \int_0^{\pi} 2 \cdot \tfrac{1}{2}(1 - \cos 6\theta)\, d\theta = \left[\theta - \tfrac{1}{6}\sin 6\theta\right]_0^{\pi} = \pi$$

13. $$A = \int_0^{2\pi} \tfrac{1}{2} r^2\, d\theta = \int_0^{2\pi} \tfrac{1}{2}(2 + \sin 4\theta)^2\, d\theta = \tfrac{1}{2}\int_0^{2\pi} (4 + 4\sin 4\theta + \sin^2 4\theta)\, d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi} \left[4 + 4\sin 4\theta + \tfrac{1}{2}(1 - \cos 8\theta)\right] d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi} \left(\tfrac{9}{2} + 4\sin 4\theta - \tfrac{1}{2}\cos 8\theta\right) d\theta = \tfrac{1}{2}\left[\tfrac{9}{2}\theta - \cos 4\theta - \tfrac{1}{16}\sin 8\theta\right]_0^{2\pi}$$

$$= \tfrac{1}{2}[(9\pi - 1) - (-1)] = \tfrac{9}{2}\pi$$

14. $$A = \int_0^{2\pi} \tfrac{1}{2} r^2\, d\theta = \int_0^{2\pi} \tfrac{1}{2}(3 - 2\cos 4\theta)^2\, d\theta = \tfrac{1}{2}\int_0^{2\pi} (9 - 12\cos 4\theta + 4\cos^2 4\theta)\, d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi} \left[9 - 12\cos 4\theta + 4 \cdot \tfrac{1}{2}(1 + \cos 8\theta)\right] d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi} (11 - 12\cos 4\theta + 2\cos 8\theta)\, d\theta = \tfrac{1}{2}\left[11\theta - 3\sin 4\theta + \tfrac{1}{4}\sin 8\theta\right]_0^{2\pi}$$

$$= \tfrac{1}{2}(22\pi) = 11\pi$$

15. $$A = \int_0^{2\pi} \tfrac{1}{2} r^2\, d\theta = \int_0^{2\pi} \tfrac{1}{2}\left(\sqrt{1 + \cos^2 5\theta}\right)^2 d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi} (1 + \cos^2 5\theta)\, d\theta = \tfrac{1}{2}\int_0^{2\pi} \left[1 + \tfrac{1}{2}(1 + \cos 10\theta)\right] d\theta$$

$$= \tfrac{1}{2}\left[\tfrac{3}{2}\theta + \tfrac{1}{20}\sin 10\theta\right]_0^{2\pi} = \tfrac{1}{2}(3\pi) = \tfrac{3}{2}\pi$$

16. $$A = \int_0^{2\pi} \tfrac{1}{2} r^2\, d\theta = \int_0^{2\pi} \tfrac{1}{2}(1 + 5\sin 6\theta)^2\, d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi} (1 + 10\sin 6\theta + 25\sin^2 6\theta)\, d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi} \left[1 + 10\sin 6\theta + 25 \cdot \tfrac{1}{2}(1 - \cos 12\theta)\right] d\theta$$

$$= \tfrac{1}{2}\int_0^{2\pi} \left[\tfrac{27}{2} + 10\sin 6\theta - \tfrac{25}{2}\cos 12\theta\right] d\theta = \tfrac{1}{2}\left[\tfrac{27}{2}\theta - \tfrac{5}{3}\cos 6\theta - \tfrac{25}{24}\sin 12\theta\right]_0^{2\pi}$$

$$= \tfrac{1}{2}\left[\left(27\pi - \tfrac{5}{3}\right) - \left(-\tfrac{5}{3}\right)\right] = \tfrac{27}{2}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. The curve passes through the pole when $r = 0 \;\Rightarrow\; 4\cos 3\theta = 0 \;\Rightarrow\; \cos 3\theta = 0 \;\Rightarrow\; 3\theta = \frac{\pi}{2} + \pi n \;\Rightarrow$ $\theta = \frac{\pi}{6} + \frac{\pi}{3}n$. The part of the shaded loop above the polar axis is traced out for $\theta = 0$ to $\theta = \pi/6$, so we'll use $-\pi/6$ and $\pi/6$ as our limits of integration.

$$A = \int_{-\pi/6}^{\pi/6} \tfrac{1}{2}(4\cos 3\theta)^2\,d\theta = 2\int_0^{\pi/6} \tfrac{1}{2}(16\cos^2 3\theta)\,d\theta$$

$$= 16\int_0^{\pi/6} \tfrac{1}{2}(1+\cos 6\theta)\,d\theta = 8\left[\theta + \tfrac{1}{6}\sin 6\theta\right]_0^{\pi/6} = 8\left(\tfrac{\pi}{6}\right) = \tfrac{4}{3}\pi$$

18. The curve given by $r^2 = 4\cos 2\theta$ passes through the pole when $r = 0 \;\Rightarrow\; 4\cos 2\theta = 0 \;\Rightarrow\; \cos 2\theta = 0 \;\Rightarrow$ $2\theta = \frac{\pi}{2} + \pi n \;\Rightarrow\; \theta = \frac{\pi}{4} + \frac{\pi}{2}n$. The part of the shaded loop above the polar axis is traced out for $\theta = 0$ to $\theta = \pi/4$, so we'll use $-\pi/4$ to $\pi/4$ as our limits of integration.

$$A = \int_{-\pi/4}^{\pi/4} \tfrac{1}{2}(4\cos 2\theta)\,d\theta = 2\int_0^{\pi/4} 2\cos 2\theta\,d\theta = 2\Big[\sin 2\theta\Big]_0^{\pi/4}$$

$$= 2\sin\tfrac{\pi}{2} = 2(1) = 2$$

19. $r = 0 \;\Rightarrow\; \sin 4\theta = 0 \;\Rightarrow\; 4\theta = \pi n \;\Rightarrow\; \theta = \frac{\pi}{4}n.$

$$A = \int_0^{\pi/4} \tfrac{1}{2}(\sin 4\theta)^2\,d\theta = \tfrac{1}{2}\int_0^{\pi/4} \sin^2 4\theta\,d\theta = \tfrac{1}{2}\int_0^{\pi/4} \tfrac{1}{2}(1-\cos 8\theta)\,d\theta$$

$$= \tfrac{1}{4}\left[\theta - \tfrac{1}{8}\sin 8\theta\right]_0^{\pi/4} = \tfrac{1}{4}\left(\tfrac{\pi}{4}\right) = \tfrac{1}{16}\pi$$

20. $r = 0 \;\Rightarrow\; 2\sin 5\theta = 0 \;\Rightarrow\; \sin 5\theta = 0 \;\Rightarrow\; 5\theta = \pi n \;\Rightarrow\; \theta = \frac{\pi}{5}n.$

$$A = \int_0^{\pi/5} \tfrac{1}{2}(2\sin 5\theta)^2\,d\theta = \tfrac{1}{2}\int_0^{\pi/5} 4\sin^2 5\theta\,d\theta$$

$$= 2\int_0^{\pi/5} \tfrac{1}{2}(1-\cos 10\theta)\,d\theta = \left[\theta - \tfrac{1}{10}\sin 10\theta\right]_0^{\pi/5} = \tfrac{\pi}{5}$$

21.

This is a limaçon, with inner loop traced out between $\theta = \frac{7\pi}{6}$ and $\frac{11\pi}{6}$ [found by solving $r = 0$].

$$A = 2\int_{7\pi/6}^{3\pi/2} \tfrac{1}{2}(1+2\sin\theta)^2\,d\theta = \int_{7\pi/6}^{3\pi/2} \left(1 + 4\sin\theta + 4\sin^2\theta\right)d\theta = \int_{7\pi/6}^{3\pi/2} \left[1 + 4\sin\theta + 4\cdot\tfrac{1}{2}(1-\cos 2\theta)\right]d\theta$$

$$= \Big[\theta - 4\cos\theta + 2\theta - \sin 2\theta\Big]_{7\pi/6}^{3\pi/2} = \left(\tfrac{9\pi}{2}\right) - \left(\tfrac{7\pi}{2} + 2\sqrt{3} - \tfrac{\sqrt{3}}{2}\right) = \pi - \tfrac{3\sqrt{3}}{2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. To determine when the strophoid $r = 2\cos\theta - \sec\theta$ passes through the pole, we solve

$$r = 0 \quad\Rightarrow\quad 2\cos\theta - \frac{1}{\cos\theta} = 0 \quad\Rightarrow\quad 2\cos^2\theta - 1 = 0 \quad\Rightarrow\quad \cos^2\theta = \frac{1}{2} \quad\Rightarrow$$

$$\cos\theta = \pm\frac{1}{\sqrt{2}} \quad\Rightarrow\quad \theta = \tfrac{\pi}{4} \text{ or } \theta = \tfrac{3\pi}{4} \text{ for } 0 \le \theta \le \pi \text{ with } \theta \ne \tfrac{\pi}{2}.$$

$$A = 2\int_0^{\pi/4} \tfrac{1}{2}(2\cos\theta - \sec\theta)^2\,d\theta = \int_0^{\pi/4}(4\cos^2\theta - 4 + \sec^2\theta)\,d\theta$$

$$= \int_0^{\pi/4}\left[4\cdot\tfrac{1}{2}(1+\cos 2\theta) - 4 + \sec^2\theta\right]d\theta = \int_0^{\pi/4}(-2 + 2\cos 2\theta + \sec^2\theta)\,d\theta$$

$$= \left[-2\theta + \sin 2\theta + \tan\theta\right]_0^{\pi/4} = \left(-\tfrac{\pi}{2} + 1 + 1\right) - 0 = 2 - \tfrac{\pi}{2}$$

23. $4\sin\theta = 2 \quad\Rightarrow\quad \sin\theta = \frac{1}{2} \quad\Rightarrow\quad \theta = \frac{\pi}{6}$ or $\frac{5\pi}{6} \quad\Rightarrow$

$$A = \int_{\pi/6}^{5\pi/6} \tfrac{1}{2}[(4\sin\theta)^2 - 2^2]\,d\theta = 2\int_{\pi/6}^{\pi/2} \tfrac{1}{2}(16\sin^2\theta - 4)\,d\theta$$

$$= \int_{\pi/6}^{\pi/2}\left[16\cdot\tfrac{1}{2}(1-\cos 2\theta) - 4\right]d\theta = \int_{\pi/6}^{\pi/2}(4 - 8\cos 2\theta)\,d\theta$$

$$= \Big[4\theta - 4\sin 2\theta\Big]_{\pi/6}^{\pi/2} = (2\pi - 0) - \left(\tfrac{2\pi}{3} - 2\sqrt{3}\right) = \tfrac{4\pi}{3} + 2\sqrt{3}$$

24. $1 - \sin\theta = 1 \quad\Rightarrow\quad \sin\theta = 0 \quad\Rightarrow\quad \theta = 0$ or $\pi \quad\Rightarrow$

$$A = \int_\pi^{2\pi} \tfrac{1}{2}\left[(1-\sin\theta)^2 - 1\right]d\theta = \frac{1}{2}\int_\pi^{2\pi}(\sin^2\theta - 2\sin\theta)\,d\theta$$

$$= \tfrac{1}{4}\int_\pi^{2\pi}(1 - \cos 2\theta - 4\sin\theta)\,d\theta = \tfrac{1}{4}\left[\theta - \tfrac{1}{2}\sin 2\theta + 4\cos\theta\right]_\pi^{2\pi}$$

$$= \tfrac{1}{4}\pi + 2$$

25. To find the area inside the leminiscate $r^2 = 8\cos 2\theta$ and outside the circle $r = 2$, we first note that the two curves intersect when $r^2 = 8\cos 2\theta$ and $r = 2$, that is, when $\cos 2\theta = \frac{1}{2}$. For $-\pi < \theta \le \pi$, $\cos 2\theta = \frac{1}{2} \quad\Leftrightarrow\quad 2\theta = \pm\pi/3$ or $\pm 5\pi/3 \quad\Leftrightarrow\quad \theta = \pm\pi/6$ or $\pm 5\pi/6$. The figure shows that the desired area is 4 times the area between the curves from 0 to $\pi/6$. Thus,

$$A = 4\int_0^{\pi/6}\left[\tfrac{1}{2}(8\cos 2\theta) - \tfrac{1}{2}(2)^2\right]d\theta = 8\int_0^{\pi/6}(2\cos 2\theta - 1)\,d\theta$$

$$= 8\Big[\sin 2\theta - \theta\Big]_0^{\pi/6} = 8\left(\sqrt{3}/2 - \pi/6\right) = 4\sqrt{3} - 4\pi/3$$

26. $1 + \cos\theta = 2 - \cos\theta \quad\Rightarrow\quad 2\cos\theta = 1 \quad\Rightarrow\quad \cos\theta = \frac{1}{2} \quad\Rightarrow$

$\theta = \frac{\pi}{3}$ or $\frac{5\pi}{3}$.

$$A = 2\int_0^{\pi/3} \tfrac{1}{2}[(1+\cos\theta)^2 - (2-\cos\theta)^2]\,d\theta \qquad \text{[by symmetry]}$$

$$= \int_0^{\pi/3}(1 + 2\cos\theta + \cos^2\theta - 4 + 4\cos\theta - \cos^2\theta)\,d\theta$$

$$= \int_0^{\pi/3}(6\cos\theta - 3)\,d\theta = \Big[6\sin\theta - 3\theta\Big]_0^{\pi/3} = 6\left(\tfrac{\sqrt{3}}{2}\right) - 3\left(\frac{\pi}{3}\right)$$

$$= 3\sqrt{3} - \pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. $3\cos\theta = 1 + \cos\theta \quad \Leftrightarrow \quad \cos\theta = \frac{1}{2} \quad \Rightarrow \quad \theta = \frac{\pi}{3}$ or $-\frac{\pi}{3}$.

$$A = 2\int_0^{\pi/3} \tfrac{1}{2}[(3\cos\theta)^2 - (1+\cos\theta)^2]\,d\theta$$
$$= \int_0^{\pi/3}(8\cos^2\theta - 2\cos\theta - 1)\,d\theta = \int_0^{\pi/3}[4(1+\cos 2\theta) - 2\cos\theta - 1]\,d\theta$$
$$= \int_0^{\pi/3}(3 + 4\cos 2\theta - 2\cos\theta)\,d\theta = \big[3\theta + 2\sin 2\theta - 2\sin\theta\big]_0^{\pi/3}$$
$$= \pi + \sqrt{3} - \sqrt{3} = \pi$$

28. $3\sin\theta = 2 - \sin\theta \quad \Rightarrow \quad 4\sin\theta = 2 \quad \Rightarrow \quad \sin\theta = \frac{1}{2} \quad \Rightarrow \quad \theta = \frac{\pi}{6}$ or $\frac{5\pi}{6}$.

$$A = 2\int_{\pi/6}^{\pi/2} \tfrac{1}{2}[(3\sin\theta)^2 - (2-\sin\theta)^2]\,d\theta$$
$$= \int_{\pi/6}^{\pi/2}(9\sin^2\theta - 4 + 4\sin\theta - \sin^2\theta]\,d\theta$$
$$= \int_{\pi/6}^{\pi/2}(8\sin^2\theta + 4\sin\theta - 4)\,d\theta$$
$$= 4\int_{\pi/6}^{\pi/2}\big[2\cdot\tfrac{1}{2}(1-\cos 2\theta) + \sin\theta - 1\big]\,d\theta$$
$$= 4\int_{\pi/6}^{\pi/2}(\sin\theta - \cos 2\theta)\,d\theta = 4\big[-\cos\theta - \tfrac{1}{2}\sin 2\theta\big]_{\pi/6}^{\pi/2}$$
$$= 4\Big[(0-0) - \Big(-\tfrac{\sqrt{3}}{2} - \tfrac{\sqrt{3}}{4}\Big)\Big] = 4\Big(\tfrac{3\sqrt{3}}{4}\Big) = 3\sqrt{3}$$

29. $3\sin\theta = 3\cos\theta \quad \Rightarrow \quad \dfrac{3\sin\theta}{3\cos\theta} = 1 \quad \Rightarrow \quad \tan\theta = 1 \quad \Rightarrow \quad \theta = \frac{\pi}{4} \quad \Rightarrow$

$$A = 2\int_0^{\pi/4} \tfrac{1}{2}(3\sin\theta)^2\,d\theta = \int_0^{\pi/4} 9\sin^2\theta\,d\theta = \int_0^{\pi/4} 9\cdot\tfrac{1}{2}(1-\cos 2\theta)\,d\theta$$
$$= \int_0^{\pi/4}\big(\tfrac{9}{2} - \tfrac{9}{2}\cos 2\theta\big)\,d\theta = \Big[\tfrac{9}{2}\theta - \tfrac{9}{4}\sin 2\theta\Big]_0^{\pi/4} = \big(\tfrac{9\pi}{8} - \tfrac{9}{4}\big) - (0-0)$$
$$= \tfrac{9\pi}{8} - \tfrac{9}{4}$$

30. $A = 4\int_0^{\pi/2} \tfrac{1}{2}(1-\cos\theta)^2\,d\theta = 2\int_0^{\pi/2}(1 - 2\cos\theta + \cos^2\theta)\,d\theta$

$$= 2\int_0^{\pi/2}\big[1 - 2\cos\theta + \tfrac{1}{2}(1+\cos 2\theta)\big]\,d\theta$$
$$= 2\int_0^{\pi/2}\big(\tfrac{3}{2} - 2\cos\theta + \tfrac{1}{2}\cos 2\theta\big)\,d\theta = \int_0^{\pi/2}(3 - 4\cos\theta + \cos 2\theta)\,d\theta$$
$$= \big[3\theta - 4\sin\theta + \tfrac{1}{2}\sin 2\theta\big]_0^{\pi/2} = \tfrac{3\pi}{2} - 4$$

31. $\sin 2\theta = \cos 2\theta \quad \Rightarrow \quad \dfrac{\sin 2\theta}{\cos 2\theta} = 1 \quad \Rightarrow \quad \tan 2\theta = 1 \quad \Rightarrow \quad 2\theta = \frac{\pi}{4} \quad \Rightarrow$

$\theta = \frac{\pi}{8} \quad \Rightarrow$

$$A = 8\cdot 2\int_0^{\pi/8} \tfrac{1}{2}\sin^2 2\theta\,d\theta = 8\int_0^{\pi/8} \tfrac{1}{2}(1-\cos 4\theta)\,d\theta$$
$$= 4\Big[\theta - \tfrac{1}{4}\sin 4\theta\Big]_0^{\pi/8} = 4\big(\tfrac{\pi}{8} - \tfrac{1}{4}\cdot 1\big) = \tfrac{\pi}{2} - 1$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. $3 + 2\cos\theta = 3 + 2\sin\theta \quad \Rightarrow \quad \cos\theta = \sin\theta \quad \Rightarrow \quad \theta = \frac{\pi}{4}$ or $\frac{5\pi}{4}$.

$$A = 2\int_{\pi/4}^{5\pi/4} \tfrac{1}{2}(3 + 2\cos\theta)^2\, d\theta = \int_{\pi/4}^{5\pi/4} (9 + 12\cos\theta + 4\cos^2\theta)\, d\theta$$

$$= \int_{\pi/4}^{5\pi/4} \left[9 + 12\cos\theta + 4 \cdot \tfrac{1}{2}(1 + \cos 2\theta)\right] d\theta$$

$$= \int_{\pi/4}^{5\pi/4} (11 + 12\cos\theta + 2\cos 2\theta)\, d\theta = \left[11\theta + 12\sin\theta + \sin 2\theta\right]_{\pi/4}^{5\pi/4}$$

$$= \left(\tfrac{55\pi}{4} - 6\sqrt{2} + 1\right) - \left(\tfrac{11\pi}{4} + 6\sqrt{2} + 1\right) = 11\pi - 12\sqrt{2}$$

33. From the figure, we see that the shaded region is 4 times the shaded region from $\theta = 0$ to $\theta = \pi/4$. $r^2 = 2\sin 2\theta$ and $r = 1 \quad \Rightarrow$

$2\sin 2\theta = 1^2 \quad \Rightarrow \quad \sin 2\theta = \frac{1}{2} \quad \Rightarrow \quad 2\theta = \frac{\pi}{6} \quad \Rightarrow \quad \theta = \frac{\pi}{12}$.

$$A = 4\int_0^{\pi/12} \tfrac{1}{2}(2\sin 2\theta)\, d\theta + 4\int_{\pi/12}^{\pi/4} \tfrac{1}{2}(1)^2\, d\theta$$

$$= \int_0^{\pi/12} 4\sin 2\theta\, d\theta + \int_{\pi/12}^{\pi/4} 2\, d\theta = \Big[-2\cos 2\theta\Big]_0^{\pi/12} + \Big[2\theta\Big]_{\pi/12}^{\pi/4}$$

$$= \left(-\sqrt{3} + 2\right) + \left(\tfrac{\pi}{2} - \tfrac{\pi}{6}\right) = -\sqrt{3} + 2 + \tfrac{\pi}{3}$$

34. $a\sin\theta = b\cos\theta \quad \Rightarrow \quad \dfrac{\sin\theta}{\cos\theta} = \dfrac{b}{a} \quad \Rightarrow \quad \tan\theta = \dfrac{b}{a}$. Let $\alpha = \tan^{-1}(b/a)$.

Then

$$A = \int_0^{\alpha} \tfrac{1}{2}(a\sin\theta)^2\, d\theta + \int_{\alpha}^{\pi/2} \tfrac{1}{2}(b\cos\theta)^2\, d\theta$$

$$= \tfrac{1}{4}a^2\left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{\alpha} + \tfrac{1}{4}b^2\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_{\alpha}^{\pi/2}$$

$$= \tfrac{1}{4}\alpha(a^2 - b^2) + \tfrac{1}{8}\pi b^2 - \tfrac{1}{4}(a^2 + b^2)(\sin\alpha\, \cos\alpha)$$

$$= \tfrac{1}{4}(a^2 - b^2)\, \tan^{-1}(b/a) + \tfrac{1}{8}\pi b^2 - \tfrac{1}{4}ab$$

35. The darker shaded region (from $\theta = 0$ to $\theta = 2\pi/3$) represents $\frac{1}{2}$ of the desired area plus $\frac{1}{2}$ of the area of the inner loop. From this area, we'll subtract $\frac{1}{2}$ of the area of the inner loop (the lighter shaded region from $\theta = 2\pi/3$ to $\theta = \pi$), and then double that difference to obtain the desired area.

$$A = 2\left[\int_0^{2\pi/3} \tfrac{1}{2}\left(\tfrac{1}{2} + \cos\theta\right)^2\, d\theta - \int_{2\pi/3}^{\pi} \tfrac{1}{2}\left(\tfrac{1}{2} + \cos\theta\right)^2\, d\theta\right]$$

$$= \int_0^{2\pi/3} \left(\tfrac{1}{4} + \cos\theta + \cos^2\theta\right) d\theta - \int_{2\pi/3}^{\pi} \left(\tfrac{1}{4} + \cos\theta + \cos^2\theta\right) d\theta$$

$$= \int_0^{2\pi/3} \left[\tfrac{1}{4} + \cos\theta + \tfrac{1}{2}(1 + \cos 2\theta)\right] d\theta$$

$$- \int_{2\pi/3}^{\pi} \left[\tfrac{1}{4} + \cos\theta + \tfrac{1}{2}(1 + \cos 2\theta)\right] d\theta$$

$$= \left[\frac{\theta}{4} + \sin\theta + \frac{\theta}{2} + \frac{\sin 2\theta}{4}\right]_0^{2\pi/3} - \left[\frac{\theta}{4} + \sin\theta + \frac{\theta}{2} + \frac{\sin 2\theta}{4}\right]_{2\pi/3}^{\pi}$$

$$= \left(\tfrac{\pi}{6} + \tfrac{\sqrt{3}}{2} + \tfrac{\pi}{3} - \tfrac{\sqrt{3}}{8}\right) - \left(\tfrac{\pi}{4} + \tfrac{\pi}{2}\right) + \left(\tfrac{\pi}{6} + \tfrac{\sqrt{3}}{2} + \tfrac{\pi}{3} - \tfrac{\sqrt{3}}{8}\right)$$

$$= \tfrac{\pi}{4} + \tfrac{3}{4}\sqrt{3} = \tfrac{1}{4}\left(\pi + 3\sqrt{3}\right)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. $r = 0 \quad \Rightarrow \quad 1 + 2\cos 3\theta = 0 \quad \Rightarrow \quad \cos 3\theta = -\frac{1}{2} \quad \Rightarrow \quad 3\theta = \frac{2\pi}{3}, \frac{4\pi}{3}$ [for $0 \le 3\theta \le 2\pi$] $\quad \Rightarrow \quad \theta = \frac{2\pi}{9}, \frac{4\pi}{9}$. The darker shaded region (from $\theta = 0$ to $\theta = 2\pi/9$) represents $\frac{1}{2}$ of the desired area plus $\frac{1}{2}$ of the area of the inner loop. From this area, we'll subtract $\frac{1}{2}$ of the area of the inner loop (the lighter shaded region from $\theta = 2\pi/9$ to $\theta = \pi/3$), and then double that difference to obtain the desired area.

$$A = 2\left[\int_0^{2\pi/9} \tfrac{1}{2}(1 + 2\cos 3\theta)^2\, d\theta - \int_{2\pi/9}^{\pi/3} \tfrac{1}{2}(1 + 2\cos 3\theta)^2\, d\theta\right]$$

Now
$$r^2 = (1 + 2\cos 3\theta)^2 = 1 + 4\cos 3\theta + 4\cos^2 3\theta = 1 + 4\cos 3\theta + 4 \cdot \tfrac{1}{2}(1 + \cos 6\theta)$$
$$= 1 + 4\cos 3\theta + 2 + 2\cos 6\theta = 3 + 4\cos 3\theta + 2\cos 6\theta$$

and $\int r^2 d\theta = 3\theta + \frac{4}{3}\sin 3\theta + \frac{1}{3}\sin 6\theta + C$, so

$$A = \left[3\theta + \tfrac{4}{3}\sin 3\theta + \tfrac{1}{3}\sin 6\theta\right]_0^{2\pi/9} - \left[3\theta + \tfrac{4}{3}\sin 3\theta + \tfrac{1}{3}\sin 6\theta\right]_{2\pi/9}^{\pi/3}$$
$$= \left[\left(\tfrac{2\pi}{3} + \tfrac{4}{3} \cdot \tfrac{\sqrt{3}}{2} + \tfrac{1}{3} \cdot \tfrac{-\sqrt{3}}{2}\right) - 0\right] - \left[(\pi + 0 + 0) - \left(\tfrac{2\pi}{3} + \tfrac{4}{3} \cdot \tfrac{\sqrt{3}}{2} + \tfrac{1}{3} \cdot \tfrac{-\sqrt{3}}{2}\right)\right]$$
$$= \tfrac{4\pi}{3} + \tfrac{4}{3}\sqrt{3} - \tfrac{1}{3}\sqrt{3} - \pi = \tfrac{\pi}{3} + \sqrt{3}$$

37. The pole is a point of intersection. $\sin\theta = 1 - \sin\theta \quad \Rightarrow \quad 2\sin\theta = 1 \quad \Rightarrow$ $\sin\theta = \frac{1}{2} \quad \Rightarrow \quad \theta = \frac{\pi}{6}$ or $\frac{5\pi}{6}$. So the other points of intersection are $\left(\frac{1}{2}, \frac{\pi}{6}\right)$ and $\left(\frac{1}{2}, \frac{5\pi}{6}\right)$.

38. The pole is a point of intersection. $1 + \cos\theta = 1 - \sin\theta \quad \Rightarrow$ $\cos\theta = -\sin\theta \quad \Rightarrow \quad \dfrac{\cos\theta}{\sin\theta} = -1 \quad \Rightarrow \quad \cot\theta = -1 \quad \Rightarrow \quad \theta = \frac{3\pi}{4}$ or $\frac{7\pi}{4}$. So the other points of intersection are $\left(1 - \frac{1}{2}\sqrt{2}, \frac{3\pi}{4}\right)$ and $\left(1 + \frac{1}{2}\sqrt{2}, \frac{7\pi}{4}\right)$.

39. $2\sin 2\theta = 1 \quad \Rightarrow \quad \sin 2\theta = \frac{1}{2} \quad \Rightarrow \quad 2\theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{13\pi}{6}$, or $\frac{17\pi}{6}$.

By symmetry, the eight points of intersection are given by

$(1, \theta)$, where $\theta = \frac{\pi}{12}, \frac{5\pi}{12}, \frac{13\pi}{12}$, and $\frac{17\pi}{12}$, and

$(-1, \theta)$, where $\theta = \frac{7\pi}{12}, \frac{11\pi}{12}, \frac{19\pi}{12}$, and $\frac{23\pi}{12}$.

[There are many ways to describe these points.]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. The pole is a point of intersection. $\cos\theta = \sin 2\theta = 2\sin\theta\cos\theta \quad \Rightarrow$ $\cos\theta(1 - 2\sin\theta) = 0 \quad \Rightarrow \quad \cos\theta = 0$ or $\sin\theta = \frac{1}{2} \quad \Rightarrow$ $\theta = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{\pi}{6}$, or $\frac{5\pi}{6}$, so the two remaining intersection points are $\left(\frac{\sqrt{3}}{2}, \frac{\pi}{6}\right)$ and $\left(-\frac{\sqrt{3}}{2}, \frac{5\pi}{6}\right)$.

41. $r = 1$ and $r^2 = 2\cos 2\theta \quad \Rightarrow \quad 1^2 = 2\cos 2\theta \quad \Rightarrow \quad \cos 2\theta = \frac{1}{2} \quad \Rightarrow$ $2\theta = \frac{\pi}{3}, \frac{5\pi}{3}, \frac{7\pi}{3}$, or $\frac{11\pi}{3} \quad \Rightarrow \quad \theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}$, or $\frac{11\pi}{6}$. Thus, the four points of intersection are $\left(1, \frac{\pi}{6}\right)$, $\left(1, \frac{5\pi}{6}\right)$, $\left(1, \frac{7\pi}{6}\right)$, and $\left(1, \frac{11\pi}{6}\right)$.

42. Clearly the pole is a point of intersection. $\sin 2\theta = \cos 2\theta \quad \Rightarrow$ $\tan 2\theta = 1 \quad \Rightarrow \quad 2\theta = \frac{\pi}{4} + 2n\pi$ [since $\sin 2\theta$ and $\cos 2\theta$ must be positive in the equations] $\quad \Rightarrow \quad \theta = \frac{\pi}{8} + n\pi \quad \Rightarrow \quad \theta = \frac{\pi}{8}$ or $\frac{9\pi}{8}$. So the curves also intersect at $\left(\frac{1}{\sqrt[4]{2}}, \frac{\pi}{8}\right)$ and $\left(\frac{1}{\sqrt[4]{2}}, \frac{9\pi}{8}\right)$.

43. The shaded region lies outside the rose $r = \sin 2\theta$ and inside the limaçon $r = 3 + 2\cos\theta$, so its area is given by

$$\begin{aligned}
A &= \int_0^{2\pi} \tfrac{1}{2}\left[(3 + 2\cos\theta)^2 - (\sin 2\theta)^2\right] d\theta = \tfrac{1}{2}\int_0^{2\pi} \left(9 + 12\cos\theta + 4\cos^2\theta - \sin^2 2\theta\right) d\theta \\
&= \tfrac{1}{2}\int_0^{2\pi} \left[9 + 12\cos\theta + 4 \cdot \tfrac{1}{2}(1 + \cos 2\theta) - \tfrac{1}{2}(1 - \cos 4\theta)\right] d\theta \\
&= \tfrac{1}{2}\int_0^{2\pi} \left[\tfrac{21}{2} + 12\cos\theta + 2\cos 2\theta + \tfrac{1}{2}\cos 4\theta\right] d\theta \\
&= \tfrac{1}{2}\left[\tfrac{21}{2}\theta + 12\sin\theta + \sin 2\theta + \tfrac{1}{8}\sin 4\theta\right]_0^{2\pi} = \tfrac{1}{2}(21\pi) = \tfrac{21\pi}{2}
\end{aligned}$$

44. $r = \sqrt{2}\cos\theta$ and $r^2 = \sqrt{3}\sin 2\theta \quad \Rightarrow \quad \left(\sqrt{2}\cos\theta\right)^2 = \sqrt{3}\sin 2\theta \quad \Rightarrow \quad 2\cos^2\theta = 2\sqrt{3}\sin\theta\cos\theta \quad \Rightarrow$ $2\cos^2\theta - 2\sqrt{3}\sin\theta\cos\theta = 0 \quad \Rightarrow \quad 2\cos\theta\left(\cos\theta - \sqrt{3}\sin\theta\right) = 0 \quad \Rightarrow \quad \cos\theta = 0$ or $\sqrt{3}\sin\theta = \cos\theta \quad \Rightarrow$ $\tan\theta = \dfrac{1}{\sqrt{3}} \quad \Rightarrow \quad \theta = \frac{\pi}{2}$ or $\frac{\pi}{6}$ for $0 \le \theta \le \frac{\pi}{2}$. The shaded region is comprised of the area swept out by the lemniscate $r^2 = \sqrt{3}\sin 2\theta$ in the interval $0 \le \theta \le \pi/6$ and the portion of the circle $r = \sqrt{2}\cos\theta$ in the interval $\pi/6 \le \theta \le \pi/2$. Thus,

$$\begin{aligned}
A &= \int_0^{\pi/6} \frac{1}{2}\left(\sqrt{3}\sin 2\theta\right) d\theta + \int_{\pi/6}^{\pi/2} \frac{1}{2}\left(\sqrt{2}\cos\theta\right)^2 d\theta = \frac{\sqrt{3}}{2}\left[-\frac{1}{2}\cos 2\theta\right]_0^{\pi/6} + \int_{\pi/6}^{\pi/2} \cos^2\theta\, d\theta \\
&= -\frac{\sqrt{3}}{4}\left(\frac{1}{2} - 1\right) + \int_{\pi/6}^{\pi/2} \frac{1}{2}(1 + \cos 2\theta)\, d\theta = \frac{\sqrt{3}}{8} + \left[\frac{1}{2}\theta + \frac{1}{4}\sin 2\theta\right]_{\pi/6}^{\pi/2} = \frac{\sqrt{3}}{8} + \left[\frac{\pi}{4} - \left(\frac{\pi}{12} + \frac{\sqrt{3}}{8}\right)\right] = \frac{\pi}{6}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. $1+\cos\theta = 3\cos\theta \;\Rightarrow\; 1 = 2\cos\theta \;\Rightarrow\; \cos\theta = \frac{1}{2} \;\Rightarrow\; \theta = \frac{\pi}{3}$. The area swept out by $r = 1+\cos\theta$, $\pi/3 \le \theta \le \pi$, contains the shaded region plus the portion of the circle $r = 3\cos\theta$, $\pi/3 \le \theta \le \pi/2$. Thus, the area of the shaded region is given by

$$\begin{aligned}
A &= \int_{\pi/3}^{\pi} \tfrac{1}{2}(1+\cos\theta)^2\,d\theta - \int_{\pi/3}^{\pi/2} \tfrac{1}{2}(3\cos\theta)^2\,d\theta = \tfrac{1}{2}\int_{\pi/3}^{\pi}(1+2\cos\theta+\cos^2\theta)\,d\theta - \tfrac{9}{2}\int_{\pi/3}^{\pi/2}\cos^2\theta\,d\theta \\
&= \tfrac{1}{2}\int_{\pi/3}^{\pi}\left[1+2\cos\theta+\tfrac{1}{2}(1+\cos 2\theta)\right]d\theta - \tfrac{9}{2}\int_{\pi/3}^{\pi/2}\tfrac{1}{2}(1+\cos 2\theta)\,d\theta \\
&= \tfrac{1}{2}\int_{\pi/3}^{\pi}\left(\tfrac{3}{2}+2\cos\theta+\tfrac{1}{2}\cos 2\theta\right)d\theta - \tfrac{9}{4}\int_{\pi/3}^{\pi/2}(1+\cos 2\theta)\,d\theta \\
&= \tfrac{1}{2}\left[\tfrac{3}{2}\theta+2\sin\theta+\tfrac{1}{4}\sin 2\theta\right]_{\pi/3}^{\pi} - \tfrac{9}{4}\left[\theta+\tfrac{1}{2}\sin 2\theta\right]_{\pi/3}^{\pi/2} \\
&= \tfrac{1}{2}\left[\tfrac{3\pi}{2} - \left(\tfrac{\pi}{2}+\sqrt{3}+\tfrac{\sqrt{3}}{8}\right)\right] - \tfrac{9}{4}\left[\tfrac{\pi}{2} - \left(\tfrac{\pi}{3}+\tfrac{\sqrt{3}}{4}\right)\right] = \tfrac{1}{2}\left(\pi - \tfrac{9\sqrt{3}}{8}\right) - \tfrac{9}{4}\left(\tfrac{\pi}{6} - \tfrac{\sqrt{3}}{4}\right) = \tfrac{\pi}{2} - \tfrac{3\pi}{8} = \tfrac{\pi}{8}
\end{aligned}$$

46. The pole is reached when $r = 1-2\sin\theta = 0 \;\Rightarrow\; \sin\theta = \frac{1}{2} \;\Rightarrow\; \theta = \frac{\pi}{6}, \frac{5\pi}{6}$, or $\frac{13\pi}{6}$. The curve's inner loop is traced from $\theta = \pi/6$ to $\theta = 5\pi/6$ (corresponding to negative r-values), while the outer loop is traced from $\theta = 5\pi/6$ to $\theta = 13\pi/6$. From the figure, we see that the area of the outer loop minus the area of the inner loop gives the area of the shaded region. Thus,

$$\begin{aligned}
A &= \int_{5\pi/6}^{13\pi/6} \tfrac{1}{2}(1-2\sin\theta)^2\,d\theta - \int_{\pi/6}^{5\pi/6} \tfrac{1}{2}(1-2\sin\theta)^2\,d\theta \\
&= \int_{5\pi/6}^{13\pi/6} \tfrac{1}{2}(1-4\sin\theta+4\sin^2\theta)\,d\theta - \int_{\pi/6}^{5\pi/6} \tfrac{1}{2}(1-4\sin\theta+4\sin^2\theta)\,d\theta \\
&= \int_{5\pi/6}^{13\pi/6} \left[\tfrac{1}{2}-2\sin\theta+2\cdot\tfrac{1}{2}(1-\cos 2\theta)\right]d\theta - \int_{\pi/6}^{5\pi/6} \left[\tfrac{1}{2}-2\sin\theta+2\cdot\tfrac{1}{2}(1-\cos 2\theta)\right]d\theta \\
&= \left[\tfrac{1}{2}\theta+2\cos\theta+\theta-\tfrac{1}{2}\sin 2\theta\right]_{5\pi/6}^{13\pi/6} - \left[\tfrac{1}{2}\theta+2\cos\theta+\theta-\tfrac{1}{2}\sin 2\theta\right]_{\pi/6}^{5\pi/6} \\
&= \left(\tfrac{13\pi}{12}+\sqrt{3}+\tfrac{13\pi}{6}-\tfrac{\sqrt{3}}{4}\right) - \left(\tfrac{5\pi}{12}-\sqrt{3}+\tfrac{5\pi}{6}+\tfrac{\sqrt{3}}{4}\right) - \left(\tfrac{5\pi}{12}-\sqrt{3}+\tfrac{5\pi}{6}+\tfrac{\sqrt{3}}{4}\right) + \left(\tfrac{\pi}{12}+\sqrt{3}+\tfrac{\pi}{6}-\tfrac{\sqrt{3}}{4}\right) \\
&= \left(\tfrac{13\pi}{4}+\tfrac{3\sqrt{3}}{4}\right) - 2\left(\tfrac{5\pi}{4}-\tfrac{3\sqrt{3}}{4}\right) + \left(\tfrac{\pi}{4}+\tfrac{3\sqrt{3}}{4}\right) = \pi + 3\sqrt{3}
\end{aligned}$$

47.

From the first graph, we see that the pole is one point of intersection. By zooming in or using the cursor, we find the θ-values of the intersection points to be $\alpha \approx 0.88786 \approx 0.89$ and $\pi - \alpha \approx 2.25$. (The first of these values may be more easily estimated by plotting $y = 1+\sin x$ and $y = 2x$ in rectangular coordinates; see the second graph.) By symmetry, the total

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

area contained is twice the area contained in the first quadrant, that is,

$$A = 2\int_0^{\alpha} \tfrac{1}{2}(2\theta)^2\,d\theta + 2\int_{\alpha}^{\pi/2} \tfrac{1}{2}(1+\sin\theta)^2\,d\theta = \int_0^{\alpha} 4\theta^2\,d\theta + \int_{\alpha}^{\pi/2} \left[1 + 2\sin\theta + \tfrac{1}{2}(1-\cos 2\theta)\right] d\theta$$

$$= \left[\tfrac{4}{3}\theta^3\right]_0^{\alpha} + \left[\theta - 2\cos\theta + \left(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right)\right]_{\alpha}^{\pi/2} = \tfrac{4}{3}\alpha^3 + \left[\left(\tfrac{\pi}{2}+\tfrac{\pi}{4}\right) - \left(\alpha - 2\cos\alpha + \tfrac{1}{2}\alpha - \tfrac{1}{4}\sin 2\alpha\right)\right] \approx 3.4645$$

48.

We need to find the shaded area A in the figure. The horizontal line representing the front of the stage has equation $y = 4 \quad \Leftrightarrow$ $r\sin\theta = 4 \quad \Rightarrow \quad r = 4/\sin\theta$. This line intersects the curve $r = 8 + 8\sin\theta$ when $8 + 8\sin\theta = \dfrac{4}{\sin\theta} \quad \Rightarrow$ $8\sin\theta + 8\sin^2\theta = 4 \quad \Rightarrow \quad 2\sin^2\theta + 2\sin\theta - 1 = 0 \quad \Rightarrow$

$$\sin\theta = \frac{-2 \pm \sqrt{4+8}}{4} = \frac{-2 \pm 2\sqrt{3}}{4} = \frac{-1+\sqrt{3}}{2} \quad \text{[the other value is less than } -1] \quad \Rightarrow \quad \theta = \sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right).$$

This angle is about 21.5° and is denoted by α in the figure.

$$A = 2\textstyle\int_{\alpha}^{\pi/2} \tfrac{1}{2}(8+8\sin\theta)^2\,d\theta - 2\int_{\alpha}^{\pi/2} \tfrac{1}{2}(4\csc\theta)^2\,d\theta = 64\int_{\alpha}^{\pi/2}(1 + 2\sin\theta + \sin^2\theta)\,d\theta - 16\int_{\alpha}^{\pi/2}\csc^2\theta\,d\theta$$

$$= 64\textstyle\int_{\alpha}^{\pi/2}\left(1 + 2\sin\theta + \tfrac{1}{2} - \tfrac{1}{2}\cos 2\theta\right)\,d\theta + 16\int_{\alpha}^{\pi/2}(-\csc^2\theta)\,d\theta = 64\left[\tfrac{3}{2}\theta - 2\cos\theta - \tfrac{1}{4}\sin 2\theta\right]_{\alpha}^{\pi/2} + 16\left[\cot\theta\right]_{\alpha}^{\pi/2}$$

$$= 16\left[6\theta - 8\cos\theta - \sin 2\theta + \cot\theta\right]_{\alpha}^{\pi/a} = 16[(3\pi - 0 - 0 + 0) - (6\alpha - 8\cos\alpha - \sin 2\alpha + \cot\alpha)]$$

$$= 48\pi - 96\alpha + 128\cos\alpha + 16\sin 2\alpha - 16\cot\alpha$$

From the figure, $x^2 + \left(\sqrt{3}-1\right)^2 = 2^2 \quad \Rightarrow \quad x^2 = 4 - \left(3 - 2\sqrt{3} + 1\right) \quad \Rightarrow$ $x^2 = 2\sqrt{3} = \sqrt{12}$, so $x = \sqrt{2\sqrt{3}} = \sqrt[4]{12}$. Using the trigonometric relationships for a right triangle and the identity $\sin 2\alpha = 2\sin\alpha\,\cos\alpha$, we continue:

$$A = 48\pi - 96\alpha + 128\cdot\frac{\sqrt[4]{12}}{2} + 16\cdot 2\cdot\frac{\sqrt{3}-1}{2}\cdot\frac{\sqrt[4]{12}}{2} - 16\cdot\frac{\sqrt[4]{12}}{\sqrt{3}-1}\cdot\frac{\sqrt{3}+1}{\sqrt{3}+1}$$

$$= 48\pi - 96\alpha + 64\sqrt[4]{12} + 8\sqrt[4]{12}\left(\sqrt{3}-1\right) - 8\sqrt[4]{12}\left(\sqrt{3}+1\right) = 48\pi + 48\sqrt[4]{12} - 96\sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right)$$

$$\approx 204.16\ \text{m}^2$$

49. $L = \displaystyle\int_a^b \sqrt{r^2 + (dr/d\theta)^2}\,d\theta = \int_0^{\pi} \sqrt{(2\cos\theta)^2 + (-2\sin\theta)^2}\,d\theta$

$$= \int_0^{\pi} \sqrt{4(\cos^2\theta + \sin^2\theta)}\,d\theta = \int_0^{\pi} \sqrt{4}\,d\theta = \left[2\theta\right]_0^{\pi} = 2\pi$$

As a check, note that the curve is a circle of radius 1, so its circumference is $2\pi(1) = 2\pi$.

50. $L = \displaystyle\int_0^{\pi/2} \sqrt{r^2 + (dr/d\theta)^2}\,d\theta = \int_0^{\pi/2} \sqrt{(e^{\theta/2})^2 + \left(\tfrac{1}{2}e^{\theta/2}\right)^2}\,d\theta = \int_0^{\pi/2} \sqrt{(e^{\theta/2})^2\left(1 + \tfrac{1}{4}\right)}\,d\theta$

$$= \sqrt{\frac{5}{4}}\int_0^{\pi/2} \left|e^{\theta/2}\right| d\theta = \frac{\sqrt{5}}{2}\int_0^{\pi/2} e^{\theta/2}\,d\theta = \frac{\sqrt{5}}{2}\left[2e^{\theta/2}\right]_0^{\pi/2} = \sqrt{5}\left(e^{\pi/4} - 1\right)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. $L = \int_a^b \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_0^{2\pi} \sqrt{(\theta^2)^2 + (2\theta)^2}\, d\theta = \int_0^{2\pi} \sqrt{\theta^4 + 4\theta^2}\, d\theta$

$= \int_0^{2\pi} \sqrt{\theta^2(\theta^2+4)}\, d\theta = \int_0^{2\pi} \theta\sqrt{\theta^2+4}\, d\theta$

Now let $u = \theta^2 + 4$, so that $du = 2\theta\, d\theta \quad \left[\theta\, d\theta = \frac{1}{2}\, du\right]$ and

$$\int_0^{2\pi} \theta\sqrt{\theta^2+4}\, d\theta = \int_4^{4\pi^2+4} \tfrac{1}{2}\sqrt{u}\, du = \tfrac{1}{2}\cdot\tfrac{2}{3}\left[u^{3/2}\right]_4^{4(\pi^2+1)} = \tfrac{1}{3}[4^{3/2}(\pi^2+1)^{3/2} - 4^{3/2}] = \tfrac{8}{3}[(\pi^2+1)^{3/2} - 1]$$

52. $L = \int_a^b \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_0^{2\pi} \sqrt{[2(1+\cos\theta)]^2 + (-2\sin\theta)^2}\, d\theta = \int_0^{2\pi} \sqrt{4 + 8\cos\theta + 4\cos^2\theta + 4\sin^2\theta}\, d\theta$

$= \int_0^{2\pi} \sqrt{8 + 8\cos\theta}\, d\theta = \sqrt{8}\int_0^{2\pi} \sqrt{1+\cos\theta}\, d\theta = \sqrt{8}\int_0^{2\pi} \sqrt{2\cdot\tfrac{1}{2}(1+\cos\theta)}\, d\theta$

$= \sqrt{8}\int_0^{2\pi} \sqrt{2\cos^2\dfrac{\theta}{2}}\, d\theta = \sqrt{8}\sqrt{2}\int_0^{2\pi} \left|\cos\dfrac{\theta}{2}\right| d\theta = 4\cdot 2\int_0^{\pi} \cos\dfrac{\theta}{2}\, d\theta$ [by symmetry]

$= 8\left[2\sin\dfrac{\theta}{2}\right]_0^{\pi} = 8(2) = 16$

53. The blue section of the curve $r = 3 + 3\sin\theta$ is traced from $\theta = -\pi/2$ to $\theta = \pi$.

$r^2 + (dr/d\theta)^2 = (3+3\sin\theta)^2 + (3\cos\theta)^2 = 9 + 18\sin\theta + 9\sin^2\theta + 9\cos^2\theta = 18 + 18\sin\theta$

$L = \int_{-\pi/2}^{\pi} \sqrt{18 + 18\sin\theta}\, d\theta = \sqrt{18}\int_{-\pi/2}^{\pi} \sqrt{1+\sin\theta}\, d\theta = \sqrt{18}\int_{-\pi/2}^{\pi} \sqrt{\dfrac{(1+\sin\theta)(1-\sin\theta)}{1-\sin\theta}}\, d\theta$

$= \sqrt{18}\int_{-\pi/2}^{\pi} \sqrt{\dfrac{1-\sin^2\theta}{1-\sin\theta}}\, d\theta = \sqrt{18}\int_{-\pi/2}^{\pi} \sqrt{\dfrac{\cos^2\theta}{1-\sin\theta}}\, d\theta = \sqrt{18}\int_{-\pi/2}^{\pi} \dfrac{|\cos\theta|}{\sqrt{1-\sin\theta}}\, d\theta$

$= \sqrt{18}\left(\int_{-\pi/2}^{\pi/2} \dfrac{\cos\theta}{\sqrt{1-\sin\theta}}\, d\theta + \int_{\pi/2}^{\pi} \dfrac{-\cos\theta}{\sqrt{1-\sin\theta}}\, d\theta\right)$

$\overset{s}{=} \sqrt{18}\left(\int_{-1}^{1} \dfrac{1}{\sqrt{1-u}}\, du - \int_1^0 \dfrac{1}{\sqrt{1-u}}\, du\right) = \sqrt{18}\left(\left[-2(1-u)^{1/2}\right]_{-1}^{1} - \left[-2(1-u)^{1/2}\right]_1^0\right)$

$= \sqrt{18}\left(2\sqrt{2} + 2\right) = 12 + 6\sqrt{2}$

54. The blue section of the curve $r = \theta + 2$ is traced from $\theta = 0$ to $\theta = 3\pi$.

$r^2 + (dr/d\theta)^2 = (\theta+2)^2 + (1)^2 = (\theta+2)^2 + 1$

$L = \int_0^{3\pi} \sqrt{(\theta+2)^2 + 1}\, d\theta$

$= \int_{\tan^{-1}2}^{\tan^{-1}(3\pi+2)} \sqrt{\tan^2 x + 1}\sec^2 x\, dx \quad \begin{bmatrix} \theta + 2 = \tan x, \\ d\theta = \sec^2 x\, dx \end{bmatrix}$

$= \int_{\tan^{-1}2}^{\tan^{-1}(3\pi+2)} \sqrt{\sec^2 x}\, \sec^2 x\, dx = \int_{\tan^{-1}2}^{\tan^{-1}(3\pi+2)} |\sec x|\sec^2 x\, dx$

$= \int_{\tan^{-1}2}^{\tan^{-1}(3\pi+2)} \sec^3 x\, dx = \tfrac{1}{2}\Big[\sec x\tan x + \ln|\sec x + \tan x|\Big]_{\tan^{-1}2}^{\tan^{-1}(3\pi+2)}$ [by Example 7.2.8]

$= \tfrac{1}{2}\left[\sqrt{1+(3\pi+2)^2}\,(3\pi+2) + \ln\left(\sqrt{1+(3\pi+2)^2} + 3\pi + 2\right)\right] - \tfrac{1}{2}\left[\sqrt{5}\,(2) + \ln\left(\sqrt{5}+2\right)\right] \quad [\approx 64.12]$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

55. The curve $r = \cos^4(\theta/4)$ is completely traced with $0 \le \theta \le 4\pi$.

$$\begin{aligned} r^2 + (dr/d\theta)^2 &= [\cos^4(\theta/4)]^2 + \left[4\cos^3(\theta/4)\cdot(-\sin(\theta/4))\cdot\tfrac{1}{4}\right]^2 \\ &= \cos^8(\theta/4) + \cos^6(\theta/4)\sin^2(\theta/4) \\ &= \cos^6(\theta/4)[\cos^2(\theta/4) + \sin^2(\theta/4)] = \cos^6(\theta/4) \end{aligned}$$

$$\begin{aligned} L &= \textstyle\int_0^{4\pi} \sqrt{\cos^6(\theta/4)}\, d\theta = \int_0^{4\pi} \left|\cos^3(\theta/4)\right| d\theta \\ &= 2\textstyle\int_0^{2\pi} \cos^3(\theta/4)\, d\theta \quad [\text{since } \cos^3(\theta/4) \ge 0 \text{ for } 0 \le \theta \le 2\pi] \quad = 8\int_0^{\pi/2} \cos^3 u\, du \quad \left[u = \tfrac{1}{4}\theta\right] \\ &= 8\textstyle\int_0^{\pi/2} (1 - \sin^2 u)\cos u\, du = 8\int_0^1 (1 - x^2)\, dx \qquad \left[\begin{array}{l} x = \sin u, \\ dx = \cos u\, du \end{array}\right] \\ &= 8\left[x - \tfrac{1}{3}x^3\right]_0^1 = 8\left(1 - \tfrac{1}{3}\right) = \tfrac{16}{3} \end{aligned}$$

56. The curve $r = \cos^2(\theta/2)$ is completely traced with $0 \le \theta \le 2\pi$.

$$\begin{aligned} r^2 + (dr/d\theta)^2 &= [\cos^2(\theta/2)]^2 + \left[2\cos(\theta/2)\cdot(-\sin(\theta/2))\cdot\tfrac{1}{2}\right]^2 \\ &= \cos^4(\theta/2) + \cos^2(\theta/2)\sin^2(\theta/2) \\ &= \cos^2(\theta/2)[\cos^2(\theta/2) + \sin^2(\theta/2)] \\ &= \cos^2(\theta/2) \end{aligned}$$

$$\begin{aligned} L &= \textstyle\int_0^{2\pi} \sqrt{\cos^2(\theta/2)}\, d\theta = \int_0^{2\pi} |\cos(\theta/2)|\, d\theta = 2\int_0^{\pi} \cos(\theta/2)\, d\theta \qquad [\text{since } \cos(\theta/2) \ge 0 \text{ for } 0 \le \theta \le \pi] \\ &= 4\textstyle\int_0^{\pi/2} \cos u\, du \quad \left[u = \tfrac{1}{2}\theta\right] \quad = 4\left[\sin u\right]_0^{\pi/2} = 4(1 - 0) = 4 \end{aligned}$$

57. The graph is symmetric about the polar axis and touches the pole when $r = \cos(\theta/5) = 0$, that is, when $\theta/5 = \pi/2$ or $\theta = 5\pi/2$. $|r|$ is a maximum when $\theta = 0$, so the top half of the red section of the curve is traced starting from the polar axis $\theta = 0$ to $\theta = \pi$. Half of the blue section of the curve is then traced from $\theta = \pi$ to $\theta = 5\pi/2$ (the pole), and, by symmetry, the entire blue section is traced from $\theta = \pi$ to $\theta = 5\pi/2 + (5\pi/2 - \pi) = 4\pi$.

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = \left(\cos\frac{\theta}{5}\right)^2 + \left(-\frac{1}{5}\sin\frac{\theta}{5}\right)^2 = \cos^2\left(\frac{\theta}{5}\right) + \frac{1}{25}\sin^2\left(\frac{\theta}{5}\right)$$

Thus, $L = \displaystyle\int_{\pi}^{4\pi} \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_{\pi}^{4\pi} \sqrt{\cos^2\left(\frac{\theta}{5}\right) + \frac{1}{25}\sin^2\left(\frac{\theta}{5}\right)}\, d\theta.$

58. $r = \dfrac{\sin\theta}{\theta}$ is positive and decreasing in the θ-interval $(0, \pi)$, so a portion of the red curve is traced out between $\theta = 0$ and $\theta = \pi/2$, followed by the blue section of the curve from $\theta = \pi/2$ to $\theta = 3\pi/2$.

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = \left(\frac{\sin\theta}{\theta}\right)^2 + \left(\frac{\theta\cdot\cos\theta - \sin\theta\cdot 1}{\theta^2}\right)^2 = \frac{\sin^2\theta}{\theta^2} + \frac{(\theta\cos\theta - \sin\theta)^2}{\theta^4}$$

Thus, $L = \displaystyle\int_{\pi/2}^{3\pi/2} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2}\, d\theta = \int_{\pi/2}^{3\pi/2} \sqrt{\frac{\sin^2\theta}{\theta^2} + \frac{(\theta\cos\theta - \sin\theta)^2}{\theta^4}}\, d\theta.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. One loop of the curve $r = \cos 2\theta$ is traced with $-\pi/4 \le \theta \le \pi/4$.

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = \cos^2 2\theta + (-2\sin 2\theta)^2 = \cos^2 2\theta + 4\sin^2 2\theta = 1 + 3\sin^2 2\theta \quad \Rightarrow$$

$$L = \int_{-\pi/4}^{\pi/4} \sqrt{1 + 3\sin^2 2\theta}\, d\theta \approx 2.4221.$$

60. $r^2 + \left(\dfrac{dr}{d\theta}\right)^2 = \tan^2\theta + (\sec^2\theta)^2 \quad \Rightarrow \quad L = \displaystyle\int_{\pi/6}^{\pi/3} \sqrt{\tan^2\theta + \sec^4\theta}\, d\theta \approx 1.2789$

61. The curve $r = \sin(6\sin\theta)$ is completely traced with $0 \le \theta \le \pi$. $\quad r = \sin(6\sin\theta) \quad \Rightarrow$

$\dfrac{dr}{d\theta} = \cos(6\sin\theta)\cdot 6\cos\theta$, so $r^2 + \left(\dfrac{dr}{d\theta}\right)^2 = \sin^2(6\sin\theta) + 36\cos^2\theta\cos^2(6\sin\theta) \quad \Rightarrow$

$$L = \int_0^{\pi} \sqrt{\sin^2(6\sin\theta) + 36\cos^2\theta\ \cos^2(6\sin\theta)}\, d\theta \approx 8.0091.$$

62. The curve $r = \sin(\theta/4)$ is completely traced with $0 \le \theta \le 8\pi$. $\quad r = \sin(\theta/4) \quad \Rightarrow \quad \dfrac{dr}{d\theta} = \dfrac{1}{4}\cos(\theta/4)$, so

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = \sin^2(\theta/4) + \tfrac{1}{16}\cos^2(\theta/4) \quad \Rightarrow \quad L = \int_0^{8\pi} \sqrt{\sin^2(\theta/4) + \tfrac{1}{16}\cos^2(\theta/4)}\, d\theta \approx 17.1568.$$

63. $r = 2\cos\theta \quad \Rightarrow \quad x = r\cos\theta = 2\cos^2\theta,\ y = r\sin\theta = 2\sin\theta\ \cos\theta = \sin 2\theta \quad \Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{2\cos 2\theta}{2\cdot 2\cos\theta\,(-\sin\theta)} = \frac{\cos 2\theta}{-\sin 2\theta} = -\cot 2\theta$$

When $\theta = \dfrac{\pi}{3}$, $\dfrac{dy}{dx} = -\cot\left(2\cdot\dfrac{\pi}{3}\right) = \cot\dfrac{\pi}{3} = \dfrac{1}{\sqrt{3}}$. [*Another method:* Use Equation 3.]

64. $r = 2 + \sin 3\theta \quad \Rightarrow \quad x = r\cos\theta = (2+\sin 3\theta)\cos\theta,\ y = r\sin\theta = (2+\sin 3\theta)\sin\theta \quad \Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{(2+\sin 3\theta)\cos\theta + \sin\theta(3\cos 3\theta)}{(2+\sin 3\theta)(-\sin\theta) + \cos\theta(3\cos 3\theta)}$$

When $\theta = \dfrac{\pi}{4}$, $\dfrac{dy}{dx} = \dfrac{\left(2+\sin\frac{3\pi}{4}\right)\cos\frac{\pi}{4} + \sin\frac{\pi}{4}\left(3\cos\frac{3\pi}{4}\right)}{\left(2+\sin\frac{3\pi}{4}\right)\left(-\sin\frac{\pi}{4}\right) + \cos\frac{\pi}{4}\left(3\cos\frac{3\pi}{4}\right)} = \dfrac{\left(2+\frac{\sqrt{2}}{2}\right)\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}\cdot 3\left(-\frac{\sqrt{2}}{2}\right)}{\left(2+\frac{\sqrt{2}}{2}\right)\left(-\frac{\sqrt{2}}{2}\right) + \frac{\sqrt{2}}{2}\cdot 3\left(-\frac{\sqrt{2}}{2}\right)}$

$= \dfrac{\sqrt{2} + \frac{1}{2} - \frac{3}{2}}{-\sqrt{2} - \frac{1}{2} - \frac{3}{2}} = \dfrac{\sqrt{2}-1}{-\sqrt{2}-2}$, or, equivalently, $2 - \dfrac{3}{2}\sqrt{2}$.

65. $r = 1/\theta \quad \Rightarrow \quad x = r\cos\theta = (\cos\theta)/\theta,\ y = r\sin\theta = (\sin\theta)/\theta \quad \Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\sin\theta(-1/\theta^2) + (1/\theta)\cos\theta}{\cos\theta(-1/\theta^2) - (1/\theta)\sin\theta}\cdot\frac{\theta^2}{\theta^2} = \frac{-\sin\theta + \theta\cos\theta}{-\cos\theta - \theta\sin\theta}$$

When $\theta = \pi$, $\dfrac{dy}{dx} = \dfrac{-0 + \pi(-1)}{-(-1) - \pi(0)} = \dfrac{-\pi}{1} = -\pi$.

66. $r = \sin\theta + 2\cos\theta \quad \Rightarrow \quad x = r\cos\theta = \sin\theta\cos\theta + 2\cos^2\theta,\ y = r\sin\theta = \sin^2\theta + 2\sin\theta\cos\theta \quad \Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{2\sin\theta\cos\theta - 2\sin^2\theta + 2\cos^2\theta}{-\sin^2\theta + \cos^2\theta - 4\sin\theta\cos\theta}$$

When $\theta = \dfrac{\pi}{2}$, $\dfrac{dy}{dx} = \dfrac{2(1)(0) - 2(1) + 2(0)}{-1 + 0 - 4(1)(0)} = \dfrac{-2}{-1} = 2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. $r = \cos 2\theta \quad \Rightarrow \quad x = r\cos\theta = \cos 2\theta\,\cos\theta,\ y = r\sin\theta = \cos 2\theta\,\sin\theta \quad \Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\cos 2\theta\,\cos\theta + \sin\theta\,(-2\sin 2\theta)}{\cos 2\theta\,(-\sin\theta) + \cos\theta\,(-2\sin 2\theta)}$$

When $\theta = \dfrac{\pi}{4}$, $\dfrac{dy}{dx} = \dfrac{0\big(\sqrt{2}/2\big) + \big(\sqrt{2}/2\big)(-2)}{0\big(-\sqrt{2}/2\big) + \big(\sqrt{2}/2\big)(-2)} = \dfrac{-\sqrt{2}}{-\sqrt{2}} = 1.$

68. $r = 1 + 2\cos\theta \quad \Rightarrow \quad x = r\cos\theta = (1 + 2\cos\theta)\cos\theta,\ y = r\sin\theta = (1 + 2\cos\theta)\sin\theta \quad \Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{(1 + 2\cos\theta)\cos\theta + \sin\theta\,(-2\sin\theta)}{(1 + 2\cos\theta)(-\sin\theta) + \cos\theta\,(-2\sin\theta)}$$

When $\theta = \dfrac{\pi}{3}$, $\dfrac{dy}{dx} = \dfrac{2\big(\frac{1}{2}\big) + \big(\sqrt{3}/2\big)\big(-\sqrt{3}\big)}{2\big(-\sqrt{3}/2\big) + \frac{1}{2}\big(-\sqrt{3}\big)} \cdot \dfrac{2}{2} = \dfrac{2 - 3}{-2\sqrt{3} - \sqrt{3}} = \dfrac{-1}{-3\sqrt{3}} = \dfrac{\sqrt{3}}{9}.$

69. $r = \sin\theta \quad \Rightarrow \quad x = r\cos\theta = \sin\theta\cos\theta,\ y = r\sin\theta = \sin^2\theta \quad \Rightarrow \quad \frac{dy}{d\theta} = 2\sin\theta\cos\theta = \sin 2\theta = 0 \quad \Rightarrow$

$2\theta = 0$ or $\pi \quad \Rightarrow \quad \theta = 0$ or $\frac{\pi}{2} \quad \Rightarrow$ horizontal tangent at $(0, 0)$, and $\big(1,\ \frac{\pi}{2}\big)$.

$\frac{dx}{d\theta} = -\sin^2\theta + \cos^2\theta = \cos 2\theta = 0 \quad \Rightarrow \quad 2\theta = \frac{\pi}{2}$ or $\frac{3\pi}{2} \quad \Rightarrow \quad \theta = \frac{\pi}{4}$ or $\frac{3\pi}{4} \quad \Rightarrow$ vertical tangent at $\left(\frac{1}{\sqrt{2}}, \frac{\pi}{4}\right)$

and $\left(\frac{1}{\sqrt{2}}, \frac{3\pi}{4}\right)$.

70. $r = 1 - \sin\theta \quad \Rightarrow \quad x = r\cos\theta = \cos\theta\,(1 - \sin\theta),\ y = r\sin\theta = \sin\theta\,(1 - \sin\theta) \quad \Rightarrow$

$\frac{dy}{d\theta} = \sin\theta\,(-\cos\theta) + (1 - \sin\theta)\,\cos\theta = \cos\theta\,(1 - 2\sin\theta) = 0 \quad \Rightarrow \quad \cos\theta = 0$ or $\sin\theta = \frac{1}{2} \quad \Rightarrow$

$\theta = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}$, or $\frac{3\pi}{2} \quad \Rightarrow$ horizontal tangent at $\big(\frac{1}{2}, \frac{\pi}{6}\big)$, $\big(\frac{1}{2}, \frac{5\pi}{6}\big)$, and $\big(2, \frac{3\pi}{2}\big)$.

$$\begin{aligned}\frac{dx}{d\theta} &= \cos\theta\,(-\cos\theta) + (1 - \sin\theta)(-\sin\theta) = -\cos^2\theta - \sin\theta + \sin^2\theta = 2\sin^2\theta - \sin\theta - 1\\ &= (2\sin\theta + 1)(\sin\theta - 1) = 0 \quad \Rightarrow\end{aligned}$$

$\sin\theta = -\frac{1}{2}$ or $1 \quad \Rightarrow \quad \theta = \frac{7\pi}{6}, \frac{11\pi}{6}$, or $\frac{\pi}{2} \quad \Rightarrow$ vertical tangent at $\big(\frac{3}{2}, \frac{7\pi}{6}\big)$, $\big(\frac{3}{2}, \frac{11\pi}{6}\big)$, and $\big(0, \frac{\pi}{2}\big)$.

Note that the tangent is vertical, not horizontal, when $\theta = \frac{\pi}{2}$, since

$$\lim_{\theta\to(\pi/2)^-} \frac{dy/d\theta}{dx/d\theta} = \lim_{\theta\to(\pi/2)^-} \frac{\cos\theta\,(1 - 2\sin\theta)}{(2\sin\theta + 1)(\sin\theta - 1)} = \infty \text{ and } \lim_{\theta\to(\pi/2)^+} \frac{dy/d\theta}{dx/d\theta} = -\infty.$$

71. $r = 1 + \cos\theta \quad \Rightarrow \quad x = r\cos\theta = \cos\theta\,(1 + \cos\theta),\ y = r\sin\theta = \sin\theta\,(1 + \cos\theta) \quad \Rightarrow$

$\frac{dy}{d\theta} = (1 + \cos\theta)\,\cos\theta - \sin^2\theta = 2\cos^2\theta + \cos\theta - 1 = (2\cos\theta - 1)(\cos\theta + 1) = 0 \quad \Rightarrow \quad \cos\theta = \frac{1}{2}$ or $-1 \quad \Rightarrow$

$\theta = \frac{\pi}{3}, \pi$, or $\frac{5\pi}{3} \quad \Rightarrow$ horizontal tangent at $\big(\frac{3}{2}, \frac{\pi}{3}\big)$, $(0, \pi)$, and $\big(\frac{3}{2}, \frac{5\pi}{3}\big)$.

$\frac{dx}{d\theta} = -(1 + \cos\theta)\sin\theta - \cos\theta\sin\theta = -\sin\theta\,(1 + 2\cos\theta) = 0 \quad \Rightarrow \quad \sin\theta = 0$ or $\cos\theta = -\frac{1}{2} \quad \Rightarrow$

$\theta = 0, \pi, \frac{2\pi}{3}$, or $\frac{4\pi}{3} \quad \Rightarrow$ vertical tangent at $(2, 0)$, $\big(\frac{1}{2}, \frac{2\pi}{3}\big)$, and $\big(\frac{1}{2}, \frac{4\pi}{3}\big)$.

Note that the tangent is horizontal, not vertical when $\theta = \pi$, since $\displaystyle\lim_{\theta\to\pi} \frac{dy/d\theta}{dx/d\theta} = 0$.

72. $r = e^\theta \quad \Rightarrow \quad x = r\cos\theta = e^\theta\cos\theta,\ y = r\sin\theta = e^\theta\sin\theta \quad \Rightarrow$

$\frac{dy}{d\theta} = e^\theta\sin\theta + e^\theta\cos\theta = e^\theta(\sin\theta + \cos\theta) = 0 \quad \Rightarrow \quad \sin\theta = -\cos\theta \quad \Rightarrow \quad \tan\theta = -1 \quad \Rightarrow$

$\theta = -\frac{1}{4}\pi + n\pi$ [n any integer] $\quad \Rightarrow$ horizontal tangents at $\Big(e^{\pi(n-1/4)}, \pi\big(n - \frac{1}{4}\big)\Big)$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\frac{dx}{d\theta} = e^\theta \cos\theta - e^\theta \sin\theta = e^\theta(\cos\theta - \sin\theta) = 0 \quad \Rightarrow \quad \sin\theta = \cos\theta \quad \Rightarrow \quad \tan\theta = 1 \quad \Rightarrow$

$\theta = \frac{1}{4}\pi + n\pi$ [n any integer] $\Rightarrow$ vertical tangents at $\left(e^{\pi(n+1/4)}, \pi\left(n + \frac{1}{4}\right)\right)$.

73. $$\tan\psi = \tan(\phi - \theta) = \frac{\tan\phi - \tan\theta}{1 + \tan\phi\,\tan\theta} = \frac{\dfrac{dy}{dx} - \tan\theta}{1 + \dfrac{dy}{dx}\tan\theta} = \frac{\dfrac{dy/d\theta}{dx/d\theta} - \tan\theta}{1 + \dfrac{dy/d\theta}{dx/d\theta}\tan\theta}$$

$$= \frac{\dfrac{dy}{d\theta} - \dfrac{dx}{d\theta}\tan\theta}{\dfrac{dx}{d\theta} + \dfrac{dy}{d\theta}\tan\theta} = \frac{\left(\dfrac{dr}{d\theta}\sin\theta + r\cos\theta\right) - \tan\theta\left(\dfrac{dr}{d\theta}\cos\theta - r\sin\theta\right)}{\left(\dfrac{dr}{d\theta}\cos\theta - r\sin\theta\right) + \tan\theta\left(\dfrac{dr}{d\theta}\sin\theta + r\cos\theta\right)} = \frac{r\cos\theta + r\cdot\dfrac{\sin^2\theta}{\cos\theta}}{\dfrac{dr}{d\theta}\cos\theta + \dfrac{dr}{d\theta}\cdot\dfrac{\sin^2\theta}{\cos\theta}}$$

$$= \frac{r\cos^2\theta + r\sin^2\theta}{\dfrac{dr}{d\theta}\cos^2\theta + \dfrac{dr}{d\theta}\sin^2\theta} = \frac{r}{dr/d\theta}$$

74. (a) $r = e^\theta \quad \Rightarrow \quad dr/d\theta = e^\theta$, so by Exercise 73, $\tan\psi = r/e^\theta = 1 \quad \Rightarrow$ $\psi = \arctan 1 = \frac{\pi}{4}$.

(b) The Cartesian equation of the tangent line at $(1, 0)$ is $y = x - 1$, and that of the tangent line at $(0, e^{\pi/2})$ is $y = e^{\pi/2} - x$.

(c) Let a be the tangent of the angle between the tangent and radial lines, that is, $a = \tan\psi$. Then, by Exercise 73, $a = \dfrac{r}{dr/d\theta} \quad \Rightarrow \quad \dfrac{dr}{d\theta} = \dfrac{1}{a}r \quad \Rightarrow$ $r = Ce^{\theta/a}$ (by Theorem 9.4.2).

75. (a) From (10.2.9),

$$S = \int_a^b 2\pi y\sqrt{(dx/d\theta)^2 + (dy/d\theta)^2}\,d\theta$$
$$= \int_a^b 2\pi y\sqrt{r^2 + (dr/d\theta)^2}\,d\theta \qquad \text{[from the derivation of Equation 10.4.6]}$$
$$= \int_a^b 2\pi r\sin\theta\sqrt{r^2 + (dr/d\theta)^2}\,d\theta$$

(b) The curve $r^2 = \cos 2\theta$ goes through the pole when $\cos 2\theta = 0 \quad \Rightarrow$ $2\theta = \frac{\pi}{2} \quad \Rightarrow \quad \theta = \frac{\pi}{4}$. We'll rotate the curve from $\theta = 0$ to $\theta = \frac{\pi}{4}$ and double this value to obtain the total surface area generated.

$r^2 = \cos 2\theta \quad \Rightarrow \quad 2r\dfrac{dr}{d\theta} = -2\sin 2\theta \quad \Rightarrow \quad \left(\dfrac{dr}{d\theta}\right)^2 = \dfrac{\sin^2 2\theta}{r^2} = \dfrac{\sin^2 2\theta}{\cos 2\theta}$.

$$S = 2\int_0^{\pi/4} 2\pi\sqrt{\cos 2\theta}\sin\theta\sqrt{\cos 2\theta + \left(\sin^2 2\theta\right)/\cos 2\theta}\,d\theta = 4\pi\int_0^{\pi/4}\sqrt{\cos 2\theta}\sin\theta\sqrt{\frac{\cos^2 2\theta + \sin^2 2\theta}{\cos 2\theta}}\,d\theta$$
$$= 4\pi\int_0^{\pi/4}\sqrt{\cos 2\theta}\sin\theta\,\frac{1}{\sqrt{\cos 2\theta}}\,d\theta = 4\pi\int_0^{\pi/4}\sin\theta\,d\theta = 4\pi\left[-\cos\theta\right]_0^{\pi/4} = -4\pi\left(\frac{\sqrt{2}}{2} - 1\right) = 2\pi\left(2 - \sqrt{2}\right)$$

76. (a) Rotation around $\theta = \frac{\pi}{2}$ is the same as rotation around the y-axis, that is, $S = \int_a^b 2\pi x\,ds$ where $ds = \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt$ for a parametric equation, and for the special case of a polar equation, $x = r\cos\theta$ and

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$ds = \sqrt{(dx/d\theta)^2 + (dy/d\theta)^2}\, d\theta = \sqrt{r^2 + (dr/d\theta)^2}\, d\theta$ [see the derivation of Equation 10.4.6]. Therefore, for a polar equation rotated around $\theta = \frac{\pi}{2}$, $S = \int_a^b 2\pi r \cos\theta \sqrt{r^2 + (dr/d\theta)^2}\, d\theta$.

(b) As in the solution for Exercise 75(b), we can double the surface area generated by rotating the curve from $\theta = 0$ to $\theta = \frac{\pi}{4}$ to obtain the total surface area.

$$\begin{aligned} S &= 2\int_0^{\pi/4} 2\pi \sqrt{\cos 2\theta}\cos\theta \sqrt{\cos 2\theta + (\sin^2 2\theta)/\cos 2\theta}\, d\theta \\ &= 4\pi \int_0^{\pi/4} \sqrt{\cos 2\theta}\cos\theta \sqrt{\frac{\cos^2 2\theta + \sin^2 2\theta}{\cos 2\theta}}\, d\theta \\ &= 4\pi \int_0^{\pi/4} \sqrt{\cos 2\theta}\cos\theta \frac{1}{\sqrt{\cos 2\theta}}\, d\theta = 4\pi \int_0^{\pi/4} \cos\theta\, d\theta \\ &= 4\pi \big[\sin\theta\big]_0^{\pi/4} = 4\pi\left(\frac{\sqrt{2}}{2} - 0\right) = 2\sqrt{2}\,\pi \end{aligned}$$

10.5 Conic Sections

1. $x^2 = 8y$ and $x^2 = 4py \quad \Rightarrow \quad 4p = 8 \quad \Leftrightarrow \quad p = 2$. The vertex is $(0, 0)$, the focus is $(0, 2)$, and the directrix is $y = -2$.

2. $9x = y^2$, so $4p = 9 \quad \Leftrightarrow \quad p = \frac{9}{4}$. The vertex is $(0, 0)$, the focus is $\left(\frac{9}{4}, 0\right)$, and the directrix is $x = -\frac{9}{4}$.

3. $5x + 3y^2 = 0 \quad \Leftrightarrow \quad y^2 = -\frac{5}{3}x$, so $4p = -\frac{5}{3} \quad \Leftrightarrow \quad p = -\frac{5}{12}$.

The vertex is $(0, 0)$, the focus is $\left(-\frac{5}{12}, 0\right)$, and the directrix is $x = \frac{5}{12}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $x^2 + 12y = 0 \quad \Leftrightarrow \quad x^2 = -12y$, so $4p = -12 \quad \Leftrightarrow \quad p = -3$.

The vertex is $(0, 0)$, the focus is $(0, -3)$, and the directrix is $y = 3$.

5. $(y + 1)^2 = 16(x - 3)$, so $4p = 16 \quad \Leftrightarrow \quad p = 4$. The vertex is $(3, -1)$, the focus is $(3 + 4, -1) = (7, -1)$, and the directrix is $x = 3 - 4 = -1$.

6. $(x - 3)^2 = 8\,(y + 1)$, so $4p = 8 \quad \Leftrightarrow \quad p = 2$. The vertex is $(3, -1)$, the focus is $(3, -1 + 2) = (3, 1)$, and the directrix is $y = -1 - 2 = -3$.

7. $y^2 + 6y + 2x + 1 = 0 \quad \Leftrightarrow \quad y^2 + 6y = -2x - 1 \quad \Leftrightarrow$ $y^2 + 6y + 9 = -2x + 8 \quad \Leftrightarrow \quad (y + 3)^2 = -2(x - 4)$, so $4p = -2 \quad \Leftrightarrow$ $p = -\frac{1}{2}$. The vertex is $(4, -3)$, the focus is $\left(\frac{7}{2}, -3\right)$, and the directrix is $x = \frac{9}{2}$.

8. $2x^2 - 16x - 3y + 38 = 0 \quad \Leftrightarrow \quad 2x^2 - 16x = 3y - 38 \quad \Leftrightarrow$ $2(x^2 - 8x + 16) = 3y - 38 + 32 \quad \Leftrightarrow \quad 2(x - 4)^2 = 3y - 6 \quad \Leftrightarrow$ $(x - 4)^2 = \frac{3}{2}(y - 2)$, so $4p = \frac{3}{2} \quad \Leftrightarrow \quad p = \frac{3}{8}$. The vertex is $(4, 2)$, the focus is $\left(4, \frac{19}{8}\right)$, and the directrix is $y = \frac{13}{8}$.

9. The equation has the form $y^2 = 4px$, where $p < 0$. Since the parabola passes through $(-1, 1)$, we have $1^2 = 4p(-1)$, so $4p = -1$ and an equation is $y^2 = -x$ or $x = -y^2$. $\quad 4p = -1$, so $p = -\frac{1}{4}$ and the focus is $\left(-\frac{1}{4}, 0\right)$ while the directrix is $x = \frac{1}{4}$.

10. The vertex is $(2, -2)$, so the equation is of the form $(x - 2)^2 = 4p(y + 2)$, where $p > 0$. The point $(0, 0)$ is on the parabola, so $4 = 4p(2) \quad \Rightarrow \quad 4p = 2$. Thus, an equation is $(x - 2)^2 = 2(y + 2)$. $\quad 4p = 2$, so $p = \frac{1}{2}$ and the focus is $\left(2, -\frac{3}{2}\right)$ while the directrix is $y = -\frac{5}{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. $\dfrac{x^2}{16}+\dfrac{y^2}{25}=1 \quad\Rightarrow\quad a=\sqrt{25}=5,\ b=\sqrt{16}=4,$

$c=\sqrt{a^2-b^2}=\sqrt{25-16}=\sqrt{9}=3$. The ellipse is centered at $(0,0)$ with vertices $(0,\pm5)$. The foci are $(0,\pm3)$.

12. $\dfrac{x^2}{4}+\dfrac{y^2}{3}=1 \quad\Rightarrow\quad a=\sqrt{4}=2,\ b=\sqrt{3},$

$c=\sqrt{a^2-b^2}=\sqrt{4-3}=\sqrt{1}=1$. The ellipse is centered at $(0,0)$ with vertices $(\pm2,0)$. The foci are $(\pm1,0)$.

13. $x^2+3y^2=9 \quad\Leftrightarrow\quad \dfrac{x^2}{9}+\dfrac{y^2}{3}=1 \quad\Rightarrow\quad a=\sqrt{9}=3,\ b=\sqrt{3},$

$c=\sqrt{a^2-b^2}=\sqrt{9-3}=\sqrt{6}$. The ellipse is centered at $(0,0)$ with vertices $(\pm3,0)$. The foci are $\left(\pm\sqrt{6},0\right)$.

14. $x^2=4-2y^2 \quad\Leftrightarrow\quad x^2+2y^2=4 \quad\Leftrightarrow\quad \dfrac{x^2}{4}+\dfrac{y^2}{2}=1 \quad\Rightarrow$

$a=\sqrt{4}=2,\ b=\sqrt{2},\ c=\sqrt{a^2-b^2}=\sqrt{4-2}=\sqrt{2}$. The ellipse is centered at $(0,0)$ with vertices $(\pm2,0)$. The foci are $\left(\pm\sqrt{2},0\right)$.

15. $4x^2+25y^2-50y=75 \quad\Leftrightarrow\quad 4x^2+25(y^2-2y+1)=75+25 \quad\Leftrightarrow$

$4x^2+25(y-1)^2=100 \quad\Leftrightarrow\quad \dfrac{x^2}{25}+\dfrac{(y-1)^2}{4}=1 \quad\Rightarrow\quad a=\sqrt{25}=5,$

$b=\sqrt{4}=2,\ c=\sqrt{25-4}=\sqrt{21}$. The ellipse is centered at $(0,1)$ with vertices $(\pm5,1)$. The foci are $\left(\pm\sqrt{21},1\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. $9x^2 - 54x + y^2 + 2y + 46 = 0 \quad \Leftrightarrow$

$9(x^2 - 6x + 9) + y^2 + 2y + 1 = -46 + 81 + 1 \quad \Leftrightarrow$

$9(x-3)^2 + (y+1)^2 = 36 \quad \Leftrightarrow \quad \dfrac{(x-3)^2}{4} + \dfrac{(y+1)^2}{36} = 1 \quad \Rightarrow$

$a = \sqrt{36} = 6$, $b = \sqrt{4} = 2$, $c = \sqrt{36-4} = \sqrt{32} = 4\sqrt{2}$. The ellipse is centered at $(3, -1)$ with vertices $(3, 5)$ and $(3, -7)$. The foci are $\left(3, -1 \pm 4\sqrt{2}\right)$.

17. The center is $(0, 0)$, $a = 3$, and $b = 2$, so an equation is $\dfrac{x^2}{4} + \dfrac{y^2}{9} = 1$. $c = \sqrt{a^2 - b^2} = \sqrt{5}$, so the foci are $\left(0, \pm\sqrt{5}\right)$.

18. The ellipse is centered at $(2, 1)$, with $a = 3$ and $b = 2$. An equation is $\dfrac{(x-2)^2}{9} + \dfrac{(y-1)^2}{4} = 1$. $c = \sqrt{a^2 - b^2} = \sqrt{5}$, so the foci are $\left(2 \pm \sqrt{5}, 1\right)$.

19. $\dfrac{y^2}{25} - \dfrac{x^2}{9} = 1 \quad \Rightarrow \quad a = 5, b = 3, c = \sqrt{a^2 + b^2} = \sqrt{25 + 9} = \sqrt{34} \quad \Rightarrow$

center $(0, 0)$, vertices $(0, \pm 5)$, foci $\left(0, \pm\sqrt{34}\right)$, asymptotes $y = \pm\frac{5}{3}x$.

Note: It is helpful to draw a $2a$-by-$2b$ rectangle whose center is the center of the hyperbola. The asymptotes are the extended diagonals of the rectangle.

20. $\dfrac{x^2}{36} - \dfrac{y^2}{64} = 1 \quad \Rightarrow \quad a = 6, b = 8, c = \sqrt{a^2 + b^2} = \sqrt{36 + 64} = 10 \quad \Rightarrow$

center $(0, 0)$, vertices $(\pm 6, 0)$, foci $(\pm 10, 0)$, asymptotes $y = \pm\frac{8}{6}x = \pm\frac{4}{3}x$

21. $x^2 - y^2 = 100 \quad \Leftrightarrow \quad \dfrac{x^2}{100} - \dfrac{y^2}{100} = 1 \quad \Rightarrow \quad a = b = 10,$

$c = \sqrt{100 + 100} = 10\sqrt{2} \quad \Rightarrow$ center $(0, 0)$, vertices $(\pm 10, 0)$,

foci $\left(\pm 10\sqrt{2}, 0\right)$, asymptotes $y = \pm\frac{10}{10}x = \pm x$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $y^2 - 16x^2 = 16 \quad \Leftrightarrow \quad \dfrac{y^2}{16} - \dfrac{x^2}{1} = 1 \quad \Rightarrow \quad a = 4, b = 1,$

$c = \sqrt{16+1} = \sqrt{17} \quad \Rightarrow \quad$ center $(0, 0)$, vertices $(0, \pm 4)$,

foci $\left(0, \pm\sqrt{17}\right)$, asymptotes $y = \pm\frac{4}{1}x = \pm 4x$

23. $x^2 - y^2 + 2y = 2 \quad \Leftrightarrow \quad x^2 - (y^2 - 2y + 1) = 2 - 1 \quad \Leftrightarrow$

$\dfrac{x^2}{1} - \dfrac{(y-1)^2}{1} = 1 \quad \Rightarrow \quad a = b = 1, c = \sqrt{1+1} = \sqrt{2} \quad \Rightarrow$

center $(0, 1)$, vertices $(\pm 1, 1)$, foci $\left(\pm\sqrt{2}, 1\right)$,

asymptotes $y - 1 = \pm\frac{1}{1}x = \pm x$.

24. $9y^2 - 4x^2 - 36y - 8x = 4 \quad \Leftrightarrow$

$9(y^2 - 4y + 4) - 4(x^2 + 2x + 1) = 4 + 36 - 4 \quad \Leftrightarrow$

$9(y-2)^2 - 4(x+1)^2 = 36 \quad \Leftrightarrow \quad \dfrac{(y-2)^2}{4} - \dfrac{(x+1)^2}{9} = 1 \quad \Rightarrow$

$a = 2, b = 3, c = \sqrt{4+9} = \sqrt{13} \quad \Rightarrow \quad$ center $(-1, 2)$, vertices

$(-1, 2 \pm 2)$, foci $\left(-1, 2 \pm \sqrt{13}\right)$, asymptotes $y - 2 = \pm\frac{2}{3}(x+1)$.

25. The hyperbola has vertices $(\pm 3, 0)$, which lie on the x-axis, so the equation has the form $\dfrac{x^2}{a^2} - \dfrac{y^2}{b^2} = 1$ with $a = 3$. The point $(5, 4)$ is on the hyperbola, so $\dfrac{5^2}{3^2} - \dfrac{4^2}{b^2} = 1 \quad \Rightarrow \quad \dfrac{25}{9} - \dfrac{16}{b^2} = 1 \quad \Rightarrow \quad \dfrac{16}{9} = \dfrac{16}{b^2} \quad \Rightarrow \quad b^2 = 9 \quad \Rightarrow \quad b = 3$. Thus, an equation is $\dfrac{x^2}{9} - \dfrac{y^2}{9} = 1$. Now, $c^2 = a^2 + b^2 = 9 + 9 = 18 \quad \Rightarrow \quad c = \sqrt{18} = 3\sqrt{2}$, so the foci are $\left(\pm 3\sqrt{2}, 0\right)$, while the asymptotes are $y = \pm\dfrac{a}{b}x = \pm\dfrac{3}{3}x = \pm x$.

26. The hyperbola has vertices $(0, \pm 2)$, which lie on the y-axis, so the equation has the form $\dfrac{y^2}{a^2} - \dfrac{x^2}{b^2} = 1$ with $a = 2$. The point $(8, 6)$ is on the hyperbola, so $\dfrac{6^2}{2^2} - \dfrac{8^2}{b^2} = 1 \quad \Rightarrow \quad 9 - \dfrac{64}{b^2} = 1 \quad \Rightarrow \quad 8 = \dfrac{64}{b^2} \quad \Rightarrow \quad b^2 = 8 \quad \Rightarrow \quad b = 2\sqrt{2}$. Thus, an equation is $\dfrac{y^2}{4} - \dfrac{x^2}{8} = 1$. Now, $c^2 = a^2 + b^2 = 4 + 8 = 12 \quad \Rightarrow \quad c = \sqrt{12} = 2\sqrt{3}$, so the foci are $\left(0, \pm 2\sqrt{3}\right)$, while the asymptotes are $y = \pm\dfrac{a}{b}x = \pm\dfrac{2}{2\sqrt{2}}x = \pm\dfrac{1}{\sqrt{2}}x$.

27. $4x^2 = y^2 + 4 \quad \Leftrightarrow \quad 4x^2 - y^2 = 4 \quad \Leftrightarrow \quad \dfrac{x^2}{1} - \dfrac{y^2}{4} = 1$. This is an equation of a *hyperbola* with vertices $(\pm 1, 0)$. The foci are at $\left(\pm\sqrt{1+4}, 0\right) = \left(\pm\sqrt{5}, 0\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $4x^2 = y + 4 \quad \Leftrightarrow \quad x^2 = \frac{1}{4}(y+4)$. This is an equation of a *parabola* with $4p = \frac{1}{4}$, so $p = \frac{1}{16}$. The vertex is $(0, -4)$ and the focus is $\left(0, -4 + \frac{1}{16}\right) = \left(0, -\frac{63}{16}\right)$.

29. $x^2 = 4y - 2y^2 \quad \Leftrightarrow \quad x^2 + 2y^2 - 4y = 0 \quad \Leftrightarrow \quad x^2 + 2(y^2 - 2y + 1) = 2 \quad \Leftrightarrow \quad x^2 + 2(y-1)^2 = 2 \quad \Leftrightarrow$ $\dfrac{x^2}{2} + \dfrac{(y-1)^2}{1} = 1$. This is an equation of an *ellipse* with vertices at $\left(\pm\sqrt{2}, 1\right)$. The foci are at $\left(\pm\sqrt{2-1}, 1\right) = (\pm 1, 1)$.

30. $y^2 - 2 = x^2 - 2x \quad \Leftrightarrow \quad y^2 - x^2 + 2x = 2 \quad \Leftrightarrow \quad y^2 - (x^2 - 2x + 1) = 2 - 1 \quad \Leftrightarrow \quad \dfrac{y^2}{1} - \dfrac{(x-1)^2}{1} = 1$. This is an equation of a *hyperbola* with vertices $(1, \pm 1)$. The foci are at $\left(1, \pm\sqrt{1+1}\right) = \left(1, \pm\sqrt{2}\right)$.

31. $3x^2 - 6x - 2y = 1 \quad \Leftrightarrow \quad 3x^2 - 6x = 2y + 1 \quad \Leftrightarrow \quad 3(x^2 - 2x + 1) = 2y + 1 + 3 \quad \Leftrightarrow \quad 3(x-1)^2 = 2y + 4 \quad \Leftrightarrow$ $(x-1)^2 = \frac{2}{3}(y+2)$. This is an equation of a *parabola* with $4p = \frac{2}{3}$, so $p = \frac{1}{6}$. The vertex is $(1, -2)$ and the focus is $\left(1, -2 + \frac{1}{6}\right) = \left(1, -\frac{11}{6}\right)$.

32. $x^2 - 2x + 2y^2 - 8y + 7 = 0 \quad \Leftrightarrow \quad (x^2 - 2x + 1) + 2(y^2 - 4y + 4) = -7 + 1 + 8 \quad \Leftrightarrow \quad (x-1)^2 + 2(y-2)^2 = 2 \quad \Leftrightarrow$ $\dfrac{(x-1)^2}{2} + \dfrac{(y-2)^2}{1} = 1$. This is an equation of an *ellipse* with vertices at $\left(1 \pm \sqrt{2}, 2\right)$. The foci are at $\left(1 \pm \sqrt{2-1}, 2\right) = (1 \pm 1, 2)$.

33. The parabola with vertex $(0, 0)$ and focus $(1, 0)$ opens to the right and has $p = 1$, so its equation is $y^2 = 4px$, or $y^2 = 4x$.

34. The parabola with focus $(0, 0)$ and directrix $y = 6$ has vertex $(0, 3)$ and opens downward, so $p = -3$ and its equation is $(x - 0)^2 = 4p(y - 3)$, or $x^2 = -12(y - 3)$.

35. The distance from the focus $(-4, 0)$ to the directrix $x = 2$ is $2 - (-4) = 6$, so the distance from the focus to the vertex is $\frac{1}{2}(6) = 3$ and the vertex is $(-1, 0)$. Since the focus is to the left of the vertex, $p = -3$. An equation is $y^2 = 4p(x+1) \quad \Rightarrow$ $y^2 = -12(x+1)$.

36. The parabola with vertex $(2, 3)$ and focus $(2, -1)$ opens downward and has $p = -1 - 3 = -4$, so its equation is $(x - 2)^2 = 4p(y - 3)$, or $(x - 2)^2 = -16(y - 3)$.

37. The parabola with vertex $(3, -1)$ having a horizontal axis has equation $[y - (-1)]^2 = 4p(x - 3)$. Since it passes through $(-15, 2)$, $(2+1)^2 = 4p(-15 - 3) \quad \Rightarrow \quad 9 = 4p(-18) \quad \Rightarrow \quad 4p = -\frac{1}{2}$. An equation is $(y+1)^2 = -\frac{1}{2}(x - 3)$.

38. The parabola with vertical axis and passing through $(0, 4)$ has equation $y = ax^2 + bx + 4$. It also passes through $(1, 3)$ and $(-2, -6)$, so

$$\begin{cases} 3 = a + b + 4 \\ -6 = 4a - 2b + 4 \end{cases} \Rightarrow \begin{cases} -1 = a + b \\ -10 = 4a - 2b \end{cases} \Rightarrow \begin{cases} -1 = a + b \\ -5 = 2a - b \end{cases}$$

Adding the last two equations gives us $3a = -6$, or $a = -2$. Since $a + b = -1$, we have $b = 1$, and an equation is $y = -2x^2 + x + 4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. The ellipse with foci $(\pm 2, 0)$ and vertices $(\pm 5, 0)$ has center $(0, 0)$ and a horizontal major axis, with $a = 5$ and $c = 2$, so $b^2 = a^2 - c^2 = 25 - 4 = 21$. An equation is $\dfrac{x^2}{25} + \dfrac{y^2}{21} = 1$.

40. The ellipse with foci $\left(0, \pm\sqrt{2}\right)$ and vertices $(0, \pm 2)$ has center $(0, 0)$ and a vertical major axis, with $a = 2$ and $c = \sqrt{2}$, so $b^2 = a^2 - c^2 = 4 - 2 = 2$. An equation is $\dfrac{x^2}{2} + \dfrac{y^2}{4} = 1$.

41. Since the vertices are $(0, 0)$ and $(0, 8)$, the ellipse has center $(0, 4)$ with a vertical axis and $a = 4$. The foci at $(0, 2)$ and $(0, 6)$ are 2 units from the center, so $c = 2$ and $b = \sqrt{a^2 - c^2} = \sqrt{4^2 - 2^2} = \sqrt{12}$. An equation is $\dfrac{(x-0)^2}{b^2} + \dfrac{(y-4)^2}{a^2} = 1 \quad \Rightarrow$ $\dfrac{x^2}{12} + \dfrac{(y-4)^2}{16} = 1$.

42. Since the foci are $(0, -1)$ and $(8, -1)$, the ellipse has center $(4, -1)$ with a horizontal axis and $c = 4$. The vertex $(9, -1)$ is 5 units from the center, so $a = 5$ and $b = \sqrt{a^2 - c^2} = \sqrt{5^2 - 4^2} = \sqrt{9}$. An equation is $\dfrac{(x-4)^2}{a^2} + \dfrac{(y+1)^2}{b^2} = 1 \quad \Rightarrow \quad \dfrac{(x-4)^2}{25} + \dfrac{(y+1)^2}{9} = 1$.

43. An equation of an ellipse with center $(-1, 4)$ and vertex $(-1, 0)$ is $\dfrac{(x+1)^2}{b^2} + \dfrac{(y-4)^2}{4^2} = 1$. The focus $(-1, 6)$ is 2 units from the center, so $c = 2$. Thus, $b^2 + 2^2 = 4^2 \quad \Rightarrow \quad b^2 = 12$, and the equation is $\dfrac{(x+1)^2}{12} + \dfrac{(y-4)^2}{16} = 1$.

44. Foci $F_1(-4, 0)$ and $F_2(4, 0) \quad \Rightarrow \quad c = 4$ and an equation is $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1$. The ellipse passes through $P(-4, 1.8)$, so $2a = |PF_1| + |PF_2| \quad \Rightarrow \quad 2a = 1.8 + \sqrt{8^2 + (1.8)^2} \quad \Rightarrow \quad 2a = 1.8 + 8.2 \quad \Rightarrow \quad a = 5$.
$b^2 = a^2 - c^2 = 25 - 16 = 9$ and the equation is $\dfrac{x^2}{25} + \dfrac{y^2}{9} = 1$.

45. An equation of a hyperbola with vertices $(\pm 3, 0)$ is $\dfrac{x^2}{3^2} - \dfrac{y^2}{b^2} = 1$. Foci $(\pm 5, 0) \quad \Rightarrow \quad c = 5$ and $3^2 + b^2 = 5^2 \quad \Rightarrow$ $b^2 = 25 - 9 = 16$, so the equation is $\dfrac{x^2}{9} - \dfrac{y^2}{16} = 1$.

46. An equation of a hyperbola with vertices $(0, \pm 2)$ is $\dfrac{y^2}{2^2} - \dfrac{x^2}{b^2} = 1$. Foci $(0, \pm 5) \quad \Rightarrow \quad c = 5$ and $2^2 + b^2 = 5^2 \quad \Rightarrow$ $b^2 = 25 - 4 = 21$, so the equation is $\dfrac{y^2}{4} - \dfrac{x^2}{21} = 1$.

47. The center of a hyperbola with vertices $(-3, -4)$ and $(-3, 6)$ is $(-3, 1)$, so $a = 5$ and an equation is $\dfrac{(y-1)^2}{5^2} - \dfrac{(x+3)^2}{b^2} = 1$. Foci $(-3, -7)$ and $(-3, 9) \quad \Rightarrow \quad c = 8$, so $5^2 + b^2 = 8^2 \quad \Rightarrow \quad b^2 = 64 - 25 = 39$ and the equation is $\dfrac{(y-1)^2}{25} - \dfrac{(x+3)^2}{39} = 1$.

48. The center of a hyperbola with vertices $(-1, 2)$ and $(7, 2)$ is $(3, 2)$, so $a = 4$ and an equation is $\dfrac{(x-3)^2}{4^2} - \dfrac{(y-2)^2}{b^2} = 1$. Foci $(-2, 2)$ and $(8, 2) \quad \Rightarrow \quad c = 5$, so $4^2 + b^2 = 5^2 \quad \Rightarrow \quad b^2 = 25 - 16 = 9$ and the equation is $\dfrac{(x-3)^2}{16} - \dfrac{(y-2)^2}{9} = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. The center of a hyperbola with vertices $(\pm 3, 0)$ is $(0, 0)$, so $a = 3$ and an equation is $\dfrac{x^2}{3^2} - \dfrac{y^2}{b^2} = 1$.

Asymptotes $y = \pm 2x \quad \Rightarrow \quad \dfrac{b}{a} = 2 \quad \Rightarrow \quad b = 2(3) = 6$ and the equation is $\dfrac{x^2}{9} - \dfrac{y^2}{36} = 1$.

50. The center of a hyperbola with foci $(2, 0)$ and $(2, 8)$ is $(2, 4)$, so $c = 4$ and an equation is $\dfrac{(y-4)^2}{a^2} - \dfrac{(x-2)^2}{b^2} = 1$.

The asymptote $y = 3 + \frac{1}{2}x$ has slope $\frac{1}{2}$, so $\dfrac{a}{b} = \dfrac{1}{2} \quad \Rightarrow \quad b = 2a$ and $a^2 + b^2 = c^2 \quad \Rightarrow \quad a^2 + (2a)^2 = 4^2 \quad \Rightarrow$

$5a^2 = 16 \quad \Rightarrow \quad a^2 = \frac{16}{5}$ and so $b^2 = 16 - \frac{16}{5} = \frac{64}{5}$. Thus, an equation is $\dfrac{(y-4)^2}{16/5} - \dfrac{(x-2)^2}{64/5} = 1$.

51. In Figure 8, we see that the point on the ellipse closest to a focus is the closer vertex (which is a distance $a - c$ from it) while the farthest point is the other vertex (at a distance of $a + c$). So for this lunar orbit, $(a - c) + (a + c) = 2a = (1728 + 110) + (1728 + 314)$, or $a = 1940$; and $(a + c) - (a - c) = 2c = 314 - 110$, or $c = 102$. Thus, $b^2 = a^2 - c^2 = 3{,}753{,}196$, and the equation is $\dfrac{x^2}{3{,}763{,}600} + \dfrac{y^2}{3{,}753{,}196} = 1$.

52. (a) Choose V to be the origin, with x-axis through V and F. Then F is $(p, 0)$, A is $(p, 5)$, so substituting A into the equation $y^2 = 4px$ gives $25 = 4p^2$ so $p = \frac{5}{2}$ and $y^2 = 10x$.

(b) $x = 11 \quad \Rightarrow \quad y = \sqrt{110} \quad \Rightarrow \quad |CD| = 2\sqrt{110}$

53. (a) Set up the coordinate system so that A is $(-200, 0)$ and B is $(200, 0)$.

$|PA| - |PB| = (1200)(980) = 1{,}176{,}000$ ft $= \frac{2450}{11}$ mi $= 2a \quad \Rightarrow \quad a = \frac{1225}{11}$, and $c = 200$ so

$b^2 = c^2 - a^2 = \dfrac{3{,}339{,}375}{121} \quad \Rightarrow \quad \dfrac{121x^2}{1{,}500{,}625} - \dfrac{121y^2}{3{,}339{,}375} = 1.$

(b) Due north of $B \quad \Rightarrow \quad x = 200 \quad \Rightarrow \quad \dfrac{(121)(200)^2}{1{,}500{,}625} - \dfrac{121y^2}{3{,}339{,}375} = 1 \quad \Rightarrow \quad y = \dfrac{133{,}575}{539} \approx 248$ mi

54. $|PF_1| - |PF_2| = \pm 2a \quad \Leftrightarrow \quad \sqrt{(x+c)^2 + y^2} - \sqrt{(x-c)^2 + y^2} = \pm 2a \quad \Leftrightarrow$

$\sqrt{(x+c)^2 + y^2} = \sqrt{(x-c)^2 + y^2} \pm 2a \quad \Leftrightarrow \quad (x+c)^2 + y^2 = (x-c)^2 + y^2 + 4a^2 \pm 4a\sqrt{(x-c)^2 + y^2} \quad \Leftrightarrow$

$4cx - 4a^2 = \pm 4a\sqrt{(x-c)^2 + y^2} \quad \Leftrightarrow \quad c^2x^2 - 2a^2cx + a^4 = a^2(x^2 - 2cx + c^2 + y^2) \quad \Leftrightarrow$

$(c^2 - a^2)x^2 - a^2y^2 = a^2(c^2 - a^2) \quad \Leftrightarrow \quad b^2x^2 - a^2y^2 = a^2b^2$ [where $b^2 = c^2 - a^2$] $\quad \Leftrightarrow \quad \dfrac{x^2}{a^2} - \dfrac{y^2}{b^2} = 1$

55. The function whose graph is the upper branch of this hyperbola is concave upward. The function is

$y = f(x) = a\sqrt{1 + \dfrac{x^2}{b^2}} = \dfrac{a}{b}\sqrt{b^2 + x^2}$, so $y' = \dfrac{a}{b}x(b^2 + x^2)^{-1/2}$ and

$y'' = \dfrac{a}{b}\left[(b^2 + x^2)^{-1/2} - x^2(b^2 + x^2)^{-3/2}\right] = ab(b^2 + x^2)^{-3/2} > 0$ for all x, and so f is concave upward.

56. We can follow exactly the same sequence of steps as in the derivation of Formula 4, except we use the points $(1, 1)$ and $(-1, -1)$ in the distance formula (first equation of that derivation) so $\sqrt{(x-1)^2 + (y-1)^2} + \sqrt{(x+1)^2 + (y+1)^2} = 4$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

will lead (after moving the second term to the right, squaring, and simplifying) to $2\sqrt{(x+1)^2+(y+1)^2} = x+y+4$, which, after squaring and simplifying again, leads to $3x^2 - 2xy + 3y^2 = 8$.

57. (a) If $k > 16$, then $k-16>0$, and $\dfrac{x^2}{k} + \dfrac{y^2}{k-16} = 1$ is an *ellipse* since it is the sum of two squares on the left side.

(b) If $0<k<16$, then $k-16<0$, and $\dfrac{x^2}{k} + \dfrac{y^2}{k-16} = 1$ is a *hyperbola* since it is the difference of two squares on the left side.

(c) If $k<0$, then $k-16<0$, and there is *no curve* since the left side is the sum of two negative terms, which cannot equal 1.

(d) In case (a), $a^2 = k$, $b^2 = k-16$, and $c^2 = a^2 - b^2 = 16$, so the foci are at $(\pm 4, 0)$. In case (b), $k-16<0$, so $a^2 = k$, $b^2 = 16-k$, and $c^2 = a^2+b^2 = 16$, and so again the foci are at $(\pm 4, 0)$.

58. (a) $y^2 = 4px \;\Rightarrow\; 2yy' = 4p \;\Rightarrow\; y' = \dfrac{2p}{y}$, so the tangent line is

$$y - y_0 = \frac{2p}{y_0}(x - x_0) \;\Rightarrow\; yy_0 - y_0^2 = 2p(x - x_0) \;\Leftrightarrow$$

$$yy_0 - 4px_0 = 2px - 2px_0 \;\Rightarrow\; yy_0 = 2p(x + x_0).$$

(b) The x-intercept is $-x_0$.

59. $x^2 = 4py \;\Rightarrow\; 2x = 4py' \;\Rightarrow\; y' = \dfrac{x}{2p}$, so the tangent line at (x_0, y_0) is $y - \dfrac{x_0^2}{4p} = \dfrac{x_0}{2p}(x - x_0)$. This line passes through the point $(a, -p)$ on the directrix, so $-p - \dfrac{x_0^2}{4p} = \dfrac{x_0}{2p}(a - x_0) \;\Rightarrow\; -4p^2 - x_0^2 = 2ax_0 - 2x_0^2 \;\Leftrightarrow$

$x_0^2 - 2ax_0 - 4p^2 = 0 \;\Leftrightarrow\; x_0^2 - 2ax_0 + a^2 = a^2 + 4p^2 \;\Leftrightarrow$

$(x_0 - a)^2 = a^2 + 4p^2 \;\Leftrightarrow\; x_0 = a \pm \sqrt{a^2+4p^2}$. The slopes of the tangent lines at $x = a \pm \sqrt{a^2+4p^2}$ are $\dfrac{a \pm \sqrt{a^2+4p^2}}{2p}$, so the product of the two slopes is

$$\frac{a+\sqrt{a^2+4p^2}}{2p} \cdot \frac{a-\sqrt{a^2+4p^2}}{2p} = \frac{a^2-(a^2+4p^2)}{4p^2} = \frac{-4p^2}{4p^2} = -1,$$

showing that the tangent lines are perpendicular.

60. Without a loss of generality, let the ellipse, hyperbola, and foci be as shown in the figure.

The curves intersect (eliminate y^2) $\Rightarrow$

$$B^2\left(\frac{x^2}{A^2} - \frac{y^2}{B^2}\right) + b^2\left(\frac{x^2}{a^2} + \frac{y^2}{b^2}\right) = B^2 + b^2 \;\Rightarrow$$

$$\frac{B^2x^2}{A^2} + \frac{b^2x^2}{a^2} = B^2 + b^2 \;\Rightarrow\; x^2\left(\frac{B^2}{A^2} + \frac{b^2}{a^2}\right) = B^2 + b^2 \;\Rightarrow$$

$$x^2 = \frac{B^2+b^2}{\dfrac{a^2B^2+b^2A^2}{A^2a^2}} = \frac{A^2a^2(B^2+b^2)}{a^2B^2+b^2A^2}.$$

Similarly, $y^2 = \dfrac{B^2b^2(a^2-A^2)}{b^2A^2+a^2B^2}$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Next we find the slopes of the tangent lines of the curves: $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1 \;\Rightarrow\; \dfrac{2x}{a^2}+\dfrac{2yy'}{b^2}=0 \;\Rightarrow\; \dfrac{yy'}{b^2}=-\dfrac{x}{a^2} \;\Rightarrow$ $y'_E=-\dfrac{b^2}{a^2}\,\dfrac{x}{y}$ and $\dfrac{x^2}{A^2}-\dfrac{y^2}{B^2}=1 \;\Rightarrow\; \dfrac{2x}{A^2}-\dfrac{2yy'}{B^2}=0 \;\Rightarrow\; \dfrac{yy'}{B^2}=\dfrac{x}{A^2} \;\Rightarrow\; y'_H=\dfrac{B^2}{A^2}\,\dfrac{x}{y}$. The product of the slopes at (x_0,y_0) is $y'_E\,y'_H=-\dfrac{b^2B^2x_0^2}{a^2A^2y_0^2}=-\dfrac{b^2B^2\left[\dfrac{A^2a^2(B^2+b^2)}{a^2B^2+b^2A^2}\right]}{a^2A^2\left[\dfrac{B^2b^2(a^2-A^2)}{b^2A^2+a^2B^2}\right]}=-\dfrac{B^2+b^2}{a^2-A^2}$. Since $a^2-b^2=c^2$ and $A^2+B^2=c^2$, we have $a^2-b^2=A^2+B^2 \;\Rightarrow\; a^2-A^2=b^2+B^2$, so the product of the slopes is -1, and hence, the tangent lines at each point of intersection are perpendicular.

61. $9x^2+4y^2=36 \;\Leftrightarrow\; \dfrac{x^2}{4}+\dfrac{y^2}{9}=1$. We use the parametrization $x=2\cos t$, $y=3\sin t$, $0\le t\le 2\pi$. The circumference is given by

$$\begin{aligned} L &= \textstyle\int_0^{2\pi}\sqrt{(dx/dt)^2+(dy/dt)^2}\,dt=\int_0^{2\pi}\sqrt{(-2\sin t)^2+(3\cos t)^2}\,dt \\ &= \textstyle\int_0^{2\pi}\sqrt{4\sin^2 t+9\cos^2 t}\,dt=\int_0^{2\pi}\sqrt{4+5\cos^2 t}\,dt \end{aligned}$$

Now use Simpson's Rule with $n=8$, $\Delta t=\dfrac{2\pi-0}{8}=\dfrac{\pi}{4}$, and $f(t)=\sqrt{4+5\cos^2 t}$ to get

$L\approx S_8=\frac{\pi/4}{3}\left[f(0)+4f\left(\frac{\pi}{4}\right)+2f\left(\frac{\pi}{2}\right)+4f\left(\frac{3\pi}{4}\right)+2f(\pi)+4f\left(\frac{5\pi}{4}\right)+2f\left(\frac{3\pi}{2}\right)+4f\left(\frac{7\pi}{4}\right)+f(2\pi)\right]\approx 15.9.$

62. The length of the major axis is $2a$, so $a=\frac{1}{2}(1.18\times 10^{10})=5.9\times 10^9$. The length of the minor axis is $2b$, so $b=\frac{1}{2}(1.14\times 10^{10})=5.7\times 10^9$. An equation of the ellipse is $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1$, or converting into parametric equations, $x=a\cos\theta$ and $y=b\sin\theta$. So

$$L=4\textstyle\int_0^{\pi/2}\sqrt{(dx/d\theta)^2+(dy/d\theta)^2}\,d\theta=4\int_0^{\pi/2}\sqrt{a^2\sin^2\theta+b^2\cos^2\theta}\,d\theta$$

Using Simpson's Rule with $n=10$, $\Delta\theta=\frac{\pi/2-0}{10}=\frac{\pi}{20}$, and $f(\theta)=\sqrt{a^2\sin^2\theta+b^2\cos^2\theta}$, we get

$$L\approx 4\cdot S_{10}=4\cdot\tfrac{\pi}{20\cdot 3}\left[f(0)+4f\left(\tfrac{\pi}{20}\right)+2f\left(\tfrac{2\pi}{20}\right)+\cdots+2f\left(\tfrac{8\pi}{20}\right)+4f\left(\tfrac{9\pi}{20}\right)+f\left(\tfrac{\pi}{2}\right)\right]\approx 3.64\times 10^{10}\text{ km}$$

63. $\dfrac{x^2}{a^2}-\dfrac{y^2}{b^2}=1 \;\Rightarrow\; \dfrac{y^2}{b^2}=\dfrac{x^2-a^2}{a^2} \;\Rightarrow\; y=\pm\dfrac{b}{a}\sqrt{x^2-a^2}$.

$$\begin{aligned} A &= 2\int_a^c \frac{b}{a}\sqrt{x^2-a^2}\,dx \overset{39}{=} \frac{2b}{a}\left[\frac{x}{2}\sqrt{x^2-a^2}-\frac{a^2}{2}\ln\left|x+\sqrt{x^2-a^2}\right|\right]_a^c \\ &= \frac{b}{a}\left[c\sqrt{c^2-a^2}-a^2\ln\left|c+\sqrt{c^2-a^2}\right|+a^2\ln|a|\right] \end{aligned}$$

Since $a^2+b^2=c^2$, $c^2-a^2=b^2$, and $\sqrt{c^2-a^2}=b$.

$$\begin{aligned} &= \frac{b}{a}\left[cb-a^2\ln(c+b)+a^2\ln a\right]=\frac{b}{a}\left[cb+a^2(\ln a-\ln(b+c))\right] \\ &= b^2c/a+ab\ln[a/(b+c)],\ \text{ where } c^2=a^2+b^2. \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. (a) $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1 \;\Rightarrow\; \dfrac{y^2}{b^2}=\dfrac{a^2-x^2}{a^2} \;\Rightarrow\; y=\pm\dfrac{b}{a}\sqrt{a^2-x^2}.$

$$V=\int_{-a}^{a}\pi\left(\frac{b}{a}\sqrt{a^2-x^2}\right)^2 dx=2\pi\frac{b^2}{a^2}\int_0^a (a^2-x^2)\,dx$$

$$=\frac{2\pi b^2}{a^2}\left[a^2x-\tfrac{1}{3}x^3\right]_0^a=\frac{2\pi b^2}{a^2}\left(\frac{2a^3}{3}\right)=\frac{4}{3}\pi b^2 a$$

(b) $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1 \;\Rightarrow\; \dfrac{x^2}{a^2}=\dfrac{b^2-y^2}{b^2} \;\Rightarrow\; x=\pm\dfrac{a}{b}\sqrt{b^2-y^2}.$

$$V=\int_{-b}^{b}\pi\left(\frac{a}{b}\sqrt{b^2-y^2}\right)^2 dy=2\pi\frac{a^2}{b^2}\int_0^b (b^2-y^2)\,dy$$

$$=\frac{2\pi a^2}{b^2}\left[b^2y-\tfrac{1}{3}y^3\right]_0^b=\frac{2\pi a^2}{b^2}\left(\frac{2b^3}{3}\right)=\frac{4}{3}\pi a^2 b$$

65. $9x^2+4y^2=36 \;\Leftrightarrow\; \dfrac{x^2}{4}+\dfrac{y^2}{9}=1 \;\Rightarrow\; a=3, b=2$. By symmetry, $\overline{x}=0$. By Example 2 in Section 7.3, the area of the top half of the ellipse is $\frac{1}{2}(\pi ab)=3\pi$. Solve $9x^2+4y^2=36$ for y to get an equation for the top half of the ellipse:

$9x^2+4y^2=36 \;\Leftrightarrow\; 4y^2=36-9x^2 \;\Leftrightarrow\; y^2=\frac{9}{4}(4-x^2) \;\Rightarrow\; y=\frac{3}{2}\sqrt{4-x^2}$. Now

$$\overline{y}=\frac{1}{A}\int_a^b\frac{1}{2}[f(x)]^2\,dx=\frac{1}{3\pi}\int_{-2}^{2}\frac{1}{2}\left(\frac{3}{2}\sqrt{4-x^2}\right)^2 dx=\frac{3}{8\pi}\int_{-2}^{2}(4-x^2)\,dx$$

$$=\frac{3}{8\pi}\cdot 2\int_0^2(4-x^2)\,dx=\frac{3}{4\pi}\left[4x-\frac{1}{3}x^3\right]_0^2=\frac{3}{4\pi}\left(\frac{16}{3}\right)=\frac{4}{\pi}$$

so the centroid is $(0, 4/\pi)$.

66. (a) Consider the ellipse $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1$ with $a>b$, so that the major axis is the x-axis. Let the ellipse be parametrized by $x=a\cos t$, $y=b\sin t$, $0\le t\le 2\pi$. Then

$$\left(\frac{dx}{dt}\right)^2+\left(\frac{dy}{dt}\right)^2=a^2\sin^2 t+b^2\cos^2 t=a^2(1-\cos^2 t)+b^2\cos^2 t=a^2+(b^2-a^2)\cos^2 t=a^2-c^2\cos^2 t,$$

where $c^2=a^2-b^2$. Using symmetry and rotating the ellipse about the major axis gives us surface area

$$S=\int 2\pi y\,ds=2\int_0^{\pi/2}2\pi(b\sin t)\sqrt{a^2-c^2\cos^2 t}\,dt=4\pi b\int_c^0\sqrt{a^2-u^2}\left(-\frac{1}{c}\,du\right)\qquad\begin{bmatrix}u=c\cos t\\ du=-c\sin t\,dt\end{bmatrix}$$

$$=\frac{4\pi b}{c}\int_0^c\sqrt{a^2-u^2}\,du\overset{30}{=}\frac{4\pi b}{c}\left[\frac{u}{2}\sqrt{a^2-u^2}+\frac{a^2}{2}\sin^{-1}\left(\frac{u}{a}\right)\right]_0^c=\frac{2\pi b}{c}\left[c\sqrt{a^2-c^2}+a^2\sin^{-1}\left(\frac{c}{a}\right)\right]$$

$$=\frac{2\pi b}{c}\left[bc+a^2\sin^{-1}\left(\frac{c}{a}\right)\right]$$

(b) As in part (a),

$$\left(\frac{dx}{dt}\right)^2+\left(\frac{dy}{dt}\right)^2=a^2\sin^2 t+b^2\cos^2 t=a^2\sin^2 t+b^2(1-\sin^2 t)=b^2+(a^2-b^2)\sin^2 t=b^2+c^2\sin^2 t.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Rotating about the minor axis gives us

$$S = \int 2\pi x\,ds = 2\int_0^{\pi/2} 2\pi(a\cos t)\sqrt{b^2 + c^2\sin^2 t}\,dt = 4\pi a\int_0^c \sqrt{b^2+u^2}\left(\frac{1}{c}\,du\right) \qquad \left[\begin{array}{c} u = c\sin t \\ du = c\cos t\,dt \end{array}\right]$$

$$\overset{21}{=} \frac{4\pi a}{c}\left[\frac{u}{2}\sqrt{b^2+u^2} + \frac{b^2}{2}\ln\left(u+\sqrt{b^2+u^2}\right)\right]_0^c = \frac{2\pi a}{c}\left[c\sqrt{b^2+c^2} + b^2\ln\left(c+\sqrt{b^2+c^2}\right) - b^2\ln b\right]$$

$$= \frac{2\pi a}{c}\left[ac + b^2\ln\left(\frac{a+c}{b}\right)\right]$$

67. Differentiating implicitly, $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 \;\Rightarrow\; \dfrac{2x}{a^2} + \dfrac{2yy'}{b^2} = 0 \;\Rightarrow\; y' = -\dfrac{b^2x}{a^2y}$ $[y \neq 0]$. Thus, the slope of the tangent line at P is $-\dfrac{b^2x_1}{a^2y_1}$. The slope of F_1P is $\dfrac{y_1}{x_1+c}$ and of F_2P is $\dfrac{y_1}{x_1-c}$. By the formula in Problem 21 in Problems Plus following Chapter 3, we have

$$\tan\alpha = \frac{\dfrac{y_1}{x_1+c} + \dfrac{b^2x_1}{a^2y_1}}{1 - \dfrac{b^2x_1y_1}{a^2y_1(x_1+c)}} = \frac{a^2y_1^2 + b^2x_1(x_1+c)}{a^2y_1(x_1+c) - b^2x_1y_1} = \frac{a^2b^2 + b^2cx_1}{c^2x_1y_1 + a^2cy_1} \qquad \left[\begin{array}{c}\text{using } b^2x_1^2 + a^2y_1^2 = a^2b^2, \\ \text{and } a^2 - b^2 = c^2\end{array}\right]$$

$$= \frac{b^2\left(cx_1 + a^2\right)}{cy_1(cx_1+a^2)} = \frac{b^2}{cy_1}$$

and $$\tan\beta = \frac{-\dfrac{b^2x_1}{a^2y_1} - \dfrac{y_1}{x_1-c}}{1 - \dfrac{b^2x_1y_1}{a^2y_1(x_1-c)}} = \frac{-a^2y_1^2 - b^2x_1(x_1-c)}{a^2y_1\,(x_1-c) - b^2x_1y_1} = \frac{-a^2b^2 + b^2cx_1}{c^2x_1y_1 - a^2cy_1} = \frac{b^2\left(cx_1 - a^2\right)}{cy_1(cx_1-a^2)} = \frac{b^2}{cy_1}$$

Thus, $\alpha = \beta$.

68. The slopes of the line segments F_1P and F_2P are $\dfrac{y_1}{x_1+c}$ and $\dfrac{y_1}{x_1-c}$, where P is (x_1, y_1). Differentiating implicitly, $\dfrac{2x}{a^2} - \dfrac{2yy'}{b^2} = 0 \;\Rightarrow\; y' = \dfrac{b^2x}{a^2y} \;\Rightarrow$ the slope of the tangent at P is $\dfrac{b^2x_1}{a^2y_1}$, so by the formula in Problem 21 in Problems Plus following Chapter 3,

$$\tan\alpha = \frac{\dfrac{b^2x_1}{a^2y_1} - \dfrac{y_1}{x_1+c}}{1 + \dfrac{b^2x_1y_1}{a^2y_1(x_1+c)}} = \frac{b^2x_1(x_1+c) - a^2y_1^2}{a^2y_1(x_1+c) + b^2x_1y_1} = \frac{b^2(cx_1+a^2)}{cy_1(cx_1+a^2)} \qquad \left[\begin{array}{c}\text{using } x_1^2/a^2 - y_1^2/b^2 = 1, \\ \text{and } a^2 + b^2 = c^2\end{array}\right] = \frac{b^2}{cy_1}$$

and $$\tan\beta = \frac{-\dfrac{b^2x_1}{a^2y_1} + \dfrac{y_1}{x_1-c}}{1 + \dfrac{b^2x_1y_1}{a^2y_1(x_1-c)}} = \frac{-b^2x_1(x_1-c) + a^2y_1^2}{a^2y_1(x_1-c) + b^2x_1y_1} = \frac{b^2(cx_1-a^2)}{cy_1(cx_1-a^2)} = \frac{b^2}{cy_1}$$

So $\alpha = \beta$.

69. Let C be the center of a circle (gray) with radius r that is tangent to both black circles (see the figure). We wish to show that $AC + BC$ is constant for all values of r, that is, for any circle drawn tangent to both black circles. The smaller black circle has

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

radius 3, so $AC = 3 + r$, and the larger black circle has radius 5, so $BC = 5 - r$. Hence, $AC + BC = 3 + r + 5 - r = 8$, which is a constant. Since the sum of the distances from C to $(-1, 0)$ and $(1, 0)$ is constant, the centers of all the circles lie on an ellipse with foci $(\pm 1, 0)$ $[c = 1]$. The sum of the distances from the foci to any point on the ellipse is $2a$, so $2a = 8 \quad \Rightarrow \quad a = 4$. Now, $c^2 = a^2 - b^2 \quad \Rightarrow \quad 1^2 = 4^2 - b^2 \quad \Rightarrow \quad b^2 = 15$. Thus, the ellipse has equation $\dfrac{x^2}{16} + \dfrac{y^2}{15} = 1$.

10.6 Conic Sections in Polar Coordinates

1. The directrix $x = 2$ is to the right of the focus at the origin, so we use the form with "$+e\cos\theta$" in the denominator. (See Theorem 6 and Figure 2.) $e = 1$ for a parabola, so an equation is $r = \dfrac{ed}{1 + e\cos\theta} = \dfrac{1 \cdot 2}{1 + 1\cos\theta} = \dfrac{2}{1 + \cos\theta}$.

2. The directrix $y = 6$ is above the focus at the origin, so we use the form with "$+e\sin\theta$" in the denominator. An equation of the ellipse is $r = \dfrac{ed}{1 + e\sin\theta} = \dfrac{\frac{1}{3} \cdot 6}{1 + \frac{1}{3}\sin\theta} = \dfrac{6}{3 + \sin\theta}$.

3. The directrix $y = -4$ is below the focus at the origin, so we use the form with "$-e\sin\theta$" in the denominator. An equation of the hyperbola is $r = \dfrac{ed}{1 - e\sin\theta} = \dfrac{2 \cdot 4}{1 - 2\sin\theta} = \dfrac{8}{1 - 2\sin\theta}$.

4. The directrix $x = -3$ is to the left of the focus at the origin, so we use the form with "$-e\cos\theta$" in the denominator. An equation of the hyperbola is $r = \dfrac{ed}{1 - e\cos\theta} = \dfrac{\frac{5}{2} \cdot 3}{1 - \frac{5}{2}\cos\theta} = \dfrac{15}{2 - 5\cos\theta}$.

5. The vertex $(2, \pi)$ is to the left of the focus at the origin, so we use the form with "$-e\cos\theta$" in the denominator. An equation of the ellipse is $r = \dfrac{ed}{1 - e\cos\theta}$. Using eccentricity $e = \dfrac{2}{3}$ with $\theta = \pi$ and $r = 2$, we get $2 = \dfrac{\frac{2}{3}d}{1 - \frac{2}{3}(-1)} \quad \Rightarrow$ $2 = \dfrac{2d}{5} \quad \Rightarrow \quad d = 5$, so we have $r = \dfrac{\frac{2}{3}(5)}{1 - \frac{2}{3}\cos\theta} = \dfrac{10}{3 - 2\cos\theta}$.

6. The directrix $r = 4\csc\theta$ (equivalent to $r\sin\theta = 4$ or $y = 4$) is above the focus at the origin, so we will use the form with "$+e\sin\theta$" in the denominator. The distance from the focus to the directrix is $d = 4$, so an equation of the ellipse is $r = \dfrac{ed}{1 + e\sin\theta} = \dfrac{(0.6)(4)}{1 + 0.6\sin\theta} \cdot \dfrac{5}{5} = \dfrac{12}{5 + 3\sin\theta}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. The vertex $\left(3, \frac{\pi}{2}\right)$ is 3 units above the focus at the origin, so the directrix is 6 units above the focus ($d = 6$), and we use the form "$+e\sin\theta$" in the denominator. $e = 1$ for a parabola, so an equation is $r = \dfrac{ed}{1 + e\sin\theta} = \dfrac{1(6)}{1 + 1\sin\theta} = \dfrac{6}{1 + \sin\theta}$.

8. The directrix $r = -2\sec\theta$ (equivalent to $r\cos\theta = -2$ or $x = -2$) is left of the focus at the origin, so we will use the form with "$-e\cos\theta$" in the denominator. The distance from the focus to the directrix is $d = 2$, so an equation of the hyperbola is $r = \dfrac{ed}{1 - e\cos\theta} = \dfrac{2(2)}{1 - 2\cos\theta} = \dfrac{4}{1 - 2\cos\theta}$.

9. $r = \dfrac{3}{1 - \sin\theta}$, where $e = 1$, so the conic is a parabola. If $\sin\theta$ appears in the denominator, use 0 and π for θ. If $\cos\theta$ appears in the denominator, use $\frac{\pi}{2}$ and $\frac{3\pi}{2}$ for θ. Thus, when $\theta = 0$ or π, $r = 3$. Hence, the equation is matched with graph VI.

10. $r = \dfrac{9}{1 + 2\cos\theta}$, where $e = 2 > 1$, so the conic is a hyperbola. When $\theta = \frac{\pi}{2}$ or $\frac{3\pi}{2}$, $r = 9$. Hence, the equation is matched with graph III.

11. $r = \dfrac{12}{8 - 7\cos\theta} \cdot \dfrac{1/8}{1/8} = \dfrac{3/2}{1 - \frac{7}{8}\cos\theta}$, where $e = \dfrac{7}{8} < 1$, so the conic is an ellipse. When $\theta = \frac{\pi}{2}$ or $\frac{3\pi}{2}$, $r = \frac{12}{8}$. Hence, the equation is matched with graph II.

12. $r = \dfrac{12}{4 + 3\sin\theta} \cdot \dfrac{1/4}{1/4} = \dfrac{3}{1 + \frac{3}{4}\sin\theta}$, where $e = \dfrac{3}{4} < 1$, so the conic is an ellipse. When $\theta = 0$ or π, $r = \frac{12}{4}$. Hence, the equation is matched with graph V.

13. $r = \dfrac{5}{2 + 3\sin\theta} \cdot \dfrac{1/2}{1/2} = \dfrac{5/2}{1 + \frac{3}{2}\sin\theta}$, where $e = \dfrac{3}{2} > 1$, so the conic is a hyperbola. When $\theta = 0$ or π, $r = \frac{5}{2}$. Hence, the equation is matched with graph IV.

14. $r = \dfrac{3}{2 - 2\cos\theta} \cdot \dfrac{1/2}{1/2} = \dfrac{3/2}{1 - \cos\theta}$, where $e = 1$, so the conic is a parabola. When $\theta = \frac{\pi}{2}$ or $\frac{3\pi}{2}$, $r = \frac{3}{2}$. Hence, the equation is matched with graph I.

15. $r = \dfrac{4}{5 - 4\sin\theta} \cdot \dfrac{1/5}{1/5} = \dfrac{4/5}{1 - \frac{4}{5}\sin\theta}$, where $e = \frac{4}{5}$ and $ed = \frac{4}{5} \quad \Rightarrow \quad d = 1$.

(a) Eccentricity $= e = \frac{4}{5}$

(b) Since $e = \frac{4}{5} < 1$, the conic is an ellipse.

(c) Since "$-\ e\sin\theta$" appears in the denominator, the directrix is below the focus at the origin, $d = |Fl| = 1$, so an equation of the directrix is $y = -1$.

(d) The vertices are $\left(4, \frac{\pi}{2}\right)$ and $\left(\frac{4}{9}, \frac{3\pi}{2}\right)$.

16. $r = \dfrac{1}{2 + \sin\theta} \cdot \dfrac{1/2}{1/2} = \dfrac{1/2}{1 + \frac{1}{2}\sin\theta}$, where $e = \dfrac{1}{2}$ and $ed = \dfrac{1}{2} \quad \Rightarrow \quad d = 1$.

(a) Eccentricity $= e = \dfrac{1}{2}$

(b) Since $e = \dfrac{1}{2} < 1$, the conic is an ellipse.

(c) Since "$+e\sin\theta$" appears in the denominator, the directrix is above the focus at the origin, $d = |Fl| = 1$, so an equation of the directrix is $y = 1$.

(d) The vertices are $\left(\frac{1}{3}, \frac{\pi}{2}\right)$ and $\left(1, \frac{3\pi}{2}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $r = \dfrac{2}{3+3\sin\theta} \cdot \dfrac{1/3}{1/3} = \dfrac{2/3}{1+1\sin\theta}$, where $e = 1$ and $ed = \frac{2}{3} \quad \Rightarrow \quad d = \frac{2}{3}$.

(a) Eccentricity $= e = 1$

(b) Since $e = 1$, the conic is a parabola.

(c) Since "$+\ e\sin\theta$" appears in the denominator, the directrix is above the focus at the origin. $d = |Fl| = \frac{2}{3}$, so an equation of the directrix is $y = \frac{2}{3}$.

(d) The vertex is at $\left(\frac{1}{3}, \frac{\pi}{2}\right)$, midway between the focus and directrix.

18. $r = \dfrac{5}{2-4\cos\theta} \cdot \dfrac{1/2}{1/2} = \dfrac{5/2}{1-2\cos\theta}$, where $e = 2$ and $ed = \dfrac{5}{2} \quad \Rightarrow \quad d = \dfrac{5}{4}$.

(a) Eccentricity $= e = 2$

(b) Since $e = 2 > 1$, the conic is a hyperbola.

(c) Since "$-e\cos\theta$" appears in the denominator, the directrix is to the left the focus at the origin. $d = |Fl| = \frac{5}{4}$, so an equation of the directrix is $x = -\frac{5}{4}$.

(d) The vertices are $\left(-\frac{5}{2}, 0\right)$ and $\left(\frac{5}{6}, \pi\right)$, so the center is midway between them, that is, $\left(\frac{5}{3}, \pi\right)$.

19. $r = \dfrac{9}{6+2\cos\theta} \cdot \dfrac{1/6}{1/6} = \dfrac{3/2}{1+\frac{1}{3}\cos\theta}$, where $e = \frac{1}{3}$ and $ed = \frac{3}{2} \quad \Rightarrow \quad d = \frac{9}{2}$.

(a) Eccentricity $= e = \frac{1}{3}$

(b) Since $e = \frac{1}{3} < 1$, the conic is an ellipse.

(c) Since "$+e\cos\theta$" appears in the denominator, the directrix is to the right of the focus at the origin. $d = |Fl| = \frac{9}{2}$, so an equation of the directrix is $x = \frac{9}{2}$.

(d) The vertices are $\left(\frac{9}{8}, 0\right)$ and $\left(\frac{9}{4}, \pi\right)$, so the center is midway between them, that is, $\left(\frac{9}{16}, \pi\right)$.

20. $r = \dfrac{1}{3-3\sin\theta} \cdot \dfrac{1/3}{1/3} = \dfrac{1/3}{1-1\sin\theta}$, where $e = 1$ and $ed = \dfrac{1}{3} \quad \Rightarrow \quad d = \dfrac{1}{3}$.

(a) Eccentricity $= e = 1$

(b) Since $e = 1$, the conic is a parabola.

(c) Since "$-e\sin\theta$" appears in the denominator, the directrix is below the focus at the origin, $d = |Fl| = \frac{1}{3}$, so an equation of the directrix is $y = -\frac{1}{3}$.

(d) The vertex is at $\left(\frac{1}{6}, \frac{3\pi}{2}\right)$, midway between the focus and the directrix.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. $r = \dfrac{3}{4 - 8\cos\theta} \cdot \dfrac{1/4}{1/4} = \dfrac{3/4}{1 - 2\cos\theta}$, where $e = 2$ and $ed = \frac{3}{4} \quad \Rightarrow \quad d = \frac{3}{8}$.

(a) Eccentricity $= e = 2$

(b) Since $e = 2 > 1$, the conic is a hyperbola.

(c) Since "$-e\cos\theta$" appears in the denominator, the directrix is to the left of the focus at the origin. $d = |Fl| = \frac{3}{8}$, so an equation of the directrix is $x = -\frac{3}{8}$.

(d) The vertices are $\left(-\frac{3}{4}, 0\right)$ and $\left(\frac{1}{4}, \pi\right)$, so the center is midway between them, that is, $\left(\frac{1}{2}, \pi\right)$.

22. $r = \dfrac{4}{2 + 3\cos\theta} \cdot \dfrac{1/2}{1/2} = \dfrac{2}{1 + \frac{3}{2}\cos\theta}$, where $e = \dfrac{3}{2}$ and $ed = 2 \quad \Rightarrow \quad d = \dfrac{4}{3}$.

(a) Eccentricity $= e = \dfrac{3}{2}$

(b) Since $e = \dfrac{3}{2} > 1$, the conic is a hyperbola.

(c) Since "$+e\cos\theta$" appears in the denominator, the directrix is to the right of the focus at the origin. $d = |Fl| = \frac{4}{3}$, so an equation of the directrix is $x = \frac{4}{3}$.

(d) The vertices are $\left(\frac{4}{5}, 0\right)$ and $(-4, \pi)$, so the center is midway between them, that is, $\left(\frac{8}{5}, 0\right)$.

23. (a) $r = \dfrac{1}{1 - 2\sin\theta}$, where $e = 2$ and $ed = 1 \quad \Rightarrow \quad d = \frac{1}{2}$. The eccentricity $e = 2 > 1$, so the conic is a hyperbola. Since "$-e\sin\theta$" appears in the denominator, the directrix is below the focus at the origin. $d = |Fl| = \frac{1}{2}$, so an equation of the directrix is $y = -\frac{1}{2}$. The vertices are $\left(-1, \frac{\pi}{2}\right)$ and $\left(\frac{1}{3}, \frac{3\pi}{2}\right)$, so the center is midway between them, that is, $\left(\frac{2}{3}, \frac{3\pi}{2}\right)$.

(b) By the discussion that precedes Example 4, the equation is $r = \dfrac{1}{1 - 2\sin\left(\theta - \frac{3\pi}{4}\right)}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $r = \dfrac{4}{5 + 6\cos\theta} = \dfrac{4/5}{1 + \frac{6}{5}\cos\theta}$, so $e = \frac{6}{5}$ and $ed = \frac{4}{5} \quad \Rightarrow \quad d = \frac{2}{3}$.

An equation of the directrix is $x = \frac{2}{3} \quad \Rightarrow \quad r\cos\theta = \frac{2}{3} \quad \Rightarrow \quad r = \dfrac{2}{3\cos\theta}$.

If the hyperbola is rotated about its focus (the origin) through an angle $\pi/3$, its equation is the same as that of the original, with θ replaced by $\theta - \frac{\pi}{3}$ (see Example 4), so $r = \dfrac{4}{5 + 6\cos\left(\theta - \frac{\pi}{3}\right)}$.

25. $r = \dfrac{e}{1 - e\cos\theta}$. For $e < 1$ the curve is an ellipse. It is nearly circular when e is close to 0. As e increases, the graph is stretched out to the right, and grows larger (that is, its right-hand focus moves to the right while its left-hand focus remains at the origin.) At $e = 1$, the curve becomes a parabola with focus at the origin.

26. (a) $r = \dfrac{ed}{1 + e\sin\theta}$. The value of d does not seem to affect the shape of the conic (a parabola) at all, just its size, position, and orientation (for $d < 0$ it opens upward, for $d > 0$ it opens downward).

(b) We consider only positive values of e. When $0 < e < 1$, the conic is an ellipse. As $e \to 0^+$, the graph approaches perfect roundness and zero size. As e increases, the ellipse becomes more elongated, until at $e = 1$ it turns into a parabola. For $e > 1$, the conic is a hyperbola, which moves downward and gets broader as e continues to increase.

$e = 0.1$

$e = 0.5$

$e = 0.9$

$e = 1$

$e = 1.1$

$e = 1.5$

$e = 10$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. $|PF| = e\,|Pl| \quad\Rightarrow\quad r = e[d - r\cos(\pi - \theta)] = e(d + r\cos\theta) \quad\Rightarrow$

$r(1 - e\cos\theta) = ed \quad\Rightarrow\quad r = \dfrac{ed}{1 - e\cos\theta}$

28. $|PF| = e\,|Pl| \quad\Rightarrow\quad r = e[d - r\sin\theta] \quad\Rightarrow\quad r(1 + e\sin\theta) = ed \quad\Rightarrow$

$r = \dfrac{ed}{1 + e\sin\theta}$

29. $|PF| = e\,|Pl| \quad\Rightarrow\quad r = e[d - r\sin(\theta - \pi)] = e(d + r\sin\theta) \quad\Rightarrow$

$r(1 - e\sin\theta) = ed \quad\Rightarrow\quad r = \dfrac{ed}{1 - e\sin\theta}$

30. The parabolas intersect at the two points where $\dfrac{c}{1 + \cos\theta} = \dfrac{d}{1 - \cos\theta} \quad\Rightarrow\quad \cos\theta = \dfrac{c - d}{c + d} \quad\Rightarrow\quad r = \dfrac{c + d}{2}$.

For the first parabola, $\dfrac{dr}{d\theta} = \dfrac{c\sin\theta}{(1 + \cos\theta)^2}$, so

$$\frac{dy}{dx} = \frac{(dr/d\theta)\sin\theta + r\cos\theta}{(dr/d\theta)\cos\theta - r\sin\theta} = \frac{c\sin^2\theta + c\cos\theta(1 + \cos\theta)}{c\sin\theta\cos\theta - c\sin\theta(1 + \cos\theta)} = \frac{1 + \cos\theta}{-\sin\theta}$$

and similarly for the second, $\dfrac{dy}{dx} = \dfrac{1 - \cos\theta}{\sin\theta} = \dfrac{\sin\theta}{1 + \cos\theta}$. Since the product of these slopes is -1, the parabolas intersect at right angles.

31. We are given $e = 0.093$ and $a = 2.28 \times 10^8$. By (7), we have

$$r = \frac{a(1 - e^2)}{1 + e\cos\theta} = \frac{2.28 \times 10^8[1 - (0.093)^2]}{1 + 0.093\cos\theta} \approx \frac{2.26 \times 10^8}{1 + 0.093\cos\theta}$$

32. We are given $e = 0.048$ and $2a = 1.56 \times 10^9 \quad\Rightarrow\quad a = 7.8 \times 10^8$. By (7), we have

$$r = \frac{a(1 - e^2)}{1 + e\cos\theta} = \frac{7.8 \times 10^8[1 - (0.048)^2]}{1 + 0.048\cos\theta} \approx \frac{7.78 \times 10^8}{1 + 0.048\cos\theta}$$

33. Here $2a =$ length of major axis $= 36.18$ AU $\quad\Rightarrow\quad a = 18.09$ AU and $e = 0.97$. By (7), the equation of the orbit is

$r = \dfrac{18.09[1 - (0.97)^2]}{1 + 0.97\cos\theta} \approx \dfrac{1.07}{1 + 0.97\cos\theta}$. By (8), the maximum distance from the comet to the sun is

$18.09(1 + 0.97) \approx 35.64$ AU or about 3.314 billion miles.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. Here $2a =$ length of major axis $= 356.5$ AU $\Rightarrow$ $a = 178.25$ AU and $e = 0.9951$. By (7), the equation of the orbit is $r = \dfrac{178.25[1-(0.9951)^2]}{1+0.9951\cos\theta} \approx \dfrac{1.7426}{1+0.9951\cos\theta}$. By (8), the minimum distance from the comet to the sun is $178.25(1-0.9951) \approx 0.8734$ AU or about 81 million miles.

35. The minimum distance is at perihelion, where $4.6\times10^7 = r = a(1-e) = a(1-0.206) = a(0.794) \Rightarrow$ $a = 4.6\times10^7/0.794$. So the maximum distance, which is at aphelion, is $r = a(1+e) = \left(4.6\times10^7/0.794\right)(1.206) \approx 7.0\times10^7$ km.

36. At perihelion, $r = a(1-e) = 4.43\times10^9$, and at aphelion, $r = a(1+e) = 7.37\times10^9$. Adding, we get $2a = 11.80\times10^9$, so $a = 5.90\times10^9$ km. Therefore $1+e = a(1+e)/a = \frac{7.37}{5.90} \approx 1.249$ and $e \approx 0.249$.

37. From Exercise 35, we have $e = 0.206$ and $a(1-e) = 4.6\times10^7$ km. Thus, $a = 4.6\times10^7/0.794$. From (7), we can write the equation of Mercury's orbit as $r = a\dfrac{1-e^2}{1+e\cos\theta}$. So since

$$\frac{dr}{d\theta} = \frac{a(1-e^2)e\sin\theta}{(1+e\cos\theta)^2} \quad \Rightarrow$$

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = \frac{a^2(1-e^2)^2}{(1+e\cos\theta)^2} + \frac{a^2(1-e^2)^2\,e^2\sin^2\theta}{(1+e\cos\theta)^4} = \frac{a^2(1-e^2)^2}{(1+e\cos\theta)^4}\left(1+2e\cos\theta+e^2\right)$$

the length of the orbit is

$$L = \int_0^{2\pi}\sqrt{r^2+(dr/d\theta)^2}\,d\theta = a(1-e^2)\int_0^{2\pi}\frac{\sqrt{1+e^2+2e\cos\theta}}{(1+e\cos\theta)^2}\,d\theta \approx 3.6\times10^8 \text{ km}$$

This seems reasonable, since Mercury's orbit is nearly circular, and the circumference of a circle of radius a is $2\pi a \approx 3.6\times10^8$ km.

10 Review

TRUE-FALSE QUIZ

1. False. Consider the curve defined by $x = f(t) = (t-1)^3$ and $y = g(t) = (t-1)^2$. Then $g'(t) = 2(t-1)$, so $g'(1) = 0$, but its graph has a *vertical* tangent when $t = 1$. *Note:* The statement is true if $f'(1) \neq 0$ when $g'(1) = 0$.

2. False. If $x = f(t)$ and $y = g(t)$ are twice differentiable, then $\dfrac{d^2y}{dx^2} = \dfrac{d}{dx}\left(\dfrac{dy}{dx}\right) = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{\dfrac{dx}{dt}}$.

3. False. For example, if $f(t) = \cos t$ and $g(t) = \sin t$ for $0 \le t \le 4\pi$, then the curve is a circle of radius 1, hence its length is 2π, but $\int_0^{4\pi}\sqrt{[f'(t)]^2+[g'(t)]^2}\,dt = \int_0^{4\pi}\sqrt{(-\sin t)^2+(\cos t)^2}\,dt = \int_0^{4\pi}1\,dt = 4\pi$, since as t increases from 0 to 4π, the circle is traversed twice.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. False. The speed of the particle with parametric equations $x = 3t + 1$, $y = 2t^2 + 1$ is $\sqrt{(dx/dt)^2 + (dy/dt)^2} = \sqrt{3^2 + (4t)^2}$. When $t = 3$, the speed $= \sqrt{9 + 16(3)^2} = \sqrt{153}$. However, $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{4t}{3}$, so $\left.\dfrac{dy}{dx}\right|_{t=3} = \dfrac{4 \cdot 3}{3} = 4 \neq \sqrt{153}$.

5. False. If $(r, \theta) = (1, \pi)$, then $(x, y) = (-1, 0)$, so $\tan^{-1}(y/x) = \tan^{-1} 0 = 0 \neq \theta$. The statement is true for points in quadrants I and IV.

6. True. The curve $r = 1 - \sin 2\theta$ is unchanged if we rotate it through $180°$ about O because $1 - \sin 2(\theta + \pi) = 1 - \sin(2\theta + 2\pi) = 1 - \sin 2\theta$. So it's unchanged if we replace r by $-r$. (See the discussion after Example 8 in Section 10.3.) In other words, it's the same curve as $r = -(1 - \sin 2\theta) = \sin 2\theta - 1$.

7. True. The polar equation $r = 2$, the Cartesian equation $x^2 + y^2 = 4$, and the parametric equations $x = 2 \sin 3t$, $y = 2 \cos 3t$ $[0 \leq t \leq 2\pi]$ all describe the circle of radius 2 centered at the origin.

8. False. The first pair of equations, $x = t^2$ and $y = t^4$, gives the portion of the parabola $y = x^2$ with $x \geq 0$, whereas the second pair of equations, $x = t^3$ and $y = t^6$, traces out the whole parabola $y = x^2$.

9. True. $y^2 = 2y + 3x \quad \Leftrightarrow \quad (y - 1)^2 = 3x + 1 = 3\left(x + \frac{1}{3}\right) = 4\left(\frac{3}{4}\right)\left(x + \frac{1}{3}\right)$, which is the equation of a parabola with vertex $\left(-\frac{1}{3}, 1\right)$ and focus $\left(-\frac{1}{3} + \frac{3}{4}, 1\right)$, opening to the right.

10. True. By rotating and translating the parabola, we can assume it has an equation of the form $y = cx^2$, where $c > 0$. The tangent at the point $\left(a, ca^2\right)$ is the line $y - ca^2 = 2ca(x - a)$; i.e., $y = 2cax - ca^2$. This tangent meets the parabola at the points $\left(x, cx^2\right)$ where $cx^2 = 2cax - ca^2$. This equation is equivalent to $x^2 = 2ax - a^2$ [since $c > 0$]. But $x^2 = 2ax - a^2 \quad \Leftrightarrow \quad x^2 - 2ax + a^2 = 0 \quad \Leftrightarrow \quad (x - a)^2 = 0 \quad \Leftrightarrow \quad x = a \quad \Leftrightarrow$ $\left(x, cx^2\right) = \left(a, ca^2\right)$. This shows that each tangent meets the parabola at exactly one point.

11. True. Consider a hyperbola with focus at the origin, oriented so that its polar equation is $r = \dfrac{ed}{1 + e\cos\theta}$, where $e > 1$. The directrix is $x = d$, but along the hyperbola we have $x = r\cos\theta = \dfrac{ed\cos\theta}{1 + e\cos\theta} = d\left(\dfrac{e\cos\theta}{1 + e\cos\theta}\right) \neq d$.

EXERCISES

1. $x = t^2 + 4t$, $y = 2 - t$, $-4 \leq t \leq 1$.

Since $y = 2 - t$ and $-4 \leq t \leq 1$, we have $1 \leq y \leq 6$. $t = 2 - y$, so $x = (2 - y)^2 + 4(2 - y) = 4 - 4y + y^2 + 8 - 4y = y^2 - 8y + 12 \quad \Leftrightarrow$ $x + 4 = y^2 - 8y + 16 = (y - 4)^2$. This is part of a parabola with vertex $(-4, 4)$, opening to the right.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. $x = 1 + e^{2t}$, $y = e^t$.

$x = 1 + e^{2t} = 1 + (e^t)^2 = 1 + y^2$, $y > 0$.

3. $x = \ln t$, $t > 0 \quad \Rightarrow \quad t = e^x$.

$y = t^2 = (e^x)^2 = e^{2x}$.

4. $x = 2\cos\theta$, $y = 1 + \sin\theta$, $\cos^2\theta + \sin^2\theta = 1 \quad \Rightarrow$

$\left(\dfrac{x}{2}\right)^2 + (y-1)^2 = 1 \quad \Rightarrow \quad \dfrac{x^2}{4} + (y-1)^2 = 1$. This is an ellipse, centered at $(0, 1)$, with semimajor axis of length 2 and semiminor axis of length 1.

5. $y = \sec\theta = \dfrac{1}{\cos\theta} = \dfrac{1}{x}$. Since $0 \le \theta \le \pi/2$, $0 < x \le 1$ and $y \ge 1$.

This is part of the hyperbola $y = 1/x$.

6. $x = 2 + 4\cos\pi t$, $y = -3 + 4\sin\pi t \quad \Rightarrow \quad \cos\pi t = \dfrac{x-2}{4}$, $\sin\pi t = \dfrac{y+3}{4}$. $\cos^2\pi t + \sin^2\pi t = 1 \quad \Rightarrow$

$\left(\dfrac{x-2}{4}\right)^2 + \left(\dfrac{y+3}{4}\right)^2 = 1$, so the motion of the particle takes place on the circle centered at $(2, -3)$ with radius 4. As t goes from 0 to 4, the particle starts at $(6, -3)$ and moves counterclockwise along the circle $(x-2)^2 + (y+3)^2 = 16$, tracing the circle twice.

7. Three different sets of parametric equations for the curve $y = \sqrt{x}$ are

(i) $x = t, \quad y = \sqrt{t}$

(ii) $x = t^4, \quad y = t^2$

(iii) $x = \tan^2 t, \quad y = \tan t, \quad 0 \le t < \pi/2$

There are many other sets of equations that also give this curve.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. For $t < -1$, $x > 0$ and $y < 0$ with x decreasing and y increasing. When $t = -1$, $(x, y) = (0, 0)$. When $-1 < t < 0$, we have $-1 < x < 0$ and $0 < y < 1/2$. When $t = 0$, $(x, y) = (-1, 0)$. When $0 < t < 1$, $-1 < x < 0$ and $-\frac{1}{2} < y < 0$. When $t = 1$, $(x, y) = (0, 0)$ again. When $t > 1$, both x and y are positive and increasing.

9. (a)

The Cartesian coordinates are $x = 4\cos\frac{2\pi}{3} = 4\left(-\frac{1}{2}\right) = -2$ and $y = 4\sin\frac{2\pi}{3} = 4\left(\frac{\sqrt{3}}{2}\right) = 2\sqrt{3}$, that is, the point $\left(-2, 2\sqrt{3}\right)$.

(b) Given $x = -3$ and $y = 3$, we have $r = \sqrt{(-3)^2 + 3^2} = \sqrt{18} = 3\sqrt{2}$. Also, $\tan\theta = \dfrac{y}{x} \quad \Rightarrow \quad \tan\theta = \dfrac{3}{-3}$, and since $(-3, 3)$ is in the second quadrant, $\theta = \frac{3\pi}{4}$. Thus, one set of polar coordinates for $(-3, 3)$ is $\left(3\sqrt{2}, \frac{3\pi}{4}\right)$, and two others are $\left(3\sqrt{2}, \frac{11\pi}{4}\right)$ and $\left(-3\sqrt{2}, \frac{7\pi}{4}\right)$.

10. $1 \le r < 2$, $\frac{\pi}{6} \le \theta \le \frac{5\pi}{6}$

11. $r = 1 + \sin\theta$. This cardioid is symmetric about the $\theta = \pi/2$ axis.

12. $r = \sin 4\theta$. This is an eight-leaved rose.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $r = \cos 3\theta$. This is a three-leaved rose. The curve is traced twice.

14. $r = 3 + \cos 3\theta$. The curve is symmetric about the horizontal axis.

15. $r = 1 + \cos 2\theta$. The curve is symmetric about the pole and both the horizontal and vertical axes.

16. $r = 2\cos(\theta/2)$. The curve is symmetric about the pole and both the horizontal and vertical axes.

17. $r = \dfrac{3}{1 + 2\sin\theta} \quad \Rightarrow \quad e = 2 > 1$, so the conic is a hyperbola. $de = 3 \quad \Rightarrow$ $d = \frac{3}{2}$ and the form "$+2\sin\theta$" imply that the directrix is above the focus at the origin and has equation $y = \frac{3}{2}$. The vertices are $\left(1, \frac{\pi}{2}\right)$ and $\left(-3, \frac{3\pi}{2}\right)$.

18. $r = \dfrac{3}{2 - 2\cos\theta} \cdot \dfrac{1/2}{1/2} = \dfrac{3/2}{1 - 1\cos\theta} \quad \Rightarrow \quad e = 1$, so the conic is a parabola. $de = \frac{3}{2} \quad \Rightarrow \quad d = \frac{3}{2}$ and the form "$-2\cos\theta$" imply that the directrix is to the left of the focus at the origin and has equation $x = -\frac{3}{2}$. The vertex is $\left(\frac{3}{4}, \pi\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $x + y = 2 \quad \Leftrightarrow \quad r\cos\theta + r\sin\theta = 2 \quad \Leftrightarrow \quad r(\cos\theta + \sin\theta) = 2 \quad \Leftrightarrow \quad r = \dfrac{2}{\cos\theta + \sin\theta}$

20. $x^2 + y^2 = 2 \quad \Rightarrow \quad r^2 = 2 \quad \Rightarrow \quad r = \sqrt{2}$. [$r = -\sqrt{2}$ gives the same curve.]

21. $r = (\sin\theta)/\theta$. As $\theta \to \pm\infty$, $r \to 0$. As $\theta \to 0$, $r \to 1$. In the first figure, there are an infinite number of x-intercepts at $x = \pi n$, n a nonzero integer. These correspond to pole points in the second figure.

22. $r = 2$ when $\theta = 0$ and when $\theta = 2\pi$. r has a maximum value of approximately 2.6 at about $\theta = \frac{\pi}{6}$ and a minimum value of approximately -2.6 at about $\theta = \frac{5\pi}{6}$. The graph touches the pole ($r = 0$) when $\theta = \frac{\pi}{2}$ and $\frac{3\pi}{2}$. Since r is positive in the θ-intervals $\left(0, \frac{\pi}{2}\right)$ and $\left(\frac{3\pi}{2}, 2\pi\right)$ and negative in the interval $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$, the graph lies entirely in the first and fourth quadrants.

23. $x = \ln t$, $y = 1 + t^2$; $t = 1$. $\dfrac{dy}{dt} = 2t$ and $\dfrac{dx}{dt} = \dfrac{1}{t}$, so $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2t}{1/t} = 2t^2$.

When $t = 1$, $(x, y) = (0, 2)$ and $dy/dx = 2$.

24. $x = t^3 + 6t + 1,\ y = 2t - t^2;\ t = -1$. $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{2 - 2t}{3t^2 + 6}$. When $t = -1$, $(x, y) = (-6, -3)$ and $\dfrac{dy}{dx} = \dfrac{4}{9}$.

25. $r = e^{-\theta} \quad \Rightarrow \quad y = r\sin\theta = e^{-\theta}\sin\theta$ and $x = r\cos\theta = e^{-\theta}\cos\theta \quad \Rightarrow$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\frac{dr}{d\theta}\sin\theta + r\cos\theta}{\frac{dr}{d\theta}\cos\theta - r\sin\theta} = \frac{-e^{-\theta}\sin\theta + e^{-\theta}\cos\theta}{-e^{-\theta}\cos\theta - e^{-\theta}\sin\theta} \cdot \frac{-e^{\theta}}{-e^{\theta}} = \frac{\sin\theta - \cos\theta}{\cos\theta + \sin\theta}.$$

When $\theta = \pi$, $\dfrac{dy}{dx} = \dfrac{0 - (-1)}{-1 + 0} = \dfrac{1}{-1} = -1$.

26. $r = 3 + \cos 3\theta \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{dy/d\theta}{dx/d\theta} = \dfrac{\frac{dr}{d\theta}\sin\theta + r\cos\theta}{\frac{dr}{d\theta}\cos\theta - r\sin\theta} = \dfrac{-3\sin 3\theta\sin\theta + (3 + \cos 3\theta)\cos\theta}{-3\sin 3\theta\cos\theta - (3 + \cos 3\theta)\sin\theta}$.

When $\theta = \pi/2$, $\dfrac{dy}{dx} = \dfrac{(-3)(-1)(1) + (3 + 0)\cdot 0}{(-3)(-1)(0) - (3 + 0)\cdot 1} = \dfrac{3}{-3} = -1$.

27. $x = t + \sin t,\ y = t - \cos t \quad \Rightarrow \quad \dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{1 + \sin t}{1 + \cos t} \quad \Rightarrow$

$$\frac{d^2y}{dx^2} = \frac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \frac{\dfrac{(1 + \cos t)\cos t - (1 + \sin t)(-\sin t)}{(1 + \cos t)^2}}{1 + \cos t} = \frac{\cos t + \cos^2 t + \sin t + \sin^2 t}{(1 + \cos t)^3} = \frac{1 + \cos t + \sin t}{(1 + \cos t)^3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $x = 1 + t^2$, $y = t - t^3$. $\dfrac{dy}{dt} = 1 - 3t^2$ and $\dfrac{dx}{dt} = 2t$, so $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{1 - 3t^2}{2t} = \frac{1}{2}t^{-1} - \frac{3}{2}t$.

$$\frac{d^2y}{dx^2} = \frac{d(dy/dx)/dt}{dx/dt} = \frac{-\frac{1}{2}t^{-2} - \frac{3}{2}}{2t} = -\tfrac{1}{4}t^{-3} - \tfrac{3}{4}t^{-1} = -\frac{1}{4t^3}\left(1 + 3t^2\right) = -\frac{3t^2 + 1}{4t^3}.$$

29. We graph the curve $x = t^3 - 3t$, $y = t^2 + t + 1$ for $-2.2 \le t \le 1.2$. By zooming in or using a cursor, we find that the lowest point is about $(1.4, 0.75)$. To find the exact values, we find the t-value at which $dy/dt = 2t + 1 = 0 \quad \Leftrightarrow \quad t = -\frac{1}{2} \quad \Leftrightarrow \quad (x, y) = \left(\frac{11}{8}, \frac{3}{4}\right)$.

30. We estimate the coordinates of the point of intersection to be $(-2, 3)$. In fact this is exact, since both $t = -2$ and $t = 1$ give the point $(-2, 3)$. So the area enclosed by the loop is

$$\int_{t=-2}^{t=1} y\,dx = \int_{-2}^{1} (t^2 + t + 1)(3t^2 - 3)\,dt = \int_{-2}^{1} (3t^4 + 3t^3 - 3t - 3)\,dt$$
$$= \left[\tfrac{3}{5}t^5 + \tfrac{3}{4}t^4 - \tfrac{3}{2}t^2 - 3t\right]_{-2}^{1} = \left(\tfrac{3}{5} + \tfrac{3}{4} - \tfrac{3}{2} - 3\right) - \left[-\tfrac{96}{5} + 12 - 6 - (-6)\right] = \tfrac{81}{20}$$

31. $x = 2a\cos t - a\cos 2t \quad \Rightarrow \quad \dfrac{dx}{dt} = -2a\sin t + 2a\sin 2t = 2a\sin t(2\cos t - 1) = 0 \quad \Leftrightarrow$

$\sin t = 0$ or $\cos t = \frac{1}{2} \quad \Rightarrow \quad t = 0, \frac{\pi}{3}, \pi$, or $\frac{5\pi}{3}$.

$y = 2a\sin t - a\sin 2t \quad \Rightarrow \quad \dfrac{dy}{dt} = 2a\cos t - 2a\cos 2t = 2a\left(1 + \cos t - 2\cos^2 t\right) = 2a(1 - \cos t)(1 + 2\cos t) = 0 \quad \Rightarrow$

$t = 0, \frac{2\pi}{3}$, or $\frac{4\pi}{3}$.

Thus the graph has vertical tangents where $t = \frac{\pi}{3}, \pi$ and $\frac{5\pi}{3}$, and horizontal tangents where $t = \frac{2\pi}{3}$ and $\frac{4\pi}{3}$. To determine what the slope is where $t = 0$, we use l'Hospital's Rule to evaluate $\displaystyle\lim_{t \to 0} \frac{dy/dt}{dx/dt} = 0$, so there is a horizontal tangent there.

t	x	y
0	a	0
$\frac{\pi}{3}$	$\frac{3}{2}a$	$\frac{\sqrt{3}}{2}a$
$\frac{2\pi}{3}$	$-\frac{1}{2}a$	$\frac{3\sqrt{3}}{2}a$
π	$-3a$	0
$\frac{4\pi}{3}$	$-\frac{1}{2}a$	$-\frac{3\sqrt{3}}{2}a$
$\frac{5\pi}{3}$	$\frac{3}{2}a$	$-\frac{\sqrt{3}}{2}a$

32. From Exercise 31, $x = 2a\cos t - a\cos 2t$, $y = 2a\sin t - a\sin 2t \quad \Rightarrow$

$$A = 2\int_{\pi}^{0} (2a\sin t - a\sin 2t)(-2a\sin t + 2a\sin 2t)\,dt = 4a^2 \int_0^{\pi} (2\sin^2 t + \sin^2 2t - 3\sin t\sin 2t)\,dt$$
$$= 4a^2 \int_0^{\pi} \left[(1 - \cos 2t) + \tfrac{1}{2}(1 - \cos 4t) - 6\sin^2 t\,\cos t\right] dt = 4a^2\left[t - \tfrac{1}{2}\sin 2t + \tfrac{1}{2}t - \tfrac{1}{8}\sin 4t - 2\sin^3 t\right]_0^{\pi}$$
$$= 4a^2\left(\tfrac{3}{2}\right)\pi = 6\pi a^2$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. The curve $r^2 = 9\cos 5\theta$ has 10 "petals." For instance, for $-\frac{\pi}{10} \le \theta \le \frac{\pi}{10}$, there are two petals, one with $r > 0$ and one with $r < 0$.

$$A = 10\int_{-\pi/10}^{\pi/10} \tfrac{1}{2}r^2\,d\theta = 5\int_{-\pi/10}^{\pi/10} 9\cos 5\theta\,d\theta = 5\cdot 9\cdot 2\int_0^{\pi/10} \cos 5\theta\,d\theta = 18\big[\sin 5\theta\big]_0^{\pi/10} = 18$$

34. $r = 1 - 3\sin\theta$. The inner loop is traced out as θ goes from $\alpha = \sin^{-1}\left(\frac{1}{3}\right)$ to $\pi - \alpha$, so

$$\begin{aligned} A &= \textstyle\int_\alpha^{\pi-\alpha} \frac{1}{2}r^2\,d\theta = \int_\alpha^{\pi/2}(1-3\sin\theta)^2\,d\theta = \int_\alpha^{\pi/2}\left[1 - 6\sin\theta + \frac{9}{2}(1-\cos 2\theta)\right]d\theta \\ &= \left[\tfrac{11}{2}\theta + 6\cos\theta - \tfrac{9}{4}\sin 2\theta\right]_\alpha^{\pi/2} = \tfrac{11}{4}\pi - \tfrac{11}{2}\sin^{-1}\left(\tfrac{1}{3}\right) - 3\sqrt{2} \end{aligned}$$

35. The curves intersect when $4\cos\theta = 2 \;\Rightarrow\; \cos\theta = \frac{1}{2} \;\Rightarrow\; \theta = \pm\frac{\pi}{3}$ for $-\pi \le \theta \le \pi$. The points of intersection are $\left(2, \frac{\pi}{3}\right)$ and $\left(2, -\frac{\pi}{3}\right)$.

36. The two curves clearly both contain the pole. For other points of intersection, $\cot\theta = 2\cos(\theta + 2n\pi)$ or $-2\cos(\theta + \pi + 2n\pi)$, both of which reduce to $\cot\theta = 2\cos\theta \;\Leftrightarrow\; \cos\theta = 2\sin\theta\cos\theta \;\Leftrightarrow\; \cos\theta(1 - 2\sin\theta) = 0 \;\Rightarrow$ $\cos\theta = 0$ or $\sin\theta = \frac{1}{2} \;\Rightarrow\; \theta = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}$ or $\frac{3\pi}{2} \;\Rightarrow$ intersection points are $\left(0, \frac{\pi}{2}\right)$, $\left(\sqrt{3}, \frac{\pi}{6}\right)$, and $\left(\sqrt{3}, \frac{11\pi}{6}\right)$.

37. The curves intersect where $2\sin\theta = \sin\theta + \cos\theta \;\Rightarrow$ $\sin\theta = \cos\theta \;\Rightarrow \theta = \frac{\pi}{4}$, and also at the origin (at which $\theta = \frac{3\pi}{4}$ on the second curve).

$$\begin{aligned} A &= \textstyle\int_0^{\pi/4} \frac{1}{2}(2\sin\theta)^2\,d\theta + \int_{\pi/4}^{3\pi/4} \frac{1}{2}(\sin\theta + \cos\theta)^2\,d\theta \\ &= \textstyle\int_0^{\pi/4}(1 - \cos 2\theta)\,d\theta + \frac{1}{2}\int_{\pi/4}^{3\pi/4}(1 + \sin 2\theta)\,d\theta \\ &= \left[\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/4} + \left[\tfrac{1}{2}\theta - \tfrac{1}{4}\cos 2\theta\right]_{\pi/4}^{3\pi/4} = \tfrac{1}{2}(\pi - 1) \end{aligned}$$

38.

$$\begin{aligned} A &= 2\textstyle\int_{-\pi/2}^{\pi/6} \frac{1}{2}\left[(2 + \cos 2\theta)^2 - (2 + \sin\theta)^2\right]d\theta \\ &= \textstyle\int_{-\pi/2}^{\pi/6}\left[4\cos 2\theta + \cos^2 2\theta - 4\sin\theta - \sin^2\theta\right]d\theta \\ &= \left[2\sin 2\theta + \tfrac{1}{2}\theta + \tfrac{1}{8}\sin 4\theta + 4\cos\theta - \tfrac{1}{2}\theta + \tfrac{1}{4}\sin 2\theta\right]_{-\pi/2}^{\pi/6} \\ &= \tfrac{51}{16}\sqrt{3} \end{aligned}$$

39. $x = 3t^2,\ y = 2t^3$.

$$\begin{aligned} L &= \textstyle\int_0^2 \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt = \int_0^2 \sqrt{(6t)^2 + (6t^2)^2}\,dt = \int_0^2 \sqrt{36t^2 + 36t^4}\,dt = \int_0^2 \sqrt{36t^2}\sqrt{1 + t^2}\,dt \\ &= \textstyle\int_0^2 6\,|t|\sqrt{1 + t^2}\,dt = 6\int_0^2 t\sqrt{1 + t^2}\,dt = 6\int_1^5 u^{1/2}\left(\frac{1}{2}du\right) \qquad \left[u = 1 + t^2,\ du = 2t\,dt\right] \\ &= 6\cdot\tfrac{1}{2}\cdot\tfrac{2}{3}\left[u^{3/2}\right]_1^5 = 2(5^{3/2} - 1) = 2\left(5\sqrt{5} - 1\right) \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $x = 2 + 3t, \quad y = \cosh 3t \quad \Rightarrow \quad (dx/dt)^2 + (dy/dt)^2 = 3^2 + (3\sinh 3t)^2 = 9(1 + \sinh^2 3t) = 9\cosh^2 3t$, so

$L = \int_0^1 \sqrt{9\cosh^2 3t}\, dt = \int_0^1 |3\cosh 3t|\ dt = \int_0^1 3\cosh 3t\, dt = \big[\sinh 3t\big]_0^1 = \sinh 3 - \sinh 0 = \sinh 3.$

41. $L = \int_\pi^{2\pi} \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_\pi^{2\pi} \sqrt{(1/\theta)^2 + (-1/\theta^2)^2}\, d\theta = \displaystyle\int_\pi^{2\pi} \frac{\sqrt{\theta^2+1}}{\theta^2}\, d\theta$

$$\overset{24}{=} \left[-\frac{\sqrt{\theta^2+1}}{\theta} + \ln\Big(\theta + \sqrt{\theta^2+1}\Big)\right]_\pi^{2\pi} = \frac{\sqrt{\pi^2+1}}{\pi} - \frac{\sqrt{4\pi^2+1}}{2\pi} + \ln\left(\frac{2\pi + \sqrt{4\pi^2+1}}{\pi + \sqrt{\pi^2+1}}\right)$$

$$= \frac{2\sqrt{\pi^2+1} - \sqrt{4\pi^2+1}}{2\pi} + \ln\left(\frac{2\pi + \sqrt{4\pi^2+1}}{\pi + \sqrt{\pi^2+1}}\right)$$

42. $L = \int_0^\pi \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_0^\pi \sqrt{\left[\sin^3\left(\frac{1}{3}\theta\right)\right]^2 + \left[\sin^2\left(\frac{1}{3}\theta\right)\cos\left(\frac{1}{3}\theta\right)\right]^2}\, d\theta$

$= \int_0^\pi \sqrt{\sin^6\left(\frac{1}{3}\theta\right) + \sin^4\left(\frac{1}{3}\theta\right)\cos^2\left(\frac{1}{3}\theta\right)}\, d\theta$

$= \int_0^\pi \sqrt{\sin^4\left(\frac{1}{3}\theta\right)\left[\sin^2\left(\frac{1}{3}\theta\right) + \cos^2\left(\frac{1}{3}\theta\right)\right]}\, d\theta = \int_0^\pi \sin^2\left(\frac{1}{3}\theta\right) d\theta$

$= \int_0^\pi \frac{1}{2}\left[1 - \cos\left(\frac{2}{3}\theta\right)\right] d\theta = \left[\frac{1}{2}\left(\theta - \frac{3}{2}\sin\left(\frac{2}{3}\theta\right)\right)\right]_0^\pi = \frac{1}{2}\pi - \frac{3}{8}\sqrt{3}$

43. (a) $x = \frac{1}{2}(t^2+3),\ y = 5 - \frac{1}{3}t^3.\ dx/dt = t$ and $dy/dt = -t^2$, so the speed at time t is the function $v(t) = s'(t) = \sqrt{t^2 + (-t^2)^2}$. At the point $(6, -4)$, $y = 5 - \frac{1}{3}t^3 = -4 \quad \Rightarrow \quad 9 = \frac{1}{3}t^3 \quad \Rightarrow \quad t^3 = 27 \quad \Rightarrow \quad t = 3$.

Thus, the speed of the particle at the point $(6, -4)$ is $v(3) = \sqrt{3^2 + 3^4} = \sqrt{90} \approx 9.49\,\text{m/s}$.

(b) To find the average speed of the particle for $0 \le t \le 8$, we find the total distance L traveled in this time, and divide it by the length of the interval. By Theorem 10.2.5,

$$L = \int_0^8 \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\, dt = \int_0^8 \sqrt{t^2 + t^4}\, dt = \int_0^8 t\sqrt{1+t^2}\, dt$$

$$= \tfrac{1}{2}\int_1^{65} \sqrt{u}\, du \qquad [u = 1 + t^2,\ du = 2t\, dt]$$

$$= \tfrac{1}{2}\left[\tfrac{2}{3}u^{3/2}\right]_1^{65} = \tfrac{1}{3}(65^{3/2} - 1)$$

Thus, the average speed is $\dfrac{L}{8} = \dfrac{1}{24}\left(65\sqrt{65} - 1\right) \approx 21.79\ \text{m/s}$.

44. (a) We see from the figure in the text that the blue section of the curve $r = 2\cos^2(\theta/2)$ is traced from $\theta = \pi/2$ to $\theta = \pi$. Thus, its length is given by

$$L = \int_{\pi/2}^{\pi} \sqrt{r^2 + (dr/d\theta)^2}\, d\theta = \int_{\pi/2}^{\pi} \sqrt{[2\cos^2(\theta/2)]^2 + [-2\sin(\theta/2)\cos(\theta/2)]^2}\, d\theta$$

$$= \int_{\pi/2}^{\pi} \sqrt{4\cos^4(\theta/2) + 4\sin^2(\theta/2)\cos^2(\theta/2)}\, d\theta = \int_{\pi/2}^{\pi} |2\cos(\theta/2)|\sqrt{\cos^2(\theta/2) + \sin^2(\theta/2)}\, d\theta$$

$$= \int_{\pi/2}^{\pi} 2\cos(\theta/2)\cdot 1\, d\theta = \Big[4\sin(\theta/2)\Big]_{\pi/2}^{\pi} = 4\cdot 1 - 4\cdot\frac{1}{\sqrt{2}} = 4 - 2\sqrt{2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $A = \int_{\pi/2}^{\pi} \frac{1}{2} r^2 \, d\theta = \int_{\pi/2}^{\pi} \frac{1}{2} [2\cos^2(\theta/2)]^2 \, d\theta = \frac{1}{2} \int_{\pi/2}^{\pi} (1+\cos\theta)^2 \, d\theta = \frac{1}{2} \int_{\pi/2}^{\pi} (1 + 2\cos\theta + \cos^2\theta) \, d\theta$

$$= \frac{1}{2} \int_{\pi/2}^{\pi} \left[1 + 2\cos\theta + \frac{1}{2}(1+\cos 2\theta)\right] d\theta = \frac{1}{2}\left[\theta + 2\sin\theta + \frac{1}{2}\left(\theta + \frac{1}{2}\sin 2\theta\right)\right]_{\pi/2}^{\pi}$$

$$= \frac{1}{2}\left[\pi + 0 + \frac{1}{2}(\pi + 0) - \frac{\pi}{2} - 2 - \frac{1}{2}\left(\frac{\pi}{2} + 0\right)\right] = \frac{1}{2}\left(\frac{3\pi}{4} - 2\right) = \frac{3\pi}{8} - 1$$

45. $x = 4\sqrt{t}, \; y = \dfrac{t^3}{3} + \dfrac{1}{2t^2}, \; 1 \le t \le 4 \quad \Rightarrow$

$$S = \int_1^4 2\pi y \sqrt{(dx/dt)^2 + (dy/dt)^2} \, dt = \int_1^4 2\pi \left(\tfrac{1}{3}t^3 + \tfrac{1}{2}t^{-2}\right) \sqrt{\left(2/\sqrt{t}\right)^2 + (t^2 - t^{-3})^2} \, dt$$

$$= 2\pi \int_1^4 \left(\tfrac{1}{3}t^3 + \tfrac{1}{2}t^{-2}\right) \sqrt{(t^2 + t^{-3})^2} \, dt = 2\pi \int_1^4 \left(\tfrac{1}{3}t^5 + \tfrac{5}{6} + \tfrac{1}{2}t^{-5}\right) dt = 2\pi \left[\tfrac{1}{18}t^6 + \tfrac{5}{6}t - \tfrac{1}{8}t^{-4}\right]_1^4 = \tfrac{471{,}295}{1024}\pi$$

46. $x = 2 + 3t, \quad y = \cosh 3t \quad \Rightarrow \quad (dx/dt)^2 + (dy/dt)^2 = 3^2 + (3\sinh 3t)^2 = 9(1 + \sinh^2 3t) = 9\cosh^2 3t$, so

$$S = \int_0^1 2\pi y \, ds = \int_0^1 2\pi \cosh 3t \sqrt{9\cosh^2 3t} \, dt = \int_0^1 2\pi \cosh 3t \, |3\cosh 3t| \, dt = \int_0^1 2\pi \cosh 3t \cdot 3\cosh 3t \, dt$$

$$= 6\pi \int_0^1 \cosh^2 3t \, dt = 6\pi \int_0^1 \tfrac{1}{2}(1 + \cosh 6t) \, dt = 3\pi \left[t + \tfrac{1}{6}\sinh 6t\right]_0^1 = 3\pi\left(1 + \tfrac{1}{6}\sinh 6\right) = 3\pi + \tfrac{\pi}{2}\sinh 6$$

47. $x = \dfrac{t^2 - c}{t^2 + 1}, \; y = \dfrac{t(t^2 - c)}{t^2 + 1}$. For all c except -1, the curve is asymptotic to the line $x = 1$. For $c < -1$, the curve bulges to the right near $y = 0$. As c increases, the bulge becomes smaller, until at $c = -1$ the curve is the straight line $x = 1$. As c continues to increase, the curve bulges to the left, until at $c = 0$ there is a cusp at the origin. For $c > 0$, there is a loop to the left of the origin, whose size and roundness increase as c increases. Note that the x-intercept of the curve is always $-c$.

48. For a close to 0, the graph of $r^a = |\sin 2\theta|$ consists of four thin petals. As a increases, the petals get wider, until as $a \to \infty$, each petal occupies almost its entire quarter-circle.

$a = 0.01$ $\quad$ $a = 0.1$ $\quad$ $a = 1$

$a = 5$ $\quad$ $a = 10$ $\quad$ $a = 25$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. $\dfrac{x^2}{9}+\dfrac{y^2}{8}=1$ is an ellipse with center $(0,0)$.
$a=3$, $b=2\sqrt{2}$, $c=1 \quad \Rightarrow$
foci $(\pm 1,0)$, vertices $(\pm 3,0)$.

50. $4x^2-y^2=16 \quad \Leftrightarrow \quad \dfrac{x^2}{4}-\dfrac{y^2}{16}=1$ is a hyperbola with center $(0,0)$, vertices $(\pm 2,0)$, $a=2$, $b=4$, $c=\sqrt{16+4}=2\sqrt{5}$, foci $\left(\pm 2\sqrt{5},0\right)$ and asymptotes $y=\pm 2x$.

51. $6y^2+x-36y+55=0 \quad \Leftrightarrow$
$6(y^2-6y+9)=-(x+1) \quad \Leftrightarrow$
$(y-3)^2=-\frac{1}{6}(x+1)$, a parabola with vertex $(-1,3)$, opening to the left, $p=-\frac{1}{24} \quad \Rightarrow \quad$ focus $\left(-\frac{25}{24},3\right)$ and directrix $x=-\frac{23}{24}$.

52. $25x^2+4y^2+50x-16y=59 \quad \Leftrightarrow$
$25(x+1)^2+4(y-2)^2=100 \quad \Leftrightarrow$
$\frac{1}{4}(x+1)^2+\frac{1}{25}(y-2)^2=1$ is an ellipse centered at $(-1,2)$ with foci on the line $x=-1$, vertices $(-1,7)$ and $(-1,-3)$; $a=5$, $b=2 \quad \Rightarrow \quad c=\sqrt{21} \quad \Rightarrow$
foci $\left(-1,2\pm\sqrt{21}\right)$.

53. The ellipse with foci $(\pm 4,0)$ and vertices $(\pm 5,0)$ has center $(0,0)$ and a horizontal major axis, with $a=5$ and $c=4$, so $b^2=a^2-c^2=5^2-4^2=9$. An equation is $\dfrac{x^2}{25}+\dfrac{y^2}{9}=1$.

54. The distance from the focus $(2,1)$ to the directrix $x=-4$ is $2-(-4)=6$, so the distance from the focus to the vertex is $\frac{1}{2}(6)=3$ and the vertex is $(-1,1)$. Since the focus is to the right of the vertex, $p=3$. An equation is $(y-1)^2=4\cdot 3[x-(-1)]$, or $(y-1)^2=12(x+1)$.

55. The center of a hyperbola with foci $(0,\pm 4)$ is $(0,0)$, so $c=4$ and an equation is $\dfrac{y^2}{a^2}-\dfrac{x^2}{b^2}=1$.

The asymptote $y=3x$ has slope 3, so $\dfrac{a}{b}=\dfrac{3}{1} \quad \Rightarrow \quad a=3b$ and $a^2+b^2=c^2 \quad \Rightarrow \quad (3b)^2+b^2=4^2 \quad \Rightarrow$

$10b^2=16 \quad \Rightarrow \quad b^2=\frac{8}{5}$ and so $a^2=16-\frac{8}{5}=\frac{72}{5}$. Thus, an equation is $\dfrac{y^2}{72/5}-\dfrac{x^2}{8/5}=1$, or $\dfrac{5y^2}{72}-\dfrac{5x^2}{8}=1$.

56. The ellipse with foci $(3,\pm 2)$ has center $(3,0)$. $a=\frac{8}{2}=4$, $c=2 \quad \Rightarrow \quad b=\sqrt{a^2-c^2}=\sqrt{4^2-2^2}=\sqrt{12} \quad \Rightarrow$ an equation of the ellipse is $\dfrac{(x-3)^2}{12}+\dfrac{y^2}{16}=1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. $x^2 + y = 100 \quad \Leftrightarrow \quad x^2 = -(y - 100)$ has its vertex at $(0, 100)$, so one of the vertices of the ellipse is $(0, 100)$. Another form of the equation of a parabola is $x^2 = 4p(y - 100)$ so $4p(y - 100) = -(y - 100) \quad \Rightarrow \quad 4p = -1 \quad \Rightarrow \quad p = -\frac{1}{4}$. Therefore the shared focus is found at $\left(0, \frac{399}{4}\right)$ so $2c = \frac{399}{4} - 0 \quad \Rightarrow \quad c = \frac{399}{8}$ and the center of the ellipse is $\left(0, \frac{399}{8}\right)$. So $a = 100 - \frac{399}{8} = \frac{401}{8}$ and $b^2 = a^2 - c^2 = \dfrac{401^2 - 399^2}{8^2} = 25$. So the equation of the ellipse is $\dfrac{x^2}{b^2} + \dfrac{\left(y - \frac{399}{8}\right)^2}{a^2} = 1 \quad \Rightarrow$ $\dfrac{x^2}{25} + \dfrac{\left(y - \frac{399}{8}\right)^2}{\left(\frac{401}{8}\right)^2} = 1$, or $\dfrac{x^2}{25} + \dfrac{(8y - 399)^2}{160{,}801} = 1$.

58. $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 \quad \Rightarrow \quad \dfrac{2x}{a^2} + \dfrac{2y}{b^2}\dfrac{dy}{dx} = 0 \quad \Rightarrow \quad \dfrac{dy}{dx} = -\dfrac{b^2}{a^2}\dfrac{x}{y}$. Therefore $\dfrac{dy}{dx} = m \quad \Leftrightarrow \quad y = -\dfrac{b^2}{a^2}\dfrac{x}{m}$. Combining this condition with $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1$, we find that $x = \pm\dfrac{a^2 m}{\sqrt{a^2m^2 + b^2}}$. In other words, the two points on the ellipse where the tangent has slope m are $\left(\pm\dfrac{a^2 m}{\sqrt{a^2m^2 + b^2}}, \mp\dfrac{b^2}{\sqrt{a^2m^2 + b^2}}\right)$. The tangent lines at these points have the equations $y \pm \dfrac{b^2}{\sqrt{a^2m^2 + b^2}} = m\left(x \mp \dfrac{a^2 m}{\sqrt{a^2m^2 + b^2}}\right)$ or $y = mx \mp \dfrac{a^2m^2}{\sqrt{a^2m^2 + b^2}} \mp \dfrac{b^2}{\sqrt{a^2m^2 + b^2}} = mx \mp \sqrt{a^2m^2 + b^2}$.

59. Directrix $x = 4 \quad \Rightarrow \quad d = 4$, so $e = \frac{1}{3} \quad \Rightarrow \quad r = \dfrac{ed}{1 + e\cos\theta} = \dfrac{4}{3 + \cos\theta}$.

60. $r = \dfrac{2}{4 - 3\cos\theta} = \dfrac{1/2}{1 - \frac{3}{4}\cos\theta} \quad \Rightarrow \quad e = \frac{3}{4}$ and $d = \frac{2}{3}$. The equation of the directrix is $x = -\frac{2}{3} \quad \Rightarrow \quad r = -2/(3\cos\theta)$. To obtain the equation of the rotated ellipse, we replace θ in the original equation with $\theta - \frac{2\pi}{3}$, and get $r = \dfrac{2}{4 - 3\cos\left(\theta - \frac{2\pi}{3}\right)}$.

61. See the end of the proof of Theorem 10.6.1. If $e > 1$, then $1 - e^2 < 0$ and Equations 10.6.4 become $a^2 = \dfrac{e^2d^2}{(e^2 - 1)^2}$ and $b^2 = \dfrac{e^2d^2}{e^2 - 1}$, so $\dfrac{b^2}{a^2} = e^2 - 1$. The asymptotes $y = \pm\dfrac{b}{a}x$ have slopes $\pm\dfrac{b}{a} = \pm\sqrt{e^2 - 1}$, so the angles they make with the polar axis are $\pm\tan^{-1}\left[\sqrt{e^2 - 1}\right] = \cos^{-1}(\pm 1/e)$.

62. (a) If (a, b) lies on the curve, then there is some parameter value t_1 such that $\dfrac{3t_1}{1 + t_1^3} = a$ and $\dfrac{3t_1^2}{1 + t_1^3} = b$. If $t_1 = 0$, the point is $(0, 0)$, which lies on the line $y = x$. If $t_1 \neq 0$, then the point corresponding to $t = \dfrac{1}{t_1}$ is given by $x = \dfrac{3(1/t_1)}{1 + (1/t_1)^3} = \dfrac{3t_1^2}{t_1^3 + 1} = b$, $y = \dfrac{3(1/t_1)^2}{1 + (1/t_1)^3} = \dfrac{3t_1}{t_1^3 + 1} = a$. So (b, a) also lies on the curve. [Another way to see this is to do part (e) first; the result is immediate.] The curve intersects the line $y = x$ when $\dfrac{3t}{1 + t^3} = \dfrac{3t^2}{1 + t^3} \quad \Rightarrow$ $t = t^2 \quad \Rightarrow \quad t = 0$ or 1, so the points are $(0, 0)$ and $\left(\frac{3}{2}, \frac{3}{2}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\dfrac{dy}{dt} = \dfrac{(1+t^3)(6t) - 3t^2(3t^2)}{(1+t^3)^2} = \dfrac{6t - 3t^4}{(1+t^3)^2} = 0$ when $6t - 3t^4 = 3t(2-t^3) = 0 \quad \Rightarrow \quad t = 0$ or $t = \sqrt[3]{2}$, so there are horizontal tangents at $(0,0)$ and $\left(\sqrt[3]{2}, \sqrt[3]{4}\right)$. Using the symmetry from part (a), we see that there are vertical tangents at $(0,0)$ and $\left(\sqrt[3]{4}, \sqrt[3]{2}\right)$.

(c) Notice that as $t \to -1^+$, we have $x \to -\infty$ and $y \to \infty$. As $t \to -1^-$, we have $x \to \infty$ and $y \to -\infty$. Also $y - (-x-1) = y + x + 1 = \dfrac{3t + 3t^2 + (1+t^3)}{1+t^3} = \dfrac{(t+1)^3}{1+t^3} = \dfrac{(t+1)^2}{t^2 - t + 1} \to 0$ as $t \to -1$. So $y = -x - 1$ is a slant asymptote.

(d) $\dfrac{dx}{dt} = \dfrac{(1+t^3)(3) - 3t(3t^2)}{(1+t^3)^2} = \dfrac{3 - 6t^3}{(1+t^3)^2}$ and from part (b) we have $\dfrac{dy}{dt} = \dfrac{6t - 3t^4}{(1+t^3)^2}$. So $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{t(2-t^3)}{1-2t^3}$.

Also $\dfrac{d^2y}{dx^2} = \dfrac{\dfrac{d}{dt}\left(\dfrac{dy}{dx}\right)}{dx/dt} = \dfrac{2(1+t^3)^4}{3(1-2t^3)^3} > 0 \quad \Leftrightarrow \quad t < \dfrac{1}{\sqrt[3]{2}}$.

So the curve is concave upward there and has a minimum point at $(0,0)$ and a maximum point at $\left(\sqrt[3]{2}, \sqrt[3]{4}\right)$. Using this together with the information from parts (a), (b), and (c), we sketch the curve.

(e) $x^3 + y^3 = \left(\dfrac{3t}{1+t^3}\right)^3 + \left(\dfrac{3t^2}{1+t^3}\right)^3 = \dfrac{27t^3 + 27t^6}{(1+t^3)^3} = \dfrac{27t^3(1+t^3)}{(1+t^3)^3} = \dfrac{27t^3}{(1+t^3)^2}$

and $3xy = 3\left(\dfrac{3t}{1+t^3}\right)\left(\dfrac{3t^2}{1+t^3}\right) = \dfrac{27t^3}{(1+t^3)^2}$, so $x^3 + y^3 = 3xy$.

(f) We start with the equation from part (e) and substitute $x = r\cos\theta$, $y = r\sin\theta$. Then $x^3 + y^3 = 3xy \quad \Rightarrow$ $r^3\cos^3\theta + r^3\sin^3\theta = 3r^2\cos\theta\,\sin\theta$. For $r \neq 0$, this gives $r = \dfrac{3\cos\theta\,\sin\theta}{\cos^3\theta + \sin^3\theta}$. Dividing numerator and denominator by $\cos^3\theta$, we obtain $r = \dfrac{3\left(\dfrac{1}{\cos\theta}\right)\dfrac{\sin\theta}{\cos\theta}}{1 + \dfrac{\sin^3\theta}{\cos^3\theta}} = \dfrac{3\sec\theta\,\tan\theta}{1+\tan^3\theta}$.

(g) The loop corresponds to $\theta \in \left(0, \frac{\pi}{2}\right)$, so its area is

$$A = \int_0^{\pi/2} \frac{r^2}{2}\,d\theta = \frac{1}{2}\int_0^{\pi/2}\left(\frac{3\sec\theta\,\tan\theta}{1+\tan^3\theta}\right)^2 d\theta = \frac{9}{2}\int_0^{\pi/2}\frac{\sec^2\theta\,\tan^2\theta}{(1+\tan^3\theta)^2}\,d\theta = \frac{9}{2}\int_0^{\infty}\frac{u^2\,du}{(1+u^3)^2} \quad [\text{let } u = \tan\theta]$$
$$= \lim_{b\to\infty}\tfrac{9}{2}\left[-\tfrac{1}{3}(1+u^3)^{-1}\right]_0^b = \tfrac{3}{2}$$

(h) By symmetry, the area between the folium and the line $y = -x - 1$ is equal to the enclosed area in the third quadrant, plus twice the enclosed area in the fourth quadrant. The area in the third quadrant is $\frac{1}{2}$, and since $y = -x - 1 \quad \Rightarrow$ $r\sin\theta = -r\cos\theta - 1 \quad \Rightarrow \quad r = -\dfrac{1}{\sin\theta + \cos\theta}$, the area in the fourth quadrant is

$\dfrac{1}{2}\displaystyle\int_{-\pi/2}^{-\pi/4}\left[\left(-\frac{1}{\sin\theta + \cos\theta}\right)^2 - \left(\frac{3\sec\theta\,\tan\theta}{1+\tan^3\theta}\right)^2\right]d\theta \overset{\text{CAS}}{=} \dfrac{1}{2}$. Therefore, the total area is $\frac{1}{2} + 2\left(\frac{1}{2}\right) = \frac{3}{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1. See the figure. The circle with center $(-1, 0)$ and radius $\sqrt{2}$ has equation $(x+1)^2 + y^2 = 2$ and describes the circular arc from $(0, -1)$ to $(0, 1)$. Converting the equation to polar coordinates gives us

$(r\cos\theta + 1)^2 + (r\sin\theta)^2 = 2 \quad \Rightarrow$

$r^2\cos^2\theta + 2r\cos\theta + 1 + r^2\sin^2\theta = 2 \quad \Rightarrow$

$r^2(\cos^2\theta + \sin^2\theta) + 2r\cos\theta = 1 \quad \Rightarrow \quad r^2 + 2r\cos\theta = 1$. Using the quadratic formula to solve for r gives us

$r = \dfrac{-2\cos\theta \pm \sqrt{4\cos^2\theta + 4}}{2} = -\cos\theta + \sqrt{\cos^2\theta + 1}$ for $r > 0$.

The darkest shaded region is $\frac{1}{8}$ of the entire shaded region A, so $\frac{1}{8}A = \int_0^{\pi/4} \frac{1}{2}r^2\,d\theta = \frac{1}{2}\int_0^{\pi/4}(1 - 2r\cos\theta)\,d\theta \quad \Rightarrow$

$$\begin{aligned}
\tfrac{1}{4}A &= \int_0^{\pi/4}\left[1 - 2\cos\theta\left(-\cos\theta + \sqrt{\cos^2\theta + 1}\right)\right]d\theta = \int_0^{\pi/4}\left(1 + 2\cos^2\theta - 2\cos\theta\sqrt{\cos^2\theta + 1}\right)d\theta \\
&= \int_0^{\pi/4}\left[1 + 2\cdot\tfrac{1}{2}(1 + \cos 2\theta) - 2\cos\theta\sqrt{(1 - \sin^2\theta) + 1}\right]d\theta \\
&= \int_0^{\pi/4}(2 + \cos 2\theta)\,d\theta - 2\int_0^{\pi/4}\cos\theta\sqrt{2 - \sin^2\theta}\,d\theta \\
&= \left[2\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{\pi/4} - 2\int_0^{1/\sqrt{2}}\sqrt{2 - u^2}\,du \qquad \left[\begin{matrix} u = \sin\theta, \\ du = \cos\theta\,d\theta \end{matrix}\right] \\
&= \left(\frac{\pi}{2} + \frac{1}{2}\right) - (0 + 0) - 2\left[\frac{u}{2}\sqrt{2 - u^2} + \sin^{-1}\frac{u}{\sqrt{2}}\right]_0^{1/\sqrt{2}} \qquad \left[\begin{matrix}\text{Formula 30,} \\ a = \sqrt{2}\end{matrix}\right] \\
&= \frac{\pi}{2} + \frac{1}{2} - 2\left(\frac{1}{2\sqrt{2}}\cdot\frac{\sqrt{3}}{\sqrt{2}} + \frac{\pi}{6}\right) = \frac{\pi}{2} + \frac{1}{2} - \frac{1}{2}\sqrt{3} - \frac{\pi}{3} = \frac{\pi}{6} + \frac{1}{2} - \frac{1}{2}\sqrt{3}.
\end{aligned}$$

Thus, $A = 4\left(\dfrac{\pi}{6} + \dfrac{1}{2} - \dfrac{1}{2}\sqrt{3}\right) = \dfrac{2\pi}{3} + 2 - 2\sqrt{3}$.

2. (a) The curve $x^4 + y^4 = x^2 + y^2$ is symmetric about both axes and about the line $y = x$ (since interchanging x and y does not change the equation) so we need only consider $y \geq x \geq 0$ to begin with. Implicit differentiation gives

$4x^3 + 4y^3y' = 2x + 2yy' \quad \Rightarrow \quad y' = \dfrac{x(1 - 2x^2)}{y(2y^2 - 1)} \quad \Rightarrow \quad y' = 0$ when $x = 0$ and when $x = \pm\frac{1}{\sqrt{2}}$. If $x = 0$, then

$y^4 = y^2 \quad \Rightarrow \quad y^2(y^2 - 1) = 0 \quad \Rightarrow \quad y = 0$ or ± 1. The point $(0, 0)$ can't be a highest or lowest point because it is isolated. [If $-1 < x < 1$ and $-1 < y < 1$, then $x^4 < x^2$ and $y^4 < y^2 \quad \Rightarrow \quad x^4 + y^4 < x^2 + y^2$, except for $(0, 0)$.]

If $x = \frac{1}{\sqrt{2}}$, then $x^2 = \frac{1}{2}$, $x^4 = \frac{1}{4}$, so $\frac{1}{4} + y^4 = \frac{1}{2} + y^2 \quad \Rightarrow \quad 4y^4 - 4y^2 - 1 = 0 \quad \Rightarrow \quad y^2 = \frac{4 \pm \sqrt{16+16}}{8} = \frac{1 \pm \sqrt{2}}{2}$.

But $y^2 > 0$, so $y^2 = \frac{1+\sqrt{2}}{2} \quad \Rightarrow \quad y = \pm\sqrt{\frac{1}{2}\left(1 + \sqrt{2}\right)}$. Near the point $(0, 1)$, the denominator of y' is positive and the numerator changes from negative to positive as x increases through 0, so $(0, 1)$ is a local minimum point. At

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\left(\frac{1}{\sqrt{2}}, \sqrt{\frac{1+\sqrt{2}}{2}}\right)$, y' changes from positive to negative, so that point gives a maximum. By symmetry, the highest points on the curve are $\left(\pm\frac{1}{\sqrt{2}}, \sqrt{\frac{1+\sqrt{2}}{2}}\right)$ and the lowest points are $\left(\pm\frac{1}{\sqrt{2}}, -\sqrt{\frac{1+\sqrt{2}}{2}}\right)$.

(b) We use the information from part (a), together with symmetry with respect to the axes and the lines $y = \pm x$, to sketch the curve.

(c) In polar coordinates, $x^4 + y^4 = x^2 + y^2$ becomes $r^4\cos^4\theta + r^4\sin^4\theta = r^2$ or $r^2 = \dfrac{1}{\cos^4\theta + \sin^4\theta}$. By the symmetry shown in part (b), the area enclosed by the curve is $A = 8\displaystyle\int_0^{\pi/4} \frac{1}{2} r^2\, d\theta = 4\int_0^{\pi/4} \frac{d\theta}{\cos^4\theta + \sin^4\theta} \overset{\text{CAS}}{=} \sqrt{2}\,\pi$.

3. In terms of x and y, we have $x = r\cos\theta = (1 + c\sin\theta)\cos\theta = \cos\theta + c\sin\theta\cos\theta = \cos\theta + \frac{1}{2}c\sin 2\theta$ and $y = r\sin\theta = (1 + c\sin\theta)\sin\theta = \sin\theta + c\sin^2\theta$. Now $-1 \le \sin\theta \le 1 \;\Rightarrow\; -1 \le \sin\theta + c\sin^2\theta \le 1 + c \le 2$, so $-1 \le y \le 2$. Furthermore, $y = 2$ when $c = 1$ and $\theta = \frac{\pi}{2}$, while $y = -1$ for $c = 0$ and $\theta = \frac{3\pi}{2}$. Therefore, we need a viewing rectangle with $-1 \le y \le 2$.

To find the x-values, look at the equation $x = \cos\theta + \frac{1}{2}c\sin 2\theta$ and use the fact that $\sin 2\theta \ge 0$ for $0 \le \theta \le \frac{\pi}{2}$ and $\sin 2\theta \le 0$ for $-\frac{\pi}{2} \le \theta \le 0$. [Because $r = 1 + c\sin\theta$ is symmetric about the y-axis, we only need to consider $-\frac{\pi}{2} \le \theta \le \frac{\pi}{2}$.] So for $-\frac{\pi}{2} \le \theta \le 0$, x has a maximum value when $c = 0$ and then $x = \cos\theta$ has a maximum value of 1 at $\theta = 0$. Thus, the maximum value of x must occur on $\left[0, \frac{\pi}{2}\right]$ with $c = 1$. Then $x = \cos\theta + \frac{1}{2}\sin 2\theta \;\Rightarrow$ $\frac{dx}{d\theta} = -\sin\theta + \cos 2\theta = -\sin\theta + 1 - 2\sin^2\theta \;\Rightarrow\; \frac{dx}{d\theta} = -(2\sin\theta - 1)(\sin\theta + 1) = 0$ when $\sin\theta = -1$ or $\frac{1}{2}$ [but $\sin\theta \ne -1$ for $0 \le \theta \le \frac{\pi}{2}$]. If $\sin\theta = \frac{1}{2}$, then $\theta = \frac{\pi}{6}$ and $x = \cos\frac{\pi}{6} + \frac{1}{2}\sin\frac{\pi}{3} = \frac{3}{4}\sqrt{3}$. Thus, the maximum value of x is $\frac{3}{4}\sqrt{3}$, and, by symmetry, the minimum value is $-\frac{3}{4}\sqrt{3}$. Therefore, the smallest viewing rectangle that contains every member of the family of polar curves $r = 1 + c\sin\theta$, where $0 \le c \le 1$, is $\left[-\frac{3}{4}\sqrt{3}, \frac{3}{4}\sqrt{3}\right] \times [-1, 2]$.

4. (a) Let us find the polar equation of the path of the bug that starts in the upper right corner of the square. If the polar coordinates of this bug, at a particular moment, are (r, θ), then the polar coordinates of the bug that it is crawling toward must be $\left(r, \theta + \frac{\pi}{2}\right)$. (The next bug must be the same distance from the origin and the angle between the lines joining the bugs to the pole must be $\frac{\pi}{2}$.) The Cartesian coordinates of the first bug are $(r\cos\theta, r\sin\theta)$ and for the second bug we have

$x = r\cos\left(\theta + \frac{\pi}{2}\right) = -r\sin\theta$, $y = r\sin\left(\theta + \frac{\pi}{2}\right) = r\cos\theta$. So the slope of the line joining the bugs is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\dfrac{r\cos\theta - r\sin\theta}{-r\sin\theta - r\cos\theta} = \dfrac{\sin\theta - \cos\theta}{\sin\theta + \cos\theta}$. This must be equal to the slope of the tangent line at (r, θ), so by Equation 10.4.7 we have $\dfrac{(dr/d\theta)\sin\theta + r\cos\theta}{(dr/d\theta)\cos\theta - r\sin\theta} = \dfrac{\sin\theta - \cos\theta}{\sin\theta + \cos\theta}$. Solving for $\dfrac{dr}{d\theta}$, we get

$\dfrac{dr}{d\theta}\sin^2\theta + \dfrac{dr}{d\theta}\sin\theta\cos\theta + r\sin\theta\cos\theta + r\cos^2\theta = \dfrac{dr}{d\theta}\sin\theta\cos\theta - \dfrac{dr}{d\theta}\cos^2\theta - r\sin^2\theta + r\sin\theta\cos\theta \Rightarrow$

$\dfrac{dr}{d\theta}\left(\sin^2\theta + \cos^2\theta\right) + r\left(\cos^2\theta + \sin^2\theta\right) = 0 \quad\Rightarrow\quad \dfrac{dr}{d\theta} = -r$. Solving this differential equation as a separable equation (as in Section 9.3), or using Theorem 9.4.2 with $k = -1$, we get $r = Ce^{-\theta}$. To determine C we use the fact that, at its starting position, $\theta = \frac{\pi}{4}$ and $r = \frac{1}{\sqrt{2}}a$, so $\frac{1}{\sqrt{2}}a = Ce^{-\pi/4} \quad\Rightarrow\quad C = \frac{1}{\sqrt{2}}ae^{\pi/4}$. Therefore, a polar equation of the bug's path is $r = \frac{1}{\sqrt{2}}ae^{\pi/4}e^{-\theta}$ or $r = \frac{1}{\sqrt{2}}ae^{(\pi/4)-\theta}$.

(b) The distance traveled by this bug is $L = \int_{\pi/4}^{\infty}\sqrt{r^2 + (dr/d\theta)^2}\,d\theta$, where $\dfrac{dr}{d\theta} = \dfrac{a}{\sqrt{2}}e^{\pi/4}(-e^{-\theta})$ and so $r^2 + (dr/d\theta)^2 = \frac{1}{2}a^2e^{\pi/2}e^{-2\theta} + \frac{1}{2}a^2e^{\pi/2}e^{-2\theta} = a^2e^{\pi/2}e^{-2\theta}$. Thus

$$L = \int_{\pi/4}^{\infty} ae^{\pi/4}e^{-\theta}\,d\theta = ae^{\pi/4}\lim_{t\to\infty}\int_{\pi/4}^{t} e^{-\theta}\,d\theta = ae^{\pi/4}\lim_{t\to\infty}\left[-e^{-\theta}\right]_{\pi/4}^{t}$$
$$= ae^{\pi/4}\lim_{t\to\infty}\left[e^{-\pi/4} - e^{-t}\right] = ae^{\pi/4}e^{-\pi/4} = a$$

5. Without loss of generality, assume the hyperbola has equation $\dfrac{x^2}{a^2} - \dfrac{y^2}{b^2} = 1$. Use implicit differentiation to get $\dfrac{2x}{a^2} - \dfrac{2y\,y'}{b^2} = 0$, so $y' = \dfrac{b^2x}{a^2y}$. The tangent line at the point (c, d) on the hyperbola has equation $y - d = \dfrac{b^2c}{a^2d}(x - c)$.

The tangent line intersects the asymptote $y = \dfrac{b}{a}x$ when $\dfrac{b}{a}x - d = \dfrac{b^2c}{a^2d}(x - c) \quad\Rightarrow\quad abdx - a^2d^2 = b^2cx - b^2c^2 \quad\Rightarrow$

$abdx - b^2cx = a^2d^2 - b^2c^2 \quad\Rightarrow\quad x = \dfrac{a^2d^2 - b^2c^2}{b(ad - bc)} = \dfrac{ad + bc}{b}$ and the y-value is $\dfrac{b}{a}\dfrac{ad + bc}{b} = \dfrac{ad + bc}{a}$.

Similarly, the tangent line intersects $y = -\dfrac{b}{a}x$ at $\left(\dfrac{bc - ad}{b}, \dfrac{ad - bc}{a}\right)$. The midpoint of these intersection points is

$\left(\dfrac{1}{2}\left(\dfrac{ad + bc}{b} + \dfrac{bc - ad}{b}\right), \dfrac{1}{2}\left(\dfrac{ad + bc}{a} + \dfrac{ad - bc}{a}\right)\right) = \left(\dfrac{1}{2}\dfrac{2bc}{b}, \dfrac{1}{2}\dfrac{2ad}{a}\right) = (c, d)$, the point of tangency.

Note: If $y = 0$, then at $(\pm a, 0)$, the tangent line is $x = \pm a$, and the points of intersection are clearly equidistant from the point of tangency.

6. (a) Since the smaller circle rolls without slipping around C, the amount of arc traversed on C ($2r\theta$ in the figure) must equal the amount of arc of the smaller circle that has been in contact with C. Since the smaller circle has radius r, it must have turned through an angle of $2r\theta/r = 2\theta$. In addition to turning through an angle 2θ, the little circle has rolled through an angle θ against C. Thus, P has turned through an angle of 3θ as shown in the figure. (If the little circle had turned through an angle of 2θ with its center pinned to the x-axis,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

then P would have turned only 2θ instead of 3θ. The movement of the little circle around C adds θ to the angle.) From the figure, we see that the center of the small circle has coordinates $(3r\cos\theta, 3r\sin\theta)$. Thus, P has coordinates (x, y), where $x = b\cos 3\theta + 3r\cos\theta$ and $y = b\sin 3\theta + 3r\sin\theta$.

(b)

$b = \frac{1}{5}r$ $\quad$ $b = \frac{2}{5}r$ $\quad$ $b = \frac{3}{5}r$ $\quad$ $b = \frac{4}{5}r$

(c) The diagram gives an alternate description of point P on the epitrochoid. Q moves around a circle of radius b, and P rotates one-third as fast with respect to Q at a distance of $3r$. Place an equilateral triangle with sides of length $3\sqrt{3}r$ so that its centroid is at Q and one vertex is at P. (The distance from the centroid to a vertex is $\frac{1}{\sqrt{3}}$ times the length of a side of the equilateral triangle.)

As θ increases by $\frac{2\pi}{3}$, the point Q travels once around the circle of radius b, returning to its original position. At the same time, P (and the rest of the triangle) rotate through an angle of $\frac{2\pi}{3}$ about Q, so P's position is occupied by another vertex. In this way, we see that the epitrochoid traced out by P is simultaneously traced out by the other two vertices as well. The whole equilateral triangle sits inside the epitrochoid (touching it only with its vertices) and each vertex traces out the curve once while the centroid moves around the circle three times.

(d) We view the epitrochoid as being traced out in the same way as in part (c), by a rotor for which the distance from its center to each vertex is $3r$, so it has radius $6r$. To show that the rotor fits inside the epitrochoid, it suffices to show that for any position of the tracing point P, there are no points on the opposite side of the rotor which are outside the epitrochoid. But the most likely case of intersection is when P is on the y-axis, so as long as the diameter of the rotor $\left(\text{which is } 3\sqrt{3}\,r\right)$ is less than the distance between the y-intercepts, the rotor will fit. The y-intercepts occur when $\theta = \frac{\pi}{2}$ or $\theta = \frac{3\pi}{2}$ $\Rightarrow$ $y = -b + 3r$ or $y = b - 3r$, so the distance between the intercepts is $(-b + 3r) - (b - 3r) = 6r - 2b$, and the rotor will fit if $3\sqrt{3}\,r \le 6r - 2b \quad \Leftrightarrow \quad 2b \le 6r - 3\sqrt{3}\,r \quad \Leftrightarrow \quad b \le \frac{3}{2}\left(2 - \sqrt{3}\right)r$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11 ☐ SEQUENCES, SERIES, AND POWER SERIES

11.1 Sequences

1. (a) A sequence is an ordered list of numbers. It can also be defined as a function whose domain is the set of positive integers.

(b) The terms a_n approach 8 as n becomes large. In fact, we can make a_n as close to 8 as we like by taking n sufficiently large.

(c) The terms a_n become large as n becomes large. In fact, we can make a_n as large as we like by taking n sufficiently large.

2. (a) From Definition 1, a convergent sequence is a sequence for which $\lim\limits_{n\to\infty} a_n$ exists. Examples: $\{1/n\}$, $\{1/2^n\}$

(b) A divergent sequence is a sequence for which $\lim\limits_{n\to\infty} a_n$ *does not* exist. Examples: $\{n\}$, $\{\sin n\}$

3. $a_n = n^3 - 1$, so the sequence is $\left\{1^3-1, 2^3-1, 3^3-1, 4^3-1, 5^3-1, \ldots\right\} = \{0, 7, 26, 63, 124, \ldots\}$.

4. $a_n = \dfrac{1}{3^n+1}$, so the sequence is $\left\{\dfrac{1}{3^1+1}, \dfrac{1}{3^2+1}, \dfrac{1}{3^3+1}, \dfrac{1}{3^4+1}, \dfrac{1}{3^5+1}, \ldots\right\} = \left\{\dfrac{1}{4}, \dfrac{1}{10}, \dfrac{1}{28}, \dfrac{1}{82}, \dfrac{1}{244}, \ldots\right\}$.

5. $\{2^n+n\}_{n=2}^{\infty}$, so the sequence is $\left\{2^2+2, 2^3+3, 2^4+4, 2^5+5, 2^6+6, \ldots\right\} = \{6, 11, 20, 37, 70, \ldots\}$.

6. $\left\{\dfrac{n^2-1}{n^2+1}\right\}_{n=3}^{\infty}$, so the sequence is

$$\left\{\frac{3^2-1}{3^2+1}, \frac{4^2-1}{4^2+1}, \frac{5^2-1}{5^2+1}, \frac{6^2-1}{6^2+1}, \frac{7^2-1}{7^2+1}, \ldots\right\} = \left\{\frac{8}{10}, \frac{15}{17}, \frac{24}{26}, \frac{35}{37}, \frac{48}{50}, \ldots\right\}.$$

7. $a_n = \dfrac{(-1)^{n-1}}{n^2}$, so the sequence is

$$\left\{\frac{(-1)^{1-1}}{1^2}, \frac{(-1)^{2-1}}{2^2}, \frac{(-1)^{3-1}}{3^2}, \frac{(-1)^{4-1}}{4^2}, \frac{(-1)^{5-1}}{5^2}, \ldots\right\} = \left\{1, -\frac{1}{4}, \frac{1}{9}, -\frac{1}{16}, \frac{1}{25}, \ldots\right\}.$$

8. $a_n = \dfrac{(-1)^n}{4^n}$, so the sequence is $\left\{\dfrac{(-1)^1}{4^1}, \dfrac{(-1)^2}{4^2}, \dfrac{(-1)^3}{4^3}, \dfrac{(-1)^4}{4^4}, \dfrac{(-1)^5}{4^5}, \ldots\right\} = \left\{-\dfrac{1}{4}, \dfrac{1}{16}, -\dfrac{1}{64}, \dfrac{1}{256}, -\dfrac{1}{1024}, \ldots\right\}$.

9. $a_n = \cos n\pi$, so the sequence is $\{\cos\pi, \cos 2\pi, \cos 3\pi, \cos 4\pi, \cos 5\pi, \ldots\} = \{-1, 1, -1, 1, -1, \ldots\}$.

10. $a_n = 1 + (-1)^n$, so the sequence is $\{1-1, 1+1, 1-1, 1+1, 1-1, \ldots\} = \{0, 2, 0, 2, 0, \ldots\}$.

11. $a_n = \dfrac{(-2)^n}{(n+1)!}$, so the sequence is

$$\left\{\frac{(-2)^1}{2!}, \frac{(-2)^2}{3!}, \frac{(-2)^3}{4!}, \frac{(-2)^4}{5!}, \frac{(-2)^5}{6!}, \ldots\right\} = \left\{-\frac{2}{2}, \frac{4}{6}, -\frac{8}{24}, \frac{16}{120}, -\frac{32}{720}, \ldots\right\} = \left\{-1, \frac{2}{3}, -\frac{1}{3}, \frac{2}{15}, -\frac{2}{45}, \ldots\right\}.$$

12. $a_n = \dfrac{2n+1}{n!+1}$, so the sequence is $\left\{\dfrac{2+1}{1+1}, \dfrac{4+1}{2+1}, \dfrac{6+1}{6+1}, \dfrac{8+1}{24+1}, \dfrac{10+1}{120+1}, \ldots\right\} = \left\{\dfrac{3}{2}, \dfrac{5}{3}, \dfrac{7}{7}, \dfrac{9}{25}, \dfrac{11}{121}, \ldots\right\}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $a_1 = 1, a_{n+1} = 2a_n + 1$. $a_2 = 2a_1 + 1 = 2 \cdot 1 + 1 = 3$. $a_3 = 2a_2 + 1 = 2 \cdot 3 + 1 = 7$. $a_4 = 2a_3 + 1 = 2 \cdot 7 + 1 = 15$.

$a_5 = 2a_4 + 1 = 2 \cdot 15 + 1 = 31$. The sequence is $\{1, 3, 7, 15, 31, \ldots\}$.

14. $a_1 = 6, a_{n+1} = \dfrac{a_n}{n}$. $a_2 = \dfrac{a_1}{1} = \dfrac{6}{1} = 6$. $a_3 = \dfrac{a_2}{2} = \dfrac{6}{2} = 3$. $a_4 = \dfrac{a_3}{3} = \dfrac{3}{3} = 1$. $a_5 = \dfrac{a_4}{4} = \dfrac{1}{4}$.

The sequence is $\left\{6, 6, 3, 1, \frac{1}{4}, \ldots\right\}$.

15. $a_1 = 2, a_{n+1} = \dfrac{a_n}{1 + a_n}$. $a_2 = \dfrac{a_1}{1 + a_1} = \dfrac{2}{1+2} = \dfrac{2}{3}$. $a_3 = \dfrac{a_2}{1 + a_2} = \dfrac{2/3}{1 + 2/3} = \dfrac{2}{5}$. $a_4 = \dfrac{a_3}{1 + a_3} = \dfrac{2/5}{1 + 2/5} = \dfrac{2}{7}$.

$a_5 = \dfrac{a_4}{1 + a_4} = \dfrac{2/7}{1 + 2/7} = \dfrac{2}{9}$. The sequence is $\left\{2, \frac{2}{3}, \frac{2}{5}, \frac{2}{7}, \frac{2}{9}, \ldots\right\}$.

16. $a_1 = 2, a_2 = 1, a_{n+1} = a_n - a_{n-1}$. Each term is defined in term of the two preceding terms.

$a_3 = a_2 - a_1 = 1 - 2 = -1$. $a_4 = a_3 - a_2 = -1 - 1 = -2$. $a_5 = a_4 - a_3 = -2 - (-1) = -1$.

The sequence is $\{2, 1, -1, -2, -1, \ldots\}$.

17. $\left\{\frac{1}{2}, \frac{1}{4}, \frac{1}{6}, \frac{1}{8}, \frac{1}{10}, \ldots\right\}$. The denominator is two times the number of the term, n, so $a_n = \dfrac{1}{2n}$.

18. $\left\{4, -1, \frac{1}{4}, -\frac{1}{16}, \frac{1}{64}, \ldots\right\}$. The first term is 4 and each term is $-\frac{1}{4}$ times the preceding one, so $a_n = 4\left(-\frac{1}{4}\right)^{n-1}$.

19. $\left\{-3, 2, -\frac{4}{3}, \frac{8}{9}, -\frac{16}{27}, \ldots\right\}$. The first term is -3 and each term is $-\frac{2}{3}$ times the preceding one, so $a_n = -3\left(-\frac{2}{3}\right)^{n-1}$.

20. $\{5, 8, 11, 14, 17, \ldots\}$. Each term is larger than the preceding term by 3, so $a_n = a_1 + d(n-1) = 5 + 3(n-1) = 3n + 2$.

21. $\left\{\frac{1}{2}, -\frac{4}{3}, \frac{9}{4}, -\frac{16}{5}, \frac{25}{6}, \ldots\right\}$. The numerator of the *n*th term is n^2 and its denominator is $n + 1$. Including the alternating signs, we get $a_n = (-1)^{n+1} \dfrac{n^2}{n+1}$.

22. $\{1, 0, -1, 0, 1, 0, -1, 0, \ldots\}$. Two possibilities are $a_n = \sin \dfrac{n\pi}{2}$ and $a_n = \cos \dfrac{(n-1)\pi}{2}$.

23.

n	$a_n = \dfrac{3n}{1+6n}$
1	0.4286
2	0.4615
3	0.4737
4	0.4800
5	0.4839
6	0.4865
7	0.4884
8	0.4898
9	0.4909
10	0.4918

It appears that $\lim\limits_{n\to\infty} a_n = 0.5$.

$$\lim_{n\to\infty} \frac{3n}{1+6n} = \lim_{n\to\infty} \frac{(3n)/n}{(1+6n)/n} = \lim_{n\to\infty} \frac{3}{1/n + 6} = \frac{3}{6} = \frac{1}{2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24.

n	$a_n = 2 + \dfrac{(-1)^n}{n}$
1	1.0000
2	2.5000
3	1.6667
4	2.2500
5	1.8000
6	2.1667
7	1.8571
8	2.1250
9	1.8889
10	2.1000

It appears that $\lim\limits_{n\to\infty} a_n = 2$.

$\lim\limits_{n\to\infty}\left(2 + \dfrac{(-1)^n}{n}\right) = \lim\limits_{n\to\infty} 2 + \lim\limits_{n\to\infty} \dfrac{(-1)^n}{n} = 2 + 0 = 2$ since $\lim\limits_{n\to\infty} \dfrac{1}{n} = 0$

and by Theorem 6, $\lim\limits_{n\to\infty} \dfrac{(-1)^n}{n} = 0$.

25.

n	$a_n = 1 + \left(-\frac{1}{2}\right)^n$
1	0.5000
2	1.2500
3	0.8750
4	1.0625
5	0.9688
6	1.0156
7	0.9922
8	1.0039
9	0.9980
10	1.0010

It appears that $\lim\limits_{n\to\infty} a_n = 1$.

$\lim\limits_{n\to\infty}\left(1 + \left(-\frac{1}{2}\right)^n\right) = \lim\limits_{n\to\infty} 1 + \lim\limits_{n\to\infty}\left(-\frac{1}{2}\right)^n = 1 + 0 = 1$ since

$\lim\limits_{n\to\infty}\left(-\frac{1}{2}\right)^n = 0$ by (9).

26.

n	$a_n = 1 + \dfrac{10^n}{9^n}$
1	2.1111
2	2.2346
3	2.3717
4	2.5242
5	2.6935
6	2.8817
7	3.0908
8	3.3231
9	3.5812
10	3.8680

It appears that the sequence does not have a limit.

$\lim\limits_{n\to\infty} \dfrac{10^n}{9^n} = \lim\limits_{n\to\infty}\left(\dfrac{10}{9}\right)^n$, which diverges by (9) since $\frac{10}{9} > 1$.

27. $a_n = \dfrac{5}{n+2} = \dfrac{5/n}{(n+2)/n} = \dfrac{5/n}{1+2/n}$, so $a_n \to \dfrac{0}{1+0} = 0$ as $n \to \infty$. Converges

28. $a_n = 5\sqrt{n+2}$, so $a_n \to \infty$ as $n \to \infty$ since $\lim\limits_{n\to\infty} \sqrt{n+2} = \infty$. Diverges

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. $a_n = \dfrac{4n^2 - 3n}{2n^2 + 1} = \dfrac{(4n^2 - 3n)/n^2}{(2n^2 + 1)/n^2} = \dfrac{4 - 3/n}{2 + 1/n^2}$, so $a_n \to \dfrac{4 - 0}{2 + 0} = 2$ as $n \to \infty$. Converges

30. $a_n = \dfrac{4n^2 - 3n}{2n + 1} = \dfrac{(4n^2 - 3n)/n}{(2n + 1)/n} = \dfrac{4n - 3}{2 + 1/n}$, so $a_n \to \infty$ as $n \to \infty$ since $\lim\limits_{n\to\infty}(4n - 3) = \infty$ and $\lim\limits_{n\to\infty}\left(2 + \dfrac{1}{n}\right) = 2$.

Diverges

31. $a_n = \dfrac{n^4}{n^3 - 2n} = \dfrac{n^4/n^3}{(n^3 - 2n)/n^3} = \dfrac{n}{1 - 2/n^2}$, so $a_n \to \infty$ as $n \to \infty$ since $\lim\limits_{n\to\infty} n = \infty$ and

$\lim\limits_{n\to\infty}\left(1 - \dfrac{2}{n^2}\right) = 1 - 0 = 1$. Diverges

32. $a_n = 2 + (0.86)^n \to 2 + 0 = 2$ as $n \to \infty$ since $\lim\limits_{n\to\infty}(0.86)^n = 0$ by (9) with $r = 0.86$. Converges

33. $a_n = 3^n 7^{-n} = \dfrac{3^n}{7^n} = \left(\dfrac{3}{7}\right)^n$, so $\lim\limits_{n\to\infty} a_n = 0$ by (9) with $r = \dfrac{3}{7}$. Converges

34. $a_n = \dfrac{3\sqrt{n}}{\sqrt{n} + 2} = \dfrac{3\sqrt{n}/\sqrt{n}}{(\sqrt{n} + 2)/\sqrt{n}} = \dfrac{3}{1 + 2/\sqrt{n}} \to \dfrac{3}{1 + 0} = 3$ as $n \to \infty$. Converges

35. Because the natural exponential function is continuous at 0, Theorem 7 enables us to write

$\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} e^{-1/\sqrt{n}} = e^{\lim\limits_{n\to\infty}(-1/\sqrt{n})} = e^0 = 1$. Converges

36. $a_n = \dfrac{4^n}{1 + 9^n} = \dfrac{4^n/9^n}{(1 + 9^n)/9^n} = \dfrac{(4/9)^n}{(1/9)^n + 1} \to \dfrac{0}{0 + 1} = 0$ as $n \to \infty$ since $\lim\limits_{n\to\infty}\left(\dfrac{4}{9}\right)^n = 0$ and

$\lim\limits_{n\to\infty}\left(\dfrac{1}{9}\right)^n = 0$ by (9). Converges

37. $a_n = \sqrt{\dfrac{1 + 4n^2}{1 + n^2}} = \sqrt{\dfrac{(1 + 4n^2)/n^2}{(1 + n^2)/n^2}} = \sqrt{\dfrac{(1/n^2) + 4}{(1/n^2) + 1}} \to \sqrt{4} = 2$ as $n \to \infty$ since $\lim\limits_{n\to\infty}(1/n^2) = 0$. Converges

38. $a_n = \cos\left(\dfrac{n\pi}{n + 1}\right) = \cos\left(\dfrac{n\pi/n}{(n + 1)/n}\right) = \cos\left(\dfrac{\pi}{1 + 1/n}\right)$, so $a_n \to \cos\pi = -1$ as $n \to \infty$ since $\lim\limits_{n\to\infty} 1/n = 0$.

Converges

39. $a_n = \dfrac{n^2}{\sqrt{n^3 + 4n}} = \dfrac{n^2/\sqrt{n^3}}{\sqrt{n^3 + 4n}/\sqrt{n^3}} = \dfrac{\sqrt{n}}{\sqrt{1 + 4/n^2}}$, so $a_n \to \infty$ as $n \to \infty$ since $\lim\limits_{n\to\infty}\sqrt{n} = \infty$ and

$\lim\limits_{n\to\infty}\sqrt{1 + 4/n^2} = 1$. Diverges

40. If $b_n = \dfrac{2n}{n + 2}$, then $\lim\limits_{n\to\infty} b_n = \lim\limits_{n\to\infty}\dfrac{(2n)/n}{(n + 2)/n} = \lim\limits_{n\to\infty}\dfrac{2}{1 + 2/n} = \dfrac{2}{1} = 2$. Since the natural exponential function is

continuous at 2, by Theorem 7, $\lim\limits_{n\to\infty} e^{2n/(n+2)} = e^{\lim_{n\to\infty} b_n} = e^2$. Converges

41. $\lim\limits_{n\to\infty}|a_n| = \lim\limits_{n\to\infty}\left|\dfrac{(-1)^n}{2\sqrt{n}}\right| = \dfrac{1}{2}\lim\limits_{n\to\infty}\dfrac{1}{n^{1/2}} = \dfrac{1}{2}(0) = 0$, so $\lim\limits_{n\to\infty} a_n = 0$ by (6). Converges

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. $\lim\limits_{n\to\infty} \dfrac{n}{n+\sqrt{n}} = \lim\limits_{n\to\infty} \dfrac{n/n}{(n+\sqrt{n}\,)/n} = \lim\limits_{n\to\infty} \dfrac{1}{1+1/\sqrt{n}} = \dfrac{1}{1+0} = 1.$ Thus, $a_n = \dfrac{(-1)^{n+1}n}{n+\sqrt{n}}$ has odd-numbered terms that approach 1 and even-numbered terms that approach -1 as $n \to \infty$, and hence, the sequence $\{a_n\}$ is divergent.

43. $a_n = \dfrac{(2n-1)!}{(2n+1)!} = \dfrac{(2n-1)!}{(2n+1)(2n)(2n-1)!} = \dfrac{1}{(2n+1)(2n)} \to 0$ as $n \to \infty$. Converges

44. $a_n = \dfrac{\ln n}{\ln(2n)} = \dfrac{\ln n}{\ln 2 + \ln n} = \dfrac{(\ln n)/\ln n}{(\ln 2 + \ln n)/\ln n} = \dfrac{1}{\frac{\ln 2}{\ln n} + 1} \to \dfrac{1}{0+1} = 1$ as $n \to \infty$. Converges

45. $a_n = \sin n$. This sequence diverges since the terms don't approach any particular real number as $n \to \infty$. The terms take on values between -1 and 1. Diverges

46. $a_n = \dfrac{\tan^{-1} n}{n}$. $\lim\limits_{n\to\infty} \tan^{-1} n = \lim\limits_{x\to\infty} \tan^{-1} x = \dfrac{\pi}{2}$ by (4), so $\lim\limits_{n\to\infty} a_n = 0$. Converges

47. $a_n = n^2 e^{-n} = \dfrac{n^2}{e^n}$. Since $\lim\limits_{x\to\infty} \dfrac{x^2}{e^x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{2x}{e^x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{2}{e^x} = 0$, it follows from Theorem 4 that $\lim\limits_{n\to\infty} a_n = 0$. Converges

48. $a_n = \ln(n+1) - \ln n = \ln\left(\dfrac{n+1}{n}\right) = \ln\left(1 + \dfrac{1}{n}\right) \to \ln(1) = 0$ as $n \to \infty$ because ln is continuous. Converges

49. $0 \le \dfrac{\cos^2 n}{2^n} \le \dfrac{1}{2^n}$ [since $0 \le \cos^2 n \le 1$], so since $\lim\limits_{n\to\infty} \dfrac{1}{2^n} = 0$, $\left\{\dfrac{\cos^2 n}{2^n}\right\}$ converges to 0 by the Squeeze Theorem.

50. $a_n = \sqrt[n]{2^{1+3n}} = (2^{1+3n})^{1/n} = (2^1 2^{3n})^{1/n} = 2^{1/n} 2^3 = 8 \cdot 2^{1/n}$, so

$\lim\limits_{n\to\infty} a_n = 8 \lim\limits_{n\to\infty} 2^{1/n} = 8 \cdot 2^{\lim_{n\to\infty}(1/n)} = 8 \cdot 2^0 = 8$ by Theorem 7, since the function $f(x) = 2^x$ is continuous at 0.

Converges

51. $a_n = n\sin(1/n) = \dfrac{\sin(1/n)}{1/n}$. Since $\lim\limits_{x\to\infty} \dfrac{\sin(1/x)}{1/x} = \lim\limits_{t\to 0^+} \dfrac{\sin t}{t}$ [where $t = 1/x$] $= 1$, it follows from Theorem 4 that $\{a_n\}$ converges to 1.

52. $a_n = 2^{-n}\cos n\pi$. $0 \le \left|\dfrac{\cos n\pi}{2^n}\right| \le \dfrac{1}{2^n} = \left(\dfrac{1}{2}\right)^n$, so $\lim\limits_{n\to\infty} |a_n| = 0$ by (9), and $\lim\limits_{n\to\infty} a_n = 0$ by (6). Converges

53. $y = \left(1 + \dfrac{2}{x}\right)^x \quad \Rightarrow \quad \ln y = x \ln\left(1 + \dfrac{2}{x}\right)$, so

$$\lim_{x\to\infty} \ln y = \lim_{x\to\infty} \frac{\ln(1+2/x)}{1/x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{\left(\dfrac{1}{1+2/x}\right)\left(-\dfrac{2}{x^2}\right)}{-1/x^2} = \lim_{x\to\infty} \frac{2}{1+2/x} = 2 \quad \Rightarrow$$

$\lim\limits_{x\to\infty} \left(1 + \dfrac{2}{x}\right)^x = \lim\limits_{x\to\infty} e^{\ln y} = e^2$, so by Theorem 4, $\lim\limits_{n\to\infty} \left(1 + \dfrac{2}{n}\right)^n = e^2$. Converges

54. $y = x^{1/x} \quad \Rightarrow \quad \ln y = \dfrac{1}{x}\ln x$, so $\lim\limits_{x\to\infty} \ln y = \lim\limits_{x\to\infty} \dfrac{\ln x}{x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1/x}{1} = \lim\limits_{x\to\infty} \dfrac{1}{x} = 0 \quad \Rightarrow$

$\lim\limits_{x\to\infty} x^{1/x} = \lim\limits_{x\to\infty} e^{\ln y} = e^0 = 1$, so by Theorem 4, $\lim\limits_{n\to\infty} n^{1/n} = 1$. Converges

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

55. $a_n = \ln(2n^2+1) - \ln(n^2+1) = \ln\left(\dfrac{2n^2+1}{n^2+1}\right) = \ln\left(\dfrac{2+1/n^2}{1+1/n^2}\right) \to \ln 2$ as $n \to \infty$. Converges

56. $\displaystyle\lim_{x\to\infty} \frac{(\ln x)^2}{x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{2(\ln x)(1/x)}{1} = 2\lim_{x\to\infty} \frac{\ln x}{x} \overset{\text{H}}{=} 2\lim_{x\to\infty} \frac{1/x}{1} = 0$, so by Theorem 4, $\displaystyle\lim_{n\to\infty} \frac{(\ln n)^2}{n} = 0$. Converges

57. $a_n = \arctan(\ln n)$. Let $f(x) = \arctan(\ln x)$. Then $\displaystyle\lim_{x\to\infty} f(x) = \tfrac{\pi}{2}$ since $\ln x \to \infty$ as $x \to \infty$ and $\arctan$ is continuous.

Thus, $\displaystyle\lim_{n\to\infty} a_n = \lim_{n\to\infty} f(n) = \tfrac{\pi}{2}$. Converges

58. $a_n = n - \sqrt{n+1}\,\sqrt{n+3} = n - \sqrt{n^2+4n+3} = \dfrac{n-\sqrt{n^2+4n+3}}{1} \cdot \dfrac{n+\sqrt{n^2+4n+3}}{n+\sqrt{n^2+4n+3}}$

$$= \frac{n^2-(n^2+4n+3)}{n+\sqrt{n^2+4n+3}} = \frac{-4n-3}{n+\sqrt{n^2+4n+3}} = \frac{(-4n-3)/n}{\left(n+\sqrt{n^2+4n+3}\right)/n} = \frac{-4-3/n}{1+\sqrt{1+4/n+3/n^2}},$$

so $\displaystyle\lim_{n\to\infty} a_n = \frac{-4-0}{1+\sqrt{1+0+0}} = \frac{-4}{2} = -2$. Converges

59. $\{0,1,0,0,1,0,0,0,1,\ldots\}$ diverges since the sequence takes on only two values, 0 and 1, and never stays arbitrarily close to either value (or any other value) for n sufficiently large.

60. $\left\{\frac{1}{1}, \frac{1}{3}, \frac{1}{2}, \frac{1}{4}, \frac{1}{3}, \frac{1}{5}, \frac{1}{4}, \frac{1}{6}, \ldots\right\}$. $a_{2n-1} = \dfrac{1}{n}$ and $a_{2n} = \dfrac{1}{n+2}$ for all positive integers n. $\displaystyle\lim_{n\to\infty} a_n = 0$ since

$\displaystyle\lim_{n\to\infty} a_{2n-1} = \lim_{n\to\infty} \frac{1}{n} = 0$ and $\displaystyle\lim_{n\to\infty} a_{2n} = \lim_{n\to\infty} \frac{1}{n+2} = 0$. For n sufficiently large, a_n can be made as close to 0 as we like. Converges

61. $a_n = \dfrac{n!}{2^n} = \dfrac{1}{2}\cdot\dfrac{2}{2}\cdot\dfrac{3}{2}\cdot\cdots\cdot\dfrac{(n-1)}{2}\cdot\dfrac{n}{2} \geq \dfrac{1}{2}\cdot\dfrac{n}{2}$ [for $n > 1$] $= \dfrac{n}{4} \to \infty$ as $n \to \infty$, so $\{a_n\}$ diverges.

62. $0 < |a_n| = \dfrac{3^n}{n!} = \dfrac{3}{1}\cdot\dfrac{3}{2}\cdot\dfrac{3}{3}\cdot\cdots\cdot\dfrac{3}{(n-1)}\cdot\dfrac{3}{n} \leq \dfrac{3}{1}\cdot\dfrac{3}{2}\cdot\dfrac{3}{n}$ [for $n > 2$] $= \dfrac{27}{2n} \to 0$ as $n \to \infty$, so by the Squeeze Theorem and Theorem 6, $\{(-3)^n/n!\}$ converges to 0.

63.

From the graph, it appears that the sequence $\{a_n\} = \left\{(-1)^n \dfrac{n}{n+1}\right\}$ is divergent, since it oscillates between 1 and -1 (approximately). To prove this, suppose that $\{a_n\}$ converges to L. If $b_n = \dfrac{n}{n+1}$, then $\{b_n\}$ converges to 1, and $\displaystyle\lim_{n\to\infty} \frac{a_n}{b_n} = \frac{L}{1} = L$. But $\dfrac{a_n}{b_n} = (-1)^n$, so $\displaystyle\lim_{n\to\infty} \frac{a_n}{b_n}$ does not exist. This contradiction shows that $\{a_n\}$ diverges.

64.

From the graph, it appears that the sequence converges to 0.

$|a_n| = \left|\dfrac{\sin n}{n}\right| = \dfrac{|\sin n|}{|n|} \leq \dfrac{1}{n}$, so $\displaystyle\lim_{n\to\infty} |a_n| = 0$. By (6), it follows that $\displaystyle\lim_{n\to\infty} a_n = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

65.

From the graph, it appears that the sequence converges to a number between 0.7 and 0.8.

$a_n = \arctan\left(\dfrac{n^2}{n^2+4}\right) = \arctan\left(\dfrac{n^2/n^2}{(n^2+4)/n^2}\right) = \arctan\left(\dfrac{1}{1+4/n^2}\right) \to$ $\arctan 1 = \dfrac{\pi}{4}$ $[\approx 0.785]$ as $n \to \infty$.

66.

From the graph, it appears that the sequence converges to 5.

$$5 = \sqrt[n]{5^n} \le \sqrt[n]{3^n+5^n} \le \sqrt[n]{5^n+5^n} = \sqrt[n]{2}\,\sqrt[n]{5^n}$$

$$= \sqrt[n]{2}\cdot 5 \to 5 \text{ as } n \to \infty \quad \left[\lim_{n\to\infty} 2^{1/n} = 2^0 = 1\right]$$

Hence, $a_n \to 5$ by the Squeeze Theorem.

Alternate solution: Let $y = (3^x+5^x)^{1/x}$. Then

$$\lim_{x\to\infty} \ln y = \lim_{x\to\infty} \frac{\ln(3^x+5^x)}{x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{3^x\ln 3 + 5^x \ln 5}{3^x+5^x} = \lim_{x\to\infty} \frac{\left(\frac{3}{5}\right)^x \ln 3 + \ln 5}{\left(\frac{3}{5}\right)^x + 1} = \ln 5,$$

so $\lim\limits_{x\to\infty} y = e^{\ln 5} = 5$, and so $\left\{\sqrt[n]{3^n+5^n}\right\}$ converges to 5.

67.

From the graph, it appears that the sequence $\{a_n\} = \left\{\dfrac{n^2\cos n}{1+n^2}\right\}$ is divergent, since it oscillates between 1 and -1 (approximately). To prove this, suppose that $\{a_n\}$ converges to L. If $b_n = \dfrac{n^2}{1+n^2}$, then $\{b_n\}$ converges to 1, and $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \dfrac{L}{1} = L$. But $\dfrac{a_n}{b_n} = \cos n$, so $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n}$ does not exist. This contradiction shows that $\{a_n\}$ diverges.

68.

From the graphs, it seems that the sequence diverges. $a_n = \dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{n!}$. We first prove by induction that $a_n \ge \left(\dfrac{3}{2}\right)^{n-1}$ for all n. This is clearly true for $n=1$, so let $P(n)$ be the statement that the above is true for n. We must show it is then true for $n+1$. $a_{n+1} = a_n \cdot \dfrac{2n+1}{n+1} \ge \left(\dfrac{3}{2}\right)^{n-1} \cdot \dfrac{2n+1}{n+1}$ (induction hypothesis). But $\dfrac{2n+1}{n+1} \ge \dfrac{3}{2}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

[since $2(2n+1) \geq 3(n+1) \quad \Leftrightarrow \quad 4n+2 \geq 3n+3 \quad \Leftrightarrow \quad n \geq 1$], and so we get that $a_{n+1} \geq \left(\frac{3}{2}\right)^{n-1} \cdot \frac{3}{2} = \left(\frac{3}{2}\right)^n$ which is $P(n+1)$. Thus, we have proved our first assertion, so since $\left\{\left(\frac{3}{2}\right)^{n-1}\right\}$ diverges [by (9)], so does the given sequence $\{a_n\}$.

69.

From the graph, it appears that the sequence approaches 0.

$$0 < a_n = \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{(2n)^n} = \frac{1}{2n} \cdot \frac{3}{2n} \cdot \frac{5}{2n} \cdot \cdots \cdot \frac{2n-1}{2n}$$

$$\leq \frac{1}{2n} \cdot (1) \cdot (1) \cdot \cdots \cdot (1) = \frac{1}{2n} \to 0 \text{ as } n \to \infty$$

So by the Squeeze Theorem, $\left\{\dfrac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{(2n)^n}\right\}$ converges to 0.

70. (a) $a_1 = 1$, $a_{n+1} = 4 - a_n$ for $n \geq 1$. $a_1 = 1$, $a_2 = 4 - a_1 = 4 - 1 = 3$, $a_3 = 4 - a_2 = 4 - 3 = 1$, $a_4 = 4 - a_3 = 4 - 1 = 3$, $a_5 = 4 - a_4 = 4 - 3 = 1$. Since the terms of the sequence alternate between 1 and 3, the sequence is divergent.

(b) $a_1 = 2$, $a_2 = 4 - a_1 = 4 - 2 = 2$, $a_3 = 4 - a_2 = 4 - 2 = 2$. Since all of the terms are 2, $\lim\limits_{n\to\infty} a_n = 2$ and hence, the sequence is convergent.

71. (a) $a_n = 1000(1.06)^n \quad \Rightarrow \quad a_1 = 1060$, $a_2 = 1123.60$, $a_3 = 1191.02$, $a_4 = 1262.48$, and $a_5 = 1338.23$.

(b) $\lim\limits_{n\to\infty} a_n = 1000 \lim\limits_{n\to\infty} (1.06)^n$, so the sequence diverges by (9) with $r = 1.06 > 1$.

72. (a) Substitute 1 to 6 for n in $I_n = 100\left(\dfrac{1.0025^n - 1}{0.0025} - n\right)$ to get $I_1 = \$0$, $I_2 = \$0.25$, $I_3 = \$0.75$, $I_4 = \$1.50$, $I_5 = \$2.51$, and $I_6 = \$3.76$.

(b) For two years, use $2 \cdot 12 = 24$ for n to get $\$70.28$.

73. (a) We are given that the initial population is 5000, so $P_0 = 5000$. The number of catfish increases by 8% per month and is decreased by 300 per month, so $P_1 = P_0 + 8\%P_0 - 300 = 1.08P_0 - 300$, $P_2 = 1.08P_1 - 300$, and so on. Thus, $P_n = 1.08P_{n-1} - 300$.

(b) Using the recursive formula with $P_0 = 5000$, we get $P_1 = 5100$, $P_2 = 5208$, $P_3 = 5325$ (rounding any portion of a catfish), $P_4 = 5451$, $P_5 = 5587$, and $P_6 = 5734$, which is the number of catfish in the pond after six months.

74. $a_{n+1} = \begin{cases} \frac{1}{2}a_n & \text{if } a_n \text{ is an even number} \\ 3a_n + 1 & \text{if } a_n \text{ is an odd number} \end{cases}$ When $a_1 = 11$, the first 40 terms are 11, 34, 17, 52, 26, 13, 40, 20, 10, 5, 16, 8, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4. When $a_1 = 25$, the first 40 terms are 25, 76, 38, 19, 58, 29, 88, 44, 22, 11, 34, 17, 52, 26, 13, 40, 20, 10, 5, 16, 8, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4, 2, 1, 4. The famous Collatz conjecture is that this sequence always reaches 1, regardless of the starting point a_1.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

75. If $|r| \geq 1$, then $\{r^n\}$ diverges by (9), so $\{nr^n\}$ diverges also, since $|nr^n| = n\,|r^n| \geq |r^n|$. If $|r| < 1$ then

$\lim\limits_{x\to\infty} xr^x = \lim\limits_{x\to\infty} \dfrac{x}{r^{-x}} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1}{(-\ln r)\,r^{-x}} = \lim\limits_{x\to\infty} \dfrac{r^x}{-\ln r} = 0$, so $\lim\limits_{n\to\infty} nr^n = 0$, and hence $\{nr^n\}$ converges whenever $|r| < 1$.

76. (a) Let $\lim\limits_{n\to\infty} a_n = L$. By Definition 2, this means that for every $\varepsilon > 0$ there is an integer N such that $|a_n - L| < \varepsilon$ whenever $n > N$. Thus, $|a_{n+1} - L| < \varepsilon$ whenever $n + 1 > N \quad \Leftrightarrow \quad n > N - 1$. It follows that $\lim\limits_{n\to\infty} a_{n+1} = L$ and so $\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} a_{n+1}$.

(b) If $L = \lim\limits_{n\to\infty} a_n$ then $\lim\limits_{n\to\infty} a_{n+1} = L$ also, so L must satisfy $L = 1/(1+L) \quad \Rightarrow \quad L^2 + L - 1 = 0 \quad \Rightarrow \quad L = \frac{-1+\sqrt{5}}{2}$ (since L has to be nonnegative if it exists).

77. Since $\{a_n\}$ is a decreasing sequence, $a_n > a_{n+1}$ for all $n \geq 1$. Because all of its terms lie between 5 and 8, $\{a_n\}$ is a bounded sequence. By the Monotonic Sequence Theorem, $\{a_n\}$ is convergent; that is, $\{a_n\}$ has a limit L. L must be less than 8 since $\{a_n\}$ is decreasing, so $5 \leq L < 8$.

78. Since $\{a_n\} = \{\cos n\} \approx \{0.54, -0.42, -0.99, -0.65, 0.28, \ldots\}$, the sequence is not monotonic. The sequence is bounded since $-1 \leq \cos n \leq 1$ for all n.

79. $a_n = \dfrac{1}{2n+3}$ is decreasing since $a_{n+1} = \dfrac{1}{2(n+1)+3} = \dfrac{1}{2n+5} < \dfrac{1}{2n+3} = a_n$ for each $n \geq 1$. The sequence is bounded since $0 < a_n \leq \frac{1}{5}$ for all $n \geq 1$. Note that $a_1 = \frac{1}{5}$.

80. $a_n > a_{n+1} \quad \Leftrightarrow \quad \dfrac{1-n}{2+n} > \dfrac{1-(n+1)}{2+(n+1)} \quad \Leftrightarrow \quad \dfrac{1-n}{2+n} > \dfrac{-n}{n+3} \quad \Leftrightarrow \quad -n^2 - 2n + 3 > -n^2 - 2n \quad \Leftrightarrow \quad 3 > 0$, which is true for all $n \geq 1$, so $\{a_n\}$ is decreasing. Since $a_1 = 0$ and $\lim\limits_{n\to\infty} \dfrac{1-n}{2+n} = \lim\limits_{n\to\infty} \dfrac{1/n - 1}{2/n + 1} = -1$, the sequence is bounded $(-1 < a_n \leq 0)$.

81. The terms of $a_n = n(-1)^n$ alternate in sign, so the sequence is not monotonic. The first five terms are -1, 2, -3, 4, and -5. Since $\lim\limits_{n\to\infty} |a_n| = \lim\limits_{n\to\infty} n = \infty$, the sequence is not bounded.

82. Since $\{a_n\} = \left\{2 + \dfrac{(-1)^n}{n}\right\} = \left\{1, 2\tfrac{1}{2}, 1\tfrac{2}{3}, \ldots\right\}$, the sequence is not monotonic. The sequence is bounded since $1 \leq a_n \leq \frac{5}{2}$ for all n.

83. $a_n = 3 - 2ne^{-n}$. Let $f(x) = 3 - 2xe^{-x}$. Then $f'(x) = 0 - 2[x(-e^{-x}) + e^{-x}] = 2e^{-x}(x-1)$, which is positive for $x > 1$, so f is increasing on $(1, \infty)$. It follows that the sequence $\{a_n\} = \{f(n)\}$ is increasing. The sequence is bounded below by $a_1 = 3 - 2e^{-1} \approx 2.26$ and above by 3, so the sequence is bounded.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

84. $a_n = n^3 - 3n + 3$. Let $f(x) = x^3 - 3x + 3$. Then $f'(x) = 3x^2 - 3 = 3(x^2 - 1)$, which is positive for $x > 1$, so f is increasing on $(1, \infty)$. It follows that the sequence $\{a_n\} = \{f(n)\}$ is increasing. The sequence is bounded below by $a_1 = 1$, but is not bounded above, so it is not bounded.

85. For $\left\{\sqrt{2}, \sqrt{2\sqrt{2}}, \sqrt{2\sqrt{2\sqrt{2}}}, \ldots\right\}$, $a_1 = 2^{1/2}$, $a_2 = 2^{3/4}$, $a_3 = 2^{7/8}, \ldots$, so $a_n = 2^{(2^n-1)/2^n} = 2^{1-(1/2^n)}$.

$\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} 2^{1-(1/2^n)} = 2^1 = 2$.

Alternate solution: Let $L = \lim\limits_{n\to\infty} a_n$. (We could show the limit exists by showing that $\{a_n\}$ is bounded and increasing.)

Then L must satisfy $L = \sqrt{2 \cdot L} \quad \Rightarrow \quad L^2 = 2L \quad \Rightarrow \quad L(L-2) = 0$. $L \neq 0$ since the sequence increases, so $L = 2$.

86. (a) Let P_n be the statement that $a_{n+1} \geq a_n$ and $a_n \leq 3$. P_1 is obviously true. We will assume that P_n is true and then show that as a consequence P_{n+1} must also be true. $a_{n+2} \geq a_{n+1} \quad \Leftrightarrow \quad \sqrt{2 + a_{n+1}} \geq \sqrt{2 + a_n} \quad \Leftrightarrow$ $2 + a_{n+1} \geq 2 + a_n \quad \Leftrightarrow \quad a_{n+1} \geq a_n$, which is the induction hypothesis. $a_{n+1} \leq 3 \quad \Leftrightarrow \quad \sqrt{2 + a_n} \leq 3 \quad \Leftrightarrow$ $2 + a_n \leq 9 \quad \Leftrightarrow \quad a_n \leq 7$, which is certainly true because we are assuming that $a_n \leq 3$. So P_n is true for all n, and so $a_1 \leq a_n \leq 3$ (showing that the sequence is bounded), and hence by the Monotonic Sequence Theorem, $\lim\limits_{n\to\infty} a_n$ exists.

(b) If $L = \lim\limits_{n\to\infty} a_n$, then $\lim\limits_{n\to\infty} a_{n+1} = L$ also, so $L = \sqrt{2 + L} \quad \Rightarrow \quad L^2 = 2 + L \quad \Leftrightarrow \quad L^2 - L - 2 = 0 \quad \Leftrightarrow$ $(L+1)(L-2) = 0 \quad \Leftrightarrow \quad L = 2$ [since L can't be negative].

87. $a_1 = 1$, $a_{n+1} = 3 - \dfrac{1}{a_n}$. We show by induction that $\{a_n\}$ is increasing and bounded above by 3. Let P_n be the proposition that $a_{n+1} > a_n$ and $0 < a_n < 3$. Clearly P_1 is true. Assume that P_n is true. Then $a_{n+1} > a_n \quad \Rightarrow \quad \dfrac{1}{a_{n+1}} < \dfrac{1}{a_n} \quad \Rightarrow$ $-\dfrac{1}{a_{n+1}} > -\dfrac{1}{a_n}$. Now $a_{n+2} = 3 - \dfrac{1}{a_{n+1}} > 3 - \dfrac{1}{a_n} = a_{n+1} \quad \Leftrightarrow \quad P_{n+1}$. This proves that $\{a_n\}$ is increasing and bounded above by 3, so $1 = a_1 < a_n < 3$, that is, $\{a_n\}$ is bounded, and hence convergent by the Monotonic Sequence Theorem.

If $L = \lim\limits_{n\to\infty} a_n$, then $\lim\limits_{n\to\infty} a_{n+1} = L$ also, so L must satisfy $L = 3 - 1/L \quad \Rightarrow \quad L^2 - 3L + 1 = 0 \quad \Rightarrow \quad L = \frac{3 \pm \sqrt{5}}{2}$.

But $L > 1$, so $L = \frac{3 + \sqrt{5}}{2}$.

88. $a_1 = 2$, $a_{n+1} = \dfrac{1}{3 - a_n}$. We use induction. Let P_n be the statement that $0 < a_{n+1} \leq a_n \leq 2$. Clearly P_1 is true, since $a_2 = 1/(3-2) = 1$. Now assume that P_n is true. Then $a_{n+1} \leq a_n \quad \Rightarrow \quad -a_{n+1} \geq -a_n \quad \Rightarrow \quad 3 - a_{n+1} \geq 3 - a_n \quad \Rightarrow$ $a_{n+2} = \dfrac{1}{3 - a_{n+1}} \leq \dfrac{1}{3 - a_n} = a_{n+1}$. Also $a_{n+2} > 0$ [since $3 - a_{n+1}$ is positive] and $a_{n+1} \leq 2$ by the induction hypothesis, so P_{n+1} is true. To find the limit, we use the fact that $\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} a_{n+1} \quad \Rightarrow \quad L = \frac{1}{3-L} \quad \Rightarrow$ $L^2 - 3L + 1 = 0 \quad \Rightarrow \quad L = \frac{3 \pm \sqrt{5}}{2}$. But $L \leq 2$, so we must have $L = \frac{3 - \sqrt{5}}{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

89. (a) Let a_n be the number of rabbit pairs in the *n*th month. Clearly $a_1 = 1 = a_2$. In the *n*th month, each pair that is 2 or more months old (that is, a_{n-2} pairs) will produce a new pair to add to the a_{n-1} pairs already present. Thus, $a_n = a_{n-1} + a_{n-2}$, so that $\{a_n\} = \{f_n\}$, the Fibonacci sequence.

(b) $a_n = \dfrac{f_{n+1}}{f_n} \quad \Rightarrow \quad a_{n-1} = \dfrac{f_n}{f_{n-1}} = \dfrac{f_{n-1} + f_{n-2}}{f_{n-1}} = 1 + \dfrac{f_{n-2}}{f_{n-1}} = 1 + \dfrac{1}{f_{n-1}/f_{n-2}} = 1 + \dfrac{1}{a_{n-2}}$. If $L = \lim\limits_{n\to\infty} a_n$, then $L = \lim\limits_{n\to\infty} a_{n-1}$ and $L = \lim\limits_{n\to\infty} a_{n-2}$, so L must satisfy $L = 1 + \dfrac{1}{L} \quad \Rightarrow \quad L^2 - L - 1 = 0 \quad \Rightarrow \quad L = \frac{1+\sqrt{5}}{2}$ [since L must be positive].

90. (a) If f is continuous, then $f(L) = f\left(\lim\limits_{n\to\infty} a_n\right) = \lim\limits_{n\to\infty} f(a_n) = \lim\limits_{n\to\infty} a_{n+1} = \lim\limits_{n\to\infty} a_n = L$ by Exercise 76(a).

(b) By repeatedly pressing the cosine key on the calculator (that is, taking cosine of the previous answer) until the displayed value stabilizes, we see that $L \approx 0.73909$.

91. (a)

From the graph, it appears that the sequence $\left\{\dfrac{n^5}{n!}\right\}$ converges to 0, that is, $\lim\limits_{n\to\infty} \dfrac{n^5}{n!} = 0$.

(b)

From the first graph, it seems that the smallest possible value of N corresponding to $\varepsilon = 0.1$ is 9, since $n^5/n! < 0.1$ whenever $n \geq 10$, but $9^5/9! > 0.1$. From the second graph, it seems that for $\varepsilon = 0.001$, the smallest possible value for N is 11 since $n^5/n! < 0.001$ whenever $n \geq 12$.

92. Let $\varepsilon > 0$ and let N be any positive integer larger than $\ln(\varepsilon)/\ln|r|$. If $n > N$, then $n > \ln(\varepsilon)/\ln|r| \quad \Rightarrow \quad n\ln|r| < \ln\varepsilon$ [since $|r| < 1 \Rightarrow \ln|r| < 0$] $\quad \Rightarrow \quad \ln(|r|^n) < \ln\varepsilon \quad \Rightarrow \quad |r|^n < \varepsilon \quad \Rightarrow \quad |r^n - 0| < \varepsilon$, and so by Definition 2, $\lim\limits_{n\to\infty} r^n = 0$.

93. **Theorem 6:** If $\lim\limits_{n\to\infty} |a_n| = 0$ then $\lim\limits_{n\to\infty} -|a_n| = 0$, and since $-|a_n| \leq a_n \leq |a_n|$, we have that $\lim\limits_{n\to\infty} a_n = 0$ by the Squeeze Theorem.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

94. Let $L = \lim\limits_{n\to\infty} a_n$ and $f(x) = x^p$, $p > 0$ and $a_n > 0$. Since f is a continuous function,

$$\lim_{n\to\infty} a_n^p = \lim_{n\to\infty} f(a_n) = f\left(\lim_{n\to\infty} a_n\right) \text{ [Theorem 7] } = f(L) = \left[\lim_{n\to\infty} a_n\right]^p.$$

95. **To Prove:** If $\lim\limits_{n\to\infty} a_n = 0$ and $\{b_n\}$ is bounded, then $\lim\limits_{n\to\infty} (a_n b_n) = 0$.

Proof: Since $\{b_n\}$ is bounded, there is a positive number M such that $|b_n| \le M$ and hence, $|a_n|\,|b_n| \le |a_n|\,M$ for all $n \ge 1$. Let $\varepsilon > 0$ be given. Since $\lim\limits_{n\to\infty} a_n = 0$, there is an integer N such that $|a_n - 0| < \dfrac{\varepsilon}{M}$ if $n > N$. Then $|a_n b_n - 0| = |a_n b_n| = |a_n|\,|b_n| \le |a_n|\,M = |a_n - 0|\,M < \dfrac{\varepsilon}{M} \cdot M = \varepsilon$ for all $n > N$. Since ε was arbitrary, $\lim\limits_{n\to\infty} (a_n b_n) = 0$.

96. (a)
$$\begin{aligned}\frac{b^{n+1} - a^{n+1}}{b - a} &= b^n + b^{n-1}a + b^{n-2}a^2 + b^{n-3}a^3 + \cdots + ba^{n-1} + a^n \\ &< b^n + b^{n-1}b + b^{n-2}b^2 + b^{n-3}b^3 + \cdots + bb^{n-1} + b^n = (n+1)b^n\end{aligned}$$

(b) Since $b - a > 0$, we have $b^{n+1} - a^{n+1} < (n+1)b^n(b-a) \;\Rightarrow\; b^{n+1} - (n+1)b^n(b-a) < a^{n+1} \;\Rightarrow$ $b^n[(n+1)a - nb] < a^{n+1}$.

(c) Substituting in part (b), $(n+1)a - nb = (n+1)\left(1 + \dfrac{1}{n+1}\right) - n\left(1 + \dfrac{1}{n}\right) = 1$, and so

$$b^n = \left(1 + \frac{1}{n}\right)^n < a^{n+1} = \left(1 + \frac{1}{n+1}\right)^{n+1}.$$

(d) Substituting in part (b), $(n+1)a - nb = (n+1)\cdot 1 - n\left(1 + \dfrac{1}{2n}\right) = \dfrac{1}{2}$, and so $\left(1 + \dfrac{1}{2n}\right)^n \left(\dfrac{1}{2}\right) < 1 \;\Rightarrow$

$$\left(1 + \frac{1}{2n}\right)^n < 2 \;\Rightarrow\; \left(1 + \frac{1}{2n}\right)^{2n} < 4.$$

(e) $a_n < a_{2n}$ since $\{a_n\}$ is increasing, so $a_n < a_{2n} < 4$.

(f) Since $\{a_n\}$ is increasing and bounded above by 4, $a_1 \le a_n \le 4$, and so $\{a_n\}$ is bounded and monotonic, and hence has a limit by the Monotonic Sequence Theorem.

97. (a) First we show that $a > a_1 > b_1 > b$.

$a_1 - b_1 = \frac{a+b}{2} - \sqrt{ab} = \frac{1}{2}\left(a - 2\sqrt{ab} + b\right) = \frac{1}{2}\left(\sqrt{a} - \sqrt{b}\right)^2 > 0$ [since $a > b$] $\;\Rightarrow\; a_1 > b_1$. Also

$a - a_1 = a - \frac{1}{2}(a+b) = \frac{1}{2}(a-b) > 0$ and $b - b_1 = b - \sqrt{ab} = \sqrt{b}\left(\sqrt{b} - \sqrt{a}\right) < 0$, so $a > a_1 > b_1 > b$. In the same way we can show that $a_1 > a_2 > b_2 > b_1$ and so the given assertion is true for $n = 1$. Suppose it is true for $n = k$, that is, $a_k > a_{k+1} > b_{k+1} > b_k$. Then

$$a_{k+2} - b_{k+2} = \tfrac{1}{2}(a_{k+1} + b_{k+1}) - \sqrt{a_{k+1}b_{k+1}} = \tfrac{1}{2}\left(a_{k+1} - 2\sqrt{a_{k+1}b_{k+1}} + b_{k+1}\right) = \tfrac{1}{2}\left(\sqrt{a_{k+1}} - \sqrt{b_{k+1}}\right)^2 > 0,$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$a_{k+1} - a_{k+2} = a_{k+1} - \frac{1}{2}(a_{k+1} + b_{k+1}) = \frac{1}{2}(a_{k+1} - b_{k+1}) > 0$, and

$$b_{k+1} - b_{k+2} = b_{k+1} - \sqrt{a_{k+1}b_{k+1}} = \sqrt{b_{k+1}}\left(\sqrt{b_{k+1}} - \sqrt{a_{k+1}}\right) < 0 \quad \Rightarrow \quad a_{k+1} > a_{k+2} > b_{k+2} > b_{k+1},$$

so the assertion is true for $n = k + 1$. Thus, it is true for all n by mathematical induction.

(b) From part (a) we have $a > a_n > a_{n+1} > b_{n+1} > b_n > b$, which shows that both sequences, $\{a_n\}$ and $\{b_n\}$, are monotonic and bounded. So they are both convergent by the Monotonic Sequence Theorem.

(c) Let $\lim\limits_{n\to\infty} a_n = \alpha$ and $\lim\limits_{n\to\infty} b_n = \beta$. Then $\lim\limits_{n\to\infty} a_{n+1} = \lim\limits_{n\to\infty} \dfrac{a_n + b_n}{2} \quad \Rightarrow \quad \alpha = \dfrac{\alpha + \beta}{2} \quad \Rightarrow$ $2\alpha = \alpha + \beta \quad \Rightarrow \quad \alpha = \beta$.

98. (a) Let $\varepsilon > 0$. Since $\lim\limits_{n\to\infty} a_{2n} = L$, there exists N_1 such that $|a_{2n} - L| < \varepsilon$ for $n > N_1$. Since $\lim\limits_{n\to\infty} a_{2n+1} = L$, there exists N_2 such that $|a_{2n+1} - L| < \varepsilon$ for $n > N_2$. Let $N = \max\{2N_1, 2N_2 + 1\}$ and let $n > N$. If n is even, then $n = 2m$ where $m > N_1$, so $|a_n - L| = |a_{2m} - L| < \varepsilon$. If n is odd, then $n = 2m + 1$, where $m > N_2$, so $|a_n - L| = |a_{2m+1} - L| < \varepsilon$. Therefore $\lim\limits_{n\to\infty} a_n = L$.

(b) $a_1 = 1$, $a_2 = 1 + \frac{1}{1+1} = \frac{3}{2} = 1.5$, $a_3 = 1 + \frac{1}{5/2} = \frac{7}{5} = 1.4$, $a_4 = 1 + \frac{1}{12/5} = \frac{17}{12} = 1.41\overline{6}$, $a_5 = 1 + \frac{1}{29/12} = \frac{41}{29} \approx 1.413793$, $a_6 = 1 + \frac{1}{70/29} = \frac{99}{70} \approx 1.414286$, $a_7 = 1 + \frac{1}{169/70} = \frac{239}{169} \approx 1.414201$, $a_8 = 1 + \frac{1}{408/169} = \frac{577}{408} \approx 1.414216$. Notice that $a_1 < a_3 < a_5 < a_7$ and $a_2 > a_4 > a_6 > a_8$. It appears that the odd terms are increasing and the even terms are decreasing. Let's prove that $a_{2n-2} > a_{2n}$ and $a_{2n-1} < a_{2n+1}$ by mathematical induction. Suppose that $a_{2k-2} > a_{2k}$. Then $1 + a_{2k-2} > 1 + a_{2k} \quad \Rightarrow \quad \dfrac{1}{1 + a_{2k-2}} < \dfrac{1}{1 + a_{2k}} \quad \Rightarrow$

$1 + \dfrac{1}{1 + a_{2k-2}} < 1 + \dfrac{1}{1 + a_{2k}} \quad \Rightarrow \quad a_{2k-1} < a_{2k+1} \quad \Rightarrow \quad 1 + a_{2k-1} < 1 + a_{2k+1} \quad \Rightarrow$

$\dfrac{1}{1 + a_{2k-1}} > \dfrac{1}{1 + a_{2k+1}} \quad \Rightarrow \quad 1 + \dfrac{1}{1 + a_{2k-1}} > 1 + \dfrac{1}{1 + a_{2k+1}} \quad \Rightarrow \quad a_{2k} > a_{2k+2}$. We have thus shown, by induction, that the odd terms are increasing and the even terms are decreasing. Also all terms lie between 1 and 2, so both $\{a_n\}$ and $\{b_n\}$ are bounded monotonic sequences and are therefore convergent by the Monotonic Sequence Theorem. Let $\lim\limits_{n\to\infty} a_{2n} = L$. Then $\lim\limits_{n\to\infty} a_{2n+2} = L$ also. We have

$$a_{n+2} = 1 + \frac{1}{1 + 1 + 1/(1 + a_n)} = 1 + \frac{1}{(3 + 2a_n)/(1 + a_n)} = \frac{4 + 3a_n}{3 + 2a_n}$$

so $a_{2n+2} = \dfrac{4 + 3a_{2n}}{3 + 2a_{2n}}$. Taking limits of both sides, we get $L = \dfrac{4 + 3L}{3 + 2L} \quad \Rightarrow \quad 3L + 2L^2 = 4 + 3L \quad \Rightarrow \quad L^2 = 2 \quad \Rightarrow$ $L = \sqrt{2}$ [since $L > 0$]. Thus, $\lim\limits_{n\to\infty} a_{2n} = \sqrt{2}$. Similarly we find that $\lim\limits_{n\to\infty} a_{2n+1} = \sqrt{2}$. So, by part (a), $\lim\limits_{n\to\infty} a_n = \sqrt{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

99. (a) Suppose $\{p_n\}$ converges to p. Then $p_{n+1} = \dfrac{bp_n}{a+p_n} \quad \Rightarrow \quad \lim\limits_{n\to\infty} p_{n+1} = \dfrac{b \lim\limits_{n\to\infty} p_n}{a + \lim\limits_{n\to\infty} p_n} \quad \Rightarrow \quad p = \dfrac{bp}{a+p} \quad \Rightarrow$

$p^2 + ap = bp \quad \Rightarrow \quad p(p+a-b) = 0 \quad \Rightarrow \quad p = 0$ or $p = b - a$.

(b) $p_{n+1} = \dfrac{bp_n}{a+p_n} = \dfrac{\left(\dfrac{b}{a}\right)p_n}{1+\dfrac{p_n}{a}} < \left(\dfrac{b}{a}\right)p_n$ since $1 + \dfrac{p_n}{a} > 1$.

(c) By part (b), $p_1 < \left(\dfrac{b}{a}\right)p_0$, $p_2 < \left(\dfrac{b}{a}\right)p_1 < \left(\dfrac{b}{a}\right)^2 p_0$, $p_3 < \left(\dfrac{b}{a}\right)p_2 < \left(\dfrac{b}{a}\right)^3 p_0$, etc. In general, $p_n < \left(\dfrac{b}{a}\right)^n p_0$,

so $\lim\limits_{n\to\infty} p_n \le \lim\limits_{n\to\infty} \left(\dfrac{b}{a}\right)^n \cdot p_0 = 0$ since $b < a$. $\left[\text{By (9), } \lim\limits_{n\to\infty} r^n = 0 \text{ if } -1 < r < 1. \text{ Here } r = \dfrac{b}{a} \in (0,1)\,.\right]$

(d) Let $a < b$. We first show, by induction, that if $p_0 < b - a$, then $p_n < b - a$ and $p_{n+1} > p_n$.

For $n = 0$, we have $p_1 - p_0 = \dfrac{bp_0}{a+p_0} - p_0 = \dfrac{p_0(b-a-p_0)}{a+p_0} > 0$ since $p_0 < b - a$. So $p_1 > p_0$.

Now we suppose the assertion is true for $n = k$, that is, $p_k < b - a$ and $p_{k+1} > p_k$. Then

$b - a - p_{k+1} = b - a - \dfrac{bp_k}{a+p_k} = \dfrac{a(b-a) + bp_k - ap_k - bp_k}{a+p_k} = \dfrac{a(b-a-p_k)}{a+p_k} > 0$ because $p_k < b - a$. So

$p_{k+1} < b - a$. And $p_{k+2} - p_{k+1} = \dfrac{bp_{k+1}}{a+p_{k+1}} - p_{k+1} = \dfrac{p_{k+1}(b-a-p_{k+1})}{a+p_{k+1}} > 0$ since $p_{k+1} < b - a$. Therefore,

$p_{k+2} > p_{k+1}$. Thus, the assertion is true for $n = k+1$. It is therefore true for all n by mathematical induction.

A similar proof by induction shows that if $p_0 > b - a$, then $p_n > b - a$ and $\{p_n\}$ is decreasing.

In either case the sequence $\{p_n\}$ is bounded and monotonic, so it is convergent by the Monotonic Sequence Theorem. It then follows from part (a) that $\lim\limits_{n\to\infty} p_n = b - a$.

DISCOVERY PROJECT Logistic Sequences

1. To write such a program in Maple it is best to calculate all the points first and then graph them. One possible sequence of commands [taking $p_0 = \frac{1}{2}$ and $k = 1.5$ for the difference equation] is

```
t:='t';p(0):=1/2;k:=1.5;
for j from 1 to 20  do p(j):=k*p(j-1)*(1-p(j-1)) od;
plot([seq([t,p(t)] t=0..20)],t=0..20,p=0..0.5,style=point);
```

In Mathematica, we can use the following program:

```
p[0]=1/2
k=1.5
p[j_]:=k*p[j-1]*(1-p[j-1])
P=Table[p[t],{t,20}]
ListPlot[P]
```

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

With $p_0 = \frac{1}{2}$ and $k = 1.5$:

n	p_n	n	p_n	n	p_n
0	0.5	7	0.3338465076	14	0.3333373303
1	0.375	8	0.3335895255	15	0.3333353318
2	0.3515625	9	0.3334613309	16	0.3333343326
3	0.3419494629	10	0.3333973076	17	0.3333338329
4	0.3375300416	11	0.3333653143	18	0.3333335831
5	0.3354052689	12	0.3333493223	19	0.3333334582
6	0.3343628617	13	0.3333413274	20	0.3333333958

With $p_0 = \frac{1}{2}$ and $k = 2.5$:

n	p_n	n	p_n	n	p_n
0	0.5	7	0.6004164790	14	0.5999967417
1	0.625	8	0.5997913269	15	0.6000016291
2	0.5859375	9	0.6001042277	16	0.5999991854
3	0.6065368651	10	0.5999478590	17	0.6000004073
4	0.5966247409	11	0.6000260637	18	0.5999997964
5	0.6016591486	12	0.5999869664	19	0.6000001018
6	0.5991635437	13	0.6000065164	20	0.5999999491

Both of these sequences seem to converge $\left(\text{the first to about } \frac{1}{3}, \text{ the second to about } 0.60\right)$.

With $p_0 = \frac{7}{8}$ and $k = 1.5$:

n	p_n	n	p_n	n	p_n
0	0.875	7	0.3239166554	14	0.3332554829
1	0.1640625	8	0.3284919837	15	0.3332943990
2	0.2057189941	9	0.3308775005	16	0.3333138639
3	0.2450980344	10	0.3320963702	17	0.3333235980
4	0.2775374819	11	0.3327125567	18	0.3333284655
5	0.3007656421	12	0.3330223670	19	0.3333308994
6	0.3154585059	13	0.3331777051	20	0.3333321164

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

With $p_0 = \frac{7}{8}$ and $k = 2.5$:

n	p_n	n	p_n	n	p_n
0	0.875	7	0.6016572368	14	0.5999869815
1	0.2734375	8	0.5991645155	15	0.6000065088
2	0.4966735840	9	0.6004159972	16	0.5999967455
3	0.6249723374	10	0.5997915688	17	0.6000016272
4	0.5859547872	11	0.6001041070	18	0.5999991864
5	0.6065294364	12	0.5999479194	19	0.6000004068
6	0.5966286980	13	0.6000260335	20	0.5999997966

The limit of the sequence seems to depend on k, but not on p_0.

2. With $p_0 = \frac{7}{8}$ and $k = 3.2$:

n	p_n	n	p_n	n	p_n
0	0.875	7	0.5830728495	14	0.7990633827
1	0.35	8	0.7779164854	15	0.5137954979
2	0.728	9	0.5528397669	16	0.7993909896
3	0.6336512	10	0.7910654689	17	0.5131681132
4	0.7428395416	11	0.5288988570	18	0.7994451225
5	0.6112926626	12	0.7973275394	19	0.5130643795
6	0.7603646184	13	0.5171082698	20	0.7994538304

It seems that eventually the terms fluctuate between two values (about 0.5 and 0.8 in this case).

3. With $p_0 = \frac{7}{8}$ and $k = 3.42$:

n	p_n	n	p_n	n	p_n
0	0.875	7	0.4523028596	14	0.8442074951
1	0.3740625	8	0.8472194412	15	0.4498025048
2	0.8007579316	9	0.4426802161	16	0.8463823232
3	0.5456427596	10	0.8437633929	17	0.4446659586
4	0.8478752457	11	0.4508474156	18	0.8445284520
5	0.4411212220	12	0.8467373602	19	0.4490464985
6	0.8431438501	13	0.4438243545	20	0.8461207931

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

With $p_0 = \frac{7}{8}$ and $k = 3.45$:

n	p_n	n	p_n	n	p_n
0	0.875	7	0.4670259170	14	0.8403376122
1	0.37734375	8	0.8587488490	15	0.4628875685
2	0.8105962830	9	0.4184824586	16	0.8577482026
3	0.5296783241	10	0.8395743720	17	0.4209559716
4	0.8594612299	11	0.4646778983	18	0.8409445432
5	0.4167173034	12	0.8581956045	19	0.4614610237
6	0.8385707740	13	0.4198508858	20	0.8573758782

From the preceding graphs, it seems that for k between 3.4 and 3.5, the terms eventually fluctuate between four values. In the graph below, the pattern followed by the terms is 0.395, 0.832, 0.487, 0.869, 0.395, ... Note that even for $k = 3.42$ (as in the first graph), there are four distinct "branches"; even after 1000 terms, the first and third terms in the pattern differ by about 2×10^{-9}, while the first and fifth terms differ by only 2×10^{-10}. With $p_0 = \frac{7}{8}$ and $k = 3.48$:

4.

$p_0 = 0.5$, $k = 3.7$

$p_0 = 0.501$, $k = 3.7$

$p_0 = 0.75$, $k = 3.9$

$p_0 = 0.749$, $k = 3.9$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$p_0 = 0.5,\ k = 3.999$

From the graphs, it seems that if p_0 is changed by 0.001, the whole graph changes completely. (Note, however, that this might be partially due to accumulated round-off error in the CAS. These graphs were generated by Maple with 100-digit accuracy, and different degrees of accuracy give different graphs.) There seem to be some some fleeting patterns in these graphs, but on the whole they are certainly very chaotic. As k increases, the graph spreads out vertically, with more extreme values close to 0 or 1.

11.2 Series

1. (a) A sequence is an ordered list of numbers whereas a series is the *sum* of a list of numbers.

(b) A series is convergent if the sequence of partial sums is a convergent sequence. A series is divergent if it is not convergent.

2. $\sum\limits_{n=1}^{\infty} a_n = 5$ means that by adding sufficiently many terms of the series we can get as close as we like to the number 5.

In other words, it means that $\lim_{n\to\infty} s_n = 5$, where s_n is the nth partial sum, that is, $\sum\limits_{i=1}^{n} a_i$.

3. $\sum\limits_{n=1}^{\infty} a_n = \lim\limits_{n\to\infty} s_n = \lim\limits_{n\to\infty} [2 - 3(0.8)^n] = \lim\limits_{n\to\infty} 2 - 3 \lim\limits_{n\to\infty} (0.8)^n = 2 - 3(0) = 2$

4. $\sum\limits_{n=1}^{\infty} a_n = \lim\limits_{n\to\infty} s_n = \lim\limits_{n\to\infty} \dfrac{n^2-1}{4n^2+1} = \lim\limits_{n\to\infty} \dfrac{(n^2-1)/n^2}{(4n^2+1)/n^2} = \lim\limits_{n\to\infty} \dfrac{1-1/n^2}{4+1/n^2} = \dfrac{1-0}{4+0} = \dfrac{1}{4}$

5. For $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^3}$, $a_n = \dfrac{1}{n^3}$. $s_1 = a_1 = \dfrac{1}{1^3} = 1$, $s_2 = s_1 + a_2 = 1 + \dfrac{1}{2^3} = 1.125$,

$s_3 = s_2 + a_3 \approx 1.1620$, $s_4 = s_3 + a_4 \approx 1.1777$, $s_5 = s_4 + a_5 \approx 1.1857$, $s_6 = s_5 + a_6 \approx 1.1903$,

$s_7 = s_6 + a_7 \approx 1.1932$, $s_8 = s_7 + a_8 \approx 1.1952$. It appears that the series is convergent.

6. For $\sum\limits_{n=1}^{\infty} \dfrac{1}{\sqrt[3]{n}}$, $a_n = \dfrac{1}{\sqrt[3]{n}}$. $s_1 = a_1 = \dfrac{1}{\sqrt[3]{1}} = 1$, $s_2 = s_1 + a_2 = 1 + \dfrac{1}{\sqrt[3]{2}} \approx 1.7937$,

$s_3 = s_2 + a_3 \approx 2.4871$, $s_4 = s_3 + a_4 \approx 3.1170$, $s_5 = s_4 + a_5 \approx 3.7018$, $s_6 = s_5 + a_6 \approx 4.2521$,

$s_7 = s_6 + a_7 \approx 4.7749$, and $s_8 = s_7 + a_8 \approx 5.2749$. It appears that the series is divergent.

7. For $\sum\limits_{n=1}^{\infty} \sin n$, $a_n = \sin n$. $s_1 = a_1 = \sin 1 \approx 0.8415$, $s_2 = s_1 + a_2 \approx 1.7508$,

$s_3 = s_2 + a_3 \approx 1.8919$, $s_4 = s_3 + a_4 \approx 1.1351$, $s_5 = s_4 + a_5 \approx 0.1762$, $s_6 = s_5 + a_6 \approx -0.1033$,

$s_7 = s_6 + a_7 \approx 0.5537$, and $s_8 = s_7 + a_8 \approx 1.5431$. It appears that the series is divergent.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. For $\sum_{n=1}^{\infty} (-1)^n n$, $a_n = (-1)^n n$. $s_1 = a_1 = (-1)^1(1) = -1$, $s_2 = s_1 + a_2 = -1 + 2 = 1$,

$s_3 = s_2 + a_3 = -2$, $s_4 = s_3 + a_4 = 2$, $s_5 = s_4 + a_5 = -3$, $s_6 = s_5 + a_6 = 3$,

$s_7 = s_6 + a_7 = -4$, $s_8 = s_7 + a_8 = 4$. It appears that the series is divergent.

9. For $\sum_{n=1}^{\infty} \dfrac{1}{n^4 + n^2}$, $a_n = \dfrac{1}{n^4 + n^2}$. $s_1 = a_1 = \dfrac{1}{1^4 + 1^2} = \dfrac{1}{2} = 0.5$, $s_2 = s_1 + a_2 = \dfrac{1}{2} + \dfrac{1}{16 + 4} = 0.55$,

$s_3 = s_2 + a_3 \approx 0.5611$, $s_4 = s_3 + a_4 \approx 0.5648$, $s_5 = s_4 + a_5 \approx 0.5663$, $s_6 = s_5 + a_6 \approx 0.5671$,

$s_7 = s_6 + a_7 \approx 0.5675$, and $s_8 = s_7 + a_8 \approx 0.5677$. It appears that the series is convergent.

10. For $\sum_{n=1}^{\infty} \dfrac{(-1)^{n-1}}{n!}$, $a_n = (-1)^{n-1}\dfrac{1}{n!}$. $s_1 = a_1 = \dfrac{1}{1!} = 1$, $s_2 = s_1 + a_2 = 1 - \dfrac{1}{2!} = 0.5$,

$s_3 = s_2 + a_3 = 0.5 + \dfrac{1}{3!} \approx 0.6667$, $s_4 = s_3 + a_4 = 0.625$, $s_5 = s_4 + a_5 \approx 0.6333$, $s_6 = s_5 + a_6 \approx 0.6319$,

$s_7 = s_6 + a_7 \approx 0.6321$, and $s_8 = s_7 + a_8 \approx 0.6321$. It appears that the series is convergent.

11.

n	s_n
1	-2
2	-1.33333
3	-1.55556
4	-1.48148
5	-1.50617
6	-1.49794
7	-1.50069
8	-1.49977
9	-1.50008
10	-1.49997

From the graph and the table, it seems that the series converges to -1.5. In fact, it is a geometric series with $a = -2$ and $r = -\dfrac{1}{3}$, so its sum is $\sum_{n=1}^{\infty} \dfrac{6}{(-3)^n} = \dfrac{-2}{1 - \left(-\frac{1}{3}\right)} = -1.5$.

Note that the point corresponding to $n = 1$ is part of both $\{a_n\}$ and $\{s_n\}$.

12.

n	s_n
1	0.54030
2	0.12416
3	-0.86584
4	-1.51948
5	-1.23582
6	-0.27565
7	0.47825
8	0.33275
9	-0.57838
10	-1.41745

The series $\sum_{n=1}^{\infty} \cos n$ diverges, since its terms do not approach 0.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13.

n	s_n
1	0.44721
2	1.15432
3	1.98637
4	2.88080
5	3.80927
6	4.75796
7	5.71948
8	6.68962
9	7.66581
10	8.64639

The series $\sum_{n=1}^{\infty} \frac{n}{\sqrt{n^2+4}}$ diverges, since its terms do not approach 0.

14.

n	s_n
1	4.90000
2	8.33000
3	10.73100
4	12.41170
5	13.58819
6	14.41173
7	14.98821
8	15.39175
9	15.67422
10	15.87196

From the graph and the table, we see that the terms are getting smaller and may approach 0, and that the series approaches a value near 16. The series is geometric with $a_1 = 4.9$ and $r = 0.7$, so its sum is $\sum_{n=1}^{\infty} \frac{7^{n+1}}{10^n} = \frac{4.9}{1-0.7} = \frac{4.9}{0.3} = 16.\overline{3}$.

15. (a) $\lim_{n\to\infty} a_n = \lim_{n\to\infty} \frac{2n}{3n+1} = \frac{2}{3}$, so the *sequence* $\{a_n\}$ is convergent by (11.1.1).

(b) Since $\lim_{n\to\infty} a_n = \frac{2}{3} \neq 0$, the *series* $\sum_{n=1}^{\infty} a_n$ is divergent by the Test for Divergence.

16. (a) Both $\sum_{i=1}^{n} a_i$ and $\sum_{j=1}^{n} a_j$ represent the sum of the first n terms of the sequence $\{a_n\}$, that is, the nth partial sum.

(b) $\sum_{i=1}^{n} a_j = \underbrace{a_j + a_j + \cdots + a_j}_{n \text{ terms}} = na_j$, which, in general, is not the same as $\sum_{i=1}^{n} a_i = a_1 + a_2 + \cdots + a_n$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. For the series $\sum_{n=1}^{\infty}\left(\frac{1}{n+2}-\frac{1}{n}\right)$,

$$s_n=\sum_{i=1}^{n}\left(\frac{1}{i+2}-\frac{1}{i}\right)$$

$$=\left(\frac{1}{3}-1\right)+\left(\frac{1}{4}-\frac{1}{2}\right)+\left(\frac{1}{5}-\frac{1}{3}\right)+\left(\frac{1}{6}-\frac{1}{4}\right)+\cdots+\left(\frac{1}{n}-\frac{1}{n-2}\right)+\left(\frac{1}{n+1}-\frac{1}{n-1}\right)+\left(\frac{1}{n+2}-\frac{1}{n}\right)$$

$$=-1-\frac{1}{2}+\frac{1}{n+1}+\frac{1}{n+2}\qquad\text{[telescoping series]}$$

Thus, $\sum_{n=1}^{\infty}\left(\frac{1}{n+2}-\frac{1}{n}\right)=\lim_{n\to\infty}s_n=\lim_{n\to\infty}\left(-1-\frac{1}{2}+\frac{1}{n+1}+\frac{1}{n+2}\right)=-1-\frac{1}{2}=-\frac{3}{2}$. Converges

18. For the series $\sum_{n=4}^{\infty}\left(\frac{1}{\sqrt{n}}-\frac{1}{\sqrt{n+1}}\right)$

$$s_n=\sum_{i=4}^{n}\left(\frac{1}{\sqrt{i}}-\frac{1}{\sqrt{i+1}}\right)=\left(\frac{1}{\sqrt{4}}-\frac{1}{\sqrt{5}}\right)+\left(\frac{1}{\sqrt{5}}-\frac{1}{\sqrt{6}}\right)+\cdots+\left(\frac{1}{\sqrt{n}}-\frac{1}{\sqrt{n+1}}\right)=\frac{1}{\sqrt{4}}-\frac{1}{\sqrt{n+1}}$$

[telescoping series]

Thus, $\sum_{n=4}^{\infty}\left(\frac{1}{\sqrt{n}}-\frac{1}{\sqrt{n+1}}\right)=\lim_{n\to\infty}s_n=\lim_{n\to\infty}\left(\frac{1}{\sqrt{4}}-\frac{1}{\sqrt{n+1}}\right)=\frac{1}{\sqrt{4}}-0=\frac{1}{2}$. Converges

19. For the series $\sum_{n=1}^{\infty}\frac{3}{n(n+3)}$, $s_n=\sum_{i=1}^{n}\frac{3}{i(i+3)}=\sum_{i=1}^{n}\left(\frac{1}{i}-\frac{1}{i+3}\right)$ [using partial fractions]. The latter sum is

$$\left(1-\tfrac{1}{4}\right)+\left(\tfrac{1}{2}-\tfrac{1}{5}\right)+\left(\tfrac{1}{3}-\tfrac{1}{6}\right)+\left(\tfrac{1}{4}-\tfrac{1}{7}\right)+\cdots+\left(\tfrac{1}{n-3}-\tfrac{1}{n}\right)+\left(\tfrac{1}{n-2}-\tfrac{1}{n+1}\right)+\left(\tfrac{1}{n-1}-\tfrac{1}{n+2}\right)+\left(\tfrac{1}{n}-\tfrac{1}{n+3}\right)$$

$$=1+\tfrac{1}{2}+\tfrac{1}{3}-\tfrac{1}{n+1}-\tfrac{1}{n+2}-\tfrac{1}{n+3}\qquad\text{[telescoping series]}$$

Thus, $\sum_{n=1}^{\infty}\frac{3}{n(n+3)}=\lim_{n\to\infty}s_n=\lim_{n\to\infty}\left(1+\tfrac{1}{2}+\tfrac{1}{3}-\tfrac{1}{n+1}-\tfrac{1}{n+2}-\tfrac{1}{n+3}\right)=1+\tfrac{1}{2}+\tfrac{1}{3}=\tfrac{11}{6}$. Converges

20. For the series $\sum_{n=1}^{\infty}\ln\frac{n}{n+1}$,

$$s_n=(\ln 1-\ln 2)+(\ln 2-\ln 3)+(\ln 3-\ln 4)+\cdots+[\ln n-\ln(n+1)]=\ln 1-\ln(n+1)=-\ln(n+1)$$

[telescoping series]

Thus, $\lim_{n\to\infty}s_n=-\infty$, so the series is divergent.

21. For the series $\sum_{n=1}^{\infty}\left(e^{1/n}-e^{1/(n+1)}\right)$,

$$s_n=\sum_{i=1}^{n}\left(e^{1/i}-e^{1/(i+1)}\right)=\left(e^{1}-e^{1/2}\right)+\left(e^{1/2}-e^{1/3}\right)+\cdots+\left(e^{1/n}-e^{1/(n+1)}\right)=e-e^{1/(n+1)}$$

[telescoping series]

Thus, $\sum_{n=1}^{\infty}\left(e^{1/n}-e^{1/(n+1)}\right)=\lim_{n\to\infty}s_n=\lim_{n\to\infty}\left(e-e^{1/(n+1)}\right)=e-e^{0}=e-1$. Converges

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. Using partial fractions, the partial sums of the series $\sum_{n=2}^{\infty} \frac{1}{n^3 - n}$ are

$$s_n = \sum_{i=2}^{n} \frac{1}{i(i-1)(i+1)} = \sum_{i=2}^{n} \left(-\frac{1}{i} + \frac{1/2}{i-1} + \frac{1/2}{i+1} \right) = \frac{1}{2} \sum_{i=2}^{n} \left(\frac{1}{i-1} - \frac{2}{i} + \frac{1}{i+1} \right)$$

$$= \frac{1}{2} \left[\left(\frac{1}{1} - \frac{2}{2} + \frac{1}{3} \right) + \left(\frac{1}{2} - \frac{2}{3} + \frac{1}{4} \right) + \left(\frac{1}{3} - \frac{2}{4} + \frac{1}{5} \right) + \left(\frac{1}{4} - \frac{2}{5} + \frac{1}{6} \right) + \cdots \right.$$

$$\left. + \left(\frac{1}{n-3} - \frac{2}{n-2} + \frac{1}{n-1} \right) + \left(\frac{1}{n-2} - \frac{2}{n-1} + \frac{1}{n} \right) + \left(\frac{1}{n-1} - \frac{2}{n} + \frac{1}{n+1} \right) \right]$$

Note: In three consecutive expressions in parentheses, the 3rd term in the first expression plus the 2nd term in the second expression plus the 1st term in the third expression sum to 0.

$$= \frac{1}{2} \left(\frac{1}{1} - \frac{2}{2} + \frac{1}{2} + \frac{1}{n} - \frac{2}{n} + \frac{1}{n+1} \right) = \frac{1}{4} - \frac{1}{2n} + \frac{1}{2n+2}$$

Thus, $\sum_{n=2}^{\infty} \frac{1}{n^3 - n} = \lim_{n \to \infty} s_n = \lim_{n \to \infty} \left(\frac{1}{4} - \frac{1}{2n} + \frac{1}{2n+2} \right) = \frac{1}{4}$.

23. $3 - 4 + \frac{16}{3} - \frac{64}{9} + \cdots$ is a geometric series with ratio $r = -\frac{4}{3}$. Since $|r| = \frac{4}{3} > 1$, the series diverges.

24. $4 + 3 + \frac{9}{4} + \frac{27}{16} + \cdots$ is a geometric series with ratio $\frac{3}{4}$. Since $|r| = \frac{3}{4} < 1$, the series converges to $\frac{a}{1-r} = \frac{4}{1 - 3/4} = 16$.

25. $10 - 2 + 0.4 - 0.08 + \cdots$ is a geometric series with ratio $-\frac{2}{10} = -\frac{1}{5}$. Since $|r| = \frac{1}{5} < 1$, the series converges to

$\frac{a}{1-r} = \frac{10}{1 - (-1/5)} = \frac{10}{6/5} = \frac{50}{6} = \frac{25}{3}$.

26. $2 + 0.5 + 0.125 + 0.03125 + \cdots$ is a geometric series with ratio $r = \frac{0.5}{2} = \frac{1}{4}$. Since $|r| = \frac{1}{4} < 1$, the series converges

to $\frac{a}{1-r} = \frac{2}{1 - 1/4} = \frac{2}{3/4} = \frac{8}{3}$.

27. $\sum_{n=1}^{\infty} 12\,(0.73)^{n-1}$ is a geometric series with first term $a = 12$ and ratio $r = 0.73$. Since $|r| = 0.73 < 1$, the series converges

to $\frac{a}{1-r} = \frac{12}{1 - 0.73} = \frac{12}{0.27} = \frac{12(100)}{27} = \frac{400}{9}$.

28. $\sum_{n=1}^{\infty} \frac{5}{\pi^n} = 5 \sum_{n=1}^{\infty} \left(\frac{1}{\pi} \right)^n$. The latter series is geometric with $a = \frac{1}{\pi}$ and ratio $r = \frac{1}{\pi}$. Since $|r| = \frac{1}{\pi} < 1$, it converges to

$\frac{1/\pi}{1 - 1/\pi} = \frac{1}{\pi - 1}$. Thus, the given series converges to $5 \left(\frac{1}{\pi - 1} \right) = \frac{5}{\pi - 1}$.

29. $\sum_{n=1}^{\infty} \frac{(-3)^{n-1}}{4^n} = \frac{1}{4} \sum_{n=1}^{\infty} \left(-\frac{3}{4} \right)^{n-1}$. The latter series is geometric with $a = 1$ and ratio $r = -\frac{3}{4}$. Since $|r| = \frac{3}{4} < 1$, it

converges to $\frac{1}{1 - (-3/4)} = \frac{4}{7}$. Thus, the given series converges to $\left(\frac{1}{4}\right)\left(\frac{4}{7}\right) = \frac{1}{7}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $\displaystyle\sum_{n=0}^{\infty} \frac{3^{n+1}}{(-2)^n} = 3\sum_{n=0}^{\infty} \left(-\tfrac{3}{2}\right)^n$ is a geometric series with ratio $r = -\tfrac{3}{2}$. Since $|r| = \tfrac{3}{2} > 1$, the series diverges.

31. $\displaystyle\sum_{n=1}^{\infty} \frac{e^{2n}}{6^{n-1}} = \sum_{n=1}^{\infty} \frac{(e^2)^n}{6^n 6^{-1}} = 6\sum_{n=1}^{\infty} \left(\frac{e^2}{6}\right)^n$ is a geometric series with ratio $r = \dfrac{e^2}{6}$. Since $|r| = \dfrac{e^2}{6}[\approx 1.23] > 1$, the series diverges.

32. $\displaystyle\sum_{n=1}^{\infty} \frac{6 \cdot 2^{2n-1}}{3^n} = \sum_{n=1}^{\infty} \frac{6(2^2)^n \cdot 2^{-1}}{3^n} = 3\sum_{n=1}^{\infty} \left(\frac{4}{3}\right)^n$ is a geometric series with ratio $r = \dfrac{4}{3}$. Since $|r| = \dfrac{4}{3} > 1$, the series diverges.

33. $\dfrac{1}{3} + \dfrac{1}{6} + \dfrac{1}{9} + \dfrac{1}{12} + \dfrac{1}{15} + \cdots = \displaystyle\sum_{n=1}^{\infty} \frac{1}{3n} = \frac{1}{3}\sum_{n=1}^{\infty} \frac{1}{n}$. This is a constant multiple of the divergent harmonic series, so it diverges.

34. $\dfrac{1}{2} + \dfrac{2}{3} + \dfrac{3}{4} + \dfrac{4}{5} + \dfrac{5}{6} + \dfrac{6}{7} + \cdots = \displaystyle\sum_{n=1}^{\infty} \frac{n}{n+1}$. This series diverges by the Test for Divergence since

$\displaystyle\lim_{n\to\infty} a_n = \lim_{n\to\infty} \frac{n}{n+1} = \lim_{n\to\infty} \frac{1}{1+1/n} = 1 \neq 0.$

35. $\dfrac{2}{5} + \dfrac{4}{25} + \dfrac{8}{125} + \dfrac{16}{625} + \dfrac{32}{3125} + \cdots = \displaystyle\sum_{n=1}^{\infty} \left(\frac{2}{5}\right)^n$. This series is geometric with $a = \dfrac{2}{5}$ and ratio $r = \dfrac{2}{5}$. Since $|r| = \dfrac{2}{5} < 1$, it converges to $\dfrac{2/5}{1-2/5} = \dfrac{2}{3}$.

36. $\frac{1}{3} + \frac{2}{9} + \frac{1}{27} + \frac{2}{81} + \frac{1}{243} + \frac{2}{729} + \cdots = \left(\frac{1}{3} + \frac{1}{27} + \frac{1}{243} + \cdots\right) + \left(\frac{2}{9} + \frac{2}{81} + \frac{2}{729} + \cdots\right)$, which are both convergent geometric series with sums $\dfrac{1/3}{1-1/9} = \dfrac{3}{8}$ and $\dfrac{2/9}{1-1/9} = \dfrac{1}{4}$, so the original series converges and its sum is $\frac{3}{8} + \frac{1}{4} = \frac{5}{8}$.

37. $\displaystyle\sum_{n=1}^{\infty} \frac{2+n}{1-2n}$ diverges by the Test for Divergence since $\displaystyle\lim_{n\to\infty} a_n = \lim_{n\to\infty} \frac{2+n}{1-2n} = \lim_{n\to\infty} \frac{2/n+1}{1/n-2} = -\frac{1}{2} \neq 0.$

38. $\displaystyle\sum_{k=1}^{\infty} \frac{k^2}{k^2-2k+5}$ diverges by the Test for Divergence since $\displaystyle\lim_{k\to\infty} \frac{k^2}{k^2-2k+5} = \lim_{k\to\infty} \frac{1}{1-2/k+5/k^2} = 1 \neq 0.$

39. $\displaystyle\sum_{n=1}^{\infty} 3^{n+1}4^{-n} = \sum_{n=1}^{\infty} \frac{3^n \cdot 3^1}{4^n} = 3\sum_{n=1}^{\infty} \left(\frac{3}{4}\right)^n$. The latter series is geometric with $a = \dfrac{3}{4}$ and ratio $r = \dfrac{3}{4}$. Since $|r| = \dfrac{3}{4} < 1$, it converges to $\dfrac{3/4}{1-3/4} = 3$. Thus, the given series converges to $3(3) = 9$.

40.
$$\begin{aligned}\sum_{n=1}^{\infty} [(-0.2)^n + (0.6)^{n-1}] &= \sum_{n=1}^{\infty}(-0.2)^n + \sum_{n=1}^{\infty}(0.6)^{n-1} \qquad \text{[sum of two geometric series]}\\ &= \frac{-0.2}{1-(-0.2)} + \frac{1}{1-0.6} = -\frac{1}{6} + \frac{5}{2} = \frac{7}{3}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. $\displaystyle\sum_{n=1}^{\infty} \frac{1}{4+e^{-n}}$ diverges by the Test for Divergence since $\displaystyle\lim_{n\to\infty} \frac{1}{4+e^{-n}} = \frac{1}{4+0} = \frac{1}{4} \neq 0$.

42. $\displaystyle\sum_{n=1}^{\infty} \frac{2^n+4^n}{e^n}$ diverges by the Test for Divergence since $\displaystyle\lim_{n\to\infty} \frac{2^n+4^n}{e^n} = \lim_{n\to\infty} \left(\frac{2^n}{e^n} + \frac{4^n}{e^n}\right) \geq \lim_{n\to\infty} \left(\frac{4}{e}\right)^n = \infty$ since $\dfrac{4}{e} > 1$.

43. $\displaystyle\sum_{k=1}^{\infty} (\sin 100)^k$ is a geometric series with first term $a = \sin 100$ [≈ -0.506] and ratio $r = \sin 100$. Since $|r| < 1$, the series converges to $\dfrac{\sin 100}{1-\sin 100} \approx -0.336$.

44. $\displaystyle\sum_{n=1}^{\infty} \frac{1}{1+\left(\frac{2}{3}\right)^n}$ diverges by the Test for Divergence since $\displaystyle\lim_{n\to\infty} \frac{1}{1+\left(\frac{2}{3}\right)^n} = \frac{1}{1+0} = 1 \neq 0$.

45. $\displaystyle\sum_{n=1}^{\infty} \ln\left(\frac{n^2+1}{2n^2+1}\right)$ diverges by the Test for Divergence since

$$\lim_{n\to\infty} a_n = \lim_{n\to\infty} \ln\left(\frac{n^2+1}{2n^2+1}\right) = \ln\left(\lim_{n\to\infty} \frac{n^2+1}{2n^2+1}\right) = \ln \tfrac{1}{2} \neq 0.$$

46. $\displaystyle\sum_{k=0}^{\infty} (\sqrt{2})^{-k} = \sum_{k=0}^{\infty} \left(\frac{1}{\sqrt{2}}\right)^k$ is a geometric series with first term $a = \left(\dfrac{1}{\sqrt{2}}\right)^0 = 1$ and ratio $r = \dfrac{1}{\sqrt{2}}$. Since $|r| < 1$, the series converges to $\dfrac{1}{1-1/\sqrt{2}} = \dfrac{\sqrt{2}}{\sqrt{2}-1} \approx 3.414$.

47. $\displaystyle\sum_{n=1}^{\infty} \arctan n$ diverges by the Test for Divergence since $\displaystyle\lim_{n\to\infty} a_n = \lim_{n\to\infty} \arctan n = \tfrac{\pi}{2} \neq 0$.

48. $\displaystyle\sum_{n=1}^{\infty} \left(\frac{3}{5^n} + \frac{2}{n}\right)$ diverges because $\displaystyle\sum_{n=1}^{\infty} \frac{2}{n} = 2\sum_{n=1}^{\infty} \frac{1}{n}$ diverges. (If it converged, then $\displaystyle\frac{1}{2} \cdot 2\sum_{n=1}^{\infty} \frac{1}{n}$ would also converge by Theorem 8(i), but we know from Example 9 that the harmonic series $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.) If the given series converges, then the difference $\displaystyle\sum_{n=1}^{\infty} \left(\frac{3}{5^n} + \frac{2}{n}\right) - \sum_{n=1}^{\infty} \frac{3}{5^n}$ must converge (since $\displaystyle\sum_{n=1}^{\infty} \frac{3}{5^n}$ is a convergent geometric series) and equal $\displaystyle\sum_{n=1}^{\infty} \frac{2}{n}$, but we have just seen that $\displaystyle\sum_{n=1}^{\infty} \frac{2}{n}$ diverges, so the given series must also diverge.

49. $\displaystyle\sum_{n=1}^{\infty} \frac{1}{e^n} = \sum_{n=1}^{\infty} \left(\frac{1}{e}\right)^n$ is a geometric series with first term $a = \dfrac{1}{e}$ and ratio $r = \dfrac{1}{e}$. Since $|r| = \dfrac{1}{e} < 1$, the series converges to $\dfrac{1/e}{1-1/e} = \dfrac{1/e}{1-1/e} \cdot \dfrac{e}{e} = \dfrac{1}{e-1}$. By Example 8, $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1$. Thus, by Theorem 8(ii),

$$\sum_{n=1}^{\infty} \left(\frac{1}{e^n} + \frac{1}{n(n+1)}\right) = \sum_{n=1}^{\infty} \frac{1}{e^n} + \sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \frac{1}{e-1} + 1 = \frac{1}{e-1} + \frac{e-1}{e-1} = \frac{e}{e-1}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50. $\sum_{n=1}^{\infty} \frac{e^n}{n^2}$ diverges by the Test for Divergence since $\lim_{n\to\infty} a_n = \lim_{n\to\infty} \frac{e^n}{n^2} = \lim_{x\to\infty} \frac{e^x}{x^2} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{e^x}{2x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{e^x}{2} = \infty \neq 0$.

51. (a) Many people would guess that $x < 1$, but note that x consists of an infinite number of 9s.

(b) $x = 0.99999\ldots = \frac{9}{10} + \frac{9}{100} + \frac{9}{1000} + \frac{9}{10{,}000} + \cdots = \sum_{n=1}^{\infty} \frac{9}{10^n}$, which is a geometric series with $a_1 = 0.9$ and $r = 0.1$. Its sum is $\frac{0.9}{1-0.1} = \frac{0.9}{0.9} = 1$, that is, $x = 1$.

(c) The number 1 has two decimal representations, $1.00000\ldots$ and $0.99999\ldots$.

(d) Except for 0, all rational numbers that have a terminating decimal representation can be written in more than one way. For example, 0.5 can be written as $0.49999\ldots$ as well as $0.50000\ldots$.

52. $a_1 = 1$, $a_n = (5-n)a_{n-1} \quad \Rightarrow \quad a_2 = (5-2)a_1 = 3(1) = 3,\ a_3 = (5-3)a_2 = 2(3) = 6,\ \ a_4 = (5-4)a_3 = 1(6) = 6,$ $a_5 = (5-5)a_4 = 0$, and all succeeding terms equal 0. Thus, $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{4} a_n = 1 + 3 + 6 + 6 = 16$.

53. $0.\overline{8} = \frac{8}{10} + \frac{8}{10^2} + \cdots$ is a geometric series with $a = \frac{8}{10}$ and $r = \frac{1}{10}$. It converges to $\frac{a}{1-r} = \frac{8/10}{1-1/10} = \frac{8}{9}$.

54. $0.\overline{46} = \frac{46}{100} + \frac{46}{100^2} + \cdots$ is a geometric series with $a = \frac{46}{100}$ and $r = \frac{1}{100}$. It converges to $\frac{a}{1-r} = \frac{46/100}{1-1/100} = \frac{46}{99}$.

55. $2.\overline{516} = 2 + \frac{516}{10^3} + \frac{516}{10^6} + \cdots$. Now $\frac{516}{10^3} + \frac{516}{10^6} + \cdots$ is a geometric series with $a = \frac{516}{10^3}$ and $r = \frac{1}{10^3}$. It converges to $\frac{a}{1-r} = \frac{516/10^3}{1-1/10^3} = \frac{516/10^3}{999/10^3} = \frac{516}{999}$. Thus, $2.\overline{516} = 2 + \frac{516}{999} = \frac{2514}{999} = \frac{838}{333}$.

56. $10.1\overline{35} = 10.1 + \frac{35}{10^3} + \frac{35}{10^5} + \cdots$. Now $\frac{35}{10^3} + \frac{35}{10^5} + \cdots$ is a geometric series with $a = \frac{35}{10^3}$ and $r = \frac{1}{10^2}$. It converges to $\frac{a}{1-r} = \frac{35/10^3}{1-1/10^2} = \frac{35/10^3}{99/10^2} = \frac{35}{990}$. Thus, $10.1\overline{35} = 10.1 + \frac{35}{990} = \frac{9999+35}{990} = \frac{10{,}034}{990} = \frac{5017}{495}$.

57. $1.234\overline{567} = 1.234 + \frac{567}{10^6} + \frac{567}{10^9} + \cdots$. Now $\frac{567}{10^6} + \frac{567}{10^9} + \cdots$ is a geometric series with $a = \frac{567}{10^6}$ and $r = \frac{1}{10^3}$. It converges to $\frac{a}{1-r} = \frac{567/10^6}{1-1/10^3} = \frac{567/10^6}{999/10^3} = \frac{567}{999{,}000} = \frac{21}{37{,}000}$. Thus,

$1.234\overline{567} = 1.234 + \frac{21}{37{,}000} = \frac{1234}{1000} + \frac{21}{37{,}000} = \frac{45{,}658}{37{,}000} + \frac{21}{37{,}000} = \frac{45{,}679}{37{,}000}$.

58. $5.\overline{71358} = 5 + \frac{71{,}358}{10^5} + \frac{71{,}358}{10^{10}} + \cdots$. Now $\frac{71{,}358}{10^5} + \frac{71{,}358}{10^{10}} + \cdots$ is a geometric series with $a = \frac{71{,}358}{10^5}$ and $r = \frac{1}{10^5}$. It converges to $\frac{a}{1-r} = \frac{71{,}358/10^5}{1-1/10^5} = \frac{71{,}358/10^5}{99{,}999/10^5} = \frac{71{,}358}{99{,}999} = \frac{23{,}786}{33{,}333}$. Thus,

$5.\overline{71358} = 5 + \frac{23{,}786}{33{,}333} = \frac{166{,}665}{33{,}333} + \frac{23{,}786}{33{,}333} = \frac{190{,}451}{33{,}333}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. $\sum_{n=1}^{\infty}(-5)^n x^n = \sum_{n=1}^{\infty}(-5x)^n$ is a geometric series with $r=-5x$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $|-5x|<1$ $\Leftrightarrow$ $|x|<\frac{1}{5}$, that is, $-\frac{1}{5}<x<\frac{1}{5}$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{-5x}{1-(-5x)}=\dfrac{-5x}{1+5x}$.

60. $\sum_{n=1}^{\infty}(x+2)^n$ is a geometric series with $r=x+2$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $|x+2|<1$ $\Leftrightarrow$ $-1<x+2<1$ $\Leftrightarrow$ $-3<x<-1$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{x+2}{1-(x+2)}=\dfrac{x+2}{-x-1}$.

61. $\sum_{n=0}^{\infty}\dfrac{(x-2)^n}{3^n}=\sum_{n=0}^{\infty}\left(\dfrac{x-2}{3}\right)^n$ is a geometric series with $r=\dfrac{x-2}{3}$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $\left|\dfrac{x-2}{3}\right|<1$ $\Leftrightarrow$ $-1<\dfrac{x-2}{3}<1$ $\Leftrightarrow$ $-3<x-2<3$ $\Leftrightarrow$ $-1<x<5$. In that case, the sum of the series is

$$\frac{a}{1-r}=\frac{1}{1-\dfrac{x-2}{3}}=\frac{1}{\dfrac{3-(x-2)}{3}}=\frac{3}{5-x}.$$

62. $\sum_{n=0}^{\infty}(-4)^n(x-5)^n=\sum_{n=0}^{\infty}[-4(x-5)]^n$ is a geometric series with $r=-4(x-5)$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $|-4(x-5)|<1$ $\Leftrightarrow$ $|x-5|<\frac{1}{4}$ $\Leftrightarrow$ $-\frac{1}{4}<x-5<\frac{1}{4}$ $\Leftrightarrow$ $\frac{19}{4}<x<\frac{21}{4}$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{1}{1-[-4(x-5)]}=\dfrac{1}{4x-19}$.

63. $\sum_{n=0}^{\infty}\dfrac{2^n}{x^n}=\sum_{n=0}^{\infty}\left(\dfrac{2}{x}\right)^n$ is a geometric series with $r=\dfrac{2}{x}$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $\left|\dfrac{2}{x}\right|<1$ $\Leftrightarrow$ $2<|x|$ $\Leftrightarrow$ $x>2$ or $x<-2$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{1}{1-2/x}=\dfrac{x}{x-2}$.

64. $\sum_{n=0}^{\infty}\dfrac{x^n}{2^n}=\sum_{n=0}^{\infty}\left(\dfrac{x}{2}\right)^n$ is a geometric series with $r=\dfrac{x}{2}$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $\left|\dfrac{x}{2}\right|<1$ $\Leftrightarrow$ $-1<\dfrac{x}{2}<1$ $\Leftrightarrow$ $-2<x<2$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{1}{1-\frac{x}{2}}=\dfrac{2}{2-x}$.

65. $\sum_{n=0}^{\infty}e^{nx}=\sum_{n=0}^{\infty}(e^x)^n$ is a geometric series with $r=e^x$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $|e^x|<1$ $\Leftrightarrow$ $-1<e^x<1$ $\Leftrightarrow$ $0<e^x<1$ $\Leftrightarrow$ $x<0$. In that case, the sum of the series is $\dfrac{a}{1-r}=\dfrac{1}{1-e^x}$.

66. $\sum_{n=0}^{\infty}\dfrac{\sin^n x}{3^n}=\sum_{n=0}^{\infty}\left(\dfrac{\sin x}{3}\right)^n$ is a geometric series with $r=\dfrac{\sin x}{3}$, so the series converges $\Leftrightarrow$ $|r|<1$ $\Leftrightarrow$ $\left|\dfrac{\sin x}{3}\right|<1$ $\Leftrightarrow$ $|\sin x|<3$, which is true for all x. Thus, the sum of the series is $\dfrac{a}{1-r}=\dfrac{1}{1-(\sin x)/3}=\dfrac{3}{3-\sin x}$.

67. After defining f, We use `convert(f,parfrac);` in Maple or `Apart` in Mathematica to find that the general term is $\dfrac{3n^2+3n+1}{(n^2+n)^3}=\dfrac{1}{n^3}-\dfrac{1}{(n+1)^3}$. So the *n*th partial sum is

$$s_n=\sum_{k=1}^{n}\left(\frac{1}{k^3}-\frac{1}{(k+1)^3}\right)=\left(1-\frac{1}{2^3}\right)+\left(\frac{1}{2^3}-\frac{1}{3^3}\right)+\cdots+\left(\frac{1}{n^3}-\frac{1}{(n+1)^3}\right)=1-\frac{1}{(n+1)^3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The series converges to $\lim_{n\to\infty} s_n = 1$. This can be confirmed by directly computing the sum using `sum(f,n=1..infinity);` (in Maple) or `Sum[f,{n,1,Infinity}]` (in Mathematica).

68. See Exercise 67 for specific CAS commands.

$\dfrac{1}{n^5-5n^3+4n} = \dfrac{1}{24(n-2)} + \dfrac{1}{24(n+2)} - \dfrac{1}{6(n-1)} - \dfrac{1}{6(n+1)} + \dfrac{1}{4n}$. So the nth partial sum is

$$\begin{aligned} s_n &= \frac{1}{24}\sum_{k=3}^{n}\left(\frac{1}{k-2} - \frac{4}{k-1} + \frac{6}{k} - \frac{4}{k+1} + \frac{1}{k+2}\right) \\ &= \frac{1}{24}\left[\left(\frac{1}{1} - \frac{4}{2} + \frac{6}{3} - \frac{4}{4} + \frac{1}{5}\right) + \cdots + \left(\frac{1}{n-2} - \frac{4}{n-1} + \frac{6}{n} - \frac{4}{n+1} + \frac{1}{n+2}\right)\right] \end{aligned}$$

The terms with denominator 5 or greater cancel, except for a few terms with n in the denominator. So as $n \to \infty$,

$s_n \to \dfrac{1}{24}\left(\dfrac{1}{1} - \dfrac{3}{2} + \dfrac{3}{3} - \dfrac{1}{4}\right) = \dfrac{1}{24}\left(\dfrac{1}{4}\right) = \dfrac{1}{96}$.

69. For $n = 1$, $a_1 = 0$ since $s_1 = 0$. For $n > 1$,

$$a_n = s_n - s_{n-1} = \frac{n-1}{n+1} - \frac{(n-1)-1}{(n-1)+1} = \frac{(n-1)n-(n+1)(n-2)}{(n+1)n} = \frac{2}{n(n+1)}$$

Also, $\sum_{n=1}^{\infty} a_n = \lim_{n\to\infty} s_n = \lim_{n\to\infty} \dfrac{1-1/n}{1+1/n} = 1$.

70. $a_1 = s_1 = 3 - \frac{1}{2} = \frac{5}{2}$. For $n \neq 1$,

$$a_n = s_n - s_{n-1} = \left(3 - n2^{-n}\right) - \left[3 - (n-1)2^{-(n-1)}\right] = -\frac{n}{2^n} + \frac{n-1}{2^{n-1}} \cdot \frac{2}{2} = \frac{2(n-1)}{2^n} - \frac{n}{2^n} = \frac{n-2}{2^n}$$

Also, $\sum_{n=1}^{\infty} a_n = \lim_{n\to\infty} s_n = \lim_{n\to\infty}\left(3 - \dfrac{n}{2^n}\right) = 3$ because $\lim_{x\to\infty} \dfrac{x}{2^x} \overset{\text{H}}{=} \lim_{x\to\infty} \dfrac{1}{2^x \ln 2} = 0$.

71. (a) The quantity of the drug in the body after the first tablet is 100 mg. After 8 hours, the body eliminates 75% of the drug, which means 25% remains. Thus, after the second tablet, there is 100 mg plus 25% of the first 100-mg tablet, that is, $100 + 0.25(100) = 125$ mg. After the third tablet, the quantity is $100 + 0.25(125)$ or, equivalently, $100 + 100(0.25) + 100(0.25)^2$. Either expression gives 131.25 mg.

(b) From part (a), we see that $Q_{n+1} = 100 + 0.25\,Q_n$.

(c) $Q_n = 100 + 100(0.25)^1 + 100(0.25)^2 + \cdots + 100(0.25)^{n-1}$

$= \sum_{i=1}^{n} 100(0.25)^{i-1}$ [geometric with $a = 100$ and $r = 0.25$]

The quantity of the antibiotic that remains in the body in the long run is $\lim_{n\to\infty} Q_n = \dfrac{100}{1-0.25} = \dfrac{100}{0.75} = 133.\overline{3}$ mg.

72. (a) The concentration of the drug after the first injection is 1.5 mg/L. "Reduced by 90%" is the same as 10% remains, so the concentration after the second injection is $1.5 + 0.10(1.5) = 1.65$ mg/L. The concentration after the third injection is $1.5 + 0.10(1.65)$, or, equivalently, $1.5 + 1.5(0.10) + 1.5(0.10)^2$. Either expression gives us 1.665 mg/L.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $C_n = 1.5 + 1.5(0.10)^1 + 1.5(0.10)^2 + \cdots + 1.5(0.10)^{n-1}$

$= \sum_{i=1}^{n} 1.5(0.10)^{i-1}$ [geometric with $a = 1.5$ and $r = 0.10$].

By (3), $C_n = \dfrac{1.5[1-(0.10)^n]}{1-0.10} = \dfrac{1.5}{0.9}[1-(0.10)^n] = \dfrac{5}{3}[1-(0.10)^n]$ mg/L.

(c) The limiting value of the concentration is $\lim_{n\to\infty} C_n = \lim_{n\to\infty} \frac{5}{3}[1-(0.10)^n] = \frac{5}{3}(1-0) = \frac{5}{3}$ mg/L.

73. (a) The quantity of the drug in the body after the first tablet is 150 mg. "Eliminates 95%" is the same as retains 5%, so after the second tablet, there is 150 mg plus 5% of the first 150- mg tablet, that is, $[150 + 150(0.05)]$ mg. After the third tablet, the quantity is $[150 + 150(0.05) + 150(0.05)^2] = 157.875$ mg. After n tablets, the quantity (in mg) is $150 + 150(0.05) + \cdots + 150(0.05)^{n-1}$. We can use Formula 3 to write this as $\dfrac{150(1-0.05^n)}{1-0.05} = \dfrac{3000}{19}(1-0.05^n)$.

(b) The number of milligrams remaining in the body in the long run is $\lim_{n\to\infty}\left[\frac{3000}{19}(1-0.05^n)\right] = \frac{3000}{19}(1-0) \approx 157.895$, only 0.02 mg more than the amount after 3 tablets.

74. (a) The residual concentration just before the second injection is De^{-aT}; before the third, $De^{-aT} + De^{-a2T}$; before the $(n+1)$st, $De^{-aT} + De^{-a2T} + \cdots + De^{-anT}$. This sum is equal to $\dfrac{De^{-aT}\left(1-e^{-anT}\right)}{1-e^{-aT}}$ [Formula 3].

(b) The limiting pre-injection concentration is $\lim_{n\to\infty} \dfrac{De^{-aT}\left(1-e^{-anT}\right)}{1-e^{-aT}} = \dfrac{De^{-aT}(1-0)}{1-e^{-aT}} \cdot \dfrac{e^{aT}}{e^{aT}} = \dfrac{D}{e^{aT}-1}$.

(c) $\dfrac{D}{e^{aT}-1} \geq C \quad \Rightarrow \quad D \geq C\left(e^{aT}-1\right)$, so the minimal dosage is $D = C\left(e^{aT}-1\right)$.

75. (a) The first step in the chain occurs when the local government spends D dollars. The people who receive it spend a fraction c of those D dollars, that is, Dc dollars. Those who receive the Dc dollars spend a fraction c of it, that is, Dc^2 dollars. Continuing in this way, we see that the total spending after n transactions is

$S_n = D + Dc + Dc^2 + \cdots + Dc^{n-1} = \dfrac{D(1-c^n)}{1-c}$ by (3).

(b) $\lim_{n\to\infty} S_n = \lim_{n\to\infty} \dfrac{D(1-c^n)}{1-c} = \dfrac{D}{1-c}\lim_{n\to\infty}(1-c^n) = \dfrac{D}{1-c}$ $\left[\text{since } 0 < c < 1 \quad \Rightarrow \quad \lim_{n\to\infty} c^n = 0\right]$

$= \dfrac{D}{s}$ [since $c + s = 1$] $= kD$ [since $k = 1/s$]

If $c = 0.8$, then $s = 1 - c = 0.2$ and the multiplier is $k = 1/s = 5$.

76. (a) Initially, the ball falls a distance H, then rebounds a distance rH, falls rH, rebounds r^2H, falls r^2H, etc. The total distance it travels is

$$H + 2rH + 2r^2H + 2r^3H + \cdots = H\left(1 + 2r + 2r^2 + 2r^3 + \cdots\right) = H\left[1 + 2r\left(1 + r + r^2 + \cdots\right)\right]$$
$$= H\left[1 + 2r\left(\frac{1}{1-r}\right)\right] = H\left(\frac{1+r}{1-r}\right) \text{ meters}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) We know that the ball falls $\frac{1}{2}gt^2$ meters in t seconds, where $g = 9.8\ \mathrm{m/s^2}$ is the gravitational acceleration. Thus, the ball falls h meters in $t = \sqrt{2h/g}$ seconds. The total travel time in seconds is

$$\begin{aligned}\sqrt{\frac{2H}{g}} + 2\sqrt{\frac{2H}{g}r} + 2\sqrt{\frac{2H}{g}r^2} + 2\sqrt{\frac{2H}{g}r^3} + \cdots &= \sqrt{\frac{2H}{g}}\left[1 + 2\sqrt{r} + 2\sqrt{r}^{\,2} + 2\sqrt{r}^{\,3} + \cdots\right] \\ &= \sqrt{\frac{2H}{g}}\left(1 + 2\sqrt{r}\left[1 + \sqrt{r} + \sqrt{r}^{\,2} + \cdots\right]\right) \\ &= \sqrt{\frac{2H}{g}}\left[1 + 2\sqrt{r}\left(\frac{1}{1-\sqrt{r}}\right)\right] = \sqrt{\frac{2H}{g}}\frac{1+\sqrt{r}}{1-\sqrt{r}}\end{aligned}$$

(c) It will help to make a chart of the time for each descent and each rebound of the ball, together with the velocity just before and just after each bounce. Recall that the time in seconds needed to fall h meters is $\sqrt{2h/g}$. The ball hits the ground with velocity $-g\sqrt{2h/g} = -\sqrt{2hg}$ (taking the upward direction to be positive) and rebounds with velocity $kg\sqrt{2h/g} = k\sqrt{2hg}$, taking time $k\sqrt{2h/g}$ to reach the top of its bounce, where its velocity is 0. At that point, its height is k^2h. All these results follow from the formulas for vertical motion with gravitational acceleration $-g$:

$$\frac{d^2y}{dt^2} = -g \quad\Rightarrow\quad v = \frac{dy}{dt} = v_0 - gt \quad\Rightarrow\quad y = y_0 + v_0t - \tfrac{1}{2}gt^2.$$

number of descent	time of descent	speed before bounce	speed after bounce	time of ascent	peak height
1	$\sqrt{2H/g}$	$\sqrt{2Hg}$	$k\sqrt{2Hg}$	$k\sqrt{2H/g}$	k^2H
2	$\sqrt{2k^2H/g}$	$\sqrt{2k^2Hg}$	$k\sqrt{2k^2Hg}$	$k\sqrt{2k^2H/g}$	k^4H
3	$\sqrt{2k^4H/g}$	$\sqrt{2k^4Hg}$	$k\sqrt{2k^4Hg}$	$k\sqrt{2k^4H/g}$	k^6H
$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$

The total travel time in seconds is

$$\begin{aligned}\sqrt{\frac{2H}{g}} + k\sqrt{\frac{2H}{g}} + k\sqrt{\frac{2H}{g}} + k^2\sqrt{\frac{2H}{g}} + k^2\sqrt{\frac{2H}{g}} + \cdots &= \sqrt{\frac{2H}{g}}\left(1 + 2k + 2k^2 + 2k^3 + \cdots\right) \\ &= \sqrt{\frac{2H}{g}}\left[1 + 2k(1 + k + k^2 + \cdots)\right] \\ &= \sqrt{\frac{2H}{g}}\left[1 + 2k\left(\frac{1}{1-k}\right)\right] = \sqrt{\frac{2H}{g}}\frac{1+k}{1-k}\end{aligned}$$

Another method: We could use part (b). At the top of the bounce, the height is $k^2h = rh$, so $\sqrt{r} = k$ and the result follows from part (b).

77. $\sum_{n=2}^{\infty}(1+c)^{-n}$ is a geometric series with $a = (1+c)^{-2}$ and $r = (1+c)^{-1}$, so the series converges when $\left|(1+c)^{-1}\right| < 1 \quad\Leftrightarrow\quad |1+c| > 1 \quad\Leftrightarrow\quad 1+c > 1$ or $1+c < -1 \quad\Leftrightarrow\quad c > 0$ or $c < -2$. We calculate the sum of the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

series and set it equal to 2: $\dfrac{(1+c)^{-2}}{1-(1+c)^{-1}} = 2 \quad \Leftrightarrow \quad \left(\dfrac{1}{1+c}\right)^2 = 2 - 2\left(\dfrac{1}{1+c}\right) \quad \Leftrightarrow \quad 1 = 2(1+c)^2 - 2(1+c) \quad \Leftrightarrow$

$2c^2 + 2c - 1 = 0 \quad \Leftrightarrow \quad c = \frac{-2 \pm \sqrt{12}}{4} = \frac{\pm\sqrt{3}-1}{2}$. However, the negative root is inadmissible because $-2 < \frac{-\sqrt{3}-1}{2} < 0$.

So $c = \frac{\sqrt{3}-1}{2}$.

78. $\sum\limits_{n=0}^{\infty} e^{nc} = \sum\limits_{n=0}^{\infty} (e^c)^n$ is a geometric series with $a = (e^c)^0 = 1$ and $r = e^c$. If $e^c < 1$, it has sum $\dfrac{1}{1-e^c}$, so $\dfrac{1}{1-e^c} = 10 \quad \Rightarrow$

$\frac{1}{10} = 1 - e^c \quad \Rightarrow \quad e^c = \frac{9}{10} \quad \Rightarrow \quad c = \ln \frac{9}{10}$.

79. Assume the harmonic series converges with sum S. Following the outlined method of proof, we have

$$S = \left(1 + \frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6}\right) + \cdots > \left(\frac{1}{2} + \frac{1}{2}\right) + \left(\frac{1}{4} + \frac{1}{4}\right) + \left(\frac{1}{6} + \frac{1}{6}\right) + \cdots = 1 + \frac{1}{2} + \frac{1}{3} + \cdots = S.$$

This indicates that $S > S \quad \Leftrightarrow \quad 0 > 0$, which is a contradiction. Thus, the assumption was incorrect and the harmonic series diverges.

80. From the hint, observe that $\dfrac{1}{n-1} + \dfrac{1}{n+1} = \dfrac{(n+1)+(n-1)}{(n-1)(n+1)} = \dfrac{2n}{n^2-1} > \dfrac{2n}{n^2} = \dfrac{2}{n}$ for $n \geq 3$. Now, assuming the harmonic series converges with sum S, we have

$$\begin{aligned} S &= 1 + \left(\frac{1}{2} + \frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7}\right) + \left(\frac{1}{8} + \frac{1}{9} + \frac{1}{10}\right) + \cdots \\ &= 1 + \left(\frac{1}{2} + \frac{1}{4} + \frac{1}{3}\right) + \left(\frac{1}{5} + \frac{1}{7} + \frac{1}{6}\right) + \left(\frac{1}{8} + \frac{1}{10} + \frac{1}{9}\right) + \cdots \\ &> 1 + \left(\frac{2}{3} + \frac{1}{3}\right) + \left(\frac{2}{6} + \frac{1}{6}\right) + \left(\frac{2}{9} + \frac{1}{9}\right) + \cdots = 1 + \left(1 + \frac{1}{2} + \frac{1}{3} + \cdots\right) = 1 + S. \end{aligned}$$

This indicates that $S > 1 + S \quad \Leftrightarrow \quad 0 > 1$, which is a contradiction. Thus, the assumption was incorrect and the harmonic series diverges.

81. From the hint, we'll show that $e^x > 1 + x$ for $x > 0$ by proving that $f(x) = e^x - (1+x) > 0$ for $x > 0$. Since $f(0) = e^0 - 1 = 0$ and $f'(x) = e^x - 1 > 0$ when $x > 0$, f is increasing on $(0, \infty) \quad \Rightarrow \quad$ when $x > 0$, $f(x) > 0 \quad \Rightarrow$ $e^x - (1+x) > 0$ for $x > 0 \quad \Rightarrow \quad e^x > 1 + x$ for $x > 0$. Now,

$$\begin{aligned} e^{1+(1/2)+(1/3)+\cdots+(1/n)} = e^1 e^{1/2} e^{1/3} \cdots e^{1/n} &> (1+1)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)\cdots\left(1+\frac{1}{n}\right) && \left[\text{since } e^x > 1+x\right] \\ &= \left(\frac{2}{1}\right)\left(\frac{3}{2}\right)\left(\frac{4}{3}\right)\cdots\left(\frac{n+1}{n}\right) && \left[\begin{array}{l}\text{each denominator cancels with the} \\ \text{numerator of the preceding fraction}\end{array}\right] \\ &= n+1 \end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Assuming the harmonic series converges with sum S, we take the limit as $n \to \infty$ to get

$\lim\limits_{n\to\infty} e^{1+(1/2)+(1/3)+\cdots+(1/n)} > \lim\limits_{n\to\infty}(n+1) \quad \Rightarrow$

$e^{\lim\limits_{n\to\infty}[1+(1/2)+(1/3)+\cdots+(1/n)]} > \lim\limits_{n\to\infty}(n+1) \quad \left[\text{since } e^x \text{ is continuous}\right] \quad \Rightarrow \quad e^S > \lim\limits_{n\to\infty}(n+1)$. The limit does not exist since $n+1 \to \infty$ as $n \to \infty$. e^S is larger than the value of the limit, so S is not finite valued, which is a contradiction. Thus, the assumption was incorrect and the harmonic series diverges.

82. The area between $y = x^{n-1}$ and $y = x^n$ for $0 \le x \le 1$ is

$$\int_0^1 (x^{n-1} - x^n)\,dx = \left[\frac{x^n}{n} - \frac{x^{n+1}}{n+1}\right]_0^1 = \frac{1}{n} - \frac{1}{n+1}$$
$$= \frac{(n+1)-n}{n(n+1)} = \frac{1}{n(n+1)}$$

We can see from the diagram that as $n \to \infty$, the sum of the areas between the successive curves approaches the area of the unit square, that is, 1. So $\sum\limits_{n=1}^{\infty} \dfrac{1}{n(n+1)} = 1$.

83. Let d_n be the diameter of C_n. We draw lines from the centers of the C_i to the center of D (or C), and using the Pythagorean Theorem, we can write

$1^2 + \left(1 - \frac{1}{2}d_1\right)^2 = \left(1 + \frac{1}{2}d_1\right)^2 \quad \Leftrightarrow$

$1 = \left(1 + \frac{1}{2}d_1\right)^2 - \left(1 - \frac{1}{2}d_1\right)^2 = 2d_1$ [difference of squares] $\quad \Rightarrow \quad d_1 = \frac{1}{2}$.

Similarly,

$1 = \left(1 + \frac{1}{2}d_2\right)^2 - \left(1 - d_1 - \frac{1}{2}d_2\right)^2 = 2d_2 + 2d_1 - d_1^2 - d_1 d_2$

$= (2 - d_1)(d_1 + d_2) \quad \Leftrightarrow$

$d_2 = \dfrac{1}{2-d_1} - d_1 = \dfrac{(1-d_1)^2}{2-d_1}$, $1 = \left(1 + \frac{1}{2}d_3\right)^2 - \left(1 - d_1 - d_2 - \frac{1}{2}d_3\right)^2 \quad \Leftrightarrow \quad d_3 = \dfrac{[1-(d_1+d_2)]^2}{2-(d_1+d_2)}$, and in general, $d_{n+1} = \dfrac{\left(1 - \sum_{i=1}^n d_i\right)^2}{2 - \sum_{i=1}^n d_i}$. If we actually calculate d_2 and d_3 from the formulas above, we find that they are $\dfrac{1}{6} = \dfrac{1}{2\cdot 3}$ and $\dfrac{1}{12} = \dfrac{1}{3\cdot 4}$ respectively, so we suspect that in general, $d_n = \dfrac{1}{n(n+1)}$. To prove this, we use induction: Assume that for all $k \le n$, $d_k = \dfrac{1}{k(k+1)} = \dfrac{1}{k} - \dfrac{1}{k+1}$. Then $\sum\limits_{i=1}^{n} d_i = 1 - \dfrac{1}{n+1} = \dfrac{n}{n+1}$ [telescoping sum]. Substituting this into our formula for d_{n+1}, we get $d_{n+1} = \dfrac{\left[1 - \dfrac{n}{n+1}\right]^2}{2 - \left(\dfrac{n}{n+1}\right)} = \dfrac{\dfrac{1}{(n+1)^2}}{\dfrac{n+2}{n+1}} = \dfrac{1}{(n+1)(n+2)}$, and the induction is complete.

Now, we observe that the partial sums $\sum_{i=1}^n d_i$ of the diameters of the circles approach 1 as $n \to \infty$; that is, $\sum\limits_{n=1}^{\infty} a_n = \sum\limits_{n=1}^{\infty} \dfrac{1}{n(n+1)} = 1$, which is what we wanted to prove.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

84. $|CD| = b\sin\theta$, $|DE| = |CD|\sin\theta = b\sin^2\theta$, $|EF| = |DE|\sin\theta = b\sin^3\theta, \ldots$. Therefore,

$|CD| + |DE| + |EF| + |FG| + \cdots = b\sum_{n=1}^{\infty}\sin^n\theta = b\left(\dfrac{\sin\theta}{1-\sin\theta}\right)$ since this is a geometric series with $r = \sin\theta$

and $|\sin\theta| < 1$ $\left[\text{because } 0 < \theta < \frac{\pi}{2}\right]$.

85. The series $1 - 1 + 1 - 1 + 1 - 1 + \cdots$ diverges (geometric series with $r = -1$), so we cannot say that

$0 = 1 - 1 + 1 - 1 + 1 - 1 + \cdots$.

86. If $\sum_{n=1}^{\infty} a_n$ is convergent, then $\lim_{n\to\infty} a_n = 0$ by Theorem 6, so $\lim_{n\to\infty}\dfrac{1}{a_n} \neq 0$, and so $\sum_{n=1}^{\infty}\dfrac{1}{a_n}$ is divergent by the Test for Divergence.

87. (a) $\sum_{n=1}^{\infty} ca_n = \lim_{n\to\infty}\sum_{i=1}^{n} ca_i = \lim_{n\to\infty} c\sum_{i=1}^{n} a_i = c\lim_{n\to\infty}\sum_{i=1}^{n} a_i = c\sum_{n=1}^{\infty} a_n$, which exists by hypothesis.

(b) $\sum_{n=1}^{\infty}(a_n - b_n) = \lim_{n\to\infty}\sum_{i=1}^{n}(a_i - b_i) = \lim_{n\to\infty}\left(\sum_{i=1}^{n} a_i - \sum_{i=1}^{n} b_i\right) = \lim_{n\to\infty}\sum_{i=1}^{n} a_i - \lim_{n\to\infty}\sum_{i=1}^{n} b_i$

$= \sum_{n=1}^{\infty} a_n - \sum_{n=1}^{\infty} b_n$, which exists by hypothesis.

88. If $\sum ca_n$ were convergent, then $\sum(1/c)(ca_n) = \sum a_n$ would be also, by Theorem 8(i). But this is not the case, so $\sum ca_n$ must diverge.

89. Suppose on the contrary that $\sum(a_n + b_n)$ converges. Then $\sum(a_n + b_n)$ and $\sum a_n$ are convergent series. So by Theorem 8(iii), $\sum[(a_n + b_n) - a_n]$ would also be convergent. But $\sum[(a_n + b_n) - a_n] = \sum b_n$, a contradiction, since $\sum b_n$ is given to be divergent.

90. No. For example, take $\sum a_n = \sum n$ and $\sum b_n = \sum(-n)$, which both diverge, yet $\sum(a_n + b_n) = \sum 0$, which converges with sum 0.

91. The partial sums $\{s_n\}$ form an increasing sequence, since $s_n - s_{n-1} = a_n > 0$ for all n. Also, the sequence $\{s_n\}$ is bounded since $s_n \leq 1000$ for all n. So by the Monotonic Sequence Theorem, the sequence of partial sums converges, that is, the series $\sum a_n$ is convergent.

92. (a) $\text{RHS} = \dfrac{1}{f_{n-1}f_n} - \dfrac{1}{f_n f_{n+1}} = \dfrac{f_n f_{n+1} - f_n f_{n-1}}{f_n^2 f_{n-1} f_{n+1}} = \dfrac{f_{n+1} - f_{n-1}}{f_n f_{n-1} f_{n+1}} = \dfrac{(f_{n-1} + f_n) - f_{n-1}}{f_n f_{n-1} f_{n+1}} = \dfrac{1}{f_{n-1}f_{n+1}} = \text{LHS}$

(b) $\sum_{n=2}^{\infty}\dfrac{1}{f_{n-1}f_{n+1}} = \sum_{n=2}^{\infty}\left(\dfrac{1}{f_{n-1}f_n} - \dfrac{1}{f_n f_{n+1}}\right)$ [from part (a)]

$= \lim_{n\to\infty}\left[\left(\dfrac{1}{f_1 f_2} - \dfrac{1}{f_2 f_3}\right) + \left(\dfrac{1}{f_2 f_3} - \dfrac{1}{f_3 f_4}\right) + \left(\dfrac{1}{f_3 f_4} - \dfrac{1}{f_4 f_5}\right) + \cdots + \left(\dfrac{1}{f_{n-1}f_n} - \dfrac{1}{f_n f_{n+1}}\right)\right]$

$= \lim_{n\to\infty}\left(\dfrac{1}{f_1 f_2} - \dfrac{1}{f_n f_{n+1}}\right) = \dfrac{1}{f_1 f_2} - 0 = \dfrac{1}{1\cdot 1} = 1$ because $f_n \to \infty$ as $n \to \infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\displaystyle\sum_{n=2}^{\infty} \frac{f_n}{f_{n-1}f_{n+1}} = \sum_{n=2}^{\infty}\left(\frac{f_n}{f_{n-1}f_n} - \frac{f_n}{f_n f_{n+1}}\right)$ [as above]

$$= \sum_{n=2}^{\infty}\left(\frac{1}{f_{n-1}} - \frac{1}{f_{n+1}}\right)$$

$$= \lim_{n\to\infty}\left[\left(\frac{1}{f_1} - \frac{1}{f_3}\right) + \left(\frac{1}{f_2} - \frac{1}{f_4}\right) + \left(\frac{1}{f_3} - \frac{1}{f_5}\right) + \left(\frac{1}{f_4} - \frac{1}{f_6}\right) + \cdots + \left(\frac{1}{f_{n-1}} - \frac{1}{f_{n+1}}\right)\right]$$

$$= \lim_{n\to\infty}\left(\frac{1}{f_1} + \frac{1}{f_2} - \frac{1}{f_n} - \frac{1}{f_{n+1}}\right) = 1 + 1 - 0 - 0 = 2 \quad \text{because } f_n \to \infty \text{ as } n \to \infty.$$

93. (a) At the first step, only the interval $\left(\frac{1}{3}, \frac{2}{3}\right)$ (length $\frac{1}{3}$) is removed. At the second step, we remove the intervals $\left(\frac{1}{9}, \frac{2}{9}\right)$ and $\left(\frac{7}{9}, \frac{8}{9}\right)$, which have a total length of $2 \cdot \left(\frac{1}{3}\right)^2$. At the third step, we remove 2^2 intervals, each of length $\left(\frac{1}{3}\right)^3$. In general, at the nth step we remove 2^{n-1} intervals, each of length $\left(\frac{1}{3}\right)^n$, for a length of $2^{n-1} \cdot \left(\frac{1}{3}\right)^n = \frac{1}{3}\left(\frac{2}{3}\right)^{n-1}$. Thus, the total length of all removed intervals is $\displaystyle\sum_{n=1}^{\infty} \frac{1}{3}\left(\frac{2}{3}\right)^{n-1} = \frac{1/3}{1-2/3} = 1$ $\left[\text{geometric series with } a = \frac{1}{3} \text{ and } r = \frac{2}{3}\right]$. Notice that at the nth step, the leftmost interval that is removed is $\left(\left(\frac{1}{3}\right)^n, \left(\frac{2}{3}\right)^n\right)$, so we never remove 0, and 0 is in the Cantor set. Also, the rightmost interval removed is $\left(1 - \left(\frac{2}{3}\right)^n, 1 - \left(\frac{1}{3}\right)^n\right)$, so 1 is never removed. Some other numbers in the Cantor set are $\frac{1}{3}$, $\frac{2}{3}$, $\frac{1}{9}$, $\frac{2}{9}$, $\frac{7}{9}$, and $\frac{8}{9}$.

(b) The area removed at the first step is $\frac{1}{9}$; at the second step, $8 \cdot \left(\frac{1}{9}\right)^2$; at the third step, $(8)^2 \cdot \left(\frac{1}{9}\right)^3$. In general, the area removed at the nth step is $(8)^{n-1}\left(\frac{1}{9}\right)^n = \frac{1}{9}\left(\frac{8}{9}\right)^{n-1}$, so the total area of all removed squares is

$$\sum_{n=1}^{\infty} \frac{1}{9}\left(\frac{8}{9}\right)^{n-1} = \frac{1/9}{1 - 8/9} = 1.$$

94. (a)

a_1	1	2	4	1	1	1000
a_2	2	3	1	4	1000	1
a_3	1.5	2.5	2.5	2.5	500.5	500.5
a_4	1.75	2.75	1.75	3.25	750.25	250.75
a_5	1.625	2.625	2.125	2.875	625.375	375.625
a_6	1.6875	2.6875	1.9375	3.0625	687.813	313.188
a_7	1.65625	2.65625	2.03125	2.96875	656.594	344.406
a_8	1.67188	2.67188	1.98438	3.01563	672.203	328.797
a_9	1.66406	2.66406	2.00781	2.99219	664.398	336.602
a_{10}	1.66797	2.66797	1.99609	3.00391	668.301	332.699
a_{11}	1.66602	2.66602	2.00195	2.99805	666.350	334.650
a_{12}	1.66699	2.66699	1.99902	3.00098	667.325	333.675

The limits seem to be $\frac{5}{3}$, $\frac{8}{3}$, 2, 3, 667, and 334. Note that the limits appear to be "weighted" more toward a_2. In general, we guess that the limit is $\dfrac{a_1 + 2a_2}{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $a_{n+1} - a_n = \frac{1}{2}(a_n + a_{n-1}) - a_n = -\frac{1}{2}(a_n - a_{n-1}) = -\frac{1}{2}\left[\frac{1}{2}(a_{n-1} + a_{n-2}) - a_{n-1}\right]$

$$= -\tfrac{1}{2}\left[-\tfrac{1}{2}(a_{n-1} - a_{n-2})\right] = \cdots = \left(-\tfrac{1}{2}\right)^{n-1}(a_2 - a_1)$$

Note that we have used the formula $a_k = \frac{1}{2}(a_{k-1} + a_{k-2})$ a total of $n-1$ times in this calculation, once for each k between 3 and $n+1$. Now we can write

$$a_n = a_1 + (a_2 - a_1) + (a_3 - a_2) + \cdots + (a_{n-1} - a_{n-2}) + (a_n - a_{n-1})$$
$$= a_1 + \sum_{k=1}^{n-1}(a_{k+1} - a_k) = a_1 + \sum_{k=1}^{n-1}\left(-\tfrac{1}{2}\right)^{k-1}(a_2 - a_1)$$

and so

$$\lim_{n\to\infty} a_n = a_1 + (a_2 - a_1)\sum_{k=1}^{\infty}\left(-\tfrac{1}{2}\right)^{k-1} = a_1 + (a_2 - a_1)\left[\frac{1}{1 - (-1/2)}\right] = a_1 + \tfrac{2}{3}(a_2 - a_1) = \frac{a_1 + 2a_2}{3}.$$

95. (a) For $\sum_{n=1}^{\infty} \frac{n}{(n+1)!}$, $s_1 = \frac{1}{1\cdot 2} = \frac{1}{2}$, $s_2 = \frac{1}{2} + \frac{2}{1\cdot 2\cdot 3} = \frac{5}{6}$, $s_3 = \frac{5}{6} + \frac{3}{1\cdot 2\cdot 3\cdot 4} = \frac{23}{24}$,

$s_4 = \frac{23}{24} + \frac{4}{1\cdot 2\cdot 3\cdot 4\cdot 5} = \frac{119}{120}$. The denominators are $(n+1)!$, so a guess would be $s_n = \frac{(n+1)! - 1}{(n+1)!}$.

(b) For $n = 1$, $s_1 = \frac{1}{2} = \frac{2! - 1}{2!}$, so the formula holds for $n = 1$. Assume $s_k = \frac{(k+1)! - 1}{(k+1)!}$. Then

$$s_{k+1} = \frac{(k+1)! - 1}{(k+1)!} + \frac{k+1}{(k+2)!} = \frac{(k+1)! - 1}{(k+1)!} + \frac{k+1}{(k+1)!(k+2)} = \frac{(k+2)! - (k+2) + k + 1}{(k+2)!}$$
$$= \frac{(k+2)! - 1}{(k+2)!}$$

Thus, the formula is true for $n = k+1$. So by induction, the guess is correct.

(c) $\lim_{n\to\infty} s_n = \lim_{n\to\infty} \frac{(n+1)! - 1}{(n+1)!} = \lim_{n\to\infty}\left[1 - \frac{1}{(n+1)!}\right] = 1$ and so $\sum_{n=1}^{\infty} \frac{n}{(n+1)!} = 1$.

96.

Let r_1 = radius of the large circle, r_2 = radius of next circle, and so on. From the figure we have $\angle BAC = 60°$ and $\cos 60° = r_1/|AB|$, so $|AB| = 2r_1$ and $|DB| = 2r_2$. Therefore, $2r_1 = r_1 + r_2 + 2r_2 \Rightarrow r_1 = 3r_2$. In general, we have $r_{n+1} = \frac{1}{3}r_n$, so the total area is

$$A = \pi r_1^2 + 3\pi r_2^2 + 3\pi r_3^2 + \cdots = \pi r_1^2 + 3\pi r_2^2\left(1 + \frac{1}{3^2} + \frac{1}{3^4} + \frac{1}{3^6} + \cdots\right)$$
$$= \pi r_1^2 + 3\pi r_2^2 \cdot \frac{1}{1 - 1/9} = \pi r_1^2 + \tfrac{27}{8}\pi r_2^2$$

Since the sides of the triangle have length 1, $|BC| = \frac{1}{2}$ and $\tan 30° = \frac{r_1}{1/2}$. Thus, $r_1 = \frac{\tan 30°}{2} = \frac{1}{2\sqrt{3}} \Rightarrow r_2 = \frac{1}{6\sqrt{3}}$,

so $A = \pi\left(\frac{1}{2\sqrt{3}}\right)^2 + \frac{27\pi}{8}\left(\frac{1}{6\sqrt{3}}\right)^2 = \frac{\pi}{12} + \frac{\pi}{32} = \frac{11\pi}{96}$. The area of the triangle is $\frac{\sqrt{3}}{4}$, so the circles occupy about 83.1% of the area of the triangle.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11.3 The Integral Test and Estimates of Sums

1. The picture shows that $a_2 = \dfrac{1}{2^{1.5}} < \displaystyle\int_1^2 \frac{1}{x^{1.5}}\,dx$,

$a_3 = \dfrac{1}{3^{1.5}} < \displaystyle\int_2^3 \frac{1}{x^{1.5}}\,dx$, and so on, so $\displaystyle\sum_{n=2}^{\infty} \frac{1}{n^{1.5}} < \int_1^{\infty} \frac{1}{x^{1.5}}\,dx$.

The integral converges by (7.8.2) with $p = 1.5 > 1$, so the series converges.

2. From the first figure, we see that $\int_1^6 f(x)\,dx < \sum_{i=1}^{5} a_i$. From the second figure, we see that $\sum_{i=2}^{6} a_i < \int_1^6 f(x)\,dx$. Thus, we have $\sum_{i=2}^{6} a_i < \int_1^6 f(x)\,dx < \sum_{i=1}^{5} a_i$.

3. The function $f(x) = x^{-3}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^{\infty} x^{-3}\,dx = \lim_{t\to\infty} \int_1^t x^{-3}\,dx = \lim_{t\to\infty} \left[\frac{x^{-2}}{-2}\right]_1^t = \lim_{t\to\infty} \left(-\frac{1}{2t^2} + \frac{1}{2}\right) = \frac{1}{2}.$$

Since this improper integral is convergent, the series $\displaystyle\sum_{n=1}^{\infty} n^{-3}$ is also convergent by the Integral Test.

4. The function $f(x) = x^{-0.3}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^{\infty} x^{-0.3}\,dx = \lim_{t\to\infty} \int_1^t x^{-0.3}\,dx = \lim_{t\to\infty} \left[\frac{x^{0.7}}{0.7}\right]_1^t = \lim_{t\to\infty} \left(\frac{t^{0.7}}{0.7} - \frac{1}{0.7}\right) = \infty.$$

Since this improper integral is divergent, the series $\displaystyle\sum_{n=1}^{\infty} n^{-0.3}$ is also divergent by the Integral Test.

5. The function $f(x) = \dfrac{2}{5x-1}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^{\infty} \frac{2}{5x-1}\,dx = \lim_{t\to\infty} \int_1^t \frac{2}{5x-1}\,dx = \lim_{t\to\infty} \left[\frac{2}{5}\ln(5x-1)\right]_1^t = \lim_{t\to\infty} \left[\frac{2}{5}\ln(5t-1) - \frac{2}{5}\ln 4\right] = \infty.$$

Since this improper integral is divergent, the series $\displaystyle\sum_{n=1}^{\infty} \frac{2}{5n-1}$ is also divergent by the Integral Test.

6. The function $f(x) = \dfrac{1}{(3x-1)^4}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^{\infty} \frac{1}{(3x-1)^4}\,dx = \lim_{t\to\infty} \int_1^t (3x-1)^{-4}\,dx = \lim_{t\to\infty} \left[\frac{1}{(-3)3}(3x-1)^{-3}\right]_1^t = \lim_{t\to\infty} \left[-\frac{1}{9(3t-1)^3} + \frac{1}{9\cdot 2^3}\right] = \frac{1}{72}.$$

Since this improper integral is convergent, the series $\displaystyle\sum_{n=1}^{\infty} \frac{1}{(3n-1)^4}$ is also convergent by the Integral Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. The function $f(x) = \dfrac{x^2}{x^3+1}$ is continuous, positive, and decreasing ($\star$) on $[2, \infty)$, so the Integral Test applies.

$$\int_2^\infty \frac{x^2}{x^3+1}\,dx = \lim_{t\to\infty}\int_2^t \frac{x^2}{x^3+1}\,dx = \lim_{t\to\infty}\left[\frac{1}{3}\ln\left|x^3+1\right|\right]_2^t = \lim_{t\to\infty}\left[\frac{1}{3}\ln\left(t^3+1\right) - \frac{1}{3}\ln 9\right] = \infty.$$

Since the improper integral is divergent, the series $\sum\limits_{n=2}^{\infty} \dfrac{n^2}{n^3+1}$ is also divergent by the Integral Test.

($\star$): $f'(x) = \dfrac{(x^3+1)(2x) - x^2(3x^2)}{(x^3+1)^2} = \dfrac{2x - x^4}{(x^3+1)^2} = -\dfrac{x(x^3-2)}{(x^3+1)^2} < 0$ for $x \geq 2$.

8. The function $f(x) = x^2 e^{-x^3}$ is continuous, positive, and decreasing ($\star$) on $[1, \infty)$, so the Integral Test applies.

$$\int_1^\infty x^2 e^{-x^3}\,dx = \lim_{t\to\infty}\int_1^t x^2 e^{-x^3}\,dx = \lim_{t\to\infty}\left[-\frac{1}{3}e^{-x^3}\right]_1^t = -\frac{1}{3}\lim_{t\to\infty}\left(e^{-t^3} - e^{-1}\right) = -\frac{1}{3}\left(0 - \frac{1}{e}\right) = \frac{1}{3e}.$$

Since this improper integral is convergent, the series $\sum\limits_{n=1}^{\infty} n^2 e^{-n^3}$ is also convergent by the Integral Test.

($\star$): $f'(x) = x^2 e^{-x^3}(-3x^2) + e^{-x^3}(2x) = xe^{-x^3}(-3x^3+2) = \dfrac{x(2-3x^3)}{e^{x^3}} < 0$ for $x > 1$

9. The function $f(x) = \dfrac{1}{x(\ln x)^3}$ is continuous, positive, and decreasing on $[2, \infty)$, so the Integral Test applies.

$$\int_2^\infty \frac{1}{x(\ln x)^3}\,dx = \lim_{t\to\infty}\int_2^t \frac{1}{x(\ln x)^3}\,dx = \lim_{t\to\infty}\int_{\ln 2}^{\ln t} \frac{du}{u^3} \quad \left[u = \ln x,\, du = \frac{dx}{x}\right] \quad = \lim_{t\to\infty}\left[-\frac{1}{2}u^{-2}\right]_{\ln 2}^{\ln t}$$
$$= \lim_{t\to\infty}\left[-\frac{1}{2(\ln t)^2} + \frac{1}{2(\ln 2)^2}\right] = 0 + \frac{1}{2(\ln 2)^2} = \frac{1}{2(\ln 2)^2}$$

Since the improper integral is convergent, the series $\sum\limits_{n=2}^{\infty} \dfrac{1}{n(\ln n)^3}$ is also convergent by the Integral Test.

10. The function $f(x) = \dfrac{\tan^{-1}x}{1+x^2}$ is continuous, positive, and decreasing ($\star$) on $[1, \infty)$, so the Integral Test applies.

$$\int_1^\infty \frac{\tan^{-1}x}{1+x^2}\,dx = \lim_{t\to\infty}\int_1^t \frac{\tan^{-1}x}{1+x^2}\,dx = \lim_{t\to\infty}\int_{\tan^{-1}1}^{\tan^{-1}t} u\,du \quad \left[u = \tan^{-1}x,\, du = \frac{dx}{1+x^2}\right]$$
$$= \lim_{t\to\infty}\left[\frac{1}{2}u^2\right]_{\tan^{-1}1}^{\tan^{-1}t} = \lim_{t\to\infty}\left[\frac{1}{2}(\tan^{-1}t)^2 - \frac{1}{2}(\tan^{-1}1)^2\right] = \frac{1}{2}\left(\frac{\pi}{2}\right)^2 - \frac{1}{2}\left(\frac{\pi}{4}\right)^2 = \frac{3\pi^2}{32}.$$

Since the improper integral is convergent, the series $\sum\limits_{n=1}^{\infty} \dfrac{\tan^{-1}n}{1+n^2}$ is also convergent by the Integral Test.

($\star$): $f'(x) = \dfrac{(1+x^2)\left(\dfrac{1}{1+x^2}\right) - (\tan^{-1}x)(2x)}{(1+x^2)^2} = \dfrac{1 - 2x\tan^{-1}x}{(1+x^2)^2} < 0$ for $2x\tan^{-1}x > 1$.

Both $2x$ and $\tan^{-1}x$ are increasing functions, and when $x = 1$, $2x\tan^{-1}x = 2(1)\tan^{-1}1 = 2\pi/4 > 1$, so it follows that $f'(x) < 0$ when $x \geq 1$.

11. $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^{\sqrt{2}}}$ is a p-series with $p = \sqrt{2} > 1$, so it converges by (1).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $\sum_{n=3}^{\infty} n^{-0.9999} = \sum_{n=3}^{\infty} \frac{1}{n^{0.9999}}$ is a p-series with $p = 0.9999 \le 1$, so it diverges by (1). The fact that the series begins with $n = 3$ is irrelevant when determining convergence.

13. $1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \frac{1}{125} + \cdots = \sum_{n=1}^{\infty} \frac{1}{n^3}$. This is a p-series with $p = 3 > 1$, so it converges by (1).

14. $\frac{1}{5} + \frac{1}{7} + \frac{1}{9} + \frac{1}{11} + \frac{1}{13} + \cdots = \sum_{n=1}^{\infty} \frac{1}{2n+3}$. The function $f(x) = \frac{1}{2x+3}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$\int_1^{\infty} \frac{1}{2x+3}\,dx = \lim_{t\to\infty} \int_1^t \frac{1}{2x+3}\,dx = \lim_{t\to\infty} \left[\frac{1}{2}\ln(2x+3)\right]_1^t = \lim_{t\to\infty} \left[\frac{1}{2}\ln(2t+3) - \frac{1}{2}\ln 5\right] = \infty$, so the series $\sum_{n=1}^{\infty} \frac{1}{2n+3}$ diverges.

15. $\frac{1}{3} + \frac{1}{7} + \frac{1}{11} + \frac{1}{15} + \frac{1}{19} + \cdots = \sum_{n=1}^{\infty} \frac{1}{4n-1}$. The function $f(x) = \frac{1}{4x-1}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$\int_1^{\infty} \frac{1}{4x-1}\,dx = \lim_{t\to\infty} \int_1^t \frac{1}{4x-1}\,dx = \lim_{t\to\infty} \left[\frac{1}{4}\ln(4x-1)\right]_1^t = \lim_{t\to\infty} \left[\frac{1}{4}\ln(4t-1) - \frac{1}{4}\ln 3\right] = \infty$, so the series $\sum_{n=1}^{\infty} \frac{1}{4n-1}$ diverges.

16. $1 + \frac{1}{2\sqrt{2}} + \frac{1}{3\sqrt{3}} + \frac{1}{4\sqrt{4}} + \frac{1}{5\sqrt{5}} + \cdots = \sum_{n=1}^{\infty} \frac{1}{n\sqrt{n}} = \sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$. This is a p-series with $p = \frac{3}{2} > 1$, so it converges by (1).

17. $\sum_{n=1}^{\infty} \frac{\sqrt{n}+4}{n^2} = \sum_{n=1}^{\infty} \left(\frac{\sqrt{n}}{n^2} + \frac{4}{n^2}\right) = \sum_{n=1}^{\infty} \frac{1}{n^{3/2}} + \sum_{n=1}^{\infty} \frac{4}{n^2}$. $\sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$ is a convergent p-series with $p = \frac{3}{2} > 1$. $\sum_{n=1}^{\infty} \frac{4}{n^2} = 4\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a constant multiple of a convergent p-series with $p = 2 > 1$, so it converges. The sum of two convergent series is convergent, so the original series is convergent.

18. The function $f(x) = \frac{\sqrt{x}}{1+x^{3/2}}$ is continuous and positive on $[1, \infty)$.

$f'(x) = \frac{(1+x^{3/2})\left(\frac{1}{2}x^{-1/2}\right) - x^{1/2}\left(\frac{3}{2}x^{1/2}\right)}{(1+x^{3/2})^2} = \frac{\frac{1}{2}x^{-1/2} + \frac{1}{2}x - \frac{3}{2}x}{(1+x^{3/2})^2} = \frac{1-2x^{3/2}}{2\sqrt{x}(1+x^{3/2})^2} < 0$ for $x \ge 1$, so f is decreasing on $[1, \infty)$, and the Integral Test applies.

$$\int_1^{\infty} \frac{\sqrt{x}}{1+x^{3/2}}\,dx = \lim_{t\to\infty} \int_1^t \frac{\sqrt{x}}{1+x^{3/2}}\,dx = \lim_{t\to\infty} \left[\tfrac{2}{3}\ln(1+x^{3/2})\right]_1^t \qquad \left[\begin{array}{c}\text{substitution}\\ \text{with } u = 1+x^{3/2}\end{array}\right]$$
$$= \lim_{t\to\infty} \left[\tfrac{2}{3}\ln(1+t^{3/2}) - \tfrac{2}{3}\ln 2\right] = \infty,$$

so the series $\sum_{n=1}^{\infty} \frac{\sqrt{n}}{1+n^{3/2}}$ diverges.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. The function $f(x) = \dfrac{1}{x^2+4}$ is continuous, positive, and decreasing on $[1, \infty)$, so we can apply the Integral Test.

$$\int_1^\infty \frac{1}{x^2+4}\,dx = \lim_{t\to\infty}\int_1^t \frac{1}{x^2+4}\,dx = \lim_{t\to\infty}\left[\frac{1}{2}\tan^{-1}\frac{x}{2}\right]_1^t = \frac{1}{2}\lim_{t\to\infty}\left[\tan^{-1}\left(\frac{t}{2}\right) - \tan^{-1}\left(\frac{1}{2}\right)\right]$$
$$= \frac{1}{2}\left[\frac{\pi}{2} - \tan^{-1}\left(\frac{1}{2}\right)\right]$$

Therefore, the series $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2+4}$ converges.

20. The function $f(x) = \dfrac{1}{x^2+2x+2}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^\infty \frac{1}{x^2+2x+2}\,dx = \lim_{t\to\infty}\int_1^t \frac{1}{(x+1)^2+1}\,dx = \lim_{t\to\infty}\Big[\arctan(x+1)\Big]_1^t$$
$$= \lim_{t\to\infty}[\arctan(t+1) - \arctan 2] = \frac{\pi}{2} - \arctan 2,$$

so the series $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2+2n+2}$ converges.

21. The function $f(x) = \dfrac{x^3}{x^4+4}$ is continuous and positive on $[2, \infty)$, and is also decreasing since

$f'(x) = \dfrac{(x^4+4)(3x^2) - x^3(4x^3)}{(x^4+4)^2} = \dfrac{12x^2 - x^6}{(x^4+4)^2} = \dfrac{x^2(12-x^4)}{(x^4+4)^2} < 0$ for $x > \sqrt[4]{12} \approx 1.86$, so we can use the Integral Test on $[2, \infty)$.

$\displaystyle\int_2^\infty \frac{x^3}{x^4+4}\,dx = \lim_{t\to\infty}\int_2^t \frac{x^3}{x^4+4}\,dx = \lim_{t\to\infty}\Big[\tfrac{1}{4}\ln(x^4+4)\Big]_2^t = \lim_{t\to\infty}\Big[\tfrac{1}{4}\ln(t^4+4) - \tfrac{1}{4}\ln 20\Big] = \infty$, so the series $\sum\limits_{n=2}^{\infty} \dfrac{n^3}{n^4+4}$ diverges, and it follows that $\sum\limits_{n=1}^{\infty} \dfrac{n^3}{n^4+4}$ diverges as well.

22. The function $f(x) = \dfrac{3x-4}{x^2-2x} = \dfrac{2}{x} + \dfrac{1}{x-2}$ [by partial fractions] is continuous, positive, and decreasing on $[3, \infty)$ since it is the sum of two such functions, so we can apply the Integral Test.

$$\int_3^\infty \frac{3x-4}{x^2-x}\,dx = \lim_{t\to\infty}\int_3^t \left[\frac{2}{x} + \frac{1}{x-2}\right]dx = \lim_{t\to\infty}\Big[2\ln x + \ln(x-2)\Big]_3^t = \lim_{t\to\infty}[2\ln t + \ln(t-2) - 2\ln 3] = \infty.$$

The integral is divergent, so the series $\sum\limits_{n=3}^{\infty} \dfrac{3n-4}{n^2-n}$ is divergent.

23. $f(x) = \dfrac{1}{x\ln x}$ is continuous and positive on $[2, \infty)$, and also decreasing since $f'(x) = -\dfrac{1+\ln x}{x^2(\ln x)^2} < 0$ for $x > 2$, so we can use the Integral Test. $\displaystyle\int_2^\infty \frac{1}{x\ln x}\,dx = \lim_{t\to\infty}[\ln(\ln x)]_2^t = \lim_{t\to\infty}[\ln(\ln t) - \ln(\ln 2)] = \infty$, so the series $\sum\limits_{n=2}^{\infty} \dfrac{1}{n\ln n}$ diverges.

24. The function $f(x) = \dfrac{\ln x}{x^2}$ is continuous and positive on $[2, \infty)$, and also decreasing since

$f'(x) = \dfrac{x^2(1/x) - (\ln x)(2x)}{(x^2)^2} = \dfrac{x - 2x\ln x}{x^4} = \dfrac{1 - 2\ln x}{x^3} < 0$ for $x > e^{1/2} \approx 1.65$, so we can use the Integral Test

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

on $[2, \infty)$.

$$\int_2^\infty \frac{\ln x}{x^2}\,dx = \lim_{t\to\infty}\int_2^t \frac{\ln x}{x^2}\,dx = \lim_{t\to\infty}\left(\left[-\frac{\ln x}{x}\right]_2^t + \int_2^t \frac{1}{x^2}\,dx\right) \qquad \left[\begin{array}{c}\text{by parts with}\\ u=\ln x,\ dv=(1/x^2)\,dx\end{array}\right]$$

$$= \lim_{t\to\infty}\left(-\frac{\ln t}{t} + \frac{\ln 2}{2} + \left[-\frac{1}{x}\right]_2^t\right) \overset{\text{H}}{=} \lim_{t\to\infty}\left(-\frac{1/t}{1} + \frac{\ln 2}{2} - \frac{1}{t} + \frac{1}{2}\right) = \frac{\ln 2 + 1}{2},$$

so the series $\sum\limits_{n=2}^{\infty} \dfrac{\ln n}{n^2}$ converges.

25. The function $f(x) = xe^{-x} = \dfrac{x}{e^x}$ is continuous and positive on $[1, \infty)$, and also decreasing since

$f'(x) = \dfrac{e^x \cdot 1 - xe^x}{(e^x)^2} = \dfrac{e^x(1-x)}{(e^x)^2} = \dfrac{1-x}{e^x} < 0$ for $x > 1$ [and $f(1) > f(2)$], so we can use the Integral Test on $[1, \infty)$.

$$\int_1^\infty xe^{-x}\,dx = \lim_{t\to\infty}\int_1^t xe^{-x}\,dx = \lim_{t\to\infty}\left(\left[-xe^{-x}\right]_1^t + \int_1^t e^{-x}\,dx\right) \qquad \left[\begin{array}{c}\text{by parts with}\\ u=x,\ dv=e^{-x}\,dx\end{array}\right]$$

$$= \lim_{t\to\infty}\left(-te^{-t} + e^{-1} + \left[-e^{-x}\right]_1^t\right) = \lim_{t\to\infty}\left(-\frac{t}{e^t} + \frac{1}{e} - \frac{1}{e^t} + \frac{1}{e}\right)$$

$$\overset{\text{H}}{=} \lim_{t\to\infty}\left(-\frac{1}{e^t} + \frac{1}{e} - 0 + \frac{1}{e}\right) = \frac{2}{e},$$

so the series $\sum\limits_{k=1}^{\infty} ke^{-k}$ converges.

26. The function $f(x) = xe^{-x^2} = \dfrac{x}{e^{x^2}}$ is continuous and positive on $[1, \infty)$, and also decreasing since

$f'(x) = \dfrac{e^{x^2} \cdot 1 - xe^{x^2} \cdot 2x}{(e^{x^2})^2} = \dfrac{1-2x^2}{e^{x^2}} < 0$ for $x > \sqrt{\frac{1}{2}} \approx 0.7$, so we can use the Integral Test on $[1, \infty)$.

$\displaystyle\int_1^\infty xe^{-x^2}\,dx = \lim_{t\to\infty}\int_1^t xe^{-x^2}\,dx = \lim_{t\to\infty}\left[-\tfrac{1}{2}e^{-x^2}\right]_1^t = \lim_{t\to\infty}\left(-\tfrac{1}{2}e^{-t^2} + \tfrac{1}{2}e^{-1}\right) = \frac{1}{2e}$, so the series $\sum\limits_{k=1}^{\infty} ke^{-k^2}$ converges.

27. The function $f(x) = \dfrac{1}{x^2+x^3} = \dfrac{1}{x^2} - \dfrac{1}{x} + \dfrac{1}{x+1}$ [by partial fractions] is continuous, positive and decreasing on $[1, \infty)$, so the Integral Test applies.

$$\int_1^\infty f(x)\,dx = \lim_{t\to\infty}\int_1^t \left(\frac{1}{x^2} - \frac{1}{x} + \frac{1}{x+1}\right)dx = \lim_{t\to\infty}\left[-\frac{1}{x} - \ln x + \ln(x+1)\right]_1^t$$

$$= \lim_{t\to\infty}\left[-\frac{1}{t} + \ln\frac{t+1}{t} + 1 - \ln 2\right] = 0 + 0 + 1 - \ln 2$$

The integral converges, so the series $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2+n^3}$ converges.

28. The function $f(x) = \dfrac{x}{x^4+1}$ is positive, continuous, and decreasing on $[1, \infty)$. [Note that

$f'(x) = \dfrac{x^4+1-4x^4}{(x^4+1)^2} = \dfrac{1-3x^4}{(x^4+1)^2} < 0$ on $[1, \infty)$.] Thus, we can apply the Integral Test.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\int_1^\infty \frac{x}{x^4+1}\,dx = \lim_{t\to\infty}\int_1^t \frac{\frac{1}{2}(2x)}{1+(x^2)^2}\,dx = \lim_{t\to\infty}\left[\frac{1}{2}\tan^{-1}(x^2)\right]_1^t = \frac{1}{2}\lim_{t\to\infty}[\tan^{-1}(t^2)-\tan^{-1}1] = \frac{1}{2}\left(\frac{\pi}{2}-\frac{\pi}{4}\right) = \frac{\pi}{8}$$

so the series $\sum\limits_{n=1}^{\infty}\frac{n}{n^4+1}$ converges.

29. The function $f(x) = \dfrac{\cos \pi x}{\sqrt{x}}$ is neither positive nor decreasing on $[1,\infty)$, so the hypotheses of the Integral Test are not satisfied for the series $\sum\limits_{n=1}^{\infty}\dfrac{\cos \pi n}{\sqrt{n}}$.

30. The function $f(x) = \dfrac{\cos^2 x}{1+x^2}$ is not decreasing on $[1,\infty)$, so the hypotheses of the Integral Test are not satisfied for the series $\sum\limits_{n=1}^{\infty}\dfrac{\cos^2 n}{1+n^2}$.

31. We have already shown (in Exercise 23) that when $p=1$ the series $\sum\limits_{n=2}^{\infty}\dfrac{1}{n(\ln n)^p}$ diverges, so assume that $p \neq 1$.

$f(x) = \dfrac{1}{x(\ln x)^p}$ is continuous and positive on $[2,\infty)$, and $f'(x) = -\dfrac{p+\ln x}{x^2(\ln x)^{p+1}} < 0$ if $x > e^{-p}$, so that f is eventually decreasing and we can use the Integral Test.

$$\int_2^\infty \frac{1}{x(\ln x)^p}\,dx = \lim_{t\to\infty}\left[\frac{(\ln x)^{1-p}}{1-p}\right]_2^t \quad [\text{for } p\neq 1] \;=\; \lim_{t\to\infty}\left[\frac{(\ln t)^{1-p}}{1-p} - \frac{(\ln 2)^{1-p}}{1-p}\right]$$

This limit exists whenever $1-p<0 \;\Leftrightarrow\; p>1$, so the series converges for $p>1$.

32. $f(x) = \dfrac{1}{x\ln x\,[\ln(\ln x)]^p}$ is positive and continuous on $[3,\infty)$. For $p \geq 0$, f clearly decreases on $[3,\infty)$; and for $p<0$, it can be verified that f is ultimately decreasing. Thus, we can apply the Integral Test.

$$I = \int_3^\infty \frac{dx}{x\ln x\,[\ln(\ln x)]^p} = \lim_{t\to\infty}\int_3^t \frac{[\ln(\ln x)]^{-p}}{x\ln x}\,dx = \lim_{t\to\infty}\left[\frac{[\ln(\ln x)]^{-p+1}}{-p+1}\right]_3^t \qquad [\text{for } p\neq 1]$$

$$= \lim_{t\to\infty}\left[\frac{[\ln(\ln t)]^{-p+1}}{-p+1} - \frac{[\ln(\ln 3)]^{-p+1}}{-p+1}\right],$$

which exists whenever $-p+1<0 \;\Leftrightarrow\; p>1$. If $p=1$, then $I = \lim\limits_{t\to\infty}\big[\ln(\ln(\ln x))\big]_3^t = \infty$. Therefore, $\sum\limits_{n=3}^{\infty}\dfrac{1}{n\ln n\,[\ln(\ln n)]^p}$ converges for $p>1$.

33. Clearly the series cannot converge if $p \geq -\frac{1}{2}$, because then $\lim\limits_{n\to\infty} n(1+n^2)^p \neq 0$. So assume $p<-\frac{1}{2}$. Then $f(x) = x(1+x^2)^p$ is continuous, positive, and eventually decreasing on $[1,\infty)$, and we can use the Integral Test.

$$\int_1^\infty x(1+x^2)^p dx = \lim_{t\to\infty}\left[\frac{1}{2}\cdot\frac{(1+x^2)^{p+1}}{p+1}\right]_1^t = \frac{1}{2(p+1)}\lim_{t\to\infty}[(1+t^2)^{p+1}-2^{p+1}].$$

This limit exists and is finite $\;\Leftrightarrow\; p+1<0 \;\Leftrightarrow\; p<-1$, so the series $\sum\limits_{n=1}^{\infty} n(1+n^2)^p$ converges whenever $p<-1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. If $p \le 0$, $\lim\limits_{n\to\infty} \dfrac{\ln n}{n^p} = \infty$ and the series diverges, so assume $p > 0$. $f(x) = \dfrac{\ln x}{x^p}$ is positive and continuous and $f'(x) < 0$ for $x > e^{1/p}$, so f is eventually decreasing and we can use the Integral Test. Integration by parts gives

$\displaystyle\int_1^\infty \frac{\ln x}{x^p}dx = \lim_{t\to\infty}\left[\frac{x^{1-p}\left[(1-p)\ln x - 1\right]}{(1-p)^2}\right]_1^t$ (for $p \ne 1$) $= \dfrac{1}{(1-p)^2}\left[\lim\limits_{t\to\infty} t^{1-p}\left[(1-p)\ln t - 1\right] + 1\right]$, which exists whenever $1 - p < 0 \;\Leftrightarrow\; p > 1$. Thus, $\sum\limits_{n=1}^{\infty} \dfrac{\ln n}{n^p}$ converges $\;\Leftrightarrow\; p > 1$.

35. Since this is a p-series with $p = x$, $\zeta(x)$ is defined when $x > 1$. Unless specified otherwise, the domain of a function f is the set of real numbers x such that the expression for $f(x)$ makes sense and defines a real number. So, in the case of a series, it's the set of real numbers x such that the series is convergent.

36. (a) $\displaystyle\sum_{n=2}^{\infty}\frac{1}{n^2} = \sum_{n=1}^{\infty}\frac{1}{n^2} - \frac{1}{1^2}$ [subtract a_1] $= \dfrac{\pi^2}{6} - 1$

(b) $\displaystyle\sum_{n=3}^{\infty}\frac{1}{(n+1)^2} = \sum_{n=4}^{\infty}\frac{1}{n^2} = \sum_{n=1}^{\infty}\frac{1}{n^2} - \left(\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2}\right) = \frac{\pi^2}{6} - \frac{49}{36}$

(c) $\displaystyle\sum_{n=1}^{\infty}\frac{1}{(2n)^2} = \sum_{n=1}^{\infty}\frac{1}{4n^2} = \frac{1}{4}\sum_{n=1}^{\infty}\frac{1}{n^2} = \frac{1}{4}\left(\frac{\pi^2}{6}\right) = \frac{\pi^2}{24}$

37. (a) $\displaystyle\sum_{n=1}^{\infty}\left(\frac{3}{n}\right)^4 = \sum_{n=1}^{\infty}\frac{81}{n^4} = 81\sum_{n=1}^{\infty}\frac{1}{n^4} = 81\left(\frac{\pi^4}{90}\right) = \frac{9\pi^4}{10}$

(b) $\displaystyle\sum_{k=5}^{\infty}\frac{1}{(k-2)^4} = \frac{1}{3^4} + \frac{1}{4^4} + \frac{1}{5^4} + \cdots = \sum_{k=3}^{\infty}\frac{1}{k^4} = \frac{\pi^4}{90} - \left(\frac{1}{1^4} + \frac{1}{2^4}\right)$ [subtract a_1 and a_2] $= \dfrac{\pi^4}{90} - \dfrac{17}{16}$

38. (a) $f(x) = 1/x^4$ is positive and continuous and $f'(x) = -4/x^5$ is negative for $x > 0$, and so the Integral Test applies.

$\displaystyle\sum_{n=1}^{\infty}\frac{1}{n^4} \approx s_{10} = \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \cdots + \frac{1}{10^4} \approx 1.082037.$

$R_{10} \le \displaystyle\int_{10}^{\infty}\frac{1}{x^4}\,dx = \lim_{t\to\infty}\left[\frac{1}{-3x^3}\right]_{10}^{t} = \lim_{t\to\infty}\left(-\frac{1}{3t^3} + \frac{1}{3\,(10)^3}\right) = \frac{1}{3000}$, so the error is at most $0.000\overline{3}$.

(b) $s_{10} + \displaystyle\int_{11}^{\infty}\frac{1}{x^4}\,dx \le s \le s_{10} + \int_{10}^{\infty}\frac{1}{x^4}\,dx \;\Rightarrow\; s_{10} + \frac{1}{3(11)^3} \le s \le s_{10} + \frac{1}{3(10)^3} \;\Rightarrow$

$1.082037 + 0.000250 = 1.082287 \le s \le 1.082037 + 0.000333 = 1.082370$, so we get $s \approx 1.08233$ with error ≤ 0.00005.

(c) The estimate in part (b) is $s \approx 1.08233$ with error ≤ 0.00005. The exact value given in Exercise 37 is $\pi^4/90 \approx 1.082323$. The difference is less than 0.00001.

(d) $R_n \le \displaystyle\int_n^{\infty}\frac{1}{x^4}\,dx = \frac{1}{3n^3}$. So $R_n < 0.00001 \;\Rightarrow\; \dfrac{1}{3n^3} < \dfrac{1}{10^5} \;\Rightarrow\; 3n^3 > 10^5 \;\Rightarrow\; n > \sqrt[3]{(10)^5/3} \approx 32.2$, that is, for $n > 32$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. (a) $f(x) = \dfrac{1}{x^2}$ is positive and continuous and $f'(x) = -\dfrac{2}{x^3}$ is negative for $x > 0$, and so the Integral Test applies.

$$\sum_{n=1}^{\infty} \frac{1}{n^2} \approx s_{10} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \cdots + \frac{1}{10^2} \approx 1.549768.$$

$$R_{10} \le \int_{10}^{\infty} \frac{1}{x^2}\,dx = \lim_{t\to\infty} \left[\frac{-1}{x}\right]_{10}^{t} = \lim_{t\to\infty}\left(-\frac{1}{t} + \frac{1}{10}\right) = \frac{1}{10},$$ so the error is at most 0.1.

(b) $s_{10} + \displaystyle\int_{11}^{\infty} \frac{1}{x^2}\,dx \le s \le s_{10} + \int_{10}^{\infty} \frac{1}{x^2}\,dx \quad \Rightarrow \quad s_{10} + \tfrac{1}{11} \le s \le s_{10} + \tfrac{1}{10} \quad \Rightarrow$

$1.549768 + 0.090909 = 1.640677 \le s \le 1.549768 + 0.1 = 1.649768$, so we get $s \approx 1.64522$ (the average of 1.640677 and 1.649768) with error ≤ 0.005 (the maximum of $1.649768 - 1.64522$ and $1.64522 - 1.640677$, rounded up).

(c) The estimate in part (b) is $s \approx 1.64522$ with error ≤ 0.005. The exact value given in Exercise 36 is $\pi^2/6 \approx 1.644934$. The difference is less than 0.0003.

(d) $R_n \le \displaystyle\int_{n}^{\infty} \frac{1}{x^2}\,dx = \frac{1}{n}$. So $R_n < 0.001$ if $\dfrac{1}{n} < \dfrac{1}{1000} \quad \Leftrightarrow \quad n > 1000$.

40. $\displaystyle\sum_{n=1}^{\infty} ne^{-2n}$. $f(x) = xe^{-2x}$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies. Using (2),

$$R_n \le \int_{n}^{\infty} xe^{-2x}\,dx = \lim_{t\to\infty}\left(\left[-\tfrac{1}{2}xe^{-2x}\right]_n^t + \int_n^t \tfrac{1}{2}e^{-2x}\,dx\right) \qquad \left[\begin{array}{c}\text{using parts with}\\ u = x,\ dv = e^{-2x}\,dx\end{array}\right]$$

$$= \lim_{t\to\infty}\left(\frac{-t}{2e^{2t}} + \frac{n}{2e^{2n}} - \frac{1}{4e^{2t}} + \frac{1}{4e^{2n}}\right) \overset{\text{H}}{=} 0 + \frac{n}{2e^{2n}} - 0 + \frac{1}{4e^{2n}} = \frac{2n+1}{4e^{2n}}$$

To be correct to four decimal places, we want $\dfrac{2n+1}{4e^{2n}} \le \dfrac{5}{10^5}$. This inequality is true for $n = 6$.

$$s_6 = \sum_{n=1}^{6} \frac{n}{e^{2n}} = \frac{1}{e^2} + \frac{2}{e^4} + \frac{3}{e^6} + \frac{4}{e^8} + \frac{5}{e^{10}} + \frac{6}{e^{12}} \approx 0.1810.$$

41. $\displaystyle\sum_{n=1}^{\infty} (2n+1)^{-6}$. $f(x) = 1/(2x+1)^6$ is continuous, positive, and decreasing on $[1, \infty)$, so the Integral Test applies.

Using (2), $R_n \le \displaystyle\int_{n}^{\infty} (2x+1)^{-6}\,dx = \lim_{t\to\infty}\left[\frac{-1}{10(2x+1)^5}\right]_n^t = \frac{1}{10(2n+1)^5}$. To be correct to five decimal places,

we want $\dfrac{1}{10(2n+1)^5} \le \dfrac{5}{10^6} \quad \Leftrightarrow \quad (2n+1)^5 \ge 20{,}000 \quad \Leftrightarrow \quad n \ge \tfrac{1}{2}\left(\sqrt[5]{20{,}000} - 1\right) \approx 3.12$, so use $n = 4$.

$$s_4 = \sum_{n=1}^{4} \frac{1}{(2n+1)^6} = \frac{1}{3^6} + \frac{1}{5^6} + \frac{1}{7^6} + \frac{1}{9^6} \approx 0.001\,446 \approx 0.00145.$$

42. $f(x) = \dfrac{1}{x(\ln x)^2}$ is positive and continuous and $f'(x) = -\dfrac{\ln x + 2}{x^2(\ln x)^3}$ is negative for $x > 1$, so the Integral Test applies.

Using (2), we need $0.01 > \displaystyle\int_{n}^{\infty} \frac{dx}{x(\ln x)^2} = \lim_{t\to\infty}\left[\frac{-1}{\ln x}\right]_n^t = \frac{1}{\ln n}$. This is true for $n > e^{100}$, so we would have to add this

many terms to find the sum of the series $\displaystyle\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^2}$ to within 0.01, which would be problematic because

$e^{100} \approx 2.7 \times 10^{43}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. $\sum\limits_{n=1}^{\infty} n^{-1.001} = \sum\limits_{n=1}^{\infty} \dfrac{1}{n^{1.001}}$ is a convergent p-series with $p = 1.001 > 1$. Using (2), we get

$R_n \le \displaystyle\int_n^{\infty} x^{-1.001}\,dx = \lim_{t\to\infty}\left[\frac{x^{-0.001}}{-0.001}\right]_n^t = -1000\lim_{t\to\infty}\left[\frac{1}{x^{0.001}}\right]_n^t = -1000\left(-\frac{1}{n^{0.001}}\right) = \frac{1000}{n^{0.001}}$. We want

$R_n < 0.000\,000\,005 \quad\Leftrightarrow\quad \dfrac{1000}{n^{0.001}} < 5\times 10^{-9} \quad\Leftrightarrow\quad n^{0.001} > \dfrac{1000}{5\times 10^{-9}} \quad\Leftrightarrow$

$n > \left(2\times 10^{11}\right)^{1000} = 2^{1000}\times 10^{11{,}000} \approx 1.07\times 10^{301}\times 10^{11{,}000} = 1.07\times 10^{11{,}301}$.

44. (a) $f(x) = \left(\dfrac{\ln x}{x}\right)^2$ is continuous and positive for $x > 1$, and since $f'(x) = \dfrac{2\ln x\,(1-\ln x)}{x^3} < 0$ for $x > e$, we can apply the Integral Test. Using a CAS, we get $\displaystyle\int_1^{\infty}\left(\frac{\ln x}{x}\right)^2 dx = 2$, so the series $\sum\limits_{n=1}^{\infty}\left(\dfrac{\ln n}{n}\right)^2$ also converges.

(b) Since the Integral Test applies, the error in $s \approx s_n$ is $R_n \le \displaystyle\int_n^{\infty}\left(\frac{\ln x}{x}\right)^2 dx = \frac{(\ln n)^2 + 2\ln n + 2}{n}$.

(c) By graphing the functions $y_1 = \dfrac{(\ln x)^2 + 2\ln x + 2}{x}$ and $y_2 = 0.05$, we see that $y_1 < y_2$ for $n \ge 1373$.

(d) Using the CAS to sum the first 1373 terms, we get $s_{1373} \approx 1.94$.

45. (a) From the figure, $a_2 + a_3 + \cdots + a_n \le \int_1^n f(x)\,dx$, so with

$f(x) = \dfrac{1}{x}$, $\dfrac{1}{2} + \dfrac{1}{3} + \dfrac{1}{4} + \cdots + \dfrac{1}{n} \le \displaystyle\int_1^n \frac{1}{x}\,dx = \ln n$.

Thus, $s_n = 1 + \dfrac{1}{2} + \dfrac{1}{3} + \dfrac{1}{4} + \cdots + \dfrac{1}{n} \le 1 + \ln n$.

(b) By part (a), $s_{10^6} \le 1 + \ln 10^6 \approx 14.82 < 15$ and

$s_{10^9} \le 1 + \ln 10^9 \approx 21.72 < 22$.

46. (a) The sum of the areas of the n rectangles in the graph to the right is

$1 + \dfrac{1}{2} + \dfrac{1}{3} + \cdots + \dfrac{1}{n}$. Now $\displaystyle\int_1^{n+1}\frac{dx}{x}$ is less than this sum because the rectangles extend above the curve $y = 1/x$, so

$\displaystyle\int_1^{n+1}\frac{1}{x}\,dx = \ln(n+1) < 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n}$, and since

$\ln n < \ln(n+1)$, $0 < 1 + \dfrac{1}{2} + \dfrac{1}{3} + \cdots + \dfrac{1}{n} - \ln n = t_n$.

(b) The area under $f(x) = 1/x$ between $x = n$ and $x = n+1$ is

$\displaystyle\int_n^{n+1}\frac{dx}{x} = \ln(n+1) - \ln n$, and this is clearly greater than the area of the inscribed rectangle in the figure to the right $\left[\text{which is } \dfrac{1}{n+1}\right]$, so

$t_n - t_{n+1} = [\ln(n+1) - \ln n] - \dfrac{1}{n+1} > 0$, and so $t_n > t_{n+1}$, so $\{t_n\}$ is a decreasing sequence.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) We have shown that $\{t_n\}$ is decreasing and that $t_n > 0$ for all n. Thus, $0 < t_n \le t_1 = 1$, so $\{t_n\}$ is a bounded monotonic sequence, and hence converges by the Monotonic Sequence Theorem.

47. $b^{\ln n} = \left(e^{\ln b}\right)^{\ln n} = \left(e^{\ln n}\right)^{\ln b} = n^{\ln b} = \dfrac{1}{n^{-\ln b}}$. $\sum\limits_{n=1}^{\infty} b^{\ln n}$ is a p-series, which converges for all b such that $-\ln b > 1 \quad \Leftrightarrow$ $\ln b < -1 \quad \Leftrightarrow \quad b < e^{-1} \quad \Leftrightarrow \quad b < 1/e$ [with $b > 0$].

48. For the series $\sum\limits_{n=1}^{\infty} \left(\dfrac{c}{n} - \dfrac{1}{n+1}\right)$,

$$s_n = \sum_{i=1}^{n} \left(\frac{c}{i} - \frac{1}{i+1}\right) = \left(\frac{c}{1} - \frac{1}{2}\right) + \left(\frac{c}{2} - \frac{1}{3}\right) + \left(\frac{c}{3} - \frac{1}{4}\right) + \cdots + \left(\frac{c}{n} - \frac{1}{n+1}\right)$$

$$= \frac{c}{1} + \frac{c-1}{2} + \frac{c-1}{3} + \frac{c-1}{4} + \cdots + \frac{c-1}{n} - \frac{1}{n+1} = c + (c-1)\left(\frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n}\right) - \frac{1}{n+1}$$

Thus, $\sum\limits_{n=1}^{\infty} \left(\dfrac{c}{n} - \dfrac{1}{n+1}\right) = \lim\limits_{n\to\infty} s_n = \lim\limits_{n\to\infty} \left[c + (c-1)\sum\limits_{i=2}^{n} \dfrac{1}{i} - \dfrac{1}{n+1}\right]$. Since a constant multiple of a divergent series is divergent, the last limit exists only if $c - 1 = 0$, so the original series converges only if $c = 1$.

11.4 The Comparison Tests

1. (a) We cannot say anything about $\sum a_n$. If $a_n > b_n$ for all n and $\sum b_n$ is convergent, then $\sum a_n$ could be convergent or divergent. (See the discusssion preceding the box titled "The Limit Comparison Test.")

(b) If $a_n < b_n$ for all n, then $\sum a_n$ is convergent. [This is part (i) of the Direct Comparison Test.]

2. (a) If $a_n > b_n$ for all n, then $\sum a_n$ is divergent. [This is part (ii) of the Direct Comparison Test.]

(b) We cannot say anything about $\sum a_n$. If $a_n < b_n$ for all n and $\sum b_n$ is divergent, then $\sum a_n$ could be convergent or divergent.

3. (a) $\dfrac{n}{n^3+5} < \dfrac{n}{n^3} = \dfrac{1}{n^2}$ for all $n \ge 2$. $\sum\limits_{n=2}^{\infty} \dfrac{1}{n^2}$ converges because it is a p-series with $p = 2 > 1$, so $\sum\limits_{n=2}^{\infty} \dfrac{n}{n^3+5}$ converges by part (i) of the Direct Comparison Test.

(b) Use the Limit Comparison Test with $a_n = \dfrac{n}{n^3-5}$ and $b_n = \dfrac{1}{n^2}$:

$$\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{n}{n^3-5} \cdot \frac{n^2}{1} = \lim_{n\to\infty} \frac{n^3}{n^3\left(1 - 5/n^3\right)} = \lim_{n\to\infty} \frac{1}{1 - 5/n^3} = \frac{1}{1-0} = 1 > 0$$

Since $\sum\limits_{n=2}^{\infty} \dfrac{1}{n^2}$ is a convergent (partial) p-series [$p = 2 > 1$], the series $\sum\limits_{n=2}^{\infty} \dfrac{n}{n^3-5}$ also converges.

4. (a) $\dfrac{n^2+n}{n^3-2} > \dfrac{n^2}{n^3-2} > \dfrac{n^2}{n^3} = \dfrac{1}{n}$ for all $n \ge 2$. $\sum\limits_{n=2}^{\infty} \dfrac{1}{n}$ diverges because it is a (partial) p-series with $p = 1 \le 1$, so $\sum\limits_{n=2}^{\infty} \dfrac{n^2+n}{n^3-2}$ diverges by part (ii) of the Direct Comparison Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Use the Limit Comparison Test with $a_n = \dfrac{n^2 - n}{n^3 + 2}$ and $b_n = \dfrac{1}{n}$:

$$\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{n^2 - n}{n^3 + 2} \cdot \frac{n}{1} = \lim_{n\to\infty} \frac{n^3 - n^2}{n^3 + 2} = \lim_{n\to\infty} \frac{1 - 1/n}{1 + 2/n^3} = \frac{1 - 0}{1 + 0} = 1 > 0$$

Since $\sum\limits_{n=2}^{\infty} \dfrac{1}{n}$ is a divergent (partial) p-series [$p = 1 \le 1$], the series $\sum\limits_{n=2}^{\infty} \dfrac{n^2 - n}{n^3 + 2}$ also diverges.

5. An inequality can be used to show that a series converges if its general term can be shown to be less than or equal to the general term of a known convergent series. The only inequality that satisfies this condition is given in part (c) since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$ is a convergent p-series [$p = 2 > 1$].

6. An inequality can be used to show that a series diverges if its general term can be shown to be greater than or equal to the general term of a known divergent series. The only inequality that satisfies this condition is given in part (c) since $\sum\limits_{n=1}^{\infty} \dfrac{1}{2n} = \dfrac{1}{2} \sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ is half of the harmonic series, which is divergent.

7. $\dfrac{1}{n^3 + 8} < \dfrac{1}{n^3}$ for all $n \ge 1$, so $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n^3 + 8}$ converges by direct comparison with $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n^3}$, which converges because it is a p-series with $p = 3 > 1$.

8. $\dfrac{1}{\sqrt{n} - 1} > \dfrac{1}{\sqrt{n}}$ for all $n \ge 2$, so $\displaystyle\sum_{n=2}^{\infty} \frac{1}{\sqrt{n} - 1}$ diverges by direct comparison with $\displaystyle\sum_{n=2}^{\infty} \frac{1}{\sqrt{n}}$, which diverges because it is a p-series with $p = \frac{1}{2} \le 1$.

9. $\dfrac{n + 1}{n\sqrt{n}} > \dfrac{n}{n\sqrt{n}} = \dfrac{1}{\sqrt{n}}$ for all $n \ge 1$, so $\displaystyle\sum_{n=1}^{\infty} \frac{n + 1}{n\sqrt{n}}$ diverges by direct comparison with $\displaystyle\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$, which diverges because it is a p-series with $p = \frac{1}{2} \le 1$.

10. $\dfrac{n - 1}{n^3 + 1} < \dfrac{n}{n^3 + 1} < \dfrac{n}{n^3} = \dfrac{1}{n^2}$ for all $n \ge 1$, so $\displaystyle\sum_{n=1}^{\infty} \frac{n - 1}{n^3 + 1}$ converges by direct comparison with $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n^2}$, which converges because it is a p-series with $p = 2 > 1$.

11. $\dfrac{9^n}{3 + 10^n} < \dfrac{9^n}{10^n} = \left(\dfrac{9}{10}\right)^n$ for all $n \ge 1$. $\sum\limits_{n=1}^{\infty} \left(\frac{9}{10}\right)^n$ is a convergent geometric series $\left(|r| = \frac{9}{10} < 1\right)$, so $\displaystyle\sum_{n=1}^{\infty} \frac{9^n}{3 + 10^n}$ converges by the Direct Comparison Test.

12. $\dfrac{6^n}{5^n - 1} > \dfrac{6^n}{5^n} = \left(\dfrac{6}{5}\right)^n$ for all $n \ge 1$. $\sum\limits_{n=1}^{\infty} \left(\frac{6}{5}\right)^n$ is a divergent geometric series $\left(|r| = \frac{6}{5} > 1\right)$, so $\displaystyle\sum_{n=1}^{\infty} \frac{6^n}{5^n - 1}$ diverges by the Direct Comparison Test.

13. For $n \ge 2$, $\ln n < n$, so $\dfrac{1}{\ln n} > \dfrac{1}{n}$. Thus, $\displaystyle\sum_{n=2}^{\infty} \frac{1}{\ln n}$ diverges by direct comparison with $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n}$, which diverges because it is a p-series with $p = 1 \le 1$ (the harmonic series).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $\dfrac{k\sin^2 k}{1+k^3} \le \dfrac{k}{1+k^3} < \dfrac{k}{k^3} = \dfrac{1}{k^2}$ for all $k \ge 1$, so $\sum\limits_{k=1}^{\infty} \dfrac{k\sin^2 k}{1+k^3}$ converges by direct comparison with $\sum\limits_{k=1}^{\infty} \dfrac{1}{k^2}$, which converges because it is a p-series with $p = 2 > 1$.

15. $\dfrac{\sqrt[3]{k}}{\sqrt{k^3+4k+3}} < \dfrac{\sqrt[3]{k}}{\sqrt{k^3}} = \dfrac{k^{1/3}}{k^{3/2}} = \dfrac{1}{k^{7/6}}$ for all $k \ge 1$, so $\sum\limits_{k=1}^{\infty} \dfrac{\sqrt[3]{k}}{\sqrt{k^3+4k+3}}$ converges by direct comparison with $\sum\limits_{k=1}^{\infty} \dfrac{1}{k^{7/6}}$, which converges because it is a p-series with $p = \frac{7}{6} > 1$.

16. $\dfrac{(2k-1)(k^2-1)}{(k+1)(k^2+4)^2} < \dfrac{2k(k^2)}{k(k^2)^2} = \dfrac{2k^3}{k^5} = \dfrac{2}{k^2}$ for all $k \ge 1$, so $\sum\limits_{k=1}^{\infty} \dfrac{(2k-1)(k^2-1)}{(k+1)(k^2+4)^2}$ converges by direct comparison with $2\sum\limits_{k=1}^{\infty} \dfrac{1}{k^2}$, which converges because it is a constant multiple of a p-series with $p = 2 > 1$.

17. $\dfrac{1+\cos n}{e^n} < \dfrac{2}{e^n}$ for all $n \ge 1$. $\displaystyle\sum_{n=1}^{\infty} \frac{2}{e^n}$ is a convergent geometric series ($|r| = \frac{1}{e} < 1$), so $\displaystyle\sum_{n=1}^{\infty} \frac{1+\cos n}{e^n}$ converges by the Direct Comparison Test.

18. $\dfrac{1}{\sqrt[3]{3n^4+1}} < \dfrac{1}{\sqrt[3]{3n^4}} < \dfrac{1}{\sqrt[3]{n^4}} = \dfrac{1}{n^{4/3}}$ for all $n \ge 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{1}{\sqrt[3]{3n^4+1}}$ converges by direct comparison with $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^{4/3}}$, which converges because it is a p-series with $p = \frac{4}{3} > 1$.

19. $\dfrac{4^{n+1}}{3^n-2} > \dfrac{4\cdot 4^n}{3^n} = 4\left(\dfrac{4}{3}\right)^n$ for all $n \ge 1$. $\sum\limits_{n=1}^{\infty} 4\left(\dfrac{4}{3}\right)^n = 4\sum\limits_{n=1}^{\infty}\left(\dfrac{4}{3}\right)^n$ is a divergent geometric series $\left(|r| = \frac{4}{3} > 1\right)$, so $\sum\limits_{n=1}^{\infty} \dfrac{4^{n+1}}{3^n-2}$ diverges by the Direct Comparison Test.

20. $\dfrac{1}{n^n} \le \dfrac{1}{n^2}$ for all $n \ge 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^n}$ converges by direct comparison with $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$, which converges because it is a p-series with $p = 2 > 1$.

21. Use the Limit Comparison Test with $a_n = \dfrac{1}{\sqrt{n^2+1}}$ and $b_n = \dfrac{1}{n}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{n}{\sqrt{n^2+1}} = \lim\limits_{n\to\infty} \dfrac{1}{\sqrt{1+(1/n^2)}} = 1 > 0$. Since the harmonic series $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ diverges, so does $\sum\limits_{n=1}^{\infty} \dfrac{1}{\sqrt{n^2+1}}$.

22. Use the Limit Comparison Test with $a_n = \dfrac{2}{\sqrt{n}+2}$ and $b_n = \dfrac{1}{\sqrt{n}}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{2\sqrt{n}}{\sqrt{n}+2} = \lim\limits_{n\to\infty} \dfrac{2}{1+2/\sqrt{n}} = 2 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{\sqrt{n}}$ is a divergent p-series [$p = \frac{1}{2} \le 1$], the series $\sum\limits_{n=1}^{\infty} \dfrac{2}{\sqrt{n}+2}$ is also divergent.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. Use the Limit Comparison Test with $a_n = \dfrac{n+1}{n^3+n}$ and $b_n = \dfrac{1}{n^2}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{(n+1)n^2}{n(n^2+1)} = \lim\limits_{n\to\infty} \dfrac{n^2+n}{n^2+1} = \lim\limits_{n\to\infty} \dfrac{1+1/n}{1+1/n^2} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$ is a convergent p-series $[p = 2 > 1]$, the series $\sum\limits_{n=1}^{\infty} \dfrac{n+1}{n^3+n}$ also converges.

24. Use the Limit Comparison Test with $a_n = \dfrac{n^2+n+1}{n^4+n^2}$ and $b_n = \dfrac{1}{n^2}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{(n^2+n+1)n^2}{n^2(n^2+1)} = \lim\limits_{n\to\infty} \dfrac{n^2+n+1}{n^2+1} = \lim\limits_{n\to\infty} \dfrac{1+1/n+1/n^2}{1+1/n^2} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$ is a convergent p-series $[p = 2 > 1]$, the series $\sum\limits_{n=1}^{\infty} \dfrac{n^2+n+1}{n^4+n^2}$ also converges.

25. Use the Limit Comparison Test with $a_n = \dfrac{\sqrt{1+n}}{2+n}$ and $b_n = \dfrac{1}{\sqrt{n}}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{\sqrt{1+n}\sqrt{n}}{2+n} = \lim\limits_{n\to\infty} \dfrac{\sqrt{n^2+n}/\sqrt{n^2}}{(2+n)/n} = \lim\limits_{n\to\infty} \dfrac{\sqrt{1+1/n}}{2/n+1} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{\sqrt{n}}$ is a divergent p-series $[\, p = \frac{1}{2} \le 1]$, the series $\sum\limits_{n=1}^{\infty} \dfrac{\sqrt{1+n}}{2+n}$ also diverges.

26. Use the Limit Comparison Test with $a_n = \dfrac{n+2}{(n+1)^3}$ and $b_n = \dfrac{1}{n^2}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{n^2(n+2)}{(n+1)^3} = \lim\limits_{n\to\infty} \dfrac{1+\frac{2}{n}}{\left(1+\frac{1}{n}\right)^3} = 1 > 0$. Since $\sum\limits_{n=3}^{\infty} \dfrac{1}{n^2}$ is a convergent (partial) p-series $[p = 2 > 1]$, the series $\sum\limits_{n=3}^{\infty} \dfrac{n+2}{(n+1)^3}$ also converges.

27. Use the Limit Comparison Test with $a_n = \dfrac{5+2n}{(1+n^2)^2}$ and $b_n = \dfrac{1}{n^3}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{n^3(5+2n)}{(1+n^2)^2} = \lim\limits_{n\to\infty} \dfrac{5n^3+2n^4}{(1+n^2)^2} \cdot \dfrac{1/n^4}{1/(n^2)^2} = \lim\limits_{n\to\infty} \dfrac{\frac{5}{n}+2}{\left(\frac{1}{n^2}+1\right)^2} = 2 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^3}$ is a convergent p-series $[p = 3 > 1]$, the series $\sum\limits_{n=1}^{\infty} \dfrac{5+2n}{(1+n^2)^2}$ also converges.

28. $\dfrac{n+3^n}{n+2^n} > \dfrac{3^n}{n+2^n} > \dfrac{3^n}{2^n+2^n} = \dfrac{3^n}{2\cdot 2^n} = \dfrac{1}{2}\left(\dfrac{3}{2}\right)^n$, so the series $\sum\limits_{n=1}^{\infty} \dfrac{n+3^n}{n+2^n}$ diverges by direct comparison with $\dfrac{1}{2}\sum\limits_{n=1}^{\infty} \left(\dfrac{3}{2}\right)^n$, which is a constant multiple of a divergent geometric series $[|r| = \frac{3}{2} > 1]$. *Or:* Use the Limit Comparison Test with $a_n = \dfrac{n+3^n}{n+2^n}$ and $b_n = \left(\dfrac{3}{2}\right)^n$.

29. $\dfrac{e^n+1}{ne^n+1} \ge \dfrac{e^n+1}{ne^n+n} = \dfrac{e^n+1}{n(e^n+1)} = \dfrac{1}{n}$ for $n \ge 1$, so the series $\sum\limits_{n=1}^{\infty} \dfrac{e^n+1}{ne^n+1}$ diverges by direct comparison with the divergent harmonic series $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$. *Or:* Use the Limit Comparison Test with $a_n = \dfrac{e^n+1}{ne^n+1}$ and $b_n = \dfrac{1}{n}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. If $a_n = \dfrac{1}{n\sqrt{n^2-1}}$ and $b_n = \dfrac{1}{n^2}$, then

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{n^2}{n\sqrt{n^2-1}} = \lim\limits_{n\to\infty} \dfrac{n/n}{\sqrt{n^2-1}/n} = \lim\limits_{n\to\infty} \dfrac{1}{\sqrt{1-1/n^2}} = \dfrac{1}{1} = 1 > 0$, so $\sum\limits_{n=2}^{\infty} \dfrac{1}{n\sqrt{n^2-1}}$ converges by the Limit Comparison Test with the convergent series $\sum\limits_{n=2}^{\infty} \dfrac{1}{n^2}$.

31. $\dfrac{2+\sin n}{n^2} \le \dfrac{2+1}{n^2} = \dfrac{3}{n^2}$, for all $n \ge 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{2+\sin n}{n^2}$ converges by direct comparison with $3\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$, which converges because it is a constant multiple of a p-series with $p = 2 > 1$.

32. $\dfrac{n^2+\cos^2 n}{n^3} \ge \dfrac{n^2}{n^3} = \dfrac{1}{n}$, for all $n \ge 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{n^2+\cos^2 n}{n^3}$ diverges by direct comparison with $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$, which diverges because it is a p-series with $p = 1 \le 1$ (the harmonic series).

33. Use the Limit Comparison Test with $a_n = \left(1+\dfrac{1}{n}\right)^2 e^{-n}$ and $b_n = e^{-n}$: $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \left(1+\dfrac{1}{n}\right)^2 = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} e^{-n} = \sum\limits_{n=1}^{\infty} \dfrac{1}{e^n}$ is a convergent geometric series $\left[|r| = \frac{1}{e} < 1\right]$, the series $\sum\limits_{n=1}^{\infty} \left(1+\dfrac{1}{n}\right)^2 e^{-n}$ also converges.

34. $\dfrac{e^{1/n}}{n} > \dfrac{1}{n}$ for all $n \ge 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{e^{1/n}}{n}$ diverges by direct comparison with the harmonic series $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$.

35. Clearly $n! = n(n-1)(n-2)\cdots(3)(2) \ge 2\cdot 2\cdot 2\cdot\cdots\cdot 2\cdot 2 = 2^{n-1}$, so $\dfrac{1}{n!} \le \dfrac{1}{2^{n-1}}$. $\sum\limits_{n=1}^{\infty} \dfrac{1}{2^{n-1}}$ is a convergent geometric series $\left[|r| = \frac{1}{2} < 1\right]$, so $\sum\limits_{n=1}^{\infty} \dfrac{1}{n!}$ converges by the Direct Comparison Test.

36. $\dfrac{n!}{n^n} = \dfrac{1\cdot 2\cdot 3\cdot\cdots\cdot(n-1)n}{n\cdot n\cdot n\cdot\cdots\cdot n\cdot n} \le \dfrac{1}{n}\cdot\dfrac{2}{n}\cdot 1\cdot 1\cdot\cdots\cdot 1$ for $n \ge 2$, so since $\sum\limits_{n=1}^{\infty} \dfrac{2}{n^2}$ converges $[p = 2 > 1]$, $\sum\limits_{n=1}^{\infty} \dfrac{n!}{n^n}$ converges also by the Direct Comparison Test.

37. Use the Limit Comparison Test with $a_n = \sin\left(\dfrac{1}{n}\right)$ and $b_n = \dfrac{1}{n}$. Then $\sum a_n$ and $\sum b_n$ are series with positive terms and

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{\sin(1/n)}{1/n} = \lim\limits_{\theta\to 0} \dfrac{\sin\theta}{\theta} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} b_n$ is the divergent harmonic series,

$\sum\limits_{n=1}^{\infty} \sin(1/n)$ also diverges. [Note that we could also use l'Hospital's Rule to evaluate the limit:

$\lim\limits_{x\to\infty} \dfrac{\sin(1/x)}{1/x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{\cos(1/x)\cdot\left(-1/x^2\right)}{-1/x^2} = \lim\limits_{x\to\infty} \cos\dfrac{1}{x} = \cos 0 = 1$.]

38. Use the Limit Comparison Test with $a_n = \sin^2\left(\dfrac{1}{n}\right)$ and $b_n = \dfrac{1}{n^2}$:

$$\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{\sin^2\left(\frac{1}{n}\right)}{\frac{1}{n^2}} = \lim_{n\to\infty} \left[\frac{\sin\left(\frac{1}{n}\right)}{\frac{1}{n}}\right]^2 = \lim_{x\to 0} \left(\frac{\sin x}{x}\right)^2 \qquad \left[\text{where } x = \frac{1}{n}\right]$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Now, $\lim\limits_{x\to 0}\dfrac{\sin x}{x}=1$ (see Equation 3.3.5) and the squaring function is continuous at $x=1$, so $\lim\limits_{x\to 0}\left(\dfrac{\sin x}{x}\right)^2=1^2=1>0$ [Theorem 2.5.8]. Since $\sum\limits_{n=1}^{\infty}\dfrac{1}{n^2}$ is a convergent p-series [$p=2>1$], the series $\sum\limits_{n=1}^{\infty}\sin^2\left(\dfrac{1}{n}\right)$ also converges.

39. Use the Limit Comparison Test with $a_n=\dfrac{1}{n}\tan\left(\dfrac{1}{n}\right)$ and $b_n=\dfrac{1}{n^2}$:

$$\lim_{n\to\infty}\frac{a_n}{b_n}=\lim_{n\to\infty}\frac{\dfrac{1}{n}\tan\left(\dfrac{1}{n}\right)}{\dfrac{1}{n^2}}=\lim_{n\to\infty}n\tan\left(\frac{1}{n}\right)=\lim_{x\to 0}\frac{\tan x}{x}\quad\left[\text{where } x=\frac{1}{n}\right]\quad\overset{\text{H}}{=}\lim_{x\to 0}\frac{\sec^2 x}{1}=\frac{1}{1}=1>0$$

Since $\sum\limits_{n=1}^{\infty}\dfrac{1}{n^2}$ is a convergent p-series [$p=2>1$], the series $\sum\limits_{n=1}^{\infty}\dfrac{1}{n}\tan\left(\dfrac{1}{n}\right)$ also converges.

40. Use the Limit Comparison Test with $a_n=\dfrac{1}{n^{1+1/n}}$ and $b_n=\dfrac{1}{n}$. $\lim\limits_{n\to\infty}\dfrac{a_n}{b_n}=\lim\limits_{n\to\infty}\dfrac{n}{n^{1+1/n}}=\lim\limits_{n\to\infty}\dfrac{1}{n^{1/n}}=1$ $\left[\text{since } \lim\limits_{x\to\infty}x^{1/x}=1 \text{ by l'Hospital's Rule}\right]$, so $\sum\limits_{n=1}^{\infty}\dfrac{1}{n}$ diverges [harmonic series] $\Rightarrow$ $\sum\limits_{n=1}^{\infty}\dfrac{1}{n^{1+1/n}}$ diverges.

41. $\sum\limits_{n=1}^{10}\dfrac{1}{5+n^5}=\dfrac{1}{5+1^5}+\dfrac{1}{5+2^5}+\dfrac{1}{5+3^5}+\cdots+\dfrac{1}{5+10^5}\approx 0.19926$. Now $\dfrac{1}{5+n^5}<\dfrac{1}{n^5}$, so the error is

$$R_{10}\le T_{10}\le\int_{10}^{\infty}\frac{1}{x^5}\,dx=\lim_{t\to\infty}\int_{10}^{t}x^{-5}\,dx=\lim_{t\to\infty}\left[\frac{-1}{4x^4}\right]_{10}^{t}=\lim_{t\to\infty}\left(\frac{-1}{4t^4}+\frac{1}{40{,}000}\right)=\frac{1}{40{,}000}=0.000\,025.$$

42. $\sum\limits_{n=1}^{10}\dfrac{e^{1/n}}{n^4}=\dfrac{e^{1/1}}{1^4}+\dfrac{e^{1/2}}{2^4}+\dfrac{e^{1/3}}{3^4}+\cdots+\dfrac{e^{1/10}}{10^4}\approx 2.84748$. Now $\dfrac{e^{1/n}}{n^4}\le\dfrac{e}{n^4}$ for $n\ge 1$, so the error is

$$R_{10}\le T_{10}\le\int_{10}^{\infty}\frac{e}{x^4}\,dx=\lim_{t\to\infty}\int_{10}^{t}ex^{-4}\,dx=\lim_{t\to\infty}\left[\frac{-e}{3x^3}\right]_{10}^{t}=\lim_{t\to\infty}\left(\frac{-e}{3t^3}+\frac{e}{3000}\right)=\frac{e}{3000}\approx 0.000\,906.$$

43. $\sum\limits_{n=1}^{10}5^{-n}\cos^2 n=\dfrac{\cos^2 1}{5}+\dfrac{\cos^2 2}{5^2}+\dfrac{\cos^2 3}{5^3}+\cdots+\dfrac{\cos^2 10}{5^{10}}\approx 0.07393$. Now $\dfrac{\cos^2 n}{5^n}\le\dfrac{1}{5^n}$, so the error is

$$R_{10}\le T_{10}\le\int_{10}^{\infty}\frac{1}{5^x}\,dx=\lim_{t\to\infty}\int_{10}^{t}5^{-x}\,dx=\lim_{t\to\infty}\left[-\frac{5^{-x}}{\ln 5}\right]_{10}^{t}=\lim_{t\to\infty}\left(-\frac{5^{-t}}{\ln 5}+\frac{5^{-10}}{\ln 5}\right)=\frac{1}{5^{10}\ln 5}<6.4\times 10^{-8}.$$

44. $\sum\limits_{n=1}^{10}\dfrac{1}{3^n+4^n}=\dfrac{1}{3^1+4^1}+\dfrac{1}{3^2+4^2}+\dfrac{1}{3^3+4^3}+\cdots+\dfrac{1}{3^{10}+4^{10}}\approx 0.19788$. Now $\dfrac{1}{3^n+4^n}<\dfrac{1}{3^n+3^n}=\dfrac{1}{2\cdot 3^n}$, so the error is

$$R_{10}\le T_{10}\le\int_{10}^{\infty}\frac{1}{2\cdot 3^x}\,dx=\lim_{t\to\infty}\int_{10}^{t}\frac{1}{2}\cdot 3^{-x}\,dx=\lim_{t\to\infty}\left[-\frac{1}{2}\frac{3^{-x}}{\ln 3}\right]_{10}^{t}=\lim_{t\to\infty}\left(-\frac{1}{2}\frac{3^{-t}}{\ln 3}+\frac{1}{2}\frac{3^{-10}}{\ln 3}\right)$$
$$=\frac{1}{2\cdot 3^{10}\ln 3}<7.7\times 10^{-6}.$$

45. Since $\dfrac{d_n}{10^n}\le\dfrac{9}{10^n}$ for each n, and since $\sum\limits_{n=1}^{\infty}\dfrac{9}{10^n}$ is a convergent geometric series $\left(|r|=\frac{1}{10}<1\right)$, $0.d_1d_2d_3\ldots=\sum\limits_{n=1}^{\infty}\dfrac{d_n}{10^n}$ will always converge by the Comparison Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. Clearly, if $p < 0$ then the series diverges, since $\lim\limits_{n\to\infty} \dfrac{1}{n^p \ln n} = \infty$. If $0 \le p \le 1$, then $n^p \ln n \le n \ln n \quad \Rightarrow$ $\dfrac{1}{n^p \ln n} \ge \dfrac{1}{n \ln n}$ and $\sum\limits_{n=2}^{\infty} \dfrac{1}{n \ln n}$ diverges (Exercise 11.3.23), so $\sum\limits_{n=2}^{\infty} \dfrac{1}{n^p \ln n}$ diverges. If $p > 1$, use the Limit Comparison Test with $a_n = \dfrac{1}{n^p \ln n}$ and $b_n = \dfrac{1}{n^p}$. $\sum\limits_{n=2}^{\infty} b_n$ converges, and $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{1}{\ln n} = 0$, so $\sum\limits_{n=2}^{\infty} \dfrac{1}{n^p \ln n}$ also converges. (Or use the Comparison Test, since $n^p \ln n > n^p$ for $n > e$.) In summary, the series converges if and only if $p > 1$.

47. Using Formula 1.5.10, we have $(\ln n)^{\ln \ln n} = e^{(\ln \ln n)(\ln \ln n)} \quad [n > 1] \quad = e^{(\ln \ln n)^2} < e^{\left(\sqrt{\ln n}\right)^2} = e^{\ln n} = n$. So $\dfrac{1}{(\ln n)^{\ln \ln n}} > \dfrac{1}{n}$ for all $n \ge 2$. Thus, the series $\sum\limits_{n=2}^{\infty} \dfrac{1}{(\ln n)^{\ln \ln n}}$ diverges by comparison with the partial harmonic series $\sum\limits_{n=2}^{\infty} \dfrac{1}{n}$, which diverges.

48. (a) Since $\lim\limits_{n\to\infty} (a_n/b_n) = 0$, there is a number $N > 0$ such that $|a_n/b_n - 0| < 1$ for all $n > N$, and so $a_n < b_n$ since a_n and b_n are positive. Thus, since $\sum b_n$ converges, so does $\sum a_n$ by the Direct Comparison Test.

(b) (i) If $a_n = \dfrac{\ln n}{n^3}$ and $b_n = \dfrac{1}{n^2}$, then $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{\ln n}{n} = \lim\limits_{x\to\infty} \dfrac{\ln x}{x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1/x}{1} = 0$. Now $\sum b_n$ is a convergent p-series $[p = 2 > 1]$, so $\sum\limits_{n=1}^{\infty} \dfrac{\ln n}{n^3}$ converges by part (a).

(ii) If $a_n = 1 - \cos\left(\dfrac{1}{n^2}\right)$ and $b_n = \dfrac{1}{n^2}$, then

$$\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{1 - \cos(1/n^2)}{1/n^2} = \lim_{x\to\infty} \frac{1 - \cos(1/x^2)}{1/x^2} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{\sin(1/x^2) \cdot (-2/x^3)}{-2/x^3} = \lim_{x\to\infty} \sin\left(\frac{1}{x^2}\right) = 0.$$

Now $\sum b_n$ is a convergent p-series $[p = 2 > 1]$, so $\sum\limits_{n=1}^{\infty} \left(1 - \cos \dfrac{1}{n^2}\right)$ converges by part (a).

49. (a) Since $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \infty$, there is an integer N such that $\dfrac{a_n}{b_n} > 1$ whenever $n > N$. (Take $M = 1$ in Definition 11.1.3.) Then $a_n > b_n$ whenever $n > N$ and since $\sum b_n$ is divergent, $\sum a_n$ is also divergent by the Comparison Test.

(b) (i) If $a_n = \dfrac{1}{\ln n}$ and $b_n = \dfrac{1}{n}$ for $n \ge 2$, then $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{n}{\ln n} = \lim\limits_{x\to\infty} \dfrac{x}{\ln x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1}{1/x} = \lim\limits_{x\to\infty} x = \infty$, so by part (a), $\sum\limits_{n=2}^{\infty} \dfrac{1}{\ln n}$ is divergent.

(ii) If $a_n = \dfrac{\ln n}{n}$ and $b_n = \dfrac{1}{n}$, then $\sum\limits_{n=1}^{\infty} b_n$ is the divergent harmonic series and $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \ln n = \lim\limits_{x\to\infty} \ln x = \infty$, so $\sum\limits_{n=1}^{\infty} a_n$ diverges by part (a).

50. Let $a_n = \dfrac{1}{n^2}$ and $b_n = \dfrac{1}{n}$. Then $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{1}{n} = 0$, but $\sum b_n$ diverges while $\sum a_n$ converges.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. $\lim\limits_{n\to\infty} na_n = \lim\limits_{n\to\infty} \dfrac{a_n}{1/n}$, so we apply the Limit Comparison Test with $b_n = \dfrac{1}{n}$. Since $\lim\limits_{n\to\infty} na_n > 0$ we know that either both series converge or both series diverge, and we also know that $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ diverges [p-series with $p = 1$]. Therefore, $\sum a_n$ must be divergent.

52. First we observe that, by l'Hospital's Rule, $\lim\limits_{x\to 0} \dfrac{\ln(1+x)}{x} = \lim\limits_{x\to 0} \dfrac{1}{1+x} = 1$. Also, if $\sum a_n$ converges, then $\lim\limits_{n\to\infty} a_n = 0$ by Theorem 11.2.6. Therefore, $\lim\limits_{n\to\infty} \dfrac{\ln(1+a_n)}{a_n} = \lim\limits_{x\to 0} \dfrac{\ln(1+x)}{x} = 1 > 0$. We are given that $\sum a_n$ is convergent and $a_n > 0$. Thus, $\sum \ln(1+a_n)$ is convergent by the Limit Comparison Test.

53. Yes. Since $\sum a_n$ is a convergent series with positive terms, $\lim\limits_{n\to\infty} a_n = 0$ by Theorem 11.2.6, and $\sum b_n = \sum \sin(a_n)$ is a series with positive terms (for large enough n). We have $\lim\limits_{n\to\infty} \dfrac{b_n}{a_n} = \lim\limits_{n\to\infty} \dfrac{\sin(a_n)}{a_n} = 1 > 0$ by Theorem 3.3.5. Thus, $\sum b_n$ is also convergent by the Limit Comparison Test.

54. Since $\sum a_n$ converges, $\lim\limits_{n\to\infty} a_n = 0$, so there exists N such that $|a_n - 0| < 1$ for all $n > N \quad \Rightarrow \quad 0 \le a_n < 1$ for all $n > N \quad \Rightarrow \quad 0 \le a_n^2 \le a_n$. Since $\sum a_n$ converges, so does $\sum a_n^2$ by the Comparison Test.

55. (a) False. The series $\sum a_n = \sum \dfrac{1}{n}$ and $\sum b_n = \sum \dfrac{1}{n}$ are both divergent, but $\sum a_n b_n = \sum \dfrac{1}{n} \cdot \dfrac{1}{n} = \sum \dfrac{1}{n^2}$ is convergent.

(b) False. The series $\sum a_n = \sum \dfrac{1}{n^2}$ converges and $\sum b_n = \sum \dfrac{1}{n}$ diverges, but $\sum a_n b_n = \sum \dfrac{1}{n^2} \cdot \dfrac{1}{n} = \sum \dfrac{1}{n^3}$ converges.

(c) True. Since $\sum a_n$ converges, for suffficiently large n, $a_n < 1 \quad \Rightarrow \quad a_n b_n < b_n$. Since $\sum b_n$ converges, it follows by the Direct Comparison Test that $\sum a_n b_n$ converges.

11.5 Alternating Series and Absolute Convergence

1. (a) An alternating series is a series whose terms are alternately positive and negative.

(b) An alternating series $\sum\limits_{n=1}^{\infty} a_n = \sum\limits_{n=1}^{\infty} (-1)^{n-1} b_n$, where $b_n = |a_n|$, converges if $0 < b_{n+1} \le b_n$ for all n and $\lim\limits_{n\to\infty} b_n = 0$. (This is the Alternating Series Test.)

(c) The error involved in using the partial sum s_n as an approximation to the total sum s is the remainder $R_n = s - s_n$ and the size of the error is smaller than b_{n+1}; that is, $|R_n| \le b_{n+1}$. (This is the Alternating Series Estimation Theorem.)

2. $\dfrac{2}{3} - \dfrac{2}{5} + \dfrac{2}{7} - \dfrac{2}{9} + \dfrac{2}{11} - \cdots = \sum\limits_{n=1}^{\infty} (-1)^{n+1} \dfrac{2}{2n+1}$. Now $b_n = \dfrac{2}{2n+1} > 0$, $\{b_n\}$ is decreasing, and $\lim\limits_{n\to\infty} b_n = 0$, so the series converges by the Alternating Series Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. $-\frac{2}{5}+\frac{4}{6}-\frac{6}{7}+\frac{8}{8}-\frac{10}{9}+\cdots=\sum_{n=1}^{\infty}(-1)^n\frac{2n}{n+4}$. Now $\lim_{n\to\infty} b_n=\lim_{n\to\infty}\frac{2n}{n+4}=\lim_{n\to\infty}\frac{2}{1+4/n}=\frac{2}{1}\neq 0$. Since $\lim_{n\to\infty} a_n\neq 0$ (in fact the limit does not exist), the series diverges by the Test for Divergence.

4. $\frac{1}{\ln 3}-\frac{1}{\ln 4}+\frac{1}{\ln 5}-\frac{1}{\ln 6}+\frac{1}{\ln 7}-\cdots=\sum_{n=1}^{\infty}\frac{(-1)^{n+1}}{\ln(n+2)}$. Now $b_n=\frac{1}{\ln(n+2)}>0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n=0$, so the series converges by the Alternating Series Test.

5. $\sum_{n=1}^{\infty} a_n=\sum_{n=1}^{\infty}\frac{(-1)^{n-1}}{3+5n}=\sum_{n=1}^{\infty}(-1)^{n-1}b_n$. Now $b_n=\frac{1}{3+5n}>0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n=0$, so the series converges by the Alternating Series Test.

6. $\sum_{n=0}^{\infty} a_n=\sum_{n=0}^{\infty}\frac{(-1)^{n+1}}{\sqrt{n+1}}=\sum_{n=0}^{\infty}(-1)^{n+1}b_n$. Now $b_n=\frac{1}{\sqrt{n+1}}>0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n=0$, so the series converges by the Alternating Series Test.

7. $\sum_{n=1}^{\infty} a_n=\sum_{n=1}^{\infty}(-1)^n\frac{3n-1}{2n+1}=\sum_{n=1}^{\infty}(-1)^n b_n$. Now $\lim_{n\to\infty} b_n=\lim_{n\to\infty}\frac{3-1/n}{2+1/n}=\frac{3}{2}\neq 0$. Since $\lim_{n\to\infty} a_n\neq 0$ (in fact the limit does not exist), the series diverges by the Test for Divergence.

8. $\sum_{n=1}^{\infty} a_n=\sum_{n=1}^{\infty}(-1)^n\frac{n^2}{n^2+n+1}=\sum_{n=1}^{\infty}(-1)^n b_n$. Now $\lim_{n\to\infty} b_n=\lim_{n\to\infty}\frac{n^2}{n^2+n+1}=\lim_{n\to\infty}\frac{1}{1+1/n+1/n^2}=1\neq 0$. Since $\lim_{n\to\infty} a_n\neq 0$, the series diverges by the Test for Divergence.

9. $\sum_{n=1}^{\infty} a_n=\sum_{n=1}^{\infty}(-1)^n e^{-n}=\sum_{n=1}^{\infty}(-1)^n b_n$. Now $b_n=\frac{1}{e^n}>0$, $\{b_n\}$ is decreasing, and $\lim_{n\to\infty} b_n=0$, so the series converges by the Alternating Series Test.

10. $b_n=\frac{\sqrt{n}}{2n+3}>0$ for $n\geq 1$. $\{b_n\}$ is decreasing for $n\geq 2$ since

$$\left(\frac{\sqrt{x}}{2x+3}\right)'=\frac{(2x+3)\left(\frac{1}{2}x^{-1/2}\right)-x^{1/2}(2)}{(2x+3)^2}=\frac{\frac{1}{2}x^{-1/2}[(2x+3)-4x]}{(2x+3)^2}=\frac{3-2x}{2\sqrt{x}\,(2x+3)^2}<0 \text{ for } x>\tfrac{3}{2}.$$

Also, $\lim_{n\to\infty} b_n=\lim_{n\to\infty}\frac{\sqrt{n}/\sqrt{n}}{(2n+3)/\sqrt{n}}=\lim_{n\to\infty}\frac{1}{2\sqrt{n}+3/\sqrt{n}}=0$. Thus, the series $\sum_{n=1}^{\infty}(-1)^n\frac{\sqrt{n}}{2n+3}$ converges by the Alternating Series Test.

11. $b_n=\frac{n^2}{n^3+4}>0$ for $n\geq 1$. $\{b_n\}$ is decreasing for $n\geq 2$ since

$$\left(\frac{x^2}{x^3+4}\right)'=\frac{(x^3+4)(2x)-x^2(3x^2)}{(x^3+4)^2}=\frac{x(2x^3+8-3x^3)}{(x^3+4)^2}=\frac{x(8-x^3)}{(x^3+4)^2}<0 \text{ for } x>2. \text{ Also,}$$

$\lim_{n\to\infty} b_n=\lim_{n\to\infty}\frac{1/n}{1+4/n^3}=0$. Thus, the series $\sum_{n=1}^{\infty}(-1)^{n+1}\frac{n^2}{n^3+4}$ converges by the Alternating Series Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $b_n = \dfrac{n}{2^n} > 0$ for $n \geq 1$. $\{b_n\}$ is decreasing for $n \geq 2$ since $\left(\dfrac{x}{2^x}\right)' = \dfrac{2^x - x\,2^x \ln 2}{(2^x)^2} = \dfrac{1 - x\ln 2}{2^x} < 0$ for $x > \dfrac{1}{\ln 2} \approx 1.4$.

Also, $\lim\limits_{n\to\infty} b_n = \lim\limits_{n\to\infty} \dfrac{n}{2^n} \overset{\text{H}}{=} \lim\limits_{n\to\infty} \dfrac{1}{2^n \ln 2} = 0$. Thus, the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n}{2^n}$ converges by the Alternating Series Test.

13. $\lim\limits_{n\to\infty} b_n = \lim\limits_{n\to\infty} e^{2/n} = e^0 = 1$, so $\lim\limits_{n\to\infty} (-1)^{n-1} e^{2/n}$ does not exist. Thus, the series $\sum\limits_{n=1}^{\infty} (-1)^{n-1} e^{2/n}$ diverges by the Test for Divergence.

14. $\lim\limits_{n\to\infty} b_n = \lim\limits_{n\to\infty} \arctan n = \frac{\pi}{2}$, so $\lim\limits_{n\to\infty} (-1)^{n-1} \arctan n$ does not exist. Thus, the series $\sum\limits_{n=1}^{\infty} (-1)^{n-1} \arctan n$ diverges by the Test for Divergence.

15. $a_n = \dfrac{\sin\left(n + \frac{1}{2}\right)\pi}{1 + \sqrt{n}} = \dfrac{(-1)^n}{1 + \sqrt{n}}$. Now $b_n = \dfrac{1}{1 + \sqrt{n}} > 0$ for $n \geq 0$, $\{b_n\}$ is decreasing, and $\lim\limits_{n\to\infty} b_n = 0$, so the series $\sum\limits_{n=0}^{\infty} \dfrac{\sin\left(n + \frac{1}{2}\right)\pi}{1 + \sqrt{n}}$ converges by the Alternating Series Test.

16. $a_n = \dfrac{n \cos n\pi}{2^n} = (-1)^n \dfrac{n}{2^n} = (-1)^n b_n$. $\{b_n\}$ is decreasing for $n \geq 2$ since

$(x2^{-x})' = x(-2^{-x} \ln 2) + 2^{-x} = 2^{-x}(1 - x\ln 2) < 0$ for $x > \dfrac{1}{\ln 2}$ [≈ 1.4]. Also, $\lim\limits_{n\to\infty} b_n = 0$ since

$\lim\limits_{x\to\infty} \dfrac{x}{2^x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1}{2^x \ln 2} = 0$. Thus, the series $\sum\limits_{n=1}^{\infty} \dfrac{n \cos n\pi}{2^n}$ converges by the Alternating Series Test.

17. $\sum\limits_{n=1}^{\infty} (-1)^n \sin\dfrac{\pi}{n}$. $b_n = \sin\dfrac{\pi}{n} > 0$ for $n \geq 2$ and $\sin\dfrac{\pi}{n} \geq \sin\dfrac{\pi}{n+1}$, and $\lim\limits_{n\to\infty} \sin\dfrac{\pi}{n} = \sin 0 = 0$, so the series converges by the Alternating Series Test.

18. $\sum\limits_{n=1}^{\infty} (-1)^n \cos\dfrac{\pi}{n}$. $\lim\limits_{n\to\infty} \cos\dfrac{\pi}{n} = \cos(0) = 1$, so $\lim\limits_{n\to\infty} (-1)^n \cos\dfrac{\pi}{n}$ does not exist and the series diverges by the Test for Divergence.

19. $b_n = \dfrac{n^2}{5^n} > 0$ for $n \geq 1$. $\{b_n\}$ is decreasing for $n \geq 2$ since

$\left(\dfrac{x^2}{5^x}\right)' = \dfrac{5^x \cdot 2x - x^2\, 5^x \ln 5}{(5^x)^2} = \dfrac{x\, 5^x (2 - x\ln 5)}{(5^x)^2} = \dfrac{x(2 - x\ln 5)}{5^x} < 0$ for $x > \dfrac{2}{\ln 5} \approx 1.2$. Also,

$\lim\limits_{n\to\infty} b_n = \lim\limits_{n\to\infty} \dfrac{n^2}{5^n} \overset{\text{H}}{=} \lim\limits_{n\to\infty} \dfrac{2n}{5^n \ln 5} \overset{\text{H}}{=} \lim\limits_{n\to\infty} \dfrac{2}{5^n (\ln 5)^2} = 0$. Thus, the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n^2}{5^n}$ converges by the Alternating Series Test.

20. $b_n = \dfrac{\sqrt{n+1} - \sqrt{n}}{1} \cdot \dfrac{\sqrt{n+1} + \sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = \dfrac{(n+1) - n}{\sqrt{n+1} + \sqrt{n}} = \dfrac{1}{\sqrt{n+1} + \sqrt{n}} > 0$ for $n \geq 1$. $\{b_n\}$ is decreasing and

$\lim\limits_{n\to\infty} b_n = 0$, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \left(\sqrt{n+1} - \sqrt{n}\right)$ converges by the Alternating Series Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. (a) A series $\sum a_n$ is called absolutely convergent if the series of absolute values $\sum |a_n|$ is convergent. If a series is absolutely convergent, then it is convergent.

(b) A series $\sum a_n$ is called conditionally convergent if it is convergent but not absolutely convergent; that is, if $\sum a_n$ converges, but $\sum |a_n|$ diverges.

(c) Suppose the series of positive terms $\sum_{n=1}^{\infty} b_n$ converges. Then $\sum |(-1)^n b_n| = \sum |b_n| = \sum b_n$ also converges, so $\sum_{n=1}^{\infty} (-1)^n b_n$ is absolutely convergent (and therefore convergent).

22. $\sum\limits_{n=1}^{\infty} |a_n| = \sum\limits_{n=1}^{\infty} \left|\dfrac{(-1)^n}{n^4}\right| = \sum\limits_{n=1}^{\infty} \dfrac{1}{n^4}$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^4}$ is a convergent p-series [$p = 4 > 1$], the series $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^n}{n^4}$ is absolutely convergent.

23. $b_n = \dfrac{1}{\sqrt[3]{n^2}} > 0$ for $n \geq 1$. $\{b_n\}$ is decreasing for $n \geq 1$, and $\lim\limits_{n\to\infty} \dfrac{1}{\sqrt[3]{n^2}} = 0$, so the series $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^{n-1}}{\sqrt[3]{n^2}}$ converges by the Alternating Series Test. Also, observe that $\sum\limits_{n=1}^{\infty} \left|\dfrac{(-1)^{n-1}}{\sqrt[3]{n^2}}\right| = \sum\limits_{n=1}^{\infty} \dfrac{1}{n^{2/3}}$ is divergent since it is a p-series with $p = \frac{2}{3} \leq 1$. Thus, the series $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^{n-1}}{\sqrt[3]{n^2}}$ is conditionally convergent.

24. Since $\lim\limits_{n\to\infty} \dfrac{n^2}{n^2+1} = \lim\limits_{n\to\infty} \dfrac{1}{1+1/n^2} = \dfrac{1}{1+0} = 1 \neq 0$ and $\lim\limits_{n\to\infty} (-1)^{n+1}$ does not exist, $\lim\limits_{n\to\infty} (-1)^{n+1} \dfrac{n^2}{n^2+1}$ does not exist, so the series $\sum\limits_{n=0}^{\infty} (-1)^{n+1} \dfrac{n^2}{n^2+1}$ diverges by the Test for Divergence.

25. $b_n = \dfrac{1}{5n+1} > 0$ for $n \geq 1$, $\{b_n\}$ is decreasing for $n \geq 1$, and $\lim\limits_{n\to\infty} b_n = 0$, so $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^n}{5n+1}$ converges by the Alternating Series Test. To determine absolute convergence, choose $a_n = \dfrac{1}{n}$ to get

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{1/n}{1/(5n+1)} = \lim\limits_{n\to\infty} \dfrac{5n+1}{n} = 5 > 0$, so $\sum\limits_{n=1}^{\infty} \dfrac{1}{5n+1}$ diverges by the Limit Comparison Test with the harmonic series. Thus, the series $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^n}{5n+1}$ is conditionally convergent.

26. $\sum\limits_{n=1}^{\infty} \dfrac{-n}{n^2+1} = -\sum\limits_{n=1}^{\infty} \dfrac{n}{n^2+1}$. Use the Limit Comparison Test with $a_n = \dfrac{n}{n^2+1}$ and $b_n = \dfrac{1}{n}$:

$$\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{n/\left(n^2+1\right)}{1/n} = \lim_{n\to\infty} \frac{n^2}{n^2+1} = \lim_{n\to\infty} \frac{1}{1+1/n^2} = \frac{1}{1+0} = 1 > 0$$

Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ is a divergent p-series [$p = 1 \leq 1$], the series $\sum\limits_{n=1}^{\infty} \dfrac{n}{n^2+1}$ also diverges, and hence, the negative of this series, $\sum\limits_{n=1}^{\infty} \dfrac{-n}{n^2+1}$, diverges.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. $\sum_{n=1}^{\infty} |a_n| = \sum_{n=1}^{\infty} \left|\frac{(-1)^n}{n^2+1}\right| = \sum_{n=1}^{\infty} \frac{1}{n^2+1}$. Since $\frac{1}{n^2+1} < \frac{1}{n^2}$ and $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a convergent p-series $[p = 2 > 1]$, the series $\sum_{n=1}^{\infty} \frac{1}{n^2+1}$ is convergent by the Direct Comparison Test. Thus, $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2+1}$ is absolutely convergent.

28. $0 < \left|\frac{\sin n}{2^n}\right| < \frac{1}{2^n}$ for $n \geq 1$ and $\sum_{n=1}^{\infty} \frac{1}{2^n}$ is a convergent geometric series $[r = \frac{1}{2} < 1]$, so $\sum_{n=1}^{\infty} \left|\frac{\sin n}{2^n}\right|$ converges by direct comparison and the series $\sum_{n=1}^{\infty} \frac{\sin n}{2^n}$ is absolutely convergent.

29. $0 < \left|\frac{1+2\sin n}{n^3}\right| < \frac{3}{n^3}$ for $n \geq 1$ and $3\sum_{n=1}^{\infty} \frac{1}{n^3}$ is a constant times a convergent p-series $[p = 3 > 1]$, so $\sum_{n=1}^{\infty} \left|\frac{1+2\sin n}{n^3}\right|$ converges by direct comparison and the series $\sum_{n=1}^{\infty} \frac{1+2\sin n}{n^3}$ is absolutely convergent.

30. $b_n = \frac{n}{n^2+4} > 0$ for $n \geq 1$, $\{b_n\}$ is decreasing for $n \geq 2$, and $\lim_{n\to\infty} b_n = 0$, so $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{n}{n^2+4}$ converges by the Alternating Series Test. To determine absolute convergence, choose $a_n = \frac{1}{n}$ to get

$\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{1/n}{n/(n^2+4)} = \lim_{n\to\infty} \frac{n^2+4}{n^2} = \lim_{n\to\infty} \frac{1+4/n^2}{1} = 1 > 0$, so $\sum_{n=1}^{\infty} \frac{n}{n^2+4}$ diverges by the Limit Comparison Test with the harmonic series. Thus, the series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{n}{n^2+4}$ is conditionally convergent.

31. $\sum_{n=2}^{\infty} \frac{(-1)^n}{\ln n}$ converges by the Alternating Series Test since $\lim_{n\to\infty} \frac{1}{\ln n} = 0$ and $\left\{\frac{1}{\ln n}\right\}$ is decreasing. Now $\ln n < n$, so $\frac{1}{\ln n} > \frac{1}{n}$, and since $\sum_{n=2}^{\infty} \frac{1}{n}$ is the divergent (partial) harmonic series, $\sum_{n=2}^{\infty} \frac{1}{\ln n}$ diverges by the Direct Comparison Test. Thus, $\sum_{n=2}^{\infty} \frac{(-1)^n}{\ln n}$ is conditionally convergent.

32. $b_n = \frac{n}{\sqrt{n^3+2}} > 0$ for $n \geq 1$, $\{b_n\}$ is decreasing for $n \geq 2$, and $\lim_{n\to\infty} b_n = 0$, so $\sum_{n=1}^{\infty} (-1)^n \frac{n}{\sqrt{n^3+2}}$ converges by the Alternating Series Test. To determine absolute convergence, choose $a_n = \frac{1}{\sqrt{n}}$ to get

$\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \left(\frac{1}{\sqrt{n}} \cdot \frac{\sqrt{n^3+2}}{n}\right) = \lim_{n\to\infty} \frac{\sqrt{n^3+2}}{\sqrt{n^3}} = \lim_{n\to\infty} \sqrt{1+\frac{2}{n^3}} = 1 > 0$, so $\sum_{n=1}^{\infty} \frac{n}{\sqrt{n^3+2}}$ diverges by limit comparision with the divergent p-series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ $\left[p = \frac{1}{2} \leq 1\right]$. Thus, $\sum_{n=1}^{\infty} (-1)^n \frac{n}{\sqrt{n^3+2}}$ is conditionally convergent.

33. $a_n = \frac{\cos n\pi}{3n+2} = (-1)^n \frac{1}{3n+2} = (-1)^n b_n$. $\{b_n\}$ is decreasing for $n \geq 1$, and $\lim_{n\to\infty} b_n = 0$, so $\sum_{n=1}^{\infty} \frac{\cos n\pi}{3n+2}$ converges by the Alternating Series Test. To determine absolute convergence, use the Limit Comparison Test with $a_n = \frac{1}{n}$:

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{1/n}{1/(3n+2)} = \lim\limits_{n\to\infty} \dfrac{3n+2}{n} = \lim\limits_{n\to\infty} \dfrac{3+2/n}{1} = 3 > 0$. Since the harmonic series diverges, so does $\sum\limits_{n=1}^{\infty} \dfrac{1}{3n+2}$. Thus, the series $\sum\limits_{n=1}^{\infty} \dfrac{\cos n\pi}{3n+2}$ is conditionally convergent.

34. The function $f(x) = \dfrac{1}{x \ln x}$ is continuous, positive, and decreasing on $[2, \infty)$.

$\displaystyle\int_2^{\infty} \frac{1}{x \ln x}\, dx = \lim_{t\to\infty} \int_2^t \frac{1}{x \ln x}\, dx = \lim_{t\to\infty} [\ln(\ln x)]_2^t = \lim_{t\to\infty} [(\ln(\ln t) - \ln(\ln 2)] = \infty$, so the series $\sum\limits_{n=2}^{\infty} \dfrac{(-1)^n}{n \ln n}$ diverges by the Integral Test. Now $\{b_n\} = \left\{ \dfrac{1}{n \ln n} \right\}$ with $n \geq 2$ is a decreasing sequence of positive terms and $\lim\limits_{n\to\infty} b_n = 0$. Thus, $\sum\limits_{n=2}^{\infty} \dfrac{(-1)^n}{n \ln n}$ converges by the Alternating Series Test. It follows that $\sum\limits_{n=2}^{\infty} \dfrac{(-1)^n}{n \ln n}$ is conditionally convergent.

35.

The graph gives us an estimate for the sum of the series $\sum\limits_{n=1}^{\infty} \dfrac{(-0.8)^n}{n!}$ of -0.55.

$b_8 = \dfrac{(0.8)^n}{8!} \approx 0.000\,004$, so

$$\sum_{n=1}^{\infty} \frac{(-0.8)^n}{n!} \approx s_7 = \sum_{n=1}^{7} \frac{(-0.8)^n}{n!}$$

$$\approx -0.8 + 0.32 - 0.085\overline{3} + 0.0170\overline{6} - 0.002\,731 + 0.000\,364 - 0.000\,042 \approx -0.5507$$

Adding b_8 to s_7 does not change the fourth decimal place of s_7, so the sum of the series, correct to four decimal places, is -0.5507.

36.

The graph gives us an estimate for the sum of the series $\sum\limits_{n=1}^{\infty} (-1)^{n-1} \dfrac{n}{8^n}$ of 0.1.

$b_6 = \dfrac{6}{8^6} \approx 0.000\,023$, so

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{n}{8^n} \approx s_5 = \sum_{n=1}^{5} (-1)^{n-1} \frac{n}{8^n}$$

$$\approx 0.125 - 0.03125 + 0.005\,859 - 0.000\,977 + 0.000\,153 \approx 0.0988$$

Adding b_6 to s_5 does not change the fourth decimal place of s_5, so the sum of the series, correct to four decimal places, is 0.0988.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. The series $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^6}$ satisfies (i) of the Alternating Series Test because $\frac{1}{(n+1)^6} < \frac{1}{n^6}$ and (ii) $\lim_{n\to\infty} \frac{1}{n^6} = 0$, so the series is convergent. Now $b_5 = \frac{1}{5^6} = 0.000064 > 0.00005$ and $b_6 = \frac{1}{6^6} \approx 0.00002 < 0.00005$, so by the Alternating Series Estimation Theorem, $n = 5$. (That is, since the 6th term is less than the desired error, we need to add the first 5 terms to get the sum to the desired accuracy.)

38. The series $\sum_{n=1}^{\infty} \frac{(-\frac{1}{3})^n}{n} = \sum_{n=1}^{\infty} (-1)^n \frac{1}{n3^n}$ satisfies (i) of the Alternating Series Test because $\frac{1}{(n+1)3^{n+1}} < \frac{1}{n3^n}$ and (ii) $\lim_{n\to\infty} \frac{1}{n3^n} = 0$, so the series is convergent. Now $b_5 = \frac{1}{5 \cdot 3^5} \approx 0.0008 > 0.0005$ and $b_6 = \frac{1}{6 \cdot 3^6} \approx 0.0002 < 0.0005$, so by the Alternating Series Estimation Theorem, $n = 5$. (That is, since the 6th term is less than the desired error, we need to add the first 5 terms to get the sum to the desired accuracy.)

39. The series $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2 2^n}$ satisfies (i) of the Alternating Series Test because $\frac{1}{(n+1)^2 2^{n+1}} < \frac{1}{n^2 2^n}$ and (ii) $\lim_{n\to\infty} \frac{1}{n^2 2^n} = 0$, so the series is convergent. Now $b_5 = \frac{1}{5^2 2^5} = 0.00125 > 0.0005$ and $b_6 = \frac{1}{6^2 2^6} \approx 0.0004 < 0.0005$, so by the Alternating Series Estimation Theorem, $n = 5$. (That is, since the 6th term is less than the desired error, we need to add the first 5 terms to get the sum to the desired accuracy.)

40. The series $\sum_{n=1}^{\infty} \left(-\frac{1}{n}\right)^n = \sum_{n=1}^{\infty} (-1)^n \frac{1}{n^n}$ satisfies (i) of the Alternating Series Test because $\frac{1}{(n+1)^{n+1}} < \frac{1}{n^n}$ and (ii) $\lim_{n\to\infty} \frac{1}{n^n} = 0$, so the series is convergent. Now $b_5 = \frac{1}{5^5} = 0.00032 > 0.00005$ and $b_6 = \frac{1}{6^6} \approx 0.00002 < 0.00005$, so by the Alternating Series Estimation Theorem, $n = 5$. (That is, since the 6th term is less than the desired error, we need to add the first 5 terms to get the sum to the desired accuracy.)

41. $b_4 = \frac{1}{8!} = \frac{1}{40{,}320} \approx 0.000\,025$, so

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!} \approx s_3 = \sum_{n=1}^{3} \frac{(-1)^n}{(2n)!} = -\frac{1}{2} + \frac{1}{24} - \frac{1}{720} \approx -0.459\,722$$

Adding b_4 to s_3 does not change the fourth decimal place of s_3, so by the Alternating Series Estimation Theorem, the sum of the series, correct to four decimal places, is -0.4597.

42. $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^6} \approx s_9 = \frac{1}{1^6} - \frac{1}{2^6} + \frac{1}{3^6} - \frac{1}{4^6} + \frac{1}{5^6} - \frac{1}{6^6} + \frac{1}{7^6} - \frac{1}{8^6} + \frac{1}{9^6} \approx 0.985\,552$. Subtracting $b_{10} = 1/10^6$ from s_9 does not change the fourth decimal place of s_9, so by the Alternating Series Estimation Theorem, the sum of the series, correct to four decimal places, is 0.9856.

43. $\sum_{n=1}^{\infty} (-1)^n n e^{-2n} \approx s_5 = -\frac{1}{e^2} + \frac{2}{e^4} - \frac{3}{e^6} + \frac{4}{e^8} - \frac{5}{e^{10}} \approx -0.105\,025$. Adding $b_6 = 6/e^{12} \approx 0.000\,037$ to s_5 does not change the fourth decimal place of s_5, so by the Alternating Series Estimation Theorem, the sum of the series, correct to four decimal places, is -0.1050.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n4^n} \approx s_6 = \frac{1}{4} - \frac{1}{2 \cdot 4^2} + \frac{1}{3 \cdot 4^3} - \frac{1}{4 \cdot 4^4} + \frac{1}{5 \cdot 4^5} - \frac{1}{6 \cdot 4^6} \approx 0.223136$. Adding $b_7 = \frac{1}{7 \cdot 4^7} \approx 0.000\,0087$ to s_6 does not change the fourth decimal place of s_6, so by the Alternating Series Estimation Theorem, the sum of the series, correct to four decimal places, is 0.2231.

45. $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots + \frac{1}{49} - \frac{1}{50} + \frac{1}{51} - \frac{1}{52} + \cdots$. The 50th partial sum of this series is an underestimate, since $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} = s_{50} + \left(\frac{1}{51} - \frac{1}{52}\right) + \left(\frac{1}{53} - \frac{1}{54}\right) + \cdots$, and the terms in parentheses are all positive. The result can be seen geometrically in Figure 1.

46. If $p > 0$, $\frac{1}{(n+1)^p} \leq \frac{1}{n^p}$ ($\{1/n^p\}$ is decreasing) and $\lim_{n\to\infty} \frac{1}{n^p} = 0$, so the series converges by the Alternating Series Test. If $p \leq 0$, $\lim_{n\to\infty} \frac{(-1)^{n-1}}{n^p}$ does not exist, so the series diverges by the Test for Divergence. Thus, $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^p}$ converges $\Leftrightarrow$ $p > 0$.

47. Clearly $b_n = \frac{1}{n+p}$ is decreasing and eventually positive and $\lim_{n\to\infty} b_n = 0$ for any p. So the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{n+p}$ converges (by the Alternating Series Test) for any p for which every b_n is defined, that is, $n + p \neq 0$ for $n \geq 1$, or p is not a negative integer.

48. Let $f(x) = \frac{(\ln x)^p}{x}$. Then $f'(x) = \frac{(\ln x)^{p-1}(p - \ln x)}{x^2} < 0$ if $x > e^p$ so f is eventually decreasing for every p. Clearly $\lim_{n\to\infty} \frac{(\ln n)^p}{n} = 0$ if $p \leq 0$, and if $p > 0$ we can apply l'Hospital's Rule $[\![p+1]\!]$ times to get a limit of 0 as well. So the series $\sum_{n=2}^{\infty} (-1)^{n-1} \frac{(\ln n)^p}{n}$ converges for all p (by the Alternating Series Test).

49. $\sum b_{2n} = \sum 1/(2n)^2$ clearly converges (by direct comparison with the p-series for $p = 2$). So suppose that $\sum (-1)^{n-1} b_n$ converges. Then by Theorem 11.2.8(ii), so does $\sum \left[(-1)^{n-1} b_n + b_n\right] = 2\left(1 + \frac{1}{3} + \frac{1}{5} + \cdots\right) = 2 \sum \frac{1}{2n-1}$. But this diverges by direct comparison with the harmonic series, a contradiction. Therefore, $\sum (-1)^{n-1} b_n$ must diverge. The Alternating Series Test does not apply since $\{b_n\}$ is not decreasing.

50. (a) We will prove this by induction. Let $P(n)$ be the proposition that $s_{2n} = h_{2n} - h_n$. $P(1)$ is the statement $s_2 = h_2 - h_1$, which is true since $1 - \frac{1}{2} = \left(1 + \frac{1}{2}\right) - 1$. So suppose that $P(n)$ is true. We will show that $P(n+1)$ must be true as a consequence.

$$\begin{aligned} h_{2n+2} - h_{n+1} &= \left(h_{2n} + \frac{1}{2n+1} + \frac{1}{2n+2}\right) - \left(h_n + \frac{1}{n+1}\right) = (h_{2n} - h_n) + \frac{1}{2n+1} - \frac{1}{2n+2} \\ &= s_{2n} + \frac{1}{2n+1} - \frac{1}{2n+2} = s_{2n+2} \end{aligned}$$

which is $P(n+1)$, and proves that $s_{2n} = h_{2n} - h_n$ for all n.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) We know that $h_{2n} - \ln(2n) \to \gamma$ and $h_n - \ln n \to \gamma$ as $n \to \infty$. So

$s_{2n} = h_{2n} - h_n = [h_{2n} - \ln(2n)] - (h_n - \ln n) + [\ln(2n) - \ln n]$, and

$\lim\limits_{n\to\infty} s_{2n} = \gamma - \gamma + \lim\limits_{n\to\infty} [\ln(2n) - \ln n] = \lim\limits_{n\to\infty} (\ln 2 + \ln n - \ln n) = \ln 2.$

51. (a) Since $\sum a_n$ is absolutely convergent, and since $\left|a_n^+\right| \le |a_n|$ and $\left|a_n^-\right| \le |a_n|$ (because a_n^+ and a_n^- each equal either a_n or 0), we conclude by the Direct Comparison Test that both $\sum a_n^+$ and $\sum a_n^-$ must be absolutely convergent. O*r:* Use Theorem 11.2.8.

(b) We will show by contradiction that both $\sum a_n^+$ and $\sum a_n^-$ must diverge. For suppose that $\sum a_n^+$ converged. Then so would $\sum\left(a_n^+ - \frac{1}{2}a_n\right)$ by Theorem 11.2.8. But $\sum\left(a_n^+ - \frac{1}{2}a_n\right) = \sum\left[\frac{1}{2}\left(a_n + |a_n|\right) - \frac{1}{2}a_n\right] = \frac{1}{2}\sum |a_n|$, which diverges because $\sum a_n$ is only conditionally convergent. Hence, $\sum a_n^+$ can't converge. Similarly, neither can $\sum a_n^-$.

52. Let $\sum b_n$ be the rearranged series constructed in the hint. [This series can be constructed by virtue of the result of Exercise 51(b).] This series will have partial sums s_n that oscillate in value back and forth across r. Since $\lim\limits_{n\to\infty} a_n = 0$ (by Theorem 11.2.6), and since the size of the oscillations $|s_n - r|$ is always less than $|a_n|$ because of the way $\sum b_n$ was constructed, we have that $\sum b_n = \lim\limits_{n\to\infty} s_n = r$.

53. Suppose that $\sum a_n$ is conditionally convergent.

(a) $\sum n^2 a_n$ is divergent: Suppose $\sum n^2 a_n$ converges. Then $\lim\limits_{n\to\infty} n^2 a_n = 0$ by Theorem 6 in Section 11.2, so there is an integer $N > 0$ such that $n > N \quad \Rightarrow \quad n^2\,|a_n| < 1$. For $n > N$, we have $|a_n| < \dfrac{1}{n^2}$, so $\sum\limits_{n>N} |a_n|$ converges by comparison with the convergent p-series $\sum\limits_{n>N} \dfrac{1}{n^2}$. In other words, $\sum a_n$ converges absolutely, contradicting the assumption that $\sum a_n$ is conditionally convergent. This contradiction shows that $\sum n^2 a_n$ diverges.

Remark: The same argument shows that $\sum n^p a_n$ diverges for any $p > 1$.

(b) $\sum\limits_{n=2}^{\infty} \dfrac{(-1)^n}{n \ln n}$ is conditionally convergent. It converges by the Alternating Series Test, but does not converge absolutely

$\Bigg[$by the Integral Test, since the function $f(x) = \dfrac{1}{x \ln x}$ is continuous, positive, and decreasing on $[2, \infty)$ and

$\displaystyle\int_2^{\infty} \frac{dx}{x \ln x} = \lim_{t\to\infty} \int_2^t \frac{dx}{x \ln x} = \lim_{t\to\infty} \Big[\ln(\ln x)\Big]_2^t = \infty\Bigg]$. Setting $a_n = \dfrac{(-1)^n}{n \ln n}$ for $n \ge 2$, we find that

$\sum\limits_{n=2}^{\infty} n a_n = \sum\limits_{n=2}^{\infty} \dfrac{(-1)^n}{\ln n}$ converges by the Alternating Series Test.

It is easy to find conditionally convergent series $\sum a_n$ such that $\sum n a_n$ diverges. Two examples are $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^{n-1}}{n}$ and $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^{n-1}}{\sqrt{n}}$, both of which converge by the Alternating Series Test and fail to converge absolutely because $\sum |a_n|$ is a p-series with $p \le 1$. In both cases, $\sum n a_n$ diverges by the Test for Divergence.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11.6 The Ratio and Root Tests

1. (a) Since $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = 8 > 1$, part (b) of the Ratio Test tells us that the series $\sum a_n$ is divergent.

(b) Since $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = 0.8 < 1$, part (a) of the Ratio Test tells us that the series $\sum a_n$ is absolutely convergent (and therefore convergent).

(c) Since $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = 1$, the Ratio Test fails and the series $\sum a_n$ might converge or it might diverge.

2. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{1}{a_n/a_{n+1}}\right| = \left|\frac{1}{2}\right| = \frac{1}{2} < 1$ Thus, the series $\sum a_n$ is absolutely convergent (and therefore convergent) by the Ratio Test.

3. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{n+1}{5^{n+1}}\cdot\frac{5^n}{n}\right| = \lim_{n\to\infty}\left|\frac{1}{5}\cdot\frac{n+1}{n}\right| = \frac{1}{5}\lim_{n\to\infty}\frac{1+1/n}{1} = \frac{1}{5}(1) = \frac{1}{5} < 1$, so the series $\sum_{n=1}^{\infty}\frac{n}{5^n}$ is absolutely convergent by the Ratio Test.

4. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-2)^{n+1}}{(n+1)^2}\cdot\frac{n^2}{(-2)^n}\right| = \lim_{n\to\infty}\left|(-2)\frac{n^2}{(n+1)^2}\right| = 2\lim_{n\to\infty}\frac{1}{(1+1/n)^2} = 2(1) = 2 > 1$, so the series $\sum_{n=1}^{\infty}\frac{(-2)^n}{n^2}$ is divergent by the Ratio Test.

5. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^n 3^{n+1}}{2^{n+1}(n+1)^3}\cdot\frac{2^n n^3}{(-1)^{n-1}3^n}\right| = \lim_{n\to\infty}\left|\left(-\frac{3}{2}\right)\frac{n^3}{(n+1)^3}\right| = \frac{3}{2}\lim_{n\to\infty}\frac{1}{(1+1/n)^3} = \frac{3}{2}(1) = \frac{3}{2} > 1$, so the series $\sum_{n=1}^{\infty}(-1)^{n-1}\frac{3^n}{2^n n^3}$ is divergent by the Ratio Test.

6. $\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-3)^{n+1}}{[2(n+1)+1]!}\cdot\frac{(2n+1)!}{(-3)^n}\right| = \lim_{n\to\infty}\left|(-3)\frac{1}{(2n+3)(2n+2)}\right| = 3\lim_{n\to\infty}\frac{1}{(2n+3)(2n+2)}$

$= 3(0) = 0 < 1$

so the series $\sum_{n=0}^{\infty}\frac{(-3)^n}{(2n+1)!}$ is absolutely convergent by the Ratio Test.

7. $\lim_{k\to\infty}\left|\frac{a_{k+1}}{a_k}\right| = \lim_{k\to\infty}\left|\frac{1}{(k+1)!}\cdot\frac{k!}{1}\right| = \lim_{k\to\infty}\frac{1}{k+1} = 0 < 1$, so the series $\sum_{k=1}^{\infty}\frac{1}{k!}$ is absolutely convergent by the Ratio Test. Since the terms of this series are positive, absolute convergence is the same as convergence.

8. $\lim_{k\to\infty}\left|\frac{a_{k+1}}{a_k}\right| = \lim_{k\to\infty}\left|\frac{(k+1)e^{-(k+1)}}{ke^{-k}}\right| = \lim_{k\to\infty}\left(\frac{k+1}{k}\cdot e^{-1}\right) = \frac{1}{e}\lim_{k\to\infty}\frac{1+1/k}{1} = \frac{1}{e}(1) = \frac{1}{e} < 1$, so the series $\sum_{k=1}^{\infty}ke^{-k}$ is absolutely convergent by the Ratio Test. Since the terms of this series are positive, absolute convergence is the same as convergence.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left[\dfrac{10^{n+1}}{(n+2)\,4^{2n+3}}\cdot\dfrac{(n+1)\,4^{2n+1}}{10^n}\right] = \lim\limits_{n\to\infty}\left(\dfrac{10}{4^2}\cdot\dfrac{n+1}{n+2}\right) = \dfrac{5}{8} < 1$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{10^n}{(n+1)4^{2n+1}}$ is absolutely convergent by the Ratio Test. Since the terms of this series are positive, absolute convergence is the same as convergence.

10. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left[\dfrac{(n+1)!}{100^{n+1}}\cdot\dfrac{100^n}{n!}\right] = \lim\limits_{n\to\infty}\dfrac{n+1}{100} = \infty$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{n!}{100^n}$ diverges by the Ratio Test.

11. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(n+1)\pi^{n+1}}{(-3)^n}\cdot\dfrac{(-3)^{n-1}}{n\pi^n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{\pi}{-3}\cdot\dfrac{n+1}{n}\right| = \dfrac{\pi}{3}\lim\limits_{n\to\infty}\dfrac{1+1/n}{1} = \dfrac{\pi}{3}(1) = \dfrac{\pi}{3} > 1$, so the series $\displaystyle\sum_{n=1}^{\infty}\frac{n\pi^n}{(-3)^{n-1}}$ diverges by the Ratio Test. *Or:* Since $\lim\limits_{n\to\infty}|a_n| = \infty$, the series diverges by the Test for Divergence.

12. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(n+1)^{10}}{(-10)^{n+2}}\cdot\dfrac{(-10)^{n+1}}{n^{10}}\right| = \lim\limits_{n\to\infty}\left|\dfrac{1}{-10}\left(\dfrac{n+1}{n}\right)^{10}\right| = \dfrac{1}{10}\lim\limits_{n\to\infty}\left(1+\dfrac{1}{n}\right)^{10} = \dfrac{1}{10}(1) = \dfrac{1}{10} < 1$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{n^{10}}{(-10)^{n+1}}$ is absolutely convergent by the Ratio Test.

13. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{\cos[(n+1)\pi/3]}{(n+1)!}\cdot\dfrac{n!}{\cos(n\pi/3)}\right| = \lim\limits_{n\to\infty}\left|\dfrac{\cos[(n+1)\pi/3]}{(n+1)\cos(n\pi/3)}\right| = \lim\limits_{n\to\infty}\dfrac{c}{n+1} = 0 < 1$ (where $0 < c \le 2$ for all positive integers n), so the series $\sum\limits_{n=1}^{\infty}\dfrac{\cos(n\pi/3)}{n!}$ is absolutely convergent by the Ratio Test.

14. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(n+1)!}{(n+1)^{n+1}}\cdot\dfrac{n^n}{n!}\right| = \lim\limits_{n\to\infty}\dfrac{(n+1)n^n}{(n+1)^{n+1}} = \lim\limits_{n\to\infty}\dfrac{n^n}{(n+1)^n} = \lim\limits_{n\to\infty}\dfrac{1}{(1+1/n)^n} = \dfrac{1}{e} < 1$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{n!}{n^n}$ is absolutely convergent by the Ratio Test.

15. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(n+1)^{100}100^{n+1}}{(n+1)!}\cdot\dfrac{n!}{n^{100}100^n}\right| = \lim\limits_{n\to\infty}\dfrac{100}{n+1}\left(\dfrac{n+1}{n}\right)^{100} = \lim\limits_{n\to\infty}\dfrac{100}{n+1}\left(1+\dfrac{1}{n}\right)^{100}$

$$= 0\cdot 1 = 0 < 1$$

so the series $\sum\limits_{n=1}^{\infty}\dfrac{n^{100}100^n}{n!}$ is absolutely convergent by the Ratio Test.

16. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{[2(n+1)]!}{[(n+1)!]^2}\cdot\dfrac{(n!)^2}{(2n)!}\right| = \lim\limits_{n\to\infty}\dfrac{(2n+2)(2n+1)}{(n+1)(n+1)} = \lim\limits_{n\to\infty}\dfrac{(2+2/n)(2+1/n)}{(1+1/n)(1+1/n)} = \dfrac{2\cdot 2}{1\cdot 1} = 4 > 1$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{(2n)!}{(n!)^2}$ diverges by the Ratio Test.

17. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(-1)^n(n+1)!}{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)(2n+1)}\cdot\dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{(-1)^{n-1}n!}\right| = \lim\limits_{n\to\infty}\dfrac{n+1}{2n+1}$

$$= \lim_{n\to\infty}\frac{1+1/n}{2+1/n} = \frac{1}{2} < 1,$$

so the series $1 - \dfrac{2!}{1\cdot 3} + \dfrac{3!}{1\cdot 3\cdot 5} - \dfrac{4!}{1\cdot 3\cdot 5\cdot 7} + \cdots + (-1)^{n-1}\dfrac{n!}{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)} + \cdots$ is absolutely convergent by the Ratio Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $\dfrac{2}{3}+\dfrac{2\cdot 5}{3\cdot 5}+\dfrac{2\cdot 5\cdot 8}{3\cdot 5\cdot 7}+\dfrac{2\cdot 5\cdot 8\cdot 11}{3\cdot 5\cdot 7\cdot 9}+\cdots=\sum\limits_{n=1}^{\infty}\dfrac{2\cdot 5\cdot 8\cdot 11\cdot\cdots\cdot(3n-1)}{3\cdot 5\cdot 7\cdot 9\cdot\cdots\cdot(2n+1)}$.

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left|\frac{2\cdot 5\cdot 8\cdot\cdots\cdot(3n-1)(3n+2)}{3\cdot 5\cdot 7\cdot\cdots\cdot(2n+1)(2n+3)}\cdot\frac{3\cdot 5\cdot 7\cdot\cdots\cdot(2n+1)}{2\cdot 5\cdot 8\cdot\cdots\cdot(3n-1)}\right|$$
$$=\lim_{n\to\infty}\frac{3n+2}{2n+3}=\lim_{n\to\infty}\frac{3+2/n}{2+3/n}=\frac{3}{2}>1,$$

so the given series diverges by the Ratio Test.

19. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right|=\lim\limits_{n\to\infty}\left|\dfrac{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)(2n+2)}{(n+1)!}\cdot\dfrac{n!}{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)}\right|=\lim\limits_{n\to\infty}\dfrac{2n+2}{n+1}=\lim\limits_{n\to\infty}\dfrac{2(n+1)}{n+1}=2>1$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{2\cdot 4\cdot 6\cdot\cdots\cdot(2n)}{n!}$ diverges by the Ratio Test.

20. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right|=\lim\limits_{n\to\infty}\left|\dfrac{2^{n+1}\,(n+1)!}{5\cdot 8\cdot 11\cdot\cdots\cdot(3n+2)\,(3n+5)}\cdot\dfrac{5\cdot 8\cdot 11\cdot\cdots\cdot(3n+2)}{2^n n!}\right|=\lim\limits_{n\to\infty}\dfrac{2(n+1)}{3n+5}=\dfrac{2}{3}<1$, so the series $\sum\limits_{n=1}^{\infty}(-1)^n\dfrac{2^n n!}{5\cdot 8\cdot 11\cdot\cdots\cdot(3n+2)}$ is absolutely convergent by the Ratio Test.

21. $\lim\limits_{n\to\infty}\sqrt[n]{|a_n|}=\lim\limits_{n\to\infty}\dfrac{n^2+1}{2n^2+1}=\lim\limits_{n\to\infty}\dfrac{1+1/n^2}{2+1/n^2}=\dfrac{1}{2}<1$, so the series $\sum\limits_{n=1}^{\infty}\left(\dfrac{n^2+1}{2n^2+1}\right)^n$ is absolutely convergent by the Root Test.

22. $\lim\limits_{n\to\infty}\sqrt[n]{|a_n|}=\lim\limits_{n\to\infty}\sqrt[n]{\left|\dfrac{(-2)^n}{n^n}\right|}=\lim\limits_{n\to\infty}\dfrac{2}{n}=0<1$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{(-2)^n}{n^n}$ is absolutely convergent by the Root Test.

23. $\lim\limits_{n\to\infty}\sqrt[n]{|a_n|}=\lim\limits_{n\to\infty}\sqrt[n]{\left|\dfrac{(-1)^{n-1}}{(\ln n)^n}\right|}=\lim\limits_{n\to\infty}\dfrac{1}{\ln n}=0<1$, so the series $\sum\limits_{n=2}^{\infty}\dfrac{(-1)^{n-1}}{(\ln n)^n}$ is absolutely convergent by the Root Test.

24. $\lim\limits_{n\to\infty}\sqrt[n]{|a_n|}=\lim\limits_{n\to\infty}\sqrt[n]{\left|\left(\dfrac{-2n}{n+1}\right)^{5n}\right|}=\lim\limits_{n\to\infty}\dfrac{2^5\,n^5}{(n+1)^5}=32\lim\limits_{n\to\infty}\dfrac{1}{\left(\dfrac{n+1}{n}\right)^5}=32\lim\limits_{n\to\infty}\dfrac{1}{(1+1/n)^5}$

$$=32(1)=32>1,$$

so the series $\sum\limits_{n=1}^{\infty}\left(\dfrac{-2n}{n+1}\right)^{5n}$ diverges by the Root Test.

25. $\lim\limits_{n\to\infty}\sqrt[n]{|a_n|}=\lim\limits_{n\to\infty}\sqrt[n]{\left(1+\dfrac{1}{n}\right)^{n^2}}=\lim\limits_{n\to\infty}\left(1+\dfrac{1}{n}\right)^n=e>1$ [by Equation 3.6.6], so the series $\sum\limits_{n=1}^{\infty}\left(1+\dfrac{1}{n}\right)^{n^2}$ diverges by the Root Test.

26. $\lim\limits_{n\to\infty}\sqrt[n]{|a_n|}=\lim\limits_{n\to\infty}\sqrt[n]{|(\arctan n)^n|}=\lim\limits_{n\to\infty}\arctan n=\frac{\pi}{2}>1$, so the series $\sum\limits_{n=0}^{\infty}(\arctan n)^n$ diverges by the Root Test.

27. $\sum\limits_{n=2}^{\infty}a_n=\sum\limits_{n=2}^{\infty}\dfrac{(-1)^n\ln n}{n}=\sum\limits_{n=2}^{\infty}(-1)^n b_n$. Now $b_n=\dfrac{\ln n}{n}>0$ for $n\geq 2$, and $\{b_n\}$ is decreasing for $n\geq 3$ since $\left(\dfrac{\ln x}{x}\right)'=\dfrac{x\cdot\frac{1}{x}-\ln x\cdot 1}{x^2}=\dfrac{1-\ln x}{x^2}<0$ when $\ln x>1$ or $x>e\approx 2.7$. Also, $\lim\limits_{n\to\infty}b_n=\lim\limits_{n\to\infty}\dfrac{\ln n}{n}\overset{\text{H}}{=}\lim\limits_{n\to\infty}\dfrac{1/n}{1}=0$,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

so the series $\sum_{n=2}^{\infty} \frac{(-1)^n \ln n}{n}$ converges by the Alternating Series Test. To determine absolute convergence, note that $\left|\frac{(-1)^n \ln n}{n}\right| = \frac{\ln n}{n} > \frac{1}{n}$ for $n \geq 3$, so $\sum_{n=2}^{\infty} \left|\frac{(-1)^n \ln n}{n}\right|$ is divergent by direct comparison with $\sum_{n=2}^{\infty} \frac{1}{n}$, which is divergent. Hence, $\sum_{n=2}^{\infty} \frac{(-1)^n \ln n}{n}$ is conditionally convergent.

28. $\lim_{n\to\infty} \sqrt[n]{|a_n|} = \lim_{n\to\infty} \sqrt[n]{\left|\left(\frac{1-n}{2+3n}\right)^n\right|} = \lim_{n\to\infty} \frac{n-1}{3n+2} = \lim_{n\to\infty} \frac{1-1/n}{3+2/n} = \frac{1}{3} < 1$, so the series $\sum_{n=1}^{\infty} \left(\frac{1-n}{2+3n}\right)^n$ is absolutely convergent by the Root Test.

29. $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-9)^{n+1}}{(n+1)10^{n+2}} \cdot \frac{n10^{n+1}}{(-9)^n}\right| = \lim_{n\to\infty} \left|\frac{(-9)n}{10(n+1)}\right| = \frac{9}{10} \lim_{n\to\infty} \frac{1}{1+1/n} = \frac{9}{10}(1) = \frac{9}{10} < 1$, so the series $\sum_{n=1}^{\infty} \frac{(-9)^n}{n10^{n+1}}$ is absolutely convergent by the Ratio Test.

30. $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(n+1)5^{2n+2}}{10^{n+2}} \cdot \frac{10^{n+1}}{n5^{2n}}\right| = \lim_{n\to\infty} \frac{5^2(n+1)}{10n} = \frac{5}{2} \lim_{n\to\infty} \left(1+\frac{1}{n}\right) = \frac{5}{2}(1) = \frac{5}{2} > 1$, so the series $\sum_{n=1}^{\infty} \frac{n5^{2n}}{10^{n+1}}$ diverges by the Ratio Test. *Or:* Since $\lim_{n\to\infty} a_n = \infty$, the series diverges by the Test for Divergence.

31. $\lim_{n\to\infty} \sqrt[n]{|a_n|} = \lim_{n\to\infty} \sqrt[n]{\left|\left(\frac{n}{\ln n}\right)^n\right|} = \lim_{n\to\infty} \frac{n}{\ln n} = \lim_{x\to\infty} \frac{x}{\ln x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{1}{1/x} = \lim_{x\to\infty} x = \infty$, so the series $\sum_{n=2}^{\infty} \left(\frac{n}{\ln n}\right)^n$ diverges by the Root Test.

32. $\left|\frac{\sin(n\pi/6)}{1+n\sqrt{n}}\right| \leq \frac{1}{1+n\sqrt{n}} < \frac{1}{n^{3/2}}$, so the series $\sum_{n=1}^{\infty} \frac{\sin(n\pi/6)}{1+n\sqrt{n}}$ converges by direct comparison with the convergent p-series $\sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$ $(p = \frac{3}{2} > 1)$. It follows that the given series is absolutely convergent.

33. $\left|\frac{(-1)^n \arctan n}{n^2}\right| < \frac{\pi/2}{n^2}$, so since $\sum_{n=1}^{\infty} \frac{\pi/2}{n^2} = \frac{\pi}{2} \sum_{n=1}^{\infty} \frac{1}{n^2}$ converges $(p = 2 > 1)$, the given series $\sum_{n=1}^{\infty} \frac{(-1)^n \arctan n}{n^2}$ converges absolutely by the Direct Comparison Test.

34. $\sum_{n=2}^{\infty} a_n = \sum_{n=2}^{\infty} \frac{(-1)^n}{\sqrt{n}\ln n} = \sum_{n=2}^{\infty} (-1)^n b_n$. Now $b_n = \frac{1}{\sqrt{n}\ln n} > 0$ for $n \geq 2$, $\{b_n\}$ is decreasing for $n \geq 2$, and $\lim_{n\to\infty} b_n = 0$, so the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{\sqrt{n}\ln n}$ converges by the Alternating Series Test. Also, observe that $\left|\frac{(-1)^n}{\sqrt{n}\ln n}\right| = \frac{1}{\sqrt{n}\ln n} > \frac{1}{\sqrt{n}\sqrt{n}} = \frac{1}{n}$ for $n \geq 2$, so the series $\sum_{n=2}^{\infty} \left|\frac{(-1)^n}{\sqrt{n}\ln n}\right|$ is divergent by direct comparison with $\sum_{n=2}^{\infty} \frac{1}{n}$, which is a divergent (partial) p-series $[p = 1 \leq 1]$. Thus, $\sum_{n=2}^{\infty} \frac{(-1)^n}{\sqrt{n}\ln n}$ is conditionally convergent.

35. By the recursive definition, $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{5n+1}{4n+3}\right| = \frac{5}{4} > 1$, so the series diverges by the Ratio Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. By the recursive definition, $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{2+\cos n}{\sqrt{n}}\right| = 0 < 1$, so the series converges absolutely by the Ratio Test.

37. The series $\sum\limits_{n=1}^{\infty}\dfrac{b_n^n\cos n\pi}{n} = \sum\limits_{n=1}^{\infty}(-1)^n\dfrac{b_n^n}{n}$, where $b_n > 0$ for $n \geq 1$ and $\lim\limits_{n\to\infty} b_n = \dfrac{1}{2}$.

$\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(-1)^{n+1}b_n^{n+1}}{n+1}\cdot\dfrac{n}{(-1)^n b_n^n}\right| = \lim\limits_{n\to\infty} b_n\dfrac{n}{n+1} = \dfrac{1}{2}(1) = \dfrac{1}{2} < 1$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{b_n^n\cos n\pi}{n}$ is absolutely convergent by the Ratio Test.

38. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(-1)^{n+1}(n+1)!}{(n+1)^{n+1}b_1b_2\cdots b_nb_{n+1}}\cdot\dfrac{n^nb_1b_2\cdots b_n}{(-1)^n\,n!}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(-1)(n+1)n^n}{b_{n+1}(n+1)^{n+1}}\right| = \lim\limits_{n\to\infty}\dfrac{n^n}{b_{n+1}(n+1)^n}$

$$= \lim_{n\to\infty}\frac{1}{b_{n+1}}\left(\frac{n}{n+1}\right)^n = \lim_{n\to\infty}\frac{1}{b_{n+1}}\left(\frac{1}{1+1/n}\right)^n = \lim_{n\to\infty}\frac{1}{b_{n+1}(1+1/n)^n} = \frac{1}{\frac{1}{2}e} = \frac{2}{e} < 1$$

so the series $\sum\limits_{n=1}^{\infty}\dfrac{(-1)^n\,n!}{n^nb_1b_2b_3\cdots b_n}$ is absolutely convergent by the Ratio Test.

39. (a) $\lim\limits_{n\to\infty}\left|\dfrac{1/(n+1)^3}{1/n^3}\right| = \lim\limits_{n\to\infty}\dfrac{n^3}{(n+1)^3} = \lim\limits_{n\to\infty}\dfrac{1}{(1+1/n)^3} = 1$. Inconclusive for $\sum\limits_{n=1}^{\infty}\dfrac{1}{n^3}$.

(b) $\lim\limits_{n\to\infty}\left|\dfrac{(n+1)}{2^{n+1}}\cdot\dfrac{2^n}{n}\right| = \lim\limits_{n\to\infty}\dfrac{n+1}{2n} = \lim\limits_{n\to\infty}\left(\dfrac{1}{2}+\dfrac{1}{2n}\right) = \dfrac{1}{2}$. Conclusive (convergent) for $\sum\limits_{n=1}^{\infty}\dfrac{n}{2^n}$.

(c) $\lim\limits_{n\to\infty}\left|\dfrac{(-3)^n}{\sqrt{n+1}}\cdot\dfrac{\sqrt{n}}{(-3)^{n-1}}\right| = 3\lim\limits_{n\to\infty}\sqrt{\dfrac{n}{n+1}} = 3\lim\limits_{n\to\infty}\sqrt{\dfrac{1}{1+1/n}} = 3$. Conclusive (divergent) for $\sum\limits_{n=1}^{\infty}\dfrac{(-3)^{n-1}}{\sqrt{n}}$.

(d) $\lim\limits_{n\to\infty}\left|\dfrac{\sqrt{n+1}}{1+(n+1)^2}\cdot\dfrac{1+n^2}{\sqrt{n}}\right| = \lim\limits_{n\to\infty}\left[\sqrt{1+\dfrac{1}{n}}\cdot\dfrac{1/n^2+1}{1/n^2+(1+1/n)^2}\right] = 1$. Inconclusive for $\sum\limits_{n=1}^{\infty}\dfrac{\sqrt{n}}{1+n^2}$.

40. We use the Ratio Test for the series $\sum\limits_{n=1}^{\infty}\dfrac{(n!)^2}{(kn)!}$:

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{[(n+1)!]^2/[k(n+1)]!}{(n!)^2/(kn)!}\right| = \lim_{n\to\infty}\left|\frac{(n+1)^2}{[k(n+1)]\,[k(n+1)-1]\cdots[kn+1]}\right|$$

Now if $k = 1$, then this is equal to $\lim\limits_{n\to\infty}\left|\dfrac{(n+1)^2}{(n+1)}\right| = \infty$, so the series diverges; if $k = 2$, the limit is

$\lim\limits_{n\to\infty}\left|\dfrac{(n+1)^2}{(2n+2)(2n+1)}\right| = \dfrac{1}{4} < 1$, so the series converges, and if $k > 2$, then the highest power of n in the denominator is larger than 2, and so the limit is 0, indicating convergence. So the series converges for $k \geq 2$.

41. (a) $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{x^{n+1}}{(n+1)!}\cdot\dfrac{n!}{x^n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{x}{n+1}\right| = |x|\lim\limits_{n\to\infty}\dfrac{1}{n+1} = |x|\cdot 0 = 0 < 1$, so by the Ratio Test the series $\sum\limits_{n=0}^{\infty}\dfrac{x^n}{n!}$ converges for all x.

(b) Since the series of part (a) always converges, we must have $\lim\limits_{n\to\infty}\dfrac{x^n}{n!} = 0$ by Theorem 11.2.6.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. (a) $R_n = a_{n+1} + a_{n+2} + a_{n+3} + a_{n+4} + \cdots = a_{n+1}\left(1 + \dfrac{a_{n+2}}{a_{n+1}} + \dfrac{a_{n+3}}{a_{n+1}} + \dfrac{a_{n+4}}{a_{n+1}} + \cdots\right)$

$$= a_{n+1}\left(1 + \frac{a_{n+2}}{a_{n+1}} + \frac{a_{n+3}}{a_{n+2}}\frac{a_{n+2}}{a_{n+1}} + \frac{a_{n+4}}{a_{n+3}}\frac{a_{n+3}}{a_{n+2}}\frac{a_{n+2}}{a_{n+1}} + \cdots\right)$$

$$= a_{n+1}(1 + r_{n+1} + r_{n+2}r_{n+1} + r_{n+3}r_{n+2}r_{n+1} + \cdots) \quad (\star)$$

$$\le a_{n+1}\left(1 + r_{n+1} + r_{n+1}^2 + r_{n+1}^3 + \cdots\right) \quad \text{[since } \{r_n\} \text{ is decreasing]} \quad = \frac{a_{n+1}}{1 - r_{n+1}}$$

(b) Note that since $\{r_n\}$ is increasing and $r_n \to L$ as $n \to \infty$, we have $r_n < L$ for all n. So, starting with equation $(\star)$,

$$R_n = a_{n+1}(1 + r_{n+1} + r_{n+2}r_{n+1} + r_{n+3}r_{n+2}r_{n+1} + \cdots) \le a_{n+1}\left(1 + L + L^2 + L^3 + \cdots\right) = \frac{a_{n+1}}{1 - L}.$$

43. (a) $s_5 = \sum\limits_{n=1}^{5} \dfrac{1}{n2^n} = \dfrac{1}{2} + \dfrac{1}{8} + \dfrac{1}{24} + \dfrac{1}{64} + \dfrac{1}{160} = \dfrac{661}{960} \approx 0.68854$. Now the ratios

$r_n = \dfrac{a_{n+1}}{a_n} = \dfrac{n2^n}{(n+1)2^{n+1}} = \dfrac{n}{2(n+1)}$ form an increasing sequence, since

$r_{n+1} - r_n = \dfrac{n+1}{2(n+2)} - \dfrac{n}{2(n+1)} = \dfrac{(n+1)^2 - n(n+2)}{2(n+1)(n+2)} = \dfrac{1}{2(n+1)(n+2)} > 0$. So by Exercise 42(b), the error

in using s_5 is $R_5 \le \dfrac{a_6}{1 - \lim\limits_{n\to\infty} r_n} = \dfrac{1/\left(6 \cdot 2^6\right)}{1 - 1/2} = \dfrac{1}{192} \approx 0.00521$.

(b) The error in using s_n as an approximation to the sum is $R_n = \dfrac{a_{n+1}}{1 - \frac{1}{2}} = \dfrac{2}{(n+1)2^{n+1}}$. We want $R_n < 0.00005 \quad \Leftrightarrow$

$\dfrac{1}{(n+1)2^n} < 0.00005 \quad \Leftrightarrow \quad (n+1)2^n > 20{,}000$. To find such an n we can use trial and error or a graph. We calculate

$(11+1)2^{11} = 24{,}576$, so $s_{11} = \sum\limits_{n=1}^{11} \dfrac{1}{n2^n} \approx 0.693109$ is within 0.00005 of the actual sum.

44. $s_{10} = \sum\limits_{n=1}^{10} \dfrac{n}{2^n} = \dfrac{1}{2} + \dfrac{2}{4} + \dfrac{3}{8} + \cdots + \dfrac{10}{1024} \approx 1.988$. The ratios $r_n = \dfrac{a_{n+1}}{a_n} = \dfrac{n+1}{2^{n+1}} \cdot \dfrac{2^n}{n} = \dfrac{n+1}{2n} = \dfrac{1}{2}\left(1 + \dfrac{1}{n}\right)$ form a

decreasing sequence, and $r_{11} = \dfrac{11+1}{2(11)} = \dfrac{12}{22} = \dfrac{6}{11} < 1$, so by Exercise 42(a), the error in using s_{10} to approximate the sum

of the series $\sum\limits_{n=1}^{\infty} \dfrac{n}{2^n}$ is $R_{10} \le \dfrac{a_{11}}{1 - r_{11}} = \dfrac{\frac{11}{2048}}{1 - \frac{6}{11}} = \dfrac{121}{10{,}240} \approx 0.0118$.

45. (i) Following the hint, we get that $|a_n| < r^n$ for $n \ge N$, and so since the geometric series $\sum_{n=1}^{\infty} r^n$ converges $[0 < r < 1]$, the series $\sum_{n=N}^{\infty} |a_n|$ converges as well by the Direct Comparison Test, and hence so does $\sum_{n=1}^{\infty} |a_n|$, so $\sum_{n=1}^{\infty} a_n$ is absolutely convergent.

(ii) If $\lim\limits_{n\to\infty} \sqrt[n]{|a_n|} = L > 1$, then there is an integer N such that $\sqrt[n]{|a_n|} > 1$ for all $n \ge N$, so $|a_n| > 1$ for $n \ge N$. Thus, $\lim\limits_{n\to\infty} a_n \ne 0$, so $\sum_{n=1}^{\infty} a_n$ diverges by the Test for Divergence.

(iii) Consider $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ [diverges] and $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$ [converges]. For each sum, $\lim\limits_{n\to\infty} \sqrt[n]{|a_n|} = 1$, so the Root Test is inconclusive.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. (a) $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{[4(n+1)]!\,[1103+26{,}390(n+1)]}{[(n+1)!]^4\,396^{4(n+1)}}\cdot\dfrac{(n!)^4\,396^{4n}}{(4n)!\,(1103+26{,}390n)}\right|$

$$= \lim_{n\to\infty}\frac{(4n+4)(4n+3)(4n+2)(4n+1)(26{,}390n+27{,}493)}{(n+1)^4\,396^4\,(26{,}390n+1103)} = \frac{4^4}{396^4} = \frac{1}{99^4} < 1,$$

so by the Ratio Test, the series $\sum\limits_{n=0}^{\infty}\dfrac{(4n)!\,(1103+26{,}390n)}{(n!)^4\,396^{4n}}$ converges.

(b) $\dfrac{1}{\pi} = \dfrac{2\sqrt{2}}{9801}\sum\limits_{n=0}^{\infty}\dfrac{(4n)!\,(1103+26{,}390n)}{(n!)^4\,396^{4n}}$

With the first term $(n=0)$, $\dfrac{1}{\pi}\approx\dfrac{2\sqrt{2}}{9801}\cdot\dfrac{1103}{1}\quad\Rightarrow\quad\pi\approx 3.141\,592\,73$, so we get 6 correct decimal places of π, which is $3.141\,592\,653\,589\,793\,238$ to 18 decimal places.

With the second term $(n=1)$, $\dfrac{1}{\pi}\approx\dfrac{2\sqrt{2}}{9801}\left(\dfrac{1103}{1}+\dfrac{4!\,(1103+26{,}390)}{396^4}\right)\quad\Rightarrow\quad\pi\approx 3.141\,592\,653\,589\,793\,878$, so we get 15 correct decimal places of π.

11.7 Strategy for Testing Series

1. (a) $\sum\limits_{n=1}^{\infty}\dfrac{1}{5^n} = \sum\limits_{n=1}^{\infty}\left(\dfrac{1}{5}\right)^n$ is a geometric series with ratio $r=\dfrac{1}{5}$. Since $|r|=\dfrac{1}{5}<1$, the series converges.

(b) $\dfrac{1}{5^n+n}<\dfrac{1}{5^n}$ for $n\ge 1$, so $\sum\limits_{n=1}^{\infty}\dfrac{1}{5^n+n}$ converges by direct comparison with $\sum\limits_{n=1}^{\infty}\dfrac{1}{5^n}$, which converges because it is a geometric series with $|r|=\dfrac{1}{5}<1$.

2. (a) $\sum\limits_{n=1}^{\infty}\dfrac{(-1)^n}{n^{3/2}} = \sum\limits_{n=1}^{\infty}(-1)^n b_n$. Now, $b_n=\dfrac{1}{n^{3/2}}>0$ for $n\ge 1$, $\{b_n\}$ is decreasing for $n\ge 1$, and $\lim\limits_{n\to\infty} b_n = 0$, so the series converges by the Alternating Series Test.

(b) $\sum\limits_{n=1}^{\infty}\dfrac{1}{n^{3/2}}$ is a p-series with $p=\dfrac{3}{2}>1$, so it converges.

3. (a) $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{n+1}{3^{n+1}}\cdot\dfrac{3^n}{n}\right| = \lim\limits_{n\to\infty}\dfrac{n+1}{3n} = \lim\limits_{n\to\infty}\left(\dfrac{n}{3n}+\dfrac{1}{3n}\right) = \lim\limits_{n\to\infty}\left(\dfrac{1}{3}+\dfrac{1}{3n}\right) = \dfrac{1}{3}<1,$

so the series $\sum\limits_{n=1}^{\infty}\dfrac{n}{3^n}$ is absolutely convergent (and therefore convergent) by the Ratio Test.

(b) $\lim\limits_{n\to\infty}\dfrac{3^n}{n} \overset{\text{H}}{=} \lim\limits_{n\to\infty}\dfrac{3^n\ln 3}{1}=\infty$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{3^n}{n}$ diverges by the Test for Divergence.

4. (a) $\lim\limits_{n\to\infty}\dfrac{n+1}{n} = \lim\limits_{n\to\infty}\left(1+\dfrac{1}{n}\right)=1\ne 0$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{n+1}{n}$ diverges by the Test for Divergence.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\lim\limits_{n\to\infty} |a_n| = \lim\limits_{n\to\infty} \left|(-1)^n \dfrac{n+1}{n}\right| = \lim\limits_{n\to\infty} \dfrac{n+1}{n} = \lim\limits_{n\to\infty} \left(1+\dfrac{1}{n}\right) = 1 \neq 0$, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n+1}{n}$ diverges by the Test for Divergence. $\left[\text{Note that } \lim\limits_{n\to\infty} (-1)^n \dfrac{n+1}{n} \text{ does not exist.}\right]$

5. (a) Use the Limit Comparison Test with $a_n = \dfrac{n}{n^2+1}$ and $b_n = \dfrac{1}{n}$.

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{n^2}{n^2+1} = \lim\limits_{n\to\infty} \dfrac{1}{1+1/n^2} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ is the divergent harmonic series, the series $\sum\limits_{n=1}^{\infty} \dfrac{n^2}{n^2+1}$ also diverges.

(b) $\lim\limits_{n\to\infty} \sqrt[n]{|a_n|} = \lim\limits_{n\to\infty} \sqrt[n]{\left(\dfrac{n}{n^2+1}\right)^n} = \lim\limits_{n\to\infty} \dfrac{n}{n^2+1} = \lim\limits_{n\to\infty} \dfrac{1/n}{1+1/n^2} = \dfrac{0}{1} = 0 < 1$, so the series $\sum\limits_{n=1}^{\infty} \left(\dfrac{n}{n^2+1}\right)^n$ converges by the Root Test.

6. (a) $\dfrac{\ln n}{n} > \dfrac{1}{n}$ for $n \geq 3$, so $\sum\limits_{n=1}^{\infty} \dfrac{\ln n}{n}$ diverges by direct comparison with $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$, which diverges because it is a p-series with $p = 1 \leq 1$.

(b) Let $f(x) = \dfrac{1}{x \ln x}$. Then f is continuous and positive on $[10, \infty)$, and also decreasing since $f'(x) = -\dfrac{1+\ln x}{x^2(\ln x)^2} < 0$ for $x > 10$, so we can use the Integral Test. $\displaystyle\int_{10}^{\infty} \frac{1}{x \ln x}\,dx = \lim_{t\to\infty} \Big[\ln|\ln x|\Big]_{10}^{t} = \lim_{t\to\infty} [\ln(\ln t) - \ln(\ln 10)] = \infty$, so the series $\sum\limits_{n=10}^{\infty} \dfrac{1}{n \ln n}$ diverges.

7. (a) Since $n! > n^2$ for $n \geq 4$, we have $\dfrac{1}{n+n!} < \dfrac{1}{n+n^2} < \dfrac{1}{n^2}$ for $n \geq 4$. Thus, $\sum\limits_{n=1}^{\infty} \dfrac{1}{n+n!}$ converges by direct comparison with $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$, which converges because it is a p-series with $p = 2 > 1$.

(b) $\dfrac{1}{n} + \dfrac{1}{n!} > \dfrac{1}{n}$ for $n \geq 1$, so $\sum\limits_{n=1}^{\infty} \left(\dfrac{1}{n} + \dfrac{1}{n!}\right)$ diverges by direct comparison with $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$, which diverges because it is a p-series with $p = 1 \leq 1$.

8. (a) Use the Limit Comparison Test with $a_n = \dfrac{1}{\sqrt{n^2+1}}$ and $b_n = \dfrac{1}{n}$.

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{n}{\sqrt{n^2+1}} = \lim\limits_{n\to\infty} \dfrac{n}{n\sqrt{1+1/n^2}} = \lim\limits_{n\to\infty} \dfrac{1}{\sqrt{1+1/n^2}} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ is the divergent harmonic series, the series $\sum\limits_{n=1}^{\infty} \dfrac{1}{\sqrt{n^2+1}}$ also diverges.

(b) $\dfrac{1}{n\sqrt{n^2+1}} < \dfrac{1}{n\sqrt{n^2}} = \dfrac{1}{n \cdot n} = \dfrac{1}{n^2}$ for $n \geq 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{1}{n\sqrt{n^2+1}}$ converges by direct comparison with $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$, which converges because it is a p-series with $p = 2 > 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. Use the Limit Comparison Test with $a_n = \dfrac{n^2-1}{n^3+1}$ and $b_n = \dfrac{1}{n}$:

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{(n^2-1)n}{n^3+1} = \lim\limits_{n\to\infty} \dfrac{n^3-n}{n^3+1} = \lim\limits_{n\to\infty} \dfrac{1-1/n^2}{1+1/n^3} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ is the divergent harmonic series, the series $\sum\limits_{n=1}^{\infty} \dfrac{n^2-1}{n^3+1}$ also diverges.

10. $\dfrac{n-1}{n^3+1} < \dfrac{n}{n^3+1} < \dfrac{n}{n^3} = \dfrac{1}{n^2}$ for $n \geq 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{n-1}{n^3+1}$ converges by direct comparison with $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$, which converges because it is a p-series with $p = 2 > 1$.

11. $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n^2-1}{n^3+1} = \sum\limits_{n=1}^{\infty} (-1)^n b_n$. Now $b_n = \dfrac{n^2-1}{n^3+1} > 0$ for $n \geq 2$, $\{b_n\}$ is decreasing for $n \geq 2$, and $\lim\limits_{n\to\infty} b_n = 0$, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n^2-1}{n^3+1}$ converges by the Alternating Series Test. By Exercise 9, $\sum\limits_{n=1}^{\infty} \dfrac{n^2-1}{n^3+1}$ diverges, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n^2-1}{n^3+1}$ is conditionally convergent.

12. $\lim\limits_{n\to\infty} |a_n| = \lim\limits_{n\to\infty} \left|(-1)^n \dfrac{n^2-1}{n^2+1}\right| = \lim\limits_{n\to\infty} \dfrac{1-1/n^2}{1+1/n^2} = 1 \neq 0$, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{n^2-1}{n^2+1}$ diverges by the Test for Divergence. $\left[\text{Note that } \lim\limits_{n\to\infty} (-1)^n \dfrac{n^2-1}{n^2+1} \text{ does not exist.}\right]$

13. $\lim\limits_{x\to\infty} \dfrac{e^x}{x^2} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{e^x}{2x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{e^x}{2} = \infty$, so $\lim\limits_{n\to\infty} \dfrac{e^n}{n^2} = \infty$. Thus, the series $\sum\limits_{n=1}^{\infty} \dfrac{e^n}{n^2}$ diverges by the Test for Divergence.

14. $\lim\limits_{n\to\infty} \sqrt[n]{|a_n|} = \lim\limits_{n\to\infty} \sqrt[n]{\dfrac{n^{2n}}{(1+n)^{3n}}} = \lim\limits_{n\to\infty} \dfrac{n^2}{(1+n)^3} = \lim\limits_{n\to\infty} \dfrac{1/n}{(1/n+1)^3} = \dfrac{0}{1} = 0 < 1$, so the series $\sum\limits_{n=1}^{\infty} \dfrac{n^{2n}}{(1+n)^{3n}}$ converges by the Root Test.

15. Let $f(x) = \dfrac{1}{x\sqrt{\ln x}}$. Then f is positive, continuous, and decreasing on $[2, \infty)$, so we can apply the Integral Test.

Since $\displaystyle\int \frac{1}{x\sqrt{\ln x}}\,dx \quad \begin{bmatrix} u = \ln x, \\ du = dx/x \end{bmatrix} = \int u^{-1/2}\,du = 2u^{1/2} + C = 2\sqrt{\ln x} + C$, we find

$\displaystyle\int_2^{\infty} \frac{dx}{x\sqrt{\ln x}} = \lim_{t\to\infty} \int_2^t \frac{dx}{x\sqrt{\ln x}} = \lim_{t\to\infty} \left[2\sqrt{\ln x}\right]_2^t = \lim_{t\to\infty} \left(2\sqrt{\ln t} - 2\sqrt{\ln 2}\right) = \infty$. Since the integral diverges, the given series $\sum\limits_{n=2}^{\infty} \dfrac{1}{n\sqrt{\ln n}}$ diverges.

16. $\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \left|\dfrac{(n+1)^4}{4^{n+1}} \cdot \dfrac{4^n}{n^4}\right| = \lim\limits_{n\to\infty} \dfrac{(n+1)^4}{4n^4} = \dfrac{1}{4} \lim\limits_{n\to\infty} \left(1 + \dfrac{1}{n}\right)^4 = \dfrac{1}{4}(1) = \dfrac{1}{4} < 1$, so the series $\sum\limits_{n=1}^{\infty} (-1)^{n-1} \dfrac{n^4}{4^n}$ is absolutely convergent (and therefore convergent) by the Ratio Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{\pi^{2n+2}}{(2n+2)!}\cdot\dfrac{(2n)!}{\pi^{2n}}\right| = \lim\limits_{n\to\infty}\dfrac{\pi^2}{(2n+2)(2n+1)} = 0 < 1$, so the series $\sum\limits_{n=0}^{\infty}(-1)^n\dfrac{\pi^{2n}}{(2n)!}$ is absolutely convergent (and therefore convergent) by the Ratio Test.

18. Let $f(x) = x^2e^{-x^3}$. Then f is continuous and positive on $[1,\infty)$, and $f'(x) = \dfrac{x(2-3x^3)}{e^{x^3}} < 0$ for $x \geq 1$, so f is decreasing on $[1,\infty)$ as well, and we can apply the Integral Test. $\int_1^\infty x^2e^{-x^3}\,dx = \lim\limits_{t\to\infty}\left[-\frac{1}{3}e^{-x^3}\right]_1^t = \frac{1}{3e}$, so the integral converges, and hence, the series $\sum\limits_{n=1}^{\infty} n^2e^{-n^3}$ converges.

19. $\sum\limits_{n=1}^{\infty}\left(\dfrac{1}{n^3}+\dfrac{1}{3^n}\right) = \sum\limits_{n=1}^{\infty}\dfrac{1}{n^3} + \sum\limits_{n=1}^{\infty}\left(\dfrac{1}{3}\right)^n$. The first series converges since it is a p-series with $p = 3 > 1$ and the second series converges since it is geometric with $|r| = \frac{1}{3} < 1$. The sum of two convergent series is convergent.

20. $\dfrac{1}{k\sqrt{k^2+1}} < \dfrac{1}{k\sqrt{k^2}} = \dfrac{1}{k^2}$, so $\sum\limits_{k=1}^{\infty}\dfrac{1}{k\sqrt{k^2+1}}$ converges by direct comparison with the convergent p-series $\sum\limits_{k=1}^{\infty}\dfrac{1}{k^2}$ ($p = 2 > 1$).

21. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{3^{n+1}(n+1)^2}{(n+1)!}\cdot\dfrac{n!}{3^nn^2}\right| = \lim\limits_{n\to\infty}\dfrac{3(n+1)^2}{(n+1)n^2} = 3\lim\limits_{n\to\infty}\dfrac{n+1}{n^2} = 0 < 1$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{3^nn^2}{n!}$ converges by the Ratio Test.

22. $\left|\dfrac{\sin 2n}{1+2^n}\right| \leq \dfrac{1}{1+2^n} < \dfrac{1}{2^n} = \left(\dfrac{1}{2}\right)^n$, so the series $\sum\limits_{n=1}^{\infty}\left|\dfrac{\sin 2n}{1+2^n}\right|$ converges by direct comparison with the geometric series $\sum\limits_{n=1}^{\infty}\left(\dfrac{1}{2}\right)^n$ with $|r| = \frac{1}{2} < 1$. Thus, the series $\sum\limits_{n=1}^{\infty}\dfrac{\sin 2n}{1+2^n}$ converges absolutely, implying convergence.

23. $a_k = \dfrac{2^{k-1}3^{k+1}}{k^k} = \dfrac{2^k2^{-1}3^k3^1}{k^k} = \dfrac{3}{2}\left(\dfrac{2\cdot 3}{k}\right)^k$. By the Root Test, $\lim\limits_{k\to\infty}\sqrt[k]{\left(\dfrac{6}{k}\right)^k} = \lim\limits_{k\to\infty}\dfrac{6}{k} = 0 < 1$, so the series $\sum\limits_{k=1}^{\infty}\left(\dfrac{6}{k}\right)^k$ converges. It follows from Theorem 8(i) in Section 11.2 that the given series, $\sum\limits_{k=1}^{\infty}\dfrac{2^{k-1}3^{k+1}}{k^k} = \sum\limits_{k=1}^{\infty}\dfrac{3}{2}\left(\dfrac{6}{k}\right)^k$, also converges.

24. Use the Limit Comparison Test with $a_n = \dfrac{\sqrt{n^4+1}}{n^3+n}$ and $b_n = \dfrac{1}{n}$:

$\lim\limits_{n\to\infty}\dfrac{a_n}{b_n} = \lim\limits_{n\to\infty}\dfrac{n\sqrt{n^4+1}}{n(n^2+1)} = \lim\limits_{n\to\infty}\dfrac{\sqrt{n^4+1}/n^2}{(n^2+1)/n^2} = \lim\limits_{n\to\infty}\dfrac{\sqrt{1+1/n^4}}{1+1/n^2} = 1 > 0$. Since $\sum\limits_{n=1}^{\infty}\dfrac{1}{n}$ is the divergent harmonic series, the series $\sum\limits_{n=1}^{\infty}\dfrac{\sqrt{n^4+1}}{n^3+n}$ also diverges.

25. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)(2n+1)}{2\cdot 5\cdot 8\cdot\cdots\cdot(3n-1)(3n+2)}\cdot\dfrac{2\cdot 5\cdot 8\cdot\cdots\cdot(3n-1)}{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}\right| = \lim\limits_{n\to\infty}\dfrac{2n+1}{3n+2}$

$$= \lim_{n\to\infty}\frac{2+1/n}{3+2/n} = \frac{2}{3} < 1,$$

so the series $\sum\limits_{n=1}^{\infty}\dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{2\cdot 5\cdot 8\cdot\cdots\cdot(3n-1)}$ converges by the Ratio Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $b_n = \dfrac{1}{\sqrt{n}-1}$ for $n \geq 2$. $\{b_n\}$ is a decreasing sequence of positive numbers and $\lim\limits_{n\to\infty} b_n = 0$, so $\sum\limits_{n=2}^{\infty} \dfrac{(-1)^{n-1}}{\sqrt{n}-1}$ converges by the Alternating Series Test.

27. Let $f(x) = \dfrac{\ln x}{\sqrt{x}}$. Then $f'(x) = \dfrac{2-\ln x}{2x^{3/2}} < 0$ when $\ln x > 2$ or $x > e^2$, so $\dfrac{\ln n}{\sqrt{n}}$ is decreasing for $n > e^2$.

By l'Hospital's Rule, $\lim\limits_{n\to\infty} \dfrac{\ln n}{\sqrt{n}} = \lim\limits_{n\to\infty} \dfrac{1/n}{1/\left(2\sqrt{n}\right)} = \lim\limits_{n\to\infty} \dfrac{2}{\sqrt{n}} = 0$, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \dfrac{\ln n}{\sqrt{n}}$ converges by the Alternating Series Test.

28. $a_k = \dfrac{\sqrt[3]{k}-1}{k(\sqrt{k}+1)} < \dfrac{\sqrt[3]{k}}{k(\sqrt{k}+1)} < \dfrac{\sqrt[3]{k}}{k\sqrt{k}} = \dfrac{k^{1/3}}{k^{3/2}} = \dfrac{1}{k^{7/6}}$, so the series $\sum\limits_{k=1}^{\infty} \dfrac{\sqrt[3]{k}-1}{k(\sqrt{k}+1)}$ converges by direct comparison with the convergent p-series $\sum\limits_{k=1}^{\infty} \dfrac{1}{k^{7/6}}$ $\left(p = \frac{7}{6} > 1\right)$.

29. $\lim\limits_{n\to\infty} |a_n| = \lim\limits_{n\to\infty} \left|(-1)^n \cos(1/n^2)\right| = \lim\limits_{n\to\infty} \left|\cos(1/n^2)\right| = \cos 0 = 1$, so the series $\sum\limits_{n=1}^{\infty} (-1)^n \cos(1/n^2)$ diverges by the Test for Divergence.

30. $\lim\limits_{k\to\infty} |a_k| = \lim\limits_{k\to\infty} \left|\dfrac{1}{2+\sin k}\right| = \lim\limits_{k\to\infty} \dfrac{1}{2+\sin k}$, which does not exist (the terms vary between $\frac{1}{3}$ and 1). Thus, the series $\sum\limits_{k=1}^{\infty} \dfrac{1}{2+\sin k}$ diverges by the Test for Divergence.

31. Using the Limit Comparison Test with $a_n = \tan\left(\dfrac{1}{n}\right)$ and $b_n = \dfrac{1}{n}$, we have

$\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{\tan(1/n)}{1/n} = \lim\limits_{x\to\infty} \dfrac{\tan(1/x)}{1/x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{\sec^2(1/x)\cdot(-1/x^2)}{-1/x^2} = \lim\limits_{x\to\infty} \sec^2(1/x) = 1^2 = 1 > 0$. Since $\sum\limits_{n=1}^{\infty} b_n$ is the divergent harmonic series, $\sum\limits_{n=1}^{\infty} a_n$ is also divergent.

32. $\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} \left(n \sin \dfrac{1}{n}\right) = \lim\limits_{n\to\infty} \dfrac{\sin(1/n)}{1/n} = \lim\limits_{x\to 0^+} \dfrac{\sin x}{x} = 1 \neq 0$, so the series $\sum\limits_{n=1}^{\infty} n\sin(1/n)$ diverges by the Test for Divergence.

33. $\dfrac{4-\cos n}{\sqrt{n}} \geq \dfrac{4-1}{\sqrt{n}} = \dfrac{3}{n^{1/2}}$ for $n \geq 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{4-\cos n}{\sqrt{n}}$ diverges by direct comparison with $\sum\limits_{n=1}^{\infty} \dfrac{3}{n^{1/2}}$, which diverges because it is a constant multiple of a p-series with $p = \frac{1}{2} \leq 1$.

34. $\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} \dfrac{8+(-1)^n n}{n} = \lim\limits_{n\to\infty} \left[\dfrac{8}{n} + (-1)^n\right]$. This limit does not exist since $8/n \to 0$ as $n \to \infty$, but $(-1)^n$ alternates between 1 and -1. Thus, the series $\sum\limits_{n=1}^{\infty} \dfrac{8+(-1)^n n}{n}$ diverges by the Test for Divergence.

35. Use the Ratio Test. $\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \left|\dfrac{(n+1)!}{e^{(n+1)^2}} \cdot \dfrac{e^{n^2}}{n!}\right| = \lim\limits_{n\to\infty} \dfrac{(n+1)n! \cdot e^{n^2}}{e^{n^2+2n+1} n!} = \lim\limits_{n\to\infty} \dfrac{n+1}{e^{2n+1}} = 0 < 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{n!}{e^{n^2}}$ converges.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\dfrac{a_{n+1}}{a_n} = \lim\limits_{n\to\infty}\left(\dfrac{n^2+2n+2}{5^{n+1}}\cdot\dfrac{5^n}{n^2+1}\right) = \lim\limits_{n\to\infty}\left(\dfrac{1+2/n+2/n^2}{1+1/n^2}\cdot\dfrac{1}{5}\right) = \dfrac{1}{5} < 1$, so $\sum\limits_{n=1}^{\infty}\dfrac{n^2+1}{5^n}$ converges by the Ratio Test.

37. $\displaystyle\int_2^{\infty}\frac{\ln x}{x^2}dx = \lim_{t\to\infty}\left[-\frac{\ln x}{x}-\frac{1}{x}\right]_1^t$ [using integration by parts] $\overset{\text{H}}{=} 1$. So $\sum\limits_{n=1}^{\infty}\dfrac{\ln n}{n^2}$ converges by the Integral Test, and since $\dfrac{k\ln k}{(k+1)^3} < \dfrac{k\ln k}{k^3} = \dfrac{\ln k}{k^2}$, the given series $\sum\limits_{k=1}^{\infty}\dfrac{k\ln k}{(k+1)^3}$ converges by the Direct Comparison Test.

38. Since $\left\{\dfrac{1}{n}\right\}$ is a decreasing sequence, $e^{1/n} \le e^{1/1} = e$ for all $n \ge 1$, and $\sum\limits_{n=1}^{\infty}\dfrac{e}{n^2}$ converges ($p = 2 > 1$), so $\sum\limits_{n=1}^{\infty}\dfrac{e^{1/n}}{n^2}$ converges by the Direct Comparison Test. (Or use the Integral Test.)

39. $\sum\limits_{n=1}^{\infty}a_n = \sum\limits_{n=1}^{\infty}(-1)^n\dfrac{1}{\cosh n} = \sum\limits_{n=1}^{\infty}(-1)^n b_n$. Now $b_n = \dfrac{1}{\cosh n} > 0$, $\{b_n\}$ is decreasing, and $\lim\limits_{n\to\infty} b_n = 0$, so the series converges by the Alternating Series Test.

Or: Write $\dfrac{1}{\cosh n} = \dfrac{2}{e^n+e^{-n}} < \dfrac{2}{e^n}$ and $\sum\limits_{n=1}^{\infty}\dfrac{1}{e^n}$ is a convergent geometric series, so $\sum\limits_{n=1}^{\infty}\dfrac{1}{\cosh n}$ is convergent by the Direct Comparison Test. So $\sum\limits_{n=1}^{\infty}(-1)^n\dfrac{1}{\cosh n}$ is absolutely convergent and therefore convergent.

40. Let $f(x) = \dfrac{\sqrt{x}}{x+5}$. Then $f(x)$ is continuous and positive on $[1,\infty)$, and since $f'(x) = \dfrac{5-x}{2\sqrt{x}\,(x+5)^2} < 0$ for $x > 5$, $f(x)$ is eventually decreasing, so we can use the Alternating Series Test. $\lim\limits_{n\to\infty}\dfrac{\sqrt{n}}{n+5} = \lim\limits_{n\to\infty}\dfrac{1}{n^{1/2}+5n^{-1/2}} = 0$, so the series $\sum\limits_{j=1}^{\infty}(-1)^j\dfrac{\sqrt{j}}{j+5}$ converges.

41. $\lim\limits_{k\to\infty}a_k = \lim\limits_{k\to\infty}\dfrac{5^k}{3^k+4^k} =$ [divide by 4^k] $\lim\limits_{k\to\infty}\dfrac{(5/4)^k}{(3/4)^k+1} = \infty$ since $\lim\limits_{k\to\infty}\left(\dfrac{3}{4}\right)^k = 0$ and $\lim\limits_{k\to\infty}\left(\dfrac{5}{4}\right)^k = \infty$.

Thus, $\sum\limits_{k=1}^{\infty}\dfrac{5^k}{3^k+4^k}$ diverges by the Test for Divergence.

42. $\lim\limits_{n\to\infty}\sqrt[n]{|a_n|} = \lim\limits_{n\to\infty}\sqrt[n]{\left|\dfrac{(n!)^n}{n^{4n}}\right|} = \lim\limits_{n\to\infty}\dfrac{n!}{n^4} = \lim\limits_{n\to\infty}\left[\dfrac{n}{n}\cdot\dfrac{n-1}{n}\cdot\dfrac{n-2}{n}\cdot\dfrac{n-3}{n}\cdot(n-4)!\right]$

$$= \lim_{n\to\infty}\left[\left(1-\frac{1}{n}\right)\left(1-\frac{2}{n}\right)\left(1-\frac{3}{n}\right)(n-4)!\right] = \infty,$$

so the series $\sum\limits_{n=1}^{\infty}\dfrac{(n!)^n}{n^{4n}}$ diverges by the Root Test.

43. $\lim\limits_{n\to\infty}\sqrt[n]{|a_n|} = \lim\limits_{n\to\infty}\left(\dfrac{n}{n+1}\right)^{n^2/n} = \lim\limits_{n\to\infty}\dfrac{1}{[(n+1)/n]^n} = \dfrac{1}{\lim\limits_{n\to\infty}(1+1/n)^n} = \dfrac{1}{e} < 1$, so the series $\sum\limits_{n=1}^{\infty}\left(\dfrac{n}{n+1}\right)^{n^2}$ converges by the Root Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. $0 \le n\cos^2 n \le n$, so $\dfrac{1}{n+n\cos^2 n} \ge \dfrac{1}{n+n} = \dfrac{1}{2n}$. Thus, $\sum\limits_{n=1}^{\infty} \dfrac{1}{n+n\cos^2 n}$ diverges by direct comparison with $\sum\limits_{n=1}^{\infty} \dfrac{1}{2n}$, which is a constant multiple of the (divergent) harmonic series.

45. $a_n = \dfrac{1}{n^{1+1/n}} = \dfrac{1}{n \cdot n^{1/n}}$, so let $b_n = \dfrac{1}{n}$ and use the Limit Comparison Test. $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{1}{n^{1/n}} = 1 > 0$ [see Exercise 4.4.63], so the series $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^{1+1/n}}$ diverges by comparison with the divergent harmonic series.

46. Note that $(\ln n)^{\ln n} = \left(e^{\ln\ln n}\right)^{\ln n} = \left(e^{\ln n}\right)^{\ln\ln n} = n^{\ln\ln n}$ and $\ln\ln n \to \infty$ as $n \to \infty$, so $\ln\ln n > 2$ for sufficiently large n. For these n we have $(\ln n)^{\ln n} > n^2$, so $\dfrac{1}{(\ln n)^{\ln n}} < \dfrac{1}{n^2}$. Since $\sum\limits_{n=2}^{\infty} \dfrac{1}{n^2}$ converges $[p = 2 > 1]$, so does $\sum\limits_{n=2}^{\infty} \dfrac{1}{(\ln n)^{\ln n}}$ by the Direct Comparison Test.

47. $\lim\limits_{n\to\infty} \sqrt[n]{|a_n|} = \lim\limits_{n\to\infty} (2^{1/n} - 1) = 1 - 1 = 0 < 1$, so the series $\sum\limits_{n=1}^{\infty} \left(\sqrt[n]{2} - 1\right)^n$ converges by the Root Test.

48. Use the Limit Comparison Test with $a_n = \sqrt[n]{2} - 1$ and $b_n = 1/n$. Then

$$\lim_{n\to\infty} \frac{a_n}{b_n} = \lim_{n\to\infty} \frac{2^{1/n} - 1}{1/n} = \lim_{x\to\infty} \frac{2^{1/x} - 1}{1/x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{2^{1/x} \cdot \ln 2 \cdot (-1/x^2)}{-1/x^2} = \lim_{x\to\infty} (2^{1/x} \cdot \ln 2) = 1 \cdot \ln 2 = \ln 2 > 0.$$

So since $\sum\limits_{n=1}^{\infty} b_n$ diverges (harmonic series), so does $\sum\limits_{n=1}^{\infty} \left(\sqrt[n]{2} - 1\right)$.

Alternate solution: $\sqrt[n]{2} - 1 = \dfrac{1}{2^{(n-1)/n} + 2^{(n-2)/n} + 2^{(n-3)/n} + \cdots + 2^{1/n} + 1}$ [rationalize the numerator] $\ge \dfrac{1}{2n}$, and since $\sum\limits_{n=1}^{\infty} \dfrac{1}{2n} = \dfrac{1}{2} \sum\limits_{n=1}^{\infty} \dfrac{1}{n}$ diverges (harmonic series), so does $\sum\limits_{n=1}^{\infty} \left(\sqrt[n]{2} - 1\right)$ by the Direct Comparison Test.

11.8 Power Series

1. A power series is a series of the form $\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots$, where x is a variable and the c_n's are constants called the coefficients of the series.

More generally, a series of the form $\sum_{n=0}^{\infty} c_n (x-a)^n = c_0 + c_1(x-a) + c_2(x-a)^2 + \cdots$ is called a power series in $(x-a)$ or a power series centered at a or a power series about a, where a is a constant.

2. (a) Given the power series $\sum_{n=0}^{\infty} c_n (x-a)^n$, the radius of convergence is:

(i) 0 if the series converges only when $x = a$

(ii) ∞ if the series converges for all x, or

(iii) a positive number R such that the series converges if $|x-a| < R$ and diverges if $|x-a| > R$.

In most cases, R can be found by using the Ratio Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The interval of convergence of a power series is the interval that consists of all values of x for which the series converges. Corresponding to the cases in part (a), the interval of convergence is: (i) the single point $\{a\}$, (ii) all real numbers; that is, the real number line $(-\infty, \infty)$, or (iii) an interval with endpoints $a - R$ and $a + R$ which can contain neither, either, or both of the endpoints. In this case, we must test the series for convergence at each endpoint to determine the interval of convergence.

3. If $a_n = \dfrac{x^n}{n}$, then $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{x^{n+1}}{n+1}\cdot\dfrac{n}{x^n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{nx}{n+1}\right| = \lim\limits_{n\to\infty}\left(\dfrac{1}{1+1/n}\,|x|\right) = |x|$. By the Ratio Test, the series $\sum\limits_{n=1}^{\infty}\dfrac{x^n}{n}$ converges when $|x| < 1$, so the radius of convergence is $R = 1$. Now we'll check the endpoints, that is, $x = \pm 1$. When $x = 1$, the series $\sum\limits_{n=1}^{\infty}\dfrac{1}{n}$ diverges since it is the harmonic series. When $x = -1$, the series $\sum\limits_{n=1}^{\infty}\dfrac{(-1)^n}{n}$ converges by the Alternating Series Test. Thus, the interval of convergence is $[-1, 1)$.

4. If $a_n = (-1)^n n x^n$, then

$\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(-1)^{n+1}(n+1)x^{n+1}}{(-1)^n\, nx^n}\right| = \lim\limits_{n\to\infty}\left|(-1)\dfrac{n+1}{n}x\right| = \lim\limits_{n\to\infty}\left[\left(1+\dfrac{1}{n}\right)|x|\right] = |x|$. By the Ratio Test, the series $\sum\limits_{n=1}^{\infty}(-1)^n n x^n$ converges when $|x| < 1$, so the radius of convergence $R = 1$. Now we'll check the endpoints, that is, $x = \pm 1$. Both series $\sum\limits_{n=1}^{\infty}(-1)^n n(\pm 1)^n = \sum\limits_{n=1}^{\infty}(\mp 1)^n n$ diverge by the Test for Divergence since $\lim\limits_{n\to\infty}|(\mp 1)^n n| = \infty$. Thus, the interval of convergence is $I = (-1, 1)$.

5. If $a_n = \sqrt{n}\,x^n$, then $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{\sqrt{n+1}\,x^{n+1}}{\sqrt{n}\,x^n}\right| = \lim\limits_{n\to\infty}\left|\sqrt{\dfrac{n+1}{n}}\,x\right| = \lim\limits_{n\to\infty}\sqrt{1+\dfrac{1}{n}}\,|x| = |x|$. By the Ratio Test, the series $\sum\limits_{n=1}^{\infty}\sqrt{n}\,x^n$ converges when $|x| < 1$, so $R = 1$. When $x = \pm 1$, both series $\sum\limits_{n=1}^{\infty}\sqrt{n}\,(\pm 1)^n$ diverge by the Test for Divergence since $\lim\limits_{n\to\infty}|\sqrt{n}\,(\pm 1)^n| = \infty$. Thus, the interval of convergence is $(-1, 1)$.

6. If $a_n = \dfrac{(-1)^n x^n}{\sqrt[3]{n}}$, then

$\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(-1)^{n+1}x^{n+1}}{\sqrt[3]{n+1}}\cdot\dfrac{\sqrt[3]{n}}{(-1)^n x^n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(-1)x\sqrt[3]{n}}{\sqrt[3]{n+1}}\right| = \lim\limits_{n\to\infty}\sqrt[3]{\dfrac{1}{1+1/n}}\,|x| = |x|$. By the Ratio Test, the series $\sum\limits_{n=1}^{\infty}\dfrac{(-1)^n x^n}{\sqrt[3]{n}}$ converges when $|x| < 1$, so $R = 1$. When $x = 1$, the series $\sum\limits_{n=1}^{\infty}\dfrac{(-1)^n}{\sqrt[3]{n}}$ converges by the Alternating Series Test. When $x = -1$, the series $\sum\limits_{n=1}^{\infty}\dfrac{1}{\sqrt[3]{n}}$ diverges since it is a p-series $\left(p = \frac{1}{3} \le 1\right)$. Thus, the interval of convergence is $(-1, 1]$.

7. If $a_n = \dfrac{n}{5^n}\,x^n$, then $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(n+1)\,x^{n+1}}{5^{n+1}}\cdot\dfrac{5^n}{n\,x^n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{n+1}{5n}\,x\right| = \lim\limits_{n\to\infty}\left(\dfrac{1}{5}+\dfrac{1}{5n}\right)|x| = \dfrac{|x|}{5}$. By the Ratio Test, the series $\sum\limits_{n=1}^{\infty}\dfrac{n}{5^n}\,x^n$ converges when $\dfrac{|x|}{5} < 1 \quad\Leftrightarrow\quad |x| < 5$, so $R = 5$. When $x = \pm 5$, both series

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\sum_{n=1}^{\infty} \frac{n(\pm 5)^n}{5^n} = \sum_{n=1}^{\infty} (\pm 1)^n n$ diverge by the Test for Divergence since $\lim_{n\to\infty} |(\pm 1)^n n| = \infty$. Thus, the interval of convergence is $(-5, 5)$.

8. If $a_n = \frac{5^n}{n} x^n$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{5^{n+1}x^{n+1}}{(n+1)} \cdot \frac{n}{5^n x^n}\right| = \lim_{n\to\infty} \left|\frac{5n}{n+1} x\right| = \lim_{n\to\infty} \left(\frac{5}{1+1/n} |x|\right) = 5|x|$. By the Ratio Test, the series $\sum_{n=2}^{\infty} \frac{5^n}{n} x^n$ converges when $5|x| < 1 \quad \Leftrightarrow \quad |x| < \frac{1}{5}$, so $R = \frac{1}{5}$. When $x = \frac{1}{5}$, the series $\sum_{n=2}^{\infty} \frac{1}{n}$ diverges since it is the (partial) harmonic series. When $x = -\frac{1}{5}$, the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n}$ converges by the Alternating Series Test. Thus, the interval of convergence is $\left[-\frac{1}{5}, \frac{1}{5}\right)$.

9. If $a_n = \frac{x^n}{n\,3^n}$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{x^{n+1}}{(n+1)\,3^{n+1}} \cdot \frac{n\,3^n}{x^n}\right| = \lim_{n\to\infty} \left|\frac{n}{3(n+1)} x\right| = \lim_{n\to\infty} \left(\frac{1}{3+3/n} |x|\right) = \frac{|x|}{3}$. By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{x^n}{n\,3^n}$ converges when $\frac{|x|}{3} < 1 \quad \Leftrightarrow \quad |x| < 3$, so $R = 3$. When $x = 3$, the series $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges since it is the harmonic series. When $x = -3$, the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges by the Alternating Series Test. Thus, the interval of convergence is $[-3, 3)$.

10. If $a_n = \frac{n}{n+1} x^n$, then

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(n+1)\,x^{n+1}}{(n+1)+1} \cdot \frac{n+1}{n\,x^n}\right| = \lim_{n\to\infty} \left|\frac{n^2+2n+1}{n^2+2n} x\right| = \lim_{n\to\infty} \left(\frac{1+2/n+1/n^2}{1+2/n} |x|\right) = |x|.$$

By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{n}{n+1} x^n$ converges when $|x| < 1$, so $R = 1$. When $x = \pm 1$, both series $\sum_{n=1}^{\infty} \frac{n(\pm 1)^n}{n+1}$ diverge by the Test for Divergence since $\lim_{n\to\infty} \frac{n}{n+1} = 1 \neq 0$ and $\lim_{n\to\infty} \frac{n(-1)^n}{n+1}$ does not exist. Thus, the interval of convergence is $(-1, 1)$.

11. If $a_n = \frac{x^n}{2n-1}$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{x^{n+1}}{2n+1} \cdot \frac{2n-1}{x^n}\right| = \lim_{n\to\infty} \left(\frac{2n-1}{2n+1} |x|\right) = \lim_{n\to\infty} \left(\frac{2-1/n}{2+1/n} |x|\right) = |x|$. By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{x^n}{2n-1}$ converges when $|x| < 1$, so $R = 1$. When $x = 1$, the series $\sum_{n=1}^{\infty} \frac{1}{2n-1}$ diverges by direct comparison with $\sum_{n=1}^{\infty} \frac{1}{2n}$ since $\frac{1}{2n-1} > \frac{1}{2n}$ and $\frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{n}$ diverges since it is a constant multiple of the harmonic series. When $x = -1$, the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{2n-1}$ converges by the Alternating Series Test. Thus, the interval of convergence is $[-1, 1)$.

12. If $a_n = \frac{(-1)^n x^n}{n^2}$, then

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-1)^{n+1}x^{n+1}}{(n+1)^2} \cdot \frac{n^2}{(-1)^n x^n}\right| = \lim_{n\to\infty} \left|\frac{(-1)xn^2}{(n+1)^2}\right| = \lim_{n\to\infty} \left[\left(\frac{n}{n+1}\right)^2 |x|\right] = 1^2 \cdot |x| = |x|.$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{(-1)^n x^n}{n^2}$ converges when $|x| < 1$, so $R = 1$. When $x = 1$, the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$ converges by the Alternating Series Test. When $x = -1$, the series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges since it is a p-series with $p = 2 > 1$. Thus, the interval of convergence is $[-1, 1]$.

13. If $a_n = \frac{x^n}{n!}$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{x^{n+1}}{(n+1)!} \cdot \frac{n!}{x^n}\right| = \lim_{n\to\infty} \left|\frac{x}{n+1}\right| = |x| \lim_{n\to\infty} \frac{1}{n+1} = |x| \cdot 0 = 0 < 1$ for *all* real x. So, by the Ratio Test, $R = \infty$ and $I = (-\infty, \infty)$.

14. Here the Root Test is easier. If $a_n = n^n x^n$, then $\lim_{n\to\infty} \sqrt[n]{|a_n|} = \lim_{n\to\infty} n\,|x| = \infty$ if $x \neq 0$, so $R = 0$ and $I = \{0\}$.

15. If $a_n = \frac{x^n}{n^4\, 4^n}$, then

$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{x^{n+1}}{(n+1)^4\, 4^{n+1}} \cdot \frac{n^4\, 4^n}{x^n}\right| = \lim_{n\to\infty} \left|\frac{n^4}{(n+1)^4} \cdot \frac{x}{4}\right| = \lim_{n\to\infty} \left(\frac{n}{n+1}\right)^4 \frac{|x|}{4} = 1^4 \cdot \frac{|x|}{4} = \frac{|x|}{4}$. By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{x^n}{n^4\, 4^n}$ converges when $\frac{|x|}{4} < 1 \quad \Leftrightarrow \quad |x| < 4$, so $R = 4$. When $x = 4$, the series $\sum_{n=1}^{\infty} \frac{1}{n^4}$ converges since it is a p-series ($p = 4 > 1$). When $x = -4$, the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^4}$ converges by the Alternating Series Test. Thus, the interval of convergence is $[-4, 4]$.

16. If $a_n = 2^n n^2 x^n$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{2^{n+1}(n+1)^2 x^{n+1}}{2^n n^2 x^n}\right| = \lim_{n\to\infty} 2\left(\frac{n+1}{n}\right)^2 |x| = 2\,|x|$. By the Ratio Test, the series $\sum_{n=1}^{\infty} 2^n n^2 x^n$ converges when $2\,|x| < 1 \quad \Leftrightarrow \quad |x| < \frac{1}{2}$, so $R = \frac{1}{2}$. When $x = \pm\frac{1}{2}$, both series $\sum_{n=1}^{\infty} 2^n n^2 \left(\pm\frac{1}{2}\right)^n = \sum_{n=1}^{\infty} (\pm 1)^n n^2$ diverge by the Test for Divergence since $\lim_{n\to\infty} \left|(\pm 1)^n n^2\right| = \infty$. Thus, the interval of convergence is $\left(-\frac{1}{2}, \frac{1}{2}\right)$.

17. If $a_n = \frac{(-1)^n\, 4^n}{\sqrt{n}}\, x^n$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-1)^{n+1}\, 4^{n+1}\, x^{n+1}}{\sqrt{n+1}} \cdot \frac{\sqrt{n}}{(-1)^n\, 4^n\, x^n}\right| = \lim_{n\to\infty} \sqrt{\frac{n}{n+1}} \cdot 4\,|x| = 4\,|x|$. By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{(-1)^n\, 4^n}{\sqrt{n}}\, x^n$ converges when $4\,|x| < 1 \quad \Leftrightarrow \quad |x| < \frac{1}{4}$, so $R = \frac{1}{4}$. When $x = \frac{1}{4}$, the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$ converges by the Alternating Series Test. When $x = -\frac{1}{4}$, the series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ diverges since it is a p-series $\left(p = \frac{1}{2} \leq 1\right)$. Thus, the interval of convergence is $\left(-\frac{1}{4}, \frac{1}{4}\right]$.

18. If $a_n = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n5^n}\, x^n$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-1)^n x^{n+1}}{(n+1)\, 5^{n+1}} \cdot \frac{n5^n}{(-1)^{n-1} x^n}\right| = \lim_{n\to\infty} \left(\frac{n}{n+1}\right) \frac{|x|}{5} = 1 \cdot \frac{|x|}{5} = \frac{|x|}{5}$. By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n5^n}\, x^n$ converges when $\frac{|x|}{5} < 1 \quad \Leftrightarrow \quad |x| < 5$, so $R = 5$. When $x = 5$, the series

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$ converges by the Alternating Series Test. When $x = -5$, the series $\sum_{n=1}^{\infty} \frac{-1}{n}$ diverges since it is a constant multiple of the harmonic series. Thus, the interval of convergence is $(-5, 5]$.

19. If $a_n = \frac{n}{2^n(n^2+1)}\, x^n$, then

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(n+1)x^{n+1}}{2^{n+1}(n^2+2n+2)} \cdot \frac{2^n(n^2+1)}{n\,x^n}\right| = \lim_{n\to\infty} \frac{n^3+n^2+n+1}{n^3+2n^2+2n} \cdot \frac{|x|}{2}$$

$$= \lim_{n\to\infty} \frac{1+1/n+1/n^2+1/n^3}{1+2/n+2/n^2} \cdot \frac{|x|}{2} = \frac{|x|}{2}$$

By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{n}{2^n(n^2+1)}\, x^n$ converges when $\frac{|x|}{2} < 1 \quad \Leftrightarrow \quad |x| < 2$, so $R = 2$. When $x = 2$, the series $\sum_{n=1}^{\infty} \frac{n}{n^2+1}$ diverges by the Limit Comparison Test with $b_n = \frac{1}{n}$. When $x = -2$, the series $\sum_{n=1}^{\infty} \frac{(-1)^n n}{n^2+1}$ converges by the Alternating Series Test. Thus, the interval of convergence is $[-2, 2)$.

20. If $a_n = \frac{x^{2n}}{n!}$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{x^{2n+2}}{(n+1)!} \cdot \frac{n!}{x^{2n}}\right| = \lim_{n\to\infty} \frac{|x^2|}{n+1} = 0 < 1$ for *all* real x. So, by the Ratio Test, $R = \infty$ and $I = (-\infty, \infty)$.

21. If $a_n = \frac{(x-2)^n}{n^2+1}$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(x-2)^{n+1}}{(n+1)^2+1} \cdot \frac{n^2+1}{(x-2)^n}\right| = |x-2| \lim_{n\to\infty} \frac{n^2+1}{(n+1)^2+1} = |x-2|$. By the Ratio Test, the series $\sum_{n=0}^{\infty} \frac{(x-2)^n}{n^2+1}$ converges when $|x-2| < 1 \quad [R = 1] \quad \Leftrightarrow \quad -1 < x-2 < 1 \quad \Leftrightarrow \quad 1 < x < 3$. When $x = 1$, the series $\sum_{n=0}^{\infty} (-1)^n \frac{1}{n^2+1}$ converges by the Alternating Series Test; when $x = 3$, the series $\sum_{n=0}^{\infty} \frac{1}{n^2+1}$ converges by direct comparison with the p-series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ $[p = 2 > 1]$. Thus, the interval of convergence is $I = [1, 3]$.

22. If $a_n = \frac{(-1)^n}{(2n-1)2^n}\,(x-1)^n$, then

$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-1)^{n+1}(x-1)^{n+1}}{(2n+1)\,2^{n+1}} \cdot \frac{(2n-1)\,2^n}{(-1)^n(x-1)^n}\right| = \lim_{n\to\infty} \frac{2n-1}{2n+1} \cdot \frac{|x-1|}{2} = \frac{|x-1|}{2}$. By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{(2n-1)\,2^n}(x-1)^n$ converges when $\frac{|x-1|}{2} < 1 \quad \Leftrightarrow \quad |x-1| < 2 \quad [R = 2] \quad \Leftrightarrow \quad -2 < x-1 < 2 \quad \Leftrightarrow$ $-1 < x < 3$. When $x = 3$, the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{2n-1}$ converges by the Alternating Series Test. When $x = -1$, the series $\sum_{n=1}^{\infty} \frac{1}{2n-1}$ diverges by the Limit Comparison Test with $b_n = \frac{1}{n}$. Thus, the interval of convergence is $(-1, 3]$.

23. If $a_n = \frac{(x+2)^n}{2^n \ln n}$, then $\lim_{n\to\infty} \left|\frac{(x+2)^{n+1}}{2^{n+1}\ln(n+1)} \cdot \frac{2^n \ln n}{(x+2)^n}\right| = \lim_{n\to\infty} \frac{\ln n}{\ln(n+1)} \cdot \frac{|x+2|}{2} = \frac{|x+2|}{2}$ since $\lim_{n\to\infty} \frac{\ln n}{\ln(n+1)} = \lim_{x\to\infty} \frac{\ln x}{\ln(x+1)} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{1/x}{1/(x+1)} = \lim_{x\to\infty} \frac{x+1}{x} = \lim_{x\to\infty} \left(1 + \frac{1}{x}\right) = 1$. By the Ratio Test, the series

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\sum_{n=2}^{\infty} \frac{(x+2)^n}{2^n \ln n}$ converges when $\frac{|x+2|}{2} < 1 \quad \Leftrightarrow \quad |x+2| < 2 \quad [R=2] \quad \Leftrightarrow \quad -2 < x+2 < 2 \quad \Leftrightarrow \quad -4 < x < 0.$

When $x = -4$, the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{\ln n}$ converges by the Alternating Series Test. When $x = 0$, the series $\sum_{n=2}^{\infty} \frac{1}{\ln n}$ diverges by the Limit Comparison Test with $b_n = \frac{1}{n}$ (or by direct comparison with the harmonic series). Thus, the interval of convergence is $[-4, 0)$.

24. If $a_n = \frac{\sqrt{n}}{8^n}(x+6)^n$, then

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{\sqrt{n+1}\,(x+6)^{n+1}}{8^{n+1}} \cdot \frac{8^n}{\sqrt{n}(x+6)^n}\right| = \lim_{n\to\infty} \sqrt{\frac{n+1}{n}} \cdot \frac{|x+6|}{8}$$
$$= \lim_{n\to\infty} \sqrt{1+\frac{1}{n}} \cdot \frac{|x+6|}{8} = \frac{|x+6|}{8}$$

By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{\sqrt{n}}{8^n}(x+6)^n$ converges when $\frac{|x+6|}{8} < 1 \quad \Leftrightarrow \quad |x+6| < 8 \quad [R=8] \quad \Leftrightarrow$ $-8 < x+6 < 8 \quad \Leftrightarrow \quad -14 < x < 2$. When $x = 2$, the series $\sum_{n=1}^{\infty} \sqrt{n}$ diverges by the Test for Divergence since $\lim_{n\to\infty} |a_n| = \lim_{n\to\infty} \sqrt{n} = \infty > 0$. Similarly, when $x = -14$, the series $\sum_{n=1}^{\infty} (-1)^n \sqrt{n}$ diverges. Thus, the interval of convergence is $(-14, 2)$.

25. If $a_n = \frac{(x-2)^n}{n^n}$, then $\lim_{n\to\infty} \sqrt[n]{|a_n|} = \lim_{n\to\infty} \frac{|x-2|}{n} = 0$, so the series converges for all x (by the Root Test). $R = \infty$ and $I = (-\infty, \infty)$.

26. If $a_n = \frac{(2x-1)^n}{5^n\sqrt{n}}$, then

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(2x-1)^{n+1}}{5^{n+1}\sqrt{n+1}} \cdot \frac{5^n\sqrt{n}}{(2x-1)^n}\right| = \lim_{n\to\infty} \frac{|2x-1|}{5}\sqrt{\frac{n}{n+1}} = \lim_{n\to\infty} \frac{|2x-1|}{5}\sqrt{\frac{1}{1+1/n}} = \frac{|2x-1|}{5}.$$

By the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{(2x-1)^n}{5^n\sqrt{n}}$ converges when $\frac{|2x-1|}{5} < 1 \quad \Leftrightarrow \quad |2x-1| < 5 \quad \Leftrightarrow \quad \left|x - \frac{1}{2}\right| < \frac{5}{2} \quad \Leftrightarrow$ $-\frac{5}{2} < x - \frac{1}{2} < \frac{5}{2} \quad \Leftrightarrow \quad -2 < x < 3$, so $R = \frac{5}{2}$. When $x = 3$, the series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ is a divergent p-series $\left(p = \frac{1}{2} \le 1\right)$. When $x = -2$, the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$ converges by the Alternating Series Test. Thus, the interval of convergence is $I = [-2, 3)$.

27. If $a_n = \frac{\ln n}{n} x^n$, then

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{\ln(n+1)\,x^{n+1}}{n+1} \cdot \frac{n}{(\ln n)\,x^n}\right| = \lim_{n\to\infty} \left|\frac{n}{n+1}\frac{\ln(n+1)}{\ln n}\,x\right|$$
$$= 1 \cdot 1 \cdot |x| = |x|$$

since $\lim_{n\to\infty} \frac{\ln(n+1)}{\ln n} \overset{\text{H}}{=} \lim_{n\to\infty} \frac{1/(n+1)}{1/n} = \lim_{n\to\infty} \frac{n}{n+1} = \lim_{n\to\infty} \frac{1}{1+1/n} = 1$. By the Ratio Test, the series $\sum_{n=4}^{\infty} \frac{\ln n}{n} x^n$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

converges when $|x| < 1$, so $R = 1$. When $x = 1$, the series $\sum_{n=4}^{\infty} \frac{\ln n}{n}$ diverges by direct comparison with the (partial) harmonic series $\sum_{n=4}^{\infty} \frac{1}{n}$. When $x = -1$, the series $\sum_{n=4}^{\infty} \frac{(-1)^n \ln n}{n}$ converges by the Alternating Series Test. Thus, the interval of convergence is $[-1, 1)$.

28. If $a_n = \frac{(-1)^n}{n \ln n} x^n$, then

$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-1)^{n+1} x^{n+1}}{(n+1)\ln(n+1)} \cdot \frac{n \ln n}{(-1)^n x^n}\right| = \lim_{n\to\infty} \left[\frac{n}{n+1} \frac{\ln n}{\ln(n+1)} |x|\right] = 1 \cdot 1 \cdot |x| = |x|$ since

$\lim_{n\to\infty} \frac{\ln n}{\ln(n+1)} \overset{\text{H}}{=} \lim_{n\to\infty} \frac{1/n}{1/(n+1)} = \lim_{n\to\infty} \frac{n+1}{n} = \lim_{n\to\infty} \left(1 + \frac{1}{n}\right) = 1$. By the Ratio Test, the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n \ln n} x^n$ converges when $|x| < 1$, so $R = 1$. When $x = 1$, the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n \ln n}$ converges by the Alternating Series Test. When $x = -1$, the series $\sum_{n=2}^{\infty} \frac{(-1)^n (-1)^n}{n \ln n} = \sum_{n=2}^{\infty} \frac{1}{n \ln n}$ diverges by the Integral Test. Thus, the interval of convergence is $(-1, 1]$.

29. $a_n = \frac{n}{b^n}(x-a)^n$, where $b > 0$.

$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \frac{(n+1)|x-a|^{n+1}}{b^{n+1}} \cdot \frac{b^n}{n|x-a|^n} = \lim_{n\to\infty} \left(1 + \frac{1}{n}\right) \frac{|x-a|}{b} = \frac{|x-a|}{b}$.

By the Ratio Test, the series converges when $\frac{|x-a|}{b} < 1 \quad \Leftrightarrow \quad |x-a| < b \quad [\text{so } R = b] \quad \Leftrightarrow \quad -b < x - a < b \quad \Leftrightarrow$ $a - b < x < a + b$. When $|x-a| = b$, $\lim_{n\to\infty} |a_n| = \lim_{n\to\infty} n = \infty$, so the series diverges. Thus, $I = (a-b, a+b)$.

30. $a_n = \frac{b^n}{\ln n}(x-a)^n$, where $b > 0$.

$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{b^{n+1}(x-a)^{n+1}}{\ln(n+1)} \cdot \frac{\ln n}{b^n (x-a)^n}\right| = \lim_{n\to\infty} \frac{\ln n}{\ln(n+1)} \cdot b|x-a| = b|x-a|$ since

$\lim_{n\to\infty} \frac{\ln n}{\ln(n+1)} = \lim_{n\to\infty} \frac{\ln x}{\ln(x+1)} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{1/x}{1/(x+1)} = \lim_{x\to\infty} \frac{x+1}{x} \overset{\text{H}}{=} \lim_{x\to\infty} \frac{1}{1} = 1$. By the Ratio Test, the series $\sum_{n=2}^{\infty} \frac{b^n}{\ln n}(x-a)^n$ converges when $b|x-a| < 1 \quad \Leftrightarrow \quad |x-a| < \frac{1}{b} \quad \Leftrightarrow \quad -\frac{1}{b} < x - a < \frac{1}{b} \quad \Leftrightarrow \quad a - \frac{1}{b} < x < a + \frac{1}{b}$, so $R = \frac{1}{b}$. When $x = a + \frac{1}{b}$, the series $\sum_{n=2}^{\infty} \frac{1}{\ln n}$ diverges by direct comparison with the divergent p-series $\sum_{n=2}^{\infty} \frac{1}{n}$ since $\frac{1}{\ln n} > \frac{1}{n}$ for $n \geq 2$. When $x = a - \frac{1}{b}$, the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{\ln n}$ converges by the Alternating Series Test. Thus, the interval of convergence is $I = \left[a - \frac{1}{b}, a + \frac{1}{b}\right)$.

31. If $a_n = n!\,(2x-1)^n$, then $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(n+1)!\,(2x-1)^{n+1}}{n!(2x-1)^n}\right| = \lim_{n\to\infty} (n+1)|2x-1| \to \infty$ as $n \to \infty$ for all $x \neq \frac{1}{2}$. Since the series diverges for all $x \neq \frac{1}{2}$, $R = 0$ and $I = \left\{\frac{1}{2}\right\}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. $a_n = \dfrac{n^2 x^n}{2 \cdot 4 \cdot 6 \cdot \cdots \cdot (2n)} = \dfrac{n^2 x^n}{2^n n!} = \dfrac{n x^n}{2^n (n-1)!}$, so

$\lim\limits_{n\to\infty} \left| \dfrac{a_{n+1}}{a_n} \right| = \lim\limits_{n\to\infty} \dfrac{(n+1)\,|x|^{n+1}}{2^{n+1} n!} \cdot \dfrac{2^n (n-1)!}{n\,|x|^n} = \lim\limits_{n\to\infty} \dfrac{n+1}{n^2} \dfrac{|x|}{2} = 0$. Thus, by the Ratio Test, the series converges for *all* real x and we have $R = \infty$ and $I = (-\infty, \infty)$.

33. If $a_n = \dfrac{(5x-4)^n}{n^3}$, then

$$\lim_{n\to\infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n\to\infty} \left| \frac{(5x-4)^{n+1}}{(n+1)^3} \cdot \frac{n^3}{(5x-4)^n} \right| = \lim_{n\to\infty} |5x-4| \left(\frac{n}{n+1} \right)^3 = \lim_{n\to\infty} |5x-4| \left(\frac{1}{1+1/n} \right)^3$$
$$= |5x-4| \cdot 1 = |5x-4|$$

By the Ratio Test, $\sum\limits_{n=1}^{\infty} \dfrac{(5x-4)^n}{n^3}$ converges when $|5x-4| < 1 \quad \Leftrightarrow \quad \left|x - \frac{4}{5}\right| < \frac{1}{5} \quad \Leftrightarrow \quad -\frac{1}{5} < x - \frac{4}{5} < \frac{1}{5} \quad \Leftrightarrow$ $\frac{3}{5} < x < 1$, so $R = \frac{1}{5}$. When $x = 1$, the series $\sum\limits_{n=1}^{\infty} \dfrac{1}{n^3}$ is a convergent p-series ($p = 3 > 1$). When $x = \frac{3}{5}$, the series $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^n}{n^3}$ converges by the Alternating Series Test. Thus, the interval of convergence is $I = \left[\frac{3}{5}, 1\right]$.

34. If $a_n = \dfrac{x^{2n}}{n\,(\ln n)^2}$, then $\lim\limits_{n\to\infty} \left| \dfrac{a_{n+1}}{a_n} \right| = \lim\limits_{n\to\infty} \left| \dfrac{x^{2n+2}}{(n+1)[\ln(n+1)]^2} \cdot \dfrac{n\,(\ln n)^2}{x^{2n}} \right| = \left|x^2\right| \lim\limits_{n\to\infty} \dfrac{n\,(\ln n)^2}{(n+1)[\ln(n+1)]^2} = x^2$.

By the Ratio Test, the series $\sum\limits_{n=2}^{\infty} \dfrac{x^{2n}}{n\,(\ln n)^2}$ converges when $x^2 < 1 \quad \Leftrightarrow \quad |x| < 1$, so $R = 1$. When $x = \pm 1$, $x^{2n} = 1$, the series $\sum\limits_{n=2}^{\infty} \dfrac{1}{n\,(\ln n)^2}$ converges by the Integral Test (see Exercise 11.3.31). Thus, the interval of convergence is $I = [-1, 1]$.

35. If $a_n = \dfrac{x^n}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}$, then

$\lim\limits_{n\to\infty} \left| \dfrac{a_{n+1}}{a_n} \right| = \lim\limits_{n\to\infty} \left| \dfrac{x^{n+1}}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)(2n+1)} \cdot \dfrac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{x^n} \right| = \lim\limits_{n\to\infty} \dfrac{|x|}{2n+1} = 0 < 1$. Thus, by the Ratio Test, the series $\sum\limits_{n=1}^{\infty} \dfrac{x^n}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}$ converges for *all* real x and we have $R = \infty$ and $I = (-\infty, \infty)$.

36. If $a_n = \dfrac{n!\,x^n}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}$, then

$\lim\limits_{n\to\infty} \left| \dfrac{a_{n+1}}{a_n} \right| = \lim\limits_{n\to\infty} \left| \dfrac{(n+1)!\,x^{n+1}}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)(2n+1)} \cdot \dfrac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{n!\,x^n} \right| = \lim\limits_{n\to\infty} \dfrac{(n+1)\,|x|}{2n+1} = \frac{1}{2}\,|x|$.

By the Ratio Test, the series $\sum\limits_{n=1}^{\infty} a_n$ converges when $\frac{1}{2}\,|x| < 1 \quad \Rightarrow \quad |x| < 2$, so $R = 2$. When $x = \pm 2$,

$|a_n| = \dfrac{n!\,2^n}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)} = \dfrac{[1 \cdot 2 \cdot 3 \cdot \cdots \cdot n]\,2^n}{[1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)]} = \dfrac{2 \cdot 4 \cdot 6 \cdot \cdots \cdot 2n}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)} > 1$, so both endpoint series diverge by the Test for Divergence. Thus, the interval of convergence is $I = (-2, 2)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. (a) We are given that the power series $\sum_{n=0}^{\infty} c_n x^n$ is convergent for $x = 4$. So by Theorem 4, it must converge for at least $-4 < x \le 4$. In particular, it converges when $x = -2$; that is, $\sum_{n=0}^{\infty} c_n(-2)^n$ is convergent.

(b) It does not follow that $\sum_{n=0}^{\infty} c_n(-4)^n$ is necessarily convergent. [See the comments after Theorem 4 about convergence at the endpoint of an interval. An example is $c_n = (-1)^n/(n4^n)$.]

38. We are given that the power series $\sum_{n=0}^{\infty} c_n x^n$ is convergent for $x = -4$ and divergent when $x = 6$. So by Theorem 4 it converges for at least $-4 \le x < 4$ and diverges for at least $x \ge 6$ and $x < -6$. Therefore:

(a) It converges when $x = 1$; that is, $\sum c_n$ is convergent.

(b) It diverges when $x = 8$; that is, $\sum c_n 8^n$ is divergent.

(c) It converges when $x = -3$; that is, $\sum c_n(-3^n)$ is convergent.

(d) It diverges when $x = -9$; that is, $\sum c_n(-9)^n = \sum(-1)^n c_n 9^n$ is divergent.

39. If $a_n = \dfrac{(n!)^k}{(kn)!}x^n$, then

$$\begin{aligned}
\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| &= \lim_{n\to\infty}\frac{[(n+1)!]^k\,(kn)!}{(n!)^k\,[k(n+1)]!}\,|x| = \lim_{n\to\infty}\frac{(n+1)^k}{(kn+k)(kn+k-1)\cdots(kn+2)(kn+1)}\,|x| \\
&= \lim_{n\to\infty}\left[\frac{(n+1)}{(kn+1)}\frac{(n+1)}{(kn+2)}\cdots\frac{(n+1)}{(kn+k)}\right]|x| \\
&= \lim_{n\to\infty}\left[\frac{n+1}{kn+1}\right]\lim_{n\to\infty}\left[\frac{n+1}{kn+2}\right]\cdots\lim_{n\to\infty}\left[\frac{n+1}{kn+k}\right]|x| \\
&= \left(\frac{1}{k}\right)^k|x| < 1 \quad\Leftrightarrow\quad |x| < k^k \text{ for convergence, and the radius of convergence is } R = k^k.
\end{aligned}$$

40. (a) Note that the four intervals in parts (a)–(d) have midpoint $m = \frac{1}{2}(p+q)$ and radius of convergence $r = \frac{1}{2}(q-p)$. We also know that the power series $\sum\limits_{n=0}^{\infty} x^n$ has interval of convergence $(-1, 1)$. To change the radius of convergence to r, we can change x^n to $\left(\dfrac{x}{r}\right)^n$. To shift the midpoint of the interval of convergence, we can replace x with $x - m$. Thus, a power series whose interval of convergence is (p, q) is $\sum\limits_{n=0}^{\infty}\left(\dfrac{x-m}{r}\right)^n$, where $m = \frac{1}{2}(p+q)$ and $r = \frac{1}{2}(q-p)$.

(b) Similar to Example 2, we know that $\sum\limits_{n=1}^{\infty}\dfrac{x^n}{n}$ has interval of convergence $[-1, 1)$. By introducing the factor $(-1)^n$ in a_n, the interval of convergence changes to $(-1, 1]$. Now change the midpoint and radius as in part (a) to get $\sum\limits_{n=1}^{\infty}(-1)^n\dfrac{1}{n}\left(\dfrac{x-m}{r}\right)^n$ as a power series whose interval of convergence is $(p, q]$.

(c) As in part (b), $\sum\limits_{n=1}^{\infty}\dfrac{1}{n}\left(\dfrac{x-m}{r}\right)^n$ is a power series whose interval of convergence is $[p, q)$.

(d) If we increase the exponent on n (to say, $n = 2$), in the power series in part (c), then when $x = q$, the power series $\sum\limits_{n=1}^{\infty}\dfrac{1}{n^2}\left(\dfrac{x-m}{r}\right)^n$ will converge by comparison to the p-series with $p = 2 > 1$, and the interval of convergence will be $[p, q]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. No. If a power series is centered at a, its interval of convergence is symmetric about a. If a power series has an infinite radius of convergence, then its interval of convergence must be $(-\infty, \infty)$, not $[0, \infty)$.

42. The partial sums of the series $\sum_{n=0}^{\infty} x^n$ definitely do not converge to $f(x) = 1/(1-x)$ for $x \geq 1$, since f is undefined at $x = 1$ and negative on $(1, \infty)$, while all the partial sums are positive on this interval. The partial sums also fail to converge to f for $x \leq -1$, since $0 < f(x) < 1$ on this interval, while the partial sums are either larger than 1 or less than 0. The partial sums seem to converge to f on $(-1, 1)$. This graphical evidence is consistent with what we know about geometric series: convergence for $|x| < 1$, divergence for $|x| \geq 1$.

43. We use the Root Test on the series $\sum c_n x^n$. We need $\lim\limits_{n\to\infty} \sqrt[n]{|c_n x^n|} = |x| \lim\limits_{n\to\infty} \sqrt[n]{|c_n|} = c\,|x| < 1$ for convergence, or $|x| < 1/c$, so $R = 1/c$.

44. Suppose $c_n \neq 0$. Applying the Ratio Test to the series $\sum c_n (x-a)^n$, we find that

$L = \lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \left|\dfrac{c_{n+1}(x-a)^{n+1}}{c_n(x-a)^n}\right| = \lim\limits_{n\to\infty} \dfrac{|x-a|}{|c_n/c_{n+1}|} (\star) = \dfrac{|x-a|}{\lim\limits_{n\to\infty} |c_n/c_{n+1}|}$ (if $\lim\limits_{n\to\infty} |c_n/c_{n+1}| \neq 0$), so the series converges when $\dfrac{|x-a|}{\lim\limits_{n\to\infty} |c_n/c_{n+1}|} < 1 \quad \Leftrightarrow \quad |x-a| < \lim\limits_{n\to\infty} \left|\dfrac{c_n}{c_{n+1}}\right|$. Thus, $R = \lim\limits_{n\to\infty} \left|\dfrac{c_n}{c_{n+1}}\right|$. If $\lim\limits_{n\to\infty} \left|\dfrac{c_n}{c_{n+1}}\right| = 0$ and $|x-a| \neq 0$, then $(\star)$ shows that $L = \infty$ and so the series diverges, and hence, $R = 0$. Thus, in all cases, $R = \lim\limits_{n\to\infty} \left|\dfrac{c_n}{c_{n+1}}\right|$.

45. For $2 < x < 3$, $\sum c_n x^n$ diverges and $\sum d_n x^n$ converges. By Exercise 11.2.89, $\sum (c_n + d_n)\, x^n$ diverges. Since both series converge for $|x| < 2$, the radius of convergence of $\sum (c_n + d_n)\, x^n$ is 2.

46. Since $\sum c_n x^n$ converges whenever $|x| < R$, $\sum c_n x^{2n} = \sum c_n \left(x^2\right)^n$ converges whenever $\left|x^2\right| < R \quad \Leftrightarrow \quad |x| < \sqrt{R}$, so the second series has radius of convergence $\sqrt{R}$.

11.9 Representations of Functions as Power Series

1. If $f(x) = \sum\limits_{n=0}^{\infty} c_n x^n$ has radius of convergence 10, then $f'(x) = \sum\limits_{n=1}^{\infty} n c_n x^{n-1}$ also has radius of convergence 10 by Theorem 2.

2. If $f(x) = \sum\limits_{n=0}^{\infty} b_n x^n$ converges on $(-2, 2)$, then $\int f(x)\, dx = C + \sum\limits_{n=0}^{\infty} \dfrac{b_n}{n+1} x^{n+1}$ has the same radius of convergence (by Theorem 2), but may not have the same interval of convergence—it may happen that the integrated series converges at an endpoint (or both endpoints).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. Our goal is to write the function in the form $\dfrac{1}{1-r}$, and then use Equation 1 to represent the function as a sum of a power series. $f(x) = \dfrac{1}{1+x} = \dfrac{1}{1-(-x)} = \sum\limits_{n=0}^{\infty}(-x)^n = \sum\limits_{n=0}^{\infty}(-1)^n x^n$ with $|-x| < 1 \quad \Leftrightarrow \quad |x| < 1$, so $R = 1$ and $I = (-1, 1)$.

4. $f(x) = \dfrac{x}{1+x} = x\left(\dfrac{1}{1-(-x)}\right) = x\sum\limits_{n=0}^{\infty}(-x)^n$, or, equivalently, $\sum\limits_{n=0}^{\infty}(-1)^n x^{n+1}$. The series converges when $|-x| < 1 \quad \Leftrightarrow \quad |x| < 1$, so $R = 1$ and $I = (-1, 1)$.

5. $f(x) = \dfrac{1}{1-x^2} = \sum\limits_{n=0}^{\infty}(x^2)^n = \sum\limits_{n=0}^{\infty} x^{2n}$. The series converges when $\left|x^2\right| < 1 \quad \Leftrightarrow \quad |x| < 1$, so $R = 1$ and $I = (-1, 1)$.

6. $f(x) = \dfrac{5}{1-4x^2} = 5\left(\dfrac{1}{1-4x^2}\right) = 5\sum\limits_{n=0}^{\infty}(4x^2)^n = 5\sum\limits_{n=0}^{\infty} 4^n x^{2n}$. The series converges when $\left|4x^2\right| < 1 \quad \Leftrightarrow$ $|x|^2 < \frac{1}{4} \quad \Leftrightarrow \quad |x| < \frac{1}{2}$, so $R = \frac{1}{2}$ and $I = \left(-\frac{1}{2}, \frac{1}{2}\right)$.

7. $f(x) = \dfrac{2}{3-x} = \dfrac{2}{3}\left(\dfrac{1}{1-x/3}\right) = \dfrac{2}{3}\sum\limits_{n=0}^{\infty}\left(\dfrac{x}{3}\right)^n$ or, equivalently, $2\sum\limits_{n=0}^{\infty}\dfrac{1}{3^{n+1}}x^n$. The series converges when $\left|\dfrac{x}{3}\right| < 1$, that is, when $|x| < 3$, so $R = 3$ and $I = (-3, 3)$.

8. $f(x) = \dfrac{4}{2x+3} = \dfrac{4}{3}\left(\dfrac{1}{1+2x/3}\right) = \dfrac{4}{3}\left(\dfrac{1}{1-(-2x/3)}\right) = \dfrac{4}{3}\sum\limits_{n=0}^{\infty}\left(-\dfrac{2x}{3}\right)^n$ or, equivalently, $\sum\limits_{n=0}^{\infty}(-1)^n\dfrac{2^{n+2}}{3^{n+1}}x^n$.

The series converges when $\left|-\dfrac{2x}{3}\right| < 1$, that is, when $|x| < \frac{3}{2}$, so $R = \frac{3}{2}$ and $I = \left(-\frac{3}{2}, \frac{3}{2}\right)$.

9. $f(x) = \dfrac{x^2}{x^4+16} = \dfrac{x^2}{16}\left(\dfrac{1}{1+x^4/16}\right) = \dfrac{x^2}{16}\left(\dfrac{1}{1-[-(x/2)]^4}\right) = \dfrac{x^2}{16}\sum\limits_{n=0}^{\infty}\left[-\left(\dfrac{x}{2}\right)^4\right]^n$ or, equivalently, $\sum\limits_{n=0}^{\infty}\dfrac{(-1)^n x^{4n+2}}{2^{4n+4}}$.

The series converges when $\left|-\left(\dfrac{x}{2}\right)^4\right| < 1 \quad \Rightarrow \quad \left|\dfrac{x}{2}\right| < 1 \quad \Rightarrow \quad |x| < 2$, so $R = 2$ and $I = (-2, 2)$.

10. $f(x) = \dfrac{x}{2x^2+1} = x\left(\dfrac{1}{1-(-2x^2)}\right) = x\sum\limits_{n=0}^{\infty}(-2x^2)^n$ or, equivalently, $\sum\limits_{n=0}^{\infty}(-1)^n 2^n x^{2n+1}$. The series converges when $\left|-2x^2\right| < 1 \quad \Rightarrow \quad \left|x^2\right| < \frac{1}{2} \quad \Rightarrow \quad |x| < \dfrac{1}{\sqrt{2}}$, so $R = \dfrac{1}{\sqrt{2}}$ and $I = \left(-\dfrac{1}{\sqrt{2}}, \dfrac{1}{\sqrt{2}}\right)$.

11. $f(x) = \dfrac{x-1}{x+2} = \dfrac{x+2-3}{x+2} = 1 - \dfrac{3}{x+2} = 1 - \dfrac{3/2}{x/2+1} = 1 - \dfrac{3}{2}\cdot\dfrac{1}{1-(-x/2)}$

$$= 1 - \frac{3}{2}\sum_{n=0}^{\infty}\left(-\frac{x}{2}\right)^n = 1 - \frac{3}{2} - \frac{3}{2}\sum_{n=1}^{\infty}\left(-\frac{x}{2}\right)^n = -\frac{1}{2} - \sum_{n=1}^{\infty}\frac{(-1)^n\,3x^n}{2^{n+1}}.$$

The geometric series $\sum\limits_{n=0}^{\infty}\left(-\dfrac{x}{2}\right)^n$ converges when $\left|-\dfrac{x}{2}\right| < 1 \quad \Leftrightarrow \quad |x| < 2$, so $R = 2$ and $I = (-2, 2)$.

Alternatively, you could write $f(x) = 1 - 3\left(\dfrac{1}{x+2}\right)$ and use the series for $\dfrac{1}{x+2}$ found in Example 2.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $f(x) = \dfrac{x+a}{x^2+a^2} \quad [a>0] \quad = \dfrac{x}{a^2}\left[\dfrac{1}{1-(-x^2/a^2)}\right] + \dfrac{a}{a^2}\left[\dfrac{1}{1-(-x^2/a^2)}\right]$

$$= \frac{x}{a^2}\sum_{n=0}^{\infty}\left(-\frac{x^2}{a^2}\right)^n + \frac{1}{a}\sum_{n=0}^{\infty}\left(-\frac{x^2}{a^2}\right)^n = \sum_{n=0}^{\infty}\frac{(-1)^n x^{2n+1}}{a^{2n+1}} + \sum_{n=0}^{\infty}\frac{(-1)^n x^{2n}}{a^{2n+1}}.$$

The geometric series $\sum\limits_{n=0}^{\infty}\left(-\dfrac{x^2}{a^2}\right)^n$ converges when $\left|-\dfrac{x^2}{a^2}\right| < 1 \quad \Leftrightarrow \quad |x| < a$, so $R = a$ and $I = (-a, a)$.

13. $f(x) = \dfrac{2x-4}{x^2-4x+3} = \dfrac{2x-4}{(x-1)(x-3)} = \dfrac{A}{x-1} + \dfrac{B}{x-3} \quad \Rightarrow \quad 2x-4 = A(x-3)+B(x-1)$. Let $x=1$ to get $-2=-2A \quad \Leftrightarrow \quad A=1$ and $x=3$ to get $2=2B \quad \Leftrightarrow \quad B=1$. Thus,

$$\frac{2x-4}{x^2-4x+3} = \frac{1}{x-1} + \frac{1}{x-3} = \frac{-1}{1-x} + \frac{1}{-3}\left[\frac{1}{1-(x/3)}\right] = -\sum_{n=0}^{\infty}x^n - \frac{1}{3}\sum_{n=0}^{\infty}\left(\frac{x}{3}\right)^n = \sum_{n=0}^{\infty}\left(-1-\frac{1}{3^{n+1}}\right)x^n.$$

We represented f as the sum of two geometric series; the first converges for $x \in (-1,1)$ and the second converges for $x \in (-3,3)$. Thus, the sum converges for $x \in (-1,1) = I$.

14. $f(x) = \dfrac{2x+3}{x^2+3x+2} = \dfrac{2x+3}{(x+1)(x+2)} = \dfrac{A}{x+1} + \dfrac{B}{x+2} \quad \Rightarrow \quad 2x+3 = A(x+2)+B(x+1)$. Let $x=-1$ to get $1=A$ and $x=-2$ to get $-1=-B \quad \Leftrightarrow \quad B=1$. Thus,

$$\frac{2x+3}{x^2+3x+2} = \frac{1}{x+1} + \frac{1}{x+2} = \frac{1}{1-(-x)} + \frac{1}{2}\left[\frac{1}{1-(-x/2)}\right]$$
$$= \sum_{n=0}^{\infty}(-x)^n + \frac{1}{2}\sum_{n=0}^{\infty}\left(-\frac{x}{2}\right)^n = \sum_{n=0}^{\infty}\left[(-1)^n\left(1+\frac{1}{2^{n+1}}\right)\right]x^n$$

We represented f as the sum of two geometric series; the first converges for $x \in (-1,1)$ and the second converges for $x \in (-2,2)$. Thus, the sum converges for $x \in (-1,1) = I$.

15. (a) $f(x) = \dfrac{1}{(1+x)^2} = \dfrac{d}{dx}\left(\dfrac{-1}{1+x}\right) = -\dfrac{d}{dx}\left[\sum\limits_{n=0}^{\infty}(-1)^n x^n\right]$ [from Exercise 3]

$= \sum\limits_{n=1}^{\infty}(-1)^{n+1}nx^{n-1}$ [from Theorem 2(i)] $= \sum\limits_{n=0}^{\infty}(-1)^n(n+1)x^n$ with $R=1$.

In the last step, note that we *decreased* the initial value of the summation variable n by 1, and then *increased* each occurrence of n in the term by 1 [also note that $(-1)^{n+2} = (-1)^n$].

(b) $f(x) = \dfrac{1}{(1+x)^3} = -\dfrac{1}{2}\dfrac{d}{dx}\left[\dfrac{1}{(1+x)^2}\right] = -\dfrac{1}{2}\dfrac{d}{dx}\left[\sum\limits_{n=0}^{\infty}(-1)^n(n+1)x^n\right]$ [from part (a)]

$= -\frac{1}{2}\sum\limits_{n=1}^{\infty}(-1)^n(n+1)nx^{n-1} = \frac{1}{2}\sum\limits_{n=0}^{\infty}(-1)^n(n+2)(n+1)x^n$ with $R=1$.

(c) $f(x) = \dfrac{x^2}{(1+x)^3} = x^2 \cdot \dfrac{1}{(1+x)^3} = x^2 \cdot \dfrac{1}{2}\sum\limits_{n=0}^{\infty}(-1)^n(n+2)(n+1)x^n$ [from part (b)]

$= \dfrac{1}{2}\sum\limits_{n=0}^{\infty}(-1)^n(n+2)(n+1)x^{n+2}$

To write the power series with x^n rather than x^{n+2}, we will *decrease* each occurrence of n in the term by 2 and *increase* the initial value of the summation variable by 2. This gives us $\dfrac{1}{2}\sum\limits_{n=2}^{\infty}(-1)^n(n)(n-1)x^n$ with $R=1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. (a) $\displaystyle\int \frac{1}{1-x}\,dx = -\ln(1-x) + C$ and

$$\int \frac{1}{1-x}\,dx = \int (1 + x + x^2 + \cdots)\,dx = \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \cdots\right) + C = \sum_{n=1}^{\infty} \frac{x^n}{n} + C \text{ for } |x| < 1.$$

So $-\ln(1-x) = \displaystyle\sum_{n=1}^{\infty} \frac{x^n}{n} + C$ and letting $x = 0$ gives $0 = C$. Thus, $f(x) = \ln(1-x) = -\displaystyle\sum_{n=1}^{\infty} \frac{x^n}{n}$ with $R = 1$.

(b) $f(x) = x\ln(1-x) = -x\displaystyle\sum_{n=1}^{\infty} \frac{x^n}{n} = -\sum_{n=1}^{\infty} \frac{x^{n+1}}{n}$.

(c) Letting $x = \dfrac{1}{2}$ gives $\ln\dfrac{1}{2} = -\displaystyle\sum_{n=1}^{\infty} \frac{(1/2)^n}{n} \quad\Rightarrow\quad \ln 1 - \ln 2 = -\sum_{n=1}^{\infty} \frac{1^n}{n2^n} \quad\Rightarrow\quad \ln 2 = \sum_{n=1}^{\infty} \frac{1}{n2^n}$.

17. We know that $\dfrac{1}{1+4x} = \dfrac{1}{1-(-4x)} = \displaystyle\sum_{n=0}^{\infty} (-4x)^n$. Differentiating, we get

$\dfrac{-4}{(1+4x)^2} = \displaystyle\sum_{n=1}^{\infty} (-4)^n n x^{n-1} = \sum_{n=0}^{\infty} (-4)^{n+1}(n+1)x^n$, so

$$f(x) = \frac{x}{(1+4x)^2} = \frac{-x}{4} \cdot \frac{-4}{(1+4x)^2} = \frac{-x}{4} \sum_{n=0}^{\infty} (-4)^{n+1}(n+1)x^n = \sum_{n=0}^{\infty} (-1)^n 4^n (n+1)x^{n+1}$$

for $|-4x| < 1 \quad\Leftrightarrow\quad |x| < \frac{1}{4}$, so $R = \frac{1}{4}$.

18. $\dfrac{1}{2-x} = \dfrac{1}{2(1-x/2)} = \dfrac{1}{2}\displaystyle\sum_{n=0}^{\infty} \left(\frac{x}{2}\right)^n = \sum_{n=0}^{\infty} \frac{1}{2^{n+1}} x^n$. Now $\dfrac{d}{dx}\left(\dfrac{1}{2-x}\right) = \dfrac{d}{dx}\left(\displaystyle\sum_{n=0}^{\infty} \frac{1}{2^{n+1}} x^n\right) \quad\Rightarrow$

$\dfrac{1}{(2-x)^2} = \displaystyle\sum_{n=1}^{\infty} \frac{1}{2^{n+1}} n x^{n-1}$ and $\dfrac{d}{dx}\left(\dfrac{1}{(2-x)^2}\right) = \dfrac{d}{dx}\left(\displaystyle\sum_{n=1}^{\infty} \frac{1}{2^{n+1}} n x^{n-1}\right) \quad\Rightarrow$

$$\frac{2}{(2-x)^3} = \sum_{n=2}^{\infty} \frac{1}{2^{n+1}} n(n-1)x^{n-2} = \sum_{n=0}^{\infty} \frac{(n+2)(n+1)}{2^{n+3}} x^n.$$

Thus, $f(x) = \left(\dfrac{x}{2-x}\right)^3 = \dfrac{x^3}{(2-x)^3} = \dfrac{x^3}{2} \cdot \dfrac{2}{(2-x)^3} = \dfrac{x^3}{2} \displaystyle\sum_{n=0}^{\infty} \frac{(n+2)(n+1)}{2^{n+3}} x^n = \sum_{n=0}^{\infty} \frac{(n+2)(n+1)}{2^{n+4}} x^{n+3}$

for $\left|\dfrac{x}{2}\right| < 1 \quad\Leftrightarrow\quad |x| < 2$, so $R = 2$.

19. By Example 4, $\dfrac{1}{(1-x)^2} = \displaystyle\sum_{n=0}^{\infty} (n+1)x^n$. Thus,

$$\begin{aligned} f(x) = \frac{1+x}{(1-x)^2} &= \frac{1}{(1-x)^2} + \frac{x}{(1-x)^2} = \sum_{n=0}^{\infty} (n+1)x^n + \sum_{n=0}^{\infty} (n+1)x^{n+1} \\ &= \sum_{n=0}^{\infty} (n+1)x^n + \sum_{n=1}^{\infty} n x^n \qquad \text{[make the starting values equal]} \\ &= 1 + \sum_{n=1}^{\infty} [(n+1) + n]x^n = 1 + \sum_{n=1}^{\infty} (2n+1)x^n = \sum_{n=0}^{\infty} (2n+1)x^n \ \text{ with } R = 1. \end{aligned}$$

20. By Example 4, $\dfrac{1}{(1-x)^2} = \displaystyle\sum_{n=0}^{\infty} (n+1)x^n$, so

$\dfrac{d}{dx}\left(\dfrac{1}{(1-x)^2}\right) = \dfrac{d}{dx}\left(\displaystyle\sum_{n=0}^{\infty} (n+1)x^n\right) \quad\Rightarrow\quad \dfrac{2}{(1-x)^3} = \displaystyle\sum_{n=1}^{\infty} (n+1)n x^{n-1}$. Thus,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$f(x) = \frac{x^2+x}{(1-x)^3} = \frac{x^2}{(1-x)^3} + \frac{x}{(1-x)^3} = \frac{x^2}{2}\cdot\frac{2}{(1-x)^3} + \frac{x}{2}\cdot\frac{2}{(1-x)^3}$$

$$= \frac{x^2}{2}\sum_{n=1}^{\infty}(n+1)nx^{n-1} + \frac{x}{2}\sum_{n=1}^{\infty}(n+1)nx^{n-1} = \sum_{n=1}^{\infty}\frac{(n+1)n}{2}x^{n+1} + \sum_{n=1}^{\infty}\frac{(n+1)n}{2}x^n$$

$$= \sum_{n=2}^{\infty}\frac{n(n-1)}{2}x^n + \sum_{n=1}^{\infty}\frac{(n+1)n}{2}x^n \qquad \text{[make the exponents on } x \text{ equal by changing an index]}$$

$$= \sum_{n=2}^{\infty}\frac{n^2-n}{2}x^n + x + \sum_{n=2}^{\infty}\frac{n^2+n}{2}x^n \qquad \text{[make the starting values equal]}$$

$$= x + \sum_{n=2}^{\infty} n^2x^n = \sum_{n=1}^{\infty} n^2x^n \ \text{ with } R = 1.$$

21. $f(x) = \ln(5-x) = -\int \frac{dx}{5-x} = -\frac{1}{5}\int \frac{dx}{1-x/5} = -\frac{1}{5}\int\left[\sum_{n=0}^{\infty}\left(\frac{x}{5}\right)^n\right]dx$

$$= C - \frac{1}{5}\sum_{n=0}^{\infty}\frac{x^{n+1}}{5^n(n+1)} = C - \sum_{n=1}^{\infty}\frac{x^n}{n5^n}$$

Putting $x = 0$, we get $C = \ln 5$. The series converges for $|x/5| < 1 \quad\Leftrightarrow\quad |x| < 5$, so $R = 5$.

22. $f(x) = x^2\tan^{-1}(x^3) = x^2\sum_{n=0}^{\infty}(-1)^n\frac{(x^3)^{2n+1}}{2n+1}$ [by Example 7] $= \sum_{n=0}^{\infty}(-1)^n\frac{x^{6n+3+2}}{2n+1} = \sum_{n=0}^{\infty}(-1)^n\frac{x^{6n+5}}{2n+1}$ for $\left|x^3\right| < 1 \quad\Leftrightarrow\quad |x| < 1$, so $R = 1$.

23. $f(x) = \frac{x^2}{x^2+1} = x^2\left(\frac{1}{1-(-x^2)}\right) = x^2\sum_{n=0}^{\infty}(-x^2)^n = \sum_{n=0}^{\infty}(-1)^n x^{2n+2}$. This series converges when $\left|-x^2\right| < 1 \quad\Leftrightarrow\quad x^2 < 1 \quad\Leftrightarrow\quad |x| < 1$, so $R = 1$. The partial sums are $s_1 = x^2$, $s_2 = s_1 - x^4$, $s_3 = s_2 + x^6$, $s_4 = s_3 - x^8$, $s_5 = s_4 + x^{10}, \ldots$. Note that s_1 corresponds to the first term of the infinite sum, regardless of the value of the summation variable and the value of the exponent. As n increases, $s_n(x)$ approximates f better on the interval of convergence, which is $(-1, 1)$.

24. From Example 5, we have $\ln(1+x) = \sum_{n=1}^{\infty}(-1)^{n-1}\frac{x^n}{n}$ with $|x| < 1$, so $f(x) = \ln(1+x^4) = \sum_{n=1}^{\infty}(-1)^{n-1}\frac{x^{4n}}{n}$ with $\left|x^4\right| < 1 \quad\Leftrightarrow\quad |x| < 1 \quad [R = 1]$. The partial sums are $s_1 = x^4$, $s_2 = s_1 - \frac{1}{2}x^8$, $s_3 = s_2 + \frac{1}{3}x^{12}$, $s_4 = s_3 - \frac{1}{4}x^{16}$, $s_5 = s_4 + \frac{1}{5}x^{20}, \ldots$. Note that s_1 corresponds to the first term of the infinite sum, regardless of the value of the summation variable and the value of the exponent. As n increases, $s_n(x)$ approximates f better on the interval of convergence, which is $[-1, 1]$. (When $x = \pm 1$, the series is the convergent alternating harmonic series.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $f(x) = \ln\left(\dfrac{1+x}{1-x}\right) = \ln(1+x) - \ln(1-x) = \displaystyle\int \frac{dx}{1+x} + \int \frac{dx}{1-x} = \int \frac{dx}{1-(-x)} + \int \frac{dx}{1-x}$

$$= \int \left[\sum_{n=0}^{\infty} (-1)^n x^n + \sum_{n=0}^{\infty} x^n\right] dx = \int [(1 - x + x^2 - x^3 + x^4 - \cdots) + (1 + x + x^2 + x^3 + x^4 + \cdots)]\, dx$$

$$= \int (2 + 2x^2 + 2x^4 + \cdots)\, dx = \int \sum_{n=0}^{\infty} 2x^{2n}\, dx = C + \sum_{n=0}^{\infty} \frac{2x^{2n+1}}{2n+1}$$

But $f(0) = \ln \frac{1}{1} = 0$, so $C = 0$ and we have $f(x) = \displaystyle\sum_{n=0}^{\infty} \frac{2x^{2n+1}}{2n+1}$ with $R = 1$. If $x = \pm 1$, then $f(x) = \pm 2 \displaystyle\sum_{n=0}^{\infty} \frac{1}{2n+1}$, which both diverge by the Limit Comparison Test with $b_n = \dfrac{1}{n}$.

The partial sums are $s_1 = \dfrac{2x}{1}$, $s_2 = s_1 + \dfrac{2x^3}{3}$, $s_3 = s_2 + \dfrac{2x^5}{5}, \ldots$.

As n increases, $s_n(x)$ approximates f better on the interval of convergence, which is $(-1, 1)$.

26. $f(x) = \tan^{-1}(2x) = 2\displaystyle\int \frac{dx}{1+4x^2} = 2\int \sum_{n=0}^{\infty} (-1)^n \left(4x^2\right)^n\, dx = 2\int \sum_{n=0}^{\infty} (-1)^n 4^n x^{2n}\, dx$

$$= C + 2\sum_{n=0}^{\infty} \frac{(-1)^n 4^n x^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} \frac{(-1)^n 2^{2n+1} x^{2n+1}}{2n+1} \qquad [f(0) = \tan^{-1} 0 = 0, \text{ so } C = 0]$$

The series converges when $\left|4x^2\right| < 1 \quad \Leftrightarrow \quad |x| < \frac{1}{2}$, so $R = \frac{1}{2}$. If $x = \pm\frac{1}{2}$, then $f(x) = \displaystyle\sum_{n=0}^{\infty} (-1)^n \frac{1}{2n+1}$ and $f(x) = \displaystyle\sum_{n=0}^{\infty} (-1)^{n+1} \frac{1}{2n+1}$, respectively. Both series converge by the Alternating Series Test. The partial sums are $s_1 = \dfrac{2x}{1}$, $s_2 = s_1 - \dfrac{2^3x^3}{3}$, $s_3 = s_2 + \dfrac{2^5x^5}{5}, \ldots$.

As n increases, $s_n(x)$ approximates f better on the interval of convergence, which is $\left[-\frac{1}{2}, \frac{1}{2}\right]$.

27. $\dfrac{t}{1-t^8} = t \cdot \dfrac{1}{1-t^8} = t \displaystyle\sum_{n=0}^{\infty} (t^8)^n = \sum_{n=0}^{\infty} t^{8n+1} \quad \Rightarrow \quad \int \frac{t}{1-t^8}\, dt = C + \sum_{n=0}^{\infty} \frac{t^{8n+2}}{8n+2}$. The series for $\dfrac{1}{1-t^8}$ converges when $\left|t^8\right| < 1 \quad \Leftrightarrow \quad |t| < 1$, so $R = 1$ for that series and also the series for $t/(1-t^8)$. By Theorem 2, the series for $\displaystyle\int \frac{t}{1-t^8}\, dt$ also has $R = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $\dfrac{t}{1+t^3} = t \cdot \dfrac{1}{1-(-t^3)} = t \sum\limits_{n=0}^{\infty} (-t^3)^n = \sum\limits_{n=0}^{\infty} (-1)^n t^{3n+1} \quad \Rightarrow \quad \displaystyle\int \frac{t}{1+t^3}\,dt = C + \sum_{n=0}^{\infty} (-1)^n \frac{t^{3n+2}}{3n+2}$. The series for $\dfrac{1}{1+t^3}$ converges when $\left|-t^3\right| < 1 \quad \Leftrightarrow \quad |t| < 1$, so $R = 1$ for that series and also for the series $\dfrac{t}{1+t^3}$. By Theorem 2, the series for $\displaystyle\int \frac{t}{1+t^3}\,dt$ also has $R = 1$.

29. From Example 5, $\ln(1+x) = \sum\limits_{n=1}^{\infty} (-1)^{n-1} \dfrac{x^n}{n}$ for $|x| < 1$, so $x^2 \ln(1+x) = \sum\limits_{n=1}^{\infty} (-1)^{n-1} \dfrac{x^{n+2}}{n}$ and $\displaystyle\int x^2 \ln(1+x)\,dx = C + \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{n+3}}{n(n+3)}$. $R = 1$ for the series for $\ln(1+x)$, so $R = 1$ for the series representing $x^2 \ln(1+x)$ as well. By Theorem 2, the series for $\displaystyle\int x^2 \ln(1+x)\,dx$ also has $R = 1$.

30. From Example 6, $\tan^{-1} x = \sum\limits_{n=0}^{\infty} (-1)^n \dfrac{x^{2n+1}}{2n+1}$ for $|x| < 1$, so $\dfrac{\tan^{-1} x}{x} = \sum\limits_{n=0}^{\infty} (-1)^n \dfrac{x^{2n}}{2n+1}$ and $\displaystyle\int \frac{\tan^{-1} x}{x}\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)^2}$. $R = 1$ for the series for $\tan^{-1} x$, so $R = 1$ for the series representing $\dfrac{\tan^{-1} x}{x}$ as well. By Theorem 2, the series for $\displaystyle\int \frac{\tan^{-1} x}{x}\,dx$ also has $R = 1$.

31. $\dfrac{x}{1+x^3} = x\left[\dfrac{1}{1-(-x^3)}\right] = x \sum\limits_{n=0}^{\infty} (-x^3)^n = \sum\limits_{n=0}^{\infty} (-1)^n x^{3n+1} \quad \Rightarrow$

$\displaystyle\int \frac{x}{1+x^3}\,dx = \int \sum_{n=0}^{\infty} (-1)^n x^{3n+1}\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{3n+2}}{3n+2}$. Thus,

$$I = \int_0^{0.3} \frac{x}{1+x^3}\,dx = \left[\frac{x^2}{2} - \frac{x^5}{5} + \frac{x^8}{8} - \frac{x^{11}}{11} + \cdots\right]_0^{0.3} = \frac{(0.3)^2}{2} - \frac{(0.3)^5}{5} + \frac{(0.3)^8}{8} - \frac{(0.3)^{11}}{11} + \cdots.$$

The series is alternating, so if we use the first three terms, the error is at most $(0.3)^{11}/11 \approx 1.6 \times 10^{-7}$.

So $I \approx (0.3)^2/2 - (0.3)^5/5 + (0.3)^8/8 \approx 0.044\,522$ to six decimal places.

32. We substitute $x/2$ for x in Example 6, and find that

$$\begin{aligned}\int \arctan\frac{x}{2}\,dx &= \int \sum_{n=0}^{\infty} (-1)^n \frac{(x/2)^{2n+1}}{2n+1}\,dx = \int \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2^{2n+1}(2n+1)}\,dx \\ &= C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+2}}{2^{2n+1}(2n+1)(2n+2)}\end{aligned}$$

Thus,

$$\begin{aligned}I = \int_0^{1/2} \arctan\frac{x}{2}\,dx &= \left[\frac{x^2}{2(1)(2)} - \frac{x^4}{2^3(3)(4)} + \frac{x^6}{2^5(5)(6)} - \frac{x^8}{2^7(7)(8)} + \frac{x^{10}}{2^9(9)(10)} - \cdots\right]_0^{1/2} \\ &= \frac{1}{2^3(1)(2)} - \frac{1}{2^7(3)(4)} + \frac{1}{2^{11}(5)(6)} - \frac{1}{2^{15}(7)(8)} + \frac{1}{2^{19}(9)(10)} - \cdots\end{aligned}$$

The series is alternating, so if we use four terms, the error is at most $1/(2^{19} \cdot 90) \approx 2.1 \times 10^{-8}$. So

$I \approx \dfrac{1}{16} - \dfrac{1}{1536} + \dfrac{1}{61{,}440} - \dfrac{1}{1{,}835{,}008} \approx 0.061\,865$ to six decimal places.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Remark: The sum of the first three terms gives us the same answer to six decimal places, but the error is at most $1/1{,}835{,}008 \approx 5.5 \times 10^{-7}$, slightly too large to guarantee the desired accuracy.

33. We substitute x^2 for x in Example 5, and find that

$$\int x\ln(1+x^2)\,dx = \int x \sum_{n=1}^{\infty}(-1)^{n-1}\frac{(x^2)^n}{n}\,dx = \int \sum_{n=1}^{\infty}(-1)^{n-1}\frac{x^{2n+1}}{n}\,dx = C + \sum_{n=1}^{\infty}(-1)^{n-1}\frac{x^{2n+2}}{n(2n+2)}$$

Thus,

$$I \approx \int_0^{0.2} x\ln(1+x^2)\,dx = \left[\frac{x^4}{1(4)} - \frac{x^6}{2(6)} + \frac{x^8}{3(8)} - \frac{x^{10}}{4(10)} + \cdots\right]_0^{0.2} = \frac{(0.2)^4}{4} - \frac{(0.2)^6}{12} + \frac{(0.2)^8}{24} - \frac{(0.2)^{10}}{40} + \cdots$$

The series is alternating, so if we use two terms, the error is at most $(0.2)^8/24 \approx 1.1 \times 10^{-7}$. So

$I \approx \dfrac{(0.2)^4}{4} - \dfrac{(0.2)^6}{12} \approx 0.000\,395$ to six decimal places.

34. $$\int_0^{0.3} \frac{x^2}{1+x^4}\,dx = \int_0^{0.3} x^2 \sum_{n=0}^{\infty}(-1)^n x^{4n}\,dx = \sum_{n=0}^{\infty}\left[\frac{(-1)^n x^{4n+3}}{4n+3}\right]_0^{0.3} = \sum_{n=0}^{\infty}\frac{(-1)^n\,3^{4n+3}}{(4n+3)10^{4n+3}}$$
$$= \frac{3^3}{3\times 10^3} - \frac{3^7}{7\times 10^7} + \frac{3^{11}}{11\times 10^{11}} - \cdots$$

The series is alternating, so if we use only two terms, the error is at most $\dfrac{3^{11}}{11\times 10^{11}} \approx 0.000\,000\,16$. So, to six decimal places, $\displaystyle\int_0^{0.3}\frac{x^2}{1+x^4}\,dx \approx \frac{3^3}{3\times 10^3} - \frac{3^7}{7\times 10^7} \approx 0.008\,969$.

35. By Example 6, $\arctan x = x - \dfrac{x^3}{3} + \dfrac{x^5}{5} - \dfrac{x^7}{7} + \cdots$, so $\arctan 0.2 = 0.2 - \dfrac{(0.2)^3}{3} + \dfrac{(0.2)^5}{5} - \dfrac{(0.2)^7}{7} + \cdots$.

The series is alternating, so if we use three terms, the error is at most $\dfrac{(0.2)^7}{7} \approx 0.000\,002$.

Thus, to five decimal places, $\arctan 0.2 \approx 0.2 - \dfrac{(0.2)^3}{3} + \dfrac{(0.2)^5}{5} \approx 0.197\,40$.

36. By Example 5, $\ln(1+x) = x - \dfrac{x^2}{2} + \dfrac{x^3}{3} - \dfrac{x^4}{4} + \cdots$, so $\ln 1.1 = \ln(1+0.1) = 0.1 - \dfrac{(0.1)^2}{2} + \dfrac{(0.1)^3}{3} - \dfrac{(0.1)^4}{4} + \cdots$.

The series is alternating, so if we use four terms, the error is at most $\dfrac{(0.1)^5}{5} = 0.000\,002$. Thus, to four decimal places,

$\ln 1.1 \approx 0.1 - \dfrac{(0.1)^2}{2} + \dfrac{(0.1)^3}{3} - \dfrac{(0.1)^4}{4} \approx 0.0953$.

37. (a) $f(x) = \displaystyle\sum_{n=0}^{\infty}\frac{x^n}{n!} \quad\Rightarrow\quad f'(x) = \sum_{n=1}^{\infty}\frac{nx^{n-1}}{n!} = \sum_{n=1}^{\infty}\frac{x^{n-1}}{(n-1)!} = \sum_{n=0}^{\infty}\frac{x^n}{n!} = f(x)$

(b) By Theorem 9.4.2, the only solution to the differential equation $df(x)/dx = f(x)$ is $f(x) = Ke^x$, but $f(0) = 1$, so $K = 1$ and $f(x) = e^x$.

Or: We could solve the equation $df(x)/dx = f(x)$ as a separable differential equation.

38. $f(x) = \displaystyle\sum_{n=0}^{\infty}\frac{(-1)^n x^{2n}}{(2n)!} \quad\Rightarrow\quad f'(x) = \sum_{n=1}^{\infty}\frac{(-1)^n 2nx^{2n-1}}{(2n)!}$ [the first term disappears], so

$$f''(x) = \sum_{n=1}^{\infty}\frac{(-1)^n(2n)(2n-1)x^{2n-2}}{(2n)!} = \sum_{n=1}^{\infty}\frac{(-1)^n x^{2(n-1)}}{[2(n-1)]!} = \sum_{n=0}^{\infty}\frac{(-1)^{n+1}x^{2n}}{(2n)!} \quad \text{[substituting } n+1 \text{ for } n\text{]}$$
$$= -\sum_{n=0}^{\infty}\frac{(-1)^n x^{2n}}{(2n)!} = -f(x) \quad\Rightarrow\quad f''(x) + f(x) = 0.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. (a) $J_0(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n}(n!)^2}$, $J_0'(x) = \sum_{n=1}^{\infty} \frac{(-1)^n 2nx^{2n-1}}{2^{2n}(n!)^2}$, and $J_0''(x) = \sum_{n=1}^{\infty} \frac{(-1)^n 2n(2n-1)x^{2n-2}}{2^{2n}(n!)^2}$, so

$$\begin{aligned}
x^2 J_0''(x) + xJ_0'(x) + x^2 J_0(x) &= \sum_{n=1}^{\infty} \frac{(-1)^n 2n(2n-1)x^{2n}}{2^{2n}(n!)^2} + \sum_{n=1}^{\infty} \frac{(-1)^n 2nx^{2n}}{2^{2n}(n!)^2} + \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+2}}{2^{2n}(n!)^2} \\
&= \sum_{n=1}^{\infty} \frac{(-1)^n 2n(2n-1)x^{2n}}{2^{2n}(n!)^2} + \sum_{n=1}^{\infty} \frac{(-1)^n 2nx^{2n}}{2^{2n}(n!)^2} + \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^{2n}}{2^{2n-2}\,[(n-1)!]^2} \\
&= \sum_{n=1}^{\infty} \frac{(-1)^n 2n(2n-1)x^{2n}}{2^{2n}(n!)^2} + \sum_{n=1}^{\infty} \frac{(-1)^n 2nx^{2n}}{2^{2n}(n!)^2} + \sum_{n=1}^{\infty} \frac{(-1)^n(-1)^{-1}2^2n^2x^{2n}}{2^{2n}(n!)^2} \\
&= \sum_{n=1}^{\infty} (-1)^n \left[\frac{2n(2n-1)+2n-2^2n^2}{2^{2n}(n!)^2}\right] x^{2n} \\
&= \sum_{n=1}^{\infty} (-1)^n \left[\frac{4n^2-2n+2n-4n^2}{2^{2n}(n!)^2}\right] x^{2n} = 0
\end{aligned}$$

(b) $$\int_0^1 J_0(x)\,dx = \int_0^1 \left[\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n}(n!)^2}\right] dx = \int_0^1 \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + \cdots\right) dx$$

$$= \left[x - \frac{x^3}{3\cdot 4} + \frac{x^5}{5\cdot 64} - \frac{x^7}{7\cdot 2304} + \cdots\right]_0^1 = 1 - \frac{1}{12} + \frac{1}{320} - \frac{1}{16{,}128} + \cdots$$

Since $\frac{1}{16{,}128} \approx 0.000062$, it follows from The Alternating Series Estimation Theorem that, correct to three decimal places, $\int_0^1 J_0(x)\,dx \approx 1 - \frac{1}{12} + \frac{1}{320} \approx 0.920$.

40. (a) If $a_n = \dfrac{(-1)^n x^{2n+1}}{n!(n+1)!\,2^{2n+1}}$, then

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{x^{2n+3}}{(n+1)!(n+2)!\,2^{2n+3}} \cdot \frac{n!(n+1)!\,2^{2n+1}}{x^{2n+1}}\right| = \left(\frac{x}{2}\right)^2 \lim_{n\to\infty} \frac{1}{(n+1)(n+2)} = 0 \text{ for all } x.$$

So $J_1(x)$ converges for all x and its domain is $(-\infty, \infty)$.

(b) $J_1(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}}$, $J_1'(x) = \sum_{n=0}^{\infty} \frac{(-1)^n (2n+1)\,x^{2n}}{n!\,(n+1)!\,2^{2n+1}}$, and $J_1''(x) = \sum_{n=1}^{\infty} \frac{(-1)^n (2n+1)(2n)\,x^{2n-1}}{n!\,(n+1)!\,2^{2n+1}}$.

$$\begin{aligned}
&x^2 J_1''(x) + xJ_1'(x) + \left(x^2-1\right)J_1(x) \\
&\quad = \sum_{n=1}^{\infty} \frac{(-1)^n (2n+1)(2n)x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} + \sum_{n=0}^{\infty} \frac{(-1)^n (2n+1)x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} \\
&\qquad + \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+3}}{n!\,(n+1)!\,2^{2n+1}} - \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} \\
&\quad = \sum_{n=1}^{\infty} \frac{(-1)^n (2n+1)(2n)x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} + \sum_{n=0}^{\infty} \frac{(-1)^n (2n+1)x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} \\
&\qquad - \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n+1}}{(n-1)!\,n!\,2^{2n-1}} - \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{n!\,(n+1)!\,2^{2n+1}} \qquad \left[\begin{array}{c}\text{Replace } n \text{ with } n-1 \\ \text{in the third term}\end{array}\right] \\
&\quad = \frac{x}{2} - \frac{x}{2} + \sum_{n=1}^{\infty} (-1)^n \left[\frac{(2n+1)(2n)+(2n+1)-(n)(n+1)2^2-1}{n!\,(n+1)!\,2^{2n+1}}\right] x^{2n+1} = 0
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $J_0(x) = \sum\limits_{n=0}^{\infty} \dfrac{(-1)^n\, x^{2n}}{2^{2n}\,(n!)^2} \quad \Rightarrow$

$$J_0'(x) = \sum_{n=1}^{\infty} \frac{(-1)^n\,(2n)x^{2n-1}}{2^{2n}\,(n!)^2} = \sum_{n=0}^{\infty} \frac{(-1)^{n+1}\,2(n+1)x^{2n+1}}{2^{2n+2}\,[(n+1)!]^2} \qquad [\text{Replace } n \text{ with } n+1]$$

$$= -\sum_{n=0}^{\infty} \frac{(-1)^n\, x^{2n+1}}{2^{2n+1}(n+1)!\,n!} \quad [\text{cancel 2 and } n+1\text{; take } -1 \text{ outside sum}] \quad = -J_1(x)$$

41. (a) $A(x) = 1 + \sum\limits_{n=1}^{\infty} a_n$, where $a_n = \dfrac{x^{3n}}{2\cdot 3\cdot 5\cdot 6\cdot\cdots\cdot(3n-1)(3n)}$, so $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = |x|^3 \lim\limits_{n\to\infty} \dfrac{1}{(3n+2)(3n+3)} = 0$ for all x, so the domain is $\mathbb{R}$.

(b), (c)

$s_0 = 1$ has been omitted from the graph. The partial sums seem to approximate $A(x)$ well near the origin, but as $|x|$ increases, we need to take a large number of terms to get a good approximation.

To plot A, we must first define $A(x)$ for the CAS. Note that for $n \geq 1$, the denominator of a_n is

$2\cdot 3\cdot 5\cdot 6\cdot\cdots\cdot(3n-1)\cdot 3n = \dfrac{(3n)!}{1\cdot 4\cdot 7\cdot\cdots\cdot(3n-2)} = \dfrac{(3n)!}{\prod_{k=1}^{n}(3k-2)}$, so $a_n = \dfrac{\prod_{k=1}^{n}(3k-2)}{(3n)!}x^{3n}$ and thus

$A(x) = 1 + \sum\limits_{n=1}^{\infty} \dfrac{\prod_{k=1}^{n}(3k-2)}{(3n)!}x^{3n}$. Both Maple and Mathematica are able to plot A if we define it this way.

Maple and Mathematica have two initially known Airy functions, called `AI·SERIES(z,m)` and `BI·SERIES(z,m)` from `AiryAi` and `AiryBi` in Maple and Mathematica (just `Ai` and `Bi` in older versions of Maple). However, it is very difficult to solve for A in terms of the CAS's Airy functions, although in fact $A(x) = \dfrac{\sqrt{3}\,\mathtt{AiryAi}(x) + \mathtt{AiryBi}(x)}{\sqrt{3}\,\mathtt{AiryAi}(0) + \mathtt{AiryBi}(0)}$.

42. $f(x) = \sum_{n=0}^{\infty} c_n x^n$, where $c_{n+4} = c_n$ for all $n \geq 0$. So

$$s_{4n-1} = c_0 + c_1x + c_2x^2 + c_3x^3 + c_0x^4 + c_1x^5 + c_2x^6 + c_3x^7 + \cdots + c_3x^{4n-1}$$

$$= \left(c_0 + c_1x + c_2x^2 + c_3x^3\right)\left(1 + x^4 + x^8 + \cdots + x^{4n-4}\right) \to \frac{c_0 + c_1x + c_2x^2 + c_3x^3}{1 - x^4} \text{ as } n \to \infty$$

[by (11.2.4) with $r = x^4$] for $\left|x^4\right| < 1 \quad \Leftrightarrow \quad |x| < 1$. Also s_{4n}, s_{4n+1}, s_{4n+2} have the same limits (for example, $s_{4n} = s_{4n-1} + c_0x^{4n}$ and $x^{4n} \to 0$ for $|x| < 1$). So if at least one of c_0, c_1, c_2, and c_3 is nonzero, then the interval of convergence is $(-1, 1)$ and $f(x) = \dfrac{c_0 + c_1x + c_2x^2 + c_3x^3}{1 - x^4}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. $f(x) = 1 + 2x + x^2 + 2x^3 + x^4 + \cdots$, where $c_{2n} = 1$ and $c_{2n+1} = 2$ for all $n \geq 0$. So

$$\begin{aligned} s_{2n-1} &= 1 + 2x + x^2 + 2x^3 + x^4 + 2x^5 + \cdots + x^{2n-2} + 2x^{2n-1} \\ &= 1(1+2x) + x^2(1+2x) + x^4(1+2x) + \cdots + x^{2n-2}(1+2x) = (1+2x)(1 + x^2 + x^4 + \cdots + x^{2n-2}) \\ &= (1+2x)\frac{1-x^{2n}}{1-x^2} \quad \text{[by (11.2.3) with } r = x^2\text{]} \quad \to \frac{1+2x}{1-x^2} \text{ as } n \to \infty \text{ by (11.2.4), when } |x| < 1. \end{aligned}$$

Also $s_{2n} = s_{2n-1} + x^{2n} \to \dfrac{1+2x}{1-x^2}$ since $x^{2n} \to 0$ for $|x| < 1$. Therefore, $s_n \to \dfrac{1+2x}{1-x^2}$ since s_{2n} and s_{2n-1} both approach $\dfrac{1+2x}{1-x^2}$ as $n \to \infty$. Thus, the interval of convergence is $(-1, 1)$ and $f(x) = \dfrac{1+2x}{1-x^2}$.

44. $\dfrac{|\sin nx|}{n^2} \leq \dfrac{1}{n^2}$, so $\sum\limits_{n=1}^{\infty} \dfrac{\sin nx}{n^2}$ converges by the Direct Comparison Test. $\dfrac{d}{dx}\left(\dfrac{\sin nx}{n^2}\right) = \dfrac{\cos nx}{n}$, so when $x = 2k\pi$ [k an integer], $\sum\limits_{n=1}^{\infty} f_n'(x) = \sum\limits_{n=1}^{\infty} \dfrac{\cos(2kn\pi)}{n} = \sum\limits_{n=1}^{\infty} \dfrac{1}{n}$, which diverges [harmonic series]. $f_n''(x) = -\sin nx$, so $\sum\limits_{n=1}^{\infty} f_n''(x) = -\sum\limits_{n=1}^{\infty} \sin nx$, which converges only if $\sin nx = 0$, or $x = k\pi$ [k an integer].

45. If $a_n = \dfrac{x^n}{n^2}$, then by the Ratio Test, $\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \left|\dfrac{x^{n+1}}{(n+1)^2} \cdot \dfrac{n^2}{x^n}\right| = |x| \lim\limits_{n\to\infty} \left(\dfrac{n}{n+1}\right)^2 = |x| < 1$ for convergence, so $R = 1$. When $x = \pm 1$, $\sum\limits_{n=1}^{\infty} \left|\dfrac{x^n}{n^2}\right| = \sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}$ which is a convergent p-series ($p = 2 > 1$), so the interval of convergence for f is $[-1, 1]$. By Theorem 2, the radii of convergence of f' and f'' are both 1, so we need only check the endpoints. $f(x) = \sum\limits_{n=1}^{\infty} \dfrac{x^n}{n^2} \quad \Rightarrow \quad f'(x) = \sum\limits_{n=1}^{\infty} \dfrac{nx^{n-1}}{n^2} = \sum\limits_{n=0}^{\infty} \dfrac{x^n}{n+1}$, and this series diverges for $x = 1$ (harmonic series) and converges for $x = -1$ (Alternating Series Test), so the interval of convergence is $[-1, 1)$. $f''(x) = \sum\limits_{n=1}^{\infty} \dfrac{nx^{n-1}}{n+1}$ diverges at both 1 and -1 (Test for Divergence) since $\lim\limits_{n\to\infty} \dfrac{n}{n+1} = 1 \neq 0$, so its interval of convergence is $(-1, 1)$.

46. (a) $\sum\limits_{n=1}^{\infty} nx^{n-1} = \sum\limits_{n=0}^{\infty} \dfrac{d}{dx} x^n = \dfrac{d}{dx}\left[\sum\limits_{n=0}^{\infty} x^n\right] = \dfrac{d}{dx}\left[\dfrac{1}{1-x}\right] = -\dfrac{1}{(1-x)^2}(-1) = \dfrac{1}{(1-x)^2}$, $|x| < 1$.

(b) (i) $\sum\limits_{n=1}^{\infty} nx^n = x \sum\limits_{n=1}^{\infty} nx^{n-1} = x\left[\dfrac{1}{(1-x)^2}\right]$ [from part (a)] $= \dfrac{x}{(1-x)^2}$ for $|x| < 1$.

(ii) Put $x = \frac{1}{2}$ in (i): $\sum\limits_{n=1}^{\infty} \dfrac{n}{2^n} = \sum\limits_{n=1}^{\infty} n\left(\frac{1}{2}\right)^n = \dfrac{1/2}{(1-1/2)^2} = 2$.

(c) (i) $\sum\limits_{n=2}^{\infty} n(n-1)x^n = x^2 \sum\limits_{n=2}^{\infty} n(n-1)x^{n-2} = x^2 \dfrac{d}{dx}\left[\sum\limits_{n=1}^{\infty} nx^{n-1}\right] = x^2 \dfrac{d}{dx} \dfrac{1}{(1-x)^2}$

$$= x^2 \frac{2}{(1-x)^3} = \frac{2x^2}{(1-x)^3} \text{ for } |x| < 1.$$

(ii) Put $x = \frac{1}{2}$ in (i): $\sum\limits_{n=2}^{\infty} \dfrac{n^2 - n}{2^n} = \sum\limits_{n=2}^{\infty} n(n-1)\left(\frac{1}{2}\right)^n = \dfrac{2(1/2)^2}{(1-1/2)^3} = 4$.

(iii) From (b)(ii) and (c)(ii), we have $\sum\limits_{n=1}^{\infty} \dfrac{n^2}{2^n} = \sum\limits_{n=1}^{\infty} \dfrac{n^2 - n}{2^n} + \sum\limits_{n=1}^{\infty} \dfrac{n}{2^n} = 4 + 2 = 6$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $f(x) = \dfrac{1}{1-x} = \sum\limits_{n=0}^{\infty} x^n \quad \Rightarrow \quad f'(x) = \dfrac{1}{(1-x)^2} = \sum\limits_{n=1}^{\infty} nx^{n-1} \quad \Rightarrow \quad f''(x) = \dfrac{2}{(1-x)^3} = \sum\limits_{n=2}^{\infty} n(n-1)x^{n-2}$

The power series representation of $h(x)$ is

$$\begin{aligned} h(x) = xf'(x) + x^2 f''(x) &= x\sum_{n=1}^{\infty} nx^{n-1} + x^2 \sum_{n=2}^{\infty} n(n-1)x^{n-2} = \sum_{n=1}^{\infty} nx^n + \sum_{n=2}^{\infty} n(n-1)x^n \\ &= x + \sum_{n=2}^{\infty} nx^n + \sum_{n=2}^{\infty} n(n-1)x^n = x + \sum_{n=2}^{\infty} [nx^n + n(n-1)x^n] \\ &= x + \sum_{n=2}^{\infty} (1+n-1)nx^n = x + \sum_{n=2}^{\infty} n^2 x^n = \sum_{n=1}^{\infty} n^2 x^n \end{aligned}$$

$h(x)$ has the same radius of convergence as the power series representation of $f(x)$, that is, $R = 1$.

Now, $h\left(\dfrac{1}{2}\right) = \sum\limits_{n=1}^{\infty} \dfrac{n^2}{2^n}$ and using the function representation, we have

$h\left(\dfrac{1}{2}\right) = \dfrac{1}{2}f'\left(\dfrac{1}{2}\right) + \left(\dfrac{1}{2}\right)^2 f''\left(\dfrac{1}{2}\right) = \dfrac{1}{2} \cdot \dfrac{1}{(1-1/2)^2} + \dfrac{1}{4} \cdot \dfrac{2}{(1-1/2)^3} = 6$. Thus, $\sum\limits_{n=1}^{\infty} \dfrac{n^2}{2^n} = 6$.

48. By Example 4, $f(x) = \dfrac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \cdots \sum\limits_{n=1}^{\infty} nx^{n-1}$, with radius of convergence $R = 1$. We wish to substitute a value of x with $|x| < 1$ that will result in a denominator of $99^2 = 9801$. Using $x = 1/100$, we get

$$\frac{1}{(1-1/100)^2} = 1 + 2\left(\frac{1}{100}\right) + 3\left(\frac{1}{100}\right)^2 + 4\left(\frac{1}{100}\right)^3 + \cdots \quad \Rightarrow$$

$$\frac{100^2}{99^2} = 1 + \frac{2}{100} + \frac{3}{100^2} + \frac{4}{100^3} + \cdots + \frac{98}{100^{97}} + \cdots \quad \Rightarrow \quad \frac{1}{9801} = \frac{1}{10^4} + \frac{2}{10^6} + \frac{3}{10^8} + \frac{4}{10^{10}} + \cdots + \frac{98}{10^{198}} + \cdots.$$

Observe that the series starts with 0.0001, that is, 1 ten-thousandths, and each subsequent term adds the next integer two decimal places deeper. This gives the initial pattern 0.00 01 02 03 04 05 06 07 08 09 10 11 . . ., which will hold until the 100th integer is reached, at which point digits will "carry down" from subsequent terms. We explore this behavior below, starting from the 98th integer.

$$\begin{aligned} &\frac{98}{10^{198}} + \frac{99}{10^{200}} + \frac{100}{10^{202}} + \frac{101}{10^{204}} + \frac{102}{10^{206}} + \cdots \\ &\qquad = \frac{98}{10^{198}} + \frac{99}{10^{200}} + \frac{1}{10^{200}} + \left(\frac{100}{10^{204}} + \frac{1}{10^{204}}\right) + \left(\frac{100}{10^{206}} + \frac{2}{10^{206}}\right) + \cdots \\ &\qquad = \frac{98}{10^{198}} + \frac{100}{10^{200}} + \left(\frac{1}{10^{202}} + \frac{1}{10^{204}}\right) + \left(\frac{1}{10^{204}} + \frac{2}{10^{206}}\right) + \cdots \\ &\qquad = \frac{98}{10^{198}} + \frac{1}{10^{198}} + \frac{1}{10^{202}} + \frac{2}{10^{204}} + \cdots = \frac{99}{10^{198}} + \frac{1}{10^{202}} + \frac{2}{10^{204}} + \cdots \end{aligned}$$

Thus, we see that the 98th integer is omitted and the sequence of integers repeats after the 99th integer (due to the repeated carrying from later terms). This gives the result $\dfrac{1}{9801} = 0.\overline{00\,01\,02\,03...96\,97\,99}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. By Example 6, $\tan^{-1} x = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{2n+1}}{2n+1}$ for $|x| < 1$. In particular, for $x = \dfrac{1}{\sqrt{3}}$, we have $\dfrac{\pi}{6} = \tan^{-1}\left(\dfrac{1}{\sqrt{3}}\right) = \sum_{n=0}^{\infty} (-1)^n \dfrac{\left(1/\sqrt{3}\right)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} (-1)^n \left(\dfrac{1}{3}\right)^n \dfrac{1}{\sqrt{3}} \dfrac{1}{2n+1}$, so

$$\pi = \frac{6}{\sqrt{3}} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)3^n} = 2\sqrt{3} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)3^n}.$$

50. (a) $\displaystyle\int_0^{1/2} \frac{dx}{x^2 - x + 1} = \int_0^{1/2} \frac{dx}{(x - 1/2)^2 + 3/4}$ $\quad \left[x - \frac{1}{2} = \frac{\sqrt{3}}{2}u,\ u = \frac{2}{\sqrt{3}}\left(x - \frac{1}{2}\right),\ dx = \frac{\sqrt{3}}{2}\,du\right]$

$$= \int_{-1/\sqrt{3}}^{0} \frac{\left(\sqrt{3}/2\right) du}{(3/4)(u^2+1)} = \frac{2\sqrt{3}}{3}\Big[\tan^{-1} u\Big]_{-1/\sqrt{3}}^{0} = \frac{2}{\sqrt{3}}\left[0 - \left(-\frac{\pi}{6}\right)\right] = \frac{\pi}{3\sqrt{3}}$$

(b) $\dfrac{1}{x^3+1} = \dfrac{1}{(x+1)(x^2 - x + 1)} \quad \Rightarrow$

$$\frac{1}{x^2 - x + 1} = (x+1)\left(\frac{1}{1+x^3}\right) = (x+1)\frac{1}{1-(-x^3)} = (x+1) \sum_{n=0}^{\infty} (-1)^n x^{3n}$$

$$= \sum_{n=0}^{\infty} (-1)^n x^{3n+1} + \sum_{n=0}^{\infty} (-1)^n x^{3n} \quad \text{for } |x| < 1 \quad \Rightarrow$$

$$\int \frac{dx}{x^2 - x + 1} = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{3n+2}}{3n+2} + \sum_{n=0}^{\infty} (-1)^n \frac{x^{3n+1}}{3n+1} \text{ for } |x| < 1 \quad \Rightarrow$$

$$\int_0^{1/2} \frac{dx}{x^2 - x + 1} = \sum_{n=0}^{\infty} (-1)^n \left[\frac{1}{4 \cdot 8^n (3n+2)} + \frac{1}{2 \cdot 8^n (3n+1)}\right] = \frac{1}{4} \sum_{n=0}^{\infty} \frac{(-1)^n}{8^n}\left(\frac{2}{3n+1} + \frac{1}{3n+2}\right).$$

By part (a), this equals $\dfrac{\pi}{3\sqrt{3}}$, so $\pi = \dfrac{3\sqrt{3}}{4} \sum_{n=0}^{\infty} \dfrac{(-1)^n}{8^n}\left(\dfrac{2}{3n+1} + \dfrac{1}{3n+2}\right)$.

51. Using the Ratio Test, the series $\sum_{n=0}^{\infty} c_n x^n$ converges when $\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| < 1 \quad \Rightarrow$

$$\lim_{n\to\infty} \left|\frac{c_{n+1}x^{n+1}}{c_n x^n}\right| = |x| \lim_{n\to\infty} \left|\frac{c_{n+1}}{c_n}\right| < 1 \quad \Rightarrow \quad |x| < \frac{1}{\lim\limits_{n\to\infty} \left|\dfrac{c_{n+1}}{c_n}\right|} = R \quad \Rightarrow \quad \lim_{n\to\infty} \left|\frac{c_{n+1}}{c_n}\right| = \frac{1}{R}.$$

Now, using the Ratio Test for the series $\sum_{n=1}^{\infty} n c_n x^{n-1}$, we find

$\lim_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\dfrac{(n+1)c_{n+1}x^n}{n c_n x^{n-1}}\right| = \lim_{n\to\infty} \left|\left(1 + \dfrac{1}{n}\right) \dfrac{c_{n+1}}{c_n}\, x\right| = |x| \lim_{n\to\infty} \left|\dfrac{c_{n+1}}{c_n}\right| = \dfrac{|x|}{R}$. Hence, the series $\sum_{n=1}^{\infty} n c_n x^{n-1}$ converges when $\dfrac{|x|}{R} < 1$ or $|x| < R$. Finally, testing the series $\sum_{n=0}^{\infty} c_n \dfrac{x^{n+1}}{n+1}$ using the Ratio Test, we have

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{c_{n+1}x^{n+2}}{n+2} \cdot \frac{n+1}{c_n x^{n+1}}\right| = \lim_{n\to\infty} \left|\frac{(n+1)c_{n+1}x}{(n+2)c_n}\right| = |x| \lim_{n\to\infty} \left|\frac{(1+1/n)c_{n+1}}{(1+2/n)c_n}\right|$$

$$= |x| \lim_{n\to\infty} \left|\frac{c_{n+1}}{c_n}\right| = \frac{|x|}{R}$$

so $\sum_{n=0}^{\infty} c_n \dfrac{x^{n+1}}{n+1}$ also converges when $|x| < R$. Thus, both $\sum_{n=1}^{\infty} n c_n x^{n-1}$ and $\sum_{n=0}^{\infty} c_n \dfrac{x^{n+1}}{n+1}$ have radius of convergence R when $\sum_{n=0}^{\infty} c_n x^n$ has radius of convergence R.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11.10 Taylor and Maclaurin Series

1. Using Theorem 5 with $\sum_{n=0}^{\infty} b_n(x-5)^n$, $b_n = \dfrac{f^{(n)}(a)}{n!}$, so $b_8 = \dfrac{f^{(8)}(5)}{8!}$.

2. (a) Using Equation 7, the Maclaurin series of f must have the form $f(0) + \dfrac{f'(0)}{1!}\,x + \dfrac{f''(0)}{2!}\,x^2 + \cdots$. Comparing to the given series, $1.1 + 0.7x^2 + 2.2x^3$, we must have $f(0) = 1.1$. But from the graph, $f(0) \approx 0.5$. Hence, the given series is *not* the Maclaurin series.

(b) Using Equation 6, a power series expansion of f at 1 must have the form $\dfrac{f'(1)}{1!}(x-1) + \dfrac{f''(1)}{2!}(x-1)^2 + \cdots$. Comparing to the given series, $1.6 - 0.8(x-1) + 0.4(x-1)^2 - \cdots$, we must have $f'(1) = -0.8$. But from the graph, f is increasing near $x = 1$, so $f'(1)$ is positive. Hence, the given series is *not* the Taylor series of f centered at 1.

(c) A power series expansion of f at 2 must have the form $f(2) + \dfrac{f'(2)}{1!}(x-2) + \dfrac{f''(2)}{2!}(x-2)^2 + \cdots$. Comparing to the given series, $2.8 + 0.5(x-2) + 1.5(x-2)^2 - 0.1(x-2)^3 + \cdots$, we must have $\frac{1}{2}f''(2) = 1.5$; that is, $f''(2)$ is positive. But from the graph, f is concave downward near $x = 2$, so $f''(2)$ must be negative. Hence, the given series is *not* the Taylor series of f centered at 2.

3. Since $f^{(n)}(0) = (n+1)!$, Equation 7 gives the Maclaurin series

$\sum_{n=0}^{\infty} \dfrac{f^{(n)}(0)}{n!}x^n = \sum_{n=0}^{\infty} \dfrac{(n+1)!}{n!}x^n = \sum_{n=0}^{\infty}(n+1)x^n$. Applying the Ratio Test with $a_n = (n+1)x^n$ gives us

$\lim_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\dfrac{(n+2)x^{n+1}}{(n+1)x^n}\right| = |x| \lim_{n\to\infty}\dfrac{n+2}{n+1} = |x|\cdot 1 = |x|$. For convergence, we must have $|x| < 1$, so the radius of convergence $R = 1$.

4. Since $f^{(n)}(4) = \dfrac{(-1)^n\, n!}{3^n(n+1)}$, Equation 6 gives the Taylor series

$\sum_{n=0}^{\infty} \dfrac{f^{(n)}(4)}{n!}(x-4)^n = \sum_{n=0}^{\infty} \dfrac{(-1)^n\, n!}{3^n(n+1)\, n!}(x-4)^n = \sum_{n=0}^{\infty} \dfrac{(-1)^n}{3^n(n+1)}(x-4)^n$, which is the Taylor series for f centered at 4. Apply the Ratio Test to find the radius of convergence R.

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^{n+1}(x-4)^{n+1}}{3^{n+1}(n+2)}\cdot\frac{3^n(n+1)}{(-1)^n(x-4)^n}\right| = \lim_{n\to\infty}\left|\frac{(-1)(x-4)(n+1)}{3(n+2)}\right|$$
$$= \frac{1}{3}\,|x-4| \lim_{n\to\infty}\frac{n+1}{n+2} = \frac{1}{3}\,|x-4|$$

For convergence, $\frac{1}{3}\,|x-4| < 1 \quad\Leftrightarrow\quad |x-4| < 3$, so $R = 3$.

5.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	xe^x	0
1	$(x+1)e^x$	1
2	$(x+2)e^x$	2
3	$(x+3)e^x$	3
4	$(x+4)e^x$	4

Using Equation 6 with $n = 0$ to 4 and $a = 0$, we get

$$\sum_{n=0}^{4} \frac{f^{(n)}(0)}{n!}(x-0)^n = \frac{0}{0!}\,x^0 + \frac{1}{1!}\,x^1 + \frac{2}{2!}\,x^2 + \frac{3}{3!}\,x^3 + \frac{4}{4!}\,x^4$$
$$= x + x^2 + \tfrac{1}{2}x^3 + \tfrac{1}{6}x^4$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6.

n	$f^{(n)}(x)$	$f^{(n)}(2)$
0	$\dfrac{1}{1+x}$	$\frac{1}{3}$
1	$-\dfrac{1}{(1+x)^2}$	$-\frac{1}{9}$
2	$\dfrac{2}{(1+x)^3}$	$\frac{2}{27}$
3	$-\dfrac{6}{(1+x)^4}$	$-\frac{6}{81}$

$$\begin{aligned}\sum_{n=0}^{3}\frac{f^{(n)}(2)}{n!}(x-2)^n &= \frac{\frac{1}{3}}{0!}(x-2)^0 - \frac{\frac{1}{9}}{1!}(x-2)^1 \\ &\quad + \frac{\frac{2}{27}}{2!}(x-2)^2 - \frac{\frac{6}{81}}{3!}(x-2)^3 \\ &= \tfrac{1}{3} - \tfrac{1}{9}(x-2) + \tfrac{1}{27}(x-2)^2 - \tfrac{1}{81}(x-2)^3\end{aligned}$$

7.

n	$f^{(n)}(x)$	$f^{(n)}(8)$
0	$\sqrt[3]{x}$	2
1	$\dfrac{1}{3x^{2/3}}$	$\frac{1}{12}$
2	$-\dfrac{2}{9x^{5/3}}$	$-\frac{2}{288}$
3	$\dfrac{10}{27x^{8/3}}$	$\frac{10}{6912}$

$$\begin{aligned}\sum_{n=0}^{3}\frac{f^{(n)}(8)}{n!}(x-8)^n &= \frac{2}{0!}(x-8)^0 + \frac{\frac{1}{12}}{1!}(x-8)^1 \\ &\quad - \frac{\frac{2}{288}}{2!}(x-8)^2 + \frac{\frac{10}{6912}}{3!}(x-8)^3 \\ &= 2 + \tfrac{1}{12}(x-8) - \tfrac{1}{288}(x-8)^2 + \tfrac{5}{20{,}736}(x-8)^3\end{aligned}$$

8.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$\ln x$	0
1	$1/x$	1
2	$-1/x^2$	-1
3	$2/x^3$	2
4	$-6/x^4$	-6

$$\begin{aligned}\sum_{n=0}^{4}\frac{f^{(n)}(1)}{n!}(x-1)^n &= \frac{0}{0!}(x-1)^0 + \frac{1}{1!}(x-1)^1 - \frac{1}{2!}(x-1)^2 \\ &\quad + \frac{2}{3!}(x-1)^3 - \frac{6}{4!}(x-1)^4 \\ &= (x-1) - \tfrac{1}{2}(x-1)^2 + \tfrac{1}{3}(x-1)^3 - \tfrac{1}{4}(x-1)^4\end{aligned}$$

9.

n	$f^{(n)}(x)$	$f^{(n)}(\pi/6)$
0	$\sin x$	$1/2$
1	$\cos x$	$\sqrt{3}/2$
2	$-\sin x$	$-1/2$
3	$-\cos x$	$-\sqrt{3}/2$

$$\begin{aligned}\sum_{n=0}^{3}\frac{f^{(n)}(\pi/6)}{n!}\left(x-\frac{\pi}{6}\right)^n &= \frac{1/2}{0!}\left(x-\frac{\pi}{6}\right)^0 + \frac{\sqrt{3}/2}{1!}\left(x-\frac{\pi}{6}\right)^1 - \frac{1/2}{2!}\left(x-\frac{\pi}{6}\right)^2 - \frac{\sqrt{3}/2}{3!}\left(x-\frac{\pi}{6}\right)^3 \\ &= \frac{1}{2} + \frac{\sqrt{3}}{2}\left(x-\frac{\pi}{6}\right) - \frac{1}{4}\left(x-\frac{\pi}{6}\right)^2 - \frac{\sqrt{3}}{12}\left(x-\frac{\pi}{6}\right)^3\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\cos^2 x$	1
1	$-2\cos x\,\sin x = -\sin 2x$	0
2	$-2\cos 2x$	-2
3	$4\sin 2x$	0
4	$8\cos 2x$	8
5	$-16\sin 2x$	0
6	$-32\cos 2x$	-32

$$\sum_{n=0}^{6} \frac{f^{(n)}(0)}{n!}(x-0)^n = \frac{1}{0!}x^0 - \frac{2}{2!}x^2 + \frac{8}{4!}x^4 - \frac{32}{6!}x^6$$
$$= 1 - x^2 + \tfrac{1}{3}x^4 - \tfrac{2}{45}x^6$$

11.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$(1-x)^{-2}$	1
1	$2(1-x)^{-3}$	2
2	$6(1-x)^{-4}$	6
3	$24(1-x)^{-5}$	24
4	$120(1-x)^{-6}$	120
$\vdots$	$\vdots$	$\vdots$

$$(1-x)^{-2} = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \cdots$$
$$= 1 + 2x + \tfrac{6}{2}x^2 + \tfrac{24}{6}x^3 + \tfrac{120}{24}x^4 + \cdots$$
$$= 1 + 2x + 3x^2 + 4x^3 + 5x^4 + \cdots = \sum_{n=0}^{\infty}(n+1)x^n$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(n+2)x^{n+1}}{(n+1)x^n}\right| = |x|\lim_{n\to\infty}\frac{n+2}{n+1} = |x|\,(1) = |x| < 1$$

for convergence, so $R = 1$.

12.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\ln(1+x)$	0
1	$(1+x)^{-1}$	1
2	$-(1+x)^{-2}$	-1
3	$2(1+x)^{-3}$	2
4	$-6(1+x)^{-4}$	-6
5	$24(1+x)^{-5}$	24
$\vdots$	$\vdots$	$\vdots$

$$\ln(1+x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2$$
$$+ \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \frac{f^{(5)}(0)}{5!}x^5 + \cdots$$
$$= 0 + x - \tfrac{1}{2}x^2 + \tfrac{2}{6}x^3 - \tfrac{6}{24}x^4 + \tfrac{24}{120}x^5 - \cdots$$
$$= x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \frac{x^5}{5} - \cdots = \sum_{n=1}^{\infty}\frac{(-1)^{n-1}}{n}x^n$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{x^{n+1}}{n+1}\cdot\frac{n}{x^n}\right| = \lim_{n\to\infty}\frac{|x|}{1+1/n} = |x| < 1 \text{ for convergence,}$$

so $R = 1$.

Notice that the answer agrees with the entry for $\ln(1+x)$ in Table 1, but we obtained it by a different method. (Compare with Example 11.9.5.)

13.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\cos x$	1
1	$-\sin x$	0
2	$-\cos x$	-1
3	$\sin x$	0
4	$\cos x$	1
$\vdots$	$\vdots$	$\vdots$

$$\cos x = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \cdots$$
$$= 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \cdots$$
$$= \sum_{n=0}^{\infty}(-1)^n\frac{x^{2n}}{(2n)!} \qquad \text{[Agrees with (16).]}$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{x^{2n+2}}{(2n+2)!}\cdot\frac{(2n)!}{x^{2n}}\right| = \lim_{n\to\infty}\frac{x^2}{(2n+2)(2n+1)} = 0 < 1$$

for all x, so $R = \infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	e^{-2x}	1
1	$-2e^{-2x}$	-2
2	$4e^{-2x}$	4
3	$-8e^{-2x}$	-8
4	$16e^{-2x}$	16
$\vdots$	$\vdots$	$\vdots$

$$e^{-2x} = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!}\, x^n = \sum_{n=0}^{\infty} \frac{(-2)^n}{n!}\, x^n.$$

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-2)^{n+1}x^{n+1}}{(n+1)!} \cdot \frac{n!}{(-2)^n x^n}\right| = \lim_{n\to\infty} \frac{2\,|x|}{n+1}$$

$$= 0 < 1 \quad \text{for all } x, \text{ so } R = \infty.$$

15.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$2x^4 - 3x^2 + 3$	3
1	$8x^3 - 6x$	0
2	$24x^2 - 6$	-6
3	$48x$	0
4	48	48
5	0	0
6	0	0
$\vdots$	$\vdots$	$\vdots$

$f^{(n)}(x) = 0$ for $n \geq 5$, so f has a finite Maclaurin series.

$$f(x) = 2x^4 - 3x^2 + 3$$

$$= f(0) + \frac{f'(0)}{1!}\, x + \frac{f''(0)}{2!}\, x^2 + \frac{f'''(0)}{3!}\, x^3 + \frac{f^{(4)}(0)}{4!}\, x^4$$

$$= 3 + 0x + \frac{-6}{2}x^2 + 0x^3 + \frac{48}{24}x^4 = 3 - 3x^2 + 2x^4$$

A finite series converges for all x, so $R = \infty$.

16.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sin 3x$	0
1	$3\cos 3x$	3
2	$-9\sin 3x$	0
3	$-27\cos 3x$	-27
4	$81\sin 3x$	0
5	$243\cos 3x$	243
6	$729\sin 3x$	0
$\vdots$	$\vdots$	$\vdots$

$$\sin 3x = f(0) + \frac{f'(0)}{1!}\, x + \frac{f''(0)}{2!}\, x^2 + \frac{f'''(0)}{3!}\, x^3 + \frac{f^{(4)}(0)}{4!}\, x^4 + \cdots$$

$$= 3x - \frac{27}{3!}\, x^3 + \frac{243}{5!}\, x^5 - \cdots = \sum_{n=0}^{\infty} \frac{(-1)^n\, 3^{2n+1}}{(2n+1)!}\, x^{2n+1}$$

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(-1)^{n+1}\, 3^{2n+3}\, x^{2n+3}}{(2n+3)!} \cdot \frac{(2n+1)!}{(-1)^n\, 3^{2n+1}\, x^{2n+1}}\right|$$

$$= x^2 \cdot \lim_{n\to\infty} \frac{3^2}{(2n+3)(2n+2)}$$

$$= 0 < 1 \text{ for all } x, \text{ so } R = \infty.$$

17.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	2^x	1
1	$2^x(\ln 2)$	$\ln 2$
2	$2^x(\ln 2)^2$	$(\ln 2)^2$
3	$2^x(\ln 2)^3$	$(\ln 2)^3$
4	$2^x(\ln 2)^4$	$(\ln 2)^4$
$\vdots$	$\vdots$	$\vdots$

$$2^x = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!}\, x^n = \sum_{n=0}^{\infty} \frac{(\ln 2)^n}{n!}\, x^n.$$

$$\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(\ln 2)^{n+1}x^{n+1}}{(n+1)!} \cdot \frac{n!}{(\ln 2)^n x^n}\right|$$

$$= \lim_{n\to\infty} \frac{(\ln 2)\,|x|}{n+1} = 0 < 1 \quad \text{for all } x, \text{ so } R = \infty.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$x\cos x$	0
1	$-x\sin x+\cos x$	1
2	$-x\cos x-2\sin x$	0
3	$x\sin x-3\cos x$	-3
4	$x\cos x+4\sin x$	0
5	$-x\sin x+5\cos x$	5
6	$-x\cos x-6\sin x$	0
7	$x\sin x-7\cos x$	-7
$\vdots$	$\vdots$	$\vdots$

$$\begin{aligned} x\cos x &= f(0)+f'(0)x+\frac{f''(0)}{2!}\,x^2+\frac{f'''(0)}{3!}\,x^3+\frac{f^{(4)}(0)}{4!}\,x^4+\cdots \\ &= 0+1x+0-\frac{3}{3!}\,x^3+0+\frac{5}{5!}\,x^5+0-\frac{7}{7!}\,x^7+\cdots \\ &= x-\frac{1}{2!}\,x^3+\frac{1}{4!}\,x^5-\frac{1}{6!}\,x^7+\cdots \\ &= \sum_{n=0}^{\infty}(-1)^n\,\frac{1}{(2n)!}\,x^{2n+1} \end{aligned}$$

$$\begin{aligned} \lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| &= \lim_{n\to\infty}\left|\frac{(-1)^{n+1}x^{2n+3}}{(2n+2)!}\cdot\frac{(2n)!}{(-1)^n x^{2n+1}}\right| \\ &= \lim_{n\to\infty}\frac{x^2}{(2n+2)(2n+1)}=0<1 \quad \text{for all } x\text{, so } R=\infty. \end{aligned}$$

19.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sinh x$	0
1	$\cosh x$	1
2	$\sinh x$	0
3	$\cosh x$	1
4	$\sinh x$	0
$\vdots$	$\vdots$	$\vdots$

$$f^{(n)}(0)=\begin{cases}0 & \text{if } n \text{ is even}\\ 1 & \text{if } n \text{ is odd}\end{cases} \qquad \text{so } \sinh x=\sum_{n=0}^{\infty}\frac{x^{2n+1}}{(2n+1)!}.$$

Use the Ratio Test to find R. If $a_n=\dfrac{x^{2n+1}}{(2n+1)!}$, then

$$\begin{aligned} \lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| &= \lim_{n\to\infty}\left|\frac{x^{2n+3}}{(2n+3)!}\cdot\frac{(2n+1)!}{x^{2n+1}}\right| = x^2\cdot\lim_{n\to\infty}\frac{1}{(2n+3)(2n+2)} \\ &= 0<1 \quad \text{for all } x\text{, so } R=\infty. \end{aligned}$$

20.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\cosh x$	1
1	$\sinh x$	0
2	$\cosh x$	1
3	$\sinh x$	0
$\vdots$	$\vdots$	$\vdots$

$$f^{(n)}(0)=\begin{cases}1 & \text{if } n \text{ is even}\\ 0 & \text{if } n \text{ is odd}\end{cases} \qquad \text{so } \cosh x=\sum_{n=0}^{\infty}\frac{x^{2n}}{(2n)!}.$$

Use the Ratio Test to find R. If $a_n=\dfrac{x^{2n}}{(2n)!}$, then

$$\begin{aligned} \lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| &= \lim_{n\to\infty}\left|\frac{x^{2n+2}}{(2n+2)!}\cdot\frac{(2n)!}{x^{2n}}\right| = x^2\cdot\lim_{n\to\infty}\frac{1}{(2n+2)(2n+1)} \\ &= 0<1 \quad \text{for all } x\text{, so } R=\infty. \end{aligned}$$

21.

n	$f^{(n)}(x)$	$f^{(n)}(2)$
0	x^5+2x^3+x	50
1	$5x^4+6x^2+1$	105
2	$20x^3+12x$	184
3	$60x^2+12$	252
4	$120x$	240
5	120	120
6	0	0
7	0	0
$\vdots$	$\vdots$	$\vdots$

$f^{(n)}(x)=0$ for $n\geq 6$, so f has a finite expansion about $a=2$.

$$\begin{aligned} f(x) &= x^5+2x^3+x=\sum_{n=0}^{5}\frac{f^{(n)}(2)}{n!}\,(x-2)^n \\ &= \frac{50}{0!}\,(x-2)^0+\frac{105}{1!}\,(x-2)^1+\frac{184}{2!}\,(x-2)^2+\frac{252}{3!}\,(x-2)^3 \\ &\qquad +\frac{240}{4!}\,(x-2)^4+\frac{120}{5!}\,(x-2)^5 \\ &= 50+105(x-2)+92(x-2)^2+42(x-2)^3 \\ &\qquad +10(x-2)^4+(x-2)^5 \end{aligned}$$

A finite series converges for all x, so $R=\infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22.

n	$f^{(n)}(x)$	$f^{(n)}(-2)$
0	$x^6 - x^4 + 2$	50
1	$6x^5 - 4x^3$	-160
2	$30x^4 - 12x^2$	432
3	$120x^3 - 24x$	-912
4	$360x^2 - 24$	1416
5	$720x$	-1440
6	720	720
7	0	0
8	0	0
$\vdots$	$\vdots$	$\vdots$

$f(n)(x) = 0$ for $n \geq 7$, so f has a finite expansion about $a = -2$.

$$f(x) = x^6 - x^4 + 2 = \sum_{n=0}^{6} \frac{f^{(n)}(-2)}{n!}(x+2)^n$$

$$= \frac{50}{0!}(x+2)^0 - \frac{160}{1!}(x+2)^1 + \frac{432}{2!}(x+2)^2 - \frac{912}{3!}(x+2)^3 + \frac{1416}{4!}(x+2)^4 - \frac{1440}{5!}(x+2)^5 + \frac{720}{6!}(x+2)^6$$

$$= 50 - 160(x+2) + 216(x+2)^2 - 152(x+2)^3 + 59(x+2)^4 - 12(x+2)^5 + (x+2)^6$$

A finite series converges for all x, so $R = \infty$.

23.

n	$f^{(n)}(x)$	$f^{(n)}(2)$
0	$\ln x$	$\ln 2$
1	$1/x$	$1/2$
2	$-1/x^2$	$-1/2^2$
3	$2/x^3$	$2/2^3$
4	$-6/x^4$	$-6/2^4$
5	$24/x^5$	$24/2^5$
$\vdots$	$\vdots$	$\vdots$

$$f(x) = \ln x = \sum_{n=0}^{\infty} \frac{f^{(n)}(2)}{n!}(x-2)^n$$

$$= \frac{\ln 2}{0!}(x-2)^0 + \frac{1}{1!\,2^1}(x-2)^1 + \frac{-1}{2!\,2^2}(x-2)^2 + \frac{2}{3!\,2^3}(x-2)^3 + \frac{-6}{4!\,2^4}(x-2)^4 + \frac{24}{5!\,2^5}(x-2)^5 + \cdots$$

$$= \ln 2 + \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(n-1)!}{n!\,2^n}(x-2)^n$$

$$= \ln 2 + \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n\,2^n}(x-2)^n$$

$$\lim_{n\to\infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n\to\infty} \left| \frac{(-1)^{n+2}(x-2)^{n+1}}{(n+1)\,2^{n+1}} \cdot \frac{n\,2^n}{(-1)^{n+1}(x-2)^n} \right| = \lim_{n\to\infty} \left| \frac{(-1)(x-2)n}{(n+1)2} \right| = \lim_{n\to\infty} \left(\frac{n}{n+1} \right) \frac{|x-2|}{2}$$

$$= \frac{|x-2|}{2} < 1 \quad \text{for convergence, so } |x-2| < 2 \text{ and } R = 2.$$

24.

n	$f^{(n)}(x)$	$f^{(n)}(-3)$
0	$1/x$	$-1/3$
1	$-1/x^2$	$-1/3^2$
2	$2/x^3$	$-2/3^3$
3	$-6/x^4$	$-6/3^4$
4	$24/x^5$	$-24/3^5$
$\vdots$	$\vdots$	$\vdots$

$$f(x) = \frac{1}{x} = \sum_{n=0}^{\infty} \frac{f^{(n)}(-3)}{n!}(x+3)^n$$

$$= \frac{-1/3}{0!}(x+3)^0 + \frac{-1/3^2}{1!}(x+3)^1 + \frac{-2/3^3}{2!}(x+3)^2 + \frac{-6/3^4}{3!}(x+3)^3 + \frac{-24/3^5}{4!}(x+3)^4 + \cdots$$

$$= \sum_{n=0}^{\infty} \frac{-n!/3^{n+1}}{n!}(x+3)^n = -\sum_{n=0}^{\infty} \frac{(x+3)^n}{3^{n+1}}$$

$$\lim_{n\to\infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n\to\infty} \left| \frac{(x+3)^{n+1}}{3^{n+2}} \cdot \frac{3^{n+1}}{(x+3)^n} \right| = \lim_{n\to\infty} \frac{|x+3|}{3} = \frac{|x+3|}{3} < 1 \quad \text{for convergence,}$$

so $|x+3| < 3$ and $R = 3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25.

n	$f^{(n)}(x)$	$f^{(n)}(3)$
0	e^{2x}	e^6
1	$2e^{2x}$	$2e^6$
2	2^2e^{2x}	$4e^6$
3	2^3e^{2x}	$8e^6$
4	2^4e^{2x}	$16e^6$
$\vdots$	$\vdots$	$\vdots$

$$\begin{aligned} f(x) = e^{2x} &= \sum_{n=0}^{\infty} \frac{f^{(n)}(3)}{n!}(x-3)^n \\ &= \frac{e^6}{0!}(x-3)^0 + \frac{2e^6}{1!}(x-3)^1 + \frac{4e^6}{2!}(x-3)^2 \\ &\qquad + \frac{8e^6}{3!}(x-3)^3 + \frac{16e^6}{4!}(x-3)^4 + \cdots \\ &= \sum_{n=0}^{\infty} \frac{2^n e^6}{n!}(x-3)^n \end{aligned}$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{2^{n+1}e^6(x-3)^{n+1}}{(n+1)!}\cdot\frac{n!}{2^n e^6(x-3)^n}\right| = \lim_{n\to\infty}\frac{2\,|x-3|}{n+1} = 0 < 1 \quad \text{for all } x\text{, so } R = \infty.$$

26.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$1/x^2$	1
1	$-2/x^3$	-2
2	$6/x^4$	6
3	$-24/x^5$	-24
4	$120/x^6$	120
$\vdots$	$\vdots$	$\vdots$

$$\begin{aligned} f(x) = \frac{1}{x^2} &= f(1) + \frac{f'(1)}{1!}(x-1) + \frac{f''(1)}{2!}(x-1)^2 + \frac{f'''(1)}{3!}(x-1)^3 + \frac{f^{(4)}(1)}{4!}(x-1)^4 + \cdots \\ &= 1 - \frac{2}{1!}(x-1) + \frac{6}{2!}(x-1)^2 - \frac{24}{3!}(x-1)^3 + \frac{120}{4!}(x-1)^4 - \cdots \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n(n+1)!}{n!}(x-1)^n = \sum_{n=0}^{\infty}(-1)^n(n+1)(x-1)^n \end{aligned}$$

$$\begin{aligned} \lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| &= \lim_{n\to\infty}\left|\frac{(-1)^{n+1}(n+2)(x-1)^{n+1}}{(-1)^n(n+1)(x-1)^n}\right| = \lim_{n\to\infty}\frac{(n+2)\,|x-1|}{(n+1)} = |x-1|\lim_{n\to\infty}\frac{1+2/n}{1+1/n} \\ &= |x-1| < 1 \text{ for convergence, so } R = 1. \end{aligned}$$

27.

n	$f^{(n)}(x)$	$f^{(n)}(\pi)$
0	$\sin x$	0
1	$\cos x$	-1
2	$-\sin x$	0
3	$-\cos x$	1
4	$\sin x$	0
5	$\cos x$	-1
6	$-\sin x$	0
7	$-\cos x$	1
$\vdots$	$\vdots$	$\vdots$

$$\begin{aligned} f(x) = \sin x &= \sum_{n=0}^{\infty} \frac{f^{(n)}(\pi)}{n!}(x-\pi)^n \\ &= \frac{-1}{1!}(x-\pi)^1 + \frac{1}{3!}(x-\pi)^3 + \frac{-1}{5!}(x-\pi)^5 + \frac{1}{7!}(x-\pi)^7 + \cdots \\ &= \sum_{n=0}^{\infty} \frac{(-1)^{n+1}}{(2n+1)!}(x-\pi)^{2n+1} \end{aligned}$$

$$\begin{aligned} \lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| &= \lim_{n\to\infty}\left|\frac{(-1)^{n+2}(x-\pi)^{2n+3}}{(2n+3)!}\cdot\frac{(2n+1)!}{(-1)^{n+1}(x-\pi)^{2n+1}}\right| \\ &= \lim_{n\to\infty}\frac{(x-\pi)^2}{(2n+3)(2n+2)} = 0 < 1 \quad \text{for all } x\text{, so } R = \infty. \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28.

n	$f^{(n)}(x)$	$f^{(n)}(\pi/2)$
0	$\cos x$	0
1	$-\sin x$	-1
2	$-\cos x$	0
3	$\sin x$	1
4	$\cos x$	0
5	$-\sin x$	-1
6	$-\cos x$	0
7	$\sin x$	1
$\vdots$	$\vdots$	$\vdots$

$$f(x) = \cos x = \sum_{n=0}^{\infty} \frac{f^{(n)}(\pi/2)}{n!}\left(x - \frac{\pi}{2}\right)^n$$

$$= \frac{-1}{1!}\left(x - \frac{\pi}{2}\right)^1 + \frac{1}{3!}\left(x - \frac{\pi}{2}\right)^3 + \frac{-1}{5!}\left(x - \frac{\pi}{2}\right)^5 + \frac{1}{7!}\left(x - \frac{\pi}{2}\right)^7 + \cdots$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n+1}}{(2n+1)!}\left(x - \frac{\pi}{2}\right)^{2n+1}$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^{n+2}\left(x - \frac{\pi}{2}\right)^{2n+3}}{(2n+3)!} \cdot \frac{(2n+1)!}{(-1)^{n+1}\left(x - \frac{\pi}{2}\right)^{2n+1}}\right|$$

$$= \lim_{n\to\infty} \frac{\left(x - \frac{\pi}{2}\right)^2}{(2n+3)(2n+2)} = 0 < 1 \quad \text{for all } x, \text{ so } R = \infty.$$

29.

n	$f^{(n)}(x)$	$f^{(n)}(\pi)$
0	$\sin 2x$	0
1	$2\cos 2x$	2
2	$-4\sin 2x$	0
3	$-8\cos 2x$	-8
4	$16\sin 2x$	0
5	$32\cos 2x$	32
$\vdots$	$\vdots$	$\vdots$

$$f(x) = \sin 2x = f(\pi) + \frac{f'(\pi)}{1!}(x-\pi) + \frac{f''(\pi)}{2!}(x-\pi)^2$$

$$+ \frac{f'''(\pi)}{3!}(x-\pi)^3 + \frac{f^{(4)}(\pi)}{4!}(x-\pi)^4 + \cdots$$

$$= 0 + \frac{2}{1!}(x-\pi) + 0 - \frac{8}{3!}(x-\pi)^3 + 0 + \frac{32}{5!}(x-\pi)^5 - \cdots$$

$$= \frac{2}{1!}(x-\pi) - \frac{8}{3!}(x-\pi)^3 + \frac{32}{5!}(x-\pi)^5 - \cdots$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n\, 2^{2n+1}}{(2n+1)!}(x-\pi)^{2n+1}$$

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^{n+1}\, 2^{2n+3}\,(x-\pi)^{2n+3}}{(2n+3)!} \cdot \frac{(2n+1)!}{(-1)^n\, 2^{2n+1}\,(x-\pi)^{2n+1}}\right|$$

$$= (x-\pi)^2 \lim_{n\to\infty} \frac{2^2}{(2n+3)(2n+2)} = 0 < 1 \text{ for all } x, \text{ so } R = \infty.$$

30.

n	$f^{(n)}(x)$	$f^{(n)}(16)$
0	$\sqrt{x}$	4
1	$\frac{1}{2}x^{-1/2}$	$\frac{1}{2}\cdot\frac{1}{4}$
2	$-\frac{1}{4}x^{-3/2}$	$-\frac{1}{4}\cdot\frac{1}{4^3}$
3	$\frac{3}{8}x^{-5/2}$	$\frac{3}{8}\cdot\frac{1}{4^5}$
4	$-\frac{15}{16}x^{-7/2}$	$-\frac{15}{16}\cdot\frac{1}{4^7}$
$\vdots$	$\vdots$	$\vdots$

$$f(x) = \sqrt{x} = \sum_{n=0}^{\infty} \frac{f^{(n)}(16)}{n!}(x-16)^n$$

$$= \frac{4}{0!}(x-16)^0 + \frac{1}{2}\cdot\frac{1}{4}\cdot\frac{1}{1!}(x-16)^1 - \frac{1}{4}\cdot\frac{1}{4^3}\cdot\frac{1}{2!}(x-16)^2$$

$$+ \frac{3}{8}\cdot\frac{1}{4^5}\cdot\frac{1}{3!}(x-16)^3 - \frac{15}{16}\cdot\frac{1}{4^7}\cdot\frac{1}{4!}(x-16)^4 + \cdots$$

$$= 4 + \frac{1}{8}(x-16) + \sum_{n=2}^{\infty}(-1)^{n-1}\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-3)}{2^n 4^{2n-1}\, n!}(x-16)^n$$

$$= 4 + \frac{1}{8}(x-16) + \sum_{n=2}^{\infty}(-1)^{n-1}\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-3)}{2^{5n-2}\, n!}(x-16)^n$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty}\left|\frac{(-1)^n\,1\cdot3\cdot5\cdot\cdots\cdot(2n-1)(x-16)^{n+1}}{2^{5n+3}(n+1)!}\cdot\frac{2^{5n-2}n!}{(-1)^{n-1}\,1\cdot3\cdot5\cdot\cdots\cdot(2n-3)(x-16)^n}\right|$$

$$= \lim_{n\to\infty}\frac{(2n-1)\,|x-16|}{2^5(n+1)} = \frac{|x-16|}{32}\lim_{n\to\infty}\frac{2-1/n}{1+1/n} = \frac{|x-16|}{32}\cdot 2$$

$$= \frac{|x-16|}{16} < 1 \quad \text{for convergence, so } |x-16| < 16 \text{ and } R = 16.$$

31. If $f(x) = \cos x$, then $f^{(n+1)}(x) = \pm\sin x$ or $\pm\cos x$. In each case, $\left|f^{(n+1)}(x)\right| \le 1$, so by Formula 9 with $a = 0$ and $M = 1$, $|R_n(x)| \le \dfrac{1}{(n+1)!}\,|x|^{n+1}$. Thus, $|R_n(x)| \to 0$ as $n \to \infty$ by Equation 10. So $\lim\limits_{n\to\infty} R_n(x) = 0$ and, by Theorem 8, the series in Exercise 13 represents $\cos x$ for all x.

32. If $f(x) = \sin x$, then $f^{(n+1)}(x) = \pm\sin x$ or $\pm\cos x$. In each case, $\left|f^{(n+1)}(x)\right| \le 1$, so by Formula 9 with $a = 0$ and $M = 1$, $|R_n(x)| \le \dfrac{1}{(n+1)!}\,|x-\pi|^{n+1}$. Thus, $|R_n(x)| \to 0$ as $n \to \infty$ by Equation 10. So $\lim\limits_{n\to\infty} R_n(x) \to 0$ and, by Theorem 8, the series in Exercise 27 represents $\sin x$ for all x.

33. If $f(x) = \sinh x$, then for all n, $f^{(n+1)}(x) = \cosh x$ or $\sinh x$. Since $|\sinh x| < |\cosh x| = \cosh x$ for all x, we have $\left|f^{(n+1)}(x)\right| \le \cosh x$ for all n. If d is any positive number and $|x| \le d$, then $\left|f^{(n+1)}(x)\right| \le \cosh x \le \cosh d$, so by Formula 9 with $a = 0$ and $M = \cosh d$, we have $|R_n(x)| \le \dfrac{\cosh d}{(n+1)!}\,|x|^{n+1}$. It follows that $|R_n(x)| \to 0$ as $n \to \infty$ for $|x| \le d$ (by Equation 10). But d was an arbitrary positive number. So by Theorem 8, the series represents $\sinh x$ for all x.

34. If $f(x) = \cosh x$, then for all n, $f^{(n+1)}(x) = \cosh x$ or $\sinh x$. Since $|\sinh x| < |\cosh x| = \cosh x$ for all x, we have $\left|f^{(n+1)}(x)\right| \le \cosh x$ for all n. If d is any positive number and $|x| \le d$, then $\left|f^{(n+1)}(x)\right| \le \cosh x \le \cosh d$, so by Formula 9 with $a = 0$ and $M = \cosh d$, we have $|R_n(x)| \le \dfrac{\cosh d}{(n+1)!}\,|x|^{n+1}$. It follows that $|R_n(x)| \to 0$ as $n \to \infty$ for $|x| \le d$ (by Equation 10). But d was an arbitrary positive number. So by Theorem 8, the series represents $\cosh x$ for all x.

35. $$\sqrt[4]{1-x} = [1+(-x)]^{1/4} = \sum_{n=0}^{\infty}\binom{1/4}{n}(-x)^n$$

$$= 1 + \tfrac{1}{4}(-x) + \frac{\frac{1}{4}\left(-\frac{3}{4}\right)}{2!}(-x)^2 + \frac{\frac{1}{4}\left(-\frac{3}{4}\right)\left(-\frac{7}{4}\right)}{3!}(-x)^3 + \cdots$$

$$= 1 - \frac{1}{4}x + \sum_{n=2}^{\infty}\frac{(-1)^{n-1}(-1)^n\cdot[3\cdot7\cdot\cdots\cdot(4n-5)]}{4^n\cdot n!}x^n$$

$$= 1 - \frac{1}{4}x - \sum_{n=2}^{\infty}\frac{3\cdot7\cdot\cdots\cdot(4n-5)}{4^n\cdot n!}x^n$$

and $|-x| < 1 \quad\Leftrightarrow\quad |x| < 1$, so $R = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. $\sqrt[3]{8+x} = \sqrt[3]{8\left(1+\dfrac{x}{8}\right)} = 2\left(1+\dfrac{x}{8}\right)^{1/3} = 2\sum\limits_{n=0}^{\infty}\dbinom{1/3}{n}\left(\dfrac{x}{8}\right)^n$

$$= 2\left[1+\frac{1}{3}\left(\frac{x}{8}\right)+\frac{\frac{1}{3}\left(-\frac{2}{3}\right)}{2!}\left(\frac{x}{8}\right)^2+\frac{\frac{1}{3}\left(-\frac{2}{3}\right)\left(-\frac{5}{3}\right)}{3!}\left(\frac{x}{8}\right)^3+\cdots\right]$$

$$= 2\left[1+\frac{1}{24}x+\sum_{n=2}^{\infty}\frac{(-1)^{n-1}\cdot[2\cdot 5\cdot\cdots\cdot(3n-4)]}{3^n\cdot 8^n\cdot n!}\,x^n\right]$$

$$= 2+\frac{1}{12}x+2\sum_{n=2}^{\infty}\frac{(-1)^{n-1}[2\cdot 5\cdot\cdots\cdot(3n-4)]}{24^n\cdot n!}\,x^n$$

and $\left|\dfrac{x}{8}\right| < 1 \quad\Leftrightarrow\quad |x| < 8$, so $R = 8$.

37. $\dfrac{1}{(2+x)^3} = \dfrac{1}{[2(1+x/2)]^3} = \dfrac{1}{8}\left(1+\dfrac{x}{2}\right)^{-3} = \dfrac{1}{8}\sum\limits_{n=0}^{\infty}\dbinom{-3}{n}\left(\dfrac{x}{2}\right)^n$. The binomial coefficient is

$$\binom{-3}{n} = \frac{(-3)(-4)(-5)\cdot\cdots\cdot(-3-n+1)}{n!} = \frac{(-3)(-4)(-5)\cdot\cdots\cdot[-(n+2)]}{n!}$$

$$= \frac{(-1)^n\cdot 2\cdot 3\cdot 4\cdot 5\cdot\cdots\cdot(n+1)(n+2)}{2\cdot n!} = \frac{(-1)^n(n+1)(n+2)}{2}$$

Thus, $\dfrac{1}{(2+x)^3} = \dfrac{1}{8}\sum\limits_{n=0}^{\infty}\dfrac{(-1)^n(n+1)(n+2)}{2}\dfrac{x^n}{2^n} = \sum\limits_{n=0}^{\infty}\dfrac{(-1)^n(n+1)(n+2)x^n}{2^{n+4}}$ for $\left|\dfrac{x}{2}\right| < 1 \quad\Leftrightarrow\quad |x| < 2$, so $R = 2$.

38. $(1-x)^{3/4} = [1+(-x)]^{3/4} = \sum\limits_{n=0}^{\infty}\dbinom{3/4}{n}(-x)^n = 1+\dfrac{3}{4}(-x)+\dfrac{\frac{3}{4}\left(-\frac{1}{4}\right)}{2!}(-x)^2+\dfrac{\frac{3}{4}\left(-\frac{1}{4}\right)\left(-\frac{5}{4}\right)}{3!}(-x)^3+\cdots$

$$= 1-\frac{3}{4}x+\sum_{n=2}^{\infty}\frac{(-1)^{n-1}(-1)^n\cdot 3\cdot[1\cdot 5\cdot 9\cdot\cdots\cdot(4n-7)]}{4^n\cdot n!}\,x^n$$

$$= 1-\frac{3}{4}x-3\sum_{n=2}^{\infty}\frac{1\cdot 5\cdot 9\cdot\cdots\cdot(4n-7)}{4^n\cdot n!}\,x^n$$

and $|-x| < 1 \quad\Leftrightarrow\quad |x| < 1$, so $R = 1$.

39. $\arctan x = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{x^{2n+1}}{2n+1}$, so $f(x) = \arctan(x^2) = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{\left(x^2\right)^{2n+1}}{2n+1} = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{1}{2n+1}x^{4n+2}$, $R = 1$.

40. $\sin x = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{x^{2n+1}}{(2n+1)!}$, so $f(x) = \sin\left(\dfrac{\pi}{4}x\right) = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{\left(\frac{\pi}{4}x\right)^{2n+1}}{(2n+1)!} = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{\pi^{2n+1}}{4^{2n+1}(2n+1)!}x^{2n+1}$, $R = \infty$.

41. $\cos x = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{x^{2n}}{(2n)!} \quad\Rightarrow\quad \cos 2x = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{(2x)^{2n}}{(2n)!} = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{2^{2n}x^{2n}}{(2n)!}$, so

$f(x) = x\cos 2x = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{2^{2n}}{(2n)!}x^{2n+1}$, $R = \infty$.

42. $e^x = \sum\limits_{n=0}^{\infty}\dfrac{x^n}{n!}$, so $f(x) = e^{3x}-e^{2x} = \sum\limits_{n=0}^{\infty}\dfrac{(3x)^n}{n!}-\sum\limits_{n=0}^{\infty}\dfrac{(2x)^n}{n!} = \sum\limits_{n=0}^{\infty}\dfrac{3^nx^n}{n!}-\sum\limits_{n=0}^{\infty}\dfrac{2^nx^n}{n!} = \sum\limits_{n=0}^{\infty}\dfrac{3^n-2^n}{n!}x^n$, $R = \infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. $\cos x = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{2n}}{(2n)!} \quad \Rightarrow \quad \cos\left(\frac{1}{2}x^2\right) = \sum_{n=0}^{\infty} (-1)^n \dfrac{\left(\frac{1}{2}x^2\right)^{2n}}{(2n)!} = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{4n}}{2^{2n}\,(2n)!}$, so

$f(x) = x\cos\left(\frac{1}{2}x^2\right) = \sum_{n=0}^{\infty} (-1)^n \dfrac{1}{2^{2n}(2n)!}\, x^{4n+1},\ R = \infty.$

44. $\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} \dfrac{x^n}{n} \quad \Rightarrow \quad \ln(1+x^3) = \sum_{n=1}^{\infty} (-1)^{n-1} \dfrac{x^{3n}}{n}$, so $f(x) = x^2 \ln(1+x^3) = \sum_{n=1}^{\infty} (-1)^{n-1} \dfrac{x^{3n+2}}{n}$,

$R = 1.$

45. We must write the binomial in the form (1+ expression), so we'll factor out a 4.

$$\frac{x}{\sqrt{4+x^2}} = \frac{x}{\sqrt{4(1+x^2/4)}} = \frac{x}{2\sqrt{1+x^2/4}} = \frac{x}{2}\left(1+\frac{x^2}{4}\right)^{-1/2} = \frac{x}{2}\sum_{n=0}^{\infty}\binom{-\frac{1}{2}}{n}\left(\frac{x^2}{4}\right)^n$$

$$= \frac{x}{2}\left[1 + \left(-\tfrac{1}{2}\right)\frac{x^2}{4} + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}\left(\frac{x^2}{4}\right)^2 + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{3!}\left(\frac{x^2}{4}\right)^3 + \cdots\right]$$

$$= \frac{x}{2} + \frac{x}{2}\sum_{n=1}^{\infty}(-1)^n \frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{2^n\cdot 4^n\cdot n!}x^{2n}$$

$$= \frac{x}{2} + \sum_{n=1}^{\infty}(-1)^n \frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{n!\,2^{3n+1}}x^{2n+1} \text{ and } \frac{x^2}{4} < 1 \quad \Leftrightarrow \quad \frac{|x|}{2} < 1 \quad \Leftrightarrow \quad |x| < 2, \quad \text{so } R = 2.$$

46. $$\frac{x^2}{\sqrt{2+x}} = \frac{x^2}{\sqrt{2\,(1+x/2)}} = \frac{x^2}{\sqrt{2}}\left(1+\frac{x}{2}\right)^{-1/2} = \frac{x^2}{\sqrt{2}}\sum_{n=0}^{\infty}\binom{-\frac{1}{2}}{n}\left(\frac{x}{2}\right)^n$$

$$= \frac{x^2}{\sqrt{2}}\left[1 + \left(-\tfrac{1}{2}\right)\left(\frac{x}{2}\right) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}\left(\frac{x}{2}\right)^2 + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{3!}\left(\frac{x}{2}\right)^3 + \cdots\right]$$

$$= \frac{x^2}{\sqrt{2}} + \frac{x^2}{\sqrt{2}}\sum_{n=1}^{\infty}(-1)^n \frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{n!\,2^{2n}}x^n$$

$$= \frac{x^2}{\sqrt{2}} + \sum_{n=1}^{\infty}(-1)^n \frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{n!\,2^{2n+1/2}}x^{n+2} \text{ and } \left|\frac{x}{2}\right| < 1 \quad \Leftrightarrow \quad |x| < 2, \quad \text{so } R = 2.$$

47. $\sin^2 x = \dfrac{1}{2}(1-\cos 2x) = \dfrac{1}{2}\left[1 - \sum_{n=0}^{\infty}\dfrac{(-1)^n(2x)^{2n}}{(2n)!}\right] = \dfrac{1}{2}\left[1 - 1 - \sum_{n=1}^{\infty}\dfrac{(-1)^n(2x)^{2n}}{(2n)!}\right] = \sum_{n=1}^{\infty}\dfrac{(-1)^{n+1}2^{2n-1}x^{2n}}{(2n)!}$,

$R = \infty$

48. $$\frac{x-\sin x}{x^3} = \frac{1}{x^3}\left[x - \sum_{n=0}^{\infty}\frac{(-1)^n x^{2n+1}}{(2n+1)!}\right] = \frac{1}{x^3}\left[x - x - \sum_{n=1}^{\infty}\frac{(-1)^n x^{2n+1}}{(2n+1)!}\right] = \frac{1}{x^3}\left[-\sum_{n=0}^{\infty}\frac{(-1)^{n+1}x^{2n+3}}{(2n+3)!}\right]$$

$$= \frac{1}{x^3}\sum_{n=0}^{\infty}\frac{(-1)^n x^{2n+3}}{(2n+3)!} = \sum_{n=0}^{\infty}\frac{(-1)^n x^{2n}}{(2n+3)!}$$

and this series also gives the required value at $x = 0$ (namely $1/6$); $R = \infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. (a) The Maclaurin series for e^x is $e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \cdots$, so

$$e^{-x} = 1 - x + \frac{1}{2!}(-x)^2 + \frac{1}{3!}(-x)^3 + \cdots = 1 - x + \frac{1}{2!}x^2 - \frac{1}{3!}x^3 + \cdots, \text{ and}$$

$$\sinh x = \frac{1}{2}(e^x - e^{-x}) = \frac{1}{2}\left[\left(1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \cdots\right) - \left(1 - x + \frac{1}{2!}x^2 - \frac{1}{3!}x^3 + \cdots\right)\right]$$

$$= \frac{1}{2}\left[(1-1) + (x+x) + \left(\frac{1}{2!}x^2 - \frac{1}{2!}x^2\right) + \left(\frac{1}{3!}x^3 + \frac{1}{3!}x^3\right) + \cdots\right]$$

$$= \frac{1}{2}\left[2x + \frac{2}{3!}x^3 + \frac{2}{5!}x^5 + \cdots\right] = x + \frac{1}{3!}x^3 + \frac{1}{5!}x^5 + \cdots = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!}$$

(b) $$\cosh x = \frac{1}{2}(e^x + e^{-x}) = \frac{1}{2}\left[\left(1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \cdots\right) + \left(1 - x + \frac{1}{2!}x^2 - \frac{1}{3!}x^3 + \cdots\right)\right]$$

$$= \frac{1}{2}\left[(1+1) + (x-x) + \left(\frac{1}{2!}x^2 + \frac{1}{2!}x^2\right) + \left(\frac{1}{3!}x^3 - \frac{1}{3!}x^3\right) + \cdots\right]$$

$$= \frac{1}{2}\left[2 + \frac{2}{2!}x^2 + \frac{2}{4!}x^4 + \cdots\right] = 1 + \frac{1}{2!}x^2 + \frac{1}{4!}x^4 + \cdots = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!}$$

50. From Table (1), $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots$, so

$$\tanh^{-1} x = \frac{1}{2}\ln\left(\frac{1+x}{1-x}\right) = \frac{1}{2}[\ln(1+x) - \ln(1-x)]$$

$$= \frac{1}{2}\left[\left(x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots\right) - \left(-x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \cdots\right)\right]$$

$$= \frac{1}{2}\left[(x+x) + \left(\frac{x^2}{2} - \frac{x^2}{2}\right) + \left(\frac{x^3}{3} + \frac{x^3}{3}\right) + \left(\frac{x^4}{4} - \frac{x^4}{4}\right) + \cdots\right]$$

$$= \frac{1}{2}\left[2x + \frac{2x^3}{3} + \frac{2x^5}{5} + \cdots\right] = x + \frac{x^3}{3} + \frac{x^5}{5} + \cdots = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{2n+1}$$

51. $\cos x \overset{(16)}{=} \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \quad \Rightarrow$

$$f(x) = \cos(x^2) = \sum_{n=0}^{\infty} \frac{(-1)^n (x^2)^{2n}}{(2n)!} = \sum_{n=0}^{\infty} \frac{(-1)^n x^{4n}}{(2n)!}$$

$$= 1 - \tfrac{1}{2}x^4 + \tfrac{1}{24}x^8 - \tfrac{1}{720}x^{12} + \cdots$$

The series for $\cos x$ converges for all x, so the same is true of the series for $f(x)$, that is, $R = \infty$. Notice that, as n increases, $T_n(x)$ becomes a better approximation to $f(x)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

52. $\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} \dfrac{x^n}{n} \quad \Rightarrow$

$$f(x) = \ln(1+x^2) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}(x^2)^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}x^{2n}}{n}$$
$$= x^2 - \tfrac{1}{2}x^4 + \tfrac{1}{3}x^6 - \tfrac{1}{4}x^8 + \cdots$$

The series for $\ln(1+x)$ has $R = 1$ and $\left|x^2\right| < 1 \quad \Leftrightarrow \quad |x| < 1$, so the series for $f(x)$ also has $R = 1$. From the graphs of f and the first few Taylor polynomials, we see that $T_n(x)$ provides a closer fit to $f(x)$ near 0 as n increases.

53. $e^x \overset{(11)}{=} \sum_{n=0}^{\infty} \dfrac{x^n}{n!}$, so $e^{-x} = \sum_{n=0}^{\infty} \dfrac{(-x)^n}{n!} = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^n}{n!}$, so

$$f(x) = xe^{-x} = \sum_{n=0}^{\infty} (-1)^n \frac{1}{n!} x^{n+1}$$
$$= x - x^2 + \tfrac{1}{2}x^3 - \tfrac{1}{6}x^4 + \tfrac{1}{24}x^5 - \tfrac{1}{120}x^6 + \cdots$$
$$= \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{(n-1)!}$$

The series for e^x converges for all x, so the same is true of the series for $f(x)$; that is, $R = \infty$. From the graphs of f and the first few Taylor polynomials, we see that $T_n(x)$ provides a closer fit to $f(x)$ near 0 as n increases.

54. From Table 1, $\tan^{-1} x = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{2n+1}}{2n+1}$, so

$$f(x) = \tan^{-1}(x^3) = \sum_{n=0}^{\infty} (-1)^n \frac{(x^3)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{6n+3}}{2n+1}$$
$$= x^3 - \tfrac{1}{3}x^9 + \tfrac{1}{5}x^{15} - \tfrac{1}{7}x^{21} + \cdots$$

The series for $\tan^{-1} x$ has $R = 1$ and $\left|x^3\right| < 1 \quad \Leftrightarrow \quad |x| < 1$, so the series for $f(x)$ also has $R = 1$. From the graphs of f and the first few Taylor polynomials, we see that $T_n(x)$ provides a closer fit to $f(x)$ near 0 as n increases.

55. $5^\circ = 5^\circ \left(\dfrac{\pi}{180^\circ}\right) = \dfrac{\pi}{36}$ radians and $\cos x = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{2n}}{(2n)!} = 1 - \dfrac{x^2}{2!} + \dfrac{x^4}{4!} - \dfrac{x^6}{6!} + \cdots$, so

$\cos \dfrac{\pi}{36} = 1 - \dfrac{(\pi/36)^2}{2!} + \dfrac{(\pi/36)^4}{4!} - \dfrac{(\pi/36)^6}{6!} + \cdots$. Now $1 - \dfrac{(\pi/36)^2}{2!} \approx 0.99619$ and adding $\dfrac{(\pi/36)^4}{4!} \approx 2.4 \times 10^{-6}$ does not affect the fifth decimal place, so $\cos 5^\circ \approx 0.99619$ by the Alternating Series Estimation Theorem.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

56. $1/\sqrt[10]{e} = e^{-1/10}$ and $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$, so

$e^{-1/10} = 1 - \frac{1}{10} + \frac{(1/10)^2}{2!} - \frac{(1/10)^3}{3!} + \frac{(1/10)^4}{4!} - \frac{(1/10)^5}{5!} + \cdots$. Now

$1 - \frac{1}{10} + \frac{(1/10)^2}{2!} - \frac{(1/10)^3}{3!} + \frac{(1/10)^4}{4!} \approx 0.90484$ and subtracting $\frac{(1/10)^5}{5!} \approx 8.3 \times 10^{-8}$ does not affect the fifth decimal place, so $e^{-1/10} \approx 0.90484$ by the Alternating Series Estimation Theorem.

57. (a) $1/\sqrt{1-x^2} = \left[1 + \left(-x^2\right)\right]^{-1/2} = 1 + \left(-\frac{1}{2}\right)\left(-x^2\right) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}\left(-x^2\right)^2 + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{3!}\left(-x^2\right)^3 + \cdots$

$$= 1 + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{2^n \cdot n!} x^{2n}$$

(b) $\sin^{-1} x = \int \frac{1}{\sqrt{1-x^2}}\, dx = C + x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{(2n+1)2^n \cdot n!} x^{2n+1}$

$$= x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2n-1)}{(2n+1)2^n \cdot n!} x^{2n+1} \quad \text{since } 0 = \sin^{-1} 0 = C.$$

58. (a) $1/\sqrt[4]{1+x} = (1+x)^{-1/4} = \sum_{n=0}^{\infty} \binom{-\frac{1}{4}}{n} x^n = 1 - \frac{1}{4}x + \frac{\left(-\frac{1}{4}\right)\left(-\frac{5}{4}\right)}{2!}x^2 + \frac{\left(-\frac{1}{4}\right)\left(-\frac{5}{4}\right)\left(-\frac{9}{4}\right)}{3!}x^3 + \cdots$

$$= 1 - \frac{1}{4}x + \sum_{n=2}^{\infty} (-1)^n \frac{1 \cdot 5 \cdot 9 \cdot \cdots \cdot (4n-3)}{4^n \cdot n!} x^n$$

(b) $1/\sqrt[4]{1+x} = 1 - \frac{1}{4}x + \frac{5}{32}x^2 - \frac{15}{128}x^3 + \frac{195}{2048}x^4 - \cdots$. $1/\sqrt[4]{1.1} = 1/\sqrt[4]{1+0.1}$, so let $x = 0.1$. The sum of the first four terms is then $1 - \frac{1}{4}(0.1) + \frac{5}{32}(0.1)^2 - \frac{15}{128}(0.1)^3 \approx 0.976$. The fifth term is $\frac{195}{2048}(0.1)^4 \approx 0.000\,009\,5$, which does not affect the third decimal place of the sum, so we have $1/\sqrt[4]{1.1} \approx 0.976$. (Note that the third decimal place of the sum of the first three terms is affected by the fourth term, so we need to use more than three terms for the sum.)

59. $\sqrt{1+x^3} = (1+x^3)^{1/2} = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} (x^3)^n = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} x^{3n} \quad \Rightarrow \quad \int \sqrt{1+x^3}\, dx = C + \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} \frac{x^{3n+1}}{3n+1}$,

with $R = 1$.

60. $\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} \quad \Rightarrow \quad \sin(x^2) = \sum_{n=0}^{\infty} (-1)^n \frac{(x^2)^{2n+1}}{(2n+1)!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{(2n+1)!} \quad \Rightarrow$

$x^2 \sin(x^2) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+4}}{(2n+1)!} \quad \Rightarrow \quad \int x^2 \sin(x^2)\, dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+5}}{(2n+1)!(4n+5)}$, with $R = \infty$.

61. $\cos x \overset{(16)}{=} \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \quad \Rightarrow \quad \cos x - 1 = \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \quad \Rightarrow \quad \frac{\cos x - 1}{x} = \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n-1}}{(2n)!} \quad \Rightarrow$

$\int \frac{\cos x - 1}{x}\, dx = C + \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n}}{2n \cdot (2n)!}$, with $R = \infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

62. $\arctan x = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{2n+1}}{2n+1} \quad \Rightarrow \quad \arctan(x^2) = \sum_{n=0}^{\infty} (-1)^n \dfrac{(x^2)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{4n+2}}{2n+1} \quad \Rightarrow$

$\displaystyle\int \arctan(x^2)\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+3}}{(2n+1)(4n+3)}$, with $R = 1$.

63. $\arctan x = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{2n+1}}{2n+1}$ for $|x| < 1$, so $x^3 \arctan x = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{2n+4}}{2n+1}$ for $|x| < 1$ and

$\displaystyle\int x^3 \arctan x\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+5}}{(2n+1)(2n+5)}$. Since $\frac{1}{2} < 1$, we have

$\displaystyle\int_0^{1/2} x^3 \arctan x\,dx = \sum_{n=0}^{\infty} (-1)^n \frac{(1/2)^{2n+5}}{(2n+1)(2n+5)} = \frac{(1/2)^5}{1\cdot 5} - \frac{(1/2)^7}{3\cdot 7} + \frac{(1/2)^9}{5\cdot 9} - \frac{(1/2)^{11}}{7\cdot 11} + \cdots$. Now

$\dfrac{(1/2)^5}{1\cdot 5} - \dfrac{(1/2)^7}{3\cdot 7} + \dfrac{(1/2)^9}{5\cdot 9} \approx 0.0059$ and subtracting $\dfrac{(1/2)^{11}}{7\cdot 11} \approx 6.3 \times 10^{-6}$ does not affect the fourth decimal place,

so $\int_0^{1/2} x^3 \arctan x\,dx \approx 0.0059$ by the Alternating Series Estimation Theorem.

64. $\sin x = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{2n+1}}{(2n+1)!}$ for all x, so $\sin(x^4) = \sum_{n=0}^{\infty} (-1)^n \dfrac{x^{8n+4}}{(2n+1)!}$ for all x and

$\displaystyle\int \sin(x^4)\,dx = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{8n+5}}{(2n+1)!\,(8n+5)}$. Thus,

$\displaystyle\int_0^1 \sin(x^4)\,dx = \sum_{n=0}^{\infty} (-1)^n \frac{1}{(2n+1)!\,(8n+5)} = \frac{1}{1!\cdot 5} - \frac{1}{3!\cdot 13} + \frac{1}{5!\cdot 21} - \frac{1}{7!\cdot 29} + \cdots$. Now

$\dfrac{1}{1!\cdot 5} - \dfrac{1}{3!\cdot 13} + \dfrac{1}{5!\cdot 21} \approx 0.1876$ and subtracting $\dfrac{1}{7!\cdot 29} \approx 6.84 \times 10^{-6}$ does not affect the fourth decimal place, so

$\int_0^1 \sin(x^4)\,dx \approx 0.1876$ by the Alternating Series Estimation Theorem.

65. $\sqrt{1+x^4} = (1+x^4)^{1/2} = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} (x^4)^n$, so $\displaystyle\int \sqrt{1+x^4}\,dx = C + \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} \frac{x^{4n+1}}{4n+1}$ and hence, since $0.4 < 1$,

we have

$$\begin{aligned} I &= \int_0^{0.4} \sqrt{1+x^4}\,dx = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} \frac{(0.4)^{4n+1}}{4n+1} \\ &= (1)\,\frac{(0.4)^1}{0!} + \frac{\frac{1}{2}}{1!}\,\frac{(0.4)^5}{5} + \frac{\frac{1}{2}\left(-\frac{1}{2}\right)}{2!}\,\frac{(0.4)^9}{9} + \frac{\frac{1}{2}\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{3!}\,\frac{(0.4)^{13}}{13} + \frac{\frac{1}{2}\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{4!}\,\frac{(0.4)^{17}}{17} + \cdots \\ &= 0.4 + \frac{(0.4)^5}{10} - \frac{(0.4)^9}{72} + \frac{(0.4)^{13}}{208} - \frac{5(0.4)^{17}}{2176} + \cdots \end{aligned}$$

Now $\dfrac{(0.4)^9}{72} \approx 3.6 \times 10^{-6} < 5 \times 10^{-6}$, so by the Alternating Series Estimation Theorem, $I \approx 0.4 + \dfrac{(0.4)^5}{10} \approx 0.40102$

(correct to five decimal places).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

66. $\displaystyle\int_0^{0.5} x^2 e^{-x^2}\,dx = \int_0^{0.5} \sum_{n=0}^{\infty} \frac{(-1)^n\, x^{2n+2}}{n!}\,dx = \sum_{n=0}^{\infty} \left[\frac{(-1)^n\, x^{2n+3}}{n!(2n+3)}\right]_0^{1/2} = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!(2n+3)2^{2n+3}}$ and since the term with $n=2$ is $\dfrac{1}{1792} < 0.001$, we use $\displaystyle\sum_{n=0}^{1} \frac{(-1)^n}{n!(2n+3)2^{2n+3}} = \frac{1}{24} - \frac{1}{160} \approx 0.0354$.

67. $$\lim_{x\to 0} \frac{x-\ln(1+x)}{x^2} = \lim_{x\to 0} \frac{x-(x-\frac{1}{2}x^2+\frac{1}{3}x^3-\frac{1}{4}x^4+\frac{1}{5}x^5-\cdots)}{x^2} = \lim_{x\to 0} \frac{\frac{1}{2}x^2-\frac{1}{3}x^3+\frac{1}{4}x^4-\frac{1}{5}x^5+\cdots}{x^2}$$
$$= \lim_{x\to 0}\left(\tfrac{1}{2}-\tfrac{1}{3}x+\tfrac{1}{4}x^2-\tfrac{1}{5}x^3+\cdots\right) = \tfrac{1}{2}$$

since power series are continuous functions.

68. $$\lim_{x\to 0} \frac{1-\cos x}{1+x-e^x} = \lim_{x\to 0} \frac{1-\left(1-\frac{1}{2!}x^2+\frac{1}{4!}x^4-\frac{1}{6!}x^6+\cdots\right)}{1+x-\left(1+x+\frac{1}{2!}x^2+\frac{1}{3!}x^3+\frac{1}{4!}x^4+\frac{1}{5!}x^5+\frac{1}{6!}x^6+\cdots\right)}$$
$$= \lim_{x\to 0} \frac{\frac{1}{2!}x^2-\frac{1}{4!}x^4+\frac{1}{6!}x^6-\cdots}{-\frac{1}{2!}x^2-\frac{1}{3!}x^3-\frac{1}{4!}x^4-\frac{1}{5!}x^5-\frac{1}{6!}x^6-\cdots}$$
$$= \lim_{x\to 0} \frac{\frac{1}{2!}-\frac{1}{4!}x^2+\frac{1}{6!}x^4-\cdots}{-\frac{1}{2!}-\frac{1}{3!}x-\frac{1}{4!}x^2-\frac{1}{5!}x^3-\frac{1}{6!}x^4-\cdots} = \frac{\frac{1}{2}-0}{-\frac{1}{2}-0} = -1$$

since power series are continuous functions.

69. $$\lim_{x\to 0} \frac{\sin x - x + \frac{1}{6}x^3}{x^5} = \lim_{x\to 0} \frac{\left(x-\frac{1}{3!}x^3+\frac{1}{5!}x^5-\frac{1}{7!}x^7+\cdots\right)-x+\frac{1}{6}x^3}{x^5}$$
$$= \lim_{x\to 0} \frac{\frac{1}{5!}x^5-\frac{1}{7!}x^7+\cdots}{x^5} = \lim_{x\to 0}\left(\frac{1}{5!}-\frac{x^2}{7!}+\frac{x^4}{9!}-\cdots\right) = \frac{1}{5!} = \frac{1}{120}$$

since power series are continuous functions.

70. $$\lim_{x\to 0} \frac{\sqrt{1+x}-1-\frac{1}{2}x}{x^2} = \lim_{x\to 0} \frac{\left(1+\frac{1}{2}x-\frac{1}{8}x^2+\frac{1}{16}x^3-\cdots\right)-1-\frac{1}{2}x}{x^2} = \lim_{x\to 0} \frac{-\frac{1}{8}x^2+\frac{1}{16}x^3-\cdots}{x^2}$$
$$= \lim_{x\to 0}\left(-\tfrac{1}{8}+\tfrac{1}{16}x-\cdots\right) = -\tfrac{1}{8} \quad \text{since power series are continuous functions.}$$

71. $$\lim_{x\to 0} \frac{x^3-3x+3\tan^{-1}x}{x^5} = \lim_{x\to 0} \frac{x^3-3x+3\left(x-\frac{1}{3}x^3+\frac{1}{5}x^5-\frac{1}{7}x^7+\cdots\right)}{x^5}$$
$$= \lim_{x\to 0} \frac{x^3-3x+3x-x^3+\frac{3}{5}x^5-\frac{3}{7}x^7+\cdots}{x^5} = \lim_{x\to 0} \frac{\frac{3}{5}x^5-\frac{3}{7}x^7+\cdots}{x^5}$$
$$= \lim_{x\to 0}\left(\tfrac{3}{5}-\tfrac{3}{7}x^2+\cdots\right) = \tfrac{3}{5} \quad \text{since power series are continuous functions.}$$

72. $$\lim_{x\to 0} \frac{\tan x - x}{x^3} = \lim_{x\to 0} \frac{\left(x+\frac{1}{3}x^3+\frac{2}{15}x^5+\cdots\right)-x}{x^3} = \lim_{x\to 0} \frac{\frac{1}{3}x^3+\frac{2}{15}x^5+\cdots}{x^3} = \lim_{x\to 0}\left(\tfrac{1}{3}+\tfrac{2}{15}x^2+\cdots\right) = \tfrac{1}{3}$$

since power series are continuous functions.

73. From Equation 11, we have $e^{-x^2} = 1 - \dfrac{x^2}{1!} + \dfrac{x^4}{2!} - \dfrac{x^6}{3!} + \cdots$ and we know that $\cos x = 1 - \dfrac{x^2}{2!} + \dfrac{x^4}{4!} - \cdots$ from Equation 16. Therefore, $e^{-x^2}\cos x = \left(1-x^2+\frac{1}{2}x^4-\cdots\right)\left(1-\frac{1}{2}x^2+\frac{1}{24}x^4-\cdots\right)$. Writing only the terms with degree ≤ 4, we get $e^{-x^2}\cos x = 1-\frac{1}{2}x^2+\frac{1}{24}x^4-x^2+\frac{1}{2}x^4+\frac{1}{2}x^4+\cdots = 1-\frac{3}{2}x^2+\frac{25}{24}x^4+\cdots$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

74. $\sec x = \dfrac{1}{\cos x} \overset{(16)}{=} \dfrac{1}{1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \cdots}$.

$$
\begin{array}{r|l}
 & 1 + \frac{1}{2}x^2 + \frac{5}{24}x^4 + \cdots \\
\hline
1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \cdots & 1 \\
 & 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \cdots \\
\hline
 & \frac{1}{2}x^2 - \frac{1}{24}x^4 + \cdots \\
 & \frac{1}{2}x^2 - \frac{1}{4}x^4 + \cdots \\
\hline
 & \frac{5}{24}x^4 + \cdots \\
 & \frac{5}{24}x^4 + \cdots \\
\hline
 & \cdots
\end{array}
$$

From the long division above, $\sec x = 1 + \frac{1}{2}x^2 + \frac{5}{24}x^4 + \cdots$.

75. $\dfrac{x}{\sin x} \overset{(15)}{=} \dfrac{x}{x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \cdots}$.

$$
\begin{array}{r|l}
 & 1 + \frac{1}{6}x^2 + \frac{7}{360}x^4 + \cdots \\
\hline
x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \cdots & x \\
 & x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \cdots \\
\hline
 & \frac{1}{6}x^3 - \frac{1}{120}x^5 + \cdots \\
 & \frac{1}{6}x^3 - \frac{1}{36}x^5 + \cdots \\
\hline
 & \frac{7}{360}x^5 + \cdots \\
 & \frac{7}{360}x^5 + \cdots \\
\hline
 & \cdots
\end{array}
$$

From the long division above, $\dfrac{x}{\sin x} = 1 + \frac{1}{6}x^2 + \frac{7}{360}x^4 + \cdots$.

76. From Table 1, we have $e^x = 1 + \dfrac{x}{1!} + \dfrac{x^2}{2!} + \dfrac{x^3}{3!} + \cdots$ and that $\ln(1+x) = x - \dfrac{x^2}{2} + \dfrac{x^3}{3} - \dfrac{x^4}{4} + \cdots$. Therefore, $y = e^x \ln(1+x) = \left(1 + \dfrac{x}{1!} + \dfrac{x^2}{2!} + \dfrac{x^3}{3!} + \cdots\right)\left(x - \dfrac{x^2}{2} + \dfrac{x^3}{3} - \dfrac{x^4}{4} + \cdots\right)$. Writing only terms with degree ≤ 3, we get $e^x \ln(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 + x^2 - \frac{1}{2}x^3 + \frac{1}{2}x^3 + \cdots = x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \cdots$.

77. $y = (\arctan x)^2 = \left(x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{7}x^7 + \cdots\right)\left(x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{7}x^7 + \cdots\right)$. Writing only the terms with degree ≤ 6, we get $(\arctan x)^2 = x^2 - \frac{1}{3}x^4 + \frac{1}{5}x^6 - \frac{1}{3}x^4 + \frac{1}{9}x^6 + \frac{1}{5}x^6 + \cdots = x^2 - \frac{2}{3}x^4 + \frac{23}{45}x^6 + \cdots$.

78. $y = e^x \sin^2 x = (e^x \sin x)\sin x = \left(x + x^2 + \frac{1}{3}x^3 + \cdots\right)\left(x - \frac{1}{6}x^3 + \cdots\right)$ [from Example 15]. Writing only the terms with degree ≤ 4, we get $e^x \sin^2 x = x^2 - \frac{1}{6}x^4 + x^3 + \frac{1}{3}x^4 + \cdots = x^2 + x^3 + \frac{1}{6}x^4 + \cdots$.

79. $\displaystyle\sum_{n=0}^{\infty} (-1)^n \frac{x^{4n}}{n!} = \sum_{n=0}^{\infty} \frac{\left(-x^4\right)^n}{n!} = e^{-x^4}$, by (11).

80. $\displaystyle\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{4n}}{n} = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{(x^4)^n}{n} = \ln(1+x^4)$ [from Table 1]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

81. $\sum_{n=0}^{\infty}(-1)^n\dfrac{x^{2n+1}}{2^{2n+1}(2n+1)} = \sum_{n=0}^{\infty}(-1)^n\dfrac{(x/2)^{2n+1}}{2n+1} = \tan^{-1}\left(\dfrac{x}{2}\right)$ [from Table 1]

82. $\sum_{n=0}^{\infty}(-1)^n\dfrac{x^{2n+1}}{2^{2n+1}(2n+1)!} = \sum_{n=0}^{\infty}(-1)^n\dfrac{(x/2)^{2n+1}}{(2n+1)!} = \sin\dfrac{x}{2}$ by (15).

83. $\sum_{n=0}^{\infty}\dfrac{(-1)^n}{n!}$ is the Maclaurin series for e^x evaluated at $x=-1$. Thus, $\sum_{n=0}^{\infty}\dfrac{(-1)^n}{n!} = e^{-1}$ by (11).

84. $\sum_{n=0}^{\infty}\dfrac{(-1)^n\pi^{2n}}{6^{2n}(2n)!} = \sum_{n=0}^{\infty}(-1)^n\dfrac{\left(\frac{\pi}{6}\right)^{2n}}{(2n)!} = \cos\frac{\pi}{6} = \frac{\sqrt{3}}{2}$, by (16).

85. $\sum_{n=1}^{\infty}(-1)^{n-1}\dfrac{3^n}{n5^n} = \sum_{n=1}^{\infty}(-1)^{n-1}\dfrac{(3/5)^n}{n} = \ln\left(1+\dfrac{3}{5}\right)$ [from Table 1] $= \ln\dfrac{8}{5}$

86. $\sum_{n=0}^{\infty}\dfrac{3^n}{5^n\,n!} = \sum_{n=0}^{\infty}\dfrac{(3/5)^n}{n!} = e^{3/5}$, by (11).

87. $\sum_{n=0}^{\infty}\dfrac{(-1)^n\pi^{2n+1}}{4^{2n+1}(2n+1)!} = \sum_{n=0}^{\infty}\dfrac{(-1)^n\left(\frac{\pi}{4}\right)^{2n+1}}{(2n+1)!} = \sin\frac{\pi}{4} = \frac{1}{\sqrt{2}}$, by (15).

88. $1-\ln 2+\dfrac{(\ln 2)^2}{2!}-\dfrac{(\ln 2)^3}{3!}+\cdots = \sum_{n=0}^{\infty}\dfrac{(-\ln 2)^n}{n!} = e^{-\ln 2} = \left(e^{\ln 2}\right)^{-1} = 2^{-1} = \frac{1}{2}$, by (11).

89. $3+\dfrac{9}{2!}+\dfrac{27}{3!}+\dfrac{81}{4!}+\cdots = \dfrac{3^1}{1!}+\dfrac{3^2}{2!}+\dfrac{3^3}{3!}+\dfrac{3^4}{4!}+\cdots = \sum_{n=1}^{\infty}\dfrac{3^n}{n!} = \sum_{n=0}^{\infty}\dfrac{3^n}{n!}-1 = e^3-1$, by (11).

90.
$$\frac{1}{1\cdot 2}-\frac{1}{3\cdot 2^3}+\frac{1}{5\cdot 2^5}-\frac{1}{7\cdot 2^7}+\cdots = \sum_{n=0}^{\infty}(-1)^n\frac{1}{(2n+1)2^{2n+1}}$$
$$= \sum_{n=0}^{\infty}(-1)^n\frac{(1/2)^{2n+1}}{2n+1} = \tan^{-1}\left(\frac{1}{2}\right) \quad \text{[from Table 1]}$$

91. If p is an nth-degree polynomial, then $p^{(i)}(x)=0$ for $i>n$, so its Taylor series at a is $p(x) = \sum_{i=0}^{n}\dfrac{p^{(i)}(a)}{i!}(x-a)^i$.

Put $x-a=1$, so that $x=a+1$. Then $p(a+1) = \sum_{i=0}^{n}\dfrac{p^{(i)}(a)}{i!}$.

This is true for any a, so replace a by x: $p(x+1) = \sum_{i=0}^{n}\dfrac{p^{(i)}(x)}{i!}$

92. Using the geometric series from Table 1, we have

$$\frac{x}{1+x^2} = x\cdot\frac{1}{1-(-x^2)} = x\cdot\sum_{n=0}^{\infty}(-x^2)^n = x\cdot\sum_{n=0}^{\infty}(-1)^n x^{2n} = \sum_{n=0}^{\infty}(-1)^n x^{2n+1}.$$

The x^{101} term is obtained when $n=50$ and is $(-1)^{50}x^{101} = x^{101}$. So the coefficient of x^{101} is 1 which, by Equation 7, must equal $\dfrac{f^{(101)}(0)}{101!}$. Thus, $\dfrac{f^{(101)}(0)}{101!} = 1 \;\Rightarrow\; f^{(101)}(0) = 101!$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

93. Using Equation 15, we have

$$x\sin(x^2) = x \cdot \sum_{n=0}^{\infty} (-1)^n \frac{(x^2)^{2n+1}}{(2n+1)!} = x \cdot \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{(2n+1)!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+3}}{(2n+1)!}$$

The x^{203} term is obtained when $n = 50$ and is $(-1)^{50} \dfrac{x^{203}}{101!} = \dfrac{1}{101!}\, x^{203}$. So the coefficient of x^{203} is $\dfrac{1}{101!}$, which, by Equation 7, must equal $\dfrac{f^{(203)}(0)}{203!}$. Thus, $\dfrac{f^{(203)}(0)}{203!} = \dfrac{1}{101!} \quad \Rightarrow \quad f^{(203)}(0) = \dfrac{203!}{101!}$.

94. The coefficient of x^{58} in the Maclaurin series of $f(x) = (1+x^3)^{30}$ is $\dfrac{f^{(58)}(0)}{58!}$. But the binomial series for $f(x)$ is $(1+x^3)^{30} = \sum_{n=0}^{\infty} \dbinom{30}{n} x^{3n}$, so it involves only powers of x that are multiples of 3 and therefore the coefficient of x^{58} is 0. So $f^{(58)}(0) = 0$.

95. Assume that $|f'''(x)| \le M$, so $f'''(x) \le M$ for $a \le x \le a+d$. Now $\int_a^x f'''(t)\,dt \le \int_a^x M\,dt \quad \Rightarrow$

$f''(x) - f''(a) \le M(x-a) \quad \Rightarrow \quad f''(x) \le f''(a) + M(x-a)$. Thus, $\int_a^x f''(t)\,dt \le \int_a^x [f''(a) + M(t-a)]\,dt \quad \Rightarrow$

$f'(x) - f'(a) \le f''(a)(x-a) + \frac{1}{2}M(x-a)^2 \quad \Rightarrow \quad f'(x) \le f'(a) + f''(a)(x-a) + \frac{1}{2}M(x-a)^2 \quad \Rightarrow$

$\int_a^x f'(t)\,dt \le \int_a^x \left[f'(a) + f''(a)(t-a) + \frac{1}{2}M(t-a)^2\right] dt \quad \Rightarrow$

$f(x) - f(a) \le f'(a)(x-a) + \frac{1}{2}f''(a)(x-a)^2 + \frac{1}{6}M(x-a)^3$. So

$f(x) - f(a) - f'(a)(x-a) - \frac{1}{2}f''(a)(x-a)^2 \le \frac{1}{6}M(x-a)^3$. But

$R_2(x) = f(x) - T_2(x) = f(x) - f(a) - f'(a)(x-a) - \frac{1}{2}f''(a)(x-a)^2$, so $R_2(x) \le \frac{1}{6}M(x-a)^3$.

A similar argument using $f'''(x) \ge -M$ shows that $R_2(x) \ge -\frac{1}{6}M(x-a)^3$. So $|R_2(x_2)| \le \frac{1}{6}M\,|x-a|^3$.

Although we have assumed that $x > a$, a similar calculation shows that this inequality is also true if $x < a$.

96. (a) $f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$ so $f'(0) = \lim\limits_{x\to 0} \dfrac{f(x)-f(0)}{x-0} = \lim\limits_{x\to 0} \dfrac{e^{-1/x^2}}{x} = \lim\limits_{x\to 0} \dfrac{1/x}{e^{1/x^2}} = \lim\limits_{x\to 0} \dfrac{x}{2e^{1/x^2}} = 0$

(using l'Hospital's Rule and simplifying in the penultimate step). Similarly, we can use the definition of the derivative and l'Hospital's Rule to show that $f''(0) = 0$, $f^{(3)}(0) = 0, \ldots, f^{(n)}(0) = 0$, so that the Maclaurin series for f consists entirely of zero terms. But since $f(x) \neq 0$ except for $x = 0$, we see that f cannot equal its Maclaurin series except at $x = 0$.

(b)

From the graph, it seems that the function is extremely flat at the origin. In fact, it could be said to be "infinitely flat" at $x = 0$, since all of its derivatives are 0 there.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

97. (a) $g(x) = \sum_{n=0}^{\infty} \binom{k}{n} x^n \quad \Rightarrow \quad g'(x) = \sum_{n=1}^{\infty} \binom{k}{n} n x^{n-1}$, so

$$\begin{aligned}
(1+x)g'(x) &= (1+x)\sum_{n=1}^{\infty}\binom{k}{n}nx^{n-1} = \sum_{n=1}^{\infty}\binom{k}{n}nx^{n-1} + \sum_{n=1}^{\infty}\binom{k}{n}nx^n \\
&= \sum_{n=0}^{\infty}\binom{k}{n+1}(n+1)x^n + \sum_{n=0}^{\infty}\binom{k}{n}nx^n \qquad \left[\begin{array}{c}\text{Replace } n \text{ with } n+1\\ \text{in the first series}\end{array}\right] \\
&= \sum_{n=0}^{\infty}(n+1)\frac{k(k-1)(k-2)\cdots(k-n+1)(k-n)}{(n+1)!}x^n + \sum_{n=0}^{\infty}\left[(n)\frac{k(k-1)(k-2)\cdots(k-n+1)}{n!}\right]x^n \\
&= \sum_{n=0}^{\infty}\frac{(n+1)k(k-1)(k-2)\cdots(k-n+1)}{(n+1)!}\left[(k-n)+n\right]x^n \\
&= k\sum_{n=0}^{\infty}\frac{k(k-1)(k-2)\cdots(k-n+1)}{n!}x^n = k\sum_{n=0}^{\infty}\binom{k}{n}x^n = kg(x)
\end{aligned}$$

Thus, $g'(x) = \dfrac{kg(x)}{1+x}$.

(b) $h(x) = (1+x)^{-k}\,g(x) \quad \Rightarrow$

$$\begin{aligned}
h'(x) &= -k(1+x)^{-k-1}g(x) + (1+x)^{-k}\,g'(x) && \text{[Product Rule]} \\
&= -k(1+x)^{-k-1}g(x) + (1+x)^{-k}\,\frac{kg(x)}{1+x} && \text{[from part (a)]} \\
&= -k(1+x)^{-k-1}g(x) + k(1+x)^{-k-1}g(x) = 0
\end{aligned}$$

(c) From part (b) we see that $h(x)$ must be constant for $x \in (-1,1)$, so $h(x) = h(0) = 1$ for $x \in (-1,1)$.

Thus, $h(x) = 1 = (1+x)^{-k}\,g(x) \quad \Leftrightarrow \quad g(x) = (1+x)^k$ for $x \in (-1,1)$.

98. Using the binomial series to expand $\sqrt{1+x}$ as a power series as in Example 9, we get

$\sqrt{1+x} = (1+x)^{1/2} = 1 + \dfrac{x}{2} + \sum_{n=2}^{\infty}\dfrac{(-1)^{n-1}1\cdot 3\cdot 5\cdot\cdots\cdot(2n-3)x^n}{2^n\cdot n!}$, so

$\left(1-x^2\right)^{1/2} = 1 - \dfrac{1}{2}x^2 - \sum_{n=2}^{\infty}\dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-3)}{2^n\cdot n!}x^{2n}$ and

$\sqrt{1-e^2\sin^2\theta} = 1 - \dfrac{1}{2}e^2\sin^2\theta - \sum_{n=2}^{\infty}\dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-3)}{2^n\cdot n!}e^{2n}\sin^{2n}\theta$. Thus,

$$\begin{aligned}
L &= 4a\int_0^{\pi/2}\sqrt{1-e^2\sin^2\theta}\,d\theta = 4a\int_0^{\pi/2}\left(1-\frac{1}{2}e^2\sin^2\theta - \sum_{n=2}^{\infty}\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-3)}{2^n\cdot n!}e^{2n}\sin^{2n}\theta\right)d\theta \\
&= 4a\left[\frac{\pi}{2} - \frac{e^2}{2}S_1 - \sum_{n=2}^{\infty}\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-3)}{n!}\left(\frac{e^2}{2}\right)^n S_n\right]
\end{aligned}$$

where $S_n = \displaystyle\int_0^{\pi/2}\sin^{2n}\theta\,d\theta = \frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{2\cdot 4\cdot 6\cdot\cdots\cdot 2n}\frac{\pi}{2}$ by Exercise 7.1.56.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$L = 4a\left(\frac{\pi}{2}\right)\left[1 - \frac{e^2}{2}\cdot\frac{1}{2} - \sum_{n=2}^{\infty}\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-3)}{n!}\left(\frac{e^2}{2}\right)^n\frac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{2\cdot 4\cdot 6\cdot\cdots\cdot 2n}\right]$$

$$= 2\pi a\left[1 - \frac{e^2}{4} - \sum_{n=2}^{\infty}\frac{e^{2n}}{2^n}\cdot\frac{1^2\cdot 3^2\cdot 5^2\cdot\cdots\cdot(2n-3)^2(2n-1)}{n!\cdot 2^n\cdot n!}\right]$$

$$= 2\pi a\left[1 - \frac{e^2}{4} - \sum_{n=2}^{\infty}\frac{e^{2n}}{4^n}\left(\frac{1\cdot 3\cdot\cdots\cdot(2n-3)}{n!}\right)^2(2n-1)\right]$$

$$= 2\pi a\left[1 - \frac{e^2}{4} - \frac{3e^4}{64} - \frac{5e^6}{256} - \cdots\right] = \frac{\pi a}{128}(256 - 64e^2 - 12e^4 - 5e^6 - \cdots)$$

DISCOVERY PROJECT An Elusive Limit

1. $f(x) = \dfrac{n(x)}{d(x)} = \dfrac{\sin(\tan x) - \tan(\sin x)}{\arcsin(\arctan x) - \arctan(\arcsin x)}$

The table of function values were obtained using Maple with 10 digits of precision. The results of this project will vary depending on the CAS and precision level. It appears that as $x \to 0^+$, $f(x) \to \frac{10}{3}$. Since f is an even function, we have $f(x) \to \frac{10}{3}$ as $x \to 0$.

x	$f(x)$
1	1.1838
0.1	0.9821
0.01	2.0000
0.001	3.3333
0.0001	3.3333

2. The graph is inconclusive about the limit of f as $x \to 0$.

3. The limit has the indeterminate form $\frac{0}{0}$. Applying l'Hospital's Rule, we obtain the form $\frac{0}{0}$ six times. Finally, on the seventh application we obtain $\lim\limits_{x\to 0}\dfrac{n^{(7)}(x)}{d^{(7)}(x)} = \dfrac{-168}{-168} = 1$.

4. $\lim\limits_{x\to 0} f(x) = \lim\limits_{x\to 0}\dfrac{n(x)}{d(x)} \overset{\text{CAS}}{=} \lim\limits_{x\to 0}\dfrac{-\frac{1}{30}x^7 - \frac{29}{756}x^9 + \cdots}{-\frac{1}{30}x^7 + \frac{13}{756}x^9 + \cdots}$

$$= \lim_{x\to 0}\frac{\left(-\frac{1}{30}x^7 - \frac{29}{756}x^9 + \cdots\right)/x^7}{\left(-\frac{1}{30}x^7 + \frac{13}{756}x^9 + \cdots\right)/x^7} = \lim_{x\to 0}\frac{-\frac{1}{30} - \frac{29}{756}x^2 + \cdots}{-\frac{1}{30} + \frac{13}{756}x^2 + \cdots} = \frac{-\frac{1}{30}}{-\frac{1}{30}} = 1$$

Note that $n^{(7)}(x) = d^{(7)}(x) = -\frac{7!}{30} = -\frac{5040}{30} = -168$, which agrees with the result in Problem 3.

5. The limit command gives the result that $\lim\limits_{x\to 0} f(x) = 1$.

6. The strange results (with only 10 digits of precision) must be due to the fact that the terms being subtracted in the numerator and denominator are very close in value when $|x|$ is small. Thus, the differences are imprecise (have few correct digits).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11.11 Applications of Taylor Polynomials

1. (a)

n	$f^{(n)}(x)$	$f^{(n)}(0)$	$T_n(x)$
0	$\sin x$	0	0
1	$\cos x$	1	x
2	$-\sin x$	0	x
3	$-\cos x$	-1	$x - \frac{1}{6}x^3$
4	$\sin x$	0	$x - \frac{1}{6}x^3$
5	$\cos x$	1	$x - \frac{1}{6}x^3 + \frac{1}{120}x^5$

Note: $T_n(x) = \sum_{k=0}^{n} \frac{f^{(k)}(0)}{k!} x^k$

(b)

x	f	$T_0(x)$	$T_1(x) = T_2(x)$	$T_3(x) = T_4(x)$	$T_5(x)$
$\frac{\pi}{4}$	0.7071	0	0.7854	0.7047	0.7071
$\frac{\pi}{2}$	1	0	1.5708	0.9248	1.0045
π	0	0	3.1416	-2.0261	0.5240

(c) As n increases, $T_n(x)$ is a good approximation to $f(x)$ on a larger and larger interval.

2. (a)

n	$f^{(n)}(x)$	$f^{(n)}(0)$	$T_n(x)$
0	$\tan x$	0	0
1	$\sec^2 x$	1	x
2	$2\sec^2 x \tan x$	0	x
3	$4\sec^2 x \tan^2 x + 2\sec^4 x$	2	$x + \frac{1}{3}x^3$

Note: $T_n(x) = \sum_{k=0}^{n} \frac{f^{(k)}(0)}{k!} x^k$

(b)

x	f	$T_0(x)$	$T_1(x) = T_2(x)$	$T_3(x)$
$\frac{\pi}{6}$	0.5774	0	0.5236	0.5714
$\frac{\pi}{4}$	1	0	0.7854	0.9469
$\frac{\pi}{3}$	1.7321	0	1.0472	1.4300

(c) As n increases, $T_n(x)$ is a good approximation to $f(x)$ on a larger and larger interval. Because the Taylor polynomials are continuous, they cannot approximate the infinite discontinuities at $x = \pm\pi/2$. They can only approximate $\tan x$ on $(-\pi/2, \pi/2)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	e^x	e
1	e^x	e
2	e^x	e
3	e^x	e

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(1)}{n!}(x-1)^n$$
$$= \frac{e}{0!}(x-1)^0 + \frac{e}{1!}(x-1)^1 + \frac{e}{2!}(x-1)^2 + \frac{e}{3!}(x-1)^3$$
$$= e + e(x-1) + \tfrac{1}{2}e(x-1)^2 + \tfrac{1}{6}e(x-1)^3$$

4.

n	$f^{(n)}(x)$	$f^{(n)}(\pi/6)$
0	$\sin x$	$1/2$
1	$\cos x$	$\sqrt{3}/2$
2	$-\sin x$	$-1/2$
3	$-\cos x$	$-\sqrt{3}/2$

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(\pi/6)}{n!}\left(x - \frac{\pi}{6}\right)^n$$
$$= \frac{1/2}{0!}\left(x-\frac{\pi}{6}\right)^0 + \frac{\sqrt{3}/2}{1!}\left(x-\frac{\pi}{6}\right)^1 - \frac{1/2}{2!}\left(x-\frac{\pi}{6}\right)^2 + \frac{\sqrt{3}/2}{3!}\left(x-\frac{\pi}{6}\right)^3$$
$$= \frac{1}{2} + \frac{\sqrt{3}}{2}\left(x-\frac{\pi}{6}\right) - \frac{1}{4}\left(x-\frac{\pi}{6}\right)^2 - \frac{\sqrt{3}}{12}\left(x-\frac{\pi}{6}\right)^3$$

5.

n	$f^{(n)}(x)$	$f^{(n)}(\pi/2)$
0	$\cos x$	0
1	$-\sin x$	-1
2	$-\cos x$	0
3	$\sin x$	1

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(\pi/2)}{n!}\left(x - \tfrac{\pi}{2}\right)^n$$
$$= -\left(x - \tfrac{\pi}{2}\right) + \tfrac{1}{6}\left(x - \tfrac{\pi}{2}\right)^3$$

6.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$e^{-x}\sin x$	0
1	$e^{-x}(\cos x - \sin x)$	1
2	$-2e^{-x}\cos x$	-2
3	$2e^{-x}(\cos x + \sin x)$	2

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(0)}{n!}x^n = x - x^2 + \tfrac{1}{3}x^3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$\ln x$	0
1	$1/x$	1
2	$-1/x^2$	-1
3	$2/x^3$	2

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(1)}{n!}(x-1)^n$$
$$= 0 + \frac{1}{1!}(x-1) + \frac{-1}{2!}(x-1)^2 + \frac{2}{3!}(x-1)^3$$
$$= (x-1) - \tfrac{1}{2}(x-1)^2 + \tfrac{1}{3}(x-1)^3$$

8.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$x\cos x$	0
1	$-x\sin x + \cos x$	1
2	$-x\cos x - 2\sin x$	0
3	$x\sin x - 3\cos x$	-3

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(0)}{n!}x^n$$
$$= 0 + \frac{1}{1!}x + 0 + \frac{-3}{3!}x^3 = x - \frac{1}{2}x^3$$

9.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	xe^{-2x}	0
1	$(1-2x)e^{-2x}$	1
2	$4(x-1)e^{-2x}$	-4
3	$4(3-2x)e^{-2x}$	12

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(0)}{n!}x^n = \tfrac{0}{1}\cdot 1 + \tfrac{1}{1}x^1 + \tfrac{-4}{2}x^2 + \tfrac{12}{6}x^3 = x - 2x^2 + 2x^3$$

10.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$\tan^{-1} x$	$\frac{\pi}{4}$
1	$\dfrac{1}{1+x^2}$	$\frac{1}{2}$
2	$\dfrac{-2x}{(1+x^2)^2}$	$-\frac{1}{2}$
3	$\dfrac{6x^2-2}{(1+x^2)^3}$	$\frac{1}{2}$

$$T_3(x) = \sum_{n=0}^{3} \frac{f^{(n)}(1)}{n!}(x-1)^n = \frac{\pi}{4} + \frac{1/2}{1}(x-1)^1 + \frac{-1/2}{2}(x-1)^2 + \frac{1/2}{6}(x-1)^3$$
$$= \tfrac{\pi}{4} + \tfrac{1}{2}(x-1) - \tfrac{1}{4}(x-1)^2 + \tfrac{1}{12}(x-1)^3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. You may be able to simply find the Taylor polynomials for $f(x) = \cot x$ using your CAS. We will list the values of $f^{(n)}(\pi/4)$ for $n = 0$ to $n = 5$.

n	0	1	2	3	4	5
$f^{(n)}(\pi/4)$	1	-2	4	-16	80	-512

$$T_5(x) = \sum_{n=0}^{5} \frac{f^{(n)}(\pi/4)}{n!}\left(x - \tfrac{\pi}{4}\right)^n$$
$$= 1 - 2\left(x - \tfrac{\pi}{4}\right) + 2\left(x - \tfrac{\pi}{4}\right)^2 - \tfrac{8}{3}\left(x - \tfrac{\pi}{4}\right)^3 + \tfrac{10}{3}\left(x - \tfrac{\pi}{4}\right)^4 - \tfrac{64}{15}\left(x - \tfrac{\pi}{4}\right)^5$$

For $n = 2$ to $n = 5$, $T_n(x)$ is the polynomial consisting of all the terms up to and including the $\left(x - \frac{\pi}{4}\right)^n$ term.

12. You may be able to simply find the Taylor polynomials for $f(x) = \sqrt[3]{1 + x^2}$ using your CAS. We will list the values of $f^{(n)}(0)$ for $n = 0$ to $n = 5$.

n	0	1	2	3	4	5
$f^{(n)}(0)$	1	0	$\frac{2}{3}$	0	$-\frac{8}{3}$	0

$$T_5(x) = \sum_{n=0}^{5} \frac{f^{(n)}(0)}{n!}\, x^n = 1 + \tfrac{1}{3}x^2 - \tfrac{1}{9}x^4$$

For $n = 2$ to $n = 5$, $T_n(x)$ is the polynomial consisting of all the terms up to and including the x^n term. Note that $T_2 = T_3$ and $T_4 = T_5$.

13. (a)

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$1/x$	1
1	$-1/x^2$	-1
2	$2/x^3$	2
3	$-6/x^4$	

$$f(x) = 1/x \approx T_2(x)$$
$$= \frac{1}{0!}(x-1)^0 - \frac{1}{1!}(x-1)^1 + \frac{2}{2!}(x-1)^2$$
$$= 1 - (x-1) + (x-1)^2$$

(b) $|R_2(x)| \le \dfrac{M}{3!}|x-1|^3$, where $|f'''(x)| \le M$. Now $0.7 \le x \le 1.3 \;\Rightarrow\; |x-1| \le 0.3 \;\Rightarrow\; |x-1|^3 \le 0.027$.

Since $|f'''(x)|$ is decreasing on $[0.7, 1.3]$, we can take $M = |f'''(0.7)| = 6/(0.7)^4$, so

$$|R_2(x)| \le \frac{6/(0.7)^4}{6}(0.027) = 0.112\,453\,1.$$

(c)

From the graph of $|R_2(x)| = \left|\dfrac{1}{x} - T_2(x)\right|$, it seems that the error is less than $0.038\,571$ on $[0.7, 1.3]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. (a)

n	$f^{(n)}(x)$	$f^{(n)}(4)$
0	$x^{-1/2}$	$\frac{1}{2}$
1	$-\frac{1}{2}x^{-3/2}$	$-\frac{1}{16}$
2	$\frac{3}{4}x^{-5/2}$	$\frac{3}{128}$
3	$-\frac{15}{8}x^{-7/2}$	

$$\begin{aligned} f(x) = x^{-1/2} &\approx T_2(x) \\ &= \frac{1/2}{0!}(x-4)^0 - \frac{1/16}{1!}(x-4)^1 + \frac{3/128}{2!}(x-4)^2 \\ &= \tfrac{1}{2} - \tfrac{1}{16}(x-4) + \tfrac{3}{256}(x-4)^2 \end{aligned}$$

(b) $|R_2(x)| \le \dfrac{M}{3!}|x-4|^3$, where $|f'''(x)| \le M$. Now $3.5 \le x \le 4.5 \;\Rightarrow\; |x-4| \le 0.5 \;\Rightarrow\; |x-4|^3 \le 0.125$.

Since $|f'''(x)|$ is decreasing on $[3.5, 4.5]$, we can take $M = |f'''(3.5)| = \dfrac{15}{8(3.5)^{7/2}}$, so

$|R_2(x)| \le \dfrac{15}{6 \cdot 8(3.5)^{7/2}}(0.125) \approx 0.000\,487$.

(c)

From the graph of $|R_2(x)| = \left|x^{-1/2} - T_2(x)\right|$, it seems that the error is less than $0.000\,343$ on $[3.5, 4.5]$.

15.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$x^{2/3}$	1
1	$\frac{2}{3}x^{-1/3}$	$\frac{2}{3}$
2	$-\frac{2}{9}x^{-4/3}$	$-\frac{2}{9}$
3	$\frac{8}{27}x^{-7/3}$	$\frac{8}{27}$
4	$-\frac{56}{81}x^{-10/3}$	

(a) $$\begin{aligned} f(x) = x^{2/3} \approx T_3(x) &= 1 + \tfrac{2}{3}(x-1) - \frac{2/9}{2!}(x-1)^2 + \frac{8/27}{3!}(x-1)^3 \\ &= 1 + \tfrac{2}{3}(x-1) - \tfrac{1}{9}(x-1)^2 + \tfrac{4}{81}(x-1)^3 \end{aligned}$$

(b) $|R_3(x)| \le \dfrac{M}{4!}|x-1|^4$, where $\left|f^{(4)}(x)\right| \le M$. Now $0.8 \le x \le 1.2 \;\Rightarrow$ $|x-1| \le 0.2 \;\Rightarrow\; |x-1|^4 \le 0.0016$. Since $\left|f^{(4)}(x)\right|$ is decreasing on $[0.8, 1.2]$, we can take $M = \left|f^{(4)}(0.8)\right| = \frac{56}{81}(0.8)^{-10/3}$, so

$|R_3(x)| \le \dfrac{\frac{56}{81}(0.8)^{-10/3}}{24}(0.0016) \approx 0.000\,096\,97$.

(c)

From the graph of $|R_3(x)| = \left|x^{2/3} - T_3(x)\right|$, it seems that the error is less than $0.000\,053\,3$ on $[0.8, 1.2]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16.

n	$f^{(n)}(x)$	$f^{(n)}(\pi/6)$
0	$\sin x$	$1/2$
1	$\cos x$	$\sqrt{3}/2$
2	$-\sin x$	$-1/2$
3	$-\cos x$	$-\sqrt{3}/2$
4	$\sin x$	$1/2$
5	$\cos x$	

(a) $f(x) = \sin x \approx T_4(x)$

$$= \tfrac{1}{2} + \tfrac{\sqrt{3}}{2}\left(x - \tfrac{\pi}{6}\right) - \tfrac{1}{4}\left(x - \tfrac{\pi}{6}\right)^2 - \tfrac{\sqrt{3}}{12}\left(x - \tfrac{\pi}{6}\right)^3 + \tfrac{1}{48}\left(x - \tfrac{\pi}{6}\right)^4$$

(b) $|R_4(x)| \le \dfrac{M}{5!}\left|x - \frac{\pi}{6}\right|^5$, where $\left|f^{(5)}(x)\right| \le M$. Now $0 \le x \le \frac{\pi}{3} \;\Rightarrow\; -\frac{\pi}{6} \le x - \frac{\pi}{6} \le \frac{\pi}{6} \;\Rightarrow\; \left|x - \frac{\pi}{6}\right| \le \frac{\pi}{6} \;\Rightarrow$ $\left|x - \frac{\pi}{6}\right|^5 \le \left(\frac{\pi}{6}\right)^5$. Since $\left|f^{(5)}(x)\right|$ is decreasing on $\left[0, \frac{\pi}{3}\right]$, we can take $M = \left|f^{(5)}(0)\right| = \cos 0 = 1$, so $|R_4(x)| \le \dfrac{1}{5!}\left(\dfrac{\pi}{6}\right)^5 \approx 0.000\,328$.

(c)

From the graph of $|R_4(x)| = |\sin x - T_4(x)|$, it seems that the error is less than $0.000\,297$ on $\left[0, \frac{\pi}{3}\right]$.

17.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sec x$	1
1	$\sec x \tan x$	0
2	$\sec x\,(2\sec^2 x - 1)$	1
3	$\sec x \tan x\,(6\sec^2 x - 1)$	

(a) $f(x) = \sec x \approx T_2(x) = 1 + \frac{1}{2}x^2$

(b) $|R_2(x)| \le \dfrac{M}{3!}|x|^3$, where $\left|f^{(3)}(x)\right| \le M$. Now $-0.2 \le x \le 0.2 \;\Rightarrow\; |x| \le 0.2 \;\Rightarrow\; |x|^3 \le (0.2)^3$.

$f^{(3)}(x)$ is an odd function and it is increasing on $[0, 0.2]$ since $\sec x$ and $\tan x$ are increasing on $[0, 0.2]$, so $\left|f^{(3)}(x)\right| \le f^{(3)}(0.2) \approx 1.085\,158\,892$. Thus, $|R_2(x)| \le \dfrac{f^{(3)}(0.2)}{3!}(0.2)^3 \approx 0.001\,447$.

(c)

From the graph of $|R_2(x)| = |\sec x - T_2(x)|$, it seems that the error is less than $0.000\,339$ on $[-0.2, 0.2]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$\ln(1+2x)$	$\ln 3$
1	$2/(1+2x)$	$\frac{2}{3}$
2	$-4/(1+2x)^2$	$-\frac{4}{9}$
3	$16/(1+2x)^3$	$\frac{16}{27}$
4	$-96/(1+2x)^4$	

(a) $f(x) = \ln(1+2x) \approx T_3(x)$

$$= \ln 3 + \tfrac{2}{3}(x-1) - \frac{4/9}{2!}(x-1)^2 + \frac{16/27}{3!}(x-1)^3$$

(b) $|R_3(x)| \le \dfrac{M}{4!}\,|x-1|^4$, where $\left|f^{(4)}(x)\right| \le M$. Now $0.5 \le x \le 1.5 \;\Rightarrow$ $-0.5 \le x-1 \le 0.5 \;\Rightarrow\; |x-1| \le 0.5 \;\Rightarrow\; |x-1|^4 \le \frac{1}{16}$, and letting $x = 0.5$ gives $M = 6$, so $|R_3(x)| \le \dfrac{6}{4!}\cdot\dfrac{1}{16} = \dfrac{1}{64} = 0.015\,625$.

(c)

From the graph of $|R_3(x)| = |\ln(1+2x) - T_3(x)|$, it seems that the error is less than 0.005 on $[0.5, 1.5]$.

19.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	e^{x^2}	1
1	$e^{x^2}(2x)$	0
2	$e^{x^2}(2+4x^2)$	2
3	$e^{x^2}(12x+8x^3)$	0
4	$e^{x^2}(12+48x^2+16x^4)$	

(a) $f(x) = e^{x^2} \approx T_3(x) = 1 + \dfrac{2}{2!}x^2 = 1 + x^2$

(b) $|R_3(x)| \le \dfrac{M}{4!}\,|x|^4$, where $\left|f^{(4)}(x)\right| \le M$. Now $0 \le x \le 0.1 \;\Rightarrow$ $x^4 \le (0.1)^4$, and letting $x = 0.1$ gives

$$|R_3(x)| \le \frac{e^{0.01}\,(12 + 0.48 + 0.0016)}{24}(0.1)^4 \approx 0.000\,053.$$

(c)

From the graph of $|R_3(x)| = \left|e^{x^2} - T_3(x)\right|$, it appears that the error is less than $0.000\,051$ on $[0, 0.1]$.

20.

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$x\ln x$	0
1	$\ln x + 1$	1
2	$1/x$	1
3	$-1/x^2$	-1
4	$2/x^3$	

(a) $f(x) = x\ln x \approx T_3(x) = (x-1) + \frac{1}{2}(x-1)^2 - \frac{1}{6}(x-1)^3$

(b) $|R_3(x)| \le \dfrac{M}{4!}\,|x-1|^4$, where $\left|f^{(4)}(x)\right| \le M$. Now $0.5 \le x \le 1.5 \;\Rightarrow$ $|x-1| \le \frac{1}{2} \;\Rightarrow\; |x-1|^4 \le \frac{1}{16}$. Since $\left|f^{(4)}(x)\right|$ is decreasing on $[0.5, 1.5]$, we can take $M = \left|f^{(4)}(0.5)\right| = 2/(0.5)^3 = 16$, so $|R_3(x)| \le \frac{16}{24}(1/16) = \frac{1}{24} = 0.041\overline{6}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c)

From the graph of $|R_3(x)| = |x \ln x - T_3(x)|$, it seems that the error is less than 0.0076 on $[0.5, 1.5]$.

21.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$x \sin x$	0
1	$\sin x + x \cos x$	0
2	$2 \cos x - x \sin x$	2
3	$-3 \sin x - x \cos x$	0
4	$-4 \cos x + x \sin x$	-4
5	$5 \sin x + x \cos x$	

(a) $f(x) = x \sin x \approx T_4(x) = \dfrac{2}{2!}(x-0)^2 + \dfrac{-4}{4!}(x-0)^4 = x^2 - \dfrac{1}{6}x^4$

(b) $|R_4(x)| \le \dfrac{M}{5!}|x|^5$, where $\left| f^{(5)}(x) \right| \le M$. Now $-1 \le x \le 1 \quad \Rightarrow$ $|x| \le 1$, and a graph of $f^{(5)}(x)$ shows that $\left| f^{(5)}(x) \right| \le 5$ for $-1 \le x \le 1$. Thus, we can take $M = 5$ and get $|R_4(x)| \le \dfrac{5}{5!} \cdot 1^5 = \dfrac{1}{24} = 0.041\overline{6}$.

(c)

From the graph of $|R_4(x)| = |x \sin x - T_4(x)|$, it seems that the error is less than 0.0082 on $[-1, 1]$.

22.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sinh 2x$	0
1	$2 \cosh 2x$	2
2	$4 \sinh 2x$	0
3	$8 \cosh 2x$	8
4	$16 \sinh 2x$	0
5	$32 \cosh 2x$	32
6	$64 \sinh 2x$	

(a) $f(x) = \sinh 2x \approx T_5(x) = 2x + \frac{8}{3!}x^3 + \frac{32}{5!}x^5 = 2x + \frac{4}{3}x^3 + \frac{4}{15}x^5$

(b) $|R_5(x)| \le \frac{M}{6!}|x|^6$, where $\left| f^{(6)}(x) \right| \le M$. For x in $[-1, 1]$, we have $|x| \le 1$. Since $f^{(6)}(x)$ is an increasing odd function on $[-1, 1]$, we see that $\left| f^{(6)}(x) \right| \le f^{(6)}(1) = 64 \sinh 2 = 32(e^2 - e^{-2}) \approx 232.119$, so we can take $M = 232.12$ and get $|R_5(x)| \le \frac{232.12}{720} \cdot 1^6 \approx 0.3224$.

(c)

From the graph of $|R_5(x)| = |\sinh 2x - T_5(x)|$, it seems that the error is less than 0.027 on $[-1, 1]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. From Exercise 5, $\cos x = -\left(x - \frac{\pi}{2}\right) + \frac{1}{6}\left(x - \frac{\pi}{2}\right)^3 + R_3(x)$, where $|R_3(x)| \le \dfrac{M}{4!}\left|x - \frac{\pi}{2}\right|^4$ with $\left|f^{(4)}(x)\right| = |\cos x| \le M = 1$. Now $x = 80^\circ = (90^\circ - 10^\circ) = \left(\frac{\pi}{2} - \frac{\pi}{18}\right) = \frac{4\pi}{9}$ radians, so the error is $\left|R_3\left(\frac{4\pi}{9}\right)\right| \le \frac{1}{24}\left(\frac{\pi}{18}\right)^4 \approx 0.000\,039$, which means our estimate would *not* be accurate to five decimal places. However, $T_3 = T_4$, so we can use $\left|R_4\left(\frac{4\pi}{9}\right)\right| \le \frac{1}{120}\left(\frac{\pi}{18}\right)^5 \approx 0.000\,001$. Therefore, to five decimal places, $\cos 80^\circ \approx -\left(-\frac{\pi}{18}\right) + \frac{1}{6}\left(-\frac{\pi}{18}\right)^3 \approx 0.17365$.

24. From Exercise 16, $\sin x = \frac{1}{2} + \frac{\sqrt{3}}{2}\left(x - \frac{\pi}{6}\right) - \frac{1}{4}\left(x - \frac{\pi}{6}\right)^2 - \frac{\sqrt{3}}{12}\left(x - \frac{\pi}{6}\right)^3 + \frac{1}{48}\left(x - \frac{\pi}{6}\right)^4 + R_4(x)$, where $|R_4(x)| \le \dfrac{M}{5!}\left|x - \frac{\pi}{6}\right|^5$ with $\left|f^{(5)}(x)\right| = |\cos x| \le M = 1$. Now $x = 38^\circ = (30^\circ + 8^\circ) = \left(\frac{\pi}{6} + \frac{2\pi}{45}\right)$ radians, so the error is $\left|R_4\left(\frac{38\pi}{180}\right)\right| \le \frac{1}{120}\left(\frac{2\pi}{45}\right)^5 \approx 0.000\,000\,44$, which means our estimate will be accurate to five decimal places. Therefore, to five decimal places, $\sin 38^\circ = \frac{1}{2} + \frac{\sqrt{3}}{2}\left(\frac{2\pi}{45}\right) - \frac{1}{4}\left(\frac{2\pi}{45}\right)^2 - \frac{\sqrt{3}}{12}\left(\frac{2\pi}{45}\right)^3 + \frac{1}{48}\left(\frac{2\pi}{45}\right)^4 \approx 0.61566$.

25. All derivatives of e^x are e^x, so $|R_n(x)| \le \dfrac{e^x}{(n+1)!}|x|^{n+1}$, where $0 < x < 0.1$. Letting $x = 0.1$, $R_n(0.1) \le \dfrac{e^{0.1}}{(n+1)!}(0.1)^{n+1} < 0.00001$, and by trial and error we find that $n = 3$ satisfies this inequality since $R_3(0.1) < 0.0000046$. Thus, by adding the four terms of the Maclaurin series for e^x corresponding to $n = 0$, 1, 2, and 3, we can estimate $e^{0.1}$ to within 0.00001. (In fact, this sum is $1.1051\overline{6}$ and $e^{0.1} \approx 1.10517$.)

26. From Table 1 in Section 11.10, $\ln(1+x) = \sum\limits_{n=1}^{\infty}(-1)^{n-1}\dfrac{x^n}{n}$ for $|x| < 1$. Thus, $\ln 1.4 = \ln(1+0.4) = \sum\limits_{n=1}^{\infty}(-1)^{n-1}\dfrac{(0.4)^n}{n}$. Since this is an alternating series, the error is less than the first neglected term by the Alternating Series Estimation Theorem, and we find that $|a_6| = (0.4)^6/6 \approx 0.0007 < 0.001$. So we need the first five (nonzero) terms of the Maclaurin series for the desired accuracy. (In fact, this sum is approximately 0.33698 and $\ln 1.4 \approx 0.33647$.)

27. $\sin x = x - \dfrac{1}{3!}x^3 + \dfrac{1}{5!}x^5 - \cdots$. By the Alternating Series Estimation Theorem, the error in the approximation $\sin x = x - \dfrac{1}{3!}x^3$ is less than $\left|\dfrac{1}{5!}x^5\right| < 0.01 \quad \Leftrightarrow$ $\left|x^5\right| < 120(0.01) \quad \Leftrightarrow \quad |x| < (1.2)^{1/5} \approx 1.037$. The curves $y = x - \frac{1}{6}x^3$ and $y = \sin x - 0.01$ intersect at $x \approx 1.043$, so the graph confirms our estimate. Since both the sine function and the given approximation are odd functions, we need to check the estimate only for $x > 0$. Thus, the desired range of values for x is $-1.037 < x < 1.037$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $\cos x = 1 - \dfrac{1}{2!}x^2 + \dfrac{1}{4!}x^4 - \dfrac{1}{6!}x^6 + \cdots$. By the Alternating Series Estimation Theorem, the error is less than $\left|-\dfrac{1}{6!}x^6\right| < 0.005 \quad \Leftrightarrow$ $x^6 < 720(0.005) \quad \Leftrightarrow \quad |x| < (3.6)^{1/6} \approx 1.238$. The curves $y = 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4$ and $y = \cos x + 0.005$ intersect at $x \approx 1.244$, so the graph confirms our estimate. Since both the cosine function and the given approximation are even functions, we need to check the estimate only for $x > 0$. Thus, the desired range of values for x is $-1.238 < x < 1.238$.

29. $\arctan x = x - \dfrac{x^3}{3} + \dfrac{x^5}{5} - \dfrac{x^7}{7} + \cdots$. By the Alternating Series Estimation Theorem, the error is less than $\left|-\frac{1}{7}x^7\right| < 0.05 \quad \Leftrightarrow$ $\left|x^7\right| < 0.35 \quad \Leftrightarrow \quad |x| < (0.35)^{1/7} \approx 0.8607$. The curves $y = x - \frac{1}{3}x^3 + \frac{1}{5}x^5$ and $y = \arctan x + 0.05$ intersect at $x \approx 0.9245$, so the graph confirms our estimate. Since both the arctangent function and the given approximation are odd functions, we need to check the estimate only for $x > 0$. Thus, the desired range of values for x is $-0.86 < x < 0.86$.

30. $f(x) = \sum\limits_{n=0}^{\infty} \dfrac{f^{(n)}(4)}{n!}(x-4)^n = \sum\limits_{n=0}^{\infty} \dfrac{(-1)^n\, n!}{3^n(n+1)\, n!}(x-4)^n = \sum\limits_{n=0}^{\infty} \dfrac{(-1)^n}{3^n(n+1)}(x-4)^n$. Now $f(5) = \sum\limits_{n=0}^{\infty} \dfrac{(-1)^n}{3^n(n+1)} = \sum\limits_{n=0}^{\infty} (-1)^n b_n$ is the sum of an alternating series that satisfies (i) $b_{n+1} \le b_n$ and (ii) $\lim\limits_{n\to\infty} b_n = 0$, so by the Alternating Series Estimation Theorem, $|R_5(5)| = |f(5) - T_5(5)| \le b_6$, and $b_6 = \dfrac{1}{3^6(7)} = \dfrac{1}{5103} \approx 0.000196 < 0.0002$; that is, the fifth-degree Taylor polynomial approximates $f(5)$ with error less than 0.0002.

31. Let $s(t)$ be the position function of the car, and for convenience set $s(0) = 0$. The velocity of the car is $v(t) = s'(t)$ and the acceleration is $a(t) = s''(t)$, so the second degree Taylor polynomial is $T_2(t) = s(0) + v(0)t + \dfrac{a(0)}{2}t^2 = 20t + t^2$. We estimate the distance traveled during the next second to be $s(1) \approx T_2(1) = 20 + 1 = 21$ m. The function $T_2(t)$ would not be accurate over a full minute, since the car could not possibly maintain an acceleration of $2\ \mathrm{m/s^2}$ for that long (if it did, its final speed would be $140\ \mathrm{m/s} \approx 313\ \mathrm{mi/h}$!).

32. (a)

n	$\rho^{(n)}(t)$	$\rho^{(n)}(20)$
0	$\rho_{20}e^{\alpha(t-20)}$	ρ_{20}
1	$\alpha\rho_{20}e^{\alpha(t-20)}$	$\alpha\rho_{20}$
2	$\alpha^2\rho_{20}e^{\alpha(t-20)}$	$\alpha^2\rho_{20}$

The linear approximation is

$$T_1(t) = \rho(20) + \rho'(20)(t-20) = \rho_{20}[1 + \alpha(t-20)]$$

The quadratic approximation is

$$T_2(t) = \rho(20) + \rho'(20)(t-20) + \frac{\rho''(20)}{2}(t-20)^2$$
$$= \rho_{20}\left[1 + \alpha(t-20) + \tfrac{1}{2}\alpha^2(t-20)^2\right]$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b)

(c)

From the graph, it seems that $T_1(t)$ is within 1% of $\rho(t)$, that is, $0.99\rho(t) \le T_1(t) \le 1.01\rho(t)$, for $-14°\text{C} \le t \le 58°\text{C}$.

33. $E = \dfrac{q}{D^2} - \dfrac{q}{(D+d)^2} = \dfrac{q}{D^2} - \dfrac{q}{D^2(1+d/D)^2} = \dfrac{q}{D^2}\left[1 - \left(1 + \dfrac{d}{D}\right)^{-2}\right]$.

We use the Binomial Series to expand $(1+d/D)^{-2}$:

$$E = \frac{q}{D^2}\left[1 - \left(1 - 2\left(\frac{d}{D}\right) + \frac{2\cdot 3}{2!}\left(\frac{d}{D}\right)^2 - \frac{2\cdot 3\cdot 4}{3!}\left(\frac{d}{D}\right)^3 + \cdots\right)\right] = \frac{q}{D^2}\left[2\left(\frac{d}{D}\right) - 3\left(\frac{d}{D}\right)^2 + 4\left(\frac{d}{D}\right)^3 - \cdots\right]$$
$$\approx \frac{q}{D^2}\cdot 2\left(\frac{d}{D}\right) = 2qd\cdot\frac{1}{D^3}$$

when D is much larger than d; that is, when P is far away from the dipole.

34. (a) $\dfrac{n_1}{\ell_o} + \dfrac{n_2}{\ell_i} = \dfrac{1}{R}\left(\dfrac{n_2 s_i}{\ell_i} - \dfrac{n_1 s_o}{\ell_o}\right)$ [Equation 1] where

$$\ell_o = \sqrt{R^2 + (s_o+R)^2 - 2R(s_o+R)\cos\phi} \quad \text{and} \quad \ell_i = \sqrt{R^2 + (s_i-R)^2 + 2R(s_i-R)\cos\phi} \quad \textbf{(2)}$$

Using $\cos\phi \approx 1$ gives

$$\ell_o = \sqrt{R^2 + (s_o+R)^2 - 2R(s_o+R)} = \sqrt{R^2 + s_o^2 + 2Rs_o + R^2 - 2Rs_o - 2R^2} = \sqrt{s_o^2} = s_o$$

and similarly, $\ell_i = s_i$. Thus, Equation 1 becomes $\dfrac{n_1}{s_o} + \dfrac{n_2}{s_i} = \dfrac{1}{R}\left(\dfrac{n_2 s_i}{s_i} - \dfrac{n_1 s_o}{s_o}\right) \quad \Rightarrow \quad \dfrac{n_1}{s_o} + \dfrac{n_2}{s_i} = \dfrac{n_2 - n_1}{R}$.

(b) Using $\cos\phi \approx 1 - \frac{1}{2}\phi^2$ in **(2)** gives us

$$\ell_o = \sqrt{R^2 + (s_o+R)^2 - 2R(s_o+R)\left(1 - \tfrac{1}{2}\phi^2\right)}$$
$$= \sqrt{R^2 + s_o^2 + 2Rs_o + R^2 - 2Rs_o + Rs_o\phi^2 - 2R^2 + R^2\phi^2} = \sqrt{s_o^2 + Rs_o\phi^2 + R^2\phi^2}$$

Anticipating that we will use the binomial series expansion $(1+x)^k \approx 1 + kx$, we can write the last expression for ℓ_o as $s_o\sqrt{1 + \phi^2\left(\dfrac{R}{s_o} + \dfrac{R^2}{s_o^2}\right)}$ and similarly, $\ell_i = s_i\sqrt{1 - \phi^2\left(\dfrac{R}{s_i} - \dfrac{R^2}{s_i^2}\right)}$. Thus, from Equation 1,

$$\frac{n_1}{\ell_o} + \frac{n_2}{\ell_i} = \frac{1}{R}\left(\frac{n_2 s_i}{\ell_i} - \frac{n_1 s_o}{\ell_o}\right) \quad \Leftrightarrow \quad n_1\ell_o^{-1} + n_2\ell_i^{-1} = \frac{n_2}{R}\cdot\frac{s_i}{\ell_i} - \frac{n_1}{R}\cdot\frac{s_o}{\ell_o} \quad \Leftrightarrow$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\frac{n_1}{s_o}\left[1+\phi^2\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\right]^{-1/2}+\frac{n_2}{s_i}\left[1-\phi^2\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)\right]^{-1/2}$$
$$=\frac{n_2}{R}\left[1-\phi^2\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)\right]^{-1/2}-\frac{n_1}{R}\left[1+\phi^2\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\right]^{-1/2}$$

Approximating the expressions for ℓ_o^{-1} and ℓ_i^{-1} by the first two terms in their binomial series, we get

$$\frac{n_1}{s_o}\left[1-\tfrac{1}{2}\phi^2\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\right]+\frac{n_2}{s_i}\left[1+\tfrac{1}{2}\phi^2\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)\right]$$
$$=\frac{n_2}{R}\left[1+\tfrac{1}{2}\phi^2\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)\right]-\frac{n_1}{R}\left[1-\tfrac{1}{2}\phi^2\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\right]\quad\Leftrightarrow$$

$$\frac{n_1}{s_o}-\frac{n_1\phi^2}{2s_o}\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)+\frac{n_2}{s_i}+\frac{n_2\phi^2}{2s_i}\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)=\frac{n_2}{R}+\frac{n_2\phi^2}{2R}\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)-\frac{n_1}{R}+\frac{n_1\phi^2}{2R}\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\quad\Leftrightarrow$$

$$\frac{n_1}{s_o}+\frac{n_2}{s_i}=\frac{n_2}{R}-\frac{n_1}{R}+\frac{n_1\phi^2}{2s_o}\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)+\frac{n_1\phi^2}{2R}\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)+\frac{n_2\phi^2}{2R}\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)-\frac{n_2\phi^2}{2s_i}\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)$$
$$=\frac{n_2-n_1}{R}+\frac{n_1\phi^2}{2}\left(\frac{R}{s_o}+\frac{R^2}{s_o^2}\right)\left(\frac{1}{s_o}+\frac{1}{R}\right)+\frac{n_2\phi^2}{2}\left(\frac{R}{s_i}-\frac{R^2}{s_i^2}\right)\left(\frac{1}{R}-\frac{1}{s_i}\right)$$
$$=\frac{n_2-n_1}{R}+\frac{n_1\phi^2R^2}{2s_o}\left(\frac{1}{R}+\frac{1}{s_o}\right)\left(\frac{1}{R}+\frac{1}{s_o}\right)+\frac{n_2\phi^2R^2}{2s_i}\left(\frac{1}{R}-\frac{1}{s_i}\right)\left(\frac{1}{R}-\frac{1}{s_i}\right)$$
$$=\frac{n_2-n_1}{R}+\phi^2R^2\left[\frac{n_1}{2s_o}\left(\frac{1}{R}+\frac{1}{s_o}\right)^2+\frac{n_2}{2s_i}\left(\frac{1}{R}-\frac{1}{s_i}\right)^2\right]$$

From Figure 8, we see that $\sin\phi=h/R$. So if we approximate $\sin\phi$ with ϕ, we get $h=R\phi$ and $h^2=\phi^2R^2$ and hence, Equation 4, as desired.

35. (a) If the water is deep, then $2\pi d/L$ is large, and we know that $\tanh x\to 1$ as $x\to\infty$. So we can approximate $\tanh(2\pi d/L)\approx 1$, and so $v^2\approx gL/(2\pi)\quad\Leftrightarrow\quad v\approx\sqrt{gL/(2\pi)}$.

(b) From the table, the first term in the Maclaurin series of $\tanh x$ is x, so if the water is shallow, we can approximate $\tanh\dfrac{2\pi d}{L}\approx\dfrac{2\pi d}{L}$, and so $v^2\approx\dfrac{gL}{2\pi}\cdot\dfrac{2\pi d}{L}\quad\Leftrightarrow\quad v\approx\sqrt{gd}$.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\tanh x$	0
1	$\operatorname{sech}^2 x$	1
2	$-2\operatorname{sech}^2 x\tanh x$	0
3	$2\operatorname{sech}^2 x\,(3\tanh^2 x-1)$	-2

(c) Since $\tanh x$ is an odd function, its Maclaurin series is alternating, so the error in the approximation $\tanh\dfrac{2\pi d}{L}\approx\dfrac{2\pi d}{L}$ is less than the first neglected term, which is $\dfrac{|f'''(0)|}{3!}\left(\dfrac{2\pi d}{L}\right)^3=\dfrac{1}{3}\left(\dfrac{2\pi d}{L}\right)^3$.

If $L>10d$, then $\dfrac{1}{3}\left(\dfrac{2\pi d}{L}\right)^3<\dfrac{1}{3}\left(2\pi\cdot\dfrac{1}{10}\right)^3=\dfrac{\pi^3}{375}$, so the error in the approximation $v^2=gd$ is less than $\dfrac{gL}{2\pi}\cdot\dfrac{\pi^3}{375}\approx 0.0132gL$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. First note that

$$\begin{aligned}
2\left(\sqrt{d^2+R^2}-d\right) &= 2\left[\sqrt{d^2}\sqrt{1+\frac{R^2}{d^2}}-d\right] \\
&\approx 2\left[d\left(1+\frac{R^2}{d^2}\cdot\frac{1}{2}+\cdots\right)-d\right] \quad \left[\text{use the binomial series } 1+\tfrac{1}{2}x+\cdots \text{ for } \sqrt{1+x}\,\right] \\
&= 2\left[\left(d+\frac{R^2}{2d}+\cdots\right)-d\right] \approx \frac{R^2}{d}
\end{aligned}$$

since for large d the other terms are comparatively small. Now $V = 2\pi k_e\sigma\left(\sqrt{d^2+R^2}-d\right) \approx \dfrac{\pi k_e R^2\sigma}{d}$ by the preceding approximation.

37. (a) L is the length of the arc subtended by the angle θ, so $L = R\theta \;\Rightarrow$ $\theta = L/R$. Now $\sec\theta = (R+C)/R \;\Rightarrow\; R\sec\theta = R+C \;\Rightarrow$ $C = R\sec\theta - R = R\sec(L/R) - R$.

(b) First we'll find a Taylor polynomial $T_4(x)$ for $f(x) = \sec x$ at $x = 0$.

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sec x$	1
1	$\sec x\tan x$	0
2	$\sec x(2\tan^2 x+1)$	1
3	$\sec x\tan x(6\tan^2 x+5)$	0
4	$\sec x(24\tan^4 x+28\tan^2 x+5)$	5

Thus, $f(x) = \sec x \approx T_4(x) = 1 + \frac{1}{2!}(x-0)^2 + \frac{5}{4!}(x-0)^4 = 1 + \frac{1}{2}x^2 + \frac{5}{24}x^4$. By part (a),

$$C \approx R\left[1+\frac{1}{2}\left(\frac{L}{R}\right)^2+\frac{5}{24}\left(\frac{L}{R}\right)^4\right]-R = R+\frac{1}{2}R\cdot\frac{L^2}{R^2}+\frac{5}{24}R\cdot\frac{L^4}{R^4}-R = \frac{L^2}{2R}+\frac{5L^4}{24R^3}.$$

(c) Taking $L = 100$ km and $R = 6370$ km, the formula in part (a) says that

$C = R\sec(L/R) - R = 6370\sec(100/6370) - 6370 \approx 0.785\,009\,965\,44$ km.

The formula in part (b) says that $C \approx \dfrac{L^2}{2R} + \dfrac{5L^4}{24R^3} = \dfrac{100^2}{2\cdot 6370} + \dfrac{5\cdot 100^4}{24\cdot 6370^3} \approx 0.785\,009\,957\,36$ km.

The difference between these two results is only $0.000\,000\,008\,08$ km, or $0.000\,008\,08$ m!

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. (a) $4\sqrt{\dfrac{L}{g}}\displaystyle\int_0^{\pi/2}\frac{dx}{\sqrt{1-k^2\sin^2 x}} = 4\sqrt{\frac{L}{g}}\int_0^{\pi/2}\left[1+\left(-k^2\sin^2 x\right)\right]^{-1/2}\,dx$

$$= 4\sqrt{\frac{L}{g}}\int_0^{\pi/2}\left[1-\frac{1}{2}\left(-k^2\sin^2 x\right)+\frac{\frac{1}{2}\cdot\frac{3}{2}}{2!}\left(-k^2\sin^2 x\right)^2-\frac{\frac{1}{2}\cdot\frac{3}{2}\cdot\frac{5}{2}}{3!}\left(-k^2\sin^2 x\right)^3+\cdots\right]dx$$

$$= 4\sqrt{\frac{L}{g}}\int_0^{\pi/2}\left[1+\left(\frac{1}{2}\right)k^2\sin^2 x+\left(\frac{1\cdot 3}{2\cdot 4}\right)k^4\sin^4 x+\left(\frac{1\cdot 3\cdot 5}{2\cdot 4\cdot 6}\right)k^6\sin^6 x+\cdots\right]dx$$

$$= 4\sqrt{\frac{L}{g}}\left[\frac{\pi}{2}+\left(\frac{1}{2}\right)\left(\frac{1}{2}\cdot\frac{\pi}{2}\right)k^2+\left(\frac{1\cdot 3}{2\cdot 4}\right)\left(\frac{1\cdot 3}{2\cdot 4}\cdot\frac{\pi}{2}\right)k^4+\left(\frac{1\cdot 3\cdot 5}{2\cdot 4\cdot 6}\right)\left(\frac{1\cdot 3\cdot 5}{2\cdot 4\cdot 6}\cdot\frac{\pi}{2}\right)k^6+\cdots\right]$$

[split up the integral and use the result from Exercise 7.1.56]

$$= 2\pi\sqrt{\frac{L}{g}}\left[1+\frac{1^2}{2^2}k^2+\frac{1^2\cdot 3^2}{2^2\cdot 4^2}k^4+\frac{1^2\cdot 3^2\cdot 5^2}{2^2\cdot 4^2\cdot 6^2}k^6+\cdots\right]$$

(b) The first of the two inequalities is true because all of the terms in the series are positive. For the second,

$$T = 2\pi\sqrt{\frac{L}{g}}\left[1+\frac{1^2}{2^2}k^2+\frac{1^2\cdot 3^2}{2^2\cdot 4^2}k^4+\frac{1^2\cdot 3^2\cdot 5^2}{2^2\cdot 4^2\cdot 6^2}k^6+\frac{1^2\cdot 3^2\cdot 5^2\cdot 7^2}{2^2\cdot 4^2\cdot 6^2\cdot 8^2}k^8+\cdots\right]$$

$$\le 2\pi\sqrt{\frac{L}{g}}\left[1+\tfrac{1}{4}k^2+\tfrac{1}{4}k^4+\tfrac{1}{4}k^6+\tfrac{1}{4}k^8+\cdots\right]$$

The terms in brackets (after the first) form a geometric series with $a=\frac{1}{4}k^2$ and $r=k^2=\sin^2\left(\frac{1}{2}\theta_0\right)<1$.

So $T \le 2\pi\sqrt{\dfrac{L}{g}}\left[1+\dfrac{k^2/4}{1-k^2}\right] = 2\pi\sqrt{\dfrac{L}{g}}\,\dfrac{4-3k^2}{4-4k^2}$.

(c) We substitute $L=1$, $g=9.8$, and $k=\sin(10^\circ/2)\approx 0.08716$, and the inequality from part (b) becomes $2.01090 \le T \le 2.01093$, so $T\approx 2.0109$. The estimate $T\approx 2\pi\sqrt{L/g}\approx 2.0071$ differs by about 0.2%.

If $\theta_0 = 42^\circ$, then $k\approx 0.35837$ and the inequality becomes $2.07153 \le T \le 2.08103$, so $T\approx 2.0763$.

The one-term estimate is the same, and the discrepancy between the two estimates increases to about 3.4%.

39. Using $f(x)=T_n(x)+R_n(x)$ with $n=1$ and $x=r$, we have $f(r)=T_1(r)+R_1(r)$, where T_1 is the first-degree Taylor polynomial of f at a. Because $a=x_n$, $f(r)=f(x_n)+f'(x_n)(r-x_n)+R_1(r)$. But r is a zero of f, so $f(r)=0$ and we have $0=f(x_n)+f'(x_n)(r-x_n)+R_1(r)$. Taking the first two terms to the left side gives us $f'(x_n)(x_n-r)-f(x_n)=R_1(r)$. Dividing by $f'(x_n)$, we get $x_n-r-\dfrac{f(x_n)}{f'(x_n)}=\dfrac{R_1(r)}{f'(x_n)}$. By the formula for Newton's method, the left side of the preceding equation is $x_{n+1}-r$, so $|x_{n+1}-r|=\left|\dfrac{R_1(r)}{f'(x_n)}\right|$. Taylor's Inequality gives us $|R_1(r)|\le\dfrac{|f''(r)|}{2!}\,|r-x_n|^2$. Combining this inequality with the facts $|f''(x)|\le M$ and $|f'(x)|\ge K$ gives us $|x_{n+1}-r|\le\dfrac{M}{2K}\,|x_n-r|^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

APPLIED PROJECT Radiation from the Stars

1. If we write $f(\lambda) = \dfrac{8\pi hc\lambda^{-5}}{e^{hc/(\lambda kT)} - 1} = \dfrac{a\lambda^{-5}}{e^{b/(\lambda T)} - 1}$, then as $\lambda \to 0^+$, it is of the form ∞/∞, and as $\lambda \to \infty$ it is of the form $0/0$, so in either case we can use l'Hospital's Rule. First of all,

$$\lim_{\lambda\to\infty} f(\lambda) \overset{\text{H}}{=} \lim_{\lambda\to\infty} \frac{a\left(-5\lambda^{-6}\right)}{-\dfrac{bT}{(\lambda T)^2}e^{b/(\lambda T)}} = 5\,\frac{aT}{b} \lim_{\lambda\to\infty} \frac{\lambda^2\lambda^{-6}}{e^{b/(\lambda T)}} = 5\,\frac{aT}{b} \lim_{\lambda\to\infty} \frac{\lambda^{-4}}{e^{b/(\lambda T)}} = 0$$

Also,
$$\lim_{\lambda\to 0^+} f(\lambda) \overset{\text{H}}{=} 5\,\frac{aT}{b} \lim_{\lambda\to 0^+} \frac{\lambda^{-4}}{e^{b/(\lambda T)}} \overset{\text{H}}{=} 5\,\frac{aT}{b} \lim_{\lambda\to 0^+} \frac{-4\lambda^{-5}}{-\dfrac{bT}{(\lambda T)^2}e^{b/(\lambda T)}} = 20\,\frac{aT}{b^2}^{\,2} \lim_{\lambda\to 0^+} \frac{\lambda^{-3}}{e^{b/(\lambda T)}}$$

This is still indeterminate, but note that each time we use l'Hospital's Rule, we gain a factor of λ in the numerator, as well as a constant factor, and the denominator is unchanged. So if we use l'Hospital's Rule three more times, the exponent of λ in the numerator will become 0. That is, for some $\{k_i\}$, all constant,

$$\lim_{\lambda\to 0^+} f(\lambda) \overset{\text{H}}{=} k_1 \lim_{\lambda\to 0^+} \frac{\lambda^{-3}}{e^{b/(\lambda T)}} \overset{\text{H}}{=} k_2 \lim_{\lambda\to 0^+} \frac{\lambda^{-2}}{e^{b/(\lambda T)}} \overset{\text{H}}{=} k_3 \lim_{\lambda\to 0^+} \frac{\lambda^{-1}}{e^{b/(\lambda T)}} \overset{\text{H}}{=} k_4 \lim_{\lambda\to 0^+} \frac{1}{e^{b/(\lambda T)}} = 0$$

2. We expand the denominator of Planck's Law using the Taylor series $e^x = 1 + x + \dfrac{x^2}{2!} + \dfrac{x^3}{3!} + \cdots$ with $x = \dfrac{hc}{\lambda kT}$, and use the fact that if λ is large, then all subsequent terms in the Taylor expansion are very small compared to the first one, so we can approximate using the Taylor polynomial T_1:

$$f(\lambda) = \frac{8\pi hc\lambda^{-5}}{e^{hc/(\lambda kT)} - 1} = \frac{8\pi hc\lambda^{-5}}{\left[1 + \dfrac{hc}{\lambda kT} + \dfrac{1}{2!}\left(\dfrac{hc}{\lambda kT}\right)^2 + \dfrac{1}{3!}\left(\dfrac{hc}{\lambda kT}\right)^3 + \cdots\right] - 1} \approx \frac{8\pi hc\lambda^{-5}}{\left(1 + \dfrac{hc}{\lambda kT}\right) - 1} = \frac{8\pi kT}{\lambda^4}$$

which is the Rayleigh-Jeans Law.

3. To convert to μm, we substitute $\lambda/10^6$ for λ in both laws. The first figure shows that the two laws are similar for large λ. The second figure shows that the two laws are very different for short wavelengths (Planck's Law gives a maximum at $\lambda \approx 0.51\ \mu$m; the Rayleigh-Jeans Law gives no minimum or maximum.).

4. From the graph in Problem 3, $f(\lambda)$ has a maximum under Planck's Law at $\lambda \approx 0.51\ \mu$m.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5.

As T gets larger, the total area under the curve increases, as we would expect: the hotter the star, the more energy it emits. Also, as T increases, the λ-value of the maximum decreases, so the higher the temperature, the shorter the peak wavelength (and consequently the average wavelength) of light emitted. This is why Sirius is a blue star and Betelgeuse is a red star: most of Sirius's light is of a fairly short wavelength; that is, a higher frequency, toward the blue end of the spectrum, whereas most of Betelgeuse's light is of a lower frequency, toward the red end of the spectrum.

11 Review

TRUE-FALSE QUIZ

1. False. See the WARNING after Theorem 11.2.6.

2. False. The series $\sum\limits_{n=1}^{\infty} n^{-\sin 1} = \sum\limits_{n=1}^{\infty} \dfrac{1}{n^{\sin 1}}$ is a p-series with $p = \sin 1 \approx 0.84 \leq 1$, so the series diverges.

3. True. If $\lim\limits_{n\to\infty} a_n = L$, then as $n \to \infty$, $2n+1 \to \infty$, so $a_{2n+1} \to L$.

4. True by Theorem 11.8.4.

Or: Use the Direct Comparison Test to show that $\sum c_n(-2)^n$ converges absolutely.

5. False. For example, take $c_n = (-1)^n/(n6^n)$.

6. True by Theorem 11.8.4.

7. False, since $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{1}{(n+1)^3}\cdot\dfrac{n^3}{1}\right| = \lim\limits_{n\to\infty}\left|\dfrac{n^3}{(n+1)^3}\cdot\dfrac{1/n^3}{1/n^3}\right| = \lim\limits_{n\to\infty}\dfrac{1}{(1+1/n)^3} = 1.$

8. True, since $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{1}{(n+1)!}\cdot\dfrac{n!}{1}\right| = \lim\limits_{n\to\infty}\dfrac{1}{n+1} = 0 < 1.$

9. False. See the paragraph preceding "The Limit Comparison Test" in Section 11.4.

10. True, since $\dfrac{1}{e} = e^{-1}$ and $e^x = \sum\limits_{n=0}^{\infty}\dfrac{x^n}{n!}$, so $e^{-1} = \sum\limits_{n=0}^{\infty}\dfrac{(-1)^n}{n!}$.

11. True. See (9) in Section 11.1.

12. True, because if $\sum |a_n|$ is convergent, then so is $\sum a_n$ by the contrapositive of Theorem 11.5.3.

13. True. By Theorem 11.10.5 the coefficient of x^3 is $\dfrac{f'''(0)}{3!} = \dfrac{1}{3} \quad\Rightarrow\quad f'''(0) = 2.$

Or: Use Theorem 11.9.2 to differentiate f three times.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. False. Let $a_n = n$ and $b_n = -n$. Then $\{a_n\}$ and $\{b_n\}$ are divergent, but $a_n + b_n = 0$, so $\{a_n + b_n\}$ is convergent.

15. False. For example, let $a_n = b_n = (-1)^n$. Then $\{a_n\}$ and $\{b_n\}$ are divergent, but $a_n b_n = 1$, so $\{a_n b_n\}$ is convergent.

16. True by the Monotonic Sequence Theorem, since $\{a_n\}$ is decreasing and $0 < a_n \le a_1$ for all $n \quad \Rightarrow \quad \{a_n\}$ is bounded.

17. True by Theorem 11.5.3. $\left[\sum (-1)^n a_n \text{ is absolutely convergent and hence convergent.}\right]$

18. True. $\lim\limits_{n\to\infty} \dfrac{a_{n+1}}{a_n} < 1 \quad \Rightarrow \quad \sum a_n$ converges (Ratio Test) $\quad \Rightarrow \quad \lim\limits_{n\to\infty} a_n = 0$ [Theorem 11.2.6].

19. True. $0.99999\ldots = 0.9 + 0.9(0.1)^1 + 0.9(0.1)^2 + 0.9(0.1)^3 + \cdots = \sum\limits_{n=1}^{\infty} (0.9)(0.1)^{n-1} = \dfrac{0.9}{1-0.1} = 1$ by the formula for the sum of a geometric series $[S = a_1/(1-r)]$ with ratio r satisfying $|r| < 1$.

20. True. Since $\lim\limits_{n\to\infty} a_n = 2$, we know that $\lim\limits_{n\to\infty} a_{n+3} = 2$. Thus, $\lim\limits_{n\to\infty} (a_{n+3} - a_n) = \lim\limits_{n\to\infty} a_{n+3} - \lim\limits_{n\to\infty} a_n = 2 - 2 = 0$.

21. True. A finite number of terms doesn't affect convergence or divergence of a series.

22. False. Let $a_n = (0.1)^n$ and $b_n = (0.2)^n$. Then $\sum\limits_{n=1}^{\infty} a_n = \sum\limits_{n=1}^{\infty} (0.1)^n = \dfrac{0.1}{1-0.1} = \dfrac{1}{9} = A$, $\sum\limits_{n=1}^{\infty} b_n = \sum\limits_{n=1}^{\infty} (0.2)^n = \dfrac{0.2}{1-0.2} = \dfrac{1}{4} = B$, and $\sum\limits_{n=1}^{\infty} a_n b_n = \sum\limits_{n=1}^{\infty} (0.02)^n = \dfrac{0.02}{1-0.02} = \dfrac{1}{49}$, but $AB = \frac{1}{9} \cdot \frac{1}{4} = \frac{1}{36}$.

EXERCISES

1. $\left\{\dfrac{2+n^3}{1+2n^3}\right\}$ converges since $\lim\limits_{n\to\infty} \dfrac{2+n^3}{1+2n^3} = \lim\limits_{n\to\infty} \dfrac{2/n^3+1}{1/n^3+2} = \dfrac{1}{2}$.

2. $a_n = \dfrac{9^{n+1}}{10^n} = 9 \cdot \left(\frac{9}{10}\right)^n$, so $\lim\limits_{n\to\infty} a_n = 9 \lim\limits_{n\to\infty} \left(\frac{9}{10}\right)^n = 9 \cdot 0 = 0$ by (11.1.9).

3. $\lim\limits_{n\to\infty} a_n = \lim\limits_{n\to\infty} \dfrac{n^3}{1+n^2} = \lim\limits_{n\to\infty} \dfrac{n}{1/n^2+1} = \infty$, so the sequence diverges.

4. $a_n = \cos(n\pi/2)$, so $a_n = 0$ if n is odd and $a_n = \pm 1$ if n is even. As n increases, a_n keeps cycling through the values $0, 1, 0, -1$, so the sequence $\{a_n\}$ is divergent.

5. $|a_n| = \left|\dfrac{n \sin n}{n^2+1}\right| \le \dfrac{n}{n^2+1} < \dfrac{1}{n}$, so $|a_n| \to 0$ as $n \to \infty$. Thus, $\lim\limits_{n\to\infty} a_n = 0$. The sequence $\{a_n\}$ is convergent.

6. $a_n = \dfrac{\ln n}{\sqrt{n}}$. Let $f(x) = \dfrac{\ln x}{\sqrt{x}}$ for $x > 0$. Then $\lim\limits_{x\to\infty} f(x) = \lim\limits_{x\to\infty} \dfrac{\ln x}{\sqrt{x}} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1/x}{1/(2\sqrt{x}\,)} = \lim\limits_{x\to\infty} \dfrac{2}{\sqrt{x}} = 0$.

Thus, by Theorem 11.1.4, $\{a_n\}$ converges and $\lim\limits_{n\to\infty} a_n = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. $\left\{\left(1+\dfrac{3}{n}\right)^{4n}\right\}$ is convergent. Let $y=\left(1+\dfrac{3}{x}\right)^{4x}$. Then

$$\lim_{x\to\infty}\ln y=\lim_{x\to\infty}4x\ln(1+3/x)=\lim_{x\to\infty}\frac{\ln(1+3/x)}{1/(4x)}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{\dfrac{1}{1+3/x}\left(-\dfrac{3}{x^2}\right)}{-1/(4x^2)}=\lim_{x\to\infty}\frac{12}{1+3/x}=12,\text{ so}$$

$$\lim_{x\to\infty}y=\lim_{n\to\infty}\left(1+\frac{3}{n}\right)^{4n}=e^{12}.$$

8. $\left\{\dfrac{(-10)^n}{n!}\right\}$ converges, since $\dfrac{10^n}{n!}=\dfrac{10\cdot10\cdot10\cdot\cdots\cdot10}{1\cdot2\cdot3\cdot\cdots\cdot10}\cdot\dfrac{10\cdot10\cdot\cdots\cdot10}{11\cdot12\cdot\cdots\cdot n}\le10^{10}\left(\dfrac{10}{11}\right)^{n-10}\to0$ as $n\to\infty$, so

$\displaystyle\lim_{n\to\infty}\frac{(-10)^n}{n!}=0$ [Squeeze Theorem]. *Or:* Use (11.10.10).

9. We use induction, hypothesizing that $a_{n-1}<a_n<2$. Note first that $1<a_2=\frac{1}{3}(1+4)=\frac{5}{3}<2$, so the hypothesis holds for $n=2$. Now assume that $a_{k-1}<a_k<2$. Then $a_k=\frac{1}{3}(a_{k-1}+4)<\frac{1}{3}(a_k+4)<\frac{1}{3}(2+4)=2$. So $a_k<a_{k+1}<2$, and the induction is complete. To find the limit of the sequence, we note that $L=\displaystyle\lim_{n\to\infty}a_n=\lim_{n\to\infty}a_{n+1}\quad\Rightarrow$ $L=\frac{1}{3}(L+4)\quad\Rightarrow\quad L=2$.

10. $\displaystyle\lim_{x\to\infty}\frac{x^4}{e^x}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{4x^3}{e^x}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{12x^2}{e^x}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{24x}{e^x}\overset{\text{H}}{=}\lim_{x\to\infty}\frac{24}{e^x}=0$

Then we conclude from Theorem 11.1.4 that $\displaystyle\lim_{n\to\infty}n^4e^{-n}=0$.

From the graph, it seems that $12^4e^{-12}>0.1$, but $n^4e^{-n}<0.1$ whenever $n>12$. So the smallest value of N corresponding to $\varepsilon=0.1$ in the definition of the limit is $N=12$.

11. $\dfrac{n}{n^3+1}<\dfrac{n}{n^3}=\dfrac{1}{n^2}$, so $\displaystyle\sum_{n=1}^{\infty}\frac{n}{n^3+1}$ converges by the Direct Comparison Test with the convergent p-series $\displaystyle\sum_{n=1}^{\infty}\frac{1}{n^2}$ $[p=2>1]$.

12. Let $a_n=\dfrac{n^2+1}{n^3+1}$ and $b_n=\dfrac{1}{n}$, so $\displaystyle\lim_{n\to\infty}\frac{a_n}{b_n}=\lim_{n\to\infty}\frac{n^3+n}{n^3+1}=\lim_{n\to\infty}\frac{1+1/n^2}{1+1/n^3}=1>0$.

Since $\displaystyle\sum_{n=1}^{\infty}b_n$ is the divergent harmonic series, $\displaystyle\sum_{n=1}^{\infty}a_n$ also diverges by the Limit Comparison Test.

13. $\displaystyle\lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right|=\lim_{n\to\infty}\left[\frac{(n+1)^3}{5^{n+1}}\cdot\frac{5^n}{n^3}\right]=\lim_{n\to\infty}\left(1+\frac{1}{n}\right)^3\cdot\frac{1}{5}=\frac{1}{5}<1$, so $\displaystyle\sum_{n=1}^{\infty}\frac{n^3}{5^n}$ converges by the Ratio Test.

14. Let $b_n=\dfrac{1}{\sqrt{n+1}}$. Then b_n is positive for $n\ge1$, the sequence $\{b_n\}$ is decreasing, and $\displaystyle\lim_{n\to\infty}b_n=0$, so the series $\displaystyle\sum_{n=1}^{\infty}\frac{(-1)^n}{\sqrt{n+1}}$ converges by the Alternating Series Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. Let $f(x) = \dfrac{1}{x\sqrt{\ln x}}$. Then f is continuous, positive, and decreasing on $[2, \infty)$, so the Integral Test applies.

$$\int_2^\infty f(x)\,dx = \lim_{t\to\infty}\int_2^t \frac{1}{x\sqrt{\ln x}}\,dx \quad \left[u = \ln x,\, du = \tfrac{1}{x}\,dx\right] \quad = \lim_{t\to\infty}\int_{\ln 2}^{\ln t} u^{-1/2}\,du = \lim_{t\to\infty}\left[2\sqrt{u}\right]_{\ln 2}^{\ln t}$$
$$= \lim_{t\to\infty}\left(2\sqrt{\ln t} - 2\sqrt{\ln 2}\right) = \infty,$$

so the series $\sum\limits_{n=2}^{\infty} \dfrac{1}{n\sqrt{\ln n}}$ diverges.

16. $\lim\limits_{n\to\infty} \dfrac{n}{3n+1} = \dfrac{1}{3}$, so $\lim\limits_{n\to\infty} \ln\left(\dfrac{n}{3n+1}\right) = \ln\frac{1}{3} \neq 0$. Thus, the series $\sum\limits_{n=1}^{\infty} \ln\left(\dfrac{n}{3n+1}\right)$ diverges by the Test for Divergence.

17. $|a_n| = \left|\dfrac{\cos 3n}{1+(1.2)^n}\right| \leq \dfrac{1}{1+(1.2)^n} < \dfrac{1}{(1.2)^n} = \left(\dfrac{5}{6}\right)^n$, so $\sum\limits_{n=1}^{\infty} |a_n|$ converges by direct comparison with the convergent geometric series $\sum\limits_{n=1}^{\infty} \left(\frac{5}{6}\right)^n$ $\left[r = \frac{5}{6} < 1\right]$. It follows that $\sum\limits_{n=1}^{\infty} a_n$ converges (by Theorem 11.5.3).

18. $\lim\limits_{n\to\infty} \sqrt[n]{|a_n|} = \lim\limits_{n\to\infty} \sqrt[n]{\left|\dfrac{n^{2n}}{(1+2n^2)^n}\right|} = \lim\limits_{n\to\infty} \dfrac{n^2}{1+2n^2} = \lim\limits_{n\to\infty} \dfrac{1}{1/n^2+2} = \dfrac{1}{2} < 1$, so $\sum\limits_{n=1}^{\infty} \dfrac{n^{2n}}{(1+2n^2)^n}$ converges by the Root Test.

19. $\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)(2n+1)}{5^{n+1}\,(n+1)!} \cdot \dfrac{5^n\,n!}{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)} = \lim\limits_{n\to\infty} \dfrac{2n+1}{5(n+1)} = \dfrac{2}{5} < 1$, so the series $\sum\limits_{n=1}^{\infty} \dfrac{1\cdot 3\cdot 5\cdot\cdots\cdot(2n-1)}{5^n\,n!}$ converges by the Ratio Test.

20. $\sum\limits_{n=1}^{\infty} \dfrac{(-5)^{2n}}{n^2\,9^n} = \sum\limits_{n=1}^{\infty} \dfrac{1}{n^2}\left(\dfrac{25}{9}\right)^n$. Now $\lim\limits_{n\to\infty} \left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty} \dfrac{25^{n+1}}{(n+1)^2\cdot 9^{n+1}} \cdot \dfrac{n^2\cdot 9^n}{25^n} = \lim\limits_{n\to\infty} \dfrac{25n^2}{9(n+1)^2} = \dfrac{25}{9} > 1$, so the series diverges by the Ratio Test.

21. $b_n = \dfrac{\sqrt{n}}{n+1} > 0$, $\{b_n\}$ is decreasing, and $\lim\limits_{n\to\infty} b_n = 0$, so the series $\sum\limits_{n=1}^{\infty} (-1)^{n-1}\dfrac{\sqrt{n}}{n+1}$ converges by the Alternating Series Test.

22. Use the Limit Comparison Test with $a_n = \dfrac{\sqrt{n+1}-\sqrt{n-1}}{n} = \dfrac{2}{n\left(\sqrt{n+1}+\sqrt{n-1}\right)}$ (rationalizing the numerator) and $b_n = \dfrac{1}{n^{3/2}}$. $\lim\limits_{n\to\infty} \dfrac{a_n}{b_n} = \lim\limits_{n\to\infty} \dfrac{2\sqrt{n}}{\sqrt{n+1}+\sqrt{n-1}} = 1$, so since $\sum\limits_{n=1}^{\infty} b_n$ converges $\left[p = \frac{3}{2} > 1\right]$, $\sum\limits_{n=1}^{\infty} a_n$ converges also.

23. Consider the series of absolute values: $\sum\limits_{n=1}^{\infty} n^{-1/3}$ is a p-series with $p = \frac{1}{3} \leq 1$ and is therefore divergent. But if we apply the Alternating Series Test, we see that $b_n = \dfrac{1}{\sqrt[3]{n}} > 0$, $\{b_n\}$ is decreasing, and $\lim\limits_{n\to\infty} b_n = 0$, so the series $\sum\limits_{n=1}^{\infty} (-1)^{n-1}\,n^{-1/3}$ converges. Thus, $\sum\limits_{n=1}^{\infty} (-1)^{n-1}\,n^{-1/3}$ is conditionally convergent.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $\sum\limits_{n=1}^{\infty} \left|(-1)^{n-1} n^{-3}\right| = \sum\limits_{n=1}^{\infty} n^{-3}$ is a convergent p-series $[p = 3 > 1]$. Therefore, $\sum\limits_{n=1}^{\infty} (-1)^{n-1} n^{-3}$ is absolutely convergent.

25. $\left|\dfrac{a_{n+1}}{a_n}\right| = \left|\dfrac{(-1)^{n+1}(n+2)3^{n+1}}{2^{2n+3}} \cdot \dfrac{2^{2n+1}}{(-1)^n(n+1)3^n}\right| = \dfrac{n+2}{n+1} \cdot \dfrac{3}{4} = \dfrac{1+(2/n)}{1+(1/n)} \cdot \dfrac{3}{4} \to \dfrac{3}{4} < 1$ as $n \to \infty$, so by the Ratio Test, $\sum\limits_{n=1}^{\infty} \dfrac{(-1)^n(n+1)3^n}{2^{2n+1}}$ is absolutely convergent.

26. $\lim\limits_{x\to\infty} \dfrac{\sqrt{x}}{\ln x} \overset{\text{H}}{=} \lim\limits_{x\to\infty} \dfrac{1/(2\sqrt{x}\,)}{1/x} = \lim\limits_{x\to\infty} \dfrac{\sqrt{x}}{2} = \infty$. Therefore, $\lim\limits_{n\to\infty} \dfrac{(-1)^n\sqrt{n}}{\ln n} \neq 0$, so the given series $\sum\limits_{n=2}^{\infty} \dfrac{(-1)^n\sqrt{n}}{\ln n}$ is divergent by the Test for Divergence.

27. $\sum\limits_{n=1}^{\infty} \dfrac{(-3)^{n-1}}{2^{3n}} = \sum\limits_{n=1}^{\infty} \dfrac{(-3)^{n-1}}{(2^3)^n} = \sum\limits_{n=1}^{\infty} \dfrac{(-3)^{n-1}}{8^n} = \dfrac{1}{8}\sum\limits_{n=1}^{\infty} \dfrac{(-3)^{n-1}}{8^{n-1}} = \dfrac{1}{8}\sum\limits_{n=1}^{\infty} \left(-\dfrac{3}{8}\right)^{n-1} = \dfrac{1}{8}\left(\dfrac{1}{1-(-3/8)}\right)$

$$= \frac{1}{8} \cdot \frac{8}{11} = \frac{1}{11}$$

28. $\sum\limits_{n=1}^{\infty} \dfrac{1}{n(n+3)} = \sum\limits_{n=1}^{\infty} \left[\dfrac{1}{3n} - \dfrac{1}{3(n+3)}\right]$ [partial fractions].

$s_n = \sum\limits_{i=1}^{n} \left[\dfrac{1}{3i} - \dfrac{1}{3(i+3)}\right] = \dfrac{1}{3} + \dfrac{1}{6} + \dfrac{1}{9} - \dfrac{1}{3(n+1)} - \dfrac{1}{3(n+2)} - \dfrac{1}{3(n+3)}$ (telescoping sum), so

$\sum\limits_{n=1}^{\infty} \dfrac{1}{n(n+3)} = \lim\limits_{n\to\infty} s_n = \dfrac{1}{3} + \dfrac{1}{6} + \dfrac{1}{9} = \dfrac{11}{18}$.

29. $\sum\limits_{n=1}^{\infty} [\tan^{-1}(n+1) - \tan^{-1} n] = \lim\limits_{n\to\infty} s_n$

$$= \lim_{n\to\infty} [(\tan^{-1} 2 - \tan^{-1} 1) + (\tan^{-1} 3 - \tan^{-1} 2) + \cdots + (\tan^{-1}(n+1) - \tan^{-1} n)]$$

$$= \lim_{n\to\infty} [\tan^{-1}(n+1) - \tan^{-1} 1] = \tfrac{\pi}{2} - \tfrac{\pi}{4} = \tfrac{\pi}{4}$$

30. $\sum\limits_{n=0}^{\infty} \dfrac{(-1)^n\pi^n}{3^{2n}\,(2n)!} = \sum\limits_{n=0}^{\infty} (-1)^n \dfrac{1}{(2n)!} \cdot \dfrac{\pi^n}{3^{2n}} = \sum\limits_{n=0}^{\infty} (-1)^n \dfrac{1}{(2n)!} \cdot \left(\dfrac{\sqrt{\pi}}{3}\right)^{2n} = \cos\left(\dfrac{\sqrt{\pi}}{3}\right)$ since $\cos x = \sum\limits_{n=0}^{\infty} (-1)^n \dfrac{x^{2n}}{(2n)!}$ for all x.

31. $1 - e + \dfrac{e^2}{2!} - \dfrac{e^3}{3!} + \dfrac{e^4}{4!} - \cdots = \sum\limits_{n=0}^{\infty} (-1)^n \dfrac{e^n}{n!} = \sum\limits_{n=0}^{\infty} \dfrac{(-e)^n}{n!} = e^{-e}$ since $e^x = \sum\limits_{n=0}^{\infty} \dfrac{x^n}{n!}$ for all x.

32. $4.17\overline{326} = 4.17 + \dfrac{326}{10^5} + \dfrac{326}{10^8} + \cdots = 4.17 + \dfrac{326/10^5}{1 - 1/10^3} = \dfrac{417}{100} + \dfrac{326}{99{,}900} = \dfrac{416{,}909}{99{,}900}$

33. $\cosh x = \dfrac{1}{2}(e^x + e^{-x}) = \dfrac{1}{2}\left(\sum\limits_{n=0}^{\infty} \dfrac{x^n}{n!} + \sum\limits_{n=0}^{\infty} \dfrac{(-x)^n}{n!}\right)$

$$= \frac{1}{2}\left[\left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots\right) + \left(1 - x + \frac{x^2}{2!} - \frac{x^3}{3!} + \frac{x^4}{4!} - \cdots\right)\right]$$

$$= \frac{1}{2}\left(2 + 2 \cdot \frac{x^2}{2!} + 2 \cdot \frac{x^4}{4!} + \cdots\right) = 1 + \frac{1}{2}x^2 + \sum_{n=2}^{\infty} \frac{x^{2n}}{(2n)!} \geq 1 + \frac{1}{2}x^2 \quad \text{for all } x$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. $\sum_{n=1}^{\infty} (\ln x)^n$ is a geometric series which converges whenever $|\ln x| < 1 \quad \Rightarrow \quad -1 < \ln x < 1 \quad \Rightarrow \quad e^{-1} < x < e$.

35. $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^5} = 1 - \frac{1}{32} + \frac{1}{243} - \frac{1}{1024} + \frac{1}{3125} - \frac{1}{7776} + \frac{1}{16{,}807} - \frac{1}{32{,}768} + \cdots$.

Since $b_8 = \frac{1}{8^5} = \frac{1}{32{,}768} < 0.000031$, $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^5} \approx \sum_{n=1}^{7} \frac{(-1)^{n+1}}{n^5} \approx 0.9721$.

36. (a) $s_5 = \sum_{n=1}^{5} \frac{1}{n^6} = 1 + \frac{1}{2^6} + \cdots + \frac{1}{5^6} \approx 1.017305$. The series $\sum_{n=1}^{\infty} \frac{1}{n^6}$ converges by the Integral Test, so we estimate the remainder R_5 with (11.3.2): $R_5 \le \int_5^{\infty} \frac{dx}{x^6} = \left[-\frac{x^{-5}}{5}\right]_5^{\infty} = \frac{5^{-5}}{5} = 0.000064$. So the error is at most 0.000064.

(b) In general, $R_n \le \int_n^{\infty} \frac{dx}{x^6} = \frac{1}{5n^5}$. If we take $n = 9$, then $s_9 \approx 1.01734$ and $R_9 \le \frac{1}{5 \cdot 9^5} \approx 3.4 \times 10^{-6}$.

So to five decimal places, $\sum_{n=1}^{\infty} \frac{1}{n^5} \approx \sum_{n=1}^{9} \frac{1}{n^5} \approx 1.01734$.

Another method: Use (11.3.3) instead of (11.3.2).

37. $\sum_{n=1}^{\infty} \frac{1}{2+5^n} \approx \sum_{n=1}^{8} \frac{1}{2+5^n} \approx 0.18976224$. To estimate the error, note that $\frac{1}{2+5^n} < \frac{1}{5^n}$, so the remainder term is

$R_8 = \sum_{n=9}^{\infty} \frac{1}{2+5^n} < \sum_{n=9}^{\infty} \frac{1}{5^n} = \frac{1/5^9}{1-1/5} = 6.4 \times 10^{-7}$ $\left[\text{geometric series with } a = \frac{1}{5^9} \text{ and } r = \frac{1}{5}\right]$.

38. (a) $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{(n+1)^{n+1}}{[2(n+1)]!} \cdot \frac{(2n)!}{n^n}\right| = \lim_{n\to\infty} \frac{(n+1)^n (n+1)^1}{(2n+2)(2n+1)n^n} = \lim_{n\to\infty} \left(\frac{n+1}{n}\right)^n \frac{1}{2(2n+1)}$

$$= \lim_{n\to\infty} \left(1 + \frac{1}{n}\right)^n \frac{1}{2(2n+1)} = e \cdot 0 = 0 < 1$$

so the given series $\sum_{n=1}^{\infty} \frac{n^n}{(2n)!}$ converges by the Ratio Test.

(b) The series in part (a) is convergent, so $\lim_{n\to\infty} a_n = \lim_{n\to\infty} \frac{n^n}{(2n)!} = 0$ by Theorem 11.2.6.

39. Use the Limit Comparison Test. $\lim_{n\to\infty} \left|\frac{\left(\frac{n+1}{n}\right)a_n}{a_n}\right| = \lim_{n\to\infty} \frac{n+1}{n} = \lim_{n\to\infty} \left(1 + \frac{1}{n}\right) = 1 > 0$.

Since $\sum |a_n|$ is convergent, so is $\sum \left|\left(\frac{n+1}{n}\right)a_n\right|$, by the Limit Comparison Test.

40. $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \left|\frac{x^{n+1}}{(n+1)^2\, 5^{n+1}} \cdot \frac{n^2 5^n}{x^n}\right| = \lim_{n\to\infty} \frac{1}{(1+1/n)^2} \frac{|x|}{5} = \frac{|x|}{5}$, so by the Ratio Test, $\sum_{n=1}^{\infty} (-1)^n \frac{x^n}{n^2\, 5^n}$ converges when $\frac{|x|}{5} < 1 \quad \Leftrightarrow \quad |x| < 5$, so $R = 5$. When $x = -5$, the series becomes the convergent p-series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ with $p = 2 > 1$. When $x = 5$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$, which converges by the Alternating Series Test. Thus, $I = [-5, 5]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left[\dfrac{|x+2|^{n+1}}{(n+1)\,4^{n+1}}\cdot\dfrac{n\,4^n}{|x+2|^n}\right] = \lim\limits_{n\to\infty}\left[\dfrac{n}{n+1}\,\dfrac{|x+2|}{4}\right] = \dfrac{|x+2|}{4} < 1 \quad\Leftrightarrow\quad |x+2| < 4$, so $R = 4$.

$|x+2| < 4 \quad\Leftrightarrow\quad -4 < x+2 < 4 \quad\Leftrightarrow\quad -6 < x < 2$. If $x = -6$, then the series $\sum\limits_{n=1}^{\infty}\dfrac{(x+2)^n}{n\,4^n}$ becomes

$\sum\limits_{n=1}^{\infty}\dfrac{(-4)^n}{n4^n} = \sum\limits_{n=1}^{\infty}\dfrac{(-1)^n}{n}$, the alternating harmonic series, which converges by the Alternating Series Test. When $x = 2$, the

series becomes the harmonic series $\sum\limits_{n=1}^{\infty}\dfrac{1}{n}$, which diverges. Thus, $I = [-6, 2)$.

42. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{2^{n+1}\,(x-2)^{n+1}}{(n+3)!}\cdot\dfrac{(n+2)!}{2^n(x-2)^n}\right| = \lim\limits_{n\to\infty}\dfrac{2}{n+3}\,|x-2| = 0 < 1$, so the series $\sum\limits_{n=1}^{\infty}\dfrac{2^n\,(x-2)^n}{(n+2)!}$

converges for all x. $R = \infty$ and $I = (-\infty, \infty)$.

43. $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{2^{n+1}(x-3)^{n+1}}{\sqrt{n+4}}\cdot\dfrac{\sqrt{n+3}}{2^n(x-3)^n}\right| = 2\,|x-3|\lim\limits_{n\to\infty}\sqrt{\dfrac{n+3}{n+4}} = 2\,|x-3| < 1 \quad\Leftrightarrow\quad |x-3| < \frac{1}{2}$,

so $R = \frac{1}{2}$. $|x-3| < \frac{1}{2} \quad\Leftrightarrow\quad -\frac{1}{2} < x-3 < \frac{1}{2} \quad\Leftrightarrow\quad \frac{5}{2} < x < \frac{7}{2}$. For $x = \frac{7}{2}$, the series $\sum\limits_{n=1}^{\infty}\dfrac{2^n(x-3)^n}{\sqrt{n+3}}$ becomes

$\sum\limits_{n=0}^{\infty}\dfrac{1}{\sqrt{n+3}} = \sum\limits_{n=3}^{\infty}\dfrac{1}{n^{1/2}}$, which diverges $\left[p = \frac{1}{2} \le 1\right]$, but for $x = \frac{5}{2}$, we get $\sum\limits_{n=0}^{\infty}\dfrac{(-1)^n}{\sqrt{n+3}}$, which is a convergent

alternating series, so $I = \left[\frac{5}{2}, \frac{7}{2}\right)$.

44. For $\sum\limits_{n=1}^{\infty}\dfrac{(2n)!}{(n!)^2}\,x^n$, $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{(2n+2)!\,x^{n+1}}{[(n+1)!]^2}\cdot\dfrac{(n!)^2}{(2n)!\,x^n}\right| = \lim\limits_{n\to\infty}\dfrac{(2n+2)(2n+1)}{(n+1)(n+1)}\,|x| = 4\,|x|$.

To converge, we must have $4\,|x| < 1 \quad\Leftrightarrow\quad |x| < \frac{1}{4}$, so $R = \frac{1}{4}$.

45.

n	$f^{(n)}(x)$	$f^{(n)}\left(\frac{\pi}{6}\right)$
0	$\sin x$	$\frac{1}{2}$
1	$\cos x$	$\frac{\sqrt{3}}{2}$
2	$-\sin x$	$-\frac{1}{2}$
3	$-\cos x$	$-\frac{\sqrt{3}}{2}$
4	$\sin x$	$\frac{1}{2}$
$\vdots$	$\vdots$	$\vdots$

$$\begin{aligned}\sin x &= f\left(\frac{\pi}{6}\right) + f'\left(\frac{\pi}{6}\right)\left(x-\frac{\pi}{6}\right) + \frac{f''\left(\frac{\pi}{6}\right)}{2!}\left(x-\frac{\pi}{6}\right)^2 + \frac{f^{(3)}\left(\frac{\pi}{6}\right)}{3!}\left(x-\frac{\pi}{6}\right)^3 + \frac{f^{(4)}\left(\frac{\pi}{6}\right)}{4!}\left(x-\frac{\pi}{6}\right)^4 + \cdots \\ &= \frac{1}{2}\left[1 - \frac{1}{2!}\left(x-\frac{\pi}{6}\right)^2 + \frac{1}{4!}\left(x-\frac{\pi}{6}\right)^4 - \cdots\right] + \frac{\sqrt{3}}{2}\left[\left(x-\frac{\pi}{6}\right) - \frac{1}{3!}\left(x-\frac{\pi}{6}\right)^3 + \cdots\right] \\ &= \frac{1}{2}\sum_{n=0}^{\infty}(-1)^n\frac{1}{(2n)!}\left(x-\frac{\pi}{6}\right)^{2n} + \frac{\sqrt{3}}{2}\sum_{n=0}^{\infty}(-1)^n\frac{1}{(2n+1)!}\left(x-\frac{\pi}{6}\right)^{2n+1}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46.

n	$f^{(n)}(x)$	$f^{(n)}\left(\frac{\pi}{3}\right)$
0	$\cos x$	$\frac{1}{2}$
1	$-\sin x$	$-\frac{\sqrt{3}}{2}$
2	$-\cos x$	$-\frac{1}{2}$
3	$\sin x$	$\frac{\sqrt{3}}{2}$
4	$\cos x$	$\frac{1}{2}$
$\vdots$	$\vdots$	$\vdots$

$$\begin{aligned}\cos x &= f\left(\frac{\pi}{3}\right) + f'\left(\frac{\pi}{3}\right)\left(x-\frac{\pi}{3}\right) + \frac{f''\left(\frac{\pi}{3}\right)}{2!}\left(x-\frac{\pi}{3}\right)^2 + \frac{f^{(3)}\left(\frac{\pi}{3}\right)}{3!}\left(x-\frac{\pi}{3}\right)^3 + \frac{f^{(4)}\left(\frac{\pi}{3}\right)}{4!}\left(x-\frac{\pi}{3}\right)^4 + \cdots \\ &= \frac{1}{2}\left[1 - \frac{1}{2!}\left(x-\frac{\pi}{3}\right)^2 + \frac{1}{4!}\left(x-\frac{\pi}{3}\right)^4 - \cdots\right] + \frac{\sqrt{3}}{2}\left[-\left(x-\frac{\pi}{3}\right) + \frac{1}{3!}\left(x-\frac{\pi}{3}\right)^3 - \cdots\right] \\ &= \frac{1}{2}\sum_{n=0}^{\infty}(-1)^n\frac{1}{(2n)!}\left(x-\frac{\pi}{3}\right)^{2n} + \frac{\sqrt{3}}{2}\sum_{n=0}^{\infty}(-1)^{n+1}\frac{1}{(2n+1)!}\left(x-\frac{\pi}{3}\right)^{2n+1}\end{aligned}$$

47. $\dfrac{1}{1+x} = \dfrac{1}{1-(-x)} = \sum\limits_{n=0}^{\infty}(-x)^n = \sum\limits_{n=0}^{\infty}(-1)^n x^n$ for $|x|<1 \quad\Rightarrow\quad \dfrac{x^2}{1+x} = \sum\limits_{n=0}^{\infty}(-1)^n x^{n+2}$ with $R=1$.

48. $\tan^{-1}x = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{x^{2n+1}}{2n+1}$ with interval of convergence $[-1,1]$, so

$\tan^{-1}(x^2) = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{(x^2)^{2n+1}}{2n+1} = \sum\limits_{n=0}^{\infty}(-1)^n\dfrac{x^{4n+2}}{2n+1}$, which converges when $x^2\in[-1,1] \quad\Leftrightarrow\quad x\in[-1,1]$.

Therefore, $R=1$.

49. $\displaystyle\int\frac{1}{4-x}\,dx = -\ln(4-x)+C$ and

$\displaystyle\int\frac{1}{4-x}\,dx = \frac{1}{4}\int\frac{1}{1-x/4}\,dx = \frac{1}{4}\int\sum_{n=0}^{\infty}\left(\frac{x}{4}\right)^n dx = \frac{1}{4}\int\sum_{n=0}^{\infty}\frac{x^n}{4^n}\,dx = \frac{1}{4}\sum_{n=0}^{\infty}\frac{x^{n+1}}{4^n(n+1)}+C$. So

$\ln(4-x) = -\dfrac{1}{4}\sum\limits_{n=0}^{\infty}\dfrac{x^{n+1}}{4^n(n+1)}+C = -\sum\limits_{n=0}^{\infty}\dfrac{x^{n+1}}{4^{n+1}(n+1)}+C = -\sum\limits_{n=1}^{\infty}\dfrac{x^n}{n4^n}+C$. Putting $x=0$, we get $C=\ln 4$.

Thus, $f(x)=\ln(4-x)=\ln 4-\sum\limits_{n=1}^{\infty}\dfrac{x^n}{n4^n}$. The series converges for $|x/4|<1 \quad\Leftrightarrow\quad |x|<4$, so $R=4$.

Another solution:

$$\begin{aligned}\ln(4-x) &= \ln[4(1-x/4)] = \ln 4+\ln(1-x/4) = \ln 4+\ln[1+(-x/4)] \\ &= \ln 4+\sum_{n=1}^{\infty}(-1)^{n+1}\frac{(-x/4)^n}{n} \quad \text{[from Table 1 in Section 11.10]} \\ &= \ln 4+\sum_{n=1}^{\infty}(-1)^{2n+1}\frac{x^n}{n4^n} = \ln 4-\sum_{n=1}^{\infty}\frac{x^n}{n4^n}.\end{aligned}$$

50. $e^x = \sum\limits_{n=0}^{\infty}\dfrac{x^n}{n!} \quad\Rightarrow\quad e^{2x} = \sum\limits_{n=0}^{\infty}\dfrac{(2x)^n}{n!} \quad\Rightarrow\quad xe^{2x} = x\sum\limits_{n=0}^{\infty}\dfrac{2^n x^n}{n!} = \sum\limits_{n=0}^{\infty}\dfrac{2^n x^{n+1}}{n!},\ R=\infty$

51. $\sin x = \sum\limits_{n=0}^{\infty}\dfrac{(-1)^n x^{2n+1}}{(2n+1)!} \quad\Rightarrow\quad \sin(x^4) = \sum\limits_{n=0}^{\infty}\dfrac{(-1)^n (x^4)^{2n+1}}{(2n+1)!} = \sum\limits_{n=0}^{\infty}\dfrac{(-1)^n x^{8n+4}}{(2n+1)!}$ for all x, so the radius of convergence is ∞.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

52. $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad \Rightarrow \quad 10^x = e^{(\ln 10)x} = \sum_{n=0}^{\infty} \frac{[(\ln 10)x]^n}{n!} = \sum_{n=0}^{\infty} \frac{(\ln 10)^n x^n}{n!}, \ R = \infty$

53. $f(x) = \dfrac{1}{\sqrt[4]{16-x}} = \dfrac{1}{\sqrt[4]{16(1-x/16)}} = \dfrac{1}{\sqrt[4]{16}\left(1-\frac{1}{16}x\right)^{1/4}} = \frac{1}{2}\left(1-\frac{1}{16}x\right)^{-1/4}$

$$= \frac{1}{2}\left[1 + \left(-\frac{1}{4}\right)\left(-\frac{x}{16}\right) + \frac{\left(-\frac{1}{4}\right)\left(-\frac{5}{4}\right)}{2!}\left(-\frac{x}{16}\right)^2 + \frac{\left(-\frac{1}{4}\right)\left(-\frac{5}{4}\right)\left(-\frac{9}{4}\right)}{3!}\left(-\frac{x}{16}\right)^3 + \cdots\right]$$

$$= \frac{1}{2} + \sum_{n=1}^{\infty} \frac{1 \cdot 5 \cdot 9 \cdot \cdots \cdot (4n-3)}{2 \cdot 4^n \cdot n! \cdot 16^n} x^n = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{1 \cdot 5 \cdot 9 \cdot \cdots \cdot (4n-3)}{2^{6n+1}\, n!} x^n$$

for $\left|-\dfrac{x}{16}\right| < 1 \quad \Leftrightarrow \quad |x| < 16$, so $R = 16$.

54. $(1-3x)^{-5} = \sum_{n=0}^{\infty} \binom{-5}{n} (-3x)^n = 1 + (-5)(-3x) + \dfrac{(-5)(-6)}{2!}(-3x)^2 + \dfrac{(-5)(-6)(-7)}{3!}(-3x)^3 + \cdots$

$$= 1 + \sum_{n=1}^{\infty} \frac{5 \cdot 6 \cdot 7 \cdot \cdots \cdot (n+4) \cdot 3^n x^n}{n!} \quad \text{for } |-3x| < 1 \quad \Leftrightarrow \quad |x| < \tfrac{1}{3}, \text{ so } R = \tfrac{1}{3}.$$

55. $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$, so $\dfrac{e^x}{x} = \dfrac{1}{x}\sum_{n=0}^{\infty} \dfrac{x^n}{n!} = \sum_{n=0}^{\infty} \dfrac{x^{n-1}}{n!} = x^{-1} + \sum_{n=1}^{\infty} \dfrac{x^{n-1}}{n!} = \dfrac{1}{x} + \sum_{n=1}^{\infty} \dfrac{x^{n-1}}{n!}$ and

$\displaystyle\int \frac{e^x}{x}\,dx = C + \ln|x| + \sum_{n=1}^{\infty} \frac{x^n}{n \cdot n!}.$

56. $(1+x^4)^{1/2} = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} (x^4)^n = 1 + \left(\frac{1}{2}\right)x^4 + \dfrac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)}{2!}(x^4)^2 + \dfrac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{3!}(x^4)^3 + \cdots$

$$= 1 + \tfrac{1}{2}x^4 - \tfrac{1}{8}x^8 + \tfrac{1}{16}x^{12} - \cdots$$

so $\int_0^1 (1+x^4)^{1/2}\,dx = \left[x + \frac{1}{10}x^5 - \frac{1}{72}x^9 + \frac{1}{208}x^{13} - \cdots\right]_0^1 = 1 + \frac{1}{10} - \frac{1}{72} + \frac{1}{208} - \cdots.$

This is an alternating series, so by the Alternating Series Test, the error in the approximation

$\int_0^1 (1+x^4)^{1/2}\,dx \approx 1 + \frac{1}{10} - \frac{1}{72} \approx 1.086$ is less than $\frac{1}{208}$, sufficient for the desired accuracy.

Thus, correct to two decimal places, $\int_0^1 (1+x^4)^{1/2}\,dx \approx 1.09$.

57. (a)

n	$f^{(n)}(x)$	$f^{(n)}(1)$
0	$x^{1/2}$	1
1	$\frac{1}{2}x^{-1/2}$	$\frac{1}{2}$
2	$-\frac{1}{4}x^{-3/2}$	$-\frac{1}{4}$
3	$\frac{3}{8}x^{-5/2}$	$\frac{3}{8}$
4	$-\frac{15}{16}x^{-7/2}$	$-\frac{15}{16}$
$\vdots$	$\vdots$	$\vdots$

$$\sqrt{x} \approx T_3(x) = 1 + \frac{1/2}{1!}(x-1) - \frac{1/4}{2!}(x-1)^2 + \frac{3/8}{3!}(x-1)^3$$
$$= 1 + \tfrac{1}{2}(x-1) - \tfrac{1}{8}(x-1)^2 + \tfrac{1}{16}(x-1)^3$$

(b)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $|R_3(x)| \le \dfrac{M}{4!}|x-1|^4$, where $\left|f^{(4)}(x)\right| \le M$ with $f^{(4)}(x) = -\frac{15}{16}x^{-7/2}$. Now $0.9 \le x \le 1.1 \quad \Rightarrow$

$-0.1 \le x-1 \le 0.1 \quad \Rightarrow \quad (x-1)^4 \le (0.1)^4$, and letting $x = 0.9$ gives $M = \dfrac{15}{16(0.9)^{7/2}}$, so

$|R_3(x)| \le \dfrac{15}{16(0.9)^{7/2}\,4!}(0.1)^4 \approx 0.000\,005\,648 \approx 0.000\,006 = 6 \times 10^{-6}$.

(d)

From the graph of $|R_3(x)| = |\sqrt{x} - T_3(x)|$, it appears that the error is less than 5×10^{-6} on $[0.9, 1.1]$.

58. (a)

$\sec x \approx T_2(x) = 1 + \frac{1}{2}x^2$

n	$f^{(n)}(x)$	$f^{(n)}(0)$
0	$\sec x$	1
1	$\sec x \tan x$	0
2	$\sec x \tan^2 x + \sec^3 x$	1
3	$\sec x \tan^3 x + 5\sec^3 x \tan x$	0
$\vdots$	$\vdots$	$\vdots$

(b)

(c) $|R_2(x)| \le \dfrac{M}{3!}|x|^3$, where $\left|f^{(3)}(x)\right| \le M$ with $f^{(3)}(x) = \sec x \tan^3 x + 5\sec^3 x \tan x$.

Now $0 \le x \le \frac{\pi}{6} \quad \Rightarrow \quad x^3 \le \left(\frac{\pi}{6}\right)^3$, and letting $x = \frac{\pi}{6}$ gives $M = \frac{14}{3}$, so $|R_2(x)| \le \dfrac{14}{3 \cdot 6}\left(\dfrac{\pi}{6}\right)^3 \approx 0.111648$.

(d)

From the graph of $|R_2(x)| = |\sec x - T_2(x)|$, it appears that the error is less than 0.02 on $\left[0, \frac{\pi}{6}\right]$.

59. $\sin x = \displaystyle\sum_{n=0}^{\infty}(-1)^n \frac{x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots$, so $\sin x - x = -\dfrac{x^3}{3!} + \dfrac{x^5}{5!} - \dfrac{x^7}{7!} + \cdots$ and

$\dfrac{\sin x - x}{x^3} = -\dfrac{1}{3!} + \dfrac{x^2}{5!} - \dfrac{x^4}{7!} + \cdots$. Thus, $\displaystyle\lim_{x\to 0}\frac{\sin x - x}{x^3} = \lim_{x\to 0}\left(-\frac{1}{6} + \frac{x^2}{120} - \frac{x^4}{5040} + \cdots\right) = -\frac{1}{6}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. (a) $F = \dfrac{mgR^2}{(R+h)^2} = \dfrac{mg}{(1+h/R)^2} = mg \sum\limits_{n=0}^{\infty} \dbinom{-2}{n} \left(\dfrac{h}{R}\right)^n$ [binomial series]

(b) We expand $F = mg\left[1 - 2\,(h/R) + 3\,(h/R)^2 - \cdots\right]$.

This is an alternating series, so by the Alternating Series Estimation Theorem, the error in the approximation $F = mg$ is less than $2mgh/R$, so for accuracy within 1% we want

$$\left|\frac{2mgh/R}{mgR^2/(R+h)^2}\right| < 0.01 \quad \Leftrightarrow \quad \frac{2h(R+h)^2}{R^3} < 0.01.$$

This inequality would be difficult to solve for h, so we substitute $R = 6{,}400$ km and plot both sides of the inequality. It appears that the approximation is accurate to within 1% for $h < 31$ km.

61. $f(x) = \sum\limits_{n=0}^{\infty} c_n\, x^n \quad \Rightarrow \quad f(-x) = \sum\limits_{n=0}^{\infty} c_n(-x)^n = \sum\limits_{n=0}^{\infty} (-1)^n c_n\, x^n$

(a) If f is an odd function, then $f(-x) = -f(x) \quad \Rightarrow \quad \sum\limits_{n=0}^{\infty} (-1)^n c_n x^n = \sum\limits_{n=0}^{\infty} -c_n x^n$. The coefficients of any power series are uniquely determined (by Theorem 11.10.5), so $(-1)^n\, c_n = -c_n$.

If n is even, then $(-1)^n = 1$, so $c_n = -c_n \quad \Rightarrow \quad 2c_n = 0 \quad \Rightarrow \quad c_n = 0$. Thus, all even coefficients are 0, that is, $c_0 = c_2 = c_4 = \cdots = 0$.

(b) If f is even, then $f(-x) = f(x) \quad \Rightarrow \quad \sum\limits_{n=0}^{\infty} (-1)^n\, c_n\, x^n = \sum\limits_{n=0}^{\infty} c_n\, x^n \quad \Rightarrow \quad (-1)^n\, c_n = c_n$.

If n is odd, then $(-1)^n = -1$, so $-c_n = c_n \quad \Rightarrow \quad 2c_n = 0 \quad \Rightarrow \quad c_n = 0$. Thus, all odd coefficients are 0, that is, $c_1 = c_3 = c_5 = \cdots = 0$.

62. $e^x = \sum\limits_{n=0}^{\infty} \dfrac{x^n}{n!} \quad \Rightarrow \quad f(x) = e^{x^2} = \sum\limits_{n=0}^{\infty} \dfrac{(x^2)^n}{n!} = \sum\limits_{n=0}^{\infty} \dfrac{x^{2n}}{n!} = \sum\limits_{n=0}^{\infty} \dfrac{1}{n!}\, x^{2n}$. By Theorem 11.10.6 with $a = 0$, we also have $f(x) = \sum\limits_{k=0}^{\infty} \dfrac{f^{(k)}(0)}{k!} x^k$. Comparing coefficients for $k = 2n$, we have $\dfrac{f^{(2n)}(0)}{(2n)!} = \dfrac{1}{n!} \quad \Rightarrow \quad f^{(2n)}(0) = \dfrac{(2n)!}{n!}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1. (a) From Formula 15a in Appendix D, with $x = y = \theta$, we get $\tan 2\theta = \dfrac{2\tan\theta}{1-\tan^2\theta}$, so $\cot 2\theta = \dfrac{1-\tan^2\theta}{2\tan\theta}$ $\Rightarrow$

$2\cot 2\theta = \dfrac{1-\tan^2\theta}{\tan\theta} = \cot\theta - \tan\theta$. Replacing θ by $\frac{1}{2}x$, we get $2\cot x = \cot\frac{1}{2}x - \tan\frac{1}{2}x$, or

$\tan\frac{1}{2}x = \cot\frac{1}{2}x - 2\cot x$.

(b) From part (a) with $\dfrac{x}{2^{n-1}}$ in place of x, $\tan\dfrac{x}{2^n} = \cot\dfrac{x}{2^n} - 2\cot\dfrac{x}{2^{n-1}}$, so the nth partial sum of $\sum\limits_{n=1}^{\infty}\dfrac{1}{2^n}\tan\dfrac{x}{2^n}$ is

$$\begin{aligned} s_n &= \frac{\tan(x/2)}{2} + \frac{\tan(x/4)}{4} + \frac{\tan(x/8)}{8} + \cdots + \frac{\tan(x/2^n)}{2^n} \\ &= \left[\frac{\cot(x/2)}{2} - \cot x\right] + \left[\frac{\cot(x/4)}{4} - \frac{\cot(x/2)}{2}\right] + \left[\frac{\cot(x/8)}{8} - \frac{\cot(x/4)}{4}\right] + \cdots \\ &\qquad + \left[\frac{\cot(x/2^n)}{2^n} - \frac{\cot(x/2^{n-1})}{2^{n-1}}\right] = -\cot x + \frac{\cot(x/2^n)}{2^n} \quad \text{[telescoping sum]} \end{aligned}$$

Now $\dfrac{\cot(x/2^n)}{2^n} = \dfrac{\cos(x/2^n)}{2^n\sin(x/2^n)} = \dfrac{\cos(x/2^n)}{x}\cdot\dfrac{x/2^n}{\sin(x/2^n)} \to \dfrac{1}{x}\cdot 1 = \dfrac{1}{x}$ as $n\to\infty$ since $x/2^n \to 0$ for $x \neq 0$. Therefore, if $x \neq 0$ and $x \neq k\pi$ where k is any integer, then

$$\sum_{n=1}^{\infty}\frac{1}{2^n}\tan\frac{x}{2^n} = \lim_{n\to\infty} s_n = \lim_{n\to\infty}\left(-\cot x + \frac{1}{2^n}\cot\frac{x}{2^n}\right) = -\cot x + \frac{1}{x}$$

If $x = 0$, then all terms in the series are 0, so the sum is 0.

2. $|AP_2|^2 = 2$, $|AP_3|^2 = 2 + 2^2$, $|AP_4|^2 = 2 + 2^2 + \left(2^2\right)^2$, $|AP_5|^2 = 2 + 2^2 + \left(2^2\right)^2 + \left(2^3\right)^2, \ldots,$

$|AP_n|^2 = 2 + 2^2 + \left(2^2\right)^2 + \cdots + (2^{n-2})^2$ [for $n \geq 3$] $= 2 + (4 + 4^2 + 4^3 + \cdots + 4^{n-2})$

$= 2 + \dfrac{4(4^{n-2}-1)}{4-1}$ [finite geometric sum with $a = 4$, $r = 4$] $= \dfrac{6}{3} + \dfrac{4^{n-1}-4}{3} = \dfrac{2}{3} + \dfrac{4^{n-1}}{3}$

So $\tan\angle P_nAP_{n+1} = \dfrac{|P_nP_{n+1}|}{|AP_n|} = \dfrac{2^{n-1}}{\sqrt{\dfrac{2}{3}+\dfrac{4^{n-1}}{3}}} = \dfrac{\sqrt{4^{n-1}}}{\sqrt{\dfrac{2}{3}+\dfrac{4^{n-1}}{3}}} = \dfrac{1}{\sqrt{\dfrac{2}{3\cdot 4^{n-1}}+\dfrac{1}{3}}} \to \sqrt{3}$ as $n\to\infty$.

Thus, $\angle P_nAP_{n+1} \to \frac{\pi}{3}$ as $n\to\infty$.

3. (a) At each stage, each side is replaced by four shorter sides, each of length $\frac{1}{3}$ of the side length at the preceding stage. Writing s_0 and ℓ_0 for the number of sides and the length of the side of the initial triangle, we generate the table at right. In general, we have $s_n = 3\cdot 4^n$ and $\ell_n = \left(\frac{1}{3}\right)^n$, so the length of the perimeter at the nth stage of construction is $p_n = s_n\ell_n = 3\cdot 4^n\cdot\left(\frac{1}{3}\right)^n = 3\cdot\left(\frac{4}{3}\right)^n$.

$s_0 = 3$	$\ell_0 = 1$
$s_1 = 3\cdot 4$	$\ell_1 = 1/3$
$s_2 = 3\cdot 4^2$	$\ell_2 = 1/3^2$
$s_3 = 3\cdot 4^3$	$\ell_3 = 1/3^3$
$\vdots$	$\vdots$

(b) $p_n = \dfrac{4^n}{3^{n-1}} = 4\left(\dfrac{4}{3}\right)^{n-1}$. Since $\frac{4}{3} > 1$, $p_n \to \infty$ as $n\to\infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) The area of each of the small triangles added at a given stage is one-ninth of the area of the triangle added at the preceding stage. Let a be the area of the original triangle. Then the area a_n of each of the small triangles added at stage n is $a_n = a \cdot \dfrac{1}{9^n} = \dfrac{a}{9^n}$. Since a small triangle is added to each side at every stage, it follows that the total area A_n added to the figure at the nth stage is $A_n = s_{n-1} \cdot a_n = 3 \cdot 4^{n-1} \cdot \dfrac{a}{9^n} = a \cdot \dfrac{4^{n-1}}{3^{2n-1}}$. Then the total area enclosed by the snowflake curve is $A = a + A_1 + A_2 + A_3 + \cdots = a + a \cdot \dfrac{1}{3} + a \cdot \dfrac{4}{3^3} + a \cdot \dfrac{4^2}{3^5} + a \cdot \dfrac{4^3}{3^7} + \cdots$. After the first term, this is a geometric series with common ratio $\dfrac{4}{9}$, so $A = a + \dfrac{a/3}{1 - \frac{4}{9}} = a + \dfrac{a}{3} \cdot \dfrac{9}{5} = \dfrac{8a}{5}$. But the area of the original equilateral triangle with side 1 is $a = \dfrac{1}{2} \cdot 1 \cdot \sin \dfrac{\pi}{3} = \dfrac{\sqrt{3}}{4}$. So the area enclosed by the snowflake curve is $\dfrac{8}{5} \cdot \dfrac{\sqrt{3}}{4} = \dfrac{2\sqrt{3}}{5}$.

4. Let the series $S = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{6} + \frac{1}{8} + \frac{1}{9} + \frac{1}{12} + \cdots$. Then every term in S is of the form $\dfrac{1}{2^m 3^n}$, $m, n \geq 0$, and furthermore each term occurs only once. So we can write

$$S = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{1}{2^m 3^n} = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{1}{2^m} \frac{1}{3^n} = \sum_{m=0}^{\infty} \frac{1}{2^m} \sum_{n=0}^{\infty} \frac{1}{3^n} = \frac{1}{1 - \frac{1}{2}} \cdot \frac{1}{1 - \frac{1}{3}} = 2 \cdot \tfrac{3}{2} = 3$$

5. (a) Let $a = \arctan x$ and $b = \arctan y$. Then, from Formula 15b in Appendix D,

$$\tan(a - b) = \frac{\tan a - \tan b}{1 + \tan a \, \tan b} = \frac{\tan(\arctan x) - \tan(\arctan y)}{1 + \tan(\arctan x) \tan(\arctan y)} = \frac{x - y}{1 + xy}$$

Now $\arctan x - \arctan y = a - b = \arctan(\tan(a - b)) = \arctan \dfrac{x - y}{1 + xy}$ since $-\frac{\pi}{2} < a - b < \frac{\pi}{2}$.

(b) From part (a) we have

$$\arctan \tfrac{120}{119} - \arctan \tfrac{1}{239} = \arctan \frac{\frac{120}{119} - \frac{1}{239}}{1 + \frac{120}{119} \cdot \frac{1}{239}} = \arctan \frac{\frac{28{,}561}{28{,}441}}{\frac{28{,}561}{28{,}441}} = \arctan 1 = \tfrac{\pi}{4}$$

(c) Replacing y by $-y$ in the formula of part (a), we get $\arctan x + \arctan y = \arctan \dfrac{x + y}{1 - xy}$. So

$$4 \arctan \tfrac{1}{5} = 2\left(\arctan \tfrac{1}{5} + \arctan \tfrac{1}{5}\right) = 2 \arctan \frac{\frac{1}{5} + \frac{1}{5}}{1 - \frac{1}{5} \cdot \frac{1}{5}} = 2 \arctan \tfrac{5}{12} = \arctan \tfrac{5}{12} + \arctan \tfrac{5}{12}$$

$$= \arctan \frac{\frac{5}{12} + \frac{5}{12}}{1 - \frac{5}{12} \cdot \frac{5}{12}} = \arctan \tfrac{120}{119}$$

Thus, from part (b), we have $4 \arctan \frac{1}{5} - \arctan \frac{1}{239} = \arctan \frac{120}{119} - \arctan \frac{1}{239} = \frac{\pi}{4}$.

(d) From Example 11.9.6 we have $\arctan x = x - \dfrac{x^3}{3} + \dfrac{x^5}{5} - \dfrac{x^7}{7} + \dfrac{x^9}{9} - \dfrac{x^{11}}{11} + \cdots$, so

$$\arctan \frac{1}{5} = \frac{1}{5} - \frac{1}{3 \cdot 5^3} + \frac{1}{5 \cdot 5^5} - \frac{1}{7 \cdot 5^7} + \frac{1}{9 \cdot 5^9} - \frac{1}{11 \cdot 5^{11}} + \cdots$$

This is an alternating series and the size of the terms decreases to 0, so by the Alternating Series Estimation Theorem, the sum lies between s_5 and s_6, that is, $0.197395560 < \arctan \frac{1}{5} < 0.197395562$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(e) From the series in part (d) we get $\arctan \dfrac{1}{239} = \dfrac{1}{239} - \dfrac{1}{3 \cdot 239^3} + \dfrac{1}{5 \cdot 239^5} - \cdots$. The third term is less than 2.6×10^{-13}, so by the Alternating Series Estimation Theorem, we have, to nine decimal places, $\arctan \frac{1}{239} \approx s_2 \approx 0.004184076$. Thus, $0.004184075 < \arctan \frac{1}{239} < 0.004184077$.

(f) From part (c) we have $\pi = 16 \arctan \frac{1}{5} - 4 \arctan \frac{1}{239}$, so from parts (d) and (e) we have $16(0.197395560) - 4(0.004184077) < \pi < 16(0.197395562) - 4(0.004184075) \quad \Rightarrow$ $3.141592652 < \pi < 3.141592692$. So, to 7 decimal places, $\pi \approx 3.1415927$.

6. (a) Let $a = \operatorname{arccot} x$ and $b = \operatorname{arccot} y$ where $0 < a - b < \pi$. Then

$$\cot(a-b) = \frac{1}{\tan(a-b)} = \frac{1 + \tan a \, \tan b}{\tan a - \tan b} = \frac{\dfrac{1}{\cot a} \cdot \dfrac{1}{\cot b} + 1}{\dfrac{1}{\cot a} - \dfrac{1}{\cot b}} \cdot \frac{\cot a \, \cot b}{\cot a \, \cot b}$$

$$= \frac{1 + \cot a \, \cot b}{\cot b - \cot a} = \frac{1 + \cot(\operatorname{arccot} x) \cot(\operatorname{arccot} y)}{\cot(\operatorname{arccot} y) - \cot(\operatorname{arccot} x)} = \frac{1 + xy}{y - x}$$

Now $\operatorname{arccot} x - \operatorname{arccot} y = a - b = \operatorname{arccot}(\cot(a-b)) = \operatorname{arccot} \dfrac{1+xy}{y-x}$ since $0 < a - b < \pi$.

(b) From part (a), we want $\operatorname{arccot}(n^2 + n + 1)$ to equal $\operatorname{arccot} \dfrac{1+xy}{y-x}$. Note that $1 + xy = n^2 + n + 1 \quad \Leftrightarrow$ $xy = n^2 + n = (n+1)n$, so if we let $x = n + 1$ and $y = n$, then $y - x = 1$. Therefore,

$$\operatorname{arccot}(n^2 + n + 1) = \operatorname{arccot}(1 + n(n+1)) = \operatorname{arccot} \frac{1 + n(n+1)}{(n+1) - n} = \operatorname{arccot} n - \operatorname{arccot}(n+1)$$

Thus, we have a telescoping series with nth partial sum

$s_n = [\operatorname{arccot} 0 - \operatorname{arccot} 1] + [\operatorname{arccot} 1 - \operatorname{arccot} 2] + \cdots + [\operatorname{arccot} n - \operatorname{arccot}(n+1)] = \operatorname{arccot} 0 - \operatorname{arccot}(n+1)$.

Thus, $\sum\limits_{n=0}^{\infty} \operatorname{arccot}(n^2 + n + 1) = \lim\limits_{n \to \infty} s_n = \lim\limits_{n \to \infty} [\operatorname{arccot} 0 - \operatorname{arccot}(n+1)] = \frac{\pi}{2} - 0 = \frac{\pi}{2}$.

7. We want $\arctan\left(\dfrac{2}{n^2}\right)$ to equal $\arctan \dfrac{x-y}{1+xy}$. Note that $1 + xy = n^2 \quad \Leftrightarrow \quad xy = n^2 - 1 = (n+1)(n-1)$, so if we let $x = n + 1$ and $y = n - 1$, then $x - y = 2$ and $xy \neq -1$. Thus, from Problem 5(a), $\arctan\left(\dfrac{2}{n^2}\right) = \arctan \dfrac{x-y}{1+xy} = \arctan x - \arctan y = \arctan(n+1) - \arctan(n-1)$. Therefore,

$$\begin{aligned}
\sum_{n=1}^{k} \arctan\left(\frac{2}{n^2}\right) &= \sum_{n=1}^{k} [\arctan(n+1) - \arctan(n-1)] \\
&= \sum_{n=1}^{k} [\arctan(n+1) - \arctan n + \arctan n - \arctan(n-1)] \\
&= \sum_{n=1}^{k} [\arctan(n+1) - \arctan n] + \sum_{n=1}^{k} [\arctan n - \arctan(n-1)] \\
&= [\arctan(k+1) - \arctan 1] + [\arctan k - \arctan 0] \quad \text{[since both sums are telescoping]} \\
&= \arctan(k+1) - \tfrac{\pi}{4} + \arctan k - 0
\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Now $\sum_{n=1}^{k} \arctan\left(\frac{2}{n^2}\right) = \lim_{k\to 0} \sum_{n=1}^{k} \arctan\left(\frac{2}{n^2}\right) = \lim_{k\to\infty} \left[\arctan(k+1) - \frac{\pi}{4} + \arctan k\right] = \frac{\pi}{2} - \frac{\pi}{4} + \frac{\pi}{2} = \frac{3\pi}{4}$.

Note: For all $n \geq 1$, $0 \leq \arctan(n-1) < \arctan(n+1) < \frac{\pi}{2}$, so $-\frac{\pi}{2} < \arctan(n+1) - \arctan(n-1) < \frac{\pi}{2}$, and the identity in Problem 5(a) holds.

8. Let's first try the case $k = 1$: $a_0 + a_1 = 0 \;\Rightarrow\; a_1 = -a_0 \;\Rightarrow$

$$\lim_{n\to\infty} \left(a_0\sqrt{n} + a_1\sqrt{n+1}\right) = \lim_{n\to\infty} \left(a_0\sqrt{n} - a_0\sqrt{n+1}\right) = a_0 \lim_{n\to\infty} \left(\sqrt{n} - \sqrt{n+1}\right) \frac{\sqrt{n}+\sqrt{n+1}}{\sqrt{n}+\sqrt{n+1}}$$
$$= a_0 \lim_{n\to\infty} \frac{-1}{\sqrt{n}+\sqrt{n+1}} = 0$$

In general we have $a_0 + a_1 + \cdots + a_k = 0 \;\Rightarrow\; a_k = -a_0 - a_1 - \cdots - a_{k-1} \;\Rightarrow$

$$\lim_{n\to\infty} \left(a_0\sqrt{n} + a_1\sqrt{n+1} + a_2\sqrt{n+2} + \cdots + a_k\sqrt{n+k}\right)$$
$$= \lim_{n\to\infty} \left(a_0\sqrt{n} + a_1\sqrt{n+1} + \cdots + a_{k-1}\sqrt{n+k-1} - a_0\sqrt{n+k} - a_1\sqrt{n+k} - \cdots - a_{k-1}\sqrt{n+k}\right)$$
$$= a_0 \lim_{n\to\infty} \left(\sqrt{n} - \sqrt{n+k}\right) + a_1 \lim_{n\to\infty} \left(\sqrt{n+1} - \sqrt{n+k}\right) + \cdots + a_{k-1} \lim_{n\to\infty} \left(\sqrt{n+k-1} - \sqrt{n+k}\right)$$

Each of these limits is 0 by the same type of simplification as in the case $k = 1$. So we have

$$\lim_{n\to\infty} \left(a_0\sqrt{n} + a_1\sqrt{n+1} + a_2\sqrt{n+2} + \cdots + a_k\sqrt{n+k}\right) = a_0(0) + a_1(0) + \cdots + a_{k-1}(0) = 0$$

9. We start with the geometric series $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$, $|x| < 1$, and differentiate:

$$\sum_{n=1}^{\infty} nx^{n-1} = \frac{d}{dx}\left(\sum_{n=0}^{\infty} x^n\right) = \frac{d}{dx}\left(\frac{1}{1-x}\right) = \frac{1}{(1-x)^2} \text{ for } |x| < 1 \;\Rightarrow\; \sum_{n=1}^{\infty} nx^n = x\sum_{n=1}^{\infty} nx^{n-1} = \frac{x}{(1-x)^2}$$

for $|x| < 1$. Differentiate again:

$$\sum_{n=1}^{\infty} n^2x^{n-1} = \frac{d}{dx}\frac{x}{(1-x)^2} = \frac{(1-x)^2 - x\cdot 2(1-x)(-1)}{(1-x)^4} = \frac{x+1}{(1-x)^3} \;\Rightarrow\; \sum_{n=1}^{\infty} n^2\,x^n = \frac{x^2+x}{(1-x)^3} \;\Rightarrow$$

$$\sum_{n=1}^{\infty} n^3x^{n-1} = \frac{d}{dx}\frac{x^2+x}{(1-x)^3} = \frac{(1-x)^3(2x+1) - (x^2+x)3(1-x)^2(-1)}{(1-x)^6} = \frac{x^2+4x+1}{(1-x)^4} \;\Rightarrow$$

$\sum_{n=1}^{\infty} n^3x^n = \frac{x^3+4x^2+x}{(1-x)^4}$, $|x| < 1$. The radius of convergence is 1 because that is the radius of convergence for the geometric series we started with. If $x = \pm 1$, the series is $\sum n^3(\pm 1)^n$, which diverges by the Test For Divergence, so the interval of convergence is $(-1, 1)$.

10. Place the y-axis as shown and let the length of each book be L. We want to show that the center of mass of the system of n books lies above the table, that is, $\overline{x} < L$. The x-coordinates of the centers of mass of the books are $x_1 = \frac{L}{2}$, $x_2 = \frac{L}{2(n-1)} + \frac{L}{2}$, $x_3 = \frac{L}{2(n-1)} + \frac{L}{2(n-2)} + \frac{L}{2}$, and so on.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Each book has the same mass m, so if there are n books, then

$$\begin{aligned}
\overline{x} &= \frac{mx_1 + mx_2 + \cdots + mx_n}{mn} = \frac{x_1 + x_2 + \cdots + x_n}{n} \\
&= \frac{1}{n}\left[\frac{L}{2} + \left(\frac{L}{2(n-1)} + \frac{L}{2}\right) + \left(\frac{L}{2(n-1)} + \frac{L}{2(n-2)} + \frac{L}{2}\right) + \cdots \right. \\
&\qquad\qquad \left. + \left(\frac{L}{2(n-1)} + \frac{L}{2(n-2)} + \cdots + \frac{L}{4} + \frac{L}{2} + \frac{L}{2}\right)\right] \\
&= \frac{L}{n}\left[\frac{n-1}{2(n-1)} + \frac{n-2}{2(n-2)} + \cdots + \frac{2}{4} + \frac{1}{2} + \frac{n}{2}\right] = \frac{L}{n}\left[(n-1)\frac{1}{2} + \frac{n}{2}\right] = \frac{2n-1}{2n}L < L
\end{aligned}$$

This shows that, no matter how many books are added according to the given scheme, the center of mass lies above the table. It remains to observe that the series $\frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \frac{1}{8} + \cdots = \frac{1}{2}\sum(1/n)$ is divergent (harmonic series), so we can make the top book extend as far as we like beyond the edge of the table if we add enough books.

11. $$\begin{aligned}
\ln\left(1 - \frac{1}{n^2}\right) &= \ln\left(\frac{n^2-1}{n^2}\right) = \ln\frac{(n+1)(n-1)}{n^2} = \ln[(n+1)(n-1)] - \ln n^2 \\
&= \ln(n+1) + \ln(n-1) - 2\ln n = \ln(n-1) - \ln n - \ln n + \ln(n+1) \\
&= \ln\frac{n-1}{n} - [\ln n - \ln(n+1)] = \ln\frac{n-1}{n} - \ln\frac{n}{n+1}.
\end{aligned}$$

Let $s_k = \sum_{n=2}^{k} \ln\left(1 - \frac{1}{n^2}\right) = \sum_{n=2}^{k}\left(\ln\frac{n-1}{n} - \ln\frac{n}{n+1}\right)$ for $k \geq 2$. Then

$s_k = \left(\ln\frac{1}{2} - \ln\frac{2}{3}\right) + \left(\ln\frac{2}{3} - \ln\frac{3}{4}\right) + \cdots + \left(\ln\frac{k-1}{k} - \ln\frac{k}{k+1}\right) = \ln\frac{1}{2} - \ln\frac{k}{k+1}$, so

$\sum_{n=2}^{\infty} \ln\left(1 - \frac{1}{n^2}\right) = \lim_{k\to\infty} s_k = \lim_{k\to\infty}\left(\ln\frac{1}{2} - \ln\frac{k}{k+1}\right) = \ln\frac{1}{2} - \ln 1 = \ln 1 - \ln 2 - \ln 1 = -\ln 2$ (or $\ln\frac{1}{2}$).

12. First notice that both series are absolutely convergent (p-series with $p > 1$.) Let the given expression be called x. Then

$$\begin{aligned}
x &= \frac{1 + \frac{1}{2^p} + \frac{1}{3^p} + \frac{1}{4^p} + \cdots}{1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots} = \frac{1 + \left(2\cdot\frac{1}{2^p} - \frac{1}{2^p}\right) + \frac{1}{3^p} + \left(2\cdot\frac{1}{4^p} - \frac{1}{4^p}\right) + \cdots}{1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots} \\
&= \frac{\left(1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots\right) + \left(2\cdot\frac{1}{2^p} + 2\cdot\frac{1}{4^p} + 2\cdot\frac{1}{6^p} + \cdots\right)}{1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots} \\
&= 1 + \frac{2\left(\frac{1}{2^p} + \frac{1}{4^p} + \frac{1}{6^p} + \frac{1}{8^p} + \cdots\right)}{1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots} = 1 + \frac{\frac{1}{2^{p-1}}\left(1 + \frac{1}{2^p} + \frac{1}{3^p} + \frac{1}{4^p} + \cdots\right)}{1 - \frac{1}{2^p} + \frac{1}{3^p} - \frac{1}{4^p} + \cdots} = 1 + 2^{1-p}x
\end{aligned}$$

Therefore, $x = 1 + 2^{1-p}x \quad\Leftrightarrow\quad x - 2^{1-p}x = 1 \quad\Leftrightarrow\quad x(1 - 2^{1-p}) = 1 \quad\Leftrightarrow\quad x = \dfrac{1}{1 - 2^{1-p}}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. If L is the length of a side of the equilateral triangle, then the area is $A = \frac{1}{2}L \cdot \frac{\sqrt{3}}{2}L = \frac{\sqrt{3}}{4}L^2$ and so $L^2 = \frac{4}{\sqrt{3}}A$.

Let r be the radius of one of the circles. When there are n rows of circles, the figure shows that

$$L = \sqrt{3}\,r + r + (n-2)(2r) + r + \sqrt{3}\,r = r\big(2n - 2 + 2\sqrt{3}\big), \text{ so } r = \frac{L}{2\big(n+\sqrt{3}-1\big)}.$$

The number of circles is $1 + 2 + \cdots + n = \dfrac{n(n+1)}{2}$, and so the total area of the circles is

$$\begin{aligned} A_n &= \frac{n(n+1)}{2}\pi r^2 = \frac{n(n+1)}{2}\,\pi\,\frac{L^2}{4\big(n+\sqrt{3}-1\big)^2} \\ &= \frac{n(n+1)}{2}\,\pi\,\frac{4A/\sqrt{3}}{4\big(n+\sqrt{3}-1\big)^2} = \frac{n(n+1)}{\big(n+\sqrt{3}-1\big)^2}\,\frac{\pi A}{2\sqrt{3}} \quad \Rightarrow \end{aligned}$$

$$\begin{aligned} \frac{A_n}{A} &= \frac{n(n+1)}{\big(n+\sqrt{3}-1\big)^2}\,\frac{\pi}{2\sqrt{3}} \\ &= \frac{1+1/n}{\big[1+\big(\sqrt{3}-1\big)/n\big]^2}\,\frac{\pi}{2\sqrt{3}} \to \frac{\pi}{2\sqrt{3}} \quad \text{as } n \to \infty \end{aligned}$$

14. Given $a_0 = a_1 = 1$ and $a_n = \dfrac{(n-1)(n-2)a_{n-1} - (n-3)a_{n-2}}{n(n-1)}$, we calculate the next few terms of the sequence:

$a_2 = \dfrac{1 \cdot 0 \cdot a_1 - (-1)a_0}{2 \cdot 1} = \dfrac{1}{2}$, $a_3 = \dfrac{2 \cdot 1 \cdot a_2 - 0 \cdot a_1}{3 \cdot 2} = \dfrac{1}{6}$, $a_4 = \dfrac{3 \cdot 2 \cdot a_3 - 1 \cdot a_2}{4 \cdot 3} = \dfrac{1}{24}$. It seems that $a_n = \dfrac{1}{n!}$,

so we try to prove this by induction. The first step is done, so assume $a_k = \dfrac{1}{k!}$ and $a_{k-1} = \dfrac{1}{(k-1)!}$. Then

$a_{k+1} = \dfrac{k(k-1)a_k - (k-2)a_{k-1}}{(k+1)k} = \dfrac{\dfrac{k(k-1)}{k!} - \dfrac{k-2}{(k-1)!}}{(k+1)k} = \dfrac{(k-1)-(k-2)}{[(k+1)(k)](k-1)!} = \dfrac{1}{(k+1)!}$ and the induction is

complete. Therefore, $\sum\limits_{n=0}^{\infty} a_n = \sum\limits_{n=0}^{\infty} \dfrac{1}{n!} = e$.

15. (a)

The x-intercepts of the curve occur where $\sin x = 0 \quad \Leftrightarrow \quad x = n\pi$, n an integer. So using the formula for disks (and either a CAS or $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$ and Formula 99 to evaluate the integral), the volume of the nth bead is

$$\begin{aligned} V_n &= \pi \textstyle\int_{(n-1)\pi}^{n\pi} \big(e^{-x/10}\sin x\big)^2\,dx = \pi \int_{(n-1)\pi}^{n\pi} e^{-x/5}\sin^2 x\,dx \\ &= \tfrac{250\pi}{101}\big(e^{-(n-1)\pi/5} - e^{-n\pi/5}\big) \end{aligned}$$

(b) The total volume is

$$\pi \textstyle\int_0^{\infty} e^{-x/5}\sin^2 x\,dx = \sum\limits_{n=1}^{\infty} V_n = \tfrac{250\pi}{101}\sum\limits_{n=1}^{\infty}\big[e^{-(n-1)\pi/5} - e^{-n\pi/5}\big] = \tfrac{250\pi}{101} \quad \text{[telescoping sum].}$$

Another method: If the volume in part (a) has been written as $V_n = \frac{250\pi}{101}e^{-n\pi/5}(e^{\pi/5} - 1)$, then we recognize $\sum\limits_{n=1}^{\infty} V_n$ as a geometric series with $a = \frac{250\pi}{101}(1 - e^{-\pi/5})$ and $r = e^{-\pi/5}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. (a) Since P_n is defined as the midpoint of $P_{n-4}P_{n-3}$, $x_n = \frac{1}{2}(x_{n-4} + x_{n-3})$ for $n \geq 5$. So we prove by induction that $\frac{1}{2}x_n + x_{n+1} + x_{n+2} + x_{n+3} = 2$. The case $n = 1$ is immediate, since $\frac{1}{2} \cdot 0 + 1 + 1 + 0 = 2$. Assume that the result holds for $n = k - 1$, that is, $\frac{1}{2}x_{k-1} + x_k + x_{k+1} + x_{k+2} = 2$. Then for $n = k$,

$$\begin{aligned}\tfrac{1}{2}x_k + x_{k+1} + x_{k+2} + x_{k+3} &= \tfrac{1}{2}x_k + x_{k+1} + x_{k+2} + \tfrac{1}{2}(x_{k+3-4} + x_{k+3-3}) \quad \text{[by above]}\\ &= \tfrac{1}{2}x_{k-1} + x_k + x_{k+1} + x_{k+2} = 2 \quad \text{[by the induction hypothesis]}\end{aligned}$$

Similarly, for $n \geq 5$, $y_n = \frac{1}{2}(y_{n-4} + y_{n-3})$, so the same argument as above holds for y, with 2 replaced by $\frac{1}{2}y_1 + y_2 + y_3 + y_4 = \frac{1}{2} \cdot 1 + 1 + 0 + 0 = \frac{3}{2}$. So $\frac{1}{2}y_n + y_{n+1} + y_{n+2} + y_{n+3} = \frac{3}{2}$ for all n.

(b) $\lim\limits_{n\to\infty} \left(\frac{1}{2}x_n + x_{n+1} + x_{n+2} + x_{n+3}\right) = \frac{1}{2}\lim\limits_{n\to\infty} x_n + \lim\limits_{n\to\infty} x_{n+1} + \lim\limits_{n\to\infty} x_{n+2} + \lim\limits_{n\to\infty} x_{n+3} = 2$. Since all the limits on the left hand side are the same, we get $\frac{7}{2}\lim\limits_{n\to\infty} x_n = 2 \quad \Rightarrow \quad \lim\limits_{n\to\infty} x_n = \frac{4}{7}$. In the same way, $\frac{7}{2}\lim\limits_{n\to\infty} y_n = \frac{3}{2} \quad \Rightarrow \quad \lim\limits_{n\to\infty} y_n = \frac{3}{7}$, so $P = \left(\frac{4}{7}, \frac{3}{7}\right)$.

17. By Table 1 in Section 11.10, $\tan^{-1} x = \sum\limits_{n=0}^{\infty} (-1)^n \dfrac{x^{2n+1}}{2n+1}$ for $|x| < 1$. In particular, for $x = \dfrac{1}{\sqrt{3}}$, we have $\dfrac{\pi}{6} = \tan^{-1}\left(\dfrac{1}{\sqrt{3}}\right) = \sum\limits_{n=0}^{\infty} (-1)^n \dfrac{\left(1/\sqrt{3}\right)^{2n+1}}{2n+1} = \sum\limits_{n=0}^{\infty} (-1)^n \left(\dfrac{1}{3}\right)^n \dfrac{1}{\sqrt{3}} \dfrac{1}{2n+1}$, so

$$\pi = \frac{6}{\sqrt{3}} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)3^n} = 2\sqrt{3} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)3^n} = 2\sqrt{3}\left(1 + \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n+1)3^n}\right) \quad \Rightarrow \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n+1)3^n} = \frac{\pi}{2\sqrt{3}} - 1.$$

18. (a) Using $s_n = a + ar + ar^2 + \cdots + ar^{n-1} = \dfrac{a(1-r^n)}{1-r}$,

$$1 - x + x^2 - x^3 + \cdots + x^{2n-2} - x^{2n-1} = \frac{1\left[1 - (-x)^{2n}\right]}{1 - (-x)} = \frac{1 - x^{2n}}{1 + x}.$$

(b) $\displaystyle\int_0^1 (1 - x + x^2 - x^3 + \cdots + x^{2n-2} - x^{2n-1})\,dx = \int_0^1 \frac{1 - x^{2n}}{1 + x}\,dx \quad \Rightarrow$

$$\left[x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots + \frac{x^{2n-1}}{2n-1} - \frac{x^{2n}}{2n}\right]_0^1 = \int_0^1 \frac{dx}{1+x} - \int_0^1 \frac{x^{2n}}{1+x}\,dx \quad \Rightarrow$$

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots + \frac{1}{2n-1} - \frac{1}{2n} = \int_0^1 \frac{dx}{1+x} - \int_0^1 \frac{x^{2n}}{1+x}\,dx$$

(c) Since $1 - \dfrac{1}{2} = \dfrac{1}{1 \cdot 2}$, $\dfrac{1}{3} - \dfrac{1}{4} = \dfrac{1}{3 \cdot 4}, \cdots, \dfrac{1}{2n-1} - \dfrac{1}{2n} = \dfrac{1}{(2n-1)(2n)}$, we see from part (b) that

$$\frac{1}{1 \cdot 2} + \frac{1}{3 \cdot 4} + \cdots + \frac{1}{(2n-1)(2n)} - \int_0^1 \frac{dx}{1+x} = -\int_0^1 \frac{x^{2n}}{1+x}\,dx. \text{ Thus,}$$

$$\left|\frac{1}{1 \cdot 2} + \frac{1}{3 \cdot 4} + \cdots + \frac{1}{(2n-1)(2n)} - \int_0^1 \frac{dx}{1+x}\right| = \int_0^1 \frac{x^{2n}}{1+x}\,dx < \int_0^1 x^{2n}\,dx$$

$$\left[\text{since } \frac{x^{2n}}{1+x} < x^{2n} \text{ for } 0 < x \leq 1\right].$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) Note that $\int_0^1 \frac{dx}{1+x} = \Big[\ln(1+x)\Big]_0^1 = \ln 2$ and $\int_0^1 x^{2n}\,dx = \left[\frac{x^{2n+1}}{2n+1}\right]_0^1 = \frac{1}{2n+1}$. So part (c) becomes

$\left|\frac{1}{1\cdot 2} + \frac{1}{3\cdot 4} + \cdots + \frac{1}{(2n-1)(2n)} - \ln 2\right| < \frac{1}{2n+1}$. In other words, the nth partial sum s_n of the given series satisfies $|s_n - \ln 2| < \frac{1}{2n+1}$. Thus, $\lim_{n\to\infty} s_n = \ln 2$, that is, $\frac{1}{1\cdot 2} + \frac{1}{3\cdot 4} + \frac{1}{5\cdot 6} + \frac{1}{7\cdot 8} + \cdots = \ln 2$.

19. Let $f(x)$ denote the left-hand side of the equation $1 + \frac{x}{2!} + \frac{x^2}{4!} + \frac{x^3}{6!} + \frac{x^4}{8!} + \cdots = 0$. If $x \geq 0$, then $f(x) \geq 1$ and there are no solutions of the equation. Note that $f(-x^2) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \cdots = \cos x$. The solutions of $\cos x = 0$ for $x < 0$ are given by $x = \frac{\pi}{2} - \pi k$, where k is a positive integer. Thus, the solutions of $f(x) = 0$ are $x = -\left(\frac{\pi}{2} - \pi k\right)^2$, where k is a positive integer.

20. Suppose the base of the first right triangle has length a. Then by repeated use of the Pythagorean theorem, we find that the base of the second right triangle has length $\sqrt{1+a^2}$, the base of the third right triangle has length $\sqrt{2+a^2}$, and in general, the nth right triangle has base of length $\sqrt{n-1+a^2}$ and hypotenuse of length $\sqrt{n+a^2}$. Thus, $\theta_n = \tan^{-1}\left(1/\sqrt{n-1+a^2}\right)$ and $\sum_{n=1}^{\infty} \theta_n = \sum_{n=1}^{\infty} \tan^{-1}\left(\frac{1}{\sqrt{n-1+a^2}}\right) = \sum_{n=0}^{\infty} \tan^{-1}\left(\frac{1}{\sqrt{n+a^2}}\right)$. We wish to show that this series diverges.

First notice that the series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n+a^2}}$ diverges by the Limit Comparison Test with the divergent p-series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ $\left[p = \frac{1}{2} \leq 1\right]$ since $\lim_{n\to\infty} \frac{1/\sqrt{n+a^2}}{1/\sqrt{n}} = \lim_{n\to\infty} \frac{\sqrt{n}}{\sqrt{n+a^2}} = \lim_{n\to\infty} \sqrt{\frac{n}{n+a^2}} = \lim_{n\to\infty} \sqrt{\frac{1}{1+a^2/n}} = 1 > 0$. Thus, $\sum_{n=0}^{\infty} \frac{1}{\sqrt{n+a^2}}$ also diverges. Now $\sum_{n=0}^{\infty} \tan^{-1}\left(\frac{1}{\sqrt{n+a^2}}\right)$ diverges by the Limit Comparison Test with $\sum_{n=0}^{\infty} \frac{1}{\sqrt{n+a^2}}$ since

$$\lim_{n\to\infty} \frac{\tan^{-1}\left(1/\sqrt{n+a^2}\right)}{1/\sqrt{n+a^2}} = \lim_{x\to\infty} \frac{\tan^{-1}\left(1/\sqrt{x+a^2}\right)}{1/\sqrt{x+a^2}} = \lim_{y\to\infty} \frac{\tan^{-1}(1/y)}{1/y} \qquad \left[y = \sqrt{x+a^2}\right]$$

$$= \lim_{z\to 0^+} \frac{\tan^{-1} z}{z} \quad \left[z = 1/y\right] \quad \overset{\text{H}}{=} \lim_{z\to 0^+} \frac{1/(1+z^2)}{1} = 1 > 0$$

Thus, $\sum_{n=1}^{\infty} \theta_n$ is a divergent series.

21. Call the series S. We group the terms according to the number of digits in their denominators:

$$S = \underbrace{\left(\tfrac{1}{1} + \tfrac{1}{2} + \cdots + \tfrac{1}{8} + \tfrac{1}{9}\right)}_{g_1} + \underbrace{\left(\tfrac{1}{11} + \cdots + \tfrac{1}{99}\right)}_{g_2} + \underbrace{\left(\tfrac{1}{111} + \cdots + \tfrac{1}{999}\right)}_{g_3} + \cdots$$

Now in the group g_n, since we have 9 choices for each of the n digits in the denominator, there are 9^n terms.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Furthermore, each term in g_n is less than $\frac{1}{10^{n-1}}$ [except for the first term in g_1]. So $g_n < 9^n \cdot \frac{1}{10^{n-1}} = 9\left(\frac{9}{10}\right)^{n-1}$.

Now $\sum\limits_{n=1}^{\infty} 9\left(\frac{9}{10}\right)^{n-1}$ is a geometric series with $a = 9$ and $r = \frac{9}{10} < 1$. Therefore, by the Comparison Test,

$S = \sum\limits_{n=1}^{\infty} g_n < \sum\limits_{n=1}^{\infty} 9\left(\frac{9}{10}\right)^{n-1} = \frac{9}{1-9/10} = 90.$

22. (a) Let $f(x) = \dfrac{x}{1-x-x^2} = \sum\limits_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots$. Then

$$\begin{aligned}
x &= (1-x-x^2)(c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots) \\
x &= c_0 + c_1 x + c_2 x^2 + c_3 x^3 + c_4 x^4 + c_5 x^5 + \cdots \\
&\qquad - c_0 x - c_1 x^2 - c_2 x^3 - c_3 x^4 - c_4 x^5 - \cdots \\
&\qquad\qquad - c_0 x^2 - c_1 x^3 - c_2 x^4 - c_3 x^5 - \cdots \\
x &= c_0 + (c_1 - c_0)x + (c_2 - c_1 - c_0)x^2 + (c_3 - c_2 - c_1)x^3 + \cdots
\end{aligned}$$

Comparing coefficients of powers of x gives us $c_0 = 0$ and

$$\begin{aligned}
c_1 - c_0 &= 1 &&\Rightarrow\quad c_1 = c_0 + 1 = 1 \\
c_2 - c_1 - c_0 &= 0 &&\Rightarrow\quad c_2 = c_1 + c_0 = 1 + 0 = 1 \\
c_3 - c_2 - c_1 &= 0 &&\Rightarrow\quad c_3 = c_2 + c_1 = 1 + 1 = 2
\end{aligned}$$

In general, we have $c_n = c_{n-1} + c_{n-2}$ for $n \geq 3$. Each c_n is equal to the nth Fibonacci number, that is,

$$\sum_{n=0}^{\infty} c_n x^n = \sum_{n=1}^{\infty} c_n x^n = \sum_{n=1}^{\infty} f_n x^n$$

Now $f(x) = \dfrac{x}{1-x-x^2} = \dfrac{x}{1-\left(x^2+x+\frac{1}{4}\right)+\frac{1}{4}} = \dfrac{x}{\frac{5}{4}-\left(x+\frac{1}{2}\right)^2} = \dfrac{4x/5}{1-\frac{4}{5}\left(x+\frac{1}{2}\right)^2} = \dfrac{4x}{5} \cdot \dfrac{1}{1-\frac{4}{5}\left(x+\frac{1}{2}\right)^2}$.

The last fraction is the sum of a geometric series with $r = \frac{4}{5}\left(x+\frac{1}{2}\right)^2$, which converges when $\left|\frac{4}{5}\left(x+\frac{1}{2}\right)^2\right| < 1 \quad \Rightarrow$

$-1 < \dfrac{2}{\sqrt{5}}\left(x+\dfrac{1}{2}\right) < 1 \quad \Rightarrow \quad -\dfrac{\sqrt{5}}{2} < x + \dfrac{1}{2} < \dfrac{\sqrt{5}}{2} \quad \Rightarrow \quad \dfrac{-\sqrt{5}-1}{2} < x < \dfrac{\sqrt{5}-1}{2}$. By Theorem 11.8.4, the Maclaurin series of $f(x)$ converges when $-R < x < R$. (We are finding the radius of convergence of the Maclaurin series of $f(x)$ as opposed to the Taylor series centered at $x = \frac{1}{2}$.) Thus, the radius of convergence must be $R = \dfrac{\sqrt{5}-1}{2}$, which ensures the inequality obtained for the geometric series is still satisfied $\left(\text{since } \left|\dfrac{\sqrt{5}-1}{2}\right| < \left|\dfrac{-\sqrt{5}-1}{2}\right|\right)$.

Another method: Applying the Ratio Test, we have $\lim\limits_{n\to\infty}\left|\dfrac{a_{n+1}}{a_n}\right| = \lim\limits_{n\to\infty}\left|\dfrac{c_{n+1}\,x^{n+1}}{c_n x^n}\right| = |x| \lim\limits_{n\to\infty} \dfrac{c_{n+1}}{c_n}$. In Exercise 11.1.89(b), the preceding limit was shown to be $\lim\limits_{n\to\infty} \dfrac{c_{n+1}}{c_n} = \dfrac{1+\sqrt{5}}{2} = \phi$ (the Golden Ratio). Thus, the Maclaurin series of $f(x)$ converges when $|x|\,\phi < 1$ or $|x| < \dfrac{1}{\phi}$, so the radius of convergence is $R = \dfrac{1}{\phi} = \dfrac{\sqrt{5}-1}{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Completing the square on x^2+x-1 gives us

$$\left(x^2+x+\frac{1}{4}\right)-1-\frac{1}{4}=\left(x+\frac{1}{2}\right)^2-\frac{5}{4}=\left(x+\frac{1}{2}\right)^2-\left(\frac{\sqrt{5}}{2}\right)^2$$
$$=\left(x+\frac{1}{2}+\frac{\sqrt{5}}{2}\right)\left(x+\frac{1}{2}-\frac{\sqrt{5}}{2}\right)=\left(x+\frac{1+\sqrt{5}}{2}\right)\left(x+\frac{1-\sqrt{5}}{2}\right)$$

So $\dfrac{x}{1-x-x^2}=\dfrac{-x}{x^2+x-1}=\dfrac{-x}{\left(x+\frac{1+\sqrt{5}}{2}\right)\left(x+\frac{1-\sqrt{5}}{2}\right)}$. The factors in the denominator are linear, so the partial fraction decomposition is

$$\frac{-x}{\left(x+\frac{1+\sqrt{5}}{2}\right)\left(x+\frac{1-\sqrt{5}}{2}\right)}=\frac{A}{x+\frac{1+\sqrt{5}}{2}}+\frac{B}{x+\frac{1-\sqrt{5}}{2}}-x=A\left(x+\tfrac{1-\sqrt{5}}{2}\right)+B\left(x+\tfrac{1+\sqrt{5}}{2}\right)$$

If $x=\frac{-1+\sqrt{5}}{2}$, then $-\frac{-1+\sqrt{5}}{2}=B\sqrt{5}\quad\Rightarrow\quad B=\frac{1-\sqrt{5}}{2\sqrt{5}}$.

If $x=\frac{-1-\sqrt{5}}{2}$, then $-\frac{-1-\sqrt{5}}{2}=A\left(-\sqrt{5}\right)\quad\Rightarrow\quad A=\frac{1+\sqrt{5}}{-2\sqrt{5}}$. Thus,

$$\frac{x}{1-x-x^2}=\frac{\dfrac{1+\sqrt{5}}{-2\sqrt{5}}}{x+\dfrac{1+\sqrt{5}}{2}}+\frac{\dfrac{1-\sqrt{5}}{2\sqrt{5}}}{x+\dfrac{1-\sqrt{5}}{2}}=\frac{\dfrac{1+\sqrt{5}}{-2\sqrt{5}}}{x+\dfrac{1+\sqrt{5}}{2}}\cdot\frac{\dfrac{2}{1+\sqrt{5}}}{\dfrac{2}{1+\sqrt{5}}}+\frac{\dfrac{1-\sqrt{5}}{2\sqrt{5}}}{x+\dfrac{1-\sqrt{5}}{2}}\cdot\frac{\dfrac{2}{1-\sqrt{5}}}{\dfrac{2}{1-\sqrt{5}}}$$
$$=\frac{-1/\sqrt{5}}{1+\dfrac{2}{1+\sqrt{5}}x}+\frac{1/\sqrt{5}}{1+\dfrac{2}{1-\sqrt{5}}x}=-\frac{1}{\sqrt{5}}\sum_{n=0}^{\infty}\left(-\frac{2}{1+\sqrt{5}}x\right)^n+\frac{1}{\sqrt{5}}\sum_{n=0}^{\infty}\left(-\frac{2}{1-\sqrt{5}}x\right)^n$$
$$=\frac{1}{\sqrt{5}}\sum_{n=0}^{\infty}\left[\left(\frac{-2}{1-\sqrt{5}}\right)^n-\left(\frac{-2}{1+\sqrt{5}}\right)^n\right]x^n$$
$$=\frac{1}{\sqrt{5}}\sum_{n=1}^{\infty}\left[\frac{(-2)^n\left(1+\sqrt{5}\right)^n-(-2)^n\left(1-\sqrt{5}\right)^n}{\left(1-\sqrt{5}\right)^n\left(1+\sqrt{5}\right)^n}\right]x^n\qquad\text{[the } n=0 \text{ term is 0]}$$
$$=\frac{1}{\sqrt{5}}\sum_{n=1}^{\infty}\left[\frac{(-2)^n\left(\left(1+\sqrt{5}\right)^n-\left(1-\sqrt{5}\right)^n\right)}{(1-5)^n}\right]x^n$$
$$=\frac{1}{\sqrt{5}}\sum_{n=1}^{\infty}\left[\frac{\left(1+\sqrt{5}\right)^n-\left(1-\sqrt{5}\right)^n}{2^n}\right]x^n\qquad[(-4)^n=(-2)^n\cdot 2^n]$$

From part (a), this series must equal $\sum_{n=1}^{\infty}f_nx^n$, so $f_n=\dfrac{\left(1+\sqrt{5}\right)^n-\left(1-\sqrt{5}\right)^n}{2^n\sqrt{5}}$, which is an explicit formula for the *n*th Fibonacci number.

23. $u=1+\dfrac{x^3}{3!}+\dfrac{x^6}{6!}+\dfrac{x^9}{9!}+\cdots$, $v=x+\dfrac{x^4}{4!}+\dfrac{x^7}{7!}+\dfrac{x^{10}}{10!}+\cdots$, $w=\dfrac{x^2}{2!}+\dfrac{x^5}{5!}+\dfrac{x^8}{8!}+\cdots$.

Use the Ratio Test to show that the series for u, v, and w have positive radii of convergence (∞ in each case), so

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Theorem 11.9.2 applies, and hence, we may differentiate each of these series:

$$\frac{du}{dx} = \frac{3x^2}{3!} + \frac{6x^5}{6!} + \frac{9x^8}{9!} + \cdots = \frac{x^2}{2!} + \frac{x^5}{5!} + \frac{x^8}{8!} + \cdots = w$$

Similarly, $\dfrac{dv}{dx} = 1 + \dfrac{x^3}{3!} + \dfrac{x^6}{6!} + \dfrac{x^9}{9!} + \cdots = u$, and $\dfrac{dw}{dx} = x + \dfrac{x^4}{4!} + \dfrac{x^7}{7!} + \dfrac{x^{10}}{10!} + \cdots = v$.

So $u' = w$, $v' = u$, and $w' = v$. Now differentiate the left-hand side of the desired equation:

$$\begin{aligned}\frac{d}{dx}(u^3 + v^3 + w^3 - 3uvw) &= 3u^2u' + 3v^2v' + 3w^2w' - 3(u'vw + uv'w + uvw') \\ &= 3u^2w + 3v^2u + 3w^2v - 3(vw^2 + u^2w + uv^2) = 0 \quad \Rightarrow\end{aligned}$$

$u^3 + v^3 + w^3 - 3uvw = C$. To find the value of the constant C, we put $x = 0$ in the last equation and get $1^3 + 0^3 + 0^3 - 3(1 \cdot 0 \cdot 0) = C \quad \Rightarrow \quad C = 1$, so $u^3 + v^3 + w^3 - 3uvw = 1$.

24. To prove: If $n > 1$, then the nth partial sum $s_n = \sum_{i=1}^{n} \dfrac{1}{i}$ of the harmonic series is not an integer.

Proof: Let 2^k be the largest power of 2 that is less than or equal to n and let M be the product of all the odd positive integers that are less than or equal to n. Suppose that $s_n = m$, an integer. Then $M2^k s_n = M2^k m$. Since $n \geq 2$, we have $k \geq 1$, and hence, $M2^k m$ is an even integer. We will show that $M2^k s_n$ is an odd integer, contradicting the equality $M2^k s_n = M2^k m$ and showing that the supposition that s_n is an integer must have been wrong.

$M2^k s_n = M2^k \sum_{i=1}^{n} \dfrac{1}{i} = \sum_{i=1}^{n} \dfrac{M2^k}{i}$. If $1 \leq i \leq n$ and i is odd, then $\dfrac{M}{i}$ is an odd integer since i is one of the odd integers that were multiplied together to form M. Thus, $\dfrac{M2^k}{i}$ is an even integer in this case. If $1 \leq i \leq n$ and i is even, then we can write $i = 2^r l$, where 2^r is the largest power of 2 dividing i and l is odd. If $r < k$, then $\dfrac{M2^k}{i} = \dfrac{2^k}{2^r} \cdot \dfrac{M}{l} = 2^{k-r}\dfrac{M}{l}$, which is an even integer, the product of the even integer 2^{k-r} and the odd integer $\dfrac{M}{l}$. If $r = k$, then $l = 1$, since $l > 1 = l \geq 2 \quad \Rightarrow$ $i = 2^k l \geq 2^k \cdot 2 = 2^{k+1}$, contrary to the choice of 2^k as the largest power of 2 that is less than or equal to n. This shows that $r = k$ only when $i = 2^k$. In that case, $\dfrac{M2^k}{i} = M$, an *odd* integer. Since $\dfrac{M2^k}{i}$ is an even integer for every i except 2^k and $\dfrac{M2^k}{i}$ is an odd integer when $i = 2^k$, we see that $M2^k s_n$ is an odd integer. This concludes the proof.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12 □ VECTORS AND THE GEOMETRY OF SPACE

12.1 Three-Dimensional Coordinate Systems

1. We start at the origin, which has coordinates $(0, 0, 0)$. First we move 4 units along the positive x-axis, affecting only the x-coordinate, bringing us to the point $(4, 0, 0)$. We then move 3 units straight downward, in the negative z-direction. Thus only the z-coordinate is affected, and we arrive at $(4, 0, -3)$.

2.

3. The distance from a point to the yz-plane is the absolute value of the x-coordinate of the point. $C(2, 4, 6)$ has the x-coordinate with the smallest absolute value, so C is the point closest to the yz-plane. $A(-4, 0, -1)$ must lie in the xz-plane since the distance from A to the xz-plane, given by the y-coordinate of A, is 0.

4. The projection of $(2, 3, 5)$ onto the xy-plane is $(2, 3, 0)$; onto the yz-plane, $(0, 3, 5)$; onto the xz-plane, $(2, 0, 5)$.

The length of the diagonal of the box is the distance between the origin and $(2, 3, 5)$, given by

$$\sqrt{(2-0)^2 + (3-0)^2 + (5-0)^2} = \sqrt{38} \approx 6.16$$

5. In $\mathbb{R}^2$, the equation $x = 4$ represents a line parallel to the y-axis and 4 units to the right of it. In $\mathbb{R}^3$, the equation $x = 4$ represents the set $\{(x, y, z) \mid x = 4\}$, the set of all points whose x-coordinate is 4. This is the vertical plane that is parallel to the yz-plane and 4 units in front of it.

6. In $\mathbb{R}^3$, the equation $y = 3$ represents a vertical plane that is parallel to the xz-plane and 3 units to the right of it. The equation $z = 5$ represents a horizontal plane parallel to the xy-plane and 5 units above it. The pair of equations $y = 3$, $z = 5$ represents the set of points that are simultaneously on both planes, or in other words, the line of intersection of the planes $y = 3$, $z = 5$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

This line can also be described as the set $\{(x, 3, 5) \mid x \in \mathbb{R}\}$, which is the set of all points in $\mathbb{R}^3$ whose x-coordinate may vary but whose y- and z-coordinates are fixed at 3 and 5, respectively. Thus the line is parallel to the x-axis and intersects the yz-plane in the point $(0, 3, 5)$.

7. The equation $x + y = 2$ represents the set of all points in $\mathbb{R}^3$ whose x- and y-coordinates have a sum of 2, or equivalently where $y = 2 - x$. This is the set $\{(x, 2 - x, z) \mid x \in \mathbb{R}, z \in \mathbb{R}\}$ which is a vertical plane that intersects the xy-plane in the line $y = 2 - x$, $z = 0$.

8. The equation $x^2 + z^2 = 9$ has no restrictions on y, and the x- and z-coordinates satisfy the equation for a circle of radius 3 with center the origin. Thus the surface $x^2 + z^2 = 9$ in $\mathbb{R}^3$ consists of all possible vertical circles (parallel to the xz-plane) $x^2 + z^2 = 9$, $y = k$, and is therefore a circular cylinder with radius 3 whose axis is the y-axis.

9. The distance between the points $P_1(3, 5, -2)$ and $P_2(-1, 1, -4)$ is

$$|P_1P_2| = \sqrt{(-1-3)^2 + (1-5)^2 + [-4-(-2)]^2} = \sqrt{16+16+4} = 6$$

10. The distance between the points $P_1(-6, -3, 0)$ and $P_2(2, 4, 5)$ is

$$|P_1P_2| = \sqrt{[2-(-6)]^2 + [4-(-3)]^2 + (5-0)^2} = \sqrt{64+49+25} = \sqrt{138}$$

11. We can find the lengths of the sides of the triangle by using the distance formula between pairs of vertices:

$$|PQ| = \sqrt{(7-3)^2 + [0-(-2)]^2 + [1-(-3)]^2} = \sqrt{16+4+16} = \sqrt{36} = 6$$
$$|QR| = \sqrt{(1-7)^2 + (2-0)^2 + (1-1)^2} = \sqrt{36+4+0} = \sqrt{40} = 2\sqrt{10}$$
$$|RP| = \sqrt{(3-1)^2 + (-2-2)^2 + (-3-1)^2} = \sqrt{4+16+16} = \sqrt{36} = 6$$

The longest side is QR, but the Pythagorean Theorem is not satisfied: $|PQ|^2 + |RP|^2 \neq |QR|^2$. Thus PQR is not a right triangle. PQR is isosceles, as two sides have the same length.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. Compute the lengths of the sides of the triangle by using the distance formula between pairs of vertices:

$$|PQ| = \sqrt{(4-2)^2 + [1-(-1)]^2 + (1-0)^2} = \sqrt{4+4+1} = \sqrt{9} = 3$$
$$|QR| = \sqrt{(4-4)^2 + (-5-1)^2 + (4-1)^2} = \sqrt{0+36+9} = \sqrt{45} = 3\sqrt{5}$$
$$|RP| = \sqrt{(2-4)^2 + [-1-(-5)]^2 + (0-4)^2} = \sqrt{4+16+16} = \sqrt{36} = 6$$

Since the Pythagorean Theorem is satisfied by $|PQ|^2 + |RP|^2 = |QR|^2$, PQR is a right triangle. PQR is not isosceles, as no two sides have the same length.

13. (a) First we find the distances between points:

$$|AB| = \sqrt{(3-2)^2 + (7-4)^2 + (-2-2)^2} = \sqrt{26}$$
$$|BC| = \sqrt{(1-3)^2 + (3-7)^2 + [3-(-2)]^2} = \sqrt{45}$$
$$|AC| = \sqrt{(1-2)^2 + (3-4)^2 + (3-2)^2} = \sqrt{3}$$

In order for the points to lie on a straight line, the sum of the two shortest distances must be equal to the longest distance. Since $\sqrt{26} + \sqrt{3} \neq \sqrt{45}$, the three points do not lie on a straight line.

(b) First we find the distances between points:

$$|DE| = \sqrt{(1-0)^2 + [-2-(-5)]^2 + (4-5)^2} = \sqrt{11}$$
$$|EF| = \sqrt{(3-1)^2 + [4-(-2)]^2 + (2-4)^2} = \sqrt{44} = 2\sqrt{11}$$
$$|DF| = \sqrt{(3-0)^2 + [4-(-5)]^2 + (2-5)^2} = \sqrt{99} = 3\sqrt{11}$$

Since $\sqrt{11} + 2\sqrt{11} = 3\sqrt{11}$, the three points lie on a straight line.

14. (a) The distance from a point to the xy-plane is the absolute value of the z-coordinate of the point. Thus, the distance from $(4, -2, 6)$ to the xy-plane is $|6| = 6$.

(b) Similarly, the distance to the yz-plane is the absolute value of the x-coordinate of the point: $|4| = 4$.

(c) The distance to the xz-plane is the absolute value of the y-coordinate of the point: $|-2| = 2$.

(d) The point on the x-axis closest to $(4, -2, 6)$ is the point $(4, 0, 0)$. (Approach the x-axis perpendicularly.)

The distance from $(4, -2, 6)$ to the x-axis is the distance between these two points:

$\sqrt{(4-4)^2 + (-2-0)^2 + (6-0)^2} = \sqrt{40} = 2\sqrt{10} \approx 6.32$.

(e) The point on the y-axis closest to $(4, -2, 6)$ is $(0, -2, 0)$. The distance between these points is

$\sqrt{(4-0)^2 + [-2-(-2)]^2 + (6-0)^2} = \sqrt{52} = 2\sqrt{13} \approx 7.21$.

(f) The point on the z-axis closest to $(4, -2, 6)$ is $(0, 0, 6)$. The distance between these points is

$\sqrt{(4-0)^2 + (-2-0)^2 + (6-6)^2} = \sqrt{20} = 2\sqrt{5} \approx 4.47$.

15. An equation of the sphere with center $(-3, 2, 5)$ and radius 4 is $[x-(-3)]^2 + (y-2)^2 + (z-5)^2 = 4^2$, or $(x+3)^2 + (y-2)^2 + (z-5)^2 = 16$. The intersection of this sphere with the yz-plane is the set of points on the sphere whose x-coordinate is 0. Putting $x = 0$ into the equation, we have $9 + (y-2)^2 + (z-5)^2 = 16$, $x = 0$ or $(y-2)^2 + (z-5)^2 = 7$, $x = 0$, which represents a circle in the yz-plane with center $(0, 2, 5)$ and radius $\sqrt{7}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. An equation of the sphere with center $(2, -6, 4)$ and radius 5 is $(x-2)^2 + [y-(-6)]^2 + (z-4)^2 = 5^2$, or $(x-2)^2 + (y+6)^2 + (z-4)^2 = 25$. The intersection of this sphere with the xy-plane is the set of points on the sphere whose z-coordinate is 0. Putting $z = 0$ into the equation, we have $(x-2)^2 + (y+6)^2 = 9, z = 0$ which represents a circle in the xy-plane with center $(2, -6, 0)$ and radius 3. To find the intersection with the xz-plane, we set $y = 0$: $(x-2)^2 + (z-4)^2 = -11$. Since no points satisfy this equation, the sphere does not intersect the xz-plane. (Also note that the distance from the center of the sphere to the xz-plane is greater than the radius of the sphere.) To find the intersection with the yz-plane, we set $x = 0$: $(y+6)^2 + (z-4)^2 = 21, x = 0$, a circle in the yz-plane with center $(0, -6, 4)$ and radius $\sqrt{21}$.

17. The radius of the sphere is the distance between $(4, 3, -1)$ and $(3, 8, 1)$: $r = \sqrt{(3-4)^2 + (8-3)^2 + [1-(-1)]^2} = \sqrt{30}$. Thus, an equation of the sphere is $(x-3)^2 + (y-8)^2 + (z-1)^2 = 30$.

18. If the sphere passes through the origin, the radius of the sphere must be the distance from the origin to the point $(1, 2, 3)$: $r = \sqrt{(1-0)^2 + (2-0)^2 + (3-0)^2} = \sqrt{14}$. Then an equation of the sphere is $(x-1)^2 + (y-2)^2 + (z-3)^2 = 14$.

19. Completing squares in the equation $x^2 + y^2 + z^2 + 8x - 2z = 8$ gives $(x^2 + 8x + 16) + y^2 + (z^2 - 2z + 1) = 8 + 16 + 1 \quad \Rightarrow \quad (x+4)^2 + y^2 + (z-1)^2 = 25$, which we recognize as an equation of a sphere with center $(-4, 0, 1)$ and radius $\sqrt{25} = 5$.

20. Completing squares in the equation $x^2 - 6x + y^2 + 4y + z^2 + 10z = 0$ gives $(x^2 - 6x + 9) + (y^2 + 4y + 4) + (z^2 + 10z + 25) = 9 + 4 + 25 \quad \Rightarrow \quad (x-3)^2 + (y+2)^2 + (z+5)^2 = 38$, which we recognize as an equation of a sphere with center $(3, -2, -5)$ and radius $\sqrt{38}$.

21. Completing squares in the equation $2x^2 - 2x + 2y^2 + 4y + 2z^2 = -1$ gives

$2\left(x^2 - x + \frac{1}{4}\right) + 2(y^2 + 2y + 1) + 2z^2 = -1 + \frac{1}{2} + 2 \quad \Rightarrow \quad 2\left(x - \frac{1}{2}\right)^2 + 2(y+1)^2 + 2z^2 = \frac{3}{2} \quad \Rightarrow$

$\left(x - \frac{1}{2}\right)^2 + (y+1)^2 + z^2 = \frac{3}{4}$, which we recognize as an equation of a sphere with center $\left(\frac{1}{2}, -1, 0\right)$ and radius $\sqrt{\frac{3}{4}} = \frac{\sqrt{3}}{2}$.

22. Completing the squares in the equation $4x^2 - 16x + 4y^2 + 6y + 4z^2 = -12$ gives

$4(x^2 - 4x + 4) + 4\left(y^2 + \frac{3}{2}y + \frac{9}{16}\right) + 4z^2 = -12 + 16 + \frac{9}{4} \quad \Rightarrow \quad 4(x-2)^2 + 4\left(y + \frac{3}{4}\right)^2 + 4z^2 = \frac{25}{4} \quad \Rightarrow$

$(x-2)^2 + \left(y + \frac{3}{4}\right)^2 + z^2 = \frac{25}{16}$, which we recognize as the equation of a sphere with center $\left(2, -\frac{3}{4}, 0\right)$ and radius $\sqrt{\frac{25}{16}} = \frac{5}{4}$.

23. If the midpoint of the line segment from $P_1(x_1, y_1, z_1)$ to $P_2(x_2, y_2, z_2)$ is $Q = \left(\dfrac{x_1 + x_2}{2}, \dfrac{y_1 + y_2}{2}, \dfrac{z_1 + z_2}{2}\right)$, then the distances $|P_1Q|$ and $|QP_2|$ are equal, and each is half of $|P_1P_2|$. We verify that this is the case:

$$|P_1P_2| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\begin{aligned}|P_1Q| &= \sqrt{\left[\tfrac{1}{2}(x_1+x_2)-x_1\right]^2+\left[\tfrac{1}{2}(y_1+y_2)-y_1\right]^2+\left[\tfrac{1}{2}(z_1+z_2)-z_1\right]^2}\\ &= \sqrt{\left(\tfrac{1}{2}x_2-\tfrac{1}{2}x_1\right)^2+\left(\tfrac{1}{2}y_2-\tfrac{1}{2}y_1\right)^2+\left(\tfrac{1}{2}z_2-\tfrac{1}{2}z_1\right)^2}=\sqrt{\left(\tfrac{1}{2}\right)^2\left[(x_2-x_1)^2+(y_2-y_1)^2+(z_2-z_1)^2\right]}\\ &= \tfrac{1}{2}\sqrt{(x_2-x_1)^2+(y_2-y_1)^2+(z_2-z_1)^2}=\tfrac{1}{2}\,|P_1P_2|\end{aligned}$$

$$\begin{aligned}|QP_2| &= \sqrt{\left[x_2-\tfrac{1}{2}(x_1+x_2)\right]^2+\left[y_2-\tfrac{1}{2}(y_1+y_2)\right]^2+\left[z_2-\tfrac{1}{2}(z_1+z_2)\right]^2}\\ &= \sqrt{\left(\tfrac{1}{2}x_2-\tfrac{1}{2}x_1\right)^2+\left(\tfrac{1}{2}y_2-\tfrac{1}{2}y_1\right)^2+\left(\tfrac{1}{2}z_2-\tfrac{1}{2}z_1\right)^2}=\sqrt{\left(\tfrac{1}{2}\right)^2\left[(x_2-x_1)^2+(y_2-y_1)^2+(z_2-z_1)^2\right]}\\ &= \tfrac{1}{2}\sqrt{(x_2-x_1)^2+(y_2-y_1)^2+(z_2-z_1)^2}=\tfrac{1}{2}\,|P_1P_2|\end{aligned}$$

So Q is indeed the midpoint of P_1P_2.

24. By Exercise 23(a), the midpoint of the diameter that has endpoints $(5,4,3)$ and $(1,6,-9)$ (and thus the center of the sphere) is $\left(\dfrac{5+1}{2},\dfrac{4+6}{2},\dfrac{3+(-9)}{2}\right)=(3,5,-3)$. The radius is half the diameter, so $r=\frac{1}{2}\sqrt{(1-5)^2+(6-4)^2+(-9-3)^2}=\frac{1}{2}\sqrt{164}=\sqrt{41}$. Therefore, an equation of the sphere is $(x-3)^2+(y-5)^2+(z+3)^2=41$.

25. (a) Since the sphere touches the xy-plane, its radius is the distance from its center, $(-1,4,5)$, to the xy-plane, which is 5. Therefore, an equation is $(x+1)^2+(y-4)^2+(z-5)^2=25$.

(b) Since the sphere touches the yz-plane, its radius is the distance from its center, $(-1,4,5)$, to the yz-plane, which is 1. Therefore, an equation is $(x+1)^2+(y-4)^2+(z-5)^2=1$.

(c) Since the sphere touches the xz-plane, its radius is the distance from its center, $(-1,4,5)$, to the xz-plane, which is 4. Therefore, an equation is $(x+1)^2+(y-4)^2+(z-5)^2=16$.

26. The shortest distance from the center, $(7,3,8)$, to any of the three coordinate planes is 3, which is the distance to the xz-plane. Therefore, an equation of the sphere is $(x-7)^2+(y-3)^2+(z-8)^2=9$.

27. The equation $z=-2$ represents a plane, parallel to the xy-plane and 2 units below it.

28. The equation $x=3$ represents a plane, parallel to the yz-plane and 3 units in front of it.

29. The inequality $y\geq 1$ represents a half-space consisting of all the points on or to the right of the plane $y=1$.

30. The inequality $x<4$ represents a half-space consisting of all the points behind the plane $x=4$.

31. The inequality $-1\leq x\leq 2$ represents all points on or between the vertical planes $x=-1$ and $x=2$.

32. The equation $z=y$ represents a plane, perpendicular to the yz-plane, and intersecting the yz-plane in the line $z=y$, $x=0$.

33. Because $z=-1$, all points in the region must lie in the horizontal plane $z=-1$. In addition, $x^2+y^2=4$, so the region consists of all points that lie on a circle with radius 2 and center on the z-axis that is contained in the plane $z=-1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. Here $x^2 + y^2 = 4$ with no restrictions on z, so a point in the region must lie on a circle of radius 2, center on the z-axis, but it could be in any horizontal plane $z = k$ (parallel to the xy-plane). Thus the region consists of all possible circles $x^2 + y^2 = 4$, $z = k$ and is therefore a circular cylinder with radius 2 whose axis is the z-axis.

35. The inequality $y^2 + z^2 \leq 25$ is equivalent to $\sqrt{y^2 + z^2} \leq 5$, which describes the set of all points in $\mathbb{R}^3$ whose distance from the x-axis is at most 5. Thus, the inequality represents the region consisting of all points on or inside a circular cylinder of radius 5 with axis the x-axis.

36. The inequality $x^2 + z^2 \leq 25$ is equivalent to $\sqrt{x^2 + z^2} \leq 5$, which describes the set of all points in $\mathbb{R}^3$ whose distance from the y-axis is at most 5. Further, $0 \leq y \leq 2$ consists of the points on or between the planes $y = 0$ and $y = 2$. Thus, the inequalities represent the region consisting of all points on or inside a circular cylinder of radius 5 with axis the y-axis from $y = 0$ to $y = 2$.

37. The equation $x^2 + y^2 + z^2 = 4$ is equivalent to $\sqrt{x^2 + y^2 + z^2} = 2$, so the region consists of those points whose distance from the origin is 2. This is the set of all points on a sphere with radius 2 and center $(0, 0, 0)$.

38. The inequality $x^2 + y^2 + z^2 \leq 4$ is equivalent to $\sqrt{x^2 + y^2 + z^2} \leq 2$, so the region consists of those points whose distance from the origin is at most 2. This is the set of all points on or inside a sphere with radius 2 and center $(0, 0, 0)$.

39. The inequalities $1 \leq x^2 + y^2 + z^2 \leq 5$ are equivalent to $1 \leq \sqrt{x^2 + y^2 + z^2} \leq \sqrt{5}$, so the region consists of those points whose distance from the origin is at least 1 and at most $\sqrt{5}$. This is the set of all points on or between spheres with radii 1 and $\sqrt{5}$ and centers $(0, 0, 0)$.

40. The inequalities $1 \leq x^2 + y^2 \leq 5$ are equivalent to $1 \leq \sqrt{x^2 + y^2} \leq \sqrt{5}$, which represents the set of all points in $\mathbb{R}^3$ whose distance is at least 1 and at most $\sqrt{5}$ from the z-axis. Thus, the region consists of all points on or between a circular cylinder of radius 1 and a circular cylinder of radius $\sqrt{5}$ with axis the z-axis.

41. The inequalities $0 \leq x \leq 3, 0 \leq y \leq 3, 0 \leq z \leq 3$ represent the set of all points in $\mathbb{R}^3$ that lie on or between the planes $x = 3$, $y = 3$, $z = 3$ in the first octant. Thus, the region is a cube with dimensions $3 \times 3 \times 3$.

42. The inequality $x^2 + y^2 + z^2 > 2z \quad \Leftrightarrow \quad x^2 + y^2 + (z - 1)^2 > 1$ is equivalent to $\sqrt{x^2 + y^2 + (z - 1)^2} > 1$, so the region consists of those points whose distance from the point $(0, 0, 1)$ is greater than 1. This is the set of all points outside the sphere with radius 1 and center $(0, 0, 1)$.

43. This describes all points whose x-coordinate is between 0 and 5, that is, $0 < x < 5$.

44. For any point on or above the disk in the xy-plane with center the origin and radius 2 we have $x^2 + y^2 \leq 4$. Also each point lies on or between the planes $z = 0$ and $z = 8$, so the region is described by $x^2 + y^2 \leq 4,\ 0 \leq z \leq 8$.

45. This describes a region all of whose points have a distance to the origin which is greater than r, but smaller than R. So inequalities describing the region are $r < \sqrt{x^2 + y^2 + z^2} < R$, or $r^2 < x^2 + y^2 + z^2 < R^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. The solid sphere itself is represented by $\sqrt{x^2+y^2+z^2} \le 2$. Since we want only the upper hemisphere, we restrict the z-coordinate to nonnegative values. Then inequalities describing the region are $\sqrt{x^2+y^2+z^2} \le 2$, $z \ge 0$, or $x^2+y^2+z^2 \le 4$, $z \ge 0$.

47. (a) To find the x- and y-coordinates of the point P, we project it onto L_2 and project the resulting point Q onto the x- and y-axes. To find the z-coordinate, we project P onto either the xz-plane or the yz-plane (using our knowledge of its x- or y-coordinate) and then project the resulting point onto the z-axis. (Or, we could draw a line parallel to QO from P to the z-axis.) The coordinates of P are $(2, 1, 4)$.

(b) A is the intersection of L_1 and L_2, B is directly below the y-intercept of L_2, and C is directly above the x-intercept of L_2.

48. Let $P = (x, y, z)$. Then $2\,|PB| = |PA| \;\Leftrightarrow\; 4\,|PB|^2 = |PA|^2 \;\Leftrightarrow$

$4\left[(x-6)^2+(y-2)^2+(z+2)^2\right] = (x+1)^2+(y-5)^2+(z-3)^2 \;\Leftrightarrow$

$4\left(x^2-12x+36\right) - x^2 - 2x + 4\left(y^2-4y+4\right) - y^2 + 10y + 4\left(z^2+4z+4\right) - z^2 + 6z = 35 \;\Leftrightarrow$

$3x^2 - 50x + 3y^2 - 6y + 3z^2 + 22z = 35 - 144 - 16 - 16 \;\Leftrightarrow\; x^2 - \frac{50}{3}x + y^2 - 2y + z^2 + \frac{22}{3}z = -\frac{141}{3}$.

By completing the square three times we get $\left(x-\frac{25}{3}\right)^2 + (y-1)^2 + \left(z+\frac{11}{3}\right)^2 = \frac{-423\,+\,625\,+\,9\,+\,121}{9} = \frac{332}{9}$, which is an equation of a sphere with center $\left(\frac{25}{3}, 1, -\frac{11}{3}\right)$ and radius $\frac{\sqrt{332}}{3}$.

49. We need to find a set of points $\left\{P(x,y,z) \;\middle|\; |AP| = |BP|\right\}$.

$\sqrt{(x+1)^2+(y-5)^2+(z-3)^2} = \sqrt{(x-6)^2+(y-2)^2+(z+2)^2} \;\Rightarrow$

$(x+1)^2+(y-5)^2+(z-3)^2 = (x-6)^2+(y-2)^2+(z+2)^2 \;\Rightarrow$

$x^2+2x+1+y^2-10y+25+z^2-6z+9 = x^2-12x+36+y^2-4y+4+z^2+4z+4 \;\Rightarrow\; 14x-6y-10z = 9$.

Thus, the set of points is a plane perpendicular to the line segment joining A and B (since this plane must contain the perpendicular bisector of the line segment AB).

50. Completing the square three times in the first equation gives $(x+2)^2+(y-1)^2+(z+2)^2 = 2^2$, a sphere with center $(-2, 1, -2)$ and radius 2. The second equation is that of a sphere with center $(0, 0, 0)$ and radius 2. The distance between the centers of the spheres is $\sqrt{(-2-0)^2+(1-0)^2+(-2-0)^2} = \sqrt{4+1+4} = 3$. Since the spheres have the same radius, the volume inside both spheres is symmetrical about the plane containing the circle of intersection of the spheres. The distance from this plane to the center of the circles is $\frac{3}{2}$. So the region inside both spheres consists of two caps of spheres of height $h = 2 - \frac{3}{2} = \frac{1}{2}$. From Exercise 6.2.61, the volume of a cap of a sphere is $V = \pi h^2\left(r - \frac{1}{3}h\right) = \pi\left(\frac{1}{2}\right)^2\left(2 - \frac{1}{3}\cdot\frac{1}{2}\right) = \frac{11\pi}{24}$. So the total volume is $2 \cdot \frac{11\pi}{24} = \frac{11\pi}{12}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. The sphere $x^2 + y^2 + z^2 = 4$ has center $(0, 0, 0)$ and radius 2. Completing squares in $x^2 - 4x + y^2 - 4y + z^2 - 4z = -11$ gives $(x^2 - 4x + 4) + (y^2 - 4y + 4) + (z^2 - 4z + 4) = -11 + 4 + 4 + 4 \quad \Rightarrow \quad (x - 2)^2 + (y - 2)^2 + (z - 2)^2 = 1$, so this is the sphere with center $(2, 2, 2)$ and radius 1. The (shortest) distance between the spheres is measured along the line segment connecting their centers. The distance between $(0, 0, 0)$ and $(2, 2, 2)$ is $\sqrt{(2-0)^2 + (2-0)^2 + (2-0)^2} = \sqrt{12} = 2\sqrt{3}$, and subtracting the radius of each circle, the distance between the spheres is $2\sqrt{3} - 2 - 1 = 2\sqrt{3} - 3$.

52. There are many different solids that fit the given description. However, any possible solid must have a circular horizontal cross-section at its top or at its base. Here we illustrate a solid with a circular base in the xy-plane. (A circular cross-section at the top results in an inverted version of the solid described below.) The vertical cross-section through the center of the base that is parallel to the xz-plane must be a square, and the vertical cross-section parallel to the yz-plane (perpendicular to the square) through the center of the base must be a triangle with two vertices on the circle and the third vertex at the center of the top side of the square. (See the figure.)

The solid can include any additional points that do not extend beyond these three "silhouettes" when viewed from directions parallel to the coordinate axes. One possibility shown here is to draw the circular base and the vertical square first. Then draw a surface formed by line segments parallel to the yz-plane that connect the top of the square to the circle.

Problem 8 in the Problems Plus section at the end of the chapter illustrates another possible solid.

12.2 Vectors

1. (a) The cost of a theater ticket is a scalar, because it has only magnitude.

(b) The current in a river is a vector, because it has both magnitude (the speed of the current) and direction at any given location.

(c) If we assume that the initial path is linear, the initial flight path from Houston to Dallas is a vector, because it has both magnitude (distance) and direction.

(d) The population of the world is a scalar, because it has only magnitude.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. If the initial point of the vector $\langle 4, 7\rangle$ is placed at the origin, then $\langle 4, 7\rangle$ is the position vector of the point $(4, 7)$.

3. Vectors are equal when they share the same length and direction (but not necessarily location). Using the symmetry of the parallelogram as a guide, we see that $\overrightarrow{AB} = \overrightarrow{DC}$, $\overrightarrow{DA} = \overrightarrow{CB}$, $\overrightarrow{DE} = \overrightarrow{EB}$, and $\overrightarrow{EA} = \overrightarrow{CE}$.

4. (a) The initial point of $\overrightarrow{BC}$ is positioned at the terminal point of $\overrightarrow{AB}$, so by the Triangle Law the sum $\overrightarrow{AB} + \overrightarrow{BC}$ is the vector with initial point A and terminal point C, namely $\overrightarrow{AC}$.

(b) By the Triangle Law, $\overrightarrow{CD} + \overrightarrow{DB}$ is the vector with initial point C and terminal point B, namely $\overrightarrow{CB}$.

(c) First we consider $\overrightarrow{DB} - \overrightarrow{AB}$ as $\overrightarrow{DB} + \left(-\overrightarrow{AB}\right)$. Then since $-\overrightarrow{AB}$ has the same length as $\overrightarrow{AB}$ but points in the opposite direction, we have $-\overrightarrow{AB} = \overrightarrow{BA}$ and so $\overrightarrow{DB} - \overrightarrow{AB} = \overrightarrow{DB} + \overrightarrow{BA} = \overrightarrow{DA}$.

(d) We use the Triangle Law twice: $\overrightarrow{DC} + \overrightarrow{CA} + \overrightarrow{AB} = \left(\overrightarrow{DC} + \overrightarrow{CA}\right) + \overrightarrow{AB} = \overrightarrow{DA} + \overrightarrow{AB} = \overrightarrow{DB}$.

5. (a)

(b)

(c)

(d)

(e)

(f)

6. (a)

(b)

(c)

(d)

(e)

(f)

7. Because the tail of $\mathbf{d}$ is the midpoint of QR we have $\overrightarrow{QR} = 2\mathbf{d}$, and by the Triangle Law, $\mathbf{a} + 2\mathbf{d} = \mathbf{b} \quad \Rightarrow$ $2\mathbf{d} = \mathbf{b} - \mathbf{a} \quad \Rightarrow \quad \mathbf{d} = \frac{1}{2}(\mathbf{b} - \mathbf{a}) = \frac{1}{2}\mathbf{b} - \frac{1}{2}\mathbf{a}$. Again by the Triangle Law, we have $\mathbf{c} + \mathbf{d} = \mathbf{b}$ so $\mathbf{c} = \mathbf{b} - \mathbf{d} = \mathbf{b} - \left(\frac{1}{2}\mathbf{b} - \frac{1}{2}\mathbf{a}\right) = \frac{1}{2}\mathbf{a} + \frac{1}{2}\mathbf{b}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. We are given $\mathbf{u}+\mathbf{v}+\mathbf{w}=\mathbf{0}$, so $\mathbf{w}=(-\mathbf{u})+(-\mathbf{v})$. (See the figure.)

Vectors $-\mathbf{u}$, $-\mathbf{v}$, and $\mathbf{w}$ form a right triangle, so from the Pythagorean Theorem we have $|-\mathbf{u}|^2+|-\mathbf{v}|^2=|\mathbf{w}|^2$. But $|-\mathbf{u}|=|\mathbf{u}|=1$ and $|-\mathbf{v}|=|\mathbf{v}|=1$, so $|\mathbf{w}|=\sqrt{|-\mathbf{u}|^2+|-\mathbf{v}|^2}=\sqrt{2}$.

9. $\mathbf{a}=\langle 1-(-2),2-1\rangle=\langle 3,1\rangle$

10. $\mathbf{a}=\langle -3-(-5),3-(-1)\rangle=\langle 2,4\rangle$

11. $\mathbf{a}=\langle 2-3,3-(-1)\rangle=\langle -1,4\rangle$

12. $\mathbf{a}=\langle 1-3,0-2\rangle=\langle -2,-2\rangle$

13. $\mathbf{a}=\langle -2-1,3-(-2),0-4\rangle=\langle -3,5,-4\rangle$

14. $\mathbf{a}=\langle 0-3,5-0,0-(-2)\rangle=\langle -3,5,2\rangle$

15. $\langle -1,4\rangle+\langle 6,-2\rangle=\langle -1+6,4+(-2)\rangle=\langle 5,2\rangle$

16. $\langle 3,-1\rangle+\langle -1,5\rangle=\langle 3+(-1),-1+5\rangle=\langle 2,4\rangle$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\langle 3,0,1\rangle + \langle 0,8,0\rangle = \langle 3+0, 0+8, 1+0\rangle$
$= \langle 3,8,1\rangle$

18. $\langle 1,3,-2\rangle + \langle 0,0,6\rangle = \langle 1+0, 3+0, -2+6\rangle$
$= \langle 1,3,4\rangle$

19. $\mathbf{a}+\mathbf{b} = \langle -3,4\rangle + \langle 9,-1\rangle = \langle -3+9, 4+(-1)\rangle = \langle 6,3\rangle$

$4\,\mathbf{a}+2\,\mathbf{b} = 4\,\langle -3,4\rangle + 2\,\langle 9,-1\rangle = \langle -12,16\rangle + \langle 18,-2\rangle = \langle 6,14\rangle$

$|\mathbf{a}| = \sqrt{(-3)^2+4^2} = \sqrt{25} = 5$

$|\mathbf{a}-\mathbf{b}| = |\langle -3-9, 4-(-1)\rangle| = |\langle -12,5\rangle| = \sqrt{(-12)^2+5^2} = \sqrt{169} = 13$

20. $\mathbf{a}+\mathbf{b} = (5\,\mathbf{i}+3\,\mathbf{j}) + (-\mathbf{i}-2\,\mathbf{j}) = 4\,\mathbf{i}+\mathbf{j}$

$4\,\mathbf{a}+2\,\mathbf{b} = 4\,(5\,\mathbf{i}+3\,\mathbf{j}) + 2\,(-\mathbf{i}-2\,\mathbf{j}) = 20\,\mathbf{i}+12\,\mathbf{j}-2\,\mathbf{i}-4\,\mathbf{j} = 18\,\mathbf{i}+8\,\mathbf{j}$

$|\mathbf{a}| = \sqrt{5^2+3^2} = \sqrt{34}$

$|\mathbf{a}-\mathbf{b}| = |(5\,\mathbf{i}+3\,\mathbf{j}) - (-\mathbf{i}-2\,\mathbf{j})| = |6\,\mathbf{i}+5\,\mathbf{j}| = \sqrt{6^2+5^2} = \sqrt{61}$

21. $\mathbf{a}+\mathbf{b} = (4\,\mathbf{i}-3\,\mathbf{j}+2\,\mathbf{k}) + (2\,\mathbf{i}-4\,\mathbf{k}) = 6\,\mathbf{i}-3\,\mathbf{j}-2\mathbf{k}$

$4\,\mathbf{a}+2\,\mathbf{b} = 4\,(4\,\mathbf{i}-3\,\mathbf{j}+2\,\mathbf{k}) + 2\,(2\,\mathbf{i}-4\,\mathbf{k}) = 16\,\mathbf{i}-12\,\mathbf{j}+8\,\mathbf{k}+4\,\mathbf{i}-8\,\mathbf{k} = 20\,\mathbf{i}-12\,\mathbf{j}$

$|\mathbf{a}| = \sqrt{4^2+(-3)^2+2^2} = \sqrt{29}$

$|\mathbf{a}-\mathbf{b}| = |(4\,\mathbf{i}-3\,\mathbf{j}+2\,\mathbf{k}) - (2\,\mathbf{i}-4\,\mathbf{k})| = |2\,\mathbf{i}-3\,\mathbf{j}+6\,\mathbf{k}| = \sqrt{2^2+(-3)^2+6^2} = \sqrt{49} = 7$

22. $\mathbf{a}+\mathbf{b} = \langle 8,1,-4\rangle + \langle 5,-2,1\rangle = \langle 8+5, 1+(-2), -4+1\rangle = \langle 13,-1,-3\rangle$

$4\,\mathbf{a}+2\,\mathbf{b} = 4\,\langle 8,1,-4\rangle + 2\,\langle 5,-2,1\rangle = \langle 32,4,-16\rangle + \langle 10,-4,2\rangle = \langle 42,0,-14\rangle$

$|\mathbf{a}| = \sqrt{8^2+1^2+(-4)^2} = \sqrt{81} = 9$

$|\mathbf{a}-\mathbf{b}| = |\langle 8-5, 1-(-2), -4-1\rangle| = |\langle 3,3,-5\rangle| = \sqrt{3^2+3^2+(-5)^2} = \sqrt{43}$

23. The vector $\langle 6,-2\rangle$ has length $|\langle 6,-2\rangle| = \sqrt{6^2+(-2)^2} = \sqrt{40} = 2\sqrt{10}$, so by Equation 4 the unit vector with the same direction is $\dfrac{1}{2\sqrt{10}}\,\langle 6,-2\rangle = \left\langle \dfrac{3}{\sqrt{10}}, -\dfrac{1}{\sqrt{10}} \right\rangle$.

24. The vector $-5\,\mathbf{i}+3\,\mathbf{j}-\mathbf{k}$ has length $|-5\,\mathbf{i}+3\,\mathbf{j}-\mathbf{k}| = \sqrt{(-5)^2+3^2+(-1)^2} = \sqrt{35}$, so by Equation 4 the unit vector with the same direction is $\dfrac{1}{\sqrt{35}}(-5\,\mathbf{i}+3\,\mathbf{j}-\mathbf{k}) = -\dfrac{5}{\sqrt{35}}\,\mathbf{i} + \dfrac{3}{\sqrt{35}}\,\mathbf{j} - \dfrac{1}{\sqrt{35}}\,\mathbf{k}$.

25. The vector $8\,\mathbf{i}-\mathbf{j}+4\,\mathbf{k}$ has length $|8\,\mathbf{i}-\mathbf{j}+4\,\mathbf{k}| = \sqrt{8^2+(-1)^2+4^2} = \sqrt{81} = 9$, so by Equation 4 the unit vector with the same direction is $\frac{1}{9}(8\,\mathbf{i}-\mathbf{j}+4\,\mathbf{k}) = \frac{8}{9}\,\mathbf{i} - \frac{1}{9}\,\mathbf{j} + \frac{4}{9}\,\mathbf{k}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $|\langle 6, 2, -3\rangle| = \sqrt{6^2 + 2^2 + (-3)^2} = \sqrt{49} = 7$, so a unit vector in the direction of $\langle 6, 2, -3\rangle$ is $\mathbf{u} = \frac{1}{7}\langle 6, 2, -3\rangle$.

A vector in the same direction but with length 4 is $4\mathbf{u} = 4 \cdot \frac{1}{7}\langle 6, 2, -3\rangle = \left\langle \frac{24}{7}, \frac{8}{7}, -\frac{12}{7}\right\rangle$.

27.

From the figure, we see that $\tan\theta = \dfrac{\sqrt{3}}{1} = \sqrt{3} \quad \Rightarrow \quad \theta = 60°$.

28.

From the figure, we see that $\tan\theta = \frac{6}{8} = \frac{3}{4}$, so $\theta = \tan^{-1}\left(\frac{3}{4}\right) \approx 36.9°$.

29. From the figure, we see that the x-component of $\mathbf{v}$ is

$v_1 = |\mathbf{v}|\cos(5\pi/6) = 4\left(-\frac{\sqrt{3}}{2}\right) = -2\sqrt{3}$ and the y-component is

$v_2 = |\mathbf{v}|\sin(5\pi/6) = 4\left(\frac{1}{2}\right) = 2$. Thus, $\mathbf{v} = \langle -2\sqrt{3}, 2\rangle$.

30. From the figure, we see that the horizontal component of the force $\mathbf{F}$ is $|\mathbf{F}|\cos 38° = 50\cos 38° \approx 39.4$ N, and the vertical component is $|\mathbf{F}|\sin 38° = 50\sin 38° \approx 30.8$ N.

31. The velocity vector $\mathbf{v}$ makes an angle of $40°$ with the horizontal and has magnitude equal to the speed at which the football was thrown. From the figure, we see that the horizontal component of $\mathbf{v}$ is $|\mathbf{v}|\cos 40° = 60\cos 40° \approx 45.96$ ft/s and the vertical component is $|\mathbf{v}|\sin 40° = 60\sin 40° \approx 38.57$ ft/s.

32. The given force vectors can be expressed in terms of their horizontal and vertical components as $20\cos 45°\,\mathbf{i} + 20\sin 45°\,\mathbf{j} = 10\sqrt{2}\,\mathbf{i} + 10\sqrt{2}\,\mathbf{j}$ and $16\cos 30°\,\mathbf{i} - 16\sin 30°\,\mathbf{j} = 8\sqrt{3}\,\mathbf{i} - 8\,\mathbf{j}$. The resultant force $\mathbf{F}$ is the sum of these two vectors: $\mathbf{F} = \left(10\sqrt{2} + 8\sqrt{3}\right)\mathbf{i} + \left(10\sqrt{2} - 8\right)\mathbf{j} \approx 28.00\,\mathbf{i} + 6.14\,\mathbf{j}$. Then we have $|\mathbf{F}| \approx \sqrt{(28.00)^2 + (6.14)^2} \approx 28.7$ lb and, letting θ be the angle $\mathbf{F}$ makes with the positive x-axis,

$$\tan\theta = \frac{10\sqrt{2} - 8}{10\sqrt{2} + 8\sqrt{3}} \quad \Rightarrow \quad \theta = \tan^{-1}\left(\frac{10\sqrt{2} - 8}{10\sqrt{2} + 8\sqrt{3}}\right) \approx 12.4°.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. The given force vectors can be expressed in terms of their horizontal and vertical components as $-300\,\mathbf{i}$ and $200\cos 60^\circ\,\mathbf{i} + 200\sin 60^\circ\,\mathbf{j} = 200\left(\frac{1}{2}\right)\,\mathbf{i} + 200\left(\frac{\sqrt{3}}{2}\right)\,\mathbf{j} = 100\,\mathbf{i} + 100\sqrt{3}\,\mathbf{j}$. The resultant force $\mathbf{F}$ is the sum of these two vectors: $\mathbf{F} = (-300+100)\,\mathbf{i} + \left(0 + 100\sqrt{3}\right)\mathbf{j} = -200\,\mathbf{i} + 100\sqrt{3}\,\mathbf{j}$. Then we have $|\mathbf{F}| \approx \sqrt{(-200)^2 + \left(100\sqrt{3}\right)^2} = \sqrt{70{,}000} = 100\sqrt{7} \approx 264.6$ N. Let θ be the angle $\mathbf{F}$ makes with the positive x-axis. Then $\tan\theta = \dfrac{100\sqrt{3}}{-200} = -\dfrac{\sqrt{3}}{2}$ and the terminal point of $\mathbf{F}$ lies in the second quadrant, so $\theta = \tan^{-1}\left(-\dfrac{\sqrt{3}}{2}\right) + 180^\circ \approx -40.9^\circ + 180^\circ = 139.1^\circ$.

34. Let $\mathbf{T}_1$ and $\mathbf{T}_2$ be the tension vectors corresponding to the support cables as shown in the figure. In terms of vertical and horizontal components,

$$\mathbf{T}_1 = |\mathbf{T}_1|\cos 60^\circ\mathbf{i} + |\mathbf{T}_1|\sin 60^\circ\mathbf{j} = \frac{1}{2}\,|\mathbf{T}_1|\;\mathbf{i} + \frac{\sqrt{3}}{2}\,|\mathbf{T}_1|\;\mathbf{j}$$

$$\mathbf{T}_2 = -\,|\mathbf{T}_2|\cos 60^\circ\mathbf{i} + |\mathbf{T}_2|\sin 60^\circ\mathbf{j} = -\frac{1}{2}\,|\mathbf{T}_2|\;\mathbf{i} + \frac{\sqrt{3}}{2}\,|\mathbf{T}_2|\;\mathbf{j}$$

The resultant of these tensions, $\mathbf{T}_1 + \mathbf{T}_2$, counterbalances the weight $\mathbf{w} = -500\,\mathbf{j}$. So $\mathbf{T}_1 + \mathbf{T}_2 = -\mathbf{w} = 500\,\mathbf{j} \;\Rightarrow$

$$\left(\frac{1}{2}\,|\mathbf{T}_1|\;\mathbf{i} + \frac{\sqrt{3}}{2}\,|\mathbf{T}_1|\;\mathbf{j}\right) + \left(-\frac{1}{2}\,|\mathbf{T}_2|\;\mathbf{i} + \frac{\sqrt{3}}{2}\,|\mathbf{T}_2|\;\mathbf{j}\right) = 500\,\mathbf{j}.$$

Equating x-components gives $\frac{1}{2}\,|\mathbf{T}_1|\;\mathbf{i} - \frac{1}{2}\,|\mathbf{T}_2|\;\mathbf{i} = 0$, so $|\mathbf{T}_1| = |\mathbf{T}_2|$ (as we would expect from the symmetry of the problem). Equating y-components, we have $\dfrac{\sqrt{3}}{2}\,|\mathbf{T}_1|\;\mathbf{j} + \dfrac{\sqrt{3}}{2}\,|\mathbf{T}_2|\;\mathbf{j} = \sqrt{3}\,|\mathbf{T}_1|\;\mathbf{j} = 500\,\mathbf{j} \;\Rightarrow\; |\mathbf{T}_1| = \dfrac{500}{\sqrt{3}}$. Thus the magnitude of each tension is $|\mathbf{T}_1| = |\mathbf{T}_2| = \dfrac{500}{\sqrt{3}} \approx 288.68$ lb. The tension vectors are

$$\mathbf{T}_1 = \frac{1}{2}\,|\mathbf{T}_1|\;\mathbf{i} + \frac{\sqrt{3}}{2}\,|\mathbf{T}_1|\;\mathbf{j} = \frac{250}{\sqrt{3}}\,\mathbf{i} + 250\,\mathbf{j} \approx 144.34\,\mathbf{i} + 250\,\mathbf{j} \text{ and } \mathbf{T}_2 = -\frac{250}{\sqrt{3}}\,\mathbf{i} + 250\,\mathbf{j} \approx -144.34\,\mathbf{i} + 250\,\mathbf{j}.$$

35. Call the two tension vectors $\mathbf{T}_2$ and $\mathbf{T}_3$, corresponding to the ropes of length 2 m and 3 m. In terms of vertical and horizontal components,

$$\mathbf{T}_2 = -\,|\mathbf{T}_2|\cos 50^\circ\mathbf{i} + |\mathbf{T}_2|\sin 50^\circ\mathbf{j} \quad \textbf{(1)} \qquad \text{and} \qquad \mathbf{T}_3 = |\mathbf{T}_3|\cos 38^\circ\mathbf{i} + |\mathbf{T}_3|\sin 38^\circ\mathbf{j} \quad \textbf{(2)}$$

The resultant of these forces, $\mathbf{T}_2 + \mathbf{T}_3$, counterbalances the weight of the hoist (which is $-350\,\mathbf{j}$), so $\mathbf{T}_2 + \mathbf{T}_3 = 350\,\mathbf{j} \;\Rightarrow$ $\left(-\,|\mathbf{T}_2|\cos 50^\circ + |\mathbf{T}_3|\cos 38^\circ\right)\mathbf{i} + \left(|\mathbf{T}_2|\sin 50^\circ + |\mathbf{T}_3|\sin 38^\circ\right)\mathbf{j} = 350\,\mathbf{j}$. Equating components, we have $-\,|\mathbf{T}_2|\cos 50^\circ + |\mathbf{T}_3|\cos 38^\circ = 0 \;\Rightarrow\; |\mathbf{T}_2| = |\mathbf{T}_3|\,\dfrac{\cos 38^\circ}{\cos 50^\circ}$ and $|\mathbf{T}_2|\sin 50^\circ + |\mathbf{T}_3|\sin 38^\circ = 350$. Substituting the first equation into the second gives $|\mathbf{T}_3|\,\dfrac{\cos 38^\circ}{\cos 50^\circ}\sin 50^\circ + |\mathbf{T}_3|\sin 38^\circ = 350 \;\Rightarrow\; |\mathbf{T}_3|\left(\cos 38^\circ\tan 50^\circ + \sin 38^\circ\right) = 350$, so the magnitudes of the tensions are $|\mathbf{T}_3| = \dfrac{350}{\cos 38^\circ\tan 50^\circ + \sin 38^\circ} \approx 225.11$ N and $|\mathbf{T}_2| = |\mathbf{T}_3|\,\dfrac{\cos 38^\circ}{\cos 50^\circ} \approx 275.97$ N. Finally, from **(1)** and **(2)**, the tension vectors are $\mathbf{T}_2 \approx -177.39\,\mathbf{i} + 211.41\,\mathbf{j}$ and $\mathbf{T}_3 \approx 177.39\,\mathbf{i} + 138.59\,\mathbf{j}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. We can consider the weight of the chain to be concentrated at its midpoint. The forces acting on the chain then are the tension vectors $\mathbf{T}_1$, $\mathbf{T}_2$ in each end of the chain and the weight $\mathbf{w}$, as shown in the figure. We know $|\mathbf{T}_1| = |\mathbf{T}_2| = 25$ N so, in terms of vertical and horizontal components, we have

$$\mathbf{T}_1 = -25\cos 37°\,\mathbf{i} + 25\sin 37°\,\mathbf{j} \qquad \mathbf{T}_2 = 25\cos 37°\,\mathbf{i} + 25\sin 37°\,\mathbf{j}$$

The resultant vector $\mathbf{T}_1 + \mathbf{T}_2$ of the tensions counterbalances the weight $\mathbf{w}$, giving $\mathbf{T}_1 + \mathbf{T}_2 = -\mathbf{w}$. Since $\mathbf{w} = -|\mathbf{w}|\,\mathbf{j}$, we have $(-25\cos 37°\,\mathbf{i} + 25\sin 37°\,\mathbf{j}) + (25\cos 37°\,\mathbf{i} + 25\sin 37°\,\mathbf{j}) = |\mathbf{w}|\,\mathbf{j} \;\Rightarrow\; 50\sin 37°\,\mathbf{j} = |\mathbf{w}|\,\mathbf{j} \;\Rightarrow$ $|\mathbf{w}| = 50\sin 37° \approx 30.1$. So the weight is 30.1 N, and since $w = mg$, the mass is $\frac{30.1}{9.8} \approx 3.07$ kg.

37. Let $\mathbf{v}_1$, $\mathbf{v}_2$, and $\mathbf{v}_3$ be the force vectors where $|\mathbf{v}_1| = 25$, $|\mathbf{v}_2| = 12$, and $|\mathbf{v}_3| = 4$. Set up coordinate axes so that the object is at the origin and $\mathbf{v}_1$, $\mathbf{v}_2$ lie in the xy-plane. We can position the vectors so that $\mathbf{v}_1 = 25\,\mathbf{i}$, $\mathbf{v}_2 = 12\cos 100°\,\mathbf{i} + 12\sin 100°\,\mathbf{j}$, and $\mathbf{v}_3 = 4\,\mathbf{k}$. The magnitude of a force that counterbalances the three given forces must match the magnitude of the resultant force. We have $\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3 = (25 + 12\cos 100°)\,\mathbf{i} + 12\sin 100°\,\mathbf{j} + 4\,\mathbf{k}$, so the counterbalancing force must have magnitude $|\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3| = \sqrt{(25 + 12\cos 100°)^2 + (12\sin 100°)^2 + 4^2} \approx 26.1$ N.

38. (a) Set up coordinate axes so that the rower is at the origin, the channel is bordered by the y-axis and the line $x = 1400$, and the current flows in the negative y direction. The rower wants to reach the point $(1400, 800)$. Let θ be the angle between the positive y-axis in the direction she should steer. (See the figure.)

In still water, the rower has velocity $\mathbf{v}_r = \langle 7\sin\theta, 7\cos\theta\rangle$ and the velocity of the current is $\mathbf{v}_c = \langle 0, -3\rangle$, so the true course of the rower is determined by the velocity vector $\mathbf{v} = \mathbf{v}_r + \mathbf{v}_c = \langle 7\sin\theta, 7\cos\theta - 3\rangle$. Let t be the time in seconds after the rower departs. Then the position of the rower is given by $t\mathbf{v}$ and the rower crosses the channel when

$$t\mathbf{v} = t\,\langle 7\sin\theta, 7\cos\theta - 3\rangle = \langle 1400, 800\rangle \quad\Rightarrow\quad 7t\sin\theta = 1400 \qquad \text{and} \qquad (7\cos\theta - 3)t = 800$$

Then $t = \dfrac{1400}{7\sin\theta} = \dfrac{200}{\sin\theta}$ and substituting gives

$$(7\cos\theta - 3)\left(\frac{200}{\sin\theta}\right) = 800 \quad\Rightarrow\quad 7\cos\theta - 3 = 4\sin\theta \qquad \textbf{(1)}$$

Squaring both sides, we have

$$49\cos^2\theta - 42\cos\theta + 9 = 16\sin^2\theta = 16(1 - \cos^2\theta)$$

$$65\cos^2\theta - 42\cos\theta - 7 = 0$$

The quadratic formula gives

$$\cos\theta = \frac{42 \pm \sqrt{(-42)^2 - 4(65)(-7)}}{2(65)} = \frac{42 \pm \sqrt{3584}}{130} \approx 0.78359 \text{ or } -0.13743$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The acute value for θ is approximately $\cos^{-1}(0.78359) \approx 38.4°$. Thus, the rower should steer in the direction that is $38.4°$ from the bank, toward upstream.

Alternate solution: We could solve **(1)** graphically by plotting $y_1 = 7\cos\theta - 3$ and $y_2 = 4\sin\theta$ on a graphing device and finding the approximate intersection point $(0.6704, 4.9702)$. Thus, $\theta \approx 0.6704$ radians, or equivalently, $38.4°$.

(b) From part (a) we know the trip is completed when $t = \dfrac{200}{\sin\theta}$. As $\theta \approx 38.4°$, the time required is approximately $\dfrac{200}{\sin(38.4°)} \approx 321.9$ seconds or 5.4 minutes.

39. Set up the coordinate axes so that north is the positive y direction and west is the negative x direction. With respect to the still air, the velocity of the plane can be written as $\mathbf{v}_{\text{plane}} = \langle -180\sin 45°, 180\cos 45° \rangle$ and the velocity of the wind is given by $\mathbf{v}_{\text{wind}} = \langle 35\sin 30°, -35\cos 30° \rangle$. (See the figure.)

Then the velocity vector of the plane relative to the ground is

$$\begin{aligned}\mathbf{v} = \mathbf{v}_{\text{plane}} + \mathbf{v}_{\text{wind}} &= \langle -180\sin 45°, 180\cos 45° \rangle + \langle 35\sin 30°, -35\cos 30° \rangle \\ &= \left\langle -90\sqrt{2} + 35/2, 90\sqrt{2} - 35\sqrt{3}/2 \right\rangle \approx \langle -109.8, 97.0 \rangle\end{aligned}$$

The ground speed is $|\mathbf{v}| \approx \sqrt{(-109.8)^2 + (97.0)^2} \approx 146.5$ mi/h. The angle the velocity vector makes with the x-axis is about $\tan^{-1}\left(\dfrac{97.0}{-109.8}\right) \approx -41.5°$ and $-41.5° + 180° = 138.5°$. Therefore, the course of the plane is about N $(138.5 - 90)°$ W or N $48.5°$ W.

40. With respect to the water's surface, the dog's velocity is the sum of the velocity of the ship with respect to the water and the velocity of the dog with respect to the ship. If we let north be the positive y direction and west be the negative x direction, we have $\mathbf{v} = \langle -32, 0 \rangle + \langle 0, 4 \rangle = \langle -32, 4 \rangle$. Then, the speed of the dog is $|\mathbf{v}| = \sqrt{(-32)^2 + 4^2} \approx 32.2$ km/h. The vector $\mathbf{v}$ makes an angle of $\tan^{-1}\left(\dfrac{4}{-32}\right) \approx -7.1°$ and $-7.1° + 180° = 172.9°$. Therefore, the dog's direction is N $(172.9 - 90)°$ W or N $82.9°$ W.

41. The slope of the tangent line to the graph of $y = x^2$ at the point $(2, 4)$ is

$$\left.\frac{dy}{dx}\right|_{x=2} = \left.2x\right|_{x=2} = 4$$

Thus, a parallel vector is $\mathbf{i} + 4\,\mathbf{j}$, which has length $|\mathbf{i} + 4\,\mathbf{j}| = \sqrt{1^2 + 4^2} = \sqrt{17}$, and so unit vectors parallel to the tangent line are $\pm\frac{1}{\sqrt{17}}\,(\mathbf{i} + 4\,\mathbf{j})$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. (a) The slope of the tangent line to the graph of $y = 2\sin x$ at the point $(\pi/6, 1)$ is

$$\left.\frac{dy}{dx}\right|_{x=\pi/6} = \left.2\cos x\right|_{x=\pi/6} = 2 \cdot \frac{\sqrt{3}}{2} = \sqrt{3}$$

Thus, a parallel vector is $\mathbf{i} + \sqrt{3}\,\mathbf{j}$, which has length $\left|\mathbf{i} + \sqrt{3}\,\mathbf{j}\right| = \sqrt{1^2 + \left(\sqrt{3}\right)^2} = \sqrt{4} = 2$, and so unit vectors parallel to the tangent line are $\pm\frac{1}{2}\left(\mathbf{i} + \sqrt{3}\,\mathbf{j}\right)$.

(b) The slope of the tangent line is $\sqrt{3}$, so the slope of a line perpendicular to the tangent line is $-\frac{1}{\sqrt{3}}$ and a vector in this direction is $\sqrt{3}\,\mathbf{i} - \mathbf{j}$. Since $\left|\sqrt{3}\,\mathbf{i} - \mathbf{j}\right| = \sqrt{\left(\sqrt{3}\right)^2 + (-1)^2} = 2$, unit vectors perpendicular to the tangent line are $\pm\frac{1}{2}\left(\sqrt{3}\,\mathbf{i} - \mathbf{j}\right)$.

(c)

43. By the Triangle Law, $\overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC}$. Then $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA} = \overrightarrow{AC} + \overrightarrow{CA}$, but $\overrightarrow{AC} + \overrightarrow{CA} = \overrightarrow{AC} + \left(-\overrightarrow{AC}\right) = \mathbf{0}$. So $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA} = \mathbf{0}$.

44. $\overrightarrow{AC} = \frac{1}{3}\overrightarrow{AB}$ and $\overrightarrow{BC} = \frac{2}{3}\overrightarrow{BA}$. $\mathbf{c} = \overrightarrow{OA} + \overrightarrow{AC} = \mathbf{a} + \frac{1}{3}\overrightarrow{AB} \quad \Rightarrow \quad \overrightarrow{AB} = 3\,\mathbf{c} - 3\,\mathbf{a}$. $\mathbf{c} = \overrightarrow{OB} + \overrightarrow{BC} = \overrightarrow{OB} + \frac{2}{3}\overrightarrow{BA} \quad \Rightarrow$ $\overrightarrow{BA} = \frac{3}{2}\,\mathbf{c} - \frac{3}{2}\,\mathbf{b}$. $\overrightarrow{BA} = -\overrightarrow{AB}$, so $\frac{3}{2}\,\mathbf{c} - \frac{3}{2}\,\mathbf{b} = 3\,\mathbf{a} - 3\,\mathbf{c} \quad \Leftrightarrow \quad \mathbf{c} + 2\,\mathbf{c} = 2\,\mathbf{a} + \mathbf{b} \quad \Leftrightarrow \quad \mathbf{c} = \frac{2}{3}\,\mathbf{a} + \frac{1}{3}\,\mathbf{b}$.

45. (a), (b)

(c) From the sketch, we estimate that $s \approx 1.3$ and $t \approx 1.6$.

(d) $\mathbf{c} = s\,\mathbf{a} + t\,\mathbf{b} \quad \Leftrightarrow \quad 7 = 3s + 2t$ and $1 = 2s - t$.
Solving these equations gives $s = \frac{9}{7}$ and $t = \frac{11}{7}$.

46. Draw $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$ emanating from the origin. Extend $\mathbf{a}$ and $\mathbf{b}$ to form lines A and B, and draw lines A' and B' parallel to these two lines through the terminal point of $\mathbf{c}$. Since $\mathbf{a}$ and $\mathbf{b}$ are not parallel, A and B' must meet (at P), and A' and B must also meet (at Q). Now we see that $\overrightarrow{OP} + \overrightarrow{OQ} = \mathbf{c}$, so if

$s = \dfrac{\left|\overrightarrow{OP}\right|}{|\mathbf{a}|}$ $\left(\text{or its negative, if } \mathbf{a} \text{ points in the direction opposite } \overrightarrow{OP}\right)$ and $t = \dfrac{\left|\overrightarrow{OQ}\right|}{|\mathbf{b}|}$ (or its negative, as in the diagram), then $\mathbf{c} = s\mathbf{a} + t\mathbf{b}$, as required.

Argument using components: Since $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$ all lie in the same plane, we can consider them to be vectors in two dimensions. Let $\mathbf{a} = \langle a_1, a_2 \rangle$, $\mathbf{b} = \langle b_1, b_2 \rangle$, and $\mathbf{c} = \langle c_1, c_2 \rangle$. We need $sa_1 + tb_1 = c_1$ and $sa_2 + tb_2 = c_2$. Multiplying the first equation by a_2 and the second by a_1 and subtracting, we get $t = \dfrac{c_2 a_1 - c_1 a_2}{b_2 a_1 - b_1 a_2}$. Similarly $s = \dfrac{b_2 c_1 - b_1 c_2}{b_2 a_1 - b_1 a_2}$. Since $\mathbf{a} \neq \mathbf{0}$ and $\mathbf{b} \neq \mathbf{0}$ and $\mathbf{a}$ is not a scalar multiple of $\mathbf{b}$, the denominator is not zero.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $|\mathbf{r}-\mathbf{r}_0|$ is the distance between the points (x, y, z) and (x_0, y_0, z_0), so the set of points is a sphere with radius 1 and center (x_0, y_0, z_0).

Alternate method: $|\mathbf{r}-\mathbf{r}_0| = 1 \quad \Leftrightarrow \quad \sqrt{(x-x_0)^2+(y-y_0)^2+(z-z_0)^2} = 1 \quad \Leftrightarrow$

$(x-x_0)^2+(y-y_0)^2+(z-z_0)^2 = 1$, which is the equation of a sphere with radius 1 and center (x_0, y_0, z_0).

48. Let P_1 and P_2 be the points with position vectors $\mathbf{r}_1$ and $\mathbf{r}_2$ respectively. Then $|\mathbf{r}-\mathbf{r}_1|+|\mathbf{r}-\mathbf{r}_2|$ is the sum of the distances from (x, y) to P_1 and P_2. Since this sum is constant, the set of points (x, y) represents an ellipse with foci P_1 and P_2. The condition $k > |\mathbf{r}_1-\mathbf{r}_2|$ assures us that the ellipse is not degenerate.

49.
$$\begin{aligned}\mathbf{a}+(\mathbf{b}+\mathbf{c}) &= \langle a_1, a_2\rangle + (\langle b_1, b_2\rangle + \langle c_1, c_2\rangle) = \langle a_1, a_2\rangle + \langle b_1+c_1, b_2+c_2\rangle \\ &= \langle a_1+b_1+c_1, a_2+b_2+c_2\rangle = \langle (a_1+b_1)+c_1, (a_2+b_2)+c_2\rangle \\ &= \langle a_1+b_1, a_2+b_2\rangle + \langle c_1, c_2\rangle = (\langle a_1, a_2\rangle + \langle b_1, b_2\rangle) + \langle c_1, c_2\rangle \\ &= (\mathbf{a}+\mathbf{b})+\mathbf{c}\end{aligned}$$

50. *Algebraically:*
$$\begin{aligned}c(\mathbf{a}+\mathbf{b}) &= c\left(\langle a_1, a_2, a_3\rangle + \langle b_1, b_2, b_3\rangle\right) = c\,\langle a_1+b_1, a_2+b_2, a_3+b_3\rangle \\ &= \langle c\,(a_1+b_1), c\,(a_2+b_2), c\,(a_3+b_3)\rangle = \langle ca_1+cb_1, ca_2+cb_2, ca_3+cb_3\rangle \\ &= \langle ca_1, ca_2, ca_3\rangle + \langle cb_1, cb_2, cb_3\rangle = c\,\mathbf{a}+c\,\mathbf{b}\end{aligned}$$

Geometrically:

According to the Triangle Law, if $\mathbf{a} = \overrightarrow{PQ}$ and $\mathbf{b} = \overrightarrow{QR}$, then $\mathbf{a}+\mathbf{b} = \overrightarrow{PR}$. Construct triangle PST as shown so that $\overrightarrow{PS} = c\,\mathbf{a}$ and $\overrightarrow{ST} = c\,\mathbf{b}$. (We have drawn the case where $c > 1$.) By the Triangle Law, $\overrightarrow{PT} = c\,\mathbf{a}+c\,\mathbf{b}$. But triangle PQR and triangle PST are similar triangles because $c\,\mathbf{b}$ is parallel to $\mathbf{b}$. Therefore, $\overrightarrow{PR}$ and $\overrightarrow{PT}$ are parallel and, in fact, $\overrightarrow{PT} = c\overrightarrow{PR}$. Thus, $c\,\mathbf{a}+c\,\mathbf{b} = c(\mathbf{a}+\mathbf{b})$.

51. Consider triangle ABC, where D and E are the midpoints of AB and BC. We know that $\overrightarrow{AB}+\overrightarrow{BC} = \overrightarrow{AC}$ **(1)** and $\overrightarrow{DB}+\overrightarrow{BE} = \overrightarrow{DE}$ **(2)**. However, $\overrightarrow{DB} = \frac{1}{2}\overrightarrow{AB}$, and $\overrightarrow{BE} = \frac{1}{2}\overrightarrow{BC}$. Substituting these expressions for $\overrightarrow{DB}$ and $\overrightarrow{BE}$ into **(2)** gives $\frac{1}{2}\overrightarrow{AB}+\frac{1}{2}\overrightarrow{BC} = \overrightarrow{DE}$. Comparing this with **(1)** gives $\overrightarrow{DE} = \frac{1}{2}\overrightarrow{AC}$. Therefore $\overrightarrow{AC}$ and $\overrightarrow{DE}$ are parallel and $\left|\overrightarrow{DE}\right| = \frac{1}{2}\left|\overrightarrow{AC}\right|$.

52. The question states that the light ray strikes all three mirrors, so it is not parallel to any of them and $a_1 \neq 0$, $a_2 \neq 0$ and $a_3 \neq 0$. Let $\mathbf{b} = \langle b_1, b_2, b_3\rangle$, as in the diagram. We can let $|\mathbf{b}| = |\mathbf{a}|$, since only its direction is important. Then

$$\frac{|b_2|}{|\mathbf{b}|} = \sin\theta = \frac{|a_2|}{|\mathbf{a}|} \quad \Rightarrow \quad |b_2| = |a_2|.$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

From the diagram $b_2\,\mathbf{j}$ and $a_2\,\mathbf{j}$ point in opposite directions, so $b_2 = -a_2$. $|AB| = |BC|$, so $|b_3| = \sin\phi\ |BC| = \sin\phi\ |AB| = |a_3|$, and $|b_1| = \cos\phi\ |BC| = \cos\phi\ |AB| = |a_1|$. $b_3\,\mathbf{k}$ and $a_3\,\mathbf{k}$ have the same direction, as do $b_1\,\mathbf{i}$ and $a_1\,\mathbf{i}$, so $\mathbf{b} = \langle a_1, -a_2, a_3\rangle$. When the ray hits the other mirrors, similar arguments show that these reflections will reverse the signs of the other two coordinates, so the final reflected ray will be $\langle -a_1, -a_2, -a_3\rangle = -\mathbf{a}$, which is parallel to $\mathbf{a}$.

DISCOVERY PROJECT The Shape of a Hanging Chain

1. As $s(x)$ is the length of the chain with uniform density ρ, the mass of the chain is given by $\rho s(x)$. Then the downward gravitational force is given by $\mathbf{w} = \langle 0, -g\rho s(x)\rangle$. Also, $\mathbf{T}_0 = \langle |\mathbf{T}_0|\cos 180°, |\mathbf{T}_0|\sin 180°\rangle = \langle -|\mathbf{T}_0|, 0\rangle$. As the system is in equilibrium, we have

$$\begin{aligned}\mathbf{T}_0 + \mathbf{T} + \mathbf{w} &= \mathbf{0}\\ \mathbf{T} &= -\mathbf{T}_0 - \mathbf{w}\\ &= -\langle -|\mathbf{T}_0|, 0\rangle - \langle 0, -g\rho s(x)\rangle\\ &= \langle |\mathbf{T}_0|, g\rho s(x)\rangle\end{aligned}$$

2. Note that the vector $\mathbf{T}$ is parallel to the tangent line to the curve at the point (x, y). Thus, the slope of the tangent line can be written as

$$\frac{dy}{dx} = \frac{g\rho s(x)}{|\mathbf{T}_0|} = \frac{s(x)}{|\mathbf{T}_0|/(g\rho)} = \frac{s(x)}{a} \qquad \text{where} \quad a = \frac{|\mathbf{T}_0|}{g\rho}$$

3. By Equation 8.1.6, $s'(x) = \displaystyle\int_0^x \sqrt{1 + \left(\frac{dy}{dt}\right)^2}$, so differentiating both sides of the equation from Problem 2 gives $\dfrac{d^2y}{dx^2} = \dfrac{1}{a}\sqrt{1 + \left(\dfrac{dy}{dx}\right)^2}$. Making the substitution $z = \dfrac{dy}{dx}$, we have $\dfrac{dz}{dx} = \dfrac{1}{a}\sqrt{1+z^2} \quad \Rightarrow \quad \dfrac{dz}{\sqrt{1+z^2}} = \dfrac{dx}{a}$.

From Table 3.11.6 we know that an antiderivative of $1/\sqrt{1+x^2}$ is $\sinh^{-1} x$, so integrating both sides of the preceding equation gives $\sinh^{-1} z = \dfrac{x}{a} + C$. We are given that $y'(0) = 0 \quad \Rightarrow \quad z(0) = 0 \quad \Rightarrow \quad C = 0$, so $\sinh^{-1} z = \dfrac{x}{a} \quad \Rightarrow$ $z = \sinh\dfrac{x}{a}$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

As $z = \dfrac{dy}{dx}$, $\dfrac{dy}{dx} = \sinh \dfrac{x}{a} \quad \Rightarrow \quad dy = \sinh \dfrac{x}{a}\, dx \quad \Rightarrow \quad \int dy = \int \sinh \dfrac{x}{a}\, dx \quad \Rightarrow \quad y = a \cosh \dfrac{x}{a} + C$. From the initial condition $y(0) = 0$, we have $0 = a \cosh 0 + C \quad \Rightarrow \quad 0 = a + C \quad \Rightarrow \quad -a = C$. Therefore, the equation of the curve is $y = a \cosh \dfrac{x}{a} - a$.

4. As the value of a increases, the graph of $y = a \cosh \dfrac{x}{a} - a$ is stretched horizontally.

12.3 The Dot Product

1. (a) $\mathbf{a} \cdot \mathbf{b}$ is a scalar, and the dot product is defined only for vectors, so $(\mathbf{a} \cdot \mathbf{b}) \cdot \mathbf{c}$ has no meaning.

(b) $(\mathbf{a} \cdot \mathbf{b})\, \mathbf{c}$ is a scalar multiple of a vector, so it does have meaning.

(c) Both $|\mathbf{a}|$ and $\mathbf{b} \cdot \mathbf{c}$ are scalars, so $|\mathbf{a}|\, (\mathbf{b} \cdot \mathbf{c})$ is an ordinary product of real numbers, and has meaning.

(d) Both $\mathbf{a}$ and $\mathbf{b} + \mathbf{c}$ are vectors, so the dot product $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c})$ has meaning.

(e) $\mathbf{a} \cdot \mathbf{b}$ is a scalar, but $\mathbf{c}$ is a vector, and so the two quantities cannot be added and $\mathbf{a} \cdot \mathbf{b} + \mathbf{c}$ has no meaning.

(f) $|\mathbf{a}|$ is a scalar, and the dot product is defined only for vectors, so $|\mathbf{a}| \cdot (\mathbf{b} + \mathbf{c})$ has no meaning.

2. $\mathbf{a} \cdot \mathbf{b} = \langle 5, -2 \rangle \cdot \langle 3, 4 \rangle = (5)(3) + (-2)(4) = 15 - 8 = 7$

3. $\mathbf{a} \cdot \mathbf{b} = \langle 1.5, 0.4 \rangle \cdot \langle -4, 6 \rangle = (1.5)(-4) + (0.4)(6) = -6 + 2.4 = -3.6$

4. $\mathbf{a} \cdot \mathbf{b} = \langle 6, -2, 3 \rangle \cdot \langle 2, 5, -1 \rangle = (6)(2) + (-2)\,(5) + (3)(-1) = 12 - 10 - 3 = -1$

5. $\mathbf{a} \cdot \mathbf{b} = \left\langle 4, 1, \frac{1}{4} \right\rangle \cdot \langle 6, -3, -8 \rangle = (4)(6) + (1)(-3) + \left(\frac{1}{4}\right)(-8) = 19$

6. $\mathbf{a} \cdot \mathbf{b} = \langle p, -p, 2p \rangle \cdot \langle 2q, q, -q \rangle = (p)(2q) + (-p)(q) + (2p)(-q) = 2pq - pq - 2pq = -pq$

7. $\mathbf{a} \cdot \mathbf{b} = (2\,\mathbf{i} + \mathbf{j}) \cdot (\mathbf{i} - \mathbf{j} + \mathbf{k}) = (2)(1) + (1)(-1) + (0)(1) = 1$

8. $\mathbf{a} \cdot \mathbf{b} = (3\,\mathbf{i} + 2\,\mathbf{j} - \mathbf{k}) \cdot (4\,\mathbf{i} + 5\,\mathbf{k}) = (3)(4) + (2)(0) + (-1)(5) = 7$

9. By Theorem 3, $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}|\, |\mathbf{b}| \cos\theta = (7)(4)\cos 30^\circ = 28\left(\frac{\sqrt{3}}{2}\right) = 14\sqrt{3}$.

10. By Theorem 3, $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}|\, |\mathbf{b}| \cos\theta = (80)(50)\cos\frac{3\pi}{4} = 4000\left(-\frac{\sqrt{2}}{2}\right) = -2000\sqrt{2}$.

11. $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ are all unit vectors, so the triangle is an equilateral triangle. Thus the angle between $\mathbf{u}$ and $\mathbf{v}$ is 60° and $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}|\, |\mathbf{v}| \cos 60^\circ = (1)(1)\left(\frac{1}{2}\right) = \frac{1}{2}$. If $\mathbf{w}$ is moved so it has the same initial point as $\mathbf{u}$, we can see that the angle between them is 120° and we have $\mathbf{u} \cdot \mathbf{w} = |\mathbf{u}|\, |\mathbf{w}| \cos 120^\circ = (1)(1)\left(-\frac{1}{2}\right) = -\frac{1}{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $\mathbf{u}$ is a unit vector, so $\mathbf{w}$ is also a unit vector, and $|\mathbf{v}|$ can be determined by examining the right triangle formed by $\mathbf{u}$ and $\mathbf{v}$. Since the angle between $\mathbf{u}$ and $\mathbf{v}$ is $45°$, we have $|\mathbf{v}| = |\mathbf{u}| \cos 45° = \frac{\sqrt{2}}{2}$. Then $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}|\,|\mathbf{v}| \cos 45° = (1)\left(\frac{\sqrt{2}}{2}\right)\frac{\sqrt{2}}{2} = \frac{1}{2}$. Since $\mathbf{u}$ and $\mathbf{w}$ are orthogonal, $\mathbf{u} \cdot \mathbf{w} = 0$.

13. (a) $\mathbf{i} \cdot \mathbf{j} = \langle 1, 0, 0\rangle \cdot \langle 0, 1, 0\rangle = (1)(0) + (0)(1) + (0)(0) = 0$. Similarly, $\mathbf{j} \cdot \mathbf{k} = (0)(0) + (1)(0) + (0)(1) = 0$ and $\mathbf{k} \cdot \mathbf{i} = (0)(1) + (0)(0) + (1)(0) = 0$.

Another method: Because $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ are mutually perpendicular, the cosine factor in each dot product (see Theorem 3) is $\cos \frac{\pi}{2} = 0$.

(b) By Property 1 of the dot product, $\mathbf{i} \cdot \mathbf{i} = |\mathbf{i}|^2 = 1^2 = 1$ since $\mathbf{i}$ is a unit vector. Similarly, $\mathbf{j} \cdot \mathbf{j} = |\mathbf{j}|^2 = 1$ and $\mathbf{k} \cdot \mathbf{k} = |\mathbf{k}|^2 = 1$.

14. The dot product $\mathbf{A} \cdot \mathbf{P}$ is

$$\begin{aligned}\langle a, b, c\rangle \cdot \langle 4, 2.5, 1\rangle &= a(4) + b(2.5) + c(1)\\ &= (\text{number of hamburgers sold})(\text{price per hamburger})\\ &\quad + (\text{number of hot dogs sold})(\text{price per hot dog})\\ &\quad\quad + (\text{number of bottles sold})(\text{price per bottle})\end{aligned}$$

so it is equal to the vendor's total revenue for that day.

15. $\mathbf{u} = \langle 5, 1\rangle$, $\mathbf{v} = \langle 3, 2\rangle \;\Rightarrow\; |\mathbf{u}| = \sqrt{5^2 + 1^2} = \sqrt{26}$, $|\mathbf{v}| = \sqrt{3^2 + 2^2} = \sqrt{13}$, and $\mathbf{u} \cdot \mathbf{v} = 5(3) + 1(2) = 17$. From Corollary 6, we have $\cos\theta = \dfrac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}||\mathbf{v}|} = \dfrac{17}{\sqrt{26}\sqrt{13}} = \dfrac{17}{13\sqrt{2}}$ and the angle between $\mathbf{u}$ and $\mathbf{v}$ is $\theta = \cos^{-1}\left(\dfrac{17}{13\sqrt{2}}\right) \approx 22°$.

16. $\mathbf{a} = \mathbf{i} - 3\,\mathbf{j}$, $\mathbf{b} = -3\,\mathbf{i} + 4\,\mathbf{j} \;\Rightarrow\; |\mathbf{a}| = \sqrt{1^2 + (-3)^2} = \sqrt{10}$, $|\mathbf{b}| = \sqrt{(-3)^2 + 4^2} = 5$, and $\mathbf{a} \cdot \mathbf{b} = 1(-3) + (-3)(4) = -15$. From Corollary 6, we have $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} = \dfrac{-15}{5\sqrt{10}} = \dfrac{-3}{\sqrt{10}}$ and the angle between $\mathbf{a}$ and $\mathbf{b}$ is $\theta = \cos^{-1}\left(\dfrac{-3}{\sqrt{10}}\right) \approx 162°$.

17. $\mathbf{a} = \langle 1, -4, 1\rangle$, $\mathbf{b} = \langle 0, 2, -2\rangle \;\Rightarrow\; |\mathbf{a}| = \sqrt{1^2 + (-4)^2 + 1^2} = \sqrt{18} = 3\sqrt{2}$, $|\mathbf{b}| = \sqrt{0^2 + 2^2 + (-2)^2} = \sqrt{8} = 2\sqrt{2}$, and $\mathbf{a} \cdot \mathbf{b} = (1)(0) + (-4)(2) + (1)(-2) = -10$. From Corollary 6, we have $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{-10}{3\sqrt{2} \cdot 2\sqrt{2}} = -\dfrac{10}{12} = -\dfrac{5}{6}$ and the angle between $\mathbf{a}$ and $\mathbf{b}$ is $\theta = \cos^{-1}\left(-\frac{5}{6}\right) \approx 146°$.

18. $\mathbf{a} = \langle -1, 3, 4\rangle$, $\mathbf{b} = \langle 5, 2, 1\rangle \;\Rightarrow\; |\mathbf{a}| = \sqrt{(-1)^2 + 3^2 + 4^2} = \sqrt{26}$, $|\mathbf{b}| = \sqrt{5^2 + 2^2 + 1^2} = \sqrt{30}$, and $\mathbf{a} \cdot \mathbf{b} = (-1)(5) + (3)(2) + (4)(1) = 5$. From Corollary 6, we have $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{5}{\sqrt{26} \cdot \sqrt{30}} = \dfrac{5}{\sqrt{780}} = \dfrac{5}{2\sqrt{195}}$ and the angle between $\mathbf{a}$ and $\mathbf{b}$ is $\theta = \cos^{-1}\left(\dfrac{5}{2\sqrt{195}}\right) \approx 80°$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $\mathbf{u} = \mathbf{i} - 4\,\mathbf{j} + \mathbf{k}$, $\mathbf{v} = -3\,\mathbf{i} + \mathbf{j} + 5\,\mathbf{k} \quad \Rightarrow \quad |\mathbf{u}| = \sqrt{1^2 + (-4)^2 + 1^2} = \sqrt{18} = 3\sqrt{2}$, $|\mathbf{v}| = \sqrt{(-3)^2 + 1^2 + 5^2} = \sqrt{35}$, and $\mathbf{u} \cdot \mathbf{v} = 1(-3) + (-4)(1) + 1(5) = -2$. From Corollary 6, we have $\cos\theta = \dfrac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}||\mathbf{v}|} = \dfrac{-2}{3\sqrt{2}\sqrt{35}} = \dfrac{-2}{3\sqrt{70}}$ and the angle between $\mathbf{u}$ and $\mathbf{v}$ is $\theta = \cos^{-1}\left(\dfrac{-2}{3\sqrt{70}}\right) \approx 95^\circ$.

20. $\mathbf{a} = 8\,\mathbf{i} - \mathbf{j} + 4\,\mathbf{k}$, $\mathbf{b} = 4\,\mathbf{j} + 2\,\mathbf{k} \quad \Rightarrow \quad |\mathbf{a}| = \sqrt{8^2 + (-1)^2 + 4^2} = \sqrt{81} = 9$, $|\mathbf{b}| = \sqrt{0^2 + 4^2 + 2^2} = \sqrt{20} = 2\sqrt{5}$, and $\mathbf{a} \cdot \mathbf{b} = (8)(0) + (-1)(4) + (4)(2) = 4$. From Corollary 6, we have $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{4}{9 \cdot 2\sqrt{5}} = \dfrac{2}{9\sqrt{5}}$ and the angle between $\mathbf{a}$ and $\mathbf{b}$ is $\theta = \cos^{-1}\left(\dfrac{2}{9\sqrt{5}}\right) \approx 84^\circ$.

21. Let p, q, and r be the angles at vertices P, Q, and R respectively. Then p is the angle between vectors $\overrightarrow{PQ}$ and $\overrightarrow{PR}$, q is the angle between vectors $\overrightarrow{QP}$ and $\overrightarrow{QR}$, and r is the angle between vectors $\overrightarrow{RP}$ and $\overrightarrow{RQ}$.

Thus $\cos p = \dfrac{\overrightarrow{PQ} \cdot \overrightarrow{PR}}{\left|\overrightarrow{PQ}\right| \left|\overrightarrow{PR}\right|} = \dfrac{\langle -2, 3\rangle \cdot \langle 1, 4\rangle}{\sqrt{(-2)^2 + 3^2}\,\sqrt{1^2 + 4^2}} = \dfrac{-2 + 12}{\sqrt{13}\,\sqrt{17}} = \dfrac{10}{\sqrt{221}}$ and $p = \cos^{-1}\left(\dfrac{10}{\sqrt{221}}\right) \approx 48^\circ$. Similarly,

$\cos q = \dfrac{\overrightarrow{QP} \cdot \overrightarrow{QR}}{\left|\overrightarrow{QP}\right| \left|\overrightarrow{QR}\right|} = \dfrac{\langle 2, -3\rangle \cdot \langle 3, 1\rangle}{\sqrt{4 + 9}\,\sqrt{9 + 1}} = \dfrac{6 - 3}{\sqrt{13}\,\sqrt{10}} = \dfrac{3}{\sqrt{130}}$ so $q = \cos^{-1}\left(\dfrac{3}{\sqrt{130}}\right) \approx 75^\circ$ and

$r \approx 180^\circ - (48^\circ + 75^\circ) = 57^\circ$.

Alternate solution: Apply the Law of Cosines three times as follows:

$$\cos p = \frac{\left|\overrightarrow{QR}\right|^2 - \left|\overrightarrow{PQ}\right|^2 - \left|\overrightarrow{PR}\right|^2}{2\left|\overrightarrow{PQ}\right| \left|\overrightarrow{PR}\right|} \qquad \cos q = \frac{\left|\overrightarrow{PR}\right|^2 - \left|\overrightarrow{PQ}\right|^2 - \left|\overrightarrow{QR}\right|^2}{2\left|\overrightarrow{PQ}\right| \left|\overrightarrow{QR}\right|} \qquad \cos r = \frac{\left|\overrightarrow{PQ}\right|^2 - \left|\overrightarrow{PR}\right|^2 - \left|\overrightarrow{QR}\right|^2}{2\left|\overrightarrow{PR}\right| \left|\overrightarrow{QR}\right|}$$

22. Let a, b, and c be the angles at vertices A, B, and C. Then a is the angle between vectors $\overrightarrow{AB}$ and $\overrightarrow{AC}$, b is the angle between vectors $\overrightarrow{BA}$ and $\overrightarrow{BC}$, and c is the angle between vectors $\overrightarrow{CA}$ and $\overrightarrow{CB}$.

Thus $\cos a = \dfrac{\overrightarrow{AB} \cdot \overrightarrow{AC}}{\left|\overrightarrow{AB}\right| \left|\overrightarrow{AC}\right|} = \dfrac{\langle 2, -2, 1\rangle \cdot \langle 0, 3, 4\rangle}{\sqrt{2^2 + (-2)^2 + 1^2}\,\sqrt{0^2 + 3^2 + 4^2}} = \dfrac{0 - 6 + 4}{3 \cdot 5} = -\dfrac{2}{15}$ and $a = \cos^{-1}\left(-\frac{2}{15}\right) \approx 98^\circ$.

Similarly, $\cos b = \dfrac{\overrightarrow{BA} \cdot \overrightarrow{BC}}{\left|\overrightarrow{BA}\right| \left|\overrightarrow{BC}\right|} = \dfrac{\langle -2, 2, -1\rangle \cdot \langle -2, 5, 3\rangle}{\sqrt{4 + 4 + 1}\,\sqrt{4 + 25 + 9}} = \dfrac{4 + 10 - 3}{3 \cdot \sqrt{38}} = \dfrac{11}{3\sqrt{38}}$ so $b = \cos^{-1}\left(\dfrac{11}{3\sqrt{38}}\right) \approx 54^\circ$ and

$c \approx 180^\circ - (98^\circ + 54^\circ) = 28^\circ$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Alternate solution: Apply the Law of Cosines three times as follows:

$$\cos a = \frac{\left|\overrightarrow{BC}\right|^2 - \left|\overrightarrow{AB}\right|^2 - \left|\overrightarrow{AC}\right|^2}{2\left|\overrightarrow{AB}\right|\left|\overrightarrow{AC}\right|} \qquad \cos b = \frac{\left|\overrightarrow{AC}\right|^2 - \left|\overrightarrow{AB}\right|^2 - \left|\overrightarrow{BC}\right|^2}{2\left|\overrightarrow{AB}\right|\left|\overrightarrow{BC}\right|} \qquad \cos c = \frac{\left|\overrightarrow{AB}\right|^2 - \left|\overrightarrow{AC}\right|^2 - \left|\overrightarrow{BC}\right|^2}{2\left|\overrightarrow{AC}\right|\left|\overrightarrow{BC}\right|}$$

23. (a) $\mathbf{a}\cdot\mathbf{b} = (9)(-2) + (3)(6) = 0$, so $\mathbf{a}$ and $\mathbf{b}$ are orthogonal (and not parallel).

(b) $\mathbf{a}\cdot\mathbf{b} = (4)(3) + (5)(-1) + (-2)(5) = -3 \neq 0$, so $\mathbf{a}$ and $\mathbf{b}$ are not orthogonal. Also, since $\mathbf{a}$ is not a scalar multiple of $\mathbf{b}$, $\mathbf{a}$ and $\mathbf{b}$ are not parallel.

(c) $\mathbf{a}\cdot\mathbf{b} = (-8)(6) + (12)(-9) + (4)(-3) = -168 \neq 0$, so $\mathbf{a}$ and $\mathbf{b}$ are not orthogonal. Because $\mathbf{a} = -\frac{4}{3}\,\mathbf{b}$, $\mathbf{a}$ and $\mathbf{b}$ are parallel.

(d) $\mathbf{a}\cdot\mathbf{b} = (3)(5) + (-1)(9) + (3)(-2) = 0$, so $\mathbf{a}$ and $\mathbf{b}$ are orthogonal (and not parallel).

24. (a) $\mathbf{u}\cdot\mathbf{v} = (-5)(3) + (4)(4) + (-2)(-1) = 3 \neq 0$, so $\mathbf{u}$ and $\mathbf{v}$ are not orthogonal. Also, $\mathbf{u}$ is not a scalar multiple of $\mathbf{v}$, so $\mathbf{u}$ and $\mathbf{v}$ are not parallel.

(b) $\mathbf{u}\cdot\mathbf{v} = (9)(-6) + (-6)(4) + (3)(-2) = -84 \neq 0$, so $\mathbf{u}$ and $\mathbf{v}$ are not orthogonal. Because $\mathbf{u} = -\frac{3}{2}\,\mathbf{v}$, $\mathbf{u}$ and $\mathbf{v}$ are parallel.

(c) $\mathbf{u}\cdot\mathbf{v} = (c)(c) + (c)(0) + (c)(-c) = c^2 + 0 - c^2 = 0$, so $\mathbf{u}$ and $\mathbf{v}$ are orthogonal (and not parallel). (Note that if $c = 0$ then $\mathbf{u} = \mathbf{v} = \mathbf{0}$, and the zero vector is considered orthogonal to all vectors. Although in this case $\mathbf{u}$ and $\mathbf{v}$ are identical, they are not considered parallel, as only nonzero vectors can be parallel.)

25. $\overrightarrow{QP} = \langle -1, -3, 2\rangle$, $\overrightarrow{QR} = \langle 4, -2, -1\rangle$, and $\overrightarrow{QP}\cdot\overrightarrow{QR} = -4 + 6 - 2 = 0$. Thus $\overrightarrow{QP}$ and $\overrightarrow{QR}$ are orthogonal, so the angle of the triangle at vertex Q is a right angle.

26. By Theorem 3, vectors $\langle 2, 1, -1\rangle$ and $\langle 1, x, 0\rangle$ meet at an angle of 45° when

$\langle 2, 1, -1\rangle \cdot \langle 1, x, 0\rangle = \sqrt{4+1+1}\,\sqrt{1+x^2+0}\,\cos 45^\circ$ or $2 + x - 0 = \sqrt{6}\,\sqrt{1+x^2}\cdot\frac{\sqrt{2}}{2} \quad\Leftrightarrow\quad 2 + x = \sqrt{3}\sqrt{1+x^2}$.

Squaring both sides gives $4 + 4x + x^2 = 3 + 3x^2 \quad\Leftrightarrow\quad 2x^2 - 4x - 1 = 0$. By the quadratic formula,

$x = \dfrac{-(-4) \pm \sqrt{(-4)^2 - 4(2)(-1)}}{2(2)} = \dfrac{4 \pm \sqrt{24}}{4} = \dfrac{4 \pm 2\sqrt{6}}{4} = 1 \pm \dfrac{\sqrt{6}}{2}$. (You can verify that both values are valid.)

27. Let $\mathbf{a} = a_1\,\mathbf{i} + a_2\,\mathbf{j} + a_3\,\mathbf{k}$ be a vector orthogonal to both $\mathbf{i}+\mathbf{j}$ and $\mathbf{i}+\mathbf{k}$. Then $\mathbf{a}\cdot(\mathbf{i}+\mathbf{j}) = 0 \quad\Leftrightarrow\quad a_1 + a_2 = 0$ and $\mathbf{a}\cdot(\mathbf{i}+\mathbf{k}) = 0 \quad\Leftrightarrow\quad a_1 + a_3 = 0$, so $a_1 = -a_2 = -a_3$. Furthermore $\mathbf{a}$ is to be a unit vector, so $1 = a_1^2 + a_2^2 + a_3^2 = 3a_1^2$ implies $a_1 = \pm\frac{1}{\sqrt{3}}$. Thus $\mathbf{a} = \frac{1}{\sqrt{3}}\,\mathbf{i} - \frac{1}{\sqrt{3}}\,\mathbf{j} - \frac{1}{\sqrt{3}}\,\mathbf{k}$ and $\mathbf{a} = -\frac{1}{\sqrt{3}}\,\mathbf{i} + \frac{1}{\sqrt{3}}\,\mathbf{j} + \frac{1}{\sqrt{3}}\,\mathbf{k}$ are two such unit vectors.

28. Let $\mathbf{u} = \langle a, b\rangle$ be a unit vector. By Theorem 3 we need $\mathbf{u}\cdot\mathbf{v} = |\mathbf{u}|\,|\mathbf{v}|\cos 60^\circ \quad\Leftrightarrow\quad 3a + 4b = (1)(5)\frac{1}{2} \quad\Leftrightarrow$

$b = \frac{5}{8} - \frac{3}{4}a$. Since $\mathbf{u}$ is a unit vector, $|\mathbf{u}| = \sqrt{a^2+b^2} = 1 \quad\Leftrightarrow\quad a^2 + b^2 = 1 \quad\Leftrightarrow\quad a^2 + \left(\frac{5}{8} - \frac{3}{4}a\right)^2 = 1 \quad\Leftrightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\frac{25}{16}a^2 - \frac{15}{16}a + \frac{25}{64} = 1 \quad \Leftrightarrow \quad 100a^2 - 60a - 39 = 0$. By the quadratic formula,

$a = \dfrac{-(-60) \pm \sqrt{(-60)^2 - 4(100)(-39)}}{2(100)} = \dfrac{60 \pm \sqrt{19{,}200}}{200} = \dfrac{3 \pm 4\sqrt{3}}{10}$. If $a = \dfrac{3 + 4\sqrt{3}}{10}$ then

$b = \dfrac{5}{8} - \dfrac{3}{4}\left(\dfrac{3 + 4\sqrt{3}}{10}\right) = \dfrac{4 - 3\sqrt{3}}{10}$, and if $a = \dfrac{3 - 4\sqrt{3}}{10}$ then $b = \dfrac{5}{8} - \dfrac{3}{4}\left(\dfrac{3 - 4\sqrt{3}}{10}\right) = \dfrac{4 + 3\sqrt{3}}{10}$. Thus the two

unit vectors are $\left\langle \dfrac{3 + 4\sqrt{3}}{10}, \dfrac{4 - 3\sqrt{3}}{10} \right\rangle \approx \langle 0.9928, -0.1196 \rangle$ and $\left\langle \dfrac{3 - 4\sqrt{3}}{10}, \dfrac{4 + 3\sqrt{3}}{10} \right\rangle \approx \langle -0.3928, 0.9196 \rangle$.

29. The line $y = 4 - 3x \quad \Leftrightarrow \quad y = -3x + 4$ has slope -3, so a vector parallel to the line is $\mathbf{a} = \langle 1, -3 \rangle$. The line $y = 3x + 2$ has slope 3, so a vector parallel to the line is $\mathbf{b} = \langle 1, 3 \rangle$. The angle between the lines is the same as the angle θ between the vectors. Here we have $\mathbf{a} \cdot \mathbf{b} = 1(1) + 3(-3) = -8$, $|\mathbf{a}| = \sqrt{1^2 + (-3)^2} = \sqrt{10}$, and $|\mathbf{b}| = \sqrt{1^2 + 3^2} = \sqrt{10}$. Then $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} = \dfrac{-8}{\sqrt{10}\sqrt{10}} = -\dfrac{4}{5}$ and $\theta = \cos^{-1}\left(-\dfrac{4}{5}\right) \approx 143.1^\circ$. Therefore, the acute angle between the two lines is approximately $180^\circ - 143.1^\circ = 36.9^\circ$.

30. The line $5x - y = 8 \quad \Leftrightarrow \quad y = 5x - 8$ has slope 5, so a vector parallel to the line is $\mathbf{a} = \langle 1, 5 \rangle$. The line $x + 3y = 15 \quad \Leftrightarrow \quad y = -\frac{1}{3}x + 5$ has slope $-\frac{1}{3}$, so a vector parallel to the line is $\mathbf{b} = \langle 3, -1 \rangle$. The angle between the lines is the same as the angle θ between the vectors. Here we have $\mathbf{a} \cdot \mathbf{b} = 1(3) + 5(-1) = -2$, $|\mathbf{a}| = \sqrt{1^2 + 5^2} = \sqrt{26}$, and $|\mathbf{b}| = \sqrt{3^2 + (-1)^2} = \sqrt{10}$. Then $\cos\theta = \dfrac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} = \dfrac{-2}{\sqrt{26}\sqrt{10}} = -\dfrac{1}{\sqrt{65}}$ and $\theta = \cos^{-1}\left(-\dfrac{1}{\sqrt{65}}\right) \approx 97.1^\circ$. Therefore, the acute angle between the two lines is approximately $180^\circ - 97.1^\circ = 82.9^\circ$.

31. The curves $y = x^2$ and $y = x^3$ meet when $x^2 = x^3 \quad \Leftrightarrow \quad x^3 - x^2 = 0 \quad \Leftrightarrow \quad x^2(x - 1) = 0 \quad \Leftrightarrow \quad x = 0, x = 1$. We have $\dfrac{d}{dx}x^2 = 2x$ and $\dfrac{d}{dx}x^3 = 3x^2$, so the tangent lines of both curves have slope 0 at $x = 0$. Thus the angle between the curves is 0° at the point $(0, 0)$. For $x = 1$, $\left.\dfrac{d}{dx}\, x^2\right|_{x=1} = 2$ and $\left.\dfrac{d}{dx}\, x^3\right|_{x=1} = 3$ so the tangent lines at the point $(1, 1)$ have slopes 2 and 3. Vectors parallel to the tangent lines are $\langle 1, 2 \rangle$ and $\langle 1, 3 \rangle$, and the angle θ between them is given by

$$\cos\theta = \frac{\langle 1, 2 \rangle \cdot \langle 1, 3 \rangle}{|\langle 1, 2 \rangle|\ |\langle 1, 3 \rangle|} = \frac{1 + 6}{\sqrt{5}\sqrt{10}} = \frac{7}{5\sqrt{2}}$$

Thus $\theta = \cos^{-1}\left(\dfrac{7}{5\sqrt{2}}\right) \approx 8.1^\circ$.

32. The curves $y = \sin x$ and $y = \cos x$ meet when $\sin x = \cos x \quad \Leftrightarrow \quad \tan x = 1 \quad \Leftrightarrow \quad x = \pi/4 \ \ [0 \le x \le \pi/2]$. Thus the point of intersection is $\left(\pi/4, \sqrt{2}/2\right)$. We have $\left.\dfrac{d}{dx}\, \sin x\right|_{x=\pi/4} = \left.\cos x\right|_{x=\pi/4} = \dfrac{\sqrt{2}}{2}$ and

$\left.\dfrac{d}{dx}\, \cos x\right|_{x=\pi/4} = -\left.\sin x\right|_{x=\pi/4} = -\dfrac{\sqrt{2}}{2}$, so the tangent lines at that point have slopes $\dfrac{\sqrt{2}}{2}$ and $-\dfrac{\sqrt{2}}{2}$. Vectors parallel to

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

the tangent lines are $\left\langle 1, \dfrac{\sqrt{2}}{2} \right\rangle$ and $\left\langle 1, -\dfrac{\sqrt{2}}{2} \right\rangle$, and the angle θ between them is given by

$$\cos\theta = \frac{\langle 1, \sqrt{2}/2\rangle \cdot \langle 1, -\sqrt{2}/2\rangle}{|\langle 1, \sqrt{2}/2\rangle|\ |\langle 1, -\sqrt{2}/2\rangle|} = \frac{1 - \frac{1}{2}}{\sqrt{\frac{3}{2}}\sqrt{\frac{3}{2}}} = \frac{1/2}{3/2} = \frac{1}{3}$$

Thus $\theta = \cos^{-1}\left(\frac{1}{3}\right) \approx 70.5°$.

33. $|\langle 4, 1, 8\rangle| = \sqrt{4^2 + 1^2 + 8^2} = \sqrt{81} = 9$. Using Equations 8 and 9, we have $\cos\alpha = \frac{4}{9}$, $\cos\beta = \frac{1}{9}$, and $\cos\gamma = \frac{8}{9}$. The direction angles are given by $\alpha = \cos^{-1}\left(\frac{4}{9}\right) \approx 63.6°$, $\beta = \cos^{-1}\left(\frac{1}{9}\right) \approx 83.6°$, and $\gamma = \cos^{-1}\left(\frac{8}{9}\right) \approx 27.3°$.

34. $|\langle -6, 2, 9\rangle| = \sqrt{(-6)^2 + 2^2 + 9^2} = \sqrt{121} = 11$. Using Equations 8 and 9, we have $\cos\alpha = -\frac{6}{11}$, $\cos\beta = \frac{2}{11}$, and $\cos\gamma = \frac{9}{11}$. The direction angles are given by $\alpha = \cos^{-1}\left(-\frac{6}{11}\right) \approx 123.1°$, $\beta = \cos^{-1}\left(\frac{2}{11}\right) \approx 79.5°$, and $\gamma = \cos^{-1}\left(\frac{9}{11}\right) \approx 35.1°$.

35. $|3\mathbf{i} - \mathbf{j} - 2\mathbf{k}| = \sqrt{3^2 + (-1)^2 + (-2)^2} = \sqrt{14}$. Using Equations 8 and 9, we have $\cos\alpha = \dfrac{3}{\sqrt{14}}$, $\cos\beta = -\dfrac{1}{\sqrt{14}}$, and $\cos\gamma = -\dfrac{2}{\sqrt{14}}$. The direction angles are given by $\alpha = \cos^{-1}\left(\dfrac{3}{\sqrt{14}}\right) \approx 36.7°$, $\beta = \cos^{-1}\left(-\dfrac{1}{\sqrt{14}}\right) \approx 105.5°$, and $\gamma = \cos^{-1}\left(-\dfrac{2}{\sqrt{14}}\right) \approx 122.3°$.

36. $|-0.7\mathbf{i} + 1.2\mathbf{j} - 0.8\mathbf{k}| = \sqrt{(-0.7)^2 + (1.2)^2 + (-0.8)^2} = \sqrt{2.57}$. Using Equations 8 and 9, we have $\cos\alpha = -\dfrac{0.7}{\sqrt{2.57}}$, $\cos\beta = \dfrac{1.2}{\sqrt{2.57}}$, and $\cos\gamma = -\dfrac{0.8}{\sqrt{2.57}}$. The direction angles are given by $\alpha = \cos^{-1}\left(-\dfrac{0.7}{\sqrt{2.57}}\right) \approx 115.9°$, $\beta = \cos^{-1}\left(\dfrac{1.2}{\sqrt{2.57}}\right) \approx 41.5°$, and $\gamma = \cos^{-1}\left(-\dfrac{0.8}{\sqrt{2.57}}\right) \approx 119.9°$.

37. $|\langle c, c, c\rangle| = \sqrt{c^2 + c^2 + c^2} = \sqrt{3c^2} = c\sqrt{3}$. Using Equations 8 and 9, we have $\cos\alpha = \cos\beta = \cos\gamma = \dfrac{c}{c\sqrt{3}} = \dfrac{1}{\sqrt{3}}$. The direction angles are given by $\alpha = \beta = \gamma = \cos^{-1}\left(\dfrac{1}{\sqrt{3}}\right) \approx 54.7°$.

38. Since $\cos^2\alpha + \cos^2\beta + \cos^2\gamma = 1$, $\cos^2\gamma = 1 - \cos^2\alpha - \cos^2\beta = 1 - \cos^2\left(\frac{\pi}{4}\right) - \cos^2\left(\frac{\pi}{3}\right) = 1 - \left(\frac{\sqrt{2}}{2}\right)^2 - \left(\frac{1}{2}\right)^2 = \frac{1}{4}$. Thus $\cos\gamma = \pm\frac{1}{2}$ and $\gamma = \frac{\pi}{3}$ or $\gamma = \frac{2\pi}{3}$.

39. $|\mathbf{a}| = \sqrt{(-5)^2 + 12^2} = \sqrt{169} = 13$. The scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{-5\cdot 4 + 12\cdot 6}{13} = 4$ and the vector projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{proj}_{\mathbf{a}}\,\mathbf{b} = \left(\dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|}\right)\dfrac{\mathbf{a}}{|\mathbf{a}|} = 4\cdot\frac{1}{13}\langle -5, 12\rangle = \left\langle -\frac{20}{13}, \frac{48}{13}\right\rangle$.

40. $|\mathbf{a}| = \sqrt{1^2 + 4^2} = \sqrt{17}$. The scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{1\cdot 2 + 4\cdot 3}{\sqrt{17}} = \dfrac{14}{\sqrt{17}}$ and the vector projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{proj}_{\mathbf{a}}\,\mathbf{b} = \left(\dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|}\right)\dfrac{\mathbf{a}}{|\mathbf{a}|} = \frac{14}{\sqrt{17}}\cdot\frac{1}{\sqrt{17}}\langle 1, 4\rangle = \left\langle \frac{14}{17}, \frac{56}{17}\right\rangle$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. $|\mathbf{a}| = \sqrt{4^2 + 7^2 + (-4)^2} = \sqrt{81} = 9$ so the scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is

$\text{comp}_{\mathbf{a}}\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{(4)(3)+(7)(-1)+(-4)(1)}{9} = \dfrac{1}{9}$. The vector projection of $\mathbf{b}$ onto $\mathbf{a}$ is

$\text{proj}_{\mathbf{a}}\mathbf{b} = \left(\dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|}\right)\dfrac{\mathbf{a}}{|\mathbf{a}|} = \frac{1}{9}\cdot\frac{1}{9}\langle 4,7,-4\rangle = \frac{1}{81}\langle 4,7,-4\rangle = \left\langle \frac{4}{81}, \frac{7}{81}, -\frac{4}{81}\right\rangle$.

42. $|\mathbf{a}| = \sqrt{1+16+64} = \sqrt{81} = 9$ so the scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \frac{1}{9}(-12+4+16) = \frac{8}{9}$, while

the vector projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{proj}_{\mathbf{a}}\,\mathbf{b} = \left(\dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|}\right)\dfrac{\mathbf{a}}{|\mathbf{a}|} = \frac{8}{9}\cdot\frac{1}{9}\langle -1,4,8\rangle = \frac{8}{81}\langle -1,4,8\rangle = \left\langle -\frac{8}{81}, \frac{32}{81}, \frac{64}{81}\right\rangle$.

43. $|\mathbf{a}| = \sqrt{9+9+1} = \sqrt{19}$ so the scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{6-12-1}{\sqrt{19}} = -\dfrac{7}{\sqrt{19}}$ while the vector

projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{proj}_{\mathbf{a}}\,\mathbf{b} = -\dfrac{7}{\sqrt{19}}\dfrac{\mathbf{a}}{|\mathbf{a}|} = -\frac{7}{\sqrt{19}}\cdot\frac{1}{\sqrt{19}}(3\,\mathbf{i} - 3\,\mathbf{j} + \mathbf{k}) = -\frac{7}{19}(3\,\mathbf{i} - 3\,\mathbf{j} + \mathbf{k}) = -\frac{21}{19}\,\mathbf{i} + \frac{21}{19}\,\mathbf{j} - \frac{7}{19}\,\mathbf{k}$.

44. $|\mathbf{a}| = \sqrt{1+4+9} = \sqrt{14}$ so the scalar projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{5+0-3}{\sqrt{14}} = \dfrac{2}{\sqrt{14}}$ while the vector

projection of $\mathbf{b}$ onto $\mathbf{a}$ is $\text{proj}_{\mathbf{a}}\,\mathbf{b} = \dfrac{2}{\sqrt{14}}\dfrac{\mathbf{a}}{|\mathbf{a}|} = \frac{2}{\sqrt{14}}\cdot\frac{1}{\sqrt{14}}\,(\mathbf{i} + 2\,\mathbf{j} + 3\,\mathbf{k}) = \frac{1}{7}(\mathbf{i} + 2\,\mathbf{j} + 3\,\mathbf{k}) = \frac{1}{7}\,\mathbf{i} + \frac{2}{7}\,\mathbf{j} + \frac{3}{7}\,\mathbf{k}$.

45. $(\text{orth}_{\mathbf{a}}\,\mathbf{b})\cdot\mathbf{a} = (\mathbf{b} - \text{proj}_{\mathbf{a}}\,\mathbf{b})\cdot\mathbf{a} = \mathbf{b}\cdot\mathbf{a} - (\text{proj}_{\mathbf{a}}\,\mathbf{b})\cdot\mathbf{a} = \mathbf{b}\cdot\mathbf{a} - \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|^2}\,\mathbf{a}\cdot\mathbf{a} = \mathbf{b}\cdot\mathbf{a} - \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|^2}\,|\mathbf{a}|^2 = \mathbf{b}\cdot\mathbf{a} - \mathbf{a}\cdot\mathbf{b} = 0$.

So they are orthogonal by (7).

46. Using the formula in Exercise 45 and the result of Exercise 40, we have

$\text{orth}_{\mathbf{a}}\,\mathbf{b} = \mathbf{b} - \text{proj}_{\mathbf{a}}\,\mathbf{b} = \langle 2,3\rangle - \left\langle \frac{14}{17}, \frac{56}{17}\right\rangle = \left\langle \frac{20}{17}, -\frac{5}{17}\right\rangle$.

47. $\text{comp}_{\mathbf{a}}\,\mathbf{b} = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = 2 \quad\Leftrightarrow\quad \mathbf{a}\cdot\mathbf{b} = 2\,|\mathbf{a}| = 2\sqrt{10}$. If $\mathbf{b} = \langle b_1, b_2, b_3\rangle$, then we need $3b_1 + 0b_2 - 1b_3 = 2\sqrt{10}$.

One possible solution is obtained by taking $b_1 = 0$, $b_2 = 0$, $b_3 = -2\sqrt{10}$. In general, $\mathbf{b} = \left\langle s, t, 3s - 2\sqrt{10}\right\rangle$, $s, t \in \mathbb{R}$.

48. (a) $\text{comp}_{\mathbf{a}}\,\mathbf{b} = \text{comp}_{\mathbf{b}}\,\mathbf{a} \quad\Leftrightarrow\quad \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|} = \dfrac{\mathbf{b}\cdot\mathbf{a}}{|\mathbf{b}|} \quad\Leftrightarrow\quad \dfrac{1}{|\mathbf{a}|} = \dfrac{1}{|\mathbf{b}|}$ or $\mathbf{a}\cdot\mathbf{b} = 0 \quad\Leftrightarrow\quad |\mathbf{b}| = |\mathbf{a}|$ or $\mathbf{a}\cdot\mathbf{b} = 0$.

That is, if $\mathbf{a}$ and $\mathbf{b}$ are orthogonal or if they have the same length.

(b) $\text{proj}_{\mathbf{a}}\,\mathbf{b} = \text{proj}_{\mathbf{b}}\,\mathbf{a} \quad\Leftrightarrow\quad \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|^2}\,\mathbf{a} = \dfrac{\mathbf{b}\cdot\mathbf{a}}{|\mathbf{b}|^2}\,\mathbf{b} \quad\Leftrightarrow\quad \mathbf{a}\cdot\mathbf{b} = 0$ or $\dfrac{\mathbf{a}}{|\mathbf{a}|^2} = \dfrac{\mathbf{b}}{|\mathbf{b}|^2}$.

But $\dfrac{\mathbf{a}}{|\mathbf{a}|^2} = \dfrac{\mathbf{b}}{|\mathbf{b}|^2} \quad\Rightarrow\quad \dfrac{|\mathbf{a}|}{|\mathbf{a}|^2} = \dfrac{|\mathbf{b}|}{|\mathbf{b}|^2} \quad\Rightarrow\quad |\mathbf{a}| = |\mathbf{b}|$. Substituting this into the previous equation gives $\mathbf{a} = \mathbf{b}$.

So $\text{proj}_{\mathbf{a}}\,\mathbf{b} = \text{proj}_{\mathbf{b}}\,\mathbf{a} \quad\Leftrightarrow\quad \mathbf{a}$ and $\mathbf{b}$ are orthogonal, or they are equal.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. The displacement vector is $\mathbf{D} = (6-0)\,\mathbf{i} + (12-10)\,\mathbf{j} + (20-8)\,\mathbf{k} = 6\,\mathbf{i} + 2\,\mathbf{j} + 12\,\mathbf{k}$ so, by Equation 12, the work done is

$W = \mathbf{F} \cdot \mathbf{D} = (8\,\mathbf{i} - 6\,\mathbf{j} + 9\,\mathbf{k}) \cdot (6\,\mathbf{i} + 2\,\mathbf{j} + 12\,\mathbf{k}) = 48 - 12 + 108 = 144$ joules.

50. Here $|\mathbf{D}| = 1000$ m, $|\mathbf{F}| = 1500$ N, and $\theta = 30°$. Thus

$W = \mathbf{F} \cdot \mathbf{D} = |\mathbf{F}|\,|\mathbf{D}| \cos\theta = (1500)(1000)\left(\frac{\sqrt{3}}{2}\right) = 750{,}000\sqrt{3}$ joules.

51. Here $|\mathbf{D}| = 80$ ft, $|\mathbf{F}| = 30$ lb, and $\theta = 40°$. Thus

$W = \mathbf{F} \cdot \mathbf{D} = |\mathbf{F}|\,|\mathbf{D}| \cos\theta = (30)(80)\cos 40° = 2400\cos 40° \approx 1839$ ft-lb.

52. $W = \mathbf{F} \cdot \mathbf{D} = |\mathbf{F}|\,|\mathbf{D}| \cos\theta = (400)(120)\cos 36° \approx 38{,}833$ ft-lb

53. First note that $\mathbf{n} = \langle a, b\rangle$ is perpendicular to the line, because if $Q_1 = (a_1, b_1)$ and $Q_2 = (a_2, b_2)$ lie on the line, then

$\mathbf{n} \cdot \overrightarrow{Q_1Q_2} = aa_2 - aa_1 + bb_2 - bb_1 = 0$, since $aa_2 + bb_2 = -c = aa_1 + bb_1$ from the equation of the line.

Let $P_2 = (x_2, y_2)$ lie on the line. Then the distance from P_1 to the line is the absolute value of the scalar projection

of $\overrightarrow{P_1P_2}$ onto $\mathbf{n}$. $\operatorname{comp}_{\mathbf{n}}\left(\overrightarrow{P_1P_2}\right) = \dfrac{|\mathbf{n} \cdot \langle x_2 - x_1, y_2 - y_1\rangle|}{|\mathbf{n}|} = \dfrac{|ax_2 - ax_1 + by_2 - by_1|}{\sqrt{a^2+b^2}} = \dfrac{|ax_1 + by_1 + c|}{\sqrt{a^2+b^2}}$

since $ax_2 + by_2 = -c$. The required distance is $\dfrac{|(3)(-2) + (-4)(3) + 5|}{\sqrt{3^2 + (-4)^2}} = \dfrac{13}{5}$.

54. $(\mathbf{r} - \mathbf{a}) \cdot (\mathbf{r} - \mathbf{b}) = 0$ implies that the vectors $\mathbf{r} - \mathbf{a}$ and $\mathbf{r} - \mathbf{b}$ are orthogonal. From the diagram (in which A, B and R are the terminal points of the vectors), we see that this implies that R lies on a sphere whose diameter is the line from A to B. The center of this circle is the midpoint of AB, that is, $\frac{1}{2}(\mathbf{a} + \mathbf{b}) = \left\langle \frac{1}{2}(a_1 + b_1), \frac{1}{2}(a_2 + b_2), \frac{1}{2}(a_3 + b_3)\right\rangle$, and its radius is $\frac{1}{2}|\mathbf{a} - \mathbf{b}| = \frac{1}{2}\sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2 + (a_3 - b_3)^2}$.

Or: Expand the given equation, substitute $\mathbf{r} \cdot \mathbf{r} = x^2 + y^2 + z^2$ and complete the squares.

55. For convenience, consider the unit cube positioned so that its back left corner is at the origin, and its edges lie along the coordinate axes. The diagonal of the cube that begins at the origin and ends at $(1, 1, 1)$ has vector representation $\langle 1, 1, 1\rangle$. The angle θ between this vector and the vector of the edge which also begins at the origin and runs along the x-axis [that is, $\langle 1, 0, 0\rangle$] is given by $\cos\theta = \dfrac{\langle 1,1,1\rangle \cdot \langle 1,0,0\rangle}{|\langle 1,1,1\rangle|\,|\langle 1,0,0\rangle|} = \dfrac{1}{\sqrt{3}} \quad \Rightarrow \quad \theta = \cos^{-1}\left(\frac{1}{\sqrt{3}}\right) \approx 54.7°$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

56. Consider a cube with sides of unit length, wholly within the first octant and with edges along each of the three coordinate axes. $\mathbf{i}+\mathbf{j}+\mathbf{k}$ and $\mathbf{i}+\mathbf{j}$ are vector representations of a diagonal of the cube and a diagonal of one of its faces. If θ is the angle between these diagonals, then $\cos\theta = \dfrac{(\mathbf{i}+\mathbf{j}+\mathbf{k})\cdot(\mathbf{i}+\mathbf{j})}{|\mathbf{i}+\mathbf{j}+\mathbf{k}|\,|\mathbf{i}+\mathbf{j}|} = \dfrac{1+1}{\sqrt{3}\sqrt{2}} = \sqrt{\dfrac{2}{3}} \quad\Rightarrow\quad \theta = \cos^{-1}\sqrt{\tfrac{2}{3}} \approx 35.3^\circ$.

57. Consider the H — C — H combination consisting of the sole carbon atom and the two hydrogen atoms that are at $(1,0,0)$ and $(0,1,0)$ (or any H — C — H combination, for that matter). Vector representations of the line segments emanating from the carbon atom and extending to these two hydrogen atoms are $\left\langle 1-\frac{1}{2}, 0-\frac{1}{2}, 0-\frac{1}{2}\right\rangle = \left\langle \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right\rangle$ and $\left\langle 0-\frac{1}{2}, 1-\frac{1}{2}, 0-\frac{1}{2}\right\rangle = \left\langle -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\right\rangle$. The bond angle, θ, is therefore given by

$$\cos\,\theta = \frac{\left\langle \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right\rangle \cdot \left\langle -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\right\rangle}{\left|\left\langle \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right\rangle\right| \left|\left\langle -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\right\rangle\right|} = \frac{-\frac{1}{4}-\frac{1}{4}+\frac{1}{4}}{\sqrt{\frac{3}{4}}\sqrt{\frac{3}{4}}} = -\frac{1}{3} \quad\Rightarrow\quad \theta = \cos^{-1}\left(-\tfrac{1}{3}\right) \approx 109.5^\circ.$$

58. Let α be the angle between $\mathbf{a}$ and $\mathbf{c}$ and β be the angle between $\mathbf{c}$ and $\mathbf{b}$. We need to show that $\alpha = \beta$. Now

$\cos\alpha = \dfrac{\mathbf{a}\cdot\mathbf{c}}{|\mathbf{a}|\,|\mathbf{c}|} = \dfrac{\mathbf{a}\cdot|\mathbf{a}|\,\mathbf{b}+\mathbf{a}\cdot|\mathbf{b}|\,\mathbf{a}}{|\mathbf{a}|\,|\mathbf{c}|} = \dfrac{|\mathbf{a}|\,\mathbf{a}\cdot\mathbf{b}+|\mathbf{a}|^2\,|\mathbf{b}|}{|\mathbf{a}|\,|\mathbf{c}|} = \dfrac{\mathbf{a}\cdot\mathbf{b}+|\mathbf{a}|\,|\mathbf{b}|}{|\mathbf{c}|}$. Similarly,

$\cos\beta = \dfrac{\mathbf{b}\cdot\mathbf{c}}{|\mathbf{b}|\,|\mathbf{c}|} = \dfrac{|\mathbf{a}|\,|\mathbf{b}|+\mathbf{b}\cdot\mathbf{a}}{|\mathbf{c}|}$. Thus $\cos\alpha = \cos\beta$. However $0^\circ \le \alpha \le 180^\circ$ and $0^\circ \le \beta \le 180^\circ$, so $\alpha = \beta$ and $\mathbf{c}$ bisects the angle between $\mathbf{a}$ and $\mathbf{b}$.

59. Let $\mathbf{a} = \langle a_1, a_2, a_3\rangle$ and $\mathbf{b} = \langle b_1, b_2, b_3\rangle$.

Property 2: $\mathbf{a}\cdot\mathbf{b} = \langle a_1, a_2, a_3\rangle \cdot \langle b_1, b_2, b_3\rangle = a_1b_1 + a_2b_2 + a_3b_3$

$= b_1a_1 + b_2a_2 + b_3a_3 = \langle b_1, b_2, b_3\rangle \cdot \langle a_1, a_2, a_3\rangle = \mathbf{b}\cdot\mathbf{a}$

Property 4: $(c\,\mathbf{a})\cdot\mathbf{b} = \langle ca_1, ca_2, ca_3\rangle \cdot \langle b_1, b_2, b_3\rangle = (ca_1)b_1 + (ca_2)b_2 + (ca_3)b_3$

$= c\,(a_1b_1 + a_2b_2 + a_3b_3) = c\,(\mathbf{a}\cdot\mathbf{b}) = a_1(cb_1) + a_2(cb_2) + a_3(cb_3)$

$= \langle a_1, a_2, a_3\rangle \cdot \langle cb_1, cb_2, cb_3\rangle = \mathbf{a}\cdot(c\,\mathbf{b})$

Property 5: $\mathbf{0}\cdot\mathbf{a} = \langle 0,0,0\rangle \cdot \langle a_1, a_2, a_3\rangle = (0)(a_1) + (0)(a_2) + (0)(a_3) = 0$

60. Let the figure be called quadrilateral $ABCD$. The diagonals can be represented by $\overrightarrow{AC}$ and $\overrightarrow{BD}$. $\overrightarrow{AC} = \overrightarrow{AB} + \overrightarrow{BC}$ and $\overrightarrow{BD} = \overrightarrow{BC} + \overrightarrow{CD} = \overrightarrow{BC} - \overrightarrow{DC} = \overrightarrow{BC} - \overrightarrow{AB}$ (Since opposite sides of the object are of the same length and parallel, $\overrightarrow{AB} = \overrightarrow{DC}$.) Thus

$$\begin{aligned}\overrightarrow{AC}\cdot\overrightarrow{BD} &= \left(\overrightarrow{AB}+\overrightarrow{BC}\right)\cdot\left(\overrightarrow{BC}-\overrightarrow{AB}\right) = \overrightarrow{AB}\cdot\left(\overrightarrow{BC}-\overrightarrow{AB}\right) + \overrightarrow{BC}\cdot\left(\overrightarrow{BC}-\overrightarrow{AB}\right)\\ &= \overrightarrow{AB}\cdot\overrightarrow{BC} - \left|\overrightarrow{AB}\right|^2 + \left|\overrightarrow{BC}\right|^2 - \overrightarrow{AB}\cdot\overrightarrow{BC} = \left|\overrightarrow{BC}\right|^2 - \left|\overrightarrow{AB}\right|^2\end{aligned}$$

But $\left|\overrightarrow{AB}\right|^2 = \left|\overrightarrow{BC}\right|^2$ because all sides of the quadrilateral are equal in length. Therefore $\overrightarrow{AC}\cdot\overrightarrow{BD} = 0$, and since both of these vectors are nonzero this tells us that the diagonals of the quadrilateral are perpendicular.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. $|\mathbf{a}\cdot\mathbf{b}| = \big|\,|\mathbf{a}|\,|\mathbf{b}|\cos\theta\big| = |\mathbf{a}|\,|\mathbf{b}|\,|\cos\theta|$. Since $|\cos\theta| \le 1$, $|\mathbf{a}\cdot\mathbf{b}| = |\mathbf{a}|\,|\mathbf{b}|\,|\cos\,\theta| \le |\mathbf{a}|\,|\mathbf{b}|$.

Note: We have equality in the case of $\cos\theta = \pm 1$, so $\theta = 0$ or $\theta = \pi$, thus equality when $\mathbf{a}$ and $\mathbf{b}$ are parallel.

62. (a)

The Triangle Inequality states that the length of the longest side of a triangle is less than or equal to the sum of the lengths of the two shortest sides.

(b) $|\mathbf{a}+\mathbf{b}|^2 = (\mathbf{a}+\mathbf{b})\cdot(\mathbf{a}+\mathbf{b}) = (\mathbf{a}\cdot\mathbf{a}) + 2(\mathbf{a}\cdot\mathbf{b}) + (\mathbf{b}\cdot\mathbf{b}) = |\mathbf{a}|^2 + 2(\mathbf{a}\cdot\mathbf{b}) + |\mathbf{b}|^2$

$$\le |\mathbf{a}|^2 + 2\,|\mathbf{a}|\,|\mathbf{b}| + |\mathbf{b}|^2 \qquad \text{[by the Cauchy-Schwartz Inequality]}$$

$$= (|\mathbf{a}| + |\mathbf{b}|)^2$$

Thus, taking the square root of both sides, $|\mathbf{a}+\mathbf{b}| \le |\mathbf{a}| + |\mathbf{b}|$.

63. (a)

The Parallelogram Law states that the sum of the squares of the lengths of the diagonals of a parallelogram equals the sum of the squares of its (four) sides.

(b) $|\mathbf{a}+\mathbf{b}|^2 = (\mathbf{a}+\mathbf{b})\cdot(\mathbf{a}+\mathbf{b}) = |\mathbf{a}|^2 + 2(\mathbf{a}\cdot\mathbf{b}) + |\mathbf{b}|^2$ and $|\mathbf{a}-\mathbf{b}|^2 = (\mathbf{a}-\mathbf{b})\cdot(\mathbf{a}-\mathbf{b}) = |\mathbf{a}|^2 - 2(\mathbf{a}\cdot\mathbf{b}) + |\mathbf{b}|^2$.

Adding these two equations gives $|\mathbf{a}+\mathbf{b}|^2 + |\mathbf{a}-\mathbf{b}|^2 = 2\,|\mathbf{a}|^2 + 2\,|\mathbf{b}|^2$.

64. If the vectors $\mathbf{u}+\mathbf{v}$ and $\mathbf{u}-\mathbf{v}$ are orthogonal, then $(\mathbf{u}+\mathbf{v})\cdot(\mathbf{u}-\mathbf{v}) = 0$. But

$$\begin{aligned}
(\mathbf{u}+\mathbf{v})\cdot(\mathbf{u}-\mathbf{v}) &= (\mathbf{u}+\mathbf{v})\cdot\mathbf{u} - (\mathbf{u}+\mathbf{v})\cdot\mathbf{v} && \text{by Property 3 of the dot product}\\
&= \mathbf{u}\cdot\mathbf{u} + \mathbf{v}\cdot\mathbf{u} - \mathbf{u}\cdot\mathbf{v} - \mathbf{v}\cdot\mathbf{v} && \text{by Property 3}\\
&= |\mathbf{u}|^2 + \mathbf{u}\cdot\mathbf{v} - \mathbf{u}\cdot\mathbf{v} - |\mathbf{v}|^2 && \text{by Properties 1 and 2}\\
&= |\mathbf{u}|^2 - |\mathbf{v}|^2
\end{aligned}$$

Thus $|\mathbf{u}|^2 - |\mathbf{v}|^2 = 0 \;\Rightarrow\; |\mathbf{u}|^2 = |\mathbf{v}|^2 \;\Rightarrow\; |\mathbf{u}| = |\mathbf{v}|$ [since $|\mathbf{u}|, |\mathbf{v}| \ge 0$].

65.

$$\begin{aligned}
\operatorname{proj}_{\mathbf{a}}\mathbf{b}\cdot\operatorname{proj}_{\mathbf{b}}\mathbf{a} &= \frac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|^2}\mathbf{a}\cdot\frac{\mathbf{b}\cdot\mathbf{a}}{|\mathbf{b}|^2}\mathbf{b} = \frac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|^2}\cdot\frac{\mathbf{b}\cdot\mathbf{a}}{|\mathbf{b}|^2}\,(\mathbf{a}\cdot\mathbf{b}) && \text{by Property 4 of the dot product}\\
&= \frac{(\mathbf{a}\cdot\mathbf{b})^2}{|\mathbf{a}|^2\,|\mathbf{b}|^2}\,(\mathbf{a}\cdot\mathbf{b}) = \left(\frac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|}\right)^2(\mathbf{a}\cdot\mathbf{b}) && \text{by Property 2}\\
&= (\cos\theta)^2\,(\mathbf{a}\cdot\mathbf{b}) = (\mathbf{a}\cdot\mathbf{b})\cos^2\theta && \text{by Corollary 6}
\end{aligned}$$

66. (a) Suppose that $\mathbf{u}$ and $\mathbf{v}$ are nonzero orthogonal vectors. Then

$$\begin{aligned}
|\mathbf{u}+\mathbf{v}|^2 &= (\mathbf{u}+\mathbf{v})\cdot(\mathbf{u}+\mathbf{v})\\
&= \mathbf{u}\cdot\mathbf{u} + \mathbf{u}\cdot\mathbf{v} + \mathbf{v}\cdot\mathbf{u} + \mathbf{v}\cdot\mathbf{v}\\
&= |\mathbf{u}|^2 + 0 + 0 + |\mathbf{v}|^2 = |\mathbf{u}|^2 + |\mathbf{v}|^2
\end{aligned}$$

(b) Suppose that $|\mathbf{u}+\mathbf{v}|^2 = |\mathbf{u}|^2 + |\mathbf{v}|^2$. From part (a), we know that $|\mathbf{u}+\mathbf{v}|^2 = |\mathbf{u}|^2 + 2(\mathbf{u}\cdot\mathbf{v}) + |\mathbf{v}|^2$. Thus, $2(\mathbf{u}\cdot\mathbf{v}) = 0$, which implies that $\mathbf{u}$ and $\mathbf{v}$ are orthogonal.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12.4 The Cross Product

1. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\2&3&0\\1&0&5\end{vmatrix}=\begin{vmatrix}3&0\\0&5\end{vmatrix}\mathbf{i}-\begin{vmatrix}2&0\\1&5\end{vmatrix}\mathbf{j}+\begin{vmatrix}2&3\\1&0\end{vmatrix}\mathbf{k}$

$=(15-0)\,\mathbf{i}-(10-0)\,\mathbf{j}+(0-3)\,\mathbf{k}=15\,\mathbf{i}-10\,\mathbf{j}-3\,\mathbf{k}$

Now $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}=\langle 15,-10,-3\rangle\cdot\langle 2,3,0\rangle=30-30+0=0$ and

$(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}=\langle 15,-10,-3\rangle\cdot\langle 1,0,5\rangle=15+0-15=0$, so $\mathbf{a}\times\mathbf{b}$ is orthogonal to both $\mathbf{a}$ and $\mathbf{b}$.

2. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\4&3&-2\\2&-1&1\end{vmatrix}=\begin{vmatrix}3&-2\\-1&1\end{vmatrix}\mathbf{i}-\begin{vmatrix}4&-2\\2&1\end{vmatrix}\mathbf{j}+\begin{vmatrix}4&3\\2&-1\end{vmatrix}\mathbf{k}$

$=(3-2)\,\mathbf{i}-[4-(-4)]\,\mathbf{j}+(-4-6)\,\mathbf{k}=\mathbf{i}-8\,\mathbf{j}-10\,\mathbf{k}$

Now $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}=\langle 1,-8,-10\rangle\cdot\langle 4,3,-2\rangle=4-24+20=0$ and

$(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}=\langle 1,-8,-10\rangle\cdot\langle 2,-1,1\rangle=2+8-10=0$, so $\mathbf{a}\times\mathbf{b}$ is orthogonal to both $\mathbf{a}$ and $\mathbf{b}$.

3. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\0&2&-4\\-1&3&1\end{vmatrix}=\begin{vmatrix}2&-4\\3&1\end{vmatrix}\mathbf{i}-\begin{vmatrix}0&-4\\-1&1\end{vmatrix}\mathbf{j}+\begin{vmatrix}0&2\\-1&3\end{vmatrix}\mathbf{k}$

$=[2-(-12)]\,\mathbf{i}-(0-4)\,\mathbf{j}+[0-(-2)]\,\mathbf{k}=14\,\mathbf{i}+4\,\mathbf{j}+2\,\mathbf{k}$

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}=(14\,\mathbf{i}+4\,\mathbf{j}+2\,\mathbf{k})\cdot(2\,\mathbf{j}-4\,\mathbf{k})=0+8-8=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{a}$.

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}=(14\,\mathbf{i}+4\,\mathbf{j}+2\,\mathbf{k})\cdot(-\mathbf{i}+3\,\mathbf{j}+\mathbf{k})=-14+12+2=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{b}$.

4. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\3&3&-3\\3&-3&3\end{vmatrix}=\begin{vmatrix}3&-3\\-3&3\end{vmatrix}\mathbf{i}-\begin{vmatrix}3&-3\\3&3\end{vmatrix}\mathbf{j}+\begin{vmatrix}3&3\\3&-3\end{vmatrix}\mathbf{k}$

$=(9-9)\,\mathbf{i}-[9-(-9)]\,\mathbf{j}+(-9-9)\,\mathbf{k}=-18\,\mathbf{j}-18\,\mathbf{k}$

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}=(-18\,\mathbf{j}-18\,\mathbf{k})\cdot(3\,\mathbf{i}+3\,\mathbf{j}-3\,\mathbf{k})=0-54+54=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{a}$.

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}=(-18\,\mathbf{j}-18\,\mathbf{k})\cdot(3\,\mathbf{i}-3\,\mathbf{j}+3\,\mathbf{k})=0+54-54=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{b}$.

5. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i}&\mathbf{j}&\mathbf{k}\\\frac{1}{2}&\frac{1}{3}&\frac{1}{4}\\1&2&-3\end{vmatrix}=\begin{vmatrix}\frac{1}{3}&\frac{1}{4}\\2&-3\end{vmatrix}\mathbf{i}-\begin{vmatrix}\frac{1}{2}&\frac{1}{4}\\1&-3\end{vmatrix}\mathbf{j}+\begin{vmatrix}\frac{1}{2}&\frac{1}{3}\\1&2\end{vmatrix}\mathbf{k}$

$=\left(-1-\frac{1}{2}\right)\mathbf{i}-\left(-\frac{3}{2}-\frac{1}{4}\right)\mathbf{j}+\left(1-\frac{1}{3}\right)\mathbf{k}=-\frac{3}{2}\,\mathbf{i}+\frac{7}{4}\,\mathbf{j}+\frac{2}{3}\,\mathbf{k}$

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}=\left(-\frac{3}{2}\,\mathbf{i}+\frac{7}{4}\,\mathbf{j}+\frac{2}{3}\,\mathbf{k}\right)\cdot\left(\frac{1}{2}\,\mathbf{i}+\frac{1}{3}\,\mathbf{j}+\frac{1}{4}\,\mathbf{k}\right)=-\frac{3}{4}+\frac{7}{12}+\frac{1}{6}=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{a}$.

Since $(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}=\left(-\frac{3}{2}\,\mathbf{i}+\frac{7}{4}\,\mathbf{j}+\frac{2}{3}\,\mathbf{k}\right)\cdot(\mathbf{i}+2\,\mathbf{j}-3\,\mathbf{k})=-\frac{3}{2}+\frac{7}{2}-2=0$, $\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{b}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ t & \cos t & \sin t\\ 1 & -\sin t & \cos t\end{vmatrix}=\begin{vmatrix}\cos t & \sin t\\ -\sin t & \cos t\end{vmatrix}\mathbf{i}-\begin{vmatrix}t & \sin t\\ 1 & \cos t\end{vmatrix}\mathbf{j}+\begin{vmatrix}t & \cos t\\ 1 & -\sin t\end{vmatrix}\mathbf{k}$

$=[\cos^2 t-(-\sin^2 t)]\,\mathbf{i}-(t\cos t-\sin t)\,\mathbf{j}+(-t\sin t-\cos t)\,\mathbf{k}=\mathbf{i}+(\sin t-t\cos t)\,\mathbf{j}+(-t\sin t-\cos t)\,\mathbf{k}$

Since
$$\begin{aligned}(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}&=[\,\mathbf{i}+(\sin t-t\cos t)\,\mathbf{j}+(-t\sin t-\cos t)\,\mathbf{k}\,]\cdot(t\,\mathbf{i}+\cos t\,\mathbf{j}+\sin t\,\mathbf{k})\\&=t+\sin t\cos t-t\cos^2 t-t\sin^2 t-\sin t\cos t\\&=t-t\left(\cos^2 t+\sin^2 t\right)=0\end{aligned}$$

$\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{a}$.

Since
$$\begin{aligned}(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}&=[\,\mathbf{i}+(\sin t-t\cos t)\,\mathbf{j}+(-t\sin t-\cos t)\,\mathbf{k}\,]\cdot(\mathbf{i}-\sin t\,\mathbf{j}+\cos t\,\mathbf{k})\\&=1-\sin^2 t+t\sin t\cos t-t\sin t\cos t-\cos^2 t\\&=1-\left(\sin^2 t+\cos^2 t\right)=0\end{aligned}$$

$\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{b}$.

7. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ t^3 & t^2 & t\\ t & 2t & 3t\end{vmatrix}=\begin{vmatrix}t^2 & t\\ 2t & 3t\end{vmatrix}\mathbf{i}-\begin{vmatrix}t^3 & t\\ t & 3t\end{vmatrix}\mathbf{j}+\begin{vmatrix}t^3 & t^2\\ t & 2t\end{vmatrix}\mathbf{k}$

$=(3t^3-2t^2)\mathbf{i}-(3t^4-t^2)\mathbf{j}+(2t^4-t^3)\mathbf{k}$

Since
$$\begin{aligned}(\mathbf{a}\times\mathbf{b})\cdot\mathbf{a}&=\left\langle 3t^3-2t^2,t^2-3t^4,2t^4-t^3\right\rangle\cdot\left\langle t^3,t^2,t\right\rangle\\&=3t^6-2t^5+t^4-3t^6+2t^5-t^4=0\end{aligned}$$

$\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{a}$.

Since
$$\begin{aligned}(\mathbf{a}\times\mathbf{b})\cdot\mathbf{b}&=\left\langle 3t^3-2t^2,t^2-3t^4,2t^4-t^3\right\rangle\cdot\langle t,2t,3t\rangle\\&=3t^4-2t^3+2t^3-6t^5+6t^5-3t^4=0\end{aligned}$$

$\mathbf{a}\times\mathbf{b}$ is orthogonal to $\mathbf{b}$.

8. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ 1 & 0 & -2\\ 0 & 1 & 1\end{vmatrix}$

$=\begin{vmatrix}0 & -2\\ 1 & 1\end{vmatrix}\mathbf{i}-\begin{vmatrix}1 & -2\\ 0 & 1\end{vmatrix}\mathbf{j}+\begin{vmatrix}1 & 0\\ 0 & 1\end{vmatrix}\mathbf{k}$

$=2\,\mathbf{i}-\mathbf{j}+\mathbf{k}$

9. According to the discussion following Example 4, $\mathbf{i}\times\mathbf{j}=\mathbf{k}$, so $(\mathbf{i}\times\mathbf{j})\times\mathbf{k}=\mathbf{k}\times\mathbf{k}=\mathbf{0}$ [by Example 2].

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $\mathbf{k} \times (\mathbf{i} - 2\,\mathbf{j}) = \mathbf{k} \times \mathbf{i} + \mathbf{k} \times (-2\,\mathbf{j})$ by Property 3 of the cross product

$= \mathbf{k} \times \mathbf{i} + (-2)\,(\mathbf{k} \times \mathbf{j})$ by Property 2

$= \mathbf{j} + (-2)(-\mathbf{i}) = 2\,\mathbf{i} + \mathbf{j}$ by the discussion following Example 4

11. $(\mathbf{j} - \mathbf{k}) \times (\mathbf{k} - \mathbf{i}) = (\mathbf{j} - \mathbf{k}) \times \mathbf{k} + (\mathbf{j} - \mathbf{k}) \times (-\mathbf{i})$ by Property 3 of the cross product

$= \mathbf{j} \times \mathbf{k} + (-\mathbf{k}) \times \mathbf{k} + \mathbf{j} \times (-\mathbf{i}) + (-\mathbf{k}) \times (-\mathbf{i})$ by Property 4

$= (\mathbf{j} \times \mathbf{k}) + (-1)(\mathbf{k} \times \mathbf{k}) + (-1)(\mathbf{j} \times \mathbf{i}) + (-1)^2(\mathbf{k} \times \mathbf{i})$ by Property 2

$= \mathbf{i} + (-1)\,\mathbf{0} + (-1)(-\mathbf{k}) + \mathbf{j} = \mathbf{i} + \mathbf{j} + \mathbf{k}$ by Example 2 and the discussion following Example 4

12. $(\mathbf{i} + \mathbf{j}) \times (\mathbf{i} - \mathbf{j}) = (\mathbf{i} + \mathbf{j}) \times \mathbf{i} + (\mathbf{i} + \mathbf{j}) \times (-\mathbf{j})$ by Property 3 of the cross product

$= \mathbf{i} \times \mathbf{i} + \mathbf{j} \times \mathbf{i} + \mathbf{i} \times (-\mathbf{j}) + \mathbf{j} \times (-\mathbf{j})$ by Property 4

$= (\mathbf{i} \times \mathbf{i}) + (\mathbf{j} \times \mathbf{i}) + (-1)(\mathbf{i} \times \mathbf{j}) + (-1)(\mathbf{j} \times \mathbf{j})$ by Property 2

$= \mathbf{0} + (-\mathbf{k}) + (-1)\,\mathbf{k} + (-1)\,\mathbf{0} = -2\,\mathbf{k}$ by Example 2 and the discussion following Example 4

13. (a) Since $\mathbf{b} \times \mathbf{c}$ is a vector, the dot product $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$ is meaningful and is a scalar.

(b) $\mathbf{b} \cdot \mathbf{c}$ is a scalar, so $\mathbf{a} \times (\mathbf{b} \cdot \mathbf{c})$ is meaningless, as the cross product is defined only for two *vectors*.

(c) Since $\mathbf{b} \times \mathbf{c}$ is a vector, the cross product $\mathbf{a} \times (\mathbf{b} \times \mathbf{c})$ is meaningful and results in another vector.

(d) $\mathbf{b} \cdot \mathbf{c}$ is a scalar, so the dot product $\mathbf{a} \cdot (\mathbf{b} \cdot \mathbf{c})$ is meaningless, as the dot product is defined only for two vectors.

(e) Since $(\mathbf{a} \cdot \mathbf{b})$ and $(\mathbf{c} \cdot \mathbf{d})$ are both scalars, the cross product $(\mathbf{a} \cdot \mathbf{b}) \times (\mathbf{c} \cdot \mathbf{d})$ is meaningless.

(f) $\mathbf{a} \times \mathbf{b}$ and $\mathbf{c} \times \mathbf{d}$ are both vectors, so the dot product $(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d})$ is meaningful and is a scalar.

14. Using Theorem 9, we have $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin\theta = (10)(8)\sin 60^\circ = 80 \cdot \frac{\sqrt{3}}{2} = 40\sqrt{3}$. By the right-hand rule, $\mathbf{u} \times \mathbf{v}$ is directed into the page.

15. If we sketch $\mathbf{u}$ and $\mathbf{v}$ starting from the same initial point, we see that the angle between them is 30°.

Using Theorem 9, we have $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin\theta = (4)(3)\sin 30^\circ = 12 \cdot \frac{1}{2} = 6$. By the right-hand rule, $\mathbf{u} \times \mathbf{v}$ is directed into the page.

16. (a) $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}|\,|\mathbf{b}| \sin\theta = 3 \cdot 2 \cdot \sin\frac{\pi}{2} = 6$

(b) $\mathbf{a} \times \mathbf{b}$ is orthogonal to $\mathbf{k}$, so it lies in the xy-plane, and its z-coordinate is 0. By the right-hand rule, its y-component is negative and its x-component is positive.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ 2 & -1 & 3\\ 4 & 2 & 1\end{vmatrix}=\begin{vmatrix}-1 & 3\\ 2 & 1\end{vmatrix}\mathbf{i}-\begin{vmatrix}2 & 3\\ 4 & 1\end{vmatrix}\mathbf{j}+\begin{vmatrix}2 & -1\\ 4 & 2\end{vmatrix}\mathbf{k}=(-1-6)\,\mathbf{i}-(2-12)\,\mathbf{j}+[4-(-4)]\,\mathbf{k}=-7\,\mathbf{i}+10\,\mathbf{j}+8\,\mathbf{k}$

$\mathbf{b}\times\mathbf{a}=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ 4 & 2 & 1\\ 2 & -1 & 3\end{vmatrix}=\begin{vmatrix}2 & 1\\ -1 & 3\end{vmatrix}\mathbf{i}-\begin{vmatrix}4 & 1\\ 2 & 3\end{vmatrix}\mathbf{j}+\begin{vmatrix}4 & 2\\ 2 & -1\end{vmatrix}\mathbf{k}=[6-(-1)]\,\mathbf{i}-(12-2)\,\mathbf{j}+(-4-4)\,\mathbf{k}=7\,\mathbf{i}-10\,\mathbf{j}-8\,\mathbf{k}$

Notice $\mathbf{a}\times\mathbf{b}=-\mathbf{b}\times\mathbf{a}$ here, as we know is always true by Property 1 of the cross product.

18. $\mathbf{b}\times\mathbf{c}=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ 2 & 1 & -1\\ 0 & 1 & 3\end{vmatrix}=\begin{vmatrix}1 & -1\\ 1 & 3\end{vmatrix}\mathbf{i}-\begin{vmatrix}2 & -1\\ 0 & 3\end{vmatrix}\mathbf{j}+\begin{vmatrix}2 & 1\\ 0 & 1\end{vmatrix}\mathbf{k}=4\,\mathbf{i}-6\,\mathbf{j}+2\,\mathbf{k}$ so

$\mathbf{a}\times(\mathbf{b}\times\mathbf{c})=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ 1 & 0 & 1\\ 4 & -6 & 2\end{vmatrix}=\begin{vmatrix}0 & 1\\ -6 & 2\end{vmatrix}\mathbf{i}-\begin{vmatrix}1 & 1\\ 4 & 2\end{vmatrix}\mathbf{j}+\begin{vmatrix}1 & 0\\ 4 & -6\end{vmatrix}\mathbf{k}=6\,\mathbf{i}+2\,\mathbf{j}-6\,\mathbf{k}.$

$\mathbf{a}\times\mathbf{b}=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ 1 & 0 & 1\\ 2 & 1 & -1\end{vmatrix}=\begin{vmatrix}0 & 1\\ 1 & -1\end{vmatrix}\mathbf{i}-\begin{vmatrix}1 & 1\\ 2 & -1\end{vmatrix}\mathbf{j}+\begin{vmatrix}1 & 0\\ 2 & 1\end{vmatrix}\mathbf{k}=-\,\mathbf{i}+3\,\mathbf{j}+\mathbf{k}$ so

$(\mathbf{a}\times\mathbf{b})\times\mathbf{c}=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ -1 & 3 & 1\\ 0 & 1 & 3\end{vmatrix}=\begin{vmatrix}3 & 1\\ 1 & 3\end{vmatrix}\mathbf{i}-\begin{vmatrix}-1 & 1\\ 0 & 3\end{vmatrix}\mathbf{j}+\begin{vmatrix}-1 & 3\\ 0 & 1\end{vmatrix}\mathbf{k}=8\,\mathbf{i}+3\,\mathbf{j}-\mathbf{k}.$

Thus $\mathbf{a}\times(\mathbf{b}\times\mathbf{c})\neq(\mathbf{a}\times\mathbf{b})\times\mathbf{c}$.

19. By Theorem 8, the cross product of two vectors is orthogonal to both vectors. So we calculate

$\langle 3,2,1\rangle\times\langle -1,1,0\rangle=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ 3 & 2 & 1\\ -1 & 1 & 0\end{vmatrix}=\begin{vmatrix}2 & 1\\ 1 & 0\end{vmatrix}\mathbf{i}-\begin{vmatrix}3 & 1\\ -1 & 0\end{vmatrix}\mathbf{j}+\begin{vmatrix}3 & 2\\ -1 & 1\end{vmatrix}\mathbf{k}=-\mathbf{i}-\mathbf{j}+5\,\mathbf{k}.$

So two unit vectors orthogonal to both given vectors are $\pm\dfrac{\langle -1,-1,5\rangle}{\sqrt{1+1+25}}=\pm\dfrac{\langle -1,-1,5\rangle}{3\sqrt{3}}$, that is, $\left\langle -\frac{1}{3\sqrt{3}},-\frac{1}{3\sqrt{3}},\frac{5}{3\sqrt{3}}\right\rangle$ and $\left\langle \frac{1}{3\sqrt{3}},\frac{1}{3\sqrt{3}},-\frac{5}{3\sqrt{3}}\right\rangle$.

20. By Theorem 8, the cross product of two vectors is orthogonal to both vectors. So we calculate

$$(\mathbf{j}-\mathbf{k})\times(\mathbf{i}+\mathbf{j})=\begin{vmatrix}\mathbf{i} & \mathbf{j} & \mathbf{k}\\ 0 & 1 & -1\\ 1 & 1 & 0\end{vmatrix}=\begin{vmatrix}1 & -1\\ 1 & 0\end{vmatrix}\mathbf{i}-\begin{vmatrix}0 & -1\\ 1 & 0\end{vmatrix}\mathbf{j}+\begin{vmatrix}0 & 1\\ 1 & 1\end{vmatrix}\mathbf{k}=\mathbf{i}-\mathbf{j}-\mathbf{k}$$

Thus two unit vectors orthogonal to both given vectors are $\pm\frac{1}{\sqrt{3}}(\mathbf{i}-\mathbf{j}-\mathbf{k})$, that is, $\frac{1}{\sqrt{3}}\,\mathbf{i}-\frac{1}{\sqrt{3}}\,\mathbf{j}-\frac{1}{\sqrt{3}}\,\mathbf{k}$ and $-\frac{1}{\sqrt{3}}\,\mathbf{i}+\frac{1}{\sqrt{3}}\,\mathbf{j}+\frac{1}{\sqrt{3}}\,\mathbf{k}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. Let $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$. Then

$$\mathbf{0} \times \mathbf{a} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0 & 0 \\ a_1 & a_2 & a_3 \end{vmatrix} = \begin{vmatrix} 0 & 0 \\ a_2 & a_3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 0 & 0 \\ a_1 & a_3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 0 & 0 \\ a_1 & a_2 \end{vmatrix} \mathbf{k} = \mathbf{0},$$

$$\mathbf{a} \times \mathbf{0} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ 0 & 0 & 0 \end{vmatrix} = \begin{vmatrix} a_2 & a_3 \\ 0 & 0 \end{vmatrix} \mathbf{i} - \begin{vmatrix} a_1 & a_3 \\ 0 & 0 \end{vmatrix} \mathbf{j} + \begin{vmatrix} a_1 & a_2 \\ 0 & 0 \end{vmatrix} \mathbf{k} = \mathbf{0}.$$

22. Let $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$ and $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$.

$$(\mathbf{a} \times \mathbf{b}) \cdot \mathbf{b} = \left\langle \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix}, \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix}, \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} \right\rangle \cdot \langle b_1, b_2, b_3 \rangle = \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} b_1 - \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} b_2 + \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} b_3$$

$$= (a_2 b_3 b_1 - a_3 b_2 b_1) - (a_1 b_3 b_2 - a_3 b_1 b_2) + (a_1 b_2 b_3 - a_2 b_1 b_3) = 0$$

23. $\mathbf{a} \times \mathbf{b} = \langle a_2 b_3 - a_3 b_2, a_3 b_1 - a_1 b_3, a_1 b_2 - a_2 b_1 \rangle$

$$= \langle (-1)(b_2 a_3 - b_3 a_2), (-1)(b_3 a_1 - b_1 a_3), (-1)(b_1 a_2 - b_2 a_1) \rangle$$

$$= -\langle b_2 a_3 - b_3 a_2, b_3 a_1 - b_1 a_3, b_1 a_2 - b_2 a_1 \rangle = -\mathbf{b} \times \mathbf{a}$$

24. $c\mathbf{a} = \langle ca_1, ca_2, ca_3 \rangle$, so

$$(c\,\mathbf{a}) \times \mathbf{b} = \langle ca_2 b_3 - ca_3 b_2, ca_3 b_1 - ca_1 b_3, ca_1 b_2 - ca_2 b_1 \rangle$$

$$= c\langle a_2 b_3 - a_3 b_2, a_3 b_1 - a_1 b_3, a_1 b_2 - a_2 b_1 \rangle = c(\mathbf{a} \times \mathbf{b})$$

$$= \langle ca_2 b_3 - ca_3 b_2, ca_3 b_1 - ca_1 b_3, ca_1 b_2 - ca_2 b_1 \rangle$$

$$= \langle a_2(cb_3) - a_3(cb_2), a_3(cb_1) - a_1(cb_3), a_1(cb_2) - a_2(cb_1) \rangle$$

$$= \mathbf{a} \times (c\,\mathbf{b})$$

25. $\mathbf{a} \times (\mathbf{b} + \mathbf{c}) = \mathbf{a} \times \langle b_1 + c_1, b_2 + c_2, b_3 + c_3 \rangle$

$$= \langle a_2(b_3 + c_3) - a_3(b_2 + c_2), a_3(b_1 + c_1) - a_1(b_3 + c_3), a_1(b_2 + c_2) - a_2(b_1 + c_1) \rangle$$

$$= \langle a_2 b_3 + a_2 c_3 - a_3 b_2 - a_3 c_2, a_3 b_1 + a_3 c_1 - a_1 b_3 - a_1 c_3, a_1 b_2 + a_1 c_2 - a_2 b_1 - a_2 c_1 \rangle$$

$$= \langle (a_2 b_3 - a_3 b_2) + (a_2 c_3 - a_3 c_2), (a_3 b_1 - a_1 b_3) + (a_3 c_1 - a_1 c_3), (a_1 b_2 - a_2 b_1) + (a_1 c_2 - a_2 c_1) \rangle$$

$$= \langle a_2 b_3 - a_3 b_2, a_3 b_1 - a_1 b_3, a_1 b_2 - a_2 b_1 \rangle + \langle a_2 c_3 - a_3 c_2, a_3 c_1 - a_1 c_3, a_1 c_2 - a_2 c_1 \rangle$$

$$= (\mathbf{a} \times \mathbf{b}) + (\mathbf{a} \times \mathbf{c})$$

26. $(\mathbf{a} + \mathbf{b}) \times \mathbf{c} = -\mathbf{c} \times (\mathbf{a} + \mathbf{b})$ by Property 1 of the cross product

$= -(\mathbf{c} \times \mathbf{a} + \mathbf{c} \times \mathbf{b})$ by Property 3

$= -(-\mathbf{a} \times \mathbf{c} + (-\mathbf{b} \times \mathbf{c}))$ by Property 1

$= \mathbf{a} \times \mathbf{c} + \mathbf{b} \times \mathbf{c}$ by Property 2

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. By plotting the vertices, we can see that the parallelogram is determined by the vectors $\overrightarrow{AB} = \langle 2, 3\rangle$ and $\overrightarrow{AD} = \langle 6, -1\rangle$. We know that the area of the parallelogram determined by two vectors is equal to the length of the cross product of these vectors. In order to compute the cross product, we consider the vector $\overrightarrow{AB}$ as the three-dimensional vector $\langle 2, 3, 0\rangle$ (and similarly for $\overrightarrow{AD}$), and then the area of parallelogram $ABCD$ is

$$\left|\overrightarrow{AB} \times \overrightarrow{AD}\right| = \left\|\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 3 & 0 \\ 6 & -1 & 0 \end{vmatrix}\right\| = |(0-0)\,\mathbf{i} - (0-0)\,\mathbf{j} + (-2-18)\,\mathbf{k}| = |-20\,\mathbf{k}| = 20$$

28. By plotting the vertices, we can see that the parallelogram is determined by the vectors $\overrightarrow{PQ} = \langle 2, 3, 1\rangle$ and $\overrightarrow{PS} = \langle 4, 2, 5\rangle$. Thus the area of parallelogram $PQRS$ is

$$\left|\overrightarrow{PQ} \times \overrightarrow{PS}\right| = \left\|\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 3 & 1 \\ 4 & 2 & 5 \end{vmatrix}\right\| = |(15-2)\,\mathbf{i} - (10-4)\,\mathbf{j} + (4-12)\,\mathbf{k}|$$

$$= |13\,\mathbf{i} - 6\,\mathbf{j} - 8\,\mathbf{k}| = \sqrt{169+36+64} = \sqrt{269} \approx 16.40$$

29. (a) Because the plane through P, Q, and R contains the vectors $\overrightarrow{PQ}$ and $\overrightarrow{PR}$, a vector orthogonal to these vectors (such as their cross product) is also orthogonal to the plane. $\overrightarrow{PQ} = \langle 2, 1, 3\rangle$ and $\overrightarrow{PR} = \langle 5, 4, 2\rangle$, so

$$\overrightarrow{PQ} \times \overrightarrow{PR} = \langle (1)(2)-(3)(4), (3)(5)-(2)(2), (2)(4)-(1)(5)\rangle = \langle -10, 11, 3\rangle$$

Therefore, $\langle -10, 11, 3\rangle$ (or any nonzero scalar multiple) is orthogonal to the plane through P, Q, and R.

(b) The area of the triangle determined by P, Q, and R is equal to half the area of the parallelogram determined by the three points. Using part (a), the area of the parallelogram is

$$|\overrightarrow{PQ} \times \overrightarrow{PR}| = |\langle -10, 11, 3\rangle| = \sqrt{(-10)^2 + 11^2 + 3^2} = \sqrt{230}$$

So the area of triangle PQR is $\frac{1}{2}\sqrt{230}$.

30. (a) Because the plane through P, Q, and R contains the vectors $\overrightarrow{PQ}$ and $\overrightarrow{PR}$, a vector orthogonal to these vectors (such as their cross product) is also orthogonal to the plane. $\overrightarrow{PQ} = \langle 3, 3, -6\rangle$ and $\overrightarrow{PR} = \langle 2, 3, 1\rangle$, so

$$\overrightarrow{PQ} \times \overrightarrow{PR} = \langle (3)(1)-(-6)(3), (-6)(2)-(3)(1), (3)(3)-(3)(2)\rangle = \langle 21, -15, 3\rangle$$

Therefore, $\langle 21, -15, 3\rangle$ (or any nonzero scalar multiple) is orthogonal to the plane through P, Q, and R.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The area of the triangle determined by P, Q, and R is equal to half the area of the parallelogram determined by the three points. Using part (a), the area of the parallelogram is

$$|\overrightarrow{PQ} \times \overrightarrow{PR}| = |\langle 21, -15, 3\rangle| = \sqrt{21^2 + (-15)^2 + 3^2} = \sqrt{675} = 15\sqrt{3}$$

So the area of triangle PQR is $\frac{15}{2}\sqrt{3}$.

31. (a) Because the plane through P, Q, and R contains the vectors $\overrightarrow{PQ}$ and $\overrightarrow{PR}$, a vector orthogonal to these vectors (such as their cross product) is also orthogonal to the plane. $\overrightarrow{PQ} = \langle -4, 3, 3\rangle$ and $\overrightarrow{PR} = \langle -3, -2, 2\rangle$, so

$$\overrightarrow{PQ} \times \overrightarrow{PR} = \langle (3)(2) - (3)(-2), (3)(-3) - (-4)(2), (-4)(-2) - (3)(-3)\rangle = \langle 12, -1, 17\rangle$$

Therefore, $\langle 12, -1, 17\rangle$ (or any nonzero scalar multiple) is orthogonal to the plane through P, Q, and R.

(b) The area of the triangle determined by P, Q, and R is equal to half the area of the parallelogram determined by the three points. Using part (a), the area of the parallelogram is

$$|\overrightarrow{PQ} \times \overrightarrow{PR}| = |\langle 12, -1, 17\rangle| = \sqrt{12^2 + (-1)^2 + 17^2} = \sqrt{434}$$

So the area of triangle PQR is $\frac{1}{2}\sqrt{434}$.

32. (a) $\overrightarrow{PQ} = \langle -3, 1, -2\rangle$ and $\overrightarrow{PR} = \langle 1, 4, -7\rangle$, so a vector orthogonal to the plane through P, Q, and R is $\overrightarrow{PQ} \times \overrightarrow{PR} = \langle (1)(-7) - (-2)(4), (-2)(1) - (-3)(-7), (-3)(4) - (1)(1)\rangle = \langle 1, -23, -13\rangle$ (or any nonzero scalar multiple).

(b) The area of the parallelogram determined by $\overrightarrow{PQ}$ and $\overrightarrow{PR}$ is $\left|\overrightarrow{PQ} \times \overrightarrow{PR}\right| = |\langle 1, -23, -13\rangle| = \sqrt{1 + 529 + 169} = \sqrt{699}$, so the area of triangle PQR is $\frac{1}{2}\sqrt{699}$.

33. By Equation 14, the volume of the parallelepiped determined by $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$ is the magnitude of their scalar triple product, which is $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} 1 & 2 & 3 \\ -1 & 1 & 2 \\ 2 & 1 & 4 \end{vmatrix} = 1\begin{vmatrix} 1 & 2 \\ 1 & 4 \end{vmatrix} - 2\begin{vmatrix} -1 & 2 \\ 2 & 4 \end{vmatrix} + 3\begin{vmatrix} -1 & 1 \\ 2 & 1 \end{vmatrix} = 1(4-2) - 2(-4-4) + 3(-1-2) = 9.$

Thus the volume of the parallelepiped is 9 cubic units.

34. $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{vmatrix} = 1\begin{vmatrix} 1 & 1 \\ 1 & 1 \end{vmatrix} - 1\begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} + 0\begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} = 0 + 1 + 0 = 1.$

So the volume of the parallelepiped determined by $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$ is 1 cubic unit.

35. $\mathbf{a} = \overrightarrow{PQ} = \langle 4, 2, 2\rangle$, $\mathbf{b} = \overrightarrow{PR} = \langle 3, 3, -1\rangle$, and $\mathbf{c} = \overrightarrow{PS} = \langle 5, 5, 1\rangle$.

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} 4 & 2 & 2 \\ 3 & 3 & -1 \\ 5 & 5 & 1 \end{vmatrix} = 4\begin{vmatrix} 3 & -1 \\ 5 & 1 \end{vmatrix} - 2\begin{vmatrix} 3 & -1 \\ 5 & 1 \end{vmatrix} + 2\begin{vmatrix} 3 & 3 \\ 5 & 5 \end{vmatrix} = 32 - 16 + 0 = 16,$$

so the volume of the parallelepiped is 16 cubic units.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. $\mathbf{a} = \overrightarrow{PQ} = \langle -4, 2, 4\rangle$, $\mathbf{b} = \overrightarrow{PR} = \langle 2, 1, -2\rangle$ and $\mathbf{c} = \overrightarrow{PS} = \langle -3, 4, 1\rangle$.

$$\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c}) = \begin{vmatrix} -4 & 2 & 4 \\ 2 & 1 & -2 \\ -3 & 4 & 1 \end{vmatrix} = -4\begin{vmatrix} 1 & -2 \\ 4 & 1 \end{vmatrix} - 2\begin{vmatrix} 2 & -2 \\ -3 & 1 \end{vmatrix} + 4\begin{vmatrix} 2 & 1 \\ -3 & 4 \end{vmatrix} = -36 + 8 + 44 = 16,$$

so the volume of the parallelepiped is 16 cubic units.

37. $\mathbf{u}\cdot(\mathbf{v}\times\mathbf{w}) = \begin{vmatrix} 1 & 5 & -2 \\ 3 & -1 & 0 \\ 5 & 9 & -4 \end{vmatrix} = 1\begin{vmatrix} -1 & 0 \\ 9 & -4 \end{vmatrix} - 5\begin{vmatrix} 3 & 0 \\ 5 & -4 \end{vmatrix} + (-2)\begin{vmatrix} 3 & -1 \\ 5 & 9 \end{vmatrix} = 4 + 60 - 64 = 0$, which says that the volume of the parallelepiped determined by $\mathbf{u}$, $\mathbf{v}$ and $\mathbf{w}$ is 0, and thus these three vectors are coplanar.

38. $\mathbf{u} = \overrightarrow{AB} = \langle 2, -4, 4\rangle$, $\mathbf{v} = \overrightarrow{AC} = \langle 4, -1, -2\rangle$ and $\mathbf{w} = \overrightarrow{AD} = \langle 2, 3, -6\rangle$.

$\mathbf{u}\cdot(\mathbf{v}\times\mathbf{w}) = \begin{vmatrix} 2 & -4 & 4 \\ 4 & -1 & -2 \\ 2 & 3 & -6 \end{vmatrix} = 2\begin{vmatrix} -1 & -2 \\ 3 & -6 \end{vmatrix} - (-4)\begin{vmatrix} 4 & -2 \\ 2 & -6 \end{vmatrix} + 4\begin{vmatrix} 4 & -1 \\ 2 & 3 \end{vmatrix} = 24 - 80 + 56 = 0$, so the volume of the parallelepiped determined by $\mathbf{u}$, $\mathbf{v}$ and $\mathbf{w}$ is 0, which says these vectors lie in the same plane. Therefore, their initial and terminal points A, B, C and D also lie in the same plane.

39. Using the notation of the text, $|\mathbf{r}| = 0.18$ m, $|\mathbf{F}| = 60$ N, and the angle between $\mathbf{r}$ and $\mathbf{F}$ is $\theta = 70^\circ + 10^\circ = 80^\circ$. (Move $\mathbf{F}$ so that both vectors start from the same point.) Then the magnitude of the torque is $|\boldsymbol{\tau}| = |\mathbf{r}\times\mathbf{F}| = |\mathbf{r}|\,|\mathbf{F}|\sin\theta = (0.18)(60)\sin 80^\circ = 10.8\sin 80^\circ \approx 10.6$ N·m.

40. (a) The position vector from the point P to the handle is $\mathbf{r} = \langle 1, 2\rangle$ and has magnitude $|\mathbf{r}| = \sqrt{1^2 + 2^2} = \sqrt{5}$ ft. Since the force vector $\mathbf{F}$ is parallel to the x-axis, the angle between $\mathbf{r}$ and $\mathbf{F}$ is $\theta = \tan^{-1}\left(\frac{2}{1}\right) \approx 63.43^\circ$ and the magnitude of the torque is $|\boldsymbol{\tau}| = |\mathbf{r}\times\mathbf{F}| = |\mathbf{r}|\,|\mathbf{F}|\sin\theta \approx \left(\sqrt{5}\right)(20)\sin 63.43^\circ \approx 40.0$ ft-lb. (Alternatively, we can observe that $\sin\theta = \frac{2}{\sqrt{5}}$, so $|\mathbf{r}|\,|\mathbf{F}|\sin\theta = \sqrt{5}\cdot 20\cdot\frac{2}{\sqrt{5}} = 40$.)

(b) In this case $\mathbf{r} = \overrightarrow{PQ} = \langle 0.6, 0.6\rangle$, so $|\mathbf{r}| = \sqrt{(0.6)^2 + (0.6)^2} = \sqrt{0.72}$ and $\theta = 45^\circ$. The magnitude of the torque is $|\boldsymbol{\tau}| = |\mathbf{r}|\,|\mathbf{F}|\sin\theta = \left(\sqrt{0.72}\right)(20)\sin 45^\circ = \left(\sqrt{0.72}\right)(20)\cdot\frac{\sqrt{2}}{2} = 10\sqrt{1.44} = 12$ ft-lb.

41. Using the notation of the text, $\mathbf{r} = \langle 0, 0.3, 0\rangle$ (measuring in meters) and $\mathbf{F}$ has direction $\langle 0, 3, -4\rangle$. The angle θ between them can be determined by $\cos\theta = \dfrac{\langle 0, 0.3, 0\rangle\cdot\langle 0, 3, -4\rangle}{|\langle 0, 0.3, 0\rangle|\,|\langle 0, 3, -4\rangle|} \;\Rightarrow\; \cos\theta = \dfrac{0.9}{(0.3)(5)} \;\Rightarrow\; \cos\theta = 0.6 \;\Rightarrow$ $\theta = \cos^{-1}(0.6) \approx 53.1^\circ$. Then $|\boldsymbol{\tau}| = |\mathbf{r}|\,|\mathbf{F}|\sin\theta \;\Rightarrow\; 100 \approx 0.3\,|\mathbf{F}|\sin 53.1^\circ \;\Rightarrow\; |\mathbf{F}| \approx \dfrac{100}{0.3\sin 53.1^\circ} \approx 417$ N.

42. Since $|\mathbf{u}\times\mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}|\sin\theta$, $0 \le \theta \le \pi$, $|\mathbf{u}\times\mathbf{v}|$ achieves its maximum value for $\sin\theta = 1 \;\Rightarrow\; \theta = \frac{\pi}{2}$, in which case $|\mathbf{u}\times\mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}| = 3\,|5\,\mathbf{j}| = 15$. The minimum value is zero, which occurs when $\sin\theta = 0 \;\Rightarrow\; \theta = 0$ or π, so when $\mathbf{u}$, $\mathbf{v}$ are parallel. Thus, when $\mathbf{u}$ points in the same direction as $\mathbf{v}$, so $\mathbf{u} = 3\,\mathbf{j}$, $|\mathbf{u}\times\mathbf{v}| = 0$. As $\mathbf{u}$ rotates counterclockwise,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{u} \times \mathbf{v}$ is directed in the negative z-direction (by the right-hand rule) and the length increases until $\theta = \frac{\pi}{2}$, in which case $\mathbf{u} = -3\,\mathbf{i}$ and $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}| = |-3\,\mathbf{i}|\,|5\,\mathbf{j}| = 15$. As $\mathbf{u}$ rotates to the negative y-axis, $\mathbf{u} \times \mathbf{v}$ remains pointed in the negative z-direction and the length of $\mathbf{u} \times \mathbf{v}$ decreases to 0, after which the direction of $\mathbf{u} \times \mathbf{v}$ reverses to point in the positive z-direction and $|\mathbf{u} \times \mathbf{v}|$ increases. When $\mathbf{u} = 3\,\mathbf{i}$ (so $\theta = \frac{\pi}{2}$), $|\mathbf{u} \times \mathbf{v}|$ again reaches its maximum of 15, after which $|\mathbf{u} \times \mathbf{v}|$ decreases to 0 as $\mathbf{u}$ rotates to the positive y-axis.

43. From Theorem 9 we have $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}|\,|\mathbf{b}| \sin\theta$, where θ is the angle between $\mathbf{a}$ and $\mathbf{b}$, and from Theorem 12.3.3 we have $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}|\,|\mathbf{b}| \cos\theta \quad \Rightarrow \quad |\mathbf{a}|\,|\mathbf{b}| = \dfrac{\mathbf{a} \cdot \mathbf{b}}{\cos\theta}$. Substituting the second equation into the first gives $|\mathbf{a} \times \mathbf{b}| = \dfrac{\mathbf{a} \cdot \mathbf{b}}{\cos\theta} \sin\theta$, so $\dfrac{|\mathbf{a} \times \mathbf{b}|}{\mathbf{a} \cdot \mathbf{b}} = \tan\theta$. Here $|\mathbf{a} \times \mathbf{b}| = |\langle 1, 2, 2\rangle| = \sqrt{1+4+4} = 3$, so $\tan\theta = \dfrac{|\mathbf{a} \times \mathbf{b}|}{\mathbf{a} \cdot \mathbf{b}} = \dfrac{3}{\sqrt{3}} = \sqrt{3} \quad \Rightarrow \quad \theta = 60°$.

44. (a) Let $\mathbf{v} = \langle v_1, v_2, v_3\rangle$. Then

$$\langle 1, 2, 1\rangle \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & 1 \\ v_1 & v_2 & v_3 \end{vmatrix} = \begin{vmatrix} 2 & 1 \\ v_2 & v_3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 1 \\ v_1 & v_3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & 2 \\ v_1 & v_2 \end{vmatrix} \mathbf{k} = (2v_3 - v_2)\,\mathbf{i} - (v_3 - v_1)\,\mathbf{j} + (v_2 - 2v_1)\,\mathbf{k}.$$

If $\langle 1, 2, 1\rangle \times \mathbf{v} = \langle 3, 1, -5\rangle$ then $\langle 2v_3 - v_2, v_1 - v_3, v_2 - 2v_1\rangle = \langle 3, 1, -5\rangle \quad \Leftrightarrow \quad 2v_3 - v_2 = 3$ **(1)**, $v_1 - v_3 = 1$ **(2)**, and $v_2 - 2v_1 = -5$ **(3)**. From **(3)** we have $v_2 = 2v_1 - 5$ and from **(2)** we have $v_3 = v_1 - 1$; substitution into **(1)** gives $2(v_1 - 1) - (2v_1 - 5) = 3 \quad \Rightarrow \quad 3 = 3$, so this is a dependent system. If we let $v_1 = a$ then $v_2 = 2a - 5$ and $v_3 = a - 1$, so $\mathbf{v}$ is any vector of the form $\langle a, 2a - 5, a - 1\rangle$.

(b) If $\langle 1, 2, 1\rangle \times \mathbf{v} = \langle 3, 1, 5\rangle$ then $2v_3 - v_2 = 3$ **(1)**, $v_1 - v_3 = 1$ **(2)**, and $v_2 - 2v_1 = 5$ **(3)**. From **(3)** we have $v_2 = 2v_1 + 5$ and from **(2)** we have $v_3 = v_1 - 1$; substitution into **(1)** gives $2(v_1 - 1) - (2v_1 + 5) = 3 \quad \Rightarrow \quad -7 = 3$, so this is an inconsistent system and has no solution.

Alternatively, if we use matrices to solve the system we could show that the determinant is 0 (and hence the system has no solution).

45. (a)

The distance between a point and a line is the length of the perpendicular from the point to the line, here $\left|\overrightarrow{PS}\right| = d$. But referring to triangle PQS, $d = \left|\overrightarrow{PS}\right| = \left|\overrightarrow{QP}\right| \sin\theta = |\mathbf{b}| \sin\theta$. But θ is the angle between $\overrightarrow{QP} = \mathbf{b}$ and $\overrightarrow{QR} = \mathbf{a}$. Thus by Theorem 9, $\sin\theta = \dfrac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|\,|\mathbf{b}|}$ and so $d = |\mathbf{b}| \sin\theta = \dfrac{|\mathbf{b}|\,|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|}$.

(b) $\mathbf{a} = \overrightarrow{QR} = \langle -1, -2, -1\rangle$ and $\mathbf{b} = \overrightarrow{QP} = \langle 1, -5, -7\rangle$. Then

$$\mathbf{a} \times \mathbf{b} = \langle (-2)(-7) - (-1)(-5), (-1)(1) - (-1)(-7), (-1)(-5) - (-2)(1)\rangle = \langle 9, -8, 7\rangle.$$

Thus the distance is $d = \dfrac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|} = \frac{1}{\sqrt{6}} \sqrt{81 + 64 + 49} = \sqrt{\frac{194}{6}} = \sqrt{\frac{97}{3}}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. (a) The distance between a point and a plane is the length of the perpendicular from the point to the plane, here $\left|\overrightarrow{TP}\right| = d$. But $\overrightarrow{TP}$ is parallel to $\mathbf{b} \times \mathbf{a}$ (because $\mathbf{b} \times \mathbf{a}$ is perpendicular to $\mathbf{b}$ and $\mathbf{a}$) and $d = \left|\overrightarrow{TP}\right| =$ the absolute value of the scalar projection of $\mathbf{c}$ along $\mathbf{b} \times \mathbf{a}$, which is $|\mathbf{c}|\,|\cos\theta|$. (Notice that this is the same setup as the development of the volume of a parallelepiped with $h = |\mathbf{c}|\,|\cos\theta|$). Thus $d = |\mathbf{c}|\,|\cos\theta| = h = V/A$ where $A = |\mathbf{a} \times \mathbf{b}|$, the area of the base. So finally $d = \dfrac{V}{A} = \dfrac{|\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})|}{|\mathbf{a} \times \mathbf{b}|}$.

(b) $\mathbf{a} = \overrightarrow{QR} = \langle -1, 2, 0\rangle$, $\mathbf{b} = \overrightarrow{QS} = \langle -1, 0, 3\rangle$ and $\mathbf{c} = \overrightarrow{QP} = \langle 1, 1, 4\rangle$. Then

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} -1 & 2 & 0 \\ -1 & 0 & 3 \\ 1 & 1 & 4 \end{vmatrix} = (-1)\begin{vmatrix} 0 & 3 \\ 1 & 4 \end{vmatrix} - 2\begin{vmatrix} -1 & 3 \\ 1 & 4 \end{vmatrix} + 0 = 17$$

and
$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 2 & 0 \\ -1 & 0 & 3 \end{vmatrix} = \begin{vmatrix} 2 & 0 \\ 0 & 3 \end{vmatrix}\mathbf{i} - \begin{vmatrix} -1 & 0 \\ -1 & 3 \end{vmatrix}\mathbf{j} + \begin{vmatrix} -1 & 2 \\ -1 & 0 \end{vmatrix}\mathbf{k} = 6\,\mathbf{i} + 3\,\mathbf{j} + 2\,\mathbf{k}$$

Thus $d = \dfrac{|\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})|}{|\mathbf{a} \times \mathbf{b}|} = \dfrac{17}{\sqrt{36+9+4}} = \dfrac{17}{7}$.

47. From Theorem 9 we have $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}|\,|\mathbf{b}|\sin\theta$ so

$$\begin{aligned} |\mathbf{a} \times \mathbf{b}|^2 &= |\mathbf{a}|^2\,|\mathbf{b}|^2 \sin^2\theta = |\mathbf{a}|^2\,|\mathbf{b}|^2\left(1 - \cos^2\theta\right) \\ &= |\mathbf{a}|^2\,|\mathbf{b}|^2 - (|\mathbf{a}|\,|\mathbf{b}|\cos\theta)^2 = |\mathbf{a}|^2\,|\mathbf{b}|^2 - (\mathbf{a} \cdot \mathbf{b})^2 \end{aligned}$$

by Theorem 12.3.3.

48. If $\mathbf{a} + \mathbf{b} + \mathbf{c} = \mathbf{0}$ then $\mathbf{b} = -(\mathbf{a} + \mathbf{c})$, so

$$\begin{aligned} \mathbf{a} \times \mathbf{b} &= \mathbf{a} \times [-(\mathbf{a} + \mathbf{c})] = -[\mathbf{a} \times (\mathbf{a} + \mathbf{c})] && \text{by Property 2 of the cross product (with } c = -1) \\ &= -[(\mathbf{a} \times \mathbf{a}) + (\mathbf{a} \times \mathbf{c})] && \text{by Property 3} \\ &= -[\mathbf{0} + (\mathbf{a} \times \mathbf{c})] = -\mathbf{a} \times \mathbf{c} && \text{by Example 2} \\ &= \mathbf{c} \times \mathbf{a} && \text{by Property 1} \end{aligned}$$

Similarly, $\mathbf{a} = -(\mathbf{b} + \mathbf{c})$ so

$$\begin{aligned} \mathbf{c} \times \mathbf{a} &= \mathbf{c} \times [-(\mathbf{b} + \mathbf{c})] = -[\mathbf{c} \times (\mathbf{b} + \mathbf{c})] \\ &= -[(\mathbf{c} \times \mathbf{b}) + (\mathbf{c} \times \mathbf{c})] = -[(\mathbf{c} \times \mathbf{b}) + \mathbf{0}] \\ &= -\mathbf{c} \times \mathbf{b} = \mathbf{b} \times \mathbf{c} \end{aligned}$$

Thus $\mathbf{a} \times \mathbf{b} = \mathbf{b} \times \mathbf{c} = \mathbf{c} \times \mathbf{a}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49.
$$\begin{aligned}
(\mathbf{a}-\mathbf{b})\times(\mathbf{a}+\mathbf{b}) &= (\mathbf{a}-\mathbf{b})\times\mathbf{a}+(\mathbf{a}-\mathbf{b})\times\mathbf{b} && \text{by Property 3 of the cross product}\\
&= \mathbf{a}\times\mathbf{a}+(-\mathbf{b})\times\mathbf{a}+\mathbf{a}\times\mathbf{b}+(-\mathbf{b})\times\mathbf{b} && \text{by Property 4}\\
&= (\mathbf{a}\times\mathbf{a})-(\mathbf{b}\times\mathbf{a})+(\mathbf{a}\times\mathbf{b})-(\mathbf{b}\times\mathbf{b}) && \text{by Property 2 (with } c=-1)\\
&= \mathbf{0}-(\mathbf{b}\times\mathbf{a})+(\mathbf{a}\times\mathbf{b})-\mathbf{0} && \text{by Example 2}\\
&= (\mathbf{a}\times\mathbf{b})+(\mathbf{a}\times\mathbf{b}) && \text{by Property 1}\\
&= 2(\mathbf{a}\times\mathbf{b})
\end{aligned}$$

50. Let $\mathbf{a}=\langle a_1,a_2,a_3\rangle$, $\mathbf{b}=\langle b_1,b_2,b_3\rangle$ and $\mathbf{c}=\langle c_1,c_2,c_3\rangle$, so $\mathbf{b}\times\mathbf{c}=\langle b_2c_3-b_3c_2,b_3c_1-b_1c_3,b_1c_2-b_2c_1\rangle$ and

$$\begin{aligned}
\mathbf{a}\times(\mathbf{b}\times\mathbf{c}) &= \langle a_2(b_1c_2-b_2c_1)-a_3(b_3c_1-b_1c_3),\, a_3(b_2c_3-b_3c_2)-a_1(b_1c_2-b_2c_1),\\
&\qquad a_1(b_3c_1-b_1c_3)-a_2(b_2c_3-b_3c_2)\rangle\\
&= \langle a_2b_1c_2-a_2b_2c_1-a_3b_3c_1+a_3b_1c_3,\, a_3b_2c_3-a_3b_3c_2-a_1b_1c_2+a_1b_2c_1,\\
&\qquad a_1b_3c_1-a_1b_1c_3-a_2b_2c_3+a_2b_3c_2\rangle\\
&= \langle (a_2c_2+a_3c_3)b_1-(a_2b_2+a_3b_3)c_1,\, (a_1c_1+a_3c_3)b_2-(a_1b_1+a_3b_3)c_2,\\
&\qquad (a_1c_1+a_2c_2)b_3-(a_1b_1+a_2b_2)c_3\rangle\\
(\star)\quad &= \langle (a_2c_2+a_3c_3)b_1-(a_2b_2+a_3b_3)c_1+a_1b_1c_1-a_1b_1c_1,\\
&\qquad (a_1c_1+a_3c_3)b_2-(a_1b_1+a_3b_3)c_2+a_2b_2c_2-a_2b_2c_2,\\
&\qquad (a_1c_1+a_2c_2)b_3-(a_1b_1+a_2b_2)c_3+a_3b_3c_3-a_3b_3c_3\rangle\\
&= \langle (a_1c_1+a_2c_2+a_3c_3)b_1-(a_1b_1+a_2b_2+a_3b_3)c_1,\\
&\qquad (a_1c_1+a_2c_2+a_3c_3)b_2-(a_1b_1+a_2b_2+a_3b_3)c_2,\\
&\qquad (a_1c_1+a_2c_2+a_3c_3)b_3-(a_1b_1+a_2b_2+a_3b_3)c_3\rangle\\
&= (a_1c_1+a_2c_2+a_3c_3)\langle b_1,b_2,b_3\rangle-(a_1b_1+a_2b_2+a_3b_3)\langle c_1,c_2,c_3\rangle\\
&= (\mathbf{a}\cdot\mathbf{c})\mathbf{b}-(\mathbf{a}\cdot\mathbf{b})\mathbf{c}
\end{aligned}$$

$(\star)$ Here we look ahead to see what terms are still needed to arrive at the desired equation. By adding and subtracting the same terms, we don't change the value of the component.

51.
$$\begin{aligned}
&\mathbf{a}\times(\mathbf{b}\times\mathbf{c})+\mathbf{b}\times(\mathbf{c}\times\mathbf{a})+\mathbf{c}\times(\mathbf{a}\times\mathbf{b})\\
&\quad= [(\mathbf{a}\cdot\mathbf{c})\mathbf{b}-(\mathbf{a}\cdot\mathbf{b})\mathbf{c}]+[(\mathbf{b}\cdot\mathbf{a})\mathbf{c}-(\mathbf{b}\cdot\mathbf{c})\mathbf{a}]+[(\mathbf{c}\cdot\mathbf{b})\mathbf{a}-(\mathbf{c}\cdot\mathbf{a})\mathbf{b}] && \text{by Exercise 50}\\
&\quad= (\mathbf{a}\cdot\mathbf{c})\mathbf{b}-(\mathbf{a}\cdot\mathbf{b})\mathbf{c}+(\mathbf{a}\cdot\mathbf{b})\mathbf{c}-(\mathbf{b}\cdot\mathbf{c})\mathbf{a}+(\mathbf{b}\cdot\mathbf{c})\mathbf{a}-(\mathbf{a}\cdot\mathbf{c})\mathbf{b}=\mathbf{0}
\end{aligned}$$

52. Let $\mathbf{c}\times\mathbf{d}=\mathbf{v}$. Then

$$\begin{aligned}
(\mathbf{a}\times\mathbf{b})\cdot(\mathbf{c}\times\mathbf{d}) &= (\mathbf{a}\times\mathbf{b})\cdot\mathbf{v}=\mathbf{a}\cdot(\mathbf{b}\times\mathbf{v}) && \text{by Property 5 of the cross product}\\
&= \mathbf{a}\cdot[\mathbf{b}\times(\mathbf{c}\times\mathbf{d})]=\mathbf{a}\cdot[(\mathbf{b}\cdot\mathbf{d})\mathbf{c}-(\mathbf{b}\cdot\mathbf{c})\mathbf{d}] && \text{by Exercise 50}\\
&= (\mathbf{b}\cdot\mathbf{d})(\mathbf{a}\cdot\mathbf{c})-(\mathbf{b}\cdot\mathbf{c})(\mathbf{a}\cdot\mathbf{d}) && \text{by Properties 3 and 4 of the dot product}\\
&= \begin{vmatrix} \mathbf{a}\cdot\mathbf{c} & \mathbf{b}\cdot\mathbf{c}\\ \mathbf{a}\cdot\mathbf{d} & \mathbf{b}\cdot\mathbf{d}\end{vmatrix}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

53. (a) No. If $\mathbf{a}\cdot\mathbf{b}=\mathbf{a}\cdot\mathbf{c}$, then $\mathbf{a}\cdot(\mathbf{b}-\mathbf{c})=0$, so $\mathbf{a}$ is perpendicular to $\mathbf{b}-\mathbf{c}$, which can happen if $\mathbf{b}\neq\mathbf{c}$. For example, let $\mathbf{a}=\langle 1,1,1\rangle$, $\mathbf{b}=\langle 1,0,0\rangle$ and $\mathbf{c}=\langle 0,1,0\rangle$.

(b) No. If $\mathbf{a}\times\mathbf{b}=\mathbf{a}\times\mathbf{c}$ then $\mathbf{a}\times(\mathbf{b}-\mathbf{c})=\mathbf{0}$, which implies that $\mathbf{a}$ is parallel to $\mathbf{b}-\mathbf{c}$, which of course can happen if $\mathbf{b}\neq\mathbf{c}$.

(c) Yes. Since $\mathbf{a}\cdot\mathbf{c}=\mathbf{a}\cdot\mathbf{b}$, $\mathbf{a}$ is perpendicular to $\mathbf{b}-\mathbf{c}$, by part (a). From part (b), $\mathbf{a}$ is also parallel to $\mathbf{b}-\mathbf{c}$. Thus since $\mathbf{a}\neq\mathbf{0}$ but is both parallel and perpendicular to $\mathbf{b}-\mathbf{c}$, we have $\mathbf{b}-\mathbf{c}=\mathbf{0}$, so $\mathbf{b}=\mathbf{c}$.

54. (a) $\mathbf{k}_i$ is perpendicular to $\mathbf{v}_j$ if $i\neq j$ by the definition of $\mathbf{k}_i$ and Theorem 8.

(b) $\mathbf{k}_1\cdot\mathbf{v}_1=\dfrac{\mathbf{v}_2\times\mathbf{v}_3}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}\cdot\mathbf{v}_1=\dfrac{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}=1$

$$\mathbf{k}_2\cdot\mathbf{v}_2=\frac{\mathbf{v}_3\times\mathbf{v}_1}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}\cdot\mathbf{v}_2=\frac{\mathbf{v}_2\cdot(\mathbf{v}_3\times\mathbf{v}_1)}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}=\frac{(\mathbf{v}_2\times\mathbf{v}_3)\cdot\mathbf{v}_1}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}=1 \qquad \text{[by Property 5 of the cross product]}$$

$$\mathbf{k}_3\cdot\mathbf{v}_3=\frac{(\mathbf{v}_1\times\mathbf{v}_2)\cdot\mathbf{v}_3}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}=\frac{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}=1 \qquad \text{[by Property 5]}$$

(c) $$\mathbf{k}_1\cdot(\mathbf{k}_2\times\mathbf{k}_3)=\mathbf{k}_1\cdot\left(\frac{\mathbf{v}_3\times\mathbf{v}_1}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}\times\frac{\mathbf{v}_1\times\mathbf{v}_2}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}\right)=\frac{\mathbf{k}_1}{[\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)]^2}\cdot[(\mathbf{v}_3\times\mathbf{v}_1)\times(\mathbf{v}_1\times\mathbf{v}_2)]$$

$$=\frac{\mathbf{k}_1}{[\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)]^2}\cdot([(\mathbf{v}_3\times\mathbf{v}_1)\cdot\mathbf{v}_2]\,\mathbf{v}_1-[(\mathbf{v}_3\times\mathbf{v}_1)\cdot\mathbf{v}_1]\,\mathbf{v}_2) \qquad \text{[by Exercise 50]}$$

But $(\mathbf{v}_3\times\mathbf{v}_1)\cdot\mathbf{v}_1=0$ since $\mathbf{v}_3\times\mathbf{v}_1$ is orthogonal to $\mathbf{v}_1$, and $(\mathbf{v}_3\times\mathbf{v}_1)\cdot\mathbf{v}_2=\mathbf{v}_2\cdot(\mathbf{v}_3\times\mathbf{v}_1)=(\mathbf{v}_2\times\mathbf{v}_3)\cdot\mathbf{v}_1=\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)$. Thus

$$\mathbf{k}_1\cdot(\mathbf{k}_2\times\mathbf{k}_3)=\frac{\mathbf{k}_1}{[\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)]^2}\cdot[\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)]\,\mathbf{v}_1=\frac{\mathbf{k}_1\cdot\mathbf{v}_1}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)}=\frac{1}{\mathbf{v}_1\cdot(\mathbf{v}_2\times\mathbf{v}_3)} \qquad \text{[by part (b)]}$$

DISCOVERY PROJECT The Geometry of a Tetrahedron

1. Set up a coordinate system so that vertex S is at the origin, $R=(0,y_1,0)$, $Q=(x_2,y_2,0)$, $P=(x_3,y_3,z_3)$.

Then $\overrightarrow{SR}=\langle 0,y_1,0\rangle$, $\overrightarrow{SQ}=\langle x_2,y_2,0\rangle$, $\overrightarrow{SP}=\langle x_3,y_3,z_3\rangle$, $\overrightarrow{QR}=\langle -x_2,y_1-y_2,0\rangle$, and $\overrightarrow{QP}=\langle x_3-x_2,y_3-y_2,z_3\rangle$. Let

$$\mathbf{v}_S=\overrightarrow{QR}\times\overrightarrow{QP}=(y_1z_3-y_2z_3)\,\mathbf{i}+x_2z_3\,\mathbf{j}+(-x_2y_3-x_3y_1+x_3y_2+x_2y_1)\,\mathbf{k}$$

Then $\mathbf{v}_S$ is an outward normal to the face opposite vertex S. Similarly,

$\mathbf{v}_R=\overrightarrow{SQ}\times\overrightarrow{SP}=y_2z_3\,\mathbf{i}-x_2z_3\,\mathbf{j}+(x_2y_3-x_3y_2)\,\mathbf{k}$, $\mathbf{v}_Q=\overrightarrow{SP}\times\overrightarrow{SR}=-y_1z_3\,\mathbf{i}+x_3y_1\,\mathbf{k}$, and

$\mathbf{v}_P=\overrightarrow{SR}\times\overrightarrow{SQ}=-x_2y_1\,\mathbf{k}\quad\Rightarrow\quad\mathbf{v}_S+\mathbf{v}_R+\mathbf{v}_Q+\mathbf{v}_P=\mathbf{0}$. Now

$$\begin{aligned}|\mathbf{v}_S|&=\text{ area of the parallelogram determined by }\overrightarrow{QR}\text{ and }\overrightarrow{QP}\\&=2\,(\text{area of triangle }RQP)=2|\mathbf{v}_1|\end{aligned}$$

So $\mathbf{v}_S=2\mathbf{v}_1$, and similarly $\mathbf{v}_R=2\mathbf{v}_2$, $\mathbf{v}_Q=2\mathbf{v}_3$, $\mathbf{v}_P=2\mathbf{v}_4$. Thus $\mathbf{v}_1+\mathbf{v}_2+\mathbf{v}_3+\mathbf{v}_4=\mathbf{0}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. (a) Let $S=(x_0,y_0,z_0)$, $R=(x_1,y_1,z_1)$, $Q=(x_2,y_2,z_2)$, $P=(x_3,y_3,z_3)$ be the four vertices. Then

$$\begin{aligned}\text{Volume} &= \tfrac{1}{3}(\text{distance from } S \text{ to plane } RQP)\times(\text{area of triangle } RQP)\\ &= \tfrac{1}{3}\frac{\left|\mathbf{N}\cdot\overrightarrow{SR}\right|}{|\mathbf{N}|}\cdot\tfrac{1}{2}\left|\overrightarrow{RQ}\times\overrightarrow{RP}\right|\end{aligned}$$

where $\mathbf{N}$ is a vector which is normal to the face RQP. Thus $\mathbf{N}=\overrightarrow{RQ}\times\overrightarrow{RP}$. Therefore

$$V=\left|\tfrac{1}{6}\left(\overrightarrow{RQ}\times\overrightarrow{RP}\right)\cdot\overrightarrow{SR}\right|=\frac{1}{6}\left|\begin{vmatrix} x_0-x & y_0-y_1 & z_0-z_1\\ x_2-x_1 & y_2-y_1 & z_2-z_1\\ x_3-x_1 & y_3-y_1 & z_3-z_1\end{vmatrix}\right|$$

(b) Using the formula from part (a), $V=\dfrac{1}{6}\left|\begin{vmatrix} 1-1 & 1-2 & 1-3\\ 1-1 & 1-2 & 2-3\\ 3-1 & -1-2 & 2-3\end{vmatrix}\right|=\dfrac{1}{6}\left|2(1-2)\right|=\dfrac{1}{3}$.

3. We define a vector $\mathbf{v}_1$ to have length equal to the area of the face opposite vertex P, so we can say $|\mathbf{v}_1|=A$, and direction perpendicular to the face and pointing outward, as in Problem 1. Similarly, we define $\mathbf{v}_2$, $\mathbf{v}_3$, and $\mathbf{v}_4$ so that $|\mathbf{v}_2|=B$, $|\mathbf{v}_3|=C$, and $|\mathbf{v}_4|=D$ and with the analogous directions. From Problem 1, we know $\mathbf{v}_1+\mathbf{v}_2+\mathbf{v}_3+\mathbf{v}_4=\mathbf{0}\ \Rightarrow$ $\mathbf{v}_4=-(\mathbf{v}_1+\mathbf{v}_2+\mathbf{v}_3)\ \Rightarrow\ |\mathbf{v}_4|=|-(\mathbf{v}_1+\mathbf{v}_2+\mathbf{v}_3)|=|\mathbf{v}_1+\mathbf{v}_2+\mathbf{v}_3|\ \Rightarrow\ |\mathbf{v}_4|^2=|\mathbf{v}_1+\mathbf{v}_2+\mathbf{v}_3|^2\ \Rightarrow$

$$\begin{aligned}\mathbf{v}_4\cdot\mathbf{v}_4 &= (\mathbf{v}_1+\mathbf{v}_2+\mathbf{v}_3)\cdot(\mathbf{v}_1+\mathbf{v}_2+\mathbf{v}_3)\\ &= \mathbf{v}_1\cdot\mathbf{v}_1+\mathbf{v}_1\cdot\mathbf{v}_2+\mathbf{v}_1\cdot\mathbf{v}_3+\mathbf{v}_2\cdot\mathbf{v}_1+\mathbf{v}_2\cdot\mathbf{v}_2+\mathbf{v}_2\cdot\mathbf{v}_3+\mathbf{v}_3\cdot\mathbf{v}_1+\mathbf{v}_3\cdot\mathbf{v}_2+\mathbf{v}_3\cdot\mathbf{v}_3\end{aligned}$$

Since the vertex S is trirectangular, we know the three faces meeting at S are mutually perpendicular, so the vectors $\mathbf{v}_1$, $\mathbf{v}_2$, $\mathbf{v}_3$ are also mutually perpendicular. Therefore, $\mathbf{v}_i\cdot\mathbf{v}_j=0$ for $i\neq j$ and $i,j\in\{1,2,3\}$. Thus we have $\mathbf{v}_4\cdot\mathbf{v}_4=\mathbf{v}_1\cdot\mathbf{v}_1+\mathbf{v}_2\cdot\mathbf{v}_2+\mathbf{v}_3\cdot\mathbf{v}_3\ \Rightarrow\ |\mathbf{v}_4|^2=|\mathbf{v}_1|^2+|\mathbf{v}_2|^2+|\mathbf{v}_3|^2\ \Rightarrow\ D^2=A^2+B^2+C^2$.

Another method: We introduce a coordinate system, as shown. Recall that the area of the parallelogram spanned by two vectors is equal to the length of their cross product, so since

$\mathbf{u}\times\mathbf{v}=\langle -q,r,0\rangle\times\langle -q,0,p\rangle=\langle pr,pq,qr\rangle$, we have

$|\mathbf{u}\times\mathbf{v}|=\sqrt{(pr)^2+(pq)^2+(qr)^2}$, and therefore

$$\begin{aligned}D^2 &= \left(\tfrac{1}{2}|\mathbf{u}\times\mathbf{v}|\right)^2=\tfrac{1}{4}[(pr)^2+(pq)^2+(qr)^2]\\ &= \left(\tfrac{1}{2}pr\right)^2+\left(\tfrac{1}{2}pq\right)^2+\left(\tfrac{1}{2}qr\right)^2=A^2+B^2+C^2.\end{aligned}$$

A third method: We draw a line from S perpendicular to QR, as shown. Now $D=\frac{1}{2}ch$, so $D^2=\frac{1}{4}c^2h^2$. Substituting $h^2=p^2+k^2$, we get $D^2=\frac{1}{4}c^2\left(p^2+k^2\right)=\frac{1}{4}c^2p^2+\frac{1}{4}c^2k^2$. But $C=\frac{1}{2}ck$, so $D^2=\frac{1}{4}c^2p^2+C^2$. Now substituting $c^2=q^2+r^2$ gives $D^2=\frac{1}{4}p^2q^2+\frac{1}{4}q^2r^2+C^2=A^2+B^2+C^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12.5 Equations of Lines and Planes

1. (a) True; each of the first two lines has a direction vector parallel to the direction vector of the third line, so these vectors are each scalar multiples of the third direction vector. Then the first two direction vectors are also scalar multiples of each other, so these vectors, and hence the two lines, are parallel.

(b) False; for example, the x- and y-axes are both perpendicular to the z-axis, yet the x- and y-axes are not parallel.

(c) True; each of the first two planes has a normal vector parallel to the normal vector of the third plane, so these two normal vectors are parallel to each other and the planes are parallel.

(d) False; for example, the xy- and yz-planes are not parallel, yet they are both perpendicular to the xz-plane.

(e) False; the x- and y-axes are not parallel, yet they are both parallel to the plane $z = 1$.

(f) True; if each line is perpendicular to a plane, then the lines' direction vectors are both parallel to a normal vector for the plane. Thus, the direction vectors are parallel to each other and the lines are parallel.

(g) False; the planes $y = 1$ and $z = 1$ are not parallel, yet they are both parallel to the x-axis.

(h) True; if each plane is perpendicular to a line, then any normal vector for each plane is parallel to a direction vector for the line. Thus, the normal vectors are parallel to each other and the planes are parallel.

(i) True; see Figure 9 and the accompanying discussion.

(j) False; they can be skew, as in Example 3.

(k) True. Consider any normal vector for the plane and any direction vector for the line. If the normal vector is perpendicular to the direction vector, the line and plane are parallel. Otherwise, the vectors meet at an angle θ, $0^\circ \le \theta < 90^\circ$, and the line will intersect the plane at an angle $90^\circ - \theta$.

2. For this line, we have $\mathbf{r}_0 = 4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$ and $\mathbf{v} = 2\mathbf{i} - \mathbf{j} + 6\mathbf{k}$, so a vector equation is $\mathbf{r} = \mathbf{r}_0 + t\mathbf{v} = (4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}) + t(2\mathbf{i} - \mathbf{j} + 6\mathbf{k}) = (4 + 2t)\,\mathbf{i} + (2 - t)\,\mathbf{j} + (-3 + 6t)\,\mathbf{k}$, and parametric equations are $x = 4 + 2t$, $y = 2 - t$, $z = -3 + 6t$.

3. For this line, we have $\mathbf{r}_0 = -\mathbf{i} + 8\mathbf{j} + 7\mathbf{k}$ and $\mathbf{v} = \frac{1}{2}\mathbf{i} + \frac{1}{3}\mathbf{j} + \frac{1}{4}\mathbf{k}$, so a vector equation is $\mathbf{r} = \mathbf{r}_0 + t\mathbf{v} = (-\mathbf{i} + 8\mathbf{j} + 7\mathbf{k}) + t\left(\frac{1}{2}\mathbf{i} + \frac{1}{3}\mathbf{j} + \frac{1}{4}\mathbf{k}\right) = \left(-1 + \frac{1}{2}t\right)\mathbf{i} + \left(8 + \frac{1}{3}t\right)\mathbf{j} + \left(7 + \frac{1}{4}t\right)\mathbf{k}$, and parametric equations are $x = -1 + \frac{1}{2}t$, $y = 8 + \frac{1}{3}t$, $z = 7 + \frac{1}{4}t$.

4. The direction vector for this line is the same as the given line, $\mathbf{v} = -3\mathbf{i} + 4\mathbf{j} + 5\mathbf{k}$. Here $\mathbf{r}_0 = 6\mathbf{i} - 2\mathbf{k}$, so a vector equation is $\mathbf{r} = \mathbf{r}_0 + t\mathbf{v} = (6\,\mathbf{i} - 2\,\mathbf{k}) + t(-3\,\mathbf{i} + 4\,\mathbf{j} + 5\,\mathbf{k}) = (6 - 3t)\,\mathbf{i} + 4t\,\mathbf{j} + (-2 + 5t)\,\mathbf{k}$, and parametric equations are $x = 6 - 3t$, $y = 4t$, $z = -2 + 5t$.

5. A line perpendicular to the given plane has the same direction as a normal vector to the plane, such as $\mathbf{n} = \langle 3, -2, 2\rangle$. So $\mathbf{r}_0 = 5\mathbf{i} + 7\mathbf{j} + \mathbf{k}$ and we can take $\mathbf{v} = 3\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$. Then a vector equation is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{r} = \mathbf{r}_0 + t\mathbf{v} = (5\,\mathbf{i} + 7\,\mathbf{j} + \mathbf{k}) + t(3\,\mathbf{i} - 2\,\mathbf{j} + 2\,\mathbf{k}) = (5 + 3t)\,\mathbf{i} + (7 - 2t)\,\mathbf{j} + (1 + 2t)\,\mathbf{k}$, and parametric equations are $x = 5 + 3t$, $y = 7 - 2t$, $z = 1 + 2t$.

6. The vector $\mathbf{v} = \langle 1 - (-5), 6 - 2, -2 - 5\rangle = \langle 6, 4, -7\rangle$ is parallel to the line. Letting $P_0 = (-5, 2, 5)$, parametric equations are $x = -5 + 6t$, $y = 2 + 4t$, $z = 5 - 7t$ and symmetric equations are $\dfrac{x+5}{6} = \dfrac{y-2}{4} = \dfrac{z-5}{-7}$.

7. The vector $\mathbf{v} = \langle 8 - 0, -1 - 0, 3 - 0\rangle = \langle 8, -1, 3\rangle$ is parallel to the line. Letting $P_0 = (0, 0, 0)$, parametric equations are $x = 8t$, $y = -t$, $z = 3t$ and symmetric equations are $\dfrac{x}{8} = \dfrac{y}{-1} = \dfrac{z}{3}$ or $\dfrac{x}{8} = -y = \dfrac{z}{3}$.

8. The vector $\mathbf{v} = \langle 1.3 - 0.4, 0.8 - (-0.2), -2.3 - 1.1\rangle = \langle 0.9, 1, -3.4\rangle$ is parallel to the line. Letting $P_0 = (0.4, -0.2, 1.1)$, parametric equations are $x = 0.4 + 0.9t$, $y = -0.2 + t$, $z = 1.1 - 3.4t$ and symmetric equations are $\dfrac{x - 0.4}{0.9} = \dfrac{y + 0.2}{1} = \dfrac{z - 1.1}{-3.4}$ or $\dfrac{x - 0.4}{0.9} = y + 0.2 = \dfrac{z - 1.1}{-3.4}$.

9. The vector $\mathbf{v} = \langle -7 - 12, 9 - 9, 11 - (-13)\rangle = \langle -19, 0, 24\rangle$ is parallel to the line. Letting $P_0 = (12, 9, -13)$, parametric equations are $x = 12 - 19t$, $y = 9$, $z = -13 + 24t$ and symmetric equations are $\dfrac{x - 12}{-19} = \dfrac{z + 13}{24}$, $y = 9$. Notice here that the direction number $b = 0$, so rather than writing $\dfrac{y - 9}{0}$ in the symmetric equation, we must write $y = 9$ separately.

10. $\mathbf{v} = (\mathbf{i} + \mathbf{j}) \times (\mathbf{j} + \mathbf{k}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{vmatrix} = \mathbf{i} - \mathbf{j} + \mathbf{k}$ is the direction of the line perpendicular to both $\mathbf{i} + \mathbf{j}$ and $\mathbf{j} + \mathbf{k}$.

With $P_0 = (2, 1, 0)$, parametric equations are $x = 2 + t$, $y = 1 - t$, $z = t$ and symmetric equations are $x - 2 = \dfrac{y - 1}{-1} = z$ or $x - 2 = 1 - y = z$.

11. The given line $\dfrac{x}{2} = \dfrac{y}{3} = \dfrac{z + 1}{1}$ has direction $\mathbf{v} = \langle 2, 3, 1\rangle$. Taking $(-6, 2, 3)$ as P_0, parametric equations are $x = -6 + 2t$, $y = 2 + 3t$, $z = 3 + t$ and symmetric equations are $\dfrac{x + 6}{2} = \dfrac{y - 2}{3} = z - 3$.

12. Setting $z = 0$ we see that $(1, 0, 0)$ satisfies the equations of both planes, so they do in fact have a line of intersection. The line is perpendicular to the normal vectors of both planes, so a direction vector for the line is $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, 2, 3\rangle \times \langle 1, -1, 1\rangle = \langle 5, 2, -3\rangle$. Taking the point $(1, 0, 0)$ as P_0, parametric equations are $x = 1 + 5t$, $y = 2t$, $z = -3t$, and symmetric equations are $\dfrac{x - 1}{5} = \dfrac{y}{2} = \dfrac{z}{-3}$.

13. Direction vectors of the lines are $\mathbf{v}_1 = \langle -2 - (-4), 0 - (-6), -3 - 1\rangle = \langle 2, 6, -4\rangle$ and $\mathbf{v}_2 = \langle 5 - 10, 3 - 18, 14 - 4\rangle = \langle -5, -15, 10\rangle$. Since $\mathbf{v}_2 = -\frac{5}{2}\mathbf{v}_1$, the direction vectors, and thus the lines, are parallel.

14. Direction vectors of the lines are $\mathbf{v}_1 = \langle 1 - (-2), 1 - 4, 1 - 0\rangle = \langle 3, -3, 1\rangle$ and $\mathbf{v}_2 = \langle 3 - 2, -1 - 3, -8 - 4\rangle = \langle 1, -4, -12\rangle$. Since $\mathbf{v}_1 \cdot \mathbf{v}_2 = 3 + 12 - 12 \neq 0$, the direction vectors, and thus the lines, are not perpendicular.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. (a) The line passes through the point $(1, -5, 6)$ and a direction vector for the line is $\langle -1, 2, -3\rangle$, so symmetric equations for the line are $\dfrac{x-1}{-1} = \dfrac{y+5}{2} = \dfrac{z-6}{-3}$.

(b) The line intersects the xy-plane when $z = 0$, so we need $\dfrac{x-1}{-1} = \dfrac{y+5}{2} = \dfrac{0-6}{-3}$ or $\dfrac{x-1}{-1} = 2 \quad\Rightarrow\quad x = -1$, $\dfrac{y+5}{2} = 2 \quad\Rightarrow\quad y = -1$. Thus the point of intersection with the xy-plane is $(-1, -1, 0)$. Similarly for the yz-plane, we need $x = 0 \quad\Rightarrow\quad 1 = \dfrac{y+5}{2} = \dfrac{z-6}{-3} \quad\Rightarrow\quad y = -3$, $z = 3$. Thus the line intersects the yz-plane at $(0, -3, 3)$. For the xz-plane, we need $y = 0 \quad\Rightarrow\quad \dfrac{x-1}{-1} = \dfrac{5}{2} = \dfrac{z-6}{-3} \quad\Rightarrow\quad x = -\frac{3}{2}$, $z = -\frac{3}{2}$. So the line intersects the xz-plane at $\left(-\frac{3}{2}, 0, -\frac{3}{2}\right)$.

16. (a) A vector normal to the plane $x - y + 3z = 7$ is $\mathbf{n} = \langle 1, -1, 3\rangle$, and since the line is to be perpendicular to the plane, $\mathbf{n}$ is also a direction vector for the line. Thus parametric equations of the line are $x = 2 + t$, $y = 4 - t$, $z = 6 + 3t$.

(b) On the xy-plane, $z = 0$. So $z = 6 + 3t = 0 \quad\Rightarrow\quad t = -2$ in the parametric equations of the line, and therefore $x = 0$ and $y = 6$, giving the point of intersection $(0, 6, 0)$. For the yz-plane, $x = 0$ so we get the same point of interesection: $(0, 6, 0)$. For the xz-plane, $y = 0$ which implies $t = 4$, so $x = 6$ and $z = 18$ and the point of intersection is $(6, 0, 18)$.

17. From Equation 4, the line segment from $\mathbf{r}_0 = 6\,\mathbf{i} - \mathbf{j} + 9\,\mathbf{k}$ to $\mathbf{r}_1 = 7\,\mathbf{i} + 6\,\mathbf{j}$ has vector equation

$$\begin{aligned}\mathbf{r}(t) &= (1-t)\,\mathbf{r}_0 + t\,\mathbf{r}_1 = (1-t)(6\,\mathbf{i} - \mathbf{j} + 9\,\mathbf{k}) + t(7\,\mathbf{i} + 6\,\mathbf{j})\\ &= (6\,\mathbf{i} - \mathbf{j} + 9\,\mathbf{k}) - t(6\,\mathbf{i} - \mathbf{j} + 9\,\mathbf{k}) + t(7\,\mathbf{i} + 6\,\mathbf{j})\\ &= (6\,\mathbf{i} - \mathbf{j} + 9\,\mathbf{k}) + t(\,\mathbf{i} + 7\,\mathbf{j} - 9\,\mathbf{k}), \quad 0 \le t \le 1.\end{aligned}$$

18. From Equation 4, the line segment from $\mathbf{r}_0 = -2\,\mathbf{i} + 18\,\mathbf{j} + 31\,\mathbf{k}$ to $\mathbf{r}_1 = 11\,\mathbf{i} - 4\,\mathbf{j} + 48\,\mathbf{k}$ has vector equation

$$\begin{aligned}\mathbf{r}(t) &= (1-t)\,\mathbf{r}_0 + t\,\mathbf{r}_1 = (1-t)(-2\,\mathbf{i} + 18\,\mathbf{j} + 31\,\mathbf{k}) + t(11\,\mathbf{i} - 4\,\mathbf{j} + 48\,\mathbf{k})\\ &= (-2\,\mathbf{i} + 18\,\mathbf{j} + 31\,\mathbf{k}) + t(13\,\mathbf{i} - 22\,\mathbf{j} + 17\,\mathbf{k}), \quad 0 \le t \le 1.\end{aligned}$$

The corresponding parametric equations are $x = -2 + 13t$, $y = 18 - 22t$, $z = 31 + 17t$, $0 \le t \le 1$.

19. Since the direction vectors $\langle 2, -1, 3\rangle$ and $\langle 4, -2, 5\rangle$ are not scalar multiples of each other, the lines aren't parallel. For the lines to intersect, we must be able to find one value of t and one value of s that produce the same point from the respective parametric equations. Thus we need to satisfy the following three equations: $3 + 2t = 1 + 4s$, $4 - t = 3 - 2s$, $1 + 3t = 4 + 5s$. Solving the last two equations we get $t = 1$, $s = 0$ and checking, we see that these values don't satisfy the first equation. Thus the lines aren't parallel and don't intersect, so they must be skew lines.

20. Since the direction vectors are $\mathbf{v}_1 = \langle -12, 9, -3\rangle$ and $\mathbf{v}_2 = \langle 8, -6, 2\rangle$, we have $\mathbf{v}_1 = -\frac{3}{2}\mathbf{v}_2$ so the lines are parallel.

21. Since the direction vectors $\langle 1, -2, -3\rangle$ and $\langle 1, 3, -7\rangle$ aren't scalar multiples of each other, the lines aren't parallel. Parametric equations of the lines are L_1: $x = 2 + t$, $y = 3 - 2t$, $z = 1 - 3t$ and L_2: $x = 3 + s$, $y = -4 + 3s$, $z = 2 - 7s$. Thus, for the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

lines to intersect, the three equations $2+t=3+s$, $3-2t=-4+3s$, and $1-3t=2-7s$ must be satisfied simultaneously. Solving the first two equations gives $t=2$, $s=1$ and checking, we see that these values do satisfy the third equation, so the lines intersect when $t=2$ and $s=1$, that is, at the point $(4,-1,-5)$.

22. The direction vectors $\langle 1,-1,3\rangle$ and $\langle 2,-2,7\rangle$ are not parallel, so neither are the lines. Parametric equations for the lines are L_1: $x=t$, $y=1-t$, $z=2+3t$ and L_2: $x=2+2s$, $y=3-2s$, $z=7s$. Thus, for the lines to interesect, the three equations $t=2+2s$, $1-t=3-2s$, and $2+3t=7s$ must be satisfied simultaneously. Solving the last two equations gives $t=-10$, $s=-4$ and checking, we see that these values don't satisfy the first equation. Thus the lines aren't parallel and don't intersect, so they must be skew.

23. $5\mathbf{i}+4\mathbf{j}+6\mathbf{k}$ is a normal vector to the plane. $(3,2,1)$ is a point on the plane. Setting $a=5$, $b=4$, $c=6$ and $x_0=3$, $y_0=2$, $z_0=1$ in Equation 7 gives $5(x-3)+4(y-2)+6(z-1)=0$, or $5x+4y+6z=29$, as an equation of the plane.

24. $\langle 6,1,-1\rangle$ is a normal vector to the plane. $(-3,4,2)$ is a point on the plane. Setting $a=6$, $b=1$, $c=-1$ and $x_0=-3$, $y_0=4$, $z_0=2$ in Equation 7 gives $6(x+3)+1(y-4)-(z-2)=0$, or $6x+y-z=-16$, as an equation of the plane.

25. Since the plane is perpendicular to the vector $-\mathbf{i}+2\,\mathbf{j}+3\,\mathbf{k}$, we can take $\langle -1,2,3\rangle$ as a normal vector to the plane. $(5,-2,4)$ is a point on the plane. Setting $a=-1$, $b=2$, $c=3$ and $x_0=5$, $y_0=-2$, $z_0=4$ in Equation 7 gives $-(x-5)+2(y+2)+3(z-4)=0$, or $-x+2y+3z=3$, as an equation of the plane.

26. Since the line is perpendicular to the plane, its direction vector, $\langle -8,-7,2\rangle$, is a normal vector to the plane. $(0,0,0)$ is a point on the plane. Setting $a=-8$, $b=-7$, $c=2$ and $x_0=0$, $y_0=0$, $z_0=0$ in Equation 7 gives $-8(x-0)-7(y-0)+2(z-0)=0$, or $-8x-7y+2z=0$, as an equation of the plane.

27. Since the line is perpendicular to the plane, its direction vector, $\langle 4,-1,5\rangle$, is a normal vector to the plane. $(1,3,-1)$ is a point on the plane. Setting $a=4$, $b=-1$, $c=5$ and $x_0=1$, $y_0=3$, $z_0=-1$ in Equation 7 gives $4(x-1)-1(y-3)+5(z+1)=0$, or $4x-y+5z=-4$, as an equation of the plane.

28. Since the two planes are parallel, they will have the same normal vectors. The plane is $z=2x-3y \quad\Leftrightarrow\quad 2x-3y-z=0$, so we can take $\mathbf{n}=\langle 2,-3,-1\rangle$, and an equation of the plane is $2(x-9)-3(y+4)-1(z+5)=0$, or $2x-3y-z=35$.

29. Since the two planes are parallel, they will have the same normal vectors. The plane is $2x-y+3z=1$, so we can take $\mathbf{n}=\langle 2,-1,3\rangle$, and an equation of the plane is $2(x-2.1)-1(y-1.7)+3(z+0.9)=0$, or $2x-y+3z=-0.2$, or $10x-5y+15z=-1$.

30. First, a normal vector for the plane $5x+2y+z=1$ is $\mathbf{n}=\langle 5,2,1\rangle$. A direction vector for the line is $\mathbf{v}=\langle 1,-1,-3\rangle$, and since $\mathbf{n}\cdot\mathbf{v}=0$ we know the line is perpendicular to $\mathbf{n}$ and hence parallel to the plane. Thus, there is a parallel plane which contains the line. By putting $t=0$, we know that the point $(1,2,4)$ is on the line and hence the new plane. We can use the same normal vector $\mathbf{n}=\langle 5,2,1\rangle$, so an equation of the plane is $5(x-1)+2(y-2)+1(z-4)=0$ or $5x+2y+z=13$.

31. The vector from $(0,1,1)$ to $(1,0,1)$, namely $\mathbf{a}=\langle 1-0,0-1,1-1\rangle=\langle 1,-1,0\rangle$, and the vector from $(0,1,1)$ to $(1,1,0)$, $\mathbf{b}=\langle 1-0,1-1,0-1\rangle=\langle 1,0,-1\rangle$, both lie in the plane, so $\mathbf{a}\times\mathbf{b}$ is a normal vector to the plane. Thus, we can take

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle (-1)((-1) - (0)(0), (0)(1) - (1)(-1), (1)(0) - (-1)(1) \rangle = \langle 1, 1, 1 \rangle$. If P_0 is the point $(0, 1, 1)$, an equation of the plane is $1(x - 0) + 1(y - 1) + 1(z - 1) = 0$ or $x + y + z = 2$.

32. Here the vectors $\mathbf{a} = \langle 3, -2, 1 \rangle$ and $\mathbf{b} = \langle 1, 1, 1 \rangle$ lie in the plane, so $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle (-2)(1) - (1)(1), (1)(1) - (3)(1), (3)(1) - (-2)(1) \rangle = \langle -3, -2, 5 \rangle$ is a normal vector to the plane. We can take the origin as P_0, so an equation of the plane is $-3(x - 0) - 2(y - 0) + 5(z - 0) = 0$ or $-3x - 2y + 5z = 0$ or $3x + 2y - 5z = 0$.

33. Here the vectors $\mathbf{a} = \langle 3 - 2, -8 - 1, 6 - 2 \rangle = \langle 1, -9, 4 \rangle$ and $\mathbf{b} = \langle -2 - 2, -3 - 1, 1 - 2 \rangle = \langle -4, -4, -1 \rangle$ lie in the plane, so a normal vector to the plane is $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle 9 + 16, -16 + 1, -4 - 36 \rangle = \langle 25, -15, -40 \rangle$ and an equation of the plane is $25(x - 2) - 15(y - 1) - 40(z - 2) = 0$ or $25x - 15y - 40z = -45$ or $5x - 3y - 8z = -9$.

34. The vectors $\mathbf{a} = \langle -2 - 3, -2 - 0, 3 - (-1) \rangle = \langle -5, -2, 4 \rangle$ and $\mathbf{b} = \langle 7 - 3, 1 - 0, -4 - (-1) \rangle = \langle 4, 1, -3 \rangle$ lie in the plane, so a normal vector to the plane is $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle 6 - 4, 16 - 15, -5 + 8 \rangle = \langle 2, 1, 3 \rangle$ and an equation of the plane is $2(x - 3) + 1(y - 0) + 3[z - (-1)] = 0$ or $2x + y + 3z = 3$.

35. If we first find two nonparallel vectors in the plane, their cross product will be a normal vector to the plane. Since the given line lies in the plane, its direction vector $\mathbf{a} = \langle -1, 2, -3 \rangle$ is one vector in the plane. We can verify that the given point $(3, 5, -1)$ does not lie on this line, so to find another nonparallel vector $\mathbf{b}$ which lies in the plane, we can pick any point on the line and find a vector connecting the points. If we put $t = 0$, we see that $(4, -1, 0)$ is on the line, so $\mathbf{b} = \langle 4 - 3, -1 - 5, 0 - (-1) \rangle = \langle 1, -6, 1 \rangle$ and $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle 2 - 18, -3 + 1, 6 - 2 \rangle = \langle -16, -2, 4 \rangle$. Thus, an equation of the plane is $-16(x - 3) - 2(y - 5) + 4[z - (-1)] = 0$ or $-16x - 2y + 4z = -62$ or $8x + y - 2z = 31$.

36. Since the line $\dfrac{x}{3} = \dfrac{y + 4}{1} = \dfrac{z}{2}$ lies in the plane, its direction vector $\mathbf{a} = \langle 3, 1, 2 \rangle$ is parallel to the plane. The point $(0, -4, 0)$ is on the line (put $t = 0$ in the corresponding parametric equations), and we can verify that the given point $(6, -1, 3)$ in the plane is not on the line. The vector connecting these two points, $\mathbf{b} = \langle 6, 3, 3 \rangle$, is therefore parallel to the plane, but not parallel to $\mathbf{a}$. Then $\mathbf{a} \times \mathbf{b} = \langle 3 - 6, 12 - 9, 9 - 6 \rangle = \langle -3, 3, 3 \rangle$ is a normal vector to the plane, and an equation of the plane is $-3(x - 0) + 3[y - (-4)] + 3(z - 0) = 0$ or $-3x + 3y + 3z = -12$ or $x - y - z = 4$.

37. Normal vectors for the given planes are $\mathbf{n}_1 = \langle 1, 2, 3 \rangle$ and $\mathbf{n}_2 = \langle 2, -1, 1 \rangle$. A direction vector, then, for the line of intersection is $\mathbf{a} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 2 + 3, 6 - 1, -1 - 4 \rangle = \langle 5, 5, -5 \rangle$, and $\mathbf{a}$ is parallel to the desired plane. Another vector parallel to the plane is the vector connecting any point on the line of intersection to the given point $(3, 1, 4)$ in the plane. Setting $z = 0$, the equations of the planes reduce to $x + 2y = 1$ and $2x - y = -3$ with simultaneous solution $x = -1$ and $y = 1$. So a point on the line is $(-1, 1, 0)$ and another vector parallel to the plane is $\mathbf{b} = \langle 3 - (-1), 1 - 1, 4 - 0 \rangle = \langle 4, 0, 4 \rangle$. Then a normal vector to the plane is $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle 20 - 0, -20 - 20, 0 - 20 \rangle = \langle 20, -40, -20 \rangle$. Equivalently, we can take $\langle 1, -2, -1 \rangle$ as a normal vector, and an equation of the plane is $1(x - 3) - 2(y - 1) - 1(z - 4) = 0$ or $x - 2y - z = -3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. The points $(0, -2, 5)$ and $(-1, 3, 1)$ lie in the desired plane, so the vector $\mathbf{v}_1 = \langle -1, 5, -4 \rangle$ connecting them is parallel to the plane. The desired plane is perpendicular to the plane $2z = 5x + 4y$ or $5x + 4y - 2z = 0$ and for perpendicular planes, a normal vector for one plane is parallel to the other plane, so $\mathbf{v}_2 = \langle 5, 4, -2 \rangle$ is also parallel to the desired plane.
A normal vector to the desired plane is $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle -10 + 16, -20 - 2, -4 - 25 \rangle = \langle 6, -22, -29 \rangle$.
Taking $(x_0, y_0, z_0) = (0, -2, 5)$, the equation we are looking for is $6(x - 0) - 22(y + 2) - 29(z - 5) = 0$ or $6x - 22y - 29z = -101$.

39. If a plane is perpendicular to two other planes, its normal vector is perpendicular to the normal vectors of the other two planes. Thus $\langle 2, 1, -2 \rangle \times \langle 1, 0, 3 \rangle = \langle 3 - 0, -2 - 6, 0 - 1 \rangle = \langle 3, -8, -1 \rangle$ is a normal vector to the desired plane. The point $(1, 5, 1)$ lies on the plane, so an equation is $3(x - 1) - 8(y - 5) - (z - 1) = 0$ or $3x - 8y - z = -38$.

40. $\mathbf{n}_1 = \langle 1, 0, -1 \rangle$ and $\mathbf{n}_2 = \langle 0, 1, 2 \rangle$. Setting $z = 0$, it is easy to see that $(1, 3, 0)$ is a point on the line of intersection of $x - z = 1$ and $y + 2z = 3$. The direction of this line is $\mathbf{v}_1 = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, -2, 1 \rangle$. A second vector parallel to the desired plane is $\mathbf{v}_2 = \langle 1, 1, -2 \rangle$, since it is perpendicular to $x + y - 2z = 1$. Therefore, a normal of the plane in question is $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 4 - 1, 1 + 2, 1 + 2 \rangle = \langle 3, 3, 3 \rangle$, or we can use $\langle 1, 1, 1 \rangle$. Taking $(x_0, y_0, z_0) = (1, 3, 0)$, the equation we are looking for is $(x - 1) + (y - 3) + z = 0 \quad \Leftrightarrow \quad x + y + z = 4$.

41. To find the x-intercept we set $y = z = 0$ in the equation $2x + 5y + z = 10$ and obtain $2x = 10 \quad \Rightarrow \quad x = 5$ so the x-intercept is $(5, 0, 0)$. When $x = z = 0$ we get $5y = 10 \quad \Rightarrow \quad y = 2$, so the y-intercept is $(0, 2, 0)$. Setting $x = y = 0$ gives $z = 10$, so the z-intercept is $(0, 0, 10)$ and we graph the portion of the plane that lies in the first octant.

42. To find the x-intercept we set $y = z = 0$ in the equation $3x + y + 2z = 6$ and obtain $3x = 6 \quad \Rightarrow \quad x = 2$ so the x-intercept is $(2, 0, 0)$. When $x = z = 0$ we get $y = 6$ so the y-intercept is $(0, 6, 0)$. Setting $x = y = 0$ gives $2z = 6 \quad \Rightarrow \quad z = 3$, so the z-intercept is $(0, 0, 3)$. The figure shows the portion of the plane that lies in the first octant.

43. Setting $y = z = 0$ in the equation $6x - 3y + 4z = 6$ gives $6x = 6 \quad \Rightarrow \quad x = 1$, when $x = z = 0$ we have $-3y = 6 \quad \Rightarrow \quad y = -2$, and $x = y = 0$ implies $4z = 6 \quad \Rightarrow \quad z = \frac{3}{2}$, so the intercepts are $(1, 0, 0)$, $(0, -2, 0)$, and $(0, 0, \frac{3}{2})$. The figure shows the portion of the plane cut off by the coordinate planes.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. Setting $y = z = 0$ in the equation $6x + 5y - 3z = 15$ gives $6x = 15 \quad \Rightarrow$ $x = \frac{5}{2}$, when $x = z = 0$ we have $5y = 15 \quad \Rightarrow \quad y = 3$, and $x = y = 0$ implies $-3z = 15 \quad \Rightarrow \quad z = -5$, so the intercepts are $\left(\frac{5}{2}, 0, 0\right)$, $(0, 3, 0)$, and $(0, 0, -5)$. The figure shows the portion of the plane cut off by the coordinate planes.

45. Substitute the parametric equations of the line into the equation of the plane: $x + 2y - z = 7 \quad \Rightarrow$ $(2 - 2t) + 2(3t) - (1 + t) = 7 \quad \Rightarrow \quad 3t + 1 = 7 \quad \Rightarrow \quad t = 2$. Therefore, the point of intersection of the line and the plane is given by $x = 2 - 2(2) = -2$, $y = 3(2) = 6$, and $z = 1 + 2 = 3$, that is, the point $(-2, 6, 3)$.

46. Substitute the parametric equations of the line into the equation of the plane: $3(t - 1) - (1 + 2t) + 2(3 - t) = 5 \quad \Rightarrow$ $-t + 2 = 5 \quad \Rightarrow \quad t = -3$. Therefore, the point of intersection of the line and the plane is given by $x = -3 - 1 = -4$, $y = 1 + 2(-3) = -5$, and $z = 3 - (-3) = 6$, that is, the point $(-4, -5, 6)$.

47. Parametric equations for the line are $x = \frac{1}{5}t$, $y = 2t$, $z = t - 2$ and substitution into the equation of the plane gives $10\left(\frac{1}{5}t\right) - 7(2t) + 3(t - 2) + 24 = 0 \quad \Rightarrow \quad -9t + 18 = 0 \quad \Rightarrow \quad t = 2$. Thus $x = \frac{1}{5}(2) = \frac{2}{5}$, $y = 2(2) = 4$, $z = 2 - 2 = 0$ and the point of intersection is $\left(\frac{2}{5}, 4, 0\right)$.

48. A direction vector for the line through $(-3, 1, 0)$ and $(-1, 5, 6)$ is $\mathbf{v} = \langle 2, 4, 6\rangle$ and, taking $P_0 = (-3, 1, 0)$, parametric equations for the line are $x = -3 + 2t$, $y = 1 + 4t$, $z = 6t$. Substitution of the parametric equations into the equation of the plane gives $2(-3 + 2t) + (1 + 4t) - (6t) = -2 \quad \Rightarrow \quad 2t - 5 = -2 \quad \Rightarrow \quad t = \frac{3}{2}$. Then $x = -3 + 2\left(\frac{3}{2}\right) = 0$, $y = 1 + 4\left(\frac{3}{2}\right) = 7$, and $z = 6\left(\frac{3}{2}\right) = 9$, and the point of intersection is $(0, 7, 9)$.

49. Setting $x = 0$, we see that $(0, 1, 0)$ satisfies the equations of both planes, so that they do in fact have a line of intersection. $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, 1, 1\rangle \times \langle 1, 0, 1\rangle = \langle 1, 0, -1\rangle$ is the direction of this line. Therefore, direction numbers of the intersecting line are 1, 0, -1.

50. The angle between the two planes is the same as the angle between their normal vectors. The normal vectors of the two planes are $\langle 1, 1, 1\rangle$ and $\langle 1, 2, 3\rangle$. The cosine of the angle θ between these two planes is

$$\cos\theta = \frac{\langle 1, 1, 1\rangle \cdot \langle 1, 2, 3\rangle}{|\langle 1, 1, 1\rangle|\,|\langle 1, 2, 3\rangle|} = \frac{1 + 2 + 3}{\sqrt{1 + 1 + 1}\sqrt{1 + 4 + 9}} = \frac{6}{\sqrt{42}} = \sqrt{\frac{6}{7}}.$$

51. Normal vectors for the planes are $\mathbf{n}_1 = \langle 1, 4, -3\rangle$ and $\mathbf{n}_2 = \langle -3, 6, 7\rangle$. The normals aren't parallel (they are not scalar multiples of each other), so neither are the planes. But $\mathbf{n}_1 \cdot \mathbf{n}_2 = -3 + 24 - 21 = 0$, so the normals, and thus the planes, are perpendicular.

52. Normal vectors for the planes are $\mathbf{n}_1 = \langle 9, -3, 6\rangle$ and $\mathbf{n}_2 = \langle 6, -2, 4\rangle$ (the plane's equation is $6x - 2y + 4z = 0$). Since $\mathbf{n}_1 = \frac{3}{2}\mathbf{n}_2$, the normals, and thus the planes, are parallel.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

53. Normal vectors for the planes are $\mathbf{n}_1 = \langle 1, 2, -1\rangle$ and $\mathbf{n}_2 = \langle 2, -2, 1\rangle$. The normals are not parallel (they are not scalar multiples of each other), so neither are the planes. Furthermore, $\mathbf{n}_1 \cdot \mathbf{n}_2 = 2 - 4 - 1 = -3 \neq 0$, so the planes aren't perpendicular. The angle between the planes is the same as the angle between the normals, given by

$$\cos\theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1|\,|\mathbf{n}_2|} = \frac{-3}{\sqrt{6}\sqrt{9}} = -\frac{1}{\sqrt{6}} \quad \Rightarrow \quad \theta = \cos^{-1}\left(-\frac{1}{\sqrt{6}}\right) \approx 114.1^\circ.$$

54. Normal vectors for the planes are $\mathbf{n}_1 = \langle 1, -1, 3\rangle$ and $\mathbf{n}_2 = \langle 3, 1, -1\rangle$. The normals are not parallel, so neither are the planes. Since $\mathbf{n}_1 \cdot \mathbf{n}_2 = 3 - 1 - 3 = -1 \neq 0$, the planes aren't perpendicular. The angle between the planes is given by

$$\cos\theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1|\,|\mathbf{n}_2|} = \frac{-1}{\sqrt{11}\sqrt{11}} = -\frac{1}{11} \quad \Rightarrow \quad \theta = \cos^{-1}\left(-\tfrac{1}{11}\right) \approx 95.2^\circ.$$

55. The planes are $2x - 3y - z = 0$ and $4x - 6y - 2z = 3$ with normal vectors $\mathbf{n}_1 = \langle 2, -3, -1\rangle$ and $\mathbf{n}_2 = \langle 4, -6, -2\rangle$. Since $\mathbf{n}_2 = 2\mathbf{n}_1$, the normals, and thus the planes, are parallel.

56. The normals are $\mathbf{n}_1 = \langle 5, 2, 3\rangle$ and $\mathbf{n}_2 = \langle 4, -1, -6\rangle$ which are not scalar multiples of each other, so the planes aren't parallel. Since $\mathbf{n}_1 \cdot \mathbf{n}_2 = 20 - 2 - 18 = 0$, the normals, and thus the planes, are perpendicular.

57. (a) To find a point on the line of intersection, set one of the variables equal to a constant, say $z = 0$. (This will fail if the line of intersection does not cross the xy-plane; in that case, try setting x or y equal to 0.) The equations of the two planes reduce to $x + y = 1$ and $x + 2y = 1$. Solving these two equations gives $x = 1$, $y = 0$. Thus a point on the line is $(1, 0, 0)$. A vector $\mathbf{v}$ in the direction of this intersecting line is perpendicular to the normal vectors of both planes, so we can take $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, 1, 1\rangle \times \langle 1, 2, 2\rangle = \langle 2 - 2, 1 - 2, 2 - 1\rangle = \langle 0, -1, 1\rangle$. By Equations 2, parametric equations for the line are $x = 1$, $y = -t$, $z = t$.

(b) The angle between the planes satisfies $\cos\theta = \dfrac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1|\,|\mathbf{n}_2|} = \dfrac{1 + 2 + 2}{\sqrt{3}\sqrt{9}} = \dfrac{5}{3\sqrt{3}}$. Therefore $\theta = \cos^{-1}\left(\dfrac{5}{3\sqrt{3}}\right) \approx 15.8^\circ$.

58. (a) If we set $z = 0$ then the equations of the planes reduce to $3x - 2y = 1$ and $2x + y = 3$ and solving these two equations gives $x = 1$, $y = 1$. Thus a point on the line of intersection is $(1, 1, 0)$. A vector $\mathbf{v}$ in the direction of this intersecting line is perpendicular to the normal vectors of both planes, so let $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 3, -2, 1\rangle \times \langle 2, 1, -3\rangle = \langle 5, 11, 7\rangle$. By Equations 2, parametric equations for the line are $x = 1 + 5t$, $y = 1 + 11t$, $z = 7t$.

(b) $\cos\theta = \dfrac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1|\,|\mathbf{n}_2|} = \dfrac{6 - 2 - 3}{\sqrt{14}\sqrt{14}} = \dfrac{1}{14} \quad \Rightarrow \quad \theta = \cos^{-1}\left(\tfrac{1}{14}\right) \approx 85.9^\circ$.

59. Setting $z = 0$, the equations of the two planes become $5x - 2y = 1$ and $4x + y = 6$. Solving these two equations gives $x = 1$, $y = 2$ so a point on the line of intersection is $(1, 2, 0)$. A vector $\mathbf{v}$ in the direction of this intersecting line is perpendicular to the normal vectors of both planes. So we can use $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 5, -2, -2\rangle \times \langle 4, 1, 1\rangle = \langle 0, -13, 13\rangle$ or equivalently we can take $\mathbf{v} = \langle 0, -1, 1\rangle$, and symmetric equations for the line are $x = 1$, $\dfrac{y - 2}{-1} = \dfrac{z}{1}$ or $x = 1$, $y - 2 = -z$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. If we set $z = 0$ then the equations of the planes reduce to $2x - y - 5 = 0$ and $4x + 3y - 5 = 0$ and solving these two equations gives $x = 2$, $y = -1$. Thus a point on the line of intersection is $(2, -1, 0)$. A vector $\mathbf{v}$ in the direction of this intersecting line is perpendicular to the normal vectors of both planes, so take $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 2, -1, -1\rangle \times \langle 4, 3, -1\rangle = \langle 4, -2, 10\rangle$ or equivalently we can take $\mathbf{v} = \langle 2, -1, 5\rangle$. Symmetric equations for the line are $\dfrac{x-2}{2} = \dfrac{y+1}{-1} = \dfrac{z}{5}$.

61. The distance from a point (x, y, z) to $(1, 0, -2)$ is $d_1 = \sqrt{(x-1)^2 + y^2 + (z+2)^2}$ and the distance from (x, y, z) to $(3, 4, 0)$ is $d_2 = \sqrt{(x-3)^2 + (y-4)^2 + z^2}$. The plane consists of all points (x, y, z) where $d_1 = d_2 \quad \Rightarrow \quad d_1^2 = d_2^2 \quad \Leftrightarrow$ $(x-1)^2 + y^2 + (z+2)^2 = (x-3)^2 + (y-4)^2 + z^2 \quad \Leftrightarrow$
$x^2 - 2x + y^2 + z^2 + 4z + 5 = x^2 - 6x + y^2 - 8y + z^2 + 25 \quad \Leftrightarrow \quad 4x + 8y + 4z = 20$ so an equation for the plane is $4x + 8y + 4z = 20$ or equivalently $x + 2y + z = 5$.
Alternatively, you can argue that the segment joining points $(1, 0, -2)$ and $(3, 4, 0)$ is perpendicular to the plane and the plane includes the midpoint of the segment.

62. The distance from a point (x, y, z) to $(2, 5, 5)$ is $d_1 = \sqrt{(x-2)^2 + (y-5)^2 + (z-5)^2}$ and the distance from (x, y, z) to $(-6, 3, 1)$ is $d_2 = \sqrt{(x+6)^2 + (y-3)^2 + (z-1)^2}$. The plane consists of all points (x, y, z) where $d_1 = d_2 \quad \Rightarrow$ $d_1^2 = d_2^2 \quad \Leftrightarrow \quad (x-2)^2 + (y-5)^2 + (z-5)^2 = (x+6)^2 + (y-3)^2 + (z-1)^2 \quad \Leftrightarrow$
$x^2 - 4x + y^2 - 10y + z^2 - 10z + 54 = x^2 + 12x + y^2 - 6y + z^2 - 2z + 46 \quad \Leftrightarrow \quad 16x + 4y + 8z = 8$ so an equation for the plane is $16x + 4y + 8z = 8$ or equivalently $4x + y + 2z = 2$.

63. The plane contains the points $(a, 0, 0)$, $(0, b, 0)$ and $(0, 0, c)$. Thus the vectors $\mathbf{a} = \langle -a, b, 0\rangle$ and $\mathbf{b} = \langle -a, 0, c\rangle$ lie in the plane, and $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle bc - 0, 0 + ac, 0 + ab\rangle = \langle bc, ac, ab\rangle$ is a normal vector to the plane. The equation of the plane is therefore $bcx + acy + abz = abc + 0 + 0$ or $bcx + acy + abz = abc$. Notice that if $a \neq 0$, $b \neq 0$ and $c \neq 0$ then we can rewrite the equation as $\dfrac{x}{a} + \dfrac{y}{b} + \dfrac{z}{c} = 1$. This is a good equation to remember!

64. (a) For the lines to intersect, we must be able to find one value of t and one value of s satisfying the three equations $1 + t = 2 - s$, $1 - t = s$ and $2t = 2$. From the third we get $t = 1$, and putting this in the second gives $s = 0$. These values of s and t do satisfy the first equation, so the lines intersect at the point $P_0 = (1 + 1, 1 - 1, 2(1)) = (2, 0, 2)$.

(b) The direction vectors of the lines are $\langle 1, -1, 2\rangle$ and $\langle -1, 1, 0\rangle$, so a normal vector for the plane is $\langle -1, 1, 0\rangle \times \langle 1, -1, 2\rangle = \langle 2, 2, 0\rangle$ and it contains the point $(2, 0, 2)$. Then an equation of the plane is $2(x - 2) + 2(y - 0) + 0(z - 2) = 0 \quad \Leftrightarrow \quad x + y = 2$.

65. Two vectors which are perpendicular to the required line are the normal of the given plane, $\langle 1, 1, 1\rangle$, and a direction vector for the given line, $\langle 1, -1, 2\rangle$. So a direction vector for the required line is $\langle 1, 1, 1\rangle \times \langle 1, -1, 2\rangle = \langle 3, -1, -2\rangle$. Thus L is given by $\langle x, y, z\rangle = \langle 0, 1, 2\rangle + t\langle 3, -1, -2\rangle$, or in parametric form, $x = 3t$, $y = 1 - t$, $z = 2 - 2t$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

66. Let L be the given line. Then $(1,1,0)$ is the point on L corresponding to $t=0$. L is in the direction of $\mathbf{a}=\langle 1,-1,2\rangle$ and $\mathbf{b}=\langle -1,0,2\rangle$ is the vector joining $(1,1,0)$ and $(0,1,2)$. Then

$\mathbf{b}-\operatorname{proj}_{\mathbf{a}}\mathbf{b}=\langle -1,0,2\rangle-\dfrac{\langle 1,-1,2\rangle\cdot\langle -1,0,2\rangle}{1^2+(-1)^2+2^2}\langle 1,-1,2\rangle=\langle -1,0,2\rangle-\frac{1}{2}\langle 1,-1,2\rangle=\left\langle -\frac{3}{2},\frac{1}{2},1\right\rangle$ is a direction vector for the required line. Thus $2\left\langle -\frac{3}{2},\frac{1}{2},1\right\rangle=\langle -3,1,2\rangle$ is also a direction vector, and the line has parametric equations $x=-3t$, $y=1+t$, $z=2+2t$. (Notice that this is the same line as in Exercise 65.)

67. Let P_i have normal vector $\mathbf{n}_i$. Then $\mathbf{n}_1=\langle 3,6,-3\rangle$, $\mathbf{n}_2=\langle 4,-12,8\rangle$, $\mathbf{n}_3=\langle 3,-9,6\rangle$, $\mathbf{n}_4=\langle 1,2,-1\rangle$. Now $\mathbf{n}_1=3\mathbf{n}_4$, so $\mathbf{n}_1$ and $\mathbf{n}_4$ are parallel, and hence P_1 and P_4 are parallel; similarly P_2 and P_3 are parallel because $\mathbf{n}_2=\frac{4}{3}\mathbf{n}_3$. However, $\mathbf{n}_1$ and $\mathbf{n}_2$ are not parallel (so not all four planes are parallel). Notice that the point $(2,0,0)$ lies on both P_1 and P_4, so these two planes are identical. The point $\left(\frac{5}{4},0,0\right)$ lies on P_2 but not on P_3, so these are different planes.

68. Let L_i have direction vector $\mathbf{v}_i$. Rewrite the symmetric equations for L_3 as $\dfrac{x-1}{1/2}=\dfrac{y-1}{-1/4}=\dfrac{z+1}{1}$; then $\mathbf{v}_1=\langle 6,-3,12\rangle$, $\mathbf{v}_2=\langle 2,1,4\rangle$, $\mathbf{v}_3=\left\langle \frac{1}{2},-\frac{1}{4},1\right\rangle$, and $\mathbf{v}_4=\langle 4,2,8\rangle$. $\mathbf{v}_1=12\mathbf{v}_3$, so L_1 and L_3 are parallel. $\mathbf{v}_4=2\mathbf{v}_2$, so L_2 and L_4 are parallel. (Note that L_1 and L_2 are not parallel.) L_1 contains the point $(1,1,5)$, but this point does not lie on L_3, so they're not identical. $(3,1,5)$ lies on L_4 and also on L_2 (for $t=1$), so L_2 and L_4 are the same line.

69. Let $Q=(1,3,4)$ and $R=(2,1,1)$, points on the line corresponding to $t=0$ and $t=1$. Let $P=(4,1,-2)$. Then $\mathbf{a}=\overrightarrow{QR}=\langle 1,-2,-3\rangle$, $\mathbf{b}=\overrightarrow{QP}=\langle 3,-2,-6\rangle$. The distance is

$$d=\frac{|\mathbf{a}\times\mathbf{b}|}{|\mathbf{a}|}=\frac{|\langle 1,-2,-3\rangle\times\langle 3,-2,-6\rangle|}{|\langle 1,-2,-3\rangle|}=\frac{|\langle 6,-3,4\rangle|}{|\langle 1,-2,-3\rangle|}=\frac{\sqrt{6^2+(-3)^2+4^2}}{\sqrt{1^2+(-2)^2+(-3)^2}}=\frac{\sqrt{61}}{\sqrt{14}}=\sqrt{\frac{61}{14}}.$$

70. Let $Q=(0,6,3)$ and $R=(2,4,4)$, points on the line corresponding to $t=0$ and $t=1$. Let $P=(0,1,3)$. Then $\mathbf{a}=\overrightarrow{QR}=\langle 2,-2,1\rangle$ and $\mathbf{b}=\overrightarrow{QP}=\langle 0,-5,0\rangle$. The distance is

$$d=\frac{|\mathbf{a}\times\mathbf{b}|}{|\mathbf{a}|}=\frac{|\langle 2,-2,1\rangle\times\langle 0,-5,0\rangle|}{|\langle 2,-2,1\rangle|}=\frac{|\langle 5,0,-10\rangle|}{|\langle 2,-2,1\rangle|}=\frac{\sqrt{5^2+0^2+(-10)^2}}{\sqrt{2^2+(-2)^2+1^2}}=\frac{\sqrt{125}}{\sqrt{9}}=\frac{5\sqrt{5}}{3}.$$

71. By Equation 9, the distance is $D=\dfrac{|ax_1+by_1+cz_1+d|}{\sqrt{a^2+b^2+c^2}}=\dfrac{|3(1)+2(-2)+6(4)-5|}{\sqrt{3^2+2^2+6^2}}=\dfrac{|18|}{\sqrt{49}}=\dfrac{18}{7}$.

72. By Equation 9, the distance is $D=\dfrac{|1(-6)-2(3)-4(5)-8|}{\sqrt{1^2+(-2)^2+(-4)^2}}=\dfrac{|-40|}{\sqrt{21}}=\dfrac{40}{\sqrt{21}}$.

73. Put $y=z=0$ in the equation of the first plane to get the point $(2,0,0)$ on the plane. Because the planes are parallel, the distance D between them is the distance from $(2,0,0)$ to the second plane. By Equation 9,

$$D=\frac{|4(2)-6(0)+2(0)-3|}{\sqrt{4^2+(-6)^2+(2)^2}}=\frac{5}{\sqrt{56}}=\frac{5}{2\sqrt{14}}\text{ or }\frac{5\sqrt{14}}{28}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

74. Put $x = y = 0$ in the equation of the first plane to get the point $(0, 0, 0)$ on the plane. Because the planes are parallel the distance D between them is the distance from $(0, 0, 0)$ to the second plane $3x - 6y + 9z - 1 = 0$. By Equation 9,

$$D = \frac{|3(0) - 6(0) + 9(0) - 1|}{\sqrt{3^2 + (-6)^2 + 9^2}} = \frac{1}{\sqrt{126}} = \frac{1}{3\sqrt{14}}.$$

75. The distance between two parallel planes is the same as the distance between a point on one of the planes and the other plane. Let $P_0 = (x_0, y_0, z_0)$ be a point on the plane given by $ax + by + cz + d_1 = 0$. Then $ax_0 + by_0 + cz_0 + d_1 = 0$ and the distance between P_0 and the plane given by $ax + by + cz + d_2 = 0$ is, from Equation 9,

$$D = \frac{|ax_0 + by_0 + cz_0 + d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|-d_1 + d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|d_1 - d_2|}{\sqrt{a^2 + b^2 + c^2}}.$$

76. The planes must have parallel normal vectors, so if $ax + by + cz + d = 0$ is such a plane, then for some $t \neq 0$, $\langle a, b, c\rangle = t\langle 1, 2, -2\rangle = \langle t, 2t, -2t\rangle$. So this plane is given by the equation $x + 2y - 2z + k = 0$, where $k = d/t$. By Exercise 75, the distance between the planes is $2 = \dfrac{|1 - k|}{\sqrt{1^2 + 2^2 + (-2)^2}} \quad \Leftrightarrow \quad 6 = |1 - k| \quad \Leftrightarrow \quad k = 7$ or -5. So the desired planes have equations $x + 2y - 2z = 7$ and $x + 2y - 2z = -5$.

77. L_1: $x = y = z \quad \Rightarrow \quad x = y$ **(1)**. L_2: $x + 1 = y/2 = z/3 \quad \Rightarrow \quad x + 1 = y/2$ **(2)**. The solution of **(1)** and **(2)** is $x = y = -2$. However, when $x = -2$, $x = z \quad \Rightarrow \quad z = -2$, but $x + 1 = z/3 \quad \Rightarrow \quad z = -3$, a contradiction. Hence the lines do not intersect. For L_1, $\mathbf{v}_1 = \langle 1, 1, 1\rangle$, and for L_2, $\mathbf{v}_2 = \langle 1, 2, 3\rangle$, so the lines are not parallel. Thus the lines are skew lines. If two lines are skew, they can be viewed as lying in two parallel planes and so the distance between the skew lines would be the same as the distance between these parallel planes. The common normal vector to the planes must be perpendicular to both $\langle 1, 1, 1\rangle$ and $\langle 1, 2, 3\rangle$, the direction vectors of the two lines. So set $\mathbf{n} = \langle 1, 1, 1\rangle \times \langle 1, 2, 3\rangle = \langle 3 - 2, -3 + 1, 2 - 1\rangle = \langle 1, -2, 1\rangle$. From above, we know that $(-2, -2, -2)$ and $(-2, -2, -3)$ are points of L_1 and L_2 respectively. So in the notation of Equation 8, $1(-2) - 2(-2) + 1(-2) + d_1 = 0 \quad \Rightarrow \quad d_1 = 0$ and $1(-2) - 2(-2) + 1(-3) + d_2 = 0 \quad \Rightarrow \quad d_2 = 1$.

By Exercise 75, the distance between these two skew lines is $D = \dfrac{|0 - 1|}{\sqrt{1 + 4 + 1}} = \dfrac{1}{\sqrt{6}}$.

Alternate solution (without reference to planes): A vector which is perpendicular to both of the lines is $\mathbf{n} = \langle 1, 1, 1\rangle \times \langle 1, 2, 3\rangle = \langle 1, -2, 1\rangle$. Pick any point on each of the lines, say $(-2, -2, -2)$ and $(-2, -2, -3)$, and form the vector $\mathbf{b} = \langle 0, 0, 1\rangle$ connecting the two points. The distance between the two skew lines is the absolute value of the scalar projection of $\mathbf{b}$ along $\mathbf{n}$, that is, $D = \dfrac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}|} = \dfrac{|1 \cdot 0 - 2 \cdot 0 + 1 \cdot 1|}{\sqrt{1 + 4 + 1}} = \dfrac{1}{\sqrt{6}}$.

78. First notice that if two lines are skew, they can be viewed as lying in two parallel planes and so the distance between the skew lines would be the same as the distance between these parallel planes. The common normal vector to the planes must be perpendicular to both $\mathbf{v}_1 = \langle 1, 6, 2\rangle$ and $\mathbf{v}_2 = \langle 2, 15, 6\rangle$, the direction vectors of the two lines, respectively. Thus set

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 36-30, 4-6, 15-12\rangle = \langle 6, -2, 3\rangle$. Setting $t = 0$ and $s = 0$ gives the points $(1, 1, 0)$ and $(1, 5, -2)$. So in the notation of Equation 8, $6 - 2 + 0 + d_1 = 0 \quad \Rightarrow \quad d_1 = -4$ and $6 - 10 - 6 + d_2 = 0 \quad \Rightarrow \quad d_2 = 10$.

Then by Exercise 75, the distance between the two skew lines is given by $D = \dfrac{|-4-10|}{\sqrt{36+4+9}} = \dfrac{14}{7} = 2$.

Alternate solution (without reference to planes): We already know that the direction vectors of the two lines are $\mathbf{v}_1 = \langle 1, 6, 2\rangle$ and $\mathbf{v}_2 = \langle 2, 15, 6\rangle$. Then $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 6, -2, 3\rangle$ is perpendicular to both lines. Pick any point on each of the lines, say $(1, 1, 0)$ and $(1, 5, -2)$, and form the vector $\mathbf{b} = \langle 0, 4, -2\rangle$ connecting the two points. Then the distance between the two skew lines is the absolute value of the scalar projection of $\mathbf{b}$ along $\mathbf{n}$, that is,

$$D = \frac{|\mathbf{n}\cdot\mathbf{b}|}{|\mathbf{n}|} = \frac{1}{\sqrt{36+4+9}}\,|0-8-6| = \frac{14}{7} = 2.$$

79. A direction vector for L_1 is $\mathbf{v}_1 = \langle 2, 0, -1\rangle$ and a direction vector for L_2 is $\mathbf{v}_2 = \langle 3, 2, 2\rangle$. These vectors are not parallel so neither are the lines. Parametric equations for the lines are L_1: $x = 2t$, $y = 0$, $z = -t$, and L_2: $x = 1 + 3s$, $y = -1 + 2s$, $z = 1 + 2s$. No values of t and s satisfy these equations simultaneously, so the lines don't intersect and hence are skew. We can view the lines as lying in two parallel planes; a common normal vector to the planes is $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 2, -7, 4\rangle$. Line L_1 passes through the origin, so $(0, 0, 0)$ lies on one of the planes, and $(1, -1, 1)$ is a point on L_2 and therefore on the other plane. Equations of the planes then are $2x - 7y + 4z = 0$ and $2x - 7y + 4z - 13 = 0$, and by Exercise 75, the distance between the two skew lines is $D = \dfrac{|0-(-13)|}{\sqrt{4+49+16}} = \dfrac{13}{\sqrt{69}}$.

Alternate solution (without reference to planes): Direction vectors of the two lines are $\mathbf{v}_1 = \langle 2, 0, -1\rangle$ and $\mathbf{v}_2 = \langle 3, 2, 2\rangle$. Then $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 2, -7, 4\rangle$ is perpendicular to both lines. Pick any point on each of the lines, say $(0, 0, 0)$ and $(1, -1, 1)$, and form the vector $\mathbf{b} = \langle 1, -1, 1\rangle$ connecting the two points. Then the distance between the two skew lines is the absolute value of the scalar projection of $\mathbf{b}$ along $\mathbf{n}$, that is, $D = \dfrac{|\mathbf{n}\cdot\mathbf{b}|}{|\mathbf{n}|} = \dfrac{|2+7+4|}{\sqrt{4+49+16}} = \dfrac{13}{\sqrt{69}}$.

80. A direction vector for the line L_1 is $\mathbf{v}_1 = \langle 1, 2, 2\rangle$. A normal vector for the plane P_1 is $\mathbf{n}_1 = \langle 1, -1, 2\rangle$. The vector from the point $(0, 0, 1)$ to $(3, 2, -1)$, $\langle 3, 2, -2\rangle$, is parallel to the plane P_2, as is the vector from $(0, 0, 1)$ to $(1, 2, 1)$, namely $\langle 1, 2, 0\rangle$. Thus a normal vector for P_2 is $\langle 3, 2, -2\rangle \times \langle 1, 2, 0\rangle = \langle 4, -2, 4\rangle$, or we can use $\mathbf{n}_2 = \langle 2, -1, 2\rangle$, and a direction vector for the line L_2 of intersection of these planes is $\mathbf{v}_2 = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, -1, 2\rangle \times \langle 2, -1, 2\rangle = \langle 0, 2, 1\rangle$. Notice that the point $(3, 2, -1)$ lies on both planes, so it also lies on L_2. The lines are skew, so we can view them as lying in two parallel planes; a common normal vector to the planes is $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle -2, -1, 2\rangle$. Line L_1 passes through the point $(1, 2, 6)$, so $(1, 2, 6)$ lies on one of the planes, and $(3, 2, -1)$ is a point on L_2 and therefore on the other plane. Equations of the planes then are $-2x - y + 2z - 8 = 0$ and $-2x - y + 2z + 10 = 0$, and by Exercise 75, the distance between the lines is

$$D = \frac{|-8-10|}{\sqrt{4+1+4}} = \frac{18}{3} = 6.$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Alternatively, direction vectors for the lines are $\mathbf{v}_1 = \langle 1, 2, 2\rangle$ and $\mathbf{v}_2 = \langle 0, 2, 1\rangle$, so $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle -2, -1, 2\rangle$ is perpendicular to both lines. Pick any point on each of the lines, say $(1, 2, 6)$ and $(3, 2, -1)$, and form the vector $\mathbf{b} = \langle 2, 0, -7\rangle$ connecting the two points. Then the distance between the two skew lines is the absolute value of the scalar projection of $\mathbf{b}$ along $\mathbf{n}$, that is, $D = \dfrac{|\mathbf{n}\cdot\mathbf{b}|}{|\mathbf{n}|} = \dfrac{|-4+0-14|}{\sqrt{4+1+4}} = \dfrac{18}{3} = 6$.

81. (a) A direction vector from tank A to tank B is $\langle 765-325, 675-810, 599-561\rangle = \langle 440, -135, 38\rangle$. Taking tank A's position $(325, 810, 561)$ as the initial point, parametric equations for the line of sight are $x = 325 + 440t$, $y = 810 - 135t$, $z = 561 + 38t$ for $0 \le t \le 1$.

(b) We divide the line of sight into 5 equal segments, corresponding to $\Delta t = 0.2$, and compute the elevation from the z-component of the parametric equations in part (a):

t	$z = 561 + 38t$	terrain elevation
0	561.0	
0.2	568.6	549
0.4	576.2	566
0.6	583.8	586
0.8	591.4	589
1.0	599.0	

Since the terrain is higher than the line of sight when $t = 0.6$, the tanks can't see each other.

82. (a) The planes $x + y + z = c$ have normal vector $\langle 1, 1, 1\rangle$, so they are all parallel. Their x-, y-, and z-intercepts are all c. When $c > 0$ their intersection with the first octant is an equilateral triangle and when $c < 0$ their intersection with the octant diagonally opposite the first is an equilateral triangle.

(b) The planes $x + y + cz = 1$ have x-intercept 1, y-intercept 1, and z-intercept $1/c$. The plane with $c = 0$ is parallel to the z-axis. As c gets larger, the planes get closer to the xy-plane.

(c) The planes $y\cos\theta + z\sin\theta = 1$ have normal vectors $\langle 0, \cos\theta, \sin\theta\rangle$, which are perpendicular to the x-axis, and so the planes are parallel to the x-axis. We look at their intersection with the yz-plane. These are lines that are perpendicular to $\langle \cos\theta, \sin\theta\rangle$ and pass through $(\cos\theta, \sin\theta)$, since $\cos^2\theta + \sin^2\theta = 1$. So these are the tangent lines to the unit circle. Thus the family consists of all planes tangent to the circular cylinder with radius 1 and axis the x-axis.

83. If $a \neq 0$, then $ax + by + cz + d = 0 \quad\Rightarrow\quad a(x + d/a) + b(y - 0) + c(z - 0) = 0$ which by (7) is the scalar equation of the plane through the point $(-d/a, 0, 0)$ with normal vector $\langle a, b, c\rangle$. Similarly, if $b \neq 0$ (or if $c \neq 0$) the equation of the plane can be rewritten as $a(x - 0) + b(y + d/b) + c(z - 0) = 0$ [or as $a(x - 0) + b(y - 0) + c(z + d/c) = 0$] which by (7) is the scalar equation of a plane through the point $(0, -d/b, 0)$ [or the point $(0, 0, -d/c)$] with normal vector $\langle a, b, c\rangle$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

DISCOVERY PROJECT Putting 3D in Perspective

1. If we view the screen from the camera's location, the vertical clipping plane on the left passes through the points $(1000, 0, 0)$, $(0, -400, 0)$, and $(0, -400, 600)$. A vector from the first point to the second is $\mathbf{v}_1 = \langle -1000, -400, 0\rangle$ and a vector from the first point to the third is $\mathbf{v}_2 = \langle -1000, -400, 600\rangle$. A normal vector for the clipping plane is $\mathbf{v}_1 \times \mathbf{v}_2 = -240{,}000\,\mathbf{i} + 600{,}000\,\mathbf{j}$ or $-2\,\mathbf{i} + 5\,\mathbf{j}$, and an equation for the plane is $-2(x - 1000) + 5(y - 0) + 0(z - 0) = 0 \quad \Rightarrow \quad 2x - 5y = 2000$. By symmetry, the vertical clipping plane on the right is given by $2x + 5y = 2000$. The lower clipping plane is $z = 0$. The upper clipping plane passes through the points $(1000, 0, 0)$, $(0, -400, 600)$, and $(0, 400, 600)$. Vectors from the first point to the second and third points are $\mathbf{v}_1 = \langle -1000, -400, 600\rangle$ and $\mathbf{v}_2 = \langle -1000, 400, 600\rangle$, and a normal vector for the plane is $\mathbf{v}_1 \times \mathbf{v}_2 = -480{,}000\,\mathbf{i} - 800{,}000\,\mathbf{k}$ or $3\,\mathbf{i} + 5\,\mathbf{k}$. An equation for the plane is $3(x - 1000) + 0(y - 0) + 5(z - 0) = 0 \quad \Rightarrow \quad 3x + 5z = 3000$.

A direction vector for the line L is $\mathbf{v} = \langle 630, 390, 162\rangle$ and taking $P_0 = (230, -285, 102)$, parametric equations are $x = 230 + 630t$, $y = -285 + 390t$, $z = 102 + 162t$. L intersects the left clipping plane when $2(230 + 630t) - 5(-285 + 390t) = 2000 \quad \Rightarrow \quad t = -\frac{1}{6}$. The corresponding point is $(125, -350, 75)$. L intersects the right clipping plane when $2(230 + 630t) + 5(-285 + 390t) = 2000 \quad \Rightarrow \quad t = \frac{593}{642}$. The corresponding point is approximately $(811.9, 75.2, 251.6)$, but this point is not contained within the viewing volume. L intersects the upper clipping plane when $3(230 + 630t) + 5(102 + 162t) = 3000 \quad \Rightarrow \quad t = \frac{2}{3}$, corresponding to the point $(650, -25, 210)$, and L intersects the lower clipping plane when $z = 0 \quad \Rightarrow \quad 102 + 162t = 0 \quad \Rightarrow \quad t = -\frac{17}{27}$. The corresponding point is approximately $(-166.7, -530.6, 0)$, which is not contained within the viewing volume. Thus L should be clipped at the points $(125, -350, 75)$ and $(650, -25, 210)$.

2. A sight line from the camera at $(1000, 0, 0)$ to the left endpoint $(125, -350, 75)$ of the clipped line has direction $\mathbf{v} = \langle -875, -350, 75\rangle$. Parametric equations are $x = 1000 - 875t$, $y = -350t$, $z = 75t$. This line intersects the screen when $x = 0 \quad \Rightarrow \quad 1000 - 875t = 0 \quad \Rightarrow \quad t = \frac{8}{7}$, corresponding to the point $\left(0, -400, \frac{600}{7}\right)$. Similarly, a sight line from the camera to the right endpoint $(650, -25, 210)$ of the clipped line has direction $\langle -350, -25, 210\rangle$ and parametric equations are $x = 1000 - 350t$, $y = -25t$, $z = 210t$. $x = 0 \quad \Rightarrow \quad 1000 - 350t = 0 \quad \Rightarrow \quad t = \frac{20}{7}$, corresponding to the point $\left(0, -\frac{500}{7}, 600\right)$. Thus the projection of the clipped line is the line segment between the points $\left(0, -400, \frac{600}{7}\right)$ and $\left(0, -\frac{500}{7}, 600\right)$.

3. From Equation 12.5.4, equations for the four sides of the screen are $\mathbf{r}_1(t) = (1 - t)\langle 0, -400, 0\rangle + t\,\langle 0, -400, 600\rangle$, $\mathbf{r}_2(t) = (1 - t)\langle 0, -400, 600\rangle + t\,\langle 0, 400, 600\rangle$, $\mathbf{r}_3(t) = (1 - t)\langle 0, 400, 0\rangle + t\,\langle 0, 400, 600\rangle$, and $\mathbf{r}_4(t) = (1 - t)\langle 0, -400, 0\rangle + t\,\langle 0, 400, 0\rangle$. The clipped line segment connects the points $(125, -350, 75)$ and $(650, -25, 210)$, so an equation for the segment is $\mathbf{r}_5(t) = (1 - t)\langle 125, -350, 75\rangle + t\,\langle 650, -25, 210\rangle$. The projection of the clipped segment connects the points

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\left(0, -400, \frac{600}{7}\right)$ and $\left(0, -\frac{500}{7}, 600\right)$, so an equation is $\mathbf{r}_6(t) = (1-t)\left\langle 0, -400, \frac{600}{7}\right\rangle + t\left\langle 0, -\frac{500}{7}, 600\right\rangle$.

The sight line on the left connects the points $(1000, 0, 0)$ and $\left(0, -400, \frac{600}{7}\right)$, so an equation is $\mathbf{r}_7(t) = (1-t)\langle 1000, 0, 0\rangle + t\left\langle 0, -400, \frac{600}{7}\right\rangle$. The other sight line connects $(1000, 0, 0)$ to $\left(0, -\frac{500}{7}, 600\right)$, so an equation is $\mathbf{r}_8(t) = (1-t)\langle 1000, 0, 0\rangle + t\left\langle 0, -\frac{500}{7}, 600\right\rangle$.

4. The vector from $(621, -147, 206)$ to $(563, 31, 242)$, $\mathbf{v}_1 = \langle -58, 178, 36\rangle$, lies in the plane of the rectangle, as does the vector from $(621, -147, 206)$ to $(657, -111, 86)$, $\mathbf{v}_2 = \langle 36, 36, -120\rangle$. A normal vector for the plane is $\mathbf{v}_1 \times \mathbf{v}_2 = \langle -1888, -142, -708\rangle$ or $\langle 8, 2, 3\rangle$, and an equation of the plane is $8x + 2y + 3z = 5292$. The line L intersects this plane when $8(230 + 630t) + 2(-285 + 390t) + 3(102 + 162t) = 5292 \quad \Rightarrow \quad t = \frac{1858}{3153} \approx 0.589$. The corresponding point is approximately $(601.25, -55.18, 197.46)$. Starting at this point, a portion of the line is hidden behind the rectangle. The line becomes visible again at the left edge of the rectangle, specifically the edge between the points $(621, -147, 206)$ and $(657, -111, 86)$. (This is most easily determined by graphing the rectangle and the line.) A plane through these two points and the camera's location, $(1000, 0, 0)$, will clip the line at the point it becomes visible. Two vectors in this plane are $\mathbf{v}_1 = \langle -379, -147, 206\rangle$ and $\mathbf{v}_2 = \langle -343, -111, 86\rangle$. A normal vector for the plane is $\mathbf{v}_1 \times \mathbf{v}_2 = \langle 10224, -38064, -8352\rangle$ and an equation of the plane is $213x - 793y - 174z = 213{,}000$. L intersects this plane when $213(230 + 630t) - 793(-285 + 390t) - 174(102 + 162t) = 213{,}000 \quad \Rightarrow \quad t = \frac{44{,}247}{203{,}268} \approx 0.2177$. The corresponding point is approximately $(367.14, -200.11, 137.26)$. Thus the portion of L that should be removed is the segment between the points $(601.25, -55.18, 197.46)$ and $(367.14, -200.11, 137.26)$.

12.6 Cylinders and Quadric Surfaces

1. (a) In $\mathbb{R}^2$, the equation $y = x^2$ represents a parabola.

(b) In $\mathbb{R}^3$, the equation $y = x^2$ doesn't involve z, so any horizontal plane with equation $z = k$ intersects the graph in a curve with equation $y = x^2$. Thus, the surface is a parabolic cylinder, made up of infinitely many shifted copies of the same parabola. The rulings are parallel to the z-axis.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) In $\mathbb{R}^3$, the equation $z = y^2$ also represents a parabolic cylinder. Since x doesn't appear, the graph is formed by moving the parabola $z = y^2$ in the direction of the x-axis. Thus, the rulings of the cylinder are parallel to the x-axis.

2. (a)

(b) Since the equation $y = e^x$ doesn't involve z, horizontal traces are copies of the curve $y = e^x$. The rulings are parallel to the z-axis.

(c) The equation $z = e^y$ doesn't involve x, so vertical traces in $x = k$ (parallel to the yz-plane) are copies of the curve $z = e^y$. The rulings are parallel to the x-axis.

3. Since y is missing from the equation, the vertical traces $x^2 + z^2 = 4$, $y = k$ are copies of the same circle in the plane $y = k$. Thus, the surface $x^2 + z^2 = 4$ is a circular cylinder of radius 2 with rulings parallel to the y-axis.

4. Since x is missing from the equation, the vertical traces $y^2 + 9z^2 = 9$, $x = k$ are copies of the same ellipse in the plane $x = k$. Thus, the surface $y^2 + 9z^2 = 9$ is an elliptic cylinder with rulings parallel to the x-axis.

5. Since z is missing from the equation, the horizontal traces $x^2 + y + 1 = 0 \quad \Rightarrow \quad y = -x^2 - 1$, $z = k$ are copies of the same parabola in the plane $z = k$. Thus, the surface $x^2 + y + 1 = 0$ is a parabolic cylinder with rulings parallel to the z-axis.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. Since y is missing from the equation, the vertical traces $z = -\sqrt{x}$, $y = k$ are copies of the curve $z = -\sqrt{x}$ with rulings parallel to the y-axis.

7. Since z is missing, each horizontal trace $xy = 1$, $z = k$, is a copy of the same hyperbola in the plane $z = k$. Thus the surface $xy = 1$ is a hyperbolic cylinder with rulings parallel to the z-axis.

8. Since x is missing, each vertical trace $z = \sin y$, $x = k$, is a copy of a sine curve in the plane $x = k$. Thus the surface $z = \sin y$ is a cylindrical surface with rulings parallel to the x-axis.

9. The trace in the xz-plane appears to be $z = \cos x$. The traces in the planes on the positive y-axis and negative y-axis are copies of the same graph. Therefore, an equation of the graph could be $z = \cos x$.

10. The trace in the yz-plane appears to be $z = y^3$. The traces in the planes on the positive x-axis and negative x-axis are copies of the same graph. Therefore, an equation of the graph could be $z = y^3$.

11. (a) The traces of $x^2 + y^2 - z^2 = 1$ in $x = k$ are $y^2 - z^2 = 1 - k^2$, a family of hyperbolas. (Note that the hyperbolas are oriented differently for $-1 < k < 1$ than for $k < -1$ or $k > 1$.) The traces in $y = k$ are $x^2 - z^2 = 1 - k^2$, a similar family of hyperbolas. The traces in $z = k$ are $x^2 + y^2 = 1 + k^2$, a family of circles. For $k = 0$, the trace in the xy-plane, the circle is of radius 1. As $|k|$ increases, so does the radius of the circle. This behavior, combined with the hyperbolic vertical traces, gives the graph of the hyperboloid of one sheet in Table 1.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) If we change the equation $x^2 + y^2 - z^2 = 1$ to $x^2 - y^2 + z^2 = 1$, the shape of the surface is unchanged, but the hyperboloid is rotated so that its axis is the y-axis. Traces in $y = k$ are circles, while traces in $x = k$ and $z = k$ are hyperbolas.

(c) Completing the square in y for $x^2 + y^2 + 2y - z^2 = 0$ gives $x^2 + (y+1)^2 - z^2 = 1$. The surface is a hyperboloid identical to the one in part (a) but shifted one unit in the negative y-direction.

12. (a) The traces of $-x^2 - y^2 + z^2 = 1$ in $x = k$ are $-y^2 + z^2 = 1 + k^2$, a family of hyperbolas, as are the traces in $y = k$, $-x^2 + z^2 = 1 + k^2$. The traces in $z = k$ are $x^2 + y^2 = k^2 - 1$, a family of circles for $|k| > 1$. As $|k|$ increases, the radii of the circles increase; the traces are empty for $|k| < 1$. This behavior, combined with the vertical traces, gives the graph of the hyperboloid of two sheets in Table 1.

(b) If the equation in part (a) is changed to $x^2 - y^2 - z^2 = 1$, the graph has the same shape as the hyperboloid in part (a) but is rotated so that its axis is the x-axis. Traces in $x = k$, $|k| > 1$, are circles, while traces in $y = k$ and $z = k$ are hyperbolas.

13. For $x = y^2 + 4z^2$, the traces in $x = k$ are $y^2 + 4z^2 = k$. When $k > 0$ we have a family of ellipses. When $k = 0$ we have just a point at the origin, and the trace is empty for $k < 0$. The traces in $y = k$ are $x = 4z^2 + k^2$, a family of parabolas opening in the positive x-direction. Similarly, the traces in $z = k$ are $x = y^2 + 4k^2$, a family of parabolas opening in the positive x-direction. We recognize the graph as an elliptic paraboloid with axis the x-axis and vertex the origin.

14. $4x^2 + 9y^2 + 9z^2 = 36$. The traces in $x = k$ are $9y^2 + 9z^2 = 36 - 4k^2 \quad \Leftrightarrow$ $y^2 + z^2 = 4 - \frac{4}{9}k^2$, a family of circles for $|k| < 3$. (The traces are a single point for $|k| = 3$ and are empty for $|k| > 3$.) The traces in $y = k$ are $4x^2 + 9z^2 = 36 - 9k^2$, a family of ellipses for $|k| < 2$. Similarly, the traces in $z = k$ are the ellipses $4x^2 + 9y^2 = 36 - 9k^2$, $|k| < 2$. The graph is an ellipsoid centered at the origin with intercepts $x = \pm 3$, $y = \pm 2$, $z = \pm 2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $x^2 = 4y^2 + z^2$. The traces in $x = k$ are the ellipses $4y^2 + z^2 = k^2$. The traces in $y = k$ are $x^2 - z^2 = 4k^2$, hyperbolas for $k \neq 0$ and two intersecting lines if $k = 0$. Similarly, the traces in $z = k$ are $x^2 - 4y^2 = k^2$, hyperbolas for $k \neq 0$ and two intersecting lines if $k = 0$. We recognize the graph as an elliptic cone with axis the x-axis and vertex the origin.

16. $z^2 - 4x^2 - y^2 = 4$. The traces in $x = k$ are the hyperbolas $z^2 - y^2 = 4 + 4k^2$, and the traces in $y = k$ are the hyperbolas $z^2 - 4x^2 = 4 + k^2$. The traces in $z = k$ are $4x^2 + y^2 = k^2 - 4$, a family of ellipses for $|k| > 2$. (The traces are a single point for $|k| = 2$ and are empty for $|k| < 2$.) The surface is a hyperboloid of two sheets with axis the z-axis.

17. $9y^2 + 4z^2 = x^2 + 36$. The traces in $x = k$ are $9y^2 + 4z^2 = k^2 + 36$, a family of ellipses. The traces in $y = k$ are $4z^2 - x^2 = 9(4 - k^2)$, a family of hyperbolas for $|k| \neq 2$ and two intersecting lines when $|k| = 2$. (Note that the hyperbolas are oriented differently for $|k| < 2$ than for $|k| > 2$.) The traces in $z = k$ are $9y^2 - x^2 = 4(9 - k^2)$, a family of hyperbolas when $|k| \neq 3$ (oriented differently for $|k| < 3$ than for $|k| > 3$) and two intersecting lines when $|k| = 3$. We recognize the graph as a hyperboloid of one sheet with axis the x-axis.

18. $3x^2 + y + 3z^2 = 0$. The traces in $x = k$ are the parabolas $y = -3z^2 - 3k^2$ which open to the left (in the negative y-direction). Traces in $y = k$ are $3x^2 + 3z^2 = -k \quad \Leftrightarrow \quad x^2 + z^2 = -\dfrac{k}{3}$, a family of circles for $k < 0$. (Traces are empty for $k > 0$ and a single point for $k = 0$.) Traces in $z = k$ are the parabolas $y = -3x^2 - 3k^2$ which open in the negative y-direction. The graph is a circular paraboloid with axis the y-axis, opening in the negative y-direction, and vertex the origin.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $\dfrac{x^2}{9}+\dfrac{y^2}{25}+\dfrac{z^2}{4}=1$. The traces in $x=k$ are $\dfrac{y^2}{25}+\dfrac{z^2}{4}=1-\dfrac{k^2}{9}$, a family of ellipses for $|k|<3$. (The traces are a single point for $|k|=3$ and are empty for $|k|>3$.) The traces in $y=k$ are the ellipses $\dfrac{x^2}{9}+\dfrac{z^2}{4}=1-\dfrac{k^2}{25}$, $|k|<5$, and the traces in $z=k$ are the ellipses $\dfrac{x^2}{9}+\dfrac{y^2}{25}=1-\dfrac{k^2}{4}$, $|k|<2$. The surface is an ellipsoid centered at the origin with intercepts $x=\pm3$, $y=\pm5$, $z=\pm2$.

20. $3x^2-y^2+3z^2=0$. The traces in $x=k$ are $y^2-3z^2=3k^2$, a family of hyperbolas for $k\neq0$ and two intersecting lines if $k=0$. Traces in $y=k$ are the circles $3x^2+3z^2=k^2 \quad\Leftrightarrow\quad x^2+z^2=\frac{1}{3}k^2$. The traces in $z=k$ are $y^2-3x^2=3k^2$, hyperbolas for $k\neq0$ and two intersecting lines if $k=0$. We recognize the surface as a circular cone with axis the y-axis and vertex the origin.

21. $y=z^2-x^2$. The traces in $x=k$ are the parabolas $y=z^2-k^2$, opening in the positive y-direction. The traces in $y=k$ are $k=z^2-x^2$, two intersecting lines when $k=0$ and a family of hyperbolas for $k\neq0$ (note that the hyperbolas are oriented differently for $k>0$ than for $k<0$). The traces in $z=k$ are the parabolas $y=k^2-x^2$ which open in the negative y-direction. Thus the surface is a hyperbolic paraboloid centered at $(0,0,0)$.

22. $x=y^2-z^2$. The traces in $x=k$ are $y^2-z^2=k$, two intersecting lines when $k=0$ and a family of hyperbolas for $k\neq0$ (oriented differently for $k>0$ than for $k<0$). The traces in $y=k$ are the parabolas $x=-z^2+k^2$, opening in the negative x-direction, and the traces in $z=k$ are the parabolas $x=y^2-k^2$ which open in the positive x-direction. The graph is a hyperbolic paraboloid centered at $(0,0,0)$.

23. This is the equation of an ellipsoid: $x^2+4y^2+9z^2=x^2+\dfrac{y^2}{(1/2)^2}+\dfrac{z^2}{(1/3)^2}=1$, with x-intercepts ±1, y-intercepts $\pm\frac{1}{2}$ and z-intercepts $\pm\frac{1}{3}$. So the major axis is the x-axis and the only possible graph is VII.

24. This is the equation of an ellipsoid: $9x^2+4y^2+z^2=\dfrac{x^2}{(1/3)^2}+\dfrac{y^2}{(1/2)^2}+z^2=1$, with x-intercepts $\pm\frac{1}{3}$, y-intercepts $\pm\frac{1}{2}$ and z-intercepts ±1. So the major axis is the z-axis and the only possible graph is IV.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $x^2 - y^2 + z^2 = 1$ is the equation of a hyperboloid of one sheet, with $a = b = c = 1$. Since the coefficient of y^2 is negative, the axis of the hyperboloid is the y-axis. Hence, the correct graph is II.

26. $-x^2 + y^2 - z^2 = 1$ is the equation of a hyperboloid of two sheets, with $a = b = c = 1$. This surface does not intersect the xz-plane at all, so the axis of the hyperboloid is the y-axis. Hence, the correct graph is III.

27. There are no real values of x and z that satisfy this equation, $y = 2x^2 + z^2$, for $y < 0$, so this surface does not extend to the left of the xz-plane. The surface intersects the plane $y = k > 0$ in an ellipse. Notice that y occurs to the first power whereas x and z occur to the second power. So the surface is an elliptic paraboloid with axis the y-axis. Its graph is VI.

28. $y^2 = x^2 + 2z^2$ is the equation of a cone with axis the y-axis. Its graph is I.

29. $x^2 + 2z^2 = 1$ is the equation of a cylinder because the variable y is missing from the equation. The intersection of the surface and the xz-plane is an ellipse. Its graph is VIII.

30. $y = x^2 - z^2$ is the equation of a hyperbolic paraboloid. The trace in the xy-plane is the parabola $y = x^2$. So the correct graph is V.

31. Vertical traces parallel to the xz-plane are circles centered at the origin whose radii increase as y decreases. (The trace in $y = 1$ is just a single point and the graph suggests that traces in $y = k$ are empty for $k > 1$.) The traces in vertical planes parallel to the yz-plane are parabolas opening to the left that shift to the left as $|x|$ increases. One surface that fits this description is a circular paraboloid, opening to the left, with vertex $(0, 1, 0)$.

32. The vertical traces parallel to the yz-plane are ellipses that are smallest in the yz-plane and increase in size as $|x|$ increases. One surface that fits this description is a hyperboloid of one sheet with axis the x-axis. The horizontal traces in $z = k$ (hyperbolas and intersecting lines) also fit this surface, as shown in the figure.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. $y^2 = x^2 + \frac{1}{9}z^2$ or $y^2 = x^2 + \dfrac{z^2}{9}$ represents an elliptic cone with vertex $(0, 0, 0)$ and axis the y-axis.

34. $4x^2 - y + 2z^2 = 0$ or $y = \dfrac{x^2}{1/4} + \dfrac{z^2}{1/2}$ or $\dfrac{y}{4} = x^2 + \dfrac{z^2}{2}$ represents an elliptic paraboloid with vertex $(0, 0, 0)$ and axis the y-axis.

35. $x^2 + 2y - 2z^2 = 0$ or $2y = 2z^2 - x^2$ or $y = z^2 - \dfrac{x^2}{2}$ represents a hyperbolic paraboloid with center $(0, 0, 0)$.

36. $y^2 = x^2 + 4z^2 + 4$ or $-x^2 + y^2 - 4z^2 = 4$ or $-\dfrac{x^2}{4} + \dfrac{y^2}{4} - z^2 = 1$ represents a hyperboloid of two sheets with axis the y-axis.

37. Completing squares in x and y gives

$\left(x^2 - 2x + 1\right) + \left(y^2 - 6y + 9\right) - z = 0 \quad \Leftrightarrow$

$(x - 1)^2 + (y - 3)^2 - z = 0$ or $z = (x - 1)^2 + (y - 3)^2$, a circular paraboloid opening upward with vertex $(1, 3, 0)$ and axis the vertical line $x = 1$, $y = 3$.

38. Completing squares in x and z gives

$\left(x^2 - 4x + 4\right) - y^2 - \left(z^2 + 2z + 1\right) + 3 = 0 + 4 - 1 \quad \Leftrightarrow$

$(x - 2)^2 - y^2 - (z + 1)^2 = 0$ or $(x - 2)^2 = y^2 + (z + 1)^2$, a circular cone with vertex $(2, 0, -1)$ and axis the horizontal line $y = 0$, $z = -1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. Completing squares in x and z gives

$\left(x^2-4x+4\right)-y^2+\left(z^2-2z+1\right)=0+4+1 \quad \Leftrightarrow$

$(x-2)^2-y^2+(z-1)^2=5$ or $\dfrac{(x-2)^2}{5}-\dfrac{y^2}{5}+\dfrac{(z-1)^2}{5}=1$, a hyperboloid of one sheet with center $(2,0,1)$ and axis the horizontal line $x=2$, $z=1$.

40. Completing squares in all three variables gives

$4(x^2-6x+9)+(y^2-8y+16)+(z^2+4z+4)=-55+36+16+4 \quad \Leftrightarrow$

$4\,(x-3)^2+(y-4)^2+(z+2)^2=1$ or

$\dfrac{(x-3)^2}{1/4}+(y-4)^2+(z+2)^2=1$, an ellipsoid with center $(3,4,-2)$.

41. Solving the equation for z we get $z=\pm\sqrt{1+4x^2+y^2}$, so we plot separately $z=\sqrt{1+4x^2+y^2}$ and $z=-\sqrt{1+4x^2+y^2}$.

To restrict the z-range as in the second graph, we can use the option `view=-4..4` in Maple's `plot3d` command, or `PlotRange->{-4,4}` in Mathematica's `Plot3D` command.

42. We plot the surface $z=x^2-y^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. Solving the equation for z we get $z = \pm\sqrt{4x^2 + y^2}$, so we plot separately $z = \sqrt{4x^2 + y^2}$ and $z = -\sqrt{4x^2 + y^2}$.

44. We plot the surface $z = x^2 - 6x + 4y^2$.

45.

46.

47. The curve $y = \sqrt{x}$ is equivalent to $x = y^2$, $y \geq 0$. Rotating the curve about the x-axis creates a circular paraboloid with vertex at the origin, axis the x-axis, opening in the positive x-direction. The trace in the xy-plane is $x = y^2$, $z = 0$, and the trace in the xz-plane is a parabola of the same shape: $x = z^2$, $y = 0$. An equation for the surface is $x = y^2 + z^2$.

48. Rotating the line $z = 2y$ about the z-axis creates a (right) circular cone with vertex at the origin and axis the z-axis. Traces in $z = k$ ($k \neq 0$) are circles with center $(0, 0, k)$ and radius $y = z/2 = k/2$, so an equation for the trace is $x^2 + y^2 = (k/2)^2$, $z = k$. Thus an equation for the surface is $x^2 + y^2 = (z/2)^2$ or $4x^2 + 4y^2 = z^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. Let $P = (x, y, z)$ be an arbitrary point equidistant from $(-1, 0, 0)$ and the plane $x = 1$. Then the distance from P to $(-1, 0, 0)$ is $\sqrt{(x+1)^2 + y^2 + z^2}$ and the distance from P to the plane $x = 1$ is $|x-1| / \sqrt{1^2} = |x-1|$ (by Equation 12.5.9). So $|x-1| = \sqrt{(x+1)^2 + y^2 + z^2} \quad \Leftrightarrow \quad (x-1)^2 = (x+1)^2 + y^2 + z^2 \quad \Leftrightarrow$ $x^2 - 2x + 1 = x^2 + 2x + 1 + y^2 + z^2 \quad \Leftrightarrow \quad -4x = y^2 + z^2$. Thus the collection of all such points P is a circular paraboloid with vertex at the origin, axis the x-axis, which opens in the negative x-direction.

50. Let $P = (x, y, z)$ be an arbitrary point whose distance from the x-axis is twice its distance from the yz-plane. The distance from P to the x-axis is $\sqrt{(x-x)^2 + y^2 + z^2} = \sqrt{y^2 + z^2}$ and the distance from P to the yz-plane ($x = 0$) is $|x| / 1 = |x|$. Thus $\sqrt{y^2 + z^2} = 2|x| \quad \Leftrightarrow \quad y^2 + z^2 = 4x^2 \quad \Leftrightarrow \quad x^2 = (y^2/2^2) + (z^2/2^2)$. So the surface is a right circular cone with vertex the origin and axis the x-axis.

51. (a) An equation for an ellipsoid centered at the origin with intercepts $x = \pm a$, $y = \pm b$, and $z = \pm c$ is $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} + \dfrac{z^2}{c^2} = 1$.

Here the poles of the model intersect the z-axis at $z = \pm 6356.523$ and the equator intersects the x- and y-axes at $x = \pm 6378.137$, $y = \pm 6378.137$, so an equation is

$$\frac{x^2}{(6378.137)^2} + \frac{y^2}{(6378.137)^2} + \frac{z^2}{(6356.523)^2} = 1$$

(b) Traces in $z = k$ are the circles $\dfrac{x^2}{(6378.137)^2} + \dfrac{y^2}{(6378.137)^2} = 1 - \dfrac{k^2}{(6356.523)^2} \quad \Leftrightarrow$

$x^2 + y^2 = (6378.137)^2 - \left(\dfrac{6378.137}{6356.523}\right)^2 k^2$.

(c) To identify the traces in $y = mx$ we substitute $y = mx$ into the equation of the ellipsoid:

$$\begin{aligned} \frac{x^2}{(6378.137)^2} + \frac{(mx)^2}{(6378.137)^2} + \frac{z^2}{(6356.523)^2} &= 1 \\ \frac{(1+m^2)x^2}{(6378.137)^2} + \frac{z^2}{(6356.523)^2} &= 1 \\ \frac{x^2}{(6378.137)^2/(1+m^2)} + \frac{z^2}{(6356.523)^2} &= 1 \end{aligned}$$

As expected, this is a family of ellipses.

52. If we position the hyperboloid on coordinate axes so that it is centered at the origin with axis the z-axis then its equation is given by $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} - \dfrac{z^2}{c^2} = 1$. Horizontal traces in $z = k$ are $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1 + \dfrac{k^2}{c^2}$, a family of ellipses, but we know that the traces are circles so we must have $a = b$. The trace in $z = 0$ is $\dfrac{x^2}{a^2} + \dfrac{y^2}{a^2} = 1 \quad \Leftrightarrow \quad x^2 + y^2 = a^2$ and since the minimum radius of 100 m occurs there, we must have $a = 100$. The base of the tower is the trace in $z = -500$ given by $\dfrac{x^2}{a^2} + \dfrac{y^2}{a^2} = 1 + \dfrac{(-500)^2}{c^2}$ but $a = 100$ so the trace is $x^2 + y^2 = 100^2 + 50{,}000^2 \dfrac{1}{c^2}$. We know the base is a circle of

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

radius 140, so we must have $100^2 + 50{,}000^2 \dfrac{1}{c^2} = 140^2 \quad \Rightarrow \quad c^2 = \dfrac{50{,}000^2}{140^2 - 100^2} = \dfrac{781{,}250}{3}$ and an equation for the tower is $\dfrac{x^2}{100^2} + \dfrac{y^2}{100^2} - \dfrac{z^2}{(781{,}250)/3} = 1$ or $\dfrac{x^2}{10{,}000} + \dfrac{y^2}{10{,}000} - \dfrac{3z^2}{781{,}250} = 1,\ -500 \le z \le 500$.

53. If (a, b, c) satisfies $z = y^2 - x^2$, then $c = b^2 - a^2$. L_1: $x = a + t$, $y = b + t$, $z = c + 2(b - a)t$, L_2: $x = a + t$, $y = b - t$, $z = c - 2(b + a)t$. Substitute the parametric equations of L_1 into the equation of the hyperbolic paraboloid in order to find the points of intersection: $z = y^2 - x^2 \quad \Rightarrow$ $c + 2(b - a)t = (b + t)^2 - (a + t)^2 = b^2 - a^2 + 2(b - a)t \quad \Rightarrow \quad c = b^2 - a^2$. As this is true for all values of t, L_1 lies on $z = y^2 - x^2$. Performing similar operations with L_2 gives: $z = y^2 - x^2 \quad \Rightarrow$ $c - 2(b + a)t = (b - t)^2 - (a + t)^2 = b^2 - a^2 - 2(b + a)t \quad \Rightarrow \quad c = b^2 - a^2$. This tells us that all of L_2 also lies on $z = y^2 - x^2$.

54. Any point on the curve of intersection must satisfy both $2x^2 + 4y^2 - 2z^2 + 6x = 2$ and $2x^2 + 4y^2 - 2z^2 - 5y = 0$. Subtracting, we get $6x + 5y = 2$, which is linear and therefore the equation of a plane. Thus the curve of intersection lies in this plane.

55.

The curve of intersection looks like a bent ellipse. The projection of this curve onto the xy-plane is the set of points $(x, y, 0)$ which satisfy $x^2 + y^2 = 1 - y^2 \quad \Leftrightarrow \quad x^2 + 2y^2 = 1 \quad \Leftrightarrow$ $x^2 + \dfrac{y^2}{\left(1/\sqrt{2}\right)^2} = 1$. This is an equation of an ellipse.

12 Review

TRUE-FALSE QUIZ

1. This is false, as the dot product of two vectors is a scalar, not a vector.

2. False. For example, if $\mathbf{u} = \mathbf{i}$ and $\mathbf{v} = -\mathbf{i}$ then $|\mathbf{u} + \mathbf{v}| = |\mathbf{0}| = 0$ but $|\mathbf{u}| + |\mathbf{v}| = 1 + 1 = 2$.

3. False. For example, if $\mathbf{u} = \mathbf{i}$ and $\mathbf{v} = \mathbf{j}$ then $|\mathbf{u} \cdot \mathbf{v}| = |0| = 0$ but $|\mathbf{u}|\,|\mathbf{v}| = 1 \cdot 1 = 1$. In fact, by Theorem 12.3.3, $|\mathbf{u} \cdot \mathbf{v}| = \big||\mathbf{u}|\,|\mathbf{v}| \cos\theta\big|$.

4. False. For example, $|\mathbf{i} \times \mathbf{i}| = |\mathbf{0}| = 0$ (see Example 12.4.2) but $|\mathbf{i}|\,|\mathbf{i}| = 1 \cdot 1 = 1$. In fact, by Theorem 12.4.9, $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}| \sin\theta$.

5. True, by Theorem 12.3.2, property 2.

6. False. Property 1 of Theorem 12.4.11 says that $\mathbf{u} \times \mathbf{v} = -\mathbf{v} \times \mathbf{u}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. True. If θ is the angle between $\mathbf{u}$ and $\mathbf{v}$, then by Theorem 12.4.9, $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}| \sin\theta = |\mathbf{v}|\,|\mathbf{u}| \sin\theta = |\mathbf{v} \times \mathbf{u}|$. (Or, by Theorem 12.4.11, $|\mathbf{u} \times \mathbf{v}| = |-\mathbf{v} \times \mathbf{u}| = |-1|\,|\mathbf{v} \times \mathbf{u}| = |\mathbf{v} \times \mathbf{u}|$.)

8. This is true by Theorem 12.3.2, property 4.

9. Theorem 12.4.11, property 2 tells us that this is true.

10. This is true by Theorem 12.4.11, property 4.

11. This is true by Theorem 12.4.11, property 5.

12. In general, this assertion is false; a counterexample is $\mathbf{i} \times (\mathbf{i} \times \mathbf{j}) \neq (\mathbf{i} \times \mathbf{i}) \times \mathbf{j}$. (See the paragraph preceding Theorem 12.4.11.)

13. This is true because $\mathbf{u} \times \mathbf{v}$ is orthogonal to $\mathbf{u}$ (see Theorem 12.4.8), and the dot product of two orthogonal vectors is 0.

14.
$$\begin{aligned}(\mathbf{u}+\mathbf{v}) \times \mathbf{v} &= \mathbf{u} \times \mathbf{v} + \mathbf{v} \times \mathbf{v} && \text{[by Theorem 12.4.11, property 4]}\\ &= \mathbf{u} \times \mathbf{v} + \mathbf{0} && \text{[by Example 12.4.2]}\\ &= \mathbf{u} \times \mathbf{v}, \text{ so this is true.}\end{aligned}$$

15. This is false. A normal vector to the plane is $\mathbf{n} = \langle 6, -2, 4 \rangle$. Because $\langle 3, -1, 2 \rangle = \frac{1}{2}\mathbf{n}$, the vector is parallel to $\mathbf{n}$ and hence perpendicular to the plane.

16. This is false, because according to Equation 12.5.8, $ax + by + cz + d = 0$ is the general equation of a plane.

17. This is false. In $\mathbb{R}^2$, $x^2 + y^2 = 1$ represents a circle, but $\{(x, y, z) \mid x^2 + y^2 = 1\}$ represents a *three-dimensional surface*, namely, a circular cylinder with axis the z-axis.

18. This is false. In $\mathbb{R}^3$ the graph of $y = x^2$ is a parabolic cylinder (see Example 12.6.1). A paraboloid has an equation such as $z = x^2 + y^2$.

19. False. For example, $\mathbf{i} \cdot \mathbf{j} = \mathbf{0}$ but $\mathbf{i} \neq \mathbf{0}$ and $\mathbf{j} \neq \mathbf{0}$.

20. This is false. By Corollary 12.4.10, $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ for any nonzero parallel vectors $\mathbf{u}$, $\mathbf{v}$. For instance, $\mathbf{i} \times \mathbf{i} = \mathbf{0}$.

21. This is true. If $\mathbf{u}$ and $\mathbf{v}$ are both nonzero, then by (7) in Section 12.3, $\mathbf{u} \cdot \mathbf{v} = 0$ implies that $\mathbf{u}$ and $\mathbf{v}$ are orthogonal. But $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ implies that $\mathbf{u}$ and $\mathbf{v}$ are parallel (see Corollary 12.4.10). Two nonzero vectors can't be both parallel and orthogonal, so at least one of $\mathbf{u}$, $\mathbf{v}$ must be $\mathbf{0}$.

22. This is true. We know $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}|\ |\mathbf{v}|\ \cos\theta$ where $|\mathbf{u}| \geq 0$, $|\mathbf{v}| \geq 0$, and $|\cos\theta| \leq 1$, so $|\mathbf{u} \cdot \mathbf{v}| = |\mathbf{u}|\ |\mathbf{v}|\ |\cos\theta| \leq |\mathbf{u}|\ |\mathbf{v}|$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

EXERCISES

1. (a) The radius of the sphere is the distance between the points $(-1, 2, 1)$ and $(6, -2, 3)$, namely, $\sqrt{[6-(-1)]^2 + (-2-2)^2 + (3-1)^2} = \sqrt{69}$. By the formula for an equation of a sphere (following Example 12.1.4), an equation of the sphere with center $(-1, 2, 1)$ and radius $\sqrt{69}$ is $(x+1)^2 + (y-2)^2 + (z-1)^2 = 69$.

(b) The intersection of this sphere with the yz-plane is the set of points on the sphere whose x-coordinate is 0. Putting $x = 0$ into the equation, we have $(y-2)^2 + (z-1)^2 = 68, x = 0$ which represents a circle in the yz-plane with center $(0, 2, 1)$ and radius $\sqrt{68}$.

(c) Completing squares gives $(x-4)^2 + (y+1)^2 + (z+3)^2 = -1 + 16 + 1 + 9 = 25$. Thus the sphere is centered at $(4, -1, -3)$ and has radius 5.

2. (a)

(b)

(c)

(d)

3. $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}|\,|\mathbf{v}| \cos 45° = (2)(3)\frac{\sqrt{2}}{2} = 3\sqrt{2}$. $\quad |\mathbf{u} \times \mathbf{v}| = |\mathbf{u}|\,|\mathbf{v}| \sin 45° = (2)(3)\frac{\sqrt{2}}{2} = 3\sqrt{2}$.
By the right-hand rule, $\mathbf{u} \times \mathbf{v}$ is directed out of the page.

4. (a) $2\,\mathbf{a} + 3\,\mathbf{b} = 2\,\mathbf{i} + 2\,\mathbf{j} - 4\,\mathbf{k} + 9\,\mathbf{i} - 6\,\mathbf{j} + 3\,\mathbf{k} = 11\,\mathbf{i} - 4\,\mathbf{j} - \mathbf{k}$

(b) $|\mathbf{b}| = \sqrt{9+4+1} = \sqrt{14}$

(c) $\mathbf{a} \cdot \mathbf{b} = (1)(3) + (1)(-2) + (-2)(1) = -1$

(d) $\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & -2 \\ 3 & -2 & 1 \end{vmatrix} = (1-4)\,\mathbf{i} - (1+6)\,\mathbf{j} + (-2-3)\,\mathbf{k} = -3\,\mathbf{i} - 7\,\mathbf{j} - 5\,\mathbf{k}$

(e) $\mathbf{b} \times \mathbf{c} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -2 & 1 \\ 0 & 1 & -5 \end{vmatrix} = 9\,\mathbf{i} + 15\,\mathbf{j} + 3\,\mathbf{k}, \quad |\mathbf{b} \times \mathbf{c}| = 3\sqrt{9+25+1} = 3\sqrt{35}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(f) $\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c}) = \begin{vmatrix} 1 & 1 & -2 \\ 3 & -2 & 1 \\ 0 & 1 & -5 \end{vmatrix} = \begin{vmatrix} -2 & 1 \\ 1 & -5 \end{vmatrix} - \begin{vmatrix} 3 & 1 \\ 0 & -5 \end{vmatrix} - 2\begin{vmatrix} 3 & -2 \\ 0 & 1 \end{vmatrix} = 9 + 15 - 6 = 18$

(g) $\mathbf{c}\times\mathbf{c} = \mathbf{0}$ for any $\mathbf{c}$.

(h) From part (e),

$$\mathbf{a}\times(\mathbf{b}\times\mathbf{c}) = \mathbf{a}\times(9\,\mathbf{i}+15\,\mathbf{j}+3\,\mathbf{k}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & -2 \\ 9 & 15 & 3 \end{vmatrix}$$

$$= (3+30)\,\mathbf{i} - (3+18)\,\mathbf{j} + (15-9)\,\mathbf{k} = 33\,\mathbf{i} - 21\,\mathbf{j} + 6\,\mathbf{k}$$

(i) The scalar projection is $\operatorname{comp}_{\mathbf{a}}\mathbf{b} = |\mathbf{b}|\cos\theta = \mathbf{a}\cdot\mathbf{b}/\,|\mathbf{a}| = -\frac{1}{\sqrt{6}}$.

(j) The vector projection is $\operatorname{proj}_{\mathbf{a}}\mathbf{b} = -\dfrac{1}{\sqrt{6}}\left(\dfrac{\mathbf{a}}{|\mathbf{a}|}\right) = -\frac{1}{6}(\mathbf{i}+\mathbf{j}-2\,\mathbf{k})$.

(k) $\cos\theta = \dfrac{\mathbf{a}\cdot\mathbf{b}}{|\mathbf{a}|\,|\mathbf{b}|} = \dfrac{-1}{\sqrt{6}\sqrt{14}} = \dfrac{-1}{2\sqrt{21}}$ and $\theta = \cos^{-1}\left(\dfrac{-1}{2\sqrt{21}}\right) \approx 96^\circ$.

5. For the two vectors to be orthogonal, we need $\langle 3,2,x\rangle\cdot\langle 2x,4,x\rangle = 0 \quad\Leftrightarrow\quad (3)(2x)+(2)(4)+(x)(x) = 0 \quad\Leftrightarrow$ $x^2+6x+8=0 \quad\Leftrightarrow\quad (x+2)(x+4)=0 \quad\Leftrightarrow\quad x=-2$ or $x=-4$.

6. We know that the cross product of two vectors is orthogonal to both given vectors. So we calculate $(\mathbf{j}+2\,\mathbf{k})\times(\mathbf{i}-2\,\mathbf{j}+3\,\mathbf{k}) = [3-(-4)]\,\mathbf{i} - (0-2)\,\mathbf{j} + (0-1)\,\mathbf{k} = 7\,\mathbf{i}+2\,\mathbf{j}-\mathbf{k}$.

Then two unit vectors orthogonal to both given vectors are $\pm\dfrac{7\,\mathbf{i}+2\,\mathbf{j}-\mathbf{k}}{\sqrt{7^2+2^2+(-1)^2}} = \pm\dfrac{1}{3\sqrt{6}}\,(7\,\mathbf{i}+2\,\mathbf{j}-\mathbf{k})$,

that is, $\frac{7}{3\sqrt{6}}\,\mathbf{i} + \frac{2}{3\sqrt{6}}\,\mathbf{j} - \frac{1}{3\sqrt{6}}\,\mathbf{k}$ and $-\frac{7}{3\sqrt{6}}\,\mathbf{i} - \frac{2}{3\sqrt{6}}\,\mathbf{j} + \frac{1}{3\sqrt{6}}\,\mathbf{k}$.

7. (a) $(\mathbf{u}\times\mathbf{v})\cdot\mathbf{w} = \mathbf{u}\cdot(\mathbf{v}\times\mathbf{w}) = 2$

(b) $\mathbf{u}\cdot(\mathbf{w}\times\mathbf{v}) = \mathbf{u}\cdot[-(\mathbf{v}\times\mathbf{w})] = -\mathbf{u}\cdot(\mathbf{v}\times\mathbf{w}) = -2$

(c) $\mathbf{v}\cdot(\mathbf{u}\times\mathbf{w}) = (\mathbf{v}\times\mathbf{u})\cdot\mathbf{w} = -(\mathbf{u}\times\mathbf{v})\cdot\mathbf{w} = -2$

(d) $(\mathbf{u}\times\mathbf{v})\cdot\mathbf{v} = \mathbf{u}\cdot(\mathbf{v}\times\mathbf{v}) = \mathbf{u}\cdot\mathbf{0} = 0$

8. $(\mathbf{a}\times\mathbf{b})\cdot[(\mathbf{b}\times\mathbf{c})\times(\mathbf{c}\times\mathbf{a})] = (\mathbf{a}\times\mathbf{b})\cdot([(\mathbf{b}\times\mathbf{c})\cdot\mathbf{a}]\,\mathbf{c} - [(\mathbf{b}\times\mathbf{c})\cdot\mathbf{c}]\,\mathbf{a})$

[by Property 6 of the cross product]

$$= (\mathbf{a}\times\mathbf{b})\cdot[(\mathbf{b}\times\mathbf{c})\cdot\mathbf{a}]\,\mathbf{c} = [\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c})]\,(\mathbf{a}\times\mathbf{b})\cdot\mathbf{c}$$

$$= [\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c})]\,[\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c})] = [\mathbf{a}\cdot(\mathbf{b}\times\mathbf{c})]^2$$

9. For simplicity, consider a unit cube positioned with its back left corner at the origin. Vector representations of the diagonals joining the points $(0,0,0)$ to $(1,1,1)$ and $(1,0,0)$ to $(0,1,1)$ are $\langle 1,1,1\rangle$ and $\langle -1,1,1\rangle$. Let θ be the angle between these two vectors. $\langle 1,1,1\rangle\cdot\langle -1,1,1\rangle = -1+1+1 = 1 = |\langle 1,1,1\rangle|\,|\langle -1,1,1\rangle|\cos\theta = 3\cos\theta \quad\Rightarrow\quad \cos\theta = \frac{1}{3} \quad\Rightarrow$ $\theta = \cos^{-1}\left(\frac{1}{3}\right) \approx 71^\circ$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $\overrightarrow{AB} = \langle 1, 3, -1 \rangle$, $\overrightarrow{AC} = \langle -2, 1, 3 \rangle$ and $\overrightarrow{AD} = \langle -1, 3, 1 \rangle$. By Equation 12.4.13,

$$\overrightarrow{AB} \cdot \left(\overrightarrow{AC} \times \overrightarrow{AD} \right) = \begin{vmatrix} 1 & 3 & -1 \\ -2 & 1 & 3 \\ -1 & 3 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 3 \\ 3 & 1 \end{vmatrix} - 3 \begin{vmatrix} -2 & 3 \\ -1 & 1 \end{vmatrix} - \begin{vmatrix} -2 & 1 \\ -1 & 3 \end{vmatrix} = -8 - 3 + 5 = -6.$$

The volume is $\left| \overrightarrow{AB} \cdot \left(\overrightarrow{AC} \times \overrightarrow{AD} \right) \right| = 6$ cubic units.

11. $\overrightarrow{AB} = \langle 1, 0, -1 \rangle$, $\overrightarrow{AC} = \langle 0, 4, 3 \rangle$, so

(a) a vector perpendicular to the plane is $\overrightarrow{AB} \times \overrightarrow{AC} = \langle 0 + 4, -(3 + 0), 4 - 0 \rangle = \langle 4, -3, 4 \rangle$.

(b) $\frac{1}{2} \left| \overrightarrow{AB} \times \overrightarrow{AC} \right| = \frac{1}{2} \sqrt{16 + 9 + 16} = \frac{\sqrt{41}}{2}$.

12. $\mathbf{D} = (5 - 1)\,\mathbf{i} + (3 - 0)\,\mathbf{j} + (8 - 2)\,\mathbf{k} = 4\,\mathbf{i} + 3\,\mathbf{j} + 6\,\mathbf{k}$.

$W = \mathbf{F} \cdot \mathbf{D} = (3\,\mathbf{i} + 5\,\mathbf{j} + 10\,\mathbf{k}) \cdot (4\,\mathbf{i} + 3\,\mathbf{j} + 6\,\mathbf{k}) = (3)(4) + (5)(3) + (10)(6) = 12 + 15 + 60 = 87$ J

13. Let F_1 be the magnitude of the force directed $20°$ away from the direction of shore, and let F_2 be the magnitude of the other force. Separating these forces into components parallel to the direction of the resultant force and perpendicular to it gives

$F_1 \cos 20° + F_2 \cos 30° = 255$ **(1)**, and $F_1 \sin 20° - F_2 \sin 30° = 0 \quad \Rightarrow \quad F_1 = F_2 \dfrac{\sin 30°}{\sin 20°}$ **(2)**. Substituting **(2)** into **(1)** gives $F_2(\sin 30° \cot 20° + \cos 30°) = 255 \quad \Rightarrow \quad F_2 \approx 114$ N. Substituting this into **(2)** gives $F_1 \approx 166$ N.

14. $|\boldsymbol{\tau}| = |\mathbf{r}|\,|\mathbf{F}| \sin\theta = (0.40)(50)\sin(90° - 30°) \approx 17.3$ N·m.

15. The line has direction $\mathbf{v} = \langle -3, 2, 3 \rangle$. Letting $P_0 = (4, -1, 2)$, parametric equations are $x = 4 - 3t,\ y = -1 + 2t,\ z = 2 + 3t$.

16. The line $\frac{1}{3}(x - 4) = \frac{1}{2}y = z + 2$, or $\dfrac{x = 4}{3} = \dfrac{y}{2} = \dfrac{z + 2}{1}$, has direction vector $\mathbf{v} = \langle 3, 2, 1 \rangle$ (or a nonzero scalar multiple). So parametric equations for the line through $(1, 0, -1)$ are $x = 1 + 3t,\ y = 2t,\ z = -1 + t$.

17. A direction vector for the line is a normal vector for the plane, $\mathbf{n} = \langle 2, -1, 5 \rangle$, and parametric equations for the line are $x = -2 + 2t,\ y = 2 - t,\ z = 4 + 5t$.

18. Since the two planes are parallel, they will have the same normal vectors. Then we can take $\mathbf{n} = \langle 1, 4, -3 \rangle$ and an equation of the plane is $1(x - 2) + 4(y - 1) - 3(z - 0) = 0$ or $x + 4y - 3z = 6$.

19. Here the vectors $\mathbf{a} = \langle 4 - 3, 0 - (-1), 2 - 1 \rangle = \langle 1, 1, 1 \rangle$ and $\mathbf{b} = \langle 6 - 3, 3 - (-1), 1 - 1 \rangle = \langle 3, 4, 0 \rangle$ lie in the plane, so $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle -4, 3, 1 \rangle$ is a normal vector to the plane and an equation of the plane is $-4(x - 3) + 3(y - (-1)) + 1(z - 1) = 0$ or $-4x + 3y + z = -14$.

20. If we first find two nonparallel vectors in the plane, their cross product will be a normal vector to the plane. Since the given line lies in the plane, its direction vector $\mathbf{a} = \langle 2, -1, 3 \rangle$ is one vector in the plane. We can verify that the given point $(1, 2, -2)$ does not lie on this line. The point $(0, 3, 1)$ is on the line (obtained by putting $t = 0$) and hence in the plane, so the vector $\mathbf{b} = \langle 0 - 1, 3 - 2, 1 - (-2) \rangle = \langle -1, 1, 3 \rangle$ lies in the plane, and a normal vector is $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \langle -6, -9, 1 \rangle$. Thus an equation of the plane is $-6(x - 1) - 9(y - 2) + (z + 2) = 0$ or $6x + 9y - z = 26$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. Substitution of the parametric equations into the equation of the plane gives $2x - y + z = 2(2-t) - (1+3t) + 4t = 2 \quad \Rightarrow$ $-t + 3 = 2 \quad \Rightarrow \quad t = 1$. When $t = 1$, the parametric equations give $x = 2 - 1 = 1$, $y = 1 + 3 = 4$ and $z = 4$. Therefore, the point of intersection is $(1, 4, 4)$.

22. Use the formula proven in Exercise 12.4.45(a). In the notation used in that exercise, $\mathbf{a}$ is just the direction of the line; that is, $\mathbf{a} = \langle 1, -1, 2\rangle$. A point on the line is $(1, 2, -1)$ (setting $t = 0$), and therefore $\mathbf{b} = \langle 1-0, 2-0, -1-0\rangle = \langle 1, 2, -1\rangle$.

Hence $d = \dfrac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}|} = \dfrac{|\langle 1,-1,2\rangle \times \langle 1,2,-1\rangle|}{\sqrt{1+1+4}} = \dfrac{|\langle -3,3,3\rangle|}{\sqrt{6}} = \sqrt{\dfrac{27}{6}} = \dfrac{3}{\sqrt{2}}$.

23. Since the direction vectors $\langle 2, 3, 4\rangle$ and $\langle 6, -1, 2\rangle$ aren't parallel, neither are the lines. For the lines to intersect, the three equations $1 + 2t = -1 + 6s$, $2 + 3t = 3 - s$, $3 + 4t = -5 + 2s$ must be satisfied simultaneously. Solving the first two equations gives $t = \frac{1}{5}$, $s = \frac{2}{5}$ and checking we see these values don't satisfy the third equation. Thus the lines aren't parallel and they don't intersect, so they must be skew.

24. (a) The normal vectors are $\langle 1, 1, -1\rangle$ and $\langle 2, -3, 4\rangle$. Since these vectors aren't parallel, neither are the planes parallel. Also $\langle 1, 1, -1\rangle \cdot \langle 2, -3, 4\rangle = 2 - 3 - 4 = -5 \neq 0$ so the normal vectors, and thus the planes, are not perpendicular.

(b) $\cos\theta = \dfrac{\langle 1,1,-1\rangle \cdot \langle 2,-3,4\rangle}{\sqrt{3}\sqrt{29}} = -\dfrac{5}{\sqrt{87}}$ and $\theta = \cos^{-1}\left(-\frac{5}{\sqrt{87}}\right) \approx 122^\circ$ [or we can say $\approx 58^\circ$].

25. $\mathbf{n}_1 = \langle 1, 0, -1\rangle$ and $\mathbf{n}_2 = \langle 0, 1, 2\rangle$. Setting $z = 0$, it is easy to see that $(1, 3, 0)$ is a point on the line of intersection of $x - z = 1$ and $y + 2z = 3$. The direction of this line is $\mathbf{v}_1 = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, -2, 1\rangle$. A second vector parallel to the desired plane is $\mathbf{v}_2 = \langle 1, 1, -2\rangle$, since it is perpendicular to $x + y - 2z = 1$. Therefore, the normal of the plane in question is $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \langle 4-1, 1+2, 1+2\rangle = 3\langle 1, 1, 1\rangle$. Taking $(x_0, y_0, z_0) = (1, 3, 0)$, the equation we are looking for is $(x-1) + (y-3) + z = 0 \quad \Leftrightarrow \quad x + y + z = 4$.

26. (a) The vectors $\overrightarrow{AB} = \langle -1-2, -1-1, 10-1\rangle = \langle -3, -2, 9\rangle$ and $\overrightarrow{AC} = \langle 1-2, 3-1, -4-1\rangle = \langle -1, 2, -5\rangle$ lie in the plane, so $\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \langle -3,-2,9\rangle \times \langle -1,2,-5\rangle = \langle -8,-24,-8\rangle$ or equivalently $\langle 1, 3, 1\rangle$ is a normal vector to the plane. The point $A(2, 1, 1)$ lies on the plane so an equation of the plane is $1(x-2) + 3(y-1) + 1(z-1) = 0$ or $x + 3y + z = 6$.

(b) The line is perpendicular to the plane so it is parallel to a normal vector for the plane, namely $\langle 1, 3, 1\rangle$. If the line passes through $B(-1, -1, 10)$ then symmetric equations are $\dfrac{x-(-1)}{1} = \dfrac{y-(-1)}{3} = \dfrac{z-10}{1}$ or $x + 1 = \dfrac{y+1}{3} = z - 10$.

(c) Normal vectors for the two planes are $\mathbf{n}_1 = \langle 1, 3, 1\rangle$ and $\mathbf{n}_2 = \langle 2, -4, -3\rangle$. The angle θ between the planes is given by

$$\cos\theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1|\ |\mathbf{n}_2|} = \frac{\langle 1,3,1\rangle \cdot \langle 2,-4,-3\rangle}{\sqrt{1^2+3^2+1^2}\sqrt{2^2+(-4)^2+(-3)^2}} = \frac{2-12-3}{\sqrt{11}\sqrt{29}} = -\frac{13}{\sqrt{319}}$$

Thus $\theta = \cos^{-1}\left(-\dfrac{13}{\sqrt{319}}\right) \approx 137^\circ$ or $180^\circ - 137^\circ = 43^\circ$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) From part (c), the point $(2, 0, 4)$ lies on the second plane, but notice that the point also satisfies the equation of the first plane, so the point lies on the line of intersection of the planes. A vector $\mathbf{v}$ in the direction of this intersecting line is perpendicular to the normal vectors of both planes, so take $\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \langle 1, 3, 1\rangle \times \langle 2, -4, -3\rangle = \langle -5, 5, -10\rangle$ or equivalently we can take $\mathbf{v} = \langle 1, -1, 2\rangle$. Parametric equations for the line are $x = 2 + t$, $y = -t$, $z = 4 + 2t$.

27. By Exercise 12.5.75, $D = \dfrac{|-2 - (-24)|}{\sqrt{3^2 + 1^2 + (-4)^2}} = \dfrac{22}{\sqrt{26}}$.

28. The equation $x = 3$ represents a plane parallel to the yz-plane and 3 units in front of it.

29. The equation $x = z$ represents a plane perpendicular to the xz-plane and intersecting the xz-plane in the line $x = z$, $y = 0$.

30. The equation $y = z^2$ represents a parabolic cylinder whose trace in the xz-plane is the x-axis and which opens to the right.

31. The equation $x^2 = y^2 + 4z^2$ represents a (right elliptical) cone with vertex at the origin and axis the x-axis.

32. $4x - y + 2z = 4$ is a plane with intercepts $(1, 0, 0)$, $(0, -4, 0)$, and $(0, 0, 2)$.

33. An equivalent equation is $-x^2 + \dfrac{y^2}{4} - z^2 = 1$, a hyperboloid of two sheets with axis the y-axis. For $|y| > 2$, traces parallel to the xz-plane are circles.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. An equivalent equation is $-x^2 + y^2 + z^2 = 1$, a hyperboloid of one sheet with axis the x-axis.

35. Completing the square in y gives

$4x^2 + 4(y-1)^2 + z^2 = 4$ or $x^2 + (y-1)^2 + \dfrac{z^2}{4} = 1$,

an ellipsoid centered at $(0, 1, 0)$.

36. Completing the square for $x = y^2 + z^2 - 2y - 4z + 5$ in y and z gives $x = (y-1)^2 + (z-2)^2$, which is a circular paraboloid with vertex $(0, 1, 2)$ and axis the horizontal line $y = 1$, $z = 2$.

37. $4x^2 + y^2 = 16 \quad \Leftrightarrow \quad \dfrac{x^2}{4} + \dfrac{y^2}{16} = 1$. The equation of the ellipsoid is $\dfrac{x^2}{4} + \dfrac{y^2}{16} + \dfrac{z^2}{c^2} = 1$, since the horizontal trace in the plane $z = 0$ must be the original ellipse. The traces of the ellipsoid in the yz-plane must be circles since the surface is obtained by rotation about the x-axis. Therefore, $c^2 = 16$ and the equation of the ellipsoid is $\dfrac{x^2}{4} + \dfrac{y^2}{16} + \dfrac{z^2}{16} = 1 \quad \Leftrightarrow$

$4x^2 + y^2 + z^2 = 16$.

38. The distance from a point $P(x, y, z)$ to the plane $y = 1$ is $|y - 1|$, so the given condition becomes

$|y-1| = 2\sqrt{(x-0)^2 + (y+1)^2 + (z-0)^2} \quad \Rightarrow \quad |y-1| = 2\sqrt{x^2 + (y+1)^2 + z^2} \quad \Rightarrow$

$(y-1)^2 = 4x^2 + 4(y+1)^2 + 4z^2 \quad \Leftrightarrow \quad -3 = 4x^2 + (3y^2 + 10y) + 4z^2 \quad \Leftrightarrow$

$\frac{16}{3} = 4x^2 + 3\left(y + \frac{5}{3}\right)^2 + 4z^2 \quad \Rightarrow \quad \frac{3}{4}x^2 + \frac{9}{16}\left(y + \frac{5}{3}\right)^2 + \frac{3}{4}z^2 = 1.$

This is the equation of an ellipsoid whose center is $\left(0, -\frac{5}{3}, 0\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

PROBLEMS PLUS

1. Since three-dimensional situations are often difficult to visualize and work with, let us first try to find an analogous problem in two dimensions. The analogue of a cube is a square and the analogue of a sphere is a circle. Thus a similar problem in two dimensions is the following: if five circles with the same radius r are contained in a square of side 1 m so that the circles touch each other and four of the circles touch two sides of the square, find r.

The diagonal of the square is $\sqrt{2}$. The diagonal is also $4r + 2x$. But x is the diagonal of a smaller square of side r. Therefore

$x = \sqrt{2}\,r \quad \Rightarrow \quad \sqrt{2} = 4r + 2x = 4r + 2\sqrt{2}\,r = \left(4 + 2\sqrt{2}\right)r \quad \Rightarrow \quad r = \frac{\sqrt{2}}{4 + 2\sqrt{2}}$.

Let's use these ideas to solve the original three-dimensional problem. The diagonal of the cube is $\sqrt{1^2 + 1^2 + 1^2} = \sqrt{3}$. The diagonal of the cube is also $4r + 2x$ where x is the diagonal of a smaller cube with edge r. Therefore

$x = \sqrt{r^2 + r^2 + r^2} = \sqrt{3}\,r \quad \Rightarrow \quad \sqrt{3} = 4r + 2x = 4r + 2\sqrt{3}\,r = \left(4 + 2\sqrt{3}\right)r$. Thus $r = \dfrac{\sqrt{3}}{4 + 2\sqrt{3}} = \dfrac{2\sqrt{3} - 3}{2}$.

The radius of each ball is $\left(\sqrt{3} - \frac{3}{2}\right)$ m.

2. Try an analogous problem in two dimensions. Consider a rectangle with length L and width W and find the area of S in terms of L and W. Since S contains B, it has area

$$A(S) = LW + \text{ the area of two } L \times 1 \text{ rectangles}$$
$$+ \text{ the area of two } 1 \times W \text{ rectangles}$$
$$+ \text{ the area of four quarter-circles of radius } 1$$

as seen in the diagram. So $A(S) = LW + 2L + 2W + \pi \cdot 1^2$.

Now in three dimensions, the volume of S is

$$LWH + 2(L \times W \times 1) + 2(1 \times W \times H) + 2(L \times 1 \times H)$$
$$+ \text{ the volume of 4 quarter-cylinders with radius 1 and height } W$$
$$+ \text{ the volume of 4 quarter-cylinders with radius 1 and height } L$$
$$+ \text{ the volume of 4 quarter-cylinders with radius 1 and height } H$$
$$+ \text{ the volume of 8 eighths of a sphere of radius 1}$$

So

$$V(S) = LWH + 2LW + 2WH + 2LH + \pi \cdot 1^2 \cdot W + \pi \cdot 1^2 \cdot L + \pi \cdot 1^2 \cdot H + \tfrac{4}{3}\pi \cdot 1^3$$
$$= LWH + 2(LW + WH + LH) + \pi(L + W + H) + \tfrac{4}{3}\pi.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a) We find the line of intersection L as in Example 12.5.7(b). Observe that the point $(-1, c, c)$ lies on both planes. Now since L lies in both planes, it is perpendicular to both of the normal vectors $\mathbf{n}_1$ and $\mathbf{n}_2$, and thus parallel to their cross product

$$\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ c & 1 & 1 \\ 1 & -c & c \end{vmatrix} = \langle 2c, -c^2+1, -c^2-1 \rangle$$

So symmetric equations of L can be written as $\dfrac{x+1}{-2c} = \dfrac{y-c}{c^2-1} = \dfrac{z-c}{c^2+1}$, provided that $c \neq 0, \pm 1$.

If $c = 0$, then the two planes are given by $y + z = 0$ and $x = -1$, so symmetric equations of L are $x = -1$, $y = -z$. If $c = -1$, then the two planes are given by $-x + y + z = -1$ and $x + y + z = -1$, and they intersect in the line $x = 0$, $y = -z - 1$. If $c = 1$, then the two planes are given by $x + y + z = 1$ and $x - y + z = 1$, and they intersect in the line $y = 0$, $x = 1 - z$.

(b) If we set $z = t$ in the symmetric equations and solve for x and y separately, we get $x + 1 = \dfrac{(t-c)(-2c)}{c^2+1}$, $y - c = \dfrac{(t-c)(c^2-1)}{c^2+1} \quad \Rightarrow \quad x = \dfrac{-2ct + (c^2-1)}{c^2+1}$, $y = \dfrac{(c^2-1)t + 2c}{c^2+1}$. Eliminating c from these equations, we have $x^2 + y^2 = t^2 + 1$. So the curve traced out by L in the plane $z = t$ is a circle with center at $(0, 0, t)$ and radius $\sqrt{t^2+1}$.

(c) The area of a horizontal cross-section of the solid is $A(z) = \pi(z^2+1)$, so $V = \int_0^1 A(z)\,dz = \pi\left[\frac{1}{3}z^3 + z\right]_0^1 = \frac{4\pi}{3}$.

4. (a) We consider velocity vectors for the plane and the wind. Let $\mathbf{v}_i$ be the initial, intended velocity for the plane and $\mathbf{v}_g$ the actual velocity relative to the ground. If $\mathbf{w}$ is the velocity of the wind, $\mathbf{v}_g$ is the resultant, that is, the vector sum $\mathbf{v}_i + \mathbf{w}$ as shown in the figure. We know $\mathbf{v}_i = 180\,\mathbf{j}$, and since the plane actually flew 80 km in $\frac{1}{2}$ hour, $|\mathbf{v}_g| = 160$. Thus $\mathbf{v}_g = (160\cos 85^\circ)\,\mathbf{i} + (160\sin 85^\circ)\,\mathbf{j} \approx 13.9\,\mathbf{i} + 159.4\,\mathbf{j}$. Finally, $\mathbf{v}_i + \mathbf{w} = \mathbf{v}_g$, so $\mathbf{w} = \mathbf{v}_g - \mathbf{v}_i \approx 13.9\,\mathbf{i} - 20.6\,\mathbf{j}$. Thus, the wind velocity is about $13.9\,\mathbf{i} - 20.6\,\mathbf{j}$, and the wind speed is $|\mathbf{w}| \approx \sqrt{(13.9)^2 + (-20.6)^2} \approx 24.9$ km/h.

(b) Let $\mathbf{v}$ be the velocity the pilot should have taken. The pilot would need to head a little west of north to compensate for the wind, so let θ be the angle $\mathbf{v}$ makes with the north direction. Then we can write $\mathbf{v} = \langle 180\cos(\theta + 90^\circ), 180\sin(\theta + 90^\circ) \rangle$. With the effect of the wind, the actual velocity (with respect to the ground) will be $\mathbf{v} + \mathbf{w}$, which we want to be due north. Equating horizontal components of the vectors, we must have $180\cos(\theta + 90^\circ) + 160\cos 85^\circ = 0 \quad \Rightarrow \quad \cos(\theta + 90^\circ) = -\frac{160}{180}\cos 85^\circ \approx -0.0775$, so $\theta \approx \cos^{-1}(-0.0775) - 90^\circ \approx 4.4^\circ$. Thus the pilot should have headed about 4.4° west of north.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $\mathbf{v}_3 = \text{proj}_{\mathbf{v}_1}\mathbf{v}_2 = \dfrac{\mathbf{v}_1 \cdot \mathbf{v}_2}{|\mathbf{v}_1|^2}\,\mathbf{v}_1 = \dfrac{5}{2^2}\,\mathbf{v}_1$ so $|\mathbf{v}_3| = \dfrac{5}{2^2}\,|\mathbf{v}_1| = \dfrac{5}{2}$,

$\mathbf{v}_4 = \text{proj}_{\mathbf{v}_2}\mathbf{v}_3 = \dfrac{\mathbf{v}_2 \cdot \mathbf{v}_3}{|\mathbf{v}_2|^2}\,\mathbf{v}_2 = \dfrac{\mathbf{v}_2 \cdot \frac{5}{2^2}\,\mathbf{v}_1}{|\mathbf{v}_2|^2}\,\mathbf{v}_2 = \dfrac{5}{2^2 \cdot 3^2}\,(\mathbf{v}_1 \cdot \mathbf{v}_2)\,\mathbf{v}_2 = \dfrac{5^2}{2^2 \cdot 3^2}\,\mathbf{v}_2 \quad \Rightarrow \quad |\mathbf{v}_4| = \dfrac{5^2}{2^2 \cdot 3^2}\,|\mathbf{v}_2| = \dfrac{5^2}{2^2 \cdot 3}$,

$\mathbf{v}_5 = \text{proj}_{\mathbf{v}_3}\mathbf{v}_4 = \dfrac{\mathbf{v}_3 \cdot \mathbf{v}_4}{|\mathbf{v}_3|^2}\,\mathbf{v}_3 = \dfrac{\frac{5}{2^2}\,\mathbf{v}_1 \cdot \frac{5^2}{2^2\,3^2}\,\mathbf{v}_2}{\left(\frac{5}{2}\right)^2}\left(\dfrac{5}{2^2}\,\mathbf{v}_1\right) = \dfrac{5^2}{2^4 \cdot 3^2}\,(\mathbf{v}_1 \cdot \mathbf{v}_2)\,\mathbf{v}_1 = \dfrac{5^3}{2^4 \cdot 3^2}\,\mathbf{v}_1 \quad \Rightarrow$

$|\mathbf{v}_5| = \dfrac{5^3}{2^4 \cdot 3^2}\,|\mathbf{v}_1| = \dfrac{5^3}{2^3 \cdot 3^2}$. Similarly, $|\mathbf{v}_6| = \dfrac{5^4}{2^4 \cdot 3^3}$, $|\mathbf{v}_7| = \dfrac{5^5}{2^5 \cdot 3^4}$, and in general, $|\mathbf{v}_n| = \dfrac{5^{n-2}}{2^{n-2} \cdot 3^{n-3}} = 3\left(\frac{5}{6}\right)^{n-2}$.

Thus

$$\sum_{n=1}^{\infty} |\mathbf{v}_n| = |\mathbf{v}_1| + |\mathbf{v}_2| + \sum_{n=3}^{\infty} 3\left(\tfrac{5}{6}\right)^{n-2} = 2 + 3 + \sum_{n=1}^{\infty} 3\left(\tfrac{5}{6}\right)^{n}$$

$$= 5 + \sum_{n=1}^{\infty} \tfrac{5}{2}\left(\tfrac{5}{6}\right)^{n-1} = 5 + \frac{\frac{5}{2}}{1 - \frac{5}{6}} \quad \text{[sum of a geometric series]} \quad = 5 + 15 = 20$$

6. Completing squares in the inequality $x^2 + y^2 + z^2 < 136 + 2(x + 2y + 3z)$ gives $(x-1)^2 + (y-2)^2 + (z-3)^2 < 150$ which describes the set of all points (x, y, z) whose distance from the point $P\,(1, 2, 3)$ is less than $\sqrt{150} = 5\sqrt{6}$. The distance from P to $Q\,(-1, 1, 4)$ is $\sqrt{4+1+1} = \sqrt{6}$, so the largest possible sphere that passes through Q and satisfies the stated conditions extends $5\sqrt{6}$ units in the opposite direction, giving a diameter of $6\sqrt{6}$. (See the figure.)

Thus the radius of the desired sphere is $3\sqrt{6}$, and its center is $3\sqrt{6}$ units from Q in the direction of P. A unit vector in this direction is $\mathbf{u} = \frac{1}{\sqrt{6}}\,\langle 2, 1, -1\rangle$, so starting at $Q(-1, 1, 4)$ and following the vector $3\sqrt{6}\,\mathbf{u} = \langle 6, 3, -3\rangle$ we arrive at the center of the sphere, $(5, 4, 1)$. An equation of the sphere then is $(x-5)^2 + (y-4)^2 + (z-1)^2 = \left(3\sqrt{6}\right)^2$ or $(x-5)^2 + (y-4)^2 + (z-1)^2 = 54$.

7. (a) When $\theta = \theta_s$, the block is not moving, so the sum of the forces on the block must be $\mathbf{0}$, thus $\mathbf{N} + \mathbf{F} + \mathbf{W} = \mathbf{0}$. This relationship is illustrated geometrically in the figure. Since the vectors form a right triangle, we have $\tan(\theta_s) = \dfrac{|\mathbf{F}|}{|\mathbf{N}|} = \dfrac{\mu_s n}{n} = \mu_s$.

(b) We place the block at the origin and sketch the force vectors acting on the block, including the additional horizontal force $\mathbf{H}$, with initial points at the origin. We then rotate this system so that $\mathbf{F}$ lies along the positive x-axis and the inclined plane is parallel to the x-axis. (See the following figure.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$|\mathbf{F}|$ is maximal, so $|\mathbf{F}| = \mu_s n$ for $\theta > \theta_s$. Then the vectors, in terms of components parallel and perpendicular to the inclined plane, are

$$\mathbf{N} = n\,\mathbf{j} \qquad \mathbf{F} = (\mu_s n)\,\mathbf{i}$$

$$\mathbf{W} = (-mg\sin\theta)\,\mathbf{i} + (-mg\cos\theta)\,\mathbf{j} \qquad \mathbf{H} = (h_{\min}\cos\theta)\,\mathbf{i} + (-h_{\min}\sin\theta)\,\mathbf{j}$$

Equating components, we have

$$\mu_s n - mg\sin\theta + h_{\min}\cos\theta = 0 \quad\Rightarrow\quad h_{\min}\cos\theta + \mu_s n = mg\sin\theta \qquad \textbf{(1)}$$

$$n - mg\cos\theta - h_{\min}\sin\theta = 0 \quad\Rightarrow\quad h_{\min}\sin\theta + mg\cos\theta = n \qquad \textbf{(2)}$$

(c) Since **(2)** is solved for n, we substitute into **(1)**:

$$h_{\min}\cos\theta + \mu_s(h_{\min}\sin\theta + mg\cos\theta) = mg\sin\theta \quad\Rightarrow$$

$$h_{\min}\cos\theta + h_{\min}\mu_s\sin\theta = mg\sin\theta - mg\mu_s\cos\theta \quad\Rightarrow$$

$$h_{\min} = mg\left(\frac{\sin\theta - \mu_s\cos\theta}{\cos\theta + \mu_s\sin\theta}\right) = mg\left(\frac{\tan\theta - \mu_s}{1 + \mu_s\tan\theta}\right)$$

From part (a) we know $\mu_s = \tan\theta_s$, so this becomes $h_{\min} = mg\left(\dfrac{\tan\theta - \tan\theta_s}{1 + \tan\theta_s\tan\theta}\right)$ and using a trigonometric identity, this is $mg\tan(\theta - \theta_s)$ as desired.

Note for $\theta = \theta_s$, $h_{\min} = mg\tan 0 = 0$, which makes sense since the block is at rest for θ_s, thus no additional force $\mathbf{H}$ is necessary to prevent it from moving. As θ increases, the factor $\tan(\theta - \theta_s)$, and hence the value of $h_{\min}$, increases slowly for small values of $\theta - \theta_s$ but much more rapidly as $\theta - \theta_s$ becomes significant. This seems reasonable, as the steeper the inclined plane, the less the horizontal components of the various forces affect the movement of the block, so we would need a much larger magnitude of horizontal force to keep the block motionless. If we allow $\theta \to 90°$, corresponding to the inclined plane being placed vertically, the value of $h_{\min}$ is quite large; this is to be expected, as it takes a great amount of horizontal force to keep an object from moving vertically. In fact, without friction (so $\theta_s = 0$), we would have $\theta \to 90° \quad\Rightarrow\quad h_{\min} \to \infty$, and it would be impossible to keep the block from slipping.

(d) Since $h_{\max}$ is the largest value of h that keeps the block from slipping, the force of friction is keeping the block from moving *up* the inclined plane; thus, $\mathbf{F}$ is directed *down* the plane. Our system of forces is similar to that in part (b), then, except that we have $\mathbf{F} = -(\mu_s n)\,\mathbf{i}$. (Note that $|\mathbf{F}|$ is again maximal.) Following our procedure in parts (b) and (c), we

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

equate components:

$$-\mu_s n - mg\sin\theta + h_{\max}\cos\theta = 0 \quad\Rightarrow\quad h_{\max}\cos\theta - \mu_s n = mg\sin\theta$$

$$n - mg\cos\theta - h_{\max}\sin\theta = 0 \quad\Rightarrow\quad h_{\max}\sin\theta + mg\cos\theta = n$$

Then substituting,

$$h_{\max}\cos\theta - \mu_s(h_{\max}\sin\theta + mg\cos\theta) = mg\sin\theta \quad\Rightarrow$$

$$h_{\max}\cos\theta - h_{\max}\mu_s\sin\theta = mg\sin\theta + mg\mu_s\cos\theta \quad\Rightarrow$$

$$h_{\max} = mg\left(\frac{\sin\theta + \mu_s\cos\theta}{\cos\theta - \mu_s\sin\theta}\right) = mg\left(\frac{\tan\theta + \mu_s}{1 - \mu_s\tan\theta}\right)$$

$$= mg\left(\frac{\tan\theta + \tan\theta_s}{1 - \tan\theta_s\tan\theta}\right) = mg\tan(\theta + \theta_s)$$

We would expect $h_{\max}$ to increase as θ increases, with similar behavior as we established for $h_{\min}$, but with $h_{\max}$ values always larger than $h_{\min}$. We can see that this is the case if we graph $h_{\max}$ as a function of θ, as the curve is the graph of $h_{\min}$ translated $2\theta_s$ to the left, so the equation does seem reasonable. Notice that the equation predicts $h_{\max} \to \infty$ as $\theta \to (90^\circ - \theta_s)$. In fact, as $h_{\max}$ increases, the normal force increases as well. When $(90^\circ - \theta_s) \le \theta \le 90^\circ$, the horizontal force is completely counteracted by the sum of the normal and frictional forces, so no part of the horizontal force contributes to moving the block up the plane no matter how large its magnitude.

8. (a) The largest possible solid is achieved by starting with a circular cylinder of diameter 1 with axis the z-axis and with a height of 1. This is the largest solid that creates a square shadow with side length 1 in the y-direction and a circular disk shadow in the z-direction. For convenience, we place the base of the cylinder on the xy-plane so that it intersects the x- and y-axes at $\pm\frac{1}{2}$. We then remove as little as possible from the solid that leaves an isosceles triangle shadow in the x-direction. This is achieved by cutting the solid with planes parallel to the x-axis that intersect the z-axis at 1 and the y-axis at $\pm\frac{1}{2}$ (see the figure).

To compute the volume of this solid, we take vertical slices parallel to the xz-plane. The equation of the base of the solid is $x^2 + y^2 = \frac{1}{4}$, so a cross-section y units to the right of the origin is a rectangle with base $2\sqrt{\frac{1}{4} - y^2}$. For $0 \le y \le \frac{1}{2}$, the solid is cut off on top by the plane $z = 1 - 2y$, so the height of the rectangular cross-section is $1 - 2y$ and the cross-sectional area is $A(y) = 2\sqrt{\frac{1}{4} - y^2}\,(1 - 2y)$. The volume of the right half of the solid is

$$\int_0^{1/2} 2\sqrt{\tfrac{1}{4} - y^2}\,(1 - 2y)\,dy = 2\int_0^{1/2}\sqrt{\tfrac{1}{4} - y^2}\,dy - 4\int_0^{1/2} y\sqrt{\tfrac{1}{4} - y^2}\,dy$$

$$= 2\left[\tfrac{1}{4}\text{ area of a circle of radius }\tfrac{1}{2}\right] - 4\left[-\tfrac{1}{3}\left(\tfrac{1}{4} - y^2\right)^{3/2}\right]_0^{1/2}$$

$$= 2\left[\tfrac{1}{4}\cdot\pi\left(\tfrac{1}{2}\right)^2\right] + \tfrac{4}{3}\left[0 - \left(\tfrac{1}{4}\right)^{3/2}\right] = \tfrac{\pi}{8} - \tfrac{1}{6}$$

Thus the volume of the solid is $2\left(\frac{\pi}{8} - \frac{1}{6}\right) = \frac{\pi}{4} - \frac{1}{3} \approx 0.45$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) There is not a smallest volume. We can remove portions of the solid from part (a) to create smaller and smaller volumes as long as we leave the "skeleton" shown in the figure intact (which has no volume at all and is not a solid). Thus we can create solids with arbitrarily small volume.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13 ☐ VECTOR FUNCTIONS

13.1 Vector Functions and Space Curves

1. The component functions $\ln(t+1)$, $\dfrac{t}{\sqrt{9-t^2}}$, and 2^t are all defined when $t+1>0 \quad \Rightarrow \quad t>-1$ and $9-t^2>0 \quad \Rightarrow$ $-3<t<3$, so the domain of $\mathbf{r}$ is $(-1,3)$.

2. The component functions $\cos t$, $\ln t$, and $\dfrac{1}{t-2}$ are all defined when $t>0$ and $t\neq 2$, so the domain of $\mathbf{r}$ is $(0,2)\cup(2,\infty)$.

3. $\lim\limits_{t\to 0} e^{-3t} = e^0 = 1$, $\lim\limits_{t\to 0} \dfrac{t^2}{\sin^2 t} = \lim\limits_{t\to 0} \dfrac{1}{\dfrac{\sin^2 t}{t^2}} = \dfrac{1}{\lim\limits_{t\to 0} \dfrac{\sin^2 t}{t^2}} = \dfrac{1}{\left(\lim\limits_{t\to 0} \dfrac{\sin t}{t}\right)^2} = \dfrac{1}{1^2} = 1$,

and $\lim\limits_{t\to 0} \cos 2t = \cos 0 = 1$. Thus

$$\lim_{t\to 0}\left(e^{-3t}\,\mathbf{i} + \frac{t^2}{\sin^2 t}\,\mathbf{j} + \cos 2t\,\mathbf{k}\right) = \left[\lim_{t\to 0} e^{-3t}\right]\mathbf{i} + \left[\lim_{t\to 0}\frac{t^2}{\sin^2 t}\right]\mathbf{j} + \left[\lim_{t\to 0}\cos 2t\right]\mathbf{k} = \mathbf{i}+\mathbf{j}+\mathbf{k}.$$

4. $\lim\limits_{t\to 1} \dfrac{t^2-t}{t-1} = \lim\limits_{t\to 1} \dfrac{t\,(t-1)}{t-1} = \lim\limits_{t\to 1} t = 1$, $\lim\limits_{t\to 1} \sqrt{t+8} = 3$, $\lim\limits_{t\to 1} \dfrac{\sin \pi t}{\ln t} = \lim\limits_{t\to 1} \dfrac{\pi\cos \pi t}{1/t} = -\pi$ [by l'Hospital's Rule].

Thus the given limit equals $\mathbf{i} + 3\,\mathbf{j} - \pi\,\mathbf{k}$.

5. $\lim\limits_{t\to\infty} \dfrac{1+t^2}{1-t^2} = \lim\limits_{t\to\infty} \dfrac{(1/t^2)+1}{(1/t^2)-1} = \dfrac{0+1}{0-1} = -1$, $\lim\limits_{t\to\infty} \tan^{-1} t = \frac{\pi}{2}$, $\lim\limits_{t\to\infty} \dfrac{1-e^{-2t}}{t} = \lim\limits_{t\to\infty} \dfrac{1}{t} - \dfrac{1}{te^{2t}} = 0-0=0$. Thus

$$\lim_{t\to\infty}\left\langle \frac{1+t^2}{1-t^2}, \tan^{-1} t, \frac{1-e^{-2t}}{t}\right\rangle = \left\langle -1, \tfrac{\pi}{2}, 0\right\rangle.$$

6. $\lim\limits_{t\to\infty} te^{-t} = \lim\limits_{t\to\infty} \dfrac{t}{e^t} = \lim\limits_{t\to\infty} \dfrac{1}{e^t} = 0$ [by l'Hospital's Rule], $\lim\limits_{t\to\infty} \dfrac{t^3+t}{2t^3-1} = \lim\limits_{t\to\infty} \dfrac{1+(1/t^2)}{2-(1/t^3)} = \dfrac{1+0}{2-0} = \dfrac{1}{2}$,

and $\lim\limits_{t\to\infty} t\sin\dfrac{1}{t} = \lim\limits_{t\to\infty} \dfrac{\sin(1/t)}{1/t} = \lim\limits_{t\to\infty} \dfrac{\cos(1/t)(-1/t^2)}{-1/t^2} = \lim\limits_{t\to\infty} \cos\dfrac{1}{t} = \cos 0 = 1$ [again by l'Hospital's Rule].

Thus $\lim\limits_{t\to\infty} \left\langle te^{-t}, \dfrac{t^3+t}{2t^3-1}, t\sin\dfrac{1}{t}\right\rangle = \left\langle 0, \tfrac{1}{2}, 1\right\rangle$.

7. The corresponding parametric equations for this curve are $x = -\cos t$, $y = t$. We can make a table of values or we can eliminate the parameter: $t = y \quad \Rightarrow \quad x = -\cos y$, with $y \in \mathbb{R}$. By comparing different values of t, we find the direction in which t increases as indicated in the graph.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. The corresponding parametric equations for this curve are $x = t^2 - 1,\ y = t$. We can make a table of values, or we can eliminate the parameter: $t = y \quad \Rightarrow \quad x = y^2 - 1$, with $y \in \mathbb{R}$. Thus the curve is a parabola with vertex $(-1, 0)$ that opens to the right. By comparing different values of t, we find the direction in which t increases as indicated in the graph.

9. The corresponding parametric equations for this curve are $x = 3\sin t$, $y = 2\cos t$. We can make a table of values, or we can eliminate the parameter: $x = 3\sin t$, $y = 2\cos t \quad \Rightarrow \quad \dfrac{x}{3} = \sin t$, $\dfrac{y}{2} = \cos t \quad \Rightarrow$ $\dfrac{x^2}{9} + \dfrac{y^2}{4} = \sin^2 t + \cos^2 t = 1$, which we recognize as the equation of an ellipse with $x \in [-3, 3]$ and $y \in [-2, 2]$. By comparing different values of t, we find the direction in which t increases as indicated in the graph.

10. The corresponding parametric equations for this curve are $x = e^t$, $y = e^{-t}$. We can make a table of values, or we can eliminate the parameter: $y = e^{-t} = 1/e^t = 1/x$ with $x, y > 0$. By comparing different values of t, we find the direction in which t increases as indicated in the graph.

11. The corresponding parametric equations are $x = t,\ y = 2 - t,\ z = 2t$, which are parametric equations of a line through the point $(0, 2, 0)$ and with direction vector $\langle 1, -1, 2\rangle$.

12. The corresponding parametric equations are $x = \sin \pi t,\ y = t,\ z = \cos \pi t$. Note that $x^2 + z^2 = \sin^2 \pi t + \cos^2 \pi t = 1$, so the curve lies on the circular cylinder $x^2 + z^2 = 1$. A point (x, y, z) on the curve lies directly to the left or right of the point $(x, 0, z)$ which moves clockwise (when viewed from the left) along the circle $x^2 + z^2 = 1$ in the xz-plane as t increases. Since $y = t$, the curve is a helix that spirals toward the right around the cylinder.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. The corresponding parametric equations are $x = 3,\ y = t,\ z = 2 - t^2$. Eliminating the parameter in y and z gives $z = 2 - y^2$. Because $x = 3$, the curve is a parabola in the vertical plane $x = 3$ with vertex $(3, 0, 2)$.

14. The corresponding parametric equations are $x = 2\cos t,\ y = 2\sin t$, $z = 1$. Eliminating the parameter in x and y gives $x^2 + y^2 = 4\cos^2 t + 4\sin^2 t = 4(\cos^2 t + \sin^2 t) = 4$. Since $z = 1$, the curve is a circle of radius 2 centered at $(0, 0, 1)$ in the horizontal plane $z = 1$.

15. The parametric equations are $x = t^2$, $y = t^4$, $z = t^6$. These are positive for $t \neq 0$ and 0 when $t = 0$. So the curve lies entirely in the first octant. The projection of the graph onto the xy-plane is $y = x^2$, $y > 0$, a half parabola. The projection onto the xz-plane is $z = x^3$, $z > 0$, a half cubic, and the projection onto the yz-plane is $y^3 = z^2$.

16. If $x = \cos t,\ y = -\cos t,\ z = \sin t$, then $x^2 + z^2 = 1$ and $y^2 + z^2 = 1$, so the curve is contained in the intersection of circular cylinders along the x- and y-axes. Furthermore, $y = -x$, so the curve is an ellipse in the plane $y = -x$, centered at the origin.

17. The projection of the curve defined by the vector function $\mathbf{r}(t) = \langle t^2, t^3, t^{-3} \rangle$ onto the yz-plane is given by $\mathbf{r}(t) = \langle 0, t^3, t^{-3} \rangle$ [we use 0 for the x-component], whose graph is the curve $z = 1/y$, $x = 0$, since $z = t^{-3} = 1/t^3$.

18. The projection of the curve defined by the vector function $\mathbf{r}(t) = \langle t + 1, 3t + 1, \cos(t/2) \rangle$ onto the xy-plane is given by $\mathbf{r}(t) = \langle t + 1, 3t + 1, 0 \rangle$ [we use 0 for the z-component], whose graph is the curve $y = 3x - 2$, $z = 0$, since $y = 3t + 1 = 3(x - 1) + 1 = 3x - 2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. The projection of the curve onto the xy-plane is given by $\mathbf{r}(t) = \langle t, \sin t, 0\rangle$ [we use 0 for the z-component] whose graph is the curve $y = \sin x$, $z = 0$. Similarly, the projection onto the xz-plane is $\mathbf{r}(t) = \langle t, 0, 2\cos t\rangle$, whose graph is the cosine wave $z = 2\cos x$, $y = 0$, and the projection onto the yz-plane is $\mathbf{r}(t) = \langle 0, \sin t, 2\cos t\rangle$ whose graph is the ellipse $y^2 + \frac{1}{4}z^2 = 1$, $x = 0$.

xy-plane

xz-plane

yz-plane

From the projection onto the yz-plane we see that the curve lies on an elliptical cylinder with axis the x-axis. The other two projections show that the curve oscillates both vertically and horizontally as we move in the x-direction, suggesting that the curve is an elliptical helix that spirals along the cylinder.

20. The projection of the curve onto the xy-plane is given by $\mathbf{r}(t) = \langle t, t, 0\rangle$ whose graph is the line $y = x$, $z = 0$.
The projection onto the xz-plane is $\mathbf{r}(t) = \langle t, 0, t^2\rangle$ whose graph is the parabola $z = x^2$, $y = 0$.
The projection onto the yz-plane is $\mathbf{r}(t) = \langle 0, t, t^2\rangle$ whose graph is the parabola $z = y^2$, $x = 0$.

xy-plane

xz-plane

yz-plane

From the projection onto the xy-plane we see that the curve lies on the vertical plane $y = x$. The other two projections show that the curve is a parabola contained in this plane.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. We take $\mathbf{r}_0 = \langle -2, 1, 0\rangle$ and $\mathbf{r}_1 = \langle 5, 2, -3\rangle$. Then, by Equation 12.5.4 we have a vector equation for the line segment:

$$\mathbf{r}(t) = (1-t)\langle -2, 1, 0\rangle + t\langle 5, 2, -3\rangle \quad \Rightarrow \quad \mathbf{r}(t) = \langle -2+7t, 1+t, -3t\rangle, \quad 0 \le t \le 1$$

with corresponding parametric equations $x = -2+7t$, $y = 1+t$, $z = -3t$, $0 \le t \le 1$.

22. We take $\mathbf{r}_0 = \langle 0, 0, 0\rangle$ and $\mathbf{r}_1 = \langle -7, 4, 6\rangle$. Then, by Equation 12.5.4 we have a vector equation for the line segment:

$$\mathbf{r}(t) = (1-t)\langle 0, 0, 0\rangle + t\langle -7, 4, 6\rangle \quad \Rightarrow \quad \mathbf{r}(t) = \langle -7t, 4t, 6t\rangle, \quad 0 \le t \le 1$$

with corresponding parametric equations $x = -7t$, $y = 4t$, $z = 6t$, $0 \le t \le 1$.

23. We take $\mathbf{r}_0 = \langle 3.5, -1.4, 2.1\rangle$ and $\mathbf{r}_1 = \langle 1.8, 0.3, 2.1\rangle$. Then, by Equation 12.5.4 we have a vector equation for the line segment:

$$\mathbf{r}(t) = (1-t)\langle 3.5, -1.4, 2.1\rangle + t\langle 1.8, 0.3, 2.1\rangle \quad \Rightarrow \quad \mathbf{r}(t) = \langle 3.5-1.7t, -1.4+1.7t, 2.1\rangle, \quad 0 \le t \le 1$$

with corresponding parametric equations $x = 3.5-1.7t$, $y = -1.4+1.7t$, $z = 2.1$, $0 \le t \le 1$.

24. We take $\mathbf{r}_0 = \langle a, b, c\rangle$ and $\mathbf{r}_1 = \langle u, v, w\rangle$. Then, by Equation 12.5.4 we have a vector equation for the line segment:

$$\mathbf{r}(t) = (1-t)\,\mathbf{r}_0 + t\,\mathbf{r}_1 = (1-t)\langle a, b, c\rangle + t\langle u, v, w\rangle$$

$$\Rightarrow \qquad \mathbf{r}(t) = \langle a+(u-a)t, b+(v-b)t, c+(w-c)t\rangle, \quad 0 \le t \le 1$$

with corresponding parametric equations $x = a+(u-a)t,\ y = b+(v-b)t,\ z = c+(w-c)t,\ 0 \le t \le 1$.

25. $x = t\cos t,\ y = t,\ z = t\sin t,\ t \ge 0$. At any point (x, y, z) on the curve, $x^2 + z^2 = t^2\cos^2 t + t^2\sin^2 t = t^2 = y^2$ so the curve lies on the circular cone $x^2 + z^2 = y^2$ with axis the y-axis. Also notice that $y \ge 0$; the graph is II.

26. $x = \cos t,\ y = \sin t,\ z = 1/(1+t^2)$. At any point on the curve we have $x^2 + y^2 = \cos^2 t + \sin^2 t = 1$, so the curve lies on the circular cylinder $x^2 + y^2 = 1$ with axis the z-axis. Notice that $0 < z \le 1$ and $z = 1$ only for $t = 0$. A point (x, y, z) on the curve lies directly above the point $(x, y, 0)$, which moves counterclockwise around the unit circle in the xy-plane as t increases, and $z \to 0$ as $t \to \pm\infty$. The graph must be VI.

27. $x = t,\ y = 1/(1+t^2),\ z = t^2$. At any point on the curve we have $z = x^2$, so the curve lies on a parabolic cylinder parallel to the y-axis. Notice that $0 < y \le 1$ and $z \ge 0$. Also the curve passes through $(0, 1, 0)$ when $t = 0$ and $y \to 0$, $z \to \infty$ as $t \to \pm\infty$, so the graph must be V.

28. $x = \cos t,\ y = \sin t,\ z = \cos 2t$. $x^2 + y^2 = \cos^2 t + \sin^2 t = 1$, so the curve lies on a circular cylinder with axis the z-axis. A point (x, y, z) on the curve lies directly above or below $(x, y, 0)$, which moves around the unit circle in the xy-plane with period 2π. At the same time, the z-value of the point (x, y, z) oscillates with a period of π. So the curve repeats itself and the graph is I.

29. $x = \cos 8t,\ y = \sin 8t,\ z = e^{0.8t},\ t \ge 0$. $x^2 + y^2 = \cos^2 8t + \sin^2 8t = 1$, so the curve lies on a circular cylinder with axis the z-axis. A point (x, y, z) on the curve lies directly above the point $(x, y, 0)$, which moves counterclockwise around the unit circle in the xy-plane as t increases. The curve starts at $(1, 0, 1)$, when $t = 0$, and $z \to \infty$ (at an increasing rate) as $t \to \infty$, so the graph is IV.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $x = \cos^2 t,\ y = \sin^2 t,\ z = t.\quad x + y = \cos^2 t + \sin^2 t = 1$, so the curve lies in the vertical plane $x + y = 1$. x and y are periodic, both with period π, and z increases as t increases, so the graph is III.

31. As $y = 4$ in the vector equation $\mathbf{r}(t) = \langle t, 4, t^2 \rangle$, the curve $z = x^2$ lies in the plane $y = 4$.

32. $\mathbf{r}(t) = \langle t, t^2, t \rangle$. Consider the projection of the curve in the xz-plane, $\mathbf{r}(t) = \langle t, 0, t \rangle$. This is the line $z = x$, $y = 0$. Thus, the curve is contained in the plane $z = x$.

33. $\mathbf{r}(t) = \langle \sin t, \cos t, -\cos t \rangle$. Consider the projection of the curve in the yz-plane, $\mathbf{r}(t) = \langle 0, \cos t, -\cos t \rangle$. This is the line $z = -y$, $x = 0$. Thus, the curve is contained in the plane $z = -y$.

34. $\mathbf{r}(t) = \langle 2t, \sin t, t + 1 \rangle$. Consider the projection in the xz-plane, $\mathbf{r}(t) = \langle 2t, 0, t + 1 \rangle$. This is the line with parametric equations $x = 2t$, $z = t + 1$, $y = 0 \quad \Rightarrow \quad x = 2t = 2(z - 1) = 2z - 2$, $y = 0$. Thus, the curve is contained in the plane $x = 2z - 2$.

35. If $x = t\cos t,\ y = t\sin t,\ z = t$, then $x^2 + y^2 = t^2\cos^2 t + t^2\sin^2 t = t^2 = z^2$, so the curve lies on the cone $z^2 = x^2 + y^2$. Since $z = t$, the curve is a spiral on this cone.

36. If $x = \sin t$, $y = \cos t$, $z = \sin^2 t$, then $x^2 = \sin^2 t = z$ and $x^2 + y^2 = \sin^2 t + \cos^2 t = 1$, so the curve is contained in the intersection of the parabolic cylinder $z = x^2$ with the circular cylinder $x^2 + y^2 = 1$. We get the complete intersection for $0 \le t \le 2\pi$.

37. Here $x = 2t,\ y = e^t,\ z = e^{2t}$. Then $t = x/2 \quad \Rightarrow \quad y = e^t = e^{x/2}$, so the curve lies on the cylinder $y = e^{x/2}$. Also $z = e^{2t} = e^x$, so the curve lies on the cylinder $z = e^x$. Since $z = e^{2t} = \left(e^t\right)^2 = y^2$, the curve also lies on the parabolic cylinder $z = y^2$.

38. Here $x = t^2,\ y = \ln t,\ z = 1/t$. The domain of $\mathbf{r}$ is $(0, \infty)$, so $x = t^2 \quad \Rightarrow \quad t = \sqrt{x} \quad \Rightarrow \quad y = \ln\sqrt{x}$. Thus one surface containing the curve is the cylinder $y = \ln\sqrt{x}$ or $y = \ln x^{1/2} = \frac{1}{2}\ln x$. Also $z = 1/t = 1/\sqrt{x}$, so the curve also lies on the cylinder $z = 1/\sqrt{x}$ or $x = 1/z^2$, $z > 0$. Finally $z = 1/t \quad \Rightarrow \quad t = 1/z \quad \Rightarrow \quad y = \ln(1/z)$, so the curve also lies on the cylinder $y = \ln(1/z)$ or $y = \ln z^{-1} = -\ln z$. Note that the surface $y = \ln(xz)$ also contains the curve, since $\ln(xz) = \ln(t^2 \cdot 1/t) = \ln t = y$.

39. Parametric equations for the curve are $x = t,\ y = 0,\ z = 2t - t^2$. Substituting into the equation of the paraboloid gives $2t - t^2 = t^2 \quad \Rightarrow \quad 2t = 2t^2 \quad \Rightarrow \quad t = 0, 1$. Since $\mathbf{r}(0) = \mathbf{0}$ and $\mathbf{r}(1) = \mathbf{i} + \mathbf{k}$, the points of intersection are $(0, 0, 0)$ and $(1, 0, 1)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. Parametric equations for the helix are $x = \sin t$, $y = \cos t$, $z = t$. Substituting into the equation of the sphere gives $\sin^2 t + \cos^2 t + t^2 = 5 \quad \Rightarrow \quad 1 + t^2 = 5 \quad \Rightarrow \quad t = \pm 2$. Since $\mathbf{r}(2) = \langle \sin 2, \cos 2, 2 \rangle$ and $\mathbf{r}(-2) = \langle \sin(-2), \cos(-2), -2 \rangle$, the points of intersection are $(\sin 2, \cos 2, 2) \approx (0.909, -0.416, 2)$ and $(\sin(-2), \cos(-2), -2) \approx (-0.909, -0.416, -2)$.

41. $\mathbf{r}(t) = \langle \cos t \sin 2t, \sin t \sin 2t, \cos 2t \rangle$. We include both a regular plot and a plot showing a tube of radius 0.08 around the curve.

42. $\mathbf{r}(t) = \left\langle te^t, e^{-t}, t \right\rangle$

43. $\mathbf{r}(t) = \left\langle \sin 3t \cos t, \frac{1}{4}t, \sin 3t \sin t \right\rangle$

44. $\mathbf{r}(t) = \langle \cos(8\cos t)\sin t, \sin(8\cos t)\sin t, \cos t \rangle$

45. $\mathbf{r}(t) = \langle \cos 2t, \cos 3t, \cos 4t \rangle$

46. $x = \sin t$, $y = \sin 2t$, $z = \cos 4t$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

We graph the projections onto the coordinate planes.

xy-plane $\qquad$ xz-plane $\qquad$ yz-plane

From the projection onto the xy-plane we see that from above the curve appears to be shaped like a "figure eight." The curve can be visualized as this shape wrapped around an almost parabolic cylindrical surface, the profile of which is visible in the projection onto the yz-plane.

47.

$x = (1 + \cos 16t)\cos t$, $y = (1 + \cos 16t)\sin t$, $z = 1 + \cos 16t$. At any point on the graph,

$$x^2 + y^2 = (1 + \cos 16t)^2 \cos^2 t + (1 + \cos 16t)^2 \sin^2 t$$
$$= (1 + \cos 16t)^2 = z^2, \text{ so the graph lies on the cone } x^2 + y^2 = z^2.$$

From the graph at left, we see that this curve looks like the projection of a leaved two-dimensional curve onto a cone.

48.

$x = \sqrt{1 - 0.25\cos^2 10t}\cos t,\ y = \sqrt{1 - 0.25\cos^2 10t}\sin t,$
$z = 0.5\cos 10t$. At any point on the graph,

$$x^2 + y^2 + z^2 = (1 - 0.25\cos^2 10t)\cos^2 t$$
$$+(1 - 0.25\cos^2 10t)\sin^2 t + 0.25\cos^2 t$$
$$= 1 - 0.25\cos^2 10t + 0.25\cos^2 10t = 1,$$

so the graph lies on the sphere $x^2 + y^2 + z^2 = 1$, and since $z = 0.5\cos 10t$ the graph resembles a trigonometric curve with ten peaks projected onto the sphere. We get the complete graph for $0 \le t \le 2\pi$.

49. If $t = -1$, then $x = 1,\ y = 4,\ z = 0$, so the curve passes through the point $(1, 4, 0)$. If $t = 3$, then $x = 9,\ y = -8,\ z = 28$, so the curve passes through the point $(9, -8, 28)$. For the point $(4, 7, -6)$ to be on the curve, we require $y = 1 - 3t = 7 \quad \Rightarrow$ $t = -2$. But then $z = 1 + (-2)^3 = -7 \neq -6$, so $(4, 7, -6)$ is not on the curve.

50. The projection of the curve C of intersection onto the xy-plane is the circle $x^2 + y^2 = 4$, $z = 0$. Then we can write $x = 2\cos t,\ y = 2\sin t,\ 0 \le t \le 2\pi$. Since C also lies on the surface $z = xy$, we have $z = xy = (2\cos t)(2\sin t) = 4\cos t\sin t$, or $2\sin(2t)$. Then parametric equations for C are $x = 2\cos t,\ y = 2\sin t,$ $z = 2\sin(2t),\ 0 \le t \le 2\pi$, and the corresponding vector function is $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j} + 2\sin(2t)\,\mathbf{k}$, $0 \le t \le 2\pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. Both equations are solved for z, so we can substitute to eliminate z: $\sqrt{x^2+y^2} = 1+y \quad \Rightarrow \quad x^2+y^2 = 1+2y+y^2 \quad \Rightarrow$ $x^2 = 1+2y \quad \Rightarrow \quad y = \frac{1}{2}(x^2-1)$. We can form parametric equations for the curve C of intersection by choosing a parameter $x = t$, then $y = \frac{1}{2}(t^2-1)$ and $z = 1+y = 1+\frac{1}{2}(t^2-1) = \frac{1}{2}(t^2+1)$. Thus a vector function representing C is $\mathbf{r}(t) = t\,\mathbf{i} + \frac{1}{2}(t^2-1)\,\mathbf{j} + \frac{1}{2}(t^2+1)\,\mathbf{k}$.

52. The projection of the curve C of intersection onto the xy-plane is the parabola $y = x^2$, $z = 0$. Then we can choose the parameter $x = t \quad \Rightarrow \quad y = t^2$. Since C also lies on the surface $z = 4x^2+y^2$, we have $z = 4x^2+y^2 = 4t^2+(t^2)^2$. Then parametric equations for C are $x = t,\ y = t^2,\ z = 4t^2+t^4$, and the corresponding vector function is $\mathbf{r}(t) = t\,\mathbf{i} + t^2\,\mathbf{j} + (4t^2+t^4)\,\mathbf{k}$.

53. The projection of the curve C of intersection onto the xy-plane is the circle $x^2+y^2 = 1$, $z = 0$, so we can write $x = \cos t$, $y = \sin t$, $0 \le t \le 2\pi$. Since C also lies on the surface $z = x^2-y^2$, we have $z = x^2-y^2 = \cos^2 t - \sin^2 t$ or $\cos 2t$. Thus parametric equations for C are $x = \cos t$, $y = \sin t$, $z = \cos 2t$, $0 \le t \le 2\pi$, and the corresponding vector function is $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j} + \cos 2t\,\mathbf{k}$, $0 \le t \le 2\pi$.

54. The projection of the curve C of intersection onto the xz-plane is the circle $x^2+z^2 = 1$, $y = 0$, so we can write $x = \cos t$, $z = \sin t$, $0 \le t \le 2\pi$. C also lies on the surface $x^2+y^2+4z^2 = 4$, and since $y \ge 0$ we can write

$$y = \sqrt{4-x^2-4z^2} = \sqrt{4-\cos^2 t - 4\sin^2 t} = \sqrt{4-\cos^2 t - 4(1-\cos^2 t)} = \sqrt{3\cos^2 t} = \sqrt{3}\,|\cos t\,|$$

Thus parametric equations for C are $x = \cos t$, $y = \sqrt{3}\,|\cos t\,|$, $z = \sin t$, $0 \le t \le 2\pi$, and the corresponding vector function is $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sqrt{3}\,|\cos t\,|\,\mathbf{j} + \sin t\,\mathbf{k}$, $0 \le t \le 2\pi$.

55.

The projection of the curve C of intersection onto the xy-plane is the circle $x^2+y^2 = 4$, $z = 0$. Then we can write $x = 2\cos t,\ y = 2\sin t,\ 0 \le t \le 2\pi$. Since C also lies on the surface $z = x^2$, we have $z = x^2 = (2\cos t)^2 = 4\cos^2 t$. Then parametric equations for C are $x = 2\cos t,\ y = 2\sin t,$ $z = 4\cos^2 t,\ 0 \le t \le 2\pi$.

56.

$x = t \quad \Rightarrow \quad y = t^2 \quad \Rightarrow \quad 4z^2 = 16 - x^2 - 4y^2 = 16 - t^2 - 4t^4 \quad \Rightarrow \quad z = \sqrt{4 - \left(\frac{1}{2}t\right)^2 - t^4}$.

Note that z is positive because the intersection is with the top half of the ellipsoid. Hence the curve is given by $x = t,\ y = t^2,\ z = \sqrt{4 - \frac{1}{4}t^2 - t^4}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. For the particles to collide, we require $\mathbf{r}_1(t) = \mathbf{r}_2(t) \quad \Leftrightarrow \quad \langle t^2, 7t-12, t^2\rangle = \langle 4t-3, t^2, 5t-6\rangle$. Equating components gives $t^2 = 4t-3$, $7t-12 = t^2$, and $t^2 = 5t-6$. From the first equation, $t^2-4t+3=0 \quad \Leftrightarrow \quad (t-3)(t-1)=0$ so $t=1$ or $t=3$. $t=1$ does not satisfy the other two equations, but $t=3$ does. The particles collide when $t=3$, at the point $(9,9,9)$.

58. The particles collide provided $\mathbf{r}_1(t) = \mathbf{r}_2(t) \quad \Leftrightarrow \quad \langle t, t^2, t^3\rangle = \langle 1+2t, 1+6t, 1+14t\rangle$. Equating components gives $t = 1+2t$, $t^2 = 1+6t$, and $t^3 = 1+14t$. The first equation gives $t=-1$, but this does not satisfy the other equations, so the particles do not collide. For the paths to intersect, we need to find a value for t and a value for s where $\mathbf{r}_1(t) = \mathbf{r}_2(s) \quad \Leftrightarrow \quad \langle t, t^2, t^3\rangle = \langle 1+2s, 1+6s, 1+14s\rangle$. Equating components, $t = 1+2s$, $t^2 = 1+6s$, and $t^3 = 1+14s$. Substituting the first equation into the second gives $(1+2s)^2 = 1+6s \quad \Rightarrow \quad 4s^2-2s=0 \quad \Rightarrow \quad 2s(2s-1)=0 \quad \Rightarrow \quad s=0$ or $s=\frac{1}{2}$. From the first equation, $s=0 \quad \Rightarrow \quad t=1$ and $s=\frac{1}{2} \quad \Rightarrow \quad t=2$. Checking, we see that both pairs of values satisfy the third equation. Thus the paths intersect twice, at the point $(1,1,1)$ when $s=0$ and $t=1$, and at $(2,4,8)$ when $s=\frac{1}{2}$ and $t=2$.

59. (a) We plot the parametric equations for $0 \le t \le 2\pi$ in the first figure. We get a better idea of the shape of the curve if we plot it simultaneously with the hyperboloid of one sheet from part (b), as shown in the second figure.

(b) Here $x = \frac{27}{26}\sin 8t - \frac{8}{39}\sin 18t,\ y = -\frac{27}{26}\cos 8t + \frac{8}{39}\cos 18t,\ z = \frac{144}{65}\sin 5t$.

For any point on the curve,

$$\begin{aligned}
x^2+y^2 &= \left(\tfrac{27}{26}\sin 8t - \tfrac{8}{39}\sin 18t\right)^2 + \left(-\tfrac{27}{26}\cos 8t + \tfrac{8}{39}\cos 18t\right)^2 \\
&= \tfrac{27^2}{26^2}\sin^2 8t - 2\cdot\tfrac{27\cdot 8}{26\cdot 39}\sin 8t\sin 18t + \tfrac{64}{39^2}\sin^2 18t \\
&\qquad + \tfrac{27^2}{26^2}\cos^2 8t - 2\cdot\tfrac{27\cdot 8}{26\cdot 39}\cos 8t\cos 18t + \tfrac{64}{39^2}\cos^2 18t \\
&= \tfrac{27^2}{26^2}\left(\sin^2 8t + \cos^2 8t\right) + \tfrac{64}{39^2}\left(\sin^2 18t + \cos^2 18t\right) - \tfrac{72}{169}\left(\sin 8t\sin 18t + \cos 8t\cos 18t\right) \\
&= \tfrac{27^2}{26^2} + \tfrac{64}{39^2} - \tfrac{72}{169}\cos(18t-8t) = \tfrac{27^2}{26^2} + \tfrac{64}{39^2} - \tfrac{72}{169}\cos 10t
\end{aligned}$$

using the trigonometric identities $\sin^2\theta + \cos^2\theta = 1$ and $\cos(x-y) = \cos x\cos y + \sin x\sin y$. Also $z^2 = \frac{144^2}{65^2}\sin^2 5t$, and the identity $\sin^2 x = \dfrac{1-\cos 2x}{2}$ gives $z^2 = \frac{144^2}{65^2}\cdot\frac{1}{2}\left[1-\cos(2\cdot 5t)\right] = \frac{144^2}{2\cdot 65^2} - \frac{144^2}{2\cdot 65^2}\cos 10t$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Then

$$\begin{aligned}144(x^2+y^2)-25z^2 &= 144\left(\tfrac{27^2}{26^2}+\tfrac{64}{39^2}-\tfrac{72}{169}\cos 10t\right)-25\left(\tfrac{144^2}{2\cdot 65^2}-\tfrac{144^2}{2\cdot 65^2}\cos 10t\right)\\ &= 144\left(\tfrac{27^2}{26^2}+\tfrac{64}{39^2}-\tfrac{25\cdot 144}{2\cdot 65^2}-\tfrac{72}{169}\cos 10t+\tfrac{25\cdot 144}{2\cdot 65^2}\cos 10t\right)\\ &= 144\left(\tfrac{27^2}{26^2}+\tfrac{64}{39^2}-\tfrac{72}{169}-\tfrac{72}{169}\cos 10t+\tfrac{72}{169}\cos 10t\right)=144\left(\tfrac{25}{36}\right)=100\end{aligned}$$

Thus the curve lies on the surface $144(x^2+y^2)-25z^2=100$ or $144x^2+144y^2-25z^2=100$, a hyperboloid of one sheet with axis the z-axis.

60. The projection of the curve onto the xy-plane is given by the parametric equations $x=(2+\cos 1.5t)\cos t$, $y=(2+\cos 1.5t)\sin t$. If we convert to polar coordinates, we have

$$\begin{aligned}r^2=x^2+y^2&=[(2+\cos 1.5t)\cos t]^2+[(2+\cos 1.5t)\sin t]^2\\ &=(2+\cos 1.5t)^2(\cos^2 t+\sin^2 t)=(2+\cos 1.5t)^2 \quad\Rightarrow\quad r=2+\cos 1.5t\end{aligned}$$

Also, $\tan\theta=\dfrac{y}{x}=\dfrac{(2+\cos 1.5t)\sin t}{(2+\cos 1.5t)\cos t}=\tan t \quad\Rightarrow\quad \theta=t.$

Thus the polar equation of the curve is $r=2+\cos 1.5\theta$. At $\theta=0$, we have $r=3$, and r decreases to 1 as θ increases to $\frac{2\pi}{3}$. For $\frac{2\pi}{3}\le\theta\le\frac{4\pi}{3}$, r increases to 3; r decreases to 1 again at $\theta=2\pi$, increases to 3 at $\theta=\frac{8\pi}{3}$, decreases to 1 at $\theta=\frac{10\pi}{3}$, and completes the closed curve by increasing to 3 at $\theta=4\pi$. We sketch an approximate graph as shown in the figure.

We can determine how the curve passes over itself by investigating the maximum and minimum values of z for $0\le t\le 4\pi$. Since $z=\sin 1.5t$, z is maximized where $\sin 1.5t=1 \quad\Rightarrow\quad 1.5t=\frac{\pi}{2}, \frac{5\pi}{2}$, or $\frac{9\pi}{2} \quad\Rightarrow$ $t=\frac{\pi}{3}, \frac{5\pi}{3}$, or 3π. z is minimized where $\sin 1.5t=-1 \quad\Rightarrow$ $1.5t=\frac{3\pi}{2}, \frac{7\pi}{2}$, or $\frac{11\pi}{2} \quad\Rightarrow\quad t=\pi, \frac{7\pi}{3}$, or $\frac{11\pi}{3}$. Note that these are precisely the values for which $\cos 1.5t=0 \quad\Rightarrow\quad r=2$, and on the graph of the projection, these six points appear to be at the three self-intersections we see. Comparing the maximum and minimum values of z at these intersections, we can determine where the curve passes over itself, as indicated in the figure.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

We show a computer-drawn graph of the curve from above, as well as views from the front and from the right side.

Top view Front view Side view

The top view graph shows a more accurate representation of the projection of the trefoil knot onto the xy-plane (the axes are rotated 90°). Notice the indentations the graph exhibits at the points corresponding to $r = 1$. Finally, we graph several additional viewpoints of the trefoil knot, along with two plots showing a tube of radius 0.2 around the curve.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. Let $\mathbf{u}(t) = \langle u_1(t), u_2(t), u_3(t)\rangle$ and $\mathbf{v}(t) = \langle v_1(t), v_2(t), v_3(t)\rangle$. In each part of this problem the basic procedure is to use Equation 1 and then analyze the individual component functions using the limit properties we have already developed for real-valued functions.

(a) $\lim\limits_{t\to a} \mathbf{u}(t) + \lim\limits_{t\to a} \mathbf{v}(t) = \left\langle \lim\limits_{t\to a} u_1(t), \lim\limits_{t\to a} u_2(t), \lim\limits_{t\to a} u_3(t)\right\rangle + \left\langle \lim\limits_{t\to a} v_1(t), \lim\limits_{t\to a} v_2(t), \lim\limits_{t\to a} v_3(t)\right\rangle$ and the limits of these component functions must each exist since the vector functions both possess limits as $t \to a$. Then adding the two vectors and using the addition property of limits for real-valued functions, we have that

$$\begin{aligned}
\lim_{t\to a} \mathbf{u}(t) + \lim_{t\to a} \mathbf{v}(t) &= \left\langle \lim_{t\to a} u_1(t) + \lim_{t\to a} v_1(t), \lim_{t\to a} u_2(t) + \lim_{t\to a} v_2(t), \lim_{t\to a} u_3(t) + \lim_{t\to a} v_3(t)\right\rangle \\
&= \left\langle \lim_{t\to a} [u_1(t) + v_1(t)], \lim_{t\to a} [u_2(t) + v_2(t)], \lim_{t\to a} [u_3(t) + v_3(t)]\right\rangle \\
&= \lim_{t\to a} \langle u_1(t) + v_1(t), u_2(t) + v_2(t), u_3(t) + v_3(t)\rangle \qquad \text{[using (1) backward]} \\
&= \lim_{t\to a} [\mathbf{u}(t) + \mathbf{v}(t)]
\end{aligned}$$

(b)
$$\begin{aligned}
\lim_{t\to a} c\mathbf{u}(t) &= \lim_{t\to a} \langle cu_1(t), cu_2(t), cu_3(t)\rangle = \left\langle \lim_{t\to a} cu_1(t), \lim_{t\to a} cu_2(t), \lim_{t\to a} cu_3(t)\right\rangle \\
&= \left\langle c\lim_{t\to a} u_1(t), c\lim_{t\to a} u_2(t), c\lim_{t\to a} u_3(t)\right\rangle = c\left\langle \lim_{t\to a} u_1(t), \lim_{t\to a} u_2(t), \lim_{t\to a} u_3(t)\right\rangle \\
&= c\lim_{t\to a} \langle u_1(t), u_2(t), u_3(t)\rangle = c\lim_{t\to a} \mathbf{u}(t)
\end{aligned}$$

(c)
$$\begin{aligned}
\lim_{t\to a} \mathbf{u}(t) \cdot \lim_{t\to a} \mathbf{v}(t) &= \left\langle \lim_{t\to a} u_1(t), \lim_{t\to a} u_2(t), \lim_{t\to a} u_3(t)\right\rangle \cdot \left\langle \lim_{t\to a} v_1(t), \lim_{t\to a} v_2(t), \lim_{t\to a} v_3(t)\right\rangle \\
&= \left[\lim_{t\to a} u_1(t)\right]\left[\lim_{t\to a} v_1(t)\right] + \left[\lim_{t\to a} u_2(t)\right]\left[\lim_{t\to a} v_2(t)\right] + \left[\lim_{t\to a} u_3(t)\right]\left[\lim_{t\to a} v_3(t)\right] \\
&= \lim_{t\to a} u_1(t)v_1(t) + \lim_{t\to a} u_2(t)v_2(t) + \lim_{t\to a} u_3(t)v_3(t) \\
&= \lim_{t\to a} [u_1(t)v_1(t) + u_2(t)v_2(t) + u_3(t)v_3(t)] = \lim_{t\to a} [\mathbf{u}(t) \cdot \mathbf{v}(t)]
\end{aligned}$$

(d)
$$\begin{aligned}
\lim_{t\to a} \mathbf{u}(t) \times \lim_{t\to a} \mathbf{v}(t) &= \left\langle \lim_{t\to a} u_1(t), \lim_{t\to a} u_2(t), \lim_{t\to a} u_3(t)\right\rangle \times \left\langle \lim_{t\to a} v_1(t), \lim_{t\to a} v_2(t), \lim_{t\to a} v_3(t)\right\rangle \\
&= \left\langle \left[\lim_{t\to a} u_2(t)\right]\left[\lim_{t\to a} v_3(t)\right] - \left[\lim_{t\to a} u_3(t)\right]\left[\lim_{t\to a} v_2(t)\right], \right. \\
&\qquad \left[\lim_{t\to a} u_3(t)\right]\left[\lim_{t\to a} v_1(t)\right] - \left[\lim_{t\to a} u_1(t)\right]\left[\lim_{t\to a} v_3(t)\right], \\
&\qquad\qquad \left. \left[\lim_{t\to a} u_1(t)\right]\left[\lim_{t\to a} v_2(t)\right] - \left[\lim_{t\to a} u_2(t)\right]\left[\lim_{t\to a} v_1(t)\right]\right\rangle \\
&= \left\langle \lim_{t\to a} [u_2(t)v_3(t) - u_3(t)v_2(t)], \lim_{t\to a} [u_3(t)v_1(t) - u_1(t)v_3(t)], \right. \\
&\qquad \left. \lim_{t\to a} [u_1(t)v_2(t) - u_2(t)v_1(t)]\right\rangle \\
&= \lim_{t\to a} \langle u_2(t)v_3(t) - u_3(t)v_2(t), u_3(t)v_1(t) - u_1(t)v_3(t), u_1(t)v_2(t) - u_2(t)v_1(t)\rangle \\
&= \lim_{t\to a} [\mathbf{u}(t) \times \mathbf{v}(t)]
\end{aligned}$$

62. Let $\mathbf{r}(t) = \langle f(t), g(t), h(t)\rangle$ and $\mathbf{b} = \langle b_1, b_2, b_3\rangle$. If $\lim\limits_{t\to a} \mathbf{r}(t) = \mathbf{b}$, then $\lim\limits_{t\to a} \mathbf{r}(t)$ exists, so by (1),

$\mathbf{b} = \lim\limits_{t\to a} \mathbf{r}(t) = \left\langle \lim\limits_{t\to a} f(t), \lim\limits_{t\to a} g(t), \lim\limits_{t\to a} h(t)\right\rangle$. By the definition of equal vectors we have $\lim\limits_{t\to a} f(t) = b_1$, $\lim\limits_{t\to a} g(t) = b_2$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and $\lim_{t\to a} h(t) = b_3$. But these are limits of real-valued functions, so by the definition of limits, for every $\varepsilon > 0$ there exists $\delta_1 > 0$, $\delta_2 > 0$, $\delta_3 > 0$ so that if $0 < |t-a| < \delta_1$ then $|f(t) - b_1| < \varepsilon/3$, if $0 < |t-a| < \delta_2$ then $|g(t) - b_2| < \varepsilon/3$, and if $0 < |t-a| < \delta_3$ then $|h(t) - b_3| < \varepsilon/3$. Letting $\delta =$ minimum of $\{\delta_1, \delta_2, \delta_3\}$, then if $0 < |t-a| < \delta$ we have $|f(t) - b_1| + |g(t) - b_2| + |h(t) - b_3| < \varepsilon/3 + \varepsilon/3 + \varepsilon/3 = \varepsilon$. But

$$\begin{aligned}|\mathbf{r}(t) - \mathbf{b}| &= |\langle f(t) - b_1, g(t) - b_2, h(t) - b_3\rangle| = \sqrt{(f(t) - b_1)^2 + (g(t) - b_2)^2 + (h(t) - b_3)^2} \\ &\le \sqrt{[f(t) - b_1]^2} + \sqrt{[g(t) - b_2]^2} + \sqrt{[h(t) - b_3]^2} = |f(t) - b_1| + |g(t) - b_2| + |h(t) - b_3|\end{aligned}$$

Thus for every $\varepsilon > 0$ there exists $\delta > 0$ such that if $0 < |t-a| < \delta$ then $|\mathbf{r}(t) - \mathbf{b}| \le |f(t) - b_1| + |g(t) - b_2| + |h(t) - b_3| < \varepsilon$.

Conversely, suppose for every $\varepsilon > 0$, there exists $\delta > 0$ such that if $0 < |t-a| < \delta$ then $|\mathbf{r}(t) - \mathbf{b}| < \varepsilon \quad \Leftrightarrow$ $|\langle f(t) - b_1, g(t) - b_2, h(t) - b_3\rangle| < \varepsilon \quad \Leftrightarrow \quad \sqrt{[f(t) - b_1]^2 + [g(t) - b_2]^2 + [h(t) - b_3]^2} < \varepsilon \quad \Leftrightarrow$ $[f(t) - b_1]^2 + [g(t) - b_2]^2 + [h(t) - b_3]^2 < \varepsilon^2$. But each term on the left side of the last inequality is positive, so if $0 < |t-a| < \delta$, then $[f(t) - b_1]^2 < \varepsilon^2$, $[g(t) - b_2]^2 < \varepsilon^2$ and $[h(t) - b_3]^2 < \varepsilon^2$ or, taking the square root of both sides in each of the above, $|f(t) - b_1| < \varepsilon$, $|g(t) - b_2| < \varepsilon$ and $|h(t) - b_3| < \varepsilon$. And by definition of limits of real-valued functions we have $\lim_{t\to a} f(t) = b_1$, $\lim_{t\to a} g(t) = b_2$, and $\lim_{t\to a} h(t) = b_3$. But by (1), $\lim_{t\to a} \mathbf{r}(t) = \left\langle \lim_{t\to a} f(t), \lim_{t\to a} g(t), \lim_{t\to a} h(t)\right\rangle$, so $\lim_{t\to a} \mathbf{r}(t) = \langle b_1, b_2, b_3\rangle = \mathbf{b}$.

13.2 Derivatives and Integrals of Vector Functions

1. (a)

(b) $\dfrac{\mathbf{r}(4.5) - \mathbf{r}(4)}{0.5} = 2[\mathbf{r}(4.5) - \mathbf{r}(4)]$, so we draw a vector in the same direction but with twice the length of the vector $\mathbf{r}(4.5) - \mathbf{r}(4)$.

$\dfrac{\mathbf{r}(4.2) - \mathbf{r}(4)}{0.2} = 5[\mathbf{r}(4.2) - \mathbf{r}(4)]$, so we draw a vector in the same direction but with 5 times the length of the vector $\mathbf{r}(4.2) - \mathbf{r}(4)$.

(c) By Definition 1, $\mathbf{r}'(4) = \lim_{h\to 0} \dfrac{\mathbf{r}(4+h) - \mathbf{r}(4)}{h}$. $\mathbf{T}(4) = \dfrac{\mathbf{r}'(4)}{|\mathbf{r}'(4)|}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $\mathbf{T}(4)$ is a unit vector in the same direction as $\mathbf{r}'(4)$, that is, parallel to the tangent line to the curve at $\mathbf{r}(4)$ with length 1.

2. (a) The curve can be represented by the parametric equations $x = t^2$, $y = t$, $0 \le t \le 2$. Eliminating the parameter, we have $x = y^2$, $0 \le y \le 2$, a portion of which we graph here, along with the vectors $\mathbf{r}(1)$, $\mathbf{r}(1.1)$, and $\mathbf{r}(1.1) - \mathbf{r}(1)$.

(b) Since $\mathbf{r}(t) = \langle t^2, t \rangle$, we differentiate components, giving $\mathbf{r}'(t) = \langle 2t, 1 \rangle$, so $\mathbf{r}'(1) = \langle 2, 1 \rangle$.

$$\frac{\mathbf{r}(1.1) - \mathbf{r}(1)}{0.1} = \frac{\langle 1.21, 1.1 \rangle - \langle 1, 1 \rangle}{0.1} = 10\,\langle 0.21, 0.1 \rangle = \langle 2.1, 1 \rangle.$$

As we can see from the graph, these vectors are very close in length and direction. $\mathbf{r}'(1)$ is defined to be

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\lim\limits_{h\to 0} \dfrac{\mathbf{r}(1+h)-\mathbf{r}(1)}{h}$, and we recognize $\dfrac{\mathbf{r}(1.1)-\mathbf{r}(1)}{0.1}$ as the expression after the limit sign with $h = 0.1$. Since h is close to 0, we would expect $\dfrac{\mathbf{r}(1.1)-\mathbf{r}(1)}{0.1}$ to be a vector close to $\mathbf{r}'(1)$.

3. $\mathbf{r}(t) = \langle t-2, t^2+1\rangle$,
$\mathbf{r}(-1) = \langle -3, 2\rangle$.
Since $(x+2)^2 = t^2 = y-1 \quad \Rightarrow$ $y = (x+2)^2+1$, the curve is a parabola.

(a), (c)

(b) $\mathbf{r}'(t) = \langle 1, 2t\rangle$,
$\mathbf{r}'(-1) = \langle 1, -2\rangle$

4. $\mathbf{r}(t) = \langle t^2, t^3\rangle$, $\mathbf{r}(1) = \langle 1, 1\rangle$.
Since $x = t^2 = (t^3)^{2/3} = y^{2/3}$, the curve is the graph of $x = y^{2/3}$.

(a), (c)

(b) $\mathbf{r}'(t) = \langle 2t, 3t^2\rangle$,
$\mathbf{r}'(1) = \langle 2, 3\rangle$

5. $\mathbf{r}(t) = e^{2t}\,\mathbf{i} + e^t\,\mathbf{j}$, $\mathbf{r}(0) = \mathbf{i}+\mathbf{j}$.
Since $x = e^{2t} = (e^t)^2 = y^2$, the curve is part of a parabola. Note that here $x > 0$, $y > 0$.

(a), (c)

(b) $\mathbf{r}'(t) = 2e^{2t}\,\mathbf{i} + e^t\,\mathbf{j}$,
$\mathbf{r}'(0) = 2\,\mathbf{i}+\mathbf{j}$

6. $\mathbf{r}(t) = e^t\,\mathbf{i} + 2t\,\mathbf{j}$, $\mathbf{r}(0) = \mathbf{i}$.
Since $x = e^t \quad \Leftrightarrow \quad t = \ln x$ and $y = 2t = 2\ln x$, the curve is the graph of $y = 2\ln x$.

(a), (c)

(b) $\mathbf{r}'(t) = e^t\,\mathbf{i} + 2\,\mathbf{j}$,
$\mathbf{r}'(0) = \mathbf{i} + 2\,\mathbf{j}$

7. $\mathbf{r}(t) = 4\sin t\,\mathbf{i} - 2\cos t\,\mathbf{j}$, $\mathbf{r}(3\pi/4) = 4(\sqrt{2}/2)\,\mathbf{i} - 2(-\sqrt{2}/2)\,\mathbf{j} = 2\sqrt{2}\,\mathbf{i} + \sqrt{2}\,\mathbf{j}$.
Here $(x/4)^2 + (y/2)^2 = \sin^2 t + \cos^2 t = 1$, so the curve is the ellipse $\dfrac{x^2}{16} + \dfrac{y^2}{4} = 1$.

(a), (c)

(b) $\mathbf{r}'(t) = 4\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j}$,
$\mathbf{r}'(3\pi/4) = -2\sqrt{2}\,\mathbf{i} + \sqrt{2}\,\mathbf{j}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $\mathbf{r}(t) = (\cos t + 1)\,\mathbf{i} + (\sin t - 1)\,\mathbf{j}$, $\mathbf{r}(-\pi/3) = \left(\frac{1}{2} + 1\right)\mathbf{i} + \left(-\frac{\sqrt{3}}{2} - 1\right)\mathbf{j} = \frac{3}{2}\,\mathbf{i} + \left(-\frac{\sqrt{3}}{2} - 1\right)\mathbf{j} \approx 1.5\,\mathbf{i} - 1.87\,\mathbf{j}$.

Here $(x-1)^2 + (y+1)^2 = \cos^2 t + \sin^2 t = 1$, so the curve is a circle of radius 1 with center $(1, -1)$.

(a), (c)

(b) $\mathbf{r}'(t) = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j}$,

$\mathbf{r}'(-\pi/3) = \frac{\sqrt{3}}{2}\mathbf{i} + \frac{1}{2}\,\mathbf{j} \approx 0.87\,\mathbf{i} + 0.5\,\mathbf{j}$

9. $\mathbf{r}(t) = \left\langle \sqrt{t-2}, 3,\ 1/t^2 \right\rangle \quad \Rightarrow$

$$\mathbf{r}'(t) = \left\langle \frac{d}{dt}\left[\sqrt{t-2}\right], \frac{d}{dt}[3], \frac{d}{dt}\left[1/t^2\right] \right\rangle = \left\langle \tfrac{1}{2}(t-2)^{-1/2}, 0, -2t^{-3} \right\rangle = \left\langle \frac{1}{2\sqrt{t-2}}, 0, -\frac{2}{t^3} \right\rangle$$

10. $\mathbf{r}(t) = \left\langle e^{-t}, t - t^3, \ln t \right\rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \left\langle -e^{-t}, 1 - 3t^2, 1/t \right\rangle$

11. $\mathbf{r}(t) = t^2\,\mathbf{i} + \cos\left(t^2\right)\,\mathbf{j} + \sin^2 t\,\mathbf{k} \quad \Rightarrow$

$$\mathbf{r}'(t) = 2t\,\mathbf{i} + \left[-\sin(t^2)\cdot 2t\right]\mathbf{j} + (2\sin t\cdot\cos t)\,\mathbf{k} = 2t\,\mathbf{i} - 2t\sin(t^2)\,\mathbf{j} + 2\sin t\cos t\,\mathbf{k}$$

12. $\mathbf{r}(t) = \dfrac{1}{1+t}\,\mathbf{i} + \dfrac{t}{1+t}\,\mathbf{j} + \dfrac{t^2}{1+t}\,\mathbf{k} \quad \Rightarrow$

$$\mathbf{r}'(t) = \frac{0 - 1(1)}{(1+t)^2}\,\mathbf{i} + \frac{(1+t)\cdot 1 - t(1)}{(1+t)^2}\,\mathbf{j} + \frac{(1+t)\cdot 2t - t^2(1)}{(1+t)^2}\,\mathbf{k} = -\frac{1}{(1+t)^2}\,\mathbf{i} + \frac{1}{(1+t)^2}\,\mathbf{j} + \frac{t^2+2t}{(1+t)^2}\,\mathbf{k}$$

13. $\mathbf{r}(t) = t\sin t\,\mathbf{i} + e^t\cos t\,\mathbf{j} + \sin t\cos t\,\mathbf{k} \quad \Rightarrow$

$$\begin{aligned}\mathbf{r}'(t) &= [t\cdot\cos t + (\sin t)\cdot 1]\,\mathbf{i} + \left[e^t(-\sin t) + (\cos t)e^t\right]\mathbf{j} + [(\sin t)(-\sin t) + (\cos t)(\cos t)]\,\mathbf{k} \\ &= (t\cos t + \sin t)\,\mathbf{i} + e^t\,(\cos t - \sin t)\,\mathbf{j} + \left(\cos^2 t - \sin^2 t\right)\mathbf{k}\end{aligned}$$

14. $\mathbf{r}(t) = \sin^2 at\,\mathbf{i} + te^{bt}\,\mathbf{j} + \cos^2 ct\,\mathbf{k} \quad \Rightarrow$

$$\begin{aligned}\mathbf{r}'(t) &= [2(\sin at)\cdot(\cos at)(a)]\,\mathbf{i} + \left[t\cdot e^{bt}(b) + e^{bt}\cdot 1\right]\mathbf{j} + [2(\cos ct)\cdot(-\sin ct)(c)]\,\mathbf{k} \\ &= 2a\sin at\cos at\,\mathbf{i} + e^{bt}\,(bt+1)\,\mathbf{j} - 2c\sin ct\cos ct\,\mathbf{k}\end{aligned}$$

15. $\mathbf{r}(t) = \mathbf{a} + t\,\mathbf{b} + t^2\,\mathbf{c} \quad \Rightarrow \quad \mathbf{r}'(t) = \mathbf{0} + \mathbf{b} + 2t\,\mathbf{c} = \mathbf{b} + 2t\,\mathbf{c}$ by Formulas 1 and 3 of Theorem 3.

16. To find $\mathbf{r}'(t)$, by Formula 5 of Theorem 3, we first expand $\mathbf{r}(t) = t\,\mathbf{a}\times(\mathbf{b} + t\,\mathbf{c}) = t(\mathbf{a}\times\mathbf{b}) + t^2(\mathbf{a}\times\mathbf{c})$, so $\mathbf{r}'(t) = \mathbf{a}\times\mathbf{b} + 2t(\mathbf{a}\times\mathbf{c})$.

17. $\mathbf{r}(t) = \left\langle t^2 - 2t, 1 + 3t, \frac{1}{3}t^3 + \frac{1}{2}t^2 \right\rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \left\langle 2t - 2, 3, t^2 + t \right\rangle \quad \Rightarrow \quad \mathbf{r}'(2) = \langle 2, 3, 6\rangle$.

So $|\mathbf{r}'(2)| = \sqrt{2^2 + 3^2 + 6^2} = \sqrt{49} = 7$ and $\mathbf{T}(2) = \dfrac{\mathbf{r}'(2)}{|\mathbf{r}'(2)|} = \frac{1}{7}\langle 2, 3, 6\rangle = \left\langle \frac{2}{7}, \frac{3}{7}, \frac{6}{7} \right\rangle$.

18. $\mathbf{r}(t) = \left\langle \tan^{-1} t, 2e^{2t}, 8te^t \right\rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \left\langle 1/(1+t^2), 4e^{2t}, 8te^t + 8e^t \right\rangle \quad \Rightarrow \quad \mathbf{r}'(0) = \langle 1, 4, 8\rangle$.

So $|\mathbf{r}'(0)| = \sqrt{1^2 + 4^2 + 8^2} = \sqrt{81} = 9$ and $\mathbf{T}(0) = \dfrac{\mathbf{r}'(0)}{|\mathbf{r}'(0)|} = \frac{1}{9}\langle 1, 4, 8\rangle = \left\langle \frac{1}{9}, \frac{4}{9}, \frac{8}{9} \right\rangle$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $\mathbf{r}(t) = \cos t\,\mathbf{i} + 3t\,\mathbf{j} + 2\sin 2t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = -\sin t\,\mathbf{i} + 3\,\mathbf{j} + 4\cos 2t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(0) = 3\,\mathbf{j} + 4\,\mathbf{k}$. So

$|\mathbf{r}'(0)| = \sqrt{0^2 + 3^2 + 4^2} = \sqrt{25} = 5$ and $\mathbf{T}(0) = \dfrac{\mathbf{r}'(0)}{|\mathbf{r}'(0)|} = \frac{1}{5}(3\,\mathbf{j} + 4\,\mathbf{k}) = \frac{3}{5}\,\mathbf{j} + \frac{4}{5}\,\mathbf{k}$.

20. $\mathbf{r}(t) = \sin^2 t\,\mathbf{i} + \cos^2 t\,\mathbf{j} + \tan^2 t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = 2\sin t\cos t\,\mathbf{i} - 2\cos t\sin t\,\mathbf{j} + 2\tan t\sec^2 t\,\mathbf{k} \;\Rightarrow$

$\mathbf{r}'\left(\frac{\pi}{4}\right) = 2\cdot\frac{\sqrt{2}}{2}\cdot\frac{\sqrt{2}}{2}\,\mathbf{i} - 2\cdot\frac{\sqrt{2}}{2}\cdot\frac{\sqrt{2}}{2}\,\mathbf{j} + 2\cdot 1\cdot(\sqrt{2})^2\,\mathbf{k} = \mathbf{i} - \mathbf{j} + 4\,\mathbf{k}$. So $\left|\mathbf{r}'\left(\frac{\pi}{4}\right)\right| = \sqrt{1^2 + 1^2 + 4^2} = \sqrt{18} = 3\sqrt{2}$

and $\mathbf{T}\left(\frac{\pi}{4}\right) = \dfrac{\mathbf{r}'\left(\frac{\pi}{4}\right)}{\left|\mathbf{r}'\left(\frac{\pi}{4}\right)\right|} = \dfrac{1}{3\sqrt{2}}\,(\mathbf{i} - \mathbf{j} + 4\,\mathbf{k}) = \dfrac{1}{3\sqrt{2}}\,\mathbf{i} - \dfrac{1}{3\sqrt{2}}\,\mathbf{j} + \dfrac{4}{3\sqrt{2}}\,\mathbf{k}$.

21. The point $(2, -2, 4)$ corresponds to $t = 1$ [note that $4/t = 4$]. Then

$$\mathbf{r}(t) = \left\langle t^3 + 1, 3t - 5, 4/t \right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \left\langle 3t^2, 3, -4/t^2 \right\rangle \;\Rightarrow\; \mathbf{r}'(1) = \langle 3, 3, -4 \rangle$$

So
$$|\mathbf{r}'(1)| = \sqrt{3^2 + 3^2 + (-4)^2} = \sqrt{34}$$

and
$$\mathbf{T}(1) = \frac{\mathbf{r}'(1)}{|\mathbf{r}'(1)|} = \frac{1}{\sqrt{34}}\,\langle 3, 3, -4 \rangle = \left\langle \frac{3}{\sqrt{34}}, \frac{3}{\sqrt{34}}, -\frac{4}{\sqrt{34}} \right\rangle$$

22. The point $(0, 0, 1)$ corresponds to $t = 0$ [note that $5t = 0$]. Then

$$\mathbf{r}(t) = \sin t\,\mathbf{i} + 5t\,\mathbf{j} + \cos t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = \cos t\,\mathbf{i} + 5\,\mathbf{j} - \sin t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(0) = \mathbf{i} + 5\,\mathbf{j}$$

So
$$|\mathbf{r}'(0)| = \sqrt{1^2 + 5^2 + 0^2} = \sqrt{26}$$

and
$$\mathbf{T}(0) = \frac{\mathbf{r}'(0)}{|\mathbf{r}'(0)|} = \frac{1}{\sqrt{26}}(\mathbf{i} + 5\mathbf{j}) = \frac{1}{\sqrt{26}}\,\mathbf{i} + \frac{5}{\sqrt{26}}\,\mathbf{j}$$

23. $\mathbf{r}(t) = \left\langle t^4, t, t^2 \right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \left\langle 4t^3, 1, 2t \right\rangle$. Then $\mathbf{r}'(1) = \langle 4, 1, 2 \rangle$, $|\mathbf{r}'(1)| = \sqrt{4^2 + 1^2 + 2^2} = \sqrt{21}$, and

$$\mathbf{T}(1) = \frac{\mathbf{r}'(1)}{|\mathbf{r}'(1)|} = \frac{1}{\sqrt{21}}\,\langle 4, 1, 2 \rangle = \left\langle \frac{4}{\sqrt{21}}, \frac{1}{\sqrt{21}}, \frac{2}{\sqrt{21}} \right\rangle$$

$\mathbf{r}''(t) = \left\langle 12t^2, 0, 2 \right\rangle$, so

$$\mathbf{r}'(t) \times \mathbf{r}''(t) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 4t^3 & 1 & 2t \\ 12t^2 & 0 & 2 \end{vmatrix} = \begin{vmatrix} 1 & 2t \\ 0 & 2 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 4t^3 & 2t \\ 12t^2 & 2 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 4t^3 & 1 \\ 12t^2 & 0 \end{vmatrix}\mathbf{k}$$
$$= (2 - 0)\,\mathbf{i} - (8t^3 - 24t^3)\,\mathbf{j} + (0 - 12t^2)\,\mathbf{k} = \left\langle 2, 16t^3, -12t^2 \right\rangle$$

24. $\mathbf{r}(t) = \left\langle e^{2t}, e^{-3t}, t \right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \left\langle 2e^{2t}, -3e^{-3t}, 1 \right\rangle$. Then $\mathbf{r}'(0) = \langle 2, -3, 1 \rangle$, $|\mathbf{r}'(0)| = \sqrt{2^2 + (-3)^2 + 1^2} = \sqrt{14}$, and

$$\mathbf{T}(0) = \frac{\mathbf{r}'(0)}{|\mathbf{r}'(0)|} = \frac{1}{\sqrt{14}}\,\langle 2, -3, 1 \rangle = \left\langle \frac{2}{\sqrt{14}}, -\frac{3}{\sqrt{14}}, \frac{1}{\sqrt{14}} \right\rangle$$

$\mathbf{r}''(t) = \left\langle 4e^{2t}, 9e^{-3t}, 0 \right\rangle \;\Rightarrow\; \mathbf{r}''(0) = \langle 4, 9, 0 \rangle$. Then

$$\mathbf{r}'(0) \times \mathbf{r}''(0) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -3 & 1 \\ 4 & 9 & 0 \end{vmatrix} = \begin{vmatrix} -3 & 1 \\ 9 & 0 \end{vmatrix}\mathbf{i} - \begin{vmatrix} 2 & 1 \\ 4 & 0 \end{vmatrix}\mathbf{j} + \begin{vmatrix} 2 & -3 \\ 4 & 9 \end{vmatrix}\mathbf{k}$$
$$= (0 - 9)\,\mathbf{i} - (0 - 4)\,\mathbf{j} + [18 - (-12)]\,\mathbf{k} = \langle -9, 4, 30 \rangle$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. The vector equation for the curve is $\mathbf{r}(t) = \left\langle t^2+1, 4\sqrt{t}, e^{t^2-t}\right\rangle$, so $\mathbf{r}'(t) = \left\langle 2t, 2/\sqrt{t}, (2t-1)e^{t^2-t}\right\rangle$. The point $(2,4,1)$ corresponds to $t=1$, so the tangent vector there is $\mathbf{r}'(1) = \langle 2,2,1\rangle$. Thus, the tangent line goes through the point $(2,4,1)$ and is parallel to the vector $\langle 2,2,1\rangle$. Parametric equations are $x = 2+2t$, $y = 4+2t$, $z = 1+t$.

26. The vector equation for the curve is $\mathbf{r}(t) = \left\langle \ln(t+1), t\cos 2t, 2^t\right\rangle$, so $\mathbf{r}'(t) = \left\langle 1/(t+1), \cos 2t - 2t\sin 2t, 2^t\ln 2\right\rangle$. The point $(0,0,1)$ corresponds to $t=0$, so the tangent vector there is $\mathbf{r}'(0) = \langle 1,1,\ln 2\rangle$. Thus, the tangent line goes through the point $(0,0,1)$ and is parallel to the vector $\langle 1,1,\ln 2\rangle$. Parametric equations are $x = 0+1\cdot t = t$, $y = 0+1\cdot t = t$, $z = 1+(\ln 2)t$.

27. The vector equation for the curve is $\mathbf{r}(t) = \left\langle e^{-t}\cos t, e^{-t}\sin t, e^{-t}\right\rangle$, so

$$\begin{aligned}\mathbf{r}'(t) &= \left\langle e^{-t}(-\sin t) + (\cos t)(-e^{-t}), e^{-t}\cos t + (\sin t)(-e^{-t}), (-e^{-t})\right\rangle \\ &= \left\langle -e^{-t}(\cos t + \sin t), e^{-t}(\cos t - \sin t), -e^{-t}\right\rangle\end{aligned}$$

The point $(1,0,1)$ corresponds to $t=0$, so the tangent vector there is $\mathbf{r}'(0) = \left\langle -e^0(\cos 0 + \sin 0), e^0(\cos 0 - \sin 0), -e^0\right\rangle = \langle -1,1,-1\rangle$. Thus, the tangent line is parallel to the vector $\langle -1,1,-1\rangle$ and parametric equations are $x = 1+(-1)t = 1-t$, $y = 0+1\cdot t = t$, $z = 1+(-1)t = 1-t$.

28. The vector equation for the curve is $\mathbf{r}(t) = \left\langle \sqrt{t^2+3}, \ln(t^2+3), t\right\rangle$, so $\mathbf{r}'(t) = \left\langle t/\sqrt{t^2+3}, 2t/(t^2+3), 1\right\rangle$. At $(2,\ln 4,1)$, $t=1$ and $\mathbf{r}'(1) = \left\langle \frac{1}{2}, \frac{1}{2}, 1\right\rangle$. Thus, parametric equations of the tangent line are $x = 2+\frac{1}{2}t$, $y = \ln 4 + \frac{1}{2}t$, $z = 1+t$.

29. First we parametrize the curve C of intersection. The projection of C onto the xy-plane is contained in the circle $x^2+y^2 = 25$, $z=0$, so we can write $x = 5\cos t$, $y = 5\sin t$. C also lies on the cylinder $y^2+z^2 = 20$, and $z \geq 0$ near the point $(3,4,2)$, so we can write $z = \sqrt{20-y^2} = \sqrt{20-25\sin^2 t}$. A vector equation then for C is

$\mathbf{r}(t) = \left\langle 5\cos t, 5\sin t, \sqrt{20-25\sin^2 t}\right\rangle \quad\Rightarrow\quad \mathbf{r}'(t) = \left\langle -5\sin t, 5\cos t, \frac{1}{2}(20-25\sin^2 t)^{-1/2}(-50\sin t\cos t)\right\rangle$.

The point $(3,4,2)$ corresponds to $t = \cos^{-1}\left(\frac{3}{5}\right)$, so the tangent vector there is

$\mathbf{r}'\left(\cos^{-1}\left(\frac{3}{5}\right)\right) = \left\langle -5\left(\frac{4}{5}\right), 5\left(\frac{3}{5}\right), \frac{1}{2}\left(20-25\left(\frac{4}{5}\right)^2\right)^{-1/2}\left(-50\left(\frac{4}{5}\right)\left(\frac{3}{5}\right)\right)\right\rangle = \langle -4,3,-6\rangle$.

The tangent line is parallel to this vector and passes through $(3,4,2)$, so a vector equation for the line is $\mathbf{r}(t) = (3-4t)\mathbf{i} + (4+3t)\mathbf{j} + (2-6t)\mathbf{k}$.

30. $\mathbf{r}(t) = \left\langle 2\cos t, 2\sin t, e^t\right\rangle \quad\Rightarrow\quad \mathbf{r}'(t) = \left\langle -2\sin t, 2\cos t, e^t\right\rangle$. The tangent line to the curve is parallel to the plane when the curve's tangent vector is orthogonal to the plane's normal vector. Thus we require $\left\langle -2\sin t, 2\cos t, e^t\right\rangle \cdot \left\langle \sqrt{3},1,0\right\rangle = 0 \quad\Rightarrow$ $-2\sqrt{3}\sin t + 2\cos t + 0 = 0 \quad\Rightarrow\quad \tan t = \frac{1}{\sqrt{3}} \quad\Rightarrow\quad t = \frac{\pi}{6}$ [since $0 \leq t \leq \pi$].

$\mathbf{r}\left(\frac{\pi}{6}\right) = \left\langle \sqrt{3}, 1, e^{\pi/6}\right\rangle$, so the point is $(\sqrt{3}, 1, e^{\pi/6})$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. $\mathbf{r}(t) = \left\langle t, e^{-t}, 2t - t^2 \right\rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \left\langle 1, -e^{-t}, 2 - 2t \right\rangle$. At $(0, 1, 0)$, $t = 0$ and $\mathbf{r}'(0) = \langle 1, -1, 2 \rangle$. Thus, parametric equations of the tangent line are $x = t$, $y = 1 - t$, $z = 2t$.

32. $\mathbf{r}(t) = \langle 2\cos t, 2\sin t, 4\cos 2t \rangle$, $\mathbf{r}'(t) = \langle -2\sin t, 2\cos t, -8\sin 2t \rangle$. At $\left(\sqrt{3}, 1, 2\right)$, $t = \frac{\pi}{6}$ and $\mathbf{r}'\left(\frac{\pi}{6}\right) = \left\langle -1, \sqrt{3}, -4\sqrt{3} \right\rangle$. Thus, parametric equations of the tangent line are $x = \sqrt{3} - t$, $y = 1 + \sqrt{3}\,t$, $z = 2 - 4\sqrt{3}\,t$.

33. $\mathbf{r}(t) = \langle t\cos t, t, t\sin t \rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \langle \cos t - t\sin t, 1, t\cos t + \sin t \rangle$. At $(-\pi, \pi, 0)$, $t = \pi$ and $\mathbf{r}'(\pi) = \langle -1, 1, -\pi \rangle$. Thus, parametric equations of the tangent line are $x = -\pi - t$, $y = \pi + t$, $z = -\pi t$.

34. (a) The tangent line to the curve $\mathbf{r}(t) = \langle \sin \pi t, 2\sin \pi t, \cos \pi t \rangle$ at $t = 0$ is the line through the point with position vector $\mathbf{r}(0) = \langle \sin 0, 2\sin 0, \cos 0 \rangle = \langle 0, 0, 1 \rangle$, and in the direction of the tangent vector, $\mathbf{r}'(0) = \langle \pi\cos 0, 2\pi\cos 0, -\pi\sin 0 \rangle = \langle \pi, 2\pi, 0 \rangle$. So an equation of the line is $\langle x, y, z \rangle = \mathbf{r}(0) + u\,\mathbf{r}'(0) = \langle 0 + \pi u, 0 + 2\pi u, 1 \rangle = \langle \pi u, 2\pi u, 1 \rangle$.

$\mathbf{r}\left(\frac{1}{2}\right) = \left\langle \sin\frac{\pi}{2}, 2\sin\frac{\pi}{2}, \cos\frac{\pi}{2} \right\rangle = \langle 1, 2, 0 \rangle$,

$\mathbf{r}'\left(\frac{1}{2}\right) = \left\langle \pi\cos\frac{\pi}{2}, 2\pi\cos\frac{\pi}{2}, -\pi\sin\frac{\pi}{2} \right\rangle = \langle 0, 0, -\pi \rangle$.

So the equation of the second line is

$\langle x, y, z \rangle = \mathbf{r}\left(\frac{1}{2}\right) + v\mathbf{r}'\left(\frac{1}{2}\right) = \langle 1, 2, 0 \rangle + v\,\langle 0, 0, -\pi \rangle = \langle 1, 2, -\pi v \rangle$.

The lines intersect where $\langle \pi u, 2\pi u, 1 \rangle = \langle 1, 2, -\pi v \rangle$, so the point of intersection is $(1, 2, 1)$.

(b)

35. The angle of intersection of the two curves is the angle between the two tangent vectors to the curves at the point of intersection. Since $\mathbf{r}_1'(t) = \left\langle 1, 2t, 3t^2 \right\rangle$ and $t = 0$ at $(0, 0, 0)$, $\mathbf{r}_1'(0) = \langle 1, 0, 0 \rangle$ is a tangent vector to $\mathbf{r}_1$ at $(0, 0, 0)$. Similarly, $\mathbf{r}_2'(t) = \langle \cos t, 2\cos 2t, 1 \rangle$ and since $\mathbf{r}_2(0) = \langle 0, 0, 0 \rangle$, $\mathbf{r}_2'(0) = \langle 1, 2, 1 \rangle$ is a tangent vector to $\mathbf{r}_2$ at $(0, 0, 0)$. If θ is the angle between these two tangent vectors, then $\cos\theta = \frac{1}{\sqrt{1}\sqrt{6}}\,\langle 1, 0, 0 \rangle \cdot \langle 1, 2, 1 \rangle = \frac{1}{\sqrt{6}}$ and $\theta = \cos^{-1}\left(\frac{1}{\sqrt{6}}\right) \approx 66°$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. To find the point of intersection, we must find the values of t and s which satisfy the following three equations simultaneously: $t = 3 - s$, $1 - t = s - 2$, $3 + t^2 = s^2$. Solving the last two equations gives $t = 1$, $s = 2$ (check these in the first equation). Thus the point of intersection is $(1, 0, 4)$. To find the angle θ of intersection, we proceed as in Exercise 35. The tangent vectors to the respective curves at $(1, 0, 4)$ are $\mathbf{r}_1'(1) = \langle 1, -1, 2\rangle$ and $\mathbf{r}_2'(2) = \langle -1, 1, 4\rangle$. So

$\cos\theta = \frac{1}{\sqrt{6}\sqrt{18}}(-1 - 1 + 8) = \frac{6}{6\sqrt{3}} = \frac{1}{\sqrt{3}}$ and $\theta = \cos^{-1}\left(\frac{1}{\sqrt{3}}\right) \approx 55°$.

Note: In Exercise 35, the curves intersect when the value of both parameters is zero. However, as seen in this exercise, it is not necessary for the parameters to be of equal value at the point of intersection.

37.
$$\begin{aligned}\int_0^2 (t\,\mathbf{i} - t^3\,\mathbf{j} + 3t^5\,\mathbf{k})\,dt &= \left(\int_0^2 t\,dt\right)\mathbf{i} - \left(\int_0^2 t^3\,dt\right)\mathbf{j} + \left(\int_0^2 3t^5\,dt\right)\mathbf{k}\\ &= \left[\tfrac{1}{2}t^2\right]_0^2\mathbf{i} - \left[\tfrac{1}{4}t^4\right]_0^2\mathbf{j} + \left[\tfrac{1}{2}t^6\right]_0^2\mathbf{k}\\ &= \tfrac{1}{2}(4-0)\,\mathbf{i} - \tfrac{1}{4}(16-0)\,\mathbf{j} + \tfrac{1}{2}(64-0)\,\mathbf{k} = 2\,\mathbf{i} - 4\,\mathbf{j} + 32\,\mathbf{k}\end{aligned}$$

38.
$$\begin{aligned}\int_1^4 \left(2t^{3/2}\,\mathbf{i} + (t+1)\sqrt{t}\,\mathbf{k}\right)dt &= \left(\int_1^4 2t^{3/2}\,dt\right)\mathbf{i} + \left[\int_1^4 (t^{3/2} + t^{1/2})\,dt\right]\mathbf{k}\\ &= \left[\tfrac{4}{5}t^{5/2}\right]_1^4\mathbf{i} + \left[\tfrac{2}{5}t^{5/2} + \tfrac{2}{3}t^{3/2}\right]_1^4\mathbf{k}\\ &= \tfrac{4}{5}(4^{5/2} - 1)\,\mathbf{i} + \left(\tfrac{2}{5}(4)^{5/2} + \tfrac{2}{3}(4)^{3/2} - \tfrac{2}{5} - \tfrac{2}{3}\right)\mathbf{k}\\ &= \tfrac{4}{5}(31)\,\mathbf{i} + \left(\tfrac{2}{5}(32) + \tfrac{2}{3}(8) - \tfrac{2}{5} - \tfrac{2}{3}\right)\mathbf{k} = \tfrac{124}{5}\,\mathbf{i} + \tfrac{256}{15}\,\mathbf{k}\end{aligned}$$

39.
$$\begin{aligned}\int_0^1 \left(\frac{1}{t+1}\,\mathbf{i} + \frac{1}{t^2+1}\,\mathbf{j} + \frac{t}{t^2+1}\,\mathbf{k}\right)dt &= \left(\int_0^1 \frac{1}{t+1}\,dt\right)\mathbf{i} + \left(\int_0^1 \frac{1}{t^2+1}\,dt\right)\mathbf{j} + \left(\int_0^1 \frac{t}{t^2+1}\,dt\right)\mathbf{k}\\ &= \left[\ln|t+1|\right]_0^1\mathbf{i} + \left[\tan^{-1}t\right]_0^1\mathbf{j} + \left[\tfrac{1}{2}\ln(t^2+1)\right]_0^1\mathbf{k}\\ &= (\ln 2 - \ln 1)\,\mathbf{i} + (\tfrac{\pi}{4} - 0)\,\mathbf{j} + \tfrac{1}{2}(\ln 2 - \ln 1)\,\mathbf{k} = \ln 2\,\mathbf{i} + \tfrac{\pi}{4}\,\mathbf{j} + \tfrac{1}{2}\ln 2\,\mathbf{k}\end{aligned}$$

40.
$$\begin{aligned}&\int_0^{\pi/4}(\sec t\tan t\,\mathbf{i} + t\cos 2t\,\mathbf{j} + \sin^2 2t\cos 2t\,\mathbf{k})\,dt\\ &= \left(\int_0^{\pi/4}\sec t\tan t\,dt\right)\mathbf{i} + \left(\int_0^{\pi/4} t\cos 2t\,dt\right)\mathbf{j} + \left(\int_0^{\pi/4}\sin^2 2t\cos 2t\,dt\right)\mathbf{k}\\ &= \left[\sec t\right]_0^{\pi/4}\mathbf{i} + \left(\left[\tfrac{1}{2}t\sin 2t\right]_0^{\pi/4} - \int_0^{\pi/4}\tfrac{1}{2}\sin 2t\,dt\right)\mathbf{j} + \left[\tfrac{1}{6}\sin^3 2t\right]_0^{\pi/4}\mathbf{k}\\ &\qquad\qquad \text{[For the } y\text{-component, integrate by parts with } u = t,\ dv = \cos 2t\,dt.]\\ &= (\sec\tfrac{\pi}{4} - \sec 0)\,\mathbf{i} + \left(\tfrac{\pi}{8}\sin\tfrac{\pi}{2} - 0 - \left[-\tfrac{1}{4}\cos 2t\right]_0^{\pi/4}\right)\mathbf{j} + \tfrac{1}{6}\left(\sin^3\tfrac{\pi}{2} - \sin^3 0\right)\mathbf{k}\\ &= (\sqrt{2} - 1)\,\mathbf{i} + \left(\tfrac{\pi}{8} + \tfrac{1}{4}\cos\tfrac{\pi}{2} - \tfrac{1}{4}\cos 0\right)\mathbf{j} + \tfrac{1}{6}(1 - 0)\,\mathbf{k} = (\sqrt{2} - 1)\,\mathbf{i} + \left(\tfrac{\pi}{8} - \tfrac{1}{4}\right)\mathbf{j} + \tfrac{1}{6}\,\mathbf{k}\end{aligned}$$

41.
$$\begin{aligned}\int \left(\frac{1}{1+t^2}\,\mathbf{i} + te^{t^2}\,\mathbf{j} + \sqrt{t}\,\mathbf{k}\right)dt &= \left(\int \frac{1}{1+t^2}\,dt\right)\mathbf{i} + \left(\int te^{t^2}\,dt\right)\mathbf{j} + \left(\int \sqrt{t}\,dt\right)\mathbf{k}\\ &= \tan^{-1}t\,\mathbf{i} + \tfrac{1}{2}e^{t^2}\,\mathbf{j} + \tfrac{2}{3}t^{3/2}\,\mathbf{k} + \mathbf{C}\end{aligned}$$

where $\mathbf{C}$ is a vector constant of integration.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. $$\int \left(t\cos t^2\,\mathbf{i} + \frac{1}{t}\,\mathbf{j} + \sec^2 t\,\mathbf{k}\right) dt = \left(\int t\cos t^2\,dt\right)\mathbf{i} + \left(\int \frac{1}{t}\,dt\right)\mathbf{j} + \left(\int \sec^2 t\,dt\right)\mathbf{k}$$
$$= \tfrac{1}{2}\sin t^2\,\mathbf{i} + \ln|t|\,\mathbf{j} + \tan t\,\mathbf{k} + \mathbf{C}$$

where $\mathbf{C}$ is a vector constant of integration.

43. $\mathbf{r}'(t) = 2t\,\mathbf{i} + 3t^2\,\mathbf{j} + \sqrt{t}\,\mathbf{k} \quad\Rightarrow\quad \mathbf{r}(t) = t^2\,\mathbf{i} + t^3\,\mathbf{j} + \frac{2}{3}t^{3/2}\,\mathbf{k} + \mathbf{C},$ where $\mathbf{C}$ is a constant vector.

But $\mathbf{i} + \mathbf{j} = \mathbf{r}\,(1) = \mathbf{i} + \mathbf{j} + \frac{2}{3}\mathbf{k} + \mathbf{C}$. Thus $\mathbf{C} = -\frac{2}{3}\mathbf{k}$ and $\mathbf{r}(t) = t^2\,\mathbf{i} + t^3\,\mathbf{j} + \left(\frac{2}{3}t^{3/2} - \frac{2}{3}\right)\mathbf{k}$.

44. $\mathbf{r}'(t) = t\,\mathbf{i} + e^t\,\mathbf{j} + te^t\,\mathbf{k} \quad\Rightarrow\quad \mathbf{r}(t) = \frac{1}{2}t^2\,\mathbf{i} + e^t\,\mathbf{j} + \left(te^t - e^t\right)\,\mathbf{k} + \mathbf{C}$. But $\mathbf{i} + \mathbf{j} + \mathbf{k} = \mathbf{r}\,(0) = \mathbf{j} - \mathbf{k} + \mathbf{C}$.

Thus $\mathbf{C} = \mathbf{i} + 2\,\mathbf{k}$ and $\mathbf{r}(t) = \left(\frac{1}{2}t^2 + 1\right)\mathbf{i} + e^t\,\mathbf{j} + (te^t - e^t + 2)\,\mathbf{k}$.

For Exercises 45–48, let $\mathbf{u}(t) = \langle u_1(t), u_2(t), u_3(t)\rangle$ and $\mathbf{v}(t) = \langle v_1(t), v_2(t), v_3(t)\rangle$. In each of these exercises, the procedure is to apply Theorem 2 so that the corresponding properties of derivatives of real-valued functions can be used.

45. $$\frac{d}{dt}\left[\mathbf{u}(t) + \mathbf{v}(t)\right] = \frac{d}{dt}\left\langle u_1(t) + v_1(t), u_2(t) + v_2(t), u_3(t) + v_3(t)\right\rangle$$
$$= \left\langle \frac{d}{dt}\left[u_1(t) + v_1\,(t)\right], \frac{d}{dt}\left[u_2(t) + v_2(t)\right], \frac{d}{dt}\left[u_3(t) + v_3(t)\right]\right\rangle$$
$$= \langle u_1'(t) + v_1'(t), u_2'(t) + v_2'(t), u_3'(t) + v_3'(t)\rangle$$
$$= \langle u_1'(t), u_2'\,(t)\,, u_3'(t)\rangle + \langle v_1'(t), v_2'(t), v_3'(t)\rangle = \mathbf{u}'(t) + \mathbf{v}'(t)$$

46. $$\frac{d}{dt}\left[f(t)\,\mathbf{u}(t)\right] = \frac{d}{dt}\left\langle f(t)u_1(t), f(t)u_2(t), f(t)u_3(t)\right\rangle$$
$$= \left\langle \frac{d}{dt}\left[f(t)u_1(t)\right], \frac{d}{dt}\left[f(t)u_2(t)\right], \frac{d}{dt}\left[f(t)u_3(t)\right]\right\rangle$$
$$= \langle f'(t)u_1(t) + f(t)u_1'(t), f'(t)u_2(t) + f(t)u_2'(t), f'(t)u_3(t) + f(t)u_3'(t)\rangle$$
$$= f'(t)\,\langle u_1(t), u_2(t), u_3(t)\rangle + f(t)\,\langle u_1'(t), u_2'(t), u_3'(t)\rangle = f'(t)\,\mathbf{u}(t) + f(t)\,\mathbf{u}'(t)$$

47. $$\frac{d}{dt}\left[\mathbf{u}(t) \times \mathbf{v}(t)\right] = \frac{d}{dt}\left\langle u_2(t)v_3(t) - u_3(t)v_2(t), u_3(t)v_1(t) - u_1(t)v_3(t), u_1(t)v_2(t) - u_2(t)v_1(t)\right\rangle$$
$$= \langle u_2'v_3(t) + u_2(t)v_3'(t) - u_3'(t)v_2(t) - u_3(t)v_2'(t),$$
$$u_3'(t)v_1(t) + u_3(t)v_1'\,(t) - u_1'(t)v_3(t) - u_1(t)v_3'(t),$$
$$u_1'(t)v_2(t) + u_1(t)v_2'(t) - u_2'(t)v_1(t) - u_2(t)v_1'(t)\rangle$$
$$= \langle u_2'(t)v_3(t) - u_3'(t)v_2\,(t)\,, u_3'(t)v_1(t) - u_1'(t)v_3(t), u_1'(t)v_2(t) - u_2'(t)v_1(t)\rangle$$
$$+ \langle u_2(t)v_3'(t) - u_3(t)v_2'(t), u_3(t)v_1'\,(t) - u_1(t)v_3'(t), u_1(t)v_2'(t) - u_2(t)v_1'(t)\rangle$$
$$= \mathbf{u}'(t) \times \mathbf{v}(t) + \mathbf{u}(t) \times \mathbf{v}'(t)$$

Alternate solution: Let $\mathbf{r}(t) = \mathbf{u}(t) \times \mathbf{v}(t)$. Then

$$\mathbf{r}(t+h) - \mathbf{r}(t) = [\mathbf{u}(t+h) \times \mathbf{v}(t+h)] - [\mathbf{u}(t) \times \mathbf{v}(t)]$$
$$= [\mathbf{u}(t+h) \times \mathbf{v}(t+h)] - [\mathbf{u}(t) \times \mathbf{v}(t)] + [\mathbf{u}(t+h) \times \mathbf{v}(t)] - [\mathbf{u}(t+h) \times \mathbf{v}(t)]$$
$$= \mathbf{u}(t+h) \times [\mathbf{v}(t+h) - \mathbf{v}(t)] + [\mathbf{u}(t+h) - \mathbf{u}(t)] \times \mathbf{v}(t)$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(Be careful of the order of the cross product.) Dividing through by h and taking the limit as $h \to 0$ we have

$$\mathbf{r}'(t) = \lim_{h\to 0} \frac{\mathbf{u}(t+h) \times [\mathbf{v}(t+h) - \mathbf{v}(t)]}{h} + \lim_{h\to 0} \frac{[\mathbf{u}(t+h) - \mathbf{u}(t)] \times \mathbf{v}(t)}{h} = \mathbf{u}(t) \times \mathbf{v}'(t) + \mathbf{u}'(t) \times \mathbf{v}(t)$$

by Exercise 13.1.61(a) and Definition 1.

48. $$\frac{d}{dt}[\mathbf{u}(f(t))] = \frac{d}{dt}\langle u_1(f(t)), u_2(f(t)), u_3(f(t))\rangle = \left\langle \frac{d}{dt}[u_1(f(t))], \frac{d}{dt}[u_2(f(t))], \frac{d}{dt}[u_3(f(t))] \right\rangle$$
$$= \langle f'(t)u_1'(f(t)), f'(t)u_2'(f(t)), f'(t)u_3'(f(t))\rangle = f'(t)\,\mathbf{u}'(t)$$

49. $$\frac{d}{dt}[\mathbf{u}(t)\cdot\mathbf{v}(t)] = \mathbf{u}'(t)\cdot\mathbf{v}(t) + \mathbf{u}(t)\cdot\mathbf{v}'(t) \qquad \text{[by Formula 4 of Theorem 3]}$$
$$= \langle \cos t, -\sin t, 1\rangle \cdot \langle t, \cos t, \sin t\rangle + \langle \sin t, \cos t, t\rangle \cdot \langle 1, -\sin t, \cos t\rangle$$
$$= t\cos t - \cos t\,\sin t + \sin t + \sin t - \cos t\,\sin t + t\cos t$$
$$= 2t\cos t + 2\sin t - 2\cos t\,\sin t$$

50. $$\frac{d}{dt}[\mathbf{u}(t)\times\mathbf{v}(t)] = \mathbf{u}'(t)\times\mathbf{v}(t) + \mathbf{u}(t)\times\mathbf{v}'(t) \qquad \text{[by Formula 5 of Theorem 3]}$$
$$= \langle \cos t, -\sin t, 1\rangle \times \langle t, \cos t, \sin t\rangle + \langle \sin t, \cos t, t\rangle \times \langle 1, -\sin t, \cos t\rangle$$
$$= \left\langle -\sin^2 t - \cos t, t - \cos t\,\sin t, \cos^2 t + t\sin t\right\rangle$$
$$+ \left\langle \cos^2 t + t\sin t, t - \cos t\,\sin t, -\sin^2 t - \cos t\right\rangle$$
$$= \left\langle \cos^2 t - \sin^2 t - \cos t + t\sin t, 2t - 2\cos t\,\sin t, \cos^2 t - \sin^2 t - \cos t + t\sin t\right\rangle$$

51. By Formula 4 of Theorem 3, $f'(t) = \mathbf{u}'(t)\cdot\mathbf{v}(t) + \mathbf{u}(t)\cdot\mathbf{v}'(t)$, and $\mathbf{v}'(t) = \left\langle 1, 2t, 3t^2\right\rangle$, so

$f'(2) = \mathbf{u}'(2)\cdot\mathbf{v}(2) + \mathbf{u}(2)\cdot\mathbf{v}'(2) = \langle 3,0,4\rangle\cdot\langle 2,4,8\rangle + \langle 1,2,-1\rangle\cdot\langle 1,4,12\rangle = 6+0+32+1+8-12 = 35.$

52. By Formula 5 of Theorem 3, $\mathbf{r}'(t) = \mathbf{u}'(t)\times\mathbf{v}(t) + \mathbf{u}(t)\times\mathbf{v}'(t)$, so

$$\mathbf{r}'(2) = \mathbf{u}'(2)\times\mathbf{v}(2) + \mathbf{u}(2)\times\mathbf{v}'(2) = \langle 3,0,4\rangle\times\langle 2,4,8\rangle + \langle 1,2,-1\rangle\times\langle 1,4,12\rangle$$
$$= \langle -16,-16,12\rangle + \langle 28,-13,2\rangle = \langle 12,-29,14\rangle$$

53. $\mathbf{r}(t) = \mathbf{a}\cos\omega t + \mathbf{b}\sin\omega t \quad\Rightarrow\quad \mathbf{r}'(t) = -\mathbf{a}\,\omega\sin\omega t + \mathbf{b}\,\omega\cos\omega t$ by Formulas 1 and 3 of Theorem 3. Then

$$\mathbf{r}(t)\times\mathbf{r}'(t) = (\mathbf{a}\cos\omega t + \mathbf{b}\sin\omega t)\times(-\mathbf{a}\,\omega\sin\omega t + \mathbf{b}\,\omega\cos\omega t)$$
$$= (\mathbf{a}\cos\omega t + \mathbf{b}\sin\omega t)\times(-\mathbf{a}\,\omega\sin\omega t) + (\mathbf{a}\cos\omega t + \mathbf{b}\sin\omega t)\times(\mathbf{b}\,\omega\cos\omega t)$$
[by Property 3 of Theorem 12.4.11]
$$= \mathbf{a}\cos\omega t\times(-\mathbf{a}\,\omega\sin\omega t) + \mathbf{b}\sin\omega t\times(-\mathbf{a}\,\omega\sin\omega t) + \mathbf{a}\cos\omega t\times\mathbf{b}\,\omega\cos\omega t + \mathbf{b}\sin\omega t\times\mathbf{b}\,\omega\cos\omega t$$
[by Property 4]
$$= (\cos\omega t)(-\omega\sin\omega t)(\mathbf{a}\times\mathbf{a}) + (\sin\omega t)(-\omega\sin\omega t)(\mathbf{b}\times\mathbf{a}) + (\cos\omega t)(\omega\cos\omega t)(\mathbf{a}\times\mathbf{b})$$
$$+ (\sin\omega t)(\omega\cos\omega t)(\mathbf{b}\times\mathbf{b}) \qquad \text{[by Property 2]}$$
$$= \mathbf{0} + \left(\omega\sin^2\omega t\right)(\mathbf{a}\times\mathbf{b}) + \left(\omega\cos^2\omega t\right)(\mathbf{a}\times\mathbf{b}) + \mathbf{0} \qquad \text{[by Property 1 and Example 12.4.2]}$$
$$= \omega\left(\sin^2\omega t + \cos^2\omega t\right)(\mathbf{a}\times\mathbf{b}) = \omega(\mathbf{a}\times\mathbf{b}) = \omega\,\mathbf{a}\times\mathbf{b} \qquad \text{[by Property 2]}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. From Exercise 53, $\mathbf{r}'(t) = -\mathbf{a}\,\omega \sin \omega t + \mathbf{b}\,\omega \cos \omega t \quad \Rightarrow \quad \mathbf{r}''(t) = -\mathbf{a}\,\omega^2 \cos \omega t - \mathbf{b}\,\omega^2 \sin \omega t$. Then

$$\begin{aligned} \mathbf{r}''(t) + \omega^2 \mathbf{r}(t) &= \left(-\mathbf{a}\,\omega^2 \cos \omega t - \mathbf{b}\,\omega^2 \sin \omega t\right) + \omega^2 \left(\mathbf{a} \cos \omega t + \mathbf{b} \sin \omega t\right) \\ &= -\mathbf{a}\,\omega^2 \cos \omega t - \mathbf{b}\,\omega^2 \sin \omega t + \mathbf{a}\,\omega^2 \cos \omega t + \mathbf{b}\,\omega^2 \sin \omega t = \mathbf{0} \end{aligned}$$

55. $\dfrac{d}{dt}\left[\mathbf{r}(t) \times \mathbf{r}'(t)\right] = \mathbf{r}'(t) \times \mathbf{r}'(t) + \mathbf{r}(t) \times \mathbf{r}''(t)$ by Formula 5 of Theorem 3. But $\mathbf{r}'(t) \times \mathbf{r}'(t) = \mathbf{0}$ (by Example 12.4.2).

Thus, $\dfrac{d}{dt}\left[\mathbf{r}(t) \times \mathbf{r}'(t)\right] = \mathbf{r}(t) \times \mathbf{r}''(t)$.

56.
$$\begin{aligned} \frac{d}{dt}\left(\mathbf{u}(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}(t)\right]\right) &= \mathbf{u}'(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}(t)\right] + \mathbf{u}(t) \cdot \frac{d}{dt}\left[\mathbf{v}(t) \times \mathbf{w}(t)\right] \\ &= \mathbf{u}'(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}(t)\right] + \mathbf{u}(t) \cdot \left[\mathbf{v}'(t) \times \mathbf{w}(t) + \mathbf{v}(t) \times \mathbf{w}'(t)\right] \\ &= \mathbf{u}'(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}(t)\right] + \mathbf{u}(t) \cdot \left[\mathbf{v}'(t) \times \mathbf{w}(t)\right] + \mathbf{u}(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}'(t)\right] \\ &= \mathbf{u}'(t) \cdot \left[\mathbf{v}(t) \times \mathbf{w}(t)\right] - \mathbf{v}'(t) \cdot \left[\mathbf{u}(t) \times \mathbf{w}(t)\right] + \mathbf{w}'(t) \cdot \left[\mathbf{u}(t) \times \mathbf{v}(t)\right] \end{aligned}$$

57. $\dfrac{d}{dt}\left|\mathbf{r}(t)\right| = \dfrac{d}{dt}\left[\mathbf{r}(t) \cdot \mathbf{r}(t)\right]^{1/2} = \frac{1}{2}\left[\mathbf{r}(t) \cdot \mathbf{r}(t)\right]^{-1/2}\left[2\mathbf{r}(t) \cdot \mathbf{r}'(t)\right] = \dfrac{1}{\left|\mathbf{r}(t)\right|}\,\mathbf{r}(t) \cdot \mathbf{r}'(t)$

58. Since $\mathbf{r}(t) \cdot \mathbf{r}'(t) = 0$, by Formula 4 of Theorem 3 we have

$\dfrac{d}{dt}\left|\mathbf{r}(t)\right|^2 = \dfrac{d}{dt}\left[\mathbf{r}(t) \cdot \mathbf{r}(t)\right] = \mathbf{r}'(t) \cdot \mathbf{r}(t) + \mathbf{r}(t) \cdot \mathbf{r}'(t) = 2\left[\mathbf{r}(t) \cdot \mathbf{r}'(t)\right] = 0$. This is true for all t, thus $\left|\mathbf{r}(t)\right|^2$, and so $\left|\mathbf{r}(t)\right|$ is a constant, and hence the curve lies on a sphere with center the origin.

59. Since $\mathbf{u}(t) = \mathbf{r}(t) \cdot \left[\mathbf{r}'(t) \times \mathbf{r}''(t)\right]$,

$$\begin{aligned} \mathbf{u}'(t) &= \mathbf{r}'(t) \cdot \left[\mathbf{r}'(t) \times \mathbf{r}''(t)\right] + \mathbf{r}(t) \cdot \frac{d}{dt}\left[\mathbf{r}'(t) \times \mathbf{r}''(t)\right] \\ &= 0 + \mathbf{r}(t) \cdot \left[\mathbf{r}''(t) \times \mathbf{r}''(t) + \mathbf{r}'(t) \times \mathbf{r}'''(t)\right] && \left[\text{since } \mathbf{r}'(t) \perp \mathbf{r}'(t) \times \mathbf{r}''(t)\right] \\ &= \mathbf{r}(t) \cdot \left[\mathbf{r}'(t) \times \mathbf{r}'''(t)\right] && \left[\text{since } \mathbf{r}''(t) \times \mathbf{r}''(t) = \mathbf{0}\right] \end{aligned}$$

60. The tangent vector $\mathbf{r}'(t)$ is defined as $\lim\limits_{h \to 0} \dfrac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$. Here we assume that this limit exists and $\mathbf{r}'(t) \neq \mathbf{0}$; then we know that this vector lies on the tangent line to the curve. As in Figure 1, let points P and Q have position vectors $\mathbf{r}(t)$ and $\mathbf{r}(t+h)$. The vector $\mathbf{r}(t+h) - \mathbf{r}(t)$ points from P to Q, so $\mathbf{r}(t+h) - \mathbf{r}(t) = \overrightarrow{PQ}$. If $h > 0$ then $t < t+h$, so Q lies "ahead" of P on the curve. If h is sufficiently small (we can take h to be as small as we like since $h \to 0$) then $\overrightarrow{PQ}$ approximates the curve from P to Q and hence points approximately in the direction of the curve as t increases. Since h is positive, $\dfrac{1}{h}\,\overrightarrow{PQ} = \dfrac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$ points in the same direction. If $h < 0$, then $t > t+h$ so Q lies "behind" P on the curve. For h sufficiently small, $\overrightarrow{PQ}$ approximates the curve but points in the direction of decreasing t. However, h is negative, so $\dfrac{1}{h}\,\overrightarrow{PQ} = \dfrac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$ points in the opposite direction, that is, in the direction of increasing t. In both cases, the difference quotient $\dfrac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$ points in the direction of increasing t. The tangent vector $\mathbf{r}'(t)$ is the limit of this difference quotient, so it must also point in the direction of increasing t.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13.3 Arc Length and Curvature

1. (a) $\mathbf{r}(t) = \langle f(t), g(t), h(t)\rangle = \langle 3-t, 2t, 4t+1\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle f'(t), g'(t), h'(t)\rangle = \langle -1, 2, 4\rangle$. Then

$$L = \int_1^3 \sqrt{(-1)^2 + 2^2 + 4^2}\,dt = \int_1^3 \sqrt{21}\,dt = \left[t\sqrt{21}\right]_1^3 = 3\sqrt{21} - \sqrt{21} = 2\sqrt{21}$$

(b) $\mathbf{r}(1) = \langle 2, 2, 5\rangle \;\Rightarrow\; P_1 = (2, 2, 5)$; $\mathbf{r}(3) = \langle 0, 6, 13\rangle \;\Rightarrow\; P_2 = (0, 6, 13)$. Then

$$|P_2P_1| = \sqrt{(0-2)^2 + (6-2)^2 + (13-5)^2} = \sqrt{84} = 2\sqrt{21}$$

2. (a) $\mathbf{r}(t) = f(t)\,\mathbf{i} + g(t)\,\mathbf{j} + h(t)\,\mathbf{k} = (t+2)\,\mathbf{i} - t\,\mathbf{j} + (3t-5)\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = \mathbf{i} - \mathbf{j} + 3\,\mathbf{k}$. Then

$$L = \int_{-1}^2 \sqrt{1^2 + (-1)^2 + 3^2}\,dt = \int_{-1}^2 \sqrt{11}\,dt = \left[t\sqrt{11}\right]_{-1}^2 = 2\sqrt{11} + \sqrt{11} = 3\sqrt{11}$$

(b) $\mathbf{r}(-1) = \mathbf{i} + \mathbf{j} - 8\,\mathbf{k} \;\Rightarrow\; P_1 = (1, 1, -8)$; $\mathbf{r}(2) = 4\,\mathbf{i} - 2\,\mathbf{j} + \mathbf{k} \;\Rightarrow\; P_2 = (4, -2, 1)$. Then

$$|P_2P_1| = \sqrt{(4-1)^2 + (-2-1)^2 + [1-(-8)]^2} = \sqrt{99} = 3\sqrt{11}$$

3. $\mathbf{r}(t) = \langle t, 3\cos t, 3\sin t\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1, -3\sin t, 3\cos t\rangle \;\Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{1^2 + (-3\sin t)^2 + (3\cos t)^2} = \sqrt{1 + 9(\sin^2 t + \cos^2 t)} = \sqrt{10}$.

Then using Formula 3, we have $L = \int_{-5}^5 |\mathbf{r}'(t)|\,dt = \int_{-5}^5 \sqrt{10}\,dt = \sqrt{10}\,t\big]_{-5}^5 = 10\sqrt{10}$.

4. $\mathbf{r}(t) = \left\langle 2t, t^2, \frac{1}{3}t^3\right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \left\langle 2, 2t, t^2\right\rangle \;\Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{2^2 + (2t)^2 + (t^2)^2} = \sqrt{4 + 4t^2 + t^4} = \sqrt{(2+t^2)^2} = 2 + t^2$ for $0 \le t \le 1$. Then using Formula 3, we have

$L = \int_0^1 |\mathbf{r}'(t)|\,dt = \int_0^1 (2+t^2)\,dt = 2t + \frac{1}{3}t^3\big]_0^1 = \frac{7}{3}$.

5. $\mathbf{r}(t) = \sqrt{2}\,t\,\mathbf{i} + e^t\mathbf{j} + e^{-t}\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = \sqrt{2}\,\mathbf{i} + e^t\mathbf{j} - e^{-t}\mathbf{k} \;\Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{\left(\sqrt{2}\right)^2 + (e^t)^2 + (-e^{-t})^2} = \sqrt{2 + e^{2t} + e^{-2t}} = \sqrt{(e^t + e^{-t})^2} = e^t + e^{-t}$ [since $e^t + e^{-t} > 0$].

Then $L = \int_0^1 |\mathbf{r}'(t)|\,dt = \int_0^1 (e^t + e^{-t})\,dt = \left[e^t - e^{-t}\right]_0^1 = e - e^{-1}$.

6. $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j} + \ln\cos t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j} + \dfrac{-\sin t}{\cos t}\,\mathbf{k} = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j} - \tan t\,\mathbf{k}$,

$|\mathbf{r}'(t)| = \sqrt{(-\sin t)^2 + \cos^2 t + (-\tan t)^2} = \sqrt{1 + \tan^2 t} = \sqrt{\sec^2 t} = |\sec t|$. Since $\sec t > 0$ for $0 \le t \le \pi/4$, here we can say $|\mathbf{r}'(t)| = \sec t$. Then

$$L = \int_0^{\pi/4} \sec t\,dt = \Big[\ln|\sec t + \tan t|\Big]_0^{\pi/4} = \ln\left|\sec\tfrac{\pi}{4} + \tan\tfrac{\pi}{4}\right| - \ln|\sec 0 + \tan 0|$$
$$= \ln\left|\sqrt{2} + 1\right| - \ln|1 + 0| = \ln(\sqrt{2} + 1)$$

7. $\mathbf{r}(t) = \mathbf{i} + t^2\,\mathbf{j} + t^3\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = 2t\,\mathbf{j} + 3t^2\,\mathbf{k} \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{4t^2 + 9t^4} = t\sqrt{4 + 9t^2}$ [since $t \ge 0$].

Then $L = \int_0^1 |\mathbf{r}'(t)|\,dt = \int_0^1 t\sqrt{4 + 9t^2}\,dt = \frac{1}{18}\cdot\frac{2}{3}(4 + 9t^2)^{3/2}\Big]_0^1 = \frac{1}{27}(13^{3/2} - 4^{3/2}) = \frac{1}{27}(13^{3/2} - 8)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $\mathbf{r}(t) = t^2\,\mathbf{i} + 9t\,\mathbf{j} + 4t^{3/2}\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = 2t\,\mathbf{i} + 9\,\mathbf{j} + 6\sqrt{t}\,\mathbf{k} \;\Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{4t^2 + 81 + 36t} = \sqrt{(2t+9)^2} = |2t+9| = 2t+9$ [since $2t+9 \geq 0$ for $1 \leq t \leq 4$]. Then

$L = \int_1^4 |\mathbf{r}'(t)|\,dt = \int_1^4 (2t+9)\,dt = \Big[t^2 + 9t\Big]_1^4 = 52 - 10 = 42.$

9. $\mathbf{r}(t) = \langle t^2, t^3, t^4\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 2t, 3t^2, 4t^3\rangle \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{(2t)^2 + (3t^2)^2 + (4t^3)^2} = \sqrt{4t^2 + 9t^4 + 16t^6}$, so

$L = \int_0^2 |\mathbf{r}'(t)|\,dt = \int_0^2 \sqrt{4t^2 + 9t^4 + 16t^6}\,dt \approx 18.6833.$

10. $\mathbf{r}(t) = \langle t, e^{-t}, te^{-t}\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1, -e^{-t}, (1-t)e^{-t}\rangle \;\Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{1^2 + (-e^{-t})^2 + [(1-t)e^{-t}]^2} = \sqrt{1 + e^{-2t} + (1-t)^2 e^{-2t}} = \sqrt{1 + (2 - 2t + t^2)e^{-2t}}$, so

$L = \int_1^3 |\mathbf{r}'(t)|\,dt = \int_1^3 \sqrt{1 + (2 + 2t + t^2)e^{-2t}}\,dt \approx 2.0454.$

11. $\mathbf{r}(t) = \langle \cos \pi t, 2t, \sin 2\pi t\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle -\pi \sin \pi t, 2, 2\pi \cos 2\pi t\rangle \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{\pi^2 \sin^2 \pi t + 4 + 4\pi^2 \cos^2 2\pi t}.$

The point $(1, 0, 0)$ corresponds to $t = 0$ and $(1, 4, 0)$ corresponds to $t = 2$, so the length is

$L = \int_0^2 |\mathbf{r}'(t)|\,dt = \int_0^2 \sqrt{\pi^2 \sin^2 \pi t + 4 + 4\pi^2 \cos^2 2\pi t}\,dt \approx 10.3311.$

12. We plot two different views of the curve with parametric equations $x = \sin t$, $y = \sin 2t$, $z = \sin 3t$. To help visualize the curve, we also include a plot showing a tube of radius 0.07 around the curve.

The complete curve is given by the parameter interval $[0, 2\pi]$ and we have $\mathbf{r}'(t) = \langle \cos t, 2\cos 2t, 3\cos 3t\rangle \;\Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{\cos^2 t + 4\cos^2 2t + 9\cos^2 3t}$, so $L = \int_0^{2\pi} |\mathbf{r}'(t)|\;dt = \int_0^{2\pi} \sqrt{\cos^2 t + 4\cos^2 2t + 9\cos^2 3t}\,dt \approx 16.0264.$

13. The projection of the curve C onto the xy-plane is the curve $x^2 = 2y$ or $y = \frac{1}{2}x^2$, $z = 0$. Then we can choose the parameter $x = t \;\Rightarrow\; y = \frac{1}{2}t^2$. Since C also lies on the surface $3z = xy$, we have $z = \frac{1}{3}xy = \frac{1}{3}(t)(\frac{1}{2}t^2) = \frac{1}{6}t^3$. Then parametric equations for C are $x = t$, $y = \frac{1}{2}t^2$, $z = \frac{1}{6}t^3$ and the corresponding vector equation is $\mathbf{r}(t) = \langle t, \frac{1}{2}t^2, \frac{1}{6}t^3\rangle$. The origin corresponds to $t = 0$ and the point $(6, 18, 36)$ corresponds to $t = 6$, so

$$L = \int_0^6 |\mathbf{r}'(t)|\,dt = \int_0^6 \left|\langle 1, t, \tfrac{1}{2}t^2\rangle\right|\,dt = \int_0^6 \sqrt{1^2 + t^2 + \left(\tfrac{1}{2}t^2\right)^2}\,dt = \int_0^6 \sqrt{1 + t^2 + \tfrac{1}{4}t^4}\,dt$$
$$= \int_0^6 \sqrt{(1 + \tfrac{1}{2}t^2)^2}\,dt = \int_0^6 (1 + \tfrac{1}{2}t^2)\,dt = \left[t + \tfrac{1}{6}t^3\right]_0^6 = 6 + 36 = 42$$

14. Let C be the curve of intersection. The projection of C onto the xy-plane is the ellipse $4x^2 + y^2 = 4$ or $x^2 + y^2/4 = 1$, $z = 0$. Then we can write $x = \cos t$, $y = 2\sin t$, $0 \leq t \leq 2\pi$. Since C also lies on the plane $x + y + z = 2$, we have

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$z = 2 - x - y = 2 - \cos t - 2\sin t$. Then parametric equations for C are $x = \cos t$, $y = 2\sin t$, $z = 2 - \cos t - 2\sin t$, $0 \le t \le 2\pi$, and the corresponding vector equation is $\mathbf{r}(t) = \langle \cos t, 2\sin t, 2 - \cos t - 2\sin t\rangle$. Differentiating gives $\mathbf{r}'(t) = \langle -\sin t, 2\cos t, \sin t - 2\cos t\rangle \quad \Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{(-\sin t)^2 + (2\cos t)^2 + (\sin t - 2\cos t)^2} = \sqrt{2\sin^2 t + 8\cos^2 t - 4\sin t\,\cos t}$. The length of C is

$L = \int_0^{2\pi} |\mathbf{r}'(t)|\,dt = \int_0^{2\pi} \sqrt{2\sin^2 t + 8\cos^2 t - 4\sin t\,\cos t}\,dt \approx 13.5191$.

15. (a) $\mathbf{r}(t) = (5-t)\,\mathbf{i} + (4t-3)\,\mathbf{j} + 3t\,\mathbf{k} \quad \Rightarrow \quad \mathbf{r}'(t) = -\mathbf{i} + 4\,\mathbf{j} + 3\,\mathbf{k}$ and $\frac{ds}{dt} = |\mathbf{r}'(t)| = \sqrt{1+16+9} = \sqrt{26}$. The point $P\,(4,1,3)$ corresponds to $t = 1$, so the arc length function from P is

$s(t) = \int_1^t |\mathbf{r}'(u)|\,du = \int_1^t \sqrt{26}\,du = \sqrt{26}\,u\big|_1^t = \sqrt{26}\,(t-1)$. Since $s = \sqrt{26}\,(t-1)$, we have $t = \dfrac{s}{\sqrt{26}} + 1$.

Substituting for t in the original equation, the reparametrization of the curve with respect to arc length is

$$\begin{aligned}\mathbf{r}(t(s)) &= \left[5 - \left(\frac{s}{\sqrt{26}} + 1\right)\right]\mathbf{i} + \left[4\left(\frac{s}{\sqrt{26}} + 1\right) - 3\right]\mathbf{j} + 3\left(\frac{s}{\sqrt{26}} + 1\right)\mathbf{k}\\ &= \left(4 - \frac{s}{\sqrt{26}}\right)\mathbf{i} + \left(\frac{4s}{\sqrt{26}} + 1\right)\mathbf{j} + \left(\frac{3s}{\sqrt{26}} + 3\right)\mathbf{k}\end{aligned}$$

(b) The point 4 units along the curve from P has position vector

$\mathbf{r}(t(4)) = \left(4 - \dfrac{4}{\sqrt{26}}\right)\mathbf{i} + \left(\dfrac{4(4)}{\sqrt{26}} + 1\right)\mathbf{j} + \left(\dfrac{3(4)}{\sqrt{26}} + 3\right)\mathbf{k}$, so the point is $\left(4 - \dfrac{4}{\sqrt{26}}, \dfrac{16}{\sqrt{26}} + 1, \dfrac{12}{\sqrt{26}} + 3\right)$.

16. (a) $\mathbf{r}(t) = e^t \sin t\,\mathbf{i} + e^t \cos t\,\mathbf{j} + \sqrt{2}\,e^t\,\mathbf{k} \quad \Rightarrow \quad \mathbf{r}'(t) = e^t\,(\cos t + \sin t)\,\mathbf{i} + e^t\,(\cos t - \sin t)\,\mathbf{j} + \sqrt{2}\,e^t\,\mathbf{k}$ and

$$\begin{aligned}\frac{ds}{dt} &= |\mathbf{r}'(t)| = \sqrt{e^{2t}(\cos t + \sin t)^2 + e^{2t}(\cos t - \sin t)^2 + 2e^{2t}}\\ &= \sqrt{e^{2t}\left[2(\cos^2 t + \sin^2 t) + 2\cos t\sin t - 2\cos t\sin t + 2\right]} = \sqrt{4e^{2t}} = 2e^t\end{aligned}$$

The point $P\left(0, 1, \sqrt{2}\right)$ corresponds to $t = 0$, so the arc length function from P is

$s(t) = \int_0^t |\mathbf{r}'(u)|\,du = \int_0^t 2e^u\,du = 2e^u\big|_0^t = 2(e^t - 1)$. Since $s = 2(e^t - 1)$, we have $e^t = \dfrac{s}{2} + 1 \quad \Leftrightarrow$

$t = \ln\left(\frac{1}{2}s + 1\right)$. Substituting for t in the original equation, the reparametrization of the curve with respect to arc length is

$\mathbf{r}(t(s)) = \left(\frac{1}{2}s + 1\right)\sin\!\left(\ln\left(\frac{1}{2}s + 1\right)\right)\mathbf{i} + \left(\frac{1}{2}s + 1\right)\cos\!\left(\ln\left(\frac{1}{2}s + 1\right)\right)\mathbf{j} + \left(\frac{\sqrt{2}}{2}s + \sqrt{2}\right)\mathbf{k}$.

(b) The point 4 units along the curve from P has position vector

$\mathbf{r}(t(4)) = \left(\frac{1}{2}(4) + 1\right)\sin\!\left(\ln\left(\frac{1}{2}(4) + 1\right)\right)\mathbf{i} + \left(\frac{1}{2}(4) + 1\right)\cos\!\left(\ln\left(\frac{1}{2}(4) + 1\right)\right)\mathbf{j} + \left(\frac{\sqrt{2}}{2}(4) + \sqrt{2}\right)\mathbf{k}$, so the point is $\left(3\sin(\ln 3), 3\cos(\ln 3), 3\sqrt{2}\right)$.

17. Here $\mathbf{r}(t) = \langle 3\sin t, 4t, 3\cos t\rangle$, so $\mathbf{r}'(t) = \langle 3\cos t, 4, -3\sin t\rangle$ and $|\mathbf{r}'(t)| = \sqrt{9\cos^2 t + 16 + 9\sin^2 t} = \sqrt{25} = 5$. The point $(0,0,3)$ corresponds to $t = 0$, so the arc length function beginning at $(0,0,3)$ and measuring in the positive direction is given by $s(t) = \int_0^t |\mathbf{r}'(u)|\,du = \int_0^t 5\,du = 5t$. $\quad s(t) = 5 \quad \Rightarrow \quad 5t = 5 \quad \Rightarrow \quad t = 1$, thus your location after moving 5 units along the curve is $(3\sin 1, 4, 3\cos 1)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $\mathbf{r}(t) = \left(\dfrac{2}{t^2+1} - 1\right)\mathbf{i} + \dfrac{2t}{t^2+1}\,\mathbf{j} \quad \Rightarrow \quad \mathbf{r}'(t) = \dfrac{-4t}{(t^2+1)^2}\,\mathbf{i} + \dfrac{-2t^2+2}{(t^2+1)^2}\,\mathbf{j},$

$$\frac{ds}{dt} = |\mathbf{r}'(t)| = \sqrt{\left[\frac{-4t}{(t^2+1)^2}\right]^2 + \left[\frac{-2t^2+2}{(t^2+1)^2}\right]^2} = \sqrt{\frac{4t^4+8t^2+4}{(t^2+1)^4}} = \sqrt{\frac{4(t^2+1)^2}{(t^2+1)^4}} = \sqrt{\frac{4}{(t^2+1)^2}} = \frac{2}{t^2+1}.$$

Since the initial point $(1, 0)$ corresponds to $t = 0$, the arc length function is

$s(t) = \displaystyle\int_0^t |\mathbf{r}'(u)|\,du = \int_0^t \frac{2}{u^2+1}\,du = 2\arctan t$. Then $\arctan t = \frac{1}{2}s \quad \Rightarrow \quad t = \tan\frac{1}{2}s$. Substituting, we have

$$\mathbf{r}(t(s)) = \left[\frac{2}{\tan^2\left(\frac{1}{2}s\right)+1} - 1\right]\mathbf{i} + \frac{2\tan\left(\frac{1}{2}s\right)}{\tan^2\left(\frac{1}{2}s\right)+1}\,\mathbf{j} = \frac{1-\tan^2\left(\frac{1}{2}s\right)}{1+\tan^2\left(\frac{1}{2}s\right)}\,\mathbf{i} + \frac{2\tan\left(\frac{1}{2}s\right)}{\sec^2\left(\frac{1}{2}s\right)}\,\mathbf{j}$$

$$= \frac{1-\tan^2\left(\frac{1}{2}s\right)}{\sec^2\left(\frac{1}{2}s\right)}\,\mathbf{i} + 2\tan\left(\tfrac{1}{2}s\right)\cos^2\left(\tfrac{1}{2}s\right)\mathbf{j} = \left[\cos^2\left(\tfrac{1}{2}s\right) - \sin^2\left(\tfrac{1}{2}s\right)\right]\mathbf{i} + 2\sin\left(\tfrac{1}{2}s\right)\cos\left(\tfrac{1}{2}s\right)\mathbf{j} = \cos s\,\mathbf{i} + \sin s\,\mathbf{j}$$

With this parametrization, we recognize the function as representing the unit circle. Note here that the curve approaches, but does not include, the point $(-1, 0)$, since $\cos s = -1$ for $s = \pi + 2k\pi$ (k an integer) but then $t = \tan\left(\frac{1}{2}s\right)$ is undefined.

19. (a) $\mathbf{r}(t) = \left\langle t^2, \sin t - t\cos t, \cos t + t\sin t\right\rangle \quad \Rightarrow$

$\mathbf{r}'(t) = \langle 2t, \cos t + t\sin t - \cos t,\ -\sin t + t\cos t + \sin t\rangle = \langle 2t, t\sin t, t\cos t\rangle \quad \Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{4t^2 + t^2\sin^2 t + t^2\cos^2 t} = \sqrt{4t^2 + t^2(\cos^2 t + \sin^2 t)} = \sqrt{5t^2} = \sqrt{5}\,t$ [since $t > 0$]. Then

$\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{5}\,t}\langle 2t, t\sin t, t\cos t\rangle = \dfrac{1}{\sqrt{5}}\langle 2, \sin t, \cos t\rangle. \quad \mathbf{T}'(t) = \dfrac{1}{\sqrt{5}}\langle 0, \cos t, -\sin t\rangle \quad \Rightarrow$

$|\mathbf{T}'(t)| = \frac{1}{\sqrt{5}}\sqrt{0 + \cos^2 t + \sin^2 t} = \dfrac{1}{\sqrt{5}}$. Thus, $\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{1/\sqrt{5}}{1/\sqrt{5}}\langle 0, \cos t, -\sin t\rangle = \langle 0, \cos t, -\sin t\rangle$.

(b) By Formula 9, the curvature is $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{1/\sqrt{5}}{\sqrt{5}\,t} = \dfrac{1}{5t}$.

20. (a) $\mathbf{r}(t) = \langle 5\sin t, t, 5\cos t\rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \langle 5\cos t, 1, -5\sin t\rangle \quad \Rightarrow \quad |\mathbf{r}'(t)| = \sqrt{25\cos^2 t + 1 + 25\sin^2 t} = \sqrt{26}$.

Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{26}}\langle 5\cos t, 1, -5\sin t\rangle. \quad \mathbf{T}'(t) = \dfrac{1}{\sqrt{26}}\langle -5\sin t, 0, -5\cos t\rangle \quad \Rightarrow$

$|\mathbf{T}'(t)| = \dfrac{1}{\sqrt{26}}\sqrt{25\sin^2 t + 0^2 + 25\cos^2 t} = \dfrac{5}{\sqrt{26}}$. Thus,

$\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{\sqrt{26}}{5}\cdot\dfrac{1}{\sqrt{26}}\langle -5\sin t, 0, -5\cos t\rangle = \langle -\sin t, 0, -\cos t\rangle$.

(b) By Formula 9, the curvature is $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{5/\sqrt{26}}{\sqrt{26}} = \dfrac{5}{26}$.

21. (a) $\mathbf{r}(t) = \left\langle t, t^2, 4\right\rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \langle 1, 2t, 0\rangle \quad \Rightarrow \quad |\mathbf{r}'(t)| = \sqrt{1 + 4t^2 + 0}$. Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{1+4t^2}}\langle 1, 2t, 0\rangle$.

$$\mathbf{T}'(t) = \frac{1}{\sqrt{1+4t^2}}\langle 0, 2, 0\rangle - \frac{4t}{(1+4t^2)^{3/2}}\langle 1, 2t, 0\rangle \qquad \text{[by Formula 3 of Theorem 13.2.3]}$$

$$= \frac{1}{(1+4t^2)^{3/2}}\left[(1+4t^2)\langle 0, 2, 0\rangle - 4t\langle 1, 2t, 0\rangle\right] = \frac{1}{(1+4t^2)^{3/2}}\langle -4t, 2, 0\rangle$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$|\mathbf{T}'(t)| = \dfrac{1}{(1+4t^2)^{3/2}}\sqrt{16t^2+4} = \dfrac{2}{1+4t^2}$. Thus,

$$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \frac{1+4t^2}{2}\cdot\frac{1}{(1+4t^2)^{3/2}}\langle -4t, 2, 0\rangle = \frac{1}{\sqrt{1+4t^2}}\langle -2t, 1, 0\rangle.$$

(b) By Formula 9, the curvature is $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{2/(1+4t^2)}{\sqrt{1+4t^2}} = \dfrac{2}{(1+4t^2)^{3/2}}$.

22. (a) $\mathbf{r}(t) = \left\langle t, t, \frac{1}{2}t^2\right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1, 1, t\rangle \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{1+1+t^2} = \sqrt{2+t^2}$.

Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{2+t^2}}\langle 1, 1, t\rangle$.

$$\mathbf{T}'(t) = \frac{1}{\sqrt{2+t^2}}\langle 0,0,1\rangle - \frac{t}{(2+t^2)^{3/2}}\langle 1,1,t\rangle \qquad \text{[by Formula 3 of Theorem 13.2.3]}$$

$$= \frac{1}{(2+t^2)^{3/2}}\left[(2+t^2)\langle 0,0,1\rangle - t\langle 1,1,t\rangle\right] = \frac{1}{(2+t^2)^{3/2}}\langle -t,-t,2\rangle$$

$|\mathbf{T}'(t)| = \dfrac{1}{(2+t^2)^{3/2}}\sqrt{4+2t^2} = \dfrac{\sqrt{2}}{2+t^2}$. Thus,

$$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \frac{2+t^2}{\sqrt{2}}\cdot\frac{1}{(2+t^2)^{3/2}}\langle -t,-t,2\rangle = \frac{1}{\sqrt{4+2t^2}}\langle -t,-t,2\rangle.$$

(b) By Formula 9, the curvature is $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{\sqrt{2}/(2+t^2)}{\sqrt{2+t^2}} = \dfrac{\sqrt{2}}{(2+t^2)^{3/2}}$.

23. (a) $\mathbf{r}(t) = \left\langle t, \frac{1}{2}t^2, t^2\right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1, t, 2t\rangle \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{1+t^2+4t^2} = \sqrt{1+5t^2}$.

Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{1+5t^2}}\langle 1, t, 2t\rangle$.

$$\mathbf{T}'(t) = \frac{-5t}{(1+5t^2)^{3/2}}\langle 1,t,2t\rangle + \frac{1}{\sqrt{1+5t^2}}\langle 0,1,2\rangle \qquad \text{[by Formula 3 of Theorem 13.2.3]}$$

$$= \frac{1}{(1+5t^2)^{3/2}}\left(\left\langle -5t,-5t^2,-10t^2\right\rangle + \left\langle 0,1+5t^2,2+10t^2\right\rangle\right) = \frac{1}{(1+5t^2)^{3/2}}\langle -5t,1,2\rangle$$

$$|\mathbf{T}'(t)| = \frac{1}{(1+5t^2)^{3/2}}\sqrt{25t^2+1+4} = \frac{1}{(1+5t^2)^{3/2}}\sqrt{25t^2+5} = \frac{\sqrt{5}\sqrt{5t^2+1}}{(1+5t^2)^{3/2}} = \frac{\sqrt{5}}{1+5t^2}.$$

Thus, $\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{1+5t^2}{\sqrt{5}}\cdot\dfrac{1}{(1+5t^2)^{3/2}}\langle -5t,1,2\rangle = \dfrac{1}{\sqrt{5+25t^2}}\langle -5t,1,2\rangle$.

(b) By Formula 9, the curvature is $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{\sqrt{5}/(1+5t^2)}{\sqrt{1+5t^2}} = \dfrac{\sqrt{5}}{(1+5t^2)^{3/2}}$.

24. (a) $\mathbf{r}(t) = \left\langle \sqrt{2}\,t, e^t, e^{-t}\right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \left\langle \sqrt{2}, e^t, -e^{-t}\right\rangle \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{2+e^{2t}+e^{-2t}} = \sqrt{(e^t+e^{-t})^2} = e^t+e^{-t}$.

Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{e^t+e^{-t}}\left\langle \sqrt{2}, e^t, -e^{-t}\right\rangle = \dfrac{1}{e^{2t}+1}\left\langle \sqrt{2}\,e^t, e^{2t}, -1\right\rangle \quad \left[\text{after multiplying by } \dfrac{e^t}{e^t}\right]$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\mathbf{T}'(t) = \frac{1}{e^{2t}+1}\left\langle \sqrt{2}\,e^t, 2e^{2t}, 0\right\rangle - \frac{2e^{2t}}{(e^{2t}+1)^2}\left\langle \sqrt{2}\,e^t, e^{2t}, -1\right\rangle \qquad \text{[by Formula 3 of Theorem 13.2.3]}$$

$$= \frac{1}{(e^{2t}+1)^2}\left[(e^{2t}+1)\left\langle \sqrt{2}\,e^t, 2e^{2t}, 0\right\rangle - 2e^{2t}\left\langle \sqrt{2}\,e^t, e^{2t}, -1\right\rangle\right] = \frac{1}{(e^{2t}+1)^2}\left\langle \sqrt{2}\,e^t\left(1-e^{2t}\right), 2e^{2t}, 2e^{2t}\right\rangle$$

$$|\mathbf{T}'(t)| = \frac{1}{(e^{2t}+1)^2}\sqrt{2e^{2t}(1-2e^{2t}+e^{4t})+4e^{4t}+4e^{4t}} = \frac{1}{(e^{2t}+1)^2}\sqrt{2e^{2t}(1+2e^{2t}+e^{4t})}$$

$$= \frac{1}{(e^{2t}+1)^2}\sqrt{2e^{2t}\left(1+e^{2t}\right)^2} = \frac{\sqrt{2}\,e^t(1+e^{2t})}{(e^{2t}+1)^2} = \frac{\sqrt{2}\,e^t}{e^{2t}+1}$$

Thus,
$$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \frac{e^{2t}+1}{\sqrt{2}\,e^t}\,\frac{1}{(e^{2t}+1)^2}\left\langle \sqrt{2}\,e^t(1-e^{2t}), 2e^{2t}, 2e^{2t}\right\rangle$$

$$= \frac{1}{\sqrt{2}\,e^t(e^{2t}+1)}\left\langle \sqrt{2}\,e^t(1-e^{2t}), 2e^{2t}, 2e^{2t}\right\rangle = \frac{1}{e^{2t}+1}\left\langle 1-e^{2t}, \sqrt{2}\,e^t, \sqrt{2}\,e^t\right\rangle$$

(b) By Formula 9, the curvature is

$$\kappa(t) = \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{\sqrt{2}\,e^t}{e^{2t}+1}\cdot\frac{1}{e^t+e^{-t}} = \frac{\sqrt{2}\,e^t}{e^{3t}+2e^t+e^{-t}} = \frac{\sqrt{2}\,e^{2t}}{e^{4t}+2e^{2t}+1} = \frac{\sqrt{2}\,e^{2t}}{(e^{2t}+1)^2}.$$

25. $\mathbf{r}(t) = t^3\,\mathbf{j} + t^2\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = 3t^2\,\mathbf{j} + 2t\,\mathbf{k}$, $\mathbf{r}''(t) = 6t\,\mathbf{j} + 2\,\mathbf{k}$, $|\mathbf{r}'(t)| = \sqrt{0^2+(3t^2)^2+(2t)^2} = \sqrt{9t^4+4t^2}$,

$\mathbf{r}'(t)\times\mathbf{r}''(t) = -6t^2\,\mathbf{i}$, $|\mathbf{r}'(t)\times\mathbf{r}''(t)| = 6t^2$. Then $\kappa(t) = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{6t^2}{\left(\sqrt{9t^4+4t^2}\right)^3} = \dfrac{6t^2}{(9t^4+4t^2)^{3/2}}$.

26. $\mathbf{r}(t) = t\,\mathbf{i} + t^2\,\mathbf{j} + e^t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = \mathbf{i} + 2t\,\mathbf{j} + e^t\,\mathbf{k}$, $\mathbf{r}''(t) = 2\,\mathbf{j} + e^t\,\mathbf{k}$,

$|\mathbf{r}'(t)| = \sqrt{1^2+(2t)^2+(e^t)^2} = \sqrt{1+4t^2+e^{2t}}$, $\mathbf{r}'(t)\times\mathbf{r}''(t) = (2t-2)e^t\,\mathbf{i} - e^t\,\mathbf{j} + 2\,\mathbf{k}$,

$|\mathbf{r}'(t)\times\mathbf{r}''(t)| = \sqrt{[(2t-2)e^t]^2+(-e^t)^2+2^2} = \sqrt{(2t-2)^2e^{2t}+e^{2t}+4} = \sqrt{(4t^2-8t+5)e^{2t}+4}$.

Then $\kappa(t) = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{\sqrt{(4t^2-8t+5)e^{2t}+4}}{\left(\sqrt{1+4t^2+e^{2t}}\right)^3} = \dfrac{\sqrt{(4t^2-8t+5)e^{2t}+4}}{(1+4t^2+e^{2t})^{3/2}}$.

27. $\mathbf{r}(t) = \sqrt{6}\,t^2\,\mathbf{i} + 2t\,\mathbf{j} + 2t^3\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = 2\sqrt{6}\,t\,\mathbf{i} + 2\,\mathbf{j} + 6t^2\,\mathbf{k}$, $\mathbf{r}''(t) = 2\sqrt{6}\,\mathbf{i} + 12t\,\mathbf{k}$,

$|\mathbf{r}'(t)| = \sqrt{24t^2+4+36t^4} = \sqrt{4(9t^4+6t^2+1)} = \sqrt{4(3t^2+1)^2} = 2(3t^2+1)$,

$\mathbf{r}'(t)\times\mathbf{r}''(t) = 24t\,\mathbf{i} - 12\sqrt{6}\,t^2\,\mathbf{j} - 4\sqrt{6}\,\mathbf{k}$,

$|\mathbf{r}'(t)\times\mathbf{r}''(t)| = \sqrt{576t^2+864t^4+96} = \sqrt{96(9t^4+6t^2+1)} = \sqrt{96(3t^2+1)^2} = 4\sqrt{6}\,(3t^2+1)$.

Then $\kappa(t) = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{4\sqrt{6}\,(3t^2+1)}{8(3t^2+1)^3} = \dfrac{\sqrt{6}}{2(3t^2+1)^2}$.

28. $\mathbf{r}(t) = \left\langle t^2, \ln t, t\ln t\right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 2t, 1/t, 1+\ln t\rangle$, $\mathbf{r}''(t) = \left\langle 2, -1/t^2, 1/t\right\rangle$. The point $(1,0,0)$ corresponds to $t=1$, and $\mathbf{r}'(1) = \langle 2,1,1\rangle$, $|\mathbf{r}'(1)| = \sqrt{2^2+1^2+1^2} = \sqrt{6}$, $\mathbf{r}''(1) = \langle 2,-1,1\rangle$, $\mathbf{r}'(1)\times\mathbf{r}''(1) = \langle 2,0,-4\rangle$,

$|\mathbf{r}'(1)\times\mathbf{r}''(1)| = \sqrt{2^2+0^2+(-4)^2} = \sqrt{20} = 2\sqrt{5}$. Then $\kappa(1) = \dfrac{|\mathbf{r}'(1)\times\mathbf{r}''(1)|}{|\mathbf{r}'(1)|^3} = \dfrac{2\sqrt{5}}{\left(\sqrt{6}\right)^3} = \dfrac{2\sqrt{5}}{6\sqrt{6}}$ or $\dfrac{\sqrt{30}}{18}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \langle 1, 2t, 3t^2 \rangle$. The point $(1, 1, 1)$ corresponds to $t = 1$, and $\mathbf{r}'(1) = \langle 1, 2, 3 \rangle \quad \Rightarrow$ $|\mathbf{r}'(1)| = \sqrt{1+4+9} = \sqrt{14}$. $\mathbf{r}''(t) = \langle 0, 2, 6t \rangle \quad \Rightarrow \quad \mathbf{r}''(1) = \langle 0, 2, 6 \rangle$. $\mathbf{r}'(1) \times \mathbf{r}''(1) = \langle 6, -6, 2 \rangle$, so $|\mathbf{r}'(1) \times \mathbf{r}''(1)| = \sqrt{36+36+4} = \sqrt{76}$. Then $\kappa(1) = \dfrac{|\mathbf{r}'(1) \times \mathbf{r}''(1)|}{|\mathbf{r}'(1)|^3} = \dfrac{\sqrt{76}}{\sqrt{14}^3} = \dfrac{1}{7}\sqrt{\dfrac{19}{14}}$.

30.

Note that we get the complete curve for $0 \le t < 2\pi$.

$\mathbf{r}(t) = \langle \cos t, \sin t, \sin 5t \rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \langle -\sin t, \cos t, 5\cos 5t \rangle$,

$\mathbf{r}''(t) = \langle -\cos t, -\sin t, -25\sin 5t \rangle$. The point $(1, 0, 0)$ corresponds to $t = 0$, and $\mathbf{r}'(0) = \langle 0, 1, 5 \rangle \quad \Rightarrow$

$|\mathbf{r}'(0)| = \sqrt{0^2 + 1^2 + 5^2} = \sqrt{26}$, $\quad \mathbf{r}''(0) = \langle -1, 0, 0 \rangle$,

$\mathbf{r}'(0) \times \mathbf{r}''(0) = \langle 0, -5, 1 \rangle \quad \Rightarrow$

$|\mathbf{r}'(0) \times \mathbf{r}''(0)| = \sqrt{0^2 + (-5)^2 + 1^2} = \sqrt{26}$. The curvature at the point $(1, 0, 0)$ is $\kappa(0) = \dfrac{|\mathbf{r}'(0) \times \mathbf{r}''(0)|}{|\mathbf{r}'(0)|^3} = \dfrac{\sqrt{26}}{\left(\sqrt{26}\right)^3} = \dfrac{1}{26}$.

31. $f(x) = x^4 \quad \Rightarrow \quad f'(x) = 4x^3 \quad \Rightarrow \quad f''(x) = 12x^2$. By Formula 11, the curvature is

$$\kappa(x) = \frac{|f''(x)|}{[1 + (f'(x))^2]^{3/2}} = \frac{\left|12x^2\right|}{[1 + (4x^3)^2]^{3/2}} = \frac{12x^2}{(1 + 16x^6)^{3/2}}.$$

32. $f(x) = \tan x \quad \Rightarrow \quad f'(x) = \sec^2 x \quad \Rightarrow \quad f''(x) = 2\sec x \cdot \sec x \, \tan x = 2\sec^2 x \, \tan x$. By Formula 11, the curvature is

$$\kappa(x) = \frac{|f''(x)|}{[1 + (f'(x))^2]^{3/2}} = \frac{\left|2\sec^2 x \, \tan x\right|}{[1 + (\sec^2 x)^2]^{3/2}} = \frac{2\sec^2 x \, |\tan x|}{(1 + \sec^4 x)^{3/2}}.$$

33. $f(x) = xe^x \quad \Rightarrow \quad f'(x) = xe^x + e^x \quad \Rightarrow \quad f''(x) = xe^x + 2e^x$. By Formula 11, the curvature is

$$\kappa(x) = \frac{|f''(x)|}{[1 + (f'(x))^2]^{3/2}} = \frac{|xe^x + 2e^x|}{[1 + (xe^x + e^x)^2]^{3/2}} = \frac{e^x \, |x + 2|}{[1 + (xe^x + e^x)^2]^{3/2}}.$$

34. $y = \ln x \quad \Rightarrow \quad y' = \dfrac{1}{x} \quad \Rightarrow \quad y'' = -\dfrac{1}{x^2}$. By Formula 11, the curvature is

$$\kappa(x) = \frac{|y''(x)|}{\left[1 + (y'(x))^2\right]^{3/2}} = \left|\frac{-1}{x^2}\right| \frac{1}{(1 + 1/x^2)^{3/2}} = \frac{1}{x^2} \frac{(x^2)^{3/2}}{(x^2+1)^{3/2}} = \frac{|x|}{(x^2+1)^{3/2}} = \frac{x}{(x^2+1)^{3/2}} \quad [\text{since } x > 0].$$

To find the maximum curvature, we first find the critical numbers of $\kappa(x)$:

$$\kappa'(x) = \frac{(x^2+1)^{3/2} - x\left(\frac{3}{2}\right)(x^2+1)^{1/2}(2x)}{\left[(x^2+1)^{3/2}\right]^2} = \frac{(x^2+1)^{1/2}[(x^2+1) - 3x^2]}{(x^2+1)^3} = \frac{1 - 2x^2}{(x^2+1)^{5/2}};$$

$\kappa'(x) = 0 \quad \Rightarrow \quad 1 - 2x^2 = 0$, so the only critical number in the domain is $x = \frac{1}{\sqrt{2}}$. Since $\kappa'(x) > 0$ for $0 < x < \frac{1}{\sqrt{2}}$ and $\kappa'(x) < 0$ for $x > \frac{1}{\sqrt{2}}$, $\kappa(x)$ attains its maximum at $x = \frac{1}{\sqrt{2}}$. Thus, the maximum curvature occurs at $\left(\frac{1}{\sqrt{2}}, \ln \frac{1}{\sqrt{2}}\right)$.

Since $\displaystyle\lim_{x \to \infty} \frac{x}{(x^2+1)^{3/2}} = 0$, $\kappa(x)$ approaches 0 as $x \to \infty$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. Since $y = y' = y'' = e^x$, the curvature is $\kappa(x) = \dfrac{|y''(x)|}{[1+(y'(x))^2]^{3/2}} = \dfrac{e^x}{(1+e^{2x})^{3/2}} = e^x(1+e^{2x})^{-3/2}$.

To find the maximum curvature, we first find the critical numbers of $\kappa(x)$:

$$\kappa'(x) = e^x(1+e^{2x})^{-3/2} + e^x\left(-\tfrac{3}{2}\right)(1+e^{2x})^{-5/2}(2e^{2x}) = e^x\frac{1+e^{2x}-3e^{2x}}{(1+e^{2x})^{5/2}} = e^x\frac{1-2e^{2x}}{(1+e^{2x})^{5/2}}.$$

$\kappa'(x) = 0$ when $1-2e^{2x} = 0$, so $e^{2x} = \frac{1}{2}$ or $x = -\frac{1}{2}\ln 2$. And since $1-2e^{2x} > 0$ for $x < -\frac{1}{2}\ln 2$ and $1-2e^{2x} < 0$ for $x > -\frac{1}{2}\ln 2$, the maximum curvature is attained at the point $\left(-\frac{1}{2}\ln 2, e^{(-\ln 2)/2}\right) = \left(-\frac{1}{2}\ln 2, \frac{1}{\sqrt{2}}\right)$.

Since $\lim\limits_{x\to\infty} e^x(1+e^{2x})^{-3/2} = 0$, $\kappa(x)$ approaches 0 as $x \to \infty$.

36. We can take the parabola as having its vertex at the origin and opening upward, so the equation is $f(x) = ax^2, a > 0$. Then by Formula 11, $\kappa(x) = \dfrac{|f''(x)|}{[1+(f'(x))^2]^{3/2}} = \dfrac{|2a|}{[1+(2ax)^2]^{3/2}} = \dfrac{2a}{(1+4a^2x^2)^{3/2}}$, thus $\kappa(0) = 2a$. We want $\kappa(0) = 4$, so $a = 2$ and the equation is $y = 2x^2$.

37. (a) C appears to be changing direction more quickly at P than Q, so we would expect the curvature to be greater at P.

(b) First we sketch approximate osculating circles at P and Q. Using the axes scale as a guide, we measure the radius of the osculating circle at P to be approximately 0.8 units, thus $\rho = \dfrac{1}{\kappa} \Rightarrow \kappa = \dfrac{1}{\rho} \approx \dfrac{1}{0.8} \approx 1.3$. Similarly, we estimate the radius of the osculating circle at Q to be 1.4 units, so $\kappa = \dfrac{1}{\rho} \approx \dfrac{1}{1.4} \approx 0.7$.

38. $y = x^4 - 2x^2 \Rightarrow y' = 4x^3 - 4x \Rightarrow y'' = 12x^2 - 4$.

$\kappa(x) = \dfrac{|y''|}{\left[1+(y')^2\right]^{3/2}} = \dfrac{\left|12x^2-4\right|}{\left[1+(4x^3-4x)^2\right]^{3/2}}$. The graph of the curvature here is what we would expect. The graph of $y = x^4 - 2x^2$ appears to be bending most sharply at the origin and near $x = \pm 1$.

39. $y = x^{-2} \Rightarrow y' = -2x^{-3} \Rightarrow y'' = 6x^{-4}$.

$\kappa(x) = \dfrac{|y''|}{\left[1+(y')^2\right]^{3/2}} = \dfrac{\left|6x^{-4}\right|}{\left[1+(-2x^{-3})^2\right]^{3/2}} = \dfrac{6}{x^4\left(1+4x^{-6}\right)^{3/2}}$.

The appearance of the two humps in this graph is perhaps a little surprising, but it is explained by the fact that $y = x^{-2}$ increases asymptotically at the origin from both directions, and so its graph has very little bend there. [Note that $\kappa(0)$ is undefined.]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $\mathbf{r}(t) = \langle t - \sin t, 1 - \cos t, 4\cos(t/2)\rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \langle 1 - \cos t, \sin t, -2\sin(t/2)\rangle \quad \Rightarrow$

$\mathbf{r}''(t) = \langle \sin t, \cos t, -\cos(t/2)\rangle$. Using a CAS, $\mathbf{r}'(t) \times \mathbf{r}''(t) = \left\langle -2\sin^3(t/2), -\sin(t/2)\sin t, \cos t - 1\right\rangle$,

$|\mathbf{r}'(t) \times \mathbf{r}''(t)| = \sqrt{3 - 4\cos t + \cos 2t}$ or $2\sqrt{2}\sin^2(t/2)$, and $|\mathbf{r}'(t)| = 2\sqrt{1 - \cos t}$ or $2\sqrt{2}\,|\sin(t/2)|$.

(To compute cross products in Maple, use the `VectorCalculus` or `LinearAlgebra` package and the `CrossProduct(a,b)` command. Here loading the `RealDomain` package will give simpler results. In Mathematica, use `Cross[a,b]`.) Then $\kappa(t) = \dfrac{|\mathbf{r}'(t) \times \mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{\sqrt{3 - 4\cos t + \cos 2t}}{8\,(1 - \cos t)^{3/2}}$ or $\dfrac{1}{4\sqrt{2 - 2\cos t}}$ or $\dfrac{1}{8\,|\sin(t/2)|}$. We plot the space curve and its curvature function for $0 \le t \le 8\pi$ below.

The asymptotes in the graph of $\kappa(t)$ correspond to the sharp cusps we see in the graph of $\mathbf{r}(t)$. The space curve bends most sharply as it approaches these cusps (mostly in the x-direction) and bends most gradually between these, near its intersections with the xy-plane, where $t = \pi + 2n\pi$ (n an integer). (The bending we see in the z-direction on the curve near these points is deceiving; most of the curvature occurs in the x-direction.) The curvature graph has local minima at these values of t.

41. $\mathbf{r}(t) = \left\langle te^t, e^{-t}, \sqrt{2}t\right\rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \left\langle (t+1)e^t, -e^{-t}, \sqrt{2}\right\rangle \quad \Rightarrow \quad \mathbf{r}''(t) = \left\langle (t+2)e^t, e^{-t}, 0\right\rangle$.

Then $\mathbf{r}'(t) \times \mathbf{r}''(t) = \left\langle -\sqrt{2}e^{-t}, \sqrt{2}(t+2)e^t, 2t+3\right\rangle$, $\quad |\mathbf{r}'(t) \times \mathbf{r}''(t)| = \sqrt{2e^{-2t} + 2(t+2)^2e^{2t} + (2t+3)^2}$,

$|\mathbf{r}'(t)| = \sqrt{(t+1)^2e^{2t} + e^{-2t} + 2}$, and $\kappa(t) = \dfrac{|\mathbf{r}'(t) \times \mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{\sqrt{2e^{-2t} + 2(t+2)^2e^{2t} + (2t+3)^2}}{\left[(t+1)^2e^{2t} + e^{-2t} + 2\right]^{3/2}}$.

We plot the space curve and its curvature function for $-5 \le t \le 5$ below.

From the graph of $\kappa(t)$ we see that curvature is maximized for $t = 0$, so the curve bends most sharply at the point $(0, 1, 0)$. The curve bends more gradually as we move away from this point, becoming almost linear. This is reflected in the curvature graph, where $\kappa(t)$ becomes nearly 0 as $|t|$ increases.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. Notice that the curve a is highest for the same x-values at which curve b is turning more sharply, and a is 0 or near 0 where b is nearly straight. So, a must be the graph of $y = \kappa(x)$, and b is the graph of $y = f(x)$.

43. Notice that the curve b has two inflection points at which the graph appears almost straight. We would expect the curvature to be 0 or nearly 0 at these values, but the curve a isn't near 0 there. Thus, a must be the graph of $y = f(x)$ rather than the graph of curvature, and b is the graph of $y = \kappa(x)$.

44. (a) The complete curve for $\mathbf{r}(t) = \langle \sin 3t, \sin 2t, \sin 3t \rangle$ is given by $0 \le t \le 2\pi$. Curvature appears to have a local (or absolute) maximum at 6 points. (Look at points where the curve appears to turn more sharply.)

(b) Using a CAS, we find (after simplifying) $\kappa(t) = \dfrac{3\sqrt{2}\,\sqrt{(5\sin t + \sin 5t)^2}}{(9\cos 6t + 2\cos 4t + 11)^{3/2}}$. (To compute cross products in Maple, use the `VectorCalculus` or `LinearAlgebra` package and the `CrossProduct(a,b)` command; in Mathematica, use `Cross[a,b]`.) The graph shows 6 local (or absolute) maximum points for $0 \le t \le 2\pi$, as observed in part (a).

45. $\mathbf{r}(t) = \left\langle t - \frac{3}{2}\sin t, 1 - \frac{3}{2}\cos t, t \right\rangle$. Using a CAS, we find (after simplifying) $\kappa(t) = \dfrac{6\sqrt{4\cos^2 t - 12\cos t + 13}}{(17 - 12\cos t)^{3/2}}$.

(To compute cross products in Maple, use the `VectorCalculus` or `LinearAlgebra` package and the `CrossProduct(a,b)` command; in Mathematica, use `Cross[a,b]`.) Curvature is largest at integer multiples of 2π.

46. Here $\mathbf{r}(t) = \langle f(t), g(t) \rangle$, $\mathbf{r}'(t) = \langle f'(t), g'(t) \rangle$, $\mathbf{r}''(t) = \langle f''(t), g''(t) \rangle$,

$|\mathbf{r}'(t)|^3 = \left[\sqrt{(f'(t))^2 + (g'(t))^2}\right]^3 = [(f'(t))^2 + (g'(t))^2]^{3/2} = (\dot{x}^2 + \dot{y}^2)^{3/2}$, and

$|\mathbf{r}'(t) \times \mathbf{r}''(t)| = |\langle 0, 0, f'(t)\,g''(t) - f''(t)\,g'(t) \rangle| = \left[(\dot{x}\ddot{y} - \ddot{x}\dot{y})^2\right]^{1/2} = |\dot{x}\ddot{y} - \dot{y}\ddot{x}|$. Thus, by Theorem 10,

$\kappa(t) = \dfrac{|\dot{x}\ddot{y} - \dot{y}\ddot{x}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}}$.

47. $x = t^2 \;\Rightarrow\; \dot{x} = 2t \;\Rightarrow\; \ddot{x} = 2$, $\;y = t^3 \;\Rightarrow\; \dot{y} = 3t^2 \;\Rightarrow\; \ddot{y} = 6t$.

Then $\kappa(t) = \dfrac{|\dot{x}\ddot{y} - \dot{y}\ddot{x}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}} = \dfrac{\left|(2t)(6t) - (3t^2)(2)\right|}{[(2t)^2 + (3t^2)^2]^{3/2}} = \dfrac{\left|12t^2 - 6t^2\right|}{(4t^2 + 9t^4)^{3/2}} = \dfrac{6t^2}{(4t^2 + 9t^4)^{3/2}}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. $x = a\cos\omega t \quad\Rightarrow\quad \dot{x} = -a\omega\sin\omega t \quad\Rightarrow\quad \ddot{x} = -a\omega^2\cos\omega t,$

$y = b\sin\omega t \quad\Rightarrow\quad \dot{y} = b\omega\cos\omega t \quad\Rightarrow\quad \ddot{y} = -b\omega^2\sin\omega t$. Then

$$\kappa(t) = \frac{|\dot{x}\ddot{y} - \dot{y}\ddot{x}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}} = \frac{\left|(-a\omega\sin\omega t)(-b\omega^2\sin\omega t) - (b\omega\cos\omega t)(-a\omega^2\cos\omega t)\right|}{[(-a\omega\sin\omega t)^2 + (b\omega\cos\omega t)^2]^{3/2}}$$

$$= \frac{\left|ab\omega^3\sin^2\omega t + ab\omega^3\cos^2\omega t\right|}{(a^2\omega^2\sin^2\omega t + b^2\omega^2\cos^2\omega t)^{3/2}} = \frac{\left|ab\omega^3\right|}{(a^2\omega^2\sin^2\omega t + b^2\omega^2\cos^2\omega t)^{3/2}}$$

49. $x = e^t\cos t \quad\Rightarrow\quad \dot{x} = e^t(\cos t - \sin t) \quad\Rightarrow\quad \ddot{x} = e^t(-\sin t - \cos t) + e^t(\cos t - \sin t) = -2e^t\sin t,$

$y = e^t\sin t \quad\Rightarrow\quad \dot{y} = e^t(\cos t + \sin t) \quad\Rightarrow\quad \ddot{y} = e^t(-\sin t + \cos t) + e^t(\cos t + \sin t) = 2e^t\cos t$. Then

$$\kappa(t) = \frac{|\dot{x}\ddot{y} - \dot{y}\ddot{x}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}} = \frac{\left|e^t(\cos t - \sin t)(2e^t\cos t) - e^t(\cos t + \sin t)(-2e^t\sin t)\right|}{\left([e^t(\cos t - \sin t)]^2 + [e^t(\cos t + \sin t)]^2\right)^{3/2}}$$

$$= \frac{\left|2e^{2t}(\cos^2 t - \sin t\cos t + \sin t\cos t + \sin^2 t)\right|}{\left[e^{2t}(\cos^2 t - 2\cos t\sin t + \sin^2 t + \cos^2 t + 2\cos t\sin t + \sin^2 t)\right]^{3/2}} = \frac{\left|2e^{2t}(1)\right|}{\left[e^{2t}(1+1)\right]^{3/2}} = \frac{2e^{2t}}{e^{3t}(2)^{3/2}} = \frac{1}{\sqrt{2}\,e^t}$$

50. $f(x) = e^{cx}, \quad f'(x) = ce^{cx}, \quad f''(x) = c^2e^{cx}$. Using Formula 11 we have

$\kappa(x) = \dfrac{|f''(x)|}{[1 + (f'(x))^2]^{3/2}} = \dfrac{\left|c^2e^{cx}\right|}{[1 + (ce^{cx})^2]^{3/2}} = \dfrac{c^2e^{cx}}{(1 + c^2e^{2cx})^{3/2}}$ so the curvature at $x = 0$ is

$\kappa(0) = \dfrac{c^2}{(1+c^2)^{3/2}}$. To determine the maximum value for $\kappa(0)$, let $f(c) = \dfrac{c^2}{(1+c^2)^{3/2}}$. Then

$f'(c) = \dfrac{2c\cdot(1+c^2)^{3/2} - c^2\cdot\frac{3}{2}(1+c^2)^{1/2}(2c)}{[(1+c^2)^{3/2}]^2} = \dfrac{(1+c^2)^{1/2}\left[2c(1+c^2) - 3c^3\right]}{(1+c^2)^3} = \dfrac{\left(2c - c^3\right)}{(1+c^2)^{5/2}}$. We have a critical number when $2c - c^3 = 0 \quad\Rightarrow\quad c(2 - c^2) = 0 \quad\Rightarrow\quad c = 0$ or $c = \pm\sqrt{2}$. $f'(c)$ is positive for $c < -\sqrt{2}$, $0 < c < \sqrt{2}$ and negative elsewhere, so f achieves its maximum value when $c = \sqrt{2}$ or $-\sqrt{2}$. In either case, $\kappa(0) = \dfrac{2}{3^{3/2}}$, so the members of the family with the largest value of $\kappa(0)$ are $f(x) = e^{\sqrt{2}x}$ and $f(x) = e^{-\sqrt{2}x}$.

51. $\mathbf{r}(t) = \left\langle t^2, \frac{2}{3}t^3, t\right\rangle$. $\left(1, \frac{2}{3}, 1\right)$ corresponds to $t = 1$.

$\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{\langle 2t, 2t^2, 1\rangle}{\sqrt{4t^2 + 4t^4 + 1}} = \dfrac{\langle 2t, 2t^2, 1\rangle}{2t^2 + 1}$, so $\mathbf{T}(1) = \left\langle \frac{2}{3}, \frac{2}{3}, \frac{1}{3}\right\rangle$.

$\mathbf{T}'(t) = -4t(2t^2+1)^{-2}\langle 2t, 2t^2, 1\rangle + (2t^2+1)^{-1}\langle 2, 4t, 0\rangle$ [by Formula 3 of Theorem 13.2.3]

$= (2t^2+1)^{-2}\left\langle -8t^2 + 4t^2 + 2, -8t^3 + 8t^3 + 4t, -4t\right\rangle = 2(2t^2+1)^{-2}\left\langle 1 - 2t^2, 2t, -2t\right\rangle$

$\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{2(2t^2+1)^{-2}\left\langle 1 - 2t^2, 2t, -2t\right\rangle}{2(2t^2+1)^{-2}\sqrt{(1-2t^2)^2 + (2t)^2 + (-2t)^2}} = \dfrac{\left\langle 1 - 2t^2, 2t, -2t\right\rangle}{\sqrt{1 - 4t^2 + 4t^4 + 8t^2}} = \dfrac{\left\langle 1 - 2t^2, 2t, -2t\right\rangle}{1 + 2t^2}$

$\mathbf{N}(1) = \left\langle -\frac{1}{3}, \frac{2}{3}, -\frac{2}{3}\right\rangle$ and $\mathbf{B}(1) = \mathbf{T}(1)\times\mathbf{N}(1) = \left\langle -\frac{4}{9} - \frac{2}{9}, -\left(-\frac{4}{9} + \frac{1}{9}\right), \frac{4}{9} + \frac{2}{9}\right\rangle = \left\langle -\frac{2}{3}, \frac{1}{3}, \frac{2}{3}\right\rangle$.

52. $(1, 0, 0)$ corresponds to $t = 0$. $\mathbf{r}(t) = \langle \cos t, \sin t, \ln\cos t\rangle$, and in Exercise 6 we found that $\mathbf{r}'(t) = \langle -\sin t, \cos t, -\tan t\rangle$ and $|\mathbf{r}'(t)| = |\sec t|$. Here we can assume $-\frac{\pi}{2} < t < \frac{\pi}{2}$ and then $\sec t > 0 \quad\Rightarrow\quad |\mathbf{r}'(t)| = \sec t$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{\langle -\sin t, \cos t, -\tan t\rangle}{\sec t} = \left\langle -\sin t\cos t, \cos^2 t, -\sin t\right\rangle$ and $\mathbf{T}(0) = \langle 0, 1, 0\rangle$.

$\mathbf{T}'(t) = \langle -[(\sin t)(-\sin t) + (\cos t)(\cos t)], 2(\cos t)(-\sin t), -\cos t\rangle = \left\langle \sin^2 t - \cos^2 t, -2\sin t\cos t, -\cos t\right\rangle$, so

$\mathbf{N}(0) = \dfrac{\mathbf{T}'(0)}{|\mathbf{T}'(0)|} = \dfrac{\langle -1, 0, -1\rangle}{\sqrt{1+0+1}} = \dfrac{1}{\sqrt{2}}\langle -1, 0, -1\rangle = \left\langle -\frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}}\right\rangle$.

Finally, $\mathbf{B}(0) = \mathbf{T}(0) \times \mathbf{N}(0) = \langle 0, 1, 0\rangle \times \left\langle -\frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}}\right\rangle = \left\langle -\frac{1}{\sqrt{2}}, 0, \frac{1}{\sqrt{2}}\right\rangle$.

53. $\mathbf{r}(t) = \langle \sin 2t, -\cos 2t, 4t\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 2\cos 2t, 2\sin 2t, 4\rangle$. The point $(0, 1, 2\pi)$ corresponds to $t = \pi/2$, and the normal plane there has normal vector $\mathbf{r}'(\pi/2) = \langle -2, 0, 4\rangle$. An equation for the normal plane is $-2(x-0) + 0(y-1) + 4(z-2\pi) = 0$ or $-2x + 4z = 8\pi$ or $x - 2z = -4\pi$.

$\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{\langle 2\cos 2t, 2\sin 2t, 4\rangle}{\sqrt{4\cos^2 2t + 4\sin^2 2t + 16}} = \dfrac{1}{2\sqrt{5}}\langle 2\cos 2t, 2\sin 2t, 4\rangle = \dfrac{1}{\sqrt{5}}\langle \cos 2t, \sin 2t, 2\rangle \;\Rightarrow$

$\mathbf{T}'(t) = \frac{1}{\sqrt{5}}\langle -2\sin 2t, 2\cos 2t, 0\rangle \;\Rightarrow\; |\mathbf{T}'(t)| = \frac{1}{\sqrt{5}}\sqrt{4\sin^2 2t + 4\cos^2 2t} = \frac{2}{\sqrt{5}}$, and

$\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \langle -\sin 2t, \cos 2t, 0\rangle$. Then $\mathbf{T}(\pi/2) = \frac{1}{\sqrt{5}}\langle -1, 0, 2\rangle$, $\mathbf{N}(\pi/2) = \langle 0, -1, 0\rangle$, and

$\mathbf{B}(\pi/2) = \mathbf{T}(\pi/2) \times \mathbf{N}(\pi/2) = \frac{1}{\sqrt{5}}\langle 2, 0, 1\rangle$. Since $\mathbf{B}(\pi/2)$ is normal to the osculating plane, so is $\langle 2, 0, 1\rangle$, and an equation of the plane is $2(x-0) + 0(y-1) + 1(z-2\pi) = 0$ or $2x + z = 2\pi$.

54. $\mathbf{r}(t) = \left\langle \ln t, 2t, t^2\right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1/t, 2, 2t\rangle$. The point $(0, 2, 1)$ corresponds to $t = 1$, and the normal plane there has normal vector $\mathbf{r}'(1) = \langle 1, 2, 2\rangle$. An equation for the normal plane is $1(x-0) + 2(y-2) + 2(z-1) = 0$ or $x + 2y + 2z = 6$.

$|\mathbf{r}'(t)| = \sqrt{1/t^2 + 4 + 4t^2} = \sqrt{[(1/t) + 2t]^2} = (1/t) + 2t$ [since $t > 0$] and then

$\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{\langle 1/t, 2, 2t\rangle}{(1/t) + 2t} = \dfrac{1}{1+2t^2}\left\langle 1, 2t, 2t^2\right\rangle$ $\left[\text{after multiplying by } \dfrac{t}{t}\right]$. By Formula 3 of Theorem 13.2.3,

$$\begin{aligned}\mathbf{T}'(t) &= -\frac{4t}{(1+2t^2)^2}\left\langle 1, 2t, 2t^2\right\rangle + \frac{1}{1+2t^2}\langle 0, 2, 4t\rangle \\ &= \frac{1}{(1+2t^2)^2}\left\langle -4t, -8t^2 + 2(1+2t^2), -8t^3 + 4t(1+2t^2)\right\rangle = \frac{1}{(1+2t^2)^2}\left\langle -4t, 2-4t^2, 4t\right\rangle\end{aligned}$$

Then

$$\begin{aligned}|\mathbf{T}'(t)| &= \frac{1}{(1+2t^2)^2}\sqrt{16t^2 + (2-4t^2)^2 + 16t^2} = \frac{1}{(1+2t^2)^2}\sqrt{16t^2 + 4 + 16t^4} \\ &= \frac{1}{(1+2t^2)^2}\cdot 2\sqrt{(1+2t^2)^2} = \frac{2}{1+2t^2}\end{aligned}$$

and $\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{1}{2(1+2t^2)}\left\langle -4t, 2-4t^2, 4t\right\rangle = \dfrac{1}{1+2t^2}\left\langle -2t, 1-2t^2, 2t\right\rangle$.

Thus $\mathbf{T}(1) = \frac{1}{3}\langle 1, 2, 2\rangle$, $\mathbf{N}(1) = \frac{1}{3}\langle -2, -1, 2\rangle$, and $\mathbf{B}(1) = \mathbf{T}(1) \times \mathbf{N}(1) = \frac{1}{9}\langle 6, -6, 3\rangle$ is normal to the osculating plane.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

We can take the parallel vector $\langle 2, -2, 1\rangle$ as a normal vector for the plane, so an equation is

$2(x-0) - 2(y-2) + 1(z-1) = 0$ or $2x - 2y + z = -3$.

Note: Since $\mathbf{r}'(1)$ is parallel to $\mathbf{T}(1)$ and $\mathbf{T}'(1)$ is parallel to $\mathbf{N}(1)$, we could have taken $\mathbf{r}'(1) \times \mathbf{T}'(1)$ as a normal vector for the plane.

55. The ellipse is given by the parametric equations $x = 2\cos t,\ y = 3\sin t$, so using the result from Exercise 46,

$$\kappa(t) = \frac{|\dot{x}\ddot{y} - \ddot{x}\dot{y}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}} = \frac{|(-2\sin t)(-3\sin t) - (3\cos t)(-2\cos t)|}{(4\sin^2 t + 9\cos^2 t)^{3/2}} = \frac{6}{(4\sin^2 t + 9\cos^2 t)^{3/2}}.$$

At $(2, 0)$, $t = 0$. Now $\kappa(0) = \frac{6}{27} = \frac{2}{9}$, so the radius of the osculating circle is $1/\kappa(0) = \frac{9}{2}$ and its center is $\left(-\frac{5}{2}, 0\right)$. Its equation is therefore $\left(x + \frac{5}{2}\right)^2 + y^2 = \frac{81}{4}$. At $(0, 3)$, $t = \frac{\pi}{2}$, and $\kappa\left(\frac{\pi}{2}\right) = \frac{6}{8} = \frac{3}{4}$. So the radius of the osculating circle is $\frac{4}{3}$ and its center is $\left(0, \frac{5}{3}\right)$. Hence its equation is $x^2 + \left(y - \frac{5}{3}\right)^2 = \frac{16}{9}$.

56. $y = \frac{1}{2}x^2 \quad \Rightarrow \quad y' = x$ and $y'' = 1$, so Formula 11 gives $\kappa(x) = \dfrac{1}{(1+x^2)^{3/2}}$. So the curvature at $(0, 0)$ is $\kappa(0) = 1$ and the osculating circle has radius 1 and center $(0, 1)$, and hence equation $x^2 + (y-1)^2 = 1$. The curvature at $\left(1, \frac{1}{2}\right)$ is $\kappa(1) = \dfrac{1}{(1+1^2)^{3/2}} = \dfrac{1}{2\sqrt{2}}$. The tangent line to the parabola at $\left(1, \frac{1}{2}\right)$ has slope 1, so the normal line has slope -1. Thus the center of the osculating circle lies in the direction of the unit vector $\left\langle -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right\rangle$. The circle has radius $2\sqrt{2}$, so its center has position vector $\left\langle 1, \frac{1}{2}\right\rangle + 2\sqrt{2}\left\langle -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right\rangle = \left\langle -1, \frac{5}{2}\right\rangle$. So the equation of the circle is $(x+1)^2 + \left(y - \frac{5}{2}\right)^2 = 8$.

57. Here $\mathbf{r}(t) = \langle t^3, 3t, t^4\rangle$, and $\mathbf{r}'(t) = \langle 3t^2, 3, 4t^3\rangle$ is normal to the normal plane for any t. The given plane has normal vector $\langle 6, 6, -8\rangle$, and the planes are parallel when their normal vectors are parallel. Thus we need to find a value for t where $\langle 3t^2, 3, 4t^3\rangle = k\,\langle 6, 6, -8\rangle$ for some $k \neq 0$. From the y-component we see that $k = \frac{1}{2}$, and $\langle 3t^2, 3, 4t^3\rangle = \frac{1}{2}\,\langle 6, 6, -8\rangle = \langle 3, 3, -4\rangle$ for $t = -1$. Thus the planes are parallel at the point $(-1, -3, 1)$.

58. To find the osculating plane, we first calculate the unit tangent and normal vectors.

In Maple, we use the `VectorCalculus` package and set `r:=<t^3,3*t,t^4>;`. After differentiating, the `Normalize` command converts the tangent vector to the unit tangent vector: `T:=Normalize(diff(r,t));`. After simplifying, we find that $\mathbf{T}(t) = \dfrac{\langle 3t^2, 3, 4t^3\rangle}{\sqrt{16t^6 + 9t^4 + 9}}$. We use a similar procedure to compute the unit normal vector,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

`N:=Normalize(diff(T,t));`. After simplifying, we have $\mathbf{N}(t) = \dfrac{\left\langle -t(8t^6-9), -3t^3(3+8t^2), 6t^2(t^4+3)\right\rangle}{\sqrt{t^2(4t^6+36t^2+9)(16t^6+9t^4+9)}}$. Then we use the command `B:=CrossProduct(T,N);`. After simplification, we find that $\mathbf{B}(t) = \dfrac{\left\langle 6t^2, -2t^4, -3t\right\rangle}{\sqrt{t^2(4t^6+36t^2+9)}}$.

In Mathematica, we define the vector function `r={t^3,3*t,t^4}` and use the command `Dt` to differentiate. We find $\mathbf{T}(t)$ by dividing the result by its magnitude, computed using the `Norm` command. (You may wish to include the option `Element[t,Reals]` to obtain simpler expressions.) $\mathbf{N}(t)$ is found similarly, and we use `Cross[T,N]` to find $\mathbf{B}(t)$.

Now $\mathbf{B}(t)$ is parallel to $\left\langle 6t^2, -2t^4, -3t\right\rangle$, so if $\mathbf{B}(t)$ is parallel to $\langle 1,1,1\rangle$ for some $t \neq 0$ [since $\mathbf{B}(0) = \mathbf{0}$], then $\left\langle 6t^2, -2t^4, -3t\right\rangle = k\,\langle 1,1,1\rangle$ for some value of k. But then $6t^2 = -2t^4 = -3t$ which has no solution for $t \neq 0$. So there is no such osculating plane.

59. First we parametrize the curve of intersection. We can choose $y = t$; then $x = y^2 = t^2$ and $z = x^2 = t^4$, and the curve is given by $\mathbf{r}(t) = \left\langle t^2, t, t^4\right\rangle$. $\mathbf{r}'(t) = \left\langle 2t, 1, 4t^3\right\rangle$ and the point $(1,1,1)$ corresponds to $t = 1$, so $\mathbf{r}'(1) = \langle 2,1,4\rangle$ is a normal vector for the normal plane. Thus an equation of the normal plane is

$2(x-1)+1(y-1)+4(z-1) = 0$ or $2x+y+4z = 7$. $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{4t^2+1+16t^6}}\left\langle 2t, 1, 4t^3\right\rangle$ and

$\mathbf{T}'(t) = -\frac{1}{2}(4t^2+1+16t^6)^{-3/2}(8t+96t^5)\left\langle 2t,1,4t^3\right\rangle + (4t^2+1+16t^6)^{-1/2}\left\langle 2,0,12t^2\right\rangle$. A normal vector for the osculating plane is $\mathbf{B}(1) = \mathbf{T}(1)\times\mathbf{N}(1)$, but $\mathbf{r}'(1) = \langle 2,1,4\rangle$ is parallel to $\mathbf{T}(1)$ and

$\mathbf{T}'(1) = -\frac{1}{2}(21)^{-3/2}(104)\langle 2,1,4\rangle + (21)^{-1/2}\langle 2,0,12\rangle = \frac{2}{21\sqrt{21}}\langle -31,-26,22\rangle$ is parallel to $\mathbf{N}(1)$ as is $\langle -31,-26,22\rangle$, so $\langle 2,1,4\rangle \times \langle -31,-26,22\rangle = \langle 126,-168,-21\rangle$ is normal to the osculating plane. Thus an equation for the osculating plane is $126(x-1)-168(y-1)-21(z-1) = 0$ or $6x-8y-z = -3$.

60. $\mathbf{r}(t) = \left\langle t+2, 1-t, \frac{1}{2}t^2\right\rangle \Rightarrow \mathbf{r}'(t) = \langle 1,-1,t\rangle$, $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{2+t^2}}\langle 1,-1,t\rangle$,

$$\begin{aligned}\mathbf{T}'(t) &= -\tfrac{1}{2}(2+t^2)^{-3/2}(2t)\langle 1,-1,t\rangle + (2+t^2)^{-1/2}\langle 0,0,1\rangle \\ &= -(2+t^2)^{-3/2}\left[t\langle 1,-1,t\rangle - (2+t^2)\langle 0,0,1\rangle\right] = \frac{-1}{(2+t^2)^{3/2}}\langle t,-t,-2\rangle\end{aligned}$$

A normal vector for the osculating plane is $\mathbf{B}(t) = \mathbf{T}(t)\times\mathbf{N}(t)$, but $\mathbf{r}'(t) = \langle 1,-1,t\rangle$ is parallel to $\mathbf{T}(t)$ and $\langle t,-t,-2\rangle$ is parallel to $\mathbf{T}'(t)$ and hence parallel to $\mathbf{N}(t)$, so $\langle 1,-1,t\rangle \times \langle t,-t,-2\rangle = \left\langle t^2+2, t^2+2, 0\right\rangle$ is normal to the osculating plane for any t. All such vectors are parallel to $\langle 1,1,0\rangle$, so at any point $\left(t+2, 1-t, \frac{1}{2}t^2\right)$ on the curve, an equation for the osculating plane is $1[x-(t+2)]+1[y-(1-t)]+0\left(z-\frac{1}{2}t^2\right) = 0$ or $x+y = 3$. Because the osculating plane at every point on the curve is the same, we can conclude that the curve itself lies in that same plane. In fact, we can easily verify that the parametric equations of the curve satisfy $x+y = 3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. $\mathbf{r}(t) = \langle e^t \cos t, e^t \sin t, e^t \rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \langle e^t(\cos t - \sin t), e^t(\cos t + \sin t), e^t \rangle$ so

$$|\mathbf{r}'(t)| = \sqrt{e^{2t}(\cos t - \sin t)^2 + e^{2t}(\cos t + \sin t)^2 + e^{2t}}$$
$$= \sqrt{e^{2t}\left[2(\cos^2 t + \sin^2 t) - 2\cos t \sin t + 2\cos t \sin t + 1\right]} = \sqrt{3e^{2t}} = \sqrt{3}\, e^t$$

and $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{3}\, e^t} \langle e^t(\cos t - \sin t), e^t(\cos t + \sin t), e^t \rangle = \dfrac{1}{\sqrt{3}} \langle \cos t - \sin t, \cos t + \sin t, 1 \rangle$. The vector $\mathbf{k} = \langle 0, 0, 1 \rangle$ is parallel to the z-axis, so for any t, the angle α between $\mathbf{T}(t)$ and the z-axis is given by

$\cos \alpha = \dfrac{\mathbf{T}(t) \cdot \mathbf{k}}{|\mathbf{T}(t)|\ |\mathbf{k}|} = \dfrac{\frac{1}{\sqrt{3}} \langle \cos t - \sin t, \cos t + \sin t, 1 \rangle \cdot \langle 0, 0, 1 \rangle}{\frac{1}{\sqrt{3}} \sqrt{(\cos t - \sin t)^2 + (\cos t + \sin t)^2 + 1}\ \sqrt{1}} = \dfrac{1}{\sqrt{2(\cos^2 t + \sin^2 t) + 1}} = \dfrac{1}{\sqrt{3}}$. Thus the angle is constant; specifically, $\alpha = \cos^{-1}(1/\sqrt{3}) \approx 54.7\,^\circ$.

$\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{(1/\sqrt{3}\,) \langle -\sin t - \cos t, -\sin t + \cos t, 0 \rangle}{(1/\sqrt{3}\,)\sqrt{2\left(\sin^2 t + \cos^2 t\right)}} = \dfrac{1}{\sqrt{2}} \langle -\sin t - \cos t, -\sin t + \cos t, 0 \rangle$, and the angle β made with the z-axis is given by $\cos \beta = \dfrac{\mathbf{N}(t) \cdot \mathbf{k}}{|\mathbf{N}(t)|\ |\mathbf{k}|} = 0$, so $\beta = 90\,^\circ$.

$\mathbf{B}(t) = \mathbf{T}(t) \times \mathbf{N}(t) = \frac{1}{\sqrt{6}} \langle \sin t - \cos t, -\sin t - \cos t, 2 \rangle$ and the angle γ made with the z-axis is given by

$\cos \gamma = \dfrac{\mathbf{B}(t) \cdot \mathbf{k}}{|\mathbf{B}(t)|\ |\mathbf{k}|} = \dfrac{\frac{1}{\sqrt{6}} \langle \sin t - \cos t, -\sin t - \cos t, 2 \rangle \cdot \langle 0, 0, 1 \rangle}{\frac{1}{\sqrt{6}} \sqrt{(\sin t - \cos t)^2 + (-\sin t - \cos t)^2 + 4}\ \sqrt{1}} = \dfrac{2}{\sqrt{6}}$ or equivalently $\dfrac{\sqrt{6}}{3}$. Again the angle is constant; specifically, $\gamma = \cos^{-1}(2/\sqrt{6}) \approx 35.3\,^\circ$.

62. If vectors $\mathbf{T}$ and $\mathbf{B}$ lie in the rectifying plane then $\mathbf{N}$ is a normal vector for the plane, as it is orthogonal to both $\mathbf{T}$ and $\mathbf{B}$. The point $\left(\sqrt{2}/2, \sqrt{2}/2, 1\right)$ corresponds to $t = \pi/4$, so we can take $\mathbf{T}'(\pi/4)$ as a normal vector for the plane [since it is parallel to $\mathbf{N}(\pi/4)$]. $\mathbf{r}(t) = \sin t\, \mathbf{i} + \cos t\, \mathbf{j} + \tan t\, \mathbf{k} \quad \Rightarrow \quad \mathbf{r}'(t) = \cos t\, \mathbf{i} - \sin t\, \mathbf{j} + \sec^2 t\, \mathbf{k}$ and

$|\mathbf{r}'(t)| = \sqrt{\cos^2 t + \sin^2 t + \sec^4 t} = \sqrt{1 + \sec^4 t}$. Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{1 + \sec^4 t}} \left(\cos t\, \mathbf{i} - \sin t\, \mathbf{j} + \sec^2 t\, \mathbf{k}\right)$.

By Formula 3 of Theorem 13.2.3,

$\mathbf{T}'(t) = -\dfrac{2\sec^4 t \tan t}{(1 + \sec^4 t)^{3/2}} \left(\cos t\, \mathbf{i} - \sin t\, \mathbf{j} + \sec^2 t\, \mathbf{k}\right) + \dfrac{1}{\sqrt{1 + \sec^4 t}} \left(-\sin t\, \mathbf{i} - \cos t\, \mathbf{j} + 2\sec^2 t \tan t\, \mathbf{k}\right)$ and

$$\mathbf{T}'(\pi/4) = -\frac{2(\sqrt{2}\,)^4(1)}{[1 + (\sqrt{2}\,)^4]^{3/2}} \left(\frac{\sqrt{2}}{2}\, \mathbf{i} - \frac{\sqrt{2}}{2}\, \mathbf{j} + (\sqrt{2}\,)^2\, \mathbf{k}\right) + \frac{1}{\sqrt{1 + (\sqrt{2}\,)^4}} \left(-\frac{\sqrt{2}}{2}\, \mathbf{i} - \frac{\sqrt{2}}{2}\, \mathbf{j} + 2(\sqrt{2}\,)^2(1)\, \mathbf{k}\right)$$
$$= -\frac{8}{5\sqrt{5}} \left(\frac{\sqrt{2}}{2}\, \mathbf{i} - \frac{\sqrt{2}}{2}\, \mathbf{j} + 2\, \mathbf{k}\right) + \frac{1}{\sqrt{5}} \left(-\frac{\sqrt{2}}{2}\, \mathbf{i} - \frac{\sqrt{2}}{2}\, \mathbf{j} + 4\, \mathbf{k}\right) = -\frac{13\sqrt{2}}{10\sqrt{5}}\, \mathbf{i} + \frac{3\sqrt{2}}{10\sqrt{5}}\, \mathbf{j} + \frac{4}{5\sqrt{5}}\, \mathbf{k}$$

We can take the parallel vector $-13\sqrt{2}\, \mathbf{i} + 3\sqrt{2}\, \mathbf{j} + 8\, \mathbf{k}$ as a normal for the plane, so an equation for the plane is

$-13\sqrt{2}\left(x - \frac{\sqrt{2}}{2}\right) + 3\sqrt{2}\left(y - \frac{\sqrt{2}}{2}\right) + 8\,(z - 1) = 0$ or $-13\sqrt{2}\, x + 3\sqrt{2}\, y + 8z = -2$ or $13x - 3y - 4\sqrt{2}\, z = \sqrt{2}$.

63. $\kappa = \left|\dfrac{d\mathbf{T}}{ds}\right| = \left|\dfrac{d\mathbf{T}/dt}{ds/dt}\right| = \dfrac{|d\mathbf{T}/dt|}{ds/dt}$ and $\mathbf{N} = \dfrac{d\mathbf{T}/dt}{|d\mathbf{T}/dt|}$, so $\kappa\mathbf{N} = \dfrac{\left|\dfrac{d\mathbf{T}}{dt}\right| \dfrac{d\mathbf{T}}{dt}}{\left|\dfrac{d\mathbf{T}}{dt}\right| \dfrac{ds}{dt}} = \dfrac{d\mathbf{T}/dt}{ds/dt} = \dfrac{d\mathbf{T}}{ds}$ by the Chain Rule.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. For a plane curve, $\mathbf{T} = |\mathbf{T}|\cos\phi\,\mathbf{i} + |\mathbf{T}|\sin\phi\,\mathbf{j} = \cos\phi\,\mathbf{i} + \sin\phi\,\mathbf{j}$. Then

$\dfrac{d\mathbf{T}}{ds} = \left(\dfrac{d\mathbf{T}}{d\phi}\right)\left(\dfrac{d\phi}{ds}\right) = (-\sin\phi\,\mathbf{i} + \cos\phi\,\mathbf{j})\left(\dfrac{d\phi}{ds}\right)$ and $\left|\dfrac{d\mathbf{T}}{ds}\right| = |-\sin\phi\,\mathbf{i} + \cos\phi\,\mathbf{j}|\left|\dfrac{d\phi}{ds}\right| = \left|\dfrac{d\phi}{ds}\right|$. Hence for a plane curve, the curvature is $\kappa = |d\phi/ds|$.

65. (a) $|\mathbf{B}| = 1 \;\Rightarrow\; \mathbf{B}\cdot\mathbf{B} = 1 \;\Rightarrow\; \dfrac{d}{ds}(\mathbf{B}\cdot\mathbf{B}) = 0 \;\Rightarrow\; 2\dfrac{d\mathbf{B}}{ds}\cdot\mathbf{B} = 0 \;\Rightarrow\; \dfrac{d\mathbf{B}}{ds} \perp \mathbf{B}$.

This shows that $d\mathbf{B}/ds$ is perpendicular to $\mathbf{B}$. Alternatively, note that this is a direct result of Theorem 13.2.4.

(b) $\mathbf{B} = \mathbf{T}\times\mathbf{N} \;\Rightarrow$

$$\begin{aligned}
\frac{d\mathbf{B}}{ds} &= \frac{d}{ds}(\mathbf{T}\times\mathbf{N}) = \frac{d}{dt}(\mathbf{T}\times\mathbf{N})\frac{1}{ds/dt} = \frac{d}{dt}(\mathbf{T}\times\mathbf{N})\frac{1}{|\mathbf{r}'(t)|} && \text{[by Formula 7]}\\
&= [(\mathbf{T}'\times\mathbf{N}) + (\mathbf{T}\times\mathbf{N}')]\frac{1}{|\mathbf{r}'(t)|} && \text{[by Formula 5 of Theorem 13.2.3]}\\
&= \left[\left(\mathbf{T}'\times\frac{\mathbf{T}'}{|\mathbf{T}'|}\right) + (\mathbf{T}\times\mathbf{N}')\right]\frac{1}{|\mathbf{r}'(t)|}\\
&= [\mathbf{0} + (\mathbf{T}\times\mathbf{N}')]\frac{1}{|\mathbf{r}'(t)|} = \frac{\mathbf{T}\times\mathbf{N}'}{|\mathbf{r}'(t)|} && [\mathbf{a}\times c\mathbf{a} = \mathbf{0}]\\
\Rightarrow\ \frac{d\mathbf{B}}{ds} &\perp \mathbf{T} && \text{[by Theorem 12.4.8]}
\end{aligned}$$

(c) $\mathbf{B} = \mathbf{T}\times\mathbf{N} \;\Rightarrow\; \mathbf{B}\perp\mathbf{T}$ and $\mathbf{B}\perp\mathbf{N}$. Since $\mathbf{T}\perp\mathbf{N}$, $\mathbf{B}$, $\mathbf{T}$, and $\mathbf{N}$ form an orthogonal set of vectors in the three-dimensional space $\mathbb{R}^3$. From parts (a) and (b), $d\mathbf{B}/ds$ is perpendicular to both $\mathbf{B}$ and $\mathbf{T}$, so $d\mathbf{B}/ds$ is parallel to $\mathbf{N}$.

66. We need to find $\mathbf{T}(t)$, $\mathbf{N}(t)$, and $\mathbf{B}(t)$ in terms of t. $\mathbf{r}(t) = \langle \sin t, 3t, \cos t\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle \cos t, 3, -\sin t\rangle \;\Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{\cos^2 t + 3^2 + \sin^2 t} = \sqrt{10}$. Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{10}}\langle \cos t, 3, -\sin t\rangle \;\Rightarrow$

$\mathbf{T}'(t) = \dfrac{1}{\sqrt{10}}\langle -\sin t, 0, -\cos t\rangle$, and $|\mathbf{T}'(t)| = \dfrac{1}{\sqrt{10}}\sqrt{\sin^2 t + 0^2 + \cos^2 t} = \dfrac{1}{\sqrt{10}}$.

$\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{\sqrt{10}}{1}\cdot\dfrac{1}{\sqrt{10}}\langle -\sin t, 0, -\cos t\rangle = \langle -\sin t, 0, -\cos t\rangle$.

Then

$$\begin{aligned}
\mathbf{B}(t) = \mathbf{T}(t)\times\mathbf{N}(t) &= \frac{1}{\sqrt{10}}\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k}\\ \cos t & 3 & -\sin t\\ -\sin t & 0 & -\cos t\end{vmatrix}\\
&= \frac{1}{\sqrt{10}}\left[-3\cos t\,\mathbf{i} - (-\cos^2 t - \sin^2 t)\,\mathbf{j} + 3\sin t\,\mathbf{k}\right] = \frac{1}{\sqrt{10}}\langle -3\cos t, 1, 3\sin t\rangle
\end{aligned}$$

$$\mathbf{B}'(t) = \frac{1}{\sqrt{10}}\langle 3\sin t, 0, 3\cos t\rangle \;\Rightarrow\; \mathbf{B}'(\pi/2) = \frac{1}{\sqrt{10}}\langle 3, 0, 0\rangle \quad\text{and}\quad \mathbf{N}(\pi/2) = \langle -1, 0, 0\rangle$$

Thus, the torsion is $\tau(t) = -\dfrac{\mathbf{B}'(\pi/2)\cdot\mathbf{N}(\pi/2)}{|\mathbf{r}'(\pi/2)|} = -\dfrac{-3/\sqrt{10}}{\sqrt{10}} = \dfrac{3}{10}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. We need to find $\mathbf{T}(t)$, $\mathbf{N}(t)$, and $\mathbf{B}(t)$ in terms of t. $\mathbf{r}(t) = \left\langle \frac{1}{2}t^2, 2t, t \right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle t, 2, 1\rangle \;\Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{t^2 + 2^2 + 1^2} = \sqrt{t^2+5}$. Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{t^2+5}}\langle t, 2, 1\rangle \;\Rightarrow$

$$\mathbf{T}'(t) = \frac{1}{\sqrt{t^2+5}}\langle 1,0,0\rangle - \frac{t}{(t^2+5)^{3/2}}\langle t,2,1\rangle = \frac{1}{(t^2+5)^{3/2}}\left[(t^2+5)\langle 1,0,0\rangle - t\langle t,2,1\rangle\right]$$
$$= \frac{1}{(t^2+5)^{3/2}}\langle 5, -2t, -t\rangle, \text{ and}$$

$$|\mathbf{T}'(t)| = \frac{1}{(t^2+5)^{3/2}}\sqrt{25+5t^2} = \frac{\sqrt{5}}{t^2+5}.$$

$$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \frac{t^2+5}{\sqrt{5}}\cdot\frac{1}{(t^2+5)^{3/2}}\langle 5,-2t,-t\rangle = \frac{1}{\sqrt{5}\sqrt{t^2+5}}\langle 5,-2t,-t\rangle.$$

Then $\mathbf{B}(t) = \mathbf{T}(t)\times\mathbf{N}(t) = \dfrac{1}{\sqrt{t^2+5}}\cdot\dfrac{1}{\sqrt{5}\sqrt{t^2+5}}\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ t & 2 & 1 \\ 5 & -2t & -t \end{vmatrix}$

$$= \frac{1}{\sqrt{5}(t^2+5)}\left[(-2t+2t)\mathbf{i} - (-t^2-5)\mathbf{j} + (-2t^2-10)\mathbf{k}\right] = \left\langle 0, \frac{1}{\sqrt{5}}, -\frac{2}{\sqrt{5}}\right\rangle$$

Since $\mathbf{B}(t)$ is constant, $\mathbf{B}'(t) = \mathbf{0}$, and $\mathbf{B}'(1)\cdot\mathbf{N}(1) = 0$. Thus, the torsion is $\tau(1) = -\dfrac{\mathbf{B}'(1)\cdot\mathbf{N}(1)}{|\mathbf{r}'(1)|} = -\dfrac{0}{\sqrt{6}} = 0$.

68. $\mathbf{r} = \left\langle t, \frac{1}{2}t^2, \frac{1}{3}t^3\right\rangle \;\Rightarrow\; \mathbf{r}' = \left\langle 1, t, t^2\right\rangle \;\Rightarrow\; \mathbf{r}'' = \langle 0, 1, 2t\rangle \;\Rightarrow\; \mathbf{r}''' = \langle 0, 0, 2\rangle$.

$\mathbf{r}'\times\mathbf{r}'' = \left\langle t^2, -2t, 1\right\rangle$. By Theorem 15, the torsion is $\tau = \dfrac{(\mathbf{r}'\times\mathbf{r}'')\cdot\mathbf{r}'''}{|\mathbf{r}'\times\mathbf{r}''|^2} = \dfrac{\left\langle t^2,-2t,1\right\rangle\cdot\langle 0,0,2\rangle}{\left(\sqrt{(t^2)^2+(-2t)^2+1^2}\right)^2} = \dfrac{2}{t^4+4t^2+1}$.

The torsion at $t=0$ is $\tau(0) = \frac{2}{1} = 2$.

69. $\mathbf{r} = \left\langle e^t, e^{-t}, t\right\rangle \;\Rightarrow\; \mathbf{r}' = \left\langle e^t, -e^{-t}, 1\right\rangle \;\Rightarrow\; \mathbf{r}'' = \left\langle e^t, e^{-t}, 0\right\rangle \;\Rightarrow\; \mathbf{r}''' = \left\langle e^t, -e^{-t}, 0\right\rangle$.

$\mathbf{r}'\times\mathbf{r}'' = \left\langle -e^{-t}, e^t, 2\right\rangle$. By Theorem 15, the torsion is

$$\tau = \frac{(\mathbf{r}'\times\mathbf{r}'')\cdot\mathbf{r}'''}{|\mathbf{r}'\times\mathbf{r}''|^2} = \frac{\left\langle -e^{-t}, e^t, 2\right\rangle\cdot\left\langle e^t, -e^{-t}, 0\right\rangle}{\left(\sqrt{(-e^{-t})^2+(e^t)^2+2^2}\right)^2} = \frac{-1-1+0}{e^{-2t}+e^{2t}+4} = \frac{-2}{e^{2t}+e^{-2t}+4}$$

The torsion at $t=0$ is $\tau(0) = \frac{-2}{1+1+4} = -\frac{1}{3}$.

70. $\mathbf{r} = \langle \cos t, \sin t, \sin t\rangle \;\Rightarrow\; \mathbf{r}' = \langle -\sin t, \cos t, \cos t\rangle \;\Rightarrow\; \mathbf{r}'' = \langle -\cos t, -\sin t, -\sin t\rangle \;\Rightarrow$

$\mathbf{r}''' = \langle \sin t, -\cos t, -\cos t\rangle$. $\mathbf{r}'\times\mathbf{r}'' = \langle 0, -1, 1\rangle$. By Theorem 15, the torsion is

$$\tau = \frac{(\mathbf{r}'\times\mathbf{r}'')\cdot\mathbf{r}'''}{|\mathbf{r}'\times\mathbf{r}''|^2} = \frac{\langle 0,-1,1\rangle\cdot\langle \sin t, -\cos t, -\cos t\rangle}{\left(\sqrt{0^2+(-1)^2+1^2}\right)^2} = \frac{0+\cos t-\cos t}{1+1} = 0$$

The torsion at $t=0$, or any value of t, is 0.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

71. $\mathbf{N} = \mathbf{B} \times \mathbf{T} \quad \Rightarrow$

$$\begin{aligned}
\frac{d\mathbf{N}}{ds} &= \frac{d}{ds}(\mathbf{B} \times \mathbf{T}) = \frac{d\mathbf{B}}{ds} \times \mathbf{T} + \mathbf{B} \times \frac{d\mathbf{T}}{ds} && \text{[by Formula 5 of Theorem 13.2.3]} \\
&= -\tau \mathbf{N} \times \mathbf{T} + \mathbf{B} \times \kappa \mathbf{N} && \text{[by Formulas 3 and 1]} \\
&= -\tau(\mathbf{N} \times \mathbf{T}) + \kappa(\mathbf{B} \times \mathbf{N}) && \text{[by Property 2 of Theorem 12.4.11]}
\end{aligned}$$

But $\mathbf{B} \times \mathbf{N} = \mathbf{B} \times (\mathbf{B} \times \mathbf{T}) = (\mathbf{B} \cdot \mathbf{T})\,\mathbf{B} - (\mathbf{B} \cdot \mathbf{B})\,\mathbf{T}$ [by Property 6 ofTheorem 12.4.11]

$$= 0 - \mathbf{T} = -\mathbf{T} \quad \Rightarrow$$

$d\mathbf{N}/ds = \tau(\mathbf{T} \times \mathbf{N}) - \kappa \mathbf{T} = -\kappa\,\mathbf{T} + \tau\,\mathbf{B}$.

72. (a) $\mathbf{r}' = s'\,\mathbf{T} \quad \Rightarrow \quad \mathbf{r}'' = s''\,\mathbf{T} + s'\,\mathbf{T}' = s''\,\mathbf{T} + s'\,\dfrac{d\mathbf{T}}{ds}s' = s''\,\mathbf{T} + \kappa(s')^2\,\mathbf{N}$ by the first Frenet-Serret formula.

(b) Using part (a), we have

$$\begin{aligned}
\mathbf{r}' \times \mathbf{r}'' &= (s'\,\mathbf{T}) \times [s''\,\mathbf{T} + \kappa(s')^2\,\mathbf{N}] \\
&= [(s'\,\mathbf{T}) \times (s''\,\mathbf{T})] + \left[(s'\mathbf{T}) \times (\kappa(s')^2\,\mathbf{N})\right] && \text{[by Property 3 of Theorem 12.4.11]} \\
&= (s's'')(\mathbf{T} \times \mathbf{T}) + \kappa(s')^3(\mathbf{T} \times \mathbf{N}) = \mathbf{0} + \kappa(s')^3\,\mathbf{B} = \kappa(s')^3\,\mathbf{B}
\end{aligned}$$

(c) Using part (a), we have

$$\begin{aligned}
\mathbf{r}''' &= [s''\,\mathbf{T} + \kappa(s')^2\,\mathbf{N}]' = s'''\,\mathbf{T} + s''\,\mathbf{T}' + \kappa'(s')^2\,\mathbf{N} + 2\kappa s's''\,\mathbf{N} + \kappa(s')^2\,\mathbf{N}' \\
&= s'''\,\mathbf{T} + s''\frac{d\mathbf{T}}{ds}\,s' + \kappa'(s')^2\,\mathbf{N} + 2\kappa s's''\,\mathbf{N} + \kappa(s')^2\,\frac{d\mathbf{N}}{ds}\,s' \\
&= s'''\,\mathbf{T} + s''s'\kappa\,\mathbf{N} + \kappa'(s')^2\,\mathbf{N} + 2\kappa s's''\,\mathbf{N} + \kappa(s')^3(-\kappa\,\mathbf{T} + \tau\,\mathbf{B}) && \text{[by Formulas 1 and 2]} \\
&= [s''' - \kappa^2(s')^3]\,\mathbf{T} + [3\kappa s's'' + \kappa'(s')^2]\,\mathbf{N} + \kappa\tau(s')^3\,\mathbf{B}
\end{aligned}$$

(d) Using parts (b) and (c) and the facts that $\mathbf{B} \cdot \mathbf{T} = 0$, $\mathbf{B} \cdot \mathbf{N} = 0$, and $\mathbf{B} \cdot \mathbf{B} = 1$, we get

$$\frac{(\mathbf{r}' \times \mathbf{r}'') \cdot \mathbf{r}'''}{|\mathbf{r}' \times \mathbf{r}''|^2} = \frac{\kappa(s')^3\,\mathbf{B} \cdot \left\{[s''' - \kappa^2(s')^3]\,\mathbf{T} + [3\kappa s's'' + \kappa'(s')^2]\,\mathbf{N} + \kappa\tau(s')^3\,\mathbf{B}\right\}}{|\kappa(s')^3\,\mathbf{B}|^2} = \frac{\kappa(s')^3\kappa\tau(s')^3}{[\kappa(s')^3]^2} = \tau.$$

73. First we find the quantities required to compute κ:

$$\mathbf{r}'(t) = \langle -a\sin t, a\cos t, b\rangle \quad \Rightarrow \quad \mathbf{r}''(t) = \langle -a\cos t, -a\sin t, 0\rangle \quad \Rightarrow \quad \mathbf{r}'''(t) = \langle a\sin t, -a\cos t, 0\rangle$$

$$|\mathbf{r}'(t)| = \sqrt{(-a\sin t)^2 + (a\cos t)^2 + b^2} = \sqrt{a^2 + b^2}$$

$$\mathbf{r}'(t) \times \mathbf{r}''(t) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -a\sin t & a\cos t & b \\ -a\cos t & -a\sin t & 0 \end{vmatrix} = ab\sin t\,\mathbf{i} - ab\cos t\,\mathbf{j} + a^2\,\mathbf{k}$$

$$|\mathbf{r}'(t) \times \mathbf{r}''(t)| = \sqrt{(ab\sin t)^2 + (-ab\cos t)^2 + (a^2)^2} = \sqrt{a^2b^2 + a^4} = a\sqrt{a^2 + b^2}$$

$$(\mathbf{r}'(t) \times \mathbf{r}''(t)) \cdot \mathbf{r}'''(t) = (ab\sin t)(a\sin t) + (-ab\cos t)(-a\cos t) + (a^2)(0) = a^2b$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

By Theorem 10, $\kappa(t) = \dfrac{|\mathbf{r}'(t) \times \mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \dfrac{a\sqrt{a^2+b^2}}{\left(\sqrt{a^2+b^2}\,\right)^3} = \dfrac{a}{a^2+b^2}$, which is a constant.

By Theorem 15, the torsion is

$$\tau = \frac{(\mathbf{r}' \times \mathbf{r}'') \cdot \mathbf{r}'''}{|\mathbf{r}' \times \mathbf{r}''|^2} = \frac{\left\langle ab\sin t, -ab\cos t, a^2\right\rangle \cdot \langle a\sin t, -a\cos t, 0\rangle}{\left(\sqrt{a^2b^2+a^4}\,\right)^2} = \frac{a^2b}{a^2(a^2+b^2)} = \frac{b}{a^2+b^2}, \text{ which is also a constant.}$$

74. $\mathbf{r} = \langle \sinh t, \cosh t, t\rangle \;\Rightarrow\; \mathbf{r}' = \langle \cosh t, \sinh t, 1\rangle$, $\mathbf{r}'' = \langle \sinh t, \cosh t, 0\rangle$, $\mathbf{r}''' = \langle \cosh t, \sinh t, 0\rangle \;\Rightarrow$

$\mathbf{r}' \times \mathbf{r}'' = \left\langle -\cosh t, \sinh t, \cosh^2 t - \sinh^2 t\right\rangle = \langle -\cosh t, \sinh t, 1\rangle \;\Rightarrow$

$$\kappa = \frac{|\mathbf{r}' \times \mathbf{r}''|}{|\mathbf{r}'|^3} = \frac{|\langle -\cosh t, \sinh t, 1\rangle|}{|\langle \cosh t, \sinh t, 1\rangle|^3} = \frac{\sqrt{\cosh^2 t + \sinh^2 t + 1}}{\left(\cosh^2 t + \sinh^2 t + 1\right)^{3/2}} = \frac{1}{\cosh^2 t + \sinh^2 t + 1} = \frac{1}{2\cosh^2 t},$$

$$\tau = \frac{(\mathbf{r}' \times \mathbf{r}'') \cdot \mathbf{r}'''}{|\mathbf{r}' \times \mathbf{r}''|^2} = \frac{\langle -\cosh t, \sinh t, 1\rangle \cdot \langle \cosh t, \sinh t, 0\rangle}{\cosh^2 t + \sinh^2 t + 1} = \frac{-\cosh^2 t + \sinh^2 t}{2\cosh^2 t} = \frac{-1}{2\cosh^2 t}$$

So at the point $(0, 1, 0)$, $t = 0$, and $\kappa = \frac{1}{2}$ and $\tau = -\frac{1}{2}$.

75. (a) At any point P on C where $\kappa(t) \neq 0$, the circle of curvature of C at P has center a distance $1/\kappa(t)$ from P in the direction of the unit normal vector $\mathbf{N}(t)$. The position vector of P is $\mathbf{r}(t)$, so we get a position vector for the center of curvature by adding $\dfrac{1}{\kappa(t)}\mathbf{N}(t)$ to $\mathbf{r}(t)$: $\mathbf{r}_e(t) = \mathbf{r}(t) + \dfrac{1}{\kappa(t)}\mathbf{N}(t)$, $\kappa(t) \neq 0$.

(b) $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j} + t\,\mathbf{k} \;\Rightarrow\; \mathbf{r}'(t) = -\sin t\mathbf{i} + \cos t\mathbf{j} + \mathbf{k} \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{\sin^2 t + \cos^2 t + 1^2} = \sqrt{2}$.

Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = -\dfrac{\sin t}{\sqrt{2}}\,\mathbf{i} + \dfrac{\cos t}{\sqrt{2}}\,\mathbf{j} + \dfrac{1}{\sqrt{2}}\,\mathbf{k} \;\Rightarrow\; \mathbf{T}'(t) = -\dfrac{\cos t}{\sqrt{2}}\,\mathbf{i} - \dfrac{\sin t}{\sqrt{2}}\,\mathbf{j} \;\Rightarrow\; |\mathbf{T}'(t)| = \dfrac{1}{\sqrt{2}}$.

So $\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = -\cos t\,\mathbf{i} - \sin t\,\mathbf{j}$ and $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{1/\sqrt{2}}{\sqrt{2}} = \dfrac{1}{2}$. Thus,

$$\mathbf{r}_e(t) = \mathbf{r}(t) + \frac{1}{\kappa(t)}\mathbf{N}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j} + t\,\mathbf{k} + 2(-\cos t\,\mathbf{i} - \sin t\,\mathbf{j}) = -\cos t\,\mathbf{i} - \sin t\,\mathbf{j} + t\,\mathbf{k}.$$

(c) The parabola $y = x^2$ can be parameterized $x = t$, $y = t^2$, which gives the corresponding vector equation

$\mathbf{r}(t) = \left\langle t, t^2\right\rangle \;\Rightarrow\; \mathbf{r}'(t) = \langle 1, 2t\rangle \;\Rightarrow\; |\mathbf{r}'(t)| = \sqrt{1+4t^2}$. Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{1+4t^2}}\langle 1, 2t\rangle \;\Rightarrow$

$$\begin{aligned}\mathbf{T}'(t) &= \frac{1}{\sqrt{1+4t^2}}\langle 0, 2\rangle - \frac{4t}{(1+4t^2)^{3/2}}\langle 1, 2t\rangle = \frac{1}{(1+4t^2)^{3/2}}\left[(1+4t^2)\langle 0, 2\rangle - 4t\langle 1, 2t\rangle\right] \\ &= \frac{1}{(1+4t^2)^{3/2}}\langle -4t, 2\rangle \;\Rightarrow\end{aligned}$$

$$|\mathbf{T}'(t)| = \frac{1}{(1+4t^2)^{3/2}}\sqrt{16t^2+4} = \frac{2}{1+4t^2}.$$

So $\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \dfrac{1+4t^2}{2} \cdot \dfrac{1}{(1+4t^2)^{3/2}}\langle -4t, 2\rangle = \dfrac{1}{\sqrt{1+4t^2}}\langle -2t, 1\rangle$ and

$\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{2/(1+4t^2)}{\sqrt{1+4t^2}} = \dfrac{2}{(1+4t^2)^{3/2}}$. Thus,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\mathbf{r}_e(t) = \mathbf{r}(t) + \frac{1}{\kappa(t)}\mathbf{N}(t) = \langle t, t^2 \rangle + \frac{(1+4t^2)^{3/2}}{2} \cdot \frac{1}{\sqrt{1+4t^2}} \langle -2t, 1 \rangle$$
$$= \langle t, t^2 \rangle + (1+4t^2) \langle -t, \tfrac{1}{2} \rangle = \langle -4t^3, \tfrac{1}{2} + 3t^2 \rangle$$

To obtain a function form of the answer, note that $x = -4t^3 \quad \Rightarrow \quad t = (-x/4)^{1/3}$, so $y_e = \frac{1}{2} + 3t^2 = \frac{1}{2} + 3[(-x/4)^{1/3}]^2 = \frac{1}{2} + 3(x/4)^{2/3}$.

76. (a) If C is planar, then it lies in a plane that we can express in the form $ax + by + cz = d$, where a, b, c, and d are not all zero. (See Equation 12.5.8.) For any t, the point $(x(t), y(t), z(t))$ lies on the curve and hence on the plane, so the equation $ax(t) + by(t) + cz(t) = d$ must be satisfied.

Conversely, if there exist scalars a, b, c, and d, not all zero, such that $ax(t) + by(t) + cz(t) = d$ for all t, then each point $(x(t), y(t), z(t))$ on C satisfies the equation $ax + by + cz = d$. By Exercise 12.5.83, this equation represents a plane, and hence C is planar.

(b) By part (a) there exist scalars a, b, c, and d, not all zero, such that for all t, $ax(t) + by(t) + cz(t) = d \quad \Rightarrow$ $\langle a, b, c \rangle \cdot \mathbf{r}(t) = d$. In addition, for all t, $ax'(t) + by'(t) + cz'(t) = 0 \quad \Rightarrow \quad \langle a, b, c \rangle \cdot \mathbf{r}'(t) = 0 \quad \Rightarrow \quad \mathbf{r}'$ is perpendicular to the normal vector $\mathbf{n} = \langle a, b, c \rangle$ of the plane containing $C \quad \Rightarrow \quad \mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|}$ is also perpendicular to $\mathbf{n}$.

Similarly, $\langle a, b, c \rangle \cdot \mathbf{r}''(t) = 0 \quad \Rightarrow \quad \mathbf{r}''$ is perpendicular to $\mathbf{n} \quad \Rightarrow$ $\mathbf{T}'(t) \cdot \mathbf{n} = \dfrac{d}{dt}\left(\dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|}\right) \cdot \mathbf{n} = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} \cdot \mathbf{n} + \dfrac{d}{dt}\left(\dfrac{\mathbf{1}}{|\mathbf{r}'(t)|}\right) \mathbf{r}'(t) \cdot \mathbf{n} = 0 + 0 = 0$. So then $\mathbf{N} \cdot \mathbf{n} = \dfrac{\mathbf{T}'(t) \cdot \mathbf{n}}{|\mathbf{T}'(t)|} = 0$, and hence $\mathbf{N}$ is perpendicular to $\mathbf{n}$. So, since $\mathbf{n} = \langle a, b, c \rangle$ is perpendicular to both $\mathbf{T}$ and $\mathbf{N}$, it follows that $\mathbf{B} = \mathbf{T} \times \mathbf{N}$ is parallel to $\mathbf{n}$ and hence is normal to the plane containing C.

(c) By part (b), $\mathbf{B}$ is normal to the plane containing C and so $\mathbf{B}(t) = \dfrac{\langle a, b, c \rangle}{\sqrt{a^2 + b^2 + c^2}}$ for all t, that is, $\mathbf{B}$ is a constant vector. Therefore, $\mathbf{B}'(t) = \mathbf{0}$ and $\tau(t) = -\dfrac{\mathbf{B}'(t) \cdot \mathbf{N}(t)}{|\mathbf{r}'(t)|} = -\dfrac{0}{|\mathbf{r}'(t)|} = 0$ for all t.

(d) The projection of the curve $\mathbf{r}(t) = \langle t, 2t, t^2 \rangle$ in the xy-plane is the curve $\mathbf{r}(t) = \langle t, 2t, 0 \rangle$, which is the line with corresponding parametric equations $x = t,\ y = 2t,\ z = 0$. Therefore, the equation of the plane containing $\mathbf{r}(t)$ is $y = 2x$, or $2x - y = 0$, with normal vector $\mathbf{n} = \langle 2, -1, 0 \rangle$. By part (b), $\mathbf{B} = \frac{1}{\sqrt{5}} \langle 2, -1, 0 \rangle$.

77. For one helix, the vector equation is $\mathbf{r}(t) = \langle 10 \cos t, 10 \sin t, 34t/(2\pi) \rangle$ (measuring in angstroms), because the radius of each helix is 10 angstroms, and z increases by 34 angstroms for each increase of 2π in t. Using the arc length formula, letting t go from 0 to $2.9 \times 10^8 \times 2\pi$, we find the approximate length of each helix to be

$$L = \int_0^{2.9\times10^8\times2\pi} |\mathbf{r}'(t)|\, dt = \int_0^{2.9\times10^8\times2\pi} \sqrt{(-10\sin t)^2 + (10\cos t)^2 + \left(\tfrac{34}{2\pi}\right)^2}\, dt = \sqrt{100 + \left(\tfrac{34}{2\pi}\right)^2}\, t \Big]_0^{2.9\times10^8\times2\pi}$$
$$= 2.9 \times 10^8 \times 2\pi \sqrt{100 + \left(\tfrac{34}{2\pi}\right)^2} \approx 2.07 \times 10^{10} \text{ Å} \text{ — more than two meters!}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

78. (a) For the function $F(x) = \begin{cases} 0 & \text{if } x < 0 \\ P(x) & \text{if } 0 < x < 1 \\ 1 & \text{if } x \geq 1 \end{cases}$ to be continuous, we must have $P(0) = 0$ and $P(1) = 1$.

For F' to be continuous, we must have $P'(0) = P'(1) = 0$. The curvature of the curve $y = F(x)$ at the point $(x, F(x))$ is $\kappa(x) = \dfrac{|F''(x)|}{\left(1 + [F'(x)]^2\right)^{3/2}}$. For $\kappa(x)$ to be continuous, we must have $P''(0) = P''(1) = 0$.

Write $P(x) = ax^5 + bx^4 + cx^3 + dx^2 + ex + f$. Then $P'(x) = 5ax^4 + 4bx^3 + 3cx^2 + 2dx + e$ and $P''(x) = 20ax^3 + 12bx^2 + 6cx + 2d$. Our six conditions are:

$$P(0) = 0 \quad \Rightarrow \quad f = 0 \quad \textbf{(1)} \qquad\qquad P(1) = 1 \quad \Rightarrow \quad a + b + c + d + e + f = 1 \quad \textbf{(2)}$$

$$P'(0) = 0 \quad \Rightarrow \quad e = 0 \quad \textbf{(3)} \qquad\qquad P'(1) = 0 \quad \Rightarrow \quad 5a + 4b + 3c + 2d + e = 0 \quad \textbf{(4)}$$

$$P''(0) = 0 \quad \Rightarrow \quad d = 0 \quad \textbf{(5)} \qquad\qquad P''(1) = 0 \quad \Rightarrow \quad 20a + 12b + 6c + 2d = 0 \quad \textbf{(6)}$$

From **(1)**, **(3)**, and **(5)**, we have $d = e = f = 0$. Thus **(2)**, **(4)** and **(6)** become **(7)** $a + b + c = 1$, **(8)** $5a + 4b + 3c = 0$, and **(9)** $10a + 6b + 3c = 0$. Subtracting **(8)** from **(9)** gives **(10)** $5a + 2b = 0$. Multiplying **(7)** by 3 and subtracting from **(8)** gives **(11)** $2a + b = -3$. Multiplying **(11)** by 2 and subtracting from **(10)** gives $a = 6$. By **(10)**, $b = -15$. By **(7)**, $c = 10$. Thus, $P(x) = 6x^5 - 15x^4 + 10x^3$.

(b)

13.4 Motion in Space: Velocity and Acceleration

1. (a) If $\mathbf{r}(t) = x(t)\,\mathbf{i} + y(t)\,\mathbf{j} + z(t)\,\mathbf{k}$ is the position vector of the particle at time t, then the average velocity over the time interval $[0, 1]$ is

$$\mathbf{v}_{\text{ave}} = \frac{\mathbf{r}(1) - \mathbf{r}(0)}{1 - 0} = \frac{(4.5\,\mathbf{i} + 6.0\,\mathbf{j} + 3.0\,\mathbf{k}) - (2.7\,\mathbf{i} + 9.8\,\mathbf{j} + 3.7\,\mathbf{k})}{1} = 1.8\,\mathbf{i} - 3.8\,\mathbf{j} - 0.7\,\mathbf{k}$$

Similarly, over the other intervals we have

$$[0.5, 1]: \quad \mathbf{v}_{\text{ave}} = \frac{\mathbf{r}(1) - \mathbf{r}(0.5)}{1 - 0.5} = \frac{(4.5\,\mathbf{i} + 6.0\,\mathbf{j} + 3.0\,\mathbf{k}) - (3.5\,\mathbf{i} + 7.2\,\mathbf{j} + 3.3\,\mathbf{k})}{0.5} = 2.0\,\mathbf{i} - 2.4\,\mathbf{j} - 0.6\,\mathbf{k}$$

$$[1, 2]: \quad \mathbf{v}_{\text{ave}} = \frac{\mathbf{r}(2) - \mathbf{r}(1)}{2 - 1} = \frac{(7.3\,\mathbf{i} + 7.8\,\mathbf{j} + 2.7\,\mathbf{k}) - (4.5\,\mathbf{i} + 6.0\,\mathbf{j} + 3.0\,\mathbf{k})}{1} = 2.8\,\mathbf{i} + 1.8\,\mathbf{j} - 0.3\,\mathbf{k}$$

$$[1, 1.5]: \quad \mathbf{v}_{\text{ave}} = \frac{\mathbf{r}(1.5) - \mathbf{r}(1)}{1.5 - 1} = \frac{(5.9\,\mathbf{i} + 6.4\,\mathbf{j} + 2.8\,\mathbf{k}) - (4.5\,\mathbf{i} + 6.0\,\mathbf{j} + 3.0\,\mathbf{k})}{0.5} = 2.8\,\mathbf{i} + 0.8\,\mathbf{j} - 0.4\,\mathbf{k}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) We can estimate the velocity at $t = 1$ by averaging the average velocities over the time intervals $[0.5, 1]$ and $[1, 1.5]$:

$\mathbf{v}(1) \approx \frac{1}{2}[(2\,\mathbf{i} - 2.4\,\mathbf{j} - 0.6\,\mathbf{k}) + (2.8\,\mathbf{i} + 0.8\,\mathbf{j} - 0.4\,\mathbf{k})] = 2.4\,\mathbf{i} - 0.8\,\mathbf{j} - 0.5\,\mathbf{k}$. Then the speed is

$|\mathbf{v}(1)| \approx \sqrt{(2.4)^2 + (-0.8)^2 + (-0.5)^2} \approx 2.58$.

2. (a) The average velocity over $2 \le t \le 2.4$ is

$\dfrac{\mathbf{r}(2.4) - \mathbf{r}(2)}{2.4 - 2} = 2.5\,[\mathbf{r}(2.4) - \mathbf{r}(2)]$, so we sketch a vector in the same direction but 2.5 times the length of $[\mathbf{r}(2.4) - \mathbf{r}(2)]$.

(b) The average velocity over $1.5 \le t \le 2$ is

$\dfrac{\mathbf{r}(2) - \mathbf{r}(1.5)}{2 - 1.5} = 2[\mathbf{r}(2) - \mathbf{r}(1.5)]$, so we sketch a vector in the same direction but twice the length of $[\mathbf{r}(2) - \mathbf{r}(1.5)]$.

(c) Using Equation 2 we have $\mathbf{v}(2) = \lim\limits_{h \to 0} \dfrac{\mathbf{r}(2+h) - \mathbf{r}(2)}{h}$.

(d) $\mathbf{v}(2)$ is tangent to the curve at $\mathbf{r}(2)$ and points in the direction of increasing t. Its length is the speed of the particle at $t = 2$. We can estimate the speed by averaging the lengths of the vectors found in parts (a) and (b) which represent the average speed over $2 \le t \le 2.4$ and $1.5 \le t \le 2$ respectively. Using the axes scale as a guide, we estimate the vectors to have lengths 2.8 and 2.7. Thus, we estimate the speed at $t = 2$ to be $|\mathbf{v}(2)| \approx \frac{1}{2}(2.8 + 2.7) = 2.75$ and we draw the velocity vector $\mathbf{v}(2)$ with this length.

3. $\mathbf{r}(t) = \left\langle -\frac{1}{2}t^2, t \right\rangle \quad \Rightarrow$ At $t = 2$:

$\mathbf{v}(t) = \mathbf{r}'(t) = \langle -t, 1 \rangle$ $\qquad \mathbf{v}(2) = \langle -2, 1 \rangle$

$\mathbf{a}(t) = \mathbf{r}''(t) = \langle -1, 0 \rangle$ $\qquad \mathbf{a}(2) = \langle -1, 0 \rangle$

$|\mathbf{v}(t)| = \sqrt{t^2 + 1}$

Notice that $x = -\frac{1}{2}y^2$, so the path is a parabola.

4. $\mathbf{r}(t) = \left\langle t^2, 1/t^2 \right\rangle \quad \Rightarrow$ At $t = 1$:

$\mathbf{v}(t) = \mathbf{r}'(t) = \left\langle 2t, -2/t^3 \right\rangle$ $\qquad \mathbf{v}(1) = \langle 2, -2 \rangle$

$\mathbf{a}(t) = \mathbf{r}''(t) = \left\langle 2, 6/t^4 \right\rangle$ $\qquad \mathbf{a}(1) = \langle 2, 6 \rangle$

$|\mathbf{v}(t)| = \sqrt{4t^2 + 4/t^6} = 2\sqrt{t^2 + 1/t^6}$

Notice that $y = 1/x$ and $x > 0$, so the path is part of the hyperbola $y = 1/x$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $\mathbf{r}(t) = 3\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j} \quad \Rightarrow$ At $t = \pi/3$:

$\mathbf{v}(t) = -3\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j}$ $\qquad \mathbf{v}\left(\frac{\pi}{3}\right) = -\frac{3\sqrt{3}}{2}\,\mathbf{i} + \mathbf{j}$

$\mathbf{a}(t) = -3\cos t\,\mathbf{i} - 2\sin t\,\mathbf{j}$ $\qquad \mathbf{a}\left(\frac{\pi}{3}\right) = -\frac{3}{2}\,\mathbf{i} - \sqrt{3}\,\mathbf{j}$

$$|\mathbf{v}(t)| = \sqrt{9\sin^2 t + 4\cos^2 t} = \sqrt{5\sin^2 t + 4\sin^2 t + 4\cos^2 t}$$
$$= \sqrt{4 + 5\sin^2 t}$$

Notice that $x^2/9 + y^2/4 = \sin^2 t + \cos^2 t = 1$, so the path is an ellipse.

6. $\mathbf{r}(t) = e^t\,\mathbf{i} + e^{2t}\,\mathbf{j} \quad \Rightarrow$ At $t = 0$:

$\mathbf{v}(t) = e^t\,\mathbf{i} + 2e^{2t}\,\mathbf{j}$ $\qquad \mathbf{v}(0) = \mathbf{i} + 2\,\mathbf{j}$

$\mathbf{a}(t) = e^t\,\mathbf{i} + 4e^{2t}\,\mathbf{j}$ $\qquad \mathbf{a}(0) = \mathbf{i} + 4\,\mathbf{j}$

$|\mathbf{v}(t)| = \sqrt{e^{2t} + 4e^{4t}} = e^t\sqrt{1 + 4e^{2t}}$

Notice that $y = e^{2t} = \left(e^t\right)^2 = x^2$, so the particle travels along a parabola, but $x = e^t$, so $x > 0$.

7. $\mathbf{r}(t) = t\,\mathbf{i} + t^2\,\mathbf{j} + 2\,\mathbf{k} \quad \Rightarrow$ At $t = 1$:

$\mathbf{v}(t) = \mathbf{i} + 2t\,\mathbf{j}$ $\qquad \mathbf{v}(1) = \mathbf{i} + 2\,\mathbf{j}$

$\mathbf{a}(t) = 2\,\mathbf{j}$ $\qquad \mathbf{a}(1) = 2\,\mathbf{j}$

$|\mathbf{v}(t)| = \sqrt{1 + 4t^2}$

Here $x = t$, $y = t^2 \quad \Rightarrow \quad y = x^2$ and $z = 2$, so the path of the particle is a parabola in the plane $z = 2$.

8. $\mathbf{r}(t) = t\,\mathbf{i} + 2\cos t\,\mathbf{j} + \sin t\,\mathbf{k} \quad \Rightarrow$ At $t = 0$:

$\mathbf{v}(t) = \mathbf{i} - 2\sin t\,\mathbf{j} + \cos t\,\mathbf{k}$ $\qquad \mathbf{v}(0) = \mathbf{i} + \mathbf{k}$

$\mathbf{a}(t) = -2\cos t\,\mathbf{j} - \sin t\,\mathbf{k}$ $\qquad \mathbf{a}(0) = -2\,\mathbf{j}$

$|\mathbf{v}(t)| = \sqrt{1 + 4\sin^2 t + \cos^2 t} = \sqrt{2 + 3\sin^2 t}$

Since $y^2/4 + z^2 = 1$, $x = t$, the path of the particle is an elliptical helix about the x-axis.

9. $\mathbf{r}(t) = \left\langle t^2 + t, t^2 - t, t^3 \right\rangle \quad \Rightarrow \quad \mathbf{v}(t) = \mathbf{r}'(t) = \left\langle 2t + 1, 2t - 1, 3t^2 \right\rangle, \quad \mathbf{a}(t) = \mathbf{v}'(t) = \langle 2, 2, 6t \rangle,$

$|\mathbf{v}(t)| = \sqrt{(2t+1)^2 + (2t-1)^2 + (3t^2)^2} = \sqrt{9t^4 + 8t^2 + 2}.$

10. $\mathbf{r}(t) = \langle 2\cos t, 3t, 2\sin t \rangle \quad \Rightarrow \quad \mathbf{v}(t) = \mathbf{r}'(t) = \langle -2\sin t, 3, 2\cos t \rangle, \quad \mathbf{a}(t) = \mathbf{v}'(t) = \langle -2\cos t, 0, -2\sin t \rangle,$

$|\mathbf{v}(t)| = \sqrt{4\sin^2 t + 9 + 4\cos^2 t} = \sqrt{13}.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. $\mathbf{r}(t) = \sqrt{2}\,t\,\mathbf{i} + e^t\,\mathbf{j} + e^{-t}\,\mathbf{k} \quad \Rightarrow \quad \mathbf{v}(t) = \mathbf{r}'(t) = \sqrt{2}\,\mathbf{i} + e^t\,\mathbf{j} - e^{-t}\,\mathbf{k}, \quad \mathbf{a}(t) = \mathbf{v}'(t) = e^t\,\mathbf{j} + e^{-t}\,\mathbf{k},$

$|\mathbf{v}(t)| = \sqrt{2 + e^{2t} + e^{-2t}} = \sqrt{(e^t + e^{-t})^2} = e^t + e^{-t}.$

12. $\mathbf{r}(t) = t^2\,\mathbf{i} + 2t\,\mathbf{j} + \ln t\,\mathbf{k} \quad \Rightarrow \quad \mathbf{v}(t) = \mathbf{r}'(t) = 2t\,\mathbf{i} + 2\,\mathbf{j} + (1/t)\,\mathbf{k}, \quad \mathbf{a}(t) = \mathbf{v}'(t) = 2\,\mathbf{i} - (1/t^2)\,\mathbf{k},$

$|\mathbf{v}(t)| = \sqrt{4t^2 + 4 + (1/t^2)} = \sqrt{[2t + (1/t)]^2} = |2t + (1/t)|.$

13. $\mathbf{r}(t) = e^t(\cos t\,\mathbf{i} + \sin t\,\mathbf{j} + t\,\mathbf{k}) = e^t \cos t\,\mathbf{i} + e^t \sin t\,\mathbf{j} + te^t\,\mathbf{k} \quad \Rightarrow$

$$\begin{aligned}\mathbf{v}(t) = \mathbf{r}'(t) &= [e^t(-\sin t) + (\cos t)e^t]\,\mathbf{i} + [e^t \cos t + (\sin t)e^t]\,\mathbf{j} + (te^t + e^t)\,\mathbf{k} \\ &= e^t[(\cos t - \sin t)\,\mathbf{i} + (\sin t + \cos t)\,\mathbf{j} + (t+1)\,\mathbf{k}]\end{aligned}$$

$$\begin{aligned}\mathbf{a}(t) = \mathbf{v}'(t) &= [e^t(-\sin t - \cos t) + (\cos t - \sin t)e^t]\,\mathbf{i} + [e^t(\cos t - \sin t) + (\sin t + \cos t)e^t]\,\mathbf{j} \\ &\qquad + [e^t \cdot 1 + (t+1)e^t]\,\mathbf{k} \\ &= e^t[-2\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j} + (t+2)\,\mathbf{k}]\end{aligned}$$

$$\begin{aligned}|\mathbf{v}(t)| &= \sqrt{e^{2t}(\cos t - \sin t)^2 + e^{2t}(\sin t + \cos t)^2 + e^{2t}(t+1)^2} \\ &= \sqrt{e^{2t}}\sqrt{\cos^2 t + \sin^2 t - 2\cos t \sin t + \sin^2 t + \cos^2 t + 2\sin t \cos t + t^2 + 2t + 1} \\ &= e^t\sqrt{t^2 + 2t + 3}\end{aligned}$$

14. $\mathbf{r}(t) = \left\langle t^2, \sin t - t\cos t, \cos t + t\sin t \right\rangle \quad \Rightarrow$

$\mathbf{v}(t) = \mathbf{r}'(t) = \langle 2t, \cos t - (-t\sin t + \cos t), -\sin t + t\cos t + \sin t\rangle = \langle 2t, t\sin t, t\cos t\rangle,$

$\mathbf{a}(t) = \mathbf{v}'(t) = \langle 2, t\cos t + \sin t, -t\sin t + \cos t\rangle,$

$|\mathbf{v}(t)| = \sqrt{4t^2 + t^2\sin^2 t + t^2\cos^2 t} = \sqrt{4t^2 + t^2} = \sqrt{5t^2} = \sqrt{5}\,t \quad$ [since $t \geq 0$].

15. $\mathbf{a}(t) = 2\,\mathbf{i} + 2t\,\mathbf{k} \quad \Rightarrow \quad \mathbf{v}(t) = \int \mathbf{a}(t)\,dt = \int (2\,\mathbf{i} + 2t\,\mathbf{k})\,dt = 2t\,\mathbf{i} + t^2\,\mathbf{k} + \mathbf{C}$. Then $\mathbf{v}(0) = \mathbf{C}$ but we were given that $\mathbf{v}(0) = 3\,\mathbf{i} - \mathbf{j}$, so $\mathbf{C} = 3\,\mathbf{i} - \mathbf{j}$ and $\mathbf{v}(t) = 2t\,\mathbf{i} + t^2\,\mathbf{k} + 3\,\mathbf{i} - \mathbf{j} = (2t+3)\,\mathbf{i} - \mathbf{j} + t^2\,\mathbf{k}$.

$\mathbf{r}(t) = \int \mathbf{v}(t)\,dt = \int \left[(2t+3)\,\mathbf{i} - \mathbf{j} + t^2\,\mathbf{k}\right] dt = (t^2 + 3t)\,\mathbf{i} - t\,\mathbf{j} + \frac{1}{3}t^3\,\mathbf{k} + \mathbf{D}$. Here $\mathbf{r}(0) = \mathbf{D}$ and we were given that $\mathbf{r}(0) = \mathbf{j} + \mathbf{k}$, so $\mathbf{D} = \mathbf{j} + \mathbf{k}$ and $\mathbf{r}(t) = (t^2 + 3t)\,\mathbf{i} + (1 - t)\,\mathbf{j} + \left(\frac{1}{3}t^3 + 1\right)\mathbf{k}$.

16. $\mathbf{a}(t) = \sin t\,\mathbf{i} + 2\cos t\,\mathbf{j} + 6t\,\mathbf{k} \quad \Rightarrow \quad \mathbf{v}(t) = \int \mathbf{a}(t)\,dt = \int (\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j} + 6t\,\mathbf{k})\,dt = -\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j} + 3t^2\,\mathbf{k} + \mathbf{C}$.

Then $\mathbf{v}(0) = -\mathbf{i} + \mathbf{C}$ but we were given that $\mathbf{v}(0) = -\mathbf{k}$, so $-\mathbf{i} + \mathbf{C} = -\mathbf{k} \quad \Rightarrow \quad \mathbf{C} = \mathbf{i} - \mathbf{k}$

and $\mathbf{v}(t) = (1 - \cos t)\,\mathbf{i} + 2\sin t\,\mathbf{j} + (3t^2 - 1)\,\mathbf{k}$.

$\mathbf{r}(t) = \int \mathbf{v}(t)\,dt = \int \left[(1 - \cos t)\,\mathbf{i} + 2\sin t\,\mathbf{j} + (3t^2 - 1)\,\mathbf{k}\right] dt = (t - \sin t)\,\mathbf{i} - 2\cos t\,\mathbf{j} + (t^3 - t)\,\mathbf{k} + \mathbf{D}$. Here $\mathbf{r}(0) = -2\,\mathbf{j} + \mathbf{D}$ and we were given that $\mathbf{r}(0) = \mathbf{j} - 4\,\mathbf{k}$, so $\mathbf{D} = 3\,\mathbf{j} - 4\,\mathbf{k}$ and

$\mathbf{r}(t) = (t - \sin t)\,\mathbf{i} + (3 - 2\cos t)\,\mathbf{j} + (t^3 - t - 4)\,\mathbf{k}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. (a) $\mathbf{a}(t) = 2t\,\mathbf{i} + \sin t\,\mathbf{j} + \cos 2t\,\mathbf{k} \quad \Rightarrow$

$\mathbf{v}(t) = \int (2t\,\mathbf{i} + \sin t\,\mathbf{j} + \cos 2t\,\mathbf{k})\,dt = t^2\,\mathbf{i} - \cos t\,\mathbf{j} + \frac{1}{2}\sin 2t\,\mathbf{k} + \mathbf{C}$

and $\mathbf{i} = \mathbf{v}(0) = -\mathbf{j} + \mathbf{C}$, so $\mathbf{C} = \mathbf{i} + \mathbf{j}$

and $\mathbf{v}(t) = \left(t^2 + 1\right)\mathbf{i} + (1 - \cos t)\,\mathbf{j} + \frac{1}{2}\sin 2t\,\mathbf{k}$.

$\mathbf{r}(t) = \int [\left(t^2 + 1\right)\,\mathbf{i} + (1 - \cos t)\,\mathbf{j} + \frac{1}{2}\sin 2t\,\mathbf{k}]dt$

$= \left(\frac{1}{3}t^3 + t\right)\,\mathbf{i} + (t - \sin t)\,\mathbf{j} - \frac{1}{4}\cos 2t\,\mathbf{k} + \mathbf{D}$

(b)

But $\mathbf{j} = \mathbf{r}(0) = -\frac{1}{4}\,\mathbf{k} + \mathbf{D}$, so $\mathbf{D} = \mathbf{j} + \frac{1}{4}\,\mathbf{k}$ and $\mathbf{r}(t) = \left(\frac{1}{3}t^3 + t\right)\mathbf{i} + (t - \sin t + 1)\,\mathbf{j} + \left(\frac{1}{4} - \frac{1}{4}\cos 2t\right)\mathbf{k}$.

18. (a) $\mathbf{a}(t) = t\,\mathbf{i} + e^t\,\mathbf{j} + e^{-t}\,\mathbf{k} \quad \Rightarrow$

$\mathbf{v}(t) = \int \left(t\,\mathbf{i} + e^t\,\mathbf{j} + e^{-t}\,\mathbf{k}\right)dt = \frac{1}{2}t^2\,\mathbf{i} + e^t\,\mathbf{j} - e^{-t}\,\mathbf{k} + \mathbf{C}$

and $\mathbf{k} = \mathbf{v}(0) = \mathbf{j} - \mathbf{k} + \mathbf{C}$, so $\mathbf{C} = -\mathbf{j} + 2\,\mathbf{k}$

and $\mathbf{v}(t) = \frac{1}{2}t^2\,\mathbf{i} + \left(e^t - 1\right)\,\mathbf{j} + \left(2 - e^{-t}\right)\,\mathbf{k}$.

$\mathbf{r}(t) = \int \left[\frac{1}{2}t^2\,\mathbf{i} + (e^t - 1)\,\mathbf{j} + (2 - e^{-t})\,\mathbf{k}\right]dt$

$= \frac{1}{6}t^3\,\mathbf{i} + (e^t - t)\,\mathbf{j} + (e^{-t} + 2t)\,\mathbf{k} + \mathbf{D}$

(b)

But $\mathbf{j} + \mathbf{k} = \mathbf{r}(0) = \mathbf{j} + \mathbf{k} + \mathbf{D}$, so $\mathbf{D} = \mathbf{0}$ and $\mathbf{r}(t) = \frac{1}{6}t^3\,\mathbf{i} + (e^t - t)\,\mathbf{j} + (e^{-t} + 2t)\,\mathbf{k}$.

19. $\mathbf{r}(t) = \left\langle t^2, 5t, t^2 - 16t\right\rangle \quad \Rightarrow \quad \mathbf{v}(t) = \langle 2t, 5, 2t - 16\rangle$, $|\mathbf{v}(t)| = \sqrt{4t^2 + 25 + 4t^2 - 64t + 256} = \sqrt{8t^2 - 64t + 281}$

and $\dfrac{d}{dt}|\mathbf{v}(t)| = \frac{1}{2}(8t^2 - 64t + 281)^{-1/2}(16t - 64)$. This is zero if and only if the numerator is zero, that is,

$16t - 64 = 0$ or $t = 4$. Since $\dfrac{d}{dt}|\mathbf{v}(t)| < 0$ for $t < 4$ and $\dfrac{d}{dt}|\mathbf{v}(t)| > 0$ for $t > 4$, the minimum speed of $\sqrt{153}$ is attained at $t = 4$ units of time.

20. Since $\mathbf{r}(t) = t^3\,\mathbf{i} + t^2\,\mathbf{j} + t^3\,\mathbf{k}$, $\mathbf{a}(t) = \mathbf{r}''(t) = 6t\,\mathbf{i} + 2\,\mathbf{j} + 6t\,\mathbf{k}$. By Newton's Second Law,

$\mathbf{F}(t) = m\,\mathbf{a}(t) = 6mt\,\mathbf{i} + 2m\,\mathbf{j} + 6mt\,\mathbf{k}$ is the required force.

21. $|\mathbf{F}(t)| = 20$ N in the direction of the positive z-axis, so $\mathbf{F}(t) = 20\,\mathbf{k}$. Also $m = 4$ kg, $\mathbf{r}(0) = \mathbf{0}$ and $\mathbf{v}(0) = \mathbf{i} - \mathbf{j}$.

Since $20\mathbf{k} = \mathbf{F}(t) = 4\,\mathbf{a}(t)$, $\mathbf{a}(t) = 5\,\mathbf{k}$. Then $\mathbf{v}(t) = 5t\,\mathbf{k} + \mathbf{c}_1$ where $\mathbf{c}_1 = \mathbf{i} - \mathbf{j}$ so $\mathbf{v}(t) = \mathbf{i} - \mathbf{j} + 5t\,\mathbf{k}$ and the speed is $|\mathbf{v}(t)| = \sqrt{1 + 1 + 25t^2} = \sqrt{25t^2 + 2}$. Also $\mathbf{r}(t) = t\,\mathbf{i} - t\,\mathbf{j} + \frac{5}{2}t^2\,\mathbf{k} + \mathbf{c}_2$ and $\mathbf{0} = \mathbf{r}(0)$, so $\mathbf{c}_2 = \mathbf{0}$

and $\mathbf{r}(t) = t\,\mathbf{i} - t\,\mathbf{j} + \frac{5}{2}t^2\,\mathbf{k}$.

22. The argument here is the same as that in the proof of Theorem 13.2.4 with $\mathbf{r}(t)$ replaced by $\mathbf{v}(t)$ and $\mathbf{r}'(t)$ replaced by $\mathbf{a}(t)$.

23. $|\mathbf{v}(0)| = 200$ m/s and, since the angle of elevation is 60°, a unit vector in the direction of the velocity is

$(\cos 60^\circ)\mathbf{i} + (\sin 60^\circ)\mathbf{j} = \frac{1}{2}\mathbf{i} + \frac{\sqrt{3}}{2}\,\mathbf{j}$. Thus $\mathbf{v}(0) = 200\left(\frac{1}{2}\mathbf{i} + \frac{\sqrt{3}}{2}\,\mathbf{j}\right) = 100\,\mathbf{i} + 100\sqrt{3}\,\mathbf{j}$ and if we set up the axes so that the projectile starts at the origin, then $\mathbf{r}(0) = \mathbf{0}$. Ignoring air resistance, the only force is that due to gravity, so

$\mathbf{F}(t) = m\mathbf{a}(t) = -mg\,\mathbf{j}$ where $g \approx 9.8$ m/s^2. Thus $\mathbf{a}(t) = -9.8\,\mathbf{j}$ and, integrating, we have $\mathbf{v}(t) = -9.8t\,\mathbf{j} + \mathbf{C}$. But

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$100\,\mathbf{i} + 100\sqrt{3}\,\mathbf{j} = \mathbf{v}(0) = \mathbf{C}$, so $\mathbf{v}(t) = 100\,\mathbf{i} + \left(100\sqrt{3} - 9.8t\right)\mathbf{j}$ and then (integrating again)

$\mathbf{r}(t) = 100\,t\,\mathbf{i} + \left(100\sqrt{3}\,t - 4.9t^2\right)\mathbf{j} + \mathbf{D}$ where $\mathbf{0} = \mathbf{r}(0) = \mathbf{D}$. Thus the position function of the projectile is

$\mathbf{r}(t) = 100\,t\,\mathbf{i} + \left(100\sqrt{3}\,t - 4.9t^2\right)\mathbf{j}$.

(a) Parametric equations for the projectile are $x(t) = 100t$, $y(t) = 100\sqrt{3}\,t - 4.9t^2$. The projectile reaches the ground when

$y(t) = 0$ (and $t > 0$) $\Rightarrow$ $100\sqrt{3}\,t - 4.9t^2 = t\left(100\sqrt{3} - 4.9t\right) = 0$ $\Rightarrow$ $t = \frac{100\sqrt{3}}{4.9} \approx 35.3$ s. So the range is

$x\left(\frac{100\sqrt{3}}{4.9}\right) = 100\left(\frac{100\sqrt{3}}{4.9}\right) \approx 3535$ m.

(b) The maximum height is reached when $y(t)$ has a critical number (or equivalently, when the vertical component of velocity is 0): $y'(t) = 0$ $\Rightarrow$ $100\sqrt{3} - 9.8t = 0$ $\Rightarrow$ $t = \frac{100\sqrt{3}}{9.8} \approx 17.7$ s. Thus the maximum height is

$y\left(\frac{100\sqrt{3}}{9.8}\right) = 100\sqrt{3}\left(\frac{100\sqrt{3}}{9.8}\right) - 4.9\left(\frac{100\sqrt{3}}{9.8}\right)^2 \approx 1531$ m.

(c) From part (a), impact occurs at $t = \frac{100\sqrt{3}}{4.9}$ s. Thus, the velocity at impact is

$\mathbf{v}\left(\frac{100\sqrt{3}}{4.9}\right) = 100\,\mathbf{i} + \left[100\sqrt{3} - 9.8\left(\frac{100\sqrt{3}}{4.9}\right)\right]\mathbf{j} = 100\,\mathbf{i} - 100\sqrt{3}\,\mathbf{j}$ and the speed is

$\left|\mathbf{v}\left(\frac{100\sqrt{3}}{4.9}\right)\right| = \sqrt{10{,}000 + 30{,}000} = 200$ m/s.

24. As in Exercise 23, $\mathbf{v}(t) = 100\,\mathbf{i} + \left(100\sqrt{3} - 9.8t\right)\mathbf{j}$ and $\mathbf{r}(t) = 100\,t\,\mathbf{i} + \left(100\sqrt{3}\,t - 4.9t^2\right)\mathbf{j} + \mathbf{D}$.

But $\mathbf{r}(0) = 100\,\mathbf{j}$, so $\mathbf{D} = 100\,\mathbf{j}$ and $\mathbf{r}(t) = 100\,t\,\mathbf{i} + \left(100 + 100\sqrt{3}\,t - 4.9t^2\right)\mathbf{j}$.

(a) $y = 0$ $\Rightarrow$ $100 + 100\sqrt{3}\,t - 4.9t^2 = 0$ or $4.9t^2 - 100\sqrt{3}\,t - 100 = 0$. From the quadratic formula we have

$t = \dfrac{100\sqrt{3} \pm \sqrt{(-100\sqrt{3})^2 - 4(4.9)(-100)}}{2(4.9)} = \dfrac{100\sqrt{3} \pm \sqrt{31{,}960}}{9.8}$. Taking the positive t-value gives

$t = \frac{100\sqrt{3} + \sqrt{31{,}960}}{9.8} \approx 35.9$ s. Thus the range is $x = 100 \cdot \frac{100\sqrt{3} + \sqrt{31{,}960}}{9.8} \approx 3592$ m.

(b) The maximum height is attained when $\dfrac{dy}{dt} = 0$ $\Rightarrow$ $100\sqrt{3} - 9.8t = 0$ $\Rightarrow$ $t = \frac{100\sqrt{3}}{9.8} \approx 17.7$ s and the

maximum height is $100 + 100\sqrt{3}\left(\frac{100\sqrt{3}}{9.8}\right) - 4.9\left(\frac{100\sqrt{3}}{9.8}\right)^2 \approx 1631$ m.

Alternate solution: Because the projectile is fired in the same direction and with the same velocity as in Exercise 23, but from a point 100 m higher, the maximum height reached is 100 m higher than that found in Exercise 23, that is, $1531 \text{ m} + 100 \text{ m} = 1631$ m.

(c) From part (a), impact occurs at $t = \frac{100\sqrt{3} + \sqrt{31{,}960}}{9.8}$ s. Thus the velocity at impact is

$\mathbf{v}\left(\frac{100\sqrt{3} + \sqrt{31{,}960}}{9.8}\right) = 100\,\mathbf{i} + \left[100\sqrt{3} - 9.8\left(\frac{100\sqrt{3} + \sqrt{31{,}960}}{9.8}\right)\right]\mathbf{j} = 100\,\mathbf{i} - \sqrt{31{,}960}\,\mathbf{j}$ and the speed is

$|\mathbf{v}| = \sqrt{10{,}000 + 31{,}960} = \sqrt{41{,}960} \approx 205$ m/s.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. As in Example 5, $\mathbf{r}(t) = (v_0 \cos 45°)t\,\mathbf{i} + \left[(v_0 \sin 45°)t - \frac{1}{2}gt^2\right]\mathbf{j} = \frac{1}{2}\left[v_0\sqrt{2}\,t\,\mathbf{i} + \left(v_0\sqrt{2}\,t - gt^2\right)\mathbf{j}\right]$. The ball lands when $y = 0$ (and $t > 0$) $\Rightarrow$ $t = \dfrac{v_0\sqrt{2}}{g}$ s. Now since it lands 90 m away, $90 = x = \frac{1}{2}v_0\sqrt{2}\,\dfrac{v_0\sqrt{2}}{g}$ or $v_0^2 = 90g$ and the initial velocity is $v_0 = \sqrt{90g} \approx 30$ m/s.

26. Let α be the angle of elevation. Here $v_0 = 400$ m/s and from Example 5, the horizontal distance traveled by the projectile is $d = \dfrac{v_0^2 \sin 2\alpha}{g}$. We want $\dfrac{400^2 \sin 2\alpha}{g} = 3000 \;\Rightarrow\; \sin 2\alpha = \dfrac{3000g}{400^2} \approx 0.1838 \;\Rightarrow\; 2\alpha \approx \sin^{-1}(0.1838) \approx 10.6°$ or $2\alpha \approx 180° - 10.6° = 169.4°$. Thus two angles of elevation are $\alpha \approx 5.3°$ and $\alpha \approx 84.7°$.

27. As in Example 5, $\mathbf{r}(t) = (v_0 \cos 36°)t\,\mathbf{i} + \left[(v_0 \sin 36°)t - \frac{1}{2}gt^2\right]\mathbf{j}$ and then $\mathbf{v}(t) = \mathbf{r}'(t) = (v_0 \cos 36°)\,\mathbf{i} + [(v_0 \sin 36°) - gt]\,\mathbf{j}$. The shell reaches its maximum height when the vertical component of velocity is zero, so $(v_0 \sin 36°) - gt = 0 \;\Rightarrow\; t = \dfrac{v_0 \sin 36°}{g}$. The vertical height of the shell at that time is 1600 ft, so

$$(v_0 \sin 36°)\left(\frac{v_0 \sin 36°}{g}\right) - \tfrac{1}{2}g\left(\frac{v_0 \sin 36°}{g}\right)^2 = 1600 \;\Rightarrow\; \left(\frac{v_0^2 \sin^2 36°}{g}\right) - \frac{1}{2}\left(\frac{v_0^2 \sin^2 36°}{g}\right) = 1600 \;\Rightarrow$$

$$\frac{v_0^2 \sin^2 36°}{2g} = 1600 \;\Rightarrow\; v_0^2 = \frac{1600(2g)}{\sin^2 36°} \;\Rightarrow\; v_0 = \sqrt{\frac{3200g}{\sin^2 36°}} \approx \frac{\sqrt{3200(32)}}{\sin 36°} \approx 544 \text{ ft/s}.$$

28. Here $v_0 = 115$ ft/s, the angle of elevation is $\alpha = 50°$, and if we place the origin at home plate, then $\mathbf{r}(0) = 3\,\mathbf{j}$. As in Example 5, we have $\mathbf{r}\,(t) = -\frac{1}{2}gt^2\,\mathbf{j} + t\,\mathbf{v}_0 + \mathbf{D}$ where $\mathbf{D} = \mathbf{r}(0) = 3\,\mathbf{j}$ and $\mathbf{v}_0 = v_0 \cos\alpha\,\mathbf{i} + v_0 \sin\alpha\,\mathbf{j}$, so $\mathbf{r}(t) = (v_0 \cos\alpha)t\,\mathbf{i} + \left[(v_0 \sin\alpha)t - \frac{1}{2}gt^2 + 3\right]\mathbf{j}$. Thus, parametric equations for the trajectory of the ball are $x = (v_0 \cos\alpha)t,\; y = (v_0 \sin\alpha)t - \frac{1}{2}gt^2 + 3$. The ball reaches the fence when $x = 400 \;\Rightarrow$ $(v_0 \cos\alpha)t = 400 \;\Rightarrow\; t = \dfrac{400}{v_0 \cos\alpha} = \dfrac{400}{115 \cos 50°} \approx 5.41$ s. At this time, the height of the ball is $y = (v_0 \sin\alpha)t - \frac{1}{2}gt^2 + 3 \approx (115 \sin 50°)(5.41) - \frac{1}{2}(32)(5.41)^2 + 3 \approx 11.2$ ft. Since the fence is 10 ft high, the ball clears the fence.

29. Place the catapult at the origin and assume the catapult is 100 meters from the city, so the city lies between $(100, 0)$ and $(600, 0)$. The initial speed is $v_0 = 80$ m/s and let θ be the angle the catapult is set at. As in Example 5, the trajectory of the catapulted rock is given by $\mathbf{r}\,(t) = (80 \cos\theta)t\,\mathbf{i} + \left[(80 \sin\theta)t - 4.9t^2\right]\mathbf{j}$. The top of the near city wall is at $(100, 15)$, which the rock will hit when $(80 \cos\theta)\,t = 100 \;\Rightarrow\; t = \dfrac{5}{4\cos\theta}$ and $(80 \sin\theta)t - 4.9t^2 = 15 \;\Rightarrow$

$80 \sin\theta \cdot \dfrac{5}{4\cos\theta} - 4.9\left(\dfrac{5}{4\cos\theta}\right)^2 = 15 \;\Rightarrow\; 100 \tan\theta - 7.65625 \sec^2\theta = 15$. Replacing $\sec^2\theta$ with $\tan^2\theta + 1$ gives $7.65625 \tan^2\theta - 100 \tan\theta + 22.65625 = 0$. Using the quadratic formula, we have $\tan\theta \approx 0.230635,\, 12.8306 \;\Rightarrow$ $\theta \approx 13.0°, 85.5°$. So for $13.0° < \theta < 85.5°$, the rock will land beyond the near city wall. The base of the far wall is located at $(600, 0)$ which the rock hits if $(80 \cos\theta)t = 600 \;\Rightarrow\; t = \dfrac{15}{2\cos\theta}$ and $(80 \sin\theta)t - 4.9t^2 = 0 \;\Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$80\sin\theta \cdot \dfrac{15}{2\cos\theta} - 4.9\left(\dfrac{15}{2\cos\theta}\right)^2 = 0 \quad\Rightarrow\quad 600\tan\theta - 275.625\sec^2\theta = 0 \quad\Rightarrow$

$275.625\tan^2\theta - 600\tan\theta + 275.625 = 0$. Solutions are $\tan\theta \approx 0.658678, 1.51819 \quad\Rightarrow\quad \theta \approx 33.4^\circ, 56.6^\circ$. Thus the rock lands beyond the enclosed city ground for $33.4^\circ < \theta < 56.6^\circ$, and the angles that allow the rock to land on city ground are $13.0^\circ < \theta < 33.4^\circ, 56.6^\circ < \theta < 85.5^\circ$. If you consider that the rock can hit the far wall and bounce back into the city, we calculate the angles that cause the rock to hit the top of the wall at $(600, 15)$: $(80\cos\theta)t = 600 \quad\Rightarrow\quad t = \dfrac{15}{2\cos\theta}$ and $(80\sin\theta)t - 4.9t^2 = 15 \quad\Rightarrow\quad 600\tan\theta - 275.625\sec^2\theta = 15 \quad\Rightarrow\quad 275.625\tan^2\theta - 600\tan\theta + 290.625 = 0$. Solutions are $\tan\theta \approx 0.727506, 1.44936 \quad\Rightarrow\quad \theta \approx 36.0^\circ, 55.4^\circ$, so the catapult should be set with angle θ where $13.0^\circ < \theta < 36.0^\circ, 55.4^\circ < \theta < 85.5^\circ$.

30. If we place the projectile at the origin then, as in Example 5, $\mathbf{r}(t) = (v_0\cos\alpha)t\,\mathbf{i} + \left[(v_0\sin\alpha)t - \frac{1}{2}gt^2\right]\mathbf{j}$ and $\mathbf{v}(t) = (v_0\cos\alpha)\,\mathbf{i} + [(v_0\sin\alpha) - gt]\,\mathbf{j}$. The maximum height is reached when the vertical component of velocity is zero, so $(v_0\sin\alpha) - gt = 0 \quad\Rightarrow\quad t = \dfrac{v_0\sin\alpha}{g}$, and the corresponding height is the vertical component of the position function:

$$(v_0\sin\alpha)t - \tfrac{1}{2}gt^2 = (v_0\sin\alpha)\left(\frac{v_0\sin\alpha}{g}\right) - \tfrac{1}{2}g\left(\frac{v_0\sin\alpha}{g}\right)^2 = \frac{1}{2g}v_0^2\sin^2\alpha$$

Half that time is $t = \dfrac{v_0\sin\alpha}{2g}$, when the height of the projectile is

$$\begin{aligned}(v_0\sin\alpha)t - \tfrac{1}{2}gt^2 &= (v_0\sin\alpha)\left(\frac{v_0\sin\alpha}{2g}\right) - \tfrac{1}{2}g\left(\frac{v_0\sin\alpha}{2g}\right)^2 \\ &= \frac{1}{2g}v_0^2\sin^2\alpha - \frac{1}{8g}v_0^2\sin^2\alpha = \frac{3}{8g}v_0^2\sin^2\alpha = \frac{3}{4}\left(\frac{1}{2g}v_0^2\sin^2\alpha\right)\end{aligned}$$

or three-quarters of the maximum height.

31. Here $\mathbf{a}(t) = -4\,\mathbf{j} - 32\,\mathbf{k}$ so $\mathbf{v}(t) = -4t\,\mathbf{j} - 32t\,\mathbf{k} + \mathbf{v}_0 = -4t\,\mathbf{j} - 32t\,\mathbf{k} + 50\,\mathbf{i} + 80\,\mathbf{k} = 50\,\mathbf{i} - 4t\,\mathbf{j} + (80 - 32t)\,\mathbf{k}$ and $\mathbf{r}(t) = 50t\,\mathbf{i} - 2t^2\,\mathbf{j} + (80t - 16t^2)\,\mathbf{k}$ (note that $\mathbf{r}_0 = \mathbf{0}$). The ball lands when the z-component of $\mathbf{r}(t)$ is zero and $t > 0$: $80t - 16t^2 = 16t(5 - t) = 0 \quad\Rightarrow\quad t = 5$. The position of the ball then is $\mathbf{r}(5) = 50(5)\,\mathbf{i} - 2(5)^2\,\mathbf{j} + [80(5) - 16(5)^2]\,\mathbf{k} = 250\,\mathbf{i} - 50\,\mathbf{j}$ or equivalently the point $(250, -50, 0)$. This is a distance of $\sqrt{250^2 + (-50)^2 + 0^2} = \sqrt{65{,}000} \approx 255$ ft from the origin at an angle of $\tan^{-1}\left(\frac{50}{250}\right) \approx 11.3^\circ$ from the eastern direction toward the south. The speed of the ball is $|\mathbf{v}(5)| = |50\,\mathbf{i} - 20\,\mathbf{j} - 80\,\mathbf{k}| = \sqrt{50^2 + (-20)^2 + (-80)^2} = \sqrt{9300} \approx 96.4$ ft/s.

32. Place the ball at the origin and consider $\mathbf{j}$ to be pointing in the northward direction with $\mathbf{i}$ pointing east and $\mathbf{k}$ pointing upward. Force = mass × acceleration $\quad\Rightarrow\quad$ acceleration = force/mass, so the wind applies a constant acceleration of $4\text{ N}/0.8\text{ kg} = 5\text{ m/s}^2$ in the easterly direction. Combined with the acceleration due to gravity, the acceleration acting on the ball is $\mathbf{a}(t) = 5\,\mathbf{i} - 9.8\,\mathbf{k}$. Then $\mathbf{v}(t) = \int \mathbf{a}(t)\,dt = 5t\,\mathbf{i} - 9.8t\,\mathbf{k} + \mathbf{C}$ where $\mathbf{C}$ is a constant vector. We know $\mathbf{v}(0) = \mathbf{C} = -30\cos 30^\circ\,\mathbf{j} + 30\sin 30^\circ\,\mathbf{k} = -15\sqrt{3}\,\mathbf{j} + 15\,\mathbf{k} \quad\Rightarrow\quad \mathbf{C} = -15\sqrt{3}\,\mathbf{j} + 15\,\mathbf{k}$ and

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{v}(t) = 5t\,\mathbf{i} - 15\sqrt{3}\,\mathbf{j} + (15 - 9.8t)\,\mathbf{k}$. $\mathbf{r}(t) = \int \mathbf{v}(t)\,dt = 2.5t^2\,\mathbf{i} - 15\sqrt{3}\,t\,\mathbf{j} + \left(15t - 4.9t^2\right)\mathbf{k} + \mathbf{D}$ but $\mathbf{r}(0) = \mathbf{D} = \mathbf{0}$ so $\mathbf{r}(t) = 2.5t^2\,\mathbf{i} - 15\sqrt{3}\,t\,\mathbf{j} + \left(15t - 4.9t^2\right)\mathbf{k}$. The ball lands when $15t - 4.9t^2 = 0 \;\Rightarrow\; t = 0, t = 15/4.9 \approx 3.0612$ s, so the ball lands at approximately $\mathbf{r}(3.0612) \approx 23.43\,\mathbf{i} - 79.53\,\mathbf{j}$ which is 82.9 m away in the direction S 16.4°E. Its speed is approximately $|\mathbf{v}(3.0612)| \approx \left|15.306\,\mathbf{i} - 15\sqrt{3}\,\mathbf{j} - 15\,\mathbf{k}\right| \approx 33.68$ m/s.

33. (a) After t seconds, the boat will be $5t$ meters west of point A. The velocity of the water at that location is $\frac{3}{400}(5t)(40 - 5t)\,\mathbf{j}$. The velocity of the boat in still water is $5\,\mathbf{i}$, so the resultant velocity of the boat is $\mathbf{v}(t) = 5\,\mathbf{i} + \frac{3}{400}(5t)(40 - 5t)\,\mathbf{j} = 5\mathbf{i} + \left(\frac{3}{2}t - \frac{3}{16}t^2\right)\mathbf{j}$. Integrating, we obtain $\mathbf{r}(t) = 5t\,\mathbf{i} + \left(\frac{3}{4}t^2 - \frac{1}{16}t^3\right)\mathbf{j} + \mathbf{C}$. If we place the origin at A (and consider $\mathbf{j}$ to coincide with the northern direction) then $\mathbf{r}(0) = \mathbf{0} \;\Rightarrow\; \mathbf{C} = \mathbf{0}$ and we have $\mathbf{r}(t) = 5t\,\mathbf{i} + \left(\frac{3}{4}t^2 - \frac{1}{16}t^3\right)\mathbf{j}$. The boat reaches the east bank when $5t = 40$, that is, when $t = 8$ s, and it is located at $\mathbf{r}(8) = 5(8)\mathbf{i} + \left(\frac{3}{4}(8)^2 - \frac{1}{16}(8)^3\right)\mathbf{j} = 40\,\mathbf{i} + 16\,\mathbf{j}$. Thus the boat is 16 m downstream.

(b) Let α be the angle north of east that the boat heads. Then the velocity of the boat in still water is given by $5(\cos\alpha)\,\mathbf{i} + 5(\sin\alpha)\,\mathbf{j}$. At t seconds, the boat is $5(\cos\alpha)t$ meters from the west bank, at which point the velocity of the water is $\frac{3}{400}[5(\cos\alpha)t][40 - 5(\cos\alpha)t]\;\mathbf{j}$. The resultant velocity of the boat is given by $\mathbf{v}(t) = 5(\cos\alpha)\,\mathbf{i} + \left[5\sin\alpha + \frac{3}{400}(5t\cos\alpha)(40 - 5t\cos\alpha)\right]\mathbf{j} = (5\cos\alpha)\,\mathbf{i} + \left(5\sin\alpha + \frac{3}{2}t\cos\alpha - \frac{3}{16}t^2\cos^2\alpha\right)\mathbf{j}$. Integrating, $\mathbf{r}(t) = (5t\cos\alpha)\,\mathbf{i} + \left(5t\sin\alpha + \frac{3}{4}t^2\cos\alpha - \frac{1}{16}t^3\cos^2\alpha\right)\mathbf{j}$ (where we have again placed the origin at A). The boat will reach the east bank when $5t\cos\alpha = 40 \;\Rightarrow\; t = \dfrac{40}{5\cos\alpha} = \dfrac{8}{\cos\alpha}$.

In order to land at point $B(40, 0)$ we need $5t\sin\alpha + \frac{3}{4}t^2\cos\alpha - \frac{1}{16}t^3\cos^2\alpha = 0 \;\Rightarrow$

$$5\left(\frac{8}{\cos\alpha}\right)\sin\alpha + \tfrac{3}{4}\left(\frac{8}{\cos\alpha}\right)^2\cos\alpha - \tfrac{1}{16}\left(\frac{8}{\cos\alpha}\right)^3\cos^2\alpha = 0 \;\Rightarrow\; \frac{1}{\cos\alpha}\,(40\sin\alpha + 48 - 32) = 0 \;\Rightarrow$$

$40\sin\alpha + 16 = 0 \;\Rightarrow\; \sin\alpha = -\frac{2}{5}$. Thus $\alpha = \sin^{-1}\left(-\frac{2}{5}\right) \approx -23.6°$, so the boat should head 23.6° south of east (upstream). The path does seem realistic. The boat initially heads upstream to counteract the effect of the current. Near the center of the river, the current is stronger and the boat is pushed downstream. When the boat nears the eastern bank, the current is slower and the boat is able to progress upstream to arrive at point B.

34. As in Exercise 33(b), let α be the angle north of east that the boat heads, so the velocity of the boat in still water is given by $5(\cos\alpha)\,\mathbf{i} + 5(\sin\alpha)\,\mathbf{j}$. At t seconds, the boat is $5(\cos\alpha)t$ meters from the west bank, at which point the velocity of the water is $3\sin(\pi x/40)\,\mathbf{j} = 3\sin[\pi \cdot 5(\cos\alpha)t/40]\,\mathbf{j} = 3\sin\left(\frac{\pi}{8}t\cos\alpha\right)\mathbf{j}$. The resultant velocity of the boat

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

then is given by $\mathbf{v}(t) = 5(\cos\alpha)\,\mathbf{i} + \left[5\sin\alpha + 3\sin\left(\frac{\pi}{8}t\cos\alpha\right)\right]\mathbf{j}$. Integrating,

$$\mathbf{r}(t) = (5t\cos\alpha)\,\mathbf{i} + \left[5t\sin\alpha - \frac{24}{\pi\cos\alpha}\cos\left(\tfrac{\pi}{8}t\cos\alpha\right)\right]\mathbf{j} + \mathbf{C}.$$

If we place the origin at A then $\mathbf{r}(0) = \mathbf{0} \;\Rightarrow\; -\dfrac{24}{\pi\cos\alpha}\,\mathbf{j} + \mathbf{C} = \mathbf{0} \;\Rightarrow\; \mathbf{C} = \dfrac{24}{\pi\cos\alpha}\,\mathbf{j}$ and

$\mathbf{r}(t) = (5t\cos\alpha)\,\mathbf{i} + \left[5t\sin\alpha - \dfrac{24}{\pi\cos\alpha}\cos\left(\tfrac{\pi}{8}t\cos\alpha\right) + \dfrac{24}{\pi\cos\alpha}\right]\mathbf{j}$. The boat will reach the east bank when

$5t\cos\alpha = 40 \;\Rightarrow\; t = \dfrac{8}{\cos\alpha}$. In order to land at point $B(40, 0)$ we need

$$5t\sin\alpha - \frac{24}{\pi\cos\alpha}\cos\left(\tfrac{\pi}{8}t\cos\alpha\right) + \frac{24}{\pi\cos\alpha} = 0 \;\Rightarrow$$

$$5\left(\frac{8}{\cos\alpha}\right)\sin\alpha - \frac{24}{\pi\cos\alpha}\cos\left[\frac{\pi}{8}\left(\frac{8}{\cos\alpha}\right)\cos\alpha\right] + \frac{24}{\pi\cos\alpha} = 0 \;\Rightarrow\; \frac{1}{\cos\alpha}\left(40\sin\alpha - \frac{24}{\pi}\cos\pi + \frac{24}{\pi}\right) = 0 \;\Rightarrow$$

$40\sin\alpha + \dfrac{48}{\pi} = 0 \;\Rightarrow\; \sin\alpha = -\dfrac{6}{5\pi}$. Thus $\alpha = \sin^{-1}\left(-\dfrac{6}{5\pi}\right) \approx -22.5^\circ$, so the boat should head 22.5° south of east.

35. If $\mathbf{r}'(t) = \mathbf{c} \times \mathbf{r}(t)$, then $\mathbf{r}'(t)$ is perpendicular to both $\mathbf{c}$ and $\mathbf{r}(t)$. Remember that $\mathbf{r}'(t)$ points in the direction of motion, so if $\mathbf{r}'(t)$ is always perpendicular to $\mathbf{c}$, the path of the particle must lie in a plane perpendicular to $\mathbf{c}$. But $\mathbf{r}'(t)$ is also perpendicular to the position vector $\mathbf{r}(t)$ which confines the path to a sphere centered at the origin. Considering both restrictions, the path must be contained in a circle that lies in a plane perpendicular to $\mathbf{c}$, and the circle is centered on a line through the origin in the direction of $\mathbf{c}$.

36. (a) From Equation 7 we have $\mathbf{a} = v'\mathbf{T} + \kappa v^2\mathbf{N}$. If a particle moves along a straight line, then $\kappa = 0$ [see Section 13.3], so the acceleration vector becomes $\mathbf{a} = v'\mathbf{T}$. Because the acceleration vector is a scalar multiple of the unit tangent vector, it is parallel to the tangent vector.

(b) If the speed of the particle is constant, then $v' = 0$ and Equation 7 gives $\mathbf{a} = \kappa v^2\mathbf{N}$. Thus the acceleration vector is parallel to the unit normal vector (which is perpendicular to the tangent vector and points in the direction that the curve is turning).

37. $\mathbf{r}(t) = (t^2 + 1)\,\mathbf{i} + t^3\,\mathbf{j} \;\Rightarrow\; \mathbf{r}'(t) = 2t\,\mathbf{i} + 3t^2\,\mathbf{j}$,

$|\mathbf{r}'(t)| = \sqrt{(2t)^2 + (3t^2)^2} = \sqrt{4t^2 + 9t^4} = t\sqrt{4 + 9t^2}$ [since $t \geq 0$], $\mathbf{r}''(t) = 2\,\mathbf{i} + 6t\,\mathbf{j}$, $\mathbf{r}'(t) \times \mathbf{r}''(t) = 6t^2\,\mathbf{k}$.

Then Equation 9 gives $a_T = \dfrac{\mathbf{r}'(t)\cdot\mathbf{r}''(t)}{|\mathbf{r}'(t)|} = \dfrac{(2t)(2) + (3t^2)(6t)}{t\sqrt{4+9t^2}} = \dfrac{4t + 18t^3}{t\sqrt{4+9t^2}} = \dfrac{4 + 18t^2}{\sqrt{4+9t^2}}$

$$\left[\begin{array}{l} \text{or by Equation 8,}\quad a_T = v' = \dfrac{d}{dt}\left[t\sqrt{4+9t^2}\right] = t\cdot\tfrac{1}{2}\left(4+9t^2\right)^{-1/2}(18t) + \left(4+9t^2\right)^{1/2}\cdot 1 \\ \qquad\qquad = \left(4+9t^2\right)^{-1/2}\left(9t^2 + 4 + 9t^2\right) = (4 + 18t^2)/\sqrt{4+9t^2} \end{array}\right]$$

and Equation 10 gives $a_N = \dfrac{|\mathbf{r}'(t) \times \mathbf{r}''(t)|}{|\mathbf{r}'(t)|} = \dfrac{6t^2}{t\sqrt{4+9t^2}} = \dfrac{6t}{\sqrt{4+9t^2}}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. $\mathbf{r}(t) = 2t^2\,\mathbf{i} + \left(\frac{2}{3}t^3 - 2t\right)\mathbf{j} \quad\Rightarrow\quad \mathbf{r}'(t) = 4t\,\mathbf{i} + (2t^2 - 2)\,\mathbf{j}$,

$|\mathbf{r}'(t)| = \sqrt{16t^2 + (2t^2-2)^2} = \sqrt{4t^4 + 8t^2 + 4} = \sqrt{4\,(t^2+1)^2} = 2(t^2+1)$,

$\mathbf{r}''(t) = 4\,\mathbf{i} + 4t\,\mathbf{j}$, $\mathbf{r}'(t) \times \mathbf{r}''(t) = (8t^2+8)\,\mathbf{k}$. Then Equation 9 gives

$a_T = \dfrac{\mathbf{r}'(t)\cdot\mathbf{r}''(t)}{|\mathbf{r}'(t)|} = \dfrac{(4t)(4) + (2t^2-2)(4t)}{2(t^2+1)} = \dfrac{8t(t^2+1)}{2(t^2+1)} = 4t \quad \left[\text{or by Equation 8, } a_T = v' = \dfrac{d}{dt}\left[2(t^2+1)\right] = 4t\right]$

and Equation 10 gives $a_N = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|} = \dfrac{8(t^2+1)}{2(t^2+1)} = 4$.

39. $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j} + t\,\mathbf{k} \quad\Rightarrow\quad \mathbf{r}'(t) = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j} + \mathbf{k}$, $|\mathbf{r}'(t)| = \sqrt{\sin^2 t + \cos^2 t + 1} = \sqrt{2}$,

$\mathbf{r}''(t) = -\cos t\,\mathbf{i} - \sin t\,\mathbf{j}$, $\mathbf{r}'(t)\times\mathbf{r}''(t) = \sin t\,\mathbf{i} - \cos t\,\mathbf{j} + \mathbf{k}$.

Then $a_T = \dfrac{\mathbf{r}'(t)\cdot\mathbf{r}''(t)}{|\mathbf{r}'(t)|} = \dfrac{\sin t\,\cos t - \sin t\,\cos t}{\sqrt{2}} = 0$ and $a_N = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|} = \dfrac{\sqrt{\sin^2 t + \cos^2 t + 1}}{\sqrt{2}} = \dfrac{\sqrt{2}}{\sqrt{2}} = 1$.

40. $\mathbf{r}(t) = t\,\mathbf{i} + 2e^t\,\mathbf{j} + e^{2t}\,\mathbf{k} \quad\Rightarrow\quad \mathbf{r}'(t) = \mathbf{i} + 2e^t\,\mathbf{j} + 2e^{2t}\,\mathbf{k}$, $|\mathbf{r}'(t)| = \sqrt{1 + 4e^{2t} + 4e^{4t}} = \sqrt{(1+2e^{2t})^2} = 1 + 2e^{2t}$,

$\mathbf{r}''(t) = 2e^t\,\mathbf{j} + 4e^{2t}\,\mathbf{k}$, $\mathbf{r}'(t)\times\mathbf{r}''(t) = 4e^{3t}\,\mathbf{i} - 4e^{2t}\,\mathbf{j} + 2e^t\,\mathbf{k}$,

$|\mathbf{r}'(t)\times\mathbf{r}''(t)| = \sqrt{16e^{6t} + 16e^{4t} + 4e^{2t}} = \sqrt{4e^{2t}(2e^{2t}+1)^2} = 2e^t(2e^{2t}+1)$. Then

$a_T = \dfrac{\mathbf{r}'(t)\cdot\mathbf{r}''(t)}{|\mathbf{r}'(t)|} = \dfrac{4e^{2t} + 8e^{4t}}{1+2e^{2t}} = \dfrac{4e^{2t}(1+2e^{2t})}{1+2e^{2t}} = 4e^{2t}$ and $a_N = \dfrac{|\mathbf{r}'(t)\times\mathbf{r}''(t)|}{|\mathbf{r}'(t)|} = \dfrac{2e^t(2e^{2t}+1)}{1+2e^{2t}} = 2e^t$.

41. $\mathbf{r}(t) = \ln t\,\mathbf{i} + (t^2+3t)\,\mathbf{j} + 4\sqrt{t}\,\mathbf{k} \quad\Rightarrow\quad \mathbf{r}'(t) = (1/t)\,\mathbf{i} + (2t+3)\,\mathbf{j} + (2/\sqrt{t}\,)\,\mathbf{k} \quad\Rightarrow$

$\mathbf{r}''(t) = (-1/t^2)\,\mathbf{i} + 2\,\mathbf{j} - (1/t^{3/2})\,\mathbf{k}$. The point $(0,4,4)$ corresponds to $t=1$, where

$\mathbf{r}'(1) = \mathbf{i} + 5\,\mathbf{j} + 2\,\mathbf{k}$, $\mathbf{r}''(1) = -\mathbf{i} + 2\,\mathbf{j} - \mathbf{k}$, and $\mathbf{r}'(1)\times\mathbf{r}''(1) = -9\,\mathbf{i} - \mathbf{j} + 7\,\mathbf{k}$. Thus at the point $(0,4,4)$,

$a_T = \dfrac{\mathbf{r}'(1)\cdot\mathbf{r}''(1)}{|\mathbf{r}'(1)|} = \dfrac{-1+10-2}{\sqrt{1+25+4}} = \dfrac{7}{\sqrt{30}}$ and $a_N = \dfrac{|\mathbf{r}'(1)\times\mathbf{r}''(1)|}{|\mathbf{r}'(1)|} = \dfrac{\sqrt{81+1+49}}{\sqrt{30}} = \sqrt{\dfrac{131}{30}}$.

42. $\mathbf{r}(t) = t^{-1}\,\mathbf{i} + t^{-2}\,\mathbf{j} + t^{-3}\,\mathbf{k} \quad\Rightarrow\quad \mathbf{r}'(t) = -t^{-2}\,\mathbf{i} - 2t^{-3}\,\mathbf{j} - 3t^{-4}\,\mathbf{k} \quad\Rightarrow\quad \mathbf{r}''(t) = 2t^{-3}\,\mathbf{i} + 6t^{-4}\,\mathbf{j} + 12t^{-5}\,\mathbf{k}$. The point $(1,1,1)$ corresponds to $t=1$, where $\mathbf{r}'(1) = -\mathbf{i} - 2\,\mathbf{j} - 3\,\mathbf{k}$, $\mathbf{r}''(1) = 2\,\mathbf{i} + 6\,\mathbf{j} + 12\,\mathbf{k}$, and

$\mathbf{r}'(1)\times\mathbf{r}''(1) = -6\,\mathbf{i} + 6\,\mathbf{j} - 2\,\mathbf{k}$. Thus at the point $(1,1,1)$, $a_T = \dfrac{\mathbf{r}'(1)\cdot\mathbf{r}''(1)}{|\mathbf{r}'(1)|} = \dfrac{-2-12-36}{\sqrt{1+4+9}} = -\dfrac{50}{\sqrt{14}}$ and

$a_N = \dfrac{|\mathbf{r}'(1)\times\mathbf{r}''(1)|}{|\mathbf{r}'(1)|} = \dfrac{\sqrt{36+36+4}}{\sqrt{14}} = \sqrt{\dfrac{76}{14}} = \sqrt{\dfrac{38}{7}}$.

43. The tangential component of $\mathbf{a}$ is the length of the projection of $\mathbf{a}$ onto $\mathbf{T}$, so we sketch the scalar projection of $\mathbf{a}$ in the tangential direction to the curve and estimate its length to be 4.5 (using the fact that $\mathbf{a}$ has length 10 as a guide). Similarly, the normal component of $\mathbf{a}$ is the length of the projection of $\mathbf{a}$ onto $\mathbf{N}$, so we sketch the scalar projection of $\mathbf{a}$ in the normal direction to the curve and estimate its length to be 9.0. Thus $a_T \approx 4.5$ cm/s^2 and $a_N \approx 9.0$ cm/s^2.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. $\mathbf{L}(t) = m\,\mathbf{r}(t) \times \mathbf{v}(t) \quad \Rightarrow$

$$\begin{aligned}\mathbf{L}'(t) &= m[\mathbf{r}'(t) \times \mathbf{v}(t) + \mathbf{r}(t) \times \mathbf{v}'(t)] \qquad \text{[by Formula 5 of Theorem 13.2.3]}\\ &= m[\mathbf{v}(t) \times \mathbf{v}(t) + \mathbf{r}(t) \times \mathbf{v}'(t)] = m[\mathbf{0} + \mathbf{r}(t) \times \mathbf{a}(t)] = \boldsymbol{\tau}(t)\end{aligned}$$

So if the torque is always $\mathbf{0}$, then $\mathbf{L}'(t) = \mathbf{0}$ for all t, and so $\mathbf{L}(t)$ is constant.

45. If the engines are turned off at time t, then the spacecraft will continue to travel in the direction of $\mathbf{v}(t)$, so we need a t such that for some scalar $s > 0$, $\mathbf{r}(t) + s\,\mathbf{v}(t) = \langle 6, 4, 9\rangle$. $\mathbf{v}(t) = \mathbf{r}'(t) = \mathbf{i} + \dfrac{1}{t}\,\mathbf{j} + \dfrac{8t}{(t^2+1)^2}\,\mathbf{k} \quad \Rightarrow$

$\mathbf{r}(t) + s\,\mathbf{v}(t) = \left\langle 3 + t + s, 2 + \ln t + \dfrac{s}{t}, 7 - \dfrac{4}{t^2+1} + \dfrac{8st}{(t^2+1)^2} \right\rangle \quad \Rightarrow \quad 3 + t + s = 6 \quad \Rightarrow \quad s = 3 - t,$

so $7 - \dfrac{4}{t^2+1} + \dfrac{8(3-t)t}{(t^2+1)^2} = 9 \quad \Leftrightarrow \quad \dfrac{24t - 12t^2 - 4}{(t^2+1)^2} = 2 \quad \Leftrightarrow \quad t^4 + 8t^2 - 12t + 3 = 0.$

It is easily seen that $t = 1$ is a root of this polynomial. Also $2 + \ln 1 + \dfrac{3-1}{1} = 4$, so $t = 1$ is the desired solution.

46. (a) $m\,\dfrac{d\mathbf{v}}{dt} = \dfrac{dm}{dt}\,\mathbf{v}_e \quad \Leftrightarrow \quad \dfrac{d\mathbf{v}}{dt} = \dfrac{1}{m}\dfrac{dm}{dt}\,\mathbf{v}_e$. Integrating both sides of this equation with respect to t gives

$$\int_0^t \frac{d\mathbf{v}}{du}\,du = \mathbf{v}_e \int_0^t \frac{1}{m}\frac{dm}{du}\,du \quad \Rightarrow \quad \int_{\mathbf{v}(0)}^{\mathbf{v}(t)} d\mathbf{v} = \mathbf{v}_e \int_{m(0)}^{m(t)} \frac{dm}{m} \quad \text{[Substitution Rule]} \quad \Rightarrow$$

$$\mathbf{v}(t) - \mathbf{v}(0) = \mathbf{v}_e \ln\frac{m(t)}{m(0)} \quad \Rightarrow \quad \mathbf{v}(t) = \mathbf{v}(0) - \mathbf{v}_e \ln\frac{m(0)}{m(t)}.$$

(b) $|\mathbf{v}(t)| = 2\,|\mathbf{v}_e|$, and $|\mathbf{v}(0)| = 0$. Therefore, by part (a), $2\,|\mathbf{v}_e| = \left| -\ln\left(\dfrac{m(0)}{m(t)}\right)\mathbf{v}_e \right| \quad \Rightarrow$

$2\,|\mathbf{v}_e| = \ln\left(\dfrac{m(0)}{m(t)}\right)|\mathbf{v}_e|$. $\left[\textit{Note: } m(0) > m(t) \text{ so that } \ln\left(\dfrac{m(0)}{m(t)}\right) > 0\right] \quad \Rightarrow \quad m(t) = e^{-2}m(0).$

Thus $\dfrac{m(0) - e^{-2}m(0)}{m(0)} = 1 - e^{-2}$ is the fraction of the initial mass that is burned as fuel.

APPLIED PROJECT Kepler's Laws

1. With $\mathbf{r} = (r\cos\theta)\,\mathbf{i} + (r\sin\theta)\,\mathbf{j}$ and $\mathbf{h} = \alpha\,\mathbf{k}$ where $\alpha > 0$,

(a) $\mathbf{h} = \mathbf{r} \times \mathbf{r}' = [(r\cos\theta)\,\mathbf{i} + (r\sin\theta)\,\mathbf{j}] \times \left[\left(r'\cos\theta - r\sin\theta\,\dfrac{d\theta}{dt}\right)\mathbf{i} + \left(r'\sin\theta + r\cos\theta\,\dfrac{d\theta}{dt}\right)\mathbf{j}\right]$

$$= \left[rr'\cos\theta\sin\theta + r^2\cos^2\theta\,\frac{d\theta}{dt} - rr'\cos\theta\sin\theta + r^2\sin^2\theta\,\frac{d\theta}{dt}\right]\mathbf{k} = r^2\frac{d\theta}{dt}\,\mathbf{k}$$

(b) Since $\mathbf{h} = \alpha\,\mathbf{k}$, $\alpha > 0$, $\alpha = |\mathbf{h}|$. But by part (a), $\alpha = |\mathbf{h}| = r^2\,(d\theta/dt)$.

(c) $A(t) = \frac{1}{2}\int_{\theta_0}^{\theta} |\mathbf{r}|^2\,d\theta = \frac{1}{2}\int_{t_0}^{t} r^2\,(d\theta/dt)\,dt$ in polar coordinates. Thus, by the Fundamental Theorem of Calculus,

$$\frac{dA}{dt} = \frac{r^2}{2}\frac{d\theta}{dt}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $\dfrac{dA}{dt} = \dfrac{r^2}{2}\dfrac{d\theta}{dt} = \dfrac{h}{2} =$ constant since $\mathbf{h}$ is a constant vector and $h = |\mathbf{h}|$.

2. (a) Since $dA/dt = \frac{1}{2}h$, a constant, $A(t) = \frac{1}{2}ht + c_1$. But $A(0) = 0$, so $A(t) = \frac{1}{2}ht$. But $A(T) =$ area of the ellipse $= \pi ab$ and $A(T) = \frac{1}{2}hT$, so $T = 2\pi ab/h$.

(b) $h^2/(GM) = ed$ where e is the eccentricity of the ellipse. But $a = ed/(1-e^2)$ or $ed = a(1-e^2)$ and $1 - e^2 = b^2/a^2$. Hence $h^2/(GM) = ed = b^2/a$.

(c) $T^2 = \dfrac{4\pi a^2 b^2}{h^2} = 4\pi^2 a^2 b^2 \dfrac{a}{GMb^2} = \dfrac{4\pi^2}{GM}a^3$.

3. From Problem 2, $T^2 = \dfrac{4\pi^2}{GM}a^3$. $T \approx 365.25$ days $\times\, 24 \cdot 60^2 \dfrac{\text{seconds}}{\text{day}} \approx 3.1558 \times 10^7$ seconds. Therefore

$a^3 = \dfrac{GMT^2}{4\pi^2} \approx \dfrac{(6.67 \times 10^{-11})(1.99 \times 10^{30})(3.1558 \times 10^7)^2}{4\pi^2} \approx 3.348 \times 10^{33}\ \text{m}^3 \quad \Rightarrow \quad a \approx 1.496 \times 10^{11}$ m. Thus, the length of the major axis of the earth's orbit (that is, $2a$) is approximately 2.99×10^{11} m $= 2.99 \times 10^8$ km.

4. We can adapt the equation $T^2 = \dfrac{4\pi^2}{GM}a^3$ from Problem 2(c) with the earth at the center of the system, so T is the period of the satellite's orbit about the earth, M is the mass of the earth, and a is the length of the semimajor axis of the satellite's orbit (measured from the earth's center). Since we want the satellite to remain fixed above a particular point on the earth's equator, T must coincide with the period of the earth's own rotation, so $T = 24$ h $= 86{,}400$ s. The mass of the earth is

$M = 5.98 \times 10^{24}$ kg, so $a = \left(\dfrac{T^2 GM}{4\pi^2}\right)^{1/3} \approx \left[\dfrac{(86{,}400)^2(6.67 \times 10^{-11})(5.98 \times 10^{24})}{4\pi^2}\right]^{1/3} \approx 4.23 \times 10^7$ m. If we assume a circular orbit, the radius of the orbit is a, and since the radius of the earth is 6.37×10^6 m, the required altitude above the earth's surface for the satellite is $4.23 \times 10^7 - 6.37 \times 10^6 \approx 3.59 \times 10^7$ m, or 35,900 km.

13 Review

TRUE-FALSE QUIZ

1. True. If we reparametrize the curve by replacing $u = t^3$, we have $\mathbf{r}(u) = u\,\mathbf{i} + 2u\,\mathbf{j} + 3u\,\mathbf{k}$, which is a line through the origin with direction vector $\mathbf{i} + 2\,\mathbf{j} + 3\,\mathbf{k}$.

2. True. Parametric equations for the curve are $x = 0$, $y = t^2$, $z = 4t$, and since $t = z/4$ we have $y = t^2 = (z/4)^2$ or $y = \frac{1}{16}z^2$, $x = 0$. This is an equation of a parabola in the yz-plane.

3. False. The vector function $\mathbf{r}(t) = \langle 2t, 3 - t, 0\rangle$ represents a line, but the line does not pass through the origin; the x-component is 0 only for $t = 0$, which corresponds to the point $(0, 3, 0)$, not $(0, 0, 0)$.

4. True. See Theorem 13.2.2.

5. False. By Formula 5 of Theorem 13.2.3, $\dfrac{d}{dt}[\mathbf{u}(t) \times \mathbf{v}(t)] = \mathbf{u}'(t) \times \mathbf{v}(t) + \mathbf{u}(t) \times \mathbf{v}'(t)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. False. For example, let $\mathbf{r}(t) = \langle \cos t, \sin t \rangle$. Then $|\mathbf{r}(t)| = \sqrt{\cos^2 t + \sin^2 t} = 1 \quad \Rightarrow \quad \dfrac{d}{dt}\,|\mathbf{r}(t)| = 0$, but

$|\mathbf{r}'(t)| = |\langle -\sin t, \cos t \rangle| = \sqrt{(-\sin t)^2 + \cos^2 t} = 1$.

7. False. κ is the magnitude of the rate of change of the unit tangent vector $\mathbf{T}$ with respect to arc length s, not with respect to t.

8. False. The binormal vector, by the definition given in Section 13.3, is $\mathbf{B}(t) = \mathbf{T}(t) \times \mathbf{N}(t) = -\,[\mathbf{N}(t) \times \mathbf{T}(t)]$.

9. True. At an inflection point where f is twice continuously differentiable we must have $f''(x) = 0$, and by Equation 13.3.11, the curvature is 0 there.

10. True. From Equation 13.3.9 , $\kappa(t) = 0 \quad \Leftrightarrow \quad |\mathbf{T}'(t)| = 0 \quad \Leftrightarrow \quad \mathbf{T}'(t) = \mathbf{0}$ for all t. But then $\mathbf{T}(t) = \mathbf{C}$, a constant vector, which is true only for a straight line.

11. False. If $\mathbf{r}(t)$ is the position of a moving particle at time t and $|\mathbf{r}(t)| = 1$ then the particle lies on the unit circle or the unit sphere, but this does not mean that the speed $|\mathbf{r}'(t)|$ must be constant. As a counterexample, let $\mathbf{r}(t) = \left\langle t, \sqrt{1-t^2} \right\rangle$, then $\mathbf{r}'(t) = \left\langle 1, -t/\sqrt{1-t^2} \right\rangle$ and $|\mathbf{r}(t)| = \sqrt{t^2 + 1 - t^2} = 1$ but $|\mathbf{r}'(t)| = \sqrt{1 + t^2/(1-t^2)} = 1/\sqrt{1-t^2}$ which is not constant.

12. True. See Theorem 13.2.4.

13. True. See the discussion preceding Example 8 in Section 13.3.

14. False. For example, $\mathbf{r}_1(t) = \langle t, t \rangle$ and $\mathbf{r}_2(t) = \langle 2t, 2t \rangle$ both represent the same plane curve (the line $y = x$), but the tangent vector $\mathbf{r}_1'(t) = \langle 1, 1 \rangle$ for all t, while $\mathbf{r}_2'(t) = \langle 2, 2 \rangle$. In fact, different parametrizations give parallel tangent vectors at a point, but their magnitudes may differ.

15. True. The projection in the xz-plane is given by $\mathbf{r}(t) = \langle \cos 2t, 0, \sin 2t \rangle$. Since $x^2 + z^2 = (\cos 2t)^2 + (\sin 2t)^2 = 1$, the projection onto the xz-plane is a circle with radius 1.

16. True. Note that the direction vector for both lines is $\mathbf{d} = \langle 1, 2, 1 \rangle$, so the lines are parallel. Further, the point $(0, 0, 1)$ is contained on both lines for $t = 0$ and $t = 1$, respectively. The lines are parallel and have a point in common; therefore, they are the same line.

EXERCISES

1. (a) The corresponding parametric equations for the curve are $x = t$, $y = \cos \pi t,\ z = \sin \pi t$. Since $y^2 + z^2 = 1$, the curve is contained in a circular cylinder with axis the x-axis. Since $x = t$, the curve is a helix.

(b) $\mathbf{r}(t) = t\,\mathbf{i} + \cos \pi t\,\mathbf{j} + \sin \pi t\,\mathbf{k} \quad \Rightarrow$

$\mathbf{r}'(t) = \mathbf{i} - \pi \sin \pi t\,\mathbf{j} + \pi \cos \pi t\,\mathbf{k} \quad \Rightarrow$

$\mathbf{r}''(t) = -\pi^2 \cos \pi t\,\mathbf{j} - \pi^2 \sin \pi t\,\mathbf{k}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. (a) The expressions $\sqrt{2-t}$, $(e^t-1)/t$, and $\ln(t+1)$ are all defined when $2-t \geq 0 \quad \Rightarrow \quad t \leq 2, t \neq 0$, and $t+1>0 \quad \Rightarrow \quad t>-1$. Thus the domain of $\mathbf{r}$ is $(-1,0) \cup (0,2]$.

(b) $\lim\limits_{t\to 0} \mathbf{r}(t) = \left\langle \lim\limits_{t\to 0} \sqrt{2-t}, \lim\limits_{t\to 0} \dfrac{e^t-1}{t}, \lim\limits_{t\to 0} \ln(t+1) \right\rangle = \left\langle \sqrt{2-0}, \lim\limits_{t\to 0} \dfrac{e^t}{1}, \ln(0+1) \right\rangle$

$= \langle \sqrt{2}, 1, 0 \rangle$ [using l'Hospital's Rule in the y-component]

(c) $\mathbf{r}'(t) = \left\langle \dfrac{d}{dt}\sqrt{2-t}, \dfrac{d}{dt}\dfrac{e^t-1}{t}, \dfrac{d}{dt}\ln(t+1) \right\rangle = \left\langle -\dfrac{1}{2\sqrt{2-t}}, \dfrac{te^t-e^t+1}{t^2}, \dfrac{1}{t+1} \right\rangle$

3. The projection of the curve C of intersection onto the xy-plane is the circle $x^2+y^2=16, z=0$. So we can write $x=4\cos t,\ y=4\sin t,\ 0 \leq t \leq 2\pi$. From the equation of the plane, we have $z=5-x=5-4\cos t$, so parametric equations for C are $x=4\cos t,\ y=4\sin t,\ z=5-4\cos t, 0 \leq t \leq 2\pi$, and the corresponding vector function is $\mathbf{r}(t) = 4\cos t\,\mathbf{i} + 4\sin t\,\mathbf{j} + (5-4\cos t)\,\mathbf{k}, 0 \leq t \leq 2\pi$.

4. The curve is given by $\mathbf{r}(t) = \langle 2\sin t, 2\sin 2t, 2\sin 3t \rangle$, so $\mathbf{r}'(t) = \langle 2\cos t, 4\cos 2t, 6\cos 3t \rangle$. The point $\left(1, \sqrt{3}, 2\right)$ corresponds to $t = \frac{\pi}{6}$ (or $\frac{\pi}{6} + 2k\pi$, k an integer), so the tangent vector there is $\mathbf{r}'(\frac{\pi}{6}) = \left\langle \sqrt{3}, 2, 0 \right\rangle$. Then the tangent line has direction vector $\left\langle \sqrt{3}, 2, 0 \right\rangle$ and includes the point $\left(1, \sqrt{3}, 2\right)$, so parametric equations are $x = 1+\sqrt{3}\,t$, $y = \sqrt{3}+2t$, $z=2$.

5. $\int_0^1 (t^2\,\mathbf{i} + t\cos\,\pi t\,\mathbf{j} + \sin\,\pi t\,\mathbf{k})\,dt = \left(\int_0^1 t^2\,dt\right)\mathbf{i} + \left(\int_0^1 t\cos\,\pi t\,dt\right)\mathbf{j} + \left(\int_0^1 \sin\,\pi t\,dt\right)\mathbf{k}$

$= \left[\frac{1}{3}t^3\right]_0^1 \mathbf{i} + \left(\left[\frac{t}{\pi}\sin\,\pi t\right]_0^1 - \int_0^1 \frac{1}{\pi}\sin\,\pi t\,dt\right)\mathbf{j} + \left[-\frac{1}{\pi}\cos\,\pi t\right]_0^1 \mathbf{k}$

$= \frac{1}{3}\,\mathbf{i} + \left[\frac{1}{\pi^2}\cos\,\pi t\right]_0^1 \mathbf{j} + \frac{2}{\pi}\,\mathbf{k} = \frac{1}{3}\,\mathbf{i} - \frac{2}{\pi^2}\,\mathbf{j} + \frac{2}{\pi}\,\mathbf{k}$

where we integrated by parts in the y-component.

6. (a) C intersects the xz-plane where $y=0 \quad \Rightarrow \quad 2t-1=0 \quad \Rightarrow \quad t=\frac{1}{2}$, so the point is $\left(2-\left(\frac{1}{2}\right)^3, 0, \ln\frac{1}{2}\right) = \left(\frac{15}{8}, 0, -\ln 2\right)$.

(b) The curve is given by $\mathbf{r}(t) = \left\langle 2-t^3, 2t-1, \ln t \right\rangle$, so $\mathbf{r}'(t) = \left\langle -3t^2, 2, 1/t \right\rangle$. The point $(1,1,0)$ corresponds to $t=1$, so the tangent vector there is $\mathbf{r}'(1) = \langle -3, 2, 1 \rangle$. Then the tangent line has direction vector $\langle -3, 2, 1 \rangle$ and includes the point $(1,1,0)$, so parametric equations are $x = 1-3t$, $y = 1+2t$, $z=t$.

(c) The normal plane has normal vector $\mathbf{r}'(1) = \langle -3, 2, 1 \rangle$ and equation $-3(x-1)+2(y-1)+z=0$ or $3x-2y-z=1$.

7. $\mathbf{r}(t) = \left\langle t^2, t^3, t^4 \right\rangle \quad \Rightarrow \quad \mathbf{r}'(t) = \left\langle 2t, 3t^2, 4t^3 \right\rangle \quad \Rightarrow \quad |\mathbf{r}'(t)| = \sqrt{4t^2+9t^4+16t^6}$ and $L = \int_0^3 |\mathbf{r}'(t)|\,dt = \int_0^3 \sqrt{4t^2+9t^4+16t^6}\,dt$. Using Simpson's Rule with $f(t) = \sqrt{4t^2+9t^4+16t^6}$ and $n=6$ we

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

have $\Delta t = \frac{3-0}{6} = \frac{1}{2}$ and

$$\begin{aligned} L &\approx \tfrac{\Delta t}{3}\left[f(0) + 4f\left(\tfrac{1}{2}\right) + 2f(1) + 4f\left(\tfrac{3}{2}\right) + 2f(2) + 4f\left(\tfrac{5}{2}\right) + f(3)\right] \\ &= \tfrac{1}{6}\left[\sqrt{0+0+0} + 4\cdot\sqrt{4\left(\tfrac{1}{2}\right)^2 + 9\left(\tfrac{1}{2}\right)^4 + 16\left(\tfrac{1}{2}\right)^6} + 2\cdot\sqrt{4(1)^2 + 9(1)^4 + 16(1)^6}\right. \\ &\qquad + 4\cdot\sqrt{4\left(\tfrac{3}{2}\right)^2 + 9\left(\tfrac{3}{2}\right)^4 + 16\left(\tfrac{3}{2}\right)^6} + 2\cdot\sqrt{4(2)^2 + 9(2)^4 + 16(2)^6} \\ &\qquad \left. + 4\cdot\sqrt{4\left(\tfrac{5}{2}\right)^2 + 9\left(\tfrac{5}{2}\right)^4 + 16\left(\tfrac{5}{2}\right)^6} + \sqrt{4(3)^2 + 9(3)^4 + 16(3)^6}\right] \\ &\approx 86.631 \end{aligned}$$

8. $\mathbf{r}(t) = \left\langle 2t^{3/2}, \cos 2t, \sin 2t\right\rangle \quad\Rightarrow\quad \mathbf{r}'(t) = \left\langle 3t^{1/2}, -2\sin 2t, 2\cos 2t\right\rangle \quad\Rightarrow$

$|\mathbf{r}'(t)| = \sqrt{9t + 4(\sin^2 2t + \cos^2 2t)} = \sqrt{9t+4}$.

Thus, $L = \int_0^1 \sqrt{9t+4}\,dt = \int_4^{13} \frac{1}{9}u^{1/2}\,du = \frac{1}{9}\left[\frac{2}{3}u^{3/2}\right]_4^{13} = \frac{2}{27}(13^{3/2} - 8)$.

9. $\mathbf{r}_1(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j} + t\,\mathbf{k}$ and $\mathbf{r}_2(t) = (1+t)\,\mathbf{i} + t^2\,\mathbf{j} + t^3\,\mathbf{k}$. The angle θ of intersection of the two curves is the angle between their respective tangents at the point of intersection. For both curves, the point $(1,0,0)$ corresponds to $t = 0$.

$\mathbf{r}_1'(t) = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j} + \mathbf{k} \quad\Rightarrow\quad \mathbf{r}_1'(0) = \mathbf{j} + \mathbf{k}$, and $\mathbf{r}_2'(t) = \mathbf{i} + 2t\,\mathbf{j} + 3t^2\,\mathbf{k} \quad\Rightarrow\quad \mathbf{r}_2'(0) = \mathbf{i}$.

$\mathbf{r}_1'(0)\cdot\mathbf{r}_2'(0) = (\mathbf{j}+\mathbf{k})\cdot\mathbf{i} = 0$. Therefore, the curves intersect in a right angle, that is, $\theta = 90°$.

10. $\mathbf{r}(t) = e^t\,\mathbf{i} + e^t\sin t\,\mathbf{j} + e^t\cos t\,\mathbf{k}$. The parametric value corresponding to the point $(1,0,1)$ is $t = 0$.

$\mathbf{r}'(t) = e^t\,\mathbf{i} + e^t(\cos t + \sin t)\,\mathbf{j} + e^t(\cos t - \sin t)\,\mathbf{k} \quad\Rightarrow\quad |\mathbf{r}'(t)| = e^t\sqrt{1^2 + (\cos t + \sin t)^2 + (\cos t - \sin t)^2} = \sqrt{3}\,e^t$

and $s(t) = \int_0^t e^u\sqrt{3}\,du = \sqrt{3}(e^t - 1) \quad\Rightarrow\quad 1 + \frac{1}{\sqrt{3}}s = e^t \quad\Rightarrow\quad t = \ln\left(1 + \frac{1}{\sqrt{3}}s\right)$.

Therefore, $\mathbf{r}(t(s)) = \left(1 + \frac{1}{\sqrt{3}}s\right)\mathbf{i} + \left(1 + \frac{1}{\sqrt{3}}s\right)\sin\ln\left(1 + \frac{1}{\sqrt{3}}s\right)\mathbf{j} + \left(1 + \frac{1}{\sqrt{3}}s\right)\cos\ln\left(1 + \frac{1}{\sqrt{3}}s\right)\mathbf{k}$.

11. (a) $\mathbf{r}(t) = \left\langle \sin^3 t, \cos^3 t, \sin^2 t\right\rangle \quad\Rightarrow\quad \mathbf{r}'(t) = \left\langle 3\sin^2 t\cos t, -3\cos^2 t\sin t, 2\sin t\cos t\right\rangle$,

$$\begin{aligned} |\mathbf{r}'(t)| &= \sqrt{9\sin^4 t\cos^2 t + 9\cos^4 t\sin^2 t + 4\sin^2 t\cos^2 t} \\ &= \sqrt{\sin^2 t\cos^2 t\left(9\sin^2 t + 9\cos^2 t + 4\right)} = \sqrt{13}\,\sin t\cos t \quad [\text{since } 0 \le t \le \pi/2 \quad\Rightarrow\quad \sin t, \cos t \ge 0] \end{aligned}$$

Then $\mathbf{T}(t) = \dfrac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \dfrac{1}{\sqrt{13}\,\sin t\cos t}\left\langle 3\sin^2 t\cos t, -3\cos^2 t\sin t, 2\sin t\cos t\right\rangle = \frac{1}{\sqrt{13}}\left\langle 3\sin t, -3\cos t, 2\right\rangle$.

(b) $\mathbf{T}'(t) = \frac{1}{\sqrt{13}}\left\langle 3\cos t, 3\sin t, 0\right\rangle$, $\quad |\mathbf{T}'(t)| = \frac{1}{\sqrt{13}}\sqrt{9\cos^2 t + 9\sin^2 t + 0} = \frac{3}{\sqrt{13}}$, and

$\mathbf{N}(t) = \dfrac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|} = \frac{1}{3}\left\langle 3\cos t, 3\sin t, 0\right\rangle = \left\langle \cos t, \sin t, 0\right\rangle$.

(c) $\mathbf{B}(t) = \mathbf{T}(t)\times\mathbf{N}(t) = \frac{1}{\sqrt{13}}\left\langle 3\sin t, -3\cos t, 2\right\rangle \times \left\langle \cos t, \sin t, 0\right\rangle = \frac{1}{\sqrt{13}}\left\langle -2\sin t, 2\cos t, 3\right\rangle$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $\kappa(t) = \dfrac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \dfrac{3/\sqrt{13}}{\sqrt{13}\,\sin t\cos t} = \dfrac{3}{13\,\sin t\cos t}$ or $\frac{3}{13}\csc t\sec t$

(e) $\tau(t) = -\dfrac{\mathbf{B}'(t)\cdot\mathbf{N}(t)}{|\mathbf{r}'(t)|} = -\dfrac{\frac{1}{\sqrt{13}}\langle -2\cos t, -2\sin t, 0\rangle \cdot \langle \cos t, \sin t, 0\rangle}{\sqrt{13}\,\sin t\cos t} = -\dfrac{-2\cos^2 t - 2\sin^2 t + 0}{13\sin t\cos t}$

$= \dfrac{2}{13\,\sin t\cos t}$ or $\frac{2}{13}\csc t\sec t$

12. See the instructions for Exercises 13.3.46–49. $x = 3\cos t,\ y = 4\sin t \ \Rightarrow\ \dot{x} = -3\sin t,\ \dot{y} = 4\cos t \ \Rightarrow$ $\ddot{x} = -3\cos t,\ \ddot{y} = -4\sin t$.

$\kappa(t) = \dfrac{|\dot{x}\ddot{y} - \dot{y}\ddot{x}|}{[\dot{x}^2 + \dot{y}^2]^{3/2}} = \dfrac{|(-3\sin t)(-4\sin t) - (4\cos t)(-3\cos t)|}{(9\sin^2 t + 16\cos^2 t)^{3/2}} = \dfrac{12}{(9\sin^2 t + 16\,\cos^2 t)^{3/2}}$.

At $(3, 0)$, $t = 0$ and $\kappa\,(0) = 12/(16)^{3/2} = \frac{12}{64} = \frac{3}{16}$. At $(0, 4)$, $t = \frac{\pi}{2}$ and $\kappa\left(\frac{\pi}{2}\right) = 12/9^{3/2} = \frac{12}{27} = \frac{4}{9}$.

13. $y = x^4,\ y' = 4x^3,\ y'' = 12x^2$, and $\kappa(x) = \dfrac{|y''|}{[1 + (y')^2]^{3/2}} = \dfrac{|12x^2|}{(1 + 16x^6)^{3/2}}$, so $\kappa(1) = \dfrac{12}{17^{3/2}}$.

14. $y = x^4 - x^2,\ y' = 4x^3 - 2x,\ y'' = 12x^2 - 2$, and

$\kappa(x) = \dfrac{|y''|}{[1 + (y')^2]^{3/2}} = \dfrac{|12x^2 - 2|}{[1 + (4x^3 - 2x)^2]^{3/2}} \ \Rightarrow\ \kappa(0) = 2.$

So the osculating circle has radius $\frac{1}{2}$ and center $\left(0, -\frac{1}{2}\right)$.

Thus, its equation is $x^2 + \left(y + \frac{1}{2}\right)^2 = \frac{1}{4}$.

15. $\mathbf{r}(t) = \langle \sin 2t, t, \cos 2t\rangle \ \Rightarrow\ \mathbf{r}'(t) = \langle 2\cos 2t, 1, -2\sin 2t\rangle \ \Rightarrow\ \mathbf{T}(t) = \frac{1}{\sqrt{5}}\langle 2\cos 2t, 1, -2\sin 2t\rangle \ \Rightarrow$ $\mathbf{T}'(t) = \frac{1}{\sqrt{5}}\langle -4\sin 2t, 0, -4\cos 2t\rangle \ \Rightarrow\ \mathbf{N}(t) = \langle -\sin 2t, 0, -\cos 2t\rangle$. So $\mathbf{N} = \mathbf{N}(\pi) = \langle 0, 0, -1\rangle$ and $\mathbf{B} = \mathbf{T}\times\mathbf{N} = \frac{1}{\sqrt{5}}\langle -1, 2, 0\rangle$. So a normal to the osculating plane is $\langle -1, 2, 0\rangle$ and an equation is $-1(x - 0) + 2(y - \pi) + 0(z - 1) = 0$ or $x - 2y + 2\pi = 0$.

16. (a) The average velocity over $[3, 3.2]$ is given by $\dfrac{\mathbf{r}(3.2) - \mathbf{r}(3)}{3.2 - 3} = 5[\mathbf{r}(3.2) - \mathbf{r}(3)]$, so we draw a vector with the same direction but 5 times the length of the vector $[\mathbf{r}(3.2) - \mathbf{r}(3)]$.

(b) $\mathbf{v}(3) = \mathbf{r}'(3) = \lim\limits_{h\to 0} \dfrac{\mathbf{r}(3 + h) - \mathbf{r}(3)}{h}$

(c) $\mathbf{T}(3) = \dfrac{\mathbf{r}'(3)}{|\mathbf{r}'(3)|}$, a unit vector in the same direction as $\mathbf{r}'(3)$, that is, parallel to the tangent line to the curve at $\mathbf{r}(3)$, pointing in the direction corresponding to increasing t, and with length 1.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\mathbf{r}(t) = t\ln t\,\mathbf{i} + t\,\mathbf{j} + e^{-t}\,\mathbf{k}$, $\quad \mathbf{v}(t) = \mathbf{r}'(t) = (1+\ln t)\,\mathbf{i} + \mathbf{j} - e^{-t}\,\mathbf{k}$,

$|\mathbf{v}(t)| = \sqrt{(1+\ln t)^2 + 1^2 + (-e^{-t})^2} = \sqrt{2 + 2\ln t + (\ln t)^2 + e^{-2t}}$, $\quad \mathbf{a}(t) = \mathbf{v}'(t) = \frac{1}{t}\,\mathbf{i} + e^{-t}\,\mathbf{k}$

18. $\mathbf{r}(t) = (2t^2 - 3)\,\mathbf{i} + 2t\,\mathbf{j} \quad \Rightarrow \quad \mathbf{v}(t) = \mathbf{r}'(t) = 4t\,\mathbf{i} + 2\,\mathbf{j}$,

speed $= |\mathbf{v}(t)| = \sqrt{16t^2+4} = 2\sqrt{4t^2+1}$, and $\mathbf{a}(t) = \mathbf{v}'(t) = \mathbf{r}''(t) = 4\,\mathbf{i}$.

At $t = 1$ we have $\mathbf{r}(1) = -\mathbf{i} + 2\,\mathbf{j}$, $\mathbf{v}(1) = 4\,\mathbf{i} + 2\,\mathbf{j}$, $\mathbf{a}(1) = 4\,\mathbf{i}$.

Notice that $y/2 = t \quad \Rightarrow \quad x = 2\,(y/2)^2 - 3 = \frac{1}{2}y^2 - 3$, so the path is a parabola.

19. $\mathbf{v}(t) = \int \mathbf{a}(t)\,dt = \int (6t\,\mathbf{i} + 12t^2\,\mathbf{j} - 6t\,\mathbf{k})\,dt = 3t^2\,\mathbf{i} + 4t^3\,\mathbf{j} - 3t^2\,\mathbf{k} + \mathbf{C}$, but $\mathbf{i} - \mathbf{j} + 3\,\mathbf{k} = \mathbf{v}(0) = \mathbf{0} + \mathbf{C}$,

so $\mathbf{C} = \mathbf{i} - \mathbf{j} + 3\,\mathbf{k}$ and $\mathbf{v}(t) = (3t^2+1)\,\mathbf{i} + (4t^3-1)\,\mathbf{j} + (3-3t^2)\,\mathbf{k}$.

$\mathbf{r}(t) = \int \mathbf{v}(t)\,dt = (t^3+t)\,\mathbf{i} + (t^4-t)\,\mathbf{j} + (3t-t^3)\,\mathbf{k} + \mathbf{D}$.

But $\mathbf{r}(0) = \mathbf{0}$, so $\mathbf{D} = \mathbf{0}$ and $\mathbf{r}(t) = (t^3+t)\,\mathbf{i} + (t^4-t)\,\mathbf{j} + (3t-t^3)\,\mathbf{k}$.

20. We set up the axes so that the shot leaves the athlete's hand 7 ft above the origin. Then we are given $\mathbf{r}(0) = 7\mathbf{j}$, $|\mathbf{v}(0)| = 43$ ft/s, and $\mathbf{v}(0)$ has direction given by a 45° angle of elevation. Then a unit vector in the direction of $\mathbf{v}(0)$ is $\frac{1}{\sqrt{2}}(\mathbf{i}+\mathbf{j}) \quad \Rightarrow \quad \mathbf{v}(0) = \frac{43}{\sqrt{2}}(\mathbf{i}+\mathbf{j})$. Assuming air resistance is negligible, the only external force is due to gravity, so as in Example 13.4.5 we have $\mathbf{a} = -g\,\mathbf{j}$ where here $g \approx 32$ ft/s^2. Since $\mathbf{v}'(t) = \mathbf{a}(t)$, we integrate, giving $\mathbf{v}(t) = -gt\,\mathbf{j} + \mathbf{C}$ where $\mathbf{C} = \mathbf{v}(0) = \frac{43}{\sqrt{2}}(\mathbf{i}+\mathbf{j}) \quad \Rightarrow \quad \mathbf{v}(t) = \frac{43}{\sqrt{2}}\,\mathbf{i} + \left(\frac{43}{\sqrt{2}} - gt\right)\mathbf{j}$. Since $\mathbf{r}'(t) = \mathbf{v}(t)$ we integrate again, so $\mathbf{r}(t) = \frac{43}{\sqrt{2}}t\,\mathbf{i} + \left(\frac{43}{\sqrt{2}}t - \frac{1}{2}gt^2\right)\mathbf{j} + \mathbf{D}$. But $\mathbf{D} = \mathbf{r}(0) = 7\,\mathbf{j} \quad \Rightarrow \quad \mathbf{r}(t) = \frac{43}{\sqrt{2}}t\,\mathbf{i} + \left(\frac{43}{\sqrt{2}}t - \frac{1}{2}gt^2 + 7\right)\mathbf{j}$.

(a) At 2 seconds, the shot is at $\mathbf{r}(2) = \frac{43}{\sqrt{2}}(2)\,\mathbf{i} + \left(\frac{43}{\sqrt{2}}(2) - \frac{1}{2}g(2)^2 + 7\right)\mathbf{j} \approx 60.8\,\mathbf{i} + 3.8\,\mathbf{j}$, so the shot is about 3.8 ft above the ground, at a horizontal distance of 60.8 ft from the athlete.

(b) The shot reaches its maximum height when the vertical component of velocity is 0: $\frac{43}{\sqrt{2}} - gt = 0 \quad \Rightarrow$

$t = \dfrac{43}{\sqrt{2}\,g} \approx 0.95$ s. Then $\mathbf{r}(0.95) \approx 28.9\,\mathbf{i} + 21.4\,\mathbf{j}$, so the maximum height is approximately 21.4 ft.

(c) The shot hits the ground when the vertical component of $\mathbf{r}(t)$ is 0, so $\frac{43}{\sqrt{2}}t - \frac{1}{2}gt^2 + 7 = 0 \quad \Rightarrow$

$-16t^2 + \frac{43}{\sqrt{2}}t + 7 = 0 \quad \Rightarrow \quad t \approx 2.11$ s. $\quad \mathbf{r}(2.11) \approx 64.2\,\mathbf{i} - 0.08\,\mathbf{j}$, thus the shot lands approximately 64.2 ft from the athlete.

21. Example 13.4.5 showed that the maximum horizontal range is achieved with an angle of elevation of 45°. In this case, however, the projectile would hit the top of the tunnel using that angle. The horizontal range will be maximized with the largest angle of elevation that keeps the projectile within a height of 30 m. From Example 13.4.5 we know that the position function of the projectile is $\mathbf{r}(t) = (v_0\cos\alpha)t\,\mathbf{i} + \left[(v_0\sin\alpha)t - \frac{1}{2}gt^2\right]\mathbf{j}$ and the velocity is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{v}(t) = \mathbf{r}'(t) = (v_0 \cos\alpha)\,\mathbf{i} + [(v_0 \sin\alpha) - gt]\,\mathbf{j}$. The projectile achieves its maximum height when the vertical component of velocity is zero, so $(v_0 \sin\alpha) - gt = 0 \;\Rightarrow\; t = \dfrac{v_0 \sin\alpha}{g}$. We want the vertical height of the projectile at that time to be 30 m: $(v_0 \sin\alpha)\left(\dfrac{v_0 \sin\alpha}{g}\right) - \frac{1}{2}g\left(\dfrac{v_0 \sin\alpha}{g}\right)^2 = 30 \;\Rightarrow$

$$\left(\frac{v_0^2 \sin^2\alpha}{g}\right) - \frac{1}{2}\left(\frac{v_0^2 \sin^2\alpha}{g}\right) = 30 \;\Rightarrow\; \frac{v_0^2 \sin^2\alpha}{2g} = 30 \;\Rightarrow\; \sin^2\alpha = \frac{30(2g)}{v_0^2} = \frac{60(9.8)}{40^2} = 0.3675 \;\Rightarrow$$

$\sin\alpha = \sqrt{0.3675}$. Thus the desired angle of elevation is $\alpha = \sin^{-1}\sqrt{0.3675} \approx 37.3\,^\circ$.

From the same example, the horizontal distance traveled is $d = \dfrac{v_0^2 \sin 2\alpha}{g} \approx \dfrac{40^2 \sin(74.6\,^\circ)}{9.8} \approx 157.4$ m.

22. $\mathbf{r}(t) = t\,\mathbf{i} + 2t\,\mathbf{j} + t^2\,\mathbf{k}$, $\mathbf{r}'(t) = \mathbf{i} + 2\,\mathbf{j} + 2t\,\mathbf{k}$, $\mathbf{r}''(t) = 2\,\mathbf{k}$, $|\mathbf{r}'(t)| = \sqrt{1 + 4 + 4t^2} = \sqrt{4t^2 + 5}$.

Then $a_T = \dfrac{\mathbf{r}'(t) \cdot \mathbf{r}''(t)}{|\mathbf{r}'(t)|} = \dfrac{4t}{\sqrt{4t^2+5}}$ and $a_N = \dfrac{|\mathbf{r}'(t) \times \mathbf{r}''(t)|}{|\mathbf{r}'(t)|} = \dfrac{|4\,\mathbf{i} - 2\,\mathbf{j}|}{\sqrt{4t^2+5}} = \dfrac{2\sqrt{5}}{\sqrt{4t^2+5}}$.

23. (a) Instead of proceeding directly, we use Formula 3 of Theorem 13.2.3: $\mathbf{r}(t) = t\,\mathbf{R}(t) \;\Rightarrow$

$\mathbf{v} = \mathbf{r}'(t) = \mathbf{R}(t) + t\,\mathbf{R}'(t) = \cos\omega t\,\mathbf{i} + \sin\omega t\,\mathbf{j} + t\,\mathbf{v}_d$.

(b) Using the same method as in part (a) and starting with $\mathbf{v} = \mathbf{R}(t) + t\,\mathbf{R}'(t)$, we have

$\mathbf{a} = \mathbf{v}' = \mathbf{R}'(t) + \mathbf{R}'(t) + t\,\mathbf{R}''(t) = 2\,\mathbf{R}'(t) + t\,\mathbf{R}''(t) = 2\,\mathbf{v}_d + t\,\mathbf{a}_d$.

(c) Here we have $\mathbf{r}(t) = e^{-t}\cos\omega t\,\mathbf{i} + e^{-t}\sin\omega t\,\mathbf{j} = e^{-t}\,\mathbf{R}(t)$. So, as in parts (a) and (b),

$\mathbf{v} = \mathbf{r}'(t) = e^{-t}\,\mathbf{R}'(t) - e^{-t}\,\mathbf{R}(t) = e^{-t}[\mathbf{R}'(t) - \mathbf{R}(t)] \;\Rightarrow$

$$\begin{aligned}\mathbf{a} = \mathbf{v}' &= e^{-t}[\mathbf{R}''(t) - \mathbf{R}'(t)] - e^{-t}[\mathbf{R}'(t) - \mathbf{R}(t)] = e^{-t}[\mathbf{R}''(t) - 2\,\mathbf{R}'(t) + \mathbf{R}(t)] \\ &= e^{-t}\,\mathbf{a}_d - 2e^{-t}\,\mathbf{v}_d + e^{-t}\,\mathbf{R}\end{aligned}$$

Thus, the Coriolis acceleration (the sum of the "extra" terms not involving $\mathbf{a}_d$) is $-2e^{-t}\,\mathbf{v}_d + e^{-t}\,\mathbf{R}$.

24. (a) $F(x) = \begin{cases} 1 & \text{if } x \le 0 \\ \sqrt{1-x^2} & \text{if } 0 < x < \frac{1}{\sqrt{2}} \\ \sqrt{2} - x & \text{if } x \ge \frac{1}{\sqrt{2}} \end{cases} \;\Rightarrow\; F'(x) = \begin{cases} 0 & \text{if } x < 0 \\ -x/\sqrt{1-x^2} & \text{if } 0 < x < \frac{1}{\sqrt{2}} \\ -1 & \text{if } x > \frac{1}{\sqrt{2}} \end{cases} \;\Rightarrow$

$$F''(x) = \begin{cases} 0 & \text{if } x < 0 \\ -1/(1-x^2)^{3/2} & \text{if } 0 < x < \frac{1}{\sqrt{2}} \\ 0 & \text{if } x > \frac{1}{\sqrt{2}} \end{cases}$$

since $\dfrac{d}{dx}[-x(1-x^2)^{-1/2}] = -(1-x^2)^{-1/2} - x^2(1-x^2)^{-3/2} = -(1-x^2)^{-3/2}$.

Now $\lim\limits_{x \to 0^+} \sqrt{1-x^2} = 1 = F(0)$ and $\lim\limits_{x \to (1/\sqrt{2})^-} \sqrt{1-x^2} = \frac{1}{\sqrt{2}} = F\left(\frac{1}{\sqrt{2}}\right)$, so F is continuous. Also, since

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\lim\limits_{x\to 0^+} F'(x) = 0 = \lim\limits_{x\to 0^-} F'(x)$ and $\lim\limits_{x\to\left(1/\sqrt{2}\right)^-} F'(x) = -1 = \lim\limits_{x\to\left(1/\sqrt{2}\right)^+} F'(x)$, F' is continuous. But

$\lim\limits_{x\to 0^+} F''(x) = -1 \neq 0 = \lim\limits_{x\to 0^-} F''(x)$, so F'' is not continuous at $x = 0$. (The same is true at $x = \frac{1}{\sqrt{2}}$.)

So F does not have continuous curvature.

(b) Set $P(x) = ax^5 + bx^4 + cx^3 + dx^2 + ex + f$. The continuity conditions on P are $P(0) = 0$, $P(1) = 1$, $P'(0) = 0$ and $P'(1) = 1$. Also the curvature must be continuous. For $x \leq 0$ and $x \geq 1$, $\kappa(x) = 0$; elsewhere

$\kappa(x) = \dfrac{|P''(x)|}{(1 + [P'(x)]^2)^{3/2}}$, so we need $P''(0) = 0$ and $P''(1) = 0$.

The conditions $P(0) = P'(0) = P''(0) = 0$ imply that $d = e = f = 0$. The other conditions imply that $a + b + c = 1$, $5a + 4b + 3c = 1$, and $10a + 6b + 3c = 0$. From these, we find that $a = 3$, $b = -8$, and $c = 6$. Therefore $P(x) = 3x^5 - 8x^4 + 6x^3$. Since there was no solution with $a = 0$, this could not have been done with a polynomial of degree 4.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1. (a) $\mathbf{r}(t) = R\cos\omega t\,\mathbf{i} + R\sin\omega t\,\mathbf{j} \quad\Rightarrow\quad \mathbf{v} = \mathbf{r}'(t) = -\omega R\sin\omega t\,\mathbf{i} + \omega R\cos\omega t\,\mathbf{j}$, so $\mathbf{r} = R(\cos\omega t\,\mathbf{i} + \sin\omega t\,\mathbf{j})$ and $\mathbf{v} = \omega R(-\sin\omega t\,\mathbf{i} + \cos\omega t\,\mathbf{j})$. $\mathbf{v}\cdot\mathbf{r} = \omega R^2(-\cos\omega t\sin\omega t + \sin\omega t\cos\omega t) = 0$, so $\mathbf{v}\perp\mathbf{r}$. Since $\mathbf{r}$ points along a radius of the circle, and $\mathbf{v}\perp\mathbf{r}$, $\mathbf{v}$ is tangent to the circle. Because it is a velocity vector, $\mathbf{v}$ points in the direction of motion.

(b) In (a), we wrote $\mathbf{v}$ in the form $\omega R\,\mathbf{u}$, where $\mathbf{u}$ is the unit vector $-\sin\omega t\,\mathbf{i} + \cos\omega t\,\mathbf{j}$. Clearly $|\mathbf{v}| = \omega R\,|\mathbf{u}| = \omega R$. At speed ωR, the particle completes one revolution, a distance $2\pi R$, in time $T = \dfrac{2\pi R}{\omega R} = \dfrac{2\pi}{\omega}$.

(c) $\mathbf{a} = \dfrac{d\mathbf{v}}{dt} = -\omega^2 R\cos\omega t\,\mathbf{i} - \omega^2 R\sin\omega t\,\mathbf{j} = -\omega^2 R(\cos\omega t\,\mathbf{i} + \sin\omega t\,\mathbf{j})$, so $\mathbf{a} = -\omega^2\mathbf{r}$. This shows that $\mathbf{a}$ is proportional to $\mathbf{r}$ and points in the opposite direction (toward the origin). Also, $|\mathbf{a}| = \omega^2\,|\mathbf{r}| = \omega^2 R$.

(d) By Newton's Second Law (see Section 13.4), $\mathbf{F} = m\mathbf{a}$, so $|\mathbf{F}| = m\,|\mathbf{a}| = mR\omega^2 = \dfrac{m\,(\omega R)^2}{R} = \dfrac{m\,|\mathbf{v}|^2}{R}$.

2. (a) Dividing the equation $|\mathbf{F}|\sin\theta = \dfrac{mv_R^2}{R}$ by the equation $|\mathbf{F}|\cos\theta = mg$, we obtain $\tan\theta = \dfrac{v_R^2}{Rg}$, so $v_R^2 = Rg\tan\theta$.

(b) $R = 400$ ft and $\theta = 12^\circ$, so $v_R = \sqrt{Rg\tan\theta} \approx \sqrt{400\cdot 32\cdot\tan 12^\circ} \approx 52.16$ ft/s ≈ 36 mi/h.

(c) We want to choose a new radius R_1 for which the new rated speed is $\frac{3}{2}$ of the old one: $\sqrt{R_1 g\tan 12^\circ} = \frac{3}{2}\sqrt{Rg\tan 12^\circ}$. Squaring, we get $R_1 g\tan 12^\circ = \frac{9}{4}Rg\tan 12^\circ$, so $R_1 = \frac{9}{4}R = \frac{9}{4}(400) = 900$ ft.

3. (a) The projectile reaches maximum height when $0 = \dfrac{dy}{dt} = \dfrac{d}{dt}\left[(v_0\sin\alpha)t - \frac{1}{2}gt^2\right] = v_0\sin\alpha - gt$; that is, when $t = \dfrac{v_0\sin\alpha}{g}$ and $y = (v_0\sin\alpha)\left(\dfrac{v_0\sin\alpha}{g}\right) - \dfrac{1}{2}g\left(\dfrac{v_0\sin\alpha}{g}\right)^2 = \dfrac{v_0^2\sin^2\alpha}{2g}$. This is the maximum height attained when the projectile is fired with an angle of elevation α. This maximum height is largest when $\alpha = 90^\circ$. In that case, $\sin\alpha = 1$ and the maximum height is $\dfrac{v_0^2}{2g}$.

(b) Let $R = v_0^2/g$. We are asked to consider the parabola $x^2 + 2Ry - R^2 = 0$ which can be rewritten as $y = -\dfrac{1}{2R}x^2 + \dfrac{R}{2}$. The points on or inside this parabola are those for which $-R \le x \le R$ and $0 \le y \le \dfrac{-1}{2R}x^2 + \dfrac{R}{2}$. When the projectile is fired at angle of elevation α, the points (x, y) along its path satisfy the relations $x = (v_0\cos\alpha)\,t$ and $y = (v_0\sin\alpha)t - \frac{1}{2}gt^2$, where $0 \le t \le (2v_0\sin\alpha)/g$ (as in Example 13.4.5). Thus $|x| \le \left|v_0\cos\alpha\left(\dfrac{2v_0\sin\alpha}{g}\right)\right| = \left|\dfrac{v_0^2}{g}\sin 2\alpha\right| \le \left|\dfrac{v_0^2}{g}\right| = |R|$. This shows that $-R \le x \le R$.

For t in the specified range, we also have $y = t\big(v_0\sin\alpha - \frac{1}{2}gt\big) = \frac{1}{2}gt\left(\dfrac{2v_0\sin\alpha}{g} - t\right) \ge 0$ and

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$y = (v_0 \sin\alpha)\dfrac{x}{v_0\cos\alpha} - \dfrac{g}{2}\left(\dfrac{x}{v_0\cos\alpha}\right)^2 = (\tan\alpha)\,x - \dfrac{g}{2v_0^2\cos^2\alpha}x^2 = -\dfrac{1}{2R\cos^2\alpha}\,x^2 + (\tan\alpha)\,x$. Thus

$$\begin{aligned} y - \left(\frac{-1}{2R}x^2 + \frac{R}{2}\right) &= \frac{-1}{2R\cos^2\alpha}\,x^2 + \frac{1}{2R}x^2 + (\tan\alpha)\,x - \frac{R}{2} \\ &= \frac{x^2}{2R}\left(1 - \frac{1}{\cos^2\alpha}\right) + (\tan\alpha)\,x - \frac{R}{2} = \frac{x^2(1-\sec^2\alpha) + 2R\,(\tan\alpha)\,x - R^2}{2R} \\ &= \frac{-(\tan^2\alpha)\,x^2 + 2R\,(\tan\alpha)\,x - R^2}{2R} = \frac{-\left[(\tan\alpha)\,x - R\right]^2}{2R} \le 0 \end{aligned}$$

We have shown that every target that can be hit by the projectile lies on or inside the parabola $y = -\dfrac{1}{2R}\,x^2 + \dfrac{R}{2}$.

Now let (a, b) be any point on or inside the parabola $y = -\dfrac{1}{2R}\,x^2 + \dfrac{R}{2}$. Then $-R \le a \le R$ and $0 \le b \le -\dfrac{1}{2R}\,a^2 + \dfrac{R}{2}$. We seek an angle α such that (a, b) lies in the path of the projectile; that is, we wish to find an angle α such that $b = -\dfrac{1}{2R\cos^2\alpha}\,a^2 + (\tan\alpha)\,a$ or equivalently $b = \dfrac{-1}{2R}\,(\tan^2\alpha + 1)a^2 + (\tan\alpha)\,a$. Rearranging this equation we get $\dfrac{a^2}{2R}\tan^2\alpha - a\tan\alpha + \left(\dfrac{a^2}{2R} + b\right) = 0$ or $a^2(\tan\alpha)^2 - 2aR(\tan\alpha) + (a^2 + 2bR) = 0 \;\; (\star)$. This quadratic equation for $\tan\alpha$ has real solutions exactly when the discriminant is nonnegative. Now $B^2 - 4AC \ge 0 \;\;\Leftrightarrow$ $(-2aR)^2 - 4a^2(a^2 + 2bR) \ge 0 \;\;\Leftrightarrow\;\; 4a^2(R^2 - a^2 - 2bR) \ge 0 \;\;\Leftrightarrow\;\; -a^2 - 2bR + R^2 \ge 0 \;\;\Leftrightarrow$ $b \le \dfrac{1}{2R}\,(R^2 - a^2) \;\;\Leftrightarrow\;\; b \le \dfrac{-1}{2R}\,a^2 + \dfrac{R}{2}$. This condition is satisfied since (a, b) is on or inside the parabola $y = -\dfrac{1}{2R}\,x^2 + \dfrac{R}{2}$. It follows that (a, b) lies in the path of the projectile when $\tan\alpha$ satisfies $(\star)$, that is, when

$$\tan\alpha = \frac{2aR \pm \sqrt{4a^2(R^2 - a^2 - 2bR)}}{2a^2} = \frac{R \pm \sqrt{R^2 - 2bR - a^2}}{a}.$$

(c)

If the gun is pointed at a target with height h at a distance D downrange, then $\tan\alpha = h/D$. When the projectile reaches a distance D downrange (remember we are assuming that it doesn't hit the ground first), we have $D = x = (v_0\cos\alpha)t$, so $t = \dfrac{D}{v_0\cos\alpha}$ and $y = (v_0\sin\alpha)t - \frac{1}{2}gt^2 = D\tan\alpha - \dfrac{gD^2}{2v_0^2\cos^2\alpha}$.

Meanwhile, the target, whose x-coordinate is also D, has fallen from height h to height $h - \frac{1}{2}gt^2 = D\tan\alpha - \dfrac{gD^2}{2v_0^2\cos^2\alpha}$. Thus the projectile hits the target.

4. (a) As in Problem 3, $\mathbf{r}(t) = (v_0\cos\alpha)t\,\mathbf{i} + \left[(v_0\sin\alpha)t - \frac{1}{2}gt^2\right]\mathbf{j}$, so $x = (v_0\cos\alpha)t$ and $y = (v_0\sin\alpha)t - \frac{1}{2}gt^2$. The difference here is that the projectile travels until it reaches a point where $x > 0$ and $y = -(\tan\theta)x$. (Here $0 \le \theta \le \frac{\pi}{2}$.)

From the parametric equations, we obtain $t = \dfrac{x}{v_0\cos\alpha}$ and $y = \dfrac{(v_0\sin\alpha)x}{v_0\cos\alpha} - \dfrac{gx^2}{2v_0^2\cos^2\alpha} = (\tan\alpha)x - \dfrac{gx^2}{2v_0^2\cos^2\alpha}$.

Thus the projectile hits the inclined plane at the point where $(\tan\alpha)x - \dfrac{gx^2}{2v_0^2\cos^2\alpha} = -(\tan\theta)x$. Since $\dfrac{gx^2}{2v_0^2\cos^2\alpha} = (\tan\alpha + \tan\theta)x$ and $x > 0$, we must have $\dfrac{gx}{2v_0^2\cos^2\alpha} = \tan\alpha + \tan\theta$. It follows that

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x = \dfrac{2v_0^2 \cos^2 \alpha}{g}(\tan\alpha + \tan\theta)$ and $t = \dfrac{x}{v_0 \cos\alpha} = \dfrac{2v_0 \cos\alpha}{g}(\tan\alpha + \tan\theta)$. This means that the parametric equations are defined for t in the interval $\left[0, \dfrac{2v_0 \cos\alpha}{g}(\tan\alpha + \tan\theta)\right]$.

(b) The downhill range (that is, the distance to the projectile's landing point as measured along the inclined plane) is $R(\alpha) = x \sec\theta$, where x is the coordinate of the landing point calculated in part (a). Thus

$$R(\alpha) = \frac{2v_0^2 \cos^2\alpha}{g}(\tan\alpha + \tan\theta)\sec\theta = \frac{2v_0^2}{g}\left(\frac{\sin\alpha\,\cos\alpha}{\cos\theta} + \frac{\cos^2\alpha\,\sin\theta}{\cos^2\theta}\right)$$
$$= \frac{2v_0^2 \cos\alpha}{g\cos^2\theta}(\sin\alpha\,\cos\theta + \cos\alpha\,\sin\theta) = \frac{2v_0^2 \cos\alpha\,\sin(\alpha+\theta)}{g\cos^2\theta}$$

$R(\alpha)$ is maximized when

$$0 = R'(\alpha) = \frac{2v_0^2}{g\cos^2\theta}[-\sin\alpha\,\sin(\alpha+\theta) + \cos\alpha\,\cos(\alpha+\theta)]$$
$$= \frac{2v_0^2}{g\cos^2\theta}\cos[(\alpha+\theta)+\alpha] = \frac{2v_0^2\cos(2\alpha+\theta)}{g\cos^2\theta}$$

This condition implies that $\cos(2\alpha+\theta) = 0 \;\Rightarrow\; 2\alpha+\theta = \frac{\pi}{2} \;\Rightarrow\; \alpha = \frac{1}{2}\left(\frac{\pi}{2} - \theta\right)$.

(c) The solution is similar to the solutions to parts (a) and (b). This time the projectile travels until it reaches a point where $x > 0$ and $y = (\tan\theta)x$. Since $\tan\theta = -\tan(-\theta)$, we obtain the solution from the previous one by replacing θ with $-\theta$. The desired angle is $\alpha = \frac{1}{2}\left(\frac{\pi}{2} + \theta\right)$.

(d) As observed in part (c), firing the projectile up an inclined plane with angle of inclination θ involves the same equations as in parts (a) and (b) but with θ replaced by $-\theta$. So if R is the distance up an inclined plane, we know from part (b) that $R = \dfrac{2v_0^2\cos\alpha\sin(\alpha-\theta)}{g\cos^2(-\theta)} \;\Rightarrow\; v_0^2 = \dfrac{Rg\cos^2\theta}{2\cos\alpha\sin(\alpha-\theta)}$. v_0^2 is minimized (and hence v_0 is minimized) with respect to α when

$$0 = \frac{d}{d\alpha}(v_0^2) = \frac{Rg\cos^2\theta}{2}\cdot\frac{-(\cos\alpha\cos(\alpha-\theta) - \sin\alpha\sin(\alpha-\theta))}{[\cos\alpha\sin(\alpha-\theta)]^2}$$
$$= \frac{-Rg\cos^2\theta}{2}\cdot\frac{\cos[\alpha+(\alpha-\theta)]}{[\cos\alpha\sin(\alpha-\theta)]^2} = \frac{-Rg\cos^2\theta}{2}\cdot\frac{\cos(2\alpha-\theta)}{[\cos\alpha\sin(\alpha-\theta)]^2}$$

Since $\theta < \alpha < \frac{\pi}{2}$, this implies $\cos(2\alpha-\theta) = 0 \;\Leftrightarrow\; 2\alpha - \theta = \frac{\pi}{2} \;\Rightarrow\; \alpha = \frac{1}{2}\left(\frac{\pi}{2}+\theta\right)$. Thus the initial speed, and hence the energy required, is minimized for $\alpha = \frac{1}{2}\left(\frac{\pi}{2}+\theta\right)$.

5. (a) $\mathbf{a} = -g\,\mathbf{j} \;\Rightarrow\; \mathbf{v} = \mathbf{v}_0 - gt\,\mathbf{j} = 2\,\mathbf{i} - gt\,\mathbf{j} \;\Rightarrow\; \mathbf{s} = \mathbf{s}_0 + 2t\,\mathbf{i} - \frac{1}{2}gt^2\,\mathbf{j} = 3.5\,\mathbf{j} + 2t\,\mathbf{i} - \frac{1}{2}gt^2\,\mathbf{j} \;\Rightarrow$ $\mathbf{s} = 2t\,\mathbf{i} + \left(3.5 - \frac{1}{2}gt^2\right)\mathbf{j}$. Therefore $y = 0$ when $t = \sqrt{7/g}$ seconds. At that instant, the ball is $2\sqrt{7/g} \approx 0.94$ ft to the right of the table top. Its coordinates (relative to an origin on the floor directly under the table's edge) are $(0.94, 0)$. At impact, the velocity is $\mathbf{v} = 2\,\mathbf{i} - \sqrt{7g}\,\mathbf{j}$, so the speed is $|\mathbf{v}| = \sqrt{4+7g} \approx 15$ ft/s.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) The slope of the curve when $t = \sqrt{\dfrac{7}{g}}$ is $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{-gt}{2} = \dfrac{-g\sqrt{7/g}}{2} = \dfrac{-\sqrt{7g}}{2}$. Thus $\cot\theta = \dfrac{\sqrt{7g}}{2}$ and $\theta \approx 7.6°$.

(c) From (a), $|\mathbf{v}| = \sqrt{4+7g}$. So the ball rebounds with speed $0.8\sqrt{4+7g} \approx 12.08$ ft/s at angle of inclination $90° - \theta \approx 82.3886°$. By Example 13.4.5, the horizontal distance traveled between bounces is $d = \dfrac{v_0^2 \sin 2\alpha}{g}$, where $v_0 \approx 12.08$ ft/s and $\alpha \approx 82.3886°$. Therefore, $d \approx 1.197$ ft. So the ball strikes the floor at about $2\sqrt{7/g} + 1.197 \approx 2.13$ ft to the right of the table's edge.

6. By the Fundamental Theorem of Calculus, $\mathbf{r}'(t) = \left\langle \sin\left(\frac{1}{2}\pi t^2\right), \cos\left(\frac{1}{2}\pi t^2\right)\right\rangle$, $|\mathbf{r}'(t)| = 1$ and so $\mathbf{T}(t) = \mathbf{r}'(t)$. Thus $\mathbf{T}'(t) = \pi t\left\langle \cos\left(\frac{1}{2}\pi t^2\right), -\sin\left(\frac{1}{2}\pi t^2\right)\right\rangle$ and the curvature is $\kappa = |\mathbf{T}'(t)| = \sqrt{(\pi t)^2(1)} = \pi\,|t|$.

7. The trajectory of the projectile is given by $\mathbf{r}(t) = (v\cos\alpha)t\,\mathbf{i} + \left[(v\sin\alpha)t - \frac{1}{2}gt^2\right]\mathbf{j}$, so $\mathbf{v}(t) = \mathbf{r}'(t) = v\cos\alpha\,\mathbf{i} + (v\sin\alpha - gt)\,\mathbf{j}$ and

$$|\mathbf{v}(t)| = \sqrt{(v\cos\alpha)^2 + (v\sin\alpha - gt)^2} = \sqrt{v^2 - (2vg\sin\alpha)\,t + g^2t^2} = \sqrt{g^2\left(t^2 - \frac{2v}{g}(\sin\alpha)\,t + \frac{v^2}{g^2}\right)}$$

$$= g\sqrt{\left(t - \frac{v}{g}\sin\alpha\right)^2 + \frac{v^2}{g^2} - \frac{v^2}{g^2}\sin^2\alpha} = g\sqrt{\left(t - \frac{v}{g}\sin\alpha\right)^2 + \frac{v^2}{g^2}\cos^2\alpha}$$

The projectile hits the ground when $(v\sin\alpha)t - \frac{1}{2}gt^2 = 0 \quad\Rightarrow\quad t = \frac{2v}{g}\sin\alpha$, so the distance traveled by the projectile is

$$L(\alpha) = \int_0^{(2v/g)\sin\alpha} |\mathbf{v}(t)|\,dt = \int_0^{(2v/g)\sin\alpha} g\sqrt{\left(t - \frac{v}{g}\sin\alpha\right)^2 + \frac{v^2}{g^2}\cos^2\alpha}\,dt$$

$$= g\left[\frac{t - (v/g)\sin\alpha}{2}\sqrt{\left(t - \frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2}\right.$$

$$\left. + \frac{[(v/g)\cos\alpha]^2}{2}\ln\left(t - \frac{v}{g}\sin\alpha + \sqrt{\left(t - \frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2}\right)\right]_0^{(2v/g)\sin\alpha}$$

[using Formula 21 in the Table of Integrals]

$$= \frac{g}{2}\left[\frac{v}{g}\sin\alpha\sqrt{\left(\frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2} + \left(\frac{v}{g}\cos\alpha\right)^2\ln\left(\frac{v}{g}\sin\alpha + \sqrt{\left(\frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2}\right)\right.$$

$$\left. + \frac{v}{g}\sin\alpha\sqrt{\left(\frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2} - \left(\frac{v}{g}\cos\alpha\right)^2\ln\left(-\frac{v}{g}\sin\alpha + \sqrt{\left(\frac{v}{g}\sin\alpha\right)^2 + \left(\frac{v}{g}\cos\alpha\right)^2}\right)\right]$$

$$= \frac{g}{2}\left[\frac{v}{g}\sin\alpha\cdot\frac{v}{g} + \frac{v^2}{g^2}\cos^2\alpha\ln\left(\frac{v}{g}\sin\alpha + \frac{v}{g}\right) + \frac{v}{g}\sin\alpha\cdot\frac{v}{g} - \frac{v^2}{g^2}\cos^2\alpha\ln\left(-\frac{v}{g}\sin\alpha + \frac{v}{g}\right)\right]$$

$$= \frac{v^2}{g}\sin\alpha + \frac{v^2}{2g}\cos^2\alpha\ln\left(\frac{(v/g)\sin\alpha + v/g}{-(v/g)\sin\alpha + v/g}\right) = \frac{v^2}{g}\sin\alpha + \frac{v^2}{2g}\cos^2\alpha\ln\left(\frac{1+\sin\alpha}{1-\sin\alpha}\right)$$

We want to maximize $L(\alpha)$ for $0 \le \alpha \le \pi/2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$L'(\alpha) = \frac{v^2}{g}\cos\alpha + \frac{v^2}{2g}\left[\cos^2\alpha \cdot \frac{1-\sin\alpha}{1+\sin\alpha} \cdot \frac{2\cos\alpha}{(1-\sin\alpha)^2} - 2\cos\alpha\,\sin\alpha\,\ln\left(\frac{1+\sin\alpha}{1-\sin\alpha}\right)\right]$$

$$= \frac{v^2}{g}\cos\alpha + \frac{v^2}{2g}\left[\cos^2\alpha \cdot \frac{2}{\cos\alpha} - 2\cos\alpha\,\sin\alpha\,\ln\left(\frac{1+\sin\alpha}{1-\sin\alpha}\right)\right]$$

$$= \frac{v^2}{g}\cos\alpha + \frac{v^2}{g}\cos\alpha\left[1-\sin\alpha\,\ln\left(\frac{1+\sin\alpha}{1-\sin\alpha}\right)\right] = \frac{v^2}{g}\cos\alpha\left[2-\sin\alpha\,\ln\left(\frac{1+\sin\alpha}{1-\sin\alpha}\right)\right]$$

$L(\alpha)$ has critical points for $0 < \alpha < \pi/2$ when $L'(\alpha) = 0 \quad \Rightarrow \quad 2 - \sin\alpha \ln\left(\dfrac{1+\sin\alpha}{1-\sin\alpha}\right) = 0$ [since $\cos\alpha \neq 0$].

Solving by graphing (or using a CAS) gives $\alpha \approx 0.9855$. Compare values at the critical point and the endpoints: $L(0) = 0$, $L(\pi/2) = v^2/g$, and $L(0.9855) \approx 1.20v^2/g$. Thus the distance traveled by the projectile is maximized for $\alpha \approx 0.9855$ or $\approx 56°$.

8. As the cable is wrapped around the spool, think of the top or bottom of the cable forming a helix of radius $R + r$. Let h be the vertical distance between coils. Then, from similar triangles,

$$\frac{2r}{\sqrt{h^2-4r^2}} = \frac{2\pi(r+R)}{h} \quad \Rightarrow \quad h^2r^2 = \pi^2(r+R)^2(h^2-4r^2) \quad \Rightarrow$$

$$h = \frac{2\pi r(r+R)}{\sqrt{\pi^2(r+R)^2 - r^2}}.$$

If we parametrize the helix by $x(t) = (R+r)\cos t$, $y(t) = (R+r)\sin t$, then we must have $z(t) = [h/(2\pi)]t$. The length of one complete cycle is

$$\ell = \int_0^{2\pi} \sqrt{[x'(t)]^2 + [y'(t)]^2 + [z'(t)]^2}\,dt = \int_0^{2\pi} \sqrt{(R+r)^2 + \left(\frac{h}{2\pi}\right)^2}\,dt = 2\pi\sqrt{(R+r)^2 + \left(\frac{h}{2\pi}\right)^2}$$

$$= 2\pi\sqrt{(R+r)^2 + \frac{r^2(R+r)^2}{\pi^2(R+r)^2 - r^2}} = 2\pi(R+r)\sqrt{1 + \frac{r^2}{\pi^2(R+r)^2 - r^2}} = \frac{2\pi^2(R+r)^2}{\sqrt{\pi^2(R+r)^2 - r^2}}$$

The number of complete cycles is $[\![L/\ell]\!]$, and so the shortest length along the spool is

$$h\left[\!\!\left[\frac{L}{\ell}\right]\!\!\right] = \frac{2\pi r(R+r)}{\sqrt{\pi^2(R+r)^2 - r^2}}\left[\!\!\left[\frac{L\sqrt{\pi^2(R+r)^2 - r^2}}{2\pi^2(R+r)^2}\right]\!\!\right]$$

9. We can write the vector equation as $\mathbf{r}(t) = \mathbf{a}t^2 + \mathbf{b}t + \mathbf{c}$ where $\mathbf{a} = \langle a_1, a_2, a_3\rangle$, $\mathbf{b} = \langle b_1, b_2, b_3\rangle$, and $\mathbf{c} = \langle c_1, c_2, c_3\rangle$. Then $\mathbf{r}'(t) = 2t\,\mathbf{a} + \mathbf{b}$ which says that each tangent vector is the sum of a scalar multiple of $\mathbf{a}$ and the vector $\mathbf{b}$. Thus the tangent vectors are all parallel to the plane determined by $\mathbf{a}$ and $\mathbf{b}$ so the curve must be parallel to this plane. [Here we assume that $\mathbf{a}$ and $\mathbf{b}$ are nonparallel. Otherwise the tangent vectors are all parallel and the curve lies along a single line.] A normal vector for the plane is $\mathbf{a} \times \mathbf{b} = \langle a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1\rangle$. The point (c_1, c_2, c_3) lies on the plane (when $t = 0$), so an equation of the plane is

$$(a_2b_3 - a_3b_2)(x - c_1) + (a_3b_1 - a_1b_3)(y - c_2) + (a_1b_2 - a_2b_1)(z - c_3) = 0$$

or

$$(a_2b_3 - a_3b_2)x + (a_3b_1 - a_1b_3)y + (a_1b_2 - a_2b_1)z = a_2b_3c_1 - a_3b_2c_1 + a_3b_1c_2 - a_1b_3c_2 + a_1b_2c_3 - a_2b_1c_3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14 ☐ PARTIAL DERIVATIVES

14.1 Functions of Several Variables

1. (a) $f(x,y) = \dfrac{x^2y}{2x-y^2} \quad \Rightarrow \quad f(1,3) = \dfrac{1^2(3)}{2(1)-3^2} = -\dfrac{3}{7}$

(b) $f(-2,-1) = \dfrac{(-2)^2(-1)}{2(-2)-(-1)^2} = \dfrac{4}{5}$

(c) $f(x+h,y) = \dfrac{(x+h)^2y}{2(x+h)-y^2}$

(d) $f(x,x) = \dfrac{x^2x}{2x-x^2} = \dfrac{x^3}{x(2-x)} = \dfrac{x^2}{2-x}$

2. (a) $g(x,y) = x\sin y + y\sin x \quad \Rightarrow \quad g(\pi,0) = \pi\sin 0 + 0\sin\pi = \pi\cdot 0 + 0\cdot 0 = 0$

(b) $g\left(\dfrac{\pi}{2},\dfrac{\pi}{4}\right) = \dfrac{\pi}{2}\sin\dfrac{\pi}{4} + \dfrac{\pi}{4}\sin\dfrac{\pi}{2} = \dfrac{\pi}{2}\left(\dfrac{\sqrt{2}}{2}\right) + \dfrac{\pi}{4}(1) = \dfrac{\pi\left(\sqrt{2}+1\right)}{4}$

(c) $g(0,y) = 0\sin y + y\sin 0 = 0 + y\cdot 0 = 0$

(d) $g(x,y+h) = x\sin(y+h) + (y+h)\sin x$

3. (a) $g(x,y) = x^2\ln(x+y) \quad \Rightarrow \quad g(3,1) = 3^2\ln(3+1) = 9\ln 4$

(b) $\ln(x+y)$ is defined only when $x+y>0 \quad \Rightarrow \quad y>-x$.

Thus, the domain of g is $\{(x,y)\mid y>-x\}$.

(c) The range of $\ln(x+y)$ is $\mathbb{R}$, so the range of g is $\mathbb{R}$.

4. (a) $h(x,y) = e^{\sqrt{y-x^2}} \quad \Rightarrow \quad h(-2,5) = e^{\sqrt{5-(-2)^2}} = e^{\sqrt{1}} = e$

(b) $\sqrt{y-x^2}$ is defined only when $y-x^2\geq 0 \quad \Rightarrow \quad y\geq x^2$.

Thus, the domain of h is $\{(x,y)\mid y\geq x^2\}$.

(c) We know $\sqrt{y-x^2}\geq 0 \quad \Rightarrow \quad e^{\sqrt{y-x^2}}\geq 1$. Thus, the range of h is $[1,\infty]$.

5. (a) $F(x,y,z) = \sqrt{y}-\sqrt{x-2z} \quad \Rightarrow \quad F(3,4,1) = \sqrt{4}-\sqrt{3-2(1)} = 2-1 = 1$

(b) $\sqrt{y}$ is defined only when $y\geq 0$. $\sqrt{x-2z}$ is defined only when $x-2z\geq 0 \quad \Rightarrow \quad z\leq \frac{1}{2}x$. Thus, the domain is $\{(x,y,z)\mid x\geq 2z, y\geq 0\}$, which is the set of points on or below the plane $z=\frac{1}{2}x$ and on or to the right of the xz-plane.

6. (a) $f(x,y,z) = \ln\left(z-\sqrt{x^2+y^2}\right) \quad \Rightarrow \quad f(4,-3,6) = \ln\left(6-\sqrt{4^2+(-3)^2}\right) = \ln\left(6-\sqrt{25}\right) = \ln 1 = 0$

(b) $\ln\left(z-\sqrt{x^2+y^2}\right)$ is defined only when $z-\sqrt{x^2+y^2}>0 \quad \Leftrightarrow \quad z>\sqrt{x^2+y^2} \quad \Rightarrow \quad z^2>x^2+y^2$. Thus, the domain is $\{(x,y,z)\mid z>\sqrt{x^2+y^2}\}$, which is the set of points inside the top half of the cone $z^2=x^2+y^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. $f(x,y) = \sqrt{x-2} + \sqrt{y-1}$. $\sqrt{x-2}$ is defined only when $x - 2 \geq 0$, or $x \geq 2$, and $\sqrt{y-1}$ is defined only when $y - 1 \geq 0$, or $y \geq 1$. So the domain of f is $\{(x,y) \mid x \geq 2,\ y \geq 1\}$.

8. $f(x,y) = \sqrt[4]{x-3y}$. $\sqrt[4]{x-3y}$ is defined only when $x - 3y \geq 0$, or $x \geq 3y$. So the domain of f is $\{(x,y) \mid x \geq 3y\}$ or equivalently $\left\{(x,y) \mid y \leq \frac{1}{3}x\right\}$.

9. $q(x,y) = \sqrt{x} + \sqrt{4 - 4x^2 - y^2}$. $\sqrt{x}$ is defined only when $x \geq 0$. $\sqrt{4 - 4x^2 - y^2}$ is defined only when $4 - 4x^2 - y^2 \geq 0 \quad \Leftrightarrow$ $1 \geq x^2 + \frac{1}{4}y^2$. So the domain of q is $\left\{(x,y) \mid x^2 + \frac{1}{4}y^2 \leq 1, x \geq 0\right\}$.

10. $g(x,y) = \ln(x^2 + y^2 - 9)$. $\ln(x^2 + y^2 - 9)$ is defined only when $x^2 + y^2 - 9 > 0 \quad \Leftrightarrow \quad x^2 + y^2 > 9$. So the domain of g is $\{(x,y) \mid x^2 + y^2 > 9\}$.

11. $g(x,y) = \dfrac{x-y}{x+y}$. g is not defined if $x + y = 0 \quad \Leftrightarrow \quad y = -x$ (and is defined otherwise). Thus, the domain of g is $\{(x,y) \mid y \neq -x\}$, the set of all points in $\mathbb{R}^2$ that are not on the line $y = -x$.

12. $g(x,y) = \dfrac{\ln(2-x)}{1 - x^2 - y^2}$. $\ln(2-x)$ is defined only when $2 - x > 0 \quad \Leftrightarrow$ $x < 2$. In addition, g is not defined if $1 - x^2 - y^2 = 0 \quad \Leftrightarrow \quad x^2 + y^2 = 1$. Thus, the domain of g is $\left\{(x,y) \mid x < 2, x^2 + y^2 \neq 1\right\}$, the set of all points to the left of the line $x = 2$ and not on the unit circle.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $p(x, y) = \dfrac{\sqrt{xy}}{x+1}$. $\sqrt{xy}$ is defined only when $xy \geq 0$. Further, p is defined only when $x + 1 \neq 0 \quad \Leftrightarrow \quad x \neq -1$. So the domain of p is $\{(x, y) \mid xy \geq 0, x \neq -1\}$.

14. $f(x, y) = \sin^{-1}(x + y)$. $\sin^{-1}(x + y)$ is defined only when $-1 \leq x + y \leq 1 \quad \Leftrightarrow \quad -1 - x \leq y \leq 1 - x$. Thus, the domain of f is $\{(x, y) \mid -1 - x \leq y \leq 1 - x\}$, which consists of those points on or between the parallel lines $y = -1 - x$ and $y = 1 - x$.

15. $f(x, y, z) = \sqrt{4 - x^2} + \sqrt{9 - y^2} + \sqrt{1 - z^2}$. f is defined only when $4 - x^2 \geq 0 \quad \Leftrightarrow \quad -2 \leq x \leq 2$, and $9 - y^2 \geq 0 \quad \Leftrightarrow \quad -3 \leq y \leq 3$, and $1 - z^2 \geq 0 \quad \Leftrightarrow \quad -1 \leq z \leq 1$. Thus, the domain of f is $\{(x, y, z) \mid -2 \leq x \leq 2,\ -3 \leq y \leq 3,\ -1 \leq z \leq 1\}$, which is a solid rectangular box with vertices $(\pm 2, \pm 3, \pm 1)$ (all 8 combinations).

16. $f(x, y, z) = \ln(16 - 4x^2 - 4y^2 - z^2)$. f is defined only when $16 - 4x^2 - 4y^2 - z^2 > 0 \quad \Rightarrow \quad \dfrac{x^2}{4} + \dfrac{y^2}{4} + \dfrac{z^2}{16} < 1$. Thus, $D = \left\{(x, y, z) \,\middle|\, \dfrac{x^2}{4} + \dfrac{y^2}{4} + \dfrac{z^2}{16} < 1\right\}$, that is, the points inside the ellipsoid $\dfrac{x^2}{4} + \dfrac{y^2}{4} + \dfrac{z^2}{16} = 1$.

17. (a) $f(160, 70) = 0.1091(160)^{0.425}(70)^{0.725} \approx 20.5$, which means that the surface area of a person 70 inches (5 feet 10 inches) tall who weighs 160 pounds is approximately 20.5 square feet.

(b) Answers will vary depending on the height and weight of the reader.

18. $P(120, 20) = 1.47(120)^{0.65}(20)^{0.35} \approx 94.2$, so when the manufacturer invests \$20 million in capital and 120,000 hours of labor are completed yearly, the monetary value of the production is about \$94.2 million.

19. (a) From Table 1, $f(-15, 40) = -27$, which means that if the temperature is $-15°\text{C}$ and the wind speed is 40 km/h, then the air would feel equivalent to approximately $-27°\text{C}$ without wind.

(b) The question is asking: when the temperature is $-20°\text{C}$, what wind speed gives a wind-chill index of $-30°\text{C}$? From Table 1, the speed is 20 km/h.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) The question is asking: when the wind speed is 20 km/h, what temperature gives a wind-chill index of $-49°\mathrm{C}$? From Table 1, the temperature is $-35°\mathrm{C}$.

(d) The function $W = f(-5, v)$ means that we fix T at -5 and allow v to vary, resulting in a function of one variable. In other words, the function gives wind-chill index values for different wind speeds when the temperature is $-5°\mathrm{C}$. From Table 1 (look at the row corresponding to $T = -5$), the function decreases and appears to approach a constant value as v increases.

(e) The function $W = f(T, 50)$ means that we fix v at 50 and allow T to vary, again giving a function of one variable. In other words, the function gives wind-chill index values for different temperatures when the wind speed is 50 km/h . From Table 1 (look at the column corresponding to $v = 50$), the function increases almost linearly as T increases.

20. (a) From Table 3, $f(95, 70) = 124$, which means that when the actual temperature is $95°\mathrm{F}$ and the relative humidity is 70%, the perceived air temperature is approximately $124°\mathrm{F}$.

(b) Looking at the row corresponding to $T = 90$, we see that $f(90, h) = 100$ when $h = 60$.

(c) Looking at the column corresponding to $h = 50$, we see that $f(T, 50) = 88$ when $T = 85$.

(d) $I = f(80, h)$ means that T is fixed at 80 and h is allowed to vary, resulting in a function of h that gives the humidex values for different relative humidities when the actual temperature is $80°\mathrm{F}$. Similarly, $I = f(100, h)$ is a function of one variable that gives the humidex values for different relative humidities when the actual temperature is $100°\mathrm{F}$. Looking at the rows of the table corresponding to $T = 80$ and $T = 100$, we see that $f(80, h)$ increases at a relatively constant rate of approximately $1°\mathrm{F}$ per 10% relative humidity, while $f(100, h)$ increases more quickly (at first with an average rate of change of $5°\mathrm{F}$ per 10% relative humidity) and at an increasing rate (approximately $12°\mathrm{F}$ per 10% relative humidity for larger values of h).

21. (a) According to Table 4, $f(40, 15) = 25$, which means that if a 40-knot wind has been blowing in the open sea for 15 hours, it will create waves with estimated heights of 25 feet.

(b) $h = f(30, t)$ means we fix v at 30 and allow t to vary, resulting in a function of one variable. Thus here, $h = f(30, t)$ gives the wave heights produced by 30-knot winds blowing for t hours. From the table (look at the row corresponding to $v = 30$), the function increases but at a declining rate as t increases. In fact, the function values appear to be approaching a limiting value of approximately 19, which suggests that 30-knot winds cannot produce waves higher than about 19 feet.

(c) $h = f(v, 30)$ means we fix t at 30, again giving a function of one variable. So, $h = f(v, 30)$ gives the wave heights produced by winds of speed v blowing for 30 hours. From the table (look at the column corresponding to $t = 30$), the function appears to increase at an increasing rate, with no apparent limiting value. This suggests that faster winds (lasting 30 hours) always create higher waves.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. (a) The cost of making x small boxes, y medium boxes, and z large boxes is $C = f(x, y, z) = 8000 + 2.5x + 4y + 4.5z$ dollars.

(b) $f(3000, 5000, 4000) = 8000 + 2.5(3000) + 4(5000) + 4.5(4000) = 53{,}500$ which means that it costs \$53,500 to make 3000 small boxes, 5000 medium boxes, and 4000 large boxes.

(c) Because no partial boxes will be produced, each of x, y, and z must be a positive integer or zero.

23. The graph of f has equation $z = y$, a plane which intersects the yz-plane in the line $z = y$, $x = 0$. The portion of this plane in the first octant is shown.

24. The graph of f has equation $z = x^2$, a parabolic cylinder.

25. $z = 10 - 4x - 5y$ or $4x + 5y + z = 10$, a plane with intercepts 2.5, 2, and 10.

26. $z = \cos y$, a cylinder.

27. $z = \sin x$, a cylinder.

28. $z = 2 - x^2 - y^2$, a circular paraboloid opening downward with vertex at $(0, 0, 2)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. $z = x^2 + 4y^2 + 1$, an elliptic paraboloid opening upward with vertex at $(0, 0, 1)$.

30. $z = \sqrt{4x^2 + y^2}$ so $4x^2 + y^2 = z^2$ and $z \geq 0$, the top half of an elliptic cone.

31. $z = \sqrt{4 - 4x^2 - y^2}$ so $4x^2 + y^2 + z^2 = 4$ or $x^2 + \dfrac{y^2}{4} + \dfrac{z^2}{4} = 1$ and $z \geq 0$, the top half of an ellipsoid.

32. (a) $f(x, y) = \dfrac{1}{1 + x^2 + y^2}$. The trace in $x = 0$ is $z = \dfrac{1}{1 + y^2}$, and the trace in $y = 0$ is $z = \dfrac{1}{1 + x^2}$. The only possibility is graph III. Notice also that the level curves of f are $\dfrac{1}{1 + x^2 + y^2} = k \quad \Leftrightarrow \quad x^2 + y^2 = \dfrac{1}{k} - 1$, a family of circles for $k < 1$.

(b) $f(x, y) = \dfrac{1}{1 + x^2 y^2}$. The trace in $x = 0$ is the horizontal line $z = 1$, and the trace in $y = 0$ is also $z = 1$. Both graphs I and II have these traces; however, notice that here $z > 0$, so the graph is I.

(c) $f(x, y) = \ln(x^2 + y^2)$. The trace in $x = 0$ is $z = \ln y^2$, and the trace in $y = 0$ is $z = \ln x^2$. The level curves of f are $\ln(x^2 + y^2) = k \quad \Leftrightarrow \quad x^2 + y^2 = e^k$, a family of circles. In addition, f is large negative when $x^2 + y^2$ is small, so this is graph IV.

(d) $f(x, y) = \cos\sqrt{x^2 + y^2}$. The trace in $x = 0$ is $z = \cos\sqrt{y^2} = \cos|y| = \cos y$, and the trace in $y = 0$ is $z = \cos\sqrt{x^2} = \cos|x| = \cos x$. Notice also that the level curve $f(x, y) = 0$ is $\cos\sqrt{x^2 + y^2} = 0 \quad \Leftrightarrow$ $x^2 + y^2 = \left(\frac{\pi}{2} + n\pi\right)^2$, a family of circles, so this is graph V.

(e) $f(x, y) = |xy|$. The trace in $x = 0$ is $z = 0$, and the trace in $y = 0$ is $z = 0$, so it must be graph VI.

(f) $f(x, y) = \cos(xy)$. The trace in $x = 0$ is $z = \cos 0 = 1$, and the trace in $y = 0$ is $z = 1$. As mentioned in part (b), these traces match both graphs I and II. Here z can be negative, so the graph is II. (Also notice that the trace in $x = 1$ is $z = \cos y$, and the trace in $y = 1$ is $z = \cos x$.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. The point $(-3, 3)$ lies between the level curves with z-values 50 and 60. Since the point is a little closer to the level curve with $z = 60$, we estimate that $f(-3, 3) \approx 56$. The point $(3, -2)$ appears to be just about halfway between the level curves with z-values 30 and 40, so we estimate $f(3, -2) \approx 35$. The graph rises as we approach the origin, gradually from above, steeply from below.

34. (a) C (Chicago) lies between level curves with pressures 1012 and 1016 mb, and since C appears to be located about one-fourth the distance from the 1012 mb isobar to the 1016 mb isobar, we estimate the pressure at Chicago to be about 1013 mb. N lies very close to a level curve with pressure 1012 mb so we estimate the pressure at Nashville to be approximately 1012 mb. S appears to be just about halfway between level curves with pressures 1008 and 1012 mb, so we estimate the pressure at San Francisco to be about 1010 mb. V lies close to a level curve with pressure 1016 mb but we can't see a level curve to its left so it is more difficult to make an accurate estimate. There are lower pressures to the right of V and V is a short distance to the left of the level curve with pressure 1016 mb, so we might estimate that the pressure at Vancouver is about 1017 mb.

(b) Winds are stronger where the isobars are closer together (see Figure 13), and the level curves are closer near S than at the other locations, so the winds were strongest at San Francisco.

35. The point $(160, 10)$, corresponding to day 160 and a depth of 10 m, lies between the isothermals with temperature values of 8 and $12°\text{C}$. Since the point appears to be located about three-fourths the distance from the $8°\text{C}$ isothermal to the $12°\text{C}$ isothermal, we estimate the temperature at that point to be approximately $11°\text{C}$. The point $(180, 5)$ lies between the 16 and $20°\text{C}$ isothermals, very close to the $20°\text{C}$ level curve, so we estimate the temperature there to be about $19.5°\text{C}$.

36. If we start at the origin and move along the x-axis, for example, the z-values of a cone centered at the origin increase at a constant rate, so we would expect its level curves to be equally spaced. A paraboloid with vertex the origin, on the other hand, has z-values which change slowly near the origin and more quickly as we move farther away. Thus, we would expect its level curves near the origin to be spaced more widely apart than those farther from the origin. Therefore contour map I must correspond to the paraboloid, and contour map II the cone.

37. Near A, the level curves are very close together, indicating that the terrain is quite steep. At B, the level curves are much farther apart, so we would expect the terrain to be much less steep than near A, perhaps almost flat.

38.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. The level curves of $B(m,h) = \dfrac{m}{h^2}$ are $\dfrac{m}{h^2} = k \quad \Leftrightarrow \quad m = kh^2$ or equivalently $h = \sqrt{m/k} = \dfrac{1}{\sqrt{k}}\sqrt{m}$ since $m > 0$, $h > 0$. We draw the level curves for $k = 18.5$, 25, 30, and 40.

The shaded region corresponds to BMI values between 18.5 and 25, those considered optimal. For a mass of 62 kg and a height of 152 cm (1.52 m), the BMI is $B(62, 1.52) = \dfrac{62}{1.52^2} \approx 26.8$, which is outside the optimal range.

40. From Exercise 39, the body mass index function is $B(m,h) = m/h^2$. The BMI for a person 200 cm (2.0 m, about 6 ft 7 in) tall and with mass 80 kg (about 176 lb) is $B(80, 2.0) = 80/(2.0)^2 = 20$. The level curve $B(m,h) = 20 \quad \Leftrightarrow \quad m = 20h^2$ is shown in the graph.

A person 1.5 m tall (about 4 ft 11 in) has a BMI on the same level curve if their mass is $m = 20(1.5)^2 = 45$ kg (about 99 lb), and a person 1.8 m (about 5 ft 11 in) tall would have mass $m = 20(1.8)^2 = 64.8$ kg (about 143 lb). (See the graph.)

41.

42.

43.

44.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. The level curves are $x^2 - y^2 = k$. When $k = 0$ the level curve is the pair of lines $y = \pm x$, and when $k \neq 0$ the level curves are a family of hyperbolas (oriented differently for $k > 0$ than for $k < 0$).

46. The level curves are $xy = k$ or $y = k/x$. When $k \neq 0$ the level curves are a family of hyperbolas. When $k = 0$ the level curve is the pair of lines $x = 0$, $y = 0$.

47. The level curves are $\sqrt{x} + y = k$ or $y = -\sqrt{x} + k$, a family of vertical translations of the graph of the root function $y = -\sqrt{x}$.

48. The level curves are $\ln(x^2 + 4y^2) = k$ or $x^2 + 4y^2 = e^k$, a family of ellipses.

49. The level curves are $ye^x = k$ or $y = ke^{-x}$, a family of exponential curves.

50. The level curves are $y - \arctan x = k$ or $y = (\arctan x) + k$, a family of vertical translations of the graph of the inverse tangent function $y = \arctan x$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. The level curves are $\sqrt[3]{x^2+y^2}=k$ or $x^2+y^2=k^3$ ($k \geq 0$), a family of circles centered at the origin with radius $k^{3/2}$.

52. For $k \neq 0$ and $(x, y) \neq (0, 0)$, $k = \dfrac{y}{x^2+y^2} \quad \Leftrightarrow$

$x^2+y^2-\dfrac{y}{k}=0 \quad \Leftrightarrow \quad x^2+\left(y-\frac{1}{2k}\right)^2=\dfrac{1}{4k^2}$, a family of circles with center $\left(0, \frac{1}{2k}\right)$ and radius $\frac{1}{2k}$ (without the origin). If $k=0$, the level curve is the x-axis.

53. The contour map consists of the level curves $k=x^2+9y^2$, a family of ellipses with major axis the x-axis. (Or, if $k=0$, the origin.)

The graph of $f(x, y)$ is the surface $z=x^2+9y^2$, an elliptic paraboloid.

If we visualize lifting each ellipse $k=x^2+9y^2$ of the contour map to the plane $z=k$, we have horizontal traces that indicate the shape of the graph of f.

54.

The contour map consists of the level curves $k=\sqrt{36-9x^2-4y^2} \quad \Rightarrow$ $9x^2+4y^2=36-k^2$, $k \geq 0$, a family of ellipses with major axis the y-axis. (Or, if $k=6$, the origin.)

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The graph of $f(x, y)$ is the surface $z = \sqrt{36 - 9x^2 - 4y^2}$, or equivalently the upper half of the ellipsoid $9x^2 + 4y^2 + z^2 = 36$. If we visualize lifting each ellipse $k = \sqrt{36 - 9x^2 - 4y^2}$ of the contour map to the plane $z = k$, we have horizontal traces that indicate the shape of the graph of f.

55. The isothermals are given by $k = 100/(1 + x^2 + 2y^2)$ or $x^2 + 2y^2 = (100 - k)/k$ $[0 < k \leq 100]$, a family of ellipses.

56. The equipotential curves are $k = \dfrac{c}{\sqrt{r^2 - x^2 - y^2}}$ or $x^2 + y^2 = r^2 - \left(\dfrac{c}{k}\right)^2$, a family of circles $(k \geq c/r)$.

Note: As $k \to \infty$, the radius of the circle approaches r.

57. $f(x, y) = xy^2 - x^3$

The traces parallel to the yz-plane (such as the left-front trace in the graph above) are parabolas; those parallel to the xz-plane (such as the right-front trace) are cubic curves. The surface is called a monkey saddle because a monkey sitting on the surface near the origin has places for both legs and tail to rest.

58. $f(x, y) = xy^3 - yx^3$

The traces parallel to either the yz-plane or the xz-plane are cubic curves.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. $f(x, y) = e^{-(x^2+y^2)/3}\left(\sin(x^2) + \cos(y^2)\right)$

60. $f(x, y) = \cos x \, \cos y$

The traces parallel to either the yz- or xz-plane are cosine curves with amplitudes that vary from 0 to 1.

61. $z = \sin(xy)$ (a) C (b) II

Reasons: This function is periodic in both x and y, and the function is the same when x is interchanged with y, so its graph is symmetric about the plane $y = x$. In addition, the function is 0 along the x- and y-axes. These conditions are satisfied only by C and II.

62. $z = e^x \cos y$ (a) A (b) IV

Reasons: This function is periodic in y but not x, a condition satisfied only by A and IV. Also, note that traces in $x = k$ are cosine curves with amplitude that increases as x increases.

63. $z = \sin(x - y)$ (a) F (b) I

Reasons: This function is periodic in both x and y but is constant along the lines $y = x + k$, a condition satisfied only by F and I.

64. $z = \sin x - \sin y$ (a) E (b) III

Reasons: This function is periodic in both x and y, but unlike the function in Exercise 63, it is not constant along lines such as $y = x + \pi$, so the contour map is III. Also notice that traces in $y = k$ are vertically shifted copies of the sine wave $z = \sin x$, so the graph must be E.

65. $z = (1 - x^2)(1 - y^2)$ (a) B (b) VI

Reasons: This function is 0 along the lines $x = \pm 1$ and $y = \pm 1$. The only contour map in which this could occur is VI. Also note that the trace in the xz-plane is the parabola $z = 1 - x^2$ and the trace in the yz-plane is the parabola $z = 1 - y^2$, so the graph is B.

66. $z = \dfrac{x - y}{1 + x^2 + y^2}$ (a) D (b) V

Reasons: This function is not periodic, ruling out the graphs in A, C, E, and F. Also, the values of z approach 0 as we use points farther from the origin. The only graph that shows this behavior is D, which corresponds to V.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. $k = 2y - z + 1$ is a family of parallel planes with normal vector $\langle 0, 2, -1 \rangle$.

68. $k = x + y^2 - z^2$ is a family of hyperbolic paraboloids with saddle point $(k, 0, 0)$.

69. Equations for the level surfaces are $k = x^2 + y^2 - z^2$. For $k = 0$, the equation becomes $z^2 = x^2 + y^2$ and the surface is a right circular cone with center the origin and axis the z-axis. For $k > 0$, we have a family of hyperboloids of one sheet with axis the z-axis. For $k < 0$ we have a family of hyperboloids of two sheets with axis the z-axis.

70. $k = x^2 + 2y^2 + 3z^2$ is a family of ellipsoids with major axis the x-axis for $k > 0$ and the origin for $k = 0$.

71. (a) The graph of g is the graph of f shifted upward 2 units.

(b) The graph of g is the graph of f stretched vertically by a factor of 2.

(c) The graph of g is the graph of f reflected about the xy-plane.

(d) The graph of $g(x, y) = -f(x, y) + 2$ is the graph of f reflected about the xy-plane and then shifted upward 2 units.

72. (a) The graph of g is the graph of f shifted 2 units in the positive x-direction.

(b) The graph of g is the graph of f shifted 2 units in the negative y-direction.

(c) The graph of g is the graph of f shifted 3 units in the negative x-direction and 4 units in the positive y-direction.

73. $f(x, y) = 3x - x^4 - 4y^2 - 10xy$

Three-dimensional view

Front view

It does appear that the function has a maximum value, at the higher of the two "hilltops." From the front view graph, the maximum value appears to be approximately 15. Both hilltops could be considered local maximum points, as the values of f there are larger than at the neighboring points. There does not appear to be any local minimum point; although the valley shape between the two peaks looks like a minimum of some kind, some neighboring points have lower function values.

74. $f(x, y) = xye^{-x^2-y^2}$

Three-dimensional view

Front view

The function does have a maximum value, which it appears to achieve at two different points (the two "hilltops"). From the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

front view graph, we can estimate the maximum value to be approximately 0.18. These same two points can also be considered local maximum points. The two "valley bottoms" visible in the graph can be considered local minimum points, as all the neighboring points give greater values of f.

75.

$f(x,y) = \dfrac{x+y}{x^2+y^2}$. As both x and y become large, the function values appear to approach 0, regardless of which direction is considered. As (x,y) approaches the origin, the graph exhibits asymptotic behavior. From some directions, $f(x,y) \to \infty$, while in others $f(x,y) \to -\infty$. (These are the vertical spikes visible in the graph.) If the graph is examined carefully, however, one can see that $f(x,y)$ approaches 0 along the line $y = -x$.

76.

$f(x,y) = \dfrac{xy}{x^2+y^2}$. The graph exhibits different limiting values as x and y become large or as (x,y) approaches the origin, depending on the direction being examined. For example, although f is undefined at the origin, the function values appear to be $\frac{1}{2}$ along the line $y = x$, regardless of the distance from the origin. Along the line $y = -x$, the value is always $-\frac{1}{2}$. Along the axes, $f(x,y) = 0$ for all values of (x,y) except the origin. Other directions, heading toward the origin or away from the origin, give various limiting values between $-\frac{1}{2}$ and $\frac{1}{2}$.

77. $f(x,y) = e^{cx^2+y^2}$. First, if $c = 0$, the graph is the cylindrical surface $z = e^{y^2}$ (whose level curves are parallel lines). When $c > 0$, the vertical trace above the y-axis remains fixed while the sides of the surface in the x-direction "curl" upward, giving the graph a shape resembling an elliptic paraboloid. The level curves of the surface are ellipses centered at the origin.

$c = 0$

For $0 < c < 1$, the ellipses have major axis the x-axis and the eccentricity increases as $c \to 0$.

$c = 0.5$ (level curves in increments of 1)

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

For $c = 1$ the level curves are circles centered at the origin.

$c = 1$ (level curves in increments of 1)

When $c > 1$, the level curves are ellipses with major axis the y-axis, and the eccentricity increases as c increases.

$c = 2$ (level curves in increments of 4)

For values of $c < 0$, the sides of the surface in the x-direction curl downward and approach the xy-plane (while the vertical trace $x = 0$ remains fixed), giving a saddle-shaped appearance to the graph near the point $(0, 0, 1)$. The level curves consist of a family of hyperbolas. As c decreases, the surface becomes flatter in the x-direction and the surface's approach to the curve in the trace $x = 0$ becomes steeper, as the graphs demonstrate.

$c = -0.5$ (level curves in increments of 0.25)

$c = -2$ (level curves in increments of 0.25)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

78. $z = (ax^2 + by^2)e^{-x^2-y^2}$. There are only three basic shapes which can be obtained (the fourth and fifth graphs are the reflections of the first and second ones in the xy-plane). Interchanging a and b rotates the graph by $90°$ about the z-axis.

$a = 1, b = 1$

$a = 2, b = 1$

$a = 1, b = -1$

$a = -1, b = -1$

$a = -2, b = -1$

If a and b are both positive ($a \neq b$), we see that the graph has two maximum points whose height increases as a and b increase. If a and b have opposite signs, the graph has two maximum points and two minimum points, and if a and b are both negative, the graph has one maximum point and two minimum points.

79. $z = x^2 + y^2 + cxy$. When $c < -2$, the surface intersects the plane $z = k \neq 0$ in a hyperbola. (See the following graph.) It intersects the plane $x = y$ in the parabola $z = (2 + c)x^2$, and the plane $x = -y$ in the parabola $z = (2 - c)x^2$. These parabolas open in opposite directions, so the surface is a hyperbolic paraboloid.

When $c = -2$ the surface is $z = x^2 + y^2 - 2xy = (x - y)^2$. So the surface is constant along each line $x - y = k$. That is, the surface is a cylinder with axis $x - y = 0$, $z = 0$. The shape of the cylinder is determined by its intersection with the

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

plane $x + y = 0$, where $z = 4x^2$, and hence the cylinder is parabolic with minimums of 0 on the line $y = x$.

$c = -5$, $z = 2$

$c = -10$

$c = -2$

When $-2 < c \leq 0$, $z \geq 0$ for all x and y. If x and y have the same sign, then $x^2 + y^2 + cxy \geq x^2 + y^2 - 2xy = (x - y)^2 \geq 0$. If they have opposite signs, then $cxy \geq 0$. The intersection with the surface and the plane $z = k > 0$ is an ellipse (see graph below). The intersection with the surface and the planes $x = 0$ and $y = 0$ are parabolas $z = y^2$ and $z = x^2$ respectively, so the surface is an elliptic paraboloid.

When $c > 0$ the graphs have the same shape, but are reflected in the plane $x = 0$, because $x^2 + y^2 + cxy = (-x)^2 + y^2 + (-c)(-x)y$. That is, the value of z is the same for c at (x, y) as it is for $-c$ at $(-x, y)$.

$c = -1$, $z = 2$

$c = 0$

$c = 10$

So the surface is an elliptic paraboloid for $0 < c < 2$, a parabolic cylinder for $c = 2$, and a hyperbolic paraboloid for $c > 2$.

80. First, we graph $f(x, y) = \sqrt{x^2 + y^2}$.

$f(x, y) = \sqrt{x^2 + y^2}$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Graphs of the other four functions follow.

$f(x, y) = e^{\sqrt{x^2 + y^2}}$

$f(x, y) = \ln \sqrt{x^2 + y^2}$

$f(x, y) = \sin\left(\sqrt{x^2 + y^2}\right)$

$f(x, y) = \dfrac{1}{\sqrt{x^2 + y^2}}$

Notice that each graph $f(x, y) = g\left(\sqrt{x^2 + y^2}\right)$ exhibits radial symmetry about the z-axis and the trace in the xz-plane for $x \geq 0$ is the graph of $z = g(x)$, $x \geq 0$. This suggests that the graph of $f(x, y) = g\left(\sqrt{x^2 + y^2}\right)$ is obtained from the graph of g by graphing $z = g(x)$ in the xz-plane and rotating the curve about the z-axis.

81. (a) $P = bL^{\alpha}K^{1-\alpha} \quad \Rightarrow \quad \dfrac{P}{K} = bL^{\alpha}K^{-\alpha} \quad \Rightarrow \quad \dfrac{P}{K} = b\left(\dfrac{L}{K}\right)^{\alpha} \quad \Rightarrow \quad \ln \dfrac{P}{K} = \ln\left(b\left(\dfrac{L}{K}\right)^{\alpha}\right) \quad \Rightarrow$

$\ln \dfrac{P}{K} = \ln b + \alpha \ln\left(\dfrac{L}{K}\right)$

(b) We list the values for $\ln(L/K)$ and $\ln(P/K)$ for the years 1899–1922. (Historically, these values were rounded to 2 decimal places.)

Year	$x = \ln(L/K)$	$y = \ln(P/K)$
1899	0	0
1900	−0.02	−0.06
1901	−0.04	−0.02
1902	−0.04	0
1903	−0.07	−0.05
1904	−0.13	−0.12
1905	−0.18	−0.04
1906	−0.20	−0.07
1907	−0.23	−0.15
1908	−0.41	−0.38
1909	−0.33	−0.24
1910	−0.35	−0.27

Year	$x = \ln(L/K)$	$y = \ln(P/K)$
1911	−0.38	−0.34
1912	−0.38	−0.24
1913	−0.41	−0.25
1914	−0.47	−0.37
1915	−0.53	−0.34
1916	−0.49	−0.28
1917	−0.53	−0.39
1918	−0.60	−0.50
1919	−0.68	−0.57
1920	−0.74	−0.57
1921	−1.05	−0.85
1922	−0.98	−0.59

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

After entering the (x, y) pairs into a calculator or CAS, the resulting least squares regression line through the points is approximately $y = 0.75136x + 0.01053$, which we round to $y = 0.75x + 0.01$.

(c) Comparing the regression line from part (b) to the equation $y = \ln b + \alpha x$ with $x = \ln(L/K)$ and $y = \ln(P/K)$, we have $\alpha = 0.75$ and $\ln b = 0.01 \quad \Rightarrow \quad b = e^{0.01} \approx 1.01$. Thus, the Cobb-Douglas production function is $P = bL^{\alpha}K^{1-\alpha} = 1.01L^{0.75}K^{0.25}$.

14.2 Limits and Continuity

1. In general, we can't say anything about $f(3, 1)$! $\lim\limits_{(x,y)\to(3,1)} f(x, y) = 6$ means that the values of $f(x, y)$ approach 6 as (x, y) approaches, but is not equal to, $(3, 1)$. If f is continuous, we know that $\lim\limits_{(x,y)\to(a,b)} f(x, y) = f(a, b)$, so $\lim\limits_{(x,y)\to(3,1)} f(x, y) = f(3, 1) = 6$.

2. (a) The outdoor temperature as a function of longitude, latitude, and time is continuous. Small changes in longitude, latitude, or time can produce only small changes in temperature, as the temperature doesn't jump abruptly from one value to another.

(b) Elevation is not necessarily continuous. If we think of a cliff with a sudden drop-off, a very small change in longitude or latitude can produce a comparatively large change in elevation, without all the intermediate values being attained. Elevation *can* jump from one value to another.

(c) The cost of a taxi ride is usually discontinuous. The cost normally increases in jumps, so small changes in distance traveled or time can produce a jump in cost. A graph of the function would show breaks in the surface.

3. We make a table of values of $f(x, y) = \dfrac{x^2y^3 + x^3y^2 - 5}{2 - xy}$ for a set of (x, y) points near the origin.

x \ y	-0.2	-0.1	-0.05	0	0.05	0.1	0.2
-0.2	-2.551	-2.525	-2.513	-2.500	-2.488	-2.475	-2.451
-0.1	-2.525	-2.513	-2.506	-2.500	-2.494	-2.488	-2.475
-0.05	-2.513	-2.506	-2.503	-2.500	-2.497	-2.494	-2.488
0	-2.500	-2.500	-2.500		-2.500	-2.500	-2.500
0.05	-2.488	-2.494	-2.497	-2.500	-2.503	-2.506	-2.513
0.1	-2.475	-2.488	-2.494	-2.500	-2.506	-2.513	-2.525
0.2	-2.451	-2.475	-2.488	-2.500	-2.513	-2.525	-2.551

As the table shows, the values of $f(x, y)$ seem to approach -2.5 as (x, y) approaches the origin from a variety of different directions. This suggests that $\lim\limits_{(x,y)\to(0,0)} f(x, y) = -2.5$. Since f is a rational function, it is continuous on its domain. f is defined at $(0, 0)$, so we can use direct substitution to establish that $\lim\limits_{(x,y)\to(0,0)} f(x, y) = \dfrac{0^2 0^3 + 0^3 0^2 - 5}{2 - 0 \cdot 0} = -\dfrac{5}{2}$, verifying our guess.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. We make a table of values of $f(x,y) = \dfrac{2xy}{x^2+2y^2}$ for a set of (x,y) points near the origin.

x \ y	-0.3	-0.2	-0.1	0	0.1	0.2	0.3
-0.3	0.667	0.706	0.545	0.000	-0.545	-0.706	-0.667
-0.2	0.545	0.667	0.667	0.000	-0.667	-0.667	-0.545
-0.1	0.316	0.444	0.667	0.000	-0.667	-0.444	-0.316
0	0.000	0.000	0.000		0.000	0.000	0.000
0.1	-0.316	-0.444	-0.667	0.000	0.667	0.444	0.316
0.2	-0.545	-0.667	-0.667	0.000	0.667	0.667	0.545
0.3	-0.667	-0.706	-0.545	0.000	0.545	0.706	0.667

It appears from the table that the values of $f(x,y)$ are not approaching a single value as (x,y) approaches the origin. For verification, if we first approach $(0,0)$ along the x-axis, we have $f(x,0)=0$, so $f(x,y)\to 0$. But if we approach $(0,0)$ along the line $y=x$, $f(x,x) = \dfrac{2x^2}{x^2+2x^2} = \dfrac{2}{3}$ $(x \neq 0)$, so $f(x,y)\to \frac{2}{3}$. Since f approaches different values along different paths to the origin, this limit does not exist.

5. $f(x,y) = x^2y^3 - 4y^2$ is a polynomial, and hence continuous, so we can find the limit by direct substitution:

$$\lim_{(x,y)\to(3,2)} f(x,y) = f(3,2) = (3)^2(2)^3 - 4(2)^2 = 56.$$

6. $f(x,y) = x^2y + 3xy^2 + 4$ is a polynomial, and hence continuous, so we can find the limit by direct substitution:

$$\lim_{(x,y)\to(5,-2)} f(x,y) = f(5,-2) = 5^2(-2) + 3(5)(-2)^2 + 4 = 14.$$

7. $f(x,y) = \dfrac{x^2y - xy^3}{x-y+2}$ is a rational function, and hence, continuous on its domain. $(-3,1)$ is in the domain of f, so we can find the limit by direct substitution: $\displaystyle\lim_{(x,y)\to(-3,1)} f(x,y) = f(-3,1) = \frac{(-3)^2(1)-(-3)(1)^3}{-3-1+2} = \frac{12}{-2} = -6.$

8. $f(x,y) = \dfrac{x^2y + xy^2}{x^2-y^2}$ is a rational function, and hence, continuous on its domain. $(2,-1)$ is in the domain of f, so we can find the limit by direct substitution: $\displaystyle\lim_{(x,y)\to(2,-1)} f(x,y) = f(2,-1) = \frac{(2)^2(-1)+(2)(-1)^2}{(2)^2-(-1)^2} = -\frac{2}{3}.$

9. $x-y$ is a polynomial and therefore continuous. Since $\sin t$ is a continuous function, the composition $\sin(x-y)$ is also continuous. The function y is a polynomial, and hence continuous, and the product of continuous functions is continuous, so $f(x,y) = y\sin(x-y)$ is a continuous function. Then $\displaystyle\lim_{(x,y)\to(\pi,\pi/2)} f(x,y) = f\left(\pi,\tfrac{\pi}{2}\right) = \tfrac{\pi}{2}\sin\left(\pi - \tfrac{\pi}{2}\right) = \tfrac{\pi}{2}\sin\tfrac{\pi}{2} = \tfrac{\pi}{2}.$

10. $2x-y$ is a polynomial and therefore continuous. Since $\sqrt{t}$ is continuous for $t\geq 0$, the composition $\sqrt{2x-y}$ is continuous where $2x-y\geq 0$. The function e^u is continuous everywhere, so the composition $f(x,y) = e^{\sqrt{2x-y}}$ is a continuous function for $2x-y\geq 0$. If $x=3$ and $y=2$ then $2x-y\geq 0$, so $\displaystyle\lim_{(x,y)\to(3,2)} f(x,y) = f(3,2) = e^{\sqrt{2(3)-2}} = e^2.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. $f(x,y) = \dfrac{x^2y^3 - x^3y^2}{x^2 - y^2} = \dfrac{x^2y^2(y-x)}{(x-y)(x+y)} = -\dfrac{x^2y^2}{x+y}$ for $x - y \neq 0$; in particular, $(x,y) \neq (1,1)$. Thus,

$$\lim_{(x,y)\to(1,1)} f(x,y) = \lim_{(x,y)\to(1,1)} \left(-\frac{x^2y^2}{x+y}\right) = -\frac{(1)^2(1)^2}{1+1} = -\frac{1}{2}.$$

12. $f(x,y) = \dfrac{\cos y - \sin 2y}{\cos x \cos y} = \dfrac{\cos y - 2\sin y \cos y}{\cos x \cos y} = \dfrac{1 - 2\sin y}{\cos x}$ for $\cos y \neq 0$; in particular, $(x,y) \neq (\pi, \pi/2)$. Thus,

$$\lim_{(x,y)\to(\pi,\pi/2)} f(x,y) = \lim_{(x,y)\to(\pi,\pi/2)} \frac{1 - 2\sin y}{\cos x} = \frac{1 - 2\sin(\pi/2)}{\cos \pi} = \frac{-1}{-1} = 1.$$

13. $f(x,y) = \dfrac{y^2}{x^2+y^2}$. First approach $(0,0)$ along the x-axis. Then $f(x,0) = 0/x^2 = 0$ for $x \neq 0$, so $f(x,y) \to 0$. Now approach $(0,0)$ along the y-axis. Then $f(0,y) = y^2/y^2 = 1$ for $y \neq 0$, so $f(x,y) \to 1$. Since f has two different limits along two different lines, the limit does not exist.

14. $f(x,y) = \dfrac{2xy}{x^2+3y^2}$. First approach $(0,0)$ along the x-axis. Then $f(x,0) = 0/x^2 = 0$ for $x \neq 0$, so $f(x,y) \to 0$. Now approach $(0,0)$ along the line $y = x$. Then $f(x,x) = 2x^2/4x^2 = 1/2$ for $x \neq 0$. Since f has two different limits along two different lines, the limit does not exist.

15. $f(x,y) = \dfrac{(x+y)^2}{x^2+y^2}$. First approach $(0,0)$ along the x-axis. Then $f(x,0) = x^2/x^2 = 1$ for $x \neq 0$, so $f(x,y) \to 1$. Now approach $(0,0)$ along the line $y = x$. Then $f(x,x) = 4x^2/(2x^2) = 2$ for $x \neq 0$, so $f(x,y) \to 2$. Since f has two different limits along two different lines, the limit does not exist.

16. $f(x,y) = \dfrac{x^2+xy^2}{x^4+y^2}$. First approach $(0,0)$ along the y-axis. Then $f(0,y) = 0/y^2 = 0$ for $y \neq 0$, so $f(x,y) \to 0$. Now approach $(0,0)$ along the line $y = x$. Then $f(x,x) = \dfrac{x^2+x^3}{x^4+x^2} = \dfrac{x^2(1+x)}{x^2(x^2+1)} = \dfrac{1+x}{1+x^2}$ for $x \neq 0$, so $f(x,y) \to 1$. Since f has two different limits along two different lines, the limit does not exist.

17. $f(x,y) = \dfrac{y^2 \sin^2 x}{x^4+y^4}$. First approach $(0,0)$ along the y-axis. Then $f(0,y) = 0/y^4 = 0$ for $y \neq 0$, so $f(x,y) \to 0$. Now approach $(0,0)$ along the line $y = x$. Then $f(x,x) = \dfrac{x^2 \sin^2 x}{2x^4} = \dfrac{1}{2}\left(\dfrac{\sin x}{x}\right)^2$ for $x \neq 0$, so by Equation 3.3.5, $f(x,y) \to \frac{1}{2}(1)^2 = \frac{1}{2}$. Since f has two different limits along two different lines, the limit does not exist.

18. $f(x,y) = \dfrac{y-x}{1-y+\ln x}$. First approach $(1,1)$ along the line $x = 1$. Then $f(1,y) = \dfrac{y-1}{1-y+0} = -1$ for $y \neq 1$, so $f(x,y) \to -1$. Now approach $(1,1)$ along the line $y = x$. Then $f(x,x) = \dfrac{x-x}{1-x+\ln x} = 0$ for $1 - x + \ln x \neq 0$. So $f(x,y) \to 0$. Since f has two different limits along two different lines, the limit does not exist.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $x^2y - xy^2 + 3$ is a polynomial and therefore continuous. t^3 is also a polynomial and therefore continuous. so the composition $(x^2y - xy^2 + 3)^3$ is continuous. Thus, $\lim\limits_{(x,y)\to(-1,-2)} (x^2y - xy^2 + 3)^3 = [(-1)^2(-2) - (-1)(-2)^2 + 3]^3 = 5^3 = 125.$

20. e^{xy} is a composition of continuous functions and therefore continuous. $\sin xy$ is also a composition of continuous functions and therefore continuous. Thus, the product $e^{xy}\sin xy$ is continuous, and $\lim\limits_{(x,y)\to(\pi,1/2)} e^{xy}\sin xy = e^{\pi/2}\sin\frac{\pi}{2} = e^{\pi/2}.$

21. $f(x,y) = \dfrac{3x-2y}{4x^2-y^2}$ is a rational function and continuous on its domain. $(2,3)$ is in the domain of f, so

$$\lim_{(x,y)\to(2,3)} f(x,y) = \frac{3(2)-2(3)}{4(2)^2-3^2} = \frac{0}{7} = 0.$$

22. $f(x,y) = \dfrac{2x-y}{4x^2-y^2} = \dfrac{2x-y}{(2x-y)(2x+y)} = \dfrac{1}{2x+y}$ for $2x-y \neq 0$. Thus, $\lim\limits_{(x,y)\to(1,2)} f(x,y) = \dfrac{1}{2(1)+2} = \dfrac{1}{4}.$

23. Let $f(x,y) = \dfrac{xy^2\cos y}{x^2+y^4}$. Then $f(x,0) = 0$ for $x \neq 0$, so $f(x,y) \to 0$ as $(x,y) \to (0,0)$ along the x-axis. Approaching $(0,0)$ along the y-axis or the line $y = x$ also gives a limit of 0. But $f\left(y^2, y\right) = \dfrac{y^2y^2\cos y}{(y^2)^2+y^4} = \dfrac{y^4\cos y}{2y^4} = \dfrac{\cos y}{2}$ for $y \neq 0$, so $f(x,y) \to \frac{1}{2}\cos 0 = \frac{1}{2}$ as $(x,y) \to (0,0)$ along the parabola $x = y^2$. Thus the limit doesn't exist.

24. $f(x,y) = \dfrac{x^3-y^3}{x^2+xy+y^2} = \dfrac{(x-y)(x^2+xy+y^2)}{x^2+xy+y^2} = x - y$ for $(x,y) \neq (0,0)$. [Note that $x^2+xy+y^2 = 0$ only when $(x,y) = (0,0)$.] Thus $\lim\limits_{(x,y)\to(0,0)} f(x,y) = \lim\limits_{(x,y)\to(0,0)} (x-y) = 0 - 0 = 0.$

25.
$$\begin{aligned}\lim_{(x,y)\to(0,0)} \frac{x^2+y^2}{\sqrt{x^2+y^2+1}-1} &= \lim_{(x,y)\to(0,0)} \frac{x^2+y^2}{\sqrt{x^2+y^2+1}-1}\cdot\frac{\sqrt{x^2+y^2+1}+1}{\sqrt{x^2+y^2+1}+1} \\ &= \lim_{(x,y)\to(0,0)} \frac{\left(x^2+y^2\right)\left(\sqrt{x^2+y^2+1}+1\right)}{x^2+y^2} = \lim_{(x,y)\to(0,0)} \left(\sqrt{x^2+y^2+1}+1\right) = 2\end{aligned}$$

26. $f(x,y) = \dfrac{xy^4}{x^2+y^8}$. On the x-axis, $f(x,0) = 0$ for $x \neq 0$, so $f(x,y) \to 0$ as $(x,y) \to (0,0)$ along the x-axis. Approaching $(0,0)$ along the curve $x = y^4$ gives $f(y^4,y) = y^8/2y^8 = \frac{1}{2}$ for $y \neq 0$, so along this path $f(x,y) \to \frac{1}{2}$ as $(x,y) \to (0,0)$. Thus the limit does not exist.

27. $x + z$ is a polynomial and therefore, continuous. $\sqrt{t}$ is continuous on its domain. Thus, the composition is continuous at $(6,1,-2)$. $\cos \pi y$ is continuous, so the product of $\sqrt{x+z}$ and $\cos \pi y$ is also continuous. Then

$$\lim_{(x,y,z)\to(6,1,-2)} \sqrt{x+z}\cos \pi y = \sqrt{6+(-2)}\cos \pi = -2.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $f(x,y,z) = \dfrac{xy+yz}{x^2+y^2+z^2}$. Then $f(x,0,0) = 0/x^2 = 0$ for $x \neq 0$, so as $(x,y,z) \to (0,0,0)$ along the x-axis, $f(x,y,z) \to 0$. But $f(x,x,0) = x^2/(2x^2) = \frac{1}{2}$ for $x \neq 0$, so as $(x,y,z) \to (0,0,0)$ along the line $y = x$, $z = 0$, $f(x,y,z) \to \frac{1}{2}$. Thus the limit doesn't exist.

29. $f(x,y,z) = \dfrac{xy+yz^2+xz^2}{x^2+y^2+z^4}$. Then $f(x,0,0) = 0/x^2 = 0$ for $x \neq 0$, so as $(x,y,z) \to (0,0,0)$ along the x-axis, $f(x,y,z) \to 0$. But $f(x,x,0) = x^2/(2x^2) = \frac{1}{2}$ for $x \neq 0$, so as $(x,y,z) \to (0,0,0)$ along the line $y = x$, $z = 0$, $f(x,y,z) \to \frac{1}{2}$. Thus the limit doesn't exist.

30. $f(x,y,z) = \dfrac{x^4+y^2+z^3}{x^4+2y^2+z}$. Then $f(x,0,0) = x^4/x^4 = 1$ for $x \neq 0$, so $f(x,y,z) \to 1$ as $(x,y,z) \to (0,0,0)$ along the x-axis. But $f(0,y,0) = y^2/(2y^2) = \frac{1}{2}$ for $y \neq 0$, so $f(x,y,z) \to \frac{1}{2}$ as $(x,y,z) \to (0,0,0)$ along the y-axis. Since f has two different limits along two different lines, the limit does not exist.

31. $-1 \leq \sin\left(\dfrac{1}{x^2+y^2}\right) \leq 1 \quad \Rightarrow \quad -xy \leq xy\sin\left(\dfrac{1}{x^2+y^2}\right) \leq xy$ for $xy > 0$. If $xy < 0$, we have $-xy \geq xy\sin\left(\dfrac{1}{x^2+y^2}\right) \geq xy$. In either case, $\displaystyle\lim_{(x,y)\to(0,0)} xy = 0$ and $\displaystyle\lim_{(x,y)\to(0,0)} (-xy) = 0$. Thus, $\displaystyle\lim_{(x,y)\to(0,0)} xy\sin\left(\dfrac{1}{x^2+y^2}\right) = 0$ by the Squeeze Theorem.

32. $f(x,y) = \dfrac{xy}{\sqrt{x^2+y^2}}$. We can see that the limit along any line through $(0,0)$ is 0, as well as along other paths through $(0,0)$ such as $x = y^2$ and $y = x^2$. So we suspect that the limit exists and equals 0; we use the Squeeze Theorem to prove our assertion. Since $|y| \leq \sqrt{x^2+y^2}$, we have $\dfrac{|y|}{\sqrt{x^2+y^2}} \leq 1$ and so $0 \leq \left|\dfrac{xy}{\sqrt{x^2+y^2}}\right| \leq |x|$. Now $|x| \to 0$ as $(x,y) \to (0,0)$, so $\left|\dfrac{xy}{\sqrt{x^2+y^2}}\right| \to 0$ and hence $\displaystyle\lim_{(x,y)\to(0,0)} f(x,y) = 0$.

33. We use the Squeeze Theorem to show that $\displaystyle\lim_{(x,y)\to(0,0)} \frac{xy^4}{x^4+y^4} = 0$:

$0 \leq \dfrac{|x|\,y^4}{x^4+y^4} \leq |x|$ since $0 \leq \dfrac{y^4}{x^4+y^4} \leq 1$, and $|x| \to 0$ as $(x,y) \to (0,0)$, so $\dfrac{|x|\,y^4}{x^4+y^4} \to 0 \quad \Rightarrow \quad \dfrac{xy^4}{x^4+y^4} \to 0$ as $(x,y) \to (0,0)$.

34. We use the Squeeze Theorem to show that $\displaystyle\lim_{(x,y,z)\to(0,0,0)} \frac{x^2y^2z^2}{x^2+y^2+z^2} = 0$:

$0 \leq \dfrac{x^2y^2z^2}{x^2+y^2+z^2} \leq x^2y^2$ since $0 \leq \dfrac{z^2}{x^2+y^2+z^2} \leq 1$, and $x^2y^2 \to 0$ as $(x,y,z) \to (0,0,0)$, so $\dfrac{x^2y^2z^2}{x^2+y^2+z^2} \to 0$ as $(x,y,z) \to (0,0,0)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35.

From the ridges on the graph, we see that as $(x, y) \to (0, 0)$ along the lines under the two ridges, $f(x, y) = \dfrac{2x^2 + 3xy + 4y^2}{3x^2 + 5y^2}$ approaches different values. Since the function approaches different values depending on the path of approach, the limit does not exist.

36.

From the graph, it appears that as we approach the origin along the lines $x = 0$ or $y = 0$, the function $f(x, y) = \dfrac{xy^3}{x^2 + y^6}$ is everywhere 0, whereas if we approach the origin along a certain curve it has a constant value of about $\frac{1}{2}$. [In fact, $f(y^3, y) = y^6/(2y^6) = \frac{1}{2}$ for $y \neq 0$, so $f(x, y) \to \frac{1}{2}$ as $(x, y) \to (0, 0)$ along the curve $x = y^3$.] Since the function approaches different values depending on the path of approach, the limit does not exist.

37. $h(x, y) = g(f(x, y)) = (2x + 3y - 6)^2 + \sqrt{2x + 3y - 6}$. Since f is a polynomial, it is continuous on $\mathbb{R}^2$ and g is continuous on its domain $\{t \mid t \geq 0\}$. Thus, h is continuous on its domain $\{(x, y) \mid 2x + 3y - 6 \geq 0\} = \left\{(x, y) \mid y \geq -\frac{2}{3}x + 2\right\}$, which consists of all points on or above the line $y = -\frac{2}{3}x + 2$.

38. $h(x, y) = g(f(x, y)) = \dfrac{1 - xy}{1 + x^2y^2} + \ln\left(\dfrac{1 - xy}{1 + x^2y^2}\right)$. f is a rational function, so it is continuous on its domain. Because $1 + x^2y^2 > 0$, the domain of f is $\mathbb{R}^2$, so f is continuous everywhere. g is continuous on its domain $\{t \mid t > 0\}$. Thus, h is continuous on its domain $\left\{(x, y) \;\middle|\; \dfrac{1 - xy}{1 + x^2y^2} > 0\right\} = \{(x, y) \mid xy < 1\}$, which consists of all points between (but not on) the two branches of the hyperbola $y = 1/x$.

39.

From the graph, it appears that f is discontinuous along the line $y = x$. If we consider $f(x, y) = e^{1/(x-y)}$ as a composition of functions, $g(x, y) = 1/(x - y)$ is a rational function and therefore continuous except where $x - y = 0 \quad \Leftrightarrow \quad y = x$. Since the function $h(t) = e^t$ is continuous everywhere, the composition $h(g(x, y)) = e^{1/(x-y)} = f(x, y)$ is continuous except along the line $y = x$, as we suspected.

40.

We can see a circular break in the graph, corresponding approximately to the unit circle, where f is discontinuous. Since $f(x, y) = \dfrac{1}{1 - x^2 - y^2}$ is a rational function, it is continuous except where $1 - x^2 - y^2 = 0 \quad \Leftrightarrow \quad x^2 + y^2 = 1$, confirming our observation that f is discontinuous on the circle $x^2 + y^2 = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. The functions xy and $1+e^{x-y}$ are continuous everywhere, and $1+e^{x-y}$ is never zero, so $F(x,y)=\dfrac{xy}{1+e^{x-y}}$ is continuous on its domain $\mathbb{R}^2$.

42. $F(x,y)=\cos\sqrt{1+x-y}=g(f(x,y))$ where $f(x,y)=\sqrt{1+x-y}$, continuous on its domain $\{(x,y)\mid 1+x-y\geq 0\}=\{(x,y)\mid y\leq x+1\}$, and $g(t)=\cos t$ is continuous everywhere. Thus F is continuous on its domain $\{(x,y)\mid y\leq x+1\}$.

43. $F(x,y)=\dfrac{1+x^2+y^2}{1-x^2-y^2}$ is a rational function and thus is continuous on its domain $\{(x,y)\mid 1-x^2-y^2\neq 0\}=\{(x,y)\mid x^2+y^2\neq 1\}$.

44. The functions e^x+e^y and $e^{xy}-1$ are continuous everywhere, so $H(x,y)=\dfrac{e^x+e^y}{e^{xy}-1}$ is continuous except where $e^{xy}-1=0 \quad\Rightarrow\quad xy=0 \quad\Rightarrow\quad x=0$ or $y=0$. Thus H is continuous on its domain $\{(x,y)\mid x\neq 0,\ y\neq 0\}$.

45. $\sqrt{x}$ is continuous on its domain $\{(x,y)\mid x\geq 0\}$ and $\sqrt{1-x^2-y^2}$ is continuous on its domain $\{(x,y)\mid 1-x^2-y^2\geq 0\}=\{(x,y)\mid x^2+y^2\leq 1\}$, so the sum $G(x,y)=\sqrt{x}+\sqrt{1-x^2-y^2}$ is continuous for $x\geq 0$ and $x^2+y^2\leq 1$, that is, $\{(x,y)\mid x^2+y^2\leq 1,\ x\geq 0\}$. This is the right half of the unit disk.

46. $G(x,y)=\ln(1+x-y)=g(f(x,y))$ where $f(x,y)=1+x-y$, a polynomial and hence continuous on $\mathbb{R}^2$, and $g(t)=\ln t$, continuous on its domain $\{t\mid t>0\}$. Thus G is continuous on its domain $\{(x,y)\mid 1+x-y>0\}=\{(x,y)\mid y<x+1\}$, the region in $\mathbb{R}^2$ below the line $y=x+1$.

47. $f(x,y,z)=h(g(x,y,z))$ where $g(x,y,z)=x^2+y^2+z^2$, a polynomial that is continuous everywhere, and $h(t)=\arcsin t$, continuous on $[-1,1]$. Thus f is continuous on its domain $\{(x,y,z)\mid -1\leq x^2+y^2+z^2\leq 1\}=\{(x,y,z)\mid x^2+y^2+z^2\leq 1\}$, so f is continuous on the unit ball.

48. $\sqrt{y-x^2}$ is continuous on its domain $\{(x,y)\mid y-x^2\geq 0\}=\{(x,y)\mid y\geq x^2\}$ and $\ln z$ is continuous on its domain $\{z\mid z>0\}$, so the product $f(x,y,z)=\sqrt{y-x^2}\ln z$ is continuous for $y\geq x^2$ and $z>0$, that is, $\{(x,y,z)\mid y\geq x^2,\ z>0\}$.

49. $f(x,y)=\begin{cases}\dfrac{x^2y^3}{2x^2+y^2} & \text{if } (x,y)\neq(0,0)\\ 1 & \text{if } (x,y)=(0,0)\end{cases}$ The first piece of f is a rational function defined everywhere except at the origin, so f is continuous on $\mathbb{R}^2$ except possibly at the origin. Since $x^2\leq 2x^2+y^2$, we have $\left|x^2y^3/(2x^2+y^2)\right|\leq\left|y^3\right|$. We know that $\left|y^3\right|\to 0$ as $(x,y)\to(0,0)$. So, by the Squeeze Theorem, $\displaystyle\lim_{(x,y)\to(0,0)} f(x,y)=\lim_{(x,y)\to(0,0)}\frac{x^2y^3}{2x^2+y^2}=0$. But $f(0,0)=1$, so f is discontinuous at $(0,0)$. Therefore, f is continuous on the set $\{(x,y)\mid (x,y)\neq(0,0)\}$.

50. $f(x,y)=\begin{cases}\dfrac{xy}{x^2+xy+y^2} & \text{if } (x,y)\neq(0,0)\\ 0 & \text{if } (x,y)=(0,0)\end{cases}$ The first piece of f is a rational function defined everywhere except at the origin, so f is continuous on $\mathbb{R}^2$ except possibly at the origin. $f(x,0)=0/x^2=0$ for $x\neq 0$, so $f(x,y)\to 0$ as

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$(x, y) \to (0, 0)$ along the x-axis. But $f(x, x) = x^2/(3x^2) = \frac{1}{3}$ for $x \neq 0$, so $f(x, y) \to \frac{1}{3}$ as $(x, y) \to (0, 0)$ along the line $y = x$. Thus $\lim\limits_{(x,y)\to(0,0)} f(x, y)$ doesn't exist, so f is not continuous at $(0, 0)$ and the largest set on which f is continuous is $\{(x, y) \mid (x, y) \neq (0, 0)\}$.

51. $$\lim_{(x,y)\to(0,0)} \frac{x^3 + y^3}{x^2 + y^2} = \lim_{r\to0^+} \frac{(r\cos\theta)^3 + (r\sin\theta)^3}{r^2} = \lim_{r\to0^+} (r\cos^3\theta + r\sin^3\theta) = 0$$

52. $$\lim_{(x,y)\to(0,0)} (x^2 + y^2)\ln(x^2 + y^2) = \lim_{r\to0^+} r^2 \ln r^2 = \lim_{r\to0^+} \frac{\ln r^2}{1/r^2} = \lim_{r\to0^+} \frac{(1/r^2)(2r)}{-2/r^3} \quad \text{[using l'Hospital's Rule]}$$
$$= \lim_{r\to0^+} (-r^2) = 0$$

53. $$\lim_{(x,y)\to(0,0)} \frac{e^{-x^2-y^2} - 1}{x^2 + y^2} = \lim_{r\to0^+} \frac{e^{-r^2} - 1}{r^2} = \lim_{r\to0^+} \frac{e^{-r^2}(-2r)}{2r} \quad \text{[using l'Hospital's Rule]}$$
$$= \lim_{r\to0^+} -e^{-r^2} = -e^0 = -1$$

54. **1.** $\lim\limits_{(x,y)\to(a,b)} x = a$: Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < \sqrt{(x-a)^2 + (y-b)^2} < \delta$, then $|x - a| < \varepsilon$.

But $|x - a| \leq \sqrt{(x-a)^2 + (y-b)^2} < \delta$, so choose $\varepsilon = \delta$. Then $|x - a| \leq \sqrt{(x-a)^2 + (y-b)^2} < \delta = \varepsilon$.

Thus, $\lim\limits_{(x,y)\to(a,b)} x = a$.

2. $\lim\limits_{(x,y)\to(a,b)} y = b$: Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < \sqrt{(x-a)^2 + (y-b)^2} < \delta$, then $|y - b| < \varepsilon$.

But $|y - b| \leq \sqrt{(x-a)^2 + (y-b)^2} < \delta$, so choose $\varepsilon = \delta$. Then $|y - b| \leq \sqrt{(x-a)^2 + (y-b)^2} < \delta = \varepsilon$.

Thus, $\lim\limits_{(x,y)\to(a,b)} y = b$.

3. $\lim\limits_{(x,y)\to(a,b)} c = c$: Given $\varepsilon > 0$, we need $\delta > 0$ such that if $0 < \sqrt{(x-a)^2 + (y-b)^2} < \delta$, then $|c - c| < \varepsilon$.

But $|c - c| = 0$, so this will be true no matter what δ we choose.

55. $\lim\limits_{(x,y)\to(0,0)} \dfrac{\sin(x^2 + y^2)}{x^2 + y^2} = \lim\limits_{r\to0^+} \dfrac{\sin(r^2)}{r^2}$, which is an indeterminate form of type $0/0$. Using l'Hospital's Rule, we get

$$\lim_{r\to0^+} \frac{\sin(r^2)}{r^2} \overset{\text{H}}{=} \lim_{r\to0^+} \frac{2r\cos(r^2)}{2r} = \lim_{r\to0^+} \cos(r^2) = 1.$$

Or: Use the fact that $\lim\limits_{\theta\to0} \dfrac{\sin\theta}{\theta} = 1$.

56. $$f(x, y) = \begin{cases} \dfrac{\sin xy}{xy} & \text{if } (x, y) \neq (0, 0) \\ 1 & \text{if } (x, y) = (0, 0) \end{cases}$$

From the graph, it appears that f is continuous everywhere. We know xy is continuous on $\mathbb{R}^2$ and $\sin t$ is continuous everywhere, so $\sin xy$ is continuous on $\mathbb{R}^2$ and $\dfrac{\sin xy}{xy}$ is continuous on $\mathbb{R}^2$ except

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

possibly where $xy = 0$. To show that f is continuous at those points, consider any point (a, b) in $\mathbb{R}^2$ where $ab = 0$. Because xy is continuous, $xy \to ab = 0$ as $(x, y) \to (a, b)$. If we let $t = xy$, then $t \to 0$ as $(x, y) \to (a, b)$ and $\lim\limits_{(x,y)\to(a,b)} \dfrac{\sin xy}{xy} = \lim\limits_{t\to 0} \dfrac{\sin t}{t} = 1$ by Equation 3.3.5. Thus $\lim\limits_{(x,y)\to(a,b)} f(x, y) = f(a, b)$ and f is continuous on $\mathbb{R}^2$.

57. (a) $f(x, y) = \begin{cases} 0 & \text{if } y \le 0 \quad \text{or} \quad y \ge x^4 \\ 1 & \text{if } 0 < y < x^4 \end{cases}$ Consider the path $y = mx^a$, $0 < a < 4$. [The path does not pass through $(0, 0)$ if $a \le 0$ except for the trivial case where $m = 0$.] If $mx^a \le 0$ then $f(x, mx^a) = 0$. If $mx^a > 0$ then $mx^a = |mx^a| = |m|\,|x^a|$ and $mx^a \ge x^4 \quad \Leftrightarrow \quad |m|\,|x^a| \ge x^4 \quad \Leftrightarrow \quad \dfrac{x^4}{|x^a|} \le |m| \quad \Leftrightarrow \quad |x|^{4-a} \le |m|$ whenever x^a is defined. Then $mx^a \ge x^4 \quad \Leftrightarrow \quad |x| \le |m|^{1/(4-a)}$ so $f(x, mx^a) = 0$ for $|x| \le |m|^{1/(4-a)}$ and $f(x, y) \to 0$ as $(x, y) \to (0, 0)$ along this path.

(b) If we approach $(0, 0)$ along the path $y = x^5$, $x > 0$ then we have $f(x, x^5) = 1$ for $0 < x < 1$ because $0 < x^5 < x^4$ there. Thus $f(x, y) \to 1$ as $(x, y) \to (0, 0)$ along this path, but in part (a) we found a limit of 0 along other paths, so $\lim\limits_{(x,y)\to(0,0)} f(x, y)$ doesn't exist and f is discontinuous at $(0, 0)$.

(c) First we show that f is discontinuous at any point $(a, 0)$ on the x-axis. If we approach $(a, 0)$ along the path $x = a$, $y > 0$ then $f(a, y) = 1$ for $0 < y < a^4$, so $f(x, y) \to 1$ as $(x, y) \to (a, 0)$ along this path. If we approach $(a, 0)$ along the path $x = a$, $y < 0$ then $f(a, y) = 0$ since $y < 0$ and $f(x, y) \to 0$ as $(x, y) \to (a, 0)$. Thus the limit does not exist and f is discontinuous on the line $y = 0$. f is also discontinuous on the curve $y = x^4$: For any point (a, a^4) on this curve, approaching the point along the path $x = a$, $y > a^4$ gives $f(a, y) = 0$ since $y > a^4$, so $f(x, y) \to 0$ as $(x, y) \to (a, a^4)$. But approaching the point along the path $x = a$, $y < a^4$ gives $f(a, y) = 1$ for $y > 0$, so $f(x, y) \to 1$ as $(x, y) \to (a, a^4)$ and the limit does not exist there.

58. Since $|\mathbf{x} - \mathbf{a}|^2 = |\mathbf{x}|^2 + |\mathbf{a}|^2 - 2\,|\mathbf{x}|\,|\mathbf{a}| \cos\theta \ge |\mathbf{x}|^2 + |\mathbf{a}|^2 - 2\,|\mathbf{x}|\,|\mathbf{a}| = (|\mathbf{x}| - |\mathbf{a}|)^2$, we have $\big||\mathbf{x}| - |\mathbf{a}|\big| \le |\mathbf{x} - \mathbf{a}|$. Let $\varepsilon > 0$ be given and set $\delta = \varepsilon$. Then if $0 < |\mathbf{x} - \mathbf{a}| < \delta$, $\big||\mathbf{x}| - |\mathbf{a}|\big| \le |\mathbf{x} - \mathbf{a}| < \delta = \varepsilon$. Hence $\lim_{\mathbf{x}\to\mathbf{a}} |\mathbf{x}| = |\mathbf{a}|$ and $f(\mathbf{x}) = |\mathbf{x}|$ is continuous on $\mathbb{R}^n$.

59. $f(\mathbf{x}) = \mathbf{c} \cdot \mathbf{x}$. Let $\varepsilon > 0$ be given. We need to find $\delta > 0$ such that if $0 < |\mathbf{x} - \mathbf{a}| < \delta$, then $|f(\mathbf{x}) - f(\mathbf{a})| = |\mathbf{c} \cdot \mathbf{x} - \mathbf{c} \cdot \mathbf{a}| < \varepsilon$. But $|\mathbf{c} \cdot \mathbf{x} - \mathbf{c} \cdot \mathbf{a}| = |\mathbf{c} \cdot (\mathbf{x} - \mathbf{a})|$ and $|\mathbf{c} \cdot (\mathbf{x} - \mathbf{a})| \le |\mathbf{c}|\,|\mathbf{x} - \mathbf{a}|$ by Exercise 12.3.61 (the Cauchy-Schwartz Inequality). Set $\delta = \varepsilon / |\mathbf{c}|$. Then if $0 < |\mathbf{x} - \mathbf{a}| < \delta$, $|f(\mathbf{x}) - f(\mathbf{a})| = |\mathbf{c} \cdot \mathbf{x} - \mathbf{c} \cdot \mathbf{a}| \le |\mathbf{c}|\,|\mathbf{x} - \mathbf{a}| < |\mathbf{c}|\,\delta = |\mathbf{c}|\,(\varepsilon / |\mathbf{c}|) = \varepsilon$. So f is continuous on $\mathbb{R}^n$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14.3 Partial Derivatives

1. By Definition 4, $f_T(92,60) = \lim_{h\to 0} \dfrac{f(92+h,60)-f(92,60)}{h}$, which we can approximate by considering $h=2$ and $h=-2$ and using the values given in Table 1: $f_T(92,60) \approx \dfrac{f(94,60)-f(92,60)}{2} = \dfrac{111-105}{2} = 3$, $f_T(92,60) \approx \dfrac{f(90,60)-f(92,60)}{-2} = \dfrac{100-105}{-2} = 2.5$. Averaging these values, we estimate $f_T\,(92,60)$ to be approximately 2.75. Thus, when the actual temperature is $92°\mathrm{F}$ and the relative humidity is 60%, the apparent temperature rises by about $2.75°\mathrm{F}$ for every degree that the actual temperature rises.

Similarly, $f_H(92,60) = \lim_{h\to 0} \dfrac{f(92,60+h)-f(92,60)}{h}$ which we can approximate by considering $h=5$ and $h=-5$: $f_H(92,60) \approx \dfrac{f(92,65)-f(92,60)}{5} = \dfrac{108-105}{5} = 0.6$, $f_H(92,60) \approx \dfrac{f(92,55)-f(92,60)}{-5} = \dfrac{103-105}{-5} = 0.4$. Averaging these values, we estimate $f_H(92,60)$ to be approximately 0.5. Thus, when the actual temperature is $92°\mathrm{F}$ and the relative humidity is 60%, the apparent temperature rises by about $0.5°\mathrm{F}$ for every percent that the relative humidity increases.

2. (a) $\partial h/\partial v$ represents the rate of change of h when we fix t and consider h as a function of v, which describes how quickly the wave heights change when the wind speed changes for a fixed time duration. $\partial h/\partial t$ represents the rate of change of h when we fix v and consider h as a function of t, which describes how quickly the wave heights change when the duration of time changes, but the wind speed is constant.

(b) By Definition 4, $f_v(40,15) = \lim_{h\to 0} \dfrac{f(40+h,15)-f(40,15)}{h}$ which we can approximate by considering $h=10$ and $h=-10$ and using the values given in the table: $f_v(40,15) \approx \dfrac{f(50,15)-f(40,15)}{10} = \dfrac{36-25}{10} = 1.1$, $f_v(40,15) \approx \dfrac{f(30,15)-f(40,15)}{-10} = \dfrac{16-25}{-10} = 0.9$. Averaging these values, we have $f_v(40,15) \approx 1.0$. Thus, when a 40-knot wind has been blowing for 15 hours, the wave heights should increase by about 1 foot for every knot that the wind speed increases (with the same time duration). Similarly, $f_t(40,15) = \lim_{h\to 0} \dfrac{f(40,15+h)-f(40,15)}{h}$ which we can approximate by considering $h=5$ and $h=-5$: $f_t(40,15) \approx \dfrac{f(40,20)-f(40,15)}{5} = \dfrac{28-25}{5} = 0.6$, $f_t(40,15) \approx \dfrac{f(40,10)-f(40,15)}{-5} = \dfrac{21-25}{-5} = 0.8$. Averaging these values, we have $f_t(40,15) \approx 0.7$. Thus, when a 40-knot wind has been blowing for 15 hours, the wave heights increase by about 0.7 feet for every additional hour that the wind blows.

(c) For fixed values of v, the function values $f(v,t)$ appear to increase in smaller and smaller increments, becoming nearly constant as t increases. Thus, the corresponding rate of change is nearly 0 as t increases, suggesting that $\lim_{t\to\infty} (\partial h/\partial t) = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a) $\partial T/\partial x$ represents the rate of change of T when we fix y and t and consider T as a function of the single variable x, which describes how quickly the temperature changes when longitude changes but latitude and time are constant. $\partial T/\partial y$ represents the rate of change of T when we fix x and t and consider T as a function of y, which describes how quickly the temperature changes when latitude changes but longitude and time are constant. $\partial T/\partial t$ represents the rate of change of T when we fix x and y and consider T as a function of t, which describes how quickly the temperature changes over time for a constant longitude and latitude.

(b) $f_x(158, 21, 9)$ represents the rate of change of temperature at longitude $158°$W, latitude $21°$N at 9:00 AM when only longitude varies. Since the air is warmer to the west than to the east, increasing longitude results in an increased air temperature, so we would expect $f_x(158, 21, 9)$ to be positive. $f_y(158, 21, 9)$ represents the rate of change of temperature at the same time and location when only latitude varies. Since the air is warmer to the south and cooler to the north, increasing latitude results in a decreased air temperature, so we would expect $f_y(158, 21, 9)$ to be negative. $f_t(158, 21, 9)$ represents the rate of change of temperature at the same time and location when only time varies. Since typically air temperature increases from the morning to the afternoon as the sun warms it, we would expect $f_t(158, 21, 9)$ to be positive.

4. (a) If we start at $(1, 2)$ and move in the positive x-direction, the graph of f increases. Thus $f_x(1, 2)$ is positive.

(b) If we start at $(1, 2)$ and move in the positive y-direction, the graph of f decreases. Thus $f_y(1, 2)$ is negative.

5. (a) The graph of f decreases if we start at $(-1, 2)$ and move in the positive x-direction, so $f_x(-1, 2)$ is negative.

(b) The graph of f decreases if we start at $(-1, 2)$ and move in the positive y-direction, so $f_y(-1, 2)$ is negative.

6. $f_x(2, 1)$ is the rate of change of f at $(2, 1)$ in the x-direction. If we start at $(2, 1)$, where $f(2, 1) = 10$, and move in the positive x-direction, we reach the next contour line [where $f(x, y) = 12$] after approximately 0.6 units. This represents an average rate of change of about $\frac{2}{0.6}$. If we approach the point $(2, 1)$ from the left (moving in the positive x-direction) the output values increase from 8 to 10 with an increase in x of approximately 0.9 units, corresponding to an average rate of change of $\frac{2}{0.9}$. A good estimate for $f_x(2, 1)$ would be the average of these two, so $f_x(2, 1) \approx 2.8$. Similarly, $f_y(2, 1)$ is the rate of change of f at $(2, 1)$ in the y-direction. If we approach $(2, 1)$ from below, the output values decrease from 12 to 10 with a change in y of approximately 1 unit, corresponding to an average rate of change of -2. If we start at $(2, 1)$ and move in the positive y-direction, the output values decrease from 10 to 8 after approximately 0.9 units, a rate of change of $\frac{-2}{0.9}$. Averaging these two results, we estimate $f_y(2, 1) \approx -2.1$.

7. $f(x, y) = 16 - 4x^2 - y^2 \quad \Rightarrow \quad f_x(x, y) = -8x$ and $f_y(x, y) = -2y \quad \Rightarrow \quad f_x(1, 2) = -8$ and $f_y(1, 2) = -4$. The graph of f is the paraboloid $z = 16 - 4x^2 - y^2$ and the vertical plane $y = 2$ intersects it in the parabola $z = 12 - 4x^2$, $y = 2$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(the curve C_1 in the first figure). The slope of the tangent line to this parabola at $(1, 2, 8)$ is $f_x(1, 2) = -8$. Similarly the plane $x = 1$ intersects the paraboloid in the parabola $z = 12 - y^2$, $x = 1$ (the curve C_2 in the second figure) and the slope of the tangent line at $(1, 2, 8)$ is $f_y(1, 2) = -4$.

8. $f(x, y) = (4 - x^2 - 4y^2)^{1/2} \quad \Rightarrow \quad f_x(x, y) = -x(4 - x^2 - 4y^2)^{-1/2}$ and $f_y(x, y) = -4y(4 - x^2 - 4y^2)^{-1/2} \quad \Rightarrow$ $f_x(1, 0) = -\frac{1}{\sqrt{3}}$, $f_y(1, 0) = 0$. The graph of f is the upper half of the ellipsoid $z^2 + x^2 + 4y^2 = 4$ and the plane $y = 0$ intersects the graph in the semicircle $x^2 + z^2 = 4$, $z \geq 0$ and the slope of the tangent line T_1 to this semicircle at $\left(1, 0, \sqrt{3}\right)$ is $f_x(1, 0) = -\frac{1}{\sqrt{3}}$. Similarly the plane $x = 1$ intersects the graph in the semi-ellipse $z^2 + 4y^2 = 3$, $z \geq 0$ and the slope of the tangent line T_2 to this semi-ellipse at $\left(1, 0, \sqrt{3}\right)$ is $f_y(1, 0) = 0$.

9. $f(x, y) = x^4 + 5xy^3 \quad \Rightarrow \quad f_x(x, y) = 4x^3 + 5y^3,\ f_y(x, y) = 0 + 5x \cdot 3y^2 = 15xy^2$

10. $f(x, y) = x^2y - 3y^4 \quad \Rightarrow \quad f_x(x, y) = 2x \cdot y - 0 = 2xy,\ f_y(x, y) = x^2 \cdot 1 - 3 \cdot 4y^3 = x^2 - 12y^3$

11. $g(x, y) = x^3 \sin y \quad \Rightarrow \quad g_x(x, y) = 3x^2 \sin y,\ g_y(x, y) = x^3 \cos y$

12. $g(x, t) = e^{xt} \quad \Rightarrow \quad g_x(x, t) = e^{xt} \cdot t = te^{xt},\ g_t(x, t) = e^{xt} \cdot x = xe^{xt}$

13. $z = \ln(x + t^2) \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = \dfrac{1}{x + t^2} \cdot 1 = \dfrac{1}{x + t^2},\ \dfrac{\partial z}{\partial t} = \dfrac{1}{x + t^2} \cdot 2t = \dfrac{2t}{x + t^2}$

14. $w = \dfrac{u}{v^2} \quad \Rightarrow \quad w_u = \dfrac{1}{v^2} \cdot 1 = \dfrac{1}{v^2},\ w_v = u \cdot (-2v^{-3}) = -\dfrac{2u}{v^3}$

15. $f(x, y) = ye^{xy} \quad \Rightarrow \quad f_x(x, y) = y \cdot e^{xy} \cdot y = y^2e^{xy},\ f_y(x, y) = y \cdot e^{xy} \cdot x + e^{xy} \cdot 1 = e^{xy} + xye^{xy}$

16. $g(x, y) = (x^2 + xy)^3 \quad \Rightarrow \quad g_x(x, y) = 3(x^2 + xy)^2(2x + y),\ g_y(x, y) = 3(x^2 + xy)^2(0 + x) = 3x(x^2 + xy)^2$

17. $g(x, y) = y(x + x^2y)^5 \quad \Rightarrow \quad g_x(x, y) = 5y(x + x^2y)^4(1 + 2xy)$,

$g_y(x, y) = y \cdot 5(x + x^2y)^4 \cdot x^2 + (x + x^2y)^5 \cdot 1 = 5x^2y(x + x^2y)^4 + (x + x^2y)^5$

18. $f(x, y) = \dfrac{x}{(x + y)^2} \quad \Rightarrow \quad f_x(x, y) = \dfrac{(x + y)^2(1) - (x)(2)(x + y)}{[(x + y)^2]^2} = \dfrac{x + y - 2x}{(x + y)^3} = \dfrac{y - x}{(x + y)^3}$,

$f_y(x, y) = \dfrac{(x + y)^2(0) - (x)(2)(x + y)}{[(x + y)^2]^2} = -\dfrac{2x}{(x + y)^3}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $f(x,y) = \dfrac{ax+by}{cx+dy} \quad \Rightarrow \quad f_x(x,y) = \dfrac{(cx+dy)(a)-(ax+by)(c)}{(cx+dy)^2} = \dfrac{(ad-bc)y}{(cx+dy)^2}$,

$f_y(x,y) = \dfrac{(cx+dy)(b)-(ax+by)(d)}{(cx+dy)^2} = \dfrac{(bc-ad)x}{(cx+dy)^2}$

20. $w = \dfrac{e^v}{u+v^2} \quad \Rightarrow \quad \dfrac{\partial w}{\partial u} = \dfrac{0(u+v^2)-e^v(1)}{(u+v^2)^2} = -\dfrac{e^v}{(u+v^2)^2}$, $\dfrac{\partial w}{\partial v} = \dfrac{e^v(u+v^2)-e^v(2v)}{(u+v^2)^2} = \dfrac{e^v(u+v^2-2v)}{(u+v^2)^2}$

21. $g(u,v) = (u^2v-v^3)^5 \quad \Rightarrow \quad g_u(u,v) = 5(u^2v-v^3)^4 \cdot 2uv = 10uv(u^2v-v^3)^4$,

$g_v(u,v) = 5(u^2v-v^3)^4(u^2-3v^2) = 5(u^2-3v^2)(u^2v-v^3)^4$

22. $u(r,\theta) = \sin(r\cos\theta) \quad \Rightarrow \quad u_r(r,\theta) = \cos(r\cos\theta)\cdot\cos\theta = \cos\theta\cos(r\cos\theta)$,

$u_\theta(r,\theta) = \cos(r\cos\theta)(-r\sin\theta) = -r\sin\theta\cos(r\cos\theta)$

23. $R(p,q) = \tan^{-1}(pq^2) \quad \Rightarrow \quad R_p(p,q) = \dfrac{1}{1+(pq^2)^2}\cdot q^2 = \dfrac{q^2}{1+p^2q^4}$, $R_q(p,q) = \dfrac{1}{1+(pq^2)^2}\cdot 2pq = \dfrac{2pq}{1+p^2q^4}$

24. $f(x,y) = x^y \quad \Rightarrow \quad f_x(x,y) = yx^{y-1}$, $f_y(x,y) = x^y\ln x$

25. $F(x,y) = \displaystyle\int_y^x \cos(e^t)\,dt \quad \Rightarrow \quad F_x(x,y) = \frac{\partial}{\partial x}\int_y^x \cos\left(e^t\right)\,dt = \cos(e^x)$ by the Fundamental Theorem of Calculus, Part 1;

$F_y(x,y) = \dfrac{\partial}{\partial y}\displaystyle\int_y^x \cos\left(e^t\right)\,dt = \frac{\partial}{\partial y}\left[-\int_x^y \cos\left(e^t\right)\,dt\right] = -\frac{\partial}{\partial y}\int_x^y \cos\left(e^t\right)\,dt = -\cos(e^y)$.

26. $F(\alpha,\beta) = \displaystyle\int_\alpha^\beta \sqrt{t^3+1}\,dt \quad \Rightarrow$

$F_\alpha(\alpha,\beta) = \dfrac{\partial}{\partial\alpha}\displaystyle\int_\alpha^\beta \sqrt{t^3+1}\,dt = \frac{\partial}{\partial\alpha}\left[-\int_\beta^\alpha \sqrt{t^3+1}\,dt\right] = -\frac{\partial}{\partial\alpha}\int_\beta^\alpha \sqrt{t^3+1}\,dt = -\sqrt{\alpha^3+1}$ by the Fundamental Theorem of Calculus, Part 1; $F_\beta(\alpha,\beta) = \dfrac{\partial}{\partial\beta}\displaystyle\int_\alpha^\beta \sqrt{t^3+1}\,dt = \sqrt{\beta^3+1}$.

27. $f(x,y,z) = x^3yz^2+2yz \quad \Rightarrow \quad f_x(x,y,z) = 3x^2yz^2$, $f_y(x,y,z) = x^3z^2+2z$, $f_z(x,y,z) = 2x^3yz+2y$

28. $f(x,y,z) = xy^2e^{-xz} \quad \Rightarrow \quad f_x(x,y,z) = y^2\left[x\cdot e^{-xz}(-z)+e^{-xz}\cdot 1\right] = (1-xz)y^2e^{-xz}$, $f_y(x,y,z) = 2xye^{-xz}$,

$f_z(x,y,z) = xy^2e^{-xz}(-x) = -x^2y^2e^{-xz}$

29. $w = \ln(x+2y+3z) \quad \Rightarrow \quad \dfrac{\partial w}{\partial x} = \dfrac{1}{x+2y+3z}$, $\dfrac{\partial w}{\partial y} = \dfrac{2}{x+2y+3z}$, $\dfrac{\partial w}{\partial z} = \dfrac{3}{x+2y+3z}$

30. $w = y\tan(x+2z) \quad \Rightarrow \quad \dfrac{\partial w}{\partial x} = y\left[\sec^2(x+2z)\right](1) = y\sec^2(x+2z)$, $\dfrac{\partial w}{\partial y} = \tan(x+2z)$,

$\dfrac{\partial w}{\partial z} = y\left[\sec^2(x+2z)\right](2) = 2y\sec^2(x+2z)$

31. $p = \sqrt{t^4+u^2\cos v} \quad \Rightarrow \quad \dfrac{\partial p}{\partial t} = \frac{1}{2}(t^4+u^2\cos v)^{-1/2}(4t^3) = \dfrac{2t^3}{\sqrt{t^4+u^2\cos v}}$,

$\dfrac{\partial p}{\partial u} = \frac{1}{2}(t^4+u^2\cos v)^{-1/2}(2u\cos v) = \dfrac{u\cos v}{\sqrt{t^4+u^2\cos v}}$, $\dfrac{\partial p}{\partial v} = \frac{1}{2}(t^4+u^2\cos v)^{-1/2}[u^2(-\sin v)] = -\dfrac{u^2\sin v}{2\sqrt{t^4+u^2\cos v}}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. $u = x^{y/z} \quad \Rightarrow \quad u_x = \dfrac{y}{z}\, x^{(y/z)-1},\ u_y = x^{y/z} \ln x \cdot \dfrac{1}{z} = \dfrac{x^{y/z}}{z} \ln x,\ \ u_z = x^{y/z} \ln x \cdot \dfrac{-y}{z^2} = -\dfrac{yx^{y/z}}{z^2} \ln x$

33. $h(x,y,z,t) = x^2 y \cos(z/t) \quad \Rightarrow \quad h_x(x,y,z,t) = 2xy\cos(z/t),\ h_y(x,y,z,t) = x^2\cos(z/t),$

$h_z(x,y,z,t) = -x^2 y \sin(z/t)(1/t) = (-x^2 y/t)\sin(z/t),\ h_t(x,y,z,t) = -x^2 y \sin(z/t)(-zt^{-2}) = (x^2 yz/t^2)\sin(z/t)$

34. $\phi(x,y,z,t) = \dfrac{\alpha x + \beta y^2}{\gamma z + \delta t^2} \quad \Rightarrow \quad \phi_x(x,y,z,t) = \dfrac{1}{\gamma z + \delta t^2}(\alpha) = \dfrac{\alpha}{\gamma z + \delta t^2},$

$\phi_y(x,y,z,t) = \dfrac{1}{\gamma z + \delta t^2}(2\beta y) = \dfrac{2\beta y}{\gamma z + \delta t^2},\ \phi_z(x,y,z,t) = \dfrac{(\gamma z + \delta t^2)(0) - (\alpha x + \beta y^2)(\gamma)}{(\gamma z + \delta t^2)^2} = \dfrac{-\gamma(\alpha x + \beta y^2)}{(\gamma z + \delta t^2)^2},$

$\phi_t(x,y,z,t) = \dfrac{(\gamma z + \delta t^2)(0) - (\alpha x + \beta y^2)(2\delta t)}{(\gamma z + \delta t^2)^2} = -\dfrac{2\delta t(\alpha x + \beta y^2)}{(\gamma z + \delta t^2)^2}$

35. $u = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2}$. For each $i = 1, \ldots, n$, $u_{x_i} = \frac{1}{2}\left(x_1^2 + x_2^2 + \cdots + x_n^2\right)^{-1/2}(2x_i) = \dfrac{x_i}{\sqrt{x_1^2 + x_2^2 + \cdots + x_n^2}}$.

36. $u = \sin(x_1 + 2x_2 + \cdots + nx_n)$. For each $i = 1, \ldots, n$, $u_{x_i} = i\cos(x_1 + 2x_2 + \cdots + nx_n)$.

37. $R(s,t) = te^{s/t} \quad \Rightarrow \quad R_t(s,t) = t \cdot e^{s/t}(-s/t^2) + e^{s/t} \cdot 1 = \left(1 - \dfrac{s}{t}\right) e^{s/t}$, so $R_t(0,1) = \left(1 - \dfrac{0}{1}\right) e^{0/1} = 1$.

38. $f(x,y) = y\sin^{-1}(xy) \quad \Rightarrow \quad f_y(x,y) = y \cdot \dfrac{1}{\sqrt{1-(xy)^2}}(x) + \sin^{-1}(xy) \cdot 1 = \dfrac{xy}{\sqrt{1-x^2y^2}} + \sin^{-1}(xy),$

so $f_y\left(1, \frac{1}{2}\right) = \dfrac{1 \cdot \frac{1}{2}}{\sqrt{1 - 1^2\left(\frac{1}{2}\right)^2}} + \sin^{-1}\left(1 \cdot \frac{1}{2}\right) = \dfrac{\frac{1}{2}}{\sqrt{\frac{3}{4}}} + \sin^{-1}\frac{1}{2} = \frac{1}{\sqrt{3}} + \frac{\pi}{6}$.

39. $f(x,y,z) = \ln \dfrac{1 - \sqrt{x^2+y^2+z^2}}{1 + \sqrt{x^2+y^2+z^2}} \quad \Rightarrow$

$$f_y(x,y,z) = \frac{1}{\dfrac{1 - \sqrt{x^2+y^2+z^2}}{1 + \sqrt{x^2+y^2+z^2}}} \cdot \frac{\left(1 + \sqrt{x^2+y^2+z^2}\right)\left(-\frac{1}{2}(x^2+y^2+z^2)^{-1/2} \cdot 2y\right) - \left(1 - \sqrt{x^2+y^2+z^2}\right)\left(\frac{1}{2}(x^2+y^2+z^2)^{-1/2} \cdot 2y\right)}{\left(1 + \sqrt{x^2+y^2+z^2}\right)^2}$$

$$= \frac{1 + \sqrt{x^2+y^2+z^2}}{1 - \sqrt{x^2+y^2+z^2}} \cdot \frac{-y(x^2+y^2+z^2)^{-1/2}\left(1 + \sqrt{x^2+y^2+z^2} + 1 - \sqrt{x^2+y^2+z^2}\right)}{\left(1 + \sqrt{x^2+y^2+z^2}\right)^2}$$

$$= \frac{-y(x^2+y^2+z^2)^{-1/2}\,(2)}{\left(1 - \sqrt{x^2+y^2+z^2}\right)\left(1 + \sqrt{x^2+y^2+z^2}\right)} = \frac{-2y}{\sqrt{x^2+y^2+z^2}\,[1 - (x^2+y^2+z^2)]}$$

so $f_y(1,2,2) = \dfrac{-2(2)}{\sqrt{1^2+2^2+2^2}\,[1 - (1^2+2^2+2^2)]} = \dfrac{-4}{\sqrt{9}\,(1-9)} = \dfrac{1}{6}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $f(x,y,z) = x^{yz} \quad \Rightarrow \quad f_z(x,y,z) = (x^{yz}\ln x)(y) = yx^{yz}\ln x$, so $f_z(e,1,0) = 1e^{(1)(0)}\ln e = 1$.

41. $x^2+2y^2+3z^2 = 1 \quad \Rightarrow \quad \dfrac{\partial}{\partial x}\left(x^2+2y^2+3z^2\right) = \dfrac{\partial}{\partial x}(1) \quad \Rightarrow \quad 2x+0+6z\dfrac{\partial z}{\partial x} = 0 \quad \Rightarrow \quad 6z\dfrac{\partial z}{\partial x} = -2x \quad \Rightarrow$

$\dfrac{\partial z}{\partial x} = \dfrac{-2x}{6z} = -\dfrac{x}{3z}$, and $\dfrac{\partial}{\partial y}\left(x^2+2y^2+3z^2\right) = \dfrac{\partial}{\partial y}(1) \quad \Rightarrow \quad 0+4y+6z\dfrac{\partial z}{\partial y} = 0 \quad \Rightarrow \quad 6z\dfrac{\partial z}{\partial y} = -4y \quad \Rightarrow$

$\dfrac{\partial z}{\partial y} = \dfrac{-4y}{6z} = -\dfrac{2y}{3z}$.

42. $x^2-y^2+z^2-2z = 4 \quad \Rightarrow \quad \dfrac{\partial}{\partial x}\left(x^2-y^2+z^2-2z\right) = \dfrac{\partial}{\partial x}(4) \quad \Rightarrow \quad 2x-0+2z\dfrac{\partial z}{\partial x} - 2\dfrac{\partial z}{\partial x} = 0 \quad \Rightarrow$

$(2z-2)\dfrac{\partial z}{\partial x} = -2x \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = \dfrac{-2x}{2z-2} = \dfrac{x}{1-z}$, and $\dfrac{\partial}{\partial y}\left(x^2-y^2+z^2-2z\right) = \dfrac{\partial}{\partial y}(4) \quad \Rightarrow$

$0-2y+2z\dfrac{\partial z}{\partial y} - 2\dfrac{\partial z}{\partial y} = 0 \quad \Rightarrow \quad (2z-2)\dfrac{\partial z}{\partial y} = 2y \quad \Rightarrow \quad \dfrac{\partial z}{\partial y} = \dfrac{2y}{2z-2} = \dfrac{y}{z-1}$.

43. $e^z = xyz \quad \Rightarrow \quad \dfrac{\partial}{\partial x}\left(e^z\right) = \dfrac{\partial}{\partial x}(xyz) \quad \Rightarrow \quad e^z\dfrac{\partial z}{\partial x} = y\left(x\dfrac{\partial z}{\partial x} + z\cdot 1\right) \quad \Rightarrow \quad e^z\dfrac{\partial z}{\partial x} - xy\dfrac{\partial z}{\partial x} = yz \quad \Rightarrow$

$\left(e^z - xy\right)\dfrac{\partial z}{\partial x} = yz$, so $\dfrac{\partial z}{\partial x} = \dfrac{yz}{e^z - xy}$.

$\dfrac{\partial}{\partial y}\left(e^z\right) = \dfrac{\partial}{\partial y}(xyz) \quad \Rightarrow \quad e^z\dfrac{\partial z}{\partial y} = x\left(y\dfrac{\partial z}{\partial x} + z\cdot 1\right) \quad \Rightarrow \quad e^z\dfrac{\partial z}{\partial y} - xy\dfrac{\partial z}{\partial y} = xz \quad \Rightarrow \quad \left(e^z - xy\right)\dfrac{\partial z}{\partial y} = xz$, so

$\dfrac{\partial z}{\partial y} = \dfrac{xz}{e^z - xy}$.

44. $yz + x\ln y = z^2 \quad \Rightarrow \quad \dfrac{\partial}{\partial x}(yz + x\ln y) = \dfrac{\partial}{\partial x}\left(z^2\right) \quad \Rightarrow \quad y\dfrac{\partial z}{\partial x} + \ln y = 2z\dfrac{\partial z}{\partial x} \quad \Rightarrow \quad \ln y = 2z\dfrac{\partial z}{\partial x} - y\dfrac{\partial z}{\partial x} \quad \Rightarrow$

$\ln y = (2z-y)\dfrac{\partial z}{\partial x}$, so $\dfrac{\partial z}{\partial x} = \dfrac{\ln y}{2z-y}$.

$\dfrac{\partial}{\partial y}(yz + x\ln y) = \dfrac{\partial}{\partial y}\left(z^2\right) \quad \Rightarrow \quad y\dfrac{\partial z}{\partial y} + z\cdot 1 + x\cdot\dfrac{1}{y} = 2z\dfrac{\partial z}{\partial y} \quad \Rightarrow \quad z + \dfrac{x}{y} = 2z\dfrac{\partial z}{\partial y} - y\dfrac{\partial z}{\partial y} \quad \Rightarrow$

$z + \dfrac{x}{y} = (2z-y)\dfrac{\partial z}{\partial y}$, so $\dfrac{\partial z}{\partial y} = \dfrac{z + (x/y)}{2z-y} = \dfrac{x+yz}{y(2z-y)}$.

45. (a) $z = f(x) + g(y) \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = f'(x), \quad \dfrac{\partial z}{\partial y} = g'(y)$

(b) $z = f(x+y)$. Let $u = x+y$. Then $\dfrac{\partial z}{\partial x} = \dfrac{df}{du}\dfrac{\partial u}{\partial x} = \dfrac{df}{du}(1) = f'(u) = f'(x+y)$,

$\dfrac{\partial z}{\partial y} = \dfrac{df}{du}\dfrac{\partial u}{\partial y} = \dfrac{df}{du}(1) = f'(u) = f'(x+y)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. (a) $z = f(x)g(y) \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = f'(x)g(y), \quad \dfrac{\partial z}{\partial y} = f(x)g'(y)$

(b) $z = f(xy)$. Let $u = xy$. Then $\dfrac{\partial u}{\partial x} = y$ and $\dfrac{\partial u}{\partial y} = x$. Hence $\dfrac{\partial z}{\partial x} = \dfrac{df}{du}\dfrac{\partial u}{\partial x} = \dfrac{df}{du} \cdot y = yf'(u) = yf'(xy)$

and $\dfrac{\partial z}{\partial y} = \dfrac{df}{du}\dfrac{\partial u}{\partial y} = \dfrac{df}{du} \cdot x = xf'(u) = xf'(xy)$.

(c) $z = f\left(\dfrac{x}{y}\right)$. Let $u = \dfrac{x}{y}$. Then $\dfrac{\partial u}{\partial x} = \dfrac{1}{y}$ and $\dfrac{\partial u}{\partial y} = -\dfrac{x}{y^2}$. Hence $\dfrac{\partial z}{\partial x} = \dfrac{df}{du}\dfrac{\partial u}{\partial x} = f'(u)\dfrac{1}{y} = \dfrac{f'(x/y)}{y}$

and $\dfrac{\partial z}{\partial y} = \dfrac{df}{du}\dfrac{\partial u}{\partial y} = f'(u)\left(-\dfrac{x}{y^2}\right) = -\dfrac{xf'(x/y)}{y^2}$.

47. $f(x,y) = x^4y - 2x^3y^2 \quad \Rightarrow \quad f_x(x,y) = 4x^3y - 6x^2y^2, \;\; f_y(x,y) = x^4 - 4x^3y$. Then $f_{xx}(x,y) = 12x^2y - 12xy^2$, $f_{xy}(x,y) = 4x^3 - 12x^2y, \;\; f_{yx}(x,y) = 4x^3 - 12x^2y$, and $f_{yy}(x,y) = -4x^3$.

48. $f(x,y) = \ln(ax+by) \quad \Rightarrow \quad f_x(x,y) = \dfrac{a}{ax+by} = a(ax+by)^{-1}, \;\; f_y(x,y) = \dfrac{b}{ax+by} = b(ax+by)^{-1}$. Then

$f_{xx}(x,y) = -a(ax+by)^{-2}(a) = -\dfrac{a^2}{(ax+by)^2}, \;\; f_{xy}(x,y) = -a(ax+by)^{-2}(b) = -\dfrac{ab}{(ax+by)^2}$,

$f_{yx}(x,y) = -b(ax+by)^{-2}(a) = -\dfrac{ab}{(ax+by)^2}$, and $f_{yy}(x,y) = -b(ax+by)^{-2}(b) = -\dfrac{b^2}{(ax+by)^2}$.

49. $z = \dfrac{y}{2x+3y} = y(2x+3y)^{-1} \quad \Rightarrow \quad z_x = y(-1)(2x+3y)^{-2}(2) = -\dfrac{2y}{(2x+3y)^2}$,

$z_y = \dfrac{(2x+3y)\cdot 1 - y\cdot 3}{(2x+3y)^2} = \dfrac{2x}{(2x+3y)^2}$. Then $z_{xx} = -2y(-2)(2x+3y)^{-3}(2) = \dfrac{8y}{(2x+3y)^3}$,

$z_{xy} = -\dfrac{(2x+3y)^2\cdot 2 - 2y\cdot 2(2x+3y)(3)}{\left[(2x+3y)^2\right]^2} = -\dfrac{(2x+3y)\,(4x+6y-12y)}{(2x+3y)^4} = \dfrac{6y-4x}{(2x+3y)^3}$,

$z_{yx} = \dfrac{(2x+3y)^2\cdot 2 - 2x\cdot 2(2x+3y)(2)}{\left[(2x+3y)^2\right]^2} = \dfrac{6y-4x}{(2x+3y)^3}, \;\; z_{yy} = 2x(-2)(2x+3y)^{-3}(3) = -\dfrac{12x}{(2x+3y)^3}$.

50. $T = e^{-2r}\cos\theta \quad \Rightarrow \quad T_r = -2e^{-2r}\cos\theta, \;\; T_\theta = -e^{-2r}\sin\theta$. Then $T_{rr} = -2e^{-2r}(-2)\cos\theta = 4e^{-2r}\cos\theta$,

$T_{r\theta} = 2e^{-2r}\sin\theta, \;\; T_{\theta r} = -e^{-2r}(-2)\sin\theta = 2e^{-2r}\sin\theta, \;\; T_{\theta\theta} = -e^{-2r}\cos\theta$.

51. $v = \sin(s^2-t^2) \quad \Rightarrow \quad v_s = \cos(s^2-t^2)\cdot 2s = 2s\cos(s^2-t^2), \;\; v_t = \cos(s^2-t^2)\cdot(-2t) = -2t\cos(s^2-t^2)$. Then

$v_{ss} = 2s\left[-\sin(s^2-t^2)\cdot 2s\right] + \cos(s^2-t^2)\cdot 2 = 2\cos(s^2-t^2) - 4s^2\sin(s^2-t^2)$,

$v_{st} = 2s\left[-\sin(s^2-t^2)\cdot(-2t)\right] + \cos(s^2-t^2)\cdot 0 = 4st\sin(s^2-t^2)$,

$v_{ts} = -2t\left[-\sin(s^2-t^2)\cdot 2s\right] + \cos(s^2-t^2)\cdot 0 = 4st\sin(s^2-t^2)$,

$v_{tt} = -2t\left[-\sin(s^2-t^2)\cdot(-2t)\right] + \cos(s^2-t^2)\cdot(-2) = -2\cos(s^2-t^2) - 4t^2\sin(s^2-t^2)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

52. $z = \arctan \dfrac{x+y}{1-xy} \quad \Rightarrow$

$$z_x = \frac{1}{1+\left(\dfrac{x+y}{1-xy}\right)^2} \cdot \frac{(1-xy)(1)-(x+y)(-y)}{(1-xy)^2} = \frac{1+y^2}{(1-xy)^2+(x+y)^2} = \frac{1+y^2}{1+x^2+y^2+x^2y^2}$$

$$= \frac{1+y^2}{(1+x^2)(1+y^2)} = \frac{1}{1+x^2},$$

$$z_y = \frac{1}{1+\left(\dfrac{x+y}{1-xy}\right)^2} \cdot \frac{(1-xy)(1)-(x+y)(-x)}{(1-xy)^2} = \frac{1+x^2}{(1-xy)^2+(x+y)^2} = \frac{1+x^2}{(1+x^2)(1+y^2)} = \frac{1}{1+y^2}.$$

Then $z_{xx} = -(1+x^2)^{-2} \cdot 2x = -\dfrac{2x}{(1+x^2)^2}$, $z_{xy} = 0$, $z_{yx} = 0$, $z_{yy} = -(1+y^2)^{-2} \cdot 2y = -\dfrac{2y}{(1+y^2)^2}$.

53. $u = x^4y^3 - y^4 \quad \Rightarrow \quad u_x = 4x^3y^3$, $u_{xy} = 12x^3y^2$ and $u_y = 3x^4y^2 - 4y^3$, $u_{yx} = 12x^3y^2$.

Thus $u_{xy} = u_{yx}$.

54. $u = e^{xy}\sin y \quad \Rightarrow \quad u_x = ye^{xy}\sin y$, $u_{xy} = ye^{xy}\cos y + (\sin y)(y \cdot xe^{xy} + e^{xy} \cdot 1) = e^{xy}(y\cos y + xy\sin y + \sin y)$,

$u_y = e^{xy}\cos y + (\sin y)(xe^{xy}) = e^{xy}(\cos y + x\sin y)$,

$u_{yx} = e^{xy} \cdot \sin y + (\cos y + x\sin y) \cdot ye^{xy} = e^{xy}(\sin y + y\cos y + xy\sin y)$. Thus $u_{xy} = u_{yx}$.

55. $u = \cos(x^2y) \quad \Rightarrow \quad u_x = -\sin(x^2y) \cdot 2xy = -2xy\sin(x^2y)$,

$u_{xy} = -2xy \cdot \cos(x^2y) \cdot x^2 + \sin(x^2y) \cdot (-2x) = -2x^3y\cos(x^2y) - 2x\sin(x^2y)$ and

$u_y = -\sin(x^2y) \cdot x^2 = -x^2\sin(x^2y)$, $u_{yx} = -x^2 \cdot \cos(x^2y) \cdot 2xy + \sin(x^2y) \cdot (-2x) = -2x^3y\cos(x^2y) - 2x\sin(x^2y)$.

Thus $u_{xy} = u_{yx}$.

56. $u = \ln(x+2y) \quad \Rightarrow \quad u_x = \dfrac{1}{x+2y} = (x+2y)^{-1}$, $u_{xy} = (-1)(x+2y)^{-2}(2) = -\dfrac{2}{(x+2y)^2}$ and

$u_y = \dfrac{1}{x+2y} \cdot 2 = 2(x+2y)^{-1}$, $u_{yx} = (-2)(x+2y)^{-2} = -\dfrac{2}{(x+2y)^2}$. Thus $u_{xy} = u_{yx}$.

57. $f(x,y) = x^4y^2 - x^3y \quad \Rightarrow \quad f_x = 4x^3y^2 - 3x^2y$, $f_{xx} = 12x^2y^2 - 6xy$, $f_{xxx} = 24xy^2 - 6y$ and

$f_{xy} = 8x^3y - 3x^2$, $f_{xyx} = 24x^2y - 6x$.

58. $f(x,y) = \sin(2x+5y) \quad \Rightarrow \quad f_y = \cos(2x+5y) \cdot 5 = 5\cos(2x+5y)$, $f_{yx} = -5\sin(2x+5y) \cdot 2 = -10\sin(2x+5y)$,

$f_{yxy} = -10\cos(2x+5y) \cdot 5 = -50\cos(2x+5y)$

59. $f(x,y,z) = e^{xyz^2} \quad \Rightarrow \quad f_x = e^{xyz^2} \cdot yz^2 = yz^2e^{xyz^2}$, $f_{xy} = yz^2 \cdot e^{xyz^2}(xz^2) + e^{xyz^2} \cdot z^2 = (xyz^4 + z^2)e^{xyz^2}$,

$f_{xyz} = (xyz^4 + z^2) \cdot e^{xyz^2}(2xyz) + e^{xyz^2} \cdot (4xyz^3 + 2z) = (2x^2y^2z^5 + 6xyz^3 + 2z)e^{xyz^2}$.

60. $g(r,s,t) = e^r\sin(st) \quad \Rightarrow \quad g_r = e^r\sin(st)$, $g_{rs} = e^r\cos(st) \cdot t = te^r\cos(st)$,

$g_{rst} = te^r(-\sin(st) \cdot s) + \cos(st) \cdot e^r = e^r[\cos(st) - st\sin(st)]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

61. $W = \sqrt{u+v^2} \quad \Rightarrow \quad \dfrac{\partial W}{\partial v} = \frac{1}{2}(u+v^2)^{-1/2}(2v) = v(u+v^2)^{-1/2}$,

$$\frac{\partial^2 W}{\partial u\,\partial v} = v\left(-\tfrac{1}{2}\right)(u+v^2)^{-3/2}(1) = -\tfrac{1}{2}v(u+v^2)^{-3/2}, \quad \frac{\partial^3 W}{\partial u^2\,\partial v} = -\tfrac{1}{2}v\left(-\tfrac{3}{2}\right)(u+v^2)^{-5/2}(1) = \tfrac{3}{4}v(u+v^2)^{-5/2}.$$

62. $V = \ln(r+s^2+t^3) \quad \Rightarrow \quad \dfrac{\partial V}{\partial t} = \dfrac{3t^2}{r+s^2+t^3} = 3t^2(r+s^2+t^3)^{-1}$,

$$\frac{\partial^2 V}{\partial s\,\partial t} = 3t^2(-1)(r+s^2+t^3)^{-2}(2s) = -6st^2(r+s^2+t^3)^{-2},$$

$$\frac{\partial^3 V}{\partial r\,\partial s\,\partial t} = -6st^2(-2)(r+s^2+t^3)^{-3}(1) = 12st^2(r+s^2+t^3)^{-3} = \frac{12st^2}{(r+s^2+t^3)^3}.$$

63. $w = \dfrac{x}{y+2z} = x(y+2z)^{-1} \quad \Rightarrow \quad \dfrac{\partial w}{\partial x} = (y+2z)^{-1}, \quad \dfrac{\partial^2 w}{\partial y\,\partial x} = -(y+2z)^{-2}(1) = -(y+2z)^{-2}$,

$$\frac{\partial^3 w}{\partial z\,\partial y\,\partial x} = -(-2)(y+2z)^{-3}(2) = 4(y+2z)^{-3} = \frac{4}{(y+2z)^3} \quad \text{and} \quad \frac{\partial w}{\partial y} = x(-1)(y+2z)^{-2}(1) = -x(y+2z)^{-2},$$

$$\frac{\partial^2 w}{\partial x\,\partial y} = -(y+2z)^{-2}, \quad \frac{\partial^3 w}{\partial x^2\,\partial y} = 0.$$

64. $u = x^a y^b z^c$. If $a=0$, or if $b=0$ or 1, or if $c=0$, 1, or 2, then $\dfrac{\partial^6 u}{\partial x\,\partial y^2\,\partial z^3} = 0$. Otherwise $\dfrac{\partial u}{\partial z} = cx^a y^b z^{c-1}$,

$$\frac{\partial^2 u}{\partial z^2} = c(c-1)x^a y^b z^{c-2}, \quad \frac{\partial^3 u}{\partial z^3} = c(c-1)(c-2)x^a y^b z^{c-3}, \quad \frac{\partial^4 u}{\partial y\,\partial z^3} = bc(c-1)(c-2)x^a y^{b-1} z^{c-3},$$

$$\frac{\partial^5 u}{\partial y^2\,\partial z^3} = b(b-1)c(c-1)(c-2)x^a y^{b-2} z^{c-3}, \text{ and } \frac{\partial^6 u}{\partial x\,\partial y^2\,\partial z^3} = ab(b-1)c(c-1)(c-2)x^{a-1} y^{b-2} z^{c-3}.$$

65. $f(x,y) = xy^2 - x^3 y \quad \Rightarrow$

$$\begin{aligned} f_x(x,y) &= \lim_{h\to 0} \frac{f(x+h,y)-f(x,y)}{h} = \lim_{h\to 0} \frac{(x+h)y^2-(x+h)^3 y-(xy^2-x^3y)}{h} \\ &= \lim_{h\to 0} \frac{h(y^2-3x^2y-3xyh-yh^2)}{h} = \lim_{h\to 0}(y^2-3x^2y-3xyh-yh^2) = y^2-3x^2y \end{aligned}$$

$$\begin{aligned} f_y(x,y) &= \lim_{h\to 0} \frac{f(x,y+h)-f(x,y)}{h} = \lim_{h\to 0} \frac{x(y+h)^2-x^3(y+h)-(xy^2-x^3y)}{h} = \lim_{h\to 0} \frac{h(2xy+xh-x^3)}{h} \\ &= \lim_{h\to 0}(2xy+xh-x^3) = 2xy-x^3 \end{aligned}$$

66. $f(x,y) = \dfrac{x}{x+y^2} \quad \Rightarrow$

$$\begin{aligned} f_x(x,y) &= \lim_{h\to 0} \frac{f(x+h,y)-f(x,y)}{h} = \lim_{h\to 0} \frac{\frac{x+h}{x+h+y^2} - \frac{x}{x+y^2}}{h} \cdot \frac{(x+h+y^2)(x+y^2)}{(x+h+y^2)(x+y^2)} \\ &= \lim_{h\to 0} \frac{(x+h)(x+y^2)-x(x+h+y^2)}{h(x+h+y^2)(x+y^2)} = \lim_{h\to 0} \frac{y^2 h}{h(x+h+y^2)(x+y^2)} \\ &= \lim_{h\to 0} \frac{y^2}{(x+h+y^2)(x+y^2)} = \frac{y^2}{(x+y^2)^2} \end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$f_y(x,y) = \lim_{h\to 0} \frac{f(x,y+h)-f(x,y)}{h} = \lim_{h\to 0} \frac{\frac{x}{x+(y+h)^2} - \frac{x}{x+y^2}}{h} \cdot \frac{\left[x+(y+h)^2\right]\left(x+y^2\right)}{\left[x+(y+h)^2\right]\left(x+y^2\right)}$$

$$= \lim_{h\to 0} \frac{x(x+y^2) - x\left[x+(y+h)^2\right]}{h[x+(y+h)^2](x+y^2)} = \lim_{h\to 0} \frac{h(-2xy-xh)}{h[x+(y+h)^2](x+y^2)}$$

$$= \lim_{h\to 0} \frac{-2xy-xh}{[x+(y+h)^2](x+y^2)} = \frac{-2xy}{(x+y^2)^2}$$

67. Assuming that the third partial derivatives of f are continuous (easily verified), we can write $f_{xzy} = f_{yxz}$. Then

$f(x,y,z) = xy^2z^3 + \arcsin\left(x\sqrt{z}\right) \quad \Rightarrow \quad f_y = 2xyz^3 + 0$, $f_{yx} = 2yz^3$, and $f_{yxz} = 6yz^2 = f_{xzy}$.

68. Let $f(x,y,z) = \sqrt{1+xz}$ and $h(x,y,z) = \sqrt{1-xy}$ so that $g = f + h$. Then $f_y = 0 = f_{yx} = f_{yxz}$ and $h_z = 0 = h_{zx} = h_{zxy}$. But (since the partial derivatives are continous on their domains) $f_{xyz} = f_{yxz}$ and $h_{xyz} = h_{zxy}$, so $g_{xyz} = f_{xyz} + h_{xyz} = 0 + 0 = 0$.

69. First of all, if we start at the point $(3,-3)$ and move in the positive y-direction, we see that both b and c decrease, while a increases. Both b and c have a low point at about $(3,-1.5)$, while a is 0 at this point. So a is definitely the graph of f_y, and one of b and c is the graph of f. To see which is which, we start at the point $(-3,-1.5)$ and move in the positive x-direction. b traces out a line with negative slope, while c traces out a parabola opening downward. This tells us that b is the x-derivative of c. So c is the graph of f, b is the graph of f_x, and a is the graph of f_y.

70. $f(x,y) = \dfrac{y}{1+x^2y^2} \quad \Rightarrow \quad f_x = \dfrac{(1+x^2y^2)(0) - y(2xy^2)}{(1+x^2y^2)^2} = -\dfrac{2xy^3}{(1+x^2y^2)^2}$,

$f_y = \dfrac{(1+x^2y^2)(1) - y(2x^2y)}{(1+x^2y^2)^2} = \dfrac{1-x^2y^2}{(1+x^2y^2)^2}$

Note that traces of f in planes parallel to the xz-plane have only one extreme value (a minimum for $y < 0$, a maximum for $y > 0$), and the traces of f_x in these planes have only one zero (going from negative to positive if $y < 0$ and from positive to negative if $y > 0$). The traces of f in planes parallel to the yz-plane have two extreme values, and the traces of f_y in these planes have two zeros.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

71. $f(x,y) = x^2y^3 \quad \Rightarrow \quad f_x = 2xy^3, \quad f_y = 3x^2y^2$

Note that traces of f in planes parallel to the xz-plane are parabolas which open downward for $y < 0$ and upward for $y > 0$, and the traces of f_x in these planes are straight lines, which have negative slopes for $y < 0$ and positive slopes for $y > 0$. The traces of f in planes parallel to the yz-plane are cubic curves, and the traces of f_y in these planes are parabolas.

72. (a) $f_{xx} = \frac{\partial}{\partial x}(f_x)$, so f_{xx} is the rate of change of f_x in the x-direction. f_x is negative at $(-1, 2)$ and if we move in the positive x-direction, the surface becomes less steep. Thus the values of f_x are increasing and $f_{xx}(-1, 2)$ is positive.

(b) f_{yy} is the rate of change of f_y in the y-direction. f_y is negative at $(-1, 2)$ and if we move in the positive y-direction, the surface becomes steeper. Thus the values of f_y are decreasing, and $f_{yy}(-1, 2)$ is negative.

(c) $f_{xy} = \frac{\partial}{\partial y}(f_x)$, so f_{xy} is the rate of change of f_x in the y-direction. f_x is positive at $(1, 2)$ and if we move in the positive y-direction, the surface becomes steeper, looking in the positive x-direction. Thus the values of f_x are increasing and $f_{xy}(1, 2)$ is positive.

(d) f_x is negative at $(-1, 2)$ and if we move in the positive y-direction, the surface gets steeper (with negative slope), looking in the positive x-direction. This means that the values of f_x are decreasing as y increases, so $f_{xy}(-1, 2)$ is negative.

73. By Definition 4, $f_x(3, 2) = \lim\limits_{h \to 0} \dfrac{f(3+h, 2) - f(3, 2)}{h}$ which we can approximate by considering $h = 0.5$ and $h = -0.5$:

$f_x(3, 2) \approx \dfrac{f(3.5, 2) - f(3, 2)}{0.5} = \dfrac{22.4 - 17.5}{0.5} = 9.8$, $f_x(3, 2) \approx \dfrac{f(2.5, 2) - f(3, 2)}{-0.5} = \dfrac{10.2 - 17.5}{-0.5} = 14.6$. Averaging these values, we estimate $f_x(3, 2)$ to be approximately 12.2.

Similarly, $f_x(3, 2.2) = \lim\limits_{h \to 0} \dfrac{f(3+h, 2.2) - f(3, 2.2)}{h}$ which we can approximate by considering $h = 0.5$ and $h = -0.5$:

$f_x(3, 2.2) \approx \dfrac{f(3.5, 2.2) - f(3, 2.2)}{0.5} = \dfrac{26.1 - 15.9}{0.5} = 20.4$, $f_x(3, 2.2) \approx \dfrac{f(2.5, 2.2) - f(3, 2.2)}{-0.5} = \dfrac{9.3 - 15.9}{-0.5} = 13.2$.

Averaging these values, we have $f_x(3, 2.2) \approx 16.8$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

To estimate $f_{xy}(3,2)$, we first need an estimate for $f_x(3,1.8)$:

$$f_x(3,1.8) \approx \frac{f(3.5,1.8)-f(3,1.8)}{0.5} = \frac{20.0-18.1}{0.5} = 3.8,\ f_x(3,1.8) \approx \frac{f(2.5,1.8)-f(3,1.8)}{-0.5} = \frac{12.5-18.1}{-0.5} = 11.2.$$

Averaging these values, we get $f_x(3,1.8) \approx 7.5$. Now $f_{xy}(x,y) = \dfrac{\partial}{\partial y}\left[f_x(x,y)\right]$ and $f_x(x,y)$ is itself a function of two variables, so Definition 4 says that $f_{xy}(x,y) = \dfrac{\partial}{\partial y}\left[f_x(x,y)\right] = \lim\limits_{h\to 0}\dfrac{f_x(x,y+h)-f_x(x,y)}{h} \quad \Rightarrow$

$f_{xy}(3,2) = \lim\limits_{h\to 0}\dfrac{f_x(3,2+h)-f_x(3,2)}{h}$. We can estimate this value using our previous work with $h=0.2$ and $h=-0.2$:

$$f_{xy}(3,2) \approx \frac{f_x(3,2.2)-f_x(3,2)}{0.2} = \frac{16.8-12.2}{0.2} = 23,\ f_{xy}(3,2) \approx \frac{f_x(3,1.8)-f_x(3,2)}{-0.2} = \frac{7.5-12.2}{-0.2} = 23.5.$$

Averaging these values, we estimate $f_{xy}(3,2)$ to be approximately 23.25.

74. (a) If we fix y and allow x to vary, the level curves indicate that the value of f decreases as we move through P in the positive x-direction, so f_x is negative at P.

(b) If we fix x and allow y to vary, the level curves indicate that the value of f increases as we move through P in the positive y-direction, so f_y is positive at P.

(c) $f_{xx} = \dfrac{\partial}{\partial x}(f_x)$, so if we fix y and allow x to vary, f_{xx} is the rate of change of f_x as x increases. Note that at points to the right of P the level curves are spaced farther apart (in the x-direction) than at points to the left of P, demonstrating that f decreases less quickly with respect to x to the right of P. So as we move through P in the positive x-direction the (negative) value of f_x increases, hence $\dfrac{\partial}{\partial x}(f_x) = f_{xx}$ is positive at P.

(d) $f_{xy} = \dfrac{\partial}{\partial y}(f_x)$, so if we fix x and allow y to vary, f_{xy} is the rate of change of f_x as y increases. The level curves are closer together (in the x-direction) at points above P than at those below P, demonstrating that f decreases more quickly with respect to x for y-values above P. So as we move through P in the positive y-direction, the (negative) value of f_x decreases, hence f_{xy} is negative.

(e) $f_{yy} = \dfrac{\partial}{\partial y}(f_y)$, so if we fix x and allow y to vary, f_{yy} is the rate of change of f_y as y increases. The level curves are closer together (in the y-direction) at points above P than at those below P, demonstrating that f increases more quickly with respect to y above P. So as we move through P in the positive y-direction the (positive) value of f_y increases, hence $\dfrac{\partial}{\partial y}(f_y) = f_{yy}$ is positive at P.

75. (a) $f(x,y) = 4 - x^2 - 2y^2$. In the plane $y=1$, $f(x,1) = 4 - x^2 - 2(1^2) = 2 - x^2$, so a vector equation for C_1 is given by $\mathbf{r}(t) = \left\langle t, 1, 2-t^2\right\rangle$ where the point $(1,1,1)$ corresponds to $t=1$. Then $\mathbf{r}'(t) = \langle 1, 0, -2t\rangle \quad \Rightarrow \quad \mathbf{r}'(1) = \langle 1, 0, -2\rangle$ and parametric equations of the tangent line are $x = t+1$, $y=1$, $z=-2t+1$. Thus, $x = t+1 \quad \Rightarrow \quad x-1 = t \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$z = -2(x-1)+1 = -2x+3$. So the equation of the tangent line can be given by $z = -2x+3$, $y = 1$ which has a slope $m = -2$. Therefore, $f_x(1,1) = -2$.

(b) In the plane $x = 1$, $f(1,y) = 4 - 1^2 - 2y^2 = 3 - 2y^2$, so a vector equation for C_2 is given by $\mathbf{r}(t) = \langle 1, t, 3-2t^2 \rangle$ where the point $(1,1,1)$ corresponds to $t = 1$. Then $\mathbf{r}'(t) = \langle 0, 1, -4t \rangle \quad \Rightarrow \quad \mathbf{r}'(1) = \langle 0, 1, -4 \rangle$ and parametric equations of the tangent line are $x = 1, y = t+1, z = -4t+1$. Thus, $y = t+1 \quad \Rightarrow \quad y - 1 = t \quad \Rightarrow$ $z = -4(y-1)+1 = -4y+5$. So the equation of the tangent line can be given by $z = -4y+5$, $x = 1$ which has a slope $m = -4$. Therefore, $f_y(1,1) = -4$.

76. For each i, $i = 1, \ldots, n$, $\partial u/\partial x_i = a_i e^{a_1x_1+a_2x_2+\cdots+a_nx_n}$ and $\partial^2 u/\partial x_i^2 = a_i^2 e^{a_1x_1+a_2x_2+\cdots+a_nx_n}$.

Then $\dfrac{\partial^2 u}{\partial x_1^2} + \dfrac{\partial^2 u}{\partial x_2^2} + \cdots + \dfrac{\partial^2 u}{\partial x_n^2} = \left(a_1^2 + a_2^2 + \cdots + a_n^2\right) e^{a_1x_1+a_2x_2+\cdots+a_nx_n} = e^{a_1x_1+a_2x_2+\cdots+a_nx_n} = u$

since $a_1^2 + a_2^2 + \cdots + a_n^2 = 1$.

77. (a) $u = \sin(kx)\,\sin(akt) \quad \Rightarrow \quad u_t = ak\sin(kx)\,\cos(akt)$, $u_{tt} = -a^2k^2\sin(kx)\,\sin(akt)$, $u_x = k\cos(kx)\,\sin(akt)$, $u_{xx} = -k^2\sin(kx)\,\sin(akt)$. Thus $u_{tt} = a^2u_{xx}$.

(b) $u = \dfrac{t}{a^2t^2 - x^2} \quad \Rightarrow \quad u_t = \dfrac{(a^2t^2 - x^2) - t(2a^2t)}{(a^2t^2 - x^2)^2} = -\dfrac{a^2t^2 + x^2}{(a^2t^2 - x^2)^2}$,

$$u_{tt} = \frac{-2a^2t(a^2t^2 - x^2)^2 + (a^2t^2 + x^2)(2)(a^2t^2 - x^2)(2a^2t)}{(a^2t^2 - x^2)^4} = \frac{2a^4t^3 + 6a^2tx^2}{(a^2t^2 - x^2)^3},$$

$$u_x = t(-1)(a^2t^2 - x^2)^{-2}(-2x) = \frac{2tx}{(a^2t^2 - x^2)^2},$$

$$u_{xx} = \frac{2t(a^2t^2 - x^2)^2 - 2tx\,(2)(a^2t^2 - x^2)(-2x)}{(a^2t^2 - x^2)^4} = \frac{2a^2t^3 - 2tx^2 + 8tx^2}{(a^2t^2 - x^2)^3} = \frac{2a^2t^3 + 6tx^2}{(a^2t^2 - x^2)^3}.$$

Thus $u_{tt} = a^2u_{xx}$.

(c) $u = (x - at)^6 + (x + at)^6 \quad \Rightarrow \quad u_t = -6a(x - at)^5 + 6a(x + at)^5$, $u_{tt} = 30a^2(x - at)^4 + 30a^2(x + at)^4$, $u_x = 6(x - at)^5 + 6(x + at)^5$, $u_{xx} = 30(x - at)^4 + 30(x + at)^4$. Thus $u_{tt} = a^2u_{xx}$.

(d) $u = \sin(x - at) + \ln(x + at) \quad \Rightarrow \quad u_t = -a\cos(x - at) + \dfrac{a}{x + at}$, $\quad u_{tt} = -a^2\sin(x - at) - \dfrac{a^2}{(x + at)^2}$,

$u_x = \cos(x - at) + \dfrac{1}{x + at}$, $\quad u_{xx} = -\sin(x - at) - \dfrac{1}{(x + at)^2}$. Thus $u_{tt} = a^2u_{xx}$.

78. (a) $u = x^2 + y^2 \quad \Rightarrow \quad u_x = 2x,\ u_{xx} = 2; \quad u_y = 2y,\ u_{yy} = 2$. Thus $u_{xx} + u_{yy} \neq 0$ and $u = x^2 + y^2$ does not satisfy Laplace's Equation.

(b) $u = x^2 - y^2$ is a solution: $u_{xx} = 2,\ u_{yy} = -2$ so $u_{xx} + u_{yy} = 0$.

(c) $u = x^3 + 3xy^2$ is not a solution: $u_x = 3x^2 + 3y^2,\ u_{xx} = 6x; \quad u_y = 6xy,\ u_{yy} = 6x$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(d) $u = \ln\sqrt{x^2+y^2}$ is a solution: $u_x = \dfrac{1}{\sqrt{x^2+y^2}}\left(\dfrac{1}{2}\right)(x^2+y^2)^{-1/2}(2x) = \dfrac{x}{x^2+y^2}$,

$u_{xx} = \dfrac{(x^2+y^2)-x(2x)}{(x^2+y^2)^2} = \dfrac{y^2-x^2}{(x^2+y^2)^2}$. By symmetry, $u_{yy} = \dfrac{x^2-y^2}{(x^2+y^2)^2}$, so $u_{xx}+u_{yy}=0$.

(e) $u = \sin x\,\cosh y + \cos x\,\sinh y$ is a solution: $u_x = \cos x\,\cosh y - \sin x\,\sinh y$, $u_{xx} = -\sin x\,\cosh y - \cos x\,\sinh y$, and $u_y = \sin x\,\sinh y + \cos x\,\cosh y$, $u_{yy} = \sin x\,\cosh y + \cos x\,\sinh y$.

(f) $u = e^{-x}\cos y - e^{-y}\cos x$ is a solution: $u_x = -e^{-x}\cos y + e^{-y}\sin x$, $\quad u_{xx} = e^{-x}\cos y + e^{-y}\cos x$, and $u_y = -e^{-x}\sin y + e^{-y}\cos x$, $\quad u_{yy} = -e^{-x}\cos y - e^{-y}\cos x$.

79. $u = \dfrac{1}{\sqrt{x^2+y^2+z^2}} \quad\Rightarrow\quad u_x = \left(-\frac{1}{2}\right)(x^2+y^2+z^2)^{-3/2}(2x) = -x(x^2+y^2+z^2)^{-3/2}$ and

$u_{xx} = -(x^2+y^2+z^2)^{-3/2} - x\left(-\frac{3}{2}\right)(x^2+y^2+z^2)^{-5/2}(2x) = \dfrac{2x^2-y^2-z^2}{(x^2+y^2+z^2)^{5/2}}$.

By symmetry, $u_{yy} = \dfrac{2y^2-x^2-z^2}{(x^2+y^2+z^2)^{5/2}}$ and $u_{zz} = \dfrac{2z^2-x^2-y^2}{(x^2+y^2+z^2)^{5/2}}$.

Thus, $u_{xx}+u_{yy}+u_{zz} = \dfrac{2x^2-y^2-z^2+2y^2-x^2-z^2+2z^2-x^2-y^2}{(x^2+y^2+z^2)^{5/2}} = 0$.

80. $u = e^{-\alpha^2k^2t}\sin kx \quad\Rightarrow\quad u_x = ke^{-\alpha^2k^2t}\cos kx$, $\quad u_{xx} = -k^2e^{-\alpha^2k^2t}\sin kx$, and $\quad u_t = -\alpha^2k^2e^{-\alpha^2k^2t}\sin kx$.

Thus, $\alpha^2u_{xx} = u_t$.

81. $c(x,t) = \dfrac{1}{\sqrt{4\pi Dt}}\,e^{-x^2/(4Dt)} \quad\Rightarrow$

$$\frac{\partial c}{\partial t} = \frac{1}{\sqrt{4\pi Dt}}\cdot e^{-x^2/(4Dt)}\left[-x^2(-1)(4Dt)^{-2}(4D)\right] + e^{-x^2/(4Dt)}\cdot\left(-\tfrac{1}{2}\right)(4\pi Dt)^{-3/2}(4\pi D)$$

$$= (4\pi Dt)^{-3/2}\left(4\pi Dt\cdot\frac{x^2}{4Dt^2} - 2\pi D\right)e^{-x^2/(4Dt)} = \frac{2\pi D}{(4\pi Dt)^{3/2}}\left(\frac{x^2}{2Dt}-1\right)e^{-x^2/(4Dt)},$$

$$\frac{\partial c}{\partial x} = \frac{1}{\sqrt{4\pi Dt}}\,e^{-x^2/(4Dt)}\cdot\frac{-2x}{4Dt} = \frac{-2\pi x}{(4\pi Dt)^{3/2}}\,e^{-x^2/(4Dt)}, \text{ and}$$

$$\frac{\partial^2 c}{\partial x^2} = \frac{-2\pi}{(4\pi Dt)^{3/2}}\left[x\cdot e^{-x^2/(4Dt)}\cdot\frac{-2x}{4Dt} + e^{-x^2/(4Dt)}\cdot 1\right]$$

$$= \frac{-2\pi}{(4\pi Dt)^{3/2}}\left(-\frac{x^2}{2Dt}+1\right)e^{-x^2/(4Dt)} = \frac{2\pi}{(4\pi Dt)^{3/2}}\left(\frac{x^2}{2Dt}-1\right)e^{-x^2/(4Dt)}.$$

Thus, $\dfrac{\partial c}{\partial t} = \dfrac{2\pi D}{(4\pi Dt)^{3/2}}\left(\dfrac{x^2}{2Dt}-1\right)e^{-x^2/(4Dt)} = D\left[\dfrac{2\pi}{(4\pi Dt)^{3/2}}\left(\dfrac{x^2}{2Dt}-1\right)e^{-x^2/(4Dt)}\right] = D\,\dfrac{\partial^2 c}{\partial x^2}$.

82. (a) $T(x,y) = \dfrac{60}{1+x^2+y^2} \quad\Rightarrow\quad \dfrac{\partial T}{\partial x} = -\dfrac{60(2x)}{(1+x^2+y^2)^2}$, so at $(2,1)$, $T_x = -\dfrac{240}{(1+4+1)^2} = -\dfrac{20}{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\dfrac{\partial T}{\partial y} = -\dfrac{60(2y)}{(1+x^2+y^2)^2}$, so at $(2,1)$, $T_y = -\dfrac{120}{36} = -\dfrac{10}{3}$. Thus, from the point $(2,1)$ the temperature is decreasing at a rate of $\frac{20}{3}$°C/m in the x-direction and is decreasing at a rate of $\frac{10}{3}$°C/m in the y-direction.

83. $\dfrac{1}{R} = \dfrac{1}{R_1} + \dfrac{1}{R_2} + \dfrac{1}{R_3}$. By the Chain Rule, taking the partial derivative of both sides with respect to R_1 gives

$\dfrac{\partial R^{-1}}{\partial R}\dfrac{\partial R}{\partial R_1} = \dfrac{\partial\left[(1/R_1)+(1/R_2)+(1/R_3)\right]}{\partial R_1}$ or $-R^{-2}\dfrac{\partial R}{\partial R_1} = -R_1^{-2}$. Thus, $\dfrac{\partial R}{\partial R_1} = \dfrac{R^2}{R_1^2}$.

84. (a) $P = \dfrac{mRT}{V}$ so $\dfrac{\partial P}{\partial V} = \dfrac{-mRT}{V^2}$; $V = \dfrac{mRT}{P}$, so $\dfrac{\partial V}{\partial T} = \dfrac{mR}{P}$; $T = \dfrac{PV}{mR}$, so $\dfrac{\partial T}{\partial P} = \dfrac{V}{mR}$.

Thus $\dfrac{\partial P}{\partial V}\dfrac{\partial V}{\partial T}\dfrac{\partial T}{\partial P} = \dfrac{-mRT}{V^2}\dfrac{mR}{P}\dfrac{V}{mR} = \dfrac{-mRT}{PV} = -1$, since $PV = mRT$.

(b) By part (a), $PV = mRT \;\Rightarrow\; P = \dfrac{mRT}{V}$, so $\dfrac{\partial P}{\partial T} = \dfrac{mR}{V}$. Also, $PV = mRT \;\Rightarrow\; V = \dfrac{mRT}{P}$ and $\dfrac{\partial V}{\partial T} = \dfrac{mR}{P}$.

Since $T = \dfrac{PV}{mR}$, we have $T\dfrac{\partial P}{\partial T}\dfrac{\partial V}{\partial T} = \dfrac{PV}{mR}\cdot\dfrac{mR}{V}\cdot\dfrac{mR}{P} = mR$.

85. $\left(P + \dfrac{n^2a}{V^2}\right)(V-nb) = nRT \;\Rightarrow\; T = \dfrac{1}{nR}\left(P + \dfrac{n^2a}{V^2}\right)(V-nb)$, so $\dfrac{\partial T}{\partial P} = \dfrac{1}{nR}(1)(V-nb) = \dfrac{V-nb}{nR}$.

We can also write $P + \dfrac{n^2a}{V^2} = \dfrac{nRT}{V-nb} \;\Rightarrow\; P = \dfrac{nRT}{V-nb} - \dfrac{n^2a}{V^2} = nRT(V-nb)^{-1} - n^2aV^{-2}$, so

$\dfrac{\partial P}{\partial V} = -nRT(V-nb)^{-2}(1) + 2n^2aV^{-3} = \dfrac{2n^2a}{V^3} - \dfrac{nRT}{(V-nb)^2}$.

86. $W = 13.12 + 0.6215T - 11.37v^{0.16} + 0.3965Tv^{0.16}$. $\dfrac{\partial W}{\partial T} = 0.6215 + 0.3965v^{0.16}$. When $T = -15$°C and $v = 30$ km/h, $\dfrac{\partial W}{\partial T} = 0.6215 + 0.3965(30)^{0.16} \approx 1.3048$, so we would expect the apparent temperature to drop by approximately 1.3°C if the actual temperature decreases by 1°C. $\dfrac{\partial W}{\partial v} = -11.37(0.16)v^{-0.84} + 0.3965T(0.16)v^{-0.84}$ and when $T = -15$°C and $v = 30$ km/h, $\dfrac{\partial W}{\partial v} = -11.37(0.16)(30)^{-0.84} + 0.3965(-15)(0.16)(30)^{-0.84} \approx -0.1592$, so we would expect the apparent temperature to drop by approximately 0.16°C if the wind speed increases by 1 km/h.

87. (a) $S = f(w,h) = 0.1091w^{0.425}h^{0.725} \;\Rightarrow\; \dfrac{\partial S}{\partial w} = 0.1091(0.425)w^{0.425-1}h^{0.725} = 0.0463675w^{-0.575}h^{0.725}$, so

$\dfrac{\partial S}{\partial w}(160,70) = 0.0463675(160)^{-0.575}(70)^{0.725} \approx 0.0545$. This means that for a person 70 inches tall who weighs 160 pounds, an increase in weight (while height remains constant) causes the surface area to increase at a rate of about 0.0545 square feet (about 7.85 square inches) per pound.

(b) $\dfrac{\partial S}{\partial h} = 0.1091(0.725)w^{0.425}h^{0.725-1} = 0.0790975w^{0.425}h^{-0.275}$, so

$\dfrac{\partial S}{\partial h}(160,70) = 0.0790975(160)^{0.425}(70)^{-0.275} \approx 0.213$. This means that for a person 70 inches tall who weighs

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

160 pounds, an increase in height (while weight remains unchanged at 160 pounds) causes the surface area to increase at a rate of about 0.213 square feet (about 30.7 square inches) per inch of height.

88. $R = C\dfrac{L}{r^4} \quad \Rightarrow \quad \dfrac{\partial R}{\partial L} = \dfrac{C}{r^4}$ and $\dfrac{\partial R}{\partial r} = CL\left(-4r^{-5}\right) = -4C\dfrac{L}{r^5}$.

$\partial R/\partial L$ is the rate at which the resistance of the flowing blood increases with respect to the length of the artery when the radius stays constant. $\partial R/\partial r$ is the rate of change of the resistance with respect to the radius of the artery when the length remains unchanged. Because $\partial R/\partial r$ is negative, the resistance decreases if the radius increases.

89. $P(v, x, m) = Av^3 + \dfrac{B(mg/x)^2}{v} = Av^3 + Bm^2g^2x^{-2}v^{-1}$.

$\partial P/\partial v = 3Av^2 - \dfrac{B(mg/x)^2}{v^2}$ is the rate of change of the power needed during flapping mode with respect to the bird's velocity when the mass and fraction of flapping time remain constant. $\partial P/\partial x = -2Bm^2g^2x^{-3}v^{-1} = -\dfrac{2Bm^2g^2}{x^3v}$ is the rate at which the power changes with respect to the fraction of time spent in flapping mode when the mass and velocity are held constant. $\partial P/\partial m = 2Bmg^2x^{-2}v^{-1} = \dfrac{2Bmg^2}{x^2v}$ is the rate of change of the power with respect to mass when the velocity and fraction of flapping time remain constant.

90. $T(x, t) = T_0 + T_1e^{-\lambda x}\sin(\omega t - \lambda x)$

(a) $\partial T/\partial x \ = T_1e^{-\lambda x}\left[\cos(\omega t - \lambda x)(-\lambda)\right] + T_1(-\lambda e^{-\lambda x})\sin(\omega t - \lambda x) = -\lambda T_1e^{-\lambda x}\left[\sin(\omega t - \lambda x) + \cos(\omega t - \lambda x)\right]$.

This quantity represents the rate of change of temperature with respect to depth below the surface, at a given time t.

(b) $\partial T/\partial t = T_1e^{-\lambda x}\left[\cos(\omega t - \lambda x)(\omega)\right] = \omega T_1e^{-\lambda x}\cos(\omega t - \lambda x)$. This quantity represents the rate of change of temperature with respect to time at a fixed depth x.

(c) $T_{xx} = \dfrac{\partial}{\partial x}\left(\dfrac{\partial T}{\partial x}\right)$

$$= -\lambda T_1\left(e^{-\lambda x}\left[\cos(\omega t - \lambda x)(-\lambda) - \sin(\omega t - \lambda x)(-\lambda)\right] + \ e^{-\lambda x}(-\lambda)\left[\sin(\omega t - \lambda x) + \cos(\omega t - \lambda x)\right]\right)$$
$$= 2\lambda^2T_1e^{-\lambda x}\cos(\omega t - \lambda x)$$

But from part (b), $T_t = \omega T_1e^{-\lambda x}\cos(\omega t - \lambda x) = \dfrac{\omega}{2\lambda^2}T_{xx}$. So with $k = \dfrac{\omega}{2\lambda^2}$, the function T satisfies the heat equation.

(d)

Note that near the surface (that is, for small x) the temperature varies greatly as t changes, but deeper (for large x) the temperature is more stable.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(e) The term $-\lambda x$ is a phase shift: it represents the fact that since heat diffuses slowly through soil, it takes time for changes in the surface temperature to affect the temperature at deeper points. As x increases, the phase shift also increases. For example, when $\lambda = 0.2$, the highest temperature at the surface is reached when $t \approx 91$, whereas at a depth of 5 feet the peak temperature is attained at $t \approx 149$, and at a depth of 10 feet, at $t \approx 207$.

91. $\dfrac{\partial K}{\partial m} = \frac{1}{2}v^2,\ \dfrac{\partial K}{\partial v} = mv,\ \dfrac{\partial^2 K}{\partial v^2} = m.$ Thus $\dfrac{\partial K}{\partial m} \cdot \dfrac{\partial^2 K}{\partial v^2} = \frac{1}{2}v^2 m = K.$

92. $E(m,v) = 2.65m^{0.66} + \dfrac{3.5m^{0.75}}{v} \quad \Rightarrow$

$E_m(m,v) = 2.65(0.66)m^{0.66-1} + \dfrac{3.5(0.75)m^{0.75-1}}{v} = 1.749m^{-0.34} + \dfrac{2.625m^{-0.25}}{v},$

$E_v(m,v) = 3.5m^{0.75}\left(-v^{-2}\right) = -\dfrac{3.5m^{0.75}}{v^2}$. Then $E_m(400,8) = 1.749(400)^{-0.34} + \dfrac{2.625(400)^{-0.25}}{8} \approx 0.301$ which means that the average energy needed for a lizard to walk or run 1 km increases at a rate of about 0.301 kcal per gram of body mass increase from 400 g if the speed is 8 km/h. $E_v(400,8) = -\dfrac{3.5(400)^{0.75}}{8^2} \approx -4.89$, which means that the average energy needed by a lizard with body mass 400 g decreases at a rate of about 4.89 kcal per km/h when the speed increases from 8 km/h.

93. By the geometry of partial derivatives, the slope of the tangent line is $f_x(1,2)$. By implicit differentiation of $4x^2 + 2y^2 + z^2 = 16$, we get $8x + 2z\,(\partial z/\partial x) = 0 \quad \Rightarrow \quad \partial z/\partial x = -4x/z$, so when $x = 1$ and $z = 2$ we have $\partial z/\partial x = -2$. So the slope is $f_x(1,2) = -2$. Thus the tangent line is given by $z - 2 = -2(x-1),\ y = 2$. Taking the parameter to be $t = x - 1$, we can write parametric equations for this line: $x = 1 + t,\ y = 2,\ z = 2 - 2t$.

94. $z = 6 - x - x^2 - 2y^2$. Setting $x = 1$, the equation of the parabola of intersection is $z = 6 - 1 - 1 - 2y^2 = 4 - 2y^2$. The slope of the tangent is $\partial z/\partial y = -4y$, so at $(1,2,-4)$ the slope is -8. Parametric equations for the line are therefore $x = 1,\ y = 2 + t,\ z = -4 - 8t$.

95. $f_x(x,y) = x + 4y \quad \Rightarrow \quad f_{xy}(x,y) = 4$ and $f_y(x,y) = 3x - y \quad \Rightarrow \quad f_{yx}(x,y) = 3$. Since f_{xy} and f_{yx} are continuous everywhere but $f_{xy}(x,y) \neq f_{yx}(x,y)$, Clairaut's Theorem implies that such a function $f(x,y)$ does not exist.

96. The Law of Cosines says that $a^2 = b^2 + c^2 - 2bc\cos A$. Thus $\dfrac{\partial(a^2)}{\partial a} = \dfrac{\partial(b^2 + c^2 - 2ab\cos A)}{\partial a}$ or $2a = -2bc\,(-\sin A)\,\dfrac{\partial A}{\partial a}$, implying that $\dfrac{\partial A}{\partial a} = \dfrac{a}{bc\sin A}$. Taking the partial derivative of both sides with respect to b gives $0 = 2b - 2c(\cos A) - 2bc\,(-\sin A)\,\dfrac{\partial A}{\partial b}$. Thus $\dfrac{\partial A}{\partial b} = \dfrac{c\cos A - b}{bc\sin A}$. By symmetry, $\dfrac{\partial A}{\partial c} = \dfrac{b\cos A - c}{bc\sin A}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

97. By Clairaut's Theorem, $f_{xyy} = (f_{xy})_y = (f_{yx})_y = f_{yxy} = (f_y)_{xy} = (f_y)_{yx} = f_{yyx}$.

98. (a) Since we are differentiating n times, with two choices of variable at each differentiation, there are 2^n nth-order partial derivatives.

(b) If these partial derivatives are all continuous, then the order in which the partials are taken doesn't affect the value of the result, that is, all nth-order partial derivatives with p partials with respect to x and $n-p$ partials with respect to y are equal. Since the number of partials taken with respect to x for an nth-order partial derivative can range from 0 to n, a function of two variables has $n+1$ distinct partial derivatives of order n if these partial derivatives are all continuous.

(c) Since n differentiations are to be performed with three choices of variable at each differentiation, there are 3^n nth-order partial derivatives of a function of three variables.

99. $f(x,y) = x(x^2+y^2)^{-3/2}e^{\sin(x^2y)}$. Let $g(x) = f(x,0) = x(x^2)^{-3/2}e^0 = x\,|x|^{-3}$. To find $f_x(1,0)$, we are using the point $(1,0)$, so near $(1,0)$, $g(x) = x^{-2}$. Then $g'(x) = -2x^{-3}$ and $g'(1) = -2$, so using Equation 1, we have $f_x(1,0) = g'(1) = -2$.

100. $f(x,y) = \sqrt[3]{x^3+y^3}$. $f_x(0,0) = \lim\limits_{h\to 0} \dfrac{f(0+h,0)-f(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{(h^3+0)^{1/3}-0}{h} = \lim\limits_{h\to 0} \dfrac{h}{h} = 1$.

Or: Let $g(x) = f(x,0) = \sqrt[3]{x^3+0} = x$. Then $g'(x) = 1$ and $g'(0) = 1$ so, by (1), $f_x(0,0) = g'(0) = 1$.

101. (a) $f(x,y) = \begin{cases} \dfrac{x^3y - xy^3}{x^2+y^2} & \text{if } (x,y) \neq (0,0) \\ 0 & \text{if } (x,y) = (0,0) \end{cases}$

(b) For $(x,y) \neq (0,0)$,

$$f_x(x,y) = \frac{(3x^2y - y^3)(x^2+y^2) - (x^3y - xy^3)(2x)}{(x^2+y^2)^2}$$

$$= \frac{x^4y + 4x^2y^3 - y^5}{(x^2+y^2)^2}$$

and, by symmetry, $f_y(x,y) = \dfrac{x^5 - 4x^3y^2 - xy^4}{(x^2+y^2)^2}$.

(c) $f_x(0,0) = \lim\limits_{h\to 0} \dfrac{f(h,0)-f(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{(0/h^2)-0}{h} = 0$ and $f_y(0,0) = \lim\limits_{h\to 0} \dfrac{f(0,h)-f(0,0)}{h} = 0$.

(d) By (3), $f_{xy}(0,0) = \dfrac{\partial f_x}{\partial y} = \lim\limits_{h\to 0} \dfrac{f_x(0,h)-f_x(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{(-h^5-0)/h^4}{h} = -1$ while by (2),

$$f_{yx}(0,0) = \frac{\partial f_y}{\partial x} = \lim_{h\to 0} \frac{f_y(h,0)-f_y(0,0)}{h} = \lim_{h\to 0} \frac{h^5/h^4}{h} = 1.$$

(e) For $(x,y) \neq (0,0)$, we use a CAS to compute

$$f_{xy}(x,y) = \frac{x^6 + 9x^4y^2 - 9x^2y^4 - y^6}{(x^2+y^2)^3}$$

Now as $(x,y) \to (0,0)$ along the x-axis, $f_{xy}(x,y) \to 1$ while as $(x,y) \to (0,0)$ along the y-axis, $f_{xy}(x,y) \to -1$. Thus f_{xy} isn't

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

continuous at $(0,0)$ and Clairaut's Theorem doesn't apply, so there is no contradiction. The graphs of f_{xy} and f_{yx} are identical except at the origin, where we observe the discontinuity.

DISCOVERY PROJECT Deriving the Cobb-Douglas Production Function

1. For $K = K_0$, we have $\dfrac{dP}{dL} = \alpha \dfrac{P}{L} \quad \Rightarrow \quad \dfrac{dP}{P} = \alpha \dfrac{dL}{L} \quad \Rightarrow \quad \displaystyle\int \frac{dP}{P} = \int \alpha \frac{dL}{L} \quad \Rightarrow \quad \ln|P| = \alpha \ln|L| + A_1(K_0)$.

Then $e^{\ln|P|} = e^{\ln|L|^\alpha + A_1(K_0)} \quad \Rightarrow \quad P = e^{A_1(K_0)} L^\alpha \quad \Rightarrow \quad P(L, K_0) = C_1(K_0) L^\alpha$, where $e^{A_1(K_0)} = C_1(K_0)$.

2. For $L = L_0$, we have $\dfrac{dP}{dK} = \beta \dfrac{P}{K} \quad \Rightarrow \quad \dfrac{dP}{P} = \beta \dfrac{dK}{K} \quad \Rightarrow \quad \displaystyle\int \frac{dP}{P} = \int \beta \frac{dK}{K} \quad \Rightarrow \quad \ln|P| = \beta \ln|K| + A_2(L_0)$.

Then $e^{\ln|P|} = e^{\ln|K|^\beta + A_2(L_0)} \quad \Rightarrow \quad P = e^{A_2(L_0)} K^\beta \quad \Rightarrow \quad P(L_0, K) = C_2(L_0) K^\beta$, where $e^{A_2(L_0)} = C_2(L_0)$.

3. Suppose both labor and capital are increased by a factor of m. Then

$$\begin{aligned} P(mL, mk) &= b(mL)^\alpha (mK)^{1-\alpha} = bm^\alpha L^\alpha m^{1-\alpha} K^{1-\alpha} = m^\alpha m^{1-\alpha} b L^\alpha K^{1-\alpha} \\ &= m(bL^\alpha K^{1-\alpha}) = mP(L, K) \end{aligned}$$

Therefore, production is also increased by a factor of m.

4. For $P(L, K) = bL^\alpha K^{1-\alpha}$,

$$\begin{aligned} L\frac{\partial P}{\partial L} + K\frac{\partial P}{\partial K} &= L(b\alpha L^{\alpha-1} K^{1-\alpha}) + K(bL^\alpha (1-\alpha) K^{1-\alpha-1}) \\ &= b\alpha L^\alpha K^{1-\alpha} + b(1-\alpha) L^\alpha K^{1-\alpha} = bL^\alpha K^{1-\alpha}[\alpha + (1-\alpha)] \\ &= bL^\alpha K^{1-\alpha} = P(L, K) \end{aligned}$$

5. $P(L, K) = 1.01 L^{0.75} K^{0.25}$. Marginal productivity of labor is given by $\partial P/\partial L = 1.01(0.75) L^{-0.25} K^{0.25}$. With $L = 194$ and $K = 407$, we have $\partial P/\partial L \approx 0.911656$. When capital is held constant at $K = 407$, as labor increases from $L = 194$, production will increase 0.911656 per unit of labor.

Marginal productivity of capital is given by $\partial P/\partial K = 1.01(0.25) L^{0.75} K^{-0.75}$. With $L = 194$ and $K = 407$, we have $\partial P/\partial K \approx 0.144849$. When labor is held constant at $L = 194$, as capital increases from $K = 407$, production will increase 0.144849 per unit of capital.

As the marginal productivity of labor is higher, it would be more beneficial to increase labor.

14.4 Tangent Planes and Linear Approximations

1. $z = f(x, y) = 16 - x^2 - y^2 \quad \Rightarrow \quad f_x(x, y) = -2x, f_y(x, y) = -2y$, so $f_x(2, 2) = -4$ and $f_y(2, 2) = -4$. By Equation 2, an equation of the tangent plane is $z - 8 = f_x(2, 2)(x - 2) + f_y(2, 2)(y - 2) \quad \Rightarrow \quad z - 8 = -4(x - 2) - 4(y - 2)$, or $z = -4x - 4y + 24$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

2. $z = f(x, y) = y^2 \sin x \quad \Rightarrow \quad f_x(x, y) = y^2 \cos x$, $f_y(x, y) = 2y \sin x$, so $f_x(\pi/2, -2) = 0$ and $f_y(\pi/2, -2) = -4$. By Equation 2, an equation of the tangent plane is $z - 4 = f_x(\pi/2, -2)(x - \pi/2) + f_y(\pi/2, -2)(y - (-2)) \quad \Rightarrow$ $z - 4 = -4(y + 2)$, or $z = -4y - 4$.

3. $z = f(x, y) = 2x^2 + y^2 - 5y \quad \Rightarrow \quad f_x(x, y) = 4x,\ f_y(x, y) = 2y - 5,$ so $f_x(1, 2) = 4$ and $f_y(1, 2) = -1$. By Equation 2, an equation of the tangent plane is $z - (-4) = f_x(1, 2)(x - 1) + f_y(1, 2)(y - 2) \quad \Rightarrow$ $z + 4 = 4(x - 1) + (-1)(y - 2)$, or $z = 4x - y - 6$.

4. $z = f(x, y) = (x + 2)^2 - 2(y - 1)^2 - 5 \quad \Rightarrow \quad f_x(x, y) = 2(x + 2),\ \ f_y(x, y) = -4(y - 1),$ so $f_x(2, 3) = 8$ and $f_y(2, 3) = -8$. By Equation 2, an equation of the tangent plane is $z - 3 = f_x(2, 3)(x - 2) + f_y(2, 3)(y - 3) \quad \Rightarrow$ $z - 3 = 8(x - 2) + (-8)(y - 3)$, or $z = 8x - 8y + 11$.

5. $z = f(x, y) = e^{x-y} \quad \Rightarrow \quad f_x(x, y) = e^{x-y}(1) = e^{x-y},\ \ f_y(x, y) = e^{x-y}(-1) = -e^{x-y},$ so $f_x(2, 2) = 1$ and $f_y(2, 2) = -1$. Thus, an equation of the tangent plane is $z - 1 = f_x(2, 2)(x - 2) + f_y(2, 2)(y - 2) \quad \Rightarrow$ $z - 1 = 1(x - 2) + (-1)(y - 2)$, or $z = x - y + 1$.

6. $z = f(x, y) = y^2 e^x \quad \Rightarrow \quad f_x(x, y) = y^2 e^x$, $f_y(x, y) = 2ye^x$, so $f_x(0, 3) = 9$ and $f_y(0, 3) = 6$. Thus, an equation of the tangent plane is $z - 9 = f_x(0, 3)(x - 0) + f_y(0, 3)(y - 3) \quad \Rightarrow \quad z - 9 = 9x + 6(y - 3)$, or $z = 9x + 6y - 9$.

7. $z = f(x, y) = \dfrac{2\sqrt{y}}{x} \quad \Rightarrow \quad f_x(x, y) = -\dfrac{2\sqrt{y}}{x^2}, f_y(x, y) = \dfrac{1}{x\sqrt{y}}$, so $f_x(-1, 1) = -2$ and $f_y(-1, 1) = -1$. Thus, an equation of the tangent plane is $z - (-2) = f_x(-1, 1)(x - (-1)) + f_y(-1, 1)(y - 1) \quad \Rightarrow$ $z + 2 = -2(x + 1) - 1(y - 1)$, or $z = -2x - y - 3$.

8. $z = f(x, y) = x/y^2 = xy^{-2} \quad \Rightarrow \quad f_x(x, y) = 1/y^2,\ \ f_y(x, y) = -2xy^{-3} = -2x/y^3,$ so $f_x(-4, 2) = \frac{1}{4}$ and $f_y(-4, 2) = 1$. Thus, an equation of the tangent plane is $z - (-1) = f_x(-4, 2)\,[x - (-4)] + f_y(-4, 2)(y - 2) \quad \Rightarrow$ $z + 1 = \frac{1}{4}(x + 4) + 1(y - 2)$, or $z = \frac{1}{4}x + y - 2$.

9. $z = f(x, y) = x \sin(x + y) \quad \Rightarrow \quad f_x(x, y) = x \cdot \cos(x + y) \cdot 1 + \sin(x + y) \cdot 1 = x \cos(x + y) + \sin(x + y)$ and $f_y(x, y) = x \cos(x + y) \cdot 1$, so $f_x(-1, 1) = (-1) \cos 0 + \sin 0 = -1,\ \ f_y(-1, 1) = (-1) \cos 0 = -1$. Thus, an equation of the tangent plane is $z - 0 = f_x(-1, 1)(x - (-1)) + f_y(-1, 1)(y - 1) \quad \Rightarrow \quad z = (-1)(x + 1) + (-1)(y - 1)$, or $x + y + z = 0$.

10. $z = f(x, y) = \ln(x - 2y) \quad \Rightarrow \quad f_x(x, y) = 1/(x - 2y),\ f_y(x, y) = -2/(x - 2y),$ so $f_x(3, 1) = 1$ and $f_y(3, 1) = -2$. Thus, an equation of the tangent plane is $z - 0 = f_x(3, 1)(x - 3) + f_y(3, 1)(y - 1) \quad \Rightarrow \quad z = 1(x - 3) + (-2)(y - 1)$, or $z = x - 2y - 1$.

11. $z = f(x, y) = x^2 + xy + 3y^2$, so $f_x(x, y) = 2x + y \quad \Rightarrow \quad f_x(1, 1) = 3,\ \ f_y(x, y) = x + 6y \quad \Rightarrow \quad f_y(1, 1) = 7$ and an equation of the tangent plane is $z - 5 = 3(x - 1) + 7(y - 1),$ or $z = 3x + 7y - 5$. After zooming in, the surface and the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

tangent plane become almost indistinguishable. (Here, the tangent plane is below the surface.) If we zoom in farther, the surface and the tangent plane will appear to coincide.

12. $z = f(x,y) = \sqrt{9 + x^2y^2} \quad \Rightarrow \quad f_x(x,y) = \frac{1}{2}\left(9 + x^2y^2\right)^{-1/2}(2xy^2) = xy^2/\sqrt{9 + x^2y^2}$,

$f_y(x,y) = \frac{1}{2}\left(9 + x^2y^2\right)^{-1/2}(2x^2y) = x^2y/\sqrt{9 + x^2y^2}$, so $f_x(2,2) = \frac{8}{5}$ and $f_y(2,2) = \frac{8}{5}$. Thus an equation of the tangent plane is $z - 5 = f_x(2,2)(x-2) + f_y(2,2)(y-2) \quad \Rightarrow \quad z - 5 = \frac{8}{5}(x-2) + \frac{8}{5}(y-2)$ or $z = \frac{8}{5}x + \frac{8}{5}y - \frac{7}{5}$. After zooming in, the surface and the tangent plane become almost indistinguishable. (Here the tangent plane is shown with fewer traces than the surface.) If we zoom in farther, the surface and the tangent plane will appear to coincide.

13. $f(x,y) = \dfrac{1 + \cos^2(x-y)}{1 + \cos^2(x+y)}$. A CAS gives

$$f_x(x,y) = -\frac{2\cos(x-y)\sin(x-y)}{1 + \cos^2(x+y)} + \frac{2\left[1 + \cos^2(x-y)\right]\cos(x+y)\sin(x+y)}{\left[1 + \cos^2(x+y)\right]^2} \quad \text{and}$$

$$f_y(x,y) = \frac{2\cos(x-y)\sin(x-y)}{1 + \cos^2(x+y)} + \frac{2\left[1 + \cos^2(x-y)\right]\cos(x+y)\sin(x+y)}{\left[1 + \cos^2(x+y)\right]^2}.$$ We use the CAS to evaluate these at $(\pi/3, \pi/6)$, giving $f_x(\pi/3, \pi/6) = -\sqrt{3}/2$ and $f_y(\pi/3, \pi/6) = \sqrt{3}/2$. Substituting into Equation 2, an equation of the tangent plane is $z = -\frac{\sqrt{3}}{2}\left(x - \frac{\pi}{3}\right) + \frac{\sqrt{3}}{2}\left(y - \frac{\pi}{6}\right) + \frac{7}{4}$. The surface and tangent plane are shown in the first graph.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

After zooming in, the surface and the tangent plane become almost indistinguishable, as shown in the second graph. (Here, the tangent plane is above the surface.) If we zoom in farther, the surface and the tangent plane will appear to coincide.

14. $f(x,y) = e^{-xy/10}\left(\sqrt{x}+\sqrt{y}+\sqrt{xy}\right)$. A CAS gives

$f_x(x,y) = -\frac{1}{10}ye^{-xy/10}\left(\sqrt{x}+\sqrt{y}+\sqrt{xy}\right) + e^{-xy/10}\left(\frac{1}{2\sqrt{x}}+\frac{y}{2\sqrt{xy}}\right)$ and

$f_y(x,y) = -\frac{1}{10}xe^{-xy/10}\left(\sqrt{x}+\sqrt{y}+\sqrt{xy}\right) + e^{-xy/10}\left(\frac{1}{2\sqrt{y}}+\frac{x}{2\sqrt{xy}}\right)$. We use the CAS to evaluate these at $(1,1)$, and then substitute the results into Equation 2 to get an equation of the tangent plane:

$z - 3e^{-0.1} = 0.7e^{-0.1}(x-1) + 0.7e^{-0.1}(y-1) \quad\Rightarrow\quad z = 0.7e^{-0.1}x + 0.7e^{-0.1}y + 1.6e^{-0.1}$. The surface and tangent plane are shown in the first graph below. After zooming in, the surface and the tangent plane become almost indistinguishable, as shown in the second graph. (Here, the tangent plane is above the surface.) If we zoom in farther, the surface and the tangent plane will appear to coincide.

15. $f(x,y) = x^3y^2$. The partial derivatives are $f_x(x,y) = 3x^2y^2$ and $f_y(x,y) = 2x^3y$, so $f_x(-2,1) = 12$ and $f_y(-2,1) = -16$. Both f_x and f_y are continuous functions, so by Theorem 8, f is differentiable at $(-2,1)$. By Equation 3, the linearization of f at $(-2,1)$ is given by

$L(x,y) = f(-2,1) + f_x(-2,1)(x-(-2)) + f_y(-2,1)(y-1) = -8 + 12(x+2) - 16(y-1) = 12x - 16y + 32.$

16. $f(x,y) = y\tan x$. The partial derivatives are $f_x(x,y) = y\sec^2 x$ and $f_y = \tan x$, so $f_x\left(\frac{\pi}{4},2\right) = 4$ and $f_y\left(\frac{\pi}{4},2\right) = 1$. Both f_x and f_y are continuous for $x \neq \frac{\pi}{2} + n\pi$, so by Theorem 8, f is differentiable at $\left(\frac{\pi}{4},2\right)$. By Equation 3, the linearization of f at $\left(\frac{\pi}{4},2\right)$ is given by $L(x,y) = f\left(\frac{\pi}{4},2\right) + f_x\left(\frac{\pi}{4},2\right)\left(x-\frac{\pi}{4}\right) + f_y\left(\frac{\pi}{4},2\right)(y-2) = 2 + 4\left(x-\frac{\pi}{4}\right) + 1(y-2) = 4x + y - \pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $f(x,y) = 1 + x\ln(xy-5)$. The partial derivatives are $f_x(x,y) = x \cdot \dfrac{1}{xy-5}\,(y) + \ln(xy-5)\cdot 1 = \dfrac{xy}{xy-5} + \ln(xy-5)$ and $f_y(x,y) = x \cdot \dfrac{1}{xy-5}\,(x) = \dfrac{x^2}{xy-5}$, so $f_x(2,3) = 6$ and $f_y(2,3) = 4$. Both f_x and f_y are continuous functions for $xy > 5$, so by Theorem 8, f is differentiable at $(2,3)$. By Equation 3, the linearization of f at $(2,3)$ is given by $L(x,y) = f(2,3) + f_x(2,3)(x-2) + f_y(2,3)(y-3) = 1 + 6(x-2) + 4(y-3) = 6x + 4y - 23$.

18. $f(x,y) = \sqrt{xy} = (xy)^{1/2}$. The partial derivatives are $f_x(x,y) = \frac{1}{2}(xy)^{-1/2}(y) = y/\left(2\sqrt{xy}\right)$ and $f_y(x,y) = \frac{1}{2}(xy)^{-1/2}(x) = x/\left(2\sqrt{xy}\right)$, so $f_x(1,4) = 4/\left(2\sqrt{4}\right) = 1$ and $f_y(1,4) = 1/\left(2\sqrt{4}\right) = \frac{1}{4}$. Both f_x and f_y are continuous functions for $xy > 0$, so f is differentiable at $(1,4)$ by Theorem 8. By Equation 3, the linearization of f at $(1,4)$ is given by $L(x,y) = f(1,4) + f_x(1,4)(x-1) + f_y(1,4)(y-4) = 2 + 1(x-1) + \frac{1}{4}(y-4) = x + \frac{1}{4}y$.

19. $f(x,y) = x^2e^y$. The partial derivatives are $f_x(x,y) = 2xe^y$ and $f_y(x,y) = x^2e^y$, so $f_x(1,0) = 2$ and $f_y(1,0) = 1$. Both f_x and f_y are continuous functions, so by Theorem 8, f is differentiable at $(1,0)$. By Equation 3, the linearization of f at $(1,0)$ is given by $L(x,y) = f(1,0) + f_x(1,0)(x-1) + f_y(1,0)(y-0) = 1 + 2(x-1) + 1(y-0) = 2x + y - 1$.

20. $f(x,y) = \dfrac{1+y}{1+x} = (1+y)(1+x)^{-1}$. The partial derivatives are $f_x(x,y) = (1+y)(-1)(1+x)^{-2} = -\dfrac{1+y}{(1+x)^2}$ and $f_y(x,y) = (1)(1+x)^{-1} = \dfrac{1}{1+x}$, so $f_x(1,3) = -1$ and $f_y(1,3) = \frac{1}{2}$. Both f_x and f_y are continuous functions for $x \neq -1$, so by Theorem 8, f is differentiable at $(1,3)$. By Equation 3, the linearization of f at $(1,3)$ is given by $L(x,y) = f(1,3) + f_x(1,3)(x-1) + f_y(1,3)(y-3) = 2 + (-1)(x-1) + \frac{1}{2}(y-3) = -x + \frac{1}{2}y + \frac{3}{2}$.

21. $f(x,y) = 4\arctan(xy)$. The partial derivatives are $f_x(x,y) = 4 \cdot \dfrac{1}{1+(xy)^2}(y) = \dfrac{4y}{1+x^2y^2}$, and $f_y(x,y) = \dfrac{4x}{1+x^2y^2}$, so $f_x(1,1) = 2$ and $f_y(1,1) = 2$. Both f_x and f_y are continuous functions, so by Theorem 8, f is differentiable at $(1,1)$. By Equation 3, the linearization of f at $(1,1)$ is given by $L(x,y) = f(1,1) + f_x(1,1)(x-1) + f_y(1,1)(y-1) = 4(\pi/4) + 2(x-1) + 2(y-1) = 2x + 2y + \pi - 4$.

22. $f(x,y) = y + \sin(x/y)$. The partial derivatives are $f_x(x,y) = (1/y)\cos(x/y)$ and $f_y(x,y) = 1 + (-x/y^2)\cos(x/y)$, so $f_x(0,3) = \frac{1}{3}$ and $f_y(0,3) = 1$. Both f_x and f_y are continuous functions for $y \neq 0$, so by Theorem 8, f is differentiable at $(0,3)$. By Equation 3, the linearization of f at $(0,3)$ is given by $L(x,y) = f(0,3) + f_x(0,3)(x-0) + f_y(0,3)(y-3) = 3 + \frac{1}{3}(x-0) + 1(y-3) = \frac{1}{3}x + y$.

23. Let $f(x,y) = e^x\cos(xy)$. Then $f_x(x,y) = e^x[-\sin(xy)](y) + e^x\cos(xy) = e^x[\cos(xy) - y\sin(xy)]$ and $f_y(x,y) = e^x[-\sin(xy)](x) = -xe^x\sin(xy)$. Both f_x and f_y are continuous functions, so by Theorem 8, f is differentiable at $(0,0)$. We have $f_x(0,0) = e^0(\cos 0 - 0) = 1$, $f_y(0,0) = 0$ and the linear approximation of f at $(0,0)$ is $f(x,y) \approx f(0,0) + f_x(0,0)(x-0) + f_y(0,0)(y-0) = 1 + 1x + 0y = x + 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. Let $f(x,y) = \dfrac{y-1}{x+1}$. Then $f_x(x,y) = (y-1)(-1)(x+1)^{-2} = \dfrac{1-y}{(x+1)^2}$ and $f_y(x,y) = \dfrac{1}{x+1}$. Both f_x and f_y are continuous functions for $x \neq -1$, so by Theorem 8, f is differentiable at $(0,0)$. We have $f_x(0,0) = 1$, $f_y(0,0) = 1$ and the linear approximation of f at $(0,0)$ is $f(x,y) \approx f(0,0) + f_x(0,0)(x-0) + f_y(0,0)(y-0) = -1 + 1x + 1y = x + y - 1$.

25. We can estimate $f(2.2, 4.9)$ using a linear approximation of f at $(2,5)$, given by $f(x,y) \approx f(2,5) + f_x(2,5)(x-2) + f_y(2,5)(y-5) = 6 + 1(x-2) + (-1)(y-5) = x - y + 9$. Thus $f(2.2, 4.9) \approx 2.2 - 4.9 + 9 = 6.3$.

26. $f(x,y) = 1 - xy\cos\pi y \;\Rightarrow\; f_x(x,y) = -y\cos\pi y$ and $f_y(x,y) = -x[y(-\pi\sin\pi y) + (\cos\pi y)(1)] = \pi xy\sin\pi y - x\cos\pi y$, so $f_x(1,1) = 1$, $f_y(1,1) = 1$. Then the linear approximation of f at $(1,1)$ is given by

$$\begin{aligned} f(x,y) &\approx f(1,1) + f_x(1,1)(x-1) + f_y(1,1)(y-1) \\ &= 2 + (1)(x-1) + (1)(y-1) = x + y \end{aligned}$$

Thus $f(1.02, 0.97) \approx 1.02 + 0.97 = 1.99$. We graph f and its tangent plane near the point $(1,1,2)$ below. Notice near $y = 1$ the surfaces are almost identical.

27. $f(x,y,z) = \sqrt{x^2+y^2+z^2} \;\Rightarrow\; f_x(x,y,z) = \dfrac{x}{\sqrt{x^2+y^2+z^2}}$, $f_y(x,y,z) = \dfrac{y}{\sqrt{x^2+y^2+z^2}}$, and $f_z(x,y,z) = \dfrac{z}{\sqrt{x^2+y^2+z^2}}$, so $f_x(3,2,6) = \frac{3}{7}$, $f_y(3,2,6) = \frac{2}{7}$, $f_z(3,2,6) = \frac{6}{7}$. Then the linear approximation of f at $(3,2,6)$ is given by

$$\begin{aligned} f(x,y,z) &\approx f(3,2,6) + f_x(3,2,6)(x-3) + f_y(3,2,6)(y-2) + f_z(3,2,6)(z-6) \\ &= 7 + \tfrac{3}{7}(x-3) + \tfrac{2}{7}(y-2) + \tfrac{6}{7}(z-6) = \tfrac{3}{7}x + \tfrac{2}{7}y + \tfrac{6}{7}z \end{aligned}$$

Thus $\sqrt{(3.02)^2 + (1.97)^2 + (5.99)^2} = f(3.02, 1.97, 5.99) \approx \frac{3}{7}(3.02) + \frac{2}{7}(1.97) + \frac{6}{7}(5.99) \approx 6.9914$.

28. From the table, $f(40,20) = 28$. To estimate $f_v(40,20)$ and $f_t(40,20)$ we follow the procedure used in Exercise 14.3.2. Since $f_v(40,20) = \lim\limits_{h\to 0} \dfrac{f(40+h,20) - f(40,20)}{h}$, we approximate this quantity with $h = \pm 10$ and use the values given in the table:

$$f_v(40,20) \approx \frac{f(50,20) - f(40,20)}{10} = \frac{40-28}{10} = 1.2, \qquad f_v(40,20) \approx \frac{f(30,20) - f(40,20)}{-10} = \frac{17-28}{-10} = 1.1$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Averaging these values gives $f_v(40,20) \approx 1.15$. Similarly, $f_t(40,20) = \lim_{h\to 0} \dfrac{f(40,20+h)-f(40,20)}{h}$, so we use $h = 10$ and $h = -5$:

$$f_t(40,20) \approx \frac{f(40,30)-f(40,20)}{10} = \frac{31-28}{10} = 0.3, \qquad f_t(40,20) \approx \frac{f(40,15)-f(40,20)}{-5} = \frac{25-28}{-5} = 0.6$$

Averaging these values gives $f_t(40,15) \approx 0.45$. The linear approximation, then, is

$$f(v,t) \approx f(40,20) + f_v(40,20)(v-40) + f_t(40,20)(t-20) \approx 28 + 1.15(v-40) + 0.45(t-20)$$

When $v = 43$ and $t = 24$, we estimate $f(43,24) \approx 28 + 1.15(43-40) + 0.45(24-20) = 33.25$, so we would expect the wave heights to be approximately 33.25 ft.

29. From the table, $f(94,80) = 127$. To estimate $f_T(94,80)$ and $f_H(94,80)$ we follow the procedure used in Section 14.3. Since $f_T(94,80) = \lim_{h\to 0} \dfrac{f(94+h,80)-f(94,80)}{h}$, we approximate this quantity with $h = \pm 2$ and use the values given in the table:

$$f_T(94,80) \approx \frac{f(96,80)-f(94,80)}{2} = \frac{135-127}{2} = 4, \qquad f_T(94,80) \approx \frac{f(92,80)-f(94,80)}{-2} = \frac{119-127}{-2} = 4$$

Averaging these values gives $f_T(94,80) \approx 4$. Similarly, $f_H(94,80) = \lim_{h\to 0} \dfrac{f(94,80+h)-f(94,80)}{h}$, so we use $h = \pm 5$:

$$f_H(94,80) \approx \frac{f(94,85)-f(94,80)}{5} = \frac{132-127}{5} = 1, \qquad f_H(94,80) \approx \frac{f(94,75)-f(94,80)}{-5} = \frac{122-127}{-5} = 1$$

Averaging these values gives $f_H(94,80) \approx 1$. The linear approximation, then, is

$$\begin{aligned} f(T,H) &\approx f(94,80) + f_T(94,80)(T-94) + f_H(94,80)(H-80) \\ &\approx 127 + 4(T-94) + 1(H-80) \qquad [\text{or } 4T + H - 329] \end{aligned}$$

Thus when $T = 95$ and $H = 78$, $f(95,78) \approx 127 + 4(95-94) + 1(78-80) = 129$, so we estimate the heat index to be approximately $129°\text{F}$.

30. From the table, $f(-15,50) = -29$. To estimate $f_T(-15,50)$ and $f_v(-15,50)$ we follow the procedure used in Section 14.3. Since $f_T(-15,50) = \lim_{h\to 0} \dfrac{f(-15+h,50)-f(-15,50)}{h}$, we approximate this quantity with $h = \pm 5$ and use the values given in the table:

$$f_T(-15,50) \approx \frac{f(-10,50)-f(-15,50)}{5} = \frac{-22-(-29)}{5} = 1.4$$

$$f_T(-15,50) \approx \frac{f(-20,50)-f(-15,50)}{-5} = \frac{-35-(-29)}{-5} = 1.2$$

Averaging these values gives $f_T(-15,50) \approx 1.3$. Similarly $f_v(-15,50) = \lim_{h\to 0} \dfrac{f(-15,50+h)-f(-15,50)}{h}$,

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

so we use $h = \pm 10$:

$$f_v(-15, 50) \approx \frac{f(-15, 60) - f(-15, 50)}{10} = \frac{-30 - (-29)}{10} = -0.1$$

$$f_v(-15, 50) \approx \frac{f(-15, 40) - f(-15, 50)}{-10} = \frac{-27 - (-29)}{-10} = -0.2$$

Averaging these values gives $f_v(-15, 50) \approx -0.15$. The linear approximation to the wind-chill index function, then, is $f(T, v) \approx f(-15, 50) + f_T(-15, 50)(T - (-15)) + f_v(-15, 50)(v - 50) \approx -29 + (1.3)(T + 15) - (0.15)(v - 50)$. Thus when $T = -17°\text{C}$ and $v = 55$ km/h, $f(-17, 55) \approx -29 + (1.3)(-17 + 15) - (0.15)(55 - 50) = -32.35$, so we estimate the wind-chill index to be approximately $-32.35°\text{C}$.

31. $m = p^5 q^3 \quad \Rightarrow \quad dm = \dfrac{\partial m}{\partial p}\, dp + \dfrac{\partial m}{\partial q}\, dq = 5p^4 q^3\, dp + 3p^5 q^2\, dq$

32. $z = x \ln(y^2 + 1) \quad \Rightarrow \quad dz = \dfrac{\partial z}{\partial x}\, dx + \dfrac{\partial z}{\partial y}\, dy = \ln(y^2 + 1)\, dx + \dfrac{x}{y^2 + 1}(2y)\, dy = \ln(y^2 + 1)\, dx + \dfrac{2xy}{y^2 + 1}\, dy$

33. $z = e^{-2x} \cos 2\pi t \quad \Rightarrow$

$$dz = \frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial t} dt = e^{-2x}(-2) \cos 2\pi t\, dx + e^{-2x}(-\sin 2\pi t)(2\pi)\, dt = -2e^{-2x} \cos 2\pi t\, dx - 2\pi e^{-2x} \sin 2\pi t\, dt$$

34. $u = \sqrt{x^2 + 3y^2} = (x^2 + 3y^2)^{1/2} \quad \Rightarrow$

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy = \tfrac{1}{2}(x^2 + 3y^2)^{-1/2}(2x)\, dx + \tfrac{1}{2}(x^2 + 3y^2)^{-1/2}(6y)\, dy = \frac{x}{\sqrt{x^2 + 3y^2}}\, dx + \frac{3y}{\sqrt{x^2 + 3y^2}}\, dy$$

35. $H = x^2 y^4 + y^3 z^5 \quad \Rightarrow \quad dH = \dfrac{\partial H}{\partial x}\, dx + \dfrac{\partial H}{\partial y}\, dy + \dfrac{\partial H}{\partial z}\, dz = 2xy^4\, dx + (4x^2 y^3 + 3y^2 z^5)\, dy + 5y^3 z^4\, dz$

36. $w = xze^{-y^2 - z^2} \quad \Rightarrow$

$$\begin{aligned} dw &= \frac{\partial w}{\partial x}\, dx + \frac{\partial w}{\partial y}\, dy + \frac{\partial w}{\partial z}\, dz = ze^{-y^2 - z^2}\, dx + xze^{-y^2 - z^2}(-2y)\, dy + x[z \cdot e^{-y^2 - z^2}(-2z) + e^{-y^2 - z^2} \cdot 1]\, dz \\ &= ze^{-y^2 - z^2}\, dx - 2xyze^{-y^2 - z^2}\, dy + x(1 - 2z^2)e^{-y^2 - z^2}\, dz \end{aligned}$$

37. $R = \alpha\beta^2 \cos\gamma \quad \Rightarrow \quad dR = \dfrac{\partial R}{\partial \alpha}\, d\alpha + \dfrac{\partial R}{\partial \beta}\, d\beta + \dfrac{\partial R}{\partial \gamma}\, d\gamma = \beta^2 \cos\gamma\, d\alpha + 2\alpha\beta \cos\gamma\, d\beta - \alpha\beta^2 \sin\gamma\, d\gamma$

38. $T = \dfrac{v}{1 + uvw} \quad \Rightarrow$

$$\begin{aligned} dT &= \frac{\partial T}{\partial u}\, du + \frac{\partial T}{\partial v}\, dv + \frac{\partial T}{\partial w}\, dw \\ &= v(-1)(1 + uvw)^{-2}(vw)\, du + \frac{1(1 + uvw) - v(uw)}{(1 + uvw)^2}\, dv + v(-1)(1 + uvw)^{-2}(uv)\, dw \\ &= -\frac{v^2 w}{(1 + uvw)^2}\, du + \frac{1}{(1 + uvw)^2}\, dv - \frac{uv^2}{(1 + uvw)^2}\, dw \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. $dx = \Delta x = 0.05$, $dy = \Delta y = 0.1$, $z = 5x^2 + y^2$, $z_x = 10x$, $z_y = 2y$. Thus when $x = 1$ and $y = 2$,

$dz = z_x(1,2)\,dx + z_y(1,2)\,dy = (10)(0.05) + (4)(0.1) = 0.9$ while

$\Delta z = f(1.05, 2.1) - f(1,2) = 5(1.05)^2 + (2.1)^2 - 5 - 4 = 0.9225.$

40. $dx = \Delta x = -0.04$, $dy = \Delta y = 0.05$, $z = x^2 - xy + 3y^2$, $z_x = 2x - y$, $z_y = 6y - x$. Thus when $x = 3$ and $y = -1$,

$dz = (7)(-0.04) + (-9)(0.05) = -0.73$ while $\Delta z = (2.96)^2 - (2.96)(-0.95) + 3(-0.95)^2 - (9 + 3 + 3) = -0.7189.$

41. $dA = \dfrac{\partial A}{\partial x}\,dx + \dfrac{\partial A}{\partial y}\,dy = y\,dx + x\,dy$ and $|\Delta x| \le 0.1$, $|\Delta y| \le 0.1$. We use $dx = 0.1$, $dy = 0.1$ with $x = 30$, $y = 24$; then the maximum error in the area is about $dA = 24(0.1) + 30(0.1) = 5.4\ \text{cm}^2$.

42. Let V be the volume. Then $V = \pi r^2 h$ and $\Delta V \approx dV = 2\pi rh\,dr + \pi r^2\,dh$ is an estimate of the amount of metal. With $dr = 0.05$ and $dh = 0.2$ (0.1 on top, 0.1 on bottom), we get $dV = 2\pi(2)(10)(0.05) + \pi(2)^2(0.2) = 2.80\pi \approx 8.8\ \text{cm}^3$.

43. The volume of a can is $V = \pi r^2 h$ and $\Delta V \approx dV$ is an estimate of the amount of tin. Here $dV = 2\pi rh\,dr + \pi r^2\,dh$, so put $dr = 0.04$, $dh = 0.08$ (0.04 on top, 0.04 on bottom) and then $\Delta V \approx dV = 2\pi(48)(0.04) + \pi(16)(0.08) \approx 16.08\ \text{cm}^3$. Thus the amount of tin is about $16\ \text{cm}^3$.

44. (a) Let A be the area. Then $A = \dfrac{1}{2}bh$ and $\Delta A \approx dA = \dfrac{\partial A}{\partial b}\,db + \dfrac{\partial A}{\partial h}\,dh = \dfrac{1}{2}h\,db + \dfrac{1}{2}b\,dh$. We have $|\Delta h| = |\Delta b| \le \varepsilon$.

So we take $dh = db = \varepsilon$ with $b = 28$ inches, $h = 16$ inches. The maximum error in the area is

$dA = \frac{1}{2}(16)\varepsilon + \frac{1}{2}(28)\varepsilon = 22\varepsilon$ square inches.

(b) With $\varepsilon = \frac{1}{4}$, we have the estimated maximum error in the area of the triangle as $22\left(\frac{1}{4}\right) = \frac{11}{2}$ square inches.

45. (a) Let V be the volume. Then $V = \pi r^2 h$ and $\Delta V \approx dV = \dfrac{\partial V}{\partial r}\,dr + \dfrac{\partial V}{\partial h}\,dh = 2\pi rh\,dr + \pi r^2\,dh$. We have

$|\Delta r| = |\Delta h| = \varepsilon$. So we take $dr = dh = \varepsilon$ with $r = 2.5$ feet and $h = 12$ feet. The maximum error in the volume is

$dV = 2\pi(2.5)(12)\varepsilon + \pi(2.5^2)\varepsilon = 66.25\pi\varepsilon\ \text{ft}^3$.

(b) We need $66.25\pi\varepsilon \le 1 \quad \Rightarrow \quad \varepsilon \lesssim 0.0048$ feet or 0.058 inches.

46. $W = 13.12 + 0.6215T - 11.37v^{0.16} + 0.3965Tv^{0.16}$, so the differential of W is

$$\begin{aligned} dW &= \frac{\partial W}{\partial T}\,dT + \frac{\partial W}{\partial v}\,dv = (0.6215 + 0.3965v^{0.16})\,dT + \left[-11.37(0.16)v^{-0.84} + 0.3965T(0.16)v^{-0.84}\right] dv \\ &= (0.6215 + 0.3965v^{0.16})\,dT + (-1.8192 + 0.06344T)v^{-0.84}\,dv \end{aligned}$$

Here we have $|\Delta T| \le 1$, $|\Delta v| \le 2$, so we take $dT = 1$, $dv = 2$ with $T = -11$, $v = 26$. The maximum error in the calculated value of W is about $dW = (0.6215 + 0.3965(26)^{0.16})(1) + (-1.8192 + 0.06344(-11))(26)^{-0.84}(2) \approx 0.96$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $T = \dfrac{mgR}{2r^2 + R^2}$, so the differential of T is

$$dT = \frac{\partial T}{\partial R}\,dR + \frac{\partial T}{\partial r}\,dr = \frac{(2r^2+R^2)(mg) - mgR(2R)}{(2r^2+R^2)^2}\,dR + \frac{(2r^2+R^2)(0) - mgR(4r)}{(2r^2+R^2)^2}\,dr$$
$$= \frac{mg(2r^2-R^2)}{(2r^2+R^2)^2}\,dR - \frac{4mgRr}{(2r^2+R^2)^2}\,dr$$

Here we have $\Delta R = 0.1$ and $\Delta r = 0.1$, so we take $dR = 0.1$, $dr = 0.1$ with $R = 3$, $r = 0.7$. Then the change in the tension T is approximately

$$dT = \frac{mg[2(0.7)^2 - (3)^2]}{[2(0.7)^2 + (3)^2]^2}\,(0.1) - \frac{4mg(3)(0.7)}{[2(0.7)^2 + (3)^2]^2}\,(0.1)$$
$$= -\frac{0.802mg}{(9.98)^2} - \frac{0.84mg}{(9.98)^2} = -\frac{1.642}{99.6004}mg \approx -0.0165mg$$

Because the change is negative, tension decreases.

48. Here $dV = \Delta V = 0.3$, $dT = \Delta T = -5$, $P = 8.31\,\dfrac{T}{V}$, so

$dP = \left(\dfrac{8.31}{V}\right)dT - \dfrac{8.31 \cdot T}{V^2}\,dV = 8.31\left[-\dfrac{5}{12} - \dfrac{310}{144}\cdot\dfrac{3}{10}\right] \approx -8.83$. Thus the pressure will drop by about $8.83\,\text{kPa}$.

49. $\dfrac{1}{R} = \dfrac{1}{R_1} + \dfrac{1}{R_2} + \dfrac{1}{R_3}$. First we find $\dfrac{\partial R}{\partial R_1}$ implicitly by taking partial derivatives of both sides with respect to R_1:

$\dfrac{\partial}{\partial R_1}\left(\dfrac{1}{R}\right) = \dfrac{\partial\,[(1/R_1) + (1/R_2) + (1/R_3)]}{\partial R_1} \quad\Rightarrow\quad -R^{-2}\dfrac{\partial R}{\partial R_1} = -R_1^{-2} \quad\Rightarrow\quad \dfrac{\partial R}{\partial R_1} = \dfrac{R^2}{R_1^2}$. Then by symmetry,

$\dfrac{\partial R}{\partial R_2} = \dfrac{R^2}{R_2^2}$, $\dfrac{\partial R}{\partial R_3} = \dfrac{R^2}{R_3^2}$. When $R_1 = 25$, $R_2 = 40$ and $R_3 = 50$, $\dfrac{1}{R} = \dfrac{17}{200} \quad\Leftrightarrow\quad R = \frac{200}{17}\ \Omega$. Since the possible error for each R_i is 0.5%, the maximum error of R is attained by setting $\Delta R_i = 0.005R_i$. So

$\Delta R \approx dR = \dfrac{\partial R}{\partial R_1}\,\Delta R_1 + \dfrac{\partial R}{\partial R_2}\,\Delta R_2 + \dfrac{\partial R}{\partial R_3}\,\Delta R_3 = (0.005)R^2\left(\dfrac{1}{R_1} + \dfrac{1}{R_2} + \dfrac{1}{R_3}\right) = (0.005)R = \frac{1}{17} \approx 0.059\ \Omega$.

50. $S = 0.1091w^{0.425}h^{0.725}$. The errors in measurement are at most 2%, so $\left|\dfrac{\Delta w}{w}\right| \le 0.02$ and $\left|\dfrac{\Delta h}{h}\right| \le 0.02$. The relative error in the calculated surface area is

$$\frac{\Delta S}{S} \approx \frac{dS}{S} = \frac{0.1091(0.425w^{0.425-1})h^{0.725}\,dw + 0.1091w^{0.425}(0.725h^{0.725-1})\,dh}{0.1091w^{0.425}h^{0.725}} = 0.425\frac{dw}{w} + 0.725\frac{dh}{h}$$

To estimate the maximum relative error, we use $\dfrac{dw}{w} = \left|\dfrac{\Delta w}{w}\right| = 0.02$ and $\dfrac{dh}{h} = \left|\dfrac{\Delta h}{h}\right| = 0.02 \quad\Rightarrow$

$\dfrac{dS}{S} = 0.425\,(0.02) + 0.725\,(0.02) = 0.023$. Thus the maximum percentage error is approximately 2.3%.

51. (a) $B(m,h) = m/h^2 \quad\Rightarrow\quad B_m(m,h) = 1/h^2$ and $B_h(m,h) = -2m/h^3$. Since $h > 0$, both B_m and B_h are continuous functions, so B is differentiable at $(23, 1.10)$. We have $B(23, 1.10) = 23/(1.10)^2 \approx 19.01$, $B_m(23, 1.10) = 1/(1.10)^2 \approx 0.8264$, and $B_h(23, 1.10) = -2(23)/(1.10)^3 \approx -34.56$, so the linear

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

approximation of B at $(23, 1.10)$ is

$B(m,h) \approx B(23,1.10)+B_m(23,1.10)(m-23)+B_h(23,1.10)(h-1.10) \approx 19.01+0.8264(m-23)-34.56(h-1.10)$

or $B(m,h) \approx 0.8264m - 34.56h + 38.02$.

(b) From part (a), for values near $m = 23$ and $h = 1.10$, $B(m,h) \approx 0.8264m - 34.56h + 38.02$. If m increases by 1 kg to 24 kg and h increases by 0.03 m to 1.13 m, we estimate the BMI to be $B(24, 1.13) \approx 0.8264(24) - 34.56(1.13) + 38.02 \approx 18.801$. This is very close to the actual computed BMI: $B(24, 1.13) = 24/(1.13)^2 \approx 18.796$.

52. $\mathbf{r}_1(t) = \left\langle 2+3t, 1-t^2, 3-4t+t^2 \right\rangle \Rightarrow \mathbf{r}_1'(t) = \langle 3, -2t, -4+2t \rangle$, $\mathbf{r}_2(u) = \left\langle 1+u^2, 2u^3-1, 2u+1 \right\rangle \Rightarrow \mathbf{r}_2'(u) = \left\langle 2u, 6u^2, 2 \right\rangle$. Both curves pass through $P(2,1,3)$ since $\mathbf{r}_1(0) = \mathbf{r}_2(1) = \langle 2,1,3 \rangle$, so the tangent vectors $\mathbf{r}_1'(0) = \langle 3,0,-4 \rangle$ and $\mathbf{r}_2'(1) = \langle 2,6,2 \rangle$ are both parallel to the tangent plane to S at P. A normal vector for the tangent plane is $\mathbf{r}_1'(0) \times \mathbf{r}_2'(1) = \langle 3,0,-4 \rangle \times \langle 2,6,2 \rangle = \langle 24,-14,18 \rangle$, so an equation of the tangent plane is $24(x-2) - 14(y-1) + 18(z-3) = 0$ or $12x - 7y + 9z = 44$.

53. To show that f is continuous at (a,b) we need to show that $\lim\limits_{(x,y)\to(a,b)} f(x,y) = f(a,b)$ or equivalently $\lim\limits_{(\Delta x,\Delta y)\to(0,0)} f(a+\Delta x, b+\Delta y) = f(a,b)$. Since f is differentiable at (a,b), $f(a+\Delta x, b+\Delta y) - f(a,b) = \Delta z = f_x(a,b)\,\Delta x + f_y(a,b)\,\Delta y + \varepsilon_1\,\Delta x + \varepsilon_2\,\Delta y$, where ϵ_1 and $\epsilon_2 \to 0$ as $(\Delta x, \Delta y) \to (0,0)$. Thus $f(a+\Delta x, b+\Delta y) = f(a,b) + f_x(a,b)\,\Delta x + f_y(a,b)\,\Delta y + \varepsilon_1\,\Delta x + \varepsilon_2\,\Delta y$. Taking the limit of both sides as $(\Delta x, \Delta y) \to (0,0)$ gives $\lim\limits_{(\Delta x,\Delta y)\to(0,0)} f(a+\Delta x, b+\Delta y) = f(a,b)$. Thus f is continuous at (a,b).

54. (a) $\lim\limits_{h\to 0} \dfrac{f(h,0)-f(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{0-0}{h} = 0$ and $\lim\limits_{h\to 0} \dfrac{f(0,h)-f(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{0-0}{h} = 0$. Thus $f_x(0,0) = f_y(0,0) = 0$.

To show that f isn't differentiable at $(0,0)$ we need only show that f is not continuous at $(0,0)$ and apply the contrapositive of Exercise 53. As $(x,y) \to (0,0)$ along the x-axis $f(x,y) = 0/x^2 = 0$ for $x \neq 0$ so $f(x,y) \to 0$ as $(x,y) \to (0,0)$ along the x-axis. But as $(x,y) \to (0,0)$ along the line $y = x$, $f(x,x) = x^2/\left(2x^2\right) = \frac{1}{2}$ for $x \neq 0$ so $f(x,y) \to \frac{1}{2}$ as $(x,y) \to (0,0)$ along this line. Thus $\lim\limits_{(x,y)\to(0,0)} f(x,y)$ doesn't exist, so f is discontinuous at $(0,0)$ and thus not differentiable there.

(b) For $(x,y) \neq (0,0)$, $f_x(x,y) = \dfrac{(x^2+y^2)y - xy(2x)}{(x^2+y^2)^2} = \dfrac{y(y^2-x^2)}{(x^2+y^2)^2}$. If we approach $(0,0)$ along the y-axis, then $f_x(x,y) = f_x(0,y) = \dfrac{y^3}{y^4} = \dfrac{1}{y}$, so $f_x(x,y) \to \pm\infty$ as $(x,y) \to (0,0)$. Thus $\lim\limits_{(x,y)\to(0,0)} f_x(x,y)$ does not exist and $f_x(x,y)$ is not continuous at $(0,0)$. Similarly, $f_y(x,y) = \dfrac{(x^2+y^2)x - xy(2y)}{(x^2+y^2)^2} = \dfrac{x(x^2-y^2)}{(x^2+y^2)^2}$ for $(x,y) \neq (0,0)$, and if we approach $(0,0)$ along the x-axis, then $f_y(x,y) = f_x(x,0) = \dfrac{x^3}{x^4} = \dfrac{1}{x}$. Thus $\lim\limits_{(x,y)\to(0,0)} f_y(x,y)$ does not exist and $f_y(x,y)$ is not continuous at $(0,0)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

APPLIED PROJECT The Speedo LZR Racer

1. $v(P,C) = \left(\dfrac{2P}{kC}\right)^{1/3} \quad \Rightarrow$

$$f(x,y) = \frac{v(P+xP, C+yC) - v(P,C)}{v(P,C)} = \frac{v(P+xP, C+yC)}{v(P,C)} - \frac{v(P,C)}{v(P,C)} = \frac{\left(\dfrac{2(P+xP)}{k(C+yC)}\right)^{1/3}}{\left(\dfrac{2P}{kC}\right)^{1/3}} - 1$$

$$= \left(\frac{2P(1+x)}{kC(1+y)} \cdot \frac{kC}{2P}\right)^{1/3} - 1 = \left(\frac{1+x}{1+y}\right)^{1/3} - 1$$

Both power and drag cannot be reduced by more than 100%, but both could be increased by any percentage, so $x \geq -1$ and $y \geq -1$. But f is undefined when $y = -1$, so the domain is $\{(x,y) \mid x \geq -1,\ y > -1\}$.

2. If x and y are small, then we can say they are near zero and we can use a linear approximation to f at $(0,0)$.

We have $f(x,y) = (1+x)^{1/3}(1+y)^{-1/3} - 1$ so the partial derivatives are

$f_x(x,y) = \frac{1}{3}(1+x)^{-2/3}(1+y)^{-1/3} = \dfrac{1}{3(1+x)^{2/3}(1+y)^{1/3}}$ and

$f_y(x,y) = -\frac{1}{3}(1+x)^{1/3}(1+y)^{-4/3} = -\dfrac{(1+x)^{1/3}}{3(1+y)^{4/3}}$. Note that f_x and f_y are continuous functions for $x > -1$, $y > -1$ so f is differentiable at $(0,0)$. Then $f_x(0,0) = \frac{1}{3}$ and $f_y(0,0) = -\frac{1}{3}$, and the linear approximation is $f(x,y) \approx f(0,0) + f_x(0,0)(x-0) + f_y(0,0)(y-0) = 0 + \frac{1}{3}(x-0) - \frac{1}{3}(y-0) = \frac{1}{3}x - \frac{1}{3}y$. According to the linear approximation, a small fractional increase in power results in 1/3 that fractional increase in speed, and a small decrease in drag has the same effect.

3. $f_{xx}(x,y) = \dfrac{1}{3(1+y)^{1/3}} \cdot \left(-\frac{2}{3}\right)(1+x)^{-5/3} = -\dfrac{2}{9(1+x)^{5/3}(1+y)^{1/3}}$,

$f_{yy}(x,y) = -\frac{1}{3}(1+x)^{1/3} \cdot \left(-\frac{4}{3}\right)(1+y)^{-7/3} = \dfrac{4(1+x)^{1/3}}{9(1+y)^{7/3}}$. Because $f_x(x,y)$ is positive in the domain of f, an increase in power results in an increase in speed, but $f_{xx}(x,y)$ is negative, so as the fractional power increases, the fractional speed increases at a declining rate. (We can say that in the positive x-direction, f is increasing and concave downward.) Thus the linear approximation gives an overestimate for an increase in power. Since $f_y(x,y)$ is negative, a *decrease* in drag *increases* speed. But $f_{yy}(x,y)$ is positive, so f_y increases as y increases and f_y decreases (f_y becomes larger and larger negative) as y decreases. (In the positive y-direction, f is decreasing and concave upward.) Thus as the fractional drag decreases, the fractional speed increases at an accelerating pace and the linear approximation gives an underestimate of the increase in power. This explains why a decrease in drag is more effective than an increase in power: Reducing drag improves speed at an increasing rate while adding power improves speed at a declining rate.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. The level curves of $f(x,y) = \left(\dfrac{1+x}{1+y}\right)^{1/3} - 1$ are

$$\left(\frac{1+x}{1+y}\right)^{1/3} - 1 = c \quad \Rightarrow \quad \frac{1+x}{1+y} = (1+c)^3 \quad \Rightarrow$$

$$y = \frac{1+x}{(1+c)^3} - 1.$$

From the level curves, we see that increasing x (from 0) by a small amount has a similar effect on the value of f as decreasing y by a small amount. However, for larger changes, a decrease in y gives greater values of f than a similar increase in x.

14.5 The Chain Rule

1. Find dz/dt using the Chain Rule: $z = x^2y + xy^2, \quad x = 3t, \quad y = t^2 \quad \Rightarrow$

$$\begin{aligned}\frac{dz}{dt} &= \frac{\partial z}{\partial x}\cdot\frac{dx}{dt} + \frac{\partial z}{\partial y}\cdot\frac{dy}{dt} = (2xy+y^2)(3) + (x^2+2xy)(2t)\\ &= 3\left[2(3t)(t^2) + (t^2)^2\right] + 2t\left[(3t)^2 + 2(3t)(t^2)\right] \qquad [\text{with } x = 3t \text{ and } y = t^2]\\ &= 18t^3 + 3t^4 + 18t^3 + 12t^4 = 36t^3 + 15t^4\end{aligned}$$

Find dz/dt by substituting first: $z\left(x(t), y(t)\right) = (3t)^2(t^2) + (3t)(t^2)^2 = 9t^4 + 3t^5 \quad \Rightarrow \quad \dfrac{dz}{dt} = 36t^3 + 15t^4$

Yes, the two answers agree.

2. Find dz/dt using the Chain Rule: $z = xye^y, \quad x = t^2, \quad y = 5t \quad \Rightarrow$

$$\begin{aligned}\frac{dz}{dt} &= \frac{\partial z}{\partial x}\cdot\frac{dx}{dt} + \frac{\partial z}{\partial y}\cdot\frac{dy}{dt} = ye^y(2t) + x(ye^y + e^y)(5)\\ &= (5t)(e^{5t})(2t) + t^2(5te^{5t} + e^{5t})(5) \qquad [\text{with } x = t^2 \text{ and } y = 5t]\\ &= 10t^2e^{5t} + 25t^3e^{5t} + 5t^2e^{5t} = 15t^2e^{5t} + 25t^3e^{5t} = 5t^2e^{5t}(3+5t)\end{aligned}$$

Find dz/dt by substituting first: $z\left(x(t), y(t)\right) = (t^2)(5t)e^{5t} = 5t^3e^{5t} \quad \Rightarrow \quad \dfrac{dz}{dt} = 5t^3e^{5t}(5) + 15t^2e^{5t} = 5t^2e^{5t}(3+5t)$

Yes, the two answers agree.

3. $z = xy^3 - x^2y, \quad x = t^2 + 1, \quad y = t^2 - 1 \quad \Rightarrow$

$$\frac{dz}{dt} = \frac{\partial z}{\partial x}\frac{dx}{dt} + \frac{\partial z}{\partial y}\frac{dy}{dt} = (y^3 - 2xy)(2t) + (3xy^2 - x^2)(2t) = 2t(y^3 - 2xy + 3xy^2 - x^2)$$

4. $z = \dfrac{x-y}{x+2y}, \quad x = e^{\pi t}, \quad y = e^{-\pi t} \quad \Rightarrow$

$$\begin{aligned}\frac{dz}{dt} &= \frac{\partial z}{\partial x}\frac{dx}{dt} + \frac{\partial z}{\partial y}\frac{dy}{dt} = \frac{(x+2y)(1) - (x-y)(1)}{(x+2y)^2}(\pi e^{\pi t}) + \frac{(x+2y)(-1) - (x-y)(2)}{(x+2y)^2}(-\pi e^{-\pi t})\\ &= \frac{3y}{(x+2y)^2}\left(\pi e^{\pi t}\right) + \frac{-3x}{(x+2y)^2}\left(-\pi e^{-\pi t}\right) = \frac{3\pi}{(x+2y)^2}\left(ye^{\pi t} + xe^{-\pi t}\right)\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $z = \sin x \cos y, \quad x = \sqrt{t}, \quad y = 1/t \quad \Rightarrow$

$$\frac{dz}{dt} = \frac{\partial z}{\partial x}\frac{dx}{dt} + \frac{\partial z}{\partial y}\frac{dy}{dt} = (\cos x \cos y)\left(\tfrac{1}{2}t^{-1/2}\right) + (-\sin x \sin y)\left(-t^{-2}\right) = \frac{1}{2\sqrt{t}}\cos x \cos y + \frac{1}{t^2}\sin x \sin y$$

6. $z = \sqrt{1+xy}, \quad x = \tan t, \quad y = \arctan t \quad \Rightarrow$

$$\begin{aligned}\frac{dz}{dt} &= \frac{\partial z}{\partial x}\frac{dx}{dt} + \frac{\partial z}{\partial y}\frac{dy}{dt} = \tfrac{1}{2}(1+xy)^{-1/2}(y)\cdot\sec^2 t + \tfrac{1}{2}(1+xy)^{-1/2}(x)\cdot\frac{1}{1+t^2}\\ &= \frac{1}{2\sqrt{1+xy}}\left(y\sec^2 t + \frac{x}{1+t^2}\right)\end{aligned}$$

7. $w = xe^{y/z}, \quad x = t^2, \quad y = 1-t, \quad z = 1+2t \quad \Rightarrow$

$$\frac{dw}{dt} = \frac{\partial w}{\partial x}\frac{dx}{dt} + \frac{\partial w}{\partial y}\frac{dy}{dt} + \frac{\partial w}{\partial z}\frac{dz}{dt} = e^{y/z}\cdot 2t + xe^{y/z}\left(\frac{1}{z}\right)\cdot(-1) + xe^{y/z}\left(-\frac{y}{z^2}\right)\cdot 2 = e^{y/z}\left(2t - \frac{x}{z} - \frac{2xy}{z^2}\right)$$

8. $w = \ln\sqrt{x^2+y^2+z^2} = \frac{1}{2}\ln(x^2+y^2+z^2), \quad x = \sin t, \quad y = \cos t, \quad z = \tan t \quad \Rightarrow$

$$\begin{aligned}\frac{dw}{dt} &= \frac{\partial w}{\partial x}\frac{dx}{dt} + \frac{\partial w}{\partial y}\frac{dy}{dt} + \frac{\partial w}{\partial z}\frac{dz}{dt} = \frac{1}{2}\cdot\frac{2x}{x^2+y^2+z^2}\cdot\cos t + \frac{1}{2}\cdot\frac{2y}{x^2+y^2+z^2}\cdot(-\sin t) + \frac{1}{2}\cdot\frac{2z}{x^2+y^2+z^2}\cdot\sec^2 t\\ &= \frac{x\cos t - y\sin t + z\sec^2 t}{x^2+y^2+z^2}\end{aligned}$$

9. First we find $\partial z/\partial s$ in two ways. $z = x^2 + y^2, \quad x = 2s+3t, \quad y = s+t \quad \Rightarrow$

$$\begin{aligned}\frac{\partial z}{\partial s} &= \frac{\partial z}{\partial x}\cdot\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\cdot\frac{\partial y}{\partial s} = 2x(2) + 2y(1)\\ &= 2(2s+3t)(2) + 2(s+t)(1) \qquad [\text{with } x = 2s+3t \text{ and } y = s+t]\\ &= 8s + 12t + 2s + 2t = 10s + 14t\end{aligned}$$

$$z(x(s,t), y(s,t)) = (2s+3t)^2 + (s+t)^2 \quad \Rightarrow \quad \frac{\partial z}{\partial s} = 2(2s+3t)(2) + 2(s+t)(1) = 8s + 12t + 2s + 2t = 10s + 14t$$

Yes, the two answers agree. Now we find $\partial z/\partial t$ in two ways.

$$\begin{aligned}\frac{\partial z}{\partial t} &= \frac{\partial z}{\partial x}\cdot\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\cdot\frac{\partial y}{\partial t} = 2x(3) + 2y(1)\\ &= 2(2s+3t)(3) + 2(s+t)(1) \qquad [\text{with } x = 2s+3t \text{ and } y = s+t]\\ &= 12s + 18t + 2s + 2t = 14s + 20t\end{aligned}$$

$$z(x(s,t), y(s,t)) = (2s+3t)^2 + (s+t)^2 \quad \Rightarrow \quad \frac{\partial z}{\partial t} = 2(2s+3t)(3) + 2(s+t)(1) = 12s + 18t + 2s + 2t = 14s + 20t$$

Yes, the two answers agree.

10. First we find $\partial z/\partial s$ in two ways. $z = x^2\sin y, \quad x = s^2 t, \quad y = st \quad \Rightarrow$

$$\begin{aligned}\frac{\partial z}{\partial s} &= \frac{\partial z}{\partial x}\cdot\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\cdot\frac{\partial y}{\partial s} = 2x\sin y(2st) + x^2\cos y(t)\\ &= 2(s^2 t)\sin(st)(2st) + (s^2 t)^2\cos(st)(t) \qquad [\text{with } x = s^2 t \text{ and } y = st]\\ &= 4s^3 t^2\sin(st) + s^4 t^3\cos(st)\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$z(x(s,t), y(s,t)) = (s^2t)^2 \sin(st) = s^4t^2 \sin(st) \quad \Rightarrow$

$\dfrac{\partial z}{\partial s} = s^4t^2 \cos(st)(t) + 4s^3t^2 \sin(st) = 4s^3t^2 \sin(st) + s^4t^3 \cos(st)$

Yes, the two answers agree. Now we find $\partial z/\partial t$ in two ways.

$$\begin{aligned}\frac{\partial z}{\partial t} &= \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial t} = 2x \sin y(s^2) + x^2 \cos y(s) \\ &= 2(s^2t)\sin(st)(s^2) + (s^2t)^2 \cos(st)(s) \qquad [\text{with } x = s^2t \text{ and } y = st] \\ &= 2s^4t \sin(st) + s^5t^2 \cos(st)\end{aligned}$$

$z(x(s,t), y(s,t)) = (s^2t)^2 \sin(st) = s^4t^2 \sin(st) \quad \Rightarrow$

$\dfrac{\partial z}{\partial t} = 2s^4t \sin(st) + s^4t^2 \cos(st)(s) = 2s^4t \sin(st) + s^5t^2 \cos(st)$

Yes, the two answers agree.

11. $z = (x-y)^5, \quad x = s^2t, \quad y = st^2 \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial s} = 5(x-y)^4(1) \cdot 2st + 5(x-y)^4(-1) \cdot t^2 = 5(x-y)^4\left(2st - t^2\right)$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t} = 5(x-y)^4(1) \cdot s^2 + 5(x-y)^4(-1) \cdot 2st = 5(x-y)^4\left(s^2 - 2st\right)$$

12. $z = \tan^{-1}(x^2+y^2), \quad x = s\ln t, \quad y = te^s \quad \Rightarrow$

$$\begin{aligned}\frac{\partial z}{\partial s} &= \frac{\partial z}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial s} = \frac{2x}{1+(x^2+y^2)^2} \cdot \ln t + \frac{2y}{1+(x^2+y^2)^2} \cdot te^s \\ &= \frac{2}{1+(x^2+y^2)^2}\left(x\ln t + yte^s\right)\end{aligned}$$

$$\begin{aligned}\frac{\partial z}{\partial t} &= \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t} = \frac{2x}{1+(x^2+y^2)^2} \cdot \frac{s}{t} + \frac{2y}{1+(x^2+y^2)^2} \cdot e^s \\ &= \frac{2}{1+(x^2+y^2)^2}\left(\frac{xs}{t} + ye^s\right)\end{aligned}$$

13. $z = \ln(3x+2y), \quad x = s\sin t, \quad y = t\cos s \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial s} = \frac{3}{3x+2y}(\sin t) + \frac{2}{3x+2y}(-t\sin s) = \frac{3\sin t - 2t\sin s}{3x+2y}$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t} = \frac{3}{3x+2y}(s\cos t) + \frac{2}{3x+2y}(\cos s) = \frac{3s\cos t + 2\cos s}{3x+2y}$$

14. $z = \sqrt{x}\,e^{xy}, \quad x = 1+st, \quad y = s^2 - t^2 \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial s} = \left(\sqrt{x} \cdot e^{xy}(y) + e^{xy} \cdot \tfrac{1}{2}x^{-1/2}\right)(t) + \sqrt{x}\,e^{xy}(x)\,(2s) = \left(yt\sqrt{x} + \frac{t}{2\sqrt{x}} + 2x^{3/2}s\right)e^{xy}$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t} = \left(\sqrt{x} \cdot e^{xy}(y) + e^{xy} \cdot \tfrac{1}{2}x^{-1/2}\right)(s) + \sqrt{x}\,e^{xy}(x)\,(-2t) = \left(ys\sqrt{x} + \frac{s}{2\sqrt{x}} - 2x^{3/2}t\right)e^{xy}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $z = (\sin\theta)/r, \quad r = st, \quad \theta = s^2 + t^2 \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial r}\cdot\frac{\partial r}{\partial s} + \frac{\partial z}{\partial \theta}\cdot\frac{\partial \theta}{\partial s} = -\frac{\sin\theta}{r^2}(t) + \frac{\cos\theta}{r}(2s) = -\frac{t\sin\theta}{r^2} + \frac{2s\cos\theta}{r}$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial r}\cdot\frac{\partial r}{\partial t} + \frac{\partial z}{\partial \theta}\cdot\frac{\partial \theta}{\partial t} = -\frac{\sin\theta}{r^2}(s) + \frac{\cos\theta}{r}(2t) = -\frac{s\sin\theta}{r^2} + \frac{2t\cos\theta}{r}$$

16. $z = \tan(u/v), \quad u = 2s + 3t, \quad v = 3s - 2t \quad \Rightarrow$

$$\begin{aligned}\frac{\partial z}{\partial s} &= \frac{\partial z}{\partial u}\frac{\partial u}{\partial s} + \frac{\partial z}{\partial v}\frac{\partial v}{\partial s} = \sec^2(u/v)(1/v)\cdot 2 + \sec^2(u/v)(-uv^{-2})\cdot 3\\ &= \frac{2}{v}\sec^2\left(\frac{u}{v}\right) - \frac{3u}{v^2}\sec^2\left(\frac{u}{v}\right) = \frac{2v-3u}{v^2}\sec^2\left(\frac{u}{v}\right)\end{aligned}$$

$$\begin{aligned}\frac{\partial z}{\partial t} &= \frac{\partial z}{\partial u}\frac{\partial u}{\partial t} + \frac{\partial z}{\partial v}\frac{\partial v}{\partial t} = \sec^2(u/v)(1/v)\cdot 3 + \sec^2(u/v)(-uv^{-2})\cdot(-2)\\ &= \frac{3}{v}\sec^2\left(\frac{u}{v}\right) + \frac{2u}{v^2}\sec^2\left(\frac{u}{v}\right) = \frac{2u+3v}{v^2}\sec^2\left(\frac{u}{v}\right)\end{aligned}$$

17. Let $x = g(t)$ and $y = h(t)$. Then $p(t) = f(x, y)$ and the Chain Rule (2) gives $\dfrac{dp}{dt} = \dfrac{\partial f}{\partial x}\dfrac{dx}{dt} + \dfrac{\partial f}{\partial y}\dfrac{dy}{dt}$. When $t = 2$, $x = g(2) = 4$ and $y = h(2) = 5$, so $p'(2) = f_x(4,5)\,g'(2) + f_y(4,5)\,h'(2) = (2)(-3) + (8)(6) = 42$.

18. $R(s,t) = G\left(u(s,t), v(s,t)\right) \quad \Rightarrow \quad \dfrac{\partial R}{\partial s} = \dfrac{\partial R}{\partial u}\dfrac{\partial u}{\partial s} + \dfrac{\partial R}{\partial v}\dfrac{\partial v}{\partial s}$ and $\dfrac{\partial R}{\partial t} = \dfrac{\partial R}{\partial u}\dfrac{\partial u}{\partial t} + \dfrac{\partial R}{\partial v}\dfrac{\partial v}{\partial t}$ by the Chain Rule (3). When $s = 1$ and $t = 2$, $u(1,2) = 5$ and $v(1,2) = 7$.

Thus $R_s(1,2) = G_u(5,7)\,u_s(1,2) + G_v(5,7)\,v_s(1,2) = (9)(4) + (-2)(2) = 32$ and

$R_t(1,2) = G_u(5,7)\,u_t(1,2) + G_v(5,7)\,v_t(1,2) = (9)(-3) + (-2)(6) = -39$.

19. $g(u,v) = f(x(u,v), y(u,v))$ where $x = e^u + \sin v,\ y = e^u + \cos v \quad \Rightarrow$

$\dfrac{\partial x}{\partial u} = e^u,\ \dfrac{\partial x}{\partial v} = \cos v,\ \dfrac{\partial y}{\partial u} = e^u,\ \dfrac{\partial y}{\partial v} = -\sin v$. By the Chain Rule (3), $\dfrac{\partial g}{\partial u} = \dfrac{\partial f}{\partial x}\dfrac{\partial x}{\partial u} + \dfrac{\partial f}{\partial y}\dfrac{\partial y}{\partial u}$. Then

$g_u(0,0) = f_x(x(0,0), y(0,0))\,x_u(0,0) + f_y(x(0,0), y(0,0))\,y_u(0,0) = f_x(1,2)(e^0) + f_y(1,2)(e^0) = 2(1) + 5(1) = 7$.

Similarly, $\dfrac{\partial g}{\partial v} = \dfrac{\partial f}{\partial x}\dfrac{\partial x}{\partial v} + \dfrac{\partial f}{\partial y}\dfrac{\partial y}{\partial v}$. Then

$$\begin{aligned}g_v(0,0) &= f_x(x(0,0), y(0,0))\,x_v(0,0) + f_y(x(0,0), y(0,0))\,y_v(0,0) = f_x(1,2)(\cos 0) + f_y(1,2)(-\sin 0)\\ &= 2(1) + 5(0) = 2\end{aligned}$$

20. $g(r,s) = f(x(r,s), y(r,s))$ where $x = 2r - s,\ y = s^2 - 4r \quad \Rightarrow \quad \dfrac{\partial x}{\partial r} = 2,\ \dfrac{\partial x}{\partial s} = -1,\ \dfrac{\partial y}{\partial r} = -4,\ \dfrac{\partial y}{\partial s} = 2s$.

By the Chain Rule (3) $\dfrac{\partial g}{\partial r} = \dfrac{\partial f}{\partial x}\dfrac{\partial x}{\partial r} + \dfrac{\partial f}{\partial y}\dfrac{\partial y}{\partial r}$. Then

$$\begin{aligned}g_r(1,2) &= f_x(x(1,2), y(1,2))\,x_r(1,2) + f_y(x(1,2), y(1,2))\,y_r(1,2) = f_x(0,0)(2) + f_y(0,0)(-4)\\ &= 4(2) + 8(-4) = -24\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Similarly, $\dfrac{\partial g}{\partial s} = \dfrac{\partial f}{\partial x}\dfrac{\partial x}{\partial s} + \dfrac{\partial f}{\partial y}\dfrac{\partial y}{\partial s}$. Then

$$\begin{aligned} g_s(1,2) &= f_x(x(1,2), y(1,2))\, x_s(1,2) + f_y(x(1,2), y(1,2))\, y_s(1,2) = f_x(0,0)(-1) + f_y(0,0)(4) \\ &= 4(-1) + 8(4) = 28 \end{aligned}$$

21.

$u = f(x,y),\ x = x(r,s,t),\ y = y(r,s,t) \quad \Rightarrow$

$$\frac{\partial u}{\partial r} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial r} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial r}, \quad \frac{\partial u}{\partial s} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial s},$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial t}$$

22.

$w = f(x,y,z),\ \ x = x(u,v),\ \ y = y(u,v),\ \ z = z(u,v) \quad \Rightarrow$

$$\frac{\partial w}{\partial u} = \frac{\partial w}{\partial x}\frac{\partial x}{\partial u} + \frac{\partial w}{\partial y}\frac{\partial y}{\partial u} + \frac{\partial w}{\partial z}\frac{\partial z}{\partial u},$$

$$\frac{\partial w}{\partial v} = \frac{\partial w}{\partial x}\frac{\partial x}{\partial v} + \frac{\partial w}{\partial y}\frac{\partial y}{\partial v} + \frac{\partial w}{\partial z}\frac{\partial z}{\partial v}$$

23.

$T = F(p,q,r),\ p = p(x,y,z),\ \ q = q(x,y,z),\ r = r(x,y,z) \quad \Rightarrow$

$$\frac{\partial T}{\partial x} = \frac{\partial T}{\partial p}\frac{\partial p}{\partial x} + \frac{\partial T}{\partial q}\frac{\partial q}{\partial x} + \frac{\partial T}{\partial r}\frac{\partial r}{\partial x},$$

$$\frac{\partial T}{\partial y} = \frac{\partial T}{\partial p}\frac{\partial p}{\partial y} + \frac{\partial T}{\partial q}\frac{\partial q}{\partial y} + \frac{\partial T}{\partial r}\frac{\partial r}{\partial y},$$

$$\frac{\partial T}{\partial z} = \frac{\partial T}{\partial p}\frac{\partial p}{\partial z} + \frac{\partial T}{\partial q}\frac{\partial q}{\partial z} + \frac{\partial T}{\partial r}\frac{\partial r}{\partial z}$$

24.

$R = F(t,u),\ \ t = t(w,x,y,z),\ \ u = u(w,x,y,z) \quad \Rightarrow$

$$\frac{\partial R}{\partial w} = \frac{\partial R}{\partial t}\frac{\partial t}{\partial w} + \frac{\partial R}{\partial u}\frac{\partial u}{\partial w}, \quad \frac{\partial R}{\partial x} = \frac{\partial R}{\partial t}\frac{\partial t}{\partial x} + \frac{\partial R}{\partial u}\frac{\partial u}{\partial x},$$

$$\frac{\partial R}{\partial y} = \frac{\partial R}{\partial t}\frac{\partial t}{\partial y} + \frac{\partial R}{\partial u}\frac{\partial u}{\partial y}, \quad \frac{\partial R}{\partial z} = \frac{\partial R}{\partial t}\frac{\partial t}{\partial z} + \frac{\partial R}{\partial u}\frac{\partial u}{\partial z}$$

25. $z = x^4 + x^2 y,\ x = s + 2t - u,\ y = stu^2 \quad \Rightarrow$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial s} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial s} = (4x^3 + 2xy)(1) + (x^2)(tu^2),$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t} = (4x^3 + 2xy)(2) + (x^2)(su^2),$$

$$\frac{\partial z}{\partial u} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial u} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial u} = (4x^3 + 2xy)(-1) + (x^2)(2stu).$$

When $s = 4$, $t = 2$, and $u = 1$ we have $x = 7$ and $y = 8$,

so $\dfrac{\partial z}{\partial s} = (1484)(1) + (49)(2) = 1582,\ \dfrac{\partial z}{\partial t} = (1484)(2) + (49)(4) = 3164,\ \dfrac{\partial z}{\partial u} = (1484)(-1) + (49)(16) = -700.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $T = v/(2u+v) = v(2u+v)^{-1},\ u = pq\sqrt{r},\ v = p\sqrt{q}\,r \quad \Rightarrow$

$$\frac{\partial T}{\partial p} = \frac{\partial T}{\partial u}\frac{\partial u}{\partial p} + \frac{\partial T}{\partial v}\frac{\partial v}{\partial p} = [-v(2u+v)^{-2}(2)](q\sqrt{r}) + \frac{(2u+v)(1)-v(1)}{(2u+v)^2}\,(\sqrt{q}\,r)$$

$$= \frac{-2v}{(2u+v)^2}\,(q\sqrt{r}) + \frac{2u}{(2u+v)^2}\,(\sqrt{q}\,r),$$

$$\frac{\partial T}{\partial q} = \frac{\partial T}{\partial u}\frac{\partial u}{\partial q} + \frac{\partial T}{\partial v}\frac{\partial v}{\partial q} = \frac{-2v}{(2u+v)^2}\,(p\sqrt{r}) + \frac{2u}{(2u+v)^2}\,\frac{pr}{2\sqrt{q}},$$

$$\frac{\partial T}{\partial r} = \frac{\partial T}{\partial u}\frac{\partial u}{\partial r} + \frac{\partial T}{\partial v}\frac{\partial v}{\partial r} = \frac{-2v}{(2u+v)^2}\,\frac{pq}{2\sqrt{r}} + \frac{2u}{(2u+v)^2}\,(p\sqrt{q}).$$

When $p = 2$, $q = 1$, and $r = 4$ we have $u = 4$ and $v = 8$,

so $\dfrac{\partial T}{\partial p} = \left(-\frac{1}{16}\right)(2) + \left(\frac{1}{32}\right)(4) = 0,\ \dfrac{\partial T}{\partial q} = \left(-\frac{1}{16}\right)(4) + \left(\frac{1}{32}\right)(4) = -\frac{1}{8},\ \dfrac{\partial T}{\partial r} = \left(-\frac{1}{16}\right)\left(\frac{1}{2}\right) + \left(\frac{1}{32}\right)(2) = \frac{1}{32}.$

27. $w = xy + yz + zx,\ x = r\cos\theta,\ y = r\sin\theta,\ z = r\theta \quad \Rightarrow$

$$\frac{\partial w}{\partial r} = \frac{\partial w}{\partial x}\frac{\partial x}{\partial r} + \frac{\partial w}{\partial y}\frac{\partial y}{\partial r} + \frac{\partial w}{\partial z}\frac{\partial z}{\partial r} = (y+z)(\cos\theta) + (x+z)(\sin\theta) + (y+x)(\theta),$$

$$\frac{\partial w}{\partial \theta} = \frac{\partial w}{\partial x}\frac{\partial x}{\partial \theta} + \frac{\partial w}{\partial y}\frac{\partial y}{\partial \theta} + \frac{\partial w}{\partial z}\frac{\partial z}{\partial \theta} = (y+z)(-r\sin\theta) + (x+z)(r\cos\theta) + (y+x)(r).$$

When $r = 2$ and $\theta = \pi/2$ we have $x = 0$, $y = 2$, and $z = \pi$, so $\dfrac{\partial w}{\partial r} = (2+\pi)(0) + (0+\pi)(1) + (2+0)(\pi/2) = 2\pi$

and $\dfrac{\partial w}{\partial \theta} = (2+\pi)(-2) + (0+\pi)(0) + (2+0)(2) = -2\pi.$

28. $P = \sqrt{u^2+v^2+w^2} = (u^2+v^2+w^2)^{1/2},\ u = xe^y,\ v = ye^x,\ w = e^{xy} \quad \Rightarrow$

$$\frac{\partial P}{\partial x} = \frac{\partial P}{\partial u}\frac{\partial u}{\partial x} + \frac{\partial P}{\partial v}\frac{\partial v}{\partial x} + \frac{\partial P}{\partial w}\frac{\partial w}{\partial x}$$

$$= \tfrac{1}{2}(u^2+v^2+w^2)^{-1/2}(2u)(e^y) + \tfrac{1}{2}(u^2+v^2+w^2)^{-1/2}(2v)(ye^x) + \tfrac{1}{2}(u^2+v^2+w^2)^{-1/2}(2w)(ye^{xy})$$

$$= \frac{ue^y + vye^x + wye^{xy}}{\sqrt{u^2+v^2+w^2}},$$

$$\frac{\partial P}{\partial y} = \frac{\partial P}{\partial u}\frac{\partial u}{\partial y} + \frac{\partial P}{\partial v}\frac{\partial v}{\partial y} + \frac{\partial P}{\partial w}\frac{\partial w}{\partial y}$$

$$= \frac{u}{\sqrt{u^2+v^2+w^2}}\,(xe^y) + \frac{v}{\sqrt{u^2+v^2+w^2}}\,(e^x) + \frac{w}{\sqrt{u^2+v^2+w^2}}\,(xe^{xy})$$

$$= \frac{uxe^y + ve^x + wxe^{xy}}{\sqrt{u^2+v^2+w^2}}.$$

When $x = 0$ and $y = 2$ we have $u = 0$, $v = 2$, and $w = 1$, so $\dfrac{\partial P}{\partial x} = \dfrac{0+4+2}{\sqrt{5}} = \dfrac{6}{\sqrt{5}}$ and $\dfrac{\partial P}{\partial y} = \dfrac{0+2+0}{\sqrt{5}} = \dfrac{2}{\sqrt{5}}.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. $N = \dfrac{p+q}{p+r},\ p = u + vw,\ q = v + uw,\ r = w + uv \quad \Rightarrow$

$$\begin{aligned}\frac{\partial N}{\partial u} &= \frac{\partial N}{\partial p}\frac{\partial p}{\partial u} + \frac{\partial N}{\partial q}\frac{\partial q}{\partial u} + \frac{\partial N}{\partial r}\frac{\partial r}{\partial u}\\ &= \frac{(p+r)(1)-(p+q)(1)}{(p+r)^2}(1) + \frac{(p+r)(1)-(p+q)(0)}{(p+r)^2}(w) + \frac{(p+r)(0)-(p+q)(1)}{(p+r)^2}(v)\\ &= \frac{(r-q)+(p+r)w-(p+q)v}{(p+r)^2},\end{aligned}$$

$$\frac{\partial N}{\partial v} = \frac{\partial N}{\partial p}\frac{\partial p}{\partial v} + \frac{\partial N}{\partial q}\frac{\partial q}{\partial v} + \frac{\partial N}{\partial r}\frac{\partial r}{\partial v} = \frac{r-q}{(p+r)^2}(w) + \frac{p+r}{(p+r)^2}(1) + \frac{-(p+q)}{(p+r)^2}(u) = \frac{(r-q)w+(p+r)-(p+q)u}{(p+r)^2},$$

$$\frac{\partial N}{\partial w} = \frac{\partial N}{\partial p}\frac{\partial p}{\partial w} + \frac{\partial N}{\partial q}\frac{\partial q}{\partial w} + \frac{\partial N}{\partial r}\frac{\partial r}{\partial w} = \frac{r-q}{(p+r)^2}(v) + \frac{p+r}{(p+r)^2}(u) + \frac{-(p+q)}{(p+r)^2}(1) = \frac{(r-q)v+(p+r)u-(p+q)}{(p+r)^2}.$$

When $u = 2$, $v = 3$, and $w = 4$ we have $p = 14$, $q = 11$, and $r = 10$, so $\dfrac{\partial N}{\partial u} = \dfrac{-1+(24)(4)-(25)(3)}{(24)^2} = \dfrac{20}{576} = \dfrac{5}{144}$,

$\dfrac{\partial N}{\partial v} = \dfrac{(-1)(4)+24-(25)(2)}{(24)^2} = \dfrac{-30}{576} = -\dfrac{5}{96}$, and $\dfrac{\partial N}{\partial w} = \dfrac{(-1)(3)+(24)(2)-25}{(24)^2} = \dfrac{20}{576} = \dfrac{5}{144}$.

30. $u = xe^{ty},\ x = \alpha^2\beta,\ y = \beta^2\gamma,\ t = \gamma^2\alpha \quad \Rightarrow$

$$\frac{\partial u}{\partial \alpha} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial \alpha} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial \alpha} + \frac{\partial u}{\partial t}\frac{\partial t}{\partial \alpha} = e^{ty}(2\alpha\beta) + xte^{ty}(0) + xye^{ty}(\gamma^2) = e^{ty}(2\alpha\beta + xy\gamma^2),$$

$$\frac{\partial u}{\partial \beta} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial \beta} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial \beta} + \frac{\partial u}{\partial t}\frac{\partial t}{\partial \beta} = e^{ty}(\alpha^2) + xte^{ty}(2\beta\gamma) + xye^{ty}(0) = e^{ty}(\alpha^2 + 2xt\beta\gamma),$$

$$\frac{\partial u}{\partial \gamma} = \frac{\partial u}{\partial x}\frac{\partial x}{\partial \gamma} + \frac{\partial u}{\partial y}\frac{\partial y}{\partial \gamma} + \frac{\partial u}{\partial t}\frac{\partial t}{\partial \gamma} = e^{ty}(0) + xte^{ty}(\beta^2) + xye^{ty}(2\gamma\alpha) = e^{ty}(xt\beta^2 + 2xy\alpha\gamma).$$

When $\alpha = -1$, $\beta = 2$, and $\gamma = 1$ we have $x = 2$, $y = 4$, and $t = -1$, so $\dfrac{\partial u}{\partial \alpha} = e^{-4}(-4+8) = 4e^{-4}$,

$\dfrac{\partial u}{\partial \beta} = e^{-4}(1-8) = -7e^{-4}$, and $\dfrac{\partial u}{\partial \gamma} = e^{-4}(-8-16) = -24e^{-4}$.

31. $y\cos x = x^2 + y^2$, so let $F(x,y) = y\cos x - x^2 - y^2 = 0$. Then by Equation 5

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{-y\sin x - 2x}{\cos x - 2y} = \frac{2x + y\sin x}{\cos x - 2y}.$$

32. $\cos(xy) = 1 + \sin y$, so let $F(x,y) = \cos(xy) - 1 - \sin y = 0$. Then by Equation 5

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{-\sin(xy)(y)}{-\sin(xy)(x) - \cos y} = -\frac{y\sin(xy)}{\cos y + x\sin(xy)}.$$

33. $\tan^{-1}(x^2y) = x + xy^2$, so let $F(x,y) = \tan^{-1}(x^2y) - x - xy^2 = 0$. Then

$$F_x(x,y) = \frac{1}{1+(x^2y)^2}(2xy) - 1 - y^2 = \frac{2xy}{1+x^4y^2} - 1 - y^2 = \frac{2xy - (1+y^2)(1+x^4y^2)}{1+x^4y^2},$$

$$F_y(x,y) = \frac{1}{1+(x^2y)^2}(x^2) - 2xy = \frac{x^2}{1+x^4y^2} - 2xy = \frac{x^2 - 2xy(1+x^4y^2)}{1+x^4y^2}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and $$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{[2xy - (1+y^2)(1+x^4y^2)]/(1+x^4y^2)}{[x^2 - 2xy(1+x^4y^2)]/(1+x^4y^2)} = \frac{(1+y^2)(1+x^4y^2) - 2xy}{x^2 - 2xy(1+x^4y^2)}$$

$$= \frac{1 + x^4y^2 + y^2 + x^4y^4 - 2xy}{x^2 - 2xy - 2x^5y^3}$$

34. $e^y \sin x = x + xy$, so let $F(x,y) = e^y \sin x - x - xy = 0$. Then $\dfrac{dy}{dx} = -\dfrac{F_x}{F_y} = -\dfrac{e^y \cos x - 1 - y}{e^y \sin x - x} = \dfrac{1 + y - e^y \cos x}{e^y \sin x - x}$.

35. $x^2 + 2y^2 + 3z^2 = 1$, so let $F(x,y,z) = x^2 + 2y^2 + 3z^2 - 1 = 0$. Then by Equations 6

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{2x}{6z} = -\frac{x}{3z} \quad \text{and} \quad \frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{4y}{6z} = -\frac{2y}{3z}.$$

36. $x^2 - y^2 + z^2 - 2z = 4$, so let $F(x,y,z) = x^2 - y^2 + z^2 - 2z - 4 = 0$. Then by Equations 6

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{2x}{2z-2} = \frac{x}{1-z} \quad \text{and} \quad \frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{-2y}{2z-2} = \frac{y}{z-1}.$$

37. $e^z = xyz$, so let $F(x,y,z) = e^z - xyz = 0$. Then $\dfrac{\partial z}{\partial x} = -\dfrac{F_x}{F_z} = -\dfrac{-yz}{e^z - xy} = \dfrac{yz}{e^z - xy}$ and

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{-xz}{e^z - xy} = \frac{xz}{e^z - xy}.$$

38. $yz + x\ln y = z^2$, so let $F(x,y,z) = yz + x\ln y - z^2 = 0$. Then $\dfrac{\partial z}{\partial x} = -\dfrac{F_x}{F_z} = -\dfrac{\ln y}{y - 2z} = \dfrac{\ln y}{2z - y}$ and

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{z + (x/y)}{y - 2z} = \frac{x + yz}{2yz - y^2}.$$

39. Since x and y are each functions of t, $T(x,y)$ is a function of t, so by the Chain Rule, $\dfrac{dT}{dt} = \dfrac{\partial T}{\partial x}\dfrac{dx}{dt} + \dfrac{\partial T}{\partial y}\dfrac{dy}{dt}$. After 3 seconds, $x = \sqrt{1+t} = \sqrt{1+3} = 2$, $y = 2 + \frac{1}{3}t = 2 + \frac{1}{3}(3) = 3$, $\dfrac{dx}{dt} = \dfrac{1}{2\sqrt{1+t}} = \dfrac{1}{2\sqrt{1+3}} = \dfrac{1}{4}$, and $\dfrac{dy}{dt} = \dfrac{1}{3}$.

Then $\dfrac{dT}{dt} = T_x(2,3)\dfrac{dx}{dt} + T_y(2,3)\dfrac{dy}{dt} = 4\left(\frac{1}{4}\right) + 3\left(\frac{1}{3}\right) = 2$. Thus the temperature is rising at a rate of $2°\text{C/s}$.

40. (a) Since $\partial W/\partial T$ is negative, a rise in average temperature (while annual rainfall remains constant) causes a decrease in wheat production at the current production levels. Since $\partial W/\partial R$ is positive, an increase in annual rainfall (while the average temperature remains constant) causes an increase in wheat production.

(b) Since the average temperature is rising at a rate of $0.15°\text{C/year}$, we know that $dT/dt = 0.15$. Since rainfall is decreasing at a rate of 0.1 cm/year, we know $dR/dt = -0.1$. Then, by the Chain Rule,

$\dfrac{dW}{dt} = \dfrac{\partial W}{\partial T}\dfrac{dT}{dt} + \dfrac{\partial W}{\partial R}\dfrac{dR}{dt} = (-2)(0.15) + (8)(-0.1) = -1.1$. Thus we estimate that wheat production will decrease at a rate of 1.1 units/year.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. $C = 1449.2 + 4.6T - 0.055T^2 + 0.00029T^3 + 0.016D$, so $\dfrac{\partial C}{\partial T} = 4.6 - 0.11T + 0.00087T^2$ and $\dfrac{\partial C}{\partial D} = 0.016$.

According to the graph, the diver is experiencing a temperature of approximately $12.5°\text{C}$ at $t = 20$ minutes, so $\dfrac{\partial C}{\partial T} = 4.6 - 0.11(12.5) + 0.00087(12.5)^2 \approx 3.36$. By sketching tangent lines at $t = 20$ to the graphs given, we estimate $\dfrac{dD}{dt} \approx \dfrac{1}{2}$ and $\dfrac{dT}{dt} \approx -\dfrac{1}{10}$. Then, by the Chain Rule, $\dfrac{dC}{dt} = \dfrac{\partial C}{\partial T}\dfrac{dT}{dt} + \dfrac{\partial C}{\partial D}\dfrac{dD}{dt} \approx (3.36)\left(-\frac{1}{10}\right) + (0.016)\left(\frac{1}{2}\right) \approx -0.33$.

Thus the speed of sound experienced by the diver is decreasing at a rate of approximately 0.33 m/s per minute.

42. $V = \pi r^2 h/3$, so $\dfrac{dV}{dt} = \dfrac{\partial V}{\partial r}\dfrac{dr}{dt} + \dfrac{\partial V}{\partial h}\dfrac{dh}{dt} = \dfrac{2\pi rh}{3}\,1.8 + \dfrac{\pi r^2}{3}(-2.5) = 20{,}160\pi - 12{,}000\pi = 8160\pi \text{ in}^3/\text{s}$.

43. (a) $V = \ell wh$, so by the Chain Rule,

$$\frac{dV}{dt} = \frac{\partial V}{\partial \ell}\frac{d\ell}{dt} + \frac{\partial V}{\partial w}\frac{dw}{dt} + \frac{\partial V}{\partial h}\frac{dh}{dt} = wh\,\frac{d\ell}{dt} + \ell h\,\frac{dw}{dt} + \ell w\,\frac{dh}{dt} = 2\cdot 2\cdot 2 + 1\cdot 2\cdot 2 + 1\cdot 2\cdot(-3) = 6 \text{ m}^3/\text{s}.$$

(b) $S = 2(\ell w + \ell h + wh)$, so by the Chain Rule,

$$\begin{aligned}\frac{dS}{dt} &= \frac{\partial S}{\partial \ell}\frac{d\ell}{dt} + \frac{\partial S}{\partial w}\frac{dw}{dt} + \frac{\partial S}{\partial h}\frac{dh}{dt} = 2(w+h)\,\frac{d\ell}{dt} + 2(\ell+h)\,\frac{dw}{dt} + 2(\ell+w)\,\frac{dh}{dt}\\ &= 2(2+2)2 + 2(1+2)2 + 2(1+2)(-3) = 10 \text{ m}^2/\text{s}\end{aligned}$$

(c) $L^2 = \ell^2 + w^2 + h^2 \quad\Rightarrow\quad 2L\,\dfrac{dL}{dt} = 2\ell\,\dfrac{d\ell}{dt} + 2w\,\dfrac{dw}{dt} + 2h\,\dfrac{dh}{dt} = 2(1)(2) + 2(2)(2) + 2(2)(-3) = 0 \quad\Rightarrow$

$dL/dt = 0$ m/s.

44. $I = \dfrac{V}{R} \quad\Rightarrow$

$$\frac{dI}{dt} = \frac{\partial I}{\partial V}\frac{dV}{dt} + \frac{\partial I}{\partial R}\frac{dR}{dt} = \frac{1}{R}\frac{dV}{dt} - \frac{V}{R^2}\frac{dR}{dt} = \frac{1}{R}\frac{dV}{dt} - \frac{I}{R}\frac{dR}{dt} = \frac{1}{400}(-0.01) - \frac{0.08}{400}(0.03) = -0.000031 \text{ A/s}$$

45. $\dfrac{dP}{dt} = 0.05$, $\dfrac{dT}{dt} = 0.15$, $V = 8.31\dfrac{T}{P}$ and $\dfrac{dV}{dt} = \dfrac{8.31}{P}\dfrac{dT}{dt} - 8.31\dfrac{T}{P^2}\dfrac{dP}{dt}$. Thus when $P = 20$ and $T = 320$,

$$\frac{dV}{dt} = 8.31\left[\frac{0.15}{20} - \frac{(0.05)(320)}{400}\right] \approx -0.27 \text{ L/s}.$$

46. $P = 1.47L^{0.65}K^{0.35}$ and considering P, L, and K as functions of time t we have

$\dfrac{dP}{dt} = \dfrac{\partial P}{\partial L}\dfrac{dL}{dt} + \dfrac{\partial P}{\partial K}\dfrac{dK}{dt} = 1.47(0.65)L^{-0.35}K^{0.35}\dfrac{dL}{dt} + 1.47(0.35)L^{0.65}K^{-0.65}\dfrac{dK}{dt}$. We are given that $\dfrac{dL}{dt} = -2$ and $\dfrac{dK}{dt} = 0.5$, so when $L = 30$ and $K = 8$, the rate of change of production $\dfrac{dP}{dt}$ is $1.47(0.65)(30)^{-0.35}(8)^{0.35}(-2) + 1.47(0.35)(30)^{0.65}(8)^{-0.65}(0.5) \approx -0.596$. Thus production at that time is decreasing at a rate of about \$596,000 per year.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. Let x be the length of the first side of the triangle and y the length of the second side. The area A of the triangle is given by $A = \frac{1}{2}xy\sin\theta$ [Formula 6 in Appendix D], where θ is the angle between the two sides. Thus A is a function of x, y, and θ, and x, y, and θ are each in turn functions of time t. We are given that $\dfrac{dx}{dt} = 3$, $\dfrac{dy}{dt} = -2$, and because A is constant, $\dfrac{dA}{dt} = 0$. By the Chain Rule, $\dfrac{dA}{dt} = \dfrac{\partial A}{\partial x}\dfrac{dx}{dt} + \dfrac{\partial A}{\partial y}\dfrac{dy}{dt} + \dfrac{\partial A}{\partial \theta}\dfrac{d\theta}{dt} \quad \Rightarrow \quad \dfrac{dA}{dt} = \frac{1}{2}y\sin\theta \cdot \dfrac{dx}{dt} + \frac{1}{2}x\sin\theta \cdot \dfrac{dy}{dt} + \frac{1}{2}xy\cos\theta \cdot \dfrac{d\theta}{dt}$.

When $x = 20$, $y = 30$, and $\theta = \pi/6$ we have

$$\begin{aligned} 0 &= \tfrac{1}{2}(30)\left(\sin\tfrac{\pi}{6}\right)(3) + \tfrac{1}{2}(20)\left(\sin\tfrac{\pi}{6}\right)(-2) + \tfrac{1}{2}(20)(30)\left(\cos\tfrac{\pi}{6}\right)\frac{d\theta}{dt} \\ &= 45 \cdot \tfrac{1}{2} - 20 \cdot \tfrac{1}{2} + 300 \cdot \frac{\sqrt{3}}{2} \cdot \frac{d\theta}{dt} = \tfrac{25}{2} + 150\sqrt{3}\,\frac{d\theta}{dt} \end{aligned}$$

Solving for $\dfrac{d\theta}{dt}$ gives $\dfrac{d\theta}{dt} = \dfrac{-25/2}{150\sqrt{3}} = -\dfrac{1}{12\sqrt{3}}$, so the angle between the sides is decreasing at a rate of $1/\left(12\sqrt{3}\right) \approx 0.048$ rad/s.

48. $f_o = \left(\dfrac{c+v_o}{c-v_s}\right) f_s = \left(\frac{332+34}{332-40}\right) 460 \approx 576.6$ Hz. v_o and v_s are functions of time t, so

$$\begin{aligned} \frac{df_o}{dt} &= \frac{\partial f_o}{\partial v_o}\frac{dv_o}{dt} + \frac{\partial f_o}{\partial v_s}\frac{dv_s}{dt} = \left(\frac{1}{c-v_s}\right) f_s \cdot \frac{dv_o}{dt} + \frac{c+v_o}{(c-v_s)^2} f_s \cdot \frac{dv_s}{dt} \\ &= \left(\tfrac{1}{332-40}\right)(460)\,(1.2) + \tfrac{332+34}{(332-40)^2}\,(460)\,(1.4) \approx 4.65 \text{ Hz/s} \end{aligned}$$

49. (a) By the Chain Rule, $\dfrac{\partial z}{\partial r} = \dfrac{\partial z}{\partial x}\cos\theta + \dfrac{\partial z}{\partial y}\sin\theta$, $\dfrac{\partial z}{\partial \theta} = \dfrac{\partial z}{\partial x}(-r\sin\theta) + \dfrac{\partial z}{\partial y}\,r\cos\theta$.

(b) $\left(\dfrac{\partial z}{\partial r}\right)^2 = \left(\dfrac{\partial z}{\partial x}\right)^2 \cos^2\theta + 2\dfrac{\partial z}{\partial x}\dfrac{\partial z}{\partial y}\cos\theta\,\sin\theta + \left(\dfrac{\partial z}{\partial y}\right)^2 \sin^2\theta$,

$\left(\dfrac{\partial z}{\partial \theta}\right)^2 = \left(\dfrac{\partial z}{\partial x}\right)^2 r^2\sin^2\theta - 2\dfrac{\partial z}{\partial x}\dfrac{\partial z}{\partial y}\,r^2\cos\theta\,\sin\theta + \left(\dfrac{\partial z}{\partial y}\right)^2 r^2\cos^2\theta$. Thus

$$\left(\frac{\partial z}{\partial r}\right)^2 + \frac{1}{r^2}\left(\frac{\partial z}{\partial \theta}\right)^2 = \left[\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2\right](\cos^2\theta + \sin^2\theta) = \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2.$$

50. By the Chain Rule, $\dfrac{\partial u}{\partial s} = \dfrac{\partial u}{\partial x}e^s\cos t + \dfrac{\partial u}{\partial y}e^s\sin t$, $\dfrac{\partial u}{\partial t} = \dfrac{\partial u}{\partial x}(-e^s\sin t) + \dfrac{\partial u}{\partial y}e^s\cos t$. Then

$\left(\dfrac{\partial u}{\partial s}\right)^2 = \left(\dfrac{\partial u}{\partial x}\right)^2 e^{2s}\cos^2 t + 2\dfrac{\partial u}{\partial x}\dfrac{\partial u}{\partial y}e^{2s}\cos t\,\sin t + \left(\dfrac{\partial u}{\partial y}\right)^2 e^{2s}\sin^2 t$ and

$\left(\dfrac{\partial u}{\partial t}\right)^2 = \left(\dfrac{\partial u}{\partial x}\right)^2 e^{2s}\sin^2 t - 2\dfrac{\partial u}{\partial x}\dfrac{\partial u}{\partial y}e^{2s}\cos t\,\sin t + \left(\dfrac{\partial u}{\partial y}\right)^2 e^{2s}\sin^2 t$. Thus

$$\left[\left(\frac{\partial u}{\partial s}\right)^2 + \left(\frac{\partial u}{\partial t}\right)^2\right]e^{-2s} = \left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. Let $u = x + at,\ v = x - at$. Then $z = f(u) + g(v)$, so $\partial z/\partial u = f'(u)$ and $\partial z/\partial v = g'(v)$.

Thus $\dfrac{\partial z}{\partial t} = \dfrac{\partial z}{\partial u}\dfrac{\partial u}{\partial t} + \dfrac{\partial z}{\partial v}\dfrac{\partial v}{\partial t} = af'(u) - ag'(v)$ and

$$\frac{\partial^2 z}{\partial t^2} = a\frac{\partial}{\partial t}\left[f'(u) - g'(v)\right] = a\left(\frac{df'(u)}{du}\frac{\partial u}{\partial t} - \frac{dg'(v)}{dv}\frac{\partial v}{\partial t}\right) = a^2 f''(u) + a^2 g''(v).$$

Similarly, $\dfrac{\partial z}{\partial x} = f'(u) + g'(v)$ and $\dfrac{\partial^2 z}{\partial x^2} = f''(u) + g''(v)$. Thus $\dfrac{\partial^2 z}{\partial t^2} = a^2\dfrac{\partial^2 z}{\partial x^2}$.

52. By the Chain Rule, $\dfrac{\partial u}{\partial s} = e^s\cos t\dfrac{\partial u}{\partial x} + e^s\sin t\dfrac{\partial u}{\partial y}$ and $\dfrac{\partial u}{\partial t} = -e^s\sin t\dfrac{\partial u}{\partial x} + e^s\cos t\dfrac{\partial u}{\partial y}$.

Then $\dfrac{\partial^2 u}{\partial s^2} = e^s\cos t\dfrac{\partial u}{\partial x} + e^s\cos t\dfrac{\partial}{\partial s}\left(\dfrac{\partial u}{\partial x}\right) + e^s\sin t\dfrac{\partial u}{\partial y} + e^s\sin t\dfrac{\partial}{\partial s}\left(\dfrac{\partial u}{\partial y}\right)$. But

$$\frac{\partial}{\partial s}\left(\frac{\partial u}{\partial x}\right) = \frac{\partial^2 u}{\partial x^2}\frac{\partial x}{\partial s} + \frac{\partial^2 u}{\partial y\,\partial x}\frac{\partial y}{\partial s} = e^s\cos t\frac{\partial^2 u}{\partial x^2} + e^s\sin t\frac{\partial^2 u}{\partial y\,\partial x} \text{ and}$$

$$\frac{\partial}{\partial s}\left(\frac{\partial u}{\partial y}\right) = \frac{\partial^2 u}{\partial y^2}\frac{\partial y}{\partial s} + \frac{\partial^2 u}{\partial x\,\partial y}\frac{\partial x}{\partial s} = e^s\sin t\frac{\partial^2 u}{\partial y^2} + e^s\cos t\frac{\partial^2 u}{\partial x\,\partial y}.$$

Also, by continuity of the partials, $\dfrac{\partial^2 u}{\partial x\,\partial y} = \dfrac{\partial^2 u}{\partial y\,\partial x}$. Thus

$$\begin{aligned}\frac{\partial^2 u}{\partial s^2} &= e^s\cos t\frac{\partial u}{\partial x} + e^s\cos t\left(e^s\cos t\frac{\partial^2 u}{\partial x^2} + e^s\sin t\frac{\partial^2 u}{\partial x\,\partial y}\right) + e^s\sin t\frac{\partial u}{\partial y} + e^s\sin t\left(e^s\sin t\frac{\partial^2 u}{\partial y^2} + e^s\cos t\frac{\partial^2 u}{\partial x\,\partial y}\right)\\ &= e^s\cos t\frac{\partial u}{\partial x} + e^s\sin t\frac{\partial u}{\partial y} + e^{2s}\cos^2 t\frac{\partial^2 u}{\partial x^2} + 2e^{2s}\cos t\sin t\frac{\partial^2 u}{\partial x\,\partial y} + e^{2s}\sin^2 t\frac{\partial^2 u}{\partial y^2}\end{aligned}$$

Similarly

$$\begin{aligned}\frac{\partial^2 u}{\partial t^2} &= -e^s\cos t\frac{\partial u}{\partial x} - e^s\sin t\frac{\partial}{\partial t}\left(\frac{\partial u}{\partial x}\right) - e^s\sin t\frac{\partial u}{\partial y} + e^s\cos t\frac{\partial}{\partial t}\left(\frac{\partial u}{\partial y}\right)\\ &= -e^s\cos t\frac{\partial u}{\partial x} - e^s\sin t\left(-e^s\sin t\frac{\partial^2 u}{\partial x^2} + e^s\cos t\frac{\partial^2 u}{\partial x\,\partial y}\right)\\ &\qquad -e^s\sin t\frac{\partial u}{\partial y} + e^s\cos t\left(e^s\cos t\frac{\partial^2 u}{\partial y^2} - e^s\sin t\frac{\partial^2 u}{\partial x\,\partial y}\right)\\ &= -e^s\cos t\frac{\partial u}{\partial x} - e^s\sin t\frac{\partial u}{\partial y} + e^{2s}\sin^2 t\frac{\partial^2 u}{\partial x^2} - 2e^{2s}\cos t\sin t\frac{\partial^2 u}{\partial x\,\partial y} + e^{2s}\cos^2 t\frac{\partial^2 u}{\partial y^2}\end{aligned}$$

Thus $e^{-2s}\left(\dfrac{\partial^2 u}{\partial s^2} + \dfrac{\partial^2 u}{\partial t^2}\right) = (\cos^2 t + \sin^2 t)\left(\dfrac{\partial^2 u}{\partial x^2} + \dfrac{\partial^2 u}{\partial y^2}\right) = \dfrac{\partial^2 u}{\partial x^2} + \dfrac{\partial^2 u}{\partial y^2}$, as desired.

53. $\dfrac{\partial z}{\partial s} = \dfrac{\partial z}{\partial x}2s + \dfrac{\partial z}{\partial y}2r$. Then

$$\begin{aligned}\frac{\partial^2 z}{\partial r\,\partial s} &= \frac{\partial}{\partial r}\left(\frac{\partial z}{\partial x}2s\right) + \frac{\partial}{\partial r}\left(\frac{\partial z}{\partial y}2r\right)\\ &= \frac{\partial^2 z}{\partial x^2}\frac{\partial x}{\partial r}2s + \frac{\partial}{\partial y}\left(\frac{\partial z}{\partial x}\right)\frac{\partial y}{\partial r}2s + \frac{\partial z}{\partial x}\frac{\partial}{\partial r}2s + \frac{\partial^2 z}{\partial y^2}\frac{\partial y}{\partial r}2r + \frac{\partial}{\partial x}\left(\frac{\partial z}{\partial y}\right)\frac{\partial x}{\partial r}2r + \frac{\partial z}{\partial y}2\\ &= 4rs\frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y\,\partial x}4s^2 + 0 + 4rs\frac{\partial^2 z}{\partial y^2} + \frac{\partial^2 z}{\partial x\,\partial y}4r^2 + 2\frac{\partial z}{\partial y}\end{aligned}$$

By the continuity of the partials, $\dfrac{\partial^2 z}{\partial r\partial s} = 4rs\dfrac{\partial^2 z}{\partial x^2} + 4rs\dfrac{\partial^2 z}{\partial y^2} + (4r^2 + 4s^2)\dfrac{\partial^2 z}{\partial x\,\partial y} + 2\dfrac{\partial z}{\partial y}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. By the Chain Rule,

(a) $\dfrac{\partial z}{\partial r} = \dfrac{\partial z}{\partial x}\cos\theta + \dfrac{\partial z}{\partial y}\sin\theta$ (b) $\dfrac{\partial z}{\partial \theta} = -\dfrac{\partial z}{\partial x}\, r\sin\theta + \dfrac{\partial z}{\partial y}\, r\cos\theta$

(c)
$$\begin{aligned}
\frac{\partial^2 z}{\partial r\,\partial\theta} &= \frac{\partial^2 z}{\partial\theta\,\partial r} = \frac{\partial}{\partial\theta}\left(\frac{\partial z}{\partial x}\cos\theta + \frac{\partial z}{\partial y}\sin\theta\right) = -\sin\theta\,\frac{\partial z}{\partial x} + \cos\theta\,\frac{\partial}{\partial\theta}\left(\frac{\partial z}{\partial x}\right) + \cos\theta\,\frac{\partial z}{\partial y} + \sin\theta\,\frac{\partial}{\partial\theta}\left(\frac{\partial z}{\partial y}\right)\\
&= -\sin\theta\,\frac{\partial z}{\partial x} + \cos\theta\left(\frac{\partial^2 z}{\partial x^2}\frac{\partial x}{\partial\theta} + \frac{\partial^2 z}{\partial y\,\partial x}\frac{\partial y}{\partial\theta}\right) + \cos\theta\,\frac{\partial z}{\partial y} + \sin\theta\,\frac{\partial^2 z}{\partial y^2}\frac{\partial y}{\partial\theta} + \frac{\partial^2 z}{\partial x\,\partial y}\frac{\partial x}{\partial\theta}\\
&= -\sin\theta\,\frac{\partial z}{\partial x} + \cos\theta\left(-r\sin\theta\,\frac{\partial^2 z}{\partial x^2} + r\cos\theta\,\frac{\partial^2 z}{\partial y\,\partial x}\right) + \cos\theta\,\frac{\partial z}{\partial y} + \sin\theta\left(r\cos\theta\,\frac{\partial^2 z}{\partial y^2} - r\sin\theta\,\frac{\partial^2 z}{\partial x\,\partial y}\right)\\
&= -\sin\theta\,\frac{\partial z}{\partial x} - r\cos\theta\,\sin\theta\,\frac{\partial^2 z}{\partial x^2} + r\cos^2\theta\,\frac{\partial^2 z}{\partial y\,\partial x} + \cos\theta\,\frac{\partial z}{\partial y} + r\cos\,\theta\sin\theta\,\frac{\partial^2 z}{\partial y^2} - r\sin^2\theta\,\frac{\partial^2 z}{\partial y\,\partial x}\\
&= \cos\theta\,\frac{\partial z}{\partial y} - \sin\theta\,\frac{\partial z}{\partial x} + r\cos\,\theta\sin\theta\left(\frac{\partial^2 z}{\partial y^2} - \frac{\partial^2 z}{\partial x^2}\right) + r(\cos^2\theta - \sin^2\theta)\,\frac{\partial^2 z}{\partial y\,\partial x}
\end{aligned}$$

55. $\dfrac{\partial z}{\partial r} = \dfrac{\partial z}{\partial x}\cos\theta + \dfrac{\partial z}{\partial y}\sin\theta$ and $\dfrac{\partial z}{\partial \theta} = -\dfrac{\partial z}{\partial x}\, r\sin\theta + \dfrac{\partial z}{\partial y}\, r\cos\theta$. Then

$$\begin{aligned}
\frac{\partial^2 z}{\partial r^2} &= \cos\theta\left(\frac{\partial^2 z}{\partial x^2}\cos\theta + \frac{\partial^2 z}{\partial y\,\partial x}\sin\theta\right) + \sin\theta\left(\frac{\partial^2 z}{\partial y^2}\sin\theta + \frac{\partial^2 z}{\partial x\,\partial y}\cos\theta\right)\\
&= \cos^2\theta\,\frac{\partial^2 z}{\partial x^2} + 2\cos\theta\,\sin\theta\,\frac{\partial^2 z}{\partial x\,\partial y} + \sin^2\theta\,\frac{\partial^2 z}{\partial y^2}
\end{aligned}$$

and

$$\begin{aligned}
\frac{\partial^2 z}{\partial \theta^2} &= -r\cos\theta\,\frac{\partial z}{\partial x} + (-r\sin\theta)\left(\frac{\partial^2 z}{\partial x^2}(-r\sin\theta) + \frac{\partial^2 z}{\partial y\,\partial x}\, r\cos\theta\right)\\
&\qquad -r\sin\theta\,\frac{\partial z}{\partial y} + r\cos\theta\left(\frac{\partial^2 z}{\partial y^2}\, r\cos\theta + \frac{\partial^2 z}{\partial x\,\partial y}(-r\sin\theta)\right)\\
&= -r\cos\theta\,\frac{\partial z}{\partial x} - r\sin\theta\,\frac{\partial z}{\partial y} + r^2\sin^2\theta\,\frac{\partial^2 z}{\partial x^2} - 2r^2\cos\theta\,\sin\theta\,\frac{\partial^2 z}{\partial x\,\partial y} + r^2\cos^2\theta\,\frac{\partial^2 z}{\partial y^2}
\end{aligned}$$

Thus

$$\begin{aligned}
\frac{\partial^2 z}{\partial r^2} + \frac{1}{r^2}\frac{\partial^2 z}{\partial \theta^2} + \frac{1}{r}\frac{\partial z}{\partial r} &= (\cos^2\theta + \sin^2\theta)\,\frac{\partial^2 z}{\partial x^2} + \left(\sin^2\theta + \cos^2\theta\right)\frac{\partial^2 z}{\partial y^2}\\
&\qquad -\frac{1}{r}\cos\theta\,\frac{\partial z}{\partial x} - \frac{1}{r}\sin\theta\,\frac{\partial z}{\partial y} + \frac{1}{r}\left(\cos\theta\,\frac{\partial z}{\partial x} + \sin\theta\,\frac{\partial z}{\partial y}\right)\\
&= \frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y^2} \text{ as desired.}
\end{aligned}$$

56. Since f is a polynomial, it has continuous second-order partial derivatives, and

$$f(tx,ty) = (tx)^2(ty) + 2(tx)(ty)^2 + 5(ty)^3 = t^3x^2y + 2t^3xy^2 + 5t^3y^3 = t^3(x^2y + 2xy^2 + 5y^3) = t^3 f\,(x,y).$$

Thus, f is homogeneous of degree 3.

57. (a) Differentiating both sides of $f(tx,ty) = t^n f(x,y)$ with respect to t using the Chain Rule, we get

$$\frac{\partial}{\partial t}\, f(tx,ty) = \frac{\partial}{\partial t}\,[t^n f(x,y)] \quad \Leftrightarrow$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\frac{\partial}{\partial(tx)} f(tx,ty)\cdot\frac{\partial(tx)}{\partial t}+\frac{\partial}{\partial(ty)} f(tx,ty)\cdot\frac{\partial(ty)}{\partial t}=x\frac{\partial}{\partial(tx)} f(tx,ty)+y\frac{\partial}{\partial(ty)} f(tx,ty)=nt^{n-1}f(x,y).$$

Setting $t=1$: $x\dfrac{\partial}{\partial x} f(x,y)+y\dfrac{\partial}{\partial y} f(x,y)=nf(x,y)$.

(b) Differentiating both sides of $f(tx,ty)=t^n f(x,y)$ with respect to t using the Chain Rule, we get

$$\frac{\partial}{\partial(tx)} f(tx,ty)\cdot\frac{\partial(tx)}{\partial t}+\frac{\partial}{\partial(ty)} f(tx,ty)\cdot\frac{\partial(ty)}{\partial t}=x\frac{\partial}{\partial(tx)} f(tx,ty)+y\frac{\partial}{\partial(ty)} f(tx,ty)=nt^{n-1}f(x,y)$$ and

differentiating again with respect to t gives

$$x\left[\frac{\partial^2}{\partial(tx)^2} f(tx,ty)\cdot\frac{\partial(tx)}{\partial t}+\frac{\partial^2}{\partial(ty)\,\partial(tx)} f(tx,ty)\cdot\frac{\partial(ty)}{\partial t}\right]$$
$$+\,y\left[\frac{\partial^2}{\partial(tx)\,\partial(ty)} f(tx,ty)\cdot\frac{\partial(tx)}{\partial t}+\frac{\partial^2}{\partial(ty)^2} f(tx,ty)\cdot\frac{\partial(ty)}{\partial t}\right]=n(n-1)t^{n-1}f(x,y).$$

Setting $t=1$ and using the fact that $f_{yx}=f_{xy}$, we have $x^2f_{xx}+2xyf_{xy}+y^2f_{yy}=n(n-1)f(x,y)$.

58. Differentiating both sides of $f(tx,ty)=t^n f(x,y)$ with respect to x using the Chain Rule, we get

$$\frac{\partial}{\partial x} f(tx,ty)=\frac{\partial}{\partial x}\left[t^n f(x,y)\right]\quad\Leftrightarrow$$
$$\frac{\partial}{\partial(tx)} f(tx,ty)\cdot\frac{\partial(tx)}{\partial x}+\frac{\partial}{\partial(ty)} f(tx,ty)\cdot\frac{\partial(ty)}{\partial x}=t^n\frac{\partial}{\partial x} f(x,y)\quad\Leftrightarrow\quad tf_x(tx,ty)=t^nf_x(x,y).$$

Thus $f_x(tx,ty)=t^{n-1}f_x(x,y)$.

59. $F(x,y,z)=0$ is assumed to define z as a function of x and y, that is, $z=f(x,y)$. So by (6), $\dfrac{\partial z}{\partial x}=-\dfrac{F_x}{F_z}$ since $F_z\neq 0$.

Similarly, it is assumed that $F(x,y,z)=0$ defines x as a function of y and z, that is $x=h(x,z)$. Then $F(h(y,z),y,z)=0$ and by the Chain Rule, $F_x\dfrac{\partial x}{\partial y}+F_y\dfrac{\partial y}{\partial y}+F_z\dfrac{\partial z}{\partial y}=0$. But $\dfrac{\partial z}{\partial y}=0$ and $\dfrac{\partial y}{\partial y}=1$, so $F_x\dfrac{\partial x}{\partial y}+F_y=0\quad\Rightarrow\quad\dfrac{\partial x}{\partial y}=-\dfrac{F_y}{F_x}$.

A similar calculation shows that $\dfrac{\partial y}{\partial z}=-\dfrac{F_z}{F_y}$. Thus $\dfrac{\partial z}{\partial x}\dfrac{\partial x}{\partial y}\dfrac{\partial y}{\partial z}=\left(-\dfrac{F_x}{F_z}\right)\left(-\dfrac{F_y}{F_x}\right)\left(-\dfrac{F_z}{F_y}\right)=-1$.

60. Given a function defined implicitly by $F(x,y)=0$, where F is differentiable and $F_y\neq 0$, we know that $\dfrac{dy}{dx}=-\dfrac{F_x}{F_y}$. Let $G(x,y)=-\dfrac{F_x}{F_y}$ so $\dfrac{dy}{dx}=G(x,y)$. Differentiating both sides with respect to x and using the Chain Rule gives

$$\frac{d^2y}{dx^2}=\frac{\partial G}{\partial x}\frac{dx}{dx}+\frac{\partial G}{\partial y}\frac{dy}{dx}\text{ where }\frac{\partial G}{\partial x}=\frac{\partial}{\partial x}\left(-\frac{F_x}{F_y}\right)=-\frac{F_yF_{xx}-F_xF_{yx}}{F_y^2},\ \frac{\partial G}{\partial y}=\frac{\partial}{\partial y}\left(-\frac{F_x}{F_y}\right)=-\frac{F_yF_{xy}-F_xF_{yy}}{F_y^2}.$$

Thus
$$\frac{d^2y}{dx^2}=\left(-\frac{F_yF_{xx}-F_xF_{yx}}{F_y^2}\right)(1)+\left(-\frac{F_yF_{xy}-F_xF_{yy}}{F_y^2}\right)\left(-\frac{F_x}{F_y}\right)$$
$$=-\frac{F_{xx}F_y^2-F_{yx}F_xF_y-F_{xy}F_yF_x+F_{yy}F_x^2}{F_y^3}$$

But F has continuous second derivatives, so by Clauraut's Theorem, $F_{yx}=F_{xy}$ and we have

$$\frac{d^2y}{dx^2}=-\frac{F_{xx}F_y^2-2F_{xy}F_xF_y+F_{yy}F_x^2}{F_y^3}$$ as desired.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14.6 Directional Derivatives and the Gradient Vector

1. We can approximate the directional derivative of the pressure function at K in the direction of S by the average rate of change of pressure between the points where the red line intersects the contour lines closest to K (extend the red line slightly at the left). In the direction of S, the pressure changes from 1000 millibars to 996 millibars and we estimate the distance between these two points to be approximately 50 km (using the fact that the distance from K to S is 300 km). Then the rate of change of pressure in the direction given is approximately $\frac{996-1000}{50} = -0.08$ millibar/km.

2. First we draw a line passing through Dubbo and Sydney. We approximate the directional derivative at Dubbo in the direction of Sydney by the average rate of change of temperature between the points where the line intersects the contour lines closest to Dubbo. In the direction of Sydney, the temperature changes from $30°\mathrm{C}$ to $27°\mathrm{C}$. We estimate the distance between these two points to be approximately 120 km, so the rate of change of maximum temperature in the direction given is approximately $\frac{27-30}{120} = -0.025°\mathrm{C/km}$.

3. $D_{\mathbf{u}}\, f(-20,30) = \nabla f\,(-20,30)\cdot\mathbf{u} = f_T(-20,30)\left(\frac{1}{\sqrt{2}}\right) + f_v(-20,30)\left(\frac{1}{\sqrt{2}}\right)$.

$f_T(-20,30) = \lim\limits_{h\to 0}\dfrac{f(-20+h,30)-f(-20,30)}{h}$, so we can approximate $f_T(-20,30)$ by considering $h=\pm 5$ and using the values given in the table: $f_T(-20,30) \approx \dfrac{f(-15,30)-f(-20,30)}{5} = \dfrac{-26-(-33)}{5} = 1.4$,

$f_T(-20,30) \approx \dfrac{f(-25,30)-f(-20,30)}{-5} = \dfrac{-39-(-33)}{-5} = 1.2$. Averaging these values gives $f_T(-20,30)\approx 1.3$.

Similarly, $f_v(-20,30) = \lim\limits_{h\to 0}\dfrac{f(-20,30+h)-f(-20,30)}{h}$, so we can approximate $f_v(-20,30)$ with $h=\pm 10$:

$f_v(-20,30) \approx \dfrac{f(-20,40)-f(-20,30)}{10} = \dfrac{-34-(-33)}{10} = -0.1$,

$f_v(-20,30) \approx \dfrac{f(-20,20)-f(-20,30)}{-10} = \dfrac{-30-(-33)}{-10} = -0.3$. Averaging these values gives $f_v(-20,30)\approx -0.2$.

Then $D_{\mathbf{u}} f(-20,30) \approx 1.3\left(\frac{1}{\sqrt{2}}\right) + (-0.2)\left(\frac{1}{\sqrt{2}}\right) \approx 0.778$.

4. $f(x,y) = xy^3 - x^2 \;\Rightarrow\; f_x(x,y) = y^3 - 2x$ and $f_y(x,y) = 3xy^2$. If $\mathbf{u}$ is a unit vector in the direction of $\theta = \pi/3$, then from Equation 6, $D_{\mathbf{u}} f(1,2) = f_x(1,2)\cos\left(\frac{\pi}{3}\right) + f_y(1,2)\sin\left(\frac{\pi}{3}\right) = 6\cdot\frac{1}{2} + 12\cdot\frac{\sqrt{3}}{2} = 3 + 6\sqrt{3}$.

5. $f(x,y) = y\cos(xy) \;\Rightarrow\; f_x(x,y) = y[-\sin(xy)](y) = -y^2\sin(xy)$ and $f_y(x,y) = y[-\sin(xy)](x) + [\cos(xy)](1) = \cos(xy) - xy\sin(xy)$. If $\mathbf{u}$ is a unit vector in the direction of $\theta = \pi/4$, then from Equation 6, $D_{\mathbf{u}} f(0,1) = f_x(0,1)\cos\left(\frac{\pi}{4}\right) + f_y(0,1)\sin\left(\frac{\pi}{4}\right) = 0\cdot\frac{\sqrt{2}}{2} + 1\cdot\frac{\sqrt{2}}{2} = \frac{\sqrt{2}}{2}$.

6. $f(x,y) = \sqrt{2x+3y} \;\Rightarrow\; f_x(x,y) = \frac{1}{2}(2x+3y)^{-1/2}(2) = 1/\sqrt{2x+3y}$ and $f_y(x,y) = \frac{1}{2}(2x+3y)^{-1/2}(3) = 3/\left(2\sqrt{2x+3y}\right)$. If $\mathbf{u}$ is a unit vector in the direction of $\theta = -\pi/6$, then from Equation 6, $D_{\mathbf{u}} f(3,1) = f_x(3,1)\cos\left(-\frac{\pi}{6}\right) + f_y(3,1)\sin\left(-\frac{\pi}{6}\right) = \frac{1}{3}\cdot\frac{\sqrt{3}}{2} + \frac{1}{2}\cdot\left(-\frac{1}{2}\right) = \frac{\sqrt{3}}{6} - \frac{1}{4}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. $f(x,y) = \arctan(xy) \quad \Rightarrow \quad f_x(x,y) = \dfrac{y}{1+(xy)^2}$ and $f_y(x,y) = \dfrac{x}{1+(xy)^2}$. If $\mathbf{u}$ is a unit vector in the direction $\theta = 3\pi/4$, then from Equation 6,

$$D_{\mathbf{u}} f(2,-3) = f_x(2,-3)\cos\left(\frac{3\pi}{4}\right) + f_y(2,-3)\sin\left(\frac{3\pi}{4}\right) = -\frac{3}{37}\left(-\frac{\sqrt{2}}{2}\right) + \frac{2}{37}\left(\frac{\sqrt{2}}{2}\right) = \frac{5\sqrt{2}}{74}.$$

8. $f(x,y) = x^2 e^y$

(a) $\nabla f(x,y) = \dfrac{\partial f}{\partial x}\,\mathbf{i} + \dfrac{\partial f}{\partial y}\,\mathbf{j} = 2xe^y\,\mathbf{i} + x^2e^y\,\mathbf{j}$

(b) $\nabla f(3,0) = 2(3)e^0\,\mathbf{i} + 3^2e^0\,\mathbf{j} = 6\,\mathbf{i} + 9\,\mathbf{j}$

(c) By Equation 9, $D_{\mathbf{u}} f(3,0) = \nabla f(3,0)\cdot\mathbf{u} = (6\mathbf{i} + 9\,\mathbf{j})\cdot\left(\frac{3}{5}\,\mathbf{i} - \frac{4}{5}\,\mathbf{j}\right) = \frac{18}{5} - \frac{36}{5} = -\frac{18}{5}$.

9. $f(x,y) = x/y = xy^{-1}$

(a) $\nabla f(x,y) = \dfrac{\partial f}{\partial x}\,\mathbf{i} + \dfrac{\partial f}{\partial y}\,\mathbf{j} = y^{-1}\,\mathbf{i} + (-xy^{-2})\,\mathbf{j} = \dfrac{1}{y}\,\mathbf{i} - \dfrac{x}{y^2}\,\mathbf{j}$

(b) $\nabla f(2,1) = \dfrac{1}{1}\,\mathbf{i} - \dfrac{2}{1^2}\,\mathbf{j} = \mathbf{i} - 2\,\mathbf{j}$

(c) By Equation 9, $D_{\mathbf{u}}\, f(2,1) = \nabla f(2,1)\cdot\mathbf{u} = (\mathbf{i} - 2\,\mathbf{j})\cdot\left(\frac{3}{5}\,\mathbf{i} + \frac{4}{5}\,\mathbf{j}\right) = \frac{3}{5} - \frac{8}{5} = -1$.

10. $f(x,y) = x^2 \ln y$

(a) $\nabla f(x,y) = \dfrac{\partial f}{\partial x}\,\mathbf{i} + \dfrac{\partial f}{\partial y}\,\mathbf{j} = 2x\ln y\,\mathbf{i} + (x^2/y)\,\mathbf{j}$

(b) $\nabla f(3,1) = 0\,\mathbf{i} + (9/1)\,\mathbf{j} = 9\,\mathbf{j}$

(c) By Equation 9, $D_{\mathbf{u}}\, f(3,1) = \nabla f(3,1)\cdot\mathbf{u} = 9\,\mathbf{j}\cdot\left(-\frac{5}{13}\,\mathbf{i} + \frac{12}{13}\,\mathbf{j}\right) = 0 + \frac{108}{13} = \frac{108}{13}$.

11. $f(x,y,z) = x^2yz - xyz^3$

(a) $\nabla f(x,y,z) = \langle f_x(x,y,z), f_y(x,y,z), f_z(x,y,z)\rangle = \left\langle 2xyz - yz^3, x^2z - xz^3, x^2y - 3xyz^2\right\rangle$

(b) $\nabla f(2,-1,1) = \langle -4+1, 4-2, -4+6\rangle = \langle -3,2,2\rangle$

(c) By Equation 14, $D_{\mathbf{u}} f(2,-1,1) = \nabla f(2,-1,1)\cdot\mathbf{u} = \langle -3,2,2\rangle\cdot\left\langle 0, \frac{4}{5}, -\frac{3}{5}\right\rangle = 0 + \frac{8}{5} - \frac{6}{5} = \frac{2}{5}$.

12. $f(x,y,z) = y^2e^{xyz}$

(a) $\nabla f(x,y,z) = \langle f_x(x,y,z), f_y(x,y,z), f_z(x,y,z)\rangle = \left\langle y^2e^{xyz}(yz), y^2\cdot e^{xyz}(xz) + e^{xyz}\cdot 2y, y^2e^{xyz}(xy)\right\rangle$

$= \left\langle y^3ze^{xyz}, (xy^2z + 2y)e^{xyz}, xy^3e^{xyz}\right\rangle$

(b) $\nabla f(0,1,-1) = \langle -1,2,0\rangle$

(c) $D_{\mathbf{u}} f(0,1,-1) = \nabla f(0,1,-1)\cdot\mathbf{u} = \langle -1,2,0\rangle\cdot\left\langle \frac{3}{13}, \frac{4}{13}, \frac{12}{13}\right\rangle = -\frac{3}{13} + \frac{8}{13} + 0 = \frac{5}{13}$

13. $f(x,y) = e^x\sin y \quad \Rightarrow \quad \nabla f(x,y) = \langle e^x\sin y, e^x\cos y\rangle$, $\nabla f(0,\pi/3) = \left\langle \frac{\sqrt{3}}{2}, \frac{1}{2}\right\rangle$, and a unit vector in the direction of $\mathbf{v}$ is $\mathbf{u} = \frac{1}{\sqrt{(-6)^2+8^2}}\langle -6,8\rangle = \frac{1}{10}\langle -6,8\rangle = \left\langle -\frac{3}{5}, \frac{4}{5}\right\rangle$, so

$D_{\mathbf{u}}\, f(0,\pi/3) = \nabla f(0,\pi/3)\cdot\mathbf{u} = \left\langle \frac{\sqrt{3}}{2}, \frac{1}{2}\right\rangle\cdot\left\langle -\frac{3}{5}, \frac{4}{5}\right\rangle = -\frac{3\sqrt{3}}{10} + \frac{4}{10} = \frac{4-3\sqrt{3}}{10}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $f(x,y) = \dfrac{x}{x^2+y^2} \quad \Rightarrow \quad \nabla f(x,y) = \left\langle \dfrac{(x^2+y^2)(1)-x(2x)}{(x^2+y^2)^2}, \dfrac{0-x(2y)}{(x^2+y^2)^2} \right\rangle = \left\langle \dfrac{y^2-x^2}{(x^2+y^2)^2}, -\dfrac{2xy}{(x^2+y^2)^2} \right\rangle$,

$\nabla f(1,2) = \left\langle \frac{3}{25}, -\frac{4}{25} \right\rangle$, and a unit vector in the direction of $\mathbf{v} = \langle 3,5 \rangle$ is $\mathbf{u} = \frac{1}{\sqrt{9+25}} \langle 3,5 \rangle = \left\langle \frac{3}{\sqrt{34}}, \frac{5}{\sqrt{34}} \right\rangle$, so

$D_{\mathbf{u}} f(1,2) = \nabla f(1,2) \cdot \mathbf{u} = \left\langle \frac{3}{25}, -\frac{4}{25} \right\rangle \cdot \left\langle \frac{3}{\sqrt{34}}, \frac{5}{\sqrt{34}} \right\rangle = \frac{9}{25\sqrt{34}} - \frac{20}{25\sqrt{34}} = -\frac{11}{25\sqrt{34}}$.

15. $g(s,t) = s\sqrt{t} \quad \Rightarrow \quad \nabla g(s,t) = \left(\sqrt{t}\,\right)\mathbf{i} + \left(s/(2\sqrt{t}\,)\right)\mathbf{j}$, $\nabla g(2,4) = 2\,\mathbf{i} + \frac{1}{2}\,\mathbf{j}$, and a unit vector in the direction of $\mathbf{v}$ is

$\mathbf{u} = \frac{1}{\sqrt{2^2+(-1)^2}}(2\,\mathbf{i} - \mathbf{j}) = \frac{1}{\sqrt{5}}(2\,\mathbf{i} - \mathbf{j})$, so $D_{\mathbf{u}}\, g(2,4) = \nabla g(2,4) \cdot \mathbf{u} = (2\,\mathbf{i} + \frac{1}{2}\,\mathbf{j}) \cdot \frac{1}{\sqrt{5}}(2\,\mathbf{i} - \mathbf{j}) = \frac{1}{\sqrt{5}}\left(4 - \frac{1}{2}\right) = \frac{7}{2\sqrt{5}}$

or $\frac{7\sqrt{5}}{10}$.

16. $g(u,v) = u^2 e^{-v} \quad \Rightarrow \quad \nabla g(u,v) = \left(2ue^{-v}\right)\mathbf{i} + \left(-u^2 e^{-v}\right)\mathbf{j}$, $\nabla g(3,0) = 6\,\mathbf{i} - 9\,\mathbf{j}$, and a unit vector in the direction of $\mathbf{v}$

is $\mathbf{u} = \frac{1}{\sqrt{3^2+4^2}}(3\,\mathbf{i} + 4\,\mathbf{j}) = \frac{1}{5}(3\,\mathbf{i} + 4\,\mathbf{j})$, so $D_{\mathbf{u}}\, g(3,0) = \nabla g(3,0) \cdot \mathbf{u} = (6\,\mathbf{i} - 9\,\mathbf{j}) \cdot \frac{1}{5}(3\,\mathbf{i} + 4\,\mathbf{j}) = \frac{1}{5}\,(18 - 36) = -\frac{18}{5}$.

17. $f(x,y,z) = x^2y + y^2z \quad \Rightarrow \quad \nabla f(x,y,z) = \left\langle 2xy, x^2 + 2yz, y^2 \right\rangle$, $\nabla f(1,2,3) = \langle 4,13,4 \rangle$, and a unit

vector in the direction of $\mathbf{v}$ is $\mathbf{u} = \frac{1}{\sqrt{4+1+4}}\langle 2,-1,2 \rangle = \frac{1}{3}\langle 2,-1,2 \rangle$, so

$D_{\mathbf{u}}\, f(1,2,3) = \nabla f(1,2,3) \cdot \mathbf{u} = \langle 4,13,4 \rangle \cdot \frac{1}{3}\langle 2,-1,2 \rangle = \frac{1}{3}\,(8 - 13 + 8) = \frac{3}{3} = 1$.

18. $f(x,y,z) = xy^2 \tan^{-1} z \quad \Rightarrow \quad \nabla f(x,y,z) = \left\langle y^2 \tan^{-1} z, 2xy \tan^{-1} z, \dfrac{xy^2}{1+z^2} \right\rangle$,

$\nabla f(2,1,1) = \left\langle 1 \cdot \frac{\pi}{4}, 4 \cdot \frac{\pi}{4}, \frac{2}{1+1} \right\rangle = \left\langle \frac{\pi}{4}, \pi, 1 \right\rangle$, and a unit vector in the direction of $\mathbf{v}$ is $\mathbf{u} = \frac{1}{\sqrt{1+1+1}}\langle 1,1,1 \rangle = \frac{1}{\sqrt{3}}\,\langle 1,1,1 \rangle$,

so $D_{\mathbf{u}}\, f(2,1,1) = \nabla f(2,1,1) \cdot \mathbf{u} = \left\langle \frac{\pi}{4}, \pi, 1 \right\rangle \cdot \frac{1}{\sqrt{3}}\,\langle 1,1,1 \rangle = \frac{1}{\sqrt{3}}\left(\frac{\pi}{4} + \pi + 1\right) = \frac{1}{\sqrt{3}}\left(\frac{5\pi}{4} + 1\right)$.

19. $h(r,s,t) = \ln(3r + 6s + 9t) \quad \Rightarrow \quad \nabla h(r,s,t) = \langle 3/(3r+6s+9t), 6/(3r+6s+9t), 9/(3r+6s+9t) \rangle$,

$\nabla h(1,1,1) = \left\langle \frac{1}{6}, \frac{1}{3}, \frac{1}{2} \right\rangle$, and a unit vector in the direction of $\mathbf{v} = 4\,\mathbf{i} + 12\,\mathbf{j} + 6\,\mathbf{k}$ is

$\mathbf{u} = \frac{1}{\sqrt{16+144+36}}\,(4\,\mathbf{i} + 12\,\mathbf{j} + 6\,\mathbf{k}) = \frac{2}{7}\,\mathbf{i} + \frac{6}{7}\,\mathbf{j} + \frac{3}{7}\,\mathbf{k}$, so

$D_{\mathbf{u}}\, h(1,1,1) = \nabla h(1,1,1) \cdot \mathbf{u} = \left\langle \frac{1}{6}, \frac{1}{3}, \frac{1}{2} \right\rangle \cdot \left\langle \frac{2}{7}, \frac{6}{7}, \frac{3}{7} \right\rangle = \frac{1}{21} + \frac{2}{7} + \frac{3}{14} = \frac{23}{42}$.

20. $D_{\mathbf{u}} f(2,2) = \nabla f(2,2) \cdot \mathbf{u}$, the scalar projection of $\nabla f(2,2)$ onto $\mathbf{u}$, so we draw a perpendicular from the tip of $\nabla f(2,2)$ to the line containing $\mathbf{u}$. We can use the point $(2,2)$ to determine the scale of the axes, and we estimate the length of the projection to be approximately 3.0 units. Since the angle between $\nabla f(2,2)$ and $\mathbf{u}$ is greater than $90°$, the scalar projection is negative. Thus $D_{\mathbf{u}}\, f(2,2) \approx -3$.

21. $f(x,y) = x^2y^2 - y^3 \quad \Rightarrow \quad \nabla f(x,y) = \left\langle 2xy^2, 2x^2y - 3y^2 \right\rangle$, so $\nabla f(1,2) = \langle 8,-8 \rangle$. The unit vector in the direction of $\overrightarrow{PQ} = \langle -3-1, 5-2 \rangle = \langle -4,3 \rangle$ is $\mathbf{u} = \frac{1}{\sqrt{(-4)^2+3^2}}\,\langle -4,3 \rangle = \left\langle -\frac{4}{5}, \frac{3}{5} \right\rangle$, so

$D_{\mathbf{u}} f(1,2) = \nabla f(1,2) \cdot \mathbf{u} = \langle 8,-8 \rangle \cdot \left\langle -\frac{4}{5}, \frac{3}{5} \right\rangle = -\frac{56}{5}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $f(x,y) = \dfrac{x}{y^2} \quad \Rightarrow \quad \nabla f(x,y) = \left\langle \dfrac{1}{y^2}, -\dfrac{2x}{y^3} \right\rangle$, so $\nabla f(3,-1) = \langle 1, 6\rangle$. The unit vector in the direction of

$\overrightarrow{PQ} = \langle -2-3, 11-(-1)\rangle = \langle -5, 12\rangle$ is $\mathbf{u} = \frac{1}{\sqrt{(-5)^2+12^2}}\langle -5, 12\rangle = \left\langle -\frac{5}{13}, \frac{12}{13}\right\rangle$, so

$D_{\mathbf{u}} f(3,-1) = \nabla f(3,-1)\cdot \mathbf{u} = \langle 1, 6\rangle \cdot \left\langle -\frac{5}{13}, \frac{12}{13}\right\rangle = \frac{67}{13}$.

23. $f(x,y) = \sqrt{xy} \quad \Rightarrow \quad \nabla f(x,y) = \left\langle \frac{1}{2}(xy)^{-1/2}(y), \frac{1}{2}(xy)^{-1/2}(x)\right\rangle = \left\langle \dfrac{y}{2\sqrt{xy}}, \dfrac{x}{2\sqrt{xy}}\right\rangle$, so $\nabla f(2,8) = \left\langle 1, \frac{1}{4}\right\rangle$.

The unit vector in the direction of $\overrightarrow{PQ} = \langle 5-2, 4-8\rangle = \langle 3, -4\rangle$ is $\mathbf{u} = \frac{1}{\sqrt{3^2+(-4)^2}}\langle 3,-4\rangle = \left\langle \frac{3}{5}, -\frac{4}{5}\right\rangle$,

so $D_{\mathbf{u}}\, f(2,8) = \nabla f(2,8)\cdot \mathbf{u} = \left\langle 1, \frac{1}{4}\right\rangle \cdot \left\langle \frac{3}{5}, -\frac{4}{5}\right\rangle = \frac{2}{5}$.

24. $f(x,y,z) = xy^2z^3 \quad \Rightarrow \quad \nabla f(x,y,z) = \left\langle y^2z^3, 2xyz^3, 3xy^2z^2\right\rangle$, so $\nabla f(2,1,1) = \langle 1,4,6\rangle$. The unit vector in the direction of $\overrightarrow{PQ} = \langle 0-2, -3-1, 5-1\rangle = \langle -2,-4,4\rangle$ is $\mathbf{u} = \frac{1}{\sqrt{(-2)^2+(-4)^2+4^2}}\langle -2,-4,4\rangle = \frac{1}{6}\langle -2,-4,4\rangle$, so

$D_{\mathbf{u}}\, f(2,1,1) = \nabla f(2,1,1)\cdot \mathbf{u} = \langle 1,4,6\rangle \cdot \frac{1}{6}\langle -2,-4,4\rangle = \frac{1}{6}(-2-16+24) = 1$.

25. $f(x,y,z) = xy - xy^2z^2 \quad \Rightarrow \quad \nabla f(x,y,z) = \left\langle y - y^2z^2, x - 2xyz^2, -2xy^2z\right\rangle$, so $\nabla f(2,-1,1) = \langle -2,6,-4\rangle$.

The unit vector in the direction of $\overrightarrow{PQ} = \langle 5-2, 1-(-1), 7-1\rangle = \langle 3,2,6\rangle$ is $\mathbf{u} = \frac{1}{\sqrt{3^2+2^2+6^2}}\langle 3,2,6\rangle = \left\langle \frac{3}{7}, \frac{2}{7}, \frac{6}{7}\right\rangle$, so

$D_{\mathbf{u}} f(2,-1,1) = \nabla f(2,-1,1)\cdot \mathbf{u} = \langle -2,6,-4\rangle \cdot \left\langle \frac{3}{7}, \frac{2}{7}, \frac{6}{7}\right\rangle = -\frac{18}{7}$.

26.

Note that the vectors drawn at P, Q, and R are perpendicular to the curves and represent the direction of maximum increase.

27. $f(x,y) = 5xy^2 \quad \Rightarrow \quad \nabla f(x,y) = \left\langle 5y^2, 10xy\right\rangle$. Then $\nabla f(3,-2) = \langle 20,-60\rangle$, or equivalently, $\langle 1,-3\rangle$ is the direction of maximum rate of change, and the maximum rate of change is $|\nabla f(3,-2)| = \sqrt{20^2+(-60)^2} = \sqrt{4000} = 20\sqrt{10}$.

28. $f(s,t) = \dfrac{s}{s^2+t^2} = s(s^2+t^2)^{-1} \quad \Rightarrow$

$\nabla f(s,t) = \left\langle s(-1)(s^2+t^2)^{-2}(2s) + (1)(s^2+t^2)^{-1},\ s(-1)(s^2+t^2)^{-2}(2t)\right\rangle = \left\langle \dfrac{t^2-s^2}{(s^2+t^2)^2}, -\dfrac{2st}{(s^2+t^2)^2}\right\rangle$.

Then $\nabla f(-1,1) = \left\langle 0, \frac{1}{2}\right\rangle$ is the direction of maximum rate of change, and the maximum rate of change is $|\nabla f(-1,1)| = \frac{1}{2}$.

29. $f(x,y) = \sin(xy) \quad \Rightarrow \quad \nabla f(x,y) = \langle y\cos(xy), x\cos(xy)\rangle$, $\nabla f(1,0) = \langle 0,1\rangle$. Thus, the maximum rate of change is $|\nabla f(1,0)| = 1$ in the direction $\langle 0,1\rangle$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $f(x,y,z) = x\ln(yz) \quad \Rightarrow \quad \nabla f(x,y,z) = \left\langle \ln(yz), x\cdot\dfrac{z}{yz}, x\cdot\dfrac{y}{yz}\right\rangle = \left\langle \ln(yz), \dfrac{x}{y}, \dfrac{x}{z}\right\rangle, \quad \nabla f(1,2,\tfrac{1}{2}) = \langle 0, \tfrac{1}{2}, 2\rangle.$

Thus, the maximum rate of change is $\left|\nabla f(1,2,\tfrac{1}{2})\right| = \sqrt{0+\tfrac{1}{4}+4} = \sqrt{\tfrac{17}{4}} = \tfrac{\sqrt{17}}{2}$ in the direction $\langle 0, \tfrac{1}{2}, 2\rangle$, or equivalently, $\langle 0, 1, 4\rangle$.

31. $f(x,y,z) = x/(y+z) = x(y+z)^{-1} \quad \Rightarrow$

$\nabla f(x,y,z) = \langle 1/(y+z), -x(y+z)^{-2}(1), -x(y+z)^{-2}(1)\rangle = \left\langle \dfrac{1}{y+z}, -\dfrac{x}{(y+z)^2}, -\dfrac{x}{(y+z)^2}\right\rangle,$

$\nabla f(8,1,3) = \langle \tfrac{1}{4}, -\tfrac{8}{4^2}, -\tfrac{8}{4^2}\rangle = \langle \tfrac{1}{4}, -\tfrac{1}{2}, -\tfrac{1}{2}\rangle$. Thus, the maximum rate of change is

$|\nabla f(8,1,3)| = \sqrt{\tfrac{1}{16}+\tfrac{1}{4}+\tfrac{1}{4}} = \sqrt{\tfrac{9}{16}} = \tfrac{3}{4}$ in the direction $\langle \tfrac{1}{4}, -\tfrac{1}{2}, -\tfrac{1}{2}\rangle$, or equivalently, $\langle 1, -2, -2\rangle$.

32. $f(p,q,r) = \arctan(pqr) \quad \Rightarrow \quad \nabla f(p,q,r) = \left\langle \dfrac{qr}{1+(pqr)^2}, \dfrac{pr}{1+(pqr)^2}, \dfrac{pq}{1+(pqr)^2}\right\rangle, \nabla f(1,2,1) = \langle \tfrac{2}{5}, \tfrac{1}{5}, \tfrac{2}{5}\rangle.$

Thus, the maximum rate of change is $|\nabla f(1,2,1)| = \sqrt{\tfrac{4}{25}+\tfrac{1}{25}+\tfrac{4}{25}} = \sqrt{\tfrac{9}{25}} = \tfrac{3}{5}$ in the direction $\langle \tfrac{2}{5}, \tfrac{1}{5}, \tfrac{2}{5}\rangle$, or equivalently, $\langle 2, 1, 2\rangle$.

33. (a) As in the proof of Theorem 15, $D_{\mathbf{u}}f = |\nabla f|\cos\theta$. Since the minimum value of $\cos\theta$ is -1, occurring when $\theta = \pi$, the maximum rate of decrease of $D_{\mathbf{u}}f$ is $-|\nabla f|$ occurring when $\theta = \pi$; that is, when $\mathbf{u}$ is in the opposite direction of ∇f (assuming $\nabla f \neq \mathbf{0}$).

(b) $f(x,y) = x^4y - x^2y^3 \quad \Rightarrow \quad \nabla f(x,y) = \langle 4x^3y - 2xy^3, x^4 - 3x^2y^2\rangle$, so f decreases fastest at the point $(2,-3)$ in the direction $-\nabla f(2,-3) = -\langle 12, -92\rangle = \langle -12, 92\rangle$. The maximum rate of decrease is $-|\nabla f(2,-3)| = -|\langle 12,-92\rangle| = -\sqrt{12^2+(-92)^2} = -\sqrt{8608} = -4\sqrt{538}$.

34. $f(x,y) = x^2 + xy^3 \quad \Rightarrow \quad \nabla f(x,y) = \langle 2x+y^3, 3xy^2\rangle$ so $\nabla f(2,1) = \langle 5, 6\rangle$. If $\mathbf{u} = \langle a, b\rangle$ is a unit vector in the desired direction then $D_{\mathbf{u}}f(2,1) = 2 \quad \Leftrightarrow \quad \langle 5,6\rangle\cdot\langle a,b\rangle = 2 \quad \Leftrightarrow \quad 5a+6b = 2 \quad \Leftrightarrow \quad b = \tfrac{1}{3} - \tfrac{5}{6}a$. But $a^2+b^2 = 1 \quad \Leftrightarrow$ $a^2 + \left(\tfrac{1}{3} - \tfrac{5}{6}a\right)^2 = 1 \quad \Leftrightarrow \quad \tfrac{61}{36}a^2 - \tfrac{5}{9}a + \tfrac{1}{9} = 1 \quad \Leftrightarrow \quad 61a^2 - 20a - 32 = 0$. By the quadratic formula, the solutions are $a = \dfrac{-(-20)\pm\sqrt{(-20)^2-4(61)(-32)}}{2(61)} = \dfrac{20\pm\sqrt{8208}}{122} = \dfrac{10\pm 6\sqrt{57}}{61}$. If $a = \dfrac{10+6\sqrt{57}}{61} \approx 0.9065$ then $b = \dfrac{1}{3} - \dfrac{5}{6}\left(\dfrac{10+6\sqrt{57}}{61}\right) \approx -0.4221$, and if $a = \dfrac{10-6\sqrt{57}}{61} \approx -0.5787$ then $b = \dfrac{1}{3} - \dfrac{5}{6}\left(\dfrac{10-6\sqrt{57}}{61}\right) \approx 0.8156$.

Thus the two directions giving a directional derivative of 2 are approximately $\langle 0.9065, -0.4221\rangle$ and $\langle -0.5787, 0.8156\rangle$.

35. For $f(x,y) = x^2+y^2-2x-4y$, the direction of greatest rate of change is $\nabla f(x,y) = (2x-2)\,\mathbf{i} + (2y-4)\,\mathbf{j}$, so we need to find all points (x,y) where $\nabla f(x,y)$ is parallel to $\mathbf{i}+\mathbf{j} \quad \Leftrightarrow \quad (2x-2)\,\mathbf{i} + (2y-4)\,\mathbf{j} = c\,(\mathbf{i}+\mathbf{j}) \quad \Leftrightarrow \quad c = 2x-2$ and $c = 2y-4$. Then $2x-2 = 2y-4 \quad \Rightarrow \quad y = x+1$, so at all points on the line $y = x+1$, the direction of greatest rate of change of f is $\mathbf{i}+\mathbf{j}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. The fisherman is traveling in the direction $\langle -80, -60\rangle$. A unit vector in this direction is $\mathbf{u} = \frac{1}{100}\langle -80, -60\rangle = \left\langle -\frac{4}{5}, -\frac{3}{5}\right\rangle$, and if the depth of the lake is given by $f(x, y) = 200 + 0.02x^2 - 0.001y^3$, then $\nabla f(x, y) = \left\langle 0.04x, -0.003y^2\right\rangle$. $D_{\mathbf{u}} f(80, 60) = \nabla f(80, 60) \cdot \mathbf{u} = \langle 3.2, -10.8\rangle \cdot \left\langle -\frac{4}{5}, -\frac{3}{5}\right\rangle = 3.92$. Since $D_{\mathbf{u}} f(80, 60)$ is positive, the depth of the lake is increasing near $(80, 60)$ in the direction toward the buoy.

37. $T = \dfrac{k}{\sqrt{x^2 + y^2 + z^2}}$ and $120 = T(1, 2, 2) = \dfrac{k}{3}$ so $k = 360$.

(a) $\mathbf{u} = \dfrac{\langle 1, -1, 1\rangle}{\sqrt{3}}$,

$$D_{\mathbf{u}}T(1, 2, 2) = \nabla T(1, 2, 2) \cdot \mathbf{u} = \left[-360\left(x^2 + y^2 + z^2\right)^{-3/2}\langle x, y, z\rangle\right]_{(1,2,2)} \cdot \mathbf{u} = -\tfrac{40}{3}\langle 1, 2, 2\rangle \cdot \tfrac{1}{\sqrt{3}}\langle 1, -1, 1\rangle = -\tfrac{40}{3\sqrt{3}}$$

(b) From (a), $\nabla T = -360\left(x^2 + y^2 + z^2\right)^{-3/2}\langle x, y, z\rangle$, and since $\langle x, y, z\rangle$ is the position vector of the point (x, y, z), the vector $-\langle x, y, z\rangle$, and thus ∇T, always points toward the origin.

38. $\nabla T = -400e^{-x^2-3y^2-9z^2}\langle x, 3y, 9z\rangle$

(a) $\mathbf{u} = \frac{1}{\sqrt{6}}\langle 1, -2, 1\rangle$, $\nabla T(2, -1, 2) = -400e^{-43}\langle 2, -3, 18\rangle$ and

$$D_{\mathbf{u}}\, T(2, -1, 2) = \left(-\frac{400e^{-43}}{\sqrt{6}}\right)(26) = -\frac{5200\sqrt{6}}{3e^{43}}\ {}^\circ\mathrm{C/m}.$$

(b) $\nabla T(2, -1, 2) = 400e^{-43}\langle -2, 3, -18\rangle$ or equivalently $\langle -2, 3, -18\rangle$.

(c) $|\nabla T| = 400e^{-x^2-3y^2-9z^2}\sqrt{x^2 + 9y^2 + 81z^2}\ {}^\circ\mathrm{C/m}$ is the maximum rate of increase. At $(2, -1, 2)$ the maximum rate of increase is $400e^{-43}\sqrt{337}\ {}^\circ\mathrm{C/m}$.

39. $\nabla V(x, y, z) = \langle 10x - 3y + yz, xz - 3x, xy\rangle$, $\nabla V(3, 4, 5) = \langle 38, 6, 12\rangle$

(a) $D_{\mathbf{u}}\, V(3, 4, 5) = \langle 38, 6, 12\rangle \cdot \frac{1}{\sqrt{3}}\langle 1, 1, -1\rangle = \frac{32}{\sqrt{3}}$

(b) $\nabla V(3, 4, 5) = \langle 38, 6, 12\rangle$, or equivalently, $\langle 19, 3, 6\rangle$.

(c) $|\nabla V(3, 4, 5)| = \sqrt{38^2 + 6^2 + 12^2} = \sqrt{1624} = 2\sqrt{406}$

40. $z = f(x, y) = 1000 - 0.005x^2 - 0.01y^2 \quad \Rightarrow \quad \nabla f(x, y) = \langle -0.01x, -0.02y\rangle$ and $\nabla f(60, 40) = \langle -0.6, -0.8\rangle$.

(a) Due south is in the direction of the unit vector $\mathbf{u} = -\mathbf{j}$ and $D_{\mathbf{u}} f(60, 40) = \nabla f\,(60, 40) \cdot \langle 0, -1\rangle = \langle -0.6, -0.8\rangle \cdot \langle 0, -1\rangle = 0.8$. Thus, if you walk due south from $(60, 40, 966)$ you will ascend at a rate of 0.8 vertical meters per horizontal meter.

(b) Northwest is in the direction of the unit vector $\mathbf{u} = \frac{1}{\sqrt{2}}\langle -1, 1\rangle$ and $D_{\mathbf{u}} f(60, 40) = \nabla f\,(60, 40) \cdot \frac{1}{\sqrt{2}}\langle -1, 1\rangle = \langle -0.6, -0.8\rangle \cdot \frac{1}{\sqrt{2}}\langle -1, 1\rangle = -\frac{0.2}{\sqrt{2}} \approx -0.14$. Thus, if you walk northwest from $(60, 40, 966)$ you will descend at a rate of approximately 0.14 vertical meters per horizontal meter.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\nabla f(60, 40) = \langle -0.6, -0.8 \rangle$ is the direction of largest slope with a rate of ascent given by

$|\nabla f(60, 40)| = \sqrt{(-0.6)^2 + (-0.8)^2} = 1$. The angle above the horizontal in which the path begins is given by

$\tan\theta = 1 \quad \Rightarrow \quad \theta = 45°$.

41. A unit vector in the direction of $\overrightarrow{AB}$ is $\mathbf{i}$ and a unit vector in the direction of $\overrightarrow{AC}$ is $\mathbf{j}$. Thus $D_{\overrightarrow{AB}} f(1, 3) = f_x(1, 3) = 3$ and $D_{\overrightarrow{AC}} f(1, 3) = f_y(1, 3) = 26$. Therefore $\nabla f(1, 3) = \langle f_x(1, 3), f_y(1, 3) \rangle = \langle 3, 26 \rangle$, and by definition, $D_{\overrightarrow{AD}} f(1, 3) = \nabla f \cdot \mathbf{u}$ where $\mathbf{u}$ is a unit vector in the direction of $\overrightarrow{AD}$, which is $\left\langle \frac{5}{13}, \frac{12}{13} \right\rangle$. Therefore, $D_{\overrightarrow{AD}} f\,(1, 3) = \langle 3, 26 \rangle \cdot \left\langle \frac{5}{13}, \frac{12}{13} \right\rangle = 3 \cdot \frac{5}{13} + 26 \cdot \frac{12}{13} = \frac{327}{13}$.

42. The curves of steepest ascent or descent are perpendicular to all of the contour lines (see Figure 13) so we sketch curves beginning at A and B that head toward lower elevations, crossing each contour line at a right angle.

43. (a) $\nabla(au + bv) = \left\langle \dfrac{\partial(au+bv)}{\partial x}, \dfrac{\partial(au+bv)}{\partial y} \right\rangle = \left\langle a\dfrac{\partial u}{\partial x} + b\dfrac{\partial v}{\partial x}, a\dfrac{\partial u}{\partial y} + b\dfrac{\partial v}{\partial y} \right\rangle = a\left\langle \dfrac{\partial u}{\partial x}, \dfrac{\partial u}{\partial y} \right\rangle + b\left\langle \dfrac{\partial v}{\partial x}, \dfrac{\partial v}{\partial y} \right\rangle$

$= a\,\nabla u + b\,\nabla v$

(b) $\nabla(uv) = \left\langle v\dfrac{\partial u}{\partial x} + u\dfrac{\partial v}{\partial x}, v\dfrac{\partial u}{\partial y} + u\dfrac{\partial v}{\partial y} \right\rangle = v\left\langle \dfrac{\partial u}{\partial x}, \dfrac{\partial u}{\partial y} \right\rangle + u\left\langle \dfrac{\partial v}{\partial x}, \dfrac{\partial v}{\partial y} \right\rangle = v\,\nabla u + u\,\nabla v$

(c) $\nabla\left(\dfrac{u}{v}\right) = \left\langle \dfrac{v\dfrac{\partial u}{\partial x} - u\dfrac{\partial v}{\partial x}}{v^2}, \dfrac{v\dfrac{\partial u}{\partial y} - u\dfrac{\partial v}{\partial y}}{v^2} \right\rangle = \dfrac{v\left\langle \dfrac{\partial u}{\partial x}, \dfrac{\partial u}{\partial y} \right\rangle - u\left\langle \dfrac{\partial v}{\partial x}, \dfrac{\partial v}{\partial y} \right\rangle}{v^2} = \dfrac{v\,\nabla u - u\,\nabla v}{v^2}$

(d) $\nabla u^n = \left\langle \dfrac{\partial(u^n)}{\partial x}, \dfrac{\partial(u^n)}{\partial y} \right\rangle = \left\langle nu^{n-1}\dfrac{\partial u}{\partial x}, nu^{n-1}\dfrac{\partial u}{\partial y} \right\rangle = nu^{n-1}\,\nabla u$

44. If we place the initial point of the gradient vector $\nabla f(4, 6)$ at $(4, 6)$, the vector is perpendicular to the level curve of f that includes $(4, 6)$, so we sketch a portion of the level curve through $(4, 6)$ (using the nearby level curves as a guideline) and draw a line perpendicular to the curve at $(4, 6)$. The gradient vector is parallel to this line, pointing in the direction of increasing

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

function values, and with length equal to the maximum value of the directional derivative of f at $(4, 6)$. We can estimate this length by finding the average rate of change in the direction of the gradient. The line intersects the contour lines corresponding to -2 and -3 with an estimated distance of 0.5 units. Thus the rate of change is approximately $\dfrac{-2-(-3)}{0.5} = 2$, and we sketch the gradient vector with length 2.

45. $f(x, y) = x^3 + 5x^2y + y^3 \quad \Rightarrow$

$D_{\mathbf{u}}f(x, y) = \nabla f(x, y) \cdot \mathbf{u} = \left\langle 3x^2 + 10xy, 5x^2 + 3y^2 \right\rangle \cdot \left\langle \frac{3}{5}, \frac{4}{5} \right\rangle = \frac{9}{5}x^2 + 6xy + 4x^2 + \frac{12}{5}y^2 = \frac{29}{5}x^2 + 6xy + \frac{12}{5}y^2$.

Then

$$\begin{aligned} D_{\mathbf{u}}^2 f(x, y) &= D_{\mathbf{u}}\left[D_{\mathbf{u}}f(x, y)\right] = \nabla\left[D_{\mathbf{u}}f(x, y)\right] \cdot \mathbf{u} = \left\langle \tfrac{58}{5}x + 6y, 6x + \tfrac{24}{5}y \right\rangle \cdot \left\langle \tfrac{3}{5}, \tfrac{4}{5} \right\rangle \\ &= \tfrac{174}{25}x + \tfrac{18}{5}y + \tfrac{24}{5}x + \tfrac{96}{25}y = \tfrac{294}{25}x + \tfrac{186}{25}y \end{aligned}$$

and $D_{\mathbf{u}}^2 f(2, 1) = \frac{294}{25}(2) + \frac{186}{25}(1) = \frac{774}{25}$.

46. (a) From Equation 9 we have $D_{\mathbf{u}}f = \nabla f \cdot \mathbf{u} = \langle f_x, f_y \rangle \cdot \langle a, b \rangle = f_x a + f_y b$ and from Exercise 39 we have

$$\begin{aligned} D_{\mathbf{u}}^2 f &= D_{\mathbf{u}}\left[D_{\mathbf{u}}f\right] = \nabla\left[D_{\mathbf{u}}f\right] \cdot \mathbf{u} \\ &= \langle f_{xx}a + f_{yx}b, f_{xy}a + f_{yy}b \rangle \cdot \langle a, b \rangle \\ &= f_{xx}a^2 + f_{yx}ab + f_{xy}ab + f_{yy}b^2 \end{aligned}$$

But $f_{yx} = f_{xy}$ by Clairaut's Theorem, so $D_{\mathbf{u}}^2 f = f_{xx}a^2 + 2f_{xy}ab + f_{yy}b^2$.

(b) $f(x, y) = xe^{2y} \quad \Rightarrow \quad f_x = e^{2y}$, $f_y = 2xe^{2y}$, $f_{xx} = 0$, $f_{xy} = 2e^{2y}$, $f_{yy} = 4xe^{2y}$ and a unit vector in the direction of $\mathbf{v}$ is $\mathbf{u} = \frac{1}{\sqrt{4^2+6^2}}\langle 4, 6 \rangle = \left\langle \frac{2}{\sqrt{13}}, \frac{3}{\sqrt{13}} \right\rangle = \langle a, b \rangle$. Then

$$\begin{aligned} D_{\mathbf{u}}^2 f &= f_{xx}a^2 + 2f_{xy}ab + f_{yy}b^2 \\ &= 0 \cdot \left(\tfrac{2}{\sqrt{13}}\right)^2 + 2 \cdot 2e^{2y}\left(\tfrac{2}{\sqrt{13}}\right)\left(\tfrac{3}{\sqrt{13}}\right) + 4xe^{2y}\left(\tfrac{3}{\sqrt{13}}\right)^2 = \tfrac{24}{13}e^{2y} + \tfrac{36}{13}xe^{2y} \end{aligned}$$

47. Let $F(x, y, z) = 2(x-2)^2 + (y-1)^2 + (z-3)^2$. Then $2(x-2)^2 + (y-1)^2 + (z-3)^2 = 10$ is a level surface of F.

$F_x(x, y, z) = 4(x-2) \quad \Rightarrow \quad F_x(3, 3, 5) = 4$, $F_y(x, y, z) = 2(y-1) \quad \Rightarrow \quad F_y(3, 3, 5) = 4$, and

$F_z(x, y, z) = 2(z-3) \quad \Rightarrow \quad F_z(3, 3, 5) = 4$.

(a) Equation 19 gives an equation of the tangent plane at $(3, 3, 5)$ as $4(x-3) + 4(y-3) + 4(z-5) = 0 \quad \Leftrightarrow$ $4x + 4y + 4z = 44$ or equivalently $x + y + z = 11$.

(b) By Equation 20, the normal line has symmetric equations $\dfrac{x-3}{4} = \dfrac{y-3}{4} = \dfrac{z-5}{4}$ or equivalently $x - 3 = y - 3 = z - 5$. Corresponding parametric equations are $x = 3 + t$, $y = 3 + t$, $z = 5 + t$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. Let $F(x, y, z) = y^2 + z^2 - x$. Then $x = y^2 + z^2 + 1 \quad \Leftrightarrow \quad y^2 + z^2 - x = -1$ is a level surface of F.

$F_x(x, y, z) = -1 \quad \Rightarrow \quad F_x(3, 1, -1) = -1$, $F_y(x, y, z) = 2y \quad \Rightarrow \quad F_y(3, 1, -1) = 2$, and $F_z(x, y, z) = 2z \quad \Rightarrow$ $F_z(3, 1, -1) = -2$.

(a) By Equation 19, an equation of the tangent plane at $(3, 1, -1)$ is $(-1)(x - 3) + 2(y - 1) + (-2)[z - (-1)] = 0$ or $-x + 2y - 2z = 1$ or $x - 2y + 2z = -1$.

(b) By Equation 20, the normal line has symmetric equations $\dfrac{x-3}{-1} = \dfrac{y-1}{2} = \dfrac{z-(-1)}{-2}$ or equivalently $x - 3 = \dfrac{y-1}{-2} = \dfrac{z+1}{2}$ and parametric equations $x = 3 - t$, $y = 1 + 2t$, $z = -1 - 2t$.

49. Let $F(x, y, z) = xy^2z^3$. Then $xy^2z^3 = 8$ is a level surface of F and $\nabla F(x, y, z) = \langle y^2z^3, 2xyz^3, 3xy^2z^2 \rangle$.

(a) $\nabla F(2, 2, 1) = \langle 4, 8, 24 \rangle$ is a normal vector for the tangent plane at $(2, 2, 1)$, so an equation of the tangent plane is $4(x - 2) + 8(y - 2) + 24(z - 1) = 0$ or $4x + 8y + 24z = 48$ or equivalently $x + 2y + 6z = 12$.

(b) The normal line has direction $\nabla F(2, 2, 1) = \langle 4, 8, 24 \rangle$ or equivalently $\langle 1, 2, 6 \rangle$, so parametric equations are $x = 2 + t$, $y = 2 + 2t$, $z = 1 + 6t$, and symmetric equations are $x - 2 = \dfrac{y-2}{2} = \dfrac{z-1}{6}$.

50. Let $F(x, y, z) = xy + yz + zx$. Then $xy + yz + zx = 5$ is a level surface of F and $\nabla F(x, y, z) = \langle y + z, x + z, x + y \rangle$.

(a) $\nabla F(1, 2, 1) = \langle 3, 2, 3 \rangle$ is a normal vector for the tangent plane at $(1, 2, 1)$, so an equation of the tangent plane is $3(x - 1) + 2(y - 2) + 3(z - 1) = 0$ or $3x + 2y + 3z = 10$.

(b) The normal line has direction $\langle 3, 2, 3 \rangle$, so parametric equations are $x = 1 + 3t$, $y = 2 + 2t$, $z = 1 + 3t$, and symmetric equations are $\dfrac{x-1}{2} = \dfrac{y-2}{1} = \dfrac{z-1}{3}$.

51. Let $F(x, y, z) = x + y + z - e^{xyz}$. Then $x + y + z = e^{xyz}$ is the level surface $F(x, y, z) = 0$, and $\nabla F(x, y, z) = \langle 1 - yze^{xyz}, 1 - xze^{xyz}, 1 - xye^{xyz} \rangle$.

(a) $\nabla F(0, 0, 1) = \langle 1, 1, 1 \rangle$ is a normal vector for the tangent plane at $(0, 0, 1)$, so an equation of the tangent plane is $1(x - 0) + 1(y - 0) + 1(z - 1) = 0$ or $x + y + z = 1$.

(b) The normal line has direction $\langle 1, 1, 1 \rangle$, so parametric equations are $x = t$, $y = t$, $z = 1 + t$, and symmetric equations are $x = y = z - 1$.

52. Let $F(x, y, z) = x^4 + y^4 + z^4 - 3x^2y^2z^2$. Then $x^4 + y^4 + z^4 = 3x^2y^2z^2$ is the level surface $F(x, y, z) = 0$, and $\nabla F(x, y, z) = \langle 4x^3 - 6xy^2z^2, 4y^3 - 6x^2yz^2, 4z^3 - 6x^2y^2z \rangle$.

(a) $\nabla F(1, 1, 1) = \langle -2, -2, -2 \rangle$ or equivalently $\langle 1, 1, 1 \rangle$ is a normal vector for the tangent plane at $(1, 1, 1)$, so an equation of the tangent plane is $1(x - 1) + 1(y - 1) + 1(z - 1) = 0$ or $x + y + z = 3$.

(b) The normal line has direction $\langle 1, 1, 1 \rangle$, so parametric equations are $x = 1 + t$, $y = 1 + t$, $z = 1 + t$, and symmetric equations are $x - 1 = y - 1 = z - 1$ or equivalently $x = y = z$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

53. $F(x,y,z) = xy + yz + zx \quad \Rightarrow \quad xy + yz + zx = 3$ is the level surface $F(x,y,z) = 3$. $\nabla F(x,y,z) = \langle y+z, x+z, y+x \rangle \quad \Rightarrow$ $\nabla F(1,1,1) = \langle 2,2,2 \rangle$, and an equation of the tangent plane is $2x + 2y + 2z = 6$ or $x + y + z = 3$. The normal line is given by $x - 1 = y - 1 = z - 1$ or $x = y = z$. To graph the surface we solve for z: $z = \dfrac{3 - xy}{x + y}$.

54. $F(x,y,z) = xyz \quad \Rightarrow \quad xyz = 6$ is the level surface $F(x,y,z) = 6$. $\nabla F(x,y,z) = \langle yz, xz, yx \rangle \quad \Rightarrow \quad \nabla F(1,2,3) = \langle 6,3,2 \rangle$, and an equation of the tangent plane is $6x + 3y + 2z = 18$. The normal line is given by $\dfrac{x-1}{6} = \dfrac{y-2}{3} = \dfrac{z-3}{2}$ or $x = 1 + 6t,\ y = 2 + 3t,$ $z = 3 + 2t$. To graph the surface we solve for z: $z = \dfrac{6}{xy}$.

55. $f(x,y) = xy \quad \Rightarrow \quad \nabla f(x,y) = \langle y, x \rangle$ and $\nabla f(3,2) = \langle 2,3 \rangle$. Since $\nabla f(3,2)$ is perpendicular to the tangent line, the tangent line has equation $\nabla f(3,2) \cdot \langle x-3, y-2 \rangle = 0 \quad \Rightarrow \quad \langle 2,3 \rangle \cdot \langle x-3, x-2 \rangle = 0 \quad \Rightarrow$ $2(x-3) + 3(y-2) = 0$ or $2x + 3y = 12$.

56. $g(x,y) = x^2 + y^2 - 4x \quad \Rightarrow \quad \nabla g(x,y) = \langle 2x-4, 2y \rangle$ and $\nabla g(1,2) = \langle -2, 4 \rangle$. Since $\nabla g(1,2)$ is perpendicular to the tangent line, the tangent line has equation $\nabla g(1,2) \cdot \langle x-1, y-2 \rangle = 0 \quad \Rightarrow$ $\langle -2,4 \rangle \cdot \langle x-1, y-2 \rangle = 0 \quad \Rightarrow \quad -2(x-1) + 4(y-2) = 0 \quad \Leftrightarrow$ $-2x + 4y = 6$ or equivalently $-x + 2y = 3$.

57. $F(x,y,z) = x^2/a^2 + y^2/b^2 + z^2/c^2$. Then $x^2/a^2 + y^2/b^2 + z^2/c^2 = 1$ is the level surface $F(x,y,z) = 1$ and $\nabla F(x_0,y_0,z_0) = \left\langle \dfrac{2x_0}{a^2}, \dfrac{2y_0}{b^2}, \dfrac{2z_0}{c^2} \right\rangle$. Thus, an equation of the tangent plane at (x_0,y_0,z_0) is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\dfrac{2x_0}{a^2}\,x + \dfrac{2y_0}{b^2}\,y + \dfrac{2z_0}{c^2}\,z = 2\left(\dfrac{x_0^2}{a^2} + \dfrac{y_0^2}{b^2} + \dfrac{z_0^2}{c^2}\right) = 2(1) = 2$ since (x_0, y_0, z_0) is a point on the ellipsoid. Hence $\dfrac{x_0}{a^2}\,x + \dfrac{y_0}{b^2}\,y + \dfrac{z_0}{c^2}\,z = 1$ is an equation of the tangent plane.

58. $F(x, y, z) = x^2/a^2 + y^2/b^2 - z^2/c^2$. Then $x^2/a^2 + y^2/b^2 - z^2/c^2 = 1$ is the level surface $F(x, y, z) = 1$ and $\nabla F(x_0, y_0, z_0) = \left\langle \dfrac{2x_0}{a^2}, \dfrac{2y_0}{b^2}, \dfrac{-2z_0}{c^2} \right\rangle$, so an equation of the tangent plane at (x_0, y_0, z_0) is $\dfrac{2x_0}{a^2}\,x + \dfrac{2y_0}{b^2}\,y - \dfrac{2z_0}{c^2}\,z = 2\left(\dfrac{x_0^2}{a^2} + \dfrac{y_0^2}{b^2} - \dfrac{z_0^2}{c^2}\right) = 2$ or $\dfrac{x_0}{a^2}\,x + \dfrac{y_0}{b^2}\,y - \dfrac{z_0}{c^2}\,z = 1$.

59. $F(x, y, z) = x^2/a^2 + y^2/b^2 - z/c$. Then $z/c = x^2/a^2 + y^2/b^2$ is the level surface $F(x, y, z) = 0$ and $\nabla F(x_0, y_0, z_0) = \left\langle \dfrac{2x_0}{a^2}, \dfrac{2y_0}{b^2}, \dfrac{-1}{c} \right\rangle$, so an equation of the tangent plane is $\dfrac{2x_0}{a^2}\,x + \dfrac{2y_0}{b^2}\,y - \dfrac{1}{c}\,z = \dfrac{2x_0^2}{a^2} + \dfrac{2y_0^2}{b^2} - \dfrac{z_0}{c}$ or $\dfrac{2x_0}{a^2}\,x + \dfrac{2y_0}{b^2}\,y = \dfrac{z}{c} + 2\left(\dfrac{x_0^2}{a^2} + \dfrac{y_0^2}{b^2}\right) - \dfrac{z_0}{c}$. But $\dfrac{z_0}{c} = \dfrac{x_0^2}{a^2} + \dfrac{y_0^2}{b^2}$, so the equation can be written as $\dfrac{2x_0}{a^2}\,x + \dfrac{2y_0}{b^2}\,y = \dfrac{z + z_0}{c}$.

60. Let $F(x, y, z) = x^2 + y^2 + 2z^2$; then the ellipsoid $x^2 + y^2 + 2z^2 = 1$ is a level surface of F. $\nabla F(x, y, z) = \langle 2x, 2y, 4z \rangle$ is a normal vector to the surface at (x, y, z) and so it is a normal vector for the tangent plane there. The tangent plane is parallel to the plane $x + 2y + z = 1$ when the normal vectors of the planes are parallel, so we need a point (x_0, y_0, z_0) on the ellipsoid where $\langle 2x_0, 2y_0, 4z_0 \rangle = k\,\langle 1, 2, 1 \rangle$ for some $k \neq 0$. Comparing components we have $2x_0 = k \;\Rightarrow\; x_0 = k/2$, $2y_0 = 2k \;\Rightarrow\; y_0 = k$, $4z_0 = k \;\Rightarrow\; z_0 = k/4$. $(x_0, y_0, z_0) = (k/2, k, k/4)$ lies on the ellipsoid, so $(k/2)^2 + k^2 + 2(k/4)^2 = 1 \;\Rightarrow\; \frac{11}{8}k^2 = 1 \;\Rightarrow\; k^2 = \frac{8}{11} \;\Rightarrow\; k = \pm 2\sqrt{\frac{2}{11}}$. Thus the tangent planes at the points $\left(\sqrt{\frac{2}{11}}, 2\sqrt{\frac{2}{11}}, \frac{1}{2}\sqrt{\frac{2}{11}}\right)$ and $\left(-\sqrt{\frac{2}{11}}, -2\sqrt{\frac{2}{11}}, -\frac{1}{2}\sqrt{\frac{2}{11}}\right)$ are parallel to the given plane.

61. The hyperboloid $x^2 - y^2 - z^2 = 1$ is a level surface of $F(x, y, z) = x^2 - y^2 - z^2$ and $\nabla F(x, y, z) = \langle 2x, -2y, -2z \rangle$ is a normal vector to the surface and hence a normal vector for the tangent plane at (x, y, z). The tangent plane is parallel to the plane $z = x + y$ or $x + y - z = 0$ if and only if the corresponding normal vectors are parallel, so we need a point (x_0, y_0, z_0) on the hyperboloid where $\langle 2x_0, -2y_0, -2z_0 \rangle = c\,\langle 1, 1, -1 \rangle$ or equivalently $\langle x_0, -y_0, -z_0 \rangle = k\,\langle 1, 1, -1 \rangle$ for some $k \neq 0$. Then we must have $x_0 = k$, $y_0 = -k$, $z_0 = k$ and substituting into the equation of the hyperboloid gives $k^2 - (-k)^2 - k^2 = 1 \;\Leftrightarrow\; -k^2 = 1$, an impossibility. Thus there is no such point on the hyperboloid.

62. First note that the point $(1, 1, 2)$ is on both surfaces. The ellipsoid $3x^2 + 2y^2 + z^2 = 9$ is a level surface of $F(x, y, z) = 3x^2 + 2y^2 + z^2$ and $\nabla F(x, y, z) = \langle 6x, 4y, 2z \rangle$. A normal vector to the surface at $(1, 1, 2)$ is $\nabla F(1, 1, 2) = \langle 6, 4, 4 \rangle$ and an equation of the tangent plane there is $6(x - 1) + 4(y - 1) + 4(z - 2) = 0$ or

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$6x+4y+4z=18$ or $3x+2y+2z=9$. The sphere is a level surface of $G(x,y,z)=x^2+y^2+z^2-8x-6y-8z+24$ and $\nabla G(x,y,z)=\langle 2x-8, 2y-6, 2z-8\rangle$. A normal vector to the sphere at $(1,1,2)$ is $\nabla G(1,1,2)=\langle -6,-4,-4\rangle$ and the tangent plane there is $-6(x-1)-4(y-1)-4(z-2)=0$ or $3x+2y+2z=9$. Since these tangent planes are identical, the surfaces are tangent to each other at the point $(1,1,2)$.

63. Let (x_0,y_0,z_0) be a point on the cone [other than $(0,0,0)$]. The cone $x^2+y^2=z^2$ is a level surface of $F(x,y,z)=x^2+y^2-z^2$ and $\nabla F(x,y,z)=\langle 2x,2y,-2z\rangle$, so $\nabla F(x_0,y_0,z_0)=\langle 2x_0,2y_0,-2z_0\rangle$ is a normal vector to the cone at this point and an equation of the tangent plane there is $2x_0\,(x-x_0)+2y_0\,(y-y_0)-2z_0\,(z-z_0)=0$ or $x_0x+y_0y-z_0z=x_0^2+y_0^2-z_0^2$. But $x_0^2+y_0^2=z_0^2$ so the tangent plane is given by $x_0x+y_0y-z_0z=0$, a plane which always contains the origin.

64. Let (x_0,y_0,z_0) be a point on the sphere and $F(x,y,z)=x^2+y^2+z^2$. Then $\nabla F(x,y,z)=\langle 2x,2y,2z\rangle$ and $\nabla F(x_0,y_0,z_0)=\langle 2x_0,2y_0,2z_0\rangle$, so the normal line is given by $\dfrac{x-x_0}{2x_0}=\dfrac{y-y_0}{2y_0}=\dfrac{z-z_0}{2z_0}$. For the center $(0,0,0)$ to be on the line, we need $-\dfrac{x_0}{2x_0}=-\dfrac{y_0}{2y_0}=-\dfrac{z_0}{2z_0}$ or equivalently $1=1=1$, which is true.

65. Let $F(x,y,z)=x^2+y^2-z$. Then the paraboloid is the level surface $F(x,y,z)=0$ and $\nabla F(x,y,z)=\langle 2x,2y,-1\rangle$, so $\nabla F(1,1,2)=\langle 2,2,-1\rangle$ is a normal vector to the surface. Thus the normal line at $(1,1,2)$ is given by $x=1+2t$, $y=1+2t$, $z=2-t$. Substitution into the equation of the paraboloid $z=x^2+y^2$ gives $2-t=(1+2t)^2+(1+2t)^2 \quad\Leftrightarrow$ $2-t=2+8t+8t^2 \quad\Leftrightarrow\quad 8t^2+9t=0 \quad\Leftrightarrow\quad t(8t+9)=0$. Thus the line intersects the paraboloid when $t=0$, corresponding to the given point $(1,1,2)$, or when $t=-\frac{9}{8}$, corresponding to the point $\left(-\frac{5}{4},-\frac{5}{4},\frac{25}{8}\right)$.

66. The ellipsoid $4x^2+y^2+4z^2=12$ is a level surface of $F(x,y,z)=4x^2+y^2+4z^2$ and $\nabla F(x,y,z)=\langle 8x,2y,8z\rangle$, so $\nabla F(1,2,1)=\langle 8,4,8\rangle$ or equivalently $\langle 2,1,2\rangle$ is a normal vector to the surface. Thus, the normal line to the ellipsoid at $(1,2,1)$ is given by $x=1+2t$, $y=2+t$, $z=1+2t$. Substitution into the equation of the sphere gives $(1+2t)^2+(2+t)^2+(1+2t)^2=102 \quad\Leftrightarrow\quad 6+12t+9t^2=102 \quad\Leftrightarrow\quad 9t^2+12t-96=0 \quad\Leftrightarrow$ $3(t+4)(3t-8)=0$. Thus, the line intersects the sphere when $t=-4$, corresponding to the point $(-7,-2,-7)$, and when $t=\frac{8}{3}$, corresponding to the point $\left(\frac{19}{3},\frac{14}{3},\frac{19}{3}\right)$.

67. Let (x_0,y_0,z_0) be a point on the surface and $F(x,y,z)=\sqrt{x}+\sqrt{y}+\sqrt{z}$. Then $\nabla F(x,y,z)=\left\langle \dfrac{1}{2\sqrt{x}},\dfrac{1}{2\sqrt{y}},\dfrac{1}{2\sqrt{z}}\right\rangle$ and $\nabla F(x_0,y_0,z_0)=\left\langle \dfrac{1}{2\sqrt{x_0}},\dfrac{1}{2\sqrt{y_0}},\dfrac{1}{2\sqrt{z_0}}\right\rangle$, so an equation of the tangent plane at the point is

$$\frac{x-x_0}{2\sqrt{x_0}}+\frac{y-y_0}{2\sqrt{y_0}}+\frac{z-z_0}{2\sqrt{z_0}}=0 \quad\Leftrightarrow\quad \frac{x}{2\sqrt{x_0}}+\frac{y}{2\sqrt{y_0}}+\frac{z}{2\sqrt{z_0}}=\frac{x_0}{2\sqrt{x_0}}+\frac{y_0}{2\sqrt{y_0}}+\frac{z_0}{2\sqrt{z_0}} \quad\Leftrightarrow$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\dfrac{x}{2\sqrt{x_0}} + \dfrac{y}{2\sqrt{y_0}} + \dfrac{z}{2\sqrt{z_0}} = \dfrac{\sqrt{x_0}+\sqrt{y_0}+\sqrt{z_0}}{2}$. But $\sqrt{x_0}+\sqrt{y_0}+\sqrt{z_0} = \sqrt{c}$, so the equation is $\dfrac{x}{\sqrt{x_0}} + \dfrac{y}{\sqrt{y_0}} + \dfrac{z}{\sqrt{z_0}} = \sqrt{c}$. The x-, y-, and z-intercepts are $\sqrt{cx_0}$, $\sqrt{cy_0}$, and $\sqrt{cz_0}$, respectively. (The x-intercept is found by setting $y = z = 0$ and solving the resulting equation for x, and the y- and z-intercepts are found similarly.) So the sum of the intercepts is $\sqrt{c}\left(\sqrt{x_0}+\sqrt{y_0}+\sqrt{z_0}\right) = c$, a constant.

68. The surface $xyz = 1$ is a level surface of $F(x,y,z) = xyz$ and $\nabla F(x,y,z) = \langle yz, xz, xy\rangle$ is normal to the surface, so a normal vector for the tangent plane to the surface at (x_0, y_0, z_0) is $\langle y_0z_0, x_0z_0, x_0y_0\rangle$. An equation for the tangent plane there is $y_0z_0(x-x_0) + x_0z_0(y-y_0) + x_0y_0(z-z_0) = 0 \;\Rightarrow\; y_0z_0x + x_0z_0y + x_0y_0z = 3x_0y_0z_0$ or $\dfrac{x}{x_0} + \dfrac{y}{y_0} + \dfrac{z}{z_0} = 3$. If (x_0, y_0, z_0) is in the first octant, then the tangent plane cuts off a pyramid in the first octant with vertices $(0,0,0)$, $(3x_0, 0, 0)$, $(0, 3y_0, 0)$, $(0, 0, 3z_0)$. The base in the xy-plane is a triangle with area $\frac{1}{2}(3x_0)(3y_0)$ and the height (along the z-axis) of the pyramid is $3z_0$. The volume of the pyramid for any point (x_0, y_0, z_0) on the surface $xyz = 1$ in the first octant is $\frac{1}{3}$ (base) (height) $= \frac{1}{3}\cdot\frac{1}{2}(3x_0)(3y_0)\cdot 3z_0 = \frac{9}{2}x_0y_0z_0 = \frac{9}{2}$ since $x_0y_0z_0 = 1$.

69. If $f(x,y,z) = z - x^2 - y^2$ and $g(x,y,z) = 4x^2 + y^2 + z^2$, then the tangent line is perpendicular to both ∇f and ∇g at $(-1,1,2)$. The vector $\mathbf{v} = \nabla f \times \nabla g$ will therefore be parallel to the tangent line.

We have $\nabla f(x,y,z) = \langle -2x, -2y, 1\rangle \;\Rightarrow\; \nabla f(-1,1,2) = \langle 2,-2,1\rangle$, and $\nabla g(x,y,z) = \langle 8x, 2y, 2z\rangle \;\Rightarrow$ $\nabla g(-1,1,2) = \langle -8, 2, 4\rangle$. Hence $\mathbf{v} = \nabla f \times \nabla g = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & 1 \\ -8 & 2 & 4 \end{vmatrix} = -10\,\mathbf{i} - 16\,\mathbf{j} - 12\,\mathbf{k}$.

Parametric equations are: $x = -1 - 10t,\; y = 1 - 16t,\; z = 2 - 12t$.

70. (a) Let $f(x,y,z) = y + z$ and $g(x,y,z) = x^2 + y^2$. The plane is a level surface of f and the cylinder is a level surface of g. Then the required tangent line is perpendicular to both ∇f and ∇g at $(1,2,1)$ and the vector $\mathbf{v} = \nabla f \times \nabla g$ is parallel to the tangent line. We have $\nabla f(x,y,z) = \langle 0,1,1\rangle \;\Rightarrow\; \nabla f(1,2,1) = \langle 0,1,1\rangle$, and $\nabla g(x,y,z) = \langle 2x, 2y, 0\rangle \;\Rightarrow\; \nabla g(1,2,1) = \langle 2,4,0\rangle$. Hence $\mathbf{v} = \nabla f \times \nabla g = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 1 & 1 \\ 2 & 4 & 0 \end{vmatrix} = -4\,\mathbf{i} + 2\,\mathbf{j} - 2\,\mathbf{k}$. So parametric equations of the desired tangent line are $x = 1 - 4t,\; y = 2 + 2t,\; z = 1 - 2t$.

(b)

71. Parametric equations for the helix are $x = \cos\pi t,\;\; y = \sin\pi t,\;\; z = t$, and substituting into the equation of the paraboloid gives $t = \cos^2\pi t + \sin^2\pi t \;\Rightarrow\; t = 1$. Thus the helix intersects the surface at the point $(\cos\pi, \sin\pi, 1) = (-1, 0, 1)$. Here $\mathbf{r}'(t) = \langle -\pi\sin\pi t, \pi\cos\pi t, 1\rangle$, so the tangent vector to the helix at that point is $\mathbf{r}'(1) = \langle -\pi\sin\pi, \pi\cos\pi, 1\rangle = \langle 0, -\pi, 1\rangle$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The paraboloid $z = x^2 + y^2 \quad \Leftrightarrow \quad x^2 + y^2 - z = 0$ is a level surface of $F(x, y, z) = x^2 + y^2 - z$ and $\nabla F(x, y, z) = \langle 2x, 2y, -1\rangle$, so a normal vector to the tangent plane at $(-1, 0, 1)$ is $\nabla F(-1, 0, 1) = \langle -2, 0, -1\rangle$. The angle θ between $\mathbf{r}'(1)$ and $\nabla F(-1, 0, 1)$ is given by

$$\cos\theta = \frac{\langle 0, -\pi, 1\rangle \cdot \langle -2, 0, -1\rangle}{|\langle 0, -\pi, 1\rangle|\ |\langle -2, 0, -1\rangle|} = \frac{0 + 0 - 1}{\sqrt{0 + \pi^2 + 1}\sqrt{4 + 0 + 1}} = \frac{-1}{\sqrt{5(\pi^2 + 1)}} \quad \Rightarrow$$

$\theta = \cos^{-1} \dfrac{-1}{\sqrt{5(\pi^2 + 1)}} \approx 97.8°$. Because $\nabla F(-1, 0, 1)$ is perpendicular to the tangent plane, the angle of intersection between the helix and the paraboloid is approximately $97.8° - 90° = 7.8°$.

72. Parametric equations for the helix are $x = \cos(\pi t/2)$, $y = \sin(\pi t/2)$, $z = t$, and substituting into the equation of the sphere gives $\cos^2(\pi t/2) + \sin^2(\pi t/2) + t^2 = 2 \quad \Rightarrow \quad 1 + t^2 = 2 \quad \Rightarrow \quad t = \pm 1$. Thus the helix intersects the sphere at two points: $(\cos(\pi/2), \sin(\pi/2), 1) = (0, 1, 1)$, when $t = 1$, and $(\cos(-\pi/2), \sin(-\pi/2), -1) = (0, -1, -1)$, when $t = -1$. Here $\mathbf{r}'(t) = \left\langle -\frac{\pi}{2}\sin(\pi t/2), \frac{\pi}{2}\cos(\pi t/2), 1\right\rangle$, so the tangent vector to the helix at $(0, 1, 1)$ is $\mathbf{r}'(1) = \langle -\pi/2, 0, 1\rangle$. The sphere $x^2 + y^2 + z^2 = 2$ is a level surface of $F(x, y, z) = x^2 + y^2 + z^2$ and $\nabla F(x, y, z) = \langle 2x, 2y, 2z\rangle$, so a normal vector to the tangent plane at $(0, 1, 1)$ is $\nabla F(0, 1, 1) = \langle 0, 2, 2\rangle$. As in Exercise 71, the angle of intersection between the helix and the sphere is the angle between the tangent vector to the helix and the tangent plane to the sphere. The angle θ between $\mathbf{r}'(1)$ and $\nabla F(0, 1, 1)$ is given by

$$\cos\theta = \frac{\langle -\pi/2, 0, 1\rangle \cdot \langle 0, 2, 2\rangle}{|\langle -\pi/2, 0, 1\rangle|\ |\langle 0, 2, 2\rangle|} = \frac{2}{\sqrt{(\pi^2/4) + 1}\sqrt{8}} = \frac{2}{\sqrt{2\pi^2 + 8}} \quad \Rightarrow \quad \theta = \cos^{-1}\frac{2}{\sqrt{2\pi^2 + 8}} \approx 67.7°$$

Because $\nabla F(0, 1, 1)$ is perpendicular to the tangent plane, the angle between $\mathbf{r}'(1)$ and the tangent plane is approximately $90° - 67.7° = 22.3°$.

At $(0, -1, -1)$, $\mathbf{r}'(-1) = \langle \pi/2, 0, 1\rangle$ and $\nabla F(0, -1, -1) = \langle 0, -2, -2\rangle$, and the angle ϕ between these vectors is given by $\cos\phi = \dfrac{\langle \pi/2, 0, 1\rangle \cdot \langle 0, -2, -2\rangle}{|\langle \pi/2, 0, 1\rangle|\ |\langle 0, -2, -2\rangle|} = \dfrac{-2}{\sqrt{2\pi^2 + 8}} \quad \Rightarrow \quad \phi = \cos^{-1}\dfrac{-2}{\sqrt{2\pi^2 + 8}} \approx 112.3°$. Thus the angle between the helix and the sphere at $(0, -1, -1)$ is approximately $112.3° - 90° = 22.3°$. (By symmetry, we would expect the angles to be identical.)

73. The direction of the normal line of F is given by ∇F, and that of G by ∇G. Assuming that $\nabla F \neq 0 \neq \nabla G$, the two normal lines are perpendicular at P if $\nabla F \cdot \nabla G = 0$ at $P \quad \Leftrightarrow \quad \langle \partial F/\partial x, \partial F/\partial y, \partial F/\partial z\rangle \cdot \langle \partial G/\partial x, \partial G/\partial y, \partial G/\partial z\rangle = 0$ at $P \quad \Leftrightarrow \quad F_x G_x + F_y G_y + F_z G_z = 0$ at P.

74. Here $F(x, y, z) = x^2 + y^2 - z^2$ and $G(x, y, z) = x^2 + y^2 + z^2 - r^2$, so $z^2 = x^2 + y^2$ is a level surface of F and $x^2 + y^2 + z^2 = r^2$ is a level surface of $G \quad \Rightarrow \quad \nabla F \cdot \nabla G = \langle 2x, 2y, -2z\rangle \cdot \langle 2x, 2y, 2z\rangle = 4x^2 + 4y^2 - 4z^2 = 4F = 0$, since the point (x, y, z) lies on the graph of $F = 0$. To see that this is true without using calculus, note that $G = 0$ is the equation of a sphere centered at the origin and $F = 0$ is the equation of a right circular cone with vertex at the origin (which is generated by lines through the origin). At any point of intersection, the sphere's normal line (which passes through the origin)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

lies on the cone, and thus is perpendicular to the cone's normal line. So the surfaces with equations $F = 0$ and $G = 0$ are everywhere orthogonal.

75. Let $\mathbf{u} = \langle a, b\rangle$ and $\mathbf{v} = \langle c, d\rangle$. Then we know that at the given point, $D_{\mathbf{u}} f = \nabla f \cdot \mathbf{u} = af_x + bf_y$ and $D_{\mathbf{v}} f = \nabla f \cdot \mathbf{v} = cf_x + df_y$. But these are just two linear equations in the two unknowns f_x and f_y, and since $\mathbf{u}$ and $\mathbf{v}$ are not parallel, we can solve the equations to find $\nabla f = \langle f_x, f_y\rangle$ at the given point. In fact,

$$\nabla f = \left\langle \frac{d\, D_{\mathbf{u}} f - b\, D_{\mathbf{v}} f}{ad - bc}, \frac{a\, D_{\mathbf{v}} f - c\, D_{\mathbf{u}} f}{ad - bc} \right\rangle.$$

76. (a) The function $f(x, y) = (xy)^{1/3}$ is continuous on $\mathbb{R}^2$ since it is a composition of a polynomial and the cube root function, both of which are continuous. (See the text just after Example 14.2.9.)

$$f_x(0,0) = \lim_{h\to 0} \frac{f(0+h,0) - f(0,0)}{h} = \lim_{h\to 0} \frac{(h\cdot 0)^{1/3} - 0}{h} = 0,$$

$$f_y(0,0) = \lim_{h\to 0} \frac{f(0,0+h) - f(0,0)}{h} = \lim_{h\to 0} \frac{(0\cdot h)^{1/3} - 0}{h} = 0.$$

Therefore, $f_x(0,0)$ and $f_y(0,0)$ do exist and are equal to 0. Now let $\mathbf{u}$ be any unit vector other than $\mathbf{i}$ and $\mathbf{j}$ (these correspond to f_x and f_y respectively.) Then $\mathbf{u} = a\,\mathbf{i} + b\,\mathbf{j}$ where $a \neq 0$ and $b \neq 0$. Thus

$D_{\mathbf{u}} f(0,0) = \lim\limits_{h\to 0} \dfrac{f(0+ha, 0+hb) - f(0,0)}{h} = \lim\limits_{h\to 0} \dfrac{\sqrt[3]{(ha)(hb)}}{h} = \lim\limits_{h\to 0} \dfrac{\sqrt[3]{ab}}{h^{1/3}}$ and this limit does not exist, so $D_{\mathbf{u}} f(0,0)$ does not exist.

(b)

Notice that if we start at the origin and proceed in the direction of the x- or y-axis, then the graph is flat. But if we proceed in any other direction, then the graph is extremely steep.

77. Since $z = f(x, y)$ is differentiable at $\mathbf{x}_0 = (x_0, y_0)$, by Definition 14.4.7 we have $\Delta z = f_x(x_0, y_0)\,\Delta x + f_y(x_0, y_0)\,\Delta y + \varepsilon_1\,\Delta x + \varepsilon_2\,\Delta y$ where $\varepsilon_1, \varepsilon_2 \to 0$ as $(\Delta x, \Delta y) \to (0, 0)$. Now $\Delta z = f(\mathbf{x}) - f(\mathbf{x}_0)$, $\langle \Delta x, \Delta y\rangle = \mathbf{x} - \mathbf{x}_0$ so $(\Delta x, \Delta y) \to (0, 0)$ is equivalent to $\mathbf{x} \to \mathbf{x}_0$ and $\langle f_x(x_0, y_0), f_y(x_0, y_0)\rangle = \nabla f(\mathbf{x}_0)$. Substituting into 14.4.7 gives $f(\mathbf{x}) - f(\mathbf{x}_0) = \nabla f(\mathbf{x}_0)\cdot(\mathbf{x} - \mathbf{x}_0) + \langle \varepsilon_1, \varepsilon_2\rangle \cdot \langle \Delta x, \Delta y\rangle$ or $\langle \varepsilon_1, \varepsilon_2\rangle \cdot (\mathbf{x} - \mathbf{x}_0) = f(\mathbf{x}) - f(\mathbf{x}_0) - \nabla f(\mathbf{x}_0)\cdot(\mathbf{x} - \mathbf{x}_0)$,

and so $\dfrac{f(\mathbf{x}) - f(\mathbf{x}_0) - \nabla f(\mathbf{x}_0)\cdot(\mathbf{x} - \mathbf{x}_0)}{|\mathbf{x} - \mathbf{x}_0|} = \dfrac{\langle \varepsilon_1, \varepsilon_2\rangle \cdot (\mathbf{x} - \mathbf{x}_0)}{|\mathbf{x} - \mathbf{x}_0|}$. But $\dfrac{\mathbf{x} - \mathbf{x}_0}{|\mathbf{x} - \mathbf{x}_0|}$ is a unit vector so

$\lim\limits_{\mathbf{x}\to\mathbf{x}_0} \dfrac{\langle \varepsilon_1, \varepsilon_2\rangle \cdot (\mathbf{x} - \mathbf{x}_0)}{|\mathbf{x} - \mathbf{x}_0|} = 0$ since $\varepsilon_1, \varepsilon_2 \to 0$ as $\mathbf{x} \to \mathbf{x}_0$. Hence $\lim\limits_{\mathbf{x}\to\mathbf{x}_0} \dfrac{f(\mathbf{x}) - f(\mathbf{x}_0) - \nabla f(\mathbf{x}_0)\cdot(\mathbf{x} - \mathbf{x}_0)}{|\mathbf{x} - \mathbf{x}_0|} = 0.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14.7 Maximum and Minimum Values

1. (a) First we compute $D(1,1) = f_{xx}(1,1)\, f_{yy}(1,1) - [f_{xy}(1,1)]^2 = (4)(2) - (1)^2 = 7$. Since $D(1,1) > 0$ and $f_{xx}(1,1) > 0$, f has a local minimum at $(1,1)$ by the Second Derivatives Test.

(b) $D(1,1) = f_{xx}(1,1)\, f_{yy}(1,1) - [f_{xy}(1,1)]^2 = (4)(2) - (3)^2 = -1$. Since $D(1,1) < 0$, f has a saddle point at $(1,1)$ by the Second Derivatives Test.

2. (a) $D(0,2) = g_{xx}(0,2)\, g_{yy}(0,2) - [g_{xy}(0,2)]^2 = (-1)(1) - (6)^2 = -37$. Since $D(0,2) < 0$, g has a saddle point at $(0,2)$ by the Second Derivatives Test.

(b) $D(0,2) = g_{xx}(0,2)\, g_{yy}(0,2) - [g_{xy}(0,2)]^2 = (-1)(-8) - (2)^2 = 4$. Since $D(0,2) > 0$ and $g_{xx}(0,2) < 0$, g has a local maximum at $(0,2)$ by the Second Derivatives Test.

(c) $D(0,2) = g_{xx}(0,2)\, g_{yy}(0,2) - [g_{xy}(0,2)]^2 = (4)(9) - (6)^2 = 0$. In this case the Second Derivatives Test gives no information about g at the point $(0,2)$.

3. In the figure, a point at approximately $(1,1)$ is enclosed by level curves which are oval in shape and indicate that as we move away from the point in any direction the values of f are increasing. Hence we would expect a local minimum at or near $(1,1)$. The level curves near $(0,0)$ resemble hyperbolas, and as we move away from the origin, the values of f increase in some directions and decrease in others, so we would expect to find a saddle point there.

To verify our predictions, we have $f(x,y) = 4 + x^3 + y^3 - 3xy \;\Rightarrow\; f_x(x,y) = 3x^2 - 3y$, $f_y(x,y) = 3y^2 - 3x$. We have critical points where these partial derivatives are equal to 0: $3x^2 - 3y = 0$, $3y^2 - 3x = 0$. Substituting $y = x^2$ from the first equation into the second equation gives $3(x^2)^2 - 3x = 0 \;\Rightarrow\; 3x(x^3 - 1) = 0 \;\Rightarrow\; x = 0$ or $x = 1$. Then we have two critical points, $(0,0)$ and $(1,1)$. The second partial derivatives are $f_{xx}(x,y) = 6x$, $f_{xy}(x,y) = -3$, and $f_{yy}(x,y) = 6y$, so $D(x,y) = f_{xx}(x,y)\, f_{yy}(x,y) - [f_{xy}(x,y)]^2 = (6x)(6y) - (-3)^2 = 36xy - 9$. Then $D(0,0) = 36(0)(0) - 9 = -9$, and $D(1,1) = 36(1)(1) - 9 = 27$. Since $D(0,0) < 0$, f has a saddle point at $(0,0)$ by the Second Derivatives Test. Since $D(1,1) > 0$ and $f_{xx}(1,1) > 0$, f has a local minimum at $(1,1)$.

4. In the figure, points at approximately $(-1,1)$ and $(-1,-1)$ are enclosed by oval-shaped level curves which indicate that as we move away from either point in any direction, the values of f are increasing. Hence we would expect local minimums at or near $(-1,\pm 1)$. Similarly, the point $(1,0)$ appears to be enclosed by oval-shaped level curves which indicate that as we move away from the point in any direction the values of f are decreasing, so we should have a local maximum there. We also show hyperbola-shaped level curves near the points $(-1,0)$, $(1,1)$, and $(1,-1)$. The values of f increase along some paths leaving these points and decrease in others, so we should have a saddle point at each of these points.

To confirm our predictions, we have $f(x,y) = 3x - x^3 - 2y^2 + y^4 \;\Rightarrow\; f_x(x,y) = 3 - 3x^2,\;\; f_y(x,y) = -4y + 4y^3$. Setting these partial derivatives equal to 0, we have $3 - 3x^2 = 0 \;\Rightarrow\; x = \pm 1$ and $-4y + 4y^3 = 0 \;\Rightarrow$ $y(y^2 - 1) = 0 \;\Rightarrow\; y = 0, \pm 1$. So our critical points are $(\pm 1, 0)$, $(\pm 1, 1)$, $(\pm 1, -1)$.

The second partial derivatives are $f_{xx}(x,y) = -6x$, $f_{xy}(x,y) = 0$, and $f_{yy}(x,y) = 12y^2 - 4$, so

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$D(x,y) = f_{xx}(x,y)\, f_{yy}(x,y) - [f_{xy}(x,y)]^2 = (-6x)(12y^2-4) - (0)^2 = -72xy^2 + 24x.$

We use the Second Derivatives Test to classify the 6 critical points:

Critical Point	D	f_{xx}	Conclusion
$(1,0)$	24	-6	$D>0,\ f_{xx}<0 \quad\Rightarrow\quad f$ has a local maximum at $(1,0)$
$(1,1)$	-48		$D<0 \quad\Rightarrow\quad f$ has a saddle point at $(1,1)$
$(1,-1)$	-48		$D<0 \quad\Rightarrow\quad f$ has a saddle point at $(1,-1)$
$(-1,0)$	-24		$D<0 \quad\Rightarrow\quad f$ has a saddle point at $(-1,0)$
$(-1,1)$	48	6	$D>0,\ f_{xx}>0 \quad\Rightarrow\quad f$ has a local minimum at $(-1,1)$
$(-1,-1)$	48	6	$D>0,\ f_{xx}>0 \quad\Rightarrow\quad f$ has a local minimum at $(-1,-1)$

5. $f(x,y) = x^2 + xy + y^2 + y \quad\Rightarrow\quad f_x = 2x + y,\ f_y = x + 2y + 1,\ f_{xx} = 2,\ f_{xy} = 1,\ f_{yy} = 2$. Then $f_x = 0$ implies $y = -2x$, and substitution into $f_y = x + 2y + 1 = 0$ gives $x + 2(-2x) + 1 = 0 \quad\Rightarrow\quad -3x = -1 \quad\Rightarrow\quad x = \frac{1}{3}$.

Then $y = -\frac{2}{3}$ and the only critical point is $\left(\frac{1}{3}, -\frac{2}{3}\right)$.

$D(x,y) = f_{xx}f_{yy} - (f_{xy})^2 = (2)(2) - (1)^2 = 3$, and since

$D\left(\frac{1}{3}, -\frac{2}{3}\right) = 3 > 0$ and $f_{xx}\left(\frac{1}{3}, -\frac{2}{3}\right) = 2 > 0$, $f\left(\frac{1}{3}, -\frac{2}{3}\right) = -\frac{1}{3}$ is a local minimum by the Second Derivatives Test.

6. $f(x,y) = xy - 2x - 2y - x^2 - y^2 \quad\Rightarrow\quad f_x = y - 2 - 2x$,

$f_y = x - 2 - 2y,\ f_{xx} = -2,\ f_{xy} = 1,\ f_{yy} = -2$. Then $f_x = 0$ implies $y = 2x + 2$, and substitution into $f_y = 0$ gives $x - 2 - 2(2x+2) = 0 \quad\Rightarrow$ $-3x = 6 \quad\Rightarrow\quad x = -2$. Then $y = -2$ and the only critical point is $(-2,-2)$. $D(x,y) = f_{xx}f_{yy} - (f_{xy})^2 = (-2)(-2) - 1^2 = 3$, and since $D(-2,-2) = 3 > 0$ and $f_{xx}(-2,-2) = -2 < 0$, $f(-2,-2) = 4$ is a local maximum by the Second Derivatives Test.

7. $f(x,y) = 2x^2 - 8xy + y^4 - 4y^3 \quad\Rightarrow\quad f_x = 4x - 8y$,

$f_y = -8x + 4y^3 - 12y^2$, $f_{xx} = 4$, $f_{xy} = -8$, $f_{yy} = 12y^2 - 24y$.

Then $f_x = 0$ implies $x = 2y$, and sustitution into $f_y = 0$ gives

$-16y + 4y^3 - 12y^2 = 0 \quad\Rightarrow\quad 4y(y^2 - 3y - 4) = 0 \quad\Rightarrow\quad y = 0, -1, 4$.

Then $x = 0, -2, 8$ and the critical points are $(0,0)$, $(-2,-1)$, and $(8,4)$.

$D(x,y) = f_{xx}f_{yy} - (f_{xy})^2$. $D(0,0) = 4(0) - 64 = -64 < 0$,

so $(0,0)$ is a saddle point. $D(-2,-1) = 4(36) - 64 = 80 > 0$ and

$f_{xx}(-2,-1) = 4 > 0$, so $f(-2,-1) = -3$ is a local minimum by the Second Derivatives Test.

$D(8,4) = 4(96) - 64 = 320 > 0$ and $f_{xx}(8,4) = 4 > 0$, so $f(8,4) = -128$ is a local minimum.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $f(x,y) = x^3 + y^3 + 3xy \quad \Rightarrow \quad f_x = 3x^2 + 3y$, $f_y = 3y^2 + 3x$, $f_{xx} = 6x$, $f_{xy} = 3$, $f_{yy} = 6y$. Then $f_x = 0$ implies $y = -x^2$ and substitution into $f_y = 0$ gives $3(-x^2)^2 + 3x = 3x^4 + 3x = 0 \quad \Rightarrow$ $3x(x^3 + 1) = 0 \quad \Rightarrow \quad x = 0$ or $x = -1$. Thus, the critical points are $(0,0)$ and $(-1,-1)$. $D(x,y) = (6x)(6y) - 3^2 = 36xy - 9$. $D(0,0) = -9 < 0$, so $(0,0)$ is a saddle point. $D(-1,-1) = 27 > 0$ and $f_{xx}(-1,-1) = -6 < 0$, so $f(-1,-1) = 1$ is a local maximum by the Second Derivatives Test.

9. $f(x,y) = (x-y)(1-xy) = x - y - x^2y + xy^2 \quad \Rightarrow \quad f_x = 1 - 2xy + y^2,\ f_y = -1 - x^2 + 2xy,\ f_{xx} = -2y,$ $f_{xy} = -2x + 2y,\ f_{yy} = 2x$. Then $f_x = 0$ implies $1 - 2xy + y^2 = 0$ and $f_y = 0$ implies $-1 - x^2 + 2xy = 0$. Adding the two equations gives $1 + y^2 - 1 - x^2 = 0 \quad \Rightarrow \quad y^2 = x^2 \quad \Rightarrow \quad y = \pm x$, but if $y = -x$ then $f_x = 0$ implies $1 + 2x^2 + x^2 = 0 \quad \Rightarrow \quad 3x^2 = -1$ which has no real solution.

If $y = x$, then substitution into $f_x = 0$ gives $1 - 2x^2 + x^2 = 0 \quad \Rightarrow$ $x^2 = 1 \quad \Rightarrow \quad x = \pm 1$, so the critical points are $(1,1)$ and $(-1,-1)$. Now $D(1,1) = (-2)(2) - 0^2 = -4 < 0$ and $D(-1,-1) = (2)(-2) - 0^2 = -4 < 0$, so $(1,1)$ and $(-1,-1)$ are saddle points.

10. $f(x,y) = y(e^x - 1) \quad \Rightarrow \quad f_x = ye^x,\ f_y = e^x - 1,\ f_{xx} = ye^x,$ $f_{xy} = e^x,\ f_{yy} = 0$. Because e^x is never zero, $f_x = 0$ only when $y = 0$, and $f_y = 0$ when $e^x = 1 \quad \Rightarrow \quad x = 0$, so the only critical point is $(0,0)$. $D(x,y) = f_{xx}f_{yy} - (f_{xy})^2 = (ye^x)(0) - (e^x)^2 = -e^{2x}$, and since $D(0,0) = -1 < 0$, $(0,0)$ is a saddle point.

11. $f(x,y) = y\sqrt{x} - y^2 - 2x + 7y \quad \Rightarrow \quad f_x = \frac{1}{2}yx^{-1/2} - 2,$ $f_y = \sqrt{x} - 2y + 7$, $f_{xx} = -\frac{1}{4}yx^{-3/2}$, $f_{xy} = \frac{1}{2}x^{-1/2}$, $f_{yy} = -2$. Then $f_x = 0 \quad \Rightarrow \quad y = 4\sqrt{x}$ and substitution into $f_y = 0$ gives $\sqrt{x} - 2\left(4\sqrt{x}\right) + 7 = -7\sqrt{x} + 7 = 0 \quad \Rightarrow \quad x = 1$, so the only critical point is $(1,4)$.

$$D(x,y) = -\frac{1}{4}yx^{-3/2}(-2) - \left(\frac{1}{2}x^{-1/2}\right)^2 = \frac{1}{2}yx^{-3/2} - \frac{1}{4x}.$$

$D(1,4) = \frac{7}{4} > 0$, and $f_{xx}(1,4) = -1 < 0$, so $f(1,4) = 14$ is a local maximum by the Second Derivatives Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $f(x,y) = 2 - x^4 + 2x^2 - y^2 \quad \Rightarrow \quad f_x = -4x^3 + 4x,\ f_y = -2y,\ f_{xx} = -12x^2 + 4,\ f_{xy} = 0,\ f_{yy} = -2$. Then $f_x = 0$ implies $-4x(x^2 - 1) = 0$, so $x = 0$ or $x = \pm 1$, and $f_y = 0$ implies $y = 0$. Thus the critical points are $(0,0)$, $(\pm 1, 0)$.

$D(0,0) = (4)(-2) - 0^2 = -8 < 0$, so $(0,0)$ is a saddle point.

$D(1,0) = D(-1,0) = (-8)(-2) - (0)^2 = 16 > 0$, and

$f_{xx}(1,0) = f_{xx}(-1,0) = -8 < 0$, so $f(1,0) = 3$ and $f(-1,0) = 3$ are local maximums.

13. $f(x,y) = x^3 - 3x + 3xy^2 \quad \Rightarrow \quad f_x = 3x^2 - 3 + 3y^2,\ f_y = 6xy,\ f_{xx} = 6x,\ f_{xy} = 6y,\ f_{yy} = 6x$. Then $f_y = 0$ implies $x = 0$ or $y = 0$. If $x = 0$, substitution into $f_x = 0$ gives $3y^2 = 3 \quad \Rightarrow \quad y = \pm 1$, and if $y = 0$, substitution into $f_x = 0$ gives $x = \pm 1$. Thus the critical points are $(0, \pm 1)$ and $(\pm 1, 0)$.

$D(0, \pm 1) = 0 - 36 < 0$, so $(0, \pm 1)$ are saddle points.

$D(\pm 1, 0) = 36 - 0 > 0$, $f_{xx}(1,0) = 6 > 0$, and $f_{xx}(-1,0) = -6 < 0$, so $f(1,0) = -2$ is a local minimum and $f(-1,0) = 2$ is a local maximum.

14. $f(x,y) = x^3 + y^3 - 3x^2 - 3y^2 - 9x \quad \Rightarrow \quad f_x = 3x^2 - 6x - 9,\ f_y = 3y^2 - 6y,\ f_{xx} = 6x - 6,\ f_{xy} = 0,\ f_{yy} = 6y - 6$. Then $f_x = 0$ implies $3(x+1)(x-3) = 0 \quad \Rightarrow \quad x = -1$ or $x = 3$, and $f_y = 0$ implies $3y(y-2) = 0 \quad \Rightarrow \quad y = 0$ or $y = 2$. Thus the critical points are $(-1,0)$, $(-1,2)$, $(3,0)$, and $(3,2)$. $D(-1,2) = (-12)(6) - (0)^2 = -72 < 0$ and

$D(3,0) = (12)(-6) - (0)^2 = -72 < 0$, so $(-1,2)$ and $(3,0)$ are saddle points. $D(-1,0) = (-12)(-6) - (0)^2 = 72 > 0$ and

$f_{xx}(-1,0) = -12 < 0$, so $f(-1,0) = 5$ is a local maximum.

$D(3,2) = (12)(6) - (0)^2 = 72 > 0$ and $f_{xx}(3,2) = 12 > 0$, so $f(3,2) = -31$ is a local minimum.

15. $f(x,y) = x^4 - 2x^2 + y^3 - 3y \quad \Rightarrow \quad f_x = 4x^3 - 4x,\ f_y = 3y^2 - 3,\ f_{xx} = 12x^2 - 4,\ f_{xy} = 0,\ f_{yy} = 6y$. Then $f_x = 0$ implies $4x(x^2 - 1) = 0 \quad \Rightarrow \quad x = 0$ or $x = \pm 1$, and $f_y = 0$ implies $3(y^2 - 1) = 0 \quad \Rightarrow \quad y = \pm 1$. Thus there are six critical points: $(0, \pm 1)$, $(\pm 1, 1)$, and $(\pm 1, -1)$.

$D(0,1) = (-4)(6) - (0)^2 = -24 < 0$ and

$D(\pm 1, -1) = (8)(-6) = -48 < 0$, so $(0,1)$ and $(\pm 1, -1)$ are saddle points. $D(0,-1) = (-4)(-6) = 24 > 0$ and $f_{xx}(0,-1) = -4 < 0$, so $f(0,-1) = 2$ is a local maximum. $D(\pm 1, 1) = (8)(6) = 48 > 0$ and $f_{xx}(\pm 1, 1) = 8 > 0$, so $f(\pm 1, 1) = -3$ are local minimums.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. $f(x,y) = x^2 + y^4 + 2xy \quad \Rightarrow \quad f_x = 2x + 2y,\ f_y = 4y^3 + 2x,\ f_{xx} = 2,\ f_{xy} = 2,\ f_{yy} = 12y^2$. Then $f_x = 0$ implies $y = -x$, and substitution into $f_y = 4y^3 + 2x = 0$ gives $-4x^3 + 2x = 0 \quad \Rightarrow \quad 2x\left(1 - 2x^2\right) = 0 \quad \Rightarrow \quad x = 0$ or $x = \pm\frac{1}{\sqrt{2}}$. Thus the critical points are $(0,0)$, $\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$, and $\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$. Now

$D(x,y) = f_{xx}f_{yy} - (f_{xy})^2 = (2)(12y^2) - (2)^2 = 24y^2 - 4$,
so $D(0,0) = -4 < 0$ and $(0,0)$ is a saddle point.

$D\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right) = D\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right) = 24\left(\frac{1}{2}\right) - 4 = 8 > 0$ and

$f_{xx}\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right) = f_{xx}\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right) = 2 > 0$, so $f\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right) = -\frac{1}{4}$

and $f\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right) = -\frac{1}{4}$ are local minimums.

17. $f(x,y) = xy - x^2y - xy^2 \quad \Rightarrow \quad f_x = y - 2xy - y^2$,

$f_y = x - x^2 - 2xy$, $f_{xx} = -2y$, $f_{xy} = 1 - 2x - 2y$, $f_{yy} = -2x$. Then

$f_x = y - 2xy - y^2 = 0 \quad \Rightarrow \quad y(1 - 2x - y) = 0 \quad \Rightarrow \quad y = 0$ or

$y = 1 - 2x$. Substituting $y = 0$ into $f_y = 0$ gives $x - x^2 = 0 \quad \Rightarrow$

$x = 0$ or $x = 1$. Next, substituting $y = 1 - 2x$ into $f_y = 0$ gives

$x - x^2 - 2x(1 - 2x) = 0 \quad \Rightarrow \quad 3x^2 - x = 0 \quad \Rightarrow \quad x = 0$ or $x = \frac{1}{3}$.

Thus, the critical points are $(0,0)$, $(1,0)$, $(0,1)$ and $\left(\frac{1}{3}, \frac{1}{3}\right)$.

$D(x,y) = (-2y)(-2x) - (1 - 2x - 2y)^2 = 4xy - (1 - 2x - 2y)^2$. $D(0,0) = D(1,0) = D(0,1) = -1 < 0$, so $(0,0)$, $(1,0)$, and $(0,1)$ are saddle points. $D\left(\frac{1}{3}, \frac{1}{3}\right) = \frac{1}{3} > 0$ and $f_{xx}\left(\frac{1}{3}, \frac{1}{3}\right) = -\frac{2}{3} < 0$, so $f\left(\frac{1}{3}, \frac{1}{3}\right) = \frac{1}{27}$ is a local maximum by the Second Derivatives Test.

18. $f(x,y) = (6x - x^2)(4y - y^2) \quad \Rightarrow \quad f_x = (6 - 2x)(4y - y^2)$,

$f_y = (6x - x^2)(4 - 2y)$, $f_{xx} = -2(4y - y^2)$, $f_{xy} = (6 - 2x)(4 - 2y)$,

$f_{yy} = -2(6x - x^2)$. Then $f_x = 0 \quad \Rightarrow \quad 6 - 2x = 0 \quad \Rightarrow \quad x = 3$ or

$4y - y^2 = 0 \quad \Rightarrow \quad y = 0$ or $y = 4$. Substituting $x = 3$ into $f_y = 0 \quad \Rightarrow$

$y = 2$. Substituting $y = 0$ into $f_y = 0 \quad \Rightarrow \quad x = 0$ or $x = 6$. $y = 4 \quad \Rightarrow$

$x = 0$ or $x = 6$ also. This gives critical points $(3,2)$, $(0,0)$, $(6,0)$, $(0,4)$,

and $(6,4)$. $D(x,y) = 4(4y - y^2)(6x - x^2) - [(6 - 2x)(4 - 2y)]^2$.

$D(0,0) = D(6,0) = D(0,4) = D(6,4) = -576 < 0$, so $(0,0)$, $(6,0)$, $(0,4)$, and $(6,4)$ are saddle points.

$D(3,2) = 144 > 0$ and $f_{xx}(3,2) = -8 < 0$, so $f(3,2) = 36$ is a local maximum by the Second Derivatives Test.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $f(x,y) = e^x \cos y \quad \Rightarrow \quad f_x = e^x \cos y,\ f_y = -e^x \sin y.$

Now $f_x = 0$ implies $\cos y = 0$ or $y = \frac{\pi}{2} + n\pi$ for n an integer.

But $\sin\left(\frac{\pi}{2} + n\pi\right) \neq 0$, so there are no critical points.

20. $f(x,y) = (x^2+y^2)e^{-x} \quad \Rightarrow \quad f_x = (x^2+y^2)(-e^{-x}) + e^{-x}(2x) = (2x - x^2 - y^2)e^{-x},\ \ f_y = 2ye^{-x},$

$f_{xx} = (2x - x^2 - y^2)(-e^{-x}) + e^{-x}(2-2x) = (x^2+y^2-4x+2)e^{-x},\ \ f_{xy} = -2ye^{-x},\ \ f_{yy} = 2e^{-x}$. Then $f_y = 0$ implies $y = 0$ and substituting into $f_x = 0$ gives $(2x - x^2)e^{-x} = 0 \quad \Rightarrow$ $x(2-x) = 0 \quad \Rightarrow \quad x = 0$ or $x = 2$, so the critical points are $(0,0)$ and $(2,0)$. $\quad D(0,0) = (2)(2) - (0)^2 = 4 > 0$ and $f_{xx}(0,0) = 2 > 0$, so $f(0,0) = 0$ is a local minimum.

$D(2,0) = (-2e^{-2})(2e^{-2}) - (0)^2 = -4e^{-4} < 0$ so $(2,0)$ is a saddle point.

21. $f(x,y) = y^2 - 2y\cos x \quad \Rightarrow \quad f_x = 2y\sin x,\ f_y = 2y - 2\cos x,$

$f_{xx} = 2y\cos x,\ f_{xy} = 2\sin x,\ f_{yy} = 2$. Then $f_x = 0$ implies $y = 0$ or $\sin x = 0 \quad \Rightarrow \quad x = 0, \pi$, or 2π for $-1 \le x \le 7$. Substituting $y = 0$ into $f_y = 0$ gives $\cos x = 0 \quad \Rightarrow \quad x = \frac{\pi}{2}$ or $\frac{3\pi}{2}$, substituting $x = 0$ or $x = 2\pi$ into $f_y = 0$ gives $y = 1$, and substituting $x = \pi$ into $f_y = 0$ gives $y = -1$. Thus the critical points are $(0,1)$, $\left(\frac{\pi}{2}, 0\right)$, $(\pi, -1)$, $\left(\frac{3\pi}{2}, 0\right)$, and $(2\pi, 1)$.

$D\left(\frac{\pi}{2}, 0\right) = D\left(\frac{3\pi}{2}, 0\right) = -4 < 0$ so $\left(\frac{\pi}{2}, 0\right)$ and $\left(\frac{3\pi}{2}, 0\right)$ are saddle points. $D(0,1) = D(\pi,-1) = D(2\pi,1) = 4 > 0$ and $f_{xx}(0,1) = f_{xx}(\pi,-1) = f_{xx}(2\pi,1) = 2 > 0$, so $f(0,1) = f(\pi,-1) = f(2\pi,1) = -1$ are local minimums.

22. $f(x,y) = \sin x\ \sin y \quad \Rightarrow \quad f_x = \cos x\ \sin y,\ f_y = \sin x\ \cos y,\ f_{xx} = -\sin x\ \sin y,\ f_{xy} = \cos x\ \cos y,$

$f_{yy} = -\sin x\ \sin y$. Here we have $-\pi < x < \pi$ and $-\pi < y < \pi$, so $f_x = 0$ implies $\cos x = 0$ or $\sin y = 0$. If $\cos x = 0$ then $x = -\frac{\pi}{2}$ or $\frac{\pi}{2}$, and if $\sin y = 0$ then $y = 0$. Substituting $x = \pm\frac{\pi}{2}$ into $f_y = 0$ gives $\cos y = 0 \quad \Rightarrow \quad y = -\frac{\pi}{2}$ or $\frac{\pi}{2}$, and substituting $y = 0$ into $f_y = 0$ gives $\sin x = 0 \quad \Rightarrow \quad x = 0$. Thus the critical points are $\left(-\frac{\pi}{2}, \pm\frac{\pi}{2}\right)$, $\left(\frac{\pi}{2}, \pm\frac{\pi}{2}\right)$, and $(0,0)$.

$D(0,0) = -1 < 0$ so $(0,0)$ is a saddle point.

$D\left(-\frac{\pi}{2}, \pm\frac{\pi}{2}\right) = D\left(\frac{\pi}{2}, \pm\frac{\pi}{2}\right) = 1 > 0$ and

$f_{xx}\left(-\frac{\pi}{2}, -\frac{\pi}{2}\right) = f_{xx}\left(\frac{\pi}{2}, \frac{\pi}{2}\right) = -1 < 0$ while

$f_{xx}\left(-\frac{\pi}{2}, \frac{\pi}{2}\right) = f_{xx}\left(\frac{\pi}{2}, -\frac{\pi}{2}\right) = 1 > 0$, so $f\left(-\frac{\pi}{2}, -\frac{\pi}{2}\right) = f\left(\frac{\pi}{2}, \frac{\pi}{2}\right) = 1$ are local maximums and $f\left(-\frac{\pi}{2}, \frac{\pi}{2}\right) = f\left(\frac{\pi}{2}, -\frac{\pi}{2}\right) = 1$ are local minimums.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. $f(x,y) = x^2 + 4y^2 - 4xy + 2 \quad \Rightarrow \quad f_x = 2x - 4y$, $f_y = 8y - 4x$, $f_{xx} = 2$, $f_{xy} = -4$, $f_{yy} = 8$. Then $f_x = 0$ and $f_y = 0$ each implies $y = \frac{1}{2}x$, so all points of the form $\left(x_0, \frac{1}{2}x_0\right)$ are critical points and for each of these we have $D\left(x_0, \frac{1}{2}x_0\right) = (2)(8) - (-4)^2 = 0$. The Second Derivatives Test gives no information, but $f(x,y) = x^2 + 4y^2 - 4xy + 2 = (x-2y)^2 + 2 \geq 2$ with equality if and only if $y = \frac{1}{2}x$. Thus $f\left(x_0, \frac{1}{2}x_0\right) = 2$ are all local (and absolute) minimums.

24. $f(x,y) = x^2 y e^{-x^2-y^2} \quad \Rightarrow$

$f_x = x^2 y e^{-x^2-y^2}(-2x) + 2xye^{-x^2-y^2} = 2xy(1-x^2)e^{-x^2-y^2}$,

$f_y = x^2 y e^{-x^2-y^2}(-2y) + x^2 e^{-x^2-y^2} = x^2(1-2y^2)e^{-x^2-y^2}$,

$f_{xx} = 2y(2x^4 - 5x^2 + 1)e^{-x^2-y^2}$,

$f_{xy} = 2x(1-x^2)(1-2y^2)e^{-x^2-y^2}$, $f_{yy} = 2x^2y(2y^2-3)e^{-x^2-y^2}$.

$f_x = 0$ implies $x = 0$, $y = 0$, or $x = \pm 1$. If $x = 0$ then $f_y = 0$ for any y-value, so all points of the form $(0, y)$ are critical points. If $y = 0$ then $f_y = 0 \quad \Rightarrow \quad x^2 e^{-x^2} = 0 \quad \Rightarrow \quad x = 0$, so $(0,0)$ (already included above) is a critical point. If $x = \pm 1$ then $(1-2y^2)e^{-1-y^2} = 0 \quad \Rightarrow \quad y = \pm\frac{1}{\sqrt{2}}$, so $\left(\pm 1, \frac{1}{\sqrt{2}}\right)$ and $\left(\pm 1, -\frac{1}{\sqrt{2}}\right)$ are critical points. Now $D\left(\pm 1, \frac{1}{\sqrt{2}}\right) = 8e^{-3} > 0$, $f_{xx}\left(\pm 1, \frac{1}{\sqrt{2}}\right) = -2\sqrt{2}\,e^{-3/2} < 0$ and $D\left(\pm 1, -\frac{1}{\sqrt{2}}\right) = 8e^{-3} > 0$, $f_{xx}\left(\pm 1, -\frac{1}{\sqrt{2}}\right) = 2\sqrt{2}\,e^{-3/2} > 0$, so $f\left(\pm 1, \frac{1}{\sqrt{2}}\right) = \frac{1}{\sqrt{2}}e^{-3/2}$ are local maximum points while $f\left(\pm 1, -\frac{1}{\sqrt{2}}\right) = -\frac{1}{\sqrt{2}}e^{-3/2}$ are local minimum points. At all critical points $(0, y)$ we have $D(0,y) = 0$, so the Second Derivatives Test gives no information. However, if $y > 0$ then $x^2 y e^{-x^2-y^2} \geq 0$ with equality only when $x = 0$, so we have local minimum values $f(0,y) = 0$, $y > 0$. Similarly, if $y < 0$ then $x^2 y e^{-x^2-y^2} \leq 0$ with equality when $x = 0$ so $f(0,y) = 0$, $y < 0$ are local maximum values, and $(0,0)$ is a saddle point.

25. $f(x,y) = x^2 + y^2 + x^{-2}y^{-2}$

From the graphs, there appear to be local minimums of about $f(1, \pm 1) = f(-1, \pm 1) \approx 3$ (and no local maximums or saddle points). $f_x = 2x - 2x^{-3}y^{-2}$, $f_y = 2y - 2x^{-2}y^{-3}$, $f_{xx} = 2 + 6x^{-4}y^{-2}$, $f_{xy} = 4x^{-3}y^{-3}$, $f_{yy} = 2 + 6x^{-2}y^{-4}$. Then

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$f_x = 0$ implies $2x^4y^2 - 2 = 0$ or $x^4y^2 = 1$ or $y^2 = x^{-4}$. Note that neither x nor y can be zero. Now $f_y = 0$ implies $2x^2y^4 - 2 = 0$, and with $y^2 = x^{-4}$ this implies $2x^{-6} - 2 = 0$ or $x^6 = 1$. Thus $x = \pm 1$ and if $x = 1$, $y = \pm 1$; if $x = -1$, $y = \pm 1$. So the critical points are $(1, 1)$, $(1, -1)$,$(-1, 1)$ and $(-1, -1)$. Now $D(1, \pm 1) = D(-1, \pm 1) = 64 - 16 > 0$ and $f_{xx} > 0$ always, so $f(1, \pm 1) = f(-1, \pm 1) = 3$ are local minimums.

26. $f(x, y) = (x - y)e^{-x^2-y^2}$

From the graphs, there appears to be a local maximum of about $f(0.5, -0.5) \approx 0.6$ and a local minimum of about $f(-0.5, 0.5) \approx -0.6$.

$f_x = (x - y)e^{-x^2-y^2}(-2x) + e^{-x^2-y^2}(1) = e^{-x^2-y^2}(1 - 2x^2 + 2xy)$,

$f_y = (x - y)e^{-x^2-y^2}(-2y) + e^{-x^2-y^2}(-1) = -e^{-x^2-y^2}(1 - 2y^2 + 2xy)$, $f_{xx} = 2e^{-x^2-y^2}(2x^3 - 3x + y - 2x^2y)$,

$f_{xy} = 2e^{-x^2-y^2}(x - y + 2x^2y - 2xy^2)$, $f_{yy} = -2e^{-x^2-y^2}(2y^3 - 3y + x - 2xy^2)$. Then $f_x = 0$ implies $1 - 2x^2 + 2xy = 0$ and $f_y = 0$ implies $1 - 2y^2 + 2xy = 0$. Subtracting these two equations gives $-2x^2 + 2y^2 = 0 \quad \Rightarrow \quad y = \pm x$. If $y = x$ then substituting into $f_x = 0$ gives $1 - 2x^2 + 2x^2 = 0$, an impossibility. Substituting $y = -x$ gives $1 - 2x^2 - 2x^2 = 0 \quad \Rightarrow \quad x^2 = \frac{1}{4} \quad \Rightarrow \quad x = \pm\frac{1}{2}$. Thus the critical points are $\left(\frac{1}{2}, -\frac{1}{2}\right)$ and $\left(-\frac{1}{2}, \frac{1}{2}\right)$. Now $D\left(\frac{1}{2}, -\frac{1}{2}\right) = (-3e^{-1/2})(-3e^{-1/2}) - (e^{-1/2})^2 = 8e^{-1} > 0$ with $f_{xx}\left(\frac{1}{2}, -\frac{1}{2}\right) = -3e^{-1/2} < 0$, so $f\left(\frac{1}{2}, -\frac{1}{2}\right) = e^{-1/2} \approx 0.607$ is a local maximum, and $D\left(-\frac{1}{2}, \frac{1}{2}\right) = (3e^{-1/2})(3e^{-1/2}) - (-e^{-1/2})^2 = 8e^{-1} > 0$ with $f_{xx}\left(-\frac{1}{2}, \frac{1}{2}\right) = 3e^{-1/2} > 0$, so $f\left(-\frac{1}{2}, \frac{1}{2}\right) = -e^{-1/2} \approx -0.607$ is a local minimum.

27. $f(x, y) = \sin x + \sin y + \sin(x + y),\ 0 \le x \le 2\pi,\ 0 \le y \le 2\pi$

From the graphs it appears that f has a local maximum at about $(1, 1)$ with value approximately 2.6, a local minimum

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

at about $(5, 5)$ with value approximately -2.6, and a saddle point at about $(3, 3)$.

$f_x = \cos x + \cos(x+y)$, $f_y = \cos y + \cos(x+y)$, $f_{xx} = -\sin x - \sin(x+y)$, $f_{yy} = -\sin y - \sin(x+y)$, $f_{xy} = -\sin(x+y)$. Setting $f_x = 0$ and $f_y = 0$ and subtracting gives $\cos x - \cos y = 0$ or $\cos x = \cos y$. Thus $x = y$ or $x = 2\pi - y$. If $x = y$, $f_x = 0$ becomes $\cos x + \cos 2x = 0$ or $2\cos^2 x + \cos x - 1 = 0$, a quadratic in $\cos x$. Thus $\cos x = -1$ or $\frac{1}{2}$ and $x = \pi$, $\frac{\pi}{3}$, or $\frac{5\pi}{3}$, giving the critical points (π, π), $\left(\frac{\pi}{3}, \frac{\pi}{3}\right)$ and $\left(\frac{5\pi}{3}, \frac{5\pi}{3}\right)$. Similarly if $x = 2\pi - y$, $f_x = 0$ becomes $(\cos x) + 1 = 0$ and the resulting critical point is (π, π). Now $D(x, y) = \sin x \sin y + \sin x \sin(x+y) + \sin y \sin(x+y)$. So $D(\pi, \pi) = 0$ and the Second Derivatives Test doesn't apply. However, along the line $y = x$ we have $f(x, x) = 2\sin x + \sin 2x = 2\sin x + 2\sin x \cos x = 2\sin x(1 + \cos x)$, and $f(x, x) > 0$ for $0 < x < \pi$ while $f(x, x) < 0$ for $\pi < x < 2\pi$. Thus every disk with center (π, π) contains points where f is positive as well as points where f is negative, so the graph crosses its tangent plane $(z = 0)$ there and (π, π) is a saddle point. $D\left(\frac{\pi}{3}, \frac{\pi}{3}\right) = \frac{9}{4} > 0$ and $f_{xx}\left(\frac{\pi}{3}, \frac{\pi}{3}\right) < 0$ so $f\left(\frac{\pi}{3}, \frac{\pi}{3}\right) = \frac{3\sqrt{3}}{2}$ is a local maximum while $D\left(\frac{5\pi}{3}, \frac{5\pi}{3}\right) = \frac{9}{4} > 0$ and $f_{xx}\left(\frac{5\pi}{3}, \frac{5\pi}{3}\right) > 0$, so $f\left(\frac{5\pi}{3}, \frac{5\pi}{3}\right) = -\frac{3\sqrt{3}}{2}$ is a local minimum.

28. $f(x, y) = \sin x + \sin y + \cos(x+y)$, $0 \le x \le \frac{\pi}{4}$, $0 \le y \le \frac{\pi}{4}$

From the graphs, it seems that f has a local maximum at about $(0.5, 0.5)$.

$f_x = \cos x - \sin(x+y)$, $f_y = \cos y - \sin(x+y)$, $f_{xx} = -\sin x - \cos(x+y)$, $f_{yy} = -\sin y - \cos(x+y)$, $f_{xy} = -\cos(x+y)$. Setting $f_x = 0$ and $f_y = 0$ and subtracting gives $\cos x = \cos y$. Thus $x = y$. Substituting $x = y$ into $f_x = 0$ gives $\cos x - \sin 2x = 0$ or $\cos x(1 - 2\sin x) = 0$. But $\cos x \neq 0$ for $0 \le x \le \frac{\pi}{4}$ and $1 - 2\sin x = 0$ implies $x = \frac{\pi}{6}$, so the only critical point is $\left(\frac{\pi}{6}, \frac{\pi}{6}\right)$. Here $f_{xx}\left(\frac{\pi}{6}, \frac{\pi}{6}\right) = -1 < 0$ and $D\left(\frac{\pi}{6}, \frac{\pi}{6}\right) = (-1)^2 - \frac{1}{4} > 0$. Thus $f\left(\frac{\pi}{6}, \frac{\pi}{6}\right) = \frac{3}{2}$ is a local maximum.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. $f(x,y) = x^4 + y^4 - 4x^2y + 2y \quad \Rightarrow \quad f_x(x,y) = 4x^3 - 8xy$ and $f_y(x,y) = 4y^3 - 4x^2 + 2$. $f_x = 0 \quad \Rightarrow$ $4x(x^2 - 2y) = 0$, so $x = 0$ or $x^2 = 2y$. If $x = 0$ then substitution into $f_y = 0$ gives $4y^3 = -2 \quad \Rightarrow \quad y = -\frac{1}{\sqrt[3]{2}}$, so $\left(0, -\frac{1}{\sqrt[3]{2}}\right)$ is a critical point. Substituting $x^2 = 2y$ into $f_y = 0$ gives $4y^3 - 8y + 2 = 0$. Using a graph, solutions are approximately $y = -1.526$, 0.259, and 1.267. (Alternatively, we could have used a calculator or a CAS to find these roots.) We have $x^2 = 2y \quad \Rightarrow \quad x = \pm\sqrt{2y}$, so $y = -1.526$ gives no real-valued solution for x, but $y = 0.259 \quad \Rightarrow \quad x \approx \pm 0.720$ and $y = 1.267 \quad \Rightarrow \quad x \approx \pm 1.592$. Thus to three decimal places, the critical points are $\left(0, -\frac{1}{\sqrt[3]{2}}\right) \approx (0, -0.794)$, $(\pm 0.720, 0.259)$, and $(\pm 1.592, 1.267)$. Now since $f_{xx} = 12x^2 - 8y$, $f_{xy} = -8x$, $f_{yy} = 12y^2$, and $D = (12x^2 - 8y)(12y^2) - 64x^2$, we have $D(0, -0.794) > 0$, $f_{xx}(0, -0.794) > 0$, $D(\pm 0.720, 0.259) < 0$, $D(\pm 1.592, 1.267) > 0$, and $f_{xx}(\pm 1.592, 1.267) > 0$. Therefore $f(0, -0.794) \approx -1.191$ and $f(\pm 1.592, 1.267) \approx -1.310$ are local minimums, and $(\pm 0.720, 0.259)$ are saddle points. There is no highest point on the graph, but the lowest points are approximately $(\pm 1.592, 1.267, -1.310)$.

30. $f(x,y) = y^6 - 2y^4 + x^2 - y^2 + y \quad \Rightarrow \quad f_x(x,y) = 2x$ and $f_y(x,y) = 6y^5 - 8y^3 - 2y + 1$. $\quad f_x = 0$ implies $x = 0$, and the graph of f_y shows that the roots of $f_y = 0$ are approximately $y = -1.273$, 0.347, and 1.211. (Alternatively, we could have found the roots of $f_y = 0$ directly, using a calculator or CAS.) So to three decimal places, the critical points are $(0, -1.273)$, $(0, 0.347)$, and $(0, 1.211)$. Now since $f_{xx} = 2$, $f_{xy} = 0$, $f_{yy} = 30y^4 - 24y^2 - 2$, and $D = 60y^4 - 48y^2 - 4$, we have $D(0, -1.273) > 0$, $f_{xx}(0, -1.273) > 0$, $D(0, 0.347) < 0$, $D(0, 1.211) > 0$, and $f_{xx}(0, 1.211) > 0$, so $f(0, -1.273) \approx -3.890$ and $f(0, 1.211) \approx -1.403$ are local minimums, and $(0, 0.347)$ is a saddle point. The lowest point on the graph is approximately $(0, -1.273, -3.890)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. $f(x,y) = x^4 + y^3 - 3x^2 + y^2 + x - 2y + 1 \quad \Rightarrow \quad f_x(x,y) = 4x^3 - 6x + 1$ and $f_y(x,y) = 3y^2 + 2y - 2$. From the graphs, we see that to three decimal places, $f_x = 0$ when $x \approx -1.301$, 0.170, or 1.131, and $f_y = 0$ when $y \approx -1.215$ or 0.549. (Alternatively, we could have used a calculator or a CAS to find these roots. We could also use the quadratic formula to find the solutions of $f_y = 0$.) So, to three decimal places, f has critical points at $(-1.301, -1.215)$, $(-1.301, 0.549)$, $(0.170, -1.215)$, $(0.170, 0.549)$, $(1.131, -1.215)$, and $(1.131, 0.549)$. Now since $f_{xx} = 12x^2 - 6$, $f_{xy} = 0$, $f_{yy} = 6y + 2$, and $D = (12x^2 - 6)(6y + 2)$, we have $D(-1.301, -1.215) < 0$, $D(-1.301, 0.549) > 0$, $f_{xx}(-1.301, 0.549) > 0$, $D(0.170, -1.215) > 0$, $f_{xx}(0.170, -1.215) < 0$, $D(0.170, 0.549) < 0$, $D(1.131, -1.215) < 0$, $D(1.131, 0.549) > 0$, and $f_{xx}(1.131, 0.549) > 0$. Therefore, to three decimal places, $f(-1.301, 0.549) \approx -3.145$ and $f(1.131, 0.549) \approx -0.701$ are local minimums, $f(0.170, -1.215) \approx 3.197$ is a local maximum, and $(-1.301, -1.215)$, $(0.170, 0.549)$, and $(1.131, -1.215)$ are saddle points. There is no highest or lowest point on the graph.

32. $f(x,y) = 20e^{-x^2-y^2} \sin 3x \cos 3y \quad \Rightarrow$

$$\begin{aligned} f_x(x,y) &= 20\cos 3y \left[e^{-x^2-y^2}(3\cos 3x) + (\sin 3x)e^{-x^2-y^2}(-2x)\right] \\ &= 20e^{-x^2-y^2} \cos 3y\,(3\cos 3x - 2x\sin 3x) \\ f_y(x,y) &= 20\sin 3x \left[e^{-x^2-y^2}(-3\sin 3y) + (\cos 3y)e^{-x^2-y^2}(-2y)\right] \\ &= -20e^{-x^2-y^2} \sin 3x\,(3\sin 3y + 2y\cos 3y) \end{aligned}$$

Now $f_x = 0$ implies $\cos 3y = 0$ or $3\cos 3x - 2x\sin 3x = 0$. For $|y| \le 1$, the solutions to $\cos 3y = 0$ are $y = \pm\frac{\pi}{6} \approx \pm 0.524$. Using a graph (or a calculator or CAS), we estimate the roots of $3\cos 3x - 2x\sin 3x$ for $|x| \le 1$ to be $x \approx \pm 0.430$. $f_y = 0$ implies $\sin 3x = 0$, so $x = 0$, or $3\sin 3y + 2y\cos 3y = 0$. From a graph (or calculator or CAS), the roots of $3\sin 3y + 2y\cos 3y$ between -1 and 1 are approximately 0 and ± 0.872. So to three decimal places, f has critical points at $(\pm 0.430, 0)$, $(0.430, \pm 0.872)$, $(-0.430, \pm 0.872)$, and $(0, \pm 0.524)$. Now

$$\begin{aligned} f_{xx} &= 20e^{-x^2-y^2} \cos 3y[(4x^2 - 11)\sin 3x - 12x\cos 3x] \\ f_{xy} &= -20e^{-x^2-y^2}(3\cos 3x - 2x\sin 3x)(3\sin 3y + 2y\cos 3y) \\ f_{yy} &= 20e^{-x^2-y^2} \sin 3x[(4y^2 - 11)\cos 3y - 12y\sin 3y] \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and $D = f_{xx}f_{yy} - f_{xy}^2$. Then $D(\pm 0.430, 0) > 0$, $f_{xx}(0.430, 0) < 0$, $f_{xx}(-0.430, 0) > 0$, $D(0.430, \pm 0.872) > 0$, $f_{xx}(0.430, \pm 0.872) > 0$, $D(-0.430, \pm 0.872) > 0$, $f_{xx}(-0.430, \pm 0.872) < 0$, and $D(0, \pm 0.524) < 0$, so $f(0.430, 0) \approx 15.973$ and $f(-0.430, \pm 0.872) \approx 6.459$ are local maximums, $f(-0.430, 0) \approx -15.973$ and $f(0.430, \pm 0.872) \approx -6.459$ are local minimums, and $(0, \pm 0.524)$ are saddle points. The highest point on the graph is approximately $(0.430, 0, 15.973)$ and the lowest point is approximately $(-0.430, 0, -15.973)$.

33. Since f is a polynomial it is continuous on D, so an absolute maximum and minimum exist. Here $f_x = 2x - 2$, $f_y = 2y$, and setting $f_x = f_y = 0$ gives $(1, 0)$ as the only critical point (which is inside D), where $f(1, 0) = -1$. Along L_1: $x = 0$ and $f(0, y) = y^2$ for $-2 \le y \le 2$, a quadratic function which attains its minimum at $y = 0$, where $f(0, 0) = 0$, and its maximum at $y = \pm 2$, where $f(0, \pm 2) = 4$. Along L_2: $y = x - 2$ for $0 \le x \le 2$, and $f(x, x - 2) = 2x^2 - 6x + 4 = 2\left(x - \frac{3}{2}\right)^2 - \frac{1}{2}$, a quadratic which attains its minimum at $x = \frac{3}{2}$, where $f\left(\frac{3}{2}, -\frac{1}{2}\right) = -\frac{1}{2}$, and its maximum at $x = 0$, where $f(0, -2) = 4$. Along L_3: $y = 2 - x$ for $0 \le x \le 2$, and $f(x, 2 - x) = 2x^2 - 6x + 4 = 2\left(x - \frac{3}{2}\right)^2 - \frac{1}{2}$, a quadratic which attains its minimum at $x = \frac{3}{2}$, where $f\left(\frac{3}{2}, \frac{1}{2}\right) = -\frac{1}{2}$, and its maximum at $x = 0$, where $f(0, 2) = 4$. Thus the absolute maximum of f on D is $f(0, \pm 2) = 4$ and the absolute minimum is $f(1, 0) = -1$.

34. Since f is a polynomial it is continuous on D, so an absolute maximum and minimum exist. $f_x = 1 - y$, $f_y = 1 - x$, and setting $f_x = f_y = 0$ gives $(1, 1)$ as the only critical point (which is inside D), where $f(1, 1) = 1$. Along L_1: $y = 0$ and $f(x, 0) = x$ for $0 \le x \le 4$, an increasing function in x, so the maximum value is $f(4, 0) = 4$ and the minimum value is $f(0, 0) = 0$. Along L_2: $y = 2 - \frac{1}{2}x$ and $f\left(x, 2 - \frac{1}{2}x\right) = \frac{1}{2}x^2 - \frac{3}{2}x + 2 = \frac{1}{2}\left(x - \frac{3}{2}\right)^2 + \frac{7}{8}$ for $0 \le x \le 4$, a quadratic function which has a minimum at $x = \frac{3}{2}$, where $f\left(\frac{3}{2}, \frac{5}{4}\right) = \frac{7}{8}$, and a maximum at $x = 4$, where $f(4, 0) = 4$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Along L_3: $x = 0$ and $f(0, y) = y$ for $0 \le y \le 2$, an increasing function in y, so the maximum value is $f(0, 2) = 2$ and the minimum value is $f(0, 0) = 0$. Thus the absolute maximum of f on D is $f(4, 0) = 4$ and the absolute minimum is $f(0, 0) = 0$.

35. $f(x, y) = x^2 + y^2 + x^2y + 4 \quad \Rightarrow \quad f_x(x, y) = 2x + 2xy$, $f_y(x, y) = 2y + x^2$, and setting $f_x = f_y = 0$ gives $(0, 0)$ as the only critical point in D, with $f(0, 0) = 4$.

On L_1: $y = -1$, $f(x, -1) = 5$, a constant.

On L_2: $x = 1$, $f(1, y) = y^2 + y + 5$, a quadratic in y which attains its maximum at $(1, 1)$, $f(1, 1) = 7$ and its minimum at $\left(1, -\frac{1}{2}\right)$, $f\left(1, -\frac{1}{2}\right) = \frac{19}{4}$.

On L_3: $f(x, 1) = 2x^2 + 5$ which attains its maximum at $(-1, 1)$ and $(1, 1)$ with $f(\pm 1, 1) = 7$ and its minimum at $(0, 1)$, $f(0, 1) = 5$.

On L_4: $f(-1, y) = y^2 + y + 5$ with maximum at $(-1, 1)$, $f(-1, 1) = 7$ and minimum at $\left(-1, -\frac{1}{2}\right)$, $f\left(-1, -\frac{1}{2}\right) = \frac{19}{4}$.

Thus the absolute maximum is attained at both $(\pm 1, 1)$ with $f(\pm 1, 1) = 7$ and the absolute minimum on D is attained at $(0, 0)$ with $f(0, 0) = 4$.

36. $f(x, y) = x^2 + xy + y^2 - 6y \quad \Rightarrow \quad f_x = 2x + y, \;\; f_y = x + 2y - 6$. Then $f_x = 0$ implies $y = -2x$, and substituting into $f_y = 0$ gives $x - 4x - 6 = 0 \quad \Rightarrow \quad x = -2$, so the only critical point is $(-2, 4)$ (which is in D) where $f(-2, 4) = -12$.

Along L_1: $y = 0$, so $f(x, 0) = x^2$, $-3 \le x \le 3$, which has a maximum value at $x = \pm 3$ where $f(\pm 3, 0) = 9$ and a minimum value at $x = 0$, where $f(0, 0) = 0$. Along L_2: $x = 3$, so $f(3, y) = 9 - 3y + y^2 = \left(y - \frac{3}{2}\right)^2 + \frac{27}{4}$, $0 \le y \le 5$, which has a maximum value at $y = 5$ where $f(3, 5) = 19$ and a minimum value at $y = \frac{3}{2}$ where $f(3, \frac{3}{2}) = \frac{27}{4}$.

Along L_3: $y = 5$, so $f(x, 5) = x^2 + 5x - 5 = \left(x + \frac{5}{2}\right)^2 - \frac{45}{4}$, $-3 \le x \le 3$, which has a maximum value at $x = 3$ where $f(3, 5) = 19$ and a minimum value at $x = -\frac{5}{2}$, where $f\left(-\frac{5}{2}, 5\right) = -\frac{45}{4}$. Along L_4: $x = -3$, so $f(-3, y) = 9 - 9y + y^2 = \left(y - \frac{9}{2}\right)^2 - \frac{45}{4}$, $0 \le y \le 5$, which has a maximum value at $y = 0$ where $f(-3, 0) = 9$ and a minimum value at $y = \frac{9}{2}$ where $f\left(-3, \frac{9}{2}\right) = -\frac{45}{4}$. Thus the absolute maximum of f on D is $f(3, 5) = 19$ and the absolute minimum is $f(-2, 4) = -12$.

37. $f(x, y) = x^2 + 2y^2 - 2x - 4y + 1 \quad \Rightarrow \quad f_x = 2x - 2, \;\; f_y = 4y - 4$. Setting $f_x = 0$ and $f_y = 0$ gives $(1, 1)$ as the only critical point (which is inside D), where $f(1, 1) = -2$. Along L_1: $y = 0$, so $f(x, 0) = x^2 - 2x + 1 = (x - 1)^2$, $0 \le x \le 2$, which has a maximum value both at $x = 0$ and $x = 2$ where $f(0, 0) = f(2, 0) = 1$ and a minimum value at $x = 1$, where $f(1, 0) = 0$. Along L_2: $x = 2$, so $f(2, y) = 2y^2 - 4y + 1 = 2(y - 1)^2 - 1$, $0 \le y \le 3$, which has a maximum value at

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$y = 3$ where $f(2,3) = 7$ and a minimum value at $y = 1$ where $f(2,1) = -1$. Along L_3: $y = 3$, so $f(x,3) = x^2 - 2x + 7 = (x-1)^2 + 6$, $0 \le x \le 2$, which has a maximum value both at $x = 0$ and $x = 2$ where $f(0,3) = f(2,3) = 7$ and a minimum value at $x = 1$, where $f(1,3) = 6$. Along L_4: $x = 0$, so $f(0,y) = 2y^2 - 4y + 1 = 2(y-1)^2 - 1$, $0 \le y \le 3$, which has a maximum value at $y = 3$ where $f(0,3) = 7$ and a minimum value at $y = 1$ where $f(0,1) = -1$. Thus the absolute maximum is attained at both $(0,3)$ and $(2,3)$, where $f(0,3) = f(2,3) = 7$, and the absolute minimum is $f(1,1) = -2$.

38. $f(x,y) = xy^2 \quad \Rightarrow \quad f_x = y^2$ and $f_y = 2xy$, and since $f_x = 0 \quad \Leftrightarrow$ $y = 0$, there are no critical points in the interior of D. Along L_1: $y = 0$ and $f(x,0) = 0$. Along L_2: $x = 0$ and $f(0,y) = 0$. Along L_3: $y = \sqrt{3 - x^2}$, so let $g(x) = f\left(x, \sqrt{3-x^2}\right) = 3x - x^3$ for $0 \le x \le \sqrt{3}$. Then $g'(x) = 3 - 3x^2 = 0 \quad \Leftrightarrow \quad x = 1$. The maximum value is $f\left(1, \sqrt{2}\right) = 2$

and the minimum occurs both at $x = 0$ and $x = \sqrt{3}$ where $f\left(0, \sqrt{3}\right) = f\left(\sqrt{3}, 0\right) = 0$. Thus the absolute maximum of f on D is $f\left(1, \sqrt{2}\right) = 2$, and the absolute minimum is 0 which occurs at all points along L_1 and L_2.

39. $f(x,y) = 2x^3 + y^4 \quad \Rightarrow \quad f_x(x,y) = 6x^2$ and $f_y(x,y) = 4y^3$. And so $f_x = 0$ and $f_y = 0$ only occur when $x = y = 0$. Hence, the only critical point inside the disk is at $x = y = 0$ where $f(0,0) = 0$. Now on the circle $x^2 + y^2 = 1$, $y^2 = 1 - x^2$ so let $g(x) = f(x,y) = 2x^3 + (1-x^2)^2 = x^4 + 2x^3 - 2x^2 + 1$, $-1 \le x \le 1$. Then $g'(x) = 4x^3 + 6x^2 - 4x = 0 \quad \Rightarrow$ $x = 0, -2$, or $\frac{1}{2}$. $f(0, \pm 1) = g(0) = 1$, $f\left(\frac{1}{2}, \pm\frac{\sqrt{3}}{2}\right) = g\left(\frac{1}{2}\right) = \frac{13}{16}$, and $(-2,-3)$ is not in D. Checking the endpoints, we get $f(-1,0) = g(-1) = -2$ and $f(1,0) = g(1) = 2$. Thus the absolute maximum and minimum of f on D are $f(1,0) = 2$ and $f(-1,0) = -2$.

Another method: On the boundary $x^2 + y^2 = 1$ we can write $x = \cos\theta$, $y = \sin\theta$, so $f(\cos\theta, \sin\theta) = 2\cos^3\theta + \sin^4\theta$, $0 \le \theta \le 2\pi$.

40. $f(x,y) = x^3 - 3x - y^3 + 12y \quad \Rightarrow \quad f_x(x,y) = 3x^2 - 3$ and $f_y(x,y) = -3y^2 + 12$ and the critical points are $(1,2)$, $(1,-2)$, $(-1,2)$, and $(-1,-2)$. But only $(1,2)$ and $(-1,2)$ are in D and $f(1,2) = 14$, $f(-1,2) = 18$. Along L_1: $x = -2$ and $f(-2,y) = -2 - y^3 + 12y$, $-2 \le y \le 3$, which has a maximum at $y = 2$ where $f(-2,2) = 14$ and a minimum at $y = -2$ where $f(-2,-2) = -18$. Along L_2: $x = 2$ and $f(2,y) = 2 - y^3 + 12y$, $2 \le y \le 3$, which has a maximum at $y = 2$ where $f(2,2) = 18$ and a minimum at $y = 3$ where $f(2,3) = 11$. Along L_3: $y = 3$ and $f(x,3) = x^3 - 3x + 9$, $-2 \le x \le 2$, which has a maximum at $x = -1$ and $x = 2$ where $f(-1,3) = f(2,3) = 11$ and a minimum at $x = 1$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and $x = -2$ where $f(1,3) = f(-2,3) = 7$. Along L_4: $y = x$ and $f(x,x) = 9x$, $-2 \le x \le 2$, which has a maximum at $x = 2$ where $f(2,2) = 18$ and a minimum at $x = -2$ where $f(-2,-2) = -18$. So the absolute maximum value of f on D is $f(2,2) = 18$ and the minimum is $f(-2,-2) = -18$.

41. $f(x,y) = -(x^2-1)^2 - (x^2y - x - 1)^2 \quad \Rightarrow \quad f_x(x,y) = -2(x^2-1)(2x) - 2(x^2y - x - 1)(2xy - 1)$ and $f_y(x,y) = -2(x^2y - x - 1)x^2$. Setting $f_y(x,y) = 0$ gives either $x = 0$ or $x^2y - x - 1 = 0$.

There are no critical points for $x = 0$, since $f_x(0,y) = -2$, so we set $x^2y - x - 1 = 0 \quad \Leftrightarrow \quad y = \dfrac{x+1}{x^2} \quad [x \ne 0]$,

so $f_x\left(x, \dfrac{x+1}{x^2}\right) = -2(x^2-1)(2x) - 2\left(x^2\,\dfrac{x+1}{x^2} - x - 1\right)\left(2x\,\dfrac{x+1}{x^2} - 1\right) = -4x(x^2-1)$. Therefore $f_x(x,y) = f_y(x,y) = 0$ at the points $(1,2)$ and $(-1,0)$. To classify these critical points, we calculate $f_{xx}(x,y) = -12x^2 - 12x^2y^2 + 12xy + 4y + 2$, $f_{yy}(x,y) = -2x^4$, and $f_{xy}(x,y) = -8x^3y + 6x^2 + 4x$. In order to use the Second Derivatives Test we calculate

$D(-1,0) = f_{xx}(-1,0)\,f_{yy}(-1,0) - [f_{xy}(-1,0)]^2 = 16 > 0$,

$f_{xx}(-1,0) = -10 < 0$, $D(1,2) = 16 > 0$, and $f_{xx}(1,2) = -26 < 0$, so both $(-1,0)$ and $(1,2)$ give local maximums.

42. $f(x,y) = 3xe^y - x^3 - e^{3y}$ is differentiable everywhere, so the requirement for critical points is that $f_x = 3e^y - 3x^2 = 0$ **(1)** and $f_y = 3xe^y - 3e^{3y} = 0$ **(2)**. From **(1)** we obtain $e^y = x^2$, and then **(2)** gives $3x^3 - 3x^6 = 0 \quad \Rightarrow \quad x = 1$ or 0, but only $x = 1$ is valid, since $x = 0$ makes **(1)** impossible. So substituting $x = 1$ into **(1)** gives $y = 0$, and the only critical point is $(1,0)$.

The Second Derivatives Test shows that this gives a local maximum, since $D(1,0) = \left[-6x(3xe^y - 9e^{3y}) - (3e^y)^2\right]_{(1,0)} = 27 > 0$ and $f_{xx}(1,0) = [-6x]_{(1,0)} = -6 < 0$. But $f(1,0) = 1$ is not an absolute maximum because, for instance, $f(-3,0) = 17$. This can also be seen from the graph.

43. Let d be the distance from $(2,0,-3)$ to any point (x,y,z) on the plane $x + y + z = 1$, so $d = \sqrt{(x-2)^2 + y^2 + (z+3)^2}$ where $z = 1 - x - y$, and we minimize $d^2 = f(x,y) = (x-2)^2 + y^2 + (4 - x - y)^2$. Then $f_x(x,y) = 2(x-2) + 2(4-x-y)(-1) = 4x + 2y - 12$, $f_y(x,y) = 2y + 2(4-x-y)(-1) = 2x + 4y - 8$. Solving $4x + 2y - 12 = 0$ and $2x + 4y - 8 = 0$ simultaneously gives $x = \frac{8}{3}$, $y = \frac{2}{3}$, so the only critical point is $\left(\frac{8}{3}, \frac{2}{3}\right)$. An absolute

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

minimum exists (since there is a minimum distance from the point to the plane) and it must occur at a critical point, so the shortest distance occurs for $x = \frac{8}{3}$, $y = \frac{2}{3}$ for which $d = \sqrt{\left(\frac{8}{3} - 2\right)^2 + \left(\frac{2}{3}\right)^2 + \left(4 - \frac{8}{3} - \frac{2}{3}\right)^2} = \sqrt{\frac{4}{3}} = \frac{2}{\sqrt{3}}$.

44. Here the distance d from a point on the plane to the point $(0, 1, 1)$ is $d = \sqrt{x^2 + (y-1)^2 + (z-1)^2}$, where $z = 2 - \frac{1}{3}x + \frac{2}{3}y$. We can minimize $d^2 = f(x, y) = x^2 + (y-1)^2 + \left(1 - \frac{1}{3}x + \frac{2}{3}y\right)^2$, so $f_x(x, y) = 2x + 2\left(1 - \frac{1}{3}x + \frac{2}{3}y\right)\left(-\frac{1}{3}\right) = \frac{20}{9}x - \frac{4}{9}y - \frac{2}{3}$ and $f_y(x, y) = 2(y-1) + 2\left(1 - \frac{1}{3}x + \frac{2}{3}y\right)\left(\frac{2}{3}\right) = -\frac{4}{9}x + \frac{26}{9}y - \frac{2}{3}$. Solving $\frac{20}{9}x - \frac{4}{9}y - \frac{2}{3} = 0$ and $-\frac{4}{9}x + \frac{26}{9}y - \frac{2}{3} = 0$ simultaneously gives $x = \frac{5}{14}$ and $y = \frac{2}{7}$, so the only critical point is $\left(\frac{5}{14}, \frac{2}{7}\right)$.

This point must correspond to the minimum distance, so the point on the plane closest to $(0, 1, 1)$ is $\left(\frac{5}{14}, \frac{2}{7}, \frac{29}{14}\right)$.

45. Let d be the distance from the point $(4, 2, 0)$ to any point (x, y, z) on the cone, so $d = \sqrt{(x-4)^2 + (y-2)^2 + z^2}$ where $z^2 = x^2 + y^2$, and we minimize $d^2 = (x-4)^2 + (y-2)^2 + x^2 + y^2 = f(x, y)$. Then $f_x(x, y) = 2(x-4) + 2x = 4x - 8$, $f_y(x, y) = 2(y-2) + 2y = 4y - 4$, and the critical points occur when $f_x = 0 \quad \Rightarrow \quad x = 2,\ f_y = 0 \quad \Rightarrow \quad y = 1$. Thus, the only critical point is $(2, 1)$. An absolute minimum exists (since there is a minimum distance from the cone to the point) which must occur at a critical point, so the points on the cone closest to $(4, 2, 0)$ are $\left(2, 1, \pm\sqrt{5}\right)$.

46. The distance from the origin to a point (x, y, z) on the surface is $d = \sqrt{x^2 + y^2 + z^2}$ where $y^2 = 9 + xz$, so we minimize $d^2 = x^2 + 9 + xz + z^2 = f(x, z)$. Then $f_x = 2x + z$, $f_z = x + 2z$, and $f_x = 0$, $f_z = 0 \quad \Rightarrow \quad x = 0$, $z = 0$, so the only critical point is $(0, 0)$. $D(0, 0) = (2)(2) - 1 = 3 > 0$ with $f_{xx}(0, 0) = 2 > 0$, so this is a minimum. Thus, $y^2 = 9 + 0 \quad \Rightarrow \quad y = \pm 3$ and the points on the surface closest to the origin are $(0, \pm 3, 0)$.

47. Let x, y, z be the positive numbers. Then $x + y + z = 100 \quad \Rightarrow \quad z = 100 - x - y$, and we want to maximize $xyz = xy(100 - x - y) = 100xy - x^2y - xy^2 = f(x, y)$ for $0 < x, y, z < 100$. $f_x = 100y - 2xy - y^2$, $f_y = 100x - x^2 - 2xy$, $f_{xx} = -2y$, $f_{yy} = -2x$, $f_{xy} = 100 - 2x - 2y$. Then $f_x = 0$ implies $y(100 - 2x - y) = 0 \quad \Rightarrow$ $y = 100 - 2x$ (since $y > 0$). Substituting into $f_y = 0$ gives $x[100 - x - 2(100 - 2x)] = 0 \quad \Rightarrow \quad 3x - 100 = 0$ (since $x > 0$) $\quad \Rightarrow \quad x = \frac{100}{3}$. Then $y = 100 - 2\left(\frac{100}{3}\right) = \frac{100}{3}$, and the only critical point is $\left(\frac{100}{3}, \frac{100}{3}\right)$. $D\left(\frac{100}{3}, \frac{100}{3}\right) = \left(-\frac{200}{3}\right)\left(-\frac{200}{3}\right) - \left(-\frac{100}{3}\right)^2 = \frac{10{,}000}{3} > 0$ and $f_{xx}\left(\frac{100}{3}, \frac{100}{3}\right) = -\frac{200}{3} < 0$. Thus $f\left(\frac{100}{3}, \frac{100}{3}\right)$ is a local maximum. It is also the absolute maximum (compare to the values of f as x, y, or $z \to 0$ or 100), so the numbers are $x = y = z = \frac{100}{3}$.

48. Let x, y, z, be the positive numbers. Then $x + y + z = 12$ and we want to minimize $x^2 + y^2 + z^2 = x^2 + y^2 + (12 - x - y)^2 = f(x, y)$ for $0 < x, y < 12$. $f_x = 2x + 2(12 - x - y)(-1) = 4x + 2y - 24$, $f_y = 2y + 2(12 - x - y)(-1) = 2x + 4y - 24$, $f_{xx} = 4$, $f_{xy} = 2$, $f_{yy} = 4$. Then $f_x = 0$ implies $4x + 2y = 24$ or $y = 12 - 2x$ and substituting into $f_y = 0$ gives $2x + 4(12 - 2x) = 24 \quad \Rightarrow \quad 6x = 24 \quad \Rightarrow \quad x = 4$ and then $y = 4$, so

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

the only critical point is $(4,4)$. $D(4,4) = 16 - 4 > 0$ and $f_{xx}(4,4) = 4 > 0$, so $f(4,4)$ is a local minimum. $f(4,4)$ is also the absolute minimum [compare to the values of f as $x, y \to 0$ or 12] so the numbers are $x = y = z = 4$.

49. Center the sphere at the origin so that its equation is $x^2 + y^2 + z^2 = r^2$, and orient the inscribed rectangular box so that its edges are parallel to the coordinate axes. Any vertex of the box satisfies $x^2 + y^2 + z^2 = r^2$, so take (x, y, z) to be the vertex in the first octant. Then the box has length $2x$, width $2y$, and height $2z = 2\sqrt{r^2 - x^2 - y^2}$ with volume given by $V(x,y) = (2x)(2y)\left(2\sqrt{r^2 - x^2 - y^2}\right) = 8xy\sqrt{r^2 - x^2 - y^2}$ for $0 < x < r$, $0 < y < r$. Then

$V_x = (8xy) \cdot \frac{1}{2}(r^2 - x^2 - y^2)^{-1/2}(-2x) + \sqrt{r^2 - x^2 - y^2} \cdot 8y = \dfrac{8y(r^2 - 2x^2 - y^2)}{\sqrt{r^2 - x^2 - y^2}}$ and $V_y = \dfrac{8x(r^2 - x^2 - 2y^2)}{\sqrt{r^2 - x^2 - y^2}}$.

Setting $V_x = 0$ gives $y = 0$ or $2x^2 + y^2 = r^2$, but $y > 0$ so only the latter solution applies. Similarly, $V_y = 0$ with $x > 0$ implies $x^2 + 2y^2 = r^2$. Substituting, we have $2x^2 + y^2 = x^2 + 2y^2 \;\Rightarrow\; x^2 = y^2 \;\Rightarrow\; y = x$. Then $x^2 + 2y^2 = r^2 \;\Rightarrow\; 3x^2 = r^2 \;\Rightarrow\; x = \sqrt{r^2/3} = r/\sqrt{3} = y$. Thus the only critical point is $\left(r/\sqrt{3}, r/\sqrt{3}\right)$. There must be a maximum volume and here it must occur at a critical point, so the maximum volume occurs when $x = y = r/\sqrt{3}$ and the maximum volume is $V\left(\frac{r}{\sqrt{3}}, \frac{r}{\sqrt{3}}\right) = 8\left(\frac{r}{\sqrt{3}}\right)\left(\frac{r}{\sqrt{3}}\right)\sqrt{r^2 - \left(\frac{r}{\sqrt{3}}\right)^2 - \left(\frac{r}{\sqrt{3}}\right)^2} = \dfrac{8}{3\sqrt{3}}\,r^3$.

50. Let x, y, and z be the dimensions of the box. We wish to minimize surface area $= 2xy + 2xz + 2yz$, but we have volume $= xyz = 1000 \;\Rightarrow\; z = \dfrac{1000}{xy}$ so we minimize

$f(x,y) = 2xy + 2x\left(\dfrac{1000}{xy}\right) + 2y\left(\dfrac{1000}{xy}\right) = 2xy + \dfrac{2000}{y} + \dfrac{2000}{x}$. Then $f_x = 2y - \dfrac{2000}{x^2}$ and $f_y = 2x - \dfrac{2000}{y^2}$. Setting $f_x = 0$ implies $y = \dfrac{1000}{x^2}$ and substituting into $f_y = 0$ gives $x - \dfrac{x^4}{1000} = 0 \;\Rightarrow\; x^3 = 1000$ [since $x \neq 0$] $\;\Rightarrow\; x = 10$. The surface area has a minimum but no maximum and it must occur at a critical point, so the minimal surface area occurs for a box with dimensions $x = 10$ cm, $y = 1000/10^2 = 10$ cm, $z = 1000/10^2 = 10$ cm.

51. The volume of the box is $V = xyz$. Since one vertex is in the plane $x + 2y + 3z = 6 \;\Leftrightarrow\; z = \frac{1}{3}(6 - x - 2y)$, the volume is given by $V(x,y) = \frac{1}{3}xy(6 - x - 2y) = \frac{1}{3}(6xy - x^2y - 2xy^2)$. Now maximize V.

$V_x = \frac{1}{3}(6y - 2xy - 2y^2) = \frac{1}{3}y(6 - 2x - 2y)$ and $V_y = \frac{1}{3}x\,(6 - x - 4y)$. Setting $f_x = 0$ and $f_y = 0$ gives $y = 3 - x$ and $x = 6 - 4y \;\Rightarrow\; y = 1$ and $x = 2$, so the critical point is $(2,1)$, which geometrically must give a maximum. Thus, the volume of the largest such box is $V = (2)(1)\left(\frac{2}{3}\right) = \frac{4}{3}$.

52. Surface area $= 2(xy + xz + yz) = 64$ cm^2, so $xy + xz + yz = 32$ or $z = \dfrac{32 - xy}{x + y}$. Maximize the volume

$f(x,y) = xy\,\dfrac{32 - xy}{x + y}$. Then $f_x = \dfrac{32y^2 - 2xy^3 - x^2y^2}{(x+y)^2} = y^2\,\dfrac{32 - 2xy - x^2}{(x+y)^2}$ and $f_y = x^2\,\dfrac{32 - 2xy - y^2}{(x+y)^2}$. Setting $f_x = 0$ implies $y = \dfrac{32 - x^2}{2x}$ and substituting into $f_y = 0$ gives $32(4x^2) - (32 - x^2)(4x^2) - (32 - x^2)^2 = 0$ or

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$3x^4 + 64x^2 - (32)^2 = 0$. Thus, $x^2 = \frac{64}{6}$ or $x = \frac{8}{\sqrt{6}}$, $y = \frac{64/3}{16/\sqrt{6}} = \frac{8}{\sqrt{6}}$ and $z = \frac{8}{\sqrt{6}}$. Thus, the box is a cube with edge length $\frac{8}{\sqrt{6}}$ cm.

53. Let the dimensions be x, y, and z; then $4x + 4y + 4z = c$ and the volume is $V = xyz = xy\left(\frac{1}{4}c - x - y\right) = \frac{1}{4}cxy - x^2y - xy^2$, $x > 0$, $y > 0$. Then $V_x = \frac{1}{4}cy - 2xy - y^2$ and $V_y = \frac{1}{4}cx - x^2 - 2xy$, so $V_x = 0 = V_y$ when $2x + y = \frac{1}{4}c$ and $x + 2y = \frac{1}{4}c$. Solving, we get $x = \frac{1}{12}c$, $y = \frac{1}{12}c$ and $z = \frac{1}{4}c - x - y = \frac{1}{12}c$. From the geometrical nature of the problem, this critical point must give an absolute maximum. Thus, the box is a cube with edge length $\frac{1}{12}c$.

54. The cost equals $5xy + 2(xz + yz)$ and $xyz = V$, so $C(x, y) = 5xy + 2V(x + y)/(xy) = 5xy + 2V(x^{-1} + y^{-1})$. Then $C_x = 5y - 2Vx^{-2}$, $C_y = 5x - 2Vy^{-2}$, $C_x = 0$ implies $y = 2V/(5x^2)$, $C_y = 0$ implies $x = \sqrt[3]{\frac{2}{5}V} = y$. Thus, the dimensions of the aquarium which minimize the cost are $x = y = \sqrt[3]{\frac{2}{5}V}$ units, $z = V^{1/3}\left(\frac{5}{2}\right)^{2/3}$.

55. Let the dimensions be x, y and z, then minimize $xy + 2(xz + yz)$ if $xyz = 32{,}000$ cm^3. Then $f(x, y) = xy + [64{,}000(x + y)/xy] = xy + 64{,}000(x^{-1} + y^{-1})$, $f_x = y - 64{,}000x^{-2}$, $f_y = x - 64{,}000y^{-2}$. And $f_x = 0$ implies $y = 64{,}000/x^2$; substituting into $f_y = 0$ implies $x^3 = 64{,}000$ or $x = 40$ and then $y = 40$. Now $D(x, y) = [(2)(64{,}000)]^2x^{-3}y^{-3} - 1 > 0$ for $(40, 40)$ and $f_{xx}(40, 40) > 0$ so this is indeed a minimum. Thus, the dimensions of the box are $x = y = 40$ cm, $z = 20$ cm.

56. Let x be the length of the north and south walls, y the length of the east and west walls, and z the height of the building. The heat loss is given by $h = 10(2yz) + 8(2xz) + 1(xy) + 5(xy) = 6xy + 16xz + 20yz$. The volume is 4000 m^3, so $xyz = 4000$, and we substitute $z = \frac{4000}{xy}$ to obtain the heat loss function $h(x, y) = 6xy + 80{,}000/x + 64{,}000/y$.

(a) Since $z = \frac{4000}{xy} \geq 4$, $xy \leq 1000 \quad \Rightarrow \quad y \leq 1000/x$. Also $x \geq 30$ and $y \geq 30$, so the domain of h is $D = \{(x, y) \mid x \geq 30, 30 \leq y \leq 1000/x\}$.

(b) $h(x, y) = 6xy + 80{,}000x^{-1} + 64{,}000y^{-1} \quad \Rightarrow$

$h_x = 6y - 80{,}000x^{-2}$, $h_y = 6x - 64{,}000y^{-2}$.

$h_x = 0$ implies $6x^2y = 80{,}000 \quad \Rightarrow \quad y = \dfrac{80{,}000}{6x^2}$ and substituting into

$h_y = 0$ gives $6x = 64{,}000\left(\dfrac{6x^2}{80{,}000}\right)^2 \quad \Rightarrow \quad x^3 = \dfrac{80{,}000^2}{6 \cdot 64{,}000} = \dfrac{50{,}000}{3}$, so

$x = \sqrt[3]{\dfrac{50{,}000}{3}} = 10\sqrt[3]{\dfrac{50}{3}} \quad \Rightarrow \quad y = \dfrac{80}{\sqrt[3]{60}}$, and the only critical point of h is $\left(10\sqrt[3]{\dfrac{50}{3}}, \dfrac{80}{\sqrt[3]{60}}\right) \approx (25.54, 20.43)$ which is not in D. Next we check the boundary of D.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

On L_1: $y = 30$, $h(x, 30) = 180x + 80{,}000/x + 6400/3$, $30 \le x \le \frac{100}{3}$. Since $h'(x, 30) = 180 - 80{,}000/x^2 > 0$ for $30 \le x \le \frac{100}{3}$, $h(x, 30)$ is an increasing function with minimum $h(30, 30) = 10{,}200$ and maximum $h\left(\frac{100}{3}, 30\right) \approx 10{,}533$.

On L_2: $y = 1000/x$, $h(x, 1000/x) = 6000 + 64x + 80{,}000/x$, $30 \le x \le \frac{100}{3}$.

Since $h'(x, 1000/x) = 64 - 80{,}000/x^2 < 0$ for $30 \le x \le \frac{100}{3}$, $h(x, 1000/x)$ is a decreasing function with minimum $h\left(\frac{100}{3}, 30\right) \approx 10{,}533$ and maximum $h\left(30, \frac{100}{3}\right) \approx 10{,}587$.

On L_3: $x = 30$, $h(30, y) = 180y + 64{,}000/y + 8000/3$, $30 \le y \le \frac{100}{3}$. $h'(30, y) = 180 - 64{,}000/y^2 > 0$ for $30 \le y \le \frac{100}{3}$, so $h(30, y)$ is an increasing function of y with minimum $h(30, 30) = 10{,}200$ and maximum $h\left(30, \frac{100}{3}\right) \approx 10{,}587$.

Thus the absolute minimum of h is $h(30, 30) = 10{,}200$, and the dimensions of the building that minimize heat loss are walls 30 m in length and height $\frac{4000}{30^2} = \frac{40}{9} \approx 4.44$ m.

(c) From part (b), the only critical point of h, which gives a local (and absolute) minimum, is approximately $h(25.54, 20.43) \approx 9396$. So a building of volume 4000 m^2 with dimensions $x \approx 25.54$ m, $y \approx 20.43$ m, $z \approx \frac{4000}{(25.54)(20.43)} \approx 7.67$ m has the least amount of heat loss.

57. Let x, y, z be the dimensions of the rectangular box. Then the volume of the box is xyz and

$L = \sqrt{x^2 + y^2 + z^2} \quad \Rightarrow \quad L^2 = x^2 + y^2 + z^2 \quad \Rightarrow \quad z = \sqrt{L^2 - x^2 - y^2}$.

Substituting, we have volume $V(x, y) = xy\sqrt{L^2 - x^2 - y^2} \quad (x, y > 0)$.

$$V_x = xy \cdot \tfrac{1}{2}(L^2 - x^2 - y^2)^{-1/2}(-2x) + y\sqrt{L^2 - x^2 - y^2} = y\sqrt{L^2 - x^2 - y^2} - \frac{x^2 y}{\sqrt{L^2 - x^2 - y^2}},$$

$V_y = x\sqrt{L^2 - x^2 - y^2} - \dfrac{xy^2}{\sqrt{L^2 - x^2 - y^2}}$. $V_x = 0$ implies $y(L^2 - x^2 - y^2) = x^2 y \quad \Rightarrow \quad y(L^2 - 2x^2 - y^2) = 0 \quad \Rightarrow$ $2x^2 + y^2 = L^2$ (since $y > 0$), and $V_y = 0$ implies $x(L^2 - x^2 - y^2) = xy^2 \quad \Rightarrow \quad x(L^2 - x^2 - 2y^2) = 0 \quad \Rightarrow$ $x^2 + 2y^2 = L^2$ (since $x > 0$). Substituting $y^2 = L^2 - 2x^2$ into $x^2 + 2y^2 = L^2$ gives $x^2 + 2L^2 - 4x^2 = L^2 \quad \Rightarrow$ $3x^2 = L^2 \quad \Rightarrow \quad x = L/\sqrt{3}$ (since $x > 0$) and then $y = \sqrt{L^2 - 2\left(L/\sqrt{3}\right)^2} = L/\sqrt{3}$.

So the only critical point is $\left(L/\sqrt{3}, L/\sqrt{3}\right)$ which, from the geometrical nature of the problem, must give an absolute maximum. Thus the maximum volume is $V\left(L/\sqrt{3}, L/\sqrt{3}\right) = \left(L/\sqrt{3}\right)^2 \sqrt{L^2 - \left(L/\sqrt{3}\right)^2 - \left(L/\sqrt{3}\right)^2} = L^3/\left(3\sqrt{3}\right)$ cubic units.

58. $Y(N, P) = kNPe^{-N-P} \quad \Rightarrow \quad Y_N = kP\left[N(-e^{-N-P}) + e^{-N-P}(1)\right] = kP(1 - N)e^{-N-P}$,

$Y_P = kN\left[P(-e^{-N-P}) + e^{-N-P}(1)\right] = kN(1 - P)e^{-N-P}$. Here $N \ge 0$ and $P \ge 0$, but if either $N = 0$ or $P = 0$ then

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

the yield is zero. Assuming that $N > 0$ and $P > 0$, $Y_N = 0$ implies $N = 1$ and $Y_P = 0$ implies $P = 1$, so the only critical point in $\{(N, P) \mid N > 0, P > 0\}$ is $(1, 1)$ where $Y(1, 1) = ke^{-2}$.

$D(N, P) = Y_{NN}Y_{PP} - (Y_{NP})^2 = \left[kP(N-2)e^{-N-P}\right]\left[kN(P-2)e^{-N-P}\right] - \left[k(1-N)(1-P)e^{-N-P}\right]^2 \quad \Rightarrow$

$D(1, 1) = \left(-ke^{-2}\right)\left(-ke^{-2}\right) - (0)^2 = k^2e^{-4} > 0$ and $Y_{NN}(1, 1) = -ke^{-2} < 0$, so $Y(1, 1) = ke^{-2}$ is a local maximum. $Y(1, 1)$ is also the absolute maximum (we have only one critical point, and $Y \to 0$ as $N \to 0$ or $P \to 0$ and $Y \to 0$ as N or P grow large) so the best yield is achieved when both the nitrogen and phosphorus levels are 1 (measured in appropriate units).

59. (a) We are given that $p_1 + p_2 + p_3 = 1 \quad \Rightarrow \quad p_3 = 1 - p_1 - p_2$, so

$H = -p_1 \ln p_1 - p_2 \ln p_2 - p_3 \ln p_3 = -p_1 \ln p_1 - p_2 \ln p_2 - (1 - p_1 - p_2) \ln (1 - p_1 - p_2)$.

(b) Because p_i is a proportion we have $0 \le p_i \le 1$, but H is undefined unless $p_1 > 0$, $p_2 > 0$, and $1 - p_1 - p_2 > 0 \quad \Leftrightarrow \quad p_1 + p_2 < 1$. This last restriction forces $p_1 < 1$ and $p_2 < 1$, so the domain of H is $\{(p_1, p_2) \mid 0 < p_1 < 1,\ p_2 < 1 - p_1\}$. It is the interior of the triangle drawn in the figure.

(c)
$$\begin{aligned} H_{p_1} &= -\left[p_1 \cdot (1/p_1) + (\ln p_1) \cdot 1\right] - \left[(1 - p_1 - p_2) \cdot (-1)/(1 - p_1 - p_2) + \ln (1 - p_1 - p_2) \cdot (-1)\right] \\ &= -1 - \ln p_1 + 1 + \ln (1 - p_1 - p_2) = \ln (1 - p_1 - p_2) - \ln p_1 \end{aligned}$$

Similarly $H_{p_2} = \ln (1 - p_1 - p_2) - \ln p_2$. Then $H_{p_1} = 0$ implies

$\ln (1 - p_1 - p_2) = \ln p_1 \quad \Rightarrow \quad 1 - p_1 - p_2 = p_1 \quad \Rightarrow \quad p_2 = 1 - 2p_1$, and $H_{p_2} = 0$ implies

$\ln (1 - p_1 - p_2) = \ln p_2 \quad \Rightarrow \quad p_1 = 1 - 2p_2$. Substituting, we have $p_1 = 1 - 2(1 - 2p_1) \quad \Rightarrow$

$3p_1 = 1 \quad \Rightarrow \quad p_1 = \frac{1}{3}$, and then $p_2 = 1 - 2\left(\frac{1}{3}\right) = \frac{1}{3}$. Thus the only critical point is $\left(\frac{1}{3}, \frac{1}{3}\right)$.

$D(p_1, p_2) = H_{p_1p_1}H_{p_2p_2} - (H_{p_1p_2})^2 = \left(\dfrac{-1}{1 - p_1 - p_2} - \dfrac{1}{p_1}\right)\left(\dfrac{-1}{1 - p_1 - p_2} - \dfrac{1}{p_2}\right) - \left(\dfrac{-1}{1 - p_1 - p_2}\right)^2$, so

$D\left(\frac{1}{3}, \frac{1}{3}\right) = (-6)(-6) - (-3)^2 = 27 > 0$ and $H_{p_1p_1}\left(\frac{1}{3}, \frac{1}{3}\right) = -6 < 0$. Thus

$H\left(\frac{1}{3}, \frac{1}{3}\right) = -\frac{1}{3}\ln\frac{1}{3} - \frac{1}{3}\ln\frac{1}{3} - \frac{1}{3}\ln\frac{1}{3} = -\ln\frac{1}{3} = \ln 3$ is a local maximum. Here it is also the absolute maximum, so the maximum value of H is $\ln 3$, which occurs for $p_1 = p_2 = p_3 = \frac{1}{3}$ (all three species have equal proportion in the ecosystem).

60. Since $p + q + r = 1$ we can substitute $p = 1 - r - q$ into P giving

$P = P(q, r) = 2(1 - r - q)q + 2(1 - r - q)r + 2rq = 2q - 2q^2 + 2r - 2r^2 - 2rq$. Since p, q and r represent proportions and $p + q + r = 1$, we know $q \ge 0$, $r \ge 0$, and $q + r \le 1$. Thus, we want to find the absolute maximum of the continuous function $P(q, r)$ on the closed set D enclosed by the lines $q = 0$, $r = 0$, and $q + r = 1$. To find any critical points, we set the partial derivatives equal to zero: $P_q(q, r) = 2 - 4q - 2r = 0$ and $P_r(q, r) = 2 - 4r - 2q = 0$. The first equation gives

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$r = 1 - 2q$, and substituting into the second equation we have $2 - 4(1 - 2q) - 2q = 0 \quad \Rightarrow \quad q = \frac{1}{3}$. Then we have one critical point, $\left(\frac{1}{3}, \frac{1}{3}\right)$, where $P\left(\frac{1}{3}, \frac{1}{3}\right) = \frac{2}{3}$. Next we find the maximum values of P on the boundary of D which consists of three line segments. For the segment given by $r = 0$, $0 \le q \le 1$, $P(q, r) = P(q, 0) = 2q - 2q^2$, $0 \le q \le 1$. This represents a parabola with maximum value $P\left(\frac{1}{2}, 0\right) = \frac{1}{2}$. On the segment $q = 0$, $0 \le r \le 1$ we have $P(0, r) = 2r - 2r^2$, $0 \le r \le 1$. This represents a parabola with maximum value $P\left(0, \frac{1}{2}\right) = \frac{1}{2}$. Finally, on the segment $q + r = 1$, $0 \le q \le 1$, $P(q, r) = P(q, 1 - q) = 2q - 2q^2$, $0 \le q \le 1$ which has a maximum value of $P\left(\frac{1}{2}, \frac{1}{2}\right) = \frac{1}{2}$. Comparing these values with the value of P at the critical point, we see that the absolute maximum value of $P(q, r)$ on D is $\frac{2}{3}$.

61. Note that here the variables are m and b, and $f(m, b) = \sum_{i=1}^{n} [y_i - (mx_i + b)]^2$. Then $f_m = \sum_{i=1}^{n} -2x_i[y_i - (mx_i + b)] = 0$ implies $\sum_{i=1}^{n} \left(x_i y_i - mx_i^2 - bx_i\right) = 0$ or $\sum_{i=1}^{n} x_i y_i = m \sum_{i=1}^{n} x_i^2 + b \sum_{i=1}^{n} x_i$ and $f_b = \sum_{i=1}^{n} -2[y_i - (mx_i + b)] = 0$ implies $\sum_{i=1}^{n} y_i = m \sum_{i=1}^{n} x_i + \sum_{i=1}^{n} b = m\left(\sum_{i=1}^{n} x_i\right) + nb$. Thus we have the two desired equations.

Now $f_{mm} = \sum_{i=1}^{n} 2x_i^2$, $f_{bb} = \sum_{i=1}^{n} 2 = 2n$ and $f_{mb} = \sum_{i=1}^{n} 2x_i$. And $f_{mm}(m, b) > 0$ always and $D(m, b) = 4n\left(\sum_{i=1}^{n} x_i^2\right) - 4\left(\sum_{i=1}^{n} x_i\right)^2 = 4\left[n\left(\sum_{i=1}^{n} x_i^2\right) - \left(\sum_{i=1}^{n} x_i\right)^2\right] > 0$ always so the solutions of these two equations do indeed minimize $\sum_{i=1}^{n} d_i^2$.

62. Any such plane must cut out a tetrahedron in the first octant. We need to minimize the volume of the tetrahedron that passes through the point $(1, 2, 3)$. Writing the equation of the plane as $\dfrac{x}{a} + \dfrac{y}{b} + \dfrac{z}{c} = 1$, the volume of the tetrahedron is given by $V = \dfrac{abc}{6}$. But $(1, 2, 3)$ must lie on the plane, so we need $\dfrac{1}{a} + \dfrac{2}{b} + \dfrac{3}{c} = 1$ $(\star)$ and thus can think of c as a function of a and b. Then $V_a = \dfrac{b}{6}\left(c + a\dfrac{\partial c}{\partial a}\right)$ and $V_b = \dfrac{a}{6}\left(c + b\dfrac{\partial c}{\partial b}\right)$. Differentiating $(\star)$ with respect to a we get $-a^{-2} - 3c^{-2}\dfrac{\partial c}{\partial a} = 0 \quad \Rightarrow$ $\dfrac{\partial c}{\partial a} = \dfrac{-c^2}{3a^2}$, and differentiating $(\star)$ with respect to b gives $-2b^{-2} - 3c^{-2}\dfrac{\partial c}{\partial b} = 0 \quad \Rightarrow \quad \dfrac{\partial c}{\partial b} = \dfrac{-2c^2}{3b^2}$. Then $V_a = \dfrac{b}{6}\left(c + a\dfrac{-c^2}{3a^2}\right) = 0 \quad \Rightarrow \quad c = 3a$, and $V_b = \dfrac{a}{6}\left(c + b\dfrac{-2c^2}{3b^2}\right) = 0 \quad \Rightarrow \quad c = \frac{3}{2}b$. Thus $3a = \frac{3}{2}b$ or $b = 2a$. Putting these into $(\star)$ gives $\frac{3}{a} = 1$ or $a = 3$ and then $b = 6$, $c = 9$. Thus the equation of the required plane is $\dfrac{x}{3} + \dfrac{y}{6} + \dfrac{z}{9} = 1$ or $6x + 3y + 2z = 18$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

DISCOVERY PROJECT Quadratic Approximations and Critical Points

1. $Q(x,y) = f(a,b) + f_x(a,b)(x-a) + f_y(a,b)(y-b) + \frac{1}{2}f_{xx}(a,b)(x-a)^2$
$+ f_{xy}(a,b)(x-a)(y-b) + \frac{1}{2}f_{yy}(a,b)(y-b)^2,$

so

$$Q_x(x,y) = f_x(a,b) + \tfrac{1}{2}f_{xx}(a,b)(2)(x-a) + f_{xy}(a,b)(y-b) = f_x(a,b) + f_{xx}(a,b)(x-a) + f_{xy}(a,b)(y-b)$$

At (a,b) we have $Q_x(a,b) = f_x(a,b) + f_{xx}(a,b)\,(a-a) + f_{xy}(a,b)(b-b) = f_x(a,b)$.

Similarly, $Q_y(x,y) = f_y(a,b) + f_{xy}(a,b)(x-a) + f_{yy}(a,b)(y-b) \quad \Rightarrow$

$Q_y(a,b) = f_y(a,b) + f_{xy}(a,b)(a-a) + f_{yy}(a,b)(b-b) = f_y(a,b)$.

For the second-order partial derivatives we have

$$Q_{xx}(x,y) = \frac{\partial}{\partial x}\left[f_x(a,b) + f_{xx}(a,b)(x-a) + f_{xy}(a,b)(y-b)\right] = f_{xx}(a,b) \quad \Rightarrow \quad Q_{xx}(a,b) = f_{xx}(a,b)$$

$$Q_{xy}(x,y) = \frac{\partial}{\partial y}\left[f_x(a,b) + f_{xx}(a,b)(x-a) + f_{xy}(a,b)(y-b)\right] = f_{xy}(a,b) \quad \Rightarrow \quad Q_{xy}(a,b) = f_{xy}(a,b)$$

$$Q_{yy}(x,y) = \frac{\partial}{\partial y}\left[f_y(a,b) + f_{xy}(a,b)(x-a) + f_{yy}(a,b)(y-b)\right] = f_{yy}(a,b) \quad \Rightarrow \quad Q_{yy}(a,b) = f_{yy}(a,b)$$

2. (a) First we find the partial derivatives and values that will be needed:

$$\begin{aligned}
f(x,y) &= e^{-x^2-y^2} & f(0,0) &= 1\\
f_x(x,y) &= -2xe^{-x^2-y^2} & f_x(0,0) &= 0\\
f_y(x,y) &= -2ye^{-x^2-y^2} & f_y(0,0) &= 0\\
f_{xx}(x,y) &= (4x^2-2)e^{-x^2-y^2} & f_{xx}(0,0) &= -2\\
f_{xy}(x,y) &= 4xye^{-x^2-y^2} & f_{xy}(0,0) &= 0\\
f_{yy}(x,y) &= (4y^2-2)e^{-x^2-y^2} & f_{yy}(0,0) &= -2
\end{aligned}$$

Then the first-degree Taylor polynomial of f at $(0,0)$ is

$$L(x,y) = f(0,0) + f_x(0,0)(x-0) + f_y(0,0)(y-0) = 1 + (0)(x-0) + (0)(y-0) = 1$$

The second-degree Taylor polynomial is given by

$$\begin{aligned}
Q(x,y) &= f(0,0) + f_x(0,0)(x-0) + f_y(0,0)(y-0) + \tfrac{1}{2}f_{xx}(0,0)(x-0)^2\\
&\qquad + f_{xy}(0,0)(x-0)(y-0) + \tfrac{1}{2}f_{yy}(0,0)(y-0)^2\\
&= 1 - x^2 - y^2
\end{aligned}$$

(b)

As we see from the graph, L approximates f well only for points (x,y) extremely close to the origin. Q is a much better approximation; the shape of its graph looks similar to that of the graph of f near the origin, and the values of Q appear to be good estimates for the values of f within a significant radius of the origin.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a) First we find the partial derivatives and values that will be needed:

$$\begin{array}{llll} f(x,y) = xe^y & f(1,0) = 1 & f_{xx}(x,y) = 0 & f_{xx}(1,0) = 0 \\ f_x(x,y) = e^y & f_x(1,0) = 1 & f_{xy}(x,y) = e^y & f_{xy}(1,0) = 1 \\ f_y(x,y) = xe^y & f_y(1,0) = 1 & f_{yy}(x,y) = xe^y & f_{yy}(1,0) = 1 \end{array}$$

Then the first-degree Taylor polynomial of f at $(1,0)$ is

$$L(x,y) = f(1,0) + f_x(1,0)(x-1) + f_y(1,0)(y-0) = 1 + (1)(x-1) + (1)(y-0) = x + y$$

The second-degree Taylor polynomial is given by

$$\begin{aligned} Q(x,y) &= f(1,0) + f_x(1,0)(x-1) + f_y(1,0)(y-0) + \tfrac{1}{2} f_{xx}(1,0)(x-1)^2 \\ &\qquad + f_{xy}(1,0)(x-1)(y-0) + \tfrac{1}{2} f_{yy}(1,0)(y-0)^2 \\ &= \tfrac{1}{2}y^2 + x + xy \end{aligned}$$

(b) $L(0.9, 0.1) = 0.9 + 0.1 = 1.0$

$Q(0.9, 0.1) = \frac{1}{2}(0.1)^2 + 0.9 + (0.9)(0.1) = 0.995$

$f(0.9, 0.1) = 0.9e^{0.1} \approx 0.9947$

(c)

As we see from the graph, L and Q both approximate f reasonably well near the point $(1,0)$. As we venture farther from the point, the graph of Q follows the shape of the graph of f more closely than L.

4. (a) $f(x,y) = ax^2 + bxy + cy^2 = a\left[x^2 + \dfrac{b}{a}xy + \dfrac{c}{a}y^2\right] = a\left[x^2 + \dfrac{b}{a}xy + \left(\dfrac{b}{2a}y\right)^2 - \left(\dfrac{b}{2a}y\right)^2 + \dfrac{c}{a}y^2\right]$

$$= a\left[\left(x + \frac{b}{2a}y\right)^2 - \frac{b^2}{4a^2}y^2 + \frac{c}{a}y^2\right] = a\left[\left(x + \frac{b}{2a}y\right)^2 + \left(\frac{4ac - b^2}{4a^2}\right)y^2\right]$$

(b) For $D = 4ac - b^2$, from part (a) we have $f(x,y) = a\left[\left(x + \dfrac{b}{2a}y\right)^2 + \left(\dfrac{D}{4a^2}\right)y^2\right]$. If $D > 0$,

$\left(\dfrac{D}{4a^2}\right)y^2 \geq 0$ and $\left(x + \dfrac{b}{2a}y\right)^2 \geq 0$, so $\left[\left(x + \dfrac{b}{2a}y\right)^2 + \left(\dfrac{D}{4a^2}\right)y^2\right] \geq 0$. Here $a > 0$, thus

$f(x,y) = a\left[\left(x + \dfrac{b}{2a}y\right)^2 + \left(\dfrac{D}{4a^2}\right)y^2\right] \geq 0$. We know $f(0,0) = 0$, so $f(0,0) \leq f(x,y)$ for all (x,y), and by definition f has a local minimum at $(0,0)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) As in part (b), $\left[\left(x+\dfrac{b}{2a}\,y\right)^2+\left(\dfrac{D}{4a^2}\right)y^2\right]\geq 0$, and since $a<0$ we have

$f(x,y)=a\left[\left(x+\dfrac{b}{2a}\,y\right)^2+\left(\dfrac{D}{4a^2}\right)y^2\right]\leq 0$. Since $f(0,0)=0$, we must have $f(0,0)\geq f(x,y)$ for all (x,y), so by definition f has a local maximum at $(0,0)$.

(d) $f(x,y)=ax^2+bxy+cy^2$, so $f_x(x,y)=2ax+by \quad\Rightarrow\quad f_x(0,0)=0$ and $f_y(x,y)=bx+2cy \quad\Rightarrow\quad f_y(0,0)=0$. Since $f(0,0)=0$ and f and its partial derivatives are continuous, we know from Equation 14.4.2 that the tangent plane to the graph of f at $(0,0)$ is the plane $z=0$. Then f has a saddle point at $(0,0)$ if the graph of f crosses the tangent plane at $(0,0)$, or equivalently, if some paths to the origin have positive function values while other paths have negative function values. Suppose we approach the origin along the x-axis; then we have $y=0 \quad\Rightarrow\quad f(x,0)=ax^2$ which has the same sign as a. We must now find at least one path to the origin where $f(x,y)$ gives values with sign opposite that of a. Since $f(x,y)=a\left[\left(x+\dfrac{b}{2a}\,y\right)^2+\left(\dfrac{D}{4a^2}\right)y^2\right]$, if we approach the origin along the line $x=-\dfrac{b}{2a}\,y$, we have $f\left(-\dfrac{b}{2a}\,y,y\right)=a\left[\left(-\dfrac{b}{2a}\,y+\dfrac{b}{2a}\,y\right)^2+\left(\dfrac{D}{4a^2}\right)y^2\right]=\dfrac{D}{4a}\,y^2$. Since $D<0$, these values have signs opposite that of a. Thus, f has a saddle point at $(0,0)$.

5. (a) Since the partial derivatives of f exist at $(0,0)$ and $(0,0)$ is a critical point, we know $f_x(0,0)=0$ and $f_y(0,0)=0$. Then the second-degree Taylor polynomial of f at $(0,0)$ can be expressed as

$$\begin{aligned}Q(x,y)&=f(0,0)+f_x(0,0)(x-0)+f_y(0,0)(y-0)+\tfrac{1}{2}f_{xx}(0,0)(x-0)^2\\&\qquad+f_{xy}(0,0)(x-0)(y-0)+\tfrac{1}{2}f_{yy}(0,0)(y-0)^2\\&=\tfrac{1}{2}f_{xx}(0,0)x^2+f_{xy}(0,0)xy+\tfrac{1}{2}f_{yy}(0,0)y^2\end{aligned}$$

(b) $Q(x,y)=\frac{1}{2}\,f_{xx}(0,0)x^2+f_{xy}(0,0)xy+\frac{1}{2}\,f_{yy}(0,0)y^2$ fits the form of the polynomial function in Problem 4 with $a=\frac{1}{2}\,f_{xx}(0,0)$, $b=f_{xy}(0,0)$, and $c=\frac{1}{2}\,f_{yy}(0,0)$. Then we know Q is a paraboloid, and that Q has a local maximum, local minimum, or saddle point at $(0,0)$. Here, $D=4ac-b^2=4\left(\frac{1}{2}\right)f_{xx}(0,0)\left(\frac{1}{2}\right)f_{yy}(0,0)-[f_{xy}(0,0)]^2=f_{xx}(0,0)f_{yy}(0,0)-[f_{xy}(0,0)]^2$, and if $D>0$ with $a=\frac{1}{2}\,f_{xx}(0,0)>0 \quad\Rightarrow\quad f_{xx}(0,0)>0$, we know from Problem 4 that Q has a local minimum at $(0,0)$. Similarly, if $D>0$ and $a<0 \quad\Rightarrow\quad f_{xx}(0,0)<0$, Q has a local maximum at $(0,0)$, and if $D<0$, Q has a saddle point at $(0,0)$.

(c) Since $f(x,y)\approx Q(x,y)$ near $(0,0)$, part (b) suggests that for $D=f_{xx}(0,0)f_{yy}(0,0)-[f_{xy}(0,0)]^2$, if $D>0$ and $f_{xx}(0,0)>0$, f has a local minimum at $(0,0)$. If $D>0$ and $f_{xx}(0,0)<0$, f has a local maximum at $(0,0)$, and if $D<0$, f has a saddle point at $(0,0)$. Together with the conditions given in part (a), this is precisely the Second Derivatives Test from Section 14.7.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14.8 Lagrange Multipliers

1. At the extreme values of f, the level curves of f just touch the curve $g(x, y) = 8$ with a common tangent line. (See Figure 1 and the accompanying discussion.) We can observe several such occurrences on the contour map, but the level curve $f(x, y) = c$ with the largest value of c which still intersects the curve $g(x, y) = 8$ is approximately $c = 59$, and the smallest value of c corresponding to a level curve which intersects $g(x, y) = 8$ appears to be $c = 30$. Thus we estimate the maximum value of f subject to the constraint $g(x, y) = 8$ to be about 59 and the minimum to be 30.

2. (a) The values $c = \pm 1$ and $c = 1.25$ seem to give curves which are tangent to the circle. These values represent possible extreme values of the function $x^2 + y$ subject to the constraint $x^2 + y^2 = 1$.

(b) $\nabla f = \langle 2x, 1\rangle$, $\lambda \nabla g = \langle 2\lambda x, 2\lambda y\rangle$. So $2x = 2\lambda x \quad \Rightarrow \quad$ either $\lambda = 1$ or $x = 0$. If $\lambda = 1$, then $y = \frac{1}{2}$ and so $x = \pm\frac{\sqrt{3}}{2}$ (from the constraint). If $x = 0$, then $y = \pm 1$. Therefore f has possible extreme values at the points $(0, \pm 1)$ and $\left(\pm\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$. We calculate $f\left(\pm\frac{\sqrt{3}}{2}, \frac{1}{2}\right) = \frac{5}{4}$ (the maximum value), $f(0, 1) = 1$, and $f(0, -1) = -1$ (the minimum value). These are our answers from part (a).

3. We want to find the extreme values of $f(x, y) = x^2 - y^2$ subject to the constraint $g(x, y) = x^2 + y^2 = 1$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2x, -2y\rangle = \lambda \langle 2x, 2y\rangle$, so we solve the equations $2x = 2\lambda x$, $-2y = 2\lambda y$, and $x^2 + y^2 = 1$. From the first equation we have $2x(\lambda - 1) = 0 \quad \Rightarrow \quad x = 0$ or $\lambda = 1$. If $x = 0$ then substitution into the constraint gives $y^2 = 1 \quad \Rightarrow \quad y = \pm 1$. If $\lambda = 1$ then substitution into the second equation gives $-2y = 2y \quad \Rightarrow \quad y = 0$, and from the constraint we must have $x = \pm 1$. Thus the possible points for the extreme values of f are $(0, \pm 1)$ and $(\pm 1, 0)$. Evaluating f at these points, we see that the maximum value of f is $f(\pm 1, 0) = 1$ and the minimum is $f(0, \pm 1) = -1$.

4. $f(x, y) = x^2 y$, $g(x, y) = x^2 + y^4 = 5$, and $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \left\langle 2xy, x^2\right\rangle = \left\langle 2\lambda x, 4\lambda y^3\right\rangle$, so we get the three equations $2xy = 2\lambda x$, $x^2 = 4\lambda y^3$, and $x^2 + y^4 = 5$. $2xy = 2\lambda x \quad \Rightarrow \quad x = 0$ or $y = \lambda$. If $x = 0$, the second equation implies $y = 0$ or $\lambda = 0$. The point $(0, 0)$ does not satisfy the constraint, so $x = \lambda = 0$ and the constraint gives a possible extreme value at the point $\left(0, \sqrt[4]{5}\right)$. Next, suppose $y = \lambda$. Substituting into the second equation $\quad \Rightarrow \quad x^2 = 4\lambda^4$ and substituting into the third equation gives $4\lambda^4 + \lambda^4 = 5 \quad \Rightarrow \quad \lambda = \pm 1$. From the second equation with $y = \lambda = 1$, we get $x = \pm 2$. From the second equation with $y = \lambda = -1$, we get $x = \pm 2$. So f also has possible extreme values at $(\pm 2, 1)$ and $(\pm 2, -1)$. Evaluating f at these 5 points, we see $f(\pm 2, 1) = 4$ is the maximum value and $f(\pm 2, -1) = -4$ is the minimum value.

5. $f(x, y) = xy$, $g(x, y) = 4x^2 + y^2 = 8$, and $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle y, x\rangle = \langle 8\lambda x, 2\lambda y\rangle$, so $y = 8\lambda x$, $x = 2\lambda y$, and $4x^2 + y^2 = 8$. First note that if $x = 0$ then $y = 0$ by the first equation, and if $y = 0$ then $x = 0$ by the second equation. But this contradicts the third equation, so $x \neq 0$ and $y \neq 0$. Then from the first two equations we have $\dfrac{y}{8x} = \lambda = \dfrac{x}{2y} \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$2y^2 = 8x^2 \quad \Rightarrow \quad y^2 = 4x^2$, and substitution into the third equation gives $4x^2 + 4x^2 = 8 \quad \Rightarrow \quad x = \pm 1$. If $x = \pm 1$ then $y^2 = 4 \quad \Rightarrow \quad y = \pm 2$, so f has possible extreme values at $(1, \pm 2)$ and $(-1, \pm 2)$. Evaluating f at these points, we see that the maximum value is $f(1, 2) = f(-1, -2) = 2$ and the minimum is $f(1, -2) = f(-1, 2) = -2$.

6. $f(x, y) = xe^y$, $g(x, y) = x^2 + y^2 = 2$, and $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle e^y, xe^y \rangle = \langle 2\lambda x, 2\lambda y \rangle$, so $e^y = 2\lambda x$, $xe^y = 2\lambda y$, and $x^2 + y^2 = 2$. First note that from the first equation $x \neq 0$. If $y = 0$, the second equation implies $x = 0$, so $y \neq 0$. Then from the first two equations we have $\dfrac{e^y}{2x} = \lambda = \dfrac{xe^y}{2y} \quad \Rightarrow \quad 2ye^y = 2x^2e^y \quad \Rightarrow \quad y = x^2$, and substituting into the third equation gives $x^2 + (x^2)^2 = 2 \quad \Rightarrow \quad x^4 + x^2 - 2 = 0 \quad \Rightarrow \quad (x^2 + 2)(x^2 - 1) = 0 \quad \Rightarrow \quad x = \pm 1$. From $y = x^2$ we have $y = 1$, so f has possible extreme values at $(\pm 1, 1)$. Evaluating f at these points, we see that the maximum value is $f(1, 1) = e$ and the minimum is $f(-1, 1) = -e$.

7. $f(x, y) = 2x^2 + 6y^2$, $g(x, y) = x^4 + 3y^4 = 1$, and $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 4x, 12y \rangle = \langle 4\lambda x^3, 12\lambda y^3 \rangle$, so we get the three equations $4x = 4\lambda x^3$, $12y = 12\lambda y^3$, and $x^4 + 3y^4 = 1$. The first equation implies that $x = 0$ or $x^2 = \dfrac{1}{\lambda}$. The second equation implies that $y = 0$ or $y^2 = \dfrac{1}{\lambda}$. Note that x and y cannot both be zero as this contradicts the third equation. If $x = 0$, the third equation implies $y = \pm \dfrac{1}{\sqrt[4]{3}}$. If $y = 0$, the third equation implies that $x = \pm 1$. Thus, f has possible extreme values at $\left(0, \pm \dfrac{1}{\sqrt[4]{3}}\right)$ and $(\pm 1, 0)$. Next, suppose $x^2 = y^2 = \dfrac{1}{\lambda}$. Then the third equation gives $\left(\dfrac{1}{\lambda}\right)^2 + 3\left(\dfrac{1}{\lambda}\right)^2 = 1 \quad \Rightarrow \quad \lambda = \pm 2$. $\lambda = -2$ results in a nonreal solution, so consider $\lambda = 2 \quad \Rightarrow \quad x = y = \pm \dfrac{1}{\sqrt{2}}$. Therefore, f also has possible extreme values at $\left(\pm \dfrac{1}{\sqrt{2}}, \pm \dfrac{1}{\sqrt{2}}\right)$ (all 4 combinations). Substituting all 8 points into f, we find the maximum value is $f\left(\pm \dfrac{1}{\sqrt{2}}, \pm \dfrac{1}{\sqrt{2}}\right) = 4$ and the minimum value is $f(\pm 1, 0) = 2$.

8. $f(x, y) = xye^{-x^2-y^2}$, $g(x, y) = 2x - y = 0$, and $\nabla f = \lambda \nabla g \quad \Rightarrow$ $\left\langle ye^{-x^2-y^2} - 2x^2ye^{-x^2-y^2}, xe^{-x^2-y^2} - 2xy^2e^{-x^2-y^2} \right\rangle = \langle 2\lambda, -\lambda \rangle$, so we get the three equations $ye^{-x^2-y^2} - 2x^2ye^{-x^2-y^2} = 2\lambda$, $xe^{-x^2-y^2} - 2xy^2e^{-x^2-y^2} = -\lambda$, and $2x - y = 0$. Multiplying the second equation by 2 and adding it to the first gives $2xe^{-x^2-y^2} - 4xy^2e^{-x^2-y^2} + ye^{-x^2-y^2} - 2x^2ye^{-x^2-y^2} = -2\lambda + 2\lambda = 0 \quad \Rightarrow$ $2x - 4xy^2 + y - 2x^2y = 0$ (as $e^{-x^2-y^2} \neq 0$). From the third equation, $2x = y$, and substituting into the new equation, we have $2x - 4x(2x)^2 + 2x - 2x^2(2x) = 0 \quad \Rightarrow \quad 4x - 20x^3 = 0 \quad \Rightarrow \quad 4x(1 - 5x^2) = 0 \quad \Rightarrow \quad x = 0$ or $x = \pm \dfrac{1}{\sqrt{5}}$, so f has possible extreme values at $(0, 0)$, $\left(\dfrac{1}{\sqrt{5}}, \dfrac{2}{\sqrt{5}}\right)$, and $\left(-\dfrac{1}{\sqrt{5}}, -\dfrac{2}{\sqrt{5}}\right)$. Substituting these into f, we see that the minimum value is $f(0, 0) = 0$ and the maximum value is $f\left(\dfrac{1}{\sqrt{5}}, \dfrac{2}{\sqrt{5}}\right) = f\left(-\dfrac{1}{\sqrt{5}}, -\dfrac{2}{\sqrt{5}}\right) = \dfrac{2}{5e}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $f(x,y,z) = 2x + 2y + z,\ g(x,y,z) = x^2 + y^2 + z^2 = 9$, and $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2,2,1\rangle = \langle 2\lambda x, 2\lambda y, 2\lambda z\rangle$, so $2\lambda x = 2$, $2\lambda y = 2$, $2\lambda z = 1$, and $x^2 + y^2 + z^2 = 9$. The first three equations imply $x = \dfrac{1}{\lambda}$, $y = \dfrac{1}{\lambda}$, and $z = \dfrac{1}{2\lambda}$. But substitution into the fourth equation gives $\left(\dfrac{1}{\lambda}\right)^2 + \left(\dfrac{1}{\lambda}\right)^2 + \left(\dfrac{1}{2\lambda}\right)^2 = 9 \quad \Rightarrow \quad \dfrac{9}{4\lambda^2} = 9 \quad \Rightarrow \quad \lambda = \pm\frac{1}{2}$, so f has possible extreme values at the points $(2,2,1)$ and $(-2,-2,-1)$. The maximum value of f on $x^2 + y^2 + z^2 = 9$ is $f(2,2,1) = 9$, and the minimum is $f(-2,-2,-1) = -9$.

10. $f(x,y,z) = e^{xyz},\ g(x,y,z) = 2x^2 + y^2 + z^2 = 24$, and $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle yze^{xyz}, xze^{xyz}, xye^{xyz}\rangle = \langle 4\lambda x, 2\lambda y, 2\lambda z\rangle$. Then $yze^{xyz} = 4\lambda x$, $xze^{xyz} = 2\lambda y$, $xye^{xyz} = 2\lambda z$, and $2x^2 + y^2 + z^2 = 24$. If any of x, y, z, or λ is zero, then the first three equations imply that two of the variables x, y, z must be zero. If $x = y = z = 0$ it contradicts the fourth equation, so exactly two are zero, and from the fourth equation the possibilities are $\left(\pm 2\sqrt{3}, 0, 0\right)$, $\left(0, \pm 2\sqrt{6}, 0\right)$, $\left(0, 0, \pm 2\sqrt{6}\right)$, all with an f-value of $e^0 = 1$. If none of x, y, z, λ is zero then from the first three equations we have $\dfrac{4\lambda x}{yz} = e^{xyz} = \dfrac{2\lambda y}{xz} = \dfrac{2\lambda z}{xy} \quad \Rightarrow \quad \dfrac{2x}{yz} = \dfrac{y}{xz} = \dfrac{z}{xy}$. This gives $2x^2 z = y^2 z \quad \Rightarrow \quad 2x^2 = y^2$ and $xy^2 = xz^2 \quad \Rightarrow$ $y^2 = z^2$. Substituting into the fourth equation, we have $y^2 + y^2 + y^2 = 24 \quad \Rightarrow \quad y^2 = 8 \quad \Rightarrow \quad y = \pm 2\sqrt{2}$, so $x^2 = 4 \quad \Rightarrow \quad x = \pm 2$ and $z^2 = y^2 \quad \Rightarrow \quad z = \pm 2\sqrt{2}$, giving possible points $\left(\pm 2, \pm 2\sqrt{2}, \pm 2\sqrt{2}\right)$ (all combinations). The value of f is e^{16} when all coordinates are positive or exactly two are negative, and the value is e^{-16} when all are negative or exactly one of the coordinates is negative. Thus the maximum of f subject to the constraint is e^{16} and the minimum is e^{-16}.

11. $f(x,y,z) = xy^2 z,\ g(x,y,z) = x^2 + y^2 + z^2 = 4$, and $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \left\langle y^2 z, 2xyz, xy^2\right\rangle = \lambda \langle 2x, 2y, 2z\rangle$. Then $y^2 z = 2\lambda x$, $2xyz = 2\lambda y$, $xy^2 = 2\lambda z$, and $x^2 + y^2 + z^2 = 4$.

Case 1: If $\lambda = 0$, then the first equation implies that $y = 0$ or $z = 0$. If $y = 0$, then any values of x and z satisfy the first three equations, so from the fourth equation all points $(x, 0, z)$ such that $x^2 + z^2 = 4$ are possible points. If $z = 0$ then from the third equation $x = 0$ or $y = 0$, and from the fourth equation, the possible points are $(0, \pm 2, 0)$, $(\pm 2, 0, 0)$. The f-value in all these cases is 0.

Case 2: If $\lambda \neq 0$ but any one of x, y, z is zero, the first three equations imply that all three coordinates must be zero, contradicting the fourth equation. Thus if $\lambda \neq 0$, none of x, y, z is zero and from the first three equations we have $\dfrac{y^2 z}{2x} = \lambda = xz = \dfrac{xy^2}{2z}$. This gives $y^2 z = 2x^2 z \quad \Rightarrow \quad y^2 = 2x^2$ and $2y^2 z^2 = 2x^2 y^2 \quad \Rightarrow \quad z^2 = x^2$. Substituting into the fourth equation, we have $x^2 + 2x^2 + x^2 = 4 \quad \Rightarrow \quad x^2 = 1 \quad \Rightarrow \quad x = \pm 1$, so $y = \pm\sqrt{2}$ and $z = \pm 1$, giving possible points $\left(\pm 1, \pm\sqrt{2}, \pm 1\right)$ (all combinations). The value of f is 2 when x and z are the same sign and -2 when they are opposite. Thus the maximum of f subject to the constraint is $f(1, \pm\sqrt{2}, 1) = f(-1, \pm\sqrt{2}, -1) = 2$ and the minimum is $f(1, \pm\sqrt{2}, -1) = f(-1, \pm\sqrt{2}, 1) = -2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $f(x,y,z) = x^2 + y^2 + z^2$, $g(x,y,z) = x^2 + y^2 + z^2 + xy = 12$, and $\nabla f = \lambda \nabla g \quad \Rightarrow$ $\langle 2x, 2y, 2z\rangle = \langle \lambda(2x+y), \lambda(2y+x), \lambda 2z\rangle$, so $(\mathbf{1})$ $2x = \lambda(2x+y)$, $(\mathbf{2})$ $2y = \lambda(2y+x)$, $(\mathbf{3})$ $2z = \lambda 2z$, and $(\mathbf{4})$ $x^2 + y^2 + z^2 + xy = 12$. First note that $\lambda = 0 \quad \Rightarrow \quad x = y = z = 0$, which contradicts $(\mathbf{4})$, so assume $\lambda \neq 0$. Then $(\mathbf{3})$ implies $\lambda = 1$ or $z = 0$. If $\lambda = 1$, then $(\mathbf{1})$ and $(\mathbf{2})$ imply $x = y = 0 \quad \Rightarrow \quad 0^2 + 0^2 + z^2 + 0 = 12 \quad \Rightarrow \quad z = \pm\sqrt{12}$. If $z = 0$, $y(\mathbf{1}) - x(\mathbf{2}) \quad \Rightarrow \quad 0 = \lambda y^2 - \lambda x^2 \quad \Rightarrow \quad x = y$ or $x = -y$. Substituting $x = y$ into $(\mathbf{4}) \quad \Rightarrow \quad y = x = \pm 2$. Substituting $x = -y$ into $(\mathbf{4}) \quad \Rightarrow \quad y = \pm\sqrt{12}$. Thus, f has possible extreme values at $\left(0, 0, \pm\sqrt{12}\right)$, $(2,2,0)$, $(-2,-2,0)$, $\left(\sqrt{12}, -\sqrt{12}, 0\right)$, and $\left(-\sqrt{12}, \sqrt{12}, 0\right)$. Evaluating f at these points, we see that the maximum value is $f\left(\sqrt{12}, -\sqrt{12}, 0\right) = f\left(-\sqrt{12}, \sqrt{12}, 0\right) = 24$ and the minimum is $f(2,2,0) = f(-2,-2,0) = 8$.

13. $f(x,y,z) = x^2 + y^2 + z^2,\ g(x,y,z) = x^4 + y^4 + z^4 = 1 \quad \Rightarrow \quad \nabla f = \langle 2x, 2y, 2z\rangle$, $\lambda \nabla g = \left\langle 4\lambda x^3, 4\lambda y^3, 4\lambda z^3\right\rangle$.

Case 1: If $x \neq 0$, $y \neq 0$, and $z \neq 0$, then $\nabla f = \lambda \nabla g$ implies $\lambda = 1/(2x^2) = 1/(2y^2) = 1/(2z^2)$ or $x^2 = y^2 = z^2$ and $3x^4 = 1$ or $x = \pm\frac{1}{\sqrt[4]{3}}$ giving the points $\left(\pm\frac{1}{\sqrt[4]{3}}, \frac{1}{\sqrt[4]{3}}, \frac{1}{\sqrt[4]{3}}\right)$, $\left(\pm\frac{1}{\sqrt[4]{3}}, -\frac{1}{\sqrt[4]{3}}, \frac{1}{\sqrt[4]{3}}\right)$, $\left(\pm\frac{1}{\sqrt[4]{3}}, \frac{1}{\sqrt[4]{3}}, -\frac{1}{\sqrt[4]{3}}\right)$, $\left(\pm\frac{1}{\sqrt[4]{3}}, -\frac{1}{\sqrt[4]{3}}, -\frac{1}{\sqrt[4]{3}}\right)$ all with an f-value of $\sqrt{3}$.

Case 2: If one of the variables equals zero and the other two are not zero, then the squares of the two nonzero coordinates are equal with common value $\frac{1}{\sqrt{2}}$ and corresponding f-value of $\sqrt{2}$.

Case 3: If exactly two of the variables are zero, then the third variable has value ± 1 with the corresponding f-value of 1.

Thus on $x^4 + y^4 + z^4 = 1$, the maximum value of f is $\sqrt{3}$ and the minimum value is 1.

14. $f(x,y,z) = x^4 + y^4 + z^4,\ g(x,y,z) = x^2 + y^2 + z^2 = 1 \quad \Rightarrow \quad \nabla f = \left\langle 4x^3, 4y^3, 4z^3\right\rangle$, $\lambda \nabla g = \langle 2\lambda x, 2\lambda y, 2\lambda z\rangle$.

Case 1: If $x \neq 0$, $y \neq 0$, and $z \neq 0$ then $\nabla f = \lambda \nabla g$ implies $\lambda = 2x^2 = 2y^2 = 2z^2$ or $x^2 = y^2 = z^2 = \frac{1}{3}$ giving 8 points each with an f-value of $\frac{1}{3}$.

Case 2: If one of the variables is 0 and the other two are not, then the squares of the two nonzero coordinates are equal with common value $\frac{1}{2}$ and the corresponding f-value is $\frac{1}{2}$.

Case 3: If exactly two of the variables are 0, then the third variable has value ± 1 with corresponding f-value of 1.

Thus on $x^2 + y^2 + z^2 = 1$, the maximum value of f is 1 and the minimum value is $\frac{1}{3}$.

15. $f(x,y,z,t) = x + y + z + t,\ g(x,y,z,t) = x^2 + y^2 + z^2 + t^2 = 1 \quad \Rightarrow \quad \langle 1,1,1,1\rangle = \langle 2\lambda x, 2\lambda y, 2\lambda z, 2\lambda t\rangle$, so $\lambda = 1/(2x) = 1/(2y) = 1/(2z) = 1/(2t)$ and $x = y = z = t$. But $x^2 + y^2 + z^2 + t^2 = 1$, so the possible points are $\left(\pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2}\right)$. Thus the maximum value of f is $f\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right) = 2$ and the minimum value is $f\left(-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right) = -2$.

16. $f(x_1, x_2, \ldots, x_n) = x_1 + x_2 + \cdots + x_n,\ g(x_1, x_2, \ldots, x_n) = x_1^2 + x_2^2 + \cdots + x_n^2 = 1 \quad \Rightarrow$ $\langle 1, 1, \ldots, 1\rangle = \langle 2\lambda x_1, 2\lambda x_2, \ldots, 2\lambda x_n\rangle$, so $\lambda = 1/(2x_1) = 1/(2x_2) = \cdots = 1/(2x_n)$ and $x_1 = x_2 = \cdots = x_n$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

But $x_1^2 + x_2^2 + \cdots + x_n^2 = 1$, so $x_i = \pm 1/\sqrt{n}$ for $i = 1, \ldots, n$. Thus the maximum value of f is $f(1/\sqrt{n}, 1/\sqrt{n}, \ldots, 1/\sqrt{n}) = \sqrt{n}$ and the minimum value is $f(-1/\sqrt{n}, -1/\sqrt{n}, \ldots, -1/\sqrt{n}) = -\sqrt{n}$.

17. If the two numbers are x and y, we want to minimize $f(x, y) = x + y$, $x > 0$, $y > 0$ subject to $g(x, y) = xy = 100$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 1, 1\rangle = \langle \lambda y, \lambda x\rangle$, so $1 = \lambda y$, $1 = \lambda x$, and $xy = 100$. The first two equations imply $y = \dfrac{1}{\lambda} = x$ and substitution into the third equation gives $\dfrac{1}{\lambda^2} = 100 \quad \Rightarrow \quad \lambda = \pm\dfrac{1}{10}$. Since $x > 0$ and $y > 0$, we have $\lambda = \dfrac{1}{10}$ and hence, $x = y = \dfrac{1}{\lambda} = 10$. Thus, the minimum value of f is $f(10, 10) = 20$. By comparing nearby values we can confirm that this gives a minimum and not a maximum. Therefore, the two numbers are 10 and 10.

18. If x and y are the dimensions of the rectangle in meters, we want to minimize $f(x, y) = 2x + 2y$, $x > 0$, $y > 0$ subject to $g(x, y) = xy = 1000$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2, 2\rangle = \langle \lambda y, \lambda x\rangle$, so $2 = \lambda y$, $2 = \lambda x$, and $xy = 1000$. The first two equations imply $y = \dfrac{2}{\lambda} = x$ and substitution into the third equation gives $\dfrac{4}{\lambda^2} = 1000 \quad \Rightarrow \quad \lambda = \pm\dfrac{1}{5\sqrt{10}}$. Since $x > 0$ and $y > 0$, we have $\lambda = \dfrac{1}{5\sqrt{10}}$ and hence, $x = y = \dfrac{2}{\lambda} = 10\sqrt{10}$. Thus, the minimum value of f is $f\left(10\sqrt{10}, 10\sqrt{10}\right) = 40\sqrt{10}$. By comparing nearby values we can confirm that this gives a minimum and not a maximum. Therefore, the dimensions of the rectangle that minimize the perimeter are $10\sqrt{10}$ m by $10\sqrt{10}$ m.

19. Let x and y be the dimensions of the rectangle in meters. Then the perimeter constraint is given by $2x + 2y = 100 \quad \Rightarrow$ $x + y = 50$. We want to maximize $f(x, y) = xy$, $x > 0$, $y > 0$ subject to $g(x, y) = x + y = 50$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow$ $\langle y, x\rangle = \langle \lambda, \lambda\rangle$, so $y = \lambda$, $x = \lambda$, and $x + y = 50$. Substituting the first two equations into the third gives $2\lambda = 50 \quad \Rightarrow$ $\lambda = 25$. Thus, the area is maximized when $x = y = 25$ meters and $f(25, 25) = 625\ \text{m}^2$.

20. Let x, y, and z be the dimensions of the box. The box has a square bottom, so $x = y$. We want to minimize $f(x, z) = x^2 + 4xz$ subject to $g(x, z) = x^2 z = 32{,}000$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2x + 4z, 4x\rangle = \left\langle 2\lambda xz, \lambda x^2\right\rangle$, so $2x + 4z = 2\lambda xz$, $4x = \lambda x^2$, and $x^2 z = 32{,}000$. The second equation implies $x = 0$ or $\lambda x = 4$. Note that $x = 0$ results in a zero volume. So let $\lambda = 4/x$. Substitution into the first equation gives $2x + 4z = 8z \quad \Rightarrow \quad x = 2z$, and substituting $2z$ for x into the third equation gives $4z^3 = 32{,}000 \quad \Rightarrow \quad z = 20 \quad \Rightarrow \quad x = 40$. Therefore, the dimensions of the box that will minimize the surface area are 40 cm by 40 cm by 20 cm.

21. The distance d from any point (x, y) in the xy-plane to the origin is given by $d = \sqrt{x^2 + y^2}$. We will minimize $d^2 = f(x, y) = x^2 + y^2$ subject to $g(x, y) = -2x + y = 3$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2x, 2y\rangle = \langle -2\lambda, \lambda\rangle$, so $2x = -2\lambda$, $2y = \lambda$, and $-2x + y = 3$. The first equation implies $x = -\lambda$, and the second, $y = \dfrac{\lambda}{2}$. Then substitution into the third

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

equation gives $2\lambda + \dfrac{\lambda}{2} = 3 \quad \Rightarrow \quad \lambda = \frac{6}{5} \quad \Rightarrow \quad x = -\frac{6}{5}$ and $y = \frac{3}{5}$. Thus, the point on the line $y = 2x + 3$ that is closest to the origin is $\left(-\frac{6}{5}, \frac{3}{5}\right)$.

22. Let r and h be the radius and height of a right circular cylinder. We want to maximize $V(r, h) = \pi r^2 h$ subject to $g(r, h) = h + 2\pi r = 108$. Then $\nabla V = \lambda \nabla g \quad \Rightarrow \quad \left\langle 2\pi rh, \pi r^2 \right\rangle = \langle 2\pi\lambda, \lambda \rangle$, so $2\pi rh = 2\pi\lambda$, $\pi r^2 = \lambda$, and $h + 2\pi r = 108$. The first equation implies $rh = \lambda$ and substitution into the second gives $\pi r^2 = rh \quad \Rightarrow \quad r = 0$ or $h = \pi r$. $r \neq 0$ (else, $V = 0$), so substitute $h = \pi r$ into the third equation. Then $\pi r + 2\pi r = 108 \quad \Rightarrow \quad \pi r = 36 \quad \Rightarrow \quad h = 36$. Therefore, the dimensions that maximize the volume are $r = \dfrac{36}{\pi}$ inches and $h = 36$ inches for a volume of $V = \dfrac{46{,}656}{\pi}$ in^3.

23. $f(x, y) = x^2 + y^2,\ g(x, y) = xy = 1$, and $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2x, 2y \rangle = \langle \lambda y, \lambda x \rangle$, so $2x = \lambda y$, $2y = \lambda x$, and $xy = 1$. From the last equation, $x \neq 0$ and $y \neq 0$, so $2x = \lambda y \quad \Rightarrow \quad \lambda = 2x/y$. Substituting, we have $2y = (2x/y)\,x \quad \Rightarrow$ $y^2 = x^2 \quad \Rightarrow \quad y = \pm x$. But $xy = 1$, so $x = y = \pm 1$ and the possible points for the extreme values of f are $(1, 1)$ and $(-1, -1)$. Here there is no maximum value, since the constraint $xy = 1 \quad \Leftrightarrow \quad y = 1/x$ allows x or y to become arbitrarily large, and hence $f(x, y) = x^2 + y^2$ can be made arbitrarily large. The minimum value is $f(1, 1) = f(-1, -1) = 2$.

24. $f(x, y, z) = x^2 + 2y^2 + 3z^2,\ g(x, y) = x + 2y + 3z = 10$, and $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2x, 4y, 6z \rangle = \langle \lambda, 2\lambda, 3\lambda \rangle$, so $2x = \lambda$, $4y = 2\lambda$, $6z = 3\lambda$, and $x + 2y + 3z = 10$. From the first three equations we have $2x = \lambda = 2y = 2z \quad \Rightarrow \quad x = y = z$, and substituting into the fourth equation gives $x + 2x + 3x = 10 \quad \Rightarrow \quad x = \frac{5}{3} = y = z$. Thus the only possible point for an extreme value of f is $\left(\frac{5}{3}, \frac{5}{3}, \frac{5}{3}\right)$. Notice here that the constraint $x + 2y + 3z = 10$ allows any of $|x|$, $|y|$, or $|z|$ to be arbitrarily large, and hence $f(x, y, z) = x^2 + 2y^2 + 3z^2$ can be made arbitrarily large. So f has no maximum value subject to the constraint. The minimum value is $f\left(\frac{5}{3}, \frac{5}{3}, \frac{5}{3}\right) = 6\left(\frac{5}{3}\right)^2 = \frac{50}{3}$.

25. $f(x, y) = e^{xy}, g(x, y) = x^3 + y^3 = 16$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle ye^{xy}, xe^{xy} \rangle = \left\langle 3\lambda x^2, 3\lambda y^2 \right\rangle$, so $ye^{xy} = 3\lambda x^2$, $xe^{xy} = 3\lambda y^2$, and $x^3 + y^3 = 16$. Multiplying the first equation by x and the second by y gives $xye^{xy} = 3\lambda x^3$ and $xye^{xy} = 3\lambda y^3 \quad \Rightarrow \quad 3\lambda x^3 = 3\lambda y^3 \quad \Rightarrow \quad x = y$ and substituting x for y into the third equation, we have $x^3 + x^3 = 16 \quad \Rightarrow$ $x = y = 2$. Therefore, f has an extreme value at the point $(2, 2)$ and evaluating f at that point we see $f(2, 2) = e^4$. Notice from the constraint that $y^3 = 16 - x^3$ and y^3 becomes increasingly negative as x^3 becomes arbitrarily large (similarly, x^3 can be increasingly negative while y^3 is arbitrarily large). Thus, for any small value $\varepsilon > 0$, we can find x and y that satisfy the constraint such that $0 < e^{xy} < \varepsilon$. Thus, f has no minimum value subject to the constraint and $f(2, 2) = e^4$ is a maximum.

26. $f(x, y, z) = 4x + 2y + z, g(x, y, z) = x^2 + y + z^2 = 1$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 4, 2, 1 \rangle = \langle 2\lambda x, \lambda, 2\lambda z \rangle$, so $4 = 2\lambda x$, $2 = \lambda$, $1 = 2\lambda z$, and $x^2 + y + z^2 = 1$. As $\lambda = 2$ by equation two, we have $x = 1$ and $z = \frac{1}{4}$ by equations one and three, respectively. Substituting these values into equation four gives $1^2 + y + \left(\frac{1}{4}\right)^2 = 1 \quad \Rightarrow \quad y = -\frac{1}{16}$. Thus, f has an extreme value at $\left(1, -\frac{1}{16}, \frac{1}{4}\right)$. Notice that $y = 1 - (x^2 + z^2) < 0$ for $x^2 + z^2 > 1$, and substituting into f, we see that

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$f(x,y,z) = 4x + 2 + z - 2(x^2 + z^2)$ can decrease without bound for values of (x,y,z) that satisfy the constraint. Thus, f has no minimum value subject to the constraint and $f\left(1, -\frac{1}{16}, \frac{1}{4}\right) = \frac{33}{8}$ is a maximum.

27. $f(x,y) = x^2 + y^2 + 4x - 4y$. For the interior of the region, we find the critical points: $f_x = 2x + 4$, $f_y = 2y - 4$, so the only critical point is $(-2,2)$ (which is inside the region) and $f(-2,2) = -8$. For the boundary, we use Lagrange multipliers. $g(x,y) = x^2 + y^2 = 9$, so $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2x+4, 2y-4\rangle = \langle 2\lambda x, 2\lambda y\rangle$. Thus $2x + 4 = 2\lambda x$ and $2y - 4 = 2\lambda y$. Adding the two equations gives $2x + 2y = 2\lambda x + 2\lambda y \quad \Rightarrow \quad x + y = \lambda(x+y) \quad \Rightarrow \quad (x+y)(\lambda - 1) = 0$, so $x + y = 0 \quad \Rightarrow \quad y = -x$ or $\lambda - 1 = 0 \quad \Rightarrow \quad \lambda = 1$. But $\lambda = 1$ leads to a contradition in $2x + 4 = 2\lambda x$, so $y = -x$ and $x^2 + y^2 = 9$ implies $2y^2 = 9 \quad \Rightarrow \quad y = \pm\frac{3}{\sqrt{2}}$. We have $f\left(\frac{3}{\sqrt{2}}, -\frac{3}{\sqrt{2}}\right) = 9 + 12\sqrt{2} \approx 25.97$ and $f\left(-\frac{3}{\sqrt{2}}, \frac{3}{\sqrt{2}}\right) = 9 - 12\sqrt{2} \approx -7.97$, so the maximum value of f on the disk $x^2 + y^2 \le 9$ is $f\left(\frac{3}{\sqrt{2}}, -\frac{3}{\sqrt{2}}\right) = 9 + 12\sqrt{2}$ and the minimum is $f(-2,2) = -8$.

28. $f(x,y) = 2x^2 + 3y^2 - 4x - 5 \quad \Rightarrow \quad \nabla f = \langle 4x - 4, 6y\rangle = \langle 0, 0\rangle \quad \Rightarrow \quad x = 1, y = 0$. Thus $(1,0)$ is the only critical point of f, and it lies in the region $x^2 + y^2 < 16$. On the boundary, $g(x,y) = x^2 + y^2 = 16 \quad \Rightarrow \quad \lambda \nabla g = \langle 2\lambda x, 2\lambda y\rangle$, so $6y = 2\lambda y \quad \Rightarrow \quad$ either $y = 0$ or $\lambda = 3$. If $y = 0$, then $x = \pm 4$; if $\lambda = 3$, then $4x - 4 = 2\lambda x \quad \Rightarrow \quad x = -2$ and $y = \pm 2\sqrt{3}$. Now $f(1,0) = -7$, $f(4,0) = 11$, $f(-4,0) = 43$, and $f\left(-2, \pm 2\sqrt{3}\right) = 47$. Thus the maximum value of $f(x,y)$ on the disk $x^2 + y^2 \le 16$ is $f\left(-2, \pm 2\sqrt{3}\right) = 47$, and the minimum value is $f(1,0) = -7$.

29. $f(x,y) = e^{-xy}$. For the interior of the region, we find the critical points: $f_x = -ye^{-xy}$, $f_y = -xe^{-xy}$, so the only critical point is $(0,0)$, and $f(0,0) = 1$. For the boundary, we use Lagrange multipliers. $g(x,y) = x^2 + 4y^2 = 1 \quad \Rightarrow \quad \lambda \nabla g = \langle 2\lambda x, 8\lambda y\rangle$, so setting $\nabla f = \lambda \nabla g$ we get $-ye^{-xy} = 2\lambda x$ and $-xe^{-xy} = 8\lambda y$. The first of these gives $e^{-xy} = -2\lambda x/y$, and then the second gives $-x(-2\lambda x/y) = 8\lambda y \quad \Rightarrow \quad x^2 = 4y^2$. Solving this last equation with the constraint $x^2 + 4y^2 = 1$ gives $x = \pm\frac{1}{\sqrt{2}}$ and $y = \pm\frac{1}{2\sqrt{2}}$. Now $f\left(\pm\frac{1}{\sqrt{2}}, \mp\frac{1}{2\sqrt{2}}\right) = e^{1/4} \approx 1.284$ and $f\left(\pm\frac{1}{\sqrt{2}}, \pm\frac{1}{2\sqrt{2}}\right) = e^{-1/4} \approx 0.779$. The former are the maximums on the region and the latter are the minimums.

30. $f(x,y,z) = z,\ g(x,y,z) = x^2 + y^2 - z^2 = 0,\ h(x,y,z) = x + y + z = 24$, and $\nabla f = \lambda \nabla g + \mu \nabla h \quad \Rightarrow \quad \langle 0,0,1\rangle = \langle 2\lambda x, 2\lambda y, -2\lambda z\rangle + \langle \mu, \mu, \mu\rangle$. Then $0 = 2\lambda x + \mu,\ \ 0 = 2\lambda y + \mu,\ \ 1 = -2\lambda z + \mu,\ \ x^2 + y^2 - z^2 = 0,$ and $x + y + z = 24$. From the first two equations we have $-2\lambda x = \mu = -2\lambda y \quad \Rightarrow \quad \lambda = 0$ or $x = y$. But $\lambda = 0 \quad \Rightarrow \quad \mu = 0$ which contradicts the third equation, so $x = y$ and substitution into the last equation gives $z = 24 - 2x$. From the fourth equation we have $x^2 + x^2 - (24 - 2x)^2 = 0 \quad \Rightarrow \quad -2x^2 + 96x - 576 = 0 \quad \Rightarrow \quad x^2 - 48x + 288 = 0 \quad \Rightarrow$

$$x = \frac{48 \pm \sqrt{1152}}{2} = 24 \pm 12\sqrt{2} = y.$$

Now $z = 24 - 2x$, so the possible points are $\left(24 + 12\sqrt{2}, 24 + 12\sqrt{2}, -24 - 24\sqrt{2}\right)$ and $\left(24 - 12\sqrt{2}, 24 - 12\sqrt{2}, -24 + 24\sqrt{2}\right)$. The maximum of f subject to the constraints is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$f\left(24-12\sqrt{2}, 24-12\sqrt{2}, -24+24\sqrt{2}\right) = -24+24\sqrt{2} \approx 9.94$ and the minimum is

$f\left(24+12\sqrt{2}, 24+12\sqrt{2}, -24-24\sqrt{2}\right) = -24-24\sqrt{2} \approx -57.94$.

31. $f(x,y,z) = x+y+z,\ g(x,y,z) = x^2+z^2 = 2,\ h(x,y,z) = x+y = 1$, and $\nabla f = \lambda\nabla g + \mu\nabla h \quad \Rightarrow$ $\langle 1,1,1\rangle = \langle 2\lambda x, 0, 2\lambda z\rangle + \langle \mu,\mu,0\rangle$. Then $1 = 2\lambda x+\mu,\ \ 1=\mu,\ \ 1 = 2\lambda z,\ \ x^2+z^2=2,$ and $x+y=1$. Substituting $\mu = 1$ into the first equation gives $\lambda = 0$ or $x = 0$. But $\lambda = 0$ contradicts $1 = 2\lambda z$, so $x = 0$. Then $x+y=1 \quad\Rightarrow\quad y=1$ and $x^2+z^2=2 \quad\Rightarrow\quad z = \pm\sqrt{2}$, so the possible points are $\left(0,1,\pm\sqrt{2}\right)$. The maximum value of f subject to the constraints is $f(0,1,\sqrt{2}) = 1+\sqrt{2} \approx 2.41$ and the minimum is $f(0,1,-\sqrt{2}) = 1-\sqrt{2} \approx -0.41$.

Note: Since $x+y=1$ is one of the constraints, we could have solved the problem by solving $f(x,z) = 1+z$ subject to $x^2+z^2=2$.

32. $f(x,y,z) = x^2+y^2+z^2,\ g(x,y,z) = x-y=1,\ h(x,y,z) = y^2-z^2=1 \quad\Rightarrow\quad \nabla f = \langle 2x,2y,2z\rangle$, $\lambda\nabla g = \langle \lambda,-\lambda,0\rangle$, and $\mu\nabla h = \langle 0, 2\mu y, -2\mu z\rangle$. Then $2x=\lambda$, $2y = -\lambda+2\mu y$, and $2z = -2\mu z \quad\Rightarrow\quad z=0$ or $\mu=-1$. If $z=0$ then $y^2-z^2=1$ implies $y^2=1 \quad\Rightarrow\quad y=\pm1$. If $y=1$, $x-y=1$ implies $x=2$, and if $y=-1$ we have $x=0$, so possible points are $(2,1,0)$ and $(0,-1,0)$. If $\mu=-1$ then $2y=-\lambda+2\mu y$ implies $4y=-\lambda$, but $\lambda=2x$ so $4y=-2x \quad\Rightarrow\quad x=-2y$ and $x-y=1$ implies $-3y=1 \quad\Rightarrow\quad y=-\frac{1}{3}$. But then $y^2-z^2=1$ implies $z^2=-\frac{8}{9}$, an impossibility. Thus the maximum value of f subject to the constraints is $f(2,1,0)=5$ and the minimum is $f(0,-1,0)=1$.

Note: Since $x-y=1 \quad\Rightarrow\quad x=y+1$ is one of the constraints we could have solved the problem by solving $f(y,z) = (y+1)^2+y^2+z^2$ subject to $y^2-z^2=1$.

33. $f(x,y,z) = yz+xy,\ g(x,y,z) = xy=1,\ h(x,y,z) = y^2+z^2=1 \quad\Rightarrow\quad \nabla f = \langle y, x+z, y\rangle$, $\lambda\nabla g = \langle \lambda y, \lambda x, 0\rangle$, $\mu\nabla h = \langle 0, 2\mu y, 2\mu z\rangle$. Then $y=\lambda y$ implies $\lambda = 1$ [$y\neq 0$ since $g(x,y,z)=1$], $x+z = \lambda x+2\mu y$ and $y = 2\mu z$. Thus $\mu = z/(2y) = y/(2y)$ or $y^2=z^2$, and so $y^2+z^2=1$ implies $y=\pm\frac{1}{\sqrt{2}}$, $z=\pm\frac{1}{\sqrt{2}}$. Then $xy=1$ implies $x=\pm\sqrt{2}$ and the possible points are $\left(\pm\sqrt{2},\pm\frac{1}{\sqrt{2}},\frac{1}{\sqrt{2}}\right)$, $\left(\pm\sqrt{2},\pm\frac{1}{\sqrt{2}},-\frac{1}{\sqrt{2}}\right)$. Hence the maximum of f subject to the constraints is $f\left(\pm\sqrt{2},\pm\frac{1}{\sqrt{2}},\pm\frac{1}{\sqrt{2}}\right) = \frac{3}{2}$ and the minimum is $f\left(\pm\sqrt{2},\pm\frac{1}{\sqrt{2}},\mp\frac{1}{\sqrt{2}}\right) = \frac{1}{2}$.

Note: Since $xy=1$ is one of the constraints we could have solved the problem by solving $f(y,z) = yz+1$ subject to $y^2+z^2=1$.

34. (a) $f(x,y) = 2x+3y,\ g(x,y) = \sqrt{x}+\sqrt{y} = 5 \quad\Rightarrow\quad \nabla f = \langle 2,3\rangle = \lambda\nabla g = \lambda\left\langle \dfrac{1}{2\sqrt{x}}, \dfrac{1}{2\sqrt{y}}\right\rangle$. Then

$2 = \dfrac{\lambda}{2\sqrt{x}}$ and $3 = \dfrac{\lambda}{2\sqrt{y}}$ so $4\sqrt{x} = \lambda = 6\sqrt{y} \quad\Rightarrow\quad \sqrt{y} = \frac{2}{3}\sqrt{x}$. With $\sqrt{x}+\sqrt{y}=5$ we have $\sqrt{x}+\frac{2}{3}\sqrt{x} = 5 \quad\Rightarrow$ $\sqrt{x}=3 \quad\Rightarrow\quad x=9$. Substituting into $\sqrt{y} = \frac{2}{3}\sqrt{x}$ gives $\sqrt{y}=2$ or $y=4$. Thus the only possible extreme value subject to the constraint is $f(9,4)=30$. (The question remains whether this is indeed the maximum of f.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $f(25,0) = 50$ which is larger than the result of part (a).

(c)

We can see from the level curves of f that the maximum occurs at the left endpoint $(0,25)$ of the constraint curve g. The maximum value is $f(0,25) = 75$.

(d) Here ∇g does not exist if $x = 0$ or $y = 0$, so the method will not locate any associated points. Also, the method of Lagrange multipliers identifies points where the level curves of f share a common tangent line with the constraint curve g. This normally does not occur at an endpoint, although an absolute maximum or minimum may occur there.

(e) Here $f(9,4)$ is the absolute *minimum* of f subject to g.

35. (a) $f(x,y) = x,\ \ g(x,y) = y^2 + x^4 - x^3 = 0 \ \ \Rightarrow\ \ \nabla f = \langle 1,0\rangle = \lambda\nabla g = \lambda\left\langle 4x^3 - 3x^2, 2y\right\rangle$. Then $1 = \lambda(4x^3 - 3x^2)$ **(1)** and $0 = 2\lambda y$ **(2)**. We have $\lambda \neq 0$ from **(1)**, so **(2)** gives $y = 0$. Then, from the constraint equation, $x^4 - x^3 = 0 \ \ \Rightarrow\ \ x^3(x-1) = 0 \ \ \Rightarrow\ \ x = 0$ or $x = 1$. But $x = 0$ contradicts **(1)**, so the only possible extreme value subject to the constraint is $f(1,0) = 1$. (The question remains whether this is indeed the minimum of f.)

(b) The constraint is $y^2 + x^4 - x^3 = 0 \ \ \Leftrightarrow\ \ y^2 = x^3 - x^4$. The left side is nonnegative, so we must have $x^3 - x^4 \geq 0$ which is true only for $0 \leq x \leq 1$. Therefore the minimum possible value for $f(x,y) = x$ is 0 which occurs for $x = y = 0$. However, $\lambda\nabla g(0,0) = \lambda\langle 0-0, 0\rangle = \langle 0,0\rangle$ and $\nabla f(0,0) = \langle 1,0\rangle$, so $\nabla f(0,0) \neq \lambda\nabla g(0,0)$ for all values of λ.

(c) Here $\nabla g(0,0) = \mathbf{0}$ but the method of Lagrange multipliers requires that $\nabla g \neq \mathbf{0}$ everywhere on the constraint curve.

36. (a) The graphs of $f(x,y) = 3.7$ and $f(x,y) = 350$ seem to be tangent to the circle, and so 3.7 and 350 are the approximate minimum and maximum values of the function $f(x,y) = x^3 + y^3 + 3xy$ subject to the constraint $(x-3)^2 + (y-3)^2 = 9$.

(b) Let $g(x,y) = (x-3)^2 + (y-3)^2$. We calculate $f_x(x,y) = 3x^2 + 3y$, $f_y(x,y) = 3y^2 + 3x$, $g_x(x,y) = 2x - 6$, and $g_y(x,y) = 2y - 6$, and use a CAS to search for solutions to the equations $g(x,y) = (x-3)^2 + (y-3)^2 = 9$, $f_x = \lambda g_x$, and $f_y = \lambda g_y$. The solutions are $(x,y) = \left(3 - \frac{3}{2}\sqrt{2}, 3 - \frac{3}{2}\sqrt{2}\right) \approx (0.879, 0.879)$ and $(x,y) = \left(3 + \frac{3}{2}\sqrt{2}, 3 + \frac{3}{2}\sqrt{2}\right) \approx (5.121, 5.121)$. These give $f\left(3 - \frac{3}{2}\sqrt{2}, 3 - \frac{3}{2}\sqrt{2}\right) = \frac{351}{2} - \frac{243}{2}\sqrt{2} \approx 3.673$ and $f\left(3 + \frac{3}{2}\sqrt{2}, 3 + \frac{3}{2}\sqrt{2}\right) = \frac{351}{2} + \frac{243}{2}\sqrt{2} \approx 347.33$, in accordance with part (a).

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. $P(L,K) = bL^{\alpha}K^{1-\alpha},\ g(L,K) = mL + nK = p \quad \Rightarrow \quad \nabla P = \left\langle \alpha bL^{\alpha-1}K^{1-\alpha}, (1-\alpha)bL^{\alpha}K^{-\alpha} \right\rangle,\ \lambda\nabla g = \langle \lambda m, \lambda n \rangle.$

Then $\alpha b(K/L)^{1-\alpha} = \lambda m$ and $(1-\alpha)b(L/K)^{\alpha} = \lambda n$ and $mL + nK = p$, so $\alpha b(K/L)^{1-\alpha}/m = (1-\alpha)b(L/K)^{\alpha}/n$ or $n\alpha/[m(1-\alpha)] = (L/K)^{\alpha}(L/K)^{1-\alpha}$ or $L = Kn\alpha/[m(1-\alpha)]$. Substituting into $mL + nK = p$ gives $K = (1-\alpha)p/n$ and $L = \alpha p/m$ for the maximum production.

38. $C(L,K) = mL + nK,\ g(L,K) = bL^{\alpha}K^{1-\alpha} = Q \quad \Rightarrow \quad \nabla C = \langle m, n \rangle,\ \lambda\nabla g = \left\langle \lambda\alpha bL^{\alpha-1}K^{1-\alpha}, \lambda(1-\alpha)bL^{\alpha}K^{-\alpha} \right\rangle.$

Then $\dfrac{m}{\alpha b}\left(\dfrac{L}{K}\right)^{1-\alpha} = \dfrac{n}{(1-\alpha)b}\left(\dfrac{K}{L}\right)^{\alpha}$ and $bL^{\alpha}K^{1-\alpha} = Q \quad \Rightarrow \quad \dfrac{n\alpha}{m(1-\alpha)} = \left(\dfrac{L}{K}\right)^{1-\alpha}\left(\dfrac{L}{K}\right)^{\alpha} \quad \Rightarrow$

$L = \dfrac{Kn\alpha}{m(1-\alpha)}$ and so $b\left[\dfrac{Kn\alpha}{m(1-\alpha)}\right]^{\alpha}K^{1-\alpha} = Q$. Hence $K = \dfrac{Q}{b\left(n\alpha/[m(1-\alpha)]\right)^{\alpha}} = \dfrac{Qm^{\alpha}(1-\alpha)^{\alpha}}{bn^{\alpha}\alpha^{\alpha}}$

and $L = \dfrac{Qm^{\alpha-1}(1-\alpha)^{\alpha-1}}{bn^{\alpha-1}\alpha^{\alpha-1}} = \dfrac{Qn^{1-\alpha}\alpha^{1-\alpha}}{bm^{1-\alpha}(1-\alpha)^{1-\alpha}}$ minimizes cost.

39. Let the sides of the rectangle be x and y. Then $f(x,y) = xy$, $g(x,y) = 2x + 2y = p \quad \Rightarrow \quad \nabla f(x,y) = \langle y, x \rangle$, $\lambda\nabla g = \langle 2\lambda, 2\lambda \rangle$. Then $\lambda = \frac{1}{2}y = \frac{1}{2}x$ implies $x = y$ and the rectangle with maximum area is a square with side length $\frac{1}{4}p$.

40. We maximize $A^2 = f(x,y,z) = s(s-x)(s-y)(s-z)$ subject to $g(x,y,z) = x + y + z$. Then $\nabla f = \langle -s(s-y)(s-z), -s(s-x)(s-z), -s(s-x)(s-y) \rangle$, $\lambda\nabla g = \langle \lambda, \lambda, \lambda \rangle$. Thus $(s-y)(s-z) = (s-x)(s-z)$ **(1)**, and $(s-x)(s-z) = (s-x)(s-y)$ **(2)**. **(1)** implies $x = y$ while **(2)** implies $y = z$, so $x = y = z = p/3$ and the triangle with maximum area is equilateral.

41. The distance from $(2,0,-3)$ to a point (x,y,z) on the plane is $d = \sqrt{(x-2)^2 + y^2 + (z+3)^2}$, so we seek to minimize $d^2 = f(x,y,z) = (x-2)^2 + y^2 + (z+3)^2$ subject to the constraint that (x,y,z) lies on the plane $x + y + z = 1$, that is, that $g(x,y,z) = x + y + z = 1$. Then $\nabla f = \lambda\nabla g \quad \Rightarrow \quad \langle 2(x-2), 2y, 2(z+3) \rangle = \langle \lambda, \lambda, \lambda \rangle$, so $x = (\lambda+4)/2$, $y = \lambda/2$, $z = (\lambda-6)/2$. Substituting into the constraint equation gives $\dfrac{\lambda+4}{2} + \dfrac{\lambda}{2} + \dfrac{\lambda-6}{2} = 1 \quad \Rightarrow \quad 3\lambda - 2 = 2 \quad \Rightarrow$ $\lambda = \frac{4}{3}$, so $x = \frac{8}{3}$, $y = \frac{2}{3}$, and $z = -\frac{7}{3}$. This must correspond to a minimum, so the shortest distance is $d = \sqrt{\left(\frac{8}{3} - 2\right)^2 + \left(\frac{2}{3}\right)^2 + \left(-\frac{7}{3} + 3\right)^2} = \sqrt{\frac{4}{3}} = \frac{2}{\sqrt{3}}$.

42. The distance from $(0,1,1)$ to a point (x,y,z) on the plane is $d = \sqrt{x^2 + (y-1)^2 + (z-1)^2}$, so we minimize $d^2 = f(x,y,z) = x^2 + (y-1)^2 + (z-1)^2$ subject to the constraint that (x,y,z) lies on the plane $x - 2y + 3z = 6$, that is, $g(x,y,z) = x - 2y + 3z = 6$. Then $\nabla f = \lambda\nabla g \quad \Rightarrow \quad \langle 2x, 2(y-1), 2(z-1) \rangle = \langle \lambda, -2\lambda, 3\lambda \rangle$, so $x = \lambda/2$, $y = 1 - \lambda$, $z = (3\lambda+2)/2$. Substituting into the constraint equation gives $\dfrac{\lambda}{2} - 2(1-\lambda) + 3 \cdot \dfrac{3\lambda+2}{2} = 6 \quad \Rightarrow \quad \lambda = \frac{5}{7}$, so $x = \frac{5}{14}$, $y = \frac{2}{7}$, and $z = \frac{29}{14}$. This must correspond to a minimum, so the point on the plane closest to the point $(0,1,1)$ is $\left(\frac{5}{14}, \frac{2}{7}, \frac{29}{14}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. Let $f(x,y,z) = d^2 = (x-4)^2 + (y-2)^2 + z^2$. Then we want to minimize f subject to the constraint $g(x,y,z) = x^2 + y^2 - z^2 = 0$. $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2(x-4), 2(y-2), 2z \rangle = \langle 2\lambda x, 2\lambda y, -2\lambda z \rangle$, so $x - 4 = \lambda x$, $y - 2 = \lambda y$, and $z = -\lambda z$. From the last equation we have $z + \lambda z = 0 \quad \Rightarrow \quad z(1+\lambda) = 0$, so either $z = 0$ or $\lambda = -1$. But from the constraint equation we have $z = 0 \quad \Rightarrow \quad x^2 + y^2 = 0 \quad \Rightarrow \quad x = y = 0$ which is not possible from the first two equations. So $\lambda = -1$ and $x - 4 = \lambda x \quad \Rightarrow \quad x = 2$, $y - 2 = \lambda y \quad \Rightarrow \quad y = 1$, and $x^2 + y^2 - z^2 = 0 \quad \Rightarrow$ $4 + 1 - z^2 = 0 \quad \Rightarrow \quad z = \pm\sqrt{5}$. This must correspond to a minimum, so the points on the cone closest to $(4,2,0)$ are $\left(2, 1, \pm\sqrt{5}\right)$.

44. Let $f(x,y,z) = d^2 = x^2 + y^2 + z^2$. Then we want to minimize f subject to the constraint $g(x,y,z) = y^2 - xz = 9$. $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2x, 2y, 2z \rangle = \langle -\lambda z, 2\lambda y, -\lambda x \rangle$, so $2x = -\lambda z$, $y = \lambda y$, and $2z = -\lambda x$. If $x = 0$ then the last equation implies $z = 0$, and from the constraint $y^2 - xz = 9$ we have $y = \pm 3$. If $x \neq 0$, then the first and third equations give $\lambda = -2x/z = -2z/x \quad \Rightarrow \quad x^2 = z^2$. From the second equation we have $y = 0$ or $\lambda = 1$. If $y = 0$ then $y^2 - xz = 9 \quad \Rightarrow$ $z = -9/x$ and $x^2 = z^2 \quad \Rightarrow \quad x^2 = 81/x^2 \quad \Rightarrow \quad x = \pm 3$. Since $z = -9/x$, $x = 3 \quad \Rightarrow \quad z = -3$ and $x = -3 \quad \Rightarrow$ $z = 3$. If $\lambda = 1$, then $2x = -z$ and $2z = -x$ which implies $x = z = 0$, contradicting the assumption that $x \neq 0$. Thus the possible points are $(0, \pm 3, 0)$, $(3, 0, -3)$, $(-3, 0, 3)$. We have $f(0, \pm 3, 0) = 9$ and $f(3, 0, -3) = f(-3, 0, 3) = 18$, so the points on the surface that are closest to the origin are $(0, \pm 3, 0)$.

45. Maximize $f(x,y,z) = xyz$ subject to $g(x,y,z) = x + y + z = 100$. $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle yz, xz, xy \rangle = \lambda \langle 1, 1, 1 \rangle$. Then $\lambda = yz = xz = xy$ implies $x = y = z = \frac{100}{3}$.

46. Minimize $f(x,y,z) = x^2 + y^2 + z^2$ subject to $g(x,y,z) = x + y + z = 12$ with $x > 0$, $y > 0$, $z > 0$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 2x, 2y, 2z \rangle = \lambda \langle 1, 1, 1 \rangle \quad \Rightarrow \quad 2x = \lambda$, $2y = \lambda$, $2z = \lambda \quad \Rightarrow \quad x = y = z$, so $x + y + z = 12 \quad \Rightarrow$ $3x = 12 \quad \Rightarrow \quad x = 4 = y = z$. By comparing nearby values we can confirm that this gives a minimum and not a maximum. Thus the three numbers are 4, 4, and 4.

47. If the dimensions are $2x$, $2y$, and $2z$, then maximize $f(x,y,z) = (2x)(2y)(2z) = 8xyz$ subject to $g(x,y,z) = x^2 + y^2 + z^2 = r^2$ $(x > 0, y > 0, z > 0)$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 8yz, 8xz, 8xy \rangle = \lambda \langle 2x, 2y, 2z \rangle \quad \Rightarrow$ $8yz = 2\lambda x$, $8xz = 2\lambda y$, and $8xy = 2\lambda z$, so $\lambda = \dfrac{4yz}{x} = \dfrac{4xz}{y} = \dfrac{4xy}{z}$. This gives $x^2 z = y^2 z \quad \Rightarrow \quad x^2 = y^2$ (since $z \neq 0$) and $xy^2 = xz^2 \quad \Rightarrow \quad z^2 = y^2$, so $x^2 = y^2 = z^2 \quad \Rightarrow \quad x = y = z$, and substituting into the constraint equation gives $3x^2 = r^2 \quad \Rightarrow \quad x = r/\sqrt{3} = y = z$. Thus the largest volume of such a box is $f\left(\frac{r}{\sqrt{3}}, \frac{r}{\sqrt{3}}, \frac{r}{\sqrt{3}}\right) = 8\left(\frac{r}{\sqrt{3}}\right)\left(\frac{r}{\sqrt{3}}\right)\left(\frac{r}{\sqrt{3}}\right) = \dfrac{8}{3\sqrt{3}} r^3$.

48. If the dimensions of the box are x, y, and z, then minimize $f(x,y,z) = 2xy + 2xz + 2yz$ subject to $g(x,y,z) = xyz = 1000$ $(x > 0, y > 0, z > 0)$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\langle 2y+2z, 2x+2z, 2x+2y\rangle = \lambda\langle yz, xz, xy\rangle \quad\Rightarrow\quad 2y+2z=\lambda yz,\ 2x+2z=\lambda xz,\ 2x+2y=\lambda xy$. Solving for λ in each equation gives $\lambda = \dfrac{2}{z}+\dfrac{2}{y}=\dfrac{2}{z}+\dfrac{2}{x}=\dfrac{2}{y}+\dfrac{2}{x} \quad\Rightarrow\quad x=y=z$. From $xyz=1000$ we have $x^3=1000 \quad\Rightarrow$ $x=10$ and the dimensions of the box are $x=y=z=10$ cm.

49. Maximize $f(x,y,z)=xyz$ subject to $g(x,y,z)=x+2y+3z=6$. $\nabla f=\lambda\nabla g \quad\Rightarrow\quad \langle yz, xz, xy\rangle=\lambda\langle 1,2,3\rangle$. Then $\lambda = yz = \frac{1}{2}xz = \frac{1}{3}xy$ implies $x=2y$, $z=\frac{2}{3}y$. But $2y+2y+2y=6$ so $y=1$, $x=2$, $z=\frac{2}{3}$ and the volume is $V=\frac{4}{3}$.

50. Maximize $f(x,y,z)=xyz$ subject to $g(x,y,z)=xy+yz+xz=32$. $\nabla f=\lambda\nabla g \quad\Rightarrow$ $\langle yz, xz, xy\rangle=\lambda\langle y+z, x+z, x+y\rangle$. Then $\lambda(y+z)=yz$ **(1)**, $\lambda(x+z)=xz$ **(2)**, and $\lambda(x+y)=xy$ **(3)**. And **(1)** minus **(2)** implies $\lambda(y-x)=z(y-x)$ so $x=y$ or $\lambda=z$. If $\lambda=z$, then **(1)** implies $z(y+z)=yz$ or $z=0$ which is false. Thus $x=y$. Similarly **(2)** minus **(3)** implies $\lambda(z-y)=x(z-y)$ so $y=z$ or $\lambda=x$. As above, $\lambda\neq x$, so $x=y=z$ and $3x^2=32$ or $x=y=z=\frac{8}{\sqrt{6}}$ cm.

51. Maximize $f(x,y,z)=xyz$ subject to $g(x,y,z)=4(x+y+z)=c$. $\nabla f=\lambda\nabla g \quad\Rightarrow\quad \langle yz, xz, xy\rangle=\lambda\langle 4,4,4\rangle$. Then $yz=4\lambda$, $xz=4\lambda$, and $xy=4\lambda$. Multiplying by x, y, and z, respectively, gives us $xyz=4\lambda x=4\lambda y=4\lambda z$, so $x=y=z$. Substituting y and z for x in g gives us $4(3x)=c \quad\Rightarrow\quad x=y=z=\frac{1}{12}c$ are the dimensions of the cube giving the maximum volume.

52. Maximize $C(x,y,z)=5xy+2xz+2yz$ subject to $g(x,y,z)=xyz=V$. $\nabla C=\lambda\nabla g \quad\Rightarrow$ $\langle 5y+2z, 5x+2z, 2x+2y\rangle=\lambda\langle yz, xz, xy\rangle$. Then $\lambda yz=5y+2z$ **(1)**, $\lambda xz=5x+2z$ **(2)**, $\lambda xy=2(x+y)$ **(3)**, and $xyz=V$ **(4)**. Now **(1)** $-$ **(2)** implies $\lambda z(y-x)=5(y-x)$, so $x=y$ or $\lambda=5/z$, but z can't be 0, so $x=y$. Then twice **(2)** minus five times **(3)** together with $x=y$ implies $\lambda y(2x-5y)=2(2z-5y)$ which gives $z=\frac{5}{2}y$ [again $\lambda\neq 2/y$ or else **(3)** implies $y=0$]. Hence $\frac{5}{2}y^3=V$ and the dimensions which minimize cost are $x=y=\sqrt[3]{\frac{2}{5}V}$ units, $z=V^{1/3}\left(\frac{5}{2}\right)^{2/3}$ units.

53. If the dimensions of the box are given by x, y, and z, then we need to find the maximum value of $f(x,y,z)=xyz$ $[x,y,z>0]$ subject to the constraint $L=\sqrt{x^2+y^2+z^2}$ or $g(x,y,z)=x^2+y^2+z^2=L^2$. $\nabla f=\lambda\nabla g \quad\Rightarrow$ $\langle yz, xz, xy\rangle=\lambda\langle 2x, 2y, 2z\rangle$, so $yz=2\lambda x \quad\Rightarrow\quad \lambda=\dfrac{yz}{2x}$, $xz=2\lambda y \quad\Rightarrow\quad \lambda=\dfrac{xz}{2y}$, and $xy=2\lambda z \quad\Rightarrow\quad \lambda=\dfrac{xy}{2z}$. Thus $\lambda=\dfrac{yz}{2x}=\dfrac{xz}{2y} \quad\Rightarrow\quad x^2=y^2$ [since $z\neq 0$] $\quad\Rightarrow\quad x=y$ and $\lambda=\dfrac{yz}{2x}=\dfrac{xy}{2z} \quad\Rightarrow\quad x=z$ [since $y\neq 0$]. Substituting into the constraint equation gives $x^2+x^2+x^2=L^2 \quad\Rightarrow\quad x^2=L^2/3 \quad\Rightarrow\quad x=L/\sqrt{3}=y=z$ and the maximum volume is $\left(L/\sqrt{3}\right)^3=L^3/\left(3\sqrt{3}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. Let the dimensions of the box be x, y, and z. Then we wish to maximize $f(x, y, z) = xyz$ subject to $g(x, y, z) = 2x + 2y + z = 108 \quad \Rightarrow \quad \nabla f = \langle yz, xz, xy \rangle$ and $\lambda \nabla g = \langle 2\lambda, 2\lambda, \lambda \rangle$. Now $yz = 2\lambda$, $xz = 2\lambda$, and $xy = \lambda$, with $x \neq 0$, $y \neq 0$, $z \neq 0$. Then $yz = xz \quad \Rightarrow \quad y = x$ **(1)** and $2xy = xz \quad \Rightarrow \quad 2y = z$ **(2)**. Substituting **(1)** and **(2)** into the constraint, we get $2y + 2y + 2y = 108 \quad \Rightarrow \quad y = x = 18$, and hence $z = 36$. Thus, the dimensions of the box that will give the largest volume and still meet USPS guidelines are 18 in by 18 in by 36 in.

55. If r and h are the radius and the height of the silo, respectively, we need to maximize $V(r, h) = \pi r^2 h + \frac{2}{3}\pi r^3$ subject to $g(r, h) = 2\pi rh + \pi r^2 + (4\pi r^2)/2 = 2\pi rh + 3\pi r^2 = S$. Then $\nabla V = \lambda \nabla g \quad \Rightarrow$ $\left\langle 2\pi rh + 2\pi r^2, \pi r^2 \right\rangle = \langle 2\lambda\pi h + 6\lambda\pi r, 2\lambda\pi r \rangle$, so the three equations are $2\pi rh + 2\pi r^2 = 2\lambda\pi h + 6\lambda\pi r$, $\pi r^2 = 2\lambda\pi r$, and $2\pi rh + 3\pi r^2 = S$. The second equation implies $r = 2\lambda$ $[r \neq 0]$. Substituting $r = 2\lambda$ into the first equation gives $2\pi(2\lambda)h + 2\pi(2\lambda)^2 = 2\lambda\pi h + 6\lambda\pi(2\lambda) \quad \Rightarrow \quad 4\pi\lambda h + 8\pi\lambda^2 = 2\lambda\pi h + 12\pi\lambda^2 \quad \Rightarrow \quad 2\pi\lambda h = 4\pi\lambda^2 \quad \Rightarrow \quad h = 2\lambda$. Thus, $r = 2\lambda = h$, and the volume of the silo is maximized, subject to a given surface area, when the radius and height are equal.

56. Let the dimensions of the box be x, y, and z, so its volume is $f(x, y, z) = xyz$, its surface area is $2xy + 2yz + 2xz = 1500$ and its total edge length is $4x + 4y + 4z = 200$. We find the extreme values of $f(x, y, z)$ subject to the constraints $g(x, y, z) = xy + yz + xz = 750$ and $h(x, y, z) = x + y + z = 50$. Then $\nabla f = \langle yz, xz, xy \rangle = \lambda \nabla g + \mu \nabla h = \langle \lambda(y + z), \lambda(x + z), \lambda(x + y) \rangle + \langle \mu, \mu, \mu \rangle$. So $yz = \lambda(y + z) + \mu$ **(1)**, $xz = \lambda(x + z) + \mu$ **(2)**, and $xy = \lambda(x + y) + \mu$ **(3)**. Notice that the box can't be a cube or else $x = y = z = \frac{50}{3}$ but then $xy + yz + xz = \frac{2500}{3} \neq 750$. Assume x is the distinct side, that is, $x \neq y$, $x \neq z$. Then **(1)** minus **(2)** implies $z\,(y - x) = \lambda(y - x)$ or $\lambda = z$, and **(1)** minus **(3)** implies $y(z - x) = \lambda(z - x)$ or $\lambda = y$. So $y = z = \lambda$ and $x + y + z = 50$ implies $x = 50 - 2\lambda$; also $xy + yz + xz = 750$ implies $x(2\lambda) + \lambda^2 = 750$. Hence $50 - 2\lambda = \dfrac{750 - \lambda^2}{2\lambda}$ or $3\lambda^2 - 100\lambda + 750 = 0$ and $\lambda = \dfrac{50 \pm 5\sqrt{10}}{3}$, giving the points $\left(\frac{1}{3}\left(50 \mp 10\sqrt{10}\right), \frac{1}{3}\left(50 \pm 5\sqrt{10}\right), \frac{1}{3}\left(50 \pm 5\sqrt{10}\right)\right)$. Thus the minimum of f is $f\left(\frac{1}{3}\left(50 - 10\sqrt{3}\right), \frac{1}{3}\left(50 + 5\sqrt{10}\right), \frac{1}{3}\left(50 + 5\sqrt{10}\right)\right) = \frac{1}{27}\left(87{,}500 - 2500\sqrt{10}\right)$, and its maximum is $f\left(\frac{1}{3}\left(50 + 10\sqrt{10}\right), \frac{1}{3}\left(50 - 5\sqrt{10}\right), \frac{1}{3}\left(50 - 5\sqrt{10}\right)\right) = \frac{1}{27}\left(87{,}500 + 2500\sqrt{10}\right)$.

Note: If either y or z is the distinct side, then symmetry gives the same result.

57. We need to find the extreme values of $f(x, y, z) = x^2 + y^2 + z^2$ subject to the two constraints $g(x, y, z) = x + y + 2z = 2$ and $h(x, y, z) = x^2 + y^2 - z = 0$. $\nabla f = \langle 2x, 2y, 2z \rangle$, $\lambda \nabla g = \langle \lambda, \lambda, 2\lambda \rangle$ and $\mu \nabla h = \langle 2\mu x, 2\mu y, -\mu \rangle$. Thus we need $2x = \lambda + 2\mu x$ **(1)**, $2y = \lambda + 2\mu y$ **(2)**, $2z = 2\lambda - \mu$ **(3)**, $x + y + 2z = 2$ **(4)**, and $x^2 + y^2 - z = 0$ **(5)**. From **(1)** and **(2)**, $2(x - y) = 2\mu(x - y)$, so if $x \neq y$, $\mu = 1$. Putting this in **(3)** gives $2z = 2\lambda - 1$ or $\lambda = z + \frac{1}{2}$, but putting $\mu = 1$ into **(1)** says $\lambda = 0$. Hence $z + \frac{1}{2} = 0$ or $z = -\frac{1}{2}$. Then **(4)** and **(5)** become $x + y - 3 = 0$ and $x^2 + y^2 + \frac{1}{2} = 0$. The

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

last equation cannot be true, so this case gives no solution. So we must have $x = y$. Then **(4)** and **(5)** become $2x + 2z = 2$ and $2x^2 - z = 0$ which imply $z = 1 - x$ and $z = 2x^2$. Thus $2x^2 = 1 - x$ or $2x^2 + x - 1 = (2x - 1)(x + 1) = 0$ so $x = \frac{1}{2}$ or $x = -1$. The two points to check are $\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$ and $(-1, -1, 2)$: $f\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right) = \frac{3}{4}$ and $f(-1, -1, 2) = 6$. Thus $\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$ is the point on the ellipse nearest the origin and $(-1, -1, 2)$ is the one farthest from the origin.

58. (a) After plotting $z = \sqrt{x^2 + y^2}$, the top half of the cone, and the plane $z = (5 - 4x + 3y)/8$ we see the ellipse formed by the intersection of the surfaces. The ellipse can be plotted explicitly using cylindrical coordinates (see Section 15.7): The cone is given by $z = r$, and the plane is $4r\cos\theta - 3r\sin\theta + 8z = 5$. Substituting $z = r$ into the plane equation gives $4r\cos\theta - 3r\sin\theta + 8r = 5 \quad \Rightarrow \quad r = \dfrac{5}{4\cos\theta - 3\sin\theta + 8}$.

Since $z = r$ on the ellipse, parametric equations (in cylindrical coordinates) are $\theta = t,\ r = z = \dfrac{5}{4\cos t - 3\sin t + 8},\ 0 \le t \le 2\pi$.

(b) We need to find the extreme values of $f(x, y, z) = z$ subject to the two constraints $g(x, y, z) = 4x - 3y + 8z = 5$ and $h(x, y, z) = x^2 + y^2 - z^2 = 0$.

$\nabla f = \lambda\nabla g + \mu\nabla h \quad \Rightarrow \quad \langle 0, 0, 1\rangle = \lambda\langle 4, -3, 8\rangle + \mu\langle 2x, 2y, -2z\rangle$, so we need $4\lambda + 2\mu x = 0 \quad \Rightarrow \quad x = -\frac{2\lambda}{\mu}$ **(1)**, $-3\lambda + 2\mu y = 0 \quad \Rightarrow \quad y = \frac{3\lambda}{2\mu}$ **(2)**, $8\lambda - 2\mu z = 1 \quad \Rightarrow \quad z = \frac{8\lambda - 1}{2\mu}$ **(3)**, $4x - 3y + 8z = 5$ **(4)**, and $x^2 + y^2 = z^2$ **(5)**. [Note that $\mu \neq 0$, else $\lambda = 0$ from **(1)**, but substitution into **(3)** gives a contradiction.]

Substituting **(1)**, **(2)**, and **(3)** into **(4)** gives $4\left(-\frac{2\lambda}{\mu}\right) - 3\left(\frac{3\lambda}{2\mu}\right) + 8\left(\frac{8\lambda - 1}{2\mu}\right) = 5 \quad \Rightarrow \quad \mu = \frac{39\lambda - 8}{10}$ and into **(5)** gives

$\left(-\frac{2\lambda}{\mu}\right)^2 + \left(\frac{3\lambda}{2\mu}\right)^2 = \left(\frac{8\lambda - 1}{2\mu}\right)^2 \quad \Rightarrow \quad 16\lambda^2 + 9\lambda^2 = (8\lambda - 1)^2 \quad \Rightarrow \quad 39\lambda^2 - 16\lambda + 1 = 0 \quad \Rightarrow \quad \lambda = \frac{1}{13}$ or $\lambda = \frac{1}{3}$.

If $\lambda = \frac{1}{13}$ then $\mu = -\frac{1}{2}$ and $x = \frac{4}{13}$, $y = -\frac{3}{13}$, $z = \frac{5}{13}$. If $\lambda = \frac{1}{3}$ then $\mu = \frac{1}{2}$ and $x = -\frac{4}{3}$, $y = 1$, $z = \frac{5}{3}$. Thus the highest point on the ellipse is $\left(-\frac{4}{3}, 1, \frac{5}{3}\right)$ and the lowest point is $\left(\frac{4}{13}, -\frac{3}{13}, \frac{5}{13}\right)$.

59. $f(x, y, z) = ye^{x-z},\ g(x, y, z) = 9x^2 + 4y^2 + 36z^2 = 36,\ h(x, y, z) = xy + yz = 1$.

$\nabla f = \lambda\nabla g + \mu\nabla h \quad \Rightarrow \quad \left\langle ye^{x-z}, e^{x-z}, -ye^{x-z}\right\rangle = \lambda\langle 18x, 8y, 72z\rangle + \mu\langle y, x + z, y\rangle$, so $ye^{x-z} = 18\lambda x + \mu y$, $e^{x-z} = 8\lambda y + \mu(x + z)$, $-ye^{x-z} = 72\lambda z + \mu y$, $9x^2 + 4y^2 + 36z^2 = 36$, $xy + yz = 1$. Using a CAS to solve these 5 equations simultaneously for x, y, z, λ, and μ (in Maple, use the `allvalues` command), we get 4 real-valued solutions:

$$\begin{array}{lllll}
x \approx 0.222444, & y \approx -2.157012, & z \approx -0.686049, & \lambda \approx -0.200401, & \mu \approx 2.108584 \\
x \approx -1.951921, & y \approx -0.545867, & z \approx 0.119973, & \lambda \approx 0.003141, & \mu \approx -0.076238 \\
x \approx 0.155142, & y \approx 0.904622, & z \approx 0.950293, & \lambda \approx -0.012447, & \mu \approx 0.489938 \\
x \approx 1.138731, & y \approx 1.768057, & z \approx -0.573138, & \lambda \approx 0.317141, & \mu \approx 1.862675
\end{array}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Substituting these values into f gives $f(0.222444, -2.157012, -0.686049) \approx -5.3506$, $f(-1.951921, -0.545867, 0.119973) \approx -0.0688$, $f(0.155142, 0.904622, 0.950293) \approx 0.4084$, $f(1.138731, 1.768057, -0.573138) \approx 9.7938$. Thus the maximum is approximately 9.7938, and the minimum is approximately -5.3506.

60. $f(x, y, z) = x + y + z,\ g(x, y, z) = x^2 - y^2 - z = 0,\ h(x, y, z) = x^2 + z^2 = 4.$

$\nabla f = \lambda \nabla g + \mu \nabla h \quad \Rightarrow \quad \langle 1, 1, 1\rangle = \lambda\langle 2x, -2y, -1\rangle + \mu\langle 2x, 0, 2z\rangle$, so $1 = 2\lambda x + 2\mu x$, $1 = -2\lambda y$, $1 = -\lambda + 2\mu z$, $x^2 - y^2 = z$, $x^2 + z^2 = 4$. Using a CAS to solve these 5 equations simultaneously for x, y, z, λ, and μ, we get 4 real-valued solutions:

$$\begin{array}{lllll} x \approx -1.652878, & y \approx -1.964194, & z \approx -1.126052, & \lambda \approx 0.254557, & \mu \approx -0.557060 \\ x \approx -1.502800, & y \approx 0.968872, & z \approx 1.319694, & \lambda \approx -0.516064, & \mu \approx 0.183352 \\ x \approx -0.992513, & y \approx 1.649677, & z \approx -1.736352, & \lambda \approx -0.303090, & \mu \approx -0.200682 \\ x \approx 1.895178, & y \approx 1.718347, & z \approx 0.638984, & \lambda \approx -0.290977, & \mu \approx 0.554805 \end{array}$$

Substituting these values into f gives $f(-1.652878, -1.964194, -1.126052) \approx -4.7431$, $f(-1.502800, 0.968872, 1.319694) \approx 0.7858$, $f(-0.992513, 1.649677, -1.736352) \approx -1.0792$, $f(1.895178, 1.718347, 0.638984) \approx 4.2525$. Thus the maximum is approximately 4.2525, and the minimum is approximately -4.7431.

61. $f(x, y) = 3x^2 + y^2$, $g(x, y) = x^2 + y^2 - 4y = 0$. Then $\nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle 6x, 2y\rangle = \langle 2\lambda x, \lambda(2y - 4)\rangle$, so the three equations are $6x = 2\lambda x$, $2y = \lambda(2y - 4)$, and $x^2 + y^2 - 4y = 0$. The first equation implies $x = 0$ or $\lambda = 3$. If $x = 0$, the third equation implies $y = 0$ or $y = 4$. If $\lambda = 3$, the second equation implies $y = 3$ and substitution into the third equation gives $x = \pm\sqrt{3}$. Thus, f has possible extreme values at $(0, 0)$, $(0, 4)$, $\left(\pm\sqrt{3}, 3\right)$. Evaluating f at these points we see that the maximum value is $f\left(\pm\sqrt{3}, 3\right) = 18$ and the minimum value is $f(0, 0) = 0$.

The minimum value of f occurs at $(0, 0)$. Substituting $y = 0$ into the second equation gives $2(0) = \lambda(2(0) - 4)$, which is true only if $\lambda = 0$. Thus, the minimum value corresponds to $\lambda = 0$.

62. (a) Let $f(x_1, \ldots, x_n, y_1, \ldots, y_n) = \sum\limits_{i=1}^{n} x_i y_i$, $g(x_1, \ldots, x_n) = \sum\limits_{i=1}^{n} x_i^2$, and $h(x_1, \ldots, x_n) = \sum\limits_{i=1}^{n} y_i^2$. Then

$\nabla f = \nabla \sum\limits_{i=1}^{n} x_i y_i = \langle y_1, y_2, \ldots, y_n, x_1, x_2, \ldots, x_n\rangle$, $\nabla g = \nabla \sum\limits_{i=1}^{n} x_i^2 = \langle 2x_1, 2x_2, \ldots, 2x_n, 0, 0, \ldots, 0\rangle$ and

$\nabla h = \nabla \sum\limits_{i=1}^{n} y_i^2 = \langle 0, 0, \ldots, 0, 2y_1, 2y_2, \ldots, 2y_n\rangle$. So $\nabla f = \lambda \nabla g + \mu \nabla h \quad \Leftrightarrow \quad y_i = 2\lambda x_i$ and $x_i = 2\mu y_i$,

$1 \le i \le n$. Then $1 = \sum\limits_{i=1}^{n} y_i^2 = \sum\limits_{i=1}^{n} 4\lambda^2 x_i^2 = 4\lambda^2 \sum\limits_{i=1}^{n} x_i^2 = 4\lambda^2 \quad \Rightarrow \quad \lambda = \pm\frac{1}{2}$. If $\lambda = \frac{1}{2}$ then $y_i = 2\left(\frac{1}{2}\right)x_i = x_i$,

$1 \le i \le n$. Thus $\sum\limits_{i=1}^{n} x_i y_i = \sum\limits_{i=1}^{n} x_i^2 = 1$. Similarly if $\lambda = -\frac{1}{2}$ we get $y_i = -x_i$ and $\sum\limits_{i=1}^{n} x_i y_i = -1$. Similarly we get

$\mu = \pm\frac{1}{2}$ giving $y_i = \pm x_i$, $1 \le i \le n$, and $\sum\limits_{i=1}^{n} x_i y_i = \pm 1$. Thus the maximum value of $\sum\limits_{i=1}^{n} x_i y_i$ is 1.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Here we assume $\sum_{i=1}^{n} a_i^2 \neq 0$ and $\sum_{i=1}^{n} b_i^2 \neq 0$. (If $\sum_{i=1}^{n} a_i^2 = 0$, then each $a_i = 0$ and so the inequality is trivially true.)

$x_i = \dfrac{a_i}{\sqrt{\sum a_j^2}} \quad \Rightarrow \quad \sum x_i^2 = \dfrac{\sum a_i^2}{\sum a_j^2} = 1$, and $y_i = \dfrac{b_i}{\sqrt{\sum b_j^2}} \quad \Rightarrow \quad \sum y_i^2 = \dfrac{\sum b_i^2}{\sum b_j^2} = 1$. Therefore, from part (a),

$\sum x_i y_i = \sum \dfrac{a_i b_i}{\sqrt{\sum a_j^2}\sqrt{\sum b_j^2}} \leq 1 \quad \Leftrightarrow \quad \sum a_i b_i \leq \sqrt{\sum a_j^2}\sqrt{\sum b_j^2}$.

63. (a) We wish to maximize $f(x_1, x_2, \ldots, x_n) = \sqrt[n]{x_1 x_2 \cdots x_n}$ subject to

$g(x_1, x_2, \ldots, x_n) = x_1 + x_2 + \cdots + x_n = c$ and $x_i > 0$.

$\nabla f = \left\langle \frac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_2 \cdots x_n), \frac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_1 x_3 \cdots x_n), \ldots, \frac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_1 \cdots x_{n-1}) \right\rangle$

and $\lambda \nabla g = \langle \lambda, \lambda, \ldots, \lambda \rangle$, so we need to solve the system of equations

$$\begin{aligned}
\tfrac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_2 \cdots x_n) = \lambda \quad &\Rightarrow \quad x_1^{1/n} x_2^{1/n} \cdots x_n^{1/n} = n\lambda x_1 \\
\tfrac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_1 x_3 \cdots x_n) = \lambda \quad &\Rightarrow \quad x_1^{1/n} x_2^{1/n} \cdots x_n^{1/n} = n\lambda x_2 \\
&\vdots \\
\tfrac{1}{n}(x_1 x_2 \cdots x_n)^{\frac{1}{n}-1}(x_1 \cdots x_{n-1}) = \lambda \quad &\Rightarrow \quad x_1^{1/n} x_2^{1/n} \cdots x_n^{1/n} = n\lambda x_n
\end{aligned}$$

This implies $n\lambda x_1 = n\lambda x_2 = \cdots = n\lambda x_n$. Note $\lambda \neq 0$, otherwise we can't have all $x_i > 0$. Thus $x_1 = x_2 = \cdots = x_n$. But $x_1 + x_2 + \cdots + x_n = c \quad \Rightarrow \quad nx_1 = c \quad \Rightarrow \quad x_1 = \dfrac{c}{n} = x_2 = x_3 = \cdots = x_n$. Then the only point where f can have an extreme value is $\left(\dfrac{c}{n}, \dfrac{c}{n}, \ldots, \dfrac{c}{n}\right)$. Since we can choose values for $(x_1, x_2, \ldots, x_n)$ that make f as close to zero (but not equal) as we like, f has no minimum value. Thus the maximum value is

$f\left(\dfrac{c}{n}, \dfrac{c}{n}, \ldots, \dfrac{c}{n}\right) = \sqrt[n]{\dfrac{c}{n} \cdot \dfrac{c}{n} \cdot \cdots \cdot \dfrac{c}{n}} = \dfrac{c}{n}$.

(b) From part (a), $\dfrac{c}{n}$ is the maximum value of f. Thus $f(x_1, x_2, \ldots, x_n) = \sqrt[n]{x_1 x_2 \cdots x_n} \leq \dfrac{c}{n}$. But $x_1 + x_2 + \cdots + x_n = c$, so $\sqrt[n]{x_1 x_2 \cdots x_n} \leq \dfrac{x_1 + x_2 + \cdots + x_n}{n}$. These two means are equal when f attains its maximum value $\dfrac{c}{n}$, but this can occur only at the point $\left(\dfrac{c}{n}, \dfrac{c}{n}, \ldots, \dfrac{c}{n}\right)$ we found in part (a). So the means are equal only when $x_1 = x_2 = x_3 = \cdots = x_n = \dfrac{c}{n}$.

APPLIED PROJECT Rocket Science

1. Initially the rocket engine has mass $M_r = M_1$ and payload mass $P = M_2 + M_3 + A$. Then the change in velocity resulting from the first stage is $\Delta V_1 = -c \ln\left(1 - \dfrac{(1-S)M_1}{M_2 + M_3 + A + M_1}\right)$. After the first stage is jettisoned we can consider the rocket engine to have mass $M_r = M_2$ and the payload to have mass $P = M_3 + A$. The resulting change in velocity from the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

second stage is $\Delta V_2 = -c\ln\left(1 - \dfrac{(1-S)M_2}{M_3 + A + M_2}\right)$. When only the third stage remains, we have $M_r = M_3$ and $P = A$, so the resulting change in velocity is $\Delta V_3 = -c\ln\left(1 - \dfrac{(1-S)M_3}{A + M_3}\right)$. Since the rocket started from rest, the final velocity attained is

$$\begin{aligned}
v_f &= \Delta V_1 + \Delta V_2 + \Delta V_3 \\
&= -c\ln\left(1 - \frac{(1-S)M_1}{M_2 + M_3 + A + M_1}\right) + (-c)\ln\left(1 - \frac{(1-S)M_2}{M_3 + A + M_2}\right) + (-c)\ln\left(1 - \frac{(1-S)M_3}{A + M_3}\right) \\
&= -c\left[\ln\left(\frac{M_1 + M_2 + M_3 + A - (1-S)M_1}{M_1 + M_2 + M_3 + A}\right) + \ln\left(\frac{M_2 + M_3 + A - (1-S)M_2}{M_2 + M_3 + A}\right)\right. \\
&\qquad\qquad \left. + \ln\left(\frac{M_3 + A - (1-S)M_3}{M_3 + A}\right)\right] \\
&= c\left[\ln\left(\frac{M_1 + M_2 + M_3 + A}{SM_1 + M_2 + M_3 + A}\right) + \ln\left(\frac{M_2 + M_3 + A}{SM_2 + M_3 + A}\right) + \ln\left(\frac{M_3 + A}{SM_3 + A}\right)\right]
\end{aligned}$$

2. Define $N_1 = \dfrac{M_1 + M_2 + M_3 + A}{SM_1 + M_2 + M_3 + A}$, $N_2 = \dfrac{M_2 + M_3 + A}{SM_2 + M_3 + A}$, and $N_3 = \dfrac{M_3 + A}{SM_3 + A}$. Then

$$\begin{aligned}
\frac{(1-S)N_1}{1 - SN_1} &= \frac{(1-S)\dfrac{M_1 + M_2 + M_3 + A}{SM_1 + M_2 + M_3 + A}}{1 - S\dfrac{M_1 + M_2 + M_3 + A}{SM_1 + M_2 + M_3 + A}} = \frac{(1-S)(M_1 + M_2 + M_3 + A)}{SM_1 + M_2 + M_3 + A - S(M_1 + M_2 + M_3 + A)} \\
&= \frac{(1-S)(M_1 + M_2 + M_3 + A)}{(1-S)(M_2 + M_3 + A)} = \frac{M_1 + M_2 + M_3 + A}{M_2 + M_3 + A}
\end{aligned}$$

as desired.

Similarly,

$$\frac{(1-S)N_2}{1 - SN_2} = \frac{(1-S)(M_2 + M_3 + A)}{SM_2 + M_3 + A - S(M_2 + M_3 + A)} = \frac{(1-S)(M_2 + M_3 + A)}{(1-S)(M_3 + A)} = \frac{M_2 + M_3 + A}{M_3 + A}$$

and
$$\frac{(1-S)N_3}{1 - SN_3} = \frac{(1-S)(M_3 + A)}{SM_3 + A - S(M_3 + A)} = \frac{(1-S)(M_3 + A)}{(1-S)(A)} = \frac{M_3 + A}{A}$$

Then

$$\begin{aligned}
\frac{M + A}{A} &= \frac{M_1 + M_2 + M_3 + A}{A} = \frac{M_1 + M_2 + M_3 + A}{M_2 + M_3 + A} \cdot \frac{M_2 + M_3 + A}{M_3 + A} \cdot \frac{M_3 + A}{A} \\
&= \frac{(1-S)N_1}{1 - SN_1} \cdot \frac{(1-S)N_2}{1 - SN_2} \cdot \frac{(1-S)N_3}{1 - SN_3} = \frac{(1-S)^3 N_1 N_2 N_3}{(1 - SN_1)(1 - SN_2)(1 - SN_3)}
\end{aligned}$$

3. Since $A > 0$, $M + A$ and consequently $\dfrac{M + A}{A}$ is minimized for the same values as M. $\ln x$ is a strictly increasing function, so $\ln\left(\dfrac{M + A}{A}\right)$ must give a minimum for the same values as $\dfrac{M + A}{A}$ and hence M. We then wish to minimize $\ln\left(\dfrac{M + A}{A}\right)$ subject to the constraint $c\,(\ln N_1 + \ln N_2 + \ln N_3) = v_f$. From Problem 2,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\ln\left(\frac{M+A}{A}\right) = \ln\left(\frac{(1-S)^3 N_1 N_2 N_3}{(1-SN_1)(1-SN_2)(1-SN_3)}\right)$$
$$= 3\ln(1-S) + \ln N_1 + \ln N_2 + \ln N_3 - \ln(1-SN_1) - \ln(1-SN_2) - \ln(1-SN_3)$$

Using the method of Lagrange multipliers, we need to solve $\nabla\left[\ln\left(\frac{M+A}{A}\right)\right] = \lambda\nabla[c(\ln N_1 + \ln N_2 + \ln N_3)]$ with $c(\ln N_1 + \ln N_2 + \ln N_3) = v_f$ in terms of N_1, N_2, and N_3. The resulting system is

$$\frac{1}{N_1} + \frac{S}{1-SN_1} = \lambda\frac{c}{N_1} \qquad \frac{1}{N_2} + \frac{S}{1-SN_2} = \lambda\frac{c}{N_2} \qquad \frac{1}{N_3} + \frac{S}{1-SN_3} = \lambda\frac{c}{N_3}$$

$$c(\ln N_1 + \ln N_2 + \ln N_3) = v_f$$

One approach to solving the system is isolating $c\lambda$ in the first three equations which gives

$1 + \frac{SN_1}{1-SN_1} = c\lambda = 1 + \frac{SN_2}{1-SN_2} = 1 + \frac{SN_3}{1-SN_3} \quad\Rightarrow\quad \frac{N_1}{1-SN_1} = \frac{N_2}{1-SN_2} = \frac{N_3}{1-SN_3} \quad\Rightarrow$

$N_1 = N_2 = N_3$ (Verify!). This says the fourth equation can be expressed as $c(\ln N_1 + \ln N_1 + \ln N_1) = v_f \quad\Rightarrow$

$3c\ln N_1 = v_f \quad\Rightarrow\quad \ln N_1 = \frac{v_f}{3c}$. Thus the minimum mass M of the rocket engine is attained for

$N_1 = N_2 = N_3 = e^{v_f/(3c)}$.

4. Using the previous results, $\frac{M+A}{A} = \frac{(1-S)^3 N_1 N_2 N_3}{(1-SN_1)(1-SN_2)(1-SN_3)} = \frac{(1-S)^3\left[e^{v_f/(3c)}\right]^3}{\left[1-Se^{v_f/(3c)}\right]^3} = \frac{(1-S)^3 e^{v_f/c}}{\left[1-Se^{v_f/(3c)}\right]^3}$.

Then $M = \frac{A(1-S)^3 e^{v_f/c}}{\left[1-Se^{v_f/(3c)}\right]^3} - A$.

5. (a) From Problem 4, $M = \frac{A(1-0.2)^3 e^{(17{,}500/6000)}}{\left(1-0.2e^{[17{,}500/(3\cdot 6000)]}\right)^3} - A \approx 90.4A - A = 89.4A$.

(b) First, $N_3 = \frac{M_3+A}{SM_3+A} \quad\Rightarrow\quad e^{[17{,}500/(3\cdot 6000)]} = \frac{M_3+A}{0.2M_3+A} \quad\Rightarrow\quad M_3 = \frac{A(1-e^{35/36})}{0.2e^{35/36}-1} \approx 3.49A$.

Then $N_2 = \frac{M_2+M_3+A}{SM_2+M_3+A} = \frac{M_2+3.49A+A}{0.2M_2+3.49A+A} \quad\Rightarrow\quad M_2 = \frac{4.49A(1-e^{35/36})}{0.2e^{35/36}-1} \approx 15.67A$ and

$N_3 = \frac{M_1+M_2+M_3+A}{SM_1+M_2+M_3+A} = \frac{M_1+15.67A+3.49A+A}{0.2M_1+15.67A+3.49A+A} \quad\Rightarrow\quad M_1 = \frac{20.16A(1-e^{35/36})}{0.2e^{35/36}-1} \approx 70.36A$.

6. As in Problem 5, $N_3 = \frac{M_3+A}{SM_3+A} \quad\Rightarrow\quad e^{24{,}700/(3\cdot 6000)} = \frac{M_3+A}{0.2M_3+A} \quad\Rightarrow\quad M_3 = \frac{A(1-e^{247/180})}{0.2e^{247/180}-1} \approx 13.9A$,

$N_2 = \frac{M_2+M_3+A}{SM_2+M_3+A} = \frac{M_2+13.9A+A}{0.2M_2+13.9A+A} \quad\Rightarrow\quad M_2 = \frac{14.9A(1-e^{247/180})}{0.2e^{247/180}-1} \approx 208A$, and

$N_3 = \frac{M_1+M_2+M_3+A}{SM_1+M_2+M_3+A} = \frac{M_1+208A+13.9A+A}{0.2M_1+208A+13.9A+A} \quad\Rightarrow\quad M_1 = \frac{222.9A(1-e^{247/180})}{0.2e^{247/180}-1} \approx 3110A$.

Here $A = 500$, so the mass of each stage of the rocket engine is approximately $M_1 = 3110(500) = 1{,}550{,}000$ lb, $M_2 = 208(500) = 104{,}000$ lb, and $M_3 = 13.9(500) = 6950$ lb.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

APPLIED PROJECT Hydro-Turbine Optimization

1. We wish to maximize the total energy production for a given total flow, so we can say Q_T is fixed and we want to maximize $KW_1 + KW_2 + KW_3$. Notice each KW_i has a constant factor $\left(170 - 1.6 \cdot 10^{-6} Q_T^2\right)$, so to simplify the computations we can equivalently maximize

$$\begin{aligned} f(Q_1, Q_2, Q_3) &= \frac{KW_1 + KW_2 + KW_3}{170 - 1.6 \cdot 10^{-6} Q_T^2} \\ &= \left(-18.89 + 0.1277 Q_1 - 4.08 \cdot 10^{-5} Q_1^2\right) + \left(-24.51 + 0.1358 Q_2 - 4.69 \cdot 10^{-5} Q_2^2\right) \\ &\qquad + \left(-27.02 + 0.1380 Q_3 - 3.84 \cdot 10^{-5} Q_3^2\right) \end{aligned}$$

subject to the constraint $g(Q_1, Q_2, Q_3) = Q_1 + Q_2 + Q_3 = Q_T$. So first we find the values of Q_1, Q_2, Q_3 where $\nabla f(Q_1, Q_2, Q_3) = \lambda \nabla g(Q_1, Q_2, Q_3)$ and $Q_1 + Q_2 + Q_3 = Q_T$ which is equivalent to solving the system

$$\begin{aligned} 0.1277 - 2(4.08 \cdot 10^{-5}) Q_1 &= \lambda \\ 0.1358 - 2(4.69 \cdot 10^{-5}) Q_2 &= \lambda \\ 0.1380 - 2(3.84 \cdot 10^{-5}) Q_3 &= \lambda \\ Q_1 + Q_2 + Q_3 &= Q_T \end{aligned}$$

Comparing the first and third equations, we have $0.1277 - 2(4.08 \cdot 10^{-5}) Q_1 = 0.1380 - 2(3.84 \cdot 10^{-5}) Q_3 \quad \Rightarrow$ $Q_1 = -126.2255 + 0.9412 Q_3$. From the second and third equations, $0.1358 - 2(4.69 \cdot 10^{-5}) Q_2 = 0.1380 - 2(3.84 \cdot 10^{-5}) Q_3 \quad \Rightarrow \quad Q_2 = -23.4542 + 0.8188 Q_3$. Substituting into $Q_1 + Q_2 + Q_3 = Q_T$ gives $(-126.2255 + 0.9412 Q_3) + (-23.4542 + 0.8188 Q_3) + Q_3 = Q_T \quad \Rightarrow$ $2.76 Q_3 = Q_T + 149.6797 \quad \Rightarrow \quad Q_3 = 0.3623 Q_T + 54.23$. Then $Q_1 = -126.2255 + 0.9412 Q_3 = -126.2255 + 0.9412(0.3623 Q_T + 54.23) = 0.3410 Q_T - 75.18$ and $Q_2 = -23.4542 + 0.8188(0.3623 Q_T + 54.23) = 0.2967 Q_T + 20.95$. As long as we maintain $250 \le Q_1 \le 1110$, $250 \le Q_2 \le 1110$, and $250 \le Q_3 \le 1225$, we can reason from the nature of the functions KW_i that these values give a maximum of f, and hence a maximum energy production, and not a minimum.

2. From Problem 1, the value of Q_1 that maximizes energy production is $0.3410 Q_T - 75.18$, but since $250 \le Q_1 \le 1110$, we must have $250 \le 0.3410 Q_T - 75.18 \le 1110 \quad \Rightarrow \quad 325.18 \le 0.3410 Q_T \le 1185.18 \quad \Rightarrow \quad 953.6 \le Q_T \le 3475.6$. Similarly, $250 \le Q_2 \le 1110 \quad \Rightarrow \quad 250 \le 0.2967 Q_T + 20.95 \le 1110 \quad \Rightarrow \quad 772.0 \le Q_T \le 3670.5$, and $250 \le Q_3 \le 1225 \quad \Rightarrow \quad 250 \le 0.3623 Q_T + 54.23 \le 1225 \quad \Rightarrow \quad 540.4 \le Q_T \le 3231.5$. Consolidating these results, we see that the values from Problem 1 are applicable only for $953.6 \le Q_T \le 3231.5$.

3. If $Q_T = 2500$, the results from Problem 1 show that the maximum energy production occurs for

$$\begin{aligned} Q_1 &= 0.3410 Q_T - 75.18 = 0.3410(2500) - 75.18 = 777.3 \\ Q_2 &= 0.2967 Q_T + 20.95 = 0.2967(2500) + 20.95 = 762.7 \\ Q_3 &= 0.3623 Q_T + 54.23 = 0.3623(2500) + 54.23 = 960.0 \end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The energy produced for these values is $KW_1 + KW_2 + KW_3 \approx 8915.2 + 8285.1 + 11{,}211.3 \approx 28{,}411.6$. We compute the energy production for a nearby distribution, $Q_1 = 770$, $Q_2 = 760$, and $Q_3 = 970$: $KW_1 + KW_2 + KW_3 \approx 8839.8 + 8257.4 + 11{,}313.5 = 28{,}410.7$. For another example, we take $Q_1 = 780$, $Q_2 = 765$, and $Q_3 = 955$: $KW_1 + KW_2 + KW_3 \approx 8942.9 + 8308.8 + 11{,}159.7 = 28{,}411.4$. These distributions are both close to the distribution from Problem 1 and both give slightly lower energy productions, suggesting that $Q_1 = 777.3$, $Q_2 = 762.7$, and $Q_3 = 960.0$ is indeed the optimal distribution.

4. First we graph each power function in its domain if all of the flow is directed to that turbine (so $Q_i = Q_T$). If we use only one turbine, the graph indicates that for a water flow of 1000 ft^3/s, Turbine 3 produces the most power, approximately 12,200 kW. In comparison, if we use all three turbines, the results of Problem 1 with $Q_T = 1000$ give $Q_1 = 265.8$, $Q_2 = 317.7$, and $Q_3 = 416.5$, resulting in a total energy production of $KW_1 + KW_2 + KW_3 \approx 8397.4$ kW. Here, using only one turbine produces significantly more energy! If the flow is only 600 ft^3/s, we do not have the option of using all three turbines, as the domain restrictions require a minimum of 250 ft^3/s in each turbine. We can use just one turbine, then, and from the graph Turbine 1 produces the most energy for a water flow of 600 ft^3.

5. If we examine the graph from Problem 4, we see that for water flows above approximately 450 ft^3/s, Turbine 2 produces the least amount of power. Therefore it seems reasonable to assume that we should distribute the incoming flow of 1500 ft^3/s between Turbines 1 and 3. (This can be verified by computing the power produced with the other pairs of turbines for comparison.) So now we wish to maximize $KW_1 + KW_3$ subject to the constraint $Q_1 + Q_3 = Q_T$ where $Q_T = 1500$.

As in Problem 1, we can equivalently maximize

$$\begin{aligned} f(Q_1, Q_3) &= \frac{KW_1 + KW_3}{170 - 1.6 \cdot 10^{-6} Q_T^2} \\ &= \left(-18.89 + 0.1277Q_1 - 4.08 \cdot 10^{-5} Q_1^2\right) + \left(-27.02 + 0.1380Q_3 - 3.84 \cdot 10^{-5} Q_3^2\right) \end{aligned}$$

subject to the constraint $g(Q_1, Q_3) = Q_1 + Q_3 = Q_T$.

Then we solve $\nabla f(Q_1, Q_3) = \lambda \nabla g(Q_1, Q_3) \quad \Rightarrow \quad 0.1277 - 2\left(4.08 \cdot 10^{-5}\right) Q_1 = \lambda$ and $0.1380 - 2(3.84 \cdot 10^{-5})Q_3 = \lambda$, thus $0.1277 - 2(4.08 \cdot 10^{-5})Q_1 = 0.1380 - 2(3.84 \cdot 10^{-5})Q_3 \quad \Rightarrow$ $Q_1 = -126.2255 + 0.9412Q_3$. Substituting into $Q_1 + Q_3 = Q_T$ gives $-126.2255 + 0.9412Q_3 + Q_3 = 1500 \quad \Rightarrow$ $Q_3 \approx 837.7$, and then $Q_1 = Q_T - Q_3 \approx 1500 - 837.7 = 662.3$. So we should apportion approximately 662.3 ft^3/s to Turbine 1 and the remaining 837.7 ft^3/s to Turbine 3. The resulting energy production is $KW_1 + KW_3 \approx 7952.1 + 10{,}256.2 = 18{,}208.3$ kW. (We can verify that this is indeed a maximum energy production by checking nearby distributions.) In comparison, if we use all three turbines with $Q_T = 1500$ we get $Q_1 = 436.3$, $Q_2 = 466.0$,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and $Q_3 = 597.7$, resulting in a total energy production of $KW_1 + KW_2 + KW_3 \approx 16{,}538.7$ kW. Clearly, for this flow level it is beneficial to use only two turbines.

6. Note that an incoming flow of 3400 ft^3/s is not within the domain we established in Problem 2, so we cannot simply use our previous work to give the optimal distribution. We will need to use all three turbines, due to the capacity limitations of each individual turbine, but 3400 is less than the maximum combined capacity of 3445 ft^3/s, so we still must decide how to distribute the flows. From the graph in Problem 4, Turbine 3 produces the most power for the higher flows, so it seems reasonable to use Turbine 3 at its maximum capacity of 1225 and distribute the remaining 2175 ft^3/s flow between Turbines 1 and 2. We can again use the technique of Lagrange multipliers to determine the optimal distribution. Following the procedure we used in Problem 5, we wish to maximize $KW_1 + KW_2$ subject to the constraint $Q_1 + Q_2 = Q_T$ where $Q_T = 2175$. We can equivalently maximize

$$\begin{aligned} f(Q_1, Q_2) &= \frac{KW_1 + KW_2}{170 - 1.6 \cdot 10^{-6} Q_T^2} \\ &= \left(-18.89 + 0.1277Q_1 - 4.08 \cdot 10^{-5} Q_1^2\right) + \left(-24.51 + 0.1358Q_2 - 4.69 \cdot 10^{-5} Q_2^2\right) \end{aligned}$$

subject to the constraint $g(Q_1, Q_2) = Q_1 + Q_2 = Q_T$. Then we solve $\nabla f(Q_1, Q_2) = \lambda \nabla g(Q_1, Q_2) \quad \Rightarrow$ $0.1277 - 2(4.08 \cdot 10^{-5})Q_1 = \lambda$ and $0.1358 - 2(4.69 \cdot 10^{-5})Q_2 = \lambda$, thus $0.1277 - 2(4.08 \cdot 10^{-5})Q_1 = 0.1358 - 2\left(4.69 \cdot 10^{-5}\right)Q_2 \quad \Rightarrow \quad Q_1 = -99.2647 + 1.1495Q_2$. Substituting into $Q_1 + Q_2 = Q_T$ gives $-99.2647 + 1.1495Q_2 + Q_2 = 2175 \quad \Rightarrow \quad Q_2 \approx 1058.0$, and then $Q_1 \approx 1117.0$. This value for Q_1 is larger than the allowable maximum flow to Turbine 1, but the result indicates that the flow to Turbine 1 should be maximized. Thus we should recommend that the company apportion the maximum allowable flows to Turbines 1 and 3, 1110 and 1225 ft^3/s, and the remaining 1065 ft^3/s to Turbine 2. Checking nearby distributions within the domain verifies that we have indeed found the optimal distribution.

14 Review

TRUE-FALSE QUIZ

1. True. $f_y(a, b) = \lim\limits_{h \to 0} \dfrac{f(a, b+h) - f(a, b)}{h}$ from Equation 14.3.3. Let $h = y - b$. As $h \to 0$, $y \to b$. Then by substituting, we get $f_y(a, b) = \lim\limits_{y \to b} \dfrac{f(a, y) - f(a, b)}{y - b}$.

2. False. If there were such a function, then $f_{xy} = 2y$ and $f_{yx} = 1$. So $f_{xy} \neq f_{yx}$, which contradicts Clairaut's Theorem.

3. False. $f_{xy} = \dfrac{\partial^2 f}{\partial y \, \partial x}$.

4. True. From Equation 14.6.14 we get $D_{\mathbf{k}} f(x, y, z) = \nabla f(x, y, z) \cdot \langle 0, 0, 1 \rangle = f_z(x, y, z)$.

5. False. See Example 14.2.3.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. False. See Exercise 14.4.54(a).

7. True. If f has a local minimum and f is differentiable at (a, b) then by Theorem 14.7.2, $f_x(a, b) = 0$ and $f_y(a, b) = 0$, so $\nabla f(a, b) = \langle f_x(a, b), f_y(a, b)\rangle = \langle 0, 0\rangle = \mathbf{0}$.

8. False. If f is not continuous at $(2, 5)$, then we can have $\lim\limits_{(x,y)\to(2,5)} f(x, y) \neq f(2, 5)$. (See Example 14.2.8.)

9. False. $\nabla f(x, y) = \langle 0, 1/y\rangle$.

10. True. This is equivalent to part (c) of the Second Derivatives Test (14.7.3).

11. True. $\nabla f = \langle \cos x, \cos y\rangle$, so $|\nabla f| = \sqrt{\cos^2 x + \cos^2 y}$. But $|\cos\theta| \leq 1$, so $|\nabla f| \leq \sqrt{2}$. Now $D_{\mathbf{u}} f(x, y) = \nabla f \cdot \mathbf{u} = |\nabla f|\,|\mathbf{u}| \cos\theta$, but $\mathbf{u}$ is a unit vector, so $|D_{\mathbf{u}} f(x, y)| \leq \sqrt{2} \cdot 1 \cdot 1 = \sqrt{2}$.

12. False. See Exercise 14.7.41.

EXERCISES

1. $f(x, y) = \ln(x + y + 1)$ is defined only when $x + y + 1 > 0 \quad \Leftrightarrow \quad y > -x - 1$, so the domain of f is $\{(x, y) \mid y > -x - 1\}$, all those points above the line $y = -x - 1$.

2. $f(x, y) = \sqrt{4 - x^2 - y^2} + \sqrt{1 - x^2}$.
$\sqrt{4 - x^2 - y^2}$ is defined only when $4 - x^2 - y^2 \geq 0 \quad \Leftrightarrow \quad x^2 + y^2 \leq 4$, and $\sqrt{1 - x^2}$ is defined only when $1 - x^2 \geq 0 \quad \Leftrightarrow \quad -1 \leq x \leq 1$, so the domain of f is $\left\{(x, y) \mid -1 \leq x \leq 1, -\sqrt{4 - x^2} \leq y \leq \sqrt{4 - x^2}\right\}$, which consists of those points on or inside the circle $x^2 + y^2 = 4$ for $-1 \leq x \leq 1$.

3. $z = f(x, y) = 1 - y^2$, a parabolic cylinder

4. $z = f(x, y) = x^2 + (y - 2)^2$, a circular paraboloid with vertex $(0, 2, 0)$ and axis parallel to the z-axis

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. The level curves are $\sqrt{4x^2+y^2}=k$ or $4x^2+y^2=k^2$, $k \geq 0$, a family of ellipses.

6. The level curves are $e^x+y=k$ or $y=-e^x+k$, a family of exponential curves.

7.

8. (a) The point $(3,2)$ lies partway between the level curves with z-values 50 and 60, and it appears that $(3,2)$ is about the same distance from either level curve. So we estimate that $f(3,2) \approx 55$.

(b) At the point $(3,2)$, if we fix y at $y=2$ and allow x to vary, the level curves indicate that the z-values decrease as x increases, so $f_x(3,2)$ is negative. In other words, if we start at $(3,2)$ and move right (in the positive x-direction), the contours show that our path along the surface $z=f(x,y)$ is descending.

(c) Both $f_y(2,1)$ and $f_y(2,2)$ are positive, because if we start from either point and move in the positive y-direction, the contour map indicates that the path is ascending. But the level curves are closer together in the y-direction at $(2,1)$ than at $(2,2)$, so the path is steeper (the z-values increase more rapidly) at $(2,1)$ and hence $f_y(2,1) > f_y(2,2)$.

9. f is a rational function, so it is continuous on its domain. Since f is defined at $(1,1)$, we use direct substitution to evaluate the limit: $\displaystyle\lim_{(x,y)\to(1,1)} \frac{2xy}{x^2+2y^2} = \frac{2(1)(1)}{1^2+2(1)^2} = \frac{2}{3}$.

10. $f(x,y)=\dfrac{2xy}{x^2+2y^2}$. As $(x,y)\to(0,0)$ along the x-axis, $f(x,0)=0/x^2=0$ for $x\neq 0$, so $f(x,y)\to 0$ along this line. As $(x,y)\to(0,0)$ along the line $x=y$, $f(x,x)=2x^2/(3x^2)=\frac{2}{3}$, so $f(x,y)\to\frac{2}{3}$. Thus, the limit doesn't exist.

11. (a) $T_x(6,4)=\displaystyle\lim_{h\to 0}\frac{T(6+h,4)-T(6,4)}{h}$, so we can approximate $T_x(6,4)$ by considering $h=\pm 2$ and using the values given in the table: $T_x(6,4)\approx \dfrac{T(8,4)-T(6,4)}{2}=\dfrac{86-80}{2}=3$,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$T_x(6,4) \approx \dfrac{T(4,4)-T(6,4)}{-2} = \dfrac{72-80}{-2} = 4$. Averaging these values, we estimate $T_x(6,4)$ to be approximately $3.5°\text{C/m}$. Similarly, $T_y\,(6,4) = \lim\limits_{h\to 0} \dfrac{T(6,4+h)-T(6,4)}{h}$, which we can approximate with $h = \pm 2$:

$T_y(6,4) \approx \dfrac{T(6,6)-T(6,4)}{2} = \dfrac{75-80}{2} = -2.5$, $T_y(6,4) \approx \dfrac{T(6,2)-T(6,4)}{-2} = \dfrac{87-80}{-2} = -3.5$. Averaging these values, we estimate $T_y(6,4)$ to be approximately $-3.0°\text{C/m}$.

(b) Here $\mathbf{u} = \left\langle \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right\rangle$, so by Equation 14.6.9, $D_{\mathbf{u}}\,T(6,4) = \nabla T(6,4)\cdot\mathbf{u} = T_x(6,4)\,\frac{1}{\sqrt{2}} + T_y(6,4)\,\frac{1}{\sqrt{2}}$. Using our estimates from part (a), we have $D_{\mathbf{u}}\,T(6,4) \approx (3.5)\,\frac{1}{\sqrt{2}} + (-3.0)\,\frac{1}{\sqrt{2}} = \frac{1}{2\sqrt{2}} \approx 0.35$. This means that as we move through the point $(6,4)$ in the direction of $\mathbf{u}$, the temperature increases at a rate of approximately $0.35°\text{C/m}$.

Alternatively, we can use Definition 14.6.2: $D_{\mathbf{u}}\,T(6,4) = \lim\limits_{h\to 0} \dfrac{T\left(6+h\,\frac{1}{\sqrt{2}}, 4+h\,\frac{1}{\sqrt{2}}\right) - T(6,4)}{h}$,

which we can estimate with $h = \pm 2\sqrt{2}$. Then $D_{\mathbf{u}}\,T(6,4) \approx \dfrac{T(8,6)-T(6,4)}{2\sqrt{2}} = \dfrac{80-80}{2\sqrt{2}} = 0$,

$D_{\mathbf{u}}\,T(6,4) \approx \dfrac{T(4,2)-T(6,4)}{-2\sqrt{2}} = \dfrac{74-80}{-2\sqrt{2}} = \dfrac{3}{\sqrt{2}}$. Averaging these values, we have $D_{\mathbf{u}}\,T(6,4) \approx \frac{3}{2\sqrt{2}} \approx 1.1°\text{C/m}$.

(c) $T_{xy}(x,y) = \dfrac{\partial}{\partial y}\left[T_x(x,y)\right] = \lim\limits_{h\to 0} \dfrac{T_x(x,y+h)-T_x(x,y)}{h}$, so $T_{xy}(6,4) = \lim\limits_{h\to 0} \dfrac{T_x(6,4+h)-T_x(6,4)}{h}$ which we can estimate with $h = \pm 2$. We have $T_x(6,4) \approx 3.5$ from part (a), but we will also need values for $T_x(6,6)$ and $T_x(6,2)$. If we use $h = \pm 2$ and the values given in the table, we have

$T_x(6,6) \approx \dfrac{T(8,6)-T(6,6)}{2} = \dfrac{80\ -\ 75}{2} = 2.5$, $T_x(6,6) \approx \dfrac{T(4,6)-T(6,6)}{-2} = \dfrac{68\ -\ 75}{-2} = 3.5$.

Averaging these values, we estimate $T_x(6,6) \approx 3.0$. Similarly,

$T_x(6,2) \approx \dfrac{T(8,2)-T_x(6,2)}{2} = \dfrac{90\ -\ 87}{2} = 1.5$, $T_x(6,2) \approx \dfrac{T(4,2)-T(6,2)}{-2} = \dfrac{74\ -\ 87}{-2} = 6.5$.

Averaging these values, we estimate $T_x(6,2) \approx 4.0$. Finally, we estimate $T_{xy}(6,4)$:

$T_{xy}(6,4) \approx \dfrac{T_x(6,6)-T_x(6,4)}{2} = \dfrac{3.0\ -\ 3.5}{2} = -0.25$, $T_{xy}(6,4) \approx \dfrac{T_x(6,2)-T_x(6,4)}{-2} = \dfrac{4.0\ -\ 3.5}{-2} = -0.25$.

Averaging these values, we have $T_{xy}\,(6,4) \approx -0.25$.

12. From the table, $T(6,4) = 80$, and from Exercise 11 we estimated $T_x(6,4) \approx 3.5$ and $T_y(6,4) \approx -3.0$. The linear approximation then is

$$T\,(x,y) \approx T(6,4) + T_x(6,4)(x-6) + T_y(6,4)(y-4) \approx 80 + 3.5(x-6) - 3(y-4) = 3.5x - 3y + 71$$

Thus at the point $(5, 3.8)$, we can use the linear approximation to estimate $T(5,3.8) \approx 3.5(5) - 3(3.8) + 71 \approx 77.1°\text{C}$.

13. $f(x,y) = (5y^3 + 2x^2y)^8 \quad\Rightarrow\quad f_x = 8(5y^3+2x^2y)^7(4xy) = 32xy(5y^3+2x^2y)^7$,

$f_y = 8(5y^3+2x^2y)^7(15y^2+2x^2) = (16x^2+120y^2)(5y^3+2x^2y)^7$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $g(u,v) = \dfrac{u+2v}{u^2+v^2} \quad \Rightarrow \quad g_u = \dfrac{(u^2+v^2)(1)-(u+2v)(2u)}{(u^2+v^2)^2} = \dfrac{v^2-u^2-4uv}{(u^2+v^2)^2}$,

$g_v = \dfrac{(u^2+v^2)(2)-(u+2v)(2v)}{(u^2+v^2)^2} = \dfrac{2u^2-2v^2-2uv}{(u^2+v^2)^2}$

15. $F(\alpha,\beta) = \alpha^2 \ln(\alpha^2+\beta^2) \quad \Rightarrow \quad F_\alpha = \alpha^2 \cdot \dfrac{1}{\alpha^2+\beta^2}\,(2\alpha) + \ln(\alpha^2+\beta^2)\cdot 2\alpha = \dfrac{2\alpha^3}{\alpha^2+\beta^2} + 2\alpha\ln(\alpha^2+\beta^2)$,

$F_\beta = \alpha^2 \cdot \dfrac{1}{\alpha^2+\beta^2}\,(2\beta) = \dfrac{2\alpha^2\beta}{\alpha^2+\beta^2}$

16. $G(x,y,z) = e^{xz}\sin(y/z) \quad \Rightarrow \quad G_x = ze^{xz}\sin(y/z),\ G_y = e^{xz}\cos(y/z)(1/z) = (e^{xz}/z)\cos(y/z)$,

$G_z = e^{xz}\cdot\cos(y/z)(-y/z^2) + \sin(y/z)\cdot xe^{xz} = e^{xz}\left[x\sin(y/z) - (y/z^2)\cos(y/z)\right]$

17. $S(u,v,w) = u\arctan(v\sqrt{w}\,) \quad \Rightarrow \quad S_u = \arctan(v\sqrt{w}\,),\ S_v = u\cdot\dfrac{1}{1+(v\sqrt{w})^2}\,(\sqrt{w}\,) = \dfrac{u\sqrt{w}}{1+v^2w}$,

$S_w = u\cdot\dfrac{1}{1+(v\sqrt{w})^2}\left(v\cdot\tfrac{1}{2}w^{-1/2}\right) = \dfrac{uv}{2\sqrt{w}\,(1+v^2w)}$

18. $C = 1449.2 + 4.6T - 0.055T^2 + 0.00029T^3 + (1.34-0.01T)(S-35) + 0.016D \quad \Rightarrow$

$\partial C/\partial T = 4.6 - 0.11T + 0.00087T^2 - 0.01(S-35)$, $\partial C/\partial S = 1.34 - 0.01T$, and $\partial C/\partial D = 0.016$. When $T = 10$, $S = 35$, and $D = 100$ we have $\partial C/\partial T = 4.6 - 0.11(10) + 0.00087(10)^2 - 0.01(35-35) \approx 3.587$, thus in 10°C water with salinity 35 parts per thousand and a depth of 100 m, the speed of sound increases by about 3.59 m/s for every degree Celsius that the water temperature rises. Similarly, $\partial C/\partial S = 1.34 - 0.01(10) = 1.24$, so the speed of sound increases by about 1.24 m/s for every part per thousand the salinity of the water increases. $\partial C/\partial D = 0.016$, so the speed of sound increases by about 0.016 m/s for every meter that the depth is increased.

19. $f(x,y) = 4x^3 - xy^2 \quad \Rightarrow \quad f_x = 12x^2 - y^2,\ f_y = -2xy,\ f_{xx} = 24x,\ f_{yy} = -2x,\ f_{xy} = f_{yx} = -2y$

20. $z = xe^{-2y} \quad \Rightarrow \quad z_x = e^{-2y},\ z_y = -2xe^{-2y},\ z_{xx} = 0,\ z_{yy} = 4xe^{-2y},\ z_{xy} = z_{yx} = -2e^{-2y}$

21. $f(x,y,z) = x^k y^l z^m \quad \Rightarrow \quad f_x = kx^{k-1}y^l z^m,\ f_y = lx^k y^{l-1} z^m,\ f_z = mx^k y^l z^{m-1},\ f_{xx} = k(k-1)x^{k-2}y^l z^m$,

$f_{yy} = l(l-1)x^k y^{l-2} z^m,\ f_{zz} = m(m-1)x^k y^l z^{m-2},\ f_{xy} = f_{yx} = klx^{k-1}y^{l-1}z^m,\ f_{xz} = f_{zx} = kmx^{k-1}y^l z^{m-1}$,

$f_{yz} = f_{zy} = lmx^k y^{l-1} z^{m-1}$

22. $v = r\cos(s+2t) \quad \Rightarrow \quad v_r = \cos(s+2t),\ v_s = -r\sin(s+2t),\ v_t = -2r\sin(s+2t),\ v_{rr} = 0,\ v_{ss} = -r\cos(s+2t)$,

$v_{tt} = -4r\cos(s+2t),\ v_{rs} = v_{sr} = -\sin(s+2t),\ v_{rt} = v_{tr} = -2\sin(s+2t),\ v_{st} = v_{ts} = -2r\cos(s+2t)$

23. $z = xy + xe^{y/x} \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = y - \dfrac{y}{x}e^{y/x} + e^{y/x},\ \dfrac{\partial z}{\partial y} = x + e^{y/x}$ and

$x\dfrac{\partial z}{\partial x} + y\dfrac{\partial z}{\partial y} = x\left(y - \dfrac{y}{x}e^{y/x} + e^{y/x}\right) + y\left(x + e^{y/x}\right) = xy - ye^{y/x} + xe^{y/x} + xy + ye^{y/x} = xy + xy + xe^{y/x} = xy + z.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $z = \sin(x + \sin t) \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = \cos(x + \sin t),\ \dfrac{\partial z}{\partial t} = \cos(x + \sin t)\cos t,$

$\dfrac{\partial^2 z}{\partial x \partial t} = -\sin(x + \sin t)\cos t,\ \dfrac{\partial^2 z}{\partial x^2} = -\sin(x + \sin t)$ and

$\dfrac{\partial z}{\partial x}\dfrac{\partial^2 z}{\partial x \partial t} = \cos(x + \sin t)\,[-\sin(x + \sin t)\cos t] = \cos(x + \sin t)\,(\cos t)\,[-\sin(x + \sin t)] = \dfrac{\partial z}{\partial t}\dfrac{\partial^2 z}{\partial x^2}.$

25. (a) $z = 3x^2 - y^2 + 2x$, $(1, -2, 1)$. $z_x = 6x + 2 \quad \Rightarrow \quad z_x(1, -2) = 8$ and $z_y = -2y \quad \Rightarrow \quad z_y(1, -2) = 4$, so an equation of the tangent plane is $z - 1 = 8(x - 1) + 4(y + 2)$, or $z = 8x + 4y + 1$.

(b) A normal vector to the tangent plane (and the surface) at $(1, -2, 1)$ is $\langle 8, 4, -1\rangle$. Then parametric equations for the normal line there are $x = 1 + 8t$, $y = -2 + 4t$, $z = 1 - t$, and symmetric equations are $\dfrac{x-1}{8} = \dfrac{y+2}{4} = \dfrac{z-1}{-1}$.

26. (a) $z = e^x \cos y$, $(0, 0, 1)$. $z_x = e^x \cos y \quad \Rightarrow \quad z_x(0, 0) = 1$ and $z_y = -e^x \sin y \quad \Rightarrow \quad z_y(0, 0) = 0$, so an equation of the tangent plane is $z - 1 = 1(x - 0) + 0(y - 0)$, or $z = x + 1$.

(b) A normal vector to the tangent plane (and the surface) at $(0, 0, 1)$ is $\langle 1, 0, -1\rangle$. Then parametric equations for the normal line there are $x = t$, $y = 0$, $z = 1 - t$, and symmetric equations are $x = 1 - z$, $y = 0$.

27. (a) Let $F(x, y, z) = x^2 + 2y^2 - 3z^2$. Then $F_x = 2x$, $F_y = 4y$, $F_z = -6z$, so $F_x(2, -1, 1) = 4$, $F_y(2, -1, 1) = -4$, $F_z(2, -1, 1) = -6$. From Equation 14.6.19, an equation of the tangent plane is $4(x - 2) - 4(y + 1) - 6(z - 1) = 0$ or, equivalently, $2x - 2y - 3z = 3$.

(b) From Equations 14.6.20, symmetric equations for the normal line are $\dfrac{x-2}{4} = \dfrac{y+1}{-4} = \dfrac{z-1}{-6}$. Parametric equations are $x = 2 + 4t$, $y = -1 - 4t$, $z = 1 - 6t$.

28. (a) Let $F(x, y, z) = xy + yz + zx$. Then $F_x = y + z$, $F_y = x + z$, $F_z = x + y$, so $F_x(1, 1, 1) = F_y(1, 1, 1) = F_z(1, 1, 1) = 2$. From Equation 14.6.19, an equation of the tangent plane is $2(x - 1) + 2(y - 1) + 2(z - 1) = 0$ or, equivalently, $x + y + z = 3$.

(b) From Equations 14.6.20, symmetric equations for the normal line are $\dfrac{x-1}{2} = \dfrac{y-1}{2} = \dfrac{z-1}{2}$ or, equivalently, $x = y = z$. Parametric equations are $x = 1 + 2t$, $y = 1 + 2t$, $z = 1 + 2t$.

29. (a) Let $F(x, y, z) = x + 2y + 3z - \sin(xyz)$. Then $F_x = 1 - yz\cos(xyz)$, $F_y = 2 - xz\cos(xyz)$, $F_z = 3 - xy\cos(xyz)$, so $F_x(2, -1, 0) = 1$, $F_y(2, -1, 0) = 2$, $F_z(2, -1, 0) = 5$. From Equation 14.6.19, an equation of the tangent plane is $1(x - 2) + 2(y + 1) + 5(z - 0) = 0$ or $x + 2y + 5z = 0$.

(b) From Equations 14.6.20, symmetric equations for the normal line are $\dfrac{x-2}{1} = \dfrac{y+1}{2} = \dfrac{z}{5}$ or $x - 2 = \dfrac{y+1}{2} = \dfrac{z}{5}$. Parametric equations are $x = 2 + t$, $y = -1 + 2t$, $z = 5t$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. Let $f(x,y) = x^2 + y^4$. Then $f_x(x,y) = 2x$ and $f_y(x,y) = 4y^3$, so $f_x(1,1) = 2$, $f_y(1,1) = 4$ and an equation of the tangent plane is $z - 2 = 2(x-1) + 4(y-1)$ or $2x + 4y - z = 4$. A normal vector to the tangent plane is $\langle 2, 4, -1\rangle$ so the normal line is given by $\dfrac{x-1}{2} = \dfrac{y-1}{4} = \dfrac{z-2}{-1}$ or $x = 1 + 2t,\ y = 1 + 4t$, $z = 2 - t$.

31. The hyperboloid is a level surface of the function $F(x,y,z) = x^2 + 4y^2 - z^2$, so a normal vector to the surface at (x_0, y_0, z_0) is $\nabla F(x_0, y_0, z_0) = \langle 2x_0, 8y_0, -2z_0\rangle$. A normal vector for the plane $2x + 2y + z = 5$ is $\langle 2, 2, 1\rangle$. For the planes to be parallel, we need the normal vectors to be parallel, so $\langle 2x_0, 8y_0, -2z_0\rangle = k\,\langle 2, 2, 1\rangle$, or $x_0 = k$, $y_0 = \frac{1}{4}k$, and $z_0 = -\frac{1}{2}k$. But $x_0^2 + 4y_0^2 - z_0^2 = 4 \quad\Rightarrow\quad k^2 + \frac{1}{4}k^2 - \frac{1}{4}k^2 = 4 \quad\Rightarrow\quad k^2 = 4 \quad\Rightarrow\quad k = \pm 2$. So there are two such points: $\left(2, \frac{1}{2}, -1\right)$ and $\left(-2, -\frac{1}{2}, 1\right)$.

32. $u = \ln(1 + se^{2t}) \quad\Rightarrow\quad du = \dfrac{\partial u}{\partial s}\,ds + \dfrac{\partial u}{\partial t}\,dt = \dfrac{e^{2t}}{1 + se^{2t}}\,ds + \dfrac{2se^{2t}}{1 + se^{2t}}\,dt$

33. $f(x,y,z) = x^3\sqrt{y^2 + z^2} \quad\Rightarrow\quad f_x(x,y,z) = 3x^2\sqrt{y^2+z^2},\ f_y(x,y,z) = \dfrac{yx^3}{\sqrt{y^2+z^2}},\ f_z(x,y,z) = \dfrac{zx^3}{\sqrt{y^2+z^2}}$,

so $f(2,3,4) = 8(5) = 40$, $f_x(2,3,4) = 3(4)\sqrt{25} = 60$, $f_y(2,3,4) = \frac{3(8)}{\sqrt{25}} = \frac{24}{5}$, and $f_z(2,3,4) = \frac{4(8)}{\sqrt{25}} = \frac{32}{5}$. Then the linear approximation of f at $(2,3,4)$ is

$$\begin{aligned} f(x,y,z) &\approx f(2,3,4) + f_x(2,3,4)(x-2) + f_y(2,3,4)(y-3) + f_z(2,3,4)(z-4) \\ &= 40 + 60(x-2) + \tfrac{24}{5}(y-3) + \tfrac{32}{5}(z-4) = 60x + \tfrac{24}{5}y + \tfrac{32}{5}z - 120 \end{aligned}$$

Then $(1.98)^3\sqrt{(3.01)^2 + (3.97)^2} = f(1.98, 3.01, 3.97) \approx 60(1.98) + \frac{24}{5}(3.01) + \frac{32}{5}(3.97) - 120 = 38.656$.

34. (a) $dA = \dfrac{\partial A}{\partial x}\,dx + \dfrac{\partial A}{\partial y}\,dy = \frac{1}{2}y\,dx + \frac{1}{2}x\,dy$ and $|\Delta x| \le 0.002$, $|\Delta y| \le 0.002$. Thus, the maximum error in the calculated area is about $dA = \frac{1}{2}(12)(0.002) + \frac{1}{2}(5)(0.002) = 0.017\ \text{m}^2$ or $170\ \text{cm}^2$.

(b) $z = \sqrt{x^2 + y^2}$, $dz = \dfrac{x}{\sqrt{x^2+y^2}}\,dx + \dfrac{y}{\sqrt{x^2+y^2}}\,dy$ and $|\Delta x| \le 0.002$, $|\Delta y| \le 0.002$. Thus, the maximum error in the calculated hypotenuse length, with $x = 5$, $y = 12$, and $z = \sqrt{5^2 + 12^2} = 13$, is about $dz = \frac{5}{13}(0.002) + \frac{12}{13}(0.002) = \frac{0.17}{65} \approx 0.0026$ m or 0.26 cm.

35. $u = x^2y^3 + z^4 \quad\Rightarrow\quad \dfrac{du}{dp} = \dfrac{\partial u}{\partial x}\dfrac{dx}{dp} + \dfrac{\partial u}{\partial y}\dfrac{dy}{dp} + \dfrac{\partial u}{\partial z}\dfrac{dz}{dp} = 2xy^3(1 + 6p) + 3x^2y^2(pe^p + e^p) + 4z^3(p\cos p + \sin p)$

36. $v = x^2\sin y + ye^{xy} \quad\Rightarrow\quad \dfrac{\partial v}{\partial s} = \dfrac{\partial v}{\partial x}\dfrac{\partial x}{\partial s} + \dfrac{\partial v}{\partial y}\dfrac{\partial y}{\partial s} = \left(2x\sin y + y^2e^{xy}\right)(1) + \left(x^2\cos y + xye^{xy} + e^{xy}\right)(t)$.

$s = 0, t = 1 \quad\Rightarrow\quad x = 2, y = 0$, so $\dfrac{\partial v}{\partial s} = 0 + (4+1)(1) = 5$.

$\dfrac{\partial v}{\partial t} = \dfrac{\partial v}{\partial x}\dfrac{\partial x}{\partial t} + \dfrac{\partial v}{\partial y}\dfrac{\partial y}{\partial t} = \left(2x\sin y + y^2e^{xy}\right)(2) + \left(x^2\cos y + xye^{xy} + e^{xy}\right)(s) = 0 + 0 = 0.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. By the Chain Rule, $\dfrac{\partial z}{\partial s} = \dfrac{\partial z}{\partial x}\dfrac{\partial x}{\partial s} + \dfrac{\partial z}{\partial y}\dfrac{\partial y}{\partial s}$. When $s = 1$ and $t = 2$, $x = g(1,2) = 3$ and $y = h(1,2) = 6$, so

$\dfrac{\partial z}{\partial s} = f_x(3,6)g_s(1,2) + f_y(3,6)\,h_s(1,2) = (7)(-1) + (8)(-5) = -47$. Similarly, $\dfrac{\partial z}{\partial t} = \dfrac{\partial z}{\partial x}\dfrac{\partial x}{\partial t} + \dfrac{\partial z}{\partial y}\dfrac{\partial y}{\partial t}$, so

$\dfrac{\partial z}{\partial t} = f_x(3,6)g_t(1,2) + f_y(3,6)\,h_t(1,2) = (7)(4) + (8)(10) = 108.$

38.

Using the tree diagram as a guide, we have

$$\frac{\partial w}{\partial p} = \frac{\partial w}{\partial t}\frac{\partial t}{\partial p} + \frac{\partial w}{\partial u}\frac{\partial u}{\partial p} + \frac{\partial w}{\partial v}\frac{\partial v}{\partial p} \qquad \frac{\partial w}{\partial q} = \frac{\partial w}{\partial t}\frac{\partial t}{\partial q} + \frac{\partial w}{\partial u}\frac{\partial u}{\partial q} + \frac{\partial w}{\partial v}\frac{\partial v}{\partial q}$$

$$\frac{\partial w}{\partial r} = \frac{\partial w}{\partial t}\frac{\partial t}{\partial r} + \frac{\partial w}{\partial u}\frac{\partial u}{\partial r} + \frac{\partial w}{\partial v}\frac{\partial v}{\partial r} \qquad \frac{\partial w}{\partial s} = \frac{\partial w}{\partial t}\frac{\partial t}{\partial s} + \frac{\partial w}{\partial u}\frac{\partial u}{\partial s} + \frac{\partial w}{\partial v}\frac{\partial v}{\partial s}$$

39. $z = y + f(x^2 - y^2) \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = 2xf'(x^2 - y^2), \quad \dfrac{\partial z}{\partial y} = 1 - 2yf'(x^2 - y^2) \quad \left[\text{where } f' = \dfrac{df}{d(x^2 - y^2)}\right]$. Then

$y\,\dfrac{\partial z}{\partial x} + x\,\dfrac{\partial z}{\partial y} = 2xyf'(x^2 - y^2) + x - 2xyf'(x^2 - y^2) = x.$

40. $A = \frac{1}{2}xy\sin\theta$ [Formula 6 in Appendix D], $dx/dt = 3$, $dy/dt = -2$, $d\theta/dt = 0.05$, and

$\dfrac{dA}{dt} = \dfrac{1}{2}\left[(y\sin\theta)\,\dfrac{dx}{dt} + (x\sin\theta)\,\dfrac{dy}{dt} + (xy\cos\theta)\,\dfrac{d\theta}{dt}\right]$. So when $x = 40$, $y = 50$ and $\theta = \frac{\pi}{6}$,

$\dfrac{dA}{dt} = \dfrac{1}{2}\left[(25)(3) + (20)(-2) + \left(1000\sqrt{3}\right)(0.05)\right] = \dfrac{35 + 50\sqrt{3}}{2} \approx 60.8 \text{ in}^2/\text{s}.$

41. $z = f(u,v)$, $u = xy$, and $v = y/x \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = \dfrac{\partial z}{\partial u}\,y + \dfrac{\partial z}{\partial v}\dfrac{-y}{x^2}$ and

$$\frac{\partial^2 z}{\partial x^2} = y\,\frac{\partial}{\partial x}\left(\frac{\partial z}{\partial u}\right) + \frac{2y}{x^3}\frac{\partial z}{\partial v} + \frac{-y}{x^2}\frac{\partial}{\partial x}\left(\frac{\partial z}{\partial v}\right) = \frac{2y}{x^3}\frac{\partial z}{\partial v} + y\left(\frac{\partial^2 z}{\partial u^2}\,y + \frac{\partial^2 z}{\partial v\,\partial u}\frac{-y}{x^2}\right) + \frac{-y}{x^2}\left(\frac{\partial^2 z}{\partial v^2}\frac{-y}{x^2} + \frac{\partial^2 z}{\partial u\,\partial v}\,y\right)$$

$$= \frac{2y}{x^3}\frac{\partial z}{\partial v} + y^2\,\frac{\partial^2 z}{\partial u^2} - \frac{2y^2}{x^2}\frac{\partial^2 z}{\partial u\,\partial v} + \frac{y^2}{x^4}\frac{\partial^2 z}{\partial v^2}$$

Also $\dfrac{\partial z}{\partial y} = x\,\dfrac{\partial z}{\partial u} + \dfrac{1}{x}\dfrac{\partial z}{\partial v}$ and

$$\frac{\partial^2 z}{\partial y^2} = x\,\frac{\partial}{\partial y}\left(\frac{\partial z}{\partial u}\right) + \frac{1}{x}\frac{\partial}{\partial y}\left(\frac{\partial z}{\partial v}\right) = x\left(\frac{\partial^2 z}{\partial u^2}x + \frac{\partial^2 z}{\partial v\,\partial u}\frac{1}{x}\right) + \frac{1}{x}\left(\frac{\partial^2 z}{\partial v^2}\frac{1}{x} + \frac{\partial^2 z}{\partial u\,\partial v}\,x\right) = x^2\,\frac{\partial^2 z}{\partial u^2} + 2\,\frac{\partial^2 z}{\partial u\,\partial v} + \frac{1}{x^2}\frac{\partial^2 z}{\partial v^2}$$

Thus

$$x^2\,\frac{\partial^2 z}{\partial x^2} - y^2\,\frac{\partial^2 z}{\partial y^2} = \frac{2y}{x}\frac{\partial z}{\partial v} + x^2y^2\,\frac{\partial^2 z}{\partial u^2} - 2y^2\,\frac{\partial^2 z}{\partial u\,\partial v} + \frac{y^2}{x^2}\frac{\partial^2 z}{\partial v^2} - x^2y^2\,\frac{\partial^2 z}{\partial u^2} - 2y^2\,\frac{\partial^2 z}{\partial u\,\partial v} - \frac{y^2}{x^2}\frac{\partial^2 z}{\partial v^2}$$

$$= \frac{2y}{x}\frac{\partial z}{\partial v} - 4y^2\,\frac{\partial^2 z}{\partial u\,\partial v} = 2v\,\frac{\partial z}{\partial v} - 4uv\,\frac{\partial^2 z}{\partial u\,\partial v}$$

since $y = xv = \dfrac{uv}{y}$ or $y^2 = uv$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. $\cos(xyz) = 1 + x^2y^2 + z^2$, so let $F(x, y, z) = 1 + x^2y^2 + z^2 - \cos(xyz) = 0$. Then by Equations 14.5.6 we have $\dfrac{\partial z}{\partial x} = -\dfrac{F_x}{F_z} = -\dfrac{2xy^2 + \sin(xyz) \cdot yz}{2z + \sin(xyz) \cdot xy} = -\dfrac{2xy^2 + yz\sin(xyz)}{2z + xy\sin(xyz)}$,

$\dfrac{\partial z}{\partial y} = -\dfrac{F_y}{F_z} = -\dfrac{2x^2y + \sin(xyz) \cdot xz}{2z + \sin(xyz) \cdot xy} = -\dfrac{2x^2y + xz\sin(xyz)}{2z + xy\sin(xyz)}$.

43. $f(x, y, z) = x^2e^{yz^2} \quad \Rightarrow \quad \nabla f = \langle f_x, f_y, f_z\rangle = \left\langle 2xe^{yz^2}, x^2e^{yz^2} \cdot z^2, x^2e^{yz^2} \cdot 2yz\right\rangle = \left\langle 2xe^{yz^2}, x^2z^2e^{yz^2}, 2x^2yze^{yz^2}\right\rangle$

44. (a) By Theorem 14.6.15, the maximum value of the directional derivative occurs when $\mathbf{u}$ has the same direction as the gradient vector.

(b) It is a minimum when $\mathbf{u}$ is in the direction opposite to that of the gradient vector (that is, $\mathbf{u}$ is in the direction of $-\nabla f$), since $D_{\mathbf{u}} f = |\nabla f| \cos\theta$ (see the proof of Theorem 14.6.15) has a minimum when $\theta = \pi$.

(c) The directional derivative is 0 when $\mathbf{u}$ is perpendicular to the gradient vector, since then $D_{\mathbf{u}} f = \nabla f \cdot \mathbf{u} = 0$.

(d) The directional derivative is half of its maximum value when $D_{\mathbf{u}} f = |\nabla f| \cos\theta = \frac{1}{2}|\nabla f| \quad \Leftrightarrow \quad \cos\theta = \frac{1}{2} \quad \Leftrightarrow$ $\theta = \frac{\pi}{3}$.

45. $f(x, y) = x^2e^{-y} \quad \Rightarrow \quad \nabla f = \left\langle 2xe^{-y}, -x^2e^{-y}\right\rangle$, $\nabla f(-2, 0) = \langle -4, -4\rangle$. The direction is given by $\langle 2 - (-2), -3 - 0\rangle = \langle 4, -3\rangle$, so $\mathbf{u} = \frac{1}{\sqrt{4^2 + (-3)^2}}\langle 4, -3\rangle = \frac{1}{5}\langle 4, -3\rangle$. Thus,

$D_{\mathbf{u}} f(-2, 0) = \nabla f(-2, 0) \cdot \mathbf{u} = \langle -4, -4\rangle \cdot \frac{1}{5}\langle 4, -3\rangle = \frac{1}{5}(-16 + 12) = -\frac{4}{5}$.

46. $f(x, y, z) = x^2y + x\sqrt{1 + z} \quad \Rightarrow \quad \nabla f = \left\langle 2xy + \sqrt{1 + z}, x^2, x/\left(2\sqrt{1 + z}\right)\right\rangle$, $\nabla f(1, 2, 3) = \left\langle 6, 1, \frac{1}{4}\right\rangle$. The direction is given by $2\,\mathbf{i} + \mathbf{j} - 2\,\mathbf{k}$, so $\mathbf{u} = \frac{1}{\sqrt{2^2 + 1^2 + (-2)^2}}\langle 2, 1, -2\rangle = \frac{1}{3}\langle 2, 1, -2\rangle$. Thus,

$D_{\mathbf{u}} f(1, 2, 3) = \nabla f(1, 2, 3) \cdot \mathbf{u} = \left\langle 6, 1, \frac{1}{4}\right\rangle \cdot \frac{1}{3}\langle 2, 1, -2\rangle = \frac{1}{3}\left(12 + 1 - \frac{1}{2}\right) = \frac{25}{6}$.

47. $f(x, y) = x^2y + \sqrt{y} \quad \Rightarrow \quad \nabla f = \left\langle 2xy, x^2 + 1/\left(2\sqrt{y}\right)\right\rangle$, $\nabla f(2, 1) = \left\langle 4, \frac{9}{2}\right\rangle$. Thus, the maximum rate of change of f at $(2, 1)$ is $|\nabla f(2, 1)| = \left|\left\langle 4, \frac{9}{2}\right\rangle\right| = \frac{\sqrt{145}}{2}$ in the direction $\left\langle 4, \frac{9}{2}\right\rangle$.

48. $f(x, y, z) = ze^{xy}$. $\nabla f = \langle zye^{xy}, zxe^{xy}, e^{xy}\rangle$, $\nabla f(0, 1, 2) = \langle 2, 0, 1\rangle$ is the direction of most rapid increase while the rate is $|\langle 2, 0, 1\rangle| = \sqrt{5}$.

49. First we draw a line passing through Homestead and the eye of the hurricane. We can approximate the directional derivative at Homestead in the direction of the eye of the hurricane by the average rate of change of wind speed between the points where this line intersects the contour lines closest to Homestead. In the direction of the eye of the hurricane, the wind speed changes from 45 to 50 knots. We estimate the distance between these two points to be approximately 8 miles, so the rate of change of wind speed in the direction given is approximately $\frac{50 - 45}{8} = \frac{5}{8} = 0.625$ knot/mi.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50. The surfaces are $f(x,y,z) = z - 2x^2 + y^2 = 0$ and $g(x,y,z) = z - 4 = 0$. The tangent line is perpendicular to both ∇f and ∇g at $(-2,2,4)$. The vector $\mathbf{v} = \nabla f \times \nabla g$ is therefore parallel to the line. $\nabla f(x,y,z) = \langle -4x, 2y, 1\rangle \quad \Rightarrow$ $\nabla f(-2,2,4) = \langle 8,4,1\rangle$, $\nabla g(x,y,z) = \langle 0,0,1\rangle \quad \Rightarrow \quad \nabla g\langle -2,2,4\rangle = \langle 0,0,1\rangle$. Hence

$\mathbf{v} = \nabla f \times \nabla g = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 8 & 4 & 1 \\ 0 & 0 & 1 \end{vmatrix} = 4\,\mathbf{i} - 8\,\mathbf{j}$. Thus, parametric equations are: $x = -2 + 4t,\ y = 2 - 8t,\ z = 4$.

51. $f(x,y) = x^2 - xy + y^2 + 9x - 6y + 10 \quad \Rightarrow \quad f_x = 2x - y + 9$, $f_y = -x + 2y - 6,\ f_{xx} = 2 = f_{yy},\ f_{xy} = -1$. Then $f_x = 0$ and $f_y = 0$ imply $y = 1, x = -4$. Thus the only critical point is $(-4,1)$ and $f_{xx}(-4,1) > 0$, $D(-4,1) = 3 > 0$, so $f(-4,1) = -11$ is a local minimum.

52. $f(x,y) = x^3 - 6xy + 8y^3 \quad \Rightarrow \quad f_x = 3x^2 - 6y,\ f_y = -6x + 24y^2,\ f_{xx} = 6x$, $f_{yy} = 48y$, $f_{xy} = -6$. Then $f_x = 0$ implies $y = x^2/2$, substituting into $f_y = 0$ implies $6x\left(x^3 - 1\right) = 0$, so the critical points are $(0,0)$, $\left(1, \frac{1}{2}\right)$. $D(0,0) = -36 < 0$ so $(0,0)$ is a saddle point while $f_{xx}\left(1, \frac{1}{2}\right) = 6 > 0$ and $D\left(1, \frac{1}{2}\right) = 108 > 0$ so $f\left(1, \frac{1}{2}\right) = -1$ is a local minimum.

53. $f(x,y) = 3xy - x^2y - xy^2 \quad \Rightarrow \quad f_x = 3y - 2xy - y^2,\ f_y = 3x - x^2 - 2xy$, $f_{xx} = -2y$, $f_{yy} = -2x,\ f_{xy} = 3 - 2x - 2y$. Then $f_x = 0$ implies $y(3 - 2x - y) = 0$ so $y = 0$ or $y = 3 - 2x$. Substituting into $f_y = 0$ implies $x(3 - x) = 0$ or $3x(-1 + x) = 0$. Hence the critical points are $(0,0)$, $(3,0)$, $(0,3)$ and $(1,1)$. $D(0,0) = D(3,0) = D(0,3) = -9 < 0$ so $(0,0)$, $(3,0)$, and $(0,3)$ are saddle points. $D(1,1) = 3 > 0$ and $f_{xx}(1,1) = -2 < 0$, so $f(1,1) = 1$ is a local maximum.

54. $f(x,y) = (x^2 + y)e^{y/2} \quad \Rightarrow \quad f_x = 2xe^{y/2}$, $f_y = e^{y/2}(2 + x^2 + y)/2$, $f_{xx} = 2e^{y/2}$, $f_{yy} = e^{y/2}(4 + x^2 + y)/4$, $f_{xy} = xe^{y/2}$. Then $f_x = 0$ implies $x = 0$, so $f_y = 0$ implies $y = -2$. But $f_{xx}(0,-2) > 0$, $D(0,-2) = e^{-2} - 0 > 0$ so $f(0,-2) = -2/e$ is a local minimum.

55. $f(x,y) = 4xy^2 - x^2y^2 - xy^3$. First solve inside D. Here $f_x = 4y^2 - 2xy^2 - y^3$, $f_y = 8xy - 2x^2y - 3xy^2$. Then $f_x = 0$ implies $y = 0$ or $y = 4 - 2x$, but $y = 0$ isn't inside D. Substituting $y = 4 - 2x$ into $f_y = 0$ implies $x = 0$, $x = 2$ or $x = 1$, but $x = 0$ isn't inside D, and when $x = 2$, $y = 0$ but $(2,0)$ isn't inside D. Thus the only critical point inside D is $(1,2)$ and $f(1,2) = 4$. Secondly we consider the boundary of D.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

On L_1: $f(x,0)=0$ and so $f=0$ on L_1. On L_2: $x=-y+6$ and $f(-y+6,y)=y^2(6-y)(-2)=-2(6y^2-y^3)$ which has critical points at $y=0$ and $y=4$. Then $f(6,0)=0$ while $f(2,4)=-64$. On L_3: $f(0,y)=0$, so $f=0$ on L_3. Thus on D the absolute maximum of f is $f(1,2)=4$ while the absolute minimum is $f(2,4)=-64$.

56. $f(x,y)=e^{-x^2-y^2}(x^2+2y^2)$. Inside D: $f_x=2xe^{-x^2-y^2}(1-x^2-2y^2)=0$ implies $x=0$ or $x^2+2y^2=1$. Then if $x=0$, $f_y=2ye^{-x^2-y^2}(2-x^2-2y^2)=0$ implies $y=0$ or $2-2y^2=0$ giving the critical points $(0,0)$, $(0,\pm1)$. If $x^2+2y^2=1$, then $f_y=0$ implies $y=0$ giving the critical points $(\pm1,0)$. Now $f(0,0)=0$, $f(\pm1,0)=e^{-1}$ and $f(0,\pm1)=2e^{-1}$. On the boundary of D: $x^2+y^2=4$, so $f(x,y)=e^{-4}(4+y^2)$ and f is smallest when $y=0$ and largest when $y^2=4$. But $f(\pm2,0)=4e^{-4}$, $f(0,\pm2)=8e^{-4}$. Thus on D the absolute maximum of f is $f(0,\pm1)=2e^{-1}$ and the absolute minimum is $f(0,0)=0$.

57. $f(x,y)=x^3-3x+y^4-2y^2$

From the graphs, it appears that f has a local maximum $f(-1,0)\approx 2$, local minimums $f(1,\pm1)\approx -3$, and saddle points at $(-1,\pm1)$ and $(1,0)$.

To find the exact quantities, we calculate $f_x=3x^2-3=0 \;\Leftrightarrow\; x=\pm1$ and $f_y=4y^3-4y=0 \;\Leftrightarrow\; y=0,\pm1$, giving the critical points estimated above. Also $f_{xx}=6x$, $f_{xy}=0$, $f_{yy}=12y^2-4$, so using the Second Derivatives Test, $D(-1,0)=24>0$ and $f_{xx}(-1,0)=-6<0$ indicating a local maximum $f(-1,0)=2$; $D(1,\pm1)=48>0$ and $f_{xx}(1,\pm1)=6>0$ indicating local minimums $f(1,\pm1)=-3$; and $D(-1,\pm1)=-48$ and $D(1,0)=-24$, indicating saddle points at $(-1,\pm1)$ and $(1,0)$.

58. $f(x,y)=12+10y-2x^2-8xy-y^4 \;\Rightarrow\; f_x(x,y)=-4x-8y$, $f_y(x,y)=10-8x-4y^3$. Now $f_x(x,y)=0 \;\Rightarrow\; x=-2x$, and substituting this into $f_y(x,y)=0$ gives $10+16y-4y^3=0 \;\Leftrightarrow\; 5+8y-2y^3=0$.

From the first graph, we see that this is true when $y\approx-1.542$, -0.717, or 2.260. (Alternatively, we could have found the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

solutions to $f_x = f_y = 0$ using a CAS.) So to three decimal places, the critical points are $(3.085, -1.542)$, $(1.434, -0.717)$, and $(-4.519, 2.260)$. Now in order to use the Second Derivatives Test, we calculate $f_{xx} = -4$, $f_{xy} = -8$, $f_{yy} = -12y^2$, and $D = 48y^2 - 64$. So since $D(3.085, -1.542) > 0$, $D(1.434, -0.717) < 0$, and $D(-4.519, 2.260) > 0$, and f_{xx} is always negative, $f(x, y)$ has local maximums $f(-4.519, 2.260) \approx 49.373$ and $f(3.085, -1.542) \approx 9.948$, and a saddle point at approximately $(1.434, -0.717)$. The highest point on the graph is approximately $(-4.519, 2.260, 49.373)$.

59. $f(x, y) = x^2y,\ g(x, y) = x^2 + y^2 = 1 \quad \Rightarrow \quad \nabla f = \langle 2xy, x^2 \rangle = \lambda \nabla g = \langle 2\lambda x, 2\lambda y \rangle$. Then $2xy = 2\lambda x$ implies $x = 0$ or $y = \lambda$. If $x = 0$ then $x^2 + y^2 = 1$ gives $y = \pm 1$ and we have possible points $(0, \pm 1)$ where $f(0, \pm 1) = 0$. If $y = \lambda$ then $x^2 = 2\lambda y$ implies $x^2 = 2y^2$ and substitution into $x^2 + y^2 = 1$ gives $3y^2 = 1 \quad \Rightarrow \quad y = \pm \frac{1}{\sqrt{3}}$ and $x = \pm\sqrt{\frac{2}{3}}$. The corresponding possible points are $\left(\pm\sqrt{\frac{2}{3}}, \pm\frac{1}{\sqrt{3}}\right)$. The absolute maximum is $f\left(\pm\sqrt{\frac{2}{3}}, \frac{1}{\sqrt{3}}\right) = \frac{2}{3\sqrt{3}}$ while the absolute minimum is $f\left(\pm\sqrt{\frac{2}{3}}, -\frac{1}{\sqrt{3}}\right) = -\frac{2}{3\sqrt{3}}$.

60. $f(x, y) = 1/x + 1/y,\ g(x, y) = 1/x^2 + 1/y^2 = 1 \quad \Rightarrow \quad \nabla f = \langle -x^{-2}, -y^{-2} \rangle = \lambda \nabla g = \langle -2\lambda x^{-3}, -2\lambda y^{-3} \rangle$. Then $-x^{-2} = -2\lambda x^3$ or $x = 2\lambda$ and $-y^{-2} = -2\lambda y^{-3}$ or $y = 2\lambda$. Thus $x = y$, so $1/x^2 + 1/y^2 = 2/x^2 = 1$ implies $x = \pm\sqrt{2}$ and the possible points are $(\pm\sqrt{2}, \pm\sqrt{2})$. The absolute maximum of f subject to $x^{-2} + y^{-2} = 1$ is then $f(\sqrt{2}, \sqrt{2}) = \sqrt{2}$ and the absolute minimum is $f(-\sqrt{2}, -\sqrt{2}) = -\sqrt{2}$.

61. $f(x, y, z) = xyz,\ g(x, y, z) = x^2 + y^2 + z^2 = 3. \quad \nabla f = \lambda \nabla g \quad \Rightarrow \quad \langle yz, xz, xy \rangle = \lambda \langle 2x, 2y, 2z \rangle$. If any of x, y, or z is zero, then $x = y = z = 0$ which contradicts $x^2 + y^2 + z^2 = 3$. Then $\lambda = \dfrac{yz}{2x} = \dfrac{xz}{2y} = \dfrac{xy}{2z} \quad \Rightarrow \quad 2y^2z = 2x^2z \quad \Rightarrow$ $y^2 = x^2$, and similarly $2yz^2 = 2x^2y \quad \Rightarrow \quad z^2 = x^2$. Substituting into the constraint equation gives $x^2 + x^2 + x^2 = 3 \quad \Rightarrow$ $x^2 = 1 = y^2 = z^2$. Thus the possible points are $(1, 1, \pm 1)$, $(1, -1, \pm 1)$, $(-1, 1, \pm 1)$, $(-1, -1, \pm 1)$. The absolute maximum is $f(1, 1, 1) = f(1, -1, -1) = f(-1, 1, -1) = f(-1, -1, 1) = 1$, and the absolute minimum is $f(1, 1, -1) = f(1, -1, 1) = f(-1, 1, 1) = f(-1, -1, -1) = -1$.

62. $f(x, y, z) = x^2 + 2y^2 + 3z^2,\ g(x, y, z) = x + y + z = 1,\ h(x, y, z) = x - y + 2z = 2 \quad \Rightarrow$ $\nabla f = \langle 2x, 4y, 6z \rangle = \lambda \nabla g + \mu \nabla h = \langle \lambda + \mu, \lambda - \mu, \lambda + 2\mu \rangle$ and $2x = \lambda + \mu$ **(1)**, $\quad 4y = \lambda - \mu$ **(2)**, $\quad 6z = \lambda + 2\mu$ **(3)**, $x + y + z = 1$ **(4)**, $\quad x - y + 2z = 2$ **(5)**. Then six times **(1)** plus three times **(2)** plus two times **(3)** implies $12(x + y + z) = 11\lambda + 7\mu$, so **(4)** gives $11\lambda + 7\mu = 12$. Also six times **(1)** minus three times **(2)** plus four times **(3)** implies $12(x - y + 2z) = 7\lambda + 17\mu$, so **(5)** gives $7\lambda + 17\mu = 24$. Solving $11\lambda + 7\mu = 12$, $7\lambda + 17\mu = 24$ simultaneously gives $\lambda = \frac{6}{23}$, $\mu = \frac{30}{23}$. Substituting into **(1)**, **(2)**, and **(3)** implies $x = \frac{18}{23}$, $y = -\frac{6}{23}$, $z = \frac{11}{23}$ giving only one point. Then $f\left(\frac{18}{23}, -\frac{6}{23}, \frac{11}{23}\right) = \frac{33}{23}$. Now since $(0, 0, 1)$ satisfies both constraints and $f(0, 0, 1) = 3 > \frac{33}{23}$, $f\left(\frac{18}{23}, -\frac{6}{23}, \frac{11}{23}\right) = \frac{33}{23}$ is an absolute minimum, and there is no absolute maximum.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

63. $f(x,y,z) = x^2 + y^2 + z^2,\ g(x,y,z) = xy^2z^3 = 2 \quad \Rightarrow \quad \nabla f = \langle 2x, 2y, 2z\rangle = \lambda \nabla g = \left\langle \lambda y^2z^3, 2\lambda xyz^3, 3\lambda xy^2z^2\right\rangle$.

Since $xy^2z^3 = 2$, $x \neq 0$, $y \neq 0$ and $z \neq 0$, so $2x = \lambda y^2z^3$ **(1)**, $\quad 1 = \lambda xz^3$ **(2)**, $\quad 2 = 3\lambda xy^2z$ **(3)**. Then **(2)** and **(3)** imply $\dfrac{1}{xz^3} = \dfrac{2}{3xy^2z}$ or $y^2 = \frac{2}{3}z^2$ so $y = \pm z\sqrt{\frac{2}{3}}$. Similarly **(1)** and **(3)** imply $\dfrac{2x}{y^2z^3} = \dfrac{2}{3xy^2z}$ or $3x^2 = z^2$ so $x = \pm\frac{1}{\sqrt{3}}z$. But $xy^2z^3 = 2$ so x and z must have the same sign, that is, $x = \frac{1}{\sqrt{3}}z$. Thus $g(x,y,z) = 2$ implies $\frac{1}{\sqrt{3}}z\left(\frac{2}{3}z^2\right)z^3 = 2$ or $z = \pm 3^{1/4}$ and the possible points are $(\pm 3^{-1/4}, 3^{-1/4}\sqrt{2}, \pm 3^{1/4})$, $(\pm 3^{-1/4}, -3^{-1/4}\sqrt{2}, \pm 3^{1/4})$. However at each of these points f takes on the same value, $2\sqrt{3}$. But $(2,1,1)$ also satisfies $g(x,y,z) = 2$ and $f(2,1,1) = 6 > 2\sqrt{3}$. Thus f has an absolute minimum value of $2\sqrt{3}$ and no absolute maximum subject to the constraint $xy^2z^3 = 2$.

Alternate solution: $g(x,y,z) = xy^2z^3 = 2$ implies $y^2 = \dfrac{2}{xz^3}$, so minimize $f(x,z) = x^2 + \dfrac{2}{xz^3} + z^2$. Then $f_x = 2x - \dfrac{2}{x^2z^3}$, $f_z = -\dfrac{6}{xz^4} + 2z$, $f_{xx} = 2 + \dfrac{4}{x^3z^3}$, $f_{zz} = \dfrac{24}{xz^5} + 2$ and $f_{xz} = \dfrac{6}{x^2z^4}$. Now $f_x = 0$ implies $2x^3z^3 - 2 = 0$ or $z = 1/x$. Substituting into $f_y = 0$ implies $-6x^3 + 2x^{-1} = 0$ or $x = \frac{1}{\sqrt[4]{3}}$, so the two critical points are $\left(\pm\frac{1}{\sqrt[4]{3}}, \pm\sqrt[4]{3}\right)$. Then $D\left(\pm\frac{1}{\sqrt[4]{3}}, \pm\sqrt[4]{3}\right) = (2+4)\left(2+\frac{24}{3}\right) - \left(\frac{6}{\sqrt{3}}\right)^2 > 0$ and $f_{xx}\left(\pm\frac{1}{\sqrt[4]{3}}, \pm\sqrt[4]{3}\right) = 6 > 0$, so each point is a minimum. Finally, $y^2 = \dfrac{2}{xz^3}$, so the four points closest to the origin are $\left(\pm\frac{1}{\sqrt[4]{3}}, \frac{\sqrt{2}}{\sqrt[4]{3}}, \pm\sqrt[4]{3}\right)$, $\left(\pm\frac{1}{\sqrt[4]{3}}, -\frac{\sqrt{2}}{\sqrt[4]{3}}, \pm\sqrt[4]{3}\right)$.

64. (a) The distance from a point (x,y) to the point $(-3,0)$ is $\sqrt{(x+3)^2+y^2}$. The distance from (x,y) to $(3,0)$ is $\sqrt{(x-3)^2+y^2}$. Then the function that gives the sum of the distances is

$f(x,y) = \sqrt{(x+3)^2+y^2} + \sqrt{(x-3)^2+y^2}$.

The graph shows several curves of $f(x,y) = k$ for $k = 9$, 10, 10.5, and 11. The smallest value of k that $g(x,y) = 16x + 15y$ intersects appears to be 10, so the point that minimizes the distance is approximately $(x,y) = (4, 2.4)$.

(b) $\nabla f = \left\langle \dfrac{x+3}{\sqrt{(x+3)^2+y^2}} + \dfrac{x-3}{\sqrt{(x-3)^2+y^2}}, \dfrac{y}{\sqrt{(x+3)^2+y^2}} + \dfrac{y}{\sqrt{(x-3)^2+y^2}} \right\rangle$. $\nabla g = \langle 16, 15\rangle$. Now $\nabla f(4, 2.4) = \left\langle \frac{7}{7.4} + \frac{1}{2.6}, \frac{2.4}{7.4} + \frac{2.4}{2.6}\right\rangle = \left\langle \frac{640}{481}, \frac{600}{481}\right\rangle$ and $\nabla g(4, 2.4) = \langle 16, 15\rangle$. ∇f and ∇g are parallel if $\nabla f = \lambda \nabla g$ for some value of λ, which is true when $\lambda = \frac{40}{481}$. Thus, ∇f and ∇g are parallel.

65.

The area of the triangle is $\frac{1}{2}ca\sin\theta$ [Formula 6 in Appendix D] and the area of the rectangle is bc. Thus, the area of the whole object is $f(a,b,c) = \frac{1}{2}ca\sin\theta + bc$. The perimeter of the object is $g(a,b,c) = 2a + 2b + c = P$. To simplify $\sin\theta$ in terms of a, b, and c notice that $a^2\sin^2\theta + \left(\frac{1}{2}c\right)^2 = a^2 \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\sin\theta = \dfrac{1}{2a}\sqrt{4a^2 - c^2}$. Thus, $f(a,b,c) = \dfrac{c}{4}\sqrt{4a^2 - c^2} + bc$. (Instead of using θ, we could just have used the Pythagorean Theorem.) As a result, by Lagrange's method, we must find a, b, c, and λ by solving $\nabla f = \lambda \nabla g$ which gives the following equations: $ca(4a^2 - c^2)^{-1/2} = 2\lambda$ **(1)**, $c = 2\lambda$ **(2)**, $\frac{1}{4}(4a^2 - c^2)^{1/2} - \frac{1}{4}c^2(4a^2 - c^2)^{-1/2} + b = \lambda$ **(3)**, and $2a + 2b + c = P$ **(4)**. From **(2)**, $\lambda = \frac{1}{2}c$ and so **(1)** produces $ca(4a^2 - c^2)^{-1/2} = c \quad \Rightarrow \quad (4a^2 - c^2)^{1/2} = a \quad \Rightarrow$ $4a^2 - c^2 = a^2 \quad \Rightarrow \quad c = \sqrt{3}\,a$ **(5)**. Similarly, since $\left(4a^2 - c^2\right)^{1/2} = a$ and $\lambda = \frac{1}{2}c$, **(3)** gives $\dfrac{a}{4} - \dfrac{c^2}{4a} + b = \dfrac{c}{2}$, so from **(5)**, $\dfrac{a}{4} - \dfrac{3a}{4} + b = \dfrac{\sqrt{3}\,a}{2} \quad \Rightarrow \quad -\dfrac{a}{2} - \dfrac{\sqrt{3}\,a}{2} = -b \quad \Rightarrow \quad b = \dfrac{a}{2}\left(1 + \sqrt{3}\right)$ **(6)**. Substituting **(5)** and **(6)** into **(4)** we get: $2a + a\left(1 + \sqrt{3}\right) + \sqrt{3}\,a = P \quad \Rightarrow \quad 3a + 2\sqrt{3}\,a = P \quad \Rightarrow \quad a = \dfrac{P}{3 + 2\sqrt{3}} = \dfrac{2\sqrt{3} - 3}{3}P$ and thus $b = \dfrac{\left(2\sqrt{3} - 3\right)\left(1 + \sqrt{3}\right)}{6}P = \dfrac{3 - \sqrt{3}}{6}P$ and $c = \left(2 - \sqrt{3}\right)P$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

PROBLEMS PLUS

1. The areas of the smaller rectangles are $A_1 = xy$, $A_2 = (L-x)y$, $A_3 = (L-x)(W-y)$, $A_4 = x(W-y)$. For $0 \le x \le L$, $0 \le y \le W$, let

$$\begin{aligned} f(x,y) &= A_1^2 + A_2^2 + A_3^2 + A_4^2 \\ &= x^2y^2 + (L-x)^2y^2 + (L-x)^2(W-y)^2 + x^2(W-y)^2 \\ &= [x^2 + (L-x)^2][y^2 + (W-y)^2] \end{aligned}$$

Then we need to find the maximum and minimum values of $f(x,y)$. Here

$f_x(x,y) = [2x - 2(L-x)][y^2 + (W-y)^2] = 0 \quad \Rightarrow \quad 4x - 2L = 0$ or $x = \frac{1}{2}L$, and

$f_y(x,y) = [x^2 + (L-x)^2][2y - 2(W-y)] = 0 \quad \Rightarrow \quad 4y - 2W = 0$ or $y = W/2$. Also

$f_{xx} = 4[y^2 + (W-y)^2]$, $f_{yy} = 4[x^2 + (L-x)^2]$, and $f_{xy} = (4x-2L)(4y-2W)$. Then

$D = 16[y^2 + (W-y)^2][x^2 + (L-x)^2] - (4x-2L)^2(4y-2W)^2$. Thus when $x = \frac{1}{2}L$ and $y = \frac{1}{2}W$, $D > 0$ and

$f_{xx} = 2W^2 > 0$. Thus a minimum of f occurs at $\left(\frac{1}{2}L, \frac{1}{2}W\right)$ and this minimum value is $f\left(\frac{1}{2}L, \frac{1}{2}W\right) = \frac{1}{4}L^2W^2$.

There are no other critical points, so the maximum must occur on the boundary. Now along the width of the rectangle let

$g(y) = f(0,y) = f(L,y) = L^2[y^2 + (W-y)^2]$, $0 \le y \le W$. Then $g'(y) = L^2[2y - 2(W-y)] = 0 \quad \Leftrightarrow \quad y = \frac{1}{2}W$.

And $g\left(\frac{1}{2}\right) = \frac{1}{2}L^2W^2$. Checking the endpoints, we get $g(0) = g(W) = L^2W^2$. Along the length of the rectangle let

$h(x) = f(x,0) = f(x,W) = W^2[x^2 + (L-x)^2]$, $0 \le x \le L$. By symmetry $h'(x) = 0 \quad \Leftrightarrow \quad x = \frac{1}{2}L$ and

$h\left(\frac{1}{2}L\right) = \frac{1}{2}L^2W^2$. At the endpoints we have $h(0) = h(L) = L^2W^2$. Therefore L^2W^2 is the maximum value of f.

This maximum value of f occurs when the "cutting" lines correspond to sides of the rectangle.

2. (a) The level curves of the function $C(x,y) = e^{-(x^2+2y^2)/10^4}$ are the curves $e^{-(x^2+2y^2)/10^4} = k$ (k is a positive constant). This equation is equivalent to $x^2 + 2y^2 = K \quad \Rightarrow \quad \dfrac{x^2}{\left(\sqrt{K}\right)^2} + \dfrac{y^2}{\left(\sqrt{K/2}\right)^2} = 1$, where $K = -10^4 \ln k$, a family of ellipses. We sketch level curves for $K = 1$, 2, 3, and 4. If the shark always swims in the direction of maximum increase of blood concentration, its direction at any point would coincide with the gradient vector. Then we know the shark's path is perpendicular to the level curves it intersects. We sketch one example of such a path.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\nabla C = -\frac{2}{10^4} e^{-(x^2+2y^2)/10^4}(x\,\mathbf{i} + 2y\,\mathbf{j})$. And ∇C points in the direction of most rapid increase in concentration; that is, ∇C is tangent to the most rapid increase curve. If $r(t) = x(t)\,\mathbf{i} + y(t)\,\mathbf{j}$ is a parametrization of the most rapid increase curve, then $\dfrac{d\mathbf{r}}{dt} = \dfrac{dx}{dt}\,\mathbf{i} + \dfrac{dy}{dt}\,\mathbf{j}$ is tangent to the curve, so $\dfrac{d\mathbf{r}}{dt} = \lambda \nabla C \quad \Rightarrow \quad \dfrac{dx}{dt} = \lambda\left[-\dfrac{2}{10^4}\, e^{-(x^2+2y^2)/10^4}\right] x$ and $\dfrac{dy}{dt} = \lambda\left[-\dfrac{2}{10^4} e^{-(x^2+2y^2)/10^4}\right](2y)$. Therefore $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = 2\,\dfrac{y}{x} \quad \Rightarrow \quad \dfrac{dy}{y} = 2\,\dfrac{dx}{x} \quad \Rightarrow \quad \ln|y| = 2\ln|x|$ so that $y = kx^2$ for some constant k. But $y(x_0) = y_0 \quad \Rightarrow \quad y_0 = kx_0^2 \quad \Rightarrow \quad k = y_0/x_0^2 \quad (x_0 = 0 \quad \Rightarrow \quad y_0 = 0 \quad \Rightarrow$ the shark is already at the origin, so we can assume $x_0 \neq 0$.) Therefore the path the shark will follow is along the parabola $y = y_0(x/x_0)^2$.

3. (a) The area of a trapezoid is $\frac{1}{2}h(b_1 + b_2)$, where h is the height (the distance between the two parallel sides) and b_1, b_2 are the lengths of the bases (the parallel sides). From the figure in the text, we see that $h = x\sin\theta$, $b_1 = w - 2x$, and $b_2 = w - 2x + 2x\cos\theta$. Therefore the cross-sectional area of the rain gutter is

$$\begin{aligned} A(x,\theta) &= \tfrac{1}{2}x\sin\theta\,[(w-2x) + (w - 2x + 2x\cos\theta)] = (x\sin\theta)(w - 2x + x\cos\theta) \\ &= wx\sin\theta - 2x^2\sin\theta + x^2\sin\theta\cos\theta,\ 0 < x \le \tfrac{1}{2}w,\ 0 < \theta \le \tfrac{\pi}{2} \end{aligned}$$

We look for the critical points of A: $\partial A/\partial x = w\sin\theta - 4x\sin\theta + 2x\sin\theta\cos\theta$ and $\partial A/\partial\theta = wx\cos\theta - 2x^2\cos\theta + x^2(\cos^2\theta - \sin^2\theta)$, so $\partial A/\partial x = 0 \quad \Leftrightarrow \quad \sin\theta\,(w - 4x + 2x\cos\theta) = 0 \quad \Leftrightarrow$ $\cos\theta = \dfrac{4x - w}{2x} = 2 - \dfrac{w}{2x} \quad (0 < \theta \le \frac{\pi}{2} \quad \Rightarrow \quad \sin\theta > 0)$. If, in addition, $\partial A/\partial\theta = 0$, then

$$\begin{aligned} 0 &= wx\cos\theta - 2x^2\cos\theta + x^2(2\cos^2\theta - 1) \\ &= wx\Big(2 - \frac{w}{2x}\Big) - 2x^2\Big(2 - \frac{w}{2x}\Big) + x^2\left[2\Big(2 - \frac{w}{2x}\Big)^2 - 1\right] \\ &= 2wx - \tfrac{1}{2}w^2 - 4x^2 + wx + x^2\left[8 - \frac{4w}{x} + \frac{w^2}{2x^2} - 1\right] = -wx + 3x^2 = x(3x - w) \end{aligned}$$

Since $x > 0$, we must have $x = \frac{1}{3}w$, in which case $\cos\theta = \frac{1}{2}$, so $\theta = \frac{\pi}{3}$, $\sin\theta = \frac{\sqrt{3}}{2}$, $k = \frac{\sqrt{3}}{6}w$, $b_1 = \frac{1}{3}w$, $b_2 = \frac{2}{3}w$, and $A = \frac{\sqrt{3}}{12}w^2$. As in Example 14.7.6, we can argue from the physical nature of this problem that we have found a local maximum of A. Now checking the boundary of A, let $g(\theta) = A(w/2,\theta) = \frac{1}{2}w^2\sin\theta - \frac{1}{2}w^2\sin\theta + \frac{1}{4}w^2\sin\theta\cos\theta = \frac{1}{8}w^2\sin 2\theta$, $0 < \theta \le \frac{\pi}{2}$. Clearly g is maximized when $\sin 2\theta = 1$ in which case $A = \frac{1}{8}w^2$. Also along the line $\theta = \frac{\pi}{2}$, let $h(x) = A\big(x, \frac{\pi}{2}\big) = wx - 2x^2,\ 0 < x < \frac{1}{2}w \quad \Rightarrow$ $h'(x) = w - 4x = 0 \quad \Leftrightarrow \quad x = \frac{1}{4}w$, and $h\big(\frac{1}{4}w\big) = w\big(\frac{1}{4}w\big) - 2\big(\frac{1}{4}w\big)^2 = \frac{1}{8}w^2$. Since $\frac{1}{8}w^2 < \frac{\sqrt{3}}{12}\,w^2$, we conclude that the local maximum found earlier was an absolute maximum; that is, the base and the sides are of equal length.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) If the metal were bent into a semicircular gutter of radius r, we would have $w = \pi r$ and $A = \frac{1}{2}\pi r^2 = \frac{1}{2}\pi\left(\frac{w}{\pi}\right)^2 = \frac{w^2}{2\pi}$.

Since $\frac{w^2}{2\pi} > \frac{\sqrt{3}\,w^2}{12}$, it *would* be better to bend the metal into a gutter with a semicircular cross-section.

4. Since $(x+y+z)^r/(x^2+y^2+z^2)$ is a rational function with domain $\{(x,y,z) \mid (x,y,z) \neq (0,0,0)\}$, f is continuous on $\mathbb{R}^3$ if and only if $\lim\limits_{(x,y,z)\to(0,0,0)} f(x,y,z) = f(0,0,0) = 0$. Recall that $(a+b)^2 \leq 2a^2 + 2b^2$ and a double application of this inequality to $(x+y+z)^2$ gives $(x+y+z)^2 \leq 4x^2 + 4y^2 + 2z^2 \leq 4(x^2+y^2+z^2)$. Now for each r,

$|(x+y+z)^r| = \left(|x+y+z|^2\right)^{r/2} = \left[(x+y+z)^2\right]^{r/2} \leq \left[4(x^2+y^2+z^2)\right]^{r/2} = 2^r(x^2+y^2+z^2)^{r/2}$

for $(x,y,z) \neq (0,0,0)$. Thus

$$|f(x,y,z) - 0| = \left|\frac{(x+y+z)^r}{x^2+y^2+z^2}\right| = \frac{|(x+y+z)^r|}{x^2+y^2+z^2} \leq 2^r\frac{(x^2+y^2+z^2)^{r/2}}{x^2+y^2+z^2} = 2^r(x^2+y^2+z^2)^{(r/2)-1}$$

for $(x,y,z) \neq (0,0,0)$. Thus if $(r/2) - 1 > 0$, that is $r > 2$, then $2^r(x^2+y^2+z^2)^{(r/2)-1} \to 0$ as $(x,y,z) \to (0,0,0)$ and so $\lim\limits_{(x,y,z)\to(0,0,0)} (x+y+z)^r/(x^2+y^2+z^2) = 0$. Hence for $r > 2$, f is continuous on $\mathbb{R}^3$. Now if $r \leq 2$, then as $(x,y,z) \to (0,0,0)$ along the x-axis, $f(x,0,0) = x^r/x^2 = x^{r-2}$ for $x \neq 0$. So when $r = 2$, $f(x,y,z) \to 1 \neq 0$ as $(x,y,z) \to (0,0,0)$ along the x-axis and when $r < 2$ the limit of $f(x,y,z)$ as $(x,y,z) \to (0,0,0)$ along the x-axis doesn't exist and thus can't be zero. Hence for $r \leq 2$ f isn't continuous at $(0,0,0)$ and thus is not continuous on $\mathbb{R}^3$.

5. Let $g(x,y) = xf\left(\frac{y}{x}\right)$. Then $g_x(x,y) = f\left(\frac{y}{x}\right) + xf'\left(\frac{y}{x}\right)\left(-\frac{y}{x^2}\right) = f\left(\frac{y}{x}\right) - \frac{y}{x}f'\left(\frac{y}{x}\right)$ and

$g_y(x,y) = xf'\left(\frac{y}{x}\right)\left(\frac{1}{x}\right) = f'\left(\frac{y}{x}\right)$. Thus the tangent plane at (x_0, y_0, z_0) on the surface has equation

$$z - x_0 f\left(\frac{y_0}{x_0}\right) = \left[f\left(\frac{y_0}{x_0}\right) - y_0 x_0^{-1} f'\left(\frac{y_0}{x_0}\right)\right](x - x_0) + f'\left(\frac{y_0}{x_0}\right)(y - y_0) \quad \Rightarrow$$

$\left[f\left(\frac{y_0}{x_0}\right) - y_0 x_0^{-1} f'\left(\frac{y_0}{x_0}\right)\right]x + \left[f'\left(\frac{y_0}{x_0}\right)\right]y - z = 0$. But any plane whose equation is of the form $ax + by + cz = 0$ passes through the origin. Thus the origin is the common point of intersection.

6. (a) At $(x_1, y_1, 0)$ the equations of the tangent planes to $z = f(x,y)$ and $z = g(x,y)$ are

$$P_1\colon\ z - f(x_1,y_1) = f_x(x_1,y_1)(x - x_1) + f_y(x_1,y_1)(y - y_1)$$

and

$$P_2\colon\ z - g(x_1,y_1) = g_x(x_1,y_1)(x - x_1) + g_y(x_1,y_1)(y - y_1)$$

respectively. P_1 intersects the xy-plane in the line given by $f_x(x_1,y_1)(x - x_1) + f_y(x_1,y_1)(y - y_1) = -f(x_1,y_1)$, $z = 0$; and P_2 intersects the xy-plane in the line given by $g_x(x_1,y_1)(x - x_1) + g_y(x_1,y_1)(y - y_1) = -g(x_1,y_1)$, $z = 0$. The point $(x_2, y_2, 0)$ is the point of intersection of these two lines, since $(x_2, y_2, 0)$ is the point where the line of

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

intersection of the two tangent planes intersects the xy-plane. Thus (x_2, y_2) is the solution of the simultaneous equations

$$f_x(x_1, y_1)(x_2 - x_1) + f_y(x_1, y_1)(y_2 - y_1) = -f(x_1, y_1)$$

and
$$g_x(x_1, y_1)(x_2 - x_1) + g_y(x_1, y_1)(y_2 - y_1) = -g(x_1, y_1)$$

For simplicity, rewrite $f_x(x_1, y_1)$ as f_x and similarly for f_y, g_x, g_y, f and g and solve the equations $(f_x)(x_2 - x_1) + (f_y)(y_2 - y_1) = -f$ and $(g_x)(x_2 - x_1) + (g_y)(y_2 - y_1) = -g$ simultaneously for $(x_2 - x_1)$ and $(y_2 - y_1)$. Then $y_2 - y_1 = \dfrac{gf_x - fg_x}{g_x f_y - f_x g_y}$ or $y_2 = y_1 - \dfrac{gf_x - fg_x}{f_x g_y - g_x f_y}$ and $(f_x)(x_2 - x_1) + \dfrac{(f_y)(gf_x - fg_x)}{g_x f_y - f_x g_y} = -f$ so $x_2 - x_1 = \dfrac{-f - [(f_y)(gf_x - fg_x)/(g_x f_y - f_x g_y)]}{f_x} = \dfrac{fg_y - f_y g}{g_x f_y - f_x g_y}$. Hence $x_2 = x_1 - \dfrac{fg_y - f_y g}{f_x g_y - g_x f_y}$.

(b) Let $f(x, y) = x^x + y^y - 1000$ and $g(x, y) = x^y + y^x - 100$. Then we wish to solve the system of equations $f(x, y) = 0$, $g(x, y) = 0$. Recall $\dfrac{d}{dx}[x^x] = x^x(1 + \ln x)$ (differentiate logarithmically), so $f_x(x, y) = x^x(1 + \ln x)$, $f_y(x, y) = y^y(1 + \ln y)$, $g_x(x, y) = yx^{y-1} + y^x \ln y$, and $g_y(x, y) = x^y \ln x + xy^{x-1}$. Looking at the graph, we estimate the first point of intersection of the curves, and thus the solution to the system, to be approximately $(2.5, 4.5)$. Then following the method of part (a), $x_1 = 2.5$, $y_1 = 4.5$ and

$$x_2 = 2.5 - \frac{f(2.5, 4.5)\, g_y(2.5, 4.5) - f_y(2.5, 4.5)\, g(2.5, 4.5)}{f_x(2.5, 4.5)\, g_y(2.5, 4.5) - f_y(2.5, 4.5)\, g_x(2.5, 4.5} \approx 2.447674117$$

$$y_2 = 4.5 - \frac{f_x(2.5, 4.5)\, g(2.5, 4.5) - f(2.5, 4.5)\, g_x(2.5, 4.5)}{f_x(2.5, 4.5)\, g_y(2.5, 4.5) - f_y(2.5, 4.5)\, g_x(2.5, 4.5)} \approx 4.555657467$$

Continuing this procedure, we arrive at the following values. (If you use a CAS, you may need to increase its computational precision.)

$x_1 = 2.5$	$y_1 = 4.5$
$x_2 = 2.447674117$	$y_2 = 4.555657467$
$x_3 = 2.449614877$	$y_3 = 4.551969333$
$x_4 = 2.449624628$	$y_4 = 4.551951420$
$x_5 = 2.449624628$	$y_5 = 4.551951420$

Thus, to six decimal places, the point of intersection is $(2.449625, 4.551951)$. The second point of intersection can be found similarly, or, by symmetry it is approximately $(4.551951, 2.449625)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. Since we are minimizing the area of the ellipse, and the circle lies above the x-axis, the ellipse will intersect the circle for only one value of y. This y-value must satisfy both the equation of the circle and the equation of the ellipse. Now $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}=1 \quad\Rightarrow\quad x^2=\dfrac{a^2}{b^2}\left(b^2-y^2\right)$. Substituting into the equation of the circle gives $\dfrac{a^2}{b^2}\left(b^2-y^2\right)+y^2-2y=0 \quad\Rightarrow\quad \left(\dfrac{b^2-a^2}{b^2}\right)y^2-2y+a^2=0$.

In order for there to be only one solution to this quadratic equation, the discriminant must be 0, so $4-4a^2\,\dfrac{b^2-a^2}{b^2}=0 \quad\Rightarrow$ $b^2-a^2b^2+a^4=0$. The area of the ellipse is $A(a,b)=\pi ab$, and we minimize this function subject to the constraint $g(a,b)=b^2-a^2b^2+a^4=0$.

Now $\nabla A=\lambda\nabla g \quad\Leftrightarrow\quad \pi b=\lambda(4a^3-2ab^2),\ \pi a=\lambda(2b-2ba^2) \quad\Rightarrow\quad \lambda=\dfrac{\pi b}{2a(2a^2-b^2)}$ **(1)**,

$\lambda=\dfrac{\pi a}{2b(1-a^2)}$ **(2)**, $b^2-a^2b^2+a^4=0$ **(3)**. Comparing **(1)** and **(2)** gives $\dfrac{\pi b}{2a(2a^2-b^2)}=\dfrac{\pi a}{2b(1-a^2)} \quad\Rightarrow$

$2\pi b^2=4\pi a^4 \quad\Leftrightarrow\quad a^2=\frac{1}{\sqrt{2}}\,b$. Substitute this into **(3)** to get $b=\frac{3}{\sqrt{2}} \quad\Rightarrow\quad a=\sqrt{\frac{3}{2}}$.

8. Let $\mathbf{u}=\langle a,b,c\rangle$ and $\mathbf{v}=\langle x,y,1\rangle$, so $|\mathbf{u}|=\sqrt{a^2+b^2+c^2}$, $|\mathbf{v}|=\sqrt{x^2+y^2+1}$, and $\mathbf{u}\cdot\mathbf{v}=ax+by+c$. Then by the Cauchy-Schwarz Inequality, $|\mathbf{u}\cdot\mathbf{v}|\le|\mathbf{u}|\ |\mathbf{v}| \quad\Rightarrow\quad |ax+by+c|\le\sqrt{a^2+b^2+c^2}\,\sqrt{x^2+y^2+1}$. Squaring both sides, we have $(ax+by+c)^2\le\left(a^2+b^2+c^2\right)\left(x^2+y^2+1\right) \quad\Rightarrow\quad \dfrac{(ax+by+c)^2}{x^2+y^2+1}\le a^2+b^2+c^2$

(since $x^2+y^2+1>0$). Thus $f(x,y)=\dfrac{(ax+by+c)^2}{x^2+y^2+1}\le a^2+b^2+c^2$. We have equality if $(ax+by+c)^2=\left(a^2+b^2+c^2\right)\left(x^2+y^2+1\right)$ or equivalently $c^2\left[(a/c)x+(b/c)y+1\right]^2=c^2\left[(a/c)^2+(b/c)^2+1\right]\left(x^2+y^2+1\right)$ which is true when $x=a/c$ and $y=b/c$. Thus the maximum value of f is $f(a/c,b/c)=a^2+b^2+c^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15 ☐ MULTIPLE INTEGRALS

15.1 Double Integrals over Rectangles

1. (a) The subrectangles are shown in the figure.

The surface is the graph of $f(x, y) = xy$ and $\Delta A = 4$, so we estimate

$$V \approx \sum_{i=1}^{3} \sum_{j=1}^{2} f(x_i, y_j)\,\Delta A$$

$$= f(2,2)\,\Delta A + f(2,4)\,\Delta A + f(4,2)\,\Delta A + f(4,4)\,\Delta A + f(6,2)\,\Delta A + f(6,4)\,\Delta A$$

$$= 4(4) + 8(4) + 8(4) + 16(4) + 12(4) + 24(4) = 288$$

(b) $V \approx \sum_{i=1}^{3} \sum_{j=1}^{2} f(\overline{x}_i, \overline{y}_j)\,\Delta A = f(1,1)\,\Delta A + f(1,3)\,\Delta A + f(3,1)\,\Delta A + f(3,3)\,\Delta A + f(5,1)\,\Delta A + f(5,3)\,\Delta A$

$$= 1(4) + 3(4) + 3(4) + 9(4) + 5(4) + 15(4) = 144$$

2. (a) The subrectangles are shown in the figure.

Here $\Delta A = 2$ and we estimate

$$\iint_R \left(1 - xy^2\right) dA \approx \sum_{i=1}^{2} \sum_{j=1}^{3} f\left(x_{ij}^*, y_{ij}^*\right) \Delta A$$

$$= f(2,-1)\,\Delta A + f(2,0)\,\Delta A + f(2,1)\,\Delta A + f(4,-1)\,\Delta A + f(4,0)\,\Delta A + f(4,1)\,\Delta A$$

$$= (-1)(2) + 1(2) + (-1)(2) + (-3)(2) + 1(2) + (-3)(2) = -12$$

(b) $\iint_R \left(1 - xy^2\right) dA \approx \sum_{i=1}^{2} \sum_{j=1}^{3} f\left(x_{ij}^*, y_{ij}^*\right) \Delta A$

$$= f(0,0)\,\Delta A + f(0,1)\,\Delta A + f(0,2)\,\Delta A + f(2,0)\,\Delta A + f(2,1)\,\Delta A + f(2,2)\,\Delta A$$

$$= 1(2) + 1(2) + 1(2) + 1(2) + (-1)(2) + (-7)(2) = -8$$

3. (a) The subrectangles are shown in the figure. Since $\Delta A = 1 \cdot \frac{1}{2} = \frac{1}{2}$, we estimate

$$\iint_R xe^{-xy}\, dA \approx \sum_{i=1}^{2} \sum_{j=1}^{2} f\left(x_{ij}^*, y_{ij}^*\right) \Delta A$$

$$= f\left(1, \tfrac{1}{2}\right)\,\Delta A + f(1,1)\,\Delta A + f\left(2, \tfrac{1}{2}\right)\,\Delta A + f(2,1)\,\Delta A$$

$$= e^{-1/2}\left(\tfrac{1}{2}\right) + e^{-1}\left(\tfrac{1}{2}\right) + 2e^{-1}\left(\tfrac{1}{2}\right) + 2e^{-2}\left(\tfrac{1}{2}\right) \approx 0.990$$

(b) $\iint_R xe^{-xy}\, dA \approx \sum_{i=1}^{2} \sum_{j=1}^{2} f(\overline{x}_i, \overline{y}_j)\,\Delta A$

$$= f\left(\tfrac{1}{2}, \tfrac{1}{4}\right)\,\Delta A + f\left(\tfrac{1}{2}, \tfrac{3}{4}\right)\,\Delta A + f\left(\tfrac{3}{2}, \tfrac{1}{4}\right)\,\Delta A + f\left(\tfrac{3}{2}, \tfrac{3}{4}\right)\,\Delta A$$

$$= \tfrac{1}{2}e^{-1/8}\left(\tfrac{1}{2}\right) + \tfrac{1}{2}e^{-3/8}\left(\tfrac{1}{2}\right) + \tfrac{3}{2}e^{-3/8}\left(\tfrac{1}{2}\right) + \tfrac{3}{2}e^{-9/8}\left(\tfrac{1}{2}\right) \approx 1.151$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a) The subrectangles are shown in the figure.

The surface is the graph of $f(x,y) = 1 + x^2 + 3y$ and $\Delta A = \frac{1}{2} \cdot \frac{3}{2} = \frac{3}{4}$, so we estimate

$$V = \iint_R (1 + x^2 + 3y)\, dA \approx \sum_{i=1}^{2} \sum_{j=1}^{2} f\left(x_{ij}^*, y_{ij}^*\right) \Delta A$$

$$= f(1,0)\,\Delta A + f\left(1, \tfrac{3}{2}\right) \Delta A + f\left(\tfrac{3}{2}, 0\right) \Delta A + f\left(\tfrac{3}{2}, \tfrac{3}{2}\right) \Delta A$$

$$= 2\left(\tfrac{3}{4}\right) + \tfrac{13}{2}\left(\tfrac{3}{4}\right) + \tfrac{13}{4}\left(\tfrac{3}{4}\right) + \tfrac{31}{4}\left(\tfrac{3}{4}\right) = \tfrac{39}{2}\left(\tfrac{3}{4}\right) = \tfrac{117}{8} = 14.625$$

(b) $V = \iint_R (1 + x^2 + 3y)\, dA \approx \sum_{i=1}^{2} \sum_{j=1}^{2} f\left(\overline{x}_i, \overline{y}_j\right) \Delta A$

$$= f\left(\tfrac{5}{4}, \tfrac{3}{4}\right) \Delta A + f\left(\tfrac{5}{4}, \tfrac{9}{4}\right) \Delta A + f\left(\tfrac{7}{4}, \tfrac{3}{4}\right) \Delta A + f\left(\tfrac{7}{4}, \tfrac{9}{4}\right) \Delta A$$

$$= \tfrac{77}{16}\left(\tfrac{3}{4}\right) + \tfrac{149}{16}\left(\tfrac{3}{4}\right) + \tfrac{101}{16}\left(\tfrac{3}{4}\right) + \tfrac{173}{16}\left(\tfrac{3}{4}\right) = \tfrac{375}{16} = 23.4375$$

5. The values of $f(x,y) = \sqrt{52 - x^2 - y^2}$ get smaller as we move farther from the origin, so on any of the subrectangles in the problem, the function will have its largest value at the lower left corner of the subrectangle and its smallest value at the upper right corner, and any other value will lie between these two. So using these subrectangles we have $U < V < L$. (Note that this is true no matter how R is divided into subrectangles.)

6. To approximate the volume, let R be the planar region corresponding to the surface of the water in the pool, and place R on coordinate axes so that x and y correspond to the dimensions given. Then we define $f(x,y)$ to be the depth of the water at (x,y), so the volume of water in the pool is the volume of the solid that lies above the rectangle $R = [0,20] \times [0,30]$ and below the graph of $f(x,y)$. We can estimate this volume using the Midpoint Rule with $m = 2$ and $n = 3$, so $\Delta A = 100$. Each subrectangle with its midpoint is shown in the figure. Then

$$V \approx \sum_{i=1}^{2} \sum_{j=1}^{3} f\left(\overline{x}_i, \overline{y}_j\right) \Delta A = \Delta A[f(5,5) + f(5,15) + f(5,25) + f(15,5) + f(15,15) + f(15,25)]$$

$$= 100(3 + 7 + 10 + 3 + 5 + 8) = 3600$$

Thus, we estimate that the pool contains 3600 cubic feet of water.

Alternatively, we can approximate the volume with a Riemann sum where $m = 4$, $n = 6$ and the sample points are taken to be, for example, the upper right corner of each subrectangle. Then $\Delta A = 25$ and

$$V \approx \sum_{i=1}^{4} \sum_{j=1}^{6} f(x_i, y_j)\, \Delta A$$

$$= 25[3 + 4 + 7 + 8 + 10 + 8 + 4 + 6 + 8 + 10 + 12 + 10 + 3 + 4 + 5 + 6 + 8 + 7 + 2 + 2 + 2 + 3 + 4 + 4]$$

$$= 25(140) = 3500$$

So we estimate that the pool contains 3500 ft^3 of water.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. (a) With $m = n = 2$, we have $\Delta A = 4$. Using the contour map to estimate the value of f at the center of each subrectangle, we have

$$\iint_R f(x,y)\ dA \approx \sum_{i=1}^{2}\sum_{j=1}^{2} f\left(\overline{x}_i, \overline{y}_j\right)\Delta A = \Delta A[f(1,1)+f(1,3)+f(3,1)+f(3,3)] \approx 4(27+4+14+17) = 248$$

(b) $f_{\text{avg}} = \frac{1}{A(R)}\iint_R f(x,y)\,dA \approx \frac{1}{16}(248) = 15.5$

8. As in Example 9, we place the origin at the southwest corner of the state. Then $R = [0,388]\times[0,276]$ (in miles) is the rectangle corresponding to Colorado and we define $f(x,y)$ to be the temperature at the location (x,y). The average temperature is given by

$$f_{\text{avg}} = \frac{1}{A(R)}\iint_R f(x,y)\,dA = \frac{1}{388\cdot 276}\iint_R f(x,y)\,dA$$

To use the Midpoint Rule with $m = n = 4$, we divide R into 16 regions of equal size, as shown in the figure, with the center of each subrectangle indicated.

The area of each subrectangle is $\Delta A = \frac{388}{4}\cdot\frac{276}{4} = 6693$, so using the contour map to estimate the function values at each midpoint, we have

$$\begin{aligned}\iint_R f(x,y)\,dA &\approx \sum_{i=1}^{4}\sum_{j=1}^{4} f\left(\overline{x}_i, \overline{y}_j\right)\Delta A\\ &\approx \Delta A\,[31+28+52+43+43+25+57+46+36+20+42+45+30+23+43+41]\\ &= 6693(605)\end{aligned}$$

Therefore, $f_{\text{avg}} \approx \dfrac{6693\cdot 605}{388\cdot 276} \approx 37.8$, so the average temperature in Colorado at 4:00 PM on a day in February was approximately $37.8°\text{F}$.

9. $z = \sqrt{2} > 0$, so we can interpret the double integral as the volume of the solid S that lies below the plane $z = \sqrt{2}$ and above the rectangle $[2,6]\times[-1,5]$. S is a rectangular solid, so $\iint_R \sqrt{2}\,dA = 4\cdot 6\cdot\sqrt{2} = 24\sqrt{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $z = 2x + 1 \geq 0$ for $0 \leq x \leq 2$, so we can interpret the integral as the volume of the solid S that lies below the plane $z = 2x + 1$ and above the rectangle $[0, 2] \times [0, 4]$. We can picture S as a rectangular solid (with height 1) surmounted by a triangular cylinder; thus

$$\iint_R (2x + 1)\, dA = (2)(4)(1) + \tfrac{1}{2}(2)(4)(4) = 24$$

11. $z = 4 - 2y \geq 0$ for $0 \leq y \leq 1$, so we can interpret the integral as the volume of the solid S that lies below the plane $z = 4 - 2y$ and above the square $[0, 1] \times [0, 1]$. We can picture S as a rectangular solid (with height 2) surmounted by a triangular cylinder; thus

$$\iint_R (4 - 2y)\, dA = (1)(1)(2) + \tfrac{1}{2}(1)(1)(2) = 3$$

12. Here $z = \sqrt{9 - y^2}$, so $z^2 + y^2 = 9$, $z \geq 0$. Thus the integral represents the volume of the top half of the part of the circular cylinder $z^2 + y^2 = 9$ that lies above the rectangle $[0, 4] \times [0, 2]$.

13. $\int_0^2 (x + 3x^2y^2)\, dx = \left[\dfrac{x^2}{2} + 3\,\dfrac{x^3}{3}\,y^2\right]_{x=0}^{x=2} = \left[\tfrac{1}{2}x^2 + x^3y^2\right]_{x=0}^{x=2} = \left[\tfrac{1}{2}(2)^2 + (2)^3y^2\right] - \left[\tfrac{1}{2}(0)^2 + (0)^3y^2\right] = 2 + 8y^2,$

$\int_0^3 (x + 3x^2y^2)\, dy = \left[xy + 3x^2\,\dfrac{y^3}{3}\right]_{y=0}^{y=3} = \left[xy + x^2y^3\right]_{y=0}^{y=3} = \left[x(3) + x^2(3)^3\right] - \left[x(0) + x^2(0)^3\right] = 3x + 27x^2$

14. $\int_0^2 y\sqrt{x+2}\, dx = \left[y \cdot \tfrac{2}{3}(x+2)^{3/2}\right]_{x=0}^{x=2} = \tfrac{2}{3}y(4)^{3/2} - \tfrac{2}{3}y(2)^{3/2} = \tfrac{16}{3}y - \tfrac{4}{3}\sqrt{2}\,y = \tfrac{4}{3}(4 - \sqrt{2}\,)\,y,$

$\int_0^3 y\sqrt{x+2}\, dy = \left[\dfrac{y^2}{2}\,\sqrt{x+2}\right]_{y=0}^{y=3} = \tfrac{1}{2}(3)^2\sqrt{x+2} - \tfrac{1}{2}(0)^2\sqrt{x+2} = \tfrac{9}{2}\sqrt{x+2}$

15. $\int_1^4 \int_0^2 (6x^2y - 2x)\, dy\, dx = \int_1^4 \left[3x^2y^2 - 2xy\right]_{y=0}^{y=2} dx = \int_1^4 \left[\left(12x^2 - 4x\right) - (0 - 0)\,\right] dx$

$= \int_1^4 (12x^2 - 4x)\, dx = \left[4x^3 - 2x^2\right]_1^4 = (256 - 32) - (4 - 2) = 222$

16. $\int_0^1 \int_0^1 (x + y)^2\, dx\, dy = \int_0^1 \int_0^1 (x^2 + 2xy + y^2)\, dx\, dy = \int_0^1 \left[\tfrac{1}{3}x^3 + x^2y + xy^2\right]_{x=0}^{x=1} dy$

$= \int_0^1 (\tfrac{1}{3} + y + y^2)\, dy = \left[\tfrac{1}{3}y + \tfrac{1}{2}y^2 + \tfrac{1}{3}y^3\right]_0^1 = \tfrac{1}{3} + \tfrac{1}{2} + \tfrac{1}{3} - 0 = \tfrac{7}{6}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\int_0^1 \int_1^2 (x + e^{-y})\,dx\,dy = \int_0^1 \left[\frac{1}{2}x^2 + xe^{-y}\right]_{x=1}^{x=2} dy = \int_0^1 \left[(2 + 2e^{-y}) - (\frac{1}{2} + e^{-y})\right] dy$

$= \int_0^1 (\frac{3}{2} + e^{-y})\,dy = \left[\frac{3}{2}y - e^{-y}\right]_0^1 = \left(\frac{3}{2} - e^{-1}\right) - (0 - 1) = \frac{5}{2} - e^{-1}$

18. $\int_{-3}^1 \int_1^2 \left(x^2 + y^{-2}\right)\,dy\,dx = \int_{-3}^1 \left[x^2 y - y^{-1}\right]_{y=1}^{y=2} dx = \int_{-3}^1 \left[\left(2x^2 - \frac{1}{2}\right) - (x^2 - 1)\right] dx$

$= \int_{-3}^1 \left(x^2 + \frac{1}{2}\right) dx = \left[\frac{1}{3}x^3 + \frac{1}{2}x\right]_{-3}^1 = \left(\frac{1}{3} + \frac{1}{2}\right) - \left(-\frac{27}{3} - \frac{3}{2}\right) = \frac{34}{3}$

19. $\int_{-3}^3 \int_0^{\pi/2} (y + y^2 \cos x)\,dx\,dy = \int_{-3}^3 \left[xy + y^2 \sin x\right]_{x=0}^{x=\pi/2} dy = \int_{-3}^3 \left(\frac{\pi}{2}y + y^2\right) dy$

$= \left[\frac{\pi}{4}y^2 + \frac{1}{3}y^3\right]_{-3}^3 = \left[\left(\frac{9\pi}{4} + 9\right) - \left(\frac{9\pi}{4} - 9\right)\right] = 18$

20. $\displaystyle\int_1^3 \int_1^5 \frac{\ln y}{xy}\,dy\,dx = \int_1^3 \frac{1}{x}\,dx \int_1^5 \frac{\ln y}{y}\,dy$ [by Equation 11]

$= \Big[\ln|x|\Big]_1^3 \left[\frac{1}{2}(\ln y)^2\right]_1^5$ [substitute $u = \ln y \quad \Rightarrow \quad du = (1/y)\,dy$]

$= (\ln 3 - 0) \cdot \frac{1}{2}[(\ln 5)^2 - 0] = \frac{1}{2}(\ln 3)(\ln 5)^2$

21. $\displaystyle\int_1^4 \int_1^2 \left(\frac{x}{y} + \frac{y}{x}\right) dy\,dx = \int_1^4 \left[x \ln|y| + \frac{1}{x} \cdot \frac{1}{2}\,y^2\right]_{y=1}^{y=2} dx = \int_1^4 \left(x \ln 2 + \frac{3}{2x}\right) dx = \left[\frac{1}{2}x^2 \ln 2 + \frac{3}{2}\ln|x|\right]_1^4$

$= \left(8 \ln 2 + \frac{3}{2}\ln 4\right) - \left(\frac{1}{2}\ln 2 + 0\right) = \frac{15}{2}\ln 2 + \frac{3}{2}\ln 4$ or $\frac{15}{2}\ln 2 + 3\ln(4^{1/2}) = \frac{21}{2}\ln 2$

22. $\int_0^1 \int_0^2 ye^{x-y}\,dx\,dy = \int_0^1 \int_0^2 ye^x e^{-y}\,dx\,dy = \int_0^2 e^x\,dx \int_0^1 ye^{-y}\,dy$ [by Equation 11]

$= [e^x]_0^2 \left[(-y - 1)e^{-y}\right]_0^1$ [by integrating by parts]

$= \left(e^2 - e^0\right)\left[-2e^{-1} - \left(-e^0\right)\right] = (e^2 - 1)(1 - 2e^{-1})$ or $e^2 - 2e + 2e^{-1} - 1$

23. $\int_0^3 \int_0^{\pi/2} t^2 \sin^3\phi\,d\phi\,dt = \int_0^{\pi/2} \sin^3\phi\,d\phi \int_0^3 t^2\,dt$ [by Equation 11] $= \int_0^{\pi/2} (1 - \cos^2\phi)\sin\phi\,d\phi \int_0^3 t^2\,dt$

$= \left[\frac{1}{3}\cos^3\phi - \cos\phi\right]_0^{\pi/2} \left[\frac{1}{3}t^3\right]_0^3 = \left[(0 - 0) - \left(\frac{1}{3} - 1\right)\right] \cdot \frac{1}{3}(27 - 0) = \frac{2}{3}(9) = 6$

24. $\int_0^1 \int_0^1 xy\sqrt{x^2 + y^2}\,dy\,dx = \int_0^1 x\left[\frac{1}{3}(x^2 + y^2)^{3/2}\right]_{y=0}^{y=1} dx = \frac{1}{3}\int_0^1 x[(x^2 + 1)^{3/2} - x^3]\,dx = \frac{1}{3}\int_0^1 [x(x^2 + 1)^{3/2} - x^4]dx$

$= \frac{1}{3}\left[\frac{1}{5}(x^2 + 1)^{5/2} - \frac{1}{5}x^5\right]_0^1 = \frac{1}{15}\left[(2^{5/2} - 1) - (1 - 0)\right] = \frac{2}{15}\left(2\sqrt{2} - 1\right)$

25. $\int_0^1 \int_0^1 v(u + v^2)^4\,du\,dv = \int_0^1 \left[\frac{1}{5}v(u + v^2)^5\right]_{u=0}^{u=1} dv = \frac{1}{5}\int_0^1 v\left[(1 + v^2)^5 - (0 + v^2)^5\right] dv$

$= \frac{1}{5}\int_0^1 \left[v(1 + v^2)^5 - v^{11}\right] dv = \frac{1}{5}\left[\frac{1}{2} \cdot \frac{1}{6}(1 + v^2)^6 - \frac{1}{12}v^{12}\right]_0^1$

[substitute $t = 1 + v^2 \quad \Rightarrow \quad dt = 2v\,dv$ in the first term]

$= \frac{1}{60}\left[(2^6 - 1) - (1 - 0)\right] = \frac{1}{60}(63 - 1) = \frac{31}{30}$

26. $\int_0^1 \int_0^1 \sqrt{s + t}\,ds\,dt = \int_0^1 \left[\frac{2}{3}(s + t)^{3/2}\right]_{s=0}^{s=1} dt = \frac{2}{3}\int_0^1 [(1 + t)^{3/2} - t^{3/2}]\,dt = \frac{2}{3}\left[\frac{2}{5}(1 + t)^{5/2} - \frac{2}{5}t^{5/2}\right]_0^1$

$= \frac{4}{15}[(2^{5/2} - 1) - (1 - 0)] = \frac{4}{15}(2^{5/2} - 2)$ or $\frac{8}{15}\left(2\sqrt{2} - 1\right)$

27. $\iint_R x \sec^2 y\,dA = \int_0^2 \int_0^{\pi/4} x \sec^2 y\,dy\,dx = \int_0^2 x\,dx \int_0^{\pi/4} \sec^2 y\,dy = \left[\frac{1}{2}x^2\right]_0^2 \Big[\tan y\Big]_0^{\pi/4}$

$= (2 - 0)\left(\tan\frac{\pi}{4} - \tan 0\right) = 2(1 - 0) = 2$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $\iint_R (y + xy^{-2})\,dA = \int_1^2 \int_0^2 (y + xy^{-2})\,dx\,dy = \int_1^2 \left[xy + \frac{1}{2}x^2y^{-2}\right]_{x=0}^{x=2} dy = \int_1^2 \left(2y + 2y^{-2}\right) dy$

$= \left[y^2 - 2y^{-1}\right]_1^2 = (4-1) - (1-2) = 4$

29. $\displaystyle\iint_R \frac{xy^2}{x^2+1}\,dA = \int_0^1 \int_{-3}^3 \frac{xy^2}{x^2+1}\,dy\,dx = \int_0^1 \frac{x}{x^2+1}\,dx \int_{-3}^3 y^2\,dy = \left[\tfrac{1}{2}\ln(x^2+1)\right]_0^1 \left[\tfrac{1}{3}y^3\right]_{-3}^3$

$= \frac{1}{2}(\ln 2 - \ln 1)\cdot\frac{1}{3}(27+27) = 9\ln 2$

30. $\displaystyle\iint_R \frac{\tan\theta}{\sqrt{1-t^2}}\,dA = \int_0^{1/2}\int_0^{\pi/3} \frac{\tan\theta}{\sqrt{1-t^2}}\,d\theta\,dt = \int_0^{1/2} \frac{1}{\sqrt{1-t^2}}\,dt \int_0^{\pi/3} \tan\theta\,d\theta = \left[\sin^{-1} t\right]_0^{1/2} \left[\ln|\sec\theta|\right]_0^{\pi/3}$

$= \left(\sin^{-1}\frac{1}{2} - \sin^{-1} 0\right)\left(\ln\left|\sec\frac{\pi}{3}\right| - \ln|\sec 0|\right) = \left(\frac{\pi}{6} - 0\right)(\ln 2 - \ln 1) = \frac{\pi}{6}\ln 2$

31. $\iint_R x\sin(x+y)\,dA = \int_0^{\pi/6}\int_0^{\pi/3} x\sin(x+y)\,dy\,dx$

$= \int_0^{\pi/6} \left[-x\cos(x+y)\right]_{y=0}^{y=\pi/3} dx = \int_0^{\pi/6} \left[x\cos x - x\cos\left(x+\frac{\pi}{3}\right)\right] dx$

$= x\left[\sin x - \sin\left(x+\frac{\pi}{3}\right)\right]_0^{\pi/6} - \int_0^{\pi/6} \left[\sin x - \sin\left(x+\frac{\pi}{3}\right)\right] dx$ $\left[\begin{array}{l}\text{by integrating by parts}\\ \text{separately for each term}\end{array}\right]$

$= \frac{\pi}{6}\left[\frac{1}{2} - 1\right] - \left[-\cos x + \cos\left(x+\frac{\pi}{3}\right)\right]_0^{\pi/6} = -\frac{\pi}{12} - \left[-\frac{\sqrt{3}}{2} + 0 - \left(-1 + \frac{1}{2}\right)\right] = \frac{\sqrt{3}-1}{2} - \frac{\pi}{12}$

32. $\displaystyle\iint_R \frac{x}{1+xy}\,dA = \int_0^1\int_0^1 \frac{x}{1+xy}\,dy\,dx = \int_0^1 \left[\ln(1+xy)\right]_{y=0}^{y=1} dx = \int_0^1 \left[\ln(1+x) - \ln 1\right] dx$

$= \int_0^1 \ln(1+x)\,dx = \left[(1+x)\ln(1+x) - x\right]_0^1$ [by integrating by parts]

$= (2\ln 2 - 1) - (\ln 1 - 0) = 2\ln 2 - 1$

33. $\iint_R ye^{-xy}\,dA = \int_0^3\int_0^2 ye^{-xy}\,dx\,dy = \int_0^3 \left[-e^{-xy}\right]_{x=0}^{x=2} dy = \int_0^3 (-e^{-2y} + 1)\,dy = \left[\frac{1}{2}e^{-2y} + y\right]_0^3$

$= \frac{1}{2}e^{-6} + 3 - \left(\frac{1}{2} + 0\right) = \frac{1}{2}e^{-6} + \frac{5}{2}$

34. $\displaystyle\iint_R \frac{1}{1+x+y}\,dA = \int_1^3\int_1^2 \frac{1}{1+x+y}\,dy\,dx = \int_1^3 \left[\ln(1+x+y)\right]_{y=1}^{y=2} dx = \int_1^3 \left[\ln(x+3) - \ln(x+2)\right] dx$

$= \left[\big((x+3)\ln(x+3) - (x+3)\big) - \big((x+2)\ln(x+2) - (x+2)\big)\right]_1^3$

[by integrating by parts separately for each term]

$= (6\ln 6 - 6 - 5\ln 5 + 5) - (4\ln 4 - 4 - 3\ln 3 + 3) = 6\ln 6 - 5\ln 5 - 4\ln 4 + 3\ln 3$

35. $z = f(x,y) = 4 - x - 2y \geq 0$ for $0 \leq x \leq 1$ and $0 \leq y \leq 1$. So the solid is the region in the first octant which lies below the plane $z = 4 - x - 2y$ and above $[0,1]\times[0,1]$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. $z = 2 - x^2 - y^2 \geq 0$ for $0 \leq x \leq 1$ and $0 \leq y \leq 1$. So the solid is the region in the first octant which lies below the circular paraboloid $z = 2 - x^2 - y^2$ and above $[0, 1] \times [0, 1]$.

37. $z = 4 - x^2 \geq 0$ for $-2 \leq x \leq 2$ and $-1 \leq y \leq 3$. So the solid is the region that lies below the parabolic cylinder $z = 4 - x^2$ and above $[-2, 2] \times [-1, 3]$.

38. (a) For any given value of $x = k$, $0 \leq k \leq 2$, we have the curve $z = k^2\sqrt{y}$, $x = k$, $1 \leq y \leq 4$.

The area below the curve and above xy-plane is given by $\displaystyle\int_1^4 k^2\sqrt{y}\,dy = \left[\frac{2k^2}{3}y^{3/2}\right]_{y=1}^{y=4} = \frac{2k^2}{3}(8-1) = \frac{14k^2}{3}$.

For $k = 1$, the area is $\dfrac{14(1^2)}{3} = \dfrac{14}{3}$. For $k = 2$, the area is $\dfrac{14(2^2)}{3} = \dfrac{56}{3}$.

(b) For any given value of $y = k$, $1 \leq k \leq 4$, we have the curve, $z = x^2\sqrt{k}$, $y = k$, $0 \leq x \leq 2$. The area below the curve and above xy-plane is given by $\displaystyle\int_0^2 x^2\sqrt{k}\,dx = \left[\frac{x^3}{3}\sqrt{k}\right]_{x=0}^{x=2} = \frac{8\sqrt{k}}{3}$. For $k = 1$, the area is $\dfrac{8\sqrt{1}}{3} = \dfrac{8}{3}$. For $k = 3$, the area is $\dfrac{8\sqrt{3}}{3}$.

(c) $\displaystyle\int_1^4\int_0^2 x^2\sqrt{y}\,dx\,dy = \int_1^4 \left[\frac{x^3}{3}\sqrt{y}\right]_{x=0}^{x=2} dy = \int_1^4 \frac{8}{3}\sqrt{y}\,dy = \left[\frac{16}{9}y^{3/2}\right]_1^4 = \frac{16}{9}(8-1) = \frac{112}{9}$

39. (a) The volume under the surface $z = xy$ and over the square $R = [0, 2] \times [0, 2]$ is given by $\int_0^2\int_0^2 xy\,dx\,dy$.

(b) $\displaystyle\int_0^2\int_0^2 xy\,dx\,dy = \int_0^2 \left[\frac{x^2}{2}\right]_{x=0}^{x=2} y\,dy = \int_0^2 \left[\frac{2^2}{2} - \frac{0^2}{2}\right] y\,dy = \int_0^2 2y\,dy = \left[y^2\right]_0^2 = 4$

40. (a) The volume under the surface $z = \cos x \cos y$ and over the square $R = \left[-\frac{\pi}{4}, \frac{\pi}{4}\right] \times \left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$ is given by $\int_{-\pi/4}^{\pi/4}\int_{-\pi/4}^{\pi/4} \cos x \cos y\,dx\,dy$.

(b) $\int_{-\pi/4}^{\pi/4}\int_{-\pi/4}^{\pi/4} \cos x \cos y\,dx\,dy = \int_{-\pi/4}^{\pi/4} \cos x\,dx \int_{-\pi/4}^{\pi/4} \cos y\,dy$ [by Equation 11]

$$= \left[\sin x\right]_{x=-\pi/4}^{x=\pi/4} \left[\sin y\right]_{y=-\pi/4}^{y=\pi/4} = \left[\sin\tfrac{\pi}{4} - \sin\left(-\tfrac{\pi}{4}\right)\right]^2$$

$$= \left[\tfrac{\sqrt{2}}{2} - \left(-\tfrac{\sqrt{2}}{2}\right)\right]^2 = \left(\sqrt{2}\right)^2 = 2$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. (a) The volume under the surface $z = 1 + ye^{xy}$ and over the rectangle $R = [0,1] \times [1,2]$ is given by

$\int_1^2 \int_0^1 \left(1 + ye^{xy}\right)\, dx\, dy.$

(b) $\displaystyle\int_1^2 \int_0^1 (1 + ye^{xy})\, dx\, dy = \int_1^2 \left[x + y \cdot \frac{1}{y} e^{xy}\right]_{x=0}^{x=1} dy = \int_1^2 \left[(1 + e^y) - (0 + 1)\right] dy = \int_1^2 e^y\, dy = e^2 - e$

42. (a) The volume under the surface $z = x^2 + y^2$ and over the square $R = [1,2] \times [1,3]$ is given by $\int_1^3 \int_1^2 (x^2 + y^2)\, dx\, dy$.

(b) $\displaystyle\int_1^3 \int_1^2 (x^2 + y^2)\, dx\, dy = \int_1^3 \left[\frac{x^3}{3} + xy^2\right]_{x=1}^{x=2} dy = \int_1^3 \left[\left(\frac{8}{3} + 2y^2\right) - \left(\frac{1}{3} + y^2\right)\right] dy = \int_1^3 \left(\frac{7}{3} + y^2\right) dy$

$$= \left[\frac{7}{3}y + \frac{y^3}{3}\right]_1^3 = (7 + 9) - \left(\frac{7}{3} + \frac{1}{3}\right) = \frac{40}{3}$$

43. The solid lies under the plane $4x + 6y - 2z + 15 = 0$ or $z = 2x + 3y + \frac{15}{2}$ so

$V = \iint_R (2x + 3y + \frac{15}{2})\, dA = \int_{-1}^1 \int_{-1}^2 (2x + 3y + \frac{15}{2})\, dx\, dy = \int_{-1}^1 \left[x^2 + 3xy + \frac{15}{2}x\right]_{x=-1}^{x=2} dy$

$= \int_{-1}^1 \left[(19 + 6y) - (-\frac{13}{2} - 3y)\right] dy = \int_{-1}^1 (\frac{51}{2} + 9y)\, dy = \left[\frac{51}{2}y + \frac{9}{2}y^2\right]_{-1}^1 = 30 - (-21) = 51$

44. $V = \iint_R (3y^2 - x^2 + 2)\, dA = \int_{-1}^1 \int_1^2 (3y^2 - x^2 + 2)\, dy\, dx = \int_{-1}^1 \left[y^3 - x^2 y + 2y\right]_{y=1}^{y=2} dx$

$= \int_{-1}^1 \left[(12 - 2x^2) - (3 - x^2)\right] dx = \int_{-1}^1 \left(9 - x^2 dx\right) = \left[9x - \frac{1}{3}x^3\right]_{-1}^1 = \frac{26}{3} + \frac{26}{3} = \frac{52}{3}$

45. $V = \int_{-2}^2 \int_{-1}^1 \left(1 - \frac{1}{4}x^2 - \frac{1}{9}y^2\right) dx\, dy = 4 \int_0^2 \int_0^1 \left(1 - \frac{1}{4}x^2 - \frac{1}{9}y^2\right) dx\, dy$

$= 4 \int_0^2 \left[x - \frac{1}{12}x^3 - \frac{1}{9}y^2 x\right]_{x=0}^{x=1} dy = 4 \int_0^2 \left(\frac{11}{12} - \frac{1}{9}y^2\right) dy = 4\left[\frac{11}{12}y - \frac{1}{27}y^3\right]_0^2 = 4 \cdot \frac{83}{54} = \frac{166}{27}$

46. The solid lies under the surface $z = x^2 + xy^2$ and above the rectangle $R = [0,5] \times [-2,2]$, so its volume is

$$V = \iint_R (x^2 + xy^2)\, dA = \int_0^5 \int_{-2}^2 (x^2 + xy^2)\, dy\, dx = \int_0^5 \left[x^2 y + \frac{1}{3}xy^3\right]_{y=-2}^{y=2} dx$$

$$= \int_0^5 \left[\left(2x^2 + \frac{8}{3}x\right) - \left(-2x^2 - \frac{8}{3}x\right)\right] dx = \int_0^5 (4x^2 + \frac{16}{3}x)\, dx$$

$$= \left[\frac{4}{3}x^3 + \frac{8}{3}x^2\right]_0^5 = \frac{500}{3} + \frac{200}{3} - 0 = \frac{700}{3}$$

47. The solid lies under the surface $z = 1 + x^2 ye^y$ and above the rectangle $R = [-1,1] \times [0,1]$, so its volume is

$$V = \iint_R (1 + x^2 ye^y)\, dA = \int_0^1 \int_{-1}^1 (1 + x^2 ye^y)\, dx\, dy = \int_0^1 \left[x + \frac{1}{3}x^3 ye^y\right]_{x=-1}^{x=1} dy$$

$$= \int_0^1 (2 + \frac{2}{3}ye^y)\, dy = \left[2y + \frac{2}{3}(y - 1)e^y\right]_0^1 \qquad \text{[by integrating by parts in the second term]}$$

$$= (2 + 0) - \left(0 - \frac{2}{3}e^0\right) = 2 + \frac{2}{3} = \frac{8}{3}$$

48. The cylinder $z = 16 - x^2$ intersects the xy-plane along the line $x = 4$, so in the first octant, the solid lies below the surface $z = 16 - x^2$ and above the rectangle $R = [0,4] \times [0,5]$ in the xy-plane.

$$V = \int_0^5 \int_0^4 (16 - x^2)\, dx\, dy = \int_0^4 (16 - x^2)\, dx \int_0^5 dy$$

$$= \left[16x - \frac{1}{3}x^3\right]_0^4 \left[y\right]_0^5 = (64 - \frac{64}{3} - 0)(5 - 0) = \frac{640}{3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. The solid lies below the surface $z = 2 + x^2 + (y-2)^2$ and above the plane $z = 1$ for $-1 \le x \le 1$, $0 \le y \le 4$. The volume of the solid is the difference in volumes between the solid that lies under $z = 2 + x^2 + (y-2)^2$ over the rectangle $R = [-1,1] \times [0,4]$ and the solid that lies under $z = 1$ over R.

$$\begin{aligned} V &= \textstyle\int_0^4 \int_{-1}^1 [2 + x^2 + (y-2)^2]\, dx\, dy - \int_0^4 \int_{-1}^1 (1)\, dx\, dy \\ &= \textstyle\int_0^4 \left[2x + \frac{1}{3}x^3 + x(y-2)^2\right]_{x=-1}^{x=1} dy - \int_{-1}^1 dx \int_0^4 dy \\ &= \textstyle\int_0^4 \left[(2 + \frac{1}{3} + (y-2)^2) - (-2 - \frac{1}{3} - (y-2)^2)\right] dy - [x]_{-1}^1\ [y]_0^4 \\ &= \textstyle\int_0^4 \left[\frac{14}{3} + 2(y-2)^2\right] dy - [1-(-1)][4-0] = \left[\frac{14}{3}y + \frac{2}{3}(y-2)^3\right]_0^4 - (2)(4) \\ &= \left[\left(\frac{56}{3} + \frac{16}{3}\right) - \left(0 - \frac{16}{3}\right)\right] - 8 = \frac{88}{3} - 8 = \frac{64}{3} \end{aligned}$$

50.

The solid lies below the plane $z = x + 2y$ and above the surface $z = \dfrac{2xy}{x^2+1}$ for $0 \le x \le 2$, $0 \le y \le 4$. The volume of the solid is the difference in volumes between the solid that lies under $z = x + 2y$ over the rectangle $R = [0,2] \times [0,4]$ and the solid that lies under $z = \dfrac{2xy}{x^2+1}$ over R.

$$\begin{aligned} V &= \int_0^2 \int_0^4 (x + 2y)dy\, dx - \int_0^2 \int_0^4 \frac{2xy}{x^2+1}\, dy\, dx = \int_0^2 \left[xy + y^2\right]_{y=0}^{y=4} dx - \int_0^2 \frac{2x}{x^2+1}\, dx \int_0^4 y\, dy \\ &= \textstyle\int_0^2 [(4x+16) - (0+0)]\, dx - \left[\ln\left|x^2+1\right|\right]_0^2 \left[\frac{1}{2}y^2\right]_0^4 = \left[2x^2 + 16x\right]_0^2 - (\ln 5 - \ln 1)(8-0) \\ &= (8 + 32 - 0) - 8\ln 5 = 40 - 8\ln 5 \end{aligned}$$

51. In Maple, we can calculate the integral by defining the integrand as `f` and then using the command `int(int(f,x=0..1),y=0..1);`. In Mathematica, we can use the command

```
Integrate[f,{x,0,1},{y,0,1}]
```

We find that $\iint_R x^5 y^3 e^{xy}\, dA = 21e - 57 \approx 0.0839$. We can use `plot3d` (in Maple) or `Plot3D` (in Mathematica) to graph the function.

52. In Maple, we can calculate the integral by defining `f:=exp(-x^2)*cos(x^2+y^2);` and `g:=2-x^2-y^2;` and then [since $2 - x^2 - y^2 > e^{-x^2}\cos(x^2+y^2)$ for $-1 \le x \le 1$, $-1 \le y \le 1$] using the command `evalf(Int(Int(g-f,x=-1..1),y=-1..1));`. Using `Int` rather than `int` forces Maple to use purely numerical techniques in evaluating the integral.

In Mathematica, we can use the command `NIntegrate[g-f,{x,-1,1},{y,-1,1}]`. We find that

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\iint_R \left[(2-x^2-y^2)-\left(e^{-x^2}\cos(x^2+y^2)\right)\right] dA \approx 3.0271$. We can use the `plot3d` command (in Maple) or `Plot3D` (in Mathematica) to graph both functions on the same screen.

53. R is the rectangle $[-1,1]\times[0,5]$. Thus, $A(R)=2\cdot 5=10$ and

$f_{\text{avg}} = \dfrac{1}{A(R)}\iint_R f(x,y)\,dA = \frac{1}{10}\int_0^5\int_{-1}^1 x^2y\,dx\,dy = \frac{1}{10}\int_0^5 \left[\frac{1}{3}x^3y\right]_{x=-1}^{x=1} dy = \frac{1}{10}\int_0^5 \frac{2}{3}y\,dy = \frac{1}{10}\left[\frac{1}{3}y^2\right]_0^5 = \frac{5}{6}$.

54. $A(R)=4\cdot 1=4$, so

$$\begin{aligned} f_{\text{avg}} &= \frac{1}{A(R)}\iint_R f(x,y)\,dA = \tfrac{1}{4}\int_0^4\int_0^1 e^y\sqrt{x+e^y}\,dy\,dx = \tfrac{1}{4}\int_0^4 \left[\tfrac{2}{3}(x+e^y)^{3/2}\right]_{y=0}^{y=1} dx \\ &= \tfrac{1}{4}\cdot\tfrac{2}{3}\int_0^4[(x+e)^{3/2}-(x+1)^{3/2}]\,dx = \tfrac{1}{6}\left[\tfrac{2}{5}(x+e)^{5/2}-\tfrac{2}{5}(x+1)^{5/2}\right]_0^4 \\ &= \tfrac{1}{6}\cdot\tfrac{2}{5}[(4+e)^{5/2}-5^{5/2}-e^{5/2}+1] = \tfrac{1}{15}[(4+e)^{5/2}-e^{5/2}-5^{5/2}+1] \approx 3.327 \end{aligned}$$

55. $\iint_R \dfrac{xy}{1+x^4}\,dA = \displaystyle\int_{-1}^1\int_0^1 \frac{xy}{1+x^4}\,dy\,dx = \int_{-1}^1 \frac{x}{1+x^4}\,dx\int_0^1 y\,dy$ [by Equation 11] but $f(x)=\dfrac{x}{1+x^4}$ is an odd function so $\displaystyle\int_{-1}^1 f(x)\,dx=0$ (by Theorem 5.5.7). Thus $\displaystyle\iint_R \frac{xy}{1+x^4}\,dA = 0\cdot\int_0^1 y\,dy = 0$.

56. $$\begin{aligned} \iint_R(1+x^2\sin y+y^2\sin x)\,dA &= \iint_R 1\,dA + \iint_R x^2\sin y\,dA + \iint_R y^2\sin x\,dA \\ &= A(R) + \int_{-\pi}^{\pi}\int_{-\pi}^{\pi} x^2\sin y\,dy\,dx + \int_{-\pi}^{\pi}\int_{-\pi}^{\pi} y^2\sin x\,dy\,dx \\ &= (2\pi)(2\pi) + \int_{-\pi}^{\pi} x^2dx\int_{-\pi}^{\pi}\sin y\,dy + \int_{-\pi}^{\pi}\sin x\,dx\int_{-\pi}^{\pi} y^2\,dy \end{aligned}$$

But $\sin x$ is an odd function, so $\int_{-\pi}^{\pi}\sin x\,dx = \int_{-\pi}^{\pi}\sin y\,dy = 0$ (by Theorem 5.5.7) and $\iint_R(1+x^2\sin y+y^2\sin x)\,dA = 4\pi^2+0+0=4\pi^2$.

57. Let $f(x,y)=\dfrac{x-y}{(x+y)^3}$. Then a CAS gives $\int_0^1\int_0^1 f(x,y)\,dy\,dx = \frac{1}{2}$ and $\int_0^1\int_0^1 f(x,y)\,dx\,dy = -\frac{1}{2}$.

To explain the seeming violation of Fubini's Theorem, note that f has an infinite discontinuity at $(0,0)$ and thus does not satisfy the conditions of Fubini's Theorem. In fact, both iterated integrals involve improper integrals that diverge at their lower limits of integration.

58. (a) Loosely speaking, Fubini's Theorem says that the order of integration of a function of two variables does not affect the value of the double integral, while Clairaut's Theorem says that the order of differentiation of such a function does not affect the value of the second-order derivative. Also, both theorems require continuity (though Fubini's allows a finite number of smooth curves to contain discontinuities).

(b) To find g_{xy}, we first hold y constant and use the single-variable Fundamental Theorem of Calculus, Part 1:

$g_x = \dfrac{d}{dx}g(x,y) = \dfrac{d}{dx}\displaystyle\int_a^x\left(\int_c^y f(s,t)\,dt\right)ds = \int_c^y f(x,t)\,dt$. Now we use the Fundamental Theorem again:

$g_{xy} = \dfrac{d}{dy}\displaystyle\int_c^y f(x,t)\,dt = f(x,y)$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

To find g_{yx}, we first use Fubini's Theorem to find that $\int_a^x \int_c^y f(s,t)\,dt\,ds = \int_c^y \int_a^x f(s,t)\,dt\,ds$, and then use the Fundamental Theorem twice, as above, to get $g_{yx} = f(x,y)$. So $g_{xy} = g_{yx} = f(x,y)$.

15.2 Double Integrals over General Regions

1. $\int_1^5 \int_0^x (8x-2y)\,dy\,dx = \int_1^5 \left[8xy - y^2\right]_{y=0}^{y=x} dx = \int_1^5 [8x(x) - (x)^2 - 8x(0) + (0)^2]dx$

$= \int_1^5 7x^2\,dx = \frac{7}{3}x^3\Big]_1^5 = \frac{7}{3}(125-1) = \frac{868}{3}$

2. $\int_0^2 \int_0^{y^2} x^2 y\,dx\,dy = \int_0^2 \left[\frac{1}{3}x^3 y\right]_{x=0}^{x=y^2} dy = \int_0^2 \frac{1}{3}y\left[(y^2)^3 - (0)^3\right] dy$

$= \int_0^2 \frac{1}{3}y^7\,dy = \frac{1}{3}\left[\frac{1}{8}y^8\right]_0^2 = \frac{1}{3}(32-0) = \frac{32}{3}$

3. $\int_0^1 \int_0^y xe^{y^3}\,dx\,dy = \int_0^1 \left[\frac{1}{2}x^2 e^{y^3}\right]_{x=0}^{x=y} dy = \int_0^1 \frac{1}{2}e^{y^3}\left[(y)^2 - (0)^2\right] dy$

$= \frac{1}{2}\int_0^1 y^2 e^{y^3}\,dy = \frac{1}{2}\left[\frac{1}{3}e^{y^3}\right]_0^1 = \frac{1}{2}\cdot\frac{1}{3}\left(e^1 - e^0\right) = \frac{1}{6}(e-1)$

4. $\int_0^{\pi/2} \int_0^x x\sin y\,dy\,dx = \int_0^{\pi/2} [x(-\cos y)]_{y=0}^{y=x}\,dx = \int_0^{\pi/2} (-x\cos x + x)\,dx = \int_0^{\pi/2} (x - x\cos x)\,dx$

$= \left[\frac{1}{2}x^2 - (x\sin x + \cos x)\right]_0^{\pi/2}$ (by integrating by parts in the second term)

$= \left(\frac{1}{2}\cdot\frac{\pi^2}{4} - \frac{\pi}{2} - 0\right) - (0-0-1) = \frac{\pi^2}{8} - \frac{\pi}{2} + 1$

5. $\int_0^1 \int_0^{s^2} \cos(s^3)\,dt\,ds = \int_0^1 \left[t\cos(s^3)\right]_{t=0}^{t=s^2} ds = \int_0^1 s^2\cos(s^3)\,ds = \frac{1}{3}\sin(s^3)\Big]_0^1 = \frac{1}{3}(\sin 1 - \sin 0) = \frac{1}{3}\sin 1$

6. $\int_0^1 \int_0^{e^v} \sqrt{1+e^v}\,dw\,dv = \int_0^1 \left[w\sqrt{1+e^v}\right]_{w=0}^{w=e^v} dv = \int_0^1 e^v\sqrt{1+e^v}\,dv = \frac{2}{3}(1+e^v)^{3/2}\Big]_0^1$

$= \frac{2}{3}(1+e)^{3/2} - \frac{2}{3}(1+1)^{3/2} = \frac{2}{3}(1+e)^{3/2} - \frac{4}{3}\sqrt{2}$

7. (a) We express the iterated integral as a Type I: $\displaystyle\int_0^2 \int_x^{3x-x^2} 2y\,dy\,dx$. A Type II would require the sum of two integrals.

(b) $\displaystyle\int_0^2 \int_x^{3x-x^2} 2y\,dy\,dx = \int_0^2 \left[y^2\right]_{y=x}^{y=3x-x^2} dx = \int_0^2 \left[(3x-x^2)^2 - (x)^2\right] dx = \int_0^2 \left(8x^2 - 6x^3 + x^4\right) dx$

$= \left[\frac{8}{3}x^3 - \frac{3}{2}x^4 + \frac{1}{5}x^5\right]_0^2 = \frac{64}{3} - 24 + \frac{32}{5} = \frac{56}{15}$

8. (a) We express the iterated integral as a Type II: $\int_0^1 \int_y^{2-y} (x+y)\,dx\,dy$. A Type I would require the sum of two integrals.

(b) $\displaystyle\int_0^1 \int_y^{2-y} (x+y)\,dx\,dy = \int_0^1 \left[\frac{x^2}{2} + xy\right]_{x=y}^{x=2-y} dy = \int_0^1 \left[\left(\frac{(2-y)^2}{2} + (2-y)y\right) - \left(\frac{y^2}{2} + y^2\right)\right] dy$

$= \int_0^1 (2-2y^2)\,dy = \left[2y - \frac{2}{3}y^3\right]_0^1 = 2 - \frac{2}{3} = \frac{4}{3}$

9. (a) We express the iterated integral as a Type II. A Type I would require the sum of two integrals. The curves intersect when $\sqrt{x} = x-2 \;\Rightarrow\; x = x^2 - 4x + 4 \;\Leftrightarrow\; 0 = x^2 - 5x + 4 \;\Leftrightarrow\; (x-4)(x-1) = 0 \;\Leftrightarrow\; x = 1$ or $x = 4$. The point for $x = 1$ is not in D. Thus, the point of intersection of the curves is $(4,2)$ and the integral is $\int_0^2 \int_{y^2}^{y+2} xy\,dx\,dy$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\displaystyle\int_0^2 \int_{y^2}^{y+2} xy\,dx\,dy = \int_0^2 y\left[\frac{x^2}{2}\right]_{x=y^2}^{x=y+2} dy = \frac{1}{2}\int_0^2 y\left[(y+2)^2-(y^2)^2\right]\,dy = \frac{1}{2}\int_0^2 [y^3+4y^2+4y-y^5]\,dy$

$\displaystyle = \tfrac{1}{2}\left[\tfrac{1}{4}y^4+\tfrac{4}{3}y^3+2y^2-\tfrac{1}{6}y^6\right]_0^2 = \tfrac{1}{2}\left(4+\tfrac{32}{3}+8-\tfrac{32}{3}\right) = 6$

10. (a) We express the iterated integral as a Type I. A Type II would require the sum of two integrals. The curves intersect when $6-x=x^2 \;\Leftrightarrow\; x^2+x-6=0 \;\Leftrightarrow\; (x+3)(x-2)=0 \;\Leftrightarrow\; x=-3$ or $x=2$. The point for $x=-3$ is not in D. Thus, the point of intersection of the two curves is $(2,4)$ and the integral is $\displaystyle\int_0^2\int_{x^2}^{6-x} x\,dy\,dx$.

(b) $\displaystyle\int_0^2\int_{x^2}^{6-x} x\,dy\,dx = \int_0^2 x\Big[y\Big]_{y=x^2}^{y=6-x} dx = \int_0^2 x[(6-x)-x^2]\,dx = \int_0^2 [6x-x^2-x^3]\,dx$

$\displaystyle = \left[3x^2-\frac{x^3}{3}-\frac{x^4}{4}\right]_0^2 = 12-\frac{8}{3}-4 = \frac{16}{3}$

11. $\displaystyle\iint_D \frac{y}{x^2+1}\,dA = \int_0^4\int_0^{\sqrt{x}} \frac{y}{x^2+1}\,dy\,dx = \int_0^4 \left[\frac{1}{x^2+1}\cdot\frac{y^2}{2}\right]_{y=0}^{y=\sqrt{x}} dx = \frac{1}{2}\int_0^4 \frac{x}{x^2+1}\,dx$

$\displaystyle = \tfrac{1}{2}\left[\tfrac{1}{2}\ln\left|x^2+1\right|\right]_0^4 = \tfrac{1}{4}\Big[\ln\left(x^2+1\right)\Big]_0^4 = \tfrac{1}{4}(\ln 17-\ln 1) = \tfrac{1}{4}\ln 17$

12. $\displaystyle\iint_D (2x+y)\,dA = \int_1^2\int_{y-1}^1 (2x+y)\,dx\,dy = \int_1^2 \left[x^2+xy\right]_{x=y-1}^{x=1} dy = \int_1^2 \left[1+y-(y-1)^2-y(y-1)\right] dy$

$\displaystyle = \int_1^2 (-2y^2+4y)\,dy = \left[-\tfrac{2}{3}y^3+2y^2\right]_1^2 = \left(-\tfrac{16}{3}+8\right)-\left(-\tfrac{2}{3}+2\right) = \tfrac{4}{3}$

13. $\displaystyle\iint_D e^{-y^2}\,dA = \int_0^3\int_0^y e^{-y^2}\,dx\,dy = \int_0^3 \left[xe^{-y^2}\right]_{x=0}^{x=y} dy = \int_0^3 \left(ye^{-y^2}-0\right) dy = \int_0^3 ye^{-y^2}\,dy$

$\displaystyle = -\tfrac{1}{2}e^{-y^2}\Big]_0^3 = -\tfrac{1}{2}\left(e^{-9}-e^0\right) = \tfrac{1}{2}\left(1-e^{-9}\right)$

14. $\displaystyle\iint_D y\sqrt{x^2-y^2}\,dA = \int_0^2\int_0^x y\sqrt{x^2-y^2}\,dy\,dx = \int_0^2 \left[-\tfrac{1}{3}(x^2-y^2)^{3/2}\right]_{y=0}^{y=x} dx = \int_0^2 \left[0+\tfrac{1}{3}(x^2)^{3/2}\right] dx$

$\displaystyle = \int_0^2 \tfrac{1}{3}x^3\,dx = \tfrac{1}{3}\cdot\tfrac{1}{4}x^4\Big]_0^2 = \tfrac{1}{12}(16-0) = \tfrac{4}{3}$

15. (a) At the right we sketch an example of a region D that can be described as lying between the graphs of two continuous functions of x (a type I region) but not as lying between graphs of two continuous functions of y (a type II region). The regions shown in Figures 6 and 8 in the text are additional examples.

(b) Now we sketch an example of a region D that can be described as lying between the graphs of two continuous functions of y but not as lying between graphs of two continuous functions of x. The first region shown in Figure 7 is another example.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16. (a) At the right we sketch an example of a region D that can be described as lying between the graphs of two continuous functions of x (a type I region) and also as lying between graphs of two continuous functions of y (a type II region). For additional examples see Figures 9–10 and 15–16 in the text.

(b) Now we sketch an example of a region D that can't be described as lying between the graphs of two continuous functions of x or between graphs of two continuous functions of y. The region shown in Figure 18 is another example.

17.

As a type I region, D lies between the lower boundary $y = 0$ and the upper boundary $y = x$ for $0 \le x \le 1$, so $D = \{(x, y) \mid 0 \le x \le 1, 0 \le y \le x\}$. If we describe D as a type II region, D lies between the left boundary $x = y$ and the right boundary $x = 1$ for $0 \le y \le 1$, so $D = \{(x, y) \mid 0 \le y \le 1, y \le x \le 1\}$.

Thus $\iint_D x\,dA = \int_0^1 \int_0^x x\,dy\,dx = \int_0^1 \left[xy\right]_{y=0}^{y=x} dx = \int_0^1 x^2\,dx = \frac{1}{3}x^3\Big]_0^1 = \frac{1}{3}(1-0) = \frac{1}{3}$ or

$\iint_D x\,dA = \int_0^1 \int_y^1 x\,dx\,dy = \int_0^1 \left[\frac{1}{2}x^2\right]_{x=y}^{x=1} dy = \frac{1}{2}\int_0^1 (1-y^2)\,dy = \frac{1}{2}\left[y - \frac{1}{3}y^3\right]_0^1 = \frac{1}{2}\left[\left(1-\frac{1}{3}\right) - 0\right] = \frac{1}{3}$.

18.

The curves $y = x^2$ and $y = 3x$ intersect at points $(0, 0)$, $(3, 9)$. As a type I region, D is enclosed by the lower boundary $y = x^2$ and the upper boundary $y = 3x$ for $0 \le x \le 3$, so $D = \left\{(x, y) \mid 0 \le x \le 3, x^2 \le y \le 3x\right\}$. If we describe D as a type II region, D is enclosed by the left boundary $x = y/3$ and the right boundary $x = \sqrt{y}$ for $0 \le y \le 9$, so $D = \left\{(x, y) \mid 0 \le y \le 9, y/3 \le x \le \sqrt{y}\right\}$. Thus

$$\iint_D xy\,dA = \int_0^3 \int_{x^2}^{3x} xy\,dy\,dx = \int_0^3 \left[x \cdot \tfrac{1}{2}y^2\right]_{y=x^2}^{y=3x} dx = \tfrac{1}{2}\int_0^3 x(9x^2 - x^4)\,dx = \tfrac{1}{2}\int_0^3 (9x^3 - x^5)\,dx$$
$$= \tfrac{1}{2}\left[9 \cdot \tfrac{1}{4}x^4 - \tfrac{1}{6}x^6\right]_0^3 = \tfrac{1}{2}\left[\left(\tfrac{9}{4} \cdot 81 - \tfrac{1}{6} \cdot 729\right) - 0\right] = \tfrac{243}{8}$$

or
$$\iint_D xy\,dA = \int_0^9 \int_{y/3}^{\sqrt{y}} xy\,dx\,dy = \int_0^9 \left[\tfrac{1}{2}x^2 y\right]_{x=y/3}^{x=\sqrt{y}} dy = \tfrac{1}{2}\int_0^9 \left(y - \tfrac{1}{9}y^2\right)y\,dy = \tfrac{1}{2}\int_0^9 \left(y^2 - \tfrac{1}{9}y^3\right)dy$$
$$= \tfrac{1}{2}\left[\tfrac{1}{3}y^3 - \tfrac{1}{9} \cdot \tfrac{1}{4}y^4\right]_0^9 = \tfrac{1}{2}\left[\left(\tfrac{1}{3} \cdot 729 - \tfrac{1}{36} \cdot 6561\right) - 0\right] = \tfrac{243}{8}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19.

The curves $y = x - 2$ or $x = y + 2$ and $x = y^2$ intersect when $y + 2 = y^2 \quad \Leftrightarrow$ $y^2 - y - 2 = 0 \quad \Leftrightarrow \quad (y-2)(y+1) = 0 \quad \Leftrightarrow \quad y = -1, y = 2$, so the points of intersection are $(1, -1)$ and $(4, 2)$. If we describe D as a type I region, the upper boundary curve is $y = \sqrt{x}$ but the lower boundary curve consists of two parts, $y = -\sqrt{x}$ for $0 \le x \le 1$ and $y = x - 2$ for $1 \le x \le 4$.

Thus $D = \{(x, y) \mid 0 \le x \le 1,\ -\sqrt{x} \le y \le \sqrt{x}\} \cup \{(x, y) \mid 1 \le x \le 4, x - 2 \le y \le \sqrt{x}\}$ and $\iint_D y\,dA = \int_0^1 \int_{-\sqrt{x}}^{\sqrt{x}} y\,dy\,dx + \int_1^4 \int_{x-2}^{\sqrt{x}} y\,dy\,dx$. If we describe D as a type II region, D is enclosed by the left boundary $x = y^2$ and the right boundary $x = y + 2$ for $-1 \le y \le 2$, so $D = \left\{(x, y) \mid -1 \le y \le 2, y^2 \le x \le y + 2\right\}$ and $\iint_D y\,dA = \int_{-1}^2 \int_{y^2}^{y+2} y\,dx\,dy$. In either case, the resulting iterated integrals are not difficult to evaluate but the region D is more simply described as a type II region, giving one iterated integral rather than a sum of two, so we evaluate the latter integral:

$$\iint_D y\,dA = \int_{-1}^2 \int_{y^2}^{y+2} y\,dx\,dy = \int_{-1}^2 \left[xy\right]_{x=y^2}^{x=y+2} dy = \int_{-1}^2 (y + 2 - y^2)y\,dy = \int_{-1}^2 (y^2 + 2y - y^3)\,dy$$

$$= \left[\tfrac{1}{3}y^3 + y^2 - \tfrac{1}{4}y^4\right]_{-1}^2 = \left(\tfrac{8}{3} + 4 - 4\right) - \left(-\tfrac{1}{3} + 1 - \tfrac{1}{4}\right) = \tfrac{9}{4}$$

20.

As a type I region, $D = \{(x, y) \mid 0 \le x \le 4, x \le y \le 4\}$ and $\iint_D y^2 e^{xy}\,dA = \int_0^4 \int_x^4 y^2 e^{xy}\,dy\,dx$. As a type II region, $D = \{(x, y) \mid 0 \le y \le 4, 0 \le x \le y\}$ and $\iint_D y^2 e^{xy}\,dA = \int_0^4 \int_0^y y^2 e^{xy}\,dx\,dy$. Evaluating $\int y^2 e^{xy}\,dy$ requires integration by parts whereas $\int y^2 e^{xy}\,dx$ does not, so the iterated integral corresponding to D as a type II region appears easier to evaluate.

$$\iint_D y^2 e^{xy}\,dA = \int_0^4 \int_0^y y^2 e^{xy}\,dx\,dy = \int_0^4 \left[ye^{xy}\right]_{x=0}^{x=y} dy = \int_0^4 \left(ye^{y^2} - y\right) dy$$

$$= \left[\tfrac{1}{2}e^{y^2} - \tfrac{1}{2}y^2\right]_0^4 = \left(\tfrac{1}{2}e^{16} - 8\right) - \left(\tfrac{1}{2} - 0\right) = \tfrac{1}{2}e^{16} - \tfrac{17}{2}$$

21.

If we describe D as a type I region, $D = \{(x, y) \mid 0 \le x \le \pi/2, 0 \le y \le \cos x\}$ and $\iint_D \sin^2 x\,dA = \int_0^{\pi/2} \int_0^{\cos x} \sin^2 x\,dy\,dx$. As a type II region, $D = \{(x, y) \mid 0 \le x \le \cos^{-1} y, 0 \le y \le 1\}$ and $\iint_D \sin^2 x\,dA = \displaystyle\int_0^1 \int_0^{\cos^{-1} y} \sin^2 x\,dx\,dy$. Evaluating $\displaystyle\int_0^{\cos^{-1} y} \sin^2 x\,dx$ will result in a very difficult integral. Therefore, we evaluate the iterated integral that describes D as a type I region because integrating $\sin^2 x$ with respect to y is easy.

$$\int_0^{\pi/2} \int_0^{\cos x} \sin^2 x\,dy\,dx = \int_0^{\pi/2} \sin^2 x \left[y\right]_{y=0}^{y=\cos x} dx = \int_0^{\pi/2} \cos x \sin^2 x\,dx$$

$$= \int_0^1 u^2\,du \quad \begin{bmatrix} u = \sin x, \\ du = \cos x\,dx \end{bmatrix} \quad = \left[\frac{u^3}{3}\right]_0^1 = \frac{1}{3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22.

By inspection, the curves $y = 2x + 4$ and $y = x^3$ intersect when $x^3 = 2x + 4 \iff x = 2$, so the point of intersection is $(2, 8)$. If we describe D as a type 1 region, $D = \{(x, y) \mid 0 \le x \le 2,\ x^3 \le y \le 2x + 4\}$ and the integral is

$$\iint_D 6x^2\,dA = \int_0^2 \int_{x^3}^{2x+4} 6x^2\,dy\,dx.$$

If we describe D as a type II region, the right boundary curve is $x = \sqrt[3]{y}$, but the left boundary curve consists of two parts, $x = 0$ for $0 \le y \le 4$ and $x = y/2 - 2$ for $4 \le y \le 8$.

In either case, the resulting iterated integrals are not difficult to evaluate, but the region D is more simply described as a type I region, giving one iterated integral rather than a sum of two, so we evaluate that integral:

$$\int_0^2 \int_{x^3}^{2x+4} 6x^2\,dy\,dx = \int_0^2 \Big[6x^2 y\Big]_{y=x^3}^{y=2x+4} dx = \int_0^2 [6x^2(2x + 4 - x^3)]\,dx = \int_0^2 (12x^3 + 24x^2 - 6x^5)\,dx$$
$$= \big[3x^4 + 8x^3 - x^6\big]_0^2 = 48 + 64 - 64 = 48$$

23.

$$\int_0^1 \int_0^{x^2} x\cos y\,dy\,dx = \int_0^1 \big[x \sin y\big]_{y=0}^{y=x^2} dx = \int_0^1 x \sin x^2\,dx$$
$$= -\tfrac{1}{2}\cos x^2\big]_0^1 = -\tfrac{1}{2}(\cos 1 - \cos 0) = \tfrac{1}{2}(1 - \cos 1)$$

24.

$$\iint_D (x^2 + 2y)\,dA = \int_0^1 \int_{x^3}^{x} (x^2 + 2y)\,dy\,dx = \int_0^1 \big[x^2 y + y^2\big]_{y=x^3}^{y=x} dx$$
$$= \int_0^1 (x^3 + x^2 - x^5 - x^6)\,dx = \big[\tfrac{1}{4}x^4 + \tfrac{1}{3}x^3 - \tfrac{1}{6}x^6 - \tfrac{1}{7}x^7\big]_0^1$$
$$= \tfrac{1}{4} + \tfrac{1}{3} - \tfrac{1}{6} - \tfrac{1}{7} = \tfrac{23}{84}$$

25.

$$\iint_D y^2\,dA = \int_1^2 \int_{y-1}^{7-3y} y^2\,dx\,dy = \int_1^2 \big[xy^2\big]_{x=y-1}^{x=7-3y} dy$$
$$= \int_1^2 [(7 - 3y) - (y - 1)]\,y^2\,dy = \int_1^2 (8y^2 - 4y^3)\,dy$$
$$= \big[\tfrac{8}{3}y^3 - y^4\big]_1^2 = \tfrac{64}{3} - 16 - \tfrac{8}{3} + 1 = \tfrac{11}{3}$$

26.

$$\iint_D xy\,dA = \int_0^1 \int_0^{\sqrt{1-x^2}} xy\,dy\,dx$$
$$= \int_0^1 \big[\tfrac{1}{2}xy^2\big]_{y=0}^{y=\sqrt{1-x^2}} dx = \int_0^1 \tfrac{1}{2}x(1 - x^2)\,dx$$
$$= \tfrac{1}{2}\int_0^1 (x - x^3)\,dx = \tfrac{1}{2}\big[\tfrac{1}{2}x^2 - \tfrac{1}{4}x^4\big]_0^1$$
$$= \tfrac{1}{2}\big(\tfrac{1}{2} - \tfrac{1}{4} - 0\big) = \tfrac{1}{8}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27.

$$\int_{-2}^{2} \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} (2x - y)\, dy\, dx$$

$$= \int_{-2}^{2} \left[2xy - \tfrac{1}{2}y^2\right]_{y=-\sqrt{4-x^2}}^{y=\sqrt{4-x^2}} dx$$

$$= \int_{-2}^{2} \left[2x\sqrt{4-x^2} - \tfrac{1}{2}\left(4-x^2\right) + 2x\sqrt{4-x^2} + \tfrac{1}{2}\left(4-x^2\right)\right] dx$$

$$= \int_{-2}^{2} 4x\sqrt{4-x^2}\, dx = -\tfrac{4}{3}\left(4-x^2\right)^{3/2}\Big]_{-2}^{2} = 0$$

[Or, note that $4x\sqrt{4-x^2}$ is an odd function, so $\int_{-2}^{2} 4x\sqrt{4-x^2}\, dx = 0$.]

28.

$$\iint_D y\, dA = \int_0^1 \int_y^{4-3y} y\, dx\, dy$$

$$= \int_0^1 \left[xy\right]_{x=y}^{x=4-3y} dy = \int_0^1 (4y - 3y^2 - y^2)\, dy$$

$$= \int_0^1 (4y - 4y^2)\, dy = \left[2y^2 - \tfrac{4}{3}y^3\right]_0^1 = 2 - \tfrac{4}{3} - 0 = \tfrac{2}{3}$$

29. (a)

As a Type I region, $D = \{(x, y) \mid 0 \le x \le 1,\ x \le y \le 1\}$ and the volume V of the solid that lies under the surface and above D is given by $V = \iint_D (1 + xy)\, dA = \int_0^1 \int_x^1 (1 + xy)\, dy\, dx$. As a Type II region, $D = \{(x, y) \mid 0 \le y \le 1,\ 0 \le x \le y\}$ and $V = \int_0^1 \int_0^y (1 + xy)\, dx\, dy$. Evaluate either integral in part (b).

(b) $$\int_0^1 \int_0^y (1 + xy)\, dx\, dy = \int_0^1 \left[x + y\,\frac{x^2}{2}\right]_{x=0}^{x=y} dy = \int_0^1 \left[\left(y + \frac{y^3}{2}\right) - 0\right] dy = \left[\frac{y^2}{2} + \frac{y^4}{8}\right]_0^1 = \frac{1}{2} + \frac{1}{8} = \frac{5}{8}$$

30. (a)

As a Type I region, $D = \{(x, y) \mid 0 \le x \le 1,\ 0 \le y \le x^2\}$ and the volume V of the solid that lies under the surface and above D is given by $V = \iint_D (x^2 + y^2)\, dA = \int_0^1 \int_0^{x^2} (x^2 + y^2)\, dy\, dx$. As a Type II region, $D = \{(x, y) \mid 0 \le y \le 1,\ \sqrt{y} \le x \le 1\}$ and $V = \int_0^1 \int_{\sqrt{y}}^1 (x^2 + y^2)\, dx\, dy$.

(b) $$\int_0^1 \int_0^{x^2} (x^2 + y^2)\, dy\, dx = \int_0^1 \left[x^2 y + \frac{y^3}{3}\right]_{y=0}^{y=x^2} dx = \int_0^1 \left[x^4 + \frac{x^6}{3}\right] dx = \left[\frac{x^5}{5} + \frac{x^7}{21}\right]_0^1 = \frac{1}{5} + \frac{1}{21} = \frac{26}{105}$$

31.

$$V = \int_0^1 \int_{x^2}^{\sqrt{x}} (3x + 2y)\, dy\, dx = \int_0^1 \left[3xy + y^2\right]_{y=x^2}^{y=\sqrt{x}} dx$$

$$= \int_0^1 \left[(3x\sqrt{x} + x) - (3x^3 + x^4)\right] dx = \int_0^1 (3x^{3/2} + x - 3x^3 - x^4)\, dx$$

$$= \left[3 \cdot \tfrac{2}{5}x^{5/2} + \tfrac{1}{2}x^2 - \tfrac{3}{4}x^4 - \tfrac{1}{5}x^5\right]_0^1 = \tfrac{6}{5} + \tfrac{1}{2} - \tfrac{3}{4} - \tfrac{1}{5} - 0 = \tfrac{3}{4}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32.

$$V = \int_{-2}^{2}\int_{y^2}^{4}(1 + x^2y^2)\,dx\,dy$$

$$= \int_{-2}^{2}\left[x + \tfrac{1}{3}x^3y^2\right]_{x=y^2}^{x=4}\,dy = \int_{-2}^{2}(4 + \tfrac{61}{3}y^2 - \tfrac{1}{3}y^8)\,dy$$

$$= \left[4y + \tfrac{61}{9}y^3 - \tfrac{1}{27}y^9\right]_{-2}^{2} = 8 + \tfrac{488}{9} - \tfrac{512}{27} + 8 + \tfrac{488}{9} - \tfrac{512}{27} = \tfrac{2336}{27}$$

33.

$$V = \int_{1}^{2}\int_{1}^{7-3y} xy\,dx\,dy = \int_{1}^{2}\left[\tfrac{1}{2}x^2y\right]_{x=1}^{x=7-3y}\,dy$$

$$= \tfrac{1}{2}\int_{1}^{2} y\left[(7-3y)^2 - 1\right]dy = \tfrac{1}{2}\int_{1}^{2}(48y - 42y^2 + 9y^3)\,dy$$

$$= \tfrac{1}{2}\left[24y^2 - 14y^3 + \tfrac{9}{4}y^4\right]_{1}^{2} = \tfrac{31}{8}$$

34.

$$V = \int_{0}^{2}\int_{0}^{2-x}(x^2 + y^2 + 1)\,dy\,dx = \int_{0}^{2}\left[x^2y + \tfrac{1}{3}y^3 + y\right]_{y=0}^{y=2-x}\,dx$$

$$= \int_{0}^{2}\left[x^2(2-x) + \tfrac{1}{3}(2-x)^3 + (2-x) - 0\right]dx$$

$$= \int_{0}^{2}\left(-\tfrac{4}{3}x^3 + 4x^2 - 5x + \tfrac{14}{3}\right)dx = \left[-\tfrac{1}{3}x^4 + \tfrac{4}{3}x^3 - \tfrac{5}{2}x^2 + \tfrac{14}{3}x\right]_{0}^{2}$$

$$= -\tfrac{16}{3} + \tfrac{32}{3} - 10 + \tfrac{28}{3} - 0 = \tfrac{14}{3}$$

35.

$$V = \int_{0}^{2}\int_{0}^{4-2x}(4 - 2x - y)\,dy\,dx = \int_{0}^{2}\left[4y - 2xy - \tfrac{1}{2}y^2\right]_{y=0}^{y=4-2x}\,dx$$

$$= \int_{0}^{2}\left[4(4-2x) - 2x(4-2x) - \tfrac{1}{2}(4-2x)^2 - 0\right]dx$$

$$= \int_{0}^{2}\left(2x^2 - 8x + 8\right)dx = \left[\tfrac{2}{3}x^3 - 4x^2 + 8x\right]_{0}^{2} = \tfrac{16}{3} - 16 + 16 - 0 = \tfrac{16}{3}$$

36.

$$V = \int_{0}^{1}\int_{x}^{2-x} x\,dy\,dx$$

$$= \int_{0}^{1}\left[xy\right]_{y=x}^{y=2-x}\,dx = \int_{0}^{1}(2x - 2x^2)\,dx$$

$$= \left[x^2 - \tfrac{2}{3}x^3\right]_{0}^{1} = \tfrac{1}{3}$$

37.

$$V = \int_{-2}^{2}\int_{x^2}^{4} x^2\,dy\,dx$$

$$= \int_{-2}^{2}\left[x^2y\right]_{y=x^2}^{y=4}\,dx = \int_{-2}^{2}(4x^2 - x^4)\,dx$$

$$= \left[\tfrac{4}{3}x^3 - \tfrac{1}{5}x^5\right]_{-2}^{2} = \tfrac{32}{3} - \tfrac{32}{5} + \tfrac{32}{3} - \tfrac{32}{5} = \tfrac{128}{15}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38.

$$V = \int_0^2 \int_0^{2y} \sqrt{4-y^2}\,dx\,dy = \int_0^2 \left[x\sqrt{4-y^2}\right]_{x=0}^{x=2y} dy$$

$$= \int_0^2 2y\sqrt{4-y^2}\,dy = \left[-\tfrac{2}{3}\left(4-y^2\right)^{3/2}\right]_0^2 = 0 + \tfrac{16}{3} = \tfrac{16}{3}$$

39.

$$V = \int_0^1 \int_0^{\sqrt{1-x^2}} y\,dy\,dx = \int_0^1 \left[\frac{y^2}{2}\right]_{y=0}^{y=\sqrt{1-x^2}} dx$$

$$= \int_0^1 \frac{1-x^2}{2}\,dx = \tfrac{1}{2}\left[x - \tfrac{1}{3}x^3\right]_0^1 = \tfrac{1}{3}$$

40.

By symmetry, the desired volume V is 8 times the volume V_1 in the first octant. Now

$$V_1 = \int_0^r \int_0^{\sqrt{r^2-y^2}} \sqrt{r^2-y^2}\,dx\,dy = \int_0^r \left[x\sqrt{r^2-y^2}\right]_{x=0}^{x=\sqrt{r^2-y^2}} dy$$

$$= \int_0^r (r^2-y^2)\,dy = \left[r^2y - \tfrac{1}{3}y^3\right]_0^r = \tfrac{2}{3}r^3$$

Thus $V = \frac{16}{3}r^3$.

41.

From the graph, it appears that the two curves intersect at $x = 0$ and at $x \approx 1.213$. Thus the desired integral is

$$\iint_D x\,dA \approx \int_0^{1.213} \int_{x^4}^{3x-x^2} x\,dy\,dx = \int_0^{1.213} \left[xy\right]_{y=x^4}^{y=3x-x^2} dx$$

$$= \int_0^{1.213} (3x^2 - x^3 - x^5)\,dx = \left[x^3 - \tfrac{1}{4}x^4 - \tfrac{1}{6}x^6\right]_0^{1.213}$$

$$\approx 0.713$$

42.

The desired solid is shown in the first graph. From the second graph, we estimate that $y = \cos x$ intersects $y = x$ at $x \approx 0.7391$. Therefore the volume of the solid is

$$V \approx \int_0^{0.7391} \int_x^{\cos x} x\,dy\,dx = \int_0^{0.7391} \left[xy\right]_{y=x}^{y=\cos x} dx$$

$$= \int_0^{0.7391} (x\cos x - x^2)\,dx = \left[\cos x + x\sin x - \tfrac{1}{3}x^3\right]_0^{0.7391} \approx 0.1024$$

Note: There is a different solid which can also be construed to satisfy the conditions stated in the exercise. This is the solid bounded by all of the given surfaces, as well as the plane $y = 0$. In case you calculated the volume of this solid and want to check your work, its volume is $V \approx \int_0^{0.7391} \int_0^x x\,dy\,dx + \int_{0.7391}^{\pi/2} \int_0^{\cos x} x\,dy\,dx \approx 0.4684$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

43. The region of integration is bounded by the curves $y = 1 - x^2$ and $y = x^2 - 1$ which intersect at $(\pm 1, 0)$ with $1 - x^2 \geq x^2 - 1$ on $[-1, 1]$. Within this region, the plane $z = 2x + 2y + 10$ is above the plane $z = 2 - x - y$, so

$$\begin{aligned}
V &= \textstyle\int_{-1}^{1} \int_{x^2-1}^{1-x^2} (2x + 2y + 10)\, dy\, dx - \int_{-1}^{1} \int_{x^2-1}^{1-x^2} (2 - x - y)\, dy\, dx \\
&= \textstyle\int_{-1}^{1} \int_{x^2-1}^{1-x^2} (2x + 2y + 10 - (2 - x - y))\, dy\, dx \\
&= \textstyle\int_{-1}^{1} \int_{x^2-1}^{1-x^2} (3x + 3y + 8)\, dy\, dx = \int_{-1}^{1} \left[3xy + \frac{3}{2}y^2 + 8y\right]_{y=x^2-1}^{y=1-x^2} dx \\
&= \textstyle\int_{-1}^{1} \left[3x(1 - x^2) + \frac{3}{2}(1 - x^2)^2 + 8(1 - x^2) - 3x(x^2 - 1) - \frac{3}{2}(x^2 - 1)^2 - 8(x^2 - 1)\right] dx \\
&= \textstyle\int_{-1}^{1} (-6x^3 - 16x^2 + 6x + 16)\, dx = \left[-\frac{3}{2}x^4 - \frac{16}{3}x^3 + 3x^2 + 16x\right]_{-1}^{1} \\
&= -\tfrac{3}{2} - \tfrac{16}{3} + 3 + 16 + \tfrac{3}{2} - \tfrac{16}{3} - 3 + 16 = \tfrac{64}{3}
\end{aligned}$$

44. The two planes intersect in the line $y = 1$, $z = 3$, so the region of integration is the plane region enclosed by the parabola $y = x^2$ and the line $y = 1$. We have $2 + y \geq 3y$ for $0 \leq y \leq 1$, so the solid region is bounded above by $z = 2 + y$ and bounded below by $z = 3y$.

$$\begin{aligned}
V &= \int_{-1}^{1} \int_{x^2}^{1} (2 + y)\, dy\, dx - \int_{-1}^{1} \int_{x^2}^{1} (3y)\, dy\, dx = \int_{-1}^{1} \int_{x^2}^{1} (2 + y - 3y)\, dy\, dx \\
&= \int_{-1}^{1} \int_{x^2}^{1} (2 - 2y)\, dy\, dx = \textstyle\int_{-1}^{1} \left[2y - y^2\right]_{y=x^2}^{y=1} dx \\
&= \textstyle\int_{-1}^{1} (1 - 2x^2 + x^4)\, dx = x - \frac{2}{3}x^3 + \frac{1}{5}x^5\big]_{-1}^{1} = \frac{16}{15}
\end{aligned}$$

45. The region of integration is bounded by the curves $y = x^2$ and $y = 1 - x^2$ which intersect at $\left(\pm\frac{1}{\sqrt{2}}, \frac{1}{2}\right)$.

The solid lies under the graph of $z = 3$ and above the graph of $z = y$, so its volume is

$$\begin{aligned}
V &= \textstyle\int_{-1/\sqrt{2}}^{1/\sqrt{2}} \int_{x^2}^{1-x^2} 3\, dy\, dx - \int_{-1/\sqrt{2}}^{1/\sqrt{2}} \int_{x^2}^{1-x^2} y\, dy\, dx = \int_{-1/\sqrt{2}}^{1/\sqrt{2}} \int_{x^2}^{1-x^2} (3 - y)\, dy\, dx \\
&= \textstyle\int_{-1/\sqrt{2}}^{1/\sqrt{2}} \left[3y - \frac{1}{2}y^2\right]_{y=x^2}^{y=1-x^2} dx = \int_{-1/\sqrt{2}}^{1/\sqrt{2}} \left[\left(3(1 - x^2) - \frac{1}{2}(1 - x^2)^2\right) - \left(3x^2 - \frac{1}{2}(x^2)^2\right)\right] dx \\
&= \textstyle\int_{-1/\sqrt{2}}^{1/\sqrt{2}} \left(\frac{5}{2} - 5x^2\right) dx = \left[\frac{5}{2}x - \frac{5}{3}x^3\right]_{-1/\sqrt{2}}^{1/\sqrt{2}} = \left(\frac{5}{2\sqrt{2}} - \frac{5}{6\sqrt{2}}\right) - \left(-\frac{5}{2\sqrt{2}} + \frac{5}{6\sqrt{2}}\right) \\
&= \tfrac{10}{3\sqrt{2}} \text{ or } \tfrac{5\sqrt{2}}{3}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. The region of integration is the portion of the first quadrant bounded by the axes and the curve $y = \sqrt{4 - x^2}$. The solid lies under the graph of $z = x + y$ and above the graph of $z = xy$, so its volume is

$$V = \int_0^2 \int_0^{\sqrt{4-x^2}} (x + y)\, dy\, dx - \int_0^2 \int_0^{\sqrt{4-x^2}} xy\, dy\, dx = \int_0^2 \int_0^{\sqrt{4-x^2}} (x + y - xy)\, dy\, dx$$

$$= \int_0^2 \left[xy + \tfrac{1}{2}y^2 - \tfrac{1}{2}xy^2\right]_{y=0}^{y=\sqrt{4-x^2}} dx = \int_0^2 \left[x\sqrt{4 - x^2} + \tfrac{1}{2}(4 - x^2) - \tfrac{1}{2}x(4 - x^2) - 0\right] dx$$

$$= \int_0^2 \left(x\sqrt{4 - x^2} + 2 - \tfrac{1}{2}x^2 - 2x + \tfrac{1}{2}x^3\right) dx = \left[-\tfrac{1}{3}(4 - x^2)^{3/2} + 2x - \tfrac{1}{6}x^3 - x^2 + \tfrac{1}{8}x^4\right]_0^2$$

$$= \left(4 - \tfrac{4}{3} - 4 + 2\right) - \left(-\tfrac{1}{3} \cdot 4^{3/2}\right) = \tfrac{2}{3} + \tfrac{8}{3} = \tfrac{10}{3}$$

47. $\int_0^1 \int_0^{1-x} (1 - x - y)\, dy\, dx$.

The solid lies below the plane $z = 1 - x - y$ or $x + y + z = 1$ and above the region $D = \{(x, y) \mid 0 \le x \le 1, 0 \le y \le 1 - x\}$ in the xy-plane. The solid is a tetrahedron.

48. $\displaystyle\int_0^1 \int_0^{1-x^2} (1 - x)\, dy\, dx$.

The solid lies below the plane $z = 1 - x$ and above the region $D = \{(x, y) \mid 0 \le x \le 1, 0 \le y \le 1 - x^2\}$ in the xy-plane.

49. $\int_0^3 \int_0^y \sqrt{9 - x^2}\, dx\, dy$.

The solid lies under the top half of the cylinder $x^2 + z^2 = 9$; that is, $z = \sqrt{9 - x^2}$, and above the region $D = \{(x, y) \mid 0 \le x \le y, 0 \le y \le 3\}$.

50. $\displaystyle\int_{-2}^{2} \int_{-1}^{3-x^2} e^{-y}\, dy\, dx$.

The solid lies below the surface $z = e^{-y}$ and above the region $D = \{(x, y) \mid -2 \le x \le 2, -1 \le y \le 3 - x^2\}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. The two bounding curves $y = x^3 - x$ and $y = x^2 + x$ intersect at the origin and at $x = 2$, with $x^2 + x > x^3 - x$ on $(0, 2)$. Using a CAS, we find that the volume of the solid is

$$V = \int_0^2 \int_{x^3 - x}^{x^2 + x} (x^3 y^4 + xy^2)\, dy\, dx = \frac{13{,}984{,}735{,}616}{14{,}549{,}535}$$

52. For $|x| \le 1$ and $|y| \le 1$, $2x^2 + y^2 < 8 - x^2 - 2y^2$. Also, the cylinder is described by the inequalities $-1 \le x \le 1$, $-\sqrt{1 - x^2} \le y \le \sqrt{1 - x^2}$. So the volume is given by

$$V = \int_{-1}^{1} \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \left[(8 - x^2 - 2y^2) - (2x^2 + y^2)\right] dy\, dx = \frac{13\pi}{2} \qquad \text{[using a CAS]}$$

53. The two surfaces intersect in the circle $x^2 + y^2 = 1$, $z = 0$ and the region of integration is the disk D: $x^2 + y^2 \le 1$. Using a CAS, the volume is $\iint_D (1 - x^2 - y^2)\, dA = \int_{-1}^{1} \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} (1 - x^2 - y^2)\, dy\, dx = \dfrac{\pi}{2}$.

54. The projection onto the xy-plane of the intersection of the two surfaces is the circle $x^2 + y^2 = 2y \;\Rightarrow$ $x^2 + y^2 - 2y = 0 \;\Rightarrow\; x^2 + (y - 1)^2 = 1$, so the region of integration is given by $-1 \le x \le 1$, $1 - \sqrt{1 - x^2} \le y \le 1 + \sqrt{1 - x^2}$. In this region, $2y \ge x^2 + y^2$ so, using a CAS, the volume is

$$V = \int_{-1}^{1} \int_{1-\sqrt{1-x^2}}^{1+\sqrt{1-x^2}} [2y - (x^2 + y^2)]\, dy\, dx = \frac{\pi}{2}$$

55.

Because the region of integration is

$D = \{(x, y) \mid 0 \le x \le y, 0 \le y \le 1\} = \{(x, y) \mid x \le y \le 1, 0 \le x \le 1\}$

we have $\int_0^1 \int_0^y f(x, y)\, dx\, dy = \iint_D f(x, y)\, dA = \int_0^1 \int_x^1 f(x, y)\, dy\, dx$.

56.

Because the region of integration is

$$\begin{aligned} D &= \left\{(x, y) \mid x^2 \le y \le 4, 0 \le x \le 2\right\} \\ &= \left\{(x, y) \mid 0 \le x \le \sqrt{y}, 0 \le y \le 4\right\} \end{aligned}$$

we have $\int_0^2 \int_{x^2}^4 f(x, y)\, dy\, dx = \iint_D f(x, y)\, dA = \int_0^4 \int_0^{\sqrt{y}} f(x, y)\, dx\, dy$.

57.

Because the region of integration is

$$\begin{aligned} D &= \{(x, y) \mid 0 \le x \le \pi/2,\ \sin x \le y \le 1\} \\ &= \{(x, y) \mid 0 \le x \le \sin^{-1} y, 0 \le y \le 1\} \end{aligned}$$

we have

$\int_0^{\pi/2} \int_{\sin x}^1 f(x, y)\, dy\, dx = \iint_D f(x, y)\, dA = \int_0^1 \int_0^{\sin^{-1} y} f(x, y)\, dx\, dy$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

58.

Because the region of integration is

$$D = \left\{(x, y) \mid 0 \le x \le \sqrt{4 - y^2}, -2 \le y \le 2\right\}$$
$$= \left\{(x, y) \mid -\sqrt{4 - x^2} \le y \le \sqrt{4 - x^2}, 0 \le x \le 2\right\}$$

we have

$$\int_{-2}^{2} \int_0^{\sqrt{4-y^2}} f(x, y)\, dx\, dy = \iint_D f(x, y)\, dA = \int_0^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} f(x, y)\, dy\, dx.$$

59.

Because the region of integration is

$$D = \{(x, y) \mid 0 \le y \le \ln x, 1 \le x \le 2\} = \{(x, y) \mid e^y \le x \le 2, 0 \le y \le \ln 2\}$$

we have

$$\int_1^2 \int_0^{\ln x} f(x, y)\, dy\, dx = \iint_D f(x, y)\, dA = \int_0^{\ln 2} \int_{e^y}^2 f(x, y)\, dx\, dy$$

60.

Because the region of integration is

$$D = \left\{(x, y) \mid \arctan x \le y \le \tfrac{\pi}{4}, 0 \le x \le 1\right\}$$
$$= \left\{(x, y) \mid 0 \le x \le \tan y, 0 \le y \le \tfrac{\pi}{4}\right\}$$

we have

$$\int_0^1 \int_{\arctan x}^{\pi/4} f(x, y)\, dy\, dx = \iint_D f(x, y)\, dA = \int_0^{\pi/4} \int_0^{\tan y} f(x, y)\, dx\, dy$$

61.

$$\int_0^1 \int_{3y}^3 e^{x^2}\, dx\, dy = \int_0^3 \int_0^{x/3} e^{x^2}\, dy\, dx = \int_0^3 \left[e^{x^2} y\right]_{y=0}^{y=x/3} dx$$
$$= \int_0^3 \left(\frac{x}{3}\right) e^{x^2}\, dx = \tfrac{1}{6}\, e^{x^2}\Big]_0^3 = \frac{e^9 - 1}{6}$$

62.

$$\int_0^1 \int_{x^2}^1 \sqrt{y}\, \sin y\, dy\, dx = \int_0^1 \int_0^{\sqrt{y}} \sqrt{y}\, \sin y\, dx\, dy = \int_0^1 \sqrt{y}\, \sin y\, [x]_{x=0}^{x=\sqrt{y}}\, dy$$
$$= \int_0^1 \left(\sqrt{y}\, \sin y\right)\left(\sqrt{y} - 0\right) dy = \int_0^1 y\, \sin y\, dy$$
$$= -y \cos y]_0^1 + \int_0^1 \cos y\, dy$$

[by integrating by parts with $u = y$, $dv = \sin y\, dy$]

$$= [-y \cos y + \sin y]_0^1 = -\cos 1 + \sin 1 - 0 = \sin 1 - \cos 1$$

63.

$$\int_0^1 \int_{\sqrt{x}}^1 \sqrt{y^3 + 1}\, dy\, dx = \int_0^1 \int_0^{y^2} \sqrt{y^3 + 1}\, dx\, dy = \int_0^1 \sqrt{y^3 + 1}\, [x]_{x=0}^{x=y^2}\, dy$$
$$= \int_0^1 y^2 \sqrt{y^3 + 1}\, dy = \tfrac{2}{9}\left(y^3 + 1\right)^{3/2}\Big]_0^1$$
$$= \tfrac{2}{9}\left(2^{3/2} - 1^{3/2}\right) = \tfrac{2}{9}\left(2\sqrt{2} - 1\right)$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64.

$$\int_0^2 \int_{y/2}^1 y\cos(x^3-1)\,dx\,dy = \int_0^1 \int_0^{2x} y\cos(x^3-1)\,dy\,dx$$
$$= \int_0^1 \cos(x^3-1)\ \left[\tfrac{1}{2}y^2\right]_{y=0}^{y=2x}\ dx$$
$$= \int_0^1 2x^2\cos(x^3-1)\,dx = \tfrac{2}{3}\sin(x^3-1)\Big]_0^1$$
$$= \tfrac{2}{3}\left[0-\sin(-1)\right] = -\tfrac{2}{3}\sin(-1) = \tfrac{2}{3}\sin 1$$

65.

$$\int_0^1 \int_{\arcsin y}^{\pi/2} \cos x\,\sqrt{1+\cos^2 x}\,dx\,dy$$
$$= \int_0^{\pi/2}\int_0^{\sin x} \cos x\,\sqrt{1+\cos^2 x}\,dy\,dx$$
$$= \int_0^{\pi/2} \cos x\,\sqrt{1+\cos^2 x}\,\big[\,y\,\big]_{y=0}^{y=\sin x}\ dx$$
$$= \int_0^{\pi/2} \cos x\,\sqrt{1+\cos^2 x}\,\sin x\,dx \qquad \left[\begin{array}{c}\text{Let } u=\cos x,\ du=-\sin x\,dx,\\ dx = du/(-\sin x)\end{array}\right]$$
$$= \int_1^0 -u\,\sqrt{1+u^2}\,du = -\tfrac{1}{3}\left(1+u^2\right)^{3/2}\Big]_1^0$$
$$= \tfrac{1}{3}\left(\sqrt{8}-1\right) = \tfrac{1}{3}\left(2\sqrt{2}-1\right)$$

66.

$$\int_0^8 \int_{\sqrt[3]{y}}^2 e^{x^4}\,dx\,dy = \int_0^2 \int_0^{x^3} e^{x^4}\,dy\,dx$$
$$= \int_0^2 e^{x^4}\Big[\,y\,\Big]_{y=0}^{y=x^3}\ dx = \int_0^2 x^3 e^{x^4}\,dx$$
$$= \tfrac{1}{4}e^{x^4}\Big]_0^2 = \tfrac{1}{4}(e^{16}-1)$$

67. $D = \{(x,y) \mid 0 \le x \le 1,\ -x+1 \le y \le 1\} \cup \{(x,y) \mid -1 \le x \le 0, x+1 \le y \le 1\}$

$\cup\{(x,y) \mid 0 \le x \le 1,\ -1 \le y \le x-1\} \cup \{(x,y) \mid -1 \le x \le 0,\ -1 \le y \le -x-1\}$, all type I.

$$\iint_D x^2\,dA = \int_0^1\int_{1-x}^1 x^2\,dy\,dx + \int_{-1}^0\int_{x+1}^1 x^2\,dy\,dx + \int_0^1\int_{-1}^{x-1} x^2\,dy\,dx + \int_{-1}^0\int_{-1}^{-x-1} x^2\,dy\,dx$$
$$= 4\int_0^1\int_{1-x}^1 x^2\,dy\,dx \qquad \text{[by symmetry of the regions and because } f(x,y)=x^2 \ge 0]$$
$$= 4\int_0^1 x^3\,dx = 4\left[\tfrac{1}{4}x^4\right]_0^1 = 1$$

68. $D = \left\{(x,y) \mid -1 \le y \le 0,\ -1 \le x \le y-y^3\right\} \cup \left\{(x,y) \mid 0 \le y \le 1, \sqrt{y}-1 \le x \le y-y^3\right\}$, both type II.

$$\iint_D y\,dA = \int_{-1}^0\int_{-1}^{y-y^3} y\,dx\,dy + \int_0^1\int_{\sqrt{y}-1}^{y-y^3} y\,dx\,dy = \int_{-1}^0 \left[xy\right]_{x=-1}^{x=y-y^3} dy + \int_0^1 \left[xy\right]_{x=\sqrt{y}-1}^{x=y-y^3} dy$$
$$= \int_{-1}^0 (y^2-y^4+y)\,dy + \int_0^1 (y^2-y^4-y^{3/2}+y)\,dy$$
$$= \left[\tfrac{1}{3}y^3 - \tfrac{1}{5}y^5 + \tfrac{1}{2}y^2\right]_{-1}^0 + \left[\tfrac{1}{3}y^3 - \tfrac{1}{5}y^5 - \tfrac{2}{5}y^{5/2} + \tfrac{1}{2}y^2\right]_0^1$$
$$= (0-\tfrac{11}{30}) + (\tfrac{7}{30}-0) = -\tfrac{2}{15}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

69. Since $x^2+y^2 \le 1$ on S, we must have $0 \le x^2 \le 1$ and $0 \le y^2 \le 1$, so $0 \le x^2y^2 \le 1 \quad \Rightarrow$ $3 \le 4 - x^2y^2 \le 4 \quad \Rightarrow \quad \sqrt{3} \le \sqrt{4-x^2y^2} \le 2$. Here we have $A(S) = \frac{1}{2}\pi(1)^2 = \frac{\pi}{2}$, so by Property 10, $\sqrt{3}A(S) \le \iint_S \sqrt{4-x^2y^2}\,dA \le 2A(S) \quad \Rightarrow \quad \frac{\sqrt{3}}{2}\pi \le \iint_S \sqrt{4-x^2y^2}\,dA \le \pi$ or we can say $2.720 < \iint_S \sqrt{4-x^2y^2}\,dA < 3.142$. (We have rounded the lower bound down and the upper bound up to preserve the inequalities.)

70. T is the triangle with vertices $(0,0)$, $(1,0)$, and $(1,2)$ so $A(T) = \frac{1}{2}(1)(2) = 1$. We have $0 \le \sin^4(x+y) \le 1$ for all x, y, and Property 10 gives $0 \cdot A(T) \le \iint_T \sin^4(x+y)dA \le 1 \cdot A(T) \quad \Rightarrow \quad 0 \le \iint_T \sin^4(x+y)dA \le 1$.

71. The average value of a function f of two variables defined on a rectangle R was defined in Section 15.1 as $f_{\text{avg}} = \frac{1}{A(R)}\iint_R f(x,y)dA$. Extending this definition to general regions D, we have $f_{\text{avg}} = \frac{1}{A(D)}\iint_D f(x,y)dA$.

Here $D = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le 3x\}$, so $A(D) = \frac{1}{2}(1)(3) = \frac{3}{2}$ and

$$f_{\text{avg}} = \frac{1}{A(D)}\iint_D f(x,y)dA = \frac{1}{3/2}\int_0^1\int_0^{3x} xy\,dy\,dx$$
$$= \frac{2}{3}\int_0^1 \left[\tfrac{1}{2}xy^2\right]_{y=0}^{y=3x} dx = \frac{1}{3}\int_0^1 9x^3\,dx = \left.\tfrac{3}{4}x^4\right]_0^1 = \frac{3}{4}$$

72. Here $D = \left\{(x,y) \mid 0 \le x \le 1, 0 \le y \le x^2\right\}$, so $A(D) = \int_0^1 x^2\,dx = \left.\frac{1}{3}x^3\right]_0^1 = \frac{1}{3}$ and

$$f_{\text{avg}} = \frac{1}{A(D)}\iint_D f(x,y)dA = \frac{1}{1/3}\int_0^1\int_0^{x^2} x\sin y\,dy\,dx$$
$$= 3\int_0^1 \left[-x\cos y\right]_{y=0}^{y=x^2} dx$$
$$= 3\int_0^1 \left[x - x\cos(x^2)\right] dx = 3\left[\tfrac{1}{2}x^2 - \tfrac{1}{2}\sin(x^2)\right]_0^1$$
$$= 3\left(\tfrac{1}{2} - \tfrac{1}{2}\sin 1 - 0\right) = \tfrac{3}{2}(1 - \sin 1)$$

73. Since $m \le f(x,y) \le M$, $\iint_D m\,dA \le \iint_D f(x,y)\,dA \le \iint_D M\,dA$ by (7) $\quad \Rightarrow$ $m\iint_D 1\,dA \le \iint_D f(x,y)\,dA \le M\iint_D 1\,dA$ by (6) $\quad \Rightarrow \quad m \cdot A(D) \le \iint_D f(x,y)\,dA \le M \cdot A(D)$ by (9).

74.

$$\iint_D f(x,y)\,dA = \int_0^1\int_0^{2y} f(x,y)\,dx\,dy + \int_1^3\int_0^{3-y} f(x,y)\,dx\,dy$$
$$= \int_0^2\int_{x/2}^{3-x} f(x,y)\,dy\,dx$$

75.

First we can write $\iint_D (x+2)\,dA = \iint_D x\,dA + \iint_D 2\,dA$. But $f(x,y) = x$ is an odd function with respect to x [that is, $f(-x,y) = -f(x,y)$] and D is symmetric with respect to x. Consequently, the volume above D and below the graph of f is the same as the volume below D and above the graph of f, so

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\iint_D x\,dA = 0$. Also, $\iint_D 2\,dA = 2 \cdot A(D) = 2 \cdot \frac{1}{2}\pi(3)^2 = 9\pi$ since D is a half disk of radius 3. Thus $\iint_D (x+2)\,dA = 0 + 9\pi = 9\pi$.

76. The graph of $f(x, y) = \sqrt{R^2 - x^2 - y^2}$ is the top half of the sphere $x^2 + y^2 + z^2 = R^2$, centered at the origin with radius R, and D is the disk in the xy-plane also centered at the origin with radius R. Thus $\iint_D \sqrt{R^2 - x^2 - y^2}\,dA$ represents the volume of a half ball of radius R which is $\frac{1}{2} \cdot \frac{4}{3}\pi R^3 = \frac{2}{3}\pi R^3$.

77. We can write $\iint_D (2x + 3y)\,dA = \iint_D 2x\,dA + \iint_D 3y\,dA$. $\iint_D 2x\,dA$ represents the volume of the solid lying under the plane $z = 2x$ and above the rectangle D. This solid region is a triangular cylinder with length b and whose cross-section is a triangle with width a and height $2a$. (See the first figure.)

Thus its volume is $\frac{1}{2} \cdot a \cdot 2a \cdot b = a^2 b$. Similarly, $\iint_D 3y\,dA$ represents the volume of a triangular cylinder with length a, triangular cross-section with width b and height $3b$, and volume $\frac{1}{2} \cdot b \cdot 3b \cdot a = \frac{3}{2}ab^2$. (See the second figure.) Thus

$$\iint_D (2x + 3y)\,dA = a^2 b + \tfrac{3}{2}ab^2$$

78.

In the first quadrant, x and y are positive and the boundary of D is $x + y = 1$. But D is symmetric with respect to both axes because of the absolute values, so the region of integration is the square shown at the left. To evaluate the double integral, we first write $\iint_D (2 + x^2y^3 - y^2 \sin x)\,dA = \iint_D 2\,dA + \iint_D x^2y^3\,dA - \iint_D y^2 \sin x\,dA$. Now $f(x, y) = x^2y^3$ is odd with respect to y [that is, $f(x, -y) = -f(x, y)$] and D is symmetric with respect to y, so $\iint_D x^2y^3\,dA = 0$.

Similarly, $g(x, y) = y^2 \sin x$ is odd with respect to x [since $g(-x, y) = -g(x, y)$] and D is symmetric with respect to x, so $\iint_D y^2 \sin x\,dA = 0$. D is a square with side length $\sqrt{2}$, so $\iint_D 2\,dA = 2 \cdot A(D) = 2\left(\sqrt{2}\right)^2 = 4$, and $\iint_D (2 + x^2y^3 - y^2 \sin x)\,dA = 4 + 0 + 0 = 4$.

79. $\iint_D \left(ax^3 + by^3 + \sqrt{a^2 - x^2}\right) dA = \iint_D ax^3\,dA + \iint_D by^3\,dA + \iint_D \sqrt{a^2 - x^2}\,dA$. Now ax^3 is odd with respect to x and by^3 is odd with respect to y, and the region of integration is symmetric with respect to both x and y, so $\iint_D ax^3\,dA = \iint_D by^3\,dA = 0$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\iint_D \sqrt{a^2 - x^2}\, dA$ represents the volume of the solid region under the graph of $z = \sqrt{a^2 - x^2}$ and above the rectangle D, namely a half circular cylinder with radius a and length $2b$ (see the figure) whose volume is $\frac{1}{2} \cdot \pi r^2 h = \frac{1}{2}\pi a^2 (2b) = \pi a^2 b$. Thus $\iint_D \left(ax^3 + by^3 + \sqrt{a^2 - x^2}\right) dA = 0 + 0 + \pi a^2 b = \pi a^2 b$.

80. By the Extreme Value Theorem (14.7.8), f has an absolute minimum value m and an absolute maximum value M in D. Then by Property 15.2.10, $mA(D) \le \iint_D f(x,y)\, dA \le MA(D)$. Dividing through by the positive number $A(D)$, we get $m \le \dfrac{1}{A(D)} \displaystyle\iint_D f(x,y)\, dA \le M$. This says that the average value of f over D lies between m and M. But f is continuous on D and takes on the values m and M, and so by the Intermediate Value Theorem must take on all values between m and M. Specifically, there exists a point (x_0, y_0) in D such that $f(x_0, y_0) = \dfrac{1}{A(D)} \displaystyle\iint_D f(x,y)\, dA$ or equivalently $\iint_D f(x,y)\, dA = f(x_0, y_0)\, A(D)$.

81. For each r such that D_r lies within the domain, $A(D_r) = \pi r^2$, and by the Mean Value Theorem for double integrals there exists (x_r, y_r) in D_r such that $f(x_r, y_r) = \dfrac{1}{\pi r^2} \displaystyle\iint_{D_r} f(x,y)\, dA$. But $\displaystyle\lim_{r \to 0^+} (x_r, y_r) = (a, b)$, so $\displaystyle\lim_{r \to 0^+} \frac{1}{\pi r^2} \iint_{D_r} f(x,y)\, dA = \lim_{r \to 0^+} f(x_r, y_r) = f(a,b)$ by the continuity of f.

82. To find the equations of the boundary curves, we require that the z-values of the two surfaces be the same. In Maple, we use the command `solve(4-x^2-y^2=1-x-y,y);` and in Mathematica, we use `Solve[4-x^2-y^2==1-x-y,y]`. We find that the curves have equations $y = \dfrac{1 \pm \sqrt{13 + 4x - 4x^2}}{2}$. To find the two points of intersection of these curves, we use the CAS to solve $13 + 4x - 4x^2 = 0$, finding that $x = \frac{1 \pm \sqrt{14}}{2}$. So, using the CAS to evaluate the integral, the volume of intersection is

$$V = \int_{(1-\sqrt{14})/2}^{(1+\sqrt{14})/2} \int_{\left(1-\sqrt{13+4x-4x^2}\right)/2}^{\left(1+\sqrt{13+4x-4x^2}\right)/2} [(4 - x^2 - y^2) - (1 - x - y)]\, dy\, dx = \frac{49\pi}{8}$$

15.3 Double Integrals in Polar Coordinates

1. The region R is more easily described with polar coordinates: $R = \{(r,\theta) \mid 0 \le r \le 4,\ 0 \le \theta \le 3\pi/2\}$. Thus, $\iint_R f(x,y)\, dA = \int_0^{3\pi/2} \int_0^4 f(r\cos\theta, r\sin\theta)\, r\, dr\, d\theta$.

2. The region R is more easily described by rectangular coordinates: $R = \{(x,y) \mid -1 \le x \le 1,\ -x \le y \le 1\}$. Thus, $\iint_R f(x,y)\, dA = \int_{-1}^1 \int_{-x}^1 f(x,y)\, dy\, dx$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. The region R is more easily described with polar coordinates: $R = \{(r,\theta) \mid 1 \le r \le 3, 0 \le \theta \le \pi\}$.

Thus, $\iint_R f(x,y)\,dA = \int_0^{\pi} \int_1^3 f(r\cos\theta, r\sin\theta)\, r\,dr\,d\theta$.

4. The region R is more easily described by polar coordinates: $R = \left\{(r,\theta) \mid 0 \le r \le 3, -\frac{\pi}{4} \le \theta \le \frac{3\pi}{4}\right\}$.

Thus, $\iint_R f(x,y)\,dA = \int_{-\pi/4}^{3\pi/4} \int_0^3 f(r\cos\theta, r\sin\theta)\, r\,dr\,d\theta$.

5. The region R is more easily described with rectangular coordinates: $R = \{(x,y) \mid 2y-2 \le x \le -2y+2, 0 \le y \le 1\}$.

Thus, $\iint_R f(x,y)\,dA = \int_0^1 \int_{2y-2}^{-2y+2} f(x,y)\,dx\,dy$.

6. The region R is more easily described with polar coordinates: $R = \{(r,\theta) \mid 8 \le r \le 10, 0 \le \theta \le 2\pi\}$.

Thus, $\iint_R f(x,y)\,dA = \int_0^{2\pi} \int_8^{10} f(r\cos\theta, r\sin\theta)\, r\,dr\,d\theta$.

7. The integral $\int_{\pi/4}^{3\pi/4} \int_1^2 r\,dr\,d\theta$ represents the area of the region $R = \{(r,\theta) \mid 1 \le r \le 2, \pi/4 \le \theta \le 3\pi/4\}$, the top quarter portion of a ring (annulus).

$$\int_{\pi/4}^{3\pi/4} \int_1^2 r\,dr\,d\theta = \left(\int_{\pi/4}^{3\pi/4} d\theta\right)\left(\int_1^2 r\,dr\right)$$
$$= \left[\theta\right]_{\pi/4}^{3\pi/4} \left[\tfrac{1}{2}r^2\right]_1^2 = \left(\tfrac{3\pi}{4} - \tfrac{\pi}{4}\right) \cdot \tfrac{1}{2}(4-1) = \tfrac{\pi}{2} \cdot \tfrac{3}{2} = \tfrac{3\pi}{4}$$

8. The integral $\int_{\pi/2}^{\pi} \int_0^{2\sin\theta} r\,dr\,d\theta$ represents the area of the region $R = \{(r,\theta) \mid 0 \le r \le 2\sin\theta, \pi/2 \le \theta \le \pi\}$. Since $r = 2\sin\theta \;\Rightarrow\; r^2 = 2r\sin\theta \;\Leftrightarrow\; x^2 + y^2 = 2y \;\Leftrightarrow$ $x^2 + (y-1)^2 = 1$, R is the portion in the second quadrant of a disk of radius 1 with center $(0,1)$.

$$\int_{\pi/2}^{\pi} \int_0^{2\sin\theta} r\,dr\,d\theta = \int_{\pi/2}^{\pi} \left[\tfrac{1}{2}r^2\right]_{r=0}^{r=2\sin\theta} d\theta = \int_{\pi/2}^{\pi} 2\sin^2\theta\,d\theta$$
$$= \int_{\pi/2}^{\pi} 2 \cdot \tfrac{1}{2}(1-\cos 2\theta)\,d\theta = \left[\theta - \tfrac{1}{2}\sin 2\theta\right]_{\pi/2}^{\pi}$$
$$= \pi - 0 - \tfrac{\pi}{2} + 0 = \tfrac{\pi}{2}$$

9. The half-disk D can be described in polar coordinates as $D = \{(r,\theta) \mid 0 \le r \le 5, 0 \le \theta \le \pi\}$. Then

$$\iint_D x^2 y\,dA = \int_0^{\pi} \int_0^5 (r\cos\theta)^2 (r\sin\theta)\, r\,dr\,d\theta = \left(\int_0^{\pi} \cos^2\theta \sin\theta\,d\theta\right)\left(\int_0^5 r^4\,dr\right)$$
$$= \left[-\tfrac{1}{3}\cos^3\theta\right]_0^{\pi} \left[\tfrac{1}{5}r^5\right]_0^5 = -\tfrac{1}{3}(-1-1) \cdot 625 = \tfrac{1250}{3}$$

10. The region R is $\frac{1}{8}$ of a disk, as shown in the figure, and can be described by $R = \{(r,\theta) \mid 0 \le r \le 2, \pi/4 \le \theta \le \pi/2\}$. Thus

$$\iint_R (2x - y)\,dA = \int_{\pi/4}^{\pi/2} \int_0^2 (2r\cos\theta - r\sin\theta)\, r\,dr\,d\theta$$
$$= \int_{\pi/4}^{\pi/2} (2\cos\theta - \sin\theta)\,d\theta \int_0^2 r^2\,dr$$
$$= \left[2\sin\theta + \cos\theta\right]_{\pi/4}^{\pi/2} \left[\tfrac{1}{3}r^3\right]_0^2$$
$$= \left(2 + 0 - \sqrt{2} - \tfrac{\sqrt{2}}{2}\right)\left(\tfrac{8}{3}\right) = \tfrac{16}{3} - 4\sqrt{2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. $\iint_R \sin(x^2+y^2)\,dA = \int_0^{\pi/2}\int_1^3 \sin(r^2)\,r\,dr\,d\theta = \int_0^{\pi/2} d\theta \int_1^3 r\sin(r^2)\,dr = \left[\theta\right]_0^{\pi/2}\left[-\frac{1}{2}\cos(r^2)\right]_1^3$

$= \left(\frac{\pi}{2}\right)\left[-\frac{1}{2}(\cos 9 - \cos 1)\right] = \frac{\pi}{4}(\cos 1 - \cos 9)$

12. $\displaystyle\iint_R \frac{y^2}{x^2+y^2}\,dA = \int_0^{2\pi}\int_a^b \frac{(r\sin\theta)^2}{r^2}\,r\,dr\,d\theta = \int_0^{2\pi}\sin^2\theta\,d\theta\int_a^b r\,dr = \int_0^{2\pi}\frac{1}{2}(1-\cos 2\theta)\,d\theta\int_a^b r\,dr$

$= \frac{1}{2}\left[\theta - \frac{1}{2}\sin 2\theta\right]_0^{2\pi}\left[\frac{1}{2}r^2\right]_a^b = \frac{1}{2}(2\pi - 0 - 0)\cdot\frac{1}{2}\left(b^2-a^2\right) = \frac{\pi}{2}(b^2-a^2)$

13. $\iint_D e^{-x^2-y^2}\,dA = \int_{-\pi/2}^{\pi/2}\int_0^2 e^{-r^2}\,r\,dr\,d\theta = \int_{-\pi/2}^{\pi/2} d\theta\int_0^2 re^{-r^2}\,dr$

$= \left[\theta\right]_{-\pi/2}^{\pi/2}\left[-\frac{1}{2}e^{-r^2}\right]_0^2 = \pi\left(-\frac{1}{2}\right)(e^{-4}-e^0) = \frac{\pi}{2}(1-e^{-4})$

14. $\iint_D \cos\sqrt{x^2+y^2}\,dA = \int_0^{2\pi}\int_0^2 \cos\sqrt{r^2}\,r\,dr\,d\theta = \int_0^{2\pi} d\theta\int_0^2 r\cos r\,dr$. For the second integral, integrate by parts with $u = r$, $dv = \cos r\,dr$. Then $\iint_D \cos\sqrt{x^2+y^2}\,dA = \left[\theta\right]_0^{2\pi}\left[r\sin r + \cos r\right]_0^2 = 2\pi(2\sin 2 + \cos 2 - 1)$.

15. R is the region shown in the figure, and can be described by $R = \{(r,\theta) \mid 0 \le \theta \le \pi/4, 1 \le r \le 2\}$. Thus $\iint_R \arctan(y/x)\,dA = \int_0^{\pi/4}\int_1^2 \arctan(\tan\theta)\,r\,dr\,d\theta$ since $y/x = \tan\theta$. Also, $\arctan(\tan\theta) = \theta$ for $0 \le \theta \le \pi/4$, so the integral becomes $\int_0^{\pi/4}\int_1^2 \theta\,r\,dr\,d\theta = \int_0^{\pi/4}\theta\,d\theta\int_1^2 r\,dr = \left[\frac{1}{2}\theta^2\right]_0^{\pi/4}\left[\frac{1}{2}r^2\right]_1^2 = \frac{\pi^2}{32}\cdot\frac{3}{2} = \frac{3}{64}\pi^2$.

16.

$$\iint_D x\,dA = \iint_{\substack{x^2+y^2\le 4\\ x\ge 0,\, y\ge 0}} x\,dA - \iint_{\substack{(x-1)^2+y^2\le 1\\ y\ge 0}} x\,dA$$

$= \int_0^{\pi/2}\int_0^2 r^2\cos\theta\,dr\,d\theta - \int_0^{\pi/2}\int_0^{2\cos\theta} r^2\cos\theta\,dr\,d\theta$

$= \int_0^{\pi/2}\frac{1}{3}(8\cos\theta)\,d\theta - \int_0^{\pi/2}\frac{1}{3}(8\cos^4\theta)\,d\theta$

$= \frac{8}{3}[\sin\theta]_0^{\pi/2} - \frac{8}{12}\left[\cos^3\theta\sin\theta + \frac{3}{2}(\theta + \sin\theta\cos\theta)\right]_0^{\pi/2}$

$= \frac{8}{3} - \frac{2}{3}\left[0 + \frac{3}{2}\left(\frac{\pi}{2}\right)\right] = \frac{16-3\pi}{6}$

17. By symmetry, the area of the region is 4 times the area of the region D in the first quadrant enclosed by the cardiod $r = 1-\cos\theta$ (see the figure). Here $D = \{(r,\theta) \mid 0 \le r \le 1-\cos\theta, 0 \le \theta \le \pi/2\}$, so the total area is

$4A(D) = 4\iint_D dA = 4\int_0^{\pi/2}\int_0^{1-\cos\theta} r\,dr\,d\theta = 4\int_0^{\pi/2}\left[\frac{1}{2}r^2\right]_{r=0}^{r=1-\cos\theta} d\theta$

$= 2\int_0^{\pi/2}(1-\cos\theta)^2 d\theta = 2\int_0^{\pi/2}(1 - 2\cos\theta + \cos^2\theta)\,d\theta$

$= 2\int_0^{\pi/2}\left[1 - 2\cos\theta + \frac{1}{2}(1+\cos 2\theta)\right]d\theta$

$= 2\left[\theta - 2\sin\theta + \frac{1}{2}\theta + \frac{1}{4}\sin 2\theta\right]_0^{\pi/2}$

$= 2\left(\frac{\pi}{2} - 2 + \frac{\pi}{4}\right) = \frac{3\pi}{2} - 4$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. The region D is described by $D = \{(r,\theta) \mid 0 \le r \le \sqrt{\theta},\, 0 \le \theta \le 2\pi\}$, so the area is

$$A(D) = \int_0^{2\pi} \int_0^{\sqrt{\theta}} r\,dr\,d\theta = \int_0^{2\pi} \left[\frac{r^2}{2}\right]_{r=0}^{r=\sqrt{\theta}} d\theta = \int_0^{2\pi} \frac{\theta}{2}\,d\theta = \left[\frac{\theta^2}{4}\right]_0^{2\pi} = \pi^2.$$

19. By symmetry, the total area is twice the area defined by

$D = \{(r,\theta) \mid 0 \le r \le \sin\theta,\, \pi/4 \le \theta \le \pi\}$ (see the figure).

The total area is

$$\begin{aligned} 2A(D) &= 2\textstyle\int_{\pi/4}^{\pi}\int_0^{\sin\theta} r\,dr\,d\theta = 2\cdot\frac{1}{2}\int_{\pi/4}^{\pi}\left[r^2\right]_{r=0}^{r=\sin\theta} d\theta = \int_{\pi/4}^{\pi}\sin^2\theta\,d\theta \\ &= \textstyle\int_{\pi/4}^{\pi}\frac{1}{2}(1-\cos 2\theta)\,d\theta = \frac{1}{2}\left[\theta - \frac{1}{2}\sin 2\theta\right]_{\pi/4}^{\pi} \\ &= \textstyle\frac{1}{2}(\pi - 0) - \frac{1}{2}\left(\frac{\pi}{4} - \frac{1}{2}\right) = \frac{3\pi}{8} + \frac{1}{4} \end{aligned}$$

20. By symmetry, the area of the region is 4 times the area of the region D in the first quadrant between the circle $r = 1/\sqrt{2}$ and the curve $r^2 = \cos 2\theta \;\Rightarrow\; r = \sqrt{\cos 2\theta}$. The curves intersect in the first quadrant when $\cos 2\theta = \left(\frac{1}{\sqrt{2}}\right)^2 \;\Rightarrow\; \cos 2\theta = \frac{1}{2} \;\Rightarrow\; 2\theta = \frac{\pi}{3} \;\Rightarrow\; \theta = \frac{\pi}{6}$. Thus, $D = \{(r,\theta) \mid 1/\sqrt{2} \le r \le \sqrt{\cos 2\theta},\, 0 \le \theta \le \pi/6\}$, so the total area is

$$\begin{aligned} 4A(D) &= 4\int_0^{\pi/6}\int_{1/\sqrt{2}}^{\sqrt{\cos 2\theta}} r\,dr\,d\theta = 4\cdot\frac{1}{2}\int_0^{\pi/6}\left[r^2\right]_{r=1/\sqrt{2}}^{r=\sqrt{\cos 2\theta}} d\theta = 2\int_0^{\pi/6}\left[\cos 2\theta - \frac{1}{2}\right] d\theta \\ &= 2\left[\textstyle\frac{1}{2}\sin 2\theta - \frac{\theta}{2}\right]_0^{\pi/6} = \frac{\sqrt{3}}{2} - \frac{\pi}{6} \end{aligned}$$

21. One loop is given by the region

$D = \{(r,\theta) \mid 0 \le r \le \sin 3\theta,\, 0 \le \theta \le \pi/3\}$, so the area is

$$\begin{aligned} \textstyle\iint_D dA &= \textstyle\int_0^{\pi/3}\int_0^{\sin 3\theta} r\,dr\,d\theta = \frac{1}{2}\int_0^{\pi/3}\left[r^2\right]_{r=0}^{r=\sin 3\theta} d\theta \\ &= \textstyle\frac{1}{2}\int_0^{\pi/3}\sin^2 3\theta\,d\theta = \frac{1}{2}\int_0^{\pi/3}\frac{1}{2}(1-\cos 6\theta)\,d\theta \\ &= \textstyle\frac{1}{4}\left[\theta - \frac{1}{6}\sin 6\theta\right]_0^{\pi/3} = \frac{\pi}{12} \end{aligned}$$

22. In polar coordinates the circle $(x-1)^2 + y^2 = 1 \;\Leftrightarrow\; x^2 + y^2 = 2x$ is $r^2 = 2r\cos\theta \;\Rightarrow\; r = 2\cos\theta$, and the circle $x^2 + y^2 = 1$ is $r = 1$. The curves intersect in the first quadrant when $2\cos\theta = 1 \;\Rightarrow\; \cos\theta = \frac{1}{2} \;\Rightarrow\; \theta = \pi/3$, so the portion of the region in the first quadrant is given by

$D = \{(r,\theta) \mid 1 \le r \le 2\cos\theta,\, 0 \le \theta \le \pi/3\}$. By symmetry, the total area is twice the area of D:

$$\begin{aligned} 2A(D) &= 2\textstyle\iint_D dA = 2\int_0^{\pi/3}\int_1^{2\cos\theta} r\,dr\,d\theta = 2\int_0^{\pi/3}\left[\frac{1}{2}r^2\right]_{r=1}^{r=2\cos\theta} d\theta \\ &= \textstyle\int_0^{\pi/3}\left(4\cos^2\theta - 1\right) d\theta = \int_0^{\pi/3}\left[4\cdot\frac{1}{2}(1+\cos 2\theta) - 1\right] d\theta \\ &= \textstyle\int_0^{\pi/3}(1 + 2\cos 2\theta)\,d\theta = \left[\theta + \sin 2\theta\right]_0^{\pi/3} = \frac{\pi}{3} + \frac{\sqrt{3}}{2} \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. (a) $V = \iint_D (1 + xy)\, dA$, where D is the portion of the circle $x^2 + y^2 = 4$ in the first quadrant. Thus,

$V = \iint_D (1 + xy)\, dA = \int_0^{\pi/2} \int_0^2 (1 + r^2 \cos\theta \sin\theta) r\, dr\, d\theta.$

(b) $$\int_0^{\pi/2} \int_0^2 (1 + r^2 \cos\theta \sin\theta) r\, dr\, d\theta = \int_0^{\pi/2} \int_0^2 \left(r + r^3 \cos\theta \sin\theta\right)\, dr\, d\theta = \int_0^{\pi/2} \left[\frac{r^2}{2} + \frac{r^4}{4} \cos\theta \sin\theta\right]_{r=0}^{r=2} d\theta$$

$$= \int_0^{\pi/2} (2 + 4\cos\theta \sin\theta)\, d\theta = \left[2\theta + 2\sin^2\theta\right]_0^{\pi/2} \qquad [u = \sin\theta,\, du = \cos\theta\, d\theta]$$

$$= \pi + 2$$

24. (a) $V = \iint_D (x^2 + y^2)\, dA$, where D is the region on or between the circles $x^2 + y^2 = 1$ and $x^2 + y^2 = 4$. Thus,

$V = \iint_{1 \le x^2 + y^2 \le 4} (x^2 + y^2)\, dA = \int_0^{2\pi} \int_1^2 r^2 r\, dr\, d\theta.$

(b) $$\int_0^{2\pi} \int_1^2 r^3\, dr\, d\theta = \int_0^{2\pi} d\theta \int_1^2 r^3\, dr = \Big[\theta\Big]_{\theta=0}^{\theta=2\pi} \cdot \frac{1}{4}\Big[r^4\Big]_{r=1}^{r=2} = 2\pi \cdot \frac{1}{4}(16 - 1) = \frac{15\pi}{2}$$

25. (a) $V = \iint_D y\, dA$, where D is the portion of the circle $x^2 + y^2 = 9$ in quadrants I–III. Thus,

$V = \iint_D y\, dA = \int_0^{3\pi/2} \int_0^3 (r \sin\theta)\, r\, dr\, d\theta.$

(b) $$\int_0^{3\pi/2} \int_0^3 r^2 \sin\theta\, dr\, d\theta = \int_0^{3\pi/2} \sin\theta\, d\theta \int_0^3 r^2\, dr = \Big[-\cos\theta\Big]_{\theta=0}^{\theta=3\pi/2} \left[\frac{r^3}{3}\right]_{r=0}^{r=3} = [0 - (-1)](9 - 0) = 9$$

26. (a) $V = \iint_D xy^2\, dA$, where D is the region on or between the circles $r = 2$ and $r = 3$ in the first quadrant. Thus,

$V = \iint_D xy^2\, dA = \int_0^{\pi/2} \int_2^3 (r\cos\theta)(r\sin\theta)^2\, r\, dr\, d\theta.$

(b) $$\int_0^{\pi/2} \int_2^3 r^4 \cos\theta \sin^2\theta\, dr\, d\theta = \int_0^{\pi/2} \cos\theta \sin^2\theta\, d\theta \int_2^3 r^4\, dr = \left[\frac{\sin^3\theta}{3}\right]_{\theta=0}^{\theta=\pi/2} \left[\frac{r^5}{5}\right]_{r=2}^{r=3}$$

$$= \left(\tfrac{1}{3} - 0\right)\left(\tfrac{243}{5} - \tfrac{32}{5}\right) = \tfrac{211}{15}$$

27. (a) The region is described by $D = \{(r, \theta) \mid 0 \le r \le \sin\theta,\, 0 \le \theta \le \pi/2\}$, so $\iint_D x\, dA = \int_0^{\pi/2} \int_0^{\sin\theta} (r\cos\theta)\, r\, dr\, d\theta.$

(b) $$\int_0^{\pi/2} \int_0^{\sin\theta} r^2 \cos\theta\, dr\, d\theta = \int_0^{\pi/2} \cos\theta \left[\frac{r^3}{3}\right]_{r=0}^{r=\sin\theta} d\theta = \frac{1}{3}\int_0^{\pi/2} \cos\theta \sin^3\theta\, d\theta$$

$$= \tfrac{1}{3}\int_0^1 u^3\, du \qquad [u = \sin\theta,\, du = \cos\theta\, d\theta]$$

$$= \tfrac{1}{3}\left[\tfrac{1}{4}u^4\right]_0^1 = \tfrac{1}{12}$$

28. (a) The region is described by $D = \{(r, \theta) \mid 0 \le r \le 1 + \cos\theta,\, 0 \le \theta \le \pi\}$, so $\iint_D dA = \int_0^{\pi} \int_0^{1+\cos\theta} 1 \cdot r\, dr\, d\theta.$

(b) $$\int_0^{\pi} \int_0^{1+\cos\theta} r\, dr\, d\theta = \int_0^{\pi} \left[\frac{r^2}{2}\right]_{r=0}^{r=1+\cos\theta} d\theta = \frac{1}{2}\int_0^{\pi} (1 + \cos\theta)^2\, d\theta = \frac{1}{2}\int_0^{\pi} \left(1 + 2\cos\theta + \cos^2\theta\right)\, d\theta$$

$$= \tfrac{1}{2}\int_0^{\pi} \left[1 + 2\cos\theta + \tfrac{1}{2}(1 + \cos 2\theta)\right]\, d\theta = \tfrac{1}{2}\int_0^{\pi} \left[\tfrac{3}{2} + 2\cos\theta + \tfrac{1}{2}\cos 2\theta\right]\, d\theta$$

$$= \tfrac{1}{2}\left[\tfrac{3\theta}{2} + 2\sin\theta + \tfrac{1}{4}\sin 2\theta\right]_0^{\pi} = \tfrac{3\pi}{4}$$

29. $V = \iint_{x^2 + y^2 \le 25} \left(x^2 + y^2\right) dA = \int_0^{2\pi} \int_0^5 r^2 \cdot r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_0^5 r^3\, dr = \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{4}r^4\right]_0^5 = 2\pi\left(\tfrac{625}{4}\right) = \tfrac{625}{2}\pi$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $V = \iint_{1 \le x^2+y^2 \le 4} \sqrt{x^2+y^2}\, dA = \int_0^{2\pi}\int_1^2 \sqrt{r^2}\; r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_1^2 r^2\, dr = \left[\,\theta\,\right]_0^{2\pi}\left[\tfrac{1}{3}r^3\right]_1^2 = 2\pi\left(\tfrac{8}{3}-\tfrac{1}{3}\right) = \tfrac{14}{3}\pi$

31. $2x+y+z=4 \quad\Leftrightarrow\quad z = 4-2x-y$, so the volume of the solid is

$$\begin{aligned} V &= \iint_{x^2+y^2\le 1} (4-2x-y)\, dA = \int_0^{2\pi}\int_0^1 (4-2r\cos\theta - r\sin\theta)\, r\, dr\, d\theta \\ &= \int_0^{2\pi}\int_0^1 \left[4r - r^2\,(2\cos\theta+\sin\theta)\right] dr\, d\theta = \int_0^{2\pi}\left[2r^2 - \tfrac{1}{3}r^3\,(2\cos\theta+\sin\theta)\right]_{r=0}^{r=1} d\theta \\ &= \int_0^{2\pi}\left[2 - \tfrac{1}{3}\,(2\cos\theta+\sin\theta)\right] d\theta = \left[2\theta - \tfrac{1}{3}\,(2\sin\theta-\cos\theta)\right]_0^{2\pi} = 4\pi + \tfrac{1}{3} - 0 - \tfrac{1}{3} = 4\pi \end{aligned}$$

32. The sphere $x^2+y^2+z^2=16$ intersects the xy-plane in the circle $x^2+y^2=16$, so

$$\begin{aligned} V &= 2\iint_{4\le x^2+y^2\le 16} \sqrt{16-x^2-y^2}\, dA \quad \text{[by symmetry]} \quad = 2\int_0^{2\pi}\int_2^4 \sqrt{16-r^2}\; r\, dr\, d\theta \\ &= 2\int_0^{2\pi} d\theta \int_2^4 r(16-r^2)^{1/2} dr = 2\left[\,\theta\,\right]_0^{2\pi}\left[-\tfrac{1}{3}(16-r^2)^{3/2}\right]_2^4 \\ &= -\tfrac{2}{3}(2\pi)(0-12^{3/2}) = \tfrac{4\pi}{3}\left(12\sqrt{12}\right) = 32\sqrt{3}\,\pi \end{aligned}$$

33. By symmetry,

$$\begin{aligned} V &= 2\iint_{x^2+y^2\le a^2} \sqrt{a^2-x^2-y^2}\, dA = 2\int_0^{2\pi}\int_0^a \sqrt{a^2-r^2}\; r\, dr\, d\theta = 2\int_0^{2\pi} d\theta \int_0^a r\sqrt{a^2-r^2}\, dr \\ &= 2\left[\,\theta\,\right]_0^{2\pi}\left[-\tfrac{1}{3}(a^2-r^2)^{3/2}\right]_0^a = 2(2\pi)\left(0+\tfrac{1}{3}a^3\right) = \tfrac{4}{3}\pi a^3 \end{aligned}$$

34. The paraboloid $z = 1+2x^2+2y^2$ intersects the plane $z=7$ when $7 = 1+2x^2+2y^2$ or $x^2+y^2=3$ and we are restricted to the first octant, so

$$\begin{aligned} V &= \iint_{\substack{x^2+y^2\le 3,\\ x\ge 0, y\ge 0}} \left[7-\left(1+2x^2+2y^2\right)\right] dA = \int_0^{\pi/2}\int_0^{\sqrt{3}} \left[7-(1+2r^2)\right] r\, dr\, d\theta \\ &= \int_0^{\pi/2} d\theta \int_0^{\sqrt{3}} \left(6r-2r^3\right) dr = \left[\,\theta\,\right]_0^{\pi/2}\left[3r^2-\tfrac{1}{2}r^4\right]_0^{\sqrt{3}} = \tfrac{\pi}{2}\cdot\tfrac{9}{2} = \tfrac{9}{4}\pi \end{aligned}$$

35. The cone $z=\sqrt{x^2+y^2}$ intersects the sphere $x^2+y^2+z^2=1$ when $x^2+y^2+\left(\sqrt{x^2+y^2}\right)^2 = 1$ or $x^2+y^2=\tfrac{1}{2}$. So

$$\begin{aligned} V &= \iint_{x^2+y^2\le 1/2} \left(\sqrt{1-x^2-y^2}-\sqrt{x^2+y^2}\right) dA = \int_0^{2\pi}\int_0^{1/\sqrt{2}} \left(\sqrt{1-r^2}-r\right) r\, dr\, d\theta \\ &= \int_0^{2\pi} d\theta \int_0^{1/\sqrt{2}} \left(r\sqrt{1-r^2}-r^2\right) dr = \left[\,\theta\,\right]_0^{2\pi}\left[-\tfrac{1}{3}(1-r^2)^{3/2}-\tfrac{1}{3}r^3\right]_0^{1/\sqrt{2}} = 2\pi\left(-\tfrac{1}{3}\right)\left(\tfrac{1}{\sqrt{2}}-1\right) = \tfrac{\pi}{3}\left(2-\sqrt{2}\right) \end{aligned}$$

36. The two paraboloids intersect when $6-x^2-y^2 = 2x^2+2y^2$ or $x^2+y^2=2$. For $x^2+y^2\le 2$, the paraboloid $z = 6-x^2-y^2$ is above $z=2x^2+2y^2$ so

$$\begin{aligned} V &= \iint_{x^2+y^2\le 2} \left[(6-x^2-y^2)-(2x^2+2y^2)\right] dA = \iint_{x^2+y^2\le 2} \left[6-3(x^2+y^2)\right] dA = \int_0^{2\pi}\int_0^{\sqrt{2}} (6-3r^2)\, r\, dr\, d\theta \\ &= \int_0^{2\pi} d\theta \int_0^{\sqrt{2}} (6r-3r^3)\, dr = \left[\,\theta\,\right]_0^{2\pi}\left[3r^2-\tfrac{3}{4}r^4\right]_0^{\sqrt{2}} = 2\pi\,(6-3) = 6\pi \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. The given solid is the region inside the cylinder $x^2+y^2=4$ between the surfaces $z=\sqrt{64-4x^2-4y^2}$ and $z=-\sqrt{64-4x^2-4y^2}$. So

$$V=\iint\limits_{x^2+y^2\le 4}\left[\sqrt{64-4x^2-4y^2}-\left(-\sqrt{64-4x^2-4y^2}\right)\right]dA=\iint\limits_{x^2+y^2\le 4}2\cdot 2\sqrt{16-x^2-y^2}\,dA$$

$$=4\int_0^{2\pi}\int_0^2\sqrt{16-r^2}\;r\,dr\,d\theta=4\int_0^{2\pi}d\theta\int_0^2 r\sqrt{16-r^2}\,dr=4\left[\theta\right]_0^{2\pi}\left[-\tfrac{1}{3}(16-r^2)^{3/2}\right]_0^2$$

$$=8\pi\left(-\tfrac{1}{3}\right)\left(12^{3/2}-16^{2/3}\right)=\tfrac{8\pi}{3}\left(64-24\sqrt{3}\right)$$

38. (a) Here the region in the xy-plane is the annular region $r_1^2\le x^2+y^2\le r_2^2$ and the desired volume is twice that above the xy-plane. Hence

$$V=2\iint\limits_{r_1^2\le x^2+y^2\le r_2^2}\sqrt{r_2^2-x^2-y^2}\,dA=2\int_0^{2\pi}\int_{r_1}^{r_2}\sqrt{r_2^2-r^2}\;r\,dr\,d\theta=2\int_0^{2\pi}d\theta\int_{r_1}^{r_2}\sqrt{r_2^2-r^2}\;r\,dr$$

$$=2\,(2\pi)\left[-\tfrac{1}{3}(r_2^2-r^2)^{3/2}\right]_{r_1}^{r_2}=\tfrac{4\pi}{3}(r_2^2-r_1^2)^{3/2}$$

(b) A cross-sectional cut is shown in the figure. So $r_2^2=\left(\frac{1}{2}h\right)^2+r_1^2$ or $\frac{1}{4}h^2=r_2^2-r_1^2$.

Thus the volume in terms of h is $V=\frac{4\pi}{3}\left(\frac{1}{4}h^2\right)^{3/2}=\frac{\pi}{6}h^3$.

39.

$$\int_0^2\int_0^{\sqrt{4-x^2}}e^{-x^2-y^2}\,dy\,dx=\int_0^{\pi/2}\int_0^2 e^{-r^2}\,r\,dr\,d\theta$$

$$=\int_0^{\pi/2}d\theta\int_0^2 re^{-r^2}dr=\left[\theta\right]_0^{\pi/2}\left[-\tfrac{1}{2}e^{-r^2}\right]_0^2$$

$$=\tfrac{\pi}{2}\left[-\tfrac{1}{2}\left(e^{-4}-1\right)\right]=\tfrac{\pi}{4}(1-e^{-4})$$

40.

$$\int_0^a\int_{-\sqrt{a^2-y^2}}^{\sqrt{a^2-y^2}}(2x+y)\,dx\,dy=\int_0^{\pi}\int_0^a(2r\cos\theta+r\sin\theta)\,r\,dr\,d\theta$$

$$=\int_0^{\pi}(2\cos\theta+\sin\theta)\,d\theta\int_0^a r^2\,dr$$

$$=\left[2\sin\theta-\cos\theta\right]_0^{\pi}\left[\tfrac{1}{3}r^3\right]_0^a$$

$$=\left[(0+1)-(0-1)\right]\cdot\tfrac{1}{3}(a^3-0)=\tfrac{2}{3}a^3$$

41. The region D of integration is shown in the figure. In polar coordinates the line $x=\sqrt{3}\,y$ is $\theta=\pi/6$, so

$$\int_0^{1/2}\int_{\sqrt{3}\,y}^{\sqrt{1-y^2}}xy^2\,dx\,dy=\int_0^{\pi/6}\int_0^1(r\cos\theta)(r\sin\theta)^2\,r\,dr\,d\theta$$

$$=\int_0^{\pi/6}\sin^2\theta\cos\theta\,d\theta\int_0^1 r^4\,dr$$

$$=\left[\tfrac{1}{3}\sin^3\theta\right]_0^{\pi/6}\left[\tfrac{1}{5}r^5\right]_0^1$$

$$=\left[\tfrac{1}{3}\left(\tfrac{1}{2}\right)^3-0\right]\left[\tfrac{1}{5}-0\right]=\tfrac{1}{120}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42.

$$\int_0^2 \int_0^{\sqrt{2x-x^2}} \sqrt{x^2+y^2}\,dy\,dx = \int_0^{\pi/2} \int_0^{2\cos\theta} r \cdot r\,dr\,d\theta$$
$$= \int_0^{\pi/2} \left[\tfrac{1}{3}r^3\right]_{r=0}^{r=2\cos\theta} d\theta$$
$$= \int_0^{\pi/2} \left(\tfrac{8}{3}\cos^3\theta\right) d\theta$$
$$= \tfrac{8}{3}\int_0^{\pi/2} (1-\sin^2\theta)\cos\theta\,d\theta$$
$$= \tfrac{8}{3}\left[\sin\theta - \tfrac{1}{3}\sin^3\theta\right]_0^{\pi/2} = \tfrac{16}{9}$$

43. $D = \{(r,\theta) \mid 0 \le r \le 1, 0 \le \theta \le 2\pi\}$, so $\iint_D e^{(x^2+y^2)^2}\,dA = \int_0^{2\pi}\int_0^1 e^{(r^2)^2}\,r\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^1 re^{r^4}\,dr = 2\pi\int_0^1 re^{r^4}\,dr$. Using a calculator, we estimate $2\pi\int_0^1 re^{r^4}\,dr \approx 4.5951$.

44. $D = \{(r,\theta) \mid 0 \le r \le 1, 0 \le \theta \le \pi/2\}$, so

$$\iint_D xy\sqrt{1+x^2+y^2}\,dA = \int_0^{\pi/2}\int_0^1 (r\cos\theta)(r\sin\theta)\sqrt{1+r^2}\;r\,dr\,d\theta$$
$$= \int_0^{\pi/2} \sin\theta\cos\theta\,d\theta \int_0^1 r^3\sqrt{1+r^2}\,dr = \left[\tfrac{1}{2}\sin^2\theta\right]_0^{\pi/2} \int_0^1 r^3\sqrt{1+r^2}\,dr$$
$$= \tfrac{1}{2}\int_0^1 r^3\sqrt{1+r^2}\,dr \approx 0.1609$$

45. The surface of the water in the pool is a circular disk D with radius 20 ft. If we place D on coordinate axes with the origin at the center of D and define $f(x,y)$ to be the depth of the water at (x,y), then the volume of water in the pool is the volume of the solid that lies above $D = \{(x,y) \mid x^2+y^2 \le 400\}$ and below the graph of $f(x,y)$. We can associate north with the positive y-direction, so we are given that the depth is constant in the x-direction and the depth increases linearly in the y-direction from $f(0,-20) = 2$ to $f(0,20) = 7$. The trace in the yz-plane is a line segment from $(0,-20,2)$ to $(0,20,7)$. The slope of this line is $\frac{7-2}{20-(-20)} = \frac{1}{8}$, so an equation of the line is $z - 7 = \frac{1}{8}(y-20) \quad\Rightarrow\quad z = \frac{1}{8}y + \frac{9}{2}$. Since $f(x,y)$ is independent of x, $f(x,y) = \frac{1}{8}y + \frac{9}{2}$. Thus the volume is given by $\iint_D f(x,y)\,dA$, which is most conveniently evaluated using polar coordinates. Then $D = \{(r,\theta) \mid 0 \le r \le 20, 0 \le \theta \le 2\pi\}$ and substituting $x = r\cos\theta$, $y = r\sin\theta$ the integral becomes

$$\int_0^{2\pi}\int_0^{20} \left(\tfrac{1}{8}r\sin\theta + \tfrac{9}{2}\right) r\,dr\,d\theta = \int_0^{2\pi} \left[\tfrac{1}{24}r^3\sin\theta + \tfrac{9}{4}r^2\right]_{r=0}^{r=20} d\theta = \int_0^{2\pi} \left(\tfrac{1000}{3}\sin\theta + 900\right) d\theta$$
$$= \left[-\tfrac{1000}{3}\cos\theta + 900\theta\right]_0^{2\pi} = 1800\pi$$

Thus the pool contains $1800\pi \approx 5655$ ft^3 of water.

46. (a) If $R \le 100$, the total amount of water supplied each hour to the region within R feet of the sprinkler is

$$V = \int_0^{2\pi}\int_0^R e^{-r} r\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^R re^{-r}\,dr = \left[\theta\right]_0^{2\pi} \left[-re^{-r} - e^{-r}\right]_0^R$$
$$= 2\pi[-Re^{-R} - e^{-R} + 0 + 1] = 2\pi(1 - Re^{-R} - e^{-R}) \text{ ft}^3$$

(b) The average amount of water per hour per square foot supplied to the region within R feet of the sprinkler is

$\dfrac{V}{\text{area of region}} = \dfrac{V}{\pi R^2} = \dfrac{2\left(1 - Re^{-R} - e^{-R}\right)}{R^2}$ ft^3 (per hour per square foot). See the definition of the average value of a function following Example 15.1.8.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. As in Exercise 15.2.71, $f_{\text{avg}} = \frac{1}{A(D)} \iint_D f(x,y)\,dA$. Here $D = \{(r,\theta) \mid a \le r \le b, 0 \le \theta \le 2\pi\}$,

so $A(D) = \pi b^2 - \pi a^2 = \pi(b^2 - a^2)$ and

$$f_{\text{avg}} = \frac{1}{A(D)} \iint_D \frac{1}{\sqrt{x^2+y^2}}\,dA = \frac{1}{\pi(b^2-a^2)} \int_0^{2\pi} \int_a^b \frac{1}{\sqrt{r^2}}\, r\,dr\,d\theta = \frac{1}{\pi(b^2-a^2)} \int_0^{2\pi} d\theta \int_a^b dr$$

$$= \frac{1}{\pi(b^2-a^2)} \left[\theta\right]_0^{2\pi} \left[r\right]_a^b = \frac{1}{\pi(b^2-a^2)}(2\pi)(b-a) = \frac{2(b-a)}{(b+a)(b-a)} = \frac{2}{a+b}$$

48. The distance from a point (x,y) to the origin is $f(x,y) = \sqrt{x^2+y^2}$, so the average distance from points in D to the origin is

$$f_{\text{avg}} = \tfrac{1}{A(D)} \iint_D \sqrt{x^2+y^2}\,dA = \tfrac{1}{\pi a^2} \int_0^{2\pi} \int_0^a \sqrt{r^2}\; r\,dr\,d\theta$$

$$= \tfrac{1}{\pi a^2} \int_0^{2\pi} d\theta \int_0^a r^2\,dr = \tfrac{1}{\pi a^2} \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{3}r^3\right]_0^a = \tfrac{1}{\pi a^2} \cdot 2\pi \cdot \tfrac{1}{3}a^3 = \tfrac{2}{3}a$$

49. $$\int_{1/\sqrt{2}}^1 \int_{\sqrt{1-x^2}}^x xy\,dy\,dx + \int_1^{\sqrt{2}} \int_0^x xy\,dy\,dx + \int_{\sqrt{2}}^2 \int_0^{\sqrt{4-x^2}} xy\,dy\,dx$$

$$= \int_0^{\pi/4} \int_1^2 r^3 \cos\theta \sin\theta\,dr\,d\theta = \int_0^{\pi/4} \left[\frac{r^4}{4}\cos\theta\sin\theta\right]_{r=1}^{r=2} d\theta$$

$$= \frac{15}{4} \int_0^{\pi/4} \sin\theta\cos\theta\,d\theta = \frac{15}{4}\left[\frac{\sin^2\theta}{2}\right]_0^{\pi/4} = \frac{15}{16}$$

50. (a) $\iint_{D_a} e^{-(x^2+y^2)} dA = \int_0^{2\pi} \int_0^a re^{-r^2}\,dr\,d\theta = 2\pi\left[-\frac{1}{2}e^{-r^2}\right]_0^a = \pi\left(1 - e^{-a^2}\right)$ for each a. Then $\lim\limits_{a\to\infty} \pi\left(1-e^{-a^2}\right) = \pi$

since $e^{-a^2} \to 0$ as $a \to \infty$. Hence $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)}\,dA = \pi$.

(b) $\iint_{S_a} e^{-(x^2+y^2)}\,dA = \int_{-a}^a \int_{-a}^a e^{-x^2} e^{-y^2}\,dx\,dy = \left(\int_{-a}^a e^{-x^2}\,dx\right)\left(\int_{-a}^a e^{-y^2}\,dy\right)$ for each a.

From part (a), $\pi = \iint_{\mathbb{R}^2} e^{-(x^2+y^2)}\,dA$, so then

$\pi = \lim\limits_{a\to\infty} \iint_{S_a} e^{-(x^2+y^2)}\,dA = \lim\limits_{a\to\infty} \left(\int_{-a}^a e^{-x^2}\,dx\right)\left(\int_{-a}^a e^{-y^2}\,dy\right) = \left(\int_{-\infty}^{\infty} e^{-x^2}\,dx\right)\left(\int_{-\infty}^{\infty} e^{-y^2}\,dy\right)$, which is

what we wish to show.

To evaluate $\lim\limits_{a\to\infty} \left(\int_{-a}^a e^{-x^2}\,dx\right)\left(\int_{-a}^a e^{-y^2}\,dy\right)$, we are using the fact that these integrals are bounded. This is true

since on $[-1,1]$, $0 < e^{-x^2} \le 1$ while on $(-\infty,-1)$, $0 < e^{-x^2} \le e^x$ and on $(1,\infty)$, $0 < e^{-x^2} < e^{-x}$. Hence

$0 \le \int_{-\infty}^{\infty} e^{-x^2}\,dx \le \int_{-\infty}^{-1} e^x\,dx + \int_{-1}^1 dx + \int_1^{\infty} e^{-x}\,dx = 2(e^{-1}+1)$.

(c) Since $\left(\int_{-\infty}^{\infty} e^{-x^2}\,dx\right)\left(\int_{-\infty}^{\infty} e^{-y^2}\,dy\right) = \pi$ and y can be replaced by x, $\left(\int_{-\infty}^{\infty} e^{-x^2}\,dx\right)^2 = \pi$ implies that

$\int_{-\infty}^{\infty} e^{-x^2}\,dx = \pm\sqrt{\pi}$. But $e^{-x^2} \ge 0$ for all x, so $\int_{-\infty}^{\infty} e^{-x^2}\,dx = \sqrt{\pi}$.

(d) Letting $t = \sqrt{2}\,x$, $\int_{-\infty}^{\infty} e^{-x^2}\,dx = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2}}\left(e^{-t^2/2}\right) dt$, so that $\sqrt{\pi} = \frac{1}{\sqrt{2}} \int_{-\infty}^{\infty} e^{-t^2/2}\,dt$ or $\int_{-\infty}^{\infty} e^{-t^2/2}\,dt = \sqrt{2\pi}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

51. (a) We integrate by parts with $u = x$ and $dv = xe^{-x^2}\,dx$. Then $du = dx$ and $v = -\frac{1}{2}e^{-x^2}$, so

$$\begin{aligned}\int_0^\infty x^2e^{-x^2}\,dx &= \lim_{t\to\infty}\int_0^t x^2e^{-x^2}\,dx = \lim_{t\to\infty}\left(-\tfrac{1}{2}xe^{-x^2}\Big]_0^t + \int_0^t \tfrac{1}{2}e^{-x^2}\,dx\right)\\ &= \lim_{t\to\infty}\left(-\tfrac{1}{2}te^{-t^2}\right) + \tfrac{1}{2}\int_0^\infty e^{-x^2}\,dx = 0 + \tfrac{1}{2}\int_0^\infty e^{-x^2}\,dx \qquad \text{[by l'Hospital's Rule]}\\ &= \tfrac{1}{4}\int_{-\infty}^\infty e^{-x^2}\,dx \qquad \text{[since } e^{-x^2} \text{ is an even function]}\\ &= \tfrac{1}{4}\sqrt{\pi} \qquad \text{[by Exercise 50(c)]}\end{aligned}$$

(b) Let $u = \sqrt{x}$. Then $u^2 = x \;\Rightarrow\; dx = 2u\,du \;\Rightarrow$

$$\int_0^\infty \sqrt{x}\,e^{-x}\,dx = \lim_{t\to\infty}\int_0^t \sqrt{x}\,e^{-x}\,dx = \lim_{t\to\infty}\int_0^{\sqrt{t}} ue^{-u^2}2u\,du = 2\int_0^\infty u^2e^{-u^2}\,du = 2\left(\tfrac{1}{4}\sqrt{\pi}\right) \quad \text{[by part(a)]} = \tfrac{1}{2}\sqrt{\pi}.$$

15.4 Applications of Double Integrals

1. $Q = \iint_D \sigma(x,y)\,dA = \int_0^5\int_2^5 (2x+4y)\,dy\,dx = \int_0^5 \left[2xy+2y^2\right]_{y=2}^{y=5}\,dx$

$= \int_0^5 (10x+50-4x-8)\,dx = \int_0^5 (6x+42)\,dx = \left[3x^2+42x\right]_0^5 = 75+210 = 285$ C

2. $Q = \iint_D \sigma(x,y)\,dA = \iint_D \sqrt{x^2+y^2}\,dA = \int_0^{2\pi}\int_0^1 \sqrt{r^2}\;r\,dr\,d\theta$

$= \int_0^{2\pi} d\theta \;\int_0^1 r^2\,dr = [\theta]_0^{2\pi}\;\left[\tfrac{1}{3}r^3\right]_0^1 = 2\pi\cdot\tfrac{1}{3} = \tfrac{2\pi}{3}$ C

3. Since the density of the lamina is higher as $x \to 1$, we might estimate $\overline{x} = 0.7$. There is no vertical change in the density, so we would be confident that $\overline{y} = 0.5$.

$m = \iint_D \rho(x,y)\,dA = \int_0^1\int_0^1 x^2\,dy\,dx = \int_0^1 dy\;\int_0^1 x^2\,dx = \frac{1}{3}$

$M_y = \iint_D x\rho(x,y)\,dA = \int_0^1\int_0^1 x^3\,dy\,dx = \int_0^1 dy\;\int_0^1 x^3\,dx = \frac{1}{4}$

$M_x = \iint_D y\rho(x,y)\,dA = \int_0^1\int_0^1 x^2y\,dy\,dx = \int_0^1 y\,dy\;\int_0^1 x^2\,dx = \frac{1}{2}\cdot\frac{1}{3} = \frac{1}{6}$

Hence, $(\overline{x},\overline{y}) = (M_y/m, M_x/m) = \left(\frac{1/4}{1/3}, \frac{1/6}{1/3}\right) = \left(\frac{3}{4}, \frac{1}{2}\right)$.

4. Since the density of the lamina increases in a uniform fashion as $(x,y) \to (1,1)$, we might estimate $(x,y) = (0.7, 0.7)$.

$m = \iint_D \rho(x,y)\,dA = \int_0^1\int_0^1 xy\,dy\,dx = \int_0^1 y\,dy\;\int_0^1 x\,dx = \frac{1}{2}\cdot\frac{1}{2} = \frac{1}{4}$

$M_y = \iint_D x\rho(x,y)\,dA = \int_0^1\int_0^1 x^2y\,dy\,dx = \int_0^1 y\,dy\;\int_0^1 x^2\,dx = \frac{1}{2}\cdot\frac{1}{3} = \frac{1}{6}$

$M_x = \iint_D y\rho(x,y)\,dA = \int_0^1\int_0^1 xy^2\,dy\,dx = \int_0^1 y^2\,dy\;\int_0^1 x\,dx = \frac{1}{3}\cdot\frac{1}{2} = \frac{1}{6}$

Hence, $(\overline{x},\overline{y}) = (M_y/m, M_x/m) = \left(\frac{1/6}{1/4}, \frac{1/6}{1/4}\right) = \left(\frac{2}{3}, \frac{2}{3}\right)$.

5. $m = \iint_D \rho(x,y)\,dA = \int_1^3\int_1^4 ky^2\,dy\,dx = k\int_1^3 dx\;\int_1^4 y^2\,dy = k\,[x]_1^3\;\left[\tfrac{1}{3}y^3\right]_1^4 = k(2)(21) = 42k,$

$\overline{x} = \frac{1}{m}\iint_D x\rho(x,y)\,dA = \frac{1}{42k}\int_1^3\int_1^4 kxy^2\,dy\,dx = \frac{1}{42}\int_1^3 x\,dx\;\int_1^4 y^2\,dy = \frac{1}{42}\left[\tfrac{1}{2}x^2\right]_1^3\;\left[\tfrac{1}{3}y^3\right]_1^4 = \frac{1}{42}(4)(21) = 2,$

$\overline{y} = \frac{1}{m}\iint_D y\rho(x,y)\,dA = \frac{1}{42k}\int_1^3\int_1^4 ky^3\,dy\,dx = \frac{1}{42}\int_1^3 dx\;\int_1^4 y^3\,dy = \frac{1}{42}\,[x]_1^3\;\left[\tfrac{1}{4}y^4\right]_1^4 = \frac{1}{42}(2)\left(\frac{255}{4}\right) = \frac{85}{28}$

Hence, $(\overline{x},\overline{y}) = \left(2, \frac{85}{28}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $m = \iint_D \rho(x,y)\,dA = \int_0^a \int_0^b (1+x^2+y^2)\,dy\,dx = \int_0^a \left[y + x^2y + \frac{1}{3}y^3\right]_{y=0}^{y=b} dx = \int_0^a \left(b + bx^2 + \frac{1}{3}b^3\right) dx$

$= \left[bx + \frac{1}{3}bx^3 + \frac{1}{3}b^3x\right]_0^a = ab + \frac{1}{3}a^3b + \frac{1}{3}ab^3 = \frac{1}{3}ab(3+a^2+b^2),$

$M_y = \iint_D x\rho(x,y)\,dA = \int_0^a \int_0^b (x + x^3 + xy^2)\,dy\,dx = \int_0^a \left[xy + x^3y + \frac{1}{3}xy^3\right]_{y=0}^{y=b} dx$

$= \int_0^a \left(bx + bx^3 + \frac{1}{3}b^3x\right) dx = \left[\frac{1}{2}bx^2 + \frac{1}{4}bx^4 + \frac{1}{6}b^3x^2\right]_0^a = \frac{1}{2}a^2b + \frac{1}{4}a^4b + \frac{1}{6}a^2b^3$

$= \frac{1}{12}a^2b(6+3a^2+2b^2)$, and

$M_x = \iint_D y\rho(x,y)\,dA = \int_0^a \int_0^b (y + x^2y + y^3)\,dy\,dx = \int_0^a \left[\frac{1}{2}y^2 + \frac{1}{2}x^2y^2 + \frac{1}{4}y^4\right]_{y=0}^{y=b} dx$

$= \int_0^a \left(\frac{1}{2}b^2 + \frac{1}{2}b^2x^2 + \frac{1}{4}b^4\right) dx = \left[\frac{1}{2}b^2x + \frac{1}{6}b^2x^3 + \frac{1}{4}b^4x\right]_0^a = \frac{1}{2}ab^2 + \frac{1}{6}a^3b^2 + \frac{1}{4}ab^4$

$= \frac{1}{12}ab^2(6+2a^2+3b^2).$

Hence, $(\overline{x},\overline{y}) = \left(\dfrac{M_y}{m}, \dfrac{M_x}{m}\right) = \left(\dfrac{\frac{1}{12}a^2b(6+3a^2+2b^2)}{\frac{1}{3}ab(3+a^2+b^2)}, \dfrac{\frac{1}{12}ab^2(6+2a^2+3b^2)}{\frac{1}{3}ab(3+a^2+b^2)}\right)$

$= \left(\dfrac{a(6+3a^2+2b^2)}{4(3+a^2+b^2)}, \dfrac{b(6+2a^2+3b^2)}{4(3+a^2+b^2)}\right).$

7. $m = \int_0^2 \int_{x/2}^{3-x} (x+y)\,dy\,dx = \int_0^2 \left[xy + \frac{1}{2}y^2\right]_{y=x/2}^{y=3-x} dx = \int_0^2 \left[x(3-x) + \frac{1}{2}(3-x)^2 - \frac{1}{2}x^2 - \frac{1}{8}x^2\right] dx$

$= \int_0^2 \left(-\frac{9}{8}x^2 + \frac{9}{2}\right) dx = \left[-\frac{9}{8}\left(\frac{1}{3}x^3\right) + \frac{9}{2}x\right]_0^2 = 6,$

$M_y = \int_0^2 \int_{x/2}^{3-x} (x^2+xy)\,dy\,dx = \int_0^2 \left[x^2y + \frac{1}{2}xy^2\right]_{y=x/2}^{y=3-x} dx = \int_0^2 \left(\frac{9}{2}x - \frac{9}{8}x^3\right) dx = \frac{9}{2},$

$M_x = \int_0^2 \int_{x/2}^{3-y} (xy+y^2)\,dy\,dx = \int_0^2 \left[\frac{1}{2}xy^2 + \frac{1}{3}y^3\right]_{y=x/2}^{y=3-x} dx = \int_0^2 \left(9 - \frac{9}{2}x\right) dx = 9.$

Hence, $m = 6$, $(\overline{x},\overline{y}) = \left(\dfrac{M_y}{m}, \dfrac{M_x}{m}\right) = \left(\dfrac{3}{4}, \dfrac{3}{2}\right).$

8. Here $D = \left\{(x,y) \mid 0 \le y \le \frac{2}{5},\ \ y/2 \le x \le 1-2y\right\}.$

$m = \int_0^{2/5} \int_{y/2}^{1-2y} x\,dx\,dy = \int_0^{2/5} \left[\frac{1}{2}x^2\right]_{x=y/2}^{x=1-2y} dy = \frac{1}{2}\int_0^{2/5} \left[(1-2y)^2 - \left(\frac{1}{2}y\right)^2\right] dy$

$= \frac{1}{2}\int_0^{2/5} \left(\frac{15}{4}y^2 - 4y + 1\right) dy = \frac{1}{2}\left[\frac{5}{4}y^3 - 2y^2 + y\right]_0^{2/5} = \frac{1}{2}\left[\frac{2}{25} - \frac{8}{25} + \frac{2}{5}\right] = \frac{2}{25},$

$M_y = \int_0^{2/5} \int_{y/2}^{1-2y} x \cdot x\,dx\,dy = \int_0^{2/5} \left[\frac{1}{3}x^3\right]_{x=y/2}^{x=1-2y} dy = \frac{1}{3}\int_0^{2/5} \left[(1-2y)^3 - \left(\frac{1}{2}y\right)^3\right] dy$

$= \frac{1}{3}\int_0^{2/5} \left(-\frac{65}{8}y^3 + 12y^2 - 6y + 1\right) dy = \frac{1}{3}\left[-\frac{65}{32}y^4 + 4y^3 - 3y^2 + y\right]_0^{2/5} = \frac{1}{3}\left[-\frac{13}{250} + \frac{32}{125} - \frac{12}{25} + \frac{2}{5}\right] = \frac{31}{750},$

$M_x = \int_0^{2/5} \int_{y/2}^{1-2y} y \cdot x\,dx\,dy = \int_0^{2/5} y\left[\frac{1}{2}x^2\right]_{x=y/2}^{x=1-2y} dy = \frac{1}{2}\int_0^{2/5} y\left(\frac{15}{4}y^2 - 4y + 1\right) dy$

$= \frac{1}{2}\int_0^{2/5} \left(\frac{15}{4}y^3 - 4y^2 + y\right) dy = \frac{1}{2}\left[\frac{15}{16}y^4 - \frac{4}{3}y^3 + \frac{1}{2}y^2\right]_0^{2/5} = \frac{1}{2}\left[\frac{3}{125} - \frac{32}{375} + \frac{2}{25}\right] = \frac{7}{750}.$

Hence, $m = \frac{2}{25}$, $(\overline{x},\overline{y}) = \left(\frac{31/750}{2/25}, \frac{7/750}{2/25}\right) = \left(\frac{31}{60}, \frac{7}{60}\right).$

9. $m = \int_{-1}^1 \int_0^{1-x^2} ky\,dy\,dx = k\int_{-1}^1 \left[\frac{1}{2}y^2\right]_{y=0}^{y=1-x^2} dx = \frac{1}{2}k\int_{-1}^1 (1-x^2)^2\,dx = \frac{1}{2}k\int_{-1}^1 (1-2x^2+x^4)\,dx$

$= \frac{1}{2}k\left[x - \frac{2}{3}x^3 + \frac{1}{5}x^5\right]_{-1}^1 = \frac{1}{2}k\left(1 - \frac{2}{3} + \frac{1}{5} + 1 - \frac{2}{3} + \frac{1}{5}\right) = \frac{8}{15}k,$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$M_y = \int_{-1}^{1}\int_0^{1-x^2} kxy\,dy\,dx = k\int_{-1}^{1}\left[\tfrac{1}{2}xy^2\right]_{y=0}^{y=1-x^2} dx = \tfrac{1}{2}k\int_{-1}^{1} x\,(1-x^2)^2\,dx = \tfrac{1}{2}k\int_{-1}^{1}(x-2x^3+x^5)\,dx$$
$$= \tfrac{1}{2}k\left[\tfrac{1}{2}x^2 - \tfrac{1}{2}x^4 + \tfrac{1}{6}x^6\right]_{-1}^{1} = \tfrac{1}{2}k\left(\tfrac{1}{2}-\tfrac{1}{2}+\tfrac{1}{6}-\tfrac{1}{2}+\tfrac{1}{2}-\tfrac{1}{6}\right) = 0,$$

$$M_x = \int_{-1}^{1}\int_0^{1-x^2} ky^2\,dy\,dx = k\int_{-1}^{1}\left[\tfrac{1}{3}y^3\right]_{y=0}^{y=1-x^2} dx = \tfrac{1}{3}k\int_{-1}^{1}(1-x^2)^3\,dx = \tfrac{1}{3}k\int_{-1}^{1}(1-3x^2+3x^4-x^6)\,dx$$
$$= \tfrac{1}{3}k\left[x - x^3 + \tfrac{3}{5}x^5 - \tfrac{1}{7}x^7\right]_{-1}^{1} = \tfrac{1}{3}k\left(1-1+\tfrac{3}{5}-\tfrac{1}{7}+1-1+\tfrac{3}{5}-\tfrac{1}{7}\right) = \tfrac{32}{105}k.$$

Hence, $m = \frac{8}{15}k$, $(\overline{x},\overline{y}) = \left(0, \frac{32k/105}{8k/15}\right) = \left(0, \frac{4}{7}\right)$.

10. The boundary curves intersect when $x+2 = x^2 \quad\Leftrightarrow\quad x^2 - x - 2 = 0 \quad\Leftrightarrow\quad x = -1, x = 2$. Thus here $D = \{(x,y) \mid -1 \le x \le 2,\ x^2 \le y \le x+2\}$.

$$m = \int_{-1}^{2}\int_{x^2}^{x+2} kx^2\,dy\,dx = k\int_{-1}^{2} x^2\left[y\right]_{y=x^2}^{y=x+2} dx = k\int_{-1}^{2}(x^3+2x^2-x^4)\,dx$$
$$= k\left[\tfrac{1}{4}x^4 + \tfrac{2}{3}x^3 - \tfrac{1}{5}x^5\right]_{-1}^{2} = k\left(\tfrac{44}{15}+\tfrac{13}{60}\right) = \tfrac{63}{20}k,$$

$$M_y = \int_{-1}^{2}\int_{x^2}^{x+2} kx^3\,dy\,dx = k\int_{-1}^{2} x^3\left[y\right]_{y=x^2}^{y=x+2} dx = k\int_{-1}^{2}(x^4+2x^3-x^5)\,dx$$
$$= k\left[\tfrac{1}{5}x^5 + \tfrac{1}{2}x^4 - \tfrac{1}{6}x^6\right]_{-1}^{2} = k\left(\tfrac{56}{15}-\tfrac{2}{15}\right) = \tfrac{18}{5}k,$$

$$M_x = \int_{-1}^{2}\int_{x^2}^{x+2} kx^2y\,dy\,dx = k\int_{-1}^{2} x^2\left[\tfrac{1}{2}y^2\right]_{y=x^2}^{y=x+2} dx = \tfrac{1}{2}k\int_{-1}^{2} x^2\,(x^2+4x+4-x^4)\,dx$$
$$= \tfrac{1}{2}k\int_{-1}^{2}(x^4+4x^3+4x^2-x^6)\,dx = \tfrac{1}{2}k\left[\tfrac{1}{5}x^5 + x^4 + \tfrac{4}{3}x^3 - \tfrac{1}{7}x^7\right]_{-1}^{2} = \tfrac{1}{2}k\left(\tfrac{1552}{105}+\tfrac{41}{105}\right) = \tfrac{531}{70}k.$$

Hence, $m = \frac{63}{20}k$, $(\overline{x},\overline{y}) = \left(\frac{18k/5}{63k/20}, \frac{531k/70}{63k/20}\right) = \left(\frac{8}{7}, \frac{118}{49}\right)$.

11. $$m = \int_0^1\int_0^{e^{-x}} xy\,dy\,dx = \int_0^1 x\left[\tfrac{1}{2}y^2\right]_{y=0}^{y=e^{-x}} dx = \tfrac{1}{2}\int_0^1 x\left(e^{-x}\right)^2\,dx = \tfrac{1}{2}\int_0^1 xe^{-2x}\,dx \qquad \left[\begin{array}{c}\text{integrate by parts with}\\ u = x,\ dv = e^{-2x}\,dx\end{array}\right]$$
$$= \tfrac{1}{2}\left[-\tfrac{1}{4}(2x+1)e^{-2x}\right]_0^1 = -\tfrac{1}{8}\left(3e^{-2}-1\right) = \tfrac{1}{8} - \tfrac{3}{8}e^{-2},$$

$$M_y = \int_0^1\int_0^{e^{-x}} x^2y\,dy\,dx = \int_0^1 x^2\left[\tfrac{1}{2}y^2\right]_{y=0}^{y=e^{-x}} dx = \tfrac{1}{2}\int_0^1 x^2e^{-2x}\,dx \qquad [\text{integrate by parts twice}]$$
$$= \tfrac{1}{2}\left[-\tfrac{1}{4}\left(2x^2+2x+1\right)e^{-2x}\right]_0^1 = -\tfrac{1}{8}\left(5e^{-2}-1\right) = \tfrac{1}{8} - \tfrac{5}{8}e^{-2},$$

$$M_x = \int_0^1\int_0^{e^{-x}} xy^2\,dy\,dx = \int_0^1 x\left[\tfrac{1}{3}y^3\right]_{y=0}^{y=e^{-x}} dx = \tfrac{1}{3}\int_0^1 xe^{-3x}\,dx$$
$$= \tfrac{1}{3}\left[-\tfrac{1}{9}(3x+1)e^{-3x}\right]_0^1 = -\tfrac{1}{27}\left(4e^{-3}-1\right) = \tfrac{1}{27} - \tfrac{4}{27}e^{-3}.$$

Hence, $m = \frac{1}{8}\left(1-3e^{-2}\right)$, $(\overline{x},\overline{y}) = \left(\dfrac{\frac{1}{8}\left(1-5e^{-2}\right)}{\frac{1}{8}(1-3e^{-2})}, \dfrac{\frac{1}{27}\left(1-4e^{-3}\right)}{\frac{1}{8}(1-3e^{-2})}\right) = \left(\dfrac{e^2-5}{e^2-3}, \dfrac{8\left(e^3-4\right)}{27(e^3-3e)}\right)$.

12. Note that $\cos x \ge 0$ for $-\pi/2 \le x \le \pi/2$.

$$m = \int_{-\pi/2}^{\pi/2}\int_0^{\cos x} y\,dy\,dx = \int_{-\pi/2}^{\pi/2}\left[\tfrac{1}{2}y^2\right]_{y=0}^{y=\cos x} dx = \tfrac{1}{2}\int_{-\pi/2}^{\pi/2}\cos^2 x\,dx = \tfrac{1}{2}\left[\tfrac{1}{2}x + \tfrac{1}{4}\sin 2x\right]_{-\pi/2}^{\pi/2} = \tfrac{\pi}{4},$$

$$M_y = \int_{-\pi/2}^{\pi/2}\int_0^{\cos x} xy\,dy\,dx = \int_{-\pi/2}^{\pi/2} x\left[\tfrac{1}{2}y^2\right]_{y=0}^{y=\cos x} dx = \tfrac{1}{2}\int_{-\pi/2}^{\pi/2} x\cos^2 x\,dx \qquad \left[\begin{array}{c}\text{integrate by parts with}\\ u = x,\ dv = \cos^2 x\,dx\end{array}\right]$$
$$= \tfrac{1}{2}\left[x\left(\tfrac{1}{2}x + \tfrac{1}{4}\sin 2x\right)\Big|_{-\pi/2}^{\pi/2} - \int_{-\pi/2}^{\pi/2}\left(\tfrac{1}{2}x + \tfrac{1}{4}\sin 2x\right)dx\right]$$
$$= \tfrac{1}{2}\left(\tfrac{1}{8}\pi^2 - \tfrac{1}{8}\pi^2 - \left[\tfrac{1}{4}x^2 - \tfrac{1}{8}\cos 2x\right]_{-\pi/2}^{\pi/2}\right) = \tfrac{1}{2}\left(0 - \left[\tfrac{1}{16}\pi^2 + \tfrac{1}{8} - \tfrac{1}{16}\pi^2 - \tfrac{1}{8}\right]\right) = 0,$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$M_x = \int_{-\pi/2}^{\pi/2}\int_0^{\cos x} y^2\,dy\,dx = \int_{-\pi/2}^{\pi/2}\left[\tfrac{1}{3}y^3\right]_{y=0}^{y=\cos x} dx = \tfrac{1}{3}\int_{-\pi/2}^{\pi/2}\cos^3 x\,dx = \tfrac{1}{3}\int_{-\pi/2}^{\pi/2}(1-\sin^2 x)\cos x\,dx$$

$$[\text{substitute } u = \sin x \quad \Rightarrow \quad du = \cos x\,dx]$$

$$= \tfrac{1}{3}\left[\sin x - \tfrac{1}{3}\sin^3 x\right]_{-\pi/2}^{\pi/2} = \tfrac{1}{3}\left(1 - \tfrac{1}{3} + 1 - \tfrac{1}{3}\right) = \tfrac{4}{9}.$$

Hence, $m = \frac{\pi}{4}$, $(\overline{x}, \overline{y}) = \left(0, \frac{4/9}{\pi/4}\right) = \left(0, \frac{16}{9\pi}\right)$.

13. "The density at any point is proportional to its distance from the x-axis" $\Rightarrow$ $\rho(x,y) = ky$.

$$m = \iint_D ky\,dA = \int_0^{\pi/2}\int_0^1 k(r\sin\theta)\,r\,dr\,d\theta = k\int_0^{\pi/2}\sin\theta\,d\theta\,\int_0^1 r^2\,dr$$

$$= k\left[-\cos\theta\right]_0^{\pi/2}\left[\tfrac{1}{3}r^3\right]_0^1 = k(1)\left(\tfrac{1}{3}\right) = \tfrac{1}{3}k,$$

$$M_y = \iint_D x\cdot ky\,dA = \int_0^{\pi/2}\int_0^1 k(r\cos\theta)(r\sin\theta)\,r\,dr\,d\theta = k\int_0^{\pi/2}\sin\theta\cos\theta\,d\theta\,\int_0^1 r^3\,dr$$

$$= k\left[\tfrac{1}{2}\sin^2\theta\right]_0^{\pi/2}\left[\tfrac{1}{4}r^4\right]_0^1 = k\left(\tfrac{1}{2}\right)\left(\tfrac{1}{4}\right) = \tfrac{1}{8}k,$$

$$M_x = \iint_D y\cdot ky\,dA = \int_0^{\pi/2}\int_0^1 k(r\sin\theta)^2\,r\,dr\,d\theta = k\int_0^{\pi/2}\sin^2\theta\,d\theta\,\int_0^1 r^3\,dr$$

$$= k\left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{\pi/2}\left[\tfrac{1}{4}r^4\right]_0^1 = k\left(\tfrac{\pi}{4}\right)\left(\tfrac{1}{4}\right) = \tfrac{\pi}{16}k.$$

Hence, $(\overline{x}, \overline{y}) = \left(\frac{k/8}{k/3}, \frac{k\pi/16}{k/3}\right) = \left(\frac{3}{8}, \frac{3\pi}{16}\right)$.

14. "The density at any point is proportional to the square of its distance from the origin" $\Rightarrow$

$$\rho(x,y) = k\left(\sqrt{x^2+y^2}\right)^2 = k(x^2+y^2) = kr^2.$$

$$m = \int_0^{\pi/2}\int_0^1 kr^3\,dr\,d\theta = \tfrac{\pi}{8}k,$$

$$M_y = \int_0^{\pi/2}\int_0^1 kr^4\cos\theta\,dr\,d\theta = \tfrac{1}{5}k\int_0^{\pi/2}\cos\theta\,d\theta = \tfrac{1}{5}k\left[\sin\theta\right]_0^{\pi/2} = \tfrac{1}{5}k,$$

$$M_x = \int_0^{\pi/2}\int_0^1 kr^4\sin\theta\,dr\,d\theta = \tfrac{1}{5}k\int_0^{\pi/2}\sin\theta\,d\theta = \tfrac{1}{5}k\left[-\cos\theta\right]_0^{\pi/2} = \tfrac{1}{5}k.$$

Hence, $(\overline{x}, \overline{y}) = \left(\frac{8}{5\pi}, \frac{8}{5\pi}\right)$.

15.

$$\rho(x,y) = k\sqrt{x^2+y^2} = kr,$$

$$m = \iint_D \rho(x,y)dA = \int_0^{\pi}\int_1^2 kr\cdot r\,dr\,d\theta$$

$$= k\int_0^{\pi}d\theta\,\int_1^2 r^2\,dr = k(\pi)\left[\tfrac{1}{3}r^3\right]_1^2 = \tfrac{7}{3}\pi k,$$

$$M_y = \iint_D x\rho(x,y)dA = \int_0^{\pi}\int_1^2 (r\cos\theta)(kr)\,r\,dr\,d\theta = k\int_0^{\pi}\cos\theta\,d\theta\,\int_1^2 r^3\,dr$$

$$= k\left[\sin\theta\right]_0^{\pi}\left[\tfrac{1}{4}r^4\right]_1^2 = k(0)\left(\tfrac{15}{4}\right) = 0$$

[this is to be expected as the region and density function are symmetric about the y-axis]

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$M_x = \iint_D y\rho(x,y)dA = \int_0^\pi \int_1^2 (r\sin\theta)(kr)\, r\, dr\, d\theta = k\int_0^\pi \sin\theta\, d\theta \int_1^2 r^3\, dr$$
$$= k\left[-\cos\theta\right]_0^\pi \left[\tfrac{1}{4}r^4\right]_1^2 = k(1+1)\left(\tfrac{15}{4}\right) = \tfrac{15}{2}k.$$

Hence, $(\overline{x},\overline{y}) = \left(0, \frac{15k/2}{7\pi k/3}\right) = \left(0, \frac{45}{14\pi}\right)$.

16. Now $\rho(x,y) = k/\sqrt{x^2+y^2} = k/r$, so

$$m = \iint_D \rho(x,y)dA = \int_0^\pi \int_1^2 (k/r)\, r\, dr\, d\theta = k\int_0^\pi d\theta \int_1^2 dr = k(\pi)(1) = \pi k,$$

$$M_y = \iint_D x\rho(x,y)dA = \int_0^\pi \int_1^2 (r\cos\theta)(k/r)\, r\, dr\, d\theta = k\int_0^\pi \cos\theta\, d\theta \int_1^2 r\, dr$$
$$= k\left[\sin\theta\right]_0^\pi \left[\tfrac{1}{2}r^2\right]_1^2 = k(0)\left(\tfrac{3}{2}\right) = 0,$$

$$M_x = \iint_D y\rho(x,y)dA = \int_0^\pi \int_1^2 (r\sin\theta)(k/r)\, r\, dr\, d\theta = k\int_0^\pi \sin\theta\, d\theta \int_1^2 r\, dr$$
$$= k\left[-\cos\theta\right]_0^\pi \left[\tfrac{1}{2}r^2\right]_1^2 = k(1+1)\left(\tfrac{3}{2}\right) = 3k.$$

Hence, $(\overline{x},\overline{y}) = \left(0, \frac{3k}{\pi k}\right) = \left(0, \frac{3}{\pi}\right)$.

17. Placing the vertex opposite the hypotenuse at $(0,0)$ as in the figure,

$\rho(x,y) = k(x^2+y^2)$. Then

$$m = \int_0^a \int_0^{a-x} k\left(x^2+y^2\right) dy\, dx$$
$$= k\int_0^a \left[ax^2 - x^3 + \tfrac{1}{3}(a-x)^3\right] dx$$
$$= k\left[\tfrac{1}{3}ax^3 - \tfrac{1}{4}x^4 - \tfrac{1}{12}(a-x)^4\right]_0^a = \tfrac{1}{6}ka^4$$

By symmetry, $$M_y = M_x = \int_0^a \int_0^{a-x} ky(x^2+y^2)\, dy\, dx = k\int_0^a \left[\tfrac{1}{2}(a-x)^2x^2 + \tfrac{1}{4}(a-x)^4\right] dx$$
$$= k\left[\tfrac{1}{6}a^2x^3 - \tfrac{1}{4}ax^4 + \tfrac{1}{10}x^5 - \tfrac{1}{20}(a-x)^5\right]_0^a = \tfrac{1}{15}ka^5$$

Hence, $(\overline{x},\overline{y}) = \left(\frac{2}{5}a, \frac{2}{5}a\right)$.

18. $\rho(x,y) = k/\sqrt{x^2+y^2} = k/r$.

$$m = \int_{\pi/6}^{5\pi/6} \int_1^{2\sin\theta} \frac{k}{r}\, r\, dr\, d\theta = k\int_{\pi/6}^{5\pi/6} \left[(2\sin\theta) - 1\right] d\theta$$
$$= k\left[-2\cos\theta - \theta\right]_{\pi/6}^{5\pi/6} = 2k\left(\sqrt{3} - \tfrac{\pi}{3}\right)$$

By symmetry of D and $f(x) = x$, $M_y = 0$, and

$$M_x = \int_{\pi/6}^{5\pi/6} \int_1^{2\sin\theta} kr\sin\theta\, dr\, d\theta = \tfrac{1}{2}k\int_{\pi/6}^{5\pi/6} (4\sin^3\theta - \sin\theta)\, d\theta$$
$$= \tfrac{1}{2}k\left[-3\cos\theta + \tfrac{4}{3}\cos^3\theta\right]_{\pi/6}^{5\pi/6} = \sqrt{3}\,k$$

Hence, $(\overline{x},\overline{y}) = \left(0, \frac{3\sqrt{3}}{2\left(3\sqrt{3}-\pi\right)}\right)$.

19. $I_x = \iint_D y^2\rho(x,y)dA = \int_1^3 \int_1^4 y^2 \cdot ky^2\, dy\, dx = k\int_1^3 dx \int_1^4 y^4\, dy = k\,[x]_1^3 \left[\tfrac{1}{5}y^5\right]_1^4 = k(2)\left(\frac{1023}{5}\right) = 409.2k,$

$I_y = \iint_D x^2\rho(x,y)\, dA = \int_1^3 \int_1^4 x^2 \cdot ky^2\, dy\, dx = k\int_1^3 x^2\, dx \int_1^4 y^2\, dy = k\left[\tfrac{1}{3}x^3\right]_1^3 \left[\tfrac{1}{3}y^3\right]_1^4 = k\left(\frac{26}{3}\right)(21) = 182k,$

and $I_0 = I_x + I_y = 409.2k + 182k = 591.2k$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. $I_x = \iint_D y^2\rho(x,y)dA = \int_0^{2/5}\int_{y/2}^{1-2y} y^2\cdot x\,dx\,dy = \int_0^{2/5} y^2\left[\frac{1}{2}x^2\right]_{x=y/2}^{x=1-2y}dy = \frac{1}{2}\int_0^{2/5} y^2(\frac{15}{4}y^2-4y+1)\,dy$

$= \frac{1}{2}\int_0^{2/5}(\frac{15}{4}y^4-4y^3+y^2)\,dy = \frac{1}{2}\left[\frac{3}{4}y^5-y^4+\frac{1}{3}y^3\right]_0^{2/5} = \frac{16}{9375}$,

$I_y = \iint_D x^2\rho(x,y)\,dA = \int_0^{2/5}\int_{y/2}^{1-2y} x^2\cdot x\,dx\,dy = \int_0^{2/5}\left[\frac{1}{4}x^4\right]_{x=y/2}^{x=1-2y}dy = \frac{1}{4}\int_0^{2/5}\left[(1-2y)^4-\frac{1}{16}y^4\right]dy$

$= \frac{1}{4}\int_0^{2/5}(\frac{255}{16}y^4-32y^3+24y^2-8y+1)\,dy = \frac{1}{4}\left[\frac{51}{16}y^5-8y^4+8y^3-4y^2+y\right]_0^{2/5} = \frac{78}{3125}$,

and $I_0 = I_x + I_y = \frac{16}{9375}+\frac{78}{3125} = \frac{2}{75}$.

21. As in Exercise 17, we place the vertex opposite the hypotenuse at $(0,0)$ and the equal sides along the positive axes.

$I_x = \int_0^a\int_0^{a-x} y^2k(x^2+y^2)\,dy\,dx = k\int_0^a\int_0^{a-x}(x^2y^2+y^4)\,dy\,dx = k\int_0^a\left[\frac{1}{3}x^2y^3+\frac{1}{5}y^5\right]_{y=0}^{y=a-x}\,dx$

$= k\int_0^a\left[\frac{1}{3}x^2(a-x)^3+\frac{1}{5}(a-x)^5\right]dx = k\left[\frac{1}{3}\left(\frac{1}{3}a^3x^3-\frac{3}{4}a^2x^4+\frac{3}{5}ax^5-\frac{1}{6}x^6\right)-\frac{1}{30}(a-x)^6\right]_0^a = \frac{7}{180}ka^6$,

$I_y = \int_0^a\int_0^{a-x} x^2k(x^2+y^2)\,dy\,dx = k\int_0^a\int_0^{a-x}(x^4+x^2y^2)\,dy\,dx = k\int_0^a\left[x^4y+\frac{1}{3}x^2y^3\right]_{y=0}^{y=a-x}\,dx$

$= k\int_0^a\left[x^4(a-x)+\frac{1}{3}x^2(a-x)^3\right]dx = k\left[\frac{1}{5}ax^5-\frac{1}{6}x^6+\frac{1}{3}\left(\frac{1}{3}a^3x^3-\frac{3}{4}a^2x^4+\frac{3}{5}ax^5-\frac{1}{6}x^6\right)\right]_0^a = \frac{7}{180}ka^6$,

and $I_0 = I_x + I_y = \frac{7}{90}ka^6$.

22. If we find the moments of inertia about the x- and y-axes, we can determine in which direction rotation will be more difficult. (See the explanation following Example 4.) The moment of inertia about the x-axis is given by

$$I_x = \iint_D y^2\rho(x,y)\,dA = \int_0^2\int_0^2 y^2(1+0.1x)\,dy\,dx = \int_0^2(1+0.1x)\left[\frac{1}{3}y^3\right]_{y=0}^{y=2}dx$$
$$= \frac{8}{3}\int_0^2(1+0.1x)\,dx = \frac{8}{3}\left[x+0.1\cdot\frac{1}{2}x^2\right]_0^2 = \frac{8}{3}(2.2)\approx 5.87$$

Similarly, the moment of inertia about the y-axis is given by

$$I_y = \iint_D x^2\rho(x,y)\,dA = \int_0^2\int_0^2 x^2(1+0.1x)\,dy\,dx = \int_0^2 x^2(1+0.1x)\left[y\right]_{y=0}^{y=2}dx$$
$$= 2\int_0^2(x^2+0.1x^3)\,dx = 2\left[\frac{1}{3}x^3+0.1\cdot\frac{1}{4}x^4\right]_0^2 = 2\left(\frac{8}{3}+0.4\right)\approx 6.13$$

Since $I_y > I_x$, more force is required to rotate the fan blade about the y-axis.

23. $I_x = \iint_D y^2\rho(x,y)dA = \int_0^h\int_0^b \rho y^2\,dx\,dy = \rho\int_0^b dx\int_0^h y^2\,dy = \rho\left[x\right]_0^b\left[\frac{1}{3}y^3\right]_0^h = \rho b\left(\frac{1}{3}h^3\right) = \frac{1}{3}\rho bh^3$,

$I_y = \iint_D x^2\rho(x,y)dA = \int_0^h\int_0^b \rho x^2\,dx\,dy = \rho\int_0^b x^2\,dx\int_0^h dy = \rho\left[\frac{1}{3}x^3\right]_0^b[y]_0^h = \frac{1}{3}\rho b^3h$,

and $m = \rho\,(\text{area of rectangle}) = \rho bh$ since the lamina is homogeneous. Hence $\overline{\overline{x}}^2 = \dfrac{I_y}{m} = \dfrac{\frac{1}{3}\rho b^3h}{\rho bh} = \dfrac{b^2}{3} \quad\Rightarrow\quad \overline{\overline{x}} = \dfrac{b}{\sqrt{3}}$

and $\overline{\overline{y}}^2 = \dfrac{I_x}{m} = \dfrac{\frac{1}{3}\rho bh^3}{\rho bh} = \dfrac{h^2}{3} \quad\Rightarrow\quad \overline{\overline{y}} = \dfrac{h}{\sqrt{3}}$.

24. Here we assume $b>0$, $h>0$ but note that we arrive at the same results if $b<0$ or $h<0$. We have $D = \left\{(x,y)\mid 0\le x\le b, 0\le y\le h-\frac{h}{b}x\right\}$, so

$I_x = \int_0^b\int_0^{h-hx/b} y^2\rho\,dy\,dx = \rho\int_0^b\left[\frac{1}{3}y^3\right]_{y=0}^{y=h-hx/b}dx = \frac{1}{3}\rho\int_0^b\left(h-\frac{h}{b}x\right)^3dx$

$= \frac{1}{3}\rho\left[-\frac{b}{h}\left(\frac{1}{4}\right)\left(h-\frac{h}{b}x\right)^4\right]_0^b = -\frac{b}{12h}\rho(0-h^4) = \frac{1}{12}\rho bh^3$,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$I_y = \int_0^b \int_0^{h-hx/b} x^2 \rho \, dy \, dx = \rho \int_0^b x^2 \left(h - \frac{h}{b}x\right) dx = \rho \int_0^b \left(hx^2 - \frac{h}{b}x^3\right) dx$

$= \rho \left[\frac{h}{3}x^3 - \frac{h}{4b}x^4\right]_0^b = \rho\left(\frac{hb^3}{3} - \frac{hb^3}{4}\right) = \frac{1}{12}\rho b^3 h,$

and $m = \int_0^b \int_0^{h-hx/b} \rho \, dy \, dx = \rho \int_0^b \left(h - \frac{h}{b}x\right) dx = \rho \left[hx - \frac{h}{2b}x^2\right]_0^b = \frac{1}{2}\rho bh$. Hence $\overline{\overline{x}}^2 = \dfrac{I_y}{m} = \dfrac{\frac{1}{12}\rho b^3 h}{\frac{1}{2}\rho bh} = \dfrac{b^2}{6} \quad \Rightarrow$

$\overline{\overline{x}} = \dfrac{b}{\sqrt{6}}$ and $\overline{\overline{y}}^2 = \dfrac{I_x}{m} = \dfrac{\frac{1}{12}\rho bh^3}{\frac{1}{2}\rho bh} = \dfrac{h^2}{6} \quad \Rightarrow \quad \overline{\overline{y}} = \dfrac{h}{\sqrt{6}}.$

25. In polar coordinates, the region is $D = \left\{(r,\theta) \mid 0 \le r \le a, 0 \le \theta \le \frac{\pi}{2}\right\}$, so

$I_x = \iint_D y^2 \rho \, dA = \int_0^{\pi/2} \int_0^a \rho (r\sin\theta)^2 \, r \, dr \, d\theta = \rho \int_0^{\pi/2} \sin^2 d\theta \int_0^a r^3 \, dr$

$= \rho\left[\frac{1}{2}\theta - \frac{1}{4}\sin 2\theta\right]_0^{\pi/2} \left[\frac{1}{4}r^4\right]_0^a = \rho\left(\frac{\pi}{4}\right)\left(\frac{1}{4}a^4\right) = \frac{1}{16}\rho a^4 \pi,$

$I_y = \iint_D x^2 \rho \, dA = \int_0^{\pi/2} \int_0^a \rho (r\cos\theta)^2 \, r \, dr \, d\theta = \rho \int_0^{\pi/2} \cos^2 d\theta \int_0^a r^3 \, dr$

$= \rho\left[\frac{1}{2}\theta + \frac{1}{4}\sin 2\theta\right]_0^{\pi/2} \left[\frac{1}{4}r^4\right]_0^a = \rho\left(\frac{\pi}{4}\right)\left(\frac{1}{4}a^4\right) = \frac{1}{16}\rho a^4 \pi,$

and $m = \rho \cdot A(D) = \rho \cdot \frac{1}{4}\pi a^2$ since the lamina is homogeneous. Hence $\overline{\overline{x}}^2 = \overline{\overline{y}}^2 = \dfrac{\frac{1}{16}\rho a^4 \pi}{\frac{1}{4}\rho a^2 \pi} = \dfrac{a^2}{4} \quad \Rightarrow \quad \overline{\overline{x}} = \overline{\overline{y}} = \dfrac{a}{2}.$

26. $m = \int_0^\pi \int_0^{\sin x} \rho \, dy \, dx = \rho \int_0^\pi \sin x \, dx = \rho\left[-\cos x\right]_0^\pi = 2\rho,$

$I_x = \int_0^\pi \int_0^{\sin x} \rho y^2 \, dy \, dx = \frac{1}{3}\rho \int_0^\pi \sin^3 x \, dx = \frac{1}{3}\rho \int_0^\pi (1 - \cos^2 x)\sin x \, dx = \frac{1}{3}\rho\left[-\cos x + \frac{1}{3}\cos^3 x\right]_0^\pi = \frac{4}{9}\rho,$

$I_y = \int_0^\pi \int_0^{\sin x} \rho x^2 \, dy \, dx = \rho \int_0^\pi x^2 \sin x \, dx = \rho\left[-x^2\cos x + 2x\sin x + 2\cos x\right]_0^\pi$ [by integrating by parts twice]

$= \rho(\pi^2 - 4).$

Then $\overline{\overline{y}}^2 = \dfrac{I_x}{m} = \dfrac{2}{9}$, so $\overline{\overline{y}} = \dfrac{\sqrt{2}}{3}$ and $\overline{\overline{x}}^2 = \dfrac{I_y}{m} = \dfrac{\pi^2 - 4}{2}$, so $\overline{\overline{x}} = \sqrt{\dfrac{\pi^2 - 4}{2}}.$

27. The right loop of the curve is given by $D = \{(r,\theta) \mid 0 \le r \le \cos 2\theta, \ -\pi/4 \le \theta \le \pi/4\}$. Using a CAS, we find $m = \iint_D \rho(x,y) \, dA = \iint_D (x^2 + y^2) \, dA = \int_{-\pi/4}^{\pi/4} \int_0^{\cos 2\theta} r^2 \, r \, dr \, d\theta = \dfrac{3\pi}{64}$. Then

$$\overline{x} = \frac{1}{m}\iint_D x\rho(x,y) \, dA = \frac{64}{3\pi}\int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} (r\cos\theta)\, r^2 \, r \, dr \, d\theta = \frac{64}{3\pi}\int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} r^4 \cos\theta \, dr \, d\theta = \frac{16384\sqrt{2}}{10395\pi} \text{ and}$$

$$\overline{y} = \frac{1}{m}\iint_D y\rho(x,y) \, dA = \frac{64}{3\pi}\int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} (r\sin\theta)\, r^2 \, r \, dr \, d\theta = \frac{64}{3\pi}\int_{-\pi/4}^{\pi/4}\int_0^{\cos 2\theta} r^4 \sin\theta \, dr \, d\theta = 0, \text{ so}$$

$(\overline{x}, \overline{y}) = \left(\dfrac{16384\sqrt{2}}{10395\pi}, 0\right).$

The moments of inertia are

$I_x = \iint_D y^2 \rho(x,y) \, dA = \int_{-\pi/4}^{\pi/4} \int_0^{\cos 2\theta} (r\sin\theta)^2 \, r^2 \, r \, dr \, d\theta = \int_{-\pi/4}^{\pi/4} \int_0^{\cos 2\theta} r^5 \sin^2\theta \, dr \, d\theta = \dfrac{5\pi}{384} - \dfrac{4}{105},$

$I_y = \iint_D x^2 \rho(x,y) \, dA = \int_{-\pi/4}^{\pi/4} \int_0^{\cos 2\theta} (r\cos\theta)^2 \, r^2 \, r \, dr \, d\theta = \int_{-\pi/4}^{\pi/4} \int_0^{\cos 2\theta} r^5 \cos^2\theta \, dr \, d\theta = \dfrac{5\pi}{384} + \dfrac{4}{105}$, and

$I_0 = I_x + I_y = \dfrac{5\pi}{192}.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. Using a CAS, we find $m = \iint_D \rho(x,y)\,dA = \int_0^2 \int_0^{xe^{-x}} x^2y^2\,dy\,dx = \frac{8}{729}(5 - 899e^{-6})$. Then

$$\overline{x} = \frac{1}{m}\iint_D x\rho(x,y)\,dA = \frac{729}{8(5-899e^{-6})}\int_0^2\int_0^{xe^{-x}} x^3y^2\,dy\,dx = \frac{2(5e^6-1223)}{5e^6-899} \text{ and}$$

$$\overline{y} = \frac{1}{m}\iint_D y\rho(x,y)\,dA = \frac{729}{8(5-899e^{-6})}\int_0^2\int_0^{xe^{-x}} x^2y^3\,dy\,dx = \frac{729(45e^6-42037e^{-2})}{32768(5e^6-899)}, \text{ so}$$

$$(\overline{x},\overline{y}) = \left(\frac{2(5e^6-1223)}{5e^6-899}, \frac{729(45e^6-42037e^{-2})}{32768(5e^6-899)}\right).$$

The moments of inertia are $I_x = \iint_D y^2\rho(x,y)\,dA = \int_0^2\int_0^{xe^{-x}} x^2y^4\,dy\,dx = \frac{16}{390625}(63 - 305593e^{-10})$,

$I_y = \iint_D x^2\rho(x,y)\,dA = \int_0^2\int_0^{xe^{-x}} x^4y^2\,dy\,dx = \frac{80}{2187}(7 - 2101e^{-6})$, and

$I_0 = I_x + I_y = \frac{16}{854296875}(13809656 - 4103515625e^{-6} - 668331891e^{-10})$.

29. (a) $f(x,y)$ is a joint density function, so we know $\iint_{\mathbb{R}^2} f(x,y)\,dA = 1$. Since $f(x,y) = 0$ outside the rectangle $[0,1]\times[0,2]$, we can say

$$\iint_{\mathbb{R}^2} f(x,y)\,dA = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f(x,y)\,dy\,dx = \int_0^1\int_0^2 Cx(1+y)\,dy\,dx$$
$$= C\int_0^1 x\left[y+\tfrac{1}{2}y^2\right]_{y=0}^{y=2}\,dx = C\int_0^1 4x\,dx = C\left[2x^2\right]_0^1 = 2C$$

Then $2C = 1 \quad\Rightarrow\quad C = \frac{1}{2}$.

(b) $P(X \le 1, Y \le 1) = \int_{-\infty}^{1}\int_{-\infty}^{1} f(x,y)\,dy\,dx = \int_0^1\int_0^1 \frac{1}{2}x(1+y)\,dy\,dx$

$$= \int_0^1 \tfrac{1}{2}x\left[y+\tfrac{1}{2}y^2\right]_{y=0}^{y=1}\,dx = \int_0^1 \tfrac{1}{2}x\left(\tfrac{3}{2}\right)dx = \tfrac{3}{4}\left[\tfrac{1}{2}x^2\right]_0^1 = \tfrac{3}{8} \text{ or } 0.375$$

(c) $P(X+Y \le 1) = P((X,Y)\in D)$ where D is the triangular region shown in the figure. Thus

$$P(X+Y\le 1) = \iint_D f(x,y)\,dA = \int_0^1\int_0^{1-x} \tfrac{1}{2}x(1+y)\,dy\,dx$$
$$= \int_0^1 \tfrac{1}{2}x\left[y+\tfrac{1}{2}y^2\right]_{y=0}^{y=1-x}\,dx = \int_0^1 \tfrac{1}{2}x\left(\tfrac{1}{2}x^2 - 2x + \tfrac{3}{2}\right)dx$$
$$= \tfrac{1}{4}\int_0^1 \left(x^3 - 4x^2 + 3x\right)dx = \tfrac{1}{4}\left[\tfrac{x^4}{4} - 4\tfrac{x^3}{3} + 3\tfrac{x^2}{2}\right]_0^1$$
$$= \tfrac{5}{48} \approx 0.1042$$

30. (a) $f(x,y) \ge 0$, so f is a joint density function if $\iint_{\mathbb{R}^2} f(x,y)\,dA = 1$. Here, $f(x,y) = 0$ outside the square $[0,1]\times[0,1]$, so $\iint_{\mathbb{R}^2} f(x,y)\,dA = \int_0^1\int_0^1 4xy\,dy\,dx = \int_0^1 \left[2xy^2\right]_{y=0}^{y=1}\,dx = \int_0^1 2x\,dx = x^2\Big]_0^1 = 1$.

Thus, $f(x,y)$ is a joint density function.

(b) (i) No restriction is placed on Y, so

$$P\left(X \ge \tfrac{1}{2}\right) = \int_{1/2}^{\infty}\int_{-\infty}^{\infty} f(x,y)\,dy\,dx = \int_{1/2}^1\int_0^1 4xy\,dy\,dx = \int_{1/2}^1 \left[2xy^2\right]_{y=0}^{y=1}\,dx = \int_{1/2}^1 2x\,dx = x^2\Big]_{1/2}^1 = \tfrac{3}{4}.$$

(ii) $P\left(X \ge \frac{1}{2}, Y \le \frac{1}{2}\right) = \int_{1/2}^{\infty}\int_{-\infty}^{1/2} f(x,y)\,dy\,dx = \int_{1/2}^1\int_0^{1/2} 4xy\,dy\,dx$

$$= \int_{1/2}^1 \left[2xy^2\right]_{y=0}^{y=1/2}\,dx = \int_{1/2}^1 \tfrac{1}{2}x\,dx = \tfrac{1}{2}\cdot\tfrac{1}{2}x^2\Big]_{1/2}^1 = \tfrac{3}{16}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) The expected value of X is given by

$$\mu_1 = \iint_{\mathbb{R}^2} x\, f(x,y)\, dA = \int_0^1 \int_0^1 x(4xy)\, dy\, dx = \int_0^1 2x^2 \left[y^2\right]_{y=0}^{y=1} dx = 2\int_0^1 x^2\, dx = 2\left[\tfrac{1}{3}x^3\right]_0^1 = \tfrac{2}{3}$$

The expected value of Y is

$$\mu_2 = \iint_{\mathbb{R}^2} y\, f(x,y)\, dA = \int_0^1 \int_0^1 y(4xy)\, dy\, dx = \int_0^1 4x\left[\tfrac{1}{3}y^3\right]_{y=0}^{y=1} dx = \tfrac{4}{3}\int_0^1 x\, dx = \tfrac{4}{3}\left[\tfrac{1}{2}x^2\right]_0^1 = \tfrac{2}{3}$$

31. (a) $f(x,y) \geq 0$, so f is a joint density function if $\iint_{\mathbb{R}^2} f(x,y)\, dA = 1$. Here, $f(x,y) = 0$ outside the first quadrant, so

$$\begin{aligned}
\iint_{\mathbb{R}^2} f(x,y)\, dA &= \int_0^\infty \int_0^\infty 0.1e^{-(0.5x+0.2y)}\, dy\, dx = 0.1\int_0^\infty \int_0^\infty e^{-0.5x}e^{-0.2y}\, dy\, dx \\
&= 0.1\int_0^\infty e^{-0.5x}\, dx \int_0^\infty e^{-0.2y}\, dy = 0.1 \lim_{t\to\infty} \int_0^t e^{-0.5x}\, dx \lim_{t\to\infty} \int_0^t e^{-0.2y}\, dy \\
&= 0.1 \lim_{t\to\infty} \left[-2e^{-0.5x}\right]_0^t \lim_{t\to\infty} \left[-5e^{-0.2y}\right]_0^t = 0.1 \lim_{t\to\infty} \left[-2(e^{-0.5t}-1)\right] \lim_{t\to\infty} \left[-5(e^{-0.2t}-1)\right] \\
&= (0.1)\cdot(-2)(0-1)\cdot(-5)(0-1) = 1
\end{aligned}$$

Thus $f(x,y)$ is a joint density function.

(b) (i) No restriction is placed on X, so

$$\begin{aligned}
P(Y \geq 1) &= \int_{-\infty}^\infty \int_1^\infty f(x,y)\, dy\, dx = \int_0^\infty \int_1^\infty 0.1e^{-(0.5x+0.2y)}\, dy\, dx \\
&= 0.1\int_0^\infty e^{-0.5x}\, dx \int_1^\infty e^{-0.2y}\, dy = 0.1 \lim_{t\to\infty} \int_0^t e^{-0.5x}\, dx \lim_{t\to\infty} \int_1^t e^{-0.2y}\, dy \\
&= 0.1 \lim_{t\to\infty} \left[-2e^{-0.5x}\right]_0^t \lim_{t\to\infty} \left[-5e^{-0.2y}\right]_1^t = 0.1 \lim_{t\to\infty} \left[-2(e^{-0.5t}-1)\right] \lim_{t\to\infty} \left[-5(e^{-0.2t}-e^{-0.2})\right] \\
&(0.1)\cdot(-2)(0-1)\cdot(-5)(0-e^{-0.2}) = e^{-0.2} \approx 0.8187
\end{aligned}$$

(ii) $$\begin{aligned}
P(X \leq 2, Y \leq 4) &= \int_{-\infty}^2 \int_{-\infty}^4 f(x,y)\, dy\, dx = \int_0^2 \int_0^4 0.1e^{-(0.5x+0.2y)}\, dy\, dx \\
&= 0.1\int_0^2 e^{-0.5x}\, dx \int_0^4 e^{-0.2y}\, dy = 0.1\left[-2e^{-0.5x}\right]_0^2 \left[-5e^{-0.2y}\right]_0^4 \\
&= (0.1)\cdot(-2)(e^{-1}-1)\cdot(-5)(e^{-0.8}-1) \\
&= (e^{-1}-1)(e^{-0.8}-1) = 1 + e^{-1.8} - e^{-0.8} - e^{-1} \approx 0.3481
\end{aligned}$$

(c) The expected value of X is given by

$$\begin{aligned}
\mu_1 &= \iint_{\mathbb{R}^2} x\, f(x,y)\, dA = \int_0^\infty \int_0^\infty x\left[0.1e^{-(0.5x+0.2y)}\right] dy\, dx \\
&= 0.1\int_0^\infty xe^{-0.5x}\, dx \int_0^\infty e^{-0.2y}\, dy = 0.1 \lim_{t\to\infty} \int_0^t xe^{-0.5x}\, dx \lim_{t\to\infty} \int_0^t e^{-0.2y}\, dy
\end{aligned}$$

To evaluate the first integral, we integrate by parts with $u = x$ and $dv = e^{-0.5x}\, dx$ (or we can use Formula 96 in the Table of Integrals): $\int xe^{-0.5x}\, dx = -2xe^{-0.5x} - \int -2e^{-0.5x}\, dx = -2xe^{-0.5x} - 4e^{-0.5x} = -2(x+2)e^{-0.5x}$.

Thus

$$\begin{aligned}
\mu_1 &= 0.1 \lim_{t\to\infty} \left[-2(x+2)e^{-0.5x}\right]_0^t \lim_{t\to\infty} \left[-5e^{-0.2y}\right]_0^t \\
&= 0.1 \lim_{t\to\infty} (-2)\left[(t+2)e^{-0.5t} - 2\right] \lim_{t\to\infty} (-5)\left[e^{-0.2t} - 1\right] \\
&= 0.1(-2)\left(\lim_{t\to\infty} \frac{t+2}{e^{0.5t}} - 2\right)(-5)(-1) = 2 \qquad \text{[by l'Hospital's Rule]}
\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The expected value of Y is given by

$$\mu_2 = \iint_{\mathbb{R}^2} y\, f(x,y)\, dA = \int_0^\infty \int_0^\infty y\left[0.1e^{-(0.5+0.2y)}\right] dy\, dx$$
$$= 0.1 \int_0^\infty e^{-0.5x}\, dx \int_0^\infty ye^{-0.2y}\, dy = 0.1 \lim_{t\to\infty} \int_0^t e^{-0.5x}\, dx \lim_{t\to\infty} \int_0^t ye^{-0.2y}\, dy$$

To evaluate the second integral, we integrate by parts with $u = y$ and $dv = e^{-0.2y}\, dy$ (or again we can use Formula 96 in the Table of Integrals) which gives $\int ye^{-0.2y}\, dy = -5ye^{-0.2y} + \int 5e^{-0.2y}\, dy = -5(y+5)e^{-0.2y}$. Then

$$\mu_2 = 0.1 \lim_{t\to\infty} \left[-2e^{-0.5x}\right]_0^t \lim_{t\to\infty} \left[-5(y+5)e^{-0.2y}\right]_0^t$$
$$= 0.1 \lim_{t\to\infty} \left[-2(e^{-0.5t}-1)\right] \lim_{t\to\infty} \left(-5\left[(t+5)e^{-0.2t}-5\right]\right)$$
$$= 0.1(-2)(-1)\cdot(-5)\left(\lim_{t\to\infty} \frac{t+5}{e^{0.2t}} - 5\right) = 5 \qquad \text{[by l'Hospital's Rule]}$$

32. (a) The lifetime of each bulb has exponential density function

$$f(t) = \begin{cases} 0 & \text{if } t < 0 \\ \frac{1}{1000}e^{-t/1000} & \text{if } t \geq 0 \end{cases}$$

If X and Y are the lifetimes of the individual bulbs, then X and Y are independent, so the joint density function is the product of the individual density functions:

$$f(x,y) = \begin{cases} 10^{-6}e^{-(x+y)/1000} & \text{if } x \geq 0, y \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

The probability that both of the bulbs fail within 1000 hours is

$$P(X \leq 1000, Y \leq 1000) = \int_{-\infty}^{1000} \int_{-\infty}^{1000} f(x,y)\, dy\, dx = \int_0^{1000} \int_0^{1000} 10^{-6}e^{-(x+y)/1000}\, dy\, dx$$
$$= 10^{-6} \int_0^{1000} e^{-x/1000}\, dx \int_0^{1000} e^{-y/1000}\, dy$$
$$= 10^{-6} \left[-1000e^{-x/1000}\right]_0^{1000} \left[-1000e^{-y/1000}\right]_0^{1000}$$
$$= \left(e^{-1}-1\right)^2 \approx 0.3996$$

(b) Now we are asked for the probability that the combined lifetimes of both bulbs is 1000 hours or less. Thus we want to find $P(X+Y \leq 1000)$, or equivalently $P((X,Y) \in D)$ where D is the triangular region shown in the figure. Then

$$P(X+Y \leq 1000) = \iint_D f(x,y)\ dA$$
$$= \int_0^{1000} \int_0^{1000-x} 10^{-6}e^{-(x+y)/1000}\, dy\, dx$$
$$= 10^{-6} \int_0^{1000} \left[-1000e^{-(x+y)/1000}\right]_{y=0}^{y=1000-x} dx = -10^{-3} \int_0^{1000} \left(e^{-1} - e^{-x/1000}\right) dx$$
$$= -10^{-3} \left[e^{-1}x + 1000e^{-x/1000}\right]_0^{1000} = 1 - 2e^{-1} \approx 0.2642$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. (a) The random variables X and Y are normally distributed with $\mu_1 = 45$, $\mu_2 = 20$, $\sigma_1 = 0.5$, and $\sigma_2 = 0.1$.

The individual density functions for X and Y, then, are $f_1(x) = \dfrac{1}{0.5\sqrt{2\pi}}\, e^{-(x-45)^2/0.5}$ and

$f_2(y) = \dfrac{1}{0.1\sqrt{2\pi}}\, e^{-(y-20)^2/0.02}$. Since X and Y are independent, the joint density function is the product

$f(x,y) = f_1(x)f_2(y) = \dfrac{1}{0.5\sqrt{2\pi}}\, e^{-(x-45)^2/0.5}\dfrac{1}{0.1\sqrt{2\pi}}\, e^{-(y-20)^2/0.02} = \frac{10}{\pi}e^{-2(x-45)^2-50(y-20)^2}$.

Then $P(40 \le X \le 50, 20 \le Y \le 25) = \int_{40}^{50}\int_{20}^{25} f(x,y)\,dy\,dx = \frac{10}{\pi}\int_{40}^{50}\int_{20}^{25} e^{-2(x-45)^2-50(y-20)^2}\,dy\,dx$.

Using a CAS or calculator to evaluate the integral, we get $P(40 \le X \le 50, 20 \le Y \le 25) \approx 0.500$.

(b) $P(4(X-45)^2 + 100(Y-20)^2 \le 2) = \iint_D \frac{10}{\pi}e^{-2(x-45)^2-50(y-20)^2}\,dA$, where D is the region enclosed by the ellipse $4(x-45)^2 + 100(y-20)^2 = 2$. Solving for y gives $y = 20 \pm \frac{1}{10}\sqrt{2-4(x-45)^2}$, the upper and lower halves of the ellipse, and these two halves meet where $y = 20$ [since the ellipse is centered at $(45, 20)$] $\Rightarrow$ $4(x-45)^2 = 2$ $\Rightarrow$ $x = 45 \pm \frac{1}{\sqrt{2}}$. Thus

$$\iint_D \frac{10}{\pi}e^{-2(x-45)^2-50(y-20)^2}\,dA = \frac{10}{\pi}\int_{45-1/\sqrt{2}}^{45+1/\sqrt{2}}\int_{20-\frac{1}{10}\sqrt{2-4(x-45)^2}}^{20+\frac{1}{10}\sqrt{2-4(x-45)^2}} e^{-2(x-45)^2-50(y-20)^2}\,dy\,dx.$$

Using a CAS or calculator to evaluate the integral, we get $P(4(X-45)^2 + 100(Y-20)^2 \le 2) \approx 0.632$.

34. Because X and Y are independent, the joint density function for Xavier's and Yolanda's arrival times is the product of the individual density functions:

$$f(x,y) = f_1(x)f_2(y) = \begin{cases} \frac{1}{50}e^{-x}y & \text{if } x \ge 0, 0 \le y \le 10 \\ 0 & \text{otherwise} \end{cases}$$

Since Xavier won't wait for Yolanda, they won't meet unless $X \ge Y$. Additionally, Yolanda will wait up to half an hour but no longer, so they won't meet unless $X - Y \le 30$. Thus the probability that they meet is $P((X,Y) \in D)$ where D is the parallelogram shown in the figure. The integral is simpler to evaluate if we consider D as a type II region, so

$$\begin{aligned} P((X,Y) \in D) &= \iint_D f(x,y)\,dx\,dy = \int_0^{10}\int_y^{y+30} \tfrac{1}{50}e^{-x}y\,dx\,dy \\ &= \tfrac{1}{50}\int_0^{10} y\left[-e^{-x}\right]_{x=y}^{x=y+30}\,dy = \tfrac{1}{50}\int_0^{10} y(-e^{-(y+30)} + e^{-y})\,dy \\ &= \tfrac{1}{50}(1-e^{-30})\int_0^{10} ye^{-y}\,dy \end{aligned}$$

By integration by parts (or Formula 96 in the Table of Integrals), this is

$\frac{1}{50}(1-e^{-30})\left[-(y+1)e^{-y}\right]_0^{10} = \frac{1}{50}(1-e^{-30})(1-11e^{-10}) \approx 0.020$. Thus there is only about a 2% chance they will meet. Such is student life!

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. (a) If $f(P, A)$ is the probability that an individual at A will be infected by an individual at P, and $k\,dA$ is the number of infected individuals in an element of area dA, then $f(P, A)k\,dA$ is the number of infections that should result from exposure of the individual at A to infected people in the element of area dA. Integration over D gives the number of infections of the person at A due to all the infected people in D. In rectangular coordinates (with the origin at the city's center), the exposure of a person at A is

$$E = \iint_D kf(P, A)\,dA = k\iint_D \tfrac{1}{20}\,[20 - d(P, A)]\;dA = k\iint_D \left[1 - \tfrac{1}{20}\sqrt{(x - x_0)^2 + (y - y_0)^2}\right] dA$$

(b) If $A = (0, 0)$, then

$$\begin{aligned} E &= k\iint_D \left[1 - \tfrac{1}{20}\sqrt{x^2 + y^2}\right] dA \\ &= k\int_0^{2\pi}\int_0^{10} \left(1 - \tfrac{1}{20}r\right) r\,dr\,d\theta = 2\pi k\left[\tfrac{1}{2}r^2 - \tfrac{1}{60}r^3\right]_0^{10} \\ &= 2\pi k\left(50 - \tfrac{50}{3}\right) = \tfrac{200}{3}\pi k \approx 209k \end{aligned}$$

For A at the edge of the city, it is convenient to use a polar coordinate system centered at A. Then the polar equation for the circular boundary of the city becomes $r = 20\cos\theta$ instead of $r = 10$, and the distance from A to a point P in the city is again r (see the figure). So

$$\begin{aligned} E &= k\int_{-\pi/2}^{\pi/2}\int_0^{20\cos\theta} \left(1 - \tfrac{1}{20}r\right) r\,dr\,d\theta = k\int_{-\pi/2}^{\pi/2} \left[\tfrac{1}{2}r^2 - \tfrac{1}{60}r^3\right]_{r=0}^{r=20\cos\theta} d\theta \\ &= k\textstyle\int_{-\pi/2}^{\pi/2} \left(200\cos^2\theta - \frac{400}{3}\cos^3\theta\right) d\theta = 200k\int_{-\pi/2}^{\pi/2} \left[\frac{1}{2} + \frac{1}{2}\cos 2\theta - \frac{2}{3}\left(1 - \sin^2\theta\right)\cos\theta\right] d\theta \\ &= 200k\left[\tfrac{1}{2}\theta + \tfrac{1}{4}\sin 2\theta - \tfrac{2}{3}\sin\theta + \tfrac{2}{3}\cdot\tfrac{1}{3}\sin^3\theta\right]_{-\pi/2}^{\pi/2} = 200k\left[\tfrac{\pi}{4} + 0 - \tfrac{2}{3} + \tfrac{2}{9} + \tfrac{\pi}{4} + 0 - \tfrac{2}{3} + \tfrac{2}{9}\right] \\ &= 200k\left(\tfrac{\pi}{2} - \tfrac{8}{9}\right) \approx 136k \end{aligned}$$

Therefore the risk of infection is much lower at the edge of the city than in the middle, so it is better to live at the edge.

15.5 Surface Area

1. Here $z = f(x, y) = 10 + x + y^2$ and D is the triangle with vertices $(0, 0)$, $(0, -2)$, and $(2, -2)$. By Formula 2, the area of the surface is

$$\begin{aligned} A(S) &= \textstyle\iint_D \sqrt{[f_x(x, y)]^2 + [f_y(x, y)]^2 + 1}\,dA = \int_{-2}^{0}\int_0^{-y} \sqrt{1^2 + (2y)^2 + 1}\,dx\,dy \\ &= \textstyle\int_{-2}^{0}\int_0^{-y} \sqrt{2 + 4y^2}\,dx\,dy = \int_{-2}^{0} \sqrt{2 + 4y^2}\,[x]_{x=0}^{x=-y}\,dy = -\int_{-2}^{0} y\sqrt{2 + 4y^2}\,dy \\ &= -\frac{1}{8}\cdot\left[\frac{2}{3}(2 + 4y^2)^{3/2}\right]_{-2}^{0} = \frac{18^{3/2} - 2^{3/2}}{12} = \frac{54\sqrt{2} - 2\sqrt{2}}{12} = \frac{13\sqrt{2}}{3} \end{aligned}$$

2. Here $z = f(x, y) = 3 + xy$ and D is the circle $x^2 + y^2 \le 1$. By Formula 2, the area of the surface is

$$\begin{aligned} A(S) &= \iint_D \sqrt{[f_x(x, y)]^2 + [f_y(x, y)]^2 + 1}\,dA = \iint_{x^2+y^2\le 1} \sqrt{y^2 + x^2 + 1}\,dA \qquad \text{[Switch to polar coordinates]} \\ &= \textstyle\int_0^{2\pi}\int_0^1 \sqrt{r^2 + 1}\,r\,dr\,d\theta = \int_0^{2\pi} d\theta\int_0^1 \sqrt{r^2 + 1}\,r\,dr = 2\pi\cdot\frac{1}{2}\left[\frac{2}{3}(r^2 + 1)^{3/2}\right]_{r=0}^{r=1} = \frac{2\pi}{3}(2^{3/2} - 1) \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. Here $z = f(x, y) = 5x + 3y + 6$ and D is the rectangle $[1, 4] \times [2, 6]$. By Formula 2, the area of the surface is

$$\begin{aligned} A(S) &= \iint_D \sqrt{[f_x(x,y)]^2 + [f_y(x,y)]^2 + 1}\, dA = \iint_D \sqrt{5^2 + 3^2 + 1}\, dA = \sqrt{35} \iint_D dA \\ &= \sqrt{35}\, A(D) = \sqrt{35}\,(4-1)(6-2) = 12\sqrt{35} \end{aligned}$$

4. Here $z = f(x, y) = \frac{1}{2} - 3x - 2y$ and D is the disk $x^2 + y^2 \le 25$. By Formula 2, the area of the surface is

$$A(S) = \iint_D \sqrt{(-3)^2 + (-2)^2 + 1}\, dA = \sqrt{14} \iint_D dA = \sqrt{14}\, A(D) = \sqrt{14}\,(\pi \cdot 5^2) = 25\sqrt{14}\,\pi$$

5. The surface S is given by $z = f(x, y) = 6 - 3x - 2y$ which intersects the xy-plane in the line $3x + 2y = 6$, so D is the triangular region given by $\left\{(x, y) \;\middle|\; 0 \le x \le 2,\, 0 \le y \le 3 - \frac{3}{2}x\right\}$. By Formula 2, the surface area of S is

$$A(S) = \iint_D \sqrt{(-3)^2 + (-2)^2 + 1}\, dA = \sqrt{14} \iint_D dA = \sqrt{14}\, A(D) = \sqrt{14}\left(\tfrac{1}{2} \cdot 2 \cdot 3\right) = 3\sqrt{14}$$

6. $z = f(x, y) = \frac{1}{4}x^2 - \frac{1}{2}y + \frac{5}{4}$, and D is the triangular region given by $\{(x, y) \mid 0 \le x \le 2,\ 0 \le y \le 2x\}$. By Formula 2,

$$\begin{aligned} A(S) &= \iint_D \sqrt{\left(\tfrac{1}{2}x\right)^2 + \left(-\tfrac{1}{2}\right)^2 + 1}\, dA = \int_0^2 \int_0^{2x} \sqrt{\tfrac{1}{4}x^2 + \tfrac{5}{4}}\, dy\, dx = \int_0^2 \tfrac{1}{2}\sqrt{x^2 + 5}\ \left[\, y \,\right]_{y=0}^{y=2x} dx \\ &= \tfrac{1}{2}\int_0^2 2x\sqrt{x^2 + 5}\, dx = \tfrac{1}{2} \cdot \tfrac{2}{3}(x^2 + 5)^{3/2}\Big]_0^2 = \tfrac{1}{3}(9^{3/2} - 5^{3/2}) = 9 - \tfrac{5}{3}\sqrt{5} \end{aligned}$$

7. The paraboloid intersects the plane $z = -2$ when $1 - x^2 - y^2 = -2 \quad \Leftrightarrow \quad x^2 + y^2 = 3$, so $D = \left\{(x, y) \mid x^2 + y^2 \le 3\right\}$. Here $z = f(x, y) = 1 - x^2 - y^2 \quad \Rightarrow \quad f_x = -2x,\ f_y = -2y$ and

$$\begin{aligned} A(S) &= \iint_D \sqrt{(-2x)^2 + (-2y)^2 + 1}\, dA = \iint_D \sqrt{4(x^2 + y^2) + 1}\, dA = \int_0^{2\pi}\int_0^{\sqrt{3}} \sqrt{4r^2 + 1}\, r\, dr\, d\theta \\ &= \int_0^{2\pi} d\theta \int_0^{\sqrt{3}} r\sqrt{4r^2 + 1}\, dr = \left[\,\theta\,\right]_0^{2\pi} \left[\tfrac{1}{12}(4r^2 + 1)^{3/2}\right]_0^{\sqrt{3}} = 2\pi \cdot \tfrac{1}{12}\left(13^{3/2} - 1\right) = \tfrac{\pi}{6}\left(13\sqrt{13} - 1\right) \end{aligned}$$

8. $x^2 + z^2 = 4 \quad \Rightarrow \quad z = \sqrt{4 - x^2}$ (since $z \ge 0$), so $f_x = -x(4 - x^2)^{-1/2},\ f_y = 0$ and

$$\begin{aligned} A(S) &= \int_0^1 \int_0^1 \sqrt{[-x(4 - x^2)^{-1/2}]^2 + 0^2 + 1}\, dy\, dx = \int_0^1 \int_0^1 \sqrt{\frac{x^2}{4 - x^2} + 1}\, dy\, dx \\ &= \int_0^1 \frac{2}{\sqrt{4 - x^2}}\, dx \int_0^1 dy = \left[2\sin^{-1}\frac{x}{2}\right]_0^1 \left[\, y \,\right]_0^1 = \left(2 \cdot \tfrac{\pi}{6} - 0\right)(1) = \tfrac{\pi}{3} \end{aligned}$$

9. $z = f(x, y) = y^2 - x^2$ with $1 \le x^2 + y^2 \le 4$. Then

$$\begin{aligned} A(S) &= \iint_D \sqrt{4x^2 + 4y^2 + 1}\, dA = \int_0^{2\pi}\int_1^2 \sqrt{4r^2 + 1}\, r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_1^2 r\sqrt{4r^2 + 1}\, dr \\ &= \left[\,\theta\,\right]_0^{2\pi} \left[\tfrac{1}{12}(4r^2 + 1)^{3/2}\right]_1^2 = \tfrac{\pi}{6}\left(17\sqrt{17} - 5\sqrt{5}\right) \end{aligned}$$

10. $z = f(x, y) = \frac{2}{3}(x^{3/2} + y^{3/2})$ and $D = \{(x, y) \mid 0 \le x \le 1, 0 \le y \le 1\}$. Then $f_x = x^{1/2}$, $f_y = y^{1/2}$ and

$$\begin{aligned} A(S) &= \iint_D \sqrt{\left(\sqrt{x}\right)^2 + \left(\sqrt{y}\right)^2 + 1}\, dA = \int_0^1 \int_0^1 \sqrt{x + y + 1}\, dy\, dx = \int_0^1 \left[\tfrac{2}{3}(x + y + 1)^{3/2}\right]_{y=0}^{y=1} dx \\ &= \tfrac{2}{3}\int_0^1 \left[(x + 2)^{3/2} - (x + 1)^{3/2}\right] dx = \tfrac{2}{3}\left[\tfrac{2}{5}(x + 2)^{5/2} - \tfrac{2}{5}(x + 1)^{5/2}\right]_0^1 \\ &= \tfrac{4}{15}(3^{5/2} - 2^{5/2} - 2^{5/2} + 1) = \tfrac{4}{15}(3^{5/2} - 2^{7/2} + 1) \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. $z = f(x, y) = xy$ with $x^2 + y^2 \le 1$, so $f_x = y$, $f_y = x \quad \Rightarrow$

$$A(S) = \iint_D \sqrt{y^2 + x^2 + 1}\, dA = \int_0^{2\pi} \int_0^1 \sqrt{r^2 + 1}\, r\, dr\, d\theta = \int_0^{2\pi} \left[\tfrac{1}{3}(r^2 + 1)^{3/2}\right]_{r=0}^{r=1} d\theta$$
$$= \int_0^{2\pi} \tfrac{1}{3}\left(2\sqrt{2} - 1\right) d\theta = \tfrac{2\pi}{3}\left(2\sqrt{2} - 1\right)$$

12. Given the sphere $x^2 + y^2 + z^2 = 4$, when $z = 1$, we get $x^2 + y^2 = 3$ so $D = \{(x, y) \mid x^2 + y^2 \le 3\}$ and $z = f(x, y) = \sqrt{4 - x^2 - y^2}$. Thus

$$A(S) = \iint_D \sqrt{[(-x)(4 - x^2 - y^2)^{-1/2}]^2 + [(-y)(4 - x^2 - y^2)^{-1/2}]^2 + 1}\, dA$$
$$= \int_0^{2\pi} \int_0^{\sqrt{3}} \sqrt{\frac{r^2}{4 - r^2} + 1}\, r\, dr\, d\theta = \int_0^{2\pi} \int_0^{\sqrt{3}} \sqrt{\frac{r^2 + 4 - r^2}{4 - r^2}}\, r\, dr\, d\theta$$
$$= \int_0^{2\pi} \int_0^{\sqrt{3}} \frac{2r}{\sqrt{4 - r^2}}\, dr\, d\theta$$
$$= \int_0^{2\pi} \left[-2(4 - r^2)^{1/2}\right]_{r=0}^{r=\sqrt{3}} d\theta = \int_0^{2\pi} (-2 + 4)\, d\theta = 2\theta\Big]_0^{2\pi} = 4\pi$$

13. $z = \sqrt{a^2 - x^2 - y^2}$, $z_x = -x(a^2 - x^2 - y^2)^{-1/2}$, $z_y = -y(a^2 - x^2 - y^2)^{-1/2}$,

$$A(S) = \iint_D \sqrt{\frac{x^2 + y^2}{a^2 - x^2 - y^2} + 1}\, dA$$
$$= \int_{-\pi/2}^{\pi/2} \int_0^{a\cos\theta} \sqrt{\frac{r^2}{a^2 - r^2} + 1}\, r\, dr\, d\theta$$
$$= \int_{-\pi/2}^{\pi/2} \int_0^{a\cos\theta} \frac{ar}{\sqrt{a^2 - r^2}}\, dr\, d\theta$$
$$= \int_{-\pi/2}^{\pi/2} \left[-a\sqrt{a^2 - r^2}\right]_{r=0}^{r=a\cos\theta} d\theta$$
$$= \int_{-\pi/2}^{\pi/2} -a\left(\sqrt{a^2 - a^2\cos^2\theta} - a\right) d\theta = 2a^2 \int_0^{\pi/2} \left(1 - \sqrt{1 - \cos^2\theta}\right) d\theta$$
$$= 2a^2 \int_0^{\pi/2} d\theta - 2a^2 \int_0^{\pi/2} \sqrt{\sin^2\theta}\, d\theta = a^2\pi - 2a^2 \int_0^{\pi/2} \sin\theta\, d\theta = a^2(\pi - 2)$$

14. To find the region D: $z = x^2 + y^2$ implies $z + z^2 = 4z$ or $z^2 - 3z = 0$. Thus $z = 0$ or $z = 3$ are the planes where the surfaces intersect. But $x^2 + y^2 + z^2 = 4z$ implies $x^2 + y^2 + (z - 2)^2 = 4$, so $z = 3$ intersects the upper hemisphere. Thus $(z - 2)^2 = 4 - x^2 - y^2$ or $z = 2 + \sqrt{4 - x^2 - y^2}$. Therefore D is the region inside the circle $x^2 + y^2 + (3 - 2)^2 = 4$, that is, $D = \{(x, y) \mid x^2 + y^2 \le 3\}$.

$$A(S) = \iint_D \sqrt{[(-x)(4 - x^2 - y^2)^{-1/2}]^2 + [(-y)(4 - x^2 - y^2)^{-1/2}]^2 + 1}\, dA$$
$$= \int_0^{2\pi} \int_0^{\sqrt{3}} \sqrt{\frac{r^2}{4 - r^2} + 1}\, r\, dr\, d\theta = \int_0^{2\pi} \int_0^{\sqrt{3}} \frac{2r\, dr}{\sqrt{4 - r^2}}\, d\theta = \int_0^{2\pi} \left[-2(4 - r^2)^{1/2}\right]_{r=0}^{r=\sqrt{3}} d\theta$$
$$= \int_0^{2\pi} (-2 + 4)\, d\theta = 2\theta\Big]_0^{2\pi} = 4\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $z = f(x, y) = (1 + x^2 + y^2)^{-1}$, $f_x = -2x(1 + x^2 + y^2)^{-2}$, $f_y = -2y(1 + x^2 + y^2)^{-2}$. Then

$$\begin{aligned} A(S) &= \iint\limits_{x^2+y^2 \le 1} \sqrt{[-2x(1 + x^2 + y^2)^{-2}]^2 + [-2y(1 + x^2 + y^2)^{-2}]^2 + 1}\ dA \\ &= \iint\limits_{x^2+y^2 \le 1} \sqrt{4(x^2 + y^2)(1 + x^2 + y^2)^{-4} + 1}\ dA \end{aligned}$$

Converting to polar coordinates we have

$$\begin{aligned} A(S) &= \int_0^{2\pi} \int_0^1 \sqrt{4r^2(1 + r^2)^{-4} + 1}\ r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_0^1 r \sqrt{4r^2(1 + r^2)^{-4} + 1}\, dr \\ &= 2\pi \int_0^1 r \sqrt{4r^2(1 + r^2)^{-4} + 1}\, dr \approx 3.6258 \ \text{ using a calculator.} \end{aligned}$$

16. $z = f(x, y) = \cos(x^2 + y^2)$, $f_x = -2x \sin(x^2 + y^2)$, $f_y = -2y \sin(x^2 + y^2)$.

$$A(S) = \iint\limits_{x^2+y^2 \le 1} \sqrt{4x^2 \sin^2(x^2 + y^2) + 4y^2 \sin^2(x^2 + y^2) + 1}\, dA = \iint\limits_{x^2+y^2 \le 1} \sqrt{4(x^2 + y^2) \sin^2(x^2 + y^2) + 1}\, dA.$$

Converting to polar coordinates gives

$$\begin{aligned} A(S) &= \int_0^{2\pi} \int_0^1 \sqrt{4r^2 \sin^2(r^2) + 1}\, r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_0^1 r \sqrt{4r^2 \sin^2(r^2) + 1}\, dr \\ &= 2\pi \int_0^1 r \sqrt{4r^2 \sin^2(r^2) + 1}\, dr \approx 4.1073 \quad \text{using a calculator.} \end{aligned}$$

17. (a) The midpoints of the four squares are $\left(\frac{1}{4}, \frac{1}{4}\right)$, $\left(\frac{1}{4}, \frac{3}{4}\right)$, $\left(\frac{3}{4}, \frac{1}{4}\right)$, and $\left(\frac{3}{4}, \frac{3}{4}\right)$. Here $f(x, y) = x^2 + y^2$, so the Midpoint Rule gives

$$\begin{aligned} A(S) &= \iint_D \sqrt{[f_x(x, y)]^2 + [f_y(x, y)]^2 + 1}\, dA = \iint_D \sqrt{(2x)^2 + (2y)^2 + 1}\, dA \\ &\approx \tfrac{1}{4}\Bigg(\sqrt{\left[2\left(\tfrac{1}{4}\right)\right]^2 + \left[2\left(\tfrac{1}{4}\right)\right]^2 + 1} + \sqrt{\left[2\left(\tfrac{1}{4}\right)\right]^2 + \left[2\left(\tfrac{3}{4}\right)\right]^2 + 1} \\ &\qquad + \sqrt{\left[2\left(\tfrac{3}{4}\right)\right]^2 + \left[2\left(\tfrac{1}{4}\right)\right]^2 + 1} + \sqrt{\left[2\left(\tfrac{3}{4}\right)\right]^2 + \left[2\left(\tfrac{3}{4}\right)\right]^2 + 1}\Bigg) \\ &= \tfrac{1}{4}\left(\sqrt{\tfrac{3}{2}} + 2\sqrt{\tfrac{7}{2}} + \sqrt{\tfrac{11}{2}}\right) \approx 1.8279 \end{aligned}$$

(b) A CAS estimates the integral to be $A(S) = \iint_D \sqrt{1 + (2x)^2 + (2y)^2}\, dA = \int_0^1 \int_0^1 \sqrt{1 + 4x^2 + 4y^2}\, dy\, dx \approx 1.8616$. This agrees with the Midpoint estimate only in the first decimal place.

18. (a) With $m = n = 2$ we have four squares with midpoints $\left(\frac{1}{2}, \frac{1}{2}\right)$, $\left(\frac{1}{2}, \frac{3}{2}\right)$, $\left(\frac{3}{2}, \frac{1}{2}\right)$, and $\left(\frac{3}{2}, \frac{3}{2}\right)$. Since $z = xy + x^2 + y^2$, the Midpoint Rule gives

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}\, dA = \iint_D \sqrt{1 + (y + 2x)^2 + (x + 2y)^2}\, dA \\ &\approx 1\left(\sqrt{1 + \left(\tfrac{3}{2}\right)^2 + \left(\tfrac{3}{2}\right)^2} + \sqrt{1 + \left(\tfrac{5}{2}\right)^2 + \left(\tfrac{7}{2}\right)^2} + \sqrt{1 + \left(\tfrac{7}{2}\right)^2 + \left(\tfrac{5}{2}\right)^2} + \sqrt{1 + \left(\tfrac{9}{2}\right)^2 + \left(\tfrac{9}{2}\right)^2}\right) \\ &= \tfrac{\sqrt{22}}{2} + \tfrac{\sqrt{78}}{2} + \tfrac{\sqrt{78}}{2} + \tfrac{\sqrt{166}}{2} \approx 17.619 \end{aligned}$$

(b) Using a CAS, we have

$A(S) = \iint_D \sqrt{1 + (y + 2x)^2 + (x + 2y)^2}\, dA = \int_0^2 \int_0^2 \sqrt{1 + (y + 2x)^2 + (x + 2y)^2}\, dy\, dx \approx 17.7165$. This is within about 0.1 of the Midpoint Rule estimate.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $z = 1 + 2x + 3y + 4y^2$, so

$$A(S) = \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}\, dA = \int_1^4 \int_0^1 \sqrt{1 + 4 + (3 + 8y)^2}\, dy\, dx = \int_1^4 \int_0^1 \sqrt{14 + 48y + 64y^2}\, dy\, dx.$$

Using a CAS, we have $\int_1^4 \int_0^1 \sqrt{14 + 48y + 64y^2}\, dy\, dx = \frac{45}{8}\sqrt{14} + \frac{15}{16}\ln\left(11\sqrt{5} + 3\sqrt{14}\sqrt{5}\right) - \frac{15}{16}\ln\left(3\sqrt{5} + \sqrt{14}\sqrt{5}\right)$ or $\frac{45}{8}\sqrt{14} + \frac{15}{16}\ln\dfrac{11\sqrt{5} + 3\sqrt{70}}{3\sqrt{5} + \sqrt{70}}$.

20. $f(x, y) = 1 + x + y + x^2 \quad \Rightarrow \quad f_x = 1 + 2x$, $f_y = 1$. We use a CAS to calculate the integral

$$A(S) = \int_{-2}^1 \int_{-1}^1 \sqrt{f_x^2 + f_y^2 + 1}\, dy\, dx$$
$$= \int_{-2}^1 \int_{-1}^1 \sqrt{(1 + 2x)^2 + 2}\, dy\, dx = 2\int_{-2}^1 \sqrt{4x^2 + 4x + 3}\, dx$$

and find that $A(S) = 3\sqrt{11} + 2\sinh^{-1}\left(\frac{3\sqrt{2}}{2}\right)$ or

$A(S) = 3\sqrt{11} + \ln\left(10 + 3\sqrt{11}\right)$.

21. $f(x, y) = 1 + x^2y^2 \quad \Rightarrow \quad f_x = 2xy^2$, $f_y = 2x^2y$. We use a CAS (with precision reduced to five significant digits, to speed up the calculation) to estimate the integral

$$A(S) = \int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \sqrt{f_x^2 + f_y^2 + 1}\, dy\, dx = \int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \sqrt{4x^2y^4 + 4x^4y^2 + 1}\, dy\, dx,$$ and find that $A(S) \approx 3.3213$.

22. Let $f(x, y) = \dfrac{1 + x^2}{1 + y^2}$. Then $f_x = \dfrac{2x}{1 + y^2}$,

$f_y = \left(1 + x^2\right)\left[-\dfrac{2y}{(1 + y^2)^2}\right] = -\dfrac{2y\left(1 + x^2\right)}{(1 + y^2)^2}$. We use a CAS to estimate $\int_{-1}^1 \int_{-(1-|x|)}^{1-|x|} \sqrt{f_x^2 + f_y^2 + 1}\, dy\, dx \approx 2.6959$. In order to graph only the part of the surface above the square, we use $-(1 - |x|) \le y \le 1 - |x|$ as the y-range in our plot command.

23. Here $z = f(x, y) = ax + by + c$, $f_x(x, y) = a$, $f_y(x, y) = b$, so

$A(S) = \iint_D \sqrt{a^2 + b^2 + 1}\, dA = \sqrt{a^2 + b^2 + 1}\iint_D dA = \sqrt{a^2 + b^2 + 1}\, A(D)$.

24. Let S be the upper hemisphere. Then $z = f(x, y) = \sqrt{a^2 - x^2 - y^2}$, so

$$A(S) = \iint_D \sqrt{[-x(a^2 - x^2 - y^2)^{-1/2}]^2 + [-y(a^2 - x^2 - y^2)^{-1/2}]^2 + 1}\, dA$$
$$= \iint_D \sqrt{\frac{x^2 + y^2}{a^2 - x^2 - y^2} + 1}\, dA = \lim_{t \to a^-} \int_0^{2\pi} \int_0^t \sqrt{\frac{r^2}{a^2 - r^2} + 1}\, r\, dr\, d\theta$$
$$= \lim_{t \to a^-} \int_0^{2\pi} \int_0^t \frac{ar}{\sqrt{a^2 - r^2}}\, dr\, d\theta = 2\pi \lim_{t \to a^-} \left[-a\sqrt{a^2 - r^2}\right]_0^t = 2\pi \lim_{t \to a^-} -a\left[\sqrt{a^2 - t^2} - a\right]$$
$$= 2\pi(-a)(-a) = 2\pi a^2.$$ Thus the surface area of the entire sphere is $4\pi a^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. If we project the surface onto the xz-plane, then the surface lies "above" the disk $x^2+z^2 \le 25$ in the xz-plane.

We have $y = f(x,z) = x^2+z^2$ and, adapting Formula 2, the area of the surface is

$$A(S) = \iint\limits_{x^2+z^2\le 25} \sqrt{[f_x(x,z)]^2+[f_z(x,z)]^2+1}\,dA = \iint\limits_{x^2+z^2\le 25} \sqrt{4x^2+4z^2+1}\,dA$$

Converting to polar coordinates $x = r\cos\theta$, $z = r\sin\theta$ we have

$$A(S) = \int_0^{2\pi}\int_0^5 \sqrt{4r^2+1}\,r\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^5 r(4r^2+1)^{1/2}\,dr = \left[\theta\right]_0^{2\pi}\left[\tfrac{1}{12}(4r^2+1)^{3/2}\right]_0^5 = \tfrac{\pi}{6}\left(101\sqrt{101}-1\right)$$

26. First we find the area of the face of the surface that intersects the positive y-axis. As in Exercise 25, we can project the face onto the xz-plane, so the surface lies "above" the disk $x^2+z^2 \le 1$. Then $y = f(x,z) = \sqrt{1-z^2}$ and the area is

$$\begin{aligned} A(S) &= \iint\limits_{x^2+z^2\le 1} \sqrt{[f_x(x,z)]^2+[f_z(x,z)]^2+1}\,dA = \iint\limits_{x^2+z^2\le 1} \sqrt{0+\left(\frac{-z}{\sqrt{1-z^2}}\right)^2+1}\,dA \\ &= \iint\limits_{x^2+z^2\le 1} \sqrt{\frac{z^2}{1-z^2}+1}\,dA = \int_{-1}^{1}\int_{-\sqrt{1-z^2}}^{\sqrt{1-z^2}} \frac{1}{\sqrt{1-z^2}}\,dx\,dz \\ &= 4\int_0^1\int_0^{\sqrt{1-z^2}} \frac{1}{\sqrt{1-z^2}}\,dx\,dz \qquad \text{[by the symmetry of the surface]} \end{aligned}$$

This integral is improper (when $z=1$), so

$$A(S) = \lim_{t\to 1^-} 4\int_0^t\int_0^{\sqrt{1-z^2}} \frac{1}{\sqrt{1-z^2}}\,dx\,dz = \lim_{t\to 1^-} 4\int_0^t \frac{\sqrt{1-z^2}}{\sqrt{1-z^2}}\,dz = \lim_{t\to 1^-} 4\int_0^t dz = \lim_{t\to 1^-} 4t = 4.$$

Since the complete surface consists of four congruent faces, the total surface area is $4(4) = 16$.

15.6 Triple Integrals

1. $$\begin{aligned} \iiint_B xyz^2\,dV &= \int_0^1\int_0^3\int_{-1}^2 xyz^2\,dy\,dz\,dx = \int_0^1\int_0^3 \left[\tfrac{1}{2}xy^2z^2\right]_{y=-1}^{y=2} dz\,dx = \int_0^1\int_0^3 \tfrac{3}{2}xz^2\,dz\,dx \\ &= \int_0^1 \left[\tfrac{1}{2}xz^3\right]_{z=0}^{z=3} dx = \int_0^1 \tfrac{27}{2}\,x\,dx = \left[\tfrac{27}{4}x^2\right]_0^1 = \tfrac{27}{4} \end{aligned}$$

2. There are six different possible orders of integration.

$$\begin{aligned} \iiint_E (xy+z^2)\,dV &= \int_0^2\int_0^1\int_0^3 (xy+z^2)\,dz\,dy\,dx = \int_0^2\int_0^1 \left[xyz+\tfrac{1}{3}z^3\right]_{z=0}^{z=3} dy\,dx = \int_0^2\int_0^1 (3xy+9)\,dy\,dx \\ &= \int_0^2 \left[\tfrac{3}{2}xy^2+9y\right]_{y=0}^{y=1} dx = \int_0^2 \left(\tfrac{3}{2}x+9\right) dx = \left[\tfrac{3}{4}x^2+9x\right]_0^2 = 21 \end{aligned}$$

$$\begin{aligned} \iiint_E (xy+z^2)\,dV &= \int_0^1\int_0^2\int_0^3 (xy+z^2)\,dz\,dx\,dy = \int_0^1\int_0^2 \left[xyz+\tfrac{1}{3}z^3\right]_{z=0}^{z=3} dx\,dy = \int_0^1\int_0^2 (3xy+9)\,dx\,dy \\ &= \int_0^1 \left[\tfrac{3}{2}x^2y+9x\right]_{x=0}^{x=2} dy = \int_0^1 (6y+18)\,dy = \left[3y^2+18y\right]_0^1 = 21 \end{aligned}$$

$$\begin{aligned} \iiint_E (xy+z^2)\,dV &= \int_0^2\int_0^3\int_0^1 (xy+z^2)\,dy\,dz\,dx = \int_0^2\int_0^3 \left[\tfrac{1}{2}xy^2+yz^2\right]_{y=0}^{y=1} dz\,dx = \int_0^2\int_0^3 \left(\tfrac{1}{2}x+z^2\right) dz\,dx \\ &= \int_0^2 \left[\tfrac{1}{2}xz+\tfrac{1}{3}z^3\right]_{z=0}^{z=3} dx = \int_0^2 \left(\tfrac{3}{2}x+9\right) dx = \left[\tfrac{3}{4}x^2+9x\right]_0^2 = 21 \end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\iiint_E (xy+z^2)\,dV = \int_0^3\int_0^2\int_0^1 (xy+z^2)\,dy\,dx\,dz = \int_0^3\int_0^2 \left[\tfrac{1}{2}xy^2+yz^2\right]_{y=0}^{y=1} dx\,dz = \int_0^3\int_0^2 \left(\tfrac{1}{2}x+z^2\right) dx\,dz$$
$$= \int_0^3 \left[\tfrac{1}{4}x^2+xz^2\right]_{x=0}^{x=2} dz = \int_0^3 \left(1+2z^2\right) dz = \left[z+\tfrac{2}{3}z^3\right]_0^3 = 21$$

$$\iiint_E (xy+z^2)\,dV = \int_0^1\int_0^3\int_0^2 (xy+z^2)\,dx\,dz\,dy = \int_0^1\int_0^3 \left[\tfrac{1}{2}x^2y+xz^2\right]_{x=0}^{x=2} dz\,dy = \int_0^1\int_0^3 \left(2y+2z^2\right) dz\,dy$$
$$= \int_0^1 \left[2yz+\tfrac{2}{3}z^3\right]_{z=0}^{z=3} dy = \int_0^1 (6y+18)\,dy = \left[3y^2+18y\right]_0^1 = 21$$

$$\iiint_E (xy+z^2)\,dV = \int_0^3\int_0^1\int_0^2 (xy+z^2)\,dx\,dy\,dz = \int_0^3\int_0^1 \left[\tfrac{1}{2}x^2y+xz^2\right]_{x=0}^{x=2} dy\,dz = \int_0^3\int_0^1 \left(2y+2z^2\right) dy\,dz$$
$$= \int_0^3 \left[y^2+2yz^2\right]_{y=0}^{y=1} dz = \int_0^3 \left(1+2z^2\right) dz = \left[z+\tfrac{2}{3}z^3\right]_0^3 = 21$$

3. $$\int_0^2\int_0^{z^2}\int_0^{y-z}(2x-y)\,dx\,dy\,dz = \int_0^2\int_0^{z^2} \left[x^2-xy\right]_{x=0}^{x=y-z} dy\,dz = \int_0^2\int_0^{z^2} \left[(y-z)^2-(y-z)y\right] dy\,dz$$
$$= \int_0^2\int_0^{z^2} \left(z^2-yz\right) dy\,dz = \int_0^2 \left[yz^2-\tfrac{1}{2}y^2z\right]_{y=0}^{y=z^2} dz = \int_0^2 \left(z^4-\tfrac{1}{2}z^5\right) dz$$
$$= \left[\tfrac{1}{5}z^5-\tfrac{1}{12}z^6\right]_0^2 = \tfrac{32}{5}-\tfrac{64}{12} = \tfrac{16}{15}$$

4. $$\int_0^1\int_y^{2y}\int_0^{x+y} 6xy\,dz\,dx\,dy = \int_0^1\int_y^{2y} \left[6xyz\right]_{z=0}^{z=x+y} dx\,dy = \int_0^1\int_y^{2y} 6xy(x+y)\,dx\,dy = \int_0^1\int_y^{2y} (6x^2y+6xy^2)\,dx\,dy$$
$$= \int_0^1 \left[2x^3y+3x^2y^2\right]_{x=y}^{x=2y} dy = \int_0^1 23y^4\,dy = \tfrac{23}{5}y^5\Big]_0^1 = \tfrac{23}{5}$$

5. $$\int_1^2\int_0^{2z}\int_0^{\ln x} xe^{-y}\,dy\,dx\,dz = \int_1^2\int_0^{2z} \left[-xe^{-y}\right]_{y=0}^{y=\ln x} dx\,dz = \int_1^2\int_0^{2z} \left(-xe^{-\ln x}+xe^0\right) dx\,dz$$
$$= \int_1^2\int_0^{2z} (-1+x)\,dx\,dz = \int_1^2 \left[-x+\tfrac{1}{2}x^2\right]_{x=0}^{x=2z} dz$$
$$= \int_1^2 \left(-2z+2z^2\right) dz = \left[-z^2+\tfrac{2}{3}z^3\right]_1^2 = -4+\tfrac{16}{3}+1-\tfrac{2}{3} = \tfrac{5}{3}$$

6. $$\int_0^{\pi/2}\int_0^{2x}\int_0^{x+z} \cos(x-2y+z)\,dy\,dz\,dx = -\tfrac{1}{2}\int_0^{\pi/2}\int_0^{2x} \left[\sin(x-2y+z)\right]_{y=0}^{y=x+z} dz\,dx$$
$$= -\tfrac{1}{2}\int_0^{\pi/2}\int_0^{2x} \left[\sin(-x-z)-\sin(x+z)\right] dz\,dx$$
$$= -\tfrac{1}{2}\int_0^{\pi/2}\int_0^{2x} \left[-2\sin(x+z)\right] dz\,dx$$
$$= \int_0^{\pi/2}\int_0^{2x} \sin(x+z)\,dz\,dx = -\int_0^{\pi/2} \left[\cos(x+z)\right]_{z=0}^{z=2x} dx$$
$$= -\int_0^{\pi/2}(\cos 3x-\cos x)\,dx = \left[\sin x-\tfrac{1}{3}\sin 3x\right]_0^{\pi/2}$$
$$= 1-\tfrac{1}{3}(-1) = \tfrac{4}{3}$$

7. $$\int_1^3\int_{-1}^2\int_{-y}^z \frac{z}{y}\,dx\,dz\,dy = \int_1^3\int_{-1}^2 \frac{z}{y}\left[x\right]_{x=-y}^{x=z} dz\,dy = \int_1^3\int_{-1}^2 \left(\frac{z^2}{y}+z\right) dz\,dy = \int_1^3 \left[\frac{z^3}{3y}+\frac{z^2}{2}\right]_{z=-1}^{z=2} dy$$
$$= \int_1^3 \left(\frac{3}{y}+\frac{3}{2}\right) dy = \left[3\ln|y|+\frac{3}{2}y\right]_1^3 = 3\ln 3+3$$

8. $$\int_0^1\int_0^1\int_0^{2-x^2-y^2} xye^z\,dz\,dy\,dx = \int_0^1\int_0^1 \left[xye^z\right]_{z=0}^{z=2-x^2-y^2} dy\,dx = \int_0^1\int_0^1 (xye^{2-x^2-y^2}-xy)\,dy\,dx$$
$$= \int_0^1 \left[-\tfrac{1}{2}xe^{2-x^2-y^2}-\tfrac{1}{2}xy^2\right]_{y=0}^{y=1} dx = \int_0^1 \left(-\tfrac{1}{2}xe^{1-x^2}-\tfrac{1}{2}x+\tfrac{1}{2}xe^{2-x^2}\right) dx$$
$$= \left[\tfrac{1}{4}e^{1-x^2}-\tfrac{1}{4}x^2-\tfrac{1}{4}e^{2-x^2}\right]_0^1 = \tfrac{1}{4}-\tfrac{1}{4}-\tfrac{1}{4}e-\tfrac{1}{4}e+0+\tfrac{1}{4}e^2 = \tfrac{1}{4}e^2-\tfrac{1}{2}e$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. (a) The solid region E can be described as $E = \{(x, y, z) \mid -1 \le x \le 1, 0 \le y \le 2 - z, 0 \le z \le 1 - x^2\}$.

Thus, $\iiint_E x\,dV = \int_{-1}^{1} \int_0^{1-x^2} \int_0^{2-z} x\,dy\,dz\,dx$.

(b) $$\int_{-1}^{1} \int_0^{1-x^2} \int_0^{2-z} x\,dy\,dz\,dx = \int_{-1}^{1} \int_0^{1-x^2} x \Big[y \Big]_{y=0}^{y=2-z} dz\,dx = \int_{-1}^{1} \int_0^{1-x^2} (2x - xz)\,dz\,dx$$
$$= \int_{-1}^{1} \left[2xz - x\frac{z^2}{2} \right]_{z=0}^{z=1-x^2} dx = \int_{-1}^{1} \left(\frac{3}{2}x - x^3 - \frac{x^5}{2} \right) dx$$
$$= \left[\frac{3x^2}{4} - \frac{x^4}{4} - \frac{x^6}{12} \right]_{-1}^{1} = 0$$

10. (a) The solid region E can be described as $E = \{(x, y, z) \mid 0 \le x \le y, 0 \le y \le 2, 0 \le z \le 4 - y^2\}$.

Thus, $\iiint_E xy\,dV = \int_0^2 \int_0^y \int_0^{4-y^2} xy\,dz\,dx\,dy$.

(b) $$\int_0^2 \int_0^y \int_0^{4-y^2} xy\,dz\,dx\,dy = \int_0^2 \int_0^y xy\,[z]_{z=0}^{z=4-y^2}\,dx\,dy = \int_0^2 \int_0^y xy(4 - y^2)\,dx\,dy = \int_0^2 \int_0^y x(4y - y^3)\,dx\,dy$$
$$= \int_0^2 (4y - y^3) \left[\tfrac{x^2}{2} \right]_{x=0}^{x=y} dy = \tfrac{1}{2} \int_0^2 (4y^3 - y^5)\,dy = \tfrac{1}{2} \left[y^4 - \tfrac{y^6}{6} \right]_0^2 = \tfrac{8}{3}$$

11. (a) The solid region E can be described as $E = \{(x, y, z) \mid 0 \le x \le 2, 0 \le y \le x^2, 0 \le z \le 2 - x\}$.

Thus, $\iiint_E (x + y)\,dV = \int_0^2 \int_0^{2-x} \int_0^{x^2} (x + y)\,dy\,dz\,dx$.

(b) $$\int_0^2 \int_0^{2-x} \int_0^{x^2} (x + y)\,dy\,dz\,dx = \int_0^2 \int_0^{2-x} \left[xy + \frac{y^2}{2} \right]_{y=0}^{y=x^2} dz\,dx = \int_0^2 \int_0^{2-x} \left(x^3 + \frac{x^4}{2} \right) dz\,dx$$
$$= \int_0^2 \left(x^3 + \frac{x^4}{2} \right) \Big[z \Big]_{z=0}^{z=2-x} dx = \int_0^2 \left(2x^3 - \frac{x^5}{2} \right) dx = \left[\frac{x^4}{2} - \frac{x^6}{12} \right]_0^2 = \frac{8}{3}$$

12. (a) The solid region E can be described as $E = \{(x, y, z) \mid z - 4 \le x \le 4 - z, -2 \le y \le 2, 0 \le z \le 4 - y^2\}$.

Thus, $\iiint_E 2\,dV = \int_{-2}^{2} \int_0^{4-y^2} \int_{z-4}^{4-z} 2\,dx\,dz\,dy$.

(b) $$\int_{-2}^{2} \int_0^{4-y^2} \int_{z-4}^{4-z} 2\,dx\,dz\,dy = \int_{-2}^{2} \int_0^{4-y^2} 2\Big[x \Big]_{x=z-4}^{x=4-z} dz\,dy = 2\int_{-2}^{2} \int_0^{4-y^2} (8 - 2z)\,dz\,dy$$
$$= 2\int_{-2}^{2} \Big[8z - z^2 \Big]_{z=0}^{z=4-y^2} dy = 2\int_{-2}^{2} (16 - y^4)\,dy = 2\Big[16y - \tfrac{y^5}{5} \Big]_{-2}^{2} = \tfrac{512}{5}$$

13. $$\iiint_E y\,dV = \int_0^3 \int_0^x \int_{x-y}^{x+y} y\,dz\,dy\,dx = \int_0^3 \int_0^x \big[yz \big]_{z=x-y}^{z=x+y} dy\,dx = \int_0^3 \int_0^x 2y^2\,dy\,dx$$
$$= \int_0^3 \big[\tfrac{2}{3}y^3 \big]_{y=0}^{y=x} dx = \int_0^3 \tfrac{2}{3}x^3\,dx = \tfrac{1}{6}x^4 \big]_0^3 = \tfrac{81}{6} = \tfrac{27}{2}$$

14. $$\iiint_E e^{z/y}\,dV = \int_0^1 \int_y^1 \int_0^{xy} e^{z/y}\,dz\,dx\,dy = \int_0^1 \int_y^1 \Big[ye^{z/y} \Big]_{z=0}^{z=xy} dx\,dy$$
$$= \int_0^1 \int_y^1 (ye^x - y)\,dx\,dy = \int_0^1 \big[ye^x - xy \big]_{x=y}^{x=1} dy = \int_0^1 \left(ey - y - ye^y + y^2 \right) dy$$
$$= \big[\tfrac{1}{2}ey^2 - \tfrac{1}{2}y^2 - (y - 1)e^y + \tfrac{1}{3}y^3 \big]_0^1 \qquad \text{[integrate by parts]}$$
$$= \tfrac{1}{2}e - \tfrac{1}{2} + \tfrac{1}{3} - 1 = \tfrac{1}{2}e - \tfrac{7}{6}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $$\iiint_E \frac{1}{x^3}\,dV = \int_0^1 \int_0^{y^2} \int_1^{z+1} \frac{1}{x^3}\,dx\,dz\,dy = \int_0^1 \int_0^{y^2} -\frac{1}{2}\left[\frac{1}{x^2}\right]_{x=1}^{x=z+1} dz\,dy$$
$$= -\frac{1}{2}\int_0^1 \int_0^{y^2} \left(\frac{1}{(z+1)^2} - 1\right) dz\,dy = -\frac{1}{2}\int_0^1 \left[-\frac{1}{z+1} - z\right]_{z=0}^{z=y^2} dy$$
$$= \frac{1}{2}\int_0^1 \left(\frac{1}{y^2+1} + y^2 - 1\right) dy = \frac{1}{2}\left[\tan^{-1} y + \frac{y^3}{3} - y\right]_0^1 = \frac{1}{2}\left[\left(\frac{\pi}{4} + \frac{1}{3} - 1\right) - 0\right]$$
$$= \frac{\pi}{8} - \frac{1}{3}$$

16. Here $E = \{(x, y, z) \mid 0 \le x \le \pi, 0 \le y \le \pi - x, 0 \le z \le x\}$, so

$$\iiint_E \sin y\,dV = \int_0^\pi \int_0^{\pi-x} \int_0^x \sin y\,dz\,dy\,dx = \int_0^\pi \int_0^{\pi-x} \left[z \sin y\right]_{z=0}^{z=x} dy\,dx = \int_0^\pi \int_0^{\pi-x} x \sin y\,dy\,dx$$
$$= \int_0^\pi \left[-x\cos y\right]_{y=0}^{y=\pi-x} dx = \int_0^\pi \left[-x\cos(\pi - x) + x\right] dx$$
$$= \left[x\sin(\pi - x) - \cos(\pi - x) + \tfrac{1}{2}x^2\right]_0^\pi \qquad \text{[integrate by parts]}$$
$$= 0 - 1 + \tfrac{1}{2}\pi^2 - 0 - 1 - 0 = \tfrac{1}{2}\pi^2 - 2$$

17. Here $E = \{(x, y, z) \mid 0 \le x \le 1, 0 \le y \le \sqrt{x}, 0 \le z \le 1 + x + y\}$, so

$$\iiint_E 6xy\,dV = \int_0^1 \int_0^{\sqrt{x}} \int_0^{1+x+y} 6xy\,dz\,dy\,dx = \int_0^1 \int_0^{\sqrt{x}} \left[6xyz\right]_{z=0}^{z=1+x+y} dy\,dx$$
$$= \int_0^1 \int_0^{\sqrt{x}} 6xy(1 + x + y)\,dy\,dx = \int_0^1 \left[3xy^2 + 3x^2y^2 + 2xy^3\right]_{y=0}^{y=\sqrt{x}} dx$$
$$= \int_0^1 (3x^2 + 3x^3 + 2x^{5/2})\,dx = \left[x^3 + \tfrac{3}{4}x^4 + \tfrac{4}{7}x^{7/2}\right]_0^1 = \tfrac{65}{28}$$

18.

Here $E = \left\{(x, y, z) \mid -1 \le x \le 1,\ 0 \le y \le 2,\ x^2 - 1 \le z \le 1 - x^2\right\}$. Thus,

$$\iiint_E (x - y)\,dV = \int_{-1}^1 \int_0^2 \int_{x^2-1}^{1-x^2} (x - y)\,dz\,dy\,dx$$
$$= \int_{-1}^1 \int_0^2 (x - y)(1 - x^2 - (x^2 - 1))\,dy\,dx$$
$$= \int_{-1}^1 \int_0^2 (2x - 2x^3 - 2y + 2x^2y)\,dy\,dx$$
$$= \int_{-1}^1 \left[2xy - 2x^3y - y^2 + x^2y^2\right]_{y=0}^{y=2} dx$$
$$= \int_{-1}^1 (4x - 4x^3 - 4 + 4x^2)\,dx$$
$$= \left[2x^2 - x^4 - 4x + \tfrac{4}{3}x^3\right]_{-1}^1 = -\tfrac{5}{3} - \tfrac{11}{3} = -\tfrac{16}{3}$$

19.

Here $T = \{(x, y, z) \mid 0 \le x \le 2,\ 0 \le y \le 2 - x,\ 0 \le z \le 2 - x - y\}$. Thus,

$$\iiint_T y^2\,dV = \int_0^2 \int_0^{2-x} \int_0^{2-x-y} y^2\,dz\,dy\,dx = \int_0^2 \int_0^{2-x} y^2(2 - x - y)\,dy\,dx$$
$$= \int_0^2 \int_0^{2-x} \left[(2 - x)y^2 - y^3\right] dy\,dx$$
$$= \int_0^2 \left[(2 - x)\left(\tfrac{1}{3}y^3\right) - \tfrac{1}{4}y^4\right]_{y=0}^{y=2-x} dx$$
$$= \int_0^2 \left[\tfrac{1}{3}(2 - x)^4 - \tfrac{1}{4}(2 - x)^4\right] dx = \int_0^2 \tfrac{1}{12}(2 - x)^4\,dx$$
$$= \left[\tfrac{1}{12}\left(-\tfrac{1}{5}\right)(2 - x)^5\right]_0^2 = -\tfrac{1}{60}(0 - 32) = \tfrac{8}{15}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20.

The projection of T onto the xz-plane is the triangle bounded by the lines $z = x$, $x = 0$, and $z = 1$. Then

$T = \{(x, y, z) \mid 0 \le x \le 1,\ x \le z \le 1,\ 0 \le y \le z - x\}$, and

$$\iiint_T xz\,dV = \int_0^1 \int_x^1 \int_0^{z-x} xz\,dy\,dz\,dx = \int_0^1 \int_x^1 xz(z-x)\,dz\,dx$$
$$= \int_0^1 \int_x^1 (xz^2 - x^2 z)\,dz\,dx = \int_0^1 \left[\tfrac{1}{3}xz^3 - \tfrac{1}{2}x^2z^2\right]_{z=x}^{z=1} dx$$
$$= \int_0^1 \left(\tfrac{1}{3}x - \tfrac{1}{2}x^2 - \tfrac{1}{3}x^4 + \tfrac{1}{2}x^4\right) dx$$
$$= \left[\tfrac{1}{6}x^2 - \tfrac{1}{6}x^3 + \tfrac{1}{30}x^5\right]_0^1 = \tfrac{1}{6} - \tfrac{1}{6} + \tfrac{1}{30} = \tfrac{1}{30}$$

21.

The projection of E onto the yz-plane is the disk $y^2 + z^2 \le 1$. Using polar coordinates $y = r\cos\theta$ and $z = r\sin\theta$, we get

$$\iiint_E x\,dV = \iint_D \left[\int_{4y^2+4z^2}^4 x\,dx\right] dA = \tfrac{1}{2}\iint_D \left[4^2 - (4y^2+4z^2)^2\right] dA$$
$$= 8\int_0^{2\pi}\int_0^1 (1 - r^4)\,r\,dr\,d\theta = 8\int_0^{2\pi} d\theta \int_0^1 (r - r^5)\,dr$$
$$= 8(2\pi)\left[\tfrac{1}{2}r^2 - \tfrac{1}{6}r^6\right]_0^1 = \tfrac{16\pi}{3}$$

22.

$$\int_0^1 \int_{3x}^3 \int_0^{\sqrt{9-y^2}} z\,dz\,dy\,dx = \int_0^1 \int_{3x}^3 \tfrac{1}{2}(9 - y^2)\,dy\,dx$$
$$= \int_0^1 \left[\tfrac{9}{2}y - \tfrac{1}{6}y^3\right]_{y=3x}^{y=3} dx$$
$$= \int_0^1 \left[9 - \tfrac{27}{2}x + \tfrac{9}{2}x^3\right] dx$$
$$= \left[9x - \tfrac{27}{4}x^2 + \tfrac{9}{8}x^4\right]_0^1 = \tfrac{27}{8}$$

23. The plane $2x + y + z = 4$ intersects the xy-plane when

$2x + y + 0 = 4 \quad \Rightarrow \quad y = 4 - 2x$, so

$E = \{(x, y, z) \mid 0 \le x \le 2, 0 \le y \le 4 - 2x, 0 \le z \le 4 - 2x - y\}$ and

$$V = \int_0^2 \int_0^{4-2x} \int_0^{4-2x-y} dz\,dy\,dx = \int_0^2 \int_0^{4-2x} (4 - 2x - y)\,dy\,dx$$
$$= \int_0^2 \left[4y - 2xy - \tfrac{1}{2}y^2\right]_{y=0}^{y=4-2x} dx$$
$$= \int_0^2 \left[4(4-2x) - 2x(4-2x) - \tfrac{1}{2}(4-2x)^2\right] dx$$
$$= \int_0^2 (2x^2 - 8x + 8)\,dx = \left[\tfrac{2}{3}x^3 - 4x^2 + 8x\right]_0^2 = \tfrac{16}{3}$$

24. The paraboloids intersect when $x^2 + z^2 = 8 - x^2 - z^2 \quad \Leftrightarrow \quad x^2 + z^2 = 4$, thus the intersection is the circle $x^2 + z^2 = 4$, $y = 4$. The projection of E onto the xz-plane is the disk $x^2 + z^2 \le 4$, so $E = \left\{(x, y, z) \mid x^2 + z^2 \le y \le 8 - x^2 - z^2, x^2 + z^2 \le 4\right\}$. Let $D = \left\{(x, z) \mid x^2 + z^2 \le 4\right\}$. Then

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

using polar coordinates $x = r\cos\theta$ and $z = r\sin\theta$, we have

$$V = \iiint_E dV = \iint_D \left(\int_{x^2+z^2}^{8-x^2-z^2} dy\right) dA = \iint_D (8 - 2x^2 - 2z^2)\, dA$$

$$= \int_0^{2\pi}\int_0^2 (8 - 2r^2)\, r\, dr\, d\theta = \int_0^{2\pi} d\theta \int_0^2 (8r - 2r^3)\, dr$$

$$= \left[\theta\right]_0^{2\pi} \left[4r^2 - \tfrac{1}{2}r^4\right]_0^2 = 2\pi(16 - 8) = 16\pi$$

25. The plane $y + z = 1$ intersects the xy-plane in the line $y = 1$, so

$E = \{(x, y, z) \mid -1 \le x \le 1, x^2 \le y \le 1, 0 \le z \le 1 - y\}$ and

$$V = \iiint_E dV = \int_{-1}^1 \int_{x^2}^1 \int_0^{1-y} dz\, dy\, dx = \int_{-1}^1 \int_{x^2}^1 (1 - y)\, dy\, dx$$

$$= \int_{-1}^1 \left[y - \tfrac{1}{2}y^2\right]_{y=x^2}^{y=1} dx = \int_{-1}^1 \left(\tfrac{1}{2} - x^2 + \tfrac{1}{2}x^4\right) dx$$

$$= \left[\tfrac{1}{2}x - \tfrac{1}{3}x^3 + \tfrac{1}{10}x^5\right]_{-1}^1 = \tfrac{1}{2} - \tfrac{1}{3} + \tfrac{1}{10} + \tfrac{1}{2} - \tfrac{1}{3} + \tfrac{1}{10} = \tfrac{8}{15}$$

26. Here $E = \{(x, y, z) \mid -1 \le y \le 4 - z, x^2 + z^2 \le 4\}$, so

$$V = \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{-1}^{4-z} dy\, dz\, dx = \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} (4 - z + 1)\, dz\, dx$$

$$= \int_{-2}^2 \left[5z - \tfrac{1}{2}z^2\right]_{z=-\sqrt{4-x^2}}^{z=\sqrt{4-x^2}} dx = \int_{-2}^2 10\sqrt{4 - x^2}\, dx$$

$$= 10\left[\tfrac{x}{2}\sqrt{4 - x^2} + 2\sin^{-1}\left(\tfrac{x}{2}\right)\right]_{-2}^2 \qquad \left[\begin{array}{l}\text{using trigonometric substitution or}\\ \text{Formula 30 in the Table of Integrals}\end{array}\right]$$

$$= 10\left[2\sin^{-1}(1) - 2\sin^{-1}(-1)\right] = 20\left(\tfrac{\pi}{2} - \left(-\tfrac{\pi}{2}\right)\right) = 20\pi$$

Alternatively, use polar coordinates to evaluate the double integral:

$$\int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} (5 - z)\, dz\, dx = \int_0^{2\pi} \int_0^2 (5 - r\sin\theta)\, r\, dr\, d\theta$$

$$= \int_0^{2\pi} \left[\tfrac{5}{2}r^2 - \tfrac{1}{3}r^3\sin\theta\right]_{r=0}^{r=2} d\theta = \int_0^{2\pi} \left(10 - \tfrac{8}{3}\sin\theta\right) d\theta$$

$$= 10\theta + \tfrac{8}{3}\cos\theta\Big]_0^{2\pi} = 20\pi$$

27. (a) The wedge can be described as the region

$$D = \{(x, y, z) \mid y^2 + z^2 \le 1, 0 \le x \le 1, 0 \le y \le x\}$$
$$= \left\{(x, y, z) \mid 0 \le x \le 1, 0 \le y \le x, 0 \le z \le \sqrt{1 - y^2}\right\}$$

So the integral expressing the volume of the wedge is

$$\iiint_D dV = \int_0^1 \int_0^x \int_0^{\sqrt{1-y^2}} dz\, dy\, dx.$$

(b) A CAS gives $\int_0^1 \int_0^x \int_0^{\sqrt{1-y^2}} dz\, dy\, dx = \frac{\pi}{4} - \frac{1}{3}$.

(Or use Formulas 30 and 87 from the Table of Integrals.)

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. Divide B into 8 cubes of size $\Delta V = 8$. With $f(x, y, z) = \sqrt{x^2 + y^2 + z^2}$, the Midpoint Rule gives

$$\begin{aligned}\iiint_B \sqrt{x^2+y^2+z^2}\,dV &\approx \sum_{i=1}^{2}\sum_{j=1}^{2}\sum_{k=1}^{2} f\left(\overline{x}_i, \overline{y}_j, \overline{z}_k\right)\Delta V \\ &= 8[f(1,1,1) + f(1,1,3) + f(1,3,1) + f(1,3,3) + f(3,1,1) \\ &\qquad + f(3,1,3) + f(3,3,1) + f(3,3,3)] \\ &\approx 239.64\end{aligned}$$

29. Here $f(x, y, z) = \cos(xyz)$ and $\Delta V = \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{8}$, so the Midpoint Rule gives

$$\begin{aligned}\iiint_B f(x,y,z)\,dV &\approx \sum_{i=1}^{l}\sum_{j=1}^{m}\sum_{k=1}^{n} f\left(\overline{x}_i, \overline{y}_j, \overline{z}_k\right)\Delta V \\ &= \tfrac{1}{8}\left[f\left(\tfrac{1}{4},\tfrac{1}{4},\tfrac{1}{4}\right) + f\left(\tfrac{1}{4},\tfrac{1}{4},\tfrac{3}{4}\right) + f\left(\tfrac{1}{4},\tfrac{3}{4},\tfrac{1}{4}\right) + f\left(\tfrac{1}{4},\tfrac{3}{4},\tfrac{3}{4}\right)\right. \\ &\qquad \left. + f\left(\tfrac{3}{4},\tfrac{1}{4},\tfrac{1}{4}\right) + f\left(\tfrac{3}{4},\tfrac{1}{4},\tfrac{3}{4}\right) + f\left(\tfrac{3}{4},\tfrac{3}{4},\tfrac{1}{4}\right) + f\left(\tfrac{3}{4},\tfrac{3}{4},\tfrac{3}{4}\right)\right] \\ &= \tfrac{1}{8}\left[\cos\tfrac{1}{64} + \cos\tfrac{3}{64} + \cos\tfrac{3}{64} + \cos\tfrac{9}{64} + \cos\tfrac{3}{64} + \cos\tfrac{9}{64} + \cos\tfrac{9}{64} + \cos\tfrac{27}{64}\right] \approx 0.985\end{aligned}$$

30. Here $f(x, y, z) = \sqrt{x}\, e^{xyz}$ and $\Delta V = 2 \cdot \frac{1}{2} \cdot 1 = 1$, so the Midpoint Rule gives

$$\begin{aligned}\iiint_B f(x,y,z)\,dV &\approx \sum_{i=1}^{l}\sum_{j=1}^{m}\sum_{k=1}^{n} f\left(\overline{x}_i, \overline{y}_j, \overline{z}_k\right)\Delta V \\ &= 1\left[f\left(1,\tfrac{1}{4},\tfrac{1}{2}\right) + f\left(1,\tfrac{1}{4},\tfrac{3}{2}\right) + f\left(1,\tfrac{3}{4},\tfrac{1}{2}\right) + f\left(1,\tfrac{3}{4},\tfrac{3}{2}\right)\right. \\ &\qquad \left. + f\left(3,\tfrac{1}{4},\tfrac{1}{2}\right) + f\left(3,\tfrac{1}{4},\tfrac{3}{2}\right) + f\left(3,\tfrac{3}{4},\tfrac{1}{2}\right) + f\left(3,\tfrac{3}{4},\tfrac{3}{2}\right)\right] \\ &= e^{1/8} + e^{3/8} + e^{3/8} + e^{9/8} + \sqrt{3}e^{3/8} + \sqrt{3}e^{9/8} + \sqrt{3}e^{9/8} + \sqrt{3}e^{27/8} \approx 70.932\end{aligned}$$

31. $E = \{(x, y, z) \mid 0 \le x \le 1, 0 \le z \le 1 - x, 0 \le y \le 2 - 2z\}$, the solid bounded by the three coordinate planes and the planes $z = 1 - x$, $y = 2 - 2z$.

32. $E = \left\{(x, y, z) \mid 0 \le y \le 2, 0 \le z \le 2 - y, 0 \le x \le 4 - y^2\right\}$, the solid bounded by the three coordinate planes, the plane $z = 2 - y$, and the cylindrical surface $x = 4 - y^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33.

If D_1, D_2, D_3 are the projections of E on the xy-, yz-, and xz-planes, then

$$D_1 = \left\{(x, y) \mid -2 \le x \le 2, 0 \le y \le 4 - x^2\right\} = \left\{(x, y) \mid 0 \le y \le 4,\ -\sqrt{4-y} \le x \le \sqrt{4-y}\,\right\}$$

$$D_2 = \left\{(y, z) \mid 0 \le y \le 4,\ -\tfrac{1}{2}\sqrt{4-y} \le z \le \tfrac{1}{2}\sqrt{4-y}\right\} = \left\{(y, z) \mid -1 \le z \le 1, 0 \le y \le 4 - 4z^2\right\}$$

$$D_3 = \left\{(x, z) \mid x^2 + 4z^2 \le 4\right\}$$

Therefore

$$\begin{aligned}
E &= \left\{(x, y, z) \mid -2 \le x \le 2, 0 \le y \le 4 - x^2,\ -\tfrac{1}{2}\sqrt{4 - x^2 - y} \le z \le \tfrac{1}{2}\sqrt{4 - x^2 - y}\,\right\} \\
&= \left\{(x, y, z) \mid 0 \le y \le 4,\ -\sqrt{4-y} \le x \le \sqrt{4-y},\ -\tfrac{1}{2}\sqrt{4 - x^2 - y} \le z \le \tfrac{1}{2}\sqrt{4 - x^2 - y}\,\right\} \\
&= \left\{(x, y, z) \mid -1 \le z \le 1, 0 \le y \le 4 - 4z^2,\ -\sqrt{4 - y - 4z^2} \le x \le \sqrt{4 - y - 4z^2}\,\right\} \\
&= \left\{(x, y, z) \mid 0 \le y \le 4,\ -\tfrac{1}{2}\sqrt{4-y} \le z \le \tfrac{1}{2}\sqrt{4-y},\ -\sqrt{4 - y - 4z^2} \le x \le \sqrt{4 - y - 4z^2}\,\right\} \\
&= \left\{(x, y, z) \mid -2 \le x \le 2,\ -\tfrac{1}{2}\sqrt{4 - x^2} \le z \le \tfrac{1}{2}\sqrt{4 - x^2}, 0 \le y \le 4 - x^2 - 4z^2\right\} \\
&= \left\{(x, y, z) \mid -1 \le z \le 1,\ -\sqrt{4 - 4z^2} \le x \le \sqrt{4 - 4z^2}, 0 \le y \le 4 - x^2 - 4z^2\right\}
\end{aligned}$$

Then

$$\begin{aligned}
\iiint_E f(x, y, z)\, dV &= \int_{-2}^{2} \int_0^{4-x^2} \int_{-\sqrt{4-x^2-y}/2}^{\sqrt{4-x^2-y}/2} f(x, y, z)\, dz\, dy\, dx \\
&= \int_0^4 \int_{-\sqrt{4-y}}^{\sqrt{4-y}} \int_{-\sqrt{4-x^2-y}/2}^{\sqrt{4-x^2-y}/2} f(x, y, z)\, dz\, dx\, dy = \int_{-1}^{1} \int_0^{4-4z^2} \int_{-\sqrt{4-y-4z^2}}^{\sqrt{4-y-4z^2}} f(x, y, z)\, dx\, dy\, dz \\
&= \int_0^4 \int_{-\sqrt{4-y}/2}^{\sqrt{4-y}/2} \int_{-\sqrt{4-y-4z^2}}^{\sqrt{4-y-4z^2}} f(x, y, z)\, dx\, dz\, dy = \int_{-2}^{2} \int_{-\sqrt{4-x^2}/2}^{\sqrt{4-x^2}/2} \int_0^{4-x^2-4z^2} f(x, y, z)\, dy\, dz\, dx \\
&= \int_{-1}^{1} \int_{-\sqrt{4-4z^2}}^{\sqrt{4-4z^2}} \int_0^{4-x^2-4z^2} f(x, y, z)\, dy\, dx\, dz
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34.

If D_1, D_2, D_3 are the projections of E on the xy-, yz-, and xz-planes, then

$$D_1 = \{(x, y) \mid -2 \le x \le 2,\ -3 \le y \le 3\}$$

$$D_2 = \left\{(y, z) \mid y^2 + z^2 \le 9\right\}$$

$$D_3 = \{(x, z) \mid -2 \le x \le 2,\ -3 \le z \le 3\}$$

Therefore

$$\begin{aligned} E &= \left\{(x, y, z) \mid -2 \le x \le 2,\ -3 \le y \le 3,\ -\sqrt{9-y^2} \le z \le \sqrt{9-y^2}\right\} \\ &= \left\{(x, y, z) \mid -3 \le y \le 3,\ -\sqrt{9-y^2} \le z \le \sqrt{9-y^2},\ -2 \le x \le 2\right\} \\ &= \left\{(x, y, z) \mid -3 \le z \le 3,\ -\sqrt{9-z^2} \le y \le \sqrt{9-z^2},\ -2 \le x \le 2\right\} \\ &= \left\{(x, y, z) \mid -2 \le x \le 2,\ -3 \le z \le 3,\ -\sqrt{9-z^2} \le y \le \sqrt{9-z^2}\right\} \end{aligned}$$

and

$$\begin{aligned} \iiint_E f(x, y, z)\, dV &= \int_{-2}^{2} \int_{-3}^{3} \int_{-\sqrt{9-y^2}}^{\sqrt{9-y^2}} f(x, y, z)\, dz\, dy\, dx = \int_{-3}^{3} \int_{-2}^{2} \int_{-\sqrt{9-y^2}}^{\sqrt{9-y^2}} f(x, y, z)\, dz\, dx\, dy \\ &= \int_{-3}^{3} \int_{-\sqrt{9-y^2}}^{\sqrt{9-y^2}} \int_{-2}^{2} f(x, y, z)\, dx\, dz\, dy = \int_{-3}^{3} \int_{-\sqrt{9-z^2}}^{\sqrt{9-z^2}} \int_{-2}^{2} f(x, y, z)\, dx\, dy\, dz \\ &= \int_{-2}^{2} \int_{-3}^{3} \int_{-\sqrt{9-z^2}}^{\sqrt{9-z^2}} f(x, y, z)\, dy\, dz\, dx = \int_{-3}^{3} \int_{-2}^{2} \int_{-\sqrt{9-z^2}}^{\sqrt{9-z^2}} f(x, y, z)\, dy\, dx\, dz \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35.

If D_1, D_2, and D_3 are the projections of E on the xy-, yz-, and xz-planes, then

$$D_1 = \left\{(x,y) \mid -2 \le x \le 2, x^2 \le y \le 4\right\} = \left\{(x,y) \mid 0 \le y \le 4, -\sqrt{y} \le x \le \sqrt{y}\right\},$$

$$D_2 = \left\{(y,z) \mid 0 \le y \le 4, 0 \le z \le 2 - \tfrac{1}{2}y\right\} = \left\{(y,z) \mid 0 \le z \le 2, 0 \le y \le 4 - 2z\right\}, \text{ and}$$

$$D_3 = \left\{(x,z) \mid -2 \le x \le 2, 0 \le z \le 2 - \tfrac{1}{2}x^2\right\} = \left\{(x,z) \mid 0 \le z \le 2, -\sqrt{4-2z} \le x \le \sqrt{4-2z}\right\}$$

Therefore

$$\begin{aligned}
E &= \left\{(x,y,z) \mid -2 \le x \le 2, x^2 \le y \le 4, 0 \le z \le 2 - \tfrac{1}{2}y\right\} \\
&= \left\{(x,y,z) \mid 0 \le y \le 4,\ -\sqrt{y} \le x \le \sqrt{y}, 0 \le z \le 2 - \tfrac{1}{2}y\right\} \\
&= \left\{(x,y,z) \mid 0 \le y \le 4, 0 \le z \le 2 - \tfrac{1}{2}y,\ -\sqrt{y} \le x \le \sqrt{y}\right\} \\
&= \left\{(x,y,z) \mid 0 \le z \le 2, 0 \le y \le 4 - 2z,\ -\sqrt{y} \le x \le \sqrt{y}\right\} \\
&= \left\{(x,y,z) \mid -2 \le x \le 2, 0 \le z \le 2 - \tfrac{1}{2}x^2, x^2 \le y \le 4 - 2z\right\} \\
&= \left\{(x,y,z) \mid 0 \le z \le 2,\ -\sqrt{4-2z} \le x \le \sqrt{4-2z}, x^2 \le y \le 4 - 2z\right\}
\end{aligned}$$

Then

$$\begin{aligned}
\iiint_E f(x,y,z)\,dV &= \int_{-2}^{2}\int_{x^2}^{4}\int_0^{2-y/2} f(x,y,z)\,dz\,dy\,dx = \int_0^4\int_{-\sqrt{y}}^{\sqrt{y}}\int_0^{2-y/2} f(x,y,z)\,dz\,dx\,dy \\
&= \int_0^4\int_0^{2-y/2}\int_{-\sqrt{y}}^{\sqrt{y}} f(x,y,z)\,dx\,dz\,dy = \int_0^2\int_0^{4-2z}\int_{-\sqrt{y}}^{\sqrt{y}} f(x,y,z)\,dx\,dy\,dz \\
&= \int_{-2}^{2}\int_0^{2-x^2/2}\int_{x^2}^{4-2z} f(x,y,z)\,dy\,dz\,dx = \int_0^2\int_{-\sqrt{4-2z}}^{\sqrt{4-2z}}\int_{x^2}^{4-2z} f(x,y,z)\,dy\,dx\,dz
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36.

If D_1, D_2, and D_3 are the projections of E on the xy-, yz-, and xz-planes, then

$$D_1 = \{(x, y) \mid 0 \le x \le 2, 2 - x \le y \le 2\} = \{(x, y) \mid 0 \le y \le 2, 2 - y \le x \le 2\},$$

$$D_2 = \left\{(y, z) \mid 0 \le y \le 2, 0 \le z \le \tfrac{1}{2}y\right\} = \{(y, z) \mid 0 \le z \le 1, 2z \le y \le 2\}, \text{ and}$$

$$D_3 = \left\{(x, z) \mid 0 \le x \le 2, 0 \le z \le \tfrac{1}{2}x\right\} = \{(x, z) \mid 0 \le z \le 1, 2z \le x \le 2\}$$

Therefore

$$\begin{aligned}
E &= \left\{(x, y, z) \mid 0 \le x \le 2, 2 - x \le y \le 2, 0 \le z \le \tfrac{1}{2}(x + y - 2)\right\} \\
&= \left\{(x, y, z) \mid 0 \le y \le 2, 2 - y \le x \le 2, 0 \le z \le \tfrac{1}{2}(x + y - 2)\right\} \\
&= \left\{(x, y, z) \mid 0 \le y \le 2, 0 \le z \le \tfrac{1}{2}y, 2 - y + 2z \le x \le 2\right\} \\
&= \{(x, y, z) \mid 0 \le z \le 1, 2z \le y \le 2, 2 - y + 2z \le x \le 2\} \\
&= \left\{(x, y, z) \mid 0 \le x \le 2, 0 \le z \le \tfrac{1}{2}x, 2 - x + 2z \le y \le 2\right\} \\
&= \{(x, y, z) \mid 0 \le z \le 1, 2z \le x \le 2, 2 - x + 2z \le y \le 2\}
\end{aligned}$$

Then

$$\begin{aligned}
\iiint_E f(x, y, z)\, dV &= \int_0^2 \int_{2-x}^2 \int_0^{(x+y-2)/2} f(x, y, z)\, dz\, dy\, dx \\
&= \int_0^2 \int_{2-y}^2 \int_0^{(x+y-2)/2} f(x, y, z)\, dz\, dx\, dy \\
&= \int_0^2 \int_0^{y/2} \int_{2-y+2z}^2 f(x, y, z)\, dx\, dz\, dy \\
&= \int_0^1 \int_{2z}^2 \int_{2-y+2z}^2 f(x, y, z)\, dx\, dy\, dz \\
&= \int_0^2 \int_0^{x/2} \int_{2-x+2z}^2 f(x, y, z)\, dy\, dz\, dx \\
&= \int_0^1 \int_{2z}^2 \int_{2-x+2z}^2 f(x, y, z)\, dy\, dx\, dz
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37.

The diagrams show the projections of E onto the xy-, yz-, and xz-planes. Therefore

$$\int_0^1\int_{\sqrt{x}}^1\int_0^{1-y} f(x,y,z)\,dz\,dy\,dx = \int_0^1\int_0^{y^2}\int_0^{1-y} f(x,y,z)\,dz\,dx\,dy = \int_0^1\int_0^{1-z}\int_0^{y^2} f(x,y,z)\,dx\,dy\,dz$$
$$= \int_0^1\int_0^{1-y}\int_0^{y^2} f(x,y,z)\,dx\,dz\,dy = \int_0^1\int_0^{1-\sqrt{x}}\int_{\sqrt{x}}^{1-z} f(x,y,z)\,dy\,dz\,dx$$
$$= \int_0^1\int_0^{(1-z)^2}\int_{\sqrt{x}}^{1-z} f(x,y,z)\,dy\,dx\,dz$$

38.

The projections of E onto the xy- and xz-planes are as in the first two diagrams and so

$$\int_0^1\int_0^{1-x^2}\int_0^{1-x} f(x,y,z)\,dy\,dz\,dx = \int_0^1\int_0^{\sqrt{1-z}}\int_0^{1-x} f(x,y,z)\,dy\,dx\,dz$$
$$= \int_0^1\int_0^{1-y}\int_0^{1-x^2} f(x,y,z)\,dz\,dx\,dy = \int_0^1\int_0^{1-x}\int_0^{1-x^2} f(x,y,z)\,dz\,dy\,dx$$

Now the surface $z = 1 - x^2$ intersects the plane $y = 1 - x$ in a curve whose projection in the yz-plane is $z = 1 - (1-y)^2$ or $z = 2y - y^2$. So we must split up the projection of E on the yz-plane into two regions as in the third diagram. For (y, z) in R_1, $0 \le x \le 1 - y$ and for (y, z) in R_2, $0 \le x \le \sqrt{1-z}$, and so the given integral is also equal to

$$\int_0^1\int_0^{1-\sqrt{1-z}}\int_0^{\sqrt{1-z}} f(x,y,z)\,dx\,dy\,dz + \int_0^1\int_{1-\sqrt{1-z}}^1\int_0^{1-y} f(x,y,z)\,dx\,dy\,dz$$
$$= \int_0^1\int_0^{2y-y^2}\int_0^{1-y} f(x,y,z)\,dx\,dz\,dy + \int_0^1\int_{2y-y^2}^1\int_0^{\sqrt{1-z}} f(x,y,z)\,dx\,dz\,dy.$$

39.

$\int_0^1\int_y^1\int_0^y f(x,y,z)\,dz\,dx\,dy = \iiint_E f(x,y,z)\,dV$ where $E = \{(x,y,z) \mid 0 \le z \le y,\ y \le x \le 1,\ 0 \le y \le 1\}$.

If D_1, D_2, and D_3 are the projections of E onto the xy-, yz- and xz-planes then

$$D_1 = \{(x,y) \mid 0 \le y \le 1,\ y \le x \le 1\} = \{(x,y) \mid 0 \le x \le 1,\ 0 \le y \le x\},$$

$$D_2 = \{(y,z) \mid 0 \le y \le 1,\ 0 \le z \le y\} = \{(y,z) \mid 0 \le z \le 1,\ z \le y \le 1\}, \text{ and}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$D_3 = \{(x, z) \mid 0 \le x \le 1, 0 \le z \le x\} = \{(x, z) \mid 0 \le z \le 1, z \le x \le 1\}.$$

Thus we also have

$$\begin{aligned} E &= \{(x, y, z) \mid 0 \le x \le 1, 0 \le y \le x, 0 \le z \le y\} = \{(x, y, z) \mid 0 \le y \le 1, 0 \le z \le y, y \le x \le 1\} \\ &= \{(x, y, z) \mid 0 \le z \le 1, z \le y \le 1, y \le x \le 1\} = \{(x, y, z) \mid 0 \le x \le 1, 0 \le z \le x, z \le y \le x\} \\ &= \{(x, y, z) \mid 0 \le z \le 1, z \le x \le 1, z \le y \le x\}. \end{aligned}$$

Then

$$\begin{aligned} \int_0^1 \int_y^1 \int_0^y f(x, y, z)\, dz\, dx\, dy &= \int_0^1 \int_0^x \int_0^y f(x, y, z)\, dz\, dy\, dx = \int_0^1 \int_0^y \int_y^1 f(x, y, z)\, dx\, dz\, dy \\ &= \int_0^1 \int_z^1 \int_y^1 f(x, y, z)\, dx\, dy\, dz = \int_0^1 \int_0^x \int_z^x f(x, y, z)\, dy\, dz\, dx \\ &= \int_0^1 \int_z^1 \int_z^x f(x, y, z)\, dy\, dx\, dz \end{aligned}$$

40.

$\int_0^1 \int_y^1 \int_0^z f(x, y, z)\, dx\, dz\, dy = \iiint_E f(x, y, z)\, dV$ where $E = \{(x, y, z) \mid 0 \le x \le z, y \le z \le 1, 0 \le y \le 1\}$.

Notice that E is bounded below by two different surfaces, so we must split the projection of E onto the xy-plane into two regions as in the second diagram. If D_1, D_2, and D_3 are the projections of E on the xy-, yz- and xz-planes then

$$\begin{aligned} D_1 = R_1 \cup R_2 &= \{(x, y) \mid 0 \le x \le 1, 0 \le y \le x\} \cup \{(x, y) \mid 0 \le x \le 1, x \le y \le 1\} \\ &= \{(x, y) \mid 0 \le y \le 1, y \le x \le 1\} \cup \{(x, y) \mid 0 \le y \le 1, 0 \le x \le y\}, \end{aligned}$$

$D_2 = \{(y, z) \mid 0 \le y \le 1, y \le z \le 1\} = \{(y, z) \mid 0 \le z \le 1, 0 \le y \le z\}$, and

$D_3 = \{(x, z) \mid 0 \le x \le 1, x \le z \le 1\} = \{(x, z) \mid 0 \le z \le 1, 0 \le x \le z\}$.

Thus we also have

$$\begin{aligned} E &= \{(x, y, z) \mid 0 \le x \le 1, 0 \le y \le x, x \le z \le 1\} \cup \{(x, y, z) \mid 0 \le x \le 1, x \le y \le 1, y \le z \le 1\} \\ &= \{(x, y, z) \mid 0 \le y \le 1, y \le x \le 1, x \le z \le 1\} \cup \{(x, y, z) \mid 0 \le y \le 1, 0 \le x \le y, y \le z \le 1\} \\ &= \{(x, y, z) \mid 0 \le z \le 1, 0 \le y \le z, 0 \le x \le z\} = \{(x, y, z) \mid 0 \le x \le 1, x \le z \le 1, 0 \le y \le z\} \\ &= \{(x, y, z) \mid 0 \le z \le 1, 0 \le x \le z, 0 \le y \le z\}. \end{aligned}$$

Then

$$\begin{aligned} \int_0^1 \int_y^1 \int_0^z f(x, y, z)\, dx\, dz\, dy &= \int_0^1 \int_0^x \int_x^1 f(x, y, z)\, dz\, dy\, dx + \int_0^1 \int_x^1 \int_y^1 f(x, y, z)\, dz\, dy\, dx \\ &= \int_0^1 \int_y^1 \int_x^1 f(x, y, z)\, dz\, dx\, dy + \int_0^1 \int_0^y \int_y^1 f(x, y, z)\, dz\, dx\, dy \\ &= \int_0^1 \int_0^z \int_0^z f(x, y, z)\, dx\, dy\, dz = \int_0^1 \int_x^1 \int_0^z f(x, y, z)\, dy\, dz\, dx \\ &= \int_0^1 \int_0^z \int_0^z f(x, y, z)\, dy\, dx\, dz \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

41. The region C is the solid bounded by a circular cylinder of radius 2 with axis the z-axis for $-2 \le z \le 2$. We can write $\iiint_C (4 + 5x^2yz^2)\,dV = \iiint_C 4\,dV + \iiint_C 5x^2yz^2\,dV$, but $f(x, y, z) = 5x^2yz^2$ is an odd function with respect to y. Since C is symmetrical about the xz-plane, we have $\iiint_C 5x^2yz^2\,dV = 0$. Thus $\iiint_C (4 + 5x^2yz^2)\,dV = \iiint_C 4\,dV = 4 \cdot V(E) = 4 \cdot \pi(2)^2(4) = 64\pi$.

42. We can write $\iiint_B (z^3 + \sin y + 3)\,dV = \iiint_B z^3\,dV + \iiint_B \sin y\,dV + \iiint_B 3\,dV$. But z^3 is an odd function with respect to z and the region B is symmetric about the xy-plane, so $\iiint_B z^3\,dV = 0$. Similarly, $\sin y$ is an odd function with respect to y and B is symmetric about the xz-plane, so $\iiint_B \sin y\,dV = 0$. Thus $\iiint_B (z^3 + \sin y + 3)\,dV = \iiint_B 3\,dV = 3 \cdot V(B) = 3 \cdot \frac{4}{3}\pi(1)^3 = 4\pi$.

43. The projection of E onto the xy-plane is the disk $D = \{(x, y) \mid x^2 + y^2 \le 1\}$.

$$\begin{aligned} m &= \iiint_E \rho(x, y, z)\,dV = \iint_D \left[\int_0^{1-x^2-y^2} 3\,dz\right] dA = \iint_D 3(1 - x^2 - y^2)\,dA \\ &= 3\int_0^1 \int_0^{2\pi} (1 - r^2)\,r\,dr\,d\theta = 3\int_0^{2\pi} d\theta \int_0^1 (r - r^3)\,dr \\ &= 3\left[\theta\right]_0^{2\pi} \left[\tfrac{1}{2}r^2 - \tfrac{1}{4}r^4\right]_0^1 = 3\,(2\pi)\left(\tfrac{1}{2} - \tfrac{1}{4}\right) = \tfrac{3}{2}\pi \end{aligned}$$

$$\begin{aligned} M_{yz} &= \iiint_E x\rho(x, y, z)\,dV = \iint_D \left[\int_0^{1-x^2-y^2} 3x\,dz\right] dA = \iint_D 3x(1 - x^2 - y^2)\,dA \\ &= 3\int_0^1 \int_0^{2\pi} (r\cos\theta)(1 - r^2)\,r\,dr\,d\theta = 3\int_0^{2\pi} \cos\theta\,d\theta \int_0^1 (r^2 - r^4)\,dr \\ &= 3\left[\sin\theta\right]_0^{2\pi} \left[\tfrac{1}{3}r^3 - \tfrac{1}{5}r^5\right]_0^1 = 3\,(0)\left(\tfrac{1}{3} - \tfrac{1}{5}\right) = 0 \end{aligned}$$

$$\begin{aligned} M_{xz} &= \iiint_E y\rho(x, y, z)\,dV = \iint_D \left[\int_0^{1-x^2-y^2} 3y\,dz\right] dA = \iint_D 3y(1 - x^2 - y^2)\,dA \\ &= 3\int_0^1 \int_0^{2\pi} (r\sin\theta)(1 - r^2)\,r\,dr\,d\theta = 3\int_0^{2\pi} \sin\theta\,d\theta \int_0^1 (r^2 - r^4)\,dr \\ &= 3\left[-\cos\theta\right]_0^{2\pi} \left[\tfrac{1}{3}r^3 - \tfrac{1}{5}r^5\right]_0^1 = 3\,(0)\left(\tfrac{1}{3} - \tfrac{1}{5}\right) = 0 \end{aligned}$$

$$\begin{aligned} M_{xy} &= \iiint_E z\rho(x, y, z)\,dV = \iint_D \left[\int_0^{1-x^2-y^2} 3z\,dz\right] dA = \iint_D \left[\tfrac{3}{2}z^2\right]_{z=0}^{z=1-x^2-y^2} dA \\ &= \tfrac{3}{2}\iint_D (1 - x^2 - y^2)^2\,dA = \tfrac{3}{2}\int_0^1 \int_0^{2\pi} (1 - r^2)^2\,r\,dr\,d\theta \\ &= \tfrac{3}{2}\int_0^{2\pi} d\theta \int_0^1 (r - 2r^3 + r^5)\,dr = \tfrac{3}{2}\left[\theta\right]_0^{2\pi} \left[\tfrac{1}{2}r^2 - \tfrac{1}{2}r^4 + \tfrac{1}{6}r^6\right]_0^1 \\ &= \tfrac{3}{2}\,(2\pi)\left(\tfrac{1}{2} - \tfrac{1}{2} + \tfrac{1}{6}\right) = \tfrac{1}{2}\pi \end{aligned}$$

Thus the mass is $\frac{3}{2}\pi$ and the center of mass is $(\overline{x}, \overline{y}, \overline{z}) = \left(\dfrac{M_{yz}}{m}, \dfrac{M_{xz}}{m}, \dfrac{M_{xy}}{m}\right) = \left(0, 0, \dfrac{1}{3}\right)$.

44. $m = \int_{-1}^1 \int_0^{1-y^2} \int_0^{1-z} 4\,dx\,dz\,dy = 4\int_{-1}^1 \int_0^{1-y^2} (1 - z)\,dz\,dy = 4\int_{-1}^1 \left[z - \frac{1}{2}z^2\right]_{z=0}^{z=1-y^2} dy = 2\int_{-1}^1 (1 - y^4)\,dy = \frac{16}{5}$,

$$\begin{aligned} M_{yz} &= \int_{-1}^1 \int_0^{1-y^2} \int_0^{1-z} 4x\,dx\,dz\,dy = 2\int_{-1}^1 \int_0^{1-y^2} (1 - z)^2\,dz\,dy = 2\int_{-1}^1 \left[-\tfrac{1}{3}(1 - z)^3\right]_{z=0}^{z=1-y^2} dy \\ &= \tfrac{2}{3}\int_{-1}^1 \left(1 - y^6\right) dy = \left(\tfrac{4}{3}\right)\left(\tfrac{6}{7}\right) = \tfrac{24}{21} \end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$M_{xz} = \int_{-1}^{1}\int_{0}^{1-y^2}\int_{0}^{1-z} 4y\,dx\,dz\,dy = \int_{-1}^{1}\int_{0}^{1-y^2} 4y(1-z)\,dz\,dy$

$= \int_{-1}^{1}\left[4y(1-y^2) - 2y(1-y^2)^2\right]dy = \int_{-1}^{1}(2y-2y^5)\,dy = 0$ [the integrand is odd]

$M_{xy} = \int_{-1}^{1}\int_{0}^{1-y^2}\int_{0}^{1-z} 4z\,dx\,dz\,dy = \int_{-1}^{1}\int_{0}^{1-y^2}(4z-4z^2)\,dz\,dy = 2\int_{-1}^{1}\left[(1-y^2)^2 - \frac{2}{3}(1-y^2)^3\right]dy$

$= 2\int_{-1}^{1}\left[\frac{1}{3} - y^4 + \frac{2}{3}y^6\right]dy = \left[\frac{4}{3}y - \frac{4}{5}y^5 + \frac{8}{21}y^7\right]_0^1 = \frac{96}{105} = \frac{32}{35}$

Thus, $(\overline{x}, \overline{y}, \overline{z}) = \left(\frac{5}{14}, 0, \frac{2}{7}\right)$

45. $m = \int_0^a\int_0^a\int_0^a (x^2+y^2+z^2)\,dx\,dy\,dz = \int_0^a\int_0^a\left[\frac{1}{3}x^3 + xy^2 + xz^2\right]_{x=0}^{x=a} dy\,dz = \int_0^a\int_0^a\left(\frac{1}{3}a^3 + ay^2 + az^2\right)dy\,dz$

$= \int_0^a\left[\frac{1}{3}a^3y + \frac{1}{3}ay^3 + ayz^2\right]_{y=0}^{y=a} dz = \int_0^a\left(\frac{2}{3}a^4 + a^2z^2\right)dz = \left[\frac{2}{3}a^4z + \frac{1}{3}a^2z^3\right]_0^a = \frac{2}{3}a^5 + \frac{1}{3}a^5 = a^5$

$M_{yz} = \int_0^a\int_0^a\int_0^a\left[x^3 + x(y^2+z^2)\right]dx\,dy\,dz = \int_0^a\int_0^a\left[\frac{1}{4}a^4 + \frac{1}{2}a^2(y^2+z^2)\right]dy\,dz$

$= \int_0^a\left(\frac{1}{4}a^5 + \frac{1}{6}a^5 + \frac{1}{2}a^3z^2\right)dz = \frac{1}{4}a^6 + \frac{1}{3}a^6 = \frac{7}{12}a^6 = M_{xz} = M_{xy}$ by symmetry of E and $\rho(x,y,z)$

Hence, $(\overline{x}, \overline{y}, \overline{z}) = \left(\frac{7}{12}a, \frac{7}{12}a, \frac{7}{12}a\right)$.

46. $m = \int_0^1\int_0^{1-x}\int_0^{1-x-y} y\,dz\,dy\,dx = \int_0^1\int_0^{1-x}\left[(1-x)y - y^2\right]dy\,dx$

$= \int_0^1\left[\frac{1}{2}(1-x)^3 - \frac{1}{3}(1-x)^3\right]dx = \frac{1}{6}\int_0^1(1-x)^3\,dx = \frac{1}{24}$

$M_{yz} = \int_0^1\int_0^{1-x}\int_0^{1-x-y} xy\,dz\,dy\,dx = \int_0^1\int_0^{1-x}\left[(x-x^2)y - xy^2\right]dy\,dx$

$= \int_0^1\left[\frac{1}{2}x(1-x)^3 - \frac{1}{3}x(1-x)^3\right]dx = \frac{1}{6}\int_0^1\left(x - 3x^2 + 3x^3 - x^4\right)dx = \frac{1}{6}\left(\frac{1}{2} - 1 + \frac{3}{4} - \frac{1}{5}\right) = \frac{1}{120}$

$M_{xz} = \int_0^1\int_0^{1-x}\int_0^{1-x-y} y^2\,dz\,dy\,dx = \int_0^1\int_0^{1-x}\left[(1-x)y^2 - y^3\right]dy\,dx$

$= \int_0^1\left[\frac{1}{3}(1-x)^4 - \frac{1}{4}(1-x)^4\right]dx = \frac{1}{12}\left[-\frac{1}{5}(1-x)^5\right]_0^1 = \frac{1}{60}$

$M_{xy} = \int_0^1\int_0^{1-x}\int_0^{1-x-y} yz\,dz\,dy\,dx = \int_0^1\int_0^{1-x}\left[\frac{1}{2}y(1-x-y)^2\right]dy\,dx$

$= \frac{1}{2}\int_0^1\int_0^{1-x}\left[(1-x)^2y - 2(1-x)y^2 + y^3\right]dy\,dx = \frac{1}{2}\int_0^1\left[\frac{1}{2}(1-x)^4 - \frac{2}{3}(1-x)^4 + \frac{1}{4}(1-x)^4\right]dx$

$= \frac{1}{24}\int_0^1(1-x)^4\,dx = -\frac{1}{24}\left[\frac{1}{5}(1-x)^5\right]_0^1 = \frac{1}{120}$

Hence, $(\overline{x}, \overline{y}, \overline{z}) = \left(\frac{1}{5}, \frac{2}{5}, \frac{1}{5}\right)$.

47. $I_x = \int_0^L\int_0^L\int_0^L k(y^2+z^2)\,dz\,dy\,dx = k\int_0^L\int_0^L\left(Ly^2 + \frac{1}{3}L^3\right)dy\,dx = k\int_0^L \frac{2}{3}L^4\,dx = \frac{2}{3}kL^5$

By symmetry, $I_x = I_y = I_z = \frac{2}{3}kL^5$.

48. $I_x = \int_{-c/2}^{c/2}\int_{-b/2}^{b/2}\int_{-a/2}^{a/2} k(y^2+z^2)\,dx\,dy\,dz = ka\int_{-c/2}^{c/2}\int_{-b/2}^{b/2}(y^2+z^2)\,dy\,dz$

$= ak\int_{-c/2}^{c/2}\left[\frac{1}{3}y^3 + z^2y\right]_{y=-b/2}^{y=b/2} dz = ak\int_{-c/2}^{c/2}\left(\frac{1}{12}b^3 + bz^2\right)dz = ak\left[\frac{1}{12}b^3z + \frac{1}{3}bz^3\right]_{-c/2}^{c/2}$

$= ak\left(\frac{1}{12}b^3c + \frac{1}{12}bc^3\right) = \frac{1}{12}kabc(b^2+c^2)$

By symmetry, $I_y = \frac{1}{12}kabc(a^2+c^2)$ and $I_z = \frac{1}{12}kabc(a^2+b^2)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

49. $I_z = \iiint_E (x^2+y^2)\,\rho(x,y,z)\,dV = \iint\limits_{x^2+y^2\le a^2} \left[\int_0^h k(x^2+y^2)\,dz\right] dA = \iint\limits_{x^2+y^2\le a^2} k(x^2+y^2)h\,dA$

$= kh\int_0^{2\pi}\int_0^a (r^2)\,r\,dr\,d\theta = kh\int_0^{2\pi} d\theta \int_0^a r^3\,dr = kh(2\pi)\left[\frac{1}{4}r^4\right]_0^a = 2\pi kh\cdot\frac{1}{4}a^4 = \frac{1}{2}\pi kha^4$

50. $I_z = \iiint_E (x^2+y^2)\rho(x,y,z)\,dV = \iint\limits_{x^2+y^2\le h^2} \left[\int_{\sqrt{x^2+y^2}}^h k(x^2+y^2)\,dz\right] dA$

$= \iint\limits_{x^2+y^2\le h^2} k(x^2+y^2)\left(h-\sqrt{x^2+y^2}\right) dA = k\int_0^{2\pi}\int_0^h r^2(h-r)\,r\,dr\,d\theta$

$= k\int_0^{2\pi} d\theta \int_0^h \left(r^3h - r^4\right) dr = k(2\pi)\left[\frac{1}{4}r^4h - \frac{1}{5}r^5\right]_0^h = 2\pi k\left(\frac{1}{4}h^5 - \frac{1}{5}h^5\right) = \frac{1}{10}\pi kh^5$

51. (a) $m = \int_{-1}^1\int_{x^2}^1\int_0^{1-y} \sqrt{x^2+y^2}\,dz\,dy\,dx$

(b) $(\overline{x},\overline{y},\overline{z})$ where $\overline{x} = \frac{1}{m}\int_{-1}^1\int_{x^2}^1\int_0^{1-y} x\sqrt{x^2+y^2}\,dz\,dy\,dx$, $\overline{y} = \frac{1}{m}\int_{-1}^1\int_{x^2}^1\int_0^{1-y} y\sqrt{x^2+y^2}\,dz\,dy\,dx$, and

$\overline{z} = \frac{1}{m}\int_{-1}^1\int_{x^2}^1\int_0^{1-y} z\sqrt{x^2+y^2}\,dz\,dy\,dx.$

(c) $I_z = \int_{-1}^1\int_{x^2}^1\int_0^{1-y}(x^2+y^2)\sqrt{x^2+y^2}\,dz\,dy\,dx = \int_{-1}^1\int_{x^2}^1\int_0^{1-y}(x^2+y^2)^{3/2}\,dz\,dy\,dx$

52. (a) $m = \int_{-1}^1\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{\sqrt{1-x^2-y^2}} \sqrt{x^2+y^2+z^2}\,dz\,dx\,dy$

(b) $(\overline{x},\overline{y},\overline{z})$ where $\overline{x} = m^{-1}\int_{-1}^1\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{\sqrt{1-x^2-y^2}} x\sqrt{x^2+y^2+z^2}\,dz\,dx\,dy$,

$\overline{y} = m^{-1}\int_{-1}^1\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{\sqrt{1-x^2-y^2}} y\sqrt{x^2+y^2+z^2}\,dz\,dx\,dy$,

$\overline{z} = m^{-1}\int_{-1}^1\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{\sqrt{1-x^2-y^2}} z\sqrt{x^2+y^2+z^2}\,dz\,dx\,dy$

(c) $I_z = \int_{-1}^1\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{\sqrt{1-x^2-y^2}}(x^2+y^2)(1+x+y+z)\,dz\,dx\,dy$

53. (a) $m = \int_0^1\int_0^{\sqrt{1-x^2}}\int_0^y (1+x+y+z)\,dz\,dy\,dx = \frac{3\pi}{32}+\frac{11}{24}$

(b) $(\overline{x},\overline{y},\overline{z}) = \Bigg(m^{-1}\int_0^1\int_0^{\sqrt{1-x^2}}\int_0^y x(1+x+y+z)\,dz\,dy\,dx,$

$m^{-1}\int_0^1\int_0^{\sqrt{1-x^2}}\int_0^y y(1+x+y+z)\,dz\,dy\,dx,$

$m^{-1}\int_0^1\int_0^{\sqrt{1-x^2}}\int_0^y z(1+x+y+z)\,dz\,dy\,dx\Bigg)$

$= \left(\dfrac{28}{9\pi+44}, \dfrac{30\pi+128}{45\pi+220}, \dfrac{45\pi+208}{135\pi+660}\right)$

(c) $I_z = \displaystyle\int_0^1\int_0^{\sqrt{1-x^2}}\int_0^y (x^2+y^2)(1+x+y+z)\,dz\,dy\,dx = \frac{68+15\pi}{240}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. (a) $m = \int_0^1 \int_{3x}^3 \int_0^{\sqrt{9-y^2}} (x^2+y^2)\,dz\,dy\,dx = \frac{56}{5} = 11.2$

(b) $(\overline{x}, \overline{y}, \overline{z})$ where $\overline{x} = m^{-1} \int_0^1 \int_{3x}^3 \int_0^{\sqrt{9-y^2}} x(x^2+y^2)\,dz\,dy\,dx \approx 0.375$,

$\overline{y} = m^{-1} \int_0^1 \int_{3x}^3 \int_0^{\sqrt{9-y^2}} y(x^2+y^2)\,dz\,dy\,dx = \frac{45\pi}{64} \approx 2.209$,

$\overline{z} = m^{-1} \int_0^1 \int_{3x}^3 \int_0^{\sqrt{9-y^2}} z(x^2+y^2)\,dz\,dy\,dx = \frac{15}{16} = 0.9375$.

(c) $I_z = \int_0^1 \int_{3x}^3 \int_0^{\sqrt{9-y^2}} (x^2+y^2)^2\,dz\,dy\,dx = \frac{10{,}464}{175} \approx 59.79$

55. (a) $f(x,y,z)$ is a joint density function, so we know $\iiint_{\mathbb{R}^3} f(x,y,z)\,dV = 1$. Here we have

$$\begin{aligned}\iiint_{\mathbb{R}^3} f(x,y,z)\,dV &= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f(x,y,z)\,dz\,dy\,dx = \int_0^2\int_0^2\int_0^2 Cxyz\,dz\,dy\,dx \\ &= C\int_0^2 x\,dx \int_0^2 y\,dy \int_0^2 z\,dz = C\left[\tfrac{1}{2}x^2\right]_0^2 \left[\tfrac{1}{2}y^2\right]_0^2 \left[\tfrac{1}{2}z^2\right]_0^2 = 8C\end{aligned}$$

Then we must have $8C = 1 \quad\Rightarrow\quad C = \frac{1}{8}$.

(b) $$\begin{aligned}P(X \le 1, Y \le 1, Z \le 1) &= \int_{-\infty}^1\int_{-\infty}^1\int_{-\infty}^1 f(x,y,z)\,dz\,dy\,dx = \int_0^1\int_0^1\int_0^1 \tfrac{1}{8}xyz\,dz\,dy\,dx \\ &= \tfrac{1}{8}\int_0^1 x\,dx \int_0^1 y\,dy \int_0^1 z\,dz = \tfrac{1}{8}\left[\tfrac{1}{2}x^2\right]_0^1 \left[\tfrac{1}{2}y^2\right]_0^1 \left[\tfrac{1}{2}z^2\right]_0^1 = \tfrac{1}{8}\left(\tfrac{1}{2}\right)^3 = \tfrac{1}{64}\end{aligned}$$

(c) $P(X+Y+Z \le 1) = P((X,Y,Z) \in E)$ where E is the solid region in the first octant bounded by the coordinate planes and the plane $x+y+z=1$. The plane $x+y+z=1$ meets the xy-plane in the line $x+y=1$, so we have

$$\begin{aligned}P(X+Y+Z \le 1) &= \iiint_E f(x,y,z)\,dV = \int_0^1\int_0^{1-x}\int_0^{1-x-y} \tfrac{1}{8}xyz\,dz\,dy\,dx \\ &= \tfrac{1}{8}\int_0^1\int_0^{1-x} xy\left[\tfrac{1}{2}z^2\right]_{z=0}^{z=1-x-y}\,dy\,dx = \tfrac{1}{16}\int_0^1\int_0^{1-x} xy(1-x-y)^2\,dy\,dx \\ &= \tfrac{1}{16}\int_0^1\int_0^{1-x}[(x^3-2x^2+x)y + (2x^2-2x)y^2 + xy^3]\,dy\,dx \\ &= \tfrac{1}{16}\int_0^1 \left[(x^3-2x^2+x)\tfrac{1}{2}y^2 + (2x^2-2x)\tfrac{1}{3}y^3 + x\left(\tfrac{1}{4}y^4\right)\right]_{y=0}^{y=1-x}\,dx \\ &= \tfrac{1}{192}\int_0^1 (x-4x^2+6x^3-4x^4+x^5)\,dx = \tfrac{1}{192}\left(\tfrac{1}{30}\right) = \tfrac{1}{5760}\end{aligned}$$

56. (a) $f(x,y,z)$ is a joint density function, so we know $\iiint_{\mathbb{R}^3} f(x,y,z)\,dV = 1$. Here we have

$$\begin{aligned}\iiint_{\mathbb{R}^3} f(x,y,z)\,dV &= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f(x,y,z)\,dz\,dy\,dx = \int_0^{\infty}\int_0^{\infty}\int_0^{\infty} Ce^{-(0.5x+0.2y+0.1z)}\,dz\,dy\,dx \\ &= C\int_0^{\infty} e^{-0.5x}\,dx \int_0^{\infty} e^{-0.2y}\,dy \int_0^{\infty} e^{-0.1z}\,dz \\ &= C\lim_{t\to\infty}\int_0^t e^{-0.5x}\,dx \lim_{t\to\infty}\int_0^t e^{-0.2y}\,dy \lim_{t\to\infty}\int_0^t e^{-0.1z}\,dz \\ &= C\lim_{t\to\infty}\left[-2e^{-0.5x}\right]_0^t \lim_{t\to\infty}\left[-5e^{-0.2y}\right]_0^t \lim_{t\to\infty}\left[-10e^{-0.1z}\right]_0^t \\ &= C\lim_{t\to\infty}\left[-2(e^{-0.5t}-1)\right] \lim_{t\to\infty}\left[-5(e^{-0.2t}-1)\right] \lim_{t\to\infty}\left[-10(e^{-0.1t}-1)\right] \\ &= C\cdot(-2)(0-1)\cdot(-5)(0-1)\cdot(-10)(0-1) = 100C\end{aligned}$$

So we must have $100C = 1 \quad\Rightarrow\quad C = \frac{1}{100}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) We have no restriction on Z, so

$$\begin{aligned} P(X \le 1, Y \le 1) &= \int_{-\infty}^{1}\int_{-\infty}^{1}\int_{-\infty}^{\infty} f(x,y,z)\,dz\,dy\,dx = \int_0^1\int_0^1\int_0^\infty \tfrac{1}{100}e^{-(0.5x+0.2y+0.1z)}\,dz\,dy\,dx \\ &= \tfrac{1}{100}\int_0^1 e^{-0.5x}\,dx \int_0^1 e^{-0.2y}\,dy \int_0^\infty e^{-0.1z}\,dz \\ &= \tfrac{1}{100}\left[-2e^{-0.5x}\right]_0^1\left[-5e^{-0.2y}\right]_0^1 \lim_{t\to\infty}\left[-10e^{-0.1z}\right]_0^t \qquad \text{[by part (a)]} \\ &= \tfrac{1}{100}(2-2e^{-0.5})(5-5e^{-0.2})(10) = (1-e^{-0.5})(1-e^{-0.2}) \approx 0.07132 \end{aligned}$$

(c) $$\begin{aligned} P(X \le 1, Y \le 1, Z \le 1) &= \int_{-\infty}^{1}\int_{-\infty}^{1}\int_{-\infty}^{1} f(x,y,z)\,dz\,dy\,dx = \int_0^1\int_0^1\int_0^1 \tfrac{1}{100}e^{-(0.5x+0.2y+0.1z)}\,dz\,dy\,dx \\ &= \tfrac{1}{100}\int_0^1 e^{-0.5x}\,dx \int_0^1 e^{-0.2y}\,dy \int_0^1 e^{-0.1z}\,dz \\ &= \tfrac{1}{100}\left[-2e^{-0.5x}\right]_0^1\left[-5e^{-0.2y}\right]_0^1\left[-10e^{-0.1z}\right]_0^1 \\ &= (1-e^{-0.5})(1-e^{-0.2})(1-e^{-0.1}) \approx 0.006787 \end{aligned}$$

57. $V(E) = L^3 \quad\Rightarrow\quad f_{\text{avg}} = \dfrac{1}{L^3}\displaystyle\int_0^L\int_0^L\int_0^L xyz\,dx\,dy\,dz = \dfrac{1}{L^3}\int_0^L x\,dx \int_0^L y\,dy \int_0^L z\,dz$

$$= \frac{1}{L^3}\left[\frac{x^2}{2}\right]_0^L\left[\frac{y^2}{2}\right]_0^L\left[\frac{z^2}{2}\right]_0^L = \frac{1}{L^3}\frac{L^2}{2}\frac{L^2}{2}\frac{L^2}{2} = \frac{L^3}{8}$$

58. The height of each point is given by its z-coordinate, so the average height of the points in $E = \{(x,y,z) \mid x^2+y^2+z^2 \le 1,\ z \ge 0\}$ is

$$\frac{1}{V(E)}\iiint_E z\,dV$$

Here $V(E) = \frac{1}{2}\cdot\frac{4}{3}\pi(1)^3 = \frac{2}{3}\pi$ [half the volume of a sphere], so

$$\begin{aligned} \tfrac{1}{V(E)}\iiint_E z\,dV &= \tfrac{1}{2\pi/3}\int_{-1}^{1}\int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}}\int_0^{\sqrt{1-x^2-y^2}} z\,dz\,dy\,dx = \tfrac{3}{2\pi}\int_{-1}^{1}\int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}}\left[\tfrac{1}{2}z^2\right]_{z=0}^{z=\sqrt{1-x^2-y^2}} dy\,dx \\ &= \tfrac{3}{2\pi}\cdot\tfrac{1}{2}\int_{-1}^{1}\int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}}(1-x^2-y^2)\,dy\,dx = \tfrac{3}{4\pi}\int_0^{2\pi}\int_0^1(1-r^2)\,r\,dr\,d\theta \\ &= \tfrac{3}{4\pi}\int_0^{2\pi}d\theta\int_0^1(r-r^3)\,dr = \tfrac{3}{4\pi}(2\pi)\left[\tfrac{1}{2}r^2-\tfrac{1}{4}r^4\right]_0^1 = \tfrac{3}{2}\left(\tfrac{1}{4}\right) = \tfrac{3}{8} \end{aligned}$$

59. (a) The triple integral will attain its maximum when the integrand $1-x^2-2y^2-3z^2$ is positive in the region E and negative everywhere else. For if E contains some region F where the integrand is negative, the integral could be increased by excluding F from E, and if E fails to contain some part G of the region where the integrand is positive, the integral could be increased by including G in E. So we require that $x^2+2y^2+3z^2 \le 1$. This describes the region bounded by the ellipsoid $x^2+2y^2+3z^2 = 1$.

(b) The maximum value of $\iiint_E (1-x^2-2y^2-3z^2)\,dV$ occurs when E is the solid region bounded by the ellipsoid $x^2+2y^2+3z^2 = 1$. The projection of E on the xy-plane is the planar region bounded by the ellipse $x^2+2y^2 = 1$, so

$$E = \left\{(x,y,z) \mid -1 \le x \le 1, -\sqrt{\tfrac{1}{2}(1-x^2)} \le y \le \sqrt{\tfrac{1}{2}(1-x^2)}, -\sqrt{\tfrac{1}{3}(1-x^2-2y^2)} \le z \le \sqrt{\tfrac{1}{3}(1-x^2-2y^2)}\right\}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and

$$\iiint_E (1 - x^2 - 2y^2 - 3z^2)\, dV = \int_{-1}^{1} \int_{-\sqrt{\frac{1}{2}(1-x^2)}}^{\sqrt{\frac{1}{2}(1-x^2)}} \int_{-\sqrt{\frac{1}{3}(1-x^2-2y^2)}}^{\sqrt{\frac{1}{3}(1-x^2-2y^2)}} (1 - x^2 - 2y^2 - 3z^2)\, dz\, dy\, dx = \frac{4\sqrt{6}}{45}\pi$$

using a CAS.

DISCOVERY PROJECT Volumes of Hyperspheres

In this project we use V_n to denote the n-dimensional volume of an n-dimensional hypersphere.

1. The interior of the circle is the set of points $\left\{(x, y) \mid -r \le y \le r,\ -\sqrt{r^2 - y^2} \le x \le \sqrt{r^2 - y^2}\right\}$. So, substituting $y = r \sin\theta$ and then using Formula 64 from the Table of Integrals to evaluate the integral, we get

$$V_2(r) = \int_{-r}^{r} \int_{-\sqrt{r^2-y^2}}^{\sqrt{r^2-y^2}} dx\, dy = \int_{-r}^{r} 2\sqrt{r^2 - y^2}\, dy = \int_{-\pi/2}^{\pi/2} 2r\sqrt{1 - \sin^2\theta}\ (r\cos\theta\, d\theta)$$

$$= 2r^2 \int_{-\pi/2}^{\pi/2} \cos^2\theta\, d\theta = 2r^2\left[\tfrac{1}{2}\theta + \tfrac{1}{4}\sin 2\theta\right]_{-\pi/2}^{\pi/2} = 2r^2\left(\tfrac{\pi}{2}\right) = \pi r^2$$

2. The region of integration is

$$\left\{(x, y, z) \mid -r \le z \le r, -\sqrt{r^2 - z^2} \le y \le \sqrt{r^2 - z^2}, -\sqrt{r^2 - z^2 - y^2} \le x \le \sqrt{r^2 - z^2 - y^2}\right\}.$$

Substituting $y = \sqrt{r^2 - z^2}\sin\theta$ and using Formula 64 to integrate $\cos^2\theta$, we get

$$V_3(r) = \int_{-r}^{r} \int_{-\sqrt{r^2-z^2}}^{\sqrt{r^2-z^2}} \int_{-\sqrt{r^2-z^2-y^2}}^{\sqrt{r^2-z^2-y^2}} dx\, dy\, dz = \int_{-r}^{r} \int_{-\sqrt{r^2-z^2}}^{\sqrt{r^2-z^2}} 2\sqrt{r^2 - z^2 - y^2}\, dy\, dz$$

$$= \int_{-r}^{r} \int_{-\pi/2}^{\pi/2} 2\sqrt{r^2 - z^2}\sqrt{1 - \sin^2\theta}\left(\sqrt{r^2 - z^2}\cos\theta\, d\theta\right) dz$$

$$= 2\left[\int_{-r}^{r} (r^2 - z^2)\, dz\right]\left[\int_{-\pi/2}^{\pi/2} \cos^2\theta\, d\theta\right] = 2\left(\frac{4r^3}{3}\right)\left(\frac{\pi}{2}\right) = \frac{4\pi r^3}{3}$$

3. The formula for 4-dimensional hypersphere is $x^2 + y^2 + z^2 + w^2 = r^2$. Here we substitute $y = \sqrt{r^2 - w^2 - z^2}\sin\theta$ and, later, $w = r\sin\phi$. Because $\int_{-\pi/2}^{\pi/2} \cos^p\theta\, d\theta$ seems to occur frequently in these calculations, it is useful to find a general formula for that integral. From Exercises 7.1.55–56, we have

$$\int_0^{\pi/2} \sin^{2k} x\, dx = \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2k-1)}{2 \cdot 4 \cdot 6 \cdot \cdots \cdot 2k}\frac{\pi}{2} \qquad \text{and} \qquad \int_0^{\pi/2} \sin^{2k+1} x\, dx = \frac{2 \cdot 4 \cdot 6 \cdot \cdots \cdot 2k}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2k+1)}$$

and from the symmetry of the sine and cosine functions, we can conclude that

$$\int_{-\pi/2}^{\pi/2} \cos^{2k} x\, dx = 2\int_0^{\pi/2} \sin^{2k} x\, dx = \frac{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2k-1)\pi}{2 \cdot 4 \cdot 6 \cdot \cdots \cdot 2k} \tag{1}$$

$$\int_{-\pi/2}^{\pi/2} \cos^{2k+1} x\, dx = 2\int_0^{\pi/2} \sin^{2k+1} x\, dx = \frac{2 \cdot 2 \cdot 4 \cdot 6 \cdot \cdots \cdot 2k}{1 \cdot 3 \cdot 5 \cdot \cdots \cdot (2k+1)} \tag{2}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Thus
$$V_4(r) = \int_{-r}^{r}\int_{-\sqrt{r^2-w^2}}^{\sqrt{r^2-w^2}}\int_{-\sqrt{r^2-w^2-z^2}}^{\sqrt{r^2-w^2-z^2}}\int_{-\sqrt{r^2-w^2-z^2-y^2}}^{\sqrt{r^2-w^2-z^2-y^2}} dx\,dy\,dz\,dw$$
$$= 2\int_{-r}^{r}\int_{-\sqrt{r^2-w^2}}^{\sqrt{r^2-w^2}}\int_{-\sqrt{r^2-w^2-z^2}}^{\sqrt{r^2-w^2-z^2}} \sqrt{r^2-w^2-z^2-y^2}\,dy\,dz\,dw$$
$$= 2\int_{-r}^{r}\int_{-\sqrt{r^2-w^2}}^{\sqrt{r^2-w^2}}\int_{-\pi/2}^{\pi/2} (r^2-w^2-z^2)\cos^2\theta\,d\theta\,dz\,dw \qquad \left[\begin{array}{l} y=\sqrt{r^2-w^2-z^2}\,\sin\theta, \\ dy=\sqrt{r^2-w^2-z^2}\,\cos\theta\,d\theta \end{array}\right]$$
$$= 2\left[\int_{-r}^{r}\int_{-\sqrt{r^2-w^2}}^{\sqrt{r^2-w^2}}(r^2-w^2-z^2)\,dz\,dw\right]\left[\int_{-\pi/2}^{\pi/2}\cos^2\theta\,d\theta\right]$$
$$= 2\left(\tfrac{\pi}{2}\right)\left[\int_{-r}^{r}\tfrac{4}{3}(r^2-w^2)^{3/2}\,dw\right] = \pi\left(\tfrac{4}{3}\right)\int_{-\pi/2}^{\pi/2} r^4\cos^4\phi\,d\phi \qquad \left[\begin{array}{l} w=r\sin\phi, \\ dw=r\cos\phi\,d\phi \end{array}\right]$$
$$= \frac{4\pi}{3}r^4\cdot\frac{1\cdot3\cdot\pi}{2\cdot4} = \frac{\pi^2r^4}{2}$$

4. By using the substitutions $x_i = \sqrt{r^2-x_n^2-x_{n-1}^2-\cdots-x_{i+1}^2}\,\cos\theta_i$ and then applying Formulas 1 and 2 from Problem 3, we can write

$$V_n(r) = \int_{-r}^{r}\int_{-\sqrt{r^2-x_n^2}}^{\sqrt{r^2-x_n^2}}\cdots\int_{-\sqrt{r^2-x_n^2-x_{n-1}^2-\cdots-x_3^2}}^{\sqrt{r^2-x_n^2-x_{n-1}^2-\cdots-x_3^2}}\int_{-\sqrt{r^2-x_n^2-x_{n-1}^2-\cdots-x_3^2-x_2^2}}^{\sqrt{r^2-x_n^2-x_{n-1}^2-\cdots-x_3^2-x_2^2}} dx_1\,dx_2\cdots dx_{n-1}\,dx_n$$
$$= 2\left[\int_{-\pi/2}^{\pi/2}\cos^2\theta_2\,d\theta_2\right]\left[\int_{-\pi/2}^{\pi/2}\cos^3\theta_3\,d\theta_3\right]\cdots\left[\int_{-\pi/2}^{\pi/2}\cos^{n-1}\theta_{n-1}\,d\theta_{n-1}\right]\left[\int_{-\pi/2}^{\pi/2}\cos^n\theta_n\,d\theta_n\right]r^n$$
$$= \begin{cases} \left[2\cdot\dfrac{\pi}{2}\right]\left[\dfrac{2\cdot2}{1\cdot3}\cdot\dfrac{1\cdot3\pi}{2\cdot4}\right]\left[\dfrac{2\cdot2\cdot4}{1\cdot3\cdot5}\cdot\dfrac{1\cdot3\cdot5\pi}{2\cdot4\cdot6}\right]\cdots\left[\dfrac{2\cdot\cdots\cdot(n-2)}{1\cdot\cdots\cdot(n-1)}\cdot\dfrac{1\cdot\cdots\cdot(n-1)\,\pi}{2\cdot\cdots\cdot n}\right]r^n & n \text{ even} \\ 2\left[\dfrac{\pi}{2}\cdot\dfrac{2\cdot2}{1\cdot3}\right]\left[\dfrac{1\cdot3\pi}{2\cdot4}\cdot\dfrac{2\cdot2\cdot4}{1\cdot3\cdot5}\right]\cdots\left[\dfrac{1\cdot\cdots\cdot(n-2)\,\pi}{2\cdot\cdots\cdot(n-1)}\cdot\dfrac{2\cdot\cdots\cdot(n-1)}{1\cdot\cdots\cdot n}\right]r^n & n \text{ odd} \end{cases}$$

By canceling within each set of brackets, we find that

$$V_n(r) = \begin{cases} \dfrac{2\pi}{2}\cdot\dfrac{2\pi}{4}\cdot\dfrac{2\pi}{6}\cdot\cdots\cdot\dfrac{2\pi}{n}r^n = \dfrac{(2\pi)^{n/2}}{2\cdot4\cdot6\cdot\cdots\cdot n}r^n = \dfrac{\pi^{n/2}}{\left(\frac{1}{2}n\right)!}r^n & n \text{ even} \\ 2\cdot\dfrac{2\pi}{3}\cdot\dfrac{2\pi}{5}\cdot\dfrac{2\pi}{7}\cdot\cdots\cdot\dfrac{2\pi}{n}r^n = \dfrac{2(2\pi)^{(n-1)/2}}{3\cdot5\cdot7\cdot\cdots\cdot n}r^n = \dfrac{2^n\left[\frac{1}{2}(n-1)\right]!\,\pi^{(n-1)/2}}{n!}r^n & n \text{ odd} \end{cases}$$

5. We need to show that $\lim\limits_{n\to\infty} V_n(1) = 0$. We'll consider the cases of n even and n odd separately.

For $n = 2k$ and $r = 1$:
$$V_n(1) = \frac{\pi^{n/2}}{\left(\frac{1}{2}n\right)!}\,r^n = \frac{\pi^{2k/2}}{\left(\frac{1}{2}\cdot2k\right)!} = \frac{\pi^k}{k!}$$

Then
$$0 \le \pi\cdot\frac{\pi}{2}\cdot\frac{\pi}{3}\cdots\frac{\pi}{k} \le \pi\cdot\frac{\pi}{2}\cdot\frac{\pi}{3}\cdot\frac{\pi}{k} = \frac{\pi^4}{6k} \qquad [\text{for } k>3]$$

$\dfrac{\pi^4}{6k} \to 0$ as $k\to\infty \quad\Rightarrow\quad V_n(1)\to 0$ as $k\to\infty$ for n even by the Squeeze Theorem.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

For $n = 2k + 1$ and $r = 1$:

$$V_n(1) = \frac{2^n \left[\frac{1}{2}(n-1)\right]! \, \pi^{(n-1)/2}}{n!} \, r^n = \frac{2^{2k+1} \, k! \, \pi^k}{(2k+1)!}$$

$$0 \le 2 \cdot \frac{2^2\pi}{3 \cdot 2} \cdot \frac{2^2(2)\pi}{5 \cdot 4} \cdot \frac{2^2(3)\pi}{7 \cdot 6} \cdot \cdots \cdot \frac{2^2(k)\pi}{(2k+1)(2k)} \le 2 \cdot \frac{2\pi}{3} \cdot \frac{2\pi}{5} \cdot \frac{2\pi}{2k+1} = \frac{2^4\pi^3}{15(2k+1)} \qquad [\text{for } k > 3]$$

$\dfrac{2^4\pi^3}{15(2k+1)} \to 0$ as $k \to \infty \quad \Rightarrow \quad V_n(1) \to 0$ as $k \to \infty$ for n odd by the Squeeze Theorem.

Thus, $\lim\limits_{n\to\infty} V_n(1) = 0$.

15.7 Triple Integrals in Cylindrical Coordinates

1. (a)

From Equations 1, $x = r\cos\theta = 5\cos\frac{\pi}{2} = 5 \cdot 0 = 0$, $y = r\sin\theta = 5\sin\frac{\pi}{2} = 5 \cdot 1 = 5$, and $z = 2$, so the point is $(0, 5, 2)$ in rectangular coordinates.

(b)

From Equations 1, $x = r\cos\theta = 6\cos\left(-\frac{\pi}{4}\right) = 6 \cdot \frac{\sqrt{2}}{2} = 3\sqrt{2}$, $y = r\sin\theta = 6\sin\left(-\frac{\pi}{4}\right) = 6\left(-\frac{\sqrt{2}}{2}\right) = -3\sqrt{2}$, and $z = -3$, so the point is $\left(3\sqrt{2}, -3\sqrt{2}, -3\right)$ in rectangular coordinates.

2. (a)

From Equations 1, $x = r\cos\theta = 2\cos\frac{5\pi}{6} = 2\left(-\frac{\sqrt{3}}{2}\right) = -\sqrt{3}$, $y = r\sin\theta = 2\sin\frac{5\pi}{6} = 2 \cdot \frac{1}{2} = 1$, and $z = 1$, so the point is $\left(-\sqrt{3}, 1, 1\right)$ in rectangular coordinates.

(b)

From Equations 1, $x = r\cos\theta = 8\cos\left(-\frac{2\pi}{3}\right) = 8\left(-\frac{1}{2}\right) = -4$, $y = r\sin\theta = 8\sin\left(-\frac{2\pi}{3}\right) = 8\left(-\frac{\sqrt{3}}{2}\right) = -4\sqrt{3}$, and $z = 5$, so the point is $\left(-4, -4\sqrt{3}, 5\right)$ in rectangular coordinates.

3. (a) $(4, 4, -3)$. From Equations 2, we have $r^2 = x^2 + y^2 = 4^2 + 4^2 = 32$, so $r = \sqrt{32}$. $\tan\theta = \frac{y}{x} = \frac{4}{4} = 1$ and the point $(4, 4)$ is in the first quadrant of the xy-plane, so $\theta = \frac{\pi}{4} + 2\pi n$. Thus, one set of cylindrical coordinates is $(r, \theta, z) = \left(4\sqrt{2}, \frac{\pi}{4}, -3\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\left(5\sqrt{3}, -5, \sqrt{3}\right)$. $r^2 = \left(5\sqrt{3}\right)^2 + (-5)^2 = 100$, so $r = 10$. $\tan\theta = \frac{-5}{5\sqrt{3}} = -\frac{1}{\sqrt{3}}$ and the point $\left(5\sqrt{3}, -5\right)$ is in the fourth quadrant of the xy-plane, so $\theta = \frac{11\pi}{6} + 2\pi n$. Thus, one set of cylindrical coordinates is $(r, \theta, z) = \left(10, -\frac{\pi}{6}, \sqrt{3}\right)$.

4. (a) $(0, -2, 9)$. $r^2 = 0^2 + (-2)^2 = 4$, so $r = 2$. $\tan\theta = 0/(-2)$ is undefined and $y < 0$, so $\theta = \frac{3\pi}{2} + 2\pi n$. Thus, one set of cylindrical coordinates is $(r, \theta, z) = \left(2, \frac{3\pi}{2}, 9\right)$.

(b) $\left(-1, \sqrt{3}, 6\right)$. $r^2 = (-1)^2 + \left(\sqrt{3}\right)^2 = 4$, so $r = 2$. $\tan\theta = \frac{\sqrt{3}}{-1} = -\sqrt{3}$ and the point $\left(-1, \sqrt{3}\right)$ is in the second quadrant of the xy-plane, so $\theta = \frac{2\pi}{3} + 2\pi n$. Thus, one set of cylindrical coordinates is $(r, \theta, z) = \left(2, \frac{2\pi}{3}, 6\right)$.

5. Since $r = 2$, the distance from any point to the z-axis is 2. Because θ and z may vary, the surface is a circular cylinder with radius 2 and axis the z-axis. (See Figure 4.)

Also, $x^2 + y^2 = r^2 = 4$, which we recognize as an equation of this cylinder.

6. Since $\theta = \frac{\pi}{6}$ but r and z may vary, the surface is a vertical plane including the z-axis and intersecting the xy-plane in the line $y = \frac{1}{\sqrt{3}}x$. (Here we are assuming that r can be negative; if we restrict $r \geq 0$, then we get a half-plane.)

7. Since $r^2 + z^2 = 4$ and $r^2 = x^2 + y^2$, we have $x^2 + y^2 + z^2 = 4$, a sphere centered at the origin with radius 2.

8. $r = 2\sin\theta \quad \Rightarrow \quad r^2 = 2r\sin\theta \quad \Rightarrow \quad x^2 + y^2 = 2y \quad \Leftrightarrow \quad x^2 + (y-1)^2 = 1$. z doesn't appear in the equation, so any horizontal trace in $z = k$ is the circle $x^2 + (y-1)^2 = 1$, $z = k$, which has center $(0, 1, k)$ and radius 1. Thus the surface is a circular cylinder with radius 1 and axis the vertical line $x = 0$, $y = 1$.

9. (a) Substituting $x^2 + y^2 = r^2$ and $x = r\cos\theta$, the equation $x^2 - x + y^2 + z^2 = 1$ becomes $r^2 - r\cos\theta + z^2 = 1$ or $z^2 = 1 + r\cos\theta - r^2$.

(b) Substituting $x = r\cos\theta$ and $y = r\sin\theta$, the equation $z = x^2 - y^2$ becomes $z = (r\cos\theta)^2 - (r\sin\theta)^2 = r^2(\cos^2\theta - \sin^2\theta)$ or $z = r^2\cos 2\theta$.

10. (a) The equation $2x^2 + 2y^2 - z^2 = 4$ can be written as $2(x^2 + y^2) - z^2 = 4$ which becomes $2r^2 - z^2 = 4$ or $z^2 = 2r^2 - 4$ in cylindrical coordinates.

(b) Substituting $x = r\cos\theta$ and $y = r\sin\theta$, the equation $2x - y + z = 1$ becomes $2r\cos\theta - r\sin\theta + z = 1$ or $z = 1 + r(\sin\theta - 2\cos\theta)$.

11.

$z = r^2 \quad \Leftrightarrow \quad z = x^2 + y^2$, a circular paraboloid opening upward with vertex the origin, and $z = 8 - r^2 \quad \Leftrightarrow \quad z = 8 - (x^2 + y^2)$, a circular paraboloid opening downward with vertex $(0, 0, 8)$. The paraboloids intersect when $r^2 = 8 - r^2 \quad \Leftrightarrow \quad r^2 = 4$. Thus $r^2 \leq z \leq 8 - r^2$ describes the solid above the paraboloid $z = x^2 + y^2$ and below the paraboloid $z = 8 - x^2 - y^2$ for $x^2 + y^2 \leq 4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12.

$z = r = \sqrt{x^2 + y^2}$ is a cone that opens upward. Thus $r \le z \le 2$ is the region above this cone and beneath the horizontal plane $z = 2$. $0 \le \theta \le \frac{\pi}{2}$ restricts the solid to that part of this region in the first octant.

13. We can position the cylindrical shell vertically so that its axis coincides with the z-axis and its base lies in the xy-plane. If we use centimeters as the unit of measurement, then cylindrical coordinates conveniently describe the shell as $6 \le r \le 7$, $0 \le \theta \le 2\pi$, $0 \le z \le 20$.

14. In cylindrical coordinates, the equations are $z = r^2$ and $z = 5 - r^2$. The curve of intersection is $r^2 = 5 - r^2$ or $r = \sqrt{5/2}$. So we graph the surfaces in cylindrical coordinates, with $0 \le r \le \sqrt{5/2}$. In Maple, we can use the `coords=cylindrical` option in a regular `plot3d` command. In Mathematica, we can use `RevolutionPlot3D` or `ParametricPlot3D`.

15. (a) In cylindrical coordinates, the region can be described as $E = \{(r, \theta, z) \mid 0 \le r \le 1, 0 \le \theta \le \pi, 0 \le z \le 2 - r^2\}$.

Thus, $\iiint_E (x^2 + y^2)\, dV = \int_0^{\pi} \int_0^1 \int_0^{2-r^2} r^2 \cdot r\, dz\, dr\, d\theta$.

(b) $$\int_0^{\pi} \int_0^1 \int_0^{2-r^2} r^3\, dz\, dr\, d\theta = \int_0^{\pi} \int_0^1 r^3 \Big[z \Big]_{z=0}^{z=2-r^2} dr\, d\theta = \int_0^{\pi} \int_0^1 (2r^3 - r^5)\, dr\, d\theta$$
$$= \int_0^{\pi} d\theta \int_0^1 (2r^3 - r^5)\, dr = \Big[\theta \Big]_{\theta=0}^{\theta=\pi} \cdot \left[\frac{r^4}{2} - \frac{r^6}{6} \right]_{r=0}^{r=1} = \frac{\pi}{3}$$

16. (a) In cylindrical coordinates, the region E is bounded above by the paraboloid $z = 6 - r^2$ and below by the cone $z = r$. The paraboloid and cone intersect when $6 - r^2 = r \;\Rightarrow\; r^2 + r - 6 = 0 \;\Rightarrow\; r = 2\ (r > 0)$, so the region can be described as $E = \{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 2, r \le z \le 6 - r^2\}$. Then

$\iiint_E (xy)\, dV = \int_0^{2\pi} \int_0^2 \int_r^{6-r^2} r\cos\theta \cdot r\sin\theta \cdot r\, dz\, dr\, d\theta$.

(b) $$\int_0^{2\pi} \int_0^2 \int_r^{6-r^2} r^3 \cos\theta \sin\theta\, dz\, dr\, d\theta = \int_0^{2\pi} \int_0^2 r^3 \cos\theta \sin\theta \Big[z \Big]_{z=r}^{z=6-r^2} dr\, d\theta$$
$$= \int_0^{2\pi} \int_0^2 r^3 \cos\theta \sin\theta\, (6r^3 - r^4 - r^5)\, dr\, d\theta$$
$$= \int_0^{2\pi} \cos\theta \sin\theta\, d\theta \int_0^2 (6r^3 - r^4 - r^5)\, dr = 0 \cdot \int_0^2 (6r^3 - r^4 - r^5)\, dr = 0$$

17.

The region of integration represents the solid enclosed by the paraboloid $z = r^2$, $(z = x^2 + y^2)$, below the plane $z = 9$ in the second and third quadrants.

$$\int_{\pi/2}^{3\pi/2} \int_0^3 \int_{r^2}^9 r\, dz\, dr\, d\theta = \int_{\pi/2}^{3\pi/2} \int_0^3 [rz]_{z=r^2}^{z=9}\, dr\, d\theta = \int_{\pi/2}^{3\pi/2} \int_0^3 \left(9r - r^3\right)\, dr\, d\theta$$
$$= \int_{\pi/2}^{3\pi/2} d\theta \int_0^3 \left(9r - r^3\right)\, dr = \pi \left[\tfrac{9}{2} r^2 - \tfrac{r^4}{4} \right]_0^3 = \tfrac{81\pi}{4}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18.

The region of integration is given in cylindrical coordinates by $E = \{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 2, 0 \le z \le r\}$. This represents the solid region enclosed by the circular cylinder $r = 2$, bounded above by the cone $z = r$, and bounded below by the xy-plane.

$$\int_0^2 \int_0^{2\pi} \int_0^r r\,dz\,d\theta\,dr = \int_0^2 \int_0^{2\pi} \left[rz\right]_{z=0}^{z=r} d\theta\,dr = \int_0^2 \int_0^{2\pi} r^2\,d\theta\,dr$$
$$= \int_0^2 r^2\,dr \int_0^{2\pi} d\theta = \left[\tfrac{1}{3}r^3\right]_0^2 \left[\theta\right]_0^{2\pi} = \tfrac{8}{3} \cdot 2\pi = \tfrac{16}{3}\pi$$

19. In cylindrical coordinates, E is given by $\{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 4, -5 \le z \le 4\}$. So

$$\iiint_E \sqrt{x^2 + y^2}\,dV = \int_0^{2\pi} \int_0^4 \int_{-5}^4 \sqrt{r^2}\,r\,dz\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^4 r^2\,dr \int_{-5}^4 dz$$
$$= \left[\theta\right]_0^{2\pi} \left[\tfrac{1}{3}r^3\right]_0^4 \left[z\right]_{-5}^4 = (2\pi)\left(\tfrac{64}{3}\right)(9) = 384\pi$$

20. The paraboloid $z = x^2 + y^2 = r^2$ intersects the plane $z = 4$ in the circle $x^2 + y^2 = 4$ or $r^2 = 4 \quad \Rightarrow \quad r = 2$, so in cylindrical coordinates, E is given by $\left\{(r, \theta, z) \,\middle|\, 0 \le \theta \le 2\pi, 0 \le r \le 2, r^2 \le z \le 4\right\}$. Thus

$$\iiint_E z\,dV = \int_0^{2\pi} \int_0^2 \int_{r^2}^4 (z)\,r\,dz\,dr\,d\theta = \int_0^{2\pi} \int_0^2 \left[\tfrac{1}{2}rz^2\right]_{z=r^2}^{z=4} dr\,d\theta$$
$$= \int_0^{2\pi} \int_0^2 \left(8r - \tfrac{1}{2}r^5\right) dr\,d\theta = \int_0^{2\pi} d\theta \int_0^2 \left(8r - \tfrac{1}{2}r^5\right) dr = 2\pi\left[4r^2 - \tfrac{1}{12}r^6\right]_0^2$$
$$= 2\pi\left(16 - \tfrac{16}{3}\right) = \tfrac{64}{3}\pi$$

21. The paraboloid $z = 4 - x^2 - y^2 = 4 - r^2$ intersects the xy-plane in the circle $x^2 + y^2 = 4$ or $r^2 = 4 \quad \Rightarrow \quad r = 2$, so in cylindrical coordinates, E is given by $\left\{(r, \theta, z) \,\middle|\, 0 \le \theta \le \pi/2, 0 \le r \le 2, 0 \le z \le 4 - r^2\right\}$. Thus

$$\iiint_E (x + y + z)\,dV = \int_0^{\pi/2} \int_0^2 \int_0^{4-r^2} (r\cos\theta + r\sin\theta + z)\,r\,dz\,dr\,d\theta$$
$$= \int_0^{\pi/2} \int_0^2 \left[r^2(\cos\theta + \sin\theta)z + \tfrac{1}{2}rz^2\right]_{z=0}^{z=4-r^2} dr\,d\theta$$
$$= \int_0^{\pi/2} \int_0^2 \left[(4r^2 - r^4)(\cos\theta + \sin\theta) + \tfrac{1}{2}r(4 - r^2)^2\right] dr\,d\theta$$
$$= \int_0^{\pi/2} \left[\left(\tfrac{4}{3}r^3 - \tfrac{1}{5}r^5\right)(\cos\theta + \sin\theta) - \tfrac{1}{12}(4 - r^2)^3\right]_{r=0}^{r=2} d\theta$$
$$= \int_0^{\pi/2} \left[\tfrac{64}{15}(\cos\theta + \sin\theta) + \tfrac{16}{3}\right] d\theta = \left[\tfrac{64}{15}(\sin\theta - \cos\theta) + \tfrac{16}{3}\theta\right]_0^{\pi/2}$$
$$= \tfrac{64}{15}(1 - 0) + \tfrac{16}{3} \cdot \tfrac{\pi}{2} - \tfrac{64}{15}(0 - 1) - 0 = \tfrac{8}{3}\pi + \tfrac{128}{15}$$

22. In cylindrical coordinates E is bounded by the planes $z = 0$, $z = r\sin\theta + 4$ and the cylinders $r = 1$ and $r = 4$, so E is given by $\{(r, \theta, z) \mid 0 \le \theta \le 2\pi,\ 1 \le r \le 4,\ 0 \le z \le r\sin\theta + 4\}$. Thus

$$\iiint_E (x - y)\,dV = \int_0^{2\pi} \int_1^4 \int_0^{r\sin\theta+4} (r\cos\theta - r\sin\theta)\,r\,dz\,dr\,d\theta = \int_0^{2\pi} \int_1^4 (r^2\cos\theta - r^2\sin\theta)\left[z\right]_{z=0}^{z=r\sin\theta+4} dr\,d\theta$$
$$= \int_0^{2\pi} \int_1^4 (r^2\cos\theta - r^2\sin\theta)(r\sin\theta + 4)\,dr\,d\theta$$
$$= \int_0^{2\pi} \int_1^4 \left[r^3(\sin\theta\cos\theta - \sin^2\theta) + 4r^2(\cos\theta - \sin\theta)\right] dr\,d\theta$$
$$= \int_0^{2\pi} \left[\tfrac{1}{4}r^4(\sin\theta\cos\theta - \sin^2\theta) + \tfrac{4}{3}r^3(\cos\theta - \sin\theta)\right]_{r=1}^{r=4} d\theta$$
$$= \int_0^{2\pi} \left[\left(64 - \tfrac{1}{4}\right)(\sin\theta\cos\theta - \sin^2\theta) + \left(\tfrac{256}{3} - \tfrac{4}{3}\right)(\cos\theta - \sin\theta)\right] d\theta$$
$$= \int_0^{2\pi} \left[\tfrac{255}{4}(\sin\theta\cos\theta - \sin^2\theta) + 84(\cos\theta - \sin\theta)\right] d\theta$$
$$= \left[\tfrac{255}{4}\left(\tfrac{1}{2}\sin^2\theta - \left(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right)\right) + 84(\sin\theta + \cos\theta)\right]_0^{2\pi} = \tfrac{255}{4}(-\pi) + 84(1) - 0 - 84(1) = -\tfrac{255}{4}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. In cylindrical coordinates, E is bounded by the cylinder $r = 1$, the plane $z = 0$, and the cone $z = 2r$. So $E = \{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 1, 0 \le z \le 2r\}$ and

$$\iiint_E x^2\, dV = \int_0^{2\pi}\int_0^1\int_0^{2r} r^2\cos^2\theta\, r\, dz\, dr\, d\theta = \int_0^{2\pi}\int_0^1 \left[r^3\cos^2\theta\, z\right]_{z=0}^{z=2r}\, dr\, d\theta = \int_0^{2\pi}\int_0^1 2r^4\cos^2\theta\, dr\, d\theta$$
$$= \int_0^{2\pi}\left[\tfrac{2}{5}r^5\cos^2\theta\right]_{r=0}^{r=1} d\theta = \tfrac{2}{5}\int_0^{2\pi}\cos^2\theta\, d\theta = \tfrac{2}{5}\int_0^{2\pi}\tfrac{1}{2}(1+\cos 2\theta)\, d\theta = \tfrac{1}{5}\left[\theta + \tfrac{1}{2}\sin 2\theta\right]_0^{2\pi} = \tfrac{2\pi}{5}$$

24. In cylindrical coordinates E is the solid region within the cylinder $r = 1$ bounded above and below by the sphere $r^2 + z^2 = 4$, so $E = \left\{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 1, -\sqrt{4-r^2} \le z \le \sqrt{4-r^2}\right\}$. Thus the volume is

$$\iiint_E dV = \int_0^{2\pi}\int_0^1\int_{-\sqrt{4-r^2}}^{\sqrt{4-r^2}} r\, dz\, dr\, d\theta = \int_0^{2\pi}\int_0^1 2r\sqrt{4-r^2}\, dr\, d\theta$$
$$= \int_0^{2\pi} d\theta \int_0^1 2r\sqrt{4-r^2}\, dr = 2\pi\left[-\tfrac{2}{3}(4-r^2)^{3/2}\right]_0^1 = \tfrac{4}{3}\pi(8 - 3^{3/2})$$

25. In cylindrical coordinates, E is bounded below by the cone $z = r$ and above by the sphere $r^2 + z^2 = 2$ or $z = \sqrt{2-r^2}$. The cone and the sphere intersect when $2r^2 = 2 \quad\Rightarrow\quad r = 1$, so $E = \left\{(r, \theta, z) \mid 0 \le \theta \le 2\pi, 0 \le r \le 1, r \le z \le \sqrt{2-r^2}\right\}$ and the volume is

$$\iiint_E dV = \int_0^{2\pi}\int_0^1\int_r^{\sqrt{2-r^2}} r\, dz\, dr\, d\theta = \int_0^{2\pi}\int_0^1 [rz]_{z=r}^{z=\sqrt{2-r^2}}\, dr\, d\theta = \int_0^{2\pi}\int_0^1 \left(r\sqrt{2-r^2} - r^2\right) dr\, d\theta$$
$$= \int_0^{2\pi} d\theta \int_0^1 \left(r\sqrt{2-r^2} - r^2\right) dr = 2\pi\left[-\tfrac{1}{3}(2-r^2)^{3/2} - \tfrac{1}{3}r^3\right]_0^1$$
$$= 2\pi\left(-\tfrac{1}{3}\right)(1 + 1 - 2^{3/2}) = -\tfrac{2}{3}\pi\left(2 - 2\sqrt{2}\right) = \tfrac{4}{3}\pi\left(\sqrt{2} - 1\right)$$

26. In cylindrical coordinates, E is bounded below by the paraboloid $z = r^2$ and above by the sphere $r^2 + z^2 = 2$ or $z = \sqrt{2-r^2}$. The paraboloid and the sphere intersect when $r^2 + r^4 = 2 \quad\Rightarrow\quad (r^2+2)(r^2-1) = 0 \quad\Rightarrow\quad r = 1$, so $E = \left\{(r, \theta, z) \mid 0 \le \theta \le 2\pi,\ 0 \le r \le 1,\ r^2 \le z \le \sqrt{2-r^2}\ \right\}$ and the volume is

$$\iiint_E dV = \int_0^{2\pi}\int_0^1\int_{r^2}^{\sqrt{2-r^2}} r\, dz\, dr\, d\theta = \int_0^{2\pi}\int_0^1 [rz]_{z=r^2}^{z=\sqrt{2-r^2}}\, dr\, d\theta = \int_0^{2\pi}\int_0^1 \left(r\sqrt{2-r^2} - r^3\right) dr\, d\theta$$
$$= \int_0^{2\pi} d\theta \int_0^1 \left(r\sqrt{2-r^2} - r^3\right) dr = 2\pi\left[-\tfrac{1}{3}(2-r^2)^{3/2} - \tfrac{1}{4}r^4\right]_0^1$$
$$= 2\pi(-\tfrac{1}{3} - \tfrac{1}{4} + \tfrac{1}{3}\cdot 2^{3/2} - 0) = 2\pi\left(-\tfrac{7}{12} + \tfrac{2}{3}\sqrt{2}\right) = \left(-\tfrac{7}{6} + \tfrac{4}{3}\sqrt{2}\right)\pi$$

27. (a) In cylindrical coordinates, E is bounded above by the paraboloid $z = 24 - r^2$ and below by the cone $z = 2\sqrt{r^2}$ or $z = 2r$ $(r \ge 0)$. The surfaces intersect when $24 - r^2 = 2r \quad\Rightarrow\quad r^2 + 2r - 24 = 0 \quad\Rightarrow\quad (r+6)(r-4) = 0 \quad\Rightarrow\quad r = 4$, so $E = \left\{(r, \theta, z) \mid 2r \le z \le 24 - r^2, 0 \le r \le 4, 0 \le \theta \le 2\pi\right\}$ and the volume is

$$\iiint_E dV = \int_0^{2\pi}\int_0^4\int_{2r}^{24-r^2} r\, dz\, dr\, d\theta = \int_0^{2\pi}\int_0^4 r\left(24 - r^2 - 2r\right) dr\, d\theta = \int_0^{2\pi} d\theta \int_0^4 \left(24r - r^3 - 2r^2\right) dr$$
$$= 2\pi\left[12r^2 - \tfrac{1}{4}r^4 - \tfrac{2}{3}r^3\right]_0^4 = 2\pi\left(192 - 64 - \tfrac{128}{3}\right) = \tfrac{512}{3}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) For constant density K, $m = KV = \frac{512}{3}\pi K$ from part (a). Since the region is homogeneous and symmetric, $M_{yz} = M_{xz} = 0$ and

$$M_{xy} = \int_0^{2\pi}\int_0^4\int_{2r}^{24-r^2}(zK)\,r\,dz\,dr\,d\theta = K\int_0^{2\pi}\int_0^4 r\left[\tfrac{1}{2}z^2\right]_{z=2r}^{z=24-r^2}dr\,d\theta$$
$$= \tfrac{K}{2}\int_0^{2\pi}\int_0^4 r[(24-r^2)^2-4r^2]\,dr\,d\theta = \tfrac{K}{2}\int_0^{2\pi}d\theta\int_0^4(576r-52r^3+r^5)\,dr$$
$$= \tfrac{K}{2}(2\pi)\left[288r^2-13r^4+\tfrac{1}{6}r^6\right]_0^4 = \pi K\left(4608-3328+\tfrac{2048}{3}\right) = \tfrac{5888}{3}\pi K$$

Thus $(\overline{x},\overline{y},\overline{z}) = \left(\dfrac{M_{yz}}{m},\dfrac{M_{xz}}{m},\dfrac{M_{xy}}{m}\right) = \left(0,0,\frac{5888\pi K/3}{512\pi K/3}\right) = \left(0,0,\frac{23}{2}\right)$.

28. (a) $V = \int_{-\pi/2}^{\pi/2}\int_0^{a\cos\theta}\int_{-\sqrt{a^2-r^2}}^{\sqrt{a^2-r^2}} r\,dz\,dr\,d\theta$

$$= 4\int_0^{\pi/2}\int_0^{a\cos\theta}\int_0^{\sqrt{a^2-r^2}} r\,dz\,dr\,d\theta$$
$$= 4\int_0^{\pi/2}\int_0^{a\cos\theta} r\sqrt{a^2-r^2}\,dr\,d\theta$$
$$= -\tfrac{4}{3}\int_0^{\pi/2}\left[(a^2-r^2)^{3/2}\right]_{r=0}^{r=a\cos\theta}d\theta$$
$$= -\tfrac{4}{3}\int_0^{\pi/2}\left[(a^2-a^2\cos^2\theta)^{3/2}-a^3\right]d\theta$$
$$= -\tfrac{4}{3}\int_0^{\pi/2}\left[(a^2\sin^2\theta)^{3/2}-a^3\right]d\theta$$
$$= -\tfrac{4}{3}\int_0^{\pi/2}(a^3\sin^3\theta-a^3)\,d\theta$$
$$= -\frac{4a^3}{3}\int_0^{\pi/2}\left[\sin\theta\,(1-\cos^2\theta)-1\right]d\theta$$
$$= -\frac{4a^3}{3}\left[-\cos\theta+\tfrac{1}{3}\cos^3\theta-\theta\right]_0^{\pi/2} = -\frac{4a^3}{3}\left(-\tfrac{\pi}{2}+\tfrac{2}{3}\right) = \tfrac{2}{9}a^3(3\pi-4)$$

(b)

To plot the cylinder and the sphere on the same screen in Maple, we can use the sequence of commands

```
sphere:=plot3d(1,theta=0..2*Pi,phi=0..Pi,coords=spherical):
cylinder:=plot3d(cos(theta),theta=-Pi/2..Pi/2,z=-1..1,coords=cylindrical):
with(plots):
display3d({sphere,cylinder});
```

In Mathematica, we can use

```
sphere=SphericalPlot3D[1,{phi,0,Pi},{theta,0,2Pi}]
cylinder=ParametricPlot3D[{(Cos[theta])^2,Cos[theta]*Sin[theta],z},
                                {theta,-Pi/2,Pi/2},{z,-1,1}]
Show[sphere,cylinder]
```

29. The paraboloid $z = 4x^2+4y^2$ intersects the plane $z = a$ when $a = 4x^2+4y^2$ or $x^2+y^2 = \frac{1}{4}a$. So, in cylindrical coordinates, $E = \left\{(r,\theta,z) \mid 0 \le r \le \frac{1}{2}\sqrt{a}, 0 \le \theta \le 2\pi, 4r^2 \le z \le a\right\}$. Thus

$$m = \int_0^{2\pi}\int_0^{\sqrt{a}/2}\int_{4r^2}^{a} Kr\,dz\,dr\,d\theta = K\int_0^{2\pi}\int_0^{\sqrt{a}/2}(ar-4r^3)\,dr\,d\theta$$
$$= K\int_0^{2\pi}\left[\tfrac{1}{2}ar^2-r^4\right]_{r=0}^{r=\sqrt{a}/2}d\theta = K\int_0^{2\pi}\tfrac{1}{16}a^2\,d\theta = \tfrac{1}{8}a^2\pi K$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Since the region is homogeneous and symmetric, $M_{yz} = M_{xz} = 0$ and

$$M_{xy} = \int_0^{2\pi}\int_0^{\sqrt{a}/2}\int_{4r^2}^{a} Krz\,dz\,dr\,d\theta = K\int_0^{2\pi}\int_0^{\sqrt{a}/2}\left(\tfrac{1}{2}a^2r - 8r^5\right)dr\,d\theta$$
$$= K\int_0^{2\pi}\left[\tfrac{1}{4}a^2r^2 - \tfrac{4}{3}r^6\right]_{r=0}^{r=\sqrt{a}/2} d\theta = K\int_0^{2\pi}\tfrac{1}{24}a^3\,d\theta = \tfrac{1}{12}a^3\pi K$$

Hence $(\overline{x}, \overline{y}, \overline{z}) = \left(0, 0, \tfrac{2}{3}a\right)$.

30. Since density is proportional to the distance from the z-axis, we can say $\rho(x, y, z) = K\sqrt{x^2 + y^2}$. Then

$$m = \int_0^{2\pi}\int_0^a\int_{-\sqrt{a^2-r^2}}^{\sqrt{a^2-r^2}}(Kr)\,r\,dz\,dr\,d\theta = 2\int_0^{2\pi}\int_0^a\int_0^{\sqrt{a^2-r^2}} Kr^2\,dz\,dr\,d\theta = 2K\int_0^{2\pi}\int_0^a r^2\sqrt{a^2-r^2}\,dr\,d\theta$$
$$= 2K\int_0^{2\pi}\left[\tfrac{1}{8}r(2r^2-a^2)\sqrt{a^2-r^2} + \tfrac{1}{8}a^4\sin^{-1}(r/a)\right]_{r=0}^{r=a} d\theta = 2K\int_0^{2\pi}\left[\left(\tfrac{1}{8}a^4\right)\left(\tfrac{\pi}{2}\right)\right] d\theta = \tfrac{1}{4}a^4\pi^2K$$

31. The region of integration is the region above the cone $z = \sqrt{x^2 + y^2}$, or $z = r$, and below the plane $z = 2$. Also, we have $-2 \le y \le 2$ with $-\sqrt{4 - y^2} \le x \le \sqrt{4 - y^2}$ which describes a circle of radius 2 in the xy-plane centered at $(0, 0)$. Thus,

$$\int_{-2}^{2}\int_{-\sqrt{4-y^2}}^{\sqrt{4-y^2}}\int_{\sqrt{x^2+y^2}}^{2} xz\,dz\,dx\,dy = \int_0^{2\pi}\int_0^2\int_r^2 (r\cos\theta)\,z\,r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^2\int_r^2 r^2(\cos\theta)\,z\,dz\,dr\,d\theta$$
$$= \int_0^{2\pi}\int_0^2 r^2(\cos\theta)\left[\tfrac{1}{2}z^2\right]_{z=r}^{z=2} dr\,d\theta = \tfrac{1}{2}\int_0^{2\pi}\int_0^2 r^2(\cos\theta)\left(4 - r^2\right)dr\,d\theta$$
$$= \tfrac{1}{2}\int_0^{2\pi}\cos\theta\,d\theta\int_0^2\left(4r^2 - r^4\right)dr = \tfrac{1}{2}\left[\sin\theta\right]_0^{2\pi}\left[\tfrac{4}{3}r^3 - \tfrac{1}{5}r^5\right]_0^2 = 0$$

32. The region of integration is the region above the plane $z = 0$ and below the paraboloid $z = 9 - x^2 - y^2$. Also, we have $-3 \le x \le 3$ with $0 \le y \le \sqrt{9 - x^2}$ which describes the upper half of a circle of radius 3 in the xy-plane centered at $(0, 0)$. Thus,

$$\int_{-3}^{3}\int_0^{\sqrt{9-x^2}}\int_0^{9-x^2-y^2}\sqrt{x^2+y^2}\,dz\,dy\,dx = \int_0^{\pi}\int_0^3\int_0^{9-r^2}\sqrt{r^2}\,r\,dz\,dr\,d\theta = \int_0^{\pi}\int_0^3\int_0^{9-r^2} r^2\,dz\,dr\,d\theta$$
$$= \int_0^{\pi}\int_0^3 r^2\left(9 - r^2\right)dr\,d\theta = \int_0^{\pi}d\theta\int_0^3\left(9r^2 - r^4\right)dr$$
$$= \left[\theta\right]_0^{\pi}\left[3r^3 - \tfrac{1}{5}r^5\right]_0^3 = \pi\left(81 - \tfrac{243}{5}\right) = \tfrac{162}{5}\pi$$

33. (a) The mountain comprises a solid conical region C. The work done in lifting a small volume of material ΔV with density $g(P)$ to a height $h(P)$ above sea level is $h(P)g(P)\,\Delta V$. Summing over the whole mountain we get $W = \iiint_C h(P)g(P)\,dV$.

(b) Here C is a solid right circular cone with radius $R = 62{,}000$ ft, height $H = 12{,}400$ ft, and density $g(P) = 200$ lb/ft^3 at all points P in C. We use cylindrical coordinates:

$$\frac{r}{R} = \frac{H - z}{H} = 1 - \frac{z}{H}$$

$$W = \int_0^{2\pi}\int_0^H\int_0^{R(1-z/H)} z\cdot 200r\,dr\,dz\,d\theta = 2\pi\int_0^H 200z\left[\tfrac{1}{2}r^2\right]_{r=0}^{r=R(1-z/H)} dz$$
$$= 400\pi\int_0^H z\,\frac{R^2}{2}\left(1 - \frac{z}{H}\right)^2 dz = 200\pi R^2\int_0^H\left(z - \frac{2z^2}{H} + \frac{z^3}{H^2}\right)dz$$
$$= 200\pi R^2\left[\frac{z^2}{2} - \frac{2z^3}{3H} + \frac{z^4}{4H^2}\right]_0^H = 200\pi R^2\left(\frac{H^2}{2} - \frac{2H^2}{3} + \frac{H^2}{4}\right)$$
$$= \tfrac{50}{3}\pi R^2H^2 = \tfrac{50}{3}\pi(62{,}000)^2(12{,}400)^2 \approx 3.1\times 10^{19}\text{ ft-lb}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

DISCOVERY PROJECT The Intersection of Three Cylinders

1. The three cylinders in the illustration in the text can be visualized as representing the surfaces $x^2 + y^2 = 1$, $x^2 + z^2 = 1$, and $y^2 + z^2 = 1$. Then we sketch the solid of intersection with the coordinate axes and equations indicated. To be more precise, we start by finding the bounding curves of the solid (shown in the first graph below) enclosed by the two cylinders $x^2 + z^2 = 1$ and $y^2 + z^2 = 1$: $x = \pm y = \pm\sqrt{1 - z^2}$ are the symmetric equations, and these can be expressed parametrically as $x = s$, $y = \pm s$, $z = \pm\sqrt{1 - s^2}$, $-1 \le s \le 1$. Now the cylinder $x^2 + y^2 = 1$ intersects these curves at the eight points $\left(\pm\frac{1}{\sqrt{2}}, \pm\frac{1}{\sqrt{2}}, \pm\frac{1}{\sqrt{2}}\right)$. The resulting solid has twelve curved faces bounded by "edges" which are arcs of circles, as shown in the third diagram. Each cylinder defines four of the twelve faces.

2. To find the volume, we split the solid into sixteen congruent pieces, one of which lies in the part of the first octant with $0 \le \theta \le \frac{\pi}{4}$. (Naturally, we use cylindrical coordinates!) This piece is described by $\left\{(r, \theta, z) \mid 0 \le r \le 1, 0 \le \theta \le \frac{\pi}{4}, 0 \le z \le \sqrt{1 - x^2}\right\}$, and so, substituting $x = r\cos\theta$, the volume of the entire solid is

$$V = 16\int_0^{\pi/4}\int_0^1\int_0^{\sqrt{1-x^2}} r\,dz\,dr\,d\theta$$

$$= 16\int_0^{\pi/4}\int_0^1 r\sqrt{1 - r^2\cos^2\theta}\,dr\,d\theta$$

$$= 16 - 8\sqrt{2} \approx 4.6863$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. To graph the edges of the solid, we use parametrized curves similar to those found in Problem 1 for the intersection of two cylinders. We must restrict the parameter intervals so that each arc extends exactly to the desired vertex. One possible set of parametric equations (with all sign choices allowed) is

$x = r,\ y = \pm r,\ z = \pm\sqrt{1-r^2},\ -\frac{1}{\sqrt{2}} \le r \le \frac{1}{\sqrt{2}}$;

$x = \pm s,\ y = \pm\sqrt{1-s^2},\ z = s,\ -\frac{1}{\sqrt{2}} \le s \le \frac{1}{\sqrt{2}}$;

$x = \pm\sqrt{1-t^2},\ y = t,\ z = \pm t,\ -\frac{1}{\sqrt{2}} \le t \le \frac{1}{\sqrt{2}}$.

4. Let the three cylinders be $x^2 + y^2 = a^2$, $x^2 + z^2 = 1$, and $y^2 + z^2 = 1$.

If $a < 1$, then the four faces defined by the cylinder $x^2 + y^2 = 1$ in Problem 1 collapse into a single face, as in the first graph. If $1 < a < \sqrt{2}$, then each pair of vertically opposed faces, defined by one of the other two cylinders, collapse into a single face, as in the second graph. If $a \ge \sqrt{2}$, then the vertical cylinder encloses the solid of intersection of the other two cylinders completely, so the solid of intersection coincides with the solid of intersection of the two cylinders $x^2 + z^2 = 1$ and $y^2 + z^2 = 1$, as illustrated in Problem 1.

If we were to vary b or c instead of a, we would get solids with the same shape, but differently oriented.

$a = 0.95,\ b = c = 1$

$a = 1.1,\ b = c = 1$

5. If $a < 1$, the solid looks similar to the first graph in Problem 4. As in Problem 2, we split the solid into sixteen congruent pieces, one of which can be described as the solid above the polar region $\left\{(r,\theta) \mid 0 \le r \le a,\ 0 \le \theta \le \frac{\pi}{4}\right\}$ in the xy-plane and below the surface $z = \sqrt{1-x^2} = \sqrt{1-r^2\cos^2\theta}$. Thus, the total volume is $V = 16\int_0^{\pi/4}\int_0^a \sqrt{1-r^2\cos^2\theta}\, r\, dr\, d\theta$.

If $a > 1$ and $a < \sqrt{2}$, we have a solid similar to the second graph in Problem 4. Its intersection with the xy-plane is graphed at the right. Again we split the solid into sixteen congruent pieces, one of which is the solid above the region shown in the second figure and below the surface $z = \sqrt{1-x^2} = \sqrt{1-r^2\cos^2\theta}$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

We split the region of integration where the outside boundary changes from the vertical line $x = 1$ to the circle $x^2 + y^2 = a^2$ or $r = a$. R_1 is a right triangle, so $\cos\theta = \frac{1}{a}$. Thus, the boundary between R_1 and R_2 is $\theta = \cos^{-1}\left(\frac{1}{a}\right)$ in polar coordinates, or $y = \sqrt{a^2 - 1}\,x$ in rectangular coordinates. Using rectangular coordinates for the region R_1 and polar coordinates for R_2, we find the total volume of the solid to be

$$V = 16\left[\int_0^1 \int_0^{\sqrt{a^2-1}\,x} \sqrt{1 - x^2}\,dy\,dx + \int_{\cos^{-1}(1/a)}^{\pi/4} \int_0^a \sqrt{1 - r^2\cos^2\theta}\,r\,dr\,d\theta\right]$$

If $a \geq \sqrt{2}$, the cylinder $x^2 + y^2 = 1$ completely encloses the intersection of the other two cylinders, so the solid of intersection of the three cylinders coincides with the intersection of $x^2 + z^2 = 1$ and $y^2 + z^2 = 1$ as illustrated in Exercise 15.5.26. Its volume is $V = 16\int_0^1 \int_0^x \sqrt{1 - x^2}\,dy\,dx$.

15.8 Triple Integrals in Spherical Coordinates

1. (a)

From Equations 1 with $(\rho, \theta, \phi) = \left(2, \frac{3\pi}{4}, \frac{\pi}{2}\right)$,

$x = \rho\sin\phi\cos\theta = 2\sin\frac{\pi}{2}\cos\frac{3\pi}{4} = 2(1)\left(-\frac{\sqrt{2}}{2}\right) = -\sqrt{2}$,

$y = \rho\sin\phi\sin\theta = 2\sin\frac{\pi}{2}\sin\frac{3\pi}{4} = 2(1)\left(\frac{\sqrt{2}}{2}\right) = \sqrt{2}$, and

$z = \rho\cos\phi = 2\cos\frac{\pi}{2} = 2 \cdot 0 = 0$, so the point is $\left(-\sqrt{2}, \sqrt{2}, 0\right)$ in rectangular coordinates.

(b)

From Equations 1 with $(\rho, \theta, \phi) = \left(4, -\frac{\pi}{3}, \frac{\pi}{4}\right)$,

$x = \rho\sin\phi\cos\theta = 4\sin\frac{\pi}{4}\cos\left(-\frac{\pi}{3}\right) = 4\left(\frac{\sqrt{2}}{2}\right)\left(\frac{1}{2}\right) = \sqrt{2}$,

$y = \rho\sin\phi\sin\theta = 4\sin\frac{\pi}{4}\sin\left(-\frac{\pi}{3}\right) = 4\left(\frac{\sqrt{2}}{2}\right)\left(-\frac{\sqrt{3}}{2}\right) = -\sqrt{6}$, and

$z = \rho\cos\phi = 4\cos\frac{\pi}{4} = 4\left(\frac{\sqrt{2}}{2}\right) = 2\sqrt{2}$, so the point is $\left(\sqrt{2}, -\sqrt{6}, 2\sqrt{2}\right)$ in rectangular coordinates.

2. (a)

From Equations 1 with $(\rho, \theta, \phi) = \left(5, \frac{\pi}{2}, \frac{\pi}{3}\right)$,

$x = \rho\sin\phi\cos\theta = 5\sin\frac{\pi}{3}\cos\frac{\pi}{2} = 5 \cdot \frac{\sqrt{3}}{2} \cdot 0 = 0$,

$y = \rho\sin\phi\sin\theta = 5\sin\frac{\pi}{3}\sin\frac{\pi}{2} = 5 \cdot \frac{\sqrt{3}}{2} \cdot 1 = \frac{5\sqrt{3}}{2}$, and

$z = \rho\cos\phi = 5\cos\frac{\pi}{3} = 5 \cdot \frac{1}{2} = \frac{5}{2}$, so the point is $\left(0, \frac{5\sqrt{3}}{2}, \frac{5}{2}\right)$ in rectangular coordinates.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b)

From Equations 1 with $(\rho, \theta, \phi) = \left(6, 0, \frac{5\pi}{6}\right)$,

$x = \rho \sin\phi \cos\theta = 6 \sin\frac{5\pi}{6} \cos 0 = 6 \cdot \frac{1}{2} \cdot 1 = 3$,

$y = \rho \sin\phi \sin\theta = 6 \sin\frac{5\pi}{6} \sin 0 = 6 \cdot \frac{1}{2} \cdot 0 = 0$, and

$z = \rho \cos\phi = 6 \cos\frac{5\pi}{6} = 6\left(-\frac{\sqrt{3}}{2}\right) = -3\sqrt{3}$, so the point is

$\left(3, 0, -3\sqrt{3}\right)$.

3. (a) From Equations 1 and 2 with $(x, y, z) = (3, 3, 0)$, $\rho = \sqrt{x^2 + y^2 + z^2} = \sqrt{3^2 + 3^2 + 0^2} = 3\sqrt{2}$,

$\cos\phi = \dfrac{z}{\rho} = \dfrac{0}{3\sqrt{2}} = 0 \quad \Rightarrow \quad \phi = \frac{\pi}{2}$ and $\cos\theta = \dfrac{x}{\rho \sin\phi} = \dfrac{3}{3\sqrt{2}} = \dfrac{1}{\sqrt{2}} \quad \Rightarrow \quad \theta = \dfrac{\pi}{4}$ [since $x > 0$ and $y > 0$].

Thus, spherical coordinates are $\left(3\sqrt{2}, \frac{\pi}{4}, \frac{\pi}{2}\right)$.

(b) $\rho = \sqrt{1^2 + (-\sqrt{3})^2 + (2\sqrt{3})^2} = 4$, $\cos\phi = \dfrac{z}{\rho} = \dfrac{2\sqrt{3}}{4} = \dfrac{\sqrt{3}}{2} \quad \Rightarrow \quad \phi = \dfrac{\pi}{6}$, and

$\cos\theta = \dfrac{x}{\rho \sin\phi} = \dfrac{1}{4 \sin\frac{\pi}{6}} = \dfrac{1}{2} \quad \Rightarrow \quad \theta = -\dfrac{\pi}{3}$ [since $x > 0$ and $y < 0$]. Thus, spherical coordinates are $\left(4, -\frac{\pi}{3}, \frac{\pi}{6}\right)$.

4. (a) $\rho = \sqrt{0^2 + 4^2 + (-4)^2} = 4\sqrt{2}$, $\cos\phi = \dfrac{z}{\rho} = \dfrac{-4}{4\sqrt{2}} = \dfrac{-1}{\sqrt{2}} \quad \Rightarrow \quad \phi = \dfrac{3\pi}{4}$, and

$\cos\theta = \dfrac{x}{\rho \sin\phi} = \dfrac{0}{4\sqrt{2} \sin\frac{3\pi}{4}} = 0 \quad \Rightarrow \quad \theta = \dfrac{\pi}{2}$ [since $y > 0$]. Thus, spherical coordinates are $\left(4\sqrt{2}, \frac{\pi}{2}, \frac{3\pi}{4}\right)$.

(b) $\rho = \sqrt{(-2)^2 + 2^2 + (2\sqrt{6})^2} = 4\sqrt{2}$, $\cos\phi = \dfrac{z}{\rho} = \dfrac{2\sqrt{6}}{4\sqrt{2}} = \dfrac{\sqrt{3}}{2} \quad \Rightarrow \quad \phi = \frac{\pi}{6}$, and

$\cos\theta = \dfrac{x}{\rho \sin\phi} = \dfrac{-2}{4\sqrt{2} \sin\frac{\pi}{6}} = -\dfrac{1}{\sqrt{2}} \quad \Rightarrow \quad \theta = \frac{3\pi}{4}$ [since $x < 0$ and $y > 0$]. Thus, spherical coordinates are

$\left(4\sqrt{2}, \frac{3\pi}{4}, \frac{\pi}{6}\right)$.

5. Since $\phi = \frac{3\pi}{4}$, but ρ and θ can vary, the surface is the bottom half of a right circular cone with vertex at the origin and axis the negative z-axis. (See Figure 4.)

6. $\rho^2 - 3\rho + 2 = 0 \quad \Rightarrow \quad (\rho - 1)(\rho - 2) = 0 \quad \Rightarrow \quad \rho = 1$ or $\rho = 2$. Thus the equation represents two surfaces. In the case $\rho = 1$, the distance from any point to the origin is 1. Because θ and ϕ can vary, the surface is a sphere centered at the origin with radius 1. (See Figure 2.) Similarly, $\rho = 2$ is a sphere centered at the origin with radius 2.

Also, $\rho = 1 \quad \Rightarrow \quad \rho^2 = 1 \quad \Rightarrow \quad x^2 + y^2 + z^2 = 1$ which we recognize as the equation of the unit sphere, and similarly, $\rho = 2 \quad \Rightarrow \quad \rho^2 = 4 \quad \Rightarrow \quad x^2 + y^2 + z^2 = 4$.

7. From Equations 1 we have $z = \rho \cos\phi$, so $\rho \cos\phi = 1 \quad \Leftrightarrow \quad z = 1$, and the surface is the horizontal plane $z = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $\rho = \cos\phi \quad \Rightarrow \quad \rho^2 = \rho\cos\phi \quad \Leftrightarrow \quad x^2 + y^2 + z^2 = z \quad \Leftrightarrow \quad x^2 + y^2 + z^2 - z + \frac{1}{4} = \frac{1}{4} \quad \Leftrightarrow \quad x^2 + y^2 + (z - \frac{1}{2})^2 = \frac{1}{4}$.

Therefore, the surface is a sphere of radius $\frac{1}{2}$ centered at $\left(0, 0, \frac{1}{2}\right)$.

9. (a) From Equation 2 we have $\rho^2 = x^2 + y^2 + z^2$, so $x^2 + y^2 + z^2 = 9 \quad \Leftrightarrow \quad \rho^2 = 9 \quad \Rightarrow \quad \rho = 3$ (since $\rho \geq 0$).

(b) From Equations 1 we have $x = \rho\sin\phi\cos\theta$, $y = \rho\sin\phi\sin\theta$, and $z = \rho\cos\phi$, so the equation $x^2 - y^2 - z^2 = 1$ becomes $(\rho\sin\phi\cos\theta)^2 - (\rho\sin\phi\sin\theta)^2 - (\rho\cos\phi)^2 = 1 \quad \Leftrightarrow \quad (\rho^2\sin^2\phi)(\cos^2\theta - \sin^2\theta) - \rho^2\cos^2\phi = 1 \quad \Leftrightarrow$ $\rho^2(\sin^2\phi\cos 2\theta - \cos^2\phi) = 1$.

10. (a) $x = \rho\sin\phi\cos\theta$, $y = \rho\sin\phi\sin\theta$, and $z = \rho\cos\phi$, so the equation $z = x^2 + y^2$ becomes $\rho\cos\phi = (\rho\sin\phi\cos\theta)^2 + (\rho\sin\phi\sin\theta)^2$ or $\rho\cos\phi = \rho^2\sin^2\phi$. If $\rho \neq 0$, this becomes $\cos\phi = \rho\sin^2\phi$ or $\rho = \cos\phi\csc^2\phi$ or $\rho = \cot\phi\csc\phi$. ($\rho = 0$ corresponds to the origin which is included in the surface.)

(b) The equation $z = x^2 - y^2$ becomes $\rho\cos\phi = (\rho\sin\phi\cos\theta)^2 - (\rho\sin\phi\sin\theta)^2$ or $\rho\cos\phi = \rho^2(\sin^2\phi)(\cos^2\theta - \sin^2\theta) \quad \Leftrightarrow \quad \rho\cos\phi = \rho^2\sin^2\phi\cos 2\theta$. If $\rho \neq 0$, this becomes $\cos\phi = \rho\sin^2\phi\cos 2\theta$. ($\rho = 0$ corresponds to the origin which is included in the surface.)

11. $\rho \leq 1$ represents the (solid) unit ball. $0 \leq \phi \leq \frac{\pi}{6}$ restricts the solid to that portion on or above the cone $\phi = \frac{\pi}{6}$, and $0 \leq \theta \leq \pi$ further restricts the solid to that portion on or to the right of the xz-plane.

12. $1 \leq \rho \leq 2$ represents the solid region between and including the spheres of radii 1 and 2, centered at the origin. $\frac{\pi}{2} \leq \phi \leq \pi$ restricts the solid to that portion on or below the xy-plane.

13. $1 \leq \rho \leq 3$ represents the solid region between and including the spheres of radii 1 and 3, centered at the origin. $0 \leq \phi \leq \frac{\pi}{2}$ restricts the solid to that portion on or above the xy-plane. $\pi \leq \theta \leq \frac{3\pi}{2}$ further restricts the solid to the portion over the third quadrant.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. $\rho \le 2$ represents the solid sphere of radius 2 centered at the origin. Notice that $x^2 + y^2 = (\rho \sin\phi \cos\theta)^2 + (\rho \sin\phi \sin\theta)^2 = \rho^2 \sin^2\phi$. Then $\rho = \csc\phi \;\Rightarrow\; \rho \sin\phi = 1 \;\Rightarrow\; \rho^2 \sin^2\phi = x^2 + y^2 = 1$, so $\rho \le \csc\phi$ restricts the solid to that portion on or inside the circular cylinder $x^2 + y^2 = 1$.

15. $x^2 + y^2 + z^2 = 4z \;\Leftrightarrow\; x^2 + y^2 + z^2 - 4z + 4 = 4 \;\Leftrightarrow\; x^2 + y^2 + (z-2)^2 = 2^2$, which is a sphere with radius 2 centered at $(0, 0, 2)$. In spherical coordinates, we have $\rho^2 = 4\rho\cos\phi \;\Leftrightarrow\; \rho^2 - 4\rho\cos\phi = 0 \;\Leftrightarrow\; \rho = 0$ or $\rho = 4\cos\phi$, so "inside the sphere" is described by $0 \le \rho \le 4\cos\phi$. The cone $z = \sqrt{x^2 + y^2}$ (see Figure 15.7.13) is described by $\phi = \frac{\pi}{4}$, so "outside the cone" is described by $\frac{\pi}{4} \le \phi \le \frac{\pi}{2}$.

16. (a) The hollow ball is a spherical shell with outer radius 15 cm and inner radius 14.5 cm. If we center the ball at the origin of the coordinate system and use centimeters as the unit of measurement, then spherical coordinates conveniently describe the hollow ball as $14.5 \le \rho \le 15$, $0 \le \theta \le 2\pi$, $0 \le \phi \le \pi$.

(b) If we position the ball as in part (a), one possibility is to take the half of the ball that is above the xy-plane. This restricts ϕ from 0 to $\pi/2$ and the hemisphere can be described by $14.5 \le \rho \le 15$, $0 \le \theta \le 2\pi$, $0 \le \phi \le \pi/2$.

17.

The region of integration is given in spherical coordinates by $E = \{(\rho, \theta, \phi) \mid 0 \le \rho \le 3,\ 0 \le \theta \le \pi/2,\ 0 \le \phi \le \pi/6\}$. This represents the solid region in the first octant bounded above by the sphere $\rho = 3$ and below by the cone $\phi = \pi/6$.

$$\begin{aligned}\int_0^{\pi/6}\int_0^{\pi/2}\int_0^3 \rho^2 \sin\phi \, d\rho\, d\theta\, d\phi &= \int_0^{\pi/6} \sin\phi\, d\phi \int_0^{\pi/2} d\theta \int_0^3 \rho^2\, d\rho \\ &= \left[-\cos\phi\right]_0^{\pi/6} \left[\theta\right]_0^{\pi/2} \left[\tfrac{1}{3}\rho^3\right]_0^3 \\ &= \left(1 - \frac{\sqrt{3}}{2}\right)\left(\frac{\pi}{2}\right)(9) = \frac{9\pi}{4}\left(2 - \sqrt{3}\right)\end{aligned}$$

18. The region of integration is given in spherical coordinates by $E = \{(\rho, \theta, \phi) \mid 0 \le \rho \le \sec\phi,\ 0 \le \theta \le 2\pi,\ 0 \le \phi \le \pi/4\}$.

$\rho = \sec\phi \;\Leftrightarrow\; \rho\cos\phi = 1 \;\Leftrightarrow\; z = 1$, so E is the solid region above the cone $\phi = \pi/4$ and below the plane $z = 1$.

$$\begin{aligned}\int_0^{\pi/4}\int_0^{2\pi}\int_0^{\sec\phi} \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi &= \int_0^{\pi/4}\int_0^{2\pi} \left[\tfrac{1}{3}\rho^3 \sin\phi\right]_{\rho=0}^{\rho=\sec\phi} d\theta\, d\phi \\ &= \int_0^{\pi/4}\int_0^{2\pi} \tfrac{1}{3}\sec^3\phi \sin\phi\, d\theta\, d\phi \\ &= \tfrac{1}{3}\int_0^{\pi/4} \sec^3\phi \sin\phi\, d\phi \int_0^{2\pi} d\theta = \tfrac{1}{3}\int_0^{\pi/4} \tan\phi \sec^2\phi\, d\phi \int_0^{2\pi} d\theta \\ &= \tfrac{1}{3}\left[\tfrac{1}{2}\tan^2\phi\right]_0^{\pi/4} \left[\theta\right]_0^{2\pi} = \tfrac{1}{3}\left(\tfrac{1}{2} - 0\right)(2\pi) = \tfrac{\pi}{3}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. The solid E is most conveniently described if we use cylindrical coordinates:

$E = \left\{(r, \theta, z) \mid 0 \le \theta \le \frac{\pi}{2},\ 0 \le r \le 3,\ 0 \le z \le 2\right\}$. Then

$\iiint_E f(x, y, z)\, dV = \int_0^{\pi/2} \int_0^3 \int_0^2 f(r\cos\theta, r\sin\theta, z)\, r\, dz\, dr\, d\theta.$

20. The solid E is most conveniently described if we use spherical coordinates:

$E = \left\{(\rho, \theta, \phi) \mid 1 \le \rho \le 2,\ \frac{\pi}{2} \le \theta \le 2\pi,\ 0 \le \phi \le \frac{\pi}{2}\right\}$. Then

$\iiint_E f(x, y, z)\, dV = \int_0^{\pi/2} \int_{\pi/2}^{2\pi} \int_1^2 f(\rho\sin\phi\cos\theta, \rho\sin\phi\sin\theta, \rho\cos\phi)\, \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi.$

21. (a) The solid can be described in spherical coordinates by $E = \{(\rho, \theta, \phi) \mid 2 \le \rho \le 3, \frac{\pi}{2} \le \theta \le \frac{3\pi}{2}, \frac{\pi}{2} \le \phi \le \pi\}$.

Thus, $\iiint_E \sqrt{x^2 + y^2 + z^2}\, dV = \int_{\pi/2}^{\pi} \int_{\pi/2}^{3\pi/2} \int_2^3 \rho \cdot \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi.$

(b) $\int_{\pi/2}^{\pi} \int_{\pi/2}^{3\pi/2} \int_2^3 \rho^3 \sin\phi\, d\rho\, d\theta\, d\phi = \int_{\pi/2}^{\pi} \sin\phi\, d\phi \int_{\pi/2}^{3\pi/2} d\theta \int_2^3 \rho^3\, d\rho$

$$= \Big[-\cos\phi\Big]_{\phi=\pi/2}^{\phi=\pi} \Big[\theta\Big]_{\theta=\pi/2}^{\theta=3\pi/2} \left[\frac{\rho^4}{4}\right]_{\rho=2}^{\rho=3} = (1)(\pi) \cdot \frac{1}{4}(81 - 16) = \frac{65\pi}{4}$$

22. (a) The solid can be described in spherical coordinates by $E = \{(\rho, \theta, \phi) \mid 0 \le \rho \le 2\sqrt{2}, 0 \le \theta \le 2\pi, 0 \le \phi \le \frac{\pi}{4}\}$.

Thus, $\iiint_E xy\, dV = \int_0^{\pi/4} \int_0^{2\pi} \int_0^{2\sqrt{2}} \rho\sin\phi\cos\theta \cdot \rho\sin\phi\sin\theta \cdot \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi.$

(b) $\int_0^{\pi/4} \int_0^{2\pi} \int_0^{2\sqrt{2}} \rho^4 \sin^3\phi\cos\theta\sin\theta\, d\rho\, d\theta\, d\phi = \int_0^{\pi/4} \sin^3\phi\, d\phi \int_0^{2\pi} \cos\theta\sin\theta\, d\theta \int_0^{2\sqrt{2}} \rho^4\, d\rho$. Since

$\int_0^{2\pi} \cos\theta\sin\theta\, d\theta = \frac{1}{2}\int_0^{2\pi} \sin 2\theta\, d\theta = \frac{1}{2}\left[-\frac{1}{2}\cos 2\theta\right]_0^{2\pi} = -\frac{1}{4}(1 - 1) = 0$, the original iterated integral equals 0.

23. In spherical coordinates, B is represented by $\{(\rho, \theta, \phi) \mid 0 \le \rho \le 5, 0 \le \theta \le 2\pi, 0 \le \phi \le \pi\}$. Thus

$$\begin{aligned}
\iiint_B (x^2 + y^2 + z^2)^2\, dV &= \int_0^{\pi} \int_0^{2\pi} \int_0^5 (\rho^2)^2 \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi = \int_0^{\pi} \sin\phi\, d\phi \int_0^{2\pi} d\theta \int_0^5 \rho^6\, d\rho \\
&= \big[-\cos\phi\big]_0^{\pi} \big[\theta\big]_0^{2\pi} \left[\tfrac{1}{7}\rho^7\right]_0^5 = (2)(2\pi)\left(\tfrac{78{,}125}{7}\right) \\
&= \tfrac{312{,}500}{7}\pi \approx 140{,}249.7
\end{aligned}$$

24. In spherical coordinates, E is represented by $\left\{(\rho, \theta, \phi) \,\middle|\, 0 \le \rho \le 1,\ 0 \le \theta \le 2\pi,\ 0 \le \phi \le \frac{\pi}{3}\right\}$. Thus

$$\begin{aligned}
\iiint_E y^2 z^2\, dV &= \int_0^{\pi/3} \int_0^{2\pi} \int_0^1 (\rho\sin\phi\sin\theta)^2 (\rho\cos\phi)^2\, \rho^2 \sin\phi\, d\rho\, d\theta\, d\phi \\
&= \int_0^{\pi/3} \sin^3\phi\cos^2\phi\, d\phi \int_0^{2\pi} \sin^2\theta\, d\theta \int_0^1 \rho^6\, d\rho \\
&= \int_0^{\pi/3} (1 - \cos^2\phi)\cos^2\phi\sin\phi\, d\phi \int_0^{2\pi} \tfrac{1}{2}(1 - \cos 2\theta)\, d\theta \int_0^1 \rho^6\, d\rho \\
&= \left[\tfrac{1}{5}\cos^5\phi - \tfrac{1}{3}\cos^3\phi\right]_0^{\pi/3} \left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi} \left[\tfrac{1}{7}\rho^7\right]_0^1 \\
&= \left[\tfrac{1}{5}\left(\tfrac{1}{2}\right)^5 - \tfrac{1}{3}\left(\tfrac{1}{2}\right)^3 - \tfrac{1}{5} + \tfrac{1}{3}\right](\pi - 0)\left(\tfrac{1}{7} - 0\right) = \tfrac{47}{480} \cdot \pi \cdot \tfrac{1}{7} = \tfrac{47}{3360}\pi
\end{aligned}$$

25. In spherical coordinates, E is represented by $\{(\rho, \theta, \phi) \mid 2 \le \rho \le 3, 0 \le \theta \le 2\pi, 0 \le \phi \le \pi\}$ and

$x^2 + y^2 = \rho^2 \sin^2\phi\cos^2\theta + \rho^2 \sin^2\phi\sin^2\theta = \rho^2 \sin^2\phi\left(\cos^2\theta + \sin^2\theta\right) = \rho^2 \sin^2\phi$. Thus

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\iiint_E (x^2+y^2)\,dV = \int_0^{\pi}\int_0^{2\pi}\int_2^3 (\rho^2\sin^2\phi)\,\rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \int_0^{\pi}\sin^3\phi\,d\phi\int_0^{2\pi}d\theta\int_2^3\rho^4\,d\rho$$
$$= \int_0^{\pi}(1-\cos^2\phi)\,\sin\phi\,d\phi\,\left[\,\theta\,\right]_0^{2\pi}\left[\tfrac{1}{5}\rho^5\right]_2^3 = \left[-\cos\phi+\tfrac{1}{3}\cos^3\phi\right]_0^{\pi}(2\pi)\cdot\tfrac{1}{5}(243-32)$$
$$= \left(1-\tfrac{1}{3}+1-\tfrac{1}{3}\right)(2\pi)\left(\tfrac{211}{5}\right) = \tfrac{1688\pi}{15}$$

26. In spherical coordinates, E is represented by $\{(\rho,\theta,\phi)\,|\,0\le\rho\le 3, 0\le\theta\le\pi, 0\le\phi\le\pi\}$. Thus

$$\iiint_E y^2\,dV = \int_0^{\pi}\int_0^{\pi}\int_0^3(\rho\sin\phi\sin\theta)^2\,\rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \int_0^{\pi}\sin^3\phi\,d\phi\int_0^{\pi}\sin^2\theta\,d\theta\int_0^3\rho^4\,d\rho$$
$$= \int_0^{\pi}(1-\cos^2\phi)\,\sin\phi\,d\phi\int_0^{\pi}\tfrac{1}{2}(1-\cos 2\theta)\,d\theta\int_0^3\rho^4\,d\rho$$
$$= \left[-\cos\phi+\tfrac{1}{3}\cos^3\phi\right]_0^{\pi}\left[\tfrac{1}{2}\left(\theta-\tfrac{1}{2}\sin 2\theta\right)\right]_0^{\pi}\left[\tfrac{1}{5}\rho^5\right]_0^3$$
$$= \left(\tfrac{2}{3}+\tfrac{2}{3}\right)\left(\tfrac{1}{2}\pi\right)\left(\tfrac{1}{5}(243)\right) = \left(\tfrac{4}{3}\right)\left(\tfrac{\pi}{2}\right)\left(\tfrac{243}{5}\right) = \tfrac{162\pi}{5}$$

27. In spherical coordinates, E is represented by $\left\{(\rho,\theta,\phi)\,\middle|\,0\le\rho\le 1, 0\le\theta\le\tfrac{\pi}{2}, 0\le\phi\le\tfrac{\pi}{2}\right\}$. Thus

$$\iiint_E xe^{x^2+y^2+z^2}\,dV = \int_0^{\pi/2}\int_0^{\pi/2}\int_0^1(\rho\sin\phi\cos\theta)e^{\rho^2}\rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \int_0^{\pi/2}\sin^2\phi\,d\phi\int_0^{\pi/2}\cos\theta\,d\theta\int_0^1\rho^3e^{\rho^2}\,d\rho$$
$$= \int_0^{\pi/2}\tfrac{1}{2}(1-\cos 2\phi)\,d\phi\int_0^{\pi/2}\cos\theta\,d\theta\left(\left.\tfrac{1}{2}\rho^2e^{\rho^2}\right]_0^1-\int_0^1\rho e^{\rho^2}\,d\rho\right)$$
$$\left[\text{integrate by parts with } u=\rho^2,\ dv=\rho e^{\rho^2}d\rho\right]$$
$$= \left[\tfrac{1}{2}\phi-\tfrac{1}{4}\sin 2\phi\right]_0^{\pi/2}\left[\sin\theta\right]_0^{\pi/2}\left[\tfrac{1}{2}\rho^2e^{\rho^2}-\tfrac{1}{2}e^{\rho^2}\right]_0^1 = \left(\tfrac{\pi}{4}-0\right)(1-0)\left(0+\tfrac{1}{2}\right) = \tfrac{\pi}{8}$$

28. In spherical coordinates, the cone $z=\sqrt{x^2+y^2}$ is equivalent to $\phi=\pi/4$ (as in Example 4) and E is represented by $\{(\rho,\theta,\phi)\,|\,1\le\rho\le 2,\ 0\le\theta\le 2\pi,\ 0\le\phi\le\pi/4\}$. Also $\sqrt{x^2+y^2+z^2}=\sqrt{\rho^2}=\rho$, so

$$\iiint_E\sqrt{x^2+y^2+z^2}\,dV = \int_0^{\pi/4}\int_0^{2\pi}\int_1^2\rho\cdot\rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \int_0^{\pi/4}\sin\phi\,d\phi\int_0^{2\pi}d\theta\int_1^2\rho^3\,d\rho$$
$$= \left[-\cos\phi\right]_0^{\pi/4}\left[\,\theta\,\right]_0^{2\pi}\left[\tfrac{1}{4}\rho^4\right]_1^2 = \left(-\tfrac{\sqrt{2}}{2}+1\right)(2\pi)\cdot\tfrac{1}{4}(16-1) = \tfrac{15}{2}\pi\left(1-\tfrac{\sqrt{2}}{2}\right)$$

29. The solid region is given by $E=\left\{(\rho,\theta,\phi)\mid 0\le\rho\le a, 0\le\theta\le 2\pi, \tfrac{\pi}{6}\le\phi\le\tfrac{\pi}{3}\right\}$ and its volume is

$$V = \iiint_E dV = \int_{\pi/6}^{\pi/3}\int_0^{2\pi}\int_0^a\rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \int_{\pi/6}^{\pi/3}\sin\phi\,d\phi\int_0^{2\pi}d\theta\int_0^a\rho^2\,d\rho$$
$$= \left[-\cos\phi\right]_{\pi/6}^{\pi/3}\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{3}\rho^3\right]_0^a = \left(-\tfrac{1}{2}+\tfrac{\sqrt{3}}{2}\right)(2\pi)\left(\tfrac{1}{3}a^3\right) = \tfrac{\sqrt{3}-1}{3}\pi a^3$$

30. If we center the ball at the origin, then the ball is given by

$B=\{(\rho,\theta,\phi)\mid 0\le\rho\le a, 0\le\theta\le 2\pi, 0\le\phi\le\pi\}$ and the distance from any point (x,y,z) in the ball to the center $(0,0,0)$ is $\sqrt{x^2+y^2+z^2}=\rho$. Thus the average distance is

$$\frac{1}{V(B)}\iiint_B\rho\,dV = \frac{1}{\frac{4}{3}\pi a^3}\int_0^{\pi}\int_0^{2\pi}\int_0^a\rho\cdot\rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \frac{3}{4\pi a^3}\int_0^{\pi}\sin\phi\,d\phi\int_0^{2\pi}d\theta\int_0^a\rho^3\,d\rho$$
$$= \frac{3}{4\pi a^3}\left[-\cos\phi\right]_0^{\pi}\left[\,\theta\,\right]_0^{2\pi}\left[\tfrac{1}{4}\rho^4\right]_0^a = \frac{3}{4\pi a^3}(2)(2\pi)\left(\tfrac{1}{4}a^4\right) = \tfrac{3}{4}a$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. (a) Since $\rho = 4\cos\phi$ implies $\rho^2 = 4\rho\cos\phi \quad \Leftrightarrow \quad x^2+y^2+z^2 = 4z \quad \Leftrightarrow \quad x^2+y^2+(z-2)^2 = 4$, the equation is that of a sphere of radius 2 with center at $(0,0,2)$. Thus

$$V = \int_0^{2\pi}\int_0^{\pi/3}\int_0^{4\cos\phi} \rho^2\sin\phi\,d\rho\,d\phi\,d\theta = \int_0^{2\pi}\int_0^{\pi/3}\left[\tfrac{1}{3}\rho^3\right]_{\rho=0}^{\rho=4\cos\phi}\sin\phi\,d\phi\,d\theta = \int_0^{2\pi}\int_0^{\pi/3}\left(\tfrac{64}{3}\cos^3\phi\right)\sin\phi\,d\phi\,d\theta$$
$$= \int_0^{2\pi}\left[-\tfrac{16}{3}\cos^4\phi\right]_{\phi=0}^{\phi=\pi/3}d\theta = \int_0^{2\pi} -\tfrac{16}{3}\left(\tfrac{1}{16}-1\right)d\theta = 5\theta\Big]_0^{2\pi} = 10\pi$$

(b) By the symmetry of the problem $M_{yz} = M_{xz} = 0$. Then

$$M_{xy} = \int_0^{2\pi}\int_0^{\pi/3}\int_0^{4\cos\phi}\rho^3\cos\phi\sin\phi\,d\rho\,d\phi\,d\theta = \int_0^{2\pi}\int_0^{\pi/3}\cos\phi\sin\phi\left(64\cos^4\phi\right)d\phi\,d\theta$$
$$= \int_0^{2\pi}64\left[-\tfrac{1}{6}\cos^6\phi\right]_{\phi=0}^{\phi=\pi/3}d\theta = \int_0^{2\pi}\tfrac{21}{2}\,d\theta = 21\pi$$

Hence, $(\overline{x},\overline{y},\overline{z}) = (0,0,21\pi/(10\pi)) = (0,0,2.1)$.

32. In spherical coordinates, the sphere $x^2+y^2+z^2=4$ is equivalent to $\rho=2$ and the cone $z=\sqrt{x^2+y^2}$ is represented by $\phi = \frac{\pi}{4}$ (as in Example 4). Thus, the solid is given by $\left\{(\rho,\theta,\phi)\,\middle|\,0\le\rho\le 2,\ 0\le\theta\le 2\pi,\ \frac{\pi}{4}\le\phi\le\frac{\pi}{2}\right\}$ and

$$V = \int_{\pi/4}^{\pi/2}\int_0^{2\pi}\int_0^2 \rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \int_{\pi/4}^{\pi/2}\sin\phi\,d\phi\int_0^{2\pi}d\theta\int_0^2\rho^2\,d\rho$$
$$= \left[-\cos\phi\right]_{\pi/4}^{\pi/2}\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{3}\rho^3\right]_0^2 = \left(\tfrac{\sqrt{2}}{2}\right)(2\pi)\left(\tfrac{8}{3}\right) = \tfrac{8\sqrt{2}\,\pi}{3}$$

33. (a) By the symmetry of the region, $M_{yz}=0$ and $M_{xz}=0$. Assuming constant density K, $m = \iiint_E K\,dV = K\iiint_E dV = \frac{\pi}{8}K$ (from Example 4). Then

$$M_{xy} = \iiint_E z\,K\,dV = K\int_0^{2\pi}\int_0^{\pi/4}\int_0^{\cos\phi}(\rho\cos\phi)\,\rho^2\sin\phi\,d\rho\,d\phi\,d\theta = K\int_0^{2\pi}\int_0^{\pi/4}\sin\phi\cos\phi\left[\tfrac{1}{4}\rho^4\right]_{\rho=0}^{\rho=\cos\phi}d\phi\,d\theta$$
$$= \tfrac{1}{4}K\int_0^{2\pi}\int_0^{\pi/4}\sin\phi\cos\phi\left(\cos^4\phi\right)d\phi\,d\theta = \tfrac{1}{4}K\int_0^{2\pi}d\theta\int_0^{\pi/4}\cos^5\phi\sin\phi\,d\phi$$
$$= \tfrac{1}{4}K\left[\theta\right]_0^{2\pi}\left[-\tfrac{1}{6}\cos^6\phi\right]_0^{\pi/4} = \tfrac{1}{4}K(2\pi)\left(-\tfrac{1}{6}\right)\left[\left(\tfrac{\sqrt{2}}{2}\right)^6-1\right] = -\tfrac{\pi}{12}K\left(-\tfrac{7}{8}\right) = \tfrac{7\pi}{96}K$$

Thus, the centroid is $(\overline{x},\overline{y},\overline{z}) = \left(\dfrac{M_{yz}}{m},\dfrac{M_{xz}}{m},\dfrac{M_{xy}}{m}\right) = \left(0,0,\dfrac{7\pi K/96}{\pi K/8}\right) = \left(0,0,\tfrac{7}{12}\right)$.

(b) As in Exercise 25, $x^2+y^2 = \rho^2\sin^2\phi$ and

$$I_z = \iiint_E (x^2+y^2)\,K\,dV = K\int_0^{2\pi}\int_0^{\pi/4}\int_0^{\cos\phi}(\rho^2\sin^2\phi)\,\rho^2\sin\phi\,d\rho\,d\phi\,d\theta = K\int_0^{2\pi}\int_0^{\pi/4}\sin^3\phi\left[\tfrac{1}{5}\rho^5\right]_{\rho=0}^{\rho=\cos\phi}d\phi\,d\theta$$
$$= \tfrac{1}{5}K\int_0^{2\pi}\int_0^{\pi/4}\sin^3\phi\cos^5\phi\,d\phi\,d\theta = \tfrac{1}{5}K\int_0^{2\pi}d\theta\int_0^{\pi/4}\cos^5\phi\left(1-\cos^2\phi\right)\sin\phi\,d\phi$$
$$= \tfrac{1}{5}K\left[\theta\right]_0^{2\pi}\left[-\tfrac{1}{6}\cos^6\phi+\tfrac{1}{8}\cos^8\phi\right]_0^{\pi/4}$$
$$= \tfrac{1}{5}K(2\pi)\left[-\tfrac{1}{6}\left(\tfrac{\sqrt{2}}{2}\right)^6+\tfrac{1}{8}\left(\tfrac{\sqrt{2}}{2}\right)^8+\tfrac{1}{6}-\tfrac{1}{8}\right] = \tfrac{2\pi}{5}K\left(\tfrac{11}{384}\right) = \tfrac{11\pi}{960}K$$

34. (a) Placing the center of the base at $(0,0,0)$, $\rho(x,y,z) = K\sqrt{x^2+y^2+z^2}$ is the density function. So

$$m = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a K\rho^3\sin\phi\,d\rho\,d\phi\,d\theta = K\int_0^{2\pi}d\theta\int_0^{\pi/2}\sin\phi\,d\phi\int_0^a\rho^3\,d\rho$$
$$= K\left[\theta\right]_0^{2\pi}\left[-\cos\phi\right]_0^{\pi/2}\left[\tfrac{1}{4}\rho^4\right]_0^a = K(2\pi)(1)\left(\tfrac{1}{4}a^4\right) = \tfrac{1}{2}\pi K a^4$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) By the symmetry of the problem $M_{yz} = M_{xz} = 0$. Then

$$\begin{aligned} M_{xy} &= \int_0^{2\pi}\int_0^{\pi/2}\int_0^a K\rho^4 \sin\phi\cos\phi\, d\rho\, d\phi\, d\theta \\ &= K\int_0^{2\pi} d\theta \int_0^{\pi/2} \sin\phi\cos\phi\, d\phi \int_0^a \rho^4\, d\rho \\ &= K\big[\,\theta\,\big]_0^{2\pi}\ \big[\tfrac{1}{2}\sin^2\phi\big]_0^{\pi/2}\ \big[\tfrac{1}{5}\rho^5\big]_0^a = K(2\pi)\big(\tfrac{1}{2}\big)\big(\tfrac{1}{5}a^5\big) = \tfrac{1}{5}\pi K a^5 \end{aligned}$$

Hence, $(\overline{x}, \overline{y}, \overline{z}) = \big(0, 0, \frac{2}{5}a\big)$.

(c) $I_z = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a (K\rho^3\sin\phi)(\rho^2\sin^2\phi)\, d\rho\, d\phi\, d\theta = K\int_0^{2\pi} d\theta \int_0^{\pi/2}\sin^3\phi\, d\phi \int_0^a \rho^5\, d\rho$

$= K\big[\,\theta\,\big]_0^{2\pi}\ \big[-\cos\phi + \frac{1}{3}\cos^3\phi\big]_0^{\pi/2}\ \big[\frac{1}{6}\rho^6\big]_0^a = K(2\pi)\big(\frac{2}{3}\big)\big(\frac{1}{6}a^6\big) = \frac{2}{9}\pi K a^6$

35. (a) The density function is $\rho(x, y, z) = K$, a constant, and by the symmetry of the problem $M_{xz} = M_{yz} = 0$. Then $M_{xy} = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a K\rho^3\sin\phi\,\cos\phi\, d\rho\, d\phi\, d\theta = \frac{1}{2}\pi K a^4 \int_0^{\pi/2}\sin\phi\,\cos\phi\, d\phi = \frac{1}{8}\pi K a^4$. But the mass is $K \cdot (\text{volume of the hemisphere}) = \frac{2}{3}\pi K a^3$, so the centroid is $\big(0, 0, \frac{3}{8}a\big)$.

(b) Place the center of the base at $(0, 0, 0)$; the density function is $\rho(x, y, z) = K$. By symmetry, the moments of inertia about any two such diameters will be equal, so we just need to find I_x:

$$\begin{aligned} I_x &= \int_0^{2\pi}\int_0^{\pi/2}\int_0^a (K\rho^2\sin\phi)\,\rho^2\,(\sin^2\phi\,\sin^2\theta + \cos^2\phi)\, d\rho\, d\phi\, d\theta \\ &= K\int_0^{2\pi}\int_0^{\pi/2}(\sin^3\phi\,\sin^2\theta + \sin\phi\,\cos^2\phi)\big(\tfrac{1}{5}a^5\big)\, d\phi\, d\theta \\ &= \tfrac{1}{5}Ka^5\int_0^{2\pi}\Big[\sin^2\theta\,\big(-\cos\phi + \tfrac{1}{3}\cos^3\phi\big) + \big(-\tfrac{1}{3}\cos^3\phi\big)\Big]_{\phi=0}^{\phi=\pi/2}\, d\theta = \tfrac{1}{5}Ka^5\int_0^{2\pi}\big[\tfrac{2}{3}\sin^2\theta + \tfrac{1}{3}\big]\, d\theta \\ &= \tfrac{1}{5}Ka^5\big[\tfrac{2}{3}\big(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\big) + \tfrac{1}{3}\theta\big]_0^{2\pi} = \tfrac{1}{5}Ka^5\big[\tfrac{2}{3}(\pi - 0) + \tfrac{1}{3}(2\pi - 0)\big] = \tfrac{4}{15}\pi K a^5 \end{aligned}$$

36. Place the center of the base at $(0, 0, 0)$, then the density is $\rho(x, y, z) = Kz$, K a constant. Then

$m = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a (K\rho\cos\phi)\,\rho^2\sin\phi\, d\rho\, d\phi\, d\theta = 2\pi K\int_0^{\pi/2}\cos\phi\sin\phi\cdot\frac{1}{4}a^4\, d\phi = \frac{1}{2}\pi K a^4\big[-\frac{1}{4}\cos 2\phi\big]_0^{\pi/2} = \frac{\pi}{4}Ka^4$.

By the symmetry of the problem $M_{xz} = M_{yz} = 0$, and

$M_{xy} = \int_0^{2\pi}\int_0^{\pi/2}\int_0^a K\rho^4\cos^2\phi\sin\phi\, d\rho\, d\phi\, d\theta = \frac{2}{5}\pi K a^5\int_0^{\pi/2}\cos^2\phi\sin\phi\, d\phi = \frac{2}{5}\pi K a^5\big[-\frac{1}{3}\cos^3\theta\big]_0^{\pi/2} = \frac{2}{15}\pi K a^5$.

Hence, $(\overline{x}, \overline{y}, \overline{z}) = \big(0, 0, \frac{8}{15}a\big)$.

37. In spherical coordinates $z = \sqrt{x^2 + y^2}$ becomes $\phi = \frac{\pi}{4}$ (as in Example 4). Then

$V = \int_0^{2\pi}\int_0^{\pi/4}\int_0^1 \rho^2\sin\phi\, d\rho\, d\phi\, d\theta = \int_0^{2\pi} d\theta \int_0^{\pi/4}\sin\phi\, d\phi \int_0^1 \rho^2\, d\rho = 2\pi\left(-\frac{\sqrt{2}}{2} + 1\right)\big(\frac{1}{3}\big) = \frac{1}{3}\pi\big(2 - \sqrt{2}\big)$,

$M_{xy} = \int_0^{2\pi}\int_0^{\pi/4}\int_0^1 \rho^3\sin\phi\cos\phi\, d\rho\, d\phi\, d\theta = 2\pi\big[-\frac{1}{4}\cos 2\phi\big]_0^{\pi/4}\big(\frac{1}{4}\big) = \frac{\pi}{8}$ and by symmetry $M_{yz} = M_{xz} = 0$.

Hence, $(\overline{x}, \overline{y}, \overline{z}) = \left(0, 0, \dfrac{3}{8\big(2 - \sqrt{2}\big)}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. Place the center of the sphere at $(0,0,0)$, let the diameter of intersection be along the z-axis, one of the planes be the xz-plane and the other be the plane whose angle with the xz-plane is $\theta = \frac{\pi}{6}$. Then in spherical coordinates the volume is given by

$V = \int_0^{\pi/6}\int_0^{\pi}\int_0^{a} \rho^2 \sin\phi \, d\rho \, d\phi \, d\theta = \int_0^{\pi/6} d\theta \int_0^{\pi} \sin\phi \, d\phi \int_0^{a} \rho^2 \, d\rho = \frac{\pi}{6}(2)\left(\frac{1}{3}a^3\right) = \frac{1}{9}\pi a^3.$

39. (a) If we orient the cylinder so that its axis is the z-axis and its base lies in the xy-plane, then the cylinder is described, in cylindrical coordinates, by $E = \{(r,\theta,z) \mid 0 \le r \le a,\ 0 \le \theta \le 2\pi,\ 0 \le z \le h\}$. Assuming constant density K, the moment of inertia about its axis (the z-axis) is

$$\begin{aligned} I_z &= \iiint_E (x^2+y^2)\,\rho(x,y,z)\,dV = \int_0^{2\pi}\int_0^{a}\int_0^{h} K(r^2)\,r\,dz\,dr\,d\theta = K\int_0^{2\pi} d\theta \int_0^{a} r^3\,dr \int_0^{h} dz \\ &= K\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^{a}\left[z\right]_0^{h} = K\,(2\pi)\left(\tfrac{1}{4}a^4\right)(h) = \tfrac{1}{2}\pi K a^4 h \end{aligned}$$

(b) By symmetry, the moments of inertia about any two diameters of the base will be equal, and one of the diameters lies on the x-axis, so we compute:

$$\begin{aligned} I_x &= \iiint_E (y^2+z^2)\,\rho(x,y,z)\,dV = \int_0^{2\pi}\int_0^{a}\int_0^{h} K(r^2\sin^2\theta + z^2)\,r\,dz\,dr\,d\theta \\ &= K\int_0^{2\pi}\int_0^{a}\int_0^{h} r^3\sin^2\theta\,dz\,dr\,d\theta + K\int_0^{2\pi}\int_0^{a}\int_0^{h} rz^2\,dz\,dr\,d\theta \\ &= K\int_0^{2\pi}\sin^2\theta\,d\theta \int_0^{a} r^3\,dr \int_0^{h} dz + K\int_0^{2\pi} d\theta \int_0^{a} r\,dr \int_0^{h} z^2\,dz \\ &= K\left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^{a}\left[z\right]_0^{h} + K\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{2}r^2\right]_0^{a}\left[\tfrac{1}{3}z^3\right]_0^{h} \\ &= K\,(\pi)\left(\tfrac{1}{4}a^4\right)(h) + K\,(2\pi)\left(\tfrac{1}{2}a^2\right)\left(\tfrac{1}{3}h^3\right) = \tfrac{1}{12}\pi K a^2 h(3a^2+4h^2) \end{aligned}$$

40.

Orient the cone so that its axis is the z-axis and its base lies in the xy-plane, as shown in the figure. (Then the z-axis is the axis of the cone and the x-axis contains a diameter of the base.) A right circular cone with axis the z-axis and vertex at the origin has equation $z^2 = c^2(x^2+y^2)$. Here we have the bottom frustum, shifted upward h units, and with $c^2 = h^2/a^2$ so that the cone includes the point $(a,0,0)$. Thus an equation of the cone in rectangular coordinates is $z = h - \frac{h}{a}\sqrt{x^2+y^2}$, $0 \le z \le h$. In cylindrical coordinates, the cone is described by

$$E = \left\{(r,\theta,z) \mid 0 \le r \le a,\ 0 \le \theta \le 2\pi,\ 0 \le z \le h\left(1 - \tfrac{1}{a}r\right)\right\}$$

(a) Assuming constant density K, the moment of inertia about its axis (the z-axis) is

$$\begin{aligned} I_z &= \iiint_E (x^2+y^2)\,\rho(x,y,z)\,dV = \int_0^{2\pi}\int_0^{a}\int_0^{h(1-r/a)} K(r^2)\,r\,dz\,dr\,d\theta \\ &= K\int_0^{2\pi}\int_0^{a}\left[r^3 z\right]_{z=0}^{z=h(1-r/a)} dr\,d\theta = K\int_0^{2\pi}\int_0^{a} r^3 h\left(1-\tfrac{1}{a}r\right) dr\,d\theta \\ &= Kh\int_0^{2\pi} d\theta \int_0^{a}\left(r^3 - \tfrac{1}{a}r^4\right) dr = Kh\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4 - \tfrac{1}{5a}r^5\right]_0^{a} \\ &= Kh\,(2\pi)\left(\tfrac{1}{4}a^4 - \tfrac{1}{5}a^4\right) = \tfrac{1}{10}\pi K a^4 h \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) By symmetry, the moments of intertia about any two diameters of the base will be equal, and one of the diameters lies on the x-axis, so we compute:

$$\begin{aligned}
I_x &= \iiint_E (y^2+z^2)\,\rho(x,y,z)\,dV = \int_0^{2\pi}\int_0^a\int_0^{h(1-r/a)} K(r^2\sin^2\theta+z^2)\,r\,dz\,dr\,d\theta \\
&= K\int_0^{2\pi}\int_0^a \left[(r^3\sin^2\theta)z+\tfrac{1}{3}rz^3\right]_{z=0}^{z=h(1-r/a)} dr\,d\theta \\
&= K\int_0^{2\pi}\int_0^a \left[(r^3\sin^2\theta)\left(h\left(1-\tfrac{1}{a}r\right)\right)+\tfrac{1}{3}r\left(h\left(1-\tfrac{1}{a}r\right)\right)^3\right] dr\,d\theta \\
&= Kh\int_0^{2\pi}\int_0^a (r^3\sin^2\theta)\left(1-\tfrac{1}{a}r\right)\,dr\,d\theta + Kh^3\int_0^{2\pi}\int_0^a \tfrac{1}{3}r\left(1-\tfrac{1}{a}r\right)^3 dr\,d\theta \\
&= Kh\int_0^{2\pi}\sin^2\theta\,d\theta\int_0^a\left(r^3-\tfrac{1}{a}r^4\right)dr + \tfrac{1}{3}Kh^3\int_0^{2\pi}d\theta\int_0^a\left(r-\tfrac{3}{a}r^2+\tfrac{3}{a^2}r^3-\tfrac{1}{a^3}r^4\right)dr \\
&= Kh\left[\tfrac{1}{2}\theta-\tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4-\tfrac{1}{5a}r^5\right]_0^a + \tfrac{1}{3}Kh^3\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{2}r^2-\tfrac{1}{a}r^3+\tfrac{3}{4a^2}r^4-\tfrac{1}{5a^3}r^5\right]_0^a \\
&= Kh(\pi)\left(\tfrac{1}{4}a^4-\tfrac{1}{5}a^4\right)+\tfrac{1}{3}Kh^3(2\pi)\left(\tfrac{1}{2}a^2-a^2+\tfrac{3}{4}a^2-\tfrac{1}{5}a^2\right) \\
&= \pi Kh\left(\tfrac{1}{20}a^4\right)+\tfrac{2}{3}\pi Kh^3\left(\tfrac{1}{20}a^2\right) = \pi Ka^2h\left(\tfrac{1}{20}a^2+\tfrac{1}{30}h^2\right)
\end{aligned}$$

41. In cylindrical coordinates the paraboloid is given by $z=r^2$ and the plane by $z=2r\sin\theta$ and the projection of the intersection onto the xy-plane is the circle $r=2\sin\theta$. Then $\iiint_E z\,dV = \int_0^{\pi}\int_0^{2\sin\theta}\int_{r^2}^{2r\sin\theta} rz\,dz\,dr\,d\theta = \frac{5\pi}{6}$ [using a CAS].

42. (a) The region enclosed by the torus is $\{(\rho,\theta,\phi)\mid 0\le\theta\le 2\pi,\ 0\le\phi\le\pi,\ 0\le\rho\le\sin\phi\}$, so its volume is

$$V=\int_0^{2\pi}\int_0^{\pi}\int_0^{\sin\phi}\rho^2\sin\phi\,d\rho\,d\phi\,d\theta = 2\pi\int_0^{\pi}\tfrac{1}{3}\sin^4\phi\,d\phi = \tfrac{2}{3}\pi\left[\tfrac{3}{8}\phi-\tfrac{1}{4}\sin 2\phi+\tfrac{1}{16}\sin 4\phi\right]_0^{\pi} = \tfrac{1}{4}\pi^2.$$

(b) In Maple, we can plot the torus using the command

```
plot3d(sin(phi),theta=0..2*Pi,
       phi=0..Pi,coords=spherical);.
```

In Mathematica, use

```
SphericalPlot3D[Sin[phi],{phi,0,Pi},{theta,0,2Pi}].
```

43. $\displaystyle\int_0^1\int_0^{\sqrt{1-x^2}}\int_{\sqrt{x^2+y^2}}^{\sqrt{2-x^2-y^2}} xy\,dz\,dy\,dx$. The region E of integration is the region above the cone $z=\sqrt{x^2+y^2}$ and below the sphere $x^2+y^2+z^2=2$ in the first octant. Because E is in the first octant we have $0\le\theta\le\frac{\pi}{2}$. The cone has equation $\phi=\frac{\pi}{4}$ (as in Example 4), so $0\le\phi\le\frac{\pi}{4}$, and $0\le\rho\le\sqrt{2}$. Then the integral becomes

$$\int_0^{\pi/4}\int_0^{\pi/2}\int_0^{\sqrt{2}}(\rho\sin\phi\cos\theta)(\rho\sin\phi\sin\theta)\rho^2\sin\phi\,d\rho\,d\theta\,d\phi$$

$$\begin{aligned}
&= \int_0^{\pi/4}\sin^3\phi\,d\phi\int_0^{\pi/2}\sin\theta\cos\theta\,d\theta\int_0^{\sqrt{2}}\rho^4\,d\rho = \left(\int_0^{\pi/4}\left(1-\cos^2\phi\right)\sin\phi\,d\phi\right)\left[\tfrac{1}{2}\sin^2\theta\right]_0^{\pi/2}\left[\tfrac{1}{5}\rho^5\right]_0^{\sqrt{2}} \\
&= \left[\tfrac{1}{3}\cos^3\phi-\cos\phi\right]_0^{\pi/4}\cdot\tfrac{1}{2}\cdot\tfrac{1}{5}\left(\sqrt{2}\right)^5 = \left[\tfrac{\sqrt{2}}{12}-\tfrac{\sqrt{2}}{2}-\left(\tfrac{1}{3}-1\right)\right]\cdot\tfrac{2\sqrt{2}}{5} = \tfrac{4\sqrt{2}-5}{15}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

44. $\int_{-a}^{a}\int_{-\sqrt{a^2-y^2}}^{\sqrt{a^2-y^2}}\int_{-\sqrt{a^2-x^2-y^2}}^{\sqrt{a^2-x^2-y^2}}(x^2z+y^2z+z^3)\,dz\,dx\,dy$. The region of integration is the solid sphere $x^2+y^2+z^2\le a^2$, so $0\le\theta\le 2\pi$, $0\le\phi\le\pi$, and $0\le\rho\le a$. Also $x^2z+y^2z+z^3=(x^2+y^2+z^2)z=\rho^2z=\rho^3\cos\phi$, so the integral becomes

$$\int_0^{\pi}\int_0^{2\pi}\int_0^{a}\left(\rho^3\cos\phi\right)\rho^2\sin\phi\,d\rho\,d\theta\,d\phi=\int_0^{\pi}\sin\phi\cos\phi\,d\phi\int_0^{2\pi}d\theta\int_0^{a}\rho^5\,d\rho=\left[\tfrac{1}{2}\sin^2\phi\right]_0^{\pi}\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{6}\rho^6\right]_0^{a}=0$$

45. $\int_{-2}^{2}\int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}}\int_{2-\sqrt{4-x^2-y^2}}^{2+\sqrt{4-x^2-y^2}}(x^2+y^2+z^2)^{3/2}\,dz\,dy\,dx$. The region of integration is the solid sphere $x^2+y^2+(z-2)^2\le 4$, or equivalently, $\rho^2\sin^2\phi+(\rho\cos\phi-2)^2=\rho^2-4\rho\cos\phi+4\le 4\quad\Rightarrow\quad\rho\le 4\cos\phi$, so $0\le\theta\le 2\pi$, $0\le\phi\le\frac{\pi}{2}$, and $0\le\rho\le 4\cos\phi$. Also $(x^2+y^2+z^2)^{3/2}=(\rho^2)^{3/2}=\rho^3$, so the integral becomes

$$\begin{aligned}\int_0^{\pi/2}\int_0^{2\pi}\int_0^{4\cos\phi}\left(\rho^3\right)\rho^2\sin\phi\,d\rho\,d\theta\,d\phi&=\int_0^{\pi/2}\int_0^{2\pi}\sin\phi\left[\tfrac{1}{6}\rho^6\right]_{\rho=0}^{\rho=4\cos\phi}d\theta\,d\phi\\&=\tfrac{1}{6}\int_0^{\pi/2}\int_0^{2\pi}\sin\phi\left(4096\cos^6\phi\right)d\theta\,d\phi\\&=\tfrac{1}{6}(4096)\int_0^{\pi/2}\cos^6\phi\sin\phi\,d\phi\int_0^{2\pi}d\theta=\tfrac{2048}{3}\left[-\tfrac{1}{7}\cos^7\phi\right]_0^{\pi/2}\left[\theta\right]_0^{2\pi}\\&=\tfrac{2048}{3}\left(\tfrac{1}{7}\right)(2\pi)=\tfrac{4096\pi}{21}\end{aligned}$$

46. The solid region between the ground and an altitude of 5 km (5000 m) is given by

$$E=\left\{(\rho,\theta,\phi)\mid 6.370\times10^6\le\rho\le 6.375\times10^6,0\le\theta\le 2\pi,0\le\phi\le\pi\right\}.$$

Then the mass of the atmosphere in this region is

$$\begin{aligned}m=\iiint_E\delta\,dV&=\int_0^{2\pi}\int_0^{\pi}\int_{6.370\times10^6}^{6.375\times10^6}(619.09-0.000097\rho)\,\rho^2\sin\phi\,d\rho\,d\phi\,d\theta\\&=\int_0^{2\pi}d\theta\int_0^{\pi}\sin\phi\,d\phi\int_{6.370\times10^6}^{6.375\times10^6}\left(619.09\rho^2-0.000097\rho^3\right)d\rho\\&=\left[\theta\right]_0^{2\pi}\left[-\cos\phi\right]_0^{\pi}\left[\tfrac{619.09}{3}\rho^3-\tfrac{0.000097}{4}\rho^4\right]_{6.370\times10^6}^{6.375\times10^6}\\&=(2\pi)(2)\left[\tfrac{619.09}{3}\left((6.375\times10^6)^3-(6.370\times10^6)^3\right)-\tfrac{0.000097}{4}\left((6.375\times10^6)^4-(6.370\times10^6)^4\right)\right]\\&\approx 4\pi\left(1.944\times10^{17}\right)\approx 2.44\times10^{18}\text{ kg}\end{aligned}$$

47. In cylindrical coordinates, the equation of the cylinder is $r=3$, $0\le z\le 10$.

The hemisphere is the upper part of the sphere radius 3, center $(0,0,10)$, equation $r^2+(z-10)^2=3^2$, $z\ge 10$. In Maple, we can use the `coords=cylindrical` option in a regular `plot3d` command. In Mathematica, we can use `ParametricPlot3D`.

48. We begin by finding the positions of Los Angeles and Montréal in spherical coordinates, using the method described in the exercise:

Montréal	Los Angeles
$\rho=3960$ mi	$\rho=3960$ mi
$\theta=360^\circ-\ 73.60^\circ=286.40^\circ$	$\theta=360^\circ-\ 118.25^\circ=241.75^\circ$
$\phi=90^\circ-\ 45.50^\circ=44.50^\circ$	$\phi=90^\circ-\ 34.06^\circ=55.94^\circ$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Now we change the above to Cartesian coordinates using $x = \rho\cos\theta\sin\phi$, $y = \rho\sin\theta\sin\phi$ and $z = \rho\cos\phi$ to get two position vectors of length 3960 mi (since both cities must lie on the surface of the earth). In particular:

$$\text{Montréal: } \langle 783.67, -2662.67, 2824.47\rangle \qquad \text{Los Angeles: } \langle -1552.80, -2889.91, 2217.84\rangle$$

To find the angle γ between these two vectors we use the dot product:

$\langle 783.67, -2662.67, 2824.47\rangle \cdot \langle -1552.80, -2889.91, 2217.84\rangle = (3960)^2\cos\gamma \quad\Rightarrow\quad \cos\gamma \approx 0.8126 \quad\Rightarrow$ $\gamma \approx 0.6223$ rad. The great circle distance between the cities is $s = \rho\,\gamma \approx 3960(0.6223) \approx 2464$ mi.

49. If E is the solid enclosed by the surface $\rho = 1 + \frac{1}{5}\sin 6\theta\,\sin 5\phi$, it can be described in spherical coordinates as $E = \left\{(\rho,\theta,\phi) \mid 0 \le \rho \le 1 + \frac{1}{5}\sin 6\theta\sin 5\phi, 0 \le \theta \le 2\pi, 0 \le \phi \le \pi\right\}$. Its volume is given by $V(E) = \iiint_E dV = \int_0^{\pi}\int_0^{2\pi}\int_0^{1+(\sin 6\theta\sin 5\phi)/5} \rho^2\sin\phi\,d\rho\,d\theta\,d\phi = \frac{136\pi}{99}$ [using a CAS].

50. $$\int_{-\infty}^{\infty}\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} \sqrt{x^2+y^2+z^2}\,e^{-(x^2+y^2+z^2)}\,dx\,dy\,dz = \lim_{R\to\infty}\int_0^{2\pi}\int_0^{\pi}\int_0^{R} \rho e^{-\rho^2}\rho^2\sin\phi\,d\rho\,d\phi\,d\theta$$
$$= \lim_{R\to\infty}\int_0^{2\pi} d\theta \int_0^{\pi}\sin\phi\,d\phi \int_0^{R}\rho^3 e^{-\rho^2}\,d\rho$$

Now use integration by parts with $u = \rho^2$, $dv = \rho e^{-\rho^2}\,d\rho$ to get

$$\lim_{R\to\infty} 2\pi(2)\left(\rho^2\left(-\tfrac{1}{2}\right)e^{-\rho^2}\Big]_0^R - \int_0^R 2\rho\left(-\tfrac{1}{2}\right)e^{-\rho^2}\,d\rho\right) = \lim_{R\to\infty} 4\pi\left(-\tfrac{1}{2}R^2e^{-R^2} + \left[-\tfrac{1}{2}e^{-\rho^2}\right]_0^R\right)$$
$$= 4\pi\lim_{R\to\infty}\left[-\tfrac{1}{2}R^2e^{-R^2} - \tfrac{1}{2}e^{-R^2} + \tfrac{1}{2}\right] = 4\pi\left(\tfrac{1}{2}\right) = 2\pi$$

(Note that $R^2e^{-R^2} \to 0$ as $R \to \infty$ by l'Hospital's Rule.)

51. (a) From the diagram, $z = r\cot\phi_0$ to $z = \sqrt{a^2 - r^2}$, $r = 0$ to $r = a\sin\phi_0$ (or use $a^2 - r^2 = r^2\cot^2\phi_0$). Thus

$$V = \int_0^{2\pi}\int_0^{a\sin\phi_0}\int_{r\cot\phi_0}^{\sqrt{a^2-r^2}} r\,dz\,dr\,d\theta$$
$$= 2\pi\int_0^{a\sin\phi_0}\left(r\sqrt{a^2-r^2} - r^2\cot\phi_0\right)dr$$
$$= \tfrac{2\pi}{3}\left[-(a^2-r^2)^{3/2} - r^3\cot\phi_0\right]_0^{a\sin\phi_0}$$
$$= \tfrac{2\pi}{3}\left[-\left(a^2 - a^2\sin^2\phi_0\right)^{3/2} - a^3\sin^3\phi_0\cot\phi_0 + a^3\right]$$
$$= \tfrac{2}{3}\pi a^3\left[1 - \left(\cos^3\phi_0 + \sin^2\phi_0\cos\phi_0\right)\right] = \tfrac{2}{3}\pi a^3(1 - \cos\phi_0)$$

(b) The wedge in question is the shaded area rotated from $\theta = \theta_1$ to $\theta = \theta_2$. Letting

V_{ij} = volume of the region bounded by the sphere of radius ρ_i and the cone with angle ϕ_j ($\theta = \theta_1$ to θ_2)

and letting V be the volume of the wedge, we have

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\begin{aligned} V &= (V_{22} - V_{21}) - (V_{12} - V_{11}) \\ &= \tfrac{1}{3}(\theta_2 - \theta_1)\left[\rho_2^3(1-\cos\phi_2) - \rho_2^3(1-\cos\phi_1) - \rho_1^3(1-\cos\phi_2) + \rho_1^3(1-\cos\phi_1)\right] \\ &= \tfrac{1}{3}(\theta_2 - \theta_1)\left[\left(\rho_2^3 - \rho_1^3\right)(1-\cos\phi_2) - \left(\rho_2^3 - \rho_1^3\right)(1-\cos\phi_1)\right] = \tfrac{1}{3}(\theta_2 - \theta_1)\left[\left(\rho_2^3 - \rho_1^3\right)(\cos\phi_1 - \cos\phi_2)\right] \end{aligned}$$

Or: Show that $V = \displaystyle\int_{\theta_1}^{\theta_2} \int_{\rho_1 \sin\phi_1}^{\rho_2 \sin\phi_2} \int_{r\cot\phi_2}^{r\cot\phi_1} r\,dz\,dr\,d\theta$.

(c) By the Mean Value Theorem with $f(\rho) = \rho^3$ there exists some $\tilde{\rho}$ with $\rho_1 \le \tilde{\rho} \le \rho_2$ such that $f(\rho_2) - f(\rho_1) = f'(\tilde{\rho})(\rho_2 - \rho_1)$ or $\rho_2^3 - \rho_1^3 = 3\tilde{\rho}^2 \Delta\rho$. Similarly, with $f(\phi) = \cos\phi$ there exists some $\tilde{\phi}$ with $\phi_1 \le \tilde{\phi} \le \phi_2$ such that $\cos\phi_2 - \cos\phi_1 = \left(-\sin\tilde{\phi}\right)\,\Delta\phi$. Substituting into the result from (b) gives $\Delta V = \frac{1}{3}(\theta_2 - \theta_1)(3\tilde{\rho}^2\,\Delta\rho)(\sin\tilde{\phi})\,\Delta\phi = \tilde{\rho}^2 \sin\tilde{\phi}\,\Delta\rho\,\Delta\theta\,\Delta\phi$.

APPLIED PROJECT Roller Derby

1. $mgh = \frac{1}{2}mv^2 + \frac{1}{2}I\omega^2 \;\Rightarrow\; 2gh = v^2 + \dfrac{I}{m}\left(\dfrac{v}{r}\right)^2 \;\Rightarrow\; 2gh = v^2\left(1 + \dfrac{I}{mr^2}\right) \;\Rightarrow\; v^2 = \dfrac{2gh}{1+I^*}$.

2. The vertical component of the speed is $dy/dt = v\sin\alpha$, so by the same reasoning as used in Problem 1,

$$\frac{dy}{dt} = \sqrt{\frac{2gy}{1+I^*}}\,\sin\alpha = \sqrt{\frac{2g}{1+I^*}}\,\sin\alpha\,\sqrt{y}.$$

3. Solving the separable differential equation, we get $\dfrac{dy}{\sqrt{y}} = \sqrt{\dfrac{2g}{1+I^*}}\,\sin\alpha\,dt \;\Rightarrow\; 2\sqrt{y} = \sqrt{\dfrac{2g}{1+I^*}}\,(\sin\alpha)t + C$.

But $y = 0$ when $t = 0$, so $C = 0$ and we have $2\sqrt{y} = \sqrt{\dfrac{2g}{1+I^*}}\,(\sin\alpha)t$. Solving for t when $y = h$ gives

$$T = \frac{2\sqrt{h}}{\sin\alpha}\sqrt{\frac{1+I^*}{2g}} = \sqrt{\frac{2h(1+I^*)}{g\sin^2\alpha}}.$$

4. Assume that the length of each cylinder is ℓ. Then the density of the solid cylinder is $\dfrac{m}{\pi r^2 \ell}$, and from Formulas 15.6.16, its moment of inertia (using cylindrical coordinates) is

$$I_z = \iiint (x^2+y^2)\frac{m}{\pi r^2 \ell}\,dV = \frac{m}{\pi r^2 \ell}\int_0^{\ell}\int_0^{2\pi}\int_0^{r} R^2 R\,dR\,d\theta\,dz = \frac{m}{\pi r^2 \ell}\cdot \ell \cdot 2\pi \cdot \left[\tfrac{1}{4}R^4\right]_0^r = \frac{mr^2}{2}$$

and so $I^* = \dfrac{I_z}{mr^2} = \dfrac{1}{2}$.

For the hollow cylinder, we consider its entire mass to lie a distance r from the axis of rotation, so $x^2 + y^2 = r^2$ is a constant. We express the density in terms of mass per unit area as $\rho = \dfrac{m}{2\pi r\ell}$, and then the moment of inertia is calculated as a double integral: $I_z = \displaystyle\iint (x^2+y^2)\frac{m}{2\pi r\ell}\,dA = \frac{mr^2}{2\pi r\ell}\iint dA = mr^2$, so $I^* = \dfrac{I_z}{mr^2} = 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. The volume of such a ball is $\frac{4}{3}\pi(r^3 - a^3) = \frac{4}{3}\pi[r^3 - (br)^3] = \frac{4}{3}\pi r^3(1 - b^3)$, and so its density is $\dfrac{m}{\frac{4}{3}\pi r^3(1-b^3)}$. Now

$$\begin{aligned}
I_z &= \iiint (x^2 + y^2)\,\frac{m}{\frac{4}{3}\pi r^3(1-b^3)}\,dV \\
&= \frac{m}{\frac{4}{3}\pi r^3(1-b^3)} \int_a^r \int_0^{2\pi} \int_0^{\pi} (\rho^2 \sin^2\phi)(\rho^2 \sin\phi)\,d\phi\,d\theta\,d\rho \qquad \text{[from Formula 15.8.3 and Exercise 15.8.25]} \\
&= \frac{m}{\frac{4}{3}\pi r^3(1-b^3)} \left[\frac{\rho^5}{5}\right]_a^r \cdot \Big[\theta\Big]_0^{2\pi} \cdot \left[-\frac{(2+\sin^2\phi)\cos\phi}{3}\right]_0^{\pi} \qquad \text{[from Formula 67 in the Table of Integrals]} \\
&= \frac{m}{\frac{4}{3}\pi r^3(1-b^3)} \cdot \frac{r^5 - a^5}{5} \cdot 2\pi \cdot \frac{4}{3} = \frac{2mr^5(1-b^5)}{5r^3(1-b^3)} = \frac{2(1-b^5)mr^2}{5(1-b^3)}
\end{aligned}$$

Therefore, $I^* = \dfrac{I_z}{mr^2} = \dfrac{2(1-b^5)}{5(1-b^3)}$. Since a represents the inner radius, $a \to 0$ corresponds to a solid ball, and $a \to r$ corresponds to a hollow ball.

6. For a solid ball, $a \to 0 \;\Rightarrow\; b \to 0$, so $I^* = \lim\limits_{b\to 0} \dfrac{2(1-b^5)}{5(1-b^3)} = \dfrac{2}{5}$. For a hollow ball, $a \to r \;\Rightarrow\; b \to 1$, so

$$I^* = \lim_{b\to 1} \frac{2(1-b^5)}{5(1-b^3)} = \frac{2}{5} \lim_{b\to 1} \frac{-5b^4}{-3b^2} = \frac{2}{5}\left(\frac{5}{3}\right) = \frac{2}{3} \qquad \text{[by l'Hospital's Rule]}$$

Note: We could instead have calculated $I^* = \lim\limits_{b\to 1} \dfrac{2(1-b)(1+b+b^2+b^3+b^4)}{5(1-b)(1+b+b^2)} = \dfrac{2\cdot 5}{5\cdot 3} = \dfrac{2}{3}$.

Thus the objects finish in the following order: solid ball $\left(I^* = \frac{2}{5}\right)$, solid cylinder $\left(I^* = \frac{1}{2}\right)$, hollow ball $\left(I^* = \frac{2}{3}\right)$, hollow cylinder $(I^* = 1)$.

15.9 Change of Variables in Multiple Integrals

1. For Exercise 1(a)–(f), we refer to the figure. Each transformation maps the boundary of S to the boundary of one of the images (I–VI).

(a) Along S_1, $v = 0$, $0 \le u \le 1$, so $x = u + v = u$ and $y = u - v = u \;\Rightarrow\; y = x$, $0 \le x \le 1$. [Note that only images V and VI have $y = x$ as a boundary.] Along S_2, $u = 1$, $0 \le v \le 1$, so $x = 1 + v$ and $y = 1 - v$. Eliminating v gives $x + y = 2$, $1 \le x \le 2$. Along S_3, $v = 1$, $0 \le u \le 1 \;\Rightarrow\; (x)x = u + 1$ and $y = u - 1$. Eliminating u gives $y - x = -2$, $1 \le x \le 2$. Finally, along S_4, $u = 0$, $0 \le v \le 1 \;\Rightarrow\; x = v$ and $y = -v \;\Rightarrow\; y = -x$, $0 \le x \le 1$. Thus, VI is the image of the transformation.

(b) Along S_1, $v = 0$, $0 \le u \le 1 \;\Rightarrow\; y = uv = 0$, $0 \le x \le 1$. Along S_2, $u = 1$, $0 \le v \le 1 \;\Rightarrow\; x = u - v = 1 - v$, $y = v \;\Rightarrow\; y = 1 - x$, $0 \le x \le 1$. Along S_3, $v = 1$, $0 \le u \le 1 \;\Rightarrow\; x = u - 1$, $y = u \;\Rightarrow\; y = x + 1$, $-1 \le x \le 0$. Finally, along S_4, $u = 0$, $0 \le v \le 1 \;\Rightarrow\; y = 0$, $-1 \le x \le 0$. Thus, I is the image of the transformation.

(c) Along S_1, $v = 0$, $0 \le u \le 1 \;\Rightarrow\; y = u\sin v = 0$ and $x = u\cos v = u\cos 0 = u \;\Rightarrow\; 0 \le x \le 1$. Along S_2, $u = 1$, $0 \le v \le 1 \;\Rightarrow\; x = \cos v$, $y = \sin v$, $0 \le v \le 1$, which are parametric equations for a circle of radius 1,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$1 \geq x \geq \cos 1$. Along S_3, $v = 1$, $0 \leq u \leq 1 \quad \Rightarrow \quad x = u\cos 1$, $y = u \sin 1 \quad \Rightarrow \quad y = \tan(1)x$, $0 \leq x \leq \cos 1$. Along S_4, $u = 0 \quad \Rightarrow \quad x = y = 0$. Thus, IV is the image of the transformation.

(d) Along S_1, $v = 0$, $0 \leq u \leq 1 \quad \Rightarrow \quad x = u - v = u$, $y = u + v^2 = u \quad \Rightarrow \quad y = x$, $0 \leq x \leq 1$. Along S_2, $u = 1$, $0 \leq v \leq 1 \quad \Rightarrow \quad x = 1 - v$, $y = 1 + v^2 \quad \Rightarrow \quad y = 1 + (1 - x)^2$, $1 \geq x \geq 0$. Along S_3, $v = 1$, $0 \leq u \leq 1 \quad \Rightarrow$ $x = u - 1$, $y = u + 1 \quad \Rightarrow \quad y = x + 2$, $-1 \leq x \leq 0$. Finally, along S_4, $u = 0$, $0 \leq v \leq 1 \quad \Rightarrow \quad x = -v$, $y = v^2 \quad \Rightarrow$ $y = x^2$, $-1 \leq x \leq 0$. Thus, V is the image of the transformation.

(e) Along S_1, $v = 0$, $0 \leq u \leq 1 \quad \Rightarrow \quad y = 2v = 0$, and since $x = u + v = u$, $0 \leq x \leq 1$. Along S_2, $u = 1$, $0 \leq v \leq 1 \quad \Rightarrow$ $x = 1 + v$, $y = 2v \quad \Rightarrow \quad y = 2x - 2$, $1 \leq x \leq 2$. Along S_3, $v = 1$, $0 \leq u \leq 1 \quad \Rightarrow \quad y = 2v = 2$, $1 \leq x \leq 2$. Finally, along S_4, $u = 0$, $0 \leq v \leq 1 \quad \Rightarrow \quad x = v$, $y = 2v \quad \Rightarrow \quad y = 2x$, $0 \leq x \leq 1$. Thus, III is the image of the transformation.

(f) Along S_1, $v = 0$, $0 \leq u \leq 1 \quad \Rightarrow \quad x = uv = 0$, and since $y = u^3 - v^3 = u^3$, $0 \leq y \leq 1$. Along S_2, $u = 1$, $0 \leq u \leq 1 \quad \Rightarrow \quad x = v$, $y = 1 - v^3 \quad \Rightarrow \quad y = 1 - x^3$, $1 \geq x \geq 0$. Along S_3, $v = 1$, $0 \leq u \leq 1 \quad \Rightarrow$ $x = u$, $y = u^3 - 1 \quad \Rightarrow \quad y = x^3 - 1$, $0 \leq x \leq 1$. Finally, along S_4, $u = 0$, $0 \leq v \leq 1 \quad \Rightarrow \quad x = 0$, $-1 \leq y \leq 0$. Thus, II is the image of the transformation.

2. The transformation maps the boundary of S to the boundary of the image R, so we first look at side S_1 in the uv-plane. S_1 is described by $v = 0$, $0 \leq u \leq 1$, so $x = u + v = u$ and $y = -v = 0$. Therefore, the image is the line segment $y = 0$, $0 \leq x \leq 1$. S_2 is the line segment $u = 1$, $0 \leq v \leq 2$, so $x = 1 + v$ and $y = -v$. Eliminating v, we have $y = 1 - x$, $1 \leq x \leq 3$. S_3 is the line segment $v = 2$, $0 \leq u \leq 1$, so $x = u + 2$ and $y = -2 \quad \Rightarrow \quad$ the image is the line segment $y = -2$, $2 \leq x \leq 3$. Finally, S_4 is the line segment $u = 0$, $0 \leq v \leq 2$, so $x = v$ and $y = -v \quad \Rightarrow \quad$ the image is the line segment $y = -x$, $0 \leq x \leq 2$. The image of the set S is the region R shown in the xy-plane, a parallelogram bounded by these four line segments.

3. The transformation maps the boundary of S to the boundary of the image R, so we first look at side S_1 in the uv-plane. S_1 is described by $v = 0$, $0 \leq u \leq 3$, so $x = 2u + 3v = 2u$ and $y = u - v = u$. Eliminating u, we have $x = 2y$, $0 \leq x \leq 6$. S_2 is the line segment $u = 3$, $0 \leq v \leq 2$, so $x = 6 + 3v$ and $y = 3 - v$. Then $v = 3 - y \quad \Rightarrow \quad x = 6 + 3(3 - y) = 15 - 3y$, $6 \leq x \leq 12$. S_3 is the line segment $v = 2$, $0 \leq u \leq 3$, so $x = 2u + 6$ and $y = u - 2$, giving $u = y + 2 \quad \Rightarrow \quad x = 2y + 10$, $6 \leq x \leq 12$. Finally, S_4 is the segment $u = 0$, $0 \leq v \leq 2$, so $x = 3v$ and $y = -v \quad \Rightarrow \quad x = -3y$, $0 \leq x \leq 6$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The image of set S is the region R shown in the xy-plane, a parallelogram bounded by these four segments.

4. S_1 is the line segment $v = 0$, $0 \le u \le 1$, so $x = v = 0$ and $y = u(1 + v^2) = u$. Since $0 \le u \le 1$, the image is the line segment $x = 0$, $0 \le y \le 1$. S_2 is the segment $u = 1$, $0 \le v \le 1$, so $x = v$ and $y = u(1 + v^2) = 1 + x^2$. Thus the image is the portion of the parabola $y = 1 + x^2$ for $0 \le x \le 1$. S_3 is the segment $v = 1$, $0 \le u \le 1$, so $x = 1$ and $y = 2u$. The image is the segment $x = 1$, $0 \le y \le 2$. S_4 is described by $u = 0$, $0 \le v \le 1$, so $0 \le x = v \le 1$ and $y = u(1 + v^2) = 0$. The image is the line segment $y = 0$, $0 \le x \le 1$. Thus, the image of S is the region R bounded by the parabola $y = 1 + x^2$, the x-axis, and the lines $x = 0$, $x = 1$.

5. S_1 is the line segment $u = v$, $0 \le u \le 1$, so $y = v = u$ and $x = u^2 = y^2$. Since $0 \le u \le 1$, the image is the portion of the parabola $x = y^2$, $0 \le y \le 1$. S_2 is the segment $v = 1$, $0 \le u \le 1$, thus $y = v = 1$ and $x = u^2$, so $0 \le x \le 1$. The image is the line segment $y = 1$, $0 \le x \le 1$. S_3 is the segment $u = 0$, $0 \le v \le 1$, so $x = u^2 = 0$ and $y = v \quad \Rightarrow \quad 0 \le y \le 1$. The image is the segment $x = 0$, $0 \le y \le 1$. Thus, the image of S is the region R in the first quadrant bounded by the parabola $x = y^2$, the y-axis, and the line $y = 1$.

6. Substituting $u = \dfrac{x}{a}$, $v = \dfrac{y}{b}$ into $u^2 + v^2 \le 1$ gives $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} \le 1$, so the image of $u^2 + v^2 \le 1$ is the elliptical region $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} \le 1$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. R is a parallelogram enclosed by the parallel lines $y = 2x - 1$, $y = 2x + 1$ and the parallel lines $y = 1 - x$, $y = 3 - x$. The first pair of equations can be written as $y - 2x = -1$, $y - 2x = 1$. If we let $u = y - 2x$ then these lines are mapped to the vertical lines $u = -1$, $u = 1$ in the uv-plane. Similarly, the second pair of equations can be written as $x + y = 1$, $x + y = 3$, and setting $v = x + y$ maps these lines to the horizontal lines $v = 1$, $v = 3$ in the uv-plane. Boundary curves are mapped to boundary curves under a transformation, so here the equations $u = y - 2x$, $v = x + y$ define a transformation T^{-1} that maps R in the xy-plane to the square S enclosed by the lines $u = -1$, $u = 1$, $v = 1$, $v = 3$ in the uv-plane. To find the transformation T that maps S to R we solve $u = y - 2x$, $v = x + y$ for x, y: Subtracting the first equation from the second gives $v - u = 3x \quad \Rightarrow \quad x = \frac{1}{3}(v - u)$ and adding twice the second equation to the first gives $u + 2v = 3y \quad \Rightarrow$ $y = \frac{1}{3}(u + 2v)$. Thus one possible transformation T (there are many) is given by $x = \frac{1}{3}(v - u)$, $y = \frac{1}{3}(u + 2v)$.

8. The boundaries of the parallelogram R are the lines $y = \frac{3}{4}x$ or $4y - 3x = 0$, $y = \frac{3}{4}x + \frac{5}{2}$ or $4y - 3x = 10$, $y = -\frac{1}{2}x$ or $x + 2y = 0$, $y = -\frac{1}{2}x + 5$ or $x + 2y = 10$. Setting $u = 4y - 3x$ and $v = x + 2y$ defines a transformation T^{-1} that maps R in the xy-plane to the square S enclosed by the lines $u = 0$, $u = 10$, $v = 0$, $v = 10$ in the uv-plane. Solving $u = 4y - 3x$, $v = x + 2y$ for x and y gives $2v - u = 5x \quad \Rightarrow \quad x = \frac{1}{5}(2v - u)$, $u + 3v = 10y \quad \Rightarrow \quad y = \frac{1}{10}(u + 3v)$. Thus one possible transformation T is given by $x = \frac{1}{5}(2v - u)$, $y = \frac{1}{10}(u + 3v)$.

9. R is a portion of an annular region (see the figure) that is easily described in polar coordinates as $R = \{(r, \theta) \mid 1 \le r \le \sqrt{2},\ 0 \le \theta \le \pi/2\}$. If we converted a double integral over R to polar coordinates the resulting region of integration is a rectangle (in the $r\theta$-plane), so we can create a transformation T here by letting u play the role of r and v the role of θ. Thus T is defined by $x = u \cos v$, $y = u \sin v$ and T maps the rectangle

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$S = \{(u, v) \mid 1 \le u \le \sqrt{2},\ 0 \le v \le \pi/2\}$ in the uv-plane to R in the xy-plane.

10. The boundaries of the region R are the curves $y = 1/x$ or $xy = 1$, $y = 4/x$ or $xy = 4$, $y = x$ or $y/x = 1$, $y = 4x$ or $y/x = 4$. Setting $u = xy$ and $v = y/x$ defines a transformation T^{-1} that maps R in the xy-plane to the square S enclosed by the lines $u = 1$, $u = 4$, $v = 1$, $v = 4$ in the uv-plane. Solving $u = xy$, $v = y/x$ for x and y gives $x^2 = u/v \quad \Rightarrow$ $x = \sqrt{u/v}$ [since x, y, u, v are all positive], $y^2 = uv \quad \Rightarrow \quad y = \sqrt{uv}$. Thus one possible transformation T is given by $x = \sqrt{u/v}$, $y = \sqrt{uv}$.

11. $x = 2u + v,\ y = 4u - v$.

The Jacobian is $\dfrac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \partial x/\partial u & \partial x/\partial v \\ \partial y/\partial u & \partial y/\partial v \end{vmatrix} = \begin{vmatrix} 2 & 1 \\ 4 & -1 \end{vmatrix} = (2)(-1) - (1)(4) = -6.$

12. $x = u^2 + uv,\ y = uv^2$.

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \partial x/\partial u & \partial x/\partial v \\ \partial y/\partial u & \partial y/\partial v \end{vmatrix} = \begin{vmatrix} 2u + v & u \\ v^2 & 2uv \end{vmatrix} = (2u + v)(2uv) - u(v^2) = 4u^2v + 2uv^2 - uv^2 = 4u^2v + uv^2$$

13. $x = s\cos t,\ y = s\sin t$.

$$\frac{\partial(x, y)}{\partial(s, t)} = \begin{vmatrix} \partial x/\partial s & \partial x/\partial t \\ \partial y/\partial s & \partial y/\partial t \end{vmatrix} = \begin{vmatrix} \cos t & -s\sin t \\ \sin t & s\cos t \end{vmatrix} = s\cos^2 t - (-s\sin^2 t) = s(\cos^2 t + \sin^2 t) = s$$

14. $x = pe^q,\ y = qe^p$.

$$\frac{\partial(x, y)}{\partial(p, q)} = \begin{vmatrix} \partial x/\partial p & \partial x/\partial q \\ \partial y/\partial p & \partial y/\partial q \end{vmatrix} = \begin{vmatrix} e^q & pe^q \\ qe^p & e^p \end{vmatrix} = e^q e^p - pe^q \cdot qe^p = e^{p+q} - pqe^{p+q} = (1 - pq)e^{p+q}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $x = uv,\ y = vw,\ z = wu.$

$$\frac{\partial(x,y,z)}{\partial(u,v,w)} = \begin{vmatrix} \partial x/\partial u & \partial x/\partial v & \partial x/\partial w \\ \partial y/\partial u & \partial y/\partial v & \partial y/\partial w \\ \partial z/\partial u & \partial z/\partial v & \partial z/\partial w \end{vmatrix} = \begin{vmatrix} v & u & 0 \\ 0 & w & v \\ w & 0 & u \end{vmatrix} = v\begin{vmatrix} w & v \\ 0 & u \end{vmatrix} - u\begin{vmatrix} 0 & v \\ w & u \end{vmatrix} + 0\begin{vmatrix} 0 & w \\ w & 0 \end{vmatrix}$$

$$= v(uw - 0) - u(0 - vw) + 0 = uvw + uvw = 2uvw$$

16. $x = u + vw,\ y = v + wu,\ z = w + uv.$

$$\frac{\partial(x,y,z)}{\partial(u,v,w)} = \begin{vmatrix} \partial x/\partial u & \partial x/\partial v & \partial x/\partial w \\ \partial y/\partial u & \partial y/\partial v & \partial y/\partial w \\ \partial z/\partial u & \partial z/\partial v & \partial z/\partial w \end{vmatrix} = \begin{vmatrix} 1 & w & v \\ w & 1 & u \\ v & u & 1 \end{vmatrix} = 1\begin{vmatrix} 1 & u \\ u & 1 \end{vmatrix} - w\begin{vmatrix} w & u \\ v & 1 \end{vmatrix} + v\begin{vmatrix} w & 1 \\ v & u \end{vmatrix}$$

$$= 1(1 - u^2) - w\,(w - uv) + v\,(uw - v)$$

$$= 1 - u^2 - w^2 + uvw + uvw - v^2 = 1 + 2uvw - u^2 - v^2 - w^2$$

17. $x = 2u + v,\ y = u + 2v \quad \Rightarrow \quad \dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} = 3$. The integrand $x - 3y = (2u + v) - 3(u + 2v) = -u - 5v$. To find the region S in the uv-plane that corresponds to R we first find the corresponding boundary under the given transformation. The line through $(0,0)$ and $(2,1)$ is $y = \frac{1}{2}x$ which is the image of $u + 2v = \frac{1}{2}(2u + v) \quad \Rightarrow \quad v = 0$; the line through $(2,1)$ and $(1,2)$ is $x + y = 3$ which is the image of $(2u + v) + (u + 2v) = 3 \quad \Rightarrow \quad u + v = 1$; the line through $(0,0)$ and $(1,2)$ is $y = 2x$ which is the image of $u + 2v = 2(2u + v) \quad \Rightarrow \quad u = 0$. Thus S is the triangle $0 \le v \le 1 - u$, $0 \le u \le 1$ in the uv-plane and

$$\iint_R (x - 3y)\,dA = \int_0^1 \int_0^{1-u} (-u - 5v)\,|3|\ dv\,du = -3\int_0^1 \left[uv + \tfrac{5}{2}v^2\right]_{v=0}^{v=1-u}\ du$$

$$= -3\int_0^1 \left(u - u^2 + \tfrac{5}{2}(1-u)^2\right)\ du = -3\left[\tfrac{1}{2}u^2 - \tfrac{1}{3}u^3 - \tfrac{5}{6}(1-u)^3\right]_0^1 = -3\left(\tfrac{1}{2} - \tfrac{1}{3} + \tfrac{5}{6}\right) = -3$$

18. $x = \frac{1}{4}(u + v),\ y = \frac{1}{4}(u - 3u) \quad \Rightarrow \quad \dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/4 & 1/4 \\ -3/4 & 1/4 \end{vmatrix} = \dfrac{1}{4}$. The integrand $4x + 8y = 4 \cdot \frac{1}{4}(u + v) + 8 \cdot \frac{1}{4}(v - 3u) = 3v - 5u$. R is a parallelogram bounded by the lines $x - y = -4$, $x - y = 4$, $3x + y = 0$, $3x + y = 8$. Since $u = x - y$ and $v = 3x + y$, R is the image of the rectangle enclosed by the lines $u = -4$, $u = 4$, $v = 0$, and $v = 8$. Thus

$$\iint_R (4x + 8y)\,dA = \int_{-4}^4 \int_0^8 (3v - 5u)\left|\tfrac{1}{4}\right|\ dv\,du = \tfrac{1}{4}\int_{-4}^4 \left[\tfrac{3}{2}v^2 - 5uv\right]_{v=0}^{v=8}\ du$$

$$= \tfrac{1}{4}\int_{-4}^4 (96 - 40u)\,du = \tfrac{1}{4}\left[96u - 20u^2\right]_{-4}^4 = 192$$

19. $x = 2u,\ y = 3v \quad \Rightarrow \quad \dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 2 & 0 \\ 0 & 3 \end{vmatrix} = 6$. The integrand $x^2 = 4u^2$. The planar ellipse $9x^2 + 4y^2 \le 36$ is the image of the disk $u^2 + v^2 \le 1$. Thus,

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\iint_R x^2\,dA = \iint\limits_{u^2+v^2\le 1} (4u^2)(6)\,du\,dv = \int_0^{2\pi}\int_0^1 (24r^2\cos^2\theta)\,r\,dr\,d\theta = 24\int_0^{2\pi}\cos^2\theta\,d\theta\int_0^1 r^3\,dr$$
$$= 24\left[\tfrac{1}{2}x + \tfrac{1}{4}\sin 2x\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^1 = 24(\pi)\left(\tfrac{1}{4}\right) = 6\pi$$

20. $x = \sqrt{2}\,u - \sqrt{2/3}\,v$, $y = \sqrt{2}\,u + \sqrt{2/3}\,v \;\Rightarrow\; \dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \sqrt{2} & -\sqrt{2/3} \\ \sqrt{2} & \sqrt{2/3} \end{vmatrix} = \dfrac{4}{\sqrt{3}}$. The integrand $x^2 - xy + y^2 = 2u^2 + 2v^2$. The planar ellipse $x^2 - xy + y^2 \le 2$ is the image of the disk $u^2 + v^2 \le 1$. Thus,

$$\iint_R (x^2 - xy + y^2)\,dA = \iint\limits_{u^2+v^2\le 1} (2u^2 + 2v^2)\left(\tfrac{4}{\sqrt{3}}\,du\,dv\right) = \int_0^{2\pi}\int_0^1 \tfrac{8}{\sqrt{3}}r^3\,dr\,d\theta = \tfrac{4\pi}{\sqrt{3}}$$

21. $x = u/v$, $y = v \;\Rightarrow\; \dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/v & -u/v^2 \\ 0 & 1 \end{vmatrix} = \dfrac{1}{v}$. The integrand $xy = u$. The line $y = x$ $[v = u/v]$ is the image of the parabola $v^2 = u$ and the line $y = 3x$ $[v = 3u/v]$ is the image of the parabola $v^2 = 3u$. The hyperbolas $xy = 1$ and $xy = 3$ are the images of the lines $u = 1$ and $u = 3$, respectively. Thus,

$$\iint_R xy\,dA = \int_1^3\int_{\sqrt{u}}^{\sqrt{3u}} u\left(\frac{1}{v}\right)dv\,du = \int_1^3 u\left(\ln\sqrt{3u} - \ln\sqrt{u}\right)du = \int_1^3 u\ln\sqrt{3}\,du = 4\ln\sqrt{3} = 2\ln 3.$$

22. $u = xy$, $v = xy^2$. To solve for x and y in terms of u and v, try dividing. $\dfrac{xy^2}{xy} = \dfrac{v}{u} \;\Rightarrow\; y = \dfrac{v}{u}$. Also, $\dfrac{(xy)^2}{xy^2} = \dfrac{u^2}{v} \;\Rightarrow\; x = \dfrac{u^2}{v}$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 2u/v & -u^2/v^2 \\ -v/u^2 & 1/u \end{vmatrix} = \dfrac{1}{v}$. The integrand $y^2 = v^2/u^2$. R is the image of the square with vertices $(1,1)$, $(2,1)$, $(2,2)$, and $(1,2)$. Thus,

$$\iint_R y^2\,dA = \int_1^2\int_1^2 \frac{v^2}{u^2}\left(\frac{1}{v}\right)du\,dv = \int_1^2 \frac{v}{2}\,dv = \frac{3}{4}$$

23. (a) $x = au$, $y = bv$, $z = cw \;\Rightarrow\; \dfrac{\partial(x,y,z)}{\partial(u,v,w)} = \begin{vmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{vmatrix} = abc$. Since $u = \dfrac{x}{a}$, $v = \dfrac{y}{b}$, $w = \dfrac{z}{c}$, the solid enclosed by the ellipsoid is the image of the ball $u^2 + v^2 + w^2 \le 1$. Thus,

$$\iiint_E dV = \iiint\limits_{u^2+v^2+w^2\le 1} abc\,du\,dv\,dw = (abc)(\text{volume of the ball}) = \tfrac{4}{3}\pi abc$$

(b) If we approximate the surface of the earth by the ellipsoid $\dfrac{x^2}{6378^2} + \dfrac{y^2}{6378^2} + \dfrac{z^2}{6356^2} = 1$, then we can estimate the volume of the earth by finding the volume of the solid E enclosed by the ellipsoid. From part (a), this is $\iiint_E dV = \frac{4}{3}\pi(6378)(6378)(6356) \approx 1.083 \times 10^{12}$ km^3.

(c) The moment of intertia about the z-axis is $I_z = \iiint_E \left(x^2 + y^2\right)\rho(x,y,z)\,dV$, where E is the solid enclosed by $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} + \dfrac{z^2}{c^2} = 1$. As in part (a), we use the transformation $x = au$, $y = bv$, $z = cw$, so $\left|\dfrac{\partial(x,y,z)}{\partial(u,v,w)}\right| = abc$ and

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$I_z = \iiint_E \left(x^2+y^2\right) k\, dV = \iiint\limits_{u^2+v^2+w^2\le 1} k(a^2u^2+b^2v^2)(abc)\, du\, dv\, dw$$

$$= abck \int_0^{\pi}\int_0^{2\pi}\int_0^1 (a^2\rho^2\sin^2\phi\cos^2\theta + b^2\rho^2\sin^2\phi\sin^2\theta)\,\rho^2\sin\phi\, d\rho\, d\theta\, d\phi$$

$$= abck\left[a^2\int_0^{\pi}\int_0^{2\pi}\int_0^1 (\rho^2\sin^2\phi\cos^2\theta)\,\rho^2\sin\phi\, d\rho\, d\theta\, d\phi + b^2\int_0^{\pi}\int_0^{2\pi}\int_0^1 (\rho^2\sin^2\phi\sin^2\theta)\,\rho^2\sin\phi\, d\rho\, d\theta\, d\phi\right]$$

$$= a^3bck\int_0^{\pi}\sin^3\phi\, d\phi\int_0^{2\pi}\cos^2\theta\, d\theta\int_0^1\rho^4\, d\rho + ab^3ck\int_0^{\pi}\sin^3\phi\, d\phi\int_0^{2\pi}\sin^2\theta\, d\theta\int_0^1\rho^4\, d\rho$$

$$= a^3bck\left[\tfrac{1}{3}\cos^3\phi-\cos\phi\right]_0^{\pi}\left[\tfrac{1}{2}\theta+\tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{5}\rho^5\right]_0^1 + ab^3ck\left[\tfrac{1}{3}\cos^3\phi-\cos\phi\right]_0^{\pi}\left[\tfrac{1}{2}\theta-\tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{5}\rho^5\right]_0^1$$

$$= a^3bck\left(\tfrac{4}{3}\right)(\pi)\left(\tfrac{1}{5}\right) + ab^3ck\left(\tfrac{4}{3}\right)(\pi)\left(\tfrac{1}{5}\right) = \tfrac{4}{15}\pi(a^2+b^2)abck$$

24. R is the region enclosed by the curves $xy=a$, $xy=b$, $xy^{1.4}=c$, and $xy^{1.4}=d$, so if we let $u=xy$ and $v=xy^{1.4}$ then R is the image of the rectangle enclosed by the lines $u=a$, $u=b$ $(a<b)$ and $v=c$, $v=d$ $(c<d)$. Now $x=u/y \;\Rightarrow\; v=(u/y)y^{1.4}=uy^{0.4} \;\Rightarrow\; y^{0.4}=u^{-1}v \;\Rightarrow\; y=(u^{-1}v)^{1/0.4}=u^{-2.5}v^{2.5}$ and $x=uy^{-1}=u(u^{-2.5}v^{2.5})^{-1}=u^{3.5}v^{-2.5}$, so

$$\frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 3.5u^{2.5}v^{-2.5} & -2.5u^{3.5}v^{-3.5} \\ -2.5u^{-3.5}v^{2.5} & 2.5u^{-2.5}v^{1.5}\end{vmatrix} = 8.75v^{-1}-6.25v^{-1} = 2.5v^{-1}$$

Thus the area of R, and the work done by the engine, is

$$\iint_R dA = \int_a^b\int_c^d \left|2.5v^{-1}\right| dv\, du = 2.5\int_a^b du\int_c^d (1/v)\, dv = 2.5\big[u\big]_a^b\big[\ln|v|\big]_c^d = 2.5(b-a)(\ln d-\ln c) = 2.5(b-a)\ln\frac{d}{c}.$$

25. Letting $u=x-2y$ and $v=3x-y$, we have $x=\frac{1}{5}(2v-u)$ and $y=\frac{1}{5}(v-3u)$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} -1/5 & 2/5 \\ -3/5 & 1/5\end{vmatrix} = \dfrac{1}{5}$.

R is the image of the rectangle enclosed by the lines $u=0$, $u=4$, $v=1$, and $v=8$. Thus,

$$\iint_R \frac{x-2y}{3x-y}\, dA = \int_0^4\int_1^8 \frac{u}{v}\left|\frac{1}{5}\right| dv\, du = \frac{1}{5}\int_0^4 u\, du\int_1^8 \frac{1}{v}\, dv = \tfrac{1}{5}\left[\tfrac{1}{2}u^2\right]_0^4\big[\ln|v|\big]_1^8 = \tfrac{8}{5}\ln 8$$

26. Letting $u=x+y$ and $v=x-y$, we have $x=\frac{1}{2}(u+v)$ and $y=\frac{1}{2}(u-v)$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/2 & 1/2 \\ 1/2 & -1/2\end{vmatrix} = -\dfrac{1}{2}$.

R is the image of the rectangle enclosed by the lines $u=0$, $u=3$, $v=0$, and $v=2$. Thus,

$$\iint_R (x+y)\, e^{x^2-y^2}\, dA = \int_0^3\int_0^2 ue^{uv}\left|-\tfrac{1}{2}\right| dv\, du = \tfrac{1}{2}\int_0^3\left[e^{uv}\right]_{v=0}^{v=2} du = \tfrac{1}{2}\int_0^3 (e^{2u}-1)\, du$$

$$= \tfrac{1}{2}\left[\tfrac{1}{2}e^{2u}-u\right]_0^3 = \tfrac{1}{2}\left(\tfrac{1}{2}e^6-3-\tfrac{1}{2}\right) = \tfrac{1}{4}(e^6-7)$$

27. Letting $u=y-x$ and $v=y+x$, we have $x=\frac{1}{2}(v-u)$ and $y=\frac{1}{2}(u+v)$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} -1/2 & 1/2 \\ 1/2 & 1/2\end{vmatrix} = -\dfrac{1}{2}$.

R is the image of the trapezoidal region with vertices $(-1,1)$, $(-2,2)$, $(2,2)$, and $(1,1)$. Thus,

$$\iint_R \cos\left(\frac{y-x}{y+x}\right) dA = \int_1^2\int_{-v}^{v}\cos\frac{u}{v}\left|-\frac{1}{2}\right| du\, dv = \frac{1}{2}\int_1^2\left[v\sin\frac{u}{v}\right]_{u=-v}^{u=v} dv$$

$$= \tfrac{1}{2}\int_1^2[v\sin(1)-v\sin(-1)]\, dv = \tfrac{1}{2}\int_1^2 2v\sin 1\, dv$$

$$= \sin 1\left[\tfrac{1}{2}v^2\right]_1^2 = \tfrac{3}{2}\sin 1$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. Letting $u = 3x$ and $v = 2y$, we have $9x^2 + 4y^2 = u^2 + v^2$, $x = \frac{1}{3}u$, and $y = \frac{1}{2}v$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/3 & 0 \\ 0 & 1/2 \end{vmatrix} = \dfrac{1}{6}$.

R is the image of the quarter-disk D given by $u^2 + v^2 \le 1$, $u \ge 0$, $v \ge 0$. Thus,

$$\iint_R \sin(9x^2 + 4y^2)\, dA = \iint_D \tfrac{1}{6} \sin(u^2 + v^2)\, du\, dv = \int_0^{\pi/2} \int_0^1 \tfrac{1}{6} \sin(r^2)\, r\, dr\, d\theta = \tfrac{\pi}{12}\left[-\tfrac{1}{2}\cos r^2\right]_0^1 = \tfrac{\pi}{24}(1 - \cos 1)$$

29. See the figure. Leting $u = x + y$ and $v = -x + y$, we have $x = \frac{1}{2}(u - v)$ and $y = \frac{1}{2}(u + v)$.

Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/2 & -1/2 \\ 1/2 & 1/2 \end{vmatrix} = \dfrac{1}{2}$. Now $|u| = |x + y| \le |x| + |y| \le 1 \quad \Rightarrow$

$-1 \le u \le 1$, and $|v| = |-x + y| \le |x| + |y| \le 1 \quad \Rightarrow \quad -1 \le v \le 1$.

R is the image of the square region with vertices $(1, 1)$, $(1, -1)$, $(-1, -1)$, and $(-1, 1)$.

Thus, $\iint_R e^{x+y}\, dA = \frac{1}{2} \int_{-1}^1 \int_{-1}^1 e^u\, du\, dv = \frac{1}{2}\left[e^u\right]_{-1}^1 \left[\, v \,\right]_{-1}^1 = e - e^{-1}$.

30. Letting $u = x + y$ and $v = y/x$, we have $x = \dfrac{u}{1+v}$ and $y = \dfrac{uv}{1+v}$. Then

$$\frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/(1+v) & -u/(1+v)^2 \\ v/(1+v) & u/(1+v)^2 \end{vmatrix} = \frac{u + uv}{(1+v)^3} = \frac{u}{(1+v)^2}$$

R is the image of the rectangle enclosed by the lines $u = 1$, $u = 3$, $v = 2$, and $v = 1/2$. Thus,

$$\begin{aligned} \iint_R \frac{y}{x}\, dA &= \int_{1/2}^2 \int_1^3 v\left(\frac{u}{(1+v)^2}\right) du\, dv = \int_{1/2}^2 \frac{v}{(1+v)^2}\, dv \int_1^3 u\, du \\ &= \left[\ln|1+v| + \frac{1}{1+v}\right]_{1/2}^2 \left[\frac{u^2}{2}\right]_1^3 \qquad \left[\text{Use the substitution } w = 1 + v \text{ for the first integral}\right] \\ &= \left(\ln 3 + \tfrac{1}{3} - \ln\tfrac{3}{2} - \tfrac{2}{3}\right) \cdot 4 = 4\ln 2 - \tfrac{4}{3} \end{aligned}$$

31. Letting $u = x + y$ and $v = y$, we have $x = u - v$ and $y = v$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1 & -1 \\ 0 & 1 \end{vmatrix} = 1$.

R is the image under T of the triangular region with vertices $(0, 0)$, $(1, 0)$ and $(1, 1)$. Thus,

$$\iint_R f(x+y)\, dA = \int_0^1 \int_0^u (1)\, f(u)\, dv\, du = \int_0^1 f(u)\left[\, v \,\right]_{v=0}^{v=u} du = \int_0^1 u f(u)\, du, \text{ as desired.}$$

15 Review

TRUE-FALSE QUIZ

1. This is true by Fubini's Theorem.

2. False. $\int_0^1 \int_0^x \sqrt{x + y^2}\, dy\, dx$ describes the region of integration as a Type I region. To reverse the order of integration, we must consider the region as a Type II region: $\int_0^1 \int_y^1 \sqrt{x + y^2}\, dx\, dy$.

3. True by Equation 15.1.11.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $\int_{-1}^{1}\int_{0}^{1} e^{x^2+y^2}\sin y\,dx\,dy = \left(\int_0^1 e^{x^2}\,dx\right)\left(\int_{-1}^{1} e^{y^2}\sin y\,dy\right) = \left(\int_0^1 e^{x^2}\,dx\right)(0) = 0$, since $e^{y^2}\sin y$ is an odd function. Therefore the statement is true.

5. True. By Equation 15.1.11 we can write $\int_0^1\int_0^1 f(x)\,f(y)\,dy\,dx = \int_0^1 f(x)\,dx\,\int_0^1 f(y)\,dy$. But $\int_0^1 f(y)\,dy = \int_0^1 f(x)\,dx$ so this becomes $\int_0^1 f(x)\,dx\,\int_0^1 f(x)\,dx = \left[\int_0^1 f(x)\,dx\right]^2$.

6. This statement is true because in the given region, $\left(x^2+\sqrt{y}\,\right)\sin(x^2y^2) \le (1+2)(1) = 3$, so $\int_1^4\int_0^1 \left(x^2+\sqrt{y}\,\right)\sin(x^2y^2)\,dx\,dy \le \int_1^4\int_0^1 3\,dA = 3A(D) = 3(3) = 9$.

7. True. $\iint_D \sqrt{4-x^2-y^2}\,dA =$ the volume under the surface $x^2+y^2+z^2=4$ and above the xy-plane

$$= \tfrac{1}{2}\left(\text{the volume of the sphere } x^2+y^2+z^2=4\right) = \tfrac{1}{2}\cdot\tfrac{4}{3}\pi(2)^3 = \tfrac{16}{3}\pi$$

8. True. The moment of inertia about the z-axis of a solid E with constant density k is $I_z = \iiint_E (x^2+y^2)\rho(x,y,z)\,dV = \iiint_E (kr^2)\,r\,dz\,dr\,d\theta = \iiint_E kr^3\,dz\,dr\,d\theta$.

9. The volume enclosed by the cone $z=\sqrt{x^2+y^2}$ and the plane $z=2$ is, in cylindrical coordinates, $V = \int_0^{2\pi}\int_0^2\int_r^2 r\,dz\,dr\,d\theta \ne \int_0^{2\pi}\int_0^2\int_r^2 dz\,dr\,d\theta$, so the assertion is false.

EXERCISES

1. As shown in the contour map, we divide R into 9 equally sized subsquares, each with area $\Delta A = 1$. Then we approximate $\iint_R f(x,y)\,dA$ by a Riemann sum with $m=n=3$ and the sample points the upper right corners of each square, so

$$\iint_R f(x,y)\,dA \approx \sum_{i=1}^{3}\sum_{j=1}^{3} f(x_i,y_j)\,\Delta A$$
$$= \Delta A\,[f(1,1)+f(1,2)+f(1,3)+f(2,1)+f(2,2)+f(2,3)+f(3,1)+f(3,2)+f(3,3)]$$

Using the contour lines to estimate the function values, we have

$$\iint_R f(x,y)\,dA \approx 1[2.7+4.7+8.0+4.7+6.7+10.0+6.7+8.6+11.9] \approx 64.0$$

2. As in Exercise 1, we have $m=n=3$ and $\Delta A = 1$. Using the contour map to estimate the value of f at the center of each subsquare, we have

$$\iint_R f(x,y)\,dA \approx \sum_{i=1}^{3}\sum_{j=1}^{3} f\left(\overline{x}_i,\overline{y}_j\right)\Delta A$$
$$= \Delta A\,[f(0.5,0.5)+(0.5,1.5)+(0.5,2.5)+(1.5,0.5)+f(1.5,1.5)$$
$$+ f(1.5,2.5)+(2.5,0.5)+f(2.5,1.5)+f(2.5,2.5)]$$
$$\approx 1[1.2+2.5+5.0+3.2+4.5+7.1+5.2+6.5+9.0] = 44.2$$

3. $\int_1^2\int_0^2 (y+2xe^y)\,dx\,dy = \int_1^2 \left[xy+x^2e^y\right]_{x=0}^{x=2} dy = \int_1^2 (2y+4e^y)\,dy = \left[y^2+4e^y\right]_1^2$
$$= 4+4e^2-1-4e = 4e^2-4e+3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. $\int_0^1 \int_0^1 ye^{xy}\,dx\,dy = \int_0^1 \left[e^{xy}\right]_{x=0}^{x=1} dy = \int_0^1 (e^y - 1)\,dy = \left[e^y - y\right]_0^1 = e - 2$

5. $\int_0^1 \int_0^x \cos(x^2)\,dy\,dx = \int_0^1 \left[\cos(x^2)y\right]_{y=0}^{y=x} dx = \int_0^1 x\cos(x^2)\,dx = \frac{1}{2}\sin(x^2)\Big]_0^1 = \frac{1}{2}\sin 1$

6. $\int_0^1 \int_x^{e^x} 3xy^2\,dy\,dx = \int_0^1 \left[xy^3\right]_{y=x}^{y=e^x} dx = \int_0^1 (xe^{3x} - x^4)\,dx = \frac{1}{3}xe^{3x}\Big]_0^1 - \int_0^1 \frac{1}{3}e^{3x}\,dx - \left[\frac{1}{5}x^5\right]_0^1$ $\left[\begin{array}{c}\text{integrate by parts}\\ \text{in the first term}\end{array}\right]$

$= \frac{1}{3}e^3 - \left[\frac{1}{9}e^{3x}\right]_0^1 - \frac{1}{5} = \frac{2}{9}e^3 - \frac{4}{45}$

7. $\int_0^\pi \int_0^1 \int_0^{\sqrt{1-y^2}} y\sin x\,dz\,dy\,dx = \int_0^\pi \int_0^1 \left[(y\sin x)z\right]_{z=0}^{z=\sqrt{1-y^2}} dy\,dx = \int_0^\pi \int_0^1 y\sqrt{1-y^2}\sin x\,dy\,dx$

$= \int_0^\pi \left[-\frac{1}{3}(1-y^2)^{3/2}\sin x\right]_{y=0}^{y=1} dx = \int_0^\pi \frac{1}{3}\sin x\,dx = -\frac{1}{3}\cos x\Big]_0^\pi = \frac{2}{3}$

8. $\int_0^1 \int_0^y \int_x^1 6xyz\,dz\,dx\,dy = \int_0^1 \int_0^y \left[3xyz^2\right]_{z=x}^{z=1} dx\,dy = \int_0^1 \int_0^y (3xy - 3x^3y)\,dx\,dy$

$= \int_0^1 \left[\frac{3}{2}x^2y - \frac{3}{4}x^4y\right]_{x=0}^{x=y} dy = \int_0^1 \left(\frac{3}{2}y^3 - \frac{3}{4}y^5\right) dy = \left[\frac{3}{8}y^4 - \frac{1}{8}y^6\right]_0^1 = \frac{1}{4}$

9. The region R is more easily described by polar coordinates: $R = \{(r,\theta) \mid 2 \le r \le 4,\, 0 \le \theta \le \pi\}$. Thus $\iint_R f(x,y)\,dA = \int_0^\pi \int_2^4 f(r\cos\theta, r\sin\theta)\,r\,dr\,d\theta$.

10. The region R is a type II region that can be described as the region enclosed by the lines $y = 4 - x$, $y = 4 + x$, and the x-axis. So using rectangular coordinates, we can say $R = \{(x,y) \mid 0 \le y \le 4,\, y - 4 \le x \le 4 - y\}$ and $\iint_R f(x,y)\,dA = \int_0^4 \int_{y-4}^{4-y} f(x,y)\,dx\,dy$.

11. $x = r\cos\theta = 2\sqrt{3}\cos\frac{\pi}{3} = 2\sqrt{3}\cdot\frac{1}{2} = \sqrt{3}$, $y = r\sin\theta = 2\sqrt{3}\sin\frac{\pi}{3} = 2\sqrt{3}\cdot\frac{\sqrt{3}}{2} = 3$, $z = 2$, so in rectangular coordinates the point is $\left(\sqrt{3}, 3, 2\right)$. $\rho = \sqrt{r^2 + z^2} = \sqrt{12 + 4} = 4$, $\theta = \frac{\pi}{3}$, and $\cos\phi = z/\rho = \frac{1}{2}$, so $\phi = \frac{\pi}{3}$ and spherical coordinates are $\left(4, \frac{\pi}{3}, \frac{\pi}{3}\right)$.

12. $r = \sqrt{4+4} = 2\sqrt{2}$; $z = -1$; $\tan\theta = \frac{2}{2} = 1$ and the point $(2,2)$ is in the first quadrant of the xy-plane, so $\theta = \frac{\pi}{4}$. Thus in cylindrical coordinates the point is $\left(2\sqrt{2}, \frac{\pi}{4}, -1\right)$. $\rho = \sqrt{4+4+1} = 3$, $\cos\phi = z/\rho = -\frac{1}{3}$, so the spherical coordinates are $\left(3, \frac{\pi}{4}, \cos^{-1}\left(-\frac{1}{3}\right)\right)$.

13. $x = \rho\sin\phi\cos\theta = 8\sin\frac{\pi}{6}\cos\frac{\pi}{4} = 8\cdot\frac{1}{2}\cdot\frac{\sqrt{2}}{2} = 2\sqrt{2}$, $y = \rho\sin\phi\sin\theta = 8\sin\frac{\pi}{6}\sin\frac{\pi}{4} = 2\sqrt{2}$, and $z = \rho\cos\phi = 8\cos\frac{\pi}{6} = 8\cdot\frac{\sqrt{3}}{2} = 4\sqrt{3}$. Thus rectangular coordinates for the point are $\left(2\sqrt{2}, 2\sqrt{2}, 4\sqrt{3}\right)$.
$r^2 = x^2 + y^2 = 8 + 8 = 16 \quad\Rightarrow\quad r = 4$, $\theta = \frac{\pi}{4}$, and $z = 4\sqrt{3}$, so cylindrical coordinates are $\left(4, \frac{\pi}{4}, 4\sqrt{3}\right)$.

14. (a) $\theta = \frac{\pi}{4}$. In cylindrical coordinates (assuming that r can be negative), this is a vertical plane that includes the z-axis and intersects the xy-plane in the line $y = x$. In spherical coordinates, because $\rho \ge 0$ and $0 \le \phi \le \pi$, we get a vertical half-plane that includes the z-axis and intersects the xy-plane in the half-line $y = x$, $x \ge 0$.

(b) $\phi = \frac{\pi}{4}$. In spherical coordinates, this is one frustum of a circular cone with vertex the origin and axis the positive z-axis.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. (a) $x^2 + y^2 + z^2 = 4$. In cylindrical coordinates, this becomes $r^2 + z^2 = 4$. In spherical coordinates, it becomes $\rho^2 = 4$ or $\rho = 2$.

(b) $x^2 + y^2 = 4$. In cylindrical coordinates: $r^2 = 4$ or $r = 2$. In spherical coordinates: $\rho^2 - z^2 = 4$ or $\rho^2 - \rho^2 \cos^2 \phi = 4$ or $\rho^2 \sin^2 \phi = 4$ or $\rho \sin \phi = 2$.

16. $\rho = 2\cos\phi \;\Rightarrow\; \rho^2 = 2\rho\cos\phi \;\Rightarrow\; x^2 + y^2 + z^2 = 2z \;\Rightarrow$ $x^2 + y^2 + (z-1)^2 = 1$. This is the equation of a sphere with radius 1, centered at $(0, 0, 1)$. Therefore, $0 \le \rho \le 2\cos\phi$ is the solid ball whose boundary is this sphere. $0 \le \theta \le \frac{\pi}{2}$ and $0 \le \phi \le \frac{\pi}{6}$ restrict the solid to the section of this ball that lies above the cone $\phi = \frac{\pi}{6}$ and is in the first octant.

17.

The region whose area is given by $\int_0^{\pi/2} \int_0^{\sin 2\theta} r\, dr\, d\theta$ is $\{(r, \theta) \mid 0 \le \theta \le \frac{\pi}{2}, 0 \le r \le \sin 2\theta\}$, which is the region inside the loop of the four-leaved rose $r = \sin 2\theta$ in the first quadrant.

18. The solid is $\{(\rho, \theta, \phi) \mid 1 \le \rho \le 2,\ 0 \le \theta \le \frac{\pi}{2},\ 0 \le \phi \le \frac{\pi}{2}\}$, which is the region in the first octant on or between the two spheres $\rho = 1$ and $\rho = 2$.

19.

$$\int_0^1 \int_x^1 \cos(y^2)\, dy\, dx = \int_0^1 \int_0^y \cos(y^2)\, dx\, dy$$
$$= \int_0^1 \cos(y^2) \left[x \right]_{x=0}^{x=y} dy = \int_0^1 y \cos(y^2)\, dy$$
$$= \left[\tfrac{1}{2} \sin(y^2) \right]_0^1 = \tfrac{1}{2} \sin 1$$

20.

$$\int_0^1 \int_{\sqrt{y}}^1 \frac{ye^{x^2}}{x^3}\, dx\, dy = \int_0^1 \int_0^{x^2} \frac{ye^{x^2}}{x^3}\, dy\, dx = \int_0^1 \frac{e^{x^2}}{x^3} \left[\tfrac{1}{2} y^2 \right]_{y=0}^{y=x^2} dx$$
$$= \int_0^1 \tfrac{1}{2} x e^{x^2}\, dx = \tfrac{1}{4} e^{x^2} \Big]_0^1 = \tfrac{1}{4}(e - 1)$$

21. $\iint_R ye^{xy}\, dA = \int_0^3 \int_0^2 ye^{xy}\, dx\, dy = \int_0^3 \left[e^{xy} \right]_{x=0}^{x=2} dy = \int_0^3 (e^{2y} - 1)\, dy = \left[\frac{1}{2} e^{2y} - y \right]_0^3 = \frac{1}{2} e^6 - 3 - \frac{1}{2} = \frac{1}{2} e^6 - \frac{7}{2}$

22. $\iint_D xy\, dA = \int_0^1 \int_{y^2}^{y+2} xy\, dx\, dy = \int_0^1 y \left[\frac{1}{2} x^2 \right]_{x=y^2}^{x=y+2} dy = \frac{1}{2} \int_0^1 y \left[(y+2)^2 - y^4 \right] dy$

$$= \tfrac{1}{2} \int_0^1 (y^3 + 4y^2 + 4y - y^5)\, dy = \tfrac{1}{2} \left[\tfrac{1}{4} y^4 + \tfrac{4}{3} y^3 + 2y^2 - \tfrac{1}{6} y^6 \right]_0^1 = \tfrac{41}{24}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23.

$$\iint_D \frac{y}{1+x^2}\,dA = \int_0^1 \int_0^{\sqrt{x}} \frac{y}{1+x^2}\,dy\,dx = \int_0^1 \frac{1}{1+x^2}\left[\tfrac{1}{2}y^2\right]_{y=0}^{y=\sqrt{x}}\,dx$$
$$= \tfrac{1}{2}\int_0^1 \frac{x}{1+x^2}\,dx = \left[\tfrac{1}{4}\ln(1+x^2)\right]_0^1 = \tfrac{1}{4}\ln 2$$

24. $$\iint_D \frac{1}{1+x^2}\,dA = \int_0^1 \int_x^1 \frac{1}{1+x^2}\,dy\,dx = \int_0^1 \frac{1}{1+x^2}\left[\,y\,\right]_{y=x}^{y=1}\,dx = \int_0^1 \frac{1-x}{1+x^2}\,dx = \int_0^1 \left(\frac{1}{1+x^2} - \frac{x}{1+x^2}\right)dx$$
$$= \left[\tan^{-1}x - \tfrac{1}{2}\ln(1+x^2)\right]_0^1 = \tan^{-1}1 - \tfrac{1}{2}\ln 2 - \left(\tan^{-1}0 - \tfrac{1}{2}\ln 1\right) = \tfrac{\pi}{4} - \tfrac{1}{2}\ln 2$$

25.

$$\iint_D y\,dA = \int_0^2 \int_{y^2}^{8-y^2} y\,dx\,dy$$
$$= \int_0^2 y\left[x\right]_{x=y^2}^{x=8-y^2}\,dy = \int_0^2 y(8-y^2-y^2)\,dy$$
$$= \int_0^2 (8y - 2y^3)\,dy = \left[4y^2 - \tfrac{1}{2}y^4\right]_0^2 = 8$$

26.

$$\iint_D y\,dA = \int_1^2 \int_{1/y}^{y} y\,dx\,dy = \int_1^2 y\left(y - \frac{1}{y}\right)dy$$
$$= \int_1^2 \left(y^2 - 1\right)dy = \left[\tfrac{1}{3}y^3 - y\right]_1^2$$
$$= \left(\tfrac{8}{3} - 2\right) - \left(\tfrac{1}{3} - 1\right) = \tfrac{4}{3}$$

27.

$$\iint_D \left(x^2+y^2\right)^{3/2}dA = \int_0^{\pi/3}\int_0^3 (r^2)^{3/2} r\,dr\,d\theta$$
$$= \int_0^{\pi/3} d\theta \int_0^3 r^4\,dr = \left[\,\theta\,\right]_0^{\pi/3}\left[\tfrac{1}{5}r^5\right]_0^3$$
$$= \frac{\pi}{3}\frac{3^5}{5} = \frac{81\pi}{5}$$

28. $$\iint_D x\,dA = \int_0^{\pi/2}\int_1^{\sqrt{2}} (r\cos\theta)\,r\,dr\,d\theta = \int_0^{\pi/2}\cos\theta\,d\theta \int_1^{\sqrt{2}} r^2\,dr = \left[\sin\theta\right]_0^{\pi/2}\left[\tfrac{1}{3}r^3\right]_1^{\sqrt{2}}$$
$$= 1\cdot\tfrac{1}{3}(2^{3/2}-1) = \tfrac{1}{3}(2^{3/2}-1)$$

29. $$\iiint_E xy\,dV = \int_0^3\int_0^x\int_0^{x+y} xy\,dz\,dy\,dx = \int_0^3\int_0^x xy\left[\,z\,\right]_{z=0}^{z=x+y}\,dy\,dx = \int_0^3\int_0^x xy(x+y)\,dy\,dx$$
$$= \int_0^3\int_0^x (x^2y + xy^2)\,dy\,dx = \int_0^3 \left[\tfrac{1}{2}x^2y^2 + \tfrac{1}{3}xy^3\right]_{y=0}^{y=x}\,dx = \int_0^3 \left(\tfrac{1}{2}x^4 + \tfrac{1}{3}x^4\right)dx$$
$$= \tfrac{5}{6}\int_0^3 x^4\,dx = \left[\tfrac{1}{6}x^5\right]_0^3 = \tfrac{81}{2} = 40.5$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $\iiint_T xy\,dV = \int_0^{1/3}\int_0^{1-3x}\int_0^{1-3x-y} xy\,dz\,dy\,dx = \int_0^{1/3}\int_0^{1-3x} xy(1-3x-y)\,dy\,dx$

$= \int_0^{1/3}\int_0^{1-3x}(xy-3x^2y-xy^2)\,dy\,dx$

$= \int_0^{1/3}\left[\frac{1}{2}xy^2-\frac{3}{2}x^2y^2-\frac{1}{3}xy^3\right]_{y=0}^{y=1-3x}dx$

$= \int_0^{1/3}\left[\frac{1}{2}x(1-3x)^2-\frac{3}{2}x^2(1-3x)^2-\frac{1}{3}x(1-3x)^3\right]dx$

$= \int_0^{1/3}\left(\frac{1}{6}x-\frac{3}{2}x^2+\frac{9}{2}x^3-\frac{9}{2}x^4\right)dx$

$= \frac{1}{12}x^2-\frac{1}{2}x^3+\frac{9}{8}x^4-\frac{9}{10}x^5\Big]_0^{1/3} = \frac{1}{1080}$

31. $\iiint_E y^2z^2\,dV = \int_{-1}^{1}\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}}\int_0^{1-y^2-z^2} y^2z^2\,dx\,dz\,dy = \int_{-1}^{1}\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} y^2z^2(1-y^2-z^2)\,dz\,dy$

$= \int_0^{2\pi}\int_0^1 (r^2\cos^2\theta)(r^2\sin^2\theta)(1-r^2)\,r\,dr\,d\theta = \int_0^{2\pi}\left(\frac{1}{2}\sin 2\theta\right)^2 d\theta\ \int_0^1(r^5-r^7)\,dr$

$= \int_0^{2\pi}\frac{1}{4}\left[\frac{1}{2}(1-\cos 4\theta)\right]d\theta\ \int_0^1(r^5-r^7)\,dr = \frac{1}{8}\left[\theta-\frac{1}{4}\sin 4\theta\right]_0^{2\pi}\ \left[\frac{1}{6}r^6-\frac{1}{8}r^8\right]_0^1$

$= \frac{1}{8}(2\pi)\left(\frac{1}{6}-\frac{1}{8}\right) = \frac{\pi}{4}\cdot\frac{1}{24} = \frac{\pi}{96}$

32. $\iiint_E z\,dV = \int_0^1\int_0^{\sqrt{1-y^2}}\int_0^{2-y} z\,dx\,dz\,dy = \int_0^1\int_0^{\sqrt{1-y^2}}(2-y)z\,dz\,dy = \int_0^1\frac{1}{2}(2-y)(1-y^2)\,dy$

$= \int_0^1\frac{1}{2}(2-y-2y^2+y^3)\,dy = \frac{13}{24}$

33. $\iiint_E yz\,dV = \int_{-2}^{2}\int_0^{\sqrt{4-x^2}}\int_0^y yz\,dz\,dy\,dx = \int_{-2}^{2}\int_0^{\sqrt{4-x^2}}\left[\frac{1}{2}yz^2\right]_{z=0}^{z=y}dy\,dx = \frac{1}{2}\int_{-2}^{2}\int_0^{\sqrt{4-x^2}} y^3\,dy\,dx$

$= \frac{1}{2}\int_0^{\pi}\int_0^2(r\sin\theta)^3\,r\,dr\,d\theta = \frac{1}{2}\int_0^{\pi}\sin^3\theta\,d\theta\ \int_0^2 r^4\,dr = \frac{1}{2}\int_0^{\pi}(1-\cos^2\theta)\sin\theta\,d\theta\ \int_0^2 r^4\,dr$

$= \frac{1}{2}\left[-\cos\theta+\frac{1}{3}\cos^3\theta\right]_0^{\pi}\ \left[\frac{1}{5}r^5\right]_0^2 = \frac{1}{2}\left(\frac{2}{3}+\frac{2}{3}\right)\left(\frac{32}{5}\right) = \frac{64}{15}$

34. $\iiint_H z^3\sqrt{x^2+y^2+z^2}\,dV = \int_0^{2\pi}\int_0^{\pi/2}\int_0^1(\rho^3\cos^3\phi)\rho(\rho^2\sin\phi)\,d\rho\,d\phi\,d\theta$

$= \int_0^{2\pi}d\theta\ \int_0^{\pi/2}\cos^3\phi\sin\phi\,d\phi\ \int_0^1\rho^6\,d\rho = 2\pi\left[-\frac{1}{4}\cos^4\phi\right]_0^{\pi/2}\left(\frac{1}{7}\right) = \frac{\pi}{14}$

35. $V = \int_0^2\int_1^4(x^2+4y^2)\,dy\,dx = \int_0^2\left[x^2y+\frac{4}{3}y^3\right]_{y=1}^{y=4}dx = \int_0^2(3x^2+84)\,dx = x^3+84x\Big]_0^2 = 176$

36.

$V = \int_0^1\int_{y+1}^{4-2y}\int_0^{x^2y}\,dz\,dx\,dy = \int_0^1\int_{y+1}^{4-2y}x^2y\,dx\,dy$

$= \int_0^1\frac{1}{3}\left[(4-2y)^3y-(y+1)^3y\right]dy$

$= \int_0^1 3(-y^4+5y^3-11y^2+7y)\,dy = 3\left(-\frac{1}{5}+\frac{5}{4}-\frac{11}{3}+\frac{7}{2}\right) = \frac{53}{20}$

37.

$V = \int_0^2\int_0^y\int_0^{(2-y)/2}\,dz\,dx\,dy = \int_0^2\int_0^y\left(1-\frac{1}{2}y\right)dx\,dy$

$= \int_0^2\left(y-\frac{1}{2}y^2\right)dy = \frac{1}{2}y^2-\frac{1}{6}y^3\Big]_0^2 = \frac{2}{3}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

38. $V = \int_0^{2\pi}\int_0^2\int_0^{3-r\sin\theta} r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^2 \Big[rz\Big]_0^{3-r\sin\theta}\,dr\,d\theta = \int_0^{2\pi}\int_0^2 (3r - r^2\sin\theta)\,dr\,d\theta$

$= \int_0^{2\pi}\left[\frac{3}{2}r^2 - \frac{1}{3}r^3\sin\theta\right]_0^2 d\theta = \int_0^{2\pi}\left[6 - \frac{8}{3}\sin\theta\right]d\theta = \left[6\theta + \frac{8}{3}\cos\theta\right]_0^{2\pi} = 12\pi$

39. Using the wedge above the plane $z = 0$ and below the plane $z = mx$ and noting that we have the same volume for $m < 0$ as for $m > 0$ (so use $m > 0$), we have

$V = 2\int_0^{a/3}\int_0^{\sqrt{a^2-9y^2}} mx\,dx\,dy = 2\int_0^{a/3}\frac{1}{2}m(a^2 - 9y^2)\,dy = m\big[a^2y - 3y^3\big]_0^{a/3} = m\big(\frac{1}{3}a^3 - \frac{1}{9}a^3\big) = \frac{2}{9}ma^3.$

40. The paraboloid and the half-cone intersect when $x^2 + y^2 = \sqrt{x^2+y^2}$, that is when $x^2 + y^2 = 1$ or 0. So

$V = \iint\limits_{x^2+y^2\le 1}\int_{x^2+y^2}^{\sqrt{x^2+y^2}} dz\,dA = \int_0^{2\pi}\int_0^1\int_{r^2}^{r} r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^1 (r^2 - r^3)\,dr\,d\theta = \int_0^{2\pi}\big(\frac{1}{3} - \frac{1}{4}\big)\,d\theta = \frac{1}{12}(2\pi) = \frac{\pi}{6}.$

41. (a) $m = \int_0^1\int_0^{1-y^2} y\,dx\,dy = \int_0^1 (y - y^3)\,dy = \frac{1}{2} - \frac{1}{4} = \frac{1}{4}$

(b) $M_y = \int_0^1\int_0^{1-y^2} xy\,dx\,dy = \int_0^1 \frac{1}{2}y(1-y^2)^2\,dy = -\frac{1}{12}(1-y^2)^3\big]_0^1 = \frac{1}{12}$,

$M_x = \int_0^1\int_0^{1-y^2} y^2\,dx\,dy = \int_0^1 (y^2 - y^4)\,dy = \frac{2}{15}$. Hence $(\overline{x}, \overline{y}) = \big(\frac{1}{3}, \frac{8}{15}\big)$.

(c) $I_x = \int_0^1\int_0^{1-y^2} y^3\,dx\,dy = \int_0^1 (y^3 - y^5)\,dy = \frac{1}{12}$,

$I_y = \int_0^1\int_0^{1-y^2} yx^2\,dx\,dy = \int_0^1 \frac{1}{3}y(1-y^2)^3\,dy = -\frac{1}{24}(1-y^2)^4\big]_0^1 = \frac{1}{24}$,

$\overline{\overline{y}}^2 = I_x/m = \frac{1/12}{1/4} = \frac{1}{3} \quad\Rightarrow\quad \overline{\overline{y}} = \frac{1}{\sqrt{3}}$, and $\overline{\overline{x}}^2 = I_y/m = \frac{1/24}{1/4} = \frac{1}{6} \quad\Rightarrow\quad \overline{\overline{x}} = \frac{1}{\sqrt{6}}$.

42. (a) In polar coordinates, the lamina occupies the region $D = \{(r,\theta) \mid 0 \le r \le a, 0 \le \theta \le \pi/2\}$. Assuming constant density K, then $m = K\,A(D) = K \cdot \frac{1}{4}\pi a^2 = \frac{1}{4}\pi K a^2$,

$M_y = \iint_D Kx\,dA = K\int_0^{\pi/2}\int_0^a (r\cos\theta)\,r\,dr\,d\theta = K\int_0^{\pi/2}\cos\theta\,d\theta\int_0^a r^2\,dr = K\,[\sin\theta]_0^{\pi/2}\left[\frac{1}{3}r^3\right]_0^a = \frac{1}{3}Ka^3$, and

$M_x = \iint_D Ky\,dA = K\int_0^{\pi/2}\sin\theta\,d\theta\int_0^a r^2\,dr = K\,[-\cos\theta]_0^{\pi/2}\left[\frac{1}{3}r^3\right]_0^a = \frac{1}{3}Ka^3$ [by symmetry $M_y = M_x$].

Thus the centroid is $(\overline{x}, \overline{y}) = (M_y/m, M_x/m) = \big(\frac{4}{3\pi}a, \frac{4}{3\pi}a\big)$.

(b) $m = \iint_D \rho(x,y)\,dA = \iint_D xy^2\,dA = \int_0^{\pi/2}\int_0^a (r\cos\theta)(r\sin\theta)^2\,r\,dr\,d\theta$

$= \int_0^{\pi/2}\sin^2\theta\,\cos\theta\,d\theta\int_0^a r^4\,dr = \left[\frac{1}{3}\sin^3\theta\right]_0^{\pi/2}\left[\frac{1}{5}r^5\right]_0^a = \frac{1}{15}a^5$,

$M_y = \int_0^{\pi/2}\int_0^a r^5\cos^2\theta\sin^2\theta\,dr\,d\theta = \frac{1}{8}\left[\theta - \frac{1}{4}\sin 4\theta\right]_0^{\pi/2}\left[\frac{1}{6}r^6\right]_0^a = \frac{1}{96}\pi a^6$, and

$M_x = \int_0^{\pi/2}\int_0^a r^5\cos\theta\sin^3\theta\,dr\,d\theta = \left[\frac{1}{4}\sin^4\theta\right]_0^{\pi/2}\left[\frac{1}{6}r^6\right]_0^a = \frac{1}{24}a^6$. Hence $(\overline{x}, \overline{y}) = \big(\frac{5}{32}\pi a, \frac{5}{8}a\big)$.

43. (a) A right circular cone with axis the z-axis and vertex at the origin has equation $z^2 = c^2(x^2 + y^2)$. Here we have the bottom frustum, shifted upward h units, and with $c^2 = h^2/a^2$ so that the cone includes the point $(a, 0, 0)$. Thus an equation of the cone in rectangular coordinates is $z = h - \frac{h}{a}\sqrt{x^2+y^2}$, $0 \le z \le h$. In cylindrical coordinates, the cone is described by $E = \big\{(r,\theta,z) \mid 0 \le r \le a,\ 0 \le \theta \le 2\pi,\ 0 \le z \le h\left(1 - \frac{1}{a}r\right)\big\}$, and its volume is $V = \frac{1}{3}\pi a^2 h$. By symmetry

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$M_{yz} = M_{xz} = 0$, and

$$\begin{aligned} M_{xy} &= \int_0^{2\pi}\int_0^a\int_0^{h(1-r/a)} z\cdot r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^a \left[\frac{1}{2}rz^2\right]_{z=0}^{z=h(1-r/a)} dr\,d\theta \\ &= \frac{1}{2}\int_0^{2\pi}\int_0^a rh^2\left(1-\frac{r}{a}\right)^2 dr\,d\theta = \frac{1}{2}h^2\int_0^{2\pi} d\theta \int_0^a \left(r - \frac{2}{a}r^2 + \frac{1}{a^2}r^3\right) dr \\ &= \tfrac{1}{2}h^2\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{2}r^2 - \tfrac{2}{3a}r^3 + \tfrac{1}{4a^2}r^4\right]_0^a = \tfrac{1}{2}h^2(2\pi)\left(\tfrac{1}{2}a^2 - \tfrac{2}{3}a^2 + \tfrac{1}{4}a^2\right) \\ &= \pi h^2\left(\tfrac{1}{12}a^2\right) = \tfrac{1}{12}\pi a^2h^2 \end{aligned}$$

Hence the centroid is $(\overline{x},\overline{y},\overline{z}) = \left(0, 0, [\pi a^2h^2/12]/[\pi a^2h/3]\right) = \left(0, 0, \frac{1}{4}h\right)$.

(b) The density function is $\rho = \sqrt{x^2+y^2} = \sqrt{r^2} = r$, so the moment of inertia about the cone's axis (the z-axis) is

$$\begin{aligned} I_z &= \iiint_E (x^2+y^2)\,\rho(x,y,z)\,dV = \int_0^{2\pi}\int_0^a\int_0^{h(1-r/a)} (r^2)(r)\,r\,dz\,dr\,d\theta \\ &= \int_0^{2\pi}\int_0^a \left[r^4z\right]_{z=0}^{z=h(1-r/a)} dr\,d\theta = \int_0^{2\pi}\int_0^a r^4h\left(1-\tfrac{1}{a}r\right) dr\,d\theta \\ &= h\int_0^{2\pi} d\theta \int_0^a \left(r^4 - \tfrac{1}{a}r^5\right) dr = h\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{5}r^5 - \tfrac{1}{6a}r^6\right]_0^a \\ &= h(2\pi)\left(\tfrac{1}{5}a^5 - \tfrac{1}{6}a^5\right) = \tfrac{1}{15}\pi a^5h \end{aligned}$$

44. $1 \le z^2 \le 4 \quad\Rightarrow\quad 1/a^2 \le x^2+y^2 \le 4/a^2$. Let $D = \left\{(x,y) \mid 1/a^2 \le x^2+y^2 \le 4/a^2\right\}$. $z = f(x,y) = a\sqrt{x^2+y^2}$, so $f_x(x,y) = ax(x^2+y^2)^{-1/2}$, $f_y(x,y) = ay(x^2+y^2)^{-1/2}$, and

$$\begin{aligned} A(S) &= \iint_D \sqrt{\frac{a^2x^2+a^2y^2}{x^2+y^2}+1}\,dA = \iint_D \sqrt{a^2+1}\,dA = \sqrt{a^2+1}\,A(D) \\ &= \sqrt{a^2+1}\left[\pi\left(\frac{2}{a}\right)^2 - \pi\left(\frac{1}{a}\right)^2\right] = \frac{3\pi}{a^2}\sqrt{a^2+1} \end{aligned}$$

45. Let D represent the given triangle; then D can be described as the area enclosed by the x- and y-axes and the line $y = 2-2x$, or equivalently $D = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le 2-2x\}$. We want to find the surface area of the part of the graph of $z = x^2+y$ that lies over D, so using Formula 15.5.3 we have

$$\begin{aligned} A(S) &= \iint_D \sqrt{1+\left(\frac{\partial z}{\partial x}\right)^2+\left(\frac{\partial z}{\partial y}\right)^2}\,dA = \iint_D \sqrt{1+(2x)^2+(1)^2}\,dA = \int_0^1\int_0^{2-2x} \sqrt{2+4x^2}\,dy\,dx \\ &= \int_0^1 \sqrt{2+4x^2}\left[y\right]_{y=0}^{y=2-2x} dx = \int_0^1 (2-2x)\sqrt{2+4x^2}\,dx = \int_0^1 2\sqrt{2+4x^2}\,dx - \int_0^1 2x\sqrt{2+4x^2}\,dx \end{aligned}$$

Using Formula 21 in the Table of Integrals with $a = \sqrt{2}$, $u = 2x$, and $du = 2\,dx$, we have

$\int 2\sqrt{2+4x^2}\,dx = x\sqrt{2+4x^2} + \ln\left(2x+\sqrt{2+4x^2}\right)$. If we substitute $u = 2+4x^2$ in the second integral, then $du = 8x\,dx$ and $\int 2x\sqrt{2+4x^2}\,dx = \frac{1}{4}\int\sqrt{u}\,du = \frac{1}{4}\cdot\frac{2}{3}u^{3/2} = \frac{1}{6}(2+4x^2)^{3/2}$. Thus

$$\begin{aligned} A(S) &= \left[x\sqrt{2+4x^2} + \ln\left(2x+\sqrt{2+4x^2}\right) - \tfrac{1}{6}(2+4x^2)^{3/2}\right]_0^1 \\ &= \sqrt{6} + \ln\left(2+\sqrt{6}\right) - \tfrac{1}{6}(6)^{3/2} - \ln\sqrt{2} + \tfrac{\sqrt{2}}{3} = \ln\tfrac{2+\sqrt{6}}{\sqrt{2}} + \tfrac{\sqrt{2}}{3} \\ &= \ln\left(\sqrt{2}+\sqrt{3}\right) + \tfrac{\sqrt{2}}{3} \approx 1.6176 \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. Using Formula 15.5.3 with $\partial z/\partial x = \sin y$, $\partial z/\partial y = x\cos y$, we get

$S = \int_{-\pi}^{\pi}\int_{-3}^{3}\sqrt{1+\sin^2 y + x^2\cos^2 y}\,dx\,dy \approx 62.9714.$

47.

$$\int_0^3\int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}}(x^3+xy^2)\,dy\,dx = \int_0^3\int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} x(x^2+y^2)\,dy\,dx$$
$$= \int_{-\pi/2}^{\pi/2}\int_0^3 (r\cos\theta)(r^2)\,r\,dr\,d\theta$$
$$= \int_{-\pi/2}^{\pi/2}\cos\theta\,d\theta\,\int_0^3 r^4\,dr$$
$$= \left[\sin\theta\right]_{-\pi/2}^{\pi/2}\left[\tfrac{1}{5}r^5\right]_0^3 = 2\cdot\tfrac{1}{5}(243) = \tfrac{486}{5} = 97.2$$

48. The region of integration is the solid hemisphere $x^2+y^2+z^2 \le 4$, $x \ge 0$.

$$\int_{-2}^{2}\int_0^{\sqrt{4-y^2}}\int_{-\sqrt{4-x^2-y^2}}^{\sqrt{4-x^2-y^2}} y^2\sqrt{x^2+y^2+z^2}\,dz\,dx\,dy$$
$$= \int_{-\pi/2}^{\pi/2}\int_0^{\pi}\int_0^2(\rho\sin\phi\sin\theta)^2\left(\sqrt{\rho^2}\right)\rho^2\sin\phi\,d\rho\,d\phi\,d\theta = \int_{-\pi/2}^{\pi/2}\sin^2\theta\,d\theta\,\int_0^{\pi}\sin^3\phi\,d\phi\,\int_0^2\rho^5\,d\rho$$
$$= \left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_{-\pi/2}^{\pi/2}\left[-\cos\phi + \tfrac{1}{3}\cos^3\phi\right]_0^{\pi}\left[\tfrac{1}{6}\rho^6\right]_0^2 = \left(\tfrac{\pi}{2}\right)\left(\tfrac{2}{3}+\tfrac{2}{3}\right)\left(\tfrac{32}{3}\right) = \tfrac{64}{9}\pi$$

49. From the graph, it appears that $1-x^2 = e^x$ at $x \approx -0.71$ and at $x = 0$, with $1-x^2 > e^x$ on $(-0.71, 0)$. So the desired integral is

$$\iint_D y^2\,dA \approx \int_{-0.71}^{0}\int_{e^x}^{1-x^2} y^2\,dy\,dx$$
$$= \tfrac{1}{3}\int_{-0.71}^{0}\left[(1-x^2)^3 - e^{3x}\right]dx$$
$$= \tfrac{1}{3}\left[x - x^3 + \tfrac{3}{5}x^5 - \tfrac{1}{7}x^7 - \tfrac{1}{3}e^{3x}\right]_{-0.71}^{0} \approx 0.0512$$

50. Let the tetrahedron be called T. The front face of T is given by the plane $x + \frac{1}{2}y + \frac{1}{3}z = 1$, or $z = 3 - 3x - \frac{3}{2}y$, which intersects the xy-plane in the line $y = 2 - 2x$. So the total mass is

$m = \iiint_T \rho(x,y,z)\,dV = \int_0^1\int_0^{2-2x}\int_0^{3-3x-3y/2}(x^2+y^2+z^2)\,dz\,dy\,dx = \frac{7}{5}$. The center of mass is

$(\overline{x},\overline{y},\overline{z}) = \left(m^{-1}\iiint_T x\rho(x,y,z)\,dV, m^{-1}\iiint_T y\rho(x,y,z)\,dV, m^{-1}\iiint_T z\rho(x,y,z)\,dV\right) = \left(\frac{4}{21}, \frac{11}{21}, \frac{8}{7}\right)$.

51. (a) $f(x,y)$ is a joint density function, so we know that $\iint_{\mathbb{R}^2} f(x,y)\,dA = 1$. Since $f(x,y) = 0$ outside the rectangle $[0,3]\times[0,2]$, we can say

$$\iint_{\mathbb{R}^2} f(x,y)\,dA = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f(x,y)\,dy\,dx = \int_0^3\int_0^2 C(x+y)\,dy\,dx$$
$$= C\int_0^3\left[xy + \tfrac{1}{2}y^2\right]_{y=0}^{y=2}dx = C\int_0^3(2x+2)\,dx = C\left[x^2+2x\right]_0^3 = 15C$$

Then $15C = 1 \quad\Rightarrow\quad C = \frac{1}{15}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $P(X \le 2, Y \ge 1) = \int_{-\infty}^{2} \int_{1}^{\infty} f(x,y)\,dy\,dx = \int_0^2 \int_1^2 \frac{1}{15}(x,y)\,dy\,dx = \frac{1}{15}\int_0^2 \left[xy + \frac{1}{2}y^2\right]_{y=1}^{y=2} dx$

$$= \frac{1}{15}\int_0^2 \left(x + \frac{3}{2}\right) dx = \frac{1}{15}\left[\frac{1}{2}x^2 + \frac{3}{2}x\right]_0^2 = \frac{1}{3}$$

(c) $P(X + Y \le 1) = P((X,Y) \in D)$ where D is the triangular region shown in the figure. Thus

$$P(X + Y \le 1) = \iint_D f(x,y)\,dA = \int_0^1 \int_0^{1-x} \frac{1}{15}(x+y)\,dy\,dx$$

$$= \frac{1}{15}\int_0^1 \left[xy + \frac{1}{2}y^2\right]_{y=0}^{y=1-x} dx$$

$$= \frac{1}{15}\int_0^1 \left[x(1-x) + \frac{1}{2}(1-x)^2\right] dx$$

$$= \frac{1}{30}\int_0^1 (1 - x^2)\,dx = \frac{1}{30}\left[x - \frac{1}{3}x^3\right]_0^1 = \frac{1}{45}$$

52. Each lamp has exponential density function

$$f(t) = \begin{cases} 0 & \text{if } t < 0 \\ \frac{1}{800}e^{-t/800} & \text{if } t \ge 0 \end{cases}$$

If X, Y, and Z are the lifetimes of the individual bulbs, then X, Y, and Z are independent, so the joint density function is the product of the individual density functions:

$$f(x,y,z) = \begin{cases} \frac{1}{800^3}e^{-(x+y+z)/800} & \text{if } x \ge 0, y \ge 0, z \ge 0 \\ 0 & \text{otherwise} \end{cases}$$

The probability that all three bulbs fail within a total of 1000 hours is $P(X + Y + Z \le 1000)$, or equivalently $P((X,Y,Z) \in E)$ where E is the solid region in the first octant bounded by the coordinate planes and the plane $x + y + z = 1000$. The plane $x + y + z = 1000$ meets the xy-plane in the line $x + y = 1000$, so we have

$$P(X + Y + Z \le 1000) = \iiint_E f(x,y,z)\,dV = \int_0^{1000}\int_0^{1000-x}\int_0^{1000-x-y} \frac{1}{800^3}e^{-(x+y+z)/800}\,dz\,dy\,dx$$

$$= \frac{1}{800^3}\int_0^{1000}\int_0^{1000-x} -800\left[e^{-(x+y+z)/800}\right]_{z=0}^{z=1000-x-y} dy\,dx$$

$$= \frac{-1}{800^2}\int_0^{1000}\int_0^{1000-x}\left[e^{-5/4} - e^{-(x+y)/800}\right] dy\,dx$$

$$= \frac{-1}{800^2}\int_0^{1000}\left[e^{-5/4}y + 800e^{-(x+y)/800}\right]_{y=0}^{y=1000-x} dx$$

$$= \frac{-1}{800^2}\int_0^{1000}\left[e^{-5/4}(1800 - x) - 800e^{-x/800}\right] dx$$

$$= \frac{-1}{800^2}\left[-\frac{1}{2}e^{-5/4}(1800-x)^2 + 800^2 e^{-x/800}\right]_0^{1000}$$

$$= \frac{-1}{800^2}\left[-\frac{1}{2}e^{-5/4}(800)^2 + 800^2 e^{-5/4} + \frac{1}{2}e^{-5/4}(1800)^2 - 800^2\right]$$

$$= 1 - \frac{97}{32}e^{-5/4} \approx 0.1315$$

53.

$$\int_{-1}^{1}\int_{x^2}^{1}\int_0^{1-y} f(x,y,z)\,dz\,dy\,dx = \int_0^1\int_0^{1-z}\int_{-\sqrt{y}}^{\sqrt{y}} f(x,y,z)\,dx\,dy\,dz$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54.

$\int_0^2 \int_0^{y^3} \int_0^{y^2} f(x,y,z)\,dz\,dx\,dy = \iiint_E f(x,y,z)\,dV$ where $E = \{(x,y,z) \mid 0 \le y \le 2, 0 \le x \le y^3, 0 \le z \le y^2\}$.

If D_1, D_2, and D_3 are the projections of E onto the xy-, yz-, and xz-planes, then

$$D_1 = \{(x,y) \mid 0 \le y \le 2, 0 \le x \le y^3\} = \{(x,y) \mid 0 \le x \le 8,\ \sqrt[3]{x} \le y \le 2\},$$

$$D_2 = \{(y,z) \mid 0 \le z \le 4,\ \sqrt{z} \le y \le 2\} = \{(y,z) \mid 0 \le y \le 2, 0 \le z \le y^2\},$$

$$D_3 = \{(x,z) \mid 0 \le x \le 8, 0 \le z \le 4\}.$$

Therefore we have

$$\begin{aligned}
\int_0^2 \int_0^{y^3} \int_0^{y^2} f(x,y,z)\,dz\,dx\,dy &= \int_0^8 \int_{\sqrt[3]{x}}^2 \int_0^{y^2} f(x,y,z)\,dz\,dy\,dx \\
&= \int_0^4 \int_{\sqrt{z}}^2 \int_0^{y^3} f(x,y,z)\,dx\,dy\,dz \\
&= \int_0^2 \int_0^{y^2} \int_0^{y^3} f(x,y,z)\,dx\,dz\,dy \\
&= \int_0^8 \int_0^{x^{2/3}} \int_{\sqrt[3]{x}}^2 f(x,y,z)\,dy\,dz\,dx + \int_0^8 \int_{x^{2/3}}^4 \int_{\sqrt{z}}^2 f(x,y,z)\,dy\,dz\,dx \\
&= \int_0^4 \int_0^{z^{3/2}} \int_{\sqrt{z}}^2 f(x,y,z)\,dy\,dx\,dz + \int_0^4 \int_{z^{3/2}}^8 \int_{\sqrt[3]{x}}^2 f(x,y,z)\,dy\,dx\,dz
\end{aligned}$$

55. Since $u = x - y$ and $v = x + y$, $x = \frac{1}{2}(u+v)$ and $y = \frac{1}{2}(v-u)$. Then $\dfrac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} 1/2 & 1/2 \\ -1/2 & 1/2 \end{vmatrix} = \dfrac{1}{2}$.

R is the image under this transformation of the square with vertices $(u,v) = (-2,2)$, $(0,2)$, $(0,4)$, and $(-2,4)$. Thus,

$$\begin{aligned}
\iint_R \frac{x-y}{x+y}\,dA &= \int_2^4 \int_{-2}^0 \frac{u}{v}\left(\frac{1}{2}\right) du\,dv = \frac{1}{2}\int_2^4 \left[\frac{u^2}{2v}\right]_{u=-2}^{u=0} dv = \frac{1}{2}\int_2^4 \left(-\frac{2}{v}\right) dv \\
&= \Big[-\ln v\Big]_2^4 = -\ln 4 + \ln 2 = -2\ln 2 + \ln 2 = -\ln 2
\end{aligned}$$

56. $\dfrac{\partial(x,y,z)}{\partial(u,v,w)} = \begin{vmatrix} 2u & 0 & 0 \\ 0 & 2v & 0 \\ 0 & 0 & 2w \end{vmatrix} = 8uvw$, so

$$\begin{aligned}
V = \iiint_E dV &= \int_0^1 \int_0^{1-u} \int_0^{1-u-v} 8uvw\,dw\,dv\,du = \int_0^1 \int_0^{1-u} 4uv(1-u-v)^2\,du \\
&= \int_0^1 \int_0^{1-u} \left[4u(1-u)^2 v - 8u(1-u)v^2 + 4uv^3\right] dv\,du \\
&= \int_0^1 \left[2u(1-u)^4 - \tfrac{8}{3}u(1-u)^4 + u(1-u)^4\right] du = \int_0^1 \tfrac{1}{3}u(1-u)^4 du \\
&= \int_0^1 \tfrac{1}{3}\left[(1-u)^4 - (1-u)^5\right] du = \tfrac{1}{3}\left[-\tfrac{1}{5}(1-u)^5 + \tfrac{1}{6}(1-u)^6\right]_0^1 = \tfrac{1}{3}\left(-\tfrac{1}{6} + \tfrac{1}{5}\right) = \tfrac{1}{90}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. See the figure. Letting $u = -x + y$ and $v = x + y$, we have $x = \frac{1}{2}(v - u)$ and $y = \frac{1}{2}(v + u)$. Then

$$\left|\frac{\partial(x,y)}{\partial(u,v)}\right| = \left|\frac{\partial x}{\partial u}\frac{\partial y}{\partial v} - \frac{\partial x}{\partial v}\frac{\partial y}{\partial u}\right| = \left|-\tfrac{1}{2}\left(\tfrac{1}{2}\right) - \tfrac{1}{2}\left(\tfrac{1}{2}\right)\right| = \left|-\tfrac{1}{2}\right| = \tfrac{1}{2}.$$

R is the image under this transformation of the square region with vertices $(u, v) = (0, 0)$, $(-2, 0)$, $(0, 2)$, and $(-2, 2)$. Thus,

$$\iint_R xy\,dA = \int_0^2 \int_{-2}^0 \frac{v^2 - u^2}{4}\left(\frac{1}{2}\right) du\,dv = \frac{1}{8}\int_0^2 \left[v^2 u - \frac{1}{3}u^3\right]_{u=-2}^{u=0} dv$$

$$= \frac{1}{8}\int_0^2 \left(2v^2 - \frac{8}{3}\right) dv = \frac{1}{8}\left[\frac{2}{3}v^3 - \frac{8}{3}v\right]_0^2 = 0$$

This result could have been anticipated by symmetry, since the integrand is an odd function of y and R is symmetric about the x-axis.

58. (a) $$\iint_D \frac{1}{(x^2+y^2)^{n/2}}\,dA = \int_0^{2\pi}\int_r^R \frac{1}{(t^2)^{n/2}} t\,dt\,d\theta = 2\pi\int_r^R t^{1-n}\,dt$$

$$= \begin{cases} \dfrac{2\pi}{2-n}t^{2-n}\Big]_r^R = \dfrac{2\pi}{2-n}\left(R^{2-n} - r^{2-n}\right) & \text{if } n \neq 2 \\ 2\pi\ln(R/r) & \text{if } n = 2 \end{cases}$$

(b) The integral in part (a) has a limit as $r \to 0^+$ for all values of n such that $2 - n > 0 \quad \Leftrightarrow \quad n < 2$.

(c) $$\iiint_E \frac{1}{(x^2+y^2+z^2)^{n/2}}\,dV = \int_r^R\int_0^\pi\int_0^{2\pi} \frac{1}{(\rho^2)^{n/2}}\rho^2\sin\phi\,d\theta\,d\phi\,d\rho = 2\pi\int_r^R\int_0^\pi \rho^{2-n}\sin\phi\,d\phi\,d\rho$$

$$= \begin{cases} \dfrac{4\pi}{3-n}\rho^{3-n}\Big]_r^R = \dfrac{4\pi}{3-n}\left(R^{3-n} - r^{3-n}\right) & \text{if } n \neq 3 \\ 4\pi\ln(R/r) & \text{if } n = 3 \end{cases}$$

(d) As $r \to 0^+$, the above integral has a limit, provided that $3 - n > 0 \quad \Leftrightarrow \quad n < 3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

PROBLEMS PLUS

1.

Let $R = \bigcup_{i=1}^{5} R_i$, where

$R_i = \{(x, y) \mid x + y \geq i + 2, x + y < i + 3, 1 \leq x \leq 3, 2 \leq y \leq 5\}$.

$\iint_R [\![x + y]\!]\, dA = \sum\limits_{i=1}^{5} \iint_{R_i} [\![x + y]\!]\, dA = \sum\limits_{i=1}^{5} [\![x + y]\!] \iint_{R_i} dA$, since

$[\![x + y]\!] = \text{constant} = i + 2$ for $(x, y) \in R_i$. Therefore

$$\begin{aligned}\iint_R [\![x + y]\!]\, dA &= \textstyle\sum_{i=1}^{5} (i + 2)\,[A(R_i)] \\ &= 3A(R_1) + 4A(R_2) + 5A(R_3) + 6A(R_4) + 7A(R_5) \\ &= 3\left(\tfrac{1}{2}\right) + 4\left(\tfrac{3}{2}\right) + 5(2) + 6\left(\tfrac{3}{2}\right) + 7\left(\tfrac{1}{2}\right) = 30\end{aligned}$$

2.

Let $R = \{(x, y) \mid 0 \leq x, y \leq 1\}$. For $x, y \in R$, $\max\{x^2, y^2\} = x^2$ if $x \geq y$, and $\max\{x^2, y^2\} = y^2$ if $x \leq y$. Therefore we divide R into two regions: $R = R_1 \cup R_2$, where $R_1 = \{(x, y) \mid 0 \leq x \leq 1, 0 \leq y \leq x\}$ and $R_2 = \{(x, y) \mid 0 \leq y \leq 1, 0 \leq x \leq y\}$. Now $\max\{x^2, y^2\} = x^2$ for $(x, y) \in R_1$, and $\max\{x^2, y^2\} = y^2$ for $(x, y) \in R_2 \quad \Rightarrow$

$$\begin{aligned}\int_0^1\!\int_0^1 e^{\max\{x^2,y^2\}}\, dy\, dx &= \iint_R e^{\max\{x^2,y^2\}}\, dA = \iint_{R_1} e^{\max\{x^2,y^2\}}\, dA + \iint_{R_2} e^{\max\{x^2,y^2\}}\, dA \\ &= \int_0^1\!\int_0^x e^{x^2}\, dy\, dx + \int_0^1\!\int_0^y e^{y^2}\, dx\, dy = \int_0^1 xe^{x^2}\, dx + \int_0^1 ye^{y^2}\, dy = e^{x^2}\Big]_0^1 = e - 1\end{aligned}$$

3. $f_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b f(x)\, dx = \dfrac{1}{1-0}\int_0^1 \left[\int_x^1 \cos(t^2)\, dt\right] dx$

$= \int_0^1\!\int_x^1 \cos(t^2)\, dt\, dx = \int_0^1\!\int_0^t \cos(t^2)\, dx\, dt$ [changing the order of integration]

$= \int_0^1 t\cos(t^2)\, dt = \frac{1}{2}\sin(t^2)\big]_0^1 = \frac{1}{2}\sin 1$

4. To show that $\int_0^2\!\int_0^x 2e^{x^2-y^2}\, dy\, dx = \int_0^2\!\int_0^x 2e^{(x+y)(x-y)}\, dy\, dx$ is equal to $\int_0^2\!\int_y^{4-y} e^{xy}\, dx\, dy$, we use a change of variables on the left-hand side. Leting $u = x + y$ and $v = x - y$, we have $x = \frac{1}{2}(u + v)$ and $y = \frac{1}{2}(u - v)$. Then

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} 1/2 & 1/2 \\ 1/2 & -1/2 \end{vmatrix} = -\frac{1}{2}$$

R is bounded by $y = x$, $x = 2$, and $y = 0$.

$y = x \quad \Rightarrow \quad \frac{1}{2}(u - v) = \frac{1}{2}(u + v) \quad \Rightarrow \quad v = 0$;

$x = 2 \quad \Rightarrow \quad \frac{1}{2}(u + v) = 2 \quad \Rightarrow$

$v = 4 - u$, or $u = 4 - v$;

$y = 0 \quad \Rightarrow \quad \frac{1}{2}(u - v) = 0 \quad \Rightarrow \quad u = v$.

So R is the image of S. Thus,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\int_0^2 \int_0^x 2e^{x^2-y^2}\,dy\,dx = \int_0^2 \int_0^x 2e^{(x+y)(x-y)}\,dy\,dx = \int_0^2 \int_v^{4-v} 2e^{uv}\left|-\frac{1}{2}\right|\,du\,dv$$

$$= \int_0^2 \int_v^{4-v} e^{uv}\,du\,dv,$$

which is the same as $\int_0^2 \int_y^{4-y} e^{xy}\,dx\,dy$.

5. Since $|xy| < 1$, except at $(1, 1)$, the formula for the sum of a geometric series gives $\dfrac{1}{1-xy} = \sum\limits_{n=0}^{\infty} (xy)^n$, so

$$\int_0^1 \int_0^1 \frac{1}{1-xy}\,dx\,dy = \int_0^1 \int_0^1 \sum_{n=0}^{\infty} (xy)^n\,dx\,dy = \sum_{n=0}^{\infty} \int_0^1 \int_0^1 (xy)^n\,dx\,dy$$

$$= \sum_{n=0}^{\infty} \left[\int_0^1 x^n\,dx\right]\left[\int_0^1 y^n\,dy\right] = \sum_{n=0}^{\infty} \frac{1}{n+1} \cdot \frac{1}{n+1}$$

$$= \sum_{n=0}^{\infty} \frac{1}{(n+1)^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \cdots = \sum_{n=1}^{\infty} \frac{1}{n^2}$$

6. Let $x = \dfrac{u-v}{\sqrt{2}}$ and $y = \dfrac{u+v}{\sqrt{2}}$. We know the region of integration in the xy-plane, so to find its image in the uv-plane we get u and v in terms of x and y, and then use the methods of Section 15.9. $x + y = \dfrac{u-v}{\sqrt{2}} + \dfrac{u+v}{\sqrt{2}} = \sqrt{2}\,u$, so $u = \dfrac{x+y}{\sqrt{2}}$, and similarly $v = \dfrac{y-x}{\sqrt{2}}$. S_1 is given by $y = 0$, $0 \le x \le 1$, so from the equations derived above, the image of S_1 is S_1': $u = \frac{1}{\sqrt{2}}x$, $v = -\frac{1}{\sqrt{2}}x$, $0 \le x \le 1$, that is, $v = -u$, $0 \le u \le \frac{1}{\sqrt{2}}$. Similarly, the image of S_2 is S_2': $v = u - \sqrt{2}$, $\frac{1}{\sqrt{2}} \le u \le \sqrt{2}$, the image of S_3 is S_3': $v = \sqrt{2} - u$, $\frac{1}{\sqrt{2}} \le u \le \sqrt{2}$, and the image of S_4 is S_4': $v - u$, $0 \le u \le \frac{1}{\sqrt{2}}$.

The Jacobian of the transformation is

$$\frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \partial x/\partial u & \partial x/\partial v \\ \partial y/\partial u & \partial y/\partial v \end{vmatrix} = \begin{vmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{vmatrix} = 1$$

From the diagram, we see that we must evaluate two integrals: one over the region $\left\{(u,v) \mid 0 \le u \le \frac{1}{\sqrt{2}},\ -u \le v \le u\right\}$ and the other over $\left\{(u,v) \mid \frac{1}{\sqrt{2}} \le u \le \sqrt{2},\ -\sqrt{2}+u \le v \le \sqrt{2}-u\right\}$. So

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\int_0^1\int_0^1 \frac{dx\,dy}{1-xy} = \int_0^{\sqrt{2}/2}\int_{-u}^{u} \frac{dv\,du}{1-\left[\frac{1}{\sqrt{2}}(u+v)\right]\left[\frac{1}{\sqrt{2}}(u-v)\right]} + \int_{\sqrt{2}/2}^{\sqrt{2}}\int_{-\sqrt{2}+u}^{\sqrt{2}-u} \frac{dv\,du}{1-\left[\frac{1}{\sqrt{2}}(u+v)\right]\left[\frac{1}{\sqrt{2}}(u-v)\right]}$$

$$= \int_0^{\sqrt{2}/2}\int_{-u}^{u} \frac{2\,dv\,du}{2-u^2+v^2} + \int_{\sqrt{2}/2}^{\sqrt{2}}\int_{-\sqrt{2}+u}^{\sqrt{2}-u} \frac{2\,dv\,du}{2-u^2+v^2}$$

$$= 2\left[\int_0^{\sqrt{2}/2} \frac{1}{\sqrt{2-u^2}}\left[\arctan\frac{v}{\sqrt{2-u^2}}\right]_{-u}^{u} du + \int_{\sqrt{2}/2}^{\sqrt{2}} \frac{1}{\sqrt{2-u^2}}\left[\arctan\frac{v}{\sqrt{2-u^2}}\right]_{-\sqrt{2}+u}^{\sqrt{2}-u} du\right]$$

$$= 4\left[\int_0^{\sqrt{2}/2} \frac{1}{\sqrt{2-u^2}}\arctan\frac{u}{\sqrt{2-u^2}}\,du + \int_{\sqrt{2}/2}^{\sqrt{2}} \frac{1}{\sqrt{2-u^2}}\arctan\frac{\sqrt{2}-u}{\sqrt{2-u^2}}\,du\right]$$

Now let $u=\sqrt{2}\sin\theta$, so $du=\sqrt{2}\cos\theta\,d\theta$ and the limits change to 0 and $\frac{\pi}{6}$ (in the first integral) and $\frac{\pi}{6}$ and $\frac{\pi}{2}$ (in the second integral). Continuing:

$$\int_0^1\int_0^1 \frac{dx\,dy}{1-xy} = 4\left[\int_0^{\pi/6} \frac{1}{\sqrt{2-2\sin^2\theta}}\arctan\left(\frac{\sqrt{2}\sin\theta}{\sqrt{2-2\sin^2\theta}}\right)\left(\sqrt{2}\cos\theta\,d\theta\right)\right.$$
$$\left. + \int_{\pi/6}^{\pi/2} \frac{1}{\sqrt{2-2\sin^2\theta}}\arctan\left(\frac{\sqrt{2}-\sqrt{2}\sin\theta}{\sqrt{2-2\sin^2\theta}}\right)\left(\sqrt{2}\cos\theta\,d\theta\right)\right]$$

$$= 4\left[\int_0^{\pi/6} \frac{\sqrt{2}\cos\theta}{\sqrt{2}\cos\theta}\arctan\left(\frac{\sqrt{2}\sin\theta}{\sqrt{2}\cos\theta}\right)d\theta + \int_{\pi/6}^{\pi/2} \frac{\sqrt{2}\cos\theta}{\sqrt{2}\cos\theta}\arctan\left(\frac{\sqrt{2}\,(1-\sin\theta)}{\sqrt{2}\cos\theta}\right)d\theta\right]$$

$$= 4\left[\int_0^{\pi/6} \arctan(\tan\theta)\,d\theta + \int_{\pi/6}^{\pi/2} \arctan\left(\frac{1-\sin\theta}{\cos\theta}\right)d\theta\right]$$

But (following the hint)

$$\frac{1-\sin\theta}{\cos\theta} = \frac{1-\cos\left(\frac{\pi}{2}-\theta\right)}{\sin\left(\frac{\pi}{2}-\theta\right)} = \frac{1-\left[1-2\sin^2\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)\right]}{2\sin\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)\cos\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)} \qquad \text{[half-angle formulas]}$$
$$= \frac{2\sin^2\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)}{2\sin\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)\cos\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)} = \tan\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)$$

Continuing:

$$\int_0^1\int_0^1 \frac{dx\,dy}{1-xy} = 4\left[\int_0^{\pi/6} \arctan(\tan\theta)\,d\theta + \int_{\pi/6}^{\pi/2} \arctan\left(\tan\left(\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right)\right)d\theta\right]$$

$$= 4\left[\int_0^{\pi/6} \theta\,d\theta + \int_{\pi/6}^{\pi/2}\left[\frac{1}{2}\left(\frac{\pi}{2}-\theta\right)\right]d\theta\right] = 4\left(\left[\frac{\theta^2}{2}\right]_0^{\pi/6} + \left[\frac{\pi\theta}{4}-\frac{\theta^2}{4}\right]_{\pi/6}^{\pi/2}\right) = 4\left(\frac{3\pi^2}{72}\right) = \frac{\pi^2}{6}$$

7. (a) Since $|xyz|<1$ except at $(1,1,1)$, the formula for the sum of a geometric series gives $\dfrac{1}{1-xyz} = \sum\limits_{n=0}^{\infty}(xyz)^n$, so

$$\int_0^1\int_0^1\int_0^1 \frac{1}{1-xyz}\,dx\,dy\,dz = \int_0^1\int_0^1\int_0^1 \sum_{n=0}^{\infty}(xyz)^n\,dx\,dy\,dz = \sum_{n=0}^{\infty}\int_0^1\int_0^1\int_0^1 (xyz)^n\,dx\,dy\,dz$$

$$= \sum_{n=0}^{\infty}\left[\int_0^1 x^n\,dx\right]\left[\int_0^1 y^n\,dy\right]\left[\int_0^1 z^n\,dz\right] = \sum_{n=0}^{\infty}\frac{1}{n+1}\cdot\frac{1}{n+1}\cdot\frac{1}{n+1}$$

$$= \sum_{n=0}^{\infty}\frac{1}{(n+1)^3} = \frac{1}{1^3}+\frac{1}{2^3}+\frac{1}{3^3}+\cdots = \sum_{n=1}^{\infty}\frac{1}{n^3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Since $|-xyz| < 1$, except at $(1, 1, 1)$, the formula for the sum of a geometric series gives $\dfrac{1}{1+xyz} = \sum\limits_{n=0}^{\infty} (-xyz)^n$, so

$$\int_0^1 \int_0^1 \int_0^1 \frac{1}{1+xyz}\,dx\,dy\,dz = \int_0^1 \int_0^1 \int_0^1 \sum_{n=0}^{\infty} (-xyz)^n\,dx\,dy\,dz = \sum_{n=0}^{\infty} \int_0^1 \int_0^1 \int_0^1 (-xyz)^n\,dx\,dy\,dz$$

$$= \sum_{n=0}^{\infty} (-1)^n \left[\int_0^1 x^n\,dx\right]\left[\int_0^1 y^n\,dy\right]\left[\int_0^1 z^n\,dz\right] = \sum_{n=0}^{\infty} (-1)^n \frac{1}{n+1} \cdot \frac{1}{n+1} \cdot \frac{1}{n+1}$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{(n+1)^3} = \frac{1}{1^3} - \frac{1}{2^3} + \frac{1}{3^3} - \cdots = \sum_{n=0}^{\infty} \frac{(-1)^{n-1}}{n^3}$$

To evaluate this sum, we first write out a few terms: $s = 1 - \dfrac{1}{2^3} + \dfrac{1}{3^3} - \dfrac{1}{4^3} + \dfrac{1}{5^3} - \dfrac{1}{6^3} \approx 0.8998$. Notice that $a_7 = \dfrac{1}{7^3} < 0.003$. By the Alternating Series Estimation Theorem from Section 11.5, we have $|s - s_6| \le a_7 < 0.003$. This error of 0.003 will not affect the second decimal place, so we have $s \approx 0.90$.

8. $$\int_0^{\infty} \frac{\arctan \pi x - \arctan x}{x}\,dx = \int_0^{\infty} \left[\frac{\arctan yx}{x}\right]_{y=1}^{y=\pi} dx = \int_0^{\infty} \int_1^{\pi} \frac{1}{1+y^2x^2}\,dy\,dx$$

$$= \int_1^{\pi} \int_0^{\infty} \frac{1}{1+y^2x^2}\,dx\,dy = \int_1^{\pi} \lim_{t\to\infty} \left[\frac{\arctan yx}{y}\right]_{x=0}^{x=t} dy$$

$$= \int_1^{\pi} \frac{\pi}{2y}\,dy = \frac{\pi}{2}\left[\ln y\right]_1^{\pi} = \frac{\pi}{2} \ln \pi$$

9. (a) $x = r\cos\theta,\ y = r\sin\theta,\ z = z.$ Then $\dfrac{\partial u}{\partial r} = \dfrac{\partial u}{\partial x}\dfrac{\partial x}{\partial r} + \dfrac{\partial u}{\partial y}\dfrac{\partial y}{\partial r} + \dfrac{\partial u}{\partial z}\dfrac{\partial z}{\partial r} = \dfrac{\partial u}{\partial x}\cos\theta + \dfrac{\partial u}{\partial y}\sin\theta$ and

$$\frac{\partial^2 u}{\partial r^2} = \cos\theta \left[\frac{\partial^2 u}{\partial x^2}\frac{\partial x}{\partial r} + \frac{\partial^2 u}{\partial y\,\partial x}\frac{\partial y}{\partial r} + \frac{\partial^2 u}{\partial z\,\partial x}\frac{\partial z}{\partial r}\right] + \sin\theta \left[\frac{\partial^2 u}{\partial y^2}\frac{\partial y}{\partial r} + \frac{\partial^2 u}{\partial x\,\partial y}\frac{\partial x}{\partial r} + \frac{\partial^2 u}{\partial z\,\partial y}\frac{\partial z}{\partial r}\right]$$

$$= \frac{\partial^2 u}{\partial x^2}\cos^2\theta + \frac{\partial^2 u}{\partial y^2}\sin^2\theta + 2\,\frac{\partial^2 u}{\partial y\,\partial x}\cos\theta\,\sin\theta$$

Similarly $\dfrac{\partial u}{\partial \theta} = -\dfrac{\partial u}{\partial x}\,r\sin\theta + \dfrac{\partial u}{\partial y}\,r\cos\theta$ and

$\dfrac{\partial^2 u}{\partial \theta^2} = \dfrac{\partial^2 u}{\partial x^2}\,r^2\sin^2\theta + \dfrac{\partial^2 u}{\partial y^2}\,r^2\cos^2\theta - 2\,\dfrac{\partial^2 u}{\partial y\,\partial x}\,r^2\sin\theta\,\cos\theta - \dfrac{\partial u}{\partial x}\,r\cos\theta - \dfrac{\partial u}{\partial y}\,r\sin\theta$. So

$$\frac{\partial^2 u}{\partial r^2} + \frac{1}{r}\frac{\partial u}{\partial r} + \frac{1}{r^2}\frac{\partial^2 u}{\partial \theta^2} + \frac{\partial^2 u}{\partial z^2} = \frac{\partial^2 u}{\partial x^2}\cos^2\theta + \frac{\partial^2 u}{\partial y^2}\sin^2\theta + 2\,\frac{\partial^2 u}{\partial y\,\partial x}\cos\theta\,\sin\theta + \frac{\partial u}{\partial x}\frac{\cos\theta}{r} + \frac{\partial u}{\partial y}\frac{\sin\theta}{r}$$

$$+ \frac{\partial^2 u}{\partial x^2}\sin^2\theta + \frac{\partial^2 u}{\partial y^2}\cos^2\theta - 2\,\frac{\partial^2 u}{\partial y\,\partial x}\sin\theta\,\cos\theta$$

$$-\frac{\partial u}{\partial x}\frac{\cos\theta}{r} - \frac{\partial u}{\partial y}\frac{\sin\theta}{r} + \frac{\partial^2 u}{\partial z^2}$$

$$= \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2}$$

(b) $x = \rho\sin\phi\cos\theta$, $y = \rho\sin\phi\sin\theta$, $z = \rho\cos\phi$. Then

$\dfrac{\partial u}{\partial \rho} = \dfrac{\partial u}{\partial x}\dfrac{\partial x}{\partial \rho} + \dfrac{\partial u}{\partial y}\dfrac{\partial y}{\partial \rho} + \dfrac{\partial u}{\partial z}\dfrac{\partial z}{\partial \rho} = \dfrac{\partial u}{\partial x}\sin\phi\,\cos\theta + \dfrac{\partial u}{\partial y}\sin\phi\,\sin\theta + \dfrac{\partial u}{\partial z}\cos\phi$, and

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$\begin{aligned}\frac{\partial^2 u}{\partial \rho^2} &= \sin\phi\cos\theta\left[\frac{\partial^2 u}{\partial x^2}\frac{\partial x}{\partial \rho}+\frac{\partial^2 u}{\partial y\,\partial x}\frac{\partial y}{\partial \rho}+\frac{\partial^2 u}{\partial z\,\partial x}\frac{\partial z}{\partial \rho}\right]\\ &\quad+\sin\phi\sin\theta\left[\frac{\partial^2 u}{\partial y^2}\frac{\partial y}{\partial \rho}+\frac{\partial^2 u}{\partial x\,\partial y}\frac{\partial x}{\partial \rho}+\frac{\partial^2 u}{\partial z\,\partial y}\frac{\partial z}{\partial \rho}\right]\\ &\quad+\cos\phi\left[\frac{\partial^2 u}{\partial z^2}\frac{\partial z}{\partial \rho}+\frac{\partial^2 u}{\partial x\,\partial z}\frac{\partial x}{\partial \rho}+\frac{\partial^2 u}{\partial y\,\partial z}\frac{\partial y}{\partial \rho}\right]\\ &= 2\,\frac{\partial^2 u}{\partial y\,\partial x}\sin^2\phi\,\sin\theta\,\cos\theta+2\,\frac{\partial^2 u}{\partial z\,\partial x}\sin\phi\,\cos\phi\,\cos\theta+2\,\frac{\partial^2 u}{\partial y\,\partial z}\sin\phi\,\cos\phi\,\sin\theta\\ &\quad+\frac{\partial^2 u}{\partial x^2}\sin^2\phi\cos^2\theta+\frac{\partial^2 u}{\partial y^2}\sin^2\phi\sin^2\theta+\frac{\partial^2 u}{\partial z^2}\cos^2\phi\end{aligned}$$

Similarly $\dfrac{\partial u}{\partial \phi}=\dfrac{\partial u}{\partial x}\,\rho\cos\phi\cos\theta+\dfrac{\partial u}{\partial y}\,\rho\cos\phi\sin\theta-\dfrac{\partial u}{\partial z}\,\rho\sin\phi$, and

$$\begin{aligned}\frac{\partial^2 u}{\partial \phi^2} &= 2\,\frac{\partial^2 u}{\partial y\,\partial x}\,\rho^2\cos^2\phi\,\sin\theta\,\cos\theta-2\,\frac{\partial^2 u}{\partial x\,\partial z}\,\rho^2\sin\phi\,\cos\phi\,\cos\theta\\ &\quad-2\,\frac{\partial^2 u}{\partial y\,\partial z}\,\rho^2\sin\phi\,\cos\phi\,\sin\theta+\frac{\partial^2 u}{\partial x^2}\,\rho^2\cos^2\phi\,\cos^2\theta+\frac{\partial^2 u}{\partial y^2}\,\rho^2\cos^2\phi\,\sin^2\theta\\ &\quad+\frac{\partial^2 u}{\partial z^2}\,\rho^2\sin^2\phi-\frac{\partial u}{\partial x}\,\rho\sin\phi\,\cos\theta-\frac{\partial u}{\partial y}\,\rho\sin\phi\,\sin\theta-\frac{\partial u}{\partial z}\,\rho\cos\phi\end{aligned}$$

And $\dfrac{\partial u}{\partial \theta}=-\dfrac{\partial u}{\partial x}\,\rho\sin\phi\sin\theta+\dfrac{\partial u}{\partial y}\,\rho\sin\phi\cos\theta$, while

$$\begin{aligned}\frac{\partial^2 u}{\partial \theta^2} &= -2\,\frac{\partial^2 u}{\partial y\,\partial x}\,\rho^2\sin^2\phi\,\cos\theta\,\sin\theta+\frac{\partial^2 u}{\partial x^2}\,\rho^2\sin^2\phi\,\sin^2\theta\\ &\quad+\frac{\partial^2 u}{\partial y^2}\,\rho^2\sin^2\phi\cos^2\theta-\frac{\partial u}{\partial x}\,\rho\sin\phi\,\cos\theta-\frac{\partial u}{\partial y}\,\rho\sin\phi\,\sin\theta\end{aligned}$$

Therefore

$$\begin{aligned}&\frac{\partial^2 u}{\partial \rho^2}+\frac{2}{\rho}\frac{\partial u}{\partial \rho}+\frac{\cot\phi}{\rho^2}\frac{\partial u}{\partial \phi}+\frac{1}{\rho^2}\frac{\partial^2 u}{\partial \phi^2}+\frac{1}{\rho^2\sin^2\phi}\frac{\partial^2 u}{\partial \theta^2}\\ &\quad=\frac{\partial^2 u}{\partial x^2}\left[(\sin^2\phi\,\cos^2\theta)+(\cos^2\phi\,\cos^2\theta)+\sin^2\theta\right]\\ &\qquad+\frac{\partial^2 u}{\partial y^2}\left[(\sin^2\phi\,\sin^2\theta)+(\cos^2\phi\,\sin^2\theta)+\cos^2\theta\right]+\frac{\partial^2 u}{\partial z^2}\left[\cos^2\phi+\sin^2\phi\right]\\ &\qquad+\frac{\partial u}{\partial x}\left[\frac{2\sin^2\phi\,\cos\theta+\cos^2\phi\,\cos\theta-\sin^2\phi\,\cos\theta-\cos\theta}{\rho\sin\phi}\right]\\ &\qquad+\frac{\partial u}{\partial y}\left[\frac{2\sin^2\phi\,\sin\theta+\cos^2\phi\,\sin\theta-\sin^2\phi\,\sin\theta-\sin\theta}{\rho\sin\phi}\right]\end{aligned}$$

But

$$2\sin^2\phi\cos\theta+\cos^2\phi\cos\theta-\sin^2\phi\cos\theta-\cos\theta=(\sin^2\phi+\cos^2\phi-1)\,\cos\theta=0$$

and similarly the coefficient of $\partial u/\partial y$ is 0. Also

$$\sin^2\phi\cos^2\theta+\cos^2\phi\cos^2\theta+\sin^2\theta=\cos^2\theta\,(\sin^2\phi+\cos^2\phi)+\sin^2\theta=1$$

and similarly the coefficient of $\partial^2 u/\partial y^2$ is 1. So Laplace's Equation in spherical coordinates is as stated.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. (a) Consider a polar division of the disk, similar to that in Figure 15.3.4, where $0 = \theta_0 < \theta_1 < \theta_2 < \cdots < \theta_n = 2\pi$, $0 = r_1 < r_2 < \cdots < r_m = R$, and where the polar subrectangle R_{ij}, as well as r_i^*, θ_j^*, Δr and $\Delta\theta$ are the same as in that figure. Thus $\Delta A_i = r_i^* \, \Delta r \, \Delta\theta$. The mass of R_{ij} is $\rho \, \Delta A_i$, and its distance from m is $s_{ij} \approx \sqrt{(r_i^*)^2 + d^2}$. According to Newton's Law of Gravitation, the force of attraction experienced by m due to this polar subrectangle is in the direction from m towards R_{ij} and has magnitude $\dfrac{Gm\rho \, \Delta A_i}{s_{ij}^2}$. The symmetry of the lamina with respect to the x- and y-axes and the position of m are such that all horizontal components of the gravitational force cancel, so that the total force is simply in the z-direction. Thus, we need only be concerned with the components of this vertical force; that is, $\dfrac{Gm\rho \, \Delta A_i}{s_{ij}^2} \sin\alpha$, where α is the angle between the origin, r_i^* and the mass m. Thus $\sin\alpha = \dfrac{d}{s_{ij}}$ and the previous result becomes $\dfrac{Gm\rho d \, \Delta A_i}{s_{ij}^3}$. The total attractive force is just the Riemann sum $\displaystyle\sum_{i=1}^{m}\sum_{j=1}^{n} \frac{Gm\rho d \, \Delta A_i}{s_{ij}^3} = \sum_{i=1}^{m}\sum_{j=1}^{n} \frac{Gm\rho d (r_i^*) \, \Delta r \, \Delta\theta}{\left[(r_i^*)^2 + d^2\right]^{3/2}}$ which becomes $\displaystyle\int_0^R \int_0^{2\pi} \frac{Gm\rho d}{(r^2 + d^2)^{3/2}} r \, d\theta \, dr$ as $m \to \infty$ and $n \to \infty$. Therefore,

$$F = 2\pi Gm\rho d \int_0^R \frac{r}{(r^2 + d^2)^{3/2}} \, dr = 2\pi Gm\rho d \left[-\frac{1}{\sqrt{r^2 + d^2}} \right]_0^R = 2\pi Gm\rho d \left(\frac{1}{d} - \frac{1}{\sqrt{R^2 + d^2}} \right)$$

(b) This is just the result of part (a) in the limit as $R \to \infty$. In this case $\dfrac{1}{\sqrt{R^2 + d^2}} \to 0$, and we are left with

$$F = 2\pi Gm\rho d \left(\frac{1}{d} - 0 \right) = 2\pi Gm\rho.$$

11. $\int_0^x \int_0^y \int_0^z f(t) \, dt \, dz \, dy = \iiint_E f(t) \, dV$, where $E = \{(t, z, y) \mid 0 \le t \le z, 0 \le z \le y, 0 \le y \le x\}$.

If we let D be the projection of E on the yt-plane then $D = \{(y, t) \mid 0 \le t \le x, t \le y \le x\}$. And we see from the diagram that $E = \{(t, z, y) \mid t \le z \le y, t \le y \le x, 0 \le t \le x\}$. So

$$\begin{aligned}
\int_0^x \int_0^y \int_0^z f(t) \, dt \, dz \, dy &= \int_0^x \int_t^x \int_t^y f(t) \, dz \, dy \, dt = \int_0^x \left[\int_t^x (y - t) \, f(t) \, dy \right] dt \\
&= \int_0^x \left[\left(\tfrac{1}{2}y^2 - ty\right) f(t) \right]_{y=t}^{y=x} dt = \int_0^x \left[\tfrac{1}{2}x^2 - tx - \tfrac{1}{2}t^2 + t^2 \right] f(t) \, dt \\
&= \int_0^x \left[\tfrac{1}{2}x^2 - tx + \tfrac{1}{2}t^2 \right] f(t) \, dt = \int_0^x \left(\tfrac{1}{2}x^2 - 2tx + t^2 \right) f(t) \, dt \\
&= \tfrac{1}{2} \int_0^x (x - t)^2 \, f(t) \, dt
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. $n^{-2}\sum_{i=1}^{n}\sum_{j=1}^{n^2}\dfrac{1}{\sqrt{n^2+ni+j}}=\sum_{i=1}^{n}\sum_{j=1}^{n^2}\dfrac{1}{\frac{1}{n}\sqrt{n^2+ni+j}}\cdot\dfrac{1}{n^3}=\sum_{i=1}^{n}\sum_{j=1}^{n^2}\dfrac{1}{\sqrt{1+\frac{i}{n}+\frac{j}{n^2}}}\cdot\dfrac{1}{n^3}$ can be considered a double Riemann sum of the function $f(x,y)=\dfrac{1}{\sqrt{1+x+y}}$ where the square region $R=\{(x,y)\mid 0\le x\le 1, 0\le y\le 1\}$ is divided into subrectangles by dividing the interval $[0,1]$ on the x-axis into n subintervals, each of width $\frac{1}{n}$, and $[0,1]$ on the y-axis is divided into n^2 subintervals, each of width $\frac{1}{n^2}$. Then the area of each subrectangle is $\Delta A=\frac{1}{n^3}$, and if we take the upper right corners of the subrectangles as sample points, we have $(x_{ij}^*, y_{ij}^*)=\left(\frac{i}{n},\frac{j}{n^2}\right)$. Finally, note that $n^2\to\infty$ as $n\to\infty$, so

$$\lim_{n\to\infty} n^{-2}\sum_{i=1}^{n}\sum_{j=1}^{n^2}\frac{1}{\sqrt{n^2+ni+j}}=\lim_{n,n^2\to\infty}\sum_{i=1}^{n}\sum_{j=1}^{n^2}\frac{1}{\sqrt{1+\frac{i}{n}+\frac{j}{n^2}}}\cdot\frac{1}{n^3}=\lim_{n,n^2\to\infty}\sum_{i=1}^{n}\sum_{j=1}^{n^2}f(x_{ij}^*,y_{ij}^*)\,\Delta A$$

But by Definition 15.1.5 this is equal to $\iint_R f(x,y)\,dA$, so

$$\begin{aligned}\lim_{n\to\infty} n^{-2}\sum_{i=1}^{n}\sum_{j=1}^{n^2}\frac{1}{\sqrt{n^2+ni+j}}&=\iint_R f(x,y)\,dA=\int_0^1\int_0^1\frac{1}{\sqrt{1+x+y}}\,dy\,dx\\&=\textstyle\int_0^1\left[2(1+x+y)^{1/2}\right]_{y=0}^{y=1}dx=2\int_0^1\left(\sqrt{2+x}-\sqrt{1+x}\right)dx\\&=2\left[\tfrac{2}{3}(2+x)^{3/2}-\tfrac{2}{3}(1+x)^{3/2}\right]_0^1=\tfrac{4}{3}(3^{3/2}-2^{3/2}-2^{3/2}+1)\\&=\tfrac{4}{3}(3\sqrt{3}-4\sqrt{2}+1)=4\sqrt{3}-\tfrac{16}{3}\sqrt{2}+\tfrac{4}{3}\end{aligned}$$

13. The volume is $V=\iiint_R dV$ where R is the solid region given. From Exercise 15.9.23(a), the transformation $x=au$, $y=bv$, $z=cw$ maps the unit ball $u^2+v^2+w^2\le 1$ to the solid ellipsoid $\dfrac{x^2}{a^2}+\dfrac{y^2}{b^2}+\dfrac{z^2}{c^2}\le 1$ with $\dfrac{\partial(x,y,z)}{\partial(u,v,w)}=abc$. The same transformation maps the plane $u+v+w=1$ to $\dfrac{x}{a}+\dfrac{y}{b}+\dfrac{z}{c}=1$. Thus the region R in xyz-space corresponds to the region S in uvw-space consisting of the smaller piece of the unit ball cut off by the plane $u+v+w=1$, a "cap of a sphere" (see the figure).

We will need to compute the volume of S, but first consider the general case where a horizontal plane slices the upper portion of a sphere of radius r to produce a cap of height h. We use spherical coordinates. From the figure, a line through the origin at angle ϕ from the z-axis intersects the plane when $\cos\phi=(r-h)/a \;\Rightarrow\; a=(r-h)/\cos\phi$, and the line passes through the outer rim of the cap when $a=r \;\Rightarrow\; \cos\phi=(r-h)/r \;\Rightarrow\; \phi=\cos^{-1}((r-h)/r)$. Thus the cap

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

is described by $\{(\rho,\theta,\phi) \mid (r-h)/\cos\phi \le \rho \le r, 0 \le \theta \le 2\pi, 0 \le \phi \le \cos^{-1}((r-h)/r)\}$ and its volume is

$$\begin{aligned}
V &= \int_0^{2\pi}\int_0^{\cos^{-1}((r-h)/r)}\int_{(r-h)/\cos\phi}^{r} \rho^2\sin\phi\,d\rho\,d\phi\,d\theta \\
&= \int_0^{2\pi}\int_0^{\cos^{-1}((r-h)/r)} \left[\tfrac{1}{3}\rho^3\sin\phi\right]_{\rho=(r-h)/\cos\phi}^{\rho=r} d\phi\,d\theta \\
&= \frac{1}{3}\int_0^{2\pi}\int_0^{\cos^{-1}((r-h)/r)} \left[r^3\sin\phi - \frac{(r-h)^3}{\cos^3\phi}\sin\phi\right] d\phi\,d\theta \\
&= \tfrac{1}{3}\int_0^{2\pi}\left[-r^3\cos\phi - \tfrac{1}{2}(r-h)^3\cos^{-2}\phi\right]_{\phi=0}^{\phi=\cos^{-1}((r-h)/r)} d\theta \\
&= \frac{1}{3}\int_0^{2\pi}\left[-r^3\left(\frac{r-h}{r}\right) - \frac{1}{2}(r-h)^3\left(\frac{r-h}{r}\right)^{-2} + r^3 + \frac{1}{2}(r-h)^3\right] d\theta \\
&= \tfrac{1}{3}\int_0^{2\pi}\left(\tfrac{3}{2}rh^2 - \tfrac{1}{2}h^3\right) d\theta = \tfrac{1}{3}\left(\tfrac{3}{2}rh^2 - \tfrac{1}{2}h^3\right)(2\pi) = \pi h^2\left(r - \tfrac{1}{3}h\right)
\end{aligned}$$

(This volume can also be computed by treating the cap as a solid of revolution and using the single variable disk method; see Exercise 6.2.61.)

To determine the height h of the cap cut from the unit ball by the plane $u+v+w=1$, note that the line $u=v=w$ passes through the origin with direction vector $\langle 1,1,1\rangle$ which is perpendicular to the plane. Therefore this line coincides with a radius of the sphere that passes through the center of the cap and h is measured along this line. The line intersects the plane at $\left(\frac{1}{3},\frac{1}{3},\frac{1}{3}\right)$ and the sphere at $\left(\frac{1}{\sqrt{3}},\frac{1}{\sqrt{3}},\frac{1}{\sqrt{3}}\right)$. (See the figure.)

The distance between these points is $h = \sqrt{3\left(\frac{1}{\sqrt{3}}-\frac{1}{3}\right)^2} = \sqrt{3}\left(\frac{1}{\sqrt{3}}-\frac{1}{3}\right) = 1-\frac{1}{\sqrt{3}}$. Thus the volume of R is

$$\begin{aligned}
V &= \iiint_R dV = \iiint_S \left|\frac{\partial(x,y,z)}{\partial(u,v,w)}\right| dV = abc\iiint_S dV = abc\,V(S) \\
&= abc\cdot\pi h^2\left(r-\tfrac{1}{3}h\right) = abc\cdot\pi\left(1-\tfrac{1}{\sqrt{3}}\right)^2\left[1-\tfrac{1}{3}\left(1-\tfrac{1}{\sqrt{3}}\right)\right] \\
&= abc\pi\left(\tfrac{4}{3}-\tfrac{2}{\sqrt{3}}\right)\left(\tfrac{2}{3}+\tfrac{1}{3\sqrt{3}}\right) = abc\pi\left(\tfrac{2}{3}-\tfrac{8}{9\sqrt{3}}\right) \approx 0.482abc
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16 ☐ VECTOR CALCULUS

16.1 Vector Fields

1. $\mathbf{F}(x, y) = \mathbf{i} + \frac{1}{2}\mathbf{j}$

All vectors in this field are identical with length $\sqrt{1^2 + \left(\frac{1}{2}\right)^2} = \sqrt{\frac{5}{4}} = \frac{\sqrt{5}}{2}$ and parallel to $\left\langle 1, \frac{1}{2} \right\rangle$, or, equivalently, $\langle 2, 1 \rangle$.

2. $\mathbf{F}(x, y) = 2\,\mathbf{i} - \mathbf{j}$

All vectors in this field are identical with length $\sqrt{2^2 + (-1)^2} = \sqrt{5}$ and parallel to $\langle 2, -1 \rangle$.

3. $\mathbf{F}(x, y) = \mathbf{i} + \frac{1}{2}y\,\mathbf{j}$

The length of the vector $\mathbf{i} + \frac{1}{2}y\,\mathbf{j}$ is $\sqrt{1 + \frac{1}{4}y^2}$. Vectors along the line $y = 0$ are horizontal with length 1.

4. $\mathbf{F}(x, y) = x\,\mathbf{i} + \frac{1}{2}y\,\mathbf{j}$

The length of the vector $x\,\mathbf{i} + \frac{1}{2}y\,\mathbf{j}$ is $\sqrt{x^2 + \frac{1}{4}y^2}$. Vectors point roughly away from the origin and vectors farther from the origin are longer.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $\mathbf{F}(x, y) = -\frac{1}{2}\,\mathbf{i} + (y - x)\,\mathbf{j}$

The length of the vector $-\frac{1}{2}\,\mathbf{i} + (y - x)\,\mathbf{j}$ is $\sqrt{\frac{1}{4} + (y - x)^2}$. Vectors along the line $y = x$ are horizontal with length $\frac{1}{2}$.

6. $\mathbf{F}(x, y) = y\,\mathbf{i} + (x + y)\,\mathbf{j}$

The length of the vector $y\,\mathbf{i} + (x + y)\,\mathbf{j}$ is $\sqrt{y^2 + (x + y)^2}$. Vectors along the x-axis are vertical, and vectors along the line $y = -x$ are horizontal with length $|y|$.

7. $\mathbf{F}(x, y) = \dfrac{y\,\mathbf{i} + x\,\mathbf{j}}{\sqrt{x^2 + y^2}}$

The length of the vector $\dfrac{y\,\mathbf{i} + x\,\mathbf{j}}{\sqrt{x^2 + y^2}}$ is

$$\sqrt{\frac{y^2}{x^2 + y^2} + \frac{x^2}{x^2 + y^2}} = 1.$$

Vectors along the x-axis are vertical, and vectors along the y-axis are horizontal. In general, vectors in Q1 and QIII point away from the origin, whereas vectors in QII and QIV point toward the origin.

8. $\mathbf{F}(x, y) = \dfrac{y\,\mathbf{i} - x\,\mathbf{j}}{\sqrt{x^2 + y^2}}$

All the vectors $\mathbf{F}(x, y)$ are unit vectors tangent to circles centered at the origin with radius $\sqrt{x^2 + y^2}$.

9. $\mathbf{F}(x, y, z) = \mathbf{i}$

All vectors in this field are identical, with length 1 and pointing in the direction of the positive x-axis.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $\mathbf{F}(x, y, z) = z\,\mathbf{i}$

At each point (x, y, z), $\mathbf{F}(x, y, z)$ is a vector of length $|z|$. For $z > 0$, all point in the direction of the positive x-axis, while for $z < 0$, all are in the direction of the negative x-axis. In each plane $z = k$, all the vectors are identical.

11. $\mathbf{F}(x, y, z) = -y\,\mathbf{i}$

At each point (x, y, z), $\mathbf{F}(x, y, z)$ is a vector of length $|y|$. For $y > 0$, all point in the direction of the negative x-axis, while for $y < 0$, all are in the direction of the positive x-axis. In each plane $y = k$, all the vectors are identical.

12. $\mathbf{F}(x, y, z) = \mathbf{i} + \mathbf{k}$

All vectors in this field have length $\sqrt{2}$ and point in the same direction, parallel to the xz-plane.

13. $\mathbf{F}(x, y) = \langle x, -y \rangle$ corresponds to graph IV. In the first quadrant all the vectors have positive x-components and negative y-components, in the second quadrant all vectors have negative x- and y-components, in the third quadrant all vectors have negative x-components and positive y-components, and in the fourth quadrant all vectors have positive x- and y-components. In addition, the vectors get shorter as we approach the origin.

14. $\mathbf{F}(x, y) = \langle y, x - y \rangle$ corresponds to graph V. All vectors in quadrants I and II have positive x-components while all vectors in quadrants III and IV have negative x-components. In addition, vectors along the line $y = x$ are horizontal, and vectors get shorter as we approach the origin.

15. $\mathbf{F}(x, y) = \langle y, y + 2 \rangle$ corresponds to graph I. As in Exercise 14, all vectors in quadrants I and II have positive x-components while all vectors in quadrants III and IV have negative x-components. Vectors along the line $y = -2$ are horizontal, and the vectors are independent of x (vectors along horizontal lines are identical).

16. $\mathbf{F}(x, y) = \langle y, 2x \rangle$ corresponds to graph VI. In the first quadrant all the vectors have positive x- and y-components. In the second quadrant all vectors have positive x-components and negative y-components. In the third quadrant all vectors have negative x- and y-components. In the fourth quadrant all vectors have negative x-components and positive y-components.

17. $\mathbf{F}(x, y) = \langle \sin y, \cos x \rangle$ corresponds to graph III. Both the x- and y-components oscillate in all four quadrants.

18. $\mathbf{F}(x, y) = \langle \cos(x + y), x \rangle$ corresponds to graph II. All vectors in quadrants I and IV have positive y-components while all vectors in quadrants II and III have negative y-components. Also, the y-components of vectors along any vertical line remain constant while the x-component oscillates.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $\mathbf{F}(x, y, z) = \mathbf{i} + 2\,\mathbf{j} + 3\,\mathbf{k}$ corresponds to graph IV, since all vectors have identical length and direction.

20. $\mathbf{F}(x, y, z) = \mathbf{i} + 2\,\mathbf{j} + z\,\mathbf{k}$ corresponds to graph I, since the horizontal vector components remain constant, but the vectors above the xy-plane point generally upward while the vectors below the xy-plane point generally downward.

21. $\mathbf{F}(x, y, z) = x\,\mathbf{i} + y\,\mathbf{j} + 3\,\mathbf{k}$ corresponds to graph III; the projection of each vector onto the xy-plane is $x\,\mathbf{i} + y\,\mathbf{j}$, which points away from the origin, and the vectors point generally upward because their z-components are all 3.

22. $\mathbf{F}(x, y, z) = x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$ corresponds to graph II; each vector $\mathbf{F}(x, y, z)$ has the same length and direction as the position vector of the point (x, y, z), and therefore the vectors all point directly away from the origin.

23.

$\mathbf{F}(x, y) = (y^2 - 2xy)\,\mathbf{i} + (3xy - 6x^2)\,\mathbf{j}$.

The vector field seems to have very short vectors near the line $y = 2x$.

For $\mathbf{F}(x, y) = \langle 0, 0\rangle$, we must have $y^2 - 2xy = 0$ and $3xy - 6x^2 = 0$. The first equation holds if $y = 0$ or $y = 2x$, and the second holds if $x = 0$ or $y = 2x$. So both equations hold [and thus $\mathbf{F}(x, y) = \mathbf{0}$] along the line $y = 2x$.

24.

$\mathbf{F}(\mathbf{x}) = (r^2 - 2r)\,\mathbf{x}$, where $\mathbf{x} = \langle x, y\rangle$ and $r = |\mathbf{x}|$.

From the graph, it appears that all of the vectors in the field lie on lines through the origin, and that the vectors have very small magnitudes near the circle $|\mathbf{x}| = 2$ and near the origin. Note that $\mathbf{F}(\mathbf{x}) = \mathbf{0} \quad \Leftrightarrow$ $r(r - 2) = 0 \quad \Leftrightarrow \quad r = 0$ or 2, so as we suspected, $\mathbf{F}(\mathbf{x}) = \mathbf{0}$ for $|\mathbf{x}| = 2$ and for $|\mathbf{x}| = 0$. Note that where $r^2 - r < 0$, the vectors point towards the origin, and where $r^2 - r > 0$, they point away from the origin.

25. $f(x, y) = y\sin(xy) \quad \Rightarrow$

$$\begin{aligned}\nabla f(x, y) &= f_x(x, y)\,\mathbf{i} + f_y(x, y)\,\mathbf{j} = (y\cos(xy)\cdot y)\,\mathbf{i} + [y\cdot x\cos(xy) + \sin(xy)\cdot 1]\,\mathbf{j}\\ &= y^2\cos(xy)\,\mathbf{i} + [xy\cos(xy) + \sin(xy)]\,\mathbf{j}\end{aligned}$$

26. $f(s, t) = \sqrt{2s + 3t} \quad \Rightarrow$

$$\nabla f(s, t) = f_s(s, t)\,\mathbf{i} + f_t(s, t)\,\mathbf{j} = \left[\tfrac{1}{2}(2s + 3t)^{-1/2}\cdot 2\right]\mathbf{i} + \left[\tfrac{1}{2}(2s + 3t)^{-1/2}\cdot 3\right]\mathbf{j} = \frac{1}{\sqrt{2s + 3t}}\,\mathbf{i} + \frac{3}{2\sqrt{2s + 3t}}\,\mathbf{j}$$

27. $f(x, y, z) = \sqrt{x^2 + y^2 + z^2} \quad \Rightarrow$

$$\begin{aligned}\nabla f(x, y, z) &= f_x(x, y, z)\,\mathbf{i} + f_y(x, y, z)\,\mathbf{j} + f_z(x, y, z)\,\mathbf{k}\\ &= \tfrac{1}{2}(x^2 + y^2 + z^2)^{-1/2}(2x)\,\mathbf{i} + \tfrac{1}{2}(x^2 + y^2 + z^2)^{-1/2}(2y)\,\mathbf{j} + \tfrac{1}{2}(x^2 + y^2 + z^2)^{-1/2}(2z)\,\mathbf{k}\\ &= \frac{x}{\sqrt{x^2 + y^2 + z^2}}\,\mathbf{i} + \frac{y}{\sqrt{x^2 + y^2 + z^2}}\,\mathbf{j} + \frac{z}{\sqrt{x^2 + y^2 + z^2}}\,\mathbf{k}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $f(x,y,z) = x^2 y e^{y/z} \quad \Rightarrow$

$$\begin{aligned}\nabla f(x,y,z) &= f_x(x,y,z)\,\mathbf{i} + f_y(x,y,z)\,\mathbf{j} + f_z(x,y,z)\,\mathbf{k} \\ &= 2xye^{y/z}\,\mathbf{i} + x^2\left[y \cdot e^{yz}(1/z) + e^{y/z}\cdot 1\right]\mathbf{j} + \left[x^2 y e^{y/z}(-y/z^2)\right]\mathbf{k} \\ &= 2xye^{y/z}\,\mathbf{i} + x^2 e^{y/z}\left(\frac{y}{z}+1\right)\mathbf{j} - \frac{x^2y^2}{z^2}e^{y/z}\,\mathbf{k}\end{aligned}$$

29. $f(x,y) = \frac{1}{2}(x-y)^2 \quad \Rightarrow$

$\nabla f(x,y) = (x-y)(1)\,\mathbf{i} + (x-y)(-1)\,\mathbf{j} = (x-y)\,\mathbf{i} + (y-x)\,\mathbf{j}$.

The length of $\nabla f(x,y)$ is $\sqrt{(x-y)^2 + (y-x)^2} = \sqrt{2}\,|x-y|$. The vectors are $\mathbf{0}$ along the line $y = x$. Elsewhere the vectors point away from the line $y = x$ with length that increases as the distance from the line increases.

30. $f(x,y) = \frac{1}{2}(x^2 - y^2) \quad \Rightarrow \quad \nabla f(x,y) = x\,\mathbf{i} - y\,\mathbf{j}$.

The length of $\nabla f(x,y)$ is $\sqrt{x^2+y^2}$. The lengths of the vectors increase as the distance from the origin increases, and the terminal point of each vector lies on the x-axis.

31. $f(x,y) = x^2 + y^2 \quad \Rightarrow \quad \nabla f(x,y) = 2x\,\mathbf{i} + 2y\,\mathbf{j}$. Thus, each vector $\nabla f(x,y)$ has the same direction and twice the length of the position vector of the point (x,y), so the vectors all point directly away from the origin and their lengths increase as we move away from the origin. Hence, ∇f is graph III.

32. $f(x,y) = x(x+y) = x^2 + xy \quad \Rightarrow \quad \nabla f(x,y) = (2x+y)\,\mathbf{i} + x\,\mathbf{j}$. The y-component of each vector is x, so the vectors point upward in quadrants I and IV and downward in quadrants II and III. Also, the x-component of each vector is 0 along the line $y = -2x$ so the vectors are vertical there. Thus, ∇f is graph IV.

33. $f(x,y) = (x+y)^2 \quad \Rightarrow \quad \nabla f(x,y) = 2(x+y)\,\mathbf{i} + 2(x+y)\,\mathbf{j}$. The x- and y-components of each vector are equal, so all vectors are parallel to the line $y = x$. The vectors are $\mathbf{0}$ along the line $y = -x$ and their length increases as the distance from this line increases. Thus, ∇f is graph II.

34. $f(x,y) = \sin\sqrt{x^2+y^2} \quad \Rightarrow$

$$\begin{aligned}\nabla f(x,y) &= \left[\cos\sqrt{x^2+y^2}\cdot\tfrac{1}{2}(x^2+y^2)^{-1/2}(2x)\right]\mathbf{i} + \left[\cos\sqrt{x^2+y^2}\cdot\tfrac{1}{2}(x^2+y^2)^{-1/2}(2y)\right]\mathbf{j} \\ &= \frac{\cos\sqrt{x^2+y^2}}{\sqrt{x^2+y^2}}\,x\,\mathbf{i} + \frac{\cos\sqrt{x^2+y^2}}{\sqrt{x^2+y^2}}\,y\,\mathbf{j} \text{ or } \frac{\cos\sqrt{x^2+y^2}}{\sqrt{x^2+y^2}}\,(x\,\mathbf{i} + y\,\mathbf{j})\end{aligned}$$

Thus each vector is a scalar multiple of its position vector, so the vectors point toward or away from the origin with length that changes in a periodic fashion as we move away from the origin. ∇f is graph I.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. $f(x, y) = \ln(1 + x^2 + 2y^2)$. We graph

$\nabla f(x, y) = \dfrac{2x}{1 + x^2 + 2y^2}\,\mathbf{i} + \dfrac{4y}{1 + x^2 + 2y^2}\,\mathbf{j}$ along with a contour map of f. The graph shows that the gradient vectors are perpendicular to the level curves. Also, the gradient vectors point in the direction in which f is increasing and are longer where the level curves are closer together.

36. $f(x, y) = \cos x - 2\sin y$.

We graph $\nabla f(x, y) = -\sin x\,\mathbf{i} - 2\cos y\,\mathbf{j}$ along with a contour map of f. The graph shows that the gradient vectors are perpendicular to the level curves. Also, the gradient vectors point in the direction in which f is increasing and are longer where the level curves are closer together.

37. $\mathbf{V}(x, y) = \langle x^2, x + y^2 \rangle$. At $t = 3$ the particle is at $(2, 1)$ so its velocity is $\mathbf{V}(2, 1) = \langle 4, 3 \rangle$. After 0.01 units of time, the particle's change in location should be approximately $0.01\,\mathbf{V}(2, 1) = 0.01\,\langle 4, 3 \rangle = \langle 0.04, 0.03 \rangle$, so the particle should be approximately at the point $(2.04, 1.03)$.

38. $\mathbf{F}(x, y) = \langle xy - 2, y^2 - 10 \rangle$. At $t = 1$ the particle is at $(1, 3)$ so its velocity is $\mathbf{F}(1, 3) = \langle 1, -1 \rangle$. After 0.05 units of time, the particle's change in location should be approximately $0.05\,\mathbf{F}(1, 3) = 0.05\,\langle 1, -1 \rangle = \langle 0.05, -0.05 \rangle$, so the particle should be approximately at the point $(1.05, 2.95)$.

39. (a) We sketch the vector field $\mathbf{F}(x, y) = x\,\mathbf{i} - y\,\mathbf{j}$ along with several approximate flow lines. The flow lines appear to be hyperbolas with shape similar to the graph of $y = \pm 1/x$, so we might guess that the flow lines have equations $y = C/x$.

(b) If $x = x(t)$ and $y = y(t)$ are parametric equations of a flow line, then the velocity vector of the flow line at the point (x, y) is $x'(t)\,\mathbf{i} + y'(t)\,\mathbf{j}$. Since the velocity vectors coincide with the vectors in the vector field, we have $x'(t)\,\mathbf{i} + y'(t)\,\mathbf{j} = x\,\mathbf{i} - y\,\mathbf{j} \;\Rightarrow\; dx/dt = x,\ dy/dt = -y$. To solve these differential equations, we know $dx/dt = x \;\Rightarrow\; dx/x = dt \;\Rightarrow\; \ln|x| = t + C \;\Rightarrow\; x = \pm e^{t+C} = Ae^t$ for some constant A, and $dy/dt = -y \;\Rightarrow\; dy/y = -dt \;\Rightarrow\; \ln|y| = -t + K \;\Rightarrow\; y = \pm e^{-t+K} = Be^{-t}$ for some constant B. Therefore $xy = Ae^t Be^{-t} = AB =$ constant. If the flow line passes through $(1, 1)$ then $(1)(1) =$ constant $= 1 \;\Rightarrow\; xy = 1 \;\Rightarrow\; y = 1/x,\ x > 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. (a) We sketch the vector field $\mathbf{F}(x, y) = \mathbf{i} + x\,\mathbf{j}$ along with several approximate flow lines. The flow lines appear to be parabolas.

(b) If $x = x(t)$ and $y = y(t)$ are parametric equations of a flow line, then the velocity vector of the flow line at the point (x, y) is $x'(t)\,\mathbf{i} + y'(t)\,\mathbf{j}$. Since the velocity vectors coincide with the vectors in the vector field, we have $x'(t)\,\mathbf{i} + y'(t)\,\mathbf{j} = \mathbf{i} + x\,\mathbf{j} \quad \Rightarrow \quad \dfrac{dx}{dt} = 1,\ \dfrac{dy}{dt} = x$. Thus, $\dfrac{dy}{dx} = \dfrac{dy/dt}{dx/dt} = \dfrac{x}{1} = x$.

(c) From part (b), $dy/dx = x$. Integrating, we have $y = \frac{1}{2}x^2 + c$. Since the particle starts at the origin, we know $(0, 0)$ is on the curve, so $0 = 0 + c \quad \Rightarrow \quad c = 0$ and the path the particle follows is $y = \frac{1}{2}x^2$.

16.2 Line Integrals

1. $x = t^2$ and $y = 2t$, $0 \le t \le 3$, so by Formula 3

$$\int_C y\,ds = \int_0^3 2t\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\,dt = \int_0^3 2t\,\sqrt{(2t)^2 + (2)^2}\,dt = \int_0^3 2t\,\sqrt{4t^2 + 4}\,dt$$

$$= \int_0^3 4t\,\sqrt{t^2 + 1}\,dt = \left[2 \cdot \tfrac{2}{3}(t^2 + 1)^{3/2}\right]_0^3 = \tfrac{4}{3}(10^{3/2} - 1)$$

2. $x = t^3$ and $y = t^4$, $1 \le t \le 2$, so by Formula 3

$$\int_C (x/y)\,ds = \int_1^2 (t^3/t^4)\sqrt{(3t^2)^2 + (4t^3)^2}\,dt = \int_1^2 (1/t) \cdot t^2\sqrt{9 + 16t^2}\,dt = \int_1^2 t\sqrt{9 + 16t^2}\,dt$$

$$= \left[\tfrac{1}{32} \cdot \tfrac{2}{3}(9 + 16t^2)^{3/2}\right]_1^2 = \tfrac{1}{48}(73^{3/2} - 25^{3/2}) \text{ or } \tfrac{1}{48}(73\sqrt{73} - 125)$$

3. Parametric equations for C are $x = 4\cos t,\ \ y = 4\sin t,\ -\frac{\pi}{2} \le t \le \frac{\pi}{2}$. Then

$$\int_C xy^4\,ds = \int_{-\pi/2}^{\pi/2} (4\cos t)(4\sin t)^4\sqrt{(-4\sin t)^2 + (4\cos t)^2}\,dt = \int_{-\pi/2}^{\pi/2} 4^5\cos t\,\sin^4 t\,\sqrt{16(\sin^2 t + \cos^2 t)}\,dt$$

$$= 4^5\int_{-\pi/2}^{\pi/2} (\sin^4 t\cos t)(4)\,dt = (4)^6\left[\tfrac{1}{5}\sin^5 t\right]_{-\pi/2}^{\pi/2} = 4^6 \cdot \tfrac{2}{5} = 1638.4$$

4. Parametric equations for C are $x = 2 + 3t,\ y = 4t,\ 0 \le t \le 1$. Then

$$\int_C xe^y\,ds = \int_0^1 (2 + 3t)\,e^{4t}\,\sqrt{3^2 + 4^2}\,dt = 5\int_0^1 (2 + 3t)\,e^{4t}\,dt$$

Integrating by parts with $u = 2 + 3t \quad \Rightarrow \quad du = 3\,dt,\ dv = e^{4t}\,dt \quad \Rightarrow \quad v = \frac{1}{4}e^{4t}$ gives

$$\int_C xe^y\,ds = 5\left[\tfrac{1}{4}(2 + 3t)e^{4t} - \tfrac{3}{16}e^{4t}\right]_0^1 = 5\left[\tfrac{5}{4}e^4 - \tfrac{3}{16}e^4 - \tfrac{1}{2} + \tfrac{3}{16}\right] = \tfrac{85}{16}e^4 - \tfrac{25}{16}$$

5. If we choose x as the parameter, parametric equations for C are $x = x,\ \ y = x^2$ for $0 \le x \le \pi$ and by Equations 7

$$\int_C \left(x^2y + \sin x\right)dy = \int_0^\pi \left[x^2(x^2) + \sin x\right] \cdot 2x\,dx = 2\int_0^\pi \left(x^5 + x\sin x\right)dx$$

$$= 2\left[\tfrac{1}{6}x^6 - x\cos x + \sin x\right]_0^\pi \quad \left[\begin{array}{c}\text{where we integrated by parts}\\ \text{in the second term}\end{array}\right]$$

$$= 2\left[\tfrac{1}{6}\pi^6 + \pi + 0 - 0\right] = \tfrac{1}{3}\pi^6 + 2\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. Choosing y as the parameter, we have $x = y^3, \quad y = y, \quad -1 \le y \le 1$. Then

$$\int_C e^x\,dx = \int_{-1}^{1} e^{y^3} \cdot 3y^2\,dy = e^{y^3}\Big]_{-1}^{1} = e^1 - e^{-1} = e - \tfrac{1}{e}.$$

7.

$C = C_1 + C_2$

On C_1: $x = x, y = \frac{1}{2}x \quad \Rightarrow \quad dy = \frac{1}{2}\,dx, \quad 0 \le x \le 2.$

On C_2: $x = x, y = 3 - x \quad \Rightarrow \quad dy = -dx, \quad 2 \le x \le 3.$

Then

$$\begin{aligned}
\int_C (x+2y)\,dx + x^2\,dy &= \int_{C_1} (x+2y)\,dx + x^2\,dy + \int_{C_2} (x+2y)\,dx + x^2\,dy \\
&= \int_0^2 \left[x + 2\left(\tfrac{1}{2}x\right) + x^2\left(\tfrac{1}{2}\right)\right] dx + \int_2^3 \left[x + 2(3-x) + x^2(-1)\right] dx \\
&= \int_0^2 \left(2x + \tfrac{1}{2}x^2\right) dx + \int_2^3 \left(6 - x - x^2\right) dx \\
&= \left[x^2 + \tfrac{1}{6}x^3\right]_0^2 + \left[6x - \tfrac{1}{2}x^2 - \tfrac{1}{3}x^3\right]_2^3 = \tfrac{16}{3} - 0 + \tfrac{9}{2} - \tfrac{22}{3} = \tfrac{5}{2}
\end{aligned}$$

8.

$C = C_1 + C_2$

On C_1: $x = 2\cos t \quad \Rightarrow \quad dx = -2\sin t\,dt,$

$y = 2\sin t \quad \Rightarrow \quad dy = 2\cos t\,dt, 0 \le t \le \frac{\pi}{2}.$

On C_2: $x = -t \quad \Rightarrow \quad dx = -dt,$

$y = 2 - t \quad \Rightarrow \quad dy = -dt, 0 \le t \le 1.$

Then

$$\begin{aligned}
\int_C x^2\,dx + y^2\,dy &= \int_{C_1} x^2\,dx + y^2\,dy + \int_{C_2} x^2\,dx + y^2\,dy \\
&= \int_0^{\pi/2} (2\cos t)^2(-2\sin t\,dt) + (2\sin t)^2(2\cos t\,dt) + \int_0^1 (-t)^2(-dt) + (2-t)^2(-dt) \\
&= \int_0^{\pi/2} (-8\cos^2 t\sin t + 8\sin^2 t\cos t)\,dt - 2\int_0^1 (t^2 - 2t + 2)\,dt \\
&= 8\left[\tfrac{1}{3}\cos^3 t + \tfrac{1}{3}\sin^3 t\right]_0^{\pi/2} - 2\left[\tfrac{1}{3}t^3 - t^2 + 2t\right]_0^1 = 8\left(\tfrac{1}{3} - \tfrac{1}{3}\right) - 2\left(\tfrac{1}{3} - 1 + 2\right) = -\tfrac{8}{3}
\end{aligned}$$

9. $x = \cos t, \quad y = \sin t, \quad z = t, \quad 0 \le t \le \pi/2$. Then by Formula 9,

$$\begin{aligned}
\int_C x^2 y\,ds &= \int_0^{\pi/2} (\cos t)^2(\sin t)\sqrt{\left(\tfrac{dx}{dt}\right)^2 + \left(\tfrac{dy}{dt}\right)^2 + \left(\tfrac{dz}{dt}\right)^2}\,dt \\
&= \int_0^{\pi/2} \cos^2 t\sin t\,\sqrt{(-\sin t)^2 + (\cos t)^2 + (1)^2}\,dt = \int_0^{\pi/2} \cos^2 t\sin t\,\sqrt{\sin^2 t + \cos^2 t + 1}\,dt \\
&= \sqrt{2}\int_0^{\pi/2} \cos^2 t\sin t\,dt = \sqrt{2}\left[-\tfrac{1}{3}\cos^3 t\right]_0^{\pi/2} = \sqrt{2}\left(0 + \tfrac{1}{3}\right) = \tfrac{\sqrt{2}}{3}
\end{aligned}$$

10. Parametric equations for the line segment C from $(3, 1, 2)$ to $(1, 2, 5)$ are $x = 3 - 2t, \; y = 1 + t, \; z = 2 + 3t, \; 0 \le t \le 1$.

Then by Formula 9,

$$\begin{aligned}
\int_C y^2 z\,ds &= \int_0^1 (1+t)^2(2+3t)\sqrt{(-2)^2 + 1^2 + 3^2}\,dt = \sqrt{14}\int_0^1 (3t^3 + 8t^2 + 7t + 2)\,dt \\
&= \sqrt{14}\left[\tfrac{3}{4}t^4 + \tfrac{8}{3}t^3 + \tfrac{7}{2}t^2 + 2t\right]_0^1 = \sqrt{14}\left(\tfrac{3}{4} + \tfrac{8}{3} + \tfrac{7}{2} + 2\right) = \tfrac{107}{12}\sqrt{14}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. Parametric equations for the line segment C from $(0,0,0)$ to $(1,2,3)$ are $x=t,\ y=2t,\ z=3t,\ 0\le t\le 1$. Then by Formula 9, $\int_C xe^{yz}\,ds=\int_0^1 te^{(2t)(3t)}\sqrt{1^2+2^2+3^2}\,dt=\sqrt{14}\int_0^1 te^{6t^2}\,dt=\sqrt{14}\left[\frac{1}{12}e^{6t^2}\right]_0^1=\frac{\sqrt{14}}{12}(e^6-1)$.

12. C: $x=t, y=\cos 2t, z=\sin 2t, 0\le t\le 2\pi$.

$\sqrt{(dx/dt)^2+(dy/dt)^2+(dz/dt)^2}=\sqrt{1^2+(-2\sin 2t)^2+(2\cos 2t)^2}=\sqrt{1+4(\sin^2 2t+\cos^2 2t)}=\sqrt{5}$.

Then by Formula 9,

$$\int_C (x^2+y^2+z^2)\,ds=\int_0^{2\pi}(t^2+\cos^2 2t+\sin^2 2t)\sqrt{5}\,dt=\sqrt{5}\int_0^{2\pi}(t^2+1)\,dt$$
$$=\sqrt{5}\left[\tfrac{1}{3}t^3+t\right]_0^{2\pi}=\sqrt{5}\left[\tfrac{1}{3}(8\pi^3)+2\pi\right]=\sqrt{5}\left(\tfrac{8}{3}\pi^3+2\pi\right)$$

13. C: $x=t, y=t^2, z=t^3, 0\le t\le 1$.

$$\int_C xye^{yz}\,dy=\int_0^1 (t)(t^2)e^{(t^2)(t^3)}\cdot 2t\,dt=\int_0^1 2t^4e^{t^5}\,dt=\left[\tfrac{2}{5}e^{t^5}\right]_0^1=\tfrac{2}{5}(e^1-e^0)=\tfrac{2}{5}(e-1)$$

14. C: $x=e^t, y=2t, z=\ln t, 1\le t\le 2$.

$$\int_C ye^z\,dz+x\ln x\,dy-y\,dx=\int_1^2 2te^{\ln t}\,\tfrac{1}{t}\,dt+e^t\ln e^t\cdot 2\,dt-2te^t\,dt=\int_1^2\left(2t+2te^t-2te^t\right)\,dt$$
$$=\int_1^2 2t\,dt=\left[t^2\right]_1^2=4-1=3$$

15. C: $x=\sin t, y=\cos t, z=\tan t, -\pi/4\le t\le \pi/4$.

$$\int_C z\,dx+xy\,dy+y^2\,dz=\int_{-\pi/4}^{\pi/4}(\tan t)(\cos t\,dt)+(\sin t)(\cos t)(-\sin t\,dt)+(\cos^2 t)(\sec^2 t\,dt)$$
$$=\int_{-\pi/4}^{\pi/4}(\sin t-\sin^2 t\cos t+1)\,dt=\left[-\cos t-\tfrac{1}{3}\sin^3 t+t\right]_{-\pi/4}^{\pi/4}$$
$$=\left[-\frac{\sqrt{2}}{2}-\frac{1}{3}\left(\frac{\sqrt{2}}{2}\right)^3+\frac{\pi}{4}\right]-\left[-\frac{\sqrt{2}}{2}-\frac{1}{3}\left(-\frac{\sqrt{2}}{2}\right)^3-\frac{\pi}{4}\right]=\frac{\pi}{2}-\frac{\sqrt{2}}{6}$$

16. C: $x=\sqrt{t}, y=t, z=t^2, 1\le t\le 4$.

$$\int_C y\,dx+z\,dy+x\,dz=\int_1^4 t\cdot\tfrac{1}{2}t^{-1/2}\,dt+t^2\cdot dt+\sqrt{t}\cdot 2t\,dt=\int_1^4\left(\tfrac{1}{2}t^{1/2}+t^2+2t^{3/2}\right)dt$$
$$=\left[\tfrac{1}{3}t^{3/2}+\tfrac{1}{3}t^3+\tfrac{4}{5}t^{5/2}\right]_1^4=\tfrac{8}{3}+\tfrac{64}{3}+\tfrac{128}{5}-\tfrac{1}{3}-\tfrac{1}{3}-\tfrac{4}{5}=\tfrac{722}{15}$$

17. Parametric equations for the line segment C from $(1,0,0)$ to $(4,1,2)$ are $x=1+3t,\ \ y=t,\ \ z=2t,\ 0\le t\le 1$. Then

$$\int_C z^2\,dx+x^2\,dy+y^2\,dz=\int_0^1(2t)^2\cdot 3\,dt+(1+3t)^2\,dt+t^2\cdot 2\,dt=\int_0^1\left(23t^2+6t+1\right)dt$$
$$=\left[\tfrac{23}{3}t^3+3t^2+t\right]_0^1=\tfrac{23}{3}+3+1=\tfrac{35}{3}$$

18.

$C=C_1+C_2$

On C_1 from $(0,0,0)$ to $(1,0,1)$: $x=t \Rightarrow dx=dt$,
$y=0 \Rightarrow dy=0\,dt$,
$z=t \Rightarrow dz=dt, 0\le t\le 1$.

On C_2 from $(1,0,1)$ to $(0,1,2)$: $x=1-t \Rightarrow dx=-dt$,
$y=t \Rightarrow dy=dt$,
$z=1+t \Rightarrow dz=dt, 0\le t\le 1$. [continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Then

$$\begin{aligned}
&\textstyle\int_C (y+z)\,dx + (x+z)\,dy + (x+y)\,dz \\
&\quad = \textstyle\int_{C_1} (y+z)\,dx + (x+z)\,dy + (x+y)\,dz + \int_{C_2} (y+z)\,dx + (x+z)\,dy + (x+y)\,dz \\
&\quad = \textstyle\int_0^1 (0+t)\,dt + (t+t)\cdot 0\,dt + (t+0)\,dt + \int_0^1 (t+1+t)(-dt) + (1-t+1+t)\,dt + (1-t+t)\,dt \\
&\quad = \textstyle\int_0^1 2t\,dt + \int_0^1 (-2t+2)\,dt = \left[t^2\right]_0^1 + \left[-t^2+2t\right]_0^1 = 1+1 = 2
\end{aligned}$$

19. (a) Along the line $x=-3$, the vectors of $\mathbf{F}$ have positive y-components, so since the path goes upward, the integrand $\mathbf{F}\cdot\mathbf{T}$ is always positive. Therefore $\int_{C_1} \mathbf{F}\cdot d\mathbf{r} = \int_{C_1} \mathbf{F}\cdot\mathbf{T}\,ds$ is positive.

(b) All of the (nonzero) field vectors along the circle with radius 3 are pointed in the clockwise direction, that is, opposite the direction to the path. So $\mathbf{F}\cdot\mathbf{T}$ is negative, and therefore $\int_{C_2} \mathbf{F}\cdot d\mathbf{r} = \int_{C_2} \mathbf{F}\cdot\mathbf{T}\,ds$ is negative.

20. Vectors starting on C_1 point in roughly the same direction as C_1, so the tangential component $\mathbf{F}\cdot\mathbf{T}$ is positive. Then $\int_{C_1} \mathbf{F}\cdot d\mathbf{r} = \int_{C_1} \mathbf{F}\cdot\mathbf{T}\,ds$ is positive. On the other hand, no vectors starting on C_2 point in the same direction as C_2, while some vectors point in roughly the opposite direction, so we would expect $\int_{C_2} \mathbf{F}\cdot d\mathbf{r} = \int_{C_2} \mathbf{F}\cdot\mathbf{T}\,ds$ to be negative.

21. $\mathbf{F}(x,y) = xy^2\,\mathbf{i} - x^2\,\mathbf{j}$ and $\mathbf{r}(t) = t^3\,\mathbf{i} + t^2\,\mathbf{j}$, $0 \le t \le 1 \quad\Rightarrow$

$\mathbf{F}(\mathbf{r}(t)) = (t^3)(t^2)^2\,\mathbf{i} - (t^3)^2\,\mathbf{j} = t^7\,\mathbf{i} - t^6\,\mathbf{j}$ and $\mathbf{r}'(t) = 3t^2\,\mathbf{i} + 2t\,\mathbf{j}$. Then

$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1 \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_0^1 (t^7\cdot 3t^2 - t^6\cdot 2t)\,dt = \int_0^1 (3t^9 - 2t^7)\,dt = \left[\frac{3}{10}t^{10} - \frac{1}{4}t^8\right]_0^1 = \frac{3}{10} - \frac{1}{4} = \frac{1}{20}$.

22. $\mathbf{F}(x,y,z) = (x+y^2)\,\mathbf{i} + xz\,\mathbf{j} + (y+z)\,\mathbf{k}$ and $\mathbf{r}(t) = t^2\,\mathbf{i} + t^3\,\mathbf{j} - 2t\,\mathbf{k}$, $0 \le t \le 2 \quad\Rightarrow$

$\mathbf{F}(\mathbf{r}(t)) = \left(t^2 + (t^3)^2\right)\mathbf{i} + (t^2)(-2t)\,\mathbf{j} + (t^3-2t)\,\mathbf{k} = (t^2+t^6)\,\mathbf{i} - 2t^3\,\mathbf{j} + (t^3-2t)\,\mathbf{k}$ and $\mathbf{r}'(t) = 2t\,\mathbf{i} + 3t^2\,\mathbf{j} - 2\,\mathbf{k}$.

Then

$$\begin{aligned}
\textstyle\int_C \mathbf{F}\cdot d\mathbf{r} &= \textstyle\int_0^2 \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_0^2 (2t^3 + 2t^7 - 6t^5 - 2t^3 + 4t)\,dt = \int_0^2 (2t^7 - 6t^5 + 4t)\,dt \\
&= \left[\tfrac{1}{4}t^8 - t^6 + 2t^2\right]_0^2 = 64 - 64 + 8 = 8
\end{aligned}$$

23. $\mathbf{F}(x,y,z) = \sin x\,\mathbf{i} + \cos y\,\mathbf{j} + xz\,\mathbf{k}$ and $\mathbf{r}(t) = t^3\,\mathbf{i} - t^2\,\mathbf{j} + t\,\mathbf{k}$, $0 \le t \le 1 \quad\Rightarrow$

$\mathbf{F}(\mathbf{r}(t)) = \sin t^3\,\mathbf{i} + \cos(-t^2)\,\mathbf{j} + t^3\cdot t\,\mathbf{k}$ and $\mathbf{r}'(t) = 3t^2\,\mathbf{i} - 2t\,\mathbf{j} + 1\,\mathbf{k}$. Then

$$\begin{aligned}
\textstyle\int_C \mathbf{F}\cdot d\mathbf{r} &= \textstyle\int_0^1 \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_0^1 (3t^2\sin t^3 - 2t\cos t^2 + t^4)\,dt \\
&= \left[-\cos t^3 - \sin t^2 + \tfrac{1}{5}t^5\right]_0^1 = \tfrac{6}{5} - \cos 1 - \sin 1
\end{aligned}$$

24. $\mathbf{F}(x,y,z) = xz\,\mathbf{i} + z^3\,\mathbf{j} + y\,\mathbf{k}$ and $\mathbf{r}(t) = e^t\,\mathbf{i} + e^{2t}\,\mathbf{j} + e^{-t}\,\mathbf{k}$, $-1 \le t \le 1 \quad\Rightarrow$

$\mathbf{F}(\mathbf{r}(t)) = e^t e^{-t}\,\mathbf{i} + (e^{-t})^3\,\mathbf{j} + e^{2t}\,\mathbf{k} = \mathbf{i} + e^{-3t}\,\mathbf{j} + e^{2t}\,\mathbf{k}$ and $\mathbf{r}'(t) = e^t\,\mathbf{i} + 2e^{2t}\,\mathbf{j} - e^{-t}\,\mathbf{k}$. Then

$$\begin{aligned}
\textstyle\int_C \mathbf{F}\cdot d\mathbf{r} &= \textstyle\int_{-1}^1 \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_{-1}^1 \left(1\cdot e^t + e^{-3t}\cdot 2e^{2t} - e^{2t}e^{-t}\right)\,dt \\
&= \textstyle\int_{-1}^1 2e^{-t}\,dt = -2\left[e^{-t}\right]_{-1}^1 = -2(e^{-1} - e)
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. $\mathbf{F}(\mathbf{r}(t)) = \sqrt{\sin^2 t + \sin t \cos t}\ \mathbf{i} + \left[(\sin t \cos t)/\sin^2 t\right]\mathbf{j} = \sqrt{\sin^2 t + \sin t \cos t}\ \mathbf{i} + \cot t\,\mathbf{j}$,

$\mathbf{r}'(t) = 2\sin t \cos t\,\mathbf{i} + (\cos^2 t - \sin^2 t)\,\mathbf{j}$. Then

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \int_{\pi/6}^{\pi/3} \mathbf{F}(\mathbf{r}(t))\cdot \mathbf{r}'(t)\,dt \\ &= \int_{\pi/6}^{\pi/3} \left[2\sin t\cos t\sqrt{\sin^2 t + \sin t\cos t} + (\cot t)(\cos^2 t - \sin^2 t)\right] dt \approx 0.5424\end{aligned}$$

26. $\mathbf{F}(\mathbf{r}(t)) = (\cos t \tan t)e^{\sin t}\,\mathbf{i} + (\tan t \sin t)e^{\cos t}\,\mathbf{j} + (\sin t \cos t)e^{\tan t}\,\mathbf{k}$

$= (\sin t)e^{\sin t}\,\mathbf{i} + (\tan t \sin t)e^{\cos t}\,\mathbf{j} + (\sin t \cos t)e^{\tan t}\,\mathbf{k}$,

$\mathbf{r}'(t) = \cos t\,\mathbf{i} - \sin t\,\mathbf{j} + \sec^2 t\,\mathbf{k}$. Then

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \int_0^{\pi/4} \mathbf{F}(\mathbf{r}(t))\cdot \mathbf{r}'(t)\,dt \\ &= \int_0^{\pi/4} \left[(\sin t\cos t)e^{\sin t} - (\tan t \sin^2 t)e^{\cos t} + (\tan t)e^{\tan t}\right] dt \approx 0.8527\end{aligned}$$

27. $x = t^2,\ \ y = t^3,\ \ z = \sqrt{t}$ so by Formula 9,

$$\begin{aligned}\int_C xy \arctan z\,ds &= \int_1^2 (t^2)(t^3)\arctan\sqrt{t}\cdot\sqrt{(2t)^2 + (3t^2)^2 + \left[1/(2\sqrt{t})\right]^2}\,dt \\ &= \int_1^2 t^5\sqrt{4t^2 + 9t^4 + 1/(4t)}\ \arctan\sqrt{t}\,dt \approx 94.8231\end{aligned}$$

28. $x = 1 + 3t,\ \ y = 2 + t^2,\ \ z = t^4$ so by Formula 9,

$$\begin{aligned}\int_C z\ln(x+y)\,ds &= \int_{-1}^1 t^4 \ln(1 + 3t + 2 + t^2)\cdot\sqrt{(3)^2 + (2t)^2 + (4t^3)^2}\,dt \\ &= \int_{-1}^1 t^4\sqrt{9 + 4t^2 + 16t^6}\ \ln(3 + 3t + t^2)\,dt \approx 1.7260\end{aligned}$$

29. We graph $\mathbf{F}(x, y) = (x - y)\,\mathbf{i} + xy\,\mathbf{j}$ and the curve C. We see that most of the vectors starting on C point in roughly the same direction as C, so for these portions of C the tangential component $\mathbf{F}\cdot\mathbf{T}$ is positive. Although some vectors in the third quadrant which start on C point in roughly the opposite direction, and hence give negative tangential components, it seems reasonable that the effect of these portions of C is outweighed by the positive tangential components. Thus, we would expect $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C \mathbf{F}\cdot\mathbf{T}\,ds$ to be positive.

To verify, we evaluate $\int_C \mathbf{F}\cdot d\mathbf{r}$. The curve C can be represented by $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j},\ \ 0 \le t \le \frac{3\pi}{2}$, so $\mathbf{F}(\mathbf{r}(t)) = (2\cos t - 2\sin t)\,\mathbf{i} + 4\cos t\sin t\,\mathbf{j}$ and $\mathbf{r}'(t) = -2\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j}$. Then

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \int_0^{3\pi/2} \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt \\ &= \int_0^{3\pi/2}[-2\sin t(2\cos t - 2\sin t) + 2\cos t(4\cos t\sin t)]\,dt \\ &= 4\int_0^{3\pi/2}(\sin^2 t - \sin t\cos t + 2\sin t\cos^2 t)\,dt \\ &= 3\pi + \tfrac{2}{3} \qquad \text{[using a CAS]}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. We graph $\mathbf{F}(x,y) = \dfrac{x}{\sqrt{x^2+y^2}}\,\mathbf{i} + \dfrac{y}{\sqrt{x^2+y^2}}\,\mathbf{j}$ and the curve C. In the first quadrant, each vector starting on C points in roughly the same direction as C, so the tangential component $\mathbf{F}\cdot\mathbf{T}$ is positive. In the second quadrant, each vector starting on C points in roughly the direction opposite to C, so $\mathbf{F}\cdot\mathbf{T}$ is negative. Here, it appears that the tangential components in the first and second quadrants counteract each other, so it seems reasonable to guess that $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C \mathbf{F}\cdot\mathbf{T}\,ds$ is zero. To verify, we evaluate $\int_C \mathbf{F}\cdot d\mathbf{r}$. The curve C can be represented by

$\mathbf{r}(t) = t\,\mathbf{i} + (1+t^2)\,\mathbf{j},\ -1 \le t \le 1$, so $\mathbf{F}(\mathbf{r}(t)) = \dfrac{t}{\sqrt{t^2+(1+t^2)^2}}\,\mathbf{i} + \dfrac{1+t^2}{\sqrt{t^2+(1+t^2)^2}}\,\mathbf{j}$ and $\mathbf{r}'(t) = \mathbf{i} + 2t\,\mathbf{j}$. Then

$$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_{-1}^{1} \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_{-1}^{1}\left(\frac{t}{\sqrt{t^2+(1+t^2)^2}} + \frac{2t(1+t^2)}{\sqrt{t^2+(1+t^2)^2}}\right)dt$$

$$= \int_{-1}^{1} \frac{t(3+2t^2)}{\sqrt{t^4+3t^2+1}}\,dt = 0 \qquad \text{[since the integrand is an odd function]}$$

31. (a) $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1 \left\langle e^{t^2-1}, (t^2)(t^3)\right\rangle \cdot \langle 2t, 3t^2\rangle\,dt = \int_0^1 \left(2te^{t^2-1} + 3t^7\right)dt = \left[e^{t^2-1} + \frac{3}{8}t^8\right]_0^1 = \frac{11}{8} - 1/e$

(b) $\mathbf{r}(0) = \mathbf{0},\ \ \mathbf{F}(\mathbf{r}(0)) = \langle e^{-1}, 0\rangle$;

$\mathbf{r}\left(\frac{1}{\sqrt{2}}\right) = \left\langle \frac{1}{2}, \frac{1}{2\sqrt{2}}\right\rangle,\ \ \mathbf{F}\left(\mathbf{r}\left(\frac{1}{\sqrt{2}}\right)\right) = \left\langle e^{-1/2}, \frac{1}{4\sqrt{2}}\right\rangle$;

$\mathbf{r}(1) = \langle 1, 1\rangle,\ \ \mathbf{F}(\mathbf{r}(1)) = \langle 1, 1\rangle$.

In order to generate the graph with Maple, we use the `line` command in the `plottools` package to define each of the vectors. For example,

```
v1:=line([0,0],[exp(-1),0]):
```

generates the vector from the vector field at the point $(0,0)$ (but without an arrowhead) and gives it the name `v1`. To show everything on the same screen, we use the `display` command. In Mathematica, we use `ListPlot` (with the `PlotJoined -> True` option) to generate the vectors, and then `Show` to show everything on the same screen.

32. (a) $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_{-1}^{1} \langle 2t, t^2, 3t\rangle \cdot \langle 2, 3, -2t\rangle\,dt = \int_{-1}^{1}(4t + 3t^2 - 6t^2)\,dt = \left[2t^2 - t^3\right]_{-1}^{1} = -2$

(b) Now $\mathbf{F}(\mathbf{r}(t)) = \langle 2t, t^2, 3t\rangle$, so $\mathbf{F}(\mathbf{r}(-1)) = \langle -2, 1, -3\rangle$, $\mathbf{F}\left(\mathbf{r}\left(-\frac{1}{2}\right)\right) = \left\langle -1, \frac{1}{4}, -\frac{3}{2}\right\rangle$, $\mathbf{F}\left(\mathbf{r}\left(\frac{1}{2}\right)\right) = \left\langle 1, \frac{1}{4}, \frac{3}{2}\right\rangle$, and $\mathbf{F}(\mathbf{r}(1)) = \langle 2, 1, 3\rangle$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. $x = e^{-t}\cos 4t,\ \ y = e^{-t}\sin 4t,\ \ z = e^{-t},\ \ 0 \le t \le 2\pi$.

Then $\dfrac{dx}{dt} = e^{-t}(-\sin 4t)(4) - e^{-t}\cos 4t = -e^{-t}(4\sin 4t + \cos 4t)$,

$\dfrac{dy}{dt} = e^{-t}(\cos 4t)(4) - e^{-t}\sin 4t = -e^{-t}(-4\cos 4t + \sin 4t)$, and $\dfrac{dz}{dt} = -e^{-t}$, so

$$\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} = \sqrt{(-e^{-t})^2[(4\sin 4t + \cos 4t)^2 + (-4\cos 4t + \sin 4t)^2 + 1]}$$
$$= e^{-t}\sqrt{16(\sin^2 4t + \cos^2 4t) + \sin^2 4t + \cos^2 4t + 1} = 3\sqrt{2}\,e^{-t}$$

So by Formula 9, $\int_C x^3y^2z\,ds = \int_0^{2\pi}(e^{-t}\cos 4t)^3(e^{-t}\sin 4t)^2(e^{-t})\,(3\sqrt{2}\,e^{-t})\,dt$

$$= \int_0^{2\pi} 3\sqrt{2}\,e^{-7t}\cos^3 4t\sin^2 4t\,dt = \tfrac{172{,}704}{5{,}632{,}705}\sqrt{2}\,(1 - e^{-14\pi})$$

34. (a) We parametrize the circle C as $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j},\ \ 0 \le t \le 2\pi$. So $\mathbf{F}(\mathbf{r}(t)) = \langle 4\cos^2 t, 4\cos t\sin t\rangle$,

$\mathbf{r}'(t) = \langle -2\sin t, 2\cos t\rangle$, and $W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{2\pi}(-8\cos^2 t\sin t + 8\cos^2 t\sin t)\,dt = 0$.

(b)

From the graph, we see that all of the vectors in the field are perpendicular to the path. This indicates that the field does no work on the particle, since the field never pulls the particle in the direction in which it is going. In other words, at any point along C, $\mathbf{F}\cdot\mathbf{T} = 0$, and so certainly $\int_C \mathbf{F}\cdot d\mathbf{r} = 0$.

35. We use the parametrization $x = 2\cos t$, $y = 2\sin t$, $-\frac{\pi}{2} \le t \le \frac{\pi}{2}$. Then

$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\,dt = \sqrt{(-2\sin t)^2 + (2\cos t)^2}\,dt = 2\,dt$, so $m = \int_C k\,ds = 2k\int_{-\pi/2}^{\pi/2} dt = 2k(\pi)$,

$\overline{x} = \frac{1}{2\pi k}\int_C xk\,ds = \frac{1}{2\pi}\int_{-\pi/2}^{\pi/2}(2\cos t)2\,dt = \frac{1}{2\pi}\big[4\sin t\big]_{-\pi/2}^{\pi/2} = \frac{4}{\pi}$, $\overline{y} = \frac{1}{2\pi k}\int_C yk\,ds = \frac{1}{2\pi}\int_{-\pi/2}^{\pi/2}(2\sin t)2\,dt = 0$.

Hence $(\overline{x}, \overline{y}) = \left(\frac{4}{\pi}, 0\right)$.

36. We use the parametrization $x = a\cos t,\ \ y = a\sin t,\ \ 0 \le t \le \frac{\pi}{2}$. Then

$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\,dt = \sqrt{(-a\sin t)^2 + (a\cos t)^2}\,dt = a\,dt$, so

$m = \int_C \rho(x,y)\,ds = \int_C kxy\,ds = \int_0^{\pi/2} k(a\cos t)(a\sin t)\,a\,dt = ka^3\int_0^{\pi/2}\cos t\sin t\,dt = ka^3\big[\frac{1}{2}\sin^2 t\big]_0^{\pi/2} = \frac{1}{2}ka^3$,

$$\overline{x} = \frac{1}{ka^3/2}\int_C x(kxy)\,ds = \frac{2}{ka^3}\int_0^{\pi/2} k(a\cos t)^2(a\sin t)a\,dt = \frac{2}{ka^3}\cdot ka^4\int_0^{\pi/2}\cos^2 t\,\sin t\,dt$$
$$= 2a\big[-\tfrac{1}{3}\cos^3 t\big]_0^{\pi/2} = 2a\left(0 + \tfrac{1}{3}\right) = \tfrac{2}{3}a, \text{ and}$$

$$\overline{y} = \frac{1}{ka^3/2}\int_C y(kxy)\,ds = \frac{2}{ka^3}\int_0^{\pi/2} k(a\cos t)(a\sin t)^2 a\,dt = \frac{2}{ka^3}\cdot ka^4\int_0^{\pi/2}\sin^2 t\,\cos t\,dt$$
$$= 2a\big[\tfrac{1}{3}\sin^3 t\big]_0^{\pi/2} = 2a\left(\tfrac{1}{3} - 0\right) = \tfrac{2}{3}a.$$

Therefore the mass is $\frac{1}{2}ka^3$ and the center of mass is $(\overline{x}, \overline{y}) = \left(\frac{2}{3}a, \frac{2}{3}a\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. (a) $\overline{x} = \dfrac{1}{m}\displaystyle\int_C x\rho(x,y,z)\,ds$, $\overline{y} = \dfrac{1}{m}\displaystyle\int_C y\rho(x,y,z)\,ds$, $\overline{z} = \dfrac{1}{m}\displaystyle\int_C z\rho(x,y,z)\,ds$, where $m = \int_C \rho(x,y,z)\,ds$.

(b) $ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2}\,dt = \sqrt{(2\cos t)^2 + (-2\sin t)^2 + 3^2}\,dt = \sqrt{4(\cos^2 t + \sin^2 t) + 9}\,dt = \sqrt{13}\,dt.$

$m = \int_C k\,ds = k\int_0^{2\pi}\sqrt{13}\,dt = k\sqrt{13}\int_0^{2\pi} dt = 2\pi k\sqrt{13},$

$$\overline{x} = \frac{1}{m}\int_C x\rho(x,y,z)\,ds = \frac{1}{2\pi k\sqrt{13}}\int_0^{2\pi} 2k\sqrt{13}\,\sin t\,dt = 0,$$

$$\overline{y} = \frac{1}{m}\int_C y\rho(x,y,z)\,ds = \frac{1}{2\pi k\sqrt{13}}\int_0^{2\pi} 2k\sqrt{13}\,\cos t\,dt = 0,$$

$$\overline{z} = \frac{1}{m}\int_C z\rho(x,y,z)\,ds = \frac{1}{2\pi k\sqrt{13}}\int_0^{2\pi}\left(k\sqrt{13}\right)(3t)\,dt = \frac{3}{2\pi}\left(2\pi^2\right) = 3\pi.$$ Hence, $(\overline{x},\overline{y},\overline{z}) = (0,0,3\pi)$.

38. $m = \int_C (x^2+y^2+z^2)\,ds = \int_0^{2\pi}(t^2+\cos^2 t+\sin^2 t)\sqrt{(1)^2+(-\sin t)^2+(\cos t)^2}\,dt$

$= \int_0^{2\pi}(t^2+1)\sqrt{2}\,dt = \sqrt{2}\left[\frac{1}{3}t^3+t\right]_0^{2\pi} = \sqrt{2}\left(\frac{8}{3}\pi^3+2\pi\right),$

$$\overline{x} = \frac{1}{\sqrt{2}\left(\frac{8}{3}\pi^3+2\pi\right)}\int_0^{2\pi}\sqrt{2}\,(t^3+t)\,dt = \frac{1}{\frac{8}{3}\pi^3+2\pi}\left[\tfrac{1}{4}t^4+\tfrac{1}{2}t^2\right]_0^{2\pi} = \frac{4\pi^4+2\pi^2}{\frac{8}{3}\pi^3+2\pi}\cdot\frac{3/(2\pi)}{3/(2\pi)}$$

$$= \frac{3\pi\left(2\pi^2+1\right)}{4\pi^2+3},$$

$$\overline{y} = \frac{1}{\sqrt{2}\left(\frac{8}{3}\pi^3+2\pi\right)}\int_0^{2\pi}\left(\sqrt{2}\cos t\right)(t^2+1)\,dt = 0,$$ and

$$\overline{z} = \frac{1}{\sqrt{2}\left(\frac{8}{3}\pi^3+2\pi\right)}\int_0^{2\pi}\left(\sqrt{2}\sin t\right)(t^2+1)\,dt = 0.$$ Hence, $(\overline{x},\overline{y},\overline{z}) = \left(\dfrac{3\pi(2\pi^2+1)}{4\pi^2+3},0,0\right)$.

39. From Example 3, $\rho(x,y) = k(1-y)$, $x = \cos t$, $y = \sin t$, and $ds = dt$, $0 \le t \le \pi \quad \Rightarrow$

$$I_x = \int_C y^2\rho(x,y)\,ds = \int_0^{\pi}\sin^2 t\,[k(1-\sin t)]\,dt = k\int_0^{\pi}(\sin^2 t - \sin^3 t)\,dt$$

$$= \tfrac{1}{2}k\int_0^{\pi}(1-\cos 2t)\,dt - k\int_0^{\pi}(1-\cos^2 t)\sin t\,dt \qquad \left[\begin{array}{c}\text{Let } u = \cos t,\ du = -\sin t\,dt \\ \text{in the second integral}\end{array}\right]$$

$$= k\left[\tfrac{\pi}{2} + \int_1^{-1}(1-u^2)\,du\right] = k\left(\tfrac{\pi}{2} - \tfrac{4}{3}\right)$$

$$I_y = \int_C x^2\rho(x,y)\,ds = k\int_0^{\pi}\cos^2 t\,(1-\sin t)\,dt = \tfrac{k}{2}\int_0^{\pi}(1+\cos 2t)\,dt - k\int_0^{\pi}\cos^2 t\sin t\,dt$$

$= k\left(\frac{\pi}{2} - \frac{2}{3}\right)$, using the same substitution as above.

40. The wire is given as $x = 2\sin t$, $y = 2\cos t$, $z = 3t$, $0 \le t \le 2\pi$ with $\rho(x,y,z) = k$. Then

$ds = \sqrt{(2\cos t)^2 + (-2\sin t)^2 + 3^2}\,dt = \sqrt{4(\cos^2 t + \sin^2 t) + 9}\,dt = \sqrt{13}\,dt$ and

$I_x = \int_C (y^2+z^2)\rho(x,y,z)\,ds = \int_0^{2\pi}(4\cos^2 t + 9t^2)(k)\sqrt{13}\,dt = \sqrt{13}\,k\left[4\left(\frac{1}{2}t + \frac{1}{4}\sin 2t\right) + 3t^3\right]_0^{2\pi}$

$= \sqrt{13}\,k(4\pi + 24\pi^3) = 4\sqrt{13}\,\pi k(1+6\pi^2)$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$I_y = \int_C (x^2+z^2)\rho(x,y,z)\,ds = \int_0^{2\pi} \left(4\sin^2 t + 9t^2\right)(k)\sqrt{13}\,dt = \sqrt{13}\,k\left[4\left(\tfrac{1}{2}t - \tfrac{1}{4}\sin 2t\right) + 3t^3\right]_0^{2\pi}$$
$$= \sqrt{13}\,k(4\pi + 24\pi^3) = 4\sqrt{13}\,\pi k(1+6\pi^2)$$

$$I_z = \int_C (x^2+y^2)\rho(x,y,z)\,ds = \int_0^{2\pi}(4\sin^2 t + 4\cos^2 t)(k)\sqrt{13}\,dt = 4\sqrt{13}\,k\int_0^{2\pi} dt = 8\pi\sqrt{13}\,k$$

41. $W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{2\pi} \langle t - \sin t, (1-\cos t)+2\rangle \cdot \langle 1-\cos t, \sin t\rangle\,dt$

$$= \int_0^{2\pi}(t - t\cos t - \sin t + \sin t\cos t + 3\sin t - \sin t\cos t)\,dt$$
$$= \int_0^{2\pi}(t - t\cos t + 2\sin t)\,dt = \left[\tfrac{1}{2}t^2 - (t\sin t + \cos t) - 2\cos t\right]_0^{2\pi} \qquad \left[\begin{array}{l}\text{integrate by parts}\\ \text{in the second term}\end{array}\right]$$
$$= 2\pi^2$$

42. Choosing y as the parameter, the curve C is parametrized by $x = y^2+1$, $y = y$, $0 \le y \le 1$. Then

$$W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1 \left\langle \left(y^2+1\right)^2, ye^{y^2+1}\right\rangle \cdot \langle 2y, 1\rangle\,dy = \int_0^1 \left[2y\left(y^2+1\right)^2 + ye^{y^2+1}\right]dy$$
$$= \left[\tfrac{1}{3}\left(y^2+1\right)^3 + \tfrac{1}{2}e^{y^2+1}\right]_0^1 = \tfrac{8}{3} + \tfrac{1}{2}e^2 - \tfrac{1}{3} - \tfrac{1}{2}e = \tfrac{1}{2}e^2 - \tfrac{1}{2}e + \tfrac{7}{3}$$

43. $\mathbf{r}(t) = \langle 2t, t, 1-t\rangle$, $0 \le t \le 1$.

$$W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1 \left\langle 2t - t^2, t - (1-t)^2, 1 - t - (2t)^2\right\rangle \cdot \langle 2, 1, -1\rangle\,dt$$
$$= \int_0^1 (4t - 2t^2 + t - 1 + 2t - t^2 - 1 + t + 4t^2)\,dt = \int_0^1 (t^2 + 8t - 2)\,dt = \left[\tfrac{1}{3}t^3 + 4t^2 - 2t\right]_0^1 = \tfrac{7}{3}$$

44. $\mathbf{r}(t) = 2\,\mathbf{i} + t\,\mathbf{j} + 5t\,\mathbf{k}$, $0 \le t \le 1$. Therefore

$$W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1 \frac{K\langle 2, t, 5t\rangle}{(4+26t^2)^{3/2}} \cdot \langle 0, 1, 5\rangle\,dt = K\int_0^1 \frac{26t}{(4+26t^2)^{3/2}}\,dt = K\left[-(4+26t^2)^{-1/2}\right]_0^1 = K\left(\tfrac{1}{2} - \tfrac{1}{\sqrt{30}}\right).$$

45. (a) $\mathbf{r}(t) = at^2\,\mathbf{i} + bt^3\,\mathbf{j} \quad\Rightarrow\quad \mathbf{v}(t) = \mathbf{r}'(t) = 2at\,\mathbf{i} + 3bt^2\,\mathbf{j} \quad\Rightarrow\quad \mathbf{a}(t) = \mathbf{v}'(t) = 2a\,\mathbf{i} + 6bt\,\mathbf{j}$, and force is mass times acceleration: $\mathbf{F}(t) = m\,\mathbf{a}(t) = 2ma\,\mathbf{i} + 6mbt\,\mathbf{j}$, $0 \le t \le 1$.

(b) $W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1 (2ma\,\mathbf{i} + 6mbt\,\mathbf{j})\cdot(2at\,\mathbf{i} + 3bt^2\,\mathbf{j})\,dt = \int_0^1 (4ma^2t + 18mb^2t^3)\,dt$

$$= \left[2ma^2t^2 + \tfrac{9}{2}mb^2t^4\right]_0^1 = 2ma^2 + \tfrac{9}{2}mb^2$$

46. $\mathbf{r}(t) = a\sin t\,\mathbf{i} + b\cos t\,\mathbf{j} + ct\,\mathbf{k} \quad\Rightarrow\quad \mathbf{v}(t) = \mathbf{r}'(t) = a\cos t\,\mathbf{i} - b\sin t\,\mathbf{j} + c\,\mathbf{k} \quad\Rightarrow\quad \mathbf{a}(t) = \mathbf{v}'(t) = -a\sin t\,\mathbf{i} - b\cos t\,\mathbf{j}$ and $\mathbf{F}(t) = m\,\mathbf{a}(t) = -ma\sin t\,\mathbf{i} - mb\cos t\,\mathbf{j}$. Thus

$$W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{\pi/2}(-ma\sin t\,\mathbf{i} - mb\cos t\,\mathbf{j})\cdot(a\cos t\,\mathbf{i} - b\sin t\,\mathbf{j} + c\,\mathbf{k})\,dt$$
$$= \int_0^{\pi/2}(-ma^2\sin t\cos t + mb^2\sin t\cos t)\,dt = m(b^2-a^2)\left[\tfrac{1}{2}\sin^2 t\right]_0^{\pi/2} = \tfrac{1}{2}m(b^2-a^2)$$

47. The combined weight of the man and the paint is 185 lb, so the force exerted (equal and opposite to that exerted by gravity) is $\mathbf{F} = 185\,\mathbf{k}$. To parametrize the staircase, let $x = 20\cos t$, $y = 20\sin t$, $z = \frac{90}{6\pi}t = \frac{15}{\pi}t$, $0 \le t \le 6\pi$. Then the work done is

$$W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{6\pi}\langle 0, 0, 185\rangle \cdot \left\langle -20\sin t, 20\cos t, \tfrac{15}{\pi}\right\rangle dt = (185)\tfrac{15}{\pi}\int_0^{6\pi} dt = (185)\left(\tfrac{15}{\pi}\right)(6\pi) \approx 1.67\times 10^4 \text{ ft-lb}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. This time m is a function of t: $m = 185 - \frac{9}{6\pi}t = 185 - \frac{3}{2\pi}t$. So let $\mathbf{F} = \left(185 - \frac{3}{2\pi}t\right)\mathbf{k}$. To parametrize the staircase, let $x = 20\cos t,\ \ y = 20\sin t,\ \ z = \frac{90}{6\pi}t = \frac{15}{\pi}t,\ \ 0 \le t \le 6\pi$. Therefore

$$W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{6\pi} \left\langle 0, 0, 185 - \tfrac{3}{2\pi}t\right\rangle \cdot \left\langle -20\sin t, 20\cos t, \tfrac{15}{\pi}\right\rangle dt = \tfrac{15}{\pi}\int_0^{6\pi}\left(185 - \tfrac{3}{2\pi}t\right) dt$$
$$= \tfrac{15}{\pi}\left[185t - \tfrac{3}{4\pi}t^2\right]_0^{6\pi} = 90\left(185 - \tfrac{9}{2}\right) \approx 1.62 \times 10^4 \text{ ft-lb}$$

49. (a) $\mathbf{r}(t) = \langle \cos t, \sin t\rangle,\ \ 0 \le t \le 2\pi$, and let $\mathbf{F} = \langle a, b\rangle$. Then

$$W = \int_C \mathbf{F}\cdot d\,\mathbf{r} = \int_0^{2\pi} \langle a, b\rangle \cdot \langle -\sin t, \cos t\rangle\ dt = \int_0^{2\pi}(-a\sin t + b\cos t)\,dt = \left[a\cos t + b\sin t\right]_0^{2\pi}$$
$$= a + 0 - a + 0 = 0$$

(b) Yes. $\mathbf{F}(x, y) = k\,\mathbf{x} = \langle kx, ky\rangle$ and

$$W = \int_C \mathbf{F}\cdot d\,\mathbf{r} = \int_0^{2\pi} \langle k\cos t, k\sin t\rangle \cdot \langle -\sin t, \cos t\rangle\, dt = \int_0^{2\pi}(-k\sin t\ \cos t + k\sin t\ \cos t)\,dt = \int_0^{2\pi} 0\,dt = 0.$$

50. Consider the base of the fence in the xy-plane, centered at the origin, with the height given by $z = h(x, y) = 4 + 0.01(x^2 - y^2)$. To graph the fence, observe that the fence is highest when $y = 0$ (where the height is 5 m) and lowest when $x = 0$ (a height of 3 m). When $y = \pm x$, the height is 4 m.

Also, the fence can be graphed using parametric equations (see Section 16.6): $x = 10\cos u,\ \ y = 10\sin u$,

$$z = v\left[4 + 0.01((10\cos u)^2 - (10\sin u)^2)\right] = v(4 + \cos^2 u - \sin^2 u)$$
$$= v(4 + \cos 2u),\ 0 \le u \le 2\pi,\ 0 \le v \le 1.$$

The surface area of one side of the fence is $\int_C h(x, y)\,ds$, where the base C of the fence is given by $x = 10\cos t,\ \ y = 10\sin t,\ \ 0 \le t \le 2\pi$. Then

$$\int_C h(x, y)\,ds = \int_0^{2\pi}\left[4 + 0.01((10\cos t)^2 - (10\sin t)^2)\right]\sqrt{(-10\sin t)^2 + (10\cos t)^2}\,dt$$
$$= \int_0^{2\pi}(4 + \cos 2t)\sqrt{100}\,dt = 10\left[4t + \tfrac{1}{2}\sin 2t\right]_0^{2\pi} = 10(8\pi) = 80\pi \text{ m}^2$$

If we paint both sides of the fence, the total surface area to cover is 160π m^2, and since 1 L of paint covers 100 m^2, we require $\frac{160\pi}{100} = 1.6\pi \approx 5.03$ L of paint.

51. Let $\mathbf{r}(t) = \langle x(t), y(t), z(t)\rangle$ and $\mathbf{v} = \langle v_1, v_2, v_3\rangle$. Then

$$\int_C \mathbf{v}\cdot d\mathbf{r} = \int_a^b \langle v_1, v_2, v_3\rangle \cdot \langle x'(t), y'(t), z'(t)\rangle\, dt = \int_a^b \left[v_1\, x'(t) + v_2\, y'(t) + v_3\, z'(t)\right] dt$$
$$= \left[v_1\, x(t) + v_2\, y(t) + v_3\, z(t)\right]_a^b = \left[v_1\, x(b) + v_2\, y(b) + v_3\, z(b)\right] - \left[v_1\, x(a) + v_2\, y(a) + v_3\, z(a)\right]$$
$$= v_1\left[x(b) - x(a)\right] + v_2\left[y(b) - y(a)\right] + v_3\left[z(b) - z(a)\right]$$
$$= \langle v_1, v_2, v_3\rangle \cdot \langle x(b) - x(a), y(b) - y(a), z(b) - z(a)\rangle$$
$$= \langle v_1, v_2, v_3\rangle \cdot \left[\langle x(b), y(b), z(b)\rangle - \langle x(a), y(a), z(a)\rangle\right] = \mathbf{v}\cdot\left[\mathbf{r}(b) - \mathbf{r}(a)\right]$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

52. If $\mathbf{r}(t) = \langle x(t), y(t), z(t)\rangle$ then

$$\begin{aligned}\int_C \mathbf{r}\cdot d\mathbf{r} &= \int_a^b \langle x(t), y(t), z(t)\rangle \cdot \langle x'(t), y'(t), z'(t)\rangle\, dt = \int_a^b [x(t)\,x'(t) + y(t)\,y'(t) + z(t)\,z'(t)]\, dt \\ &= \left[\tfrac{1}{2}[x(t)]^2 + \tfrac{1}{2}[y(t)]^2 + \tfrac{1}{2}[z(t)]^2\right]_a^b \\ &= \tfrac{1}{2}\left\{\left([x(b)]^2 + [y(b)]^2 + [z(b)]^2\right) - \left([x(a)]^2 + [y(a)]^2 + [z(a)]^2\right)\right\} \\ &= \tfrac{1}{2}\left[|\mathbf{r}(b)|^2 - |\mathbf{r}(a)|^2\right]\end{aligned}$$

53. The work done in moving the object is $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C \mathbf{F}\cdot\mathbf{T}\, ds$. We can approximate this integral by dividing C into 7 segments of equal length $\Delta s = 2$ and approximating $\mathbf{F}\cdot\mathbf{T}$, that is, the tangential component of force, at a point (x_i^*, y_i^*) on each segment. Since C is composed of straight line segments, $\mathbf{F}\cdot\mathbf{T}$ is the scalar projection of each force vector onto C. If we choose (x_i^*, y_i^*) to be the point on the segment closest to the origin, then the work done is

$$\int_C \mathbf{F}\cdot\mathbf{T}\, ds \approx \sum_{i=1}^{7} [\mathbf{F}(x_i^*, y_i^*)\cdot\mathbf{T}(x_i^*, y_i^*)]\,\Delta s = [2+2+2+2+1+1+1](2) = 22$$

Thus, we estimate the work done to be approximately 22 J.

54. Use the orientation pictured in the figure. Then since $\mathbf{B}$ is tangent to any circle that lies in the plane perpendicular to the wire, $\mathbf{B} = |\mathbf{B}|\,\mathbf{T}$ where $\mathbf{T}$ is the unit tangent to the circle C: $x = r\cos\theta,\ \ y = r\sin\theta$. Thus $\mathbf{B} = |\mathbf{B}|\,\langle -\sin\theta, \cos\theta\rangle$. Then $\int_C \mathbf{B}\cdot d\mathbf{r} = \int_0^{2\pi} |\mathbf{B}|\,\langle -\sin\theta, \cos\theta\rangle \cdot \langle -r\sin\theta, r\cos\theta\rangle\, d\theta = \int_0^{2\pi} |\mathbf{B}|\, r\, d\theta = 2\pi r\,|\mathbf{B}|$. (Note that $|\mathbf{B}|$ here is the magnitude of the field at a distance r from the wire's center.) But by Ampere's Law $\int_C \mathbf{B}\cdot d\mathbf{r} = \mu_0 I$. Hence $|\mathbf{B}| = \mu_0 I/(2\pi r)$.

16.3 The Fundamental Theorem for Line Integrals

1. C appears to be a smooth curve, and since ∇f is continuous, we know f is differentiable. Then Theorem 2 says that the value of $\int_C \nabla f\cdot d\mathbf{r}$ is simply the difference of the values of f at the terminal and initial points of C. From the graph, this is $50 - 10 = 40$.

2. C is represented by the vector function $\mathbf{r}(t) = (t^2+1)\,\mathbf{i} + (t^3+t)\,\mathbf{j}$, $0 \le t \le 1$, so $\mathbf{r}'(t) = 2t\,\mathbf{i} + (3t^2+1)\,\mathbf{j}$. Since $3t^2 + 1 \neq 0$, we have $\mathbf{r}'(t) \neq \mathbf{0}$, thus C is a smooth curve. ∇f is continuous, and hence f is differentiable, so by Theorem 2 we have $\int_C \nabla f\cdot d\mathbf{r} = f(\mathbf{r}(1)) - f(\mathbf{r}(0)) = f(2,2) - f(1,0) = 9 - 3 = 6$.

3. Let $P(x,y) = xy + y^2$ and $Q(x,y) = x^2 + 2xy$. Then $\partial P/\partial y = x + 2y$ and $\partial Q/\partial x = 2x + 2y$. Since $\partial P/\partial y \neq \partial Q/\partial x$, $\mathbf{F}(x,y) = P\,\mathbf{i} + Q\,\mathbf{j}$ is not conservative by Theorem 5.

4. $\partial(y^2 - 2x)/\partial y = 2y = \partial(2xy)/\partial x$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$ which is open and simply-connected, so $\mathbf{F}$ is conservative by Theorem 6. Thus, there exists a function f such that $\nabla f = \mathbf{F}$, that is, $f_x(x,y) = y^2 - 2x$ and $f_y(x,y) = 2xy$. But $f_x(x,y) = y^2 - 2x$ implies $f(x,y) = xy^2 - x^2 + g(y)$ and differentiating both sides of this equation with respect to y gives $f_y(x,y) = 2xy + g'(y)$. Thus $2xy = 2xy + g'(y)$ so $g'(y) = 0$ and $g(y) = K$ where K is a constant. Hence $f(x,y) = xy^2 - x^2 + K$ is a potential function for $\mathbf{F}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $\dfrac{\partial}{\partial y}\left(y^2e^{xy}\right) = y^2 \cdot xe^{xy} + 2ye^{xy} = (xy^2 + 2y)e^{xy}$,

$\dfrac{\partial}{\partial x}\left[(1+xy)e^{xy}\right] = (1+xy)\cdot ye^{xy} + ye^{xy} = ye^{xy} + xy^2e^{xy} + ye^{xy} = (xy^2+2y)e^{xy}$.

Since these partial derivatives are equal and the domain of $\mathbf{F}$ is $\mathbb{R}^2$ which is open and simply-connected, $\mathbf{F}$ is conservative by Theorem 6. Thus, there exists a function f such that $\nabla f = \mathbf{F}$, that is, $f_x(x,y) = y^2e^{xy}$ and $f_y(x,y) = (1+xy)e^{xy}$. But $f_x(x,y) = y^2e^{xy}$ implies $f(x,y) = ye^{xy} + g(y)$ and differentiating both sides of this equation with respect to y gives $f_y(x,y) = (1+xy)e^{xy} + g'(y)$. Thus $(1+xy)e^{xy} = (1+xy)e^{xy} + g'(y)$ so $g'(y) = 0$ and $g(y) = K$ where K is a constant. Hence $f(x,y) = ye^{xy} + K$ is a potential function for $\mathbf{F}$.

6. $\partial(ye^x)/\partial y = e^x = \partial(e^x + e^y)/\partial x$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$ which is open and simply-connected, so $\mathbf{F}$ is conservative. Hence there exists a function f such that $\nabla f = \mathbf{F}$. Here $f_x(x,y) = ye^x$ implies $f(x,y) = ye^x + g(y)$ and then $f_y(x,y) = e^x + g'(y)$. But $f_y(x,y) = e^x + e^y$ so $g'(y) = e^y \quad \Rightarrow \quad g(y) = e^y + K$ and $f(x,y) = ye^x + e^y + K$ is a potential function for $\mathbf{F}$.

7. $\partial(ye^x + \sin y)/\partial y = e^x + \cos y = \partial(e^x + x\cos y)/\partial x$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$. Hence $\mathbf{F}$ is conservative so there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x,y) = ye^x + \sin y$ implies $f(x,y) = ye^x + x\sin y + g(y)$ and $f_y(x,y) = e^x + x\cos y + g'(y)$. But $f_y(x,y) = e^x + x\cos y$ so $g(y) = K$ and $f(x,y) = ye^x + x\sin y + K$ is a potential function for $\mathbf{F}$.

8. $\partial(2xy + y^{-2})/\partial y = 2x - 2y^{-3} = \partial(x^2 - 2xy^{-3})/\partial x$ and the domain of $\mathbf{F}$ is $\{(x,y) \mid y > 0\}$ which is open and simply-connected. Hence $\mathbf{F}$ is conservative, so there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x,y) = 2xy + y^{-2}$ implies $f(x,y) = x^2y + xy^{-2} + g(y)$ and $f_y(x,y) = x^2 - 2xy^{-3} + g'(y)$. But $f_y(x,y) = x^2 - 2xy^{-3}$ so $g'(y) = 0 \quad \Rightarrow \quad g(y) = K$. Then $f(x,y) = x^2y + xy^{-2} + K$ is a potential function for $\mathbf{F}$.

9. $\partial(y^2\cos x + \cos y)/\partial y = 2y\cos x - \sin y = \partial(2y\sin x - x\sin y)/\partial x$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$ which is open and simply-connected. Hence $\mathbf{F}$ is conservative so there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x,y) = y^2\cos x + \cos y$ implies $f(x,y) = y^2\sin x + x\cos y + g(y)$ and $f_y(x,y) = 2y\sin x - x\sin y + g'(y)$. But $f_y(x,y) = 2y\sin x - x\sin y$ so $g'(y) = 0 \quad \Rightarrow \quad g(y) = K$ and $f(x,y) = y^2\sin x + x\cos y + K$ is a potential function for $\mathbf{F}$.

10. $\partial(\ln y + y/x)/\partial y = 1/y + 1/x = \partial(\ln x + x/y)/\partial x$ and the domain of $\mathbf{F}$ is $\{(x,y) \mid x > 0,\ y > 0\}$ which is open and simply-connected. Hence $\mathbf{F}$ is conservative so there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x,y) = \ln y + y/x$ implies $f(x,y) = x\ln y + y\ln x + g(y)$ and $f_y(x,y) = x/y + \ln x + g'(y)$. But $f_y(x,y) = \ln x + x/y$ so $g'(y) = 0 \quad \Rightarrow$ $g(y) = K$ and $f(x,y) = x\ln y + y\ln x + K$ is a potential function for $\mathbf{F}$.

11. (a) $\mathbf{F}$ has continuous first-order partial derivatives and $\dfrac{\partial}{\partial y}(2xy) = 2x = \dfrac{\partial}{\partial x}(x^2)$ on $\mathbb{R}^2$, which is open and simply-connected. Thus, $\mathbf{F}$ is conservative by Theorem 6. Then we know that the line integral of $\mathbf{F}$ is independent of path;

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

in particular, the value of $\int_C \mathbf{F} \cdot d\mathbf{r}$ depends only on the endpoints of C. Since all three curves have the same initial and terminal points, $\int_C \mathbf{F} \cdot d\mathbf{r}$ will have the same value for each curve.

(b) We first find a potential function f, so that $\nabla f = \mathbf{F}$. We know $f_x(x, y) = 2xy$ and $f_y(x, y) = x^2$. Integrating $f_x(x, y)$ with respect to x, we have $f(x, y) = x^2 y + g(y)$. Differentiating both sides with respect to y gives $f_y(x, y) = x^2 + g'(y)$, so we must have $x^2 + g'(y) = x^2 \quad \Rightarrow \quad g'(y) = 0 \quad \Rightarrow \quad g(y) = K$, a constant. Thus $f(x, y) = x^2 y + K$, and we can take $K = 0$. All three curves start at $(1, 2)$ and end at $(3, 2)$, so by Theorem 2, $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f(3, 2) - f(1, 2) = 3^2(2) - 1^2(2) = 16$ for each curve.

12. (a) $\mathbf{F}(x, y) = 2xy\,\mathbf{i} + (x^2 + \sin y)\,\mathbf{j}$.

Solution 1: $\mathbf{F}$ has continuous first-order partial derivatives and $\dfrac{\partial(2xy)}{\partial y} = 2x = \dfrac{\partial(x^2 + \sin y)}{\partial x}$ on $\mathbb{R}^2$, which is open and simply-connected. Therefore, the vector field is conservative and there exists a function $f(x, y)$ such that $\nabla f = \mathbf{F}$. Here, $f_x(x, y) = 2xy$ implies $f(x, y) = x^2 y + g(y)$ and $f_y(x, y) = x^2 + g'(y)$, but $f_y(x, y) = x^2 + \sin y$, which implies $g'(y) = \sin y \quad \Rightarrow \quad g(y) = -\cos y + K$ and $f(x, y) = x^2 y - \cos y + K$ is a potential function for $\mathbf{F}$. Then

$$\int_C \mathbf{F} \cdot d\mathbf{r} = f\left(2, \tfrac{\pi}{2}\right) - f(0, 0) = 2^2\left(\tfrac{\pi}{2}\right) - \cos\left(\tfrac{\pi}{2}\right) + K - 0 + \cos 0 - K = 2\pi + 1$$

Solution 2: As in Example 4, since $\mathbf{F}$ is conservative, we know that $\int_C \mathbf{F} \cdot d\mathbf{r}$ is independent of path, so we replace the curve C by the simpler curve C_1 consisting of the line segment connecting the two endpoints of C. Thus, C_1 can be represented by $\mathbf{r}(t) = 2t\,\mathbf{i} + \frac{\pi}{2}t\,\mathbf{j}$, $0 \le t \le 1$. Then

$$\begin{aligned}
\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\,dt \\
&= \int_0^1 \left\{2(2t)\left(\tfrac{\pi}{2}t\right)(2) + \left[(2t)^2 + \sin\left(\tfrac{\pi}{2}t\right)\right]\left(\tfrac{\pi}{2}\right)\right\} dt = \left[\int_0^1 6\pi t^2 + \tfrac{\pi}{2}\sin\left(\tfrac{\pi}{2}t\right)\right] dt \\
&= \left[2\pi t^3 - \cos\left(\tfrac{\pi}{2}t\right)\right]_0^1 = (2\pi - 0) - (0 - 1) = 2\pi + 1
\end{aligned}$$

(b) From part (a), we know that $\mathbf{F}$ is conservative and therefore independent of path. Thus, since C is a closed path, by Theorem 3, $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$.

13. (a) $\mathbf{F}(x, y) = (3x^2 + y^2)\,\mathbf{i} + 2xy\,\mathbf{j}$. $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j}$, $\pi \le t \le 2\pi$. Then

$$\mathbf{F}(\mathbf{r}(t)) = [3(2\cos t)^2 + (2\sin t)^2]\,\mathbf{i} + 2(2\cos t)(2\sin t)\,\mathbf{j} = (12\cos^2 t + 4\sin^2 t)\,\mathbf{i} + 8\cos t\sin t\,\mathbf{j}$$

and $\mathbf{r}'(t) = -2\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j}$, so

$$\begin{aligned}
\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_\pi^{2\pi} \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\,dt = \int_\pi^{2\pi} \left[-2\sin t(12\cos^2 t + 4\sin^2 t) + 2\cos t(8\cos t\sin t)\right] dt \\
&= \int_\pi^{2\pi} (-8\sin t\cos^2 t - 8\sin^3 t)\,dt = -8\int_\pi^{2\pi} \sin t\,dt = 8\Big[\cos t\Big]_\pi^{2\pi} = 8[1 - (-1)] = 16
\end{aligned}$$

(b) $\dfrac{\partial(3x^2 + y^2)}{\partial y} = 2y = \dfrac{\partial(2xy)}{\partial x}$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$, which is open and simply-connected. Thus, $\mathbf{F}$ is conservative and there exists a function $f(x, y)$ such that $\nabla f = \mathbf{F}$. Then $f_x(x, y) = 3x^2 + y^2$ implies $f(x, y) = x^3 + xy^2 + g(y)$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Differentiating both sides with respect to y gives $f_y(x,y) = 2xy + g'(y)$, so we must have $2xy + g'(y) = 2xy \quad \Rightarrow$ $g'(y) = 0 \quad \Rightarrow \quad g(y) = K$, a constant. Thus, $f(x,y) = x^3 + xy^2 + K$ is a potential function for $\mathbf{F}$.

(c) From part (b), $f(x,y) = x^3 + xy^2 + K$. Thus,

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f(2,0) - f(-2,0) = 2^3 + 0 + K - (-2)^3 - 0 - K = 16$$

(d) We replace C with the line segment from $(-2,0)$ to $(2,0)$ so $\mathbf{r}(t) = \langle t, 0\rangle$, $-2 \le t \le 2$, which implies $\mathbf{r}'(t) = \langle 1, 0\rangle$ and $\mathbf{F}(\mathbf{r}(t)) = \langle 3t^2, 0\rangle$. Thus, $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_{-2}^{2} \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\, dt = \int_{-2}^{2} 3t^2\, dt = \left[t^3\right]_{-2}^{2} = 2^3 - (-2)^3 = 16$.

14. (a) $\mathbf{F}(x,y) = \langle \sin y + e^x, x\cos y\rangle$ and C: $x = t$, $y = t(3-t)$, $0 \le t \le 3$.

$\dfrac{\partial(\sin y + e^x)}{\partial y} = \cos y = \dfrac{\partial(x\cos y)}{\partial x}$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$, which is open and simply-connected. Thus, $\mathbf{F}$ is conservative and there exists a function $f(x,y)$ such that $\nabla f = \mathbf{F}$. Then $f_x(x,y) = \sin y + e^x$ implies $f(x,y) = x\sin y + e^x + g(y)$, so $f_y(x,y) = x\cos y + g'(y)$, which implies $g'(y) = 0 \quad \Rightarrow \quad g(y) = K$. Therefore, $f(x,y) = x\sin y + e^x + K$ is a potential function for $\mathbf{F}$.

(b) The endpoints of C are $(0,0)$ and $(3,0)$. Thus, by Theorem 2,

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_C \nabla f \cdot d\mathbf{r} = f(3,0) - f(0,0)\\ &= 3 \cdot \sin 0 + e^3 + K - (0 \cdot \sin 0 + e^0 + K)\\ &= e^3 - 1\end{aligned}$$

(c) We replace C with the line segment that connects the endpoints of C along the x-axis. So $x = t$, $y = 0$, $0 \le t \le 3$. Then $\mathbf{r}(t) = \langle t, 0\rangle$, $\mathbf{r}'(t) = \langle 1, 0\rangle$, and $\mathbf{F}(\mathbf{r}(t)) = \langle \sin 0 + e^t, t\cos 0\rangle = \langle e^t, t\rangle$. Therefore, $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^3 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\, dt = \int_0^3 e^t\, dt = \left[e^t\right]_0^3 = e^3 - 1$.

15. (a) $\mathbf{F}(x,y) = \langle ye^{xy}, xe^{xy}\rangle$ and C: $x = \sin \frac{\pi}{2}t$, $y = e^{t-1}(1 - \cos \pi t)$, $0 \le t \le 1$.

$\dfrac{\partial(ye^{xy})}{\partial y} = e^{xy} + xye^{xy} = \dfrac{\partial(xe^{xy})}{\partial x}$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$, which is open and simply connected. Thus, $\mathbf{F}$ is conservative and there exists a function $f(x,y)$ such that $\nabla f = \mathbf{F}$. Then $f_x(x,y) = ye^{xy}$ implies $f(x,y) = e^{xy} + g(y)$, so $f_y(x,y) = xe^{xy} + g'(y)$, which implies $g'(y) = 0 \quad \Rightarrow \quad g(y) = K$. Therefore, $f(x,y) = e^{xy} + K$ is a potential function for $\mathbf{F}$.

(b) The endpoints of C are $(0,0)$ and $(1,2)$. Thus, by Theorem 2,

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_C \nabla f \cdot d\mathbf{r} = f(1,2) - f(0,0)\\ &= e^2 + K - e^0 - K = e^2 - 1\end{aligned}$$

(c) We replace C with the line segment that connects the endpoints of C. So $x = t$ and $y = 2t$, $0 \le t \le 1$. Then $\mathbf{r}(t) = \langle t, 2t\rangle$, $\mathbf{r}'(t) = \langle 1, 2\rangle$, and

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{F}(\mathbf{r}(t)) = \left\langle 2te^{2t^2}, te^{2t^2} \right\rangle$. Therefore,

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\, dt = \int_0^1 4te^{2t^2}\, dt \\ &= \int_0^2 e^u\, du \qquad [u = 2t^2,\ du = 4t\, dt] \\ &= \left[e^u\right]_0^2 = e^2 - 1 \end{aligned}$$

16. $f(x, y, z) = xy^2z + x^2$ and C: $x = t^2$, $y = e^{t^2-1}$, $z = t^2 + t$, $-1 \le t \le 1$. ∇f is the gradient vector field of f and therefore conservative. Thus, by Theorem 2, $\int_C \nabla f \cdot d\mathbf{r} = f(1, 1, 2) - f(1, 1, 0) = [(1)(1^2)(2) + 1^2] - (0 + 1^2) = 2$.

17. (a) $\mathbf{F}(x, y) = \langle 2x, 4y \rangle$. If $\mathbf{F} = \nabla f$, then $f_x(x, y) = 2x$ and $f_y(x, y) = 4y$. $f_x(x, y) = 2x$ implies that $f(x, y) = x^2 + g(y)$ and $f_y(x, y) = g'(y) = 4y$, so $g(y) = 2y^2 + K$. We can take $K = 0$, so $f(x, y) = x^2 + 2y^2$.

(b) C is a smooth curve with initial point $(4, -2)$ and terminal point $(1, 1)$, so by Theorem 2, $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f(1, 1) - f(4, -2) = (1 + 2) - (16 + 8) = -21$.

18. (a) $\mathbf{F}(x, y) = (3 + 2xy^2)\,\mathbf{i} + 2x^2y\,\mathbf{j}$. If $\mathbf{F} = \nabla f$, then $f_x(x, y) = 3 + 2xy^2$ and $f_y(x, y) = 2x^2y$. $f_x(x, y) = 3 + 2xy^2$ implies $f(x, y) = 3x + x^2y^2 + g(y)$ and $f_y(x, y) = 2x^2y + g'(y)$. But $f_y(x, y) = 2x^2y$ so $g'(y) = 0 \quad \Rightarrow \quad g(y) = K$. We can take $K = 0$, so $f(x, y) = 3x + x^2y^2$.

(b) C is a smooth curve with initial point $(1, 1)$ and terminal point $\left(4, \frac{1}{4}\right)$, so by Theorem 2, $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f\left(4, \frac{1}{4}\right) - f(1, 1) = (12 + 1) - (3 + 1) = 9$.

19. (a) $\mathbf{F}(x, y) = x^2y^3\,\mathbf{i} + x^3y^2\,\mathbf{j}$. If $\mathbf{F} = \nabla f$, then $f_x(x, y) = x^2y^3$ and $f_y(x, y) = x^3y^2$. $f_x(x, y) = x^2y^3$ implies $f(x, y) = \frac{1}{3}x^3y^3 + g(y)$ and $f_y(x, y) = x^3y^2 + g'(y)$. But $f_y(x, y) = x^3y^2$, so $g'(y) = 0 \quad \Rightarrow$ $g(y) = K$, a constant. We can take $K = 0$, so $f(x, y) = \frac{1}{3}x^3y^3$.

(b) C is a smooth curve with initial point $\mathbf{r}(0) = (0, 0)$ and terminal point $\mathbf{r}(1) = (-1, 3)$, so by Theorem 2, $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f(-1, 3) - f(0, 0) = -9 - 0 = -9$.

20. (a) $\mathbf{F}(x, y) = (1 + xy)e^{xy}\,\mathbf{i} + x^2e^{xy}\,\mathbf{j}$. $f_y(x, y) = x^2e^{xy}$ implies $f(x, y) = xe^{xy} + g(x) \quad \Rightarrow$ $f_x(x, y) = xye^{xy} + e^{xy} + g'(x) = (1 + xy)e^{xy} + g'(x)$. But $f_x(x, y) = (1 + xy)e^{xy}$ so $g'(x) = 0 \quad \Rightarrow \quad g(x) = K$. We can take $K = 0$, so $f(x, y) = xe^{xy}$.

(b) The initial point of C is $\mathbf{r}(0) = (1, 0)$ and the terminal point is $\mathbf{r}(\pi/2) = (0, 2)$, so by Theorem 2, $\int_C \mathbf{F} \cdot d\mathbf{r} = f(0, 2) - f(1, 0) = 0 - e^0 = -1$.

21. (a) $\mathbf{F}(x, y, z) = 2xy\,\mathbf{i} + (x^2 + 2yz)\,\mathbf{j} + y^2\,\mathbf{k}$. $f_x(x, y, z) = 2xy$ implies that $f(x, y, z) = x^2y + g(y, z)$ and so $f_y(x, y, z) = x^2 + g_y(y, z)$. But $f_y(x, y, z) = x^2 + 2yz$, which implies $g_y(y, z) = 2yz + h_y(z) \quad \Rightarrow$ $g(y, z) = y^2z + h(z)$. So $f(x, y, z) = x^2y + y^2z + h(z)$ and $f_z(x, y, z) = y^2 + h'(z)$. But $f_z(x, y, z) = y^2 \quad \Rightarrow$ $h'(z) = 0 \quad \Rightarrow \quad h(z) = K$. We can take $K = 0$, so $f(x, y, z) = x^2y + y^2z$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) C is a smooth curve with initial point $(2, -3, 1)$ and terminal point $(-5, 1, 2)$, so by Theorem 2,

$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f(-5, 1, 2) - f(2, -3, 1) = (25 + 2) - (-12 + 9) = 30.$

22. (a) $\mathbf{F}(x, y, z) = (y^2 z + 2xz^2)\,\mathbf{i} + 2xyz\,\mathbf{j} + (xy^2 + 2x^2 z)\,\mathbf{k}$. $f_x(x, y, z) = y^2 z + 2xz^2$ implies $f(x, y, z) = xy^2 z + x^2 z^2 + g(y, z)$ and so $f_y(x, y, z) = 2xyz + g_y(y, z)$. But $f_y(x, y, z) = 2xyz$ so $g_y(y, z) = 0 \quad \Rightarrow \quad g(y, z) = h(z)$. Thus, $f(x, y, z) = xy^2 z + x^2 z^2 + h(z)$ and $f_z(x, y, z) = xy^2 + 2x^2 z + h'(z)$. But $f_z(x, y, z) = xy^2 + 2x^2 z$, so $h'(z) = 0 \quad \Rightarrow \quad h(z) = K$. Hence, $f(x, y, z) = xy^2 z + x^2 z^2$ (taking $K = 0$).

(b) $t = 0$ corresponds to the point $(0, 1, 0)$ and $t = 1$ corresponds to $(1, 2, 1)$, so by Theorem 2,

$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f(1, 2, 1) - f(0, 1, 0) = 5 - 0 = 5.$

23. (a) $\mathbf{F}(x, y, z) = yze^{xz}\,\mathbf{i} + e^{xz}\,\mathbf{j} + xye^{xz}\,\mathbf{k}$. $f_x(x, y, z) = yze^{xz}$ implies $f(x, y, z) = ye^{xz} + g(y, z)$ and so $f_y(x, y, z) = e^{xz} + g_y(y, z)$. But $f_y(x, y, z) = e^{xz}$ so $g_y(y, z) = 0 \quad \Rightarrow \quad g(y, z) = h(z)$. Thus, $f(x, y, z) = ye^{xz} + h(z)$ and $f_z(x, y, z) = xye^{xz} + h'(z)$. But $f_z(x, y, z) = xye^{xz}$, so $h'(z) = 0 \quad \Rightarrow \quad h(z) = K$. Hence $f(x, y, z) = ye^{xz}$ (taking $K = 0$).

(b) $\mathbf{r}(0) = \langle 1, -1, 0\rangle$, $\mathbf{r}(2) = \langle 5, 3, 0\rangle$ so $\int_C \mathbf{F} \cdot d\mathbf{r} = f(5, 3, 0) - f(1, -1, 0) = 3e^0 + e^0 = 4$.

24. (a) $\mathbf{F}(x, y, z) = \sin y\,\mathbf{i} + (x\cos y + \cos z)\,\mathbf{j} - y\sin z\,\mathbf{k}$. $f_x(x, y, z) = \sin y$ implies $f(x, y, z) = x\sin y + g(y, z)$ and so $f_y(x, y, z) = x\cos y + g_y(y, z)$. But $f_y(x, y, z) = x\cos y + \cos z$ so $g_y(y, z) = \cos z \quad \Rightarrow \quad g(y, z) = y\cos z + h(z)$. Thus, $f(x, y, z) = x\sin y + y\cos z + h(z)$ and $f_z(x, y, z) = -y\sin z + h'(z)$. But $f_z(x, y, z) = -y\sin z$, so $h'(z) = 0 \quad \Rightarrow \quad h(z) = K$. Hence, $f(x, y, z) = x\sin y + y\cos z$ (taking $K = 0$).

(b) $\mathbf{r}(0) = \langle 0, 0, 0\rangle$, $\mathbf{r}(\pi/2) = \langle 1, \pi/2, \pi\rangle$ so $\int_C \mathbf{F} \cdot d\mathbf{r} = f(1, \pi/2, \pi) - f(0, 0, 0) = 1 - \frac{\pi}{2} - 0 = 1 - \frac{\pi}{2}$.

25. The functions $2xe^{-y}$ and $2y - x^2 e^{-y}$ have continuous first-order derivatives on $\mathbb{R}^2$ and $\dfrac{\partial}{\partial y}\left(2xe^{-y}\right) = -2xe^{-y} = \dfrac{\partial}{\partial x}\left(2y - x^2 e^{-y}\right)$, so $\mathbf{F}(x, y) = 2xe^{-y}\,\mathbf{i} + \left(2y - x^2 e^{-y}\right)\mathbf{j}$ is a conservative vector field by Theorem 6 and hence the line integral is independent of path. Thus a potential function f exists, and $f_x(x, y) = 2xe^{-y}$ implies $f(x, y) = x^2 e^{-y} + g(y)$ and $f_y(x, y) = -x^2 e^{-y} + g'(y)$. But $f_y(x, y) = 2y - x^2 e^{-y}$ so $g'(y) = 2y \quad \Rightarrow \quad g(y) = y^2 + K$. We can take $K = 0$, so $f(x, y) = x^2 e^{-y} + y^2$. Then $\int_C 2xe^{-y}\,dx + (2y - x^2 e^{-y})\,dy = f(2, 1) - f(1, 0) = 4e^{-1} + 1 - 1 = 4/e$.

26. The functions $\sin y$ and $x\cos y - \sin y$ have continuous first-order derivatives on $\mathbb{R}^2$ and $\dfrac{\partial}{\partial y}(\sin y) = \cos y = \dfrac{\partial}{\partial x}(x\cos y - \sin y)$, so $\mathbf{F}(x, y) = \sin y\,\mathbf{i} + (x\cos y - \sin y)\,\mathbf{j}$ is a conservative vector field by Theorem 6 and hence the line integral is independent of path. Thus a potential function f exists, and $f_x(x, y) = \sin y$ implies $f(x, y) = x\sin y + g(y)$ and $f_y(x, y) = x\cos y + g'(y)$. But $f_y(x, y) = x\cos y - \sin y$, so $g'(y) = -\sin y \quad \Rightarrow$ $g(y) = \cos y + K$. We can take $K = 0$, so $f(x, y) = x\sin y + \cos y$. Then $\int_C \sin y\,dx + (x\cos y - \sin y)\,dy = f(1, \pi) - f(2, 0) = -1 - 1 = -2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. If $\mathbf{F}$ is conservative, then $\int_C \mathbf{F} \cdot d\mathbf{r}$ is independent of path. This means that the work done along all piecewise-smooth curves that have the described initial and terminal points is the same. Your reply: It doesn't matter which curve is chosen.

28. The curves C_1 and C_2 connect the same two points but $\int_{C_1} \mathbf{F} \cdot d\mathbf{r} \neq \int_{C_2} \mathbf{F} \cdot d\mathbf{r}$. Thus $\mathbf{F}$ is not independent of path, and therefore is not conservative.

29. $\mathbf{F}(x, y) = x^3\,\mathbf{i} + y^3\,\mathbf{j}$, $W = \int_C \mathbf{F} \cdot d\mathbf{r}$. Since $\partial(x^3)/\partial y = 0 = \partial(y^3)/\partial x$, there exists a function f such that $\nabla f = \mathbf{F}$. In fact, $f_x(x, y) = x^3 \quad \Rightarrow \quad f(x, y) = \frac{1}{4}x^4 + g(y) \quad \Rightarrow \quad f_y(x, y) = 0 + g'(y)$. But $f_y(x, y) = y^3$ so $g'(y) = y^3 \quad \Rightarrow \quad g(y) = \frac{1}{4}y^4 + K$. We can take $K = 0$, so $f(x, y) = \frac{1}{4}x^4 + \frac{1}{4}y^4$. Thus $W = \int_C \mathbf{F} \cdot d\mathbf{r} = f(2, 2) - f(1, 0) = (4 + 4) - \left(\frac{1}{4} + 0\right) = \frac{31}{4}$.

30. $\mathbf{F}(x, y) = (2x + y)\,\mathbf{i} + x\,\mathbf{j}$, $W = \int_C \mathbf{F} \cdot d\mathbf{r}$. Since $\partial(2x + y)/\partial y = 1 = \partial(x)/\partial x$, there exists a function f such that $\nabla f = \mathbf{F}$. In fact, $f_x(x, y) = 2x + y \quad \Rightarrow \quad f(x, y) = x^2 + xy + g(y) \quad \Rightarrow \quad f_y(x, y) = x + g'(y)$. But $f_y(x, y) = x$ so $g'(y) = 0 \quad \Rightarrow \quad g(y) = K$. We can take $K = 0$, so $f(x, y) = x^2 + xy$. Thus $W = \int_C \mathbf{F} \cdot d\mathbf{r} = f(4, 3) - f(1, 1) = (16 + 12) - (1 + 1) = 26$.

31. We know that if the vector field (call it $\mathbf{F}$) is conservative, then around any closed path C, $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$. But take C to be a circle centered at the origin, oriented counterclockwise. All of the field vectors that start on C are roughly in the direction of motion along C, so the integral around C will be positive. Therefore the field is not conservative.

32. We know that if the vector field (call it $\mathbf{F}$) is conservative, then around any closed path C, $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$. For any closed path we draw in the field, it appears that some vectors on the curve point in approximately the same direction as the curve and a similar number point in roughly the opposite direction. (Some appear perpendicular to the curve as well.) Therefore it is plausible that $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$ for every closed curve C which means $\mathbf{F}$ is conservative.

33.

From the graph, it appears that $\mathbf{F}$ is conservative, since around all closed paths, the number and size of the field vectors pointing in directions similar to that of the path seem to be roughly the same as the number and size of the vectors pointing in the opposite direction. To check, we calculate $\dfrac{\partial}{\partial y}(\sin y) = \cos y = \dfrac{\partial}{\partial x}(1 + x\cos y)$. Thus, $\mathbf{F}$ is conservative, by Theorem 6.

34. $f(x, y) = \sin(x - 2y) \quad \Rightarrow \quad \mathbf{F} = \nabla f(x, y) = \cos(x - 2y)\,\mathbf{i} - 2\cos(x - 2y)\,\mathbf{j}$

(a) We use Theorem 2: $\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_{C_1} \nabla f \cdot d\mathbf{r} = f(\mathbf{r}(b)) - f(\mathbf{r}(a))$ where C_1 starts at $t = a$ and ends at $t = b$. So because $f(0, 0) = \sin 0 = 0$ and $f(\pi, \pi) = \sin(\pi - 2\pi) = 0$, one possible curve C_1 is the straight line from $(0, 0)$ to (π, π); that is, $\mathbf{r}(t) = \pi t\,\mathbf{i} + \pi t\,\mathbf{j}$, $0 \le t \le 1$.

(b) From (a), $\int_{C_2} \mathbf{F} \cdot d\mathbf{r} = f(\mathbf{r}(b)) - f(\mathbf{r}(a))$. So because $f(0, 0) = \sin 0 = 0$ and $f\left(\frac{\pi}{2}, 0\right) = 1$, one possible curve C_2 is $\mathbf{r}(t) = \frac{\pi}{2}t\,\mathbf{i}$, $0 \le t \le 1$, the straight line from $(0, 0)$ to $\left(\frac{\pi}{2}, 0\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. Since $\mathbf{F}$ is conservative, there exists a function f such that $\mathbf{F} = \nabla f$, that is, $P = f_x$, $Q = f_y$, and $R = f_z$. Since P, Q, and R have continuous first-order partial derivatives, Clairaut's Theorem says that $\partial P/\partial y = f_{xy} = f_{yx} = \partial Q/\partial x$, $\partial P/\partial z = f_{xz} = f_{zx} = \partial R/\partial x$, and $\partial Q/\partial z = f_{yz} = f_{zy} = \partial R/\partial y$.

36. Here $\mathbf{F}(x, y, z) = y\,\mathbf{i} + x\,\mathbf{j} + xyz\,\mathbf{k}$. Then using the notation of Exercise 35, $\partial P/\partial z = 0$ while $\partial R/\partial x = yz$. Since these aren't equal, $\mathbf{F}$ is not conservative. Thus by Theorem 4, the line integral of $\mathbf{F}$ is not independent of path.

37. $D = \{(x, y) \mid 0 < y < 3\}$ consists of those points between, but not on, the horizontal lines $y = 0$ and $y = 3$.

(a) Since D does not include any of its boundary points, it is open. More formally, at any point in D there is a disk centered at that point that lies entirely in D.

(b) Any two points chosen in D can always be joined by a path that lies entirely in D, so D is connected. (D consists of just one "piece.")

(c) D is connected and it has no holes, so it's simply-connected. (Every simple closed curve in D encloses only points that are in D.)

38. $D = \{(x, y) \mid 1 < |x| < 2\}$ consists of those points between, but not on, the vertical lines $x = 1$ and $x = 2$, together with the points between the vertical lines $x = -1$ and $x = -2$.

(a) The region does not include any of its boundary points, so it is open.

(b) D consists of two separate pieces, so it is not connected. [For instance, both the points $(-1.5, 0)$ and $(1.5, 0)$ lie in D but they cannot be joined by a path that lies entirely in D.]

(c) Because D is not connected, it's not simply-connected.

39. $D = \{(x, y) \mid 1 \le x^2 + y^2 \le 4,\ y \ge 0\}$ is the semiannular region in the upper half-plane between circles centered at the origin of radii 1 and 2 (including all boundary points).

(a) D includes boundary points, so it is not open. [Note that at any boundary point, $(1, 0)$ for instance, any disk centered there cannot lie entirely in D.]

(b) The region consists of one piece, so it's connected.

(c) D is connected and has no holes, so it's simply-connected.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $D = \{(x, y) \mid (x, y) \neq (2, 3)\}$ consists of all points in the xy-plane except for $(2, 3)$.

(a) D has only one boundary point, namely $(2, 3)$, which is not included, so the region is open.

(b) D is connected, as it consists of only one piece.

(c) D is not simply-connected, as it has a hole at $(2, 3)$. Thus any simple closed curve that encloses $(2, 3)$ lies in D but includes a point that is not in D.

41. (a) $P = -\dfrac{y}{x^2 + y^2}$, $\dfrac{\partial P}{\partial y} = \dfrac{y^2 - x^2}{(x^2 + y^2)^2}$ and $Q = \dfrac{x}{x^2 + y^2}$, $\dfrac{\partial Q}{\partial x} = \dfrac{y^2 - x^2}{(x^2 + y^2)^2}$. Thus $\dfrac{\partial P}{\partial y} = \dfrac{\partial Q}{\partial x}$.

(b) C_1: $x = \cos t$, $y = \sin t$, $0 \le t \le \pi$, C_2: $x = \cos t$, $y = \sin t$, $t = 2\pi$ to $t = \pi$. Then

$\displaystyle\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_0^{\pi} \frac{(-\sin t)(-\sin t) + (\cos t)(\cos t)}{\cos^2 t + \sin^2 t}\, dt = \int_0^{\pi} dt = \pi$ and $\displaystyle\int_{C_2} \mathbf{F} \cdot d\mathbf{r} = \int_{2\pi}^{\pi} dt = -\pi$ Since these aren't equal, the line integral of $\mathbf{F}$ isn't independent of path. (Or notice that $\int_{C_3} \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} dt = 2\pi$ where C_3 is the circle $x^2 + y^2 = 1$, and apply the contrapositive of Theorem 3.) This doesn't contradict Theorem 6, since the domain of $\mathbf{F}$, which is $\mathbb{R}^2$ except the origin, isn't simply-connected.

42. (a) Here $\mathbf{F}(\mathbf{r}) = c\mathbf{r}/|\mathbf{r}|^3$ and $\mathbf{r} = x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$. Then $f(\mathbf{r}) = -c/|\mathbf{r}|$ is a potential function for $\mathbf{F}$, that is, $\nabla f = \mathbf{F}$. (See the discussion of gradient fields in Section 16.1.) Hence $\mathbf{F}$ is conservative and its line integral is independent of path. Let $P_1 = (x_1, y_1, z_1)$ and $P_2 = (x_2, y_2, z_2)$.

$$W = \int_C \mathbf{F} \cdot d\mathbf{r} = f(P_2) - f(P_1) = -\frac{c}{(x_2^2 + y_2^2 + z_2^2)^{1/2}} + \frac{c}{(x_1^2 + y_1^2 + z_1^2)^{1/2}} = c\left(\frac{1}{d_1} - \frac{1}{d_2}\right).$$

(b) In this case, $c = -(mMG) \quad \Rightarrow$

$$\begin{aligned} W &= -mMG\left(\frac{1}{1.52 \times 10^{11}} - \frac{1}{1.47 \times 10^{11}}\right) \\ &= -(5.97 \times 10^{24})(1.99 \times 10^{30})(6.67 \times 10^{-11})(-2.2377 \times 10^{-13}) \\ &\approx 1.77 \times 10^{32}\ \text{J} \end{aligned}$$

(c) In this case, $c = \epsilon q Q \quad \Rightarrow$

$$W = \epsilon q Q\left(\frac{1}{10^{-12}} - \frac{1}{5 \times 10^{-13}}\right) = \left(8.985 \times 10^9\right)(1)\left(-1.6 \times 10^{-19}\right)\left(-10^{12}\right) \approx 1400\ \text{J}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16.4 Green's Theorem

1. (a)

C_1: $x = t \;\Rightarrow\; dx = dt,\; y = 0 \;\Rightarrow\; dy = 0\,dt,\; 0 \le t \le 5.$

C_2: $x = 5 \;\Rightarrow\; dx = 0\,dt,\; y = t \;\Rightarrow\; dy = dt,\; 0 \le t \le 4.$

C_3: $x = 5 - t \;\Rightarrow\; dx = -dt,\; y = 4 \;\Rightarrow\; dy = 0\,dt,\; 0 \le t \le 5.$

C_4: $x = 0 \;\Rightarrow\; dx = 0\,dt,\; y = 4 - t \;\Rightarrow\; dy = -dt,\; 0 \le t \le 4$

Thus $\oint_C y^2\,dx + x^2y\,dy = \oint\limits_{C_1+C_2+C_3+C_4} y^2\,dx + x^2y\,dy = \int_0^5 0\,dt + \int_0^4 25t\,dt + \int_0^5(-16+0)\,dt + \int_0^4 0\,dt$

$$= 0 + \left[\tfrac{25}{2}t^2\right]_0^4 + [-16t]_0^5 + 0 = 200 + (-80) = 120$$

(b) Note that C as given in part (a) is a positively oriented, piecewise-smooth, simple closed curve. Then by Green's Theorem,

$$\oint_C y^2\,dx + x^2y\,dy = \iint_D \left[\tfrac{\partial}{\partial x}(x^2y) - \tfrac{\partial}{\partial y}(y^2)\right] dA = \int_0^5\int_0^4 (2xy - 2y)\,dy\,dx = \int_0^5 \left[xy^2 - y^2\right]_{y=0}^{y=4} dx$$

$$= \int_0^5 (16x - 16)\,dx = \left[8x^2 - 16x\right]_0^5 = 200 - 80 = 120$$

2. (a) Parametric equations for C are $x = 4\cos t,\;\; y = 4\sin t,\;\; 0 \le t \le 2\pi$. Then $dx = -4\sin t\,dt,\; dy = 4\cos t\,dt$ and

$$\oint_C y\,dx - x\,dy = \int_0^{2\pi}[(4\sin t)(-4\sin t) - (4\cos t)(4\cos t)]\,dt$$

$$= -16\int_0^{2\pi}(\sin^2 t + \cos^2 t)\,dt = -16\int_0^{2\pi} 1\,dt = -16(2\pi) = -32\pi$$

(b) Note that C as given in part (a) is a positively oriented, smooth, simple closed curve. Then by Green's Theorem,

$$\oint_C y\,dx - x\,dy = \iint_D \left[\tfrac{\partial}{\partial x}(-x) - \tfrac{\partial}{\partial y}(y)\right] dA = \iint_D (-1-1)\,dA = -2\iint_D dA$$

$$= -2(\text{area of } D) = -2\cdot\pi(4)^2 = -32\pi$$

3. (a)

C_1: $x = t \;\Rightarrow\; dx = dt,\; y = 0 \;\Rightarrow\; dy = 0\,dt,\; 0 \le t \le 1.$

C_2: $x = 1 \;\Rightarrow\; dx = 0\,dt,\; y = t \;\Rightarrow\; dy = dt,\; 0 \le t \le 2.$

C_3: $x = 1 - t \;\Rightarrow\; dx = -dt,\; y = 2 - 2t \;\Rightarrow\; dy = -2\,dt,\; 0 \le t \le 1.$

Thus $\oint_C xy\,dx + x^2y^3\,dy = \oint\limits_{C_1+C_2+C_3} xy\,dx + x^2y^3\,dy$

$$= \int_0^1 0\,dt + \int_0^2 t^3\,dt + \int_0^1 \left[-(1-t)(2-2t) - 2(1-t)^2(2-2t)^3\right] dt$$

$$= 0 + \left[\tfrac{1}{4}t^4\right]_0^2 + \int_0^1 \left[-2(1-t)^2 - 16(1-t)^5\right] dt$$

$$= 4 + \left[\tfrac{2}{3}(1-t)^3 + \tfrac{8}{3}(1-t)^6\right]_0^1 = 4 + 0 - \tfrac{10}{3} = \tfrac{2}{3}$$

(b) $\oint_C xy\,dx + x^2y^3\,dy = \iint_D \left[\tfrac{\partial}{\partial x}(x^2y^3) - \tfrac{\partial}{\partial y}(xy)\right] dA = \int_0^1\int_0^{2x}(2xy^3 - x)\,dy\,dx$

$$= \int_0^1 \left[\tfrac{1}{2}xy^4 - xy\right]_{y=0}^{y=2x} dx = \int_0^1 (8x^5 - 2x^2)\,dx = \tfrac{4}{3} - \tfrac{2}{3} = \tfrac{2}{3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a)

$$C_1\colon\ x = t \quad\Rightarrow\quad dx = dt,\ y = t^2 \quad\Rightarrow\quad dy = 2t\,dt,\ \ 0 \le t \le 1$$

$$C_2\colon\ x = 1 - t \quad\Rightarrow\quad dx = -dt,\ y = 1 \quad\Rightarrow\quad dy = 0\,dt,\ 0 \le t \le 1$$

$$C_3\colon\ x = 0 \quad\Rightarrow\quad dx = 0\,dt,\ y = 1 - t \quad\Rightarrow\quad dy = -dt,\ \ 0 \le t \le 1$$

Thus

$$\begin{aligned}
\oint_C x^2y^2\,dx + xy\,dy &= \oint_{C_1+C_2+C_3} x^2y^2\,dx + xy\,dy \\
&= \int_0^1 \left[t^2(t^2)^2\,dt + t(t^2)(2t\,dt)\right] + \int_0^1 \left[(1-t)^2(1)^2(-dt) + (1-t)(1)(0\,dt)\right] \\
&\qquad + \int_0^1 \left[(0)^2(1-t)^2(0\,dt) + (0)(1-t)(-dt)\right] \\
&= \int_0^1 \left(t^6 + 2t^4\right) dt + \int_0^1 \left(-1 + 2t - t^2\right) dt + \int_0^1 0\,dt \\
&= \left[\tfrac{1}{7}t^7 + \tfrac{2}{5}t^5\right]_0^1 + \left[-t + t^2 - \tfrac{1}{3}t^3\right]_0^1 + 0 = \left(\tfrac{1}{7} + \tfrac{2}{5}\right) + \left(-1 + 1 - \tfrac{1}{3}\right) = \tfrac{22}{105}
\end{aligned}$$

(b) $$\begin{aligned}
\oint_C x^2y^2\,dx + xy\,dy &= \iint_D \left[\tfrac{\partial}{\partial x}(xy) - \tfrac{\partial}{\partial y}(x^2y^2)\right] dA = \int_0^1 \int_{x^2}^1 (y - 2x^2y)\,dy\,dx \\
&= \int_0^1 \left[\tfrac{1}{2}y^2 - x^2y^2\right]_{y=x^2}^{y=1} dx = \int_0^1 \left(\tfrac{1}{2} - x^2 - \tfrac{1}{2}x^4 + x^6\right) dx \\
&= \left[\tfrac{1}{2}x - \tfrac{1}{3}x^3 - \tfrac{1}{10}x^5 + \tfrac{1}{7}x^7\right]_0^1 = \tfrac{1}{2} - \tfrac{1}{3} - \tfrac{1}{10} + \tfrac{1}{7} = \tfrac{22}{105}
\end{aligned}$$

5. The region D enclosed by C is $[0,3] \times [0,4]$, so

$$\begin{aligned}
\int_C ye^x\,dx + 2e^x\,dy &= \iint_D \left[\tfrac{\partial}{\partial x}(2e^x) - \tfrac{\partial}{\partial y}(ye^x)\right] dA = \int_0^3 \int_0^4 (2e^x - e^x)\,dy\,dx \\
&= \int_0^3 e^x\,dx \int_0^4 dy = \left[e^x\right]_0^3 \left[y\right]_0^4 = (e^3 - e^0)(4 - 0) = 4(e^3 - 1)
\end{aligned}$$

6.

The region D enclosed by C can be given by $\{(x,y) \mid 1 \le x \le 2, 1 \le y \le 4\}$, so

$$\begin{aligned}
\int_C \ln(xy)\,dx + \frac{y}{x}\,dy &= \iint_D \left[\frac{\partial}{\partial x}\left(\frac{y}{x}\right) - \frac{\partial}{\partial y}\ln(xy)\right] dA \\
&= \int_1^4 \int_1^2 \left(-\frac{y}{x^2} - \frac{1}{y}\right) dx\,dy = \int_1^4 \left[\frac{y}{x} - \frac{x}{y}\right]_{x=1}^{x=2} dy \\
&= \int_1^4 \left(\frac{y}{2} - \frac{2}{y} - y + \frac{1}{y}\right) dy = \int_1^4 \left(-\frac{y}{2} - \frac{1}{y}\right) dy \\
&= \left[-\frac{y^2}{4} - \ln y\right]_1^4 = -\frac{16}{4} - \ln 4 + \frac{1}{4} + 0 = -\frac{15}{4} - \ln 4
\end{aligned}$$

7.

The region D enclosed by C can be given by $\{(x,y) \mid 0 \le y \le 3x, 0 \le x \le 1\}$, so

$$\begin{aligned}
\int_C x^2y^2\,dx + y\tan^{-1} y\,dy &= \iint_D \left[\frac{\partial}{\partial x}\left(y\tan^{-1} y\right) - \frac{\partial}{\partial y}\left(x^2y^2\right)\right] dA \\
&= \int_0^1 \int_0^{3x} (-2x^2y)\,dy\,dx = -\int_0^1 \left[x^2y^2\right]_{y=0}^{y=3x} dx \\
&= -9\int_0^1 x^4\,dx = -\tfrac{9}{5}\left[x^5\right]_0^1 = -\tfrac{9}{5}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8.

The region D enclosed by C is given by $\{(x,y) \mid 0 \le y \le 1,\ 0 \le x \le 2y\}$, so

$$\begin{aligned}\int_C (x^2+y^2)\,dx + (x^2-y^2)\,dy &= \iint_D \left[\tfrac{\partial}{\partial x}(x^2-y^2) - \tfrac{\partial}{\partial y}(x^2+y^2)\right] dA \\ &= \int_0^1 \int_0^{2y} (2x-2y)\,dx\,dy \\ &= \int_0^1 \left[x^2-2xy\right]_{x=0}^{x=2y} dy \\ &= \int_0^1 (4y^2-4y^2)\,dy = \int_0^1 0\,dy = 0\end{aligned}$$

9. $$\begin{aligned}\int_C \left(y+e^{\sqrt{x}}\right) dx + (2x+\cos y^2)\,dy &= \iint_D \left[\tfrac{\partial}{\partial x}(2x+\cos y^2) - \tfrac{\partial}{\partial y}\left(y+e^{\sqrt{x}}\right)\right] dA \\ &= \int_0^1 \int_{x^2}^{\sqrt{x}} (2-1)\,dy\,dx = \int_0^1 (\sqrt{x}-x^2)\,dx = \left[\tfrac{2}{3}x^{3/2} - \tfrac{1}{3}x^3\right]_0^1 = \tfrac{1}{3}\end{aligned}$$

10. $$\begin{aligned}\int_C y^4\,dx + 2xy^3\,dy &= \iint_D \left[\tfrac{\partial}{\partial x}(2xy^3) - \tfrac{\partial}{\partial y}(y^4)\right] dA = \iint_D (2y^3-4y^3)\,dA \\ &= -2\iint_D y^3\,dA = 0\end{aligned}$$

because $f(x,y) = y^3$ is an odd function with respect to y and D is symmetric about the x-axis.

11. $$\begin{aligned}\int_C y^3\,dx - x^3\,dy &= \iint_D \left[\tfrac{\partial}{\partial x}(-x^3) - \tfrac{\partial}{\partial y}(y^3)\right] dA = \iint_D (-3x^2-3y^2)\,dA = \int_0^{2\pi}\int_0^2 (-3r^2)\,r\,dr\,d\theta \\ &= -3\int_0^{2\pi} d\theta \int_0^2 r^3\,dr = -3\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^2 = -3(2\pi)(4) = -24\pi\end{aligned}$$

12. $$\begin{aligned}\int_C (1-y^3)\,dx + (x^3+e^{y^2})\,dy &= \iint_D \left[\tfrac{\partial}{\partial x}(x^3+e^{y^2}) - \tfrac{\partial}{\partial y}(1-y^3)\right] dA = \iint_D (3x^2+3y^2)\,dA \\ &= \int_0^{2\pi}\int_2^3 (3r^2)\,r\,dr\,d\theta = 3\int_0^{2\pi} d\theta \int_2^3 r^3\,dr \\ &= 3\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_2^3 = 3(2\pi)\cdot\tfrac{1}{4}(81-16) = \tfrac{195}{2}\pi\end{aligned}$$

13. The region D enclosed by C is given by $\{(r,\theta) \mid 1 \le r \le 2, 0 \le \theta \le \pi/2\}$ (in polar coordinates),which is traversed counterclockwise, so C has positive orientation. Thus,

$$\begin{aligned}\int_C \left(3+e^{x^2}\right) dx + \left(\tan^{-1} y + 3x^2\right) dy &= \iint_D \left[\tfrac{\partial}{\partial x}(\tan^{-1} y + 3x^2) - \tfrac{\partial}{\partial y}(3+e^{x^2})\right] dA \\ &= \iint_D 6x\,dA = 6\int_0^{\pi/2}\int_1^2 r\cos\theta\,r\,dr\,d\theta \qquad \text{[Switching to polar coordinates]} \\ &= 6\int_0^{\pi/2}\cos\theta\,d\theta \int_1^2 r^2\,dr = 6\left[\sin\theta\right]_{\theta=0}^{\theta=\pi/2}\left[\tfrac{r^3}{3}\right]_{r=1}^{r=2} \\ &= 6(1-0)\left(\tfrac{8}{3}-\tfrac{1}{3}\right) = 14\end{aligned}$$

14. The region D enclosed by C is given by $\{(x,y) \mid y^2 \le x \le 4, 0 \le y \le 2\}$. C is traversed clockwise, so $-C$ gives the positive orientation. Then

$$\begin{aligned}-\int_{-C} (x^{2/3}+y^2)\,dx + (y^{4/3}-x^2)\,dy &= -\iint_D \left[\tfrac{\partial}{\partial x}(y^{4/3}-x^2) - \tfrac{\partial}{\partial y}(x^{2/3}+y^2)\right] dA \\ &= -\int_0^2\int_{y^2}^4 (-2x-2y)\,dx\,dy = \int_0^2\int_{y^2}^4 (2x+2y)\,dx\,dy \\ &= \int_0^2 \left[x^2+2xy\right]_{x=y^2}^{x=4} dy = \int_0^2 (16+8y-y^4-2y^3)\,dy \\ &= \left[16y+4y^2-\tfrac{y^5}{5}-\tfrac{y^4}{2}\right]_0^2 = 16(2)+4(2^2)-\tfrac{2^5}{5}-\tfrac{2^4}{2}-0 = \tfrac{168}{5}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $\mathbf{F}(x,y) = \langle y\cos x - xy\sin x, xy + x\cos x\rangle$ and the region D enclosed by C is given by $\{(x,y) \mid 0 \le x \le 2,\ 0 \le y \le 4-2x\}$. C is traversed clockwise, so $-C$ gives the positive orientation.

$$\begin{aligned}
\textstyle\int_C \mathbf{F}\cdot d\mathbf{r} &= -\textstyle\int_{-C}(y\cos x - xy\sin x)\,dx + (xy + x\cos x)\,dy \\
&= -\iint_D \left[\tfrac{\partial}{\partial x}(xy + x\cos x) - \tfrac{\partial}{\partial y}(y\cos x - xy\sin x)\right] dA \\
&= -\iint_D (y - x\sin x + \cos x - \cos x + x\sin x)\,dA = -\int_0^2\int_0^{4-2x} y\,dy\,dx \\
&= -\int_0^2 \left[\tfrac{1}{2}y^2\right]_{y=0}^{y=4-2x} dx = -\int_0^2 \tfrac{1}{2}(4-2x)^2\,dx = -\int_0^2 (8 - 8x + 2x^2)\,dx \\
&= -\left[8x - 4x^2 + \tfrac{2}{3}x^3\right]_0^2 = -\left(16 - 16 + \tfrac{16}{3} - 0\right) = -\tfrac{16}{3}
\end{aligned}$$

16. $\mathbf{F}(x,y) = \left\langle e^{-x} + y^2, e^{-y} + x^2\right\rangle$ and the region D enclosed by C is given by $\{(x,y) \mid -\pi/2 \le x \le \pi/2, 0 \le y \le \cos x\}$. C is traversed clockwise, so $-C$ gives the positive orientation.

$$\begin{aligned}
\textstyle\int_C \mathbf{F}\cdot d\mathbf{r} &= -\textstyle\int_{-C}\left(e^{-x} + y^2\right)dx + \left(e^{-y} + x^2\right)dy = -\iint_D \left[\tfrac{\partial}{\partial x}\left(e^{-y} + x^2\right) - \tfrac{\partial}{\partial y}\left(e^{-x} + y^2\right)\right] dA \\
&= -\int_{-\pi/2}^{\pi/2}\int_0^{\cos x}(2x - 2y)\,dy\,dx = -\int_{-\pi/2}^{\pi/2}\left[2xy - y^2\right]_{y=0}^{y=\cos x} dx \\
&= -\int_{-\pi/2}^{\pi/2}(2x\cos x - \cos^2 x)\,dx = -\int_{-\pi/2}^{\pi/2}\left[2x\cos x - \tfrac{1}{2}(1 + \cos 2x)\right] dx \\
&= -\left[2x\sin x + 2\cos x - \tfrac{1}{2}\left(x + \tfrac{1}{2}\sin 2x\right)\right]_{-\pi/2}^{\pi/2} \qquad \text{[integrate by parts in the first term]} \\
&= -\left(\pi - \tfrac{1}{4}\pi - \pi - \tfrac{1}{4}\pi\right) = \tfrac{1}{2}\pi
\end{aligned}$$

17. $\mathbf{F}(x,y) = \langle y - \cos y, x\sin y\rangle$ and the region D enclosed by C is the disk with radius 2 centered at $(3,-4)$. C is traversed clockwise, so $-C$ gives the positive orientation.

$$\begin{aligned}
\textstyle\int_C \mathbf{F}\cdot d\mathbf{r} &= -\textstyle\int_{-C}(y - \cos y)\,dx + (x\sin y)\,dy = -\iint_D \left[\tfrac{\partial}{\partial x}(x\sin y) - \tfrac{\partial}{\partial y}(y - \cos y)\right] dA \\
&= -\iint_D (\sin y - 1 - \sin y)\,dA = \iint_D dA = \text{area of } D = \pi(2)^2 = 4\pi
\end{aligned}$$

18. $\mathbf{F}(x,y) = \left\langle \sqrt{x^2+1}, \tan^{-1} x\right\rangle$ and the region D enclosed by C is given by $\{(x,y) \mid 0 \le x \le 1, x \le y \le 1\}$. C is oriented positively, so

$$\begin{aligned}
\textstyle\int_C \mathbf{F}\cdot d\mathbf{r} &= \textstyle\int_C \sqrt{x^2+1}\,dx + \tan^{-1} x\,dy = \displaystyle\iint_D \left[\frac{\partial}{\partial x}\left(\tan^{-1} x\right) - \frac{\partial}{\partial y}\left(\sqrt{x^2+1}\right)\right] dA \\
&= \int_0^1\int_x^1 \left(\frac{1}{1+x^2} - 0\right) dy\,dx = \int_0^1 \frac{1}{1+x^2}\left[y\right]_{y=x}^{y=1} dx = \int_0^1 \frac{1}{1+x^2}(1-x)\,dx \\
&= \int_0^1 \left(\frac{1}{1+x^2} - \frac{x}{1+x^2}\right) dx = \left[\tan^{-1} x - \frac{1}{2}\ln(1+x^2)\right]_0^1 = \frac{\pi}{4} - \frac{1}{2}\ln 2
\end{aligned}$$

19. Here $C = C_1 + C_2$ where

C_1 can be parametrized as $x = t,\ \ y = 0,\ \ -\pi/2 \le t \le \pi/2$, and

C_2 is given by $x = -t,\ \ y = \cos t,\ \ -\pi/2 \le t \le \pi/2$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Then the line integral is

$$\oint_{C_1+C_2} x^3y^4\,dx + x^5y^4\,dy = \int_{-\pi/2}^{\pi/2}(0+0)\,dt + \int_{-\pi/2}^{\pi/2}[(-t)^3(\cos t)^4(-1) + (-t)^5(\cos t)^4(-\sin t)]\,dt$$
$$= 0 + \int_{-\pi/2}^{\pi/2}(t^3\cos^4 t + t^5\cos^4 t\sin t)\,dt = \tfrac{1}{15}\pi^4 - \tfrac{4144}{1125}\pi^2 + \tfrac{7{,}578{,}368}{253{,}125} \approx 0.0779$$

according to a CAS. The double integral is

$$\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) dA = \int_{-\pi/2}^{\pi/2}\int_0^{\cos x}(5x^4y^4 - 4x^3y^3)\,dy\,dx = \tfrac{1}{15}\pi^4 - \tfrac{4144}{1125}\pi^2 + \tfrac{7{,}578{,}368}{253{,}125} \approx 0.0779,$$

verifying Green's Theorem in this case.

20. We can parametrize C as $x = \cos\theta,\ \ y = 2\sin\theta,\ \ 0 \le \theta \le 2\pi$. Then the line integral is

$$\oint_C P\,dx + Q\,dy = \int_0^{2\pi}\left[2\cos\theta - (\cos\theta)^3(2\sin\theta)^5\right](-\sin\theta)\,d\theta + \int_0^{2\pi}(\cos\theta)^3(2\sin\theta)^8\cdot 2\cos\theta\,d\theta$$
$$= \int_0^{2\pi}(-2\cos\theta\sin\theta + 32\cos^3\theta\sin^6\theta + 512\cos^4\theta\sin^8\theta)\,d\theta = 7\pi,$$

according to a CAS. The double integral is $\displaystyle\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) dA = \int_{-1}^{1}\int_{-\sqrt{4-4x^2}}^{\sqrt{4-4x^2}}(3x^2y^8 + 5x^3y^4)\,dy\,dx = 7\pi.$

21. By Green's Theorem, $W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_C x(x+y)\,dx + xy^2\,dy = \iint_D (y^2 - x)\,dA$ where C is the path described in the question and D is the triangle bounded by C. So

$$W = \int_0^1\int_0^{1-x}(y^2 - x)\,dy\,dx = \int_0^1\left[\tfrac{1}{3}y^3 - xy\right]_{y=0}^{y=1-x}dx = \int_0^1\left(\tfrac{1}{3}(1-x)^3 - x(1-x)\right)dx$$
$$= \left[-\tfrac{1}{12}(1-x)^4 - \tfrac{1}{2}x^2 + \tfrac{1}{3}x^3\right]_0^1 = \left(-\tfrac{1}{2} + \tfrac{1}{3}\right) - \left(-\tfrac{1}{12}\right) = -\tfrac{1}{12}$$

22. By Green's Theorem, $W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_C \sin x\,dx + \left(\sin y + xy^2 + \frac{1}{3}x^3\right)dy = \iint_D (y^2 + x^2 - 0)\,dA$, where D is the region (a quarter-disk) bounded by C. Converting to polar coordinates, we have

$W = \int_0^{\pi/2}\int_0^5 r^2\cdot r\,dr\,d\theta = \left[\theta\right]_0^{\pi/2}\left[\frac{1}{4}r^4\right]_0^5 = \frac{1}{2}\pi\left(\frac{625}{4}\right) = \frac{625}{8}\pi.$

23. Let C_1 be the arch of the cycloid from $(0,0)$ to $(2\pi,0)$, which corresponds to $0 \le t \le 2\pi$, and let C_2 be the segment from $(2\pi,0)$ to $(0,0)$, so C_2 is given by $x = 2\pi - t$, $y = 0$, $0 \le t \le 2\pi$. Then $C = C_1 \cup C_2$ is traversed clockwise, so $-C$ is oriented positively. Thus $-C$ encloses the area under one arch of the cycloid and from (5) we have

$$A = -\oint_{-C} y\,dx = \int_{C_1} y\,dx + \int_{C_2} y\,dx = \int_0^{2\pi}(1-\cos t)(1-\cos t)\,dt + \int_0^{2\pi} 0\,(-dt)$$
$$= \int_0^{2\pi}(1 - 2\cos t + \cos^2 t)\,dt + 0 = \left[t - 2\sin t + \tfrac{1}{2}t + \tfrac{1}{4}\sin 2t\right]_0^{2\pi} = 3\pi$$

24.

$$A = \oint_C x\,dy = \int_0^{2\pi}(5\cos t - \cos 5t)(5\cos t - 5\cos 5t)\,dt$$
$$= \int_0^{2\pi}(25\cos^2 t - 30\cos t\cos 5t + 5\cos^2 5t)\,dt$$
$$= \left[25\left(\tfrac{1}{2}t + \tfrac{1}{4}\sin 2t\right) - 30\left(\tfrac{1}{8}\sin 4t + \tfrac{1}{12}\sin 6t\right) + 5\left(\tfrac{1}{2}t + \tfrac{1}{20}\sin 10t\right)\right]_0^{2\pi}$$

[Use Formula 80 in the Table of Integrals]

$$= 30\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. (a) Using Equation 16.2.8, we write parametric equations of the line segment as $x = (1-t)x_1 + tx_2$, $y = (1-t)y_1 + ty_2$, $0 \le t \le 1$. Then $dx = (x_2 - x_1)\,dt$ and $dy = (y_2 - y_1)\,dt$, so

$$\begin{aligned}\int_C x\,dy - y\,dx &= \int_0^1 [(1-t)x_1 + tx_2](y_2 - y_1)\,dt + [(1-t)y_1 + ty_2](x_2 - x_1)\,dt \\ &= \int_0^1 \left(x_1(y_2 - y_1) - y_1(x_2 - x_1) + t[(y_2 - y_1)(x_2 - x_1) - (x_2 - x_1)(y_2 - y_1)]\right) dt \\ &= \int_0^1 (x_1y_2 - x_2y_1)\,dt = x_1y_2 - x_2y_1\end{aligned}$$

(b) We apply Green's Theorem to the path $C = C_1 \cup C_2 \cup \cdots \cup C_n$, where C_i is the line segment that joins (x_i, y_i) to (x_{i+1}, y_{i+1}) for $i = 1, 2, \ldots, n-1$, and C_n is the line segment that joins (x_n, y_n) to (x_1, y_1). From (5), $\frac{1}{2}\int_C x\,dy - y\,dx = \iint_D dA$, where D is the polygon bounded by C. Therefore

$$\begin{aligned}\text{area of polygon} = A(D) &= \iint_D dA = \tfrac{1}{2}\int_C x\,dy - y\,dx \\ &= \tfrac{1}{2}\left(\int_{C_1} x\,dy - y\,dx + \int_{C_2} x\,dy - y\,dx + \cdots + \int_{C_{n-1}} x\,dy - y\,dx + \int_{C_n} x\,dy - y\,dx\right)\end{aligned}$$

To evaluate these integrals we use the formula from (a) to get

$A(D) = \frac{1}{2}[(x_1y_2 - x_2y_1) + (x_2y_3 - x_3y_2) + \cdots + (x_{n-1}y_n - x_ny_{n-1}) + (x_ny_1 - x_1y_n)]$.

(c) $A = \frac{1}{2}[(0\cdot 1 - 2\cdot 0) + (2\cdot 3 - 1\cdot 1) + (1\cdot 2 - 0\cdot 3) + (0\cdot 1 - (-1)\cdot 2) + (-1\cdot 0 - 0\cdot 1)]$
$= \frac{1}{2}(0 + 5 + 2 + 2) = \frac{9}{2}$

26. By Green's Theorem, $\frac{1}{2A}\oint_C x^2\,dy = \frac{1}{2A}\iint_D 2x\,dA = \frac{1}{A}\iint_D x\,dA = \overline{x}$ and $-\frac{1}{2A}\oint_C y^2\,dx = -\frac{1}{2A}\iint_D(-2y)\,dA = \frac{1}{A}\iint_D y\,dA = \overline{y}$.

27. We orient the quarter-circular region as shown in the figure.

$A = \frac{1}{4}\pi a^2$ so $\overline{x} = \dfrac{1}{\pi a^2/2}\displaystyle\oint_C x^2\,dy$ and $\overline{y} = -\dfrac{1}{\pi a^2/2}\displaystyle\oint_C y^2\,dx$.

Here $C = C_1 + C_2 + C_3$ where

C_1: $x = t,\ \ y = 0,\ \ 0 \le t \le a$;

C_2: $x = a\cos t,\ \ y = a\sin t,\ \ 0 \le t \le \frac{\pi}{2}$; and

C_3: $x = 0, y = a - t, 0 \le t \le a$. Then

$$\begin{aligned}\oint_C x^2\,dy &= \int_{C_1} x^2\,dy + \int_{C_2} x^2\,dy + \int_{C_3} x^2\,dy = \int_0^a 0\,dt + \int_0^{\pi/2}(a\cos t)^2(a\cos t)\,dt + \int_0^a 0\,dt \\ &= \int_0^{\pi/2} a^3\cos^3 t\,dt = a^3\int_0^{\pi/2}(1 - \sin^2 t)\cos t\,dt = a^3\left[\sin t - \tfrac{1}{3}\sin^3 t\right]_0^{\pi/2} = \tfrac{2}{3}a^3\end{aligned}$$

so $\overline{x} = \dfrac{1}{\pi a^2/2}\displaystyle\oint_C x^2\,dy = \dfrac{4a}{3\pi}$.

$$\begin{aligned}\oint_C y^2\,dx &= \int_{C_1} y^2\,dx + \int_{C_2} y^2\,dx + \int_{C_3} y^2\,dx = \int_0^a 0\,dt + \int_0^{\pi/2}(a\sin t)^2(-a\sin t)\,dt + \int_0^a 0\,dt \\ &= \int_0^{\pi/2}(-a^3\sin^3 t)\,dt = -a^3\int_0^{\pi/2}(1 - \cos^2 t)\sin t\,dt = -a^3\left[\tfrac{1}{3}\cos^3 t - \cos t\right]_0^{\pi/2} = -\tfrac{2}{3}a^3,\end{aligned}$$

so $\overline{y} = -\dfrac{1}{\pi a^2/2}\displaystyle\oint_C y^2\,dx = \dfrac{4a}{3\pi}$. Thus $(\overline{x}, \overline{y}) = \left(\dfrac{4a}{3\pi}, \dfrac{4a}{3\pi}\right)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. Here $A = \frac{1}{2}ab$ and $C = C_1 + C_2 + C_3$, where C_1: $x = x$, $y = 0$, $0 \le x \le a$;

C_2: $x = a$, $y = y$, $0 \le y \le b$; and C_3: $x = x$, $y = \frac{b}{a}x$, $x = a$ to $x = 0$. Then

$$\oint_C x^2\,dy = \int_{C_1} x^2\,dy + \int_{C_2} x^2\,dy + \int_{C_3} x^2\,dy = 0 + \int_0^b a^2\,dy + \int_a^0 (x^2)\left(\tfrac{b}{a}\,dx\right)$$
$$= a^2b + \tfrac{b}{a}\left[\tfrac{1}{3}x^3\right]_a^0 = a^2b - \tfrac{1}{3}a^2b = \tfrac{2}{3}a^2b.$$

Similarly, $\oint_C y^2\,dx = \int_{C_1} y^2\,dx + \int_{C_2} y^2\,dx + \int_{C_3} y^2\,dx = 0 + 0 + \int_a^0 \left(\frac{b}{a}x\right)^2\,dx = \frac{b^2}{a^2}\cdot\frac{1}{3}x^3\Big]_a^0 = -\frac{1}{3}ab^2$. Thus

$\overline{x} = \frac{1}{2A}\oint_C x^2\,dy = \frac{1}{ab}\cdot\frac{2}{3}a^2b = \frac{2}{3}a$ and $\overline{y} = -\frac{1}{2A}\oint_C y^2\,dx = -\frac{1}{ab}\left(-\frac{1}{3}ab^2\right) = \frac{1}{3}b$, so $(\overline{x}, \overline{y}) = \left(\frac{2}{3}a, \frac{1}{3}b\right)$.

29. By Green's Theorem, $-\frac{1}{3}\rho\oint_C y^3\,dx = -\frac{1}{3}\rho\iint_D(-3y^2)\,dA = \iint_D y^2\rho\,dA = I_x$ and

$\frac{1}{3}\rho\oint_C x^3\,dy = \frac{1}{3}\rho\iint_D(3x^2)\,dA = \iint_D x^2\rho\,dA = I_y$.

30. By symmetry the moments of inertia about any two diameters are equal. Centering the disk at the origin, the moment of inertia about a diameter equals

$$I_y = \tfrac{1}{3}\rho\oint_C x^3\,dy = \tfrac{1}{3}\rho\int_0^{2\pi}(a\cos t)^3(a\cos t\,dt) = \tfrac{1}{3}\rho\int_0^{2\pi}(a^4\cos^4 t)\,dt$$
$$= \tfrac{1}{3}a^4\rho\int_0^{2\pi}\left[\tfrac{1}{2}(1+\cos 2t)\right]^2 dt = \tfrac{1}{3}a^4\rho\int_0^{2\pi}\left(\tfrac{3}{8} + \tfrac{1}{2}\cos 2t + \tfrac{1}{8}\cos 4t\right)dt$$
$$= \tfrac{1}{3}a^4\rho\left[\tfrac{3}{8}t + \tfrac{1}{4}\sin 2t + \tfrac{1}{32}\sin 4t\right]_0^{2\pi} = \tfrac{1}{3}a^4\rho\cdot\tfrac{3(2\pi)}{8} = \tfrac{1}{4}\pi a^4\rho$$

31. As in Example 5, let C' be a counterclockwise-oriented circle with center the origin and radius a, where a is chosen to be small enough so that C' lies inside C, and D the region bounded by C and C'. Here

$$P = \frac{2xy}{(x^2+y^2)^2} \quad\Rightarrow\quad \frac{\partial P}{\partial y} = \frac{2x(x^2+y^2)^2 - 2xy\cdot 2(x^2+y^2)\cdot 2y}{(x^2+y^2)^4} = \frac{2x^3-6xy^2}{(x^2+y^2)^3} \text{ and}$$

$$Q = \frac{y^2-x^2}{(x^2+y^2)^2} \quad\Rightarrow\quad \frac{\partial Q}{\partial x} = \frac{-2x(x^2+y^2)^2 - (y^2-x^2)\cdot 2(x^2+y^2)\cdot 2x}{(x^2+y^2)^4} = \frac{2x^3-6xy^2}{(x^2+y^2)^3}.$$ Thus, as in the example,

$$\int_C P\,dx + Q\,dy + \int_{-C'} P\,dx + Q\,dy = \iint_D\left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right)dA = \iint_D 0\,dA = 0$$

and $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_{C'}\mathbf{F}\cdot d\mathbf{r}$. We parametrize C' as $\mathbf{r}(t) = a\cos t\,\mathbf{i} + a\sin t\,\mathbf{j}$, $0 \le t \le 2\pi$. Then

$$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_{C'}\mathbf{F}\cdot d\mathbf{r} = \int_0^{2\pi}\frac{2\,(a\cos t)\,(a\sin t)\,\mathbf{i} + \left(a^2\sin^2 t - a^2\cos^2 t\right)\mathbf{j}}{\left(a^2\cos^2 t + a^2\sin^2 t\right)^2}\cdot\Big(-a\sin t\,\mathbf{i} + a\cos t\,\mathbf{j}\Big)\,dt$$
$$= \frac{1}{a}\int_0^{2\pi}\left(-\cos t\sin^2 t - \cos^3 t\right)dt = \frac{1}{a}\int_0^{2\pi}\left(-\cos t\sin^2 t - \cos t\left(1-\sin^2 t\right)\right)dt$$
$$= -\frac{1}{a}\int_0^{2\pi}\cos t\,dt = -\frac{1}{a}\sin t\Big]_0^{2\pi} = 0$$

32. P and Q have continuous partial derivatives on $\mathbb{R}^2$, so by Green's Theorem we have

$$\int_C \mathbf{F}\cdot d\mathbf{r} = \iint_D\left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right)dA = \iint_D\left[\frac{\partial}{\partial x}(3x - y^2) - \frac{\partial}{\partial y}(x^2+y)\right]dA$$
$$= \iint_D(3-1)\,dA = 2\iint_D dA = 2\cdot A(D) = 2\cdot 6 = 12$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

33. Since C is a simple closed path which doesn't pass through or enclose the origin, there exists an open region that doesn't contain the origin but does contain D. Thus $P = -y/(x^2+y^2)$ and $Q = x/(x^2+y^2)$ have continuous partial derivatives on this open region containing D and we can apply Green's Theorem. But by Exercise 16.3.41(a), $\partial P/\partial y = \partial Q/\partial x$, so $\oint_C \mathbf{F}\cdot d\mathbf{r} = \iint_D 0\,dA = 0$.

34. We express D as a type II region: $D = \{(x,y) \mid f_1(y) \le x \le f_2(y),\ c \le y \le d\}$ where f_1 and f_2 are continuous functions. Then $\displaystyle\iint_D \frac{\partial Q}{\partial x}\,dA = \int_c^d \int_{f_1(y)}^{f_2(y)} \frac{\partial Q}{\partial x}\,dx\,dy = \int_c^d [Q(f_2(y),y) - Q(f_1(y),y)]\,dy$ by the Fundamental Theorem of Calculus. But referring to the figure, $\displaystyle\oint_C Q\,dy = \oint_{C_1+C_2+C_3+C_4} Q\,dy$.

Then $\int_{C_1} Q\,dy = \int_d^c Q(f_1(y),y)\,dy$, $\int_{C_2} Q\,dy = \int_{C_4} Q\,dy = 0$, and $\int_{C_3} Q\,dy = \int_c^d Q(f_2(y),y)\,dy$. Hence

$\oint_C Q\,dy = \int_c^d [Q(f_2(y),y) - Q(f_1(y),y)]\,dy = \iint_D (\partial Q/\partial x)\,dA$.

35. Using the first part of Equation 5, we have that $\iint_R dx\,dy = A(R) = \int_{\partial R} x\,dy$. But $x = g(u,v)$, and $dy = \dfrac{\partial h}{\partial u}\,du + \dfrac{\partial h}{\partial v}\,dv$, and we orient ∂S by taking the positive direction to be that which corresponds, under the mapping, to the positive direction along ∂R, so

$$\begin{aligned}
\int_{\partial R} x\,dy &= \int_{\partial S} g(u,v)\left(\frac{\partial h}{\partial u}\,du + \frac{\partial h}{\partial v}\,dv\right) = \int_{\partial S} g(u,v)\,\frac{\partial h}{\partial u}\,du + g(u,v)\,\frac{\partial h}{\partial v}\,dv \\
&= \pm\iint_S \left[\tfrac{\partial}{\partial u}\left(g(u,v)\,\tfrac{\partial h}{\partial v}\right) - \tfrac{\partial}{\partial v}\left(g(u,v)\,\tfrac{\partial h}{\partial u}\right)\right] dA \qquad \text{[using Green's Theorem in the } uv\text{-plane]} \\
&= \pm\iint_S \left(\tfrac{\partial g}{\partial u}\tfrac{\partial h}{\partial v} + g(u,v)\,\tfrac{\partial^2 h}{\partial u\,\partial v} - \tfrac{\partial g}{\partial v}\tfrac{\partial h}{\partial u} - g(u,v)\,\tfrac{\partial^2 h}{\partial v\,\partial u}\right) dA \qquad \text{[using the Chain Rule]} \\
&= \pm\iint_S \left(\tfrac{\partial x}{\partial u}\tfrac{\partial y}{\partial v} - \tfrac{\partial x}{\partial v}\tfrac{\partial y}{\partial u}\right) dA \quad \text{[by the equality of mixed partials]} \quad = \pm\iint_S \tfrac{\partial(x,y)}{\partial(u,v)}\,du\,dv
\end{aligned}$$

The sign is chosen to be positive if the orientation that we gave to ∂S corresponds to the usual positive orientation, and it is negative otherwise. In either case, since $A(R)$ is positive, the sign chosen must be the same as the sign of $\dfrac{\partial(x,y)}{\partial(u,v)}$.

Therefore, $A(R) = \displaystyle\iint_R dx\,dy = \iint_S \left|\frac{\partial(x,y)}{\partial(u,v)}\right| du\,dv$.

16.5 Curl and Divergence

1. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ xy^2z^2 & x^2yz^2 & x^2y^2z \end{vmatrix}$

$$\begin{aligned}
&= \left[\frac{\partial}{\partial y}(x^2y^2z) - \frac{\partial}{\partial z}(x^2yz^2)\right]\mathbf{i} - \left[\frac{\partial}{\partial x}(x^2y^2z) - \frac{\partial}{\partial z}(xy^2z^2)\right]\mathbf{j} + \left[\frac{\partial}{\partial x}(x^2yz^2) - \frac{\partial}{\partial y}(xy^2z^2)\right]\mathbf{k} \\
&= (2x^2yz - 2x^2yz)\,\mathbf{i} - (2xy^2z - 2xy^2z)\,\mathbf{j} + (2xyz^2 - 2xyz^2)\,\mathbf{k} = \mathbf{0}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}(xy^2z^2) + \dfrac{\partial}{\partial y}(x^2yz^2) + \dfrac{\partial}{\partial z}(x^2y^2z) = y^2z^2 + x^2z^2 + x^2y^2$

2. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ 0 & x^3yz^2 & y^4z^3 \end{vmatrix}$

$$= \left[\frac{\partial}{\partial y}(y^4z^3) - \frac{\partial}{\partial z}(x^3yz^2)\right]\mathbf{i} - \left[\frac{\partial}{\partial x}(y^4z^3) - \frac{\partial}{\partial z}(0)\right]\mathbf{j} + \left[\frac{\partial}{\partial x}(x^3yz^2) - \frac{\partial}{\partial y}(0)\right]\mathbf{k}$$

$$= (4y^3z^3 - 2x^3yz)\,\mathbf{i} - (0-0)\,\mathbf{j} + (3x^2yz^2 - 0)\,\mathbf{k}$$

$$= (4y^3z^3 - 2x^3yz)\,\mathbf{i} + 3x^2yz^2\,\mathbf{k}$$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}(0) + \dfrac{\partial}{\partial y}(x^3yz^2) + \dfrac{\partial}{\partial z}(y^4z^3) = 0 + x^3z^2 + 3y^4z^2 = x^3z^2 + 3y^4z^2$

3. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ xye^z & 0 & yze^x \end{vmatrix} = (ze^x - 0)\,\mathbf{i} - (yze^x - xye^z)\,\mathbf{j} + (0 - xe^z)\,\mathbf{k}$

$$= ze^x\,\mathbf{i} + (xye^z - yze^x)\,\mathbf{j} - xe^z\,\mathbf{k}$$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}(xye^z) + \dfrac{\partial}{\partial y}(0) + \dfrac{\partial}{\partial z}(yze^x) = ye^z + 0 + ye^x = y(e^z + e^x)$

4. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \sin yz & \sin zx & \sin xy \end{vmatrix}$

$$= (x\cos xy - x\cos zx)\,\mathbf{i} - (y\cos xy - y\cos yz)\,\mathbf{j} + (z\cos zx - z\cos yz)\,\mathbf{k}$$

$$= x(\cos xy - \cos zx)\,\mathbf{i} + y(\cos yz - \cos xy)\,\mathbf{j} + z(\cos zx - \cos yz)\,\mathbf{k}$$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}(\sin yz) + \dfrac{\partial}{\partial y}(\sin zx) + \dfrac{\partial}{\partial z}(\sin xy) = 0 + 0 + 0 = 0$

5. (a) $\operatorname{curl}\mathbf{F} = \nabla\times\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \dfrac{\sqrt{x}}{1+z} & \dfrac{\sqrt{y}}{1+x} & \dfrac{\sqrt{z}}{1+y} \end{vmatrix}$

$$= \left[\sqrt{z}\,(-1)(1+y)^{-2} - 0\right]\mathbf{i} - \left[0 - \sqrt{x}(-1)(1+z)^{-2}\right]\mathbf{j} + \left[\sqrt{y}\,(-1)(1+x)^{-2} - 0\right]\mathbf{k}$$

$$= -\frac{\sqrt{z}}{(1+y)^2}\,\mathbf{i} - \frac{\sqrt{x}}{(1+z)^2}\,\mathbf{j} - \frac{\sqrt{y}}{(1+x)^2}\,\mathbf{k}$$

(b) $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \dfrac{\partial}{\partial x}\left(\dfrac{\sqrt{x}}{1+z}\right) + \dfrac{\partial}{\partial y}\left(\dfrac{\sqrt{y}}{1+x}\right) + \dfrac{\partial}{\partial z}\left(\dfrac{\sqrt{z}}{1+y}\right)$

$$= \frac{1}{2\sqrt{x}\,(1+z)} + \frac{1}{2\sqrt{y}\,(1+x)} + \frac{1}{2\sqrt{z}\,(1+y)}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. (a) $\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \ln(2y+3z) & \ln(x+3z) & \ln(x+2y) \end{vmatrix}$

$$= \left(\frac{2}{x+2y} - \frac{3}{x+3z}\right)\mathbf{i} - \left(\frac{1}{x+2y} - \frac{3}{2y+3z}\right)\mathbf{j} + \left(\frac{1}{x+3z} - \frac{2}{2y+3z}\right)\mathbf{k}$$

$$= \left(\frac{2}{x+2y} - \frac{3}{x+3z}\right)\mathbf{i} + \left(\frac{3}{2y+3z} - \frac{1}{x+2y}\right)\mathbf{j} + \left(\frac{1}{x+3z} - \frac{2}{2y+3z}\right)\mathbf{k}$$

(b) $\operatorname{div}\mathbf{F} = \nabla \cdot \mathbf{F} = \dfrac{\partial}{\partial x}\left[\ln(2y+3z)\right] + \dfrac{\partial}{\partial y}\left[\ln(x+3z)\right] + \dfrac{\partial}{\partial z}\left[\ln(x+2y)\right] = 0 + 0 + 0 = 0$

7. (a) $\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ e^x \sin y & e^y \sin z & e^z \sin x \end{vmatrix} = (0 - e^y \cos z)\,\mathbf{i} - (e^z \cos x - 0)\,\mathbf{j} + (0 - e^x \cos y)\,\mathbf{k}$

$$= \langle -e^y \cos z, -e^z \cos x, -e^x \cos y \rangle$$

(b) $\operatorname{div}\mathbf{F} = \nabla \cdot \mathbf{F} = \dfrac{\partial}{\partial x}(e^x \sin y) + \dfrac{\partial}{\partial y}(e^y \sin z) + \dfrac{\partial}{\partial z}(e^z \sin x) = e^x \sin y + e^y \sin z + e^z \sin x$

8. (a) $\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \arctan(xy) & \arctan(yz) & \arctan(zx) \end{vmatrix}$

$$= \left(0 - \frac{y}{1+(yz)^2}\right)\mathbf{i} - \left(\frac{z}{1+(zx)^2} - 0\right)\mathbf{j} + \left(0 - \frac{x}{1+(xy)^2}\right)\mathbf{k}$$

$$= \left\langle -\frac{y}{1+y^2z^2}, -\frac{z}{1+x^2z^2}, -\frac{x}{1+x^2y^2} \right\rangle$$

(b) $\operatorname{div}\mathbf{F} = \nabla \cdot \mathbf{F} = \dfrac{\partial}{\partial x}\left[\arctan(xy)\right] + \dfrac{\partial}{\partial y}\left[\arctan(yz)\right] + \dfrac{\partial}{\partial z}\left[\arctan(zx)\right] = \dfrac{y}{1+x^2y^2} + \dfrac{z}{1+y^2z^2} + \dfrac{x}{1+x^2z^2}$

9. (a) $\operatorname{div}\mathbf{F}$ is negative because the vectors that start near P are shorter than those that end near P. Intuitively, if $\mathbf{F}$ represents a velocity field of fluid flow, then the net flow at P is inward.

(b) $\operatorname{curl}\mathbf{F}$ is zero because we can see that if $\mathbf{F}$ represents a velocity field of fluid flow, then a paddle wheel placed at P moves with the fluid, but does not rotate.

10. (a) $\operatorname{div}\mathbf{F}$ is positive because the vectors that start near P are longer than those that end near P. Intuitively, if $\mathbf{F}$ represents a velocity field of fluid flow, then the net flow at P is outward.

(b) $\operatorname{curl}\mathbf{F}$ is zero because we can see that if $\mathbf{F}$ represents a velocity field of fluid flow, then a paddle wheel placed at P moves with the fluid, but does not rotate.

11. (a) $\operatorname{div}\mathbf{F}$ is zero because the vectors that start near P are the same length as those that end near P. Intuitively, if $\mathbf{F}$ represents a velocity field of fluid flow, then the net flow at P is zero.

(b) $\operatorname{curl}\mathbf{F} \neq 0$ because we can see that if $\mathbf{F}$ represents a velocity field of fluid flow, then a paddle wheel placed at P would rotate clockwise about its axis, and hence the curl vector there points in the direction of $-\mathbf{k}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

12. (a) $\operatorname{div}\mathbf{F}$ is zero because the vectors that start near P are the same length as those that end near P. Intuitively, if $\mathbf{F}$ represents a velocity field of fluid flow, then the net flow at P is zero.

(b) $\operatorname{curl}\mathbf{F} \neq 0$ because we can see that if $\mathbf{F}$ represents a velocity field of fluid flow, then a paddle wheel placed at P would rotate counterclockwise about its axis, and hence the curl vector there points in the direction of $\mathbf{k}$.

13. (a) We need to verify $\operatorname{curl}(\nabla f) = \mathbf{0}$ for $f(x, y, z) = \sin xyz$. First, $\nabla f = yz\cos xyz\,\mathbf{i} + xz\cos xyz\,\mathbf{j} + xy\cos xyz\,\mathbf{k}$. Then

$$\begin{aligned}
\operatorname{curl}(\nabla f) &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ yz\cos xyz & xz\cos xyz & xy\cos xyz \end{vmatrix} \\
&= [-x^2yz\sin xyz + x\cos xyz - (-x^2yz\sin xyz + x\cos xyz)]\,\mathbf{i} \\
&\quad - [-xy^2z\sin xyz + y\cos xyz - (-xy^2z\sin xyz + y\cos xyz)]\,\mathbf{j} \\
&\quad + [-xyz^2\sin xyz + z\cos xyz - (-xyz^2\sin xyz + z\cos xyz)]\,\mathbf{k} = \mathbf{0}
\end{aligned}$$

(b) We need to verify that $\operatorname{div}\operatorname{curl}\mathbf{F} = 0$ for $\mathbf{F}(x, y, z) = xyz^2\,\mathbf{i} + x^2yz^3\,\mathbf{j} + y^2\,\mathbf{k}$. First,

$$\begin{aligned}
\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ xyz^2 & x^2yz^3 & y^2 \end{vmatrix} \\
&= (2y - 3x^2yz^2)\,\mathbf{i} - (0 - 2xyz)\,\mathbf{j} + (2xyz^3 - xz^2)\,\mathbf{k} \\
&= (2y - 3x^2yz^2)\,\mathbf{i} + 2xyz\,\mathbf{j} + (2xyz^3 - xz^2)\,\mathbf{k}
\end{aligned}$$

Then

$$\begin{aligned}
\operatorname{div}\operatorname{curl}\mathbf{F} = \nabla \cdot (\nabla \times \mathbf{F}) &= \tfrac{\partial}{\partial x}(2y - 3x^2yz^2) + \tfrac{\partial}{\partial y}(2xyz) + \tfrac{\partial}{\partial z}(2xyz^3 - xz^2) \\
&= -6xyz^2 + 2xz + 6xyz^2 - 2xz = 0
\end{aligned}$$

14. (a) $\operatorname{curl} f = \nabla \times f$ is meaningless because f is a scalar field.

(b) $\operatorname{grad} f$ is a vector field.

(c) $\operatorname{div}\mathbf{F}$ is a scalar field.

(d) $\operatorname{curl}(\operatorname{grad} f)$ is a vector field.

(e) $\operatorname{grad}\mathbf{F}$ is meaningless because $\mathbf{F}$ is not a scalar field.

(f) $\operatorname{grad}(\operatorname{div}\mathbf{F})$ is a vector field.

(g) $\operatorname{div}(\operatorname{grad} f)$ is a scalar field.

(h) $\operatorname{grad}(\operatorname{div} f)$ is meaningless because f is a scalar field.

(i) $\operatorname{curl}(\operatorname{curl}\mathbf{F})$ is a vector field.

(j) $\operatorname{div}(\operatorname{div}\mathbf{F})$ is meaningless because $\operatorname{div}\mathbf{F}$ is a scalar field.

(k) $(\operatorname{grad} f) \times (\operatorname{div}\mathbf{F})$ is meaningless because $\operatorname{div}\mathbf{F}$ is a scalar field.

(l) $\operatorname{div}(\operatorname{curl}(\operatorname{grad} f))$ is a scalar field.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. $\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ 2xy^3z^2 & 3x^2y^2z^2 & 2x^2y^3z \end{vmatrix}$

$$= (6x^2y^2z - 6x^2y^2z)\,\mathbf{i} - (4xy^3z - 4xy^3z)\,\mathbf{j} + (6xy^2z^2 - 6xy^2z^2)\,\mathbf{k} = \mathbf{0}$$

and $\mathbf{F}$ is defined on all of $\mathbb{R}^3$ whose component functions have continuous partial derivatives, so, by Theorem 4, $\mathbf{F}$ is conservative. Thus, there exists a function f such that $\nabla f = \mathbf{F}$. Now $f_x(x, y, z) = 2xy^3z^2$ implies that $f(x, y, z) = x^2y^3z^2 + g(y, z)$ and then $f_y(x, y, z) = 3x^2y^2z^2 + g_y(y, z)$. But $f_y(x, y, z) = 3x^2y^2z^2$, so $g(y, z) = h(z)$ and $f(x, y, z) = x^2y^3z^2 + h(z)$. Thus, $f_z(x, y, z) = 2x^2y^3z + h'(z)$, but $f_z(x, y, z) = 2x^2y^3z$, so $h(z) = K$, a constant. Hence, a potential function for $\mathbf{F}$ is $f(x, y, z) = x^2y^3z^2 + K$.

16. $\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ yz & xz + y & xy - x \end{vmatrix} = (x - x)\,\mathbf{i} - (y - 1 - y)\,\mathbf{j} + (z - z)\,\mathbf{k} = \mathbf{j} \neq \mathbf{0},$

so $\mathbf{F}$ is not conservative.

17. $\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \ln y & (x/y) + \ln z & y/z \end{vmatrix} = (1/z - 1/z)\,\mathbf{i} - (0 - 0)\,\mathbf{j} + (1/y - 1/y)\,\mathbf{k} = \mathbf{0}$

and the partial derivatives of the component functions are defined and continuous on the open set $\{(x, y, z) \mid y, z > 0\}$, so, by Theorem 4, $\mathbf{F}$ is conservative. Thus, there exists a function f such that $\nabla f = \mathbf{F}$. Now $f_x(x, y, z) = \ln y$ implies that $f(x, y, z) = x \ln y + g(y, z)$ and then $f_y(x, y, z) = (x/y) + g_y(y, z)$. But $f_y(x, y, z) = (x/y) + \ln z$, so $g(y, z) = y \ln z + h(z)$ and $f(x, y, z) = x \ln y + y \ln z + h(z)$. Thus, $f_z(x, y, z) = (y/z) + h'(z)$, but $f_z(x, y, z) = y/z$, so $h(z) = K$, a constant. Hence, a potential function for $\mathbf{F}$ is $f(x, y, z) = x \ln y + y \ln z + K$.

18. $\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ yz\sin(xy) & xz\sin(xy) & -\cos(xy) \end{vmatrix}$

$$\begin{aligned} &= [x\sin(xy) - x\sin(xy)]\,\mathbf{i} - [y\sin(xy) - y\sin(xy)]\,\mathbf{j} \\ &\qquad + [z\sin(xy) + xyz\cos(xy) - z\sin(xy) - xyz\cos(xy)]\,\mathbf{k} \\ &= \mathbf{0} \end{aligned}$$

and $\mathbf{F}$ is defined on all of $\mathbb{R}^3$ whose component functions have continuous partial derivatives, so, by Theorem 4, $\mathbf{F}$ is conservative. Thus, there exists a function f such that $\nabla f = \mathbf{F}$. Now $f_x(x, y, z) = yz\sin(xy)$ implies that $f(x, y, z) = -z\cos(xy) + g(y, z)$ and then $f_y(x, y, z) = xz\sin(xy) + g_y(y, z)$. But $f_y(x, y, z) = xz\sin(xy)$, so $g(y, z) = h(z)$ and $f(x, y, z) = -z\cos(xy) + h(z)$. Thus, $f_z(x, y, z) = -\cos(xy) + h'(z)$, but $f_z(x, y, z) = -\cos(xy)$, so $h(z) = K$, a constant. Hence, a potential function for $\mathbf{F}$ is $f(x, y, z) = -z\cos(xy) + K$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. $\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ yz^2e^{xz} & ze^{xz} & xyze^{xz} \end{vmatrix}$

$$= (xze^{xz} - e^{xz} - xze^{xz})\,\mathbf{i} - (yze^{xz} + xyz^2e^{xz} - 2yze^{xz} - xyz^2e^{xz})\,\mathbf{j} + (z^2e^{xz} - z^2e^{xz})\,\mathbf{k}$$

$$= -e^{xz}\,\mathbf{i} + yze^{xz}\,\mathbf{j} \neq \mathbf{0},$$

so $\mathbf{F}$ is not conservative.

20. $\operatorname{curl}\mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ e^z\cos x & e^y\cos z & e^z\sin x - e^y\sin z \end{vmatrix}$

$$= [-e^y\sin z - (-e^y\sin z)]\,\mathbf{i} - (e^z\cos x - e^z\cos x)\,\mathbf{j} + (0-0)\,\mathbf{k} = \mathbf{0}$$

and $\mathbf{F}$ is defined on all of $\mathbb{R}^3$ whose component functions have continuous partial derivatives, so, by Theorem 4, $\mathbf{F}$ is conservative. Thus, there exists a function f such that $\nabla f = \mathbf{F}$. Now $f_x(x,y,z) = e^z\cos x$ implies that $f(x,y,z) = e^z\sin x + g(y,z)$ and then $f_y(x,y,z) = g_y(y,z)$. But $f_y(x,y,z) = e^y\cos z$, so $g(y,z) = e^y\cos z + h(z)$ and $f(x,y,z) = e^z\sin x + e^y\cos z + h(z)$. Thus, $f_z(x,y,z) = e^z\sin x - e^y\sin z + h'(z)$, but $f_z(x,y,z) = e^z\sin x - e^y\sin z$, so $h(z) = K$, a constant. Hence, a potential function for $\mathbf{F}$ is $f(x,y,z) = e^z\sin x + e^y\cos z + K$.

21. No. Assume there is such a $\mathbf{G}$. Then $\operatorname{div}(\operatorname{curl}\mathbf{G}) = \dfrac{\partial}{\partial x}(x\sin y) + \dfrac{\partial}{\partial y}(\cos y) + \dfrac{\partial}{\partial z}(z - xy) = \sin y - \sin y + 1 \neq 0$, which contradicts Theorem 11.

22. No. Assume there is such a $\mathbf{G}$. Then $\operatorname{div}(\operatorname{curl}\mathbf{G}) = \dfrac{\partial}{\partial x}(x) + \dfrac{\partial}{\partial y}(y) + \dfrac{\partial}{\partial z}(z) = 1 + 1 + 1 \neq 0$ which contradicts Theorem 11.

23. $\operatorname{curl}\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ f(x) & g(y) & h(z) \end{vmatrix} = (0-0)\,\mathbf{i} + (0-0)\,\mathbf{j} + (0-0)\,\mathbf{k} = \mathbf{0}$. Hence $\mathbf{F} = f(x)\,\mathbf{i} + g(y)\,\mathbf{j} + h(z)\,\mathbf{k}$ is irrotational.

24. $\operatorname{div}\mathbf{F} = \dfrac{\partial}{\partial x}(f(y,z)) + \dfrac{\partial}{\partial y}(g(x,z)) + \dfrac{\partial}{\partial z}(h(x,y)) = 0$ so $\mathbf{F}$ is incompressible.

For Exercises 25–31, let $\mathbf{F}(x,y,z) = P_1\,\mathbf{i} + Q_1\,\mathbf{j} + R_1\,\mathbf{k}$ and $\mathbf{G}(x,y,z) = P_2\,\mathbf{i} + Q_2\,\mathbf{j} + R_2\,\mathbf{k}$.

25. $\operatorname{div}(\mathbf{F}+\mathbf{G}) = \operatorname{div}\langle P_1+P_2, Q_1+Q_2, R_1+R_2\rangle = \dfrac{\partial(P_1+P_2)}{\partial x} + \dfrac{\partial(Q_1+Q_2)}{\partial y} + \dfrac{\partial(R_1+R_2)}{\partial z}$

$$= \frac{\partial P_1}{\partial x} + \frac{\partial P_2}{\partial x} + \frac{\partial Q_1}{\partial y} + \frac{\partial Q_2}{\partial y} + \frac{\partial R_1}{\partial z} + \frac{\partial R_2}{\partial z} = \left(\frac{\partial P_1}{\partial x} + \frac{\partial Q_1}{\partial y} + \frac{\partial R_1}{\partial z}\right) + \left(\frac{\partial P_2}{\partial x} + \frac{\partial Q_2}{\partial y} + \frac{\partial R_2}{\partial z}\right)$$

$$= \operatorname{div}\langle P_1, Q_1, R_1\rangle + \operatorname{div}\langle P_2, Q_2, R_2\rangle = \operatorname{div}\mathbf{F} + \operatorname{div}\mathbf{G}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $$\begin{aligned}\operatorname{curl}\mathbf{F}+\operatorname{curl}\mathbf{G} &= \left[\left(\frac{\partial R_1}{\partial y}-\frac{\partial Q_1}{\partial z}\right)\mathbf{i}+\left(\frac{\partial P_1}{\partial z}-\frac{\partial R_1}{\partial x}\right)\mathbf{j}+\left(\frac{\partial Q_1}{\partial x}-\frac{\partial P_1}{\partial y}\right)\mathbf{k}\right]\\ &\qquad+\left[\left(\frac{\partial R_2}{\partial y}-\frac{\partial Q_2}{\partial z}\right)\mathbf{i}+\left(\frac{\partial P_2}{\partial z}-\frac{\partial R_2}{\partial x}\right)\mathbf{j}+\left(\frac{\partial Q_2}{\partial x}-\frac{\partial P_2}{\partial y}\right)\mathbf{k}\right]\\ &= \left[\frac{\partial(R_1+R_2)}{\partial y}-\frac{\partial(Q_1+Q_2)}{\partial z}\right]\mathbf{i}+\left[\frac{\partial(P_1+P_2)}{\partial z}-\frac{\partial(R_1+R_2)}{\partial x}\right]\mathbf{j}\\ &\qquad+\left[\frac{\partial(Q_1+Q_2)}{\partial x}-\frac{\partial(P_1+P_2)}{\partial y}\right]\mathbf{k}=\operatorname{curl}(\mathbf{F}+\mathbf{G})\end{aligned}$$

27. $$\begin{aligned}\operatorname{div}(f\mathbf{F}) &= \operatorname{div}(f\langle P_1,Q_1,R_1\rangle)=\operatorname{div}\langle fP_1,fQ_1,fR_1\rangle=\frac{\partial(fP_1)}{\partial x}+\frac{\partial(fQ_1)}{\partial y}+\frac{\partial(fR_1)}{\partial z}\\ &= \left(f\frac{\partial P_1}{\partial x}+P_1\frac{\partial f}{\partial x}\right)+\left(f\frac{\partial Q_1}{\partial y}+Q_1\frac{\partial f}{\partial y}\right)+\left(f\frac{\partial R_1}{\partial z}+R_1\frac{\partial f}{\partial z}\right)\\ &= f\left(\frac{\partial P_1}{\partial x}+\frac{\partial Q_1}{\partial y}+\frac{\partial R_1}{\partial z}\right)+\langle P_1,Q_1,R_1\rangle\cdot\left\langle\frac{\partial f}{\partial x},\frac{\partial f}{\partial y},\frac{\partial f}{\partial z}\right\rangle=f\operatorname{div}\mathbf{F}+\mathbf{F}\cdot\nabla f\end{aligned}$$

28. $$\begin{aligned}\operatorname{curl}(f\mathbf{F}) &= \left[\frac{\partial(fR_1)}{\partial y}-\frac{\partial(fQ_1)}{\partial z}\right]\mathbf{i}+\left[\frac{\partial(fP_1)}{\partial z}-\frac{\partial(fR_1)}{\partial x}\right]\mathbf{j}+\left[\frac{\partial(fQ_1)}{\partial x}-\frac{\partial(fP_1)}{\partial y}\right]\mathbf{k}\\ &= \left[f\frac{\partial R_1}{\partial y}+R_1\frac{\partial f}{\partial y}-f\frac{\partial Q_1}{\partial z}-Q_1\frac{\partial f}{\partial z}\right]\mathbf{i}+\left[f\frac{\partial P_1}{\partial z}+P_1\frac{\partial f}{\partial z}-f\frac{\partial R_1}{\partial x}-R_1\frac{\partial f}{\partial x}\right]\mathbf{j}\\ &\qquad+\left[f\frac{\partial Q_1}{\partial x}+Q_1\frac{\partial f}{\partial x}-f\frac{\partial P_1}{\partial y}-P_1\frac{\partial f}{\partial y}\right]\mathbf{k}\\ &= f\left[\frac{\partial R_1}{\partial y}-\frac{\partial Q_1}{\partial z}\right]\mathbf{i}+f\left[\frac{\partial P_1}{\partial z}-\frac{\partial R_1}{\partial x}\right]\mathbf{j}+f\left[\frac{\partial Q_1}{\partial x}-\frac{\partial P_1}{\partial y}\right]\mathbf{k}\\ &\qquad+\left[R_1\frac{\partial f}{\partial y}-Q_1\frac{\partial f}{\partial z}\right]\mathbf{i}+\left[P_1\frac{\partial f}{\partial z}-R_1\frac{\partial f}{\partial x}\right]\mathbf{j}+\left[Q_1\frac{\partial f}{\partial x}-P_1\frac{\partial f}{\partial y}\right]\mathbf{k}\\ &= f\operatorname{curl}\mathbf{F}+(\nabla f)\times\mathbf{F}\end{aligned}$$

29. $$\begin{aligned}\operatorname{div}(\mathbf{F}\times\mathbf{G}) &= \nabla\cdot(\mathbf{F}\times\mathbf{G})=\begin{vmatrix}\partial/\partial x & \partial/\partial y & \partial/\partial z\\ P_1 & Q_1 & R_1\\ P_2 & Q_2 & R_2\end{vmatrix}=\frac{\partial}{\partial x}\begin{vmatrix}Q_1 & R_1\\ Q_2 & R_2\end{vmatrix}-\frac{\partial}{\partial y}\begin{vmatrix}P_1 & R_1\\ P_2 & R_2\end{vmatrix}+\frac{\partial}{\partial z}\begin{vmatrix}P_1 & Q_1\\ P_2 & Q_2\end{vmatrix}\\ &= \left[Q_1\frac{\partial R_2}{\partial x}+R_2\frac{\partial Q_1}{\partial x}-Q_2\frac{\partial R_1}{\partial x}-R_1\frac{\partial Q_2}{\partial x}\right]-\left[P_1\frac{\partial R_2}{\partial y}+R_2\frac{\partial P_1}{\partial y}-P_2\frac{\partial R_1}{\partial y}-R_1\frac{\partial P_2}{\partial y}\right]\\ &\qquad+\left[P_1\frac{\partial Q_2}{\partial z}+Q_2\frac{\partial P_1}{\partial z}-P_2\frac{\partial Q_1}{\partial z}-Q_1\frac{\partial P_2}{\partial z}\right]\\ &= \left[P_2\left(\frac{\partial R_1}{\partial y}-\frac{\partial Q_1}{\partial z}\right)+Q_2\left(\frac{\partial P_1}{\partial z}-\frac{\partial R_1}{\partial x}\right)+R_2\left(\frac{\partial Q_1}{\partial x}-\frac{\partial P_1}{\partial y}\right)\right]\\ &\qquad-\left[P_1\left(\frac{\partial R_2}{\partial y}-\frac{\partial Q_2}{\partial z}\right)+Q_1\left(\frac{\partial P_2}{\partial z}-\frac{\partial R_2}{\partial x}\right)+R_1\left(\frac{\partial Q_2}{\partial x}-\frac{\partial P_2}{\partial y}\right)\right]\\ &= \mathbf{G}\cdot\operatorname{curl}\mathbf{F}-\mathbf{F}\cdot\operatorname{curl}\mathbf{G}\end{aligned}$$

30. $\operatorname{div}(\nabla f\times\nabla g)=\nabla g\cdot\operatorname{curl}(\nabla f)-\nabla f\cdot\operatorname{curl}(\nabla g)$ [by Exercise 29] $=0$ [by Theorem 3]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. $\operatorname{curl}(\operatorname{curl}\mathbf{F}) = \nabla \times (\nabla \times \mathbf{F}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \partial R_1/\partial y - \partial Q_1/\partial z & \partial P_1/\partial z - \partial R_1/\partial x & \partial Q_1/\partial x - \partial P_1/\partial y \end{vmatrix}$

$$= \left(\frac{\partial^2 Q_1}{\partial y \partial x} - \frac{\partial^2 P_1}{\partial y^2} - \frac{\partial^2 P_1}{\partial z^2} + \frac{\partial^2 R_1}{\partial z \partial x}\right)\mathbf{i} + \left(\frac{\partial^2 R_1}{\partial z \partial y} - \frac{\partial^2 Q_1}{\partial z^2} - \frac{\partial^2 Q_1}{\partial x^2} + \frac{\partial^2 P_1}{\partial x \partial y}\right)\mathbf{j}$$
$$+ \left(\frac{\partial^2 P_1}{\partial x \partial z} - \frac{\partial^2 R_1}{\partial x^2} - \frac{\partial^2 R_1}{\partial y^2} + \frac{\partial^2 Q_1}{\partial y \partial z}\right)\mathbf{k}$$

Now let's consider $\operatorname{grad}(\operatorname{div}\mathbf{F}) - \nabla^2\mathbf{F}$ and compare with the above. (Note that $\nabla^2\mathbf{F}$ is defined in the discussion following Example 16.5.5.)

$$\operatorname{grad}(\operatorname{div}\mathbf{F}) - \nabla^2\mathbf{F} = \left[\left(\frac{\partial^2 P_1}{\partial x^2} + \frac{\partial^2 Q_1}{\partial x \partial y} + \frac{\partial^2 R_1}{\partial x \partial z}\right)\mathbf{i} + \left(\frac{\partial^2 P_1}{\partial y \partial x} + \frac{\partial^2 Q_1}{\partial y^2} + \frac{\partial^2 R_1}{\partial y \partial z}\right)\mathbf{j} + \left(\frac{\partial^2 P_1}{\partial z \partial x} + \frac{\partial^2 Q_1}{\partial z \partial y} + \frac{\partial^2 R_1}{\partial z^2}\right)\mathbf{k}\right]$$
$$- \left[\left(\frac{\partial^2 P_1}{\partial x^2} + \frac{\partial^2 P_1}{\partial y^2} + \frac{\partial^2 P_1}{\partial z^2}\right)\mathbf{i} + \left(\frac{\partial^2 Q_1}{\partial x^2} + \frac{\partial^2 Q_1}{\partial y^2} + \frac{\partial^2 Q_1}{\partial z^2}\right)\mathbf{j}\right.$$
$$\left. + \left(\frac{\partial^2 R_1}{\partial x^2} + \frac{\partial^2 R_1}{\partial y^2} + \frac{\partial^2 R_1}{\partial z^2}\right)\mathbf{k}\right]$$
$$= \left(\frac{\partial^2 Q_1}{\partial x \partial y} + \frac{\partial^2 R_1}{\partial x \partial z} - \frac{\partial^2 P_1}{\partial y^2} - \frac{\partial^2 P_1}{\partial z^2}\right)\mathbf{i} + \left(\frac{\partial^2 P_1}{\partial y \partial x} + \frac{\partial^2 R_1}{\partial y \partial z} - \frac{\partial^2 Q_1}{\partial x^2} - \frac{\partial^2 Q_1}{\partial z^2}\right)\mathbf{j}$$
$$+ \left(\frac{\partial^2 P_1}{\partial z \partial x} + \frac{\partial^2 Q_1}{\partial z \partial y} - \frac{\partial^2 R_1}{\partial x^2} - \frac{\partial^2 R_2}{\partial y^2}\right)\mathbf{k}$$

Then applying Clairaut's Theorem to reverse the order of differentiation in the second partial derivatives as needed and comparing, we have $\operatorname{curl}\operatorname{curl}\mathbf{F} = \operatorname{grad}\operatorname{div}\mathbf{F} - \nabla^2\mathbf{F}$ as desired.

32. (a) $\nabla \cdot \mathbf{r} = \left(\dfrac{\partial}{\partial x}\mathbf{i} + \dfrac{\partial}{\partial y}\mathbf{j} + \dfrac{\partial}{\partial z}\mathbf{k}\right) \cdot (x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}) = 1 + 1 + 1 = 3$

(b) $\nabla \cdot (r\mathbf{r}) = \nabla \cdot \sqrt{x^2 + y^2 + z^2}\,(x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k})$

$$= \left(\frac{x^2}{\sqrt{x^2+y^2+z^2}} + \sqrt{x^2+y^2+z^2}\right) + \left(\frac{y^2}{\sqrt{x^2+y^2+z^2}} + \sqrt{x^2+y^2+z^2}\right)$$
$$+ \left(\frac{z^2}{\sqrt{x^2+y^2+z^2}} + \sqrt{x^2+y^2+z^2}\right)$$
$$= \frac{1}{\sqrt{x^2+y^2+z^2}}\,(4x^2 + 4y^2 + 4z^2) = 4\sqrt{x^2+y^2+z^2} = 4r$$

Another method:

By Exercise 27, $\nabla \cdot (r\mathbf{r}) = \operatorname{div}(r\mathbf{r}) = r \operatorname{div}\mathbf{r} + \mathbf{r} \cdot \nabla r = 3r + \mathbf{r} \cdot \dfrac{\mathbf{r}}{r}$ [see Exercise 33(a) below] $= 4r$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) $\nabla^2 r^3 = \nabla^2 \left(x^2+y^2+z^2\right)^{3/2}$

$$= \tfrac{\partial}{\partial x}\left[\tfrac{3}{2}(x^2+y^2+z^2)^{1/2}(2x)\right] + \tfrac{\partial}{\partial y}\left[\tfrac{3}{2}(x^2+y^2+z^2)^{1/2}(2y)\right] + \tfrac{\partial}{\partial z}\left[\tfrac{3}{2}(x^2+y^2+z^2)^{1/2}(2z)\right]$$

$$= 3\left[\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2x)(x) + (x^2+y^2+z^2)^{1/2}\right]$$
$$+ 3\left[\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2y)(y) + (x^2+y^2+z^2)^{1/2}\right]$$
$$+ 3\left[\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2z)(z) + (x^2+y^2+z^2)^{1/2}\right]$$

$$= 3(x^2+y^2+z^2)^{-1/2}(4x^2+4y^2+4z^2) = 12(x^2+y^2+z^2)^{1/2} = 12r$$

Another method: $\frac{\partial}{\partial x}\left(x^2+y^2+z^2\right)^{3/2} = 3x\sqrt{x^2+y^2+z^2} \quad \Rightarrow \quad \nabla r^3 = 3r(x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k}) = 3r\,\mathbf{r}$, so $\nabla^2 r^3 = \nabla\cdot\nabla r^3 = \nabla\cdot(3r\,\mathbf{r}) = 3(4r) = 12r$ by part (b).

33. (a) $\nabla r = \nabla\sqrt{x^2+y^2+z^2} = \dfrac{x}{\sqrt{x^2+y^2+z^2}}\,\mathbf{i} + \dfrac{y}{\sqrt{x^2+y^2+z^2}}\,\mathbf{j} + \dfrac{z}{\sqrt{x^2+y^2+z^2}}\,\mathbf{k} = \dfrac{x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k}}{\sqrt{x^2+y^2+z^2}} = \dfrac{\mathbf{r}}{r}$

(b) $\nabla\times\mathbf{r} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \dfrac{\partial}{\partial x} & \dfrac{\partial}{\partial y} & \dfrac{\partial}{\partial z} \\ x & y & z \end{vmatrix} = \left[\dfrac{\partial}{\partial y}(z) - \dfrac{\partial}{\partial z}(y)\right]\mathbf{i} + \left[\dfrac{\partial}{\partial z}(x) - \dfrac{\partial}{\partial x}(z)\right]\mathbf{j} + \left[\dfrac{\partial}{\partial x}(y) - \dfrac{\partial}{\partial y}(x)\right]\mathbf{k} = \mathbf{0}$

(c) $\nabla\left(\dfrac{1}{r}\right) = \nabla\left(\dfrac{1}{\sqrt{x^2+y^2+z^2}}\right)$

$$= \frac{-\dfrac{1}{2\sqrt{x^2+y^2+z^2}}(2x)}{x^2+y^2+z^2}\,\mathbf{i} - \frac{\dfrac{1}{2\sqrt{x^2+y^2+z^2}}(2y)}{x^2+y^2+z^2}\,\mathbf{j} - \frac{\dfrac{1}{2\sqrt{x^2+y^2+z^2}}(2z)}{x^2+y^2+z^2}\,\mathbf{k}$$

$$= -\frac{x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k}}{(x^2+y^2+z^2)^{3/2}} = -\frac{\mathbf{r}}{r^3}$$

(d) $\nabla\ln r = \nabla\ln(x^2+y^2+z^2)^{1/2} = \frac{1}{2}\nabla\ln(x^2+y^2+z^2)$

$$= \frac{x}{x^2+y^2+z^2}\,\mathbf{i} + \frac{y}{x^2+y^2+z^2}\,\mathbf{j} + \frac{z}{x^2+y^2+z^2}\,\mathbf{k} = \frac{x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k}}{x^2+y^2+z^2} = \frac{\mathbf{r}}{r^2}$$

34. $\mathbf{r} = x\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k} \quad\Rightarrow\quad r = |\mathbf{r}| = \sqrt{x^2+y^2+z^2}$, so

$$\mathbf{F} = \frac{\mathbf{r}}{r^p} = \frac{x}{(x^2+y^2+z^2)^{p/2}}\,\mathbf{i} + \frac{y}{(x^2+y^2+z^2)^{p/2}}\,\mathbf{j} + \frac{z}{(x^2+y^2+z^2)^{p/2}}\,\mathbf{k}$$

Then $\dfrac{\partial}{\partial x}\dfrac{x}{(x^2+y^2+z^2)^{p/2}} = \dfrac{(x^2+y^2+z^2)-px^2}{(x^2+y^2+z^2)^{1+p/2}} = \dfrac{r^2-px^2}{r^{p+2}}$. Similarly,

$\dfrac{\partial}{\partial y}\dfrac{y}{(x^2+y^2+z^2)^{p/2}} = \dfrac{r^2-py^2}{r^{p+2}}$ and $\dfrac{\partial}{\partial z}\dfrac{z}{(x^2+y^2+z^2)^{p/2}} = \dfrac{r^2-pz^2}{r^{p+2}}$. Thus

$$\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F} = \frac{r^2-px^2}{r^{p+2}} + \frac{r^2-py^2}{r^{p+2}} + \frac{r^2-pz^2}{r^{p+2}} = \frac{3r^2-px^2-py^2-pz^2}{r^{p+2}}$$

$$= \frac{3r^2-p(x^2+y^2+z^2)}{r^{p+2}} = \frac{3r^2-pr^2}{r^{p+2}} = \frac{3-p}{r^p}$$

Consequently, if $p=3$ we have $\operatorname{div}\mathbf{F} = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. By (13), $\oint_C f(\nabla g)\cdot\mathbf{n}\,ds = \iint_D \operatorname{div}(f\nabla g)\,dA = \iint_D [f\operatorname{div}(\nabla g) + \nabla g\cdot\nabla f]\,dA$ by Exercise 27. But $\operatorname{div}(\nabla g) = \nabla^2 g$. Hence $\iint_D f\nabla^2 g\,dA = \oint_C f(\nabla g)\cdot\mathbf{n}\,ds - \iint_D \nabla g\cdot\nabla f\,dA$.

36. By Exercise 35, $\iint_D f\nabla^2 g\,dA = \oint_C f(\nabla g)\cdot\mathbf{n}\,ds - \iint_D \nabla g\cdot\nabla f\,dA$ and $\iint_D g\nabla^2 f\,dA = \oint_C g(\nabla f)\cdot\mathbf{n}\,ds - \iint_D \nabla f\cdot\nabla g\,dA$. Hence $\iint_D \left(f\nabla^2 g - g\nabla^2 f\right)dA = \oint_C [f(\nabla g)\cdot\mathbf{n} - g(\nabla f)\cdot\mathbf{n}]\,ds + \iint_D (\nabla f\cdot\nabla g - \nabla g\cdot\nabla f)\,dA = \oint_C [f\nabla g - g\nabla f]\cdot\mathbf{n}\,ds$.

37. Let $f(x,y) = 1$. Then $\nabla f = \mathbf{0}$ and Green's first identity (see Exercise 35) says $\iint_D \nabla^2 g\,dA = \oint_C (\nabla g)\cdot\mathbf{n}\,ds - \iint_D \mathbf{0}\cdot\nabla g\,dA \quad\Rightarrow\quad \iint_D \nabla^2 g\,dA = \oint_C \nabla g\cdot\mathbf{n}\,ds$. But g is harmonic on D, so $\nabla^2 g = 0 \quad\Rightarrow\quad \oint_C \nabla g\cdot\mathbf{n}\,ds = 0$ and $\oint_C D_{\mathbf{n}} g\,ds = \oint_C (\nabla g\cdot\mathbf{n})\,ds = 0$.

38. Let $g = f$. Then Green's first identity (see Exercise 35) says $\iint_D f\,\nabla^2 f\,dA = \oint_C (f)(\nabla f)\cdot\mathbf{n}\,ds - \iint_D \nabla f\cdot\nabla f\,dA$. But f is harmonic, so $\nabla^2 f = 0$, and $\nabla f\cdot\nabla f = |\nabla f|^2$, so we have $0 = \oint_C (f)(\nabla f)\cdot\mathbf{n}\,ds - \iint_D |\nabla f|^2\,dA \quad\Rightarrow$ $\iint_D |\nabla f|^2\,dA = \oint_C (f)(\nabla f)\cdot\mathbf{n}\,ds = 0$ since $f(x,y) = 0$ on C.

39. (a) We know that $\omega = v/d$ (where v is the tangential speed), and from the diagram $\sin\theta = d/r$ (where $r = |\mathbf{r}|$) $\Rightarrow$ $v = d\omega = (\sin\theta)r\omega = |\mathbf{w}\times\mathbf{r}|$ (by 12.4.9). But $\mathbf{v}$ is perpendicular to both $\mathbf{w}$ and $\mathbf{r}$, so that $\mathbf{v} = \mathbf{w}\times\mathbf{r}$.

(b) From part (a), $\mathbf{v} = \mathbf{w}\times\mathbf{r} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0 & \omega \\ x & y & z \end{vmatrix} = (0\cdot z - \omega y)\,\mathbf{i} - (0\cdot z - \omega x)\,\mathbf{j} + (0\cdot y - 0\cdot x)\,\mathbf{k} = -\omega y\,\mathbf{i} + \omega x\,\mathbf{j}$.

(c) $\operatorname{curl}\mathbf{v} = \nabla\times\mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ -\omega y & \omega x & 0 \end{vmatrix}$

$$= \left[\frac{\partial}{\partial y}(0) - \frac{\partial}{\partial z}(\omega x)\right]\mathbf{i} + \left[\frac{\partial}{\partial z}(-\omega y) - \frac{\partial}{\partial x}(0)\right]\mathbf{j} + \left[\frac{\partial}{\partial x}(\omega x) - \frac{\partial}{\partial y}(-\omega y)\right]\mathbf{k}$$
$$= [\omega - (-\omega)]\,\mathbf{k} = 2\omega\,\mathbf{k} = 2\mathbf{w}$$

40. Let $\mathbf{H} = \langle h_1, h_2, h_3\rangle$ and $\mathbf{E} = \langle E_1, E_2, E_3\rangle$.

(a) $\nabla\times(\nabla\times\mathbf{E}) = \nabla\times(\operatorname{curl}\mathbf{E}) = \nabla\times\left(-\dfrac{1}{c}\dfrac{\partial\mathbf{H}}{\partial t}\right) = -\dfrac{1}{c}\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \partial h_1/\partial t & \partial h_2/\partial t & \partial h_3/\partial t \end{vmatrix}$

$$= -\frac{1}{c}\left[\left(\frac{\partial^2 h_3}{\partial y\,\partial t} - \frac{\partial^2 h_2}{\partial z\,\partial t}\right)\mathbf{i} + \left(\frac{\partial^2 h_1}{\partial z\,\partial t} - \frac{\partial^2 h_3}{\partial x\,\partial t}\right)\mathbf{j} + \left(\frac{\partial^2 h_2}{\partial x\,\partial t} - \frac{\partial^2 h_1}{\partial y\,\partial t}\right)\mathbf{k}\right]$$

$$= -\frac{1}{c}\frac{\partial}{\partial t}\left[\left(\frac{\partial h_3}{\partial y} - \frac{\partial h_2}{\partial z}\right)\mathbf{i} + \left(\frac{\partial h_1}{\partial z} - \frac{\partial h_3}{\partial x}\right)\mathbf{j} + \left(\frac{\partial h_2}{\partial x} - \frac{\partial h_1}{\partial y}\right)\mathbf{k}\right]$$

[assuming that the partial derivatives are continuous so that the order of differentiation does not matter]

$$= -\frac{1}{c}\frac{\partial}{\partial t}\operatorname{curl}\mathbf{H} = -\frac{1}{c}\frac{\partial}{\partial t}\left(\frac{1}{c}\frac{\partial\mathbf{E}}{\partial t}\right) = -\frac{1}{c^2}\frac{\partial^2\mathbf{E}}{\partial t^2}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) $\nabla \times (\nabla \times \mathbf{H}) = \nabla \times (\operatorname{curl} \mathbf{H}) = \nabla \times \left(\dfrac{1}{c}\dfrac{\partial \mathbf{E}}{\partial t}\right) = \dfrac{1}{c}\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ \partial E_1/\partial t & \partial E_2/\partial t & \partial E_3/\partial t \end{vmatrix}$

$$= \frac{1}{c}\left[\left(\frac{\partial^2 E_3}{\partial y\,\partial t} - \frac{\partial^2 E_2}{\partial z\,\partial t}\right)\mathbf{i} + \left(\frac{\partial^2 E_1}{\partial z\,\partial t} - \frac{\partial^2 E_3}{\partial x\,\partial t}\right)\mathbf{j} + \left(\frac{\partial^2 E_2}{\partial x\,\partial t} - \frac{\partial^2 E_1}{\partial y\,\partial t}\right)\mathbf{k}\right]$$

$$= \frac{1}{c}\frac{\partial}{\partial t}\left[\left(\frac{\partial E_3}{\partial y} - \frac{\partial E_2}{\partial z}\right)\mathbf{i} + \left(\frac{\partial E_1}{\partial z} - \frac{\partial E_3}{\partial x}\right)\mathbf{j} + \left(\frac{\partial E_2}{\partial x} - \frac{\partial E_1}{\partial y}\right)\mathbf{k}\right]$$ [assuming that the partial derivatives are continuous so that the order of differentiation does not matter]

$$= \frac{1}{c}\frac{\partial}{\partial t}\operatorname{curl}\mathbf{E} = \frac{1}{c}\frac{\partial}{\partial t}\left(-\frac{1}{c}\frac{\partial \mathbf{H}}{\partial t}\right) = -\frac{1}{c^2}\frac{\partial^2 \mathbf{H}}{\partial t^2}$$

(c) Using Exercise 31, we have that $\operatorname{curl}\operatorname{curl}\mathbf{E} = \operatorname{grad}\operatorname{div}\mathbf{E} - \nabla^2\mathbf{E} \quad \Rightarrow$

$$\nabla^2\mathbf{E} = \operatorname{grad}\operatorname{div}\mathbf{E} - \operatorname{curl}\operatorname{curl}\mathbf{E} = \operatorname{grad} 0 + \frac{1}{c^2}\frac{\partial^2\mathbf{E}}{\partial t^2} \quad \text{[from part (a)]} \quad = \frac{1}{c^2}\frac{\partial^2\mathbf{E}}{\partial t^2}.$$

(d) As in part (c), $\nabla^2\mathbf{H} = \operatorname{grad}\operatorname{div}\mathbf{H} - \operatorname{curl}\operatorname{curl}\mathbf{H} = \operatorname{grad} 0 + \dfrac{1}{c^2}\dfrac{\partial^2\mathbf{H}}{\partial t^2}$ [using part (b)] $= \dfrac{1}{c^2}\dfrac{\partial^2\mathbf{H}}{\partial t^2}$.

41. For any continuous function f on $\mathbb{R}^3$, define a vector field $\mathbf{G}(x,y,z) = \langle g(x,y,z), 0, 0\rangle$ where $g(x,y,z) = \int_0^x f(t,y,z)\,dt$. Then $\operatorname{div}\mathbf{G} = \dfrac{\partial}{\partial x}(g(x,y,z)) + \dfrac{\partial}{\partial y}(0) + \dfrac{\partial}{\partial z}(0) = \dfrac{\partial}{\partial x}\int_0^x f(t,y,z)\,dt = f(x,y,z)$ by the Fundamental Theorem of Calculus. Thus every continuous function f on $\mathbb{R}^3$ is the divergence of some vector field.

16.6 Parametric Surfaces and Their Areas

1. $P(4,-5,1)$ lies on the parametric surface $\mathbf{r}(u,v) = \langle u+v, u-2v, 3+u-v\rangle$ if and only if there are values for u and v where $u+v=4$, $u-2v=-5$, and $3+u-v=1$. From the first equation we have $u=4-v$ and substituting into the second equation gives $4-v-2v=-5 \quad \Leftrightarrow \quad v=3$. Then $u=1$, and these values satisfy the third equation, so P does lie on the surface.

$Q(0,4,6)$ lies on $\mathbf{r}(u,v)$ if and only if $u+v=0$, $u-2v=4$, and $3+u-v=6$, but solving the first two equations simultaneoulsy gives $u=\frac{4}{3}$, $v=-\frac{4}{3}$ and these values do not satisfy the third equation, so Q does not lie on the surface.

2. $P(1,2,1)$ lies on the parametric surface $\mathbf{r}(u,v) = \left\langle 1+u-v, u+v^2, u^2-v^2\right\rangle$ if and only if there are values for u and v where $1+u-v=1$, $u+v^2=2$, and $u^2-v^2=1$. From the first equation we have $u=v$ and substituting into the third equation gives $0=1$, an impossibility, so P does not lie on the surface.

$Q(2,3,3)$ lies on $\mathbf{r}(u,v)$ if and only if $1+u-v=2$, $u+v^2=3$, and $u^2-v^2=3$. From the first equation we have $u=v+1$ and substituting into the second equation gives $v+1+v^2=3 \quad \Leftrightarrow \quad v^2+v-2=0 \quad \Leftrightarrow \quad (v+2)(v-1)=0$, so $v=-2 \quad \Rightarrow \quad u=-1$ or $v=1 \quad \Rightarrow \quad u=2$. The third equation is satisfied by $u=2$, $v=1$ so Q does lie on the surface.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. $\mathbf{r}(u,v) = (u+v)\,\mathbf{i} + (3-v)\,\mathbf{j} + (1+4u+5v)\,\mathbf{k} = \langle 0,3,1\rangle + u\,\langle 1,0,4\rangle + v\,\langle 1,-1,5\rangle$. From Example 3, we recognize this as a vector equation of a plane through the point $(0,3,1)$ and containing vectors $\mathbf{a} = \langle 1,0,4\rangle$ and $\mathbf{b} = \langle 1,-1,5\rangle$. If we wish to find a more conventional equation for the plane, a normal vector to the plane is $\mathbf{a}\times\mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 4 \\ 1 & -1 & 5 \end{vmatrix} = 4\,\mathbf{i} - \mathbf{j} - \mathbf{k}$ and an equation of the plane is $4(x-0)-(y-3)-(z-1)=0$ or $4x-y-z=-4$.

4. $\mathbf{r}(u,v) = u^2\,\mathbf{i} + u\cos v\,\mathbf{j} + u\sin v\,\mathbf{k}$, so the corresponding parametric equations for the surface are $x=u^2,\ y=u\cos v$, $z=u\sin v$. For any point (x,y,z) on the surface, we have $y^2+z^2 = u^2\cos^2 v + u^2\sin^2 v = u^2 = x$. Since no restrictions are placed on the parameters, the surface is $x = y^2+z^2$, which we recognize as a circular paraboloid whose axis is the x-axis.

5. $\mathbf{r}(s,t) = \langle s\cos t, s\sin t, s\rangle$, so the corresponding parametric equations for the surface are $x = s\cos t,\ \ y = s\sin t,\ \ z = s$. For any point (x,y,z) on the surface, we have $x^2+y^2 = s^2\cos^2 t + s^2\sin^2 t = s^2 = z^2$. Since no restrictions are placed on the parameters, the surface is $z^2 = x^2+y^2$, which we recognize as a circular cone with axis the z-axis.

6. $\mathbf{r}(s,t) = \langle 3\cos t, s, \sin t\rangle$, so the corresponding parametric equations for the surface are $x = 3\cos t,\ \ y = s,\ \ z = \sin t$. For any point (x,y,z) on the surface, we have $(x/3)^2 + z^2 = \cos^2 t + \sin^2 t = 1$, so vertical cross-sections parallel to the xz-plane are all identical ellipses. Since $y = s$ and $-1 \le s \le 1$, the surface is the portion of the elliptic cylinder $\frac{1}{9}x^2 + z^2 = 1$ corresponding to $-1 \le y \le 1$.

7. $\mathbf{r}(u,v) = \left\langle u^2, v^2, u+v\right\rangle,\ \ -1 \le u \le 1,\ \ -1 \le v \le 1.$

The surface has parametric equations $x = u^2$, $y = v^2$, $z = u+v$, $-1 \le u \le 1$, $-1 \le v \le 1$.

In Maple, the surface can be graphed by entering

```
plot3d([u^2,v^2,u+v],u=-1..1,v=-1..1);
```

In Mathematica we use the `ParametricPlot3D` command.

If we keep u constant at u_0, $x = u_0^2$, a constant, so the corresponding grid curves must be the curves parallel to the yz-plane. If v is constant, we have $y = v_0^2$, a constant, so these grid curves are the curves parallel to the xz-plane.

8. $\mathbf{r}(u,v) = \left\langle u, v^3, -v\right\rangle,\ \ -2 \le u \le 2,\ -2 \le v \le 2.$

The surface has parametric equations $x = u$, $y = v^3$, $z = -v$, $-2 \le u \le 2$, $-2 \le v \le 2$. If $u = u_0$ is constant, $x = u_0 =$ constant, so the corresponding grid curves are the curves parallel to the yz-plane. If $v = v_0$ is constant, $y = v_0^3 =$ constant, so the corresponding grid curves are the curves parallel to the xz-plane.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $\mathbf{r}(u,v) = \langle u^3, u\sin v, u\cos v\rangle,\ -1 \le u \le 1,\ 0 \le v \le 2\pi$

The surface has parametric equations $x = u^3,\ y = u\sin v$, $z = u\cos v,\ -1 \le u \le 1,\ 0 \le v \le 2\pi$. Note that if $u = u_0$ is constant then $x = u_0^3$ is constant and $y = u_0 \sin v$, $z = u_0 \cos v$ describe a circle in y, z of radius $|u_0|$, so the corresponding grid curves are circles parallel to the yz-plane. If $v = v_0$, a constant, the parametric equations become $x = u^3$, $y = u\sin v_0$, $z = u\cos v_0$. Then $y = (\tan v_0)z$, so these are the grid curves we see that lie in planes $y = kz$ that pass through the x-axis.

10. $\mathbf{r}(u,v) = \langle u, \sin(u+v), \sin v\rangle,\ -\pi \le u \le \pi,\ -\pi \le v \le \pi.$

The surface has parametric equations $x = u$, $y = \sin(u+v)$, $z = \sin v$, $-\pi \le u \le \pi$, $-\pi \le v \le \pi$. If $u = u_0$ is constant, $x = u_0 =$ constant, so the corresponding grid curves are the curves parallel to the yz-plane. If $v = v_0$ is constant, $z = \sin v_0 =$ constant, so the corresponding grid curves are the curves parallel to the xy-plane.

11. $x = \sin v,\ \ y = \cos u \sin 4v,\ \ z = \sin 2u\ \sin 4v,\ \ 0 \le u \le 2\pi,\ \ -\frac{\pi}{2} \le v \le \frac{\pi}{2}.$

Note that if $v = v_0$ is constant, then $x = \sin v_0$ is constant, so the corresponding grid curves must be parallel to the yz-plane. These are the vertically oriented grid curves we see, each shaped like a "figure-eight." When $u = u_0$ is held constant, the parametric equations become $x = \sin v$, $y = \cos u_0 \sin 4v$, $z = \sin 2u_0 \sin 4v$. Since z is a constant multiple of y, the corresponding grid curves are the curves contained in planes $z = ky$ that pass through the x-axis.

12. $x = \cos u,\ y = \sin u\ \sin v,\ z = \cos v,\ 0 \le u \le 2\pi,\ \ 0 \le v \le 2\pi.$

If $u = u_0$ is constant, then $x = \cos u_0 =$ constant, so the corresponding grid curves are the curves parallel to the yz-plane. If $v = v_0$ is constant, then $z = \cos v_0 =$ constant, so the corresponding grid curves are the curves parallel to the xy-plane.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $\mathbf{r}(u,v) = u\cos v\,\mathbf{i} + u\sin v\,\mathbf{j} + v\,\mathbf{k}$. The parametric equations for the surface are $x = u\cos v$, $y = u\sin v$, $z = v$. We look at the grid curves first; if we fix v, then x and y parametrize a straight line in the plane $z = v$ which intersects the z-axis. If u is held constant, the projection onto the xy-plane is circular; with $z = v$, each grid curve is a helix. The surface is a spiraling ramp, graph IV.

14. $\mathbf{r}(u,v) = uv^2\,\mathbf{i} + u^2v\,\mathbf{j} + (u^2 - v^2)\,\mathbf{k}$. The parametric equations for the surface are $x = uv^2,\ y = u^2v,\ z = u^2 - v^2$. If $u = u_0$ is held constant, then $x = u_0v^2$, $y = u_0^2v$ so $x = u_0(y/u_0^2)^2 = (1/u_0^3)y^2$, and $z = u_0^2 - v^2 = u_0^2 - (1/u_0)x$. Thus each grid curve corresponding to $u = u_0$ lies in the plane $z = u_0^2 - (1/u_0)x$ and its projection onto the xy-plane is a parabola $x = ky^2$ with axis the x-axis. Similarly, if $v = v_0$ is held constant, then $x = uv_0^2$, $y = u^2v_0 \quad \Rightarrow$ $y = (x/v_0^2)^2v_0 = (1/v_0^3)x^2$, and $z = u^2 - v_0^2 = (1/v_0)y - v_0^2$. Each grid curve lies in the plane $z = (1/v_0)y - v_0^2$ and its projection onto the xy-plane is a parabola $y = kx^2$ with axis the y-axis. The surface is graph VI.

15. $\mathbf{r}(u,v) = (u^3 - u)\,\mathbf{i} + v^2\,\mathbf{j} + u^2\,\mathbf{k}$. The parametric equations for the surface are $x = u^3 - u,\ y = v^2,\ z = u^2$. If we fix u then x and z are constant so each corresponding grid curve is contained in a line parallel to the y-axis. (Since $y = v^2 \geq 0$, the grid curves are half-lines.) If v is held constant, then $y = v^2 =$ constant, so each grid curve is contained in a plane parallel to the xz-plane. Since x and z are functions of u only, the grid curves all have the same shape. The surface is the cylinder shown in graph I.

16. $x = (1-u)(3+\cos v)\cos 4\pi u$, $y = (1-u)(3+\cos v)\sin 4\pi u$, $z = 3u + (1-u)\sin v$. These equations correspond to graph V: when $u = 0$, then $x = 3 + \cos v$, $y = 0$, and $z = \sin v$, which are equations of a circle with radius 1 in the xz-plane centered at $(3,0,0)$. When $u = \frac{1}{2}$, then $x = \frac{3}{2} + \frac{1}{2}\cos v$, $y = 0$, and $z = \frac{3}{2} + \frac{1}{2}\sin v$, which are equations of a circle with radius $\frac{1}{2}$ in the xz-plane centered at $\left(\frac{3}{2}, 0, \frac{3}{2}\right)$. When $u = 1$, then $x = y = 0$ and $z = 3$, giving the topmost point shown in the graph. This suggests that the grid curves with u constant are the vertically oriented circles visible on the surface. The spiralling grid curves correspond to keeping v constant.

17. $x = \cos^3 u\,\cos^3 v$, $y = \sin^3 u\,\cos^3 v$, $z = \sin^3 v$. If $v = v_0$ is held constant then $z = \sin^3 v_0$ is constant, so the corresponding grid curve lies in a horizontal plane. Several of the graphs exhibit horizontal grid curves, but the curves for this surface are neither ellipses nor straight lines, so graph III is the only possibility. (In fact, the horizontal grid curves here are members of the family $x = a\cos^3 u$, $y = a\sin^3 u$ and are called astroids.) The vertical grid curves we see on the surface correspond to $u = u_0$ held constant, as then we have $x = \cos^3 u_0\,\cos^3 v$, $y = \sin^3 u_0\,\cos^3 v$ so the corresponding grid curve lies in the vertical plane $y = (\tan^3 u_0)x$ through the z-axis.

18. $x = \sin u,\ y = \cos u\,\sin v,\ z = \sin v$. If $v = v_0$ is fixed, then $z = \sin v_0$ is constant, and $x = \sin u$, $y = (\sin v_0)\cos u$ describe an ellipse that is contained in the horizontal plane $z = \sin v_0$. If $u = u_0$ is fixed, then $x = \sin u_0$ is constant, and $y = (\cos u_0)\sin v$, $z = \sin v \quad \Rightarrow \quad y = (\cos u_0)z$, so the grid curves are portions of lines through the x-axis contained in the plane $x = \sin u_0$ (parallel to the yz-plane). The surface is graph II.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. From Example 3, parametric equations for the plane through the point $(0, 0, 0)$ that contains the vectors $\mathbf{a} = \langle 1, -1, 0\rangle$ and $\mathbf{b} = \langle 0, 1, -1\rangle$ are $x = 0 + u(1) + v(0) = u,\ y = 0 + u(-1) + v(1) = v - u,\ z = 0 + u(0) + v(-1) = -v$.

20. From Example 3, parametric equations for the plane through the point $(0, -1, 5)$ that contains the vectors $\mathbf{a} = \langle 2, 1, 4\rangle$ and $\mathbf{b} = \langle -3, 2, 5\rangle$ are $x = 0 + u(2) + v(-3) = 2u - 3v,\ y = -1 + u(1) + v(2) = -1 + u + 2v$, $z = 5 + u(4) + v(5) = 5 + 4u + 5v$.

21. Solving the equation $4x^2 - 4y^2 - z^2 = 4$ for x gives $x^2 = 1 + y^2 + \frac{1}{4}z^2 \quad \Rightarrow \quad x = \sqrt{1 + y^2 + \frac{1}{4}z^2}$. (We choose the positive root since we want the part of the hyperboloid that corresponds to $x \geq 0$.) If we let y and z be the parameters, parametric equations are $y = y,\ z = z,\ x = \sqrt{1 + y^2 + \frac{1}{4}z^2}$.

22. Solving the equation $x^2 + 2y^2 + 3z^2 = 1$ for y gives $y^2 = \frac{1}{2}(1 - x^2 - 3z^2) \quad \Rightarrow \quad y = -\sqrt{\frac{1}{2}(1 - x^2 - 3z^2)}$ (since we want the part of the ellipsoid that corresponds to $y \leq 0$). If we let x and z be the parameters, parametric equations are $x = x,\ z = z,\ y = -\sqrt{\frac{1}{2}(1 - x^2 - 3z^2)}$.

Alternate solution: The equation can be rewritten as $x^2 + \dfrac{y^2}{\left(1/\sqrt{2}\right)^2} + \dfrac{z^2}{\left(1/\sqrt{3}\right)^2} = 1$, and if we let $x = u\cos v$ and $z = \dfrac{1}{\sqrt{3}}\,u\sin v$, then $y = -\sqrt{\frac{1}{2}(1 - x^2 - 3z^2)} = -\sqrt{\frac{1}{2}(1 - u^2\cos^2 v - u^2\sin^2 v)} = -\sqrt{\frac{1}{2}(1 - u^2)}$, where $0 \leq u \leq 1$ and $0 \leq v \leq 2\pi$.

Second alternate solution: We can adapt the formulas for converting from spherical to rectangular coordinates as follows. We let $x = \sin\phi\cos\theta,\ y = \dfrac{1}{\sqrt{2}}\sin\phi\sin\theta,\ z = \dfrac{1}{\sqrt{3}}\cos\phi$; the surface is generated for $0 \leq \phi \leq \pi,\ \pi \leq \theta \leq 2\pi$.

23. Since the cone $z = \sqrt{x^2 + y^2}$ intersects the sphere $x^2 + y^2 + z^2 = 4$ in the circle $x^2 + y^2 = 2$, $z = \sqrt{2}$ and we want the portion of the sphere above this, we can parametrize the surface as $x = x$, $y = y$, $z = \sqrt{4 - x^2 - y^2}$ where $x^2 + y^2 \leq 2$.
Alternate solution: Using spherical coordinates, $x = 2\sin\phi\cos\theta$, $y = 2\sin\phi\sin\theta$, $z = 2\cos\phi$ where $0 \leq \phi \leq \frac{\pi}{4}$ and $0 \leq \theta \leq 2\pi$.

24. We can parametrize the cylinder as $x = 3\cos\theta,\ y = y,\ z = 3\sin\theta$. To restrict the surface to that portion above the xy-plane and between the planes $y = -4$ and $y = 4$ we require $0 \leq \theta \leq \pi$, $-4 \leq y \leq 4$.

25. In spherical coordinates, parametric equations are $x = 6\sin\phi\cos\theta$, $y = 6\sin\phi\sin\theta$, $z = 6\cos\phi$. The intersection of the sphere with the plane $z = 3\sqrt{3}$ corresponds to $z = 6\cos\phi = 3\sqrt{3} \quad \Rightarrow \quad \cos\phi = \frac{\sqrt{3}}{2} \quad \Rightarrow \quad \phi = \frac{\pi}{6}$, and the plane $z = 0$ (the xy-plane) corresponds to $\phi = \frac{\pi}{2}$. Thus the surface is described by $\frac{\pi}{6} \leq \phi \leq \frac{\pi}{2},\ 0 \leq \theta \leq 2\pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. Using x and y as the parameters, $x = x$, $y = y$, $z = x + 3$ where $0 \le x^2 + y^2 \le 1$. Also, since the plane intersects the cylinder in an ellipse, the surface is a planar ellipse in the plane $z = x + 3$. Thus, parametrizing with respect to s and θ, we have $x = s\cos\theta$, $y = s\sin\theta$, $z = 3 + s\cos\theta$ where $0 \le s \le 1$ and $0 \le \theta \le 2\pi$.

27. The surface appears to be a portion of a circular cylinder of radius 3 with axis the x-axis. An equation of the cylinder is $y^2 + z^2 = 9$, and we can impose the restrictions $0 \le x \le 5$, $y \le 0$ to obtain the portion shown. To graph the surface on a CAS, we can use parametric equations $x = u$, $y = 3\cos v$, $z = 3\sin v$ with the parameter domain $0 \le u \le 5$, $\frac{\pi}{2} \le v \le \frac{3\pi}{2}$. Alternatively, we can regard x and z as parameters. Then parametric equations are $x = x$, $z = z$, $y = -\sqrt{9 - z^2}$, where $0 \le x \le 5$ and $-3 \le z \le 3$.

28. The surface appears to be a portion of a sphere of radius 1 centered at the origin. In spherical coordinates, the sphere has equation $\rho = 1$, and imposing the restrictions $\frac{\pi}{2} \le \theta \le 2\pi$, $\frac{\pi}{4} \le \phi \le \pi$ will give the portion of the sphere shown. Thus, to graph the surface on a CAS we can either use spherical coordinates with the stated restrictions, or we can use parametric equations: $x = \sin\phi\cos\theta$, $y = \sin\phi\sin\theta$, $z = \cos\phi$, $\frac{\pi}{2} \le \theta \le 2\pi$, $\frac{\pi}{4} \le \phi \le \pi$.

29. Using Equations 3, we have the parametrization $x = x$,

$$y = \frac{1}{1+x^2}\cos\theta, \quad z = \frac{1}{1+x^2}\sin\theta, \quad -2 \le x \le 2, \quad 0 \le \theta \le 2\pi.$$

30. Letting θ be the angle of rotation about the y-axis (adapting Equations 3), we have the parametrization $x = (1/y)\cos\theta$, $y = y$, $z = (1/y)\sin\theta$, $y \ge 1$, $0 \le \theta \le 2\pi$.

31. (a) Replacing $\cos u$ by $\sin u$ and $\sin u$ by $\cos u$ gives parametric equations $x = (2 + \sin v)\sin u$, $y = (2 + \sin v)\cos u$, $z = u + \cos v$. From the graph, it appears that the direction of the spiral is reversed. We can verify this observation by noting that the projection of the spiral grid curves onto the xy-plane, given by $x = (2 + \sin v)\sin u$, $y = (2 + \sin v)\cos u$, $z = 0$, draws a circle in the clockwise direction for each value of v. The original equations, on the other hand, give circular projections drawn in the counterclockwise direction. The equation for z is identical in both surfaces, so as z increases, these grid curves spiral up in opposite directions for the two surfaces.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Replacing $\cos u$ by $\cos 2u$ and $\sin u$ by $\sin 2u$ gives parametric equations $x = (2 + \sin v)\cos 2u$, $y = (2 + \sin v)\sin 2u$, $z = u + \cos v$. From the graph, it appears that the number of coils in the surface doubles within the same parametric domain. We can verify this observation by noting that the projection of the spiral grid curves onto the xy-plane, given by $x = (2 + \sin v)\cos 2u$, $y = (2 + \sin v)\sin 2u$, $z = 0$ (where v is constant), complete circular revolutions for $0 \le u \le \pi$ while the original surface requires $0 \le u \le 2\pi$ for a complete revolution. Thus, the new surface winds around twice as fast as the original surface, and since the equation for z is identical in both surfaces, we observe twice as many circular coils in the same z-interval.

32. First we graph the surface as viewed from the front, then from two additional viewpoints.

The surface appears as a twisted sheet, and is unusual because it has only one side. (The Möbius strip is discussed in more detail in Section 16.7.)

33. $\mathbf{r}(u, v) = (u + v)\,\mathbf{i} + 3u^2\,\mathbf{j} + (u - v)\,\mathbf{k}$.

$\mathbf{r}_u = \mathbf{i} + 6u\,\mathbf{j} + \mathbf{k}$ and $\mathbf{r}_v = \mathbf{i} - \mathbf{k}$, so $\mathbf{r}_u \times \mathbf{r}_v = -6u\,\mathbf{i} + 2\,\mathbf{j} - 6u\,\mathbf{k}$. Since the point $(2, 3, 0)$ corresponds to $u = 1$, $v = 1$, a normal vector to the surface at $(2, 3, 0)$ is $-6\,\mathbf{i} + 2\,\mathbf{j} - 6\,\mathbf{k}$, and an equation of the tangent plane is $-6x + 2y - 6z = -6$ or $3x - y + 3z = 3$.

34. $\mathbf{r}(u, v) = (u^2 + 1)\,\mathbf{i} + (v^3 + 1)\,\mathbf{j} + (u + v)\,\mathbf{k}$.

$\mathbf{r}_u = 2u\,\mathbf{i} + \mathbf{k}$ and $\mathbf{r}_v = 3v^2\,\mathbf{j} + \mathbf{k}$, so $\mathbf{r}_u \times \mathbf{r}_v = -3v^2\,\mathbf{i} - 2u\,\mathbf{j} + 6uv^2\,\mathbf{k}$. Since the point $(5, 2, 3)$ corresponds to $u = 2$, $v = 1$, a normal vector to the surface at $(5, 2, 3)$ is $-3\,\mathbf{i} - 4\,\mathbf{j} + 12\,\mathbf{k}$, and an equation of the tangent plane is $-3(x - 5) - 4(y - 2) + 12(z - 3) = 0$ or $3x + 4y - 12z = -13$.

35. $\mathbf{r}(u, v) = u\cos v\,\mathbf{i} + u\sin v\,\mathbf{j} + v\,\mathbf{k} \quad \Rightarrow \quad \mathbf{r}\left(1, \frac{\pi}{3}\right) = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}, \frac{\pi}{3}\right)$.

$\mathbf{r}_u = \cos v\,\mathbf{i} + \sin v\,\mathbf{j}$ and $\mathbf{r}_v = -u\sin v\,\mathbf{i} + u\cos v\,\mathbf{j} + \mathbf{k}$, so a normal vector to the surface at the point $\left(\frac{1}{2}, \frac{\sqrt{3}}{2}, \frac{\pi}{3}\right)$ is $\mathbf{r}_u\left(1, \frac{\pi}{3}\right) \times \mathbf{r}_v\left(1, \frac{\pi}{3}\right) = \left(\frac{1}{2}\,\mathbf{i} + \frac{\sqrt{3}}{2}\,\mathbf{j}\right) \times \left(-\frac{\sqrt{3}}{2}\,\mathbf{i} + \frac{1}{2}\,\mathbf{j} + \mathbf{k}\right) = \frac{\sqrt{3}}{2}\,\mathbf{i} - \frac{1}{2}\,\mathbf{j} + \mathbf{k}$. Thus an equation of the tangent plane at $\left(\frac{1}{2}, \frac{\sqrt{3}}{2}, \frac{\pi}{3}\right)$ is $\frac{\sqrt{3}}{2}\left(x - \frac{1}{2}\right) - \frac{1}{2}\left(y - \frac{\sqrt{3}}{2}\right) + 1\left(z - \frac{\pi}{3}\right) = 0$ or $\frac{\sqrt{3}}{2}x - \frac{1}{2}y + z = \frac{\pi}{3}$.

36. $\mathbf{r}(u, v) = \sin u\,\mathbf{i} + \cos u\,\sin v\,\mathbf{j} + \sin v\,\mathbf{k} \quad \Rightarrow \quad \mathbf{r}\left(\frac{\pi}{6}, \frac{\pi}{6}\right) = \left(\frac{1}{2}, \frac{\sqrt{3}}{4}, \frac{1}{2}\right)$.

$\mathbf{r}_u = \cos u\,\mathbf{i} - \sin u\sin v\,\mathbf{j}$ and $\mathbf{r}_v = \cos u\cos v\,\mathbf{j} + \cos v\,\mathbf{k}$, so a normal vector to the surface at the point $\left(\frac{1}{2}, \frac{\sqrt{3}}{4}, \frac{1}{2}\right)$ is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\mathbf{r}_u\left(\frac{\pi}{6}, \frac{\pi}{6}\right) \times \mathbf{r}_v\left(\frac{\pi}{6}, \frac{\pi}{6}\right) = \left(\frac{\sqrt{3}}{2}\,\mathbf{i} - \frac{1}{4}\,\mathbf{j}\right) \times \left(\frac{3}{4}\,\mathbf{j} + \frac{\sqrt{3}}{2}\,\mathbf{k}\right) = -\frac{\sqrt{3}}{8}\,\mathbf{i} - \frac{3}{4}\,\mathbf{j} + \frac{3\sqrt{3}}{8}\,\mathbf{k}.$

Thus an equation of the tangent plane at $\left(\frac{1}{2}, \frac{\sqrt{3}}{4}, \frac{1}{2}\right)$ is $-\frac{\sqrt{3}}{8}\left(x - \frac{1}{2}\right) - \frac{3}{4}\left(y - \frac{\sqrt{3}}{4}\right) + \frac{3\sqrt{3}}{8}\left(z - \frac{1}{2}\right) = 0$ or

$\sqrt{3}\,x + 6y - 3\sqrt{3}\,z = \frac{\sqrt{3}}{2}$ or $2x + 4\sqrt{3}\,y - 6z = 1$.

37. $\mathbf{r}(u, v) = u^2\,\mathbf{i} + 2u \sin v\,\mathbf{j} + u \cos v\,\mathbf{k} \quad \Rightarrow \quad \mathbf{r}(1, 0) = (1, 0, 1)$.

$\mathbf{r}_u = 2u\,\mathbf{i} + 2 \sin v\,\mathbf{j} + \cos v\,\mathbf{k}$ and $\mathbf{r}_v = 2u \cos v\,\mathbf{j} - u \sin v\,\mathbf{k}$,

so a normal vector to the surface at the point $(1, 0, 1)$ is

$\mathbf{r}_u(1, 0) \times \mathbf{r}_v(1, 0) = (2\,\mathbf{i} + \mathbf{k}) \times (2\,\mathbf{j}) = -2\,\mathbf{i} + 4\,\mathbf{k}$.

Thus an equation of the tangent plane at $(1, 0, 1)$ is

$-2(x - 1) + 0(y - 0) + 4(z - 1) = 0$ or $-x + 2z = 1$.

38. $\mathbf{r}(u, v) = (1 - u^2 - v^2)\,\mathbf{i} - v\,\mathbf{j} - u\,\mathbf{k}$.

$\mathbf{r}_u = -2u\,\mathbf{i} - \mathbf{k}$ and $\mathbf{r}_v = -2v\,\mathbf{i} - \mathbf{j}$. Since the point $(-1, -1, -1)$ corresponds to $u = 1$, $v = 1$, a normal vector to the surface at $(-1, -1, -1)$ is

$\mathbf{r}_u(1, 1) \times \mathbf{r}_v(1, 1) = (-2\,\mathbf{i} - \mathbf{k}) \times (-2\,\mathbf{i} - \mathbf{j}) = -\mathbf{i} + 2\,\mathbf{j} + 2\,\mathbf{k}$.

Thus an equation of the tangent plane is $-1(x + 1) + 2(y + 1) + 2(z + 1) = 0$ or $-x + 2y + 2z = -3$.

39. The surface S is given by $z = f(x, y) = 6 - 3x - 2y$ which intersects the xy-plane in the line $3x + 2y = 6$, so D is the triangular region given by $\left\{(x, y) \;\middle|\; 0 \le x \le 2,\ 0 \le y \le 3 - \frac{3}{2}x\right\}$. By Formula 9, the surface area of S is

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}\, dA \\ &= \iint_D \sqrt{1 + (-3)^2 + (-2)^2}\, dA = \sqrt{14} \iint_D dA = \sqrt{14}\, A(D) = \sqrt{14}\left(\tfrac{1}{2} \cdot 2 \cdot 3\right) = 3\sqrt{14}. \end{aligned}$$

40. $\mathbf{r}(u, v) = \langle u + v, 2 - 3u, 1 + u - v \rangle \quad \Rightarrow \quad \mathbf{r}_u = \langle 1, -3, 1 \rangle$, $\mathbf{r}_v = \langle 1, 0, -1 \rangle$, and $\mathbf{r}_u \times \mathbf{r}_v = \langle 3, 2, 3 \rangle$. Then by Definition 6,

$$A(S) = \iint_D |\,\mathbf{r}_u \times \mathbf{r}_v\,|\; dA = \int_0^2 \int_{-1}^1 |\,\langle 3, 2, 3 \rangle\,|\; dv\, du = \sqrt{22} \int_0^2 du \int_{-1}^1 dv = \sqrt{22}\,(2)(2) = 4\sqrt{22}$$

41. Here we can write $z = f(x, y) = \frac{1}{3} - \frac{1}{3}x - \frac{2}{3}y$ and D is the disk $x^2 + y^2 \le 3$, so by Formula 9 the area of the surface is

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}\, dA = \iint_D \sqrt{1 + \left(-\tfrac{1}{3}\right)^2 + \left(-\tfrac{2}{3}\right)^2}\, dA = \tfrac{\sqrt{14}}{3} \iint_D dA \\ &= \tfrac{\sqrt{14}}{3}\, A(D) = \tfrac{\sqrt{14}}{3} \cdot \pi\left(\sqrt{3}\right)^2 = \sqrt{14}\,\pi \end{aligned}$$

42. $z = f(x, y) = \sqrt{x^2 + y^2} \quad \Rightarrow \quad \dfrac{\partial z}{\partial x} = \dfrac{1}{2}\left(x^2 + y^2\right)^{-1/2} \cdot 2x = \dfrac{x}{\sqrt{x^2 + y^2}}$, $\dfrac{\partial z}{\partial y} = \dfrac{y}{\sqrt{x^2 + y^2}}$, and

$$\sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} = \sqrt{1 + \frac{x^2}{x^2 + y^2} + \frac{y^2}{x^2 + y^2}} = \sqrt{1 + \frac{x^2 + y^2}{x^2 + y^2}} = \sqrt{2}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Here D is given by $\{(x,y)\ \big|\ 0 \le x \le 1,\ x^2 \le y \le x\}$, so by Formula 9, the surface area of S is

$$A(S) = \iint_D \sqrt{2}\,dA = \int_0^1 \int_{x^2}^{x} \sqrt{2}\,dy\,dx = \sqrt{2}\int_0^1 \left(x - x^2\right) dx = \sqrt{2}\ \left[\tfrac{1}{2}x^2 - \tfrac{1}{3}x^3\right]_0^1 = \sqrt{2}\left(\tfrac{1}{2} - \tfrac{1}{3}\right) = \tfrac{\sqrt{2}}{6}$$

43. $z = f(x,y) = \frac{2}{3}(x^{3/2} + y^{3/2})$ and $D = \{(x,y)\,|\,0 \le x \le 1,\ 0 \le y \le 1\}$. Then $f_x = x^{1/2}$, $f_y = y^{1/2}$ and

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + \left(\sqrt{x}\right)^2 + \left(\sqrt{y}\right)^2}\,dA = \int_0^1 \int_0^1 \sqrt{1 + x + y}\,dy\,dx \\ &= \int_0^1 \left[\tfrac{2}{3}(x + y + 1)^{3/2}\right]_{y=0}^{y=1} dx = \tfrac{2}{3}\int_0^1 \left[(x+2)^{3/2} - (x+1)^{3/2}\right] dx \\ &= \tfrac{2}{3}\left[\tfrac{2}{5}(x+2)^{5/2} - \tfrac{2}{5}(x+1)^{5/2}\right]_0^1 = \tfrac{4}{15}(3^{5/2} - 2^{5/2} - 2^{5/2} + 1) = \tfrac{4}{15}(3^{5/2} - 2^{7/2} + 1) \end{aligned}$$

44. $z = f(x,y) = 4 - 2x^2 + y$ and $D = \{(x,y)\,|\,0 \le x \le 1,\ 0 \le y \le x\}$. Thus, by Formula 9,

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + (-4x)^2 + (1)^2}\,dA = \int_0^1 \int_0^x \sqrt{16x^2 + 2}\,dy\,dx = \int_0^1 x\sqrt{16x^2 + 2}\,dx \\ &= \tfrac{1}{32} \cdot \tfrac{2}{3}(16x^2 + 2)^{3/2}\Big]_0^1 = \tfrac{1}{48}(18^{3/2} - 2^{3/2}) = \tfrac{1}{48}(54\sqrt{2} - 2\sqrt{2}) = \tfrac{13}{12}\sqrt{2} \end{aligned}$$

45. $z = f(x,y) = xy$ with $x^2 + y^2 \le 1$, so $f_x = y$, $f_y = x \quad \Rightarrow$

$$\begin{aligned} A(S) &= \iint_D \sqrt{1 + y^2 + x^2}\,dA = \int_0^{2\pi}\int_0^1 \sqrt{r^2 + 1}\,r\,dr\,d\theta = \int_0^{2\pi} \left[\tfrac{1}{3}\left(r^2 + 1\right)^{3/2}\right]_{r=0}^{r=1} d\theta \\ &= \int_0^{2\pi} \tfrac{1}{3}\left(2\sqrt{2} - 1\right) d\theta = \tfrac{2\pi}{3}\left(2\sqrt{2} - 1\right) \end{aligned}$$

46. A parametric representation of the surface is $x = z^2 + y$, $y = y$, $z = z$ with $0 \le y \le 2,\ 0 \le z \le 2$.

Hence $\mathbf{r}_y \times \mathbf{r}_z = (\mathbf{i} + \mathbf{j}) \times (2z\,\mathbf{i} + \mathbf{k}) = \mathbf{i} - \mathbf{j} - 2z\,\mathbf{k}$. Then

$$\begin{aligned} A(S) &= \iint_D |\,\mathbf{r}_y \times \mathbf{r}_z\,|\ dA = \int_0^2\int_0^2 \sqrt{1 + 1 + 4z^2}\,dy\,dz = \int_0^2 2\sqrt{2 + 4z^2}\,dz \\ &= \left[2 \cdot \tfrac{1}{2}\left(z\sqrt{2 + 4z^2} + \ln\left(2z + \sqrt{2 + 4z^2}\right)\right)\right]_0^2 \quad \left[\begin{array}{c}\text{Use trigonometric substitution} \\ \text{or Formula 21 in the Table of Integrals}\end{array}\right] \\ &= 6\sqrt{2} + \ln\left(4 + 3\sqrt{2}\right) - \ln\sqrt{2} \ \text{ or } \ 6\sqrt{2} + \ln\frac{4 + 3\sqrt{2}}{\sqrt{2}} = 6\sqrt{2} + \ln\left(2\sqrt{2} + 3\right) \end{aligned}$$

Note: In general, if $x = f(y,z)$ then $\mathbf{r}_y \times \mathbf{r}_z = \mathbf{i} - \dfrac{\partial f}{\partial y}\,\mathbf{j} - \dfrac{\partial f}{\partial z}\,\mathbf{k}$ and $A\,(S) = \displaystyle\iint_D \sqrt{1 + \left(\frac{\partial f}{\partial y}\right)^2 + \left(\frac{\partial f}{\partial z}\right)^2}\,dA$.

47. A parametric representation of the surface is $x = x$, $y = x^2 + z^2$, $z = z$ with $0 \le x^2 + z^2 \le 16$.

Hence $\mathbf{r}_x \times \mathbf{r}_z = (\mathbf{i} + 2x\,\mathbf{j}) \times (2z\,\mathbf{j} + \mathbf{k}) = 2x\,\mathbf{i} - \mathbf{j} + 2z\,\mathbf{k}$.

Note: In general, if $y = f(x,z)$ then $\mathbf{r}_x \times \mathbf{r}_z = \dfrac{\partial f}{\partial x}\,\mathbf{i} - \mathbf{j} + \dfrac{\partial f}{\partial z}\,\mathbf{k}$, and $A(S) = \displaystyle\iint_D \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial z}\right)^2}\,dA$.

Then

$$\begin{aligned} A(S) &= \iint_{0 \le x^2 + z^2 \le 16} \sqrt{1 + 4x^2 + 4z^2}\,dA = \int_0^{2\pi}\int_0^4 \sqrt{1 + 4r^2}\ r\,dr\,d\theta \\ &= \int_0^{2\pi} d\theta \int_0^4 r\sqrt{1 + 4r^2}\,dr = 2\pi\left[\tfrac{1}{12}(1 + 4r^2)^{3/2}\right]_0^4 = \tfrac{\pi}{6}\left(65^{3/2} - 1\right) \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

48. $\mathbf{r}(u,v) = u\cos v\,\mathbf{i} + u\sin v\,\mathbf{j} + v\,\mathbf{k}$, $0 \le u \le 1$, $0 \le v \le \pi \quad \Rightarrow$

$\mathbf{r}_u = \langle \cos v, \sin v, 0\rangle$, $\mathbf{r}_v = \langle -u\sin v, u\cos v, 1\rangle$, and $\mathbf{r}_u \times \mathbf{r}_v = \langle \sin v, -\cos v, u\rangle$. Then

$$A(S) = \int_0^\pi \int_0^1 \sqrt{1+u^2}\,du\,dv = \int_0^\pi dv \int_0^1 \sqrt{1+u^2}\,du$$
$$= \pi\left[\tfrac{u}{2}\sqrt{u^2+1} + \tfrac{1}{2}\ln\left|u+\sqrt{u^2+1}\,\right|\right]_0^1 = \tfrac{\pi}{2}\left[\sqrt{2} + \ln\left(1+\sqrt{2}\,\right)\right]$$

49. $x = u^2$, $y = uv$, $z = \frac{1}{2}v^2$, $0 \le u \le 1$, $0 \le v \le 2 \quad \Rightarrow \quad \mathbf{r}_u = \langle 2u, v, 0\rangle$, $\mathbf{r}_v = \langle 0, u, v\rangle$, and $\mathbf{r}_u \times \mathbf{r}_v = \left\langle v^2, -2uv, 2u^2\right\rangle$.

Then

$$A(S) = \iint_D |\mathbf{r}_u \times \mathbf{r}_v|\,dA = \int_0^1\int_0^2 \sqrt{v^4 + 4u^2v^2 + 4u^4}\,dv\,du = \int_0^1\int_0^2 \sqrt{(v^2+2u^2)^2}\,dv\,du$$
$$= \int_0^1\int_0^2 (v^2+2u^2)\,dv\,du = \int_0^1 \left[\tfrac{1}{3}v^3 + 2u^2v\right]_{v=0}^{v=2} du = \int_0^1 \left(\tfrac{8}{3} + 4u^2\right) du = \left[\tfrac{8}{3}u + \tfrac{4}{3}u^3\right]_0^1 = 4$$

50. The cylinder encloses separate portions of the sphere in the upper and lower halves. The top half of the sphere is $z = f(x,y) = \sqrt{b^2 - x^2 - y^2}$ and D is given by $\left\{(x,y)\ \middle|\ x^2+y^2 \le a^2\right\}$. By Formula 9, the surface area of the upper enclosed portion is

$$A = \iint_D \sqrt{1 + \left(\frac{-x}{\sqrt{b^2-x^2-y^2}}\right)^2 + \left(\frac{-y}{\sqrt{b^2-x^2-y^2}}\right)^2}\,dA = \iint_D \sqrt{1 + \frac{x^2+y^2}{b^2-x^2-y^2}}\,dA$$
$$= \iint_D \sqrt{\frac{b^2}{b^2-x^2-y^2}}\,dA = \int_0^{2\pi}\int_0^a \frac{b}{\sqrt{b^2-r^2}}\,r\,dr\,d\theta = b\int_0^{2\pi} d\theta \int_0^a \frac{r}{\sqrt{b^2-r^2}}\,dr$$
$$= b\left[\theta\right]_0^{2\pi}\left[-\sqrt{b^2-r^2}\,\right]_0^a = 2\pi b\left(-\sqrt{b^2-a^2} + \sqrt{b^2-0}\,\right) = 2\pi b\left(b - \sqrt{b^2-a^2}\,\right)$$

The lower portion of the sphere enclosed by the cylinder has identical shape, so the total area is $2A = 4\pi b\left(b - \sqrt{b^2-a^2}\,\right)$.

51. From Formula 9 with $z = f(x,y)$, we have $A(S) = \iint_D \sqrt{1+(f_x)^2+(f_y)^2}\,dA$. Since $|f_x| \le 1$ and $|f_y| \le 1$, we know $0 \le (f_x)^2 \le 1$ and $0 \le (f_y)^2 \le 1$, so $1 \le 1 + (f_x)^2 + (f_y)^2 \le 3 \quad \Rightarrow \quad 1 \le \sqrt{1+(f_x)^2+(f_y)^2} \le \sqrt{3}$. By Property 15.2.10, $\iint_D 1\,dA \le \iint_D \sqrt{1+(f_x)^2+(f_y)^2}\,dA \le \iint_D \sqrt{3}\,dA \quad \Rightarrow \quad A(D) \le A(S) \le \sqrt{3}\,A(D) \quad \Rightarrow$ $\pi R^2 \le A(S) \le \sqrt{3}\pi R^2$.

52. $z = f(x,y) = \cos(x^2+y^2)$ with $x^2+y^2 \le 1$.

$$A(S) = \iint_D \sqrt{1 + (-2x\sin(x^2+y^2))^2 + (-2y\sin(x^2+y^2))^2}\,dA$$
$$= \iint_D \sqrt{1 + 4x^2\sin^2(x^2+y^2) + 4y^2\sin^2(x^2+y^2)}\,dA = \iint_D \sqrt{1 + 4(x^2+y^2)\sin^2(x^2+y^2)}\,dA$$
$$= \int_0^{2\pi}\int_0^1 \sqrt{1+4r^2\sin^2(r^2)}\,r\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^1 r\sqrt{1+4r^2\sin^2(r^2)}\,dr$$
$$= 2\pi\int_0^1 r\sqrt{1+4r^2\sin^2(r^2)}\,dr \approx 4.1073$$

53. $z = f(x,y) = \ln(x^2+y^2+2)$ with $x^2+y^2 \le 1$.

$$A(S) = \iint_D \sqrt{1 + \left(\frac{2x}{x^2+y^2+2}\right)^2 + \left(\frac{2y}{x^2+y^2+2}\right)^2}\,dA = \iint_D \sqrt{1 + \frac{4x^2+4y^2}{(x^2+y^2+2)^2}}\,dA$$
$$= \int_0^{2\pi}\int_0^1 \sqrt{1 + \frac{4r^2}{(r^2+2)^2}}\,r\,dr\,d\theta = \int_0^{2\pi} d\theta \int_0^1 r\sqrt{\frac{(r^2+2)^2+4r^2}{(r^2+2)^2}}\,dr = 2\pi\int_0^1 \frac{r\sqrt{r^4+8r^2+4}}{r^2+2}\,dr$$
$$\approx 3.5618$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. Let $f(x,y) = \dfrac{1+x^2}{1+y^2}$. Then $f_x = \dfrac{2x}{1+y^2}$,

$f_y = (1+x^2)\left[-\dfrac{2y}{(1+y^2)^2}\right] = -\dfrac{2y(1+x^2)}{(1+y^2)^2}$.

We use a CAS to estimate

$\int_{-1}^{1}\int_{-(1-|x|)}^{1-|x|} \sqrt{1+f_x^2+f_y^2}\,dy\,dx \approx 2.6959$.

In order to graph only the part of the surface above the square, we use $-(1-|x|) \le y \le 1-|x|$ as the y-range in our plot command.

55. (a) $z = \dfrac{1}{1+x^2+y^2} \quad\Rightarrow\quad \dfrac{\partial z}{\partial x} = \dfrac{-2x}{(1+x^2+y^2)^2}$ and $\dfrac{\partial z}{\partial y} = \dfrac{-2y}{(1+x^2+y^2)^2}$.

$$A(S) = \iint_D \sqrt{1+\left(\frac{\partial z}{\partial x}\right)^2+\left(\frac{\partial z}{\partial y}\right)^2}\,dA = \int_0^6\int_0^4 \sqrt{1+\frac{4x^2+4y^2}{(1+x^2+y^2)^4}}\,dy\,dx.$$

Using the Midpoint Rule with $f(x,y) = \sqrt{1+\dfrac{4x^2+4y^2}{(1+x^2+y^2)^4}}$, $m=3$, $n=2$ we have

$$A(S) \approx \sum_{i=1}^{3}\sum_{j=1}^{2} f\left(\overline{x}_i,\overline{y}_j\right)\Delta A = 4\left[f(1,1)+f(1,3)+f(3,1)+f(3,3)+f(5,1)+f(5,3)\right] \approx 24.2055$$

(b) Using a CAS, we have $A(S) = \displaystyle\int_0^6\int_0^4 \sqrt{1+\frac{4x^2+4y^2}{(1+x^2+y^2)^4}}\,dy\,dx \approx 24.2476$. This agrees with the estimate in part (a) to the first decimal place.

56. $\mathbf{r}(u,v) = \left\langle \cos^3 u\,\cos^3 v, \sin^3 u\,\cos^3 v, \sin^3 v\right\rangle$, so $\mathbf{r}_u = \left\langle -3\cos^2 u\,\sin u\cos^3 v, 3\sin^2 u\,\cos u\cos^3 v, 0\right\rangle$,

$\mathbf{r}_v = \left\langle -3\cos^3 u\,\cos^2 v\,\sin v, -3\sin^3 u\,\cos^2 v\,\sin v, 3\sin^2 v\,\cos v\right\rangle$, and

$\mathbf{r}_u \times \mathbf{r}_v = \left\langle 9\cos u\,\sin^2 u\,\cos^4 v\,\sin^2 v, 9\cos^2 u\,\sin u\,\cos^4 v\,\sin^2 v, 9\cos^2 u\,\sin^2 u\,\cos^5 v\,\sin v\right\rangle$. Then

$$\begin{aligned}|\mathbf{r}_u \times \mathbf{r}_v| &= 9\sqrt{\cos^2 u\,\sin^4 u\,\cos^8 v\,\sin^4 v + \cos^4 u\,\sin^2 u\,\cos^8 v\,\sin^4 v + \cos^4 u\,\sin^4 u\,\cos^{10} v\,\sin^2 v}\\ &= 9\sqrt{\cos^2 u\,\sin^2 u\,\cos^8 v\,\sin^2 v\,(\sin^2 v + \cos^2 u\,\sin^2 u\,\cos^2 v)}\\ &= 9\cos^4 v\,|\cos u\,\sin u\,\sin v|\sqrt{\sin^2 v + \cos^2 u\,\sin^2 u\,\cos^2 v}\end{aligned}$$

Using a CAS, we have $A(S) = \int_0^{\pi}\int_0^{2\pi} 9\cos^4 v\,|\cos u\,\sin u\,\sin v|\sqrt{\sin^2 v + \cos^2 u\,\sin^2 u\,\cos^2 v}\,dv\,du \approx 4.4506$.

57. $z = 1+2x+3y+4y^2$, so

$$A(S) = \iint_D \sqrt{1+\left(\frac{\partial z}{\partial x}\right)^2+\left(\frac{\partial z}{\partial y}\right)^2}\,dA = \int_1^4\int_0^1 \sqrt{1+4+(3+8y)^2}\,dy\,dx = \int_1^4\int_0^1 \sqrt{14+48y+64y^2}\,dy\,dx.$$

Using a CAS, we have

$\int_1^4\int_0^1 \sqrt{14+48y+64y^2}\,dy\,dx = \frac{45}{8}\sqrt{14} + \frac{15}{16}\ln\left(11\sqrt{5}+3\sqrt{14}\sqrt{5}\right) - \frac{15}{16}\ln\left(3\sqrt{5}+\sqrt{14}\sqrt{5}\right)$

or $\frac{45}{8}\sqrt{14} + \frac{15}{16}\ln\frac{11\sqrt{5}+3\sqrt{70}}{3\sqrt{5}+\sqrt{70}}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

58. (a) $x = au\cos v$, $y = bu\sin v$, $z = u^2$, $0 \le u \le 2$, $0 \le v \le 2\pi \quad \Rightarrow \quad \mathbf{r}_u = a\cos v\,\mathbf{i} + b\sin v\,\mathbf{j} + 2u\,\mathbf{k}$,

$\mathbf{r}_v = -au\sin v\,\mathbf{i} + bu\cos v\,\mathbf{j} + 0\,\mathbf{k}$, and $\mathbf{r}_u \times \mathbf{r}_v = -2bu^2\cos v\,\mathbf{i} - 2au^2\sin v\,\mathbf{j} + abu\,\mathbf{k}$.

$A(S) = \int_0^{2\pi}\int_0^2 |\mathbf{r}_u \times \mathbf{r}_v|\,du\,dv = \int_0^{2\pi}\int_0^2 \sqrt{4b^2u^4\cos^2 v + 4a^2u^4\sin^2 v + a^2b^2u^2}\,du\,dv$

(b) $x^2 = a^2u^2\cos^2 v$, $y^2 = b^2u^2\sin^2 v$, $z = u^2 \quad \Rightarrow \quad x^2/a^2 + y^2/b^2 = u^2 = z$ which is an elliptic paraboloid. To find D, notice that $0 \le u \le 2 \quad \Rightarrow \quad 0 \le z \le 4 \quad \Rightarrow \quad 0 \le x^2/a^2 + y^2/b^2 \le 4$. Therefore, using Formula 9, we have

$$A(S) = \int_{-2a}^{2a}\int_{-b\sqrt{4-(x^2/a^2)}}^{b\sqrt{4-(x^2/a^2)}} \sqrt{1 + (2x/a^2)^2 + (2y/b^2)^2}\,dy\,dx.$$

(c)

(d) We substitute $a = 2$, $b = 3$ in the integral in part (a) to get $A(S) = \int_0^{2\pi}\int_0^2 2u\sqrt{9u^2\cos^2 v + 4u^2\sin^2 v + 9}\,du\,dv$. We use a CAS to estimate the integral accurate to four decimal places. To speed up the calculation, we can set `Digits:=7;` (in Maple) or use the approximation command `N` (in Mathematica). We find that $A(S) \approx 115.6596$.

59. (a) $x = a\sin u\cos v$, $y = b\sin u\sin v$, $z = c\cos u \quad \Rightarrow$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = (\sin u\cos v)^2 + (\sin u\sin v)^2 + (\cos u)^2$$
$$= \sin^2 u + \cos^2 u = 1$$

and since the ranges of u and v are sufficient to generate the entire graph, the parametric equations represent an ellipsoid.

(b)

(c) From the parametric equations (with $a = 1$, $b = 2$, and $c = 3$), we calculate $\mathbf{r}_u = \cos u\cos v\,\mathbf{i} + 2\cos u\sin v\,\mathbf{j} - 3\sin u\,\mathbf{k}$ and $\mathbf{r}_v = -\sin u\sin v\,\mathbf{i} + 2\sin u\cos v\,\mathbf{j}$. So $\mathbf{r}_u \times \mathbf{r}_v = 6\sin^2 u\cos v\,\mathbf{i} + 3\sin^2 u\sin v\,\mathbf{j} + 2\sin u\cos u\,\mathbf{k}$, and the surface area is given by $A(S) = \int_0^{2\pi}\int_0^{\pi} |\mathbf{r}_u \times \mathbf{r}_v|\,du\,dv = \int_0^{2\pi}\int_0^{\pi} \sqrt{36\sin^4 u\cos^2 v + 9\sin^4 u\sin^2 v + 4\cos^2 u\sin^2 u}\,du\,dv$

60. (a) $x = a\cosh u\,\cos v$, $\;y = b\cosh u\,\sin v$, $\;z = c\sinh u \quad \Rightarrow$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = \cosh^2 u\,\cos^2 v + \cosh^2 u\,\sin^2 v - \sinh^2 u$$
$$= \cosh^2 u - \sinh^2 u = 1$$

and the parametric equations represent a hyperboloid of one sheet.

(b)

(c) $\mathbf{r}_u = \sinh u\,\cos v\,\mathbf{i} + 2\sinh u\,\sin v\,\mathbf{j} + 3\cosh u\,\mathbf{k}$ and $\mathbf{r}_v = -\cosh u\,\sin v\,\mathbf{i} + 2\cosh u\,\cos v\,\mathbf{j}$, so $\mathbf{r}_u \times \mathbf{r}_v = -6\cosh^2 u\,\cos v\,\mathbf{i} - 3\cosh^2 u\,\sin v\,\mathbf{j} + 2\cosh u\,\sinh u\,\mathbf{k}$. We integrate between $u = \sinh^{-1}(-1) = -\ln\left(1 + \sqrt{2}\right)$ and $u = \sinh^{-1} 1 = \ln\left(1 + \sqrt{2}\right)$, since then z varies between

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

-3 and 3, as desired. So the surface area is

$$A(S) = \int_0^{2\pi} \int_{-\ln\left(1+\sqrt{2}\right)}^{\ln\left(1+\sqrt{2}\right)} |\mathbf{r}_u \times \mathbf{r}_v| \, du \, dv$$
$$= \int_0^{2\pi} \int_{-\ln\left(1+\sqrt{2}\right)}^{\ln\left(1+\sqrt{2}\right)} \sqrt{36\cosh^4 u \, \cos^2 v + 9\cosh^4 u \, \sin^2 v + 4\cosh^2 u \, \sinh^2 u} \, du \, dv$$

61. To find the region D: $z = x^2 + y^2$ implies $z + z^2 = 4z$ or $z^2 - 3z = 0$. Thus $z = 0$ or $z = 3$ are the planes where the surfaces intersect. But $x^2 + y^2 + z^2 = 4z$ implies $x^2 + y^2 + (z-2)^2 = 4$, so $z = 3$ intersects the upper hemisphere. Thus $(z-2)^2 = 4 - x^2 - y^2$ or $z = 2 + \sqrt{4 - x^2 - y^2}$. Therefore D is the region inside the circle $x^2 + y^2 + (3-2)^2 = 4$, that is, $D = \left\{(x, y) \mid x^2 + y^2 \le 3\right\}$.

$$A(S) = \iint_D \sqrt{1 + [(-x)(4 - x^2 - y^2)^{-1/2}]^2 + [(-y)(4 - x^2 - y^2)^{-1/2}]^2} \, dA$$
$$= \int_0^{2\pi} \int_0^{\sqrt{3}} \sqrt{1 + \frac{r^2}{4 - r^2}} \, r \, dr \, d\theta = \int_0^{2\pi} \int_0^{\sqrt{3}} \frac{2r \, dr}{\sqrt{4 - r^2}} \, d\theta = \int_0^{2\pi} \left[-2(4 - r^2)^{1/2}\right]_{r=0}^{r=\sqrt{3}} d\theta$$
$$= \int_0^{2\pi} (-2 + 4) \, d\theta = 2\theta\Big]_0^{2\pi} = 4\pi$$

62. We first find the area of the face of the surface that intersects the positive y-axis. A parametric representation of the surface is $x = x, \;\; y = \sqrt{1 - z^2}, \;\; z = z$ with $x^2 + z^2 \le 1$. Then $\mathbf{r}(x, z) = \left\langle x, \sqrt{1 - z^2}, z \right\rangle \;\Rightarrow\; \mathbf{r}_x = \langle 1, 0, 0 \rangle$, $\mathbf{r}_z = \left\langle 0, -z/\sqrt{1 - z^2}, 1 \right\rangle$ and $\mathbf{r}_x \times \mathbf{r}_z = \left\langle 0, -1, -z/\sqrt{1 - z^2} \right\rangle \;\Rightarrow\; |\,\mathbf{r}_x \times \mathbf{r}_z\,| = \sqrt{1 + \dfrac{z^2}{1 - z^2}} = \dfrac{1}{\sqrt{1 - z^2}}$.

$$A(S) = \iint\limits_{x^2+z^2 \le 1} |\,\mathbf{r}_x \times \mathbf{r}_z\,| \, dA = \int_{-1}^{1} \int_{-\sqrt{1-z^2}}^{\sqrt{1-z^2}} \frac{1}{\sqrt{1 - z^2}} \, dx \, dz = 4 \int_0^1 \int_0^{\sqrt{1-z^2}} \frac{1}{\sqrt{1 - z^2}} \, dx \, dz \quad \left[\begin{array}{c}\text{by the symmetry} \\ \text{of the surface}\end{array}\right]$$

This integral is improper [when $z = 1$], so

$$A(S) = \lim_{t \to 1^-} 4 \int_0^t \int_0^{\sqrt{1-z^2}} \frac{1}{\sqrt{1 - z^2}} \, dx \, dz = \lim_{t \to 1^-} 4 \int_0^t \frac{\sqrt{1 - z^2}}{\sqrt{1 - z^2}} \, dz = \lim_{t \to 1^-} 4 \int_0^t dz = \lim_{t \to 1^-} 4t = 4$$

Since the complete surface consists of four congruent faces, the total surface area is $4(4) = 16$.

Alternate solution: The face of the surface that intersects the positive y-axis can also be parametrized as $\mathbf{r}(x, \theta) = \langle x, \cos\theta, \sin\theta \rangle$ for $-\frac{\pi}{2} \le \theta \le \frac{\pi}{2}$ and $x^2 + z^2 \le 1 \;\Leftrightarrow\; x^2 + \sin^2\theta \le 1 \;\Leftrightarrow$ $-\sqrt{1 - \sin^2\theta} \le x \le \sqrt{1 - \sin^2\theta} \;\Leftrightarrow\; -\cos\theta \le x \le \cos\theta$. Then $\mathbf{r}_x = \langle 1, 0, 0 \rangle$, $\mathbf{r}_\theta = \langle 0, -\sin\theta, \cos\theta \rangle$ and $\mathbf{r}_x \times \mathbf{r}_\theta = \langle 0, -\cos\theta, -\sin\theta \rangle \;\Rightarrow\; |\,\mathbf{r}_x \times \mathbf{r}_\theta\,| = 1$, so $A(S) = \int_{-\pi/2}^{\pi/2} \int_{-\cos\theta}^{\cos\theta} 1 \, dx \, d\theta = \int_{-\pi/2}^{\pi/2} 2\cos\theta \, d\theta = 2\sin\theta\Big]_{-\pi/2}^{\pi/2} = 4$. Again, the area of the complete surface is $4(4) = 16$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

63. Let $A(S_1)$ be the surface area of that portion of the surface which lies above the plane $z = 0$. Then $A(S) = 2A(S_1)$.

Following Example 10, a parametric representation of S_1 is $x = a\sin\phi\cos\theta$, $y = a\sin\phi\sin\theta$, $z = a\cos\phi$ and $|\mathbf{r}_\phi \times \mathbf{r}_\theta| = a^2\sin\phi$. For D, $0 \le \phi \le \frac{\pi}{2}$ and for each fixed ϕ, $\left(x - \frac{1}{2}a\right)^2 + y^2 \le \left(\frac{1}{2}a\right)^2$ or $\left[a\sin\phi\cos\theta - \frac{1}{2}a\right]^2 + a^2\sin^2\phi\sin^2\theta \le (a/2)^2$ implies $a^2\sin^2\phi - a^2\sin\phi\cos\theta \le 0$ or $\sin\phi\,(\sin\phi - \cos\theta) \le 0$. But $0 \le \phi \le \frac{\pi}{2}$, so $\cos\theta \ge \sin\phi$ or $\sin\left(\frac{\pi}{2} + \theta\right) \ge \sin\phi$ or $\phi - \frac{\pi}{2} \le \theta \le \frac{\pi}{2} - \phi$.

Hence $D = \left\{(\phi, \theta) \mid 0 \le \phi \le \frac{\pi}{2}, \phi - \frac{\pi}{2} \le \theta \le \frac{\pi}{2} - \phi\right\}$. Then

$$A(S_1) = \int_0^{\pi/2}\int_{\phi-(\pi/2)}^{(\pi/2)-\phi} a^2\sin\phi\,d\theta\,d\phi = a^2\int_0^{\pi/2}(\pi - 2\phi)\sin\phi\,d\phi$$
$$= a^2\left[(-\pi\cos\phi) - 2(-\phi\cos\phi + \sin\phi)\right]_0^{\pi/2} = a^2(\pi - 2)$$

Thus $A(S) = 2a^2(\pi - 2)$.

Alternate solution: Working on S_1 we could parametrize the portion of the sphere by $x = x$, $y = y$, $z = \sqrt{a^2 - x^2 - y^2}$.

Then $|\mathbf{r}_x \times \mathbf{r}_y| = \sqrt{1 + \dfrac{x^2}{a^2 - x^2 - y^2} + \dfrac{y^2}{a^2 - x^2 - y^2}} = \dfrac{a}{\sqrt{a^2 - x^2 - y^2}}$ and

$$A(S_1) = \iint\limits_{0 \le (x-(a/2))^2 + y^2 \le (a/2)^2} \frac{a}{\sqrt{a^2 - x^2 - y^2}}\,dA = \int_{-\pi/2}^{\pi/2}\int_0^{a\cos\theta} \frac{a}{\sqrt{a^2 - r^2}}\,r\,dr\,d\theta$$
$$= \int_{-\pi/2}^{\pi/2} -a(a^2 - r^2)^{1/2}\Big]_{r=0}^{r=a\cos\theta}\,d\theta = \int_{-\pi/2}^{\pi/2} a^2[1 - (1 - \cos^2\theta)^{1/2}]\,d\theta$$
$$= \int_{-\pi/2}^{\pi/2} a^2(1 - |\sin\theta|)\,d\theta = 2a^2\int_0^{\pi/2}(1 - \sin\theta)\,d\theta = 2a^2\left(\tfrac{\pi}{2} - 1\right)$$

Thus $A(S) = 4a^2\left(\frac{\pi}{2} - 1\right) = 2a^2(\pi - 2)$.

Notes:

(1) Perhaps working in spherical coordinates is the most obvious approach here. However, you must be careful in setting up D.

(2) In the alternate solution, you can avoid having to use $|\sin\theta|$ by working in the first octant and then multiplying by 4. However, if you set up S_1 as above and arrived at $A(S_1) = a^2\pi$, you now see your error.

64. (a) Here $z = a\sin\alpha$, $y = |AB|$, and $x = |OA|$. But $|OB| = |OC| + |CB| = b + a\cos\alpha$ and $\sin\theta = \dfrac{|AB|}{|OB|}$ so that $y = |OB|\sin\theta = (b + a\cos\alpha)\sin\theta$. Similarly $\cos\theta = \dfrac{|OA|}{|OB|}$ so $x = (b + a\cos\alpha)\cos\theta$. Hence a parametric representation for the torus is $x = b\cos\theta + a\cos\alpha\,\cos\theta$, $y = b\sin\theta + a\cos\alpha\,\sin\theta$, $z = a\sin\alpha$, where $0 \le \alpha \le 2\pi$, $0 \le \theta \le 2\pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b)

$a = 1, b = 8$

$a = 3, b = 8$

$a = 3, b = 4$

(c) $x = b\cos\theta + a\cos\alpha\,\cos\theta$, $y = b\sin\theta + a\cos\alpha\,\sin\theta$, $z = a\sin\alpha$, so $\mathbf{r}_\alpha = \langle -a\sin\alpha\,\cos\theta, -a\sin\alpha\,\sin\theta, a\cos\alpha\rangle$, $\mathbf{r}_\theta = \langle -(b + a\cos\alpha)\sin\theta, (b + a\cos\alpha)\cos\theta, 0\rangle$ and

$$\begin{aligned}\mathbf{r}_\alpha \times \mathbf{r}_\theta &= \left(-ab\cos\alpha\,\cos\theta - a^2\cos\alpha\,\cos^2\theta\right)\mathbf{i} + \left(-ab\sin\alpha\,\cos\theta - a^2\sin\alpha\,\cos^2\theta\right)\mathbf{j} \\ &\qquad + \left(-ab\cos^2\alpha\,\sin\theta - a^2\cos^2\alpha\,\sin\theta\,\cos\theta - ab\sin^2\alpha\,\sin\theta - a^2\sin^2\alpha\,\sin\theta\,\cos\theta\right)\mathbf{k} \\ &= -a\,(b + a\cos\alpha)\,[(\cos\theta\,\cos\alpha)\,\mathbf{i} + (\sin\theta\,\cos\alpha)\,\mathbf{j} + (\sin\alpha)\,\mathbf{k}]\end{aligned}$$

Then $|\mathbf{r}_\alpha \times \mathbf{r}_\theta| = a(b + a\cos\alpha)\sqrt{\cos^2\theta\,\cos^2\alpha + \sin^2\theta\,\cos^2\alpha + \sin^2\alpha} = a(b + a\cos\alpha)$.

Note: $b > a$, $-1 \le \cos\alpha \le 1$ so $|b + a\cos\alpha| = b + a\cos\alpha$. Hence

$A(S) = \int_0^{2\pi}\int_0^{2\pi} a(b + a\cos\alpha)\,d\alpha\,d\theta = 2\pi\left[ab\alpha + a^2\sin\alpha\right]_0^{2\pi} = 4\pi^2 ab.$

16.7 Surface Integrals

1. The box is a cube where each face has surface area 4. The centers of the faces are $(\pm1, 0, 0)$, $(0, \pm1, 0)$, $(0, 0, \pm1)$. For each face we take the point P_{ij}^* to be the center of the face and $f(x, y, z) = \cos(x + 2y + 3z)$, so by Definition 1,

$$\begin{aligned}\iint_S f(x, y, z)\,dS &\approx [f(1, 0, 0)](4) + [f(-1, 0, 0)](4) + [f(0, 1, 0)](4) \\ &\qquad + [f(0, -1, 0)](4) + [f(0, 0, 1)](4) + [f(0, 0, -1)](4) \\ &= 4\,[\cos 1 + \cos(-1) + \cos 2 + \cos(-2) + \cos 3 + \cos(-3)] \approx -6.93\end{aligned}$$

2. Each quarter-cylinder has surface area $\frac{1}{4}[2\pi(1)(2)] = \pi$, and the top and bottom disks have surface area $\pi(1)^2 = \pi$. We can take $(0, 0, 1)$ as a sample point in the top disk, $(0, 0, -1)$ in the bottom disk, and $(\pm1, 0, 0)$, $(0, \pm1, 0)$ in the four

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

quarter-cylinders. Then $\iint_S f(x,y,z)\,dS$ can be approximated by the Riemann sum

$$f(1,0,0)(\pi)+f(-1,0,0)(\pi)+f(0,1,0)\,(\pi)+f(0,-1,0)(\pi)+f(0,0,1)(\pi)+f(0,0,-1)(\pi)$$
$$=(2+2+3+3+4+4)\pi=18\pi\approx 56.5.$$

3. We can use the xz- and yz-planes to divide H into four patches of equal size, each with surface area equal to $\frac{1}{8}$ the surface area of a sphere with radius $\sqrt{50}$, so $\Delta S=\frac{1}{8}(4)\pi\left(\sqrt{50}\right)^2=25\pi$. Then $(\pm3,\pm4,5)$ are sample points in the four patches, and using a Riemann sum as in Definition 1, we have

$$\iint_H f(x,y,z)\,dS\approx f(3,4,5)\,\Delta S+f(3,-4,5)\,\Delta S+f(-3,4,5)\,\Delta S+f(-3,-4,5)\,\Delta S$$
$$=(7+8+9+12)(25\pi)=900\pi\approx 2827$$

4. On the surface, $f(x,y,z)=g\left(\sqrt{x^2+y^2+z^2}\right)=g(2)=-5$. So since the area of a sphere is $4\pi r^2$,

$\iint_S f(x,y,z)\,dS=\iint_S g(2)\,dS=-5\iint_S\,dS=-5[4\pi(2)^2]=-80\pi.$

5. $\mathbf{r}(u,v)=(u+v)\,\mathbf{i}+(u-v)\,\mathbf{j}+(1+2u+v)\,\mathbf{k}$, $0\le u\le 2$, $0\le v\le 1$ and

$\mathbf{r}_u\times\mathbf{r}_v=(\mathbf{i}+\mathbf{j}+2\,\mathbf{k})\times(\mathbf{i}-\mathbf{j}+\mathbf{k})=3\,\mathbf{i}+\mathbf{j}-2\,\mathbf{k}\quad\Rightarrow\quad|\mathbf{r}_u\times\mathbf{r}_v|=\sqrt{3^2+1^2+(-2)^2}=\sqrt{14}$. Then by Formula 2,

$$\iint_S(x+y+z)\,dS=\iint_D(u+v+u-v+1+2u+v)\;|\mathbf{r}_u\times\mathbf{r}_v|\;dA=\int_0^1\int_0^2(4u+v+1)\cdot\sqrt{14}\,du\,dv$$
$$=\sqrt{14}\int_0^1\left[2u^2+uv+u\right]_{u=0}^{u=2}dv=\sqrt{14}\int_0^1(2v+10)\,dv=\sqrt{14}\left[v^2+10v\right]_0^1=11\sqrt{14}$$

6. $\mathbf{r}(u,v)=u\cos v\,\mathbf{i}+u\sin v\,\mathbf{j}+u\,\mathbf{k}$, $0\le u\le 1$, $0\le v\le\pi/2$ and

$\mathbf{r}_u\times\mathbf{r}_v=(\cos v\,\mathbf{i}+\sin v\,\mathbf{j}+\mathbf{k})\times(-u\sin v\,\mathbf{i}+u\cos v\,\mathbf{j})=-u\cos v\,\mathbf{i}-u\sin v\,\mathbf{j}+u\,\mathbf{k}\quad\Rightarrow$

$|\mathbf{r}_u\times\mathbf{r}_v|=\sqrt{u^2\cos^2 v+u^2\sin^2 v+u^2}=\sqrt{2u^2}=\sqrt{2}\,u$ [since $u\ge 0$]. Then by Formula 2,

$$\iint_S xyz\,dS=\iint_D(u\cos v)(u\sin v)(u)\;|\mathbf{r}_u\times\mathbf{r}_v|\;dA=\int_0^1\int_0^{\pi/2}(u^3\sin v\cos v)\cdot\sqrt{2}\,u\,dv\,du$$
$$=\sqrt{2}\int_0^1 u^4\,du\int_0^{\pi/2}\sin v\cos v\,dv=\sqrt{2}\left[\tfrac{1}{5}u^5\right]_0^1\left[\tfrac{1}{2}\sin^2 v\right]_0^{\pi/2}=\sqrt{2}\cdot\tfrac{1}{5}\cdot\tfrac{1}{2}=\tfrac{1}{10}\sqrt{2}$$

7. $\mathbf{r}(u,v)=\langle u\cos v,u\sin v,v\rangle$, $0\le u\le 1$, $0\le v\le\pi$ and

$\mathbf{r}_u\times\mathbf{r}_v=\langle\cos v,\sin v,0\rangle\times\langle -u\sin v,u\cos v,1\rangle=\langle\sin v,-\cos v,u\rangle\quad\Rightarrow$

$|\mathbf{r}_u\times\mathbf{r}_v|=\sqrt{\sin^2 v+\cos^2 v+u^2}=\sqrt{u^2+1}$. Then

$$\iint_S y\,dS=\iint_D(u\sin v)\;|\mathbf{r}_u\times\mathbf{r}_v|\;dA=\int_0^1\int_0^{\pi}(u\sin v)\cdot\sqrt{u^2+1}\,dv\,du=\int_0^1 u\sqrt{u^2+1}\,du\int_0^{\pi}\sin v\,dv$$
$$=\left[\tfrac{1}{3}(u^2+1)^{3/2}\right]_0^1\left[-\cos v\right]_0^{\pi}=\tfrac{1}{3}(2^{3/2}-1)\cdot 2=\tfrac{2}{3}(2\sqrt{2}-1)$$

8. $\mathbf{r}(u,v)=\left\langle 2uv,u^2-v^2,u^2+v^2\right\rangle$, $u^2+v^2\le 1$ and

$\mathbf{r}_u\times\mathbf{r}_v=\langle 2v,2u,2u\rangle\times\langle 2u,-2v,2v\rangle=\left\langle 8uv,4u^2-4v^2,-4u^2-4v^2\right\rangle$, so

$$|\mathbf{r}_u\times\mathbf{r}_v|=\sqrt{(8uv)^2+(4u^2-4v^2)^2+(-4u^2-4v^2)^2}=\sqrt{64u^2v^2+32u^4+32v^4}$$
$$=\sqrt{32(u^2+v^2)^2}=4\sqrt{2}\,(u^2+v^2)$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Then

$$\iint_S (x^2+y^2)\,dS = \iint_D \left[(2uv)^2+(u^2-v^2)^2\right]\,|\mathbf{r}_u\times\mathbf{r}_v|\,dA = \iint_D (4u^2v^2+u^4-2u^2v^2+v^4)\cdot 4\sqrt{2}\,(u^2+v^2)\,dA$$
$$= 4\sqrt{2}\iint_D (u^4+2u^2v^2+v^4)\,(u^2+v^2)\,dA = 4\sqrt{2}\iint_D (u^2+v^2)^3\,dA = 4\sqrt{2}\int_0^{2\pi}\int_0^1 (r^2)^3\,r\,dr\,d\theta$$
$$= 4\sqrt{2}\int_0^{2\pi} d\theta \int_0^1 r^7\,dr = 4\sqrt{2}\,\left[\theta\right]_0^{2\pi}\,\left[\tfrac{1}{8}r^8\right]_0^1 = 4\sqrt{2}\cdot 2\pi\cdot\tfrac{1}{8} = \sqrt{2}\,\pi$$

9. $z = 1+2x+3y$, so $\dfrac{\partial z}{\partial x} = 2$ and $\dfrac{\partial z}{\partial y} = 3$. The surface is the graph of a function, so by Formula 4,

$$\iint_S x^2yz\,dS = \iint_D x^2yz\sqrt{\left(\frac{\partial z}{\partial x}\right)^2+\left(\frac{\partial z}{\partial y}\right)^2+1}\,dA = \int_0^3\int_0^2 x^2y(1+2x+3y)\,\sqrt{4+9+1}\,dy\,dx$$
$$= \sqrt{14}\int_0^3\int_0^2 (x^2y+2x^3y+3x^2y^2)\,dy\,dx = \sqrt{14}\int_0^3 \left[\tfrac{1}{2}x^2y^2+x^3y^2+x^2y^3\right]_{y=0}^{y=2}\,dx$$
$$= \sqrt{14}\int_0^3 (10x^2+4x^3)\,dx = \sqrt{14}\left[\tfrac{10}{3}x^3+x^4\right]_0^3 = 171\sqrt{14}$$

10. The surface S is given by $z = 4-2x-2y$, which intersects the xy-plane in the line $2x+2y=4$, so $D = \{(x,y)\mid 0\le x\le 2,\ 0\le y\le 2-x\}$. The surface is the graph of a function, so by Formula 4,

$$\iint_S xz\,dS = \iint_D x(4-2x-2y)\,\sqrt{(-2)^2+(-2)^2+1}\,dA = 3\int_0^2\int_0^{2-x}\left(4x-2x^2-2xy\right)dy\,dx$$
$$= 3\int_0^2\left[4xy-2x^2y-xy^2\right]_{y=0}^{y=2-x}dx = 3\int_0^2\left[4x(2-x)-2x^2(2-x)-x(2-x)^2\right]dx$$
$$= 3\int_0^2\left(x^3-4x^2+4x\right)dx = 3\left[\tfrac{1}{4}x^4-\tfrac{4}{3}x^3+2x^2\right]_0^2 = 3\left(4-\tfrac{32}{3}+8\right) = 4$$

11. An equation of the plane through the points $(1,0,0)$, $(0,-2,0)$, and $(0,0,4)$ is $4x-2y+z=4$ (see Example 12.5.5), so S is the region in the plane $z = 4-4x+2y$ over $D = \{(x,y)\mid 0\le x\le 1,\ 2x-2\le y\le 0\}$. Thus, by Formula 4,

$$\iint_S x\,dS = \iint_D x\,\sqrt{(-4)^2+(2)^2+1}\,dA = \sqrt{21}\int_0^1\int_{2x-2}^0 x\,dy\,dx = \sqrt{21}\int_0^1 [xy]_{y=2x-2}^{y=0}\,dx$$
$$= \sqrt{21}\int_0^1(-2x^2+2x)\,dx = \sqrt{21}\left[-\tfrac{2}{3}x^3+x^2\right]_0^1 = \sqrt{21}\left(-\tfrac{2}{3}+1\right) = \tfrac{\sqrt{21}}{3}$$

12. $z = \frac{2}{3}(x^{3/2}+y^{3/2})$ and by Formula 4,

$$\iint_S y\,dS = \iint_D y\sqrt{\left(\sqrt{x}\right)^2+\left(\sqrt{y}\right)^2+1}\,dA = \int_0^1\int_0^1 y\sqrt{x+y+1}\,dx\,dy$$
$$= \int_0^1 y\left[\tfrac{2}{3}(x+y+1)^{3/2}\right]_{x=0}^{x=1}dy = \int_0^1 \tfrac{2}{3}y\left[(y+2)^{3/2}-(y+1)^{3/2}\right]dy$$

Substituting $u = y+2$ in the first term and $t = y+1$ in the second, we have

$$\iint_S y\,dS = \tfrac{2}{3}\int_2^3(u-2)u^{3/2}\,du - \tfrac{2}{3}\int_1^2(t-1)t^{3/2}\,dt = \tfrac{2}{3}\left[\tfrac{2}{7}u^{7/2}-\tfrac{4}{5}u^{5/2}\right]_2^3 - \tfrac{2}{3}\left[\tfrac{2}{7}t^{7/2}-\tfrac{2}{5}t^{5/2}\right]_1^2$$
$$= \tfrac{2}{3}\left[\tfrac{2}{7}(3^{7/2}-2^{7/2})-\tfrac{4}{5}(3^{5/2}-2^{5/2})-\tfrac{2}{7}(2^{7/2}-1)+\tfrac{2}{5}(2^{5/2}-1)\right]$$
$$= \tfrac{2}{3}\left(\tfrac{18}{35}\sqrt{3}+\tfrac{8}{35}\sqrt{2}-\tfrac{4}{35}\right) = \tfrac{4}{105}\left(9\sqrt{3}+4\sqrt{2}-2\right)$$

13. Using y and z as parameters, we have $\mathbf{r}(y,z) = (y^2+z^2)\,\mathbf{i}+y\,\mathbf{j}+z\,\mathbf{k},\ y^2+z^2\le 1$. Then $\mathbf{r}_y\times\mathbf{r}_z = (2y\,\mathbf{i}+\mathbf{j})\times(2z\,\mathbf{i}+\mathbf{k}) = \mathbf{i}-2y\,\mathbf{j}-2z\,\mathbf{k}$ and $|\mathbf{r}_y\times\mathbf{r}_z| = \sqrt{1+4y^2+4z^2} = \sqrt{1+4(y^2+z^2)}$. Thus, by

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Formula 2,

$$\iint_S z^2\,dS = \iint\limits_{y^2+z^2\le 1} z^2\sqrt{1+4(y^2+z^2)}\,dA = \int_0^{2\pi}\int_0^1 (r\sin\theta)^2\sqrt{1+4r^2}\,r\,dr\,d\theta$$

$$= \int_0^{2\pi}\sin^2\theta\,d\theta\int_0^1 r^3\sqrt{1+4r^2}\,dr \qquad \left[\text{let } u=1+4r^2 \quad\Rightarrow\quad r^2=\tfrac{1}{4}(u-1) \text{ and } r\,dr=\tfrac{1}{8}du\right]$$

$$= \left[\tfrac{1}{2}\theta-\tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\int_1^5\tfrac{1}{4}(u-1)\sqrt{u}\cdot\tfrac{1}{8}du = \pi\cdot\tfrac{1}{32}\int_1^5(u^{3/2}-u^{1/2})\,du = \tfrac{1}{32}\pi\left[\tfrac{2}{5}u^{5/2}-\tfrac{2}{3}u^{3/2}\right]_1^5$$

$$= \tfrac{1}{32}\pi\left[\tfrac{2}{5}(5)^{5/2}-\tfrac{2}{3}(5)^{3/2}-\tfrac{2}{5}+\tfrac{2}{3}\right] = \tfrac{1}{32}\pi\left(\tfrac{20}{3}\sqrt{5}+\tfrac{4}{15}\right) = \tfrac{1}{120}\pi\left(25\sqrt{5}+1\right)$$

14. Using x and z as parameters, we have $\mathbf{r}(x,z) = x\,\mathbf{i}+\sqrt{x^2+z^2}\,\mathbf{j}+z\,\mathbf{k}$, $x^2+z^2\le 25$. Then

$$\mathbf{r}_x\times\mathbf{r}_z = \left(\mathbf{i}+\frac{x}{\sqrt{x^2+z^2}}\,\mathbf{j}\right)\times\left(\frac{z}{\sqrt{x^2+z^2}}\,\mathbf{j}+\mathbf{k}\right) = \frac{x}{\sqrt{x^2+z^2}}\,\mathbf{i}-\mathbf{j}+\frac{z}{\sqrt{x^2+z^2}}\,\mathbf{k} \text{ and}$$

$|\mathbf{r}_x\times\mathbf{r}_z| = \sqrt{\dfrac{x^2}{x^2+z^2}+1+\dfrac{z^2}{x^2+z^2}} = \sqrt{\dfrac{x^2+z^2}{x^2+z^2}+1} = \sqrt{2}$. Thus, by Formula 2,

$$\iint_S y^2z^2\,dS = \iint\limits_{x^2+z^2\le 25}(x^2+z^2)z^2\sqrt{2}\,dA = \sqrt{2}\int_0^{2\pi}\int_0^5 r^2(r\sin\theta)^2\,r\,dr\,d\theta$$

$$= \sqrt{2}\int_0^{2\pi}\sin^2\theta\,d\theta\int_0^5 r^5\,dr = \sqrt{2}\left[\tfrac{1}{2}\theta-\tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{6}r^6\right]_0^5$$

$$= \sqrt{2}\,(\pi)\cdot\tfrac{1}{6}(15{,}625-0) = \frac{15{,}625\sqrt{2}}{6}\pi$$

15. Using x and z as parameters, we have $\mathbf{r}(x,z) = x\,\mathbf{i}+(x^2+4z)\,\mathbf{j}+z\,\mathbf{k}$, $0\le x\le 1$, $0\le z\le 1$. Then

$\mathbf{r}_x\times\mathbf{r}_z = (\mathbf{i}+2x\,\mathbf{j})\times(4\,\mathbf{j}+\mathbf{k}) = 2x\,\mathbf{i}-\mathbf{j}+4\,\mathbf{k}$ and $|\mathbf{r}_x\times\mathbf{r}_z| = \sqrt{4x^2+1+16} = \sqrt{4x^2+17}$. Thus, by Formula 2,

$$\iint_S x\,dS = \int_0^1\int_0^1 x\sqrt{4x^2+17}\,dz\,dx = \int_0^1 x\sqrt{4x^2+17}\,dx = \left[\tfrac{1}{8}\cdot\tfrac{2}{3}(4x^2+17)^{3/2}\right]_0^1$$

$$= \tfrac{1}{12}(21^{3/2}-17^{3/2}) = \tfrac{1}{12}\left(21\sqrt{21}-17\sqrt{17}\right) = \tfrac{7}{4}\sqrt{21}-\tfrac{17}{12}\sqrt{17}$$

16. The sphere intersects the cone in the circle $x^2+y^2=\frac{1}{2}$, $z=\frac{1}{\sqrt{2}}$, so S is the portion of the sphere where $z\ge\frac{1}{\sqrt{2}}$.

Using spherical coordinates to parametrize the sphere we have $\mathbf{r}(\phi,\theta) = \sin\phi\,\cos\theta\,\mathbf{i}+\sin\phi\,\sin\theta\,\mathbf{j}+\cos\phi\,\mathbf{k}$, and

$|\mathbf{r}_\phi\times\mathbf{r}_\theta| = \sin\phi$ (as in Example 1). The portion where $z\ge\frac{1}{\sqrt{2}}$ corresponds to $0\le\phi\le\frac{\pi}{4}$, $0\le\theta\le 2\pi$, so by Formula 2,

$$\iint_S y^2\,dS = \int_0^{2\pi}\int_0^{\pi/4}(\sin\phi\sin\theta)^2(\sin\phi)\,d\phi\,d\theta = \int_0^{2\pi}\sin^2\theta\,d\theta\int_0^{\pi/4}\sin^3\phi\,d\phi = \int_0^{2\pi}\sin^2\theta\,d\theta\int_0^{\pi/4}(1-\cos^2\phi)\sin\phi\,d\phi$$

$$= \left[\tfrac{1}{2}\theta-\tfrac{1}{4}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{3}\cos^3\phi-\cos\phi\right]_0^{\pi/4} = \pi\left(\tfrac{\sqrt{2}}{12}-\tfrac{\sqrt{2}}{2}-\tfrac{1}{3}+1\right) = \left(\tfrac{2}{3}-\tfrac{5\sqrt{2}}{12}\right)\pi$$

17. Using spherical coordinates to parametrize the sphere (see Example 16.6.4), we have

$\mathbf{r}(\phi,\theta) = 2\sin\phi\cos\theta\,\mathbf{i}+2\sin\phi\sin\theta\,\mathbf{j}+2\cos\phi\,\mathbf{k}$ and $|\mathbf{r}_\phi\times\mathbf{r}_\theta| = 4\sin\phi$ (see Example 16.6.10). Here S is the portion of the sphere corresponding to $0\le\phi\le\pi/2$, so by Formula 2,

$$\iint_S(x^2z+y^2z)\,dS = \iint_S(x^2+y^2)z\,dS = \int_0^{2\pi}\int_0^{\pi/2}(4\sin^2\phi)(2\cos\phi)(4\sin\phi)\,d\phi\,d\theta$$

$$= 32\int_0^{2\pi}d\theta\int_0^{\pi/2}\sin^3\phi\cos\phi\,d\phi = 32\,(2\pi)\left[\tfrac{1}{4}\sin^4\phi\right]_0^{\pi/2} = 16\pi(1-0) = 16\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. S is given by $\mathbf{r}(u,v) = \cos v\,\mathbf{i} + u\,\mathbf{j} + \sin v\,\mathbf{k}$, $0 \le u \le 2$, $0 \le v \le \pi$ (see Example 16.6.5). Then

$\mathbf{r}_u \times \mathbf{r}_v = \mathbf{j} \times (-\sin v\,\mathbf{i} + \cos v\,\mathbf{k}) = \cos v\,\mathbf{i} + \sin v\,\mathbf{k}$ and $|\mathbf{r}_u \times \mathbf{r}_v| = \sqrt{\cos^2 v + \sin^2 v} = 1$, so

$$\begin{aligned}\iint_S (x+y+z)\,dS &= \int_0^\pi \int_0^2 (\cos v + u + \sin v)(1)\,du\,dv = \int_0^\pi \left[u(\cos v + \sin v) + \tfrac{1}{2}u^2\right]_{u=0}^{u=2} dv \\ &= \int_0^\pi (2\cos v + 2\sin v + 2)\,dv = [2\sin v - 2\cos v + 2v]_0^\pi = 2 + 2\pi + 2 = 4 + 2\pi\end{aligned}$$

19. Here S consists of three surfaces: S_1, the lateral surface of the cylinder; S_2, the front formed by the plane $x + y = 5$; and the back, S_3, in the plane $x = 0$.

On S_1: the surface is given by $\mathbf{r}(u,v) = u\,\mathbf{i} + 3\cos v\,\mathbf{j} + 3\sin v\,\mathbf{k}$, $0 \le v \le 2\pi$ (see Example 16.6.5), and $0 \le x \le 5 - y \;\Rightarrow\; 0 \le u \le 5 - 3\cos v$. Then $\mathbf{r}_u \times \mathbf{r}_v = -3\cos v\,\mathbf{j} - 3\sin v\,\mathbf{k}$ and $|\mathbf{r}_u \times \mathbf{r}_v| = \sqrt{9\cos^2 v + 9\sin^2 v} = 3$, so

$$\begin{aligned}\iint_{S_1} xz\,dS &= \int_0^{2\pi}\int_0^{5-3\cos v} u(3\sin v)(3)\,du\,dv = 9\int_0^{2\pi} \left[\tfrac{1}{2}u^2\right]_{u=0}^{u=5-3\cos v} \sin v\,dv \\ &= \tfrac{9}{2}\int_0^{2\pi} (5 - 3\cos v)^2 \sin v\,dv = \tfrac{9}{2}\left[\tfrac{1}{9}(5-3\cos v)^3\right]_0^{2\pi} = 0.\end{aligned}$$

On S_2: $\mathbf{r}(y,z) = (5-y)\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$ and $|\mathbf{r}_y \times \mathbf{r}_z| = |\mathbf{i} + \mathbf{j}| = \sqrt{2}$, where $y^2 + z^2 \le 9$ and

$$\begin{aligned}\iint_{S_2} xz\,dS &= \iint_{y^2+z^2\le 9} (5-y)z\sqrt{2}\,dA = \sqrt{2}\int_0^{2\pi}\int_0^3 (5 - r\cos\theta)(r\sin\theta)\,r\,dr\,d\theta \\ &= \sqrt{2}\int_0^{2\pi}\int_0^3 (5r^2 - r^3\cos\theta)(\sin\theta)\,dr\,d\theta = \sqrt{2}\int_0^{2\pi} \left[\tfrac{5}{3}r^3 - \tfrac{1}{4}r^4\cos\theta\right]_{r=0}^{r=3} \sin\theta\,d\theta \\ &= \sqrt{2}\int_0^{2\pi} \left(45 - \tfrac{81}{4}\cos\theta\right)\sin\theta\,d\theta = \sqrt{2}\left(\tfrac{4}{81}\right)\cdot\tfrac{1}{2}\left(45 - \tfrac{81}{4}\cos\theta\right)^2\Big]_0^{2\pi} = 0\end{aligned}$$

On S_3: $x = 0$ so $\iint_{S_3} xz\,dS = 0$. Hence $\iint_S xz\,dS = 0 + 0 + 0 = 0$.

20. Let S_1 be the lateral surface, S_2 the top disk, and S_3 the bottom disk.

On S_1: $\mathbf{r}(\theta,z) = 3\cos\theta\,\mathbf{i} + 3\sin\theta\,\mathbf{j} + z\,\mathbf{k}$, $0 \le \theta \le 2\pi$, $0 \le z \le 2$, $|\mathbf{r}_\theta \times \mathbf{r}_z| = 3$,

$\iint_{S_1} (x^2 + y^2 + z^2)\,dS = \int_0^{2\pi}\int_0^2 (9 + z^2)\,3\,dz\,d\theta = 2\pi(54 + 8) = 124\pi$.

On S_2: $\mathbf{r}(\theta,r) = r\cos\theta\,\mathbf{i} + r\sin\theta\,\mathbf{j} + 2\,\mathbf{k}$, $0 \le r \le 3$, $0 \le \theta \le 2\pi$, $|\mathbf{r}_\theta \times \mathbf{r}_r| = r$,

$\iint_{S_2} (x^2 + y^2 + z^2)\,dS = \int_0^{2\pi}\int_0^3 (r^2 + 4)\,r\,dr\,d\theta = 2\pi\left(\frac{81}{4} + 18\right) = \frac{153}{2}\pi$.

On S_3: $\mathbf{r}(\theta,r) = r\cos\theta\,\mathbf{i} + r\sin\theta\,\mathbf{j}$, $0 \le r \le 3$, $0 \le \theta \le 2\pi$, $|\mathbf{r}_\theta \times \mathbf{r}_r| = r$,

$\iint_{S_3} (x^2 + y^2 + z^2)\,dS = \int_0^{2\pi}\int_0^3 (r^2 + 0)\,r\,dr\,d\theta = 2\pi\left(\frac{81}{4}\right) = \frac{81}{2}\pi$.

Hence $\iint_S \left(x^2 + y^2 + z^2\right) dS = 124\pi + \frac{153}{2}\pi + \frac{81}{2}\pi = 241\pi$.

21. From Exercise 5, $\mathbf{r}(u,v) = (u+v)\,\mathbf{i} + (u-v)\,\mathbf{j} + (1 + 2u + v)\,\mathbf{k}$, $0 \le u \le 2$, $0 \le v \le 1$, and $\mathbf{r}_u \times \mathbf{r}_v = 3\,\mathbf{i} + \mathbf{j} - 2\,\mathbf{k}$. Then

$$\begin{aligned}\mathbf{F}(\mathbf{r}(u,v)) &= (1 + 2u + v)e^{(u+v)(u-v)}\,\mathbf{i} - 3(1 + 2u + v)e^{(u+v)(u-v)}\,\mathbf{j} + (u+v)(u-v)\,\mathbf{k} \\ &= (1 + 2u + v)e^{u^2 - v^2}\,\mathbf{i} - 3(1 + 2u + v)e^{u^2 - v^2}\,\mathbf{j} + (u^2 - v^2)\,\mathbf{k}\end{aligned}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Because the z-component of $\mathbf{r}_u \times \mathbf{r}_v$ is negative we use $-(\mathbf{r}_u \times \mathbf{r}_v)$ in Formula 9 for the upward orientation:

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_D \mathbf{F} \cdot (-(\mathbf{r}_u \times \mathbf{r}_v))\, dA = \int_0^1 \int_0^2 \left[-3(1+2u+v)e^{u^2-v^2} + 3(1+2u+v)e^{u^2-v^2} + 2(u^2-v^2)\right] du\, dv$$
$$= \int_0^1 \int_0^2 2(u^2-v^2)\, du\, dv = 2\int_0^1 \left[\tfrac{1}{3}u^3 - uv^2\right]_{u=0}^{u=2} dv = 2\int_0^1 \left(\tfrac{8}{3} - 2v^2\right) dv$$
$$= 2\left[\tfrac{8}{3}v - \tfrac{2}{3}v^3\right]_0^1 = 2\left(\tfrac{8}{3} - \tfrac{2}{3}\right) = 4$$

22. $\mathbf{r}(u,v) = \langle u\cos v, u\sin v, v\rangle$, $0 \le u \le 1$, $0 \le v \le \pi$ and

$\mathbf{r}_u \times \mathbf{r}_v = \langle \cos v, \sin v, 0\rangle \times \langle -u\sin v, u\cos v, 1\rangle = \langle \sin v, -\cos v, u\rangle$. Since $\mathbf{F}(x,y,z) = z\,\mathbf{i} + y\,\mathbf{j} + x\,\mathbf{k}$,

$\mathbf{F}(\mathbf{r}(u,v)) = v\,\mathbf{i} + u\sin v\,\mathbf{j} + u\cos v\,\mathbf{k}$, and by Formula 9,

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_D \mathbf{F} \cdot (\mathbf{r}_u \times \mathbf{r}_v)\, dA = \int_0^1 \int_0^\pi (v\sin v - u\sin v\cos v + u^2\cos v)\, dv\, du$$
$$= \int_0^1 \left[\sin v - v\cos v - \tfrac{1}{2}u\sin^2 v + u^2\sin v\right]_{v=0}^{v=\pi} du = \int_0^1 \pi\, du = \pi u\Big]_0^1 = \pi$$

23. $\mathbf{F}(x,y,z) = xy\,\mathbf{i} + yz\,\mathbf{j} + zx\,\mathbf{k}$, $z = g(x,y) = 4 - x^2 - y^2$, and D is the square $[0,1] \times [0,1]$, so by Equation 10

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_D [-xy(-2x) - yz(-2y) + zx]\, dA = \int_0^1 \int_0^1 [2x^2y + 2y^2(4-x^2-y^2) + x(4-x^2-y^2)]\, dy\, dx$$
$$= \int_0^1 \left[x^2y^2 + \tfrac{8}{3}y^3 - \tfrac{2}{3}x^2y^3 - \tfrac{2}{5}y^5 + 4xy - x^3y - \tfrac{1}{3}xy^3\right]_{y=0}^{y=1} dx$$
$$= \int_0^1 \left(\tfrac{1}{3}x^2 + \tfrac{11}{3}x - x^3 + \tfrac{34}{15}\right) dx = \left[\tfrac{1}{9}x^3 + \tfrac{11}{6}x^2 - \tfrac{1}{4}x^4 + \tfrac{34}{15}x\right]_0^1 = \tfrac{713}{180}$$

24. $\mathbf{F}(x,y,z) = -x\,\mathbf{i} - y\,\mathbf{j} + z^3\,\mathbf{k}$, $z = g(x,y) = \sqrt{x^2+y^2}$, and D is the annular region $\{(x,y) \mid 1 \le x^2+y^2 \le 9\}$. Since S has downward orientation, we have, by Equation 10,

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = -\iint_D \left[-(-x)\left(\frac{x}{\sqrt{x^2+y^2}}\right) - (-y)\left(\frac{y}{\sqrt{x^2+y^2}}\right) + z^3\right] dA$$
$$= -\iint_D \left[\frac{x^2+y^2}{\sqrt{x^2+y^2}} + \left(\sqrt{x^2+y^2}\right)^3\right] dA = -\int_0^{2\pi} \int_1^3 \left(\frac{r^2}{r} + r^3\right) r\, dr\, d\theta$$
$$= -\int_0^{2\pi} d\theta \int_1^3 (r^2 + r^4)\, dr = -\left[\theta\right]_0^{2\pi} \left[\tfrac{1}{3}r^3 + \tfrac{1}{5}r^5\right]_1^3$$
$$= -2\pi\left(9 + \tfrac{243}{5} - \tfrac{1}{3} - \tfrac{1}{5}\right) = -\tfrac{1712}{15}\pi$$

25. $\mathbf{F}(x,y,z) = x\,\mathbf{i} + y\,\mathbf{j} + z^2\,\mathbf{k}$, and using spherical coordinates, S is given by $x = \sin\phi\cos\theta$, $y = \sin\phi\sin\theta$, $z = \cos\phi$, $0 \le \theta \le 2\pi$, $0 \le \phi \le \pi$. $\mathbf{F}(\mathbf{r}(\phi,\theta)) = (\sin\phi\cos\theta)\,\mathbf{i} + (\sin\phi\sin\theta)\,\mathbf{j} + (\cos^2\phi)\,\mathbf{k}$ and, from Example 4,

$\mathbf{r}_\phi \times \mathbf{r}_\theta = \sin^2\phi\cos\theta\,\mathbf{i} + \sin^2\phi\sin\theta\,\mathbf{j} + \sin\phi\cos\phi\,\mathbf{k}$. Thus,

$$\mathbf{F}(\mathbf{r}(\phi,\theta)) \cdot (\mathbf{r}_\phi \times \mathbf{r}_\theta) = \sin^3\phi\cos^2\theta + \sin^3\phi\sin^2\theta + \sin\phi\cos^3\phi = \sin^3\phi + \sin\phi\cos^3\phi$$

and

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_D [\mathbf{F}(\mathbf{r}(\phi,\theta)) \cdot (\mathbf{r}_\phi \times \mathbf{r}_\theta)]\, dA = \int_0^{2\pi} \int_0^\pi (\sin^3\phi + \sin\phi\cos^3\phi)\, d\phi\, d\theta$$
$$= \int_0^{2\pi} d\theta \int_0^\pi (1 - \cos^2\phi + \cos^3\phi)\sin\phi\, d\phi = (2\pi)\left[-\cos\phi + \tfrac{1}{3}\cos^3\phi - \tfrac{1}{4}\cos^4\phi\right]_0^\pi$$
$$= 2\pi\left(1 - \tfrac{1}{3} - \tfrac{1}{4} + 1 - \tfrac{1}{3} + \tfrac{1}{4}\right) = \tfrac{8}{3}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $\mathbf{F}(x,y,z) = y\,\mathbf{i} - x\,\mathbf{j} + 2z\,\mathbf{k}$, $z = g(x,y) = \sqrt{4 - x^2 - y^2}$ and D is the disk $\{(x,y) \mid x^2 + y^2 \le 4\}$. S has downward orientation, so by Equation 10,

$$\begin{aligned}
\iint_S \mathbf{F}\cdot d\mathbf{S} &= -\iint_D \left[-y\cdot\tfrac{1}{2}(4-x^2-y^2)^{-1/2}(-2x) - (-x)\cdot\tfrac{1}{2}(4-x^2-y^2)^{-1/2}(-2y) + 2z\right] dA \\
&= -\iint_D \left(\frac{xy}{\sqrt{4-x^2-y^2}} - \frac{xy}{\sqrt{4-x^2-y^2}} + 2\sqrt{4-x^2-y^2}\right) dA \\
&= -\iint_D 2\sqrt{4-x^2-y^2}\,dA = -2\int_0^{2\pi}\int_0^2 \sqrt{4-r^2}\,r\,dr\,d\theta = -2\int_0^{2\pi} d\theta \int_0^2 r\sqrt{4-r^2}\,dr \\
&= -2(2\pi)\left[-\tfrac{1}{2}\cdot\tfrac{2}{3}(4-r^2)^{3/2}\right]_0^2 = -4\pi\left[0 + \tfrac{1}{3}(4)^{3/2}\right] = -4\pi\cdot\tfrac{8}{3} = -\tfrac{32}{3}\pi
\end{aligned}$$

27. $\mathbf{F}(x,y,z) = y\,\mathbf{j} - z\,\mathbf{k}$. Let S_1 be the paraboloid $y = x^2 + z^2$, $0 \le y \le 1$ and S_2 the disk $x^2 + z^2 \le 1$, $y = 1$. On S_1 we have $\mathbf{r}(x,z) = x\,\mathbf{i} + (x^2+z^2)\,\mathbf{j} + z\,\mathbf{k}$. Since S is a closed surface, we use the outward orientation.

On S_1: $\mathbf{F}(\mathbf{r}(x,z)) = (x^2+z^2)\,\mathbf{j} - z\,\mathbf{k}$ and $\mathbf{r}_x \times \mathbf{r}_z = 2x\,\mathbf{i} - \mathbf{j} + 2z\,\mathbf{k}$ (since the **j**-component must be negative on S_1). Then by Formula 9,

$$\begin{aligned}
\iint_{S_1} \mathbf{F}\cdot d\mathbf{S} &= \iint_{x^2+z^2\le 1} [0 - (x^2+z^2) - 2z^2]\,dA = -\int_0^{2\pi}\int_0^1 (r^2 + 2r^2\sin^2\theta)\,r\,dr\,d\theta \\
&= -\int_0^{2\pi}\int_0^1 r^3(1+2\sin^2\theta)\,dr\,d\theta = -\int_0^{2\pi}(1+1-\cos 2\theta)\,d\theta \int_0^1 r^3\,dr \\
&= -\left[2\theta - \tfrac{1}{2}\sin 2\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^1 = -4\pi\cdot\tfrac{1}{4} = -\pi
\end{aligned}$$

On S_2: $\mathbf{F}(\mathbf{r}(x,z)) = \mathbf{j} - z\,\mathbf{k}$ and $\mathbf{r}_z \times \mathbf{r}_x = \mathbf{j}$. Then $\iint_{S_2} \mathbf{F}\cdot d\mathbf{S} = \iint_{x^2+z^2\le 1} (1)\,dA = \pi$.

Hence $\iint_S \mathbf{F}\cdot d\mathbf{S} = -\pi + \pi = 0$.

28. $\mathbf{F}(x,y,z) = yz\,\mathbf{i} + zx\,\mathbf{j} + xy\,\mathbf{k}$, $z = g(x,y) = x\sin y$, and D is the rectangle $[0,2]\times[0,\pi]$, so by Equation 10

$$\begin{aligned}
\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iint_D [-yz(\sin y) - zx(x\cos y) + xy]\,dA = \int_0^{\pi}\int_0^2 (-xy\sin^2 y - x^3\sin y\cos y + xy)\,dx\,dy \\
&= \int_0^{\pi}\left[-\tfrac{1}{2}x^2 y\sin^2 y - \tfrac{1}{4}x^4\sin y\cos y + \tfrac{1}{2}x^2 y\right]_{x=0}^{x=2} dy \\
&= \int_0^{\pi}\left(-2y\sin^2 y - 4\sin y\cos y + 2y\right) dy \qquad \text{[integrate by parts in the first term]} \\
&= \left[\left(-\tfrac{1}{2}y^2 + \tfrac{1}{2}y\sin 2y + \tfrac{1}{4}\cos 2y\right) - 2\sin^2 y + y^2\right]_0^{\pi} = -\tfrac{1}{2}\pi^2 + \tfrac{1}{4} + \pi^2 - \tfrac{1}{4} = \tfrac{1}{2}\pi^2
\end{aligned}$$

29. Here S consists of the six faces of the cube as labeled in the figure. On S_1:

$\mathbf{F} = \mathbf{i} + 2y\,\mathbf{j} + 3z\,\mathbf{k}$, $\mathbf{r}_y \times \mathbf{r}_z = \mathbf{i}$ and $\iint_{S_1} \mathbf{F}\cdot d\mathbf{S} = \int_{-1}^1\int_{-1}^1 dy\,dz = 4$;

S_2: $\mathbf{F} = x\,\mathbf{i} + 2\,\mathbf{j} + 3z\,\mathbf{k}$, $\mathbf{r}_z \times \mathbf{r}_x = \mathbf{j}$ and $\iint_{S_2} \mathbf{F}\cdot d\mathbf{S} = \int_{-1}^1\int_{-1}^1 2\,dx\,dz = 8$;

S_3: $\mathbf{F} = x\,\mathbf{i} + 2y\,\mathbf{j} + 3\,\mathbf{k}$, $\mathbf{r}_x \times \mathbf{r}_y = \mathbf{k}$ and $\iint_{S_3} \mathbf{F}\cdot d\mathbf{S} = \int_{-1}^1\int_{-1}^1 3\,dx\,dy = 12$;

S_4: $\mathbf{F} = -\mathbf{i} + 2y\,\mathbf{j} + 3z\,\mathbf{k}$, $\mathbf{r}_z \times \mathbf{r}_y = -\mathbf{i}$ and $\iint_{S_4} \mathbf{F}\cdot d\mathbf{S} = 4$;

S_5: $\mathbf{F} = x\,\mathbf{i} - 2\,\mathbf{j} + 3z\,\mathbf{k}$, $\mathbf{r}_x \times \mathbf{r}_z = -\mathbf{j}$ and $\iint_{S_5} \mathbf{F}\cdot d\mathbf{S} = 8$;

S_6: $\mathbf{F} = x\,\mathbf{i} + 2y\,\mathbf{j} - 3\,\mathbf{k}$, $\mathbf{r}_y \times \mathbf{r}_x = -\mathbf{k}$ and $\iint_{S_6} \mathbf{F}\cdot d\mathbf{S} = \int_{-1}^1\int_{-1}^1 3\,dx\,dy = 12$.

Hence $\iint_S \mathbf{F}\cdot d\mathbf{S} = \sum_{i=1}^{6} \iint_{S_i} \mathbf{F}\cdot d\mathbf{S} = 48$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $\mathbf{F}(x, y, z) = x\,\mathbf{i} + y\,\mathbf{j} + 5\,\mathbf{k}$. Here S consists of three surfaces: S_1, the lateral surface of the cylinder $x^2 + z^2 = 1$; S_2, the front formed by the plane $x + y = 2$; and the back, S_3, in the plane $y = 0$.

On S_1: $\mathbf{r}(\theta, y) = \sin\theta\,\mathbf{i} + y\,\mathbf{j} + \cos\,\theta\,\mathbf{k}$. $\mathbf{F}(\mathbf{r}(\theta, y)) = \sin\theta\,\mathbf{i} + y\,\mathbf{j} + 5\,\mathbf{k}$ and $\mathbf{r}_\theta \times \mathbf{r}_y = \sin\theta\,\mathbf{i} + \cos\theta\,\mathbf{k} \quad \Rightarrow$

$$\iint_{S_1} \mathbf{F} \cdot d\mathbf{S} = \int_0^{2\pi} \int_0^{2-\sin\theta} (\sin^2\theta + 5\cos\theta)\,dy\,d\theta$$
$$= \int_0^{2\pi} (2\sin^2\theta + 10\cos\theta - \sin^3\theta - 5\sin\theta\,\cos\theta)\,d\theta = 2\pi$$

On S_2: $\mathbf{r}(x, z) = x\,\mathbf{i} + (2 - x)\,\mathbf{j} + z\,\mathbf{k}$. $\mathbf{F}(\mathbf{r}(x, z)) = x\,\mathbf{i} + (2 - x)\,\mathbf{j} + 5\,\mathbf{k}$ and $\mathbf{r}_z \times \mathbf{r}_x = \mathbf{i} + \mathbf{j}$.

$$\iint_{S_2} \mathbf{F} \cdot d\mathbf{S} = \iint_{x^2 + z^2 \le 1} [x + (2 - x)]\,dA = 2\pi$$

On S_3: $\mathbf{F}(\mathbf{r}(x, z)) = x\,\mathbf{i} + 5\,\mathbf{k}$. The surface is oriented in the negative y-direction so that $\mathbf{n} = -\mathbf{j}$ and by (8), $\iint_{S_3} \mathbf{F} \cdot d\mathbf{S} = \iint_{S_3} \mathbf{F} \cdot \mathbf{n}\,d\mathbf{S} = 0$. Hence, $\iint_S \mathbf{F} \cdot d\mathbf{S} = 4\pi$.

31. $\mathbf{F}(x, y, z) = x^2\,\mathbf{i} + y^2\,\mathbf{j} + z^2\,\mathbf{k}$. Here S consists of four surfaces: S_1, the top surface (a portion of the circular cylinder $y^2 + z^2 = 1$); S_2, the bottom surface (a portion of the xy-plane); S_3, the front half-disk in the plane $x = 2$, and S_4, the back half-disk in the plane $x = 0$.

On S_1: The surface is $z = \sqrt{1 - y^2}$ for $0 \le x \le 2$, $-1 \le y \le 1$ with upward orientation, so by Equation 10,

$$\iint_{S_1} \mathbf{F} \cdot d\mathbf{S} = \int_0^2 \int_{-1}^1 \left[-x^2\,(0) - y^2 \left(-\frac{y}{\sqrt{1 - y^2}}\right) + z^2\right] dy\,dx = \int_0^2 \int_{-1}^1 \left(\frac{y^3}{\sqrt{1 - y^2}} + 1 - y^2\right) dy\,dx$$
$$= \int_0^2 \left[-\sqrt{1 - y^2} + \tfrac{1}{3}(1 - y^2)^{3/2} + y - \tfrac{1}{3}y^3\right]_{y=-1}^{y=1} dx = \int_0^2 \tfrac{4}{3}\,dx = \tfrac{8}{3}$$

On S_2: The surface is $z = 0$ for $0 \le x \le 2$, $-1 \le y \le 1$ with downward orientation, so that $\mathbf{n} = -\mathbf{k}$ and by (8),

$$\iint_{S_2} \mathbf{F} \cdot d\mathbf{S} = \iint_{S_2} \mathbf{F} \cdot \mathbf{n}\,d\mathbf{S} = \int_0^2 \int_{-1}^1 \left(-z^2\right) dy\,dx = \int_0^2 \int_{-1}^1 (0)\,dy\,dx = 0$$

On S_3: The surface is $x = 2$ for $-1 \le y \le 1$, $0 \le z \le \sqrt{1 - y^2}$, oriented in the positive x-direction, so that $\mathbf{n} = \mathbf{i}$ and by (8),

$$\iint_{S_3} \mathbf{F} \cdot d\mathbf{S} = \iint_{S_3} \mathbf{F} \cdot \mathbf{n}\,d\mathbf{S} = \int_{-1}^1 \int_0^{\sqrt{1-y^2}} x^2\,dz\,dy = \int_{-1}^1 \int_0^{\sqrt{1-y^2}} 4\,dz\,dy = 4A\,(S_3) = 2\pi$$

On S_4: The surface is $x = 0$ for $-1 \le y \le 1$, $0 \le z \le \sqrt{1 - y^2}$, oriented in the negative x-direction, so that $\mathbf{n} = -\mathbf{i}$ and by (8),

$$\iint_{S_4} \mathbf{F} \cdot d\mathbf{S} = \iint_{S_4} \mathbf{F} \cdot \mathbf{n}\,d\mathbf{S} = \int_{-1}^1 \int_0^{\sqrt{1-y^2}} (-x^2)\,dz\,dy = \int_{-1}^1 \int_0^{\sqrt{1-y^2}} (0)\;dz\,dy = 0$$

Thus, $\iint_S \mathbf{F} \cdot d\mathbf{S} = \frac{8}{3} + 0 + 2\pi + 0 = 2\pi + \frac{8}{3}$.

32. $\mathbf{F}(x, y, z) = y\,\mathbf{i} + (z - y)\,\mathbf{j} + x\,\mathbf{k}$. Here S consists of four surfaces: S_1, the triangular face with vertices $(1, 0, 0)$, $(0, 1, 0)$, and $(0, 0, 1)$; S_2, the face of the tetrahedron in the xy-plane; S_3, the face in the xz-plane; and S_4, the face in the yz-plane.

On S_1: The face is the portion of the plane $z = 1 - x - y$ for $0 \le x \le 1$, $0 \le y \le 1 - x$ with upward orientation,

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

so by Equation 10,

$$\iint_{S_1} \mathbf{F}\cdot d\mathbf{S} = \int_0^1\int_0^{1-x}\left[-y\,(-1)-(z-y)\,(-1)+x\right]dy\,dx = \int_0^1\int_0^{1-x}(z+x)\,dy\,dx = \int_0^1\int_0^{1-x}(1-y)\,dy\,dx$$
$$= \int_0^1\left[y-\tfrac{1}{2}y^2\right]_{y=0}^{y=1-x}dx = \tfrac{1}{2}\int_0^1\left(1-x^2\right)dx = \tfrac{1}{2}\left[x-\tfrac{1}{3}x^3\right]_0^1 = \tfrac{1}{3}$$

On S_2: The surface is $z=0$ for $0\le x\le 1$, $0\le y\le 1-x$ with downward orientation, so that that $\mathbf{n}=-\mathbf{k}$ and by (8),

$$\iint_{S_2}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_2}\mathbf{F}\cdot\mathbf{n}\,dS = \int_0^1\int_0^{1-x}(-x)\,dy\,dx = -\int_0^1 x\,(1-x)\,dx = -\left[\tfrac{1}{2}x^2-\tfrac{1}{3}x^3\right]_0^1 = -\tfrac{1}{6}$$

On S_3: The surface is $y=0$ for $0\le x\le 1$, $0\le z\le 1-x$, oriented in the negative y-direction, so that $\mathbf{n}=-\mathbf{j}$ and by (8),

$$\iint_{S_3}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_3}\mathbf{F}\cdot\mathbf{n}\,dS = \int_0^1\int_0^{1-x}-(z-y)\,dz\,dx = -\int_0^1\int_0^{1-x}z\,dz\,dx = -\int_0^1\left[\tfrac{1}{2}z^2\right]_{z=0}^{z=1-x}dx$$
$$= -\tfrac{1}{2}\int_0^1(1-x)^2\,dx = \tfrac{1}{6}\left[(1-x)^3\right]_0^1 = -\tfrac{1}{6}$$

On S_4: The surface is $x=0$ for $0\le y\le 1$, $0\le z\le 1-y$, oriented in the negative x-direction, so that $\mathbf{n}=-\mathbf{i}$ and by (8),

$$\iint_{S_4}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_4}\mathbf{F}\cdot\mathbf{n}\,dS = \int_0^1\int_0^{1-y}(-y)\,dz\,dy = -\int_0^1 y\,(1-y)\,dy = -\left[\tfrac{1}{2}y^2-\tfrac{1}{3}y^3\right]_0^1 = -\tfrac{1}{6}$$

Thus, $\iint_S \mathbf{F}\cdot d\mathbf{S} = \frac{1}{3}-\frac{1}{6}-\frac{1}{6}-\frac{1}{6} = -\frac{1}{6}$.

33. $z = xe^y \;\Rightarrow\; \partial z/\partial x = e^y$, $\partial z/\partial y = xe^y$, so by Formula 4, a CAS gives

$\iint_S (x^2+y^2+z^2)\,dS = \int_0^1\int_0^1 (x^2+y^2+x^2e^{2y})\sqrt{e^{2y}+x^2e^{2y}+1}\,dx\,dy \approx 4.5822.$

34. $z = x^2y^2 \;\Rightarrow\; \partial z/\partial x = 2xy^2,\ \partial z/\partial y = 2x^2y$, so by Formula 4, a CAS gives

$$\iint_S xyz\,dS = \int_0^2\int_0^1 xy(x^2y^2)\sqrt{(2xy^2)^2+(2x^2y)^2+1}\,dx\,dy$$
$$= \int_0^2\int_0^1 x^3y^3\sqrt{4x^2y^4+4x^4y^2+1}\,dx\,dy = -\tfrac{151}{33}-\tfrac{1}{220}\sqrt{3}\,\pi+\tfrac{1977}{176}\ln 7-\tfrac{9891}{880}\ln 3+\tfrac{3}{440}\sqrt{3}\tan^{-1}\tfrac{5}{\sqrt{3}}$$

35. We use Formula 4 with $z = 3-2x^2-y^2 \;\Rightarrow\; \partial z/\partial x = -4x$, $\partial z/\partial y = -2y$. The boundaries of the region $3-2x^2-y^2\ge 0$ are $-\sqrt{\frac{3}{2}}\le x\le\sqrt{\frac{3}{2}}$ and $-\sqrt{3-2x^2}\le y\le\sqrt{3-2x^2}$, so we use a CAS (with precision reduced to seven or fewer digits; otherwise the calculation may take a long time) to calculate

$$\iint_S x^2y^2z^2\,dS = \int_{-\sqrt{3/2}}^{\sqrt{3/2}}\int_{-\sqrt{3-2x^2}}^{\sqrt{3-2x^2}} x^2y^2(3-2x^2-y^2)^2\sqrt{16x^2+4y^2+1}\,dy\,dx \approx 3.4895$$

36. The flux of $\mathbf{F}$ across S is given by $\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_S \mathbf{F}\cdot\mathbf{n}\,dS$. Now on S, $z = g(x,y) = 2\sqrt{1-y^2}$, so $\partial g/\partial x = 0$ and $\partial g/\partial y = -2y(1-y^2)^{-1/2}$. Therefore, by Equation 10,

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \int_{-2}^2\int_{-1}^1\left(-x^2y\left[-2y(1-y^2)^{-1/2}\right]+\left[2\sqrt{1-y^2}\right]^2 e^{x/5}\right)dy\,dx = \tfrac{1}{3}(16\pi+80e^{2/5}-80e^{-2/5})$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. If S is given by $y = h(x, z)$, then S is also the level surface $f(x, y, z) = y - h(x, z) = 0$.

$\mathbf{n} = \dfrac{\nabla f(x, y, z)}{|\nabla f(x, y, z)|} = \dfrac{-h_x\,\mathbf{i} + \mathbf{j} - h_z\,\mathbf{k}}{\sqrt{h_x^2 + 1 + h_z^2}}$, and $-\mathbf{n}$ is the unit normal that points to the left. Now we proceed as in the derivation of (10), using Formula 4 to evaluate

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S \mathbf{F} \cdot (-\mathbf{n})\, dS = \iint_D (P\,\mathbf{i} + Q\,\mathbf{j} + R\,\mathbf{k}) \cdot \frac{\dfrac{\partial h}{\partial x}\,\mathbf{i} - \mathbf{j} + \dfrac{\partial h}{\partial z}\,\mathbf{k}}{\sqrt{\left(\dfrac{\partial h}{\partial x}\right)^2 + 1 + \left(\dfrac{\partial h}{\partial z}\right)^2}} \sqrt{\left(\frac{\partial h}{\partial x}\right)^2 + 1 + \left(\frac{\partial h}{\partial z}\right)^2}\, dA$$

where D is the projection of S onto the xz-plane. Therefore $\displaystyle\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_D \left(P\,\frac{\partial h}{\partial x} - Q + R\,\frac{\partial h}{\partial z}\right) dA$.

38. If S is given by $x = k(y, z)$, then S is also the level surface $f(x, y, z) = x - k(y, z) = 0$.

$\mathbf{n} = \dfrac{\nabla f(x, y, z)}{|\nabla f(x, y, z)|} = \dfrac{\mathbf{i} - k_y\,\mathbf{j} - k_z\,\mathbf{k}}{\sqrt{1 + k_y^2 + k_z^2}}$, and since the x-component is positive this is the unit normal that points forward.

Now we proceed as in the derivation of (10), using Formula 4 for

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S \mathbf{F} \cdot \mathbf{n}\, dS = \iint_D (P\,\mathbf{i} + Q\,\mathbf{j} + R\,\mathbf{k}) \cdot \frac{\mathbf{i} - \dfrac{\partial k}{\partial y}\,\mathbf{j} - \dfrac{\partial k}{\partial z}\,\mathbf{k}}{\sqrt{1 + \left(\dfrac{\partial k}{\partial y}\right)^2 + \left(\dfrac{\partial k}{\partial z}\right)^2}} \sqrt{1 + \left(\frac{\partial k}{\partial y}\right)^2 + \left(\frac{\partial k}{\partial z}\right)^2}\, dA$$

where D is the projection of S onto the yz-plane. Therefore $\displaystyle\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_D \left(P - Q\,\frac{\partial k}{\partial y} - R\,\frac{\partial k}{\partial z}\right) dA$.

39. $m = \iint_S K\, dS = K \cdot 4\pi\left(\frac{1}{2}a^2\right) = 2\pi a^2 K$; by symmetry $M_{xz} = M_{yz} = 0$, and

$M_{xy} = \iint_S zK\, dS = K \int_0^{2\pi} \int_0^{\pi/2} (a\cos\phi)(a^2 \sin\phi)\, d\phi\, d\theta = 2\pi K a^3 \left[-\frac{1}{4}\cos 2\phi\right]_0^{\pi/2} = \pi K a^3$.

Hence $(\overline{x}, \overline{y}, \overline{z}) = \left(0, 0, \frac{1}{2}a\right)$.

40. S is given by $\mathbf{r}(x, y) = x\,\mathbf{i} + y\,\mathbf{j} + \sqrt{x^2 + y^2}\,\mathbf{k}$, $|\mathbf{r}_x \times \mathbf{r}_y| = \sqrt{1 + \dfrac{x^2 + y^2}{x^2 + y^2}} = \sqrt{2}$ so

$$\begin{aligned} m &= \iint_S (10 - z)\, dS = \iint_S \left(10 - \sqrt{x^2 + y^2}\right) dS = \iint_{1 \le x^2 + y^2 \le 16} \left(10 - \sqrt{x^2 + y^2}\right) \sqrt{2}\, dA \\ &= \int_0^{2\pi} \int_1^4 \sqrt{2}\,(10 - r)\, r\, dr\, d\theta = 2\pi\sqrt{2}\left[5r^2 - \tfrac{1}{3}r^3\right]_1^4 = 108\sqrt{2}\,\pi \end{aligned}$$

41. (a) $I_z = \iint_S (x^2 + y^2)\rho(x, y, z)\, dS$ (see 15.6.16).

(b) $I_z = \iint_S (x^2 + y^2)\left(10 - \sqrt{x^2 + y^2}\right) dS = \displaystyle\iint_{1 \le x^2 + y^2 \le 16} (x^2 + y^2)\left(10 - \sqrt{x^2 + y^2}\right) \sqrt{2}\, dA$

$= \int_0^{2\pi} \int_1^4 \sqrt{2}\,(10r^3 - r^4)\, dr\, d\theta = 2\sqrt{2}\,\pi\left(\frac{4329}{10}\right) = \frac{4329}{5}\sqrt{2}\,\pi$

42. Using spherical coordinates to parametrize the sphere we have $\mathbf{r}(\phi, \theta) = 5\sin\phi\cos\theta\,\mathbf{i} + 5\sin\phi\sin\theta\,\mathbf{j} + 5\cos\phi\,\mathbf{k}$, and $|\mathbf{r}_\phi \times \mathbf{r}_\theta| = 25\sin\phi$ (see Example 16.6.10). S is the portion of the sphere where $z \ge 4$, so $0 \le \phi \le \tan^{-1}(3/4)$ and $0 \le \theta \le 2\pi$.

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(a) $m = \iint_S \rho(x,y,z)dS = \int_0^{2\pi}\int_0^{\tan^{-1}(3/4)} k(25\sin\phi)\,d\phi\,d\theta = 25k\int_0^{2\pi} d\theta \int_0^{\tan^{-1}(3/4)} \sin\phi\,d\phi$

$= 25k(2\pi)\left[-\cos\left(\tan^{-1}\frac{3}{4}\right)+1\right] = 50\pi k\left(-\frac{4}{5}+1\right) = 10\pi k.$

Because S has constant density, $\overline{x} = \overline{y} = 0$ by symmetry, and

$\overline{z} = \frac{1}{m}\iint_S z\rho(x,y,z)dS = \frac{1}{10\pi k}\int_0^{2\pi}\int_0^{\tan^{-1}(3/4)} k(5\cos\phi)(25\sin\phi)\,d\phi\,d\theta$

$= \frac{1}{10\pi k}(125k)\int_0^{2\pi} d\theta \int_0^{\tan^{-1}(3/4)} \sin\phi\cos\phi\,d\phi = \frac{1}{10\pi k}(125k)(2\pi)\left[\frac{1}{2}\sin^2\phi\right]_0^{\tan^{-1}(3/4)} = 25\cdot\frac{1}{2}\left(\frac{3}{5}\right)^2 = \frac{9}{2},$

so the center of mass is $(\overline{x},\overline{y},\overline{z}) = \left(0,0,\frac{9}{2}\right)$.

(b) $I_z = \iint_S (x^2+y^2)\rho(x,y,z)dS = \int_0^{2\pi}\int_0^{\tan^{-1}(3/4)} k(25\sin^2\phi)(25\sin\phi)\,d\phi\,d\theta$

$= 625k\int_0^{2\pi} d\theta \int_0^{\tan^{-1}(3/4)} \sin^3\phi\,d\phi = 625k(2\pi)\left[\frac{1}{3}\cos^3\phi - \cos\phi\right]_0^{\tan^{-1}(3/4)}$

$= 1250\pi k\left[\frac{1}{3}\left(\frac{4}{5}\right)^3 - \frac{4}{5} - \frac{1}{3} + 1\right] = 1250\pi k\left(\frac{14}{375}\right) = \frac{140}{3}\pi k$

43. The rate of flow through the cylinder is the flux $\iint_S \rho\mathbf{v}\cdot\mathbf{n}\,dS = \iint_S \rho\mathbf{v}\cdot d\mathbf{S}$ (see Formula 7). We use the parametric representation of the cylinder $\mathbf{r}(u,v) = 2\cos u\,\mathbf{i} + 2\sin u\,\mathbf{j} + v\,\mathbf{k}$ for S, where $0 \le u \le 2\pi$, $0 \le v \le 1$, so $\mathbf{r}_u = -2\sin u\,\mathbf{i} + 2\cos u\,\mathbf{j}$, $\mathbf{r}_v = \mathbf{k}$, and the outward orientation is given by $\mathbf{r}_u \times \mathbf{r}_v = 2\cos u\,\mathbf{i} + 2\sin u\,\mathbf{j}$. Then

$$\begin{aligned}\iint_S \rho\mathbf{v}\cdot d\mathbf{S} &= \rho\int_0^{2\pi}\int_0^1 \left(v\,\mathbf{i} + 4\sin^2 u\,\mathbf{j} + 4\cos^2 u\,\mathbf{k}\right)\cdot(2\cos u\,\mathbf{i} + 2\sin u\,\mathbf{j})\,dv\,du \\ &= \rho\int_0^{2\pi}\int_0^1 \left(2v\cos u + 8\sin^3 u\right)dv\,du = \rho\int_0^{2\pi}\left(\cos u + 8\sin^3 u\right)du \\ &= \rho\left[\sin u + 8\left(-\tfrac{1}{3}\right)(2+\sin^2 u)\cos u\right]_0^{2\pi} = 0\text{ kg/s}\end{aligned}$$

44. A parametric representation for the hemisphere S is $\mathbf{r}(\phi,\theta) = 3\sin\phi\,\cos\theta\,\mathbf{i} + 3\sin\phi\,\sin\theta\,\mathbf{j} + 3\cos\phi\,\mathbf{k}$, $0 \le \phi \le \pi/2$, $0 \le \theta \le 2\pi$. Then $\mathbf{r}_\phi = 3\cos\phi\,\cos\theta\,\mathbf{i} + 3\cos\phi\,\sin\theta\,\mathbf{j} - 3\sin\phi\,\mathbf{k}$, $\mathbf{r}_\theta = -3\sin\phi\sin\theta\,\mathbf{i} + 3\sin\phi\cos\theta\,\mathbf{j}$, and the outward orientation is given by $\mathbf{r}_\phi \times \mathbf{r}_\theta = 9\sin^2\phi\,\cos\theta\,\mathbf{i} + 9\sin^2\phi\,\sin\theta\,\mathbf{j} + 9\sin\phi\,\cos\phi\,\mathbf{k}$. The rate of flow through S is [by (7)]

$$\begin{aligned}\iint_S \rho\mathbf{v}\cdot d\mathbf{S} &= \rho\int_0^{\pi/2}\int_0^{2\pi}(3\sin\phi\,\sin\theta\,\mathbf{i} + 3\sin\phi\,\cos\theta\,\mathbf{j})\cdot\left(9\sin^2\phi\,\cos\theta\,\mathbf{i} + 9\sin^2\phi\sin\theta\,\mathbf{j} + 9\sin\phi\,\cos\phi\,\mathbf{k}\right)d\theta\,d\phi \\ &= 27\rho\int_0^{\pi/2}\int_0^{2\pi}\left(\sin^3\phi\,\sin\theta\,\cos\theta + \sin^3\phi\,\sin\theta\,\cos\theta\right)d\theta\,d\phi = 54\rho\int_0^{\pi/2}\sin^3\phi\,d\phi\int_0^{2\pi}\sin\theta\,\cos\theta\,d\theta \\ &= 54\rho\left[-\tfrac{1}{3}(2+\sin^2\phi)\cos\phi\right]_0^{\pi/2}\left[\tfrac{1}{2}\sin^2\theta\right]_0^{2\pi} = 0\text{ kg/s}\end{aligned}$$

45. S consists of the hemisphere S_1 given by $z = \sqrt{a^2 - x^2 - y^2}$ and the disk S_2 given by $0 \le x^2 + y^2 \le a^2$, $z = 0$.

On S_1: As in Example 4, we use a parametric representation to get $\mathbf{E} = a\sin\phi\,\cos\theta\,\mathbf{i} + a\sin\phi\,\sin\theta\,\mathbf{j} + 2a\cos\phi\,\mathbf{k}$, $\mathbf{T}_\phi \times \mathbf{T}_\theta = a^2\sin^2\phi\,\cos\theta\,\mathbf{i} + a^2\sin^2\phi\,\sin\theta\,\mathbf{j} + a^2\sin\phi\,\cos\phi\,\mathbf{k}$. Thus

$$\begin{aligned}\iint_{S_1}\mathbf{E}\cdot d\mathbf{S} &= \int_0^{2\pi}\int_0^{\pi/2}(a^3\sin^3\phi + 2a^3\sin\phi\,\cos^2\phi)\,d\phi\,d\theta \\ &= \int_0^{2\pi}\int_0^{\pi/2}(a^3\sin\phi + a^3\sin\phi\,\cos^2\phi)\,d\phi\,d\theta = (2\pi)a^3\left(1+\tfrac{1}{3}\right) = \tfrac{8}{3}\pi a^3\end{aligned}$$

On S_2: $\mathbf{E} = x\,\mathbf{i} + y\,\mathbf{j}$, and $\mathbf{r}_y \times \mathbf{r}_x = -\mathbf{k}$ so $\iint_{S_2}\mathbf{E}\cdot d\mathbf{S} = 0$.

Hence, by (11), the total charge is $q = \varepsilon_0\iint_S \mathbf{E}\cdot d\mathbf{S} = \frac{8}{3}\pi a^3\varepsilon_0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

46. Referring to the figure, on

S_1: $\mathbf{E} = \mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$, $\mathbf{r}_y \times \mathbf{r}_z = \mathbf{i}$ and $\iint_{S_1} \mathbf{E} \cdot d\mathbf{S} = \int_{-1}^{1}\int_{-1}^{1} dy\,dz = 4$;

S_2: $\mathbf{E} = x\,\mathbf{i} + \mathbf{j} + z\,\mathbf{k}$, $\mathbf{r}_z \times \mathbf{r}_x = \mathbf{j}$ and $\iint_{S_2} \mathbf{E} \cdot d\mathbf{S} = \int_{-1}^{1}\int_{-1}^{1} dx\,dz = 4$;

S_3: $\mathbf{E} = x\,\mathbf{i} + y\,\mathbf{j} + \mathbf{k}$, $\mathbf{r}_x \times \mathbf{r}_y = \mathbf{k}$ and $\iint_{S_3} \mathbf{E} \cdot d\mathbf{S} = \int_{-1}^{1}\int_{-1}^{1} dx\,dy = 4$;

S_4: $\mathbf{E} = -\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$, $\mathbf{r}_z \times \mathbf{r}_y = -\mathbf{i}$ and $\iint_{S_4} \mathbf{E} \cdot d\mathbf{S} = 4$.

Similarly, $\iint_{S_5} \mathbf{E} \cdot d\mathbf{S} = \iint_{S_6} \mathbf{E} \cdot d\mathbf{S} = 4$. Hence, by (11), $q = \varepsilon_0 \iint_S \mathbf{E} \cdot d\mathbf{S} = \varepsilon_0 \sum_{i=1}^{6} \iint_{S_i} \mathbf{E} \cdot d\mathbf{S} = 24\varepsilon_0$.

47. The heat flow for $K = 6.5$ and $u(x, y, z) = 2y^2 + 2z^2$ is $-K\,\nabla u = -6.5(4y\,\mathbf{j} + 4z\,\mathbf{k})$. As in Example 16.6.5, we can parametrize the cylindrical surface S: $y^2 + z^2 = 6$, $0 \le x \le 4$ as $\mathbf{r}(x, \theta) = x\,\mathbf{i} + \sqrt{6}\,\cos\theta\,\mathbf{j} + \sqrt{6}\,\sin\theta\,\mathbf{k}$, $0 \le x \le 4$, $0 \le \theta \le 2\pi$. Since we want the inward heat flow, we use $\mathbf{r}_x \times \mathbf{r}_\theta = -\sqrt{6}\,\cos\theta\,\mathbf{j} - \sqrt{6}\,\sin\theta\,\mathbf{k}$. Then the rate of heat flow inward is given by

$$\begin{aligned}
\iint_S (-K\,\nabla u) \cdot d\mathbf{S} &= -K \iint \nabla u \cdot (\mathbf{r}_x \times \mathbf{r}_\theta)\,dA \\
&= -6.5 \int_0^{2\pi}\int_0^4 \left(4\sqrt{6}\cos\theta\,\mathbf{j} + 4\sqrt{6}\sin\theta\,\mathbf{k}\right) \cdot \left(-\sqrt{6}\cos\theta\,\mathbf{j} - \sqrt{6}\sin\theta\,\mathbf{k}\right) dx\,d\theta \\
&= -6.5 \int_0^{2\pi}\int_0^4 (-24\cos^2\theta - 24\sin^2\theta)\,dx\,d\theta = -6.5(-24)(4)(2\pi) = 1248\pi
\end{aligned}$$

48. $u(x, y, z) = c/\sqrt{x^2 + y^2 + z^2}$,

$$\begin{aligned}
\mathbf{F} = -K\,\nabla u &= -K\left[-\frac{cx}{(x^2+y^2+z^2)^{3/2}}\,\mathbf{i} - \frac{cy}{(x^2+y^2+z^2)^{3/2}}\,\mathbf{j} - \frac{cz}{(x^2+y^2+z^2)^{3/2}}\,\mathbf{k}\right] \\
&= \frac{cK}{(x^2+y^2+z^2)^{3/2}}\,(x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k})
\end{aligned}$$

and the outward unit normal is $\mathbf{n} = \dfrac{1}{a}\,(x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k})$.

Thus $\mathbf{F} \cdot \mathbf{n} = \dfrac{cK}{a(x^2+y^2+z^2)^{3/2}}\,(x^2 + y^2 + z^2)$, but on S, $x^2 + y^2 + z^2 = a^2$ so $\mathbf{F} \cdot \mathbf{n} = \dfrac{cK}{a^2}$. Hence the rate of heat flow across S is $\displaystyle\iint_S \mathbf{F} \cdot d\mathbf{S} = \frac{cK}{a^2}\iint_S dS = \frac{cK}{a^2}\,(4\pi a^2) = 4\pi K c$.

49. Let S be a sphere of radius a centered at the origin. Then $|\mathbf{r}| = a$ and $\mathbf{F}(\mathbf{r}) = c\mathbf{r}/\,|\mathbf{r}|^3 = \left(c/a^3\right)(x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k})$. A parametric representation for S is $\mathbf{r}(\phi, \theta) = a\sin\phi\,\cos\theta\,\mathbf{i} + a\sin\phi\,\sin\theta\,\mathbf{j} + a\cos\phi\,\mathbf{k}$, $0 \le \phi \le \pi$, $0 \le \theta \le 2\pi$. Then $\mathbf{r}_\phi = a\cos\phi\,\cos\theta\,\mathbf{i} + a\cos\phi\,\sin\theta\,\mathbf{j} - a\sin\phi\,\mathbf{k}$, $\mathbf{r}_\theta = -a\sin\phi\,\sin\theta\,\mathbf{i} + a\sin\phi\,\cos\theta\,\mathbf{j}$, and the outward orientation is given by $\mathbf{r}_\phi \times \mathbf{r}_\theta = a^2\sin^2\phi\,\cos\theta\,\mathbf{i} + a^2\sin^2\phi\,\sin\theta\,\mathbf{j} + a^2\sin\phi\,\cos\phi\,\mathbf{k}$. The flux of $\mathbf{F}$ across S is

$$\begin{aligned}
\iint_S \mathbf{F} \cdot d\mathbf{S} &= \int_0^{\pi}\int_0^{2\pi} \frac{c}{a^3}\,(a\sin\phi\,\cos\theta\,\mathbf{i} + a\sin\phi\,\sin\theta\,\mathbf{j} + a\cos\phi\,\mathbf{k}) \\
&\qquad \cdot \left(a^2\sin^2\phi\,\cos\theta\,\mathbf{i} + a^2\sin^2\phi\,\sin\theta\,\mathbf{j} + a^2\sin\phi\,\cos\phi\,\mathbf{k}\right) d\theta\,d\phi \\
&= \frac{c}{a^3}\int_0^{\pi}\int_0^{2\pi} a^3\left(\sin^3\phi + \sin\phi\,\cos^2\phi\right) d\theta\,d\phi = c\int_0^{\pi}\int_0^{2\pi} \sin\phi\,d\theta\,d\phi = 4\pi c
\end{aligned}$$

Thus the flux does not depend on the radius a.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16.8 Stokes' Theorem

1. Both H and P are oriented piecewise-smooth surfaces that are bounded by the simple, closed, smooth curve $x^2 + y^2 = 4$, $z = 0$ (which we can take to be oriented positively for both surfaces). Then H and P satisfy the hypotheses of Stokes' Theorem, so by (3) we know $\iint_H \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} = \int_C \mathbf{F} \cdot d\mathbf{r} = \iint_P \operatorname{curl} \mathbf{F} \cdot d\mathbf{S}$ (where C is the boundary curve).

2. $\mathbf{F}(x, y, z) = x^2 \sin z\, \mathbf{i} + y^2\, \mathbf{j} + xy\, \mathbf{k}$. The paraboloid $z = 1 - x^2 - y^2$ intersects the xy-plane in the circle $x^2 + y^2 = 1$, $z = 0$. This boundary curve C should be oriented in the counterclockwise direction when viewed from above, so a vector equation of C is $\mathbf{r}(t) = \cos t\, \mathbf{i} + \sin t\, \mathbf{j}$, $0 \le t \le 2\pi$. Then $\mathbf{r}'(t) = -\sin t\, \mathbf{i} + \cos t\, \mathbf{j}$,

$$\mathbf{F}(\mathbf{r}(t)) = (\cos t)^2 (\sin 0)\, \mathbf{i} + (\sin t)^2\, \mathbf{j} + (\cos t)(\sin t)\, \mathbf{k} = \sin^2 t\, \mathbf{j} + \sin t\, \cos t\, \mathbf{k},$$

and by Stokes' Theorem,

$$\begin{aligned}\iint_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} &= \int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\, dt = \int_0^{2\pi} (\sin^2 t\, \mathbf{j} + \sin t\, \cos t\, \mathbf{k}) \cdot (-\sin t\, \mathbf{i} + \cos t\, \mathbf{j})\, dt \\ &= \int_0^{2\pi} (0 + \sin^2 t \cos t + 0)\, dt = \left[\tfrac{1}{3} \sin^3 t\right]_0^{2\pi} = 0\end{aligned}$$

3. $\mathbf{F}(x, y, z) = ze^y\, \mathbf{i} + x \cos y\, \mathbf{j} + xz \sin y\, \mathbf{k}$. The boundary curve C is the circle $x^2 + z^2 = 16$, $y = 0$ where the hemisphere intersects the xz-plane. The curve should be oriented in the counterclockwise direction when viewed from the right (from the positive y-axis), so a vector equation of C is $\mathbf{r}(t) = 4\cos(-t)\, \mathbf{i} + 4\sin(-t)\, \mathbf{k} = 4\cos t\, \mathbf{i} - 4\sin t\, \mathbf{k}$, $0 \le t \le 2\pi$. Then $\mathbf{r}'(t) = -4\sin t\, \mathbf{i} - 4\cos t\, \mathbf{k}$ and

$$\mathbf{F}(\mathbf{r}(t)) = (-4\sin t)e^0\, \mathbf{i} + (4\cos t)(\cos 0)\, \mathbf{j} + (4\cos t)(-4\sin t)(\sin 0)\, \mathbf{k} = -4\sin t\, \mathbf{i} + 4\cos t\, \mathbf{j},$$

and by Stokes' Theorem,

$$\begin{aligned}\iint_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} &= \int_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\, dt = \int_0^{2\pi} (-4\sin t\, \mathbf{i} + 4\cos t\, \mathbf{j}) \cdot (-4\sin t\, \mathbf{i} - 4\cos t\, \mathbf{k})\, dt \\ &= \int_0^{2\pi} (16\sin^2 t + 0 + 0)\, dt = 16\left[\tfrac{1}{2}t - \tfrac{1}{4}\sin 2t\right]_0^{2\pi} = 16\pi\end{aligned}$$

4. $\mathbf{F}(x, y, z) = \tan^{-1}(x^2yz^2)\, \mathbf{i} + x^2y\, \mathbf{j} + x^2z^2\, \mathbf{k}$. The boundary curve C is the circle $y^2 + z^2 = 4$, $x = 2$ which should be oriented in the counterclockwise direction when viewed from the front, so a vector equation of C is $\mathbf{r}(t) = 2\, \mathbf{i} + 2\cos t\, \mathbf{j} + 2\sin t\, \mathbf{k}$, $0 \le t \le 2\pi$. Then $\mathbf{F}(\mathbf{r}(t)) = \tan^{-1}(32\cos t \sin^2 t)\, \mathbf{i} + 8\cos t\, \mathbf{j} + 16\sin^2 t\, \mathbf{k}$, $\mathbf{r}'(t) = -2\sin t\, \mathbf{j} + 2\cos t\, \mathbf{k}$, and $\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = -16\sin t \cos t + 32\sin^2 t \cos t$. Thus,

$$\begin{aligned}\iint_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} &= \oint_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\, dt = \int_0^{2\pi} (-16\sin t \cos t + 32\sin^2 t \cos t)\, dt \\ &= \left[-8\sin^2 t + \tfrac{32}{3}\sin^3 t\right]_0^{2\pi} = 0\end{aligned}$$

5. $\mathbf{F}(x, y, z) = xyz\, \mathbf{i} + xy\, \mathbf{j} + x^2yz\, \mathbf{k}$. C is the square in the plane $z = -1$. Rather than evaluating a line integral around C we can use Equation 3: $\iint_{S_1} \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} = \oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_{S_2} \operatorname{curl} \mathbf{F} \cdot d\mathbf{S}$ where S_1 is the original cube without the bottom and S_2 is the bottom face of the cube. $\operatorname{curl} \mathbf{F} = x^2z\, \mathbf{i} + (xy - 2xyz)\, \mathbf{j} + (y - xz)\, \mathbf{k}$. For S_2, we choose $\mathbf{n} = \mathbf{k}$ so that C has the same orientation for both surfaces. Then $\operatorname{curl} \mathbf{F} \cdot \mathbf{n} = y - xz = x + y$ on S_2, where $z = -1$. Thus, by (16.7.8), $\iint_{S_2} \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} = \int_{-1}^{1} \int_{-1}^{1} (x + y)\, dx\, dy = 0$ so $\iint_{S_1} \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $\mathbf{F}(x,y,z) = e^{xy}\,\mathbf{i} + e^{xz}\,\mathbf{j} + x^2 z\,\mathbf{k}$. The boundary curve C is the circle $x^2+z^2=1$, $y=0$ which should be oriented in the counterclockwise direction when viewed from the right, so a vector equation of C is

$\mathbf{r}(t) = \cos(-t)\,\mathbf{i} + \sin(-t)\,\mathbf{k} = \cos t\,\mathbf{i} - \sin t\,\mathbf{k}$, $0 \le t \le 2\pi$. Then $\mathbf{F}(\mathbf{r}(t)) = \mathbf{i} + e^{-\cos t \sin t}\,\mathbf{j} - \cos^2 t\,\sin t\,\mathbf{k}$,

$\mathbf{r}'(t) = -\sin t\,\mathbf{i} - \cos t\,\mathbf{k}$, and $\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = -\sin t + \cos^3 t\,\sin t$. Thus,

$\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \oint_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_0^{2\pi}(-\sin t + \cos^3 t\,\sin t)\,dt = \left[\cos t - \frac{1}{4}\cos^4 t\right]_0^{2\pi} = 0.$

7. $\mathbf{F}(x,y,z) = (x+y^2)\,\mathbf{i} + (y+z^2)\,\mathbf{j} + (z+x^2)\,\mathbf{k}$ and $\operatorname{curl}\mathbf{F} = -2z\,\mathbf{i} - 2x\,\mathbf{j} - 2y\,\mathbf{k}$. We take the surface to be the planar region enclosed by C and D to be the projection of S onto the xy-plane, so S is the portion of the plane $x+y+z=1$ over $D = \{(x,y) \mid 0 \le x \le 1, 0 \le y \le 1-x\}$. Since C is oriented counterclockwise, we orient S upward. Using Equation 16.7.10, we have $z = g(x,y) = 1-x-y$, $P=-2z$, $Q=-2x$, $R=-2y$, and

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_D [-(-2z)(-1) - (-2x)(-1) + (-2y)]\,dA \\ &= \int_0^1\int_0^{1-x}(-2)\,dy\,dx = -2\int_0^1(1-x)\,dx = -1\end{aligned}$$

8. $\mathbf{F}(x,y,z) = \mathbf{i} + (x+yz)\,\mathbf{j} + \left(xy - \sqrt{z}\right)\mathbf{k}$ and $\operatorname{curl}\mathbf{F} = (x-y)\,\mathbf{i} - y\,\mathbf{j} + \mathbf{k}$. We take the surface S to be the planar region enclosed by C and D to be the projection of S onto the xy-plane, so S is the portion of the plane $3x+2y+z=1$ over $D = \left\{(x,y) \mid 0 \le x \le \frac{1}{3}, 0 \le y \le \frac{1}{2}(1-3x)\right\}$. We orient S upward and use Equation 16.7.10 with $z = g(x,y) = 1-3x-2y$:

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_D [-(x-y)(-3) - (-y)(-2) + 1]\,dA = \int_0^{1/3}\int_0^{(1-3x)/2}(1+3x-5y)\,dy\,dx \\ &= \int_0^{1/3}\left[(1+3x)y - \tfrac{5}{2}y^2\right]_{y=0}^{y=(1-3x)/2}dx = \int_0^{1/3}\left[\tfrac{1}{2}(1+3x)(1-3x) - \tfrac{5}{2}\cdot\tfrac{1}{4}(1-3x)^2\right]dx \\ &= \int_0^{1/3}\left(-\tfrac{81}{8}x^2 + \tfrac{15}{4}x - \tfrac{1}{8}\right)dx = \left[-\tfrac{27}{8}x^3 + \tfrac{15}{8}x^2 - \tfrac{1}{8}x\right]_0^{1/3} = -\tfrac{1}{8} + \tfrac{5}{24} - \tfrac{1}{24} = \tfrac{1}{24}\end{aligned}$$

9. $\mathbf{F}(x,y,z) = xy\,\mathbf{i} + yz\,\mathbf{j} + zx\,\mathbf{k}$. $\operatorname{curl}\mathbf{F} = -y\,\mathbf{i} - z\,\mathbf{j} - x\,\mathbf{k}$ and we take S to be the part of the paraboloid $z = 1-x^2-y^2$ in the first octant. Since C is oriented counterclockwise (from above), we orient S upward. Then using Equation 16.7.10 with $z = g(x,y) = 1-x^2-y^2$ we have

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_D [-(-y)(-2x) - (-z)(-2y) + (-x)]\,dA \\ &= \iint_D \left[-2xy - 2y(1-x^2-y^2) - x\right]dA \\ &= \int_0^{\pi/2}\int_0^1 \left[-2(r\cos\theta)(r\sin\theta) - 2(r\sin\theta)(1-r^2) - r\cos\theta\right] r\,dr\,d\theta \\ &= \int_0^{\pi/2}\int_0^1 \left[-2r^3\sin\theta\cos\theta - 2(r^2-r^4)\sin\theta - r^2\cos\theta\right] dr\,d\theta \\ &= \int_0^{\pi/2}\left[-\tfrac{1}{2}r^4\sin\theta\cos\theta - 2\left(\tfrac{1}{3}r^3 - \tfrac{1}{5}r^5\right)\sin\theta - \tfrac{1}{3}r^3\cos\theta\right]_{r=0}^{r=1} d\theta \\ &= \int_0^{\pi/2}\left(-\tfrac{1}{2}\sin\theta\cos\theta - \tfrac{4}{15}\sin\theta - \tfrac{1}{3}\cos\theta\right)d\theta = \left[-\tfrac{1}{4}\sin^2\theta + \tfrac{4}{15}\cos\theta - \tfrac{1}{3}\sin\theta\right]_0^{\pi/2} \\ &= -\tfrac{1}{4} - \tfrac{4}{15} - \tfrac{1}{3} = -\tfrac{17}{20}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. $\mathbf{F}(x,y,z) = 2y\,\mathbf{i} + xz\,\mathbf{j} + (x+y)\,\mathbf{k}$. The curve of intersection is an ellipse in the plane $z = y + 2$. $\operatorname{curl}\mathbf{F} = (1-x)\,\mathbf{i} - \mathbf{j} + (z-2)\,\mathbf{k}$ and we take the surface S to be the planar region enclosed by C with upward orientation. From Equation 16.7.10 with $z = g(x,y) = y + 2$ we have

$$\begin{aligned}\oint_C \mathbf{F}\cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_{x^2+y^2\le 1} [-(1-x)(0) - (-1)(1) + (y+2-2)]\,dA \\ &= \iint_{x^2+y^2\le 1} (y+1)\,dA = \int_0^{2\pi}\int_0^1 (r\sin\theta + 1)\,r\,dr\,d\theta = \int_0^{2\pi} \left[\tfrac{1}{3}r^3\sin\theta + \tfrac{1}{2}r^2\right]_{r=0}^{r=1} d\theta \\ &= \int_0^{2\pi} \left(\tfrac{1}{3}\sin\theta + \tfrac{1}{2}\right) d\theta = \left[-\tfrac{1}{3}\cos\theta + \tfrac{1}{2}\theta\right]_0^{2\pi} = \pi\end{aligned}$$

11. $\mathbf{F}(x,y,z) = \left\langle -yx^2, xy^2, e^{xy}\right\rangle$ and $\operatorname{curl}\mathbf{F} = xe^{xy}\,\mathbf{i} - ye^{xy}\,\mathbf{j} + (x^2+y^2)\,\mathbf{k}$. C is the circle in the xy-plane centered at the origin with radius 2. Choose S to be the portion of the xy-plane enclosed by C. So $S = D = \{(x,y) \mid x^2+y^2 \le 4\}$. C is oriented counterclockwise, so we orient S upward and the normal vector to S is $\mathbf{n} = \mathbf{k}$. By Stokes' Theorem (See Solution 2 of Example 2), we get

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,d\mathbf{S} = \iint_S [xe^{xy}\,\mathbf{i} - ye^{xy}\,\mathbf{j} + (x^2+y^2)\,\mathbf{k}]\cdot\mathbf{k}\,dS \\ &= \iint_D (x^2+y^2)\,dA = \int_0^2\int_0^{2\pi} r^2\,r\,d\theta\,dr = 2\pi\int_0^2 r^3\,dr = 2\pi\left[\frac{r^4}{4}\right]_0^2 = 8\pi\end{aligned}$$

12. $\mathbf{F}(x,y,z) = ze^x\,\mathbf{i} + (z-y^3)\,\mathbf{j} + (x-z^3)\,\mathbf{k}$ and $\operatorname{curl}\mathbf{F} = -\,\mathbf{i} - (1-e^x)\,\mathbf{j}$. C is the circle $y^2+z^2 = 4$, $x = 3$, and we choose the surface S to be the portion of the plane $x = 3$ enclosed by C. The projection of S onto the yz-plane is the disk $D = \{(y,z) \mid y^2+z^2 \le 4\}$. C is oriented clockwise, so we orient S to have normal vector $\mathbf{n} = \mathbf{i}$. By Stokes' Theorem (see Solution 2 of Example 2), we get

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,d\mathbf{S} = \iint_S [-\mathbf{i} - (1-e^x)\,\mathbf{j}]\cdot\mathbf{i}\,dS \\ &= -\iint_D dA = -A(D) = -\pi(2^2) = -4\pi\end{aligned}$$

13. $\mathbf{F}(x,y,z) = x^2y\,\mathbf{i} + x^3\,\mathbf{j} + e^z\tan^{-1}z\,\mathbf{k}$ and $\operatorname{curl}\mathbf{F} = 2x^2\,\mathbf{k}$. Note that the curve $C = \langle\cos t, \sin t, \sin t\rangle$ is contained in the plane $y = z$ because the $\mathbf{j}$ and $\mathbf{k}$ components of the curve are equal. We choose the surface S to be the portion of the plane $y = z$ enclosed by C. The projection of C onto the xy-plane is the circle $\langle\cos t, \sin t, 0\rangle$, and D is the disk in the xy-plane enclosed by C. We orient S upward and use Equation 16.7.10 with $z = g(x,y) = y$:

$$\begin{aligned}\int_C \mathbf{F}\cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_D [-(0)(0) - (0)(1) + 2x^2]\,dA = \int_0^{2\pi}\int_0^1 2(r\cos\theta)^2\,r\,dr\,d\theta \\ &= \int_0^{2\pi} (1+\cos 2\theta)\,d\theta \int_0^1 r^3\,dr = \left[\theta + \frac{\sin 2\theta}{2}\right]_0^{2\pi}\left[\frac{r^4}{4}\right]_0^1 = (2\pi + 0 - 0)\frac{1}{4} = \frac{\pi}{2}\end{aligned}$$

14. $\mathbf{F}(x,y,z) = \left\langle x^3 - z, xy, y + z^2\right\rangle$ and $\operatorname{curl}\mathbf{F} = \mathbf{i} - \mathbf{j} + y\,\mathbf{k}$. C is the curve of intersection of the paraboloid $z = x^2+y^2$ and the plane $z = x$. Let S be the portion of the plane $z = x$ enclosed by C. To find the projection of S onto the xy-plane, note that $x = x^2+y^2$. Converting to polar coordinates, we get $r\cos\theta = r^2 \quad\Rightarrow\quad r = \cos\theta$. So D is the region in the

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

xy-plane enclosed by the circle $r = \cos\theta$; that is, $D = \{(r,\theta) \mid 0 \le r \le \cos\theta,\ 0 \le \theta \le \pi\}$. We orient S upward and use Equation 16.7.10 with $z = g(x,y) = x$:

$$\int_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_D [-(1)(1) - (-1)(0) + y]\,dA = \iint_D (-1+y)\,dA$$

$$= \int_0^{\pi}\int_0^{\cos\theta} (-1 + r\sin\theta)\,r\,dr\,d\theta = \int_0^{\pi}\left[-\frac{r^2}{2} + \frac{r^3}{3}\sin\theta\right]_0^{\cos\theta} d\theta$$

$$= \int_0^{\pi}\left(-\frac{\cos^2\theta}{2} + \frac{\cos^3\theta}{3}\sin\theta\right) d\theta = \int_0^{\pi}\left(-\frac{1+\cos 2\theta}{4} + \frac{\cos^3\theta}{3}\sin\theta\right) d\theta$$

$$= \left[-\frac{1}{4}\theta - \frac{\sin 2\theta}{8} - \frac{\cos^4\theta}{12}\right]_0^{\pi} = \left(-\frac{\pi}{4} - 0 - \frac{1}{12}\right) - \left(0 - 0 - \frac{1}{12}\right) = -\frac{\pi}{4}$$

15. (a) $\mathbf{F}(x,y,z) = x^2 z\,\mathbf{i} + xy^2\,\mathbf{j} + z^2\,\mathbf{k}$. The curve of intersection is an ellipse in the plane $x + y + z = 1$. The unit normal is $\mathbf{n} = \frac{1}{\sqrt{3}}(\mathbf{i} + \mathbf{j} + \mathbf{k})$, $\operatorname{curl}\mathbf{F} = x^2\,\mathbf{j} + y^2\,\mathbf{k}$, and $\operatorname{curl}\mathbf{F}\cdot\mathbf{n} = \frac{1}{\sqrt{3}}(x^2 + y^2)$. Then, by Stokes' Theorem,

$$\oint_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,d\mathbf{S} = \iint_S \tfrac{1}{\sqrt{3}}\left(x^2 + y^2\right) dS$$

$$= \iint_{x^2+y^2\le 9} \left(x^2 + y^2\right) dx\,dy = \int_0^{2\pi}\int_0^3 r^3\,dr\,d\theta = 2\pi\left(\tfrac{81}{4}\right) = \tfrac{81\pi}{2}$$

(b)

(c) One possible parametrization is $x = 3\cos t$, $y = 3\sin t$, $z = 1 - 3\cos t - 3\sin t$, $0 \le t \le 2\pi$.

16. (a) $\mathbf{F}(x,y,z) = x^2 y\,\mathbf{i} + \frac{1}{3}x^3\,\mathbf{j} + xy\,\mathbf{k}$. S is the part of the surface $z = y^2 - x^2$ that lies above the unit disk D. $\operatorname{curl}\mathbf{F} = x\,\mathbf{i} - y\,\mathbf{j} + (x^2 - x^2)\,\mathbf{k} = x\,\mathbf{i} - y\,\mathbf{j}$. Using Equation 16.7.10 with $z = g(x,y) = y^2 - x^2$, by Stokes' Theorem we have

$$\int_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_D [-x(-2x) - (-y)(2y)]\,dA = 2\iint_D (x^2 + y^2)\,dA$$

$$= 2\int_0^{2\pi}\int_0^1 r^2 r\,dr\,d\theta = 2(2\pi)\left[\tfrac{1}{4}r^4\right]_0^1 = \pi$$

(b)

(c) One possible set of parametric equations is $x = \cos t$, $y = \sin t$, $z = \sin^2 t - \cos^2 t$, $0 \le t \le 2\pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\mathbf{F}(x, y, z) = -y\,\mathbf{i} + x\,\mathbf{j} - 2\,\mathbf{k}$. The boundary curve C is the circle $x^2 + y^2 = 16$, $z = 4$ oriented in the clockwise direction as viewed from above (since S is oriented downward). We can parametrize C by $\mathbf{r}(t) = 4\cos t\,\mathbf{i} - 4\sin t\,\mathbf{j} + 4\,\mathbf{k}$, $0 \le t \le 2\pi$, and then $\mathbf{r}'(t) = -4\sin t\,\mathbf{i} - 4\cos t\,\mathbf{j}$. Thus $\mathbf{F}(\mathbf{r}(t)) = 4\sin t\,\mathbf{i} + 4\cos t\,\mathbf{j} - 2\,\mathbf{k}$, $\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = -16\sin^2 t - 16\cos^2 t = -16$, and

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t)\,dt = \int_0^{2\pi} (-16)\,dt = -16\,(2\pi) = -32\pi$$

Now curl $\mathbf{F} = 2\,\mathbf{k}$, and the projection D of S on the xy-plane is the disk $x^2 + y^2 \le 16$, so by Equation 16.7.10 with $z = g(x, y) = \sqrt{x^2 + y^2}$ [and multiplying by -1 for the downward orientation] we have

$$\iint_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} = -\iint_D (-0 - 0 + 2)\,dA = -2 \cdot A(D) = -2 \cdot \pi(4^2) = -32\pi$$

18. $\mathbf{F}(x, y, z) = -2yz\,\mathbf{i} + y\,\mathbf{j} + 3x\,\mathbf{k}$. The paraboloid intersects the plane $z = 1$ when $1 = 5 - x^2 - y^2 \quad \Leftrightarrow \quad x^2 + y^2 = 4$, so the boundary curve C is the circle $x^2 + y^2 = 4$, $z = 1$ oriented in the counterclockwise direction as viewed from above. We can parametrize C by $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j} + \mathbf{k}$, $0 \le t \le 2\pi$, and then $\mathbf{r}'(t) = -2\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j}$. Thus $\mathbf{F}(\mathbf{r}(t)) = -4\sin t\,\mathbf{i} + 2\sin t\,\mathbf{j} + 6\cos t\,\mathbf{k}$, $\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = 8\sin^2 t + 4\sin t\cos t$, and

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} (8\sin^2 t + 4\sin t\cos t)\,dt = 8\left(\tfrac{1}{2}t - \tfrac{1}{4}\sin 2t\right) + 2\sin^2 t\Big]_0^{2\pi} = 8\pi$$

Now curl $\mathbf{F} = (-3 - 2y)\,\mathbf{j} + 2z\,\mathbf{k}$, and the projection D of S on the xy-plane is the disk $x^2 + y^2 \le 4$, so by Equation 16.7.10 with $z = g(x, y) = 5 - x^2 - y^2$ we have

$$\begin{aligned}
\iint_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} &= \iint_D [-0 - (-3 - 2y)(-2y) + 2z]\,dA = \iint_D [-6y - 4y^2 + 2(5 - x^2 - y^2)]\,dA \\
&= \int_0^{2\pi} \int_0^2 \left[-6r\sin\theta - 4r^2\sin^2\theta + 2(5 - r^2)\right] r\,dr\,d\theta \\
&= \int_0^{2\pi} \left[-2r^3\sin\theta - r^4\sin^2\theta + 5r^2 - \tfrac{1}{2}r^4\right]_{r=0}^{r=2} d\theta = \int_0^{2\pi} \left(-16\sin\theta - 16\sin^2\theta + 20 - 8\right) d\theta \\
&= 16\cos\theta - 16\left(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right) + 12\theta\Big]_0^{2\pi} = 8\pi
\end{aligned}$$

19. $\mathbf{F}(x, y, z) = y\,\mathbf{i} + z\,\mathbf{j} + x\,\mathbf{k}$. The boundary curve C is the circle $x^2 + z^2 = 1$, $y = 0$ oriented in the counterclockwise direction as viewed from the positive y-axis. Then C can be described by $\mathbf{r}(t) = \cos t\,\mathbf{i} - \sin t\,\mathbf{k}$, $0 \le t \le 2\pi$, and $\mathbf{r}'(t) = -\sin t\,\mathbf{i} - \cos t\,\mathbf{k}$. Thus $\mathbf{F}(\mathbf{r}(t)) = -\sin t\,\mathbf{j} + \cos t\,\mathbf{k}$, $\mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) = -\cos^2 t$, and $\oint_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} (-\cos^2 t)\,dt = -\frac{1}{2}t - \frac{1}{4}\sin 2t\Big]_0^{2\pi} = -\pi$. Now curl $\mathbf{F} = -\mathbf{i} - \mathbf{j} - \mathbf{k}$, and S can be parametrized (see Example 16.6.10) by $\mathbf{r}(\phi, \theta) = \sin\phi\,\cos\theta\,\mathbf{i} + \sin\phi\,\sin\theta\,\mathbf{j} + \cos\phi\,\mathbf{k}$, $0 \le \theta \le \pi$, $0 \le \phi \le \pi$. Then $\mathbf{r}_\phi \times \mathbf{r}_\theta = \sin^2\phi\,\cos\theta\,\mathbf{i} + \sin^2\phi\,\sin\theta\,\mathbf{j} + \sin\phi\,\cos\phi\,\mathbf{k}$ and

$$\begin{aligned}
\iint_S \operatorname{curl} \mathbf{F} \cdot d\mathbf{S} &= \iint_{x^2+z^2 \le 1} \operatorname{curl} \mathbf{F} \cdot (\mathbf{r}_\phi \times \mathbf{r}_\theta)\,dA = \int_0^{\pi} \int_0^{\pi} (-\sin^2\phi\,\cos\theta - \sin^2\phi\,\sin\theta - \sin\phi\,\cos\phi)\,d\theta\,d\phi \\
&= \int_0^{\pi} (-2\sin^2\phi - \pi\sin\phi\,\cos\phi)\,d\phi = \left[\tfrac{1}{2}\sin 2\phi - \phi - \tfrac{\pi}{2}\sin^2\phi\right]_0^{\pi} = -\pi
\end{aligned}$$

20. Let S be the surface in the plane $x + y + z = 1$ with upward orientation enclosed by C. Then an upward unit normal vector for S is $\mathbf{n} = \frac{1}{\sqrt{3}}(\mathbf{i} + \mathbf{j} + \mathbf{k})$. Orient C in the counterclockwise direction, as viewed from above. $\int_C z\,dx - 2x\,dy + 3y\,dz$ is equivalent to $\int_C \mathbf{F} \cdot d\mathbf{r}$ for $\mathbf{F}(x, y, z) = z\,\mathbf{i} - 2x\,\mathbf{j} + 3y\,\mathbf{k}$, and the components of $\mathbf{F}$ are polynomials, which have continuous

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

partial derivatives throughout $\mathbb{R}^3$. We have curl $\mathbf{F} = 3\,\mathbf{i} + \mathbf{j} - 2\,\mathbf{k}$, so by Stokes' Theorem,

$$\int_C z\,dx - 2x\,dy + 3y\,dz = \int_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,dS = \iint_S (3\,\mathbf{i} + \mathbf{j} - 2\,\mathbf{k})\cdot\tfrac{1}{\sqrt{3}}\,(\mathbf{i}+\mathbf{j}+\mathbf{k})\;dS$$
$$= \tfrac{2}{\sqrt{3}}\iint_S dS = \tfrac{2}{\sqrt{3}}\,(\text{surface area of } S)$$

Thus the value of $\int_C z\,dx - 2x\,dy + 3y\,dz$ is always $\frac{2}{\sqrt{3}}$ times the area of the region enclosed by C, regardless of its shape or location. [Notice that because $\mathbf{n}$ is normal to a plane, it is constant. But curl $\mathbf{F}$ is also constant, so the dot product $\operatorname{curl}\mathbf{F}\cdot\mathbf{n}$ is constant and we could have simply argued that $\iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,dS$ is a constant multple of $\iint_S dS$, the surface area of S.]

21. $\mathbf{F}(x,y,z) = z^2\,\mathbf{i} + 2xy\,\mathbf{j} + 4y^2\,\mathbf{k}$. It is easier to use Stokes' Theorem than to compute the work directly. Let S be the planar region enclosed by the path of the particle, so S is the portion of the plane $z = \frac{1}{2}y$ for $0 \le x \le 1$, $0 \le y \le 2$, with upward orientation. $\operatorname{curl}\mathbf{F} = 8y\,\mathbf{i} + 2z\,\mathbf{j} + 2y\,\mathbf{k}$ and by Stokes' Theorem and Equation 16.7.10, the work done is

$$W = \oint_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_D \left[-8y\,(0) - 2z\left(\tfrac{1}{2}\right) + 2y\right] dA$$
$$= \int_0^1\int_0^2 \left(2y - \tfrac{1}{2}y\right) dy\,dx = \int_0^1\int_0^2 \tfrac{3}{2}y\,dy\,dx = \int_0^1 \left[\tfrac{3}{4}y^2\right]_{y=0}^{y=2} dx = \int_0^1 3\,dx = 3$$

22. $\int_C (y+\sin x)\,dx + (z^2+\cos y)\,dy + x^3\,dz = \int_C \mathbf{F}\cdot d\mathbf{r}$, where $\mathbf{F}(x,y,z) = (y+\sin x)\,\mathbf{i} + (z^2+\cos y)\,\mathbf{j} + x^3\,\mathbf{k} \quad\Rightarrow$ $\operatorname{curl}\mathbf{F} = -2z\,\mathbf{i} - 3x^2\,\mathbf{j} - \mathbf{k}$. Since $\sin 2t = 2\sin t\,\cos t$, C lies on the surface $z = 2xy$. Let S be the part of this surface that is bounded by C. Then the projection of S onto the xy-plane is the unit disk $D\ \ [x^2+y^2 \le 1]$. C is traversed clockwise (when viewed from above) so S is oriented downward. Using Equation 16.7.10 with $g(x,y) = 2xy$, $P = -2z = -2(2xy) = -4xy$, $Q = -3x^2$, $R = -1$ and multiplying by -1 for the downward orientation, we have

$$\int_C \mathbf{F}\cdot d\mathbf{r} = -\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = -\iint_D \left[-(-4xy)(2y) - (-3x^2)(2x) - 1\right] dA$$
$$= -\iint_D (8xy^2 + 6x^3 - 1)\,dA = -\int_0^{2\pi}\int_0^1 (8r^3\cos\theta\sin^2\theta + 6r^3\cos^3\theta - 1)\,r\,dr\,d\theta$$
$$= -\int_0^{2\pi}\left(\tfrac{8}{5}\cos\theta\sin^2\theta + \tfrac{6}{5}\cos^3\theta - \tfrac{1}{2}\right)d\theta = -\left[\tfrac{8}{15}\sin^3\theta + \tfrac{6}{5}\left(\sin\theta - \tfrac{1}{3}\sin^3\theta\right) - \tfrac{1}{2}\theta\right]_0^{2\pi} = \pi$$

23. Assume S is a sphere centered at the origin with radius a and let H_1 and H_2 be the upper and lower hemispheres, respectively, of S. Then $\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_{H_1} \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} + \iint_{H_2} \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \oint_{C_1}\mathbf{F}\cdot d\mathbf{r} + \oint_{C_2}\mathbf{F}\cdot d\mathbf{r}$ by Stokes' Theorem. But C_1 is the circle $x^2+y^2 = a^2$ oriented in the counterclockwise direction while C_2 is the same circle oriented in the clockwise direction. Hence $\oint_{C_2}\mathbf{F}\cdot d\mathbf{r} = -\oint_{C_1}\mathbf{F}\cdot d\mathbf{r}$ so $\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = 0$ as desired.

24. (a) By Exercise 16.5.28, $\operatorname{curl}(f\nabla g) = f\operatorname{curl}(\nabla g) + \nabla f\times\nabla g = \nabla f\times\nabla g$ since $\operatorname{curl}(\nabla g) = \mathbf{0}$. Hence by Stokes' Theorem $\int_C (f\nabla g)\cdot d\mathbf{r} = \iint_S (\nabla f\times\nabla g)\cdot d\mathbf{S}$.

(b) As in (a), $\operatorname{curl}(f\nabla f) = \nabla f\times\nabla f = \mathbf{0}$, so by Stokes' Theorem, $\int_C (f\nabla f)\cdot d\mathbf{r} = \iint_S [\operatorname{curl}(f\nabla f)]\cdot d\mathbf{S} = 0$.

(c) As in part (a),

$$\operatorname{curl}(f\nabla g + g\nabla f) = \operatorname{curl}(f\nabla g) + \operatorname{curl}(g\nabla f) \qquad \text{[by Exercise 16.5.26]}$$
$$= (\nabla f\times\nabla g) + (\nabla g\times\nabla f) = \mathbf{0} \qquad [\text{since } \mathbf{u}\times\mathbf{v} = -(\mathbf{v}\times\mathbf{u})]$$

Hence by Stokes' Theorem, $\int_C (f\nabla g + g\nabla f)\cdot d\mathbf{r} = \iint_S \operatorname{curl}(f\nabla g + g\nabla f)\cdot d\mathbf{S} = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

16.9 The Divergence Theorem

1. $\mathbf{F}(x,y,z) = 3x\,\mathbf{i} + xy\,\mathbf{j} + 2xz\,\mathbf{k} \quad\Rightarrow\quad \operatorname{div}\mathbf{F} = 3 + x + 2x = 3 + 3x$, so

$\iiint_E \operatorname{div}\mathbf{F}\,dV = \int_0^1\int_0^1\int_0^1 (3x+3)\,dx\,dy\,dz = \frac{9}{2}$ (notice the triple integral is three times the volume of the cube plus three times $\overline{x}$).

To compute $\iint_S \mathbf{F}\cdot d\mathbf{S}$, on

S_1: $x = 1$, $\mathbf{F} = 3\,\mathbf{i} + y\,\mathbf{j} + 2z\,\mathbf{k}$, $\mathbf{n} = \mathbf{i}$, and $\iint_{S_1} \mathbf{F}\cdot d\mathbf{S} = \iint_{S_1} \mathbf{F}\cdot\mathbf{i}\,d\mathbf{S} = \iint_{S_1} 3\,dS = 3$;

S_2: $y = 1$, $\mathbf{F} = 3x\,\mathbf{i} + x\,\mathbf{j} + 2xz\,\mathbf{k}$, $\mathbf{n} = \mathbf{j}$, and $\iint_{S_2} \mathbf{F}\cdot d\mathbf{S} = \iint_{S_2} \mathbf{F}\cdot\mathbf{j}\,d\mathbf{S} = \iint_{S_2} x\,dS = \frac{1}{2}$;

S_3: $z = 1$, $\mathbf{F} = 3x\,\mathbf{i} + xy\,\mathbf{j} + 2x\,\mathbf{k}$, $\mathbf{n} = \mathbf{k}$, and $\iint_{S_3} \mathbf{F}\cdot d\mathbf{S} = \iint_{S_3} \mathbf{F}\cdot\mathbf{k}\,d\mathbf{S} = \iint_{S_3} 2x\,dS = 1$;

S_4: $x = 0$, $\mathbf{F} = \mathbf{0}$, $\mathbf{n} = -\mathbf{i}$, $\iint_{S_4} \mathbf{F}\cdot d\mathbf{S} = 0$;

S_5: $y = 0$, $\mathbf{F} = 3x\,\mathbf{i} + 2xz\,\mathbf{k}$, $\mathbf{n} = -\mathbf{j}$, and $\iint_{S_5} \mathbf{F}\cdot d\mathbf{S} = \iint_{S_5} \mathbf{F}\cdot(-\mathbf{j})\,d\mathbf{S} = \iint_{S_5} 0\,dS = 0$;

S_6: $z = 0$, $\mathbf{F} = 3x\,\mathbf{i} + xy\,\mathbf{j}$, $\mathbf{n} = -\mathbf{k}$, and $\iint_{S_6} \mathbf{F}\cdot d\mathbf{S} = \iint_{S_6} \mathbf{F}\cdot(-\mathbf{k})\,d\mathbf{S} = \iint_{S_6} 0\,dS = 0$.

Thus, $\iint_S \mathbf{F}\cdot d\mathbf{S} = 3 + \frac{1}{2} + 1 + 0 + 0 + 0 = \frac{9}{2}$.

2. $\mathbf{F}(x,y,z) = y^2z^3\,\mathbf{i} + 2yz\,\mathbf{j} + 4z^2\,\mathbf{k} \quad\Rightarrow\quad \operatorname{div}\mathbf{F} = 0 + 2z + 8z = 10z$,

so, using cylindrical coordinates,

$$\begin{aligned}
\iiint_E \operatorname{div}\mathbf{F}\,dV &= \iiint_E 10z\,dV = \int_0^{2\pi}\int_0^3\int_{r^2}^9 (10z)\,r\,dz\,dr\,d\theta \\
&= \int_0^{2\pi}\int_0^3 \left[5rz^2\right]_{z=r^2}^{z=9} dr\,d\theta = \int_0^{2\pi}\int_0^3 (405r - 5r^5)\,dr\,d\theta \\
&= \int_0^{2\pi} d\theta \int_0^3 (405r - 5r^5)\,dr = \left[\theta\right]_0^{2\pi}\left[\tfrac{405}{2}r^2 - \tfrac{5}{6}r^6\right]_0^3 \\
&= 2\pi\left(\tfrac{3645}{2} - \tfrac{1215}{2}\right) = 2430\pi
\end{aligned}$$

On S_1: The surface is $z = x^2 + y^2$, $x^2 + y^2 \le 9$, with downward orientation. Then, by Equation 16.7.10,

$$\begin{aligned}
\iint_{S_1} \mathbf{F}\cdot d\mathbf{S} &= -\iint_D [-(y^2z^3)(2x) - (2yz)(2y) + (4z^2)]\,dA \\
&= \iint_D \left[2xy^2(x^2+y^2)^3 + 4y^2(x^2+y^2) - 4(x^2+y^2)^2\right] dA \\
&= \int_0^{2\pi}\int_0^3 (2r^3\cos\theta\sin^2\theta\cdot r^6 + 4r^2\sin^2\theta\cdot r^2 - 4r^4)\,r\,dr\,d\theta \\
&= \int_0^{2\pi}\int_0^3 (2r^{10}\sin^2\theta\cos\theta + 4r^5\sin^2\theta - 4r^5)\,dr\,d\theta \\
&= \int_0^{2\pi} \left[\tfrac{2}{11}r^{11}\sin^2\theta\cos\theta + \tfrac{2}{3}r^6\sin^2\theta - \tfrac{2}{3}r^6\right]_{r=0}^{r=3} d\theta \\
&= \int_0^{2\pi} \left(\tfrac{354{,}294}{11}\sin^2\theta\cos\theta + 486\sin^2\theta - 486\right) d\theta \\
&= \left[\tfrac{354{,}294}{11}\cdot\tfrac{1}{3}\sin^3\theta + 486\left(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right) - 486\theta\right]_0^{2\pi} \\
&= 0 + 486(\pi - 0) - 486(2\pi) = -486\pi
\end{aligned}$$

On S_2: The surface is $z = 9$, $x^2 + y^2 \le 9$, with upward orientation and $\mathbf{n} = \mathbf{k}$, so

$$\begin{aligned}
\iint_{S_2} \mathbf{F}\cdot d\mathbf{S} &= \iint_{S_2} \mathbf{F}\cdot\mathbf{n}\,d\mathbf{S} = \iint_{S_2} 4z^2\,d\mathbf{S} = \iint_{x^2+y^2\le 9} 4(9)^2\,dA \\
&= 324(\text{area of circle}) = 324\cdot\pi(3)^2 = 2916\pi
\end{aligned}$$

Thus $\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_{S_1} \mathbf{F}\cdot d\mathbf{S} + \iint_{S_2} \mathbf{F}\cdot d\mathbf{S} = -486\pi + 2916\pi = 2430\pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. $\mathbf{F}(x,y,z) = \langle z, y, x\rangle \quad \Rightarrow \quad \operatorname{div}\mathbf{F} = 0+1+0 = 1$, so $\iiint_E \operatorname{div}\mathbf{F}\,dV = \iiint_E 1\,dV = V(E) = \frac{4}{3}\pi\cdot 4^3 = \frac{256}{3}\pi$. S is a sphere of radius 4 centered at the origin which can be parametrized by $\mathbf{r}(\phi,\theta) = \langle 4\sin\phi\cos\theta, 4\sin\phi\sin\theta, 4\cos\phi\rangle$, $0 \le \phi \le \pi$, $0 \le \theta \le 2\pi$ (similar to Example 16.6.10). Then

$$\begin{aligned}\mathbf{r}_\phi \times \mathbf{r}_\theta &= \langle 4\cos\phi\cos\theta, 4\cos\phi\sin\theta, -4\sin\phi\rangle \times \langle -4\sin\phi\sin\theta, 4\sin\phi\cos\theta, 0\rangle \\ &= \left\langle 16\sin^2\phi\cos\theta, 16\sin^2\phi\sin\theta, 16\cos\phi\sin\phi\right\rangle\end{aligned}$$

and $\mathbf{F}(\mathbf{r}(\phi,\theta)) = \langle 4\cos\phi, 4\sin\phi\sin\theta, 4\sin\phi\cos\theta\rangle$. Thus,

$$\begin{aligned}\mathbf{F}\cdot(\mathbf{r}_\phi \times \mathbf{r}_\theta) &= 64\cos\phi\sin^2\phi\cos\theta + 64\sin^3\phi\sin^2\theta + 64\cos\phi\sin^2\phi\cos\theta \\ &= 128\cos\phi\sin^2\phi\cos\theta + 64\sin^3\phi\sin^2\theta\end{aligned}$$

and

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_D \mathbf{F}\cdot(\mathbf{r}_\phi \times \mathbf{r}_\theta)\,dA &= \int_0^{2\pi}\int_0^{\pi}\left(128\cos\phi\sin^2\phi\cos\theta + 64\sin^3\phi\sin^2\theta\right)d\phi\,d\theta \\ &= \int_0^{2\pi}\left[\tfrac{128}{3}\sin^3\phi\cos\theta + 64\left(\tfrac{1}{3}\cos^3\phi - \cos\phi\right)\sin^2\theta\right]_{\phi=0}^{\phi=\pi}d\theta \\ &= \int_0^{2\pi}\tfrac{256}{3}\sin^2\theta\,d\theta = \tfrac{256}{3}\left[\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right]_0^{2\pi} = \tfrac{256}{3}\pi\end{aligned}$$

4. $F(x,y,z) = \left\langle x^2, -y, z\right\rangle \quad \Rightarrow \quad \operatorname{div}\mathbf{F} = 2x - 1 + 1 = 2x$, so

$$\iiint_E \operatorname{div}\mathbf{F}\,dV = \iint_{y^2+z^2\le 9}\left[\int_0^2 2x\,dx\right]dA = \iint_{y^2+z^2\le 9} 4\,dA = 4(\text{area of circle}) = 4(\pi\cdot 3^2) = 36\pi$$

Let S_1 be the front of the cylinder (in the plane $x = 2$), S_2 the back (in the yz-plane), and S_3 the lateral surface of the cylinder. S_1 is the disk $x = 2$, $y^2 + z^2 \le 9$. A unit normal vector is $\mathbf{n} = \langle 1,0,0\rangle$ and $\mathbf{F} = \langle 4, -y, z\rangle$ on S_1, so $\iint_{S_1}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_1}\mathbf{F}\cdot\mathbf{n}\,dS = \iint_{S_1} 4\,dS = 4(\text{surface area of } S_1) = 4(\pi\cdot 3^2) = 36\pi$. S_2 is the disk $x = 0$, $y^2 + z^2 \le 9$. Here $\mathbf{n} = \langle -1,0,0\rangle$ and $\mathbf{F} = \langle 0, -y, z\rangle$, so $\iint_{S_2}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_2}\mathbf{F}\cdot\mathbf{n}\,dS = \iint_{S_2} 0\,dS = 0$.

S_3 can be parametrized by $\mathbf{r}(x,\theta) = \langle x, 3\cos\theta, 3\sin\theta\rangle$, $0 \le x \le 2$, $0 \le \theta \le 2\pi$. Then $\mathbf{r}_x \times \mathbf{r}_\theta = \langle 1,0,0\rangle \times \langle 0, -3\sin\theta, 3\cos\theta\rangle = \langle 0, -3\cos\theta, -3\sin\theta\rangle$. For the outward (positive) orientation we use $-(\mathbf{r}_x \times \mathbf{r}_\theta)$ and $\mathbf{F}(\mathbf{r}(x,\theta)) = \left\langle x^2, -3\cos\theta, 3\sin\theta\right\rangle$, so

$$\begin{aligned}\iint_{S_3}\mathbf{F}\cdot d\mathbf{S} = \iint_D \mathbf{F}\cdot(-(\mathbf{r}_x \times \mathbf{r}_\theta))\,dA &= \int_0^2\int_0^{2\pi}\left(0 - 9\cos^2\theta + 9\sin^2\theta\right)d\theta\,dx \\ &= -9\int_0^2 dx\int_0^{2\pi}\cos 2\theta\,d\theta = -9\,(2)\left[\tfrac{1}{2}\sin 2\theta\right]_0^{2\pi} = 0\end{aligned}$$

Thus $\iint_S \mathbf{F}\cdot d\mathbf{S} = 36\pi + 0 + 0 = 36\pi$.

5. $\mathbf{F}(x,y,z) = xye^z\,\mathbf{i} + xy^2z^3\,\mathbf{j} - ye^z\,\mathbf{k} \quad \Rightarrow$

$\operatorname{div}\mathbf{F} = \frac{\partial}{\partial x}(xye^z) + \frac{\partial}{\partial y}(xy^2z^3) + \frac{\partial}{\partial z}(-ye^z) = ye^z + 2xyz^3 - ye^z = 2xyz^3$, so by the Divergence Theorem,

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}\mathbf{F}\,dV &= \int_0^3\int_0^2\int_0^1 2xyz^3\,dz\,dy\,dx = 2\int_0^3 x\,dx\int_0^2 y\,dy\int_0^1 z^3\,dz \\ &= 2\left[\tfrac{1}{2}x^2\right]_0^3\left[\tfrac{1}{2}y^2\right]_0^2\left[\tfrac{1}{4}z^4\right]_0^1 = 2\left(\tfrac{9}{2}\right)(2)\left(\tfrac{1}{4}\right) = \tfrac{9}{2}\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $\mathbf{F}(x,y,z) = x^2yz\,\mathbf{i} + xy^2z\,\mathbf{j} + xyz^2\,\mathbf{k} \quad \Rightarrow$

$\operatorname{div}\mathbf{F} = \frac{\partial}{\partial x}(x^2yz) + \frac{\partial}{\partial y}(xy^2z) + \frac{\partial}{\partial z}(xyz^2) = 2xyz + 2xyz + 2xyz = 6xyz$, so by the Divergence Theorem,

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}\mathbf{F}\,dV = \int_0^a\int_0^b\int_0^c 6xyz\,dz\,dy\,dx = 6\int_0^a x\,dx\int_0^b y\,dy\int_0^c z\,dz$$
$$= 6\left[\tfrac{1}{2}x^2\right]_0^a\left[\tfrac{1}{2}y^2\right]_0^b\left[\tfrac{1}{2}z^2\right]_0^c = 6\left(\tfrac{1}{2}a^2\right)\left(\tfrac{1}{2}b^2\right)\left(\tfrac{1}{2}c^2\right) = \tfrac{3}{4}a^2b^2c^2$$

7. $\mathbf{F}(x,y,z) = 3xy^2\,\mathbf{i} + xe^z\,\mathbf{j} + z^3\,\mathbf{k} \quad \Rightarrow \quad \operatorname{div}\mathbf{F} = 3y^2 + 0 + 3z^2$, so using cylindrical coordinates with $y = r\cos\theta$, $z = r\sin\theta$, $x = x$ we have, by the Divergence Theorem,

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E (3y^2+3z^2)\,dV = \int_0^{2\pi}\int_0^1\int_{-1}^2 (3r^2\cos^2\theta + 3r^2\sin^2\theta)\,r\,dx\,dr\,d\theta$$
$$= 3\int_0^{2\pi}d\theta\int_0^1 r^3\,dr\int_{-1}^2 dx = 3\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{4}r^4\right]_0^1\left[x\right]_{-1}^2 = 3(2\pi)\left(\tfrac{1}{4}\right)(3) = \tfrac{9\pi}{2}$$

8. $\mathbf{F}(x,y,z) = (x^3+y^3)\,\mathbf{i} + (y^3+z^3)\,\mathbf{j} + (z^3+x^3)\,\mathbf{k} \quad \Rightarrow \quad \operatorname{div}\mathbf{F} = 3x^2 + 3y^2 + 3z^2$, so by the Divergence Theorem,

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E 3(x^2+y^2+z^2)\,dV = \int_0^{\pi}\int_0^{2\pi}\int_0^2 3\rho^2\cdot\rho^2\sin\phi\,d\rho\,d\theta\,d\phi = 3\int_0^{\pi}\sin\phi\,d\phi\int_0^{2\pi}d\theta\int_0^2\rho^4\,d\rho$$
$$= 3\left[-\cos\phi\right]_0^{\pi}\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{5}\rho^5\right]_0^2 = 3\,(2)\,(2\pi)\left(\tfrac{32}{5}\right) = \tfrac{384}{5}\pi$$

9. $\mathbf{F}(x,y,z) = xe^y\,\mathbf{i} + (z-e^y)\,\mathbf{j} - xy\,\mathbf{k} \quad \Rightarrow \quad \operatorname{div}\mathbf{F} = e^y + (-e^y) + 0 = 0$, so by the Divergence Theorem, $\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E 0\,dV = 0$.

10. $\mathbf{F}(x,y,z) = e^y\tan z\,\mathbf{i} + x^2y\,\mathbf{j} + e^x\cos y\,\mathbf{k} \quad \Rightarrow \quad \operatorname{div}\mathbf{F} = x^2$, so by the Divergence Theorem,

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}\mathbf{F}\,dV = \int_{-1}^1\int_{-1}^1\int_0^{2-x-y^3} x^2\,dz\,dx\,dy = \int_{-1}^1\int_{-1}^1 x^2(2-x-y^3)\,dx\,dy$$
$$= \int_{-1}^1\int_{-1}^1 (2x^2 - x^3 - x^2y^3)\,dx\,dy = \int_{-1}^1\int_{-1}^1 (2x^2 - x^2y^3)\,dx\,dy \qquad [x^3 \text{ is odd}]$$
$$= \int_{-1}^1\left[\frac{2}{3}x^3 - \frac{x^3}{3}y^3\right]_{x=-1}^{x=1} dy = \int_{-1}^1\left(\frac{4}{3} - \frac{2}{3}y^3\right)dy = 2\int_0^1\frac{4}{3}\,dy \qquad \left[\begin{matrix}\tfrac{4}{3}\text{ is even,}\\ y^3\text{ is odd}\end{matrix}\right]$$
$$= \frac{8}{3}$$

11. $\mathbf{F}(x,y,z) = (2x^3+y^3)\,\mathbf{i} + (y^3+z^3)\,\mathbf{j} + 3y^2z\,\mathbf{k} \quad \Rightarrow \quad \operatorname{div}\mathbf{F} = 6x^2 + 3y^2 + 3y^2 = 6x^2 + 6y^2$, so

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E 6(x^2+y^2)\,dV = \int_0^{2\pi}\int_0^1\int_0^{1-r^2} 6r^2\cdot r\,dz\,dr\,d\theta = \int_0^{2\pi}\int_0^1 6r^3(1-r^2)\,dr\,d\theta$$
$$= \int_0^{2\pi}d\theta\int_0^1(6r^3 - 6r^5)\,dr = \left[\theta\right]_0^{2\pi}\left[\tfrac{3}{2}r^4 - r^6\right]_0^1 = 2\pi\left(\tfrac{3}{2}-1\right) = \pi$$

12. $\mathbf{F}(x,y,z) = (xy+2xz)\,\mathbf{i} + (x^2+y^2)\,\mathbf{j} + (xy-z^2)\,\mathbf{k}$. For $x^2+y^2 \le 4$ the plane $z = y-2$ is below the xy-plane, so the solid E bounded by S is $E = \left\{(x,y,z) \mid x^2+y^2 \le 4,\ y-2 \le z \le 0\right\}$. Here $\operatorname{div}\mathbf{F} = y + 2z + 2y - 2z = 3y$, so

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E 3y\,dV = \int_0^{2\pi}\int_0^2\int_{r\sin\theta-2}^0 (3r\sin\theta)\,r\,dz\,dr\,d\theta$$
$$= \int_0^{2\pi}\int_0^2(3r^2\sin\theta)(0 - r\sin\theta + 2)\,dr\,d\theta = \int_0^{2\pi}\int_0^2\left(-3r^3\sin^2\theta + 6r^2\sin\theta\right)dr\,d\theta$$
$$= \int_0^{2\pi}\left[-\tfrac{3}{4}r^4\sin^2\theta + 2r^3\sin\theta\right]_{r=0}^{r=2}d\theta = \int_0^{2\pi}\left(-12\sin^2\theta + 16\sin\theta\right)d\theta$$
$$= \left[-12\left(\tfrac{1}{2}\theta - \tfrac{1}{4}\sin 2\theta\right) - 16\cos\theta\right]_0^{2\pi} = -12\pi - 16 + 16 = -12\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $\mathbf{F}(x,y,z) = x^2 z\,\mathbf{i} + xz^3\,\mathbf{j} + y\ln(x+1)\,\mathbf{k} \quad\Rightarrow\quad \operatorname{div}\mathbf{F} = 2xz$, so by the Divergence Theorem,

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iiint_E \operatorname{div}\mathbf{F}\,dV = \int_0^4\int_0^3\int_0^{2-(1/2)x} 2xz\,dz\,dy\,dx = \int_0^4\int_0^3 x\left(2-\tfrac{1}{2}x\right)^2\,dy\,dx\\ &= \int_0^4\int_0^3\left(4x-2x^2+\tfrac{1}{4}x^3\right)\,dy\,dx = 3\int_0^4\left(4x-2x^2+\tfrac{1}{4}x^3\right)dx\\ &= 3\left[2x^2-\tfrac{2}{3}x^3+\tfrac{1}{16}x^4\right]_0^4 = 3\left(32-\tfrac{128}{3}+16\right) = 16\end{aligned}$$

14. $\mathbf{F}(x,y,z) = (xy-z^2)\,\mathbf{i} + x^3\sqrt{z}\,\mathbf{j} + (xy+z^2)\,\mathbf{k} \quad\Rightarrow\quad \operatorname{div}\mathbf{F} = y+2z$, so by the Divergence Theorem,

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iiint_E \operatorname{div}\mathbf{F}\,dV = \int_{-1}^1\int_{y^2}^1\int_0^{1-x}(y+2z)\,dz\,dx\,dy\\ &= \int_{-1}^1\int_{y^2}^1\left[y(1-x)+(1-x)^2\right]dx\,dz = \int_{-1}^1\int_{y^2}^1(y-yx+1-2x+x^2)\,dx\,dy\\ &= \int_{-1}^1\left[yx-y\frac{x^2}{2}+x-x^2+\frac{x^3}{3}\right]_{x=y^2}^{x=1}dy = \int_{-1}^1\left(-\frac{y^6}{3}+\frac{y^5}{2}+y^4-y^3-y^2+\frac{y}{2}+\frac{1}{3}\right)dy\\ &= 2\int_0^1\left(-\frac{y^6}{3}+y^4-y^2+\frac{1}{3}\right)dy = 2\left[-\frac{y^7}{21}+\frac{y^5}{5}-\frac{y^3}{3}+\frac{y}{3}\right]_0^1 = \frac{32}{105}\end{aligned}$$

15. The tetrahedron has vertices $(0,0,0)$, $(a,0,0)$, $(0,b,0)$, $(0,0,c)$ and is described by

$$E = \left\{(x,y,z)\mid 0\le x\le a,\ 0\le y\le b\left(1-\tfrac{x}{a}\right),\ 0\le z\le c\left(1-\tfrac{x}{a}-\tfrac{y}{b}\right)\right\}$$

Here we have $\mathbf{F}(x,y,z) = z\,\mathbf{i} + y\,\mathbf{j} + zx\,\mathbf{k} \quad\Rightarrow\quad \operatorname{div}\mathbf{F} = 0+1+x = x+1$, so

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iiint_E (x+1)\,dV = \int_0^a\int_0^{b\left(1-\frac{x}{a}\right)}\int_0^{c\left(1-\frac{x}{a}-\frac{y}{b}\right)}(x+1)\,dz\,dy\,dx\\ &= \int_0^a\int_0^{b\left(1-\frac{x}{a}\right)}(x+1)\left[c\left(1-\tfrac{x}{a}-\tfrac{y}{b}\right)\right]dy\,dx\\ &= c\int_0^a(x+1)\left[\left(1-\tfrac{x}{a}\right)y-\tfrac{1}{2b}y^2\right]_{y=0}^{y=b\left(1-\frac{x}{a}\right)}dx\\ &= c\int_0^a(x+1)\left[\left(1-\tfrac{x}{a}\right)\cdot b\left(1-\tfrac{x}{a}\right)-\tfrac{1}{2b}\cdot b^2\left(1-\tfrac{x}{a}\right)^2\right]dx = \tfrac{1}{2}bc\int_0^a(x+1)\left(1-\tfrac{x}{a}\right)^2dx\\ &= \tfrac{1}{2}bc\int_0^a\left(\tfrac{1}{a^2}x^3+\tfrac{1}{a^2}x^2-\tfrac{2}{a}x^2+x-\tfrac{2}{a}x+1\right)dx\\ &= \tfrac{1}{2}bc\left[\tfrac{1}{4a^2}x^4+\tfrac{1}{3a^2}x^3-\tfrac{2}{3a}x^3+\tfrac{1}{2}x^2-\tfrac{1}{a}x^2+x\right]_0^a\\ &= \tfrac{1}{2}bc\left(\tfrac{1}{4}a^2+\tfrac{1}{3}a-\tfrac{2}{3}a^2+\tfrac{1}{2}a^2-a+a\right) = \tfrac{1}{2}bc\left(\tfrac{1}{12}a^2+\tfrac{1}{3}a\right) = \tfrac{1}{24}abc(a+4)\end{aligned}$$

16. $\mathbf{r} = x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k} \quad\Rightarrow\quad |\mathbf{r}| = \sqrt{x^2+y^2+z^2} \quad\Rightarrow$

$\mathbf{F}(x,y,z) = |\mathbf{r}|^2\,\mathbf{r} = x(x^2+y^2+z^2)\,\mathbf{i} + y(x^2+y^2+z^2)\,\mathbf{j} + z(x^2+y^2+z^2)\,\mathbf{k} \quad\Rightarrow$

$\operatorname{div}\mathbf{F} = x\cdot 2x + (x^2+y^2+z^2) + y\cdot 2y + (x^2+y^2+z^2) + z\cdot 2z + (x^2+y^2+z^2) = 5(x^2+y^2+z^2)$. Then

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iiint_E 5(x^2+y^2+z^2)\,dV = \int_0^\pi\int_0^{2\pi}\int_0^R 5\rho^2\cdot\rho^2\sin\phi\,d\rho\,d\theta\,d\phi\\ &= 5\int_0^\pi\sin\phi\,d\phi\int_0^{2\pi}d\theta\int_0^R\rho^4\,d\rho = 5\left[-\cos\phi\right]_0^\pi\left[\theta\right]_0^{2\pi}\left[\tfrac{1}{5}\rho^5\right]_0^R = 5\,(2)\,(2\pi)\left(\tfrac{1}{5}R^5\right) = 4\pi R^5\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $\mathbf{r} = x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k} \quad \Rightarrow \quad |\mathbf{r}| = \sqrt{x^2+y^2+z^2} \quad \Rightarrow$

$$\mathbf{F}(x,y,z) = |\mathbf{r}|\,\mathbf{r} = x\sqrt{x^2+y^2+z^2}\,\mathbf{i} + y\sqrt{x^2+y^2+z^2}\,\mathbf{j} + z\sqrt{x^2+y^2+z^2}\,\mathbf{k} \quad \Rightarrow$$

$$\begin{aligned}
\operatorname{div}\mathbf{F} &= x\cdot\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2x) + (x^2+y^2+z^2)^{1/2} \\
&\qquad + y\cdot\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2y) + (x^2+y^2+z^2)^{1/2} \\
&\qquad + z\cdot\tfrac{1}{2}(x^2+y^2+z^2)^{-1/2}(2z) + (x^2+y^2+z^2)^{1/2} \\
&= (x^2+y^2+z^2)^{-1/2}\left[x^2+(x^2+y^2+z^2)+y^2+(x^2+y^2+z^2)+z^2+(x^2+y^2+z^2)\right] \\
&= \frac{4(x^2+y^2+z^2)}{\sqrt{x^2+y^2+z^2}} = 4\sqrt{x^2+y^2+z^2}
\end{aligned}$$

Then
$$\begin{aligned}
\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iiint_E 4\sqrt{x^2+y^2+z^2}\,dV = \int_0^{\pi/2}\int_0^{2\pi}\int_0^1 4\sqrt{\rho^2}\cdot\rho^2\sin\phi\,d\rho\,d\theta\,d\phi \\
&= \textstyle\int_0^{\pi/2}\sin\phi\,d\phi\int_0^{2\pi}d\theta\int_0^1 4\rho^3\,d\rho = [-\cos\phi]_0^{\pi/2}\,[\theta]_0^{2\pi}\,\left[\rho^4\right]_0^1 = (1)\,(2\pi)\,(1) = 2\pi
\end{aligned}$$

18.

By the Divergence Theorem, the flux of $\mathbf{F}$ across the surface of the cube is

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \int_0^{\pi/2}\int_0^{\pi/2}\int_0^{\pi/2}\left[\cos x\,\cos^2 y + 3\sin^2 y\,\cos y\,\cos^4 z + 5\sin^4 z\,\cos z\,\cos^6 x\right]dz\,dy\,dx = \tfrac{19}{64}\pi^2$$

19. For S_1 we have $\mathbf{n} = -\mathbf{k}$, so $\mathbf{F}\cdot\mathbf{n} = \mathbf{F}\cdot(-\mathbf{k}) = -x^2z - y^2 = -y^2$ (since $z = 0$ on S_1). So if D is the unit disk, we get $\iint_{S_1}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_1}\mathbf{F}\cdot\mathbf{n}\,dS = \iint_D(-y^2)\,dA = -\int_0^{2\pi}\int_0^1 r^2\,(\sin^2\theta)\,r\,dr\,d\theta = -\frac{1}{4}\pi$. Now since S_2 is closed, we can use the Divergence Theorem. Since $\operatorname{div}\mathbf{F} = \frac{\partial}{\partial x}\left(z^2x\right) + \frac{\partial}{\partial y}\left(\frac{1}{3}y^3 + \tan^{-1}z\right) + \frac{\partial}{\partial z}\left(x^2z + y^2\right) = z^2 + y^2 + x^2$, we use spherical coordinates to get $\iint_{S_2}\mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}\mathbf{F}\,dV = \int_0^{2\pi}\int_0^{\pi/2}\int_0^1 \rho^2\cdot\rho^2\sin\phi\,d\rho\,d\phi\,d\theta = \frac{2}{5}\pi$. Finally, $\iint_S\mathbf{F}\cdot d\mathbf{S} = \iint_{S_2}\mathbf{F}\cdot d\mathbf{S} - \iint_{S_1}\mathbf{F}\cdot d\mathbf{S} = \frac{2}{5}\pi - \left(-\frac{1}{4}\pi\right) = \frac{13}{20}\pi$.

20. As in the hint to Exercise 19, we create a closed surface $S_2 = S\cup S_1$, where S is the part of the paraboloid $x^2+y^2+z = 2$ that lies above the plane $z = 1$, and S_1 is the disk $x^2+y^2 = 1$ on the plane $z = 1$ oriented downward, and we then apply the Divergence Theorem. Since the disk S_1 is oriented downward, its unit normal vector is $\mathbf{n} = -\mathbf{k}$ and $\mathbf{F}\cdot(-\mathbf{k}) = -z = -1$ on S_1. So $\iint_{S_1}\mathbf{F}\cdot d\mathbf{S} = \iint_{S_1}\mathbf{F}\cdot\mathbf{n}\,dS = \iint_{S_1}(-1)\,dS = -A(S_1) = -\pi$. Let E be the region bounded by S_2. Then $\iint_{S_2}\mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}\mathbf{F}\,dV = \iiint_E 1\,dV = \int_0^1\int_0^{2\pi}\int_1^{2-r^2} r\,dz\,d\theta\,dr = \int_0^1\int_0^{2\pi}(r-r^3)\,d\theta\,dr = (2\pi)\frac{1}{4} = \frac{\pi}{2}$. Thus the flux of $\mathbf{F}$ across S is $\iint_S\mathbf{F}\cdot d\mathbf{S} = \iint_{S_2}\mathbf{F}\cdot d\mathbf{S} - \iint_{S_1}\mathbf{F}\cdot d\mathbf{S} = \frac{\pi}{2} - (-\pi) = \frac{3\pi}{2}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. The vectors that end near P_1 are longer than the vectors that start near P_1, so the net flow is inward near P_1 and $\operatorname{div}\mathbf{F}(P_1)$ is negative. The vectors that end near P_2 are shorter than the vectors that start near P_2, so the net flow is outward near P_2 and $\operatorname{div}\mathbf{F}(P_2)$ is positive.

22. (a) The vectors that end near P_1 are shorter than the vectors that start near P_1, so the net flow is outward and P_1 is a source. The vectors that end near P_2 are longer than the vectors that start near P_2, so the net flow is inward and P_2 is a sink.

(b) $\mathbf{F}(x,y) = \langle x, y^2 \rangle \quad \Rightarrow \quad \operatorname{div}\mathbf{F} = \nabla \cdot \mathbf{F} = 1 + 2y$. The y-value at P_1 is positive, so $\operatorname{div}\mathbf{F} = 1 + 2y$ is positive, thus P_1 is a source. At P_2, $y < -1$, so $\operatorname{div}\mathbf{F} = 1 + 2y$ is negative, and P_2 is a sink.

23.

From the graph it appears that for points above the x-axis, vectors starting near a particular point are longer than vectors ending there, so divergence is positive. The opposite is true at points below the x-axis, where divergence is negative.

$\mathbf{F}(x,y) = \langle xy, x + y^2 \rangle \quad \Rightarrow \quad \operatorname{div}\mathbf{F} = \frac{\partial}{\partial x}(xy) + \frac{\partial}{\partial y}\left(x + y^2\right) = y + 2y = 3y.$

Thus $\operatorname{div}\mathbf{F} > 0$ for $y > 0$, and $\operatorname{div}\mathbf{F} < 0$ for $y < 0$.

24.

From the graph it appears that for points above the line $y = -x$, vectors starting near a particular point are longer than vectors ending there, so divergence is positive. The opposite is true at points below the line $y = -x$, where divergence is negative.

$\mathbf{F}(x,y) = \langle x^2, y^2 \rangle \quad \Rightarrow \quad \operatorname{div}\mathbf{F} = \frac{\partial}{\partial x}(x^2) + \frac{\partial}{\partial y}(y^2) = 2x + 2y$. Then $\operatorname{div}\mathbf{F} > 0$ for $2x + 2y > 0 \quad \Rightarrow \quad y > -x$, and $\operatorname{div}\mathbf{F} < 0$ for $y < -x$.

25. Since $\dfrac{\mathbf{x}}{|\mathbf{x}|^3} = \dfrac{x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}}{(x^2+y^2+z^2)^{3/2}}$ and $\dfrac{\partial}{\partial x}\left(\dfrac{x}{(x^2+y^2+z^2)^{3/2}}\right) = \dfrac{(x^2+y^2+z^2) - 3x^2}{(x^2+y^2+z^2)^{5/2}}$ with similar expressions for $\dfrac{\partial}{\partial y}\left(\dfrac{y}{(x^2+y^2+z^2)^{3/2}}\right)$ and $\dfrac{\partial}{\partial z}\left(\dfrac{z}{(x^2+y^2+z^2)^{3/2}}\right)$, we have

$\operatorname{div}\left(\dfrac{\mathbf{x}}{|\mathbf{x}|^3}\right) = \dfrac{3(x^2+y^2+z^2) - 3(x^2+y^2+z^2)}{(x^2+y^2+z^2)^{5/2}} = 0$, except at $(0,0,0)$ where it is undefined.

26. We first need to find $\mathbf{F}$ so that $\iint_S \mathbf{F}\cdot\mathbf{n}\,dS = \iint_S (2x + 2y + z^2)\,dS$, so $\mathbf{F}\cdot\mathbf{n} = 2x + 2y + z^2$. But for S,

$\mathbf{n} = \dfrac{x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}}{\sqrt{x^2+y^2+z^2}} = x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$. Thus $\mathbf{F} = 2\,\mathbf{i} + 2\,\mathbf{j} + z\,\mathbf{k}$ and $\operatorname{div}\mathbf{F} = 1$.

If $B = \left\{(x,y,z) \mid x^2+y^2+z^2 \le 1\right\}$, then, by the Divergence Theorem,

$$\iint_S (2x + 2y + z^2)\,dS = \iiint_B dV = V(B) = \tfrac{4}{3}\pi(1)^3 = \tfrac{4}{3}\pi$$

27. $\iint_S \mathbf{a}\cdot\mathbf{n}\,dS = \iiint_E \operatorname{div}\mathbf{a}\,dV = 0$ since $\operatorname{div}\mathbf{a} = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. $\frac{1}{3}\iint_S \mathbf{F}\cdot d\mathbf{S} = \frac{1}{3}\iiint_E \operatorname{div}\mathbf{F}\,dV = \frac{1}{3}\iiint_E 3\,dV = V(E)$

29. $\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}(\operatorname{curl}\mathbf{F})\,dV = 0$ by Theorem 16.5.11.

30. $\iint_S D_{\mathbf{n}} f\,dS = \iint_S (\nabla f\cdot\mathbf{n})\,dS = \iiint_E \operatorname{div}(\nabla f)\,dV = \iiint_E \nabla^2 f\,dV$

31. $\iint_S (f\nabla g)\cdot\mathbf{n}\,dS = \iiint_E \operatorname{div}(f\,\nabla g)\,dV$ [by (1) with $\mathbf{F} = f\,\nabla g$]

$= \iiint_E [f\operatorname{div}(\nabla g) + \nabla g\cdot\nabla f]\,dV$ [by Exercise 16.5.27]

$= \iiint_E (f\,\nabla^2 g + \nabla f\cdot\nabla g)\,dV$

32. $\iint_S (f\nabla g - g\nabla f)\cdot\mathbf{n}\,dS = \iiint_E \left[(f\,\nabla^2 g + \nabla f\cdot\nabla g) - (g\nabla^2 f + \nabla g\cdot\nabla f)\right]dV$ [by Exercise 31].

But $\nabla g\cdot\nabla f = \nabla f\cdot\nabla g$, so that $\iint_S (f\,\nabla g - g\nabla f)\cdot\mathbf{n}\,dS = \iiint_E (f\nabla^2 g - g\nabla^2 f)\,dV$.

33. If $\mathbf{c} = c_1\,\mathbf{i} + c_2\,\mathbf{j} + c_3\,\mathbf{k}$ is an arbitrary constant vector, we define $\mathbf{F} = f\mathbf{c} = fc_1\,\mathbf{i} + fc_2\,\mathbf{j} + fc_3\,\mathbf{k}$. Then

$\operatorname{div}\mathbf{F} = \operatorname{div} f\mathbf{c} = \dfrac{\partial f}{\partial x}\,c_1 + \dfrac{\partial f}{\partial y}\,c_2 + \dfrac{\partial f}{\partial z}\,c_3 = \nabla f\cdot\mathbf{c}$ and the Divergence Theorem says $\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}\mathbf{F}\,dV \quad\Rightarrow$

$\iint_S \mathbf{F}\cdot\mathbf{n}\,dS = \iiint_E \nabla f\cdot\mathbf{c}\,dV$. In particular, if $\mathbf{c} = \mathbf{i}$ then $\iint_S f\mathbf{i}\cdot\mathbf{n}\,dS = \iiint_E \nabla f\cdot\mathbf{i}\,dV \quad\Rightarrow$

$\displaystyle\iint_S fn_1\,dS = \iiint_E \frac{\partial f}{\partial x}\,dV$ (where $\mathbf{n} = n_1\,\mathbf{i} + n_2\,\mathbf{j} + n_3\,\mathbf{k}$). Similarly, if $\mathbf{c} = \mathbf{j}$ we have $\displaystyle\iint_S fn_2\,dS = \iiint_E \frac{\partial f}{\partial y}\,dV$,

and $\mathbf{c} = \mathbf{k}$ gives $\displaystyle\iint_S fn_3\,dS = \iiint_E \frac{\partial f}{\partial z}\,dV$. Then

$$\begin{aligned}\iint_S f\mathbf{n}\,dS &= \left(\iint_S fn_1\,dS\right)\mathbf{i} + \left(\iint_S fn_2\,dS\right)\mathbf{j} + \left(\iint_S fn_3\,dS\right)\mathbf{k}\\ &= \left(\iiint_E \frac{\partial f}{\partial x}\,dV\right)\mathbf{i} + \left(\iiint_E \frac{\partial f}{\partial y}\,dV\right)\mathbf{j} + \left(\iiint_E \frac{\partial f}{\partial z}\,dV\right)\mathbf{k} = \iiint_E \left(\frac{\partial f}{\partial x}\,\mathbf{i} + \frac{\partial f}{\partial y}\,\mathbf{j} + \frac{\partial f}{\partial z}\,\mathbf{k}\right)dV\\ &= \iiint_E \nabla f\,dV \quad \text{as desired.}\end{aligned}$$

34. By Exercise 33, $\iint_S p\mathbf{n}\,dS = \iiint_E \nabla p\,dV$, so

$\mathbf{F} = -\iint_S p\mathbf{n}\,dS = -\iiint_E \nabla p\,dV = -\iiint_E \nabla(\rho g z)\,dV = -\iiint_E (\rho g\,\mathbf{k})\,dV = -\rho g\left(\iiint_E dV\right)\mathbf{k} = -\rho g V(E)\,\mathbf{k}$.

But the weight of the displaced liquid is volume $\times$ density $\times\ g = \rho g V(E)$, thus $\mathbf{F} = -W\mathbf{k}$ as desired.

16 Review

TRUE-FALSE QUIZ

1. False. $\operatorname{div}\mathbf{F}$ is a scalar field.

2. True. See Definition 16.5.1.

3. True. Use Theorem 16.5.3 and the fact that $\operatorname{div}\mathbf{0} = 0$.

4. True. See Theorem 16.3.2.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. False. See Exercise 16.3.41. (But the assertion is true if D is simply-connected; see Theorem 16.3.6.)

6. False. See the discussion accompanying Figure 8 in Section 16.2.

7. False. For example, $\operatorname{div}(y\,\mathbf{i}) = 0 = \operatorname{div}(x\,\mathbf{j})$ but $y\,\mathbf{i} \neq x\,\mathbf{j}$.

8. True. Line integrals of conservative vector fields are independent of path, and by Theorem 16.3.3, work $= \int_C \mathbf{F} \cdot d\mathbf{r} = 0$ for any closed path C.

9. True. See Exercise 16.5.26.

10. False. $\mathbf{F} \cdot \mathbf{G}$ is a scalar field, so $\operatorname{curl}(\mathbf{F} \cdot \mathbf{G})$ has no meaning.

11. True. Apply the Divergence Theorem and use the fact that $\operatorname{div} \mathbf{F} = 0$ since $\mathbf{F}$ is a constant vector field.

12. False. See Theorem 16.5.11. If the statement were true, then $\operatorname{div} \operatorname{curl} \mathbf{F}$ would equal $1 + 1 + 1 = 3 \neq 0$.

13. False. By Formulas 16.4.5, the area is given by $-\oint_C y\,dx$ or $\oint_C x\,dy$.

EXERCISES

1. (a) Vectors starting on C point in roughly the direction opposite to C, so the tangential component $\mathbf{F} \cdot \mathbf{T}$ is negative. Thus $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \mathbf{F} \cdot \mathbf{T}\,ds$ is negative.

(b) The vectors that end near P are shorter than the vectors that start near P, so the net flow is outward near P and $\operatorname{div} \mathbf{F}(P)$ is positive.

2. We can parametrize C by $x = x$, $y = x^2$, $0 \le x \le 1$ so

$$\int_C x\,ds = \int_0^1 x\sqrt{1 + (2x)^2}\,dx = \tfrac{1}{12}(1 + 4x^2)^{3/2}\Big]_0^1 = \tfrac{1}{12}\left(5\sqrt{5} - 1\right).$$

3.
$$\begin{aligned}\int_C yz\cos x\,ds &= \int_0^\pi (3\cos t)(3\sin t)\cos t\sqrt{(1)^2 + (-3\sin t)^2 + (3\cos t)^2}\,dt = \int_0^\pi (9\cos^2 t\,\sin t)\sqrt{10}\,dt \\ &= 9\sqrt{10}\left(-\tfrac{1}{3}\cos^3 t\right)\Big]_0^\pi = -3\sqrt{10}\,(-2) = 6\sqrt{10}\end{aligned}$$

4. $x = 3\cos t \quad\Rightarrow\quad dx = -3\sin t\,dt$, $y = 2\sin t \quad\Rightarrow\quad dy = 2\cos t\,dt$, $0 \le t \le 2\pi$, so

$$\begin{aligned}\int_C y\,dx + \left(x + y^2\right)dy &= \int_0^{2\pi}\left[(2\sin t)(-3\sin t) + (3\cos t + 4\sin^2 t)(2\cos t)\right]dt \\ &= \int_0^{2\pi}(-6\sin^2 t + 6\cos^2 t + 8\sin^2 t\,\cos t)\,dt = \int_0^{2\pi}\left[6(\cos^2 t - \sin^2 t) + 8\sin^2 t\,\cos t\right]dt \\ &= \int_0^{2\pi}(6\cos 2t + 8\sin^2 t\,\cos t)\,dt = 3\sin 2t + \tfrac{8}{3}\sin^3 t\Big]_0^{2\pi} = 0\end{aligned}$$

Or: Notice that $\frac{\partial}{\partial y}(y) = 1 = \frac{\partial}{\partial x}\left(x + y^2\right)$, so $\mathbf{F}(x, y) = \left\langle y, x + y^2\right\rangle$ is a conservative vector field. Since C is a closed curve, $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C y\,dx + (x + y^2)\,dy = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

5. $\int_C y^3\,dx + x^2\,dy = \int_{-1}^{1}\left[y^3(-2y) + (1-y^2)^2\right]dy = \int_{-1}^{1}(-y^4 - 2y^2 + 1)\,dy$

$= \left[-\frac{1}{5}y^5 - \frac{2}{3}y^3 + y\right]_{-1}^{1} = -\frac{1}{5} - \frac{2}{3} + 1 - \frac{1}{5} - \frac{2}{3} + 1 = \frac{4}{15}$

6. $\int_C \sqrt{xy}\,dx + e^y\,dy + xz\,dz = \int_0^1 \left(\sqrt{t^4 \cdot t^2}\cdot 4t^3 + e^{t^2}\cdot 2t + t^4 \cdot t^3 \cdot 3t^2\right)dt = \int_0^1 (4t^6 + 2te^{t^2} + 3t^9)\,dt$

$= \left[\frac{4}{7}t^7 + e^{t^2} + \frac{3}{10}t^{10}\right]_0^1 = e - \frac{9}{70}$

7. C: $x = 1 + 2t \;\Rightarrow\; dx = 2\,dt$, $y = 4t \;\Rightarrow\; dy = 4\,dt$, $z = -1 + 3t \;\Rightarrow\; dz = 3\,dt$, $0 \le t \le 1$.

$\int_C xy\,dx + y^2\,dy + yz\,dz = \int_0^1 [(1+2t)(4t)(2) + (4t)^2(4) + (4t)(-1+3t)(3)]\,dt$

$= \int_0^1 (116t^2 - 4t)\,dt = \left[\frac{116}{3}t^3 - 2t^2\right]_0^1 = \frac{116}{3} - 2 = \frac{110}{3}$

8. $\mathbf{F}(x,y) = xy\,\mathbf{i} + x^2\,\mathbf{j}$ and $\mathbf{r}(t) = \sin t\,\mathbf{i} + (1+t)\,\mathbf{j}$, $0 \le t \le \pi \;\Rightarrow\; \mathbf{F}(\mathbf{r}(t)) = (\sin t)(1+t)\,\mathbf{i} + (\sin^2 t)\,\mathbf{j}$,

$\mathbf{r}'(t) = \cos t\,\mathbf{i} + \mathbf{j}$ and

$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^\pi ((1+t)\sin t\cos t + \sin^2 t)\,dt = \int_0^\pi \left(\frac{1}{2}(1+t)\sin 2t + \sin^2 t\right)dt$

$= \left[\frac{1}{2}\left((1+t)\left(-\frac{1}{2}\cos 2t\right) + \frac{1}{4}\sin 2t\right) + \frac{1}{2}t - \frac{1}{4}\sin 2t\right]_0^\pi = \frac{\pi}{4}$

9. $\mathbf{F}(x,y,z) = e^z\,\mathbf{i} + xz\,\mathbf{j} + (x+y)\,\mathbf{k}$ and $\mathbf{r}(t) = t^2\,\mathbf{i} + t^3\,\mathbf{j} - t\,\mathbf{k}$, $0 \le t \le 1 \;\Rightarrow$

$\mathbf{F}(\mathbf{r}(t)) = e^{-t}\,\mathbf{i} + t^2(-t)\,\mathbf{j} + (t^2 + t^3)\,\mathbf{k}$, $\mathbf{r}'(t) = 2t\,\mathbf{i} + 3t^2\,\mathbf{j} - \mathbf{k}$ and

$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1 (2te^{-t} - 3t^5 - (t^2 + t^3))\,dt = \left[-2te^{-t} - 2e^{-t} - \frac{1}{2}t^6 - \frac{1}{3}t^3 - \frac{1}{4}t^4\right]_0^1 = \frac{11}{12} - \frac{4}{e}$.

10. (a) $\mathbf{F}(x,y,z) = z\,\mathbf{i} + x\,\mathbf{j} + y\,\mathbf{k}$ and C: $x = 3 - 3t$, $y = \frac{\pi}{2}t$, $z = 3t$, $0 \le t \le 1$. Then

$W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^1 \left[3t\,\mathbf{i} + (3-3t)\,\mathbf{j} + \frac{\pi}{2}t\,\mathbf{k}\right]\cdot\left[-3\,\mathbf{i} + \frac{\pi}{2}\,\mathbf{j} + 3\,\mathbf{k}\right]dt = \int_0^1 \left[-9t + \frac{3\pi}{2}\right]dt = \frac{1}{2}(3\pi - 9)$.

(b) $W = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{\pi/2}(3\sin t\,\mathbf{i} + 3\cos t\,\mathbf{j} + t\,\mathbf{k})\cdot(-3\sin t\,\mathbf{i} + \mathbf{j} + 3\cos t\,\mathbf{k})\,dt$

$= \int_0^{\pi/2}(-9\sin^2 t + 3\cos t + 3t\cos t)\,dt = \left[-9\left(\frac{1}{2}t - \frac{1}{4}\sin 2t\right) + 3\sin t + 3(t\sin t + \cos t)\right]_0^{\pi/2}$

$= -\frac{9\pi}{4} + 3 + \frac{3\pi}{2} - 3 = -\frac{3\pi}{4}$

11. $\mathbf{F}(x,y) = (1+xy)e^{xy}\,\mathbf{i} + (e^y + x^2e^{xy})\,\mathbf{j} \;\Rightarrow\; \frac{\partial}{\partial y}\left[(1+xy)e^{xy}\right] = 2xe^{xy} + x^2ye^{xy} = \frac{\partial}{\partial x}\left[e^y + x^2e^{xy}\right]$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$, so $\mathbf{F}$ is conservative. Thus there exists a function f such that $\mathbf{F} = \nabla f$. Then $f_y(x,y) = e^y + x^2e^{xy}$ implies $f(x,y) = e^y + xe^{xy} + g(x)$ and then $f_x(x,y) = xye^{xy} + e^{xy} + g'(x) = (1+xy)e^{xy} + g'(x)$. But $f_x(x,y) = (1+xy)e^{xy}$, so $g'(x) = 0 \;\Rightarrow\; g(x) = K$. Thus $f(x,y) = e^y + xe^{xy} + K$ is a potential function for $\mathbf{F}$.

12. $\mathbf{F}(x,y,z) = \sin y\,\mathbf{i} + x\cos y\,\mathbf{j} - \sin z\,\mathbf{k}$ is defined on all of $\mathbb{R}^3$, its components have continuous partial derivatives, and $\operatorname{curl}\mathbf{F} = (0-0)\,\mathbf{i} - (0-0)\,\mathbf{j} + (\cos y - \cos y)\,\mathbf{k} = \mathbf{0}$, so $\mathbf{F}$ is conservative by Theorem 16.5.4. Thus there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x,y,z) = \sin y$ implies $f(x,y,z) = x\sin y + g(y,z)$ and then

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$f_y(x,y,z) = x\cos y + g_y(y,z)$. But $f_y(x,y,z) = x\cos y$, so $g_y(y,z) = 0 \quad \Rightarrow \quad g(y,z) = h(z)$. Then $f(x,y,z) = x\sin y + h(z)$ implies $f_z(x,y,z) = h'(z)$. But $f_z(x,y,z) = -\sin z$, so $h(z) = \cos z + K$. Thus a potential function for $\mathbf{F}$ is $f(x,y,z) = x\sin y + \cos z + K$.

13. $\mathbf{F}(x,y) = (4x^3y^2 - 2xy^3)\,\mathbf{i} + (2x^4y - 3x^2y^2 + 4y^3)\,\mathbf{j} \quad \Rightarrow$

$\frac{\partial}{\partial y}(4x^3y^2 - 2xy^3) = 8x^3y - 6xy^2 = \frac{\partial}{\partial x}(2x^4y - 3x^2y^2 + 4y^3)$ and the domain of $\mathbf{F}$ is $\mathbb{R}^2$, so $\mathbf{F}$ is conservative. Furthermore, $f(x,y) = x^4y^2 - x^2y^3 + y^4$ is a potential function for $\mathbf{F}$. $t = 0$ corresponds to the point $(0,1)$ and $t = 1$ corresponds to $(1,1)$, so $\int_C \mathbf{F}\cdot d\mathbf{r} = f(1,1) - f(0,1) = 1 - 1 = 0$.

14. $\mathbf{F}(x,y,z) = e^y\,\mathbf{i} + (xe^y + e^z)\,\mathbf{j} + ye^z\,\mathbf{k} \quad \Rightarrow \quad \operatorname{curl}\mathbf{F} = (e^z - e^z)\,\mathbf{i} - (0-0)\,\mathbf{j} + (e^y - e^y)\,\mathbf{k} = \mathbf{0}$. The domain of $\mathbf{F}$ is $\mathbb{R}^3$ and the components of $\mathbf{F}$ have continuous partial derivatives, so $\mathbf{F}$ is conservative. Furthermore, we can show that $f(x,y,z) = xe^y + ye^z$ is a potential function for $\mathbf{F}$. Then $\int_C \mathbf{F}\cdot d\mathbf{r} = f(4,0,3) - f(0,2,0) = 4 - 2 = 2$.

15. C_1: $\mathbf{r}(t) = t\,\mathbf{i} + t^2\,\mathbf{j}$, $-1 \le t \le 1$;

C_2: $\mathbf{r}(t) = -t\,\mathbf{i} + \mathbf{j}$, $-1 \le t \le 1$.

Then

$$\begin{aligned}\int_C xy^2\,dx - x^2y\,dy &= \int_{-1}^{1}[t(t^2)^2(1) - t^2(t^2)(2t)]\,dt \\ &\qquad + \int_{-1}^{1}[(-t)(1)^2(-1) - (-t)^2(1)(0)]\,dt \\ &= \int_{-1}^{1}(-t^5)\,dt + \int_{-1}^{1} t\,dt \\ &= 0 \qquad [-t^5 \text{ and } t \text{ are odd}]\end{aligned}$$

Using Green's Theorem, we have

$$\begin{aligned}\int_C xy^2\,dx - x^2y\,dy &= \iint_D \left[\frac{\partial}{\partial x}(-x^2y) - \frac{\partial}{\partial y}(xy^2)\right] dA = \iint_D (-2xy - 2xy)\,dA = \int_{-1}^{1}\int_{x^2}^{1} -4xy\,dy\,dx \\ &= \int_{-1}^{1}\left[-2xy^2\right]_{y=x^2}^{y=1} dx = \int_{-1}^{1}(2x^5 - 2x)\,dx = 0 \qquad [2x^5 - 2x \text{ is odd}]\end{aligned}$$

16. $\int_C \sqrt{1+x^3}\,dx + 2xy\,dy = \iint_D \left[\frac{\partial}{\partial x}(2xy) - \frac{\partial}{\partial y}\left(\sqrt{1+x^3}\right)\right] dA = \int_0^1\int_0^{3x}(2y - 0)\,dy\,dx = \int_0^1 9x^2\,dx = \left[3x^3\right]_0^1 = 3$

17. $\int_C x^2y\,dx - xy^2\,dy = \iint\limits_{x^2+y^2\le 4}\left[\frac{\partial}{\partial x}(-xy^2) - \frac{\partial}{\partial y}(x^2y)\right] dA = \iint\limits_{x^2+y^2\le 4}(-y^2 - x^2)\,dA = -\int_0^{2\pi}\int_0^2 r^3\,dr\,d\theta = -8\pi$

18. $\mathbf{F}(x,y,z) = e^{-x}\sin y\,\mathbf{i} + e^{-y}\sin z\,\mathbf{j} + e^{-z}\sin x\,\mathbf{k} \quad \Rightarrow$

$\operatorname{curl}\mathbf{F} = (0 - e^{-y}\cos z)\,\mathbf{i} - (e^{-z}\cos x - 0)\,\mathbf{j} + (0 - e^{-x}\cos y)\,\mathbf{k} = -e^{-y}\cos z\,\mathbf{i} - e^{-z}\cos x\,\mathbf{j} - e^{-x}\cos y\,\mathbf{k}$, and

$\operatorname{div}\mathbf{F} = -e^{-x}\sin y - e^{-y}\sin z - e^{-z}\sin x$.

19. If we assume there is such a vector field $\mathbf{G}$, then $\operatorname{div}(\operatorname{curl}\mathbf{G}) = \operatorname{div}(2x\,\mathbf{i} + 3yz\,\mathbf{j} - xz^2\,\mathbf{k}) = 2 + 3z - 2xz$. Since $\operatorname{div}(\operatorname{curl}\mathbf{F}) = 0$ for all vector fields $\mathbf{F}$ [by (16.5.11)], such a $\mathbf{G}$ cannot exist.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. Let $\mathbf{F} = P_1\,\mathbf{i} + Q_1\,\mathbf{j} + R_1\,\mathbf{k}$ and $\mathbf{G} = P_2\,\mathbf{i} + Q_2\,\mathbf{j} + R_2\,\mathbf{k}$ be vector fields whose first partials exist and are continuous. Then

$$\begin{aligned}\mathbf{F}\operatorname{div}\mathbf{G} - \mathbf{G}\operatorname{div}\mathbf{F} = {} & \left[P_1\left(\frac{\partial P_2}{\partial x} + \frac{\partial Q_2}{\partial y} + \frac{\partial R_2}{\partial z}\right)\mathbf{i} + Q_1\left(\frac{\partial P_2}{\partial x} + \frac{\partial Q_2}{\partial y} + \frac{\partial R_2}{\partial z}\right)\mathbf{j} + R_1\left(\frac{\partial P_2}{\partial x} + \frac{\partial Q_2}{\partial y} + \frac{\partial R_2}{\partial z}\right)\mathbf{k}\right] \\ & - \left[P_2\left(\frac{\partial P_1}{\partial x} + \frac{\partial Q_1}{\partial y} + \frac{\partial R_1}{\partial z}\right)\mathbf{i} + Q_2\left(\frac{\partial P_1}{\partial x} + \frac{\partial Q_1}{\partial y} + \frac{\partial R_1}{\partial z}\right)\mathbf{j}\right. \\ & \left. + R_2\left(\frac{\partial P_1}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R_1}{\partial z}\right)\mathbf{k}\right]\end{aligned}$$

and

$$\begin{aligned}(\mathbf{G}\cdot\nabla)\,\mathbf{F} - (\mathbf{F}\cdot\nabla)\,\mathbf{G} = {} & \left[\left(P_2\,\frac{\partial P_1}{\partial x} + Q_2\,\frac{\partial P_1}{\partial y} + R_2\,\frac{\partial P_1}{\partial z}\right)\mathbf{i} + \left(P_2\,\frac{\partial Q_1}{\partial x} + Q_2\,\frac{\partial Q_1}{\partial y} + R_2\,\frac{\partial Q_1}{\partial z}\right)\mathbf{j}\right. \\ & \left. + \left(P_2\,\frac{\partial R_1}{\partial x} + Q_2\,\frac{\partial R_1}{\partial y} + R_2\,\frac{\partial R_1}{\partial z}\right)\mathbf{k}\right] \\ & - \left[\left(P_1\,\frac{\partial P_2}{\partial x} + Q_1\,\frac{\partial P_2}{\partial y} + R_1\,\frac{\partial P_2}{\partial z}\right)\mathbf{i} + \left(P_1\,\frac{\partial Q_2}{\partial x} + Q_1\,\frac{\partial Q_2}{\partial y} + R_1\,\frac{\partial Q_2}{\partial z}\right)\mathbf{j}\right. \\ & \left. + \left(P_1\,\frac{\partial R_2}{\partial x} + Q_1\,\frac{\partial R_2}{\partial y} + R_1\,\frac{\partial R_2}{\partial z}\right)\mathbf{k}\right]\end{aligned}$$

Hence

$$\begin{aligned}&\mathbf{F}\operatorname{div}\mathbf{G} - \mathbf{G}\operatorname{div}\mathbf{F} + (\mathbf{G}\cdot\nabla)\,\mathbf{F} - (\mathbf{F}\cdot\nabla)\,\mathbf{G} \\ &\quad = \left[\left(P_1\,\frac{\partial Q_2}{\partial y} + Q_2\,\frac{\partial P_1}{\partial x}\right) - \left(P_2\,\frac{\partial Q_1}{\partial y} + Q_1\,\frac{\partial P_2}{\partial y}\right)\right. \\ &\qquad\qquad \left. - \left(P_2\,\frac{\partial R_1}{\partial z} + R_1\,\frac{\partial P_2}{\partial z}\right) + \left(P_1\,\frac{\partial R_2}{\partial z} + R_2\,\frac{\partial P_1}{\partial z}\right)\right]\mathbf{i} \\ &\qquad + \left[\left(Q_1\,\frac{\partial R_2}{\partial z} + R_2\,\frac{\partial Q_1}{\partial z}\right) - \left(Q_2\,\frac{\partial R_1}{\partial z} + R_1\,\frac{\partial Q_2}{\partial z}\right)\right. \\ &\qquad\qquad \left. - \left(P_1\,\frac{\partial Q_2}{\partial x} + Q_2\,\frac{\partial P_1}{\partial x}\right) + \left(P_2\,\frac{\partial Q_1}{\partial x} + Q_1\,\frac{\partial P_2}{\partial x}\right)\right]\mathbf{j} \\ &\qquad + \left[\left(P_2\,\frac{\partial R_1}{\partial x} + R_1\,\frac{\partial P_2}{\partial x}\right) - \left(P_1\,\frac{\partial R_2}{\partial x} + R_2\,\frac{\partial P_1}{\partial x}\right)\right. \\ &\qquad\qquad \left. - \left(Q_1\,\frac{\partial R_2}{\partial y} + R_2\,\frac{\partial Q_1}{\partial y}\right) + \left(Q_2\,\frac{\partial R_1}{\partial y} + R_1\,\frac{\partial Q_2}{\partial y}\right)\right]\mathbf{k} \\ &\quad = \left[\frac{\partial}{\partial y}\,(P_1Q_2 - P_2Q_1) - \frac{\partial}{\partial z}\,(P_2R_1 - P_1R_2)\right]\mathbf{i} \\ &\qquad + \left[\frac{\partial}{\partial z}\,(Q_1R_2 - Q_2R_1) - \frac{\partial}{\partial x}\,(P_1Q_2 - P_2Q_1)\right]\mathbf{j} \\ &\qquad\qquad + \left[\frac{\partial}{\partial x}\,(P_2R_1 - P_1R_2) - \frac{\partial}{\partial y}\,(Q_1R_2 - Q_2R_1)\right]\mathbf{k} \\ &\quad = \operatorname{curl}(\mathbf{F}\times\mathbf{G})\end{aligned}$$

21. For any piecewise-smooth simple closed plane curve C bounding a region D, we can apply Green's Theorem to $\mathbf{F}(x,y) = f(x)\,\mathbf{i} + g(y)\,\mathbf{j}$ to get $\int_C f(x)\,dx + g(y)\,dy = \iint_D \left[\frac{\partial}{\partial x}\,g(y) - \frac{\partial}{\partial y}\,f(x)\right] dA = \iint_D 0\,dA = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $\nabla^2(fg) = \dfrac{\partial^2(fg)}{\partial x^2} + \dfrac{\partial^2(fg)}{\partial y^2} + \dfrac{\partial^2(fg)}{\partial z^2}$

$$= \frac{\partial}{\partial x}\left(\frac{\partial f}{\partial x}\,g + f\,\frac{\partial g}{\partial x}\right) + \frac{\partial}{\partial y}\left(\frac{\partial f}{\partial y}\,g + f\,\frac{\partial g}{\partial y}\right) + \frac{\partial}{\partial z}\left(\frac{\partial f}{\partial z}\,g + f\,\frac{\partial g}{\partial z}\right) \quad \text{[Product Rule]}$$

$$= \frac{\partial^2 f}{\partial x^2}\,g + 2\,\frac{\partial f}{\partial x}\frac{\partial g}{\partial x} + f\,\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}\,g + 2\,\frac{\partial f}{\partial y}\frac{\partial g}{\partial y}$$

$$+ f\,\frac{\partial^2 g}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}\,g + 2\,\frac{\partial f}{\partial z}\frac{\partial g}{\partial z} + f\,\frac{\partial^2 g}{\partial z^2} \quad \text{[Product Rule]}$$

$$= f\left(\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2} + \frac{\partial^2 g}{\partial z^2}\right) + g\left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}\right) + 2\left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}\right\rangle \cdot \left\langle \frac{\partial g}{\partial x}, \frac{\partial g}{\partial y}, \frac{\partial g}{\partial z}\right\rangle$$

$$= f\nabla^2 g + g\nabla^2 f + 2\nabla f \cdot \nabla g$$

Another method: Using the rules in Exercises 14.6.43(b), 16.5.25, and 16.5.27 [with $\operatorname{div}\mathbf{F} = \nabla\cdot\mathbf{F}$], we have

$$\nabla^2(fg) = \nabla\cdot\nabla(fg) = \nabla\cdot(f\,\nabla g + g\,\nabla f) = f\nabla^2 g + \nabla g\cdot\nabla f + g\nabla^2 f + \nabla f\cdot\nabla g$$
$$= f\,\nabla^2 g + g\,\nabla^2 f + 2\nabla f\cdot\nabla g$$

23. $\nabla^2 f = 0$ means that $\dfrac{\partial^2 f}{\partial x^2} + \dfrac{\partial^2 f}{\partial y^2} = 0$. Now if $\mathbf{F} = f_y\,\mathbf{i} - f_x\,\mathbf{j}$ and C is any closed path in D, then applying Green's Theorem, we get

$$\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C f_y\,dx - f_x\,dy = \iint_D \left[\tfrac{\partial}{\partial x}(-f_x) - \tfrac{\partial}{\partial y}(f_y)\right] dA$$
$$= -\iint_D (f_{xx} + f_{yy})\,dA = -\iint_D 0\,dA = 0$$

Therefore the line integral is independent of path, by Theorem 16.3.3.

24. (a) $x^2 + y^2 = \cos^2 t + \sin^2 t = 1$, so C lies on the circular cylinder $x^2 + y^2 = 1$. But also $y = z$, so C lies on the plane $y = z$. Thus C is contained in the intersection of the plane $y = z$ and the cylinder $x^2 + y^2 = 1$; with $0 \le t \le 2\pi$ we get the entire intersection (an ellipse).

(b) Apply Stokes' Theorem, $\int_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S}$:

$$\operatorname{curl}\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial/\partial x & \partial/\partial y & \partial/\partial z \\ 2xe^{2y} & 2x^2e^{2y} + 2y\cot z & -y^2\csc^2 z \end{vmatrix} = \left\langle -2y\csc^2 z - (-2y\csc^2 z), 0, 4xe^{2y} - 4xe^{2y}\right\rangle = \mathbf{0}$$

Therefore $\int_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \mathbf{0}\cdot d\mathbf{S} = 0$.

25. $z = f(x, y) = x^2 + 2y$ with $0 \le x \le 1$, $0 \le y \le 2x$. Thus

$$A(S) = \iint_D \sqrt{1 + 4x^2 + 4}\,dA = \int_0^1\int_0^{2x} \sqrt{5 + 4x^2}\,dy\,dx = \int_0^1 2x\sqrt{5 + 4x^2}\,dx = \tfrac{1}{6}(5 + 4x^2)^{3/2}\Big]_0^1 = \tfrac{1}{6}\left(27 - 5\sqrt{5}\right).$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. (a) $\mathbf{r}(u,v) = v^2\,\mathbf{i} - uv\,\mathbf{j} + u^2\,\mathbf{k}$, $0 \le u \le 3$, $-3 \le v \le 3 \quad \Rightarrow$

$\mathbf{r}_u = -v\,\mathbf{j} + 2u\,\mathbf{k}$, $\mathbf{r}_v = 2v\,\mathbf{i} - u\,\mathbf{j}$ and

$\mathbf{r}_u \times \mathbf{r}_v = 2u^2\,\mathbf{i} + 4uv\,\mathbf{j} + 2v^2\,\mathbf{k}$. Since the point $(4,-2,1)$ corresponds to $u=1$, $v=2$ (or $u=-1$, $v=-2$ but $\mathbf{r}_u \times \mathbf{r}_v$ is the same for both), a normal vector to the surface at $(4,-2,1)$ is $2\,\mathbf{i} + 8\,\mathbf{j} + 8\,\mathbf{k}$ and an equation of the tangent plane is $2x + 8y + 8z = 0$ or $x + 4y + 4z = 0$.

(b)

(c) By Definition 16.6.6, the area of S is given by

$$A(S) = \iint_D |\,\mathbf{r}_u \times \mathbf{r}_v\,|\,dA = \int_0^3\int_{-3}^3 \sqrt{(2u^2)^2 + (4uv)^2 + (2v^2)^2}\,dv\,du = 2\int_0^3\int_{-3}^3 \sqrt{u^4 + 4u^2v^2 + v^4}\,dv\,du.$$

(d) $\mathbf{F}(x,y,z) = \dfrac{z^2}{1+x^2}\,\mathbf{i} + \dfrac{x^2}{1+y^2}\,\mathbf{j} + \dfrac{y^2}{1+z^2}\,\mathbf{k}$. By Equation 16.7.9, the surface integral is

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iint_D \mathbf{F}\cdot(\,\mathbf{r}_u \times \mathbf{r}_v\,)\,dA = \int_0^3\int_{-3}^3 \left\langle \frac{(u^2)^2}{1+(v^2)^2}, \frac{(v^2)^2}{1+(-uv)^2}, \frac{(-uv)^2}{1+(u^2)^2} \right\rangle \cdot \left\langle 2u^2, 4uv, 2v^2 \right\rangle\,dv\,du$$

$$= \int_0^3\int_{-3}^3 \left(\frac{2u^6}{1+v^4} + \frac{4uv^5}{1+u^2v^2} + \frac{2u^2v^4}{1+u^4} \right) dv\,du \approx 1524.0190$$

27. $z = g(x,y) = x^2 + y^2$ with $0 \le x^2 + y^2 \le 4$. By Formula 16.7.4,

$$\iint_S z\,dS = \iint_{x^2+y^2\le 4} (x^2+y^2)\sqrt{(2x)^2 + (2y)^2 + 1}\,dA = \int_0^{2\pi}\int_0^2 r^2\sqrt{1+4r^2}\,r\,dr\,d\theta$$

$$= \int_0^{2\pi}\int_1^{17} \left(\tfrac{u-1}{4}\right)\sqrt{u}\,\left(\tfrac{1}{8}\,du\right)d\theta \quad \begin{bmatrix} u = 1+4r^2, \\ du = 8r\,dr \end{bmatrix} \quad = \tfrac{1}{32}(2\pi)\left[\tfrac{2}{5}u^{5/2} - \tfrac{2}{3}u^{3/2}\right]_1^{17}$$

$$= \tfrac{\pi}{16}\cdot\tfrac{2}{15}\left[(3u^2 - 5u)\sqrt{u}\right]_1^{17} = \tfrac{\pi}{120}\left[782\sqrt{17} + 2\right] = \tfrac{1}{60}\pi\left(391\sqrt{17} + 1\right)$$

28. $z = g(x,y) = 4 + x + y$ with $0 \le x^2 + y^2 \le 4$. By Formula 16.7.4,

$$\iint_S (x^2z + y^2z)\,dS = \iint_{x^2+y^2\le 4} (x^2+y^2)(4+x+y)\sqrt{1^2+1^2+1}\,dA$$

$$= \int_0^2\int_0^{2\pi} \sqrt{3}\,r^3(4 + r\cos\theta + r\sin\theta)\,d\theta\,dr = \int_0^2 8\pi\sqrt{3}\,r^3\,dr = 32\pi\sqrt{3}$$

29. $\mathbf{F}(x,y,z) = xz\,\mathbf{i} - 2y\,\mathbf{j} + 3x\,\mathbf{k}$. Since the sphere bounds a simple solid region, the Divergence Theorem applies and

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}\mathbf{F}\,dV = \iiint_E (z-2)\,dV = \iiint_E z\,dV - 2\iiint_E dV$$

$$= 0 \begin{bmatrix} \text{odd function in } z \\ \text{and } E \text{ is symmetric} \end{bmatrix} - 2\cdot V(E) = -2\cdot\tfrac{4}{3}\pi(2)^3 = -\tfrac{64}{3}\pi$$

Alternate solution: $\mathbf{F}(\mathbf{r}(\phi,\theta)) = 4\sin\phi\,\cos\theta\,\cos\phi\,\mathbf{i} - 4\sin\phi\,\sin\theta\,\mathbf{j} + 6\sin\phi\,\cos\theta\,\mathbf{k}$,

$\mathbf{r}_\phi \times \mathbf{r}_\theta = 4\sin^2\phi\,\cos\theta\,\mathbf{i} + 4\sin^2\phi\,\sin\theta\,\mathbf{j} + 4\sin\phi\,\cos\phi\,\mathbf{k}$, and

$\mathbf{F}\cdot(\mathbf{r}_\phi \times \mathbf{r}_\theta) = 16\sin^3\phi\,\cos^2\theta\,\cos\phi - 16\sin^3\phi\,\sin^2\theta + 24\sin^2\phi\,\cos\phi\,\cos\theta$. Then

$$\iint_S \mathbf{F}\cdot d\mathbf{S} = \int_0^{2\pi}\int_0^{\pi} (16\sin^3\phi\,\cos\phi\,\cos^2\theta - 16\sin^3\phi\,\sin^2\theta + 24\sin^2\phi\,\cos\phi\,\cos\theta)\,d\phi\,d\theta$$

$$= \int_0^{2\pi} \tfrac{4}{3}(-16\sin^2\theta)\,d\theta = -\tfrac{64}{3}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $\mathbf{F}(x,y,z) = x^2\,\mathbf{i} + xy\,\mathbf{j} + z\,\mathbf{k}$, $z = f(x,y) = x^2 + y^2$, $\mathbf{r}(x,y) = x\,\mathbf{i} + y\,\mathbf{j} + (x^2+y^2)\,\mathbf{k}$, $\mathbf{r}_x \times \mathbf{r}_y = -2x\,\mathbf{i} - 2y\,\mathbf{j} + \mathbf{k}$ (because of upward orientation) and $\mathbf{F}(\mathbf{r}(x,y)) \cdot (\mathbf{r}_x \times \mathbf{r}_y) = -2x^3 - 2xy^2 + x^2 + y^2$. Then, by Formula 16.7.9,

$$\begin{aligned}\iint_S \mathbf{F}\cdot d\mathbf{S} &= \iint\limits_{x^2+y^2\le 1} (-2x^3 - 2xy^2 + x^2 + y^2)\,dA \\ &= \int_0^1\int_0^{2\pi}(-2r^3\cos^3\theta - 2r^3\cos\theta\,\sin^2\theta + r^2)\,r\,dr\,d\theta = \int_0^1 r^3(2\pi)\,dr = \tfrac{\pi}{2}\end{aligned}$$

31. $F(x,y,z) = x^2\,\mathbf{i} + y^2\,\mathbf{j} + z^2\,\mathbf{k}$. Since $\operatorname{curl}\mathbf{F} = \mathbf{0}$, $\iint_S (\operatorname{curl}\mathbf{F})\cdot d\mathbf{S} = 0$. We parametrize C: $\mathbf{r}(t) = \cos t\,\mathbf{i} + \sin t\,\mathbf{j}$, $0 \le t \le 2\pi$. Then $\mathbf{r}'(t) = -\sin t\,\mathbf{i} + \cos t\,\mathbf{j}$ and

$\oint_C \mathbf{F}\cdot d\mathbf{r} = \int_0^{2\pi}\mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t)\,dt = \int_0^{2\pi}(-\cos^2 t\,\sin t + \sin^2 t\,\cos t)\,dt = \left[\frac{1}{3}\cos^3 t + \frac{1}{3}\sin^3 t\right]_0^{2\pi} = 0.$

32. By Stokes' Theorem, $\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \oint_C \mathbf{F}\cdot d\mathbf{r}$, where C is $\mathbf{r}(t) = 2\cos t\,\mathbf{i} + 2\sin t\,\mathbf{j} + \mathbf{k}$, $0 \le t \le 2\pi$.

So $\mathbf{r}'(t) = -2\sin t\,\mathbf{i} + 2\cos t\,\mathbf{j}$, $\mathbf{F}(\mathbf{r}(t)) = 8\cos^2 t\,\sin t\,\mathbf{i} + 2\sin t\,\mathbf{j} + e^{4\cos t\sin t}\,\mathbf{k}$, and

$\mathbf{F}(\mathbf{r}(t))\cdot\mathbf{r}'(t) = -16\cos^2 t\sin^2 t + 4\sin t\cos t$. Thus,

$$\begin{aligned}\oint_C \mathbf{F}\cdot d\mathbf{r} &= \int_0^{2\pi}(-16\cos^2 t\,\sin^2 t + 4\sin t\,\cos t)\,dt \\ &= \left[-16\left(-\tfrac{1}{4}\sin t\,\cos^3 t + \tfrac{1}{16}\sin 2t + \tfrac{1}{8}t\right) + 2\sin^2 t\right]_0^{2\pi} = -4\pi\end{aligned}$$

33. $\mathbf{F}(x,y,z) = xy\,\mathbf{i} + yz\,\mathbf{j} + zx\,\mathbf{k}$. The surface is given by $x+y+z=1$ or $z = 1-x-y$, $0 \le x \le 1$, $0 \le y \le 1-x$. $\mathbf{r}(x,y) = x\,\mathbf{i} + y\,\mathbf{j} + (1-x-y)\,\mathbf{k}$ and $\mathbf{r}_x \times \mathbf{r}_y = \mathbf{i} + \mathbf{j} + \mathbf{k}$. Then, by Formula 16.7.9,

$$\begin{aligned}\oint_C \mathbf{F}\cdot d\mathbf{r} &= \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_D (-y\,\mathbf{i} - z\,\mathbf{j} - x\,\mathbf{k})\cdot(\mathbf{i}+\mathbf{j}+\mathbf{k})\,dA \\ &= \iint_D (-1)\,dA = -(\text{area of } D) = -\tfrac{1}{2}.\end{aligned}$$

34. By the Divergence Theorem,

$\iint_S \mathbf{F}\cdot d\mathbf{S} = \iiint_E 3(x^2+y^2+z^2)\,dV = \int_0^{2\pi}\int_0^1\int_0^2 (3r^2+3z^2)\,r\,dz\,dr\,d\theta = 2\pi\int_0^1(6r^3+8r)\,dr = 11\pi.$

35. $\iiint_E \operatorname{div}\mathbf{F}\,dV = \iiint\limits_{x^2+y^2+z^2\le 1} 3\,dV = 3(\text{volume of sphere}) = 4\pi$. Then

$\mathbf{F}(\mathbf{r}(\phi,\theta))\cdot(\mathbf{r}_\phi \times \mathbf{r}_\theta) = \sin^3\phi\,\cos^2\theta + \sin^3\phi\,\sin^2\theta + \sin\phi\,\cos^2\phi = \sin\phi$ and

$\iint_S \mathbf{F}\cdot d\mathbf{S} = \int_0^{2\pi}\int_0^{\pi}\sin\phi\,d\phi\,d\theta = (2\pi)(2) = 4\pi.$

36. Here we must use Equation 16.9.7 since $\mathbf{F}$ is not defined at the origin. Let S_1 be the sphere of radius 1 with center at the origin and outer unit normal $\mathbf{n}_1 = \mathbf{r}/|\mathbf{r}|$. Let S_2 be the surface of the ellipsoid with outer unit normal $\mathbf{n}_2$ and let E be the solid region between S_1 and S_2. Then the outward flux of $\mathbf{F}$ through the ellipsoid is given by

$\iint_{S_2}\mathbf{F}\cdot\mathbf{n}_2\,dS = -\iint_{S_1}\mathbf{F}\cdot(-\mathbf{n}_1)\,dS + \iiint_E \operatorname{div}\mathbf{F}\,dV$. But $\mathbf{F} = \mathbf{r}/|\mathbf{r}|^3$ with $\mathbf{r} = x\,\mathbf{i} + y\,\mathbf{j} + z\,\mathbf{k}$ and $\mathbf{r} = |\mathbf{r}|$, so

$\operatorname{div}\mathbf{F} = \nabla\cdot\left(|\mathbf{r}|^{-3}\,\mathbf{r}\right) = |\mathbf{r}|^{-3}(\nabla\cdot\mathbf{r}) + \mathbf{r}\cdot\left(\nabla|\mathbf{r}|^{-3}\right) = |\mathbf{r}|^{-3}(3) + \mathbf{r}\cdot\left(-3|\mathbf{r}|^{-4}\right)\left(\mathbf{r}|\mathbf{r}|^{-1}\right) = 0$

[Exercises 16.5.32 and 16.5.33]. And $\mathbf{F}\cdot\mathbf{n}_1 = \dfrac{\mathbf{r}}{|\mathbf{r}|^3}\cdot\dfrac{\mathbf{r}}{|\mathbf{r}|} = |\mathbf{r}|^{-2} = 1$ on S_1.

Thus $\iint_{S_2}\mathbf{F}\cdot\mathbf{n}_2\,dS = \iint_{S_1} dS + \iiint_E 0\,dV = (\text{surface area of the unit sphere}) = 4\pi(1)^2 = 4\pi.$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. Because $\operatorname{curl}\mathbf{F} = \mathbf{0}$, $\mathbf{F}$ is conservative, so there exists a function f such that $\nabla f = \mathbf{F}$. Then $f_x(x,y,z) = 3x^2yz - 3y$ implies $f(x,y,z) = x^3yz - 3xy + g(y,z) \quad \Rightarrow \quad f_y(x,y,z) = x^3z - 3x + g_y(y,z)$. But $f_y(x,y,z) = x^3z - 3x$, so $g(y,z) = h(z)$ and $f(x,y,z) = x^3yz - 3xy + h(z)$. Then $f_z(x,y,z) = x^3y + h'(z)$ but $f_z(x,y,z) = x^3y + 2z$, so $h(z) = z^2 + K$ and a potential function for $\mathbf{F}$ is $f(x,y,z) = x^3yz - 3xy + z^2$. Hence $\int_C \mathbf{F}\cdot d\mathbf{r} = \int_C \nabla f \cdot d\mathbf{r} = f(0,3,0) - f(0,0,2) = 0 - 4 = -4$.

38. Let C' be the circle with center at the origin and radius a as in the figure. Let D be the region bounded by C and C'. Then D's positively oriented boundary is $C \cup (-C')$. Hence by Green's Theorem

$$\int_C \mathbf{F}\cdot d\mathbf{r} + \int_{-C'} \mathbf{F}\cdot d\mathbf{r} = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) dA = 0,$$

so

$$\begin{aligned}
\int_C \mathbf{F}\cdot d\mathbf{r} &= -\int_{-C'} \mathbf{F}\cdot d\mathbf{r} = \int_{C'} \mathbf{F}\cdot d\mathbf{r} = \int_0^{2\pi} \mathbf{F}(\mathbf{r}(t))\cdot \mathbf{r}'(t)\,dt \quad [\mathbf{r}(t) = a\cos t\,\mathbf{i} + a\sin t\,\mathbf{j}] \\
&= \int_0^{2\pi} \left[\frac{2a^3\cos^3 t + 2a^3\cos t\,\sin^2 t - 2a\sin t}{a^2}(-a\sin t) + \frac{2a^3\sin^3 t + 2a^3\cos^2 t\,\sin t + 2a\cos t}{a^2}(a\cos t)\right] dt \\
&= \int_0^{2\pi} \frac{2a^2}{a^2}\,dt = 4\pi
\end{aligned}$$

39. By the Divergence Theorem, $\iint_S \mathbf{F}\cdot\mathbf{n}\,dS = \iiint_E \operatorname{div}\mathbf{F}\,dV = 3(\text{volume of } E) = 3(8-1) = 21$.

40. The stated conditions allow us to use the Divergence Theorem. Hence $\iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iiint_E \operatorname{div}(\operatorname{curl}\mathbf{F})\,dV = 0$ since $\operatorname{div}(\operatorname{curl}\mathbf{F}) = 0$.

41. Let $\mathbf{F} = \mathbf{a}\times\mathbf{r} = \langle a_1, a_2, a_3\rangle \times \langle x,y,z\rangle = \langle a_2z - a_3y, a_3x - a_1z, a_1y - a_2x\rangle$. Then curl $\mathbf{F} = \langle 2a_1, 2a_2, 2a_3\rangle = 2\mathbf{a}$, and $\iint_S 2\mathbf{a}\cdot d\mathbf{S} = \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \int_C \mathbf{F}\cdot d\mathbf{r} = \int_C (\mathbf{a}\times\mathbf{r})\cdot d\mathbf{r}$ by Stokes' Theorem.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

☐ PROBLEMS PLUS

1. Let S_1 be the portion of $\Omega(S)$ between $S(a)$ and S, and let ∂S_1 be its boundary. Also let S_L be the lateral surface of S_1 [that is, the surface of S_1 except S and $S(a)$]. Applying the Divergence Theorem we have $\iint_{\partial S_1} \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = \iiint_{S_1} \nabla\cdot\frac{\mathbf{r}}{r^3}\,dV$.

But
$$\nabla\cdot\frac{\mathbf{r}}{r^3} = \left\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}\right\rangle \cdot \left\langle \frac{x}{(x^2+y^2+z^2)^{3/2}}, \frac{y}{(x^2+y^2+z^2)^{3/2}}, \frac{z}{(x^2+y^2+z^2)^{3/2}}\right\rangle$$
$$= \frac{(x^2+y^2+z^2-3x^2)+(x^2+y^2+z^2-3y^2)+(x^2+y^2+z^2-3z^2)}{(x^2+y^2+z^2)^{5/2}} = 0$$

$\Rightarrow \iint_{\partial S_1} \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = \iiint_{S_1} 0\,dV = 0$. On the other hand, notice that for the surfaces of ∂S_1 other than $S(a)$ and S, $\mathbf{r}\cdot\mathbf{n} = 0 \quad\Rightarrow$

$0 = \iint_{\partial S_1} \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = \iint_S \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS + \iint_{S(a)} \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS + \iint_{S_L} \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = \iint_S \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS + \iint_{S(a)} \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS \quad\Rightarrow$

$\iint_S \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = -\iint_{S(a)} \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS$. Notice that on $S(a)$, $r = a \quad\Rightarrow\quad \mathbf{n} = -\frac{\mathbf{r}}{r} = -\frac{\mathbf{r}}{a}$ and $\mathbf{r}\cdot\mathbf{r} = r^2 = a^2$, so

that $-\iint_{S(a)} \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS = \iint_{S(a)} \frac{\mathbf{r}\cdot\mathbf{r}}{a^4}\,dS = \iint_{S(a)} \frac{a^2}{a^4}\,dS = \frac{1}{a^2}\iint_{S(a)} dS = \frac{\text{area of } S\,(a)}{a^2} = |\Omega(S)|$.

Therefore $|\Omega(S)| = \iint_S \frac{\mathbf{r}\cdot\mathbf{n}}{r^3}\,dS$.

2. By Green's Theorem
$$\int_C (y^3 - y)\,dx - 2x^3\,dy = \iint_D \left[\frac{\partial(-2x^3)}{\partial x} - \frac{\partial(y^3-y)}{\partial y}\right] dA = \iint_D (1 - 6x^2 - 3y^2)\,dA$$

Notice that for $6x^2 + 3y^2 > 1$, the integrand is negative. The integral has maximum value if it is evaluated only in the region where the integrand is positive, which is within the ellipse $6x^2 + 3y^2 = 1$. So the simple closed curve that gives a maximum value for the line integral is the ellipse $6x^2 + 3y^2 = 1$.

3. The given line integral $\frac{1}{2}\int_C (bz - cy)\,dx + (cx - az)\,dy + (ay - bx)\,dz$ can be expressed as $\int_C \mathbf{F}\cdot d\mathbf{r}$ if we define the vector field $\mathbf{F}$ by $\mathbf{F}(x,y,z) = P\,\mathbf{i} + Q\,\mathbf{j} + R\,\mathbf{k} = \frac{1}{2}(bz - cy)\,\mathbf{i} + \frac{1}{2}(cx - az)\,\mathbf{j} + \frac{1}{2}(ay - bx)\,\mathbf{k}$. Then define S to be the planar interior of C, so S is an oriented, smooth surface. Stokes' Theorem says $\int_C \mathbf{F}\cdot d\mathbf{r} = \iint_S \operatorname{curl}\mathbf{F}\cdot d\mathbf{S} = \iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,dS$. Now
$$\operatorname{curl}\mathbf{F} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z}\right)\mathbf{i} + \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x}\right)\mathbf{j} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right)\mathbf{k}$$
$$= \left(\tfrac{1}{2}a + \tfrac{1}{2}a\right)\mathbf{i} + \left(\tfrac{1}{2}b + \tfrac{1}{2}b\right)\mathbf{j} + \left(\tfrac{1}{2}c + \tfrac{1}{2}c\right)\mathbf{k} = a\,\mathbf{i} + b\,\mathbf{j} + c\,\mathbf{k} = \mathbf{n}$$

so $\operatorname{curl}\mathbf{F}\cdot\mathbf{n} = \mathbf{n}\cdot\mathbf{n} = |\mathbf{n}|^2 = 1$, hence $\iint_S \operatorname{curl}\mathbf{F}\cdot\mathbf{n}\,dS = \iint_S dS$ which is simply the surface area of S. Thus, $\int_C \mathbf{F}\cdot d\mathbf{r} = \frac{1}{2}\int_C (bz - cy)\,dx + (cx - az)\,dy + (ay - bx)\,dz$ is the plane area enclosed by C.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. The surface given by $x = \sin u$, $y = \sin v$, $z = \sin(u + v)$ is difficult to visualize, so we first graph the surface from three different points of view.

The trace in the horizontal plane $z = 0$ is given by $z = \sin(u + v) = 0 \quad \Rightarrow \quad u + v = k\pi$ [k an integer]. Then we can write $v = k\pi - u$, and the trace is given by the parametric equations $x = \sin u$, $y = \sin v = \sin(k\pi - u) = \sin k\pi \cos u - \cos k\pi \sin u = \pm \sin u$, and since $\sin u = x$, the trace consists of the two lines $y = \pm x$.

If $z = 1$, $z = \sin(u + v) = 1 \quad \Rightarrow \quad u + v = \frac{\pi}{2} + 2k\pi$. So $v = \left(\frac{\pi}{2} + 2k\pi\right) - u$ and the trace in $z = 1$ is given by the parametric equations $x = \sin u$, $y = \sin v = \sin\left(\left(\frac{\pi}{2} + 2k\pi\right) - u\right) = \sin\left(\frac{\pi}{2} + 2k\pi\right)\cos u - \cos\left(\frac{\pi}{2} + 2k\pi\right)\sin u = \cos u$. This curve is equivalent to $x^2 + y^2 = 1$, $z = 1$, a circle of radius 1. Similarly, in $z = -1$ we have $z = \sin(u + v) = -1 \quad \Rightarrow$ $u + v = \frac{3\pi}{2} + 2k\pi \quad \Rightarrow \quad v = \left(\frac{3\pi}{2} + 2k\pi\right) - u$, so the trace is given by the parametric equations $x = \sin u$, $y = \sin v = \sin\left(\left(\frac{3\pi}{2} + 2k\pi\right) - u\right) = \sin\left(\frac{3\pi}{2} + 2k\pi\right)\cos u - \cos\left(\frac{3\pi}{2} + 2k\pi\right)\sin u = -\cos u$, which again is a circle, $x^2 + y^2 = 1$, $z = -1$.

If $z = \frac{1}{2}$, $z = \sin(u + v) = \frac{1}{2} \quad \Rightarrow \quad u + v = \alpha + 2k\pi$ where $\alpha = \frac{\pi}{6}$ or $\frac{5\pi}{6}$. Then $v = (\alpha + 2k\pi) - u$ and the trace in $z = \frac{1}{2}$ is given by the parametric equations $x = \sin u$, $y = \sin v = \sin[(\alpha + 2k\pi) - u] = \sin(\alpha + 2k\pi)\cos u - \cos(\alpha + 2k\pi)\sin u = \frac{1}{2}\cos u \pm \frac{\sqrt{3}}{2}\sin u$. In rectangular coordinates, $x = \sin u$ so $y = \frac{1}{2}\cos u \pm \frac{\sqrt{3}}{2}x \quad \Rightarrow \quad y \pm \frac{\sqrt{3}}{2}x = \frac{1}{2}\cos u \quad \Rightarrow \quad 2y \pm \sqrt{3}\,x = \cos u$. But then $x^2 + \left(2y \pm \sqrt{3}\,x\right)^2 = \sin^2 u + \cos^2 u = 1 \quad \Rightarrow \quad x^2 + 4y^2 \pm 4\sqrt{3}\,xy + 3x^2 = 1 \quad \Rightarrow \quad 4x^2 \pm 4\sqrt{3}\,xy + 4y^2 = 1$, which may be recognized as a conic section. In particular, each equation is an ellipse rotated $\pm 45^\circ$ from the standard orientation (see the following graph). The trace in $z = -\frac{1}{2}$ is similar: $z = \sin(u + v) = -\frac{1}{2} \quad \Rightarrow \quad u + v = \beta + 2k\pi$ where $\beta = \frac{7\pi}{6}$ or $\frac{11\pi}{6}$. Then $v = (\beta + 2k\pi) - u$ and the trace is given by the parametric equations $x = \sin u$, $y = \sin v = \sin[(\beta + 2k\pi) - u] = \sin(\beta + 2k\pi)\cos u - \cos(\beta + 2k\pi)\sin u = -\frac{1}{2}\cos u \pm \frac{\sqrt{3}}{2}\sin u$. If we convert to rectangular coordinates, we arrive at the same pair of equations, $4x^2 \pm 4\sqrt{3}\,xy + 4y^2 = 1$, so the trace is identical to the trace in $z = \frac{1}{2}$.

Graphing each of these, we have the following 5 traces.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$z = -1$ $\qquad$ $z = -\frac{1}{2}$ $\qquad$ $z = 0$

$z = \frac{1}{2}$ $\qquad$ $z = 1$

Visualizing these traces on the surface reveals that horizontal cross sections are pairs of intersecting ellipses whose major axes are perpendicular to each other. At the bottom of the surface, $z = -1$, the ellipses coincide as circles of radius 1. As we move up the surface, the ellipses become narrower until at $z = 0$ they collapse into line segments, after which the process is reversed, and the ellipses widen to again coincide as circles at $z = 1$.

5. $$(\mathbf{F} \cdot \nabla)\,\mathbf{G} = \left(P_1 \frac{\partial}{\partial x} + Q_1 \frac{\partial}{\partial y} + R_1 \frac{\partial}{\partial z}\right)(P_2\,\mathbf{i} + Q_2\,\mathbf{j} + R_2\,\mathbf{k})$$

$$= \left(P_1 \frac{\partial P_2}{\partial x} + Q_1 \frac{\partial P_2}{\partial y} + R_1 \frac{\partial P_2}{\partial z}\right)\mathbf{i} + \left(P_1 \frac{\partial Q_2}{\partial x} + Q_1 \frac{\partial Q_2}{\partial y} + R_1 \frac{\partial Q_2}{\partial z}\right)\mathbf{j}$$

$$+ \left(P_1 \frac{\partial R_2}{\partial x} + Q_1 \frac{\partial R_2}{\partial y} + R_1 \frac{\partial R_2}{\partial z}\right)\mathbf{k}$$

$$= (\mathbf{F} \cdot \nabla P_2)\,\mathbf{i} + (\mathbf{F} \cdot \nabla Q_2)\,\mathbf{j} + (\mathbf{F} \cdot \nabla R_2)\,\mathbf{k}.$$

Similarly, $(\mathbf{G} \cdot \nabla)\,\mathbf{F} = (\mathbf{G} \cdot \nabla P_1)\,\mathbf{i} + (\mathbf{G} \cdot \nabla Q_1)\,\mathbf{j} + (\mathbf{G} \cdot \nabla R_1)\,\mathbf{k}$. Then

$$\mathbf{F} \times \operatorname{curl} \mathbf{G} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ P_1 & Q_1 & R_1 \\ \partial R_2/\partial y - \partial Q_2/\partial z & \partial P_2/\partial z - \partial R_2/\partial x & \partial Q_2/\partial x - \partial P_2/\partial y \end{vmatrix}$$

$$= \left(Q_1 \frac{\partial Q_2}{\partial x} - Q_1 \frac{\partial P_2}{\partial y} - R_1 \frac{\partial P_2}{\partial z} + R_1 \frac{\partial R_2}{\partial x}\right)\mathbf{i} + \left(R_1 \frac{\partial R_2}{\partial y} - R_1 \frac{\partial Q_2}{\partial z} - P_1 \frac{\partial Q_2}{\partial x} + P_1 \frac{\partial P_2}{\partial y}\right)\mathbf{j}$$

$$+ \left(P_1 \frac{\partial P_2}{\partial z} - P_1 \frac{\partial R_2}{\partial x} - Q_1 \frac{\partial R_2}{\partial y} + Q_1 \frac{\partial Q_2}{\partial z}\right)\mathbf{k}$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

and

$$\mathbf{G} \times \operatorname{curl} \mathbf{F} = \left(Q_2 \frac{\partial Q_1}{\partial x} - Q_2 \frac{\partial P_1}{\partial y} - R_2 \frac{\partial P_1}{\partial z} + R_2 \frac{\partial R_1}{\partial x}\right) \mathbf{i} + \left(R_2 \frac{\partial R_1}{\partial y} - R_2 \frac{\partial Q_1}{\partial z} - P_2 \frac{\partial Q_1}{\partial x} + P_2 \frac{\partial P_1}{\partial y}\right) \mathbf{j} + \left(P_2 \frac{\partial P_1}{\partial z} - P_2 \frac{\partial R_1}{\partial x} - Q_2 \frac{\partial R_1}{\partial y} + Q_2 \frac{\partial Q_1}{\partial z}\right) \mathbf{k}.$$

Then

$$(\mathbf{F} \cdot \nabla)\, \mathbf{G} + \mathbf{F} \times \operatorname{curl} \mathbf{G} = \left(P_1 \frac{\partial P_2}{\partial x} + Q_1 \frac{\partial Q_2}{\partial x} + R_1 \frac{\partial R_2}{\partial x}\right) \mathbf{i} + \left(P_1 \frac{\partial P_2}{\partial y} + Q_1 \frac{\partial Q_2}{\partial y} + R_1 \frac{\partial R_2}{\partial y}\right) \mathbf{j} + \left(P_1 \frac{\partial P_2}{\partial z} + Q_1 \frac{\partial Q_2}{\partial z} + R_1 \frac{\partial R_2}{\partial z}\right) \mathbf{k}$$

and

$$(\mathbf{G} \cdot \nabla)\, \mathbf{F} + \mathbf{G} \times \operatorname{curl} \mathbf{F} = \left(P_2 \frac{\partial P_1}{\partial x} + Q_2 \frac{\partial Q_1}{\partial x} + R_2 \frac{\partial R_1}{\partial x}\right) \mathbf{i} + \left(P_2 \frac{\partial P_1}{\partial y} + Q_2 \frac{\partial Q_1}{\partial y} + R_2 \frac{\partial R_1}{\partial y}\right) \mathbf{j} + \left(P_2 \frac{\partial P_1}{\partial z} + Q_2 \frac{\partial Q_1}{\partial z} + R_2 \frac{\partial R_1}{\partial z}\right) \mathbf{k}.$$

Hence

$$\begin{aligned}
&(\mathbf{F} \cdot \nabla)\, \mathbf{G} + \mathbf{F} \times \operatorname{curl} \mathbf{G} + (\mathbf{G} \cdot \nabla)\, \mathbf{F} + \mathbf{G} \times \operatorname{curl} \mathbf{F} \\
&\quad = \left[\left(P_1 \frac{\partial P_2}{\partial x} + P_2 \frac{\partial P_1}{\partial x}\right) + \left(Q_1 \frac{\partial Q_2}{\partial x} + Q_2 \frac{\partial Q_1}{\partial y}\right) + \left(R_1 \frac{\partial R_2}{\partial x} + R_2 \frac{\partial R_1}{\partial x}\right)\right] \mathbf{i} \\
&\qquad + \left[\left(P_1 \frac{\partial P_2}{\partial y} + P_2 \frac{\partial P_1}{\partial y}\right) + \left(Q_1 \frac{\partial Q_2}{\partial y} + Q_2 \frac{\partial Q_1}{\partial y}\right) + \left(R_1 \frac{\partial R_2}{\partial y} + R_2 \frac{\partial R_1}{\partial y}\right)\right] \mathbf{j} \\
&\qquad + \left[\left(P_1 \frac{\partial P_2}{\partial z} + P_2 \frac{\partial P_1}{\partial z}\right) + \left(Q_1 \frac{\partial Q_2}{\partial z} + Q_2 \frac{\partial Q_1}{\partial z}\right) + \left(R_1 \frac{\partial R_2}{\partial z} + R_2 \frac{\partial R_1}{\partial z}\right)\right] \mathbf{k} \\
&\quad = \nabla(P_1 P_2 + Q_1 Q_2 + R_1 R_2) = \nabla(\mathbf{F} \cdot \mathbf{G}).
\end{aligned}$$

6. (a) First we place the piston on coordinate axes so the top of the cylinder is at the origin and $x(t) \geq 0$ is the distance from the top of the cylinder to the piston at time t. Let C_1 be the curve traced out by the piston during one four-stroke cycle, so C_1 is given by $\mathbf{r}(t) = x(t)\,\mathbf{i}$, $a \leq t \leq b$. (Thus, the curve lies on the positive x-axis and reverses direction several times.) The force on the piston is $AP(t)\,\mathbf{i}$, where A is the area of the top of the piston and $P(t)$ is the pressure in the cylinder at time t. As in Section 16.2, the work done on the piston is $\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_a^b AP(t)\,\mathbf{i} \cdot x'(t)\,\mathbf{i}\,dt = \int_a^b AP(t)\,x'(t)\,dt$. Here, the volume of the cylinder at time t is $V(t) = Ax(t) \quad \Rightarrow \quad V'(t) = Ax'(t) \quad \Rightarrow \quad \int_a^b AP(t)\,x'(t)\,dt = \int_a^b P(t)\,V'(t)\,dt$. Since the curve C in the PV-plane corresponds to the values of P and V at time t, $a \leq t \leq b$, we have

$$W = \int_a^b AP(t)\,x'(t)\,dt = \int_a^b P(t)\,V'(t)\,dt = \int_C P\,dV$$

Another method: If we divide the time interval $[a, b]$ into n subintervals of equal length Δt, the amount of work done on the piston in the ith time interval is approximately $AP(t_i)[x(t_i) - x(t_{i-1})]$. Thus we estimate the total work done during

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

one cycle to be $\sum_{i=1}^{n} AP(t_i)[x(t_i) - x(t_{i-1})]$. If we allow $n \to \infty$, we have

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} AP(t_i)[x(t_i) - x(t_{i-1})] = \lim_{n\to\infty} \sum_{i=1}^{n} P(t_i)[Ax(t_i) - Ax(t_{i-1})]$$

$$= \lim_{n\to\infty} \sum_{i=1}^{n} P(t_i)[V(t_i) - V(t_{i-1})] = \int_C P\,dV$$

(b) Let C_L be the lower loop of the curve C and C_U the upper loop. Then $C = C_L \cup C_U$. C_L is positively oriented, so from Formula 16.4.5 we know the area of the lower loop in the PV-plane is given by $-\oint_{C_L} P\,dV$. C_U is negatively oriented, so the area of the upper loop is given by $-\left(-\oint_{C_U} P\,dV\right) = \oint_{C_U} P\,dV$. From part (a),

$$W = \int_C P\,dV = \int_{C_L \cup C_U} P\,dV = \oint_{C_L} P\,dV + \oint_{C_U} P\,dV$$

$$= \oint_{C_U} P\,dV - \left(-\oint_{C_L} P\,dV\right),$$

the difference of the areas enclosed by the two loops of C.

7. (a) For each value of $u = u_0$, $\mathbf{X}(u_0, v) = \mathbf{r}(u_0) + q\cos v\,\mathbf{N}(u_0) + q\sin v\,\mathbf{B}(u_0)$ is a circle of radius q that is perpendicular to the tangent vector, $\mathbf{r}'(u_0)$. Thus, the union of all circles, $a \le u \le b$ gives a Tube(C, q) around the curve C.

(b) $\mathbf{X}(u, v) = \mathbf{r}(u) + q\cos v\,\mathbf{N}(u) + q\sin v\,\mathbf{B}(u)$, $a \le u \le b$, $0 \le v \le 2\pi$.

First, we find $\mathbf{X}_u(u, v)$ and $\mathbf{X}_v(u, v)$. Note that, as $\mathbf{r}(u)$ is parametrized with respect to arc length, $|\mathbf{r}'(u)| = 1$ and thus, $\mathbf{r}'(u) = \mathbf{T}(u)$. With the Frenet-Serret Formulas, this gives

$$\begin{aligned}
\mathbf{X}_u(u, v) &= \mathbf{r}'(u) + q\cos v\,\mathbf{N}'(u) + q\sin v\,\mathbf{B}'(u) \\
&= \mathbf{T}(u) + q\cos v\,\left(-\kappa\mathbf{T}(u) + \tau\mathbf{B}(u)\right) + q\sin v\,\left(-\tau\mathbf{N}(u)\right) \\
&= \mathbf{T}(u) - q\kappa\cos v\,\mathbf{T}(u) + q\tau\cos v\,\mathbf{B}(u) - q\tau\sin v\,\mathbf{N}(u) \\
&= (1 - q\kappa\cos v)\mathbf{T}(u) + q\tau\cos v\,\mathbf{B}(u) - q\tau\sin v\,\mathbf{N}(u)
\end{aligned}$$

$$\begin{aligned}
\mathbf{X}_v(u, v) &= \mathbf{0} - q\sin v\,\mathbf{N}(u) + q\cos v\,\mathbf{B}(u) \\
&= -q\sin v\,\mathbf{N}(u) + q\cos v\,\mathbf{B}(u)
\end{aligned}$$

Then

$$\begin{aligned}
\mathbf{X}_u(u, v) \times \mathbf{X}_v(u, v) &= [(1 - q\kappa\cos v)\mathbf{T}(u) + q\tau\cos v\,\mathbf{B}(u) - q\tau\sin v\,\mathbf{N}(u)] \\
&\qquad \times\ [-q\sin v\,\mathbf{N}(u) + q\cos v\,\mathbf{B}(u)] \\
&= (1 - q\kappa\cos v)(-q\sin v)\mathbf{T}(u) \times \mathbf{N}(u) + (1 - q\kappa\cos v)(q\cos v)\mathbf{T}(u) \times \mathbf{B}(u) \\
&\qquad - q^2\tau\cos v\sin v\,\mathbf{B}(u) \times \mathbf{N}(u) + q^2\tau\cos^2 v\,\mathbf{B}(u) \times \mathbf{B}(u) \\
&\qquad + q^2\tau\sin^2 v\,\mathbf{N}(u) \times \mathbf{N}(u) - q^2\tau\sin v\cos v\,\mathbf{N}(u) \times \mathbf{B}(u)
\end{aligned}$$

The last four terms drop out since $\mathbf{N}(u) \times \mathbf{N}(u) = \mathbf{B}(u) \times \mathbf{B}(u) = \mathbf{0}$ and $\mathbf{N}(u) \times \mathbf{B}(u) = -(\mathbf{B}(u) \times \mathbf{N}(u))$, so

$$\mathbf{X}_u(u, v) \times \mathbf{X}_v(u, v) = q(1 - q\kappa\cos v)[-\sin v\,\mathbf{B}(u) - \cos v\,\mathbf{N}(u)] \qquad [\mathbf{B} = \mathbf{T} \times \mathbf{N}, \mathbf{N} = -\mathbf{T} \times \mathbf{B}]$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Next,

$$\begin{aligned}|\mathbf{X}_u(u,v)\times\mathbf{X}_v(u,v)|^2 &= |q(1-q\kappa\cos v)[-\sin v\,\mathbf{B}(u)-\cos v\,\mathbf{N}(u)]|^2\\ &= q^2(1-q\kappa\cos v)^2[(-\sin v)^2\,|\mathbf{B}(u)|^2+(-\cos v)^2\,|\mathbf{N}(u)|^2]\\ &\qquad\qquad\text{[by the Pythagorean Theorem for vectors]}\\ &= q^2(1-q\kappa\cos v)^2\left[\sin^2 v\,(1)+\cos^2 v\,(1)\right]=q^2(1-q\kappa\cos v)^2\end{aligned}$$

Thus, $|\mathbf{X}_u(u,v)\times\mathbf{X}_v(u,v)|=q(1-q\kappa\cos v)$. Therefore, the surface area of $\mathrm{Tube}(C,q)$ is

$$\begin{aligned}S(q)&=\int_a^b\int_0^{2\pi}|\mathbf{X}_u(u,v)\times\mathbf{X}_v(u,v)|\,dv\,du=\int_a^b\int_0^{2\pi}q(1-q\kappa\cos v)\,dv\,du\\ &=q\int_a^b\left[v-q\kappa\sin v\right]_{v=0}^{v=2\pi}\,du=q\int_a^b 2\pi\,du=2\pi q(b-a)=2\pi qL\quad[\text{as } b-a=L]\end{aligned}$$

(c) Volume V of $\mathrm{Tube}(C,r)=\int_0^r S(q)\,dq=\int_0^r 2\pi qL\,dq=\left[\pi q^2L\right]_0^r=\pi r^2L$

(d) First find the length L of the helix: $\mathbf{r}(t)=\langle\cos t,\sin t,t\rangle\ \Rightarrow\ \mathbf{r}'(t)=\langle-\sin t,\cos t,1\rangle\ \Rightarrow\ |\mathbf{r}'(t)|=\sqrt{2}\ \Rightarrow$ $L=\int_a^b|\mathbf{r}'(t)|\,dt=\int_0^{4\pi}\sqrt{2}\,dt=4\sqrt{2}\pi$. Then by part (c), with $r=0.2$ and $L=4\sqrt{2}\pi$, we have $V=\pi r^2L=\pi(0.2)^2\left(4\sqrt{2}\pi\right)=\frac{4}{25}\sqrt{2}\pi^2$.

(e) We create the torus by forming a tube of radius r around the circle of radius R. The length of the curve is $L=2\pi R$ and by part (c), $V=\pi r^2L=\pi r^2(2\pi R)=2\pi^2r^2R$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

APPENDIXES

A Numbers, Inequalities, and Absolute Values

1. $|5-23| = |-18| = 18$

2. $|5| - |-23| = 5 - 23 = -18$

3. $|-\pi| = \pi$ because $\pi > 0$.

4. $|\pi - 2| = \pi - 2$ because $\pi - 2 > 0$.

5. $\left|\sqrt{5} - 5\right| = -\left(\sqrt{5} - 5\right) = 5 - \sqrt{5}$ because $\sqrt{5} - 5 < 0$.

6. $\Big|\,|-2| - |-3|\,\Big| = |2-3| = |-1| = 1$

7. If $x < 2$, $x - 2 < 0$, so $|x-2| = -(x-2) = 2 - x$.

8. If $x > 2$, $x - 2 > 0$, so $|x-2| = x - 2$.

9. $|x+1| = \begin{cases} x+1 & \text{if } x+1 \geq 0 \\ -(x+1) & \text{if } x+1 < 0 \end{cases} = \begin{cases} x+1 & \text{if } x \geq -1 \\ -x-1 & \text{if } x < -1 \end{cases}$

10. $|2x-1| = \begin{cases} 2x-1 & \text{if } 2x-1 \geq 0 \\ -(2x-1) & \text{if } 2x-1 < 0 \end{cases} = \begin{cases} 2x-1 & \text{if } x \geq \frac{1}{2} \\ 1-2x & \text{if } x < \frac{1}{2} \end{cases}$

11. $\left|x^2+1\right| = x^2+1$ [since $x^2+1 \geq 0$ for all x].

12. Determine when $1 - 2x^2 < 0 \;\Leftrightarrow\; 1 < 2x^2 \;\Leftrightarrow\; x^2 > \frac{1}{2} \;\Leftrightarrow\; \sqrt{x^2} > \sqrt{\frac{1}{2}} \;\Leftrightarrow\; |x| > \sqrt{\frac{1}{2}} \;\Leftrightarrow$

$x < -\frac{1}{\sqrt{2}}$ or $x > \frac{1}{\sqrt{2}}$. Thus, $\left|1-2x^2\right| = \begin{cases} 1-2x^2 & \text{if } -\frac{1}{\sqrt{2}} \leq x \leq \frac{1}{\sqrt{2}} \\ 2x^2-1 & \text{if } x < -\frac{1}{\sqrt{2}} \text{ or } x > \frac{1}{\sqrt{2}} \end{cases}$

13. $2x + 7 > 3 \;\Leftrightarrow\; 2x > -4 \;\Leftrightarrow\; x > -2$, so $x \in (-2, \infty)$.

14. $3x - 11 < 4 \;\Leftrightarrow\; 3x < 15 \;\Leftrightarrow\; x < 5$, so $x \in (-\infty, 5)$.

15. $1 - x \leq 2 \;\Leftrightarrow\; -x \leq 1 \;\Leftrightarrow\; x \geq -1$, so $x \in [-1, \infty)$.

16. $4 - 3x \geq 6 \;\Leftrightarrow\; -3x \geq 2 \;\Leftrightarrow\; x \leq -\frac{2}{3}$, so $x \in \left(-\infty, -\frac{2}{3}\right]$.

17. $2x + 1 < 5x - 8 \;\Leftrightarrow\; 9 < 3x \;\Leftrightarrow\; 3 < x$, so $x \in (3, \infty)$.

18. $1 + 5x > 5 - 3x \;\Leftrightarrow\; 8x > 4 \;\Leftrightarrow\; x > \frac{1}{2}$, so $x \in \left(\frac{1}{2}, \infty\right)$.

19. $-1 < 2x - 5 < 7 \;\Leftrightarrow\; 4 < 2x < 12 \;\Leftrightarrow\; 2 < x < 6$, so $x \in (2, 6)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

20. $1 < 3x + 4 \leq 16 \quad \Leftrightarrow \quad -3 < 3x \leq 12 \quad \Leftrightarrow \quad -1 < x \leq 4$, so $x \in (-1, 4]$.

21. $0 \leq 1 - x < 1 \quad \Leftrightarrow \quad -1 \leq -x < 0 \quad \Leftrightarrow \quad 1 \geq x > 0$, so $x \in (0, 1]$.

22. $-5 \leq 3 - 2x \leq 9 \quad \Leftrightarrow \quad -8 \leq -2x \leq 6 \quad \Leftrightarrow \quad 4 \geq x \geq -3$, so $x \in [-3, 4]$.

23. $4x < 2x + 1 \leq 3x + 2$. So $4x < 2x + 1 \quad \Leftrightarrow \quad 2x < 1 \quad \Leftrightarrow \quad x < \frac{1}{2}$, and $2x + 1 \leq 3x + 2 \quad \Leftrightarrow \quad -1 \leq x$. Thus, $x \in \left[-1, \frac{1}{2}\right)$.

24. $2x - 3 < x + 4 < 3x - 2$. So $2x - 3 < x + 4 \quad \Leftrightarrow \quad x < 7$, and $x + 4 < 3x - 2 \quad \Leftrightarrow \quad 6 < 2x \quad \Leftrightarrow \quad 3 < x$, so $x \in (3, 7)$.

25. $(x - 1)(x - 2) > 0$.

Case 1: (both factors are positive, so their product is positive) $x - 1 > 0 \quad \Leftrightarrow \quad x > 1$, and $x - 2 > 0 \quad \Leftrightarrow \quad x > 2$, so $x \in (2, \infty)$.

Case 2: (both factors are negative, so their product is positive) $x - 1 < 0 \quad \Leftrightarrow \quad x < 1$, and $x - 2 < 0 \quad \Leftrightarrow \quad x < 2$, so $x \in (-\infty, 1)$.

Thus, the solution set is $(-\infty, 1) \cup (2, \infty)$.

26. $(2x + 3)(x - 1) \geq 0$.

Case 1: $2x + 3 \geq 0 \quad \Leftrightarrow \quad x \geq -\frac{3}{2}$, and $x - 1 \geq 0 \quad \Leftrightarrow \quad x \geq 1$, so $x \in [1, \infty)$.

Case 2: $2x + 3 \leq 0 \quad \Leftrightarrow \quad x \leq -\frac{3}{2}$, and $x - 1 \leq 0 \quad \Leftrightarrow \quad x \leq 1$, so $x \in \left(-\infty, -\frac{3}{2}\right]$.

Thus, the solution set is $\left(-\infty, -\frac{3}{2}\right] \cup [1, \infty)$.

27. $2x^2 + x \leq 1 \quad \Leftrightarrow \quad 2x^2 + x - 1 \leq 0 \quad \Leftrightarrow \quad (2x - 1)(x + 1) \leq 0$.

Case 1: $2x - 1 \geq 0 \quad \Leftrightarrow \quad x \geq \frac{1}{2}$, and $x + 1 \leq 0 \quad \Leftrightarrow \quad x \leq -1$, which is an impossible combination.

Case 2: $2x - 1 \leq 0 \quad \Leftrightarrow \quad x \leq \frac{1}{2}$, and $x + 1 \geq 0 \quad \Leftrightarrow \quad x \geq -1$, so $x \in \left[-1, \frac{1}{2}\right]$.

Thus, the solution set is $\left[-1, \frac{1}{2}\right]$.

28. $x^2 < 2x + 8 \quad \Leftrightarrow \quad x^2 - 2x - 8 < 0 \quad \Leftrightarrow \quad (x - 4)(x + 2) < 0$.

Case 1: $x > 4$ and $x < -2$, which is impossible.

Case 2: $x < 4$ and $x > -2$.

Thus, the solution set is $(-2, 4)$.

29. $x^2 + x + 1 > 0 \quad \Leftrightarrow \quad x^2 + x + \frac{1}{4} + \frac{3}{4} > 0 \quad \Leftrightarrow \quad \left(x + \frac{1}{2}\right)^2 + \frac{3}{4} > 0$. But since $\left(x + \frac{1}{2}\right)^2 \geq 0$ for every real x, the original inequality will be true for all real x as well.

Thus, the solution set is $(-\infty, \infty)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

30. $x^2 + x > 1 \quad \Leftrightarrow \quad x^2 + x - 1 > 0$. Using the quadratic formula, we obtain

$x^2 + x - 1 = \left(x - \frac{-1-\sqrt{5}}{2}\right)\left(x - \frac{-1+\sqrt{5}}{2}\right) > 0.$

Case 1: $x - \frac{-1-\sqrt{5}}{2} > 0$ and $x - \frac{-1+\sqrt{5}}{2} > 0$, so that $x > \frac{-1+\sqrt{5}}{2}$.

Case 2: $x - \frac{-1-\sqrt{5}}{2} < 0$ and $x - \frac{-1+\sqrt{5}}{2} < 0$, so that $x < \frac{-1-\sqrt{5}}{2}$.

Thus, the solution set is $\left(-\infty, \frac{-1-\sqrt{5}}{2}\right) \cup \left(\frac{-1+\sqrt{5}}{2}, \infty\right)$.

31. $x^2 < 3 \quad \Leftrightarrow \quad x^2 - 3 < 0 \quad \Leftrightarrow \quad \left(x - \sqrt{3}\right)\left(x + \sqrt{3}\right) < 0.$

Case 1: $x > \sqrt{3}$ and $x < -\sqrt{3}$, which is impossible.

Case 2: $x < \sqrt{3}$ and $x > -\sqrt{3}$.

Thus, the solution set is $\left(-\sqrt{3}, \sqrt{3}\right)$.

Another method: $x^2 < 3 \quad \Leftrightarrow \quad |x| < \sqrt{3} \quad \Leftrightarrow \quad -\sqrt{3} < x < \sqrt{3}$.

32. $x^2 \geq 5 \quad \Leftrightarrow \quad x^2 - 5 \geq 0 \quad \Leftrightarrow \quad \left(x - \sqrt{5}\right)\left(x + \sqrt{5}\right) \geq 0.$

Case 1: $x \geq \sqrt{5}$ and $x \geq -\sqrt{5}$, so $x \in \left[\sqrt{5}, \infty\right)$.

Case 2: $x \leq \sqrt{5}$ and $x \leq -\sqrt{5}$, so $x \in \left(-\infty, -\sqrt{5}\right]$.

Thus, the solution set is $\left(-\infty, -\sqrt{5}\right] \cup \left[\sqrt{5}, \infty\right)$.

Another method: $x^2 \geq 5 \quad \Leftrightarrow \quad |x| \geq \sqrt{5} \quad \Leftrightarrow \quad x \geq \sqrt{5}$ or $x \leq -\sqrt{5}$.

33. $x^3 - x^2 \leq 0 \quad \Leftrightarrow \quad x^2(x-1) \leq 0$. Since $x^2 \geq 0$ for all x, the inequality is satisfied when $x - 1 \leq 0 \quad \Leftrightarrow \quad x \leq 1$.

Thus, the solution set is $(-\infty, 1]$.

34. $(x+1)(x-2)(x+3) = 0 \quad \Leftrightarrow \quad x = -1, 2,$ or -3. Construct a chart:

Interval	$x+1$	$x-2$	$x+3$	$(x+1)(x-2)(x+3)$
$x < -3$	$-$	$-$	$-$	$-$
$-3 < x < -1$	$-$	$-$	$+$	$+$
$-1 < x < 2$	$+$	$-$	$+$	$-$
$x > 2$	$+$	$+$	$+$	$+$

Thus, $(x+1)(x-2)(x+3) \geq 0$ on $[-3, -1]$ and $[2, \infty)$, and the solution set is $[-3, -1] \cup [2, \infty)$.

35. $x^3 > x \quad \Leftrightarrow \quad x^3 - x > 0 \quad \Leftrightarrow \quad x\left(x^2 - 1\right) > 0 \quad \Leftrightarrow \quad x(x-1)(x+1) > 0$. Construct a chart:

Interval	x	$x-1$	$x+1$	$x(x-1)(x+1)$
$x < -1$	$-$	$-$	$-$	$-$
$-1 < x < 0$	$-$	$-$	$+$	$+$
$0 < x < 1$	$+$	$-$	$+$	$-$
$x > 1$	$+$	$+$	$+$	$+$

Since $x^3 > x$ when the last column is positive, the solution set is $(-1, 0) \cup (1, \infty)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

36. $x^3 + 3x < 4x^2 \quad \Leftrightarrow \quad x^3 - 4x^2 + 3x < 0 \quad \Leftrightarrow \quad x\left(x^2 - 4x + 3\right) < 0 \quad \Leftrightarrow \quad x(x-1)(x-3) < 0.$

Interval	x	$x-1$	$x-3$	$x(x-1)(x-3)$
$x < 0$	$-$	$-$	$-$	$-$
$0 < x < 1$	$+$	$-$	$-$	$+$
$1 < x < 3$	$+$	$+$	$-$	$-$
$x > 3$	$+$	$+$	$+$	$+$

Thus, the solution set is $(-\infty, 0) \cup (1, 3)$.

37. $1/x < 4$. This is clearly true for $x < 0$. So suppose $x > 0$. then $1/x < 4 \quad \Leftrightarrow$ $1 < 4x \quad \Leftrightarrow \quad \frac{1}{4} < x$. Thus, the solution set is $(-\infty, 0) \cup \left(\frac{1}{4}, \infty\right)$.

38. $-3 < 1/x \le 1$. We solve the two inequalities separately and take the intersection of the solution sets. First, $-3 < 1/x$ is clearly true for $x > 0$. So suppose $x < 0$. Then $-3 < 1/x \quad \Leftrightarrow \quad -3x > 1 \quad \Leftrightarrow \quad x < -\frac{1}{3}$, so for this inequality, the solution set is $\left(-\infty, -\frac{1}{3}\right) \cup (0, \infty)$. Now $1/x \le 1$ is clearly true if $x < 0$. So suppose $x > 0$. Then $1/x \le 1 \quad \Leftrightarrow \quad 1 \le x$, and the solution set here is $(-\infty, 0) \cup [1, \infty)$.

Taking the intersection of the two solution sets gives the final solution set: $\left(-\infty, -\frac{1}{3}\right) \cup [1, \infty)$.

39. $C = \frac{5}{9}(F - 32) \quad \Rightarrow \quad F = \frac{9}{5}C + 32$. So $50 \le F \le 95 \quad \Rightarrow \quad 50 \le \frac{9}{5}C + 32 \le 95 \quad \Rightarrow \quad 18 \le \frac{9}{5}C \le 63 \quad \Rightarrow$ $10 \le C \le 35$. So the interval is $[10, 35]$.

40. Since $20 \le C \le 30$ and $C = \frac{5}{9}(F - 32)$, we have $20 \le \frac{5}{9}(F - 32) \le 30 \quad \Rightarrow \quad 36 \le F - 32 \le 54 \quad \Rightarrow \quad 68 \le F \le 86$. So the interval is $[68, 86]$.

41. (a) Let T represent the temperature in degrees Celsius and h the height in km. $T = 20$ when $h = 0$ and T decreases by $10°\text{C}$ for every km ($1°\text{C}$ for each 100-m rise). Thus, $T = 20 - 10h$ when $0 \le h \le 12$.

(b) From part (a), $T = 20 - 10h \quad \Rightarrow \quad 10h = 20 - T \quad \Rightarrow \quad h = 2 - T/10$. So $0 \le h \le 5 \quad \Rightarrow \quad 0 \le 2 - T/10 \le 5 \quad \Rightarrow$ $-2 \le -T/10 \le 3 \quad \Rightarrow \quad -20 \le -T \le 30 \quad \Rightarrow \quad 20 \ge T \ge -30 \quad \Rightarrow \quad -30 \le T \le 20$. Thus, the range of temperatures (in $°\text{C}$) to be expected is $[-30, 20]$.

42. The ball will be at least 32 ft above the ground if $h \ge 32 \quad \Leftrightarrow \quad 128 + 16t - 16t^2 \ge 32 \quad \Leftrightarrow \quad 16t^2 - 16t - 96 \le 0 \quad \Leftrightarrow$ $16(t-3)(t+2) \le 0$. $t = 3$ and $t = -2$ are endpoints of the interval we're looking for, and constructing a table gives $-2 \le t \le 3$. But $t \ge 0$, so the ball will be at least 32 ft above the ground in the time interval $[0, 3]$.

43. $|2x| = 3 \quad \Leftrightarrow \quad$ either $2x = 3$ or $2x = -3 \quad \Leftrightarrow \quad x = \frac{3}{2}$ or $x = -\frac{3}{2}$.

44. $|3x + 5| = 1 \quad \Leftrightarrow \quad$ either $3x + 5 = 1$ or -1. In the first case, $3x = -4 \quad \Leftrightarrow \quad x = -\frac{4}{3}$, and in the second case, $3x = -6 \quad \Leftrightarrow \quad x = -2$. So the solutions are -2 and $-\frac{4}{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45. $|x+3| = |2x+1| \quad \Leftrightarrow \quad$ either $x+3 = 2x+1$ or $x+3 = -(2x+1)$. In the first case, $x = 2$, and in the second case, $x+3 = -2x-1 \quad \Leftrightarrow \quad 3x = -4 \quad \Leftrightarrow \quad x = -\frac{4}{3}$. So the solutions are $-\frac{4}{3}$ and 2.

46. $\left|\dfrac{2x-1}{x+1}\right| = 3 \quad \Leftrightarrow \quad$ either $\dfrac{2x-1}{x+1} = 3$ or $\dfrac{2x-1}{x+1} = -3$. In the first case, $2x-1 = 3x+3 \quad \Leftrightarrow \quad x = -4$, and in the second case, $2x-1 = -3x-3 \quad \Leftrightarrow \quad x = -\frac{2}{5}$.

47. By Property 5 of absolute values, $|x| < 3 \quad \Leftrightarrow \quad -3 < x < 3$, so $x \in (-3, 3)$.

48. By Properties 4 and 6 of absolute values, $|x| \geq 3 \quad \Leftrightarrow \quad x \leq -3$ or $x \geq 3$, so $x \in (-\infty, -3] \cup [3, \infty)$.

49. $|x-4| < 1 \quad \Leftrightarrow \quad -1 < x-4 < 1 \quad \Leftrightarrow \quad 3 < x < 5$, so $x \in (3, 5)$.

50. $|x-6| < 0.1 \quad \Leftrightarrow \quad -0.1 < x-6 < 0.1 \quad \Leftrightarrow \quad 5.9 < x < 6.1$, so $x \in (5.9, 6.1)$.

51. $|x+5| \geq 2 \quad \Leftrightarrow \quad x+5 \geq 2$ or $x+5 \leq -2 \quad \Leftrightarrow \quad x \geq -3$ or $x \leq -7$, so $x \in (-\infty, -7] \cup [-3, \infty)$.

52. $|x+1| \geq 3 \quad \Leftrightarrow \quad x+1 \geq 3$ or $x+1 \leq -3 \quad \Leftrightarrow \quad x \geq 2$ or $x \leq -4$, so $x \in (-\infty, -4] \cup [2, \infty)$.

53. $|2x-3| \leq 0.4 \quad \Leftrightarrow \quad -0.4 \leq 2x-3 \leq 0.4 \quad \Leftrightarrow \quad 2.6 \leq 2x \leq 3.4 \quad \Leftrightarrow \quad 1.3 \leq x \leq 1.7$, so $x \in [1.3, 1.7]$.

54. $|5x-2| < 6 \quad \Leftrightarrow \quad -6 < 5x-2 < 6 \quad \Leftrightarrow \quad -4 < 5x < 8 \quad \Leftrightarrow \quad -\frac{4}{5} < x < \frac{8}{5}$, so $x \in \left(-\frac{4}{5}, \frac{8}{5}\right)$.

55. $1 \leq |x| \leq 4$. So either $1 \leq x \leq 4$ or $1 \leq -x \leq 4 \quad \Leftrightarrow \quad -1 \geq x \geq -4$. Thus, $x \in [-4, -1] \cup [1, 4]$.

56. $0 < |x-5| < \frac{1}{2}$. Clearly $0 < |x-5|$ for $x \neq 5$. Now $|x-5| < \frac{1}{2} \quad \Leftrightarrow \quad -\frac{1}{2} < x-5 < \frac{1}{2} \quad \Leftrightarrow \quad 4.5 < x < 5.5$. So the solution set is $(4.5, 5) \cup (5, 5.5)$.

57. $a(bx-c) \geq bc \quad \Leftrightarrow \quad bx - c \geq \dfrac{bc}{a} \quad \Leftrightarrow \quad bx \geq \dfrac{bc}{a} + c = \dfrac{bc+ac}{a} \quad \Leftrightarrow \quad x \geq \dfrac{bc+ac}{ab}$

58. $a \leq bx + c < 2a \quad \Leftrightarrow \quad a - c \leq bx < 2a - c \quad \Leftrightarrow \quad \dfrac{a-c}{b} \leq x < \dfrac{2a-c}{b}$ (since $b > 0$)

59. $ax + b < c \quad \Leftrightarrow \quad ax < c - b \quad \Leftrightarrow \quad x > \dfrac{c-b}{a}$ [since $a < 0$]

60. $\dfrac{ax+b}{c} \leq b \quad \Leftrightarrow \quad ax + b \geq bc$ [since $c < 0$] $\quad \Leftrightarrow \quad ax \geq bc - b \quad \Leftrightarrow \quad x \leq \dfrac{b(c-1)}{a}$ [since $a < 0$]

61. $|(x+y) - 5| = |(x-2) + (y-3)| \leq |x-2| + |y-3| < 0.01 + 0.04 = 0.05$

62. Use the Triangle Inequality: $|x+3| < \frac{1}{2} \quad \Rightarrow$

$|4x+13| = |4(x+3)+1| \leq |4(x+3)| + |1| = 4|x+3| + 1 < 4\left(\frac{1}{2}\right) + 1 = 3$

Another method: $|x+3| < \frac{1}{2} \quad \Rightarrow \quad -\frac{1}{2} < x+3 < \frac{1}{2} \quad \Rightarrow \quad -2 < 4x+12 < 2 \quad \Rightarrow \quad -1 < 4x+13 < 3 \quad \Rightarrow$ $|4x+13| < 3$

63. If $a < b$ then $a + a < a + b$ and $a + b < b + b$. So $2a < a + b < 2b$. Dividing by 2, we get $a < \frac{1}{2}(a+b) < b$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. If $0 < a < b$, then $\dfrac{1}{ab} > 0$. So $a < b \quad \Rightarrow \quad \dfrac{1}{ab} \cdot a < \dfrac{1}{ab} \cdot b \quad \Leftrightarrow \quad \dfrac{1}{b} < \dfrac{1}{a}$.

65. $|ab| = \sqrt{(ab)^2} = \sqrt{a^2b^2} = \sqrt{a^2}\,\sqrt{b^2} = |a|\,|b|$

66. $\left|\dfrac{a}{b}\right| |b| = \left|\dfrac{a}{b} \cdot b\right| = |a|$ [using the result of Exercise 65]. Dividing the equation through by $|b|$ gives $\left|\dfrac{a}{b}\right| = \dfrac{|a|}{|b|}$.

67. If $0 < a < b$, then $a \cdot a < a \cdot b$ and $a \cdot b < b \cdot b$ [using Rule 3 of Inequalities]. So $a^2 < ab < b^2$ and hence $a^2 < b^2$.

68. Following the hint, the Triangle Inequality becomes $|(x - y) + y| \le |x - y| + |y| \quad \Leftrightarrow \quad |x| \le |x - y| + |y| \quad \Leftrightarrow$ $|x - y| \ge |x| - |y|$.

69. Observe that the sum, difference and product of two integers is always an integer. Let the rational numbers be represented by $r = m/n$ and $s = p/q$ (where m, n, p and q are integers with $n \neq 0$, $q \neq 0$). Now $r + s = \dfrac{m}{n} + \dfrac{p}{q} = \dfrac{mq + pn}{nq}$, but $mq + pn$ and nq are both integers, so $\dfrac{mq + pn}{nq} = r + s$ is a rational number by definition. Similarly, $r - s = \dfrac{m}{n} - \dfrac{p}{q} = \dfrac{mq - pn}{nq}$ is a rational number. Finally, $r \cdot s = \dfrac{m}{n} \cdot \dfrac{p}{q} = \dfrac{mp}{nq}$ but mp and nq are both integers, so $\dfrac{mp}{nq} = r \cdot s$ is a rational number by definition.

70. (a) No. Consider the case of $\sqrt{2}$ and $-\sqrt{2}$. Both are irrational numbers, yet $\sqrt{2} + \left(-\sqrt{2}\right) = 0$ and 0, being an integer, is not irrational.

(b) No. Consider the case of $\sqrt{2}$ and $\sqrt{2}$. Both are irrational numbers, yet $\sqrt{2} \cdot \sqrt{2} = 2$ is not irrational.

B Coordinate Geometry and Lines

1. Use the distance formula with $P_1(x_1, y_1) = (1, 1)$ and $P_2(x_2, y_2) = (4, 5)$ to get

$$|P_1P_2| = \sqrt{(4-1)^2 + (5-1)^2} = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$$

2. The distance from $(1, -3)$ to $(5, 7)$ is $\sqrt{(5-1)^2 + [7-(-3)]^2} = \sqrt{4^2 + 10^2} = \sqrt{116} = 2\sqrt{29}$.

3. The distance from $(6, -2)$ to $(-1, 3)$ is $\sqrt{-1-6)^2 + [3-(-2)]^2} = \sqrt{(-7)^2 + 5^2} = \sqrt{74}$.

4. The distance from $(1, -6)$ to $(-1, -3)$ is $\sqrt{(-1-1)^2 + [-3-(-6)]^2} = \sqrt{(-2)^2 + 3^2} = \sqrt{13}$.

5. The distance from $(2, 5)$ to $(4, -7)$ is $\sqrt{(4-2)^2 + (-7-5)^2} = \sqrt{2^2 + (-12)^2} = \sqrt{148} = 2\sqrt{37}$.

6. The distance from (a, b) to (b, a) is $\sqrt{(b-a)^2 + (a-b)^2} = \sqrt{(a-b)^2 + (a-b)^2} = \sqrt{2(a-b)^2} = \sqrt{2}\,|a-b|$.

7. The slope m of the line through $P(1, 5)$ and $Q(4, 11)$ is $m = \dfrac{11-5}{4-1} = \dfrac{6}{3} = 2$.

8. The slope m of the line through $P(-1, 6)$ and $Q(4, -3)$ is $m = \dfrac{-3-6}{4-(-1)} = -\dfrac{9}{5}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. The slope m of the line through $P(-3,3)$ and $Q(-1,-6)$ is $m = \dfrac{-6-3}{-1-(-3)} = -\dfrac{9}{2}$.

10. The slope m of the line through $P(-1,-4)$ and $Q(6,0)$ is $m = \dfrac{0-(-4)}{6-(-1)} = \dfrac{4}{7}$.

11. Using $A(0,2)$, $B(-3,-1)$, and $C(-4,3)$, we have $|AC| = \sqrt{(-4-0)^2+(3-2)^2} = \sqrt{(-4)^2+1^2} = \sqrt{17}$ and $|BC| = \sqrt{[-4-(-3)]^2+[3-(-1)]^2} = \sqrt{(-1)^2+4^2} = \sqrt{17}$, so the triangle has two sides of equal length, and is isosceles.

12. (a) Using $A(6,-7)$, $B(11,-3)$, and $C(2,-2)$, we have

$|AB| = \sqrt{(11-6)^2+[-3-(-7)]^2} = \sqrt{5^2+4^2} = \sqrt{41}$,

$|AC| = \sqrt{(2-6)^2+[-2-(-7)]^2} = \sqrt{(-4)^2+5^2} = \sqrt{41}$, and

$|BC| = \sqrt{(2-11)^2+[-2-(-3)]^2} = \sqrt{(-9)^2+1^2} = \sqrt{82}$.

Thus, $|AB|^2 + |AC|^2 = 41 + 41 = 82 = |BC|^2$ and so $\triangle ABC$ is a right triangle.

(b) $m_{AB} = \dfrac{-3-(-7)}{11-6} = \dfrac{4}{5}$ and $m_{AC} = \dfrac{-2-(-7)}{2-6} = -\dfrac{5}{4}$. Thus $m_{AB} \cdot m_{AC} = -1$ and so AB is perpendicular to AC and $\triangle ABC$ must be a right triangle.

(c) Taking lengths from part (a), the base is $\sqrt{41}$ and the height is $\sqrt{41}$. Thus the area is $\frac{1}{2}bh = \frac{1}{2}\sqrt{41}\sqrt{41} = \frac{41}{2}$.

13. Using $A(-2,9)$, $B(4,6)$, $C(1,0)$, and $D(-5,3)$, we have

$|AB| = \sqrt{[4-(-2)]^2+(6-9)^2} = \sqrt{6^2+(-3)^2} = \sqrt{45} = \sqrt{9}\sqrt{5} = 3\sqrt{5}$,

$|BC| = \sqrt{(1-4)^2+(0-6)^2} = \sqrt{(-3)^2+(-6)^2} = \sqrt{45} = \sqrt{9}\sqrt{5} = 3\sqrt{5}$,

$|CD| = \sqrt{(-5-1)^2+(3-0)^2} = \sqrt{(-6)^2+3^2} = \sqrt{45} = \sqrt{9}\sqrt{5} = 3\sqrt{5}$, and

$|DA| = \sqrt{[-2-(-5)]^2+(9-3)^2} = \sqrt{3^2+6^2} = \sqrt{45} = \sqrt{9}\sqrt{5} = 3\sqrt{5}$. So all sides are of equal length and we have a rhombus. Moreover, $m_{AB} = \dfrac{6-9}{4-(-2)} = -\dfrac{1}{2}$, $m_{BC} = \dfrac{0-6}{1-4} = 2$, $m_{CD} = \dfrac{3-0}{-5-1} = -\dfrac{1}{2}$, and $m_{DA} = \dfrac{9-3}{-2-(-5)} = 2$, so the sides are perpendicular. Thus, A, B, C, and D are vertices of a square.

14. (a) Using $A(-1,3)$, $B(3,11)$, and $C(5,15)$, we have

$|AB| = \sqrt{[3-(-1)]^2+(11-3)^2} = \sqrt{4^2+8^2} = \sqrt{80} = 4\sqrt{5}$,

$|BC| = \sqrt{(5-3)^2+(15-11)^2} = \sqrt{2^2+4^2} = \sqrt{20} = 2\sqrt{5}$, and

$|AC| = \sqrt{[5-(-1)]^2+(15-3)^2} = \sqrt{6^2+12^2} = \sqrt{180} = 6\sqrt{5}$. Thus, $|AC| = |AB| + |BC|$.

(b) $m_{AB} = \dfrac{11-3}{3-(-1)} = \dfrac{8}{4} = 2$ and $m_{AC} = \dfrac{15-3}{5-(-1)} = \dfrac{12}{6} = 2$. Since the segments AB and AC have the same slope, A, B and C must be collinear.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

15. For the vertices $A(1,1)$, $B(7,4)$, $C(5,10)$, and $D(-1,7)$, the slope of the line segment AB is $\dfrac{4-1}{7-1}=\dfrac{1}{2}$, the slope of CD is $\dfrac{7-10}{-1-5}=\dfrac{1}{2}$, the slope of BC is $\dfrac{10-4}{5-7}=-3$, and the slope of DA is $\dfrac{1-7}{1-(-1)}=-3$. So AB is parallel to CD and BC is parallel to DA. Hence $ABCD$ is a parallelogram.

16. For the vertices $A(1,1)$, $B(11,3)$, $C(10,8)$, and $D(0,6)$, the slopes of the four sides are $m_{AB}=\dfrac{3-1}{11-1}=\dfrac{1}{5}$, $m_{BC}=\dfrac{8-3}{10-11}=-5$, $m_{CD}=\dfrac{6-8}{0-10}=\dfrac{1}{5}$, and $m_{DA}=\dfrac{1-6}{1-0}=-5$. Hence $AB \parallel CD$, $BC \parallel DA$, $AB \perp BC$, $BC \perp CD$, $CD \perp DA$, and $DA \perp AB$, and so $ABCD$ is a rectangle.

17. The graph of the equation $x=3$ is a vertical line with x-intercept 3. The line does not have a slope.

18. The graph of the equation $y=-2$ is a horizontal line with y-intercept -2. The line has slope 0.

19. $xy=0 \quad\Leftrightarrow\quad x=0$ or $y=0$. The graph consists of the coordinate axes.

20. $|y|=1 \quad\Leftrightarrow\quad y=1$ or $y=-1$

21. By the point-slope form of the equation of a line, an equation of the line through $(2,-3)$ with slope 6 is $y-(-3)=6(x-2)$ or $y=6x-15$.

22. $y-4=-3[x-(-1)]$ or $y=-3x+1$

23. $y-7=\frac{2}{3}(x-1)$ or $y=\frac{2}{3}x+\frac{19}{3}$

24. $y-(-5)=-\frac{7}{2}[x-(-3)]$ or $y=-\frac{7}{2}x-\frac{31}{2}$

25. The slope of the line through $(2,1)$ and $(1,6)$ is $m=\dfrac{6-1}{1-2}=-5$, so an equation of the line is $y-1=-5(x-2)$ or $y=-5x+11$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. For $(-1,-2)$ and $(4,3)$, $m = \dfrac{3-(-2)}{4-(-1)} = 1$. An equation of the line is $y-3 = 1(x-4)$ or $y = x-1$.

27. By the slope-intercept form of the equation of a line, an equation of the line is $y = 3x-2$.

28. By the slope-intercept form of the equation of a line, an equation of the line is $y = \frac{2}{5}x+4$.

29. Since the line passes through $(1,0)$ and $(0,-3)$, its slope is $m = \dfrac{-3-0}{0-1} = 3$, so an equation is $y = 3x-3$.

Another method: From Exercise 61, $\dfrac{x}{1} + \dfrac{y}{-3} = 1 \quad\Rightarrow\quad -3x+y=-3 \quad\Rightarrow\quad y = 3x-3$.

30. For $(-8,0)$ and $(0,6)$, $m = \dfrac{6-0}{0-(-8)} = \dfrac{3}{4}$. So an equation is $y = \frac{3}{4}x+6$.

Another method: From Exercise 61, $\dfrac{x}{-8} + \dfrac{y}{6} = 1 \quad\Rightarrow\quad -3x+4y=24 \quad\Rightarrow\quad y = \frac{3}{4}x+6$.

31. The line is parallel to the x-axis, so it is horizontal and must have the form $y = k$. Since it goes through the point $(x,y) = (4,5)$, the equation is $y = 5$.

32. The line is parallel to the y-axis, so it is vertical and must have the form $x = k$. Since it goes through the point $(x,y) = (4,5)$, the equation is $x = 4$.

33. Putting the line $x+2y=6$ into its slope-intercept form gives us $y = -\frac{1}{2}x+3$, so we see that this line has slope $-\frac{1}{2}$. Thus, we want the line of slope $-\frac{1}{2}$ that passes through the point $(1,-6)$: $y-(-6) = -\frac{1}{2}(x-1) \quad\Leftrightarrow\quad y = -\frac{1}{2}x - \frac{11}{2}$.

34. $2x+3y+4=0 \quad\Leftrightarrow\quad y = -\frac{2}{3}x - \frac{4}{3}$, so $m = -\frac{2}{3}$ and the required line is $y = -\frac{2}{3}x+6$.

35. $2x+5y+8=0 \quad\Leftrightarrow\quad y = -\frac{2}{5}x - \frac{8}{5}$. Since this line has slope $-\frac{2}{5}$, a line perpendicular to it would have slope $\frac{5}{2}$, so the required line is $y-(-2) = \frac{5}{2}[x-(-1)] \quad\Leftrightarrow\quad y = \frac{5}{2}x + \frac{1}{2}$.

36. $4x-8y=1 \quad\Leftrightarrow\quad y = \frac{1}{2}x - \frac{1}{8}$. Since this line has slope $\frac{1}{2}$, a line perpendicular to it would have slope -2, so the required line is $y - \left(-\frac{2}{3}\right) = -2\left(x-\frac{1}{2}\right) \quad\Leftrightarrow\quad y = -2x+\frac{1}{3}$.

37. $x+3y=0 \quad\Leftrightarrow\quad y = -\frac{1}{3}x$, so the slope is $-\frac{1}{3}$ and the y-intercept is 0.

38. $2x-5y=0 \quad\Leftrightarrow\quad y = \frac{2}{5}x$, so the slope is $\frac{2}{5}$ and the y-intercept is 0.

39. $y=-2$ is a horizontal line with slope 0 and y-intercept -2.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $2x - 3y + 6 = 0 \quad \Leftrightarrow$
$y = \frac{2}{3}x + 2$, so the slope is $\frac{2}{3}$ and the y-intercept is 2.

41. $3x - 4y = 12 \quad \Leftrightarrow$
$y = \frac{3}{4}x - 3$, so the slope is $\frac{3}{4}$ and the y-intercept is -3.

42. $4x + 5y = 10 \quad \Leftrightarrow$
$y = -\frac{4}{5}x + 2$, so the slope is $-\frac{4}{5}$ and the y-intercept is 2.

43. $\{(x, y) \mid x < 0\}$

44. $\{(x, y) \mid y > 0\}$

45. $\{(x, y) \mid xy < 0\} =$
$\{(x, y) \mid x < 0 \text{ and } y > 0\}$
$\cup \{(x, y) \mid x > 0 \text{ and } y < 0\}$

46. $\{(x, y) \mid x \geq 1 \text{ and } y < 3\}$

47. $\left\{(x, y) \,\middle|\, |x| \leq 2\right\} =$
$\{(x, y) \mid -2 \leq x \leq 2\}$

48. $\left\{(x, y) \,\middle|\, |x| < 3 \text{ and } |y| < 2\right\}$

49. $\{(x, y) \mid 0 \leq y \leq 4, x \leq 2\}$

50. $\{(x, y) \mid y > 2x - 1\}$

51. $\{(x, y) \mid 1 + x \leq y \leq 1 - 2x\}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

52. $\{(x, y) \mid -x \le y < \frac{1}{2}(x+3)\}$

53. Let $P\,(0, y)$ be a point on the y-axis. The distance from P to $(5, -5)$ is $\sqrt{(5-0)^2 + (-5-y)^2} = \sqrt{5^2 + (y+5)^2}$. The distance from P to $(1, 1)$ is $\sqrt{(1-0)^2 + (1-y)^2} = \sqrt{1^2 + (y-1)^2}$. We want these distances to be equal: $\sqrt{5^2 + (y+5)^2} = \sqrt{1^2 + (y-1)^2} \;\Leftrightarrow\; 5^2 + (y+5)^2 = 1^2 + (y-1)^2 \;\Leftrightarrow\; 25 + (y^2 + 10y + 25) = 1 + (y^2 - 2y + 1) \;\Leftrightarrow\; 12y = -48 \;\Leftrightarrow\; y = -4$. So the desired point is $(0, -4)$.

54. Let M be the point $\left(\dfrac{x_1 + x_2}{2}, \dfrac{y_1 + y_2}{2}\right)$. Then

$$|MP_1|^2 = \left(x_1 - \frac{x_1 + x_2}{2}\right)^2 + \left(y_1 - \frac{y_1 + y_2}{2}\right)^2 = \left(\frac{x_1 - x_2}{2}\right)^2 + \left(\frac{y_1 - y_2}{2}\right)^2$$

$$|MP_2|^2 = \left(x_2 - \frac{x_1 + x_2}{2}\right)^2 + \left(y_2 - \frac{y_1 + y_2}{2}\right)^2 = \left(\frac{x_2 - x_1}{2}\right)^2 + \left(\frac{y_2 - y_1}{2}\right)^2$$

Hence, $|MP_1| = |MP_2|$; that is, M is equidistant from P_1 and P_2.

55. (a) Using the midpoint formula from Exercise 54 with $(1, 3)$ and $(7, 15)$, we get $\left(\frac{1+7}{2}, \frac{3+15}{2}\right) = (4, 9)$.

(b) Using the midpoint formula from Exercise 54 with $(-1, 6)$ and $(8, -12)$, we get $\left(\frac{-1+8}{2}, \frac{6+(-12)}{2}\right) = \left(\frac{7}{2}, -3\right)$.

56. With $A(1, 0)$, $B(3, 6)$, and $C(8, 2)$, the midpoint M_1 of AB is $\left(\frac{1+3}{2}, \frac{0+6}{2}\right) = (2, 3)$, the midpoint M_2 of BC is $\left(\frac{3+8}{2}, \frac{6+2}{2}\right) = \left(\frac{11}{2}, 4\right)$, and the midpoint M_3 of CA is $\left(\frac{8+1}{2}, \frac{2+0}{2}\right) = \left(\frac{9}{2}, 1\right)$. The lengths of the medians are

$$|AM_2| = \sqrt{\left(\tfrac{11}{2} - 1\right)^2 + (4-0)^2} = \sqrt{\left(\tfrac{9}{2}\right)^2 + 4^2} = \sqrt{\tfrac{145}{4}} = \tfrac{\sqrt{145}}{2}$$

$$|BM_3| = \sqrt{\left(\tfrac{9}{2} - 3\right)^2 + (1-6)^2} = \sqrt{\left(\tfrac{3}{2}\right)^2 + (-5)^2} = \sqrt{\tfrac{109}{4}} = \tfrac{\sqrt{109}}{2}$$

$$|CM_1| = \sqrt{(2-8)^2 + (3-2)^2} = \sqrt{(-6)^2 + 1^2} = \sqrt{37}$$

57. $2x - y = 4 \;\Leftrightarrow\; y = 2x - 4 \;\Rightarrow\; m_1 = 2$ and $6x - 2y = 10 \;\Leftrightarrow\; 2y = 6x - 10 \;\Leftrightarrow\; y = 3x - 5 \;\Rightarrow\; m_2 = 3$. Since $m_1 \ne m_2$, the two lines are not parallel. To find the point of intersection: $2x - 4 = 3x - 5 \;\Leftrightarrow\; x = 1 \;\Rightarrow\; y = -2$. Thus, the point of intersection is $(1, -2)$.

58. $3x - 5y + 19 = 0 \;\Leftrightarrow\; 5y = 3x + 19 \;\Leftrightarrow\; y = \frac{3}{5}x + \frac{19}{5} \;\Rightarrow\; m_1 = \frac{3}{5}$ and $10x + 6y - 50 = 0 \;\Leftrightarrow\; 6y = -10x + 50 \;\Leftrightarrow\; y = -\frac{5}{3}x + \frac{25}{3} \;\Rightarrow\; m_2 = -\frac{5}{3}$. Since $m_1 m_2 = \frac{3}{5}\left(-\frac{5}{3}\right) = -1$, the two lines are perpendicular. To find the point of intersection: $\frac{3}{5}x + \frac{19}{5} = -\frac{5}{3}x + \frac{25}{3} \;\Leftrightarrow\; 9x + 57 = -25x + 125 \;\Leftrightarrow\; 34x = 68 \;\Leftrightarrow\; x = 2 \;\Rightarrow\; y = \frac{3}{5} \cdot 2 + \frac{19}{5} = \frac{25}{5} = 5$. Thus, the point of intersection is $(2, 5)$.

59. With $A(1, 4)$ and $B(7, -2)$, the slope of segment AB is $\frac{-2-4}{7-1} = -1$, so its perpendicular bisector has slope 1. The midpoint of AB is $\left(\frac{1+7}{2}, \frac{4+(-2)}{2}\right) = (4, 1)$, so an equation of the perpendicular bisector is $y - 1 = 1(x - 4)$ or $y = x - 3$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. (a) Side PQ has slope $\frac{4-0}{3-1}=2$, so its equation is $y-0=2(x-1) \quad \Leftrightarrow \quad y=2x-2$. Side QR has slope $\frac{6-4}{-1-3}=-\frac{1}{2}$, so its equation is $y-4=-\frac{1}{2}(x-3) \quad \Leftrightarrow \quad y=-\frac{1}{2}x+\frac{11}{2}$. Side RP has slope $\frac{0-6}{1-(-1)}=-3$, so its equation is $y-0=-3(x-1) \quad \Leftrightarrow \quad y=-3x+3$.

(b) M_1 (the midpoint of PQ) has coordinates $\left(\frac{1+3}{2}, \frac{0+4}{2}\right)=(2,2)$. M_2 (the midpoint of QR) has coordinates $\left(\frac{3-1}{2}, \frac{4+6}{2}\right)=(1,5)$. M_3 (the midpoint of RP) has coordinates $\left(\frac{1-1}{2}, \frac{0+6}{2}\right)=(0,3)$. RM_1 has slope $\frac{2-6}{2-(-1)}=-\frac{4}{3}$ and hence equation $y-2=-\frac{4}{3}(x-2) \quad \Leftrightarrow \quad y=-\frac{4}{3}x+\frac{14}{3}$. PM_2 is a vertical line with equation $x=1$. QM_3 has slope $\frac{3-4}{0-3}=\frac{1}{3}$ and hence equation $y-3=\frac{1}{3}(x-0) \quad \Leftrightarrow \quad y=\frac{1}{3}x+3$. PM_2 and RM_1 intersect where $x=1$ and $y=-\frac{4}{3}(1)+\frac{14}{3}=\frac{10}{3}$, or at $\left(1, \frac{10}{3}\right)$. PM_2 and QM_3 intersect where $x=1$ and $y=\frac{1}{3}(1)+3=\frac{10}{3}$, or at $\left(1, \frac{10}{3}\right)$, so this is the point where all three medians intersect.

61. (a) Since the x-intercept is a, the point $(a,0)$ is on the line, and similarly since the y-intercept is b, $(0,b)$ is on the line. Hence, the slope of the line is $m=\dfrac{b-0}{0-a}=-\dfrac{b}{a}$. Substituting into $y=mx+b$ gives $y=-\dfrac{b}{a}x+b \quad \Leftrightarrow \quad \dfrac{b}{a}x+y=b \quad \Leftrightarrow \quad \dfrac{x}{a}+\dfrac{y}{b}=1$.

(b) Letting $a=6$ and $b=-8$ gives $\dfrac{x}{6}+\dfrac{y}{-8}=1 \quad \Leftrightarrow \quad -8x+6y=-48$ [multiply by -48] $\quad \Leftrightarrow \quad 6y=8x-48 \quad \Leftrightarrow \quad 3y=4x-24 \quad \Leftrightarrow \quad y=\frac{4}{3}x-8$.

62. (a) Let $d=$ distance traveled (in miles) and $t=$ time elapsed (in hours). At $t=0$, $d=0$ and at $t=50$ minutes $=50\cdot\frac{1}{60}=\frac{5}{6}$ h, $d=40$. Thus, we have two points: $(0,0)$ and $\left(\frac{5}{6},40\right)$, so $m=\dfrac{40-0}{5/6-0}=48$ and $d=48t$.

(b)

(c) The slope is 48 and represents the car's speed in mi/h.

C Graphs of Second-Degree Equations

1. An equation of the circle with center $(3,-1)$ and radius 5 is $(x-3)^2+(y+1)^2=5^2=25$.

2. An equation of the circle with center $(-2,-8)$ and radius 10 is $(x+2)^2+(y+8)^2=10^2=100$.

3. The equation has the form $x^2+y^2=r^2$. Since $(4,7)$ lies on the circle, we have $4^2+7^2=r^2 \quad \Rightarrow \quad r^2=65$. So the required equation is $x^2+y^2=65$.

4. The equation has the form $(x+1)^2+(y-5)^2=r^2$. Since $(-4,-6)$ lies on the circle, we have $r^2=(-4+1)^2+(-6-5)^2=130$. So an equation is $(x+1)^2+(y-5)^2=130$.

5. $x^2+y^2-4x+10y+13=0 \quad \Leftrightarrow \quad x^2-4x+y^2+10y=-13 \quad \Leftrightarrow$ $(x^2-4x+4)+(y^2+10y+25)=-13+4+25=16 \quad \Leftrightarrow \quad (x-2)^2+(y+5)^2=4^2$. Thus, we have a circle with center $(2,-5)$ and radius 4.

6. $x^2+y^2+6y+2=0 \quad \Leftrightarrow \quad x^2+(y^2+6y+9)=-2+9 \quad \Leftrightarrow \quad x^2+(y+3)^2=7$. Thus, we have a circle with center $(0,-3)$ and radius $\sqrt{7}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

7. $x^2+y^2+x=0 \quad\Leftrightarrow\quad \left(x^2+x+\frac{1}{4}\right)+y^2=\frac{1}{4} \quad\Leftrightarrow\quad \left(x+\frac{1}{2}\right)^2+y^2=\left(\frac{1}{2}\right)^2$. Thus, we have a circle with center $\left(-\frac{1}{2},0\right)$ and radius $\frac{1}{2}$.

8. $16x^2+16y^2+8x+32y+1=0 \quad\Leftrightarrow\quad 16\left(x^2+\frac{1}{2}x+\frac{1}{16}\right)+16(y^2+2y+1)=-1+1+16 \quad\Leftrightarrow$ $16\left(x+\frac{1}{4}\right)^2+16(y+1)^2=16 \quad\Leftrightarrow\quad \left(x+\frac{1}{4}\right)^2+(y+1)^2=1$. Thus, we have a circle with center $\left(-\frac{1}{4},-1\right)$ and radius 1.

9. $2x^2+2y^2-x+y=1 \quad\Leftrightarrow\quad 2\left(x^2-\frac{1}{2}x+\frac{1}{16}\right)+2\left(y^2+\frac{1}{2}y+\frac{1}{16}\right)=1+\frac{1}{8}+\frac{1}{8} \quad\Leftrightarrow$ $2\left(x-\frac{1}{4}\right)^2+2\left(y+\frac{1}{4}\right)^2=\frac{5}{4} \quad\Leftrightarrow\quad \left(x-\frac{1}{4}\right)^2+\left(y+\frac{1}{4}\right)^2=\frac{5}{8}$. Thus, we have a circle with center $\left(\frac{1}{4},-\frac{1}{4}\right)$ and radius $\frac{\sqrt{5}}{2\sqrt{2}}=\frac{\sqrt{10}}{4}$.

10. $x^2+y^2+ax+by+c=0 \quad\Leftrightarrow\quad \left(x^2+ax+\frac{1}{4}a^2\right)+\left(y^2+by+\frac{1}{4}b^2\right)=-c+\frac{1}{4}a^2+\frac{1}{4}b^2 \quad\Leftrightarrow$ $\left(x+\frac{1}{2}a\right)^2+\left(y+\frac{1}{2}b\right)^2=\frac{1}{4}(a^2+b^2-4c)$. For this to represent a nondegenerate circle, $\frac{1}{4}(a^2+b^2-4c)>0$ or $a^2+b^2>4c$. If this condition is satisfied, the circle has center $\left(-\frac{1}{2}a,-\frac{1}{2}b\right)$ and radius $\frac{1}{2}\sqrt{a^2+b^2-4c}$.

11. $y=-x^2$. Parabola

12. $y^2-x^2=1$. Hyperbola

13. $x^2+4y^2=16 \quad\Leftrightarrow\quad \dfrac{x^2}{16}+\dfrac{y^2}{4}=1$. Ellipse

14. $x=-2y^2$. Parabola

15. $16x^2-25y^2=400 \quad\Leftrightarrow\quad \dfrac{x^2}{25}-\dfrac{y^2}{16}=1$. Hyperbola

16. $25x^2+4y^2=100 \quad\Leftrightarrow\quad \dfrac{x^2}{4}+\dfrac{y^2}{25}=1$. Ellipse

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $4x^2 + y^2 = 1 \quad \Leftrightarrow \quad \dfrac{x^2}{1/4} + y^2 = 1$. Ellipse

18. $y = x^2 + 2$. Parabola with vertex at $(0, 2)$

19. $x = y^2 - 1$. Parabola with vertex at $(-1, 0)$

20. $9x^2 - 25y^2 = 225 \quad \Leftrightarrow \quad \dfrac{x^2}{25} - \dfrac{y^2}{9} = 1$. Hyperbola

21. $9y^2 - x^2 = 9 \quad \Leftrightarrow \quad y^2 - \dfrac{x^2}{9} = 1$. Hyperbola

22. $2x^2 + 5y^2 = 10 \quad \Leftrightarrow \quad \dfrac{x^2}{5} + \dfrac{y^2}{2} = 1$. Ellipse

23. $xy = 4$. Hyperbola

24. $y = x^2 + 2x = \left(x^2 + 2x + 1\right) - 1 = (x + 1)^2 - 1$.
Parabola with vertex at $(-1, -1)$

25. $9(x - 1)^2 + 4(y - 2)^2 = 36 \quad \Leftrightarrow$
$\dfrac{(x - 1)^2}{4} + \dfrac{(y - 2)^2}{9} = 1$. Ellipse centered at $(1, 2)$

26. $16x^2 + 9y^2 - 36y = 108 \quad \Leftrightarrow$
$16x^2 + 9(y^2 - 4y + 4) = 108 + 36 = 144 \quad \Leftrightarrow$
$\dfrac{x^2}{9} + \dfrac{(y - 2)^2}{16} = 1$. Ellipse centered at $(0, 2)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

27. $y = x^2 - 6x + 13 = \left(x^2 - 6x + 9\right) + 4 = (x-3)^2 + 4$. Parabola with vertex at $(3, 4)$

28. $x^2 - y^2 - 4x + 3 = 0 \quad \Leftrightarrow$
$\left(x^2 - 4x + 4\right) - y^2 = -3 + 4 = 1 \quad \Leftrightarrow$
$(x-2)^2 - y^2 = 1$. Hyperbola centered at $(2, 0)$

29. $x = 4 - y^2 = -y^2 + 4$. Parabola with vertex at $(4, 0)$

30. $y^2 - 2x + 6y + 5 = 0 \quad \Leftrightarrow \quad y^2 + 6y + 9 = 2x + 4 \quad \Leftrightarrow$
$(y+3)^2 = 2(x+2)$. Parabola with vertex $(-2, -3)$

31. $x^2 + 4y^2 - 6x + 5 = 0 \quad \Leftrightarrow$
$(x^2 - 6x + 9) + 4y^2 = -5 + 9 = 4 \quad \Leftrightarrow$
$\dfrac{(x-3)^2}{4} + y^2 = 1$. Ellipse centered at $(3, 0)$

32. $4x^2 + 9y^2 - 16x + 54y + 61 = 0 \quad \Leftrightarrow$
$4(x^2 - 4x + 4) + 9(y^2 + 6y + 9) = -61 + 16 + 81 = 36$
$\Leftrightarrow \quad \dfrac{(x-2)^2}{9} + \dfrac{(y+3)^2}{4} = 1$. Ellipse centered at $(2, -3)$

33. $y = 3x$ and $y = x^2$ intersect where $3x = x^2 \quad \Leftrightarrow$
$0 = x^2 - 3x = x(x-3)$, that is, at $(0, 0)$ and $(3, 9)$.

34. $y = 4 - x^2$, $x - 2y = 2$. Substitute y from the first equation into the second: $x - 2(4 - x^2) = 2 \quad \Leftrightarrow$
$2x^2 + x - 10 = 0 \quad \Leftrightarrow \quad (2x+5)(x-2) = 0 \quad \Leftrightarrow$
$x = -\frac{5}{2}$ or 2. So the points of intersection are $\left(-\frac{5}{2}, -\frac{9}{4}\right)$ and $(2, 0)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. The parabola must have an equation of the form $y = a(x-1)^2 - 1$. Substituting $x = 3$ and $y = 3$ into the equation gives $3 = a(3-1)^2 - 1$, so $a = 1$, and the equation is $y = (x-1)^2 - 1 = x^2 - 2x$. Note that using the other point $(-1, 3)$ would have given the same value for a, and hence the same equation.

36. The ellipse has an equation of the form $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1$. Substituting $x = 1$ and $y = -\dfrac{10\sqrt{2}}{3}$ gives $\dfrac{1^2}{a^2} + \dfrac{\left(-10\sqrt{2}/3\right)^2}{b^2} = \dfrac{1}{a^2} + \dfrac{200}{9b^2} = 1$. Substituting $x = -2$ and $y = \dfrac{5\sqrt{5}}{3}$ gives $\dfrac{(-2)^2}{a^2} + \dfrac{\left(5\sqrt{5}/3\right)^2}{b^2} = \dfrac{4}{a^2} + \dfrac{125}{9b^2} = 1$. From the first equation, $\dfrac{1}{a^2} = 1 - \dfrac{200}{9b^2}$. Putting this into the second equation gives $4\left(1 - \dfrac{200}{9b^2}\right) + \dfrac{125}{9b^2} = 1 \quad \Leftrightarrow$ $3 = \dfrac{675}{9b^2} \quad \Leftrightarrow \quad b^2 = \dfrac{675}{27} = 25$, so $b = 5$. Hence $\dfrac{1}{a^2} = 1 - \dfrac{200}{9(5)^2} = \dfrac{1}{9}$ and so $a = 3$. The equation of the ellipse is $\dfrac{x^2}{9} + \dfrac{y^2}{25} = 1$.

37. $\{(x, y) \mid x^2 + y^2 \le 1\}$

38. $\{(x, y) \mid x^2 + y^2 > 4\}$

39. $\{(x, y) \mid y \ge x^2 - 1\}$

40. $\{(x, y) \mid x^2 + 4y^2 \le 4\}$

D Trigonometry

1. $210^\circ = 210^\circ\left(\frac{\pi}{180^\circ}\right) = \frac{7\pi}{6}$ rad

2. $300^\circ = 300^\circ\left(\frac{\pi}{180^\circ}\right) = \frac{5\pi}{3}$ rad

3. $9^\circ = 9^\circ\left(\frac{\pi}{180^\circ}\right) = \frac{\pi}{20}$ rad

4. $-315^\circ = -315^\circ\left(\frac{\pi}{180^\circ}\right) = -\frac{7\pi}{4}$ rad

5. $900^\circ = 900^\circ\left(\frac{\pi}{180^\circ}\right) = 5\pi$ rad

6. $36^\circ = 36^\circ\left(\frac{\pi}{180^\circ}\right) = \frac{\pi}{5}$ rad

7. $4\pi \text{ rad} = 4\pi\left(\frac{180^\circ}{\pi}\right) = 720^\circ$

8. $-\frac{7\pi}{2} \text{ rad} = -\frac{7\pi}{2}\left(\frac{180^\circ}{\pi}\right) = -630^\circ$

9. $\frac{5\pi}{12} \text{ rad} = \frac{5\pi}{12}\left(\frac{180^\circ}{\pi}\right) = 75^\circ$

10. $\frac{8\pi}{3} \text{ rad} = \frac{8\pi}{3}\left(\frac{180^\circ}{\pi}\right) = 480^\circ$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. $-\frac{3\pi}{8}$ rad $= -\frac{3\pi}{8}\left(\frac{180^\circ}{\pi}\right) = -67.5^\circ$

12. 5 rad $= 5\left(\frac{180^\circ}{\pi}\right) = \left(\frac{900}{\pi}\right)^\circ$

13. Using Formula 3, $a = r\theta = 36 \cdot \frac{\pi}{12} = 3\pi$ cm.

14. Using Formula 3, $a = r\theta = 10 \cdot 72^\circ\left(\frac{\pi}{180^\circ}\right) = 4\pi$ cm.

15. Using Formula 3, $\theta = a/r = \frac{1}{1.5} = \frac{2}{3}$ rad $= \frac{2}{3}\left(\frac{180^\circ}{\pi}\right) = \left(\frac{120}{\pi}\right)^\circ \approx 38.2^\circ$.

16. $a = r\theta \quad \Rightarrow \quad r = \frac{a}{\theta} = \frac{6}{3\pi/4} = \frac{8}{\pi}$ cm

17.

18.

19.

20.

21.

22.

23.

From the diagram we see that a point on the terminal side is $P(-1, 1)$. Therefore, taking $x = -1$, $y = 1$, $r = \sqrt{2}$ in the definitions of the trigonometric ratios, we have $\sin\frac{3\pi}{4} = \frac{1}{\sqrt{2}}$, $\cos\frac{3\pi}{4} = -\frac{1}{\sqrt{2}}$, $\tan\frac{3\pi}{4} = -1$, $\csc\frac{3\pi}{4} = \sqrt{2}$, $\sec\frac{3\pi}{4} = -\sqrt{2}$, and $\cot\frac{3\pi}{4} = -1$.

24.

From the diagram and Figure 9, we see that a point on the terminal side is $P\left(-1, -\sqrt{3}\right)$. Therefore, taking $x = -1$, $y = -\sqrt{3}$, $r = 2$ in the definitions of the trigonometric ratios, we have $\sin\frac{4\pi}{3} = -\frac{\sqrt{3}}{2}$, $\cos\frac{4\pi}{3} = -\frac{1}{2}$, $\tan\frac{4\pi}{3} = \sqrt{3}$, $\csc\frac{4\pi}{3} = -\frac{2}{\sqrt{3}}$, $\sec\frac{4\pi}{3} = -2$, and $\cot\frac{4\pi}{3} = \frac{1}{\sqrt{3}}$.

25.

From the diagram we see that a point on the terminal side is $P(0, 1)$. Therefore taking $x = 0$, $y = 1$, $r = 1$ in the definitions of the trigonometric ratios, we have $\sin\frac{9\pi}{2} = 1$, $\cos\frac{9\pi}{2} = 0$, $\tan\frac{9\pi}{2} = y/x$ is undefined since $x = 0$, $\csc\frac{9\pi}{2} = 1$, $\sec\frac{9\pi}{2} = r/x$ is undefined since $x = 0$, and $\cot\frac{9\pi}{2} = 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26.

From the diagram, we see that a point on the terminal side is $P(-1, 0)$. Therefore taking $x = -1$, $y = 0$, $r = 1$ in the definitions of the trigonometric ratios we have $\sin(-5\pi) = 0$, $\cos(-5\pi) = -1$, $\tan(-5\pi) = 0$, $\csc(-5\pi)$ is undefined, $\sec(-5\pi) = -1$, and $\cot(-5\pi)$ is undefined.

27.

Using Figure 9 we see that a point on the terminal side is $P\left(-\sqrt{3}, 1\right)$. Therefore taking $x = -\sqrt{3}$, $y = 1$, $r = 2$ in the definitions of the trigonometric ratios, we have $\sin\frac{5\pi}{6} = \frac{1}{2}$, $\cos\frac{5\pi}{6} = -\frac{\sqrt{3}}{2}$, $\tan\frac{5\pi}{6} = -\frac{1}{\sqrt{3}}$, $\csc\frac{5\pi}{6} = 2$, $\sec\frac{5\pi}{6} = -\frac{2}{\sqrt{3}}$, and $\cot\frac{5\pi}{6} = -\sqrt{3}$.

28.

From the diagram, we see that a point on the terminal side is $P(-1, 1)$. Therefore taking $x = -1$, $y = 1$, $r = \sqrt{2}$ in the definitions of the trigonometric ratios we have $\sin\frac{11\pi}{4} = \frac{1}{\sqrt{2}}$, $\cos\frac{11\pi}{4} = -\frac{1}{\sqrt{2}}$, $\tan\frac{11\pi}{4} = -1$, $\csc\frac{11\pi}{4} = \sqrt{2}$, $\sec\frac{11\pi}{4} = -\sqrt{2}$, and $\cot\frac{11\pi}{4} = -1$.

29. $\sin\theta = y/r = \frac{3}{5} \quad\Rightarrow\quad y = 3$, $r = 5$, and $x = \sqrt{r^2 - y^2} = 4$ (since $0 < \theta < \frac{\pi}{2}$). Therefore taking $x = 4$, $y = 3$, $r = 5$ in the definitions of the trigonometric ratios, we have $\cos\theta = \frac{4}{5}$, $\tan\theta = \frac{3}{4}$, $\csc\theta = \frac{5}{3}$, $\sec\theta = \frac{5}{4}$, and $\cot\theta = \frac{4}{3}$.

30. Since $0 < \alpha < \frac{\pi}{2}$, α is in the first quadrant where x and y are both positive. Therefore, $\tan\alpha = y/x = \frac{2}{1} \quad\Rightarrow\quad y = 2$, $x = 1$, and $r = \sqrt{x^2 + y^2} = \sqrt{5}$. Taking $x = 1$, $y = 2$, $r = \sqrt{5}$ in the definitions of the trigonometric ratios, we have $\sin\alpha = \frac{2}{\sqrt{5}}$, $\cos\alpha = \frac{1}{\sqrt{5}}$, $\csc\alpha = \frac{\sqrt{5}}{2}$, $\sec\alpha = \sqrt{5}$, and $\cot\alpha = \frac{1}{2}$.

31. $\frac{\pi}{2} < \phi < \pi \quad\Rightarrow\quad \phi$ is in the second quadrant, where x is negative and y is positive. Therefore $\sec\phi = r/x = -1.5 = -\frac{3}{2} \quad\Rightarrow\quad r = 3$, $x = -2$, and $y = \sqrt{r^2 - x^2} = \sqrt{5}$. Taking $x = -2$, $y = \sqrt{5}$, and $r = 3$ in the definitions of the trigonometric ratios, we have $\sin\phi = \frac{\sqrt{5}}{3}$, $\cos\phi = -\frac{2}{3}$, $\tan\phi = -\frac{\sqrt{5}}{2}$, $\csc\phi = \frac{3}{\sqrt{5}}$, and $\cot\theta = -\frac{2}{\sqrt{5}}$.

32. Since $\pi < x < \frac{3\pi}{2}$, x is in the third quadrant where x and y are both negative. Therefore $\cos x = x/r = -\frac{1}{3} \quad\Rightarrow\quad x = -1$, $r = 3$, and $y = -\sqrt{r^2 - x^2} = -\sqrt{8} = -2\sqrt{2}$. Taking $x = -1$, $r = 3$, $y = -2\sqrt{2}$ in the definitions of the trigonometric ratios, we have $\sin x = -\frac{2\sqrt{2}}{3}$, $\tan x = 2\sqrt{2}$, $\csc x = -\frac{3}{2\sqrt{2}}$, $\sec x = -3$, and $\cot x = \frac{1}{2\sqrt{2}}$.

33. $\pi < \beta < 2\pi$ means that β is in the third or fourth quadrant where y is negative. Also since $\cot\beta = x/y = 3$ which is positive, x must also be negative. Therefore $\cot\beta = x/y = \frac{3}{1} \quad\Rightarrow\quad x = -3$, $y = -1$, and $r = \sqrt{x^2 + y^2} = \sqrt{10}$. Taking $x = -3$, $y = -1$ and $r = \sqrt{10}$ in the definitions of the trigonometric ratios, we have $\sin\beta = -\frac{1}{\sqrt{10}}$, $\cos\beta = -\frac{3}{\sqrt{10}}$, $\tan\beta = \frac{1}{3}$, $\csc\beta = -\sqrt{10}$, and $\sec\beta = -\frac{\sqrt{10}}{3}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. Since $\frac{3\pi}{2} < \theta < 2\pi$, θ is in the fourth quadrant where x is positive and y is negative. Therefore $\csc\theta = r/y = -\frac{4}{3} \Rightarrow$ $r = 4$, $y = -3$, and $x = \sqrt{r^2 - y^2} = \sqrt{7}$. Taking $x = \sqrt{7}$, $y = -3$, and $r = 4$ in the definitions of the trigonometric ratios, we have $\sin\theta = -\frac{3}{4}$, $\cos\theta = \frac{\sqrt{7}}{4}$, $\tan\theta = -\frac{3}{\sqrt{7}}$, $\sec\theta = \frac{4}{\sqrt{7}}$, and $\cot\theta = -\frac{\sqrt{7}}{3}$.

35. $\sin 35^\circ = \dfrac{x}{10} \quad\Rightarrow\quad x = 10\sin 35^\circ \approx 5.73576$ cm

36. $\cos 40^\circ = \dfrac{x}{25} \quad\Rightarrow\quad x = 25\cos 40^\circ \approx 19.15111$ cm

37. $\tan\frac{2\pi}{5} = \dfrac{x}{8} \quad\Rightarrow\quad x = 8\tan\frac{2\pi}{5} \approx 24.62147$ cm

38. $\cos\dfrac{3\pi}{8} = \dfrac{22}{x} \quad\Rightarrow\quad x = \dfrac{22}{\cos\dfrac{3\pi}{8}} \approx 57.48877$ cm

39.

(a) From the diagram we see that $\sin\theta = \dfrac{y}{r} = \dfrac{a}{c}$, and $\sin(-\theta) = \dfrac{-a}{c} = -\dfrac{a}{c} = -\sin\theta$.

(b) Again from the diagram we see that $\cos\theta = \dfrac{x}{r} = \dfrac{b}{c} = \cos(-\theta)$.

40. (a) Using (13a) and (13b), we have

$$\tan(x+y) = \frac{\sin(x+y)}{\cos(x+y)} = \frac{\sin x\,\cos y + \cos x\,\sin y}{\cos x\,\cos y - \sin x\,\sin y} = \frac{\dfrac{\sin x\,\cos y}{\cos x\,\cos y} + \dfrac{\cos x\,\sin y}{\cos x\,\cos y}}{\dfrac{\cos x\,\cos y}{\cos x\,\cos y} - \dfrac{\sin x\,\sin y}{\cos x\,\cos y}} = \frac{\tan x + \tan y}{1 - \tan x\,\tan y}$$

(b) From (11a) and (11b), we have $\tan(-\theta) = -\tan\theta$, so (15a) implies that

$$\tan(x-y) = \tan(x+(-y)) = \frac{\tan x + \tan(-y)}{1 - \tan x\,\tan(-y)} = \frac{\tan x - \tan y}{1 + \tan x\,\tan y}$$

41. (a) Using (13a) and (14a), we have

$$\tfrac{1}{2}[\sin(x+y) + \sin(x-y)] = \tfrac{1}{2}[\sin x\,\cos y + \cos x\,\sin y + \sin x\,\cos y - \cos x\,\sin y] = \tfrac{1}{2}(2\sin x\,\cos y) = \sin x\,\cos y.$$

(b) This time, using (13b) and (14b), we have

$$\tfrac{1}{2}[\cos(x+y) + \cos(x-y)] = \tfrac{1}{2}[\cos x\,\cos y - \sin x\,\sin y + \cos x\,\cos y + \sin x\,\sin y] = \tfrac{1}{2}(2\cos x\,\cos y) = \cos x\,\cos y.$$

(c) Again using (13b) and (14b), we have

$$\begin{aligned}\tfrac{1}{2}[\cos(x-y) - \cos(x+y)] &= \tfrac{1}{2}[\cos x\,\cos y + \sin x\,\sin y - \cos x\,\cos y + \sin x\,\sin y] \\ &= \tfrac{1}{2}(2\sin x\,\sin y) = \sin x\,\sin y\end{aligned}$$

42. Using (14b), $\cos\left(\frac{\pi}{2} - x\right) = \cos\frac{\pi}{2}\cos x + \sin\frac{\pi}{2}\sin x = 0\cdot\cos x + 1\cdot\sin x = \sin x$.

43. Using (13a), we have $\sin\left(\frac{\pi}{2} + x\right) = \sin\frac{\pi}{2}\cos x + \cos\frac{\pi}{2}\sin x = 1\cdot\cos x + 0\cdot\sin x = \cos x$.

44. Using (14a), we have $\sin(\pi - x) = \sin\pi\cos x - \cos\pi\,\sin x = 0\cdot\cos x - (-1)\sin x = \sin x$.

45. Using (7), we have $\sin\theta\cot\theta = \sin\theta\cdot\dfrac{\cos\theta}{\sin\theta} = \cos\theta$.

46. $(\sin x + \cos x)^2 = \sin^2 x + 2\sin x\,\cos x + \cos^2 x = (\sin^2 x + \cos^2 x) + \sin 2x$ [by (16a)] $= 1 + \sin 2x$ [by (8)]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $\sec y - \cos y = \dfrac{1}{\cos y} - \cos y$ [by (7)] $= \dfrac{1-\cos^2 y}{\cos y} = \dfrac{\sin^2 y}{\cos y}$ [by (8)] $= \dfrac{\sin y}{\cos y}\sin y = \tan y \sin y$ [by (7)]

48. $\tan^2\alpha - \sin^2\alpha = \dfrac{\sin^2\alpha}{\cos^2\alpha} - \sin^2\alpha = \dfrac{\sin^2\alpha - \sin^2\alpha\cos^2\alpha}{\cos^2\alpha} = \dfrac{\sin^2\alpha\,(1-\cos^2\alpha)}{\cos^2\alpha} = \tan^2\alpha\sin^2\alpha$ [by (7), (8)]

49. $\cot^2\theta + \sec^2\theta = \dfrac{\cos^2\theta}{\sin^2\theta} + \dfrac{1}{\cos^2\theta}$ [by (7)] $= \dfrac{\cos^2\theta\cos^2\theta + \sin^2\theta}{\sin^2\theta\cos^2\theta}$

$$= \frac{(1-\sin^2\theta)(1-\sin^2\theta)+\sin^2\theta}{\sin^2\theta\cos^2\theta} \text{ [by (8)] } = \frac{1-\sin^2\theta+\sin^4\theta}{\sin^2\theta\cos^2\theta}$$

$$= \frac{\cos^2\theta+\sin^4\theta}{\sin^2\theta\cos^2\theta} \text{ [by (7)] } = \frac{1}{\sin^2\theta} + \frac{\sin^2\theta}{\cos^2\theta} = \csc^2\theta + \tan^2\theta \text{ [by (7)]}$$

50. $2\csc 2t = \dfrac{2}{\sin 2t} = \dfrac{2}{2\sin t\cos t}$ [by (16a)] $= \dfrac{1}{\sin t\cos t} = \sec t\csc t$

51. Using (15a), we have $\tan 2\theta = \tan(\theta+\theta) = \dfrac{\tan\theta+\tan\theta}{1-\tan\theta\tan\theta} = \dfrac{2\tan\theta}{1-\tan^2\theta}$.

52. $\dfrac{1}{1-\sin\theta} + \dfrac{1}{1+\sin\theta} = \dfrac{1+\sin\theta+1-\sin\theta}{(1-\sin\theta)(1+\sin\theta)} = \dfrac{2}{1-\sin^2\theta} = \dfrac{2}{\cos^2\theta}$ [by (8)] $= 2\sec^2\theta$

53. Using (16a) and (17a),

$$\sin x\sin 2x + \cos x\cos 2x = \sin x\,(2\sin x\cos x) + \cos x\,(2\cos^2 x - 1) = 2\sin^2 x\cos x + 2\cos^3 x - \cos x$$

$$= 2(1-\cos^2 x)\cos x + 2\cos^3 x - \cos x \text{ [by (8)]}$$

$$= 2\cos x - 2\cos^3 x + 2\cos^3 x - \cos x = \cos x$$

Or: $\sin x\sin 2x + \cos x\cos 2x = \cos(2x - x)$ [by 14(b)] $= \cos x$

54. We start with the right side using equations (13a) and (14a):

$$\sin(x+y)\sin(x-y) = (\sin x\cos y + \cos x\sin y)(\sin x\cos y - \cos x\sin y)$$

$$= \sin^2 x\cos^2 y - \sin x\cos y\cos x\sin y + \cos x\sin y\sin x\cos y - \cos^2 x\sin^2 y$$

$$= \sin^2 x\,(1-\sin^2 y) - (1-\sin^2 x)\sin^2 y \quad \text{[by (8)]}$$

$$= \sin^2 x - \sin^2 x\sin^2 y - \sin^2 y + \sin^2 x\sin^2 y = \sin^2 x - \sin^2 y$$

55. $\dfrac{\sin\phi}{1-\cos\phi} = \dfrac{\sin\phi}{1-\cos\phi}\cdot\dfrac{1+\cos\phi}{1+\cos\phi} = \dfrac{\sin\phi\,(1+\cos\phi)}{1-\cos^2\phi} = \dfrac{\sin\phi\,(1+\cos\phi)}{\sin^2\phi}$ [by (8)]

$$= \frac{1+\cos\phi}{\sin\phi} = \frac{1}{\sin\phi} + \frac{\cos\phi}{\sin\phi} = \csc\phi + \cot\phi \quad \text{[by (7)]}$$

56. $\tan x + \tan y = \dfrac{\sin x}{\cos x} + \dfrac{\sin y}{\cos y} = \dfrac{\sin x\cos y + \cos x\sin y}{\cos x\cos y} = \dfrac{\sin(x+y)}{\cos x\cos y}$ [by (13a)]

57. Using (13a),

$$\sin 3\theta + \sin\theta = \sin(2\theta+\theta) + \sin\theta = \sin 2\theta\cos\theta + \cos 2\theta\sin\theta + \sin\theta$$

$$= \sin 2\theta\cos\theta + (2\cos^2\theta - 1)\sin\theta + \sin\theta \quad \text{[by (17a)]}$$

$$= \sin 2\theta\cos\theta + 2\cos^2\theta\sin\theta - \sin\theta + \sin\theta = \sin 2\theta\cos\theta + \sin 2\theta\cos\theta \quad \text{[by (16a)]}$$

$$= 2\sin 2\theta\cos\theta$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

58. We use (13b) with $x = 2\theta$, $y = \theta$ to get

$$\begin{aligned}\cos 3\theta &= \cos(2\theta + \theta) = \cos 2\theta\ \cos\theta - \sin 2\theta\ \sin\theta \\ &= (2\cos^2\theta - 1)\cos\theta - 2\sin^2\theta\ \cos\theta \quad \text{[by (17a) and (16a)]} \\ &= (2\cos^2\theta - 1)\cos\theta - 2(1 - \cos^2\theta)\ \cos\theta \quad \text{[by (8)]} \\ &= 2\cos^3\theta - \cos\theta - 2\cos\theta + 2\cos^3\theta = 4\cos^3\theta - 3\cos\theta\end{aligned}$$

59. Since $\sin x = \frac{1}{3}$ we can label the opposite side as having length 1, the hypotenuse as having length 3, and use the Pythagorean Theorem to get that the adjacent side has length $\sqrt{8}$. Then, from the diagram, $\cos x = \frac{\sqrt{8}}{3}$. Similarly we have that $\sin y = \frac{3}{5}$. Now use (13a):

$\sin(x + y) = \sin x\ \cos y + \cos x\ \sin y = \frac{1}{3} \cdot \frac{4}{5} + \frac{\sqrt{8}}{3} \cdot \frac{3}{5} = \frac{4}{15} + \frac{3\sqrt{8}}{15} = \frac{4 + 6\sqrt{2}}{15}$.

60. Use (13b) and the values for $\sin y$ and $\cos x$ obtained in Exercise 59 to get

$$\cos(x + y) = \cos x\ \cos y - \sin x\ \sin y = \frac{\sqrt{8}}{3} \cdot \frac{4}{5} - \frac{1}{3} \cdot \frac{3}{5} = \frac{8\sqrt{2} - 3}{15}$$

61. Using (14b) and the values for $\cos x$ and $\sin y$ obtained in Exercise 59, we have

$$\cos(x - y) = \cos x\ \cos y + \sin x\ \sin y = \frac{\sqrt{8}}{3} \cdot \frac{4}{5} + \frac{1}{3} \cdot \frac{3}{5} = \frac{8\sqrt{2} + 3}{15}$$

62. Using (14a) and the values for $\sin y$ and $\cos x$ obtained in Exercise 59, we get

$$\sin(x - y) = \sin x\ \cos y - \cos x\ \sin y = \frac{1}{3} \cdot \frac{4}{5} - \frac{\sqrt{8}}{3} \cdot \frac{3}{5} = \frac{4 - 6\sqrt{2}}{15}$$

63. Using (16a) and the values for $\sin y$ and $\cos y$ obtained in Exercise 59, we have $\sin 2y = 2\sin y\ \cos y = 2 \cdot \frac{3}{5} \cdot \frac{4}{5} = \frac{24}{25}$.

64. Using (17a) with $\cos y = \frac{4}{5}$, we have $\cos 2y = 2\cos^2 y - 1 = 2\left(\frac{4}{5}\right)^2 - 1 = \frac{32}{25} - 1 = \frac{7}{25}$.

65. $2\cos x - 1 = 0 \quad \Leftrightarrow \quad \cos x = \frac{1}{2} \quad \Rightarrow \quad x = \frac{\pi}{3}, \frac{5\pi}{3}$ for $x \in [0, 2\pi]$.

66. $3\cot^2 x = 1 \quad \Leftrightarrow \quad 3 = 1/\cot^2 x \quad \Leftrightarrow \quad \tan^2 x = 3 \quad \Leftrightarrow \quad \tan x = \pm\sqrt{3} \quad \Rightarrow \quad x = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}$, and $\frac{5\pi}{3}$.

67. $2\sin^2 x = 1 \quad \Leftrightarrow \quad \sin^2 x = \frac{1}{2} \quad \Leftrightarrow \quad \sin x = \pm\frac{1}{\sqrt{2}} \quad \Rightarrow \quad x = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$.

68. $|\tan x| = 1 \quad \Leftrightarrow \quad \tan x = -1$ or $\tan x = 1 \quad \Leftrightarrow \quad x = \frac{3\pi}{4}, \frac{7\pi}{4}$ or $x = \frac{\pi}{4}, \frac{5\pi}{4}$.

69. Using (16a), we have $\sin 2x = \cos x \quad \Leftrightarrow \quad 2\sin x\ \cos x - \cos x = 0 \quad \Leftrightarrow \quad \cos x(2\sin x - 1) = 0 \quad \Leftrightarrow \quad \cos x = 0$ or $2\sin x - 1 = 0 \quad \Rightarrow \quad x = \frac{\pi}{2}, \frac{3\pi}{2}$ or $\sin x = \frac{1}{2} \quad \Rightarrow \quad x = \frac{\pi}{6}$ or $\frac{5\pi}{6}$. Therefore, the solutions are $x = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}, \frac{3\pi}{2}$.

70. By (16a), $2\cos x + \sin 2x = 0 \quad \Leftrightarrow \quad 2\cos x + 2\sin x \cos x = 0 \quad \Leftrightarrow \quad 2\cos x\,(1 + \sin x) = 0 \quad \Leftrightarrow \quad \cos x = 0$ or $1 + \sin x = 0 \quad \Leftrightarrow \quad x = \frac{\pi}{2}, \frac{3\pi}{2}$ or $\sin x = -1 \quad \Rightarrow \quad x = \frac{3}{2}\pi$. So the solutions are $x = \frac{\pi}{2}, \frac{3\pi}{2}$.

71. $\sin x = \tan x \quad \Leftrightarrow \quad \sin x - \tan x = 0 \quad \Leftrightarrow \quad \sin x - \dfrac{\sin x}{\cos x} = 0 \quad \Leftrightarrow \quad \sin x\left(1 - \dfrac{1}{\cos x}\right) = 0 \quad \Leftrightarrow \quad \sin x = 0$ or $1 - \dfrac{1}{\cos x} = 0 \quad \Rightarrow \quad x = 0, \pi, 2\pi$ or $1 = \dfrac{1}{\cos x} \quad \Rightarrow \quad \cos x = 1 \quad \Rightarrow \quad x = 0, 2\pi$. Therefore the solutions are $x = 0, \pi, 2\pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

72. By (17a), $2+\cos 2x = 3\cos x \quad \Leftrightarrow \quad 2+2\cos^2 x - 1 = 3\cos x \quad \Leftrightarrow \quad 2\cos^2 x - 3\cos x + 1 = 0 \quad \Leftrightarrow$ $(2\cos x - 1)(\cos x - 1) = 0 \quad \Leftrightarrow \quad \cos x = 1$ or $\cos x = \frac{1}{2} \quad \Rightarrow \quad x = 0, 2\pi$ or $x = \frac{\pi}{3}, \frac{5\pi}{3}$.

73. We know that $\sin x = \frac{1}{2}$ when $x = \frac{\pi}{6}$ or $\frac{5\pi}{6}$, and from Figure 18(a), we see that $\sin x \le \frac{1}{2} \quad \Rightarrow \quad 0 \le x \le \frac{\pi}{6}$ or $\frac{5\pi}{6} \le x \le 2\pi$ for $x \in [0, 2\pi]$.

74. $2\cos x + 1 > 0 \quad \Rightarrow \quad 2\cos x > -1 \quad \Rightarrow \quad \cos x > -\frac{1}{2}$. $\cos x = -\frac{1}{2}$ when $x = \frac{2\pi}{3}, \frac{4\pi}{3}$ and from Figure 18(b), we see that $\cos\, x > -\frac{1}{2}$ when $0 \le x < \frac{2\pi}{3}, \frac{4\pi}{3} < x \le 2\pi$.

75. $\tan x = -1$ when $x = \frac{3\pi}{4}, \frac{7\pi}{4}$, and $\tan x = 1$ when $x = \frac{\pi}{4}$ or $\frac{5\pi}{4}$. From Figure 19(a) we see that $-1 < \tan x < 1 \quad \Rightarrow$ $0 \le x < \frac{\pi}{4}, \frac{3\pi}{4} < x < \frac{5\pi}{4}$, and $\frac{7\pi}{4} < x \le 2\pi$.

76. We know that $\sin x = \cos x$ when $x = \frac{\pi}{4}, \frac{5\pi}{4}$, and from the diagram we see that $\sin x > \cos x$ when $\frac{\pi}{4} < x < \frac{5\pi}{4}$.

77. $\angle C = 180^\circ - 50^\circ - 68^\circ = 62^\circ$.

Then $a = \dfrac{230\sin 50^\circ}{\sin 62^\circ} \approx 200$ and $b = \dfrac{230\sin 68^\circ}{\sin 62^\circ} \approx 242$.

78. $c^2 = 3^2 + 4^2 - 2\cdot 3\cdot 4\cdot \cos 53^\circ = 9 + 16 - 24\cos 53^\circ \approx 10.556 \quad \Leftrightarrow \quad c \approx \sqrt{10.556} \approx 3.2$. Then $\sin B = \dfrac{4\sin 53^\circ}{3.25} \approx 0.983 \quad \Leftrightarrow \quad \angle B \approx \sin^{-1}(0.983) \approx 79^\circ$ and $\angle A \approx 180^\circ - 53^\circ - 79^\circ = 48^\circ$.

79. $|AB|^2 = |AC|^2 + |BC|^2 - 2\,|AC|\,|BC|\cos\angle C = (820)^2 + (910)^2 - 2(820)(910)\cos 103^\circ \approx 1{,}836{,}217 \quad \Rightarrow$ $|AB| \approx 1355$ m

80. By the Law of Cosines we have

$b^2 = 200^2 + 100^2 - 2\cdot 200\cdot 100\cos 160^\circ \approx 87{,}587.70$. Thus $b \approx 295.95$.

To find $\angle A$ we first use the Law of Sines:

$\dfrac{\sin A}{100} \approx \dfrac{\sin 160^\circ}{295.95} \quad \Leftrightarrow \quad \sin A = 100\cdot\dfrac{\sin 160^\circ}{295.95} \approx 0.11557 \quad \Leftrightarrow$

$A = \sin^{-1}(0.11557) \approx 6.636^\circ$

81. Using the formula from (6), the area of the triangle is $\frac{1}{2}(10)(3)\sin 107^\circ \approx 14.34457\ \text{cm}^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

82. By Heron's formula we have $\mathcal{A} = \sqrt{s(s-a)(s-b)(s-c)}$, where $s = \frac{1}{2}(4+5+7) = 8$. So then, $\mathcal{A} = \sqrt{8(8-4)(8-5)(8-7)} = 4\sqrt{6} \approx 9.80$.

83. $y = \cos\left(x - \frac{\pi}{3}\right)$. We start with the graph of $y = \cos x$ and shift it $\frac{\pi}{3}$ units to the right.

84. $y = \tan 2x$. Start with the graph of $y = \tan x$ with period π and compress it to a period of $\frac{\pi}{2}$.

85. $y = \frac{1}{3}\tan\left(x - \frac{\pi}{2}\right)$. We start with the graph of $y = \tan x$, shift it $\frac{\pi}{2}$ units to the right and compress it to $\frac{1}{3}$ of its original vertical size.

86. $y = 1 + \sec x$. Start with the graph of $y = \sec x$ and raise it by one unit.

87. $y = |\sin x|$. We start with the graph of $y = \sin x$ and reflect the parts below the x-axis about the x-axis.

88. $y = 2 + \sin\left(x + \frac{\pi}{4}\right)$. Start with the graph of $y = \sin x$, and shift it $\frac{\pi}{4}$ units to the left and 2 units up.

89. Using the Law of Cosines, we have $c^2 = 1^2 + 1^2 - 2(1)(1)\cos(\alpha - \beta) = 2[1 - \cos(\alpha - \beta)]$. Now, using the distance formula, $c^2 = |AB|^2 = (\cos\alpha - \cos\beta)^2 + (\sin\alpha - \sin\beta)^2$. Equating these two expressions for c^2, we get

$2[1 - \cos(\alpha - \beta)] = \cos^2\alpha + \sin^2\alpha + \cos^2\beta + \sin^2\beta - 2\cos\alpha\,\cos\beta - 2\sin\alpha\,\sin\beta \quad \Rightarrow$

$1 - \cos(\alpha - \beta) = 1 - \cos\alpha\,\cos\beta - \sin\alpha\,\sin\beta \quad \Rightarrow \quad \cos(\alpha - \beta) = \cos\alpha\,\cos\beta + \sin\alpha\,\sin\beta.$

90. $\cos(x + y) = \cos(x - (-y)) = \cos x\,\cos(-y) + \sin x\,\sin(-y)$

$= \cos x\,\cos y - \sin x\,\sin y$ [using Equations (11a) and (11b)]

91. In Exercise 90 we used the subtraction formula for cosine to prove the addition formula for cosine. Using that formula with $x = \frac{\pi}{2} - \alpha$, $y = \beta$, we get $\cos\left[\left(\frac{\pi}{2} - \alpha\right) + \beta\right] = \cos\left(\frac{\pi}{2} - \alpha\right)\cos\beta - \sin\left(\frac{\pi}{2} - \alpha\right)\sin\beta \quad \Rightarrow$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\cos\left[\frac{\pi}{2}-(\alpha-\beta)\right]=\cos\left(\frac{\pi}{2}-\alpha\right)\cos\beta-\sin\left(\frac{\pi}{2}-\alpha\right)\sin\beta$. Now we use the identities given in the problem, $\cos\left(\frac{\pi}{2}-\theta\right)=\sin\theta$ and $\sin\left(\frac{\pi}{2}-\theta\right)=\cos\theta$, to get $\sin(\alpha-\beta)=\sin\alpha\,\cos\beta-\cos\alpha\,\sin\beta$.

E Sigma Notation

1. $\sum_{i=1}^{5}\sqrt{i}=\sqrt{1}+\sqrt{2}+\sqrt{3}+\sqrt{4}+\sqrt{5}$

2. $\sum_{i=1}^{6}\frac{1}{i+1}=\frac{1}{2}+\frac{1}{3}+\frac{1}{4}+\frac{1}{5}+\frac{1}{6}+\frac{1}{7}$

3. $\sum_{i=4}^{6}3^i=3^4+3^5+3^6$

4. $\sum_{i=4}^{6}i^3=4^3+5^3+6^3$

5. $\sum_{k=0}^{4}\frac{2k-1}{2k+1}=-1+\frac{1}{3}+\frac{3}{5}+\frac{5}{7}+\frac{7}{9}$

6. $\sum_{k=5}^{8}x^k=x^5+x^6+x^7+x^8$

7. $\sum_{i=1}^{n}i^{10}=1^{10}+2^{10}+3^{10}+\cdots+n^{10}$

8. $\sum_{j=n}^{n+3}j^2=n^2+(n+1)^2+(n+2)^2+(n+3)^2$

9. $\sum_{j=0}^{n-1}(-1)^j=1-1+1-1+\cdots+(-1)^{n-1}$

10. $\sum_{i=1}^{n}f(x_i)\,\Delta x_i=f(x_1)\,\Delta x_1+f(x_2)\,\Delta x_2+f(x_3)\,\Delta x_3+\cdots+f(x_n)\,\Delta x_n$

11. $1+2+3+4+\cdots+10=\sum_{i=1}^{10}i$

12. $\sqrt{3}+\sqrt{4}+\sqrt{5}+\sqrt{6}+\sqrt{7}=\sum_{i=3}^{7}\sqrt{i}$

13. $\frac{1}{2}+\frac{2}{3}+\frac{3}{4}+\frac{4}{5}+\cdots+\frac{19}{20}=\sum_{i=1}^{19}\frac{i}{i+1}$

14. $\frac{3}{7}+\frac{4}{8}+\frac{5}{9}+\frac{6}{10}+\cdots+\frac{23}{27}=\sum_{i=3}^{23}\frac{i}{i+4}$

15. $2+4+6+8+\cdots+2n=\sum_{i=1}^{n}2i$

16. $1+3+5+7+\cdots+(2n-1)=\sum_{i=1}^{n}(2i-1)$

17. $1+2+4+8+16+32=\sum_{i=0}^{5}2^i$

18. $\frac{1}{1}+\frac{1}{4}+\frac{1}{9}+\frac{1}{16}+\frac{1}{25}+\frac{1}{36}=\sum_{i=1}^{6}\frac{1}{i^2}$

19. $x+x^2+x^3+\cdots+x^n=\sum_{i=1}^{n}x^i$

20. $1-x+x^2-x^3+\cdots+(-1)^n\,x^n=\sum_{i=0}^{n}(-1)^i x^i$

21. $\sum_{i=4}^{8}(3i-2)=[3(4)-2]+[3(5)-2]+[3(6)-2]+[3(7)-2]+[3(8)-2]=10+13+16+19+22=80$

22. $\sum_{i=3}^{6}i(i+2)=3\cdot5+4\cdot6+5\cdot7+6\cdot8=15+24+35+48=122$

23. $\sum_{j=1}^{6}3^{j+1}=3^2+3^3+3^4+3^5+3^6+3^7=9+27+81+243+729+2187=3276$

(For a more general method, see Exercise 47.)

24. $\sum_{k=0}^{8}\cos k\pi=\cos 0+\cos\pi+\cos 2\pi+\cos 3\pi+\cos 4\pi+\cos 5\pi+\cos 6\pi+\cos 7\pi+\cos 8\pi$

$=1-1+1-1+1-1+1-1+1=1$

25. $\sum_{n=1}^{20}(-1)^n=-1+1-1+1-1+1-1+1-1+1-1+1-1+1-1+1-1+1-1+1-1+1=0$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. $\sum_{i=1}^{100} 4 = \underbrace{4+4+4+\cdots+4}_{\text{(100 summands)}} = 100 \cdot 4 = 400$

27. $\sum_{i=0}^{4} (2^i + i^2) = (1+0)+(2+1)+(4+4)+(8+9)+(16+16) = 61$

28. $\sum_{i=-2}^{4} 2^{3-i} = 2^5 + 2^4 + 2^3 + 2^2 + 2^1 + 2^0 + 2^{-1} = 63.5$

29. $\sum_{i=1}^{n} 2i = 2\sum_{i=1}^{n} i = 2 \cdot \dfrac{n(n+1)}{2}$ [by Theorem 3(c)] $= n(n+1)$

30. $\sum_{i=1}^{n} (2-5i) = \sum_{i=1}^{n} 2 - \sum_{i=1}^{n} 5i = 2n - 5\sum_{i=1}^{n} i = 2n - \dfrac{5n(n+1)}{2} = \dfrac{4n}{2} - \dfrac{5n^2+5n}{2} = -\dfrac{n(5n+1)}{2}$

31. $\sum_{i=1}^{n} (i^2+3i+4) = \sum_{i=1}^{n} i^2 + 3\sum_{i=1}^{n} i + \sum_{i=1}^{n} 4 = \dfrac{n(n+1)(2n+1)}{6} + \dfrac{3n(n+1)}{2} + 4n$

$= \frac{1}{6}[(2n^3+3n^2+n)+(9n^2+9n)+24n] = \frac{1}{6}(2n^3+12n^2+34n) = \frac{1}{3}n(n^2+6n+17)$

32. $\sum_{i=1}^{n} (3+2i)^2 = \sum_{i=1}^{n} (9+12i+4i^2) = \sum_{i=1}^{n} 9 + 12\sum_{i=1}^{n} i + 4\sum_{i=1}^{n} i^2 = 9n + 6n(n+1) + \dfrac{2n(n+1)(2n+1)}{3}$

$= \dfrac{27n+18n^2+18n+4n^3+6n^2+2n}{3} = \frac{1}{3}(4n^3+24n^2+47n) = \frac{1}{3}n(4n^2+24n+47)$

33. $\sum_{i=1}^{n} (i+1)(i+2) = \sum_{i=1}^{n} (i^2+3i+2) = \sum_{i=1}^{n} i^2 + 3\sum_{i=1}^{n} i + \sum_{i=1}^{n} 2 = \dfrac{n(n+1)(2n+1)}{6} + \dfrac{3n(n+1)}{2} + 2n$

$= \dfrac{n(n+1)}{6}[(2n+1)+9] + 2n = \dfrac{n(n+1)}{3}(n+5) + 2n$

$= \dfrac{n}{3}[(n+1)(n+5)+6] = \dfrac{n}{3}(n^2+6n+11)$

34. $\sum_{i=1}^{n} i(i+1)(i+2) = \sum_{i=1}^{n} (i^3+3i^2+2i) = \sum_{i=1}^{n} i^3 + 3\sum_{i=1}^{n} i^2 + 2\sum_{i=1}^{n} i$

$= \left[\dfrac{n(n+1)}{2}\right]^2 + \dfrac{3n(n+1)(2n+1)}{6} + \dfrac{2n(n+1)}{2}$

$= n(n+1)\left[\dfrac{n(n+1)}{4} + \dfrac{2n+1}{2} + 1\right] = \dfrac{n(n+1)}{4}(n^2+n+4n+2+4)$

$= \dfrac{n(n+1)}{4}(n^2+5n+6) = \dfrac{n(n+1)(n+2)(n+3)}{4}$

35. $\sum_{i=1}^{n} (i^3-i-2) = \sum_{i=1}^{n} i^3 - \sum_{i=1}^{n} i - \sum_{i=1}^{n} 2 = \left[\dfrac{n(n+1)}{2}\right]^2 - \dfrac{n(n+1)}{2} - 2n$

$= \frac{1}{4}n(n+1)[n(n+1)-2] - 2n = \frac{1}{4}n(n+1)(n+2)(n-1) - 2n$

$= \frac{1}{4}n[(n+1)(n-1)(n+2)-8] = \frac{1}{4}n[(n^2-1)(n+2)-8] = \frac{1}{4}n(n^3+2n^2-n-10)$

36. By Theorem 3(c) we have that $\sum_{i=1}^{n} i = \dfrac{n(n+1)}{2} = 78 \quad\Leftrightarrow\quad n(n+1) = 156 \quad\Leftrightarrow\quad n^2+n-156 = 0 \quad\Leftrightarrow$

$(n+13)(n-12) = 0 \quad\Leftrightarrow\quad n = 12$ or -13. But $n = -13$ produces a negative answer for the sum, so $n = 12$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

37. By Theorem 2(a) and Example 3, $\sum_{i=1}^{n} c = c\sum_{i=1}^{n} 1 = cn$.

38. Let S_n be the statement that $\sum_{i=1}^{n} i^3 = \left[\frac{n(n+1)}{2}\right]^2$.

1. S_1 is true because $1^3 = \left(\frac{1\cdot 2}{2}\right)^2$.

2. Assume S_k is true. Then $\sum_{i=1}^{k} i^3 = \left[\frac{k(k+1)}{2}\right]^2$, so

$$\sum_{i=1}^{k+1} i^3 = \left[\frac{k(k+1)}{2}\right]^2 + (k+1)^3 = \frac{(k+1)^2}{4}\left[k^2+4(k+1)\right] = \frac{(k+1)^2}{4}(k+2)^2 = \left(\frac{(k+1)[(k+1)+1]}{2}\right)^2$$

showing that S_{k+1} is true.

Therefore, S_n is true for all n by mathematical induction.

39. $\sum_{i=1}^{n}[(i+1)^4 - i^4] = (2^4-1^4)+(3^4-2^4)+(4^4-3^4)+\cdots+\left[(n+1)^4-n^4\right]$

$$= (n+1)^4 - 1^4 = n^4+4n^3+6n^2+4n$$

On the other hand,

$$\sum_{i=1}^{n}[(i+1)^4-i^4] = \sum_{i=1}^{n}(4i^3+6i^2+4i+1) = 4\sum_{i=1}^{n} i^3 + 6\sum_{i=1}^{n} i^2 + 4\sum_{i=1}^{n} i + \sum_{i=1}^{n} 1$$

$$= 4S + n(n+1)(2n+1) + 2n(n+1) + n \qquad \left[\text{where } S = \sum_{i=1}^{n} i^3\right]$$

$$= 4S + 2n^3 + 3n^2 + n + 2n^2 + 2n + n = 4S + 2n^3 + 5n^2 + 4n$$

Thus, $n^4+4n^3+6n^2+4n = 4S+2n^3+5n^2+4n$, from which it follows that

$4S = n^4+2n^3+n^2 = n^2(n^2+2n+1) = n^2(n+1)^2$ and $S = \left[\frac{n(n+1)}{2}\right]^2$.

40. The area of G_i is

$$\left(\sum_{k=1}^{i} k\right)^2 - \left(\sum_{k=1}^{i-1} k\right)^2 = \left[\frac{i(i+1)}{2}\right]^2 - \left[\frac{(i-1)i}{2}\right]^2 = \frac{i^2}{4}\left[(i+1)^2-(i-1)^2\right]$$

$$= \frac{i^2}{4}\left[(i^2+2i+1)-(i^2-2i+1)\right] = \frac{i^2}{4}(4i) = i^3$$

Thus, the area of $ABCD$ is $\sum_{i=1}^{n} i^3 = \left[\frac{n(n+1)}{2}\right]^2$.

41. (a) $\sum_{i=1}^{n}\left[i^4-(i-1)^4\right] = \left(1^4-0^4\right)+\left(2^4-1^4\right)+\left(3^4-2^4\right)+\cdots+\left[n^4-(n-1)^4\right] = n^4-0 = n^4$

(b) $\sum_{i=1}^{100}\left(5^i-5^{i-1}\right) = \left(5^1-5^0\right)+\left(5^2-5^1\right)+\left(5^3-5^2\right)+\cdots+\left(5^{100}-5^{99}\right) = 5^{100}-5^0 = 5^{100}-1$

(c) $\sum_{i=3}^{99}\left(\frac{1}{i}-\frac{1}{i+1}\right) = \left(\frac{1}{3}-\frac{1}{4}\right)+\left(\frac{1}{4}-\frac{1}{5}\right)+\left(\frac{1}{5}-\frac{1}{6}\right)+\cdots+\left(\frac{1}{99}-\frac{1}{100}\right) = \frac{1}{3}-\frac{1}{100} = \frac{97}{300}$

(d) $\sum_{i=1}^{n}(a_i-a_{i-1}) = (a_1-a_0)+(a_2-a_1)+(a_3-a_2)+\cdots+(a_n-a_{n-1}) = a_n-a_0$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

42. Summing the inequalities $-|a_i| \le a_i \le |a_i|$ for $i = 1, 2, \ldots, n$, we get $-\sum_{i=1}^{n} |a_i| \le \sum_{i=1}^{n} a_i \le \sum_{i=1}^{n} |a_i|$. Since $|x| \le c \quad \Leftrightarrow$

$-c \le x \le c$, we have $\left|\sum_{i=1}^{n} a_i\right| \le \sum_{i=1}^{n} |a_i|$. *Another method:* Use mathematical induction.

43. $\lim_{n\to\infty} \sum_{i=1}^{n} \frac{1}{n}\left(\frac{i}{n}\right)^2 = \lim_{n\to\infty} \frac{1}{n^3} \sum_{i=1}^{n} i^2 = \lim_{n\to\infty} \frac{1}{n^3} \frac{n(n+1)(2n+1)}{6} = \lim_{n\to\infty} \frac{1}{6}\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right) = \frac{1}{6}(1)(2) = \frac{1}{3}$

44. $\lim_{n\to\infty} \sum_{i=1}^{n} \frac{1}{n}\left[\left(\frac{i}{n}\right)^3 + 1\right] = \lim_{n\to\infty} \sum_{i=1}^{n} \left[\frac{i^3}{n^4} + \frac{1}{n}\right] = \lim_{n\to\infty} \left[\frac{1}{n^4} \sum_{i=1}^{n} i^3 + \frac{1}{n} \sum_{i=1}^{n} 1\right] = \lim_{n\to\infty} \left[\frac{1}{n^4}\left(\frac{n(n+1)}{2}\right)^2 + \frac{1}{n}(n)\right]$

$$= \lim_{n\to\infty} \frac{1}{4}\left(1+\frac{1}{n}\right)^2 + 1 = \tfrac{1}{4} + 1 = \tfrac{5}{4}$$

45. $\lim_{n\to\infty} \sum_{i=1}^{n} \frac{2}{n}\left[\left(\frac{2i}{n}\right)^3 + 5\left(\frac{2i}{n}\right)\right] = \lim_{n\to\infty} \sum_{i=1}^{n} \left[\frac{16}{n^4} i^3 + \frac{20}{n^2} i\right] = \lim_{n\to\infty} \left[\frac{16}{n^4} \sum_{i=1}^{n} i^3 + \frac{20}{n^2} \sum_{i=1}^{n} i\right]$

$$= \lim_{n\to\infty} \left[\frac{16}{n^4} \frac{n^2(n+1)^2}{4} + \frac{20}{n^2} \frac{n(n+1)}{2}\right] = \lim_{n\to\infty} \left[\frac{4(n+1)^2}{n^2} + \frac{10n(n+1)}{n^2}\right]$$

$$= \lim_{n\to\infty} \left[4\left(1+\frac{1}{n}\right)^2 + 10\left(1+\frac{1}{n}\right)\right] = 4 \cdot 1 + 10 \cdot 1 = 14$$

46. $\lim_{n\to\infty} \sum_{i=1}^{n} \frac{3}{n}\left[\left(1+\frac{3i}{n}\right)^3 - 2\left(1+\frac{3i}{n}\right)\right] = \lim_{n\to\infty} \sum_{i=1}^{n} \frac{3}{n}\left[1 + \frac{9i}{n} + \frac{27i^2}{n^2} + \frac{27i^3}{n^3} - 2 - \frac{6i}{n}\right]$

$$= \lim_{n\to\infty} \sum_{i=1}^{n} \left[\frac{81}{n^4} i^3 + \frac{81}{n^3} i^2 + \frac{9}{n^2} i - \frac{3}{n}\right]$$

$$= \lim_{n\to\infty} \left[\frac{81}{n^4} \frac{n^2(n+1)^2}{4} + \frac{81}{n^3} \frac{n(n+1)(2n+1)}{6} + \frac{9}{n^2} \frac{n(n+1)}{2} - \frac{3}{n} n\right]$$

$$= \lim_{n\to\infty} \left[\frac{81}{4}\left(1+\frac{1}{n}\right)^2 + \frac{27}{2}\left(1+\frac{1}{n}\right)\left(2+\frac{1}{n}\right) + \frac{9}{2}\left(1+\frac{1}{n}\right) - 3\right]$$

$$= \tfrac{81}{4} + \tfrac{54}{2} + \tfrac{9}{2} - 3 = \tfrac{195}{4}$$

47. Let $S = \sum_{i=1}^{n} ar^{i-1} = a + ar + ar^2 + \cdots + ar^{n-1}$. Multiplying both sides by r gives us

$rS = ar + ar^2 + \cdots + ar^{n-1} + ar^n$. Subtracting the first equation from the second, we find

$(r-1)S = ar^n - a = a(r^n - 1)$, so $S = \dfrac{a(r^n - 1)}{r - 1}$ [since $r \ne 1$].

48. $\sum_{i=1}^{n} \frac{3}{2^{i-1}} = 3 \sum_{i=1}^{n} \left(\frac{1}{2}\right)^{i-1} = \dfrac{3\left[\left(\frac{1}{2}\right)^n - 1\right]}{\frac{1}{2} - 1}$ [using Exercise 47 with $a = 3$ and $r = \frac{1}{2}$] $= 6\left[1 - \left(\frac{1}{2}\right)^n\right]$

49. $\sum_{i=1}^{n} (2i + 2^i) = 2 \sum_{i=1}^{n} i + \sum_{i=1}^{n} 2 \cdot 2^{i-1} = 2\frac{n(n+1)}{2} + \frac{2(2^n - 1)}{2 - 1} = 2^{n+1} + n^2 + n - 2.$

For the first sum we have used Theorems 2(a) and 3(c), and for the second, Exercise 47 with $a = r = 2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50. $\displaystyle\sum_{i=1}^{m}\left[\sum_{j=1}^{n}(i+j)\right] = \sum_{i=1}^{m}\left[\sum_{j=1}^{n} i + \sum_{j=1}^{n} j\right]$ [Theorem 2(b)] $\displaystyle = \sum_{i=1}^{m}\left[ni + \frac{n(n+1)}{2}\right]$ [Theorem 3(b) and 3(c)]

$$= \sum_{i=1}^{m} ni + \sum_{i=1}^{m}\frac{n(n+1)}{2} = \frac{nm(m+1)}{2} + \frac{nm(n+1)}{2} = \frac{nm}{2}(m+n+2)$$

G The Logarithm Defined as an Integral

1. (a)

We interpret $\ln 1.5$ as the area under the curve $y = 1/x$ from $x = 1$ to $x = 1.5$. The area of the rectangle $BCDE$ is $\frac{1}{2}\cdot\frac{2}{3} = \frac{1}{3}$. The area of the trapezoid $ABCD$ is $\frac{1}{2}\cdot\frac{1}{2}\left(1+\frac{2}{3}\right) = \frac{5}{12}$. Thus, by comparing areas, we observe that $\frac{1}{3} < \ln 1.5 < \frac{5}{12}$.

(b) $\ln x = \int_1^x (1/t)\,dt$, so $\ln 1.5 = \int_1^{1.5}(1/t)\,dt$. With $f(t) = 1/t$, $n = 10$, and $\Delta t = \frac{1.5-1}{10} = 0.05$, we have

$$\ln 1.5 = \int_1^{1.5}(1/t)\,dt \approx (0.05)[f(1.025) + f(1.075) + \cdots + f(1.475)] = (0.05)\left[\tfrac{1}{1.025} + \tfrac{1}{1.075} + \cdots + \tfrac{1}{1.475}\right]$$
$$\approx 0.4054$$

2. (a) $y = \dfrac{1}{t}$, $y' = -\dfrac{1}{t^2}$. The slope of the line through $A(1,1)$ and $D\left(2, \frac{1}{2}\right)$ is $\dfrac{1/2-1}{2-1} = -\dfrac{1}{2}$. Let c be the t-coordinate of the point on $y = \dfrac{1}{t}$ with slope $-\dfrac{1}{2}$. Then $-\dfrac{1}{c^2} = -\dfrac{1}{2} \;\Rightarrow\; c^2 = 2 \;\Rightarrow\; c = \sqrt{2}$ since $c > 0$. Therefore, the tangent line is given by $y - \frac{1}{\sqrt{2}} = -\frac{1}{2}\left(t - \sqrt{2}\right)$, or $y = -\frac{1}{2}t + \sqrt{2}$.

(b)

Since the graph of $y = 1/t$ is concave upward, the graph lies above the tangent line, that is, above the line segment BC. Now $|AB| = -\frac{1}{2} + \sqrt{2}$ and $|CD| = -1 + \sqrt{2}$. The area of the trapezoid $ABCD$ is $\frac{1}{2}\left[\left(-\frac{1}{2} + \sqrt{2}\right) + \left(-1 + \sqrt{2}\right)\right]1 = -\frac{3}{4} + \sqrt{2} \approx 0.6642$. So $\ln 2 >$ area of trapezoid $ABCD > 0.66$.

3.

The area of R_i is $\dfrac{1}{i+1}$ and so $\dfrac{1}{2} + \dfrac{1}{3} + \cdots + \dfrac{1}{n} < \displaystyle\int_1^n \frac{1}{t}\,dt = \ln n$.

The area of S_i is $\dfrac{1}{i}$ and so $1 + \dfrac{1}{2} + \cdots + \dfrac{1}{n-1} > \displaystyle\int_1^n \frac{1}{t}\,dt = \ln n$.

Thus, $\dfrac{1}{2} + \dfrac{1}{3} + \cdots + \dfrac{1}{n} < \ln n < 1 + \dfrac{1}{2} + \cdots + \dfrac{1}{n-1}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a) From the diagram, we see that the area under the graph of $y = 1/x$ between $x = 1$ and $x = 2$ is less than the area of the square, which is 1. So $\ln 2 = \int_1^2 (1/x)\,dx < 1$. To show the other side of the inequality, we must find an area larger than 1 which lies under the graph of $y = 1/x$ between $x = 1$ and $x = 3$. One way to do this is to partition the interval $[1, 3]$ into 8 intervals of equal length and calculate the resulting Riemann sum, using the right endpoints:

$$\frac{1}{4}\left(\frac{1}{5/4} + \frac{1}{3/2} + \frac{1}{7/4} + \frac{1}{2} + \frac{1}{9/4} + \frac{1}{5/2} + \frac{1}{11/4} + \frac{1}{3}\right) = \frac{28{,}271}{27{,}720} > 1$$

and therefore $1 < \int_1^3 (1/x)\,dx = \ln 3$.

A slightly easier method uses the fact that since $y = 1/x$ is concave upward, it lies above all its tangent lines. Drawing two such tangent lines at the points $\left(\frac{3}{2}, \frac{2}{3}\right)$ and $\left(\frac{5}{2}, \frac{2}{5}\right)$, we see that the area under the curve from $x = 1$ to $x = 3$ is more than the sum of the areas of the two trapezoids, that is, $\frac{2}{3} + \frac{2}{5} = \frac{16}{15}$. Thus, $1 < \frac{16}{15} < \int_1^3 (1/x)\,dx = \ln 3$.

(b) By part (a), $\ln 2 < 1 < \ln 3$. But e is defined such that $\ln e = 1$, and because the natural logarithm function is increasing, we have $\ln 2 < \ln e < \ln 3 \quad \Leftrightarrow \quad 2 < e < 3$.

5. If $f(x) = \ln(x^r)$, then $f'(x) = (1/x^r)\,(rx^{r-1}) = r/x$. But if $g(x) = r \ln x$, then $g'(x) = r/x$. So f and g must differ by a constant: $\ln(x^r) = r \ln x + C$. Put $x = 1$: $\ln(1^r) = r \ln 1 + C \quad \Rightarrow \quad C = 0$, so $\ln(x^r) = r \ln x$.

6. Using the second law of logarithms and Equation 10, we have $\ln(e^x/e^y) = \ln e^x - \ln e^y = x - y = \ln(e^{x-y})$. Since $\ln$ is a one-to-one function, it follows that $e^x/e^y = e^{x-y}$.

7. Using the third law of logarithms and Equation 10, we have $\ln e^{rx} = rx = r \ln e^x = \ln(e^x)^r$. Since $\ln$ is a one-to-one function, it follows that $e^{rx} = (e^x)^r$.

8. Using Definition 13 and the second law of exponents for e^x, we have $a^{x-y} = e^{(x-y)\ln a} = e^{x \ln a - y \ln a} = \dfrac{e^{x \ln a}}{e^{y \ln a}} = \dfrac{a^x}{a^y}$.

9. Using Definition 13, the first law of logarithms, and the first law of exponents for e^x, we have $(ab)^x = e^{x \ln(ab)} = e^{x(\ln a + \ln b)} = e^{x \ln a + x \ln b} = e^{x \ln a} e^{x \ln b} = a^x b^x$.

10. Let $\log_a x = r$ and $\log_a y = s$. Then $a^r = x$ and $a^s = y$.

(a) $xy = a^r a^s = a^{r+s} \quad \Rightarrow \quad \log_a(xy) = r + s = \log_a x + \log_a y$

(b) $\dfrac{x}{y} = \dfrac{a^r}{a^s} = a^{r-s} \quad \Rightarrow \quad \log_a \dfrac{x}{y} = r - s = \log_a x - \log_a y$

(c) $x^y = (a^r)^y = a^{ry} \quad \Rightarrow \quad \log_a(x^y) = ry = y \log_a x$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Cut here and keep for reference

CHAPTER 1 CONCEPT CHECK ANSWERS

1. (a) What is a function? What are its domain and range?

A function f is a rule that assigns to each element x in a set D exactly one element, called $f(x)$, in a set E. The domain is the set D and the range is the set of all possible values of $f(x)$ as x varies throughout the domain.

(b) What is the graph of a function?

The graph of a function f consists of all points (x, y) such that $y = f(x)$ and x is in the domain of f.

(c) How can you tell whether a given curve is the graph of a function?

Use the Vertical Line Test: a curve in the xy-plane is the graph of a function of x if and only if no vertical line intersects the curve more than once.

2. Discuss four ways of representing a function. Illustrate your discussion with examples.

A function can be represented verbally, numerically, visually, or algebraically. An example of each is given below.

Verbally: An assignment of students to chairs in a classroom (a description in words)

Numerically: A tax table that assigns an amount of tax to an income (a table of values)

Visually: A graphical history of the Dow Jones average (a graph)

Algebraically: A relationship between the area A and side length s of a square: $A = s^2$ (an explicit formula)

3. (a) What is an even function? How can you tell if a function is even by looking at its graph? Give three examples of an even function.

A function f is even if it satisfies $f(-x) = f(x)$ for every number x in its domain. If the graph of a function is symmetric with respect to the y-axis, then f is even. Examples are $f(x) = x^2$, $f(x) = \cos x$, $f(x) = |x|$.

(b) What is an odd function? How can you tell if a function is odd by looking at its graph? Give three examples of an odd function.

A function f is odd if it satisfies $f(-x) = -f(x)$ for every number x in its domain. If the graph of a function is symmetric with respect to the origin, then f is odd. Examples are $f(x) = x^3$, $f(x) = \sin x$, $f(x) = 1/x$.

4. What is an increasing function?

A function f is increasing on an interval I if $f(x_1) < f(x_2)$ whenever $x_1 < x_2$ in I.

5. What is a mathematical model?

A mathematical model is a mathematical description (often by means of a function or an equation) of a real-world phenomenon. (See the discussion on pages 23–24.)

6. Give an example of each type of function.

(a) Linear function: $f(x) = 2x + 1$, $f(x) = ax + b$

(b) Power function: $f(x) = x^2$, $f(x) = x^n$

(c) Exponential function: $f(x) = 2^x$, $f(x) = b^x$

(d) Quadratic function: $f(x) = x^2 + x + 1$, $f(x) = ax^2 + bx + c$

(e) Polynomial of degree 5: $f(x) = x^5 + 2x^4 - 3x^2 + 7$

(f) Rational function: $f(x) = \dfrac{x}{x+2}$, $f(x) = \dfrac{P(x)}{Q(x)}$

where $P(x)$ and $Q(x)$ are polynomials

7. Sketch by hand, on the same axes, the graphs of the following functions.

(a) $f(x) = x$ **(b)** $g(x) = x^2$

(c) $h(x) = x^3$ **(d)** $j(x) = x^4$

8. Draw, by hand, a rough sketch of the graph of each function.

(a) $y = \sin x$

(b) $y = \tan x$

(c) $y = e^x$

(d) $y = \ln x$

(continued)

(e) $y = 1/x$

(f) $y = |x|$

(g) $y = \sqrt{x}$

(h) $y = \tan^{-1}x$

9. Suppose that f has domain A and g has domain B.

(a) What is the domain of $f + g$?

The domain of $f + g$ is the intersection of the domain of f and the domain of g; that is, $A \cap B$.

(b) What is the domain of fg?

The domain of fg is also $A \cap B$.

(c) What is the domain of f/g?

The domain of f/g must exclude values of x that make g equal to 0; that is, $\{x \in A \cap B \mid g(x) \neq 0\}$.

10. How is the composite function $f \circ g$ defined? What is its domain?

The composition of f and g is defined by $(f \circ g)(x) = f(g(x))$. The domain is the set of all x in the domain of g such that $g(x)$ is in the domain of f.

11. Suppose the graph of f is given. Write an equation for each of the graphs that are obtained from the graph of f as follows.

(a) Shift 2 units upward: $y = f(x) + 2$

(b) Shift 2 units downward: $y = f(x) - 2$

(c) Shift 2 units to the right: $y = f(x - 2)$

(d) Shift 2 units to the left: $y = f(x + 2)$

(e) Reflect about the x-axis: $y = -f(x)$

(f) Reflect about the y-axis: $y = f(-x)$

(g) Stretch vertically by a factor of 2: $y = 2f(x)$

(h) Shrink vertically by a factor of 2: $y = \frac{1}{2}f(x)$

(i) Stretch horizontally by a factor of 2: $y = f\left(\frac{1}{2}x\right)$

(j) Shrink horizontally by a factor of 2 $y = f(2x)$

12. (a) What is a one-to-one function? How can you tell if a function is one-to-one by looking at its graph?

A function f is one-to-one if it never takes on the same value twice; that is $f(x_1) \neq f(x_2)$ whenever $x_1 \neq x_2$. When looking at a graph, use the Horizontal Line Test: a function is one-to-one if and only if no horizontal line intersects its graph more than once.

(b) If f is a one-to-one function, how is its inverse function f^{-1} defined? How do you obtain the graph of f^{-1} from the graph of f?

If f is a one-to-one function with domain A and range B, then its inverse function f^{-1} has domain B and range A and is defined by

$$f^{-1}(y) = x \iff f(x) = y$$

for any y in B.

The graph of f^{-1} is obtained by reflecting the graph of f about the line $y = x$.

13. (a) How is the inverse sine function $f(x) = \sin^{-1}x$ defined? What are its domain and range?

The inverse sine function $f(x) = \sin^{-1}x$ is defined as

$$\sin^{-1}x = y \iff \sin y = x$$

and $$-\pi/2 \leqslant y \leqslant \pi/2$$

Its domain is $-1 \leqslant x \leqslant 1$ and its range is $-\pi/2 \leqslant y \leqslant \pi/2$.

(b) How is the inverse cosine function $f(x) = \cos^{-1}x$ defined? What are its domain and range?

The inverse cosine function $f(x) = \cos^{-1}x$ is defined as

$$\cos^{-1}x = y \iff \cos y = x$$

and $$0 \leqslant y \leqslant \pi$$

Its domain is $-1 \leqslant x \leqslant 1$ and its range is $0 \leqslant y \leqslant \pi$.

(c) How is the inverse tangent function $f(x) = \tan^{-1}x$ defined? What are its domain and range?

The inverse tangent function $f(x) = \tan^{-1}x$ is defined as

$$\tan^{-1}x = y \iff \tan y = x$$

and $$-\pi/2 < y < \pi/2$$

Its domain is $\mathbb{R}$ and its range is $-\pi/2 < y < \pi/2$.

Cut here and keep for reference

CHAPTER 2 CONCEPT CHECK ANSWERS

1. Explain what each of the following means and illustrate with a sketch.

(a) $\lim_{x \to a} f(x) = L$ means that the values of $f(x)$ approach L as the values of x approach a (but $x \neq a$).

(b) $\lim_{x \to a^+} f(x) = L$ means that the values of $f(x)$ approach L as the values of x approach a through values greater than a.

(c) $\lim_{x \to a^-} f(x) = L$ means that the values of $f(x)$ approach L as the values of x approach a through values less than a.

(d) $\lim_{x \to a} f(x) = \infty$ means that the values of $f(x)$ can be made arbitrarily large by taking x sufficiently close to a (but not equal to a).

(e) $\lim_{x \to \infty} f(x) = L$ means that the values of $f(x)$ can be made arbitrarily close to L by requiring x to be sufficiently large.

2. Describe several ways in which a limit can fail to exist. Illustrate with sketches.

In general, the limit of a function fails to exist when the function values do not approach a fixed number. For each of the following functions, the limit fails to exist at $x = 2$.

The left and right limits are not equal.

There is an infinite discontinuity.

The function values oscillate between 1 and −1 infinitely often.

3. State the following Limit Laws.

(a) Sum Law

The limit of a sum is the sum of the limits:
$\lim_{x \to a} [f(x) + g(x)] = \lim_{x \to a} f(x) + \lim_{x \to a} g(x)$

(b) Difference Law

The limit of a difference is the difference of the limits:
$\lim_{x \to a} [f(x) - g(x)] = \lim_{x \to a} f(x) - \lim_{x \to a} g(x)$

(c) Constant Multiple Law

The limit of a constant times a function is the constant times the limit of the function: $\lim_{x \to a} [cf(x)] = c \lim_{x \to a} f(x)$

(d) Product Law

The limit of a product is the product of the limits:
$\lim_{x \to a} [f(x)g(x)] = \lim_{x \to a} f(x) \cdot \lim_{x \to a} g(x)$

(e) Quotient Law

The limit of a quotient is the quotient of the limits, provided that the limit of the denominator is not 0:

$$\lim_{x \to a} \frac{f(x)}{g(x)} = \frac{\lim_{x \to a} f(x)}{\lim_{x \to a} g(x)} \quad \text{if } \lim_{x \to a} g(x) \neq 0$$

(continued)

(f) Power Law

The limit of a power is the power of the limit:
$\lim_{x \to a} [f(x)]^n = \left[\lim_{x \to a} f(x)\right]^n$ (for n a positive integer)

(g) Root Law

The limit of a root is the root of the limit:
$\lim_{x \to a} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \to a} f(x)}$ (for n a positive integer)

4. What does the Squeeze Theorem say?

If $f(x) \leqslant g(x) \leqslant h(x)$ when x is near a (except possibly at a) and $\lim_{x \to a} f(x) = \lim_{x \to a} h(x) = L$, then $\lim_{x \to a} g(x) = L$. In other words, if $g(x)$ is squeezed between $f(x)$ and $h(x)$ near a, and if f and h have the same limit L at a, then g is forced to have the same limit L at a.

5. (a) What does it mean to say that the line $x = a$ is a vertical asymptote of the curve $y = f(x)$? Draw curves to illustrate the various possibilities.

It means that the limit of $f(x)$ as x approaches a from one or both sides is positive or negative infinity.

(b) What does it mean to say that the line $y = L$ is a horizontal asymptote of the curve $y = f(x)$? Draw curves to illustrate the various possibilities.

It means that $\lim_{x \to \infty} f(x) = L$ or $\lim_{x \to -\infty} f(x) = L$.

6. Which of the following curves have vertical asymptotes? Which have horizontal asymptotes?

(a) $y = x^4$: No asymptotes

(b) $y = \sin x$: No asymptotes

(c) $y = \tan x$: Vertical asymptotes $x = \pi/2 + \pi n$, n an integer

(d) $y = \tan^{-1} x$: Horizontal asymptotes $y = \pm\pi/2$

(e) $y = e^x$: Horizontal asymptote $y = 0$ $\left(\lim_{x \to -\infty} e^x = 0\right)$

(f) $y = \ln x$: Vertical asymptote $x = 0$ $\left(\lim_{x \to 0^+} \ln x = -\infty\right)$

(g) $y = 1/x$: Vertical asymptote $x = 0$, horizontal asymptote $y = 0$

(h) $y = \sqrt{x}$: No asymptotes

7. (a) What does it mean for f to be continuous at a?

A function f is continuous at a number a if the value of the function at $x = a$ is the same as the limit when x approaches a; that is, $\lim_{x \to a} f(x) = f(a)$.

(b) What does it mean for f to be continuous on the interval $(-\infty, \infty)$? What can you say about the graph of such a function?

A function f is continuous on the interval $(-\infty, \infty)$ if it is continuous at every real number a.

The graph of such a function has no hole or break in it.

8. (a) Give examples of functions that are continuous on $[-1, 1]$.

$f(x) = x^3 - x$, $g(x) = 2^x$, $y = \sin x$, $y = \tan x$, $y = 1/(x - 3)$, and $h(x) = |x|$ are all continuous on $[-1, 1]$.

(b) Give an example of a function that is not continuous on $[0, 1]$.

$$f(x) = \frac{1}{x - \frac{1}{2}} \qquad \left(f(x) \text{ is not defined at } x = \tfrac{1}{2}\right)$$

9. What does the Intermediate Value Theorem say?

If f is continuous on $[a, b]$ and N is any number between $f(a)$ and $f(b)$ $[f(a) \neq f(b)]$, Then there exists a number c in (a, b) such that $f(c) = N$. In other words, a continuous function takes on every intermediate value between the function values $f(a)$ and $f(b)$.

10. Write an expression for the slope of the tangent line to the curve $y = f(x)$ at the point $(a, f(a))$.

The slope of the tangent line is given by

$$\lim_{x \to a} \frac{f(x) - f(a)}{x - a} \quad \text{or} \quad \lim_{h \to 0} \frac{f(a + h) - f(a)}{h}$$

(continued)

Cut here and keep for reference

11. Suppose an object moves along a straight line with position $f(t)$ at time t. Write an expression for the instantaneous velocity of the object at time $t = a$. How can you interpret this velocity in terms of the graph of f?

The instantaneous velocity at time $t = a$ is

$$v(a) = \lim_{h \to 0} \frac{f(a + h) - f(a)}{h}$$

It is equal to the slope of the tangent line to the graph of f at the point $P(a, f(a))$.

12. If $y = f(x)$ and x changes from x_1 to x_2, write expressions for the following.

(a) The average rate of change of y with respect to x over the interval $[x_1, x_2]$:

$$\frac{\Delta y}{\Delta x} = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

(b) The instantaneous rate of change of y with respect to x at $x = x_1$:

$$\lim_{\Delta x \to 0} \frac{\Delta y}{\Delta x} = \lim_{x_2 \to x_1} \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

13. Define the derivative $f'(a)$. Discuss two ways of interpreting this number.

$$f'(a) = \lim_{h \to 0} \frac{f(a + h) - f(a)}{h}$$

or, equivalently,

$$f'(a) = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$$

The derivative $f'(a)$ is the instantaneous rate of change of $y = f(x)$ (with respect to x) when $x = a$ and also represents the slope of the tangent line to the graph of f at the point $P(a, f(a))$.

14. Define the second derivative of f. If $f(t)$ is the position function of a particle, how can you interpret the second derivative?

The second derivative of f, f'', is the derivative of f'.

If $f(t)$ is the position function of a particle, then the first derivative is velocity and the second derivative is the instantaneous rate of change of velocity, namely, acceleration.

15. **(a)** What does it mean for f to be differentiable at a?

f is differentiable at a if the derivative $f'(a)$ exists.

(b) What is the relation between the differentiability and continuity of a function?

If f is differentiable at a, then f is continuous at a.

(c) Sketch the graph of a function that is continuous but not differentiable at $a = 2$.

The graph of f changes direction abruptly at $x = 2$, so f has no tangent line there.

16. Describe several ways in which a function can fail to be differentiable. Illustrate with sketches.

A function is not differentiable at any value where the graph has a "corner," where the graph has a discontinuity, or where it has a vertical tangent line.

A corner

A discontinuity

A vertical tangent

Cut here and keep for reference

CHAPTER 3 CONCEPT CHECK ANSWERS

1. State each differentiation rule both in symbols and in words.

(a) The Power Rule

If n is any real number, then $\frac{d}{dx}(x^n) = nx^{n-1}$.

To find the derivative of a variable raised to a constant power, we multiply the expression by the exponent and then subtract one from the exponent.

(b) The Constant Multiple Rule

If c is a constant and f is a differentiable function, then

$$\frac{d}{dx}[cf(x)] = c\frac{d}{dx}f(x)$$

The derivative of a constant times a function is the constant times the derivative of the function.

(c) The Sum Rule

If f and g are both differentiable, then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)$$

The derivative of a sum of functions is the sum of the derivatives.

(d) The Difference Rule

If f and g are both differentiable, then

$$\frac{d}{dx}[f(x) - g(x)] = \frac{d}{dx}f(x) - \frac{d}{dx}g(x)$$

The derivative of a difference of functions is the difference of the derivatives.

(e) The Product Rule

If f and g are both differentiable, then

$$\frac{d}{dx}[f(x)g(x)] = f(x)\frac{d}{dx}[g(x)] + g(x)\frac{d}{dx}[f(x)]$$

The derivative of a product of two functions is the first function times the derivative of the second function plus the second function times the derivative of the first function.

(f) The Quotient Rule

If f and g are both differentiable, then

$$\frac{d}{dx}\left[\frac{f(x)}{g(x)}\right] = \frac{g(x)\frac{d}{dx}[f(x)] - f(x)\frac{d}{dx}[g(x)]}{[g(x)]^2}$$

The derivative of a quotient of functions is the denominator times the derivative of the numerator minus the numerator times the derivative of the denominator, all divided by the square of the denominator.

(g) The Chain Rule

If g is differentiable at x and f is differentiable at $g(x)$, then the composite function defined by $F(x) = f(g(x))$ is differentiable at x and F' is given by the product

$$F'(x) = f'(g(x))g'(x)$$

The derivative of a composite function is the derivative of the outer function evaluated at the inner function times the derivative of the inner function.

2. State the derivative of each function.

(a) $y = x^n$: $y' = nx^{n-1}$

(b) $y = e^x$: $y' = e^x$

(c) $y = b^x$: $y' = b^x \ln b$

(d) $y = \ln x$: $y' = 1/x$

(e) $y = \log_b x$: $y' = 1/(x \ln b)$

(f) $y = \sin x$: $y' = \cos x$

(g) $y = \cos x$: $y' = -\sin x$

(h) $y = \tan x$: $y' = \sec^2 x$

(i) $y = \csc x$: $y' = -\csc x \cot x$

(j) $y = \sec x$: $y' = \sec x \tan x$

(k) $y = \cot x$: $y' = -\csc^2 x$

(l) $y = \sin^{-1} x$: $y' = 1/\sqrt{1 - x^2}$

(m) $y = \cos^{-1} x$: $y' = -1/\sqrt{1 - x^2}$

(n) $y = \tan^{-1} x$: $y' = 1/(1 + x^2)$

(o) $y = \sinh x$: $y' = \cosh x$

(p) $y = \cosh x$: $y' = \sinh x$

(q) $y = \tanh x$: $y' = \operatorname{sech}^2 x$

(r) $y = \sinh^{-1} x$: $y' = 1/\sqrt{1 + x^2}$

(s) $y = \cosh^{-1} x$: $y' = 1/\sqrt{x^2 - 1}$

(t) $y = \tanh^{-1} x$: $y' = 1/(1 - x^2)$

3. (a) How is the number e defined?

e is the number such that $\lim_{h \to 0} \frac{e^h - 1}{h} = 1$.

(b) Express e as a limit.

$$e = \lim_{x \to 0}(1 + x)^{1/x}$$

(c) Why is the natural exponential function $y = e^x$ used more often in calculus than the other exponential functions $y = b^x$?

The differentiation formula for $y = b^x$ $[dy/dx = b^x \ln b]$ is simplest when $b = e$ because $\ln e = 1$.

(continued)

(d) Why is the natural logarithmic function $y = \ln x$ used more often in calculus than the other logarithmic functions $y = \log_b x$?

The differentiation formula for $y = \log_b x$ $[dy/dx = 1/(x \ln b)]$ is simplest when $b = e$ because $\ln e = 1$.

4. (a) Explain how implicit differentiation works.

Implicit differentiation consists of differentiating both sides of an equation with respect to x, treating y as a function of x. Then we solve the resulting equation for y'.

(b) Explain how logarithmic differentiation works.

Logarithmic differentiation consists of taking natural logarithms of both sides of an equation $y = f(x)$, simplifying using the Laws of Logarithms, differentiating implicitly with respect to x, and then solving the resulting equation for y'.

5. Give several examples of how the derivative can be interpreted as a rate of change in physics, chemistry, biology, economics, or other sciences.

In physics, interpretations of the derivative include velocity, linear density, electrical current, power (the rate of change of work), and the rate of radioactive decay. Chemists can use derivatives to measure reaction rates and the compressibility of a substance under pressure. In biology the derivative measures rates of population growth and blood flow. In economics, the derivative measures marginal cost (the rate of change of cost as more items are produced) and marginal profit. Other examples include the rate of heat flow in geology, the rate of performance improvement in psychology, and the rate at which a rumor spreads in sociology.

6. (a) Write a differential equation that expresses the law of natural growth.

If $y(t)$ is the value of a quantity y at time t, then

$$\frac{dy}{dt} = ky \qquad \text{where } k > 0 \text{ is a constant}$$

(b) Under what circumstances is this an appropriate model for population growth?

The equation in part (a) is an appropriate model when there is enough room and nutrition to support growth.

(c) What are the solutions of this equation?

If $y(0) = y_0$, then the solutions are $y(t) = y_0 e^{kt}$.

7. (a) Write an expression for the linearization of f at a.

$$L(x) = f(a) + f'(a)(x - a)$$

(b) If $y = f(x)$, write an expression for the differential dy.

$$dy = f'(x)\,dx$$

(c) If $dx = \Delta x$, draw a picture showing the geometric meanings of Δy and dy.

Cut here and keep for reference

CHAPTER 4 CONCEPT CHECK ANSWERS

1. Explain the difference between an absolute maximum and a local maximum. Illustrate with a sketch.

The function value $f(c)$ is the absolute maximum value of f if $f(c)$ is the largest function value on the entire domain of f, whereas $f(c)$ is a local maximum value if it is the largest function value when x is near c.

2. (a) What does the Extreme Value Theorem say?

If f is a continuous function on a closed interval $[a, b]$, then it always attains an absolute maximum and an absolute minimum value on that interval.

(b) Explain how the Closed Interval Method works.

To find the absolute maximum and minimum values of a continuous function f on a closed interval $[a, b]$, we follow these three steps:

- Find the critical numbers of f in the interval (a, b) and compute the values of f at these numbers.
- Find the values of f at the endpoints of the interval.
- The largest of the values from the previous two steps is the absolute maximum value; the smallest of these values is the absolute minimum value.

3. (a) State Fermat's Theorem.

If f has a local maximum or minimum at c, and if $f'(c)$ exists, then $f'(c) = 0$.

(b) Define a critical number of f.

A critical number of a function f is a number c in the domain of f such that either $f'(c) = 0$ or $f'(c)$ does not exist.

4. (a) State Rolle's Theorem.

Let f be a function that satisfies the following three hypotheses:

- f is continuous on the closed interval $[a, b]$.
- f is differentiable on the open interval (a, b).
- $f(a) = f(b)$

Then there is a number c in (a, b) such that $f'(c) = 0$.

(b) State the Mean Value Theorem and give a geometric interpretation.

If f is continuous on the interval $[a, b]$ and differentiable on (a, b), then there exists a number c between a and b such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

Geometrically, the theorem says that there is a point $P(c, f(c))$, where $a < c < b$, on the graph of f where the tangent line is parallel to the secant line that connects $(a, f(a))$ and $(b, f(b))$.

5. (a) State the Increasing/Decreasing Test.

If $f'(x) > 0$ on an interval, then f is increasing on that interval.

If $f'(x) < 0$ on an interval, then f is decreasing on that interval.

(b) What does it mean to say that f is concave upward on an interval I?

f is concave upward on an interval if the graph of f lies above all of its tangents on that interval.

(c) State the Concavity Test.

If $f''(x) > 0$ on an interval, then the graph of f is concave upward on that interval.

If $f''(x) < 0$ on an interval, then the graph of f is concave downward on that interval.

(d) What are inflection points? How do you find them?

Inflection points on the graph of a continuous function f are points where the curve changes from concave upward to concave downward or from concave downward to concave upward. They can be found by determining the values at which the second derivative changes sign.

6. (a) State the First Derivative Test.

Suppose that c is a critical number of a continuous function f.

- If f' changes from positive to negative at c, then f has a local maximum at c.
- If f' changes from negative to positive at c, then f has a local minimum at c.
- If f' is positive to the left and right of c, or negative to the left and right of c, then f has no local maximum or minimum at c.

(b) State the Second Derivative Test.

Suppose f'' is continuous near c.

- If $f'(c) = 0$ and $f''(c) > 0$, then f has a local minimum at c.
- If $f'(c) = 0$ and $f''(c) < 0$, then f has a local maximum at c.

(continued)

(c) What are the relative advantages and disadvantages of these tests?

The Second Derivative Test is sometimes easier to use, but it is inconclusive when $f''(c) = 0$ and fails if $f''(c)$ does not exist. In either case the First Derivative Test must be used.

7. (a) What does l'Hospital's Rule say?

L'Hospital's Rule says that *if* the limit of a quotient of functions is an indeterminate form of type $0/0$ or ∞/∞, then the limit is equal to the limit of the quotient of their derivatives:

$$\lim_{x \to a} \frac{f(x)}{g(x)} = \lim_{x \to a} \frac{f'(x)}{g'(x)}$$

(b) How can you use l'Hospital's Rule if you have a product $f(x)\,g(x)$ where $f(x) \to 0$ and $g(x) \to \infty$ as $x \to a$?

Write fg as $\dfrac{f}{1/g}$ or $\dfrac{g}{1/f}$ so that the limit becomes an indeterminate form of type $0/0$ or ∞/∞.

(c) How can you use l'Hospital's Rule if you have a difference $f(x) - g(x)$ where $f(x) \to \infty$ and $g(x) \to \infty$ as $x \to a$?

Convert the difference into a quotient by using a common denominator, rationalizing, factoring, or by some other method.

(d) How can you use l'Hospital's Rule if you have a power $[f(x)]^{g(x)}$ where $f(x) \to 0$ and $g(x) \to 0$ as $x \to a$?

By taking the natural logarithm of both sides of $y = [f(x)]^{g(x)}$, we can convert the power to a product by writing $\ln y = g(x) \ln f(x)$. Alternatively, we can write the function as an exponential: $[f(x)]^{g(x)} = e^{g(x) \ln f(x)}$.

8. State whether each of the following limit forms is indeterminate. Where possible, state the limit.

(a) $\dfrac{0}{0}$: indeterminate. L'Hospital's Rule can be applied to this form. Note that every derivative is a limit of this form.

(b) $\dfrac{\infty}{\infty}$: indeterminate. L'Hospital's Rule can be applied to this form.

(c) $\dfrac{0}{\infty}$: not indeterminate. A limit of this form has value 0.

(d) $\dfrac{\infty}{0}$: not indeterminate. A limit of this form could equal ∞, $-\infty$, or may not exist (but it cannot equal a finite value).

(e) $\infty + \infty$: not indeterminate. A limit of this form is equal to ∞.

(f) $\infty - \infty$: indeterminate

(g) $\infty \cdot \infty$: not indeterminate. A limit of this form is equal to ∞.

(h) $\infty \cdot 0$: indeterminate

(i) 0^0: indeterminate

(j) 0^∞: not indeterminate. A limit of this form has value 0.

(k) ∞^0: indeterminate

(l) 1^∞: indeterminate

9. If you have a graphing calculator or computer, why do you need calculus to graph a function?

Calculus reveals all the important aspects of a graph, such as local extreme values and inflection points, that can be missed when relying solely on technology. In many cases we can find exact locations of these key points rather than approximations. Using derivatives to identify the behavior of the graph also helps us choose an appropriate viewing window and alerts us to where we may wish to zoom in on a graph.

10. (a) Given an initial approximation x_1 to a root of the equation $f(x) = 0$, explain geometrically, with a diagram, how the second approximation x_2 in Newton's method is obtained.

We find the tangent line L to the graph of $y = f(x)$ at the point $(x_1, f(x_1))$. Then x_2 is the x-intercept of L.

(b) Write an expression for x_2 in terms of x_1, $f(x_1)$, and $f'(x_1)$.

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

(c) Write an expression for x_{n+1} in terms of x_n, $f(x_n)$, and $f'(x_n)$.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

(d) Under what circumstances is Newton's method likely to fail or to work very slowly?

Newton's method is likely to fail or to work very slowly when $f'(x_1)$ is close to 0. It also fails when $f'(x_i)$ is undefined.

11. (a) What is an antiderivative of a function f?

A function F is an antiderivative of f if $F'(x) = f(x)$.

(b) Suppose F_1 and F_2 are both antiderivatives of f on an interval I. How are F_1 and F_2 related?

They are identical or they differ by a constant.

Cut here and keep for reference

CHAPTER 5 CONCEPT CHECK ANSWERS

1. (a) Write an expression for a Riemann sum of a function f. Explain the meaning of the notation that you use.

A Riemann sum of f is

$$\sum_{i=1}^{n} f(x_i^*)\,\Delta x$$

where x_i^* is a point in the ith subinterval $[x_{i-1}, x_i]$ and Δx is the length of the subintervals.

(b) If $f(x) \geqslant 0$, what is the geometric interpretation of a Riemann sum? Illustrate with a diagram.

If f is positive, then a Riemann sum can be interpreted as the sum of areas of approximating rectangles, as shown in the figure.

(c) If $f(x)$ takes on both positive and negative values, what is the geometric interpretation of a Riemann sum? Illustrate with a diagram.

If f takes on both positive and negative values then the Riemann sum is the sum of the areas of the rectangles that lie above the x-axis and the negatives of the areas of the rectangles that lie below the x-axis (the areas of the blue rectangles minus the areas of the gray rectangles).

2. (a) Write the definition of the definite integral of a continuous function from a to b.

If f is a continuous function on the interval $[a, b]$, then we divide $[a, b]$ into n subintervals of equal width $\Delta x = (b - a)/n$. We let $x_0\ (= a), x_1, x_2, \ldots, x_n\ (= b)$ be the endpoints of these subintervals. Then

$$\int_a^b f(x)\,dx = \lim_{n\to\infty} \sum_{i=1}^{n} f(x_i^*)\,\Delta x$$

where x_i^* is any sample point in the ith subinterval $[x_{i-1}, x_i]$.

(b) What is the geometric interpretation of $\int_a^b f(x)\,dx$ if $f(x) \geqslant 0$?

If f is positive, then $\int_a^b f(x)\,dx$ can be interpreted as the area under the graph of $y = f(x)$ and above the x-axis for $a \leqslant x \leqslant b$.

(c) What is the geometric interpretation of $\int_a^b f(x)\,dx$ if $f(x)$ takes on both positive and negative values? Illustrate with a diagram.

In this case $\int_a^b f(x)\,dx$ can be interpreted as a "net area," that is, the area of the region above the x-axis and below the graph of f (labeled "+" in the figure) minus the area of the region below the x-axis and above the graph of f (labeled "−").

3. State the Midpoint Rule.

If f is a continuous function on the interval $[a, b]$ and we divide $[a, b]$ into n subintervals of equal width $\Delta x = (b - a)/n$, then

$$\int_a^b f(x)\,dx \approx \sum_{i=1}^{n} f(\bar{x}_i)\,\Delta x$$

where $\bar{x}_i$ = midpoint of $[x_{i-1}, x_i] = \frac{1}{2}(x_{i-1} + x_i)$.

4. State both parts of the Fundamental Theorem of Calculus.

Suppose f is continuous on $[a, b]$.

Part 1. If $g(x) = \int_a^x f(t)\,dt$, then $g'(x) = f(x)$.

Part 2. $\int_a^b f(x)\,dx = F(b) - F(a)$, where F is any antiderivative of f, that is, $F' = f$.

5. (a) State the Net Change Theorem.

The integral of a rate of change is the net change:

$$\int_a^b F'(x)\,dx = F(b) - F(a)$$

(b) If $r(t)$ is the rate at which water flows into a reservoir, what does $\int_{t_1}^{t_2} r(t)\,dt$ represent?

$\int_{t_1}^{t_2} r(t)\,dt$ represents the change in the amount of water in the reservoir between time t_1 and time t_2.

6. Suppose a particle moves back and forth along a straight line with velocity $v(t)$, measured in feet per second, and acceleration $a(t)$.

(a) What is the meaning of $\int_{60}^{120} v(t)\,dt$?

$\int_{60}^{120} v(t)\,dt$ represents the net change in position (the displacement) of the particle from $t = 60$ to $t = 120$ seconds, in other words, in the second minute.

(b) What is the meaning of $\int_{60}^{120} |v(t)|\,dt$?

$\int_{60}^{120} |v(t)|\,dt$ represents the total distance traveled by the particle in the second minute.

(continued)

(c) What is the meaning of $\int_{60}^{120} a(t)\,dt$?

$\int_{60}^{120} a(t)\,dt$ represents the change in velocity of the particle in the second minute.

7. (a) Explain the meaning of the indefinite integral $\int f(x)\,dx$.

The indefinite integral $\int f(x)\,dx$ is another name for an antiderivative of f, so $\int f(x)\,dx = F(x)$ means that $F'(x) = f(x)$.

(b) What is the connection between the definite integral $\int_a^b f(x)\,dx$ and the indefinite integral $\int f(x)\,dx$?

The connection is given by Part 2 of the Fundamental Theorem:

$$\int_a^b f(x)\,dx = \int f(x)\,dx\Big]_a^b$$

if f is continuous on $[a, b]$.

8. Explain exactly what is meant by the statement that "differentiation and integration are inverse processes."

Part 1 of the Fundamental Theorem of Calculus can be rewritten as

$$\frac{d}{dx}\int_a^x f(t)\,dt = f(x)$$

which says that if f is integrated and then the result is differentiated, we arrive back at the original function f.

Since $F'(x) = f(x)$, Part 2 of the theorem (or, equivalently, the Net Change Theorem) states that

$$\int_a^b F'(x)\,dx = F(b) - F(a)$$

This says that if we take a function F, first differentiate it, and then integrate the result, we arrive back at the original function, but in the form $F(b) - F(a)$.

Also, the indefinite integral $\int f(x)\,dx$ represents an antiderivative of f, so

$$\frac{d}{dx}\int f(x)\,dx = f(x)$$

9. State the Substitution Rule. In practice, how do you use it?

If $u = g(x)$ is a differentiable function and f is continuous on the range of g, then

$$\int f(g(x))\,g'(x)\,dx = \int f(u)\,du$$

In practice, we make the substitutions $u = g(x)$ and $du = g'(x)\,dx$ in the integrand to make the integral simpler to evaluate.

Cut here and keep for reference

CHAPTER 6 CONCEPT CHECK ANSWERS

1. (a) Draw two typical curves $y = f(x)$ and $y = g(x)$, where $f(x) \geqslant g(x)$ for $a \leqslant x \leqslant b$. Show how to approximate the area between these curves by a Riemann sum and sketch the corresponding approximating rectangles. Then write an expression for the exact area.

A Riemann sum that approximates the area between these curves is $\sum_{i=1}^{n} [f(x_i^*) - g(x_i^*)]\,\Delta x$. A sketch of the corresponding approximating rectangles:

An expression for the exact area is

$$\lim_{n\to\infty} \sum_{i=1}^{n} [f(x_i^*) - g(x_i^*)]\,\Delta x = \int_a^b [f(x) - g(x)]\,dx$$

(b) Explain how the situation changes if the curves have equations $x = f(y)$ and $x = g(y)$, where $f(y) \geqslant g(y)$ for $c \leqslant y \leqslant d$.

Instead of using "top minus bottom" and integrating from left to right, we use "right minus left" and integrate from bottom to top: $A = \int_c^d [f(y) - g(y)]\,dy$

2. Suppose that Sue runs faster than Kathy throughout a 1500-meter race. What is the physical meaning of the area between their velocity curves for the first minute of the race?

It represents the number of meters by which Sue is ahead of Kathy after 1 minute.

3. (a) Suppose S is a solid with known cross-sectional areas. Explain how to approximate the volume of S by a Riemann sum. Then write an expression for the exact volume.

We slice S into n "slabs" of equal width Δx. The volume of the ith slab is approximately $A(x_i^*)\,\Delta x$, where x_i^* is a sample point in the ith slab and $A(x_i^*)$ is the cross-sectional area of S at x_i^*. Then the volume of S is approximately $\sum_{i=1}^{n} A(x_i^*)\,\Delta x$ and the exact volume is

$$V = \lim_{n\to\infty} \sum_{i=1}^{n} A(x_i^*)\,\Delta x = \int_a^b A(x)\,dx$$

(b) If S is a solid of revolution, how do you find the cross-sectional areas?

If the cross-section is a disk, find the radius in terms of x or y and use $A = \pi(\text{radius})^2$. If the cross-section is a washer, find the inner radius r_{in} and outer radius r_{out} and use $A = \pi(r_{\text{out}}^2) - \pi(r_{\text{in}}^2)$.

4. (a) What is the volume of a cylindrical shell?

$$V = 2\pi rh\,\Delta r = (\text{circumference})(\text{height})(\text{thickness})$$

(b) Explain how to use cylindrical shells to find the volume of a solid of revolution.

We approximate the region to be revolved by rectangles, oriented so that revolution forms cylindrical shells rather than disks or washers. For a typical shell, find the circumference and height in terms of x or y and calculate

$$V = \int_a^b (\text{circumference})(\text{height})(dx \text{ or } dy)$$

(c) Why might you want to use the shell method instead of slicing?

Sometimes slicing produces washers or disks whose radii are difficult (or impossible) to find explicitly. On other occasions, the cylindrical shell method leads to an easier integral than slicing does.

5. Suppose that you push a book across a 6-meter-long table by exerting a force $f(x)$ at each point from $x = 0$ to $x = 6$. What does $\int_0^6 f(x)\,dx$ represent? If $f(x)$ is measured in newtons, what are the units for the integral?

$\int_0^6 f(x)\,dx$ represents the amount of work done. Its units are newton-meters, or joules.

6. (a) What is the average value of a function f on an interval $[a, b]$?

$$f_{\text{ave}} = \frac{1}{b-a}\int_a^b f(x)\,dx$$

(b) What does the Mean Value Theorem for Integrals say? What is its geometric interpretation?

If f is continuous on $[a, b]$, then there is a number c in $[a, b]$ at which the value of f is exactly equal to the average value of the function, that is, $f(c) = f_{\text{ave}}$. This means that for positive functions f, there is a number c such that the rectangle with base $[a, b]$ and height $f(c)$ has the same area as the region under the graph of f from a to b.

CHAPTER 7 CONCEPT CHECK ANSWERS

1. State the rule for integration by parts. In practice, how do you use it?

To integrate $\int f(x)g'(x)\,dx$, let $u = f(x)$ and $v = g(x)$. Then $\int u\,dv = uv - \int v\,du$.

In practice, try to choose $u = f(x)$ to be a function that becomes simpler when differentiated (or at least not more complicated) at long as $dv = g'(x)\,dx$ can be readily integrated to give v.

2. How do you evaluate $\int \sin^m x \cos^n x\,dx$ if m is odd? What if n is odd? What if m and n are both even?

If m is odd, use $\sin^2 x = 1 - \cos^2 x$ to write all sine factors except one in terms of cosine. Then substitute $u = \cos x$.

If n is odd, use $\cos^2 x = 1 - \sin^2 x$ to write all cosine factors except one in terms of sine. Then substitute $u = \sin x$.

If m and n are even, use the half-angle identities

$$\sin^2 x = \tfrac{1}{2}(1 - \cos 2x) \qquad \cos^2 x = \tfrac{1}{2}(1 + \cos 2x)$$

3. If the expression $\sqrt{a^2 - x^2}$ occurs in an integral, what substitution might you try? What if $\sqrt{a^2 + x^2}$ occurs? What if $\sqrt{x^2 - a^2}$ occurs?

If $\sqrt{a^2 - x^2}$ occurs, try $x = a \sin\theta$; if $\sqrt{a^2 + x^2}$ occurs, try $x = a \tan\theta$, and if $\sqrt{x^2 - a^2}$ occurs, try $x = a \sec\theta$.

4. What is the form of the partial fraction decomposition of a rational function $P(x)/Q(x)$ if the degree of P is less than the degree of Q and $Q(x)$ has only distinct linear factors? What if a linear factor is repeated? What if $Q(x)$ has an irreducible quadratic factor (not repeated)? What if the quadratic factor is repeated?

For distinct linear factors,

$$\frac{P(x)}{Q(x)} = \frac{A_1}{a_1x + b_1} + \frac{A_2}{a_2x + b_2} + \cdots + \frac{A_k}{a_kx + b_k}$$

If the linear factor $a_1x + b_1$ is repeated r times, then we must include all the terms

$$\frac{B_1}{a_1x + b_1} + \frac{B_2}{(a_1x + b_1)^2} + \cdots + \frac{B_r}{(a_1x + b_1)^r}$$

If $Q(x)$ has an irreducible quadratic factor (not repeated), then we include a term of the form

$$\frac{Ax + B}{ax^2 + bx + c}$$

If the irreducible quadratic factor is repeated r times, then we include all the terms

$$\frac{A_1x + B_1}{ax^2 + bx + c} + \frac{A_2x + B_2}{(ax^2 + bx + c)^2} + \cdots + \frac{A_rx + B_r}{(ax^2 + bx + c)^r}$$

5. State the rules for approximating the definite integral $\int_a^b f(x)\,dx$ with the Midpoint Rule, the Trapezoidal Rule, and Simpson's Rule. Which would you expect to give the best estimate? How do you approximate the error for each rule?

Let $a \leqslant x \leqslant b$, $I = \int_a^b f(x)\,dx$, and $\Delta x = (b - a)/n$.

Midpoint Rule:

$$I \approx M_n = \Delta x\,[f(\bar{x}_1) + f(\bar{x}_2) + \cdots + f(\bar{x}_n)]$$

where $\bar{x}_i$ is the midpoint of $[x_{i-1}, x_i]$.

Trapezoidal Rule:

$$I \approx T_n$$
$$= \frac{\Delta x}{2}[f(x_0) + 2f(x_1) + 2f(x_2) + \cdots + 2f(x_{n-1}) + f(x_n)]$$

where $x_i = a + i\,\Delta x$.

Simpson's Rule:

$$I \approx S_n$$
$$= \frac{\Delta x}{3}[f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \cdots + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)]$$

where n is even.

We would expect the best estimate to be given by Simpson's Rule.

Suppose $|f''(x)| \leqslant K$ and $|f^{(4)}(x)| \leqslant L$ for $a \leqslant x \leqslant b$. The errors in the Midpoint, Trapezoidal, and Simpson's Rules are given by, respectively,

$$|E_M| \leqslant \frac{K(b - a)^3}{24n^2} \qquad |E_T| \leqslant \frac{K(b - a)^3}{12n^2}$$

$$|E_S| \leqslant \frac{L(b - a)^5}{180n^4}$$

6. Define the following improper integrals.

(a) $\int_a^\infty f(x)\,dx = \lim_{t\to\infty} \int_a^t f(x)\,dx$

(b) $\int_{-\infty}^b f(x)\,dx = \lim_{t\to-\infty} \int_t^b f(x)\,dx$

(c) $\int_{-\infty}^\infty f(x)\,dx = \int_{-\infty}^a f(x)\,dx + \int_a^\infty f(x)\,dx$, where a is any real number (assuming that both integrals are convergent)

7. Define the improper integral $\int_a^b f(x)\,dx$ for each of the following cases.

(a) f has an infinite discontinuity at a.

If f is continuous on $(a, b]$, then

$$\int_a^b f(x)\,dx = \lim_{t\to a^+} \int_t^b f(x)\,dx$$

if this limit exists (as a finite number).

(b) f has an infinite discontinuity at b.

If f is continuous on $[a, b)$, then

$$\int_a^b f(x)\,dx = \lim_{t\to b^-} \int_a^t f(x)\,dx$$

if this limit exists (as a finite number).

(c) f has an infinite discontinuity at c, where $a < c < b$.

If both $\int_a^c f(x)\,dx$ and $\int_c^b f(x)\,dx$ are convergent, then

$$\int_a^b f(x)\,dx = \int_a^c f(x)\,dx + \int_c^b f(x)\,dx$$

8. State the Comparison Theorem for improper integrals.

Suppose that f and g are continuous functions with $f(x) \geqslant g(x) \geqslant 0$ for $x \geqslant a$.

(a) If $\int_a^\infty f(x)\,dx$ is convergent, then $\int_a^\infty g(x)\,dx$ is convergent.

(b) If $\int_a^\infty g(x)\,dx$ is divergent, then $\int_a^\infty f(x)\,dx$ is divergent.

Cut here and keep for reference

CHAPTER 8 CONCEPT CHECK ANSWERS

1. (a) How is the length of a curve defined?

We can approximate a curve C by a polygon with vertices P_i along C. The length L of C is defined to be the limit of the lengths of these inscribed polygons:

$$L = \lim_{n \to \infty} \sum_{i=1}^{n} |P_{i-1}P_i|$$

(b) Write an expression for the length of a smooth curve given by $y = f(x)$, $a \leqslant x \leqslant b$.

$$L = \int_a^b \sqrt{1 + [f'(x)]^2}\, dx$$

(c) What if x is given as a function of y?

If $x = g(y)$, $c \leqslant y \leqslant d$, then $L = \int_c^d \sqrt{1 + [g'(y)]^2}\, dy$.

2. (a) Write an expression for the surface area of the surface obtained by rotating the curve $y = f(x)$, $a \leqslant x \leqslant b$, about the x-axis.

$$S = \int_a^b 2\pi f(x)\sqrt{1 + [f'(x)]^2}\, dx$$

(b) What if x is given as a function of y?

If $x = g(y)$, $c \leqslant y \leqslant d$, then $S = \int_c^d 2\pi y\sqrt{1 + [g'(y)]^2}\, dy$.

(c) What if the curve is rotated about the y-axis?

$$S = \int_a^b 2\pi x\sqrt{1 + [f'(x)]^2}\, dx$$

or

$$S = \int_c^d 2\pi g(y)\sqrt{1 + [g'(y)]^2}\, dy$$

3. Describe how we can find the hydrostatic force against a vertical wall submersed in a fluid.

We divide the wall into horizontal strips of equal height Δx and approximate each by a rectangle with horizontal length $f(x_i)$ at depth x_i. If δ is the weight density of the fluid, then the hydrostatic force is

$$F = \lim_{n \to \infty} \sum_{i=1}^{n} \delta x_i f(x_i)\, \Delta x = \int_a^b \delta x f(x)\, dx$$

4. (a) What is the physical significance of the center of mass of a thin plate?

The center of mass is the point at which the plate balances horizontally.

(b) If the plate lies between $y = f(x)$ and $y = 0$, where $a \leqslant x \leqslant b$, write expressions for the coordinates of the center of mass.

$$\bar{x} = \frac{1}{A}\int_a^b x f(x)\, dx \qquad \text{and} \qquad \bar{y} = \frac{1}{A}\int_a^b \tfrac{1}{2}[f(x)]^2\, dx$$

where $A = \int_a^b f(x)\, dx$.

5. What does the Theorem of Pappus say?

If a plane region $\mathcal{R}$ that lies entirely on one side of a line ℓ in its plane is rotated about ℓ, then the volume of the resulting solid is the product of the area of $\mathcal{R}$ and the distance traveled by the centroid of $\mathcal{R}$.

6. Given a demand function $p(x)$, explain what is meant by the consumer surplus when the amount of a commodity currently available is X and the current selling price is P. Illustrate with a sketch.

The consumer surplus represents the amount of money saved by consumers in purchasing the commodity at price P [when they were willing to purchase it at price $p(x)$], corresponding to an amount demanded of X.

7. (a) What is cardiac output?

It is the volume of blood pumped by the heart per unit time, that is, the rate of flow into the aorta.

(b) Explain how the cardiac output can be measured by the dye dilution method.

An amount A of dye is injected into part of the heart and its concentration $c(t)$ leaving the heart is measured over a time interval $[0, T]$ until the dye has cleared. The cardiac output is given by $A/\int_0^T c(t)\, dt$.

8. What is a probability density function? What properties does such a function have?

Given a random variable X, its probability density function f is a function such that $\int_a^b f(x)\, dx$ gives the probability that X lies between a and b. The function f has the properties that $f(x) \geqslant 0$ for all x, and $\int_{-\infty}^{\infty} f(x)\, dx = 1$.

9. Suppose $f(x)$ is the probability density function for the weight of a female college student, where x is measured in pounds.

(a) What is the meaning of the integral $\int_0^{130} f(x)\, dx$?

It represents the probability that a randomly chosen female college student weighs less than 130 pounds.

(b) Write an expression for the mean of this density function.

$$\mu = \int_{-\infty}^{\infty} x f(x)\, dx = \int_0^{\infty} x f(x)\, dx$$

[since $f(x) = 0$ for $x < 0$]

(c) How can we find the median of this density function?

The median of f is the number m such that

$$\int_m^{\infty} f(x)\, dx = \frac{1}{2}$$

10. What is a normal distribution? What is the significance of the standard deviation?

A normal distribution corresponds to a random variable X that has a probability density function with a bell-shaped graph and equation given by

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}}\, e^{-(x-\mu)^2/(2\sigma^2)}$$

where μ is the mean and the positive constant σ is the standard deviation. σ measures how spread out the values of X are.

CHAPTER 9 CONCEPT CHECK ANSWERS

1. (a) What is a differential equation?

It is an equation that contains an unknown function and one or more of its derivatives.

(b) What is the order of a differential equation?

It is the order of the highest derivative that occurs in the equation.

(c) What is an initial condition?

It is a condition of the form $y(t_0) = y_0$.

2. What can you say about the solutions of the equation $y' = x^2 + y^2$ just by looking at the differential equation?

The equation tells us that the slope of a solution curve at any point (x, y) is $x^2 + y^2$. Note that $x^2 + y^2$ is always positive except at the origin, where $y' = x^2 + y^2 = 0$. Thus there is a horizontal tangent at $(0, 0)$ but nowhere else and the solution curves are increasing everywhere.

3. What is a direction field for the differential equation $y' = F(x, y)$?

A direction field (or slope field) for the differential equation $y' = F(x, y)$ is a two-dimensional graph consisting of short line segments with slope $F(x, y)$ at point (x, y).

4. Explain how Euler's method works.

Euler's method says to start at the point given by the initial value and proceed in the direction indicated by the direction field. Stop after a short time, look at the slope at the new location, and proceed in that direction. Keep stopping and changing direction according to the direction field until the approximation is complete.

5. What is a separable differential equation? How do you solve it?

It is a differential equation in which the expression for dy/dx can be factored as a function of x times a function of y, that is, $dy/dx = g(x)\,f(y)$. We can solve the equation by rewriting it as $[1/f(y)]\,dy = g(x)\,dx$, integrating both sides, and solving for y.

6. What is a first-order linear differential equation? How do you solve it?

A first-order linear differential equation is a differential equation that can be put in the form

$$\frac{dy}{dx} + P(x)y = Q(x)$$

where P and Q are continuous functions on a given interval. To solve such an equation, we multiply both sides by the integrating factor $I(x) = e^{\int P(x)\,dx}$ to put it in the form $\big(I(x)\,y\big)' = I(x)\,Q(x)$. We then integrate both sides and solve for y.

7. (a) Write a differential equation that expresses the law of natural growth. What does it say in terms of relative growth rate?

If $P(t)$ is the value of a quantity y at time t and if the rate of change of P with respect to t is proportional to its size $P(t)$ at any time, then $\dfrac{dP}{dt} = kP$.

In this case the relative growth rate, $\dfrac{1}{P}\dfrac{dP}{dt}$, is constant.

(b) Under what circumstances is this an appropriate model for population growth?

It is an appropriate model under ideal conditions: unlimited environment, adequate nutrition, absence of predators and disease.

(c) What are the solutions of this equation?

If $P(0) = P_0$, the initial value, then the solutions are $P(t) = P_0 e^{kt}$.

8. (a) Write the logistic differential equation.

The logistic differential equation is

$$\frac{dP}{dt} = kP\left(1 - \frac{P}{M}\right)$$

where M is the carrying capacity.

(b) Under what circumstances is this an appropriate model for population growth?

It is an appropriate model for population growth if the population grows at a rate proportional to the size of the population in the beginning, but eventually levels off and approaches its carrying capacity because of limited resources.

9. (a) Write Lotka-Volterra equations to model populations of food-fish (F) and sharks (S).

$$\frac{dF}{dt} = kF - aFS \qquad \text{and} \qquad \frac{dS}{dt} = -rS + bFS$$

(b) What do these equations say about each population in the absence of the other?

In the absence of sharks, an ample food supply would support exponential growth of the fish population, that is, $dF/dt = kF$, where k is a positive constant. In the absence of fish, we assume that the shark population would decline at a rate proportional to itself, that is $dS/dt = -rS$, where r is a positive constant.

Cut here and keep for reference

CHAPTER 10 CONCEPT CHECK ANSWERS

1. (a) What is a parametric curve?

A parametric curve is a set of points of the form $(x, y) = (f(t), g(t))$, where f and g are functions of a variable t, the parameter.

(b) How do you sketch a parametric curve?

Sketching a parametric curve, like sketching the graph of a function, is difficult to do in general. We can plot points on the curve by finding $f(t)$ and $g(t)$ for various values of t, either by hand or with a calculator or computer. Sometimes, when f and g are given by formulas, we can eliminate t from the equations $x = f(t)$ and $y = g(t)$ to get a Cartesian equation relating x and y. It may be easier to graph that equation than to work with the original formulas for x and y in terms of t.

2. (a) How do you find the slope of a tangent to a parametric curve?

You can find dy/dx as a function of t by calculating

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} \qquad \text{if } dx/dt \neq 0$$

(b) How do you find the area under a parametric curve?

If the curve is traced out once by the parametric equations $x = f(t)$, $y = g(t)$, $\alpha \leqslant t \leqslant \beta$, then the area is

$$A = \int_a^b y\,dx = \int_\alpha^\beta g(t)f'(t)\,dt$$

[or $\int_\beta^\alpha g(t)f'(t)\,dt$ if the leftmost point is $(f(\beta), g(\beta))$ rather than $(f(\alpha), g(\alpha))$].

3. Write an expression for each of the following:

(a) The length of a parametric curve

If the curve is traced out once by the parametric equations $x = f(t)$, $y = g(t)$, $\alpha \leqslant t \leqslant \beta$, then the length is

$$L = \int_\alpha^\beta \sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt$$

$$= \int_\alpha^\beta \sqrt{[f'(t)]^2 + [g'(t)]^2}\,dt$$

(b) The area of the surface obtained by rotating a parametric curve about the x-axis

$$S = \int_\alpha^\beta 2\pi y\sqrt{(dx/dt)^2 + (dy/dt)^2}\,dt$$

$$= \int_\alpha^\beta 2\pi g(t)\sqrt{[f'(t)]^2 + [g'(t)]^2}\,dt$$

(c) The speed of a particle traveling along a parametric curve

$$v(t) = s'(t) = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$$

4. (a) Use a diagram to explain the meaning of the polar coordinates (r, θ) of a point.

(b) Write equations that express the Cartesian coordinates (x, y) of a point in terms of the polar coordinates.

$$x = r\cos\theta \qquad y = r\sin\theta$$

(c) What equations would you use to find the polar coordinates of a point if you knew the Cartesian coordinates?

To find a polar representation (r, θ) with $r \geqslant 0$ and $0 \leqslant \theta < 2\pi$, first calculate $r = \sqrt{x^2 + y^2}$. Then θ is specified by $\tan\theta = y/x$. Be sure to choose θ so that (r, θ) lies in the correct quadrant.

5. (a) How do you find the area of a region bounded by a polar curve?

$$A = \int_a^b \tfrac{1}{2}r^2\,d\theta = \int_a^b \tfrac{1}{2}[f(\theta)]^2\,d\theta$$

(b) How do you find the length of a polar curve?

$$L = \int_a^b \sqrt{(dx/d\theta)^2 + (dy/d\theta)^2}\,d\theta$$

$$= \int_a^b \sqrt{r^2 + (dr/d\theta)^2}\,d\theta$$

$$= \int_a^b \sqrt{[f(\theta)]^2 + [f'(\theta)]^2}\,d\theta$$

(c) How do you find the slope of a tangent line to a polar curve?

$$\frac{dy}{dx} = \frac{\dfrac{dy}{d\theta}}{\dfrac{dx}{d\theta}} = \frac{\dfrac{d}{d\theta}(y)}{\dfrac{d}{d\theta}(x)} = \frac{\dfrac{d}{d\theta}(r\sin\theta)}{\dfrac{d}{d\theta}(r\cos\theta)}$$

$$= \frac{\left(\dfrac{dr}{d\theta}\right)\sin\theta + r\cos\theta}{\left(\dfrac{dr}{d\theta}\right)\cos\theta - r\sin\theta} \qquad \text{where } r = f(\theta)$$

6. (a) Give a geometric definition of a parabola.

A parabola is a set of points in a plane whose distances from a fixed point F (the focus) and a fixed line l (the directrix) are equal.

(b) Write an equation of a parabola with focus $(0, p)$ and directrix $y = -p$. What if the focus is $(p, 0)$ and the directrix is $x = -p$?

In the first case an equation is $x^2 = 4py$ and in the second case, $y^2 = 4px$.

(continued)

7. (a) Give a definition of an ellipse in terms of foci.

An ellipse is a set of points in a plane the sum of whose distances from two fixed points (the foci) is a constant.

(b) Write an equation for the ellipse with foci $(\pm c, 0)$ and vertices $(\pm a, 0)$.

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

where $a \geqslant b > 0$ and $c^2 = a^2 - b^2$.

8. (a) Give a definition of a hyperbola in terms of foci.

A hyperbola is a set of points in a plane the difference of whose distances from two fixed points (the foci) is a constant. This difference should be interpreted as the larger distance minus the smaller distance.

(b) Write an equation for the hyperbola with foci $(\pm c, 0)$ and vertices $(\pm a, 0)$.

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

where $c^2 = a^2 + b^2$.

(c) Write equations for the asymptotes of the hyperbola in part (b).

$$y = \pm \frac{b}{a} x$$

9. (a) What is the eccentricity of a conic section?

If a conic section has focus F and corresponding directrix l, then the eccentricity e is the fixed ratio $|PF|/|Pl|$ for points P of the conic section.

(b) What can you say about the eccentricity if the conic section is an ellipse? A hyperbola? A parabola?

$e < 1$ for an ellipse; $e > 1$ for a hyperbola; $e = 1$ for a parabola

(c) Write a polar equation for a conic section with eccentricity e and directrix $x = d$. What if the directrix is $x = -d$? $y = d$? $y = -d$?

$$\text{directrix } x = d: \quad r = \frac{ed}{1 + e\cos\theta}$$

$$x = -d: \quad r = \frac{ed}{1 - e\cos\theta}$$

$$y = d: \quad r = \frac{ed}{1 + e\sin\theta}$$

$$y = -d: \quad r = \frac{ed}{1 - e\sin\theta}$$