bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Methane Cycling Microbes are Important Predictors of Methylmercury Accumulation in Rice Paddies

Rui Zhang[a], Alexandre J Poulain[a], Qiang Pu[b], Jiang Liu[b*], Mahmoud A. Abdelhafiz[b**], Xinbin Feng[b], Bo Meng[b#], Daniel S Grégoire[c#]

[a] Department of Biology, University of Ottawa, Ottawa, Canada

[b] State Key Laboratory of Environmental Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, Guiyang, China

[c] Department of Chemistry, Carleton University, Ottawa, Canada

Running Head: Methane Cycling Microbes and MeHg Accumulation

#Address correspondence to Daniel S Grégoire, danielgregoire@cunet.carleton.ca and Bo Meng: mengbo@vip.skleg.cn.

*Present address: College of Resources, Sichuan Agricultural University, Chengdu, China

**Present address: Geology Department, Faculty of Science, Al-Azhar University, Assiut, Egypt

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Abstract

Microbial production of methylmercury from inorganic mercury in rice paddies poses health risks to consumers of this essential dietary staple. Although mercury-methylating communities are well characterized, the microbial guilds contributing to methylmercury accumulation in rice paddies remain unclear. Here, we collected paddy soils across a mercury concentration gradient throughout the rice growing season to identify microbial and environmental factors influencing methylmercury dynamics. We show that *hgcA* gene abundance, the key gene required for methylation, was not a significant predictor of methylmercury concentration in paddy soils. We also show that *merB* gene abundance correlated with methylmercury in mercury-polluted rhizosphere samples. Methane cycling genes were actively expressed, and their beta-diversity was significantly associated with methylmercury levels. Methanogen abundance correlated with higher methylmercury under elevated total mercury concentrations. Analysis of the methanotroph-associated *mbnT* gene, implicated in demethylation, revealed an unexpected positive correlation with methylmercury. Multiple regression and machine learning models converged on mercury bioavailability and methanogen/methanotroph abundances as key predictors of methylmercury, with methanogen-associated *hgcA* gene abundance and methanogen-methanotroph interactions highlighted under flooded, low-redox conditions. These findings suggest that methane-cycling microbes play key roles in methylmercury cycling dynamics and point to management strategies that could simultaneously mitigate mercury pollution and greenhouse gas emissions.

Importance

Methylmercury is a microbially-derived neurotoxin that accumulates in rice, which is a global food staple. Predicting mercury's fate in rice paddies is challenging because of the interplay between microbes responsible for methylmercury cycling and variables that control mercury availability. Our study coupled genomic and geochemical measurements with machine learning to identify the key predictors of methylmercury accumulation in paddy soils. We demonstrate that methanogen and methanotroph abundance, and mercury bioavailability, are major predictors of methylmercury variability in paddies. We show that considering interactions between methane cycling guilds improves our capacity to predict methylmercury accumulation in soils compared to approaches that rely solely on mercury cycling genes. This work can inform remediation strategies for mercury in rice paddies but also wetlands and permafrost where methane and mercury cycling are tightly coupled. Such strategies could provide a solution to simultaneously mitigate methylmercury exposure and reduce greenhouse gas emissions amid global environmental change.

Keywords: methanogen, methanotroph, mercury, methylmercury, methylation, demethylation, metagenomics

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Introduction

Mercury (Hg) is a metal primarily emitted as gaseous elemental Hg (Hg(0)) from natural and anthropogenic sources and deposits in ecosystems worldwide, where it undergoes oxidation to Hg(II) and subsequent microbial conversion to methylmercury (MeHg), a potent neurotoxin (1). MeHg accumulates in rice, a global food staple, with particularly elevated concentrations in paddies near mining, smelting, and industrial areas with elevated Hg(0) emissions (2). The global atmospheric transportation of Hg could also affect rice paddies away from pollution sources (3). Since paddy soil represents a major source of MeHg in rice grains (4), predicting MeHg accumulation in rice paddies is critical for mitigating Hg exposure through diet.

Net MeHg accumulation in rice paddies represents an interplay between methylation and demethylation processes (5). Hg methylation is facilitated by anaerobic microbes harboring the *hgcAB* gene cluster (6–8), while demethylation occurs via the Hg resistance (*mer*) operon–dependent pathways (e.g., *merB*/*merA*) and *mer*-independent microbial pathways (5). These processes are also connectively influenced by environmental variables such as Hg bioavailability, redox, dissolved organic matter (DOM), sulfide (5,9–12) and other microbial players indirectly (13).

While Hg methylating communities have been extensively characterized in rice paddies and other ecosystems (14–19), MeHg poorly correlates with Hg-cycling genes (e.g., *hgcA*, *merA*, *merB*) and *hgcA* transcripts (20). This disconnect highlights the need for integrated approaches that consider multiple variables simultaneously when predicting what controls MeHg accumulation in rice.

Recent evidence suggests that methane-cycling microbes play underappreciated roles in mediating MeHg dynamics. In rice paddies near Hg-mining areas, methanogens emerged as critical players controlling MeHg levels (14,15,21), potentially contributing to both methylation and demethylation, simultaneously (14,22,23). Furthermore, methanotrophs were experimentally shown to degrade MeHg at picomolar to nanomolar concentrations under near-neutral pH conditions typical of rice paddies, using a methanobactin-mediated mechanism distinct from the conventional *mer*-dependent pathway (24,25). However, the ecosystem-level significance of methanotrophic MeHg demethylation and its interaction with methanogenic processes in determining net MeHg accumulation has not been evaluated in rice paddy systems.

To address these critical gaps, we aim to (i) identify and rank the relative importance of microbial communities and environmental variables relevant to MeHg turnover as predictors of MeHg accumulation patterns in mining-impacted rice paddies; (ii) characterize the abundance and distribution of methane-cycling microorganisms in relation to MeHg concentrations; and (iii) develop predictive models to inform future mechanistic studies on Hg cycling.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

By integrating metagenomic, geochemical, and statistical approaches, we show that methanogen and methanotroph abundances are the leading biological predictor, while Hg bioavailability is the primary environmental determinant of MeHg variability. Machine learning analyses further highlighted the importance of methanogen methylator abundance and a methanogen–methanotroph interaction, particularly under flooded conditions where model performance improved. Together, these findings provide environmental-scale data that guide hypothesis development for broader geographical testing, and inform strategies that may jointly mitigate greenhouse gas emissions and Hg exposure.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Materials and Methods

Soil collection, geochemical analyses, and sequencing

Rice paddy soil samples were collected from three sites in Guizhou province, Southwest China: Huaxi (HX, background site), Gouxi (GX, artisanal Hg smelting area), and Sikeng (SK, abandoned Hg mining area). Sampling occurred from May 2021 to April 2022, spanning pre-plantation through post-harvest periods. Surface and rhizosphere soils were collected and processed as follows: individual replicate samples were analyzed separately for geochemical parameters, while replicate samples were combined into composite samples for DNA extraction and subsequent metagenomic analyses. Soil geochemical parameters were analyzed, including Hg species (THg by Cold Vapor Atomic Fluorescence Spectrometry (CVAFS), MeHg by gas chromatography-CVAFS following USEPA methods, and bio-available fraction Hg by the $Mg(NO_3)_2$ extraction method, anions (by ion chromatography) (26–28), elemental sulfur (by high-performance liquid chromatography), dissolved organic carbon, soil organic matter, and redox-sensitive species (S^{2-}, Fe^{2+}, Fe^{3+}). Recovery rates of the standard reference materials for THg and MeHg analyses were provided in Supplementary Data 1. DNA and RNA extractions were performed using the PowerSoil DNA Isolation Kit and the RNeasy PowerSoil Total RNA Kit (QIAGEN) following the manufacturer's instructions. Both metagenomic and metatranscriptomic sequencing were performed on an Illumina NovaSeq 6000 platform, producing 150 bp paired-end reads. Sequencing was conducted at Azenta (Suzhou, China). The sequencing aimed for a read depth of approximately 15 Gbp per sample (range: 10.9 – 20 Gbp). Quality metrics are provided in Supplementary Data 2. In total, 50 metagenomic and 6 metatranscriptomic samples were obtained. Details on sample collection, geochemical analyses and sequencing are provided in Supplementary Method 1.

Genome assembly and binning

Metagenomic data were processed using fastp and FastQC for quality control (Supplementary Data 3) (29,30). Individual assemblies were performed with MetaSPAdes, while site-specific co-assemblies used MEGAHIT (31). Contigs>2000 bp were retained and mapped with BWA-MEM to generate coverage data. Key assembly metrics, such as N50, total contigs, length distribution and mapping rates are provided in Supplementary Data 4. Four binning algorithms (MetaBAT 2, MaxBin 2, CONCOCT, and VAMB) were applied (32–35), yielding 5307 initial bins. Bin quality was assessed with CheckM2 (36), and DAS-Tool (37) was used to select the highest quality bins, followed by dereplication with dRep (38) (ANI threshold 98%). We retained 267 metagenome-assembled genomes (MAGs) with>50% completeness and<10% redundancy following manual refinement in Anvi'o (39). Taxonomic classification was performed using GTDB-Tk (40) based on GTDB (release 214) (Supplementary Data 5()) . The co-assembly approach was chosen over individual assembly due to higher MAG yield and

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

greater taxonomic diversity. Details on genome assembly, binning and software versions are provided in Supplementary Method 2.

Metatranscriptomic sequence processing

The RNA sequences were trimmed and filtered to a minimum mean quality score of 30 using fastp (29). Forward and reverse sequences were overlapped with a minimum and maximum of 10 and 150 bp, respectively, using FLASH (41). Sequences were subsequently sorted into SSU ribosomal RNA (rRNA), LSU rRNA, and non-rRNA using SortMeRNA (42) against the default reference database (smr_v4.3_default_db). The non-rRNA fraction was mapped to the ORFs identified on the contigs from the corresponding sample and linked with featureCounts results to determine the transcript abundance of *mcrA*, *pmoA*, *mmoX*, *hgcA*, and *merB*. Taxon-specific transcript abundance was calculated by dividing the RNA reads mapped to transcripts by the total RNA reads in the sample, normalized to HMM length.

Functional gene abundance analyses

For microbial abundance estimation of various functional guilds, including methanogens, methanotrophs, Hg-methylating microbes, and demethylating microbes, we employed a gene-centric approach at the contig level using individual assemblies. We selected individual assemblies over co-assembly to preserve sample-specific information and enable more accurate quantification of functional genes within each sample. While co-assembly is valuable for generating high-quality MAGs by aggregating data across samples, it inherently homogenizes information from multiple samples, potentially obscuring subtle variations in gene abundance at the individual sample level.

Open reading frames (ORFs) were identified using Prodigal (43), short reads were mapped using BWA MEM (44), and reads mapped to ORFs were quantified using featureCounts from Subread package (45). Functional genes were retrieved using Hidden Markov Models (HMMs) through hmmsearch of HMMER software (46). Specifically, the McrA HMM (PF02249) was sourced from PFAM, while PmoA and MmoX HMMs were obtained from the GraftM gene packages (https://data.ace.uq.edu.au/public/graftm/7) with an E-value threshold of 1×10^{-5}. We conducted protein homology searches using BLASTp against the NCBI non-redundant database (as of Dec 5, 2023) to confirm their identity and refine search results. Queries consisted of HMM hits of the corresponding genes, each limited to the single best hit (-max_target_seqs 1) to focus on the most relevant homolog. McrA, PmoA, and MmoX hits were filtered based on sequence titles indicative of their respective functions (e.g., McrA hits containing "methyl coenzyme M reductase"; PmoA and MmoX hits classified as "methane monooxygenase"). Extra precautions were taken with methane-cycling genes, as PmoA sequence is homologous to other alkane monooxygenases, and it shares homology with the

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

AmoA of ammonia oxidizers. The highest similarity to a known methanotrophic isolate was prioritized.

Methanobactin biosynthesis and transport genes from references genomes, MAGs and metagenomes were identified using HMMs listed from a previous study with predefined threshold values (47).

Hg-cycling genes required specialized validation procedures. *hgcAB* genes were identified using Hg-MATE-Db HMMs (ver. 1.01142021) (48) and filtered for conserved motifs [N(V/I)WCA(A/G)GK] and [C(M/I)ECGA] (6). MerB sequences were retrieved using pre-compiled HMMs (49) and validated for characteristic cysteine and aspartic acid signatures (50).

For additional functional guilds, IRB detection employed HMM profiles from FeGenie (51) to identify genes related to iron reduction, with gene-specific domain E-value cutoffs as defined in the database configuration files. Abundance for IRB was calculated by summing coverage counts of all identified genes and normalizing against total coding sequence (CDS) counts in each metagenome. For SRB quantification, we targeted the *dsrA* gene using the TIGR02064.1 HMM profile with a domain E-value cutoff of 392.9.

Taxonomic classification was conducted using MMseqs2 (52) against GTDB (release 214) (53) with lowest common ancestor strategy. To facilitate cross-sample comparison, we converted taxon-specific coverage values to the percentage of reads normalized to HMM lengths. Total reads numbers in each sample were referred to effective library size values calculated using the edgeR package (54). Details on functional gene analyses and software versions are provided in Supplementary Method 3.

$$Normalized\ Taxon\ Coverage = \frac{Taxon\ Coverage}{(Total\ reads\ number * HMM\ length)}$$

Complementary metagenomic approaches

Our study employed two complementary metagenomic strategies. Gene-centric analysis quantifies functional potential across all community members, capturing the full breadth of metabolic capacities including genes from low-abundance taxa that may not assemble into complete genomes. Genome-resolved analysis using MAGs provides taxonomic context, linking functional genes to specific organisms and enabling network analyses of population-level responses to environmental changes. Together, these approaches address different scales: community-wide functional potential (gene-centric) and organism-specific contributions (genome-resolved), both essential for understanding Hg cycling processes.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Statistical analyses

Principal Component Analysis (PCA) of geochemical variables was performed with standardized data, and distance-based redundancy analysis (db-RDA) was used to assess relationships between methane-cycling communities and environmental factors based on Bray-Curtis dissimilarity. Permutational multivariate ANOVA (PERMANOVA) evaluated effects of site and soil compartment on community composition (i.e., beta-diversity), with pairwise comparisons using Bonferroni correction. Multiple linear regression with AICc model selection was employed to evaluate the relative importance of methylation potential (*hgcA* abundance) and Hg bioavailability (F1-Hg) in predicting MeHg concentrations. Seven candidate models representing different hypotheses were tested, and model assumptions were verified through standard diagnostic tests.

MAG relative abundances were calculated by aligning reads to dereplicated genome sets using CoverM (v0.7.0) (55). Time-dependent correlations between MAG abundance and MeHg concentrations were identified using extended Local Similarity Analysis (eLSA) across 11 time points for each site (Supplementary Data 6) (56,57). False Discovery Rate correction was applied using the Benjamini-Hochberg procedure, with significant associations defined as FDR-adjusted $p<0.05$. Random forest regression models were developed to predict MeHg concentrations from 12 predictor variables (9 microbial functional guilds and 3 environmental parameters) using 100-fold cross-validation with 70:30 train-test splits. Model performance was evaluated using RMSE, MAE, R^2, and Pearson correlation coefficients. All analyses were conducted in R (v4.3.0) with statistical significance set at $\alpha=0.05$. Detailed methodological parameters are provided in Supplementary Method 4.

Code and data availability

The code used for the bioinformatics pipeline is available at https://github.com/rzhan186/gy2021_bioinformatics. Supplementary Methods, Discussions, and Figures are provided in the Supplementary Materials. Supplementary Data and Results can be accessed at https://doi.org/10.6084/m9.figshare.29847869.v3. Raw sequencing data have been deposited in the NCBI Sequence Read Archive under BioProject ID PRJNA1179437.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Results and Discussion

Spatial and temporal dynamics of MeHg concentration across the rice paddies




Figure 1. Hg dynamics in sampled rice paddies over the growing season. (A) Total mercury (THg), (B) methylmercury (MeHg), and (C) water-soluble Hg fraction (F1) in surface soils (left panels) and rhizosphere soils (right panels). Data points represent individual measurements from replicate soil samples collected at three sites (HX, GX, and SK), with smoothed trend lines (±95% confidence interval). Note that surface soils were sampled throughout the entire study period (including post-harvest), while rhizosphere soils were only collected during the rice growing season when plants were present, resulting in different x-axis scales. THg is shown in mg/kg, while MeHg and F1-fraction are in μg/kg. Day 0 corresponds to the beginning of the sampling period.

THg and MeHg levels varied significantly across rice paddies (Fig. 1, Fig. S3). Average THg concentrations across the entire sampling period were significantly elevated at the mining-impacted sites SK (42.03±21.5 mg/kg) and GX (21.41±4.04 mg/kg), compared to the background site HX (0.17±0.02 mg/kg). However, average MeHg concentrations over the sampling period were significantly higher at GX (3.44±0.70 μg/kg), followed by SK (2.42±0.76 μg/kg) and HX (0.78±0.21 μg/kg) (Fig. S3). This supports prior studies indicating that THg is a poor predictor of the net MeHg production (58).

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Distinct temporal trends were observed for MeHg at the mining-impacted sites, which exhibited different patterns compared to the relatively stable levels of THg (Fig. 1A). At GX and SK, MeHg peaked at cultivation onset (day 0) and declined throughout the rice-growing season (day 0-80), with further reductions observed post-harvest after drainage. MeHg concentrations were typically higher during the water-saturated season (day 0-100) than in the post-harvest period (day 100-260). A notable increase in MeHg occurred at GX on day 290, coinciding with pre-planting and re-drainage for the next growing season. In contrast, at HX, MeHg levels remained relatively stable overall. Throughout the three sampling sites, no significant differences in MeHg were detected between surface soil and rhizosphere (Mann-Whitney U test, $p>0.05$). This pattern of fluctuating MeHg in mining-affected sites during cultivation and stability at the background site is previously undocumented, due to the lack of extensive longitudinal sampling. Subsequent sections consider the potential microbial and geochemical drivers contributing to these trends in Hg concentrations.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Geochemical factors partially explain site-specific MeHg patterns

Geochemical factors were assessed to contextualize their potential impacts on MeHg accumulation (please refer to Fig. S1, Supplementary Result 1 throughout this section). Redox potential measured in overlying water revealed that HX is more oxidizing (109.29±19.12 mV) compared to GX (57.41±37.09 mV) and SK (60.96±20.3 mV). Regardless of soil compartments, DOM concentrations differed significantly between sites, with HX showing the highest levels (0.72±0.31 mg/g). MeHg correlated positively with DOM in surface soil at GX (Spearman's ρ=0.33) and SK (ρ=0.58), while correlating negatively with DOM $SUVA_{254}$ in SK surface (ρ=-0.56) and rhizosphere (ρ=-0.60) (Fig. S2). The bioavailable fraction of Hg (F1-Hg) exhibited temporal fluctuations that partially paralleled MeHg changes, primarily in GX surface soil (ρ=0.54, p=0.0034, n=27) and SK rhizosphere (ρ=0.62, p=0.014, n=15) (Fig. 1C, Fig. S2). Noticeably, GX initially showed higher F1-Hg than SK, but these differences diminished over the growing season with both sites reaching comparable levels after day 100 (Fig. 1C). This decline aligns with the post-harvest period when rice fields are dried and aerated, promoting immobilization of dissolved Hg species through redox-driven precipitation and adsorption processes (e.g., Fe-oxyhydroxide precipitation and adsorption) (12,59).

Several key factors showed no significant differences between sites. Soil pH displayed similar temporal trends at HX and GX, while SK showed more apparent fluctuation, but with no significant site differences overall (Kruskal-Wallis test, p>0.05, Fig. S3). Terminal electron acceptors including sulfate (avg. ~100 mg/kg or 100 μM) and nitrate (avg. ~ 4 mg/kg or 6.45 μM) demonstrated similar temporal patterns across all sites. Hydrogen sulfide remained consistently low (0-5 μM) with no significant site differences (Kruskal-Wallis test, p>0.05, Fig. S3). This suggests similar rates of microbial sulfate reduction and constraints on Hg methylation or bioavailability as sulfide can readily react with Hg(II), forming HgS (60,61). Fe^{2+}/total Fe ratios showed similar temporal trends except in HX subsurface porewater, which displayed lower ratios suggesting reduced iron reduction at the depth of rice roots.

The three sites differ primarily in bioavailable Hg content (*i.e.,* F1-Hg) and DOM, which are both variable known to control Hg methylation (62–64). Since DOM affects Hg(II) bioavailability at low sulfide conditions ($\lesssim$30 μM) (65,66), the consistently low sulfide levels and differences in F1-Hg suggest DOM potentially contributes to observed MeHg fluctuations. Interestingly, DOM aromaticity (i.e., $SUVA_{254}$) showed a pronounced effect at SK, where more lower DOM $SUVA_{254}$, indicating higher DOM bioavailability (67), correlated with higher MeHg in both soil compartments, consistent with previous findings in rice paddy environments (62).

Taken together, our geochemical data suggest that iron and sulfur cycling processes have limited influence on net MeHg accumulation and are insufficient to explain the observed

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

299 spatiotemporal patterns of MeHg fluctuation across our study sites. While ORP measurements
300 indicate that HX conditions favor microaerophilic to anaerobic processes (denitrification,
301 manganese reduction), and GX and SK are more conducive to iron reduction and sulfate
302 reduction, these differences in redox environments do not adequately account for the MeHg
303 distribution patterns observed. Rather, our findings suggest that net MeHg accumulation is
304 likely controlled by methylating microbes beyond SRB and IRB and active demethylation
305 processes. These results highlight the need to assess the potential contributions of alternative
306 microbial methylators and demethylators to MeHg dynamics in rice paddy environments under
307 the geochemical conditions observed in this study.
308

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

MeHg levels show limited association with *hgcA* gene abundance across metabolic guilds




Figure 2. Analysis of *hgcA* genes and their correlation with MeHg. (A) Taxonomic distribution and abundance of *hgcA* genes across specific metabolic guilds throughout the sampling period. Color patterns represent iron-reducing bacteria (IRB, blue patterns), sulfate-reducing bacteria (SRB, green patterns), methanogens (orange/red patterns), and syntrophs (purple patterns). Note that sequences not classified to known functional groups were excluded from panel A for visualization purposes but are included in total abundance calculations. (B) Spearman correlation coefficients between MeHg concentrations and gene abundances for different metabolic guilds. Syntrophs are not shown due to insufficient data. Asterisks indicate statistical significance levels (* $p<0.05$; ns: not significant).

As the key marker gene of Hg methylation, we searched for and classified *hgcA* in the assembled metagenomes (Fig. 2) to assess methylation potential and identify the microbial players involved. Total *hgcA* abundance was, on average, higher at mining-impacted sites GX (mean=$8.4\times10^{-8}\pm3.0\times10^{-8}$; n=16) and SK (mean=$8.4\times10^{-8}\pm2.7\times10^{-8}$; n=17) compared to the background site (mean=$5.5\times10^{-8}\pm1.8\times10^{-8}$; n=16), aligning with the higher MeHg levels observed at contaminated sites (Fig.1, Fig. 2). Metatranscriptomic analysis confirmed active *hgcA* expression across all sites (n=6), with similar expression in surface soils ($\sim1.0\times10^{-8}$) but substantially higher expression in GX and SK rhizospheres ($\sim1.4–1.7\times10^{-8}$), supporting the functional relevance of the metagenomic potential (Fig. S4A).

Temporally, *hgcA* abundance at HX remained relatively stable, mirroring MeHg trends (Fig. 1B). Across the growing season, correlations between total *hgcA* abundance and soil

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

MeHg were inconsistent with guild-specific differences (Fig. 2B). Taxonomically, *hgcA* was primarily assigned to *Deferrimicrobiaceae* at HX, whereas *Methanoregulaceae*-related *hgcA* increased at GX and SK, particularly in rhizosphere samples.

As an additional test of drivers of MeHg accumulation, we fitted multiple linear regression models with model selection by AICc (Supplementary Result 2). The best-supported model included additive effects of bioavailable Hg(II) (F1-Hg) and overall *hgcA*, while controlling for site and soil compartment. In this model, F1-Hg was a major positive predictor of MeHg concentrations (β=0.73, $p<0.001$), whereas total *hgcA* abundance was not significant. Adding an interaction between F1-Hg and *hgcA* did not improve fit (ΔAICc=3). Consistent with this, the explained variance (adjusted R^2) increased from ~60% (additive model without site/compartment) to ~84% when site and compartment effects were included, underscoring other environmental influences on MeHg.

These results suggest that net MeHg accumulation is not primarily controlled by methylation potential in Hg-contaminated rice paddies, but rather by Hg substrate availability and other environmental factors. This contrasts with findings from river systems where direct correlations between *hgcA* abundance and MeHg production have been observed under more uniform geochemical conditions (68–70). In rice paddies, the relationship between *hgcA* and MeHg content is complicated by several factors: (i) Pronounced geochemical gradients in DOM characteristics, redox conditions, and Hg bioavailability; (ii) the presence of multiple MeHg sinks including biotic and abiotic demethylation processes (20,71); and (iii) MeHg uptake by rice plants, which demonstrate efficient MeHg bioaccumulation from paddy soil (i.e., bioaccumulation factors (BAFs) of 5.5-5.6) (72,73). Although the quantitative effect of plant uptake on soil MeHg concentrations was not considered in this study, plant-mediated removal may partially contribute to soil MeHg content we observed. The interplay between MeHg production, degradation, and plant uptake processes operating simultaneously in rice paddies underscores the need to consider the genetic potential of methylating microbes alongside broader ecosystem variables that operate at larger spatial scales to better predict the fate of MeHg in rice paddy systems.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

merB-mediated demethylation and MeHg dynamics

Following *hgcA* analyses, we investigated demethylation potential represented by *merB*, which encodes the organomercurial lyase that performs reductive demethylation (RD) (74,75). Total *merB* gene abundance in rhizosphere was consistently higher compared to surface soil, which were similar across the three sites (HX:mean=$5.1\times10^{-8}\pm5.5\times10^{-8}$, n=11; GX: $3.5\times10^{-8}\pm5.2\times10^{-8}$, n=11; SK:$5.8\times10^{-8}\pm3.9\times10^{-8}$, n=11) (Fig. S5). The consistently higher *merB* abundance in rhizosphere samples is likely attributed to radial oxygen loss (ROL) (76), creating a more oxidizing condition conducive to aerobic *merB*-carrying microbes (77), although direct soil ORP measurements were not obtained in this study. The highest rhizosphere *merB* was observed at day 0, followed by a gradual decline, aligning with previous findings that ROL rates decrease over the rice growth period (77). This temporal trend further supports the potential role of plant-mediated oxygen dynamics in structuring the *merB*-carrying community

Multiple regression analysis revealed that *merB* gene abundance has a significant positive association with MeHg levels ($\beta=2.51\times10^{6}$, p=0.0113) after accounting for site-specific effects. The positive association between gene abundance and absolute MeHg concentrations suggest adaptation of microbial communities to Hg-contaminated environments. In highly polluted aquatic systems, increased THg load drives higher absolute MeHg concentrations (74). The microbial communities respond by enriching for Hg-resistant bacteria carrying the *mer*-operon, including *merB* genes (74).

Crucially, this correlation between *merB* and MeHg is particularly strong in GX (Spearman's $\rho=0.6$; n=5) and SK (Spearman's $\rho=0.83$; n=6) rhizosphere samples. This suggests that the ROL-driven oxidizing conditions select for the aerobic *merB*-carrying organisms, and where THg loads are high (GX & SK), these adapted populations are most enriched, leading to the strong co-variance between *merB* gene abundance and MeHg concentration in this specific micro-environment.

At the transcriptomic level, *merB* transcripts were detected only in HX surface soils (n=6, Fig. S4B). This disconnect from gene abundance likely reflects: (i) site-and compartment-specific conditions regulating *merB* induction (e.g., oxygen and DOM), which may have been favorable only at HX surface at the sampling time (78); (ii) the transient nature of *merB* transcription and the short half-lives of *mer* transcripts, making them easy to miss in single time-point sampling; and (iii) temporal fluctuations in Hg(II) bioavailability that narrow the window for detectable *merB* expression (74,79). These findings highlight the need for temporal transcriptomic sampling to better establish relationships between *merB* expression and MeHg dynamics *in situ*.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Methane-cycling communities show coupled dynamics and shared responses to Hg contamination




Figure 3. Community structure and environmental drivers of methanogens and aerobic methanotrophs. (A) Taxonomic distribution and relative abundance of *mcrA* genes across sampling days. (B) Taxonomic distribution and relative abundance of *pmoA*/*mmoX* genes across sampling days. (C&D): Distance-based redundancy analysis (db-RDA) of beta-diversity in relation to geochemical variables, with panel C showing methanogens and panel D showing methanotrophs. Points represent individual samples colored by site. Arrows indicate environmental variables with significance levels (*** $p<0.001$, ** $p<0.01$, * $p<0.05$).

Recently, the role of methane-cycling microbes in mediating MeHg dynamics has become increasingly evident through controlled laboratory studies (14,22,80–82) However, the environmental relevance and field-scale implications of these laboratory-derived mechanisms remain largely unexplored in complex agricultural systems.

We show that despite contrasting Hg contamination sources (GX: atmospheric deposition; SK: historical smelting residues), methane-cycling communities at GX and SK exhibited similar beta-diversity (PERMANOVA, $p<0.001$; Fig. 3C&D). Constrained ordinations showed that geochemical variables explained 27.67% of the beta-diversity in methanogens and 22.14% in methanotrophs ($p=0.001$ and 0.004, respectively; Fig. 3C–D).

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

418 THg and MeHg were significant predictors for both groups, and DOM concentration showed
419 an additional significant association with methanogen beta-diversity (permutation tests, $p<0.05$;
420 Supplementary Result 3).

421 Methanogen and methanotroph abundances were significantly and positively correlated
422 ($\beta=0.51$, $p=0.003$, $n=50$; Fig. S6), indicating that their populations tended to co-vary across
423 samples. However, methanotrophs were consistently present at roughly an order of magnitude
424 lower abundance than methanogens (Fig. 3A&B). This observation aligns with studies
425 demonstrating positive co-occurrence networks between these groups (83) and synchronized
426 responses to environmental factors in rice paddies (84), likely reflecting their metabolic
427 interdependence. Methanotrophs rely on methane produced by methanogens and may, in turn,
428 help regulate local oxygen concentrations (83,84). This interdependence is further supported
429 by the notably lower abundances of both groups at HX, the site with the highest average ORP,
430 where conditions appeared too oxidizing for robust methanogenesis and suboptimal for aerobic
431 methanotrophy, resulting in reduced populations of both metabolically linked functional groups.

432 The similarity of methane-cycling community structures at Hg-impact sites, in addition
433 to their significant association with THg and MeHg suggests the presence of a core methane-
434 cycling community that adapts similarly to Hg impacts regardless of contamination source.
435

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Context-dependent methanogen-MeHg relationships links to DOM quality and compartment effects

We subsequently examined how methanogen abundance (via *mcrA*) links to MeHg concentrations across sites and soil compartments. Overall methanogen abundance showed marginal association with MeHg (β=-1.30×10^{7}, p=0.064), but this relationship varied with THg concentration (β=4.57×10^{2}, p=0.049), indicating context-dependent relationships where higher methanogen abundance amplifies MeHg production under elevated THg conditions.

Methanogen community beta-diversity correlated significantly with DOM (Fig. 3C). Additionally, methanogen abundance showed negative correlations with both DOM quantity (β=-5.94×10^{-8}, p=0.007) and $SUVA_{254}$ (β=-2.88×10^{-8}, p=0.003), suggesting lower abundance where DOM is less bioavailable. Similarly, DOM $SUVA_{254}$ correlated negatively with MeHg (β=-0.48, p=0.005) while DOM quantity showed no significant relationship (β=0.60, p=0.140), aligning with previous work showing that DOM bioavailability more critically determines MeHg concentrations than quantity (10,67).

Our correlational field data reveal significant associations between DOM, methanogens, and MeHg, these observational relationships that are consistent with established experimental evidence for DOM's dual role in methylation processes. Higher $SUVA_{254}$ values indicate aromatic, humified DOM that reduces MeHg through strong Hg complexation (10,67). Conversely, bioavailable DOM fractions (autochthonous DOM, fresh organic matter, low-molecular-weight acids, algal organic matter) enhance MeHg production by providing carbon sources and stimulating methylating microorganisms, including methanogens (10,63,67). The negative correlations with $SUVA_{254}$ support this framework: Highly aromatic DOM suppresses both methanogenic proliferation and MeHg formation, while bioavailable DOM promote both processes.

Specific methanogen taxa showed compartment-dependent patterns. *Methanoregulaceae*, the dominant family across sites with diverse Hg methylators (7), (Fig. 3A, Supplementary Result 4), mirrored overall methanogen trends with positive MeHg correlations in SK surface soil (ρ=0.745, p=0.011; n=11) but negative correlations in SK rhizosphere (ρ=-0.943, p=0.017; n=6). These family-specific *mcrA* patterns differ from the relationships shown in Fig. 2, which reports methanogen-associated *hgcA*; methanogens may directly or indirectly contribute to methylation and demethylation, so *mcrA*–MeHg associations do not necessarily mirror *hgcA*–MeHg patterns.

Minor taxa showed different compartment preferences. *Methanospirillaceae* (ρ=0.974, p=0.005; n=6) and *Methanosarcinaceae* (ρ=0.941, p=0.005; n=6) correlated positively with MeHg in SK rhizosphere, with weaker patterns at HX and GX. Metatranscriptomic analysis (n=6) detected active *mcrA* expression across sites, with higher levels at Hg-contaminated sites,

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

473 particularly by *Methanoregulaceae*, providing functional support for gene abundance data (Fig.
474 S4C).

475 Interestingly, we recovered sequences from *Methanoperedenaceae*, an archaeal group
476 (ANME), which performs anaerobic oxidation of methane (AOM) via reverse methanogenesis
477 (85). This finding corroborates the detection of *Methanoperedenaceae*-associated *hgcA* from
478 our rice paddy metagenomes (Fig. 2A, Supplementary Result 7). Although *mcrA* and *hgcA*
479 transcripts associated with *Methanoperedenaceae* were detected at the Hg-impacted sites, (Fig.
480 S4C), we exclude this group from subsequent analyses due to the lack of empirical evidence
481 for their methylation capacity. Their potential role in Hg cycling are discussed in
482 Supplementary Discussion 1.
483

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Flooding associates higher methanotroph abundance with lower MeHg

Methanotroph abundance relationships with MeHg followed similar site-specific patterns to those observed for methanogens, consistent with the two guilds co-varying across samples (Fig. A&B). However, when focusing specifically on the flooded period, multiple regression revealed a significant negative association between methanotroph abundance and MeHg concentrations ($\beta = -3.21 \times 10^7$, $p=0.009$), after adjusting for site and soil compartment, an effect not observed in the methanogen data.

Metagenomic analyses revealed that *Methylomonadaceae*, *Beijerinckiaceae*, and *Methylococcaceae* were the most abundant methanotroph taxa (Fig. 3B, Supplementary Result 5). Metatranscriptomic data further confirmed active methanotrophy at all sites, with particularly high *pmoA* expression at SK, where *Methylomonadaceae* was the most highly expressed methanotroph family (Fig. S4D). *Methylomonadaceae* showed the strongest negative correlations with MeHg at GX ($\rho=-0.8$, $p=0.133$; $n=5$) and SK ($\rho=-0.9$, $p=0.083$; $n=5$) rhizosphere, while other methanotroph taxa showed weaker correlations (Supplementary Result 6).

It is likely that the negative methanotroph–MeHg relationship reflects either (i) flooding-induced oxygen depletion that suppresses aerobic methanotrophs while stimulating anaerobic Hg methylation, or (ii) the influence of taxa such as *Methylomonadaceae*, which are adapted to low-oxygen conditions via anaerobic denitrification (86–88), allowing persistence in flooded, microaerophilic rhizospheres and potential methanotroph-mediated demethylation via a methanobactin-related pathway, which we explore in the next section.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Exploratory analysis of *mbnT* as a potential demethylation biomarker




Figure 4. Presence/Absence of Methanobactin-Related Genes in Methanotroph Genomes. This heatmap illustrates the presence and absence of methanobactin biosynthesis and transport genes across experimentally studied methanotrophs (24) and MAGs recovered in this study, along with their respective quality metrics. The x-axis denotes the methanobactin genes, while the left y-axis represents the taxonomic classification of each genome, and the right y-axis lists the RefSeq or MAG IDs. All RefSeq genomes lack MBn biosynthesis, except OB3B (Note *mbnB* is present is OB3B as a pseudogene, which is not recovered through the HMM based analyses, but confirmed with RefSeq annotation).

Recent studies indicate a methanotroph-specific MeHg demethylation pathway facilitated by methanobactins (24,25). Methanobactins are copper-sequestering chalkophores essential for particulate methane monooxygenase (pMMO) activity in some methanotrophs (89). While primarily binding copper with high affinity, methanobactins also bind other metals, including Hg(II), albeit with lower affinity (89).

The cellular uptake of methanobactins is mediated by MbnT, a TonB-dependent transporter (TBDT) , which is an outer membrane protein (90). This uptake is crucial as methanobactin appears to deliver MeHg inside the cell, and a study shows that deleting *mbnT* impairs methanobactin uptake and, consequently, MeHg degradation (24,25). The degradation of MeHg is thought to be carried out by the periplasmic methanol dehydrogenase, targeting the carbon-Hg bond, and is distinct from the *merB*-dependent organomercurial lyase pathway, as these methanotrophs generally lack the *merB* gene (24,25)

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Given the prevalence of methanotrophs in rice paddies, we investigated whether their observed association with MeHg dynamics is linked to MbnT-mediated demethylation. We examined the genomic context of previously studied demethylating methanotrophs and methanotroph-MAGs recovered from rice paddies to determine if they encode genes related to methanobactin biosynthesis (*i.e., mbnA, mbnB, mbnC* as core components, and additional enzymes like *mbnD, mbnF, mbnN, mbnS, mbnX*) and transport (*mbnE, mbnM, mbnT*) (47). Aligning with previous research, we confirmed that the demethylating *Methylosinus trichosporium OB3b*, encodes the full suite of genes for methanobactin biosynthesis and uptake/transport (Fig. 4). *Methylomicrobium album BG8* and *Methylocystis sp. strain Rockwell*, also able to demethylate, encode *mbnT* but lack genes necessary for methanobactin biosynthesis. Conversely, *Methylococcus capsulatus Bath* lacks genes for both methanobactin biosynthesis and uptake (*mbnT*) and was shown to not demethylate (Fig. 4) (24). Subsequently, we show that none of the methanotroph MAGs possessed *mbnT* genes sufficiently similar to those found in the experimentally confirmed demethylating methanotrophs, and these MAGs also lacked all essential genes for methanobactin biosynthesis. Therefore, the rice paddy MAGs are unlikely to exhibit MeHg demethylation activity, although some of these MAGs showed significant correlations with MeHg via network analyses (Supplementary Result 8 & Discussion 2).

Based on experimental evidence, we hypothesized that *mbnT* could serve as a marker for MeHg demethylation capacity as the demethylating methanotrophs are shown to possess the gene and subsequently analyzed *mbnT* from assembled metagenomes of rice paddies. We recovered *mbnT* that resemble those found in known demethylating methanotrophs using an HMM (TIGR01783, sequence cutoff 223.5). However, recognizing that TBDTs, including *mbnT*, have homologs in diverse organisms beyond methanotrophs, we employed a conservative approach, retrieving sequences classified to known demethylating methanotroph genera: *Methylomicrobium*, *Methylocystis*, and *Methylosinus* (Fig. S7). Transcriptomic analyses show that *mbnT* transcripts from these genera were present in Hg-contaminated rice paddies, though to a limited extent (n=6, Fig. S4E). Multiple regression analyses revealed that total *mbnT* gene abundance from these genera represents a significant positive predictor of MeHg concentration ($\beta=4.579 \times 10^{7}$, $p=0.0136$) after accounting for site and compartment specific effects (Supplementary Result 9), contrary to an expected negative correlation if *mbnT* represents effective demethylation.

Our findings suggest that while methanotroph demethylation capacity appears to be present in rice paddies, utilizing *mbnT* as a marker to assess net MeHg accumulation requires consideration of several limitations: (i) *mbnT* is not single-copy gene (91), and it belongs to the family of TBDTs, which are responsible for the uptake of diverse substrates, including metal chelates, and have numerous homologs in organisms other than methanotrophs (47).

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Despite using stringent cutoff, the presence of *mbnT* homologs with high similarity could lead to biased abundance estimations. Furthermore, gene abundance dose not directly equate to activity. (ii) Methanotrophs exhibit varied demethylation rates (24). These rates depend on various biological factors, including the structure of methanobactin (24), the efficiency of the MbnT transport system (89), and the presence of additional copper-handling proteins, like MbnH and MbnP (90), as well as environmental factors such as redox conditions, pH, and the presence of competing metals (92–94). (iii) Methanobactins can influence Hg cycling through secondary mechanisms. This includes altering microbial community composition (*e.g.,* through copper competition impacting denitrifying bacteria (89)) or affecting the bioavailability of Hg species via competitive binding (94,95). These broader effects complicate the interpretation of *mbnT*'s role in MeHg demethylation.

Future studies should prioritize establishing direct relationships between *mbnT* expression and demethylation rates *in situ* through quantitative approaches with adequate sample sizes, before linking *mbnT* abundance to environmental MeHg dynamics. Additionally, broader experimental validation across diverse methanotrophic taxa is needed to confirm the universal applicability of this biomarker. Given the complexity revealed by our environmental data, controlled laboratory studies examining *mbnT* expression under varying Hg, redox, and competing metal conditions would help define when *mbnT* serves as a reliable indicator of methanotroph-mediated MeHg demethylation.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Methane-cycling microbes as key biological predictors of MeHg variability




Figure 5. Variable importance and model performance for predicting MeHg concentrations using random forest analysis (full model). (A) Variable importance plot showing the mean percent increase in mean squared error (%IncMSE) for each predictor variable when permuted, with 95% confidence intervals across 100 different data splits. Larger point size and green coloration indicate higher significance rates. Values show the percentage of data splits where the variable was significant ($p<0.05$) followed by the mean p-value. (B) Model performance metrics displaying mean values with 95% confidence intervals across all data splits. The model was built using 2001 trees with mtry=4 and validated through 100 permutation tests with random 70/30 train/test splits.

Our analyses demonstrate that relying on single factors has limited explanatory power for net MeHg accumulation, despite significant associations found for individual variables. The complexity expected from concurrent methylation and demethylation processes is reflected in the increase in model R^2 when accounting for site and compartment effects (e.g., from 60% to 84% in hgcA models). To address the limitations of single-factor analyses and identify the key, potentially non-linear, variables driving MeHg patterns across diverse functional genes and varying environmental conditions, we employed Random Forest (RF) analysis.

The RF models built using all variables achieved moderate predictive performance ($R^2=0.70\pm0.13$, range: 0.40-0.95, Fig. 5). Bioavailable Hg(II) (F1-Hg) consistently emerged as the dominant predictor across all data splits (100% significance rate, p=0.010). Methanogen abundance ranked as the second most important variable (Mean IncMSE=20.8, 100% significance rate, p=0.013), followed by methanogen-hgcA abundance (Mean IncMSE=9.3, 49% significance rate), substantially outperforming other microbial indicators like total Hg methylators (*hgcA*), sulfate-reducing bacteria (SRB), and iron-reducing bacteria (IRB) abundances.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Methanotroph abundance was identified as the third most important biological variable overall (Mean IncMSE = 8.8; 31% significance rate). Crucially, an explicit interaction term between methanogen and methanotroph abundances (i.e., their product) was a critical predictor of MeHg (Mean IncMSE = 18.3; 80% significance rate; p=0.038). This result suggests that the methanotrophic influence is primarily expressed through community-level coupling with methane-producing populations. Given that methanogens are key methylators that amplify MeHg under elevated THg, and methanotrophs showed a significant negative association with MeHg during the flooded period (suggesting a role in MeHg consumption/demethylation), the interaction term highlights that the net MeHg outcome is highly sensitive to the balance and synchronous activity of these two metabolically interdependent microbial guilds.

Constraining the RF model solely to the flooded period substantially improved predictive accuracy (R^2=0.86±0.08; Fig. S8B). This improved performance was accompanied by a modest increase in the relative importance of methanogens (mean %IncMSE rose from approximately 21 to 23) and a decrease for F1-Hg (mean %IncMSE fell from approximately 32 to 26), suggesting that the predictive power of methanogens is comparatively greater under flooded, reduced conditions than that of Hg bioavailability alone. Integrating genomic and geochemical data through RF modeling revealed that methanogen, methanogen methylator, and methanotroph abundance were the top biological factors contributing to net MeHg accumulation, while Hg bioavailability and dissolved organic matter (DOM) aromaticity were the dominant geochemical factors. Prediction accuracy further improved when removing less influential variables, reinforcing the importance of these key factors.

While previous studies have associated methanogenesis with MeHg content in rice paddies (14,21), our work advances this understanding by identifying methanotrophy as an important biological correlate of net MeHg accumulation and characterizing specific methane-cycling populations associated with MeHg variability. Methanotrophic involvement is particularly noteworthy, as it suggests a previously underappreciated potential demethylation pathway in rice paddies. Our findings indicate that studying the relationship between microbial methane metabolism and Hg cycling provides critical insights into the key biogeochemical factors associated with net MeHg accumulation in Hg-contaminated rice paddies.

While these findings advance our understanding of microbial associations with MeHg dynamics, several considerations should guide their interpretation and application. The observed coupling between methane and Hg cycling, though mechanistically plausible, requires validation across diverse environmental contexts before broader generalization to other rice paddy systems. Our RF approach, while effective at identifying statistical associations, cannot establish causal directionality between predictors and MeHg accumulation, highlighting the need for controlled experimental validation of these relationships. We recommend validation across diverse rice paddy systems with (i) varying flooding patterns

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

649 and water management practices (12,96), (ii) different rice cultivars (97), and (iii) diverse
650 geographical span (3) and contamination sources (98–100) to benchmark model transferability.

651 Our results underscore that effective assessment of MeHg risk requires integrating
652 microbial community data with physicochemical gradients that govern both Hg bioavailability
653 and microbial niche occupation. The importance of methane-cycling communities in our
654 predictive models highlights promising avenues for management strategies that could
655 simultaneously address climate change mitigation and Hg pollution through targeted
656 manipulation of shared microbial pathways (101).
657

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Acknowledgement

This research was funded by the National Natural Science Foundation of China (Grant No. 42394092), the Guizhou Provincial Major Scientific and Technological Program ([2024]013) to BM, and the Natural Sciences and Engineering Research Council of Canada to AJP and DSG. We thank Tianli Zhan for contributing to the geochemical analyses. The bioinformatics analyses were conducted using computing clusters at the Digital Research Alliance of Canada.

Conflict of Interest

The authors declare no competing financial interest.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Reference:

1. Lindberg S et al. A Synthesis of Progress and Uncertainties in Attributing the Sources of Mercury in Deposition. AMBIO: A Journal of the Human Environment. 2007 Feb;36(1):19–33.

2. Zhao L et al. Mercury methylation in rice paddy and accumulation in rice plant: A review. Ecotoxicology and Environmental Safety. 2020 June;195:110462.

3. Liu M et al. Rice life cycle-based global mercury biotransport and human methylmercury exposure. Nat Commun. 2019 Nov 14;10(1):5164.

4. Meng B et al. The Process of Methylmercury Accumulation in Rice (*Oryza sativa* L.). Environ Sci Technol. 2011 Apr;45(7):2711–7.

5. Barkay T et al. Demethylation─The Other Side of the Mercury Methylation Coin: A Critical Review. ACS Environ Au. 2021 Nov 2;acsenvironau.1c00022.

6. Parks JM et al. The genetic basis for bacterial mercury methylation. Science. 2013;339(6125):1332–5.

7. Gilmour CC et al. Robust Mercury Methylation across Diverse Methanogenic Archaea. MBio. 2018 May 2;9(2):10-128.

8. Gilmour CC et al. Mercury Methylation by Novel Microorganisms from New Environments. Environ Sci Technol. 2013 Oct 15;47(20):11810–20.

9. Poulin BA et al. A comprehensive sulfate and DOM framework to assess methylmercury formation and risk in subtropical wetlands. Nat Commun. 2025 May 7;16(1):4253.

10. Bravo AG et al. The interplay between total mercury, methylmercury and dissolved organic matter in fluvial systems: A latitudinal study across Europe. Water Research. 2018 Nov 1;144:172–82.

11. Abdelhafiz MA et al. DOM influences Hg methylation in paddy soils across a Hg contamination gradient. Environmental Pollution. 2023 Apr 1;322:121237.

12. Wang X et al. Growing Rice Aerobically Markedly Decreases Mercury Accumulation by Reducing Both Hg Bioavailability and the Production of MeHg. Environ Sci Technol. 2014 Feb 4;48(3):1878–85.

13. Liu YR et al. Overlooked Role of Putative Non-Hg Methylators in Predicting Methylmercury Production in Paddy Soils. Environ Sci Technol. 2019 Nov 5;53(21):12330–8.

14. Wu Q et al. Methanogenesis Is an Important Process in Controlling MeHg Concentration in Rice Paddy Soils Affected by Mining Activities. Environ Sci Technol. 2020 Nov 3;54(21):13517–26.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

15. Zhang R et al. Mining-impacted rice paddies select for Archaeal methylators and reveal a putative (Archaeal) regulator of mercury methylation. ISME COMMUN. 2023 July 15;3(1):74.

16. Liu YR et al. Analysis of the Microbial Community Structure by Monitoring an Hg Methylation Gene (hgcA) in Paddy Soils along an Hg Gradient. Applied and Environmental Microbiology. 2014 May 1;80(9):2874–9.

17. Liu YR et al. Unraveling Microbial Communities Associated with Methylmercury Production in Paddy Soils. Environ Sci Technol. 2018 Nov 20;52(22):13110–8.

18. Si Y et al. Mercury methylation coupled to iron reduction by dissimilatory iron-reducing bacteria. Chemosphere. 2015 Mar 1;122:206–12.

19. A. Vishnivetskaya T et al. Microbial community structure with trends in methylation gene diversity and abundance in mercury-contaminated rice paddy soils in Guizhou, China. Environmental Science: Processes & Impacts. 2018;20(4):673–85.

20. Bravo AG et al. Persistent Hg contamination and occurrence of Hg-methylating transcript (hgcA) downstream of a chlor-alkali plant in the Olt River (Romania). Environ Sci Pollut Res. 2016 June 1;23(11):10529–41.

21. Hao Z et al. Relative importance of aceticlastic methanogens and hydrogenotrophic methanogens on mercury methylation and methylmercury demethylation in paddy soils. Science of The Total Environment. 2024 Jan;906:167601.

22. Kronberg RM et al. Mechanisms of Methyl Mercury Net Degradation in Alder Swamps: The Role of Methanogens and Abiotic Processes. Environ Sci Technol Lett. 2018 Apr 10;5(4):220–5.

23. Marvin-DiPasquale MC et al. Bacterial Methylmercury Degradation in Florida Everglades Peat Sediment. Environ Sci Technol. 1998 Sept 1;32(17):2556–63.

24. Kang-Yun CS et al. Evidence for methanobactin “Theft” and novel chalkophore production in methanotrophs: impact on methanotrophic-mediated methylmercury degradation. ISME J. 2022 Jan;16(1):211–20.

25. Lu X et al. Methylmercury uptake and degradation by methanotrophs. Sci Adv. 2017 May 31;3(5):e1700041.

26. Jeyakumar P et al. Copper and zinc spiking of biosolids: effect of incubation period on metal fractionation and speciation and microbial activity. Environ Chem. 2008;5(5):347.

27. Tessier A et al. Sequential extraction procedure for the speciation of particulate trace metals. Anal Chem. 1979 June 1;51(7):844–51.

28. Wang J et al. Ammonium thiosulphate enhanced phytoextraction from mercury contaminated soil – Results from a greenhouse study. Journal of Hazardous Materials. 2011 Feb 15;186(1):119–27.

29. Chen S et al. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018 Sept 1;34(17):i884–90.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

30. Andrews S. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data [Internet]. 2010 [cited 2021 June 8]. Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

31. Li D et al. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics. 2015 May 15;31(10):1674–6.

32. Kang DD et al. MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. PeerJ. 2019 July 26;7.

33. Wu YW et al. MaxBin 2.0: an automated binning algorithm to recover genomes from multiple metagenomic datasets. Bioinformatics. 2016 Feb 15;32(4):605–7.

34. Alneberg J et al. Binning metagenomic contigs by coverage and composition. Nat Methods. 2014 Nov;11(11):1144–6.

35. Nissen JN et al. Improved metagenome binning and assembly using deep variational autoencoders. Nat Biotechnol. 2021 May;39(5):555–60.

36. Chklovski A et al. CheckM2: a rapid, scalable and accurate tool for assessing microbial genome quality using machine learning. Nat Methods. 2023 Aug;20(8):1203–12.

37. Sieber CMK et al. Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. Nat Microbiol. 2018 July;3(7):836–43.

38. Olm MR et al. dRep: a tool for fast and accurate genomic comparisons that enables improved genome recovery from metagenomes through de-replication. ISME J. 2017 Dec;11(12):2864–8.

39. Eren AM et al. Community-led, integrated, reproducible multi-omics with anvi'o. Nat Microbiol. 2021 Jan;6(1):3–6.

40. Chaumeil PA et al. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. Hancock J, editor. Bioinformatics. 2019 Nov 15;btz848.

41. Magoč T et al. FLASH: fast length adjustment of short reads to improve genome assemblies. Bioinformatics. 2011 Nov 1;27(21):2957–63.

42. Kopylova E et al. SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. Bioinformatics. 2012;28(24):3211–7.

43. Hyatt D et al. Prodigal: prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics. 2010 Mar 8;11(1):119.

44. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. 2013 Mar 16 [cited 2022 Feb 4]; Available from: http://arxiv.org/abs/1303.3997v2

45. Liao Y et al. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. 2014 Apr 1;30(7):923–30.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

774 46. Eddy SR. HMMER User's Guide Biological sequence analysis using profile hidden
775 Markov models. 2020 Nov;227.

776 47. Dassama LMK et al. Methanobactins: from genome to function. Metallomics. 2017 Jan
777 25;9(1):7–20.

778 48. Gionfriddo C et al. Hg-MATE-Db.v1.01142021 [Internet]. The Smithsonian Institution;
779 2021 [cited 2021 Jan 31]. p. 8700889 Bytes. Available from:
780 https://smithsonian.figshare.com/articles/dataset/Hg-MATE-
781 Db_v1_01142021/13105370

782 49. Christakis CA et al. Expanded Diversity and Phylogeny of mer Genes Broadens
783 Mercury Resistance Paradigms and Reveals an Origin for MerA Among Thermophilic
784 Archaea. Frontiers in Microbiology. 2021;12.

785 50. Lafrance-Vanasse J et al. Crystal Structures of the Organomercurial Lyase MerB in Its
786 Free and Mercury-bound Forms: INSIGHTS INTO THE MECHANISM OF
787 METHYLMERCURY DEGRADATION*. Journal of Biological Chemistry. 2009 Jan
788 9;284(2):938–44.

789 51. Garber AI et al. FeGenie: A Comprehensive Tool for the Identification of Iron Genes
790 and Iron Gene Neighborhoods in Genome and Metagenome Assemblies. Frontiers in
791 Microbiology. 2020;11.

792 52. Steinegger M et al. MMseqs2 enables sensitive protein sequence searching for the
793 analysis of massive data sets. Nat Biotechnol. 2017 Nov;35(11):1026–8.

794 53. Parks DH et al. A complete domain-to-species taxonomy for Bacteria and Archaea. Nat
795 Biotechnol. 2020 Sept;38(9):1079–86.

796 54. Robinson MD et al. edgeR: a Bioconductor package for differential expression analysis
797 of digital gene expression data. Bioinformatics. 2010;26(1):139–40.

798 55. Aroney STN et al. CoverM: read alignment statistics for metagenomics. Bioinformatics.
799 2025 Apr 1;41(4):btaf147.

800 56. Ruan Q et al. Local similarity analysis reveals unique associations among marine
801 bacterioplankton species and environmental factors. Bioinformatics. 2006 Oct
802 15;22(20):2532–8.

803 57. Xia LC et al. Extended local similarity analysis (eLSA) of microbial community and
804 other time series data with replicates. BMC Syst Biol. 2011 Dec 14;5(2):S15.

805 58. Kelly CA et al. Is total mercury concentration a good predictor of methyl mercury
806 concentration in aquatic systems? Water Air Soil Pollut. 1995 Feb 1;80(1):715–24.

807 59. Wang J et al. Mobilization, Methylation, and Demethylation of Mercury in a Paddy Soil
808 Under Systematic Redox Changes. Environ Sci Technol. 2021 July 20;55(14):10133–
809 41.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

60. Ullrich SM et al. Mercury in the Aquatic Environment: A Review of Factors Affecting Methylation. Critical Reviews in Environmental Science and Technology. 2001 July 1;31(3):241–93.

61. Zhang Y et al. Biochar and nitrate reduce risk of methylmercury in soils under straw amendment. Science of The Total Environment. 2018 Apr 1;619–620:384–90.

62. Tang W et al. Increased Methylmercury Accumulation in Rice after Straw Amendment. Environ Sci Technol. 2019 June 4;53(11):6144–53.

63. Pu Q et al. Dissolved organic matter fosters core mercury-methylating microbiomes for methylmercury production in paddy soils. Biogeosciences. 2025 Mar 24;22(6):1543–56.

64. Bravo AG et al. Biotic formation of methylmercury: A bio–physico–chemical conundrum. Limnol Oceanogr. 2020 May;65(5):1010–27.

65. Poulin BA et al. Effects of Sulfide Concentration and Dissolved Organic Matter Characteristics on the Structure of Nanocolloidal Metacinnabar. Environ Sci Technol. 2017 Nov 21;51(22):13133–42.

66. Graham AM et al. Effect of Dissolved Organic Matter Source and Character on Microbial Hg Methylation in Hg–S–DOM Solutions. Environ Sci Technol. 2013 June 4;47(11):5746–54.

67. Hansen AM et al. Optical properties of dissolved organic matter (DOM): Effects of biological and photolytic degradation. Limnology and Oceanography. 2016;61(3):1015–32.

68. Peterson BD et al. Metabolically diverse microorganisms mediate methylmercury formation under nitrate-reducing conditions in a dynamic hydroelectric reservoir. The ISME Journal. 2023 Oct 1;17(10):1705–18.

69. Jones DS et al. Diverse Communities of hgcAB+ Microorganisms Methylate Mercury in Freshwater Sediments Subjected to Experimental Sulfate Loading. Environ Sci Technol. 2020 Nov 17;54(22):14265–74.

70. Capo E et al. Expression Levels of *hgcAB* Genes and Mercury Availability Jointly Explain Methylmercury Formation in Stratified Brackish Waters. Environ Sci Technol. 2022 Sept 20;56(18):13119–30.

71. Christensen GA et al. Determining the Reliability of Measuring Mercury Cycling Gene Abundance with Correlations with Mercury and Methylmercury Concentrations. Environ Sci Technol. 2019 Aug 6;53(15):8649–63.

72. Zhang H et al. Bioaccumulation of Methylmercury versus Inorganic Mercury in Rice (Oryza sativa L.) Grain. Environ Sci Technol. 2010 June 15;44(12):4499–504.

73. Liu J et al. Methylmercury production in a paddy soil and its uptake by rice plants as affected by different geochemical mercury pools. Environment International. 2019 Aug 1;129:461–9.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

74. Schaefer JK et al. Role of the Bacterial Organomercury Lyase (MerB) in Controlling Methylmercury Accumulation in Mercury-Contaminated Natural Waters. Environ Sci Technol. 2004 Aug 1;38(16):4304–11.

75. Barkay T et al. Bacterial mercury resistance from atoms to ecosystems. FEMS Microbiol Rev. 2003 June;27(2–3):355–84.

76. Li Y et al. Root-induced changes in radial oxygen loss, rhizosphere oxygen profile, and nitrification of two rice cultivars in Chinese red soil regions. Plant Soil. 2013 Apr 1;365(1):115–26.

77. Wang X et al. Radial oxygen loss has different effects on the accumulation of total mercury and methylmercury in rice. Plant Soil. 2014 Dec 1;385(1):343–55.

78. Sanz-Sáez I et al. Microorganisms Involved in Methylmercury Demethylation and Mercury Reduction are Widely Distributed and Active in the Bathypelagic Deep Ocean Waters. Environ Sci Technol. 2024 Aug 6;58(31):13795–807.

79. Barkay T et al. Effects of dissolved organic carbon and salinity on bioavailability of mercury. Applied and Environmental Microbiology. 1997 Nov;63(11):4267–71.

80. Barkay T et al. Microbial Transformations of Mercury: Potentials, Challenges, and Achievements in Controlling Mercury Toxicity in the Environment. In: Advances in Applied Microbiology. Elsevier; 2005. p. 1–52.

81. Oremland RS et al. Methylmercury Decomposition in Sediments and Bacterial Cultures: Involvement of Methanogens and Sulfate Reducers in Oxidative Demethylation. Appl Environ Microbiol. 1991 Jan;57(1):130–7.

82. Lu X et al. Anaerobic Mercury Methylation and Demethylation by *Geobacter bemidjiensis* Bem. Environ Sci Technol. 2016 Apr 19;50(8):4366–73.

83. Wang W et al. Methanogen–methanotroph community has a more consistent and integrated structure in rice rhizosphere than in bulk soil and rhizoplane. Molecular Ecology. 2024;33(13):e17416.

84. Lee HJ et al. Methane emission and dynamics of methanotrophic and methanogenic communities in a flooded rice field ecosystem. FEMS Microbiology Ecology. 2014 Apr 1;88(1):195–212.

85. Haroon MF et al. Anaerobic oxidation of methane coupled to nitrate reduction in a novel archaeal lineage. Nature. 2013 Aug;500(7464):567–70.

86. Deng Y et al. Methylomonadaceae was the active and dominant methanotroph in Tibet lake sediments. ISME Communications. 2024 Jan 1;4(1):ycae032.

87. Grégoire DS et al. Microbial methane cycling in a landfill on a decadal time scale. Nat Commun. 2023 Nov 16;14(1):7402.

88. Kits KD et al. Methane oxidation coupled to nitrate reduction under hypoxia by the Gammaproteobacterium ethylomonas denitrificans, sp. nov. type strain FJG1. Environmental Microbiology. 2015;17(9):3219–32.

bioRxiv preprint doi: https://doi.org/10.1101/2025.10.09.681528; this version posted October 11, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

89. Reyes RM et al. Methanobactins: Structures, Biosynthesis, and Microbial Diversity. Annual Review of Microbiology. 2024 Nov 10;78(1):383–401.

90. Dassama LMK et al. Methanobactin transport machinery. Proc Natl Acad Sci USA. 2016 Nov 15;113(46):13027–32.

91. Peng P et al. Two TonB-Dependent Transporters in Methylosinus trichosporium OB3b Are Responsible for Uptake of Different Forms of Methanobactin and Are Involved in the Canonical "Copper Switch." Applied and Environmental Microbiology. 2022 Jan 11;88(1):e01793-21.

92. Tian HJ et al. Ecological drivers of methanotrophic communities in paddy soils around mercury mining areas. Science of The Total Environment. 2020 June;721:137760.

93. Choi DW et al. Spectral and thermodynamic properties of Ag(I), Au(III), Cd(II), Co(II), Fe(III), Hg(II), Mn(II), Ni(II), Pb(II), U(IV), and Zn(II) binding by methanobactin from *Methylosinus trichosporium* OB3b. Journal of Inorganic Biochemistry. 2006 Dec 1;100(12):2150–61.

94. Yin X et al. Synergistic Effects of a Chalkophore, Methanobactin, on Microbial Methylation of Mercury. Applied and Environmental Microbiology. 2020 May 19;86(11):e00122-20.

95. Stenzler BR et al. Diffusion of H2S from anaerobic thiolated ligand biodegradation rapidly generates bioavailable mercury. Environmental Microbiology. 2022;24(7):3212–28.

96. Qian H et al. Greenhouse gas emissions and mitigation in rice agriculture. Nat Rev Earth Environ. 2023 Sept 26;4(10):716–32.


97. Peng X et al. Reducing total mercury and methylmercury accumulation in rice grains through water management and deliberate selection of rice cultivars. Environmental Pollution. 2012 Mar 1;162:202–8.


98. Horvat M et al. Total mercury, methylmercury and selenium in mercury polluted areas in the province Guizhou, China. Science of The Total Environment. 2003 Mar;304(1–3):231–56.


99. Xu X et al. The local impact of a coal-fired power plant on inorganic mercury and methyl-mercury distribution in rice (Oryza sativa L.). Environmental Pollution. 2017 Apr;223:11–8.


100. Liang P et al. Human exposure to mercury in a compact fluorescent lamp manufacturing area: By food (rice and fish) consumption and occupational exposure. Environmental Pollution. 2015 Mar 1;198:126–32.


101. Li C et al. Aligning global mercury mitigation with climate action. Nat Commun. 2025 Sept 28;16(1):7826.


A
Surface soil
Rhizosphere (Rice growing period)
THg concentration (mg/kg)
Rice growing period
Post harvest
B
MeHg concentration (μg/kg)
C
F1−fraction Hg (μg/kg)
Days after rice planting
Site
HX
GX
SK


A
Surface
Rhizosphere
HX
GX
SK
% total reads normalised by HMM length
5e−08
4e−08
3e−08
2e−08
1e−08
0e+00
−20
0
20
40
60
80
100
120
200
260
290
Days after rice planting
Family
Deferrimicrobiaceae
Geoalkalibacteraceae
Geobacteraceae
Desulfatiglandaceae
Desulfobulbaceae
Desulfocapsaceae
Desulfomonilaceae
Desulfosalsimonadaceae
Desulfovibrionaceae
Desulfurivibrionaceae
Methanocellaceae
Methanomassiliicoccaceae
Methanomicrobiaceae
Methanoregulaceae
Methanosarcinaceae
Methanosphaerulaceae
Methanospirillaceae
Methanoperedenaceae
Smithellaceae
Syntrophaceae
Syntrophobacteraceae
B
Surface
Rhizosphere
Total
SRB
IRB
Methanogen
Correlation Coefficient
0.5
0.0
−0.5
−1.0
ns
*
HX
GX
SK
Site
Site HX GX SK


A
B
C
D
Surface
Rhizosphere
HX
GX
SK
% total reads normalised by HMM length
Days post rice plantation
Methanoregulaceae
Methanotrichaceae
Methanosphaerulaceae
Methanobacteriaceae
Methanocellaceae
Methanosarcinaceae
uc_Methanomicrobiales
Methanocorpusculaceae
Methanomassiliicoccaceae
Methanofastidiosaceae
Methanofollaceae
Methylomonadaceae
Beijerinckiaceae
Methylococcaceae
Methyloligellaceae
Burkholderiaceae.B
Methylophilaceae
RDA2 (15.94% of fitted, 4.41% of total)
RDA1 (60.27% of fitted, 16.68% of total)
DOM *
MeHg *
THg ***
pH
DOM SUVA254
sulfate
nitrate
Variance explained by constrained axes: 27.67%
RDA2 (28.13% of fitted, 6.23% of total)
RDA1 (50.56% of fitted, 11.2% of total)
nitrate
pH
DOM SUVA254
sulfate
THg **
DOM
MeHg **
Variance explained by constrained axes: 22.14%
Site HX GX SK


Methanobactin Biosynthesis
Transport
RefSeq or MAG ID
Methylomicrobium album BG8
Methylocystis sp. strain Rockwell
Methylosinus trichosporium OB3b
Methylococcus capsulatus str. Bath
GCF_000214275.2
GCF_000188155.2
GCF_000178815.2
GCA_000008325.1
Methylomirabilia Class
CSP1-6 Family
Alphaproteobacteria Class
Methyloligellaceae Family
Methylomirabilia Class
2-02-FULL-66-22 Family
HX–017
HX–041
SK–072
SK–005
SK–083
HX–029
SK–064
GX–048
MeHg Demethylation
No
Yes
Gene Presence
Absence
Presence
Completeness
100–90
90–80
80–70
70–60
Redundancy
7.5–5
5–2.5
2.5–0
mbnA
mbnB
mbnC
mbnD
mbnF
mbnN
mbnS
mbnX
mbnT
mbnE
mbnM
MDH


A
F1Hg
Total Methanogen
Total Methanogen hgcA
Total Methanotroph
DOM SUVA254
DOM
Total SRB
Total hgcA
Total SRB hgcA
Total IRB hgcA
Total merB
Total IRB
100.0% (p=0.010)
100.0% (p=0.013)
49.0% (p=0.082)
31.0% (p=0.091)
17.0% (p=0.126)
5.0% (p=0.195)
6.0% (p=0.236)
3.0% (p=0.282)
2.0% (p=0.248)
0.0% (p=0.328)
0.0% (p=0.346)
1.0% (p=0.407)
Predictor Variable
0
10
20
30
Mean Percent Increase in Mean Squared Error (%IncMSE)
Significance
Consistently Significant (p<0.05 in >=90% of splits)
Sometimes Significant (p<0.05 in 10–89% of splits)
Rarely Significant (p<0.05 in <10% of splits)
B
Value
1.2
0.9
0.6
0.3
0.0
0.704
(95% CI: 0.466–0.907)
0.835
(95% CI: 0.682–0.952)
0.719
(95% CI: 0.465–1.022)
0.561
(95% CI: 0.376–0.792)
R²
Pearson r
RMSE
MAE