An independently detected and vetted 5.45-day, Earth-sized transit candidate in the TOI-913 system (TIC 407126408)

Josh Gold
Independent researcher, Acton, Massachusetts, USA

Preprint, version 2, 12 October 2026 (version 1: 11 October 2026). Not peer reviewed.
Version 1 DOI: 10.5281/zenodo.23290579

Abstract

We report a transit-like signal with period $P = 5.450138 \pm 0.000016$ d and depth 196^{+24}_{-21} ppm on the K dwarf TOI-913 (TIC 407126408; $V = 10.5$, $d = 65.2$ pc), which hosts the confirmed sub-Neptune TOI-913 b ($P = 11.0994$ d). The signal was found by an automated multi-sector search of TESS 2-minute photometry spanning Sectors 12–102 (2019–2026; 34 transits) and corresponds to a radius of $1.12 \pm 0.09\,R_\oplus$ if it arises on the target. It passes ten automated false-positive tests, including odd/even depth, secondary eclipse, shape, per-sector consistency and a season-holdout prediction test. It is independently recovered as the strongest periodic signal in Quick-Look Pipeline full-frame-image photometry. In 200 block-shuffled copies of the light curve, no transit search reached the observed signal strength (empirical false-alarm probability $< 1.5\%$ at 95% confidence), although this test does not capture systematics coherent over many days. A transit fit with free stellar density yields $\rho_\star = 2.43^{+0.96}_{-1.02}\,\rho_\odot$, consistent with the catalogue value $2.09 \pm 0.45\,\rho_\odot$. The only Gaia neighbour previously flagged as able to produce the depth, TIC 407126405 at 9.5″, is a background giant ($R \approx 10\,R_\odot$, $d \approx 5$ kpc) whose geometry cannot reproduce the observed 2.4 h duration. TESS difference-image centroids cannot separate it from the target, so this exclusion rests on the duration alone. With archival Gemini/Zorro speckle constraints, TRICERATOPS gives FPP = 0.054 and NFPP = 0.041, which meets neither the "validated" nor the "likely planet" criteria of Giacalone et al. (both require NFPP < 0.001). The residual nearby-eclipsing-binary probability is dominated by three faint stars at 26.5″, 56.9″ and 58.0″; per-sector centroids from an independent pipeline disfavour the first two. Seeing-limited ground-based photometry could test all three. The signal was also detected by the independent open-source search `eumemic/exoplanets` (2026 October 8), which set it aside because of the 9.5″ neighbour, and at a detection strength (MES 8.0–9.6) where that search finds about a third of vetted signals to be false alarms. We present it as a planet candidate, not a confirmed planet. If real, it lies near the 2:1 period commensurability with TOI-913 b ($P_b/P_c = 2.037$).

1 Introduction

Faint transit signals from small planets can become detectable only once several years of TESS photometry are combined. This project was inspired by the approach of P. Rabtsevich [1], who stacked TESS sectors from several years for a single star and then tried systematically to falsify the resulting candidate, including by hiding one observing season and predicting its transit times. We automated that approach for a sample of nearby K and M dwarfs. In a 200-star pilot, one

previously unreported signal survived all automated tests. This note documents that signal and our attempts to rule it out.

TOI-913 (TIC 407126408) is a K dwarf ($T_{\rm eff} = 4969\,$K, $R_\star = 0.733\,R_\odot$, $M_\star = 0.82\,M_\odot$; TIC 8 [8]) at 65.2 pc (Gaia DR3 parallax $15.31 \pm 0.02\,$mas [9]). It hosts TOI-913 b, a $2.45\,R_\oplus$ planet at 11.0986 d, statistically validated by Mistry et al. [4], and the long-period candidate TOI-913.02, seen in two transits.

2 Data and search

We used SPOC [6] PDCSAP 2-minute light curves for Sectors 12, 13, 65, 66, 93, 100, 101 and 102, retrieved with `lightkurve` [10]. Quality-flagged cadences were removed and only upward outliers were clipped. Each sector was normalized and detrended with a biweight filter (`wotan` [11]) with a 0.61 d window. We searched with transit least squares (TLS [12]) over 0.5–40 d on 10-minute bins, iteratively masking each detection. Detection required SDE ≥ 9, SNR ≥ 7 and ≥ 3 transits. TLS 2.0 multiplies trial durations by a "fill factor" (data span over baseline). On multi-year data with large gaps, this shortened durations about 20× and suppressed SNR about 4×, so we disabled it.

After TOI-913 b (SDE 125.5, SNR 34.3), the search found a 27.3 d systematic, which the kill tests rejected, and then the 5.45 d signal (SDE 23.6, SNR 8.9, 34 transits). The 5.45 d signal is therefore not the strongest peak in the search until that systematic is masked; in an unmasked search of the standard light curve it ranks sixth (Table 2). Before the kill tests, the light curve was re-detrended with all detected transits masked.

Pipeline validation. Before the pilot, the pipeline was run on 20 hosts of 26 confirmed small planets and on 10 detached eclipsing binaries from the TESS EB catalogue [16]. It recovered 23/26 planets at the correct period, and all 23 survived every hard test. The three misses had only 1–2 sectors. It rejected 8/9 detected binaries; the exception is a twin-eclipse binary detected at half its period. In an injection-recovery grid on two quiet stars, planets $\geq 1.5\,R_\oplus$ were recovered out to 15 d, and planets $\geq 2\,R_\oplus$ out to 30 d.

3 The signal

Figure 1 shows the folded transit. We fit both planets jointly with `batman` [13] and `emcee` [14]. Each planet had a free period, epoch, $R_p/R_\star$ and impact parameter, on circular orbits, with quadratic limb darkening fixed to $(0.451, 0.189)$ and a jitter term. We ran two fits: one with a shared stellar density and a catalogue prior, and one with an independent density per planet and no prior. Table 1 lists the results.

4 Vetting

4.1 Automated tests

The signal passed all hard tests: odd/even depth (211 vs 245 ppm, 0.4σ); secondary eclipse at phase 0.5 and across phases 0.3–0.7 (maximum 2.4σ); duration plausibility; size; per-sector depth consistency ($p = 0.17$); harmonic relation to TOI-913 b ($P_b/P = 2.037$, not an integer ratio); and




Figure 1: Phase-folded TESS photometry of TOI-913 (Sectors 12–102) at the candidate period (left) and at TOI-913 b's period (right), each with the other planet's model removed. Grey: 2-minute data; black: 20-minute bins; orange: joint transit model.

Parameter	Candidate (".03")	TOI-913 b
Period (d)	5.450138 ± 0.000016	11.09938 (fit)
T_0 ($\mathrm{BJD_{TDB}}$)	$2458628.6479^{+0.0046}_{-0.0073}$	—
Depth (ppm)	196^{+24}_{-21}	912^{+27}_{-24}
$R_p/R_\star$	0.0140 ± 0.0008	—
R_p ($R_\oplus$)	1.12 ± 0.09	2.4
Impact parameter	$0.42^{+0.13}_{-0.21}$	$0.19^{+0.16}_{-0.13}$
T_{14} (h)	$2.37^{+0.16}_{-0.15}$	3.27
$\rho_\star$, free fit ($\rho_\odot$)	$2.43^{+0.96}_{-1.02}$	$1.61^{+0.40}_{-0.75}$
a (AU), T_{eq} (K, $A = 0$)	0.057, ~ 860	0.091, ~ 680

Table 1: Transit parameters. The joint fit has a shared density with a catalogue prior of $2.09 \pm 0.45\,\rho_\odot$; the free fit has no prior. The R_p uncertainty includes a 5% uncertainty on $R_\star$.

an injection test at the same period and depth. The trapezoid shape test was inconclusive at this SNR. In a season-holdout test, each of the four observing seasons was hidden, the ephemeris was fitted on the rest, and depth was measured only at the predicted times. Three of four seasons showed the transit at $> 2\sigma$. The fourth (Sector 93 alone) gave 1.8σ. Figure 2 shows the diagnostic page. A fresh, read-only AI review of that page flagged the $9.5''$ neighbour and the depth scatter between sectors, both addressed below.

4.2 Robustness

We remeasured the depth over the central 60% of the transit, with red-noise-aware errors, across variants (Table 2). Every season and sector jackknife gives 196–244 ppm. No transit departs by $> 3\sigma$ from the mean, and removing the three deepest leaves 190 ± 25 ppm. One of 34 transits lies within 0.5 d of a data gap (8% expected by chance), and none lie within 0.1 d of a momentum dump (43 dump events, comprising 231 flagged cadences; version 1 called these "231 momentum dumps"). The only non-confirming variant is uncorrected SAP flux. There the signal is weaker (159 ± 24 ppm, SNR 6.6) and a narrow search drifts to 5.427 d.




Figure 2: Pipeline diagnostic page: full light curve with predicted transits (red), fold, odd/even, phase 0.5, depth per sector, the holdout test, and folds per season.

Variant	Depth (ppm)	SNR	Full 0.5–40 d TLS search
PDCSAP, biweight 0.61 d	225 ± 24	9.5	rank 6 (systematics unmasked)
Biweight 0.5 / 0.75 / 1.0 / 1.5 d	230 / 232 / 231 / 235	9.0–9.9	—
Cosine detrending	297 ± 23	13.1	**5.4501 d top peak** (SDE 12.8)
QLP FFI photometry (8 sectors)	250 ± 23	10.6	**5.4501 d top peak** (SDE 15.2)
TESS-SPOC FFI	185 ± 31	6.0	—
SAP flux (uncorrected)	159 ± 24	6.6	not recovered

Table 2: Signal depth across data products and detrending choices, with TOI-913 b and TOI-913.02 masked.

4.3 Is the signal on the target?

Stellar density. With no prior, the transit shape implies $\rho_\star = 2.43^{+0.96}_{-1.02}\,\rho_\odot$. That agrees with the TIC value (2.09 ± 0.45) and with the density from TOI-913 b's transit.

The 9.5″ neighbour. TIC 407126405 ($T = 14.05$, $\Delta T = 4.4$) lies within one TESS pixel. It is a background giant: Gaia parallax 0.16 ± 0.02 mas, TIC $R = 10.0\,R_\odot$, luminosity class giant. To produce the observed dilution it would need a $\approx 1.1\%$ eclipse. An orbit with $P = 5.45$ d around a $10\,R_\odot$, 1–3 $M_\odot$ star has $a/R_\star \approx 1.4$–1.9, so a central transit lasts ≈ 23–33 h. A grazing chord short enough to last 2.4 h would occult at most $\sim$0.02% of the stellar disk. We therefore exclude this star as the source. TRICERATOPS assigns it negligible probability ($\sim 10^{-26}$), but this follows from the same catalogue radius and is not independent evidence. No pixel-level test separates the two stars (below), so the exclusion rests on the duration argument and the TIC radius.

Pixel-level tests. We stacked per-sector difference images and compared fits with the source at the target or at a neighbour (Figure 3). For the 9.5″ star, the test has little power at this SNR. A control on TOI-913 b prefers the target at only $\approx 2\sigma$, and the candidate gives no preference (0.5σ). It is likewise inconclusive for the two faint neighbours below (0.8σ and 0.2σ). In one sector (S65), the fit prefers the 56.9″ star at $\approx 3\sigma$, but this is not significant over all sectors. We report these as non-constraining.

Independent centroids. The `eumemic/exoplanets` team fitted PRF centroids to difference images in all eight sectors [2]. Per-sector offsets from the target range from 0.4″ to 26.8″, and the quality-weighted mean lies about 5″ from the target, toward TIC 407126405. They therefore do not separate the target from the 9.5″ giant, and we do not use them for that purpose. (An earlier statement of a 0.36″ offset referred to the best single sector and was withdrawn by that team.) They do argue against the two faint stars that dominate the NFPP: no sector places the centroid within 11″ of TIC 407126406 or within 30″ of TIC 10001766874. In that team's calibration on known TOIs, signals from stars 21–63″ away produced offsets above 15″ in 98–100% of cases. This does not cover TIC 407126397, and it is TESS-pixel evidence, not a substitute for ground-based photometry.

High-resolution imaging. Archival speckle imaging on ExoFOP shows no companions to contrasts of $\Delta m = 6.47$ at 0.5″ (Gemini/Zorro, 832 nm, 2020 March 14) and $\Delta m = 6.9$ at 1″ (SOAR/HRCam, I, 2019 August 12).

4.4 Periodograms and empirical false-alarm probability

The search threshold (SDE ≥ 9) is calibrated on white noise, so we measured the false-alarm probability on this star's own light curve (Figure 4; code `periodogram_fap.py`). All tests use the light curve of Section 2.




Figure 3: Per-sector difference images (top: lost flux over noise) and out-of-transit images (bottom). ×: target; ○: TIC 407126405; ⋆: best-fit location. TESS's 21″ pixels cannot separate the 9.5″ pair at this SNR.

Matched-filter test. We reran the same TLS search on 200 copies of the light curve in which 1-day blocks were shuffled within each sector. This keeps noise on timescales below a day and destroys any coherent period. Each copy used the masks in place when the pipeline found the signal (TOI-913 b, the two TOI-913.02 events and the 27.3 d systematic). On the real data, 5.4501 d is the top peak, at SDE 21.8. The highest SDE anywhere in 0.5–40 d across the shuffled copies had a median of 13.1, a 99th percentile of 18.1 and a maximum of 18.8. None reached 21.8, so the false-alarm probability is $< 1.5\%$ (95% upper limit). Block shuffling does not reproduce systematics coherent over many days, such as the 27.3 d signal that outranks the candidate in the unmasked search, so the true false-alarm probability may be higher than this test indicates. For comparison, the `eumemic/exoplanets` search of inverted light curves finds that $37 \pm 4\%$ of vetted signals at MES 9–10 are false alarms, averaged over $\sim 8 \times 10^5$ stars [2].

Generalized Lomb–Scargle. A GLS periodogram [17] does not detect the signal: at 5.45 d the power is 6.0×10^{-5}, with Baluev false-alarm probability [18] ≈ 1. It does not detect TOI-913 b either. With b left in the light curve, b's GLS power is 1.8×10^{-4} (rank 1653, false-alarm probability ≈ 1), although TLS detects it at SDE 125. A transit of 2–3 h fills 1–2% of the orbit and is poorly matched by a sinusoid. The only GLS peaks above the 0.1% level lie below 0.8 d, at the scale of the detrending window. GLS therefore cannot decide this signal either way.

Spectral window and momentum dumps. The window function has power 1.1×10^{-3} at 5.45 d, against 0.055 at its strongest peak (38 d). An alias of TOI-913 b at 5.45 d would require window power at 10.7 d, where it is 7×10^{-4}. The 43 momentum-dump events are spaced 3.4–6.8 d apart (median 6.4 d). In their periodogram, 5.45 d lies at the 43rd percentile of power. Folded at 5.45 d, no dump falls in transit (0.8 expected), and their phases show no clustering (Rayleigh $Z = 0.04$).

4.5 False-positive probability

We ran TRICERATOPS v1.1 [15] on the folded light curve, using the SPOC apertures of all eight sectors and 8 runs of $N = 10^6$. Without the contrast curve, FPP $= 0.155 \pm 0.008$ and NFPP $= 0.032 \pm 0.001$. With the Zorro 832 nm curve, FPP $= 0.054 \pm 0.002$ and NFPP $= 0.041 \pm 0.001$. The probability of a planet on the target is 0.92. NFPP is dominated by NEB scenarios on TIC 10001766874 ($T = 15.91$, 56.9″; would need a 31.7% eclipse; 0.020), TIC 407126406 ($T = 16.97$, 26.5″; 22.7%; 0.017) and TIC 407126397 ($T = 14.9$, 58.0″; about 10%; 0.004). Every other star contributes less than 0.001. Giacalone et al. [15] define a "validated planet" as FPP < 0.015 and




Figure 4: From top: GLS periodogram with TOI-913 b left in, with Baluev false-alarm levels; spectral window; periodogram of the momentum-dump times; and the maximum TLS SDE in each of 200 block-shuffled light curves, against the observed 5.45 d signal (orange).

NFPP < 0.001, and a "likely planet" as FPP < 0.5 and NFPP < 0.001. With NFPP $= 0.041$, the candidate meets neither: it is a planet candidate, not a likely or validated planet. (Version 1 said it met the FPP criterion for a likely planet, which was misleading, since that category also requires the NFPP criterion.) TIC 10001766874 has an unusual TIC identifier and may be a catalogue artefact.

5 Prior detections

No CTOI exists for this star on ExoFOP. SPOC multi-sector searches through Sectors 65, 69 and 96 produced TCEs only for TOI-913 b and TOI-913.02. Scaled to the five sectors available through S96, our signal would have SNR ≈ 7.0, near the SPOC threshold of 7.1. In Sectors 12–13 alone, it has SNR 5.4, which explains its absence from searches limited to early data. The RAVEN FFI survey [20] (Sectors 1–55) lists a weak second signal on this star at 3.70 d, which we do not recover in eight sectors (TLS SNR < 6 near 3.70 d). The independent open-source search `eumemic/exoplanets` [3] detected the same signal ($P = 5.450121$ d, 220 ppm, MES 8.0; MES 9.6 at 5.45017 d, 224 ppm in a second search with b masked [2]) in its 2026 October 8 release, with the status "close neighbour could produce the depth". Our duration argument (Section 4.3) addresses this concern; that team's pipeline requires MES ≥ 10 for candidates.

6 Discussion

If confirmed, the candidate would be a $\sim 1.1\,R_\oplus$ planet interior to TOI-913 b, with $P_b/P_c = 2.037$, just wide of the 2:1 commensurability. For a near-resonant pair, transit-timing variations are expected with a super-period of $|1/P_c - 2/P_b|^{-1}$; this ratio places it near 304 d. Its mass, with an expected RV semi-amplitude of order $0.5\,\mathrm{m\,s^{-1}}$, is beyond most current spectrographs. The most direct next step is seeing-limited photometry of TIC 10001766874, TIC 407126406 and TIC 407126397 during a predicted transit, to exclude the remaining nearby-eclipsing-binary scenarios. The ephemeris uncertainty in late 2026 is about ± 15 min. The 200 ppm transit on the target itself is beyond typical ground-based precision, so the strongest test of the signal is TESS Sector 114 (February–March 2027), for which we pre-registered predicted transit times before the data exist [19].

7 Conclusion

We present TIC 407126408.03 (proposed designation, pending any CTOI assignment) as a planet candidate. It is a $1.12 \pm 0.09\,R_\oplus$, 5.450138 d transit signal that survives automated vetting, independent-product recovery, stellar-density consistency and high-resolution imaging constraints, with FPP $= 0.054$ and an empirical false-alarm probability $< 1.5\%$ in block-shuffled light curves. It is not statistically validated, nor a "likely planet" (NFPP $= 0.041$). Ground-based photometry of three faint neighbours, and the pre-registered TESS Sector 114 transits, are the next tests.

Acknowledgements

This work was inspired by P. Rabtsevich (@p_rabtsevich), who shared his TESS candidate search and methods openly [1]. We thank the `eumemic/exoplanets` project for publishing its independent search openly, and for its all-sector centroid analysis and false-alarm statistics. We thank P. Mistry for comments that prompted the periodogram and false-alarm tests in version 2. This paper includes data collected by the TESS mission, funded by NASA's Science Mission Directorate, and obtained from MAST at STScI. It uses data and services from ExoFOP-TESS, including high-resolution imaging contributed by S. Howell (Gemini/Zorro) and A. Tokovinin (SOAR/HRCam), and data from ESA's Gaia mission. We used QLP and SPOC data products, `lightkurve`, `wotan`, `transitleastsquares`, `batman`, `emcee`, `astropy`, `astroquery` and TRICERATOPS.

AI assistance. The search pipeline, vetting code and much of this analysis were written and run with substantial assistance from Anthropic's Claude AI coding agent, directed and reviewed by the author. Independent fresh AI sessions were used as adversarial reviewers of diagnostic plots. All numbers in this note come from the released code and data.

Data availability. Code, per-test result files and figures accompany this preprint in the same Zenodo record, and the code is at `https://github.com/killercatfish/tess-hunt`.

Changes from version 1. Added Section 4.4 (periodograms and empirical false-alarm probability) and Figure 4. Corrected the TRICERATOPS classification (Section 4.5): the candidate is not a "likely planet". Corrected "231 momentum dumps" to 43 dump events (231 flagged cadences). Reported NFPP as 0.041 ± 0.001 (the run mean, 0.0415; version 1 gave 0.042) and added the third faint star, TIC 407126397. Stated that the $9.5''$ giant is excluded by transit duration alone, and added the independent centroid analysis and the search rank of the signal.

References

[1] Rabtsevich, P. 2026, preprint, Zenodo, doi:10.5281/zenodo.22967456

[2] eumemic 2026, comments on GitHub issue *TIC 407126408 (TOI-913) 5.45 d: independent detection plus vetting of the* $9.5''$ *neighbour*, 12 October 2026, `https://github.com/eumemic/exoplanets/issues/2`

[3] eumemic 2026, *Multi-sector TESS transit searches of nearby K and M dwarfs and of TOI hosts*, `https://github.com/eumemic/exoplanets`

[4] Mistry, P., et al. 2023, AJ, 166, 9 (VaTEST II)

[5] Ricker, G. R., et al. 2015, JATIS, 1, 014003

[6] Jenkins, J. M., et al. 2016, Proc. SPIE, 9913, 99133E

[7] Huang, C. X., et al. 2020, RNAAS, 4, 204

[8] Stassun, K. G., et al. 2019, AJ, 158, 138

[9] Gaia Collaboration, Vallenari, A., et al. 2023, A&A, 674, A1

[10] Lightkurve Collaboration 2018, ASCL, 1812.013

[11] Hippke, M., David, T. J., Mulders, G. D., & Heller, R. 2019, AJ, 158, 143

[12] Hippke, M., & Heller, R. 2019, A&A, 623, A39

[13] Kreidberg, L. 2015, PASP, 127, 1161

[14] Foreman-Mackey, D., et al. 2013, PASP, 125, 306

[15] Giacalone, S., et al. 2021, AJ, 161, 24

[16] Prša, A., et al. 2022, ApJS, 258, 16

[17] Zechmeister, M., & Kürster, M. 2009, A&A, 496, 577

[18] Baluev, R. V. 2008, MNRAS, 385, 1279

[19] Gold, J. 2026, pre-registration, Zenodo, doi:10.5281/zenodo.23290795

[20] Lafarga, M., et al. 2026, arXiv:2603.22597