

7.8

Improper Integrals

Improper Integrals

In this section we extend the concept of a definite integral to the case where the interval is infinite and also to the case where f has an infinite discontinuity in $[a, b]$. In either case the integral is called an *improper* integral.



Type 1: Infinite Intervals

Type 1: Infinite Intervals

Consider the infinite region S that lies under the curve $y = 1/x^2$, above the x -axis, and to the right of the line $x = 1$.

You might think that, since S is infinite in extent, its area must be infinite, but let's take a closer look.

The area of the part of S that lies to the left of the line $x = t$ (shaded in Figure 1) is

$$\begin{aligned} A(t) &= \int_1^t \frac{1}{x^2} dx = \left. -\frac{1}{x} \right|_1^t \\ &= 1 - \frac{1}{t} \end{aligned}$$




Figure 1

Type 1: Infinite Intervals

Notice that $A(t) < 1$ no matter how large t is chosen. We also observe that

$$\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left(1 - \frac{1}{t} \right) = 1$$

The area of the shaded region approaches 1 as $t \rightarrow \infty$ (see Figure 2), so we say that the area of the infinite region S is equal to 1 and we write

$$\int_1^{\infty} \frac{1}{x^2} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2} dx = 1$$




Figure 2

Type 1: Infinite Intervals

Using this example as a guide, we define the integral of f (not necessarily a positive function) over an infinite interval as the limit of integrals over finite intervals.

Type 1: Infinite Intervals

1 Definition of an Improper Integral of Type 1

(a) If $\int_a^t f(x) dx$ exists for every number $t \geq a$, then

$$\int_a^\infty f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx$$

provided this limit exists (as a finite number).

(b) If $\int_t^b f(x) dx$ exists for every number $t \leq b$, then

$$\int_{-\infty}^b f(x) dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$$

provided this limit exists (as a finite number).

The improper integrals $\int_a^\infty f(x) dx$ and $\int_{-\infty}^b f(x) dx$ are called **convergent** if the corresponding limit exists and **divergent** if the limit does not exist.

(c) If both $\int_a^\infty f(x) dx$ and $\int_{-\infty}^a f(x) dx$ are convergent, then we define

$$\int_{-\infty}^\infty f(x) dx = \int_{-\infty}^a f(x) dx + \int_a^\infty f(x) dx$$

In part (c) any real number a can be used.

Type 1: Infinite Intervals

Any of the improper integrals in Definition 1 can be interpreted as an area provided that f is a positive function.

For instance, in case (a) if $f(x) \geq 0$ and the integral $\int_a^\infty f(x) dx$ is convergent, then we define the area of the region

$S = \{(x, y) \mid x \geq a, 0 \leq y \leq f(x)\}$ in Figure 3 to be

$$A(S) = \int_a^\infty f(x) dx$$

This is appropriate because $\int_a^\infty f(x) dx$ is the limit as $t \rightarrow \infty$ of the area under the graph of f from a to t .




Figure 3

Examples

Example 1. Determine whether the integral $\int_1^{\infty} (1/x) dx$ is convergent or divergent.

Example 2. Evaluate $\int_{-\infty}^0 x e^x dx$.

Example 3. Evaluate $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$.

Example 4. For what values of p is the integral

$$\int_1^{\infty} \frac{1}{x^p} dx$$

convergent?

Type 1: Infinite Intervals

We summarize the findings of Example 4 as follows:

2 $\int_1^{\infty} \frac{1}{x^p} dx$ is convergent if $p > 1$ and divergent if $p \leq 1$.



Type 2: Discontinuous Integrands

Type 2: Discontinuous Integrands

Suppose that f is a positive continuous function defined on a finite interval $[a, b)$ but has a vertical asymptote at b .

Let S be the unbounded region under the graph of f and above the x -axis between a and b . (For Type 1 integrals, the regions extended indefinitely in a horizontal direction. Here the region is infinite in a vertical direction.)

The area of the part of S between a and t (the shaded region in Figure 7) is

$$A(t) = \int_a^t f(x) \, dx$$




Figure 7

Type 2: Discontinuous Integrands

If it happens that $A(t)$ approaches a definite number A as $t \rightarrow b^-$, then we say that the area of the region S is A and we write

$$\int_a^b f(x) \, dx = \lim_{t \rightarrow b^-} \int_a^t f(x) \, dx$$

We use this equation to define an improper integral of Type 2 even when f is not a positive function, no matter what type of discontinuity f has at b .

Type 2: Discontinuous Integrands

3 Definition of an Improper Integral of Type 2

(a) If f is continuous on $[a, b)$ and is discontinuous at b , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx$$

if this limit exists (as a finite number).

(b) If f is continuous on $(a, b]$ and is discontinuous at a , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx$$

if this limit exists (as a finite number).

The improper integral $\int_a^b f(x) dx$ is called **convergent** if the corresponding limit exists and **divergent** if the limit does not exist.

(c) If f has a discontinuity at c , where $a < c < b$, and both $\int_a^c f(x) dx$ and $\int_c^b f(x) dx$ are convergent, then we define

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

Examples

Example 5. Find $\int_2^5 \frac{1}{\sqrt{x-2}} dx$.

Example 6. Determine whether $\int_0^{\frac{\pi}{2}} \sec x dx$ is convergent.

Example 7. Evaluate $\int_0^3 \frac{1}{x-1} dx$ if possible.

Example 8. Evaluate $\int_0^1 \ln x dx$.

Example. For what values of p does the integral $\int_0^1 \frac{1}{x^p} dx$ converge?



A Comparison Test for Improper Integrals

A Comparison Test for Improper Integrals

Sometimes it is impossible to find the exact value of an improper integral and yet it is important to know whether it is convergent or divergent.

In such cases the following theorem is useful. Although we state it for Type 1 integrals, a similar theorem is true for Type 2 integrals.

Comparison Theorem Suppose that f and g are continuous functions with $f(x) \geq g(x) \geq 0$ for $x \geq a$.

(a) If $\int_a^\infty f(x) dx$ is convergent, then $\int_a^\infty g(x) dx$ is convergent.

(b) If $\int_a^\infty g(x) dx$ is divergent, then $\int_a^\infty f(x) dx$ is divergent.

A Comparison Test for Improper Integrals

We omit the proof of the Comparison Theorem, but Figure 12 makes it seem plausible.




Figure 12

If the area under the top curve $y = f(x)$ is finite, then so is the area under the bottom curve $y = g(x)$.

A Comparison Test for Improper Integrals

And if the area under $y = g(x)$ is infinite, then so is the area under $y = f(x)$. [Note that the reverse is not necessarily true:

If $\int_a^\infty g(x) dx$ is convergent, $\int_a^\infty f(x) dx$ may or may not be convergent, and if $\int_a^\infty f(x) dx$ is divergent, $\int_a^\infty g(x) dx$ may or may not be divergent.]

Examples

Example 9. Show that $\int_0^{\infty} e^{-x^2} dx$ is convergent.

Example 10. Show that $\int_1^{\infty} \frac{1+e^{-x}}{x} dx$ is divergent.