

Week 4. Problems for class:

8. A 40.0-kg box initially at rest is pushed 5.00 m along a rough, horizontal floor with a constant applied horizontal force of 130 N. The coefficient of friction between box and floor is 0.300. Find (a) the work done by the applied force, (b) the increase in internal energy in the box–floor system as a result of friction, (c) the work done by the normal force, (d) the work done by the gravitational force, (e) the change in kinetic energy of the box, and (f) the final speed of the box.
16. **Q/C** The electric motor of a model train accelerates the train from rest to 0.620 m/s in 21.0 ms. The total mass of the train is 875 g. (a) Find the minimum power delivered to the train by electrical transmission from the metal rails during the acceleration. (b) Why is it the minimum power?

26. **Q/C** **Review.** As shown in Figure P8.26, a light string that does not stretch changes from horizontal to vertical as it passes over the edge of a table. The string connects m_1 , a 3.50-kg block originally at rest on the horizontal table at a height $h = 1.20$ m above the floor, to m_2 , a hanging 1.90-kg block originally a distance $d = 0.900$ m above the floor. Neither the surface of the table nor its edge exerts a force of kinetic friction. The blocks start to move from rest. The sliding block m_1 is projected horizontally after reaching the edge of the table. The hanging block m_2 stops without bouncing when it strikes the floor. Consider the two blocks plus the Earth as the system. (a) Find the speed at which m_1 leaves the edge of the table. (b) Find the impact speed of m_1 on the floor. (c) What is the shortest length of the string so that it does not go taut while m_1 is in flight? (d) Is the energy of the system when it is released from rest equal to the energy of the system just before m_1 strikes the ground? (e) Why or why not?




Figure P8.26

40. **S** A pendulum, comprising a light string of length L and a small sphere, swings in the vertical plane. The string hits a peg located a distance d below the point of suspension

(Fig. P8.40). (a) Show that if the sphere is released from a height below that of the peg, it will return to this height after the string strikes the peg. (b) Show that if the pendulum is released from rest at the horizontal position ($\theta = 90^\circ$) and is to swing in a complete circle centered on the peg, the minimum value of d must be $3L/5$.




Figure P8.40


41. A ball whirls around in a *vertical* circle at the end of a string. The other end of the string is fixed at the center of the circle. Assuming the total energy of the ball–Earth system remains constant, show that the tension in the string at the bottom is greater than the tension at the top by six times the ball’s weight.

S

44. Starting from rest, a 64.0-kg person bungee jumps from a tethered hot-air balloon 65.0 m above the ground. The bungee cord has negligible mass and unstretched length 25.8 m. One end is tied to the basket of the balloon and the other end to a harness around the person’s body. The cord is modeled as a spring that obeys Hooke’s law with a spring constant of 81.0 N/m, and the person’s body is modeled as a particle. The hot-air balloon does not move. (a) Express the gravitational potential energy of the person–Earth system as a function of the person’s variable height y above the ground. (b) Express the elastic potential energy of the cord as a function of y . (c) Express the total potential energy of the person–cord–Earth system as a function of y . (d) Plot a graph of the gravitational, elastic, and total potential energies as functions of y . (e) Assume air resistance is negligible. Determine the minimum height of the person above the ground during his plunge. (f) Does the potential energy graph show any equilibrium position or positions? If so, at what elevations? Are they stable or unstable? (g) Determine the jumper’s maximum speed.

Q/C

45. Review. A uniform board of length L is sliding along a smooth, frictionless, horizontal plane as shown in Figure P8.45a. The board then slides across the boundary with a rough horizontal surface. The coefficient of kinetic friction between the board and the second surface is μ_k . (a) Find the acceleration of the board at the moment its front end has traveled a distance x beyond the boundary. (b) The board stops at the moment its back end reaches the boundary as shown in Figure P8.45b. Find the initial speed v of the board.



Problems for practice at home:

9. A smooth circular hoop with a radius of 0.500 m is placed flat on the floor. A 0.400-kg particle slides around the inside edge of the hoop. The particle is given an initial speed of 8.00 m/s. After one revolution, its speed has dropped to 6.00 m/s because of friction with the floor. (a) Find the energy transformed from mechanical to internal in the particle–hoop–floor system as a result of friction in one revolution. (b) What is the total number of revolutions the particle makes before stopping? Assume the friction force remains constant during the entire motion.

[(a) 5.60J, (b) 2.28 revolutions]

32. **Q/C** As it plows a parking lot, a snowplow pushes an ever-growing pile of snow in front of it. Suppose a car moving through the air is similarly modeled as a cylinder of area A pushing a growing disk of air in front of it. The originally stationary air is set into motion at the constant speed v of the cylinder as shown in Figure P8.32.

In a time interval Δt , a new disk of air of mass Δm must be moved a distance $v \Delta t$ and hence must be given a kinetic energy $\frac{1}{2}(\Delta m)v^2$.

Using this model, show that the car's power loss owing to air resistance is $\frac{1}{2}\rho A v^3$

and that the resistive force acting on the car is $\frac{1}{2}\rho A v^2$, where ρ is the density of air. Compare this result with the empirical expression $\frac{1}{2}D\rho A v^2$ for the resistive force.




Figure P8.32