

Week 5. Problems for class:

- 23.** A proton, moving with a velocity of $v_i \hat{i}$, collides elastically with another proton that is initially at rest. Assuming that the two protons have equal speeds after the collision, find (a) the speed of each proton after the collision in terms of v_i and (b) the direction of the velocity vectors after the collision.

- 24.** A uniform piece of sheet metal is shaped as shown in Figure P9.24. Compute the x and y coordinates of the center of mass of the piece.




Figure P9.24

- 28.** The vector position of a 3.50-g particle moving in the xy plane varies in time according to $\vec{r}_1 = (3\hat{i} + 3\hat{j})t + 2\hat{j}t^2$, where t is in seconds and \vec{r} is in centimeters. At the same time, the vector position of a 5.50 g particle varies as $\vec{r}_2 = 3\hat{i} - 2\hat{i}t^2 - 6\hat{j}t$. At $t = 2.50$ s, determine (a) the vector position of the center of mass of the system, (b) the linear momentum of the system, (c) the velocity of the center of mass, (d) the acceleration of the center of mass, and (e) the net force exerted on the two-particle system.

- 33.** A rocket for use in deep space is to be capable of boosting a total load (payload plus rocket frame and engine) of 3.00 metric tons to a speed of 10 000 m/s. (a) It has an engine and fuel designed to produce an exhaust speed of 2 000 m/s. How much fuel plus oxidizer is required? (b) If a different fuel and engine design could give an exhaust speed of 5 000 m/s, what amount of fuel and oxidizer would be required for the same task? (c) Noting that the exhaust speed in part (b) is 2.50 times higher than that in part (a), explain why the required fuel mass is not simply smaller by a factor of 2.50.

- 52.** Sand from a stationary hopper falls onto a moving conveyor belt at the rate of 5.00 kg/s as shown in Figure P9.52. The conveyor belt is supported by frictionless rollers and moves at a constant speed of $v = 0.750$ m/s under the action of a constant horizontal external force \vec{F}_{ext} supplied by the motor that drives the belt. Find (a) the sand's rate of change of momentum in the horizontal direction, (b) the force of friction exerted by the belt on the sand, (c) the external force \vec{F}_{ext} , (d) the work done by \vec{F}_{ext} in 1 s, and (e) the kinetic energy acquired by the falling sand each second due to the change in its horizontal motion. (f) Why are the answers to parts (d) and (e) different?




Figure P9.52

53. Two particles with masses m and $3m$ are moving toward each other along the x axis with the same initial speeds v_i . Particle m is traveling to the left, and particle $3m$ is traveling to the right. They undergo an elastic glancing collision such that particle m is moving in the negative y direction after the collision at a right angle from its initial direction. (a) Find the final speeds of the two particles in terms of v_i . (b) What is the angle θ at which the particle $3m$ is scattered?

Problems for practice at home:

26. A rod of length 30.0 cm has linear density (mass per length) given by

$$\lambda = 50.0 + 20.0x$$

where x is the distance from one end, measured in meters, and λ is in grams/meter. (a) What is the mass of the rod? (b) How far from the $x = 0$ end is its center of mass?

[(a) 15.9g, (b) 0.153m]

41. Two gliders are set in motion on a horizontal air track. A light spring of force constant k is attached to the back end of the second glider. As shown in Figure P9.41, the first glider, of mass m_1 , moves to the right with speed v_1 , and the second glider, of mass m_2 , moves more slowly to the right with speed v_2 . When m_1 collides with the spring attached to m_2 , the spring compresses by a distance x_{\max} , and the gliders then move apart again. In terms of v_1 , v_2 , m_1 , m_2 , and k , find (a) the speed v at maximum compression, (b) the maximum compression x_{\max} , and (c) the velocity of each glider after m_1 has lost contact with the spring.




Figure P9.41

$$[(a) \frac{m_1 v_1 + m_2 v_2}{m_1 + m_2}, (b) (v_1 - v_2) \sqrt{\frac{m_1 m_2}{k(m_1 + m_2)}}, (c) \frac{2m_1 v_1 + (m_2 - m_1)v_2}{m_1 + m_2}, \dots]$$