

When possible you should work on the questions in advance and be prepared for discussions on them with your colleagues and tutor. Some questions are straightforward calculation while others require more discussion and thought. The computer icon  indicates where computer calculation or programming may be of use. Further questions can be found in the lecture notes as well as the recommended unit resources (see Moodle).

Sequence notation and defining formulas

(1) List the initial five terms of the following sequences.

(a) $\{a_n\}_{n=0}^{\infty}$, where $a_n = \frac{3n}{n^2+1}$.

The first five terms of the sequence are

$$\left[0, \frac{3}{2}, \frac{6}{5}, \frac{9}{10}, \frac{12}{17}\right]$$

(b) $\{b_n\}_{n=1}^{\infty}$, where $b_n = \frac{2^n}{2+2^n}$.

The first five terms of the sequence are

$$\left[\frac{1}{2}, \frac{2}{3}, \frac{4}{5}, \frac{8}{9}, \frac{16}{17}\right]$$

(c) $\{c_n\}_{n=1}^{\infty}$, where $c_n = \frac{(-1)^n}{3n}$.

The first five terms of the sequence are

$$\left[-\frac{1}{3}, \frac{1}{6}, -\frac{1}{9}, \frac{1}{12}, -\frac{1}{15}\right]$$

(d) $\{d_n\}_{n=1}^{\infty}$, where $d_n = \frac{(-1)^{n-1}}{2n}$.

The first five terms of the sequence are

$$\left[\frac{1}{2}, -\frac{1}{4}, \frac{1}{6}, -\frac{1}{8}, \frac{1}{10}\right]$$

(e) $\{e_n\}_{n=0}^{\infty}$, where $e_0 = e_1 = 1$ and for all $n \geq 2$, $e_n = e_{n-1} + e_{n-2}$.

The first five terms of the sequence are

$$[1, 1, 2, 3, 5]$$

(f) $\{f_n\}_{n=1}^{\infty}$, where $f_1 = 3$ and for all $n \geq 1$, $f_{n+1} = \frac{f_n}{n}$.

The first five terms of the sequence are

$$\left[3, 3, \frac{3}{2}, \frac{1}{2}, \frac{1}{8}\right]$$

(g) $\{g_n\}_{n=0}^{\infty}$, where $g_0 = 2$ and for all $n \geq 0$, $g_{n+1} = \frac{g_n}{1+g_n}$.

The first five terms of the sequence are

$$\left[2, \frac{2}{3}, \frac{2}{5}, \frac{2}{7}, \frac{2}{9}\right]$$

(h) $\{h_n\}_{n=0}^{\infty}$, where $h_n = \frac{3^n}{(2n)!}$.

The first five terms of the sequence are

$$\left[1, \frac{3}{2}, \frac{3}{8}, \frac{3}{80}, \frac{9}{4480}\right]$$

(2) For each of the sequences in question 1 find an expression for the $(2m+3)^{\text{rd}}$ element, simplified as far as possible.

The element a_{2m+3} is given by

$$a_{2m+3} = \frac{3(2m+3)}{(2m+3)^2+1}$$

The element b_{2m+3} is given by

$$b_{2m+3} = \frac{2^{2m+3}}{2^{2m+3} + 2}$$

The element c_{2m+3} is given by

$$c_{2m+3} = \frac{(-1)^{2m+3}}{3(2m+3)} = \frac{-1}{3(2m+3)}$$

The element d_{2m+3} is given by

$$d_{2m+3} = \frac{(-1)^{2m+2}}{2(2m+3)} = \frac{1}{2(2m+3)}$$

The element e_{2m+3} could be expressed as

$$e_{2m+3} = e_{2m+2} + e_{2m+1}.$$

Getting a more useful closed form expression would involve determining the closed form expression for the element e_n . This can generally be done for linear recurrence relations with enough initial conditions (as here), by finding the roots of the characteristic equation. Refer to section 7.4 of *Modern Engineering Mathematics* by Glyn James, see link to e-book version of this via the unit's Moodle area.

It is relatively straightforward to see that a nice solution in terms of factorials for the sequence $\{f_n\}_{n=1}^{\infty}$ is given by

$$f_n = \frac{3}{(n-1)!},$$

so the appropriate expression for f_{2m+3} is

$$f_{2m+3} = \frac{3}{(2m+2)!}.$$

The element g_{2m+3} can be expressed as

$$g_{2m+3} = \frac{g_{2m+2}}{1 + g_{2m+2}}.$$

The element h_{2m+3} can be written as

$$h_{2m+3} = \frac{3^{2m+3}}{(4m+6)!}.$$


Increasing / decreasing sequences

(3) Consider the sequences in question 1.


- (a) Decide whether they are increasing, decreasing or neither. Computer plots of sections of the sequence will help with this.




The plot of the beginning of $\{a_n\}$ is




The plot of the beginning of $\{b_n\}$ is




The plot of the beginning of $\{c_n\}$ is




The plot of the beginning of $\{d_n\}$ is




This $\{e_n\}$ (the Fibonacci sequence) begins positive and after the second term each term comes from adding a positive quantity to the previous term – so it is clearly increasing after the second term. The plot of $\{e_n\}$ looks like




The factorial form of f_n clearly shows that the sequence will be decreasing after the first couple of terms. The plot looks like



The plot of the beginning of $\{g_n\}$ looks like



The plot of the beginning of $\{h_n\}$ looks like



- (b) For those that are increasing or decreasing, try and construct a clear argument (proof) for your conclusion using the ratio and/or difference of consecutive terms.

The sequences a , b , $|c|$, $|d|$, h can all be shown to be decreasing/increasing as appropriate by examining the ratio of consecutive terms and showing it is always less than or greater than 1 as appropriate. An equivalent approach could be taken by considering the difference of consecutive terms and comparing to 0. Let's look at the sequence a as an example:

$$\begin{aligned} \frac{a_{n+1}}{a_n} &= \frac{3n+3}{(n+1)^2+1} \cdot \frac{n^2+1}{3n} \\ &= \frac{3n^3+3n^2+3n+3}{3n^3+6n^2+6n} \\ &= 1 + \frac{-3n^2-3n+3}{3n^3+6n^2+6n} \\ &< 1, \text{ for all } n \geq 1. \end{aligned}$$

The last inequality follows because the numerator in the fraction term is clearly negative and denominator positive, for $n \geq 1$. Then from the overall inequality above, $a_{n+1}/a_n < 1$, we get $a_{n+1} < a_n$ as required, since all the a_n terms are positive.



For the sequences e , f and g we can give more direct arguments based on the initial term(s) and the nature of the recurrence relation.

Convergence definition for sequences

- (4) How well do you understand the definition of convergence for a sequence?

- (a) Can you quote it accurately from memory?
- (b) Can you draw a suitable diagram illustrating sequence convergence and attach the parameters from the definition to it in the correct way?
- (c) Don't be too attached to the symbols x_n, N, ϵ used in the definition in the notes. Can you reliably quote the definition using entirely different symbols?

These are discussion questions for the students and class. The type of diagrams I have in mind though for part (b) are like the following



or

- (5) Consider the sequence $\{a_n\}_{n=1}^{\infty}$, where $a_n = \frac{1}{n^2}$. Does this have a limit and if so prove this by showing that the definition of convergence is satisfied.

In the sequence of terms $1/n^2$ the denominator is clearly increasing and unbounded as $n \rightarrow \infty$. Therefore the term will be decreasing and convergent with limit 0. To prove this formally by the definition of convergence we proceed as follows.

Let $\epsilon > 0$ be given. We set N to be equal to the smallest integer which is greater than or equal to $1/\sqrt{\epsilon}$. Then if $n > N$ then $1/n^2 < 1/(1/\sqrt{\epsilon})^2 = \epsilon$. This is the required inequality and so we conclude that $1/n^2 \rightarrow 0$ as $n \rightarrow \infty$.

- (6) Consider the following statements about a sequence $\{y_n\}_{n=1}^{\infty}$ which at first glance appear similar to the definition of sequence convergence.

S1: There exists a real value $\lambda > 0$ and a positive integer M such that for all $m > M$, $|y_m - L| < \lambda$.

This statement says that for a quantity L there is a fixed distance λ and a point y_M in the sequence after which the term y_m is always within a distance λ of L .

Considering this for a while we see that this implies that the sequence $\{y_n\}$ is bounded and moreover any bounded sequence will satisfy this statement S1 for any real quantity L whatsoever. For if a sequence is bounded then it is always within a set distance of zero, and hence of any other number.

So the sequences that satisfy S1 are exactly the bounded sequences.

S2: There exists an integer $M > 0$ such that for all real values $\lambda > 0$, if $m > M$ then $|y_m - L| < \lambda$.

This statement says that after some point y_M , the sequence terms are within the distance λ of this quantity L , and that this is true for **any** real $\lambda > 0$. How can sequence terms be arbitrarily close to some quantity L ? This is only possible if all these sequence elements are actually equal to L .

So the statement S2 is true, only of the sequences which are *eventually constant*, i.e. after some point all the sequence elements are equal to this quantity L .

For each of these statements can you give some examples of sequences (and values L) that satisfy them? For each statement can you characterize all the sequences that satisfy them?

Applying convergence results

(7) Determine the limit of the following convergent sequences. You might need the algebra of limits theorem and other results. Your solutions should explain clearly the reasoning used.

Parts (a) - (e) can be solved by applying the algebra of limits theorem. The proofs are summarised below but should be supplemented with some commentary as well.

(a) $\{a_n\}_{n=0}^{\infty}$, where $a_n = \frac{3n}{n^2+1}$.

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{3/n}{1 + 1/n^2} = \frac{0}{1+0} = 0.$$

(b) $\{b_n\}_{n=1}^{\infty}$, where $b_n = \frac{2^n}{2+2^n}$.

$$\lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \frac{1}{1/2^{n-1} + 1} = \frac{1}{0+1} = 1.$$

(c) $\{c_n\}_{n=1}^{\infty}$, where $c_n = \frac{5n^2-2}{3n^2-n}$.

$$\lim_{n \rightarrow \infty} c_n = \lim_{n \rightarrow \infty} \frac{5 - 2/n^2}{3 - 1/n} = \frac{5-0}{3-0} = 5/3.$$

(d) $\{d_n\}_{n=1}^{\infty}$, where $d_n = \frac{2n}{3n-1} - \frac{n^2+1}{2-n^2}$.

$$\lim_{n \rightarrow \infty} d_n = \lim_{n \rightarrow \infty} \left(\frac{2}{3 - 1/n} \right) - \lim_{n \rightarrow \infty} \left(\frac{1 + 1/n^2}{2/n^2 - 1} \right) = \frac{2}{3-0} - \frac{1+0}{0-1} = 5/3.$$

(e) $\{e_n\}_{n=1}^{\infty}$, where $e_n = \frac{n^3}{n^3+1}$.

$$\lim_{n \rightarrow \infty} e_n = \lim_{n \rightarrow \infty} \frac{1}{1 + 1/n^3} = \frac{1}{1+0} = 1.$$

(f) $\{f_n\}_{n=1}^{\infty}$, where $f_n = \frac{\cos(3n)}{n^{1/3}+1}$.

This questions requires the use of the Squeeze Theorem (not in the notes, but I will present it in the lectures) First we use the boundedness of the cosine function to say

$$\frac{-1}{n^{1/3}+1} \leq f_n \leq \frac{1}{n^{1/3}+1}.$$

Then we notice that the sequences at the ends of this inequality both converge to zero as the denominators are increasing and unbounded. Therefore by the Squeeze Theorem $f_n \rightarrow 0$ as $n \rightarrow \infty$.

Confirm your results by comparing to computer plots of the sequences.

(8) Consider the following sequence (which continues in the suggested way)

$$\sqrt{2}, \sqrt{2\sqrt{2}}, \sqrt{2\sqrt{2\sqrt{2}}}, \sqrt{2\sqrt{2\sqrt{2\sqrt{2}}}}, \dots$$

(a) Formulate this as a sequence defined by a recurrence relation.

This is the sequence $\{a_n\}_1^{\infty}$ where $a_1 = \sqrt{2}$ and for all $n \geq 1$, $a_{n+1} = \sqrt{2a_n}$.

- (b) With the aid of the Monotone Convergence Theorem prove that the sequence is convergent.

First we establish the monotonicity of the sequence. Numerical investigations suggest the sequence is increasing. Certainly $a_1 < 2$ and thereafter $a_n = \sqrt{a_n}\sqrt{a_n} < \sqrt{2}\sqrt{a_n} = a_{n+1}$ so the sequence is always increasing. (This is really a quick proof by induction. You might mention as much although the students will not necessarily have covered it yet).

Next we establish that the sequence is bounded. Numerical investigation suggests increasing convergence to 2, so 2 seems a likely upper bound. Certainly $a_1 < 2$, and thereafter $a_{n+1} = \sqrt{2}\sqrt{a_n} < \sqrt{2}\sqrt{2} = 2$. (Again, this is really proof by induction.)

So the sequence is increasing and bounded above, therefore it is convergent by the Monotone Convergence Theorem.

- (c) Determine the limit.

Since we now know the limit exists let us denote it by a . Then by the algebra of limits theorem the limit a satisfies

$$\begin{aligned} a &= \sqrt{2a} \\ \Leftrightarrow a &= 2. \end{aligned}$$

- (9) Consider the sequence $\{a_n\}_{n=0}^{\infty}$ defined by the recurrence relation

$$a_0 = 1, \quad \text{for all } n \geq 1, \quad a_n = 3 - \frac{1}{a_{n-1}}.$$

Prove that this sequence is increasing, bounded and hence convergent. Determine its limit.

Firstly, all sequence elements are bounded below by 1, as $a_0 = 1$ and thereafter $a_n = 3 - 1/a_{n-1} \geq 3 - 1 = 2$. Hence all the elements a_n are positive and so sequence is bounded above by 3.

Next we prove that the sequence is increasing, again taking an inductive approach. By inspection we see that $a_2 > a_1$, then if we assume that $a_m > a_{m-1}$ and argue from here as follows

$$\begin{aligned} a_m > a_{m-1} > 0 &\Rightarrow 1/a_m < 1/a_{m-1} \\ &\Rightarrow -1/a_m > -1/a_{m-1} \\ &\Rightarrow 3 - 1/a_m > 3 - 1/a_{m-1} \\ &\Rightarrow a_{m+1} > a_m, \end{aligned}$$

and hence the sequence is always increasing.

So by the Monotone Convergence Theorem the sequence is convergent. If we let $a = \lim_{n \rightarrow \infty} a_n$ and apply the algebra of limits theorem to the recurrence relation we get

$$a = 3 - 1/a.$$

This is equivalent to

$$a^2 - 3a + 1 = 0.$$

This has the solutions

$$a = -\frac{1}{2}\sqrt{5} + \frac{3}{2}, \quad a = \frac{1}{2}\sqrt{5} + \frac{3}{2}.$$

But only the second one of these is in the range $1 \leq a < 3$ so we conclude that the limit of the sequence is

$$a = \frac{1}{2}\sqrt{5} + \frac{3}{2} \approx 2.62.$$

- (10) Consider the sequence $\{x_n\}_{n=0}^{\infty}$ defined by the recurrence relation

$$x_0 = 2, \quad \text{for all } n \geq 1, \quad x_n = \frac{1}{3 - x_{n-1}}.$$

Is this sequence convergent? If so, determine the limit.

Can be solved using similar arguments to the previous two questions. Numerical investigation suggests a decreasing convergent sequence with limit of approximately 1/4.

- establish that the sequence elements are bounded above by 2 and below by 0.
- prove decreasing nature
- conclude limit exists, by MCT.
- then solve the equation $x = 1/(3 - x)$ and identify which solution is the limit.

- (11) Analyze the convergence/divergence of the sequence $\{y_n\}_{n=0}^{\infty}$ defined by $y_n = nr^n$, where $r \in \mathbb{R}$. Your conclusion should say it diverges for these values of r , converges for those, and preferably determine the limit when it exists.

We need separate arguments for various cases of $r \in \mathbb{R}$.

If $r = 0$ then for all $n \geq 1$, $y_n = 0$ and the sequence is trivially convergent.

If $|r| \geq 1$ then $|y_n|$ is increasing and unbounded and so these sequences are divergent.

If $0 < |r| < 1$ then the sequence is eventually decreasing since the ratio of consecutive terms satisfies

$$\frac{|y_{n+1}|}{|y_n|} = \left(1 + \frac{1}{n}\right) |r|.$$

Now the $1 + 1/n$ factor is decreasing and converging to 1, so eventually it will pass below the value $1/|r|$ and from then on we have $|y_{n+1}| < |y_n|$. So the sequence is decreasing in absolute value and has absolute value bounded below by 0. So by the MCT the sequence is converging.

Let $y = \lim_{n \rightarrow \infty} y_n$. There is a recurrence type relation satisfied by the sequence which is

$$y_{n+1} = \frac{n+1}{n} r y_n = (1 + 1/n) r y_n,$$

and so by the algebra of limits theorem we have

$$y = r y,$$

which can only be true if $y = 0$.

More convergence results

- (12) Suppose that the sequence $\{x_n\}_{n=1}^{\infty}$ is bounded and that $y_n \rightarrow 0$ as $n \rightarrow \infty$. Prove that the product sequence $\{x_n y_n\}_{n=1}^{\infty}$ also converges to 0, i.e. $x_n y_n \rightarrow 0$ as $n \rightarrow \infty$.

Let M be a bound for the sequence $\{x_n\}$, i.e. for all n , $|x_n| \leq M$. Let $\epsilon > 0$ be given. Since y_n converges to 0 there will exist an N such that if $n \geq N$ then $|y_n| < \epsilon/M$. Then for all $n \geq N$ we have

$$\begin{aligned} |x_n y_n| &= |x_n| |y_n| \\ &\leq M |y_n| \\ &< M \epsilon / M = \epsilon. \end{aligned}$$

This proves that $x_n y_n \rightarrow 0$ as required.

- (13) Suppose that $\{n_j\}_{j=1}^{\infty} = n_1, n_2, n_3, \dots$ is a strictly increasing sequence of positive integers, i.e. for all $j \geq 1$, $n_{j+1} > n_j \geq 1$. Prove that if $a_m \rightarrow L$ as $m \rightarrow \infty$ then the *subsequence* $\{a_{n_j}\}_{j=1}^{\infty}$ also converges to the same limit L , i.e.

$$a_{n_j} \rightarrow L \text{ as } j \rightarrow \infty.$$

This result says that every subsequence of a convergent sequence is also convergent and tending to the same limit as the sequence itself. It's an example of something that kind of takes more work to state the result carefully than it does to prove it.

Simply put, for all $\epsilon > 0$ there exists a N_ϵ such that $n \geq N_\epsilon$ implies $|a_n - L| < \epsilon$. This is the definition of convergence for $a_n \rightarrow L$. The very same choice of N_ϵ will work for the subsequence as if $j \geq N_\epsilon$ then $n_j \geq N_\epsilon$ since $n_j \geq j$. Then $a_{n_j} \rightarrow L$ as required.

Estimating π

- (14) This question relates to the Buffon's needle activity in induction week where we carried out a needle dropping experiment to estimate the value of π . Here we will produce some better estimates of π using a precise geometric method based on the technique used by Archimedes to estimate π .

Consider figure 1 which shows a unit circle (i.e. with radius 1) together with two n -sided regular polygons. The inner polygon is the largest such one that fits inside the circle, i.e. its vertices lie on the circle, and the outer polygon is the smallest such one containing the circle, i.e. the midpoints of its sides lie on the circle. The angle formed at the centre of the circle by two adjacent polygon vertices is thus $2\pi/n$ as shown. The circle has area π . Let a_n denote the area of the inner n -sided polygon and b_n denote the area of the outer polygon. By studying a_n and b_n we will obtain a sequence of successively narrower intervals containing π .

- Make sure you appreciate why for every $n \geq 3$ we have $a_n < \pi < b_n$, and that as n increases we will obtain successively narrower intervals estimating π .
- Carefully consider the triangles making up the polygons and prove that

$$a_n = \frac{n}{2} \sin\left(\frac{2\pi}{n}\right) \quad \text{and} \quad b_n = n \tan\left(\frac{\pi}{n}\right).$$

- With the help of various trigonometric formulas derive two recurrence relations, one giving a_{2n} in terms of a_n , the other giving b_{2n} in terms of b_n . The recurrence relations you are aiming for are quadratic in nature, i.e. involving constants, the usual arithmetic operations




Figure 1: Unit circle with inscribed and circumscribed n -sided regular polygons, shown here with $n = 8$.

of addition/subtraction/multiplication/division and squaring and square roots.

The recurrence relation for a_n is

$$a_{2n} = n \sqrt{\frac{1}{2} \left(1 - \sqrt{1 - \left(\frac{2a_n}{n} \right)^2} \right)}.$$

- (d) By considering the inscribed and circumscribed squares prove the initial values $a_4 = 2$ and $b_4 = 4$ and then use your recurrence formulas to generate some interval estimates of π .
- (e) Code your recurrence formulas in a suitable programming language (Matlab, Sage, Python, Excel, ...) and see how accurately you can estimate π .



Suggestions for further work

You may get curious results as you compute a_{2^m} and b_{2^m} for larger and larger m , such as the sequence failing to consistently increase/decrease or failing to return a value at all. This is because you have run up against rounding errors due to the computer only being able to store numbers to a certain precision (i.e. number of digits accuracy). Increase the accuracy of your estimates by increasing the precision of the numbers in your computation. Some information available from:

Matlab – <http://uk.mathworks.com/help/symbolic/numbers-and-precision.html>

SageMath – http://doc.sagemath.org/pdf/en/reference/rings_numerical/rings_numerical.pdf

When possible you should work on the questions in advance and be prepared for discussions on them with your colleagues and tutor. Some questions are straightforward calculation while others require more discussion and thought. The computer icon  indicates where computer calculation or programming may be of use. Further questions can be found in the lecture notes as well as the recommended unit resources (see Moodle).

Series notation and defining formulas

- (1) Write down some initial terms of the following series

$$(i) \sum_{n=1}^{\infty} \frac{(-1)^n}{2n-1}, \quad (ii) \sum_{n=1}^{\infty} \frac{n+1}{n^3+2}, \quad (iii) \sum_{n=1}^{\infty} \frac{3}{n!}, \quad (iv) \sum_{n=1}^{\infty} \left(\sum_{i=1}^n i^{-2} \right).$$

$$\begin{aligned} (i) \sum_{n=1}^{\infty} \frac{(-1)^n}{2n-1} &= -1 + \frac{1}{3} - \frac{1}{5} + \frac{1}{7} - \frac{1}{9} + \dots \\ (ii) \sum_{n=1}^{\infty} \frac{n+1}{n^3+2} &= \frac{2}{3} + \frac{3}{10} + \frac{4}{29} + \frac{5}{66} + \frac{6}{127} + \dots \\ (iii) \sum_{n=1}^{\infty} \frac{3}{n!} &= 3 + \frac{3}{2} + \frac{1}{2} + \frac{1}{8} + \frac{1}{40} + \dots \\ (iv) \sum_{n=1}^{\infty} \left(\sum_{i=1}^n i^{-2} \right) &= 1 + \frac{5}{4} + \frac{49}{36} + \frac{205}{144} + \frac{5269}{3600} + \frac{5369}{3600} + \dots \end{aligned}$$

Series convergence definition

- (2) Make sure you can accurately quote the definition of convergence for a series.

The series $\sum_{n=1}^{\infty} a_n$ converges if and only if the associated sequence of partial sums, $\{S_k\}_{k=1}^{\infty}$ where $S_k = \sum_{n=1}^k a_n$, converges. If $S_k \rightarrow S$ as $k \rightarrow \infty$ then we write $S = \sum_{n=1}^{\infty} a_n$. Otherwise we say that $\sum_{n=1}^{\infty} a_n$ diverges.

- (3) Can you confidently give the proof that the harmonic series $\sum_{j=1}^{\infty} 1/j$ diverges?

See notes.

- (4) Can you prove the General Term Test using the definition of convergence?

Suppose that $S = \sum_{n=1}^{\infty} a_n$, i.e. $S_k = \sum_{n=1}^k a_n \rightarrow S$ as $k \rightarrow \infty$. Note that $a_k = S_k - S_{k-1}$. Let $\epsilon > 0$ be given. By the convergence of the sequence $\{S_k\}$ there exists a point K such that for all $k \geq K$ we have $|S_k - S| < \frac{\epsilon}{2}$. Then for every $k \geq K + 1$ we have

$$|a_k| = |S_k - S_{k-1}| = |S_k - S - (S_{k-1} - S)| \leq |S_k - S| + |S_{k-1} - S| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

This proves the convergence of a_k to 0 as $k \rightarrow \infty$.

It's a good idea to use a diagram of the number line here to explain the choice of interval radius $\frac{\epsilon}{2}$ and explain the idea behind this.

Geometric series test

- (5) Ensure that you can derive the partial sum formula for a general geometric series of the form

$$\sum_{n=1}^{\infty} ar^{n-1}.$$

1

Can you then prove the convergence/divergence of this series for all possible combinations of values of the initial term a and the common ratio r ?

Exploiting the structure of the terms of the geometric series we see that the partial sums S_k satisfy

$$S_k - rS_k = a - ar^k$$

which implies that for $r \neq 1$ we have

$$S_k = \frac{a(1 - r^k)}{1 - r}.$$

Using this we can decide the convergence status of the geometric series.

If $a = 0$ then all terms are zero and the series is trivially convergent. So we assume $a \neq 0$ in the following.

If $r = 1$ then the series is $\sum_{n=1}^{\infty} a = a + a + a + \dots$ which is clearly divergent as $a \neq 0$.

If $r = -1$ then the series is $\sum_{n=1}^{\infty} a(-1)^{n-1} = a - a + a - \dots$ which again is divergent as $a \neq 0$.

In all other cases we can make use of the partial sums formula to say that

$$S = \sum_{n=1}^{\infty} ar^{n-1} = \lim_{k \rightarrow \infty} S_k = \lim_{k \rightarrow \infty} \frac{a(1 - r^k)}{1 - r}.$$

If $|r| > 1$ then the term r^k is divergent and so the series is divergent also. If $|r| < 1$ then the term r^k is convergent to 0 and so the series is also convergent and

$$S = \sum_{n=1}^{\infty} ar^{n-1} = \lim_{k \rightarrow \infty} S_k = \lim_{k \rightarrow \infty} \frac{a(1 - r^k)}{1 - r} = \frac{a}{1 - r}.$$

- (6) Use the analysis of the ratio test to determine the convergence status and sum if possible, of the series

$$(i) \sum_{n=1}^{\infty} \frac{2}{3} \left(\frac{1}{3}\right)^{n+1}, \quad (ii) \sum_{n=1}^{\infty} 3 \left(\frac{1}{2}\right)^n, \quad (iii) \sum_{n=1}^{\infty} \frac{1}{5} \left(\frac{6}{5}\right)^{n+2}.$$

We just need to adjust the term in each case to match the general form of the geometric series used in the previous question and then apply the convergence criteria and formula to get

$$(i) \frac{1}{9}, \quad (ii) 3, \quad (iii) \text{divergent}.$$

- (7) Consider the number a with decimal expansion

$$a = 0.1234123412341234 \dots,$$

continuing in the suggested way. Use the geometric series summation formula to express a as a rational number, i.e. a ratio of integers.

We can express a in series form by

$$a = \sum_{n=1}^{\infty} \frac{1234}{10000^n}.$$

This is a convergent geometric series and applying the sum formula gives

$$a = \frac{1234}{9999}.$$

Application of the ratio test

- (8) Determine the convergence or divergence of the series

$$\sum_{m=0}^{\infty} \frac{2^m}{m!}.$$

Writing a_m for the series term the ratio of consecutive terms satisfies

$$\begin{aligned} \lim_{m \rightarrow \infty} \frac{a_{m+1}}{a_m} &= \lim_{m \rightarrow \infty} \left(\frac{2^{m+1}}{(m+1)!} \frac{m!}{2^m} \right) \\ &= \lim_{m \rightarrow \infty} \frac{2}{m+1} \\ &= 0. \end{aligned}$$

So by the ratio test the series $\sum a_m$ converges.

- (9) Determine the convergence or divergence of the series

$$\sum_{m=0}^{\infty} \frac{2^m}{(m+1)^2}.$$

Writing a_m for the series term the ratio of consecutive terms satisfies

$$\begin{aligned}\lim_{m \rightarrow \infty} \frac{a_{m+1}}{a_m} &= \lim_{m \rightarrow \infty} \left(\frac{2^{m+1}}{(m+2)^2} \frac{(m+1)^2}{2^m} \right) \\ &= \lim_{m \rightarrow \infty} 2 \left(\frac{m+1}{m+2} \right)^2 \\ &= 2 \lim_{m \rightarrow \infty} \left(1 - \frac{1}{m+2} \right)^2 \\ &= 2, \text{ by the algebra of limits theorem for seqs.}\end{aligned}$$

So by the ratio test the series $\sum a_m$ diverges.

- (10) Does the ratio test give a conclusion about the series

$$\sum_{n=1}^{\infty} \frac{n}{n+1}?$$

If not, determine the convergence/divergence of this series by more direct means.

Writing a_n for the series term the ratio of consecutive terms satisfies

$$\begin{aligned}\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} &= \lim_{n \rightarrow \infty} \left(\frac{n+1}{n+2} \frac{n+1}{n} \right) \\ &= \lim_{n \rightarrow \infty} \frac{n^2 + 2n + 1}{n^2 + 2n} \\ &= \lim_{n \rightarrow \infty} \frac{1 + 2/n + 1/n^2}{1 + 2/n} \\ &= 1.\end{aligned}$$

So the ratio test is inconclusive. Looking at the series again we notice that the terms do not converge to zero, in fact

$$\lim_{n \rightarrow \infty} a_n = 1.$$

So the series does not pass the General Term Test and so the series is divergent.

Application of comparison tests

I tend to prefer the application of the limit comparison test, but anything doable with the limit comparison test can normally be achieved with an equivalent use of the standard comparison test.

- (11) Use a suitable comparison test to determine the convergence status of the series

$$\sum_{k=1}^{\infty} \frac{k^2 + 2}{k(k^2 + 5)}.$$

The dominant behaviour of the term here is k^2 in the numerator and k^3 in the denominator so the we should compare to the term $1/k$ which is the term of a divergent series.

Writing a_k for the series term in the question and examining the limit of the ratio we get

$$\begin{aligned}\lim_{k \rightarrow \infty} \frac{a_k}{1/k} &= \lim_{k \rightarrow \infty} \left(\frac{k^3 + 2k}{k^3 + 5k} \right) \\ &= 1\end{aligned}$$

So the series in question compares with the divergent harmonic series and so by the limit comparison test the series is also divergent.

- (12) Use a suitable comparison test to determine the convergence status of the series

$$\sum_{k=1}^{\infty} \frac{k+3}{k(k^2+2)}.$$

The dominant behaviour of the term here is k^1 in the numerator and k^3 in the denominator so the we should compare to the term $1/k^2$ which is the term of a convergent series.

Writing a_k for the series term in the question and examining the limit of the ratio we get

$$\begin{aligned}\lim_{k \rightarrow \infty} \frac{a_k}{1/k^2} &= \lim_{k \rightarrow \infty} \left(\frac{k^3 + 3k^2}{k^3 + 2k} \right) \\ &= 1\end{aligned}$$

So the series in question compares with a convergent series and so by the limit comparison test the series is also convergent.

Application of partial fraction expansions

- (13) Consider the series $\sum_{n=1}^{\infty} a_n$ where the term is defined by

$$a_n = \frac{9n^2 + 33n + 12}{(n+4)(n+3)(n+1)n}.$$

Obtain the partial fraction expansion of a_n and use this to obtain the sum of the series.

The partial fraction expansion of a_n is $-\frac{2}{n+4} - \frac{1}{n+3} + \frac{2}{n+1} + \frac{1}{n}$ and the sum of the series is 4.

- (14) Consider the series $\sum_{n=1}^{\infty} b_n$ where the term is defined by

$$b_n = \frac{8}{n^2 + 6n + 5}.$$

Obtain the partial fraction expansion of b_n and use this to obtain the sum of the series.

The partial fraction expansion of b_n is $-\frac{2}{n+5} + \frac{2}{n+1}$ and the sum of the series is $\frac{77}{30}$.

- (15) Consider the series $\sum_{n=2}^{\infty} c_n$ where the term is defined by

$$c_n = \frac{9}{n^2 + n - 2}.$$

Obtain the partial fraction expansion of c_n and use this to obtain the sum of the series. (n.b. take note of the starting value of the index n)

The partial fraction expansion of c_n is $-\frac{3}{n+2} + \frac{3}{n-1}$ and the sum of the series is $\frac{11}{2}$.

- (16) Consider the series $\sum_{n=1}^{\infty} d_n$ where the term is defined by

$$d_n = \frac{11-n}{n^2 + 6n + 5}.$$

Obtain the partial fraction expansion of d_n and use this to decide whether the series is convergent or divergent.

The partial fraction expansion of d_n is $-\frac{4}{n+5} + \frac{3}{n+1}$. Because the numerators are different we do not get complete cancellation. Considering the partial sum S_k we see that

$$S_k = \frac{3}{2} + \frac{3}{3} + \frac{3}{4} + \frac{3}{5} - \left(\sum_{n=6}^{k+1} \frac{1}{n} \right) - \frac{4}{k+2} - \frac{4}{k+3} - \frac{4}{k+4} - \frac{4}{k+5}.$$

We see that S_k contains, essentially, the partial sum of the harmonic series. Since the harmonic series diverges then so does the sequence S_k and hence the series in question is divergent.

Alternatively we can apply the limit comparison test to this series, comparing with the harmonic series, and this will also show that it is divergent.

Computer work



- (17) You should ensure that you are able to use Matlab to build an array containing a specified number of initial terms of a series.
- (18) You should then be able to build an array containing the partial sums of that series and plot them as visual aid to determining the long term behaviour of the series.
- (19) Investigate the capabilities of Matlab's `symsum` command from its symbolic toolbox. See the information at <http://uk.mathworks.com/help/symbolic/symsum.html>.
- (20) Apply these techniques to the series mentioned in this problem sheet, the questions in the lecture notes, etc
- (21) Investigate the capabilities of other computer systems, such as SageMath in this regard. Register for a free account at <https://cocalc.com> and check out Gregory Bard's book *Sage for Undergraduates* available from <http://www.gregorybard.com/Sage.html>.