$$\begin{tikzcd}
\bullet_{0} \arrow[r] & \bullet_{0} & \bullet_{\ker A_3} \arrow[l, "0"'] \\
 & \bullet_{0} \arrow[u] &
\end{tikzcd}.$$

These representations are multiples of the indecomposable objects

$$\begin{tikzcd}
\bullet_{1} \arrow[r, "0"] & \bullet_{0} & \bullet_{0} \arrow[l] \\
 & \bullet_{0} \arrow[u] &
\end{tikzcd},$$

$$\begin{tikzcd}
\bullet_{0} \arrow[r] & \bullet_{0} & \bullet_{0} \arrow[l] \\
 & \bullet_{1} \arrow[u, "0"] &
\end{tikzcd},$$

$$\begin{tikzcd}
\bullet_{0} \arrow[r] & \bullet_{0} & \bullet_{1} \arrow[l, "0"'] \\
 & \bullet_{0} \arrow[u] &
\end{tikzcd}.$$

So we get to a situation where all of the maps A_1, A_2, A_3 are injective:

$$\begin{tikzcd}
\bullet_{V_1} \arrow[r, hook, "A_1"] & \bullet_{V} & \bullet_{V_3} \arrow[l, hook', "A_3"'] \\
 & \bullet_{V_2} \arrow[u, hook, "A_2"] &
\end{tikzcd}.$$

As in Example 6.2.4, we can then identify the spaces V_1, V_2, V_3 with subspaces of V. So we get to the **triple of subspaces problem** of classifying triples of subspaces of a given space V.