Theory and Methods in Causal Inference

Lecture Notes

Jae Kwang Kim

Department of Statistics

Iowa State University

STAT 5900B

Last updated: April 11, 2026

Contents

Preface **1**
Organization . . . 1
Notation . . . 1

1 Introduction **3**
1.1 Motivation: Why Causal Inference Is Hard . . . 3
1.1.1 The Central Question . . . 3
1.2 The Causal Trinity: Three Languages for One Idea . . . 4
1.2.1 Language 1: Structural Equation Models . . . 4
1.2.2 Language 2: Directed Acyclic Graphs . . . 5
1.2.3 Language 3: Potential Outcomes . . . 6
1.2.4 Roadmap: How the Trinity Organizes These Notes . . . 6
1.2.5 Historical Roots of the Causal Trinity . . . 6
1.3 The Core Distinction . . . 7
1.3.1 Two Scenarios: Observed vs. Unobserved Confounder . . . 8
1.4 The Gaussian Linear Confounded Model . . . 8
1.4.1 Setup . . . 8
1.4.2 Two Explicit Densities . . . 9
1.4.3 The Endogeneity Gap . . . 9
1.4.4 Lab: Simulating the Two Densities . . . 10
1.4.5 The IV DAG Versus the Mediation DAG . . . 10
1.5 The Two-Step Paradigm . . . 12
1.5.1 Step 1: Identification . . . 12
1.5.2 Step 2: Estimation . . . 12
1.5.3 A Worked Example: Flu Vaccination and Infection . . . 12
1.6 Summary . . . 14
1.7 Problems . . . 14

2 DAGs and d-Separation **17**
2.1 Directed Acyclic Graphs . . . 17
2.1.1 Basic Definition . . . 17
2.1.2 Structural Relationships . . . 18
2.2 Paths, Blocking, and d-Separation . . . 19
2.2.1 Path Structures . . . 19
2.2.2 Blocking Rules . . . 19
2.2.3 The d-Separation Criterion . . . 20
2.2.4 Practical Ways to Check d-Separation . . . 21
2.3 Collider Bias and the IV DAG . . . 22
2.3.1 Berkson's Bias . . . 22
2.3.2 Full d-Separation Analysis of the IV DAG . . . 23
2.4 The Markov Property and Factorization . . . 23
2.5 Worked Example: The Education–Earnings DAG . . . 24
2.6 The Big Picture . . . 25
2.7 Summary . . . 25
2.8 Problems . . . 26

3 The Do-Calculus and Identification Criteria **29**
3.1 From d-Separation to Intervention: Intervention Graphs . . . 29
3.1.1 What Does Intervention Mean? . . . 29
3.1.2 The Two Graph Operations . . . 30
3.2 The Back-Door Criterion . . . 31
3.2.1 Motivation: When the Confounder Is Unobserved . . . 31
3.2.2 Definition and Theorem . . . 31
3.2.3 The Adjustment Formula: Standardization . . . 33
3.3 The Front-Door Criterion . . . 34
3.3.1 The Configuration and Criterion . . . 34
3.3.2 The Front-Door Formula . . . 35
3.4 The Three Rules of Do-Calculus . . . 36
3.4.1 Statement of the Rules . . . 36
3.4.2 Intuition for Each Rule . . . 37
3.4.3 Proofs of the Identification Formulas . . . 38
3.5 The Do-Calculus is Complete . . . 39
3.6 The Big Picture . . . 40
3.7 Summary . . . 40
3.8 Problems . . . 42

References **45**

Appendices **47**

A Update Log **47**
A.1 April 2026 . . . 47
A.1.1 April 11, 2026 . . . 47
A.1.2 April 9, 2026 . . . 47

Preface

These lecture notes accompany the graduate course **STAT 5900B: Theory and Methods in Causal Inference** at the Department of Statistics, Iowa State University.

The course targets graduate students in statistics. It assumes familiarity with probability theory, mathematical statistics, and linear models at the level of a graduate sequence, but no prior exposure to causal inference.

Download full PDF

Organization

The notes are organized in four parts.

Part I — Foundations (Chapters 1–4) builds the graphical and conceptual foundation: the causal trinity (SEM, DAG, potential outcomes), DAGs and d-separation, the do-calculus and identification criteria, and the connection between potential outcomes and the do-calculus.

Part II — Identification (Chapters 5–8) bridges the graphical framework and observed data: randomization and back-door adjustment, propensity score methods, instrumental variables, and mediation and front-door identification.

Part III — Estimation (Chapters 9–12) covers semiparametric efficiency theory: estimating equations and influence functions, doubly robust estimation, flexible nuisance estimation with cross-fitting, and estimation under instrumental variables.

Part IV — Policy Learning and Sequential Decision Making (Chapters 13–15) covers policy evaluation, optimal policy learning, and dynamic treatment regimes.

Notation

Throughout these notes:

- $\mathrm{do}(T = t)$ denotes the do-operator (intervention).
- $Y(t)$ denotes the potential outcome under $T = t$.
- $\mathbb{E}[\,\cdot\,]$ denotes expectation; $\perp\!\!\!\perp$ denotes conditional independence.
- $\mathcal{G}$ denotes a DAG; $\mathcal{G}_T$ denotes the mutilated graph after intervening on T.

Chapter 1

Introduction

Learning Objectives

By the end of this chapter, students should be able to:

1. Articulate the distinction between an interventional distribution $f(y \mid \mathrm{do}(T{=}t))$ and an observational conditional distribution $f(y \mid T{=}t)$, and explain why they differ whenever T is endogenous.
2. Describe the same causal question in all three languages — SEM, DAG graph surgery, and potential outcomes — and begin to translate between them.
3. Derive the two densities explicitly in the Gaussian linear confounded model, and identify the endogeneity bias $\alpha\gamma\sigma_U^2/\sigma_T^2$ as the gap between them.
4. State the two-step paradigm (identification then estimation) and locate each step in the causal inference pipeline.
5. Distinguish the IV DAG from the mediation DAG and explain why conflating an instrument with a mediator leads to identification failure.

This chapter is an orientation, not a complete treatment. It introduces four ideas — the interventional/observational distinction, the causal trinity, the Gaussian confounded model as a running example, and the identification-versus-estimation paradigm — that will be developed rigorously in Chapters 2–4 and beyond. After one reading, students should have a map of the territory; the details will become clear as the course proceeds.

1.1 Motivation: Why Causal Inference Is Hard

Causal inference studies when and how the observational distribution can be used to recover the distribution that would arise under interventions.

1.1.1 The Central Question

Causal inference is concerned with a deceptively simple question: what would happen to Y if we were to *set* $T = t$? This is fundamentally different from the question that standard statistical procedures answer: what is the distribution of Y among units *observed* to have $T = t$?

Definition: Interventional vs. Observational Distribution

- The *interventional distribution* $f(y \mid \mathrm{do}(T{=}t))$ is the distribution of Y in a hypothetical world where T has been *externally set* to t, regardless of the usual data-generating mechanism for T.
- The *observational conditional distribution* $f(y \mid T{=}t)$ is the distribution of Y among the subpopulation of units for whom $T = t$ was *observed* to hold.

These two distributions coincide only when T is randomized (exogenous). In all other cases they differ, and the difference is *endogeneity bias*.

Definition: Exogeneity and Endogeneity

Suppose the outcome is generated by the structural equation

$$Y = f_Y(T, X, U),$$

where T is the treatment, X are observed covariates, and U represents unobserved common causes of T and Y (confounders).

- The treatment T is *exogenous given* X if $T \perp\!\!\!\perp U \mid X$.
- The treatment T is *endogenous given* X if $T \not\perp\!\!\!\perp U \mid X$, so that some unobserved variable U affects both T and Y.

Example: Drug Trial with Unmeasured Severity

Does a new drug ($T = 1$) reduce blood pressure (Y)? Suppose sicker patients are more likely to take the drug, so that unmeasured severity U is a common cause of T and Y. Then:

- *Observational study*: $\mathbb{E}[Y \mid T{=}1] > \mathbb{E}[Y \mid T{=}0]$. Drug users have higher blood pressure on average. A naïve analyst concludes the drug is harmful.
- *Randomized trial*: $\mathbb{E}[Y \mid \mathrm{do}(T{=}1)] < \mathbb{E}[Y \mid \mathrm{do}(T{=}0)]$. Forcing everyone to take the drug reduces blood pressure. The drug is beneficial.

Same data. Different questions. Opposite answers. The entire discrepancy is caused by the back-door path $T \leftarrow U \rightarrow Y$, which the observational quantity $\mathbb{E}[Y \mid T{=}1]$ cannot remove.

The Fundamental Problem of Causal Inference

We observe $f(y \mid T{=}t)$ directly from data. We want $f(y \mid \mathrm{do}(T{=}t))$. The goal of *identification theory* is to find conditions under which the former can be used to recover the latter.

1.2 The Causal Trinity: Three Languages for One Idea

The same causal question — what happens when we set $T = t$? — can be expressed in three distinct but equivalent languages. Collectively, we call these the **causal trinity**: the *structural equation model* (SEM), the *directed acyclic graph* (DAG) with graph surgery, and the *potential outcomes* framework. Each member of the trinity illuminates a different facet of the same underlying idea, and fluency in all three is essential for modern causal reasoning. Figure 1.1 depicts their relationships.

1.2.1 Language 1: Structural Equation Models

A structural equation model (SEM) represents the causal data-generating mechanism as a system of equations, one per variable, together with a joint distribution over exogenous errors.

Definition: Structural Equation Model (SEM)

A recursive SEM over variables (U, T, Y) is a system

$$U \sim p(u), \qquad T = f_T(U, Z, \delta), \qquad Y = f_Y(T, U, \varepsilon),$$

where δ and ε are mutually independent exogenous disturbances, Z is an observed instrument, and U is an unobserved confounder.

Remark: Error Independence in the NPSEM-IE

The assumption that the disturbances δ and ε are *mutually independent* is a substantive restriction, not merely a technical convenience. Because U is unobserved and enters both f_T and f_Y, error independence does not remove the dependence between T and $Y(t)$ induced by U; it restricts only the additional channel of dependence running through the equation-specific errors (δ, ε). Richardson



Figure 1.1: The **causal trinity**: three languages that express the same causal question from different vantage points. Potential outcomes define *what* we want to estimate; the DAG with graph surgery specifies *how* to identify it from observed data; the SEM provides the *generative mechanism* for likelihood construction. Double-headed arrows show the translation assumptions connecting adjacent languages. All three converge on the same central question (ellipse).

> and Robins (2014) call SEMs with this structure *Nonparametric Structural Equation Models with Independent Errors* (NPSEM-IEs).
> These notes adopt the NPSEM-IE framework throughout for two practical reasons: error independence ensures the DAG's Markov condition holds, making the Markov factorization of the joint density immediate; and it integrates cleanly with the graphical identification theory developed in later chapters.

Observational regime. T is determined by f_T. Because U appears in both f_T and f_Y, we have $\mathrm{Cov}(T, \varepsilon) \neq 0$ in general: this is endogeneity.

Interventional regime. The do-operator replaces the equation for T with a constant t, yielding the *mutilated system*:

$$U \sim p(u), \qquad T = t \text{ (fixed)}, \qquad Y = f_Y(t, U, \varepsilon).$$

Now T and U are independent by construction: the back-door path through U has been severed.

1.2.2 Language 2: Directed Acyclic Graphs

A DAG makes the same surgery graphically visible.

> Definition: Graph Surgery
>
> In the observational DAG, arrows into T encode the mechanisms that determine T (including back-door paths through unobserved variables). Under do(T=t), all arrows into T are *deleted*. The resulting mutilated graph $\mathcal{G}_T$ has no back-door paths from T to Y: the causal effect flows only along the direct path $T \to Y$.

The observational and mutilated DAGs for the Gaussian confounded model are shown side by side in Figure 1.3 in Section 1.4.

1.2.3 Language 3: Potential Outcomes

Definition: Potential Outcome

(Rubin 1974) $Y(t)$ is the outcome that *would have been observed* had T been set to t, possibly contrary to fact.
SUTVA (stable unit treatment value assumption): $Y_i = Y_i(T_i)$, i.e., the observed outcome for unit i equals the potential outcome at the actual treatment received. This rules out interference between units and multiple versions of treatment.

Connection to the do-operator. Under standard structural assumptions (consistency + no interference), the potential outcome $Y(t)$ and the interventional distribution are linked by

$$Y(t) \;\sim\; f(y \mid \mathrm{do}(T{=}t)).$$

Remark

Potential outcomes define what we want to estimate (e.g., $\mathbb{E}[Y(1)] - \mathbb{E}[Y(0)]$), but they do not specify how to write an observed-data likelihood for estimation. The do-operator fills this gap by providing an explicit link between the interventional density and the observational distribution via identification rules (back-door, front-door, do-calculus).

1.2.4 Roadmap: How the Trinity Organizes These Notes

The three languages are not used equally throughout the course.

Part I — Foundations: the language of DAGs. The DAG and the do-operator are the primary language here because they make the interventional/observational distinction *syntactically explicit*: confounding is visible as an open back-door path, and identifying assumptions are visible as graph criteria.

Part II — Identification: bridging DAGs and observed data. Ignorability ($Y(t) \perp\!\!\!\perp T \mid X$) is the potential-outcomes counterpart of the back-door criterion; the propensity score $\pi(X) = P(T{=}1 \mid X)$ operationalizes it in observed data.

Part III — Estimation: semiparametric and machine-learning methods. The semiparametric efficiency framework — efficient influence functions, doubly robust estimators, double machine learning — is largely language-neutral: it works with whatever moment condition the identification step produces.

1.2.5 Historical Roots of the Causal Trinity

The three languages grew independently in different disciplines over more than a century, driven by different scientific questions and different communities. Understanding this history explains why the three schools sometimes use incompatible terminology for the same idea.

Structural Equation Models: genetics and econometrics. The oldest strand begins with the biologist Sewall Wright (1921), who introduced *path analysis* to decompose correlations among hereditary traits into direct and indirect causal contributions. The framework was transplanted into economics by Haavelmo (1943), who argued that economic relationships should be modeled as autonomous structural equations robust to policy interventions. The Cowles Commission formalized simultaneous equation systems throughout the 1940s–50s. Lucas's (1976) critique — that reduced-form equations break down when policy changes the data-generating process — is essentially a statement of the interventional/observational distinction, decades before that phrase existed.

Potential Outcomes: experimental statistics and epidemiology. Neyman (1923) introduced the notation $Y_i(t)$ in the context of randomized agricultural experiments. Fisher (1935) placed randomization at the foundation of causal inference. The crucial extension to *observational* studies was made by Rubin (1974), who formalized the assignment mechanism and stated SUTVA explicitly. Robins (1986) extended potential outcomes to longitudinal treatments and time-varying confounding via g-computation. Imbens and Angrist (1994) connected the framework to instrumental variables and defined the LATE.

DAGs and do-calculus: computer science and philosophy. Pearl developed Bayesian networks through the 1980s as a representation for probabilistic reasoning in AI. The pivotal step came in Pearl (1995), where the *do-operator* was introduced as a formal syntax for distinguishing interventional from observational distributions within a graph-based algebra. The 2000 monograph *Causality* synthesised DAGs, the do-calculus, identification theory, and the connections to potential outcomes into a unified framework.

Convergence. By the 2000s it became clear that all three frameworks are, under standard assumptions, *logically equivalent*: any result expressible in one can be translated into the others. The causal trinity is therefore not three competing theories but three dialects of the same language, each optimized for a different task. Figure 1.2 summarizes the key milestones.



Figure 1.2: Key milestones in the three schools of causal inference. SEM (top, blue) originates in genetics and econometrics; PO (middle, purple) in experimental statistics; DAG (bottom, green) in computer science and philosophy. The three developed largely in isolation until converging in the 2000s.

1.3 The Core Distinction

The single most important inequality of this course is:

$$\underbrace{f(y \mid x, \mathrm{do}(T{=}t))}_{\text{interventional}} \neq \underbrace{f(y \mid x, T{=}t)}_{\text{observational}} \qquad \text{whenever } T \text{ is endogenous.} \tag{1.1}$$

The key assumption that connects the two densities is *conditional exchangeability*.

Definition: Conditional Exchangeability

Given observed covariates X and confounders U, the treatment assignment is *conditionally exchangeable with the intervention* if

$$Y(t) \perp\!\!\!\perp T \mid X, U,$$

or equivalently,

$$f(y \mid x, T{=}t, U{=}u) \;=\; f(y \mid x, \mathrm{do}(T{=}t), U{=}u).$$

This says that once we condition on all common causes (X, U) of T and Y, observing $T{=}t$ and intervening to set $T{=}t$ produce the same distribution of Y: there is no residual confounding given (X, U).

Under conditional exchangeability, the interventional density has a direct representation in terms of observable quantities.

Proposition: Back-Door Adjustment Formula

Under conditional exchangeability (Definition above),

$$f(y \mid x, \mathrm{do}(T{=}t)) \;=\; \int f(y \mid x, T{=}t, U{=}u)\, p(u \mid x)\, du. \tag{1.2}$$

Proof

Step (i): weight under the intervention. Under $\mathrm{do}(T{=}t)$, the arrow $U \to T$ is severed in the mutilated graph $\mathcal{G}_T$, so $U \perp\!\!\!\perp T \mid X$ under the intervention. The law of total probability therefore gives:

$$f(y \mid x,\, \mathrm{do}(T{=}t)) = \int f(y \mid x,\, \mathrm{do}(T{=}t),\, U{=}u)\; p(u \mid x)\; du.$$

Step (ii): replacing the interventional kernel. By conditional exchangeability, $f(y \mid x,\, \mathrm{do}(T{=}t),\, U{=}u) = f(y \mid x,\, T{=}t,\, U{=}u)$. Substituting completes the proof. □

Remark: Confounding Discrepancy

What OLS estimates is the observational density $f(y \mid x, T{=}t)$, not the interventional density. The gap

$$f(y \mid x,\, T{=}t) - f(y \mid x,\, \mathrm{do}(T{=}t)) = \int f(y \mid x,\, T{=}t,\, U{=}u)\big[\, p(u \mid x,\, T{=}t) - p(u \mid x)\,\big]\, du$$

is non-zero whenever U and T are dependent — the endogeneity condition.

1.3.1 Two Scenarios: Observed vs. Unobserved Confounder

Scenario 1: U is observed (no unmeasured confounding). When every component of U is recorded in the data, the back-door adjustment formula (Equation 1.2) becomes usable from data:

$$f(y \mid x,\, \mathrm{do}(T{=}t)) = \int f(y \mid x,\, T{=}t,\, U{=}u)\; p(u \mid x)\; du. \tag{1.3}$$

The confounder is averaged over its *population* distribution $p(u \mid x)$, not over the treatment-conditional distribution. A special case is *unconfoundedness*: if $U \perp\!\!\!\perp T \mid X$ already holds observationally, formula Equation 1.3 reduces to the familiar regression $\mathbb{E}[Y \mid X{=}x, T{=}t]$. This is the key assumption behind regression adjustment and propensity score methods studied in Chapter 6.

Scenario 2: U is unobserved (unmeasured confounding). When U is latent, formula Equation 1.3 is not usable. Additional structure beyond (Y, T, X) must be imposed. The main strategies studied in this course are **instrumental variables** (Chapter 7), the **front-door criterion** (Chapter 3), and **sensitivity analysis**.

Table 1.1: The two scenarios arising from the back-door adjustment proposition.

	Scenario 1: U observed	**Scenario 2: U unobserved**
Confounding type	No unmeasured confounding	Unmeasured confounding
Back-door formula usable?	Yes	No
Identification route	Back-door / standardization	IV, front-door, …
Key assumption	Unconfoundedness: $U \perp\!\!\!\perp T \mid X$	Exclusion restriction, monotonicity, …
Chapters	2–6	7

1.4 The Gaussian Linear Confounded Model

1.4.1 Setup

The Gaussian linear confounded model is our primary running example throughout this course. It is simple enough to permit closed-form calculations yet rich enough to illustrate every key concept.

Example: Gaussian Linear Confounded Model

The structural equations are

$$Y = \beta T + \gamma U + \varepsilon, \qquad T = \alpha U + \delta, \tag{1.4}$$

where U is an unobserved confounder and the three primitive variables are mutually independent:

$$U \sim \mathcal{N}(0, \sigma_U^2), \qquad \varepsilon \sim \mathcal{N}(0, \sigma_\varepsilon^2), \qquad \delta \sim \mathcal{N}(0, \sigma_\delta^2).$$

Here β is the causal effect of T on Y, γ is the direct effect of the confounder U on Y, and α is the direct effect of U on T.

The endogeneity of T is immediate. From the analyst's perspective, the composite error in the Y equation is $\gamma U + \varepsilon$, so:

$$\mathrm{Cov}(T,\ \gamma U + \varepsilon) = \gamma\,\mathrm{Cov}(\alpha U + \delta,\ U) = \gamma\alpha\sigma_U^2 \ \neq\ 0 \quad (\text{when } \alpha \neq 0 \text{ and } \gamma \neq 0).$$

Consequently, OLS regressing Y on T does not estimate β.

The causal structure is captured by the DAG pair in Figure 1.3. In the observational DAG, the back-door path $T \leftarrow U \rightarrow Y$ is open: unobserved confounder U causes both T and Y, making T and the composite error $\gamma U + \varepsilon$ correlated. Under the intervention do(T=t), all arrows into T are deleted (shown faded), severing the back-door path.

Figure 1.3: Graph surgery on the Gaussian confounded model. The faded arrow in the mutilated DAG has been severed by the intervention do(T=t). The back-door path $T \leftarrow U \rightarrow Y$ is eliminated; only the causal path $T \rightarrow Y$ remains active. The treatment node becomes a fixed constant (rectangle).

1.4.2 Two Explicit Densities

Write $\sigma_T^2 = \alpha^2\sigma_U^2 + \sigma_\delta^2$ for the marginal variance of T.

Interventional density. Set $T = t$ externally. The back-door path through U is severed, so U is independent of T under the intervention and is drawn from its marginal $\mathcal{N}(0, \sigma_U^2)$. Then $Y \mid \mathrm{do}(T{=}t) = \beta t + \gamma U + \varepsilon$ with $U \perp\!\!\!\perp \varepsilon$. Hence

$$f(y \mid \mathrm{do}(T{=}t)) = \mathcal{N}\big(\beta t,\ \ \gamma^2\sigma_U^2 + \sigma_\varepsilon^2\big)\,. \tag{1.5}$$

Observational density. Observe $T = t$. Since Y and T are both linear combinations of the independent normals (U, ε, δ), the pair (Y, T) is jointly normal. Their covariance is $\mathrm{Cov}(Y, T) = \beta\sigma_T^2 + \gamma\alpha\sigma_U^2$. Applying the conditional normal formula:

$$f(y \mid T{=}t) = \mathcal{N}\left(\left(\beta + \frac{\gamma\alpha\sigma_U^2}{\sigma_T^2}\right)t,\ \ \frac{\gamma^2\sigma_U^2\sigma_\delta^2}{\sigma_T^2} + \sigma_\varepsilon^2\right). \tag{1.6}$$

1.4.3 The Endogeneity Gap

Equations Equation 1.5 and Equation 1.6 differ in both mean and variance:

Table 1.2: *Equal when $\alpha = 0$ or $\gamma = 0$ (no confounding)*

	Interventional	**Observational**
Mean	βt	$\left(\beta + \frac{\gamma\alpha\sigma_U^2}{\sigma_T^2}\right) t$
Variance	$\gamma^2\sigma_U^2 + \sigma_\varepsilon^2$	$\frac{\gamma^2\sigma_U^2\sigma_\delta^2}{\sigma_T^2} + \sigma_\varepsilon^2$
Residual	$(\gamma U{+}\varepsilon) \perp\!\!\!\perp T$	$(\gamma U{+}\varepsilon)$ correlated with T
Identified by	RCT (or IV, Chapter 7)	OLS

The *endogeneity bias* of OLS is $\gamma\alpha\sigma_U^2/\sigma_T^2$: the coefficient of T in the observational mean exceeds the causal coefficient β by this amount. Numerically, with $\alpha = \gamma = 1$ and $\sigma_U^2 = \sigma_\delta^2 = 1$:

$$\text{bias} = \frac{1 \times 1 \times 1}{1^2 \times 1 + 1} = \frac{1}{2} = 0.5.$$

1.4.4 Lab: Simulating the Two Densities

The analytical results of this section were verified by simulation using $n = 10{,}000$ draws from the structural equations with parameters $\beta = 2$, $\alpha = 1$, $\gamma = 1$, $\sigma_U = \sigma_\varepsilon = \sigma_\delta = 1$. The R code is distributed separately as `chapter1_lab.R`.

Experiment 1: OLS bias. By the omitted-variable bias formula, the OLS probability limit is $\beta + \gamma\alpha\sigma_U^2/\sigma_T^2 = 2 + 0.5 = 2.5$.

	Simulated	**Theory**	**Formula**
OLS slope	2.5147	2.5000	$\beta + \gamma\alpha\sigma_U^2/\sigma_T^2$
True β	—	2.0000	β
Bias	0.5147	0.5000	$\gamma\alpha\sigma_U^2/\sigma_T^2$

OLS overestimates the causal effect by 25%.

Experiment 2: Two-sample comparison at $t_0 = 1$. Theoretical values are from Equation 1.5 and Equation 1.6 at $t_0 = 1$.

	Mean (Simulated)	**Mean (Theory)**	**SD (Simulated)**	**SD (Theory)**
$\mathbb{E}[Y \mid T{=}1]$	2.567	2.500	1.240	$\sqrt{1.5} = 1.225$
$\mathbb{E}[Y \mid \text{do}(T{=}1)]$	1.999	2.000	1.423	$\sqrt{2} = 1.414$

The variance reduction in the observational distribution reflects the ratio $\sigma_{\text{obs}}^2/\sigma_{\text{int}}^2 = 1.5/2 = 0.75$.

Experiment 3: Density overlay for varying α. Figure 1.4 shows the interventional density $\mathcal{N}(\beta t_0, \gamma^2\sigma_U^2 + \sigma_\varepsilon^2)$ (solid) and the observational density Equation 1.6 (dashed), both evaluated at $t_0 = 1$, for $\alpha \in \{0, 0.3, 0.7, 1.0\}$.

1.4.5 The IV DAG Versus the Mediation DAG

The IV DAG introduced above looks, at first glance, structurally similar to the mediation DAG that will appear in Chapter 8. The similarity is superficial; the conceptual difference is fundamental. Confusing the two is one of the most common errors in applied causal inference.

Figure 1.5 places the two DAGs side by side.

Table 1.3 makes the comparison systematic.



Figure 1.4: Interventional density $\mathcal{N}(2,2)$ (solid black) versus observational density for $\alpha \in \{0, 0.3, 0.7, 1.0\}$ (dashed), evaluated at $t_0 = 1$. As α increases, the observational curve shifts rightward (growing bias) and narrows (reduced variance). At $\alpha = 0$ there is no confounding and the two densities coincide. The interventional curve (solid) is invariant to α.



Figure 1.5: The IV DAG (a) and the mediation DAG (b) both involve a three-node chain, but they are structurally and conceptually distinct. In (a), the instrument Z is pre-treatment and exogenous: it satisfies $Z \perp\!\!\!\perp U$ and has no direct path to Y (exclusion restriction). In (b), the mediator M is post-treatment and endogenous: it is caused by T and transmits part of T's effect to Y.

Table 1.3: Instrumental variable versus mediator: a systematic comparison.

	Instrument Z	**Mediator** M
Position relative to T	Pre-treatment (upstream)	Post-treatment (downstream)
Causal relationship to T	Z causes T	T causes M
Direct path to Y?	**No** (exclusion restriction)	**Yes** ($M \to Y$ on causal path)
Independence from U?	**Yes** ($Z \perp\!\!\!\perp U$)	Not required
Primary purpose	Identify $T \to Y$ with unmeasured confounding	Decompose $T \to Y$ into direct and indirect
Key assumptions	Relevance, exogeneity, exclusion	Cross-world independence; no M–Y unmeasured confounding
Course coverage	Chapter 7	Chapter 8

The two chains look similar but are not interchangeable:

$$\underbrace{Z \longrightarrow T \longrightarrow Y}_{\text{IV chain: } Z \text{ is the identification device}} \qquad \text{versus} \qquad \underbrace{T \longrightarrow M \longrightarrow Y}_{\text{Mediation chain: } M \text{ is on the causal path}}$$

> Remark
>
> There is one important configuration where an observed post-treatment variable serves a role *analogous* to an instrument: the *front-door criterion* (Chapter 3). When $T \to M \to Y$ and $T \leftarrow U \to Y$ with U unobserved, but M is itself unconfounded, the causal effect of T on Y can still be identified by combining two back-door adjustments. This is the one case where a mediator also plays an identification role.

1.5 The Two-Step Paradigm

Causal inference is a two-step discipline. The two steps are logically distinct and require different tools.

1.5.1 Step 1: Identification

> Definition: Identification
>
> A causal quantity $P(y \mid \mathrm{do}(T{=}t))$ is *identified* from the observational distribution P if there exists a functional Ψ such that
> $$P(y \mid \mathrm{do}(T{=}t)) = \Psi(P),$$
> and $\Psi(P)$ depends only on the observable joint distribution.

Identification is a purely mathematical question about the structure of the causal model: given the assumed graph and assumptions, can the interventional density be expressed as a function of the observed-data distribution? It does not depend on sample size.

The main tools are: the back-door criterion (Chapters 2–3, applied in Chapters 5–6), the front-door criterion (Chapter 3), the three rules of the do-calculus (Chapter 3), and IV assumptions (Chapter 7).

1.5.2 Step 2: Estimation

Once $\Psi(P)$ has been identified, the statistical problem is to estimate $\Psi(P)$ from n observations $O_1, \dots, O_n$ as efficiently as possible. The main tools are efficient influence functions (EIF), inverse probability weighting (IPW), doubly robust estimators, and double machine learning (DML), all covered in Part III.

1.5.3 A Worked Example: Flu Vaccination and Infection

Scenario. A public health agency observes 1,000 individuals over one flu season. Each person either received a flu vaccine ($T = 1$) or did not ($T = 0$), and either became infected ($Y = 1$) or did not ($Y = 0$).

Age group X (elderly $= E$, young $= Y$) is recorded for all individuals. Elderly people are both more likely to be vaccinated and more susceptible to infection, so X confounds the relationship between T and Y.

The causal DAG is shown in Figure 1.6.



Figure 1.6: Causal DAG for the vaccination example. The green arrow is the causal path $T \to Y$. The red dashed arrows form the back-door path $T \leftarrow X \to Y$: age X causes both vaccination T and infection Y, creating a spurious association. Because X is observed, conditioning on it blocks the back-door path.

Observed data.

	Vaccinated ($T = 1$)		**Unvaccinated** ($T = 0$)		Total
	n	Infected	n	Infected	
Elderly ($X = E$)	360	108 (30%)	40	20 (50%)	400
Young ($X = Y$)	60	3 (5%)	540	54 (10%)	600
Total	420	111 (26.4%)	580	74 (12.8%)	1,000

Simpson's paradox. Within every stratum the vaccine reduces infection. Yet in the aggregate the vaccinated group has a *higher* infection rate (26.4% vs 12.8%). The reversal occurs because the vaccinated group is disproportionately elderly (86% of vaccinated vs. 7% of unvaccinated).

Stratum	Vaccinated	Unvaccinated	Difference	Conclusion
Elderly ($X = E$)	30%	50%	-20 pp	vaccine helps
Young ($X = Y$)	5%	10%	-5 pp	vaccine helps
Aggregate	**26.4%**	**12.8%**	$+13.6$ **pp**	**vaccine harms??**

Step 1: Identification. The set $\{X\}$ satisfies the back-door criterion: it blocks the path $T \leftarrow X \to Y$ and contains no descendant of T. By the back-door formula:

$$P(Y{=}1 \mid \mathrm{do}(T{=}t)) = \sum_{x \in \{E,Y\}} P(Y{=}1 \mid T{=}t, X{=}x)\, P(X{=}x). \tag{1.7}$$

Step 2: Estimation. Plugging sample proportions into Equation 3.1, with $\hat{P}(X{=}E) = 0.4$ and $\hat{P}(X{=}Y) = 0.6$:

$$\hat{P}(Y{=}1 \mid \mathrm{do}(T{=}1)) = 0.30 \times 0.4 + 0.05 \times 0.6 = 0.15, \qquad \hat{P}(Y{=}1 \mid \mathrm{do}(T{=}0)) = 0.50 \times 0.4 + 0.10 \times 0.6 = 0.26.$$

The estimated causal risk difference is $\hat{\Delta}_{\text{causal}} = 0.15 - 0.26 = -0.11$: the vaccine causally reduces infection probability by 11 percentage points.

The Two Steps, Concretely

Step 1 (identification) established — from the DAG alone — that the back-door formula (Equation 3.1) expresses the causal quantity as a functional of observable data.
Step 2 (estimation) replaced the population probabilities in that formula with sample proportions. These two steps are logically separate: Step 1 is a mathematical argument about the causal model that does not depend on sample size; Step 2 is a statistical argument about finite-sample estimation that does not depend on the causal model. A researcher who skips Step 1 and reports the raw difference conflates the two and draws the wrong conclusion.

1.6 Summary

1. **Interventional vs. observational distribution.** Causal inference answers questions about interventions $f(y \mid \mathrm{do}(T{=}t))$, not about observations $f(y \mid T{=}t)$. These two distributions coincide only under exogeneity; whenever T is endogenous they differ, and the difference is endogeneity bias.
2. **The causal trinity.** The same causal question can be expressed in three languages — SEM (equation surgery), DAG (graph surgery), and potential outcomes ($Y(t)$). The do-operator makes the interventional/observational distinction syntactically explicit. Fluency in all three is essential for modern causal reasoning.
3. **Endogeneity bias in the Gaussian confounded model.** In the Gaussian linear confounded model, the coefficient of T in the observational mean $\mathbb{E}[Y \mid T{=}t]$ exceeds the causal coefficient β by $\gamma\alpha\sigma_U^2/\sigma_T^2$, where $\sigma_T^2 = \alpha^2\sigma_U^2 + \sigma_\delta^2$. This is the omitted-variable bias.
4. **Two-step paradigm.** Causal inference is a two-step discipline: *identification* (expressing $\Psi(P)$ as a functional of observable data, a mathematical question about the causal model) followed by *estimation* (constructing $\hat{\Psi}_n$ efficiently from n observations, a statistical question about finite samples). The two steps are logically independent.
5. **Instrument vs. mediator.** The IV DAG ($Z \to T \to Y$ with $Z \perp\!\!\!\perp U$) and the mediation DAG ($T \to M \to Y$) are structurally distinct. Conflating the two leads to violations of the exclusion restriction and identification failure.

Causal inference is the study of when the observational distribution contains enough information to recover the interventional distribution. Everything in this course — the back-door criterion, do-calculus, instrumental variables, doubly robust estimation — is an answer to that question in a particular setting.

1.7 Problems

1. Do-notation fundamentals. For each statement below, decide whether it refers to an interventional or an observational distribution, and rewrite it unambiguously using do-notation where appropriate.

(a) "The probability that a patient recovers given that they took the drug."
(b) "The probability that a patient would recover if we prescribed the drug to everyone."
(c) "Among students who attended tutoring, the average exam score was 85."
(d) "If we enrolled all students in tutoring, the average exam score would be 85."

2. Deriving the observational density. In the Gaussian confounded model, verify Equation 1.6 by carrying out the following steps. Let $\sigma_T^2 = \alpha^2\sigma_U^2 + \sigma_\delta^2$.

(a) Show that (Y, T) is jointly normal by writing both as linear combinations of the independent normals (U, ε, δ).
(b) Compute $\mathrm{Cov}(Y, T)$ and $\mathrm{Var}(T)$.
(c) Apply the conditional normal formula to derive $\mathbb{E}[Y \mid T{=}t]$ and $\mathrm{Var}[Y \mid T{=}t]$, and hence verify Equation 1.6.
(d) Show that the OLS probability limit is $\beta + \gamma\alpha\sigma_U^2/\sigma_T^2$, and interpret this as an omitted-variable bias formula.

3. The causal trinity. Consider the following verbal causal claim: "Aspirin (T) reduces the risk of heart attack (Y) because it inhibits platelet aggregation (M), but patients with pre-existing cardiovascular disease (U, unobserved) are both more likely to take aspirin and more likely to have a heart attack."

(a) Draw the causal DAG implied by this description.
(b) Write down the recursive SEM with appropriate structural functions f_T, f_M, f_Y.
(c) State the SUTVA assumption and discuss whether it is plausible in this setting.
(d) Explain why $\mathbb{E}[Y \mid T{=}1] - \mathbb{E}[Y \mid T{=}0]$ does not identify the causal effect $\mathbb{E}[Y \mid \mathrm{do}(T{=}1)] - \mathbb{E}[Y \mid \mathrm{do}(T{=}0)]$ in this DAG.

4. IV versus mediation. For each scenario below, determine whether the intermediate variable described is an *instrument* or a *mediator*, and state which identifying assumption it satisfies or violates.

(a) A researcher uses distance to the nearest college (Z) to study the effect of college attendance (T) on earnings (Y).

(b) A nutritionist studies how a high-fiber diet (T) reduces cholesterol (M), which in turn reduces heart disease risk (Y).
(c) A political scientist uses rainfall on election day (Z) to estimate the effect of voter turnout (T) on electoral outcomes (Y).
(d) A pharmacologist studies whether a drug (T) reduces blood pressure (Y) by inhibiting a specific enzyme (M), and proposes to use M as an instrument for T. Evaluate this proposal.

Chapter 2

DAGs and d-Separation

Learning Objectives

By the end of this chapter, students should be able to:

1. Construct a DAG from a verbal description of a causal model and identify parents, descendants, ancestors, and non-descendants of any node.
2. Classify every intermediate node on a path as a chain, fork, or collider, and apply the blocking rule for each.
3. Apply the d-separation criterion to determine all conditional independence relationships implied by a DAG.
4. Recognize and avoid collider bias, including the case where a descendant of a collider is conditioned upon.
5. State the Markov factorization and use it to write the joint density of a DAG in terms of local conditional densities.
6. Interpret d-separation results as conditional independence assumptions on observed variables, and recognize their role as the graphical expression of causal assumptions.

2.1 Directed Acyclic Graphs

DAGs allow us to translate qualitative causal assumptions into quantitative statistical restrictions. This single idea underlies everything in this chapter: once we draw a graph encoding what we believe about the causal structure of a problem, the graph tells us exactly which variables are independent, which are confounded, and what we need to condition on to identify a causal effect.

Remark

In this chapter, the word *graph* does not mean a plot of data or a graph of a function. It means a collection of nodes and edges used to represent relationships among variables. Our objective is to understand how such graphs encode statistical structure — especially conditional independence, confounding, and path blocking — and how this graphical structure will later support causal identification.

2.1.1 Basic Definition

Definition: Directed Acyclic Graph

A *directed acyclic graph* (DAG) $\mathcal{G} = (V, E)$ consists of a finite set of nodes V and a set of directed edges $E \subseteq V \times V$ such that there is no directed cycle: no node V_i has a directed path to itself.

In a causal DAG each node represents a variable and each directed edge $A \to B$ encodes "A is a direct cause of B, given all other variables in the model." The acyclicity condition is essential: it guarantees that

the variables can be ordered topologically so that the joint density admits the recursive factorization

$$p(v_1, \dots, v_k) = \prod_{i=1}^{k} p(v_i \mid \mathrm{Pa}(v_i)).$$

2.1.2 Structural Relationships

Definition: Structural Relationships in a DAG

For a node $v \in \mathcal{V}$ in $\mathcal{G} = (\mathcal{V}, E)$:

- $\mathrm{Pa}(v) = \{w \in \mathcal{V} : w \to v \in E\}$ are the *parents* of v.
- $\mathrm{De}(v) =$ all nodes reachable from v by directed paths are the *descendants* of v.
- $\mathrm{An}(v) =$ all nodes with a directed path to v are the *ancestors* of v.
- $\mathrm{Nd}(v) = \mathcal{V} \ (\mathrm{De}(v) \cup \{v\})$ are the *non-descendants* of v.

Note that $\mathrm{Nd}(v)$ includes the parents of v; when stating the local Markov property in Section 2.4, we will exclude the parents explicitly.

Quick Terminology Summary

A *node* represents a variable, and an *edge* represents a relationship between two variables. If $A \to B$, then A is a *parent* of B and B is a *child* of A. Two nodes are *adjacent* if they are connected by an edge. A *path* is any sequence of connected nodes, regardless of arrow direction; a *directed path* is a path whose arrows all point in the same direction. A *directed acyclic graph* (DAG) is a directed graph with no directed cycles.

Example: Education, Earnings, and Family Background

Consider the causal story: education (E) affects earnings (Y); family background (B) is a common cause of both; and neighborhood (N) affects education but has no direct effect on earnings.



The Markov factorization is $p(n, b, e, y) = p(n)\, p(b)\, p(e \mid n, b)\, p(y \mid e, b)$. Reading off structural relationships: $\mathrm{Pa}(E) = \{N, B\}$, $\mathrm{Pa}(Y) = \{E, B\}$, $\mathrm{Pa}(N) = \mathrm{Pa}(B) = \emptyset$. The descendants of N are $\{E, Y\}$; B and N are non-descendants of each other.

Example: The IV DAG

The canonical instrumental variable model involves four nodes: Z (instrument), T (treatment), Y (outcome), and U (unobserved confounder).



$\mathrm{Pa}(T) = \{Z, U\}$, $\mathrm{Pa}(Y) = \{T, U\}$, $\mathrm{Pa}(Z) = \mathrm{Pa}(U) = \emptyset$. The descendants of Z are $\{T, Y\}$; U and Z are non-descendants of each other.

Example: The Mediation DAG

In a mediation model, treatment T affects outcome Y both directly and through an intermediate variable M (the mediator). An unobserved confounder U creates a back-door path $T \leftarrow U \to Y$,

making T endogenous. The mediator M is itself unconfounded.



$\mathrm{Pa}(T) = \{U\}$, $\mathrm{Pa}(M) = \{T\}$, $\mathrm{Pa}(Y) = \{M, T, U\}$, $\mathrm{Pa}(U) = \emptyset$. The total effect of T on Y travels along two paths: the direct path $T \to Y$ and the indirect path $T \to M \to Y$. Because there is a direct path $T \to Y$ bypassing M, the front-door criterion does not apply here.

2.2 Paths, Blocking, and d-Separation

A DAG encodes direct causal relationships through its edges, but variables can also be statistically related through longer routes that mix directed and reversed edges. The concept of a *path* formalises these routes, and the question of whether a path transmits or blocks statistical dependence — depending on which variables are conditioned upon — is precisely what d-separation answers. The key insight is that conditioning can both *close* channels that were open (by blocking a chain or fork) and *open* channels that were closed (by activating a collider). Figure 2.1 illustrates all three cases.

2.2.1 Path Structures

Definition: Path

A *path* between nodes X and Y in $\mathcal{G}$ is any sequence of distinct nodes $X = V_0, V_1, \dots, V_k = Y$ such that each consecutive pair is connected by an edge in either direction.

Every intermediate node V_m on a path plays one of three structural roles. In a **chain** ($V_{m-1} \to V_m \to V_{m+1}$), V_m is a causal intermediary through which influence is transmitted. In a **fork** ($V_{m-1} \leftarrow V_m \to V_{m+1}$), V_m is a common cause that creates confounding between the two endpoints. In a **collider** ($V_{m-1} \to V_m \leftarrow V_{m+1}$), both arrows point *into* V_m; this is the crucial asymmetric case whose behavior under conditioning is the opposite of the other two.



Figure 2.1: The three fundamental path structures in a DAG. Chains and forks are *open by default* and are blocked by conditioning on M. Colliders are *closed by default* and are opened by conditioning on M (or any descendant of M).

2.2.2 Blocking Rules

Structure	Pattern	Effect of conditioning	Key intuition
Chain	$A \to M \to B$	Blocks the path	Causal transmission
Fork	$A \leftarrow M \to B$	Blocks the path	Common cause
Collider	$A \to M \leftarrow B$	**Opens** the path	Common effect

The critical asymmetry — colliders are *closed by default* and *opened* by conditioning, while chains and

forks are *open by default* and *closed* by conditioning — is what makes collider bias so easy to overlook in practice.

Chain: Smoking → Tar → Cancer. Let A = smoking, M = tar deposits, B = lung cancer. Marginally, heavy smokers have elevated cancer rates — the path is open. Conditioning on tar level blocks the path: once tar level is fixed, the causal channel is fully accounted for.

Fork: Poverty → Poor Diet; Poverty → Lack of Exercise. Let M = poverty, A = poor diet, B = lack of exercise. Unconditionally, diet quality and exercise are correlated through poverty. Conditioning on income level removes this spurious association.

Collider: Accident → Hospitalization ← Cancer. Let A = traffic accident, B = cancer, M = hospitalization. In the general population, $A \perp\!\!\!\perp B$. But among hospitalized patients — conditioning on M — the two become negatively associated. This is Berkson's bias (Berkson 1946).

Three-Node Motifs at a Glance

In a *chain* $A \to M \to B$, conditioning on M blocks the path. In a *fork* $A \leftarrow M \to B$, conditioning on M removes the spurious association. In a *collider* $A \to M \leftarrow B$, the path is blocked by default, but conditioning on M — or on any descendant of M — opens the path and induces association. These three motifs are the local building blocks of d-separation in arbitrary DAGs.

2.2.3 The d-Separation Criterion

Definition: d-Blocking and d-Separation

A path π is *d-blocked* by a set $\mathbf{S}$ if:

1. π contains a chain $A \to M \to B$ or fork $A \leftarrow M \to B$ with $M \in \mathbf{S}$; or
2. π contains a collider $A \to C \leftarrow B$ with $C \notin \mathbf{S}$ and no descendant of C is in $\mathbf{S}$.

Nodes X and Y are *d-separated* by $\mathbf{S}$, written $(X \perp\!\!\!\perp Y \mid \mathbf{S})_{\mathcal{G}}$, if every path between X and Y in $\mathcal{G}$ is d-blocked by $\mathbf{S}$.

Example: Applying the d-Separation Definition to the Three Toy Settings

(i) Chain. A = smoking, M = tar, B = cancer. Only path: $A \to M \to B$.

- $\mathbf{S} = \emptyset$: not d-blocked. $\Rightarrow A \not\perp\!\!\!\perp B$.
- $\mathbf{S} = \{M\}$: clause (1) applies. $\Rightarrow A \perp\!\!\!\perp B \mid M$.

(ii) Fork. M = poverty, A = poor diet, B = lack of exercise. Only path: $A \leftarrow M \to B$.

- $\mathbf{S} = \emptyset$: not d-blocked. $\Rightarrow A \not\perp\!\!\!\perp B$.
- $\mathbf{S} = \{M\}$: clause (1) applies. $\Rightarrow A \perp\!\!\!\perp B \mid M$.

(iii) Collider. A = accident, M = hospitalization, B = cancer. Only path: $A \to M \leftarrow B$.

- $\mathbf{S} = \emptyset$: collider $M \notin \mathbf{S}$, clause (2) applies. $\Rightarrow A \perp\!\!\!\perp B$.
- $\mathbf{S} = \{M\}$: clause (2) no longer blocks. $\Rightarrow A \not\perp\!\!\!\perp B \mid M$ (Berkson's bias).

Example: A Direct Probability Proof for the Chain

Consider the chain $X_1 \to X_2 \to X_3$. By the Markov factorization, $p(x_1, x_2, x_3) = p(x_1)\,p(x_2 \mid x_1)\,p(x_3 \mid x_2)$. Conditioning on $X_2 = x_2$:

$$p(x_1, x_3 \mid x_2) = \frac{p(x_1)\,p(x_2 \mid x_1)\,p(x_3 \mid x_2)}{p(x_2)} = p(x_1 \mid x_2)\,p(x_3 \mid x_2).$$

Hence $X_1 \perp\!\!\!\perp X_3 \mid X_2$. The fork case follows by an identical calculation.

Theorem: Verma & Pearl (1988)

(Pearl 2009, Theorem 1.2.4) If $(X \perp\!\!\!\perp Y \mid \mathbf{S})_{\mathcal{G}}$, then $X \perp\!\!\!\perp Y \mid \mathbf{S}$ in every distribution that is Markov with respect to $\mathcal{G}$.

Remark: Soundness, Not Converse

Theorem 2.1 (Verma & Pearl) states that d-separation implies conditional independence for every distribution Markov with respect to $\mathcal{G}$. The converse need not hold without additional assumptions such as *faithfulness*.

Table 2.1 summarizes the Markov factorization and its independence implication for each of the three path structures.

Table 2.1: Markov factorization and d-separation result for each path structure.

Structure	**Markov factorization** $p(a, m, b)$	**d-separation**
Chain $A \to M \to B$	$p(a)\,p(m \mid a)\,p(b \mid m)$	$(A \perp\!\!\!\perp B \mid M)_{\mathcal{G}}$
Fork $A \leftarrow M \to B$	$p(m)\,p(a \mid m)\,p(b \mid m)$	$(A \perp\!\!\!\perp B \mid M)_{\mathcal{G}}$
Collider $A \to M \leftarrow B$	$p(a)\,p(b)\,p(m \mid a, b)$	$(A \perp\!\!\!\perp B)_{\mathcal{G}}$

2.2.4 Practical Ways to Check d-Separation

Bayes-Ball Intuition. Imagine releasing a ball from X asking whether it can reach Y, given that nodes in $\mathbf{S}$ are observed. At a chain or fork node not in $\mathbf{S}$: ball passes through. At a chain or fork node in $\mathbf{S}$: ball stops. At a collider not in $\mathbf{S}$ (and no descendant in $\mathbf{S}$): ball stops. At a collider in $\mathbf{S}$ (or with a descendant in $\mathbf{S}$): ball passes through (Shachter 1998).

Moral Graph Transformation. (1) Restrict to the ancestral set $\mathrm{An}(X \cup Y \cup \mathbf{S})$. (2) Moralise: for every collider $A \to C \leftarrow B$, add an undirected edge $A - B$. (3) Remove all edge directions. (4) Delete all nodes in $\mathbf{S}$. If X and Y are disconnected, then $(X \perp\!\!\!\perp Y \mid \mathbf{S})_{\mathcal{G}}$ (Lauritzen 1996, Ch. 3).

Example: Checking d-Separation in a Four-Node DAG

Consider the DAG with edges $Z \to T$, $U \to T$, $T \to Y$, $U \to Y$.



Query 1. Is $(Z \perp\!\!\!\perp U)_{\mathcal{G}}$? The only path $Z \to T \leftarrow U$ has T as a collider with $T \notin \emptyset$ — blocked. $(Z \perp\!\!\!\perp U)_{\mathcal{G}}$.
Query 2. Is $(Z \perp\!\!\!\perp U \mid T)_{\mathcal{G}}$? Now $T \in \mathbf{S}$: the collider is observed, the path is open. $Z \not\perp\!\!\!\perp U \mid T$.
Comparing the two queries: Z and U are marginally independent but become dependent once we condition on T. This is Berkson's bias.

Example: Practice DAG — Eight-Node Graph

Consider the DAG with nodes W, X_1, T, M_1, M_2, Y, C, D and edges $W \to X_1$, $W \to T$, $W \to Y$, $T \to M_1$, $M_1 \to Y$, $M_1 \to M_2$, $X_1 \to C$, $M_2 \to C$, $C \to D$.



(1) Is $X_1 \perp\!\!\!\perp Y$? No. Two open paths: $X_1 \leftarrow W \rightarrow Y$ and $X_1 \leftarrow W \rightarrow T \rightarrow M_1 \rightarrow Y$.
(2) Is $X_1 \perp\!\!\!\perp Y \mid W$? Yes. Both open paths pass through the fork W; conditioning on W blocks them.
(3) Is $T \perp\!\!\!\perp M_2 \mid M_1$? Yes. The path $T \rightarrow M_1 \rightarrow M_2$ is blocked by M_1; all other paths are also blocked.
(4) Does conditioning on C create collider bias? Yes. C is a collider on $X_1 \rightarrow C \leftarrow M_2$. Conditioning on C opens this path, inducing a spurious association between X_1 and M_2.
(5) Does conditioning on D open the path $X_1 \rightarrow C \leftarrow M_2$? Yes. D is a descendant of C. By clause (2) of Definition 2.1, the collider at C is activated.

2.3 Collider Bias and the IV DAG

2.3.1 Berkson's Bias

Definition: Collider Bias

Collider bias is the spurious association induced between two variables when one conditions on their common effect (a collider), or on a descendant of that collider. In a path $A \rightarrow C \leftarrow B$, the variables A and B may be marginally independent, but conditioning on C — or on any descendant of C — can make them statistically dependent.

Collider Bias vs. Confounding

Unlike confounding, which is *removed* by conditioning on a common cause, collider bias is *created* by conditioning on a common effect. This is why conditioning on more variables is not always safer in causal analysis: adding a collider or a descendant of a collider to the adjustment set introduces bias rather than reducing it.

Example: Collider Bias in the IV DAG

In the IV DAG, T is a collider on the path $Z \rightarrow T \leftarrow U$. Marginally, $Z \perp\!\!\!\perp U$ (the instrument is exogenous). However,

$$Z \perp\!\!\!\perp U \qquad \text{but} \qquad Z \not\perp\!\!\!\perp U \mid T.$$

Conditioning on T — restricting to the treated subpopulation — makes the instrument Z appear associated with U, invalidating the IV argument within that stratum.

Example: Collider Bias through Selection: Talent and Wealth

Suppose both talent (A) and wealth (B) increase the probability of admission to an elite school, so admission (S) is a collider: $A \to S \leftarrow B$.



In the general population, $A \perp\!\!\!\perp B$. But among admitted students — conditioning on S — lower talent makes higher wealth more likely, and vice versa. This is collider bias: restricting to admitted students is precisely conditioning on a common effect.

2.3.2 Full d-Separation Analysis of the IV DAG

We work through the IV DAG with edges $Z \to T$, $T \to Y$, $U \to T$, $U \to Y$, where U is unobserved.

Endpoints	**Path**	**Type**	**Why**
$Z \leftrightarrow Y$	$Z \to T \to Y$	*Causal*	Directed from Z to Y; carries the IV signal
$Z \leftrightarrow Y$	$Z \to T \leftarrow U \to Y$	*Associational*	Collider at T; activates back-door via U
$Z \leftrightarrow U$	$Z \to T \leftarrow U$	*Associational*	Collider at T; dormant unless T conditioned on

Question 1. Is $Z \perp\!\!\!\perp Y$? There are two paths. The directed path $Z \to T \to Y$ is open, since it is a chain and we are not conditioning on T. The path $Z \to T \leftarrow U \to Y$ contains a collider at T; since we are not conditioning on T, that path is blocked. Because the causal path remains open, $Z \not\perp\!\!\!\perp Y$.

Question 2. Is $Z \perp\!\!\!\perp Y \mid T$? The causal path $Z \to T \to Y$ is blocked because conditioning on T blocks the chain. The path $Z \to T \leftarrow U \to Y$ contains a collider at T, and conditioning on T opens it. Hence $Z \not\perp\!\!\!\perp Y \mid T$. The residual association is entirely spurious — a collider artefact, not a causal effect.

Question 3. Is $Z \perp\!\!\!\perp Y \mid \{T, U\}$? The causal path is blocked by T. The path $Z \to T \leftarrow U \to Y$ has its collider at T opened by conditioning on T, but then blocked at U because we also condition on U. Hence $Z \perp\!\!\!\perp Y \mid \{T, U\}$.

Question 4. Is $Z \perp\!\!\!\perp U$? The only path $Z \to T \leftarrow U$ contains a collider at T. Since we are not conditioning on T or any descendant of T, the path is blocked. Hence $Z \perp\!\!\!\perp U$. This is *exogeneity*.

The four questions give a complete graphical account of the three IV assumptions. Question 1 establishes *relevance*; Question 4 establishes *exogeneity*; Question 3 establishes the *exclusion restriction*. Question 2 is the warning: conditioning on T alone simultaneously closes the causal channel and opens the confounded channel.

2.4 The Markov Property and Factorization

Proposition: Conditional Independence in the Three-Node Chain

Let $X_1 \to X_2 \to X_3$ be a chain with Markov factorization $p(x_1, x_2, x_3) = p(x_1)\, p(x_2 \mid x_1)\, p(x_3 \mid x_2)$. Then $X_1 \perp\!\!\!\perp X_3 \mid X_2$. The analogous result for the fork $X_1 \leftarrow X_2 \to X_3$ is left as an exercise.

Definition: Local Markov Property

A distribution P satisfies the *local Markov property* with respect to $\mathcal{G}$ if, for every node $V_i \in \mathcal{V}$:

$$V_i \perp\!\!\!\perp (\mathrm{Nd}(V_i) \;\; \mathrm{Pa}(V_i)) \mid \mathrm{Pa}(V_i).$$

That is, once we condition on the direct causes of a node, that node is independent of all variables that are neither its descendants nor its parents.

Remark

The *global Markov property* is the statement that every d-separation in $\mathcal{G}$ implies a conditional independence in P — precisely the content of Theorem 2.1 (Verma & Pearl). For DAGs, the local and global Markov properties are equivalent (Lauritzen 1996, Proposition 3.27).

An immediate consequence is the *Markov factorization*:

$$p(v_1, \dots, v_k) \;=\; \prod_{i=1}^{k} p(v_i \mid \mathrm{Pa}(v_i))\,. \tag{2.1}$$

Example: Markov Factorization in the Education Example

For the DAG with edges $N \to E$, $B \to E$, $B \to Y$, $E \to Y$:

$$p(n, b, e, y) = p(n)\, p(b)\, p(e \mid n, b)\, p(y \mid e, b).$$

A regression of Y on E alone is confounded by B. Conditioning on B blocks the back-door path $E \leftarrow B \to Y$ and identifies $P(y \mid \mathrm{do}(E{=}e))$ via the back-door formula (Chapter 3).

Example: Markov Factorization in the IV DAG

Including the latent U: $p(z, t, y, u) = p(z)\, p(u)\, p(t \mid z, u)\, p(y \mid t, u)$. Since U is unobserved, the observable factorization is obtained by marginalizing:

$$p(z, t, y) = \int p(z)\, p(u)\, p(t \mid z, u)\, p(y \mid t, u)\, \mathrm{d}u.$$

This cannot be simplified to $p(z)\, p(t \mid z)\, p(y \mid t)$ when U is a common cause of T and Y: this is precisely the endogeneity problem.

2.5 Worked Example: The Education–Earnings DAG

This section is the template for how we will use DAGs throughout the course: specify the DAG, read off the factorization, use d-separation to identify implied conditional independences, and interpret the result in causal terms.

The causal story. Education (E) affects earnings (Y); family background (B) is a common cause of both; neighborhood (N) affects education but has no direct effect on earnings. All four variables are observed; there are no latent confounders.



Step 1 — Markov factorization. $\mathrm{Pa}(N) = \mathrm{Pa}(B) = \emptyset$, $\mathrm{Pa}(E) = \{N, B\}$, $\mathrm{Pa}(Y) = \{E, B\}$.

$$p(n, b, e, y) \;=\; p(n)\, p(b)\, p(e \mid n, b)\, p(y \mid e, b).$$

Step 2 — d-Separation. Two paths between N and Y: Path 1: $N \to E \to Y$ (chain). Path 2: $N \to E \leftarrow B \to Y$ (collider at E).

- *Query (i):* $(N \perp\!\!\!\perp Y)_{\mathcal{G}}$? Path 1 is a chain with nothing conditioned on — open. $N \not\perp\!\!\!\perp Y$.
- *Query (ii):* $(N \perp\!\!\!\perp Y \mid E)_{\mathcal{G}}$? Path 1 blocked by E; Path 2 collider E opened, path continues through B which is open. $N \not\perp\!\!\!\perp Y \mid E$. This is a collider trap: conditioning on E opens a spurious association between N and Y through B.

- *Query (iii):* $(N \perp\!\!\!\perp Y \mid \{E, B\})_{\mathcal{G}}$? Path 1 blocked by E; Path 2 opened at E but blocked at B. $(N \perp\!\!\!\perp Y \mid E, B)_{\mathcal{G}}$.
- *Query (iv):* $(N \perp\!\!\!\perp B)_{\mathcal{G}}$? Only path $N \to E \leftarrow B$ has collider E with $E \notin \emptyset$ — blocked. $N \perp\!\!\!\perp B$.

Step 3 — Conditional Independence. By Theorem 2.1 (Verma & Pearl): $N \perp\!\!\!\perp B$, $N \perp\!\!\!\perp Y \mid E, B$, $N \not\perp\!\!\!\perp Y$, $N \not\perp\!\!\!\perp Y \mid E$.

Step 4 — Identification. The only back-door path is $E \leftarrow B \to Y$. With $\mathbf{S} = \{B\}$:

$$P(y \mid \mathrm{do}(E{=}e)) \;=\; \sum_{b} P(y \mid e, b)\, P(b).$$

2.6 The Big Picture



Figure 2.2: The four-step pipeline from causal assumptions to identification formula.

The logical chain in compact form:

structural assumptions $\Longrightarrow$ DAG $\Longrightarrow$ d-separation $\Longrightarrow$ CI under the Markov property $\Longrightarrow$ identification formulas.

2.7 Summary

1. **DAGs as causal structure.** A DAG is a directed acyclic graph whose nodes represent variables and whose directed edges represent direct causal relationships. Once a DAG is specified, it encodes which variables are causally connected, which are potential confounders, and which may block or open paths.

2. **Three-node motifs.** Every path is built from three local structures: chains, forks, and colliders. Chains and forks are open by default and blocked by conditioning on the middle node. Colliders are blocked by default and opened by conditioning on the collider or one of its descendants.

3. **d-Separation and conditional independence.** The d-separation criterion is a graphical rule: X and Y are d-separated by $\mathbf{S}$ exactly when every path between them is blocked by $\mathbf{S}$. Under the Markov property, this implies the corresponding conditional independence. Some such implications may be assessed empirically, but agreement with data does not by itself verify the graph.

4. **Markov factorization.** The local Markov property yields

$$p(v_1, \dots, v_k) \;=\; \prod_{i=1}^{k} p(v_i \mid \mathrm{Pa}(v_i))\,,$$

which expresses the joint distribution in terms of local conditional distributions.

5. **Collider bias.** Conditioning on a collider — or on a descendant of a collider — can induce a spurious association between otherwise independent variables (Berkson 1946). This is the *opposite* of confounding adjustment: conditioning on a common cause removes spurious association, whereas conditioning on a common effect creates it.

6. **A practical workflow.** To analyze a DAG, proceed in four steps: identify the graph structure, determine which paths are open or blocked, translate d-separation statements into conditional independences under the Markov property, and interpret those independences in light of the causal question.

2.8 Problems

1. Warm-up: a single collider. Consider the DAG $X \to Y \leftarrow Z$, where X and Z have no other connections.

(a) Identify the structural role of Y on the path $X \to Y \leftarrow Z$.
(b) Is $X \perp\!\!\!\perp Z$? Apply the d-separation criterion and state which blocking rule applies.
(c) Is $X \perp\!\!\!\perp Z \mid Y$? Explain what happens to the path when Y is conditioned on.

2. d-Separation practice. Consider the DAG: $A \to B \to D$, $A \to C \to D$, $B \to E$, $C \to E$.

(a) List all paths between A and E. (*Hint*: there are four paths; two pass through D.)
(b) For each path, identify the role of each intermediate node.
(c) Does $\{B, C\}$ d-separate A and E?
(d) Does $\{D\}$ d-separate B and C? What type of node is D on the path $B \to D \leftarrow C$?

3. Markov factorization and collider activation. Consider the DAG with edges $A \to E$, $A \to W$, $F \to E$, $E \to W$, where A = ability, F = family income, E = education, W = wages.

(a) Write down the Markov factorization $p(a, f, e, w)$.
(b) Is $(F \perp\!\!\!\perp W)_{\mathcal{G}}$? List all paths between F and W.
(c) Is $(F \perp\!\!\!\perp W \mid E)_{\mathcal{G}}$? Identify the role of E on each path.
(d) A researcher regresses W on E and F, omitting A. Is the coefficient on E a causal effect?

4. Theorem for the collider. Consider the collider $A \to M \leftarrow B$ with Markov factorization $p(a, m, b) = p(a)\, p(b)\, p(m \mid a, b)$.

(a) By marginalizing over M, show that $A \perp\!\!\!\perp B$ in the joint distribution.
(b) Show that conditioning on M breaks this independence.
(c) Explain why parts (a) and (b) together are consistent with Theorem 2.1 (Verma & Pearl).

5. Terminology check. Consider the DAG with edges $U \to X$, $U \to Z$, $X \to W$, $Z \to W$, $W \to Y$.

(a) Identify the parents, children, ancestors, descendants, and non-descendants of node W.
(b) List all pairs of adjacent nodes.
(c) Which pairs of nodes are connected by a directed path?
(d) Write the Markov factorization $p(u, x, z, w, y)$.

6. Markov factorization and local Markov property. Consider the DAG with edges $X_1 \to X_2$, $X_1 \to X_3$, $X_2 \to X_4$, $X_3 \to X_4$.

(a) Write the joint density $p(x_1, x_2, x_3, x_4)$ implied by the Markov factorization.
(b) State the local Markov property for each of the four nodes.
(c) Is $(X_2 \perp\!\!\!\perp X_3)_{\mathcal{G}}$? Is $(X_2 \perp\!\!\!\perp X_3 \mid X_1)_{\mathcal{G}}$?

7. Toy proof: conditional independence in the fork. Consider the fork $X_1 \leftarrow X_2 \to X_3$ with Markov factorization $p(x_1, x_2, x_3) = p(x_2)\, p(x_1 \mid x_2)\, p(x_3 \mid x_2)$.

(a) Show directly that $X_1 \perp\!\!\!\perp X_3 \mid X_2$.
(b) Is $X_1 \perp\!\!\!\perp X_3$ marginally? Justify both graphically and algebraically.
(c) Explain why conditioning on the common cause X_2 removes the association.

8. d-Separation in the practice DAG. Refer to the eight-node DAG in the eight-node example above. For each query, state whether it is true or false and justify by listing all relevant paths.

(a) $X_1 \perp\!\!\!\perp Y$

(b) $X_1 \perp\!\!\!\perp Y \mid W$
(c) $T \perp\!\!\!\perp M_2 \mid M_1$
(d) Does conditioning on C induce collider bias between X_1 and M_2?
(e) Does conditioning on D open the path $X_1 \to C \leftarrow M_2$? State which clause of Definition 2.1 applies.

Chapter 3

The Do-Calculus and Identification Criteria

Learning Objectives

By the end of this chapter, students should be able to:

1. Define the two intervention graph operations $\mathcal{G}_{\overline{X}}$ and $\mathcal{G}_{\underline{X}}$ and explain their causal interpretation.
2. Apply the back-door criterion to determine whether a set $\mathbf{S}$ identifies a causal effect, and write the back-door adjustment formula.
3. Apply the front-door criterion in graphs where the confounder is unobserved but a suitable mediator exists.
4. State all three rules of do-calculus, identify the graph condition that licenses each rule, and use them to prove the back-door and front-door formulas.
5. State the completeness theorem of Shpitser and Pearl (2006) and explain its practical implication: if the do-calculus cannot identify a quantity from the observational distribution relative to the assumed graph, then no purely observational method can do so without additional assumptions or new data.

How to Read This Chapter

This chapter has two pedagogical layers. Sections 3.1–3.3 develop the two most important concrete identification criteria — back-door and front-door adjustment — using intervention graphs and direct graphical reasoning. These sections are the core material for a first reading.
Sections 3.4–3.5 then place these criteria inside the more general theory of the do-calculus and identifiability. These later sections are conceptually important, but they are more abstract and may be read as a second pass after the concrete criteria are understood.

3.1 From d-Separation to Intervention: Intervention Graphs

In Chapter 2 we learned to read conditional independence from a DAG using d-separation. This chapter takes the next step: translating the graphical language into *identification formulas* — expressions that write the interventional distribution $P(y \mid \mathrm{do}(T{=}t))$ entirely in terms of quantities observable from data.

3.1.1 What Does Intervention Mean?

Before introducing any graph machinery, it is worth being precise about what the $\mathrm{do}(\cdot)$ operator means and why it is not the same as conditioning.

Conditioning versus intervening. The conditional distribution $P(Y \mid T{=}t)$ describes the subpopulation of units for whom T was *observed* to equal t. Because treatment assignment may be influenced by confounders, this subpopulation is not representative of the full population. The interventional distribution

$P(Y \mid \mathrm{do}(T{=}t))$, by contrast, describes the population that would result if T were *externally set* to t for everyone — removing the natural process that determines T and replacing it with the forced value. The two distributions coincide only when T has no unmeasured common causes with Y.

A simple illustration: suppose temperature Z causes both ice-cream sales T and crime Y.

- $P(Y \mid T{=}t)$ is the crime rate on days when ice-cream sales happen to equal t — high sales happen on hot days, which also drive higher crime, so the correlation is confounded by Z.
- $P(Y \mid \mathrm{do}(T{=}t))$ is the crime rate we would observe if we *fixed* sales at level t by decree. Forcing sales does not change the temperature, so crime remains driven only by Z, not by the sales level we mandated. If T has no direct causal path to Y, then $P(Y \mid \mathrm{do}(T{=}t)) = P(Y)$ for all t.

The structural basis of the do-operator. In the SEM framework of Chapter 1, every variable is generated by a structural equation. For the treatment node:

$$T \;=\; f_T(\mathrm{Pa}(T), U_T),$$

where $\mathrm{Pa}(T)$ are the causal parents of T and U_T is exogenous noise. An *intervention* $\mathrm{do}(T{=}t)$ **replaces this entire equation** with the constant $T = t$. The replacement has two consequences: (1) T is no longer influenced by its parents — its dependence on $\mathrm{Pa}(T)$ and U_T is severed; (2) all other structural equations remain unchanged.

Graph surgery as the graphical realization. Because $\mathrm{Pa}(T)$ no longer affects T after the intervention, every arrow pointing *into* T in the DAG becomes meaningless and should be removed. The resulting graph — the *intervention graph* $\mathcal{G}_{\overline{T}}$ — represents the post-intervention world. D-separation in $\mathcal{G}_{\overline{T}}$ therefore encodes conditional independence in $P(\,\cdot \mid \mathrm{do}(T{=}t))$, not in the original observational distribution P.

3.1.2 The Two Graph Operations

Definition: Intervention Graphs $\mathcal{G}_{\overline{X}}$ and $\mathcal{G}_{\underline{X}}$

Let $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ be a DAG and $X \subseteq \mathcal{V}$ a set of nodes.

- $\mathcal{G}_{\overline{X}}$ (*intervention graph*, or *mutilated graph*) is obtained by *deleting all edges into* X: remove every arrow $V \to X_i$ for all $X_i \in X$. This represents the intervention $\mathrm{do}(X{=}x)$, which replaces the structural equation for X by the constant x, thereby severing X's dependence on all its former parents.
- $\mathcal{G}_{\underline{X}}$ (*auxiliary observation graph*) is obtained by *deleting all edges out of* X: remove every arrow $X_i \to V$ for all $X_i \in X$. This is a *technical device* used in Rule 1 of the do-calculus to determine when an observation on X can be inserted into or removed from a probability expression. It does *not* represent a probabilistic model of conditioning on X.

Mnemonic.

$$\mathcal{G}_{\overline{X}} : \text{cut } \textit{incoming} \text{ arrows into } X, \qquad \mathcal{G}_{\underline{X}} : \text{cut } \textit{outgoing} \text{ arrows from } X.$$

Figure 3.1 illustrates both operations on the IV DAG $\{Z \to T,\ T \to Y,\ U \to T,\ U \to Y\}$.



Figure 3.1: The two graph operations on the IV DAG. Faded arrows have been removed. $\mathcal{G}_{\overline{T}}$ models external intervention $\mathrm{do}(T{=}t)$: the back-door path $T \leftarrow U \to Y$ is severed. $\mathcal{G}_{\underline{T}}$ deactivates T's influence on its descendants, used by Rule 1 to add or delete T from a conditioning set.

The Intervention Principle

In the original DAG $\mathcal{G}$, the node T is determined by its parents $\mathrm{Pa}(T)$ via the structural equation $T = f_T(\mathrm{Pa}(T), \varepsilon_T)$. Intervening on T removes the structural mechanism that normally determines T; therefore the parents of T no longer influence it. The intervention $\mathrm{do}(T{=}t)$ *replaces this equation* with the constant $T = t$, making T independent of all its former parents. Graphically, this severs every arrow into T. The remaining structure of the graph is unchanged.
A d-separation statement in $\mathcal{G}_{\overline{T}}$ is therefore a conditional independence statement in the *post-intervention* distribution $P(\cdot \mid \mathrm{do}(T{=}t))$, not in the observational distribution P.

3.2 The Back-Door Criterion

The back-door criterion is the most widely used identification strategy. It gives a simple graphical condition under which conditioning on an observed set $\mathbf{S}$ is sufficient to identify the causal effect of T on Y.

3.2.1 Motivation: When the Confounder Is Unobserved

If all common causes of T and Y were observed, identification would be trivial: adjust for all of them. The interesting case is when some confounders are *unobserved*, yet an observed set $\mathbf{S}$ can still block every spurious path.

Consider the following DAG. A drug T affects recovery Y. Socioeconomic status U is an unobserved confounder ($U \to T$, $U \to Y$), but U also affects an observed proxy W ($U \to W$), which in turn affects Y ($W \to Y$).



The only back-door path is $T \leftarrow U \to W \to Y$. Although U is unobserved, W sits on this path and *is* observed. Conditioning on W blocks the path at the chain node W, and W is not a descendant of T, so $\mathbf{S} = \{W\}$ satisfies the back-door criterion. The key insight: the criterion does not require adjusting for the confounders themselves, only for an observed set that *intercepts* every back-door path.

3.2.2 Definition and Theorem

Definition: Back-Door Criterion

A set of observed variables $\mathbf{S}$ satisfies the *back-door criterion* for the causal effect of T on Y in $\mathcal{G}$ if:

1. No node in $\mathbf{S}$ is a descendant of T.
2. $\mathbf{S}$ blocks every *back-door path* from T to Y, i.e., every path that begins with an arrow pointing *into* T.

Remark

The terminology reflects the geometry of the graph relative to T. A *back-door path* enters T from behind — it begins with an arrow pointing *into* T — and represents a confounding route through which T and Y become spuriously associated without any causal mechanism from T to Y. The front-door criterion (Section Section 3.3) is named for the complementary idea: it identifies the causal effect by intercepting the *front-door* (causal) paths at a mediator M, rather than blocking the back-door (confounding) paths directly.
Condition 2 is equivalent to requiring $(Y \perp\!\!\!\perp T \mid \mathbf{S})_{\mathcal{G}_{\underline{T}}}$: $\mathbf{S}$ d-separates T and Y in the graph $\mathcal{G}_{\underline{T}}$, obtained by *deleting all arrows out of* T.

Example: A Single Confounder

Consider the DAG $T \leftarrow C \rightarrow Y$ with $T \rightarrow Y$, where C is an observed confounder. Does $\mathbf{S} = \{C\}$ satisfy the back-door criterion?

1. C is not a descendant of T.
2. $\{C\}$ blocks $T \leftarrow C \rightarrow Y$ (fork at C).

The back-door formula gives: $P(y \mid \mathrm{do}(T{=}t)) = \sum_c P(y \mid t, c)\, P(c)$.

Example: Two Back-Door Paths — Both Must Be Blocked

Consider treatment T, outcome Y, observed variables C_1 and C_2, and an unobserved confounder U with edges $C_1 \rightarrow T$, $C_1 \rightarrow Y$, $U \rightarrow T$, $U \rightarrow C_2$, $C_2 \rightarrow Y$, $T \rightarrow Y$.
There are exactly two back-door paths: (1) $T \leftarrow C_1 \rightarrow Y$ and (2) $T \leftarrow U \rightarrow C_2 \rightarrow Y$.

- $\mathbf{S} = \{C_1\}$: blocks path 1 but leaves path 2 open. **Fails.**
- $\mathbf{S} = \{C_2\}$: blocks path 2 but leaves path 1 open. **Fails.**
- $\mathbf{S} = \{C_1, C_2\}$: C_1 blocks path 1; C_2 blocks path 2. **Succeeds.**

$$P(y \mid \mathrm{do}(T{=}t)) \;=\; \sum_{c_1,\, c_2} P(y \mid t,\, c_1,\, c_2)\, P(c_1,\, c_2).$$

The unobserved U never appears in the formula: what matters is that C_2 sits on the back-door path and is observed.

Why Condition 1 Is Essential: A Worked Counterexample

Consider a drug T that affects recovery Y via direct effect $T \rightarrow Y$ and indirect effect $T \rightarrow M \rightarrow Y$, with unobserved confounder $U \rightarrow T$, $U \rightarrow Y$. If we naively try $\mathbf{S} = \{M\}$:

1. It violates Condition 1: M is a descendant of T.
2. It *blocks causal information*: the stratum-specific association $P(y \mid t, m)$ holds M fixed, so it cannot see the effect of T that flows through M. The result estimates the direct effect $T \rightarrow Y$ only, not the total causal effect.

Why Condition 2 Can Be Violated by a Collider: M-Bias

Consider the DAG with $T \rightarrow Y$, $U_1 \rightarrow T$, $U_1 \rightarrow C$, $U_2 \rightarrow C$, $U_2 \rightarrow Y$, where U_1, U_2 are unobserved and C is observed. The only back-door path is $T \leftarrow U_1 \rightarrow C \leftarrow U_2 \rightarrow Y$. Node C is a *collider* on this path, which is blocked by default: $P(y \mid \mathrm{do}(t)) = P(y \mid t)$.
Now suppose a researcher naively includes C in the adjustment set. Adding C to $\mathbf{S}$ *activates* the collider, opening the back-door path. The set $\{C\}$ fails Condition 2: it creates a back-door path rather than blocking one. This is called *M-bias*.

Theorem: Back-Door Adjustment Formula

If $\mathbf{S}$ satisfies the back-door criterion for the effect of T on Y in $\mathcal{G}$ (Pearl 1993), and

$$P(T{=}t \mid \mathbf{S}{=}\mathbf{s}) \;>\; 0 \quad \text{for all } \mathbf{s} \text{ with } P(\mathbf{S}{=}\mathbf{s}) > 0, \tag{positivity}$$

then:

$$P(y \mid \mathrm{do}(T{=}t)) \;=\; \int P(y \mid T{=}t,\; \mathbf{S}{=}\mathbf{s})\, dP(\mathbf{s}). \tag{3.1}$$

Equivalently, for any measurable function h:

$$\mathbb{E}[h(Y) \mid \mathrm{do}(T{=}t)] \;=\; \mathbb{E}_{\mathbf{S}}\Big[\mathbb{E}[h(Y) \mid T{=}t,\; \mathbf{S}]\Big]. \tag{3.2}$$

Remark: Positivity

The positivity (or overlap) condition requires every value of the treatment $T{=}t$ to be observable within each stratum of $\mathbf{S}$. It is a *data-support* condition, not a graphical one: the back-door criterion is a property of the DAG, but positivity is a property of the joint distribution P. Both must hold simultaneously for the adjustment formula to be well-defined.

3.2.3 The Adjustment Formula: Standardization

Equation Equation 3.1 is also known as the *standardization formula* or, in the epidemiological literature, the *g-formula* (Robins 1986). The interpretation is important:

- $P(y \mid t, \mathbf{s})$ is the stratum-specific conditional distribution of Y given $T{=}t$ and $\mathbf{S}{=}\mathbf{s}$. Within each stratum, T is not confounded, so this conditional distribution *is* causal.
- The outer integration $\int \cdots \, dP(\mathbf{s})$ weights the strata by the *population marginal* distribution of $\mathbf{S}$, rather than the treatment-conditional distribution.
- In a randomized experiment, $\mathbf{S} = \emptyset$ satisfies the back-door criterion, and Equation 3.1 reduces to $P(y \mid \mathrm{do}(t)) = P(y \mid t)$.

Example: Back-Door Adjustment — A Numerical Walkthrough

Use the single-confounder graph $T \leftarrow C \rightarrow Y$, $T \rightarrow Y$ with all variables binary: C = disease severity, T = treatment, Y = recovery.

C	$P(C)$	$P(T{=}1 \mid C)$	$P(Y{=}1 \mid T{=}0, C)$	$P(Y{=}1 \mid T{=}1, C)$
0	0.50	0.20	0.10	0.40
1	0.50	0.80	0.50	0.80

The true ACE is 0.30 (treatment effect within each stratum).
Naïve estimate: $P(Y{=}1 \mid T{=}1) - P(Y{=}1 \mid T{=}0) = 0.72 - 0.18 = \mathbf{0.54}$.
Back-door adjusted estimate: weighting by $P(C)$:

$$P(Y{=}1 \mid \mathrm{do}(T{=}1)) = 0.40 \times 0.5 + 0.80 \times 0.5 = 0.60,$$

$$P(Y{=}1 \mid \mathrm{do}(T{=}0)) = 0.10 \times 0.5 + 0.50 \times 0.5 = 0.30.$$

Adjusted difference: $0.60 - 0.30 = \mathbf{0.30}$ — matching the true ACE.
The naïve estimate overstates the causal effect by 0.24 because treated patients are predominantly severe ($C{=}1$), who recover at higher rates even without treatment.

Example: Back-Door Identification in the Education Example

The Education–Earnings DAG from Chapter 2: edges $N \rightarrow E$, $B \rightarrow E$, $B \rightarrow Y$, $E \rightarrow Y$.
Back-door paths from E to Y: Only $E \leftarrow B \rightarrow Y$.
Does $\{B\}$ satisfy the criterion? (1) B is not a descendant of E. (2) $\{B\}$ blocks $E \leftarrow B \rightarrow Y$ (fork at B).

$$P(y \mid \mathrm{do}(E{=}e)) = \sum_b P(y \mid e, b)\, P(b).$$

Does $\{N\}$ work? N is not a descendant of E, but $\{N\}$ does *not* block $E \leftarrow B \rightarrow Y$. **Fails.**
Does $\{N, B\}$ work? Both conditions hold. **Valid** (though redundant).

Remark: Multiple Valid Adjustment Sets

(a) Both $\{B\}$ and $\{N, B\}$ satisfy the back-door criterion here. Any such set yields the same interventional distribution; the identification target is unique even though the adjustment sets are

not.
(b) Supersets of a valid adjustment set are not automatically valid. Adding a variable can violate Condition 2 if that variable is a collider on some back-door path (as in the M-bias example). This must be verified from the graph, not assumed.
(c) Smaller valid sets are often preferable in practice. Every variable added to $\mathbf{S}$ must also satisfy positivity. Adjusting for fewer variables reduces variance and avoids practical positivity failure.

3.3 The Front-Door Criterion

The back-door criterion requires that all confounders be observed. The front-door criterion provides identification in a different configuration: when there exists a measured *mediator* M on the causal path from T to Y with specific independence properties.

3.3.1 The Configuration and Criterion

Definition: Front-Door Criterion

A set of variables $\mathbf{M}$ satisfies the *front-door criterion* for the effect of T on Y in $\mathcal{G}$ if:
1. $\mathbf{M}$ intercepts all directed paths from T to Y (all causal paths pass through $\mathbf{M}$).
2. There are no unblocked back-door paths from T to $\mathbf{M}$ (the effect of T on $\mathbf{M}$ is not confounded).
3. All back-door paths from $\mathbf{M}$ to Y are blocked by T (conditioning on T closes all confounding of the $\mathbf{M} \to Y$ effect).

Example: A Single Mediator with Unobserved Confounding

Consider the DAG $T \to M \to Y$ with $U \to T$ and $U \to Y$, where M is observed and U is unobserved.



Does M satisfy the front-door criterion?
1. *M intercepts all directed paths from T to Y.* The only directed path is $T \to M \to Y$.
2. *No unblocked back-door paths from T to M.* There is no edge $U \to M$, so no back-door path exists.
3. *All back-door paths from M to Y are blocked by T.* The only back-door path is $M \leftarrow T \leftarrow U \to Y$. Conditioning on T blocks this path (chain at T).

All three conditions hold, so the front-door formula applies:

$$P(y \mid \mathrm{do}(T{=}t)) = \sum_m P(m \mid t) \sum_{t'} P(y \mid t', m)\, P(t').$$

Example: When the Front-Door Criterion Fails

Case (a): Condition 2 fails — the $T \to M$ link is confounded. Adding the edge $U \to M$ to the prototypical graph creates an unblocked back-door path $T \leftarrow U \to M$ from T to M. Stage 1 of the front-door formula uses $P(M{=}m \mid T{=}t)$ as if the $T \to M$ link were unconfounded. But with $U \to M$, the association between T and M is partly due to their shared cause U, so Stage 1 no longer estimates $P(m \mid \mathrm{do}(T{=}t))$.
Case (b): Condition 3 fails — the $M \to Y$ link has an extra unobserved confounder. Adding a second unobserved variable V with $V \to M$ and $V \to Y$ creates a back-door path $M \leftarrow V \to Y$ from M to Y that is *not* blocked by conditioning on T. Stage 2 uses $\sum_{t'} P(y \mid t', m)\, P(t')$ as the back-door-adjusted effect of M on Y with T as adjustment, but T is no longer a valid adjustment set.

Modification	Condition violated	Stage broken
Add $U \to M$	Cond. 2: $T \to M$ confounded	Stage 1
Add $V \to M$ and $V \to Y$	Cond. 3: $M \to Y$ confounded by V	Stage 2

3.3.2 The Front-Door Formula

Theorem: Front-Door Adjustment Formula

If M satisfies the front-door criterion for the effect of T on Y (Pearl 1995), and $P(t) > 0$ for all t, then:

$$P(y \mid \mathrm{do}(T{=}t)) \;=\; \int \underbrace{\Big[\int P(y \mid T{=}t',\; M{=}m)\, dP(t')\Big]}_{\text{Stage 2}} \underbrace{dP(m \mid t)}_{\text{Stage 1}}. \tag{3.3}$$

Equivalently, writing $\mu(t', m) = \mathbb{E}[Y \mid T{=}t',\; M{=}m]$:

$$\mathbb{E}[Y \mid \mathrm{do}(T{=}t)] \;=\; \mathbb{E}_{M|T=t}\Big[\mathbb{E}_T[\mu(T,\, M)]\Big]. \tag{3.4}$$

Stage 1: $dP(m \mid t)$ is the distribution of the mediator M given $T{=}t$. Front-door condition 2 guarantees no confounding between T and M.
Stage 2: $\int P(y \mid t',\, m)\, dP(t')$ is the back-door-adjusted causal effect of M on Y, with T serving as the adjustment variable.

The formula is the composition of two back-door applications. (1) Identify $P(m \mid \mathrm{do}(T{=}t))$: by condition 2, no back-door paths from T to M exist, so $P(m \mid \mathrm{do}(T{=}t)) = P(m \mid T{=}t)$. (2) Identify $P(y \mid \mathrm{do}(M{=}m))$: by condition 3, T is a valid back-door adjustment set for the $M \to Y$ effect, so $P(y \mid \mathrm{do}(M{=}m)) = \int P(y \mid T{=}t',\; M{=}m)\, dP(t')$. (3) Chain the two stages via condition 1.

Example: Smoking, Tar, and Cancer

The classic front-door example (Pearl 1995): $T =$ smoking, $M =$ tar deposits in lungs, $Y =$ lung cancer, $U =$ genetic predisposition (unobserved). The graph is $T \to M \to Y$ with $U \to T$ and $U \to Y$. There is no direct $T \to Y$ path.
Conditions 1–3 are satisfied. The front-door formula gives:

$$P(\text{cancer} \mid \mathrm{do}(T{=}\text{smoke})) = \int \Big[\int P(\text{cancer} \mid t', m)\, dP(t')\Big]\, dP(m \mid \text{smoke}),$$

which is entirely computable from the observed joint distribution of (T, M, Y), without any data on U.

Example: Front-Door Formula — Binary Numerical Computation

Let $T, M, Y \in \{0, 1\}$ with U unobserved and the graph $T \to M \to Y$, $U \to T$, $U \to Y$.

T	M	Y	$P(T, M, Y)$
0	0	0	0.360
0	0	1	0.040
0	1	0	0.050
0	1	1	0.050
1	0	0	0.070
1	0	1	0.030
1	1	0	0.120
1	1	1	0.280

The inner sum (Stage 2) for each value of M:

$$S_0 = \sum_{t'} P(Y{=}1 \mid t', M{=}0)\, P(t') = 0.1 \times 0.5 + 0.3 \times 0.5 = 0.20,$$

$$S_1 = \sum_{t'} P(Y{=}1 \mid t', M{=}1)\, P(t') = 0.5 \times 0.5 + 0.7 \times 0.5 = 0.60.$$

Applying the outer sum (Stage 1):

$$P(Y{=}1 \mid \mathrm{do}(T{=}1)) = 0.2 \times 0.20 + 0.8 \times 0.60 = 0.52,$$

$$P(Y{=}1 \mid \mathrm{do}(T{=}0)) = 0.8 \times 0.20 + 0.2 \times 0.60 = 0.28.$$

ACE $= 0.52 - 0.28 = 0.24$. The naïve estimate $0.62 - 0.18 = 0.44$ is almost twice the true causal effect.

Comparing Back-Door and Front-Door

The two criteria are *not* nested. Neither implies the other.

- The back-door criterion requires all confounders to be observed (or blocked by observed variables), but places no restriction on mediators.
- The front-door criterion requires an observed mediator intercepting all causal paths, but the confounder U can be entirely unobserved.

There exist graphs where neither criterion alone applies, but where the full do-calculus (or the ID algorithm) can still identify the causal effect.

3.4 The Three Rules of Do-Calculus

Having seen two concrete identification strategies, we now develop the general algebraic machinery that underlies both. Pearl (1995) introduced the three rules of do-calculus as a system for manipulating expressions involving the do-operator: each rule licenses a transformation that removes or exchanges terms, bringing the expression one step closer to a purely observational formula.

Remark: First Reading / Second Reading

On a first reading, focus on the qualitative role of the three rules: Rule 1 adds or removes observations, Rule 2 exchanges an action for an observation, and Rule 3 adds or removes actions. The detailed graph subscripts and formal statements can be absorbed gradually. Each rule tests whether a variable becomes irrelevant after the appropriate arrows have been deleted from the graph — exactly the same kind of d-separation check introduced in Chapter 2.

3.4.1 Statement of the Rules

Each rule has the form "*expression* A = *expression* B *if* $(Y \perp\!\!\!\perp Z \mid X, W)_{\mathcal{G}'}$", where $\mathcal{G}'$ is a specific intervention graph built from $\mathcal{G}$ by two deletions.

Subscript	**Step 1: delete**	**Step 2: delete**
$\mathcal{G}_{\overline{X}\,\underline{Z}}$	all arrows *into* X	all arrows *out of* Z
$\mathcal{G}_{\overline{X}\,\overline{Z(W)}}$	all arrows *into* X	arrows *into* Z except those from W
$\mathcal{G}_{\overline{X}\,\overline{Z(\overline{W})}}$	all arrows *into* X	arrows *into* Z that are not from ancestors of W in $\mathcal{G}_{\overline{X}}$

Theorem: The Three Rules of Do-Calculus

Let $\mathcal{G}$ be a DAG over variables $\mathcal{V}$ (Pearl 1995), and let $X, Y, Z, W \subseteq \mathcal{V}$ be disjoint sets. The rules hold for any distribution P Markov with respect to $\mathcal{G}$.
Rule 1 (Insertion/Deletion of Observations).

$$P(y \mid \mathrm{do}(x), z, w) \;=\; P(y \mid \mathrm{do}(x), w) \quad \text{if} \quad (Y \perp\!\!\!\perp Z \mid X, W)_{\mathcal{G}_{\overline{X}\,\underline{Z}}}.$$

If Y and Z are d-separated in the graph with arrows into X deleted *and* arrows out of Z deleted, then conditioning on Z has no effect on the distribution of Y.
Rule 2 (Action/Observation Exchange).

$$P(y \mid \mathrm{do}(x), \mathrm{do}(z), w) \;=\; P(y \mid \mathrm{do}(x), z, w) \quad \text{if} \quad (Y \perp\!\!\!\perp Z \mid X, W)_{\mathcal{G}_{\overline{X}\,\overline{Z(W)}}}.$$

When the condition holds, intervening on Z is equivalent to conditioning on Z.
Rule 3 (Insertion/Deletion of Actions).

$$P(y \mid \mathrm{do}(x), \mathrm{do}(z), w) \;=\; P(y \mid \mathrm{do}(x), w) \quad \text{if} \quad (Y \perp\!\!\!\perp Z \mid X, W)_{\mathcal{G}_{\overline{X}\,\overline{Z(W)}}}.$$

When the condition holds, the intervention $\mathrm{do}(z)$ can be deleted entirely.

Remark

Each rule is a *conditional independence statement in a specific intervention graph.* The three rules cover all ways of adding or removing terms from a do-expression: Rule 1 handles observations; Rule 2 exchanges actions for observations; Rule 3 deletes actions. Any sequence of such operations that converts a do-expression into a purely observational expression is an *identification proof.*

Rule	**Interpretation**
Rule 1	Insert or delete *observations* from a conditioning set
Rule 2	Exchange an *intervention* $\mathrm{do}(z)$ for an *observation* z
Rule 3	Insert or delete *interventions* from an expression

3.4.2 Intuition for Each Rule

Rule 1 — Adding/Removing Observations. The condition $(Y \perp\!\!\!\perp Z \mid X, W)_{\mathcal{G}_{\overline{X}\,\underline{Z}}}$ says: in the post-$\mathrm{do}(x)$ world, if we further deactivate Z's causal influence (delete its outgoing arrows), Y and Z become independent given W. This means Z carries no information about Y that is not already in W, so it can be freely inserted or removed from the conditioning set.

Example: Rule 1 — Deleting a Redundant Observation

Graph: $Z \to T \to Y$, no other edges. Suppose we intervene on T.
In $\mathcal{G}_{\overline{T}\,\underline{Z}}$: delete arrows into T (removes $Z \to T$), then delete arrows out of Z (nothing remains). Node Z is now isolated, so $(Y \perp\!\!\!\perp Z \mid T)$ holds trivially. Rule 1 gives:

$$P(y \mid \mathrm{do}(t), z) \;=\; P(y \mid \mathrm{do}(t)).$$

Once we fix T externally, knowing Z (the former cause of T) carries no additional information about Y.

Rule 2 — Swapping Action for Observation. This is the most frequently used rule. If the only paths from Z to Y are causal (forward) paths, then intervening on Z and conditioning on Z have the same effect on Y — there is no confounding between Z and Y.

Example: Rule 2 — Action/Observation Exchange

Graph: $X \to Z \to Y$. Form $\mathcal{G}_{\overline{X}\,\overline{Z}}$: delete arrows into X (none) and all arrows into Z (removes $X \to Z$). Node Z has no incoming arrows. The only path from Z to Y is the causal $Z \to Y$; $(Y \perp\!\!\!\perp Z)_{\mathcal{G}_{\overline{X}\,\overline{Z}}}$ holds. Rule 2 gives:

$$P(y \mid \mathrm{do}(x), \mathrm{do}(z)) \;=\; P(y \mid \mathrm{do}(x), z).$$

Y depends on Z only through Z's value, not through the mechanism that generated Z.

Rule 3 — Deleting an Intervention. The condition says that Z has no causal path to Y that is not blocked by X or W in the appropriate intervention graph. Setting Z to any value has no effect on Y, so $\mathrm{do}(z)$ can be dropped.

Example: Rule 3 — Deleting an Irrelevant Intervention

Graph: $Z \to X \to Y$. Form $\mathcal{G}_{\overline{X}\,\overline{Z}}$: delete all arrows into X (removes $Z \to X$), then delete all arrows into Z (none). Node Z is now isolated; $(Y \perp\!\!\!\perp Z)_{\mathcal{G}_{\overline{X}\,\overline{Z}}}$ holds trivially. Rule 3 gives:

$$P(y \mid \mathrm{do}(x), \mathrm{do}(z)) \;=\; P(y \mid \mathrm{do}(x)).$$

Once X is fixed externally, Z's only route to Y (through X) is severed.

Example: A Two-Rule Derivation — the Proxy Variable Graph

Graph. $T \to Y$, $U \to T$, $U \to S$, $S \to Y$ (U unobserved, S observed).
Step 1. Introduce S by the law of total probability:

$$P(y \mid \mathrm{do}(t)) \;=\; \sum_s P(y \mid \mathrm{do}(t),\, s)\; P(s \mid \mathrm{do}(t)).$$

Step 2. Rule 3 applied to $P(s \mid \mathrm{do}(t))$. Form $\mathcal{G}_{\overline{T}}$ by deleting $U \to T$. In this graph, T has no parents and no edge connects T to S. $(S \perp\!\!\!\perp T)_{\mathcal{G}_{\overline{T}}}$ holds, so $P(s \mid \mathrm{do}(t)) = P(s)$.
Step 3. Back-door d-separation applied to $P(y \mid \mathrm{do}(t), s)$. Form $\mathcal{G}_{\underline{T}}$ by deleting $T \to Y$. The only back-door path $T \leftarrow U \to S \to Y$ is blocked by conditioning on S. Therefore $P(y \mid \mathrm{do}(t), s) = P(y \mid t, s)$.
Combining:

$$P(y \mid \mathrm{do}(t)) \;=\; \sum_s P(y \mid t,\, s)\; P(s).$$

3.4.3 Proofs of the Identification Formulas

Proof of the Back-Door Formula (Theorem 3.1)

Graph structure: $T \to Y$, unobserved $U \to T$, $U \to Y$, and observed $\mathbf{S}$ blocking the back-door path $T \leftarrow U \to Y$.
Step 1: Introduce S by the law of total probability.

$$P(y \mid \mathrm{do}(t)) = \int P(y \mid \mathrm{do}(t),\ \mathbf{s})\, dP(\mathbf{s} \mid \mathrm{do}(t)).$$

Step 2: Simplify $P(\mathbf{s} \mid \mathrm{do}(t))$. Because $\mathbf{S}$ contains no descendants of T (condition 1), the graphical condition $(\mathbf{S} \perp\!\!\!\perp T)_{\mathcal{G}_{\overline{T}}}$ holds. Rule 3 gives $P(\mathbf{s} \mid \mathrm{do}(t)) = P(\mathbf{s})$.
Step 3: Convert $P(y \mid \mathrm{do}(t), \mathbf{s})$ to an observation. Condition 2 implies $(Y \perp\!\!\!\perp T \mid \mathbf{S})_{\mathcal{G}_{\underline{T}}}$. By the truncated factorization: $P(y \mid \mathrm{do}(t), \mathbf{s}) = P(y \mid t, \mathbf{s})$.
Combining:

$$P(y \mid \mathrm{do}(t)) = \int P(y \mid t, \mathbf{s})\, dP(\mathbf{s}). \qquad \square$$

Proof of the Front-Door Formula (Theorem 3.2)

Graph structure: $T \to M \to Y$, unobserved $U \to T$, $U \to Y$, no direct $T \to Y$ edge.
Step 1: Introduce M by the law of total probability.

$$P(y \mid \mathrm{do}(t)) = \int P(y \mid \mathrm{do}(t),\ m)\, dP(m \mid \mathrm{do}(t)).$$

Step 2: Simplify $P(m \mid \mathrm{do}(t))$. In $\mathcal{G}_{\overline{T}}$, there are no back-door paths from T to M by front-door condition 2. Rule 2 applies: $P(m \mid \mathrm{do}(t)) = P(m \mid t)$.
Step 3a: $P(y \mid \mathrm{do}(t), m) = P(y \mid \mathrm{do}(m))$. In $\mathcal{G}_{\overline{T}}$, U and T become independent, and since there is no arrow $U \to M$ (condition 2), U is also independent of M. Therefore $P(y \mid \mathrm{do}(t), m) = \int P(y \mid m, u)\, dP(u) = P(y \mid \mathrm{do}(m))$.
Step 3b: $P(y \mid \mathrm{do}(m)) = \int P(y \mid m, t')\, dP(t')$. By front-door condition 3, $\{T\}$ satisfies the back-door criterion for the effect of M on Y, and the back-door formula gives the result.
Assembling:

$$P(y \mid \mathrm{do}(t)) = \int \left[\int P(y \mid t',\, m)\, dP(t') \right] dP(m \mid t). \qquad \square$$

3.5 The Do-Calculus is Complete

A natural question after seeing the three rules is: are they enough? Could there be a graph where the causal effect is identifiable in principle, but none of the rules can derive a purely observational expression? Shpitser and Pearl (2006) answered this definitively.

Throughout Sections Section 3.1–Section 3.4 we have represented latent common causes as explicit unobserved nodes. The completeness theorem uses a more general graph class that encodes latent common causes compactly as *bidirected edges*: a bidirected edge $X \leftrightarrow Y$ stands for an unobserved variable U with $U \to X$ and $U \to Y$. A DAG augmented with such bidirected edges is called a *semi-Markovian DAG*.

Theorem: Completeness of the Do-Calculus

Let $\mathcal{G}$ be a semi-Markovian DAG. A causal quantity $P(y \mid \mathrm{do}(t))$ is identifiable from P relative to $\mathcal{G}$ if and only if the do-calculus can derive a purely observational expression for it (Shpitser and Pearl 2006).

Example: The Bow Graph — The Simplest Non-Identifiable Structure

The *bow graph* is the semi-Markovian DAG with one directed edge $T \to Y$ and one bidirected edge $T \leftrightarrow Y$ (representing an unobserved U with $U \to T$ and $U \to Y$).



explicit U

semi-Markovian notation

Because U is unobserved and there are no other variables, no observed set can block the back-door path $T \leftarrow U \to Y$, and no observed mediator exists. We show $P(y \mid \mathrm{do}(t))$ is not identified by constructing two models.
Let $T, Y, U \in \{0, 1\}$ with $U \sim \mathrm{Bernoulli}(1/2)$.
Model $\mathcal{M}_1$ (pure confounding; no causal effect of T). $T = U$, $Y = U$. Both are driven entirely by U. The observed joint: $P(T{=}0, Y{=}0) = P(T{=}1, Y{=}1) = 1/2$. Under $\mathrm{do}(T{=}1)$, $Y = U$ is unchanged: $P_1(Y{=}1 \mid \mathrm{do}(T{=}1)) = 1/2$.
Model $\mathcal{M}_2$ (full causal effect). $T = U$, $Y = T$. Y is determined entirely by T. The observed joint is *identical* to $\mathcal{M}_1$. Under $\mathrm{do}(T{=}1)$, $Y = T = 1$ deterministically: $P_2(Y{=}1 \mid \mathrm{do}(T{=}1)) = 1$.
Conclusion. The two models produce identical $P(T, Y)$ yet assign different values to $P(Y{=}1 \mid$

do(T=1)) — 1/2 versus 1. Therefore $P(y \mid \mathrm{do}(t))$ is *not identified* from $P(T, Y)$ in the bow graph.

Proof Sketch and the ID Algorithm

The *ID algorithm* (Shpitser and Pearl 2006) is a procedure that takes as input a semi-Markovian DAG $\mathcal{G}$, a target intervention distribution $P(y \mid \mathrm{do}(t))$, and the observed joint distribution P, and either returns an explicit observational formula or reports non-identifiability.
Sufficiency: proved constructively via the ID algorithm. The algorithm proceeds recursively: (1) *Simplification* — remove nodes not ancestral to Y in $\mathcal{G}_{\overline{T}}$; (2) *Decomposition by c-components* — factorise over confounded components; (3) *Recursive identification* — attempt to identify each c-component factor; (4) *Failure* — if any subproblem cannot be reduced, return FAIL.
Necessity: proved by showing that whenever the ID algorithm fails, the causal effect is genuinely non-identifiable. The key graphical obstruction is the *hedge*: a pair of subgraphs (F, F') encoding a "loop" of confounding that the do-calculus cannot break.
Shpitser and Pearl (2006) prove that the ID algorithm fails if and only if $\mathcal{G}$ contains a hedge. When a hedge exists, one can explicitly construct two observationally indistinguishable models with different causal effects. Combining sufficiency and necessity: the do-calculus succeeds if and only if no hedge exists. □

Remark: Practical Implication of Completeness

Theorem 3.3 has a crucial practical implication: *if the ID algorithm reports non-identification, then no estimation method — however clever — can recover the causal effect from observational data alone*, given the assumed graph structure. Non-identification is not a limitation of a particular technique; it is a fundamental property of the causal model. Three remedies exist: impose additional structural assumptions (e.g., linearity or monotonicity) that go beyond the graph; collect additional data that break the obstruction; or target a different, identified causal quantity instead.

3.6 The Big Picture

This chapter completes the *theoretical* identification machinery of the course. The chain of reasoning begun in Chapter 1 and formalised in Chapter 2 now terminates in concrete identification formulas. Chapters 4–7 then apply this machinery to specific research designs.

The completeness theorem (Section Section 3.5) closes the loop: every causal effect that is identifiable from the observational distribution relative to the assumed graph can be derived by the do-calculus, and every effect that is not identifiable cannot be recovered by any observational method without additional assumptions or new data.

3.7 Summary

1. Two graph operations underpin the do-calculus: $\mathcal{G}_{\overline{X}}$ (delete arrows *into* X, modeling the intervention $\mathrm{do}(x)$) and $\mathcal{G}_{\underline{X}}$ (delete arrows *out of* X, used as a technical device for Rule 1 — it does not model conditioning on X).

2. The back-door criterion identifies $P(y \mid \mathrm{do}(t))$ when an observed set $\mathbf{S}$ blocks all back-door paths and contains no descendant of T. The general formula is $\int P(y \mid t, \mathbf{s})\, dP(\mathbf{s})$, equivalently $\mathbb{E}_{\mathbf{S}}[\mathbb{E}[h(Y) \mid T{=}t, \mathbf{S}]]$ in iterated-expectation form (standardization; the g-formula for $K = 1$).

3. The front-door criterion identifies $P(y \mid \mathrm{do}(t))$ via an observed mediator M intercepting all causal paths from T to Y, even when the confounder U is entirely unobserved, provided the three front-door conditions hold.

4. The do-calculus has three rules, each licensed by a d-separation condition in an appropriate intervention graph. Rule 1 adds/removes observations; Rule 2 exchanges actions for observations; Rule 3 deletes actions. The back-door and front-door formulas are each derivable as three-step sequences of these rules.



Figure 3.2: The six-level pipeline from causal assumptions to estimation.

5. **Completeness.** The do-calculus is complete for identification from observational data relative to the assumed graph (Shpitser and Pearl 2006): a causal effect is identifiable if and only if the do-calculus can derive a purely observational expression for it. If the do-calculus cannot identify the quantity in that setting, then no purely observational method can do so without additional assumptions or new data.

3.8 Problems

1. Warm-up: graph operations. Let $\mathcal{G}$ be the IV DAG with edges $Z \to T$, $T \to Y$, $U \to T$, $U \to Y$ (U unobserved).

(a) Draw $\mathcal{G}_{\overline{T}}$ and identify all paths from Z to Y that remain open.
(b) Draw $\mathcal{G}_{\underline{T}}$ and identify all paths from Z to Y that remain. For each path, classify it as a causal chain, fork, or collider path, and state whether it is open or closed by default.
(c) In $\mathcal{G}_{\overline{T}}$, is $Z \perp\!\!\!\perp Y \mid \emptyset$? Justify by listing all paths from Z to Y in $\mathcal{G}_{\overline{T}}$ and checking whether each is open or blocked.
(d) How does the absence of a direct edge $Z \to Y$ in the DAG relate to the exclusion restriction?

2. Back-door practice. Consider the DAG: $X \to T$, $X \to Y$, $T \to M$, $M \to Y$, $T \to Y$, where X is observed.

(a) List all back-door paths from T to Y.
(b) Does $\{X\}$ satisfy the back-door criterion for the effect of T on Y? Write the resulting adjustment formula.
(c) Does $\{M\}$ satisfy the back-door criterion? Explain why or why not.
(d) Does $\{X, M\}$ satisfy the back-door criterion? Identify which condition M violates.
(e) Even if the formal criterion issue in (d) were somehow set aside, explain the substantive bias that arises from conditioning on a mediator M that lies on a causal path $T \to M \to Y$.

3. Front-door identification. Consider the DAG: $T \to M \to Y$, with $U \to T$ and $U \to Y$ (unobserved U, no direct $T \to Y$ edge).

(a) Verify that M satisfies all three front-door conditions.
(b) Derive the front-door formula step by step, citing the do-calculus rule used at each step.
(c) Now add a direct edge $T \to Y$ to the graph. Does M still satisfy the front-door criterion? Explain which condition fails.

4. Identification or non-identification? For each of the following DAGs, determine whether $P(y \mid \mathrm{do}(t))$ is identified non-parametrically. If identified, state the formula; if not, explain the structural obstruction.

(a) $T \to Y$, $U \to T$, $U \to Y$, U unobserved.
(b) Same as (a) with instrument $Z \to T$ and $Z \perp\!\!\!\perp U$ (no direct $Z \to Y$ edge). Show that $P(y \mid \mathrm{do}(t))$ is **not** identified non-parametrically by constructing two binary SEMs $\mathcal{M}_1$ and $\mathcal{M}_2$ that agree on $P(Z, T, Y)$ but disagree on $P(Y{=}1 \mid \mathrm{do}(T{=}1))$.
(c) $T \to M \to Y$, $U \to T$, $U \to Y$, $U \to M$, U unobserved.
(d) $T \to Y$, $X \to T$, $X \to Y$, X observed.

5. Back-door formula: proof and uniqueness.

(a) Let $\mathcal{M}$ be a structural causal model Markov with respect to $\mathcal{G}$, and let $\mathbf{S}$ satisfy the back-door criterion for (T, Y) in $\mathcal{G}$ with positivity. Prove that $P(y \mid \mathrm{do}(t)) = \int P(y \mid t, \mathbf{s})\, dP(\mathbf{s})$.
(b) Suppose $\mathbf{S}_1$ and $\mathbf{S}_2$ both satisfy the back-door criterion for (T, Y) in $\mathcal{G}$. Deduce from part (a) that $\int P(y \mid t, \mathbf{s}_1)\, dP(\mathbf{s}_1) = \int P(y \mid t, \mathbf{s}_2)\, dP(\mathbf{s}_2)$. Interpret this result.

6. Rule sequencing on an unseen graph. Consider the graph from the two-path example: $C_1 \to T$, $C_1 \to Y$, $U \to T$, $U \to C_2$, $C_2 \to Y$, $T \to Y$, with C_1 and C_2 observed and U unobserved. Derive $P(y \mid \mathrm{do}(t)) = \sum_{c_1, c_2} P(y \mid t, c_1, c_2)\, P(c_1, c_2)$ from first principles using the three rules. For each step: (i) name the rule applied, (ii) state which arrows are deleted, and (iii) verify the required d-separation condition.

7. Non-identifiability by construction. Consider the graph $T \to M \to Y$, $U \to T$, $U \to M$, $U \to Y$ (front-door condition 2 violated because U confounds $T \to M$). Let $T, M, Y, U \in \{0, 1\}$ with

$U \sim \text{Bern}(1/2)$.

(a) Construct two SEMs $\mathcal{M}_1$ and $\mathcal{M}_2$, each compatible with the graph, such that $P_{\mathcal{M}_1}(T, M, Y) = P_{\mathcal{M}_2}(T, M, Y)$ but $P_{\mathcal{M}_1}(Y{=}1 \mid \text{do}(T{=}1)) \neq P_{\mathcal{M}_2}(Y{=}1 \mid \text{do}(T{=}1))$.

(b) Explain precisely which identification strategy fails in this graph and why, referencing (i) the back-door criterion, (ii) the front-door criterion, and (iii) the completeness theorem.

References

Berkson, Joseph. 1946. "Limitations of the Application of Fourfold Table Analysis to Hospital Data." *Biometrics Bulletin* 2 (3): 47–53.

Lauritzen, Steffen L. 1996. *Graphical Models.* Oxford University Press.

Pearl, Judea. 1993. "Comment: Graphical Models, Causality and Intervention." *Statistical Science* 8 (3): 266–69.

Pearl, Judea. 1995. "Causal Diagrams for Empirical Research." *Biometrika* 82 (4): 669–88.

Pearl, Judea. 2009. *Causality: Models, Reasoning, and Inference.* 2nd ed. Cambridge University Press.

Richardson, Thomas S., and James M. Robins. 2014. *ACE Bounds; Single World Intervention Graphs (SWIGs) and Identification of Causal Effects.* University of Washington.

Robins, James M. 1986. "A New Approach to Causal Inference in Mortality Studies with a Sustained Exposure Period—Application to Control of the Healthy Worker Survivor Effect." *Mathematical Modelling* 7 (9–12): 1393–512.

Rubin, Donald B. 1974. "Estimating Causal Effects of Treatments in Randomized and Nonrandomized Studies." *Journal of Educational Psychology* 66 (5): 688–701.

Shachter, Ross D. 1998. "Bayes-Ball: The Rational Pastime (for Determining Irrelevance and Requisite Information in Belief Networks and Influence Diagrams)." *Proceedings of the Fourteenth Conference on Uncertainty in Artificial Intelligence (UAI)*, 480–87.

Shpitser, Ilya, and Judea Pearl. 2006. "Identification of Joint Interventional Distributions in Recursive Semi-Markovian Causal Models." *Proceedings of the Twenty-First National Conference on Artificial Intelligence (AAAI)* 21: 1219–26.

Appendix A

Update Log

This page tracks updates to the lecture notes. Check here to see what has changed each week.

A.1 April 2026

A.1.1 April 11, 2026

- **Chapter 3** (The Do-Calculus and Identification Criteria) published online.
  - Intervention graphs ($\mathcal{G}_{\overline{T}}$, $\mathcal{G}_{\underline{T}}$)
  - Back-door and front-door criteria with worked numerical examples
  - Three rules of do-calculus with proofs
  - Completeness theorem (Shpitser & Pearl 2006) and the bow graph

A.1.2 April 9, 2026

- **Chapter 2** (DAGs and d-Separation) published online.
  - Chain, fork, and collider structures
  - d-Separation criterion with Bayes-ball and moral graph methods
  - Full d-separation analysis of the IV DAG
  - Collider bias and M-bias examples
- **Chapter 1** (Introduction) published online.
  - Interventional vs. observational distributions
  - The causal trinity: SEM, DAG, potential outcomes
  - Gaussian linear confounded model
  - Two-step paradigm (identification then estimation)