

Patient Name: _____ D.O.B.: _____ Date: _____

Doctor's Name: _____ Tech Name: _____

1. Place the following supplies on a clean, dry surface.
 - A. Dosage Log & Emergency Instructions
 - B. Custom Serum Vials
 - C. Two prescribed unused sterile syringes
 - D. Alcohol pads
 - E. Epi-Pen
 - F. Benadryl
 - G. Sharps Container
2. Wash hands thoroughly.
3. Refer to the Dosage Log Worksheet to determine amount of treatment.
4. Clean the selected treatment area thoroughly with an alcohol swab prior to treatment.
5. The appropriate areas to give a treatment are: (R) Tricep, (L) Tricep, Abdomen or Thigh.
6. Wipe the top of vial A with an alcohol pad before inserting needle.
7. Remove the needle cover, insert needle into vial. Use extreme caution when handling needles.
8. Turn vial upside down and draw appropriate dosage of treatment stated on your Dosage Log and remove needle from vial.
9. Make sure the bevel of the needle is up and insert the needle into injection site at 45° angle.
10. Once you have pierced the skin with the needle check the treatment site for blood.
 - A. If no blood is visible push in the plunger to inject the treatment.
 - B. If you see blood you must attempt to aspirate (Pull back on the plunger) to ensure you have not entered a blood vessel. If blood is visible in the syringe, do not push down on the syringe plunger, instead remove the needle and discard syringe and start over.
11. Remove the needle from the site at the same angle you inserted.
12. Discard syringe into a sharps container. Once a syringe pierces the skin, it is not reusable.
13. Wipe down treatment site and apply bandage if needed.
14. Keep your Epi-Pen on hand for three (3) hours after you have administered treatment.
15. Repeat these steps for vial B.

Mild allergy reactions to immunotherapy injections infrequently occur. Symptoms can be treated by the application of an ice pack to the affected area and the administration of an appropriate dose of oral Benadryl.

1. Itching at or around the injection site.
2. A limited skin flushing or redness at or around the injections site.
3. A limited number of welts (urticaria) at or around the injections site.
4. Swelling (edema) at the injection site or other limited areas (e.g. extremities, lips, etc.)

- ❖ Shortness of breath
- ❖ Wheezing
- ❖ Feel like your throat is closing
- ❖ Swelling of the tongue that causes difficulty in breathing or swallowing

1. **Use the Epi-Pen**
2. **Call "911"** *Inform the "911" dispatcher that you used your Epi-Pen due to a severe allergic reaction.*
3. **Once the "911" call has been initiated, immediately take by mouth liquid 12.5mg/5ml Children's Benadryl (Diphenhydramine).** Follow the dosage instructions for your age group and/or weight listed on the back of the medication bottle.

Make sure "911" has already been informed prior to taking Benadryl.

Follow the above instructions even if you begin to feel better because the relieving effects of Benadryl and Epinephrine may only be temporary.

[illegible]

***Note: “cc” and “ml” equal same dosage amount.**

THIS DOSAGE LOG MUST BE COMPLETED, INITIALED AND BROUGHT WITH YOU TO YOUR NEXT ALLERGY APPOINTMENT. IF PATIENT LOG IS NOT BROUGHT TO APPOINTMENT, APPOINTMENT WILL BE RESCHEDULED. WE APOLOGIZE FOR ANY INCONVENIENCE BUT THIS IS A REQUIREMENT OF YOUR INSURANCE CARRIER.

Patient Name: _____ D.O.B.: _____ Date: _____

Doctor's Name: _____ Tech Name: _____

1. Place the following supplies on a clean, dry surface.
 - A. Dosage Log & Emergency Instructions
 - B. Custom Serum Vials
 - C. Two prescribed unused sterile syringes
 - D. Alcohol pads
 - E. Epi-Pen
 - F. Benadryl
 - G. Sharps Container
2. Wash hands thoroughly.
3. Refer to the Dosage Log Worksheet to determine amount of treatment.
4. Clean the selected treatment area thoroughly with an alcohol swab prior to treatment.
5. The appropriate areas to give a treatment are: (R) Tricep, (L) Tricep, Abdomen or Thigh.
6. Wipe the top of vial A with an alcohol pad before inserting needle.
7. Remove the needle cover, insert needle into vial. Use extreme caution when handling needles.
8. Turn vial upside down and draw appropriate dosage of treatment stated on your Dosage Log and remove needle from vial.
9. Make sure the bevel of the needle is up and insert the needle into injection site at 45° angle.
10. Once you have pierced the skin with the needle check the treatment site for blood.
 - A. If no blood is visible push in the plunger to inject the treatment.
 - B. If you see blood you must attempt to aspirate (Pull back on the plunger) to ensure you have not entered a blood vessel. If blood is visible in the syringe, do not push down on the syringe plunger, instead remove the needle and discard syringe and start over.
11. Remove the needle from the site at the same angle you inserted.
12. Discard syringe into a sharps container. Once a syringe pierces the skin, it is not reusable.
13. Wipe down treatment site and apply bandage if needed.
14. Keep your Epi-Pen on hand for three (3) hours after you have administered treatment.
15. Repeat these steps for vial B.

Mild allergy reactions to immunotherapy injections infrequently occur. Symptoms can be treated by the application of an ice pack to the affected area and the administration of an appropriate dose of oral Benadryl.

1. Itching at or around the injection site.
2. A limited skin flushing or redness at or around the injections site.
3. A limited number of welts (urticaria) at or around the injections site.
4. Swelling (edema) at the injection site or other limited areas (e.g. extremities, lips, etc.)

- ❖ Shortness of breath
- ❖ Wheezing
- ❖ Feel like your throat is closing
- ❖ Swelling of the tongue that causes difficulty in breathing or swallowing

1. **Use the Epi-Pen**
2. **Call “911”** Inform the “911” dispatcher that you used your Epi-Pen due to a severe allergic reaction.
3. **Once the “911” call has been initiated, immediately take by mouth liquid 12.5mg/5ml Children’s Benadryl (Diphenhydramine).** Follow the dosage instructions for your age group and/or weight listed on the back of the medication bottle.

Follow the above instructions even if you begin to feel better because the relieving effects of Benadryl and Epinephrine may only be temporary.

[illegible]

***Note: “cc” and “ml” equal same dosage amount.**

THIS DOSAGE LOG MUST BE COMPLETED, INITIALED AND BROUGHT WITH YOU TO YOUR NEXT ALLERGY APPOINTMENT. IF PATIENT LOG IS NOT BROUGHT TO APPOINTMENT, APPOINTMENT WILL BE RESCHEDULED. WE APOLOGIZE FOR ANY INCONVENIENCE BUT THIS IS A REQUIREMENT OF YOUR INSURANCE CARRIER.

Patient Name: _____ D.O.B.: _____ Date: _____

Doctor's Name: _____ Tech Name: _____

1. Place the following supplies on a clean, dry surface.
 - A. Dosage Log & Emergency Instructions
 - B. Custom Serum Vials
 - C. Two prescribed unused sterile syringes
 - D. Alcohol pads
 - E. Epi-Pen
 - F. Benadryl
 - G. Sharps Container
2. Wash hands thoroughly.
3. Refer to the Dosage Log Worksheet to determine amount of treatment.
4. Clean the selected treatment area thoroughly with an alcohol swab prior to treatment.
5. The appropriate areas to give a treatment are: (R) Tricep, (L) Tricep, Abdomen or Thigh.
6. Wipe the top of vial A with an alcohol pad before inserting needle.
7. Remove the needle cover, insert needle into vial. Use extreme caution when handling needles.
8. Turn vial upside down and draw appropriate dosage of treatment stated on your Dosage Log and remove needle from vial.
9. Make sure the bevel of the needle is up and insert the needle into injection site at 45° angle.
10. Once you have pierced the skin with the needle check the treatment site for blood.
 - A. If no blood is visible push in the plunger to inject the treatment.
 - B. If you see blood you must attempt to aspirate (Pull back on the plunger) to ensure you have not entered a blood vessel. If blood is visible in the syringe, do not push down on the syringe plunger, instead remove the needle and discard syringe and start over.
11. Remove the needle from the site at the same angle you inserted.
12. Discard syringe into a sharps container. Once a syringe pierces the skin, it is not reusable.
13. Wipe down treatment site and apply bandage if needed.
14. Keep your Epi-Pen on hand for three (3) hours after you have administered treatment.
15. Repeat these steps for vial B.

MILD ALLERGIC REACTION INSTRUCTIONS

Mild allergy reactions to immunotherapy injections infrequently occur. Symptoms can be treated by the application of an ice pack to the affected area and the administration of an appropriate dose of oral Benadryl.

1. Itching at or around the injection site.
2. A limited skin flushing or redness at or around the injections site.
3. A limited number of welts (urticaria) at or around the injections site.
4. Swelling (edema) at the injection site or other limited areas (e.g. extremities, lips, etc.)

- ❖ Shortness of breath
- ❖ Wheezing
- ❖ Feel like your throat is closing
- ❖ Swelling of the tongue that causes difficulty in breathing or swallowing

1. **Use the Epi-Pen**
2. **Call "911"** *Inform the "911" dispatcher that you used your Epi-Pen due to a severe allergic reaction.*
3. **Once the "911" call has been initiated, immediately take by mouth liquid 12.5mg/5ml Children's Benadryl (Diphenhydramine).** Follow the dosage instructions for your age group and/or weight listed on the back of the medication bottle.

Make sure "911" has already been informed prior to taking Benadryl.

Follow the above instructions even if you begin to feel better because the relieving effects of Benadryl and Epinephrine may only be temporary.

[illegible]

***Note: “cc” and “ml” equal same dosage amount.**

THIS DOSAGE LOG MUST BE COMPLETED, INITIALED AND BROUGHT WITH YOU TO YOUR NEXT ALLERGY APPOINTMENT. IF PATIENT LOG IS NOT BROUGHT TO APPOINTMENT, APPOINTMENT WILL BE RESCHEDULED. WE APOLOGIZE FOR ANY INCONVENIENCE BUT THIS IS A REQUIREMENT OF YOUR INSURANCE CARRIER.

Doctor's Name: _____ **Tech Name:** _____

THIS DOSAGE LOG MUST BE COMPLETED, INITIALED AND BROUGHT WITH YOU TO YOUR NEXT ALLERGY APPOINTMENT. IF PATIENT LOG IS NOT BROUGHT TO APPOINTMENT, APPOINTMENT WILL BE RESCHEDULED. WE APOLOGIZE FOR ANY INCONVENIENCE BUT THIS IS A REQUIREMENT OF YOUR INSURANCE CARRIER.

D.O.B.: _____ **Date:** _____

Tech Name: _____

VIAL 6 INSTRUCTIONS

1. Place the following supplies on a clean, dry surface.
 - A. Dosage Log & Emergency Instructions
 - B. Custom Serum Vials
 - C. Two prescribed unused sterile syringes
 - D. Alcohol pads
 - E. Epi-Pen
 - F. Benadryl
 - G. Sharps Container
2. Wash hands thoroughly.
3. Refer to the Dosage Log Worksheet to determine amount of treatment.
4. Clean the selected treatment area thoroughly with an alcohol swab prior to treatment.
5. The appropriate areas to give a treatment are: (R) Tricep, (L) Tricep, Abdomen or Thigh.
6. Wipe the top of vial A with an alcohol pad before inserting needle.
7. Remove the needle cover, insert needle into vial. Use extreme caution when handling needles.
8. Turn vial upside down and draw appropriate dosage of treatment stated on your Dosage Log and remove needle from vial.
9. Make sure the bevel of the needle is up and insert the needle into injection site at 45° angle.
10. Once you have pierced the skin with the needle check the treatment site for blood.
 - A. If no blood is visible push in the plunger to inject the treatment.
 - B. If you see blood you must attempt to aspirate (Pull back on the plunger) to ensure you have not entered a blood vessel. If blood is visible in the syringe, do not push down on the syringe plunger, instead remove the needle and discard syringe and start over.
11. Remove the needle from the site at the same angle you inserted.
12. Discard syringe into a sharps container. Once a syringe pierces the skin, it is not reusable.
13. Wipe down treatment site and apply bandage if needed.
14. Keep your Epi-Pen on hand for three (3) hours after you have administered treatment.
15. Repeat these steps for vial B.

KEEP VIALS IN REFRIGERATOR (36-46°F / 2-8°C) WHEN NOT IN USE.

MILD ALLERGIC REACTION INSTRUCTIONS

Mild allergy reactions to immunotherapy injections infrequently occur. Symptoms can be treated by the application of an ice pack to the affected area and the administration of an appropriate dose of oral Benadryl. **These reactions may include:**

1. Itching at or around the injection site.
2. A limited skin flushing or redness at or around the injections site.
3. A limited number of welts (urticaria) at or around the injections site.
4. Swelling (edema) at the injection site or other limited areas (e.g. extremities, lips, etc.).

EMERGENCY SEVERE ALLERGIC REACTION INSTRUCTIONS

Immediately follow the emergency instructions below if you develop any of the following symptoms:

- Shortness of breath
- Wheezing
- Feel like your throat is closing
- Swelling of the tongue that causes difficulty in breathing or swallowing

1. Use the Epi-Pen
2. Call "911" Inform the "911" dispatcher that you used your Epi-Pen due to a severe allergic reaction.
3. Once the "911" call has been initiated, immediately take by mouth liquid 12.5mg/5ml Children's Benadryl (Diphenhydramine). Follow the dosage instructions for your age group and/or weight listed on the back of the medication bottle.

Make sure “911” has already been informed prior to taking Benadryl.

Follow the above instructions even if you begin to feel better because the relieving effects of Benadryl and Epinephrine may only be temporary.

IMMUNOTHERAPY DOSAGE SCHEDULE, LOG & EMERGENCY INSTRUCTIONS - VIAL 6

[illegible]