

A Sample Beamer Presentation

Eric Towne

Bates College

June 24, 2015

Presentation Outline

A Sample
Beamer
Presentation

Eric Towne

The Usual Suspects

A Sample
Beamer
Presentation

Eric Towne

You might think that if $f'(0) = 0$ (and f is not a constant function) then at $x = 0$, f must have

The Usual Suspects

You might think that if $f'(0) = 0$ (and f is not a constant function) then at $x = 0$, f must have

- a local maximum, or

The Usual Suspects

You might think that if $f'(0) = 0$ (and f is not a constant function) then at $x = 0$, f must have

- a local maximum, or
- a local minimum, or

The Usual Suspects

You might think that if $f'(0) = 0$ (and f is not a constant function) then at $x = 0$, f must have

- a local maximum, or
- a local minimum, or
- an inflection point.

The Usual Suspects

You might think that if $f'(0) = 0$ (and f is not a constant function) then at $x = 0$, f must have

- a local maximum, or
- a local minimum, or
- an inflection point.

If that's what you think, then you are ...

The Usual Suspects

A Sample
Beamer
Presentation

Eric Towner

You might think that if $f'(0) = 0$ (and f is not a constant function) then at $x = 0$, f must have

- a local maximum, or
- a local minimum, or
- an inflection point.

If that's what you think, then you are ... (notice that we're giving you time to reconsider!) ...

The Usual Suspects

You might think that if $f'(0) = 0$ (and f is not a constant function) then at $x = 0$, f must have

- a local maximum, or
- a local minimum, or
- an inflection point.

If that's what you think, then you are ... (notice that we're giving you time to reconsider!) ... wrong.

A Counterexample

Consider the function

$$f(x) = \begin{cases} x^2 \sin(1/x), & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

Let's see what $f'(0)$ is.

Finding $f'(0)$

A Sample
Beamer
Presentation

Eric Towne

By the definition of derivative,

$$f'(0) =$$

Finding $f'(0)$

By the definition of derivative,

$$f'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$$

Finding $f'(0)$

By the definition of derivative,

$$\begin{aligned} f'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{h^2 \sin(1/h) - 0}{h} \end{aligned}$$

Finding $f'(0)$

By the definition of derivative,

$$\begin{aligned} f'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{h^2 \sin(1/h) - 0}{h} \\ &= \lim_{h \rightarrow 0} h \sin(1/h) \end{aligned}$$

Since $-h \leq h \sin(1/h) \leq h$

Finding $f'(0)$

By the definition of derivative,

$$\begin{aligned} f'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{h^2 \sin(1/h) - 0}{h} \\ &= \lim_{h \rightarrow 0} h \sin(1/h) \end{aligned}$$

Since $-h \leq h \sin(1/h) \leq h$ and $\lim_{h \rightarrow 0} (-h) = \lim_{h \rightarrow 0} (h) = 0$,

Finding $f'(0)$

By the definition of derivative,

$$\begin{aligned} f'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{h^2 \sin(1/h) - 0}{h} \\ &= \lim_{h \rightarrow 0} h \sin(1/h) \end{aligned}$$

Since $-h \leq h \sin(1/h) \leq h$ and $\lim_{h \rightarrow 0} (-h) = \lim_{h \rightarrow 0} (h) = 0$, the
Theorem says

Finding $f'(0)$

By the definition of derivative,

$$\begin{aligned} f'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{h^2 \sin(1/h) - 0}{h} \\ &= \lim_{h \rightarrow 0} h \sin(1/h) \end{aligned}$$

Since $-h \leq h \sin(1/h) \leq h$ and $\lim_{h \rightarrow 0} (-h) = \lim_{h \rightarrow 0} (h) = 0$, the Squeeze Theorem says $f'(0) = 0$.