

Contents

1	My solutions for exercises	2
1.1	Exercise 2.2.3	2
1.1.1	(a) $a \geq a$	2
1.1.2	(b) If $a \geq b$ and $b \geq c$, then $a \geq c$	2
1.1.3	(c) If $a \geq b$ and $b \geq a$, then $a = b$	2
1.1.4	(d) $a \geq b$ iff $a + c \geq b + c$	2
1.1.5	(e) $a < b$ iff $a++ \leq b$	3
1.1.6	(f) $a < b$ iff $b = a + d$ for some positive number d	3
1.2	Exercise 2.2.4	3
1.3	Exercise 2.2.5	4
1.3.1	Lemma a. $a \neq 0 \rightarrow \exists m(N(m) \wedge a = 1 + m)$	4
1.3.2	Lemma b. $m \neq 0 \wedge n \neq 0 \rightarrow m + n \neq 1$	4
1.3.3	Lemma c. $a < b++ \wedge a \neq b \rightarrow a < b$	5
1.3.4	Proof	5
1.4	Exercise 2.2.6	5
1.4.1	Initial solution	6
1.4.2	Solution with revised writing & structure	6
1.5	The product of two natural numbers is a natural number. . .	6
1.6	Exercise 2.3.1	7
1.6.1	Lemma. $n \cdot 0 = 0$	7
1.6.2	Lemma. $m \cdot (n++) = (m \cdot n) + m$	7
1.6.3	Solution	7
1.7	Exercise 2.3.2	8
1.8	Proposition 2.3.4	8
1.9	Exercise 2.3.3	8
1.10	Proposition 2.3.6	9
1.11	Corollary 2.3.7	9
1.12	Exercise 2.3.4	9
1.13	Exercise 2.3.5	10
1.14	Exercise 3.1.1	10
1.15	Exercise 3.1.2	11
1.16	Exercise 3.1.3	11
1.16.1	If a and b are objects, then $\{a, b\} = \{a\} \cup \{b\}$	11
1.16.2	If A and B are sets, then $A \cup B = B \cup A$	11
1.16.3	If A is a set, then $A \cup A = A \cup \emptyset = \emptyset \cup A = A$	12
1.17	Exercise 3.1.4	12
1.18	Exercise 3.1.5	13
1.19	Exercise 3.1.8.	13

1.20 Exercise 3.1.11	14
1.21 Exercise 3.2.2.	14

1 My solutions for exercises

I'm practicing proof writing as well as problem solving here, so the writing style can vary from one solution to another.

1.1 Exercise 2.2.3

1.1.1 (a) $a \geq a$.

Proof. We seek an m such that $a = a + m$. Clearly 0 is such an m , and hence $a \geq a$.

1.1.2 (b) If $a \geq b$ and $b \geq c$, then $a \geq c$.

Proof. Let m and n be natural numbers such that $a = b + m$ and $b = c + n$. Then we have $a = (c + n) + m$. By the associative law for addition, $a = c + (n + m)$. Therefore $a \geq c$.

1.1.3 (c) If $a \geq b$ and $b \geq a$, then $a = b$.

Proof. Let m and n be natural numbers such that $a = b + m$ and $b = a + n$. Then we have $a = (a + n) + m$ and $a = a + (n + m)$, and so $n + m$ must be zero. Thus $n = m = 0$ by Corollary 2.2.9. Substituting 0 for m into $a = b + m$, we conclude that $a = b$.

1. Lemma. If $a + m = a$ then $m = 0$.

Proof. $a + m = a = a + 0$. Using the cancellation law gives $m = 0$.

1.1.4 (d) $a \geq b$ iff $a + c \geq b + c$.

Proof.

1. (i) If $a \geq b$ then $a + c \geq b + c$.

Let $a = b + m$ for some natural number m . Then $a + c = (b + c) + m$, hence $a + c \geq b + c$.

2. (ii) If $a + c \geq b + c$ then $a \geq b$.

Since $a + c \geq b + c$, there exists a natural number m such that $a + c = (b + c) + m$. Then $a + c = (b + m) + c$. By the cancellation law $a = b + m$. Hence $a \geq b$.

1.1.5 (e) $a < b$ iff $a++ \leq b$.

Proof. First, suppose $a < b$. Then there exists a natural number m such that

$$b = a + m$$

Since $a < b$ implies $a \neq b$, $m \neq 0$. Thus by Lemma 2.2.10, there is a natural number n such that $n++ = m$. Hence

$$b = a + (n++) = (a + n)++ = (a++) + n$$

Therefore $a++ \leq b$.

Conversely, suppose that $a++ \leq b$. Let m be a natural number such that $b = (a++) + m$. Then

$$b = (a++) + m = (a + m)++ = a + (m++)$$

Suppose, for contradiction, that $a = b$. Then

$$a = a + (m++)$$

and cancelling a gives $m++ = 0$. But that contradicts Axiom 2.3, which says $m++ \neq 0$. Thus $a \neq b$.

Since $b = a + (m++)$, we conclude that $a < b$.

1.1.6 (f) $a < b$ iff $b = a + d$ for some positive number d .

Proof. First, assume that $a < b$. Then $b = a + d$ for some natural number d . Suppose, for contradiction, that $d = 0$. then $b = a + 0 = a$, contradicting the fact that $a \neq b$. Hence $d \neq 0$, and we conclude d is positive.

Conversely, assume that $b = a + d$ for some positive number d . If $a = b$, $d = 0$ by the same argument used in (e), contradicting the fact that $d \neq 0$. Thus $a \neq b$ and we conclude that $a < b$.

1.2 Exercise 2.2.4

Proposition 2.2.13. Let a and b be natural numbers. Then exactly one of the following statements is true: $a < b$, $a = b$, $a > b$.

Proof. We first prove that at most one of the statements is true. If $a = b$, by definition $a < b$ and $a > b$ is false. If $a < b$ or $a > b$, then by definition $a \neq b$. Finally, if $a < b$ and $a > b$, then $a = b$ by Proposition 2.2.12.(c), a contradiction. Hence at most one of the statements is true.

We now prove that at least one of the statements is true. We proceed by induction on a . Consider the base case $a = 0$. Since $b = 0 + b$, we have $0 \leq b$ for all b . Thus either $0 < b$ or $0 = b$, and the base case is done. Now suppose inductively that $a < b \vee a = b \vee a > b$. We need to show that the proposition still holds on $a++$. If $a = b$, we have $a++ = b++ = b + 1$. Since $a++ \neq b$, $a++ > b$. If $a < b$, $a++ \leq b$ by Proposition 2.2.12.(e). Then either $a++ < b$ or $a++ = b$, and in either case we are done. Finally, if $a > b$ then for some natural number m , $a = b + m$. Now $a++ = (b + m)++ = b + (m++)$. Hence $a++ > b$. This closes the induction.

1.3 Exercise 2.2.5

Proposition 2.2.14 (Strong principle of induction). Let m_0 be a natural number, and let $P(m)$ be a property pertaining to an arbitrary natural number m . Suppose that for each $m \geq m_0$, we have the following implication: If $P(m')$ is true for all natural numbers $m_0 \leq m' < m$, then $P(m)$ is also true. (In particular, this means that $P(m_0)$ is true, since in this case the hypothesis is vacuous.) Then $P(m)$ is true for all natural numbers $m \geq m_0$.

The solution became unnecessarily long here because I tried to give a different proof for each lemma.

1.3.1 Lemma a. $a \neq 0 \rightarrow \exists m(N(m) \wedge a = 1 + m)$.

Proof. Since a is positive, by Lemma 2.2.10 there exists a natural number m such that $a = m++$. Then we have $a = m++ = 0 + m++ = (0 + m)++ = (0++) + m = 1 + m$.

1.3.2 Lemma b. $m \neq 0 \wedge n \neq 0 \rightarrow m + n \neq 1$.

Proof 1. Suppose $m + n = 1$. by Lemma 2.2.10, there exist natural numbers p and q such that $p++ = m$ and $q++ = n$. then $m + n = (p++) + (q++) = ((p + q)++) = 1 = 0++$. by Axiom 2.4. we infer $(p + q)++ = 0$, contradicting axiom 2.3., i.e., $(p + q) \neq 0$. hence $m + n \neq 1$.

Proof 2. Suppose $m + n = 1$. by Lemma 2.2.10, there exists a natural number p such that $p++ = m$. then $m + n = (p++) + n = (p + n)++ = 1 = 0++$. by axiom 2.4 we infer $p + n = 0$. but since n is positive, $p + n \neq 0$. By Proposition 2.2.8, a contradiction. Hence $m + n \neq 1$.

1.3.3 Lemma c. $a < b++ \wedge a \neq b \rightarrow a < b$.

Proof 1 (direct). By Proposition 2.2.12.(f), $b++ = a + n$ for some positive number n . Since n is positive, by Lemma a there exists an m such that $n = m + 1$. Now $b++ = b + 1 = a + n = a + (m + 1) = (a + m) + 1$. Cancelling 1 from both sides yields $b = a + m$. Thus $b \geq a$. Since we also have $a \neq b$, we conclude that $b > a$.

Proof 2 (indirect). Suppose $a < b$ is false. Then by Proposition 2.2.13, $a > b$ must hold. Now choose two positive numbers n and m such that $b++ = a + n$ and $a = b + m$. Now $b++ = b + 1 = a + n = (b + m) + n = b + (m + n)$. Cancelling b from both sides gives $m + n = 1$. But this contradicts Lemma b, since m and n are both positive. Hence $b > a$.

1.3.4 Proof

Proof. Let $Q(n)$ be the property that $P(m)$ is true for all $m_0 \leq m < n$. We prove by induction that $Q(n)$ is true for every natural number n .

If $n = 0$ then $Q(0)$ is vacuously true, since there's no m such that $m < 0$.

Now suppose inductively that $Q(n)$. Let m satisfy $m_0 \leq m < n++$. We must show $P(m)$. We can think of two cases here: $m = n$ and $m \neq n$.

First, consider the case $m = n$. Notice that by the inductive hypothesis, $P(m')$ for all $m_0 \leq m' < n$. It follows that $P(n)$, i.e., $P(m)$, from the hypothesis of the theorem, and we are done.

Now if $m \neq n$, by Lemma c $m < n$. Thus by the inductive hypothesis we infer that $P(m)$.

This closes the induction, and we have proven that $Q(n)$ holds for every natural number n .

Using this result, we now show the conclusion of the theorem, that is, $P(m)$ for all $m \geq m_0$. For every natural number $m \geq m_0$, $Q(m++)$ holds, since $m++$ is a natural number. Together with the fact that $m_0 \leq m < m++$, we conclude that $P(m)$. This demonstrates the strong principle of induction.

1.4 Exercise 2.2.6

Let n be a natural number, and let $P(m)$ be a property pertaining to the natural numbers such that whenever $P(m++)$ is true, then $P(m)$ is true. Suppose that $P(n)$ is also true. Prove that $P(m)$ is true for all natural numbers $m \leq n$; this is known as the principle of backwards induction.

1.4.1 Initial solution

1. Lemma. If $m \leq 0$ then $m = 0$.

Proof. Let $m \leq 0$. By definition $0 = m + n$ for some natural number n . But by Corollary 2.2.9, m and n must both be zero. Hence $m = 0$.

2. Solution

We induct on n . We first prove the principle holds on $n = 0$. Let m be a natural number such that $m \leq 0$. Then m must be 0, and from $P(0)$ we immediately obtain that $P(m)$.

Now suppose inductively that the principle holds on n . We have to prove the principle on $n++$. We first obtain the conclusion of the inductive hypothesis. We have $P(n++)$, so we infer $P(n)$ from the hypothesis. we have the requirements of backwards induction on n , and thus $P(m)$ is true for every natural number $m \leq n$.

Now let m be a natural number such that $m \leq n++$. If $m = n++$, then $P(n++)$, i.e., $P(m)$ and we are done. Otherwise, $m \leq n$ and from the earlier result we obtain $P(m)$. This closes the induction, and we have demonstrated the principle of backwards induction.

1.4.2 Solution with revised writing & structure

Using the same lemma as above.

Let $Q(n)$ be the property that: "if $P(n)$ is true, then $P(m)$ is true for all $m \leq n$ ". We prove $Q(n)$ for every natural number n , using induction.

Consider the base case $n = 0$. Let $m \leq 0$. Then m must be 0 by the above lemma, and since $P(0)$, we conclude that $P(m)$.

Suppose inductively that $Q(n)$. We have to prove $Q(n++)$. From $P(n++)$ and the definition of P , we have $P(n)$. Thus by the inductive hypothesis, $P(m)$ for all $m \leq n$.

Now let m be a natural number such that $m \leq n++$. If $m = n++$, then we substitute m for $n++$ in $P(n++)$, and we are done. Otherwise, $m \leq n$ and thus $P(m)$. This closes the induction.

1.5 The product of two natural numbers is a natural number.

Proof. Let n, m be natural numbers. We induct on n , keeping m fixed.

The base case is obvious, since $0 \cdot m = 0$, and 0 is a natural number.

Now suppose inductively that $n \cdot m$ is a natural number. Then $(n++) \cdot m = (n \cdot m) + n$ by Definition 2.3.1. Since $n \cdot m$ and m are natural numbers, $(n \cdot m) + n$ is so by the property of addition. This closes the induction.

1.6 Exercise 2.3.1

Lemma 2.3.2 (Multiplication is commutative). Let n, m be natural numbers. Then $n \cdot m = m \cdot n$.

1.6.1 Lemma. $n \cdot 0 = 0$.

Proof. We prove by induction. The base case $0 \cdot 0 = 0$ is straightforward from the definition. Now if $n \cdot 0 = 0$, then $(n++) \cdot 0 = (n \cdot 0) + 0 = 0 + 0 = 0$.

1.6.2 Lemma. $m \cdot (n++) = (m \cdot n) + m$.

Proof. We use induction on m . For the base case, by definition of multiplication we have $0 \cdot (n++) = 0 = 0 + 0 = (0 \cdot n) + 0$, and we are done.

Now suppose inductively that $m \cdot (n++) = (m \cdot n) + m$. Then by definition of multiplication

$$(m++) \cdot (n++) = (m \cdot (n++)) + n++ = (m \cdot n + m) + n++$$

And the properties of addition gives

$$(m \cdot n + m) + n++ = m \cdot n + (m + n)++ = m \cdot n + n + (m++)$$

Finally, applying the definition of multiplication we obtain

$$(m++) \cdot n + m++.$$

Thus $(m++) \cdot (n++) = (m++) \cdot n + m++$. This closes the induction.

1.6.3 Solution

We induct on n . Consider the base case $n = 0$. Then by definition of multiplication and the first lemma, $0 \cdot m = 0 = m \cdot 0$.

Now we prove the inductive step $(n++) \cdot m = m \cdot (n++)$. We start from the inductive hypothesis $n \cdot m = m \cdot n$. We add m to both sides and obtain $(n \cdot m) + m = (m \cdot n) + m$. By definition, LHS is equal to $(n++) \cdot m$. Also, the above lemma says that $(m \cdot n) + m$ is equal to $m \cdot (n++)$. Thus we conclude that $(n++) \cdot m = m \cdot (n++)$, and this closes the induction.

1.7 Exercise 2.3.2

Lemma 2.3.3 (Positive natural numbers have no zero divisors). Let n, m be natural numbers. Then $n \cdot m = 0$ if and only if at least one of n, m is equal to zero. In particular, if n and m are both positive, then nm is also positive.

Proof. First, suppose that $n = 0 \vee m = 0$. If $n = 0$, then by Definition 2.3.1 $n \cdot m = 0 \cdot m = 0$. Otherwise, $m = 0$, and $m \cdot n = 0$ by the same argument. Since multiplication is commutative, that leads to $n \cdot m = 0$.

Conversely, suppose that $nm = 0$, and for contradiction both n and m are positive. Then by Lemma 2.2.10, $n = p++$ for some natural p . Now $n \cdot m = (p++) \cdot m$, and by definition of multiplication $n \cdot m = p \cdot m + m$. Since $n \cdot m = 0$, we have $p \cdot m + m = 0$. By Corollary 2.2.9 it follows that $m = 0$, a contradiction. Thus at least one of n and m must be zero.

1.8 Proposition 2.3.4

Proposition 2.3.4 (Distributive Law). For any natural numbers a, b, c , we have $a(b + c) = ab + ac$ and $(b + c)a = ba + ca$.

Proof. Since multiplication is commutative, we only have to show the first identity. We induct on a , keeping b and c fixed.

For the base case $a = 0$, we have the following identities:

$$0(b + c) = 0 = 0 + 0 = 0b + 0c$$

Hence the base case is done.

Now suppose inductively that $a(b + c) = ab + ac$. Then

$$\begin{aligned} (a++) (b + c) &= a(b + c) + b + c && \text{Definition of multiplication} \\ &= (ab + ac) + b + c && \text{Inductive hypothesis} \\ &= (ab + b) + (ac + c) && \text{Commutativity and associativity of addition} \\ &= (a++)b + (a++)c && \text{Definition of multiplication} \end{aligned}$$

This proves the inductive step, and thus $a(b + c) = ab + ac$ for all natural numbers a, b, c .

1.9 Exercise 2.3.3

Proposition 2.3.5 (Multiplication is associative). For any natural numbers a, b, c , we have $(a \cdot b) \cdot c = a \cdot (b \cdot c)$.

Proof. We use induction on a . The base case $a = 0$ is straightforward: both $(0 \cdot b) \cdot c$ and $0 \cdot (b \cdot c)$ are equal to 0 by the definition of multiplication.

Now suppose inductively that $(a \cdot b) \cdot c = a \cdot (b \cdot c)$. We need to show that $((a++) \cdot b) \cdot c = (a++) \cdot (b \cdot c)$. The left-hand side is $(a \cdot b + b) \cdot c$ by definition, which is by distributive law equal to $(a \cdot b) \cdot c + b \cdot c$. The right-hand side is $a \cdot (b \cdot c) + b \cdot c$. But by the inductive hypothesis $(a \cdot b) \cdot c = a \cdot (b \cdot c)$, so we conclude that both sides are equal. This closes the induction.

1.10 Proposition 2.3.6

Proposition 2.3.6 (Multiplication preserves order). If a, b are natural numbers such that $a < b$, and c is positive, then $ac < bc$.

Proof. By Proposition 2.2.12.(f), there exists a positive number m such that $b = a + m$. Then $bc = (a + m)c = ac + mc$. Since m and c are both positive, $mc > 0$ by Lemma 2.3.3. Thus, using Proposition 2.2.12.(f) again, we can conclude that $bc > ac$.

1.11 Corollary 2.3.7

Corollary 2.3.7 (Cancellation law). Let a, b, c be natural numbers such that $ac = bc$ and c is non-zero. Then $a = b$.

Proof. We proceed by contradiction. Suppose $a \neq b$. Then by the trichotomy of order, either $a > b$ or $a < b$. If $a > b$, by Proposition 2.3.6 $ac > bc$, which contradicts $ac = bc$. We can also obtain the same result when $a < b$. Hence $a = b$.

1.12 Exercise 2.3.4

Prove the identity $(a + b)^2 = a^2 + 2ab + b^2$ for all natural numbers a, b .

Lemma. $a^1 = a$ for all natural number a .

Proof. $a^1 = a^{0++} = a^0 \cdot a = 1 \cdot a = a$.

We now prove the identity $(a + b)^2 = a^2 + 2ab + b^2$ by the following identities:

$$\begin{aligned}
(a+b)^2 &= (a+b)^1 \cdot (a+b) && \text{Definition of exponentiation} \\
&= (a+b) \cdot (a+b) && \text{Lemma} \\
&= (a+b) \cdot a + (a+b) \cdot b && \text{Distributive law} \\
&= aa + ba + ab + bb && \text{Distributive law again} \\
&= aa + ab + ab + bb && \text{Commutativity of multiplication} \\
&= aa + 1ab + ab + bb && \text{Identity of multiplication} \\
&= aa + 2ab + bb && \text{Definition of multiplication} \\
&= a^1a + 2ab + b^1b && \text{Lemma} \\
&= a^2 + 2ab + b^2 && \text{Definition of exponentiation}
\end{aligned}$$

(Since the exercise comes immediately after the definition of exponentiation, I decided to be verbose here.)

1.13 Exercise 2.3.5

Proposition 2.3.9 (Euclidean algorithm). Let n be a natural number, and let q be a positive number. Then there exist natural numbers m, r such that $0 \leq r < q$ and $n = mq + r$.

Proof. We induct on n , keeping q fixed. Consider the base case $n = 0$. Then we can choose $m = 0$ and $r = 0$, since $0 < q$ and $0 = 0q + 0$.

Now suppose inductively that there exist natural numbers m and r satisfying the conditions. We have to find m' and r' such that $0 \leq r' < q$ and $n++ = m'q + r'$.

Since $n = mq + r$, we have $n++ = (mq + r)++ = m' + (r++)$.

By the inductive hypothesis $r < q$, and thus $r++ \leq q$. So we have two cases here: $r++ < q$ or $r++ = q$.

If $r++ < q$, then $m' = m$ and $r' = r++$, so we are done. Otherwise $r++ = q$, and then $mq + (r++) = mq + q = (m+1)q$. Hence we can choose $m' = m+1$ and $r = 0$. This closes the induction.

1.14 Exercise 3.1.1

Show that the definition of equality in Definition 3.1.4 is reflexive, symmetric, and transitive.

Proof. Let A, B, C be sets. First, for any object x we have $x \in A \leftrightarrow x \in A$, since it's a tautology. Hence $A = A$. If $A = B$ then for any object x , $x \in A \leftrightarrow x \in B$. But that is equivalent to $x \in B \leftrightarrow x \in A$, which leads

to $B = A$. Finally, suppose that $A = B$ and $B = C$. Then for any object x , $x \in A \leftrightarrow x \in B$ and $x \in B \leftrightarrow x \in C$. Thus we have $x \in A \leftrightarrow x \in C$, which is equivalent to $A = C$.

1.15 Exercise 3.1.2

Using only Definition 3.1.4, Axiom 3.1, Axiom 3.2, and Axiom 3.3, prove that the sets $\emptyset$, $\{\emptyset\}$, $\{\{\emptyset\}\}$, $\{\emptyset, \{\emptyset\}\}$ are all distinct (i.e., no two of them are equal to each other).

1. $\emptyset$ is not equal to any other set listed. By Axiom 3.3, the other sets contain at least one element: $\emptyset \in \{\emptyset\}$, $\{\emptyset\} \in \{\{\emptyset\}\}$, $\emptyset \in \{\emptyset, \{\emptyset\}\}$. But by Axiom 3.2, $\emptyset$ has no elements, specifically $\emptyset \notin \emptyset$ and $\{\emptyset\} \notin \emptyset$. Hence by Definition 3.1.4, no sets listed are equal to $\emptyset$.
2. $\{\emptyset\} \neq \{\{\emptyset\}\}$. By Axiom 3.3, $\{\{\emptyset\}\}$ contains $\{\emptyset\}$, and $\{\emptyset\}$ contains $\{\emptyset\}$ if and only if $\{\emptyset\} = \emptyset$. We already have shown that $\emptyset \neq \{\emptyset\}$. Hence $\{\emptyset\}$ is in $\{\{\emptyset\}\}$ but it is not an element of $\{\emptyset\}$, and by Definition 3.1.4 $\{\emptyset\}$ is not equal to $\{\{\emptyset\}\}$.
3. $\{\emptyset\} \neq \{\emptyset, \{\emptyset\}\}$. By Axiom 3.3, $x \in \{\emptyset, \{\emptyset\}\}$ if and only if $x = \emptyset$ or $x = \{\emptyset\}$. Thus, $\{\emptyset\} \in \{\emptyset, \{\emptyset\}\}$, but we know that $\{\emptyset\}$ is not an element of $\{\emptyset\}$. Thus the two sets are distinct.
4. $\{\{\emptyset\}\} \neq \{\emptyset, \{\emptyset\}\}$. By a similar argument, $\emptyset$ is in $\{\emptyset, \{\emptyset\}\}$, but since $\emptyset \neq \{\emptyset\}$, it's not in $\{\{\emptyset\}\}$. Hence the two sets are also distinct.

1.16 Exercise 3.1.3

Prove the remaining claims in Lemma 3.1.13.

1.16.1 If a and b are objects, then $\{a, b\} = \{a\} \cup \{b\}$.

We have $x \in \{a, b\} \leftrightarrow x = a \vee x = b$. But $x = a$ is equivalent to $x \in \{a\}$ and $x = b$ is equivalent to $x \in \{b\}$. Thus $x \in \{a, b\} \leftrightarrow x \in \{a\} \vee x \in \{b\}$, and by Axiom 3.4 this is equivalent to $x \in \{a, b\} \leftrightarrow x \in \{a\} \cup \{b\}$. Hence $\{a, b\}$ and $\{a\} \cup \{b\}$ are equal sets.

1.16.2 If A and B are sets, then $A \cup B = B \cup A$.

For any object x we have the following (by Axiom 3.4):

$$\begin{aligned}
x \in A \cup B &\leftrightarrow x \in A \vee x \in B \\
&\leftrightarrow x \in B \vee x \in A \\
&\leftrightarrow x \in B \cup A
\end{aligned}$$

Hence $A \cup B = B \cup A$ by Definition 3.1.4.

1.16.3 If A is a set, then $A \cup A = A \cup \emptyset = \emptyset \cup A = A$.

Let x be an object. Then $x \in A \cup A$ if and only if $x \in A$ or $x \in A$, which is equivalent to $x \in A$. Thus $A \cup A$ and A are equal sets.

Also, $x \in A$ is equivalent to $x \in A \vee x \in \emptyset$, since $x \in \emptyset$ is always false by Axiom 3.2. So A and $A \cup \emptyset$ are equal.

Finally, by the previous argument we know that the union operation for sets is commutative, so $A \cup \emptyset$ and $\emptyset \cup A$ are also equal.

1.17 Exercise 3.1.4

Prove the remaining claims in Proposition 3.1.18.

Lemma. Let A and B be sets. If $A \subset B$, then there is an object x such that $x \in B$ and $x \notin A$.

Proof. Since $A \neq B$, there is an object x which belongs to exactly one of A and B . But if $x \in A$ and $x \notin B$, that would contradict $A \subseteq B$, which states every $x \in A$ also satisfies $x \in B$. Thus $x \in B$ and $x \notin A$.

We now prove the two remaining claims in the proposition. Let A, B, C be sets.

1. If $A \subseteq B$ and $B \subseteq A$, then $A = B$. Let x be an object. Then by Definition 3.1.15 we have $x \in A \rightarrow x \in B$ and $x \in B \rightarrow x \in A$. These two statements form the equivalence $x \in A \leftrightarrow x \in B$, which is precisely the definition of $A = B$.
2. If $A \subset B$ and $B \subset C$ then $A \subset C$.
 - Since the hypothesis implies that $A \subseteq B$ and $B \subseteq C$, $A \subseteq C$ by the preceding claim (which is in the book). Now let us prove $A \neq C$. Since $B \subset C$ implies $B \neq C$, either $B \subseteq C$ or $C \subseteq B$ is false by (1). We already have $B \subseteq C$, so $C \subseteq B$. Now, suppose for the sake contradiction that $A = C$. Then we can substitute C for A in $A \subseteq B$, obtaining $C \subseteq B$, which is a contradiction. Hence $A \neq C$.

- Alternative: Proving $A \subseteq C$ is the same as above, and we prove that $A \neq C$. Since $A \subset B$, by the above Lemma there is an element x in B which is not an element of A . Also, since $B \subset C$, x is an element of C . Thus, we have x , which is an element of C but not an element of A , demonstrating $A \neq C$.

1.18 Exercise 3.1.5

Let A, B be sets. Show that the statements $A \subseteq B$, $A \cup B = B$, $A \cap B = A$ are logically equivalent (any one of them implies the other two).

Proof. We prove the equivalences by proving three implications: $A \subseteq B \rightarrow A \cup B = B$, $A \cup B = B \rightarrow A \cap B = A$, and $A \cap B = A \rightarrow A \subseteq B$.

1. $A \subseteq B \rightarrow A \cup B = B$. Let x be an object. It is logically true that $x \in B$ implies that $x \in A$ or $x \in B$. Also, suppose $x \in A$ or $x \in B$. Since all elements of A are elements of B , in either case $x \in B$. Hence we have the equivalence $x \in A \vee x \in B \leftrightarrow x \in B$, and so the two sets are equal.
2. $A \cup B = B \rightarrow A \cap B = A$. It is clear that $x \in A$ and $x \in B$ implies $x \in A$. Now let x be an element of A . since $x \in A$ or $x \in B$ implies $x \in B$, x must also be an element of B . Hence every element of A is an element of $A \cap B$, which demonstrates the equality $A \cap B = A$.
3. $A \cap B = A \rightarrow A \subseteq B$. We have $x \in A$ and $x \in B$ if and only if $x \in A$. Thus, if $x \in A$, then also $x \in B$, which demonstrates $x \subseteq B$.

1.19 Exercise 3.1.8.

Let A, B be sets. Prove the absorption laws $A \cap (A \cup B) = A$ and $A \cup (A \cap B) = A$.

Proof. We use the results of Exercise 3.1.7 here.

1. $A \cap (A \cup B) = A$. For any set A and B , we have $A \subseteq A$ as well as $A \subseteq A \cup B$, so it follows that $A \subseteq A \cap (A \cup B)$. Also, since $A \cap B \subseteq A$ for every B , we have $A \cap (A \cup B) \subseteq A$. Hence the two sets A and $A \cap (A \cup B)$ are subsets of each other, demonstrating their equality.
2. $A \cup (A \cap B) = A$. Similarly, since $A \subseteq A$ and $A \cap B \subseteq A$, it follows that their union is also a subset of A , i.e., $A \cup (A \cap B) \subseteq A$. Also, since $A \subseteq (A \cup B)$ for any B , we have $A \subseteq A \cup (A \cap B)$. Therefore $A \cup (A \cap B) = A$.

1.20 Exercise 3.1.11

Show that the axiom of replacement implies the axiom of specification.

Proof. Let A be a set, and let $P(x)$ be a property pertaining to the elements of A . We have to show that there exists a set X such that for any object x , $x \in X$ if and only if $x \in A$ and $P(x)$.

Let $Q(x, y)$ be the statement " $x = y$ and $P(x)$ ".

We claim that for each $x \in A$, there is at most one y for which $Q(x, y)$ is true. Let $x \in A$, and let y_1, y_2 satisfy $Q(x, y_1)$ and $Q(x, y_2)$. Then $x = y_1$ and $x = y_2$, which then implies $y_1 = y_2$. Hence our claim is true, and by Axiom of replacement there exists a set Y such that for any object z , $z \in Y$ if and only if $Q(x, z)$ for some $x \in A$.

We now prove that Y is the X we are seeking, i.e., for every z , $z \in Y$ if and only if $z \in A$ and $P(z)$.

First, suppose $z \in A$ and $P(z)$. To show that $z \in Y$, we have to find an $x \in A$ such that $Q(x, z)$ holds. Indeed, $Q(z, z)$ holds, since $z = z$ and $P(z)$. Thus, z is such an x , and therefore $z \in Y$.

Conversely, let $z \in Y$. Then there exists an $x \in A$ such that $Q(x, z)$, i.e., $x = z$ and $P(x)$. Since $x = z$, it follows that $P(z)$ and $z \in A$.

Hence, Y is the set we were looking for, and this completes our proof.

1.21 Exercise 3.2.2.

Use the axiom of regularity (and the singleton set axiom) to show that if A is a set, then $A \notin A$. Furthermore, show that if A and B are two sets, then either $A \notin B$ or $B \notin A$.

1. For every set A , $A \notin A$.

Proof. Suppose, for the sake of contradiction, that there exists a set A such that $A \in A$. Consider the singleton set $\{A\}$. Since it is nonempty, axiom of regularity states that there exists an $x \in \{A\}$ such that x is either not a set, or is disjoint from $\{A\}$. By definition of $\{A\}$, x is equal to A , which is clearly is a set. It is also not disjoint from $\{A\}$, since there is an element, A , which is both elements of A and $\{A\}$. This contradicts the properties of x . Hence $A \notin A$ for every set A .

2. For every set A and B , either $A \notin B$ or $B \notin A$.

Proof. Let A and B be sets. Suppose, for the sake of contradiction, that $A \in B$ and $B \in A$. By Axiom 3.3 there exists a pair set $\{A, B\}$ such that $x \in \{A, B\}$ if and only if $x = A$ or $x = B$. Notice that neither A

nor B is disjoint from $\{A, B\}$, since B is both in A and $\{A, B\}$, and A is both in B and $\{A, B\}$. Thus, every element of $\{A, B\}$ is a set, and it is not disjoint from $\{A, B\}$, contradicting the axiom of regularity. Hence either $A \notin B$ or $B \notin A$.