KAISER PERMANENTE
HEALTHCONNECT

INPATIENT

Key Message

Effective: Immediately
Those Affected Include: ED and Inpatient Physicians, Advanced Practice Providers, Nursing, and Pharmacy

New Diabetic Ketoacidosis (DKA) Adult Two-Bag System Order Set

We replaced the DKA ADULT ED SCAL and DKA ICU IP SCAL order sets with **DKA TWO-BAG SYSTEM ENDO ICU IP SCAL**. It is to be used in both ED and Inpatient, and has an associated protocol (https://cl.kp.org/scal/cpg/kphc/dka-two-bag-system.html).

In the DKA Two-Bag System, a non-dextrose and a dextrose 10% IV fluid bag are administered simultaneously and titrated based on the Blood Glucose POCT and the protocol table below, while the insulin drip remains constant.

Blood Glucose (MG/dL) Measured Hourly	Bag 1 (Non-Dextrose)	Bag 2 (Dextrose 10%)
Greater than 250	100% of total IV fluid rate	0% of total IV fluid rate
200-250	50% of total IV fluid rate	50% of total IV fluid rate
Less than 200	0% of total IV fluid rate	100% of total IV fluid rate

A two-year pilot at Riverside and Moreno Valley hospitals shows that initiating the DKA Two-Bag System in the ED improves continuity of care, simplifies the management of insulin drip and IV fluids, allows for early closure of anion gap, and decreases the incidence of hypoglycemic events.

SSD Systems Solutions & Deployment
Intellectual Property of Kaiser Permanente Southern California
Proprietary and Confidential
For Internal Use Only
SSD Communications
Jul 2021

Important Ordering Points:

- Initiate the DKA Two-Bag System in the ED using the **DKA ADULT TWO-BAG SYSTEM ENDO ICU IP SCAL** order set.
- Specify the **Total IV Fluid Rate** in the weblink protocol order.



- The insulin drip dose must be ordered in **units/hr** to match the units used in the adult IV pump. Do not order insulin drip in weight-based units.



SSD Systems Solutions & Deployment
Intellectual Property of Kaiser Permanente Southern California
Proprietary and Confidential
For Internal Use Only
SSD Communications
Jul 2021