4. The patient is called by the Radiology Nurse and assessed for a history of allergic reaction to contrast injection, or allergy to iodine. If allergic, the patient is educated regarding the need for prophylactic treatment.

5. RN will give instructions on the medication dosing, and clarify which pharmacy of choice the patient would like to pick up the medication from.

6. The RN will enter the order under the PCP or ordering physician and make sure to check the right pharmacy for that patient when ordering the prep. RN will send an in-basket message to that physician that the steroid prep order has been placed for the ordered exam.

For Hospital Outpatient Visits (Hov):

- Diphenhydramine 50 mg orally 1 hour prior to the Imaging Study
- Prednisone 50 mg orally at the following intervals: 50 mg Thirteen (13) hours prior; 50 mg Seven (7) hours prior, and 50mg One (1) hour prior to the Imaging Study
- The patient is called by the Interventional Radiology Nurse and given the following pre-procedure instructions:
  - Date and time of the Imaging Study
  - Why Diphenhydramine and Prednisone are ordered
  - What pharmacy to pick up the medications
  - Instructions on how to take the medication
  - To have a driver available to take him/her to the Imaging Study and home after the study
  - The phone number to the nurses' office for follow-up for any additional concerns

For Hospital Inpatients Or Emergency Room Patients:

1. The Radiology Nurse will contact the patient's nurse with pre-procedure instructions as followed:

   A. Provider will place order for patient to follow schedule below.

   - Prednisone 50 mg orally at the following intervals: 50 mg Thirteen (13) hours prior; 50 mg Seven (7) hours prior, and 50mg One (1) hour prior to the Imaging Study and Diphenhydramine 50 mg orally 1 hour prior to the Imaging Study.

2. If patient is unable to take medications orally, or the study must be performed emergently, methylprednisolone 125 mg IV Push should be ordered to be given 10 minutes before the procedure only.

Pediatrics:

1. The pediatric patient is defined up to age fifteen. The formula below is the recommended guideline but dose may be modified for the patient by a Radiologist.