



Current COVID and Paxlovid Procedures


Clinical Case Quiz

Treating a COVID-Positive Patient

Today is August 30, and a 30-year-old female began experiencing a sore throat on August 25. She took a COVID test at home today, and it was positive. She calls Gcn inquiring about Paxlovid. She weighs 60 kg, and she has no other medical history. Her symptoms are currently mild. The best advice to provide her is:

- She should isolate and use supportive care measures, as she does not qualify for Paxlovid.
- Today is Day 6 of her illness, so she does not qualify for Paxlovid. Paxlovid must be given within the first 5 days.
- She should visit Kaiser for a PCR Test.
- She qualifies for Paxlovid, as she meets criteria for the medication. Today is Day 5 of her illness.

The answer is d.



Note. From *Positive Covid-19 Rapid Antigen Test* [Photograph], by Alan Tennyson, 2022, Wikimedia Commons (https://commons.wikimedia.org/wiki/File:Positive_Covid-19_Rapid_Antigen_Test.jpg).

Current Criteria for Paxlovid

Handling a Patient's Request for Paxlovid

If a member desires to treat COVID-19 with Paxlovid, please keep in mind the following criteria:

- Presence of mild to moderate symptoms within five days or less (Day 0 is the first day of symptoms) *and* age equal to or greater than 12 years old *and* weight over 40 kg.
- Contraindications include severe hepatic impairment, GFR<30, history of clinically significant hypersensitivity to Paxlovid and drugs highly dependent on CYP3A for clearance.
- Patients over the age of 70 with no history of hypertension, diabetes, or kidney disease will need an updated GFR within 6 months.
- Patients with a GFR under 60, kidney disease, hypertension, or diabetes will need a GFR within 3 months.
- Those under 70 with no history of kidney disease, hypertension, or diabetes do not need a GFR checked prior to Paxlovid treatment.
- If a member requires an updated GFR, order a stat creatinine and order the Paxlovid. Direct the member to a Kaiser facility that is capable of processing stat labs (typically those that have an urgent care).

Documentation for Paxlovid Consent

Using the Paxlovid Consent Sheet

When providing counseling for Paxlovid, please document using the Paxlovid consent sheet. This can also serve as your note, as it is quite comprehensive. Use the .RVMCPAXLOVIDCONSENT SmartPhrase. The SmartPhrase includes guidance on medications that interact with Paxlovid.

Calling in a Prescription



Note. From *Package of Paxlovid* [Photograph], by James Heilman, 2022, Wikimedia Commons (<https://commons.wikimedia.org/wiki/File:Paxlovid2022.jpg>).

Ensure Paxlovid Availability

Verify that the patient's chosen pharmacy carries Paxlovid. [Use this link to determine which pharmacy carries Paxlovid.](#)

Paxlovid Work Note Process



Handling a Member's Work Note Request

If a Member needs a work note, *do not provide an activity/Rx form. Provide a letter* indicating they have COVID. This can then be sent to their employer so they can provide further guidance. (You can also view the procedure in the images on the bottom.)

To provide a letter, please do the following:

- Go to Communications.
- Choose to send the letter to kp.org if the patient is active on kp.org.
- Enter "COVID" in the "Choose Letter" section.
- Select "COVID-19 positive result need to isolate letter."

Special thanks to Dr. Jerrica Chen for providing information regarding the letter workflow.



References

Heilman, J. (2022). *Package of Paxlovid* [Photograph]. Wikimedia Commons. <https://commons.wikimedia.org/wiki/File:Paxlovid2022.jpg>

Tennyson, A. (2022). *Positive COVID-19 Rapid Antigen Test* [Photograph]. Wikimedia Commons. https://commons.wikimedia.org/wiki/File:Positive_Covid-19_Rapid_Antigen_Test.jpg