




Tripledemic Preparedness

COVID




Note. From SARS-CoV-2 [Artistic Rendering], by Centers for Disease Control and Prevention, 2019, Wikimedia Commons (https://commons.wikimedia.org/wiki/File:SARS-CoV-2_without_background.png).

- Maternal vaccination helps provide protection against infant hospitalization during the first 3 to 6 months of life.

- COVID-19 cases and hospitalizations are increasing across the country.
- The dominant variants are Omicron subvariants.
- A new monovalent COVID-19 booster is expected to be available for KP members around 9/28/2023.
- This vaccine is directed against Omicron subvariant XBB.1.5, but it should offer protection against other Omicron subvariants.
- Indicated for ages 6 months and older.
- KP will primarily be offering the Pfizer monovalent vaccines. Moderna availability will be limited.

Influenza

- Influenza vaccines for this flu season are now available.
- Patients with egg allergies may get any influenza vaccine (egg-based or non-egg-based).
- Anyone 6 months or older should get the influenza vaccine annually.
- Pregnant patients can get vaccinated during any trimester.
- Maternal vaccination offers protection via maternal antibodies to protect against influenza during the first few months of the baby's life.



Note. From Influenza Virus [Artistic Rendering], by NIAID, 2022, Flickr (<https://www.flickr.com/photos/54591706@N02/52461389748/>).


Other influenza vaccine products

- Fluzone: High-dose vaccine for ages 65 and older.
- Flumist: Live attenuated intranasal spray for ages 2 to 49. This vaccine is not recommended for use in pregnant people, those who are immunocompromised, or people with certain medical conditions.

RSV: Vaccines and Monoclonal Antibody

- Two RSV vaccines (**Arexvy: GSK** and **Abrysvo: Pfizer**) are FDA-approved for adults age 60 and older. At this time, KP has a limited amount due to a supply chain delay.
 - Both are recombinant protein vaccines.
- It has been shown to provide some protection against severe disease for at least two RSV seasons.
- Adults 60 years and older may receive one dose of RSV vaccine using shared clinical decision-making with a provider.

Adults at Higher Risk of Severe Illness from RSV Disease



Note. From Human Respiratory Syncytial Virus [Artistic Rendering], by NIAID, 2023, Flickr (<https://www.flickr.com/photos/54591706@N02/52869900362/>).

- Chronic lung disease: asthma, COPD
- Cardiovascular disease: CHF and CAD
- Immune compromise
- Hematologic disorders
- Neurologic disorders
- Endocrine disorders: diabetes
- Kidney disease
- Liver disease
- Any other condition that puts the patient at risk of severe respiratory illness if RSV is contracted.
- Residents of nursing homes or long-term care facilities.

Higher incidence of inflammatory neurologic conditions (Guillain-Barré syndrome, acute disseminated encephalomyelitis) and atrial fibrillation were observed in RSV vaccine recipients during clinical trials. Due to the small number of inflammatory neurologic events and atrial fibrillation events in the clinical trials, it is not yet known whether these events occurred due to random chance, or whether the RSV vaccination increases the risk of these events.

Maternal RSV Vaccine (**Abrysvo: Pfizer**)

- One dose given at 32-36 weeks gestation.
- Protects infants after birth from RSV.
- Prevents RSV-associated lower respiratory tract disease from birth through 6 months old.

Long-acting monoclonal antibody (**Nirsevimab Beyfortus**): Expected to be available in October

- Reduce risk of severe RSV disease.
- Given to infants during or entering their first RSV season and less than 8 months old.
- Also consider giving to children less than 20 months old at risk of severe RSV disease entering their second RSV season.
- No increased incidence of serious adverse events among vaccine recipients.

Influenza vaccine can be given with other vaccines, including the COVID-19 vaccine, preferably in a different limb to decrease local injection site reaction.

All vaccines can be given during the same visit.

References

- Centers for Disease Control and Prevention. (2023). Clinical vaccination guidance for pregnant people. Centers for Disease Control and Prevention. https://emergency.cdc.gov/coca/calls/2023/callinfo_081023.asp
- Centers for Disease Control and Prevention. (2023). Influenza vaccination: A summary for clinicians. Centers for Disease Control and Prevention. <https://www.cdc.gov/flu/professionals/vaccination/vax-summary.htm#vaccinated>
- Centers for Disease Control and Prevention. (2019). SARS-CoV-2 [Artistic Rendering]. Wikimedia Commons. https://commons.wikimedia.org/wiki/File:SARS-CoV-2_without_background.png
- National Institute of Allergy and Infectious Diseases. (2023). Human Respiratory Syncytial Virus [Artistic Rendering]. Flickr. <https://www.flickr.com/photos/54591706@N02/52869900362/>
- National Institute of Allergy and Infectious Diseases. (2022). Influenza Virus [Artistic Rendering]. Flickr. <https://www.flickr.com/photos/54591706@N02/52461389748/>
- SCPMG News Center. (2023). Daily COVID-19 town halls for physicians. SCPMG. <https://sp-cloud.kp.org/sites/SCPMGNewsCenter/SitePages/Daily-COVID-19-Town-Halls-for-Physicians.aspx>