

Timeline Check-In

Set Up

Packages

```
# general use and cleaning
library(tidyverse)
library(here)
library(janitor)
library(snakecase)

# wrap axis labels
library(scales)

# read .xlsx files
library(readxl)

# visualize missing data
library(naniar)

# arrange plots
library(patchwork)

library(dplyr)      # data manipulation
library(broom)      # tidy statistical output
library(corrplot)
library(ggrepel)
```

Data

```
# taxonomic information
taxon_list <- read_csv(here("data", "taxon_list.csv"))

# aquatic invertebrates
aq_ins <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "Aquatic Insects")

# read in site information
sites <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "sites")

# read in water quality information
water_quality <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "Water Quality",
  na = c("", "n/a", "N/A", "over", "173+"))
```

Cleaning

sites object

```
# creating new object from sites
ncos_sites <- sites |>
  # filter to only include NCOS sites sampled four times a year
  filter(sector == "NCOS" & sampling_frequency == "Quarterly")
```

Cleaning taxon_list

```
# creating new clean object from taxon_list
taxon_list_clean <- taxon_list |>
  # converting taxon name to snake case (no spaces or capital letters,
  # only underscores)
  mutate(verbatim_name = to_any_case(verbatim_name, case = "snake"))
```

Cleaning water_quality

```
# creating new clean object from water quality
water_quality_clean <- water_quality |>
  # clean column names
  clean_names() |>
  # filtering join: only include sites in ncos_sites
  semi_join(ncos_sites, by = "site") |>
  # create new column to join with later data frames
  unite("date_site", date, site, remove = TRUE) |>
  # group by date_site column
  group_by(date_site) |>
  # calculate median pH, salinity, DO
  summarize(
    med_ph = median(dissolved_oxygen_mg_l, na.rm = TRUE),
    med_sal = median(salinity_ppt, na.rm = TRUE),
    med_do = median(dissolved_oxygen_mg_l, na.rm = TRUE)) |>
  # ungroup data frame
  ungroup() |>
  # filter out salinity outliers
  filter(date_site != "2022-08-12_NVBR")
```

Cleaning aq_ins

```
# creating new clean object from aquatic inverts
aq_ins_clean <- aq_ins |>
  # clean column names
  clean_names() |>
  # filtering join: only include sites occurring in ncos_sites
  semi_join(ncos_sites, by = c("site")) |>
  # renaming columns
  rename(date = date_on_vial) |>
  # select columns of interest
  select(
    date, sample_type, site,
    ostracod,
    copepod,
    hemiptera_corixidae_boatman,
    cladocera,
    nematode,
    diptera_ceratopogonidae,
    annelida_oligochaete,
```

```

    diptera_chironomid,
    annelida_polychaete) |>
# filter to only include "filtered beaker" samples
filter(sample_type %in% c("FB 250", "FB250")) |>
# convert the data frame to long format
pivot_longer(cols = ostracod:annelida_polychaete,
              names_to = "taxon",
              values_to = "abundance") |>
# group by date, site, taxon
group_by(date, site, taxon) |>
# calculate average abundance per liter (sum abundance divided by 7.5 liters)
summarize(ave_lit = sum(abundance, na.rm = TRUE)/7.5) |>
# ungroup data frame
ungroup() |>
# create a new column called `date_site` to join with water quality
unite("date_site", date, site, remove = FALSE) |>
# join with water quality
left_join(water_quality_clean, by = c("date_site")) |>
# join with taxon list
left_join(taxon_list_clean, by = c("taxon" = "verbatim_name")) |>
# add full names of sites
mutate(site_full = case_when(
  site == "NVBR" ~ "North of Venoco Bridge",
  site == "NEC" ~ "South of Venoco Bridge",
  site == "NMC" ~ "Main Channel",
  site == "NPB" ~ "South of Phelps Bridge",
  site == "NPB1" ~ "North of Phelps Bridge",
  site == "NPB2" ~ "Phelps Road",
  site == "NWP" ~ "West Pond",
  site == "NDC" ~ "Devereux Creek"
)) |>
# setting factor levels for site
mutate(site_full = fct_reorder(site_full, med_do, .fun = "median", .na_rm = TRUE),
       site_full = fct_rev(site_full))

```

Focus Taxa (focus_taxa)

```

# taxa to include in all figures
focus_taxa <- c("Ostracod", "Copepod", "Cladocera",
                "Hemiptera Corixidae Boatman", "Nematode")

```

Creating DO bins (creating do_category column)

```
aq_ins_clean_bins <- aq_ins_clean |>
  mutate(bins = case_when(site_full == "Phelps Road" ~ "low",
                           site_full == "North of Phelps Bridge" ~ "low",
                           site_full == "South of Venoco Bridge" ~ "moderate",
                           site_full == "South of Phelps Bridge" ~ "moderate",
                           site_full == "Devereux Creek" ~ "moderate",
                           site_full == "North of Venoco Bridge" ~ "high",
                           site_full == "Main Channel" ~ "high")) |>
  mutate(taxon = str_replace_all(taxon, "_", " "), taxon = str_to_title(taxon)) |>
  filter(taxon %in% focus_taxa) |>
  # create column that calculates log-transformed abundance
  mutate(log_ave_lit = log(ave_lit + 1)) |>
  mutate(bins = factor(bins, levels = c("low", "moderate", "high")))

# create object DO_taxa from aq_ins_clean
DO_taxa <- aq_ins_clean |>
  # remove rows that are missing median salinity values
  filter(!is.na(med_do)) |>
  # create column that calculates log-transformed abundance
  mutate(log_ave_lit = log(ave_lit + 1)) |>
  # mutates to title-case taxon names
  mutate(taxon = str_replace_all(taxon, "_", " "), taxon = str_to_title(taxon)) |>
  # only include focus taxa
  filter(taxon %in% focus_taxa) |>
  # order taxon levels by maximum average abundance per liter
  mutate(taxon = fct_reorder(taxon, ave_lit, .fun = max, .desc = TRUE)) |>
  # keep only the columns needed for the figure
  select(date, site, site_full, taxon, ave_lit, log_ave_lit, med_do)
total_abundance <- DO_taxa |>
  filter(taxon %in% focus_taxa) |>
  group_by(date, site, med_do) |>
  summarize(
    total_abundance = sum(ave_lit, na.rm = TRUE),
    log_total_abundance = log10(total_abundance + 1),
    .groups = "drop"
  )
total_abundance <- total_abundance |>
  mutate(
    do_category = case_when(
      med_do <= quantile(med_do, 0.25, na.rm = TRUE) ~ "Low DO",
```

```

    med_do <= quantile(med_do, 0.75, na.rm = TRUE) ~ "Medium DO",
    TRUE ~ "High DO"
  )
)
total_abundance$do_category <- factor(
  total_abundance$do_category,
  levels = c("Low DO", "Medium DO", "High DO")
)

```

Ideas for analysis and background

- should include some citations (not all)
- should include ideas for analyses to answer questions and some preliminary code to run those analyses

Citations:

NEED TO ADD

Spearman Correlation Analysis for Total Abundance and DO levels

```

cor.test(
  total_abundance$log_total_abundance,
  total_abundance$med_do,
  method = "spearman",
  exact = FALSE)

```

Spearman's rank correlation rho

data: total_abundance\$log_total_abundance and total_abundance\$med_do

S = 8526.3, p-value = 0.08182

alternative hypothesis: true rho is not equal to 0

sample estimates:

rho

-0.3027201

Total log-transformed invertebrate abundance had a **weak negative** relationship with dissolved oxygen, but this relationship was not statistically significant using Spearman correlation.

Correlation analysis for Each Focus Taxa and DO levels

```
# create wide dataset (one column per taxon)
taxa_wide <- DO_taxa |>
  select(date, site, taxon, log_ave_lit, med_do) |>
  pivot_wider(
    names_from = taxon,
    values_from = log_ave_lit
  )

# calculate correlation of each taxon with dissolved oxygen
# calculate Spearman correlation of each taxon with dissolved oxygen
cor_do <- taxa_wide |>
  select(-date, -site) |>
  summarize(across(-med_do, ~ cor(.x,
                                med_do,
                                method = "spearman",
                                use = "complete.obs"))) |>
  pivot_longer(cols = everything(),
    names_to = "taxon",
    values_to = "correlation")

ggplot(cor_do, aes(x = taxon, y = "Dissolved oxygen", fill = correlation)) +
  geom_tile(color = "grey70") +
  geom_point(aes(size = abs(correlation)), shape = 21, fill = "white") +


  # move labels ABOVE circles
  geom_text(aes(label = round(correlation, 2)),
    nudge_y = 0.15, # <-- this moves text up
    size = 4) +

  scale_fill_gradient2(
    low = "red",
    mid = "white",
    high = "blue",
    midpoint = 0,
    limits = c(-1, 1),
```

```

    name = "r"
  ) +
  scale_size(range = c(2, 10), guide = "none") +
  labs(
    x = "Taxon",
    y = NULL,
    title = "Spearman correlation between dissolved oxygen and invertebrate abundance"
  ) +
  theme_minimal() +
  theme(
    panel.grid = element_blank(),
    axis.text.x = element_text(angle = 45, hjust = 1)
  )

```



Most focus taxa showed **weak negative** correlations with dissolved oxygen, with Cladocera

and Copepod showing the strongest negative relationships and Nematode showing almost no relationship.

Pearson correlation for continuous DO and total community abundance

```
# Pearson correlation for continuous DO and total community abundance
cor.test(total_abundance$log_total_abundance,
         total_abundance$med_do,
         method = "pearson")
```

Pearson's product-moment correlation

```
data: total_abundance$log_total_abundance and total_abundance$med_do
t = -2.2882, df = 32, p-value = 0.02887
alternative hypothesis: true correlation is not equal to 0
95 percent confidence interval:
 -0.63289820 -0.04217141
sample estimates:
      cor
-0.3749894
```

```
# r = -0.3749894, p = 0.02887
# there is a significant negative correlation between abundance and continous DO
```

There was a **statistically significant negative linear relationship** between median dissolved oxygen and log-transformed total invertebrate abundance, suggesting that abundance decreased as DO increased.

Kruskal-Wallis test comparing abundance across DO categories

```
kruskal.test(log_total_abundance ~ do_category,
             data = total_abundance)
```

Kruskal-Wallis rank sum test

```
data: log_total_abundance by do_category
Kruskal-Wallis chi-squared = 2.0001, df = 2, p-value = 0.3679
```

```
# Kruskal-Wallis chi-squared = 3.2212, df = 2, p-value = 0.1998
# there is not a significant difference between abundance and DO categories
```

Total log-transformed invertebrate abundance **did not differ significantly** across low, moderate, and high DO categories.

Kruskal-Wallis test between abundance and bins for Copepods

```
kruskal.test(log_ave_lit ~ bins,
             data = aq_ins_clean_bins,
             subset = taxon == "Copepod")
```

Kruskal-Wallis rank sum test

data: log_ave_lit by bins

Kruskal-Wallis chi-squared = 4.405, df = 2, p-value = 0.1105

```
# Kruskal-Wallis chi-squared = 4.405, df = 2, p-value = 0.1105
# no significant difference
```

Copepod abundance **did not differ significantly across DO bins**, suggesting DO category was not a strong predictor of Copepod abundance.

Kruskal-Wallis test between abundance and bins for Ostracods

```
kruskal.test(log_ave_lit ~ bins,
             data = aq_ins_clean_bins,
             subset = taxon == "Ostracod")
```

Kruskal-Wallis rank sum test

data: log_ave_lit by bins

Kruskal-Wallis chi-squared = 6.7235, df = 2, p-value = 0.03467

```
# Kruskal-Wallis chi-squared = 6.7235, df = 2, p-value = 0.03467
# significant difference of abundance btwn DO level bins
```

Ostracod abundance **differed significantly across DO bins**, indicating that Ostracods may respond differently to low, moderate, and high DO conditions.

Pairwise test between abundance and individual bins for Ostracods

```
bins_ostracods <- aq_ins_clean_bins |>
  filter(taxon == "Ostracod")
pairwise.wilcox.test(
  bins_ostracods$log_ave_lit,
  bins_ostracods$bins,
  p.adjust.method = "bonferroni")
```

Pairwise comparisons using Wilcoxon rank sum test with continuity correction

data: bins_ostracods\$log_ave_lit and bins_ostracods\$bins

	low	moderate
moderate	0.387	-
high	1.000	0.051

P value adjustment method: bonferroni

```
# p = 0.051 for comparison btwn moderate and high DO level bins
```

After Bonferroni correction, the moderate vs. high DO comparison was close to significant, suggesting a possible difference in Ostracod abundance between these bins but not strong enough to conclude significance at $\alpha = 0.05$.

Chi-Square

```
# create abundance table for chi-square test
chi_square_table <- aq_ins_clean_bins |>
  filter(taxon %in% focus_taxa) |>
  group_by(bins, taxon) |>
  summarise(
    total_abundance = sum(ave_lit, na.rm = TRUE),
    .groups = "drop"
  ) |>
  pivot_wider(
    names_from = taxon,
    values_from = total_abundance,
    values_fill = 0
  ) |>
  column_to_rownames("bins") |>
  as.matrix()

view(chi_square_table)

# show table used in chi-square test
chi_square_table
```

	Cladocera	Copepod	Hemiptera	Corixidae	Boatman	Nematode	Ostracod
low	980.2667	115.8667			0.2666667	0.1333333	1202.933333
moderate	510.6667	1170.4000			83.8666667	0.2666667	1057.866667
high	264.2667	1453.4667			8.8000000	0.0000000	7.866667

```
# chi-square test for differences in taxon composition across DO bins
chi_square_test <- chisq.test(chi_square_table)

# show chi-square result
chi_square_test
```

Pearson's Chi-squared test

```
data: chi_square_table
X-squared = 2859.2, df = 8, p-value < 2.2e-16
```

Taxon composition **differed significantly across DO bins**, suggesting that the invertebrate community structure changes depending on dissolved oxygen category.

Visualizations directly relevant to answering questions

Directly Relevant Figure 1: DO levels vs invert. abundance for each focus taxa

```
DO_taxa_medians <- DO_taxa |>
  # groups data by taxon so each species gets one median point
  group_by(taxon) |>
  # calculates median DO and median log abundance for each taxon
  summarise(
    med_do_point = median(med_do, na.rm = TRUE),
    med_log_ave_lit = median(log_ave_lit, na.rm = TRUE),
    .groups = "drop"
  )

DO_taxa_plot <- ggplot(data = DO_taxa, # assigns plot to object do_taxa_plot
  # uses DO_taxa data frame
  mapping = aes(x = med_do, # x-axis: med_do
    y = log_ave_lit, # y-axis: log_ave_lit
    color = taxon)) + # color by taxon


# first layer: points show individual abundance measurements
geom_point(shape = 16, # shape is circle
  size = 2, # size is 2
  alpha = 0.7) + # opacity is 0.7

# second layer: larger point shows the median location for each taxon
geom_point(data = DO_taxa_medians,
  mapping = aes(x = med_do_point,
    y = med_log_ave_lit),
  shape = 18,
  size = 5) +

# separate panels for each taxon with free x- and y-axis scales
facet_wrap(~ taxon, scales = "free") +
# use non-default colors for points
scale_color_viridis_d() +
labs(x = "Median DO (mg/L)", # relabel x-axis
  y = "log + 1 transformed average abundance/L", # relabel y-axis
  # creates descriptive title
  title = "Invertebrate abundance as a function of DO") +

# uses non-default theme
```

```
theme_minimal() +
# remove redundant legend
theme(legend.position = "none")
DO_taxa_plot
```



Directly Relevant Figure 2: DO levels vs total invert. abundance

```
# Summed abundance across focus taxa vs DO
focus_taxa <- c("Cladocera", "Copepod",
                 "Hemiptera Corixidae Boatman",
                 "Nematode", "Ostracod")

total_abundance <- DO_taxa |>
```

```

filter(taxon %in% focus_taxa) |>
group_by(date, site, med_do) |>
summarize(
  total_abundance = sum(ave_lit, na.rm = TRUE),
  .groups = "drop"
) |>
mutate(log_total_abundance = log10(total_abundance + 1))

# fit linear model
total_abundance_lm <- lm(log_total_abundance ~ med_do,
  data = total_abundance)

# create new data frame for predicted values
total_abundance_predictions <- data.frame(
  med_do = seq(min(total_abundance$med_do, na.rm = TRUE),
    max(total_abundance$med_do, na.rm = TRUE),
    length.out = 100)
)

# generate predicted values and confidence intervals
total_abundance_predictions <- total_abundance_predictions |>
  bind_cols(
    predict(total_abundance_lm,
      newdata = total_abundance_predictions,
      interval = "confidence") |>
      as.data.frame()
  )

# plot raw data and model predictions
ggplot(total_abundance,
  aes(x = med_do,
    y = log_total_abundance)) +

# raw data points
geom_point(color = "magenta3",
  size = 3,
  alpha = 0.7) +

# confidence interval around model predictions
geom_ribbon(data = total_abundance_predictions,
  aes(x = med_do,
    ymin = lwr,

```

```


        ymax = upr),
        inherit.aes = FALSE,
        fill = "cyan2",
        alpha = 0.35) +

# predicted line from linear model
geom_line(data = total_abundance_predictions,
          aes(x = med_do,
              y = fit),
          inherit.aes = FALSE,
          color = "purple4",
          linewidth = 1) +

labs(
  x = "Dissolved oxygen (mg/L)",
  y = "Log total invertebrate abundance",
  title = "Relationship between dissolved oxygen and total invertebrate abundance"
) +

theme_classic()

```

```
broom::tidy(total_abundance_lm)
```

```
# A tibble: 2 x 5
```

	term	estimate	std.error	statistic	p.value
	<chr>	<dbl>	<dbl>	<dbl>	<dbl>
1	(Intercept)	2.01	0.302	6.65	0.000000170
2	med_do	-0.0723	0.0316	-2.29	0.0289

Directly Relevant Figure 3: Abundance vs DO with binned levels

```
total_abundance <- total_abundance |>
mutate(
  do_category = case_when(
```

```

    med_do <= quantile(med_do, 0.25, na.rm = TRUE) ~ "Low DO",
    med_do <= quantile(med_do, 0.75, na.rm = TRUE) ~ "Medium DO",
    TRUE ~ "High DO"
  )
)
total_abundance$do_category <- factor(
  total_abundance$do_category,
  levels = c("Low DO", "Medium DO", "High DO")
)

ggplot(total_abundance,
  aes(x = do_category,
    y = log_total_abundance,
    fill = do_category)) +

  geom_boxplot(alpha = 0.7,
    outlier.color = "black",
    outlier.size = 2) +


  # optional: add jittered points to show raw data
  geom_jitter(width = 0.15,
    alpha = 0.5,
    size = 2,
    color = "black") +

  labs(
    x = "Dissolved oxygen level",
    y = "Log total invertebrate abundance",
    title = "Invertebrate abundance across dissolved oxygen levels"
  ) +

  scale_fill_manual(values = c(
    "Low DO" = "red3",
    "Medium DO" = "grey60",
    "High DO" = "blue3"
  )) +

  theme_classic() +
  theme(legend.position = "none")

```



Other exploratory visualizations

Exploratory Figure 1: Variance of DO across sights


```
# create object aq_ins_clean_fig2 from aq_ins_clean
aq_ins_clean_fig2 <- aq_ins_clean |>
  # convert taxon to title case
  mutate(taxon = str_replace_all(taxon, "_", " "),
         taxon = str_to_title(taxon)) |>
  # filter to include focus taxa
  filter(taxon %in% focus_taxa)

# base layer
DO <- ggplot(data = aq_ins_clean_fig2,
  # x-axis: site
  # y-axis: mean salinity
  mapping = aes(x = site_full,
                y = med_do)) +
  # first layer: boxplot
  geom_boxplot(width = 0.35,
```

```

        outlier.shape = NA) +
# second layer: salinity measurements at each survey
geom_jitter(height = 0,
            width = 0.12,
            shape = 21,
            alpha = 0.6) +
# wrapping x-axis text
scale_x_discrete(labels = label_wrap(width = 15)) +
# second layer: points to represent means
stat_summary(fun = median,
            color = "red",
            size = 1) +
# relabelling x- and y-axes
labs(x = "Site",
     y = "Median DO (mh/L)",
     title = "Site DO levels") +
# different theme
theme_bw()
# show plot
DO

```



Exploratory Figure 2: Composition plot across DO bins


```
composition_plot <- DO_taxa |>
  mutate(
    do_category = case_when(
      med_do <= quantile(med_do, 0.25, na.rm = TRUE) ~ "Low DO",
      med_do <= quantile(med_do, 0.75, na.rm = TRUE) ~ "Medium DO",
      TRUE ~ "High DO"
    ),
    do_category = factor(do_category,
                        levels = c("Low DO",
                                   "Medium DO",
                                   "High DO"))
  )
```

```

) |>
group_by(do_category, taxon) |>
summarize(
  total_abundance = sum(ave_lit, na.rm = TRUE),
  .groups = "drop"
) |>
ggplot(aes(x = do_category,
           y = total_abundance,
           fill = taxon)) +
geom_col(position = "fill") +
scale_y_continuous(labels = scales::percent_format()) +
labs(
  x = "Dissolved oxygen category",
  y = "Relative community composition",
  fill = "Taxon",
  title = "Invertebrate community composition shifts across dissolved oxygen levels"
) +
theme_classic()

# display plot
composition_plot

```



Exploratory Figure 3: Time-series plot of total abundance (all species)

```
# species/taxa to include in the total abundance figure
species_interest <- c(
  "Ostracod",
  "Copepod",
  "Cladocera",
  "Hemiptera Corixidae Boatman",
  "Nematode"
)


# create total abundance over time
total_abundance_time <- DO_taxa |>
```

```

# keeps only the taxa of interest
filter(taxon %in% species_interest) |>
# makes sure date is treated as a date
mutate(date = as.Date(date)) |>
# groups by date only, so abundance is summed across sites and taxa
group_by(date) |>
# sums average abundance per liter across all taxa of interest
summarise(
  total_abundance = sum(ave_lit, na.rm = TRUE),
  .groups = "drop"
)

# plot total abundance over time
ggplot(total_abundance_time, aes(x = date, y = total_abundance)) +
  # adds a line connecting total abundance values over time
  geom_line(linewidth = 1) +
  # adds points for each sampling date
  geom_point(size = 2) +
  # formats y-axis labels with commas
  scale_y_continuous(labels = scales::comma) +
  # adds clear labels
  labs(
    title = "Total Abundance of Species of Interest Over Time",
    x = "Sampling Date",
    y = "Total Abundance"
  ) +
  # uses a clean theme
  theme_minimal()

```

Exploratory Figure 4: Time-series area plot

```
# species/taxa to include in the area plot
species_interest <- c(
  "Ostracod",
  "Copepod",
  "Cladocera",
  "Hemiptera Corixidae Boatman",
  "Nematode"
)


# create abundance over time by taxon
taxon_abundance_time <- DO_taxa |>
  # keeps only the taxa of interest
  filter(taxon %in% species_interest) |>
  # makes sure date is treated as a date
  mutate(date = as.Date(date)) |>
  # groups by date and taxon so each species stays separate
  group_by(date, taxon) |>
  # sums across sites, but not across taxa
  summarise(
```

```

    ave_lit = sum(ave_lit, na.rm = TRUE),
    .groups = "drop"
  )

# create time series area plot by taxon
ggplot(taxon_abundance_time, aes(x = date, y = ave_lit, fill = taxon)) +
  # creates stacked area plot with a different color for each taxon
  geom_area(alpha = 0.8) +
  # formats y-axis labels with commas
  scale_y_continuous(labels = scales::comma) +
  # adds clear labels
  labs(
    title = "Abundance of Species of Interest Over Time",
    x = "Sampling Date",
    y = "Average Abundance per Liter",
    fill = "Taxon"
  ) +
  # uses a clean theme
  theme_minimal()

```



Exploratory Figure 5: Distribution of Median Dissolved Oxygen Across Sites

```
# create new object to store medians of median do levels
median_fig2 <- aq_ins_clean_fig2 |>
  group_by(site_full) |>
  summarize(median_do = median(med_do, na.rm = TRUE))

# assign quartiles for medians for each site
median_fig2$quartile <- cut(
  median_fig2$median_do,
  breaks = quantile(median_fig2$median_do, probs = seq(0, 1, 0.25)),
  labels = c("Q1", "Q2", "Q3", "Q4"),
  include.lowest = TRUE,
  right = FALSE)


# boundaries of quartiles
q <- quantile(median_fig2$median_do, probs = seq(0, 1, 0.25))

# distribution plot of DO levels across sites
ggplot(median_fig2, aes(x = median_do)) +
  # density curve
  geom_density(fill = "lightblue",
               alpha = 0.4,
               color = "blue",
               linewidth = 1.2) +
  # vertical quartile lines
  geom_vline(xintercept = q,
             linetype = "dashed",
             color = "gray40") +
  # site labels
  geom_text_repel(aes(y = 0.01,
                     label = str_wrap(site_full, width = 10)),
                 size = 3,
                 box.padding = 0.5,      # space around the text box
                 show.legend = FALSE) +
  # quartile dashed line labels
  geom_text(data = data.frame(x = q, label = names(q)),
            aes(x = x, y = 0.14, label = paste0(label, "\n", round(x, 2))),
            color = "gray40",
            size = 3.5,
            hjust = -0.1) + # shift text slightly to the right of the line
  # remove space between bottom of plot and y-axis, set y-axis limits
```

```

scale_y_continuous(expand = c(0,0),
                   limits = c(0, 0.15)) +
# axes labels and title
labs(x = "Median Dissolved Oxygen (mg/L)",
     y = "Density",
     title = "Distribution of Median Dissolved Oxygen Aross Sites") +
# theme
theme_bw() +
theme(legend.position = "none")

```



These bins were created by taking the median value of the median dissolved oxygen levels at each site. Low DO is the 25th percentile, moderate DO is the 75th percentile, and high DO is the 100th percentile.

- Low DO: Phelps Road and North of Phelps Bridge

- Moderate DO: South of Venoco Bridge, South of Phelps Bridge, Devereux Creek
- High DO: North of Venoco Bridge, Main Channel

Exploratory Figure 6: Invertebrate Abundance in DO Bins

```
aq_ins_clean_bins <- aq_ins_clean |>
  mutate(bins = case_when(site_full == "Phelps Road" ~ "low",
                          site_full == "North of Phelps Bridge" ~ "low",
                          site_full == "South of Venoco Bridge" ~ "moderate",
                          site_full == "South of Phelps Bridge" ~ "moderate",
                          site_full == "Devereux Creek" ~ "moderate",
                          site_full == "North of Venoco Bridge" ~ "high",
                          site_full == "Main Channel" ~ "high")) |>
  mutate(taxon = str_replace_all(taxon, "_", " "), taxon = str_to_title(taxon)) |>
  filter(taxon %in% focus_taxa) |>
  # create column that calculates log-transformed abundance
  mutate(log_ave_lit = log(ave_lit + 1)) |>
  mutate(bins = factor(bins, levels = c("low", "moderate", "high")))

# base layer: ggplot
ins_bins_plot <- ggplot(data = aq_ins_clean_bins,
                       mapping = aes(x = bins, # x-axis: DO bins
                                     y = log_ave_lit, # y-axis: log_ave_lit
                                     color = taxon)) + # color by taxon

# first layer: points show individual abundance measurements
geom_jitter(width = 0.15,
            height = 0,
            shape = 16,
            size = 2,
            alpha = 0.7) +

# calculate median for each bin for each taxon
stat_summary(fun = median,
            geom = "point",
            color = "red",
            size = 3) +

# separate panels for each taxon
facet_wrap(~ taxon) +

# use non-default colors for points
scale_color_viridis_d() +


labs(x = "DO levels", # relabel x-axis
```

```

y = "log + 1 transformed average abundance/L", # relabel y-axis
# creates descriptive title
title = "Invertebrate abundance vs DO levels") +
# uses non-default theme
theme_bw() +
# remove redundant legend
theme(legend.position = "none")

# shows plot
ins_bins_plot

```




Plan for elective (if completing)

- should include some description/image of drafts
- should include plan for making progress in the next two weeks

Elective idea: add microscopic images of each of the 5 main inverts


Water color bubbles/ blow colored bubbles with watercolor around each to demonstrate how much dissolved oxygen each likes

- More bubbles = likes more DO
- Less bubbles = likes less DO




Most common inverts

1. Ostracod




a.

2. Copepod



a.

3. Cladocera



a.

4. Corixidae (Water Boatmen)



a.

5. Nematode

Plan for next 2 weeks: NEED TO ADD